
Digitized by the Internet Archive
in 2025

https://archive.org/details/index-kewensis_1895_2

ROYAL ZOOLOGICAL INSTITUTE
16 NOV 1970

IMPERIAL BUREAU
13 MAR. 1925
OF MYCOLOGY

INDEX KEWENSIS

INDEX KEWENSIS

AN ENUMERATION OF THE GENERA AND SPECIES

OF

FLOWERING PLANTS

FROM THE TIME OF LINNAEUS TO THE YEAR 1885 INCLUSIVE
TOGETHER WITH THEIR AUTHORS' NAMES, THE WORKS IN
WHICH THEY WERE FIRST PUBLISHED, THEIR NATIVE COUNTRIES
AND THEIR SYNONYMS

COMPILED AT THE EXPENSE OF
THE LATE CHARLES ROBERT DARWIN
UNDER THE DIRECTION OF JOSEPH D. HOOKER

BY

B. DAYDON JACKSON

VOLUME II

OXFORD
AT THE CLARENDON PRESS
M DCCC XCV

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM MDCCCLXXXV

COMPLECTENS

NOMINE RECEPTO AUCTORE PATRIA

UNICUIQUE PLANTAE SUBJECTIS

SUMPTIBUS BEATI CAROLI ROBERTI DARWIN

DUCTU ET CONSILIO JOSEPHI D. HOOKER

CONFECIT B. DAYDON JACKSON

TOMUS II

OXONII

E PRELO CLARENDONIANO

M DCCCXCV

LONDINI
APUD HENRICUM FROWDE



NOVI EBORACI
APUD A. MACMILLAN ET SOCIOS

OXONII: EXCUDEBAT HORATIUS HART, TYPOGRAPHICUS ACADEMICUS

ADDENDA ET EMENDANDA

QUAEDAM

III

(MENDA NON COMMEMORATA LECTOR BENEVOLENS IPSE CORRIGET)

a. = *adde.* d. = *dele.* l. = *lege.* p. = *pone.*

PAG.

KAKOSMANTHUS.
2[1]. l. *macrophyllus*, . . . 578 = P. macrophylla.

3[3]. insere KATAKIDOZAMIA, Haage & Schmidt, ex Regel, Gartenfl. (1876) 5 = CATAKIDOZAMIA, T. Hill = **Macrozamia**, Miq. (Cycad.).

KERATEPHORUS.
5[3]. l. *Leerii*, . . . 579 = P. Leerii.

KLEINIA.
8[1]. a. *Mandraliscae*, Tineo, in Ann. Agr. Sicil.; ex Lojac. Isole Eolie, 111.—Hab.?

10[2]. l. KODDA-PAIL, Adans. Fam. ii. 75 (non 25).

KOELREUTERIA.
11[3]. l. **Marsdensia**, R. Br. (Asclep.).

KOPSIA, Dum.
12[3]. l. *aegyptiaca*, . . . = O. aegyptiaca.
arenaria, . . . = O. arenaria.
Borkhauseni, . . . = O. arenaria.
lavandulacea, . . . = O. lavandulacea.
longiflora, . . . = O. aegyptiaca.
ramosa, . . . = O. ramosa.
a. *stricta*, Caruel, in Parl. Fl. Ital. vi. 356 = O. Schulzii.

LACHENALIA.
18[2]. l. pallida, *Soland.* etc.

LACHNOPETALUM.
19[1]. l. *glabrum*, . . . l. c. = R. Lachnopetala.

LACHNOSTOMA.
19[2]. a. cynanchoides, *Benth. & Hook. f. Gen.* ii. 766.—Am. bor. occ.
l. maritimum, *Benth. & Hook. f.* (ead. cit.); *Nichols.* etc.

LANNEOMA.
28[2] a. *triphylla*, March. Rev. Anacard. 157, sphalm. = seq.

LASIANTHERA.
34[1]. a. malaccensis, *Mast. l. c.* 584.—Malacca.

LATHYRUS.
37[3]. l. *ensifolius*, Badarro, etc.
38[1]. l. *macrorrhizus*, Wimm. . . . = montanus.
39[1]. a. Tommasinii, *Sprun. ex Marches. in Boll. Soc. Adriat. Sc. Nat. Trieste*, v. (1880) p. xxviii, nomen.—Europ.

PAG.

LAURIDIA.
40[1]. l. *rupicola*, . . . (= *Myroxylum reticulatum*).—Afr. austr.

LAVATERA.
44[1]. l. *rotundata*, Láz. . . . (1881), etc.

LEDUM.
47[3]. l. *californicum*, Kellogg, . . . 14 = glandulosum.

LEIANTHUS.
49[1]. l. OMPHALOSTIGMA, Reichb. Nom. 132 (1841).

LEIDESIA.
49[2]. a. *Sonderiana*, Muell. Arg. in DC. Prod. xv. II. 699 = capensis.

LEOCHILUS.
51[2]. a. pygmaeus, *Benth. & Hook. f. Gen.* iii. 565.—Guatemala.

LEPIDAGATHIS.
55[3]. l. *nervosa*, Wight, etc.
56[1]. l. subuninervia, *C. B. Clarke*, etc.

LEPIDIUM.
56[2]. a. *bipinnatum*, Gibert, Enum. Pl. Montevid. 66 = praec.

LEPIDOTHECA.
59[1]. a. *Santolina*, Walp. Rep. ii. 639, sphalm. = seq.

LEPIGONUM.
59[3]. a. *roseum*, Rafin. Fl. Tellur. iii. 81 = Arenaria geniculata?

LEPTADENIA.
60[3]. l. Spartium, *Wight* (et in synon. passim).

66[3]. a. LERISCA, Schlecht. in Bot. Zeit. iii. (1845) 476 = **Criptangium**, Schrad. (Cyperac.).

LETTSOMIA.
68[2]. l. *sumatrana*, . . . = Ipomoea staphylina.

LEUCADENDRON.
68[2]. a. *conocarpum*, Steud. Nom. ed. II. ii. 399, in syn. = Leucospermum conocarpum.
68[3]. a. *ellipticum*, Steud. (ead. cit.) = Leucospermum ellipticum.

LEUCAS.
70[1]. l. *dimidiata*, Benth. in Wall. Cat. n. 2521; et in DC. etc.
70[2]. a. *pallida*, Steud. Nom. ed. II. ii., in syn. = Leonotis pallida.

PAG.

LEUCOJUM.
71[3]. a. *aestivale*, Steud. Nom. ed. II. ii. 196, in syn. = seq.

LEUCOMERIS.
71[3]. a. *scandens*, Sch. Bip. in Flora, xxxvi. (1854) 275 = Erigeron scandens.

72[1]. a. LEUCONOCARPUS, Spruce, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 295 = **Moronobea**, Aubl. (Guttif.).
riparius, Spruce, l. c. = M. riparia.

LEUCOSPERMUM.
74[1]. l. involucratum, *Roem. & Schult.*
a. *tortum*, Steud. Nom. ed. II. ii. 401, in syn., sphalm = Leucadendron tortum.

74[2]. linea prima l. **LEUCOSYKE.**

LIBANOTIS.
76[2]. l. *chachroides*, Steud. . . . = *cachroides*.

LIBIDIBIA.
76[3]. l. *coriaria*, . . . = C. coriaria.

79[1]. l. LIGIA, Fasano, in Att. Acc. Nap. 1787 (1788) 245 [t. 18. f. 2].

LIGUSTICUM.
79[2]. l. *angustilobum*, Phil. . . . 95 = Pimpinella angustilobum.

LILIORHIZA.
81[2]. a. *Piattiana*, Kellogg, ex Curran, in Bull. Calif. Acad. i. IV. (1885) 150 = F. pluriflora?
a. *viridia*, Kellogg, in Proc. Calif. Acad. ii. (1863) 48 = F. lanceolata.

LIMODORUM.
85[2]. a. *mauritanicum*, Sw. Adnot. Bot. 59 = Angraecum gladiifolium?

LIMOSELLA.
86[2]. a. *lacustris*, Steud. Nom. ed. II. ii. 350, in syn., sphalm. = Littorella juncea.

LINARIA.
87[2]. l. italica, *Trevir. Ind. Sem. Hort. Vratisl. App.* ii. (1820) 2; *et in*, etc.

LIPARIS.
98[3]. a. *diphyllos*, Nimmo, in J. Grah. Cat. Bomb. Fl. 252 = Habenaria diphylla.
l. Galeottiana, *Hemsl.* . . . 559.—Mexic.

LIPOCHAETA.
94[3]. l. *hastata*, Kellogg, . . . = Verbesina remota.

PAG.

LIQUIDAMBAR.
96[2]. l. *peregrina*, . . . = Myrica asplenifolia.

LISTERA.
98[1]. a. *multinervia*, Peterm. in Flora, xxvii. (1844) 369 = ovata.

LOASA.
102[1]. l. *aurantiaca*, Hort. ex Lilja, in Linnaea, xv. (1841) 263, etc.
102[2]. l. punicea, *Phil.* . . . (1884), etc.

LOBELIA.
102 . l. *adenophora*, . . . = Lysipomia aretioides.
104[2]. l. *glandulosa*, . . . (eadem).
104[3]. l. *limoselloides*, . . . (eadem).
105[3]. l. Rhynchopetalum, *Hemsl.*
106[2]. l. *subulata*, Benth. . . . 137 = Rhizocephalium subulatum.

LOMATIA.
109[2]. l. *chilensis*, C. Gay, . . . 310 = Orites myrtoidea.

LONCHOCARPUS.
110[1]. l. Neuroscapha, *Benth.*
110[2]. l. *Pterocarpus*, . . . 260 = Phyllocarpus Pterocarpus.
a. *scandens*, Steud. Nom. ed. II. ii. 461, in syn., sphalm. = L. Nicou.

LOPHOPHYLLUM.
113[2]. l. bicristatum, *Griff.*

LUMNITZERA.
124[2]. l. coccinea, *Wight & Arn. Prod.* 316, etc.

LUNDIA.
124[3]. l. *monachanthe*, ... = O. spinosa.

LYCOPUS.
132[2]. l. *laciniatus*, Rouy, in La Natural. ii. (1882) (Matér. Fl. Portug.) 507 = europaeus.

MACROCNEMIUM.
141[3]. l. *exsertum*, Oerst. . . . 45 = Pogonopus exsertus.

143[1]. a. MACROSCAPA, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 148 = **Stropholirion**, Torr. (Liliac.).
volubilis, Kellogg, l. c. = S. californicum.

MAIANTHEMUM.
147[1]. l. *Bifolium*, DC. Fl. Fr. iii. 177; [DC.] etc.

MALACHRA.
148[1]. a. *diversifolia*, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 134; xii. (1845) 79 = Pavonia diversifolia.

MALESHERBIA.
150[1]. l. *paniculata*, . . . 323, etc.

MALLOTUS.
150[2]. l. Apelta, *Muell. Arg.*
l. *chinensis*, Muell. Arg. in DC. Prod. xv. II. 965, in syn. etc.
japonicus, *Muell. Arg.* . . .—Japon.
150[3]. l. *micranthus*, Muell. Arg.
tiliaefolius, *Muell. Arg.*

MANDEVILLA.
160[3]. a. Wrightiana, *Benth. & Hook. f. Gen.* ii. 727.—Cuba.

MAPPA.
164[3]. l. *peltata*, Wight, etc.

MAPPIA.
l. *montana*, Miers, l. c. 398 (Apodytis sp.).—Java.

MARAH.
165[1]. l. *muricata*, Kellogg,

MARIANTHUS.
168[1]. l. ONCOSPORUM, Putterl. in Endl. Nov. Stirp. Dec. 9 (1839).

MARSDENIA.
171[2]. l. tinctoria, *R. Br.* . . . 30.—As. trop. (non Afr.).

MARTYNIA.
172[3]. l. Proboscidea, *Glox.*

MASSONIA.
174[3]. a. *nivea*, Burch. ex Salisb. Gen. Pl. Fragm. 17 = jasminiflora?

PAG.

185[3]. a. MEGAPHYLLUM, Spruce, ex Baill. Hist. Pl. vii. 457, in nota (1880) = **Pentagonia**, Benth. (Rubiac.).

189[2]. l. MELANDRIUM, . . . ed. II. ii. 37, 274 (1812).
l. *Flos-cuculi*, . . . ed. II. ii. 275 =
189[3]. l. *pratense*, . . . ed. II. ii. 274 =
sylvestre, . . . ed. II. ii. 274 =

MELASTOMA.
192[2]. l. *chulesis*, . . . = Osbeckia nepalensis.

MELIA.
197[1]. l. *Bambolo*, C. DC. in DC. Monog. i. 767 = seq. *Bombolo*, Welw. etc.

MELICOCCA.
198[3]. a. *javanica*, Hassk. Hort. Bog. Desc. i. 138 = Otophora javanica.

MELOCACTUS.
201[2]. a. approximatus, *Suring. in Versl. en Med. Kon. Akad. Wetensch.* III. ii. (1885) 187.—Ind. occ.
cornutus, *Suring.* (*l. c.*) 191.—Ind. occ.
Evertizianus, *Suring.* (*l. c.*) 190.—Ind. occ.
ferox, Suring. (l. c.) 185 = rubellus.
hexacanthus, Suring. (l. c.) = rubellus.
201[3]. intermedius, *Suring.* (*l. c.*) 192.—Ind. occ.
Koolwijkianus, *Suring.* (*l. c.*) 184.—Ind. occ.
parvispinus, *Suring.* (*l. c.*) 183.—Ind. occ.
patens, *Suring.* (*l. c.*) 190.—Ind. occ.
pusillus, *Suring.* (*l. c.*) 192.—Ind. occ
rectiusculus, *Suring.* (*l. c.*) 187.—Ind. occ.
reversus, *Suring.* (*l. c.*).—Ind. occ.
rubellus, *Suring.* (*l. c.*) 184.—Ind. occ.
spatanginus, *Suring.* (*l. c.*) 193.—Ind. occ.
stramineus, *Suring.* (*l. c.*) 185.—Ind. occ.
202[1]. *trichacanthus*, Suring. (l. c.) 186 = stramineus?

MELOCHIA.
202[2]. a. corymbosa, *Hemsl. Biol. Centr. Am. Bot.* i. 130.—Mexic.

MENTHA.
210[2]. l. Welwitschii, *Rouy, in La Natural.* ii. (1882) (*Matér. Fl. Portug.*) 47.—Europ.

212[3]. l. MERIDIANA, [Linn.] Linn. f. etc.

MERTENSIA.
213[3]. l. *virginica*, . . . = pulmonarioides.

MESPILUS.
218[3]. a. *media*, Bechst. Forstb. 150, 334 = Crataegus Oxyacantha.

METACHILUM.
220[1]. l. *cyathiferum*, Lindl. l. c. etc.
cyathiforme, Steud. Nom. ed. II. ii. 136 = praec.

METROXYLON.
222[2]. l. Pigafettia, in synon. passim.—Cf. *elatum*, *filare*, *microcarpum*, *microspermum*.

MICRANTHUS.
228[1]. l. *oppositifolius*, Wendl.

MICROCARPAEA.
228[2]. a. *chamaedrifolia*, Steud. Nom. ed. II. ii. 246, in syn. = seq.

MICROGLOSSA.
229[2]. l. altissima, *DC.* . . . = Psiadia altissima.
hispida, DC. . . . = Psiadia hispida.

MICROLICIA.
230[3]. a. *thymifolia*, Steud. Nom. ed. II. ii. 447, in syn., sphalm. = serpyllifolia.

MICROLOMA.
a. *hastatum*, Decne. in DC. Prod. viii. 499 = sagittatum.
231[1]. l. *pyrotechnicum*, . . . = Leptadenia Spartium.

247[3]. a. MIRKOOA, Wight, Ic. i. t. 217, in textu (1840) = **Ammannia**, Houst. (Lythrar.).
rotundifolia, Wight, l. c. = A. rotundifolia.

MITREOLA.
250[1]. a. *lanceolata*, Torr. ex Croom, Cat. Pl. New Bern, 45 = petiolata.

PAG.

MNASIUM.
250[2]. l. *unilaterale*, . . . = Spathanthus unilateralis.

MOEHRINGIA.
251[3]. a. *sedifolia*, Freyn, in Oestr. Bot. Zeitschr. xxvi. (1876) 227 = *papulosa*.

MONNIERA.
257[2]. a. *longifolia*, Steud. Nom. ed. II. ii. 373, sphalm. = Monnina longifolia.

MONNIERIA.
257[2]. a. *trifoliata*, Baill. Adansonia, xi. (1875) 276 = praec.

MUEHLENBECKIA.
268[3]. l. *Benthamii*, Endl. Gen. Suppl. iv. II. 51, etc.
quadrangulata, (eadem).
volcanica, (eadem).

MURALTIA.
271[1]. a. *ericoides*, Steud. Nom. ed. II. ii. 371, in syn., sphalm. = Burmanni.
l. laxa, *DC.* etc.

MUSA.
272[1]. a. Charlioi, *E. Hill*, *Rep. Brisbane Bot. Gard.* (1874) 7.—Hab.?
a. *Rhinozerotis*, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 64 = nana?

MYCETIA.
275[2]. a. *javanica*, Hook. f. Fl. Brit. Ind. iii. 95, sphalm. = praec.

NARCISSUS.
293[2]. a. *biternaceus*, Salisb. Gen. Pl. Fragm. 100 = incomparabilis?

NECTANDRA.
299[2]. a. *decandra*, Roxb. Fl. Ind. i. 425 = Linostoma decandrum.

NEPETA.
305[3]. l. dasyangula, . . . 12.—Reg. Himal.
309[3]. a. NESOEA, Wight, Ic., in textu sub t. 217 (1840) = **Ammannia**, Houst. (Lythrar.).
octandra, Wight, l. c. = A. octandra.

NIDORELLA.
315[1]. l. *altissima*, Benth. & Hook. f. . . . = Psiadia altissima.
l. *hispida*, Benth. & Hook. f. . . . = Psiadia hispida.

NIDULARIUM.
315[1]. l. *caeruleum*, Lem.

NIVENIA.
315[3]. l. PARANOMUS, . . . (1807).

NOLANA.
316[2]. l. triquetra, . . . *App.* 12.

NYCTAGO.
320[3]. l. *angulata*, . . . (= *Quamoclidion angulatum*).—Mexic.

NYMPHAEA.
321[2]. l. *Fenzliana*, . . . = Rudgeana.

OCOTEA.
326[2]. l. Bofo, *H. B. & K.*

ODONTOGLOSSUM.
330[3]. l. *pygmaeum*, Lindl. . . . = Leochilus pygmaeus.

OENANTHE.
331[3]. l. *amoena*, Sweet, Hort. Brit. ed. II. 247, sphalm. = Oenothera amoena.

OENOTHERA.
333[3]. l. erosa, *Lehm. Sem. Hort. Hamb.* (1824) 20; *et in*, etc.

OLEARIA.
338[1]. l. Allomii, *T. Kirk*, . . . i. ed. II. (1871) 92, etc.

ONCIDIUM.
341[3]. l. *acrobotryum*, Klotzsch, . . . = Harrisonianum.

OPHRYOSPORUS.
354[1]. l. Pachychaete, *Baker.*

PAG.

OPLISMENUS.

355[3]. l. PANICULUM, Ard.

359[2]. l. ORCHIDIUM, [Sw.] Summa Veg. Scand. 32 (1814); et in, etc.
l. *arcticum*, [Sw.] (ead. cit.).

OSBECKIA.

379[3]. a. *pendulifolia*, Steud. Nom. ed. II. ii. 447, in syn. = Aciotis pendulifolia.

OXYTROPIS.

396[1]. l. Middendorffii, *Trautv. in Middend. Sibir. Reise* (*Fl. Taimyr.* 49. t. 7).—Sibir.

396[3]. l. *Trautvetteri*, Meinsh. . . . = Middendorffii.

PALICOUREA.

404[1]. l. quadrifolia, *DC. Prod.* iv. 526.—Guian.

PANAX.

406[1]. l. *micranthum*, Wall. Cat. n. 4938 = Leschenaultii.

PANICUM.

411[2]. l. *capillaceum*, Lam. . . . = brevifolium, *Linn.*

412[2]. l. decompositum, *R. Br. Prod.* 191.—Austral.

412[2]. a. *dentispica*, Kunth, Rév. Gram. i. sub t. 39 = *densispica.*

413[3]. l. *Gardneri*, Thw. . . . = Isachne Gardneri.

414[2]. l. *hirtum*, Nees, . . . = brevifolium, *Linn.*

414[3]. l. *insulicola*, Steud. . . . = auritum.

415[3]. l. *lanuginosum*, . . . = mollicomum.

416[1]. l. mollicomum, *Kunth.*

PAG.

PANICUM :—

417[1]. d. *oxyzoides*, Nees, etc.
a. *panis*, Jess. Deutschl. Graes. 240 = Setaria italica.

417[3]. l. *plicatum*, Roxb.
l. *polystachion*, Linn. . . . = setosum.

418[3]. l. *Roxburghianum*, Schult. . . . = repens.

419[1]. l. sericeum, [*Soland.*] . . . Ind. occ.

419[3]. l. tomentosum, *Roxb.* . . . *Fl. Ind.* i. 301.

PARAPOTTSIA.

424[1]. l. = Pottsia cantonensis (*bis*).

PARIETARIA.

425[3]. l. *microphylla*, Linn. . . . = Pilea muscosa.

PASPALUM.

429[3]. l. arundinaceum, *Poir.* . . . Am. bor.

430[2]. l. *ctenostachyum*, Trin. . . . = Eriochloa grandiflora.

PATRINIA.

438[2]. l. scabiosaefolia, *Fisch. ex Trevir. Ind. Sem. Hort. Vratisl. App.* ii. (1820) 2.—Dahuria.

PATRISIA.
l. *bicolor*, DC. . . . = R. Candollei.
l. *parviflora*, DC. . . . = R. Patrisii.

PAVETTA.

440[2]. l. Gerrardi, *Harv.*

PAG.

PEDALIUM.

444[2]. l. *filiforme*, Pav. . . . 267 (= *Ischnia verbenacea*). —Mexic.

PELARGONIUM.

452[1]. l. *viastum*, Steud. . . . = australe.

PELIOSTOMUM.

456[2]. l. *scoparium*, . . . Anticharis scoparia (in Addend.).

PENTAGONIA.

459[3]. l. MEGAPHYLLUM, Spruce . . . 457, etc.

473[1]. a. PEROSTIS, Beauv. Agrost. 5 (1812) = Perotis, Ait. (Gramin.).
laxifolia, Beauv. l. c. 6 = Perotis latifolia.

PETALIDIUM.

476[2]. a. Currori, *Benth. ex S. Moore, l. c.* 226.—Afr. austr.

PIMPINELLA.

529[1]. l. *ramosissima*, Fisch. . . . Hort. Vratisl. (1819) 3, etc.

PINANGA.

529[2]. a. caudata, *Becc. Malesia*, i. (1877) 101, in obs.—N. Guin.

PLEURANTHIUM.

569[1]. l. Adendrobium, *Benth. & Hook. f.*

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM M.DCCC.LXXXV

COMPLECTENS

KABLIKIA—KAGENECKIA.

KABLIKIA, Opiz, Seznam, 55 (1852) = **Primula**, Linn.
minima, Opiz, l. c. = P. minima.

KADAKIA, Rafin. Fl. Tellur. ii. 9 (1836) = **Pontederia**, Linn.

KADALI, Adans. Fam. ii. 234 (1763) = **Osbeckia**, Linn. (Melast.).

KADALIA, Rafin. Sylva Tellur. 101 (1838) = **Dissotis**, Benth. (1849) (Melastom.).
antennina, Rafin. l. c. = D. antennina.
rotundifolia, Rafin. l. c. = D. rotundifolia.

KADSURA, Kaempf. ex Juss. in Ann. Mus. Par. xvi. (1810) 340. *MAGNOLIACEAE*, Benth. & Hook. f. i. 19.
SARCOCARPON, Blume, Bijdr. 21 (1825).
cauliflora, *Blume, Fl. Jav. Schiz.* 11.—Java.
chinensis, Hance, ex Benth. Fl. Hongk. 8 = Schisandra Hanceana.
chinensis, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 149 = Schisandra chinensis.
grandiflora, Wall. Tent. Fl. Nap. 10. t. 14 = Schisandra grandiflora.
japonica, Benth. in Hook. Kew Journ. iii. (1851) 258 = Schisandra chinensis.
japonica, Benth. Fl. Hongk. 8 = Schisandra Hanceana.
japonica, *Juss. in Ann. Mus. Par.* xvi. (1810) 340; *Dun. Monog.* 57.—Japon.
japonica, Wall. Tent. Fl. Nap. 12 = Roxburghiana.
propinqua, Wall. l. c. 11. t. 15 = Schisandra propinqua.
pubescens, *Miq. Fl. Ind. Bat. Suppl.* 620.—Sumatra.
Roxburghiana, *Arn. in Mag. Zool. & Bot.* ii. (1838) 546.—Reg. Himal.
scandens, *Blume, Fl. Jav. Schiz.* 9.—Java.
Wightiana, *Arn. in Mag. Zool. & Bot.* ii. (1838) 546.—Ind. or.

KADUA, Cham. & Schlecht. in Linnaea, iv. (1829) 157. *RUBIACEAE*, Benth. & Hook. f. ii. 61.
WIEGMANNIA, Meyen, Reise, ii. 139 (1835).
acuminata, *Cham. & Schlecht. in Linnaea*, iv. (1829) 163.—Ins. Sandvic.
affinis, Cham. & Schlecht. l. c. 164 = Gouldia sandwicensis.
Arnottii, G. Don, Gen. Syst. iii. 533 = Menziesiana.
centranthoides, *Hook. & Arn. Bot. Beech. Voy.* 85.—Ins. Sandvic.
Cookiana, *Cham. & Schlecht. in Linnaea*, iv. (1829) 158.—Ins. Sandvic.
cordata, *Cham. & Schlecht. l. c.* 160.—Ins. Sandvic.
glaucifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 318.—Ins. Sandvic.

KADUA:—
glomerata, *Hook. & Arn. Bot. Beech. Voy.* 85.—Ins. Sandvic.
grandis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 318.—Ins. Sandvic.
Kaalae, Wawra, in Flora, lvii. (1874) 273 = grandis.
laxiflora, *H. Mann, in Proc. Am. Acad.* vii. (1867) 171.—Ins. Sandvic.
Menziesiana, *Cham. & Schlecht. in Linnaea*, iv. (1829) 160.—Ins. Sandvic.
parvula, *A. Gray, in Proc. Am. Acad.* iv. (1860) 317.—Ins. Sandvic.
petiolata, A. Gray, l. c. 318 = acuminata, grandis.
Romanzoffiensis, Cham. & Schlecht. in Linnaea, iv. (1829) 162 = Gouldia Romanzoffiensis.
Smithii, Hook. & Arn. Bot. Beech. Voy. 86 = Menziesiana.
Waimeae, *Wawra, in Flora*, lvii. (1874) 264.—Ins. Sandvic.

KADULA, Rafin. Fl. Tellur. iv. 90 (1836) = **Cuscuta**, Tourn. (Convolv.).
corymbosa, Rafin. l. c. = C. corymbosa.

KADURIAS, Rafin. Fl. Tellur. iv. 91 (1836) = **Cuscuta**, Tourn. (Convolv.).
reflexa, Rafin. l. c. = C. reflexa.

KAELERIA, Boiss. Diagn. Ser. II. iv. 134 (1856) sphalm. = **Koeleria**, Pers. (Gramin.).

KAEMPFERA, Houst. Reliq. 3. t. 2 (1781) = **Tamonea**, Aubl. (Verben.).

KAEMPFERA, Spreng. Syst. i. 9 (1825) = seq.

KAEMPFERIA, Linn. Gen. ed. I. 331 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 641.
CADALVENA, Fenzl, in Sitzb. Akad. Math.-Nat. Wien, li. Abth. II. (1865) 139.
CIENKOWSKYA, Solms, in Schweinf. Beitr. Fl. Aethiop. 197 (1867).
MONOLOPHUS, Wall. Pl. As. Rar. i. 24. t. 24 (1830).
TRITOPHUS, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 341.
angustifolia, *Rosc. in Trans. Linn. Soc.* viii. (1807) 351.—As. trop.
candida, *Wall. Pl. As. Rar.* i. 47. t. 56.—Burma.
Crawfurdia, Wall. ex Horan. Prod. Scitam. 22 = elegans.
diversifolia, Link, ex A. Dietr. Sp. Pl. i. 57 = ovalifolia.
elegans, *Wall. Cat.* n. 6593.—Burma.
Galanga, *Linn. Sp. Pl.* 3.—As. trop.
Gilberti, *Hort. Bull.—Cf. Gard. Chron.* (1882) I. 712.—Ind. or.

KAEMPFERIA:—
Hedychium, Lam. Tabl. Encyc. i. 7 = Hedychium coronarium.
humilis, Salisb. Prod. 6 = K. Galanga.
lanceolata, *Lodd. ex Steud. Nom.* ed. II. i. 843 (Quid?).—Ind. or.
latifolia, Donn, Hort. Cantab. ed. III. 1; ed. VII. 3 = K. Galanga.
linearis, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 19.—Reg. Himal.
longa, Jacq. Hort. Schoenb. iii. 37 = rotunda.
marginata, *Carey, in Rosc. Scit. Pl.* t. 93.—Burma.
ornata, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) 159. t. 537.—Borneo.
ovalifolia, *Roxb. Hort. Beng.* 1; *Pl. Corom.* iii. 75.—Burma; Malaya.
ovata, Rosc. in Trans. Linn. Soc. viii. (1807) 351 = pandurata.
pandurata, *Roxb. in As. Res.* xi. (1810) 328.—As. trop.
Parishii, Hook. f. Bot. Mag. t. 5763 = ovalifolia.
parviflora, *Wall. Cat.* n. 6587.—Burma.
plantaginea, Steud. Nom. ed. II. i. 843 = seq.
plantaginifolia, Salisb. in Trans. Hort. Soc. i. (1812) 286 = K. Galanga.
procumbens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = K. Galanga.
purpurea, *Koen. in Retz. Obs.* iii. 57.—Ind. or.
Roscoeana, *Wall. in Bot. Reg.* t. 1212.—Burma.
Roscoei, Wall. Cat. n. 6584 = praec.
rosea, *Schweinf. ex Benth. & Hook. f. Gen.* iii. 642.—Afr. trop.
rotunda, Blanco, Fl. Filip. ed. I. 5 = K. Galanga.
rotunda, Donn, Hort. Cantab. ed. III. 1 = pandurata.
rotunda, *Linn. Sp. Pl.* 3.—As. trop.
Roxburghiana, Schult. Mant. i. 33 = angustifolia.
scaposa, *Benth. & Hook. f. Gen.* iii. 642.—Ind. or.
secunda, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 19.—Reg. Himal.; Burma.
speciosa, Thunb. Fl. Javan. 8 = Costus speciosus?
undulata, Link, Enum. Hort. Berol. i. 3 = angustifolia.
undulata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 391.—Hab.?
versicolor, Salisb. in Trans. Hort. Soc. i. (1812) 286 = rotunda.
vittata, *N. E. Br. in Gard. Chron.* (1882) II. 264.—Sumatra.

KAGENACKIA, Steud. Nom. ed. II. i. 844 (1840), sphalm. = seq.

KAGENECKIA, Ruiz & Pav. Prod. 145. t. 37 (1794). *ROSACEAE*, Benth. & Hook. f. i. 614.
LYDEA, Molina, Sagg. Chile, ed. II. 14 (1810).
amygdalifolia, *Presl, Epim. Bot.* 195.—Peruv.

KAGENECKIA :—
angustifolia, *D. Don, in Edinb. N. Phil. Journ.* xiii. (Oct.–Dec. 1831) 111.—Chili.
crataegifolia, Lindl. Bot. Reg. t. 1836 = oblonga.
crataegoides, D. Don, in Edinb. N. Phil. Journ. (Oct.–Dec. 1831) 111 = oblonga.
glutinosa, *H. B. & K. Nov. Gen. et Sp.* vi. 237.—Peruv.
lanceolata, *Ruiz & Pav. Syst. Veg.* 290.—Peruv.
oblonga, *Ruiz & Pav. l. c.* 289.—Chili.
ovata, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 64.—Chili.

KAHIRIA, Forsk. Fl. Aegypt. Arab. 153 (1775) = **Ethulia**, Linn. (Compos.).

KAKILE, Desf. Fl. Atlant. ii. 77 (1798) = **Cakile**, Tourn. (Crucif.).

KAKOSMANTHUS, Hassk. Retzia, 97 (1855); et in Flora, xxxviii. (1855) 577 = **Payena**, Linn. (Sapotac.).
macrophyllus, Hassk. ll. cc. 97, 578.—Java.

ALABOTIS, Rafin. Fl. Tellur. ii. 19 (1826) = **Allium**, Tourn. (Liliac.).
canadensis, Rafin. l. c. = A canadense.
Clusianum, Rafin. l. c. = A. subhirsutum.
flavum, Rafin. l. c. = A. flavum.
Moly, Rafin. l. c. = A. Moly.
nigrum, Rafin. l. c. = A. nigrum.
pallens, Rafin. l. c. = A. oleraceum.
paniculatum, Rafin. l. c. = A. paniculatum.

KALANCHOË, Adans. Fam. ii. 248 (1763) *CRASSULACEAE*, Benth. & Hook. f. i. 659.
CALANCHOË, Pers. Syn. i. 445 (1805).
VEREIA, Andr. Bot. Rep. t. 21 (1797).
abrupta, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 512.—Ins. Socotr.
acutiflora, *Haw. Syn. Pl. Succ.* 109.—Ind. or.?
acutiflora, Kurz, in Journ. As. Soc. Beng. (1876) II. 309 = spathulata.
aegyptiaca, DC. Pl. Grass. t. 64 = *crenata*.
Afzeliana, *Britten, in Oliver, Fl. Trop. Afr.* ii. 393.—Afr. trop.
alternans, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. ii. 380 = thyrsiflora.
alternans, *Pers. Syn.* i. 446.—Arab.
amplexicaulis, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia amplexicaulis.
amplexicaulis, Heyne, in Wall. Cat. n. 7221 = floribunda, *Wight*.
angustifolia, *A. Rich. Tent. Fl. Abyss.* i. 313.—Afr. trop.
biternata, Wight, ex Wall. Cat. n. 7221 = laciniata.
brachycalyx, *A. Rich. Tent. Fl. Abyss.* i. 312.—Afr. trop.
brachyloba, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 392.—Afr. trop.
brasiliensis, *Cambess. in St. Hil. Fl. Bras. Pl. Mer.* ii. 196.—Bras.
campanulata, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia campanulata.
ceratophylla, *Haw. Rev. Pl. Succ.* 23.—China.
coccinea, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 395.—Afr. trop.
cochleatum, *Lem. Illustr. Hortic.* vi. (1859), *Misc.* 100.—Hab.?
corymbosa, Herb. Madr. ex Wall. Cat. n. 7224 = spathulata?
crenata, Britten, in Oliver, Fl. Trop. Afr. ii. 394, partim = spathulata, brasiliensis.
crenata, *Haw. Syn. Pl. Succ.* 109.—Afr. trop.
delagoënsis, Eckl. & Zeyh. Enum. 305 = Bryophyllum tubiflorum.
eriophylla, *Hilsenb. & Boj. ex Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 149.—Madag.
farinacea, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 512.—Ins. Socotr.
floribunda, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 150.—Madag.
floribunda, *Wight & Arn. Prod.* 359.—Ind. or.
floripendula, Steud. Nom. ed. II. i. 252 = Bryophyllum calycinum.
glandulosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 312.—Ind. or.; Afr. trop.
glaucescens, *Britten, in Oliver, Fl. Trop. Afr.* ii. 393.—Afr. trop.

KALANCHOË :—
gracilipes, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia gracilipes.
gracilis, *Hance, in Journ. Bot.* viii. (1870) 6.—Ins. Formosa.
grandiflora, *A. Rich. Tent. Fl. Abyss.* i. 310.—Afr. trop.
grandiflora, *Wight & Arn. Prod.* 359; *Wall. Cat.* n. 7226.—Ind. or.
heterophylla, Wight & Arn. l. c. 360 = glandulosa.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 468.—Madag.
hirta, *Harv. in Harv. & Sond. Fl. Cap.* ii. 397.—Afr. austr.
humilis, *Britten, in Oliver, Fl. Trop. Afr.* ii. 397.—Afr. trop.
hybrida, Desf. ex Steud. Nom. ed. I. 135 = spathulata.
integerrima, *Lange, Ind. Sem. Hort. Haun.* (1872) 5, adn.; *et in Bot. Tidsskr.* (1878) 139.—Afr. austr.?
laciniata, *DC. Pl. Grass.* t. 100.—As. et Afr. trop.
lanceolata, *Pers. Syn.* i. 446.—Arab.
macrosepala, *Hance, in Journ. Bot.* viii. (1870) 5.—Ins. Hongkong.
miniata, Hilsenb. & Boj. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 149 = Kitchingia miniata.
modesta, Kotschy & Peyr. Pl. Tinn. 18 = brachycalyx.
multiceps, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 469.—Madag.
nudicaulis, Buch.-Ham. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 414 = spathulata.
oblongifolia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 379.—Afr. austr.
orgyalis, *Baker, in Journ. Bot.* xx. (1882) 110.—Madag.
panduriformis, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia panduriformis.
paniculata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 380.—Afr. austr.
parviflora, Baill. in Bull. Soc. Linn. Par. i. (1885) 468 = Kitchingia parviflora.
peltata, Baill. l. c. = Kitchingia peltata.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 311 = crenata.
pinnata, Pers. Syn. i. 446 = Bryophyllum calycinum.
platysepala, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 393.—Afr. trop.
porphyrocalyx, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia porphyrocalyx.
pubescens, R. Br. ex Oliver, Fl. Trop. Afr. ii. 396 = brachycalyx.
pumila, *Baker, in Journ. Linn. Soc.* xx. (1883) 139.—Madag.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* i. 311.—Abyssin.
Ritchieana, Dalz. in Hook. Kew Journ. iv. (1852) 346 = glandulosa.
robusta, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 512.—Ins. Socotr.
rotundifolia, *Haw. in Phil. Mag.* (1825) 31.—Afr. austr.
scapigera, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 397.—Afr. trop.
Schimperiana, *A. Rich. Tent. Fl. Abyss.* i. 310.—Abyssin.
schizophylla, Baill. in Bull. Soc. Linn. Par. i. (1885) 469 = Kitchingia schizophylla.
spathulata, *DC. Pl. Grass.* t. 65.—As. trop.
spathulata, Wall. Cat. n. 7224 = floribunda.
stenosiphon, *Britten, in Oliver, Fl. Trop. Afr.* ii. 395.—Afr. trop.
subamplectens, Wall. Pl. As. Rar. ii. t. 167 = spathulata.
synsepala, *Baker, in Journ. Bot.* xx. (1882) 110.—Madag.
teretifolia, Haw. in Wall. Pl. As. Rar. ii. 53. t. 166 = laciniata.
thyrsiflora, *Harv. in Harv. & Sond. Fl. Cap.* ii. 380.—Afr. austr.
tomentosa, *Baker, in Journ. Bot.* xx. (1882) 110.—Madag.
trichantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 140.—Madag.
varians, Haw. in Phil. Mag. N. S. ix. (1829) 302 = spathulata.
velutina, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 396.—Afr. trop.
Verea, Pers. Syn. i. 446 = crenata.
Welwitschii, *Britten, in Oliver, Fl. Trop. Afr.* ii. 394.—Afr. trop.
Wightianum, Wall. Cat. n. 7225 = floribunda.

KALAWAEL, Adans. Fam. ii. 344 (1763) = **Connarus**, Linn.

KALBFUSSIA, Sch. Bip. in Flora, xvi. (1833) 723 = **Leontodon**, Linn. (Compos.).
algeriensis, Pomel, Nouv. Mat. Fl. Atl. 10.—Alger.
Kralikii, Pomel, l. c. 11.—Alger.
Muelleri, Sch. Bip. in Ann. Sc. Nat. Sér. I. i. (1834) 378 = L. Salzmanni.
oranensis, Pomel, Nouv. Mat. Fl. Atl. 268.—Alger.
orientalis, Jaub. & Spach, Illustr. iii. 117 = L. hispidulus.
parvifolia, Pomel, Nouv. Mat. Fl. Atl. 11.—Alger.
Reboudiana, Pomel, l. c. 268.—Alger.
Salzmanni, Pomel, l. c. 10, 268 = *oranensis*.
Sultzmanni, Sch. Bip. in Ann. Sc. Nat. Sér. I. i. (1834) 378 = L. Salzmanni.

KALENICZENKIA, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 252 = **Brachysema**, R. Br. (Legumin.).
daviesioides, Turcz. l. c. = B. daviesioides.

KALERIA, Adans. Fam. ii. 506 (1763) = **Silene**, Linn. (Caryoph.).

KALI, Tourn. ex Adans. Fam. ii. 261 (1763) = **Salsola**, Linn. (Chenopod.).
inermis, Moench, Meth. 331 = S. Soda.
rosacea, Moench, Meth. Suppl. 115 = S. rosacea.
Soda, Moench, Meth. 331 = S. Kali.
Soda, Scop. Fl. Carn. ed. II. i. 175 = S. Soda.
Tragus, Scop. l. c. = S. Kali.

KALIDIUM, Moq. in DC. Prod. xiii. II. 146 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 63.
arabicum, *Moq. l. c.* 147.—Arab.
caspicum, *Ung.-Sternb. in Atti Congr. Bot. Firenz.* 1874 (1876) 317.—Reg. Casp.; Turkest.
circassianum, Ledeb. ex Ung.-Sternb. l. c. 320 = Schrenkianum.
foliatum, *Moq. in DC. Prod.* xiii. II. 147.—Ross.; As. bor.
gracile, *Fenzl, in Ledeb. Fl. Ross.* iii. 769.—Mongolia.
Pallasianum, Bunge, ex Ung.-Sternb. Verz. Syst. Salic. 95 = Schrenkianum.
Schrenkianum, *Bunge, ex Ung.-Sternb. l. c.*—Turkest.

KALIMARES, Rafin. Fl. Tellur. ii. 46 (1836) = **Aster**, Tourn. (Compos.).
Amellus, Rafin. l. c. = A. Amellus.

KALIMERIS, Cass. in Dict. Sc. Nat. xxiv. 324 (1822) = **Aster**, Tourn. (Compos.).
platycephala, Cass. l. c. 325 = A. incisus.

KALIPHORA, Hook. f. in Benth. & Hook. f. Gen. i. 951 (1867). *CORNACEAE*, Benth. & Hook. f. l. c.
madagascariensis, *Hook. f. in Hook. Ic. Pl.* t. 1023.—Madag.

KALLIAS, Cass. in Dict. Sc. Nat. xxiv. 326 (1822) = **Heliopsis**, Pers. (Compos.).
dubia, Cass. l. c. 333 = H. scabra.
ovata, Cass. l. c. 327 = H. buphthalmoides.

KALLOPHYLLON, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 365 (1876) = **Symphyopappus**, Turcz. (Compos.).
crenatum, Pohl, ex Baker, l. c. 367 = S. reticulatus.
ovatum, Pohl, ex Baker, l. c. 368 = S. reticulatus.

KALLSTROEMIA, Scop. Introd. 212 (1777) = **Tribulus**, Linn. (Zygophyll.).
cistoides, Endl. in Ann. Wien. Mus. i. (1836) 184 = T. cistoides.
grandiflora, Torr. ex A. Gray, Pl. Wright. i. 28 = T. grandiflorus.
maxima, Wight & Arn. Prod. 145 = T. maximus.
minor, Hook. f. in Hook. Niger Fl. 269 = T. maximus.
tribuloides, Wight & Arn. Prod. 145 = T. brasiliensis.
Tribulus, Meissn. Gen. Comm. 43 = T. maximus.

KALMIA, Linn. Diss. Chen. 18 (1750); Amoen. Acad. iii. 13 (1756). *ERICACEAE*, Benth. & Hook. f. ii. 596.

KALMIA:—
angustifolia, *Linn. Sp. Pl.* 391.—Am. bor.
ciliata, Bartr. ex Dum.-Cours. Bot. Cult. ed. II. iii. 323 = hirsuta.
cuneata, *Michx. Fl. Bor. Am.* i. 257.—Am. bor.
ericoides, *Griseb. Cat. Pl. Cub.* 51.—Cuba.
glauca, *Ait. Hort. Kew.* ed. I. ii. 64. t. 8.—Am. bor.
hirsuta, *Walt. Fl. Carol.* 138.—Am. bor.
intermedia, *Lange, Ind. Sem. Hort. Haun. App.* (1873) 3.—Hab.?
latifolia, *Linn. Sp. Pl.* 391.—Am. bor.
myrtifolia, Andr. in Rev. Hortic. (1883) 10 = latifolia.
nitida, Forbes, Hort. Woburn. 93 = latifolia?
oleaefolia, Dum.-Cours. Bot. Cult. ed. II. iii. 322 = glauca.
polifolia, Wangenh. in Ges. Naturf. Fr. Berl. Schr. (1788) II. 130. t. 5 = glauca.
rosmarinifolia, Dum.-Cours. Bot. Cult. ed. I. ii. 250; ed. II. iii. 250 = glauca.
serotina, Hoffmgg. Verz. Pfl. 70 = latifolia?

KALOMIKTA, Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 219 = **Actinidia**, Lindl. (Ternstroem.).
mandshurica, Regel, l. c. = A. callosa.

KALOPANAX, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 16 (1863) = **Acanthopanax**, Decne. (Araliac.).
divaricatum, Miq. l. c. 17 = A. divaricatum.
innovans, Miq. l. c. = A. innovans.
resectum, Miq. l. c. 18 = Macropanax oreophilum?
ricinifolium, Miq. l. c. 16 = A. ricinifolium.
sumatranum, Miq. l. c. 17.—Sumatra.

KALOSANTHES, Haw. Rev. Pl. Succ. 6 (1821) = **Rochea**, DC. (Crassulac.).
bicolor, Haw. l. c. 7 = R. odoratissima.
biconvexa, Haw. in Phil. Mag. (1824) 185 = R. biconvexa.
capitata, Sweet, Hort. Brit. ed. II. 224 = R. odoratissima.
coccinea, Haw. Rev. Pl. Succ. 18 = R. coccinea.
cymosa, Haw. l. c. 7 = Crassula cymosa.
fascicularis, G. Don, Gen. Syst. iii. 108 = R. fascicularis.
flava, Sweet, Hort. Brit. ed. I. 178 = Crassula flava.
jasminea, Haw. Rev. Pl. Succ. 6 = R. jasminea.
media, Haw. l. c. 7 = R. versicolor.
odoratissima, Haw. l. c. 7 = R. odoratissima.
versicolor, Haw. l. c. = R. versicolor.

KALPANDRIA, Walp. Rep. i. 435 (1842) = CALPANDRIA, Blume = **Camellia**, Linn. (Ternstroem.).

KALYMOPETALON, Pohl, ex Flora, viii. (1825) 183, nomen (Quid?).

KAMBALA, Rafin. Sylva Tellur. 19 (1838) = **Sonneratia**, Linn. f. (Lythrar.).
pendula, Rafin. l. c. = S. apetala.

KAMELIA, Steud. Nom. ed. I. 455 (1821) = **Camellia**, Linn. (Ternstroem.).

KAMETTIA, Kostel. Allg. Med. Pharm. Fl. iii. 1062 (1834) = **Ellertonia**, Wight (1848) (Apocyn.).
malabarica, Kostel. l. c. = E. Rheedii.

KAMPMANNIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Zanthoxylum**, Linn. (Rutac.).
fraxinifolium, Rafin. l. c. = Z. Clava-Herculis.

KAMPMANNIA, Steud. Syn. Pl. Gram. 34 (1855) = **Arundo**, Tourn.? (Gramin.).
zeelandica, Steud. l. c. 35 = A. conspicua?

KAMPTZIA, Nees, Prog. Gratul. Kamptzia (1840); et in Nov. Act. Nat. Cur. xviii. Suppl. I. (1841) p. ix = **Syncarpia**, Tenore (Myrtac.).
albens, Nees, ll. cc. = S. laurifolia.

KANAHIA, R. Br. in Mem. Wern. Soc. i. (1809) 39. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 760.
Delilii, Kotschy, ex Decne. in DC. Prod. viii. 537 = laniflora.
Forskålii, Decne. l. c. = laniflora.
laniflora, *R. Br. in Salt, Abyss. App.* 64; *G. Don, Gen. Syst.* iv. 149.—Arab.; Abyssin.

KANDELIA, Wight & Arn. Prod. 310 (1834). *RHIZOPHOREAE*, Benth. & Hook. f. i. 679.
Rheedii, *Wight & Arn. l. c.* 311.—Ind. or.; Malaya.

KANDENA, Rafin. Sylva Tellur. 138 (1838) = **Azima**, Lam. (Salvador.).
spinosa, Rafin. l. c. = A. tetracantha.

KANDIS, Adans. Fam. ii. 422 (1763) = **Lepidium**, Tourn. (Crucif.).

KANILIA, Blume, Mus. Bot. Lugd. Bat. i. 140 (1849) = **Bruguiera**, Lam. (Rhizoph.).
caryophylloides, Blume, l. c. 141 = B. caryophylloides.
parviflora, Blume, l. c. 140 = B. parviflora.

KANIMIA, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 446. *COMPOSITAE*, Benth. & Hook. f. ii. 246.
corymbulosa, *Benth. & Hook. f. Gen.* ii. 247.—N. Granat.
erythralina, *Benth. & Hook. f. l. c.*—Bras.
gracilis, *Baker, in Mart. Fl. Bras.* vi. II. 371.—Bras.
nitida, Baker, l. c. 370 = erythralina.
oblongifolia, *Baker, l. c.* 369.—Bras.
palustris, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 447.—Bras.
Pohlii, *Baker, in Mart. Fl. Bras.* vii. II. 370.—Bras.
purpurascens, *Baker, l. c.* 371.—Bras.
strobolifera, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 447.—Bras.
thapsoides, *Benth. & Hook. f. Gen.* ii. 247.—Bras.

KANOPIKON, Rafin. Fl. Tellur. iv. 114 (1836) = **Euphorbia**, Linn.
atropurpureum, Rafin. l. c. = E. atropurpurea.

KANTEMON, Rafin. New Fl. Am. iii. 19 (1836) = **Lonicera**, Linn. (Caprifol.).
angustifolium, Rafin. l. c. = L. sempervirens.
ciliosum, Rafin. l. c. = L. ciliosa.

KANTUFFA, Bruce, Trav. t. 14 (1790) = **Pterolobium**, R. Br. (Legumin.).

KARA-ANGOLAM, Adans. Fam. ii. 84 (1763) = **Alangium**, Lam. (Cornac.).

KARAKA, Rafin. Sylva Tellur. 72 (1838) = **Sterculia**, Linn.
colorata, Rafin. l. c. = S. colorata.

KARAMYSCHEWIA, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 266 = **Oldenlandia**, Linn. (Rubiac.).
hedyotoides, Fisch. & Mey. l. c. = O. ramosissima.

KARAMYSCHOVIA, Fisch. ex Steud. Nom. ed. II. i. 844 (1840) = praec.

KARATAS, [Plum.] Mill. Gard. Dict. ed. VI (1752). *BROMELIACEAE*, Benth. & Hook. f. iii. 660.
NIDULARIUM, Lam. Jard. Fleur. (1846) Misc. iv. 60.
REGELIA, Lem. Illustr. Hortic. vii. (1860) sub t. 245, in nota.
acanthocrater, *Antoine, Bromel.* 49. tt. 29, 30.—Bras.
agavaefolia, *Devans. in Fl. des Serres*, Sér. II. xii. (1877) 233.—Guiana.
Binoti, *Antoine, Bromel.* t. 34.—Bras.
Carolinae, *Antoine, l. c.* 52. t. 31.—Bras.
coriacea, *Antoine, l. c.* 51. t. 30.—Bras.
cruenta, *Nichols. Dict. Gard.* ii. 216.—Bras.
fulgens, *Antoine, Bromel.* 41. t. 24.—Bras.
humilis, *E. Morr. in Belg. Hortic.* (1872) 131.—Mexic.; N. Granat.
Innocentii, *Antoine, Bromel.* 44. t. 24.—Bras.
Lagopus, E. Morr. ex Devans. in Fl. des Serres, Sér. II. xii. (1877) 233 = Plumieri?
Laurentii, *Antoine, Bromel.* 48. t. 28.—Bras.
Legrellae, *E. Morr. in Belg. Hortic.* (1872) 129.—Bras.
Meyendorffii, *Antoine, Bromel.* 54. t. 32.—Bras.
Morreniana, *Antoine, l. c.* t. 35.—Guiana.
olens, Nichols. Dict. Gard. ii. 216 = Carolinae.
Penguin, Mill. Gard. Dict. ed. VIII = Bromelia Pinguin.
Plumieri, *E. Morr. in Belg. Hortic.* (1872) 131.—Panama.
purpurea, *Antoine, Bromel.* 42. t. 25.—Bras.
Scheremetiewii, *Antoine, l. c.* 46. t. 27.—Bras.
spectabilis, *Antoine, l. c.* t. 33.—Bras.

KARDANOGLYPHOS, Schlecht. in Linnaea, xxviii. (1856) 472 = **Cardamine**, Linn. (Crucif.).
nana, Schlecht. l. c. = C. nana, *C. Muell.*

KARE-KANDEL, Adans. Fam. ii. 88 (1763) = **Carallia**, Roxb. (Rhizoph.).

KARELINIA, Less. in Linnaea, ix. (1834) 187 = **Pluchea**, Cass. (Compos.).
caspia, Less. l. c.—Reg. Casp.

KARIVIA, Arn. in Hook. Journ. Bot. iii. (1841) 275 = **Melothria**, Linn. (Cucurb.).
amplexicaulis, Arn. l. c. = M. amplexicaulis.
javanica, Miq. Fl. Ind. Bat. i. I. 661 = M. javanica.
longicirrha, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 80 = Actinostemma lobatum.
Rheedii, M. Roem. Syn. Pepon. 45 = M. heterophylla.
samoënsis, A. Gray, Bot. U. St. Expl. Exped. i. 643 = M. Grayana.
sinuosa, M. Roem. Syn. Pepon. 46 = M. heterophylla.
umbellata, Arn. in Hook. Journ. Bot. iii. (1841) 275 = M. heterophylla.

KARKANDELA, Rafin. Sylva Tellur. 106 (1838) = **Musa**, Linn.
malabarica, Rafin. l. c. = M. paradisiaca.

KARKINETRON, Rafin. Fl. Tellur. iii. 11 (1836) = **Muehlenbeckia**, Meissn. (Polygon.).
hastatum, Rafin. l. c. = M. sagittifolia.
undulatum, Rafin. l. c. = M. sagittifolia.

KARPATON, Rafin. Fl. Ludov. 78 (1817) (Ordo Ignot.). —Cf. Benth. & Hook. f. ii. 2.
hastatum, Rafin. l. c. 79.—Am. bor.

KARTHEMIA, Sch. Bip. in Walp. Rep. ii. 953 (1843) = VARTHEIMIA, DC. = **Iphiona**, Cass. (Compos.).

KARWINSKIA, Zucc. Nov. Stirp. i. 349. t. 16 (1832). *RHAMNEAE*, Benth. & Hook. f. i. 377.
affinis, Schlecht. in Linnaea, xv. (1841) 460 = Humboldtiana.
biniflora, Schlecht. l. c. = Humboldtiana.
glandulosa, Zucc. in Flora, vii. (1832) II. Beibl. 71 = Humboldtiana.
Humboldtiana, S. Wats. Pl. Wheeler. 7 = Rhamnus californica.
Humboldtiana, *Zucc. Nov. Stirp.* i. 351.—Am. bor. occ.
mollis, *Schlecht. in Linnaea*, xv. (1841) 461.—Mexic.
sessilifolia, *Schlecht. l. c.*—Mexic.
subcordata, *Schlecht. l. c.* 462.—Mexic.
umbellata, Schlecht. l. c. 460 = Humboldtiana?

KASAILO, Dennst. Schluess. Hort. Malab. 30 (1818) = **Ambelania**?, Aubl. (Apocyn.).
racemosa, Dennst. l. c.—Malabar.

KASTNERA, Sch. Bip. in Flora, xxxvi. (1853) 37 = **Liabum**, Adans. (Compos.).
tenera, Sch. Bip. l. c. 38.—N. Granat.

KATARSIS, Medic. Malv. 111 (1787) = **Gypsophila**, Linn. (Caryoph.).
arborea, Medic. l. c. = G. arborea.

KATOUTHEKA, Adans. Fam. ii. 159 (1763) = **Ardisia**, Sw. (Myrsin.).

KATOUTHEXA, Steud. Nom. ed. II. i. 844 (1840) = praec.

KATUBALA, Adans. Fam. ii. 67 (1763) = **Canna**, Linn. (Scitam.).

KAUFMANNIA, Regel, in Act. Hort. Petrop. iii. II. (1875) 293. *PRIMULACEAE*, Benth. & Hook. f. ii. 1239.
Semenovi, *Regel, l. c.*—Turkest.

KAULFUSSIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Xanthophyllum**, Roxb. (Polygal.).
geminiflora, Dennst. l. c. = X. flavescens.

KAULFUSSIA, Nees, Hor. Phys. Berol. 53. t. 11 (1820) = **Charieis**, Cass. (Compos.).
amelloides, Eckl. ex DC. Prod. v. 223 = Aster Bergerianus.

KAULFUSSIA:—
amelloides, Nees, Hor. Phys. Berol. 53. t. 11 = Charieis heterophylla.
ciliata, Spreng. f. Tent. Suppl. 25 = Gymnostephium ciliare.
strigosa, Spreng. f. l. c. = Aster strigosus.

KAVALAMA, Rafin. Sylva Tellur. 72 (1838) = **Sterculia**, Linn.
urens, Rafin. l. c. = S. urens.

KAYEA, Wall. Pl. As. Rar. iii. 5. t. 210 (1832). *GUTTIFERAE*, Benth. & Hook. f. i. 176.
Beccariana, *Baill. Adansonia*, xi. (1876) 369.—Borneo.
cuspidata, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 296 = stylosa.
eugeniaefolia, *Pierre, Fl. For. Cochinch.* t. 98.—Cambodia.
ferruginea, *Pierre, l. c.* t. 99.—Cambodia.
floribunda, *Wall. Pl. As. Rar.* iii. 5.—Reg. Himal.
hexapetala, Pierre, Fl. For. Cochinch. sub t. 98, in obs. = Calophyllum hexapetalum.
Korthalsiana, *Pierre, l. c.*—Malaya.
lepidota, *Pierre, l. c.*—Cochinch.?
macrantha, *Baill. Adansonia*, xi. (1876) 369.—Borneo.
macrocarpa, *Pierre, Fl. For. Cochinch.* t. 100.—Cochinch.
myrtifolia, *Baill. Adansonia*, xi. (1876) 368.—Borneo.
nervosa, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 277.—Burma; Malaya.
philippinensis, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 298 = floribunda.
racemosa, *Planch. & Triana, l. c.* 297.—Malaya.
stylosa, *Thw. Enum. Pl. Zeyl.* 50.—Zeylan.

KEDROSTIS, Medic. in Staatsw. Vorles. Churpf. Phys.-Ges. i. (1791) 229; et Phil. Bot. ii. 69 (1791). *CUCURBITACEAE*.
CONIANDRA, Schrad. in Eckl. & Zeyh. Enum. 275 (1836).
CYRTONEMA, Schrad. l. c. 276 (1836).
RHYNCHOCARPA, Schrad. in Linnaea, xii. (1838) 403.
africana, *Cogn. in DC. Monog. Phan.* iii. 643.—Afr. austr.
Bainesii, *Cogn. l. c.* 644.—Afr. trop.
digitata, *Cogn. l. c.* 639.—Afr. austr.
foetidissima, *Cogn. l. c.* 634.—Afr. trop.
glauca, Cogn. l. c. 640 = grossulariaefolia.
grossulariaefolia, *Presl, Bot. Bemerk.* 72.—Afr. austr.
hirtella, *Cogn. in DC. Monog. Phan.* iii. 644.—Abyssin.
mollis, *Cogn. l. c.* 637.—Afr. austr.
nana, *Cogn. l. c.*—Afr. austr.
punctulata, *Cogn. l. c.* 642.—Afr. austr.
rostrata, *Cogn. l. c.* 636.—Ind. or.
Zeyheri, *Cogn. l. c.* 641.—Afr. austr.

KEENANIA, Hook. f. Fl. Brit. Ind. iii. 101 (1881). *RUBIACEAE*.
modesta, *Hook. f. l. c.*—Reg. Himal.

KEERLIA, DC. Prod. v. 309 (1836), partim = **Aphanostephus**, DC. (Compos.).
linearifolia, DC. l. c. 310 = Xanthocephalum Alamani.
ramosa, DC. l. c. = Aphanostephus ramosus.
skirrhobasis, DC. in Deless. Ic. Sel. iv. t. 18 = Leucopsis arkansana.

KEERLIA, A. Gray & Engelm. in Proc. Am. Acad. i. (1848) 47. *COMPOSITAE*, Benth. & Hook. f. ii. 262.
bellidifolia, *A. Gray & Engelm. l. c.*—N. Mexic.
effusa, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1857) 222.—N. Mexic.

KEFERSTEINIA, Reichb. f. in Bot. Zeit. x. (1852) 633 = **Zygopetalum**, Hook. (Orchid.).
bicallosa, Reichb. f. Otia Bot. Hamb. 31, Ind. = Z. bicallosum.
expansa, Reichb. f. l. c. 9, in syn. = Z. expansum.
Gemma, Reichb. f. in Gard. Chron. (1874) i. 406 = Z. Gemma.
graminea, Reichb. f. in Bot. Zeit. x. (1852) 634 = Z. gramineum.
lactea, Reichb. f. in Gard. Chron. (1872) 1290 = Z. lacteum.
laminata, Reichb. f. l. c. (1885) ii. 70 = Z. laminatum.

KEFERSTEINIA:—
leucantha, Reichb. f. ex Linden, Illustr. Hortic. xxix. (1882) 52.—Venezuela.
Mystacina, Reichb. f. in Gard. Chron. (1881) i. 530 = Zygopetalum Mystacinum.
sanguinea, Pritz. Ic. Ind. ii. 161 = seq.
sanguinolenta, Reichb. f. in Bot. Zeit. x. (1852) 635 = Z. sanguinolentum.
stapelioides, Reichb. f. in Bot. Zeit. x. (1852) 634 = Z. Moritzii.
taurina, Reichb. f. in Linnaea, xli. (1877) 5 = Z. taurinum.

KEGELIA, Reichb. f. in Bot. Zeit. x. (1852) 670. *ORCHIDEAE*, Benth. & Hook. f. iii. 477.
Houtteana, *Reichb. f. l. c.* 671.—Guiana.

KEGELIA, Sch. Bip. in Linnaea, xxi. (1848) 245 = **Eleutheranthera**, Poit. (Compos.).
ruderalis, Sch. Bip. l. c. = E. ovata.

KEIRIA, Bowdich, Madeira, 259 (1825) (an Oleac.?).
lutea, Bowdich, l. c.—Ins. Mader.

KEISKEA, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 105 (1865). *LABIATAE*, Benth. & Hook. f. ii. 1181.
japonica, *Miq. l. c.*—Japon.

KEITHIA, Benth. Lab. Gen. et Sp. 409 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1189.
ERIOTHYMUS, J. A. Schmidt, in Mart. Fl. Bras. viii. 171. t. 32 (1858).
ciliata, *Benth. Lab. Gen. et Sp.* 732.—Mexic.
coccinea, *Benth. l. c.* 412.—Bras.
denudata, *Benth. l. c.*—Bras.
gracilis, *Benth. l. c.*—Bras.
lavanduloides, Benth. l. c. 411 = villosa.
marifolia, Schau. in Linnaea, xx. (1847) 705 = Poliomintha marifolia.
minima, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 173.—Bras.
nitida, *Benth. in DC. Prod.* xii. 247.—Bras.
pilosa, Benth. Lab. Gen. et Sp. 411 = scutellarioides.
rubiacea, *Benth. l. c.* 410.—Bras.
scutellarioides, *Benth. l. c.*—Bras.
villosa, *Benth. l. c.* 411.—Bras.

KEITHIA, Spreng. Neue Entdeck. iii. 57 (1822) (an Capparid.?).
brasiliensis, Spreng. l. c.—Bras.

KEITIA, Regel, in Act. Hort. Petrop. v. (1877) 639. *IRIDEAE*, Benth. & Hook. f. iii. 697.
natalensis, *Regel, l. c.* 640.—Afr. austr.

KELLAUA, A.DC. in Ann. Sc. Nat. Sér. II. xvi. (1842) 96 = **Euclea**, Linn. (Ebenac.).
Schimperi, A. DC. l. c. xviii. (1842) 209 = E. Kellau.

KELLERIA, Endl. Gen. Suppl. iv. ii. 61 (1847) = **Drapetes**, Banks (Thymel.).
Dieffenbachii, Endl. l. c. = D. Dieffenbachii.
villosa, Bergg. in Minneskr. Fisiog. Sällks. Lund (1877) n. viii. 17.—N. Zel.

KELLETTIA, Seem. Bot. Voy. Herald, 85, 254 (1853) = **Prockia**, Linn. (Tiliac.).
odorata, Seem. l. c. 85 = P. Crucis.
ovata, C. Muell. in Walp. Ann. vi. 1202, sphalm. = praec.

KELLOGGIA, Torr. ex Benth. & Hook. f. Gen. ii. 137 (1873); et in Bot. U. St. Expl. Exp. ii. 332. t. 6 (1874). *RUBIACEAE*, Benth. & Hook. f. ii. 137.
galioides, *Torr. l. c.*—Calif.

KEMELIA, Rafin. Sylva Tellur. 138, 139 (1838) = **Camellia**, Linn. (Ternstroem.).

KEMOXIS, Rafin. Sylva Tellur. 86 (1838) = **Vitis**, Tourn. (Ampelid.).
acida, Rafin. l. c. = V. acida.

KENDRICKIA, Hook. f. in Benth. & Hook. f. Gen. i. 751 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
Walkeri, *Hook. f. l. c.* 752.—Zeylan.

KENNEDYA, Vent. Jard. Malm. 104. t. 104 (1804) (Kennedia). *LEGUMINOSAE*, Benth. & Hook. f. i. 531.
AMPHODUS, Lindl. Bot. Reg. t. 1101 (1829).
CAULINIA, Moench, Meth. Suppl. 47 (1802).
PHYSOLOBIUM, Hueg. Bot. Archiv, t. 2 (1837).
ZICHYA, Hueg. l. c. t. 1 (1837).
arabica, Hochst. ex Benth. in Journ. Linn. Soc. viii. (1865) 265 = Teramnus labialis.
arenaria, Benth. in Enum. Pl. Hueg. 38 = prostrata.
Baumanni, Meissn. in Lehm. Pl. Preiss. i. 89 (Camptosematis sp.).—Austral.
Beckxiana, *F. Muell. Fragm.* xi. 98.—Austral.
bracteata, Gaudich. in Freyc. Voy. Bot. 486 = prostrata.
coccinea, *Vent. Jard. Malm.* 105. t. 105.—Austral.
Comptoniana, Link, Enum. Hort. Berol. ii. 235 = Hardenbergia Comptoniana.
cordata, Lindl. Bot. Reg. t. 944 = Hardenbergia monophylla.
dilatata, A. Cunn. ex Lindl. l. c. t. 1526 = coccinea.
eximia, *Lindl. in Paxt. Mag. Bot.* xvi. (1849) 35.—Austral.
Fredwoodii, *Hort. ex Verlot, in Rev. Hortic.* (1866) 279, 332.—Austral.
glabrata, *Lindl. Bot. Reg.* t. 1838.—Austral.
heterophylla, Mackay, ex Sweet, Hort. Brit. ed. I. 481 = coccinea.
inophylla, Lindl. Bot. Reg. t. 1421 = coccinea.
lateritia, F. Muell. Fragm. iv. 78 = macrophylla.
latifolia, Lindl. ex Steud. Nom. ed. II. i. 844 = Hardenbergia monophylla.
longiracemosa, Lodd. Bot. Cab. t. 1940 = Hardenbergia monophylla.
macrophylla, *Benth. Fl. Austral.* ii. 252.—Austral.
macrophylla, Lindl. Bot. Reg. t. 1862 = Hardenbergia Comptoniana.
Marryattae, Lindl. l. c. t. 1790 = prostrata.
Marryattiana, Maund, Botanist, i. 83 = praec.
microphylla, *Meissn. in Lehm. Pl. Preiss.* i. 91.—Austral.
microphylla, Sieber, ex Steud. Nom. ed. II. i. 845 = Glycine clandestina.
monophylla, Vent. Jard. Malm. 106. t. 106 = Hardenbergia monophylla.
monosperma, Steud. Nom. ed. II. i. 845 = Erythrina monosperma.
nigricans, *Lindl. Bot. Reg.* t. 1715.—Austral.
oblonga, *Lem. Hortic. Univ.* v. 127 (fide DC. Sched.).—Hab.?
oblongata, *Lam. ex Regel, Ind. Sem. Hort. Petrop.* (1861) 53 (an praec.?).—Hab.?
ovata, Sims, Bot. Mag. t. 2169 = Hardenbergia monophylla.
parviflora, *Meissn. in Lehm. Pl. Preiss.* i. 91.—Austral.
phaseolifolia, Hoffm. in Linnaea, xvi. (1842) Litt. 281 = rubicunda.
physolobioides, Walp. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 186 = parviflora?
procurrens, *Benth. in Mitch. Journ. Trop. Austral.* 365.—Austral.
prorepens, *F. Muell. Fragm.* viii. 225.—Austral.
prostrata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 299.—Austral.
retusa, *F. Muell. Fragm.* v. 106.—Austral.
rubicunda, *Vent. Jard. Malm.* 104. t. 104.—Austral.
sericea, *G. Don, Gen. Syst.* ii. 343.—Austral.
splendens, Meissn. in Lehm. Pl. Preiss. i. 89 (Camptosematis sp.).—Austral.
stipularis, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 421 = prostrata.
Stirlingii, *Lindl. Bot. Reg.* t. 1845.—Austral.
tabacina, Labill. Sert. Austr. Caled. 70. t. 70 = Glycine tabacina.

KENNEDYNELLA, Steud. Nom. ed. II. i. 845 (1840) = **Glycine**, Linn. (Legumin.).
clandestina, Steud. l. c. = G. clandestina.
elongata, Steud. l. c. = G. tabacina.
microphylla, Steud. l. c. = G. clandestina.
tabacina, Steud. l. c. = G. tabacina.
tomentosa, Steud. l. c. = G. tomentosa.

KENTIA, Adans. Fam. ii. 508 (1763) = **Trigonella**, Linn. (Legumin.).

KENTIA, Blume, in Bull. Néerl. (1838) 64; et Rumphia, ii. 94 (1843). *PALMAE*, Benth. & Hook. f. iii. 884.
acuminata, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 207 = Carpenteria acuminata.

KENTIA:—
australis, *Hort.—Cf. Gard. Chron.* (1873) 6.—Ins. D. Howe.
Balmoreana, C. Moore, in Illustr. Hortic. xxi. (1874) t. 191 = *Belmoreana*.
Baueri, Seem. Fl. Vit. 269 = Rhopalostylis Baueri.
Beccarii, F. Muell. Extra-Trop. Pl. Ind. ed. (1880) 163 = Nengella montana.
Belmoreana, F. Muell. Fragm. vii. 99 = Howea Belmoreana.
Canterburyana, F. Muell. l. c. 101 = Hedyscepe Canterburyana.
concinna, T. Moore, in Gard. Chron. (1877) i. 43 = Microkentia gracilis.
costata, Becc. Malesia, i. (1877) 36 = Gulubia costata.
Deplanchei, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 314 = Microkentia Deplanchei.
divaricata, Planch. ex Brongn. in Compt. Rend. Par. lxxvii. (1873) 398 = Kentiopsis divaricata.
elegans, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 312 = Cyphophoenix elegans.
Exorrhiza, H. Wendl. in Bonplandia, x. (1862) 191 = Exorrhiza Wendlandiana.
Forsteriana, F. Muell. ex H. Wendl. in Kerch. Palm. 248 = Howea Forsteriana.
fulcita, Brongn. in Compt. Rend. Par. lxxvii. (1873) 399 = Cyphophoenix fulcita.
gracilis, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 315 = Microkentia gracilis.
Humboldtiana, Brongn. ex H. Wendl. in Kerch. Palm. 248 = Cyphokentia Humboldtiana.
Joannis, F. Muell. ex H. Wendl. l. c. = Veitchia Joannis.
Lindeni, Hort. Linden, ex Illustr. Hortic. xxiv. (1877) t. 276 = Kentiopsis macrocarpa.
Luciani, *Linden.—Cf. Gard. Chron.* (1878) i. 440; *et Illustr. Hortic.* xxix. (1882) t. 451.—Hab.?
macrocarpa, Vieill. ex Brongn. in Compt. Rend. Par. lxxvii. (1873) 398 = Kentiopsis macrocarpa.
macrostachya, Panch. ex Brongn. l. c. 399 = Cyphokentia macrostachya.
minor, *F. Muell. Fragm.* viii. 235.—Austral.
moluccana, Becc. Malesia, i. (1877) 35 = Gulubia moluccana.
monostachya, F. Muell. Fragm. vii. 82 = Bacularia monostachya.
Mooreana, F. Muell. l. c. 101 = Clinostigma Mooreanum.
oleracea, Seem. ex H. Wendl. in Kerch. Palm. 248 = Oreodoxa oleracea?
olivaeformis, Brongn. & Gris, in Compt. Rend. Par. lxxvii. (1873) 399 = Kentiopsis olivaeformis.
Pancheri, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 316 = Microkentia Pancheri.
paradoxa, Mart. Hist. Nat. Palm. iii. 312 = Ophiria paradoxa.
polystemon, Panch. ex H. Wendl. in Kerch. Palm. 248 = Kentiopsis divaricata.
procera, *Blume, in Bull. Néerl.* (1838) 65.—N. Guinea.
robusta, Hort. Linden, ex H. Wendl. in Kerch. Palm. 248 = Cyphokentia Vieillardi.
rubricaulis, *Hort.—Cf. Gard. Chron.* (1876) i. 603.—Austral.
rupicola, *Hort. Linden, l. c.* (1878) i. 440.—Hab.?
sapida, Mart. Hist. Nat. Palm. iii. 312 = Rhopalostylis sapida.
Storckii, F. Muell. ex H. Wendl. in Kerch. Palm. 248 = Veitchia Storckii.
subglobosa, F. Muell. ex H. Wendl. l. c. = Veitchia subglobosa.
Vieillardi, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 313 = Cyphosperma Vieillardi.
Wendlandiana, F. Muell. Fragm. vii. 102 = Hydriastele Wendlandiana.

KENTIA, Steud. Nom. ed. II. i. 845 (1840) = **Fagraea**, Thunb. (Logan.).
morindaefolia, Steud. l. c. = F. racemosa.

KENTIOPSIS, Brongn. in Comp. Rend. Par. lxxvii. (1873) 398. *PALMAE*, Benth. & Hook. f. iii. 887.
divaricata, *Brongn. l. c.* (an al. gen. ?).—N. Caled.
Luciani, Rodigas, in Illustr. Hortic. xxix. (1882) sub t. 451 = Kentia Luciani.
macrocarpa, *Brongn. in Comp. Rend. Par.* lxxvii. (1873) 398.—N. Caled.
olivaeformis, *Brongn. l. c.*—N. Caled.

KENTRANTHUS, Neck. Elem. i. 122 (1790) = **Centranthus**, DC. (Valerian.).

KENTROPHYLLUM, Neck. Elem. i. 86 (1790) = **Carthamnus**, Linn. (Compos.).
aegyptiacum, Tausch, in Flora, xii. (1829) 71, nomen.—Aegypt.
album, Cass. in Dict. Sc. Nat. xxiv. 389 = C. leucocaulos.
alexandrinum, Boiss. Diagn. Ser. II. vi. 115 = C. glaucus.
anatolicum, Boiss. l. c. I. vi. 113 = C. glaucus.
arborescens, Hook. Bot. Mag. t. 3302 = C. arborescens.
aureum, Boiss. Diagn. Ser. I. x. 99 = C. flavescens.
baeticum, Boiss. & Reut. Pugill. Pl. Nov. 65 = C. lanatus.
canescens, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1855) 13 = C. canescens?
coeruleum, Gren. & Godr. Fl. Fr. ii. 264 = Carduncellus caeruleus.
creticum, Boiss. Diagn. Ser. I. x. 99 = C. glaucus.
creticum, Tausch, in Flora, xii. (1829) 71 = C. lanatus.
dentatum, DC. Prod. vi. 611 = C. dentatus.
elatum, Gasp. in Rendic. Soc. Borb. Nap. i. (1842) 186 = C. arborescens.
flavescens, Tausch, in Flora, xii. (1879) 71 = C. flavescens.
glaucum, Tausch, l. c. = C. glaucus.
incanum, Tausch, l. c. = C. lanatus.
lanatum, DC. & Duby, Bot. Gall. i. 293 = C. lanatus.
leucocaulon, DC. Prod. vi. 610 = C. leucocaulos.
luteum, Cass. in Dict. Sc. Nat. xxiv. 381 = C. lanatus.
mareoticum, DC. Prod. vi. 611 = C. mareoticus.
montanum, Pomel, Nouv. Mat. Fl. Atl. 22.—Alger.
purpureum, Tausch, in Flora, xii. (1829) 71, nomen.—Hab.?
rubrum, Link, in Linnaea, ix. (1834) 580 = C. dentatus.
syriacum, Boiss. Diagn. Ser. I. x. 98 = C. glaucus.
tauricum, C. A. Mey. Verz. Pfl. Cauc. 66 = C. lanatus.
tenue, Boiss. & Blanche, Diagn. Ser. II. iii. 50 = C. glaucus.
turbinatum, Gasp. ex Guss. Fl. Sic. Syn. ii. 430 = C. turbinatus.

KENTROPHYTA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 353 (1840) = **Astragalus**, Linn. (Legumin.).
montana, Nutt. ex Torr. & Gray, l. c. = A. Kentrophyta.
viridis, Nutt. l. c. = A. Kentrophyta.

KENTROPSIS, Moq. Chenop. Enum. 83 (1840) = **Sclerolaena**, R. Br. (Chenopod.).
brevicuspis, F. Muell. Fragm. iv. 150 = Anisacantha brevicuspis.
diacantha, Moq. in DC. Prod. xiii. ii. 138 = S. diacantha.
eriacantha, F. Muell. Fragm. ii. 140 = S. lanicuspis.
glabra, F. Muell. l. c. i. 139 = Anisacantha glabra.
lanata, Moq. Chenop. Enum. 83 = S. bicornis.

KEPA, Tourn. ex Rafin. Fl. Tellur. ii. 19 (1836) = **Allium**, Tourn. (Liliac.).
esculenta, Rafin. l. c. 20 = A. Cepa.
fistulosa, Rafin. l. c. = A. fistulosum.

KEPPLERIA, Mart. ex Endl. Gen. 251 (1837) = **Bentinckia**, Berry (Palm.).

KEPPLERIA, Meissn. Gen 355 (1842) = **Oncosperma**, Blume (Palm.).
tigillaria, Meissn. l. c.; Comm. 266 = O. filamentosa.

KERAMANTHUS, Hook. f. Bot. Mag. t. 6271 (1876). *PASSIFLORACEAE*.
Kirkii, *Hook. f. l. c.*—Afr. trop.

KERAMOCARPUS, Fenzl, in Russegg. Reise, ii. 966 (1843) = **Coriandrum**, Linn. (Umbellif.).
Tordylium, Fenzl, l. c. = C. tordylioides.

KERANDRENIA, Steud. Nom. ed. II. i. 845 (1840), sphalm. = **Keraudrenia**, J. Gay (Stercul.).

KERANTHUS, Lour. ex Endl. Gen. 193, in syn. (1836) = **Dendrobium**, Sw. (Orchid.).

KERASELMA, Neck. Elem. ii. 353 (1790) = **Euphorbia**, Linn.
corifolia, Rafin. Fl. Tellur. iv. 116 = E. corifolia.
Cyparissias, Rafin. l. c. = E. Cyparissias.
diversifolia, Rafin. l. c. = E. Terracina?
Esula, Rafin. l. c. = E. Esula.
exigua, Rafin. l. c. = E. exigua.

KERASELMA:—
falcata, Rafin. Fl. Tellur. iv. 116 = Euphorbia falcata.
genistoides, Rafin. l. c. = E. genistoides.
lathyroides, Rafin. l. c. = E. lathyroides.
leptophylla, Rafin. l. c. = E. tenuifolia.
lucida, Rafin. l. c. = E. lucida.
oleracea, Rafin. l. c. = E. Peplus.
pallida, Rafin. l. c. = E. salicifolia, lucida.
Peplus, Rafin. l. c. = E. Peplus.
provincialis, Rafin. l. c. = E. Terracina.
retusa, Rafin. l. c. = E. cornuta?
rubra, Rafin. l. c. = E. falcata?
segetalis, Rafin. l. c. = E. segetalis.
Seguieri, Rafin. l. c. = E. Gerardi.
squamosa, Rafin. l. c. = E. aspera.
sylvatica, Rafin. l. c. = E. sylvatica.

KERASKOMION, Rafin. New Fl. Am. iv. 21 (1836) = **Cicuta**, Tourn. (Umbellif.).
bulbiferum, Rafin. l. c. = C. bulbifera.

KERATEPHORUS, Hassk. Retzia, 100 (1855); et in Flora, xxxviii. (1855) 578 = **Payena**, A. DC. (Sapotac.).
Leerii, Hassk. ll. cc. 101; 579.—Sumatra.
Wightii, Hassk. ll. cc. 100; 579 = P. lucida.
Wightii, Maing. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 547 = P. malaccensis.

KERATOPHORUS, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 547 (1882) sphalm. = praec.

KERAUDRENIA, J. Gay, in Mém. Mus. Par. vii. (1821) 461. t. 23. *STERCULIACEAE*, Benth. & Hook. f. i. 227.
adenolasia, F. Muell. Fragm. x. 96 = Seringia adenolasia.
hermanniaefolia, *J. Gay, in Mém. Mus. Par.* vii. (1821) 462. t. 23.—Austral.
Hillii, *F. Muell. ex Benth. Fl. Austral.* i. 246.—Austral.
Hookeri, F. Muell. Fragm. i. 28, 242 = seq.
Hookeriana, *Walp. Ann.* ii. 164.—Austral.
integrifolia, Hook. in Mitch. Trop. Austral. 341 = Hookeriana.
integrifolia, *Steud. in Lehm. Pl. Preiss.* i. 236.—Austral.
lanceolata, *Benth. Fl. Austral.* i. 245.—Austral.
microphylla, Steetz, in Lehm. Pl. Preiss. ii. 347 = hermanniaefolia.
nephrosperma, *Benth. Fl. Austral.* i. 246.—Austral.
velutina, Steetz, in Lehm. Pl. Preiss. ii. 348 = integrifolia.

KERBERA, Fourn. in Mart. Fl. Bras. vi. iv. 290 (1885). *ASCLEPIADEAE*.
Eichleri, *Fourn. l. c.*—Bras.

KERCHOVEA, Joriss. in Belg. Hortic. xxxii. (1882) 201. t. 8 = **Stromanthe**, Sond. (Scitam.).
floribunda, Joriss. l. c.—Bras.

KERIA, Spreng. Anleit. ii. ii. 869 (1818) = **Kerria**, DC. (Rosac.).

KERINGA, Rafin. Sylva Tellur 144 (1838) = **Vernonia**, Schreb. (Compos.).
amygdalina, Rafin. l. c. = V. amygdalina.

KERINOZOMA, Steud. ex Zoll. Syst. Verz. Ind. Archip. 57 (1854); et Syn. Pl. Gram. 358 (1855) = **Ischaemum**?, Linn. (Gramin.).—Cf. Benth. & Hook. f. Gen. iii. 1096.
Cheribon, Steud. Syn. Pl. Gram. 358 = *collina*.
collina, Steud. in Zoll. Syst. Verz. Ind. Archip. 57, nomen.—Java.
littoralis, Steud. l. c. nomen.—Java.
Suraboja, Steud. Syn. Pl. Gram. 358 = *littoralis*.

KERMADECIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 228. *PROTEACEAE*, Benth. & Hook. f. iii. 178.
austro-caledonica, *Benth. & Hook. f. Gen.* iii. 178.—N. Caled.
Bleasdali, Benth. & Hook. f. l. c. = Roupala Bleasdalei.
elliptica, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 228.—N. Caled.
rotundifolia, *Brongn. & Gris, l. c.*—N. Caled.
sinuata, *Brongn. & Gris, l. c.*—N. Caled.

KERMULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = **Fleurya**, Gaudich. (Urtic.).

KERNERA, Medic. in Pflanzeng. 71 (1792) = **Cochlearia**, Linn. (Crucif.).
auriculata, Reichb. in Moessl. Handb. ed. II. ii. 1142 = C. auriculata.
Boissieri, Reut. in Boiss. & Reut. Pugill. Pl. Nov. 9 = C. saxatilis?
decipiens, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 842 = C. decipiens.
glastifolia, Reichb. Fl. Germ. Excurs. 669 = C. glastifolia.
myagrodes, Medic. in Usteri, Neue Ann. Bot. ii. (1794) 42 = C. saxatilis.
polysperma, Willk. & Cost. in Linnaea, xxx. (1859–60) 86 = C. polysperma.
sagittata, Miégev. in Bull. Soc. Bot. Fr. xiv. (1867) 145 = Neslia paniculata.
saxatilis, Reichb. in Moessl. Handb. ed. II. ii. 1142 = C. saxatilis.
simplex, Rafin. Atl. Journ. 178 (Quid?).—Am. bor.

KERNERA, Schrank, Baier. Reise, 50 (1786) = **Tozzia**, Linn. (Scrophul.).
bavarica, Schrank, l. c. = T. alpina.

KERNERA, Willd. Sp. Pl. iv. 947 (1805) = **Posidonia**, Koen. (Naiad.).
antarctica, Schult. f. Syst. vii. 170 = Cymodocea antarctica.
nodosa, Schult. f. l. c. = Cymodocea aequorea.
oceanica, Willd. Sp. Pl. iv. 947 = P. oceanica.
ovalis, Schult. f. Syst. vii. 170 = Halophila ovata.
serrulata, Schult. f. l. c. = Cymodocea serrulata.
spinulosa, Schult. f. l. c. = Halophila spinulosa.

KERNERIA, Moench, Meth. 595 (1794) = **Bidens**, Tourn. (Compos.).
bipinnata, Gren. & Godr. Fl. Fr. ii. 169 = B. bipinnata.
coreopsoides, Cass. in Dict. Sc. Nat. lix. 328 = B. chrysanthemoides.
dubia, Cass. l. c. xxiv. 398 = B. pilosa.
ferulaefolia, Cass. l. c. li. 43 = B. ferulaefolia.
helianthoides, Cass. l. c. xxiv. 399 = B. chrysanthemoides.
leucantha, Cass. l. c. 398 = B. pilosa.
pilosa, Lowe, Man. Fl. Mader. i. 474 = B. pilosa.
rubifolia, Cass. in Dict. Sc. Nat. xxiv. 399 = B. rubifolia.
serrulata, Cass. l. c. li. 474 = B. grandiflora.
tetragona, Moench, Meth. 595 = B. pilosa.

KERRIA, DC. in Trans. Linn. Soc. xii. (1817) 156. *ROSACEAE*, Benth. & Hook. f. i. 613.
japonica, *DC. l. c.* 157.—Japon.
tetrapetala, Siebold, in Verh. Batav. Gen. xii. (1830) 69; Lem. in Illustr. Hortic. xv. (1868) sub t. 556 = japonica.

KETELEERIA, Carr. in Rev. Hortic. (1866) 449. *CONIFERAE* [Benth. & Hook. f. iii. 442].
Fortunei, *Carr. l. c.*—Japon.

KETHOSIA, Rafin. Fl. Tellur. iv. 78 (1836) = **Hewittia**, Wight & Arn. (Convolv.).
involucrata, Rafin. l. c. = H. bicolor.

KETMIA, Tourn. ex Burm. Thes. Zeyl. 133 (1737) = **Hibiscus**, Linn. (Malvac.).
arborea, Moench, Meth. 617 = H. syriacus.
glandulosa, Moench, Meth. Suppl. 202 = H. cannabinus.
mutabilis, Moench, Meth. 617 = H. mutabilis.
syriaca, Scop. Fl. Carn. ed. II. ii. 45 = H. syriacus.
syrorum, Medic. Malv. 45 = H. syriacus.
Trionum, Scop. Fl. Carn. ed. II. ii. 44 = H. Trionum.

KETMIASTRUM, Heist. Syst. 8 (1748) = praec.?

KETTMIA, Medic. in Vorles. Churpf. Phys.-Oek. Ges. iv. I. (1789) 240 = KETMIA, Tourn. = **Hibiscus**, Linn. (Malvac.).

KEULIA, Molina, Sagg. Chile, ed. II. 159 (1810) = **Gomortega**, Ruiz & Pav. (Laurin.).
chilensis, Molina, l. c. = G. nitida.

KEURA, Forsk. Fl. Aegypt. Arab. 95, 122, 172 (1775) = **Pandanus**, Linn.
odora, Thunb. Nov. Gen. 26 = seq.
odorifera, Forsk. Fl. Aegypt. Arab. 172 = P. odoratissimus.

KEURVA, Endl. Gen. 242 (1837) = praec.

KEYSERLINGIA, Bunge, ex Boiss. Fl. Orient. ii. 629 (1872). *LEGUMINOSAE.*
Buxbaumii, *Bunge, ex Boiss. l. c.* 630.—Persia.
Griffithii, *Boiss. l. c.*—Afghan.

KHAYA, A. Juss. in Mém. Mus. Par. xix. (1830) 249; (Meliac. 97. t. 10). *MELIACEAE*, Benth. & Hook. f. i. 338.
anthotheca, *C. DC. in DC. Monog. Phan.* i. 721.—Afr. trop.
senegalensis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 249 (*Meliac.* 98. t. 10).—Afr. trop.

KHAYEA, Planch. & Triana, in Bull. Soc. Bot. Fr. viii. (1861) 99 = **Kayea**, Wall. (Guttif.).

KHYTIGLOSSA, Nees, in DC. Prod. xi. 159, in syn. (1847) err. typ. = RHYTIGLOSSA, Nees = **Dianthera**, Gronov. (Acanthac.).

KIBARA, Endl. Gen. 314 (1837). *MONIMIACEAE*, Benth. & Hook. f. iii. 140.
BRONGNIARTIA, Blume, Bijdr. 435 (1825).
SARCODISCUS, Griff. Notul. iv. 380 (1854).
SCIADICARPUS, Hassk. in Flora, xxv. (1842) II. Beibl. 20.
WILKIEA, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 64.
aruensis, *Becc. Malesia*, i. (1877) 188.—N. Guinea.
Blumei, *Steud. Nom.* ed. II. i. 846.—Malaya.
chartacea, *Blume, Mus. Bot. Lugd. Bat.* ii. 89.—Sumatra.
coriacea, Hook. f. & Thoms. Fl. Ind. i. 166 = Blumei.
cuspidata, *Blume, Mus. Bot. Lugd. Bat.* ii. 89.—Sumatra.
formicarum, *Becc. Malesia*, i. (1877) 188.—N. Guinea.
hospitans, *Becc. l. c.* 189.—N. Guinea.
laxiflora, *Benth. Fl. Austral.* v. 289.—Austral.
longipes, *Benth. l. c.*—Austral.
macrophylla, *Benth. l. c.* 288.—Austral.
obtusa, *Blume, Mus. Bot. Lugd. Bat.* ii. 89.—Ins. Celebes.
olivaeformis, *Becc. Malesia*, i. (1877) 187.—N. Guinea.
pubescens, *Benth. Fl. Austral.* v. 290.—Austral.

KIBATALIA, G. Don, Gen. Syst. iv. 86 (1837) = **Kickxia**, Blume (Apocyn.).
arborea, G. Don. l. c. = K. arborea.

KIBBESSIA, Walp. Rep. ii. 148 (1843), sphalm. = **Kibessia**, DC. (Melastom.).

KIBERA, Adans. Fam. ii. 417 (1763) = **Sisymbrium**, Tourn. (Crucif.).

KIBESSIA, DC. Prod. iii. 196 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 771.
MACROPLACIS, Blume, Mus. Bot. Lugd. Bat. i. 7 (1849).
RECTOMITRA, Blume, l. c. 6 (1849).
acuminata, *Decne. in Ann. Sc. Nat.* Sér. III. v. (1846) 316.—Singapore.
angustifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 9.—Borneo.
azurea, *DC. Prod.* iii. 196.—Java.
cordata, *Korth. Verh. Nat. Gesch. Bot.* 253.—Sumatra.
cupularis, Decne. in Deless. Ic. Sel. iii. 3. t. 5 = simplex.
pubescens, *Decne. in Ann. Sc. Nat.* Sér. III. v. (1846) 318.—Penang.
sessilis, *Blume, Mus. Bot. Lugd. Bat.* i. 8.—Java.
simplex, *Korth. Verh. Nat. Gesch. Bot.* 253.—Malaya.
subalata, *Blume, Mus. Bot. Lugd. Bat.* i. 8.—Sumatra.
subulata, Triana, in Trans. Linn. Soc. xxviii. (1871) 152 sphalm. = praec.
tetraptera, *Miq. Fl. Ind. Bat.* i. I. 571.—Java.
tuberculata, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 553.—Malacca.

KICKXIA, Blume, Fl. Jav. Praef. p. vii (1828) (Kixia); et Rumphia, iv. 25 (1848). *APOCYNACEAE*, Benth. & Hook. f. ii. 721.
HASSELTIA, Blume, Bijdr. 1045 (1826).

KICKXIA:—
KIBATALIA, G. Don, Gen. Syst. iv. 86 (1837).
africana, *Benth. in Hook. Ic. Pl.* t. 1276.—Afr. trop.
arborea, *Steud. Nom.* ed. II. i. 846.—Java.
arborea, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 132. t. 428 = Blancoi.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 318.—Ins. Philipp.

KICKXIA, Dum. Fl. Belg. 35 (1827) = **Linaria**, Tourn. (Scrophul.).
Elatine, Dum. l. c. = L. Elatine.
spuria, Dum. l. c. = L. spuria.

KIELBOUL, Adans. Fam. ii. 31 (1763) = **Aristida**, Linn. (Gramin.).

KIELMEYERA, Mart. Nov. Gen. et Sp. i. 109. t. 68–72 (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 188.
MARTINIERIA, Vell. Fl. Flum. 232 (1825); v. t. 114 (1827).
angustifolia, *Pohl, Pl. Bras. Ic.* ii. 46.—Bras.
argentea, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 165.—Bras.
coriacea, *Mart. Nov. Gen. et Sp.* i. 112.—Bras.
corymbosa, *Mart. l. c.* 113. t. 72.—Bras.
excelsa, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 308.—Bras.
falcata, *Cambess. in St. Hil. l. c.* 304.—Bras.
humifusa, *St. Hil. l. c.* 307.—Bras.
membranacea, *Casar. Dec. Stirp. Nov. Bras.* v. 39.—Bras.
microphylla, *Pohl, Pl. Bras. Ic.* ii. 49.—Bras.
neriifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 306.—Bras.
oblonga, *Pohl, Pl. Bras. Ic.* ii. 47.—Bras.
petiolaris, *Mart. Nov. Gen. et Sp.* i. 111. t. 69.—Bras.
pumila, *Pohl, Pl. Bras. Ic.* ii. 48.—Bras.
rosea, *Mart. Nov. Gen. et Sp.* i. 110.—Bras.
rubriflora, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 305.—Bras.
rugosa, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 165.—Bras.
speciosa, *St. Hil. Pl. Us. Bras.* t. 58.—Bras.
tomentosa, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 308. t. 61.—Bras.
variabilis, *Mart. Nov. Gen. et Sp.* i. 112. t. 71.—Bras.

KIELMIERA, G. Don, Gen. Syst. i. 564 (1831) = praec.

KIERSCHLEGERIA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 176 = **Fuchsia**, Linn. (Onagrar.).
Lindleyi, Spach, l. c. = F. parviflora.
lycioides, Spach, l. c. = F. rosea.

KIESERA, Reinw. ex Blume, Cat. Gew. Buitenz. 93 (1823); et in Syll. Ratisb. ii. (1828) 11 = **Tephrosia**, Pers. (Legumin.).
gracilis, Miq. Fl. Ind. Bat. i. I. 291.—Java.
minor, Miq. l. c.—Java.
sericea, Reinw. ex Blume, Cat. Gew. Buitenz. 93; et in Syll. Ratisb. ii. (1828) 11 = T. candida.
sumatrana, Miq. Fl. Ind. Bat. i. I. 1083.—Sumatra.

KIESERIA, Nees, in Neuwied, Reise nach Bras. ii. 338 (1821) = **Bonnetia**, Mart. (Ternstroem.).
stricta, Nees, l. c. = B. stricta.

KIGELIA, DC. Prod. ix. 247 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1053.
KIGELKEIA, Rafin. Sylva Tellur. 166 (1838), nomen prius.
SOTOR, Fenzl, in Ber. 21. Verssaml. Deutsch. Naturf. 1843 (1844) 168.
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 60.—Abyss.
aethiopica, *Decne. in Deless. Ic. Sel. Pl.* v. 39. t. 93.—Afr. trop.
africana, *Benth. in Hook. Niger Fl.* 463.—Afr. austr.
madagascariensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 274.—Madag.
pinnata, *DC. Prod.* ix. 247.—Afr. trop.

KIGELKEIA, Rafin. Sylva Tellur. 166 (1838) = praec.

KIGELLARIA, Endl. Ench. 479 (1841) = **Kiggelaria**, Linn. (Bixin.).

KIGGELARIA, Linn. Gen. ed. I. 309 (1737).
BIXINEAE, Benth. & Hook. f. i. 130.
africana, *Linn. Sp. Pl.* 1037.—Afr. austr.
Dregeana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 333.—Afr. austr.
ferruginea, *Eckl. & Zeyh. Enum.* 15.—Afr. austr.
glandulosa, Salisb. Prod. 321 = africana.
integrifolia, Jacq. Coll. ii. 296; Ic. Pl. Rar. t. 628 = africana.

KIGGELARIA, Linn. Syst. ed. I (1735) = **Cliffortia** ? Linn. (Rosac.).

KILBERA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = **Sisymbrium**, Tourn. (Crucif.).
supina, Fourr. l. c. = S. supinum.

KILIANA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) I. Intell. 26, nomen = **Pulicaria**, Gaertn. (Compos.).
chrysopsioides, Sch. Bip. l. c. = P. petiolaris.

KILIANIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 336 (1873) = praec.

KILLINGA, Adans. Fam. ii. 498 (1763) = **Athamanta**, Linn. (Umbellif.).

KILLINGA, Lestib. Essai Cyp. 28 (1819) = **Kyllinga**, Rottb. (Cyperac.).

KILLINGIA, Juss. Gen. 2 1789) = praec.

KILLYNGIA, Ham. Prod. Pl. Ind. Occ. 13 (1825) = praec.

KINGIA, R. Br. in King, Narr. Surv. Austral. Coast, ii. 535 (1827). *JUNCACEAE*, Benth. & Hook. f. iii. 865.
argentea, Preiss, in Lehm. Pl. Preiss. ii. 52 = australis.
australis, *R. Br. in King, Narr. Surv. Austral. Coast*, ii. 535.—Austral.

KINGSBOROUGHIA, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 67 = **Meliosma**, Blume (Sabiac.).
alba, Liebm. l. c. = M. alba.

KINGSTONIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 531 (1821) = **Saxifraga**, Linn.
guttata, S. F. Gray, l. c. = S. Hirculus.

KINGSTONIA, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 93 (1872). *ANONACEAE*.
nervosa, *Hook. f. & Thoms. l. c.*—Malaya.

KINIA, Rafin. Specch. i. 87 (1814) (Gen. ex auctore Medeolae et Trillio affine).
biflora, Rafin. l. c. 88.—Borneo.

KINKINA, Adans. Fam. ii. 147 (1763) = **Cinchona**, Linn. (Rubiac.).

KIOSMINA, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
hispanica, Rafin. l. c. = S. hispanica.

KIPPISTIA, Miers, in Trans. Linn. Soc. xxviii. (1872) 414, 416 = **Salacia**, Linn. (Hippocrat.).
cognata, Miers, l. c. 417 = S. cognata.
diffusiflora, Miers, l. c. 418 = S. diffusiflora.
organensis, Miers, l. c. 416 = S. serrata.
serrata, Miers, l. c. 419 = S. serrata.

KIPPISTIA, F. Muell. Rep. Babb. Exped. 12 (1858) = **Minuria**, DC. (Compos.).
suaedifolia, F. Muell. l. c. 13 = M. suaedifolia.

KIRCHNERA, Opiz, in Lotos, viii. (1858) 65 = **Astragalus**, Tourn. (Legumin.).
arenaria, Opiz, l. c. = A. arenarius.

KIRGANELIA, Juss. Gen. 387 (1789) = **Phyllanthus**, Linn. (Euphorb.).
alba, Blanco, Fl. Filip. ed. I. 713 = P. albus.
Boiviniana, Baill. Adansonia, ii. (1861–62) 49.—Ins. Mayotte.
Bojeriana, Baill. l. c. 47 = P. Bojerianus.
decipiens, Baill. l. c. 49 = P. decipiens.
dubia, Baill. Etud. Gén. Euph. 614 = P. reticulatus.
dumetosa, Spreng. Syst. iii. 48 = P. dumetosus.
eglandulosa, Baill. Etud. Gén. Euph. 614 = P. reticulatus.
elegans, Juss. ex Spreng. Syst. iii. 48 = P. Casticum.

KIRGANELIA :—
elliptica, Spreng. Syst. iii. 48 = Securinega elliptica.
floribunda, Baill. Adansonia, i. (1860–61) 83 = Phyllanthus floribundus.
floribunda, Spreng. Syst. iii. 48 = P. salviaefolius.
glaucescens, Baill. Adansonia, ii. (1861–62) 48.—Madag.
intermedia, Baill. Etud. Gén. Euph. 614 = P. reticulatus.
multiflora, Baill. l. c. = P. reticulatus.
nigrescens, Blanco, Fl. Filip. ed. I. 712 = P. nigrescens.
Pervilleana, Baill. Adansonia, ii. (1861–62) 50 = P. Pervilleanus.
phyllanthoides, A. Juss. Tent. Euph. 108 = P. Casticum.
Prieuriana, Baill. Adansonia, i. (1860–61) 82 = P. reticulatus.
puberula, Baill. Etud. Gén. Euph. 614 = P. reticulatus.
pumila, Blanco, Fl. Filip. ed. I. 712 = P. pumilus.
reticulata, Baill. Etud. Gén. Euph. 613 = P. reticulatus.
salviaefolia, Spreng. Syst. iii. 48 = P. salviaefolius.
sinensis, Baill. Etud. Gén. Euph. 614, nomen = P. reticulatus.
timorensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 483 = P. Casticum.
triandra, Blanco, Fl. Filip. ed. I. 711 = P. triandrus.
trilocularis, Baill. Etud. Gén. Euph. 614 = P. decipiens.
Vieillardi, Baill. Adansonia, ii. (1861–62) 231.—N. Caled.
villosa, Blanco, Fl. Filip. ed. I. 712 = P. Llanosi.
virginea, J. F. Gmel. Syst. 1008 = P. Casticum.
Wightiana, Baill. Etud. Gén. Euph. 613, 614 = P. reticulatus.
zanzibariensis, Baill. Adansonia, ii. (1861–62) 48.—Afr. trop.

KIRGANELLA, J. F. Gmel. Syst. 1008 (1791) = praec.

KIRGANELLIA, Dum. Anal. Fam. 45 (1829) = praec.

KIRILOVIA, Lindl. Veg. Kingd. 513 (1847) = seq.

KIRILOWIA, Bunge, Del. Sem. Hort. Dorpat. (1843) 7. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 57.
eriantha, *Bunge, l. c.* 8.—Sibir.
pilosa, Bunge, l. c. = Panderia pilosa.
villosa, *Benth. & Hook. f. Gen.* iii. 57.—Afghan.

KIRKIA, Oliver, Fl. Trop. Afr. i. 310 (1868). *SIMARUBEAE*.
acuminata, *Oliver, l. c.* 311; *et in Hook. Ic. Pl.* t. 1036.—Afr. trop.

KIRSCHLEGERA, Reichb. Nom. 170 (1841) = seq.

KIRSCHLEGERIA, Reichb. Handb. 246 (1837) = KIERSCHLEGERIA, Spach = **Fuchsia**, Linn. (Onagrar.).

KISSENIA, R. Br. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 43. *LOASEAE*, Benth. & Hook. f. i. 803.
CNIDONE, E. Mey. ex Endl. Gen. Suppl. ii. 76, in syn. (1842).
FISSENIA, R. Br. ex Endl. l. c. (1842).
spathulata, *R. Br. ex T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 43.—Arab.; Afr. trop.

KISSODENDRON, Seem. Journ. Bot. iii. (1865) 201 = **Hedera**, Tourn. (Araliac.).
australianum, Seem. l. c. = H. australiana.

KITAIBELA, Batsch, Tab. 19 (1802) = seq.

KITAIBELIA, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 107. *MALVACEAE*, Benth. & Hook. f. i. 200.
Balansae, *Boiss. Fl. Orient.* i. 817.—Cilicia.
vitifolia, *Willd. in Ges. Naturf. Fr. Neue Schr.* ii. (1799) 108. f. 4.—Slavon.

KITCHINGIA, Baker, in Journ. Linn. Soc. xviii. (1881) 268. *CRASSULACEAE*.
amplexicaulis, *Baker, l. c.* xx. (1883) 142.—Madag.
campanulata, *Baker, l. c.* xviii. (1881) 269.—Madag.
gracilipes, *Baker, l. c.* 268.—Madag.
miniata, *Baker, in Journ. Bot.* xx. (1882) 109.—Madag.

KITCHINGIA :—
panduriformis, *Baker, in Journ. Linn. Soc.* xx. (1883) 141.—Madag.
parviflora, *Baker, l. c.*—Madag.
peltata, *Baker, l. c.* 140.—Madag.
porphyrocalyx, *Baker, l. c.* 142.—Madag.
schizophylla, *Baker, l. c.* xxi. (1884) 340.—Madag.

KITTELIA, Reichb. Handb. 186 (1837) = **Cyanea**, Gaudich. (Campanulac.).

KITTELOCHARIS, Alef. in Bot. Zeit. xxi. (1863) 282 = **Reinwardtia**, Dum. (Lineae).
trigyna, Alef. l. c. = R. trigyna.

KIXIA, Blume, Fl. Jav. Praef. p. vii (1830) = **Kickxia**, Blume (Apocynac.).

KIXIA, Meissn. Comm. 215, in syn. (1843?) = KICKXIA, Dum. = **Linaria**, Tourn. (Scrophul.).

KLADNIA, Schur, Enum. Pl. Transs. 53 (1866) = **Hesperis**, Linn. (Crucif.).
tristis, Schur, l. c. = H. tristis.

KLANDERIA, F. Muell. in Linnaea, xxv. (1852) 426 = **Prostanthera**, Labill. (Labiat.).
chlorantha, F. Muell. l. c. = P. chlorantha.

KLAPROTHIA, H. B. & K. Nov. Gen. et Sp. vi. 121. t. 537 (1823). *LOASEAE*, Benth. & Hook. f. i. 804.
mentzelioides, *H. B. & K. l. c.*—Ecuador.

KLASEA, Cass. in Dict. Sc. Nat. xxxv. 173 (1825); xli. 321 = **Serratula**, Dill. (Compos.).
centauroides, Cass. l. c. xxv. 173 = S. centauroides.
cichoracea, Webb, Iter Hisp. 34 = S. flavescens.
heterophylla, Cass. in Dict. Sc. Nat. xxxv. 173; xli. 323 = S. heterophylla.
nudicaulis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 98 = S. nudicaulis.
pinnatifida, Cass. in Dict. Sc. Nat. xxxv. 173; xli. 321 = S. radiata.
pinnatifida, Webb, Iter Hisp. 34 = S. pinnatifida.
quinquefolia, Cass. in Dict. Sc. Nat. xxxv. 173; xli. 321 = S. quinquefolia.

KLATTIA, Baker, in Journ. Linn. Soc. xvi. (1877) 109. *IRIDEAE*, Benth. & Hook. f. iii. 702.
partita, *Baker, l. c.* 110.—Afr. austr.

KLAUSEA, Endl. Gen. 480, in syn. (1838), sphalm. = KLASEA, Cass. = **Serratula**, Dill. (Compos.).

KLEINHOFIA, Gisek. Prael. 452 (1792) = seq.

KLEINHOVEA, Roxb. Fl. Ind. iii. 140 (1832) = seq.

KLEINHOVIA, Linn. Sp. Pl. ed. II. 1365 (1763). *STERCULIACEAE*, Benth. & Hook. f. i. 219.
Hospita, *Linn. l. c.*—As. trop.

KLEINIA, Crantz, Inst. ii. 488 (1766) = **Quisqualis**, Linn. (Combret.).
quadricolor, Crantz, l. c. 489 = Q. indica.

KLEINIA, Jacq. Stirp. Select. Am. 215. t. 127 (1763) = **Porophyllum**, Vaill. (Compos.).
brevifolia, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 52 = P. linifolium.
cacalioides, Less. in Linnaea, vi. (1831) 132 = P. cacalioides.
colorata, H. B. & K. Nov. Gen. et Sp. iv. 158 = P. coloratum.
decumbens, Heynh. Nom. i. 437 = P. decumbens.
exserta, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 52 = P. exsertum.
filifolia, Spreng. Syst. iii. 438 = P. linifolium.
glandulosa, [Moç. & Sesse] ex DC. Prod. v. 648 = P. macrocephalum.
jorullensis, H. B. & K. Nov. Gen. et Sp. iv. 156. t. 356 = P. jorullense.
linearis, Hort. Par. ex DC. Prod. v. 650 = P. decumbens.
linifolia, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 52 = P. lineare.
obscura, Spreng. Syst. iii. 438 = P. lineare.
oppositifolia, Spreng. l. c. = P. oppositifolium.
Porophyllum, Willd. Sp. Pl. iii. 1738 = P. ruderale.

KLEINIA :—
pumila, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 52 = Porophyllum lineare.
ruderalis, Jacq. Enum. Pl. Carib. 28 = P. ruderale.
Selloi, Spreng. Syst. iii. 438 = Stevia oxylaena.
suffruticosa, Lodd. Bot. Cab. t. 1561 = P. decumbens.
suffruticosa, Willd. Sp. Pl. iii. 1739 = P. linifolium.
tagetoides, H. B. & K. Nov. Gen. et Sp. iv. 158 = P. tagetoides.
viridiflora, H. B. & K. l. c. 157 = P. viridiflorum.

KLEINIA, Juss. in Ann. Mus. Par. ii. (1803) 424. t. 61. f. 1 = **Jaumea**, Pers. (Compos.).
carnosa, Vis. Hort. Patav. (1842) 143 = J. carnosa.
linearifolia, Juss. in Ann. Mus. Par. ii. (1803) 424 = J. linearis.

KLEINIA, Linn. Syst. ed. I (1735) = **Senecio**, Tourn. (Compos.).
acaulis, DC. Prod. vi. 339 = S. acaulis, *Sch. Bip.*
aizoidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = seq.
aizoides, DC. Prod. vi. 337 = S. aizoides.
alata, G. F. W. Mey. Prim. Fl. Esseq. 249 = Mikania scandens.
angulata, Willd. Sp. Pl. iii. 1739 = Emilia angulata.
Anteuphorbia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = seq.
Anteuphorbium, DC. Prod. vi. 338 = S. Anteuphorbium.
articulata, Haw. Syn. Pl. Succ. 315 = S. articulatus.
breviscapa, DC. Prod. vi. 337 = S. cicatricosus.
cana, DC. l. c. 338 = S. quinquangulatus.
crassulaefolia, DC. l. c. 336 = S. crassulaefolius.
cuneifolia, DC. l. c. 337 = S. cuneifolius.
Cusimbua, Less. in Linnaea, vi. (1831) 133 = Gynura angulosa.
Ecklonis, Harv. in Harv. & Sond. Fl. Cap. iii. 318.—Afr. austr.
ficoides, Haw. Syn. Pl. Succ. 313 = S. ficoides.
fulgens, Hook. f. Bot. Mag. t. 5590.—Afr. austr.
gonoclada, DC. Prod. vi. 336 = S. gonocladus.
Haworthii, DC. l. c. 338 = S. Haworthii, *Sch. Bip.*
hieracioides, Less. in Linnaea, vi. (1831) 133 = Gynura auriculata.
japonica, Less. l. c. 134 = Gynura pinnatifida.
longiflora, DC. Prod. vi. 337 = S. longiflorus.
neriifolia, Haw. Syn. Pl. Succ. 312 = S. Kleinia.
odora, DC. Prod. vi. 339 = S. odorus.
papillaris, Haw. Syn. Pl. Succ. 313 = S. papillaris.
pendula, DC. Prod. vi. 339 = S. pendulus.
pinguifolia, DC. l. c. 336 = S. pinguifolius.
pteroneura, DC. l. c. 338 = S. pteroneurus.
pugioniformis, DC. l. c. = S. pugioniformis.
radicans, Haw. in Till. Phil. Mag. (1823); DC. Prod. vi. 337 = S. radicans.
repens, Haw. Syn. Pl. Succ. 313 = S. succulentus, *Sch. Bip.*
rigida, DC. Prod. vi. 338 = S. spinescens.
sempervivа, DC. l. c. 339 = S. sempervivus.
subradiata, DC. l. c. = S. subradiatus.
talinoides, DC. l. c. 337 = S. talinoides.
teres, Wall. Cat. n. 3164 = Emilia prenanthoidea.
tomentosd, Haw. Syn. Pl. Succ. 314 = S. Haworthii, *Sch. Bip.*

KLEISTROCALYX, Steud. in Flora, xxxiii. (1850) 229, nomen = **Rynchospora**, Vahl (Cyperac.).
junciformis, Steud. l. c. = R. oligantha.

KLENZEA, Sch. Bip. ex Steud. Nom. ed. II. i. 847 (1840); et ex Hochst. in Flora, xxiv. (1841) 1. Intell. 26 = **Athrixia**, Ker-Gawl. (Compos.) partim.
abyssinica, Sch. Bip. ex Steud. l. c.; Hochst. l. c. = Antithrixia abyssinica.
lycopodioides, Sch. Bip. ex Walp. Rep. ii. 973 = Bryomorpha Zeyheri.
rosmarinifolia, Sch. Bip. ex Steud. Nom. ed. II. i. 847; Hochst. in Flora, xxiv. (1841) 1. Intell. 26 = A. rosmarinifolia.

KLONION, Rafin. New Fl. Am. iv. 35 (1836) = **Eryngium**, Tourn. (Umbellif.).
gracile, Rafin. l. c. 36 = E. Baldwini.
tenuifolium, Rafin. l. c. = E. Baldwini.

KLOPSTOCKIA, Karst. in Linnaea, xxviii. (1856) 251 = **Ceroxylon**, Humb. (Palm.).
cerifera, Karst. l. c. 251 = C. Klopstockiae.
coarctata, Engel, l. c. xxxiii. (1864–65) 676 = C. coarctatum.

KLOPSTOCKIA :—
interrupta, Karst. in Linnaea, xxviii. (1856) 252 = Ceroxylon interruptum.
parvifrons, Engel, l. c. xxxiii. (1864–65) 674 = C. parvifrons.
quindiuensis, Karst. Fl. Colomb. i. 1. t. 1 = C. quindiuense.
utilis, Karst. in Linnaea, xxviii. (1856) 252 = C. utile.
Vogeliana, Engel, l. c. xxxiii. (1864–65) 673 = C. Vogelianum.

KLOTSCHIA, Endl. Gen. 767 (1839) = seq.

KLOTZSCHIA, Cham. in Linnaea, viii. (1833) 327. *UMBELLIFERAE*, Benth. & Hook. f. i. 880.
brasiliensis, *Cham. l. c.*—Bras.

KLOTZSCHIPHYTUM, Baill. Etud. Gén. Euph. 382 (1858) = **Croton**, Linn. (Euphorb.).
mauritianum, Baill. l. c. 383 = C. mauritianus.

KLUGIA, Schlecht. in Linnaea, viii. (1833) 248. *GESNERACEAE*, Benth. & Hook. f. ii. 1019.
GLOSSANTHUS, Klein, ex Benth. Scroph. Ind. 57 (1835).
ampliata, *C. B. Clarke, in DC. Monog. Phan.* v. 160.—Reg. Himal.
azurea, *Schlecht. in Linnaea*, viii. (1833) 248.—Mexic.
glabra, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 489 = Notoniana.
Notoniana, *A. DC. in DC. Prod.* ix. 276.—Ind. or.
Notoniana, Hook. Bot. Mag. t. 4620 = zeylanica.
scabra, Dalz. in Dalz. & Gibs. Bomb. Fl. 134 = Notoniana.
zeylanica, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 490.—Zeylan.

KLUKIA, Andrz. ex DC. Syst. ii. 459 (1821) = **Malcolmia**, R. Br. (Crucif.).
rigida, Andrz. ex DC. l. c. 480 = M. torulosa.

KNAFIA, Opiz, Seznam, 56 (1852) = **Salix**, Linn.
Helix, Opiz, l. c. = S. purpurea.
purpurea, Opiz, l. c. = S. purpurea.

KNANTIA, Hill, Veg. Syst. v. 18 (1772) sphalm. = KNAUTIA, Linn. = **Scabiosa**, Tourn. (Dipsac.).

KNAPPIA, Ferd. Bauer, ex Steud. Nom. ed. II. i. 846 (1840); ii. 75 = **Rhinchoglossum**, Blume (Gesnerac.).

KNAPPIA, Sm. Engl. Bot. t. 1127 (1803); Fl. Brit. iii. 1387 (1804) = **Mibora**, Adans. (Gramin.).
agrostidea, Sm. ll. cc. = M. verna.
verna, Trin. Ic. Gram. ii. t. 17 = M. verna.

KNAUTIA, Linn. Syst. ed. I (1735) = **Scabiosa**, Tourn. (Dipsac.).
alpigena, Schur, Enum. Pl. Transs. 297 = S. arvensis.
ambigua, Boiss. & Orph. in Boiss. Diagn. Ser. II. vi. 95 = S. ambigua.
angustifolia, Reut. ex Willk. & Lange, Prod. Fl. Hisp. ii. 16 = S. legionensis.
arvensis, Coult. Dipsac. 29 = S. arvensis.
bidens, Boiss. Diagn. Ser. I. ii. 108 = S. bidens.
Brandзai, Porc. in Magyar Növényt. Lapok, ix. (1885) 128 = S. sylvatica.
calycina, Guss. Fl. Sic. Syn. i. 170.—Sicil.
carpathica, Heuff. in Flora, xxxix. (1856) 50 = S. arvensis.
carpophylax, Jord. Cat. Gr. Jard. Gren. (1853) 12; F. Schultz, Arch. Fl. Fr. et Allem. 319 = S. sylvatica.
centauroides, Pomel, Nouv. Mat. Fl. Atl. 65.—Alger.
ciliata, Coult. Dipsac. 30 = S. ciliata.
ciliata, Schur, Enum. Pl. Transs. 296 = S. arvensis.
collina, Gren. & Godr. Fl. Fr. ii. 75 = S. collina.
communis, Godr. Fl. Lorr. ed. I. i. 322 = S. arvensis.
craciunelensis, Porc. in Magyar Növényt. Lapok, ix. (1885) 128 = S. longifolia.
cuspidata, Jord. Cat. Gr. Jard. Gren. (1853); F. Schultz, Arch. Fl. Fr. et Allem. 320 = S. sylvatica.
diodon, Guss. Fl. Sic. Syn. i. 171.—Sicil.
dipsacifolia, Heuff. in Flora, xxxix. (1856) 54 = S. sylvatica.
diversifolia, DC. Prod. iv. 652 = S. banatica.
Drymeja, Heuff. in Flora, xxxix. (1856) 53 = S. sylvatica?
dumetorum, Heuff. l. c. 51 = S. arvensis.
exaltata, Schur, Enum. Pl. Transs. 296 = S. arvensis.

KNAUTIA :—
Godetii, Reut. Ind. Sem. Hort. Genev. (1857) 4.—Cf. Linnaea, xxix. (1857–58) 723 = S. longifolia.
heterotricha, C. Koch, in Linnaea, xxiv. (1851) 444 = S. montana.
hybrida, Coult. Dipsac. 30 = S. integrifolia.
indivisa, Bor. Fl. Cent. Fr. ed. III. ii. 316 = S. arvensis.
integrifolia, Bertol. Fl. Ital. ii. 32 = S. integrifolia.
integrifolia, C. Koch, in Linnaea, xxiv. (1851) 443.—As. Min.
integrifolia, Schur, Enum. Pl. Transs. 295 = S. arvensis.
Kossuthii, Pantoč. in Magyar Növényt. Lapok, vi. (1882) 162 = S. pubescens.
lanceolata, Pomel, Nouv. Mat. Fl. Atl. 65.—Alger.
legionensis, DC. Prod. iv. 652 = S. legionensis.
longifolia, Koch, Syn. Fl. Germ. ed. I. 343 = S. longifolia.
lyrophylla, Panč. ex Boiss. Fl. Orient. iii. 128 = *macedonica*?
macedonica, Griseb. Spicil. Fl. Rumel. ii. 178.—Macedon.
magnifica, Boiss. & Orph. ex Orph. in Bull. Congr. Bot. Pétersb. 1869 (1870) 138.—Macedon.
mauritanica, Pomel, Nouv. Mat. Fl. Atl. 64.—Alger.
mollis, Jord. Cat. Jard. Dijon, (1848) 25 = S. gramuntia.
montana, DC. Prod. iv. 651 = S. montana.
neglecta, Meurer, in Oestr. Bot. Zeitschr. xxvii. (1877) 366 = S. arvensis?
nympharum, Boiss. & Heldr. Diagn. Ser. II. ii. 124 = S. ciliata.
occidentalis, Hill, Veg. Syst. v. 18 (Quid?).—'Am. bor.'
ochroleuca, Boiss. Fl. Orient. iii. 129 = S. ambigua.
orientalis, Linn. Sp. Pl. 101 = S. orientalis.
orientalis, Urv. in Mém. Soc. Linn. Par. i. (1822) 270 = S. integrifolia.
ovatifolia, G. Don, Gen. Syst. iii. 688 = S. sylvatica.
palaestina, Linn. Mant. ii. 197 = S. brachiata.
pannonica, Heuff. in Flora, xxxix. (1856) 52 = S. sylvatica.
plumosa, Linn. Mant. ii. 197 = S. plumosa.
propontica, Linn. Sp. Pl. ed. II. 1666 = S. propontica.
puberula, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 320 = S. sylvatica.
sagittata, Blanco, Fl. Filip. ed. I. 54 = Mikania scandens.
Salcedi, G. Don, Gen. Syst. iii. 687.—Hispan.
silvatica, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 360 = S. sylvatica.
spathulata, Lag. Gen. et Sp. Nov. 9 = S. spathulata, *D. Dietr.*
subacaulis, Schur, Enum. Pl. Transs. 295 = S. arvensis.
subcanescens, Cat. Jard. Jord. Gren. (1853) 12; F. Schultz, Arch. Fl. Fr. et Allem. 319 = S. sylvatica.
subscaposa, Boiss. & Reut. Pugill. Pl. Nov. (1852) 53.—Hispan.
sylvatica, Duby, Bot. Gall. i. 257 = S. sylvatica.
Timeroyi, Jord. Cat. Jard. Dijon, (1848) 25 = S. arvensis.
tomentosa, Payot, Fl. Mont-Blanc.—Cf. Bull. Soc. Bot. Fr. xxix. (1882) Bibl. 43 = S. tomentosa?
transalpina, Dalla Torre, Atl. Alpenfl. 122.—Europ.
transsilvanica, Schur, Enum. Pl. Transs. 296 = S. sylvatica.
trichotoma, Moench, Meth. 487 = S. orientalis.
Urvillei, Coult. Dipsac. 29. t. 1. f. 10 = S. integrifolia.
variabilis, F. Schultz, in Arch. Fl. Fr. et Allem. 67 = S. arvensis.
virgata, Jord. Cat. Jard. Dijon (1848) 26.—Pedem.

KNAVEL, Adans. Fam. ii. 506 (1763) = **Scleranthus**, Linn. (Illecebrac.).
annuum, Scop. Fl. Carn. ed. II. i. 298 = S. annuus.

KNEIFFIA, Spach, Hist. Vég. Phan. iv. (1835) 73 = **Oenothera**, Linn. (Onagrar.).
angustifolia, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 367 = Oe. fruticosa.
chrysantha, Spach, l. c. 368 = Oe. pterosperma.
floribunda, Spach, Hist. Vég. Phan. 376 = Oe. fruticosa.
Fraseri, Spach, l. c. 375 = Oe. glauca.
glauca, Spach, l. c. 374 = Oe. glauca.
linearifolia, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 167 = Oe. linifolia.
linearis, Spach, Hist. Vég. Phan. iv. 376 = Oe. fruticosa.

KORTHALSIA, Blume, Rumphia, ii. 166. t. 130. f. 2 (1843). *PALMAE*, Benth. & Hook. f. iii. 932.
CALAMOSAGUS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 22.
andamanensis, *Becc. Malesia*, ii. (1884) 76.—Ins. Andaman.
angustifolia, *Blume, Rumphia*, ii. 172.—Borneo.
celebica, H. Wendl. in Kerch. Palm. 248 = Licuala celebica.
Cheb, *Becc. Malesia*, ii. (1884) 67.—Borneo.
debilis, *Blume, Rumphia*, ii. 169.—Sumatra.
echinometra, *Becc. Malesia*, ii. (1884) 66.—Borneo.
ferox, *Becc. l. c.* 73.—Borneo.
Flabellum, Miq. ex H. Wendl. in Kerch. Palm. 248 = Licuala Flabellum.
flagellaris, Miq. Fl. Ind. Bat. Suppl. 591 = angustifolia.
hispida, *Becc. Malesia*, ii. (1884) 71.—Sumatra.
horrida, *Becc. l. c.* 66.—Borneo.
Junghuhnii, *Miq. Pl. Jungh.* 162.—Java.
laciniosa, *Mart. Hist. Nat. Palm.* iii. 211.—Burma.
Lobbiana, H. Wendl. in Bot. Zeit. xvii. (1859) 174 = rigida.
penduliflora, Miq. ex H. Wendl. in Kerch. Palm. 248 = Licuala penduliflora.
polystachya, *Mart. Hist. Nat. Palm.* iii. 210. t. 172. —Malacca.
rigida, *Blume, Rumphia*, ii. 167.—Malaya.
robusta, *Blume, l. c.* 170.—Malaya.
rostrata, *Blume, l. c.* 168.—Borneo.
rubiginosa, *Becc. Malesia*, ii. (1884) 72.—Borneo.
scaphigera, Kurz, For. Fl. Brit. Burma, ii. 513 = andamanensis.
scaphigera, *Mart. Hist. Nat. Palm.* iii. 211.—Malacca.
Teysmanni, *Miq. Fl. Ind. Bat. Suppl.* 591.—Sumatra.
wallichaefolia, *H. Wendl. in Kerch. Palm.* 248.—Ind. or.
Zippelii, *Blume, Rumphia*, ii. 171.—N. Guinea.

KORYCARPUS, Zea, in Act. Matr. (1806); ex Lag. Gen. et Sp. Nov. 4. (1816) = **Diarrhena**, Rafin. (Gramin.).
arundinaceus, Zea, l. c. ex Lag. l. c. = D. americana.

KOSARIA, Forsk. Fl. Aegypt. Arab. 164 (1775) = **Dorstenia**, Linn. (Urticac.).
Barnimiana, Schweinf. in Sitzb. Naturf. Ges. Berl. (Juli 21, 1863) = D. Barnimiana.
obovata, Schweinf. Beitr. Fl. Aethiop. 190 = D. obovata.
palmata, Schweinf. Heart of Africa, i. 220.—Afr. trop.
tropaeolifolia, Schweinf. in Verh. Zool. Bot. Ges. Wien, xviii. (1868) 687 = D. tropaeolifolia.

KOSTELETSKYA, Brongn. Enum. Gen. 77 (1843), sphalm. = seq.

KOSTELETZKYA, Presl, Rel. Haenk. ii. 130 t. 70. (1831). *MALVACEAE*, Benth. & Hook. f. i. 206.
PENTAGONOCARPUS, Michx. ex Parl. Fl. Ital. v. 105 (1872).
acuminata, *Britten, Journ. Bot.* xviii. (1880) 91.—Mexic.
adoënsis, *Mast. in Oliver, Fl. Trop. Afr.* i. 192.—Afr. trop.
altheaefolia, *A. Gray, Pl. Wright.* i. 23.—Am. bor. occ.
asterocarpa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 191 = sagittata.
batacensis, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 24.—Ins. Philipp.
cordata, *Presl, Rel. Haenk.* ii. 132.—Mexic.
Coulteri, *A. Gray, Pl. Wright.* i. 23.—Mexic.
digitata, *A. Gray, in Proc. Am. Acad.* viii. (1870) 289.—Am. bor.
Grantii, *Garcke.—Cf. Linnaea*, xxxviii. (1874) 697; xliii. (1880–82) 53.—Afr. trop.
hastata, *Garcke, in Eichl. Fahrb.* i. (1881) 222, in syn. —Ind. or.?
hastata, *Presl, Rel. Haenk.* ii. 130.—Mexic.
hispida, *Baker, in Journ. Linn. Soc.* xx. (1883) 98.—Madag.
hispida, *Presl, Rel. Haenk.* ii. 132.—Mexic.
hispidula, *Garcke, in Eichl. Fahrb.* i. (1881) 223, in syn.—Hab.?

KOSTELETZKYA:—
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 46.—Madag.
malacosperma, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 192.—Afr. austr.
mexicana, *Garcke, in Eichl. Fahrb.* i. (1881) 223, in syn.—Mexic.
palmata, *Presl, Bot. Bemerk.* 19.—Mexic.
paniculata, *Benth. Pl. Hartw.* 285.—Mexic.
pentacarpa, *Ledeb. Fl. Ross.* i. 437.—Rossia.
pentasperma, *Griseb. Fl. Brit. W. Ind.* 83.—Am. trop.
sagittata, *Presl, Rel. Haenk.* ii. 131. t. 70.—Mexic.
smilacifolia, *A. Gray, Pl. Wright.* i. 23.—N. Mexic.
Thouarsiana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 541.—Madag.
velutina, *Garcke, in Bremen, Abh.* vii. (1881) 198. —Madag.
virginica, *Presl, ex A. Gray. Gen. Illustr.* ii. 80. t. 132. —Am. bor.

KOTCHUBAEA, Regel, ex Benth. & Hook. f. Gen. i. 98 (1862) = **Kutchubaea**, Fisch. (Rubiac.).
laxiflora, Benth. & Hook. f. l. c. = Kutchubaea insignis.

KOTSCHYA, Endl. Stirp. Nov. Dec. 4 (1839) = **Smithia**, Ait. (Legumin.).
africana, Endl. l. c. 6 = S. Kotschyi.

KOTSJILETTI, Adans. Fam. ii. 60 (1763) = **Xyris**, Linn.

KOWALEWSKIA, Turcz. in Bull. Soc. Nat. Mosc. (1859) I. 263 = **Clethra**, Linn. (Ericac.).
integerrima, Turcz. l. c. = C. mexicana.
serrulata, Turcz. l. c. = C. mexicana.

KOZOLA, Rafin. Fl. Tellur. ii. 25 (1836) = **Scilla**, Linn. (Liliac.).
japonica, Rafin. l. c. = S. japonica.

KRALIKIA, Coss. & Dur. in Bull. Soc. Bot. Fr. xiv. (1867) 89. *GRAMINEAE*, Benth. & Hook. f. iii. 1205.
africana, *Coss. & Dur. l. c.* 90.—Afr. bor.

KRALIKIA, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 151 (1867) = **Chiliocephalum**, Benth. (Compos.).
myriocephala, Sch. Bip. l. c., nomen = C. Schimperi.
Schultzii, Buching. ex Sch. Bip. l. c., nomen = C. Schimperi.

KRAMERIA, Loefl. It. Hisp. 195 (1758). *POLYGALEAE*, Benth. & Hook. f. i. 140.
IXINA et STEMEIENA, Rafin. Atl. Journ. 144 (1832).
argentea, *Mart. ex Spreng. Syst.* ii. 844.—Bras.
arida, *O. Berg, in Bot. Zeit.* xiv. (1856) 762.—Venezuela.
Beyrichii, Sporled. ex O. Berg, l. c. 752 = lanceolata, *Torr.*
canescens, *A. Gray, Pl. Wright.* i. 42.—Am. bor. occ.
canescens, Willd. ex Schult. Mant. iii. 303 = triandra.
cinerea, *Schau. in Linnaea*, xx. (1847) 725.—Mexic.
cistoides, *Hook. & Arn. Bot. Beech. Voy.* 8. t. 5.—Chili.
cuspidata, *Presl, Rel. Haenk.* ii. 103.—Mexic.
cytisoides, *Cav. Ic.* iv. 61. t. 390.—Mexic.
didymandra, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 361.—Bras.
erecta, Willd. ex Schult. Mant. iii. 303 = lanceolata, *Torr.*?
evolvuloides, Triana & Planch. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 72 = spartioides.
glabra, Spreng. Neue Entdeck. ii. 157 = Zollernia falcata.
grandiflora, *A. St. Hil. Fl. Bras. Mer.* ii. 73.—Bras.
grandifolia, *O. Berg, in Bot. Zeit.* xiv. (1856) 764.—Hab.?
Illuca, *Phil. Fl. Atac.* 9.—Chili.
Ixina, Benth. Pl. Hartw. 13 = lanceolata, *Torr.*?
Ixina, H. B. & K. Nov. Gen. et Sp. vii. 244 = tomentosa.
Ixina, *Linn. Sp. Pl.* ed. II. 177.—Am. austr.
Ixine, Linn. Syst. ed. X. 899 = praec.
lanceolata, *O. Berg, in Bot. Zeit.* xiv. (1856) 765.—Venezuela.
lanceolata, *Torr. in Ann. Lyc. N. York*, ii. (1828) 166. —Am. bor. occ.

KRAMERIA:—
latifolia, *Moric. Pl. Nouv. Am.* 63. t. 43.—Bras.
linearis, Poir. Encyc. Suppl. iii. 226 = pentapetala.
linifolia, Willd. ex Schult. Mant. iii. 303 = K. Ixina.
longipes, *O. Berg, in Bot. Zeit.* xiv. (1856) 765.—Bras.
ovata, *O. Berg, l. c.* 764 = tomentosa.
parviflora, *Benth. Bot. Voy. Sulph.* 6. t. 2.—Am. bor. occ.
parvifolia, *Benth. l. c.* 6. t. 1.—Mexic.
pauciflora, [Moç. & Sesse,] ex DC. Prod. i. 341 = lanceolata, *Torr.*
pentapetala, *Ruiz & Pav. Fl. Per.* i. 62. t. 794. f. a.—Peruv.
ramosissima, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 326.—Mexic.
revoluta, *O. Berg, in Bot. Zeit.* xiv. (1856) 751.—Mexic.
ruscifolia, A. St. Hil. Fl. Bras. Mer. ii. 74 = latifolia.
secundiflora, [Moç. & Sesse,] ex DC. Prod. i. 341 = lanceolata, *Torr.*?
spartioides, *Klotzsch, ex O. Berg, in Bot. Zeit.* xiv. (1856) 761.—N. Granat.
tomentosa, *A. St. Hil. Fl. Bras. Mer.* ii. 74.—Bras.
triandra, *Ruiz & Pav. Fl. Per.* i. 61.—Peruv.

KRANIKOFA, Rafin. Princ. Somiol. 30 (1814) = KRASCHENINNIKOVIA, Gueldenst. = **Eurotia**, Adans. (Chenop.).

KRAPFIA, DC. Syst. i. 228 (1818) = **Ranunculus**, Linn.
ranunculacea, Steud. Nom. ed. II. i. 850 = seq.
ranunculina, DC. Syst. i. 228 = R. Krapfia.

KRASCHENINIKOVIA, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 9 = **Stellaria**, Linn. (Caryophyll.).
Davidi, Franch. Pl. David. i. (1884) 51. t. 10.—China.
heterantha, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 376.—Japon.
heterophylla, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 187. —Japon.
Maximowicziana, Franch. & Sav. Enum. Pl. Jap. ii. 297.—Japon.
rupestris, Turcz. in Bull. Soc. Nat. Mosc. (1838) 89, nomen = S. bulbosa.
sylvatica, Maxim. Prim. Fl. Amur. 57.—Reg. Amur.

KRASCHENINNIKOVIA, Gueldenst. in Nov. Comm. Acad. Petrop. xvi. (1772) 548. t. 17 = **Eurotia**, Adans. (Chenopod.).
ceratoides, Gueldenst. l. c. 548, 555 = E. ceratoides.
latens, J. F. Gmel. Syst. 274 = E. ceratioides.
lutens, Steud. Nom. ed. II. i. 850 = praec.

KRASCHNIKOWIA, Turcz. ex Ledeb. Fl. Ross. iii. 402, in syn. (1849) = **Marrubium**, Tourn. (Labiat.).
geraniifolia, Turcz. ex Ledeb. l. c. = M. incisum.

KRATZMANNIA, Opiz, Seznam, 56 (1852) = **Agropyron**, J. Gaertn. (Gramin.).
imbricata, Opiz, l. c. = A. cristatum.

KRAUHNIA, Steud. Nom. ed. II. i. 850 (1840) = seq.

KRAUNHIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Wistaria**, Nutt. (1818) (Legumin.).
frutescens. Rafin. l. c. = W. frutescens.

KRAUSSIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 21. *RUBIACEAE*, Benth. & Hook. f. ii. 95.
CARPOTHALIS, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1843).
congesta, *Oliver, in Journ. Linn. Soc.* xv. (1876) 95. —Afr. trop.
coriacea, Sond. in Linnaea, xxiii. (1850) 53 = Tricalysia Sonderiana.
floribunda, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 21.—Afr. austr.
lanceolata, *Sond. in Linnaea*, xxiii. (1850) 53.—Afr. austr.
pavettoides, Harv. in Harv. & Sond. Fl. Cap. iii. 22 (= *Webera pavettoides*).—Afr. austr.

KRAUSSIA, Sch. Bip. in Flora, xxvii. (1844) 672 = **Amellus**, Linn. (Compos.).
capensis, Sch. Bip. l. c. = A. Lychnitis.

KREBSIA, Eckl. & Zeyh. Enum. 179 (1836) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
argentea, Eckl. & Zeyh. l. c. = Lebeckia microphylla.
Capnitidis, Steud. Nom. ed. II. i. 850 = L. Capnitis.
carnosa, Eckl. & Zeyh. Enum. 180 = L. carnosa.
cytisoides, Steud. Nom. ed. II. i. 850 = L. cytisoides.
divaricata, Eckl. & Zeyh. Enum. 179 = L. divaricata.
eriocarpa, Steud. Nom. ed. II. i. 850 = L. cytisoides.
genuflexa, Steud. l. c. = L. genuflexa.
striata, Steud. l. c. = L. carnosa.
stricta, Eckl. & Zeyh. Enum. 179 = L. cytisoides.

KREBSIA, Harv. Gen. S. Afr. Pl. ed. II. 233 (1868) = **Gomphocarpus**, R. Br. (Asclep.).

KREIDEK, Adans. Fam. ii. 225 (1763) = **Scoparia**, Linn. (Scrophul.).

KREMERIA, Coss. & Dur. in Bull. Soc. Bot. Fr. iii. (1856) 671 = **Muricaria**, Desv. (Crucif.).
Cordylocarpus, Coss. & Dur. l. c. 672 = M. prostrata?

KREMERIA, Dur. in Duch. Rev. Bot. i. (1846) 64 = **Chrysanthemum**, Linn. (Compos.).
paludosa, Dur. l. c. 365 = C. paludosum.

KREYSIGIA, Reichb. Ic. Exot. iii. 11. t. 229 (1829).
LILIACEAE, Benth. & Hook. f. iii. 829.
TRIPLADENIA, D. Don, in Proc. Linn. Soc. i. (1839) 46.
Cunninghami, *F. Muell. Fragm.* vii. 71.—Austral.
multiflora, *Reichb. Ic. Exot.* iii. 11. t. 229.—Austral.

KRIGIA, Schreb. Gen. 532 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 507.
ADOPOGON, Neck. Elem. i. 55 (1790).
CYNTHIA, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 309.
LUTHERA, Sch. Bip. in Linnaea, x. (1836) 257.
SERINIA, Rafin. Fl. Ludov. 149 (1817).
amplexicaulis, *Nutt. Gen. Am.* ii. 127.—Am. bor.
bellidioides, Scheele, in Linnaea, xxv. (1852) 257 = occidentalis.
caroliniana, Hook. Comp. Bot. Mag. i. (1835) 100 = K. Dandelion.
caroliniana, Nutt. Gen. Am. ii. 126 = virginica.
chilensis, *Bert. in 'Merc. Chil.'* (1829); *et in Bull. Férussac*, xx. (1830) 112.—Chili.
chilensis, Nees, ex DC. Prod. vii. 89 = Microseris pygmaea.
Dandelion, *Nutt. Gen. Am.* ii. 127.—Am. bor.
dichotoma, Nutt. l. c. = virginica.
integrifolia, Rafin. New Fl. Am. iv. 86 (Quid?).—Am. bor.
leptophylla, DC. Prod. vii. 88 = virginica.
montana, *Nutt. Gen. Am.* ii. 127.—Am. bor.
nervosa, Hook. Ic. Pl. t. 237 = occidentalis.
occidentalis, *Nutt. in Journ. Acad. Philad.* vii. (1834) 104.—Am. bor.
oppositifolia, Rafin. Fl. Ludov. 57 = virginica.
petiolaris, Rafin. New Fl. Am. iv. 86 (Quid?).—Am. bor.
pinnatifida, Bert. ex DC. Prod. vii. 89 = Microseris pygmaea.
pusilla, Hort. ex DC. l. c. = Microseris pygmaea.
tenella, Reichb. Ic. Fl. Germ. i. 72 = virginica.
virginica, *Willd. Sp. Pl.* iii. 1618.—Am. bor.

KROCKERIA, Neck. Elem. ii. 317 (1790) = **Xylopia**, Linn. (Anonac.).

KROCKERIA, Steud. Nom. ed. II. i. 458 (1821) = KROKERIA, Moench = **Lotus**, Linn. (Legumin.).

KROKERIA, Endl. Gen. 832, in syn. (1841) = KROCKERIA, Neck. = **Xylopia**, Linn. (Anonac.).

KROKERIA, Moench, Meth. 143 (1794) = **Lotus**, Linn. (Legumin.).
edulis, Willk. & Lange, Prod. Fl. Hisp. iii. 340 = L. edulis.
oligoceratos, Moench, Meth. 143 = L. edulis.

KROMBHOLTZIA, Benth. in Journ. Linn. Soc.. xix. (1881) 121 = seq.

KROMBHOLZIA, Rupr. ex Gal. in Bull. Acad. Brux. ix. II. (1844) 247 = **Zeugites**, Schreb. (Gramin.).
latifolia, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 464 = Z. latifolia.
mexicana, Rupr. in Bull. Acad. Brux. ix. II. (1844) 247 = Z. Galeottiana.

KROMON, Rafin. Fl. Tellur. ii. 21 (1836) = **Allium**, Tourn. (Liliac.).
parviflorum, Rafin. l. c. = A. parviflorum.

KRUBERA, Hoffm. Gen. Umbell. 103 (1814) = **Capnophyllum**, Gaertn. (Umbellif.).
caffra, Eckl. & Zeyh. Enum. 353 (= *Stenosemis caffra*). —Afr. austr.
leptophylla, Hoffm. Gen. Umbell. 104. t. 3 = C. dichotomum.
peregrina, Lowe, Man. Fl. Mad. i. 361 = C. dichotomum.

KRUEGERIA, Scop. Introd. 314 (1777) = **Macrolobium**, Schreb. (Legumin.).

KRUHSEA, Regel, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 122 = **Streptopus**, Michx. (Liliac.).
Tilingiana, Regel, l. c. = S. roseus.
Tilingii, Maxim. in Mém. Sav. Etr. Pétersb. ix. (1859) 278 = praec.

KRYNITZIA, Reichb. Nom. 112 (1841) = seq.

KRYNITZKIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. vii. 52 (1841). *BORAGINEAE*, [Benth. & Hook. f. ii. 851.]
PIPTOCALYX, Torr. in Bot. U. St. Expl. Exped. ii. 413. t. 12 (1874).
affinis, *A. Gray, in Proc. Am. Acad.* xx. (1885) 270.—Am. bor. occ.
alyssoides, *A. Gray, l. c.* 280.—Chili.
ambigua, *A. Gray, l. c.* 273.—Am. bor. occ.
angustifolia, *A. Gray, l. c.* 272.—Calif.
barbigera, *A. Gray, l. c.* 273.—Am. bor. occ.
californica, *A. Gray, l. c.* 266.—Calif.
cedrosensis, Greene, in Bull. Calif. Acad. i. IV. (1885) 204 = maritima.
Chorisiana, *A. Gray, in Proc. Am. Acad.* xx. (1885) 267.—Calif.
circumcissa, *A. Gray, l. c.* 275.—Am. bor. occ.
clandestina, *A. Gray, l. c.* 274.—Chili.
Cooperi, *A. Gray, l. c.* 267.—Arizona.
crassisepala, *A. Gray, l. c.* 268.—Am. bor. occ.
cycloptera, Greene, in Bull. Calif. Acad. i. IV. (1885) 207 = pterocarya.
denticulata, Greene, l. c. 205 = muriculata.
dichotoma, *Greene, l. c.* 206.—Nevada.
dumetorum, *Greene, ex A. Gray, in Proc. Am. Acad.* xx. (1885) 272.—Calif.
Fendleri, *A. Gray, l. c.* 268.—Am. bor. occ.
floribunda, *A. Gray, l. c.* 265.—Mexic.
foliosa, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 205. —Calif.
fulvocanescens, *A. Gray, in Proc. Am. Acad.* xx. (1885) 280.—Am. bor. occ.
glomerata, *A. Gray, l. c.* 279.—Am. bor. occ.
heliotropioides, *A. Gray, l. c.* 265.—Texas.
holoptera, *A. Gray, l. c.* 276.—Arizona.
intermedia, *A. Gray, l. c.* 273.—Calif.
Jamesii, *A. Gray, l. c.* 278.—Am. bor. occ.
Jonesii, *A. Gray, l. c.* 274.—Calif.
leiocarpa, Benth. Pl. Hartw. 326 = oxycarya.
leiocarpa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vii. 52.—Am. bor. occ.
leucophaea, *A. Gray, in Proc. Am. Acad.* xx. (1885) 280.—Am. bor. occ.
linearis, *A. Gray, l. c.*—Chili.
linifolia, *A. Gray, l. c.* 266.—Reg. Andin.
lithocarya, *Greene, ex A. Gray, l. c.* 265.—Calif.
maritima, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 204.—Calif.
micrantha, *A. Gray, in Proc. Am. Acad.* xx. (1885) 275.—Am. bor. occ.
micromeres, *A. Gray, l. c.* 274.—Calif.
microstachys, *Greene, ex A. Gray, l. c.* 269.—Calif.
mohavensis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 207.—Arizona.
mollis, *A. Gray, in Proc. Am. Acad.* xx. (1885) 267.—Calif.
muriculata, *A. Gray, l. c.* 273.—Am. bor. occ.

KRYNITZKIA:—
obovata, *A. Gray, in Proc. Am. Acad.* xx. (1885) 265. —Sibir.
oxycarya, *A. Gray, l. c.* 269.—Am. bor. occ.
oxygona, *A. Gray, l. c.* 277.—Arizona.
Palmeri, A. Gray, l. c. 278 = Jamesii.
Parryi, *A. Gray, l. c.* 265.—Mexic.
Pattersoni, *A. Gray, l. c.* 268.—Colorado.
phacelioides, Fisch. & Mey. ex Phil. f. Cat. Pl. Vasc. Chil. 211 = Eritrichium phacelioides.
plebeia, *A. Gray, in Proc. Am. Acad.* xx. (1885) 266. —Ins. Unalaschka.
pterocarya, *A. Gray, l. c.* 276.—Am. bor. occ.
pusilla, *A. Gray, l. c.* 274.—Texas.
racemosa, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 208.—Calif.
ramosa, *A. Gray, in Proc. Am. Acad.* xx. (1885) 274. —Mexic.
ramosissima, *A. Gray, l. c.* 277.—Am. bor. occ.
rostellata, Greene, in Bull. Calif. Acad. i. IV. (1885) 203 = microstachys.
Scouleri, *A. Gray, in Proc. Am. Acad.* xx. (1885) 267. —Am. bor. occ.
sericea, *A. Gray, l. c.* 279.—Am. bor. occ.
setosissima, *A. Gray, l. c.* 276.—Am. bor. occ.
sparsiflora, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 203.—Calif.
tenuifolia, *A. Gray, in Proc. Am. Acad.* xx. (1885) 267. —Mexic.
texana, *A. Gray, l. c.* 268.—Texas.
Torreyana, *A. Gray, l. c.* 271.—Am. bor. occ.
trachycarpa, *A. Gray, l. c.* 266.—Calif.; Chili.
virgata, *A. Gray, l. c.* 279.—Am. bor. occ.
Watsoni, *A. Gray, l. c.* 271.—Am. bor. occ.

KTENOSACHNE, Steud. Syn. Pl. Gram. 150 (1855) = **Prionachne**, Nees (Gramin.).
tenerrima, Steud. l. c.—Hab.?

KTENOSPERMUM, Lehm. Del. Sem. Hort. Hamb. (1837) 7.—Cf. Linnaea, xiii. (1838) Litt. 84 = **Pectocarya**, DC. (Boragin.).
linifolium, Lehm. ll. cc.—Hab.?

KUA, Rheede, ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 394, cum ic. = **Curcuma**, Linn. (Scitam.).
domestica, Medic. l. c. 396.—Ind. or.

KUALA, Karst. & Triana, in Linnaea, xxviii. (1856) 428 = **Esenbeckia**, Kunth (Rutac.).
alata, Karst. & Triana, l. c. 429 = E. alata.
laevis, Karst. & Triana, l. c. = E. alata.

KUESTERA, Regel, Gartenfl. vi. (1857) 345 = **Beloperone**, Nees (Acanthac.).
aurantiaca, Regel, l. c. = B. aurantiaca.

KUGIA, Bert. ex Lindl. Nat. Syst. ed. II. 438 (1836), nomen (Quid?).

KUHLIA, H. B. & K. Nov. Gen. et Sp. vii. 234. t. 652 (1825). *SAMYDACEAE*, Benth. & Hook. f. i. 798.
brasiliensis, A. Gray, Bot. U. St. Expl. Exped. i. 74 = Banara brasiliensis.
glauca, *H. B. & K. Nov. Gen. et Sp.* vii. 234.—N. Granat.
mollis, Poepp. & Endl. Nov. Gen. et Sp. iii. 74. t. 285 = Banara guianensis.
parviflora, A. Gray, Bot. U. St. Expl. Exped. i. 73 = Banara parviflora.
ulmifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 236. t. 653. —Am. austr.

KUHLIA, Reinw. in Blume, Cat. Gew. Buitenz. 51 (1823), nomen; et ex Blume, Bijdr. 777 (1826) = **Fagraea**, Thunb. (Logan.).
morindaefolia, Reinw. ll. cc. = F. morindaefolia.

KUHNIA, Linn. Sp. Pl. ed. II. 1662 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 248.
albicaulis, Rafin. New Fl. Am. iv. 105 (Quid?).—Am. bor.
altaica, Rafin. l. c. 79 = eupatorioides.
arabica, Hochst. & Steud. ex DC. Prod. vii. 267 = Pegolettia senegalensis.
arguta, *H. B. & K. Nov. Gen. et Sp.* iv. 105.—N. Granat.

KUHNIA:—
baccharoides, DC. Prod. v. 127 = Symphyopappus viscosus.
cinerea, Rafin. New Fl. Am. iv. 106 (Quid?).—Am. bor.
Critonia, Willd. Sp. Pl. iii. 1773 = eupatorioides.
dasypia, Rafin. New Fl. Am. i. 79 = eupatorioides.
divaricata, Rafin. l. c. iv. 105 (Quid?).—Am. bor.
elliptica, Rafin. l. c. i. 76, 79 = eupatorioides.
eupatorioides, *Linn. Sp. Pl.* ed. II. 1662.—Am. bor.
frutescens, Hornem. Hort. Hafn. ii. 79 = rosmarinifolia.
fruticosa, Vent. ex Steud. Nom. ed. I. 322, in syn. = praec.
fulva, Rafin. New Fl. Am. i. 80 = eupatorioides.
glabra, Rafin. l. c. 79 = eupatorioides.
glabrata, DC. Prod. v. 127 = Eupatorium oblongifolium.
glomerata, Rafin. New Fl. Am. iv. 105 (Quid?).—Am. bor.
glutinosa, DC. Prod. v. 127 = Eupatorium altissimum.
glutinosa, Ell. Sketch, ii. 292 = eupatorioides.
hirtiflora, DC. Prod. v. 127 = Trichogonia hirtiflora.
icosantha, Sch. Bip. in Linnaea, xxii. (1849) 572 = Trichogonia podocarpa.
latifolia, Rafin. New Fl. Am. iv. 105 (Quid?).—Am. bor.
leptophylla, Scheele, in Linnaea, xxi. (1848) 598 = rosmarinifolia.
linearifolia, Spreng. Syst. iii. 439 = Jaumea linearis.
macrantha, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = eupatorioides.
Maximiliani, Sinning, in Otto & Dietr. Allg. Gartenz. vii. (1839) 138 = eupatorioides.
media, Rafin. New Fl. Am. i. 76 = eupatorioides.
multiramea, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 168 = Ophryosporus triangularis.
paniculata, Cass. in Dict. Sc. Nat. xxiv. 516 = eupatorioides.
pinnata, J. F. Gmel. Syst. 375 = Petalostemon corymbosum.
podocarpa, DC. Prod. v. 127 = Trichogonia podocarpa.
pubescens, Rafin. New Fl. Am. i. 80 = eupatorioides.
pyrifolia, *Mart. Hort. Erlang.* 175; *ex Steud. Nom.* ed. II. i. 851.—Hab.?
rosmarinifolia, *Vent. Jard. Cels*, t. 91.—Mexic.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 207.—Mexic.
spicata, Baill. Hist. Pl. viii. 298 = Liatris spicata.
suaveolens, Fresen. Enum. Sem. Hort. Francof. (1838). —Cf. Linnaea, xiii. (1839) Litt. 94 = eupatorioides.
tuberosa, Rafin. New Fl. Am. i. 80 = eupatorioides.
villosa, Spreng. Syst. iii. 439 = Trichogonia villosa.
virgata, Rafin. New Fl. Am. iv. 105 (Quid?).—Am. bor.

KUHNISTERA, Lam. Encyc. iii. 370 (1789) = **Petalostemon**, Michx. (Legumin.).
carolinensis, Lam. l. c. = P. corymbosum.
violacea, Ait. ex Steud. Nom. ed. II. i. 851 = P. violaceum.

KUHNISTRA, Endl. Gen. 1270 (1840) = praec.

KUKOLIS, Rafin. Sylva Tellur. 55 (1838) = **Hebecladus**, Miers (1845) (Solanac.).
bicolor, Rafin. l. c. = H. bicolor.

KUMARA, Medic. Theodora, 69. t. 4 (1786) = **Aloë**, Tourn. (Liliac.).
disticha, Medic. l. c. 70. t. 4 = A. plicatilis.

KUMBAYA, Endl. ex Steud. Nom. ed. II. i. 663, in syn. (1840) = **Gardenia**, Ellis (Rubiac.).

KUMLIENIA, Greene, in Bull. Calif. Acad. i. iv. (1885) 337. *RANUNCULACEAE.*
hystricula, *Greene, l. c.*—Am. bor. occ.

KUMMERIA, Mart. Herb. Fl. Bras. n. 1276 (1840?) = **Discophora**, Miers (Olacin.).
brasiliensis, Mart. l. c. = D. guianensis.

KUNDA, Rafin. Fl. Tellur. ii. 82 (1836) = **Amorphophallus**, Blume (Aroid.).
verrucosa, Rafin. l. c. = A. campanulatus.

KUNDMANNIA, Scop. Introd. 116 (1777). *UMBELLIFERAE*, Benth. & Hook. f. i. 903.
BRIGNOLIA, Bertol. in Desv. Journ. Bot. iv. (1815) 76.
CAMPDERIA, Lag. Amen. Nat. ii. 99 (1821).
sicula, *DC. Prod.* iv. 143.—Reg. Mediterr.
syriaca, *Boiss. Diagn.* Ser. I. x. 31.—Syria.

KUNIRIA, Rafin. Fl. Tellur. iv. 61 (1836) = **Diclíptera**, Juss. (Acanthac.).
falcata, Rafin. l. c. = D. laevigata.
malabarica, Rafin. l. c. (Quid?).—Ind. or.

KUNOKALE, Rafin. Fl. Tellur. iii. 12 (1836) = **Fagopyrum**, Tourn. (Polygonac.).
carneum, Rafin. l. c. = F. esculentum.

KUNTHIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Garuga**, Roxb. (Burser.).
cochinensis, Dennst. l. c. = G. pinnata.

KUNTHIA, Humb. & Bonpl. Pl. Aequin. ii. 128. t. 122 (1809) = **Chamaedorea**, Willd. (Palm.).
caryotoides, Warsc. in Otto & Dietr. Allg. Gartenz. xiii. (1845) 253, nomen.—N. Granat.
Deppii, Hort. ex Zucc. l. c. ii. (1834) 245 = C. elegans.
montana, Humb. & Bonpl. Pl. Aequin. ii. 128. t. 122 (= *Morenia Lindeniana*).—N. Granat.
xalapensis, Otto & Dietr. ex H. Wendl. in Kerch. Palm. 249 = seq.
xalappensis, Hort. ex H. Wendl. Index Palm. (1854) 15 = C. Schiedeana.

KUNTIA, Dum. Anal. Fam. 56 (1829) = praec.

KUNZEA, Reichb. Consp. 175 (1828). *MYRTACEAE*, Benth. & Hook. f. i. 703.
PENTAGONASTER, Klotzsch, in Otto & Dietr. Allg. Gartenz. iv. (1836) 113.
SALISIA, Lindl. Swan River App. 10 (1839).
Baxteri, *Schau. in Lehm. Pl. Preiss.* i. 123.—Austral.
brachyandra, F. Muell. Fragm. ii. 27 = Leptospermum abnorme.
calida, *F. Muell. l. c.* vi. 23.—Austral.
capitata, *Reichb. Consp.* 175.—Austral.
corifolia, *Reichb. l. c.*—Austral.
ericifolia, F. Muell. in Trans. Vict. Inst. i. (1855) 123 = Muelleri.
ericifolia, *Reichb. Consp.* 175.—Austral.
eriocalyx, *F. Muell. Fragm.* ii. 28.—Austral.
hirsuta, Turcz. in Bull. Soc. Nat. Mosc. (1862) ii. 326 = capitata.
leptospermoides, F. Muell. in Nederl. Kruidk. Arch. iv. (1859) 146 = peduncularis.
micrantha, *Schau. in Lehm. Pl. Preiss.* i. 125.—Austral.
micromera, *Schau. l. c.* 223.—Austral.
Muelleri, *Benth. Fl. Austral.* iii. 113.—Austral.
oligandra, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 336.—Austral.
opposita, *F. Muell. Fragm.* vi. 24.—Austral.
parvifolia, *Schau. in Lehm. Pl. Preiss.* i. 123.—Austral.
pauciflora, *Schau. l. c.* 124.—Austral.
peduncularis, *F. Muell. in Trans. Vict. Inst.* i. (1855) 124.—Austral.
pelagia, F. Muell. in Nederl. Kruidk. Arch. iv. (1859) 145 = corifolia.
podantha, F. Muell. Fragm. ii. 28 = Leptospermum erubescens.
pomifera, *F. Muell. in Trans. Vict. Inst.* i. (1855) 124. —Austral.
praestans, Schau. in Lehm. Pl. Preiss. i. 124 = recurva.
Preissiana, *Schau. l. c.* 125.—Austral.
propinqua, Schau. l. c. 126 = ericifolia.
recurva, *Schau. l. c.* 125.—Austral.
Schaueri, Schau. l. c. 123 = capitata.
sericea, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) i. 162.—Austral.
sprengelioides, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 336.—Austral.
squarrosa, Turcz. l. c. 335 = recurva.
vestita, Schau. in Lehm. Pl. Preiss. i. 126 = ericifolia.
villiceps, Schau. l. c. 125 = Preissiana.

KUNZIA, Spreng. Syst. ii. 475 (1825) = **Purshia**, DC. (Rosac.).
tridentata, Spreng. l. c. = P. tridentata.

KUNZMANNIA, Klotzsch & Schomb. Fauna & Fl. Brit. Gui. 1098 (1843), nomen. *TEREBINTHACEAE.*
Roraimae, *Klotzsch & Schomb. l. c.* nomen.—Guian.

KURITIS, Rafin. Fl. Tellur. ii. 63 (1836) = **Selago**, Linn.
ciliata, Rafin. l. c. 64 = S. ciliata.
Gilii, Rafin. l. c. = S. myrtifolia.

KURKAS, Rafin. Sylva Tellur. 62 (1838) = **Croton**, Linn. (Euphorb.).
acuminatum, Rafin. l. c. = Mallotus japonicus.
congestum, Rafin. l. c. = C. congestus.
laxiflorum, Rafin. l. c. = C. laxiflorus.
populifolium, Rafin. l. c. = C. populifolius.
quadrisetosum, Rafin. l. c. = C. quadrisetosus.
Tiglium, Rafin. l. c. = C. Tiglium.

KURRIA, Hochst. & Steud. in Flora, xxiv. (1841) i. Intell. 28; xxv. (1842) 233 = **Hymenodictyon**, Wall. (Rubiac.).
floribunda, Hochst. & Steud. ll. cc. 28; 234 = H. Kurria.

KURRIA, Steud. Nom. ed. II. i. 851 (1840) = **Hermannia**, Linn. (Sterculiac.).
arabica, Steud. l. c. = H. arabica.

KURRIMIA, Wall. Cat. n. 4334 (1831). *CELASTRINEAE*, Benth. & Hook. f. i. 365.
BHESA, Buch.-Ham. in Edinb. Phil. Journ.. xvi. (Apr. 1834) 315, partim.
NOTHOCNESTIS, Miq. Fl. Ind. Bat. Suppl. 530 (1860).
PYROSPERMUM, Miq. l. c. 402 (1860).
TROCHISANDRA, Bedd. Fl. Sylv. i. t. 120 (1871).
bipartita, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 622.—Ind. or.
calophylla, Wall. Cat. n. 4335 = pulcherrima.
ceylanica, *Arn. in Nov. Act. Nat. Cur.* xviii. i. (1836) 328.—Zeylan.
macrophylla, Wall. Cat. n. 7200 = Itea macrophylla.
Maingayi, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 622. —Singapore.
paniculata, *Wall. Cat.* n. 4336.—Malaya.
pulcherrima, *Wall. l. c.*—Ind. or.; Malaya.
robusta, Kurz, ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 622 = pulcherrima.

KUSCHAKEWICZIA, Regel & Smirn. in Act. Hort. Petrop. v. (1877) 625. *BORAGINEAE.*
turkestanica, *Regel & Smirn. l. c.* 626.—Turkest.

KUSTERA, Benth. & Hook. f. Gen. ii. 1111 (1876) = KUESTERA, Regel = **Beloperone**, Nees (Acanth.).

KUTCHUBAEA, Fisch. ex DC. Prod. iv. 373 (1830). *RUBIACEAE* [Benth. & Hook. f. ii. 98].
KOTCHUBAEA, Regel, ex Benth. & Hook. f. l. c. (1862).
insignis, *Fisch. ex DC. l. c.*—Guiana.

KYBERIA, Neck. Elem. i. 42 (1790) = **Bellis**, Linn. (Compos.).

KYDIA, Roxb. Hort. Beng. 50 [97] (1814); Pl. Corom. iii. 11. tt. 215, 216 (1819). *MALVACEAE*, Benth. & Hook. f. i. 203.
angustifolia, Arn. in Nov. Act. Nat. Cur. xviii. i. (1836) 322 = Julostyles angustifolia.
axillaris, Thw. Enum. Pl. Zeyl. 30 = Dicellostyles axillaris.
calycina, *Roxb. Hort. Beng.* 50; *Pl. Corom.* iii. 11. t. 215.—Ind. or.; Burma.
fraterna, Roxb. Pl. Corom. iii. t. 216 = calycina.
glabrescens, *Mast. in Hook. f. Fl. Brit. Ind.* i. 348.—Reg. Himal.; Malaya.
jujubifolia, *Griff. Notul.* iv. 534.—Zeylan.
paterna, Roxb. Hort. Beng. [97] = calycina.
pulverulenta, Buch.-Ham. in Wall. Cat. n. 1176 = calycina.
Roxburghiana, Wight, Ic. iii. t. 881 = calycina.

KYLINGA, Roem. & Schult. Syst. ii. 236 (1817) = seq.

KYLINGIA, Stokes, Bot. Mat. Med. i. 120 (1812) = seq.

KYLLINGA, Rottb. Desc. et Ic. 12. t. 4 (1773). *CYPERACEAE*, Benth. & Hook. f. iii. 1045.
CYPROLEPIS, Steud. in Flora, xxxiii. (1850) 229.
HEDYCHLOË, Rafin. Ann. Nat. (1820) 16.
LYPROLEPIS, Steud. Syn. Pl. Cyp. 130 (1855).
THYOCEPHALON, Forst. Char. Gen. 129. t. 65 (1776).

KYLLINGA:—
alata, *Nees, in Linnaea*, ix. (1834) 286, nomen; *et* x. (1836) 139.—Afr. austr.
alba, *Nees, l. c.* x. (1836) 140.—Afr. austr.
alba, Steud. in Flora, xvi. (1842) 597 = controversa.
albescens, Steud. Syn. Pl. Cyp. 68 = Lipocarpha argentea.
allata, Nees, in Linnaea, x. (1836) 139 = alata.
ambigua, Steud. Syn. Pl. Cyp. 70 = Scirpus Steudneri.
aphylla, Kunth, Enum. Pl. ii. 127 = vaginata.
aromatica, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 146 = elatior.
atrosanguinea, Steud. in Flora, xxv. (1842) 599 = pulchella.
aurata, *Nees, in Linnaea*, x. (1836) 139.—Afr.; Madag.
aurea, T. Thoms. in App. Speke, Journ. 654 = macrocephala.
bifolia, Miq. Fl. Ind. Bat. iii. 293 = obtusata.
biglumis, Steud. Nom. ed. II. i. 851 = Mariscus biglumis.
bracheilema, *Steud. in Flora*, xxv. (1842) 598.—Abyssin.
brevifolia, *Rottb. Desc. et Ic.* 13. t. 4. f. 3.—Amphig. trop.
bulbocaulis, *Boeck. in Flora*, lviii. (1875) 258.—Afr. trop.
bulbosa, *Beauv. Fl. Owar.* i. 11. t. 8. f. 1.—Afr. trop.
bulbosa, Koen. ex Vahl, Enum. ii. 376, in nota = Mariscus umbellata.
bulbosa, Steud. Syn. Pl. Cyp. 62 = Mariscus bulbocaulis.
caespitosa, Nees, in Mart. Fl. Bras. ii. 1. 12 (syn. Kunth excl.) = pumila.
cajennensis, Auct. vide *cayennensis*.
campestris, Salzm. ex Steud. Syn. Pl. Cyp. 68 = odorata.
capensis, *Steud. in Flora*, xii. (1829) 153.—Afr. austr.
capillaris, Sw. Prod. Veg. Ind. Occ. 20 = Mariscus filiformis.
capitata, Beauv. Fl. Owar. i. t. 31 = vaginata.
cayennensis, Lam. Illustr. i. 149 = Mariscus elatus.
chlorotropis, *Steud. in Flora*, xxv. (1842) 598.—Abyssin.
ciliata, Kunth, Enum. Pl. ii. 136 = Lehmanni.
consanguinea, Kunth, l. c. 135 = aurata.
controversa, *Steud. Syn. Pl. Cyp.* 70.—Abyssin.
crassipes, *Boeck. in Flora*, xlii. (1859) 444.—Afr. trop.
cristata, Afzel. ex A. Rich. Tent. Fl. Abyss. ii. 493 = squamulata.
cristata, Kunth, Enum. Pl. ii. 136 = alba.
cruciata, Nees, in Linnaea, ix. (1834) 286 = brevifolia.
cruciformis, Schrad. ex Schult. Mant. ii. 107 = brevifolia.
cylindrica, *Nees, in Wight, Contrib.* 91.—Geront. trop.
cyperina, Retz. Obs. vi. 21 = Mariscus umbellatus.
cyperoides, Roxb. Hort. Beng. 81; Fl. Ind. i. 182 = Courtoisia cyperoides.
decora, Steud. Syn. Pl. Cyp. 317 = Ascolepis brasiliensis.
dentata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 493 = squamulata.
dipsacoides, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 41 = squamulata.
dorsocrena, Nees.—Cf. Linnaea, xxxv. (1867–68) 430 = controversa.
elata, Steud. Syn. Pl. Cyp. 70 = polyphylla.
elatior, *Kunth, Enum. Pl.* ii. 135.—Afr. austr.
elongata, H. B. & K. Nov. Gen. et Sp. i. 211 = odorata.
erecta, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 42 = obtusata.
eriocauloides, Steud. in Flora, xxv. (1842) 597 = Ascolepis eriocauloides.
exigua, *Boeck. in Bremen, Abh.* vii. (1880) 36.—Madag.
filiformis, Sw. Prod. Veg. Ind. Occ. 20 = Mariscus filiformis.
fraterna, Steud. in Lechl. Berb. Am. Austr. 56 = pumila.
fuscata, Miq. Fl. Ind. Bat. iii. 294 = brevifolia.
fuscescens, Boeck. in Linnaea, xxxv. (1867–68) 421 = melanosperma, brevifolia.
geminiflora, Steud. Syn. Pl. Cyp. 70 = triceps.
globosa, Beauv. Fl. Owar. i. t. 31 = vaginata.
gracilis, Kunth, Enum. Pl. ii. 134 = brevifolia.
gracilis, Zoll. Syst. Verz. 63 = monocephala.
gracillima, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 142 = brevifolia.
granularis, Desf. ex Link, Jahrb. i. III. 91 = Mariscus gracilis.
Hohenackeri, Hochst. ex Steud. Syn. Pl. Cyp. 60 = brevifolia.

KYLLINGA:—
Honolulu, Steud. ex Jard. in Bull. Soc. Linn. Normand. Sér. II. ix. (1875) 278, 281 = brevifolia.
hortensis, Salzm. ex Schlecht. in Bot. Zeit. vii. (1849) 117 = pumila.
inaurata, Nees, ex Boeck. in Linnaea, xxxv. (1867–68) 406 = tetragona.
incompleta, Jacq. Coll. iv. 101; et Ic. t. 300 = Mariscus elatus.
intermedia, R. Br. Prod. 219 = brevifolia.
intermedia, Seem. in Bonplandia, ix. (1861) 261 = monocephala.
involucrata, Boj. ex Baker, Fl. Maurit. 415 = polyphylla.
laxa, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 14 = brevifolia.
Lehmanni, *Nees, in Linnaea*, x. (1835–36) 139.—Afr. austr.
leucantha, Boeck. l. c. xxxviii. (1874) 356 = macrocephala.
leucocephala, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 170 = odorata.
leucocephala, Boeck. in Flora, lviii. (1875) 257 = macrocephala.
Liebmanni, Steud. Syn. Pl. Cyp. 67 = Mariscus gracilis.
longiculmis, Miq. Fl. Ind. Bat. iii. 292 = brevifolia.
macrantha, Boeck. in Linnaea, xxxv. (1867–68) 420 = polyphylla.
macrocephala, *A. Rich. Tent. Fl. Abyss.* ii. 491.—Afr. trop.
maculata, Michx. Fl. Bor. Am. i. 29 = Lipocarpha Humboldtiana.
Mariae, Steud. Syn. Pl. Cyp. 69 (Marisci sp.).—Madag.
Martiana, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 14 = odorata.
melanosperma, *Nees, in Wight, Contrib.* 91.—As. et Afr. trop.
Metzii, Hochst. ex Steud. Syn. Pl. Cyp. 70 = squamulata.
microcephala, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 234 = Mariscus gracilis.
microcephala, Steud. Syn. Pl. Cyp. 70 = Scirpus kyllingioides.
mindorensis, Steud. l. c. 67 = monocephala.
monocephala, Linn. f. Suppl. 104 = monocephala, brevifolia.
monocephala, Muhl. Desc. Gram. 3 = odorata.
monocephala, Nees, in Wight, Contrib. 91 = triceps.
monocephala, *Rottb. Desc. et Ic.* 13. t. 4. f. 4.—Geront. trop.
monocephala, Sieber, ex Steud. Nom. ed. II. i. 851 = pumila.
monocephala, Stokes, Bot. Mat. Med. i. 120 = Scirpus Cephalotes.
monocephala, Thunb. Fl. Jap. 35 = brevifolia.
multinervia, Steud. Syn. Pl. Cyp. 317 = Cyperus dubius.
nana, Nees, in Hook. & Arn. Bot. Beech. Voy. 224 = triceps.
Naumanniana, Boeck. in Flora, lxii. (1879) 516 = pumila.
nervosa, *Steud. l. c.* xxv. (1842) 597.—Abyssin.
nivea, Pers. Syn. i. 57 = brevifolia.
obtusata, J. & C. Presl, Rel. Haenk. i. 183.—Amphig. trop.
odorata, H. B. & K. Nov. Gen. et Sp. i. 211 = pumila.
odorata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 233 = brevifolia.
odorata, Thou. ex Steud. Nom. ed. II. i. 851 = polyphylla.
odorata, *Vahl, Enum. Pl.* ii. 382.—Am. trop.
oligostachya, *Boeck. in Linnaea*, xxxv. (1867–68) 407.—Ins. Fiji.
ovularis, Michx. Fl. Bor. Am. i. 29 = Mariscus ovularis.
panicea, Rottb. Desc. et Ic. 15. t. 4. f. 1 = Mariscus umbellatus.
pauciflora, *Ridl. in Trans. Linn. Soc.* II. ii. (1884) 147. t. 23. ff. 1–4.—Afr. trop.
peruviana, *Lam. Encyc.* iii. 366.—Afr. et Am. trop.
polycephala, Link, Enum. Hort. Berol. i. 47 = Cyperus Luzulae.
polyphylla, *Thou. ex Link, Jahrb.* i. III. 91; *Willd. ex Kunth, Enum. Pl.* ii. 134.—Afr. trop.; Maurit.
pulchella, *Kunth, l. c.* 137.—Afr. trop. et austr.
pumila, *Michx. Fl. Bor. Am.* i. 28.—Afr. et Am. trop.
pumila, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 270 = odorata.
pumila, Steud. in Flora, xxv. (1842) 596 = triceps.
Pumilio, *Steud. Syn. Pl. Cyp.* 67.—Malaya.

KYLLINGA:—
pungens, Link, Hort. Berol. i. 326 = obtusata.
punicea, J. F. Gmel. Syst. 122, sphalm. = *panicea*.
repens, Salzm. ex Steud. Syn. Pl. Cyp. 67 = obtusata.
rigida, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 169 = pungens.
rigida, Link, Hort. Berol. ii. 323 = Cyperus umbellatus.
rigidifolia, Boeck. in Linnaea, xxxv. (1867–68) 432 (Marisci sp.).—Cuba.
rigidula, Steud. Syn. Pl. Cyp. 71 (pl. Nubica) = pumila.
robusta, *Boeck. in Linnaea*, xxxv. (1867–68) 409.—Bras.
scirpina, Reichb. ex Steud. Nom. ed. II. i. 470, 852 = Cyperus Luzulae, etc.
sesquiflora, Torr. in Ann. Lyc. N. York, iii. (1836) 287 = odorata.
Sojauxii, Boeck. in Flora, lxii. (1879) 515 = aurata.
sororia, Kunth, Enum. Pl. ii. 131 = brevifolia.
sphaerocephala, *Boeck. in Flora*, lviii. (1875) 258.—Afr. trop.
squamulata, *Vahl, Enum.* ii. 381.—As. et Afr. trop.
squamulosa, Kunth, Enum. Pl. ii. 135 = squamulata.
squarrosa, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 170 = Mariscus flavus.
squarrosa, Steud. Syn. Pl. Cyp. 68 (Lipocarphae sp.).—Ins. Molucc.
stricta, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 13 = obtusata.
sumatrensis, Retz. Obs. iv. 13 = Mariscus umbellatus.
tenuifolia, Steud. Syn. Pl. Cyp. 69 = pumila.
tenuis, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 168 = brevifolia.
tenuis, Boeck. in Linnaea, xxxv. (1867–68) 423 = obtusata.
tenuissima, Steud. Syn. Pl. Cyp. 67 = brevifolia.
tetragona, *Spreng. ex Nees, in Linnaea*, vii. (1832) 512.—Afr. austr.
tibialis, Poit. ex Ledeb. Diss. Pl. Doming. n. 1.—Cf. Schrad. N. Journ. ii. (1807) II, III. 291 = vaginata.
tricephala, Salisb. Prod. 32 = brevifolia.
triceps, Forst. f. Prod. 7 = monocephala.
triceps, Linn. f. Suppl. 104 = monocephala, triceps.
triceps, *Rottb. Desc. et Ic.* 14; *Progr.* 15.—Geront. trop.
triceps, Sw. Obs. Bot. 33 = odorata.
triceps, Thunb. Fl. Jap. 35 = brevifolia.
tuberosa, Scheele, in Linnaea, xvii. (1843) 342 = odorata.
umbellata, Rottb. Desc. et Ic. 15. t. 4. f. 2; Linn. f. Suppl. 105 = Mariscus umbellatus.
umbellata, Sw. ex Kunth, Enum. Pl. ii. 117 = Mariscus elatus.
uncinata, Link, Enum. Hort. Berol. i. 47 = pumila.
vaginata, Lam. Tabl. Encyc. i. 148 = peruviana.
vaginata, Reichb. f. ex Kunth, Enum. Pl. ii. 128 = obtusata.
vaginata, Zoll. Syst. Verz. 63 = melanosperma.
viridiflora, Link, Hort. Berol. ii. 322, in obs. = pumila.
viridiflora, *Roxb. ex Spreng. Cat. Sem. Hal.; ex Steud. Nom.* ed. II. i. 852.—Hab. ?
viridula, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 492 = cylindrica.
Welwitschii, Ridl. in Trans. Linc. Soc. Ser. II. ii. (1884) 147 = triceps.

KYLLINGIA, Linn. f. Suppl. ii. (1781) = praec.

KYMAPLEURA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) errat. [455] = **Troximon**, Nutt. (Compos.).
heterophylla, Nutt. l. c. = T. heterophyllum.

KYRSTENIA, Neck. Elem. i. 81 (1790) = **Eupatorium**, Tourn. (Compos.).

KYRTANDRA, J. F. Gmel. Syst. 37 (1791) = **Cyrtandra**, Forst. (Gesnerac.).

KYRTANTHUS, J. F. Gmel. Syst. 362 (1791) = **Posoqueria**, Aubl. (Rubiac.).
longiflorus, J. F. Gmel. l. c. = P. longiflora.

L

LABATIA, Mart. Nov. Gen. et Sp. ii. 70. tt. 161, 162 (1826). *SAPOTACEAE*, Benth. & Hook. f. ii. 657.
Caimito, Mart. in Flora, xxi. (1838) II. Beibl. 90 = Lucuma Caimito.
chrysophyllifolia, *Griseb. Cat. Pl. Cub.* 166.—Cuba.

LABATIA:—
chrysophylloides, Mart. in Flora, xxi. (1838) II. Beibl. 92 = Lucuma chrysophylloides.
conica, Vell. Fl. Flum. 49; i. t. 125 = macrocarpa, *Mart.*
dictyonura, Griseb. Cat. Pl. Cub. 166 = Lucuma dictyoneura.
elliptica, Pohl, ex Miq. in Mart. Fl. Bras. vii. 75 = Lucuma ramiflora.
glomerata, *Pohl, ex Miq. l. c.* 81.—Bras.
laevigata, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 92.—Bras.
lasiocarpa, Mart. l. c. 91 = Lucuma lasiocarpa.
macrocarpa, *Mart. Nov. Gen. et Sp.* ii. 71. tt. 161, 162.—Bras.
macrocarpa, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 196.—N. Caled.
oblonga, Pohl, ex Mart. Fl. Bras. vii. 74 = Lucuma torta.
parinarioides, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 451.—Bras.
pedunculata, *Willd. Sp. Pl.* i. 624.—Guian.
psammophila, Mart. in Flora, xxi. (1838) II. Beibl. 93 = Lucuma psammophila.
ramiflora, Mart. l. c. = Lucuma ramiflora.
reticulata, Mart. l. c. 90 = Lucuma laurifolia.
salicifolia, *Klotzsch, ex Walp. Rep.* vi. 454.—Bras.
salicifolia, Mart. in Flora, xxi. (1838) II. Beibl. 92 = Lucuma Sellowii.
torta, Mart. l. c. 94 = Lucuma torta.

LABATIA, Scop. Introd. 197 (1777) = **Ilex**, Linn.

LABATIA, Sw. Prod. Veg. Ind. Occ. 32 (1788); Fl. Ind. Occ. 263. t. 6 (1797) = **Lucuma**, Juss. (Sapot.).
sessiliflora, Sw. ll. cc.—Ind. occ.

LABICHEA, Gaudich. ex DC. Prod. ii. 507 (1825); et in Freyc. Voy. Bot. 485. t. 112 (1826). *LEGUMINOSAE*, Benth. & Hook. f. i. 573.
bipunctata, Paxt. Mag. Bot. x. (1843) 150 = lanceolata.
Buettneriana, *F. Muell. in Melb. Chemist* (June 1882).—Austral.
cassioides, *Gaudich. ex DC. Prod.* ii. 507; *et in Freyc. Voy. Bot.* 486. t. 16.—Austral.
digitata, Benth. in Mitch. Journ. Trop. Austral. 273 = rupestris.
diversifolia, Meissn. in Lehm. Pl. Preiss. i. 23 = lanceolata.
lanceolata, *Benth. in Enum. Pl. Hueg.* 41.—Austral.
nitida, *Benth. Fl. Austral.* ii. 293.—Austral.
punctata, *Benth. in Lindl. Swan River App.* 15.—Austral.
rupestris, *Benth. in Mitch. Journ. Trop. Austral.* 342.—Austral.
tephrosiaefolia, *Meissn. in Bot. Zeit.* xiii. (1855) 12.—Austral.

LABILLARDIERA, Roem. & Schult. Syst. v. pp. xxvi, 330 (1819) = **Billardiera**, Sm. (Pittosp.).
canariensis, Roem. & Schult. l. c. 330 = B. scandens.
fusiformis, Roem. & Schult. l. c. = Sollya heterophylla.
longiflora, Roem. & Schult. l. c. = B. longiflora.
mutabilis, Roem. & Schult. l. c. = B. scandens.
scandens, Roem. & Schult. l. c. = B. scandens.

LABISIA, Lindl. Bot. Reg. (1845) t. 48. *MYRSINEAE*, Benth. & Hook. f. ii. 645.
Malouiana, *Linden & Rod. in Illustr. Hortic.* xxxii. (1885) 186. t. 580.—Borneo.
pothoina, *Lindl. Bot. Reg.* (1845) t. 48.—Malaya.
pumila, Benth. & Hook. f. Gen. ii. 645 = pothoina.

LABLAB, Adans. Fam. ii. 325 (1763) = **Dolichos**, Linn. (Legumin.).
benghalensis, Medic. Phil. Bot. i. 205 = D. Lablab.
cultratus, DC. Prod. ii. 402 = D. Lablab.
ferrugineus, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 354 = D. Lablab.
javanicus, Miq. Fl. Ind. Bat. i. 1. 191 = D. Lablab.
leucocarpus, Savi, Diss. (1821) 22 = D. Lablab.
lignosus, R. Grah. in Wall. Cat. n. 5537 = D. Lablab.
martinicensis, Medic. Phil. Bot. i. 205 = D. Lablab.
microcarpus, DC. Prod. ii. 402 = D. Ladlab.
nankinicus, Savi, Diss. (1821) 22 = D. Lablab.
niger, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 354 = D. Lablab.

LABLAB:—
perennans, DC. Prod. ii. 402 = Dolichos Lablab.
purpureus, Sweet, Hort. Brit. ed. I. 481 = D. Lablab.
rufus, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 354 = D. Lablab.
texanus, Scheele, in Linnaea, xxi. (1848) 467 = Galactia texana.
uncinatus, A. Br. ex A. Rich. Tent. Fl. Abyss. i. 225, in nota = D. Lablab.
vulgaris, Savi, Diss. (1821) 19 = D. Lablab.

LABLAVIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 236 (1834) = praec.
vulgaris, D. Don, l. c. = D. Lablab.

LABORDEA, Benth. in Journ. Linn. Soc. i. (1857) 82, sphalm. = seq.

LABORDIA, Gaudich. in Freyc. Voy. Bot. 449. t. 60 (1826). *LOGANIACEAE*, Benth. & Hook. f. ii. 792.
Echitis, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 239.—Ins. Sandvic.
fagraeoidea, *Gaudich. in Freyc. Voy. Bot.* 450. t. 60 (fragraeoidea).—Ins. Sandvic.
fagraeoidea, *A. Gray, in Proc. Am. Acad.* iv. (1858–59) 323 (non praec.).—Ins. Sandvic.
hedyosmifolia, Baill. in Bull. Soc. Linn. Par. i. (1880) 240 = fagraeoidea, *Gaudich.*
hirtella, *H. Mann, in Proc. Am. Acad.* vii. (1867) 196.—Ins. Sandvic.
membranacea, *H. Mann, l. c.* 197.—Ins. Sandvic.
molokaiana, Baill. in Bull. Soc. Linn. Par. i. (1880) 240 = fagraeoidea, *Gaudich.*
pallida, H. Mann, in Proc. Am. Acad. vii. (1867) 196 = fagraeoidea.
sessilis, A. Gray, l. c. iv. (1858–59) 323 = fagraeoide, *Gaudich.*
tinifolia, *A. Gray. l. c.* 322.—Ins. Sandvic.
Waialealae, *Wawra, in Flora*, lv. (1872) 516.—Ins. Sandvic.
Waiolani, Wawra, l. c. = fagraeoidea, *Gaudich.*

LABOUCHERIA, F. Muell. in Journ. Linn. Soc. iii. (1859) 158 = **Erythrophloeum**, Afzel. (Legumin.).
chlorostachya, F. Muell. l. c. 159 = E. Laboucherii.

LABOURDONNAISIA, Boj. in Mém. Soc. Phys. Genèv. ix. (1841) 295, cum ic. *SAPOTACEAE*, Benth. & Hook. f. ii. 659.
albescens, *Benth. in Benth. & Hook. f. Gen.* ii. 660.—Cuba.
calophylloides, *Boj. in Mém. Soc. Phys. Genèv.* ix. (1841) 298.—Ins. Mascar.
discolor, *Sond. in Linnaea*, xxiii. (1850) 73.—Afr. austr.
glauca, *Boj. in Mém. Soc. Phys. Genèv.* ix. (1841) 299.—Ins. Mascar.
revoluta, *Boj. l. c.* 297.—Ins. Mascar.
sarcophleia, *Boj. l. c.*—Ins. Mascar.
sericea, Benth. & Hook. f. Gen. ii. 660 = discolor.

LABOURDONNEIA, Boj. Hort. Maurit. 199 (1837) = praec.

LABRADIA, Swediaur, Arzneymitt. i. (1801) 164 = **Mucuna**, Adans. (Legumin.).
pruriens, Swediaur, l. c. = M. pruriens.
urens, Swediaur, l. c. = M. urens.

LABRAMIA, A. DC. in DC. Prod. viii. 672 (1844). *SAPOTACEAE*, Benth. & Hook. f. ii. 660.
DELASTREA, A. DC. in DC. Prod. viii. 195 (1844).
Bojeri, *A. DC. l. c.* 673.—Cf. 196.—Madag.

LABURNUM, Linn. Syst. ed. I (1735); Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 362. *LEGUMINOSAE*, Benth. & Hook. f. i. 481.
PODOCYTISUS, Boiss. & Heldr. Diagn. Ser. I. ix. 7 (1849).
Adami ×, Lavall. Arb. Segrez. 60 = *Cytisus Adami.*
album, J. Presl, Rostl. iii. 99 = Cytisus albus.
alpinum, *J. Presl, l. c.*—Europ.
Alschingeri, C. Koch & Fint. Wochenschr. ii. (1859) 405 = Cytisus Alschingeri.
anagyroides, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 363.—Europ.
arboreum, J. Presl, Rostl. iii. 100 = Cytisus arboreus.
caramanicum, *Benth. & Hook. f. Gen.* i. 481.—Graecia; As. Min.

LABURNUM:—
fragrans, Griseb. Spicil. Fl. Rumel. i. 7 = Petteria ramentacea.
grandiflorum, J. Presl, Rostl. iii. 100 = Cytisus grandiflorus.
Heuffelii, Wierzb. ex Fuss, Fl. Transs. 152 = Cytisus austriacus.
lanigerum, J. Presl, Rostl. iii. 100 = Calycotome villosa.
molle, J. Presl, l. c. = Cytisus mollis.
monadelphum, Pritz. Ic. Ind. ii. 162, sphalm. = Lacis monadelpha.
nigricanum, Fuss, Fl. Transs. 151 = seq.?
nigricans, J. Presl, Rostl. iii. 99 = Cytisus nigricans.
nubigenum, J. Presl, l. c. = Petteria ramentacea.
patens, J. Presl, l. c. 100 = Cytisus patens.
pendulum, Rafin. Sylva Tellur. 24 = anagyroides.
praecox, Fuss, Fl. Transs. 151 = anagyroides?
purpureum, Drapiez, Encyc. et Fl. Serr. Angl. v. (1837) 6 = Cytisus purpureus.
ramentaceum, C. Koch, Dendrol. i. 20 = Petteria ramentacea.
Rochelii, Wierzb. ex Fuss, Fl. Transs. 153 = Cytisus austriacus.
sessilifolium, J. Presl, Rostl. iii. 99 = Cytisus sessilifolius.
spinosum, J. Presl, l. c. 100 = Calycotome spinosa.
tardiflorum, Hort. in Rev. Hortic. (1878) 120 = anagyroides.
triflorum, J. Presl, Rostl. iii. 100 = Cytisus triflorus.
vulgare, J. Presl, l. c. 99 = anagyroides.
Weldeni, Griseb. ex Lavall. Arb. Segrez. 60 = Petteria ramentacea.

LACAENA, Lindl. Bot. Reg. (1843) Misc. 68. *ORCHIDEAE*, Benth. & Hook. f. iii. 547.
NAUENIA, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 193.
bicolor, *Lindl. Bot. Reg.* (1843) *Misc.* 68.—Mexic.
spectabilis, *Reichb. f. in Bonplandia*, ii. (1854) 92.—Costa Rica.

LACANTHIS, Rafin. Fl. Tellur. ii. 94 (1836) = **Euphorbia**, Linn.
splendens, Rafin. l. c. = E. splendens.

LACARA, Rafin. Fl. Tellur. iv. 70 (1836) = **Campanula**, Linn.
capitata, Rafin. l. c. = C. lingulata.
saxatilis, Rafin. l. c. = C. saxatilis.
scabra, Rafin. l. c. = C. lingulata.
subtriloba, Rafin. l. c.—Hab.?
villosa, Rafin. l. c. = C. lingulata?

LACARA, Spreng. Neue Entdeck. iii. 56 (1822) = **Bauhinia**, Linn. (Legumin.).
triplinervia, Spreng. l. c.—Bras.

LACARIS, Buch.-Ham. ex Pfeiff. Nom. ii. 4 (1871) = **Zanthoxylum**, Linn. (Rutac.).

LACATHEA, Salisb. Parad. Lond. t. 56 (1806) = **Gordonia**, Ellis (Rutac.).
florida, Salisb. l. c. = G. pubescens.

LACCODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 496 = **Cupania**, Plum. (Sapindac.).
ferrugineus, Radlk. l. c. 514 = C. ferruginea.

LACCOSPADIX, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 205 = **Calyptrocalyx**, Blume (Palm.).
australasicus, H. Wendl. & Drude, l. c. 206 = C. australasicus.

LACCOSPERMA, G. Mann & H. Wendl. in Kerch. Palm. 249 (1878) = **Ancistrophyllum**, G. Mann & H. Wendl. (Palm.).
laeve, G. Mann & H. Wendl. l. c.—Afr. trop.
opacum, G. Mann & H. Wendl. l. c.—Afr. trop.

LACELLIA, Viv. Fl. Lib. Spec. 58. t. 22. f. 2 (1824) = **Volutarella**, Cass. (Compos.).
lybica, Viv. l. c. = V. bicolor.

LACEPEDEA, H. B. & K. Nov. Gen. et Sp. v. 142. t. 144 (1821) = **Turpinia**, Vent. (Sapindac.).
insignis, H. B. & K. l. c. = T. insignis.
pinnata, Schlecht. in Linnaea, x. (1836) 240 = T. pinnata.

LACERDAEA, Berg, in Linnaea, xxvii. (1854) 437 = **Campomanesia**, Ruiz & Pav. (Myrtac.).
Luschnathiana, Berg, l. c.—Bras.

LACERPITIUM, Thunb. Prod. Pl. Cap. 50 (1794) = **Laserpitium**, Tourn. (Umbellif.).

LACHANODENDRON, Reinw. ex Blume, Cat. Gew. Buitenz. 70 (1823), nomen = **Lansium**, Rumph. (Meliac.).
album, Reinw. l. c. = seq.?
domesticum, Nees, in Flora, viii. (1825) i. 103 = L. domesticum.

LACHANODES, DC. in Guill. Arch. Bot. ii. (1833) 332 = **Senecio**, Tourn. (Compos.).
cuneifolia, DC. Prod. vi. 443 = S. Leucadendron.
Leucadendron, DC. l. c. = S. Leucadendron.
Pladaroxylon, Endl. Gen. 461 = S. Leucadendron.
prenanthiflora, Burch. ex DC. Prod. vi. 443 = S. prenanthiflorus.

LACHANOSTACHYS, Endl. Gen. Suppl. iii. 64 (1843) = **Lachnostachys**, Hook. (Compos.).

LACHARA, Linn. Syst. ed. II. 22 (1740) = **Lachnaea**, Linn. (Thymel.).

LACHENALIA, Jacq. in Nov. Act. Helv. i. (1787) 39. *LILIACEAE*, Benth. & Hook. f. iii. 807.
BRACHYSCYPHA, Baker, in Journ. Linn. Soc. xi. (1870) 393.
CHLORIZA, Salisb. Gen. Pl. Fragm. 21 (1866).
COELANTHUS, Willd. ex Schult. f. Syst. vii. (1830) p. xlvii, obs.
DIPCADIOIDES, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 432.
HIMAS, Salisb. Gen. Pl. Fragm. 21 (1866).
MONOESTES, Salisb. l. c. (1866).
ORCHIASTRUM, Lem. in Illustr. Hortic. ii. (1855) Misc. 96.
ORCHIOPS, Salisb. Gen. Pl. Fragm. 21 (1866).
PLATYESTES, Salisb. l. c. (1866).
SCILLOPSIS, Lem. in Illustr. Hortic. ii. (1855) Misc. 96; iii. (1856) 33.
TRIALLOSIA, Rafin. Fl. Tellur. iii. 60 (1836).
albida, Tratt. Obs. Bot. Tab. 162 = contaminata.
aloides, Pers. Syn. i. 377 = tricolor.
anguinea, *Sweet, Hort. Brit.* ed. I. 420.—Afr. austr.
angustifolia, Jacq. Coll. Suppl. 47 = contaminata.
aurea, Lindl. in Loud. Encyc. Pl. 1543 = tricolor.
bicolor, Lodd. Bot. Cab. t. 1129 = violacea.
bifolia, Ker-Gawl. in Bot. Mag. t. 1611 = isopetala.
botryoides, Tratt. Obs. Bot. Tab. t. 140 = purpureo-coerulea.
Bowieana, *Baker, in Journ. Linn. Soc.* xi. (1871) 410. —Afr. austr.
campanulata, *Baker, in Journ. Bot.* xii. (1874) 6.—Afr. austr.
carnosa, *Baker, in Journ. Linn. Soc.* xi. (1871) 407.—Afr. austr.
contaminata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 461. —Afr. austr.
convallarioides, *Baker, in Journ. Linn. Soc.* xi. (1871) 407.—Afr. austr.
Cooperi, *Baker, l. c.* 409.—Afr. austr.
declinata, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. iv. 292 = pustulata.
fistulosa, *Baker, in Gard. Chron.* (1884) i. 668.—Afr. austr.
flava, Andr. Bot. Rep. t. 456 = tricolor.
fragrans, *Andr. l. c.* t. 302.—Afr. austr.
fragrans, Jacq. Hort. Schoenb. i. 43. t. 82 = versicolor.
glaucina, *Jacq. Coll. Suppl.* 39.—Afr. austr.
graminifolia, Soland. ex Baker, in Journ. Linn. Soc. xi. (1871) 398 = Dipcadi hyacinthoides.
hirta, *Thunb. Prod. Pl. Cap.* 64.—Afr. austr.
hyacinthoides, Jacq. Coll. Suppl. 58 = contaminata.
hyacinthoides, Lam. Encyc. iii. 373 = orchioides.
isopetala, *Jacq. l. c.* 68.—Afr. austr.
juncifolia, *Baker, in Journ. Linn. Soc.* xi. (1871) 409 —Afr. austr.
lanceaefolia, Jacq. Coll. Suppl. 69; Ic. t. 402 = Scilla lanceaefolia.
latifolia, *Tratt. Obs. Bot. Tab.* 142.—Afr. austr.?
lilacina, *Baker, in Gard. Chron.* (1884) i. 668.—Afr. austr.
liliflora, *Jacq. Coll. Suppl.* 66.—Afr. austr.
linguiformis, Lam. Encyc. iii. 372 = pendula, rubida.
lucida, *Ker-Gawl. in Bot. Mag.* t. 1372.—Afr. austr.
lutea, Sims, Bot. Mag. t. 1907 = tricolor.
luteola, Jacq. Coll. iv. 148 = tricolor.

LACHENALIA:—
macrophylla, Lem. Illustr. Hortic. ii. (1855) Misc. 99 = tricolor.
maculata, Tratt. Obs. Bot. Tab. t. 168 = Scilla lanceaefolia.
mediana, *Jacq. Coll.* iii. 242.—Afr. austr.
mutabilis, Sweet, Hort. Brit. ed. I. 420 = orchioides.
Nelsoni ×, Hort.—Cf. Gard. Chron. (1882) 366; (1884) i. 412.
nervosa, *Ker-Gawl. in Bot. Mag.* t. 1497.—Afr. austr.
odoratissima, *Baker, in Gard. Chron.* (1884) i. 668.—Afr. austr.
orchioides, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 460. —Afr. austr.
orthopetala, *Jacq. Coll.* iii. 240.—Afr. austr.
pallida, [DC. in] Red. Lil. t. 22 = lucida.
pallida, [Soland. in] Ait. Hort. Kew. ed. I. i. 460.—Afr. austr.
patens, *Hort. ex Andr. Bot. Rep.* sub t. 202.—Hab.?
patula, *Jacq. Coll.* iv. 149.—Afr. austr.
pendula, [*Soland. in*] *Ait. Hort. Kew.* i. 461.—Afr. austr.
pulchella, Kunth, Enum. Pl. iv. 284 = orchioides.
punctata, Jacq. Coll. ii. 323 = rubida.
purpurea, Jacq. l. c. Suppl. 65 = versicolor.
purpureo-coerulea, *Jacq. l. c.* 63.— Afr. austr.
pusilla, Eckl. ex Kunth, Enum. Pl. iv. 283 = reflexa.
pusilla, *Jacq. Coll. Suppl.* 71.—Afr. austr.
pustulata, Berg. ex Kunth, Enum. Pl. iv. 284 = orchioides.
pustulata, *Jacq. Coll.* iii. 244.—Afr. austr.
pyramidalis, *Dehnh. in Rev. Napol.* i. iii. 162; *ex Walp. Ann.* i. 853.—Afr. austr.
quadricolor, Jacq. Coll. Suppl. 62 = tricolor.
racemosa, *Ker-Gawl. in Bot. Mag.* t. 1517.—Afr. austr.
ramosa, Lam. Encyc. iii. 373 = Phormium tenax.
reclinata, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. iv. 292 = pustulata.
reflexa, Andr. Bot. Rep. t. 299 = Scilla lanceolata.
reflexa, *Thunb. Prod. Pl. Cap.* 64.—Afr. austr.
rosea, Andr. Bot. Rep. t. 296 = isopetala.
rubida, *Jacq. Coll. Suppl.* 60.—Afr. austr.
sanguinolenta, Willd. ex Kunth, Enum. Pl. iv. 286 = isopetala.
serosa, Steud. Nom. ed. II. ii. 2, sphalm. = seq.
serotina, Willd. Sp. Pl. ii. 175 = Dipcadi erythraeum.
sessiliflora, Andr. Bot. Rep. t. 460 = glaucina.
sessilifolia, Baker, in Journ. Linn. Soc. xi. (1871) 405, err. typ. = praec.
speciosa, F. G. Dietr. Vollst. Lexic. Gaertn. ix. 358 = Dipcadi glaucum.
tenella, *Schneev. ex Steud. Nom.* ed. I. 459 (Quid?).—Hab.?
tigrina, Jacq. Coll. Suppl. 67 = rubida.
trichophylla, *Baker, in Journ. Bot.* xii. (1874) 368.—Afr. austr.
tricolor, *Jacq. in Nov. Act. Helv.* i. (1787) 39.—Afr. austr.
unicolor, Eckl. ex Kunth, Enum. Pl. iv. 284 = orchioides.
unicolor, Jacq. Coll. Suppl. 61 = versicolor.
unifolia, *Jacq. Hort. Schoenb.* i. 43.—Afr. austr.
ustulata, Banks, ex Schult. f. Syst. vii. 601 = orthopetala.
versicolor, *Baker, in Journ. Linn. Soc.* xi. (1871) 409. —Afr. austr.
violacea, *Jacq. Coll.* iv. 147.—Afr. austr.
viridis, Thunb. Prod. Pl. Cap. 64 = Dipcadi filamentosum.
Wrightii, *Baker, in Journ. Bot.* xvi. (1878) 322.—Afr. austr.
Zeyheri, *Baker, in Journ. Linn. Soc.* xi. (1871) 407.—Afr. austr.

LACHNAEA, Linn. Gen. ed. II. 165 (1742). *THYMELAEACEAE*, Benth. & Hook. f. iii. 196.
LACHARA, Linn. Syst. ed. II. 22 (1740).
RADOJITSKYA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 176.
alpina, *Meissn. in DC. Prod.* xiv. 578.—Afr. austr.
ambigua, *Meissn. in Linnaea*, xiv. (1840) 417.—Afr. austr.
aurea, *Eckl. & Zeyh. ex Meissn. in DC. Prod.* xiv. 576. —Afr. austr.
axillaris, *Meissn. in Linnaea*, xiv. (1840) 422.—Afr. austr.
Burchellii, *Meissn. l. c.* 420.—Afr. austr.
buxifolia, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 575 = macrantha.

LACHNAEA:—
buxifolia, Lam. Encyc. iii. 373 = Gnidia filamentosa.
capitata, *Crantz, Inst.* i. 129.—Afr. austr.
ciliata, Crantz, l. c. = Cryptadenia ciliata.
conglomerata, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 578 = capitata.
conglomerata, Linn. Sp. Pl. 560 = Passerina ericoides.
crassifolia, Meissn. in DC. Prod. xiv. 579 = diosmoides.
densiflora, *Meissn. l. c.* 578.—Afr. austr.
diosmoides, *Meissn. in Linnaea*, xiv. (1840) 418.—Afr. austr.
dodecandra, Crantz, Inst. i. 129 = Struthiola erecta.
eriocephala, Linn. Sp. Pl. 560 = Lasiosiphon eriocephalus.
ericoides, *Meissn. in DC. Prod.* xiv. 579.—Afr. austr.
filamentosa, *Meissn. in Linnaea*, xiv. (1840) 410.—Afr. austr.
filiformis, Crantz, Inst. i. 129 = Passerina filiformis.
glauca, Salisb. Parad. Lond. t. 109 = filamentosa.
globulifera, *Meissn. in Linnaea*, xiv. (1840) 412.—Afr. austr.
gracilis, *Meissn. l. c.* 421.—Afr. austr.
grandiflora, Baill. Hist. Pl. vi. 107 = Cryptadenia grandiflora.
hirsuta, Crantz, Inst. i. 129 = Thymelaea hirsuta.
laevigata, Crantz, l. c. = Gnidia oppositifolia.
macrantha, *Meissn. in DC. Prod.* xiv. 575.—Afr. austr.
nervosa, *Meissn. in Linnaea*, xiv. (1840) 579.—Afr. austr.
paleacea, Banks, ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 324 = Passerina paleacea.
penicillata, *Meissn. in Linnaea*, xiv. (1840) 421.—Afr. austr.
phylicoides, Lam. Encyc. iii. 374 = Passerina glomerata.
purpurea, Andr. Bot. Rep. t. 293 = Lasiosiphon eriocephalus.
sericea, Crantz, Inst. i. 129 = Gnidia sericea.
sphaerocephala, Burm. f. Fl. Cap. Prod. 12 = Lasiosiphon eriocephalus.
striata, *Meissn. in Linnaea*, xiv. (1840) 415.—Afr. austr.
uniflora, Crantz, Inst. i. 130 = Cryptadenia uniflora.

LACHNAGROSTIS, Trin. Fund. Agrost. 128 (1820) = **Deyeuxia**, Clar. (Gramin.).
aemula, Nees, ex Hook. f. Handb. N. Zeal. Fl. 329 = D. Forsteri.
Billardieri, Trin. Gram. Unifl. 217 = D. Billardieri.
Chamissonis, Trin. l. c. 216 = D. Chamissonis.
ciliata, Nees, ex Steud. Syn. Pl. Gram. 193 = Calamagrostis ciliata.
ciliata, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 373 = Agrostis ciliata.
filiformis, Trin. Fund. Agrost. 128 = D. Forsteri.
Forsteri, Trin. Gram. Unifl. 217 = D. Forsteri.
Gussonis, Trin. Ic. xx. t. 243 = Triplachne nitens.
Hookeri, Nees, ex Steud. Syn. Pl. Gram. 173 = D. Royleana (in Addend.).
Hookeriana, Nees, ex Steud. l. c. 192 = Calamagrostis Hookeriana.
plebeia, Trin. Gran. Unifl. 217 = D. Forsteri.
phleoides, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 146 = Calamagrostis phleoides.
Preissii, Nees, in Lehm. Pl. Preiss. ii. 97 = D. Forsteri.
retrofracta, Trin. Fund. Agrost. 128 = D. Forsteri.
Royleana, Steud. Syn. Pl. Gram. 173 = D. Royleana (in Addend.).
Roylei, Nees, ex Steud. l. c. 193 = Calamagrostis Royleana.
rudis, Trin. Gram. Unifl. 217 = D. scabra.
scabra, Nees, ex Steud. Nom. ed. II. i. 250 = D. scabra.
Willdenowii, Trin. in Hook. Lond. Journ. Bot. ii. (1843) 412 = D. Forsteri.

LACHNANTHES, Ell. Sketch, i. 47 (1817). *HAEMODORACEAE*, Benth. & Hook. f. iii. 674.
CAMDERIA, Dum. Anal. Fam. 80 (1829).
GYROTHECA, Salisb. in Trans. Hort. Soc. i. (1812) 327.
HERITIERA, J. F. Gmel. Syst. ii. 113 (1791).
tinctoria, *Ell. Sketch*, i. 47.—Am. bor.

LACHNASTOMA, Korth. in Nederl. Kruidk. Arch. ii. ii. (1851) 201 = **Coffea**, Linn. (Rubiac.).
triflorum, Korth. l. c. = C. khasiana.

LACHNEA, Linn. Phil. Bot. 33 (1751) = **Lachnaea**, Linn. (Thymel.).

LACHNIA, Baill. Hist. Pl. vi. 107 (1875) = praec.

LACHNOCAPSA, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 500. *CRUCIFERAE.*
spathulata, *Balf. f. l. c.* 501.—Ins. Socotra.

LACHNOCAULON, Kunth, Enum. Pl. iii. 497 (1841). *ERIOCAULEAE*, Benth. & Hook. f. iii. 1024.
Beyrichianum, Sporl. ex Koern. in Linnaea, xxvii. (1854) 567 = Michauxii.
diandrum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 108.—Texas.
digynum, Koern. in Linnaea, xxvii. (1854) 570 = Michauxii.
glabrum, *Koern. l. c.* 568.—Am. bor.
Michauxii, *Kunth, Enum. Pl.* iii. 497.—Am. bor.

LACHNOCEPHALUS, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 36 = **Mallophora**, Endl. (Verbenac.).
lepidotus, Turcz. l. c. = M. globiflora.

LACHNOCHLOA, Steud. in Flora, xxxiii. (1850), nomen; Syn. Pl. Gram. 5 (1855) ('Genus delendum') Benth. & Hook. f. Gen. iii. 1096.
pilosa, Steud. ll. cc.—Afr. trop.

LACHNOLEPIS, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 132 (1863) = **Gyrinops**, Gaertn. (Thymel.).
moluccana, Miq. l. c.—Ins. Molucc.

LACHNOLOMA, Bunge, Del. Sem. Hort. Dorpat. (1843), p. viii; et in Linnaea, xviii. (1844) 154. *CRUCIFERAE*, Benth. & Hook. f. i. 98.
Lehmanni, *Bunge, ll. cc.*—Turkest.

LACHNOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 571 = **Ratonia**, DC. (Sapind.).
glabrum, Turcz. l. c.—Ins. Philipp.

LACHNOPHYLLUM, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 327 (Rel. Lehm. 151). *COMPOSITAE*, Benth. & Hook. f. ii. 279.
gossypinum, *Bunge, ll. cc.*—Turkest.
Noëanum, *Boiss. Diagn.* Ser. II. iii. 17.—Persia.

LACHNOPODIUM, Blume, in Flora, xiv. (1831) 477 = **Otanthera**, Blume (Melastom.).
bracteatum, Blume, Mus. Bot. Lugd. Bat. i. 56 = O. bracteata.
rubro-limbatum, Blume, in Flora, xiv. (1831) 477 = O. rubro-limbata.

LACHNOPYLIS, Hochst. in Flora, xxvi. (1843) 77 = **Nuxia**, Lam. (Loganiac.).
oppositifolia, Hochst. l. c. = N. oppositifolia.
ternifolia, Hochst. l. c. = N. congesta.

LACHNORHIZA, A. Rich. Fl. Cub. Fanerog. ii. 34 (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 231.
asteroides, *Griseb. Cat. Pl. Cub.* 152.—Cuba.
piloselloides, *A. Rich. Fl. Cub. Fanerog.* ii. 34.—Cuba.

LACHNOSIPHONIUM, Hochst. in Flora, xxv. (1842) 236 = **Randia**, Linn. (Rubiac.).
obovatum, Hochst. l. c. 238 = R. Lachnosiphonium.

LACHNOSPERMUM, Willd. Sp. Pl. iii. 1787 (1803). *COMPOSITAE*, Benth. & Hook. f. ii. 325.
CARPHOLOMA, D. Don, in Mem. Wern. Soc. v. (1826) 555.
ericifolium, *Willd. Sp. Pl.* iii. 1787.—Afr. austr.
ericoides, Harv. in Harv. & Sond. Fl. Cap. iii. 272 = praec.
fasciculatum, Baill. Hist. Pl. viii. 183 = ericifolium.

LACHNOSTACHYS, Hook. Ic. Pl. tt. 414, 415 (1842). *VERBENACEAE*, Benth. & Hook. f. ii. 1139.
PYCNOLACHNE, Turcz. in Bull. Soc. Nat. Mosc. (1863) II. 215.
WALCOTTIA, F. Muell. Fragm. i. 241 (1859).
albicans, *Hook. Ic. Pl.* t. 414.—Austral.
bracteosa, F. Muell. Fragm. viii. 49 = Newcastlia bracteosa.
Cliftoni, *F. Muell. l. c.* ix. 3.—Austral.
ferruginea, *Hook. Ic. Pl.* t. 415.—Austral.
verbascifolia, *F. Muell. Fragm.* vi. 158.—Austral.
Walcottii, *F. Muell. l. c.* ii. 140.—Austral.

LACHNOSTOMA, H. B. & K. Nov. Gen. et Sp. iii. 198. t. 232 (1818). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 767.
IBATIA, Decne. in DC. Prod. viii. 599 (1844).
arizonicum, *A. Gray, in Proc. Am. Acad.* xx. (1885) 296.—Arizona.
Balbisii, Decne. in DC. Prod. viii. 602 = Pherotrichis villosa.
hastulatum, A. Gray, in Proc. Am. Acad. xi. (1876) 87 = Gonolobus hastulatus.
lasiostemma, *Hemsl. Biol. Centr. Am. Bot.* ii. 335.—Guatemala.
maritimum, *Nichols. Dict. Gard.* ii. 226.—Venezuela.
nigrum, *Decne. in DC. Prod.* viii. 602.—Bras.
ovatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 318.—Cuba.
parviflorum, Torr. Bot. Mex. Bound. 165 = Gonolobus parviflorus.
prostratum, Decne. in DC. Prod. viii. 602 = Gonolobus prostratus.
tigrinum, *H. B. & K. Nov. Gen. et Sp.* iii. 199. t. 232.—N. Granat.

LACHNOSTOMA, Korth. ex Flora, xxxiv. (1851) II. 536 = LACHNASTOMA, Korth. = **Coffea**, Linn. (Rubiac.).

LACHNOSTYLIS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 503. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 269.
capensis, *Turcz. l. c.*—Afr. austr.
hirta, Muell. Arg. in DC. Prod. xv. II. 225 = capensis.
minor, Sond. in Linnaea, xxiii. (1850) 132 = capensis.

LACHNOTHALAMUS, F. Muell. Fragm. iii. 156 (1863) = **Chthonocephalus**, Steetz (Compos.).
tomentellus, F. Muell. l. c. = C. tomentellus.

LACHRYMA-JOB, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = **Coix**, Linn. (Gramin.).

LACHRYMARIA, Heist. Syst. 12 (1748) = praec.

LACIMARIA, Hill, Veg. Syst. iv. 49 (1763) = **Liatris**, Schreb. (Compos.).
scariosa, Hill, Hort. Kew. 70 = L. scariosa.
squarrosa, Hill, l. c. = L. squarrosa.

LACIS, Dulac, Fl. Hautes-Pyr. 347 (1867) = **Trinia**, Hoffm. (Umbellif.).
vulgaris, Dulac, l. c. = T. vulgaris.

LACIS, Schreb. Gen. 366 (1789), partim; Lindl. Introd. Nat. Syst. ed. II. 442 (1836). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 109.
aspera, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 73 = Mourera aspera.
aspera, Raeusch. Nom. Bot. ed. III. 158 = Mourera fluviatilis.
Bongardi, Tul. Podost. Monog. 69 = monadelpha.
ceratophylla, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 78 = Podostemon Ceratophyllum.
Chrysanthemum, Schnizl. Ic. t. 85 (= *Neolacis Richardiana*).—Guian.
Dicraea, Steud. Nom. ed. II. ii. 2 = Podostemon minutiflorus, *Dicraeia imbricata.*
disticha, Cham. in Linnaea, viii. (1833) 653 = Podostemon distichus.
fallacissima, Bisch. in Nov. Act. Nat. Cur. xvii. II. (1835) 963 = Apinagia Willdenowii.
fluviatilis, J. G. Gmel. Syst. 834; Willd. Sp. Pl. ii. 1225 = Mourera fluviatilis.
foeniculacea, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 74 = Marathrum foeniculaceum.
foeniculacea, Schlecht. & Cham. in Linnaea, vi. (1831) 42 = Marathrum Schiedeanum.
fucoides, Mart. Nov. Gen. et Sp. i. 5. t. 2 = Apinagia fucoides.
membranacea, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 76 = Apinagia membranacea.
monadelpha, *Bong. l. c.* 78.—Bras.
pygmaea, Bong. l. c. 77 = Apinagia pygmaea.
Riedelii, Bong. l. c. 75 = Apinagia Riedelii.
ruppioides, Bong. l. c. 78 = Apinagia ruppioides.
Schiedeana, Cham. in Linnaea, ix. (1834) 504. t. 6 = Marathrum Schiedeanum.
Wallichii, Steud. Nom. ed. II. ii. 2 = Podostemon Wallichii.

LACISTEMA, Sw. Prod. Veg. Ind. Occ. 12 (1788). *LACISTEMACEAE*, Benth. & Hook. f. iii. 413.
DIDYMANDRA, Willd. Sp. Pl. iv. 971 (1805).
NEMATOSPERMUM, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105.
SYNZYGANTHERA, Ruiz & Pav. Prod. Fl. Per. et Chil. 137. t. 50 (1794).
alterum, *Spreng. Neue Entdeck.* iii. 20.—Am. trop.
angustum, *Schnizl. in Mart. Fl. Bras.* iv. I. 283. t. 80.—Bras.
Berterianum, Schult. Mant. i. 66 = alterum.
Blancheti, *A. DC. in DC. Prod.* xvi. II. 592.—Bras.
coriaceum, *A. DC. l. c.* 593.—Bras.
ellipticum, Schnizl. in Mart. Fl. Bras. iv. I. 285.—Bras.
elongatum, Schnizl. l. c. 282 = myricoides.
floribundum, *Miq. in Linnaea*, xviii. (1844) 24.—Am. trop.
grandifolium, *Schnizl. in Mart. Fl. Bras.* iv. I. 284.—Bras.
intermedium, *Schnizl. l. c.* 281. t. 81. f. 2.—Bras.
lucidum, *Schnizl. l. c.* 282. t. 79. f. 5.—Bras.
myricoides, *Sw. Prod. Veg. Ind. Occ.* 12.—Am. trop.
oblongum, Spreng. Syst. i. 124 = myricoides.
Poeppigii, *A. DC. in DC. Prod.* xvi. II. 593.—Am. trop.
polystachyum, *Schnizl. in Mart. Fl. Bras.* iv. I. 284. t. 78.—Bras.
pubescens, *Mart. Nov. Gen. et Sp.* i. 155. t. 94.—Bras.
purpureum, *A. DC. in DC. Prod.* xvi. II. 593.—Peruv.
recurvum, *Schnizl. in Mart. Fl. Bras.* iv. I. 283. t. 81. f. 4.—Bras.
robustum, *Schnizl. l. c.* 281. t. 77.—Bras.
rostratum, Schnizl. ex A. DC. in DC. Prod. xvi. II. 593 = recurvum.
serrulatum, *Mart. Nov. Gen. et Sp.* i. 156. t. 95.—Bras.

LACMELLIA, Karst. in Linnaea, xxviii. (1856) 449. *APOCYNACEAE*, Benth. & Hook. f. ii. 694.
ZSCHOKKEA, Muell. Arg. in Mart. Fl. Bras. vi. I. 20. tt. 6, 7 (1860).
edulis, *Karst. l. c.* 450.—N. Granat.
floribunda, *Benth. & Hook. f. Gen.* ii. 674.—Am. trop.
lucida, *Miers, Apocyn. S. Am.* 14.—Am. trop.

LACRYMA, Medic. Phil. Bot. i. 177 (1789) = **Coix**, Linn. (Gram.).

LACRYMA-JOB, Tourn. ex Linn. Gen. ed. I. 280 (1737) = praec.

LACTARIA, Rumpf, ex Rafin. Sylva Tellur. 162 (1838); Miq. Fl. Ind. Bat. ii. 414 (1856) = **Ochrosia**, Juss. (Apocyn.).
Ackeringae, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 249 = O. Ackeringae.
calocarpa, Hassk. in Nederl. Kruidk. Arch. iv. (1859) 9 = O. elliptica.
coccinea, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 249 = O. coccinea.
Moorei, F. Muell. Fragm. iii. 110 = O. Moorei.
salubris, Rafin. Sylva Tellur. 162 = O. borbonica.

LACTORIS, Phil. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 521. t. 13. *PIPERACEAE*, Benth. & Hook. f. iii. 127.
ANSONIA, Bert. ex Hemsl. Bot. Challenger, i. III. 53 (1884).
fernandeziana, *Phil. in Verh. Zool.-Bot. Ges. Wien*, xv. (1865) 523.—Ins. J. Fernand.

LACTUCA, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 524.
AGATHYRSUS, D. Don, in Edinb. N. Phil. Journ. (1828–29) 310.
BRACHYRAMPHUS, DC. Prod. vii. 176 (1838).
CEPHALORRHYNCHUS, Boiss. Diagn. Ser. I. iv. 28 (1844).
CHORISMA, D. Don, in Edinb. N. Phil. Journ. (1828–29) 308.
CHORSIS, DC. Prod. vii. 177 (1838).
CICERBITA, Wallr. Sched. Crit. Fl. Hal. 433 (1822).
CYANOSERIS, Schur, Enum. Pl. Transs. 369 (1866).
DUBYAEA, DC. Prod. vii. 247 (1838).

LACTUCA :—
GALATHENIUM, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 442.
IXERIS, Cass. in Dict. Sc. Nat. xxiv. 49 (1822).
LACTUCOPSIS, Sch. Bip. ex Vis. in Mem. Ist. Venet. xv. (1870) 5 (Vis. & Panç. Pl. Serb. Rar. Dec. iii. 5).
MELANOSERIS, Decne. in Jacquem. Voy. Bot. 101. t. 109 (1844).
MICRAUCHENIA, Froel. in DC. Prod. vii. 239 (1838).
MULGEDIUM, Cass. in Dict. Sc. Nat. xxxiii. 296 (1824).
MYCELIS, Cass. l. c. 483 (1824).
PHAENIXOPUS, Cass. l. c. xxxix. 391 (1826).
PHAENOPUS, DC. Prod. vii. 176 (1838).
PYRRHOPAPPUS, A. Rich. Tent. Fl. Abyss. i. 463 (1847).
STEPTORHAMPHUS, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 381 (Rel. Lehm. 205).
WIESTIA, Sch. Bip. in Herb. & Winckl. Jahrb. Pharm. iv. (1841) 154, in nota.
abyssinica, *Fresen. in Mus. Senckenb.* iii. (1839) 72. —Abyssin.
acanthifolia, *Benth. & Hook. f. Gen.* ii. 525.—Ins. Creta.
aculeata, *Boiss. Fl. Orient.* iii. 809.—As. Min.
acuminata, *Spreng. Syst.* iii. 659.—Am. bor.
adenophora, *Boiss. & Kotschy, in Boiss. Fl. Orient.* iii. 814.—Armen.
adulteriana, Gren. & Godr. Fl. Fr. ii. 319 = saligna.
alabamensis, *Bertol. in Nov. Comm. Acad. Bonon.* vii. (1844) 200.—Am. bor.
albana, C. A. Mey. Verz. Pfl. Cauc. 56 = racemosa.
albicaulis, Boiss. Fl. Orient. iii. 809 = L. Scariola.
albiflora, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 531.—Japon.
alpina, *Benth. & Hook. f. Gen.* ii. 525.—Europ.
altaica, *Fisch. & Mey. in Ind. Sem. Hort. Petrop.* xi. 73.—Reg. altaic.
altissima, Bieb. Fl. Taur. Cauc. ii. 242 = sagittata.
ambigua, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 410 = deltoidea.
ambigua, Schrad. Obs. in Cat. Hort. Gotting. (1834) = virosa.
amurensis, Herder, Fl. Radd. iii. 4. p. 29 = Raddeana.
amurensis, Regel & Maxim. in Ind. Sem. Hort. Petrop. (1857) 42 = brevirostris, *Champ.*
angulata, Rafin. New Fl. Am. iv. 83 (Quid?).—Am. bor.
angustana, Auct. = *augustana*.
angustifolia, Gilib. Fl. Lituan. i. 234 = saligna.
angustifolia, Lam. ex Steud. Nom. ed. I. 460 = L. Scariola.
arabica, *Jaub. & Spach, Illustr.* iii. 115.—Arab.
armena, Boiss. Fl. Orient. iii. 808 = quercina.
arvensis, Edgew. in Trans. Linn. Soc. xx. (1846) 79 = dissecta.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 265.—Alger.
augustana, All. Fl. Pedem. i. 224 = L. Scariola.
augustana, Reichb. ex Nym. Consp. 435 = sagittata.
auriculata, Aitch. in Journ. Linn. Soc. xix. (1882) 173 = L. Scariola.
auriculata, *DC. Prod.* vii. 140.—Reg. Himal.
autumnalis, Bernh. Syst. Verz. Erf. 140 = Leontodon autumnalis.
axillaris, *Rafin. New Fl. Am.* iv. 83 (Quid?).—Am. bor.
Benthamii, *C. B. Clarke, Comp. Ind.* 273.—Reg. Himal.
bialata, Griff. Notul. iv. 247 = brevirostris, *Champ.*
bracteata, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 270.—Reg. Himal.
bracteata, Wall. Cat. n. 3243 = L. Scariola.
brassicaefolia, *Boiss. Diagn.* Ser. I. xi. 49.—Persia.
brevirostra, Dulac, Fl. Hautes-Pyr. 494 = L. Plumieri.
brevirostris, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 237.—As. or.
brevirostris, *Fenzl, in Kotschy, Reise Cilic. Taur.* 384. —As. Min.
Brunoniana, C. B. Clarke, Comp. Ind. 266 = graciliflora.
canadensis, A. Gray, Man. Bot. N. U. St. 803 = hirsuta.
canadensis, *Linn. Syst.* ed. X. 1193.—Am. bor.
canescens, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 324.—Turkest.
capensis, *Thunb. Prod. Pl. Cap.* 139.—Afr. austr.

LACTUCA :—
capitata, DC. Prod. vii. 138 = sativa.
caroliniana, Walt. Fl. Carol. 193 = canadensis.
cataonica, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 815.—As. Min.
caucasica, C. Koch, in Linnaea, xvii. (1843) 275 = saligna.
cernua, Jacquem. Voy. Bot. 100 = Prenanthes violaefolia.
Chaixii, Vill. Prosp. 32 = sagittata.
chondrillaeflora, *Bor. Fl. Centr. Fr.* ed. II. 312.—Europ. austr.
chondrilloides, Steud. Nom. ed. II. ii. 3 = *prenanthoides*.
cichoriifolia, DC. Fl. Fr. Suppl. 434 = perennis.
Clarkei, *Hook. f. Fl. Brit. Ind.* iii. 406.—Reg. Himal.
coerulea, *Reichb. Fl. Germ. Excurs.* 855.—Tyrol.
coriacea, Sch. Bip. in Linnaea, xv. (1841) 725 = L. Scariola.
cracoviensis, Bueck, ex Nym. Consp. 435 = saligna.
crambifolia, *Boiss. Fl. Orient.* iii. 806.—Turkest.
crassifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 842.—Ins. Socotr.
cretica, *Desf. in Ann. Mus. Par.* xi. (1808) 160.—Graecia; As. Min.
crispa, Roth, Bot. Abh. 24 = L. Scariola.
cyanea, C. Koch, in Linnaea, xxiii. (1850) 671 = saligna.
debilis, *Benth. & Hook. f. Gen.* ii. 526.—Japon.
decipiens, *C. B. Clarke, Comp. Ind.* 266.—Reg. Himal.
decorticata, *Forsk. Fl. Aegypt. Arab.* 216.—Reg. Mediterr.
Deggleriana, *Hegetschw. Fl. Schw.* 759.—Helvet.
deltoidea, *DC. ex C. A. Mey. Verz. Pfl. Cauc.* 56.—Reg. Caucas.
denticulata, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 529.—Japon.
dissecta, *D. Don, Prod. Fl. Nep.* 164.—Reg. Himal.
Dregeana, *DC. Prod.* vii. 137.—Afr. austr.
dubia, Jord. Pugill. 119 = L. Scariola.
Dubyaea, *C. B. Clarke, Comp. Ind.* 271.—Reg. Himal.
elata, Salisb. Prod. 180 = L. Scariola.
elongata, Mulh. ex Willd. Sp. Pl. iii. 1525 = canadensis.
erysimifolia, DC. Prod. vii. 140 = muralis.
esculenta, Salisb. Prod. 180 = sativa.
Fischeriana, *DC. Prod.* vii. 135.—Sibir.
fistulosa, *Rafin. Fl. Ludov.* 58.—Am. bor.
flava, Forsk. Fl. Aegypt. Arab. 143 = Launaea nudicaulis.
flavida, Jord. Cat. Jard. Dijon (1848), 26 = virosa.
floridana, *Gaertn. Fruct.* ii. 362.—Am. bor.
formosana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 525.—China; Formosa.
fragilis, Asso, Syn. Stirp. Arag. 108 (= *Zollikoferia resedifolia*).—Hispan.
glabra, DC. in Wight, Contrib. 26 = Crepis acaulis.
glandulifera, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 208.—Afr. trop.
glareosa, *Schott & Kotschy, in Boiss. Fl. Orient.* iii. 812.—Cilicia.
glauciifolia, *Boiss. l. c.* iii. 813.—Persia.
goraeënsis, *Sch. Bip. in Flora*, xxv. (1842) 422.—Afr. trop.
graciliflora, *DC. Prod.* vii. 139.—Reg. Himal.; Malaya.
gracilis, *DC. l. c.* 140.—Reg. Himal.
graeca, *Boiss. Fl. Orient.* iii. 812.—Graecia.
graminea, Spreng. Syst. iii. 659 = sèq.
graminifolia, *Michx. Fl. Bor. Am.* ii. 85.—Am. bor.
hastata, *DC. Prod.* vii. 139.—Ind. or.
Heyneana, *DC. l. c.* 140.—Ind. or.
hirsuta, *Muhl. ex Nutt. Gen. Am.* ii. 124.—Am. bor.
hispida, *DC. Prod.* vii. 139.—As. Min.; Syria.
Hochstetteri, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 27 = capensis.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 81 = macrorhiza.
humilis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 463.—Abyssin.
indica, *Linn. Mant.* ii. 278.—Malaya.
inermis, *Forsk. Fl. Aegypt. Arab.* 144.—Arab.
integrifolia, *Bigel. Fl. Bost.* ed. II. 287.—Am. bor.
integrifolia, Nutt. Gen. Am. ii. 124 = pulchella.
intricata, *Boiss. Diagn.* Ser. I. iv. 27.—As. Min.
intybacea, *Jacq. Ic. Pl. Rar.* i. t. 162.—Nicaragua.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 384.—Ind. occ.
japonica, *Regel, Ind. Sem. Hort. Petrop.* (1866) 91.—Japon.
Kanitziana, *Martelli, in Nuov. Giorn. Bot.* xv. (1883) 302.—Borneo.
Kotschyana, Reut. Cat. Hort. Genev. (1867) 4 = scarioloides.

LACTUCA :—
Kotschyi, Sch. Bip. in Linnaea, xv. (1841) 725 = taraxacifolia.
laciniata, Roth, Catalect. i. 96 = L. Scariola.
Lactucarii, *Lamotte, in Soc. Dauph.* (1880) 271.—Gallia.
laevigata, C. B. Clarke, Comp. Ind. 269 = macrorhiza.
laevigata, *DC. Prod.* vii. 140.—Java.
latifolia, Gilib. Fl. Lituan. i. 234 = L. Scariola.
Lessertiana, *C. B. Clarke, Comp. Ind.* 270.—Reg. Himal.
leucophaea, A. Gray, in Proc. Am. Acad. xix. (1883) 73 = alpina.
leucophaea, Sibth. Fl. Graec. t. 794 = cretica.
livida, *Boiss. & Reut. in Boiss. Voy. Bot. Espagne*, ii. 742.—Hispan.
longidentata, *Moris, Pl. Sard. Elench.* 2.—Sardoa.
longifolia, *DC. Prod.* vii. 135.—Reg. Himal.
longifolia, Michx. Fl. Bor. Am. ii. 85 = canadensis.
ludoviciana, *Riddell, Syn. Fl. W. St.* 51; *DC. Prod.* vii. 141.—Am. bor. occ.
lutea, Dulac, Fl. Hautes-Pyr. 495 = chondrillaeflora.
macrantha, *C. B. Clarke, Comp. Ind.* 267.—Reg. Himal.
macrophylla, *A. Gray, Syn. Fl. N. Am.* i. II. 444.—Hab.?
macrorhiza, *Hook. f. Fl. Brit. Ind.* iii. 408.—Reg. Himal.
maculata, *Hornem. Hort. Hafn.* 755.—Hab.?
maritima, *Rafin. New Fl. Am.* iv. 83.—Am. bor.
massaviensis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 460.—Arab.; Abyssin.
mauritiana, Poir. Encyc. Suppl. iii. 292 = indica.
microcephala, *DC. Prod.* vii. 134.—As. Min.
mulgedioides, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 815.—Kurdistan.
muralis, *E. Mey. Chlor. Hannov.* 431; *Fresen. Taschenb.* (1832) 484.—Europ.; As. occ.
napifolia, DC. in Wight, Contrib. 27 = Crepis japonica.
nubica, Sch. Bip. in Linnaea, xv. (1841) 725 = goraeënsis.
nudicaulis, Murr. in Comm. Gotting. iii. (1780) 74. t. 4 = Launaea nudicaulis.
oblongifolia, Nutt. in Fras. Cat.; ex ej. Gen. Am. ii. 124 = pulchella.
obtusa, C. B. Clarke, Comp. Ind. 261 = Launaea nudicaulis.
octophylla, Sch. Bip. in Linnaea, xv. (1841) 725 = goraeënsis.
officinarum, Crantz, Inst. i. 227 = L. Scariola.
Oldhami, Maxim. in Bull. Acad. Pétersb. xix. (1874) 532 = L. Thunbergiana.
orientalis, *Boiss. Fl. Orient.* iii. 819.—Reg. Himal.; As. Min.
ovatifolia, *Rafin. New Fl. Am.* iv. 83.—Am. bor.
palmata, Willd. Phyt. 10 = L. Scariola.
palmensis, *C. Bolle, in Bonplandia*, vii. (1859) 298.—Ins. Canar.
paradoxa, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 460.—Abyssin.
pentaphylla, Sch. Bip. in Linnaea, xv. (1841) 725 = taraxacifolia.
perennis, *Linn. Sp. Pl.* 796.—Europ. austr.
persica, *Boiss. Diagn.* Ser. I. vii. 9.—Persia; Beluch.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 462 = goraeënsis.
petraea, Fisch. & Mey. in DC. Prod. vii. 134 = tuberosa.
picridiformis, *Boiss. Fl. Orient.* iii. 807.—Beluch.
pilosa, *Rafin. New Fl. Am.* iv. 83 (Quid?).—Am. bor.
plicata, Balb. Cat. Hort. Taur. (1813) 45 = L. Scariola.
Plumieri, *Gren. & Godr. Fl. Fr.* ii. 322.—Gallia.
polycephala, *Benth. & Hook. f. Gen.* ii. 526.—Ind. or.; Burma.
polyclada, *Boiss. Diagn.* Ser. I. vii. 10.—Persia.
prenanthoides, Scop. Fl. Carn. ed. II. ii. 100 = Chondrilla prenanthoides.
procera, Schrad. ex DC. Prod. vii. 137 = canadensis.
procumbens, *Roxb. Hort. Beng.* 60, nomen.—Ind. or.
Pseudo-virosa, Sch. Bip. ex Nym. Consp. 435 = L. Scariola.
pulchella, *DC. Prod.* vii. 134.—Am. bor. occ.
pygmaea, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 565.—Java.
quercina, *Linn. Sp. Pl.* 795.—Europ. austr.; Reg. Caucas.
querna, *Pall. Reise Statth. Suedl. Russl.* i. 221 (an praec. ?).—Reg. Casp.
racemosa, *Willd. Sp. Pl.* iii. 1524.—Armenia.

LACTUCA:—
Raddiana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 526.—Japon.
ramosissima, Benth. & Hook. f. Gen. ii. 526 = denticulata.
ramosissima, *Gren. & Godr. Fl. Fr.* ii. 318.—Gallia.
rapunculoides, *C. B. Clarke, Comp. Ind.* 268.—Reg. Himal.
rariflora, *Fresen. in Mus. Senckenb.* iii. (1839) 73.—Abyss.
remotiflora, *DC. in Wight, Contrib.* 26.—Ind. occ.; Arab.
repens, *Benth. & Hook. f. Gen.* ii. 526.—Japon.
rhynchocarpa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 841.—Ins. Socotr.
rigida, DC. Prod. vii. 135 = Crepis acaulis.
Roborowskii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 177.—China.
rostrata, Benth. & Hook. f. Gen. ii. 525 = graciliflora.
rosularis, *Boiss. Fl. Orient.* iii. 813.—Persia.
runcinata, DC. in Wight, Contrib. 26 = Heyneana.
sagittarioides, *C. B. Clarke, Comp. Ind.* 265.—Reg. Himal.
sagittata, Hook. f. & Thoms. ex C. B. Clarke, l. c. = sagittarioides.
sagittata, *Waldst. & Kit. Pl. Rar. Hung.* i. 1. t. 1.—Europ. austr.; As. Min.
sagittifolia, Ell. Sketch, ii. 253 = integrifolia.
salicifolia, Salisb. Prod. 180 = saligna.
saligna, *Linn. Sp. Pl.* 796.—Europ.; Afr. bor.; Oriens.
sanguinea, Bigel. Fl. Bost. ed. II. 287 = hirsuta.
sarmentosa, Wight, Contrib. 27 = Launaea pinnatifida.
sativa, Linn. Sp. Pl. ed. I. 795 = L. Scariola.
Scariola, *Linn. l. c.* ed. II. 1119.—Europ.; Oriens.
scarioloides, *Boiss. Diagn.* Ser. I. xi. 49.—Persia.
Schimperi, *Jaub. & Spach, Illustr.* iii. 114 = remotiflora.
Schimperi, Jord. Pugill. 119 = L. Scariola.
Schweinfurthii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 452.—Afr. trop.
segusiana, Balb. Elench. Fl. Taur. 94 = tenerrima.
Serriola, *Linn. Cent. Pl.* ii. 29.—Afr. trop.
seticuspis, *Boiss. Fl. Orient.* iii. 816.—Syria.
sibirica, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 528.—Sibiria.
sinuata, Forsk. Fl. Aegypt. Arab. 215 = virosa.
sonchifolia, Debeaux, Fl. Tchéf. 90 = denticulata.
sonchifolia, DC. Prod. vii. 134 = tatarica.
sonchifolia, *Panč. in Verh. Zool.-Bot. Ver. Wien*, vi. (1856) 559.—Europ.
sonchifolia, Willd. Sp. Pl. iii. 1530 = tuberosa.
sonchoides, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 816.—As. Min.
sonchoides, Lapeyr. Hist. Abr. Pl. Pyr. 461 = perennis.
soongarica, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 174.—Soongar.
sororia, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 189.—Japon; China.
spiciformis, Dulac, Fl. Hautes-Pyr. 495 = saligna.
spinosa, *Benth. & Hook. f. Gen.* ii. 525.—Oriens.
spinosa, Lam. Encyc. iii. 408 = Launaea spinosa.
squarrosa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 189 = brevirostris, *Champ.*
Stocksii, Boiss. Diagn. Ser. II. iii. 96 = auriculata.
stolonifera, *Benth. & Hook. f. Gen.* ii. 526.—Japon.
stricta, Waldst. & Kit. Pl. Rar. Hung. i. 47 = quercina.
sylvestris, Lam. Fl. Fr. ii. 84 = L. Scariola.
taraxacifolia, *Schum. & Thonn. Beskr. Guin. Pl.* 380.—Afr. trop.
tatarica, *C. A. Mey. Verz. Pfl. Cauc.* 56.—As. occ.; Ind. or.
tenerrima, *Pourr. in Mém. Acad. Toul.* iii. (1788) 321.—Europ. austr.; Marocc.
tephrocarpa, C. Koch, in Linnaea, xxiii. (1850) 672 = L. Scariola.
Thunbergii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 530.—Japon.
Tommasiniana, Sch. Bip. in Flora, xvi. (1842) 160 = saligna.
triangulata, *Maxim. Prim. Fl. Amur.* 177.—Reg. Amur.
triquetra, *Benth. & Hook. f. Gen.* ii. 525.—Syria.
tuberosa, *Jacq. Hort. Vindob.* i. 18.—As. Min. Persia.
undulata, *Ledeb. Fl. Alt.* iv. 156.—As. occ. et bor.

LACTUCA:—
versicolor, Sch. Bip. in Bonplandia, x. (1862) 331 = Fischeriana.
vialea, Bell. in Mém. Acad. Turin, x–xi. (1802–3) 450 = quercina.
villosa, Jacq. Hort. Schoenb. iii. t. 367 = acuminata.
villosa, Muhl. ex Torr. & Gray, Fl. Am. ii. 497 = canadensis.
viminea, C. B. Clarke, Comp. Ind. 262 = orientalis.
viminea, *J. & C. Presl, Fl. Čech.* 160.—Reg. Mediterr.
violaefolia, C. B. Clarke, Comp. Ind. 269 = Prenanthes violaefolia.
virgata, Tausch, in Flora, iv. (1829) 36 = intybacea.
virosa, Habl. Taur. 160 = saligna.
virosa, *Linn. Sp. Pl.* 795.—Europ.; As. bor.
virosa, Lucé, Fl. Osil. 260 = L. Scariola.
virosa, Thunb. Prod. Pl. Cap. 139 = Dregeana.
Wallrothii, Spreng. Pugill. i. 53 = saligna.
Wilhelmsiana, *Fisch. & Mey. ex DC. Prod.* vii. 134.—Armenia.

LACTUCOPSIS, Sch. Bip. ex Vis. in Mem. 1st. Venet. xv. (1870) 5 (Vis. & Panč. Pl. Serb. Rar. Dec. iii. 5) = **Lactuca**, Tourn. (Compos.).
altissima, Sch. Bip. ex Vis. l. c. 6 = L. sagittata.
aurea, Sch. Bip. ex Vis. l. c. 7 = L. sonchifolia.
brevirostris, Sch. Bip. ex Vis. l. c. 6 = L. brevirostris, *Fenzl.*
Chaixi, Sch. Bip. ex Vis. l. c. = L. sagittata.
mulgedioides, Sch. Bip. ex Vis. l. c. 7 = L. mulgedioides.
Plumierii, Sch. Bip. ex Vis. l. c. 8 = L. Plumieri.
quercina, Sch. Bip. ex Vis. l. c. 7 = L. quercina.
Wilhelmsiana, Sch. Bip. ex Vis. l. c. 6 = L. Wilhelmsiana.

LACURIS, Buch.-Ham. in Wall. Cat. n. 7119 (1849) = **Zanthoxylum**, Linn. (Rutac.).
illicioides, Buch.-Ham. l. c. = Z. Rhetza.

LADANIUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 67 = **Cistus**, Linn.
cyprium, Spach, l. c. = C. cyprius.
laurifolium, Spach, l. c. = C. laurifolius.
officinarum, Spach, l. c. = C. ladaniferus.

LADANUM, Dill. ex Linn. Syst. ed. I (1735) = **Galeopsis**, Tourn. (Labiat.).
luteum, Gilib. Fl. Lituan. i. 83 = G. ochroleuca.
purpureum, Gilib. l. c. 82 = G. Ladanum.

LADANUM, Rafin. Sylva Tellur. 131 (1838) = **Cistus**, Linn.
verum, Rafin. l. c. = C. ladaniferus.

LADENBERGIA, Klotzsch, in Hayne, Arzneigew. xiv. adnot. ad t. 15 (excl. sp.) (1846). *RUBIACEAE*, Benth. & Hook. f. ii. 34.
Joosia, Karst. Fl. Columb. i. 9. t. 5 (1858).
acutifolia, Klotzsch, in Hayne, Arzneigew. xiv. t. 15, adnot. = Cascarilla acutifolia.
Bergeniana, Klotzsch, l. c. t. 15 = Remijia Bergeniana.
cava, Klotzsch, l. c. t. 15, adnot. = Cascarilla Pavonii.
cujabensis, Klotzsch, l. c. = Remijia cujabensis.
densiflora, Klotzsch, l. c. = Remijia densiflora.
dichotoma, *Klotzsch, l. c.*—Peruv.
dissimiliflora, Klotzsch, l. c. = Exostemma dissimiliflorum.
firmula, Klotzsch, l. c. = Remijia firmula.
hexandra, Klotzsch, l. c. = Cascarilla hexandra.
Lambertiana, Klotzsch, l. c. = Cascarilla Lambertiana.
macrocarpa, Klotzsch, l. c. = Cascarilla macrocarpa.
macrocnemia, Klotzsch, l. c. = Remijia macrocnemia.
magnifolia, Klotzsch, l. c. = Cascarilla magnifolia.
Moritziana, *Klotzsch, l. c.*—N. Granat.
nitida, Klotzsch, l. c. = Cascarilla nitida.
ovalifolia, Klotzsch, l. c. = Macrocnemum Humboldtianum.
Riedeliana, Klotzsch, l. c. = Cascarilla Riedeliana.
Roraimae, Klotzsch, l. c. = Cascarilla Roraimae.
Schomburgkii, *Klotzsch, l. c.*—Guiana.
stenocarpa, Klotzsch, l. c. = Cascarilla stenocarpa.
tenuiflora, Klotzsch, l. c. = Remijia tenuiflora.
undata, Klotzsch, l. c. = Cascarilla undata.

LADOICEA, Miq. Fl. Ind. Bat. iii. 48 (1855), sphalm. = **Lodoicea**, Comm. (Palm.).

LAEA, Brongn. Enum. Gen. 88 (1843) = **Leea**, Linn. (Ampelid.).

LAECHHARDTIA, Archer, ex Gord. Pinet. Suppl. 40 (1862), sphalm. = LEICHHARDTIA, Sheph. = **Callitris**, Vent. (Conif.).
Macleayana, Archer, ex Gord. l. c. = C. Macleayana.

LAELIA, Adans. Fam. ii. 423 (1763) = **Bunias**, Tourn. (Crucif.).
cochlearioides, Pers. Syn. ii. 185 = B. cochlearioides.
iberioides, Pers. l. c. = Calepina Corvini.
orientalis, Desv. Journ. Bot. iii. (1814) 160 = B. orientalis.
podocarpa, C. A. Mey. ex Rupr. Fl. Cauc. 136, in textu = B. orientalis?
prostrata, Pers. Syn. ii. 185 = Muricaria prostrata.

LAELIA, Lindl. Gen. et Sp. Orch. 115 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 533.
Amalia, Reichb. Nom. 52 (1841).
Amalias, Hoffmgg. Preiss.-Verz. Orch. 20; ex Linnaea, xvi. (1842), Litt. 228.
acuminata, Lindl. Bot. Reg. (1841), Misc. 17 = rubescens.
albida, *Batem. ex Lindl. l. c.* (1839) t. 54. *Misc.* 2.—Mexic.
amanda ×, Reichb. f. in Gard. Chron. (1882) 776.
Amesiana ×, Reichb. f. l. c. (1884) 109.
anceps, *Lindl. Bot. Reg.* t. 1751.—Mexic.
aurantiaca, Beer, Prakt. Orch. 208, 256 = Epidendrum aurantiacum.
autumnalis, *Lindl. Gen. et Sp. Orch.* 115.—Mexic.
Barkeriana, Knowles & Westc. Flor. Cab. i. (1837) 63 = anceps.
bella ×, Reichb. f. in Gard. Chron. (1884) i. 174.
Boothiana, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 322.—Bras.
Brysiana, Lem. Illustr. Hortic. iii. (1856) 48 = elegans.
callistoglossa ×, Reichb. f. in Gard. Chron. (1882) i. 76.
caloglossa ×, Reichb. f. l. c. (1877) i. 202.
candida, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 573.—Mexic.
Canhamiana ×, Reichb. f. l. c. (1885) ii. 6.
Casperiana, Reichb. f. in C. Koch, Wochenschr. ii. (1859) 336 = purpurata.
cattleyoides, *Rodrig. Orch. Nov.* i. 67.—Bras.
caulescens, *Lindl. Bot. Reg.* (1841) sub t. 1.—Bras.
cinnabarina, *Batem. in Bot. Mag.* t. 4302.—Bras.
cinnamomea, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 180.—Bras.
Crawshayana, *Reichb. f. in Gard. Chron.* (1883) i. 142.—Hab.?
crispa, *Reichb. f. in Fl. des Serres*, Sér. I. ix. (1853–54) 102.—Bras.
crispilabia, *A. Rich. ex Warner, Sel. Orch.* Ser. II. t. 6.—Bras.
Dayana, Reichb. f. in Gard. Chron. (1876) ii. 772 = pumila.
Digbyana, *Benth. in Journ. Linn. Soc.* xviii. (1880) 314.—Honduras.
discolor, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = albida.
Dormaniana, Reichb. f. in Gard. Chron. (1878) 332, 627 = Cattleya Dormaniana.
elegans, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 242.—Bras.
euspatha, Reichb. f. in Hamb. Gartenz. xxi. (1860) 420 = *Schilleriana*.
flammea ×, Reichb. f. in Gard. Chron. (1874) i. 599.
flava, *Lindl. Bot. Reg.* (1839) *Misc.* 88.—Mexic.
fulva, Lindl. ex Heynh. Nom. ii. 29 = flava?
furfuracea, *Lindl. Bot. Reg.* (1839) t. 26.—Mexic.
Galeottiana, *Lindl. ex Heynh. Nom.* ii. 29.—Mexic.
geraënsis, *Rodrig. Orch. Nov.* i. 66.—Bras.
gigantea, Warner, Sel. Orch. Ser. I. t. 6 = elegans.
glauca, *Benth. in Journ. Linn. Soc.* xviii. (1880) 314.—Mexic.
grandiflora, *Lindl. Gen. et Sp. Orch.* 115.—Mexic.
grandis, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 60.—Bras.
harpophylla ×, Reichb. f. in Gard. Chron. (1873) 542.
irrorata, Reichb. f. in Hamb. Gartenz. xv. (1859) 57 = *Schilleriana*.
Jongheana, *Reichb. f. in Flora*, lv. (1872) 158; *et in Gard. Chron.* (1872) 425.—Bras.
Lawrenceana, Warner, Sel. Orch. Ser. II. sub t. 6 = crispilabia.
Leeana ×, Reichb. f. in Gard. Chron. (1882) i. 492.

LAELIA:—
Lindenii, Lindl. Orchid. Linden. 10 = Cattleyopsis delicatula.
Lindleyana, Nichols. Dict. Gard. ii. 229 = Cattleya Lindleyana.
longipes, *Reichb. f. Xenia Orch.* ii. 59, in obs.—Bras.
Lundii, *Reichb. f. in Otia Bot. Hamb.* 92, in obs.—Bras.
majalis, Lindl. in Bot. Reg. (1839) Misc. 35 = grandiflora.
monophylla, *N. E. Br. in Gard. Chron.* (1882) II. 782.—Jamaica.
Mylamiana ×, Reichb. f. l. c. (1876) II. 740.
pedunculatis, Lindl. in Bot. Reg. (1842) Misc. 9 = rubescens.
Perrinii, *Batem. in Paxt. Mag. Bot.* xiii. (1847) 5.—Bras.
Philbrickiana ×, Reichb. f. in Gard. Chron. (1879) II. 102.
Pilcheri ×, Dombr. Flor. Mag. t. 340 = seq.
Pilcheriana ×, Reichb. f. in Gard. Chron. (1868) 375.
praestans, Lindl. & Reichb. f. in Koch, Allg. Gartenz. (1857) 336 = pumila.
primulina, Lindl. Bot. Reg. (1840), sub t. 41, in obs. (Quid?).—Hab.?
pubescens, Lem. Jard. Fleur. ii. (1852) Misc. 79, 99 = rubescens.
pumila, *Reichb. f. in Fl. des Serres*, Sér. I. ix. (1853–54) 102.—Bras.
purpurata, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 111.—Bras.
Regnellii, *Rodrig. Orch. Nov.* ii. 154.—Bras.
Rivieri, Carr. in Rev. Hortic. (1874) 331 = Boothiana.
rubescens, *Lindl. Bot. Reg.* (1840) *Misc.* 20. t. 41.—Mexic.
rupestris, *Lindl. l. c.* xxviii. (1842) sub t. 62.—Bras.
Schilleriana ×, Reichb. f. in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 323.
Sedeni ×, Reichb. f. in Gard. Chron. (1877) II. 424.
Stetzneriana, Reichb. f. in Hamb. Gartenz. xvi. (1860) 282 = *Schilleriana*.
superbiens, *Lindl. in Bot. Reg.* (1840) *Misc.* 46.—Mexic.
Turneri, Hort. ex Warner, Sel. Orch. Ser. I. t. 12 = elegans.
Veitchiana ×, Reichb. f.—Cf. Gard. Chron. (1874) I. 566.
Veitchii, Hort. ex Nichols. Dict. Gard. ii. 229 = *flammea*.
violacea, Reichb. f. in Bonplandia, ii. (1854) 89 = rubescens?
virens, *Lindl. Bot. Reg.* (1844) *Misc.* 1.—Bras.
Wallisi, Linden, in Belg. Hortic. xv. (1865) 102 = Cattleya labiata.
Warnerii ×, Hort. ex Warner, Orch. Sel. Ser. III. t. 1 = *Schilleriana*.
Wendlandi, *Reichb. f. in Walp. Ann.* vi. 431.—Guatemala.
Wyattiana, Reichb. f. in Gard. Chron. (1883) II. 426 = purpurata.
xanthina, *Lindl. in Bot. Mag.* t. 5144.—Bras.

LAELIOPSIS, Lindl. in Paxt. Flow. Gard. iii. (1852–53) 155. t. 105 = **Epidendrum**, Linn. &c. (Orchid.).
chinensis, Lindl. l. c. 156 = E. nonchinense.
cubensis, Lindl. l. c. = E. cubense.
domingensis, Lindl. & Paxt. l. c. 155 = Broughtonia lilacina.
Lindenii, Lindl. l. c. 156 = Cattleyopsis delicatula.

LAEMELLIA, Pfeiff. Nom. ii. 10 (1871), sphalm. = **Lacmellia**, Karst. (Apocyn.).

LAENECIA, Sch. Bip. in Walp. Rep. ii. 952 (1843) = seq.

LAENNECIA, Cass. in Dict. Sc. Nat. xxv. 91 (1822) = **Conyza**, Linn. (Compos.).
alpina, Poepp. & Endl. Nov. Gen. et Sp. ii. 56. t. 262.—Chili.
filaginoides, DC. Prod. v. 376.—Mexic.
gnaphalioides, Cass. in Dict. Sc. Nat. xxv. 92 = C. gnaphalioides.
parvifolia, DC. Prod. v. 376.—Mexic.
pinnatifida, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 178.—Peruv.

LAERTIA, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 354 = **Leersia**, Sw. (Gramin.).

LAESTADIA, Kunth, in Less. Syn. Comp. 203 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 260.
LESTADIA, Spach, Hist. Vég. Phan. x. 28 (1841).
Lechleri, *Wedd. Chlor. And.* i. 184.—Peruv.
muscicola, *Wedd. l. c.* t. 32.—N. Granat.
pinifolia, *Kunth, in Less. Syn. Comp.* 203.—Am. austr.
rupestris, *Benth. Pl. Hartw.* 203.—N. Granat.

LAETIA, Loefl. It. Hisp. 190 (1758). *BIXINEAE*, Benth. & Hook. f. i. 126.
CASINGA, Griseb. in Goett. Abh. ix. (1861) 27, 29.
GUIDONIA, P. Br. Hist. Jamaic. 249 (1756).
LIGHTFOOTIA, Sw. Prod. Veg. Ind. Occ. 83 (1788).
THIODIA, Benn. Pl. Jav. Rar. 192 (1838).
acuminata, *Bonpl. ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 104.—N. Granat.
americana, Linn. Syst. ed. X. 1074 = apetala.
apetala, *Jacq. Select. Am.* 167. t. 108.—Am. austr.
calophylla, *Eichl. in Mart. Fl. Bras.* xiii. I. 454.—Am. austr.
casearioides, *Sagot, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 85.—Guiana.
completa, *Jacq. Enum. Pl. Carib.* 24.—Am. austr.
coriacea, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. I. 456.—Bras.
corymbulosa, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 83.—Bras.
crenata, *A. Rich. Ess. Fl. Cub.* 87.—Cuba.
cupulata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 84.—Bras.
floribunda, Spruce, l. c. 83 = corymbulosa.
glandulosa, Poepp. ex Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 288 = Banara mollis.
guazumaefolia, *H. B. & K. Nov. Gen. et Sp.* v. 356.—Mexic.
Guidonia, *Sw. Prod. Veg. Ind. Occ.* 83.—Jamaic.
hirtella, *H. B. & K. Nov. Gen. et Sp.* v. 355.—Mexic.
longifolia, *A. Rich. Ess. Fl. Cub.* 85.—Cuba.
lucida, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 286.—N. Granat.
parviflora, *Spreng. Syst.* ii. 609.—Bras.
procera, *Eichl. in Mart. Fl. Bras.* xiii. I. 453.—Bras.
scabra, *Spreng. Syst.* ii. 609.—Ins. S. Doming.
serrulata, *Ruiz & Pav. Ic. Ined.* x; *ex Pritz. Ic. Ind.* 615.—Hab.?
suaveolens, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 85.—Am. austr.
ternstroemioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 156.—Cuba.
Thamnia, Linn. Pl. Jam. Rar. 31; Amoen. Acad. v. 413 = apetala?
Thamnia, *Sw. Prod. Veg. Ind. Occ.* 83.—Jamaic.

LAETJI, Osb. ex Steud. Nom. ed. I. 460 (1821) = **Nephelium**, Linn. (Sapind.).
chinensis, Osb. ex Steud. l. c. = N. Litchi.

LAFOENSIA, Vand. Fl. Lusit. 33. t. 2. f. 13 (1788); et in Roem. Script. 112. t. 7. f. 13 (1796). *LYTHRARIEAE*, Benth. & Hook. f. i. 781.
CALYPLECTUS, Ruiz & Pav. Prod. 73. t. 13 (1794).
PTYCHODON, Klotzsch, ex Reichb. Nom. 172 (1841).
acuminata, *DC. in Mém. Soc. Phys. Genèv.* iii. II. (1826) 73; *Prod.* iii. 94.—Peruv.
adenophylla, Steud. Nom. ed. II. ii. 4 = replicata.
densiflora, *Pohl, Ic. Pl. Bras.* ii. 142.—Bras.
emarginata, *Koehne, in Mart. Fl. Bras.* xiii. II. 355.—Bras.
glyptocarpa, *Koehne, l. c.* 353.—Bras.
illustris, Vell. Fl. Flum. 218; v. t. 78 = Vandelliana.
kielmeyeraefolia, St. Hil. Fl. Bras. Mer. iii. 160 = Vandelliana.
lucida, Klotzsch, ex Koehne, in Mart. Fl. Bras. xiii. II. 350 = L. Pacari.
mexicana, [Moç. & Sesse,] ex DC. Prod. iii. 94 = punicaefolia.
microphylla, Pohl, Ic. Pl. Bras. ii. 145 = Vandelliana.
nummularifolia, *St. Hil. Fl. Bras. Mer.* iii. 158.—Bras.
Pacari, *St. Hil. l. c.* 159.—Bras.
petiolata, Klotzsch, ex Koehne, in Mart. Fl. Bras. xiii. II. 350 = L. Pacari.
punicaefolia, *DC. in Mém. Soc. Phys. Genèv.* iii. II. (1826) 86. t. 1.—N. Granat.
replicata, *Pohl, Ic. Pl. Bras.* ii. 145.—Bras.
scaberrima, Poepp. ex Koehne, in Engl. Bot. Jahrb. iii. (1882) 149 = Physocalymma scaberrimum.

LAFOENSIA:—
Sellowiana, Klotzsch, ex Koehne, in Engl. Bot. Jahrb. iii. (1882) 151 = L. Pacari.
sessilifolia, Klotzsch, ex Koehne, in Mart. Fl. Bras. xiii. II. 351 = L. Pacari.
speciosa, *DC. in Mém. Soc. Phys. Genèv.* iii. II. (1826) 73; *Prod.* iii. 94.—N. Granat.
Vandelliana, *Cham. & Schlecht. in Linnaea*, ii. (1827) 346; *DC. Prod.* iii. 94.—Bras.

LAFUENTEA, Lag. Gen. et Sp. Nov. 19 (1816) *SCROPHULARINEAE*, Benth. & Hook. f. ii. 962.
DURIEURA, Mérat, in Mém. Soc. Acad. Lille (1829), 432. t. 4.
rotundifolia, *Lag. Gen. et Sp. Nov.* 19.—Hispan.

LAFUENTIA, Benth. Scroph. Rev. 7 (1835) = praec.

LAGANSA, Rumpf, ex Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Cappar.).
alba, Rafin. l. c. = C. graveolens.

LAGARINTHUS, E. Mey. Comm. Pl. Afr. Austr. 202 (1837) = **Schizoglossum**, E. Mey. (Asclepiad.).
abyssinicus, Hochst. ex Benth. & Hook. f. Gen. ii. 753 = S. interruptum.
barbatus, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 257.—Afr. austr.
brevicuspis, E. Mey. Comm. Pl. Afr. Austr. 204 = Gomphocarpus brevicuspis.
corniculatus, G. Mey. l. c. 208 = Gomphocarpus corniculatus.
eustegioides, E. Mey. l. c. 207 = Gomphocarpus eustegioides.
exilis, Decne. in DC. Prod. viii. 556.—Afr. austr.
expansus, E. Mey. in Drège, Zwei Pfl. Docum. 196 = Gomphocarpus expansus.
filiformis, E. Mey. Comm. Pl. Afr. Austr. 203 = Gomphocarpus filiformis (in Addend.).
flexuosus, E. Mey. ex Decne. in DC. Prod. viii. 556 = Gomphocarpus flexuosus.
gibbus, E. Mey. Comm. Pl. Afr. Austr. 204 = Gomphocarpus gibbus.
gracilis, E. Mey. l. c. 206.—Afr. austr.
hispidus, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 257.—Afr. austr.
interruptus, E. Mey. Comm. Pl. Afr. Austr. 208 = Schizoglossum interruptum.
involucratus, E. Mey. l. c. 203 = Xysmalobium involucratum.
linearis, E. Mey. l. c. 207 = Gomphocarpus linearis.
macer, E. Mey. l. c. 206.—Afr. austr.
microdon, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 317.—Afr. austr.
multicaulis, E. Mey. Comm. Pl. Afr. Austr. 205 = Gomphocarpus multicaulis.
navicularis, E. Mey. l. c. 204 = Gomphocarpus navicularis.
peltigerus, E. Mey. l. c. 205 = Gomphocarpus peltigerus.
revolutus, E. Mey. l. c. = Gomphocarpus revolutus.
tenellus, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 256.—Afr. austr.
tenuis, E. Mey. Comm. Pl. Afr. Austr. 203.—Afr. austr.
truncatus, E. Mey. l. c. 206.—Afr. austr.
virgatus, E. Mey. l. c. 208.—Afr. austr.

LAGAROPYXIS, Miq. Ann. Mus. Lugd. Bat. i. 198 (1863–64) = **Stereospermum**, Cham. (Bignon.).
gigantea, Miq. l. c. = S. hypostictum.
glandulosa, Miq. l. c. 199 = S. glandulosum.

LAGAROSIPHON, Harv. in Hook. Journ. Bot. iv. (1842) 230. t. 22. *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 450.
NECHAMANDRA, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 78.
cordofanum, *Casp. in Pringsh. Jahrb.* i. (1858) 504.—Afr. trop.
madagascariensis, *Casp. in Bremen, Abh.* vii. (1882) 252.—Madag.
muscoides, *Harv. in Hook. Journ. Bot.* iv. (1842) 230.—Afr. austr.
Roxburghii, *Benth. in Benth. & Hook. f. Gen.* iii. 451.—Ind. or.
Schweinfurthii, *Casp. in Bot. Zeit.* xxviii. (1870) 88.—Afr. trop.
Steudneri, *Casp. in Schweinf. Beitr. Fl. Aethiop.* 200.—Afr. trop.

LAGASCEA, Cav. in Anal. Cienc. Nat. vi. (1803) 331 (Lagasca); Willd. Enum. Hort. Berol. 941 (1809). *COMPOSITAE*, Benth. & Hook. f. ii. 342.
NOCCA, Cav. Ic. iii. 12. t. 224 (1794).
NOCCAEA, Willd. Sp. Pl. iii. 2393 (1803); Jacq. Fragm. 58. t. 85 (1805).
angustifolia, *DC. Prod.* v. 92.—Mexic.
biflora, *Hemsl. Diagn. Pl. Nov.* 33.—Mexic.
campestris, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 238 = mollis.
decipiens, *Hemsl. Diagn. Pl. Nov.* 33.—Mexic.
helianthifolia, H. B. & K. Nov. Gen. et Sp. iv. 24. t. 311 = suaveolens.
heteropappus, *Hemsl. Diagn. Pl. Nov.* 33.—Mexic.
Kunthiana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 238 = mollis.
latifolia, DC. Prod. v. 92 = suaveolens.
Liebmannii, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
macrophylla, Steud. Nom. ed. II. i. 4 = suaveolens.
Moçinniana, *DC. Prod.* v. 92.—Mexic.
mollis, *Cav. in Anal. Cienc. Nat.* vi. 333 (1803) t. 44. —Cuba.
mollis, Sch. Bip. in Seem. Bot. Voy. Herald, 298 = decipiens.
rubra, *H. B. & K. Nov. Gen. et Sp.* iv. 24. t. 311.—Mexic.
suaveolens, *H. B. & K. l. c.*—Mexic.

LAGENANDRA, Dalz. in Hook. Kew Journ. iv. (1852) 289. *AROIDEAE*, Benth. & Hook. f. iii. 963.
insignis, *Trimen, in Journ. Bot.* xxiii. (1885) 269. —Zeylan.
Koenigii, *Thw. Enum. Pl. Zeyl.* 334.—Zeylan.
lancifolia, *Thw. l. c.*—Zeylan.
ovata, Thw. l. c. = toxicaria.
Thwaitesii, *Engl. in DC. Monog. Phan.* ii. 621.—Zeylan.
toxicaria, *Dalz. in Hook. Kew Journ.* iv. (1852) 289. —Ind. or.

LAGENARIA, Ser. in Mém. Soc. Phys. Genèv. iii. 1. (1825) 25. t. 2. *CUCURBITACEAE*, Benth. & Hook. f. i. 823.
angolensis, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 10 = Adenopus breviflorus.
cochinchinensis, M. Roem. Syn. Pepon. 61 = vulgaris.
dasystemon, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 81 = Benincasa cerifera.
hispida, Ser. in DC. Prod. iii. 299 = vulgaris.
idolatrica, Ser. in DC. l. c. = vulgaris.
mascarena, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 187 = Sphaerosicyos Meyeri.
microcarpa, Naud. in Rev. Hortic. (1855) 65 = vulgaris.
sagittata, Harv. in Harv. & Sond. Fl. Cap. ii. 489 = Trochomeria sagittata.
sphaerica, E. Mey. Zwei Pfl. Docum. 197 = Sphaerosicyos Meyeri.
sphaerocarpa, Arn. in Hook. Journ. Bot. iii. (1841) 277 = Sphaerosicyos Meyeri.
toxicaria, Cogn. in DC. Monog. Phan. iii. 418, in syn. = Lagenandra toxicaria.
vittata, Ser. in DC. Prod. iii. 299 = vulgaris.
vulgaris, *Ser. in Mém. Soc. Phys. Genèv.* iii. 1. (1825) 25.—Reg. trop.

LAGENIA, Fourn. in Mart. Fl. Bras. vi. iv. 293 (1885) = **Araujia**, Brot. (Asclep.).
angustifolia, Fourn. l. c. 294 = A. angustifolia.
megapotamica, Fourn. l. c. = A. megapotamica.

LAGENIAS, E. Mey. Comm. Pl. Afr. Austr. 186 (1837) = **Sebaea**, R. Br. (Gentian.).
pusillus, E. Mey. l. c. = S. pusilla.

LAGENIFERA, Cass. in Bull. Soc. Philom. (1815) 199 = **Lagenophora**, Cass. (1818) (Compos.).
magellanica, Cass. l. c. = L. Commersonii.

LAGENOCARPUS, Klotzsch, in Linnaea, xii. (1838) 215 = **Salaxis**, Salisb. (Ericac.).
imbricatus, Klotzsch, l. c.—Afr. austr.
tenuis, Benth. in DC. Prod. vii. 710.—Afr. austr.

LAGENOCARPUS, Nees, in Linnaea, ix. (1834) 304. *CYPERACEAE*, Benth. & Hook. f. iii. 1067.
ADAMOGETON, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 165 (1842).

LAGENOCARPUS:—
ANOGYNA, Nees, in Hook. Journ. Bot. ii. (1840) 395.
MICROLEPIS, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 164 (1842).
OROBIUM, Schrad. ex Nees, l. c. (1842).
PHAENOPYRUM, Schrad. ex Nees, l. c. 166 (1842).
adamantinus, *Nees, in Mart. Fl. Bras.* ii. 1. 165.—Bras.
crassipes, Boeck. in Flora, lxiii. (1880) 453 = Martii.
guianensis, *Nees, in Linnaea*, ix. (1834) 303.—Bras.; Guiana.
Martii, *Nees, in Mart. Fl. Bras.* ii. 1. 169.—Bras.
Neesii, Boeck. in Linnaea, xxxviii. (1874) 426 = rigidus.
rigidus, *Nees, in Mart. Fl. Bras.* ii. 1. 167.—Bras.
topazinus, *Nees, l. c.* 168.—Bras.
tremulus, *Nees, l. c.*—Ins. Trinit.
velutinus, *Nees, l. c.* 166.—Bras.

LAGENOPHORA, Cass. in Bull. Soc. Philom. (1818) 34. *COMPOSITAE*, Benth. & Hook. f. ii. 263.
EMPHYSOPUS, Hook. f. in Hook. Lond. Journ. vi. (1847) 113.
IXAUCHENUS, Cass. in Dict. Sc. Nat. lvi. 176 (1826).
LAGENIFERA, Cass. in Bull. Soc. Philom. (1815) 199.
MICROCALIA, A. Rich. Fl. Nouv. Zél. 230. t. 30 (1832).
SOLENOGYNE, Cass. in Dict. Sc. Nat. lvi. 174 (1826).
Billardieri, *Cass. l. c.* xxv. 111.—As. trop.; Austral.
Commersonii, *Cass. l. c.* 110.—Reg. Magell.
emphysopus, *Hook. f. Fl. Tasm.* i. 189.—Austral.
Forsteri, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 126 = seq. ?.
Forsteri, *DC. Prod.* v. 307.—N. Zel.
gracilis, Steetz, in Lehm. Pl. Preiss. i. 431 = Billardieri.
Gunniana, Steetz, l. c. = Hnegelii.
Harveyi, Thw. Enum. Pl. Zeyl. 162 = Billardieri.
hirsuta, *Poepp. ex Less. in Linnaea*, vi. (1831) 131. —Chili.
Huegelii, *Benth. Enum. Pl. Hueg.* i. 56.—Austral.
javanica, Blume, ex DC. Prod. v. 308 = Myriactis Wightii.
lanata, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 126.—N. Zel.
latifolia, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 113 = Billardieri.
Lechleri, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
maviensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 172.—Ins. Sandvic.
montana, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 113 = Billardieri.
muscicola, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
petiolata, *Hook. f. Fl. N. Zel.* 125.—N. Zel.
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 121.—Ins. Fiji.
pilosa, Blume, ex DC. Prod. v. 308 = Myriactis pilosa.
pinnatifida, *Hook. f. Fl. N. Zel.* 126.—N. Zel.
purpurascens, *Phil. in Anal. Univ. Chil.* (1873) 489. —Chili.
Solenogyne, *F. Muell. Fragm.* v. 62.—Austral.
sundana, Miq. Fl. Ind. Bat. ii. 32 = Billardieri.

LAGENULA, Lour. Fl. Cochinch. 88 (1790) = **Fusanus**, R. Br. (Santal.).
pedata, Lour. l. c. = F. pedatus.

LAGERSTROEMIA, Linn. Syst. ed. X. 1076 (1759). *LYTHRARIEAE*, Benth. & Hook. f. i. 783.
ADAMBEA, Lam. Encyc. i. 39 (1783).
FATIOA, DC. Prod. iii. 88 (1828).
MUNCHAUSIA, Linn. Mant. ii. 153 (1771).
PTEROCALYMNA, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 508.
SOTULARIA, Rafin. Sylva Tellur. 98 (1838).
TSJINKIN, Adans. Fam. ii. 401 (1763).
acuminata, Wall. ex Koehne, in Engl. Bot. Jahrb. iv. (1883) 27 = villosa.
apetala, *T. Anders. ex Teijsm. & Binn. Cat. Hort. Bog.* 241, nomen.—Burma.
Archeriana, *Bailey, Synops. Queensl. Flora*, 196.—Austral.
augusta, Wall. Cat. n. 2113 = L. Flos-reginae.
calycina, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 25. —Ins. Philipp.
calyculata, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 307.—Burma.

LAGERSTROEMIA:—
carinata, Wall. Cat. n. 1599 = floribunda.
celebica, Blume, Mus. Bot. Lugd. Bat. ii. 127 = ovalifolia.
chinensis, Lam. Encyc. iii. 375 = indica.
cuspidata, Wall. Cat. n. 2116 = floribunda.
elegans, Wall. ex Paxt. Mag. Bot. xiv. (1848) 269 = indica.
Engleriana, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 24. —Ins. Timor.
Fatioa, Blume, Mus. Bot. Lugd. Bat. ii. 125 = parviflora.
floribunda, *Jack, in Malay. Misc.* i. (1820) v. 38.—Malaya.
floribunda, Wall. Cat. n. 2115 A = turbinata.
Flos-reginae, *Retz. Obs.* v. 25.—As. trop.
grandiflora, Roxb. Hort. Beng. 38; Fl. Ind. ii. 503 = Duabanga sonneratioides.
hexaptera, *Miq. Fl. Ind. Bat.* i. 1. 623.—Malaya.
hirsuta, Rottl. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 576 = Rottleri.
hirsuta, *Willd. Sp. Pl.* ii. 1178.—Ind. or.
hypoleuca, *Kurz, Andam. Report, App.* p. viii; *et in Journ. As. Soc. Beng.* xli. (1872) II. 307.—Ins. Andaman.
indica, *Linn. Syst.* ed. X. 1076.—As. trop.
javanica, *Thunb. Fl. Javan.* 21 (Quid?).—Java.
lanceolata, *Wall. Cat.* n. 2120.—Ind. or.
lanceolata, Wight & Arn. Prod. 309 = parviflora.
Loudoni, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 425.—Siam.
macrocarpa, Wall. Cat. n. 2114 A; Voigt, Hort. Suburb. Calc. 132 = L. Flos-reginae.
madagascariensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 270.—Madag.
major, Retz. Obs. i. 20 = L. Flos-reginae.
microcarpa, Hance, in Journ. Bot. xvi. (1878) 107 = subcostata.
microcarpa, Wight, Ic. i. t. 109 = lanceolata.
minor, Retz. Obs. i. 20 = L. Flos-reginae ?
Munchausia, Willd. Sp. Pl. ii. 1178 = L. Flos-reginae.
ovalifolia, *Teijsm. & Binn. in Naturk. Tijdschr. Nederl. Ind.* ii. (1840) 306.—Malaya.
paniculata, *Vidal, Phan. Cuming. Philipp.* 39, 115.—Ins. Philipp.
parviflora, *Roxb. Pl. Corom.* i. 48. t. 66.—Ind. or.
piriformis, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 23.—Ins. Philipp.
plicifolia, Stokes, Bot. Mat. Med. iii. 105 = L. Flos-reginae.
pubescens, Wall. Cat. n. 2112 = tomentosa.
pulchra, Salisb. Prod. 365 = indica.
punctata, *Blume, Mus. Bot. Lugd. Bat.* ii. 126.—Malaya.
reginae, Roxb. Pl. Corom. i. 46. t. 65 = L. Flos-reginae.
Riedeliana, Oliver, in Journ. Linn. Soc. xv. (1876) 99 = ovalifolia.
Rottleri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 576.—Ind. or.
speciosa, Pers. Syn. ii. 72 = L. Flos-reginae.
subcostata, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 20.—China.
tomentosa, *Presl, Bot. Bemerk.* 142.—Burma.
turbinata, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 34. —Siam.
venusta, *Wall. Cat.* n. 2117.—Burma.
villosa, *Wall. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 234.—Burma.

LAGETTA, Juss. Gen. 77 (1789). *THYMELAEACEAE*, Benth. & Hook. f. iii. 195.
alpestris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 135 = Daphnopsis alpestris.
funifera, Mart. Nov. Gen. et Sp. i. 64. t. 39 = Funifera utilis.
lintearia, *Lam. Encyc.* iii. 376, 440.—Ind. occ.
malabarica, Poir. in Dict. Sc. Nat. xxv. 115 = Cansjera Rheedii.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 193.—Cuba.

LAGGERA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 26; Steud. Nom. ed. II. ii. 5 (1841); et in Walp. Rep. ii. 953 (1843). *COMPOSITAE*, Benth. & Hook. f. ii. 290.
alata, *Sch. Bip. ex Oliver, in Trans. Linn. Soc.* xxxix. (1873) 94.—As. et Afr. trop.

LAGGERA:—
arida, C. B. Clarke, Comp. Ind. 92 = Pluchea arguta.
aurita, *Sch. Bip. ex C. B. Clarke, l. c.*—As. et Afr. trop.
Braunii, *Vatke, in Linnaea*, xxxix. (1875) 486.—Afr. trop.
brevipes, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 327.—Afr. trop.
crassifolia, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. Intell. 26.—Abyss.
divaricata, Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 325 = alata.
flava, *Benth. & Hook. f. Gen.* ii. 290.—As. trop.
heteromalla, Vatke, in Linnaea, xxxix. (1875) 487 = Inula Mannii.
Musra, Benth. & Hook. f. Gen. 290 = Blumea lacera.
oblonga, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 327.—Afr. trop.
pterodonta, Sch. Bip. ex Oliver, in Trans. Linn. Soc. xxix. (1873) 94 = pupurascens.
purpurascens, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 26.—As. et Afr. trop.
senecionidea, Benth. & Hook. f. Gen. ii. 290 = Blumea membranacea.
sordida, Vatke, in Oestr. Bot. Zeitschr. (1875) 325 = Pluchea sordida.
squarrosa, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 325.—Afr. trop.
tomentosa, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. Intell. 26.—Abyssin.

LAGOCHILIUM, Nees, in Mart. Fl. Bras. ix. 85. t. 10 (1847) = **Aphelandra**, R. Br. (Acanthac.).
aurantiacum, Nees, in DC. Prod. xi. 292.—Bras.
bahiense, Nees, l. c. 291.—Bras.
decoratum, Nees, l. c.—Bras.
Hydromestus, Nees, l. c. = A. Hydromestus.
igneum, Nees, l. c. 292 = Stenandrium igneum.
luteum, Nees, in Mart. Fl. Bras. ix. 87 = A. lutea.
maculatum, Nees, in DC. Prod. xi. 290.—Peruv.
Maximilianum, Nees, in Mart. Fl. Bras ix. 85.—Bras
molle, Nees, in DC. Prod. xi. 292.—N. Granat.
montanum, Nees, in Mart. Fl. Bras. ix. 87.—Bras.
mucronatum, Nees, l. c. 86.—Bras.
obtusifolium, Nees, in DC. Prod. xi. 291.—Bras.
obtusum, Nees, in Mart. Fl. Bras. ix. 86.—Bras.
ornatum, Nees, in DC. Prod. xi. 291 = A. ornata.
pictum, Nees, l. c. 290 = A. picta?
repandum, Nees, in Mart. Fl. Bras. ix. 86.—Bras.
Schiedeanum, Nees, l. c. 87 = A. Schiedeana.
spicatum, Nees, in DC. Prod. xi. 293.—Mexic.

LAGOCHILUS, Bunge, in Benth. Lab. Gen. et Sp. 640 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1211.
YERMOLOFIA, Endl. Gen. 625 (1838).
acutilobus, *Fisch. & Mey. Enum. Pl. Schrenk*, 31.—Turcoman.
affinis, *Rupr. Sert. Tiansch.* 68.—As. centr.
alutaceus, *Bunge, Lab. Pers.* 74.—Khorassan.
Aucheri, *Boiss. Diagn.* Ser. I. v. 38.—Persia.
Bungei, *Benth. Lab. Gen. et Sp.* 641.—Soongar.
cabulicus, *Benth. in DC. Prod.* xii. 515.—Afghan.
cuneatus, *Benth. l. c.*—Afghan.
diacanthophyllus, *Benth. Lab. Gen. et Sp.* 641.—Mongol.
diacanthophyllus, Benth. l. c. = Bungei.
diacanthophyllus, Bong. et Mey. Verz. Saisang-nor. Pfl. 55. = leiacanthus.
glaberrimus, *C. Koch, in Linnaea*, xxi. (1848) 679.—Tauria.
hirsutus, Bunge, ex Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 1. 425 = hirtus.
hirtus, *Fisch. & Mey. Enum. Pl. Schrenk*, 32.—Soongar.
hispidus, *Boiss. Fl. Orient.* iv. 769.—Persia.
hispidus, *Fisch. & Mey. Enum. Pl. Schrenk*, 33.—Ind. or.
ilicifolius, *Bunge, ex Benth. Lab. Gen. et Sp.* 641.—Mongol.
inebrians, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 438.—As. centr.
insignis, Boiss. Fl. Orient. iv. 770 = macranthus.
insignis, *Fisch. & Mey. Enum. Pl. Schrenk*, 33.—Persia.
kaschgaricus, *Rupr. Sert. Tiansch.* 67.—Turkestan.
Kotschyanus, *Boiss. Diagn.* Ser. I. xii. 87.—Persia.
leiacanthus, *Fisch. & Mey. Enum. Pl. Schrenk*, 29.—Soongar.
macracanthus, *Fisch. & Mey. l. c.* 30.—Persia.

LAGOCHILUS:—
occultiflorus, *Rupr. Sert. Tiansch.* 67.—As. centr.
platyacanthus, *Rupr. l. c.* 68.—As. centr.
platycalyx, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 13.—Soongaria.
pungens, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 195.—Soongaria.

LAGOCODES, Rafin. Fl. Tellur. ii. 24 (1836) = **Scilla**, Linn. (Liliac.).
cernua, Rafin. l. c. 25 = S. sibirica.
hyacinthoides, Rafin. l. c. = S. hyacinthoides.
nutans, Rafin. l. c. = S. festalis.
patula, Rafin. l. c. = S. hispanica.

LAGOËCIA, Linn. Gen. ed. I. 61 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 880.
CUMINOIDES, Tourn. ex Moench, Meth. 93 (1794).
Cuminoides, *Linn. Sp. Pl.* 203.—Hispan.; Oriens.
Cuminoides, Soyer-Willem. Phytol. i. 254 = Carum Carvi.

LAGONYCHIUM, Bieb. Fl. Taur. Cauc. iii. 288 (1819) = **Prosopis**, Linn. (Legumin.).
Stephanianum, Bieb. l. c. = P. Stephaniana.

LAGOPHYLLA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 390. *COMPOSITAE*, Benth. & Hook. f. ii. 395.
HOLOZONIA, Greene, in Bull. Torrey Bot. Club, ix. (1882) 122.
congesta, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 86.—Am. bor. occ.
dichotoma, *Benth. Pl. Hartw.* 317.—Calif.
filipes, *A. Gray, in Pacif. Rail. Rep.* iv. 109.—Calif.
glandulosa, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 219.—Am. bor. occ.
minima, Kellogg, in Proc. Calif. Acad. v. (1873) 53 = seq.
ramosissima, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 391.—Am. bor. occ.
serrata, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 280.—Am. bor. occ.

LAGOPSIS, Bunge, ex Benth. Lab. Gen. et Sp. 586 (1834) = **Marrubium**, Tourn. (Labiat.).
flava, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 425 = M. flavum.
incana, Bunge, Verz. Suppl. Fl. Alt. 59 = M. lanatum.
viridis, Bunge, ex Benth. Lab. Gen. et Sp. 586 = M. eriostachyum.

LAGOPUS, Bernh. Syst. Verz. Erf. 229 (1800) = **Trifolium**, Linn. (Legumin.).
flexuosus, Bernh. l. c. 238 = T. medium.
glaber, Bernh. l. c.—Europ.
montanus, Bernh. l. c. 239 = T. montanum.
pratensis, Bernh. l. c. = T. pratense.

LAGOPUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 140 = **Plantago**, Tourn.
albicans, Fourr. l. c. = P. albicans.
arvensis, Fourr. l. c. = P. Lagopus.
Bellardi, Fourr. l. c. = P. Bellardi.
lanatus, Fourr. l. c. = P. Bellardi.
lanceolatus, Fourr. l. c. = P. lanceolata.
Timbali, Fourr. l. c. = P. lanceolata.

LAGOSERIS, Hoffmgg. & Link, Fl. Port. ii. 149 (1820); Link, Enum. Hort. Berol. ii. 289 (1822) = **Crepis**, Linn. (Compos.).
alata, Nym. Syll. 50 = Pterotheca nemausensis.
apargioides, Link, Handb. i. 798 = Chondrilla Peltidium.
aralensis, Boiss. Fl. Orient. iii. 882.—Turkestan.
bifida, Koch, Syn. Fl. Germ. ed. I. 435 = Pterotheca bifida.
bursifolia, Link, Enum. Hort. Berol. ii. 289 = C. bursifolia.
calycina, Hoffmgg. & Link, Fl. Port. ii. 150.—Lusit.
cernua, Link, Enum. Hort. Berol. ii. 289 = C. neglecta.
crepoides, Bieb. Suppl. ad Cent. Pl. Ross. t. 30 = Pterotheca bifida.
frigida, Boiss. Fl. Orient. iii. 882.—As. Min.
hieracioides, Boiss. & Haussk. ex Boiss. l. c. 883.—Persia.
intybacea, Hoffmgg. & Link, Fl. Port. ii. 149 = C. taraxacifolia.

LAGOSERIS:—
leontodontoides, Link, Enum. Hort. Berol. ii. 289 = Crepis leontodontoides.
nemausensis, Bieb. Fl. Taur. Cauc. iii. 538 = Pterotheca bifida.
orientalis, Boiss. Fl. Orient. iii. 882.—As. occ. et bor.
purpurea, Stev. ex DC. Prod. vii. 180; Boiss. l. c. 883 = Pterotheca bifida.
raphanifolia, Link, Enum. Hort. Berol. ii. 289 = C. taraxacifolia.
Ruppelii, Sch. Bip. in Flora, xxii. (1839) 19 = Pterotheca bifida.
taraxacifolia, Reichb. Ic. Fl. Germ. i. 28 = C. taraxacifolia.
taraxacoides, Link, Handb. i. 797 = C. taraxacoides.
taurica, Bieb. Fl. Taur. Cauc. iii. 539 = Pterotheca bifida.
taurinensis, Link, Enum. Hort. Berol. ii. 289 = C. Marschallii.
tenuifolia, Reichb. Ic. Fl. Germ. i. 11 = C. tenuifolia.
versicolor, Fisch. ex Link, Enum. Hort. Berol. ii. 289 = Lactuca Fischeriana.

LAGOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 416 = **Tetradymia**, DC. (Compos.).
ambiguus, Nutt. l. c. 417 = T. spinosa.
microphyllus, Nutt. l. c. 416 = T. spinosa.

LAGOTIA, C. Muell. in Walp. Ann. iv. 409 (1857), sphalm. = SAGOTIA, Duchass. & Walp. = **Desmodium**, Desv. (Legum.).

LAGOTIS, J. Gaertn. in Nov. Comm. Acad. Petrop. xiv. (1770) 533. t. 18. *SELAGINEAE* [Benth. & Hook. f. ii. 1129].
GERBERIA, Stell. ex Choisy, in DC. Prod. xii. 24 (1848).
GYMNANDRA, Pall. Reise, iii. 710 (1776).
brachystachya, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 525.—China.
brevituba, *Maxim. l. c.* 524.—China.
cashmeriana, Rupr. Sert. Tiansch. 64, in textu = glauca.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 559.—Reg. Himal.
decumbens, *Rupr. Sert. Tiansch.* 64.—Turkestan.
glauca, *J. Gaertn. in Nov. Comm. Petrop.* xiv. (1770) 533. t. 18. f. 2.—As. et Am. bor. et arct.
globosa, *Hook. f. Fl. Brit. Ind.* iv. 558.—Reg. Himal.
Gmelini, Rupr. Sert. Tiansch. 64 = glauca.
Korolkowi, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 522, 525.—Turkestan.
kunawurensis, Rupr. Sert. Tiansch. 64 = glauca.
Pallasii, Rupr. l. c. = glauca.
spectabilis, *Hook. f. Fl. Brit. Ind.* iv. 560.—Reg. Himal.
Stelleri, Rupr. Fl. Samojed. Cis. 49. n. 214 = glauca.
stolonifera, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 525.—As. Min.

LAGOTIS, E. Mey. in Drège, Zwei Pfl. Docum. 197 (1843) = **Carpacoce**, Sond. (Rubiac.).
spermacea, E. Mey. l. c. = C. Spermacoce.
spermacocea, E. Mey. ex Hook. Lond. Journ. Bot. ii. (1843) 541 = C. Spermacoce.

LAGOWSKIA, Trautv. in Bull. Phys. Math. Pétersb. xvi. (1858) 321 = **Coluteocarpus**, Boiss. (Crucif.).
physocarpa, Trautv. l. c. 322 = C. reticulatus.

LAGREZIA, Moq. in DC. Prod. xiii. II. 252 (1849) = **Celosia**, Linn. (Amarant.).
altissima, Moq. l. c. 253 = Deeringia altissima.
madagascariensis, Moq. l. c. 252 = C. madagascariensis.

LAGUNA, Cav. Diss. iii. 175. t. 71. f. 1 (1787) = **Hibiscus**, Linn. (Malvac.).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 71 = H. abyssinicus.
aculeata, Cav. Diss. iii. 173. t. 71. f. 1 = H. ficulneus.
angulata, Hort. ex G. Don, Gen. Syst. i. 505 = H. ficulneus.
lanceolata, Gaudich. in Freyc. Voy. Bot. 476 = H. lanceolatus.
lobata, Willd. Sp. Pl. iii. 733 = H. Solandra.
Patersonia, Sims, Bot. Mag. t. 769; Pers. Syn. ii. 259 = Lagunaria Patersonii.
sileniflora, Wall. Cat. n. 1882 = H. Solandra.
sinuata, Hornem. Hort. Hafn. ii. 645 = H. ficulneus.

LAGUNA :—
spathulata, Gaudich. in Freyc. Voy. Bot. 476 = Hibiscus spathulatus, *Gaudich.*
squamea, Vent. Jard. Malm. t. 42 = Lagunaria Patersonii.
ternata, Willd. Sp. Pl. iii. 733 = H. ternatus.
tricolor, G. Don, Gen. Syst. i. 505.—Afr. trop.

LAGUNAEA, Ag. Aph. 182 (1822) = LAGUNEA, Lour. = **Polygonum**, Linn.

LAGUNAEA, Schreb. Gen. ii. 463 (1791) = LAGUNA, Cav. = **Hibiscus**, Linn. (Malv.).

LAGUNAENA, Ritg. in Marb. Schrift. ii. (1831) 135 = LAGUNEA, Lour. = **Polygonum**, Linn.

LAGUNARIA, G. Don, Gen. Syst. i. 485 (1831). *MALVACEAE*, Benth. & Hook. f. i. 208.
cuneiformis, G. Don, l. c. = Fugosia cuneiformis.
lilacina, G. Don, in Sweet, Hort. Brit. ed. III. 77 = Fugosia hakeaefolia.
Patersonii, *G. Don, Gen. Syst.* i. 485.—Austral.

LAGUNCULARIA, Gaertn. f. Fruct. iii. 209. t. 217 (1805). *COMBRETACEAE*, Benth. & Hook. f. i. 688.
HORAM, Adans. Fam. ii. 80 (1763).
SPHENOCARPUS, Rich. Anal. Fr. 92 (1802).
coccinea, Gaudich. in Freyc. Voy. Bot. t. 104 = Lumnitzera coccinea.
glabriflora, Presl, Rel. Haenk. ii. 22 = racemosa.
Haenkei, Endl. in Ann. Wien. Mus. i. (1836) 180 = Lumnitzera pedicellata.
lutea, Gaudich. in Freyc. Voy. Bot. 481. t. 105. f. 1 = Lumnitzera lutea.
obovata, *Miq. in Linnaea*, xviii. (1844) 752.—Guian.
pedicellata, Steud. Nom. ed. II. ii. 5 = Lumnitzera pedicellata.
purpurea, Gaudich. in Freyc. Voy. Bot. 481 = Lumnitzera coccinea.
racemosa, *Gaertn. f. Fruct.* iii. 209. t. 217. f. 2.—Ind. occ.; Guian.
rosea, Gaudich. in Freyc. Voy. Bot. 481. t. 105. f. 2 = Lumnitzera rosea.

LAGUNEA, Lour. Fl. Cochinch. 220 (1790) = **Polygonum**, Linn.
cochinchinensis, Lour. l. c. = P. orientale.

LAGUNEA, Pers. Syn. ii. 259 (1807) = LAGUNA, Cav. = **Hibiscus**, Linn. (Malvac.).

LAGUNIZIA, Scop. Introd. 216 (1777) = **Homalium**, Jacq. (Samyd.).

LAGUNOA, Poir. in Dict. Sc. Nat. xxv. 128 (1822) = **Llagunoa**, Ruiz & Pav. (Sapind.).

LAGUROSTEMON, Cass. in Dict. Sc. Nat. liii. 466 (1828) = **Saussurea**, DC. (Compos.).
liatroides, Fisch. ex Wydl. in Linnaea, v. (1830) 427 = S. pycnocephala.
pygmaeus, Cass. in Dict. Sc. Nat. liii. 467 = S. pygmaea.

LAGURUS, Linn. Gen. ed. I. 353 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1154.
cylindricus, Linn. Syst. ed. X. 878 = Imperata arundinacea.
ovatus, *Linn. Sp. Pl.* 81.—Europ. austr.; Afr. bor.
Schoenanthus, Steud. Nom. ed. I. 461, 46 = Andropogon Schoenanthus.

LAGYNIAS, E. Mey. in Drège, Zwei Pfl. Docum. 197 (1843) = **Vangueria**, Juss. (Rubiac.).
discolor, E. Mey. l. c. = V. lasiantha.

LAHAYA, Roem. & Schult. Syst. v. 402 (1819) = **Polycarpaea**, Lam. (Caryophyll.).
alsinefolia, Roem. & Schult. l. c. 405 = Polycarpon alsinefolium.
aristata, Roem. & Schult. l. c. 403 = P. aristata.
corymbòsa, Roem. & Schult. l. c. 404 = P. corymbosa.
diffusa, Roem. & Schult. l. c. 402 = P. Teneriffae.
fragilis, Roem. & Schult. l. c. 406 = P. fragilis.
gnaphalioides, Roem. & Schult. l. c. 405 = P. nivea.
latifolia, Roem. & Schult. l. c. 403 = P. latifolia.
memphitica, Roem. & Schult. l. c. 406 = Polycarpon depressum.

LAHAYA :—
minuartoides, Roem. & Schult. Syst. v. 406 = Polycarpaea minuartioides.
polycarpoides, Roem. & Schult. l. c. 404 = Polycarpon peploides.
spadicea, Roem. & Schult. l. c. = P. corymbosa.
stellata, Roem. & Schult. l. c. 403 = P. stellata.
tenuifolia, Roem. & Schult. l. c. 404 = P. tenuifolia.

LAHIA, Hassk. Hort. Bogor. ed. Nov. 99 (1858). *MALVACEAE*, Benth. & Hook. f. i. 213.
kutejensis, *Hassk. l. c.* 100.—Borneo.

LAIS, Salisb. Gen. Pl. Fragm. 134 (1866) = **Hippeastrum**, Herb. (Amaryll.).
crocata, Salisb. l. c. = H. rutilum.
fulgida, Salisb. l. c. = H. rutilum.

LALAGE, Lindl. Bot. Reg. t. 1722 (1834) = **Bossiaea**, Vent. (Legumin.).
acuminata, Meissn. in Lehm. Pl. Preiss. i. 86 = B. ornata.
angustifolia, Meissn. l. c. = B. ornata.
hoveaefolia, Benth. in Lindl. Swan River App. 15 = B. ornata.
ornata, Lindl. Bot. Reg. t. 1722 = B. ornata.
stipularis, Meissn. in Lehm. Pl. Preiss. i. 87 = B. ornata.

LALLEMANDIA, Walp. Rep. iii. 986 (1845) = seq.

LALLEMANTIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 52 (1840). *LABIATAE*, Benth. & Hook. f. ii. 1200.
ZORNIA, Moench, Meth. 410 (1794).
azurea, Boiss. & Huet, ex Boiss. Fl. Orient. iv. 675 = canescens.
canescens, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 53.—As. Min.; Persia.
iberica, *Fisch. & Mey. l. c.*—As. Min.; Syria.
peltata, *Fisch. & Mey. l. c.*—Reg. Caucas.; Persia.
Royleana, *Benth. in DC. Prod.* xii. 404.—Persia; Reg. Himal.
sulphurea, C. Koch, in Linnaea, xxi. (1848) 679 = iberica.

LAMANONIA, Vell. Fl. Flum. 228 (1825); v. t. 104 (1827) = **Belangera**, Cambess. (Saxifr.).
ternata, Vell. ll. cc. = B. Lamanonia.

LAMARCHEA, Gaudich. in Freyc. Voy. Bot. 483. t. 110 (1826). *MYRTACEAE*, Benth. & Hook. f. i. 704.
hakeaefolia, *Gaudich. l. c.* 484. t. 110.—Austral.

LAMARCKEA, Steud. Nom. ed. II. ii. 6 (1841) = seq.

LAMARCKIA, DC. Fl. Fr. iii. 30 (1805) = **Lamarkia**, Moench (Gramin.).

LAMARCKIA, Hort. ex Endl. Gen. 1088, in syn. (1841) = **Elaeodendron**, Jacq. f. (Celastr.).

LAMARCKIA, Vahl, in Danske Nat. Selsk. Skrivt. vi. (1810) 93 = **Markea**, Rich. (Solanac.).

LAMARKEA, Pers. Syn. i. 218 (1805) = **Markea**, Rich. (Solanac.).
coccinea, Pers. l. c. = M. coccinea.

LAMARKEA, Reichb. Consp. 175 (1828) = **Lamarchea**, Gaudich. (Myrtac.).

LAMARKIA, G. Don, Gen. Syst. iv. 487 (1837) = **Markea**, Rich. (Solanac.).

LAMARKIA, Medic. Phil. Bot. i. 28 (1789) = **Sida**, Linn. (Malvac.).
morifolia, Medic. l. c. = S. humilis.

LAMARKIA, Moench, Meth. 201 (1794). *GRAMINEAE*, Benth. & Hook. f. iii. 1183.
CHRYSURUS, Pers. Syn. i. 80 (1805).
PTERIUM, Desv. in Journ. Bot. i. (1813) 75.
TINAEA, Garzia, in Rel. Accad. Zel. Aci Reale, ann. 3–4, 24; ex Parl. Fl. Palerm. i. 138 (1845).
aurea, *Moench, Meth.* 201.—Reg. Mediterr.; Oriens.
Hookeriana, Griff. Itin. Not. 349; Notul. iii. 94 = aurea.
tenella, DC. Cat. Hort. Monsp. 120 = Aegopogon geminiflorus.

LAMBERTIA, Sm. in Trans. Linn. Soc. iv. (1798) 214. t. 20. *PROTEACEAE*, Benth. & Hook. f. iii. 179.
Drummondii, Gardn. in Gardn. & Field. Sert. Pl. t. 22 = inermis.
echinata, *R. Br. in Trans. Linn. Soc.* x. (1810) 189.—Austral.
ericifolia, *R. Br. Prot. Nov.* 30.—Austral.
formosa, *Sm. in Trans. Linn. Soc.* iv. (1798) 223.—Austral.
ilicifolia, *Hook. Ic. Pl.* t. 553.—Austral.
inermis, *R. Br. in Trans. Linn. Soc.* x. (1810) 188.—Austral.
longifolia, *Lodd. ex Steud. Nom.* ed. II. ii. 6.—Austral.
multiflora, *Lindl. Swan River App.* 32.—Austral.
ovalifolia, *Heynh. Nom.* i. 444 (Quid?).—Austral.
propinqua, R. Br. Prot. Nov. 30 = echinata.
rariflora, *Meissn. in Lehm. Pl. Preiss.* ii. 263.—Austral.
teretifolia, Gaertn. f. Fruct. iii. 213. t. 217. f. 7 = Hakea pugioniformis.
uniflora, *R. Br. in Trans. Linn. Soc.* x. (1810) 188.—Austral.

LAMIA, Endl. Gen. 949, in syn. (1841) = LEMIA, Vand. = **Portulaca**, Linn.

LAMIELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = **Lamium**, Tourn. (Labiat.).
amplexicaulis, Fourr. l. c. = Lamium amplexicaule.

LAMIOPSIS, Opiz, Seznam, 56 (1852) = **Lamium**, Tourn. (Labiat.).
amplexicaulis, Opiz, l. c. = Lamium amplexicaule.
purpurea, Opiz, l. c. = Lamium purpureum.

LAMIUM, [Tourn.] Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1210.
ERIANTHERA, Benth. in Hook. Bot. Misc. iii. (1833) 380.
GALEOBDOLON, Adans. Fam. ii. 190 (1763).
LAMIELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134.
LAMIOPSIS, Opiz, Seznam, 56 (1852).
ORVALA, Linn. Gen. ed. I. 174 (1737).
POLLICHIA, Schrank, in Act. Erf. iii. (1781) 35.
PSILOPSIS, Neck. Elem. i. 319 (1790).
affine, *Guss. & Tenore, Fl. Nap. App.* iv. p. x.—Italia.
albiflorum, Schur, Enum. Pl. Transs. 534 = purpureum.
album, Desf. Fl. Atlant. ii. 18 = flexuosum.
album, *Linn. Sp. Pl.* 579.—Europ.; Oriens.
aleppicum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 761.—Syria.
alpestre, Trautv. in Act. Hort. Petrop. ii. (1873) 481 = tomentosum.
amplexicaule, *Linn. Sp. Pl.* 579.—Europ.; As. bor.
armenum, *Boiss. Diagn.* Ser. I. vii. 55.—Armenia.
barbatum, Sieb. & Zucc. Fl. Jap. Fam. Nat. 34 = album.
bifidum, *Cyrilli, Pl. Rar. Neap. Fasc.* i. 22. t. 7.—Europ. austr.
bithynicum, *Benth. in DC. Prod.* xii. 505.—Bithynia.
calycinum, Urv. in Mém. Soc. Linn. Par. i. (1822) 323 = moschatum.
capitatum, Sm. in Rees, Cycl. xx. n. 7 = album.
Cardiaca ×, Cogn. in Bull. Soc. Bot. Fr. xx. (1873) p. xxiv.—Gallia.
cathariaefolium, *Schott & Kotschy, in Tchihat. Asie Min. Bot.* ii. 178, nomen.—As. Min.
chinense, *Benth. in DC. Prod.* xii. 512.—China.
Coesfeldiae, Weihe, ex Reichb. Ic. Pl. Crit. iii. 21 = purpureum.
Columnae, *Tenore, Ind. Sem. Hort. Neap.* (1827) 16; *Syll.* 285.—Italia.
confertum, Fries, Summa Veg. Skand. 15, 198 = purpureum.
confusum, Martr. Fl. Tarn, 561 = purpureum.
corsicum, Gren. & Godr. Fl. Fr. ii. 679 = longiflorum.
crinitum, *Montb. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 48.—As. Min.; Persia.
cryptanthum, Guss. Fl. Sicul. Prod. ii. 97 = bifidum.
cupreum, *Schott, in Oestr. Bot. Wochenbl.* (1855) 357. —Transsylv.
cylleneum, Boiss. & Orph. Diagn. Ser. II. iv. 45 = striatum.
cymbalariaefolium, *Boiss. Fl. Orient.* iv. 759.—Lycia.
decipiens, Sond. ex Martr. Fl. Tarn, 560 = purpureum.
dilatatum, Schur, Enum. Pl. Transs. 535 = cupreum.
dissectum, With. Bot. Arr. ed. III. iii. 527 = purpureum.
Durandoi, *Pomel, Nouv. Mat. Fl. Atl.* 117.—Alger.

LAMIUM:—
Ehrenbergii, *Boiss. & Reut. ex Boiss. Fl. Orient.* iv. 761.—Syria.
elegantissimum, Schur, Enum. Pl. Transs. 535 = cupreum.
eriocephalum, *Benth. in DC. Prod.* xii. 506.—As. Min.
erythrotrichum, Boiss. Diagn. Ser. I. v. 26 = Wiedmannia orientalis.
filicaule, Boiss. l. c. xii. 86 = tomentosum.
flexuosum, *Tenore, Fl. Nap.* ii. 19. t. 52.—Reg. Mediterr.
foetidum, Gilib. Exercit. i. 97 = purpureum.
foliosum, Crantz, Stirp. Austr. ed. II. fasc. iv. 258 = album.
galactophyllum, *Boiss. & Reut. ex Boiss. Fl. Orient.* iv. 766.—Armenia.
Galeobdolon, *Crantz, Stirp. Austr.* ed. II. fasc. iv. 262. —Europ.
garganicum, *Linn. Sp. Pl.* ed. II. 808.—Italia; Graecia; China.
glechomoides, Sm. in Rees, Cycl. xx. n. 11 = striatum.
grandiflorum, Herb. Willd. ex Benth. in Linnaea, xi. (1837) 334 = maculatum.
grandiflorum, Pourr. in Mém. Acad. Toul. iii. (1788) 322 = longiflorum.
grandiflorum, Salisb. Prod. 81 = L. Orvala.
Grenieri, Mutel, Fl. Fr. iii. 25 = maculatum.
guestphalicum, Weihe, ex Nym. Consp. 575 = purpureum.
Gundelsheimeri, *C. Koch, in Linnaea*, xvii. (1843) 297. —Armenia.
heterophyllum, *Scheele, in Linnaea*, xxii. (1849) 594.—Dalmat.
hirsutum, Lam. Encyc. iii. 410 = maculatum.
hirtum, Poir. Encyc. Suppl. 296 = Stachys Macraei.
hispidulum, Michx. Fl. Bor. Am. ii. 4 = Synandra grandiflora.
humile, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 179.—Japon.
hybridum, Vill. Hist. Pl. Dauph. i. 251 = purpureum.
incisum, Willd. Sp. Pl. iii. 89 = purpureum.
inflatum, *Heuff. in Verh. Zool.-Bot. Ges. Wien*, viii. (1858) 179.—Europ. or.
intermedium, *Fries, Novit. Fl. Suec.* ed. I. 105; ed. II. 192.—Europ. bor.
Ladanum, Crantz, Stirp. Austr. ed. II. fasc. iv. 259 = Galeopsis Ladanum.
laevigatum, DC. Fl. Fr. iii. 541 = longiflorum.
laevigatum, Linn. Sp. Pl. ed. II. 808 = maculatum.
lasioclades, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 101.—Mesopot.
latifolium, *Gromow, ex Trautv. in Act. Hort. Petrop.* ix. (1884) 109 (sp. ignot.).—Rossia.
longiflorum, *Tenore, Fl. Nap. Prod.* p. xxxiv.—Europ. austr.
luteum, Krock. Fl. Siles. Suppl. ii. 148 = L. Galeobdolon.
lycium, *Boiss. Fl. Orient.* iv. 762.—Lycia.
macrodon, *Boiss. & Huet, Diagn.* Ser. II. iv. 45.—Kurdistan.
maculatum, *Linn. Sp. Pl.* ed. II. 809.—Europ.; Oriens.
Manganottianum, Clem. in Mem. Acc. Sc. Torino, Ser. II. xvi. (1857) 311 = veronicaefolium.
melissaefolium, Mill. Gard. Dict. ed. VIII. n. 5 = maculatum.
mesogaeon, Heldr. ex Boiss. Fl. Orient. iv. 760 = amplexicaule.
Michaelianum, Tausch, in Flora, xiv. (1831) 215 = garganicum.
micranthum, *Weinm. in Syll. Ratisb.* i. (1824) 68.—Bras.?
microphyllum, *Boiss. Diagn.* Ser. I. v. 25.—As. Min.
molle, *Boiss. & Orph. ex Boiss. Fl. Orient.* iv. 756.—Macedon.
molle, [Soland. in] Ait. Hort. Kew. ed. I. ii. 297 = purpureum.
moluccaefolium, Link, Handb. i. 471, sphalm. = seq.
moluccellaefolium, Fries, Nov. Fl. Suec. 72; ex Reichb. Ic. Pl. Crit. iii. 72 = intermedium.
moschatum, *Mill. Gard. Dict.* ed. VIII. n. 4.—Graecia; As. Min.
multifidum, Linn. Sp. Pl. 579 = Wiedemannia multifida.
mutabile, Dum. Fl. Belg. 45 = maculatum.
nepetaefolium, Boiss. Fl. Orient. iv. 757 = striatum.
nivale, Boiss. & Heldr. Diagn. Ser. I. vii. 54 = striatum.
niveum, Hort. ex Reichb. f. Ic. Fl. Germ. xxviii. 3 = album.

LAMIUM:—
niveum, Schrad. in Linnaea, xv. (1841) Litt. 94 = maculatum.
nudum, Crantz, Stirp. Austr. ed. II. fasc. iv. 259; Moench, Meth. 393 = purpureum.
numidicum, Noé, in Bull. Soc. Bot. Fr. ii. (1855) 584 = longiflorum.
ochroleucum, Link, in Linnaea, ix. (1834) 574 = moschatum.
ocimifolium, Sm. in Rees, Cycl. xx. n. 14 = purpureum.
orthodon, Boiss. & Huet, Diagn. Ser. II. iv. 45 = macrodon.
Orvala, *Linn. Syst.* ed. X. 1099.—Europ. austr.
pallidum, Salisb. Prod. 81 = garganicum.
palmatum, Sm. in Rees, Cyclop. xx. n. 18 = amplexicaule.
pannonicum, Scop. Fl. Carn. ed. II. i. 406 = L. Orvala.
parietariaefolium, Benth. Lab. Gen. et Sp. 739 = album.
pedemontanum, Reichb. Fl. Germ. Excurs. 322 = longiflorum.
persicum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 180 = crinitum.
petiolatum, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 381 = album.
Petitinum, J. Gay, ex Benth. Lab. Gen. et Sp. 515 = flexuosum.
phalacranthera, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = L. Orvala.
pictum, Boiss. & Heldr. Diagn. Ser. II. iv. 44 = striatum.
plebeium, Boiss. l. c. 45 = macrodon.
plebeium, Boiss. & Huet, ex Boiss. Fl. Orient. iv. 758 = striatum.
ponticum, *Boiss. & Bal. ex Boiss. l. c.* 766.—As. Min.
pubescens, *Sibth. ex Benth. Lab. Gen. et Sp.* 515.—Ital.
purpureum, *Linn. Sp. Pl.* 579.—Europ.; As. bor.
rectum, Schenk, Pl. Nov. Sp. Aegypt. 19 = striatum.
reniforme, Montbr. & Auch. in Ann. Sc. Nat. Sér. II. vi. (1836) 47 = striatum.
rhomboideum, *Benth. Lab. Gen. et Sp.* 509.—Reg. Himal.
Robertsonii, *Boiss. Diagn.* Ser. I. vii. 54.—Persia.
rubrum, Jenkinson, Brit. Pl. 128; Wallr. Sched. Crit. 300 = maculatum.
rugosum, Sibth. & Sm. Fl. Graec. vi. 45. t. 555 = pubescens.
rugosum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 296 = maculatum.
stenosiphon, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 760.—Persia.
stoloniferum, Lapeyr. Hist. Abr. Pl. Pyr. 333 = maculatum.
striatum, *Sibth. & Sm. Fl. Graec.* vi. 46. t. 557.—Graecia; As. Min.
tenuiflorum, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 34.—As. Min.
Tetrahit, Crantz, Stirp. Austr. ed. II. fasc. iv. 261 = Galeopsis Tetrahit.
tomentosum, Benth. ex Nym. Consp. 575 = L. Columnae.
tomentosum, *Willd. Sp. Pl.* iii. 90.—As. Min.; Persia.
truncatum, *Boiss. Diagn.* Ser. I. xii. 86.—Syria.
urticaefolium, Weihe, ex Reichb. Ic. Pl. Crit. iii. 21 = purpureum.
veronicaefolium, Benth. in Ann. Sc. Nat. Sér. II. vi. (1835) 47 = armenum.
veronicaefolium, *Benth. Lab. Gen. et Sp.* 510.—Bithyn.
vestitum, Benth. in DC. Prod. xii. 507 = tomentosum.
vulgatum, Benth. Lab. Gen. et Sp. 514 = album, maculatum.
westfalicum, Weihe, ex Boenn. in Flora, vi. (1823) 539 = seq.
westphalium, Bluff & Fingerh. Comp. Fl. Germ. ed. II. i. II. 353 = purpureum.

LAMOTTEA, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235 = **Munbya**, Pomel (Legumin.).
polystachya, Pomel, l. c. = M. polystachya.

LAMOTTEA, Pomel, Mat. Fl. Atl. 3 (1860) = **Carduncellus**, Adans. (Compos.).
arenosa, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235 (= *Onobroma arenosum*).—Alger.
caerulea, Pomel, Mat. Fl. Atl. 4 = C. caeruleus.
calva, Pomel, l. c. 7 (= *Onobroma calvum*).—Alger.
carlinoides, Pomel, l. c. 5 (= *Onobroma carlinoides*). —Alger.

LAMOTTEA:—
carthamoides, Pomel, Mat. Fl. Atl. 6 (= *Onobroma carthamoides*).—Alger.
centaureoides, Pomel, l. c. 5.—Alger.
Choulettiana, Pomel, l. c. 3 (= *Onobroma Choulettianum*).—Alger.
depauperata, Pomel, l. c. 6 (= *Onobroma depauperatum*).—Alger.
hispanica, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235 (= *Onobroma hispanicum*).—Alger.
incrassata, Pomel, Mat. Fl. Atl. 5.—Alger.
montana, Pomel, l. c. 4.—Alger.
multifida, Pomel, l. c.—Alger.
pectinata, Pomel, l. c. 6 = C. pectinatus.

LAMOUROUXIA, H. B. & K. Nov. Gen. et Sp. ii. 335. t. 167 ad 169 (1827). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 978.
andicola, Gill. ex Benth. in DC. Prod. x. 542, 554 = Euphrasia andicola.
bartsioides, Hook. Bot. Misc. ii. (1831) 234 = Bartsia densiflora.
betonicaefolia, Benth. in DC. Prod. x. 539 = macrantha.
breviflora, *Benth. l. c.* 540.—Peruv.
cordata, Schlecht. in Linnaea, v. (1830) 103 = viscosa.
dependens, *Benth. in DC. Prod.* x. 539.—Guatemal.
euphrasioides, Bert. ex Benth. in DC. Prod. x. 542 = Bartsia densiflora.
grandiflora, *Benth. ex Linden, Cat.* n. 10 (1855) 6.—Mexic.
Gutierrezii, *Oerst. in Kjoeb. Vidensk. Meddel.* (1 53) 29.—Venezuela.
laciniata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 32 = multifida.
lanceolata, *Benth. in DC. Prod.* x. 542.—Mexic.
linearis, Benth. l. c. 540 = tenuifolia.
longiflora, *Benth. Pl. Hartw.* 22.—Mexic.
loxensis, *Benth. l. c.* 147.—Ecuador.
macrantha, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 32.—Mexic.
microphylla, *Mart. & Gal. l. c.* 31.—Mexic.
multifida, *H. B. & K. Nov. Gen. et Sp.* ii. 339.—Mexic.
ovata, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 33.—Mexic.
parviflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 465.—Mexic.
parvifolia, Benth. in DC. Prod. x. 541 = microphylla.
rhinanthifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 337. t. 169.—Mexic.
scabra, Benth. in Seem. Bot. Voy. Herald, 177. t. 33 = Gutierrezii.
serratifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 336. t. 168. —N. Granat.
spathacea, *Benth. in DC. Prod.* x. 539.—Mexic.
subincisa, *Benth. l. c.* 540.—Peruv.
sylvatica, *H. B. & K. Nov. Gen. et Sp.* ii. 337. t. 169. —Peruv.
tenuifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 32.—Mexic.
viejensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 28. —Nicaragua.
virgata, *H. B. & K. Nov. Gen. et Sp.* ii. 336. t. 167.—Ecuador.
viscosa, *H. B. & K. l. c.* 338.—Mexic.
xalapensis, *H. B. & K. l. c.*—Mexic.

LAMPETIA, Rafin. Fl. Tellur. iii. 24 (1836) = **Mollugo**, Linn. (Ficoid.).
nudicaulis, Rafin. l. c. = M. nudicaulis.

LAMPETIA, M. Roem. Syn. Hesper. 42 (1846) = **Atalantia**, Correa (Rutac.).
racemosa, M. Roem. l. c. = A. racemosa.
sarmentosa, M. Roem. l. c. = Luvunga eleutherandra.

LAMPOCARPYA, Spreng. Anleit. ii. I. 133 (1817) = seq.

LAMPOCARYA, R. Br. Prod. 238 (1810) = **Gahnia**, Forst. (Cyperac.).
affinis, Brongn. in Duperr. Voy. Coq. Bot. 166. t. 29 = G. Gaudichaudii.
aspera, R. Br. Prod. 238 = G. aspera.
Gaudichaudii, Brongn. in Duperr. Voy. Coq. Bot. 166 = G. Gaudichaudii.
hexandra, R. Br. Prod. 238 = G. trifida.
lacera, Lesson, ex A. Rich. Voy. Astrol. 109 = G. lacera.

LAMPOCARYA :—
leptostachya, Schrad. ex Boeck. in Linnaea, xxxviii. (1874) 354 = Gahnia leptostachya.
rawacensis, Kunth, Enum. Pl. ii. 333 = G. aspera.
schoenoides, R. Br. Prod. 238 = G. schoenoides.
setifolia, Lesson & Rich. ex Hook. f. Handb. N. Zeal. Fl. 306 = G. setifolia.
tenax, Hook. f. Fl. N. Zel. i. 277 = Cladium Gunnii.
xanthocarpa, Hook. f. l. c. 278 = G. xanthocarpa.

LAMPRA, Benth. Pl. Hartw. 95 (1842) = **Weldenia**, Schult. (Commelin.).
volcanica, Benth. l. c. = W. candida.

LAMPRA, Lindl. ex DC. Prod. iv. 72 (1830) = **Trachymene**, Rudge (Umbellif.).

LAMPRACHAENIUM, Benth. in Benth. & Hook. f. Gen. ii. 225 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
microcephalum, *Benth. l. c.* 226.—Ind. or.

LAMPROCAPNOS, Endl. Gen. Suppl. v. 32 (1850) = **Dicentra**, Bernh. (Papav.).

LAMPROCARPUS, Blume, in Schult. f. Syst. vii. 1615, 1726 (1830) = **Pollia**, Thunb. (Commelin.).
thyrsiflorus, Blume, l. c. 1726 = P. thyrsiflora.

LAMPROCARYA, Nees, in Linnaea, ix. (1834) 301 = LAMPOCARYA, R. Br. = **Gahnia**, Forst. (Cyperac.).

LAMPROCAULOS, Mast. in DC. Monog. Phan. i. 349. t. 3 (1878). *RESTIACEAE*, [Benth. & Hook. f. iii. 1032].
grandis, *Mast. l. c.*—Afr. austr.
Neesii, *Mast. l. c.* 350.—Afr. austr.

LAMPROCHLAENA, F. Muell. Fragm. iii. 157 (1863) = **Myriocephalus**, Benth. (Compos.).
Oldfieldii, F. Muell. l. c. = M. nudus.

LAMPROCOCCUS, Beer, Bromel. 103 (1857) = **Aechmea**, Ruiz & Pav. (Bromel.).
caerulescens, Regel, Gartenfl. (1871) 225. t. 694 = A. caerulescens.
chlorocarpus, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 382 = A. parviflora.
corallinus, Beer, Bromel. 106 = A. corallina.
farinosus, Regel, Ind. Sem. Hort. Petrop. (1868) 79 ; et in Ann. Sc. Nat. Sér. V. x. (1869) 381.—Bras. ?
fulgens, Beer, Bromel. 103 = A. fulgens.
glomeratus, Beer, l. c. 105 = A. glomerata ?
Laurentianus, C. Koch, Wochenschr. iii. (1860) 73 = A. Weilbachii.
miniatus, Beer, Bromel. 104 = A. fulgens.
ramosus, Beer, l. c. 106.—Ind. occ.
Vallerandi, Carr. in Rev. Hortic. (1877) 129 = Streptocalyx Vallerandi.
Weilbachii, E. Morr. in Belg. Hortic. (1861) 305 = A. Weilbachii.

LAMPROCONUS, Lem. Jard. Fleur. ii. (1852) sub t. 127 = **Pitcairnia**, L'Hérit. (Bromel.).
Altensteinii, Lem. l. c. = P. Altensteinii.
giganteus, Lem. l. c. = P. Altensteinii.
Jacksoni, Hemsl. Biol. Centr. Am. Bot. iii. 316 = P. Jacksoni.
undulatus, Lem. Jard. Fleur. i (1852) sub t. 127 = P. undulata.
Warscewiczii, Lem. Illustr. Hortic. viii. (1861) Misc. 14 = P. atrorubens.

LAMPRODITHYROS, Hassk. in Flora, xlvi. (1863) 388 = **Aneilema**, R. Br. (Commelin.).
acuminatus, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 223 = A. acuminatum.
adhaerens, Hassk. in Schweinf. Beitr. Fl. Aethiop. 211 = A. aequinoctiale.
aequinoctialis, Hassk. l. c. = A. aequinoctiale.
beniniensis, Hassk. in Peters, Reise Mossamb. Bot. 529 = A. beniniense.
biflorus, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 224 = A. biflorum.
Dregeanus, Hassk. in Peters, Reise Mossamb. Bot. 529 = A. Dregeanum.

LAMPRODITHYROS :—
Ehrenbergii, Hassk. in Schweinf. Beitr. Fl. Aethiop. 210 = Aneilema Ehrenbergii.
gracilis, Kotschy & Peyr. Pl. Tinn. 47. t. 23 = A. gracile.
lanceolatus, Hassk. in Schweinf. Beitr. Fl. Aethiop. 211 = A. lanceolatum.
paniculatus, Hassk. in Flora, xlvi. (1863) 389 = Floscopa scandens.
Petersii, Hassk. l. c. = A. Petersii.
protensus, Hassk. in Peters, Reise Mossamb. Bot. ii. 529 = A. protensum.
rivularis, Hassk. in Schweinf. Beitr. Fl. Aethiop. 211 = Floscopa rivularis.
Russegeri, Fenzl, in Sitzb. Acad. Wien, Math. Nat. t. i. Abth. II. (1865) 138 = A. lanceolatum.
Schomburgkianus, Hassk. in Flora, xlvi. (1863) 389 = A. ovato-oblongum.
stenorhachys, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 226 = A. ovato-oblongum.
tacazzeanus, Hassk. in Peters, Reise Mossamb. Bot. 531 = A. aequinoctiale.

LAMPROLOBIUM, Benth. Fl. Austral. ii. 202 (1864). *LEGUMINOSAE*, Benth. & Hook. f. i. 495.
fruticosum, *Benth. l. c.*—Austral.
megalophyllum, *F. Muell. Fragm.* ix. 67.—Austral.

LAMPROPHYLLUM, Miers, in Proc. Linn. Soc. ii. (1854) 338 ; et in Trans. Linn. Soc. xxi. (1855) 249. t. 26 = **Rheedia**, Linn. (Guttif.).
Gardnerianum, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xiv. (1860) 322 = R. Gardneriana.
laetum, Miers, in Proc. Linn. Soc. ii. (1854) 338 ; et in Trans. Linn. Soc. xxi. (1855) 249 (sp. dub.).—Am. trop.

LAMPROSPERMUM, Klotzsch, in Schomb. Fauna et Fl. Brit. Gui. 1181 (1848), nomen. *SAPINDACEAE*.
guianense, *Klotzsch, l. c.*—Guiana.
Schomburgkii, *Klotzsch, l. c.*—Guiana.

LAMPROSTACHYS, Boj. ex Benth. in DC. Prod. xii. 457 (1848) = **Achyrospermum**, Blume (Labiat.).
rosea, Boj. ex Benth. l. c. 458 = A. fruticosum.

LAMPROTHAMNUS, Hiern, in Oliver, Fl. Trop. Afr. iii. 130 (1877). *RUBIACEAE*.
zanguebaricus, *Hiern, l. c.*—Afr. trop.

LAMPROTIS, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = **Erica**, Tourn.
aperta, G. Don, Gen. Syst. iii. 812 = E. borboniaefolia.
Aphanes, G. Don, l. c. = E. Aphanes.
Bedfordiana, G. Don, l. c. 811 = E. leucantha.
biflora, G. Don, l. c. 812 = E. dianthifolia.
borboniaefolia, G. Don, l. c. = E. borboniaefolia.
brevifolia, G. Don, l. c. 811 = E. brevifolia.
calycina, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = E. bracteata.
calycinoides, G. Don, Gen. Syst. iii. 811 = E. glomerata.
chlamydiflora, G. Don, l. c. = E. chlamydiflora.
corifolia, G. Don, l. c. 810 = E. corifolia.
dianthifolia, G. Don, l. c. 812 = E. dianthifolia.
fabrilis, G. Don, l. c. 811 = E. fabrilis.
flagellaris, G. Don, l. c. = E. teretiuscula.
flagelliformis, G. Don, l. c. = E. teretiuscula.
gnaphalodes, G. Don, l. c. = E. gnaphalodes.
holosericea, G. Don, l. c. = E. andromedaeflora.
hyssopifolia, G. Don, l. c. = E. bracteata.
Loddigesii, G. Don, l. c. = E. spumosa.
lucida, G. Don, l. c. = E. lucida.
lutea, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = E. lutea.
modesta, G. Don, Gen. Syst. iii. 811 = E. modesta.
mucronata, Steud. Nom. ed. II. ii. 7 = E. mucronata.
munda, G. Don, Gen. Syst. iii. 811 = E. nigrita.
oppositifolia, G. Don, l. c. 812 = E. lutea.
pallida, G. Don, l. c. 810 = E. palliiflora.
paniculata, G. Don, l. c. 811 = E. paniculata.
phylicoides, G. Don, l. c. = E. pulchella.
rubella, G. Don, l. c. 812 = E. glomerata.
rubida, G. Don, l. c. 811 = E. glomerata.
selaginifolia, G. Don, l. c. = E. selaginifolia.
spumosa, G. Don, l. c. 810 = E. spumosa.
squamosa, G. Don, l. c. 812 = E. squamosa.

LAMPROTIS :—
taxifolia, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = Erica taxifolia.
tenuifolia, D. Don, l. c. = E. tennifolia.
togata, G. Don, Gen. Syst. iii. 812 = E. borboniaefolia.
turgida, G. Don, l. c. 810 = E. taxifolia.

LAMPSANA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 209 (1745) = **Lapsana**, Linn. (Compos.).

LAMPUJANG, Koen. in Retz. Obs. iii. 62 (1783) = **Zingiber**, Adans. (Scitam.).
majus, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 392.—Ind. or.

LAMYRA, Cass. in Dict. Sc. Nat. xxv. 218 (1822) = **Cnicus**, Linn. (Compos.).
alata, Cass. l. c. 222 ; et l. 470 = C. strictus.
angustifolia, Cass. l. c. xxv. 221 = C. echinocephalus.
diacantha, Cass. l. c. = C. diacanthus.
glabella, Cass. l. c. 222 ; et l. 470 = C. strictus.
hispanica, Webb, Iter Hisp. 34 = C. hispanicus.
lobata, Cass. in Dict. Sc. Nat. xxv. 221 = C. afer.
pinnatifida, Cass. l. c. = C. horridus.
stipulacea, Cass. l. c. 222 = C. stellatus.
triacantha, Cass. l. c. 220 = C. Casabonae.
undulata, Cass. l. c. = C. hispanicus.

LANARIA, Adans. Fam. ii. 255 (1763) = **Gypsophila**, Linn. (Caryophyll.).

LANARIA, Ait. Hort. Kew. ed. I. i. 462 (1789). *HAEMODORACEAE*, Benth. & Hook. f. iii. 675.
ARGOLASIA, Juss. Gen. 60 (1789).
AUGEA, Retz. Obs. v. 3 (1789).
plumosa, *Ait. Hort. Kew.* ed. I. i. 462.—Afr. austr.

LANCEA, Hook. f. & Thoms. in Hook. Kew Journ. ix. (1857) 244. t. 7. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 948.
tibetica, *Hook. f. & Thoms. l. c.*—Reg. Himal.

LANCISIA, Lam. Illustr. Gen. t. 701. ff. 2, 3 (1798 ?) = **Lidbeckia**, Berg (Compos.).
anthemoides, Moench, Meth. Suppl. 232 = Cotula anthemoides.
bipinnata, Pers. Syn. ii. 463 = Lasiospermum radiatum.
lobata, Pers. l. c. = L. lobata.
pectinata, Lam. ex Pers. l. c. = L. pectinata.

LANCISIA, Ponted. ex Adans. Fam. ii. 126 (1763) ; Gaertn. Fruct. ii. 422 (1791) = **Cenia**, Juss. (Compos.).
turbinata, Gaertn. l. c. = C. turbinata.

LANCRETIA, Delile, Fl. Egypte, 213. t. 25 (1813) = **Bergia**, Linn. (Elatin.)
suffruticosa, Delile, l. c. = B. odorata.

LANDERSIA, Macfad. Fl. Jamaic. ii. 142 (1837) = **Melothria**, Linn. (Cucurb.).
pervaga, Macfad. l. c. = M. pervaga.

LANDIA, Comm. ex Juss. Gen. 201, in obs. (1789) = **Mussaenda**, Burm. (Rubiac.).
astrographa, Comm. ex DC. Prod. iv. 372 = M. arcuata.
lappacea, Domb. ex Vitm. Summa, iv. 340 = M. Landia.
stelligera, Comm. ex DC. Prod. iv. 372 = M. arcuata.

LANDOLFIA, D. Dietr. Syn. Pl. i. 561, 625 (1839) = seq.

LANDOLPHIA, Beauv. Fl. Owar. i. 54. t. 34 (1806). *APOCYNACEAE*, Benth. & Hook. f. ii. 692.
ALSTONIA, Scop. Introd. 198 (1777).
FATERNA, Noronha, ex A. DC. in DC. Prod. viii. 327 (1844).
PACOURIA, Aubl. Pl. Gui. i. 268. t. 105 (1775).
VAHEA, Lam. Illustr. Gen. t. 169 (1798 ?)
WILLUGHBEIA, Klotzsch, in Peters, Reise Mossamb. Bot. 281 (1861).
capensis, *Oliver, in Hook. Ic. Pl.* t. 1228.—Afr. austr.
comorensis, *Benth. & Hook. f. Gen.* ii. 693.—Ins. Comor.
florida, *Benth. in Hook. Niger Fl.* 444.—Afr. trop.

LANDOLPHIA:—
Heudelotii, *A. DC. in DC. Prod.* viii. 320.—Afr. trop.
Kirkii, [*Dyer, in*] *Kew Gard. Rep.* 1880 (1881) 39.—Afr. trop.
madagascariensis, *Benth. & Hook. f. Gen.* ii. 693.—Madag.
Mannii, [*Dyer, in*] *Kew Gard. Rep.* 1880 (1881) 39.—Afr. trop.
Michelini, *Benth. in Hook. Niger Fl.* 444.—Afr. trop.
owariensis, *Beauv. Fl. Owar.* i. 54. t. 34.—Afr. trop.
Petersiana, [*Dyer, in*] *Kew Gard. Rep.* 1880 (1881) 42.—Afr. trop.
scandens, *Didr. in Kjoeb. Vidensk. Meddel.* (1854) 190.—Afr. trop.

LANDTIA, Less. Syn. Comp. 37 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 457.
SCHNITTSPAHNIA, Sch. Bip. in Flora, xxv. (1842) 436.
UBIAEA, J. Gay, in A. Rich. Tent. Fl. Abyss. i. 447 (1847).
hirsuta, *Less. Syn. Comp.* 37.—Afr. austr.
media, DC. Prod. vi. 495 = hirsuta, nervosa.
nervosa, *Less. Syn. Comp.* 38.—Afr. austr.
Rueppellii, *Benth. & Hook. f. Gen.* ii. 457 (Ruppellii); *et ex Vatke, in Linnaea*, xxxix. (1875) 509.—Afr. trop.
Schimperi, *Benth. & Hook. f. l. c.*; *et ex Vatke, l. c.*—Abyssin.

LANEASAGUM, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 71 = **Cyclostemon**, Blume (Euphorb.).
oblongifolium, Bedd. l. c. = C. macrophyllus.

LANESSANIA, Baill. Adansonia, xi. (1875) 298. *URTICACEAE*, Benth. & Hook. f. iii. 370.
turbinata, *Baill. l. c.*—Bras.

LANGEFELDIA, Steud. Nom. ed. II. ii. 7 (1841) = **Langeveldia**, Gaudich. (Urtic.).

LANGERSTROEMIA, Cram. Disp. 126 (1803), sphalm. = **Lagerstroemia**, Linn. (Lythrar.).

LANGEVELDIA, Gaudich. in Freyc. Voy. Bot. 494 (1826) = **Elatostema**, Forst. (Urtic.).
acuminata, Gaudich. l. c. = E. acuminatum.

LANGIA, Endl. Gen. 304 (1837) = **Hermbstaedtia**, Reichb. (Amaranth.).
glauca, Endl. ex Moq. in DC. Prod. xiii. II. 247 = H. glauca.

LANGLEIA, Scop. Introd. 231 (1777) = **Casearia**, Jacq. (Samyd.).

LANGSDORFFIA, Fisch. ex Regel, Ind. Sem. Hort. Petrop. (1863) 26 = **Chloris**, Sw. (Gramin.).
repens, Fisch. l. c. = C. distichophylla.

LANGSDORFFIA, Mart. in Eschweg. Journ. Bras. ii. (1818) 178. *BALANOPHOREAE*, Benth. & Hook. f. iii. 236.
SENFTENBERGIA, Klotzsch & Karst. ex Eichl. in DC. Prod. xvii. 140 (1873).
hypogaea, *Mart. in Eschweg. Journ. Bras.* ii. (1818) 178. t. 5.—Mexic.
indica, Wight & Arn. in Hook. Ic. Pl. tt. 205, 206 = Balanophora indica.
janeirensis, Rich. in Mém. Mus. Par. viii. (1822) 412. t. 19 = hypogaea.
Moritziana, *Klotzsch & Karst. in Linnaea*, xx. (1847) 461.—N. Granat.
rubiginosa, Wedd. in Ann. Sc. Nat. Sér. III. xiv. (1850) 187 = hypogaea.

LANGSDORFFIA, Raddi, in Mem. Soc. Ital. Mod. xviii. (1820) 347. t. 1 = **Cocos**, Linn. (Palm.).
Pseudo-cocos, Raddi, l. c. = C. Mikaniana.

LANGSDORFFIA, Steud. Nom. ed. II. ii. 7 (1841) = LANGSDORFIA, Leandro = **Zanthoxylum**, Linn. (Rutac.).

LANGSDORFFIA, Willd. ex Steud. Nom. ed. II. ii. 7 (1841) = **Lycoseris**, Cass. (Compos.).

LANGSDORFIA, Leandro, in Denkschr. Akad. Muench. vii. (1819) 229 = **Zanthoxylum**, Linn. (Rutac.).
instrumentaria, Leandro, l. c. = Z. monogynum.

LANGSDORFIA, 'Raddi', ex Pfeiff. Nom. ii. 23 (1874) = LANGSDORFFIA, = **Cocos**, Linn. (Palm.).

LANGSDORFIA, Rafin. Fl. Tellur. iii. 74 (1836) = **Nicotiana**, Tourn. (Solanac.).
viridiflora, Rafin. l. c. = N. Langsdorffii.

LANGSDORFIA, Willd. ex Less. Syn. Comp. 97 (1832) = **Lycoseris**, Cass. (Compos.).
melastomoides, Willd. ex Less. l. c. = L. latifolia.

LANGUAS, Koen. in Retz. Obs. iii. 64 (1783) = **Alpinia**, Linn. (Scitam.).
aquaticum, Koen. l. c. 67 = A. aquatica.
chinensis, Koen. l. c. 65 = A. chinensis.
vulgare, Koen. l. c. 64 = A. alba.

LANIGEROSTEMMA, Chapel. ex Endl. Gen. 1035 (1840) = **Eliaea**, Cambess. (Hypericin.).

LANIPILA, Burch. Trav. i. 259 (1824) = **Lasiospermum**, Lag. (Compos.).
capensis, Burch. ex Steud. Nom. ed. II. ii. 7 = Lasiospermum radiatum.

LANIUM, Lindl. ex Benth. in Hook. Ic. Pl. t. 1334 (1881). *ORCHIDEAE*, Benth. & Hook. f. iii. 522.
Avicula, *Lindl. ex Benth. l. c.* t. 1335.—Bras.
microphyllum, *Lindl. ex Benth. l. c.* t. 1334.—Guiana.

LANKESTERIA, Lindl. Bot. Reg. (1845) Misc. 86. *ACANTHACEAE*, Benth. & Hook. f. ii. 1083.
Barteri, *Hook. f. Bot. Mag.* t. 5533.—Afr. trop.
elegans, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 33.—Afr. trop.
hispida, T. Anders. l. c. 32 = parviflora.
longiflora, Lindl. Bot. Reg. (1846) t. 12 = parviflora.
parviflora, *Lindl. l. c.* (1845) *Misc.* 86.—Afr. trop.

LANNARIA, Heist. Syst. 7 (1748) (gen. dub.).

LANNEA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 153. t. 42 (1832) = **Odina**, Roxb. (Anacard.).
acida, A. Rich. l. c. 154 = O. acida.
velutina, A. Rich. l. c. = O. velutina.

LANNEOMA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 91. t. 1. f. 2. *ANACARDIACEAE*, Benth. & Hook. f. i. 428.
velutina, *Delile, l. c.*—Abyssin.

LANSBERGIA, De Vriese, Epim. Ind. Sem. Hort. Lugd. Bat. (1846) 2 = **Trimezia**, Salisb. (Irid.).
caracasana, De Vriese, l. c.—Cf. Linnaea, xxiv. (1851) 166 = T. lurida.
cathartica, Klatt, in Linnaea, xxxi. (1861–62) 549 = T. juncifolia.
juncifolia, Klatt, l. c. = T. juncifolia.
martinicensis, Baker, in Journ. Linn. Soc. xvi. (1877) 127 = T. lurida.
meridensis, Klatt, in Abh. Naturf. Ges. Halle, xiii. (1882) 364 = T. lurida.
monticola, Klatt, in Mart. Fl. Bras. iii. I. 526 = T. juncifolia.
purgans, Klatt, l. c. 527 = T. juncifolia.
setacea, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 364.—Bras.?
spathata, Klatt, in Mart. Fl. Bras. iii. I. 528.—Bras.
violacea, Klatt, l. c. 526; Warm. in Kjoeb. Vidensk. Meddel. (1872) 129 = T. juncifolia.

LANSIUM, Rumph. Amb. i. 151. t. 54 (1741); Jack, in Trans. Linn. Soc. xiv. I. (1823) 115. *MELIACEAE*, Benth. & Hook. f. i. 334.
LACHANODENDRON, Reinw. in Blume, Cat. Gew. Buitenz. 70 (1823), nomen.
anamalaiense, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 40.—Ind. or.
anamalayanum, Bedd. in Trans. Linn. Soc. xxv. (1865) 212 = praec.
aqueum, *Jack, l. c.* xiv. I. (1823) 116.—Malaya.
cinereum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 558.—Malacc.

LANSIUM:—
domesticum, *Jack, in Trans. Linn. Soc.* xiv. I. (1823) 115.—Malaya.
humile, *Hassk. Hort. Bog. Desc.* i. 121.—Sumatra.
javanicum, M. Roem. Syn. Hesper. 99 = domesticum.
montanum, Steud. Nom. ed. II. ii. 8 (= *Milnea montana*).—Sumatra.
pedicellatum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 558.—Malacc.
silvestre, M. Roem. Syn. Hesperid. 99 = Clausena Wampi.

LANTANA, Linn. Gen. ed. I. 185 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1142.
RIEDELIA, Cham. in Linnaea, vii. (1832) 240.
TAMONOPSIS, Griseb. [Pl. Lorentz.] in Abh. Goett. (1874) 198.
abyssinica, Otto & Dietr. Allg. Gartenz. ix. (1841) 378 = Lippia adoënsis.
achyrantifolia, Desf. Cat. Hort. Par. ed. III. 392 = trifolia.
aculeata, Linn. Sp. Pl. 627 = L. Camara.
africana, Linn. l. c. 628 = Oftia africana.
alba, *Mill. Gard. Dict.* ed. VIII. n. 8.—Am. austr.
alba, Schau. in DC. Prod. xi. 606 = indica.
albopurpurea, Desf. Cat. Hort. Par. ed. III. 392 = trifolia.
amethystina, Otto & Dietr. Allg. Gartenz. ix. (1841) 370 = nivea.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 10 = stricta.
annua, Hort. Calc. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 563 = crenulata.
annua, Linn. Sp. Pl. 627 = trifolia.
antidotalis, *Schum. & Thonn. Beskr. Guin. Pl.* 276.—Afr. trop.
antillana, Rafin. Sylva Tellur. 82 = L. Camara.
armata, *Schau. in DC. Prod.* xi. 597.—Venezuela.
asperata, Hort. Par. ex Vis. Hort. Patav. (1842) 142 = nivea.
bahiensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 206.—Bras.
Bartramii, *Baldw. Reliq. Baldwin.* 247 (Quid?).—Am. bor.
bracteosa, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1845) 326 = Lippia callicarpaefolia.
brasiliensis, *Link, Enum. Hort. Berol.* ii. 126.—Bras.
bullata, Linn. Sp. Pl. 627 = Cordia curassavica.
Camara, *Linn. l. c.*—Am. trop.
canescens, Hort. Calc. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 564 = Lippia geminata.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 259.—Am. trop.
capensis, Thunb. Prod. Pl. Cap. 98 = Oftia africana.
caracasana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 206.—Venezuela.
celtidifolia, H. B. & K. Nov. Gen. et Sp. ii. 259 = trifolia.
chamaedrifolia, *Cham. in Linnaea*, vii. (1832) 127.—Bras.
Chamissonis, *Benth. & Hook. f. Gen.* ii. 1142.—Bras.
cinerea, Lam. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 378 = brasiliensis.
Clarazii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 229.—Patagon.
coccinea, *Lodd. ex G. Don, in Loud. Hort. Brit.* 245, nomen.—Am. austr.
collina, Decne. in Jacquem. Voy. Bot. 136. t. 141 = indica.
corymbosa, Linn. Sp. Pl. 628 = Cordia ulmifolia.
crenulata, *Otto & Dietr. Allg. Gartenz.* ix. (1841) 363.—Ind. or.?
crispa, Thunb. Prod. Pl. Cap. 98 = Oftia africana.
crocea, *Jacq. Hort. Schoenb.* iv. t. 473.—Ind. occ.
cujabensis, *Schau. in DC. Prod.* xi. 599.—Bras.
cuneatifolia, Klotzsch, ex Walp. Rep. iv. 45 = Lippia citrata.
decumbens, Sellow, ex A. DC. in DC. Prod. xi. 604 = Sellowiana.
delicatissima, *Poit. in Rev. Hortic.* Sér. IV. i. (1852) 461.—Hab.?
dubia, Royle, Illustr. Bot. Himal. 300. t. 73 = alba.
dubia, Wall. ex Walp. Rep. iv. 63 = trifolia, indica.
filiformis, Domb. ex Schau. in DC. Prod. xi. 602 = scabiosaeflora.
flava, *Medic. in Act. Acad. Theod. Palat.* iii. *Phys.* (1775) 225.—Hab.?
floridana, *Rafin. Atl. Journ.* 148 (Quid?).—Am. bor.

LANTANA:—
formosa, Hort. ex C. Koch, Wochenschr. i. (1858) 322 = crocea.
fucata, Lindl. Bot. Reg. t. 798 = lilacina.
fuscata, C. Koch, Wochenschr. i. (1858) 322 = praec.
geminata, Spreng. Syst. ii. 763 = Lippia geminata.
Geroldiana, Otto & Dietr. Allg. Gartenz. ix. (1841) 372 = hispida.
glutinosa, Poepp. in Otto & Dietr. l. c. x. (1842) 315 = tiliaefolia.
Gogchana, Buch.-Ham. ex Voigt, Hort. Suburb. Calc. 472 = alba.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 326.—Mexic.
hirta, R. Grah. in Edinb. N. Phil. Journ. (Oct.-Dec. 1826) 186 = hispida.
hispida, *H. B. & K. Nov. Gen. et Sp.* ii. 260.—Mexic.
horrida, *H. B. & K. l. c.* 261.—Mexic.
incana, Otto & Dietr. Allg. Gartenz. ix. (1841) 371 = involucrata.
incarnata, *Rafin. Sylva Tellur.* 83.—Hab.?
inconspicua, Tausch, in Flora, xix. (1836) 391 = lilacina.
indica, *Roxb. Hort. Beng.* 46 ; *Fl. Ind.* iii. 89.—As. et Afr. trop.
indica, Wall. Cat. n. 1823 = trifolia.
inermis, *Mill. Gard. Dict.* ed. VIII. n. 2.—Jamaic.
involucrata, Forsk. ex Sw. Fl. Ind. Occ. ii. 1055 = stricta.
involucrata, *Linn. Cent. Pl.* ii. 22; *Amoen. Acad.* iv. 319.—Am. calid.
Kisi, *A. Rich. Tent. Fl. Abyss.* ii. 169.—Abyss.
lamiifolia, Otto & Dietr. Allg. Gartenz. ix. (1841) 372 = canescens.
lanuginosa, Mill. Gard. Dict. ed VIII. n. 3 = involucrata.
latifolia, Tausch, in Flora, xix. (1836) 391 = indica.
lavandulacea, Willd. Sp. Pl. iii. 319 = Lippia asperifolia.
leucantha, Hort. ex Schau. in DC. Prod. xi. 606 = indica.
lilacina, *Desf. Cat. Hort. Par.* ed. III. 392.—Bras.
lippioides, Hook. & Arn. Bot. Beech. Voy. 305 = Lippia geminata.
lippioides, *Spreng. Syst.* ii. 763.—Bras.
Lockhardtii, D. Don, ex Schau. in Mart. Fl. Bras. ix. 232 = Chamissonis.
lucida, *Schau. in DC. Prod.* xi. 602.—Bras.
Lundiana, *Schau. l. c.* 605.—Bras.
macrophylla, *Schau. l. c.* 595.—Bras.
macropoda, *Torr. Bot. Mex. Bound.* 127.—Mexic.
Maximiliani, *Schau. in DC. Prod.* xi. 595.—Bras.
melissaefolia, [Soland. in] Ait. Hort. Kew. ed. I. ii. 352 = flava.
melissaeodorifera, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 124.—Guiān.
microcephala, *A. Rich. Fl. Cub. Fanerog.* ii. 141.—Cuba.
microphylla, *Franch. in Révoil, Çomali*, 49.—Afr. trop.
microphylla, *Mart. Reise Bras.* ii. 792.—Bras.
mista, *Linn. Syst.* ed. XII. 417.—Ind. occ.
mixta, Hort. ex Schau. in DC. Prod. xi. 598 = horrida.
mixta, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 227 = mista.
mollis, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.-Oct. 1829) 184.—Mexic.
mollissima, Desf. Cat. Hort. Par. ed. III. 393 = Lippia geminata.
Moritziana, *Otto & Dietr. Allg. Gartenz.* ix. (1841) 369.—Venezuela.
multicolor, *Lem. Fl. des Serres*, Sér. I. iii. (1847) 239 *Misc.* 5.—Mexic.
multiflora, *Hort. ex Otto & Dietr. Allg. Gartenz.* ix. (1841) 370.—Ind. occ.
mutabilis, Lippold, ex Otto & Dietr. l. c. x. (1842) 314 = mista.
mutabilis, Salisb. Prod. 107 = L. Camara.
mutabilis, Weigel, in Phys. Sällsk. Handl. i. (1776) 46 = L. Camara?
nivea, *Vent. Jard. Malm.* 8. t. 8.—Am. trop.
odorata, Linn. Syst. ed. XII. 418 = involucrata.
odorata, Weigelt, ex Cham. in Linnaea, vii. (1832) 215 = Lippia geminata.
origanoides, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 327 = Lippia graveolens.
parvifolia, *Desf. Cat. Hort. Par.* ed. III. 392.—Hab.?
parvifolia, Rafin. Herb. 21; Atl. Journ. 176 = involucrata?

LANTANA:—
parvifolia, Salisb. Prod. 107 = involucrata.
peduncularis, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1854) 200.—Ins. Galap.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 169.—Abyssin.
pilosa, H. B. & K. Nov. Gen. et Sp. ii. 260 = trifolia.
Pohliana, *Schau. in DC. Prod.* xi. 601.—Bras.
polyacantha, Schau. l. c. 597 = L. Camara.
polycephala, *R. Br. in Salt, Abyss. App.* 65, nomen.—Abyssin.
procurrens, *Schau. in DC. Prod.* xi. 596.—Bras.
Pseudo-thea, St. Hil. Pl. Us. Bras. t. 70 = Lippia Pseudo-thea.
purpurea, *Benth. & Hook. f. Gen.* ii. 1142.—Mexic.; Venezuela.
purpurea, *Hornem. Hort. Hafn.* ii. 583.—Am. austr.
racemosa, *Lour. Fl. Cochinch.* ii. 376.—Cochinch.
Radula, *Sw. Prod. Veg. Ind. Occ.* 92.—Ind. occ.
recta, Cham. in Linnaea, vii. (1832) 125 = lilacina.
recta, [Soland. in] Ait. Hort. Kew. ed. I. ii. 351 = involucrata.
repens, Spreng. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 563, sphalm. = Lippia nodiflora.
reticulata, Pers. Syn. ii. 141 = Sellowiana.
reticulata, *Rafin. Sylva Tellur.* 82.—Hab.?
Riedeliana, *Schau. in DC. Prod.* xi. 601.—Bras.
robusta, *Schau. l. c.*—Bras.
rosea, *Rafin. Sylva Tellur.* 83.—Hab.?
rubra, Perr. in Mém. Soc. Linn. Par. iii. (1825) 124 = indica?
rugosa, Thunb. Prod. Pl. Cap. 98 = salvifolia.
rugosa, Willd. ex Schau. in Mart. Fl. Bras. ix. 263 = L. Radula.
rugulosa, *H. B. & K. Nov. Gen. et Sp.* ii. 261.—Ecuador.
salicifolia, *H. B. & K. l. c.* 259.—N. Granat.
salviaefolia, Cham. in Linnaea, vii. (1832) 125 = lilacina.
salvifolia, *Jacq. Hort. Schoenb.* iii. 18. t. 285.—Afr. austr.
salvifolia, Linn. Syst. ed. X. 1116 = Buddleia salvifolia.
Salzmanni, *Schau. in DC. Prod.* xi. 602.—Bras.
sanguinea, *Medic. in Act. Acad. Theod. Palat.* iii. *Phys.* (1775) 229.—Hab.?
sarmentosa, Spreng. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 563, sphalm. = Lippia nodiflora.
scabiosaeflora, *H. B. & K. Nov. Gen. et Sp.* ii. 261.—Peruv.
scabra, *Wall. Cat.* n. 1822.—Ind. or.
scabrida, [Soland. in] Ait. Hort. Kew. ed. I. ii. 352 = L. Camara.
Selloi, Steud. Nom. ed. II. ii. 8 = seq.
Sellowiana, *Link & Otto, Ic. Pl. Sel. Berol.* 107. t. 50.—Am. austr.
spicata, Vell. Fl. Flum. 254; vi. t. 58 = brasiliensis.
Spiraeastrum, *Mart. & Schau. in DC. Prod.* xi. 596.—Bras.
splendens, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 226 = crenulata.
stricta, *Sw. Prod. Veg. Ind. Occ.* 92.—Jamaic.
suaveolens, Desf. Cat. Hort. Par. ed. III. 393 = involucrata.
suaveolens, Poir. ex Spreng. Syst. ii. 762 = stricta.
teucriifolia, Otto & Dietr. Allg. Gartenz. ix. (1841) 371 = hispida.
tiliaefolia, *Cham. in Linnaea*, vii. (1832) 122.—Bras.
trifolia, Cham. in Linnaea, vii. (1832) 124, ex parte = undulata, *Schrank*.
trifolia, *Linn. Sp. Pl.* 626.—Am. trop.
triplinervia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 205.—Java.
undulata, *Rafin. Sylva Tellur.* 82 (Quid?).—Am. bor.?
undulata, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 56.—Bras.
urticaefolia, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
variegata, Otto & Dietr. Allg. Gartenz. x. (1842) 314 = purpurea.
velutina, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 325.—Mexic.
viburnoides, Blanco, Fl. Filip. ed. II. 345 = L. Camara.
viburnoides, *Vahl, Symb. Bot.* i. 45.—Abyssin.; Arab.
violacea, Desf. Cat. Hort. Par. ed. III. 392 = salvifolia.
viscosa, *Pohl, ex Schau. in DC. Prod.* xi. 601.—Bras.
Wightiana, Wall. Cat. n. 2655 = indica.

LANTANOPSIS, Wright, ex Griseb. in Mem. Am. Acad. N. S. viii. (1863) 513. *COMPOSITAE*, Benth. & Hook. f. ii. 344.
hispidula, *Wright, l. c.*—Cuba.

LANTHORUS, Presl, Epim. Bot. 256 (1849) = **Loranthus**, Linn.
spicifer, Presl, l. c. 257 = Loranthus spicifer.

LANZANA, Stokes, Bot. Mat. Med. ii. 543 (1812) = **Buchanania**, Roxb. (Anacard.).
solitaria, Stokes, l. c. = B. Lanzan.

LAOTHOË, Rafin. Fl. Tellur. iii. 53 (1836) = **Chlorogalum**, Kunth (1843) (Liliac.).
pomeridiana, Rafin. l. c. = C. pomeridianum.

LAPAGERIA, Ruiz & Pav. Fl. Per. iii. 64. t. 297 (1802). *LILIACEAE*, Benth. & Hook. f. iii. 767.
CAPIA, Domb. ex Juss. in Dict. Sc. Nat. vi. 500 (1805).
PAGERIA, Juss. in Dict. Sc. Nat. iii. 53 (1816).
PHAENOCODON, Salisb. Gen. Pl. Fragm. 58 (1866).
alba, Decne. in Rev. Hortic. Sér. IV. i. (1852) 441 = rosea.
Hookeri, Bridges, ex Hook. Journ. Bot. i. (1834) 178 = rosea.
rosea, *Ruiz & Pav. Fl. Per.* iii. 65. t. 297.—Chili.

LAPASATHUS, Presl, Bot. Bemerk. 132 (1844); in Abh. Boehm. Ges. V. iii. (1845) 562 = **Aspalathus**, Linn. (Legumin.).
lanceolata, Presl, l. c. = A. lanceolata.

LAPATHON, Rafin. New Fl. Am. iv. 51 (1836) = **Rumex**, Linn. (Polygon.).
sylvaticum, Rafin. l. c. = R. sanguineus.

LAPATHUM, [Tourn.] Adans. Fam. ii. 277 (1763) = **Rumex**, Linn. (Polygon.).
Acetosa, Scop. Fl. Carn. ed. II. i. 260 = R. Acetosa.
Acetosella, Scop. l. c. 261 = R. Acetosella.
acutum, Scop. l. c. 262 = R. acutus.
alpestre, Scop. l. c. 261 = R. scutatus.
alpinum, Lam. Fl. Fr. iii. 7 = R. alpinus.
aquaticum, Scop. Fl. Carn. ed. II. i. 263 = R. aquaticus.
arborescens, Moench, Meth. 355 = R. Lunaria.
arvense, Lam. Fl. Fr. iii. 8 = R. Acetosella.
aureum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 275 = R. maritimus.
bienne, Moench, Meth. 354 = R. Britannica.
bucephalophorum, Lam. Fl. Fr. iii. 7 = R. bucephalophorus.
bucephalorum, Moench, Meth. 356 = praec.
conglomeratum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 273 = R. conglomeratus.
crispum, Scop. Fl. Carn. ed. II. i. 261 = R. crispus.
Daiwoo, Siebold, in Verh. Bat. Gen. xii. (1830) 19.—Japon.
digynum, Lam. Fl. Fr. iii. 6 = Oxyria digyna.
divaricatum, Rénault, Fl. Dép. Orne, 60 = R. pulcher.
giganteum, Opiz, Seznam, 57 = R. Hydrolapathum.
glomeratum, Gilib. Exercit. ii. 444 = R. conglomeratus.
hortense, Lam. Fl. Fr. iii. 2 = R. Patientia.
Hydrolapathum, Scop. Fl. Carn. ed. II. i. 262 = R. Hydrolapathum.
limosum, Rénault, Fl. Dép. Orne, 60 = R. palustris.
maritimum, Moench, Meth. 355 = R. maritimus.
maximum, Gilib. Exercit. ii. 443 = R. aquaticus.
minus, Lam. Fl. Fr. iii. 4 = R. maritimus.
nemorosum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 145 = R. sanguineus.
obtusifolium, Moench, Meth. 356 = R. obtusifolius.
palustre, S. F. Gray, Nat. Arr. Brit. Pl. ii. 275 = R. palustris.
persicarioides, Moench, Meth. 355 = R. maritimus.
pratense, Lam. Fl. Fr. iii. 8 = R. Acetosa.
pulchrum, Moench, Meth. Suppl. 121 = R. pulcher.
rivale, Rénault, Fl. Dép. Orne, 60 = R. Hydrolapathum?
rubellum, Moench, Meth. 355 = R. aegyptiacus, ucranicus.
sanguineum, Moench, l. c. 354 = R. sanguineus.
scutatum, Lam. Fl. Fr. iii. 6 = R. scutatus.
sinuatum, Lam. l. c. 5 = R. pulcher.
sylvestre, Lam. l. c. 4 = R. obtusifolius.
vesicarium, Moench, Meth. 356 = R. vesicarius.
virgatum, Moench, l. c. 355 = R. acutus.
viride, S. F. Gray, Nat. Arr. Brit. Pl. ii. 274 = R. sanguineus.

LAPEIROUSIA, Thunb. Prod. Pl. Cap. 163 (1800) = **Peyrousea**, DC. (Compos.).
calycina, Thunb. l. c. = P. calycina.
Thunbergii, Cass. in Dict. Sc. Nat. xxv. 251 = P. calycina.

LAPEYROUSA, Poir. in Dict. Sc. Nat. xxv. 252 (1822) = seq.

LAPEYROUSIA, Pourr. in Mém. Acad. Toul. iii. (1788) 79. t. 6 (Lapeirousia). *IRIDEAE*, Benth. & Hook. f. iii. 705.
ANOMATHECA, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 227.
ANOMAZA, Salisb. in Trans. Hort. Soc. i. (1812) 323.
MERISTOSTIGMA, A. Dietr. Sp. Pl. ii. 593 (1833).
OVIEDA, Spreng. Syst. i. 147 (1825).
PEYROUSIA, Poir. in Dict. Sc. Nat. xxxix. 363 (1826).
PSILOSIPHON, Welw. ex Baker, in Trans. Linn. Soc. Bot. Ser. II. i. (1878) 272.
SOPHRONIA, Licht. in Roem. & Schult. Syst. i. 482 (1817).
abyssinica, *Baker, in Journ. Linn. Soc.* xvi. (1877) 155. —Afr. trop.
aculeata, Sweet, Hort. Brit. ed. I. 396 = compressa.
anceps, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 238 = compressa.
azurea, Eckl. Top. Verz. 31 = corymbosa.
Bainesii, *Baker, in Journ. Bot.* xiv. (1876) 338.—Afr. austr.
bracteata, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 238; Bot. Mag. sub t. 1246 = fissifolia.
compressa, *Pourr. in Mém. Acad. Toul.* iii. (1788) 80. t. 6.—Afr. austr.
corymbosa, *Ker-Gawl. in Bot. Mag.* t. 595.—Afr. austr.
cyanescens, *Baker, in Trans. Linn. Soc. Bot.* Ser. II. i. (1878) 272.—Afr. trop.
divaricata, *Baker, in Journ. Bot.* xiv. (1876) 337.—Afr. austr.
erythrantha, *Baker, in Journ. Linn. Soc.* xvi. (1877) 155.—Afr. austr. et trop.
Fabricii, *Ker-Gawl. in Bot. Mag.* sub t. 1246.—Afr. austr.
falcata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 238.—Afr. austr.
fasciculata, *Ker-Gawl. l. c.*—Afr. austr.
fastigiata, Ker-Gawl. l. c. = corymbosa.
fissifolia, *Ker-Gawl. l. c.*—Afr. austr.
fistulosa, *Baker, in Journ. Linn. Soc.* xvi. (1877) 155. —Afr. austr.
fragrans, *Baker, in Trans. Linn. Soc. Bot.* Ser. II. i. (1878) 272.—Afr. trop.
juncea, *Pourr. in Mém. Acad. Toul.* iii. (1788) 79; *Ker-Gawl. in Bot. Mag.* t. 606.—Afr. austr.
littoralis, *Baker, in Trans. Linn. Soc. Bot.* Ser. II. i. (1878) 272.—Afr. trop.
macrochlamys, *Baker, in Journ. Bot.* xiv. (1876) 338. —Afr. austr.
manuleaeflora, Eckl. Top. Verz. 31 = micrantha.
marmorata, Steud. Nom. ed. II. ii. 9 = juncea.
micrantha, *Baker, in Journ. Linn. Soc.* xvi. (1877) 156.—Afr. austr.
montana, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 359.—Afr. austr.
odoratissima, *Baker, in Trans. Linn. Soc. Bot.* Ser. II. i. (1878) 273.—Afr. trop.
purpureo-lutea, *Baker, in Journ. Linn. Soc.* xvi. (1877) 155.—Afr. austr.
silenoides, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 238.—Afr. austr.

LAPEYROUSIA, Spreng. Anleit. ii. 552 (1818) = LAPEIROUSIA, Thunb. = **Peyrousia**, DC. (Compos.).

LAPHAMIA, A. Gray, Pl. Wright. i. 99. t. 9 (1852). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
MONOTHRIX, Torr. in Stansb. Utah Exped. 389. t. 7 (1852).
aglossa, *Benth. & Hook. f. Gen.* ii. 398.—Mexic.
angustifolia, *A. Gray, Pl. Wright.* i. 100.—N. Mexic.
bisetosa, *Torr. ex A. Gray, l. c.* ii. 106.—Mexic.
cinerea, *A. Gray, in Torr. Bot. Mex. Bound.* 82.—Mexic.
coronopifolia, *Hemsl. Biol. Cent. Am. Bot.* ii. 210.—Mexic.
dissecta, Torr. ex A. Gray, Pl. Wright. ii. 81 = Perityle dissecta.

LAPHAMIA:—
Emoryi, Benth. & Hook. f. Gen. ii. 398 = Perityle californica.
halimifolia, *A. Gray, Pl. Wright.* i. 100.—N. Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 101.—Mexic.
Lindheimeri, *A. Gray, Pl. Wright.* i. 101.—Texas.
megacephala, *S. Wats. in Am. Natural.* vii. (1873) 301. —Am. bor. occ.
nuda, Benth. & Hook. f. Gen. ii. 398 = Perityle californica.
Palmeri, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 372. —Arizona.
Parryi, *Benth. & Hook. f. Gen.* ii. 398.—Mexic.
peninsularis, *Greene, in Bull. Calif. Acad.* i. n. 1. (1884) 8.—Calif.
rupestris, *A. Gray, Pl. Wright.* i. 100.—Mexic.
Stansburii, *A. Gray, l. c.* 101.—Utah.

LAPIEDRA, Lag. Gen. et Sp. Nov. 14 (1816). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 720.
gracilis, Baker, in Journ. Linn. Soc. xvi. (1878) 678 = Tapeinanthus humilis.
Martinezii, *Lag. Gen. et Sp. Nov.* 14.—Hispan.
Placiana, Herb. Amaryll. 188 = Martinezii.

LAPITHEA, Griseb. in DC. Prod. ix. 48 (1845) = **Sabbatia**, Adans. (Gentian.).
gentianoides, Griseb. l. c. = S. gentianoides.

LAPLACEA, H. B. & K. Nov. Gen. et Sp. v. 207. t. 461 (1821). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 186.
CLOSASCHIMA, Korth. Verh. Nat. Gesch. Bot. 139. t. 28 (1842).
HAEMOCHARIS, Salisb. Parad. Lond. sub n. 56 (1806).
LINDLEYA, Nees, in Flora, iv. (1821) 299.
WIKSTROEMIA, Schrad. in Goett. Gel. Anz. i. (1821) 710.
amboinensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 114. —Malaya.
aromatica, *Miq. Fl. Ind. Bat. Suppl.* 482.—Malaya.
barbinervis, *Moric. in Mém. Soc. Phys. Genèv.* vii. (1830) 256; *Pl. Nouv. Am.* 16. t. 11.—Peruv.
buxifolia, *Miq. Fl. Ind. Bat. Suppl.* 482.—Sumatra.
cameniaefolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 270.—N. Granat.
camenioides, *Sond. in Linnaea*, xxii. (1849) 549.—Bras.
Curtyana, *A. Rich. Ess. Fl. Cub.* 225.—Cuba.
Haematoxylon, *G. Don, Gen. Syst.* i. 569.—Jamaic.
inaequalilatera, Schott, in Spreng. Syst. iv. App. 408 = semiserrata.
integerrima, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 113. —Java.
intermedia, *Benth. Pl. Hartw.* 126.—Am. austr.
marginata, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 148.—Borneo.
ovalis, *Choisy, l. c.*—Sumatra.
parviflora, *Mart. Reise Bras.* i. 207, nomen.—Bras.
praemorsa, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 100.—Guian.
pubescens, Planch. & Linden, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 269.—N. Granat.
quinoderma, *Wedd. Hist. Nat. Quin.* 33.—Boliv.; Peruv.
semiserrata, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 300. —Bras.
speciosa, *H. B. & K. Nov. Gen. et Sp.* v. 207. t. 461.—Peruv.
subintegerrima, *Miq. Fl. Ind. Bat. Suppl.* 483.—Sumatra.
symplocoides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 269.—N. Granat.
tomentosa, *G. Don, Gen. Syst.* i. 569.—Bras.
viscosa, *Griseb. Fl. Brit. W. Ind.* 104.—Jamaic.
vulcanica, *Korth. Verh. Nat. Gesch. Bot.* 136.—Sumatra.
Wrightii, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 166.—Cuba.

LAPORTEA, Gaudich. in Freyc. Voy. Bot. 498 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 383.
DENDROCNIDE, Miq. Pl. Jungh. i. 29 (1857).
DISCOCARPUS, Liebm. in Vidensk. Selsk. Skrift. V. ii. (1852) 308.
OBLIXILIS, Rafin. Fl. Tellur. iii. 49 (1836).
SCLEPSION, Rafin. ex Wedd. in Ann. Sc. Nat. Sér. IV. vii. (1857) 355.

LAPORTEA:—
alatipes, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 215. —Afr. occ.
americana, *Gaudich. in Freyc. Voy. Bot.* 498.—Am. austr.
amplissima, *Miq. Fl. Ind. Bat.* i. II. 232.—Java.
bulbifera, *Wedd. Monog. Urtic.* 139.—Japon.
canadensis, *Gaudich. in Freyc. Voy. Bot.* 498.—Am. bor.
carolinensis, *Gaudich. l. c.*—Am. bor.
contracta, Blume, ex Miq. Fl. Ind. Bat. i. II. 232 = costata.
costata, *Miq. l. c.* 231.—Java.
crenulata, *Gaudich. in Freyc. Voy. Bot.* 498.—As. trop.
decumana, *Wedd. Monog. Urtic.* 129.—Malaya.
evitata, Wedd. in DC. Prod. xvi. 79 = terminalis.
Gaudichaudiana, *Wedd. Monog. Urtic.* 137.—Ins. Philipp.
gigantea, Gaudich. in Freyc. Voy. Bot. 498 = crenulata.
Gigas, *Wedd. Monog. Urtic.* 129. t. 3.—Austral.
Harveyi, *Seem. in Bonplandia*, ix. (1861) 259; *Fl. Vit.* 238.—Ins. Fiji.
latifolia, Gaudich. Bot. Voy. Bonite, t. 81 = crenulata.
laxiflora, *Wedd. in DC. Prod.* xvi. I. 81.—Java.
Liebmannii, Wedd. Monog. Urtic. 133 = nicaraguensis.
mexicana, Wedd. in DC. Prod. xvi. I. 84 = platycarpa.
microstigma, *Gaudich. ex Wedd. l. c.* 138.—Java.
Milnei, *Seem. Fl. Vit.* 238.—Ins. Fiji.
moroides, *Wedd. Monog. Urtic.* 142.—Austral.
nicaraguensis, *Wedd. in DC. Prod.* xvi. I. 84.—Nicaragua.
oblongata, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 103, 106.—Java.
oleracea, *Wedd. Monog. Urtic.* 141.—Reg. Himal.
peltata, *Gaudich. in Freyc. Voy. Bot.* 498.—Malaya.
photiniphylla, *Urtic. Monog. Wedd.* 138.—Austral.; Ins. Pacif.
platycarpa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 182.—Mexic.
pterostigma, *Wedd. in DC. Prod.* xvi. I. 87.—Ins. Formosa.
pustulata, Wedd. Monog. Urtic. 140 = canadensis.
repens, *Wedd. in DC. Prod.* xvi. I. 81.—Afr. occ.
sinuata, Blume, ex Wedd. l. c. 86 = stimulans.
stimulans, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 103. —Java.
terminalis, *Wight, Ic.* vi. t. 1972.—Ind. or.
ternatense, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 302.—Ins. Ternate.
vitiensis, Seem. in Bonplandia, ix. (1861) 259; Fl. Vit. 239 = photiniphylla.
Vriesiana, *Wedd. in DC. Prod.* xvi. I. 81.—Java.

LAPPA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 189 (1745) = **Arctium**, Linn. (Compos.).
amplissima, Boiss. Diagn. Ser. I. vi. 108 = Cousinia amplissima.
amplissima, Kotschy, Reise Taur. 412 = A. majus.
Arctium, Hill, Veg. Syst. iv. 28 = A. majus.
atlantica, Pomel, Nouv. Mat. Fl. Atl. 285.—Alger.
Bardana, Moench, Meth. 552 = A. majus.
carduelis, Lam. Encyc. i. 378 = Carduus arctioides.
conglomerata, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 79 = A. minus.
edulis, Siebold, ex Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 185 = A. majus.
glabra, Lam. Fl. Fr. ii. 37 = A. majus.
intermedia, Reichb. f. Ic. Fl. Germ. xv. 54. t. 812 = A. nemorosum.
intermedia ×, Simk. in Termész. Füzetek, ii. (1878) 145.—Europ.
Kotschyi, Hort. Vind. ex Boiss. Fl. Orient. iii. 458 = A. majus.
macrosperma, Wallr. in Linnaea, xiv. (1840) 639; et Beitr. Fl. Hercyn. 269 = A. nemorosum.
major, Gaertn. Fruct. ii. 379. t. 162 = A. majus.
minor, Hill, Veg. Syst. iv. 28; DC. Fl. Fr. iv. 77 = A. minus.
nemorosa, Koern. ex Griewank, in Arch. Ver. Naturg. Mecklenb. xxiii. (1870) 189 = A. nemorosum.
notha ×, Ruhm. in Eichl. Jahrb. Berl. (1881) 238.—Europ.
officinalis, All. Fl. Pedem. i. 145 = A. majus.
Personata, Lam. Encyc. i. 378 = Carduus Personata.
platylepis, Boiss. & Balansa, ex Boiss. Fl. Orient. iii. 458.—As. Min.

LAPPA:—
puberis, Bor. Fl. Cent. Fr. ed. III. ii. 758 = Arctium minus.
repens, DC. ex Kar. in Bull. Soc. Nat. Mosc. (1839) 160, nomen.—Turcoman.
rosacea, Wallr. Beitr. Fl. Hercyn. 270 = A. majus?
sylvestris, Pomel, Nouv. Mat. Fl. Atl. 284.—Alger.
tenella, Fisch. & Mey. in DC. Prod. vi. 662 = Cousinia tenella.
tomentosa, Lam. Fl. Fr. ii. 37 = A. majus.
vulgaris, Hill, Veg. Syst. iv. 28 = A. majus.

LAPPAGO, Schreb. Gen. 55 (1789) = **Tragus**, Hall. (Gramin.).
aliena, Spreng. Neue Entdeck. iii. 15 = T. racemosus.
Berteroniana, Schult. ex Steud. Syn. Pl. Gram. 112, in syn. = T. racemosus.
biflora, Roxb. Hort. Beng. 82; Fl. Ind. i. 281 = T. racemosus.
decipiens, Fig. & De Not. in Mem. Acc. Torin. (1854) 387 = T. breviflorus.
latipes, Steud. Nom. ed. II. ii. 9 = Latipes senegalensis.
oplismenoides, Speg. in Anal. Soc. Cient. Argent. xvi. (1883) 132.—Reg. Argent.
phleoides, Fig. & De Not. in Ann. Sc. Nat. Sér. III. xix. (1853) 373.—Nubia.
racemosa, Honck. Syn. i. 440 = T. racemosus.
setacea, Spreng. ex Steud. Nom. ed. II. ii. 9, 11 = Lasiochloa hispida.
setiformis, Spreng. ex Steud. ll. cc. = Lasiochloa hirta.

LAPPAGOPSIS, Steud. Syn. Pl. Gram. 112 (1855) = **Paspalum**, Linn. (Gramin.).
bijuga, Steud. l. c. = P. dissitiflorum.

LAPPARIA, Heist. Syst. 11 (1748) = **Xanthium**, Tourn. (Compos.).

LAPPULA, Moench, Meth. 416 (1794) = **Echinospermum**, Sw. (Boragin.).
concava, F. Muell. Census, 100 = E. concavum.
deflexa, Garcke, Fl. Deutschl. ed. VI. 275 = E. deflexum.
echinata, Gilib. Fl. Lituan. i. 25 = E. Lappula.
Myosotis, Moench, Meth. 417 = E. Lappula.
squarrosa, Dum. Fl. Belg. 40 = E. Lappula.

LAPPULARIA, Pomel, Nouv. Mat. Fl. Atl. 149 (1874) = **Caucalis**, Linn. (Umbellif.).
bifrons, Pomel, l. c. 150 = C. bifrons.
infesta, Pomel, l. c. = C. infesta.
neglecta, Pomel, l. c. = C. infesta.
nodosa, Pomel, l. c. = C. nodosa.

LAPSANA, Linn. Meth. Sex. (1737); Syst. ed. II. 28 (1740) pro parte. *COMPOSITAE*, Benth. & Hook. f. ii. 509.
LAMPSANA, Tourn. ex Rupp. Fl. Jen. ed. Hall. 209 (1745).
adenophora, *Boiss. Diagn.* Ser. I. iv. 423.—As. Min.
alpina, *Boiss. & Bal. l. c.* Ser. II. vi. 117.—As. Min.
apogonoides, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 288.—Japon.
capillaris, Linn. Sp. Pl. 812 = Crepis virens.
Cassia, Boiss. Diagn. Ser. I. xi. 33 = pedunculatis.
chondrilloides, Linn. Sp. Pl. 812 = Phaecasium lampsanoides.
communis, Linn. l. c. 811.—Europ.; Oriens; As. bor.
crassifolia, Vahl, ex DC. Prod. vii. 73 = Pamphalea Commersonii.
crispa, Willd. Sp. Pl. iii. 1624 = communis.
foetida, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 264 = Hyoseris foetida.
foetida, Scop. Fl. Carn. ed. II. ii. 118 = Crepis foetida.
glandulifera, Cass. in Dict. Sc. Nat. xxv. 211 = grandiflora.
gracilis, Lam. Fl. Fr. ii. 102 = Arnoseris pusilla.
grandiflora, *Bieb. Fl. Taur. Cauc.* ii. 261.—Reg. Caucas.
intermedia, *Bieb. Fl. Taur. Cauc.* iii. 540.—Byzant.; Reg. Caucas.
japonica, *Burm. f. Fl. Ind.* 174.—Japon.
Koelpinia, Linn. f. Suppl. 248 = Koelpinia linearis.
lampsanaefolia, Mill. Gard. Dict. ed. VIII. n. 3.—Rhagadiolus edulis.
leontodontoides, Scop. ex DC. Prod. vii. 82 = Crepis foetida.

LAPSANA:—
lyrata, Willd. Enum. Hort. Berol. 832 = grandiflora.
macrocarpa, Coss. in Bull. Soc. Fr. ix. (1862) 173 = communis.
minima, All. Fl. Pedem. i. 206 = Arnoseris pusilla.
parviflora, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 396.—Japon.
peduncularis, *Boiss. Fl. Orient.* iii. 720.—As. Min.; Syria.
pisidica, Boiss. & Heldr. Diagn. Ser. I. xi. 33 = peduncularis.
pubescens, Hornem. Hort. Hafn. Suppl. 91 = communis.
pulchra, Vill. Hist. Pl. Dauph. iii. 163 = Phaecasium lampsanoides.
pusilla, Willd. Sp. Pl. iii. 1623 = Arnoseris pusilla.
ramosissima, Boiss. & Heldr. Diagn. Ser. I. xi. 34 = peduncularis.
ramosissima, Crantz, Inst. i. 236 = Rhagadiolus edulis.
Rhagadiolus, Linn. Sp. Pl. 812 = Rhagadiolus edulis.
silvatica, Nym. Consp. 474 = communis.
sonchifolia, Gilib. Fl. Lituan. i. 243 = communis?
stellata, Linn. Sp. Pl. 811 = Rhagadiolus edulis.
sylvatica, Wallr. Beitr. Fl. Hercyn. ii. 125; ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 420 = communis.
taraxacoides, Forsk. Fl. Aegypt. Arab. 145 = Hyoseris lucida.
taraxaconoides, J. F. Gmel. Syst. 1571 = praec.
virgata, *Desf. Fl. Atlant.* ii. 235.—Afr. bor.
vulgaris, Gromow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 513, nomen = communis.
Zacintha, Linn. Sp. Pl. 811 = Zacintha versicosa.
Zasyntha, Steud. Nom. ed. II. ii. 9 = praec.

LAPULA, Gilib. Fl. Lituan. i. 25 (1781) = LAPPULA Moench = **Echinospermum**, Sw. (Boragin.).

LARBRAEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 348 = LARBREA, A. St. Hil. = **Stellaria**, Linn. (Caryophyll.).

LARBREA, A. St. Hil. in Mém. Mus. Par. ii. (1815) 287 = **Stellaria**, Linn. (Caryoph.).
aquatica, A. St. Hil. l. c. 261 = S. uliginosa.
aquatica, Ser. in DC. Prod. i. 395 = S. aquatica.
bracteata, Reichb. Ic. Fl. Germ. xv. 36 t. 226 = S. uliginosa.
crassifolia, Reichb. l. c. = S. crassifolia.
graminea, Fuss, Fl. Transs. 119 = S. graminea.
linoides, Opiz, Seznam, 57 = S. linoides.
palustris, Fuss, Fl. Transs. 118 = S. palustris.
uliginosa, Reichb. Ic. Fl. Germ. xv. 36 = S. uliginosa.

LARDIZABALA, Ruiz & Pav. Prod. 143. t. 37 (1794). *BERBERIDEAE*, Benth. & Hook. f. i. 42.
BOISSIERA, Domb. ex DC. Syst. i. 512, in syn. (1818).
COGYLIA, Molina, Sagg. Chile, ed. II. 300 (1810).
biternata, *Ruiz & Pav. Syst. Veg.* 286.—Chili.
discolor, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 64. t. 189.—Am. austr.
infusiata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ii. (1858) 431.—Chili.
silvicola, *Miers, l. c.* 191.—Chili.
ternata, *Molina, Sagg. Chile*, ed. II. 137.—Chili.
trifoliata, Ruiz & Pav. Syst. Veg. 286 = Boquila trifoliata.
trifoliolata, DC. Syst. i. 513 = praec.
triternata, *Ruiz & Pav. Syst. Veg.* 287.—Chili.

LARENTIA, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 362 = **Cypella**, Herb. (Irid.).
linearis, Klatt, l. c. (= *Alophia linearis*).—Venezuela.

LAREPHES, Rafin. Fl. Tellur. iv. 86 (1836) = **Echium**, Tourn. (Boragin.).
parviflorum, Rafin. l. c. = E. tenue.

LARETIA, Gill & Hook. in Hook. Bot. Misc. i. (1830) 329. t. 65. *UMBELLIFERAE*, Benth. & Hook. f. i. 875.
acaulis, *Gill & Hook. l. c.*—Chili.

LARIADENIA, [Schlecht.] in Bot. Zeit. iv. (1846) 284, sphalm. = **Lasiadenia**, Benth. (Thymel.).

LARIX, Tourn. ex Adans. Fam. ii. 480 (1763). *CONIFERAE*, Benth. & Hook. f. iii. 442.
altaica, Fisch. ex Gord. Pinet. 127 = sibirica.
amabilis, [Nels.] Senilis, Pinac. 84 = Pseudolarix Kaempferi.
americana, *Michx. Fl. Bor. Am.* ii. 203; *Arb. Forest. Am. Sept.* iii. 37. t. 4.—Am. bor.
archangelica, Laws. Agric. Man. 389 = sibirica.
caducifolia, Gilib. Exercit. ii. 413 = europaea.
Cedrus, Mill. Gard. Dict. ed. VIII. n. 3 = Cedrus Libani.
chinensis, Mill. l. c. n. 2 = Cunninghamia sinensis.
communis, Laws. ex Gord. Pinet. 124 = europaea.
dahurica, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 101.—Sibir.
decidua, Mill. Gard. Dict. ed. VIII. n. 1 = europaea.
Deodara, C. Koch, Wochenschr. x. (1867) 308 = Cedrus Libani.
europaea, *DC. Fl. Fr.* iii. 277.—Europ.; As. bor.
europaea, Middend. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 255 = dahurica.
excelsa, Link, in Abh. Berl. Akad. (1827) 182 = europaea.
Fraseri, Curt. ex Gord. Pinet. 129 = americana.
Gmelini, Ledeb. ex Gord. l. c. 123 = dahurica.
Griffithiana, Carr. Conif. ed. I. 278 = seq.
Griffithii, *Hook. f. & Thoms. in Hook. f. Illustr. Himal. Pl.* t. 21.—Reg. Himal.
intermedia, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 101 = sibirica.
intermedia, Lodd. Cat. (1836) 50; Link, in Linnaea, xv. (1841) 535 = americana.
japonica, Hort. ex Carr. Conif. ed. I. 272 = leptolepis.
japonica, *A. Murr. Pines & Firs of Japan*, 94. f. 178.—Japon.
Kaempferi, Fortune, ex Gord. Pinet. 292 = Pseudolarix Kaempferi.
kamtschatica, Hort. ex Carr. Conif. ed. I. 279 = dahurica.
laricina, C. Koch, Dendrol. ii. II. 263 = americana.
Ledebourii, Rupr. Beitr. Pfl. Russ. Reich. ii. 56 = sibirica.
leptolepis, *Hort. ex Endl. Conif.* 338.—Japon.
Lyallii, *Parl. in Journ. Bot.* i. (1863) 35; *Gard. Chron.* (1863) 916.—Am. bor. occ.
microcarpa, Forbes, Pinet. Woburn. 139. t. 47 = americana.
Nuttallii, Parl. ex Gord. Pinet. ed. II. 176 = occidentalis.
occidentalis, *Nutt. Sylv. N. Am.* ii. 199.—Am. bor. occ.
orientalis, Thunb. Fl. Jap. 275 = leptolepis.
patula, Salisb. in Trans. Linn. Soc. viii. (1808) 314 = Cedrus Libani.
pendula, Salisb. l. c. 313 = americana.
Pseudo-larix, Lodd. ex Gord. Pinet. 127 = sibirica.
pyramidalis, Salisb. in Trans. Linn. Soc. viii. (1808) 313 = europaea.
repens, Laws. ex Steud. Nom. ed. II. ii. 10 = europaea.
rossica, Sabine, ex Loud. Encyc. Trees, 1054 = europaea.
rubra, Hort. ex Steud. Nom. ed. II. ii. 10 = americana.
sibirica, *Ledeb. Fl. Alt.* iv. 204.—Sibiria.
sibirica, Nois. ex Steud. Nom. ed. I. 463; ed. II. ii. 10 = europaea, leptolepis.
sikkimensis, Hook. ex Gord. Pinet. ed. II. 171 = Griffithii.
tenuifolia, Salisb. in Trans. Linn. Soc. viii. (1808) 313 = americana.
vulgaris, Fisch. ex Spach, Hist. Vég. Phan. xi. 432 = europaea.

LARMZON, Roxb. Fl. Ind. ii. 385, in syn. (1832) = **Buchanania**, Roxb. (Anacard.).

LARNALLES, Rafin. Fl. Tellur. iii. 70 (1836) = **Commelina**?, Linn.
dichotoma, Rafin. l. c. 71 (Quid?).—Hab.?
glauca, Rafin. l. c. (Quid?).—Hab.?

LARNANDRA, Rafin. Neogenyt. 4 (1825) = **Epidendrum**, Linn. (Orchid.).
conopsea, Rafin. Fl. Tellur. ii. 40 = E. conopseum.
Magnolia, Rafin. Neogenyt. 4 = E. conopseum.

LARNASTYRA, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
Claytoni, Rafin. l. c. = S. Claytoni.
indica, Rafin. l. c. = S. indica.
lyrata, Rafin. l. c. = S. lyrata.
urticifolia, Rafin. l. c. = S. urticifolia.
verbenacea, Rafin. l. c. = S. Verbenaca.

LARNAX, Miers, in Ann. & Mag. Mat. Hist. Ser. II. iv. (1849) 37 = **Athenaea**, Sendtn. (Solanac.).
Morrisoni, Miers, Illustr. S. Am. Pl. ii. App. 60 = Withania Morrisoni.
orinocensis, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 38 = Withania orinocensis.
ramosa, Miers, Illustr. S. Am. Pl. ii. App. 61 = ithania ramosa.
sordida, Miers, l. c. = Chamaesaracha sordida.
subtriflora, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. 1849) 38 = Withania subtriflora.
xalapensis, Miers, l. c. = A. xalapensis.

LAROCHEA, Pers. Syn. i. 337 (1805) = **Crassula**, Dill.
coccinea, Pers. Syn. i. 337 = Rochea coccinea.
cymosa, Haw. Syn. Pl. Succ. 50 = C. cymosa.
falcata, Pers. Syn. i. 337 = C. falcata.
fascicularis, Schult. Syst. vi. 709 = Rochea fascicularis.
flava, Haw. Syn. Pl. Succ. 50 = C. flava.
odoratissima, Haw. l. c. 51 = Rochea odoratissima.
tiniflora, Lem. Illustr. Hortic. viii. (1861) sub t. 269 = Rochea tiniflora.

LARRADIA, Pritz. Ic. Ind. i. 619 (1855), sphalm. = **Lavradia**, Vell. (Violar.).

LARREA, Cav. in Anal. Hist. Nat. ii. (1800) 119. tt. 18, 19. *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 267.
cuneifolia, *Cav. l. c.* 123.—Reg. Argent.
divaricata, *Cav. l. c.* 122.—Reg. Argent.
glutinosa, *Engelm. in Wisliz. Tour N. Mexic.* 93.—Mexic.
mexicana, *Moric. Pl. Nouv. Am.* 71. t. 48.—Mexic.
nitida, *Cav. in Anal. Hist. Nat.* ii. (1800) 120.—Reg. Argent.
trigyna, Kunze, ex Klotzsch, in Linnaea, x. (1835–36) 432 = Wendtia Poeppigiana.

LARREA, Orteg. Hort. Matr. Dec. 15. t. 2 (1797) = **Hoffmanseggia**, Cav. (Legumin.).
glauca, Orteg. l. c. = H. falcaria.
gracilis, Ruiz & Pav. Fl. Per. iv. t. 377 = H. gracilis.
viscosa, Ruiz & Pav. l. c. = H. viscosa.

LARYSACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 123 = **Ruellia**, Linn. (Acanthac.).
ciliatus, Oerst. l. c. = R. ciliata.

LASCADIUM, Rafin. Fl. Ludov. 114 (1817) 'nomen delendum,' Benth. & Hook. f. Gen. iii. 256.
lanatum, Rafin. Fl. Ludov. 114.—Lousiana.
lanuginosum, Rafin. ex Steud. Nom. ed. II. ii. 10 = praec.

LASEGUEA, A. DC. in DC. Prod. viii. 481 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 725.
acutifolia, *DC. in Ann. Sc. Nat.* Sér. III. i. (1844) 261. —Bras.
antennacea, Miers, Apocyn. S. Am. 251 = Echites antennacea.
bicolor, Miers, l. c. 251 = Echites bicolor.
emarginata, *A. DC. in DC. Prod.* viii. 481.—Bras.
erecta, Muell. Arg. in Mart. Fl. Bras. vi. 1. 135 = acutifolia.
foliosa, Miers, Apocyn. S. Am. 253 = Mandevilla foliosa.
glabra, DC. in Ann. Sc. Nat. Sér. III. i. (1844) 262 = acutifolia.
Guilleminiana, A. DC. in DC. Prod. viii. 481 = acutifolia.
Hookeri, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 136.—Bras.
Jaegeri, Miers, Apocyn. S. Am. 254 = Urechites Jaegeri.
latiuscula, *Miers, l. c.* 251.—Guian.
leptocarpa, Miers, l. c. 254 = Parsonsia leptocarpa.
obliquinervia, DC. in Ann. Sc. Nat. Sér. III. i. (1844) 261 = acutifolia.
Pentlandiana, *DC. l. c.* 262.—Bras.
pubiflora, *Miers, Apocyn. S. Am.* 253.—Ind. occ.
subspicata, Miers, l. c. 252 = Echites subspicata.
venustula, *Miers, l. c.*—Guian.
villosa, *Miers, l. c.* 250.—Nicaragua.

LASER, Borck. ex Gaertn. Mey. & Scherb. Fl. Wett. i. 244, 384 (1799); Baumg. Enum. Stirp. Transsylv. i. 227 (1816) = **Laserpitium**, Linn. (Umbellif.).
aquilegifolium, Roehl. ex Steud. Nom. ed. I. 463; ed. II. ii. 10 = Siler trilobum.

LASER:—
trilobum, Baumg. Enum. Stirp. Transsylv. i. 228 = Laserpitium marginatum.
trilobum, Borck. ex Gaertn. Mey. & Scherb. Fl. Wett. i. 384 = Siler trilobum.

LASERPICIUM, Rivin. ex Aschers. Fl. Brandenb. 256 (1864) = seq.

LASERPITIUM, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 929.
ASPITIUM, Neck. ex Steud. Nom. ed. II. ii. 10 (1841).
GUILLONEA, Coss. Pl. Nouv. Esp. 109 (1851).
LASER, Borck. ex Gaertn. Mey. & Scherb. Fl. Wett. i. 244, 384 (1799).
Aciphylla, Linn. f. Suppl. 181 = Ligusticum Aciphylla.
affine, *Ledeb. Fl. Ross.* ii. 335.—Reg. Caucas.
africanum, F. G. Dietr. Vollst. Lexik. Gaertn. v. 315 (= *Cnidium Fontanesii*).—Afr. bor.
alpinum, Waldst. & Kit. Pl. Rar. Hung. ii. 281 = Siler trilobum.
ambiguum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 31 = marginatum.
angustifolium, *Georgi, Beschr. Russ. Reich.* iv. 847 (Quid?).—Polon.
angustifolium, Linn. Sp. Pl. 248 = gallicum.
angustifolium, Scop. Fl. Carn. ed. II. i. 197 = peucedanoides.
angustissimum, Willd. Sp. Pl. i. 1416 = gallicum.
aquilegiaefolium, Jacq. Fl. Austr. ii. 29. t. 147 = Siler trilobum.
aquilegifolium, DC. Fl. Fr. Suppl. 510 = Nestleri.
aquilegifolium, Moretti, Notiz. in Mem. i. 255; ex Bertol. Fl. Ital. iii. 404 = latifolium.
aquilegifolium, Murr. Syst. ed. XIV. 281 = Siler trilobum.
aquilegifolium, Tenore, Fl. Nap. Prod. 19; ex ej. Fl. Nap. iii. 345 = Trochiscanthes nodiflora.
Archangelica, Hort. ex Link, Enum. Hort. Berol. i. 273 = hispidum.
Archangelica, *Wulf. in Jacq. Coll.* i. 214; *Ic. Pl. Rar.* i. t. 58.—Europ.
asperrimum, Schur, Enum. Pl. Transs. 269 = latifolium.
asperum, Crantz, Stirp. Austr. ed. I. fasc. iii. 50; ed. II. iii. 179 = latifolium.
Athamantae, *Spreng. in Schult. Syst.* vi. 624.—Sibiria.
atlanticum, Poir. Encyc. Suppl. iii. 304 = Elaeoselinum foetidum.
aureum, *Willd. Sp. Pl.* i. 1416.—As. Min.
bavaricum, *Schrank, Baier. Fl.* i. 548.—Bavar.
canescens, Boiss. Elench. 49 = scabrum.
capense, Thunb. Prod. Pl. Cap. 50 = Peucedanum capense.
carniolicum, Bernh. ex Steud. Nom. ed. II. ii. 10 = Siler trilobum.
Carota, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 46 = Thapsia polygama.
caucasicum, Bieb. Fl. Cauc. i. 222 = Carum caucasicum.
Cervaria, C. C. Gmel. Fl. Bad. i. 657 = latifolium.
chironicum, Hill, Veg. Syst. vi. 30 = seq.
Chironium, Linn. Sp. Pl. 249 = Opoponax Chironium.
Chironium, Scop. Fl. Carn. ed. II. i. 197 = L. Archangelica.
commune, Mill. Gard. Dict. ed. VIII. n. 1 = latifolium.
coniifolium, Wall. Cat. n. 584 = Selinum tenuifolium.
crispum, Turra, Fl. Ital. Prod. 65 = latifolium.
crithminum, Link, in Buch, Beschr. Canar. Ins. 151 = Astydamia canariensis.
cuneatum, Moench, Meth. 79 = gallicum.
cuneifolium, Nestler, ex Soyer-Willem. Obs. 87 = Nestleri.
cyanapiifolium, Marschl. in Flora, xvi. (1833) 456 = Ligusticum cynapiifolium.
daucoides, Desf. Fl. Atlant. i. 253 = Daucus laserpitioides.
daucoides, Duf. ex DC. Prod. iv. 206 = prutenicum.
dauricum, Jacq. Hort. Vindob. iii. 22. t. 38 (= *Cnidium dahuricum*).—Dahuria.
elegans, Clarke, Trav. ii. 262 = aureum.
ferulaceum, *Lapeyr. Hist. Abr. Pl. Pyr.* 152.—Mont. Pyren.
ferulaceum, Linn. Sp. Pl. ed. II. 358 = Prangos ferulacea.

LASERPITIUM:—
Fontanesii, Pers. Syn. i. 313 = Elaeoselinum foetidum.
formosum, Willd. Sp. Pl. i. 1416 = gallicum.
gallicum, *Linn. Sp. Pl.* 248.—Europ.
gallicum, Scop. Fl. Carn. ed. II. i. 196 = prutenicum.
garganicum, Bertol. Fl. Ital. iii. 398 = L. Siler.
garganicum, *Tenore, Fl. Nap.* i. 122. t. 22.—Ital.; Graec.
Gaudinii, Moretti, in Comol. Prod. Fl. Com. 52 = marginatum.
glabrum, Crantz, Stirp. Austr. ed. I. fasc. iii. 54; ed. II. 181 = latifolium.
glaucum, Cav. Ic. ii. 72. t. 190 lapsu = scabrum.
gracile, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 30 = latifolium.
gummiferum, *Desf. Fl. Atlant.* i. 254. t. 72.—Hispan.; Afr. bor.
Halleri, All. Fl. Pedem. ii. 11 = L. Panax.
hirsutum, Gilib. Fl. Lituan. ii. 13 = prutenicum.
hirsutum, Lam. Fl. Fr. iii. 648 = L. Panax.
hirtellum, Gaud. Fl. Helv. ii. 350 = nitidum.
hispidum, *Bieb. Fl. Taur. Cauc.* i. 221.—Reg. Caucas.
Krapffii, *Crantz, Cl. Umb. Emend.* 67.—Istria.
latifolium, *Linn. Sp. Pl.* 248; *Lam. Encyc.* iii. 423.—Europ.
laxiflorum, *Fisch. ex Sweet, Hort. Brit.* ed. II. 253, nomen.—Sibiria.
leucolaenum, Boiss. & Bal. ex Boiss. Fl. Orient. ii. 1065 = affine.
Libanotis, Lam. Encyc. iii. 423 = latifolium.
lineatum, Tausch, in Flora, xiv. (1831) 667 = L. Siler.
Lobelii, Vill. Prosp. 24 = Carum rigidulum.
longiradium, *Boiss. Voy. Bot. Espagne*, ii. 734.—Hispan.
lucidum, [Soland. in] Ait. Hort. Kew. ed. I. i. 345 = Ligusticum simplex.
luteolum, Gaud. Fl. Helvet. ii. 348 = marginatum.
macrophyllum, *Borb. in Rad Jugos. Akad. Zagreb.* li. (1880) 53.—Europ.
marginatum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 210.—Europ. austr.
meifolium, Vent. ex DC. Prod. iv. 208 = Lophosciadum meifolium.
meoides, Desf. Fl. Atlant. i. 253 = Elaeoselinum meoides.
montanum, Lam. Fl. Fr. iii. 415 = L. Siler.
multifidum, *Kit. in Linnaea*, xxxii. (1863) 464.—Hungar.
multipartitum, Kit. l. c. = multifidum.
mutellinoides, Crantz, Cl. Umb. Emend. 67 = Ligusticum simplex.
nebrodense, Jan, ex DC. Prod. iv. 205 = L. Siler.
Nestleri, *Soyer-Willem. Obs.* 87.—Gallia.
nitidum, *Zanted. in Comm. Ateneo Brescia*, 1815 (1818) 89.—Europ. austr.
obliquum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 30 (Quid?).—Transsylv.
paludapifolium, Mill. Gard. Dict. ed. VIII. n. 3 = latifolium.
Panax, *Gouan, Illustr.* 13.—Europ.
petrophilum, Boiss. & Heldr. Diagn. Ser. I. x. 46 = Polylophium thalictroides.
peucedanoides, Brot. Fl. Lusit. i. 428 = Peucedanum lancifolium.
peucedanoides, Desf. Fl. Atlant. i. 254 = Elaeoselinum foetidum.
peucedanoides, *Linn. Cent. Pl.* ii. 13.—Europ.
pilosum, Comol. Prod. Fl. Com. 53 = nitidum.
pilosum, Willd. Enum. Hort. Berol. 310 = hispidum.
podolicum, *Rehm. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 497.—Europ.
polygamum, Lam. Encyc. iii. 425 = gummiferum.
pratense, *Ledeb. ex Sweet, Hort. Brit.* ed. II. 253 (Quid?).—Sibiria.
prutenicum, *Linn. Sp. Pl.* 248.—Europ.
prutenicum, Pall. ex Bieb. Casp. App. 209 = hispidum.
Pseudo-meum, *Orph. Heldr. & Sart. in Boiss. Diagn.* Ser. II. ii. 95.—Graec.
Pseudo-siler, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 30 = latifolium.
pubescens, Lag. Gen. et Sp. Nov. 12 = latifolium.
pubescens, Willd. ex Ledeb. Fl. Ross. ii. 329 = Zosimia orientalis.
resinosum, Presl, Delic. Prag. 137 = Ferula nudicaulis.
scabrum, *Cav. Ic.* ii. 72. t. 190.—Hispan.
selinoides, Crantz, Stirp. Austr. ed. I. fasc. ii. 182; ed. II. 182 = prutenicum.

LASERPITIUM:—
selinoides, Mill. Gard. Dict. ed. VIII. n. 6 = gallicum?
selinoides, Scop. Fl. Carn. ed. II. i. 198 = Selinum Carvifolia.
siculum, Spreng. Syst. i. 918 = L. Siler.
silaifolium, Jacq. Fl. Austr. v. 52. t. 44 = Carum rigidulum.
Siler, *Linn. Sp. Pl.* 249.—Europ.
Siler, Sibth. & Sm. Fl. Graec. Prod. i. 191 = garganicum.
simplex, Boeber, in Pall. N. Nord. Beitr. vi. 261 = Selinum sylvestre.
simplex, Lapeyr. Hist. Abr. Pl. Pyr. 152 = Meum pyrenaicum.
simplex, Linn. Mant. i. 56 = Ligusticum simplex.
simplex, Pourr. ex Nym. Consp. 291 = Ligusticum pyrenaicum.
Steveni, *Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop.* iv. 40.—Reg. Caucas.
striatum, Wall. Cat. n. 583 = Selinum striatum.
ternatum, Willd. ex Spreng. in Schult. Syst. vi. 555 = Oreosciadium montanum.
thapsiaeforme, Brot. Fl. Lusit. i. 427 = gummiferum.
thapsioides, Desf. Fl. Atlant. i. 252 = Elaeoselinum foetidum.
trifoliatum, Sieber, ex Spreng. in Schult. Syst. vi. 619 = L. Siler.
trifurcatum, Lam. Fl. Fr. iii. 416 = gallicum.
trilobum, Lapeyr. Hist. Abr. Pl. Pyr. 151 = Nestleri.
trilobum, Linn. Sp. Pl. 248 = Siler trilobum.
trilobum, Suter, Fl. Helvet. i. 163 = marginatum.
triquetrum, Vent. Jard. Cels, 97 = Prangos triquetra.
triternatisectum, Dulac, Fl. Hautes-Pyr. 358 = Nestleri.
ulopterum, *Trevir. in Linnaea*, x. (1836) *Litt.* 73.—Gallia.
verticillatum, Waldst. & Kit. Pl. Rar. Hung. ii. 186 = Trochiscanthes nodiflora.
Winkleri, *Herbich, Sel. Pl. Rar. Galic. et Bukov.* 11.—Austria.

LASIA, Lour. Fl. Cochinch. 81 (1790). *AROIDEAE*, Benth. & Hook. f. iii. 995.
LASIUS, Hassk. Cat. Hort. Bog. Alt. 59 (1844).
aculeata, *Lour. Fl. Cochinch.* 81.—As. trop.
desciscens, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 127 = aculeata.
Hermannii, Schott, in Bonplandia, v. (1857) 125 = heterophylla.
heterophylla, Schott, Melet. 21 = aculeata.
Jenkinsii, Schott, in Bonplandia, v. (1857) 125 = aculeata.
Loureirii, Schott, Melet. i. 21 = aculeata.
Merkusii, Hassk. Cat. Hort. Bog. Alt. 59 = Cyrtosperma Merkusii.
Roxburghii, Griff. Notul. iii. 155 = aculeata.
spinosa, Thw. Enum. Pl. Zeyl. 336 = aculeata.
stipitata, N. E. Br. in Gard. Chron. (1882) II. 70 = Podolasia stipitata.
stipulata, Hort. ex N. E. Br. l. c. 71, sphalm. = praec.
Zollingeri, Schott, in Bonplandia, v. (1857) 125 = aculeata.

LASIADENIA, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 632. *THYMELAEACEAE*, Benth. & Hook. f. iii. 192.
LINODENDRON, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 187.
cubensis, *Benth. & Hook. f. Gen.* iii. 192.—Cuba.
rupestris, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 632.—Guian.

LASIAGROSTIS, Link, Hort. Berol. i. 99 (1827) = **Stipa**, Linn. (Gramin.).
alpina, F. Schmidt, Reisen. Amurl. 73.—Reg. Amur.
australis, Zoll. & Mor. Syst. Verz. Zoll. 101 = S. australis.
Calamagrostris, Link, Hort. Berol. i. 99 = S. Calamagrostris.
capensis, Nees, Fl. Afr. Austr. 167 = Trachypogon polymorphus.
Caragana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 90 = S. Caragana.
elongata, Nees, l. c. = S. elongata.
gigantea, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 96 = S. gigantea.
Jacquemontii, Munro, ex Boiss. Fl. Orient. v. 506 = S. Jacquemontii.

LASIAGROSTIS:—
mongholica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 87 = Stipa mongolica.
splendens, Kunth, Rév. Gram. i. 58 = Muehlenbergia alpestris.
subsessiliflora, Rupr. Sert. Tiansch. 35.—Sibir.
tenacissima, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 94 = S. tenacissima.
tremula, Rupr. Sert. Tiansch. 35.—Sibir.
variegata, G. F. W. Mey. ex Steud. Nom. ed. II. ii. 11 = Melica ciliata.

LASIAKE, Rafin. Fl. Tellur. iv. 89 (1836) = **Verbascum**, Tourn. (Scrophul.).
lyratum, Rafin. l. c. = Celsia cretica.
plicatum, Rafin. l. c. = V. plicatum.

LASIANDRA, DC. Prod. iii. 127 (1828) = **Tibouchina**, Aubl. (Melastom.).
adenostemon, DC. l. c. 130 = T. adenostemon.
adenostemon, Miq. in Linnaea, xxii. (1849) 539 = T. multiflora.
aegopogon, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 153 = T. aegopogon.
aemula, DC. Prod. iii. 128 = T. aemula.
albiflora, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 159 = T. hospita.
andina, Poepp. ex Naud. l. c. 149 = T. ochypetala.
angustifolia, Naud. l. c. 135 = T. angustifolia.
annularis, Naud. l. c. 138 = T. setulosa.
arborea, Naud. l. c. 159 = T. arborea.
argentea, DC. Prod. iii. 130 = T. holosericea.
aspera, Salzm. ex Triana, in Trans. Linn. Soc. xxviii. (1873) 44 = T. sulcata.
asperior, Cham. in Linnaea, ix. (1834) 435 = T. asperior.
asperior, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 132 = T. intermedia.
barbigera, Naud. l. c. 155 = T. barbigera.
Benthamiana, Naud. l. c. 159 = T. Benthamiana.
bipenicillata, Naud. l. c. 156 = T. bipenicillata.
calcarata, Naud. l. c. 127 = Desmoscelis calcarata.
calyptrata, Bunbury, in Proc. Linn. Soc. i. (1841) 108 = T. canescens.
Candolleana, DC. Prod. iii. 129 = T. Candolleana.
cardinalis, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 132 = T. cardinalis.
Claussenii, Naud. l. c. 152 = T. Claussenii.
coerulea, Reichb. Hort. Bot. iii. 18. t. 250 = T. viminea.
collina, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 148 = T. collina.
collina, Salzm. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. Lhotskyana.
cordiformis, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 143 = T. Lhotzkyana.
corymbosa, Naud. l. c. 159 = T. Lindeniana.
dendroides, Naud. l. c. 132 = T. dendroides.
dimorphandra, Miq. in Linnaea, xxii. (1849) 538 = T. Fothergillae.
Diplostegium, Cham. in Linnaea, ix. (1834) 438 = T. canescens.
dubia, Cham. l. c. 437 = T. dubia.
echinata, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 159 = T. Gardneriana.
elegans, Naud. l. c. 159 = T. elegans.
Endlicheri, Beer & Fenzl, ex Seem. in Harting. Parad. Vindob. 47. t. 78 = Monochaetina Humboldtiana.
exasperata, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 154 = T. exasperata.
fissinervia, DC. Prod. iii. 130 = T. fissinervia.
Fontanesiana, DC. l. c. = T. granulosa.
Fontanesiana, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 150 = T. Langsdorffiana.
Fothergillae, Cham. in Linnaea, ix. (1834) 437 = T. Fothergillae.
foveolata, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 148 = T. foveolata.
frigidula, DC. Prod. iii. 127 = T. frigidula.
Gardneri, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 140 = T. Gardneri.
Gaudichaudiana, DC. Prod. iii. 128 = T. Gaudichaudiana.
Gaudichaudiana, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 136 = T. corymbosa.
gracilis, Naud. l. c. 128 = T. gracilis.
heteromalla, Naud. l. c. 158 = T. heteromalla.
hieracioides, Naud. l. c. 127 = T. hieracioides.
hirsuta, Naud. l. c. 159 = T. gracilis.
holosericea, Naud. l. c. 139 = T. villosissima.

LASIANDRA:—
hospita, DC. Prod. iii. 129 = Tibouchina hospita.
hydrophila, DC. l. c. 128 = T. hygrophila.
Imperatoris, Wawra, in Oestr. Bot. Zeitschr. xiii. (1863) 87 = T. elegans.
Kunthiana, DC. Prod. iii. 128 = T. Moricandiana.
lanceolata, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 140 = T. Candolleana.
Langsdorffiana, DC. Prod. iii. 130 = T. Langsdorffiana.
laxa, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 159 (= *Pleroma laxum*).—Peruv.
ledifolia, Naud. l. c. = Brachyotum ledifolium.
lepidota, Naud. l. c. 153 = T. ochypetala.
Lhotzskyana, Presl, Symb. Bot. ii. 2 = T. Lhotzkyana.
longistyla, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 143 = T. urceolaris.
lutescens, Naud. l. c. 137 = T. Gaudichaudiana.
macrantha, Linden & Seem. in Journ. Bot. ii. (1864) 361 = T. semidecandra.
macrantha, Rich. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 44 = T. Candolleana.
macrantha, Vukot. in Rad Jugos. Akad. Zagreb. xxvii. (1874) 212 (Lasyandra) (Quid?).—Hab.?
macrocarpa, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 152 = T. semidecandra.
macrochiton, DC. Prod. iii. 129 = T. macrochiton.
macrophylla, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 140 = T. heteromalla.
martialis, Cham. in Linnaea, ix. (1834) 433 = T. martialis.
martialis, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 134 = T. Lindeniana.
Martiana, Cham. in Linnaea, ix. (1834) 431 = T. Martiusiana.
Martiusiana, Cham. l. c. 431 = T. frigidula.
Martiusiana, DC. Prod. iii. 127 = T. Martiusiana.
Matthaei, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 142 = T. ochypetala.
Maximiliana, DC. Prod. iii. 128 = T. Maximiliana.
Maximiliana, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 138 = T. Gaudichaudiana.
melastomoides, Naud. l. c. 154 = T. melastomoides.
mollis, Cham. in Linnaea, ix. (1834) 444 = T. Chamissoana.
Moricandiana, DC. Prod. iii. 128 = T. Moricandiana.
mucorifera, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 141 = T. Fothergillae.
multiceps, Naud. l. c. 149 = T. multiceps.
multiflora, Naud. l. c. 159 = T. multiflora.
muricata, Presl, Symb. Bot. i. 68 = T. estrellensis.
mutabilis, Riedel, ex Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 145 = T. mutabilis.
nigrescens, Naud. l. c. 147 = T. trichopoda.
nudicaulis, Naud. l. c. 128 = T. hieracioides.
obscura, Cham. in Linnaea, ix. (1834) 435 = T. corymbosa.
ochypetala, DC. Prod. iii. 128 = T. ochypetala.
oleaefolia, Mart. Nov. Gen. et Sp. iii. 97. t. 244 = Microlepis oleaefolia.
osbeckioides, Steud. in Flora, xxvii. (1844) II. 720 = T. Gaudichaudiana.
oxypetala, Walp. Ann. ii. 561 = *ochypetala*.
paniculata, Rich. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. adenostemon.
papyrifera, Pohl, ex Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 157 = T. papyrifera.
petiolata, R. Grah. in Hook. Bot. Mag. t. 3766 = T. Gaudichaudiana.
phalacrostemon, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 137 = T. Gaudichaudiana.
pleromoides, Naud. l. c. 131 (= *Pleroma mauricarpum*).—Peruv.
pogonanthera, Naud. l. c. 153 = T. pogonanthera.
proteaeformis, DC. Prod. iii. 130 = T. holosericea.
pulchella, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 129 = T. gracilis.
pulchra, Cham. in Linnaea, ix. (1834) 439 = T. pulchra.
Raddiana, DC. Prod. iii. 129 = T. Raddiana.
Regnellii, Miq. in Linnaea, xxii. (1849) 539 = T. martialis.
Riedelii, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 135 = T. martialis.
rigidula, Naud. l. c. 150 = T. aemula.
salviaefolia, Cham. in Linnaea, ix. (1834) 441 = T. salviaefolia.
salviaefolia, Miq. l. c. xxii. (1849) 538 = T. stenocarpa.

LASIANDRA :—
sarmentosa, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 130 (=*Pleroma laxum*).—Peruv.
Sellowiana, Cham. in Linnaea, ix. (1834) 440 = Tibouchina Sellowiana.
semidecandra, Schrank & Mart. ex DC. Prod. iii. 129 = T. semidecandra.
sericans, Miq. in Linnaea, xxii. (1849) 538 = T. stenocarpa.
sericea, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 144 = T. Lhotzkyana.
setulosa, Spring, ex Mart. in Flora, xxii. (1839) 1. Beibl. 9 = T. setulosa.
spathularis, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 147 = T. macrochiton.
spoliata, Naud. l. c. 137 = T. Gaudichaudiana.
stenocarpa, DC. Prod. iii. 130 = T. stenocarpa.
strigillosa, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 158 = T. gracilis.
sulcata, Naud. l. c. 137 = T. sulcata.
terminalis, Naud. l. c. 142 = T. Moricandiana.
tetraptera, Naud. l. c. 133 = T. Benthamiana.
Thereminiana, DC. Prod. iii. 129 = T. Thereminiana.
tibouchinoides, Cham. in Linnaea, ix. (1834) 432 = T. multiceps.
tibouchinoides, DC. Prod. iii. 128 = T. trichopoda.
Tibuchina, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 153 = T. aspera.
Trichopoda, Naud. l. c. 159 = T. trichopoda.
trifolia, DC. Prod. iii. 127 = T. frigidula.
urceolaris, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 130 = T. australis.
ursina, Cham. in Linnaea, ix. (1834) 443 = T. ursina.
Urvilleana, DC. Prod. iii. 130 = T. Urvilleana.
velutina, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 144 = T. velutina.
velutina, Pohl, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. urceolaris.
versicolor, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 146 = T. Raddiana.
verticillaris, Naud. l. c. 134 = T. frigidula.
villosa, Naud. l. c. 128 = T. ursina.
villosissima, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. villosissima.
viminea, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 159 = T. viminea.
virgata, Naud. l. c. 159 = T. virgata.
Weddellii, Naud. l. c. 151 = T. Weddellii.

LASIANDROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = praec.

LASIANTHAEA, DC. Prod. v. 607 (1836) = **Zexmenia**, La Llave (Compos.).
helianthoides, DC. l. c. 608 = Z. helianthoides, *A. Gray*.

LASIANTHEA, Endl. Ench. 239 (1841) = praec.

LASIANTHEMUM, Klotzsch, in Schomb. Faun. et Fl. Brit. Gui. 1181 (1848), nomen. *SAPINDACEAE*.
bijugum, *Klotzsch, l. c.* nomen.—Guiana.
unijugum, *Klotzsch, l. c.* nomen.—Guiana.

LASIANTHERA, Beauv. Fl. Owar. i. 85. t. 51 (1805). *OLACINEAE*, Benth. & Hook. f. i. 350.
MEDUSANTHERA, Seem. in Journ. Bot. ii. (1864) 74.
STEMONURUS, Blume, Bijdr. 648 (1825), partim.
URANDRA, Thw. in Hook. Kew Journ. vii. (1855) 211.
africana, *Beauv. Fl. Owar.* i. 85. t. 51.—Afr. trop.
apicalis, *Benth. & Hook. f. Gen.* i. 350.—Zeylan.
australiana, F. Muell. Fragm. vi. 3 = Gomphandra australiana.
austro-caledonica, *Baill. Adansonia*, xi. (1873–76) 193.—N. Caled.
capitulata, *Miq. Fl. Ind. Bat.* i. 1. 791.—Sumatra.
javanica, *Miq. l. c.* 790.—Malaya.
lanceolata, *Mast. in Hook. f. Fl. Brit. Ind.* i. 585.—Malacca.
littoralis, *Miq. Fl. Ind. Bat.* i. 1. 792.—N. Guin.
macrocarpa, *Miq. l. c.*—Borneo.
macrophylla, *Miq. l. c.*—Malaya.
Maingayi, *Mast. in Hook. f. Fl. Brit. Ind.* i. 585.—Malacca.
membranacea, *Miq. Fl. Ind. Bat.* i. 1. 792.—Java.
ovalifolia, *Miq. l. c. Suppl.* 342.—Sumatra.
papuana, *Becc. Malesia*, i. (1877) 108.—N. Guin.

LASIANTHERA :—
parviflora, *Miq. Fl. Ind. Bat.* i. 1. 791.—Sumatra.
prasina, Miq. l. c. = Gomphandra affinis.
secundiflora, Miq. l. c. Suppl. 342. t. 2 = apicalis.
tetrandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 327 = Gomphandra axillaris.

LASIANTHUS, Gronov. ex Linn. Syst. ed. I (1735) = **Gordonia**, Ellis (Ternstroem.).

LASIANTHUS, Jack, in Trans. Linn. Soc. xiv. (1823) 125. *RUBIACEAE*, Benth. & Hook. f. ii. 129.
MEPHITIDIA, Reinw. ex Blume, Cat. Gew. Buitenz. 51 (1823).
NOVATILEA, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 503.
OCTAVIA, DC. Prod. iv. 464 (1830).
SANTIA, Wight & Arn. Prod. 422 (1834).
acuminatus, Wight, ex Hook. f. Fl. Brit. Ind. iii. 185 = strigillosus.
acuminatus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 511.—Ind. or.
acutatus, Miq. Fl. Ind. Bat. Suppl. 549 = mucronulatus.
africanus, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 228.—Afr. trop.
andamanicus, *Hook. f. Fl. Brit. Ind.* iii. 189.—Burma.
apiocarpus, *Miq. Fl. Ind. Bat.* ii. 318.—Sumatra.
appressus, *Hook. f. Fl. Brit. Ind.* iii. 181.—Malaya.
attenuatus, *Jack, in Trans. Linn. Soc.* xiv. (1823) 126.—Malaya.
Biermanni, *King, ex Hook. f. Fl. Brit. Ind.* iii. 190.—Reg. Himal.
Blumeanus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 507.—Ind. or.
Boivinianus, *Baill. Adansonia*, xii. (1876–79) 232.—Ins. Comor.
bracteatus, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 501 = cyanocarpus.
bracteolatus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 246.—Sumatra.
capitatus, *Blume, Bijdr.* 996.—Java.
capitulatus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 511.—Ind. or.
chinensis, *Benth. Fl. Hongk.* 160.—Malaya; China.
chrysoneurus, *Miq. Fl. Ind. Bat.* ii. 322.—Java.
chrysophyllus, *Miq. l. c.*—Sumatra.
ciliatus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 509.—Ind. or.
cinnamomoides, Miq. Fl. Ind. Bat. ii. 325 = reticulatus.
constrictus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 515.—Burma.
crinitus, *Hook. f. Fl. Brit. Ind.* iii. 182.—Malaya.
cyanocarpus, *Jack, in Trans. Linn. Soc.* xiv. (1823) 125.—Sumatra.
densifolius, *Miq. Fl. Ind. Bat.* ii. 321.—Malaya.
dichotomus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 508.—Ind. or.
dubius, Wight, l. c. 516 = Psychotria Wightiana.
ellipticus, *Wight, l. c.* 507.—Malacca.
firmus, *Miq. Fl. Ind. Bat. Suppl.* 549.—Sumatra.
foetens, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 517 = Saprosma ceylanicum.
Fordii, *Hance, in Journ. Bot.* xxiii. (1885) 324.—China.
Gardneri, *Hook. f. Fl. Brit. Ind.* iii. 186.—Zeylan.
glabrescens, *Miq. Fl. Ind. Bat.* ii. 317.—Java.
Griffithii, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 505.—Malacca.
Helferi, *Hook. f. Fl. Brit. Ind.* iii. 189.—Burma.
hexandrus, *Blume, Bijdr.* 998.—Java.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 184.—Reg. Himal.
inaequalis, Blume, Bijdr. 996 = cyanocarpus.
inconspicuus, *Hook. f. Fl. Brit. Ind.* iii. 187.—Reg. Himal.
inodorus, *Blume, Bijdr.* 998.—Java.
iteophyllus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 247.—Java.
Jackianus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 502.—Ind. or.
japonicus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 110.—Japon.
Kurzii, *Hook. f. Fl. Brit. Ind.* iii. 183.—Burma.
laevicaulis, Kurz, in Journ. Bot. xiii. (1875) 327 = cyanocarpus.
laevigatus, *Blume, Bijdr.* 998.—Java.

LASIANTHUS :—
lancifolius, *Hook. f. Fl. Brit. Ind.* iii. 187.—Reg. Himal.
latifolius, *Blume, ex Miq. Fl. Ind. Bat.* ii. 320.—Java.
longicauda, *Hook. f. Fl. Brit. Ind.* iii. 190.—Reg. Himal.
longifolius, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 514.—Malacca.
lucidus, *Blume, Bijdr.* 997.—Java.
lucidus, Zoll. ex Miq. Fl. Ind. Bat. ii. 322 = rigidus.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 188.—Malaya.
micranthus, *Hook. f. l. c.* 190.—Reg. Himal.
Moonii, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 504.—Zeylan.
Moralesii, *Wright, in Sauv. Fl. Cub.* 67.—Cuba.
mucronulatus, *Miq. Fl. Ind. Bat.* ii. 325.—Borneo.
obliquus, *Thw. Enum. Pl. Zeyl.* 420.—Zeylan.
oblongifolius, *Bedd. Ic. Pl. Ind. Or.* i. 2.—Ind. or.
obovatus, *Bedd. l. c.*—Ind. or.
obscurus, *Blume, ex Miq. Fl. Ind. Bat.* ii. 317.—Java.
Oculus-cati, Miq. l. c. 315 = cyanocarpus.
okelensis, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 122 = Tricalysia okelensis.
oliganthus, *Thw. Enum. Pl. Zeyl.* 420.—Zeylan.
ovatus, *Miq. Fl. Ind. Bat.* ii. 321.—Sumatra.
palembanicus, *Miq. Fl. Ind. Bat. Suppl.* 549.—Sumatra.
parvifolius, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 512.—Ind. or.
pauciflorus, *Wight, l. c.* 515.—Burma.
pilosus, *Wight, l. c.* 506.—Malacca.
plagiophyllus, Hance, in Journ. Bot. xiii. (1875) 196 = Wallichii.
platyphyllus, *Miq. Fl. Ind. Bat.* ii. 318.—Sumatra.
polycarpus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 247.—Borneo.
protractus, Thw. Enum. Pl. Zeyl. 420 = strigosus.
pterospermus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 510.—Malacca.
purpureus, *Blume, Bijdr.* 1000.—Java.
reticulatus, *Blume, l. c.*—Java.
retosus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 506.—Malacca.
rhinophyllus, *Thw. Enum. Pl. Zeyl.* 420.—Zeylan.
rhinozerotis, *Blume, Bijdr.* 996.—Java.
rhizophyllus, Hook. f. Fl. Brit. Ind. iii. 185, sphalm. = rhinophyllus.
rigidus, *Miq. Fl. Ind. Bat.* ii. 321.—Java.
rostratus, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 510 = ciliatus.
Roxburghii, Wight, l. c. 502 = cyanocarpus.
rufus, *Miq. Fl. Ind. Bat.* ii. 325.—Java.
sikkimensis, *Hook. f. Fl. Brit. Ind.* iii. 180.—Reg. Himal.
singalensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 247.—Sumatra.
spathulatus, *Miq. l. c.* 246.—Borneo.
stercorarius, *Blume, Bijdr.* 1000.—Java.
stercorarius, Kurz, For. Fl. Br. Burm. ii. 31 = Kurzii.
stipularis, *Blume, Bijdr.* 997.—Burma; Malaya.
strigillosus, *Hook. f. Fl. Brit. Ind.* iii. 185.—Ind. or.
strigosus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 512.—Zeylan.
subsessilis, *Miq. Fl. Ind. Bat. Suppl.* 550.—Sumatra.
sylvestris, *Blume, Bijdr.* 999.—Java.
tentaculatus, *Hook. f. Fl. Brit. Ind.* iii. 181.—Reg. Himal.
Thwaitesii, *Hook. f. l. c.* 185.—Zeylan.
tobingensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 249.—Sumatra.
tomentosus, *Blume, Bijdr.* 997.—Java.
tomentosus, Zoll. ex Miq. Fl. Ind. Bat. ii. 317 = obscurus.
truncatus, *Bedd. Ic. Pl. Ind. Or.* i. 2.—Ind. or.
tubiferus, *Hook. f. Fl. Brit. Ind.* iii. 183.—Reg. Himal.
tubiflorus, Blume, Bijdr. 999 = Hamiltonia suaveolens.
varians, *Thw. Enum. Pl. Zeyl.* 420.—Zeylan.
venosus, *Blume, Bijdr.* 999.—Java.
venosus, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 514 = Wightianus.
venulosus, *Wight, l. c.* 508.—Ind. or.
Vriesianus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 249.—Borneo.
Walkerianus, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 513.—Zeylan.
Wallichii, *Wight, l. c.* 503.—Reg. Himal.; Burma.
Wightianus, *Hook. f. Fl. Brit. Ind.* iii. 188.—Malacca.

LASIANTHUS, Zucc. ex DC. Prod. v. 607 (1836) = **Zexmenia**, La Llave (Compos.).

LASIERPA, Torr. in Geol. Rep. N. York (1839), 152 = **Chiogenes**, Salisb. (Oleac.).
hispidula, Torr. l. c. = C. serpyllifolia.

LASIMORPHA, Schott, in Bonplandia, v. (1857) 127 = **Cyrtosperma**, Griff. (Aroid.).
Afzelii, Schott, Gen. Aroid. t. 85. f. 11 = C. Afzelii.
senegalensis, Schott, in Bonplandia, v. (1857) 127 = C. senegalense.

LASINEMA, Steud. Nom. ed. II. ii. 85, in syn. (1841) sphalm. = **Lysinema**, R. Br. (Epacrid.).

LASINGROSTIS, Link, Handb. i. 39 (1829), sphalm. = LASIAGROSTIS, Link = **Stipa**, Linn. (Gramin.).

LASINIA, Rafin. New Fl. Am. ii. 48 (1836) = **Baptisia**, Vent. (Legumin.).
bicolor, Rafin. l. c. 49 = Thermopsis mollis.
bracteata, Rafin. l. c. 50 = B. leucophaea.
cinerea, Rafin. l. c. = B. villosa.
fulva, Rafin. l. c. 49 = B. villosa.
grandifolia, Rafin. l. c. = Thermopsis mollis.
mollis, Rafin. l. c. 50 = Thermopsis mollis.
pallens, Rafin. l. c. 49 = Thermopsis mollis.
reticulata, Rafin. l. c. 48 = B. lanceolata.

LASIOBEMA, Miq. Fl. Ind. Bat. i. 1. 71 (1855) = **Bauhinia**, Linn. (Legumin.).
anguina, Miq. l. c. = B. anguina.
Horsfieldii, Miq. l. c. = B. anguina.

LASIOCARPHUS, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 138 (1873) = **Stilpnopappus**, Mart. (Compos.).
ferrugineus, Pohl, ex Baker, l. c. = S. ferrugineus.

LASIOCARPUS, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 264 (1847) = **Acaena**, Vahl (Rosac.).
humilis, Banks & Soland. ex Hook. f. l. c. = A. pumila.

LASIOCARPUS, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 90. *MALPIGHIACEAE*, Benth. & Hook. f. i. 255.
salicifolius, *Liebm. l. c.*—Mexic.

LASIOCEPHALUS, Willd. ex Schlecht. in Ges. Nat. Fr. Berl. Mag. viii. (1814) 308 = **Culcitium**, Humb. & Bonpl. (Compos.).
lingulatus, Schlecht. l. c. 309.—Am. mer.
ovatus, Schlecht. l. c.—Am. mer.

LASIOCHLOA, Kunth, Rév. Gram. ii. 555. t. 192 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1194. TRIBOLIUM, Desv. Opusc. 64 (1831).
adscendens, *Kunth, Enum. Pl.* i. 388.—Afr. austr.
ciliaris, Kunth, Rév. Gram. ii. 555. t. 192, excl. syn. = adscendens.
ciliaris, *Nees, Fl. Afr. Austr.* 432.—Afr. austr.
hirta, *Kunth, Rév. Gram.* ii. 559. t. 194.—Afr. austr.
hispida, *Kunth, Enum. Pl.* i. 388.—Afr. austr.
laevis, *Kunth, l. c.*—Afr. austr.
longifolia, *Kunth, Rév. Gram.* ii. 557. t. 193.—Afr. austr.
obtusifolia, *Nees, Fl. Afr. Austr.* 430.—Afr. austr.
ovata, *Nees, l. c.* 431.—Afr. austr.
pectinata, *Trin. Sp. Gram.* i. t. 73; *ex Pritz. Ic. Ind.* i. 620.—Hab.?
planifolia, Trin. ex Steud. Nom. ed. II. ii. 12 (Quid?).—Hab.?
serrata, *Kunth, Enum. Pl.* i. 388.—Afr. austr.
utriculosa, *Nees, Fl. Afr. Austr.* 436.—Afr. austr.
villosa, *Kunth, Enum. Pl.* i. 388.—Afr. austr.

LASIOCLADUS, Boj. ex Nees, in DC. Prod. xi. 510 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1122.
anthospermifolius, *Boj. ex Nees, in DC. Prod.* xi. 510.—Madag.
boerhaaviaefolius, Boj. l. c. = Adhatoda boerhaviifolia.
chlorotrichos, *Boj. l. c.*—Madag.
spicatus, Boj. l. c. = Adhatoda spicata.

LASIOCORYS, Benth. Lab. Gen. et Sp. 600 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1213.
abyssinica, *Benth. l. c.*—Abyssin.
arabica, *Jaub. & Spach, Illustr.* iv. 126.—Arab.; Afr. trop.
capensis, *Benth. Lab. Gen. et Sp.* 600.—Afr. austr.

LASIOCORYS:—
hyssopifolia, *Franch. in Révoil, Çomali*, 57.—Afr. trop.
flagellifera, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotr.
royleoides, Vatke, in Linnaea, xliii. (1880–82) 98 = arabica.
spiculifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotr.
stachydiformis, *Benth. in DC. Prod.* xii. 534.—Abyssin.

LASIOCROTON, Griseb. Fl. Brit. W. Ind. 46 (1859). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 317.
macrophyllus, *Griseb. l. c.*—Jamaic.
prunifolius, *Griseb. in Goett. Nachr.* (1865) 175 (al. gen. ?).—Cuba.

LASIODISCUS, Hook. f. in Benth. & Hook. f. Gen. i. 381 (1862). *RHAMNEAE*, Benth. & Hook. f. l. c.
Mannii, *Hook. f. l. c.*—Afr. trop.
Pervillei, *Baill. Adansonia*, viii. (1867–68) 202.—Madag.

LASIOGYNE, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 418 (Pl. Meyen.) = **Croton**, Linn. (Euphorb.).
brasiliensis, Klotzsch, l. c. = C. compressus.
phlomoides, Griseb. in Goett. Abh. vii. (1857) 171 = C. astroites.
Pottsii, Klotzsch, in Seem. Bot. Voy. Herald, 278 = C. Pottsii.

LASIOLEPIS, Benn. Pl. Jav. Rar. 202. t. 42 (1838) = **Harrisonia**, R. Br. (Rutac.).
Bennetii, Planch. in Hook. Lond. Journ. Bot. v. (1846) 570 = H. paucijuga.
multijuga, Benn. Pl. Jav. Rar. 202. t. 42 = H. paucijuga.
paucijuga, Benn. l. c. = H. paucijuga.

LASIOLEPIS, Boeck. in Flora, lvi. (1873) 90 = **Eriocaulon**, Linn.
aquatica, Boeck. l. c. 91 = E. melanocephalum.
brevifolia, Boeck. l. c. 90.—Ind. or.
pilosa, Boeck. l. c. 91 = Paepalanthus Lamarckii.

LASIOLYTRUM, Steud. in Flora, xxix. (1846) 18 = **Arthraxon**, Beauv. (Gramin.).
hispidum, Steud. l. c. = A. ciliaris.
pilosum, Steud. ex Jord. in Mém. Soc. Sc. Nat. Cherb. v. (1857) 299 = A. ciliaris.

LASIONEMA, D. Don, in Lond. & Edinb. Phil. Mag. ii. (1833) 377; et in Trans. Linn. Soc. xvii. (1834) 141 = **Macrocnemum**, R. Br. (Rubiac.).
cinchonoides, Wedd. Hist. Nat. Quin. 99 = M. cinchonoides.
dubium, Didr. in Kjoeb. Vidensk. Meddel. (1854) 188.—Bras.
glabrescens, Benth. Bot. Voy. Sulph. 105 = M. glabrescens.
grandiflorum, Wedd. Hist. Nat. Quin. 98 = M. grandiflorum.
Humboldtianum, Wedd. l. c. 99 = M. Humboldtianum.
pubescens, Benth. Pl. Hartw. 191 = M. pubescens.
roseum, D. Don, in Lond. & Edinb. Phil. Mag. ii. (1833) 377 = M. roseum.

LASIOPERA, Hoffmgg. & Link, Fl. Port. i. 298 (1809) = **Bartsia**, Linn. (Scrophular.).
aspera, Hoffmgg. & Link, l. c. 302. t. 59 = B. aspera.
Bocconi, Presl, Fl. Sic. i. 34 = Euphrasia Bocconi.
frutescens, Presl, Bot. Bemerk. 91 = Euphrasia Bocconi.
rhinanthina, Hoffmgg. & Link, Fl. Port. i. 299. t. 58 = B. Trixago.
rigidifolia, Presl, Fl. Sic. i. 34 = Euphrasia rigidifolia.
squamosa, Presl, Bot. Bemerk. 91 = B. purpurea.
tenuifolia, Hoffmgg. & Link, Fl. Port. i. 303. t. 60 = Euphrasia tenuifolia.
viscosa, Hoffmgg. & Link, l. c. 301. t. 61 = B. viscosa.

LASIOPETALUM, Sm. in Trans. Linn. Soc. iv. (1798) 216. *STERCULIACEAE*, Benth. & Hook. f. i. 228.
ASTEROCHITON, Turcz. in Bull. Soc. Nat. Mosc. (1852) II. 138.
CORETHROSTYLIS, Endl. Nov. Stirp. Dec. 1 (1839).
acutiflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 145.—Austral.

LASIOPETALUM:—
arborescens, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 36 = Seringia platyphylla.
Baueri, *Steetz, in Lehm. Pl. Preiss.* ii. 339.—Austral.
Behrii, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 36.—Austral.
bracteatum, *Benth. Fl. Austral.* i. 266.—Austral.
capitellatum, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 148 = discolor.
confertiflorum, F. Muell. in Linnaea, xxv. (1852) 377 = discolor.
cordifolium, *Endl. in Enum. Pl. Hueg.* 10.—Austral.
dasyphyllum, *Sieber, ex Hook. Journ. Bot.* ii. (1840) 414.—Austral.
discolor, *Hook. Comp. Bot. Mag.* i. (1835) 276.—Austral.
Drummondii, *Benth. Fl. Austral.* i. 264.—Austral.
dumosum, Hort. ex Steetz, in Lehm. Pl. Preiss. ii. 356 = Rulingia parviflora.
dumosum, Lodd. Bot. Cab. t. 1564 = Rulingia hermanniaefolia.
ferrugineum, *Sm. in Andr. Bot. Rep.* t. 208.—Austral.
Fitzgibbonii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 72.—Austral.
floribundum, *Benth. Fl. Austral.* i. 265.—Austral.
glutinosum, *F. Muell. Fragm.* xi. 113.—Austral.
Gunnii, Steetz, in Lehm. Pl. Preiss. ii. 342 = dasyphyllum.
indutum, *Steud. l. c.* i. 235.—Austral.
laxiflorum, *F. Muell. Fragm.* xi. 112.—Austral.
ledifolium, Vent. Jard. Malm. sub n. 59 = Boronia ledifolia.
macrophyllum, *R. Grah. in Bot. Mag.* t. 3908.—Austral.
Maxwelli, *F. Muell. Fragm.* xi. 107.—Austral.
membranaceum, *Benth. Fl. Austral.* i. 266.—Austral.
micranthum, *Hook. f. Fl. Tasm.* i. 51.—Tasman.
molle, *Benth. Fl. Austral.* i. 266.—Austral.
Ogilvieanum, *F. Muell. Fragm.* xi. 107.—Austral.
Oldfieldii, F. Muell. l. c. ii. 6 = acutiflorum.
oppositifolium, *F. Muell. l. c.* 5.—Austral.
parviflorum, *Rudge, in Trans. Linn. Soc.* x. (1811) 297. t. 12. f. 2.—Austral.
parvuliflorum, *F. Muell. Fragm.* vi. 174.—Austral.
prostratum, Hort. ex Steetz, in Lehm. Pl. Preiss. ii. 356 = Rulingia parviflora.
purpureum, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 36 = Thomasia purpurea.
pygmaeum, *F. Muell. Census*, 16.—Austral.
quercifolium, Andr. Bot. Rep. t. 459 = Thomasia quercifolia.
quinquenervium, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 146 = acutiflorum.
rosmarinifolium, *Benth. Fl. Austral.* i. 264.—Austral.
rubiginosum, A. Cunn. in Field, N. S. Wales, 354 = ferrugineum.
rufum, *R. Br. ex Benth. Fl. Austral.* i. 263.—Austral.
Schulzenii, *Benth. l. c.* 265.—Austral.
Sieberi, Steetz, in Lehm. Pl. Preiss. ii. 338 = ferrugineum.
solanaceum, Sims, Bot. Mag. t. 1486 = Thomasia solanacea.
stelligerum, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 147 = Thomasia stelligera.
Tepperi, *F. Muell. Fragm.* xi. 109.—Austral.
tomentosum, Cels, ex J. Gay, in Mém. Mus. Par. x. (1823) 200 = Rulingia pannosa.
triphyllum, Labill. Nov. Holl. Pl. i. 63. t. 88 = Thomasia triphylla.
Wilhelmii, F. Muell. in Trans. Phil. Soc. Vict. ii. (1858) 65 = dasyphyllum.

LASIOPHYTON, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 44 = **Micropsis**, DC. (Compos.).
pusillum, Hook. & Arn. l. c. = M. nana.

LASIOPOA, Ehrh. Beitr. iv. 147 (1789) = **Bromus**, Linn. (Gramin.).

LASIOPOGON, Cass. in Bull. Soc. Philom. (1818) 75. *COMPOSITAE*, Benth. & Hook. f. ii. 304.
lanatum, *Cass. l. c.*—Reg. Mediterr.; Persia; Afr. austr.
micropoides, *DC. Prod.* vi. 246.—Afr. austr.
molluginoides, DC. l. c. = micropoides.
muscoides, DC. l. c. = lanatum.

LASIOPTERA, Andrz. ex DC. Syst. ii. 534 (1821) = **Lepidium**, Tourn. (Crucif.).
campestris, Andrz. ex DC. l. c. 535 = L. campestre.
hirta, Andrz. ex DC. l. c. 536 = L. hirtum.

LASIOPUS, Cass. in Bull. Soc. Philom. (1817) 152 = **Gerbera**, Gronov. (Compos.).
ambiguus, Cass. in Dict. Sc. Nat. xxv. 299 = G. ambigua.
Bojeri, DC. Prod. vii. 19 = G. Bojeri.
Candollei, DC. l. c. = G. Candollei.
coriaceus, DC. l. c. = G. ambigua.
viridifolius, DC. l. c. = G. viridifolia.

LASIOPUS, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 346 (1836) = **Taraxacum**, Hall. (Compos.).
sonchoides, D. Don, l. c. = T. montanum.

LASIORRHIZA, Lag. Amen. Nat. i. 32 (1811) = **Leuceria**, Lag. (Compos.).
ceterachifolia, Cass. in Dict. Sc. Nat. xliii. 80 = Leuceria gossypina.
cinerea, Less. Syn. Comp. 404 = Leuceria cinerea.
glacialis, Poepp. ex Less. l. c. 407 = Leuceria pulchella.
lithospermifolia, Less. l. c. 406 (= *Chabraea lithospermifolia*).—Chili.
pulchella, Steud. Nom. ed. II. ii. 12 = Leuceria pulchella.
purpurea, Less. in Linnaea, v. (1830) 11 = Leuceria purpurea.
rosea, Less. Syn. Comp. 405 = Leuceria runcinata.
tomentosa, Less. in Linnaea, v. (1830) 12 = Leuceria Bridgesii.
viscosa, Cass. in Dict. Sc. Nat. xliii. 80 = Leuceria gossypina.

LASIOSIPHON, Fresen. in Flora, xx. (1838) 602. *THYMELAEACEAE*, Benth. & Hook. f. iii. 197.
affinis, *Kotschy & Peyr. Pl. Tinn.* 39. t. 19 B.—Afr. trop.
anthylloides, *Meissn. in DC. Prod.* xiv. 595.—Afr. austr.
Bojerianus, *Decne. in Jacquem. Voy. Bot.* 149.—Madag.
Burchellii, *Meissn. in DC. Prod.* xiv. 594.—Afr. austr.
caffer, *Meissn. l. c.* 593.—Afr. austr.
cuneatus, *Decne. in Jacquem. Voy. Bot.* 149.—Madag.
Dregeanus, *Endl. Gen. Suppl.* iv. II. 67.—Afr. austr.
eriocephalus, *Decne. in Jacquem. Voy. Bot.* 148.—Ind. or.
glaucus, *Fresen. in Flora*, xxi. (1838) 603.—Abyssin.
Huegelii, Meissn. in DC. Prod. xiv. 598 = eriocephalus.
insularis, Meissn. l. c. = eriocephalus.
Kraussii, *Meissn. l. c.* 596.—Afr. austr.
linifolius, *Decne. in Jacquem. Voy. Bot.* 148.—Afr. austr.
macropetalus, *Meissn. in DC. Prod.* xiv. 594.—Afr. austr.
madagascariensis, *Decne. in Jacquem. Voy. Bot.* 148.—Madag.
Meissnerianus, *Endl. Gen. Suppl.* iv. II. 67.—Afr. austr.
Metzianus, Miq. Analect. Bot. Ind. ii. 3. t. 1 = eriocephalus.
microphyllus, *Meissn. in DC. Prod.* xiv. 593.—Afr. austr.
pubescens, *Decne. in Jacquem. Voy. Bot.* 148.—Madag.
pulchellus, *Decne. l. c.* 149.—Afr. austr.
rostratus, *Meissn. in DC. Prod.* xiv. 597.—Madag.
scandens, Endl. Gen. Suppl. iv. II. 67 = Linostoma scandens.
sisparensis, Meissn. in DC. Prod. xiv. 598 = eriocephalus.
socotranus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotr.
speciosus, Decne. in Jacquem. Voy. Bot. 147. t. 150 = eriocephalus.
splendens, *Endl. Gen. Suppl.* iv. II. 67.—Afr. austr.
triplinervis, *Decne. in Jacquem. Voy. Bot.* 149.—Afr. austr.

LASIOSPERMUM, Fisch. Hort. Gorenk. ed. II. 34 (1812) = **Scorzonera**, Tourn. (Compos.).
angustifolium, Fisch. l. c. = E. eriosperma.
ensifolium, Fisch. l. c. = E. ensifolia.
hirsutum, Steud. Nom. ed. II. i. 12 = seq.
hirtum, Fisch. Hort. Gorenk. ed. II. 34.—Hab. ?

LASIOSPERMUM, Lag. Gen. et Sp. Nov. 31 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 416.
ERIOCARPA, Lag. ex DC. Prod. vi. 38 (1837).
ERIOSPHAERA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 221 (1817).

LASIOSPERMUM :—
LANIPILA, Burch. Trav. i. 259 (1824).
MATAXA, Spreng. Syst. iv. Cur. Post. 297 (1827).
alpinum, Reichb. Fl. Germ. Excurs. ii. 225 = pedunculare.
anthemoides, G. Don, ex Loud. Hort. Brit. 337 = Anthemis cretica.
brachyglossum, *DC. Prod.* vi. 38.—Afr. austr.
crithmifolia, G. Don, ex Loud. Hort. Brit. 337 = Santolina crithmifolia.
eriospermum, G. Don, l. c. = pedunculare.
pedunculare, *Lag. Gen. et Sp. Nov.* 31.—Afr. austr.
radiatum, *Trevir. in Nov. Act. Nat. Cur.* xiii. I. (1826) 205.—Afr. austr.
rigidum, G. Don, ex Loud. Hort. Brit. 337 = Anthemis cretica.

LASIOSPORA, Cass. in Dict. Sc. Nat. xxv. 306 (1822) = **Scorzonera**, Tourn. (Compos.).
angustifolia, Cass. in Dict. Sc. Nat. xxv. 306 = S. eriosperma.
candidissima, Sch. Bip. in Nov. Act. Nat. Cur. xxi. I. (1845) 166 = S. veratrifolia.
cretica, Cass. in Dict. Sc. Nat. xxv. 308 = S. cretica.
divaricata, Sch. Bip. in Nov. Act. Nat. Cur. xxi. I. (1845) 166 = S. Aucheriana.
ensifolia, Cass. in Dict. Sc. Nat. xxv. 307 = S. ensifolia.
eriophora, Sch. Bip. in Nov. Act. Nat. Cur. xxi. I. (1845) 166 = S. eriophora.
eriosperma, Sch. Bip. l. c. = S. eriosperma.
hirsuta, Cass. in Dict. Sc. Nat. xxv. 308 = S. hirsuta.
ispahanica, Walp. Rep. vi. 347 = S. ispahanica.
lanata, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 10 = S. lanata.
lasiosperma, Sch. Bip. ex Walp. Rep. vi. 346 = S. eriosperma.
latifolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 10 = S. nervosa.
macrophylla, Sch. Bip. in Nov. Act. Nat. Cur. xxi. I. (1845) 166 = S. macrophylla.
nervosa, Sch. Bip. l. c. = S. nervosa.
paniculata, Walp. Rep. vi. 347 = S. nervosa.
tuberosa, Cass. in Dict. Sc. Nat. xxxv. 307 = S. tuberosa.
veratrifolia, Walp. Rep. vi. 347 = S. veratrifolia.

LASIOSTEGA, Rupr. ex Benth. Pl. Hartw. 347 (1857) = **Buchloë**, Engelm. (1859) (Gramin.).
humilis, Rupr. ex Benth. l. c. = B. dactyloides.

LASIOSTELMA, Benth. in Benth. & Hook. f. Gen. ii. 776 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
Sandersoni, *Oliver, in Hook. Ic. Pl.* t. 1449.—Afr. austr.

LASIOSTEMON, Benth. & Hook. f. Gen. i. 285 (1862) = LASIOSTEMUM, Nees & Mart. = **Cusparia**, Humb. (Rutac.).

LASIOSTEMON, Schott, ex Endl. Gen. 690, in syn. (1839) = **Esterhazya**, Mikan. (Scrophular.).

LASIOSTEMUM, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 152, 171 = **Cusparia**, Humb. (Rutac.).
arboreum, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 239 = Geniostoma Lasiostemon.
silvestre, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 171 = C. silvestris.

LASIOSTOMA, Schreb. Gen. 75 (1789) = **Strychnos**, Linn. (Loganiac.).
Bredemeyeri, Schult. Mant. iii. 64.—Venezuela.
cirrhosa, Willd. Sp. Pl. i. 624 = S. Rouhamon.
cornifolium, Spreng. Neue Entdeck. i. 281.—Bras.
Curare, H. B. & K. Nov. Gen. et Sp. vii. 210 = S. Rouhamon.
divaricata, G. F. W. Mey. Prim. Fl. Esseq. 83 = S. Rouhamon.
glabrum, Willd. ex DC. Prod. ix. 18 = *Bredemeyeri*.
Rouhamon, Roem. & Schult. Syst. iii. 105 = S. Rouhamon.
Ruhamon, J. F. Gmel. Syst. 248 = S. Rouhamon.
stans, Willd. ex Schult. Mant. iii. 64.—Bras.

LASIOSTOMA, Spreng. Syst. i. 422 (1825) = **Hydnophytum**, Jack (Rubiac.).
formicarum, Spreng. Syst. i. 423 = H. formicarum.
loranthifolium, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 224 = H. loranthifolium.

LASIOSTOMA :—
oblongum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 223 = Hydnophytum oblongum.
tuberosum, Spreng. Syst. i. 423 = H. Gaudichaudii.

LASIOSTOMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. (1850) 239 = **Geniostoma**, Forst. (Logan.).
arboreum, Zipp. ex Blume, l. c. = G. Lasiostemon.

LASIOSTYLES, Presl, Bot. Bemerk. 149 (1844); in Abh. Boehm. Ges. V. iii. (1845) 579 = **Cleidion**, Blume (Euphorb.).
salicifolia, Presl, ll. cc. = C. javanicum.

LASIOTRICHOS, Lehm. Cat. Lem. Hort. Hamb. (1834).—Cf. Linnaea, x. (1836) Litt. 113 = **Fingerhuthia**, Nees (Gramin.).

LASIPANA, Rafin. Sylva Tellur. 21 (1838) = **Mallotus**, Lour. (Euphorb.).
tricuspis, Rafin. l. c. 22 = M. paniculatus.

LASIURUS, Boiss. Diagn. Ser. II. iv. (1859) 145 = **Rottboellia**, Linn. f. (Gramin.).
hirsutus, Boiss. l. c. 146 = R. hirsuta.

LASIUS, Hassk. Cat. Hort. Bog. Alt. 59 (1844) = **Lasia**, Lour. (Aroid.).

LASS, Adans. Fam. ii. 400 (1763) = **Abutilon**, Tourn. (Malvac.).

LASSIA, Baill. Etud. Gén. Euph. 464. t. 4 (1858) = **Tragia**, Linn. (Euphorb.).
scandens, Baill. l. c. 465 = T. scandens.

LASTARRIACA, Remy, in C. Gay, Fl. Chil. v. 289. t. 58 (1849). *POLYGONACEAE*, Benth. & Hook. f. iii. 94.
DONATIA, Bert. ex Remy, in C. Gay, Fl. Chil. v. 289 (1849).
chilensis, *Remy, in C. Gay, Fl. Chil.* v. 290.—Chili.

LASTHENIA, Cass. Opusc. iii. 88 (1834). *COMPOSITAE*, Benth. & Hook. f. ii. 400.
HOLOGYMNE, Bartl. Ind. Sem. Hort. Gotting.; ex Linnaea, xii. (1838) Litt. 81.
RANCAGUA, Poepp. & Endl. Nov. Gen. et Sp. i. 15. tt. 24, 25 (1835).
XANTHO, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 191.
ambigua, A. Gray. in Proc. Am. Acad. vi. (1864–65) 547 = Eriophyllum ambiguum.
aurea, *Passer. Ind. Sem. Hort. Parm.* (1855) 3.—Hab. ?
Bridgesii, Steud. Nom. ed. II. ii. 12 = obtusifolia.
californica, DC. ex Lindl. Bot. Reg. sub t. 1780 = glabrata.
canescens, *Regel, Gartenfl.* (1870) 82.—Am. bor. ?
Coulteri, Greene, in Bull. Calif. Acad. i. iv. (1885) 192 = glabrata.
glaberrima, DC. Prod. v. 664 = glabrata.
glabrata, *Lindl. Bot. Reg.* t. 1780.—Calif.
Kunthii, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 319 = obtusifolia.
obtusifolia, *Cass. Opusc.* iii. 89.—Chili.

LASTILA, Alef. in Bonplandia, ix. (1861) 146 = **Lathyrus**, Tourn. (Legumin.).
hirsuta, Alef. l. c. = L. hirsutus.

LASYNEMA, Poir. Encyc. Suppl. ii. 553, 555 (1811) = **Lysinema**, R. Br. (Epacrid.).

LATANA, Robin, Voy. Louis. iii. 385 (1807) = **Lantana**, Linn. (Verben.).

LATANIA, Comm. ex Juss. Gen. 39 (1789). *PALMAE*, Benth. & Hook. f. iii. 940.
CLEOPHORA, Gaertn. Fruct. ii. 185. t. 120 (1791).
aurea, *Duncan, Cat. Hort. Maur.* 56.—Ins. Rodriguez.
borbonica, Lam. Encyc. iii. 427 = Commersonii.
chinensis, Jacq. Fragm. 16. t. 16. f. 1 = Commersonii.
Commersonii, *J. F. Gmel. Syst.* 1035.—Ins. Mascar.
glaucophylla, Hort. ex Baker, Fl. Maurit. 381 = Loddigesii.
Loddigesii, *Mart. Hist. Nat. Palm.* iii. 224. t. 161. f. ii. 10–14.—Ins. Maurit.
plagaecoma, Comm. ex Baker, Fl. Maurit. 381 = Commersonii.
rubra, Jacq. Fragm. 13. t. 8 = Commersonii.
Verschaffeltii, Lem. Illustr. Hortic. vi. (1859) t. 229 = aurea.

LATERIFISSUM, Dulac, Fl. Hautes-Pyr. 366 (1867) = **Montia**, Linn. (Portulac.).
minus, Dulac, l. c. = M. fontana.
rivulare, Dulac, l. c. = M. fontana.

LATHIRUS, Neck. Delic. Gallo-Belg. ii. 305 (1768) = **Lathyrus**, Tourn. (Legum.).

LATHRAEA, Linn. Syst. ed. I (1735); Gen. ed. I. 5 (1737). *OROBANCHACEAE*, Benth. & Hook. f. ii. 985.
ANBLATUM, Tourn. ex Adans. Fam. ii. 207 (1763).
CLANDESTINA, Tourn. ex Adans. l. c. (1763).
SQUAMARIA, Riv. ex Linn. Gen. ed. I. 183 (1737).
amentacea, Schlecht. ex Ledeb. Fl. Ross. iii. 323 = Boschniakia glabra.
Anblatum, Linn. Sp. Pl. 606 = Phelypaea coccinea?
Clandestina, Georgi, Reise, i. 223 = Boschniakia glabra.
Clandestina, *Linn. Sp. Pl.* 605.—Europ. austr.
Clandestina, Uspenski, Jekaterinenb. 262 = L. Squamaria.
japonica, Maxim. ex Franch. & Sav. Enum. Pl. Jap. ii. 461, 462 = Miqueliana.
Miqueliana, *Franch. & Sav. l. c.* 461.—Japon.
Phelipaea, Gueldenst. Reisen, i. 422 = Phelypaea coccinea.
Phelypaea, Linn. Sp. Pl. 606 = Phelypaea tinctoria.
Phelypea, Forsk. Fl. Aegypt. Arab. 111 = Orobanche ramosa.
quinquefida, Forsk. l. c. = Phelypaea lutea.
rhodopea, *Dingl. in Bot. Zeit.* xxxv. (1877) 74.—Thracia.
simplex, S. F. Gray, Nat. Arr. Brit. Pl. ii. 316 = L. Squamaria.
Squamaria, *Linn. Sp. Pl.* 606.—Europ.; As. bor.
strobilacea, Schlecht. ex Ledeb. Fl. Ross. iii. 323 = Boschniakia glabra.

LATHRAEOPHILA, Hook. f. in Trans. Linn. Soc. xxii. (1859) 55 = LATRAEOPHILA, Leandr. Sacram. = **Helosis**, Rich. (Balanoph.).

LATHRIOGYNA, Eckl. & Zeyh. Enum. 170 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 472.
HEUDUSA, E. Mey. Comm. Pl. Afr. Austr. 153 (1835).
candicans, *Eckl. & Zeyh. Enum.* 171.—Afr. austr.
parvifolia, *Eckl. & Zeyh. l. c.* 170.—Afr. austr.

LATHRISIA, Sw. Annot. Bot. 48 (1829) = **Bartholina**, R. Br. (Orchid.).
pectinata, Sw. l. c. = B. pectinata.

LATHROPHYTUM, Eichl. in Bot. Zeit. xxvi. (1868) 513. t. 9. *BALANOPHOREAE*, Benth. & Hook. f. iii. 239.
Peckolti, *Eichl. l. c.* 550.—Bras.

LATHYRAEA, Gled. in Act. Berl. v. (1749) 129, sphalm. = **Lathraea**, Linn. (Orobanch.).

LATHYROS, St. Lag. in Ann. Soc. Bot Lyon, vii. (1880) 129 = **Lathyrus**, Tourn. (Legum.).
clymenus, St. Lag. l. c. = Lathyrus Clymenum.

LATHYRUS, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 526.
ANURUS, Presl, in Weitenw. Beitr. ii. (1837) 24.
APHACA, Tourn. ex Linn. Gen. ed. I. 217 (1737).
ASTROPHIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 278 (1838).
CICERCULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358.
CICERELLA, DC. Prod. ii. 369 (1825).
CLYMENUM, Tourn. ex Adans. Fam. ii. 330 (1763).
GRAPHIOSA, Alef. in Bonplandia, ix. (1861) 128.
LASTILA, Alef. l. c. 146 (1861).
LENS, Tourn. ex Hall. Enum. Stirp. Helv. ii. 601 (1742) partim.
NAVIDURA, Alef. in Bonplandia, ix. (1861) 147.
OCHRUS, Tourn. ex Adans. Fam. ii. 330 (1763).
OROBUS, Linn. Syst. ed. I (1735).
PISUM, Tourn. ex Linn. Gen. ed. I. 222 (1737) partim.
PLATYSTYLIS, Sweet, Brit. Flow. Gard. t. 289 (1828).
SPATULIMA, Rafin. Fl. Tellur. ii. 88 (1836).
acirrosus, Schur, Enum. Pl. Transs. 931, Index = hirsutus.

LATHYRUS:—
acutifolius, Vog. in Linnaea, xiii. (1839) 27 = pubescens.
acutus, *Gromow, ex Trautv. in Act. Hort. Petrop.* iii. 1. (1873) 75, nomen.—Ross.
aegyptiacus, Hasselq. Iter Palaest. 482 = tingetanus.
affinis, Guss. Fl. Sic. Syn. ii. 853 = L. Aphaca.
alatus, Sibth. & Sm. Fl. Graec. Prod. ii. 66 = L. Clymenum.
alatus, *Tenore, Fl. Nap. Prod.* 42.—Ital.
Alberjilla, *Steud. Nom.* ed. II. ii. 13.—Chili.
albidus, Aikin, ex Eaton, Man. Bot. N. Am. ed. VI. 198 = ochroleucus.
albus, Garcke, Fl. Deutschl. ed. IV. 94 = pannonicus.
albus, *A. Gray, in S. Wats. Bot. Calif.* ii. 442.—Calif.
Allardi, *Battand. in Bull. Soc. Bot. Fr.* xxix. (1882) 288.—Alger.
alpestris, *Kit. ex Reichb. Ic. Fl. Germ.* t. 2270.—Europ.
altaicus, *Ledeb. Fl. Alt.* iii. 355.—As. bor.
americanus, Mill. Gard. Dict. ed. VIII. n. 19 = Baptisia perfoliata?
amoenus, *Fenzl, in Russegg. Reis.* i. 890. t. 2; *Pugill. Pl. Nov. Syr.* 3.—Syria.
amphicarpos, Gouan, Hort. Monsp. 368 = setifolius.
amphicarpos, Linn. Sp. Pl. 729 = sativus?
amplicarpos, Sibth. & Sm. Fl. Graec. vii. 62. t. 693 = blepharicarpus.
angulatus, Linn. Mant. ii. 443 = sphaericus.
angulatus, *Linn. Sp. Pl.* 731.—Europ.
angustifolius, Martr. Fl. Tarn, 184 = latifolius.
angustifolius, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = sylvestris.
annuus, *Linn. Amoen. Acad.* iii. 417, nota.—Reg. Medit.; Oriens.
anomalus, Phil. in Linnaea, xxviii. (1856) 627 = Vicia bijuga.
Aphaca, *Linn. Sp. Pl.* 729.—Europ.; Oriens.
aphyllus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 609 = L. Aphaca.
aphyllus, *Link, ex Webb & Berth. Phyt.-Canar.* ii. 103.—Ins. Canar.
aristatus, Vis. Ind. Sem. Hort. Patav. (1826); Fl. Dalmat. ii. 330, in syn. t. 47 = ciliatus.
Armitageanus, Knowles & Westc. Flor. Cab. iii. ('1840') [1839] 81 = nervosus, *Lam.*
articulatus, *Linn. Sp. Pl.* 731.—Reg. Mediterr. occ.
articulatus, Loisel. Fl. Gall. ii. 144 = L. Clymenum.
arvensis, Phil. in Linnaea, xxviii. (1856) 624 = stipularis.
asphodeloides, *Gren. & Godr. Fl. Fr.* i. 488.—Gallia.
attenuatus, Tenore, Fl. Nap. Prod. 41 = sessilifolius.
attenuatus, Viv. Fragm. t. 19 = tuberosus.
aurantius, *C. Koch, in Linnaea*, xv. (1841) 723.—Reg. Cauc.
auriculatus, Bertol. Rar. Ital. Pl. Dec. ii. ed. I. 38 = L. Clymenum.
axillaris, Lam. Encyc. ii. 706 = inconspicuus, sphaericus.
barcinonensis, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 299 = Vicia bithynica.
Berterianus, Colla, in Mem. Acc. Torin. xxxvii. (1834) 61 = stipularis.
bijugus, *Boiss. & Noé, Diagn.* Ser. II. v. 94.—Persia.
bithynicus, Linn. Sp. Pl. 731 = Vicia bithynica.
blepharicarpus, *Boiss. Diagn.* Ser. I. ix. 126.—Syria.
Bolanderi, *S. Wats. in Proc. Am. Acad.* xx. (1885) 363.—Calif.
brachyphyllus, Schur, in Oestr. Bot. Zeitschr. ix. (1861) 85 = latifolius.
brachyterus, *Alef. in Bonplandia*, ix. (1861) 151.—As. Min.?
brutius, Tenore, Ind. Sem. Hort. Neap. (1825) 12 = grandiflorus.
californicus, Dougl. in Lindl. Bot. Reg. t. 1144 = maritimus.
campestris, *Phil. in Linnaea*, xxviii. (1856) 626.—Chili.
canescens, *Gren. & Godr. Fl. Fr.* i. 489.—Gallia.
cashmericus, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 180 = pratensis.
cassius, *Boiss. Diagn.* Ser. I. ix. 128.—Syria.
chilensis, Steud. Nom. ed. II. ii. 13 = magellanicus.
chius, Boiss. & Orph. in Boiss. Diagn. Ser. II. vi. 68 = hierosolymitanus.
chloranthus, *Boiss. & Bal. l. c.* 67.—As. Min.; Persia.
chrysanthus, *Boiss. l. c.* Ser. I. vi. 46.—Mesopot.
Cicer, Habl. Taur. 125; ex Bieb. Fl. Taur. Cauc. ii. 156 = hirsutus.
Cicera, All. Fl. Pedem. i. 329 = articulatus.

LATHYRUS:—
Cicera, *Linn. Sp. Pl.* 730.—Europ.; Oriens.
ciliatus, *Guss. Pl. Rar.* 296. t. 49.—Europ. austr.
cirrhosus, *Ser. in DC. Prod.* ii. 374.—Mont. Pyren.
clymenoides, *Ser. l. c.*—Ins. Maurit.
Clymenum, *Linn. Sp. Pl.* 732.—Reg. Mediterr.
coccineus, All. Fl. Pedem. i. 330 = sphaericus.
coerulescens, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 43.—Marocc.
controversus, Loret & Barr. Fl. Montp. ii. 808, in obs. = sphaericus.
cornutus, Spreng. Fl. Hal. Mant. 47 = L. Clymenum.
crassipes, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 198 = stipularis.
crassipes, *Phil. in Linnaea*, xxxiii. (1864-65) 54.—Chili.
currentifolius, Lam. Fl. Fr. ii. 571 = L. Ochrus.
curvus, Roth, Bot. Abh. 13 = incurvus.
Cusickii, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 371.—Am. bor. occ.
cyaneus, *C. Koch, in Linnaea*, xv. (1841) 723.—Rossia.
Davidii, *Hance, in Journ. Bot.* ix. (1871) 130.—Manchur.
debilis, *Clos, in C. Gay, Fl. Chil.* ii. 142.—Chili.
debilis, Vog. in Linnaea, xiii. (1839) 26 = subulatus.
decaphyllus, Hook. Fl. Bor. Am. i. 159 = venosus?
decaphyllus, Pursh, Fl. Am. Sept. ii. 471 = polymorphus.
denudatus, Gilib. Fl. Lituan. ii. 98 = pratensis.
dicirrhus, *Clos, in C. Gay, Fl. Chil.* ii. 143.—Chili.
diffusus, G. Don, Gen. Syst. ii. 336 = Vicia americana.
dissitifolius, Nutt. ex Torr. & Gray, Fl. N. Am. i. 277 = Vicia americana.
diversifolius, *Presl, Delic. Prag.* 38.—Sicilia.
Dolichos, Medic. Beob. (1783) 215 (Quid?).—Hab.?
Drummondii, *Hort.—Cf. Gard. Chron.* (1876) II. 16.—Hab.?
dubius, Tenore, Ind. Sem. Hort. Neap. (1825) 11. n. 12 = L. Cicera.
dumetorum, *Ledeb. Comm. in J. G. Gmel. Fl. Sib.* 110.—Sibir.
dumetorum, *Phil. in Linnaea*, xxviii. (1856) 626.—Chili.
elegans, Vog. l. c. xiii. (1839) 30 = nervosus, *Lam.*
Engelmanni, *Bischoff, Del. Sem. Hort. Heidelb.* 1839.—*Cf. Linnaea*, xiv. (1840) *Litt.* 132.—Am. bor. occ.
ensifolius, Badarso, in Brugnat. Giorn. Fis. Dec. II. vii. (1824) 369 = latifolius.
ensifolius, J. Gay, in Ann. Sc. Nat. Sér. IV. viii. (1857) 313 = canescens.
epetiolaris, *Clos, in C. Gay, Fl. Chil.* ii. 146.—Chili.
erectus, Lag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 318 = angulatus.
erectus, *Lag. Gen. et Sp. Nov.* 22.—Europ. austr.; Oriens.
erythrinus, Presl, Fl. Sicul. p. xxiii = L. Cicera.
Ewaldi, *Meinsh. Fl. Ingr.* 90.—Ingria.
filiformis, *J. Gay, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 315.—Europ. austr.
filipetiolatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Aphaca.
fissus, *Ball, in Journ. Bot.* xi. (1873) 332.—Marocc.
foliaceopetiolatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Nissolia.
frigidus, Schott & Kotschy, in Boiss. Diagn. Ser. II. v. 94 = Pisum formosum.
fruticosus, *Cav. Ic.* i. 58. t. 84.—Peruv.
Gerardi, Roth, Catalect. fasc. ii. 96 = sphaericus.
gladiatus, Hook. Ic. Pl. t. 72 = magellanicus.
glaucifolius, Beck, Bot. N. & Mid. St. 90 = ochroleucus.
Gorgoni, Heldr. ex Nym. Consp. 203 = annuus.
Gorgoni, *Parl. in Giorn. Sc. Sicil.* lxii. (1838) 3.—Sicil.
gracilis, *Phil. in Linnaea*, xxviii. (1856) 627.—Chili.
gramineus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 610 = L. Nissolia.
grandiflorus, Láng, in Syll. Ratisb. i. (1824) 182 = sylvestris.
grandiflorus, *Sibth. & Sm. Fl. Graec. Prod.* ii. 67.—Ital.; Graec.
grandifolius, Schur, Enum. Pl. Transs. 176 = latifolius.
granulatus, Schur, l. c. = palustris.
Hallersteinii, Baumg. Enum. Stirp. Transs. ii. 333 = pratensis.
helodes, *Link, ex DC. Prod.* ii. 376, nomen.—Hab.?

LATHYRUS:—

heterophyllus, Gouan, Hort. Monsp. 370 = latifolius.
heterophyllus, Lapeyr. Hist. Abr. Pl. Pyr. 416 = sylvestris.
heterophyllus, *Linn. Sp. Pl.* 733.—Europ.
hexaedrus, Chaub. in 'Ann. Sc. Obs. (Juin, 1830)'; ex Bory & Chaub. Fl. Pélop. 47 = angulatus.
hierosolymitanus, *Boiss. Diagn.* Ser. I. ix. 127.—Syria.
himalayensis, Cambess. in Jacquem. Voy. i. 44 = Vicia rigidula.
hirsutus, *Linn. Sp. Pl.* 732.—Europ.; Oriens.
hirtus, *Lam. Encyc.* ii. 706.—Hab.?
hispanicus, Hasselq. ex Steud. Nom. ed. I. 466 = tingitanus.
hispanicus, Mill. Gard. Dict. ed. VIII. n. 5 = articulatus.
hispidulus, Boiss. Diagn. Ser. I. vi. 46 = inconspicuus.
Hookeri, G. Don, Gen. Syst. ii. 332 = magellanicus.
Hookerianus, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 198 = subulatus.
humilis, Fisch. ex Ser. in DC. Prod. ii. 378 = altaicus.
hydrophilus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 276 = palustris.
inconspicuus, *Linn. Sp. Pl.* 730.—Europ.; Oriens.
incurvus, Rafin. Atl. Journ. 151 (Quid?).—Am. bor.
incurvus, Reichb. Fl. Germ. Excurs. 536 = palustris.
incurvus, *Willd. Sp. Pl.* iii. 1091.—Rossia; As. Min.
inermis, Rochel, ex Frivald. in Magyar Tud. Tár. Evkön. ii. (1835) 250. t. 2 = hirsutus.
intermedius, Eichw. Skizze, 168 = heterophyllus.
intermedius, Wallr. Sched. Crit. 386 = sylvestris.
italicus, *Fisch. ex DC. Prod.* ii. 376, nomen.—Ital.
italicus, Hort. ex Hornem. Enum. Pl. Hort. Hafn. (1807) 41 = Vicia articulata.
japonicus, Willd. Sp. Pl. iii. 1092 = maritimus.
Lanszwertii, Kellogg, in Proc. Calif. Acad. ii. (1863) 150 = palustris.
latifolius, *Linn. Sp. Pl.* 733.—Europ.
latifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 69 = undulatus.
latifolius, Vis. Stirp. Dalmat. 39 = sylvestris.
Ledebourii, *Trautv. in Act. Hort. Petrop.* iii. (1875) 72.—Rossia.
Lens, Bernh. Syst. Verz. Erf. 247 = Lens esculenta.
Lenticula, Peterm. Deutschl. Fl. 155 = Lens Lenticula.
leptophyllus, *Bieb. Fl. Taur. Cauc.* ii. 155.—Reg. Caucas.
leptophyllus, C. Koch, ex Boiss. Fl. Orient. ii. 609 = hirsutus.
leptophyllus, Maly, Enum. Pl. Austr. 365 = annuus.
linearifolius, Nym. Consp. 202 = palustris.
linearifolius, *Vog. in Linnaea*, xiii. (1839) 28.—Reg. Argent.
linearis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 276 = Vicia americana.
litoralis, *Phil. in Linnaea*, xxviii. (1856) 625.—Chili.
littoralis, *Endl. ex Walp. Rep.* i. 722.—Am. bor. occ.
longepedunculatus, Ledeb. Ind. Hort. Dorpat. (1825) Suppl. 5 = angulatus.
longifolius, Tenore, Ind. Sem. Hort. Neap. (1825) 12 = latifolius.
longipes, *Phil. in Linnaea*, xxviii. (1856) 625.—Chili.
lusitanicus, *Mart. ex DC. Prod.* ii. 376.—Hispan.
Lusseri, Heer, ex Koch, Syn. Fl. Germ. ed. II. 1021 = pratensis.
luteus, Baker, in Hook. f. Fl. Brit. Ind. ii. 180 = montanus?
luteus, Moench, Meth. 138 = annuus.
luteus, *Munby, Fl. Algér.* 73.—Algeria.
lycius, *Boiss. & Heldr. Diagn.* Ser. I. ix. 128.—Lycia.
Maccaguenii, Tod. ex Nym. Consp. 201 = odoratus.
Macraei, *Hook. & Arn. Bot. Beech. Voy.* 21.—Chili.
macropus, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 198 = pubescens.
macrorrhizus, *Wimm. Fl. Schles.* (1841) 166.—Europ.
macrostachys, Vog. in Linnaea, xiii. (1839) 23 = magellanicus.
magellanicus, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 344 = nervosus, *Lam.*
magellanicus. *Lam. Encyc.* ii. 708.—Reg. Magellan.
magniflorus, *Mill. Gard. Dict.* ed. VIII. n. 14.—Hab.?
maritimus, *Bigel. Fl. Bost.* ed. III. 286.—Europ.; Am. bor.
maritimus, Torr. Bot. Mex. Bound. 49 = vestitus.

LATHYRUS:—

marmoratus, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 606.—Syria; Aegypt.
megalanthos, Steud. Nom. ed. II. ii. 14 = sylvestris.
membranaceus, Presl, Delic. Prag. 40 = latifolius.
Messerschmidii, *Franch. & Sav. Enum. Pl. Jap.* i. 106.—Japon.
mexicanus, [*Wender.*] *Ind. Sem. Hort. Marb.* (1837).—*Cf. Linnaea*, xii. (1838) *Litt.* 85.—Mexic.
micranthus, Gérard, ex Loisel. in Desv. Journ. Bot. ii. (1809) 362 = inconspicuus.
microphyllus, *Scheele, in Linnaea*, xxi. (1848) 575.—Dalmat.
miniatus, Bieb. ex Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 161 = rotundifolius.
mollis, *Steud. Nom.* ed. II. ii. 14.—Chili.
monanthos, Baumg. Enum. Stirp. Transs. ii. 332 = Vicia tetrasperma.
monanthos, Willd. Sp. Pl. iii. 1083 = Vicia monantha.
monspeliensis, Delile, ex Lorr. & Barr. Fl. Montp. 195 = latifolius.
montanus, *Bernh. Syst. Verz. Erf.* 247; *Gren. & Godr. Fl. Fr.* i. 486.—Europ.; As. bor.
montevidensis, Vog. in Linnaea, xiii. (1839) 22 = stipularis.
multiceps, *Clos, in C. Gay, Fl. Chil.* ii. 149.—Chili.
multiflorus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 274 = venosus.
multiflorus, *Peterm. Deutschl. Fl.* 155.—German.
mutabilis, Klinggr. ex Nym. Consp. 202 = pisiformis.
mutabilis, *Sweet, Brit. Flow. Gard.* Ser. II. t. 194.—Sibiria.
myrtifolius, Muhl. ex Willd. Sp. Pl. iii. 1091 = palustris.
myrtifolius, Spreng. Syst. iii. 264 = polymorphus.
neglectus, Puel, Cat. Pl. Lot, 179 = sylvestris.
nervatus, *Presl, Delic. Prag.* 39.—Sicilia.
nervosus, *Boiss. Fl. Orient.* ii. 609.—Mesopot.
nervosus, *Lam. Encyc.* ii. 708.—Reg. Argent.
neurolobus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 125.—Ins. Creta.
nevadensis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 133.—Am. bor. occ.
niger, *Bernh. Syst. Verz. Erf.* 247; *Wimm. Fl. Schles.* (1841) 166.—Europ.; Reg. Cauc.
nigricans, Peterm. Deutschl. Fl. 155 = Lens nigricans.
Nissolia, *Linn. Sp. Pl.* 729.—Europ.; Oriens.
nitens, *Vog. in Linnaea*, xiii. (1839) 25.—Reg. Argent.
Noëanus, Alef. in Bonplandia, ix. (1861) 152 = sylvestris.
occidentalis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 276 = palustris.
ochraceus, *Kittel, Fl. Deutschl.* ed. II. 1301.—Europ.
ochroleucus, *Hook. Fl. Bor. Am.* i. 159.—Am. bor. occ.
ochroleucus, Torr. in Bot. U. St. Expl. Exped. ii. 267 = venosus.
Ochrus, *DC. Fl. Fr.* iv. 578.—Reg. Mediterr.
odoratus, *Linn. Sp. Pl.* 732.—Ital.
odoratus-zeylanicus, Burm. f. Fl. Ind. 162 = odoratus.
odorus, Thomas, ex Tenore, Syll. 355 = grandiflorus.
oleraceus, Lam. Fl. Fr. ii. 580 = Pisum arvense.
ornatus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 277.—Am. bor. occ.
ovalifolius, *Phil. in Anal. Univ. Chil.* (1872) 694.—Chili.
ovatus, Royle, Illustr. Bot. Himal. 200 = altaicus.
pallescens, *C. Koch, in Linnaea*, xv. (1841) 723.—Rossia.
paluster, Auct. = seq.
palustris, Lang, ex Tenore, Syll. 356 = L. Clymenum.
palustris, *Linn. Sp. Pl.* 733.—Amphig. bor.
pannonicus, *Garcke, Fl. Deutschl.* ed. VI. 112.—Europ.; As. bor.
parisiensis, *Mill. Gard. Dict.* ed. VIII. n. 4.—Europ.
Parreiszii, *Presl, in Weitenw. Beitr.* ii. (1837) 30.—Tauria.
parviflorus, Roth, Catalect. fasc. i. 88 = inconspicuus.
parvifolius, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 345.—Am. bor.
pedunculatus, Poir. Encyc. Suppl. ii. 775 = rotundifolius.
pentaphyllus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 (Quid?).—Java.
petiolaris, Vog. in Linnaea, xiii. (1839) 29 = pubescens.
Philippii, Alef. ex Phil. in Anal. Univ. Chil. (1872) 694 = magellanicus.
pilosus, Cham. in Linnaea, vi. (1831) 548 = palustris.

LATHYRUS:—

pilosus, Steud. & Hochst. ex Reichb. Fl. Germ. Excurs. 534 = L. Cicera.
pisaster, *Clos, in C. Gay, Fl. Chil.* ii. 148.—Chili.
pisiformis, Houtt. Handleid. x. 197. t. 85. f. 1 = maritimus.
pisiformis, *Linn. Sp. Pl.* 734.—Europ.; As. bor.
pisiformis, Richards. in Frankl. Journ. App. 17 = ochroleucus.
platyphyllus, A. Gouault, in Rev. Hortic. Sér. IV. iii. (1854) 321 = seq.?
platyphyllus, Retz. Prod. Fl. Scand. ed. II. 170 = sylvestris.
polyanthus, *Boiss. & Blanche, Diagn.* Ser. II. ii. 43, in nota.—Syria; Mesopot.; Oriens.
polymorphus, Hook. Lond. Journ. Bot. vi. (1847) 207 = nevadensis.
polymorphus, *Nutt. Gen. Am.* ii. 97.—Am. bor. occ.
polyphyllus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 274.—Am. bor. occ.
polyphyllus, S. Wats. in Rep. King's Exp. v. 78 = palustris.
pratensis, *Linn. Sp. Pl.* 733.—Reg. bor. temp.
prostratus, Brign. Fasc. Pl. Forojul. 28 = setifolius.
Pseudo-aphaca, *Boiss. Diagn.* Ser. I. ii. 105.—As. Min.
pterocaulos, *Phil. in Anal. Univ. Chil.* (1872) 695.—Chili.
pubescens, Hook. Bot. Mag. t. 3996 = tomentosus.
pubescens, *Hook. & Arn. Bot. Beech. Voy.* 21.—Chili.
pubescens, Nutt. ex Torr. & Gray, Fl. N. Am. i. 274 = venosus.
pubescens, Porter & Coult. Fl. Colerad. 32 = palustris.
pulcher, J. Gay, in Ann. Sc. Nat. Sér. IV. viii. (1857) 311 = latifolius.
pulcher, Salisb. Prod. 338 = tingitanus.
purpureo-coeruleus, *Knowles & Westc. Flor. Cab.* ii. (1838) 126, 177. t. 88.—Bras.
purpureus, Desf. in Ann. Mus. Par. xii. (1808) 56. t. 7 = L. Clymenum.
purpureus, Gilib. Fl. Lituan. ii. 99 = latifolius.
purpureus, *Presl, Delic. Prag.* 39.—Sicilia.
pusillus, *Ell. Sketch*, ii. 223.—Am. bor.
pyrenaicus, Jord. Cat. Jard. Dijon (1848), 27 = sylvestris.
quadrimarginatus, *Bory & Chaub. Fl. Pélop.* 47.—Graecia.
roseus, *Phil. in Linnaea*, xxviii. (1856) 626.—Chili.
roseus, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 92.—Reg. Caucas.
rotundifolius, Janka, in Linnaea, xxx. (1859–60) 565 = latifolius.
rotundifolius, *Willd. Sp. Pl.* iii. 1088.—Ross.; Oriens.
sagittatus, Molina, Sagg. Chile, ed. II. 293 = magellanicus.
sanguineus, Haberle, ex Steud. Nom. ed. II. ii. 14 = pisiformis.
sativus, *Linn. Sp. Pl.* 730.—Europ.; Afr. bor.; Oriens.
sativus, Sibth. & Sm. ex Steud. Nom. ed. II. ii. 14 = quadrimarginatus.
saxatilis, Vis. Fl. Dalmat. iii. 330 = ciliatus.
segetum, Lam. Fl. Fr. ii. 571 = L. Aphaca.
sepium, Scop. Fl. Carn. ed. II. 64 = pratensis.
sericeus, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 60.—Chili.
sericeus, *Lam. Encyc.* ii. 709.—Bras.
sessilifolius, Hook. & Arn. Bot. Beech. Voy. 20 = magellanicus.
sessilifolius, *Tenore, Fl. Neap. Prod. App.* v. 21.—Italia.
setifolius, Gouan, Hort. Monsp. 368 = sphaericus.
setifolius, *Linn. Sp. Pl.* 731.—Europ. austr.
setiger, *Phil. in Linnaea*, xxxiii. (1864–65) 53.—Chili.
siculus, *Steud. Nom.* ed. II. ii. 14 = purpureus.
sinensis, *Mart. ex DC. Prod.* ii. 376.—China.
speciosus, *G. Don, Gen. Syst.* ii. 333.—Mexic.
speciosus, Hort. ex Steud. Nom. ed. II. ii. 14 = articulatus.
spectabilis, *Forsk. Fl. Aegypt. Arab.* 135.—Arab.
sphaericus, Bieb. Tabl. Prov. Casp. 210 = leptophyllus.
sphaericus, *Retz. Obs.* iii. 39.—Europ. austr.
sphaericus, Schur, Enum. Pl. Transs. 174 = L. Nissolia.
spissus, Steph. ex Ledeb. Fl. Ross. i. 677 = Vicia costata.
splendens, *Kellogg, in Proc. Calif. Acad.* vii. 1. (1877) 90.—Calif.

LATHYRUS:—
spurius, Willd. Enum. Hort. Berol. 760 = articulatus.
stans, Vis. in Flora, xii. (1829) I. Erg. 19 = inconspicuus.
stenolobus, *Boiss. Diagn.* Ser. I. ix. 124.—Syria.
stenophyllus, *Boiss. & Heldr. l. c.* 126.—As. Min.; Syria.
stipulaceus, Le Conte, in Torr. Cat. Pl. N. York, 60, 92 = palustris.
stipularis, *Presl, in Weitenw. Beitr.* ii. (1837) 30.—Peruv.
strictus, *Grauer, Dec.* 12.—Hab.?
strictus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 276 = vestitus.
subandinus, *Phil. in Linnaea*, xxxiii. (1864–65) 52.—Chili.
subrotundus, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 390.—Manchur.
subulatus, *Lam. Encyc.* ii. 707.—Bras.
sulphureus, *Brewer, ex A. Gray, in Proc. Am. Acad.* vii. (1868) 399.—Calif.
sylvestris, *Linn. Sp. Pl.* 733.—Europ.
Szowitsii, *Boiss. Fl. Orient.* ii. 605.—Persia.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* i. 105; ii. 326.—Japon.
tataricus, Schreb. ex Steud. Nom. ed. II. ii. 14 = L. Clymenum.
tenuifolius, Desf. Fl. Atlant. ii. 160 = L. Clymenum.
tetrapterus, *Pomel, Nouv. Mat. Fl. Atl.* 194.—Alger.
Thirkaenus, C. Koch, in Linnaea, xix. (1847) 65 = L. Ochrus.
tingitanus, *Linn. Sp. Pl.* 732.—Reg. Mediterr. occ.
tomentosus, Forsk. Fl. Aegypt. Arab. 135 = Tephrosia tomentosa.
tomentosus, *Lam. Encyc.* ii. 709.—Bras.
tomentosus, Steud. Nom. ed. II. ii. 14 = Coursetia tomentosa.
Torreyi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 337.—Calif.
trachycarpus, *Boiss. Fl. Orient.* ii. 608.—Mesopot.
trachyspermus, *Webb, ex Rodr. in Bull. Soc. Bot. Fr.* xxv. (1878) 239.—Ins. Canar.
trichocalyx, *Phil. in Anal. Univ. Chil.* (1872) 695.—Chili.
trigonus, Vog. in Linnaea, xiii. (1839) 31 = nervosus, *Lam.*
tuberculatus, Gaud. Fl. Helv. iv. 488. in obs. = seq.?
tuberosus, *Linn. Sp. Pl.* 732.—Geront. bor.
tumidus, Willd. Sp. Pl. iii. 1082 = Vicia bithynica.
turgidus, Lam. Encyc. ii. 706 = Vicia bithynica.
undulatus, *Boiss. Diagn.* Ser. II. ii. 41.—Byzant.
variegatus, Gilib. Fl. Lituan. ii. 99 = sylvestris.
variegatus, Gren. & Godr. Fl. Fr. i. 485 = multiflorus.
variegatus, Host, Fl. Austr. ii. 327 = hirsutus.
varius, *C. Koch, in Linnaea*, xv. (1841) 723.—Italia.
venosus, *Muhl. ex Willd. Sp. Pl.* iii. 1092.—Am. bor.
venosus, Sweet, Brit. Flow. Gard. Ser. I. t. 37 = maritimus.
vernus, *Bernh. Syst. Verz. Erf.* 247; *Wimm. Fl. Schles.* (1841) 166.—Europ.
vestitus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 276.—Am. bor. occ.
viciaefolius, Ledeb. Fl. Ross. i. 686 = palustris.
viciaeformis, Gilib. Fl. Lituan. ii. 100; Wallr. Sched. Crit. 388 = palustris.
villosus, *Frivald. in Flora*, xix. (1836) 437.—Rumelia.
vinealis, *Boiss. & Noé, Diagn.* Ser. II. ii. 42.—Armen.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1862) II. 443.—Chili.

LATICOMA, Rafin. Fl. Tellur. iv. 12 (1836) = **Nerine**, Herb. (Amaryll.).
compressa, Rafin. l. c. = N. lucida.

LATIPES, Kunth, Rév. Gram. i. 261. t. 42 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1123.
senegalensis, *Kunth, l. c.*—Afr. trop.

LATOUREA, Benth. & Hook. f. Gen. iii. 501 (1883) = seq.

LATOURIA, Blume, Rumphia, iv. 41. t. 195, 199 (1850) = **Dendrobium**, Sw. (Orchid.)
spectabilis, Blume, l. c. = D. spectabile.

LATOURIA, Lindl. Veg. Kingd. 695 (1847) = **Leschenaultia**, R. Br. (Goodenov.).
filiformis, De Vriese, in Verh. Haarl. x. (1854) 187 = Leschenaultia filiformis.

LATRAEOPHILA, Leandr. Sacram. ex A. St. Hil. in Ann. Sc. Nat. Sér. II. vii. (1837) 32 = **Helosis**, Rich. (Balanoph.).

LATREILLEA, DC. Prod. v. 504 (1836) = **Ichthyothere**, Mart. (Compos.).
dicarpa, Toro, ex Ernst, in Flora, lvii. (1874) 212.—Venezuela.
glabrata, Benth. in Hook. Journ. Bot. ii. (1840) 42 = I. Cunabi.
integrifolia, DC. Prod. v. 504 = I. integrifolia.
latifolia, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 110 = I. latifolia.
linearis, Benth. l. c. = I. linearis.
peruviana, Poepp. & Endl. Nov. Gen. et Sp. iii. 46. t. 46 = I. peruviana.
serrata, DC. Prod. v. 504 = I. Cunabi.

LATRIENDA, Rafin. Fl. Tellur. iv. 81 (1836) = **Convolvulus**, Tourn., etc.
brasiliensis, Rafin. l. c. = Ipomoea Pes-caprae.
Imperati, Rafin. l. c. = Ipomoea carnosa.
multiflora, Rafin. l. c. = Ipomoea sidaefolia.
palmata, Rafin. l. c. = Ipomoea quinquefolia.
Soldanella, Rafin. l. c. = C. Soldanella.

LATROBEA, Meissn. in Lehm. Pl. Preiss. ii. 219 (1847). *LEGUMINOSAE*, Benth. & Hook. f. i. 471.
LEPTOCYTISUS, Meissn. l. c. 211 (1847).
Brunonis, *Meissn. l. c.* 219.—Austral.
diosmifolia, *Benth. Fl. Austral.* ii. 142.—Austral.
genistoides, *Meissn. in Lehm. Pl. Preiss.* ii. 219.—Austral.
hirtella, *Benth. Fl. Austral.* ii. 142.—Austral.
pungens, *Benth. l. c.* 140.—Austral.
tenella, *Benth. l. c.* 142.—Austral.

LATUA, Phil. in Bot. Zeit. xvi. (1858) 242. *SOLANACEAE*, Benth. & Hook. f. ii. 896.
venenosa, *Phil. l. c.* 242.—Chili.

LATYRUS, Gren. in Acad. Sc. Besanç. Séances Publ. (1838) 131 = **Lathyrus**, Tourn. (Legumin.).

LAUBERTIA, A. DC. in DC. Prod. viii. 486 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 724.
Boissieri, *A. DC. l. c.* 487.—Peruv.?

LAUCHEA, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 121 = **Begonia**, Linn.
verticillata, Klotzsch, l. c. 122 = B. verticillata.

LAUGERIA, Linn. Syst. ed. XII. 177 (1767) = LAUGIERIA, Jacq. = **Guettarda**, Linn. (Rubiac.).

LAUGERIA, Vahl, Eclog. Am. 26. t. 10 (1796) partim. *RUBIACEAE*, Benth. & Hook. f. ii. 100.
coriacea, Vahl, l. c. = Guettarda coriacea.
densiflora, *Benth. & Hook. f. Gen.* ii. 101.—Ind. occ.
dependens, Ruiz & Pav. Fl. Per. ii. 23 = Guettarda dependens.
dichotoma, [Moç. & Sesse,] ex DC. Prod. iv. 461 = Antirrhoea dichotoma.
hirsuta, Ruiz & Pav. Fl. Per. ii. 22 = Guettarda hirsuta.
lucida, Sw. Prod. Veg. Ind. Occ. 48 = Antirrhoea lucida.
resinosa, *Vahl, Eclog. Am.* i. 27.—Am. austr.
tomentosa, Sw. Prod. Veg. Ind. Occ. 48 (= *Stenostomum tomentosum*).—Jamaic.
tubulosa, Fors, ex DC. Prod. iv. 458 = Guettarda coriacea.

LAUGIERIA, Jacq. Enum. Pl. Carib. 2 (1760) = **Guettarda**, Linn. (Rubiac.).
Edechia, Crantz, Inst. ii. 381 = G. odorata.
odorata, Jacq. Enum. Pl. Carib. 16 = G. odorata.

LAUNAEA, Cass. in Dict. Sc. Nat. xxv. 321 (1822). *COMPOSITAE* [Benth. & Hook. f. ii. 520].
AMMOSERIS, Endl. Gen. 500 (1838).
LOMATOLEPIS, Cass. in Dict. Sc. Nat. xlviii. 422 (1827).
MICRORHYNCHUS, Less. Syn. Comp. 139 (1832).
RHABDOTHECA, Cass. in Dict. Sc. Nat. xlviii. 424 (1827).
ZOLLIKOFERIA, DC. Prod. vii. 183 (1830).
aspleniifolia, *Hook. f. Fl. Brit. Ind.* iii. 415.—Ind. or.
bellidifolia, *Cass. in Dict. Sc. Nat.* xxv. 321.—Afr. trop.

LAUNAEA:—
chondrilloides, *Hook. f. Fl. Brit. Ind.* iii. 415.—Oriens; Ind. or.
crepoides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 842.—Ins. Socotr.
glomerata, *Hook. f. Fl. Brit. Ind.* iii. 417.—Afr. bor.; Oriens.
microcephala, *Hook. f. l. c.* 415.—Reg. Himal.
nudicaulis, *Hook. f. l. c.* 416.—Oriens; Ind. or.
pinnatifida, *Cass. in Ann. Sc. Nat.* Ser. I. xxiii. (1831) 85.—As. et Afr. trop.
secunda, *Hook. f. Fl. Brit. Ind.* iii. 416.—Reg. Himal.
spinosa, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 428.—Marocc.

LAUNAYA, Reichb. Nom. 95 (1841) = praec.

LAUNEA, Endl. Gen. Suppl. i. 1388, in syn. (1841) = **Launaea**, Cass. (Compos.).

LAUNZAN, Buch.-Ham. in As. Res. v. (1799) 123 = **Buchanania**, Roxb. (Anacard.).

LAUNZEA, Endl. Ench. 599 (1841) = praec.

LAURADIA, Vand. Fl. Lusit. 15 (1788) = **Lavradia**, Vell. (Violar.).

LAUREA, Gaudich. Bot. Voy. Bonite, 513. t. 88 (1826) **Bagassa**, Aubl. (Urticac.).
tiliaefolia, Gaudich. l. c. = B. Sagotiana.

LAURELIA, Juss. in Ann. Mus. Par. xiv. (1809) 134. *MONIMIACEAE*, Benth. & Hook. f. iii. 145.
PAVONIA, Ruiz & Pav. Fl. Per. Prod. 127. t. 28 (1794).
THIGA, Molina, Sagg. Chile, ed. II. 297 (1810).
aromatica, *Juss. ex Poir. Encyc. Suppl.* iii. 313.—Chili.
crenata, Poepp. ex Tul. in Arch. Mus. Par. viii. (1855–56) 416 = aromatica.
Novae-Zelandiae, *A. Cunn. in Ann. Nat. Hist.* Ser. I. i. (1838) 381.—N. Zel.
sempervirens, Tul. in Arch. Mus. Par. viii. (1855–56) 416 = aromatica.
serrata, Bert. Merc. Chil. 685; ex Am. Journ. Sc. xxiii. (1833) 89 = aromatica.

LAUREMBERGIA, Berg. Desc. Pl. Cap. 350 (1767) = **Serpicula**, Linn. (Halorag.).
capensis, J. F. Gmel. Syst. 284 = S. repens.
coccinea, Kanitz, in Mart. Fl. Bras. viii. II. 377.—Bras.
repens, Berg. Desc. Pl. Cap. 350 = S. repens.
tetrandra, Kanitz, in Mart. Fl. Bras. xiii. II. 378.—Bras.

LAURENCELLIA, Neum. in Rev. Hortic. Sér. II. iv. (1845) 26 = LAWRENCELLIA, Lindl. = **Helichrysum**, Vaill. (Compos.).

LAURENTA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 207 = seq.

LAURENTIA, Michx. ex Adans. Fam. ii. 134 (1763); Neck. Elem. i. 131 (1790). *CAMPANULACEAE*, Benth. & Hook. f. ii. 549.
ENCHYSIA, Presl, Prod. Monog. Lobel. 40 (1836).
PORTERELLA, Torr. in Hayd. Rep. Geol. Surv. Montana, 488 (1872).
SOLENOPSIS, Presl, Prod. Monog. Lobel. 32 (1836).
arabidea, *A. DC. in DC. Prod.* vii. 410.—Afr. austr.
Baueri, A. DC. l. c. 411 = Isotoma fluviatilis?
bifida, Sond. in Harv. & Sond. Fl. Cap. iii. 552 = Dregeana.
canariensis, *A. DC. in DC. Prod.* vii. 410.—Ins. Canar.
carnosula, *Benth. & Hook. f. Gen.* ii. 549.—Am. bor. occ.
commutata, *Tod. Ind. Sem. Hort. Panorm.* (1872); *ex Nuov. Giorn. Bot.* v. (1873) 160.—Sicil.?
Dregeana, *A. DC. in DC. Prod.* vii. 411.—Afr. austr.
erinoides, *Nichols. Dict. Gard.* ii. 238.—Afr. austr.
etbaica, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 683.—Afr. trop.
Gaudichaudii, A. DC. in DC. Prod. vii. 411 = Isotoma fluviatilis?
hederacea, *Sond. in Harv. & Sond. Fl. Cap.* iii. 553.—Afr. austr.

LAURENTIA:—
Michelii, *A. DC. in DC. Prod.* vii. 409.—Reg. Mediterr.
Michelii, Ind. Sem. Hort. Berol. (1866) 17; ex Tod. in Nuov. Giorn. Bot. v. (1873) 160 = commutata.
micrantha, *A. DC. in DC. Prod.* vii. 411.—Afr. austr.
minuta, *A. DC. l. c.* 410.—Afr. austr.
platycalyx, F. Muell. in Trans. Vict. Inst. i. (1855) 39 = Pratia platycalyx.
pusilla, A. DC. in DC. Prod. vii. 411 = Isotoma pusilla.
pygmaea, Sond. in Harv. & Sond. Fl. Cap. iii. 552 = micrantha.
ramosissima, *Benth. & Hook. f. Gen.* ii. 549.—Mexic.
repens, *Benth. & Hook. f. l. c.*—Afr. austr.
tenella, *A. DC. in DC. Prod.* vii. 410.—Reg. Mediterr.

LAURENTIA, Steud. Nom. ed. I. 466 (1821) = LORENTEA, Orteg. = **Sanvitalia**, Gualt. (Compos.).
atropurpurea, Steud. l. c. = S. procumbens.

LAUREOLA, M. Roem. Syn. Hesper. 74 (1846) = **Skimmia**, Thunb. (Rutac.).
fragrans, M. Roem. l. c. = S. Laureola.

LAUREOLA, Rupp. Fl. Jen. ed. Hall. 40 (1745) = **Daphne**, Linn. (Thymel.).
sempervirens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 147 = D. Laureola.

LAURERIA, Schlecht. in Linnaea, viii. (1833) 513 = **Juanulloa**, Ruiz & Pav. (Solanac.).
mexicana, Schlecht. l. c. = J. mexicana.

LAURIDIA, Eckl. & Zeyh. Enum. 124 (1835) = **Elaeodendron**, Jacq. (Celastrin.).
reticulata, Eckl. & Zeyh. l. c. 125.—Afr. austr.
rupicola, Eckl. & Zeyh. l. c. = *reticulata*?

LAUROCERASUS, [Tourn.] M. Roem. Syn. Rosifl. 5, 89 (1847) = **Prunus**, Linn. (Rosac.).
acuminata, M. Roem. l. c. 92 = P. acuminata.
brachybotrys, M. Roem. l. c. 91 = P. brachybotrya.
brasiliensis, M. Roem. l. c. 90 = P. brasiliensis.
caroliniana, M. Roem. l. c. = P. caroliniana.
ferruginea, M. Roem. l. c. 91 = P. ferruginea.
Hixa, M. Roem. l. c. 92 = P. Hixa.
ilicifolia, M. Roem. l. c. = P. ilicifolia.
laurifolia, M. Roem. l. c. 91 = P. laurifolia, *Decne.*
lusitanica, M. Roem. l. c. 92 = P. lusitanica.
mexicana, M. Roem. l. c. 90 = P. laurifolia, *Schlecht.*
multiglandulosa, M. Roem. l. c. 92 = P. Hixa.
occidentalis, M. Roem. l. c. 89 = P. occidentalis.
officinalis, M. Roem. l. c. 91 = P. Laurocerasus.
opaca, M. Roem. l. c. = P. opaca.
Ottinii, Carr. in Rev. Hortic. (1877) 159 = P. Laurocerasus.
reflexa, M. Roem. Syn. Rosifl. 91 = P. reflexa.
salicifolia, M. Roem. l. c. 89 = P. salicifolia.
samydoides, M. Roem. l. c. 92 = P. samydoides.
sphaerocarpa, M. Roem. l. c. 89 = P. sphaerocarpa.
undulata, M. Roem. l. c. 92 = P. undulata.
vulgaris, Carr. in Rev. Hortic. (1877) 160 = P. Laurocerasus.

LAUROPHILLUS, Roem. & Schult. Syst. iii. 30 (1818) = seq.

LAUROPHYLLUS, Thunb. Nov. Gen. vi. 104 (1792) = **Botryceras**, Willd. (Anacard.).
capensis, Thunb. l. c. = B. laurinum.

LAURUS, [Tourn.] Linn. Gen. ed. I. 120 (1737).
LAURINEAE, Benth. & Hook. f. iii. 163.
AJOVEA, Rafin. Sylva Tellur. 134 (1838).
BALANOPSIS, Rafin. l. c. (1838).
abbreviata, *Salisb. Prod.* 344.—Am. bor.
acuminata, R. Br. ex Meissn. in DC. Prod. xv. i. 78 = Cryptocarya triplinervis.
acuminata, Reinw. ex Nees, Syst. Laurin. 611 = Lindera acuminata.
acuminata, Ruiz & Pav. ex Meissn. in DC. Prod. xv. i. 117 (= *Oreodaphne sublanuginosa*).—Peruv.
acuminata, Wall. ex Meissn. l. c. 34 = Phoebe pallida.
acutifolia, Pav. Laurin. Peruv. t. 17 = Nectandra berchemiaefolia.
aestivalis, Lam. Encyc. iii. 453 = Litsea geniculata.
aestivalis, Linn. Sp. Pl. 370 = Lindera Benzoin.

LAURUS:—
aestivalis, Mill. ex Willd. Sp. Pl. ii. 484 = Nectandra Willdenoviana.
aestivalis, Sesse, ex Meissn. in DC. Prod. xv. i. 193 = Litsea glaucescens.
aestivalis, Vell. Fl. Flum. 164; iv. t. 54 = Nectandra villosa.
aggregata, *Sims, ex Meissn. in DC. Prod.* xv. i. 516. —China.
alba, *Reinw. ex Miq. Fl. Ind. Bat.* i. ii. 582.—Java.
albida, Loud. ex Meissn. in DC. Prod. xv. i. 513, 516 = Sassafras officinale.
albiflora, Wall. Cat. n. 2569 = Cinnamomum Tamala.
alpigena, Sw. Fl. Ind. Occ. ii. 704 = Persea alpigena.
americana, Herb. Berol. ex Nees, Syst. Laurin. 135 = Persea indica.
americana, Mill. Gard. Dict. ed. VIII. n. 10 = Daphnopsis tinifolia, *Meissn.*
amygdalina, Buch.-Ham. ex Wall. Cat. n. 2585 = Cryptocarya amygdalina.
angusta, Rafin. Sylva Tellur. 136 = nobilis?
angustifolia, Willd. ex Nees, Syst. Laurin. 401 = Acrodiclidium salicifolium.
armeniaca, Poepp. ex Nees, l. c. 264 (= *Oreodaphne evonymodaphne*).—Peruv.
atra, Vell. Fl. Flum. 163; iv. t. 51 = Nectrandra puberula?
attenuata, Wall. Cat. n. 2600 = Phoebe attenuata.
aurantiodora, Ruiz & Pav. Fl. Per. iv. t. 349 (= *Mespilidaphne aurantiodora*).—Peruv.
australis, A. Cunn. ex Hook. Journ. Bot. iv. (1842) 436 = Cryptocarya australis.
axillaris, Lam. Encyc. iii. 453 = Litsea geniculata.
bulanocarpa, Ruiz & Pav. Fl. Per. iv. t. 354 (= *Mespilodaphne balanocarpa*.).—Peruv.
Barbujana, Cav. in Anal. Cienc. Nat. iii. (1801) 52 = Apollonias canariensis.
Barbusana, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 16 = canariensis.
Barbusano, Hochst. ex Webb & Berth. Phyt. Canar. iii. 229 = praec.
Bazania, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 560 = Cinnamomum obtusifolium.
Bejolghota, Buch.-Ham. l. c. 559 = Cinnamomum obtusiflorum.
Belloto, Miers, ex Nees, Syst. Laurin. 178 = Peumus Boldus.
Benzoin, Buch.-Ham. ex Wall. Cat. n. 2574 B = Cinnamomum obtusifolium.
Benzoin, Linn. Sp. Pl. 370 = Lindera Benzoin.
Bezolghota, Buch.-Ham. ex Wall. Cat. n. 2574 C = Cinnamomum obtusifolium.
bibracteata, Blume, Bijdr. 553 (= *Daphnidium bibracteatum*).—Java.
biflora, Pav. ex Meissn. in DC. Prod. xv. i. 119, 234 (= *Oreodaphne cuneata*).—Peruv.
bilocularis, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 311 = Beilschmiedia Roxburghiana.
Bofo, Willd. ex Nees, Syst. Laurin. 441 = Ocotea Bofo.
bogotensis, Willd. ex Nees, l. c. 163 = Persea Mutisii.
bombycina, Buch.-Ham. ex Wall. Cat. n. 2607 D = Machilus odoratissima.
Borbonia, Hort. ex Meissn. in DC. Prod. xv. i. 233 = canariensis.
Borbonia, Lam. Encyc. iii. 450 (= *Oreodaphne Martiniana*).—Guian.
Borbonia, Linn. Sp. Pl. 370 = Persea caroliniensis.
Borbonia, Sw. Fl. Ind. Occ. ii. 714 = Nectandra antillana.
Bowiei, Hook. Journ. Bot. iv. (1842) 419. t. 23 = Cryptocarya australis.
bullata, Burch. ex Nees, Syst. Laurin. 449 = Ocotea bullata.
Burmanni, Nees, Cinn. Disp. 57 = Cinnamomum Burmanni.
caerulea, Ruiz & Pav. Fl. Per. iv. t. 350 = Persea laevigata.
caesia, Reinw. ex Blume, Bijdr. 553 = Lindera caesia.
calicaris, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 380 (= *Tetranthera Tangao*).—N. Zel.
calophylla, Reinw. ex Nees, Cinn. Disp. 63 = Cinnamomum camphoratum.
calycaris, Soland. ex Meissn. in DC. Prod. xv. i. 191, 234 (= *Tetranthera Tangao*).—N. Zel.
calycina, Stokes, Bot. Mat. Med. ii. 421 = sumatrensis.
Camelilla, Willd. ex Meissn. in Mart. Fl. Bras. v. ii. 199 = Ocotea pretiosa.

LAURUS:—
Camphora, Linn. Sp. Pl. 369 = Cinnamomum Camphora.
Camphora, Thunb. ex Blume, Mus. Bot. Lugd. Bat. i. 346 = Litsea zeylanica.
camphorata, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe lanceolata.
camphorifera, Salisb. Prod. 343 = Cinnamomum Camphora.
canaliculata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 (= *Oreodaphne cayennensis*).—Guian.
canariensis, *Webb & Berth. Phyt. Canar.* iii. 3, 229.—In. Canar.
canariensis, Willd. Enum. Hort. Berol. i. 435 = Apollonias canariensis.
Canella, Mill. Gard. Dict. ed. VIII. n. 12 = Litsea zeylanica.
Canella, Vell. Fl. Flum. 164; iv. t. 56 (Quid?).—Bras.
Canellila, Willd. ex Nees, Syst. Laurin. 237 = Ocotea pretiosa.
carolinensis, Catesby, Carol. i. 63; Michx. Fl. Bor. Am. i. 265 = Persea carolinensis.
carolinensis, Hort. ex Meissn. in DC. Prod. xv. i. 233 = canariensis.
caryophyllata, Reinw. ex Meissn. in DC. l. c. 20, 235 = Cinnamomum iners.
Caryophyllus, Lour. Fl. Cochinch. 250 = Cinnamomum Culilawan.
Cassia, Burm. f. Fl. Ind. 91 = Cinnamomum zeylanicum.
Cassia, Heyne, ex Wall. Cat. n. 2583 D = Cinnamomum sulphuratum.
Cassia, Hort. ex Nees, Syst. Laurin. 42 = Cinnamomum iners.
Cassia, Linn. Sp. Pl. 369 = Litsea zeylanica, Cinnamomum zeylanicum.
Cassia, Nees, Diss. Laurin. 53 = Cinnamomum Cassia.
Cassia, Roxb. Hort. Beng. 30; Fl. Ind. ii. 297 = Cinnamomum Tamala.
Cassia, Wight, ex Nees, Syst. Laurin. 41 = Cinnamomum obtusifolium.
Catesbaei, Pers. Syn. i. 449 = Nectandra Willdenowiana.
Catesbyana, Michx. Fl. Bor. Am. i. 244 = Nectandra Willdenowiana.
Cathia, Buch.-Ham. ex Meissn. in DC. Prod. xv. i. 23, 235 = Phoebe paniculata.
caudata, Wall. Cat. n. 2603 = Cinnamomum caudatum.
caustica, Molina, Sagg. Chile, 176 = Mauria simplicifolia.
Cervantesii, H. B. & K. Nov. Gen. et Sp. i. 169 = Litsea glaucescens.
Champa, Buch.-Ham. ex Wall. Cat. n. 2607 C = Machilus odoratissima.
chinensis, Hort. Berol. ex Nees, Syst. Laurin. 92 = Cinnamomum Parthenoxylon.
chinensis, Meissn. in DC. Prod. xv. i. 211 = Actinodaphne chinensis.
Chloroxylon, Linn. Syst. ed. X. 1010 = Ceanothus Chloroxylon.
Chloroxylon, Sw. Prod. Veg. Ind. Occ. 260 (Cocculi sp.). —Ind. occ.
cinnamifera, Stokes, Bot. Mat. Med. ii. 412 = Cinnamomum zeylanicum?
cinnamomea, Salisb. Prod. 343 = Cinnamomum zeylanicum.
cinnamomoides, Hort. Berol. ex Nees, Syst. Laurin. 62 = Cinnamomum Burmanni.
cinnamomoides, Mutis, ex H. B. & K. Nov. Gen. et Sp. ii. 169 = Nectandra cinnamomoides.
Cinnamomum, Andr. Bot. Rep. t. 596 = Cinnamomum Cassia.
Cinnamomum, Blanco, Fl. Filip. ed. I. 319 = Cinnamomum Burmanni.
Cinnamomum, Linn. Sp. Pl. 369 = Cinnamomum zeylanicum.
Cinnamomum, Lour. Fl. Cochinch. i. 305 = Cinnamomum Loureirii.
Cinnamomum, Wight, ex Nees, in Wall. Pl. As. Rar. ii. 74 = Cinnamomum obtusifolium.
citriformis, Vell. Fl. Flum. 164; iv. t. 53 (Quid?).— Bras.
citriodora, Sieb. & Zucc. ex Meissn. in DC. Prod. xv. i. 516 (= *Aperula citriodora*).—Japon.
coccineus, Boj. ex Meissn. l. c. 511 (= *Mespilodaphne Lindleyana*).—Madag.
coerulea, Ruiz & Pav. ex Meissn. l. c. 49, 235 = Persea laevigata.
cordata, Vell. Fl. Flum. 163; iv. t. 50 = Persea pyrifolia.
coriacea, Sw. Prod. Veg. Ind. Occ. 65 = Nectandra Willdenoviana.

LAURUS:—
coruscans, Willd. ex Nees, Syst. Laurin. 305 = Nectandra cymbarum.
crassifolia, Poir. Encyc. Suppl. iii. 323 = Ocotea puberula.
crucifolia, Norenha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = Litsea sebifera.
Cubeba, Lour. Fl. Cochinch. 252 = Litsea Cubeba.
Cuipala, Buch.-Ham. ex Meissn. in DC. Prod. xv. I. 221, 235 = Litsea lanuginosa.
Culilaban, Blanco, Fl. Filip. ed. I. 315 = Cinnamomum pauciflorum.
Culilaban, Linn. Mant. ii. 237 = Cinnamomum Culilawan.
Culilawan, Hort. ex Nees, Syst. Laurin. 613 (= *Daphnidium gracile*).—Hab. ?
Culilawan, Reinw. ex Meissn. in DC. Prod. xv. I. 18 = Cinnamomum Tamala.
Culilawang, Nees, Disp. Cinnam. 61 = Cinnamomum Culilawan.
Culillaban, Buch.-Ham. in Wall. Cat. n. 2583 C = Cinnamomum iners ?
Culitlaban, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. ii. 74 = Cinnamomum zeylanicum.
Culitlaban, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 299 = Cinnamomum Culilawan.
cuneata, Domb. ex Nees, Syst. Laurin. 385 (= *Oreodaphne cuneata*).—Peruv.
cuneifolia, Ruiz & Pav. Fl. Per. iv. t. 359 (= *Oreodaphne cuneata*).—Peruv.
cuneifolia, Tafalla, ex Walp. Ann. iii. 312 (= *Oreodaphne obovata*).—Peruv.
cupularis, Lam. Encyc. iii. 447 (= *Mespilodaphne cupularis*).—Ins. Mascar.
cupularis, Pers. ex Meissn. in DC. Prod. xv. I. 105 (= *Mespilodaphne Persooniana*).—Ins. Mascar. ?
cupularis, Schott, ex Nees, Syst. Laurin. 308 = Nectandra paterifera.
curvifolia, Lour. Fl. Cochinch. 252 = Cinnamomum curvifolium.
cuspidata, D. Don, Prod. Fl. Nep. 64 = Lindera melastomacea.
cyathifera, Vahl, ex Meissn. in DC. Prod. xv. I. 165 = Nectandra Willdenowiana, sanguinea.
cymosa, Willd. ex Nees, Syst. Laurin. 437 (= *Mespilodaphne cupularis*).—Ins. Mascar.
cynamomum, *Vell. Fl. Flum.* 163; iv. t. 47.—Bras.
declinata, Blume, Cat. Gew. Buitenz. 66 = Phoebe declinata.
difformis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 (Ocoteae sp. ?).—Guian.
dioica, Wall. Cat. n. 7983 = Gelonium lanceolatum.
diospyroides, Michx. Fl. Bor. Am. i. 243 = Lindera melissaefolia.
Diospyros, Pursh, Fl. Am. Sept. i. 276 = Lindera melissaefolia.
discolor, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 (Quid ?).—Guian.
discolor, Willd. ex Nees, Syst. Laurin. 286 = Nectandra discolor.
divaricata, Poir. Encyc. Suppl. iii. 322 (= *Oreodaphne divaricata*).—Guian.
dubia, Wall. Cat. n. 2571 = Cinnamomun iners, multiflorum.
dulcis, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 303 = Cinnamomum Burmanni.
dysodantha, Ruiz & Pav. Fl. Per. iv. t. 355 (= *Goeppertia dysodantha*).—Peruv.
elegans, Blume, in Flora, vii. (1824) 291 (Quid ?).—Java.
elongata, Salisb. Prod. 344 = Persea carolinensis.
elongata, Vahl, ex Nees, Syst. Laurin. 116 = Phoebe elongata.
elongata, Willd. ex Nees, l. c. 117 = Nectandra patens.
enervia, *Mill. Gard. Dict.* ed. VIII. n. 8.—Am. bor.
Engelmanni, Steud. Nom. ed. II. ii. 16, lapsu = Lathyrus Engelmanni.
erythropus, Mart. Reise Bras. i. 401, nomen = Persea erythropus.
exaltata, Rudolph, ex Meissn. in DC. Prod. xv. I. 143 = Ocotea floribunda.
exaltata, Schott, ex Nees, Syst. Laurin. 322 = Nectandra concinna.
exaltata, Spreng. ex Nees, l. c. 310 = Nectandra leucantha.
exaltata, Sw. Prod. Veg. Ind. Occ. 65 = Nectandra sanguinea.

LAURUS:—
exaltata, Willd. ex Nees, Syst. Laurin. 412 = Ocotea botryophylla.
ferruginea, Ruiz & Pav. Fl. Per. iv. t. 372 = Persea ferruginea.
firma, Wall. Cat. n. 2597 = Endiandra firma.
flava, Reinw. ex Blume, Cat. Gew. Buitenz. 66 = Cryptocarya costata.
floribunda, Sw. Prod. Veg. Ind. Occ. 65 = Ocotea floribunda.
floribunda, Wall. Cat. n. 2593 = Cryptocarya amygdalina.
floribunda, Wall. l. c. n. 2593 B = Phoebe pallida.
foetens, Brouss. ex Meissn. in DC. Prod. xv. I. 65 = Apollonias canariensis.
foetens, [Soland. in] Ait. Hort. Kew. ed. I. ii. 39 = Ocotea foetens.
foetens, Willd. ex Nees, Syst. Laurin. 151 = Persea carolinensis.
fragrans, Salisb. Prod. 344 = Lindera Benzoin.
fruticosa, Reinw. ex Nees, Syst. Laurin. 543 = Litsea amara.
furcata, Ruiz & Pav. Fl. Per. iv. t. 367 = Nectandra furcata.
geminiflora, Desv. in Ham. Prod. Pl. Ind. Occ. 37 (= *Goeppertia geminiflora*).—Bras.
geminiflora, Reinw. ex Nees, Syst. Laurin. 102 = Phoebe excelsa.
geminiflora, Steud. Nom. ed. II. ii. 16, sphalm. = seq.
gemmiflora, Blume, Bijdr. 555 (= *Daphnidium gemmiflorum*).—Java.
gemmiflora, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 328 = Phoebe cuspidata.
geniculata, Michx. Fl. Bor. Am. i. 204; Walt. Fl. Carol. 133 = Litsea geniculata.
glandulifera, Jenkins, ex Meissn. in DC. Prod. xv. I. 26 = Cinnamomum Parthenoxylon.
glandulifera, Wall. in Act. Soc. Med. Phys. Calc. i. (1825) 45, 51. t. 1 = Cinnamomum glanduliferum.
glauca, Hort. Wuerz. ex Nees, Syst. Laurin. 613 (= *Daphnidium gracile*).—Hab. ?
glauca, Thunb. Fl. Jap. 173 = Litsea glauca.
glaucescens, Buch.-Ham. ex Wall. Pl. As. Rar. ii. 70 = Cinnamomum Cecidodaphne.
glaucescens, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 307 = Phoebe pallida.
glaucescens, Willd. ex Nees, Syst. Laurin. 538 = Litsea glaucescens.
globosa, Aubl. Pl. Gui. i. 364 = Nectandra sanguinea.
globosa, Lam. Encyc. iii. 451 = Nectandra antillana.
glomerata, Blume, Cat. Gew. Buitenz. 66 = Actinodaphne glomerata.
glomerata, Lam. Encyc. iii. 452 = Litsea geniculata.
gracilis, Hort. Berol. ex Meissn. in DC. Prod. xv. i. 230 (= *Daphnidium gracile*).—Hab. ?
gracilis, Hort. ex G. Don, in Loud. Hort. Brit. 160 = Cinnamomum Camphora.
granadensis, Forsyth, ex Meissn. in DC. Prod. xv. I. 164 = Nectandra sanguinea.
grandis, Wall. Cat. n. 2594 A = Phoebe opaca.
grandis, Wall. Cat. n. 2594 B = Alseodaphne costalis.
Gullavara, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. ii. 65 = Actinodaphne angustifolia.
Hamiltonii, *Spreng. Syst.* iv. *Cur. Post.* 155.—Reg. Himal.
heteranthera, Ruiz & Pav. Fl. Per. iv. t. 364 = Phoebe Ruiziana.
hexandra, Blanco, Fl. Filip. ed. II. 222 = Iteadaphne confusa.
hexandra, Spreng. ex Nees, Syst. Laurin. 420 (= *Mespiladaphne notata*).—Bras.
hexandra, Willd. Sp. Pl. ii. 482 = Aiouea guianensis.
Heyneana, Wall. Cat. n. 2576 = Cinnamomum Heyneanum, iners.
Hihua, Ruiz & Pav. Fl. Per. iv. t. 366 = Nectandra leucanthra.
hundaris, Nees, Syst. Laurin. 711, Index = seq.
hundensis, Willd. ex Nees, l. c. 139 = Phoebe cinnamomifolia.
hypericifolia, Willd. ex Nees, l. c. 165 = Persea hypericifolia.
iavitensis, Spreng. vide *javitensis*.
incrassata, Jack, in Malay. Misc. ii. (1822) n. VII. 33 = Dehaasia microcarpa.
indica, Linn. Sp. Pl. 370 = Persea indica.
indica, Lour. Fl. Cochinch. i. 253 = Machilus odoratissima.

LAURUS:—
indica, Sieber, ex Nees, Syst. Laurin. 128 = Persea gratissima.
indica, Thunb. Fl. Jap. 173 = Machilus Thunbergii.
iners, Reinw. ex Nees, in Wall. Pl. As. Rar. ii. 73 = Cinnamomum iners.
involucrata, Koen. in Retz. Obs. vi. 27 = Litsea sebifera.
involucrata, Lam. Encyc. iii. 445 = Litsea zeylanica.
javitensis, *H. B. & K. Nov. Gen. et Sp.* i. 169.—Venezuela.
javitensis, Spreng. ex Nees, Syst. Laurin. 145 = Persea pyrifolia.
Keule, Domb. ex Lam. Encyc. iii. 455 = Gomostega nitida.
laevigata, Willd. ex Meissn. in DC. Prod. xv. I. 49, 237 = Persea laevigata.
lanceifolia, Willd. ex Nees, Syst. Laurin. 316 = Nectandra turbacensis.
lanceolaria, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 309 = Phoebe lanceolata.
lanceolata, Wall. Cat. n. 2599, sphalm. = *lanceolaria*.
lanosa, Blanco, Fl. Filip. ed. I. 318 = Litsea villosa.
latifolia, Herb. Banks, ex Meissn. in DC. Prod. xv. I. 164, 237 = Nectandra sanguinea.
latifolia, Salisb. Prod. 343 = Persea indica.
leptobotra, Ruiz & Pav. Fl. Per. iv. t. 368 (= *Oreodaphne leptobotra*).—Peruv.
leucocarpa, Salzm. ex Meissn. in DC. Prod. xv. I. 250 = Sparattanthelium botocudorum.
Leucoxylon, Sw. Prod. Veg. Ind. Occ. 65 = Ocotea Leucoxylon.
Leucoxylon, Willd. ex Nees, Syst. Laurin. 141 = Phoebe antillana.
Ligi, Domb. ex Nees, l. c. 157 = Persea Lingue.
ligustrina, Wall. Cat. n. 2588 = Phoebe lanceolata.
ligustrina, Wall. l. c. n. 2588 A = Phoebe declinata.
limbosa, Ruiz & Pav. Fl. Per. iv. t. 361 = Nectandra limbata.
lineatifolia, Ruiz & Pav. l. c. t. 371 = Nectandra Amazonum.
Linge, Ruiz, ex Meissn. in DC. Prod. xv. I. 48 = seq.
Lingue, Ruiz & Pav. Fl. Per. iv. t. 358 = Persea Lingue.
Lingui, Bridges, ex Nees, Syst. Laurin. 672 = Peumus Boldus.
Lingui, Gill. ex Meissn. in DC. Prod. xv. I. 48 = Persea Meyeniana.
Linguy, Miers, ex Nees, Syst. Laurin. 157 = Persea Lingue.
longifolia, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 327 = Phoebe parviflora.
longifolia, Ruiz & Pav. Fl. Per. iv. t. 370 = Nectandra crassipes, longifolia, *Nees*.
longifolia, *Vahl, in West, Bidr. Ste-Croix*, 282, nomen (Quid ?).—Ind. occ.
longifolia, Willd. ex Meissn. in DC. Prod. xv. I. 112, 237 = Ocotea longifolia.
longipes, Schlecht. in Linnaea, vii. (1832) 390 = Persea longipes.
lucida, Thunb. Fl. Jap. 174 (non Laurinea).—Japon.
lucida, Wall. Cat. n. 2590 = Alseodaphne lucida.
ludoviciana, Rafin. Fl. Ludov. 27 (Quid ?).—Am. bor.
Machilus, Reinw. ex Miq. Fl. Ind. Bat. i. I. 929 = Dehaasia media.
macrantha, Wall. Cat. n. 2587 = Machilus macrantha.
macrophylla, A. DC. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 379 = Nesodaphne Tarairi.
macrophylla, D. Don, Prod. Fl. Nep. 64 = Lindera Neesiana, Dodecadenia grandiflora.
macrophylla, Wall. Cat. n. 2575 = Cinnamomum obtusifolium.
madagascariensis, Juss. ex Baill. in Bull. Soc. Linn. Par. i. (1885) 448 = Ravensara Topak.
Madang, Blume, Bijdr. 555 = Beilschmiedia Madang.
maderensis, Lam. Encyc. iii. 449 = Ocotea foetens.
malabathrica, Roxb. Hort. Beng. [30]; Soland. ex Roxb. Fl. Ind. ii. 297 = Cinnamomum iners.
Malabathrum, Reinw. ex Nees, in Wall. Pl. As. Rar. ii. 74 = Cinnamomum Cassia.
Malabathrum, Wall. Cat. n. 2583 = Cinnamomum iners.
Malabatrum, Burm. f. Fl. Ind. 92 = Cinnamomum javanicum.
Malabatrum, Lam. Encyc. iii. 445 = Cinnamomum iners.
martinicensis, Blanco, Fl. Filip. ed. I. 317 = Beilschmiedia Madang.
martinicensis, Jacq. Coll. ii. 109 = Oreodaphne Jacquini, Nectandra sanguinea.

LAURUS:—
martinicensis, Sieber, ex Meissn. in DC. Prod. xv. 1. 137 = Ocotea Sieberi.
mauritiana, Stadm. ex Soyer-Willem. in Usteri, Ann. Bot. xviii. (1796) 31; Boj. ex Meissn. l. c. 104 (= *Mespilodaphne cupularis*).—Ins. Mascar.
maxima, Salzm. ex Meissn. l. c. 83, 237 = Aiouea brasiliensis.
melissaefolia, Walt. Fl. Carol. 134 = Lindera melissaefolia.
membranacea, Sw. Prod. Veg. Ind. Occ. 65 = Nectandra membranacea.
microphylla, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 351 = Lindera acuminata.
Mimosa, Vell. Fl. Flum. 163; iv. t. 49 (Quid?).—Bras.
montana, Link, ex Meissn. in DC. Prod. xv. 1. 13, 237 = Cinnamomum zeylanicum.
montana, Poepp. ex Meissn. l. c. 31 = Phoebe antillana.
montana, Sw. Prod. Veg. Ind. Occ. 65 = Phoebe montana.
moschata, Pav. ex Meissn. in DC. Prod. xv. 1. 106, 237 (= *Mespilodaphne moschata*).—Peruv.
Muca, Ruiz & Pav. Fl. Per. iv. t. 360 = Aydendron Muca.
mucronata, Poir. Encyc. Suppl. iii. 323 = Cryptocarya mucronata.
multiflora, Roxb. Hort. Beng. [89]; Fl. Ind. ii. 298 = Cinnamomum multiflorum.
multiflora, Ruiz, ex Meissn. in DC. Prod. xv. 1. 150 = Nectandra Amazonum.
multiflora, Ruiz, ex Nees, in Linnaea, xxi. (1848) 501 = Nectandra canescens.
multiglandulosa, Ruiz & Pav. Fl. Per. iv. t. 365 (= *Oreodaphne Poeppigiana*).—Peruv.
Myrrha, Lour. Fl. Cochinch. 251 = Litsea trinervia?
Nacushia, Meissn. in DC. Prod. xv. 1. 231 = seq.
Nacusua, D. Don, Prod. Fl. Nep. 64 (Linderae sp.).—Nepal.
Naucushia, Buch.-Ham. ex D. Don, l. c. = praec.
Neesiana, Schott, ex Nees, Syst. Laurin. 488 (= *Mespilodaphne cupularis*).—Ins. Mascar.
nervosa, Pav. ex Meissn. in DC. Prod. xv. 1. 165 = Nectandra Willdenowiana.
nitida, Domb. ex Nees, Syst. Laurin. 385 (= *Oreodaphne cuneata*).—Peruv.
nitida, Heyne, ex Wall. Cat. n. 2582 B = Litsea zeylanica.
nitida, Masson, ex Meissn. in DC. Prod. xv. 1. 506 = Apollonias canariensis.
nitida, Roxb. Hort. Beng. [30]; Fl. Ind. iii. 300 = Cinnamomum iners.
nitida, Ruiz & Pav. Fl. Per. iv. t. 353 (= *Mespilodaphne nitida*).—Peruv.
nivea, Lodd. ex Loud. Hort. Brit. 161 (Quid?).—Hab.?
nobilis, Cav. in Anal. Cienc. Nat. iii. (1801) 55 = canariensis.
nobilis, *Linn. Sp. Pl.* 369.—Reg. Mediterr.
nutans, Hornem. ex Nees, Syst. Laurin. 424 = Ocotea Sieberi.
nutans, Isert, ex Nees, l. c. 141 = Phoebe cubensis.
oblongo-obovata, Ruiz, ex Nees, in Linnaea, xxi. (1848) 518 (= *Oreodaphne cuneata*).—Peruv.
obovata, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. ii. 64 = Actinodaphne obovata.
obovata, Ruiz & Pav. Fl. Per. iv. t. 357 (= *Oreodaphne obovata*).—Peruv.
obtusifolia, Pav. ex Meissn. in DC. Prod. xv. 1. 119 = Oreodaphne obovata.
obtusifolia, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 302 = Cinnamomum obtusifolium.
obtusifolia, Wall. Cat. n. 2574 = Cinnamomum Bazania.
obtusifolia, Willd. ex Meissn. in DC. Prod. xv. i. 94, 238 = Aydendron obtusifolium.
Ocotea, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Ocotea guianensis.
octandra, Vell. Fl. Flum. 164; iv. t. 57 (Quid?).—Bras.
odoratissima, Steud. Nom. ed. II. ii. 16 = *odorifera*.
odoratissima, Wall. Cat. n. 2607 = Machilus odoratissima.
odorifera, Vell. Fl. Flum. 164; iv. t. 58 = Aydendron suaveolens.
opifera, Meissn. in Mart. Fl. Bras. v. II. 195 = Ocotea opifera.
ovalifolia, Ruiz & Pav. Fl. Per. iv. t. 346 (= *Oreodaphne sublanuginosa*).—Peruv.

LAURUS:—
pallens, Meissn. in DC. Prod. xv. 1. 515 = Dodecadenia grandiflora.
palustris, Poepp. ex Meissn. l. c. 48 = Persea Lingue.
paniculata, Poir. Encyc. Suppl. iii. 323 = Phoebe cubensis.
paniculata, Wall. Cat. n. 2598 = Phoebe paniculata.
Parthenoxylon, Jack, in Malay. Misc. i. (1820) v. 28 = Cinnamomum Parthenoxylon.
parviflora, Balb. ex Meissn. in DC. Prod. xv. 1. 3 = Phoebe antillana.
parviflora, Poepp. ex Nees, Syst. Laurin. 120 = Acrodiclidium jamaicense.
parviflora, Pohl, ex Meissn. in DC. Prod. xv. 1. 98 (= *Mespilodaphne notata*).—Bras.
parviflora, Sw. Fl. Ind. Occ. ii. 717 (= *Oreodaphne parviflora*).—Jamaic.
parvifolia, *Lam. Encyc.* iii. 451 = Acrodiclidium jamaciense.
patens, Sw. Prod. Veg. Ind. Occ. 65 = Phoebe antillana.
pauciflora, Blume, Bijdr. 555 (= *Aperula confusa*).—Java.
pauciflora, Schlecht. ex Nees, Syst. Laurin. 405 = Nectandra nitidula.
pauciflora, Wall. Cat. n. 2579 = Cinnamomum pauciflorum.
pauperata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 (Quid?).—Guiana.
pedunculariis, Wall. Cat. n. 2596 (= *Haasea pedunculariis*).—Ind. or.
pedunculata, Thunb. Fl. Jap. 174 = Cinnamomum pedunculatum.
pendens, Willd. ex Nees, Syst. Laurin. 302 = Nectandra sanguinea.
pendula, Sw. Prod. Veg. Ind. Occ. 65 = Beilschmiedia pendula.
Persea, Linn. Sp. Pl. 370 = Persea gratissima.
Persea, Willd. ex Nees, Syst. Laurin. 441 = Ocotea Bofa.
Peumo, Domb. ex Lam. Encyc. iii. 455 = Cryptocarya Peumus.
Peumus, Miers, in Hook. & Arn. Bot. Beech. Voy. 45 = Cryptocarya Peumus.
Pichurim, Herb. Berol. ex Nees, Syst. Laurin. 458 = Ocotea polyantha.
Pichurim, Willd. ex Nees, Syst. Laurin. 249 = Aydendron Laurel.
pilosa, Lour. Fl. Cochinch. 253 = Machilus pilosa.
piperita, Meissn. in DC. Prod. xv. 1. 238 = Litsea Cubeba.
polyadelpha, *Lour. Fl. Cochinch.* 251.—Cochinch.
polyantha, Blume, Bijdr. 554 (= *Aperula polyantha*).—Java.
porrecta, Roxb. Hort. Beng. [30]; Fl. Ind. ii. 308 = Cinnamomum Parthenoxylon.
pruinosa, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 323 = Cinnamomum Parthenoxylon.
pruinosa, Wall. Cat. n. 2584 = Actinodaphne pruinosa.
pruinosa, Willd. ex Nees, Syst. Laurin. 146 = Persea laevigata.
Pseudo-Benzoin, Michx. Fl. Bor. Am. i. 243 = Lindera Benzoin.
Pseudo-Cassia, Herb. Zucc. ex Meissn. in DC. Prod. xv. 1. 503 = Cinnamomum Burmanni.
Pseudo-Cassia, Reinw. ex Nees, Syst. Laurin. 663 = Cinnamomum javanicum, camphoratum.
Pseudo-Sassafras, Blume, Bijdr. 573 = Cinnamomum Parthenoxylon.
psychotrioides, Willd. ex Nees, Syst. Laurin. 453 = Ocotea psychotrioides.
puberula, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Ocotea puberula.
pubescens, D. Don, ex Nees, Syst. Laurin. 588 = Dodecadenia grandiflora.
pubescens, Wall. Cat. n. 2595 = Phoebe pallida, paniculata.
pubescens, Willd. ex Nees, Syst. Laurin. 293 = Nectandra macrophylla.
Pucheri, Humb. ex Willd. l. c. 250 = Aydendron firmulum, Sellowii.
Puchiry, Ruiz & Pav. Fl. Per. iv. t. 352 = Nectandra limbata.
punctata, Ruiz & Pav. l. c. t. 347 = Nectandra punctata.
punctata, Spreng. Neue Entdeck. ii. 112 (al. ordin.).—Bras.
purpurea, Ruiz & Pav. Fl. Per. iv. t. 351 = Nectandra polita.
pyrifolia, Willd. ex Meissn. in Mart. Fl. Bras. v. II. 276 = Ocotea squarrosa.

LAURUS:—
pyrifolia, Willd. ex Nees, Syst. Laurin. 146 = Persea laevigata.
Quixos, Lam. Encyc. iii. 455 = Ocotea pretiosa.
recurvata, Roxb. ex Wall. Cat. n. 2564 B = Litsea umbrosa.
recurvata, Roxb. Hort. Beng [30]; Fl. Ind. ii. 301 = Cinnamomum pauciflorum.
reticulata, Poir. Encyc. Suppl. iii. 322 = Apollonias canariensis.
reticulata, Ruiz & Pav. Fl. Per. iv. t. 348 = Nectandra villosa.
retroflexa, Poir. Encyc. Suppl. iii. 323 = Ocotea floribunda.
revoluta, Willd. ex Nees, Syst. Laurin. 250 = Aydendron firmulum.
rigida, Wall. Cat. n. 2570 = Cinnamomum zeylanicum.
rigida, Willd. ex Nees, Syst. Laurin. 284 = Nectandra rigida.
rimosa, Blume, Bijdr. 574 = Machilus rimosa.
rimosa, Zoll. ex Meissn. in DC. Prod. xv. 1. 35 = Phoebe sumatrana.
robusta, Reinw. ex Zoll. in Flora, xxx. (1847) 474 = Litsea robusta.
rotundifolia, *Wall. Cat.* n. 2591.—China.
rubescens, Buek, Ind. ad DC. Prod. iv. 198, cit. falsa = *pubescens*?
Sailiana, Spreng. Syst. ii. 266 = seq.
Sailyana, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 558 = Cinnamomum Tamala.
salicifolia, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 379 (= *Nesodaphne Tawa*).—N. Zel.
salicifolia, Buch.-Ham. ex Nees, Syst. Laurin. 109 = Phoebe lanceolata.
salicifolia, Hort. ex Nees, l. c. 580 = nobilis.
salicifolia, Schott, ex Nees, l. c. 254 = Aydendron microbotryum.
salicifolia, Sw. Fl. Ind. Occ. ii. 709 = Acrodiclidium salicifolium.
salicifolia, Trevir. ex Nees, Syst. Laurin. 439 (= *Oreodaphne domingensis*).
salicifolia, Wight, ex Nees, l. c. 560 (= *Tetranthera glabrata*).—Ind. or.
salicifolia, Willd. ex Nees, l. c. 290 = Nectandra Willdenowiana.
Salsafraz, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19, sphalm.? = *L. Sassafras*
sanguinea, Sw. Fl. Ind. Occ. ii. 707 = Nectandra sanguinea.
sarcocalyx, Mart. ex Meissn. in DC. Prod. xv. 1. 167, 239 = Nectandra nitidula.
Sassafras, Linn. Sp. Pl. 371 = Sassafras officinale.
Sassafras, Lour. Fl. Cochinch. 254 = Sassafras Parthenoxylon.
semecarpifolia, Wall. Cat. n. 2586 = Alseodaphne semicarpifolia.
sericea, *Blume, Bijdr.* 554.—Java.
sericea, Hook. ex Nees, Syst. Laurin. 369 = Aydendron sericeum.
sericea, Wall. Cat. n. 2606 = Machilus sericea.
sericea, Willd. ex Nees, Syst. Laurin. 162 = Persea sericea.
serrata, Blanco, Fl. Filip. ed. I. 319 = Casearia glomerata.
smilacifolia, Wall. Cat. n. 2572 = Grewia laurifolia.
Soilyana, Buch.-Ham. ex Wall. l. c. n. 2569 D = Cinnamomum Tamala.
solaniformis, Vell. Fl. Flum. 163; iv. t. 52 (Quid?).—Bras.
Soncaurium, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 557 = Cinnamomum Tamala.
splendens, Forsyth, ex Meissn. in DC. Prod. xv. 1. 174, 239 = Aydendron sericeum.
splendens, Willd. ex Meissn. in Mart. Fl. Bras. v. II. 276 = Nectandra sanguinea.
splendens, Willd. ex Nees, Syst. Laurin. 302 = Nectandra salicifolia.
subcordata, Ruiz & Pav. Fl. Per. iv. t. 369 = Persea subcordata.
sublanuginosa, Ruiz, ex Nees, in Linnaea, xxi. (1848) 515 (= *Oreodaphne sublanuginosa*).—Peruv.
subscandens, Vell. Fl. Flum. 164; iv. t. 59 (Quid?).—Bras.
subumbelliflora, Blume, Bijdr. 534 (= *Aperula confusa*).—Java.
sulcata, Ruiz & Pav. Fl. Per. iv. t. 356 = Persea sulcata.
sumatrensis, J. F. Gmel. Syst. 650 = Cinnamomum Camphora?

LAURUS:—
Surca, Willd. ex Nees, Syst. Laurin. 478 = Ocotea guianensis.
surinamensis, Sw. Fl. Ind. Occ. ii. 708 = Ocotea guianensis.
sylvestris, Buch.-Ham. ex Roxb. Fl. Ind. ii. 307 = Phoebe pallida.
Tama, Meissn. in DC. Prod. xv. 1. 66, sphalm. = *Tawa*.
Tamala, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 533, excl. syn. = Cinnamomum Tamala.
Tarairi, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 379 (= *Nesodaphne Tarairi*).—N. Zel.
Tawa, A. Cunn. l. c. (= *Nesodaphne Tawa*).—N. Zel.
Tazia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 557 = Cinnamomum Tamala.
Teneriffae, Poir. Encyc. Suppl. iii. 322 = Persea indica.
tenuifolia, Mill. Gard. Dict. ed. VIII. n. 3 = nobilis.
thyrsiflora, Poir. Encyc. iii. 322 (= *Hufelandia thyrsiflora*).—Madag.
tigricida, Mutis, ex Nees, Syst. Laurin. 286 = Nectandra discolor.
Tihua, Ruiz, ex Meissn. in DC. Prod. xv. 1. 151, sphalm. = *L. Hihua*.
Till, Poir. Encyc. Suppl. iii. 324 = Ocotea foetens.
tomentosa, Buch.-Ham. ex Meissn. in DC. Prod. xv. 1. 23, 240 = Cinnamomum tomentosum.
tomentosa, Hort. ex Meissn. l. c. 233 = canariensis.
triandra, Sw. Prod. Veg. Ind. Occ. 65 = Acrodiclidium jamaicense.
triplinervia, Reinw. ex Wall. Pl. As. Rar. ii. 75 = Cinnamomum Tamala.
triplinervis, Ruiz & Pav. Fl. Per. iv. t. 363 = Phoebe peruviana.
umbellata, Buch.-Ham. ex Spreng. Syst. iv. Cur. Post. 155 = Hamiltonii.
umbellata, Thunb. Fl. Jap. 175 = Rhaphiolepis japonica.
umbrosa, Willd. ex Nees, Syst. Laurin. 456 = Ocotea umbrosa, *H. B. & K.*
undulata, Mill. Gard. Dict. ed. VIII. n. 2 = nobilis.
variifolia, Salisb. Prod. 344 = Sassafras officinale.
venosa, Domb. ex Nees, Syst. Laurin. 291 = Nectandra villosa.
verticillata, Royen, ex Blume, Mus. Bot. Lugd. Bat. i. 346 = Litsea zeylanica.
viburnoides, Willd. ex Nees, Syst. Laurin. 148 = Persea laevigata.
Victoriana, Colenso, ex Hook. f. Handb. N. Zeal. Fl. 239 (= *Nesodaphne Tawa*).—N. Zel.
villosa, Roxb. Hort. Beng. [89]; Fl. Ind. ii. 310 = Phoebe pallida.
villosa, Willd. ex Nees, Syst. Laurin. 287 = Nectandra villosa.
viminea, Wall. Cat. n. 2578 = Cinnamomum vimineum.
virgultosa, Mart. ex Nees, Syst. Laurin. 304 = Ocotea virgultosa.
Winterana, Linn. Sp. Pl. 371 = Canella alba.
zeylanica, Lam. ex Meissn. in DC. Prod. xv. 1. 240 = Litsea zeylanica.

LAUSONIA, Juss. Gen. 331 (1789) = **Lawsonia**, Linn. (Lythrar.).

LAUTEMBERGIA, Baill. Etud. Gén. Euph. 451 (1858) = **Alchornea**, Sw. (Euphorb.).
multispicata, Baill. l. c. 452 = A. multispicata.

LAVALLEA, Baill. Adansonia, ii. (1862) 359 = **Strombosia**, Blume (Olacin.).
philippinensis, Baill. l. c. = S. philippinensis.
zeylanica, Baill. l. c. = S. ceylanica.

LAVANDULA, Tourn. ex Linn. Syst. ed. II. 25 (1740). *LABIATAE*, Benth. & Hook. f. ii. 1179.
CHAETOSTACHYS, Benth. in Wall. Pl. As. Rar. ii. 19 (1831).
FABRICIA, Adans. Fam. ii. 188 (1763).
STOECHAS, Tourn. ex Linn. Syst. ed. I (1735).
STYPHONIA, Medic. Phil. Bot. ii. 70 (1791).
abrotanoides, *Lam. Encyc.* iii. 429.—Ins. Canar.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = vera.
angustifolia, Moench, Meth. 389; Risso, Hist. Nat. Eur. Mér. ii. 448 = L. Spica.
apiifolia, C. Sm. in Tuckey, Congo, 250, nomen = rotundifolia?
atriplicifolia, *Benth. in DC. Prod.* xii. 146.—Aegypt.
Buchii, Webb & Berth. Phyt. Canar. iii. 58 = pinnata.

LAVANDULA:—
Burmanni, *Benth. Lab. Gen. et Sp.* 151.—Ind. or.
candriensis, Mill. Gard. Dict. ed. VIII. n. 4 = abrotanoides.
cariensis, *Boiss. Diagn.* Ser. I. v. 3.—As. Min.
carnosa, Linn. f. Suppl. 273 = Anisochilus carnosus.
coronipifolia, *Poir. Encyc.* iii. 308.—Aegypt.; Oriens.
delphinensis, Jord. ex Billot, Cent. 27, 28 (1859) [text], 4 = L. Spica.
dentata, *Linn. Sp. Pl.* 572.—Reg. Mediterr.
elegans, Desf. Tabl. ed. I. 59 = abrotanoides.
erigens, Jord. & Fourr. Brev. Pl. Nov. fas. ii. 88 = L. Spica.
formosa, F. G. Dietr. ex Link, Enum. Hort. Berol. ii. 103 = pinnata.
fragrans, Salisb. Prod. 78 = L. Spica.
Gibsoni, *J. Grah. Cat. Bomb. Pl.* 149.—Ind. or.
heterophylla, Hort. Par. ex Poir. Encyc. Suppl. iii. 308 = dentata, vera.
heterophylla, *Viv. Elench. Pl. Hort. Dinegro*, 23.—Hab.?
incana, Salisb. Prod. 78 = L. Stoechas.
inclinans, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 88 = L. Spica.
interrupta, Jord. & Fourr. l. c. 89 = L. Spica.
lanata, *Boiss. Elench.* 72.—Hispan.
latifolia, Medic. Beobacht. (1783) 135; Vill. Hist. Pl. Dauph. ii. 363; Ehrh. Beitr. vii. 55 = L. Spica.
Lawii, Wight, Ic. t. 1439 = Gibsoni.
Massoni, Cels, ex Ging. Hist. Lav. 137 = viridis.
Minutolii, *C. Bolle, in Bonplandia*, viii. (1860) 279.—Ins. Canar.
multifida, Burm. f. Fl. Ind. 126 = Burmanni.
multifida, Burm. f. l. c. t. 38. f. 1 = coronopifolia.
multifida, *Linn. Sp. Pl.* 572.—Reg. Mediterr. occ.
Nimmoi, *Benth. in DC. Prod.* xii. 148.—Ins. Socotr.?
officinalis, Chaix, in Vill. Hist. Pl. Dauph. i. 355; iii. 363 = vera.
ovata, Steud. Nom. ed. II. ii. 17 = L. Spica.
pedunculata, *Cav. Desc.* 70.—Hispan.; Ins. Mader.
Perrottetii, Benth. in DC. Prod. xii. 147 = Gibsoni.
pinnata, *Linn. f. Diss. Lav.* 55.—Ins. Mader. et Canar.
pinnata, Moench, Meth. Suppl. 135 = dentata.
pinnatifida, Salisb. Prod. 79 = pinnata.
pinnatifida, Webb, Iter Hisp. 19 = multifida.
Pseudostoechas, Reichb. ex Holl, in Flora, xii. (1829) 691 = L. Stoechas.
pubescens, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 246.—Aegypt.; Arab.
pyrenaica, DC. Fl. Fr. v. 398 = vera.
rodundifolia, *Benth. Lab. Gen. et Sp.* 150.—Ins. Cap. Virid.
santolinaefolia, *Spach, in Faub. & Spach, Illustr.* iv. iii. t. 373.—Oriens.
setifera, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* 1. (1860) 29.—Arab.
spectabilis, C. Koch, in Linnaea, xxi. (1848) 646 = cariensis.
Spica, *Cav. Desc.* 69; *DC. Fl. Fr.* v. 397.—Reg. Mediterr.
Spica, Loisel. Fl. Gall. ii. 346 = vera.
stoechadensis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = seq.
Stoechas, *Linn. Sp. Pl.* 573.—Reg. Mediterr.
striata, Boiss. Fl. Orient. iv. 542, sphalm. = seq.
stricta, Delile, Fl. Egypte, 238. t. 32 = coronopifolia.
subnuda, *Benth. in DC. Prod.* xii. 148.—Arab.
tenuisecta, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 65, nomen.—Marocc.
vera, *DC. Fl. Fr. Suppl.* 398.—Reg. Mediterr.
viridis, *L'Hérit. Sert. Angl.* 19.—Lusitan.; Ins. Mader.
vulgaris, Lam. Fl. Fr. ii. 403 = vera.

LAVANGA, Meissn. Gen. 46; Comm. 34 (1837), sphalm. = **Luvunga**, Buch.-Ham. (Rutac.).

LAVATERA, Linn. Gen. ed. I. 205 (1737). *MALVACEAE*, Benth. & Hook. f. i. 200.
ANTHEMA, Medic. Malv. 42 (1787).
AXOLOPHA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258.
NAVAEA, Webb, Phyt. Canar. i. 32. t. 1 C (1836).
OLBIA, Medic. Malv. 41 (1787).
SAVINIONA, Webb, Phyt. Canar. i. 30. t. 1 B (1836).
STEGIA, DC. Fl. Fr. iv. 835 (1805).
acerifolia, *Cav. in Anal. Cienc. Nat.* vi. (1803) 339.—Ins. Canar.

LAVATERA:—
acutifolia, Lam. Fl. Fr. iii. 137 = L. Olbia.
africana, Cav. Diss. v. 282. t. 139. f. 1 = L. Olbia.
africana, Mill. Gard. Dict. ed. VIII. n. 2 = trimestris.
agrigentina, Tineo, Pl. Rar. Sic. Pugill. i. 13 = flava.
alba, Medic. Malv. 41 = trimestris.
albida, Boj. Hort. Maurit. 26 (Quid?).—Hab.?
altheaefolia, Mill. Gard. Dict. ed. VIII. n. 1 = trimestris.
ambigua, DC. Prod. i. 440 = thuringiaca.
americana, Linn. Syst. ed. X. 1148 = Abutilon Jacquini.
arborea, *Linn. Sp. Pl.* 690.—Europ. occ.; Afr. bor.
assurgentiflora, *Kellogg, in Proc. Calif. Acad.* i. (1854) 4.—Calif.
australis, A. Cunn. ex Hook. f. in Hook. Journ. Bot. ii. (1840) 412 = plebeia.
Behriana, Schlecht. in Linnaea, xxiv. (1851) 699 = plebeia.
biennis, *Bieb. Fl. Taur. Cauc.* ii. 143.—Reg. Caucas.
biennis, Guss. Fl. Sic. Prod. ii. 347 = punctata.
biflora, E. Mey. ex Harv. & Sond. Fl. Cap. i. 166 (= *Sphaeroma Julii*).—Afr. austr.
brachyfolia, *Walp. Rep.* i. 291.—Ins. Canar.
bryoniaefolia, Mill. Gard. Dict. ed. VIII. n. 11 = unguiculata.
cachemiriana, *Cambess. in Jacquem. Voy.* iv. 29.—Reg. Himal.
calicina, Poir. Encyc. Suppl. iii. 310 = triloba.
coccinea, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. iv. 329 = phoenicea.
cretica, *Linn. Sp. Pl.* 691.—Reg. Mediterr.
Empedoclis, Ucria, in Roem. Arch. i. 1. 69 = cretica.
eriocalyx, Steud. in Flora, xxxix. (1856) 438 = arborea.
flava, *Desf. Fl. Atlant.* ii. 119. t. 172.—Reg. Mediterr.
gallica, Hort. Par. ex Nym. Consp. 128 = maritima.
grandiflora, Lam. Fl. Fr. iii. 137; Moench, Meth. 615 = trimestris.
hirsuta, Mill. Gard. Dict. ed. VIII. n. 5 = Malva plicata.
hispanica, Mill. l. c. n. 9 = maritima.
hispida, Desf. Fl. Atlant. ii. 118. t. 171 = L. Olbia.
hispida, Sims, Bot. Mag. t. 2269 = plebeia.
insularis, *S. Wats. in Proc. Am. Acad.* xii. (1877) 249.—Calif.
Julii, Burch. ex DC. Prod. i. 438 (= *Sphaeroma Julii*).—Afr. austr.
kashmiriana, Mast. in Hook. f. Fl. Brit. Ind. i. 319 = cachemiriana.
lanata, Steud. Nom. ed. II. ii. 18, sphalm. = Lavandula lanata.
lanceolata, *Willd. Enum. Hort. Berol.* 733.—Hab.?
lusitanica, Auct. ex Hoffmgg. Verz. Pfl. Nachtr. 146 = septembola.
lusitanica, *Linn. Sp. Pl.* 691.—Lusitan.
maritima, *Gouan, Illustr.* 46. t. 11. f. 2.—Reg. Mediterr. occ.
mauritanica, *Duch. Revue Bot.* ii. (1846–47) 436.—Afr. bor.
meonantha, Link, Handb. ii. 359 = trimestris?
micans, *Linn. Sp. Pl.* 690.—Hispan.
minoricensis, Cambess. in Mém. Mus. Par. xiv. (1827) 334 = triloba.
mollis, Ehrenb. ex Sweet, Hort. Brit. ed. II. 60 = cretica?
moschata, Moris, Elench. Fl. Sard. 9 = triloba.
muricata, *Panč. in Verh. Zool. Bot. Ver. Wien*, vi. (1856) 497.—Servia.
neapolitana, Tenore, Prod. Fl. Nap. p. lxii = cretica.
oblongifolia, *Boiss. in Bibl. Univ. Genèv.* (1838) 111. t. 33.—Hispan.
occidentalis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 113, 125.—Calif.
Olbia, *Linn. Sp. Pl.* 690.—Europ. austr.
Olbia, Sibth. & Sm. Fl. Graec. Prod. ii. 46 = unguiculata.
Olbia, Steph. Fl. Mosq. n. 469 = thuringiaca.
pallescens, Moris, Fl. Sard. i. 301 = triloba.
phoenicea, *Vent. Jard. Malm.* 120. t. 120.—Ins. Madera.
phoenicea, Willd. Enum. Hort. Berol. Suppl. 49 = acerifolia.
plebeia, *Sims, Bot. Mag.* t. 2269.—Austral.
prostrata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 166 (= *Sphaeroma prostrata*).—Afr. austr.
Pseudo-olbia, Desf. Tabl. ed. I. 145 = L. Olbia?
Pseudo-olbia, Poir. Encyc. Suppl. iii. 309 = L. Olbia.
punctata, *All. Auctuar.* 26.—Europ. austr.; As. Min.

LAVATERA:—
ramosissima, *Opiz, Natural.* x. (1825) 192; *et ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 187. —Hab.?
rosea, Medic. Malv. 40 = trimestris.
rotundata, Láz. in Trab. Soc. Linn. Matr. (1880) 33 = triloba.
rotundifolia, Lam. Fl. Fr. iii. 138 = maritima.
rupestris, Pomel, Nouv. Mat Fl. Atl. 343 = triloba.
salvitellensis, Brign. ex Loud. Hort. Brit. Suppl. ii. 644; Hort. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 186 = thuringiaca.
scabra, *Hort. ex Turcz. l. c.* 187.—Hab.?
septemloba, *Hoffmgg. Verz. Pfl. Nachtr.* 146.—Hab.?
sicula, Tineo, Pl. Rar. Sic. Pugill. i. 14 = cretica.
subovata, *DC. Prod.* i. 439.—Mauritan.
sylvestris, Brot. Fl. Lusit. ii. 277 = cretica.
sylvestris, Cyrilli, ex Tenore, Prod. p. xl = thuringiaca.
Thirkeana, C. Koch, in Linnaea, xix. (1847) 58 = thuringiaca.
thuringiaca, All. Fl. Pedem. ii. 42 = L. Olbia.
thuringiaca, *Linn. Sp. Pl.* 691.—Europ. austr.; As. Min.
thuringiaca, Savi, Fl. Pis. ii. 126 = punctata.
tomentosa, Dum.-Cours. Bot. Cult. ed. II. v. 74 = unguiculata.
triloba, Gouan, Fl. Monsp. 48 = maritima.
triloba, Jacq. Hort. Vindob. 30. t. 74 = L. Olbia.
triloba, *Linn. Sp. Pl.* 691.—Hispan.
triloba, Sebast. & Mauri, Fl. Rom. 227 = cretica.
trimestris, *Linn. Sp. Pl.* 692.—Reg. Mediterr.
tripartita, *DC. Prod.* i. 440.—Hab.?
undulata, Desf. Arb. i. 471 = L. Olbia.
undulata, Mill. Gard. Dict. ed. VIII. n. 10 = lusitanica.
unguiculata, *Desf. Arb.* i. 471.—Sicil.; Graecia; Ins. Samos.
veneta, Mill. Gard. Dict. ed. VIII. n. 6 = arborea.
venosa, *S. Wats. in Proc. Am. Acad.* xii. (1877) 249. —Am. bor. occ.
vitifolia, Wierzb. ex Schur, Enum. Pl. Transs. 129 = thuringiaca.
vulgaris, H. Mart. Fl. Mosq. 121 = thuringiaca.
Weinmanniana, Bess. Hort. Cremen. (1823); ex Reichb. Ic. Bot. Exot. 45. t. 60 = plebeia.
Weinmanniana, Trev. ex DC. Prod. i. 440 = cretica.

LAVAUXIA, Spach, Hist. Vég. Phan. iv. 366 (1835) = **Oenothera**, Linn. (Onagrar.).
centaurifolia, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 359 = O. centaurifolia.
cuspidata, Spach, l. c. 358 = O. acaulis.
mutica, Spach, l. c. 359 = O. acaulis.
Nuttalliana, Spach, l. c. 358 = O. triloba.

LAVENDULA, Linn. Syst. ed. I (1735) = **Lavandula**, Tourn. (Labiat.).

LAVENIA, Sw. Prod. Veg. Ind. Occ. 112 (1788); Fl. Ind. Occ. 1329 (1806) = **Adenostemma**, Forst. (Compos.).
alba, Wall. Cat. n. 3220 = A. viscosum.
aquatica, Spreng. Syst. iii. 445 = A. viscosum.
carnosa, Wall. Cat. n. 3219 = A. viscosum.
decumbens, Spreng. Syst. iii. 445 = A. Berterii.
decumbens, Sw. Prod. Veg. Ind. Occ. 112 = A. Swartzii.
dentata, Wall. Cat. n. 3216 = Myriactis nepalensis.
divaricata, Buch.-Ham. ex DC. Prod. v. 111 = A. viscosum.
elata, Spreng. Syst. iii. 445 = A. viscosum.
erecta, Sw. Prod. Veg. Ind. Occ. 112 = A. viscosum.
fastigiata, Blume, Bijdr. 905 = A. viscosum.
glutinosa, Gaudich. in Freyc. Voy. Bot. 471 = A. glutinosum.
hirsuta, Blume, Bijdr. 905 = A. hirsutum.
latifolia, Spreng. Syst. iii. 445 = A. viscosum.
macrophylla, Blume, Bijdr. 905 = A. viscosum.
montana, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 185 = A. viscosum.
nudiflora, Hort. Calc. ex DC. Prod. v. 112 = A. viscosum.
parviflora, Blume, Bijdr. 905 = A. viscosum.
rigida, Wall. Cat. n. 3215 = Rhynchospermum verticillatum.
rugosa, Wight, in Wall. l. c. n. 3221 = A. viscosum.
sphaerantha, Wall. l. c. n. 3217 = Myriactis nepalensis.
viscida, Buch.-Ham. in Wall. l. c. n. 3222 = A. viscosum.

LAVOISERIA, Spreng. Gen. i. 341 (1830) = seq.

LAVOISIERA, DC. Prod. iii. 102 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 738.
LAVOISERIA, Spreng. Gen. i. 341 (1830).
alba, *DC. Prod.* iii. 103.—Bras.
australis, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 151.—Bras.
Bergii, *Cogn. in Mart. Fl. Bras.* xiv. iii. 154.—Bras.
bicolor, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 151.—Bras.
Blanchetiana, Naud. l. c. 150 = Microlicia Blanchetiana.
blepharocentra, Mart. in DC. Prod. iii. 103 = imbricata.
caryophyllea, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 150.—Bras.
cataphracta, DC. Prod. iii. 102 = imbricata.
centiformis, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 151 = imbricata.
cerifera, *Gardn. in Field. Sert. Pl.* t. 63.—Bras.
Chamaepitys, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 153.—Bras.
compta, DC. Prod. iii. 103 = imbricata.
confertiflora, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 149.—Bras.
cordata, *Cogn. in Mart. Fl. Bras.* xiv. iii. 140.—Bras.
crassifolia, *DC. Prod.* iii. 104.—Bras.
elegans, *Cogn. in Mart. Fl. Bras.* xiv. iii. 160.—Bras.
firmula, *DC. Prod.* iii. 104.—Bras.
Francavillana, *Cogn. in Mart. Fl. Bras.* xiv. iii. 158. —Bras.
Gardneri, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 221 = mucorifera.
gentianoides, *DC. Prod.* iii. 104.—Bras.
glandulifera, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 149.—Bras.
glandulifolia, Naud. l. c. iii. (1845) 192 = praec.
Glazioviana, *Cogn. in Mart. Fl. Bras.* xiv. iii. 157.— Bras.
glutinosa, *Cogn. l. c.* 145.—Bras.
grandiflora, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 148.—Bras.
humilis, *Naud. l. c.* 153.—Bras.
imbricata, Benth. in Hook. Journ. Bot. ii. (1840) 300 = Glazioviana.
imbricata, *DC. Prod.* iii. 103.—Bras.
insignis, *DC. l. c.*—Bras.
itambana, *DC. l. c.* 104.—Bras.
linifolia, DC. l. c. = Microlicia linifolia.
lycopodioides, Gardn. ex Hook. Ic. Pl. t. 502 = tetragona.
macrocarpa, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 148.—Bras.
microlicioides, *Naud. l. c.* xii. (1849) 221.—Bras.
microphylla, Naud. l. c. ii. (1844) 150 = Microlicia decipiens.
mucorifera, *DC. Prod.* iii. 103.—Bras.
nervulosa, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 149.—Bras.
pectinata, *Cogn. in Mart. Fl. Bras.* xiv. iii. 155.— Bras.
phyllocalycina, *Cogn. l. c.* 156.—Bras.
Pohliana, *Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 30.—Bras.
pulchella, *Champ. in Linnaea*, ix. (1834) 370.—Bras.
pulchella, Triana, in Trans. Linn. Soc. xxviii. (1871) 30, partim = australis.
pulcherrima, *DC. Prod.* iii. 104.—Bras.
punctata, *DC. l. c.*—Bras.
reticulata; Moric. Pl. Nouv. Am. 170. t. 99 = nervulosa.
Riedeliana, *Cogn. in Mart. Fl. Bras.* xiv. iii. 141.— Bras.
rigida, *Cogn. l. c.* 144.—Bras.
scaberula, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 151.—Bras.
scabrella, Naud. l. c. xii. (1849) 226 = scaberula.
Selloana, *Cogn. in Mart. Fl. Bras.* xiv. iii. 152.— Bras.
serrulata, *Cogn. l. c.* 153.—Bras.
subulata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 30.—Bras.
tetragona, *DC. Prod.* iii. 103.—Bras.
viminalis, DC. l. c. = Microlicia viminalis.

LAVRADIA, Swediaur, Arzneymitt. i. (1801) 111, in syn. = LABRADIA, Swediaur = **Mucuna**, Adans. (Legumin.).

LAVRADIA, Vell. ex Vand. Fl. Lusit. 15. t. 1. f. 6 (1788); et Roem. Script. Lus. et Bras. 88. t. 6. f. 6. *VIOLARIEAE*, Benth. & Hook. f. i. 120.
LAURADIA, Vand. l. c., in textu (1788).
alpestris, *Mart. Nov. Gen. et Sp.* i. 32. t. 22.—Bras.
capillaris, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 113.—Bras.
elegantissima, *A. St. Hil. l. c.* 109.—Bras.
ericoides, *A. St. Hil. l. c.* 108.—Bras.
glandulosa, A. St. Hil. l. c. 112 = montana.
montana, *Mart. Nov. Gen. et Sp.* 33. t. 23.—Bras.
Vellozia, A. St. Hil. in Mém. Mus. Par. xi. (1824) 110 = Velloziana.
Velloziana, *Vand. ex Steud. Nom.* ed. I. 466.—Bras.

LAWIA, Griff. ex Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 112 = **Terniola**, Tul. (Podost.).
longipes, Tul. l. c. 113 = T. longipes.
pulchella, Tul. l. c. = T. pulchella.
zeylanica, Tul. l. c. 112 = T. zeylanica.

LAWIA, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 14 = **Adenosacme**, Wall. (Rubiac.).
acuminata, Wight, l. c. 15; et Ic. t. 1070 = A. Lawii.

LAWRENCELLA, Lindl. Swan River App. 23 (1839) = **Helichrysum**, Vaill. (Compos.).
rosea, Lindl. l. c. = H. Lawrencella.

LAWRENCIA, Hook. Ic. Pl. t. 261 (1840) = **Plagianthus**, Forst. (Malvac.).
glomerata, Hook. l. c. t. 417 = P. glomeratus.
spicata, Hook. l. c. t. 262 = P. spicatus.
squamata, Nees, in Lehm. Pl. Preiss. i. 242 = P. squamatus.

LAWSONIA, Linn. Gen. ed. I. 111 (1737). *LYTHRARIEAE*, Benth. & Hook. f. i. 782.
ALCANNA, Gaertn. Fruct. ii. 133. t. 110 (1791).
PONTALETSJE, Adans. Fam. ii. 159 (1763).
Achronychia, Linn. f. Suppl. 219; Labill. Sert. Austro-Caled. 66. t. 65 = Acronychia laevis.
alba, *Lam. Encyc.* iii. 106.—Oriens.
coccinea, Sm. in Rees, Cycl. xx. n. 3 = alba.
falcata, Lour. Fl. Cochinch. 229 = alba?
falcifolia, Stokes, Bot. Mat. Med. ii. 364 = alba?
inermis, Linn. Sp. 349 = alba.
purpurea, Lam. Encyc. iii. 107 = alba.
spinosa, Linn. Sp. Pl. 349 = alba.

LAXANON, Rafin. New Fl. Am. ii. 16 (1836) = **Apogon**? Ell. (Compos.).
diversifolium, Rafin. l. c. ii. 46.—Am. bor.
heterophylum, Rafin. l. c. iv. 106.—Am. bor.
parviflorum, Rafin. l. c. ii. 16.—Am. bor.

LAXMANNIA, R. Br. Prod. 285 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 796.
acuta, Endl. in Lehm. Pl. Preiss. ii. 42 = squarrosa.
brachyphylla, *F. Muell. Fragm.* i. 158.—Austral.
gracilis, *R. Br. Prod.* 286.—Austral.
grandiflora, *Lindl. Swan River App.* 56. t. 7.— Austral.
grandiflora, F. Muell. Fragm. vii. 88 = squarrosa.
illecebrosa, Reichb. f. Beitr. Syst. Pfl. 72 = gracilis.
minor, *R. Br. Prod.* 286.—Austral.
minor, Hook. f. Fl. Tasm. ii. 60 = sessiliflora.
paleacea, F. Muell. Fragm. i. 159 = grandiflora.
pauciflora, Endl. in Lehm. Pl. Preiss. ii. 42 = squarrosa.
ramosa, *Lindl. Swan River App.* 56.—Austral.
Roei, Endl. in Lehm. Pl. Preiss. ii. 42 = minor.
sessiliflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 363.—Austral.
sessilis, Endl. in Lehm. Pl. Preiss. ii. 42 = squarrosa.
sessilis, *Lindl. Swan River App.* 56.—Austral.
squarrosa, Endl. in Lehm. Pl. Preiss. ii. 42 = grandiflora.
squarrosa, *Lindl. Swan River App.* 56.—Austal.
sylvestris, Endl. in Lehm. Pl. Preiss. ii. 42 = squarrosa.

LAXMANNIA, Fisch. Hort. Gorenk. ed. II. 67 (1812) = **Coluria**, R. Br. (Rosac.).
geoides, Fisch. ex Ledeb. Fl. Alt. ii. 263 = C. geoides.
potentilloides, Fisch. Cat. Hort. Gorenk. ed. II. 67 = C. potentilloides.

LAXMANNIA, Forst. Char. Gen. 94. t. 47 (1776) = **Petrobium**, R. Br. (Compos.).
arborea, Forst. l. c. = P. arboreum.

LAXMANNIA, S. G. Gmel. ex Trin. in Mém. Acad. Pétersb. vi. (1818) 492. t. 11 = **Crucianella**, Linn. (Rubiac.).
fasciculata, Trin. in Mém. Acad. Pétersb. vi. (1818) 492 = C. stylosa.

LAXMANNIA, Schreb. Gen. ii. 800 (1791) = **Acronychia**, Forst. (Rutac.).
Ankenda, Raeusch. Nom. ed. III. 99 = A. laurifolia.

LAXOPETALUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 163 (1873) = **Eremanthus**, Less. (Compos.).
albidum, Pohl, l. c. = E. goyazensis.

LAYA, Endl. Gen. 1309, in syn. (1841) = seq.

LAYIA, Hook. & Arn. Bot. Beech. Voy. 148 in nota, 357 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 395.
CALLIACHYRIS, Torr. & Gray, in Journ. Bost. Soc. Nat. Hist. v. (1844) 110.
CALLICHROA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 (1835).
CALLIGLOSSA, Hook. & Arn. Bot. Beech. Voy. 356 (1840).
ERIOPAPPUS, Arn. in Lindl. Introd. Nat. Syst. ed. II. 443 (1836).
MADAROGLOSSA, DC. Prod. v. 694 (1836).
OXYURA, DC. l. c. 693 (1836).
TOLLATIA, Endl. Gen. 428 (1838).
UROSTYLIS, Meissn. Gen. 207 (1839).
Calliglossa, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 103.—Calif.
carnosa, *Torr. & Gray, Fl. N. Am.* ii. 394.—Calif.
chrysanthemoides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 360.—Calif.
Douglasii, *Hook. & Arn. Bot. Beech. Voy.* 358.—Am. bor. occ.
elegans, *Torr. & Gray, Fl. N. Am.* ii. 394.—Calif.
Fremonti, *A. Gray, in Mem. Am. Ac.* N. S. iv. (1849) 103.—Calif.
gaillardioides, *Hook. & Arn. Bot. Beech. Voy.* 148.—Calif.
glandulosa, *Hook. & Arn. l. c. Suppl.* 358.—Am. bor. occ.
graveolens, Greene, in Bull. Calif. Acad. i. n. III. (1885) 92 = heterotricha.
heterotricha, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 358.—Calif.
hieracioides, *Hook. & Arn. l. c.*—Calif.
Jonesii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 18.—Calif.
neo-mexicana, A. Gray, Pl. Wright. ii. 98 = glandulosa.
pentachaeta, *A. Gray, in Pacif. Rail. Rep.* iv. 108. t. 16.—Calif.
platyglossa, *A. Gray, in Mem. Am. Ac.* N. S. iv. (1849) 103.—Calif.

LAYIA, Hook. & Arn. Bot. Beech. Voy. 183. t. 38 (1833) = **Ormosia**, Jacks. (Legumin.).
emarginata, Hook. & Arn. l. c. = O. emarginata.

LAZAROLUS, Medic. Phil. Bot. i. 15 (1789) = **Pyrus**, Tourn. (Rosac.).
Aria, Borckh. in Roem. Arch. i. III. 88 = P. Aria.
Chamaemespilus, Borckh. l. c. = P. Chamaemespilus.
intermedia, Borckh. l. c. = P. intermedia.
oxyacanthoides, Borckh. l. c. = Crataegus Azarolus.
pinnatifida, Borckh. l. c. = P. Aucuparia.
Pollveria, Medic. Geschich. 81 = P. Pollveria.
Pollvilleriana, Borckh. in Roem. Arch. i. III. 88 = P. Pollveria.
torminalis, Borckh. l. c. = P. torminalis.

LEA, Stokes, Bot. Mat. Med. i. 455 (1812) = **Leea**, Royen (Ampelid.).

LEACHIA, Cass. in Dict. Sc. Nat. xxv. 388 (1822) = **Coreopsis**, Linn. (Compos.).
crassifolia, Cass. l. c. = C. lanceolata.
lanceolata, Cass. l. c. = C. lanceolata.
trifoliata, Cass. l. c. 389 = C. lanceolata.

LEAEBA, Forsk. Fl. Aegypt. Arab. 172 (1775) = **Cocculus**, DC. (Menisperm.).
dubia, J. F. Gmel. Syst. 567 = C. Leaeba.

LEANDRA, Raddi, in Mem. Mod. xviii. Fis. (1820) 385. *MELASTOMACEAE*, Benth. & Hook. f. i. 760.
agrestis, *Raddi, l. c.* xx. (1829) 157.—Bras.
amplexicaulis, *DC. Prod.* iii. 153.—Bras.
angustifolia, DC. l. c. 154 = Ossaea angustifolia.

LEANDRA:—
asperifolia, Cham. in Linnaea, x. (1835–36) 33 = melastomoides.
auricoma, Spring, ex Mart. in Flora, xx. (1837) Beibl. II. 74 = Oxymeris aurea.
bullosa, *Raddi, in Mem. Mod.* xx. (1829) 160.—Bras.
capillaris, *Raddi, l. c.* 153.—Bras.
ciliata, *Cham. in Linnaea*, x. (1835–36) 36.—Bras.
corcovadensis, *Raddi, in Mem. Mod.* xx. (1829) 141.—Bras.
dubia, DC. Prod. iii. 154 = Oxymeris dubia.
estrellensis, *Raddi, in Mem. Mod.* xx. (1829) 148.—Bras.
fimbriata, *Raddi, l. c.* 158.—Bras.
hirsutissima, *Raddi, l. c.* 152.—Bras.
hirta, Raddi, l. c. xviii. Fis. (1820) 387 = Oxymeris hirta.
hispida, Schrank & Mart. ex DC. Prod. iii. 153 = amplexicaulis.
involucrata, *DC. l. c.* 154.—Bras.
involucrata, Raddi, in Mem. Mod. xx. (1829) 145 = melastomoides.
melastomoides, *Raddi, l. c.* xviii. *Fis.* (1820) 386.—Bras.
paulina, *DC. Prod.* iii. 153.—Bras.
punicea, *Raddi, in Mem. Mod.* xx. (1829) 156.—Bras.
racemifera, DC. Prod iii. 155 = Miconia racemifera.
Raddii, Mart. ex DC. l. c. 154 = sericea?
rubella, *Raddi, in Mem. Mod.* xx. (1829) 147.—Bras.
salicifolia, *Raddi, l. c.* 142.—Bras.
scabra, *DC. Prod.* iii. 154.—Bras.
sericea, *DC. l. c.*—Bras.
staminea, *Raddi, in Mem. Mod.* xx. (1829) 154.—Bras.
sylvestris, DC. Prod. iii. 154 = Oxymeris sylvestris.
umbellata, *DC. l. c.* 153.—Bras.
variabilis, *Raddi, in Mem. Mod.* xx. (1829) 150.—Bras.
villosa, DC. Prod. iii. 154 = scabra.

LEANTRIA, Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 45 = **Myrtus**, Tourn.
myrtoides, Soland. ex Forst. f. l. c. nomen (Quid?).—Hab.?

LEAROSA, Reichb. Nom. 69 (1841) = **Doryphora**, Endl. (Monimiac.).

LEAVENWORTHIA, Torr. in Ann. Lyc. N. York, iii. (1837) 87. t. 5. *CRUCIFERAE*, Benth. & Hook. f. i. 70.
aurea, Torr. l. c. 88 = Michauxii.
Michauxii, *Torr. l. c.* 89.—Am. bor. occ.
stylosa, *A. Gray, ex Chapm. Fl. S. U. St. Suppl.* 605.—Am. bor.
torulosa, *A. Gray, ex Chapm. l. c.*—Am. bor.

LEBECKIA, Thunb. Nov. Gen. 139 (1800); Prod. Pl. Cap. 122 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 477.
ACANTHOBOTRYA, Eckl. & Zeyh. Enum. 192 (1836).
CALOBOTA, Eckl. & Zeyh. l. c. 191 (1836).
SARCOPHYLLUM, E. Mey. Comm. Pl. Afr. Austr. 32 (1835).
STIZA, E. Mey. l. c. 31 (1835).
ambigua, *E. Mey. l. c.* 34.—Afr. austr.
ambigua, E. Mey. l. c. quoad litt. C = sepiaria.
angustifolia, E. Mey. in Linnaea, viii. (1833) 155 = multiflora.
aphylla, Thunb. Nov. Gen. 140 = Indigofera filifolia?
armata, E. Mey. Comm. Pl. Afr. Austr. 35 = spinescens.
armata, Thunb. Nov. Gen. 141 = Wiborgia spinescens.
armata, Willd. ex Walp. in Linnaea, xiii. (1839) 478 = microphylla.
Boureana, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 362.—Afr. austr.
Burchellii, Spreng. Syst. iv. Cur. Post. 273 = Vigna Burchellii.
Candolleana, Walp. in Linnaea, xiii. (1839) 477 = contaminata.
canescens, E. Mey. Comm. Pl. Afr. Austr. 34 = multiflora.
cinerea, *E. Mey. l. c.* 35.—Afr. austr.
contaminata, *DC. Prod.* ii. 136.—Afr. austr.
contaminata, Eckl. & Zeyh. ex Presl, Bot. Bemerk. 126 = ambigua.
contaminata, Thunb. Nov. Gen. 140 = Aspalathus corymbosa.
cytisoides, *Thunb. l. c.* 143.—Afr. austr.
decipiens, E. Mey. Comm. Pl. Afr. Austr. 35 = sericea.
decutiens, E. Mey. l. c. 34 = multiflora.

LEBECKIA:—
densa, *Thunb. Nov. Gen.* 142.—Afr. austr.
disticha, *Steud. Nom.* ed. II. ii. 19.—Afr. austr.
epiaria, Thunb. Prod. Pl. Cap. 122, err. typ. = sepiaria.
fasciculata, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 361 = humilis.
flexuosa, E. Mey. in Linnaea, viii. (1833) 34 = sericea.
gracilis, Eckl. & Zeyh. Enum. 192 = sepiaria.
grandiflora, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 357.—Afr. austr.
humilis, *Thunb. Nov. Gen.* 142.—Afr. austr.
leptophylla, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 359.—Afr. austr.
linearifolia, *E. Mey. Comm. Pl. Afr. Austr.* 33.—Afr. austr.
linearis, *DC. Prod.* ii. 137.—Afr. austr.
macrantha, *Harv. & Sond. Fl. Cap.* ii. 83.—Afr. austr.
marginata, *E. Mey. Comm. Pl. Afr. Austr.* 35.—Afr. austr.
Meyeriana, *Eckl. & Zeyh. Enum.* 192.—Afr. austr.
microphylla, *E. Mey. Comm. Pl. Afr. Austr.* 155.—Afr. austr.
mucronata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 359.—Afr. austr.
multiflora, *E. Mey. Comm. Pl. Afr. Austr.* 34.—Afr. austr.
nuda, Sims, Bot. Mag. t. 2214 = Indigofera filifolia?
pauciflora, Benth. ex Harv. & Sond. Fl. Cap. ii. 85 = contaminata.
pauciflora, *Eckl. & Zeyh. Enum.* 192.—Afr. austr.
Plukenetiana, *E. Mey. Comm. Pl. Afr. Austr.* 33.—Afr. austr.
psiloloba, *Walp. in Linnaea*, xiii. (1839) 478.—Afr. austr.
pulchella, Walp. l. c. 477 = sericea.
pungens, *Thunb. Nov. Gen.* 141.—Afr. austr.
retamoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 123.—Madag.
sarcophylloides, E. Mey. in Linnaea, vii. (1832) 155 = Simsiana.
Scorpius, Thunb. Nov. Gen. 141 = Wiborgia spinescens.
sepiaria, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 357 = Simsiana.
sepiaria, *Thunb. Nov. Gen.* 140.—Afr. austr.
sericea, *Thunb. l. c.* 143.—Afr. austr.
sessilifolia, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 362.—Afr. austr.
Simsiana, *Eckl. & Zeyh. Enum.* 192.—Afr. austr.
simsioides, Steud. Nom. ed. II. ii. 19 = praec.
spinescens, *Harv. in Harv. & Sond. Fl. Cap.* ii. 88.—Afr. austr.
subnuda, *DC. Prod.* ii. 136.—Afr. austr.
subternata, Link, Enum. Hort. Berol. ii. 225 = Cyclopia latifolia.

LEBETANTHUS, Endl. Gen. Suppl. i. 1411 (1841). *EPACRIDEAE*, Benth. & Hook. f. ii. 616.
ALLODAPE, Endl. Gen. 749 (1839), nomen prius.
JACQUINOTIA, Hombr. & Jacquinot, Bot. Astrol. et Zél. Dicot. Phan. t. 22 (1846); et ex Decne. Bot. Voy. Astrol. et Zél. 32 (1853).
americanus, *Endl. Ench.* 384.—Reg. Magellan.

LEBETHANTHUS, Reichb. Nom. 123 (1841) = praec.

LEBETINA, Cass. in Dict. Sc. Nat. xxv. 394 (1822) = **Adenophyllum**, Pers. (Compos.).
cancellata, Cass. l. c. = A. Porophyllum.

LEBIDIBIA, Griseb. Cat. Pl. Cub. 78 (1866) = LIBIDIBIA, Schlecht. = **Caesalpinia**, Linn. (Legumin.).

LEBIDIERA, Baill. Etud. Gén. Euph. Atlas, 50. t. 27 (1858) = **Cleistanthus**, Hook. f. (Euphorb.).
Cunninghamii, Muell. Arg. in Linnaea, xxxii. (1863) 80 = C. Cunninghamii.
ferruginea, Baill. Etud. Gén. Euph. Atlas, 50 = C. ferrugineus.
laurina, Muell. Arg. in Linnaea, xxxii. (1863) 80 = C. stipitatus.
malabarica, Muell. Arg. l. c. 81 = C. malabaricus.
pallida, Muell. Arg. l. c. 80 = C. pallidus.
patula, Muell. Arg. l. c. 81 = C. patulus.
stipularis, Muell. Arg. l. c. = C. stipularis.
Tulasneana, Muell. Arg. l. c. = Bridelia Tulasneana.

LEBIDIEROPSIS, Muell. Arg. in Linnaea, xxxii. (1863) 79 = **Cleistanthus**, Hook. f. (Euphorb.).
collina, Muell. Arg. l. c. 80 = C. collinus.
orbicularis, Muell. Arg. in DC. Prod. xv. II. 509 = C. collinus.

LEBRETONIA, Schranck, Pl. Rar. Hort. Monac. t. 90 (1819) = **Pavonia**, Cav. (Malvac.).
acuminata, A. Rich. Tent. Fl. Abyss. i. 54 = P. Kraussiana.
cernua, Span. in Linnaea, xv. (1841) 168 = P. cernua.
coccinea, Schrank, Pl. Rar. Hort. Monac. t. 90 = P. Schrankii.
cordata, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 190 = P. glechomifolia.
crenata, Hochst. ex Walp. Rep. v. 89; iv. Index, 506 = P. Kraussiana.
flava, Wall. Cat. n. 1883 = P. glechomifolia.
glechomaefolia, A. Rich. Tent. Fl. Abyss. i. 54 = P. glechomifolia.
latifolia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 98 = P. latifolia.
procumbens, Wight & Arn. Prod. 47 = P. glechomifolia.
semiserrata, DC. Prod. i. 446 = P. semiserrata.

LEBRETONNIA, Brongn. Enum. Gen. 77 (1843) = praec.

LECANANTHUS, Jack, in Malay. Misc. ii. (1822) n. VII. 83. *RUBIACEAE*, Benth. & Hook. f. ii. 73.
erubescens, *Jack, l. c.*—Malaya.

LECANIODISCUS, Planch. ex Benth. in Hook. Niger Fl. 250 (1849). *SAPINDACEAE*, Benth. & Hook. f. i. 402.
cupanioides, *Planch. l. c.* 251.—Afr. trop.
fraxinifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 429.—Afr. trop.

LECANOCARPUS, Nees, in Nees & Sinning, Amoen. Bot. Bonn. ii. (1824) 4. t. 2 = **Acroglochin**, Schrad. (Chenop.).
aristata, Zucc. in Mart. Hort. Monac. (1829) 56 = A. chenopodioides?
cauliflorus, Nees, in Nees & Sinning, Amoen. Bot. Bonn. ii. (1824) 5 = A. chenopodioides.
nepalensis, Nees, l. c. t. 2 = A. chenopodioides.

LECANOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 193. t. 12 (1856) = **Maoutia**, Wedd. (Urtic.).
borbonica, Blume, l. c. 194 = Pouzolzia viminea.
diversifolia, Blume, l. c. t. 11 = M. diversifolia.
heterophylla, Blume, l. c. 194 = M. diversifolia.
rubricaulis, Blume, l. c. = Pouzolzia rubricaulis.
scaberrima, Blume, l. c. t. 12.—Hab.?

LECANORCHIS, Blume, Mus. Bot. Lugd. Bat. ii. 188 (1856). *ORCHIDEAE*, Benth. & Hook. f. iii. 606.
japonica, *Blume, l. c.*—Japon.
javanica, *Blume, l. c.*—Java.

LECANTHUS, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 187. *URTICACEAE*, Benth. & Hook. f. iii. 385.
major, Wedd. l. c. = Wightii.
pedunculatis, Wedd. in DC. Prod. xvi. I. 164 = Wightii.
Solandri, *Seem. Fl. Vit.* 236.—Ins. Fiji.
Wallichii, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 187 = Wightii.
Wightii, *Wedd. l. c.*—Ind. or.

LECHEA, Kalm, ex Linn. Diss. Chen. 13. '15' (1751); Amoen. Acad. iii. 10 (1756). *CISTINEAE*, Benth. & Hook. f. i. 114.
GAURA, Lam. Illustr. t. 281. f. 3 (1793?).
HORANTHES, Rafin. New Fl. Am. iii. 30 (1836).
LECHIDIUM, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 286.
brevifolia, Rafin. New Fl. Am. i. 96 = thymifolia?
cinerea, Rafin. l. c. 91 = Torreyi.
corymbosa, *Rafin. l. c.* 94.—Am. bor.
cubensis, *Leggett, in Bull. Torrey Bot. Club*, vi. (1875) 252.—Cuba.
divaricata, Shuttlew. ex Leggett, l. c. 251 = major?
Drummondii, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 284 = major?

LECHEA:—
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 154. t. 88.—Am. bor.
floridana, Rafin. New Fl. Am. i. 95 (Quid?).—Am. bor.
furfuracea, Rafin. l. c. 92 (Quid?).—Am. bor.
glomerata, Rafin. l. c. 94 (Quid?).—Am. bor.
heterophylla, Rafin. l. c. 93 (Quid?).—Am. bor.
juncifolia, Walt. Fl. Carol. 83 = tenuifolia?
laxiflora, Rafin. New Fl. Am. i. 92 = minor?
major, Linn. Sp. Pl. 90 = Helianthemum canadense.
major, *Michx. Fl. Bor. Am.* i. 76.—Am. bor.
mexicana, Hort. Berol. ex Spach, in Hook. Comp. Bot. Mag. ii. (1836) 289 = Helianthemum glomeratum.
minor, Hook. Journ. Bot. i. (1834) 192 = Novae-Cesareae?
minor, Linn. Sp. Pl. 90; Amoen. Acad. iii. 10 = major.
minor, *Walt. Fl. Carol.* 83.—Am. bor.
mucronata, Rafin. in Am. Monthly Mag. (1817) 359, nomen; New Fl. Am. i. 93 = major.
Novae-Cesareae, *C. F. Austin, ex A. Gray, Man. Bot. N. U. St.* 81.—Am. bor.
patula, *Leggett, in Bull. Torrey Bot. Club*, vi. (1875) 251, nomen.—Am. bor.
pauciflora, Rafin. New Fl. Am. i. 96 = thymifolia?
pulchella, Rafin. l. c. 91 = minor?
racemulosa, Hook. Journ. Bot. i. (1834) 193 = Torreyi.
racemulosa, *Lam. Illustr.* ii. 423. t. 281. f. 3.—Am. bor.
recurvata, Rafin. New Fl. Am. i. 93 (Quid?).—Am. bor.
revoluta, Rafin. l. c. 94 (Quid?).—Am. bor.
secundiflora, Rafin. l. c. 95 = thymifolia?
sessiliflora, Rafin. l. c. 97 = patula?
Skinneri, *Benth. Bot. Voy. Sulph.* 66.—Am. bor. occ.
stellata, Rafin. New Fl. Am. i. 96 = thymifolia, Novae-Cesareae.
surculosa, Rafin. l. c. 94 = minor?
tenuifolia, *Michx. Fl. Bor. Am.* i. 77.—Am. bor.
ternifolia, Rafin. New Fl. Am. i. 91 = racemulosa?
thesioides, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 285 = racemulosa.
thymifolia, *Michx. Fl. Bor. Am.* i. 77.—Am. bor.
Torreyi, *Leggett, in S. Wats. Bibl. Ind. N. Am. Bot.* i. 81.—Am. bor.
uniflora, Rafin. New Fl. Am. i. 96 = thymifolia?
verna, Rafin. l. c. 92 = tenuifolia.
verticillata, Willd. Sp. Pl. i. 495 = Bergia ammannioides.
villosa, Ell. Sketch, i. 184 = major.
virgata, Rafin. New Fl. Am. i. 95 (Quid?).—Am. bor.

LECHEA, Lour. Fl. Cochinch. 60 (1790) = **Commelina**, Linn.
chinensis, Lour. l. c. = C. Loureiri.

LECHENAULTIA, R. Br. Prod. 581 (1810) = **Leschenaultia**, R. Br. (Gooden.).

LECHEOIDES, Endl. Gen. 905 (1839) sphalm. = seq.

LECHIDIUM, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 286 = **Lechea**, Kalm (Cistin.).
Drummondii, Spach, l. c. = Lechea Drummondii.

LECHLERA, Griseb. in Lech. Berb. Am. Austr. 53 (1857) = **Solenomelus**, Miers (Irid.).
Sisyrinchium, Griseb. l. c. = S. Lechleri.

LECHLERA, Miq. ex Steud. Syn. Pl. Gram. 101 in nota sub RELCHELA (1855). 'Xanthoxylearum. Quid?'

LECHLERA, Steud. Syn. Pl. Gram. 101, in nota (1855) = RELCHELA, Steud. = **Calamagrostis**, Roth (Gramin.).

LECHLERIA, Phil. ex eod. in Linnaea, xxviii. (1856) 654 = **Huanaca**, Cav. (Umbellif.).
palmata, Phil. l. c. = H. Cavanillesii.

LECISCIUM, Gaertn. f. Fruct. iii. 221. t. 220 (1805). An Myrtacea?
drupaceum, Gaertn. f. l. c. (Quid?).—Hab.?

LECOCARPUS, Decne. Voy. Vénus, 19. t. 14 (1864). *COMPOSITAE*, Benth. & Hook. f. ii. 348.
foliosus, *Decne. l. c.* 19.—Ins. Galap.
pinnatifidus, Decne. l. c. t. 14 = foliosus.

LECOCKIA, Meissn. Gen. Comm. 108 (1843) = seq.

LECOKIA, DC. Coll. Mém. v. 67 (1829); Prod. iv. 240 (1830). *UMBELLIFERAE*, Benth & Hook. f. i. 882.
cretica, *DC. ll. cc.*—Oriens.

LECONTEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 195. t. 20, f. 1 bis. *RUBIACEAE*, Benth. & Hook. f. ii. 134.
argentea, *A. Rich. l. c.* 195.—Madag.
Bojeriana, *A. Rich. l. c.*—Madag.

LECONTIA, W. Cooper, ex Torr. Compend. Fl. 358 [bis] (1826) = **Peltandra**, Rafin. (Aroid.).
virginica, Torr. l. c. = P. alba.

LECOSTEMON, [Moç. & Sesse,] ex DC. Prod. ii. 639 (1825). *ROSACEAE*, Benth. & Hook. f. i. 609.
amazonicum, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 295.—Bras.
crassipes, *Spruce, ex Benth. l. c.*—Bras.
Gardnerianum, *Benth. l. c.*—Bras.
macrophyllum, *Spruce, ex Benth. l. c.* 296.—Bras.
terniflorum, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 639.—Mexic.

LECOSTOMUM, Steud. Nom. ed. II. ii. 20 (1841) = praec.

LECYTHIS, Loefl. It. Hisp. 189 (1758). *MYRTACEAE*, Benth & Hook. f. i. 723.
CHYTROMA, Miers, in Trans. Linn. Soc. xxx. (1874) 229.
ESCHWEILERA, Mart. in DC. Prod. iii. 293 (1825).
JUGASTRUM, Miers, in Trans. Linn. Soc. xxx. (1874) 275.
acuminata, *DC. Prod.* iii. 292.—Bras.
acuminatissima, *Berg, in Mart. Fl. Bras.* xiv. I. 494.—Bras.
albiflora, *DC. Prod.* iii. 292.—Bras.
amara, *Aubl. Pl. Gui.* ii. 716. t. 286, et t. 285. f. 1.—Guian.
Amazonii, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. I. 484.—Bras.
ampla, *Miers, in Trans. Linn. Soc.* xxx. (1874) 204.—N. Granat.
ampullaria, *Miers, l. c.* 201.—N. Granat.
angustifolia, *Endl. ex Walp. Rep.* ii. 193.—Bras.
basizone, *Berg, in Mart. Fl. Bras.* xiv. I. 617.—Guian.
biserrata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 213.—Bras.
Blanchetiana, *Berg, in Mart. Fl. Bras.* xiv. I. 495.—Bras.
bracteata, *Willd. Sp. Pl.* ii. 1174 = Couroupita guianensis.
bracteosa, *Poepp. ex Berg, in Mart. Fl Bras.* xiv. I. 495.—Bras.
brasiliensis, *Spreng, Syst.* ii. 603.—Bras.
chartacea, *Berg, in Linnaea,* xxvii. (1854) 450.—Guian.
cognata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 215.—Guian.
compressa, Vell. Fl. Flum. 222; v. t. 87 = angustifolia.
cordata, *Berg, in Linnaea,* xxvii. (1854) 457.—Bras.
coriacea, *DC. Prod.* iii. 291.—Bras.
corrugata, *Poit. in Mém. Mus. Par.* xiii. (1825) 146. f. 3.—Guian.
Couratari, Spreng. Syst. iv. Cur. Post. 208 = Couratari guianensis.
Coxiana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 217.—Bras.
crassinoda, *Miers, l. c.* 216.—Guian.
cupularis, *Miers, l. c.* 224.—Bras.
densa, *Miers, l. c.* 211.—Bras.
dubia, *H. B. & K. Nov. Gen. et Sp.* vii. 259.—N. Granat.
elegans, *Berg, in Mart. Fl. Bras.* xiv. I. 499.—Bras.
elliptica, *H. B. & K. Nov. Gen. et Sp.* vii. 259.—N. Granat.
fagifolia, *Miq. ex Berg, in Linnaea,* xxvii. (1854) 451.—Guian.
glabra, Berg, l. c. 452 = Lecythopsis glabra.
gracilipes, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 203.—Guian.
grandiflora *Aubl. Pl. Gui.* ii. 712, tt. 283, 284, 285.—Guian.

LECYTHIS:—
grandifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 494.—Bras.
Idatimon, *Aubl. Pl. Gui.* ii. 721. t. 289.—Guian.
Idatimon, DC. ex Berg, in Mart. Fl. Bras. xiv. I. 498 = ovata.
idatimonoides, *Berg, l. c.* 496.—Guian.
integrifolia, *Ruiz & Pav. Fl. Per.* 529; *ex Miers, in Trans. Linn. Soc.* xxx. (1874) 225.—Ecuador.
laevicula, *Miers, l. c.* 224.—Guian.
laevifolia, *Griseb. Fl. Brit. W. Ind.* 711, nomen; *Miers, l. c.* 256.—Ins. Trinit.
lanceolata, *Poir. Encyc.* vi. 27.—Bras.
Langsdorffii, *Berg, in Mart. Fl. Bras.* xiv. I. 615.—Bras.
limbata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 214.—Bras.
longifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 259.—Venezuela.
longipes, *Poit. in Mém. Mus. Par.* xiii. (1825) 144. t. 2.—Guian.
Luschnathii, *Berg, in Mart. Fl. Bras.* xiv. I. 499.—Bras.
lutea, *Aubl. Pl. Gui.* ii. 721.—Guian.
macrophylla, *Berg, in Mart. Fl. Bras.* xiv. I. 616.—Guian.
marawynensis, *Berg, l. c.* 489.—Guian.
Marcgraaviana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 210.—Bras.
Martiana, *Berg, in Mart. Fl. Bras.* xiv. I. 487.—Bras.
Martini, *Berg, l. c.* 616.—Guian.
Melinonis, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 202.—Guian.
micaraguensis, Steud. Nom. ed. II. ii. 20, sphalm. = *nicaraguarensis*.
micrantha, *Berg, in Linnaea*, xxvii. (1854) 454.—Guian.
minor, *Jacq. Stirp. Select. Am.* 168. t. 109.—N. Granat.
minor, Vell. Fl. Flum. 222; v. t. 85 = lanceolata.
monosperma, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. I. 485.—Bras.
multiflora, *Sm. in Rees, Cycl.* xx. n. 8.—Guian.
nana, *Berg, in Mart. Fl. Bras.* xiv. I. 617.—Bras.
nicaraguarensis, [Moç. & Sesse,] ex DC. Prod. iii. 294 = Couroupita nicaraguarensis.
obversa, *Berg, in Mart. Fl. Bras.* xiv. I. 503.—Bras.
odora, *Poepp. ex Berg, l. c.* 492.—Bras.
odoratissima, *Salzm. ex Miers, in Trans. Linn. Soc.* xxx. (1874) 255.—Bras.
Ollaria, *Linn. Syst.* ed. X. 1071.—Am. trop.
Ollaria, Saldanha, Configuração, 70 = biserrata.
Ollaria, Spruce, in Hook. Kew Journ. ii. (1850) 74 = usitata.
Ollaria, Vell. Fl. Flum. 222; v. t. 88 = Velloziana.
ovalifolia, *DC. Prod.* iii. 292.—Bras.
ovata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 378.—Bras.
pachysepala, *Spruce, ex Berg, in Mart. Fl. Bras.* xiv. I. 494.—Bras.
paniculata, *Berg, l. c.* 501.—Bras.
parviflora, *Aubl. Pl. Gui.* ii. 717.—Guiana.
parviflora, *Sagot, ex Miers, in Trans. Linn. Soc.* xxx. (1874) 262.—Guian.
parvifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 496.—Bras.
pedicellaris, *DC. Prod.* iii. 292.—Bras.
pedicellata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
persistens, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 201.—Guian.
pilaris, *Miers, in Trans. Linn. Soc.* xxx. (1874) 223.—Bras.
pilosa, *Poepp. ex Berg, in Mart. Fl. Bras.* xiv. I. 500.—Bras.
Pisonis, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 377.—Bras.
platycarpa, *Poit. in Mém. Mus. Par.* xiii. (1825) 146 (platicarpa).—Bras.
platyzone, *Berg, in Mart. Fl. Bras.* xiv. I. 485.—Bras.
Pohlii, *Berg, l. c.* 484.—Bras.
Poiteaui, *Berg, l. c.* 615.—Guian.
pyramidata, Vell. Fl. Flum. 222; v. t. 86 = Lecythopsis rufescens.
racemiflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 200.—Guian.
retusa, *Spruce, ex Berg, in Mart. Fl. Bras.* xiv. I. 487.—Bras.
Riedeliana, *Berg, l. c.* 618.—Bras.

LECYTHIS:—
rorida, *Berg, in Mart. Fl. Bras.* xiv. I. 488.—Bras.
rosea, *Spruce, ex Berg, l. c.*—Bras.
rubicunda, *Miers, in Trans. Linn. Soc.* xxx. (1874) 226.—Guian.
rufescens, Berg, in Linnaea, xxvii. (1854) 451 = Lecythopsis rufescens.
salebrosa, *Berg, in Mart. Fl. Bras.* xiv. I. 488.—Guian.
Schomburgkii, *Berg, in Linnaea*, xxvii. (1854) 456.—Bras.
setifera, *Miers, in Trans. Linn. Soc.* xxx. (1874) 227.—Bras.
Siberiana, *Berg, in Mart. Fl. Bras.* xiv. I. 501.—Bras.
sphaeroides, *Miers, in Trans. Linn. Soc.* xxx. (1874) 212.—Bras.
Spruceana, *Berg, in Mart. Fl. Bras.* xiv. I. 487.—Bras.
subbiflora, *Ruiz & Pav. Fl. Per.* iv. t. 461.—Peruv.
subglandulosa, *Steud. ex Berg, in Linnaea*, xxvii. (1854) 459.—Guian.
tenax, *Moritz, ex Berg, l. c.* 456.—Venezuela.
tenuifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 502.—Bras.
tuberculata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 218.—Bras.
tumefacta, *Miers, l. c.* 216.—Guian.
turbinata, *Berg, in Mart. Fl. Bras.* xiv. I. 493.—Bras.
urnigera, *Mart. l. c.* 483.—Bras.
usitata, *Miers, in Trans. Linn. Soc.* xxx. (1874) 208.—Bras.
validissima, *Miers, l. c.* 205.—Guian.
variolata, *Miers, l. c.* 220.—Bras.
vasiformis, *Miers, l. c.* 204.—Bras.
Velloziana, *Miers, l. c.* 203.—Bras.
venusta, *Miers, l. c.* 214.—Guian.
verrucosa, Hoffm. ex Berg, in Mart. Fl. Bras. xiv. I. 498 = ovata.
verruculosa, *Berg, l. c.* 502.—Bras.
Wullschlaegeliana, *Berg, l. c.* 493.—Guian.
Zabucajo, *Aubl. Pl. Gui.* ii. 719. tt. 288, 284.—Guian.
Zabucayo, Hook. Kew Journ. ii. (1850) 74 = usitata.

LECYTHOPSIS, Schrank, in Denkschr. Akad. Muench. vii. (1821) 241. *MYRTACEAE*, Benth. & Hook. f. i. 723.
COURATARI, Cambess. in St. Hil. Fl. Pav. Mer. ii. 379 (1829).
glabra, *Berg, in Mart. Fl. Bras.* xiv. I. 505.—Bras.
rufescens, *Berg, l. c.* 504.—Bras.

LEDEBOURIA, Reichb. Consp. 143 (1828) = LEDEBURIA, Link = **Pimpinella**, Linn. (Umbellif.).

LEDEBOURIA, Roth, Nov. Pl. Sp. 194 (1821) = **Scilla**, Linn. (Liliac.).
hyacinthina, Roth, l. c. 195 = S. indica.
hyacinthoides, Steud. Nom. ed. II. ii. 20 = praec.
maculata, Dalz. in Hook. Kew Journ. ii. (1850) 143 = S. indica.

LEDEBURIA, Link, Enum. Hort. Berol. i. 286 (1821) = **Pimpinella**, Linn. (Umbellif.).
pimpinelloides, Link, l. c. = P. peregrina.

LEDELIA, Rafin. Sylva Tellur. 154 (1838) = **Pomaderris**, Labill. (Rhamnac.).
betulina, Rafin. l. c. = P. betulina.

LEDENBERGIA, Klotzsch, ex Moq. in DC. Prod. xiii. II. 14 (1849). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 81.
rosea-aenea, *Lem. Illustr. Hortic.* xvi. (1869) t. 591.—Bras.
seguierioides, *Klotzsch, ex Moq. in DC. Prod.* xiii. II. 14.—Am. austr.

LEDGERIA, F. Muell. Fragm. i. 238 (1859) = **Galeola**, Lour. (Orchid.).
aphylla, F. Muell. l. c. 239 = G. cassythoides.
foliata, F. Muell. l. c. ii. 167 = G. foliata.

LEDOCARPON, Desf. in Mém. Mus. Par. iv. (1818) 250 = **Balbisia**, Cav. (Geraniac.).
argenteum, Presl, Symb. Bot. v. 67 = B. verticillata.
chilense, Steud. Nom. ed. II. ii. 20 = seq.
chiloënse, Desf. in Mém. Mus. Par. iv. (1818) 551. t. 13 = B. verticillata.
cistiflorum, Meyen, Reise, i. 470 = B. peduncularis.
Meyenianum, Steud. Nom. ed. II. ii. 20 = B. Meyeniana.
pedunculare, Lindl. Bot. Reg. t. 1392 = B. verticillata.
Reynoldsii, Hook. Ic. Pl. t. 14 = Wendtia Reynoldsii.

LEDOCARPUM, DC. Prod. i. 702 (1824) = praec.

LEDONIA, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 369 = **Cistus**, Tourn.
aprica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 18 = C. salvifolius.
arrigens, Jord. & Fourr. l. c. 17 = C. salvifolius.
elegans, Jord. & Fourr. l. c. 18 = C. salvifolius.
fruticans, Jord. & Fourr. l. c. = C. salvifolius.
heterophylla, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 369 = C. longifolius, *Lam.*
hirsuta, Spach, l. c. = C. hirsutus.
humilis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 17 = C. salvifolius.
microphylla, Jord. & Fourr. l. c. = C. salvifolius.
peduncularis, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 369 = C. salvifolius.
platyphylla, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 18 = C. salvifolius.
populifolia, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 369 = C. populifolius.
rhodanensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 16 = C. salvifolius.
salviaefolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 339 = C. salvifolius.
velutina, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 18 = C. salvifolius.

LEDOTHAMNUS, Meissn. in. Mart. Fl. Bras. vii. 172 (1863). *ERICACEAE*, Benth. & Hook. f. ii. 597.
guyanensis, *Meissn. l. c.*—Am. trop.

LEDUM, Reichb. in Moessl. Handb. ed. II. i. p. 1 (1827), sphalm. = **Sedum**, Tourn. (Crassul.).

LEDUM, Rupp. ex Linn. Syst. ed. I (1735). *ERICACEAE*, Benth. & Hook. f. ii. 599.
DULIA, Adans. Fam. ii. 165 (1763).
buxifolium, Berg. in Act. Acad. Petrop. i. I. (1777) 213 = Leiophyllum buxifolium.
californicum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 14.—Calif.
canadense, Lodd. Bot. Cab. t. 1049 = latifolium.
decumbens, Lodd. ex Steud. Nom. ed. II. ii. 20 = palustre.
dilatatum, Rupr. in Beitr. Pfl. Russ. Reich. iv. 57 = palustre.
glandulosum, *Nutt. in Trans. Am. Phil. Soc.* viii. (1843) 270.—Am. bor. occ.
graveolens, Gilib. Fl. Lituan. ii. 199 = palustre.
groenlandicum, Retz. Fl. Scand. Prod. i. 77 = latifolium.
latifolium, *Jacq. Coll.* ii. 308.—Am. bor.
palustre, *Linn. Sp. Pl.* 391.—Reg. bor. et arct.
palustre, Michx. Fl. Bor. Am. i. 259 = latifolium.
serpyllifolium, L'Hérit. ex DC. Prod. vii. 730 = Leiophyllum buxifolium.
thymifolium, Lam. Encyc. iii. 459 = Leiophyllum buxifolium.
tomentosum, Stokes, Bot. Mat. Med. ii. 502 = palustre.

LEEA, Royen, ex Linn. Mant. i. 17, 124 (1767). *AMPELIDEAE*, Benth. & Hook. f. i. 388.
AQUILICIA, Linn. Mant. ii. 146, 211 (1771).
OTILLIS, Gaertn. Fruct. i. 275. t. 57 (1788).
LEEANIA, Rafin. Princ. Somiol. 30 (1814).
SANSOVINIA, Scop. Introd. 228 (1777).
aculeata, *Blume, Bijdr.* 197; *Blanco, Fl. Filip.* ed. II. 127.—Malaya; Ins. Philipp.
acuminata, Herb. Kew. ex C. B. Clarke, in Journ. Bot. xix. (1881) 141 = umbraculifera.
acuminata, *Wall. Cat.* n. 6830.—Reg. Himal.; Malaya.
aequata, *Linn. Mant.* i. 124.—Bengal; Burma; Malaya.
aequata, Roxb. ex Wight & Arn. Prod. 132; Wall. Cat. n. 6824 D = sambucina.
alata, *Edgew. in Trans. Linn. Soc.* xx. (1846) 36.—Reg. Himal.
amabilis, *Hort. Veitch. ex Mast. in Gard. Chron.* (1882) I. 492.—Borneo.
ancolana, *Miq. Fl. Ind. Bat.* i. II. 611.—Sumatra.
angulata, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 97.—Java.
angustifolia, M. Laws. in Hook. f. Fl. Brit. Ind. i. 665 = parallela.
arborea, Boj. Hort. Maurit. 61 = guineensis?
arborea, Sieber, ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = sambucina.
aspera, *Edgew. in Trans. Linn. Soc.* xx. (1846) 36.—Ind. or.

LEEA:—
aspera, Kurz, in Journ. As. Soc. xliv. (1873) II. 178 = herbacea.
aspera, Wall. ex Roxb. Fl. Ind. ed. Carey, ii. 468 = robusta.
aurantiaca, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 577.—Java.
biserrata, *Miq. Fl. Ind. Bat. Suppl.* 518.—Ins. Banca.
bracteata, *C. B. Clarke, in Journ. Bot.* xix. (1881) 164.—Reg. Himal.
Brunoniana, *C. B. Clarke, l. c.* 166.—Austral.
callosa, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 96 = sundaica.
celebica, *C. B. Clarke, in Journ. Bot.* xix. (1881) 166.—Ins. Celebes.
cinerea, M. Laws. in Hook. f. Fl. Brit. Ind. i. 665 = latifolia.
coccinea, Boj. Hort. Maurit. 61 = sambucina.
coccinea, *Planch. in Hort. Donat.* 6; *et in Hook. Bot. Mag.* t. 5299.—Burma.
compactiflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 65.—Burma.
cordata, Wall. Cat. n. 6819 (Vitis sp.).—Ind. or.
coriacea, M. Laws. in Hook. f. Fl. Brit. Ind. i. 665 = latifolia.
crispa, M. Laws. l. c. partim = herbacea.
crispa, *Linn. Mant.* i. 124.—Ind. or.; Burma.
Cumingii, *C. B. Clarke, in Journ. Bot.* xix. (1881) 166.—Ins. Philipp.
diffusa, M. Laws. in Hook. f. Fl. Brit. Ind. i. 667 = robusta.
divaricata, *Teijsm. & Binn. Cat. Hort. Bog.* 388, nomen.—Hab. ?
fuliginosa, Miq. Fl. Ind. Bat. Suppl. 518 = sundaica.
gigantea, *Griff. Notul.* iv. 697.—Burma; Malaya.
gigantea, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 65 = tubercolosemen.
grandifolia, *Kurz, in Journ. Bot.* xiii. (1875) 325.—Ins. Nicobar.
guineense, *G. Don, Gen. Syst.* i. 712.—Afr. trop.
herbacea, *Buch.-Ham. in Wall. Cat.* n. 6829, 6824, partim.—Reg. Himal.
heterophylla, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = sambucina.
hirsuta, Blume, Bijdr. 197 = aequata.
hirta, *Banks, ex Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 655.—Reg. Himal.; Burma; Malaya.
hirta, Hornem. Hort. Hafn. i. 231 = aequata.
horrida, *Teijsm. & Binn. Cat. Hort. Bog.* 169 = angulata.
humilis, *Hassk. Cat. Hort. Bog. Alt.* 168.—Java.
integrifolia, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 659.—Ind. or.
javanica, *Blume, Bijdr.* 197.—Malaya.
Kurzii, *C. B. Clarke, in Journ. Bot.* xix. (1881) 165.—Ins. Andam.
laeta, *Wall. Cat.* n. 6831.—Burma; Java.
laevis, Heyne, ex Wall. Cat. n. 1258 = Heynea trijuga.
latifolia, *Wall. l. c.* n. 6821.—Ind. or.
linearifolia, *C. B. Clarke, in Journ. Bot.* xix. (1881) 165.—Cambodia.
lucida, Linden, ex Hook. Bot. Mag. sub t. 5299 = coccinea
macrophylla, DC. Prod. i. 635 = latifolia.
macrophylla, M. Laws. in Hook. f. Fl. Brit. Ind. i. 664, partim = latifolia.
macrophylla, *Roxb. ex Hornem. Hort. Hafn.* i. 251; *Wight, Ic.* t. 1154.—Ind. or.; Burma.
maculata, *Desf. Cat. Hort. Par.* ed. III. 406.—Hab. ?
manillensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 314.—Ins. Philipp.
Mastersii, *C. B. Clarke, in Journ. Bot.* xix. (1881) 142.—Reg. Himal.
moluccana, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = aculeata.
odontophylla, Wall. Cat. n. 6820 = Vitis lanata ?
Ottilis, DC. Prod. i. 636 = sambucina.
palembanica, Miq. Fl. Ind. Bat. Suppl. 519 = sundaica.
parallela, *Wall. Cat.* n. 6828.—Bengal; Burma.
pinnata, Andr. Bot. Rep. v. t. 355 = crispa.
polyphylla, *Miq. Fl. Ind. Bat.* i. II. 610.—Sumatra.
pubescens, *Zipp. ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 97.—Ins. Timor.
pumila, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 302.—Burma.
robusta, Blume, Bijdr. 198 = sundaica.
robusta, M. Laws. in Hook. f. Fl. Brit. Ind. i. 667 = bracteata.
robusta, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 655.—Ind. or.; Burma.

LEEA:—
robusta, Wight, ex C. B. Clarke, in Journ. Bot. xix. (1881) 105 = Wightii.
rubra, *Blume, Bijdr.* 197.—Burma; Malaya.
rubra, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 66 = praec. ?
rubra, Royle, Illustr. Bot. Himal. 145 = alata.
sambucifolia, Salisb. Prod. 144 = sambucina.
sambucina, Benth. Fl. Austral. i. 451 = Brunoniana.
sambucina, Blanco, Fl. Filip. ed. II. 126 = rubra ?
sambucina, M. Laws. in Hook. f. Fl. Brit. Ind. i. 166 = aequata.
sambucina, Schum. & Thonn. Beskr. Guin. Pl. 134 = guineense.
sambucina, *Willd. Sp. Pl.* i. 1177.—Geront. trop.
sanguinea, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 66 = rubra.
sanguinea, M'Lell. ex C. B. Clarke, in Journ. Bot. xix. (1881) 102 = pumila.
sanguinea, Wall. Cat. n. 6824 M = laeta.
sanguinea, *Wall. ex Boj. Hort. Maurit.* 61.—Ind. or.
scabra, Roxb. ex Roem. & Schult. Syst. iv. 814, in obs. = hirta.
serrulata, Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = acuminata.
setuligera, *C. B. Clarke, in Journ. Bot.* xix. (1881) 105.—Ind. or.
simplex, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = sambucina.
simplicifolia, Griff. Notul. iv. 697 = macrophylla.
simplicifolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 101 = Zippeliana.
simplicifolia, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 578.—Sumatra.
speciosa, Sieber, ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 99 = sambucina.
spinosa, Spreng. Syst. i. 670 = Aralia spinosa.
Staphylea, Roxb. Hort. Beng. 18; Fl. Ind. i. 658 = sambucina.
Staphylea, Wall. Cat. n. 6824 = acuminata, alata.
Staphylea, Wight, Illustr. t. 58 = Wightii.
striata, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 96 = sundaica.
sumatrana, *Miq. Fl. Ind. Bat.* i. II. 610.—Sumatra.
sundaica, *Miq. l. c.*—Java.
tinctoria, *Lindl. ex Baker, in Oliver, Fl. Trop. Afr.* i. 416.—Afr. trop.
trifoliata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 666.—Reg. Himal.
tuberculosemen, *C. B. Clarke, in Journ. Bot.* xix. (1881) 141.—Burma.
umbraculifera, *C. B. Clarke, l. c.*—Reg. Himal.
viridiflora, Planch. Hort. Donat. 6 = sambucina.
Wrightii, *C. B. Clarke, in Journ. Bot.* xix. (1881) 105.—Malabar.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 101.—N. Guin.

LEEANIA, Rafin. Princ. Somiol. 30 (1814) = praec.

LEERIA, Steud. Nom. ed. I. 469 (1821) = LERIA, DC. = **Chaptalia**, Vent. (Compos.).

LEERSIA, Soland. ex Sw. Prod. Veg. Ind. Occ. 21 (1788). *GRAMINEAE*, Benth. & Hook. f. iii. 1117.
ASPRELLA, Schreb. Gen. 45 (1789).
BLEPHAROCHLOA, Endl. Gen. 1352 (1841).
EHRHARTIA, [Web. in] Wigg. Prim. Fl. Holsat. 63 (1780).
ENDODIA, Rafin. Neogenyt. 4 (1825).
HOMALOCENCHRUS, Mieg. ex Hall. Stirp. Helv. ii. 201 (1768), nomen prius.
LAERTIA, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 354.
PSEUDORYZA, Griff. Ic. Pl. Asiat. t. 144 (1847).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 356 = hexandra.
aegyptiaca, Figar. & De Not. in Mem. Acc. Torin. Ser. II. xiv. (1853) 317 = hexandra.
aristata, Roxb. Hort. Beng. 26; Fl. Ind. ii. 207 = Hygroryza aristata.
aspera, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 168 = monandra.
asperrima, Willd. ex Trin. l. c. 171 = oryzoides.
australis, R. Br. Prod. 210 = hexandra.
brasiliensis, *Spreng. Nov. Prov.* 47.—Bras.
capensis, *C. Muell. in Bot. Zeit.* xiv. (1856) 345.—Afr. austr.

LEERSIA:—
ciliata, Roxb. Hort. Beng. 26; Fl. Ind. ii. 207 = hexandra.
contracta, Nees, Agrost. Bras. 516 = hexandra.
debilis, *Bal. & Poitrass. in Bull. Soc. Hist. Nat. Toul.* xii. (1878) 220.—Paraguay.
digitata, *Poir. Encyc. Suppl.* iii. 329.—Am. austr.
disticha, *Benth. in Hook. Niger Fl.* 556.—Afr. trop.
distichophylla, *Bal. & Poitrass. in Bull. Soc. Hist. Nat. Toul.* xii. (1878) 221.—Paraguay.
elongata, Willd. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 172 = hexandra.
ferox, *Figar. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1853) 319.—Nubia.
glaberrima, *Trin. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 173.—Texas.
Gouini, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 514.—Mexic.
gracilis, Willd. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 173 = hexandra.
Griffithiana, *C. Muell. in Bot. Zeit.* xiv. (1856) 345.—Ind. or.
hexandra, *Sw. Prod. Veg. Ind. Occ.* 21.—Amphig. trop.
imbricata, Poir. Encyc. Suppl. iii. 329 = virginica.
latifolia, Willd. ex Steud. Nom. ed. II. ii. 21 = Maltebrunia leersioides.
lenticularis, *Michx. Fl. Bor. Am.* i. 39.—Am. bor.
ligularis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 168.—Mexic.
luzonensis, J. & C. Presl, Rel. Haenk. i. 207 = hexandra.
mauritanica, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 174 = oryzoides.
mexicana, H. B. & K. Nov. Gen. et Sp. i. 195 = hexandra.
mexicana, Rupr. in Bull. Acad. Brux. ix. (1842) 245 = Gouini.
monandra, *Sw. Prod. Veg. Ind. Occ.* 21.—Jamaic.
oryzoides, Michx. Fl. Bor. Am. i. 37 = virginica.
oryzoides, *Sw. Prod. Veg. Ind. Occ.* 21.—Amphig.
ovata, *Poir. Encyc. Suppl.* iii. 329.—Am. bor.
parviflora, *Desv. Opusc.* 61.—Aegypt.
Triniana, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 174 = hexandra.
virginica, *Willd. Sp. Pl.* i. 325.—Am. bor.

LEEUWENHOCKIA, Steud. Nom. ed. II. ii. 21 (1841) = **Levenhookia**, R. Br. (Stylid.).

LEEUWENHOECKIA, E. Mey. ex Endl. Nov. Stirp. Dec. 37 (1839) = **Dombeya**, Cav. (Stercul.).
tiliacea, E. Mey. l. c. = D. tiliacea.

LEEUWENHOEKIA, Spreng. Anleit. ii. I. 300 (1817) = **Levenhookia**, R. Br. (Stylid.).

LEEUWENHOOKIA, Reichb. Consp. 91 (1828) = praec.

LEFEBUREA, Endl. Gen. Suppl. ii. 69 (1842) = **Lefebvrea**, A. Rich. (Umbellif.).

LEFEBURIA, Lindl. Veg. Kingd. 778 (1847) = praec.

LEFEBVREA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 260. t. 15. f. 1. *UMBELLIFERAE*, Benth. & Hook. f. i. 921.
abyssinica, *A. Rich. l. c.*—Abyss.
brachystyla, *Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 80.—Afr. trop.

LEGAZPIA, Blanco, Fl. Filip. ed. II. 338 (1845) = **Torenia**, Linn. (Scrophul.).
triptera, Blanco, l. c. 339 = T. cardiosepala.

LEGENDREA, Webb & Berth. Phyt. Canar. iii. 26 (1836–50). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 870.
mollissima, *Webb & Berth. l. c.* 27.—Ins. Canar.

LEGNEPHORA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 88 = **Pericampylus**, Miers (Menisp.).
Moorii, Miers, l. c. 90 = P. incanus.

LEGNOTIS, Sw. Prod. Veg. Ind. Occ. 84 (1788) = **Cassipourea**, Aubl. (Rhizoph.).
Cassipourea, Sw. l. c. = C. macrophylla.
elliptica, Sw. l. c. = C. elliptica.
macrophylla, Mart. ex DC. Prod. iii. 34 = C. macrophylla.

LEGOUIXIA, Heurck. & Muell. Arg. in Heurck. Obs. Bot. 145 (1871) = **Epigynum**, Wight (Apocyn.).
amabilis, Heurck. & Muell. Arg. l. c. 146 = E. Griffithianum.

LEGOUSIA, Durand, Fl. Bourg. i. 37; ii. 26 (1782) = **Specularia**, Heist. (Campanul.).
arvensis, Durand, ll. cc. = S. Speculum.
Durandi, Delarb. Fl. Auv. ed. II. 45 = S. Speculum.
ericoides, Buek, Ind. ad DC. Prod. iii. 255 = Prismatocarpus paniculatus.
hybrida, Delarb. Fl. Auv. ed. II. 47 = S. hybrida.
parviflora, S. F. Gray, Nat. Arr. Brit. Pl. ii. 410 = S. hybrida.
Speculum, Fisch. ex A. DC. Monog. Camp. 347 = S. Speculum.

LEGOUZIA, Delarb. Fl. Auv. ed. II. 45 (1800) = praec.

LEGUMINARIA, Bur. Monog. Bign. 42 (1864) = **Adenocalymma**, Mart. (Bignon.).

LEHMANNA, Casseb. & Theob. Fl. Wett. ed. II. 32 (1847) = **Gentiana**, Linn.
ciliata, Casseb. & Theob. l. c. = G. ciliata.

LEHMANNIA, Jacq. ex Steud. Nom. ed. II. ii. 21 (1841); Jacq. f. Ecl. ii. t. 108 (1844) = **Moschosma**, Reichb. (Labiat.).
ocymoidea, Jacq. ll. cc. = M. polystachya.

LEHMANNIA, Spreng. Anleit. ii. 1. 458 (1817) = **Nicotiana**, Linn. (Solan.).
tomentosa, Spreng. l. c. = N. tomentosa.

LEHMANNIA, Tratt. Ros. Monog. iv. 144 (1824) = **Potentilla**, Linn. (Rosac.).
lignosa, Tratt. l. c. 145 = P. lignosa.

LEIACHENIS, Rafin. Fl. Tellur. ii. 45 (1836) = **Aster**, Tourn. (Compos.).
cordifolia, Rafin. l. c. = A. cordifolius.
corymbosa, Rafin. l. c. = A. corymbosus.
dumosa, Rafin. l. c. = A. dumosus.
foliolosus, Rafin. l. c. = A. vimineus.
paludosa, Rafin. l. c. = A. paludosus.
paniculata, Rafin. l. c. = A. paniculatus.
sparsiflorus, Rafin. l. c. = A. sparsiflorus.

LEIANDRA, Rafin. Fl. Tellur. ii. 17 (1836) = **Tradescantia**, Linn. (Commel.).
cordifolia, Rafin. l. c. = T. cordifolia.
divaricata, Rafin. l. c. = T. virginica.

LEIANTHOSTEMON, Miq. Stirp. Surin. Sel. 147 (1850) = **Voyria**, Aubl. (Gentian.).
corymbosus, Miq. l. c. = V. corymbosa.
niveus, Miq. l. c. 148 = V. nivea.

LEIANTHUS, Griseb. Gen. et Sp. Gent. 196 (1839). *GENTIANEAE*, Benth. & Hook. f. ii. 807.
OMPHALOSTIGMA, Griseb. Gen. et Sp. Gent. 198 (1839).
PETASOSTYLIS, Griseb. in DC. Prod. ix. 71 (1845).
axillaris, *Hemsl. Biol. Centr. Am. Bot.* ii. 344.—Am. centr.
brevidentatus, *Hemsl. l. c.*—Guatemala.
cordifolius, *Griseb. Gen. et Sp. Gent.* 197.—Ind. occ.
cuspidatus, Griseb. in DC. Prod. ix. 82 = nigrescens.
exsertus, *Griseb. Gen. et Sp. Gent.* 197.—Jamaic.
lanceolatus, Benth. Pl. Hartw. 24 = Gentiana lanceolata.
latifolius, *Griseb. Gen. et Sp. Gent.* 199.—Jamaic.
longifolius, *Griseb. l. c.* 196.—Jamaic.
nigrescens, *Griseb. l. c.* 199.—Mexic.
saponarioides, *Griseb. l. c.* 198.—Mexic.
Seemanni, *Griseb. in Seem. Bot. Voy. Herald*, 170.—Costa Rica.
silenifolius, *Griseb. Cat. Pl. Cub.* 180.—Cuba.
Skinneri, *Hemsl. Biol. Centr. Am. Bot.* ii. 345.—Guatemala.
umbellatus, *Griseb. Gen. et Sp. Gent.* 199.—Jamaic.

LEIBNITZIA, Cass. in Dict. Sc. Nat. xxv. 420 (1822) = **Gerbera**, Gronov. (Compos.).
cryptogama, Cass. l. c. 421 = G. Anandria.
phanerogama, Cass. l. c. 422 = G. Anandria.

LEIBOLDIA, Schlecht. in Linnaea, xix. (1847) 742 = **Vernonia**, Schreb. (Compos.).
arctioides, Schlecht. l. c. 743 = V. arctioides.
ovata, Schlecht. l. c. 742 = V. Leiboldiana.

LEICESTERIA, Pritz. Ic. Ind. i. 628 (1855) = **Leycesteria**, Wall. (Caprifol.).

LEICHARDTIA, R. Br. in Sturt, Exped. Centr.-Austr. App. 81 (1849) = seq.

LEICHHARDTIA, R. Br. in Sturt, Exped. Centr. Austr. ii. App. 81 (1849) = **Marsdenia**, R. Br. (Asclep.).
australis, R. Br. l. c. = M. Leichhardtiana.

LEICHHARDTIA, F. Muell. Fragm. x. 67 (1876). *MENISPERMACEAE*.
clamboides, *F. Muell. l. c.* 68.—Austral.

LEICHHARDTIA, Sheph. Cat. Pl. Cult. Sydn. (1851) 15 = **Callitris**, Vent. (Conif.).
Macleayana, Sheph. l. c. = C. Macleayana.

LEIDESIA, Muell. Arg. in DC. Prod. xv. II. 792 (1866). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 310.
capensis, *Muell. Arg. l. c.* 793.—Afr. austr.
obtusa, *Muell. Arg. l. c.*—Afr. austr.

LEIENA, Rafin. Fl. Tellur. iv. 33 (1836) = **Restio**, Linn.

LEIGHIA, Cass. in Dict. Sc. Nat. xxv. 435 (1822) = **Viguiera**, H. B. & K. (Compos.).
anchusaefolia, DC. Prod. v. 580 = V. anchusaefolia.
angustifolia, DC. l. c. 582 = Aspilia buphthalmiflora.
arillata, DC. l. c. = Aspilia buphthalmiflora.
Baldwiniana, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 365.—Am. austr.
bicolor, Cass. in Dict. Sc. Nat. xxv. 436 = Helianthus angustifolius.
buphthalmiflora, DC. Prod. v. 582 = Aspilia buphthalmiflora.
calendulacea, DC. l. c. 582 = Aspilia buphthalmiflora.
debilis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 365.—Reg. Argent.
dissitifolia, DC. Prod. v. 581 = V. dissitifolia.
ecliptaefolia, DC. l. c. 583 = Aspilia ecliptaefolia.
elegans, Cass. in Dict. Sc. Nat. xxv. 435 = V. linearis.
Gilliesii, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 313 = V. Gilliesii.
grandiflora, Gardn. in Field. Sert. Pl. tt. 54, 55 = V. grandiflora.
Hookeriana, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 365 = Wyethia angustifolia.
immarginata, DC. Prod. v. 581 = V. stenophylla.
lanceolata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 365 = Helianthella uniflora ?
laxa, DC. Prod. v. 580 = V. laxa.
leptocephala, DC. l. c. 582.—Mexic. ?
linearis, DC. l. c. 581 = V. linearis.
lomatoneura, DC. l. c. = V. anchusaefolia.
longifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 365 = Helianthus longifolius.
microphylla, Cass. in Dict. Sc. Nat. xxv. 436 = Helianthus microphyllus.
nonneaefolia, DC. Prod. v. 581 = V. nonneaefolia.
ovatifolia, DC. l. c. 583 = V. ovatifolia.
silphioides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 314 = Aspilia silphioides.
speciosa, DC. Prod. v. 583 = Tithonia speciosa.
sphaerocephala, DC. l. c. 582 = V. sphaerocephala.
stenophylla, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 313 = V. stenophylla.
tucumanensis, Hook. & Arn. l. c. 314 = V. tucumanensis.
uniflora, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 364 = Helianthella uniflora.
urticiformis, DC. Prod. v. 582 = V. urticiformis.

LEIGHIA, Scop. Introd. 128 (1777) = **Ethulia**, Linn. (Compos.).

LEIMANISA, Rafin. New Fl. Am. iv. 93 (1836) = **Gentiana**, Linn.
pratensis, Rafin. l. c. = G. Amarella.

LEIMANTHEMUM, Ritg. in Marb. Schrift. ii. (1830) 130 = seq.

LEIMANTHIUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 24 = **Melanthium**, Linn. (Liliac.).
glaucum, Schult. f. Syst. vii. 1551 = Zygadenus glaucus.
hybridum, Sweet, Hort. Brit. ed. I. 429 = M. virginicum.
laetum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 24 = Zygadenus muscaetoxicus.
monoicum, Sweet, Hort. Brit. ed. I. 429 = Veratrum parviflorum.
Nuttallii, Hook. Fl. Bor. Am. ii. 177 = Zygadenus Nuttallii.
pallidum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 24 = Zygadenus muscaetoxicus.
sibiricum, Schult. f. Syst. vii. 1551 = Zygadenus sibiricus.
virginicum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 24 = M. virginicum.

LEINCKERIA, Neck. Elem. ii. 172 (1790) = seq.

LEINKERIA, Scop. Introd. 345 (1777) = **Roupala**, Aubl. (Proteac.).

LEIOCALYX, Planch. ex Hook. Niger Fl. 350 (1849) = **Dissotis**, Benth. (Melastom.).

LEIOCARPUS, Blume, Bijdr. 581 (1825) = **Aporosa**, Blume (Euphorb.).
arborescens, Hassk. Cat. Hort. Bog. Desc. 59 = A. arborescens.
arboreus, Blume, Bijdr. 582 = A. arborea.
arboreus, Miq. Fl. Ind. Bat. Suppl. 443 = A. Miqueliana.
fruticosus, Blume, Bijdr. 582 = A. fruticosa.
quadrilocularis, Miq. Fl. Ind. Bat. Suppl. 443 = A. quadrilocularis.
serratus, Hassk. Cat. Hort. Bog. Desc. 55, in obs. = A. microcalyx.

LEIOCARYA, Hochst. in Flora, xxvii. (1844) 30 = **Trichodesma**, R. Br. (Borag.).
Kotschyana, Hochst. l. c. = T. zeylanicum.

LEIOCHILUS, Benth. in Journ. Linn. Soc. xviii. (1881) 328 = **Leochilus**, Knowles & Westc. (Orchid.).

LEIOCHILUS, Hook. f. in Benth. & Hook. f. Gen. ii. 116 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
resinosus, *Hook. f. l. c.*—Madag.

LEIOCLUSIA, Baill. in Bull. Soc. Linn. Par. i. (1880) 244. *GUTTIFERAE*.
Boiviniana, *Baill. l. c.*—Madag.

LEIOLIGO, Rafin. Fl. Tellur. ii. 42 (1836) = **Solidago**, Vaill. (Compos.).
cesia, Rafin. l. c. = S. caesia.
corymbosa, Rafin. l. c. = S. Virgaurea ?
elata, Rafin. l. c. = S. petiolaris.
erecta, Rafin. l. c. = S. petiolaris.
lithospermifolia, Rafin. l. c. = S. sempervirens.
petiolaris, Rafin. l. c. = S. petiolaris.
pulverulenta, Rafin. l. c. = S. puberula.
pyramidata, Rafin. l. c. = S. pilosa.
rigida, Rafin. l. c. = S. rigida.
salicina, Rafin. l. c. = S. patula.
speciosa, Rafin. l. c. = S. speciosa.

LEIOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 94 = **Dalbergia**, Linn. (Legumin.)
nigrum, Benth. l. c. = D. nigra.

LEIOLOBIUM, Reichb. Consp. 184 (1828) = **Nasturtium**, R. Br. (Crucif.).
austriacum, Opiz, Seznam, 58 = N. austriacum.

LEIOPHYLLUM, Ehrh. Beitr. iv. 146 (1789) = **Scirpus**, Tourn. (Cyperac.).

LEIOPHYLLUM, Hedw. f. Gen. 313 (1806). *ERICACEAE*, Benth. & Hook. f. ii. 597.
AMMYRSINE, Pursh, Fl. Am. Sept. i. 301 (1814).
DENDRIUM, Desv. Journ. Bot. iii. (1814) 36.
FISCHERA, Sw. in Mém. Soc. Nat. Mosc. v. (1817) 14.
buxifolium, *Ell. Sketch*, i. 483.—Am. bor.
Lyoni, Sweet, Hort. Brit. ed. III. 444 = buxifolium.
prostratum, Loud. Arboret. 1155 = buxifolium.
serpyllifolium, DC. Prod. vii. 730 = buxifolium.
thymifolium, Steud. Nom. ed. I. 469 = buxifolium.

LEIOPYXIS, Miq. Fl. Ind. Bat. Suppl. 445 (1860) = **Cleistanthus**, Hook. f. (Euphorb.).
sumatrana, Miq. l. c. 446 = C. sumatranus.

LEIOSANDRA, Rafin. Fl. Tellur. iv. 89 (1836) = **Verbascum**, Tourn. (Scrophul.).
crassifolia, Rafin. l. c. = V. crassifolium.
cuspidata, Rafin. l. c. = V. Thapsus.

LEIOSPERMUM, D. Don, in Edinb. N. Phil. Journ. (Apr.–June 1830) 91 = **Weinmannia**, Linn. (Saxifrag.).
parviflorum, D. Don, l. c. = W. parviflora.
racemosum, D. Don, l. c. = W. racemosa.

LEIOSPERMUM, Wall. Cat. n. 6923 (1832?) = **Psilotrichum**, Blume (Amarant.).
ferrugineum, Wall. Cat. n. 6923 = Ps. ferrugineum.
ferrugineum, Wight, Ic. t. 721 = Ps. trichotomum.

LEIOSTEGIA, Benth. in Hook. Journ. Bot. ii. (1840) 294 = **Comolia**, DC. (Melastom.).
vernicosa, Benth. l. c. = C. vernicosa.

LEIOSTEMON, Rafin. Neogenyt. 2 (1825) = **Pentstemon**, Mitch. (Scrophul.).
frutescens, Rafin. l. c. = P. frutescens.
purpureum, Rafin. Atl. Journ. 145.—Am. bor.

LEIOTHAMNUS, Griseb. Gen. et Sp. Gent. 205 (1839) = **Lisianthus**, Linn. (Gentian.).
anomalus, Griseb. l. c. = Lisianthus anomalus.
Elisabethae, Schomb. in Verh. Preuss. Ver. Gartenb. xviii. (1847) 155. t. 1 = Lisianthus Elisabethae.

LEIOTULUS, Ehrenb. in Linnaea, iv. (1829) 399 = **Malabaila**, Hoffm. (Umbellif.).
alexandrinus, Ehrenb. l. c. = M. Sekakul.

LEIPHAIMOS, Schlecht. & Cham. in Linnaea, vi. (1831) 387 = **Voyria**, Aubl. (Gentian.).
aurantiacus, Miq. Stirp. Surin. Sel. 148 = V. aurantiaca.
mexicanus, Miq. l. c. = V. mexicana?
nudus, Miq. l. c. = V. nuda.
pallidus, Walp. Ann. iii. 83 = V. aphylla.
parasitica, Schlecht. & Cham. in Linnaea, vi. (1831) 387 = V. mexicana.
spathaceus, Miq. Stirp. Surin. Sel. 148 = V. spathacea.
tenellus, Miq. l. c. = V. tenella.
tenuiflorus, Miq. l. c. = V. tenuiflora.
uniflorus, Miq. l. c. = V. aphylla.

LEITGEBIA, Eichl. in Mart. Fl. Bras. xiii. 1. 413 (1871). *VIOLARIEAE.*
guianensis, *Eichl. l. c.*—Guiana.

LEITNERIA, Chapm. Fl. S. U. St. 427 (1860). *LEITNERIEAE*, Benth. & Hook. f. iii. 397.
floridana, *Chapm. l. c.* 428.—Am. bor.

LEJICA, DC. Prod. v. 534, in syn. (1836) = LEPIA, Hill = **Zinnia**, Linn. (Compos.).

LELOUTRIA, Gaudich. Bot. Voy. Bonite, t. 110. (1844) 52 = **Dolia**, Lindl. (Convolv.).
aplocaryoides, Gaudich. l. c.—Am. austr.

LEMAIREA, De Vriese, in Nat. Verh. Haarl. x. (1854) (Gooden.) 189. 'Genus delendum,' Benth. & Hook. f. ii. 537.
amboinensis, De Vriese, l. c.—Ins. Amboina.

LEMBOTROPIS, Griseb. Spicil. Fl. Rumel. i. 10 (1843) = **Cytisus**, Linn. (Umbell.).
affinis, Presl, Bot. Bemerk. 138 = C. affinis.
nigricans, Griseb. Spicil. Fl. Rumel. i. 10 = C. nigricans.
sessilifolia, C. Koch, Dendrol. i. 22 = C. sessilifolius.
triflora, Presl, Bot. Bemerk. 138 = C. triflorus.

LEMIA, Vand. Fl. Lusit. 35. t. 2. f. 15 (1788); et in Roem. Script. 116. t. 7. f. 15 (1796) = **Portulaca**, Tourn.

LEMMATIUM, DC. Prod. v. 669 (1836) = **Calea**, Linn. (Compos.).
rotundifolium, DC. l. c.; et in Deless. Ic. Sel. iv. t. 44 = C. rotundifolia.

LEMMONIA, A. Gray, in Proc. Am. Acad. xii. (1877) 162. *HYDROPHYLLACEAE.*
californica, *A. Gray, l. c.*—Calif.

LEMNA, Linn. Syst. ed. I (1735). *LEMNACEAE*, Benth. & Hook. f. iii. 1001.
HYDROPHACE, Hall. Enum. Stirp. Helv. iii. 68 (1768).
LENTICULA, Mich. ex Adans. Fam. ii. 471 (1763).
SPIRODELA, Schleid. in Linnaea, xiii. (1839) 391.
STAUROGETON, Reichb. Nom. 33 (1841).
TELMATOPHACE, Schleid. in Linnaea, xiii. (1839) 391.
aequinoctialis, Welw. Apont. 578 = angolensis.
angolensis, *Welw. in Journ. Bot.* iii. (1865) 112.—Afr. trop.
arrhiza, Linn. Mant. ii. 294 = Wolffia Michelii.
arrhiza, Willd. ex Schleid. in Linnaea, xiii. (1839) 390 = Wolffia Delilii.
bannatica, Waldst. & Kit. ex Schleid. l. c. 392 = polyrrhiza.
bisulca, Veesenm. in Beitr. Pfl. Russ. Reich. ix. 104 = trisulca?
conjugata, Willd. ex Schleid. in Linnaea, xiii. (1839) 391 = minor.
cruciata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 566 = trisulca.
cyclostasa, Ell. ex Chev. Fl. Par. ii. 256; Schleid. in Linnaea, viii. (1839) 390 = minor.
dimidiata, Rafin. in Am. Monthly Mag. ii. (1817) 43 Ricciae sp. (Crypt.).
gibba, Blanco, Fl. Filip. ed. I. 672 = oligorrhiza.
gibba, *Linn. Sp. Pl.* 970.—Reg. bor. et austr.
gibba, *Seem. ex Kurz, in Journ. Linn. Soc.* ix. (1867) 266 (an praec.?).—Ins. Fiji.
gibba, Weber, ex Mert. & Koch, Deutschl. Fl. i. 295 = minor.
globosa, Roxb. Hort. Beng. 66; Fl. Ind. iii. 565 = Wolffia Michelii.
hyalina, Delile, Fl. Aegypt. 27 = Wolffia Delilii.
intermedia, Ruthe, ex Schleid. in Linnaea, xiii. (1839) 391 = trisulca.
javanica, Bauer, ex Hegelm. Lemnac. 150 = oligorrhiza.
major, Griff. Notul. iii. 216 = polyrhiza.
melanorrhiza, F. Muell. ex Kurz, in Journ. Bot. v. (1867) 115 = oligorrhiza.
microscopica, Schur, Enum. Pl. Transs. 635 = Wolffia Michelii.
minima, Chev. Fl. Par. ii. 256 = minor.
minima, Phil. in Linnaea, xxxiii. (1864–65) 239, in syn. = *minuta*.
minor, Griff. Notul. iii. 216 = paucicostata.
minor, *Linn. Sp. Pl.* 970.—Amphig.
minuta, Desv. ex Steud. Nom. ed. I. 469 = Marsilea minuta (Crypt.).
minuta, H. B. & K. Nov. Gen. et Sp. i. 372 = minor.
minuta, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid?).—Ind. or.
obcordata, Boj. Hort. Maurit. 358 = minor.
obcordata, Buch.-Ham. in Wall. Cat. n. 5201 (Quid?).—Ind. or.
obcordata, Vahl, Symb. Bot. ii. 95 = Ricciae sp. (Crypt.).
oblonga, *Phil. in Linnaea*, xxix. (1857–58) 45.—Chili.
oligorrhiza, *Kurz, in Journ. Linn. Soc.* ix. (1867) 267.—Austral.; As. et Am. trop.
orbicularis, Kit. ex Schult. Oestr. Fl. ed. II. i. 64 = polyrhiza.
orbiculata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 565 = polyrhiza.
ovata, A. Br. ex Krauss, in Flora, xxviii. (1845) 344 = minor.
palustris, Haenke, ex Mert. & Koch, Deutschl. Fl. i. 295 = minor.
paucicostata, *Hegelm. Lemnac.* 138. t. 8.—As. et Am. trop.
perpusilla, *Torr. Fl. N. York*, i. 245.—Am. bor.
pleiorrhiza, F. Muell. in Journ. Bot. v. (1867) 115 = oligorrhiza.
polyrhiza, *Linn. Sp. Pl.* 970.—Reg. bor. et austr.
punctata, *G. F. W. Mey. Prim. Fl. Esseq.* 262.—Guian.
quadrifolia, Steud. Nom. ed. I. 469 = Marsilia quadrifolia (Crypt.).
tenera, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 78.—Ind. or.
thermalis, Beauv. ex Nutt. Gen. Am. i. 19 = polyrhiza.
Torreyi, Austin, in A. Gray, Man. Bot. N. U. St. (1867) 479 = valdiviana.

LEMNA:—
transsilvanica, Schur, Enum. Pl. Transs. 635 = polyrhiza.
trichorrhiza, Thuill. ex Schleid. in Linnaea, xiii. (1839) 391 = gibba, *Linn.*
trisulca, *Linn. Sp. Pl.* 970.—Reg. bor. et austr.
umbonata, A. Br. ex Hegelm. Lemnac. 156 = polyrhiza.
Valdesiana, S. Wats. in Bot. King's Exp. 336, sphalm. = seq.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864) 239.—Chili; Mexic.
vulgaris, Lam. Encyc. iii. 464 = gibba, minor.

LEMNESCIA, Willd. Sp. Pl. ii. 1172 (1799) = seq.

LEMNISCIA, Schreb. Gen. 358 (1789) = **Vantanea**, Aubl. (Humiriac.).
floribunda, Willd. Sp. Pl. ii. 1172 = V. guianensis.
guianensis, J. F. Gmel. Syst. 817 = V. guianensis.

LEMNOPSIS, Zipp. in Flora, xii. (1829) i. 285 (Quid?). *PODOSTEMACEAE*?
major, Zoll. Syst. Verz. Ind. Archip. 75.—Malaya.
minioides, Zipp. in Flora, xii. (1829) i. 285; et in Alg. Konst. en Letter-bode, i. (1829) 297.—Malaya.
minor, Zoll. Syst. Verz. Ind. Archip. 75.—Malaya.

LEMONIA, Lindl. Bot. Reg. (1840) t. 59 = **Ravenia**, Vell. (Rutac.).
spectabilis, Lindl. l. c. = R. spectabilis.

LEMONIA, Pers. Syn. i. 44 (1805), ut sect. = LOMENIA, Pourr. = **Watsonia**, Mill. (Irid.).

LEMOTRIS, Rafin. Fl. Tellur. ii. 26 (1836) = seq.

LEMOTRYS, Rafin. Fl. Tellur. iii. 50 (1836) = **Camassia**, Lindl. (Liliac.).
hyacinthina, Rafin. l. c. 51 = C. esculenta.

LENCYMMAEA, Presl, Epim. Bot. 211 (1849). 'Genus dubium,' Benth. & Hook. f. i. 735.
salicifolia, *Presl, l. c.*—Burma.

LENIDIA, Thou. Gen. Nov. Madag. 17 (1806) = **Wormia**, Rottb. (Dillen.).
alata, Poir. in Dict. Sc. Nat. xxv. 448 = W. alata.
dentata, Poir. l. c. = W. triquetra.
madagascariensis, Poir. Encyc. Suppl. iii. 330 = W. madagascariensis.
triquetra, Poir. in Dict. Sc. Nat. xxv. 448 = W. triquetra.

LENNEA, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Hort. Berol. ii. 65. t. 26 (1842). *LEGUMINOSAE*, Benth. & Hook. f. i. 500.
robinioides, *Klotzsch, l. c.*—Mexic.
viridiflora, *Seem. Bot. Voy. Herald*, 107.—Uruguay.

LENNOA, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 7 (1824). *LENNOACEAE*, Benth. & Hook. f. ii. 622.
CORALLOPHYLLUM, H. B. & K. Nov. Gen. et Sp. vii. 275. t. 660 bis (1825).
arenaria, Fourn. in Bull. Soc. Bot. Fr. xvi. (1869) 11 = Pholisma arenarium.
caerulea, *Fourn. l. c.; Solms, in Abh. Naturf. Ges. Halle*, xi. (1870) 176.—Mexic.
madrepoides, Steud. Nom. ed. II. ii. 22, sphalm. = seq.
madreporoides, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 7.—Mexic.

LENORMANDIA, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Gram. 359 (1855) = **Andropogon**, Linn. (Gramin.).
insignis, Steud. ll. cc. = A. squarrosus.

LENS, [Tourn.] Linn. Syst. ed. I (1735); Moench, Meth. 131 (1794); Gren. & Godr. Fl. Fr. i. 476 (1848). *LEGUMINOSAE*, Benth. & Hook. f. i. 525.
abyssinica, Alef. in Bonplandia, ix. (1861) 131 = esculenta.
Biebersteinii, Lamotte, Prod. i. 220 = nigricans.
camelorum, Webb & Berth. Phyt. Canar. ii. 97 = esculenta.
culinare, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 361 = esculenta.
cyanea, *Alef. in Bonplandia*, ix. (1861) 129.—Persia.

LENS:—
disperma, Webb & Berth. Phyt. Canar. ii. 97 = esculenta.
esculenta, *Moench, Meth.* 131.—Oriens.
himalayensis, *Alef. in Bonplandia*, ix. (1861) 130 (Quid ?).—Reg. Himal.
Hohenackeri, Webb & Berth. Phyt. Canar. ii. 97 = L. Lenticula.
Lenticula, *Alef. in Bonplandia*, ix. (1861) 129.—Europ.
lentoides, Webb & Berth. Phyt. Canar. ii. 97 = nigricans.
monantha, Moench, Meth. 131 = Vicia monanthos.
nigricans, *Godr. Fl. Lorr.* ed. I. i. 173.—Reg. Mediterr.
Nummularia, Alef. in Bonplandia, ix. (1861) 130 = esculenta.
sativa, Hell. Fl. Wirceb. ii. 169 = esculenta.
Schnittspahni, Alef. in Bonplandia, ix. (1861) 129 (= *Ervum orientale*).—Persia; Ind. or.
Tenorei, Lamotte, Prod. i. 220, in obs. = nigricans.
uniflora, Schur, Enum. Pl. Transs. 171 = L. Lenticula.
vulgaris, Delarb. Fl. Auv. ed. II. 472 = esculenta.

LENTAGO, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87 = **Viburnum**, Tourn. (Caprifol.).

LENTIBULARIA, Adans. Fam. ii. 208 (1763) = **Utricularia**, Linn. (Lentib.).
major, Gilib. Fl. Lituan. i. 139 = U. vulgaris.
minor, Rafin. Fl. Tellur. iv. 108 = U. minor.
vulgaris, Moench, Meth. 520 = U. vulgaris.

LENTICULA, Mich. ex Adans. Fam. ii. 471 (1763) = **Lemna**, Linn.
arrhiza, Lam. Fl. Fr. ii. 190 = Wolffia Michelii.
cyclostasa, Kurz, in Journ. Linn. Soc. ix. (1867) 266 = Lemna minor.
gibba, Moench, Meth. 319 = Lemna gibba.
gibbosa, Renault, Fl. Dép. Orne, 40 = Lemna gibba.
minima, Kurz, in Journ. Linn. Soc. ix. (1867) 266 = Lemna minor.
minor, Scop. Fl. Carn. ed. II. ii. 213 = Lemna minor.
polyrhiza, Lam. Fl. Fr. ii. 189 = Lemna polyrhiza.
ramosa, Lam. l. c. = Lemna trisulca.
trisulca, Scop. Fl. Carn. ed. II. ii 213 = Lemna trisulca.
vulgaris, Lam. Fl. Fr. ii. 189 = Lemna minor, gibba.

LENTISCUS, [Tourn.] Linn. Syst. ed. I (1735) = **Pistacia**, Linn. (Anacard.).
massiliensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = P. Lentiscus.
vulgaris, Fourr. l. c. xvi. (1868) 356 = P. Lentiscus.

LENZIA, Phil. in Linnaea, xxxiii. (1864–65) 222. Gen. dub. *AMARANTAC.*—Cf. Benth. & Hook. f. iii. 23.
Chamaepitys, *Phil. l. c.*—Chili.

LEOBARDIA, Pomel, Nouv. Mat. Fl. Atl. 162 (1874) = seq.

LEOBORDEA, Delile, in Laborde, Voy. Arab. Pétr. 86 (1833) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 161 = Lot. Leobordea.
argyrolobioides, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 237 = Lot. genistoides.
clandestina, Steud. Nom. ed. II. ii. 23 = Lot. clandestina.
cytisoides, Bal. ex Boiss. Fl. Orient. ii. 31 = Lot. genistoides.
genistoides, Fenzl, Pugill. Pl. Nov. Syr. 6, adnot. = Lot. genistoides.
lotoidea, Delile, in Laborde, Voy. Arab. Pétr. 86 = Lot. Leobordea.
lotoides, Fenzl, Pugill. Pl. Nov. Syr. 6, sphalm. = Lot. genistoides.
lupinifolia, Boiss. ex Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 237 = Lot. lupinifolia.
persica, Jaub. & Spach, l. c. 235 = Lot. Leobordea.
porrecta, Steud. Nom. ed. II. ii. 23 = Lot. porrecta.
sericea, Ledeb. Fl. Ross. i. 512 = Lot. genistoides.
sphaerosperma, Jaub. & Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 235 = Lot. Leobordea.
villosa, Pomel, Nouv. Mat. Fl. Atl. 162.—Alger.

LEOCHILUS, Knowles & Westc. Flor. Cab. ii. (1838) 143. *ORCHIDEAE*, Benth. & Hook. f. iii. 564.
Cryptosaccus, Reichb. f. Xenia Orch. i. 15 (1858).
Cryptosanus, Scheidw. in Otto & Dietr. Allg. Gartenz. xi. (1843) 101.

LEOCHILUS:—
Leiochilus, Benth. in Journ. Linn. Soc. xviii. (1881) 328.
Rhynchostelis, Reichb. f. in Bot. Zeit. x. (1852) 770.
carinatus, *Lindl. Bot. Reg.* (1842). *Misc.* 23.—Mexic.
cochlearis, *Lindl. l. c.*—Jamaic.
herbaceus, *Lindl. l. c.* (1844). *Misc.* 90.—Venezuel.
Leiboldi, Reichb. f. in Linnaea, xviii. (1844) 404 = Papperitzia Leiboldi.
oncidioides, *Knowles & Westc. Flor. Cab.* ii. (1838) 143.—Mexic.
salvus, *Griseb. Cat. Pl. Cub.* 267.—Cuba.
sanguinolentus, *Lindl. Bot. Reg.* (1844). *Misc.* 91.—Peruv.
scriptus, *Reichb. f. Xenia Orch.* 15.—Mexic.

LEONHARDIA, Opiz, in Lotos, vii. (1857) 88 = **Ulex**, Linn. (Legumin.).
Boivini, Opiz, l. c. 89 = U. Boivini.
Cossonii, Opiz, l. c. = U. Cossonii.
Escayracii, Opiz, l. c. = U. Escayracii.
lurida, Opiz, l. c. = U. Webbianus.
megalorites, Opiz, l. c. = U. Boivini.
Salzmanni, Opiz, l. c. = U. Boivini.
Vaillantii, Opiz, l. c. = U. Boivini.
Webbiana, Opiz, l. c. = U. Webbianus.

LEONIA, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 6 (1825) = **Salvia**, Tourn. (Labiat.).
salvifolia, Cerv. l. c. = S. Leonia.

LEONIA, Mutis, ex Kunth, Syn. 462 (1822) = **Siparuna**, Aubl. (Monimiac.).

LEONIA, Ruiz & Pav. Fl. Per. ii. 69. t. 222 (1799). *VIOLARIEAE*, Benth. & Hook. f. i. 119.
Steudelia, Mart. Nov. Gen. et Sp. t. 168 (1827).
cymosa, *Mart. Nov. Gen. et Sp.* ii. 88.—Bras.
glycycarpa, *Ruiz & Pav. Fl. Per.* ii. 69.—Bras.
paniculata, Mart. in Flora, xxiv. (1841) II. Beibl. 26 = Villaresia paniculata.
racemosa, Mart. Nov. Gen. et Sp. ii. 86. t. 168 = glycycarpa.

LEONICENIA, Scop. Introd. 212 (1777) = **Miconia**, Ruiz & Pav. (Melastom.).

LEONITIS, Spach, Hist. Vég. Phan. ix. 210 (1840) = seq.

LEONOTIS, R. Br. in Ait. Hort. Kew. ed. II. iii. 409 (1811). *LABIATAE*, Benth. & Hook. f. ii. 1214.
Hemisodon, Rafin. Fl. Tellur. iii. 88 (1836).
capensis, Rafin. Fl. Tellur. iii. 88 = L. Leonitis.
caribea, Rafin. l. c. = Leucas martinicensis.
dubia, *E. Mey. Comm. Pl. Afr. Austr.* 242.—Afr. austr.
dysophylla, *Benth. in E. Mey. l. c.*—Afr. austr.
hirtiflora, *Benth. in DC. Prod.* xii. 536.—Afr. austr.
intermedia, *Lindl. Bot. Reg.* t. 850.—Afr. austr.
Leonitis, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iii. 410.—Afr. austr.
Leonurus, [*R. Br.*] *l. c.*—Afr. austr.
mollis, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 242.—Afr. austr.
nepetaefolia, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iii. 409.—Amphig. trop.
nepetaefolia, Schimp. ex Benth. in DC. Prod. x. 535 = rugosa.
ovata, Boj. Hort. Maurit. 250 = nepetaefolia.
ovata, Spreng. Syst. ii. 744 = L. Leonitis.
pallida, *Benth. Lab. Gen. et Sp.* 619.—Afr. trop.
parvifolia, Benth. l. c. = dubia.
Raineriana, *Vis. Ort. Bot. Padov.* (1842) 142.—Afr. bor.
rugosa, *Benth. in DC. Prod.* xii. 535.—Abyssin.
tuberosa, *Hoffmgg. Verz. Pfl.* 71.—Sibir.
velutina, Fenzl, in Flora, xxvii. (1844) 312 = Raineriana.

LEONTIA, Reichb. Consp. 195 (1828) = LUNTIA, Neck. = **Croton**, Linn. (Euphorb.).

LEONTICE, Linn. Syst. ed. I (1735). *BERBERIDEAE*, Benth. & Hook. f. i. 43.
Gymnospermium, Spach, Hist. Vég. Phan. viii. 66 (1839).

LEONTICE:—
Leontopetalon, Tourn. ex Adans. Fam. ii. 433 (1763).
Alberti, *Regel, Gartenfl.* (1881) 293.—Turkestan.
altaica, Pall. in Act. Acad. Petrop. i. (1779) II. 257. t. 8. f. 1–3 = Bongardia Rauwolfii.
armeniaca, Boiv. in Bélang. Voy. Ind. Or. Icon. absque num. (1846) = minor.
brevibracteata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Leontopetalum.
chrysogonia, St. Lag. l. c. = seq.
Chrysogonum, Linn. Sp. Pl. 312 = Bongardia Rauwolfii.
darwasica, *Regel, in Act. Hort. Petrop.* viii. (1884) 692.—Bokhara.
Eversmanni, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1848) 131.—Persia; Turkest.
incerta, Pall. Reise, iii. 726 = L. Leontopetalum.
leontopetaloides, Linn. Sp. Pl. 313 = Tacca pinnatifida.
Leontopetalum, Hook. f. & Thoms. Fl. Ind. i. 229 = Eversmanni.
Leontopetalum, *Linn. Sp. Pl.* 312.—Ital.; Oriens.
microrrhyncha, *S. Moore, in Journ. Linn. Soc.* xvii. (1879) 377.—China.
minor, Boiss. Fl. Orient. i. 100.—Persia.
odessana, Fisch. ex G. Don, Gen. Syst. i. 119 = Bongardia Rauwolfii.
Oliverii, Steud. Nom. ed. II. ii. 23 = Bongardia Rauwolfii.
secunda, Griff. Notul. iv. 701; Ic. Pl. Asiat. t. 647 = Bongardia Rauwolfii.
Smirnowii, *Trautv. in Act. Hort. Petrop.* vii. (1880) 405.—Reg. Caucas.
thalictrifolium, Salisb. Prod. 252 = seq.
thalictroides, Linn. Sp. Pl. 312 = Caulophyllum thalictroides.
triphylla, Sm. in Rees, Cycl. xx. n. 5 = Achlys triphylla.
vesicaria, Willd. Sp. Pl. ii. 148 = L. Leontopetalum.

LEONTOCHIR, Phil. in Anal. Univ. Chil. (1873) 545 (Descr. Nuev. Pl. ii. 68). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 737.
Ovallei, *Phil. ll. cc.*—Chili.

LEONTODON, Adans. Fam. ii. 112 (1763) = **Taraxacum**, Hall. (Compos.).

LEONTODON, Linn. Gen. ed. I. 239 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 520.
Antodon, Neck. Elem. i. 58 (1790).
Apargia, Scop. Fl. Carn. ed. II. ii. 113 (1772).
Apargidium, Torr. & Gray, Fl. N. Am. ii. 474 (1841).
Asterothrix, Cass. in Dict. Sc. Nat. xlviii. 422, 434 (1827).
Baldingeria, F. W. Schmidt, Samml. Phys. Aufs. i. 277 (1795).
Colobium, Roth, in Roem. Archiv, i. I. 36 (1796).
Deloderium, Cass. in Dict. Sc. Nat. xlviii. 430 (1827).
Fidelia, Sch. Bip. in Flora, xvii. (1834) 482.
Hedypnois, [Tourn.]. Scop. Fl. Carn. ed. II. ii. 99 (1772).
Hemilepis, Kunze, Ind. Sem. Hort. Lips. (1838); ex Schlecht. in Bot. Zeit. x. (1852) 875.
Kalbfussia, Sch. Bip. in Flora, xvi. (1833) 723.
Millina, Cass. in Dict. Sc. Nat. xxxi. 89 (1824).
Oporinia, D. Don, in Edinb. N. Phil. Journ. (1828-29) 309.
Plancia, Neck. Elem. i. 49 (1790).
Scorzoneroides, Moench, Meth. 549 (1794).
Streckera, Sch. Bip. in Flora, xvii. (1834) 483.
Thrica, S. F. Gray, Nat. Arr. Brit. Pl. ii. 428 (1821).
Thrincia, Roth, in Roem. Archiv, i. I. 36 (1796); et Catal. i. 97 (1797).
Thrixia, Dulac, Fl. Hautes-Pyr. 495 (1867).
Virea, Adans. Fam. ii. 112 (1763).
acutissimus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = Lactuca indica.
adhaerens, *Fenzl, ex Ball, in Ann. & Mag. Nat. Hist.* Ser. II. vi. (1850) 15.—As. Min.
alpestris, Tausch, in Flora, iv. (1821) 564 = Taraxacum officinale.
alpinus, Guss. Pl. Rar. Sic. 395 = Taraxacum apenninum.
alpinus, Jacq. Fl. Austr. i. 58 = hastilis.
alpinus, Stev. in Mém. Soc. Nat. Mosc. iv. (1812–13) 100 = Taraxacum officinale.
alpinus, Vill. Prosp. 34 = incanus.

LEONTODON:—
annuus, *DC. Prod.* vii. 104.—Aegypt.
anomalus, *Ball, in Ann. & Mag. Nat. Hist.* Ser. II. vi. (1850) 9.—Italia.
apenninus, Tenore, Rel. Viagg. Abruzz. 83 = Taraxacum appenninum.
apiculatus, Ledeb. ex Reichb. Fl. Germ. Excurs. 270 = Taraxacum officinale.
arabicus, *Boiss. Fl. Orient.* iii. 728.—Arab.
arcuatus, Tausch, in Flora, iv. (1821) 567 = Taraxacum officinale.
asper, Forsk. Fl. Aegypt. Arab. 145 = annuus.
asper, *Reichb. Fl. Germ. Excurs.* 252.—Europ. austr.; Oriens.
asperrimus, *Boiss. ex Ball, in Ann. & Mag. Nat. Hist.* Ser. II. vi. (1850) 15.—As. Min.; Persia.
asperrimus, Schur, Enum. Pl. Transs. 358 = hispidus.
aurantiacus, Reichb. Fl. Germ. Excurs. 853 = pyrenaicus.
aureus, Georgi, Beschr. Russ. Reich. iv. 1209 (Quid?).—Ross. med.
aureus, Linn. Syst. ed. X. 1193 = Crepis aurea.
autumnalis, *Linn. Sp. Pl.* 798.—Europ.
autumnalis, Oed. Fl. Dan. t. 501 = Crepis tectorum.
Balansae, *Boiss. Diagn.* Ser. II. iii. 89.—Algeria.
Berinii, *Reichb. Fl. Germ. Excurs.* 253.—Europ. austr.
bessarabicus, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 34 = Taraxacum palustre.
bicolor, Turcz. ex DC. Prod. vii. 149 = Taraxacum bicolor.
biscutellaefolius, DC. Prod. vii. 103 = asper.
biscutellaefolius, Heldr. ex Boiss. Diagn. Ser. I. xi. 41 = oxylepis.
borealis, *DC. Prod.* vii. 102.—Am. bor. occ.
Boryi, *Boiss. in DC. l. c.* 103.—Hispan.
Bourgaeanus, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 217.—Hispan.
brachyglossus, Peterm. Fl. Bienitz, 53 = autumnalis.
Brumati, Reichb. Fl. Germ. Excurs. 253 = hastilis.
bulbosus, Linn. Sp. Pl. 798 = Crepis bulbosa.
carolinianus, Walt. Fl. Carol. 192 = Pyrrhopappus carolinianus.
carpetanus, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 96; (*Pugill.* 146).—Hispan.
caucasicus, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 34 = hastilis.
caucasicus, Stév. in Mém. Soc. Nat. Mosc. iv. (1812–13) 100; Bieb. Fl. Taur. Cauc. iii. 530 = Taraxacum officinale.
ceratophorus, Ledeb. Fl. Alt. iv. 149, in nota = Taraxacum officinale.
chilensis, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 31 = Taraxacum Gilliesii.
chillensis, *DC. Prod.* vii. 105 (chilensis).—Ecuador.
chondrillaefolius, Scop. Fl. Carn. ed. II. ii. 112 = hispidus.
cichoraceus, *Boiss. Fl. Orient.* iii. 729.—Europ. austr.
cichoriaceus. Schleich. Cat. Pl. Helv. (1821) 22; et ex Reichb. Fl. Germ. Excurs. 270 = Taraxacum officinale.
ciliatus, Scop. Fl. Carn. ed. II. ii. 112 = Hypochoeris radicata.
collinus, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) II. 102.—Sibir. baical.
contortus, Dum. Fl. Belg. 63 = autumnalis?
corniculatus, Kit. in Schult. Oestr. Fl. ed. II. ii. 406 = Taraxacum officinale.
coronopifolius, *Comm. ex Sch. Bip. in Nov. Act. Nat. Cur.* xii. I. (1845) 124.—Reg. Magellan.
coronopifolius, Desf. Fl. Atlant. ii. 229. t. 214 = Picris radicata.
corymbosus, Borkh. in Roem. Neu. Mag. Bot. 24 = Picris hieracioides.
crepidiformis, Lam. Encyc. iii. 530 = pyrenaicus.
creticus, *Boiss. Fl. Orient.* iii. 728.—Ins. Creta.
crispatus, Griseb. & Schenk, in Linnaea, xxv. (1852) 610 = hispidus.
crispus, DC. ex Nym. Consp. 468 = hastilis.
crispus, *Vill. Prosp.* 34; *Hist. Pl. Dauph.* iii. 84. t. 25.—Europ. austr.
croceus, Haenke, in Jacq. Coll. ii. 16 = pyrenaicus.
dandaleus, Sieber, ex Steud. Nom. ed. II. ii. 23 (Quid?).—Creta.
Dandelion, Hill, Veg. Syst. iii. 161 = Taraxacum officinale.
Dandelion, Linn. Sp. Pl. 798 = Kirgia Dandelion.
danubialis, Jacq. Enum. Hort. Vindob. 139, 270 = hastilis.
dentatus, Linn. Mant. i. 107 = Crepis pygmaea.

LEONTODON:—
dentatus, Tausch, in Flora, vii. (1824) 248 = Taraxacum palustre.
dissectus, Ledeb. in Mém. Acad. Pétersb. v. 1812 (1814) 555 = Taraxacum dissectum.
divaricatus, Boiss. Diagn. Ser. I. xi. 41 = adhaerens.
dubius, Reichb. Fl. Germ. Excurs. 252 = hastilis.
dubius, *Timb. in Bull. Soc. Sc. Phys.-Nat. Toul.* vi. (1883–84) 177.—Gallia.
elegans, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 153 = Hieracium fragile, *Banks & Soland.*
erectus, Mayer, Arb. Eintr. Fr. i. III. 69; ex Schrank, Baier. Fl. ii. 315 = Taraxacum palustre.
eriopodon, D. Don, in Mem. Wern. Soc. iii. (1820) 413 = Taraxacum officinale.
eriopus, Spreng. Syst. iii. 658 = praec.
erythrospermus, Eichw. Skizze, 150 = Taraxacum officinale.
fasciculatus, *Nym. Syll.* 53.—Europ. austr.
glaber, Baill. Hist. Pl. viii. 294 = Hypochoeris glabra.
glaber, Gilib. Fl. Lituan. i. 229 = autumnalis.
glaber, *Ucria, in Roem. Arch.* i. 1. 70.—Sicilia.
glaber, Vukot. in Rad Jugos. Akad. Zagreb. xxxiv. (1876) 120 = hastilis?
glandulosus, Pav. ex Fries, Symb. Hierac. 136 = Hieracium peruanum.
glaucanthus, Ledeb. Fl. Alt. iv. 151 = Taraxacum officinale.
glaucescens, Bieb. Fl. Taur. Cauc. iii. 530 = Taraxacum officinale.
Gouani, DC. Prod. vii. 101 = pyrenaicus.
graecus, *Boiss. & Heldr. Diagn.* Ser. I. xi. 39.—Graecia.
Gussoni, Spreng. Syst. iii. 658 = Crepis bursifolia.
gymnanthus, Link, in Linnaea, ix. (1834) 582 = Taraxacum gymnanthum.
hastilis, *Linn. Sp. Pl.* ed. II. 1123.—Europ.; Oriens.
Heldreichianus, Boiss. Fl. Orient. iii. 731 = graecus.
helminthioides, *Coss. & Dur. in Bull. Soc. Bot. Fr.* xxii. (1875) 62; *et ex Ball, in Journ. Linn. Soc.* xvi. (1878) 544.—Afr. bor.
helveticus, Mérat, in Ann. Sc. Nat. Sér. I. xxii. (1831) 108 = pyrenaicus.
hieracioides, DC. Prod. vii. 105 = Picris hieracioides.
hirsutus, Hook. Fl. Bor. Am. i. 296 = Troximon humile.
hirtus, *Linn. Syst.* ed. X. 1194.—Europ.
hirtus, Pall. ex Georgi, Beschr. Russ. Reich. iv. 1210 = asper.
hispanicus, Bess. Enum. Pl. Volh. 31 = asper.
hispanicus, *Poir. Encyc. Suppl.* iii. 453; *Mérat, in Ann. Sc. Nat.* Sér. I. xxii. (1831) 106.—Hispan.
hispidulus, *Boiss. Fl. Orient.* iii. 727.—Afr. bor.; Oriens.
hispidus, Cav. Ic. t. 149 = hispanicus.
hispidus, Coss. & Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 217 = Bourgaeanus.
hispidus, *Linn. Sp. Pl.* 799.—Europ.
hispidus, Poll. Hist. Pl. Palat. ii. 382 = hirtus.
hispidus, Scop. Fl. Carn. ed. II. ii. 112 = crispus.
hispidus, Vand. Fasc. Pl. 18 = Hieracium intybaceum.
Huguenini ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 110.—Europ.
hyoseroides, DC. Prod. vii. 104 = hirtus?
hyoseroides, Welw. ex Reichb. Fl. Germ. Excurs. 863 = hastilis.
incanus, *Schrank, Baier. Reise*, 14; *DC. Fl. Fr.* iv. 56.—Europ.
infestus, Salisb. Prod. 181 = Taraxacum officinale.
keretinus, F. Nyland. Spicil. Pl. Fennic. i. n. 75 = pyrenaicus.
Kotschyi, Boiss. Diagn. Ser. I. xi. 40 = asper.
Kunthianus, Wall. Cat. n. 3292 = Saussurea Kunthiana.
lactucoides, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 151 = Microseris Forsteri.
laevigatus, Willd. Sp. Pl. iii. 1546 = Taraxacum officinale.
lanatus, Ledeb. in Mém. Acad. Sc. Pétersb. v. 1812 (1814), 554 = Taraxacum serotinum.
lanatus, Linn. Cent. Pl. i. 26; Amoen. Acad. iv. 287 = Scorzonera lanata.
leptocephalus, Koch, Syn. Fl. Germ. ed. I. 429, in syn. = Taraxacum officinale.
leucanthus, Ledeb. Fl. Alt. iv. 154 = Taraxacum bicolor.
libanoticus, *Boiss. Diagn.* Ser. I. xi. 39.—Syria.
Linkii, Wallr. in Linnaea, xiv. (1840) 662; Beitr. Fl. Hercyn. 292 = autumnalis.
liveus, All. Fl. Pedem. i. 210 = Taraxacum palustre.

LEONTODON:—
lividus, Waldst. & Kit. Pl. Rar. Hung. ii. 120 = Taraxacum palustre.
longifolius, Schur, Enum. Pl. Transs. 358 = hastilis.
lucidus, *DC. Prod.* vii. 104.—Italia.
lyratus, Ledeb. Fl. Alt. iv. 152 = Taraxacum lyratum.
macrorrhizus, *Steud. Nom.* ed. II. ii. 24.—Italia.
major, Mérat, in Ann. Sc. Nat. Sér. I. xxii. (1831) 106 = hispidus.
maroccanus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 544.—Hispan.; Marocc.
masmenaeus, *Boiss. & Bal. Diagn.* Ser. II. v. 117.—Cappadoc.
medius, Chaill. ex DC. Prod. vii. 146 = Taraxacum officinale.
megalorrhizus, *Forsk. Fl. Aegypt. Arab.* 216.—Byzant.
microcephalus, *Boiss. Voy. Bot. Espagne*, ii. 380. t. 115.—Hispan.
minimus, Brigant. ex Guss. Fl. Sic. Syn. ii. 397 = Taraxacum minimum.
montanus, Lam. Fl. Fr. iii. 640 = Taraxacum officinale.
montanus, C. A. Mey. Verz. Pfl. Cauc. 58 = Taraxacum montanum.
mucronatus, Forsk. Fl. Aegypt. Arab. 144 = Launaea chondrilloides.
muricatus, L. Hérit. Stirp. Nov. 178 = Picris asplenioides.
nevadensis, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 230.—Hispan.
nigricans, Kit. in Schult. Oestr. Fl. ed. II. ii. 405 = Taraxacum officinale.
nivalis, Hegetschw. Fl. Schw. 755 = L. Taraxaci.
nudicaulis, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28; Soland. ex Lowe, Man. Fl. Mad. i. 535 = Rothii.
nudicaulis, Mérat, in Ann. Sc. Nat. Sér. I. xxii. (1831) 109 = hirtus.
obliquus, Fries, Nov. Fl. Suec. ed. II. 243 = Taraxacum officinale.
obovatus, Willd. Sp. Pl. iii. 1546 = Taraxacum officinale.
officinalis, J. F. Gmel. Syst. 1174; With. Arr. Brit. Pl. ed. III. 679 = Taraxacum officinale.
oligocephalus, Schur, Enum. Pl. Transs. 357 = autumnalis?
oxylepis, *Boiss. & Heldr. Diagn.* Ser. I. xi. 40.—As. Min.
paludosus, Vitm. Summa Pl. iv. 399 = Taraxacum palustre.
palustris, Sm. Fl. Brit. ii. 823 = Taraxacum palustre.
parviflorus, Tausch, in Flora, xiv. (1831) 224 = Taraxacum palustre.
parvulus, Wall. Cat. n. 3247 = Taraxacum officinale.
Pavonii, *Boiss. Diagn.* Ser. II. iii. 87.—Hispan.
pectinatus, Boiss. Fl. Orient. iii. 733 = masmenaeus.
permiatus, Schur, Enum. Pl. Transs. 359 = hastilis.
pinnatifidus, *Ball, in Ann. & Mag. Nat. Hist.* Ser. II. vi. (1850) 17.—Italia.
pinnatifidus, Tausch, in Flora, vii. (1824) 248 = Taraxacum palustre.
Pollinii, Welw. ex Reichb. Fl. Germ. Excurs. 853 = pyrenaicus.
praecox, Kit. ex Reichb. l. c. 270 = Taraxacum officinale.
pratensis, Lam. Fl. Fr. ii. 115 = crispus.
pratensis, Reichb. Fl. Germ. Excurs. 253 = autumnalis.
Preslii, *DC. Prod.* vii. 105.—Sicilia.
proteiformis, Vill. Prosp. 34 = hastilis.
protheiformis, Balb. ex Pers. Syn. ii. 369 = praec.
Pseudocrispus, Sch. Bip. ex Nym. Consp. 468 = hastilis.
Pseudo-Taraxaci, Schur, Enum. Pl. Transs. 357 = pyrenaicus.
psilocalix, Mérat, in Ann. Sc. Nat. Sér. I. xxii. (1831) 106 = hirtus.
pycnocephalus, Ledeb. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 185 = Taraxacum officinale.
pyrenaicus, *Gouan, Illustr.* 55. t. 22.—Mont. Pyren.
radiatus, Lam. Fl. Fr. ii. 114 = Hyoseris radiata.
radicatus, Baill. Hist. Pl. viii. 294 = Hypochoeris radicata.
Raii, Gouan, Illustr. 55 = Taraxacum palustre.
recognitus, Monn. Ess. Hierac. 82 = hastilis.
repens, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 148 = Taraxacum officinale.
Rosani, DC. Prod. vii. 104 = hirtus.
Rothii, Ball, in Journ. Linn. Soc. xvi. (1878) 543 = saxatilis.

LEONTODON :—
runcinatus, Gilib. Fl. Lituan. i. 228 = Taraxacum officinale.
salinus, Ledeb. Fl. Alt. iv. 153 = Taraxacum collinum.
salinus, Poll. Hist. Pl. Palat. ii. 380 = Taraxacum palustre.
salsus, Link, Enum. Hort. Berol. ii. 283 = Taraxacum palustre.
Salzmanni, *Ball, in Journ. Linn. Soc.* xvi. (1878) 545.—Reg. Mediterr.
saxatilis, *Lam. Fl. Fr.* ii. 115.—Reg. Mediterr. occ.
saxatilis, Reichb. Fl. Germ. Excurs. 252 = crispus.
scaber, Mielich. in Flora, xxii. (1839) 492 = hispidus.
Scorzonera, Roth, Neue Beitr. i. 229 = Taraxacum palustre.
serotinus, Sibth. & Sm. Fl. Graec. viii. 71. t. 796 = Taraxacum officinale.
serotinus, Waldst. & Kit. Pl. Rar. Hung. ii. 119. t. 144 = Taraxacum serotinum.
setosus, Schur, Enum. Pl. Transs. 359 = hastilis.
siculus, Nym. Syll. 54 = hastilis.
sinensis, Lour. Fl. Cochinch. ii. 479 = Taraxacum officinale.
spatulifolius, Schloss. & Vuk. Syll. 59; et in Bull. Soc. Bot. Fr. xxix. (1882) Bibl. 6 = incanus.
squamosus, Lam. Encyc. iii. 529 = pyrenaicus.
Stevenii, Spreng. Syst. iii. 658 = Taraxacum officinale.
sublyratus, *Mérat, in Ann. Sc. Nat.* Sér. I. xxii. (1831) 108.—Europ.
Taraxaci, *Loisel. Fl. Gall.* ed. I. 513.—Europ.
taraxacifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = praec.
taraxacoides, Hoppe, in Sturm, Deutschl. Fl. Heft 41 = Taraxacum officinale.
taraxacoides, *Mérat, in Ann. Sc. Nat.* Sér. I. xxii. (1831) 108.—Gall.; Hispan.
Taraxacon, Neck. Delic. Gallo-Belg. ii. 329 = Taraxacum officinale.
Taraxacon, With. Arr. Brit. Pl. ed. III. 679 = Taraxacum palustre.
Taraxaconastrum, Vand. Fasc. Pl. 18 = seq.?
taraxaconoides, Vand. l. c. 17 = hirtus.
Taraxacum, Linn. Sp. Pl. 798 = Taraxacum officinale.
Tenorii, Nym. Consp. 468 = hastilis.
tenuiflorus, *DC. Prod.* vii. 102.—Helvet.
tenuiflorus, Reichb. Fl. Germ. Excurs. 853 = alpinus.
tenuifolius, Hoppe, ex Sturm, Deutschl. Fl. Heft 41 = Taraxacum tenuifolium.
terglouensis, Hacq. Pl. Carn. 11. t. 2. f. 5 = Crepis hyoseridifolia.
tingitanus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 544.—Hispan.; Marocc.
tomentosus, Linn. f. Suppl. 37 = Chaptalia albicans.
tomentosus, Vahl, ex Lange, in Kjoeb. Vidensk. Meddel. (1861) 101 = Taraxacum serotinum.
trivialis, Ball, in Journ. Bot. xi. (1873) 372 = hispidulus.
tuberosus, Georgi, Beschr. Russ. Reich. iv. 1210 (Quid?).—Reg. Caucas.
tuberosus, *Linn. Sp. Pl.* 799.—Europ. austr.; Oriens.
uliginosus, *Schult. ex Steud. Nom.* ed. I. 470.—Hab.?
umbellatus, Schrank, Baier. Fl. ii. 334 = Picris hieracioides.
validus, Peterm. Fl. Lips. Excurs. 578 = autumnalis?
variegatus, Poir. Encyc. Suppl. iii. 455; DC. Prod. vii. 105 = Hypochoeris variegata.
Villarsii, Loisel. Fl. Gall. ed. I. 514 = hirtus.
vulgaris, Kitt. Taschenb. ed. II. 534 = hispidus, hastilis, &c.
vulgaris, Lam. Fl. Fr. ii. 113 = Taraxacum officinale.

LEONTOGLOSSUM, Hance, in Walp. Ann. ii. 18 (1851) = **Tetracera**, Linn. (Dilleniac.).
scabrum, Hance, l. c. = T. sarmentosa.

LEONTONDON, Robin, Voy. Louis. iii. 425 (1807), sphalm. = **Leontodon**, Linn. (Compos.).

LEONTONIX, Heynh. Nom. i. 453 (1840) = seq.

LEONTONYX, Cass. in Dict. Sc. Nat. xxv. 466 (1822). *COMPOSITAE*, Benth. & Hook. f. ii. 312.
SPIRALEPIS, D. Don, in Mem. Wern. Soc. v. (1826) 551.
angustifolius, *DC. Prod.* vi. 168.—Afr. austr.
bicolor, *DC. l. c.*—Afr. austr.
candidissimus, *DC. l. c.* 169 = spathulatus.
coloratus, *Cass. in Dict. Sc. Nat.* xxv. 467.—Afr. austr.
declinatus, Heynh. Nom. i. 453; ii. 349 = Helichrysum declinatum.
glomeratus, DC. Prod. vi. 168 = coloratus.
modestus, Heynh. Nom. i. 453; ii. 349 = Helipterum gnaphalioides.
pusillus, Drège, ex Harv. & Sond. Fl. Cap. iii. 207 = spathulatus.
pusillus, *Less. Syn. Consp.* 327.—Afr. austr.
spathulatus, *Less. l. c.*—Afr. austr.
squarrosus, DC. Prod. vi. 167 = tomentosus.
stramineus, *DC. l. c.* 168 (sp. dub.).—Afr. austr.
tinctus, DC. l. c. = tomentosus.
tomentosus, *Cass. in Dict. Sc. Nat.* xxv. 466.—Afr. austr.

LEONTOPETALON, Tourn. ex Adans. Fam. ii. 433 (1763) = **Leontice**, Linn. (Berberid.).

LEONTOPHTHALMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140 = **Calea**, Linn. (Compos.).
peruvianum, H. B. & K. Nov. Gen. et Sp. iv. 296. t. 409 = C. Leontophthalmum.

LEONTOPODION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 85 = seq.

LEONTOPODIUM, R. Br. in Trans. Linn. Soc. xii. (1817) 124. *COMPOSITAE*, Benth. & Hook. f. ii. 302.
alpinum, *Cass. in Dict. Sc. Nat.* xxv. 474.—Europ.; Reg. Himal.
alpinum, Colm. ex Willk. & Lange, Prod. Fl. Hisp. ii. 64 = Evax exigua.
Andersoni, C. B. Clarke, Comp. Ind. 100 = alpinum.
catipes, F. Muell. in Papers Roy. Soc. Tasm. (1882) 44 = Raoulia catipes.
helveticum, D. Don, ex G. Don, in Loud. Hort. Brit. 340 = alpinum.
himalayanum, DC. Prod. vi. 276 = alpinum.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 178.—Japon.
javanicum, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind. ii. (1845) 263 = Gnaphalium japonicum.
linearifolium, *Benth. & Hook. f. Gen.* ii. 303.—Bolivia.
Meredithae, *F. Muell. Census*, 80.—Tasman.
monocephalum, Edgew. in Trans. Linn. Soc. xx. (1846) 73 = alpinum.
monoicum, *Benth. & Hook. f. Gen.* ii. 303.—N. Granat.
sibiricum, *Cass. in Dict. Sc. Nat.* xxv. 475.—As. bor. occ.
umbellatum, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 346 = alpinum.

LEONURA, Usteri, ex Steud. Nom. ed. II. ii. 25 (1841) = **Salvia**, Tourn. (Labiat.).
spaendochea, Usteri, ex Steud. l. c. = S. leonuroides.

LEONUROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = seq.

LEONURUS, Linn. Syst. ed. I (1735); Gen. ed. I. 176 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1210.
CARDIACA, Moench, Meth. 401 (1794).
CHAITURUS, Willd. Prod. Fl. Berl. 200 (1787).
PANZERIA, Moench, Meth. 402 (1794).
aconitifolius, Schlecht. ex Ledeb. Fl. Ross. iii. 423 = L. Cardiaca.
africanus, Mill. Gard. Dict. ed. VIII. n. 1 = Leonotis Leonurus.
alpinus, Schang. in Pall. N. Nord. Beitr. vi. 64 = Phlomis alpina.
altaicus, Spreng. Syst. ii. 738 = sibiricus.
altissimus, Bunge, ex Benth. Lab. Gen. et Sp. 521 = sibiricus.
campestris, Andrz. ex Benth. l. c. 518 = L. Cardiaca.
canescens, Dum. Fl. Belg. 46; Steud. Nom. ed. II. ii. 25 = L. Cardiaca.
capitatus, *Fresen. in Linnaea*, xv. (1841) *Litt.* 91.—Am. bor.
Cardiaca, *Linn. Sp. Pl.* 584.—Reg. bor. temp.
condensatus, Dum.-Cours. Bot. Cult. ed. II. iii. 67 = Phlomis condensata.
condensatus, Hort. ex Hornem. Hort. Hafn. ii. 557 = L. Cardiaca.
cordifolius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 (Quid?).—Java.

LEONURUS :—
crispus, Murr. in Nov. Comm. Gotting. viii. (1777) 44. t. 4 = L. Cardiaca.
discolor, Koch, in Linnaea, xxi. (1848) 680 = L. Cardiaca.
dschungaricus, *Regel, in Act. Hort. Petrop.* vi. (1879) 367.—Turkestan.
eriophorus, Turcz. ex Steud. Nom. ed. II. ii. 25 = seq.
eriostachys, Turcz. ex Benth. Lab. Gen. et Sp. 586 = Marrubium eriostachyum.
Galeobdolon, Scop. Fl. Carn. ed. II. i. 409 = Lamium Galeobdolon.
germanica, Gueldenst. Reisen, ii. 20 = Stachys germanica?
glabra, Gilib. Fl. Lithuan. ed. Usteri, Delect. ii. 321 (Quid?).—Europ.
glaucescens, *Bunge, in Ledeb. Fl. Alt.* ii. 409.—Sibir. altaic.
globosus, Moench, Meth. 400 = Leonotis nepetaefolia.
grandiflorus, Moench, l. c. = Leonotis Leonurus.
heterophyllus, Sweet, Hort. Brit. ed. I. 321; Brit. Flow. Gard. Ser. I. t. 197 = sibiricus.
illyricus, Hort. ex Benth. Lab. Gen. et Sp. 518 = L. Cardiaca.
indicus, Linn. Syst. ed. X. 1101; Burm. f. Fl. Ind. 127 = Leucas linifolia.
indicus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = javanicus?
intermedius, *Didr. Ind. Sem. Hort. Haun.* (1849) 14.—Ins. Tahiti.
japonicus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 112 = macranthus.
javanicus, *Blume, Bijdr.* 828.—Java.
javanicus, Span. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 339 = Gironniera nervosa.
lacerus, Lindl. in Trans. Hort. Soc. vi. (1826) 298 = L. Cardiaca.
lanatus, *Pers. Syn.* ii. 126; *Spreng. Syst.* ii. 738.—Sibir.; China.
macranthus, *Maxim. Prim. Fl. Amur.* 476.—Reg. Amur.
Marrubiastrum, Burm. f. Fl. Ind. 127 = Leucas zeylanica.
Marrubiastrum, *Linn. Sp. Pl.* 584.—Europ.; As. bor.
Marrubiastrum, Lour. Fl. Cochinch. 436 = Leonotis nepetaefolia.
marrubifolius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = oblongifolius.
marrubifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Marrubiastrum.
mollis, Ehrenb. ex Boiss. Fl. Orient. iv. 721 = Stachys Ehrenbergii.
multifidus, Desf. Tabl. ed. II. 270; Cat. Hort. Par. ed. III. 395 = sibiricus.
neglectus, Schrank, in Syll. Ratisb. ii. (1828) 61 = L. Cardiaca.
nepetaefolius, Mill. Gard. Dict. ed. VIII. n. 2 = Leonotis nepetaefolia.
oblongifolius, *Blume, Bijdr.* 828.—Java.
occidentalis, Colla, in Mem. Acc. Torin. xxxiii. (1829) 154 = sibiricus.
parviflorus, Moench, Meth. 401 = Leucas zeylanica.
parviflorus, Salisb. Prod. 84 = L. Marrubiastrum.
persicus, *Boiss. Diagn.* Ser. I. v. 27.—Persia.
pubescens, Benth. in Wall. Pl. As. Rar. i. 63 = L. Cardiaca.
quinquelobatus, Gilib. Fl. Lithuan. ed. Usteri, Delect. ii. 321 (Quid?).—Europ.
Royleanus, Benth. in Wall. Pl. As. Rar. i. 63 = L. Cardiaca.
ruderalis, Salisb. Prod. 84 = L. Cardiaca.
sibiricus, *Linn. Sp. Pl.* 584.—Sibir.; China.
sibiricus, Schang. in Pall. N. Nord. Beitr. vi. 97 = glaucescens.
superbus, Medic. Beobacht. (1783) 127 = Leonotis Leonurus.
supinus, Steph. ex Willd. Sp. Pl. iii. 116 = Marrubium incisum.
tataricus, Burm. f. Fl. Ind. 127 = sibiricus.
tataricus, *Linn. Sp. Pl.* 584.—Asia med.
tataricus, Tchihat. Asie Min. Bot. ii. 172 = L. Cardiaca.
trilobatus, Dulac, Fl. Hautes-Pyr. 409 = L. Cardiaca.
villosus, Desf. Tabl. ed. II. 73 = L. Cardiaca.

LEOPARDANTHUS, Blume, Rumphia, iv. 47 (1848); et Mus. Bot. Lugd. Bat. i. 47. t. 15 (1849) = **Dipodium**, R. Br. (Orchid.).
scandens, Blume, ll. cc. 47; 48. t. 15.—Malaya.

LEOPOLDIA, Herb. in Bot. Mag. sub t. 2113, in nota (1819) = **Hippeastrum**, Herb. (Amaryll.).
Belladonna, M. Roem. Syn. Ensat. 129 = Amaryllis Belladonna.
illustris, M. Roem. l. c. 130 = H. psittacinum.
Principis, M. Roem. l. c. 129 = H. reticulatum.
reticulata, Herb. in Bot. Mag. sub t. 2113, in nota = H. reticulatum.
striatifolia, Herb. in Trans. Hort. Soc. iv. (1822) 181 = H. reticulatum.

LEOPOLDIA, Parl. Fl. Palerm. i. 435 (1845); et in Giorn. Bot. Ital. ii. II. (1847) 57 = **Muscari**, Mill. (Liliac.).
abortiva, Jord. ex Nym. Consp. 732 = Hyacinthus trifoliatus.
Bouviana, Heldr. ex Nym. Consp. 733 (= *Bellevalia Bouviana*).—Graecia.
Calandriniana, Parl. Fl. Ital. ii. 496 = M. comosum.
Clusiana, Heldr. ex Nym. Consp. 732 = M. comosum.
comosa, Parl. in Giorn. Bot. Ital. ii. II. (1847) 160 = M. comosum.
constricta, Heldr. in Bull. Soc. Nat. Mosc. liii. (1878) I. 71 = M. constrictum.
Cupaniana, Parl. Fl. Ital. ii. 497 = M. Cupanianum.
curta, Heldr. in Bull. Soc. Nat. Mosc. liii. (1878) I. 65 = M. Holzmanni.
graeca, Heldr. l. c. 67 = M. graecum.
graminifolia, Heldr. ex Nym. Consp. 733 (= *Bellevalia graminifolia*).—Ins. Graec.
Gussonii, Parl. Fl. Ital. ii. 498 = M. Gussonii.
Holzmanni, Heldr. in Bull. Soc. Nat. Mosc. liii. (1878) I. 65 = M. Holzmanni.
maritima, Parl. in Giorn. Bot. Ital. ii. II. (1847) 162 = M. maritimum.
Neumayerii, Heldr. in Bull. Soc. Nat. Mosc. liii. (1878) I. 70 = M. Neumayeri.
pharmacusana, Heldr. l. c. 66 = M. pharmacusanum.
Pinardi, Parl. in Giorn. Bot. Ital. ii. II. (1847) 162 = M. Pinardi.
pyramidalis, Heldr. in Bull. Soc. Nat. Mosc. liii. (1878) I. 71 = M. pyramidale.
Sartoriana, Heldr. l. c. 66 = M. Sartorianum.
tenuiflora, Heldr. l. c. 70 = M. caucasicum.
theraea, Heldr. l. c. 69 = M. theraeum.
trojana, Heldr. l. c. 68 = M. trojanum.
Weissii, Freyn, in Oestr. Bot. Zeitschr. xxviii. (1878) 87 = M. Weissii.

LEOPOLDINIA, Mart. Hist. Nat. Palm. ii. 58 (1824). *PALMAE*, Benth. & Hook. f. iii. 920.
insignis, *Mart. l. c.* 60. t. 53. f. 16, 17.—Bras.
major, *Wallace, Palm-trees, Amaz.* 15. t. 5.—Bras.
Piassaba, *Wallace, ex Archer, in Hook. Kew Journ.* vii. (1855) 213.—Am. trop.
pulchra, Hort. Linden, ex Seem. in Gard. Chron. (1870) 495 = Cocos Weddelliana.
pulchra, *Mart. Hist. Nat. Palm.* ii. 59. tt. 52, 53.—Bras.

LEPACHIS, Rafin. in Am. Monthly Mag. (1819) 195 = **Echinacea**, Moench (Compos.).
purpurea, Rafin. l. c. = E. purpurea.

LEPACHYS, Rafin. in Journ. Phys. lxxxix. (1819) 100. *COMPOSITAE* [Benth. & Hook. f. ii. 366].
OBELISCARIA, Cass. in Dict. Sc. Nat. xxxv. 272 (1825).
RATIBIDA, Rafin. in Am. Monthly Mag. (1818) 268.
angustifolia, Torr. & Gray, Fl. N. Am. ii. 314, in syn. = pinnatifida.
columnaris, *Torr. & Gray, l. c.* 315.—Am. bor. occ.
peduncularis, *Torr. & Gray, l. c.*—Texas.
pinnata, Torr. & Gray, l. c. 314 = pinnatifida.
pinnatifida, *Rafin. in Journ. Phys.* lxxxix. (1819) 100.—Am. bor.
serratus, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457 = peduncularis.
Tagetes, *A. Gray, in Pacif. Rail. Rep.* iv. 103.—Am. bor. occ.

LEPADENA, Rafin. Fl. Tellur. iv. 113 (1836) = **Euphorbia**, Linn.
leucoloma, Rafin. l. c. 114 = E. marginata.

LEPANTES, Sw. in Schrad. Journ. ii. (1799) 249 = seq.

LEPANTHES, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 85. *ORCHIDEAE*, Benth. & Hook. f. iii. 491.
andrenoglossa, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Ecuador.
Aquila-Borussiae, *Reichb. f. l. c.* ii. (1854) 22.—Venezuela.
armeniaca, *Rodrig. Orch. Nov.* ii. 58.—Bras.
Avis, *Reichb. f. Xenia Orch.* i. 144. t. 50; ii. 5–7.—Mexic.
bicarinata, *Rodrig. Orch. Nov.* ii. 54.—Bras.
bicolor, *Rodrig. l. c.* 49; *et in Rev. de Engenh.* iii. (1881) 110. n. 7.—Bras.
bicornuta, *Rodrig. l. c.* 61.—Bras.
biloba, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 356.—Peruv.
blepharistes, *Reichb. f. Beitr. Orch. Centr.-Am.* 92.—Costa Rica.
calodictyon, *Hook. Bot. Mag.* t. 5259.—Ecuador.
capitana, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Peruv.
carinifera, *Rodrig. in Rev. de Engenh.* iii. (1881) 110. n. 8.—Bras.
cassidea, *Reichb. f. Xenia Orch.* i. 146.—Peruv.
chrysosepala, *Rodrig. Orch. Nov.* ii. 44.—Bras.
chrysostigma, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 329.—Cuba.
ciliata, Lindl. ex Griseb. Cat. Pl. Cub. 260 = Pleurothallis blepharophylla.
cochlearifolia, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 86.—Jamaic.
concinna, *Sw. l. c.* 85.—Jamaic.
crassicaulis, *Rodrig. Orch. Nov.* ii. 60.—Bras.
crassifolia, *Reichb. f. in Linnaea*, xli. (1877) 47.—Jamaic.?
crebifolia, *Rodrig. Orch. Nov.* ii. 67.—Bras.
crinita, *Rodrig. l. c.* 66.—Bras.
cristata, *Rodrig. l. c.* 57.—Bras.
cryptophyta, *Rodrig. l. c.* 65.—Bras.
cyanoptera, *Reichb. f. in Linnaea*, xxii. (1849) 819.—Venezuela.
dorsalis, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 329.—Cuba.
elata, *Reichb. f. Beitr. Orch. Centr.-Am.* 90.—Costa Rica.
erinacea, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Costa Rica.
Ferdinandiana, *Rodrig. Orch. Nov.* ii. 64.—Bras.
flaminea, *Rodrig. in Rev. de Engenh.* iii. (1881) 109. n. 4.—Bras.
fulva, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 329.—Cuba.
Gargantua, *Reichb. f. Xenia Orch.* i. 150.—Peruv.
granulosa, *Rodrig. Orch. Nov.* ii. 45.—Bras.
Gunningiana, *Rodrig. l. c.* 48.—Bras.
helicocephala, *Reichb. f. Xenia Orch.* i. 150.—Guian.
horrida, *Reichb. f. in Bonplandia*, iii. (1855) 70.—Costa Rica.
imbricata, *Rodrig. Orch. Nov.* ii. 56.—Bras.
ionoptera, *Reichb. f. in Bonplandia*, iii. (1855) 70.—N. Granat.
Jamiesoni, *Lindl. ex Reichb. f. Xenia Orch.* i. 152.—Ecuador.
laciniata, *Rodrig. Orch. Nov.* ii. 68.—Bras.
Lindleyana, *Oerst. ex Reichb. f. in Bonplandia*, iii. (1855) 225; *Xenia Orch.* i. 149. t. 50.—Costa Rica.
lineolata, *Rodrig. Orch. Nov.* ii. 53.—Bras.
lobisserrata, *Rodrig. l. c.* 63.—Bras.
Loddigesiana, *Lindl. Bot. Reg.* t. 1762; *Reichb. f. Xenia Orch.* i. 145.—Jamaic.
mandibularis, *Reichb. f. in Flora*, xlviii. (1865) 275.—Ind. occ.
marginata, *Rodrig. Orch. Nov.* ii. 68.—Bras.
marmorata, *Rodrig. l. c.* 42.—Bras.
mesochlora, *Reichb. f. Xenia Orch.* i. 147.—Peruv.
microphyta, *Rodrig. Orch. Nov.* ii. 67.—Bras.
modesta, *Rodrig. l. c.* 62.—Bras.
monoptera, *Lindl. in Hook. Journ. Bot.* i. (1834) 10.—Peruv.
montana, *Rodrig. Orch. Nov.* i. 22, nomen.—Bras.
mucronata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 356.—Peruv.
muscicola, *Rodrig. Orch. Nov.* ii. 56.—Bras.
muscosa, *Rodrig. l. c.* 43.—Bras.
nanegalensis, *Reichb. f. in Linnaea*, xli. (1877) 46.—Ecuador.
nemorosa, *Rodrig. Orch. Nov.* ii. 54.—Bras.

LEPANTHES:—
nubicola, *Reichb. f. Xenia Orch.* i. 149.—N. Granat.
nummularia, *Reichb. f. l. c.* 142.—Peruv.
orbiculata, *Lindl. ex Reichb. f. l. c.* 152.—Hab.?
otostalix, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Ecuador.
Paivaeana, *Reichb. f. Xenia Orch.* iii. 26.—Chili.
papyrophylla, *Reichb. f. l. c.* i. 151.—Ecuador.
parahybunensis, *Rodrig. Orch. Nov.* ii. 57.—Bras.
pellifeloides, *Rodrig. in Rev. de Engenh.* iii. (1881) 110 n. 5.—Bras.
picnostachia, *Rodrig. Orch. Nov.* ii. *Ind.* p. vii, nomen (Quid?).—Bras.
pristidis, *Reichb. f. in Linnaea*, xxii. (1849) 820.—Mexic.
pteropogon, *Reichb. f. Xenia Orch.* i. 146.—Peruv.
pulchella, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 86.—Jamaic.
punctata, *Rodrig. Orch. Nov.* ii. 52.—Bras.
punctatifolia, *Rodrig. l. c.* 55.—Bras.
pusilla, *Rodrig. in Rev. de Engenh.* iii. (1881) 110. n. 6.—Bras.
quadridentata, *Rodrig. Orch. Nov.* ii. 50.—Bras.
ramphastorhyncha, *Rodrig. l. c.* 55.—Bras.
recurvipetala, *Rodrig. l. c.* 62.—Bras.
renipetala, *Rodrig. l. c.* 66.—Bras.
roseola, *Reichb. f. in Linnaea*, xli. (1877) 46.—Ecuador.
rotundata, *Griseb. Fl. Brit. W. Ind.* 610.—Ind. occ.
ruscifolia, *Reichb. f. in Linnaea*, xxii. (1849) 819.—Venezuela.
sanguinea, *Hook. Bot. Mag.* t. 4112.—Jamaic.
sarcopetala, *Rodrig. Orch. Nov.* ii. 46.—Bras.
Schiedei, *Reichb. f. in Linnaea*, xxii. (1849) 820.—Mexic.
secunda, *Rodrig. Orch. Nov.* ii. 70.—Bras.
selenitepala, *Reichb. f. in Ber. Deutsch. Bot. Ges.* iii. (1885) 279.—Ind. occ.
striata, *Rodrig. Orch. Nov.* ii. 59.—Bras.
tabacina, *Rodrig. l. c.* 47.—Bras.
tenera, *Rodrig. l. c.* 51.—Bras.
tipulifera, *Reichb. f. Beitr. Orch. Centr.-Am.* 91. t. 10.—Costa Rica.
tricarinata, *Rodrig. Orch. Nov.* ii. 43.—Bras.
trichodactyla, *Lindl. in Mem. Acad. Am.* N. S. viii. (1860) 220.—Cuba.
tricolor, *Rodrig. Orch. Nov.* ii. 52.—Bras.
tridentata, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 86.—Jamaic.
trilineata, *Rodrig. Orch. Nov.* ii. 42.—Bras.
Turialvae, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Costa Rica.
umbrosa, *Rodrig. Orch. Nov.* ii. 60.—Bras.
urotepala, *Reichb. f. Xenia Orch.* i. 151.—Hab.?
variegata, *Rodrig. Orch. Nov.* ii. 61.—Bras.
Vespertilio, *Reichb. f. Xenia Orch.* i. 143.—Peruv.
viridula, *Rodrig. Orch. Nov.* ii. 51.—Bras.
Wageneri, *Reichb. in Bonplandia*, iii. (1855) 70.—Mexic.
Wawraeana, *Rodrig. Orch. Nov.* ii. 46.—Bras.
Wendlandi, *Reichb. f. Beitr. Orch. Centr.-Am.* 91. t. 9.—Costa Rica.
Wrightii, *Reichb. f. in Flora*, xlviii. (1865) 275.—Cuba.

LEPANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

LEPARGYRAEA, Steud. Nom. ed. II. ii. 25 (1841) = seq.

LEPARGYREA, Rafin. in Am. Monthly Mag. (1818) 176 = **Elaeagnus**, Linn.

LEPECHINIA, Willd. Hort. Berol. i. 20. t. 21 (1806). *LABIATAE*, Benth. & Hook. f. ii. 1192.
ULRICIA, Jacq. ex Steud. Nom. ed. I. 862 (1821).
clinopodiifolia, Willd. Hort. Berol. sub t. 21 = Mentha dahurica.
procumbens, *Benth. Lab. Gen. et Sp.* 415.—Mexic.
Schiedeana, Vatke, in Verh. Bot. Ver. Brandenb. (1875) 36 = procumbens.
spicata, *Willd. Hort. Berol.* i. 21. t. 21.—Mexic.

LEPEOCERCIS, Trin. Fund. Agrost. 203 (1820) = **Andropogon**, Linn. (Gramin.).
annulata, Nees, Fl. Afr. Austr. 98 = A. annulatus.
Bladhii, Nees, ex Steud. Syn. Pl. Gram. 380 = A. annulatus.
digitata, Nees, Fl. Afr. Austr. 99 = Trachypogon dactyloides?

LEPEOCERCIS:—
mollissima, Nees, ex Baker, Fl. Maurit. 444 = Andropogon caricosus.
pertusa, Hassk. Pl. Jav. Rar. 52 = A. pertusus.
serrata, Trin. Fund. Agrost. 203. t. 18 = A. caricosus.

LEPEOSTEGERES, Blume, in Schult. f. Syst. vii. 1731 (1830) = **Loranthus**, Linn.
bahajensis, Miq. Fl. Ind. Bat. i. II. 833 = Lor. bahajensis.
gemmiflorus, Blume, in Schult. f. Syst. vii. 1731 = Lor. gemmiflorus.
sankumbangensis, Miq. Fl. Ind. Bat. i. I. 834.—Borneo.

LEPERIZA, Herb. App. 41 (1821) = **Urceolina**, Reichb. (Amaryll.).
eucrosioides, Baker, in Gard. Chron. (1878) II. 170.—Am. trop.
latifolia, Herb. App. 41 = U. latifolia.

LEPIA, Desv. Journ. Bot. iii. (1814) 166 = **Lepidium**, Linn. (Crucif.).
accedens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 338.—Gallia.
Bonanniana, Presl, Fl. Sic. 84 = Lepidium hirtum.
campestris, Desv. Journ. Bot. iii. (1814) 165 = Lepidium campestre.
campicola, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 338 = Lepidium campestre.
errabunda, Fourr. l. c. = Lepidium campestre.
glastifolia, Desv. Journ. Bot. iii. (1814) 166 = Lepidium glastifolium.
hirta, Desv. l. c. = Lepidium hirtum.
latifolia, Desv. l. c. = Thlaspi latifolium.
linifolia, Desv. l. c. = Lepidium linifolium.
marginata, Desv. l. c. = Aethionema saxatile.
membranacea, Desv. l. c. 166, 181 = Aethionema membranaceum.
oppositifolia, Desv. l. c. 166 = Aethionema oppositifolium.
rotunda, Desv. l. c. = Lepidium rotundum.
sativa, Desv. l. c. 165 = Lepidium sativum.
stylata, Webb, Iter Hisp. 76 = Lepidium nebrodense?

LEPIA, Hill, Veg. Syst. ii. 97 (1761) = **Zinnia**, Linn. (Compos.).
multiflora, Hill, Hort. Kew. 18 = Z. pauciflora.
pauciflora, Hill, l. c. = Z. pauciflora.

LEPIACTIS, Rafin. Fl. Tellur. ii. 43 (1836) = **Solidago**, Vaill. (Compos.).
virgata, Rafin. l. c. = S. stricta.

LEPIACTIS, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentibul.).
stellaris, Rafin. l. c. = U. stellaris.

LEPIANTHES, Rafin. Sylva Tellur. 84 (1838) = **Piper**, Linn.
granulatum, Rafin. l. c. 85 = P. angustifolium.
maculosum, Rafin. l. c. = Peperomia maculosa.
peltatum, Rafin. l. c. = P. peltatum.
umbellatum, Rafin. l. c. = P. umbellatum.

LEPICAUNE, Lapeyr. Hist. Abr. Pl. Pyr. 478 (1813) = **Crepis**, Linn. (Compos.).
albida, Lapeyr. ex DC. Prod. vii. 152 = C. albida.
balsamea, Lapeyr. Hist. Abr. Pl. Pyr. 478 = Hieracium amplexicaule.
blattarioides, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 104 = C. blattarioides.
ciliata, C. Koch, in Linnaea, xxiii. (1850) 683 = C. ciliata.
dshimilensis, C. Koch, l. c. = C. djimilensis.
grandiflora, Lapeyr. Hist. Abr. Pl. Pyr. 479 = C. grandiflora.
intybacea, Lapeyr. l. c. = C. grandiflora.
multicaulis, Lapeyr. l. c. 480 = C. blattarioides.
prunellaefolium, Lapeyr. ex DC. Prod. vii. 169 = C. pygmaea.
sibirica, C. Koch, in Linnaea, xxiii. (1850) 683 = C. sibirica.
spinulosa, Lapeyr. Hist. Abr. Pl. Pyr. 480 = Sonchus oleraceus.
tomentosa, Lapeyr. l. c. 481 = Senecio Doronicum.
turbinata, Lapeyr. l. c. 480, 481 = C. blattarioides.

LEPICEPHALUS, Lag. Gen. et Sp. Nov. 7 (1816) = **Cephalaria**, Schrad. (Dipsac.).
alpinus, Lag. l. c. 8 = C. alpina.
atratus, Lag. l. c. 7 = C. tatarica.

LEPICEPHALUS:—
attenuatus, Lag. Gen. et Sp. Nov. 8 = Cephalaria attenuata.
centaurioides, Lag. l. c. 7 = C. corniculata.
ciliata, Lag. l. c. 8 = Scabiosa ciliata.
coriaceus, Lag. l. c. = C. corniculata.
corniculatus, Lag. l. c. = C. corniculata.
cretaceus, Lag. ex Eichw. Skizze, 152 = C. corniculata.
decurrens, Lag. Gen. et Sp. Nov. 8 = C. attenuata.
humilis, Lag. l. c. = C. attenuata.
inflexus, Eichw. Skizze, 151 = Scabiosa australis.
inflexus, Steud. Nom. ed. II. ii. 26 = Scabiosa Succisa.
leucanthemus, Lag. Gen. et Sp. Nov. 7 = C. corniculata.
leucanthus, Lag. l. c. = C. leucantha.
rigidus, Lag. l. c. 8 = C. rigida.
scabra, Lag. l. c. = C. scabra.
Succisa, Eichw. Skizze, 151 = Scabiosa Succisa.
syriacus, Lag. Gen. et Sp. Nov. 8 = C. syriaca.
tartarica, Lag. l. c. = C. tatarica.
transsylvanicus, Lag. l. c. = C. transsylvanica.
uralensis, Lag. l. c. = C. corniculata.
ustulatus, Lag. l. c. = C. ustulata.

LEPICLINE, Less. Syn. Comp. 296. (1832) = LEPISCLINE, Cass. = **Helichrysum**, Vaill. (Compos.).

LEPIDACANTHUS, Presl, Bot. Bemerk. 96 (1844); in Abh. Boehm. Ges. V. iii. (1845) 526. *ACANTHACEAE*.
comosus, Presl, ll. cc.—Bras.
latibracteatus, Presl, ll. cc.—Bras.

LEPIDADENIA, Nees, in Edinb. N. Phil. Journ. xv. (Apr.–Oct. 1833) 379; Laurin. Expos. 19 (1833) = **Litsea**, Lam. (Laurin.).
costata, Miq. Fl. Ind. Bat. i. I. 938.—Arch. Sunda.
cuneata, Miq. l. c. 936.—Sumatra.
diversifolia, Miq. l. c. = Litsea diversifolia.
ferruginea, Miq. l. c. 935 = Litsea ferruginea.
fusca, Miq. l. c. 937.—Sumatra.
fuscescens, Miq. l. c. 939.—Ins. Banca.
glabrata, Wight, Ic. t. 1838.—Ind. or.
Griffithii, Wight, l. c. t. 1846 (= *Cylicodaphne myristicaefolia*).—Ind. or.
grisea, Miq. Fl. Ind. Bat. i. I. 939.—Sumatra.
hirta, Miq. l. c. 940.—Java.
lucida, Miq. l. c. 937.—Sumatra.
magnifica, Miq. l. c. 936.—Sumatra.
mollis, Miq. l. c. 940.—Java?
Neesiana, Wight, Ic. t. 1840 (= *Cylicodaphne Neesiana*).—Malacca.
Noronhiana, Miq. Fl. Ind. Bat. i. I. 934 = Litsea Noronhae.
ochracea, Miq. l. c. 938.—Sumatra.
ovalifolia, Wight, Ic. t. 1839 = Litsea ovalifolia.
pallida, Miq. Fl. Ind. Bat. i. I. 935.—Ins. Molucc.
polyneura, Miq. l. c. 938.—Java?
punctata, Miq. l. c. 939.—Borneo.
robusta, Miq. l. c. 934 = Litsea robusta.
rubiginosa, Miq. l. c. 938.—Sumatra.
rubra, Miq. l. c. 934 = Litsea rubra.
Seloany, Miq. l. c. Suppl. 361.—Sumatra.
sumatrana, Miq. l. c. i. I. 937.—Sumatra.
tephrophylla, Miq. l. c. 940.—Java.
tuberculata, Miq. l. c. 937.—Sumatra.
velutina, Miq. l. c. 935.—Java.
Wightiana, Nees, in Edinb. N. Phil. Journ. xv. (Apr.–Oct. 1833) 379 = Litsea Sebifera.

LEPIDAGATHIS, Willd. Sp. Pl. iii. 400 (1800). *ACANTHACEAE*, Benth. & Hook. f. ii. 1101.
APOLEPSIS, Hassk. Cat. Hort. Bog. Alt. 150 (1844).
RUSSEGERA, Endl. Nov. Stirp. Dec. 38 (1839).
TELIOSTACHYA, Nees, in Mart. Fl. Bras. ix. 71, t. 8. (1847).
alopecuroides, *R. Br. ex Griseb. Fl. Brit. W. Ind.* 453.—Am. trop.
anobrya, *Nees, in DC. Prod.* xi. 255.—Afr. trop.
aristata, Nees, in Wall. Pl. As. Rar. iii. 95 = hyalina.
attenuata, Zipp. ex Span. in Linnaea, xv. (1841) 329 = humifusa.
axillaris, Wight, ex Nees, in DC. Prod. xi. 259 = trinervis.
Beddomei, T. Anders. in Journ. Linn. Soc. ix. (1867) 499 = Hamiltoniana.
Billardieriana, *Nees, in DC. Prod.* xi. 253.—Java.
calycina, *Hochst. ex DC. l. c.* 252.—Scind.; Abyssin.
ceylainca, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* ii. (1841) p. li.—Zeylan.
chlorostachya, *Nees, in DC. Prod,* xi. 256.—Burma.

LEPIDAGATHIS:—
clavata, *Dalz. in Hook. Kew Journ.* ii. (1850) 340.—Ind. or.
cristata, *Willd. Sp. Pl.* iii. 400.—Ind. or.
cuspidata, *Nees, in Wall. Pl. As. Rar.* iii. 97.—Ind. or.
diffusa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 518.—Ind. or.
discolor, Stocks, ex Boiss. Fl. Orient. iv. 525 = calycina.
dulcis, *Nees, in Wall. Pl. As. Rar.* iii. 96.—Burma.
eucephala, *Miq. Fl. Ind. Bat.* ii. 814.—Malaya.
falcata, *Nees, in Wall. Pl. As. Rar.* iii. 96.—Burma.
fasciculata, *Nees, l. c.* 95.—Ind. or.
glandulosa, *Nees, in DC. Prod.* xi. 243.—Abyssin.
goënsis, Dalz. in Hook. Kew Journ. ii. (1850) 340 = fasciculata.
grandiflora, Dalz. l. c. 138 = Calacantha Dalzellianus.
Hamiltoniana, *Wall. Pl. As. Rar.* iii. 96; *Cat.* n. 2422.—Ind. or.
Heudelotiana, *Nees, in DC. Prod.* xi. 254.—Afr. trop.
hirta, Nees, in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) p. liii = fasciculata.
humifusa, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 386.—Ins. Timor.
hyalina, *Nees, in Wall. Pl. As. Rar.* iii. 95.—As. trop.
hyssopifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 34.—Afr. trop.
incana, *Nees, in DC. Prod.* xi. 259.—Java.
incurva, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 119 = hyalina.
iridescens, T. Anders. in Journ. Linn. Soc. ix. (1867) 496 = hyalina.
javanica, *Blume, Bijdr.* 801.—Java.
javanica, Hort. Bogor. ex Miq. Fl. Ind. Bat. ii. 812 = straminea.
laciniata, Wall. ex Nees, in DC. Prod. xi. 227, in syn. = Barleria cristata.
laguroidea, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 34.—Afr. trop.
lanata, Wall. ex Nees, in DC. Prod. xi. 252, in syn. = scariosa.
laxa, *Nees, l. c.* 251.—Ins. Philipp.
linearis, *T. Anders. Cat. Hort. Calc.* 43, nomen; *et in Journ. Linn. Soc.* ix. (1867) 496.—Burma.
longifolia, *Wight, Ic.* iv. t. 1564.—Ind. or.
lutea, *Dalz. in Hook. Kew Journ.* ii. (1850) 139.—Ind. or.
luzona, *Nees, in DC. Prod.* xi. 259.—Ins. Philipp.
Medusae, *S. Moore, in Journ. Bot.* xviii. (1880) 39.—Afr. trop.
mitis, *Dalz. in Hook. Kew Journ.* iii. (1851) 226.—Ind. or.
mollis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 33.—Afr. trop.
mucronata, Nees, in Wall. Pl. As. Rar. iii. 95 = hyalina.
myrtifolia, *S. Moore, in Journ. Bot.* xviii. (1880) 38.—Afr. trop.
Neesiana, Wight, ex T. Anders. in Journ. Linn. Soc. ix. (1867) 499 = Neuracanthus Neesianus.
nervata, Wight, Ic. iv. t. 1620 = ceylanica.
neurophylla, T. Anders. in Journ. Linn. Soc. ix. (1867) 497 = hyalina.
pallescens, *S. Moore, in Journ. Bot.* xviii. (1880) 308.—Afr. trop.
parviflora, *Blume, Bijdr.* 801.—Java.
parviflora, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 387 = javanica.
Pavala, *Nees, in DC. Prod.* xi. 160.—Ind. or.
peniculifera, *S. Moore, in Journ. Bot.* xviii. (1880) 39.—Afr. trop.
prostrata, *Dalz. in Hook. Kew Journ.* ii. (1850) 138.—Ind. or.
psilantha, *Nees, in DC. Prod.* xi. 251.—Ins. Philipp.
pungens, *Nees, in Wall. Pl. As. Rar.* iii. 97.—Ind. or.
purpuricaulis, *Nees, l. c.* 96.—Reg. Himal.; Burma.
radicalis, *Hochst. ex Nees, in DC. Prod.* xi. 255.—Abyssin.
repanda, *Blume, Bijdr.* 802.—Java.
repens, *Spreng. Syst.* ii. 827.—Cochinch.
Riedeliana, *Nees, in Mart. Fl. Bras.* ix. 71.—Bras.
rigida, *Dalz. in Hook. Kew Journ.* ii. (1850) 340.—Ind. or.
riparia, Nees, in DC. Prod. xi. 253 = hyalina.
rupestris, Bedd. Ic. Pl. Ind. Or. t. 229 = mitis.
rupestris, Nees, in Wall. Pl. As. Rar. iii. 96 = cristata.
scariosa, *Nees, l. c.* 95.—Ind. or.; Abyssin.
secunda, *Nees, in DC. Prod.* xi. 259.—Ins. Philipp.
semiherbacea, Nees, in Wall. Pl. As. Rar. iii. 96 = hyalina.

LEPIDAGATHIS:—
setigera, *Blume, Bijdr.* 802.—Java.
Shuteri, T. Anders. in Journ. Linn. Soc. ix. (1867) 494 = cristata.
simplex, *T. Anders. l. c.* 497.—Burma.
sphaerostachya, Nees, in DC. Prod. xi. 254 = Neuracanthus sphaerostachyus.
spinosa, *Wight, ex Nees, in Wall. Pl. As. Rar.* iii. 97.—Ind. or.
straminea, *Miq. Fl. Ind. Bat.* ii. 812.—Java?
striata, Nees, in Wall. Pl. As. Rar. iii. 96 = purpuricaulis.
strobilifera, Stocks, in Hook. Kew Journ. iv. (1852) 177 = calycina.
strobilina, *T. Anders. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 96.—Burma.
subuninervia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 519.—Burma.
terminalis, Hochst. ex Nees, in DC. Prod. xi. 251 = scariosa.
timorensis, Steud. Nom. ed. II. ii. 26 = javanica.
trinervis, *Nees, in Wall. Pl. As. Rar.* iii. 96.—Ind. or.
ustulata, Nees, l. c. 95 = hyalina.
Walkeriana, *Nees, in DC. Prod.* xi. 260.—Zeylan.

LEPIDAMPHORA, Zoll. ex Miq. Fl. Ind. Bat. i. I. 217 (1855) = **Dioclea**, H. B. & K. (Legumin.).
volubilis, Zoll. l. c. = D. reflexa.

LEPIDANCHE, Engelm. in Am. Journ. Sc. xliii. (1842) 343. t. 6 = **Cuscuta**, Linn. (Convolv.).
adpressa, Engelm. in Hook. Lond. Journ. Bot. ii. (1842) 194, 198 = C. compacta.
compositarum, Engelm. in Am. Journ. Sc. xliii. (1842) 344. t. 6. f. 30–35 = C. glomerata.

LEPIDANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 64 (1861) = **Dissotis**, Benth. (Melastom.).
triplinervium, Klotzsch, l. c. = D. prostrata.

LEPIDANTHUS, Nees, in Linnaea, v. (1830) 665 = **Hypodiscus**, Nees (Restiac.).
Willdenowia, Nees, l. c. 665 = H. Willdenovia.

LEPIDANTHUS, Nutt. in Trans. Am. Phil. Soc. v. (1837) 175 = **Andrachne**, Linn. (Euphorb.).
phyllanthoides, Nutt. l. c. = A. Roemeriana.

LEPIDANTHUS, Nutt. in Trans. Am. Phil. Soc. N. S vii. (1841) 396 = **Matricaria**, Tourn. (Compos.).
suaveolens, Nutt. l. c. 397 = M. discoidea.

LEPIDAPLOA, Cass. in Bull. Soc. Philom. (1817) 66; et in Dict. Sc. Nat. xxvi. 16 (1823) = **Vernonia**, Schreb. (Compos.).
albicaulis, Cass. in Dict. Sc. Nat. xxvi. 17 = V. longifolia.
aristata, Cass. l. c. = V. aristata.
buxifolia, Cass. l. c. 18 = V. buxifolia.
canescens, Cass. l. c. = V. canescens.
lanceolata, Cass. l. c. = V. longifolia.
phyllostachya, Cass. l. c. 16 = V. arborescens.
scorpioides, Cass. l. c. = V. scorpioides.

LEPIDEILEMA, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 93 = **Streptochaeta**, Schrad. (Gramin.).
lancifolium, Trin. l. c. = S. spicata.

LEPIDEREMA, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 498. *SAPINDACEAE.*
papuana, *Radlk. l. c.* 535.—N. Guin.

LEPIDIBERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S xvi. (1868) 338 = **Lepidium**, Linn. (Crucif.).
graminifolia, Fourr. l. c. = Lepidium graminifolium.
polyclada, Fourr. l. c. = Lepidium graminifolium.

LEPIDION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = seq.

LEPIDIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 192 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 87.
CARDAMON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 338.
CARDARIA, Desv. Journ. Bot. iii. (1814) 163.
CYNOCARDAMUM, Webb & Berth. Phyt. Canar. i. 96 (1836).
DILEPTIUM, Rafin. Fl. Ludov. 85 (1817).
DISCOVIUM, Rafin. in Journ. Phys. lxxxix. (1819) 96.
IBERIS, Adans. Fam. ii. 422 (1763).
JUNDZILLIA, Andrz. ex DC. Syst. ii. 529 (1821).

LEPIDIUM:—
KANDIS, Adans. Fam. ii. 422 (1763).
LASIOPTERA, Andrz. ex DC. Syst. ii. 534 (1821).
LEPIA, Desv. Journ. Bot. iii. (1814) 166.
LEPIDIBERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 338.
MONOPLOCA, Bunge, in Lehm. Pl. Preiss. i. 259 (1845).
NASTURTIOIDES, Medic. Pflanzeng. i. 81 (1792).
PHYSOLEPIDIUM, Schrenk, Enum. Pl. 97 (1841).
SENCKENBERGIA, Gaertn. Mey. & Schreb. Fl. Wett. ii. 413 (1800).
THLASPIDIUM, Spach, Hist. Vég. Phan. vi. 557 (1838).
abrotanifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 308.—Ecuador.
abyssinicum, *Hochst. in A. Rich. Tent. Fl. Abyss.* i. 21.—Abyss.
affghanum, *Boiss. Fl. Orient.* i. 358.—Afghanist.
affine, Ledeb. Ind. Sem. Hort. Dorpat. (1821) 22; et in Link, Enum. Hort. Berol. ii. 152 = latifolium.
affine, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 284.—Bolivia.
africanum, DC. Syst. ii. 552 = ruderale.
alpigenum, *A. Rich. Tent. Fl. Abyss.* i. 22.—Abyss.
alpinum, Linn. Cent. Pl. ii. 23; Amoen. Acad. iv. 321 = Hutchinsia alpina.
alpinum, O.F. Muell. in Fl. Dan. t. 569 = groenlandicum.
alpinum, Sievers, in Pall. N. Nord. Beitr. vii. 350 = Smelowskia calycina.
alyssoides, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 10.—N. Mexic.
ambiguum, Lange, in Kjoeb. Vidensk. Meddel. (1865) 74 = L. Cardamines.
ambiguum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 34 = ruderale.
americanum, Vell. Fl. Flum. 271; vi. t. 108 = Senebiera pinnatifida.
amplexicaule, Stev. ex DC. Syst. ii. 554 = cordatum.
amplexicaule, *Willd. Sp. Pl.* iii. 436.—Sibir.
anglicum, Huds. Fl. Angl. ed. II. i. 280 = Senebiera pinnatifida.
angulosum, Urv. in Mém. Soc. Linn. Par. i. (1822) 334 = incisum.
apetalos, Gilib. Fl. Lituan. ii. 54 = ruderale.
apetalum, Willd. Sp. Pl. iii. 439 = incisum.
arcuatum, DC. in Mém. Soc. Hist. Nat. Par. (1799) 145 = L. Iberis.
armoracium, *Fisch. & Mey Ind. Sem. Hort. Petrop.* ix. 77.—Abyss.
arvense, Mill. Gard. Dict. ed. VIII. n. 2 = L. Draba.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 195.—Persia; Afghan.
auriculatum, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1857) 51.—Hab.?
auritum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 307.—Ecuador.
australe, *T. Kirk, in Trans. N. Z. Inst.* xiv. (1882) 381.—N. Zel.
Berteronianum, *Steud. Nom.* ed. II. ii. 26.—Chili.
bidentatum, Montin, in Nov. Act. Nat. Cur. vi. (1778) 324. t. 5 = piscidium.
bipinnatifidum, *Desv. Journ. Bot.* iii. (1814) 165.—Am. austr.
bipinnatum, *Thunb. Prod. Pl. Cap.* 107.—Afr. austr.
Bonannianum, Guss. Fl. Sic. Prod. ii. 211 = nebrodense.
bonariense, Bert. ex Steud. Nom. ed. II. ii. 26 = Berteronianum.
bonariense, *Linn. Sp. Pl.* 645.—Bras.
bonariense, Mill. Gard. Dict. ed. VIII. n. 13 = Senebiera pinnatifida.
bonariense, Pall. Reise, ii. 329 = incisum.
brevicaule, *Barn. in C. Gay, Fl Chil.* i. 165.—Chili.
brevicaule, Hoppe, ex Mert. & Koch, Deutschl. Fl. iv. 519 = Hutchinsia alpina.
Bursa-pastoris, Willd. Prod. Fl. Berl. 211 = Capsella Bursa-pastoris.
caespitosum, *Desv. in Journ. Bot.* iii. (1814) 165, 178.—Armenia.
californicum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 115 = Menziesii.
calycinum, *Godr. in Mém. Acad. Montp.* (*Sect. Medic.*) i. (1853) 416.—Hab.?
calycinum, Steph. ex Willd. Sp. Pl. iii. 433 = Smelowskia calycina.
calycotrichum, Kunze, in Flora, xxix. (1846) 756 = nebrodense.
campestre, Brot. ex Nym. Consp. 65 = hirtum.

LEPIDIUM:—
campestre, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 88.—Europ.
campicolum, Jord. Diagn. i. 329 = campestre.
Candollei, Desv. Journ. Bot. iii. (1814) 165 = crassifolium.
capense, R. Br. ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 175 = ruderale.
capense, Eckl. & Zeyh. Enum. 6 = africanum.
capense, E. Mey. ex Harv. & Sond. Fl. Cap. i. 30 = divaricatum.
capense, *Thunb. Prod. Pl. Cap.* 107.—Afr. austr.
capitatum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 175.—Reg. Himal.
Cardamines, *Linn. Cent. Pl.* i. 17; *Amoen. Acad.* iv. 278.—Hispan.
Carrerasii, *Rodrig. in Anal. Soc. Esp. Hist. Nat.* iii. (1874) 7.—Hispan.
cepeaefolium, Reichb. in Moessl. Handb. ed. II. ii. 1125 = Thlaspi cepeaefolium.
ceratocarpum, Pall. Reise, ii. 740 = Thlaspi ceratocarpon.
chalepense, Fisch. ex DC. Syst. ii. 388 = Thlaspi hastulatum.
chalepense, Ledeb. Fl. Ross. i. 203 = L. Draba.
chalepense, *Linn. Cent. Pl.* ii. 23; *Amoen. Acad.* iv. 321.—Oriens.
Chichicara, *Desv. Journ. Bot.* iii. (1814) 165.—Bras.
chilense, *Kunze, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* I. (1843) 250.—Chili.
chinense, *Franch. Pl. Davidian.* i. (1884) 39.—China.
chinense, Stokes, Bot. Mat. Med. iii. 429 = Nasturtium sinense.
chloraefolium, Willd. ex Spreng. Syst. ii. 865 = Aethionema chloraefolium.
cordatum, *Willd. ex DC. Syst.* ii. 554.—Turkest.; Sibir.
cornutum, Sibth. & Sm. Fl. Graec. Prod. ii. 6 = spinosum.
cornutum, Urv. in Mém. Soc. Linn. Par. i. (1822) 334 = Andrzeiowskia Cardamine.
coronopifolium, [*Fisch.*] *Cat. Hort. Gorenk.* ed. I. (1808) 79.—Persia; Turkest.
coronopifolium, H. B. & K. ex DC. Syst. ii. 532 = Humboldtii.
corymbosum, Hook. & Arn. Bot. Beech. Voy. 323 = montanum.
crassifolium, *Waldst. & Kit. Pl. Rar. Hung.* i. 4. t. 4.—Hungar.; Oriens.
crenatifolium, *Boiss. & Bal. Diagn.* Ser. II. vi. 22.—Cappadoc.
crispum, Desv. Journ. Bot. iii. (1814) 165, 176 = ruderale.
cristatum, Lapeyr. Hist. Abr. Pl. Pyr. 366 = Senebiera Coronopus.
Cumingianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 30.—Chili.
cuneifolium, DC. Syst. ii. 545 = foliosum.
decumbens, Desv. Journ. Bot. iii. (1814) 165, 176 = incisum.
densiflorum, Schrad. Ind. Sem. Hort. Gotting. (1832) 4 = incisum.
dentatum, *Bieb. ex Georgi, Beschr. Russ. Reich.* III. iv. 1021.—Ross.
Descemetri, Rayneval, in Bull. Soc. Bot. Fr. ii. (1855) 738 = crassifolium.
desertorum, *Eckl. & Zeyh. Enum.* 6.—Afr. austr.
desertorum, Schrenk, Enum. Pl. Nov. 99 = Stroganowia brachyota.
dhayense, Munby, in Bull. Soc. Bot. Fr. ii. (1855) 282 = nebrodense.
diandrum, Medic. in Usteri, Ann. Bot. viii. (1794) 45 = L. Iberis.
dictyotum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 329.—Am. bor. occ.
didymum, Linn. Mant. i. 92 = Senebiera didyma.
diffusum, *DC. Syst.* ii. 554.—Am. bor.
divaricatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 375.—Afr. austr.
divaricatum, Willd. ex DC. Syst. ii. 553 = incisum.
diversifolium, Pers. Syn. ii. 187 = capense.
Draba, *Linn. Sp. Pl.* 645.—Europ.; Oriens.
drabifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = praec.
Eckloni, Schrad. ex Regel, Ind. Sem. Hort. Petrop. (1856) 33 = capense.
eremophilum, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 199.—As. centr.
erinaceum, *Boiss. Diagn.* Ser. I. vi. 17.—Persia.
errabundum, Jord. Diagn. i. 330 = campestris.

LEPIDIUM:—
exiguiflorum, [Clairv.] Man. Herb. 214 = L. Iberis.
fastigiatum, *Ledeb. Fl. Ross.* 205.—Ross. austr.
fenestratum, E. Mey. ex Harv. & Sond. Fl. Cap. i. 129 = desertorum.
flaccidum, *Roth, ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 309.—Reg. Casp.
flavum, *Torr. in Pacif. Rail. Rep.* iv. 67.—Am. bor. occ.
flexicaule, *T. Kirk, in Trans. N. Z. Inst.* xiv. (1882) 380.—N. Zel.
flexuosum, Eckl. & Zeyh. Enum. 6 = capense.
flexuosum, *Thunb. Prod. Pl. Cap.* 107.—Afr. austr.
foliosum, *Desv. Journ. Bot.* iii. (1814) 164, 180.—Austral.
fragrans, Willd. in Roem. & Usteri, Mag. iv. (1790) 11tes St. 36 = Alyssum maritimum.
Fremontii, *S. Wats. in Bot. King's Exped.* 30. t. 4. ff. 3, 4.—Am. bor. occ.
fruticosum, DC. Prod. i. 208 = seq.
fruticulosum, Desv. Journ. Bot. iii. (1814) 165, 180 = ruderale.
gelidum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 283.—Bolivia.
glastifolium, *Desf. Fl. Atlant.* ii. 66. t. 147.—Alger.
glaucescens, Dum. Fl. Belg. 119 = ruderale?
gramineum, Lam. Fl. Fr. ii. 469 = L. Iberis.
graminifolium, Cav. Ic. ii. 41. t. 151. f. 2 = suffruticosum, *Linn.*
graminifolium, *Linn. Syst.* ed. X. 1127.—Europ.; Syria.
granatense, Coss. Notes Crit. ii. 27 = nebrodense.
groenlandicum, *Hornem. in Fl. Dan.* t. 569.—Groenland.
Gussoni, Schrad. ex Steud. Nom. ed. II. ii. 27 = nebrodense.
Halleri, Crantz, Stirp. Austr. ed I. fasc. i. 5 = Hutchinsia alpina.
heterophyllum, Benth. Cat. Pl. Pyr. 95 = hirtum.
heterophyllum, Boiss. Voy. Bot. Espagne, ii. 51 = nebrodense.
hirtellum, *Sond. in Harv. & Sond. Fl. Cap.* i. 30.—Afr. austr.
hirtum, *Sm. Comp. Fl. Brit.* ed. III. 98.—Europ.
Humboldtii, *DC. Syst.* ii. 532.—Ecuador.
humifusum, *Reg. Cat.* 9; *et ex Spreng. Syst.* iv. *Cur. Post.* 240.—Corsica.
hyssopifolium, Desv. Journ. Bot. iii. (1814) 164, 179 = ruderale.
ibericum, Schrad. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 39 = ruderale.
iberifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Iberis.
iberioides, Desv. Journ. Bot. iii. (1814) 165, 176 = ruderale.
Iberis, *Linn. Sp. Pl.* 645.—Europ. austr.
Iberis, Poll. Hist. Pl. Palat. ii. 209 = graminifolium.
Iberis, Schkuhr, Handb. ii. 222. t. 180 = virginicum.
impressum, Bunge, in Lehm. Pl. Preiss. i. 260 = foliosum.
incisum, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 15.—N. Zel.
incisum, Edgew. in Trans. Linn. Soc. xx. (1851) 33 = capitatum.
incisum, *Roth, Neue Beitr.* i. 224; *Bieb. Fl. Taur. Cauc.* ii. 98.—Rossia; Sibir.
integrifolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 116.—Am. bor. occ.
intermedium, *A. Gray, Pl. Wright.* ii. 15.—Am. bor. occ.
intermedium, *A. Rich. Tent. Fl. Abyss.* i. 21.—Abyssin.
intermixtum, *Tenore, Fl. Neap. Syll. App.* v. 18.—Ital.
intricatum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 26 = Moriera spinosa.
karataviense, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 242.—Turkestan.
Kawarau, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 270.—N. Zel.
Kirilowii, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) I. 130.—As. centr.
lacerum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 191.—Persia; Sibir.
laciniatum, Willd. ex DC. Syst. ii. 532 = coronopifolium.
lanatum, *Barn. in C. Gay, Fl. Chil.* i. 167.—Chili.
lanceolatum, *Presl, Fl. Sic.* i. 82.—Sicilia.
lanceolatum, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 250.—Peruv.
lasiocarpum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 115.—Am. bor. occ.

LEPIDIUM:—
latifolium, *Linn. Sp. Pl.* 644.—Europ.; As. bor.; Oriens.
latipes, *Hook. Ic. Pl.* t. 41.—Am. bor. occ.
leiocarpum, DC. Syst. ii. 536 = Aethionema coridifolium.
leiocarpum, Hook. & Arn. Bot. Beech. Voy. 324 = nitidum.
leptopetalum, *F. Muell. Vict. Pl.* i. 48.—Austral.
lineare, *DC. in Lam. Encyc.* v. 46.—Hispan.
linifolium, *Steud. Nom.* ed. II. ii. 27; *Benth. Fl. Austral.* i. 84.—Austral.
Linnaei, Crantz, Stirp. Austr. ed. I. fasc. i. 7 = Hutchinsia petraea.
linoides, *Thunb. Prod. Pl. Cap.* 107.—Afr. austr.
lyratum, *Linn. Sp. Pl.* 644.—Armenia.
majus, *Darracq, in Bull. Soc. Bot. Fr.* xv. (1868) p. xiii.—Gallia.
marginatum, *Griseb. in Goett. Abh.* xix. (1874) 72.—Reg. Argent.
marginatum, Lapeyr. Hist. Abr. Pl. Pyr. i. 365 = Aethionema ovalifolium.
membranaceum, Steud. Nom. ed. II. ii. 27 = Aethionema membranaceum.
mendocinum, *Phil. Sert. Mendoc. Alt.* 2.—Chili.
Menziesii, *DC. Syst.* ii. 539.—Am. bor. occ.
Meyenii, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 249.—Bolivia.
Meyeri, *Claus, in Beitr. Pfl. Russ. Reich.* vii. 286.—Ross.
micranthum, Ledeb. Ic. Pl. Ross. i. 22. t. 92 = incisum, *Roth.*
micropetalum, Stev. ex Ledeb. Fl. Ross. i. 205 = incisum, *Roth.*
micropterum, *Miq. Stirp. Surin. Sel.* 106.—Guian.
microstylum, Boiss. & Heldr. Diagn. Ser. II. vi. 1 = nebrodense.
miscellaneorum, Schultz, Obs. Bot. 122 = L. Iberis.
mixtum, Jord. Diagn. i. 332 = graminifolium.
monoplocoides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 35.—Austral.
montanum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 116.—Am. bor. occ.
myriocarpum, *Sond. in Linnaea*, xxiii. (1850) 4.—Afr. austr.
nanum, *S. Wats. in Bot. King's Exped.* 30. t. 4. ff. 5–7.—Am. bor. occ.
nebrodense, *Guss. Fl. Sic. Syn.* ii. 154.—Sicil.; Ins. Graec.
niloticum, *Sieber, ex Steud. Nom.* ed. II. ii. 26.—Aegypt.
nitidum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 116.—Am. bor. occ.
Novae-Hollandiae, Desv. Journ. Bot. iii. (1814) 177 = ruderale.
nudicaule, Linn. Sp. Pl. 643 = Teesdalia nudicaulis.
obovatum, Kit. ex Steud. Nom. ed. II. ii. 27 = incisum.
obtusum, *Basiner, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 203.—As. med.
oleraceum, *Forst. f. Prod.* 46.—N. Zel.
oleraceum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 374 = piscidium.
oppositifolium, Labill. Ic. Pl. Syr. Dec. v. 14. t. 9 = Aethionema oppositifolium.
owaihiense, *Cham. & Schlecht. in Linnaea*, i. (1826) 32.—Ins. Sandvic.
oxycarpum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 116.—Am. bor. occ.
oxyotum, DC. Syst. ii. 530 = humifusum.
papillosum, *F. Muell. in Linnaea*, xxv. (1852) 370.—Austral.
parviflorum, *Pomel, Nouv. Mat. Fl. Atl.* 378.—Alger.
pedicellosum, *F. Muell. Fragm.* xi. 27.—Austral.
perfoliatum, *Linn. Sp. Pl.* 643.—Europ. austr.; Oriens.
persicum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 196.—Persia.
petraeum, Linn. Sp. Pl. 644 = Hutchinsia petraea.
petraeum, Lour. Fl. Cochinch. 395 = Nasturtium sinense.
petrophilum, Coss. Notes Crit. 148 = nebrodense?
phlebopetalum, *F. Muell. Vict. Pl.* i. 47.—Austral.
pinnatifidum, *Ledeb. Fl. Ross.* i. 206.—Reg. Casp.
pinnatum, Eckl. & Zeyh. Enum. 7; et ex Harv. & Sond. Fl. Cap. i. 30 = hirtellum.
pinnatum, Thunb. Prod. Pl. Cap. 107 = divaricatum.
piscidium, *Forst. f. Prod.* 46.—Ins. Pacif.
Pollichii, Roth, Tent. Fl. Germ. ii. 91 = graminifolium.
polycladum, Jord. Diagn. i. 331 = graminifolium.

LEPIDIUM:—
pratense, Serres, ex Billot, in F. Schultz, Arch. Fl. Fr. et Allem. 197 = hirtum.
praecox, DC. Syst. ii. 554 = virginicum.
procumbens, Linn. Sp. Pl. 643 = Capsella procumbens.
propinquum, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 378.—Reg. Caucas.
prostratum, Savi, in Santi, Viag. ii. 18. t. 1 = Senebiera pinnatifida.
puberulum, Bunge, in Lehm. Pl. Preiss. i. 261 = ruderale.
pubescens, *Desv. Journ. Bot.* iii. (1814) 165, 180.—Bras.
pubescens, Tineo, Cat. Hort. Panorm. 150 = nebrodense.
pumilum, *Boiss. & Bal. Diagn.* Ser. II. vi. 21.—Cappadoc.
pusillum, Lam. Fl. Fr. ii. 468 = Capsella procumbens.
quitense, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 309.—Ecuador.
racemosum, *Griseb. in Goett. Abh.* vi. (1854) 116.—Reg. Magellan.
Ramburii, *Boiss. Voy. Bot. Espagne*, ii. 52. t. 14.—Hispan.
repens, Boiss. Fl. Orient. i. 356 = L. Draba.
rigidum, *Pomel, Nouv. Mat. Fl. Atl.* 378.—Alger.
rotundifolium, All. Fl. Pedem. i. 252 = Thlaspi rotundifolium.
rotundum, *DC. Syst.* ii. 537.—Austral.
ruderale, Burm. f. Prod. Fl. Cap. 17 = Iberis pinnata.
ruderale, *Linn. Sp. Pl.* 645.—Europ.; Oriens; Austral.
salinum, Georgi, Beschr. Russ. Reich. III. iv. 1121 = seq.
salsum, Pall. Reise in Suedl. Statth. Russ. Reich. ii. 336 = crassifolium.
sativum, Cambess. Enum. Pl. Balear. 38 = Carrerasii.
sativum, *Linn. Sp. Pl.* 644.—Oriens.
scapiferum, Wallr. Sched. Crit. 344 = Teesdalia nudicaulis.
sectifolium, *Steud. in Flora*, xxxix. (1856) 412.—Peruv.
Serra, *H. Mann, in Proc. Am. Acad.* vii. (1867) 149.—Ins. Sandvic.
setosum, *Boiss. Diagn.* Ser. II. i. 45.—Afghan.
sibiricum, Pall. Reise, iii. 34 = Draba repens.
sibiricum, Schweigg. Enum. Pl. Hort. Regiom. (1812) 43 = latifolium.
Sieberi, *W. Mann, in Syll. Ratisb.* i. (1824) 171.—Creta.
sisymbrioides, *Hook. f. Handb. N. Zel. Fl.* 14.—N. Zel.
Smithii, Hook. Brit. Fl. ed. III. 300 = hirtum.
Solandri, T. Kirk, in Trans. N. Z. Inst. xiv. (1882) 380 = sisymbrioides.
songaricum, Schrenk, Enum. Pl. Nov. 98 = coronopifolium.
sordidum, *A. Gray, Pl. Wright.* i. 10; ii. 15.—Texas.
spathulatum, *Phil. Fl. Atac.* 8.—Chili.
spicatum, *Desv. Journ. Bot.* iii. (1814) 164, 178.—Chili.
spinescens, *DC. Syst.* ii. 534.—Syria; Mesopot.
spinosum, *Linn. Mant.* ii. 253.—Oriens.
squamatum, Forsk. Fl. Aegypt. Arab. 117 = Senebiera Coronopus.
strongylophyllum, *F. Muell. ex Benth. Fl. Austral.* i. 84.—Austral.
stylatum, Lag. & Rodr. in Cienc. Anal. Nat. v. (1802) 276 = nebrodense?
stylosum, Pers. Syn. ii. 187 = Nasturtium pyrenaicum.
subdentatum, Burch. ex DC. Syst. ii. 545 = linoides.
subdentatum, Eckl. & Zeyh. Enum. 6 = africanum.
subdentatum, Meissn. ex Harv. & Sond. Fl. Cap. i. 29 = capense.
subulatum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 90 = ruderale.
subulatum, *Linn. Sp. Pl.* 644.—Europ.
subvaginatum, *Steud. Nom.* ed. II. ii. 28.—Chili.
suffruticosum, *Linn. Mant.* i. 91.—Hispan.
suffruticosum, *Phil. in Linnaea*, xxviii. (1856) 670.—Chili.
sylvaticum, Eckl. & Zeyh. Enum. 6 = capense.
tenuicaule, *T. Kirk, in Trans. N. Z. Inst.* xiv. (1882) 381.—N. Zel.
tenuissimum, *Steud. Nom.* ed. II. ii. 28.—Chili.
tetraspermum, Dufour, ex DC. Syst. ii. 390 = Hutchinsia petraea.
texanum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 449 = intermedium, *A. Gray*.
Thlaspi, Roxb. Hort. Beng. 48; Fl. Ind. iii. 116 = Thlaspi arvense.

LEPIDIUM:—
thlaspidioides, Pall. Reise, iii. 161 = Thlaspi alpestre.
Tournefortii, Spreng. Syst. ii. 866 = Aethionema cordatum.
triandrum, Stokes, Bot. Mat. Med. iii. 426 = virginicum.
trifurcum, *Sond. in Linnaea*, xxiii. (1850) 4.—Afr. austr.
tuberosum, DC. Syst. ii. 554 = Cardamine rhomboidea?
utaviense, Regel, in Act. Hort. Petrop. i. (1871) 92 = montanum.
vagum, Jord. Diagn. i. 331 = campestre.
verrucosum, DC. in Mém. Soc. Hist. Nat. Par. (1799) 145 = crassifolium.
vesicarium, *Linn. Sp. Pl.* 643.—Reg. Cauc.; Persia.
Villarsii, Gren. & Godr. Fl. Fr. i. 150 = hirtum.
violiforme, DC. in Mém. Soc. Hist. Nat. Par. (1799) 145 = Cochlearia acaulis.
virgatum, Jord. Diagn. i. 333 = graminifolium.
virginicum, Gren. & Godr. Fl. Fr. i. 151 = majus.
virginicum, *Linn. Sp. Pl.* 645.—Am. bor.
Wrightii, *A. Gray, Pl. Wright.* ii. 15.—Am. bor. occ.

LEPIDOBOLUS, Nees, in Lehm. Pl. Preiss. ii. 66 (1846). *RESTIACEAE*, Benth. & Hook. f. iii. 1033.
chaetocephalus, *F. Muell. Fragm.* viii. 84.—Austral.
drapetocoleus, *F. Muell. l. c.*—Austral.
Preissianus, *Nees, in Lehm. Pl. Preiss.* ii. 66.—Austral.

LEPIDOCARPA, Korth. in. Nederl. Kruidk. Arch. iii. (1855) 385 = **Parinarium**, Juss. (Rosac.).
costata, Korth. l. c. 387 = P. costatum.
ovalis, Korth. l. c. 386 = P. costatum.

LEPIDOCARPUS, Adans. Fam. ii. 284 (1763) = **Protea**, Linn.

LEPIDOCARYA, Korth. ex Miq. Fl. Ind. Bat. i. 1. 353 (1855) = praec.

LEPIDOCARYON, Spreng. Syst. ii. 138 (1825) = seq.

LEPIDOCARYUM, Mart. Hist. Nat. Palm. ii. 49 (1824). *PALMAE*, Benth. & Hook. f. iii. 938.
casiquiarense, Spruce, ex Drude, in Mart. Fl. Bras. iii. II. 300 = tenue.
enneaphyllum, Rodrig. Enum. Palm. Nov. 19 = tenue.
gracile, *Mart. Hist. Nat. Palm.* ii. 50. tt. 45, 46.—Bras.
guainiense, *Spruce, ex Drude, in Mart. Fl. Bras.* iii. II. 300.—Bras.
quadripartitum, Spruce, l. c. 298 = tenue.
sexpartitum, Rodrig. Enum. Palm. Nov. 19 = tenue.
tenue, *Mart. Hist. Nat. Palm.* ii. 51. t. 47.—Bras.

LEPIDOCERAS, Hook. f. Fl. Antarct. ii. 293 (1846). *LORANTHACEAE*, Benth. & Hook. f. iii. 216.
MYRTOBIUM, Miq. in Linnaea, xxv. (1852) 652.
Dombeyi, *Hook. f. Fl. Antarct.* ii. 293.—Chili.
Kingii, *Hook. f. l. c.*—Chili.
punctulatum, Clos, in C. Gay, Fl. Chil. iii. 165 = Eremolepis punctulata.
punctulatum, Griseb. in Goett. Abh. vi. (1854) 124 = Eremolepis verrucosa.
squamifer, *Clos, in C. Gay, Fl. Chil.* iii. 166.—Chili.

LEPIDOCOCCA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 588 = **Caperonia**, St. Hil. (Euphorb.).
serrata, Turcz. l. c. 589 = Chrozophora senegalensis.
Sieberi, Turcz. l. c. = Caperonia palustris.

LEPIDOCOCCUS, H. Wendl. & Drude, in Kerch. Palm. 249 (1878) = **Mauritia**, Linn. (Palm.).
aculeatus, H. Wendl. & Drude, l. c. = M. aculeata.
armatus, H. Wendl. & Drude, l. c. = M. armata.
pumilus, H. Wendl. & Drude, l. c. = M. pumila.

LEPIDOCOMA, Jungh. Reisen durch Java, 338 (1845) = **Flemingia**, Roxb. (Legumin.).
trifoliatum, Jungh. l. c. = F. involucrata.

LEPIDOCROTON, Klotzsch, in Schomb. Faun. et Fl. Gui. 1162, nomen (1848). *MENISPERMACEAE*?
latifolius, *Klotzsch, l. c.* (Quid?).—Guian.

LEPIDOCROTON, Presl, Epim. Bot. 213 (1850) = **Chrozophora**, Neck. (Euphorb.).
serratus, Presl, l. c. = C. senegalensis.

LEPIDOGYNE, Blume, Orch. Archip. Ind. 93. t. 25 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 603.
longifolia, *Blume, l. c.*—Java.

LEPIDONEMA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 31 (1835) = **Microseris**, D. Don (Compos.).
chilense, Fisch. & Mey. l. c. = M. pygmaea.

LEPIDOPAPPUS, Moç. & Sesse, ex DC. Prod. v. 655 (1836) = **Florestina**, Cass. (Compos.).
tripteris, Moç. & Sesse, l. c. = F. tripteris.

LEPIDOPELMA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 118. t. 22 (1862) = **Sarcococca**, Lindl. (Euphorb.).
podocarpifolia, Klotzsch, l. c. = S. pruniformis.

LEPIDOPETALUM, Blume, Rumphia, iii. 171 (1847) = **Ratonia**, DC. (Sapind.).
Jackianum, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 623 = Cupania Jackiana.
montanum, Radlk. l. c.—Sumatra.
Perrottetii, Blume, Rumphia, iii. 172.—Ins. Philipp.

LEPIDOPHORUM, Neck. Elem. i. 14 (1790) = **Anthemis**, Mich. (Compos.).
repandum, DC. Prod. vi. 19 = A. repanda.

LEPIDOPHYLLUM, Cass. in Bull. Soc. Philom. (1816) 199. *COMPOSITAE*, Benth. & Hook. f. ii. 257.
TOLA, Wedd. ex Benth. & Hook. f. Gen. ii. 257 (1873).
acaule, *Benth. & Hook. f. Gen.* ii. 259.—Bolivia.
cupressiforme, *Cass. in Dict. Sc. Nat.* xxvi. 37.—Reg. Magell.
Meyeni, *A. Gray, in Proc. Am. Acad.* v. (1862) 123.—Reg. Andin.
quadrangulare, Benth. & Hook. f. Gen. ii. 258 = Meyeni.
rigidum, *A. Gray, in Proc. Am. Acad.* v. (1862) 123.—Reg. Andin.
rupestre, *A. Gray, l. c.*—Reg. Andin.

LEPIDOPHYTON, Benth. & Hook. f. Gen. iii. 238 (1880) = seq.

LEPIDOPHYTUM, Hook. f. in Lindl. Veg. Kingd. ed. III. 90 (1853), nomen = **Lophophytum**, Schott (Balanoph.).

LEPIDOPIRONIA, A. Rich. Tent. Fl. Abyss. ii. 442. t. 101 (1850). *GRAMINEAE*, Benth. & Hook. f. iii. 1170.
cenchriformis, *A. Rich. l. c.*—Abyssin.

LEPIDOPLOA, Sch. Bip. in Linnaea, xx. (1847) 508 sphalm. = LEPIDAPLOA, Cass. = **Vernonia**, Schreb. (Compos.).

LEPIDOPOGON, Tausch, in Flora, xii. (1829) 1. Erg. 37 = **Cylindrocline**, Cass. (Compos.).
gnaphalodes, Tausch, l. c. xiv. (1831) 224 = C. Commersonii.
Ponae, Tausch, l. c. xii. (1829) 1. Erg. 37 = C. Commersonii.

LEPIDOPYRONIA, Benth. in Journ. Linn. Soc. xix. (1881) 106 = **Lepidopironia**, A. Rich. (Gramin.).

LEPIDOSERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 103 = **Crepis**, Linn. (Compos.).
recognita, Fourr. l. c. = C. recognita.
taraxacifolia, Fourr. l. c. = C. taraxacifolia.

LEPIDOSPARTUM, A. Gray, in Proc. Am. Acad. xix. (1883) 50. *COMPOSITAE*.
squamatum, *A. Gray, l. c.*—Calif.

LEPIDOSPERMA, Labill. Pl. Nov. Holl. i. 14. t. 11 (1804). *CYPERACEAE*, Benth. & Hook. f. iii. 1065.
LEPIDOTOSPERMA, Roem. & Schult. Syst. ii. 78 (1817).

LEPIDOSPERMA:—
MACHAERINA, Nees, in Lehm. Pugill. viii. 54 (1844) (Macherina).
anceps, Willd. ex Link, Jahrb. i. III. 75 = Cladium iridifolium.
angustatum, *R. Br. Prod.* 235.—Austral.
angustifolium, Hook. f. Fl. Tasm. ii. 92. t. 147 = laterale.
aphyllum, *R. Br. Prod.* 235.—Austral.
australe, Hook. f. Fl. N. Zel. i. 279 = tetragonum.
Brunonianum, *Nees, in Lehm. Pl. Preiss.* ii. 92.—Austral.
Burmanni, Spreng. Syst. i. 194 (= *Schoenus Burmanni*).—Afr. austr.
canescens, *Boeck. in Linnaea*, xxxviii. (1874) 330.—Austral.
carphoides, *F. Muell. ex Benth. Fl. Austral.* vii. 400.—Austral.
chinense, *Nees, ex Meyen, in Linnaea*, ix. (1834) 302.—China; Malacca.
Colensoi, Boeck. l. c. xxxviii. (1874) 328 = Cladium junceum.
concavum, *R. Br. Prod.* 234.—Austral.
concavum, Hook. f. Fl. Tasm. ii. 91. t. 146 = laterale.
confine, Nees, in Lehm. Pl. Preiss. ii. 93 = striatum.
congestum, *R. Br. Prod.* 234.—Austral.
costale, *Nees, in Lehm. Pl. Preiss.* ii. 92.—Austral.
Drummondii, *Benth. Fl. Austral.* vii. 391.—Austral.
effusum, *Benth. l. c.* 387.—Austral.
elatius, *A. Cunn. ex Hook. f. Handb. N. Zeal. Fl.* 307.—N. Zel.
elatius, *Labill. Nov. Holl. Pl.* i. 15. t. 11.—Austral.
ensatum, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 47 = gladiatum.
ensifolium, Willd. ex Kunth, Enum. Pl. ii. 315 = Cladium ensifolium.
exaltatum, *R. Br. Prod.* 234.—Austral.
filiforme, *Labill. Pl. Nov. Holl.* i. 17. t. 15.—Austral.
fimbriatum, Nees, in Lehm. Pl. Preiss. ii. 91 = angustatum.
flexuosum, *R. Br. Prod.* 235.—Austral.
gladiatum, *Labill. Nov. Holl. Pl.* i. 15. t. 12.—Austral.
gladiatum, Nees, ex Benth. Fl. Austral. vii. 390 = squamatum.
globosum, *Labill. Nov. Holl. Pl.* i. 16. t. 14.—Austral.
gracile, *R. Br. Prod.* 235.—Austral.
Gunnii, Boeck. in Linnaea, xxxviii. (1874) 325 = lineare.
humile, *Boeck. l. c.*—Austral.
involucratum, Hort. Lucae, ex Steud. Nom. ed. II. i. 341, in syn. = Schoenus flexuosus.
involucratum, *Roem. & Schult. Syst.* ii. 81.—Austral.
involucratum, Schrad. ex Schult. Mant. ii. 474 = Rottboellii.
iridifolium, Willd. ex Link, Jahrb. i. III. 75 = Cladium iridifolium.
laeve, R. Br. Prod. 235 = globosum.
laterale, *R. Br. l. c.* 234.—Austral.
laterale, Hook. f. Fl. Tasm. ii. 91. t. 147 = squamatum.
leptophyllum, *Benth. Fl. Austral.* vii. 398.—Austral.
leptostachyum, *Benth. l. c.* 397.—Austral.
lineare, *R. Br. Prod.* 235.—Austral.
lineare, Nees, ex Kunth, Enum. Pl. ii. 318 = laterale.
longitudinale, R. Br. Prod. 234 = laterale.
longitudinale, *Labill. Nov. Holl. Pl.* i. 16. t. 13.—Austral.
Muelleri, Boeck. in Linnaea, xxxviii. (1874) 320 = angustatum.
Neesii, *Kunth, Enum. Pl.* ii. 319.—Austral.
Oldfieldii, *Hook. Fl. Tasm.* ii. 91. t. 146.—Austral.
pauciflorum, F. Muell. Fragm. ix. 23 = Tricostularia pauciflora.
pubisquameum, *Steud. Syn. Pl. Cyp.* 158.—Austral.
resinosum, *F. Muell. ex Benth. Fl. Austral.* vii. 392.—Austral.
Rottboellii, *Schrad. ex Schult. Mant.* ii. 474.—Afr. austr.
rupestre, *Benth. Fl. Austral.* vii. 388.—Austral.
scabrum, *Nees, in Lehm. Pl. Preiss.* ii. 92.—Austral.
semiteres, *F. Muell. ex Boeck. in Linnaea*, xxxviii. (1874) 327.—Austral.
Sieberi, Kunth, Enum. Pl. ii. 320 = squamatum.
Sieberi, Nees, in Lehm. Pl. Preiss. ii. 90 = resinosum.
squamatum, *Labill. Nov. Holl. Pl.* i. 17. t. 16.—Austral.
striatum, *R. Br. Prod.* 235.—Austral.

LEPIDOSPERMA:—
striatum, F. Muell. Fragm. ix. 27 = carphoides.
tenue, *Benth. Fl. Austral.* vii. 397.—Austral.
tetragonum, *R. Br. Prod.* 235.—Austral.
tetragonum, Labill. Nov. Holl. Pl. i. 17. t. 17 = Cladium tetraquetrum.
tetragonum, Nees, ex Kunth, Enum. Pl. ii. 319 = Neesii.
tetragynum, R. Br. Prod. 234 = laterale.
tetraquetrum, *Nees, in Lehm. Pl. Preiss.* ii. 90.—Austral.
thermale, Schrad. in Goett. Gel. Anz. (1821) III. 2070 (Tetrariae sp.).—Afr. austr.
tortuosum, *F. Muell. Fragm.* ix. 23.—Austral.
triangulare, Boeck. in Linnaea, xxxviii. (1874) 336 (Tetrariae sp.).—Afr. austr.
tuberculatum, *Nees, in Lehm. Pl. Preiss.* ii. 90.—Austral.
uncinatum, Nees, l. c. 93 = Mesomelaena stygia.
ustulatum, Steud. Syn. Pl. Cyp. 157 = angustatum.
viscidum, R. Br. Prod. 234 = angustatum.
waigiuensis, *Steud. Syn. Pl. Cyp.* 158.—Ins. Waigiu.
zeylanicum, Lindl. ex Boeck. in Linnaea, xxxviii. (1874) 332 = Cladium undulatum.

LEPIDOSPORA, F. Muell. Fragm. ix. 34 (1875). *CYPERACEAE.*
tenuissima, *F. Muell. l. c.*—Austral.

LEPIDOSTACHYS, Wall. Cat. n. 6816 (1832) = **Aporosa**, Blume (Euphorb.).
grandifolia, Planch. ex Muell. Arg. in DC. Prod. xv. II. 471 = A. fusiformis.
lanceolata, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 254 = A. lanceolata.
macrophylla, Tul. l. c. 253 = A. macrophylla.
oblonga, Wall. Cat. n. 7299 = A. oblonga.
parviflora, Planch. ex Baill. Etud. Gén. Euph. 645, nomen = A. Planchoniana.
Roxburghii, Wall. Cat. n. 6816 = A. Roxburghii.
villosa, Wall. l. c. 7298 = A. villosa.

LEPIDOSTEMON, Hassk. Cat. Hort. Bog. Alt. 140 (1844) = **Lepistemon**, Blume (Convolv.).

LEPIDOSTEMON, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 131. *CRUCIFERAE*, Benth. & Hook. f. i. 77.
pedunculosus, *Hook. f. & Thoms. l. c.* 156.—Reg. Himal.

LEPIDOSTEMON, Lem. Illustr. Hortic. ix. (1862) t. 315 = **Pentstemon**, Mitch. (Scrophul.).
penstemonoides, Lem. l. c. = P. antirrhinoides.

LEPIDOSTEPHANUS, Bartl. Ind. Sem. Hort. Gotting. (1837); ex Linnaea, xii. (1838) Litt. 82 = **Achyrachaena**, Schau. (Compos.).
madioides, Bartl. l. c. = A. mollis.

LEPIDOSTEPHIUM, Oliver, in Hook. Ic. Pl. xi. 22. t. 1030 (1868). *COMPOSITAE*, Benth. & Hook. f. ii. 422.
denticulatum, *Oliver, l. c.*—Afr. austr.

LEPIDOTHAMNUS, Phil. in Linnaea, xxx. (1859–60) 730 = **Dacrydium**, Soland. (Conif.).
Fonki, Phil. l. c. 731.—Chili.

LEPIDOTHECA, Nutt. in Am. Phil. Trans. vii. (1841) [454] = **Matricaria**, Tourn. (Compos.).
suaveolens, Nutt. l. c. [454,] 397 = M. discoidea.

LEPIDOTOSPERMA, Roem. & Schult. Syst. ii. 78 (1817) = **Lepidosperma**, Labill. (Cyperac.).

LEPIDOTURUS, Baill. Etud. Gén. Euph. 448 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 316.
alnifolius, *Baill. l. c.* 449.—Madag.
laxiflorus, *Benth. in Hook. Ic. Pl.* xiii. 76. t. 1297.—Afr. trop.
occidentalis, *Muell. Arg. in Journ. Bot.* ii. (1864) 332.—Afr. occ.

LEPIDOTURUS, Boj. Hort. Maurit. 285 (1837), nomen. Gen. dub. Euphorbiac.
alnifolius, Boj. l. c. (Acalyphae sp. ?).—Madag.

LEPIDOZAMIA, Regel, Gartenfl. vi. (1857) 11 = **Macrozamia**, Miq. (Cycad.).
Denisoni, Regel, l. c. (1876) 3. t. 854 = M. Denisonii.
Hopei, Regel, l. c. 5, 6 = M. Peroffskyana.
minor, Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 56 = M. Peroffskyana.
Peroffskyana, Regel, in Bull. Soc. Nat. Mosc. (1857) I. 182 = M. Peroffskyana.

LEPIGONUM, Wahlb. Fl. Gothob. 45 (1820) = **Spergularia**, J. & C. Presl (Caryoph.).
anceps, Bartl. in Lehm. Pl. Preiss. i. 244 = S. rubra.
arenarium, Kindb. Syn. Framst. Lepig. 13 = S. grandis.
australe, F. Muell. ex Kindb. Monog. Lepig. in Ind. = S. media.
azoricum, Kindb. l. c. 30. t. 11. f. 15 = S. media.
brevifolium, Bartl. in Lehm. Pl. Preiss. i. 243 = S. rubra.
caespitosum, Kindb. Syn. Framst. Lepig. 13 = S. grandis.
campestre, Kindb. Monog. Lepig. 35 = S. diandra.
canadense, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 14 = S. media, salina.
caninum, Leffl. in Oestr. Bot. Zeitschr. xix. (1869) 103 = S. salina, media, rubra.
capense, Fisch. & Mey. Ind. Sem. Hort. Petrop. I. 10; Regel, l. c. (1856) 34 = S. media.
capillaceum, Kindb. & Lange, Pl. Hisp. Nov. fasc. i. 4 = S. capillacea.
chilense, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 14 = S. macrotheca.
depauperatum, Kindb. Monog. Lepig. 25.—Chili.
depressum, Kindb. l. c. 17 = S. grandis.
diandrum, Kindb. Syn. Framst. Lepig. 35 = S. campestris.
diandrum, Nym. Syll. 250; Fries, Ind. Sem. Hort. Upsal. (1855); ex Kindb. l. c. 7 = S. diandra.
dubium, Kindb. Monog. Lepig. 18. t. 1. f. 6 = S. media.
eximium, Kindb. l. c. 32.—Syria.
fasciculare, Loennr. Obs. Crit. Pl. Suec. (1854) 13 = S. media.
fimbriatum, Kindb. Monog. Lepig. 21 = S. fimbriata.
firmum, Kindb. l. c. 33 = S. purpureum.
floribundum, Kindb. Syn. Framst. Lepig. 5 = *depauperatum*.
glandulosum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 14; Regel, l. c. (1856) 35 = S. diandra.
glandulosum, Liebm. Ind. Sem. Hort. Haun. (1853) 21. —Cf. Linnaea, xxix. (1857–58) 737 = S. Liebmanniana.
glutinosum, G. Don, in Sweet, Hort. Brit. ed. III. 69 = Holosteum liniflorum.
gracile, S. Wats. in Proc. Am. Acad. xvii. (1882) 367 = Am. bor. occ.
grande, Kindb. Syn. Framst. Lepig. 15 = S. grandis.
grandiflorum, Kindb. Monog. Lepig. 34 = S. grandiflora.
halophilum, Fisch. & Mey. ex Avé-Lall. Ind. Sem. Hort. Petrop. x. 22 = S. media.
heterospermum, Guss. Enum. Pl. Inar. 51; Grech-Delic. Fl. Melit. 6.—Europ. austr.
heterospermum, Schur, Enum. Pl. Transs. 126 = S. media.
laeve, Kindb. Monog. Lepig. 32. t. 11. f. 18 = Spergula laevis.
largiflorum, F. Muell. ex Kindb. Monog. Lepig. 29 = S. rubra.
laxiflorum, Bartl. in Lehm. Pl. Preiss. i. 244 = S. rubra.
leiospermum, Kindb. Monog. Lepig. 23 = S. salina.
Liebmannianum, Lange, Ind. Sem. Hort. Haun. (1858) App. 2 = S. Liebmanniana.
macrorhizon, Nym. Syll. 249 = S. macrorhiza.
macrorhizum, Kindb. Monog. Lepig. 22 = praec.
macrothecum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 14 = S. macrotheca.
marginatum, Koch, Syn. Fl. Germ. ed. II. 121 = S. media.
marinum, Kindb. Monog. Lepig. 18 = S. marina.
marinum, Wahlb. Fl. Gothob. 45; Fries, Mant. iii. 35 = S. media.
medium, Fries, l. c. 33 = S. salina.
medium, Wahlb. Fl. Gothob. 45 = S. marina.
melanospermum, Lange, ex Kindb. Monog. Lepig. 42 = S. diandra.

LEPIGONUM:—
microspermum, Kindb. Monog. Lepig. 26.—Abyssin.
molluginеum, Fries, Ind. Sem. Hort. Upsal. (1856).—Cf. Linnaea, xxix. (1857–58) 738 = Arenaria molluginea.
murale, Kindb. Monog. Lepig. 31 = S. villosa.
neglectum, Kindb. Syn. Framst. Lepig. 6 = S. salina.
paradoxum, Kindb. Monog. Lepig. 44 = Arenaria lanuginosa.
prostratum, Kindb. l. c. = S. prostrata.
purpureum, Kindb. l. c. 33 = S. capillacea.
radicans, Kindb. Syn. Framst. Lepig. 10 = S. rubra.
rubrum, Wahlb. Fl. Gothob. 45 = S. rubra.
rupestre, Lebel, ex Kindb. Monog. Lepig. 29 = S. rupestris.
rupicola, Lebel, ex Kindb. Syn. Framst. Lepig. 8 = S. rupestris.
salinum, G. Don, in Sweet, Hort. Brit. ed. III. 69; Fries, Mant. iii. 34 = S. salina.
salsugineum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 10 = S. salsugineum.
salsugineum, Kindb. Monog. Lepig. 42. t. 3 = S. diandra.
segetale, Koch, Syn. Fl. Germ. ed. II. 120 = S. segetalis.
sperguloides, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 10 = S. rubra.
tasmanicum, Kindb. Monog. Lepig. 31 = S. media ?
trachyspermum, Kindb. l. c. = S. villosa.
villosum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 15 = Arenaria villosa.

LEPILAENA, J. Drum. ex Harv. in Hook. Kew Journ. vii. (1855) 57. *NAIADACEAE*, Benth. & Hook. f. iii. 1016.
HEXATHECA, Sond. ex F. Muell. Fragm. viii. 217 (1874).
australis, *J. Drum. ex Harv. in Hook. Kew Journ.* vii. (1855) 58.—Austral.
cylindrocarpa, *Benth. Fl. Austral.* vii. 180.—Austral.
Preissii, *F. Muell. Fragm.* viii. 217.—Austral.

LEPIMENES, Rafin. Fl. Tellur. iv. 91 (1836) = **Cuscuta**, Linn. (Convolv.).
Epithymum, Rafin. l. c. = C. Epithymum.

LEPINEMA, Rafin. Fl. Tellur. iii. 26 (1836) = **Enicostema**, Blume (Gentian.).
verticillata, Rafin. l. c. 27 = E. littorale.

LEPINIA, Decne. in Ann. Sc. Nat. Sér. III. xii. (1849) 194. t. 9. *APOCYNACEAE*, Benth. & Hook. f. ii. 698.
taitensis, *Decne. l. c.*—Ins. Tahiti.

LEPIONURUS, Blume, Bijdr. 1148 (1826). *OLACINEAE*, Benth. & Hook. f. i. 349.
LEPTONIUM, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 236.
javanicus, G. Don, Gen. Syst. ii. 16 = sylvestris.
oblongifolius, *Mast. in Hook. f. Fl. Brit. Ind.* i. 583. —Ind. or.; Malaya.
sylvestris, *Blume, Bijdr.* 1148.—Java.

LEPIOSTEGERES, Benth. & Hook. f. Gen. iii. 211 (1880) err. typ. = LEPEOSTEGERES, Blume = **Loranthus**, Linn.

LEPIPOGON, Bertol. f. in Mem. Acc. Sc. Bologn. iv. (1853) 539. t. 21. *RUBIACEAE.*
obovatum, *Bertol. f. l. c.*—Afr. trop.

LEPIRODIA, Juss. in Dict. Sc. Nat. xlv. 272 (1827) = **Lepyrodia**, R. Br. (Restiac.).

LEPIRONIA, Rich. in Pers. Syn. i. 70 (1805). *CYPERACEAE*, Benth. & Hook. f. iii. 1056.
CHONDRACHNE, R. Br. Prod. 220 (1810).
CHORICARPHA, Boeck. in Flora, xli. (1858) 19.
bancana, Miq. Fl. Ind. Bat. Suppl. 604 = Mapania bancana.
ceylanica, Miq. Illustr. Fl. Arch. Ind. 61. t. 22 = Mapania zeylanica.
cuspidata, Miq. Fl. Ind. Bat. Suppl. 603 = Mapania humilis.
enodis, Miq. l. c. (Mapaniae sp.).—Ins. Malay.
foliosa, Miq. l. c. (Mapaniae sp.).—Malaya.
humilis, Miq. Illustr. Fl. Arch. Ind. 61. t. 23 = Mapania humilis.

LEPIRONIA:—
macrocephala, Miq. l. c. 64. t. 27 (Mapaniae sp.).—Malaya.
mucronata, *Rich. in Pers. Syn.* i. 70.—Madag.; As. trop.; Austral.; Polynesia.
palustris, Miq. Illustr. Fl. Arch. Ind. 63. t. 25 = Mapania palustris.
squamata, Miq. l. c. 64. t. 26 (Mapaniae sp.).—Malaya.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 604 = Mapania sumatrana.

LEPISANTHES, Blume, Bijdr. 238 (1825). *SAPINDACEAE*, Benth. & Hook. f. i. 403.
angustifolia, *Blume, Rumphia*, iii. 154.—Sumatra.
Browniana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 680.—Burma.
burmanica, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 183.—Burma.
confinis, *Blume, Rumphia*, iii. 153.—Java.
cuneata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 680.—Penang.
deficiens, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 276 = Hemigyrosa deficiens.
eriolepis, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 106.—Ins. Philipp.
frutescens, *Blume, Rumphia*, iii. 152.—Java.
heterolepis, *Blume, l. c.* 153.—Borneo.
hirtella, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 105.—Borneo.
longifolia, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 276 = Hemigyrosa longifolia.
montana, *Blume, Bijdr.* 238.—Java.
pallens, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 300, 309.—Ins. Timor.
schizolepis, *Radlk. Sapind. Holland.-Ind. Nachtr.* 87.—Malaya.
sessiliflora, *Blume, Rumphia*, iii. 153.—Java.
tetraphylla, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 276 = Hemigyrosa canescens.

LEPISCLINE, Cass. in Bull. Soc. Philom. (1818) 31 = **Helichrysum**, Vaill. (Compos.).
cymosa, Cass. in Dict. Sc. Nat. xxvi. 49 = H. cymosum.
nudifolia, Cass. l. c. = H. nudifolium.

LEPISELINUM, Presl, Bot. Bemerk. 75 (1844) = LEPISMA, E. Mey. (Umbellif.).
paniculatum, Presl, l. c. = *Lepisma paniculatum*.

LEPISIA, Presl, Lepis. Nov. Pl. Gen. (1829); et Symb. Bot. i. 9. t. 5 (1830) = **Tetraria**, Beauv. (Cyperac.).
ustulata, Presl, l. c. (= *Elynanthus ustulatus*).—Afr. austr.

LEPISIPHON, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 180 = **Osteospermum**, Linn. (Compos.).
dentatus, Turcz. l. c. = O. moniliferum.

LEPISMA, E. Mey. in Drège, Zwei Pfl. Docum. 198 (1848) (Umbellif.).
paniculatum, E. Mey. l. c. = Rhyticarpus rugosus, et Annesorhizae sp. (*Glia gummifera*).—Afr. austr.
verticillatum, E. Mey. l. c. = Polemannia verticillata.

LEPISMIUM, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 315, 380 = **Rhipsalis**, Gaertn. (Cacteae).
alternatum, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 576 = R. alternata.
commune, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 315 = R. Mittleri.
crusiforme, Miq. in Bull. Néerl. (1838) 49 = R. cereiformis?
Duprei, Hort. Par. ex Foerst. Handb. Cact. ed. I. 456.—Hab.?
fluminense, Miq. in Bull. Néerl. (1838) 48.—Bras.
Knightii, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 380 = R. Knightii.
laevigatum, Salm-Dyck, Cat. Hort. Dyck, ed. I. 41, nomen.—Hab.?
Myosurus, Pfeiff. Enum. Diagn. Cact. 139 = R. Myosurus.
paradoxum, Salm-Dyck, ex Pfeiff. l. c. 140 = R. paradoxa.
ramosissimum, Lem. Illustr. Hortic. xi. (1864) Misc. 73.—Bras.
tenue, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 315.—Hab.?

LEPISTACHYA, Zipp. ex Miq. Illustr. Fl. Ind. Arch. 61 (1871) = **Mapania**, Aubl. (Cyperac.).
praemorsa, Zipp. ex Miq. l. c. = M. humilis.

LEPISTEMON, Blume, Bijdr. 722 (1825). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 873.
LEPIDOSTEMON, Hassk. Cat. Hort. Bog. Alt. 140 (1844).
NEMODON, Griff. Notul. iv. 286 (1854).
africanum, *Oliver, in Hook. Ic. Pl.* t. 1270.—Afr. trop.
Fitzalani, *F. Muell. Fragm.* x. 111.—Austral.
flavescens, *Blume, Bijdr.* 722.—Java.
flavescens, Wight, Ic. t. 1862 = Wallichii.
Lucae, *F. Muell. in Victoria Naturalist*, (Oct. 1885).—Austral.
muricatum, Span. in Linnaea, xv. (1841) 339 = Ipomoea chryseides.
reniformis, Hassk. Pl. Jav. Rar. 524 = Ipomoea reniformis.
urceolatus, F. Muell. Census, 94 = Fitzalani.
Wallichii, *Choisy, Conv. Or.* 61.—Ind. or.

LEPISTEMUM, Reichb. Nom. 113 (1841) = praec.

LEPISTOMA, Blume, Fl. Jav. Praef. 7 (1828) = **Cryptolepis**, R. Br. (Asclep.).
javanicum, Blume, ex Steud. Nom. ed. II. ii. 29 = C. javanica.

LEPITOMA, Torr. ex Steud. Nom. ed. II. ii. 29 (1841) = **Pleuropogon**, R. Br. (Gramin.).
brevifolia, Torr. ex Steud. l. c. = P. Douglassii.

LEPIURUS, Dum. Obs. Gram. Belg. 140 (1823) = **Lepturus**, R. Br. (Gramin.).

LEPOSMA, Blume, Bijdr. 1049 (1826) = **Cryptolepis**, R. Br. (Asclep.).
javanicum, Blume, l. c. = C. javanica.

LEPSIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 123; et in Abh. Akad. Berl. (1855) 61. t. 5 A = **Begonia**, Linn.
foliosa, Klotzsch, in Abh. Akad. Berl. (1855) 62 = B. foliosa.
microphylla, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 123 = B. microphylla.
Poeppigiana, Klotzsch, in Abh. Akad. Berl. (1855) 63 = B. Poeppigiana.

LEPTA, Lour. Fl. Cochinch. 82 (1790) = **Evodia**, Forst. (Rutac.).
triphylla, Lour. l. c. = E. triphylla.

LEPTACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75, 169 (1832) = **Strobilanthes**, Blume (Acanth.).
alatus, Wight, Ic. iv. t. 1527 = S. perfoliatus.
helicoides, Nees, in DC. Prod. xi. 170 = S. helicoides.
rubicundus, Nees, in Wall. Pl. As. Rar. iii. 90 = S. rubicundus.
Walkeri, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 313 = S. pulcherrimus.
Walkeri, Wight, Ic. t. 1507 = S. amabilis.

LEPTACTINA, Hook. f. Ic. Pl. t. 1092 (1871). *RUBIACEAE*, Benth. & Hook. f. ii. 85.
densiflora, *Hook. f. Ic. Pl.* xi. 74, sub t. 1092.—Afr. trop.
heinsioides, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 88.—Afr. trop.
involucrata, *Hook. f. Ic. Pl.* iv. 74, sub t. 1092.—Afr. trop.
Mannii, *Hook. f.* t. 1092.—Afr. trop.
senegambica, *Hook. f.* xi. 74.—Afr. trop.
tetraloba, *N. E. Br. in Gard. Chron.* (1885) II. 391.—Afr. trop.

LEPTACTINIA, Hook. f. in Benth. & Hook. f. Gen. ii. 85 (1873) sphalm. = praec.

LEPTADENIA, R. Br. in Mem. Wern. Soc. i. (1809) 34. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 778.
abyssinica, *Decne. in DC. Prod.* viii. 628.—Abyssin.
appendiculata, Decne. in DC. Prod. viii. 628 = reticulata.
Bojeriana, *Decne. in DC. Prod.* viii. 629.—Madag.
brevipes, Wight, ex Hook. f. Fl. Brit. Ind. iv. 63 = reticulata.
cordata, Hochst. ex Decne. in DC. Prod. viii. 628 = lancifolia.
Delilei, Decne. l. c. = heterophylla.

LEPTADENIA:—
elliptica, *Blume, Bijdr.* 1066.—Java.
Forskålii, *G. Don, Gen. Syst.* iv. 124; *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 269.—Arab.
gracilis, Decne. l. c. = L. Spartum.
hastata, Decne. in DC. Prod. viii. 551; Vatke, in Linnaea, xl. (1876) 217 = lancifolia.
heterophylla, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 270.—Oriens.
imberbis, Wight, Contrib. 48 = reticulata.
Jacquemontiana, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 270 = L. Spartum.
lancifolia, *Decne. l. c.* 269.—Afr. trop.
madagascariensis, *Decne. in DC. Prod.* viii. 629.—Madag.
pallida, *Hochst. ex Decne. l. c.*—Abyssin.
pyrotechnica, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 269 = L. Spartum.
reticulata, *Wight, Contrib.* 47.—Ind. or.; Burma.
Spartum, *Wight, l. c.* 48.—Ind. bor. occ.; Arab.; Afr. bor.
visciformis, *Vatke, in Oestr. Bot. Zeitschr.* xxvi. (1876) 146.—Afr. trop.

LEPTALEA, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1835) 102 = **Facelis**, Cass. (Compos.).
apiculata, D. Don, l. c. = F. apiculata.

LEPTALEUM, DC. Syst. ii. 510 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 81.
filifolium, *DC. l. c.* 511.—Sibir.
pygmaeum, DC. l. c. = filifolium.

LEPTALIUM, Sweet, Hort. Brit. ed. III. 38 (1839) = praec.

LEPTALIX, Rafin. New Fl. Am. iii. 93 (1836) = **Fraxinus**, Tourn. (Oleac.).
acuminata, Rafin. l. c. = F. americana.
alba, Rafin. Alsog. Am. 32 = F. americana.
atrovirens, Rafin. l. c. 37 = F. excelsa.
canaliculata, Rafin. l. c. 35 = F. canaliculata.
caroliniana, Rafin. New Fl. Am. iii. 93 = F. platycarpa.
cinerea, Rafin. Alsog. Am. 32 = F. cinerea.
cordata, Rafin. l. c. 34 = F. cordata.
elliptica, Rafin. l. c. 32 = F. pubescens.
epiptera, Rafin. l. c. 33 = F. americana.
expansa, Rafin. l. c. = F. viridis.
fusca, Rafin. l. c. = F. fusca.
glauca, Rafin. l. c. = F. glauca.
grandifolia, Rafin. l. c. 38 = F. grandifolia.
juglandifolia, Rafin. New Fl. Am. iii. 93 = F. americana.
lacera, Rafin. Alsog. Am. 38 = F. lacera.
lancifolia, Rafin. l. c. 35 = F. americana.
longifolia, Rafin. l. c. 32 = F. pubescens.
media, Rafin. l. c. 36 = F. media.
mixta, Rafin. l. c. 32 = F. parvifolia.
mollis, Rafin. l. c. 34 = F. mollis.
multifolia, Rafin. l. c. 37 = F. multifolia.
nana, Rafin. l. c. 33 = F. excelsior.
nigra, Rafin. l. c. = F. nigra.
obliqua, Rafin. l. c. 38 = F. oxyphylla.
ovata, Rafin. l. c. 32 = F. pubescens.
pallida, Rafin. l. c. 33 = F. platycarpa.
palustris, Rafin. l. c. 35 = F. palustris.
pannosa, Rafin. l. c. 34 = F. pannosa.
platycarpa, Rafin. l. c. 35 = F. platycarpa.
pleurodes, Rafin. l. c. = F. pleurodes.
porosa, Rafin. l. c. 34 = F. porosa.
pubescens, Rafin. New Fl. Am. iii. 93 = F. pubescens.
pulverulenta, Rafin. Alsog. Am. 32 = F. pulverulenta.
quadrangulata, Rafin. l. c. 37 = F. quadrangulata.
repanda, Rafin. l. c. = F. repanda.
reticulata, Rafin. l. c. 34 = F. reticulata.
retusa, Rafin. l. c. = F. retusa, *Rafin.*
Richardi, Rafin. l. c. 32 = F. Richardi.
rubicunda, Rafin. l. c. = F. rubicunda.
rufa, Rafin. l. c. 33 = F. rufa.
teres, Rafin. l. c. 36 = F. teres.
undulata, Rafin. l. c. 37 = F. undulata.
viridis, Rafin. l. c. 32 = F. viridis.

LEPTAMINIUM, Steud. Nom. ed. II. ii. 29 (1841) = seq.

LEPTAMNIUM, Rafin. in Am. Monthly Mag. (1818) 267 = **Epifagus**, Nutt. (Orobanch.).
virginianum, Rafin. l. c. = E. americanus.

LEPTANDRA, Nutt. Gen. Am. i. 7 (1818) = **Veronica**, Tourn. (Scrophul.).
alba, Rafin. Med. Fl. ii. 21 = V. virginica.
angustifolia, Lehm. Del. Sem. Hort. Hamb. (1839). —Cf. Linnaea, xiv. (1840) Litt. 130 = V. tubiflora.
caerulea, Rafin. Med. Fl. ii. 21, in obs. = V. virginica.
japonica, Rafin. l. c. = V. virginica?
Meyeri, G. Don, Gen. Syst. iv. 579 = V. tubiflora.
oppositifolia, Rafin. New Fl. Am. ii. 21 = V. virginica.
purpurea, Rafin. Med. Fl. ii. 20 = V. virginica.
sibirica, Nutt. Gen. Am. i. 7 = V. virginica.
tubiflora, Fisch. & Mey. in Ann. Sc. Nat. Sér. II. v. (1836) 301 = V. tubiflora.
villosa, Rafin. Med. Fl. ii. 21 = V. virginica.
virginica, Nutt. Gen. Am. i. 7 = V. virginica.

LEPTANTHE, Klotzsch, in Bot. Ergeb. Waldem. Reise, 95. t. 63 (1862) = **Macrotomia**, DC. (Boragin.).
macrostachya, Klotzsch, l. c. = M. Benthami.

LEPTANTHES, Wight, ex Wall. Cat. sub n. 5048 (1831) = **Hydrilla**, Rich. (Hydroch.).
verticillatus, Wight, l. c. = H. verticillata.

LEPTANTHUS, Michx. Fl. Bor. Am. i. 24. t. 5 (1803) = **Heteranthera**, Ruiz & Pav. (Ponted.).
gramineus, Michx. l. c. 25 = H. graminea.
ovalis, Michx. l. c. = H. limosa.
peruvianus, Pers. Syn. i. 56 = H. reniformis.
reniformis, Michx. Fl. Bor. Am. i. 24 = H. reniformis.
virginicus, Pers. Syn. i. 56 = H. reniformis.

LEPTARGYREIA, Schlecht. in DC. Prod. xiv. 607 (1857) = **Shepherdia**, Nutt. (Elaeagn.).

LEPTARRHENA, R. Br. Chlor. Melv. 15 (1824); in Parry's First Voy. Suppl. 273. *SAXIFRAGEAE*, Benth. & Hook. f. i. 634.
amplexifolia, *R. Br. l. c.*—Am. arct.
inundata, Behr, ex Kellogg, in Proc. Calif. Acad. i. (1855) 45 = Saxifraga peltata.
micrantha, 'Fisch.; Rafin. Fl. Tellur. ii. 72'; ex S. Wats. Bibl. Ind. N. Am. Bot. 328 = amplexifolia.
pyrolifolia, Ser. in DC. Prod. iv. 48 = amplexifolia.

LEPTASEA, Haw. Saxifr. Enum. 39 (1821) = **Saxifraga**, Tourn.
aizoides, Haw. l. c. 40 = S. aizoides.
mutata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 385 = S. mutata.
tricuspidata, Haw. Saxifr. Enum. 40 = S. tricuspidata.

LEPTASPIS, R. Br. Prod. 211 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1110.
Banksii, *R. Br. l. c.* 211.—Austral.
cochleata, *Thw. Enum. Pl. Zeyl.* 357.—Zeylan.
Cumingii, *Steud. Syn. Pl. Gram.* 416.—Ins. Philipp.
lanceolata, *Zoll. Syst. Verz. Ind. Archip.* 53.—Java.
manillensis, Steud. Syn. Pl. Gram. 8 = urceolata.
umbrosa, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 326.—N. Caled.
urceolata, *R. Br. in Benn. Pl. Jav. Rar.* 23.—Ind. or.; Malaya.
zeylanica, Nees, ex Steud. Syn. Pl. Gram. 8 = urceolata.

LEPTATHERUM, Nees, in Proc. Linn. Soc. i. (1841) 92 = **Pollinia**, Trin. (Gramin.).
japonicum, Franch. & Sav. Enum. Pl. Jap. ii. 609 = P. nuda.
Royleanum, Nees, in Proc. Linn. Soc. i. (1841) 93 = P. nuda.

LEPTAULUS, Benth. in Benth. & Hook. f. Gen. i. 351, 996 (1862). *OLACINEAE*, Benth. & Hook. f. i. 351.
citrioides, *Baill. Adansonia*, iii. (1862–63) 375.—Madag.
daphnoides, *Benth. in Benth. & Hook. f. Gen.* i. 351.—Afr. trop.

LEPTAXIS, Rafin. Fl. Tellur. ii. 75 (1836) = **Tolmiea**, Torr. & Gray (1840) (Saxifr.).
Menziesi, Rafin. l. c. 76 = T. Menziesii.

LEPTEIRIS, Rafin. New Fl. Am. ii. 73 (1836) = **Pentstemon**, Mitch. (Scrophul.).
parviflora, Rafin. l. c. = P. confertus.

LEPTEMON, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Crotonopsis**, Michx. (Euphorb.).
lineare, Rafin. Sylva Tellur. 67 = C. linearis.

LEPTERANTHUS, Neck. Elem. i. 73 (1790) = **Centaurea**, Linn. (Compos.).
acutifolius, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 96 = C. pectinata.
comatus, Fourr. l. c. = C. pectinata.
fuscatus, Fourr. l. c. = C. pectinata.
hygrometicus, Cass. in Dict. Sc. Nat. xxvi. 65 = C. phrygia.
incoronatus, Cass. l. c. 66 = C. flosculosa.
linifolius, Webb, Iter Hisp. 32 = C. linifolia.
pectinatus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 96 = C. pectinata.
rufescens, Fourr. l. c. = C. pectinata.
supinus, Fourr. l. c. = C. pectinata.

LEPTICA, E. Mey. ex DC. Prod. vii. 16 (1838) = **Gerbera**, Gronov. (Compos.).
ciliata, E. Mey. ex DC. l. c. 17 = G. piloselloides.
sinuata, E. Mey. ex Steud. Nom. ed. II. ii. 29 = G. piloselloides.

LEPTIDIUM, Presl, Bot. Bemerk. 49 (1844); in Abh. Boehm. Ges. V. iii. (1845) 479 = LEPTIS, E. Mey. = **Lotononis**, Eckl. & Zeyh. (Legumin.).
brachyolobum, Presl, ll. cc. = Loton. brachyloba.
calycinum, Presl, ll. cc. = Loton. calycina.
debile, Presl, ll. cc. = Loton. porrecta.
divaricatum, Presl, ll. cc. = Loton. calycina.
erubescens, Presl, ll. cc. = Loton. pumila.
falcatum, Presl, ll. cc. = Loton. falcata.
filicaule, Presl, ll. cc. = Loton. versicolor.
molle, Presl, ll. cc. = Loton. mollis.
tenellum, Presl, ll. cc. = Loton. tenella.
versicolor, Presl, ll. cc. = Loton. versicolor.

LEPTILIX, Rafin. Neogent. 3 (1825) = **Tofieldia**, Huds. (Liliac.).

LEPTILON, Rafin. in Am. Monthly Mag. (1818) 268 = **Erigeron**, Linn. (Compos.).
divaricatum, Rafin. l. c. = E. divaricatus.

LEPTINELLA, Cass. in Bull. Soc. Philom. (1822) 127 = **Cotula**, Tourn. (Compos.).
acaenoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 325 = C. reptans?
bogotensis, DC. Prod. vi. 142 = Plagiocheilus bogotensis.
dioica, Hook. f. Fl. N. Zel. i. 129 = C. dioica.
Featherstonii, F. Muell. Veg. Pl. Chath. Isl. 27. t. 5 = C. Featherstonii.
Filicula, Hook. f. Fl. Tasm. i. 194. t. 51 B = C. Filicula.
intricata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 117 = C. reptans.
lanata, Hook. f. Fl. Antarct. i. 26 = C. lanata.
longipes, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 117 = C. reptans.
minor, Hook. f. Fl. N. Zel. i. 129 = C. minor.
multifida, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 118 = C. reptans.
pedunculariis, DC. Prod. vi. 141 = Plagiocheilus pedunculariis.
perpusilla, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 14.—Persia.
pinnata, Cass. in Bull. Soc. Philom. (1822) 127.—Am. austr.
plumosa, Hook. f. Fl. Antarct. i. 26. t. 20 = C. plumosa.
potentillina, F. Muell. Veg. Pl. Chath. Isl. 28. t. 6 = C. lanata?
propinqua, Hook. f. Fl. Antarct. i. 27.—Ins. Antarct.
pusilla, Hook. f. Fl. N. Zel. i. 129 = C. perpusilla.
scariosa, Cass. in Bull. Soc. Philom. (1822) 127 = C. reptans.
squalida, Hook. f. Fl. N. Zel. i. 129 = C. squalida.

LEPTIS, E. Mey. ex Eckl. & Zeyh. Enum. 174 (1836) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
argentea, Walp. Rep. ii. 837 = Loton. argentea.
brachygloba, Steud. Nom. ed. II. ii. 29 = seq.
brachyloba, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Loton. brachyloba.

LEPTIS:—
calycina, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Lotononis calycina.
debilis, Eckl. & Zeyh. Enum. 175 = Loton. porrecta.
divaricata, Eckl. & Zeyh. l. c. = Loton. calycina.
erubescens, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Loton. pumila.
falcata, Benth. l. c. = Loton. falcata.
filicaulis, Eckl. & Zeyh. Enum. 175 = Loton. versicolor.
Kraussiana, Walp. Rep. ii. 837 = Loton. versicolor.
mollis, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Loton. mollis.
prolifera, Eckl. & Zeyh. Enum. 175 = Loton. porrecta.
tenella, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Loton. tenella.
versicolor, Eckl. & Zeyh. Enum. 175 = Loton. versicolor.

LEPTOBAEA, Benth. in Benth. & Hook. f. Gen. ii. 1025 (1876). *GESNERACEAE*, Benth. & Hook. f. l. c.
CHAMPIONIA, C. B. Clarke, Comm. et Cyrt. Beng. 98. t. 68 (1874).
glabra, *C. B. Clarke, in DC. Monog. Phan.* v. 165.—Ind. or.
multiflora, *Gamble, Darjeeling List*, 58.—Reg. Himal.

LEPTOBASIS, Dulac, Fl. Hautes-Pyr. 198 (1867) = **Sisymbrium**, Tourn. (Crucif.).
pinnatipartita, Dulac, l. c. = S. tanacetifolia.

LEPTOBOTRYS, Baill. Etud. Gén. Euph. 478. t. 2 (1858) = **Tragia**, Plum. (Euphorb.).
discolor, Baill. l. c. = T. innocua.

LEPTOCALLIS, G. Don, Gen. Syst. iv. 260 (1837) = **Ipomoea**, Linn. (Convolv.).
armata, G. Don, in Sweet, Hort. Brit. ed. III. 482 = I. armata.
quinata, G. Don, Gen. Syst. iv. 260 = I. armata.
ternata, G. Don, l. c. = I. ternifolia.

LEPTOCARPAEA, DC. Syst. ii. 201 (1821) = **Sisymbrium**, Tourn. (Crucif.).
Loeselii, DC. l. c. 202 = S. Loeselii.

LEPTOCARPHA, DC. Prod. v. 495 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 361.
rivularis, *DC. l. c.*—Chili; Peruv.

LEPTOCARPHA, Endl. Gen. Suppl. i. 1383 in syn. (1841), sphalm. = LEPTOPHORA, Rafin. = **Helenium**, Linn. (Compos.).

LEPTOCARPUS, R. Br. Prod. 250 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1032.
CALOPSIS, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 44. t. 3.
SCHOENODUM, Labill. Nov. Holl. Pl. ii. 79. t. 229 (1806), partim.
aristatus, *R. Br. Prod.* 250.—Austral.
aristatus, F. Muell. Fragm. viii. 91 = erianthus.
Brownii, *Hook. f. Fl. Tasm.* ii. 73. t. 136.—Austral.
Burchellii, *Mast. in Journ. Linn. Soc.* x. (1869) 222.—Afr. austr.
canus, *Nees, in Ann. & Mag. Nat. Hist.* Ser. I. vi. (1841) 50.—Austral.
chilensis, *Mast. in DC. Monog. Phan.* i. 341.—Chili.
ciliaris, Nees, in Lehm. Pl. Preiss. ii. 64 = canus.
coangustatus, *Nees, l. c.* 65.—Austral.
desertus, F. Muell. Fragm. viii. 93 = spathaceus.
dichotomus, Heynh. Nom. i. 455 = *distachyos*?
disjunctus, *Mast. in Journ. Linn. Soc.* xvii. (1879) 344.—Cochinch.
distachyos, R. Br. Prod. 250 = Thamnochortus umbellatus.
elatior, *R. Br. l. c.* 250.—Austral.
erianthus, *Benth. Fl. Austral.* vii. 235.—Austral.
glaucus, Nees, in Lehm. Pl. Preiss. ii. 64 = Lepyrodia glauca.
imbricatus, R. Br. Prod. 250 = Thamnochortus imbricatus.
imbricatus, Sieber, ex Kunth, Enum. Pl. iii. 400 = Restio bifidus.
incurvatus, *Mast. in Journ. Linn. Soc.* x. (1869) 223.—Afr. austr.
modestus, *Mast. l. c.* 225.—Afr. austr.
neglectus, *Mast. l. c.*—Afr. austr.
oxylepis, *Mast. l. c.* 223.—Afr. austr.
paniculatus, *Mast. l. c.* 221.—Afr. austr.

LEPTOCARPUS:—
peronatus, *Mast. in Journ. Linn. Soc.* x. (1869) 224. —Afr. austr.
ramosus, *R. Br. Prod.* 250.—Austral.
scariosus, *R. Br. l. c.*—Austral.
Schultzii, *Benth. Fl. Austral.* vii. 237.—Austral.
setuligerus, F. Muell. Fragm. viii. 97 = tenax.
simplex, R. Br. Prod. 250, partim = Brownii.
simplex, *A. Rich. Fl. N. Zél.* 142.—N. Zel.
spathaceus, *R. Br. l. c.*—Austral.
squarrosus, [Nees, in] Sieber, ex Kunth, Enum. Pl. iii. 419 = Hypolaena lateriflora.
tenax, *R. Br. Prod.* 250.—Austral.
tenellus, F. Muell. Fragm. viii. 90 = aristatus.
thamnochortoides, F. Muell. l. c. 96 = tenax.

LEPTOCARPUS, Willd. ex Link, Jahrb. i. III. 51 (1820) = **Tamonea**, Aubl. (Verben.).
chamaedrifolius, Willd. l. c. = T. spicata.

LEPTOCARYDION, Hochst. ex Benth. & Hook. f. Gen. iii. 1176 (1883) = **Triodia**, R. Br. (Gramin.).

LEPTOCAULIS, Nutt. ex DC. Coll. Mém. v. 39. t. 10 (1829); et Prod. iv. 107 (1830) = **Apium**, Linn. (Umbell.).
diffusus, Nutt. ex DC. Prod. iv. 107 = seq.
divaricatus, DC. Coll. Mém. v. 39. t. 10 = A. divaricatum.
echinatus, Nutt. ex DC. Prod. iv. 107 = A. echinatum.
inermis, Nutt. ex DC. l. c. = A. patens?
patens, Nutt. ex DC. l. c. = A. patens.

LEPTOCERAS, Lindl. Swan River App. 53 (1839) = **Caladenia**, R. Br. (Orchid.).
fimbriata, Lindl. l. c. = C. fimbriata.
macrophylla, Lindl. Gen. et Sp. Orch. 416 = C. Menziesii.
Menziesii, Lindl. l. c. = C. Menziesii.
oblonga, Lindl. Swan River App. 53 = C. Menziesii.
pectinata, Endl. in Lehm. Pl. Preiss. ii. 6 = Lyperanthus nigricans.
pectinata, Lindl. Swan River App. 53 = C. fimbriata.
sulphurea, Lindl. Gen. et Sp. Orch. 416 = C. suaveolens.

LEPTOCERCUS, Rafin. in Am. Monthly Mag. (1819) 190 = **Lepturus**, R. Br. (Gramin.).

LEPTOCHLAENA, Spreng. Gen. i. 355 (1830) = **Leptolaena**, Thou. (Chlaenac.).

LEPTOCHLOA, Beauv. Agrost. 71. t. 15. f. 1 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1172.
LEPTOSTACHYS, G. F. W. Mey. Prim. Fl. Esseq. 73 (1818).
OXYDENIA, Nutt. Gen. Am. i. 76 (1818).
RABDOCHLOA, Beauv. Agrost. 84. t. 17. f. 3 (1812).
anoplia, Nees, in Linnaea, xix. (1847) 691 = Chloris anoplia.
arabica, Kunth, Rév. Gram. i. 91 = Dinebra arabica.
arenaria, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 29 = Diplachne arenaria.
attenuata, Steud. Syn. Pl. Gram. 209 = mucronata.
brachiata, *Steud. l. c.*—Ind. occ.
Burchellii, *Munro, ex Doell, in Mart. Fl. Bras.* ii. III. 94.—Bras.
calycina, *Kunth, Rév. Gram.* i. 91.—Ind. or.
capillacea, *Beauv. Agrost.* 71.—As. trop.; Austral.
chinensis, Nees, in Syll. Ratisb. i. (1824) 4; Agrost. Bras. 432 = capillacea.
coerulescens, *Steud. Syn. Pl. Gram.* 209.—Ind. or.
coromandelina. *Steud. l. c.*—Ind. or.
cynosuroides, *Roem. & Schult. Syst.* ii. 579.—Ind. or.
Digitaria, *Nees, Agrost. Bras.* 433.—Ecuador.
domingensis, *Trin. Fund. Agrost.* 133; *Link, Enum. Hort. Berol.* i. 103.—Am. austr.
dubia, Nees, in Syll. Ratisb. i. (1824) 4; Agrost. Bras. 433 = Diplachne dubia.
eragrostoides, *Steud. Syn. Pl. Gram.* 208.—Japon.
fascicularis, A. Gray, Man. Bot. N. U. St. ed. V. 623 = Diplachne fascicularis.
fascicularis, Griseb. ex Benth. in Journ. Linn. Soc. xix. (1881) 108 = Panicum sanguinale.
filiformis, *Beauv. Agrost.* 71.—Am. austr.; Ins. Phil.
filiformis, J. & C. Presl, Rel. Haenk. i. 288 = mucronata.

LEPTOCHLOA:—
filiformis, Roem. & Schult. Syst. ii. 580 = cynosuroides.
floribunda, *Doell, in Mart. Fl. Bras.* ii. III. 89.—Bras.
fusca, Kunth, Rév. Gram. i. 91 = Diplachne fusca.
gracilis, *Nees, in Syll. Ratisb.* i. (1824) 4 = Am. austr.
gracilis, Wight, ex Steud. Syn. Pl. Gram. 213 = uniflora.
grandiglumis, Nees, Fl. Afr. Austr. 252 (Diplachne sp.). —Afr. austr.
hirta, *Steud. Syn. Pl. Gram.* 209.—Ins. Galap.
humilis, Nees, ex Steud. l. c. 210 = Chloris anoplia.
imbricata, Thurb. in S. Wats. Bot. Calif. ii. 293 = Diplachne imbricata.
Langloisii, *Vasey, in Bull. Torrey Bot. Club.* xii. (1885) 7.—Am. bor.
laxa, Trin. ex Steud. Nom. ed. II. ii. 30 (Quid?). —Hab.?
Liebmannii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 565.—Mexic.
Lindleyana, *Kunth, Enum. Pl.* i. 525.—Ins. Galap.
loliiformis, F. Muell. Fragm. viii. 128 = Diplachne loliiformis.
longa, *Griseb. Fl. Brit. W. Ind.* 538.—Ind. occ.
mollis, *Kunth, Rév. Gram.* ii. 443. t. 135.—Afr. trop.
monostachya, *Roem. & Schult. Syst.* ii. 580.—Ind. or.
mucronata, *Kunth, Rév. Gram.* i. 91.—Am. bor.
mutica, Steud. Syn. Pl. Gram. 208 = virgata.
Nealleyi, *Vasey, in Bull. Torrey Bot. Club*, xii. (1885) 7.—Am. bor.
Neesii, *Benth. in Journ. Linn. Soc.* xix. (1881) 108. —Zeylan.
obtusiflora, Trin. ex Steud. Nom. ed. II. ii. 30 = Diplachne dubia.
obtusifolia, *Hochst. in Flora*, xxxviii. (1855) 203.— Abyss.
panicoides, *Wight, ex Steud. Syn. Pl. Gram.* 62.— Ind. or.
paniculata, *Fourn. in Bull. Soc. Bot. Fr.* xxvii. (1880) 296.—Nicaragua.
patens, *Kunth, Enum. Pl.* i. 271.—Chili.
pectinata, *Kunth, Rév. Gram.* i. 91.—Ind. or.
pellucida, Steud. Syn. Pl. Gram. 209 = mucronata.
phleoidea, Willk. & Lange, Prod. Fl. Hisp. i. 74, in syn. = Koeleria phleoides.
plumosa, Anderss. in Peters, Reise Mossamb. Bot. 557 = Diplachne alopecuroides.
polystachya, *Benth. Fl. Austral.* vii. 617.—Austral.
polystachya, Kunth, Rév. Gram. i. 91 = Diplachne fascicularis.
procera, *Nees, in Syll. Ratisb.* i. (1824) 2.—Am. austr.
racemosa, Kunth, Rév. Gram. i. 91 = Eleusine verticillata.
rigida, Munro, ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 336 = Diplachne rigida.
scabra, *Nees, Agrost. Bras.* 435.—Bras.
Schimperiana, *Hochst. in Flora*, xxxviii. (1855) 203.— Abyss.
setacea, *Hochst. l. c.* xxiv. (1841) 1. *Intell.* 20, nomen. —Abyss.
stricta, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 566. —Mexic.
stricta, Trin. ex Steud. Nom. ed. II. ii. 30, nomen (Quid?).—Hab.?
subdigitata, *Trin. ex Steud. Syn. Pl. Gram.* 210.— Austral.
tenerrima, Roem. & Schult. Syst. ii. 581 = capillacea.
tetraquetra, *J. & C. Presl, Rel. Haenk.* i. 288.—Ins. Philipp.
uniflora, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 409.— Abyss.
uniflora, Trin. ex Steud. Nom. ed. II. ii. 463, in syn. = Diplachne fascicularis.
verticillata, Kunth, Rév. Gram. i. 91 = Eleusine verticillata.
virgata, *Beauv. Agrost.* 71.—Am. austr.
virgata, Spreng. Syst. i. 350 = domingensis.
virgata, *Wight, ex Steud. Syn. Pl. Gram.* 213.— Ind. or.
Virletii. *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 566. —Mexic.
Wightiana, Nees, ex Steud. Syn. Pl. Gram. 209 (Eragrostidis sp.?).—Ind. or.

LEPTOCLADUS, Oliver, in Journ. Linn. Soc. viii. (1864) 160. t. 12 = **Mostuea**, Didr. (Logan.).
Thomsoni, Oliver, l. c. = M. Thomsoni.

LEPTOCLINIUM, Benth. in Benth. & Hook. f. Gen. ii. 244 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 244.
fruticosum, A. Gray, in Proc. Am. Acad. xv. (1880) 48 = Garberia fruticosa.
trichotomum, *Benth. ex Baker, in Mart. Fl. Bras.* vi. II. 272.—Bras.

LEPTOCNEMIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 624 (1840) = **Cymopterus**, Rafin. (Umbellif.).
campestris, Nutt. ex Torr. & Gray, l. c. = C. campestris.

LEPTOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 193. t. 57 (1856) = **Pouzolzia**, Gaudich. (Urtic.).
borbonica, Blume, l. c. 194 = P. viminea.
heterophylla, Blume, l. c. = P. heterophylla.
rubricaulis, Blume, l. c. = P. rubricaulis.

LEPTOCODON, Lem. Illustr. Hortic. iii. (1856) Misc. 49; Hook. f. & Tho ms. in Journ. Linn. Soc. ii. (1858) 17. *CAMPANULACEAE*, Benth. & Hook. f. ii. 557.
gracilis, *Lem. l. c.*—Reg. Himal.

LEPTOCODON, Sond. in Fl. Harv. & Sond. Fl. Cap. iii. 584 (1865) = **Treichelia**, Vatke (Campan.).
longebracteatum, Sond. l. c. 585 = T. longebracteata.

LEPTOCOMA, Less. in Linnaea, vi. (1831) 130 = **Rhynchospermum**, Reinw. (Compos.).
racemosa, Less. l. c. = R. verticillatum.

LEPTOCORYPHIUM, Nees, Agrost. Bras. 83 (1829) = **Anthaenantia**, Beauv. (Gramin.).
Drummondi, C. Muell. in Bot. Zeit. xix. (1861) 314.— Am. bor.
lanatum, Nees, Agrost. Bras. 84 = A. lanata.
molle, Nees, l. c. 85 = A. lanata.
obtusum, Steud. Syn. Pl. Gram. 34.—Am. bor.
penicilligerum, Speg. in Anal. Soc. Cient. Argent. xvi. (1883) 102.—Reg. Argent.

LEPTOCYAMUS, Benth. in Trans. Linn. Soc. xviii. (1839) 209, adnot. = **Glycine**, Linn. (Legumin.).
clandestinus, Benth. in Hook. f. Fl. Tasm. i. 102 = G. clandestina.
elongatus, Benth. in Trans. Linn. Soc. xviii. (1839) 209 = G. tabacina.
latifolius, Benth. in Mitch. Journ. Trop. Austral. 361 = G. tabacina.
microphyllus, Benth. in Trans. Linn. Soc. xviii. (1839) 209 = G. clandestina.
sericeus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 40 = G. sericea.
tasmanicus, Benth. in Hook. f. Fl. Tasm. i. 102. t. 17 = G. Latrobeana.
tomentosus, Benth. in Trans. Linn. Soc. xviii. (1839) 209 = G. tomentosa.

LEPTOCYTISUS, Meissn. in Lehm. Pl. Preiss. ii. 211 (1847) = **Latrobea**, Meissn. (Legumin.).
hirtellus, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 258 = Latr. hirtella.
tenellus, Meissn. in Lehm. Pl. Preiss. ii. 211 = Latr. tenella.

LEPTODACTYLON, Hook. & Arn. Bot. Beech. Voy. 369. t. 89 (1841) = **Gilia**, Ruiz & Pav. (Polem.).
caespitosum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 157 = G. pungens.
californicum, Hook. & Arn. Bot. Beech. Voy. Suppl. 369 = G. californica.
Hookeri, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 157 = G. pungens.
pungens, Nutt. l. c. = G. pungens.

LEPTODAPHNE, Nees, Laurin. Expos. 16 (1833) = **Ocotea**, Aubl. (Laurin.).
Pseudopsychotria, Nees, ex Meissn. in Mart. Fl. Bras. v. II. 238 = O. Sieberi.
subalpina, Nees, Syst. Laurin. 361.—Bras.
tenuiflora, Nees, l. c. 359.—Bras.

LEPTODERMIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 191 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 135.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 198.—Reg. Himal.
lanceolata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 191. —Reg. Himal.

LEPTODERMIS:—
oblonga, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 108.—China.
scabrida, *Hook. f. Fl. Brit. Ind.* iii. 199.—Reg. Himal.
virgata, *Edgew. ex Aitch. Cat. Punjab Pl.* 70; *et ex Hook. f. Fl. Brit. Ind.* iii. 198.—Reg. Himal.

LEPTODESMIA, Benth. in Benth. & Hook. f. Gen. i. 522 (1875). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
congesta, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 152.—Ind. or.

LEPTOGLOSSIS, Benth. Bot. Voy. Sulph. 143 (1844). *SOLANACEAE*, Benth. & Hook. f. ii. 908.
CYCLOSTIGMA, Phil. Sert. Mendoc. Alt. 39 (1871).
Coulteri, A. Gray, in Proc. Am. Acad. xii. (1877) 165 = Nierembergia Coulteri.
linifolia, *Benth. & Hook. f. Gen.* ii. 909.—Chili.
schwenkiodes, *Benth. Bot. Voy. Sulph.* 143.—Peruv.
tenuis, *Benth. & Hook. f. Gen.* ii. 909.—Chili.
texana, A. Gray, in Proc. Am. Acad. xii. (1877) 164 = Nierembergia viscosa.

LEPTOGLOTTIS, DC. Mém. Lég. 451 (1825) = **Schrankia**, Willd. (Legumin.).
Nuttallii, DC. l. c. = S. uncinata.

LEPTOGONUM, Benth. in Benth. & Hook. f. Gen. iii. 103 (1880). *POLYGONACEAE*, Benth. & Hook. f. l. c.
domingensis, *Benth. l. c.* 104.—Ins. S. Doming.

LEPTOGYNE, Less. in Linnaea, vi. (1831) 151 = **Cotula**, Tourn. (Compos.).
rosea, Less. l. c. 152 = C. rosea.

LEPTOLAENA, Thou. Hist. Vég. Afr. Austr. 41. t. 11 (1807). *CHLAENACEAE*, Benth. & Hook. f. i. 195.
LEPTOCHLAENA, Spreng. Gen. i. 355 (1830).
multiflora, *Thou. Hist. Vég. Afr. Austr.* 41. t. 11.—Madag.
pauciflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 96.—Madag.
turbinata, *Baker, l. c.* 97.—Madag.

LEPTOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 124 = **Glycine**, Linn. (Legumin.).
clandestinum, Benth. l. c. 125 = G. clandestina.
elongatum, Benth. l. c. = G. tabacina.
microphyllum, Benth. l. c. = G. clandestina.
tabacinum, Benth. l. c. = G. tabacina.
tomentosum, Benth. l. c. = G. tomentosa.

LEPTOLOBIUM, Vog. in Linnaea, xi. (1837) 388 = **Sweetia**, Spreng. (Legumin.).
aurantiacum, Mart. Herb. Fl. Bras. n. 1149 = Diptychandra aurantiaca.
bijugum, Vog. in Linnaea, xi. (1837) 391 = S. bijuga.
costulatum, Miq. Stirp. Surin. Sel. 17 (Ormosiae sp.).—Guian.
dasycarpum, Vog. in Linnaea, xi. (1837) 388 = S. dasycarpa.
elegans, Vog. l. c. 390 = S. elegans.
glabrifolium, Tul. in Arch. Mus. Par. iv. (1844) 119 = S. dasycarpa.
lanceolatum, Tul. l. c. 118 = S. dasycarpa.
leiocarpum, Vog. in Linnaea, xi. (1837) 393 = Apuleia praecox.
luteum, Mart. Herb. Fl. Bras. n. 1148 = Sclerolobium aureum.
nitens, Vog. in Linnaea, xi. (1837) 394 = S. nitens.
nitidulum, Miq. Stirp. Surin. Sel. 18 = S. nitens.
punctatum, Benth. in Linnaea, xxii. (1849) 526 = Myrocarpus frondosus.
tenuifolium, Vog. l. c. xi. (1837) 390 = S. tenuifolia.
tortum, Mart. Herb. Fl. Bras. n. 1151 = S. dasycarpa.
tortuosum, Mart. ex Benth. in Journ. Linn. Soc. viii. (1865) 262 = S. glabrifolia.

LEPTOMERIA, R. Br. Prod. 353 (1810). *SANTALACEAE*, Benth. & Hook. f. iii. 229.
acerba, R. Br. l. c. 354 = Omphacomeria acerba.
acerba, A. Cunn. ex A. DC. in DC. Prod. xiv. 680 = Omphacomeria psilotoides.
acerba, Sieber, ex Benth. Fl. Austral. vi. 229 = Exocarpus cupressiformis.

LEPTOMERIA:—
acida, *R. Br. Prod.* 353.—Austral.
aphylla, *R. Br. l. c.* 354.—Austral.
aphylla, A. DC. in DC. Prod. xiv. 677 = pauciflora.
aphylla, Sieber, ex Benth. Fl. Austral. vi. 219 = Choretrum lateriflorum.
axillaris, *R. Br. Prod.* 354.—Austral.
Billardieri, Sieber, ex Benth. Fl. Austral. vi. 219 = Choretrum Candollei.
Billardierii, *R. Br. Prod.* 354.—Austral.
Brownii, Miq. in Lehm. Pl. Preiss. i. 612 = squarrulosa.
chrysadena, Miq. l. c. = scrobiculata.
Cunninghamii, *Miq. l. c.* 611.—Austral.
empetriformis, *Miq. l. c.* 610.—Austral.
ericoides, Miq. l. c. 611 = scrobiculata.
glomerata, *F. Muell. ex Hook. f. Fl. Tasm.* ii. 370.—Austral.
hirtella, Miq. in Lehm. Pl. Preiss. i. 610 = empetriformis.
laxa, *Miq. l. c.* 612.—Austral.
Lehmanni, Miq. l. c. 614 = pauciflora.
obovata, *Miq. l. c.* 613.—Austral.
odorata, Miq. l. c. = Exocarpus odoratus.
pauciflora, *R. Br. Prod.* 354.—Austral.
Preissiana, *A. DC. in DC. Prod.* xiv. 678.—Austral.
pungens, F. Muell. in Trans. Vict. Inst. i. (1855) 41 = aphylla.
scrobiculata, *R. Br. Prod.* 354.—Austral.
spinosa, *A. DC. in DC. Prod.* xiv. 678.—Austral.
squarrulosa, *R. Br. Prod.* 354.—Austral.
xiphoclada, Sieber, ex Spreng. Syst. iv. Cur. Post. 109 = Amperea spartioides.

LEPTOMON, Steud. Nom. ed. II. i. 447, in syn. (1840) = LEPTEMON, Rafin. = **Crotonopsis**, Michx. (Euphorb.).

LEPTONEMA, Hook. Ic. Pl. t. 692. (1844) = **Dolichostylis**, Turcz. (Crucif.).
Lindeni, Hook. l. c. = D. Lindenii.

LEPTONEMA, A. Juss. Euph. Tent. 19. t. 4 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 275.
melanthesoides, F. Muell. in Hook. Kew Journ. ix. (1857) 17 = Fluggea microcarpa.
venosum, *A. Juss. Euph. Tent.* 19. t. 4. f. 12.—Madag.

LEPTONIUM, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 236 = **Lepionurus**, Blume (Olacin.).
oblongifolium, Griff. l. c. 237 = Lepionurus oblongifolius.

LEPTONYCHIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 222. *STERCULIACEAE*, Benth. & Hook. f. i. 983.
BINNENDYKIA, Kurz, in Tijdschr. Nederl. Ind. xxviii. (1865) 164.
acuminata, *Mast. in Hook. f. Fl. Brit. Ind.* i. 379.—Malacca.
glabra, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 222.—Burma; Malaya.
heteroclita, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 67 = glabra.
lanceolata, *Mast. in Oliver, Fl. Trop. Afr.* i. 239.—Afr. trop.
moacurroides, *Bedd. Fl. Sylvat.* i. t. 114.—Ind. or.
urophylla, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* i. 238.—Afr. trop.

LEPTOPAETIA, Harv. Gen. Pl. S. Afr. ed. II. f. 231 (1868) = **Tacazzea**, Decne. (Asclep.).

LEPTOPETALUM, Hook. & Arn. Bot. Beech. Voy. 295 (1841) = **Hedyotis**, Linn. (Rubiac.).
mexicanum, Hook. & Arn. l. c. = H. leptopetala.

LEPTOPETION, Schott, Gen. Aroid. t. 8 (1856) = **Biarum**, Schott (Aroid.).
alexandrinum, Schott, l. c. = B. Olivieri.

LEPTOPHOBA, Ehrh. Beitr. iv. 146 (1789) = **Aira**, Linn. (Gramin.).

LEPTOPHOENIX, Becc. in Ann. Jard. Buitenz. ii. (1885) 82. *PALMAE*.
affinis, *Becc. l. c.*—N. Guin.
pinangoides, *Becc. l. c.*—N. Guin.

LEPTOPHORA, Rafin. in Am. Monthly Mag. (1819) 195 = **Helenium**, Linn. (Compos.)

LEPTOPHRAGMA, Benth. ex Dun. in DC. Prod. xiii. i. 578 (1852) = **Petunia**, Juss. (Solan.).
montevidensis, Benth. ex Dun. l. c. 579 = P. parviflora?
prostrata, Benth. ex Dun. l. c. 578 = P. parviflora.

LEPTOPHRAGMA, R. Br. ex Benn. Pl. Jav. Rar. 185 (1840?). *MELIACEAE*?—Cf. Benth. & Hook. f. Gen. i. 330.

LEPTOPHYLLUM, Ehrh. Beitr. iv. 147 (1784) = **Arenaria**, Rupp. (Caryoph.).

LEPTOPHYTUS, Cass. in Bull. Soc. Philom. (1817) 11; et in Dict. Sc. Nat. xxvi. 77 (1823) = **Leysera**, Linn. (Compos.).
leyseroides, Cass. in Dict. Sc. Nat. xxiii. 560; xxvi. 78 = Leysera capillifolia.

LEPTOPODA, Nutt. Gen. Am. ii. 174 (1818) = **Helenium**, Linn. (Compos.).
acaulis, DC. Prod. v. 663 = Actinella acaulis.
brachypoda, Torr. & Gray, Fl. N. Am. ii. 388 = H. nudiflorum.
brevifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 373 = H. brevifolium.
decurrens, Macbr. in Ell. Sketch, ii. 446 = H. Nuttallii.
denticulata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 373 = H. Nuttallii.
fimbriata, Torr. & Gray, Fl. N. Am. ii. 387 = H. fimbriatum.
floridana, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
helenioides, Cass. in Dict. Sc. Nat. xxvi. 80 = H. Nuttallii.
Helenium, Nutt. Gen. Am. ii. 174 = H. Nuttallii.
incisa, Torr. & Gray, Fl. N. Am. ii. 387 = H. Nuttallii.
integrifolia, M. A. Curt. ex Torr. & Gray, l. c. = H. Curtisii.
pinnatifida, Schweinitz, ex Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 372 = H. vernale.
puberula, Macbr. in Ell. Sketch, ii. 445 = H. vernale.

LEPTOPTERIS, Blume, Mus. Bot. Lugd. Bat. i. 240. t. 34 (1850) = **Gelsemium**, Juss. (Logan.).
sumatrana, Blume, l. c. = G. elegans.

LEPTOPUS, Decne. in Jacquem. Voy. Bot. 155. t. 156 (1844) = **Andrachne**, Linn. (Euphorb.).
cordifolius, Decne. l. c. = A. cordifolius.

LEPTOPUS, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 249 = **Euphorbia**, Linn.
adiantoides, Klotzsch & Garcke, l. c. = E. adianthoides.
brasiliensis, Klotzsch & Garcke, l. c. = E. sciadophila.
Hartwegii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 46 = E. adianthoides.
ocymoides, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 249 = E. ocymoides.
orinocensis, Klotzsch & Garcke, l. c. 250.—Bras.
Poeppigii, Klotzsch & Garcke, l. c. = E. Poeppigii.
segoviensis, Klotzsch & Garcke, l. c. = E. segoviensis.

LEPTOPYRUM, Rafin. in Med. Repos. N. York, v. (1808) 351 = **Avena**, ? Linn. (Gramin.).
tenellum, Rafin. l. c. (Quid?).—Am. bor.

LEPTOPYRUM, Reichb. Consp. 192 (1828) = **Isopyrum**, Linn. (Ranunc.).
fumarioides, Reichb. l. c. = I. fumarioides.

LEPTORACHIS, Baill. Etud. Gén. Euph. 495 (1858) = LEPTORHACHIS, Klotzsch = **Tragia**, Plum. (Euphorb.).

LEPTORDERMIS, DC. Prod. iv. 462 (1830) sphalm. = **Leptodermis**, Wall. (Rubiac.).

LEPTORHABDOS, Schrenk, in Fisch. & Mey. Pl. Schrenk. i. 23 (1841). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 971.
DARGERIA, Decne. in Jacquem. Voy. Bot. 115. t. 121 (1844).

LEPTORHABDOS:—
Benthamiana, *Walp. Rep.* iii. 387.—Reg. Himal.
brevidens, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 13.—Sibir. altaic.
linifolia, *Walp. Rep.* iii. 388.—Reg. Himal.
micrantha, *Schrenk, in Fisch. & Mey. Pl. Schrenk*, i. 24.—Soongaria.
parviflora, Benth. in DC. Prod. x. 510 = Benthamiana.
virgata, Benth. l. c. = Benthamiana.

LEPTORHACHIS, Klotzsch, in Wiegm. Archiv, vii. (1841) 189 = **Tragia**, Plum. (Euphorb.).
capensis, Muell. Arg. in DC. Prod. xv. II. 926 = T. capensis.
hastata, Klotzsch, in Wiegm. Archiv, vii. (1841) 189 = T. hastata.

LEPTORHACHYS, Meissn. Comm. 369 (1843) = praec.

LEPTORHOËS, C. B. Clarke, in Hemsl. Diagn. Pl. Nov. 55 (1880). *COMMELINACEAE*, Benth. & Hook. f. iii. 856.
filiformis, *C. B. Clarke, l. c.*—Mexic.

LEPTORHYNCHOS, Less. Syn. Comp. 273 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 307.
APHANORYNCHUS, Endl. Gen. 445 (1838).
DORATOLEPIS, Schlecht. in Linnaea, xx. (1847) 593.
RHYTIDANTHE, Benth. in Enum. Pl. Hueg. 63 (1837).
ambiguus, *Benth. Fl. Austral.* iii. 609.—Austral.
aureus, Benth. in Enum. Pl. Hueg. 64 = Waitzia aurea.
Baileyi, *F. Muell. Fragm.* x. 101.—Austral.
citrinus, Benth. in Enum. Pl. Hueg. 64 = Waitzia Steetziana.
elongatus, *DC. Prod.* vi. 160.—Austral.
gracilis, Walp. in Linnaea, xiv. (1840) 317 = squamatus.
hemisphaericus, DC. Prod. vi. 160 = squamatus.
Lhotskyanus, Walp. in Linnaea, xiv. (1840) 317 = squamatus.
linearis, *Less. Syn. Comp.* 273.—Austral.
medius, A. Cunn. in DC. Prod. vi. 160 = elongatus.
nitidulus, DC. Prod. vi. 160 = squamatus.
pannaetioides, *Benth. Fl. Austral.* iii. 609.—Austral.
Podolepis, DC. Prod. vi. 160 = Waitzia Podolepis.
pulchellus, *F. Muell. Fragm.* i. 53.—Austral.
Rhytidanthe, Benth. in DC. Prod. vi. 679 = elongatus.
squamatus, *Less. Syn. Comp.* 273.—Austral.
suaveolens, Benth. in Enum. Pl. Hueg. 64 = Waitzia nivea.
tenuifolius, *F. Muell. Fragm.* i. 52.—Austral.
Waitzia, *Sond. in Linnaea*, xxv. (1852) 501.—Austral.

LEPTORMUS, Eckl. & Zeyh. Enum. 8 (1835) = **Heliophila**, Linn. (Crucif.).
acuminatus, Eckl. & Zeyh. Enum. 8 = H. acuminata.
caledonicus, l. c. 9 = H. caledonica.
dissectus, Eckl. & Zeyh. l. c. 8 = H. dissecta.
longifolius, Eckl. & Zeyh. l. c. 9 = H. longifolia.
pendulus, Eckl. & Zeyh. l. c. = H. pendula.
rivalis, Eckl. & Zeyh. l. c. = H. rivalis.
tenellus, Eckl. & Zeyh. l. c. = H. dissecta.
trifidus, Eckl. & Zeyh. l. c. = H. trifida.
tripartitus, Eckl. & Zeyh. l. c. = H. Zeyheri.

LEPTORRHYNCHUS, F. Muell. Fragm. x. 101 (1877) = **Leptorhynchos**, Less. (Comp.).

LEPTOSCELA, Hook. f. in Benth. & Hook. f. Gen. ii. 59 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c. 59.
ruellioides, *Hook. f. Ic. Pl.* t. 1149.—Bras.

LEPTOSCHOENUS, Nees, in Hook. Journ. Bot. ii. (1840) 393 = **Rynchospora**, Vahl (Cyperac.).
prolifer, Nees, l. c.—Am. trop.

LEPTOSEMA, Benth. in Ann. Wien. Mus. ii. (1838) 84 = **Brachysema**, R. Br. (Legumin.).
bossiaeoides, Benth. l. c. = B. bossiaeoides.
Chambersii, F. Muell. Pl. Fitzalan. 8 = B. Chambersii.
oxyloboides, F. Muell. l. c. = B. oxyloboides.

LEPTOSERIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 438 = **Malacothrix**, DC. (Compos.).
californica, A. Gray, Syn. Fl. N. Am. i. II. 465 (Index), sphalm. = M. saxatilis.
saxatilis, A. Gray, l. c. = M. saxatilis.
sonchoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 439 = M. sonchoides.

LEPTOSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = **Silene**, Linn. (Caryoph.).
inaperta, Fourr. l. c. = S. inaperta.

LEPTOSIPHON, Benth. in Bot. Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polem.).
androsaceus, Benth. l. c. = G. androsacea.
bicolor, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 156 = G. tenella.
densiflorus, Benth. in Bot. Reg. sub t. 1622 = G. densiflora.
grandiflorus, Benth. l. c. = G. densiflora.
luteus, Benth. l. c. = G. micrantha.
parviflorus, Benth. l. c. = G. micrantha.
parviflorus, Hook. f. Bot. Mag. t. 5863 = G. androsacea.
roseus, Thomps. ex Gard. Chron. (1871) 1264 = Gilia micrantha.

LEPTOSKELA, Hook. f. Ic. Pl. t. 1149 (1873), lapsu = **Leptoscela**, Hook. f. (Rubiac.).

LEPTOSOLENA, Presl, Rel. Haenk. i. 111. t. 18 (1827). *SCITAMINEAE*, Benth. & Hook. f. iii. 647.
Haenkei, *Presl, l. c.*—Ins. Philipp.

LEPTOSOMUS, Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 174 = **Eichhornia**, Kunth (Ponted.).
natans, Schlecht. l. c. = E. natans.

LEPTOSPARTION, Griff. Notul. iv. 511 (1854); Ic. Pl. Asiat. t. 591 = **Duabanga**, Buch.-Ham. (Lythrar.).
grandiflora, Griff. l. c. = D. sonneratioides.

LEPTOSPERMUM, Forst. Char. Gen. 71. t. 36 (1776). *MYRTACEAE*, Benth. & Hook. f. i. 703.
FABRICIA, Gaertn. Fruct. i. 175. t. 35. f. 4 (1788).
GLAPHYRIA, Jack, in Trans. Linn. Soc. xiv. (1823) 128.
HOMALOSPERMUM, Schau. in Linnaea, xvii. (1843) 242.
MACKLOTTIA, Korth. in Nederl. Kruidk. Arch. i. (1847) 196.
PERICALYMMA, Endl. Gen. 1230 (1840).
abnorme, *F. Muell. ex Benth. Fl. Austral.* iii. 109.—Austral.
acerosum, Schau. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 250 = seq.?
aciculare, Schau. in Linnaea, xv. (1841). 429 = scoparium.
acuminatum, Wendl. ex Steud. Nom. ed. I. 473 = stellatum.
acutifolium, Hort. Vratisl. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 225 = flavescens.
alpestre, *Blume, Bijdr.* 1100.—Java.
ambiguum, Sm. in Trans. Linn. Soc. iii. (1797) 264 = Kunzea corifolia.
amboinense, Blume, Bijdr. 1100; DC. Prod. iii. 229 = flavescens.
angustifolium, Endl. in Enum. Pl. Hueg. 50 = Hypocalymna angustifolia.
Annae, *Stein, in Regel, Gartenfl.* (1885) 66.—Ins. Philipp.
aquaticum, Hort. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 243 = flavescens.
arachnoides, Gaertn. Fruct. i. 175. t. 35 = arachnoideum.
arachnoideum, Hort. Berol. ex Schau. in Linnaea, xv. (1841) 433 = scoparium.
arachnoideum, *Sm. in Trans. Linn. Soc.* iii. (1797) 263. —Austral.
attenuatum, *Sm. in Trans. Linn. Soc.* iii. (1797) 262. —Autsral.
australe, Salisb. Prod. 350 = pubescens.
baccatum, Schau. in Linnaea, xv. (1841) 433 = scoparium.
baccatum, Sm. in Trans. Linn. Soc. iii. (1797) 264 = arachnoideum.
Blumei, Steud. Nom. ed. II. ii. 31 = flavescens.
brevifolium, Rudge, in Trans. Linn. Soc. viii. (1807) 299. t. 14 = Baeckea brevifolia.
brevifolium, Schau. in Walp. Rep. ii. 168, sphalm. = *buxifolium*, Wendl. f.
brevipes, F. Muell. in Trans. Vict. Inst. i. (1855) 125 = attenuatum.
bullatum, Hort. ex Loud. Hort. Brit. Suppl. iii. 576 = scoparium.
buxifolium, Dehn. Revist. Napol. i. III. 172; ex Walp. Rep. ii. 169 (sp. dub.).—Austral.

LEPTOSPERMUM:—
buxifolium, Wendl. f. in Otto & Dietr. Allg. Gartenz. i. (1833) 186 = flavescens.
Candollei, Schau. in Linnaea, xv. (1841) 441 = pubescens.
canescens, Wendl. ex Steud. Nom. ed. I. 493 (sp. dub.). —Hab.?
Celsianum, *Tenore, Ind. Sem. Hort. Neap.* (1825) 11. n. 13.—Austral.?
ciliatum, Forst. Char. Gen. 72 = Metrosideros ciliata.
ciliolatum, Otto & Dietr. Allg. Gartenz. ix. (1841) 241 (sp. dub.).—Austral.
collinum, Forst. Char Gen. 72 = Metrosideros collina.
crassipes, *Lehm. Ind. Sem. Hort. Hamb.* (1842).—Austral.
cuneatum, Hoffmgg. Verz. Pfl. 72 = pubescens.
cuneiforme, Otto & Dietr. Allg. Gartenz. ix. (1841) 251 (sp. dub.).—Austral.
Cunninghamii, Schau. in Linnaea, xv. (1841) 420 = pubescens.
cupressinum, Otto & Dietr. Allg. Gartenz. ix. (1841) 250 (sp. dub.).—Austral.
cuspidatum, Hoffmgg. Verz. Pfl. 72 = pubescens?
divaricatum, Schau. in Walp. Rep. ii. 923 = scoparium.
dubium, Spreng. Syst. ii. 492 = Astartea fascicularis.
ellipticum, *Endl. in Enum. Pl. Hueg.* 51.—Austral.
emarginatum, Wendl. f. ex Link, Enum. Hort. Berol. ii. 25 = flavescens.
ericoides, *A. Rich. Bot. Voy. Astrolab.* i. 338.—N. Zel.
eriocalyx, Sieber, ex Spreng. Syst. iv. Cur. Post. 194 = parvifolium.
erubescens, *Schau. in Lehm. Pl. Preiss.* i. 121.—Austral.
Fabricia, *Benth. Fl. Austral.* iii. 102.—Austral.
firmum, *Benth. l. c.* 104.—Austral.
flavescens, *Sm. in Trans. Linn. Soc.* iii. (1797) 262.—Austral.
flexuosum, Spreng. Nov. Prov. 25 = Agonis flexuosa.
floribundum, Jungh. Java, i. 578; et in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 37 = javanicum.
floribundum, Salisb. Prod. 349 = scoparium.
floridum, *Benth. Fl. Austral.* iii. 110.—Austral.
glaucescens, Schau. in Linnaea, xv. (1841) 421 = pubescens.
glomeratum, Wendl. f. in Flora, ii. (1819) 678 = Agonis flexuosa.
gnidiaefolium, DC. Prod. iii. 228 = attenuatum.
gnidiaefolium, Hort. ex Benth. Fl. Austral. iii. 197 = stellatum.
grandiflorum, Lodd. Bot. Cab. t. 514 = flavescens.
grandifolium, Sm. in Trans. Linn. Soc. vi. (1802) 299 = pubescens.
humifusum, A. Cunn. ex Schau. in Linnaea, xv. (1841) 425 = scoparium.
hypericifolium, Otto & Dietr. Allg. Gartenz. ix. (1841) 243, nomen = Agonis hypericifolia.
imbricatum, Sm. in Trans. Linn. Soc. vi. (1802) 300 = Baeckea camphorata.
incanum, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 335.—Austral.
javanicum, *Blume, Bijdr.* 1100.—Burma; Java.
juniperifolium, Cav. Ic. iv. 18 = scoparium.
juniperinum, Sm. in Trans. Linn. Soc. iii. (1797) 263 = scoparium.
laevigatum, *F. Muell. Ann. Rep.* (1858) 22; *Fragm.* iv. 60.—Austral.
lanigerum, Sm. in Trans. Linn. Soc. iii. (1797) 263 = pubescens.
laricifolium, A. Cunn. ex Schau. in Walp. Rep. ii. 922 = Astartea fascicularis.
Leucadendrum, Forst. Char. Gen. 72 = flavescens, pubescens.
linearifolium, DC. Prod. iii. 227 = Agonis linearifolia.
linifolium, Dum.-Cours. Bot. Cult. ed. II. v. 384 = scoparium.
lucidum, Hort. Vinar. ex Schau. in Linnaea, xv. (1841) 435 = attenuatum.
macrocarpum, A. Cunn. ex Schau. l. c. 425 = scoparium.
marginatum, Labill. Nov. Holl. Pl. ii. 10. t. 148 = Agonis marginata.
micromyrtus, Miq. in Nederl. Kruidk. Arch. iv. (1859) 145 = flavescens.
microphyllum, Hoffmgg. Verz. Pfl. 72 = pubescens?
microphyllum, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 142 = pubescens.
mucronatum, *Cels, ex DC. Prod.* iii. 229, nomen.—Hab.?

LEPTOSPERMUM:—
multicaule, A. Cunn. in Field, N. S. Wales, 349 = myrtifolium.
multiflorum, Cav. Ic. iv. 17. t. 331. f. 1 = scoparium.
myrsinoides, *Schlecht. in Linnaea*, xx. (1847) 653.—Austral.
myrtifolium, *Sieber, ex DC. Prod.* iii. 228.—Austral.
nervosum, Hort. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 260 = flavescens.
nitens, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 335.—Austral.
nitidum, Hook. f. Fl. Tasm. i. 139 = pubescens.
nobile, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 145 = flavescens.
obliquum, Colla, Hort. Ripul. App. ii. 351 = scoparium.
obovatum, Sweet, Fl. Austral. t. 36 = flavescens.
obtusum, Lodd. Cat. ex G. Don, in Loud. Hort. Brit. 195, nomen = flavescens.
oligandrum, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 335.—Austral.
ovale, Dum.-Cours. Bot. Cult. ed. II. v. 385 = parvifolium.
ovalifolium, Sweet, Hort. Brit. ed. I. 155 = pubescens?
Oxycedrus, Schau. in Linnaea, xv. (1841) 432 = scoparium.
paludosum, Hort. Berol. ex Schau. l. c. 410 = flavescens.
parvifolium, *Sm. in Trans. Linn. Soc.* iii. (1797) 263. —Austral.
parvulum, Labill. Sert. Austro-Caled. 62. t. 61 = Baeckia virgata.
pendulum, Sieber, ex Benth. Fl. Austral. iii. 108 = attenuatum.
perforatum, Forst. Char. Gen. 72 = Metrosideros scandens.
persiciflorum, Reichb. Hort. Bot. iii. 8. t. 220 = scoparium.
pilosum, Schau. in Walp. Rep. ii. 923 = pubescens.
pinifolium, Labill. Sert. Austro-Caled. 63. t. 62 = Baeckea pinifolia.
polygalaefolium, Salisb. Prod. 350 = flavescens.
porophyllum, Cav. Ic. iv. 17 = flavescens.
pubescens, *Lam. Encyc.* iii. 466.—Austral.
pungens, *Banks, ex Dum.-Cours. Bot. Cult.* ed. II. v. 385.—Austral.
pungens, *Otto & Dietr. Allg. Gartenz.* ix. (1841) 260. —Austral.
recurvifolium, Salisb. Prod. 350 = scoparium.
recurvum, *Hook. f. Ic. Pl.* t. 893.—Borneo.
resiniferum, Bertol. Amoen. Ital. 29 = Agonis flexuosa.
retusum, Hort. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 243 = flavescens.
robustum, Endl. in Enum. Pl. Hueg. 50 = Hypocalymma robustum.
Roei, *Benth. Fl. Austral.* iii. 110.—Austral.
roseum, Hort. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 243 = flavescens.
rubricaule, Cels, ex Link, Enum. Hort. Berol. ii. 25 = scoparium.
rupestre, *Hook. f. Ic. Pl.* t. 308.—Tasman.
salicifolium, *Lam. Encyc.* iii. 467.—Java.
scabrum, Wall. Cat. sub n. 3650 = Deutzia staminea.
scandens, Forst. Char. Gen. 72 = Metrosideros florida.
scoparium, *Forst. l. c.*—Austral.; N. Zel.
sericatum, Lindl. in Mitch. Journ. Trop. Austral. 298 = stellatum.
sericeum, Hort. Berol. ex Schau. in Linnaea, xv. (1841) 421 = pubescens.
sericeum, Labill. Nov. Holl. Pl. ii. 9. t. 147 = Kunzea sericea.
sericeum, Schau. in Lehm. Pl. Preiss. i. 121 = erubescens.
sericophyllum, Schau. in Linnaea, xv. (1841) 420 = pubescens.
speciosum, Schau. in Walp. Rep. ii. 923 = Melaleuca Leucadendron.
spinescens, *Endl. in Enum. Pl. Hueg.* 51.—Austral.
splendens, Schau. in Walp. Rep. ii. 923 = pubescens.
squarrosum, Gaertn. Fruct. i. 174. t. 35; Sieber, ex Spreng. Syst. iv. Cur. Post. 194 = scoparium.
stamineum, Wall. Cat. sub n. 3651 = Deutzia staminea.
stellatum, *Cav. Ic.* iv. 16. t. 330. f. 1.—Austral.
styphelioides, Schau. in Linnaea, xv. (1841) 423 = scoparium.
tenuifolium, Cav. ex Steud. Nom. ed. I. 474; ed. II. ii. 32 = scoparium.
Thea, Willd. Sp. Pl. ii. 949 = flavescens.

LEPTOSPERMUM:—
theaeforme, A. Cunn. ex Schau. in Walp. Rep. v. 739 = Agonis hypericifolia.
thymifolium, A. Cunn. in Field, N. S. Wales, 349 = pubescens.
thymifolium, Hoffmgg. Verz. Pfl. 72 = praec.?
tomentosum, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 384 = pubescens.
tonsum, Schau. in Linnaea, xv. (1841) 422 = pubescens.
tortuosum, Dehnh. Revist. Napol. i. III. 171; ex Walp. Rep. ii. 169 (sp. dub.).—Austral.
triloculare, Vent. Jard. Malm. 88. t. 84 = arachnoideum.
trinerve, White, ex Sm. in Trans. Linn. Soc. iii. (1797) 263, err. = Melaleuca trinervia, *White*.
tuberculatum, Poir. Encyc. Suppl. iii. 338 = flavescens.
umbellatum, Gaertn. Fruct. i. 174. t. 35. f. 3 = Eucalyptus tereticornis.
villosum, Fisch. Hort. Gorenk. ed. II. (1812) 63 (sp. dub.).—Hab.?
villosum, Otto & Dietr. Allg. Gartenz. ix. (1841) 242 = pubescens.
virgatum, Forst. Char. Gen. 72 = Baeckea virgata.
virgatum, Willd. Sp. Pl. ii. 951 = flavescens.

LEPTOSTACHIA, Mitch. in Act. Phys. Med. Soc. Nat. Cur. viii. (1748), App. 212 = **Phryma**, Linn. (Verbenac.).

LEPTOSTACHYA, Benth. & Hook. f. Gen. ii. 1137 (1876) = praec.

LEPTOSTACHYA, Nees, in Wall. Pl. As. Rar. iii. 105 (1832) = **Dianthera** et **Justicia**, Linn. (Acanth.).
antirrhina, Nees, in Mart. Fl. Bras. ix. 150 = J. antirrhina.
campylostemon, Nees, in DC. Prod. xi. 378 = J. campylostemon.
comata, Kostel. Allg. Med.-Pharm. Flora, iii. 932 = D. comata.
cordata, Nees, in DC. Prod. xi. 378 = J. cordata.
crinita, Nees, in Benth. Bot. Voy. Sulph. 147.—N. Granat.
dichotoma, Nees, in DC. Prod. xi. 379 = D. dichotoma.
heterophylla, Nees, in Mart. Fl. Bras. ix. 150 = D. heterophylla.
laxa, Nees, l. c. 151.—Bras.
lucida, Nees, l. c. 149.—Bras.
Martiana, Nees, in Benth. Bot. Voy. Sulph. 147 = D. comata.
parviflora, Nees, in Mart. Fl. Bras. ix. 151 = D. comata.
pectoralis, Nees & Mart. in Flora, xxii. (1839) I. Beibl. 8; et ex Kostel. Allg. Med.-Pharm. Flora, iii. 932 = D. pectoralis.
Poeppigiana, Nees, in Mart. Fl. Bras. ix. 150.—Peruv.
reptans, Kostel. Allg. Med.-Pharm. Flora, iii. 932 = D. reptans?
rupestris, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 532.—Bras.
secundiflora, Nees, in DC. Prod. xi. 378.—Peruv.
virens, Nees, l. c.—Afr. trop.
virgata, Nees, in Wall. Pl. As. Rar. iii. 105 = D. virgata?
Wallichii, Nees, l. c. = D. Wallichii.
zeylanica, Nees, in DC. Prod. xi. 379 = J. Hookeriana.

LEPTOSTACHYS, Ehrh. Beitr. iv. 147 (1789) = **Carex**, Linn. (Cyperac.).

LEPTOSTACHYS, G. F. W. Mey. Prim. Fl. Esseq. 73 (1818) = **Leptochloa**, Beauv. (Gramin.).
Digitaria, G. F. W. Mey. l. c. 74 = Leptochloa Digitaria.
domingensis, G. F. W. Mey. l. c. = Leptochloa domingensis.
dubia, G. F. W. Mey. l. c. = Diplachne dubia.
filiformis, G. F. W. Mey. l. c. = Leptochloa filiformis.
gracilis, G. F. W. Mey. l. c. = Leptochloa gracilis.
virgata, G. F. W. Mey. l. c. = Leptochloa virgata.

LEPTOSTELMA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 38 (1830) = **Erigeron**, Linn. (Compos.).
Donianum, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 243 = Eupatorium macrocephalum, leptolepis.
maxima, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 38 = Erigeron maximus.

LEPTOSTEMMA, Blume, Bijdr. 1057 (1826) = **Dischidia**, R. Br. (Asclep.).
fasciculatum, Blume, l. c. 1058 = D. hirsuta.
hirsutum, Blume, l. c. = D. hirsuta.
lanceolatum, Blume, l. c. 1057 = D. lanceolata.
latifolium, Blume, l. c. 1058 = D. latifolia.
peltatum, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 148 = D. peltata.
pictum, Zipp. ex Blume, l. c. = D. picta.
punctatum, Blume, Bijdr. 1058 = D. punctata.
sagittatum, Blume, l. c. 1057 = D. sagittata.
truncatum, Blume, l. c. 1058 = D. truncata.
truncatum, Zoll. ex Miq. Fl. Ind. Bat. ii. 519 = D. rhombifolia.

LEPTOSTIGMA, Arn. in Hook. Journ. Bot. iii. (1841) 270 = **Nertera**, Banks & Soland. (Rubiac.).
Arnottianum, Walp. Rep. ii. 463 (= *Coprosma calycina*).—Chili.

LEPTOSTYLIS, Benth. in Benth. & Hook. f. Gen. ii. 659 (1876). *SAPOTACEAE*, Benth. & Hook. f. l. c.
filipes, *Benth. l. c.*—N. Caled.
longiflora, *Benth. l. c.*—N. Caled.

LEPTOSYNE, DC. Prod. v. 531 (1836). *COMPOSITAE*, [Benth. & Hook. f. ii. 385].
ACOMA, Benth. Bot. Voy. Sulph. 29. t. 17 (1844).
PUGIOPAPPUS, A. Gray, ex Torr. in Pacif. Rail. Rep. iv. 104 (1857).
TUCKERMANNIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 363.
arizonica, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 218.—Arizona.
Bigelovii, *A. Gray, Syn. Fl. N. Am.* i. II. 300.—Calif.
californica, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 363 = Douglasii.
calliopsidea, *A. Gray, Syn. Fl. N. Am.* i. II. 300.—Calif.
dissecta, *A. Gray, l. c.* 301.—Calif.
Douglasii, *DC. Prod.* x. 531.—Calif.
gigantea, *Kellogg, in Proc. Calif. Acad.* iv. (1870) 198.—Calif.
heterocarpa, A. Gray, Syn. Fl. N. Am. i. II. 301 = Coreocarpus heterocarpus.
maritima, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358. —Calif.
Newberryi, A. Gray, l. c. = Douglasii.
parthenioides, A. Gray, Syn. Fl. N. Am. i. II. 301 = Coreocarpus parthenioides.
Stillmannii, *A. Gray, in Torr. Bot. Mex. Bound.* 92.—Calif.

LEPTOTAENIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 629 (1840) = **Ferula**, Tourn. (Umbellif.).
californica, Nutt. l. c. 630 = F. californica.
dissecta, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 408 = Peucedanum nudicaule.
dissecta, Nutt. ex Torr. & Gray. Fl. N. Am. i. 630 = F. dissoluta.
multifida, Nutt. l. c. = F. multifida.

LEPTOTES, Lindl. Bot. Reg. t. 1625 (1833) = **Tetramicra**, Lindl. (Orchid.).
bicolor, Lindl. l. c. = T. bicolor.
glaucophylla, Hoffmgg. in Bot. Zeit. i. (1843) 833 = T. bicolor.
paranaënsis, Rodrig. Orch. Nov. ii. 163.—Bras.
serrulata, Lindl. Sert. Orch. (1838) t. 11 = T. bicolor.
tenuis, Reichb. f. in Hamb. Gartenz. xxi. (1865) 296. —Bras.
unicolor, Rodrig. Orch. Nov. i. 74.—Bras.

LEPTOTHAMNUS, DC. Prod. v. 367 (1836) = **Nolletia**, Cass. (Compos.).
ciliaris, DC. l. c. = N. ciliaris.
rarifolius, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 65 = N. rarifolia.

LEPTOTHERIUM, D. Dietr. Syn. Pl. i. 398 (1839) = seq.

LEPTOTHRIUM, Kunth, in H. B. & K. Nov. Gen. et Sp. i. 340 (1815) = **Isochilus**, R. Br. (Orchid.).
lineare, Kunth, l. c. = I. linearis.

LEPTOTHRIUM, Kunth, Rév. Gram. 156 (1830); et Enum. Pl. i. 470. t. 38. f. 2 (1833). *GRAMINEAE*, Benth. & Hook. f. iii. 1124.
rigidum, *Kunth, ll. cc.* 156; 471.—Am. trop.

LEPTOTHYRSA, Hook. f. in Benth. & Hook. f. Gen. i. 284 (1862). *RUTACEAE*, Benth. & Hook. f. l. c.
Sprucei, *Hook. f. l. c.*—Bras.

LEPTOTIS, Hoffmgg. Verz. Pfl. 72 (1824) = **Ursinia**, Gaertn. (Compos.).
anthemoides, Hoffmgg. l. c. = U. anthemoides.

LEPTOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 73 = **Gnephosis**, Cav. (Compos.).
perpusilla, Turcz. l. c. = G. pygmaea.

LEPTRANTHUS, Steud. Nom. ed. II. ii. 32 (1841) = LEPTERANTHUS, Neck. = **Centaurea**, Linn. (Compos.).

LEPTRINA, Rafin. in Am. Monthly Mag. (1819) 192 = **Montia**, Linn. (Portulac.).
autumnalis, Rafin. in Journ. Phys. (1819) 96 = M. fontana.

LEPTRINIA, Schult. Mant. ii. 38 (1824) = praec.

LEPTUNIS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 366, in textu = **Asperula**, Linn. (Rubiac.).
tenuis, Stev. l. c. = A. trichodes.

LEPTUROPSIS, Steud. Syn. Pl. Gram. 357 (1855) *GRAMINEARUM* gen. dub.—Cf. Benth. & Hook. f. iii. 1096.
triaristata, Steud. l. c. 358.—'Ins. Loss Guineae.'

LEPTURUS, R. Br. Prod. 207 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1204.
LEPIURUS, Dum. Obs. Gram. Belg. 140 (1823).
LEPTOCERCUS, Rafin. in Am. Monthly Mag. (1819) 190.
PHOLIURUS, Trin. Fund. Agrost. 131 (1820).
ROTTBOELLIA, Host, Gram. Austr. i. t. 24 (1801).
aciculatus, *Steud. Syn. Pl. Gram.* 357.—Ind. or.
acutiglumis, *Steud. l. c.*—Ins. Tahiti.
Bolanderi, *Thurb. ex A. Gray, in Proc. Am. Acad.* vii. (1868) 401.—Calif.
compressus, Steud. Syn. Pl. Gram. 357 = filiformis.
cylindricus, *Trin. Fund. Agrost.* 123.—Reg. bor. et austr.
erectus, Szov. ex Griseb. in Ledeb. Fl. Ross. iv. 325 = incurvatus.
fasciculatus, Trin. Fund. Agrost. 123 = Hemarthria compressa.
filiformis, *Trin. l. c.*—Europ.; Afr. bor.
hirtulus, *Regel, in Act. Hort. Petrop.* vii. (1880) 576. —Turkestan.
incurvatus, *Trin. Fund. Agrost.* 123.—Reg. bor. et austr.
minimus, *Hochst. in Flora*, xxxviii. (1855) 333.—Abyssin.
paniculatus, Nutt. Gen. Am. i. 81 = Schedonnardus texanus.
pannonicus, *Kunth, Rév. Gram.* i. 151.—Europ. austr.; Reg. Caucas.
persicus, *Boiss. Diagn.* Ser. I. xiii. 71.—Persia.
pubescens, *Boiss. Fl. Orient.* v. 685.—Mesopot.
ramosus, *Kunth, Rév. Gram.* i. 152.—Afr. bor.
repens, Boj. Hort. Maurit. 372 = Stenotaphrum subulatum.
repens, *R. Br. Prod.* 207.—Austral.; Ins. Pacif.
Roxburghianus, *Steud. Syn. Pl. Gram.* 357.—Ind. or.
simplex, Kunth, Enum. Pl. i. 463 = cylindricus?
strigosus, Dum. Obs. Gram. Belg. 146 = filiformis.
subulatus, Kunth, Rév. Gram. i. 151 = cylindricus.
tenuis, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 97.—Ins. Socotr.

LEPURANDRA, Nimmo, in J. Grah. Cat. Bomb. Pl. 193 (1839) = **Antiaris**, Lesch. (Urticac.).
saccidora, Nimmo, l. c. = A. toxicaria.

LEPUROPETALON, Ell. Sketch, i. 37 (1817). *SAXIFRAGEAE*, Benth. & Hook. f. i. 639.
CRYPTOPETALUM, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 344.
pusillum, *C. Gay, Fl. Chil.* iii. 42.—Chili.
spathulatum, *Ell. Sketch*, i. 370.—Am. bor.

LEPYRODIA, R. Br. Prod. 247 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1030.
SPORADANTHUS, F. Muell. ex J. Buch. in Trans. N. Z. Inst. vi. (1874) 340; x. (1878) App. 41.

LEPYRODIA:—
anaectocolea, *F. Muell. Fragm.* viii. 78.—Austral.
Anarthria, *F. Muell. l. c.* 73.—Austral.
Drummondiana, *Steud. Syn. Pl. Cyp.* 248.—Austral.
elongata, Spreng. Syst. iv. Cur. Post. 36 = Hypolaena lateriflora.
glauca, *F. Muell. Fragm.* viii. 77.—Austral.
gracilis, *R. Br. Prod.* 247.—Austral.
hermaphrodita, *R. Br. Prod.* 248.—Austral.
hermaphrodita, Nees, ex Benth. Fl. Austral. vii. 216 = scariosa.
interrupta, *F. Muell. Fragm.* viii. 74.—Austral.
macra, *Nees, in Ann. & Mag. Nat. Hist.* Ser. I. vi. (1841) 50.—Austral.
monoica, *F. Muell. Fragm.* viii. 76.—Austral.
Muelleri, *Benth. Fl. Austral.* vii. 215.—Austral.
Muirii, *F. Muell. Fragm.* viii. 78.—Austral.
paniculata, F. Muell. 2nd Gen. Rep. 16 = tasmanica.
scariosa, *R. Br. Prod.* 248.—Austral.
stricta, *R. Br. l. c.* 247.—Austral.
stricta, F. Muell. ex Benth. Fl. Austral. vii. 216 = Muelleri.
stricta, Nees, ex Benth. l. c. 216 = gracilis.
succedanea, Nees, in Lehm. Pl. Preiss. ii. 60 = stricta.
tasmanica, *Hook. f. Fl. Tasm.* ii. 72. t. 135.—Austral.
tasmanica, F. Muell. Fragm. viii. 75 = Muelleri.
Traversii, *F. Muell. l. c.* 79.—Ins. Chath.

LEPYRODICLIS, Fenzl, ex Endl. Gen. 966 (1840) = **Arenaria**, Rupp. (Caryoph.).
cerastioides, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 170.—As. centr.
holosteoides, Fenzl, ex Fisch. & Mey. Enum. Pl. Nov. Schrenk, 93, in nota = A. holosteoides.
stellarioides, Fisch. & Mey. l. c. = A. stellarioides.
tenera, Boiss. Diagn. Ser. II. i. 84 = A. tenera.

LEPYRONIA, Lestib. Ess. Cyp. 25 (1819) = **Lepironia**, Rich. (Cyperac.).

LERCHEA, Hall. Hort. Gotting. 21 (1743) = **Suaeda**, Forsk. (Chenop.).
altissima, Medic. Phil. Bot. i. 33 = S. altissima.
fruticosa, Medic. l. c. = S. fruticosa.
obtusifolia, Steud. Nom. ed. I. 187, in syn. = S. fruticosa.
salsa, Medic. Phil. Bot. i. 33 = S. salsa.

LERCHEA, Linn. Mant. ii. 155 (1767). *RUBIACEAE*, Benth. & Hook. f. ii. 53.
CODARIA, Linn. ex Benn. Pl. Jav. Rar. 99 (1838).
calycina, *A. Gray, in Proc. Am. Acad.* iv. (1858–59) 311.—Ins. Pacif.
fruticulosa, Reinw. ex Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 165 = Xanthophytum fruticulosum.
interrupta, *Korth. l. c.* 164.—Sumatra.
longicauda, *Linn. Mant.* ii. 256.—Java.

LERCHENFELDIA, Schur, Enum. Pl. Transs. 753 (1866) = **Deschampsia**, Beauv. (Gramin.).
cuprina, Schur, l. c. = D. flexuosa.
flexuosa, Schur, l. c. = D. flexuosa.
uliginosa, Schur, l. c. 754 = D. flexuosa.

LERCHIA, Endl. Gen. 298, in syn. (1837) = **Lerchea**, Linn. (Rubiac.).

LERCHIA, Reichb. Consp. 109 (1828) = LEACHIA, Cass. = **Coreopsis**, Linn. (Compos.).

LERESCHIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 127 = **Pimpinella**, Linn. (Umbellif.).
Thomasii, Boiss. l. c. 128 = P. Thomasii.

LERETIA, Vell. Fl. Flum. 99; iii. t. 2 (1825) = **Mappia**, Jacq. (Olac.).
ampla, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iv. (1859) 364 = M. ampla.
cordata, Vell. Fl. Flum. 99; iii. t. 2 = M. cordata.
nitida, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iv. (1859) 365 = M. nitida.
paniculata, Mart. Fl. Bras. vii. 17 = Villaresia paniculata.
Velloxii, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 392 = M. cordata.

LERIA, DC. in Ann. Mus. Par. xix. (1812) 68 = **Chaptalia**, Vent. (Compos.).
albicans, DC. l. c. = C. albicans.

LERIA:—
araneosa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 374 = Trichocline araneosa.
caespitosa, Spreng. Syst. iv. 502 = Chevreulia stolonifera.
dentata, Spreng. l. c. = Chaptalia dentata.
exscapa, DC. in Ann. Mus. Par. xix. (1812) 68 = C. exscapa.
integrifolia, Cass. in Dict. Sc. Nat. xxvi. 103 = C. integrifolia.
leiocarpa, DC. Prod. vi. 42 = C. integrifolia.
lutescens, Sch. Bip. in Linnaea, xxxiv. (1865–66) 527.—Bolivia.
lyrata, DC. in Ann. Mus. Par. xix. (1812) 68 = C. nutans.
Mandonii, Sch. Bip, in Linnaea, xxxiv. (1865–66) 527, nomen = C. piloselloides.
media, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 515.—Ind. occ.
nervosa, Spreng. Syst. iii. 501 = Landtia nervosa.
nutans, DC. in Ann. Mus. Par. xix. (1812) 68 = C. nutans.
nutans, Less. in Linnaea, v. (1830) 154, partim = C. integrifolia.
pumila, DC. in Ann. Mus. Par. xix. (1812) 68 = C. sinuata.
rotundifolia, Sch. Bip. in Linnaea, xxxiv. (1865–66) 527, nomen = C. rotundifolia.
sarmentosa, DC. in Ann. Mus. Par. xix. (1812) 68 = Chevreulia stolonifera.
sinuata, DC. Prod. vii. 42 = C. sinuata.
spathulata, D. Don, in Trans. Linn. Soc. xvi. (1830) 249 = C. spathulata.
stenocephala, Griseb. Cat. Pl. Cub. 158.—Cuba.
tomentosa, Griseb. Fl. Brit. W. Ind. 383, in obs. = C. tomentosa.

LEROUXIA, Mérat, Nouv. Fl. Par. 77 (1812) = **Lysimachia**, Linn. (Primul.).
nemorum, Mérat, l. c. = Lysim. nemorum.

LESCAILLEA, Griseb. Cat. Pl. Cub. 156 (1866). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
equisetiformis, *Griseb. l. c.* 157.—Cuba.

LESCHENAULTIA, R. Br. Prod. 581 (1810) (Lechenaultia). *GOODENOVIEAE*, Benth. & Hook. f. ii. 537.
LATOURIA, Lindl. Veg. Kingd. 695 (1847).
acutiloba, *Benth. Fl. Austral.* iv. 41.—Austral.
agrostophylla, *F. Muell. Fragm.* vi. 8.—Austral.
arcuata, De Vriese, in Lehm. Pl. Preiss. i. 416 = linarioides.
Baxteri, G. Don, in Loud. Hort. Brit. 79 = formosa.
biloba, *Lindl. Swan River App.* 27.—Austral.
chlorantha, *F. Muell. Fragm.* ii. 20.—Austral.
divaricata, *F. Muell. l. c.* iii. 33.—Austral.
Drummondi, De Vriese, Gooden. (in Verh. Haarl. x.) 182 = biloba.
expansa, *R. Br. Prod.* 581.—Austral.
expansa, De Vriese, Gooden. 184 = floribunda.
filiformis, *R. Br. Prod.* 581.—Austral.
floribunda, *Benth. in Enum. Pl. Hueg.* 70.—Austral.
formosa, *R. Br. Prod.* 581.—Austral.
glauca, Lindl. Swan River App. 27 = floribunda.
grandiflora, DC. Prod. vii. 519 = biloba.
heteromera, *Benth. Fl. Austral.* iv. 43.—Austral.
hirsuta, *F. Muell. Fragm.* vi. 9.—Austral.
humilis, Spreng. Syst. i. 720 = Anthotium humile.
laricina, *Lindl. Swan River App.* 27.—Austral.
linarioides, *DC. Prod.* vii. 519.—Austral.
longiloba, *F. Muell. Fragm.* vi. 10.—Austral.
multiflora, Lodd. Bot. Cab. t. 1579 = formosa.
oblata, Sweet, Fl. Austral. t. 46 = formosa.
pallescens, De Vriese, in Lehm. Pl. Preiss. i. 415 = floribunda.
parviflora, De Vriese, l. c. i. 416 = laricina.
pinastroides, Lehm. l. c. ii. 244 = tubiflora.
splendens, Hook. Bot. Mag. t. 4256 = laricina.
stricta, *F. Muell. Fragm.* viii. 245.—Austral.
superba, *F. Muell. l. c.* vi. 10.—Austral.
tenuifolia, De Vriese, in Lehm. Pl. Preiss. i. 415 = laricina.
tubiflora, *R. Br. Prod.* 581.—Austral.

LESEMIA, Rafin. Fl. Tellur. iii. 91 (1836) = **Salvia**, Tourn. (Labiat.).
coccinea, Rafin. l. c. = S. Grahami.

LESOURDIA, Fourn. in Bull. Soc. Bot. Fr. xxvii. (1880) 102. tt. 3, 4 = **Scleropogon**, Phil. (Gramin.).
Karwinskyana, Fourn. l. c. = S. brevifolius.
multiflora, Fourn. l. c. = S. brevifolius.

LESPEDEZA, Michx. Fl. Bor. Am. ii. 70. t. 39 (1803).
LEGUMINOSAE, Benth. & Hook. f. i. 524.
CAMPYLOTROPIS, Bunge, Pl. Monghol.-Chin. 6 (1835).
OXYRAMPHIS, Wall. Cat. n. 5348 (1831).
PHLEBOSPORIUM, Jungh. Reise, 346 (1845).
angustifolia, Darl. Florul. Cestr. 81; et Fl. Cestr. ed. I. 419; Rafin. ex DC. Prod. ii. 349 = reticulata.
angustifolia, *Ell. Sketch*, ii. 206.—Am. bor.
angustifolia, Nutt. ex Maxim. in Act. Hort. Petrop. ii. (1873) 379 = polystachya.
argyraea, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 120 = juncea.
bicolor, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 69.—China.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 47.—Japon.
capitata, *Michx. Fl. Bor. Am.* ii. 71.—Am. bor.
Caraganae, *Bunge, Pl. Monghol.-Chin.* 11.—China.
chinensis, *G. Don, Gen. Syst.* ii. 307.—China.
ciliata, *Benth. in Hook. Kew Journ.* iv. (1852) 48.—China.
coriacea, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 417 = Desmodium ciliare?
cuneata, G. Don, Gen. Syst. ii. 307 = juncea.
cyrtobotrya, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 48.—Japon.
cytisoides, *Benth. in Miq. Pl. Jungh.* 230.—Java.
cytisoides, Bertol. Misc. Bot. ix. 14 = Rhynchosia galactioides.
Davidi, *Franch. Pl. Davidian.* i. (1884) 94.—China.
decora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 231.—Burma.
divergens, Pursh, Fl. Am. Sept. ii. 481 = violacea.
elegans, *Cambess. in Jacquem. Voy. Bot.* 43. t. 52.—Reg. Himal.
elliptica, Benth. ex Maxim. in Act. Hort. Petrop. ii. (1873) 353 = Sieboldi.
eriocarpa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 102.—Reg. Himal.
floribunda, *Bunge, Pl. Monghol.-Chin.* 13.—China.
frutescens, DC. Prod. ii. 349 (excl. syn.) = reticulata.
frutescens, Ell. Sketch, ii. 206 = capitata.
fruticosa, Pers. Syn. ii. 318 = capitata.
Gerardiana, *R. Grah. in Wall. Cat.* n. 5744.—Reg. Himal.
glomerata, Hornem. Hort. Hafn. Suppl. 81 = villosa.
hirta, Ell. Sketch, ii. 207 = polystachya.
hirta, Hornem. Hort. Hafn. 699 = villosa.
indica, Spreng. Syst. iii. 202 = Rhynchosia rufescens.
involucrata, Wall. Cat. n. 5742 = Flemingia involucrata.
juncea, *Pers. Syn.* ii. 318.—Reg. Himal.; As. bor.
juncea, Wall. Cat. n. 5743 B = Indigofera aspalathoides.
kanaorensis, Cambess. in Jacquem. Voy. Bot. 42 = juncea.
lagopoides, Pers. Syn. ii. 318 = Uraria lagopoides.
lanata, *Benth. Fl. Austral.* ii. 241.—Austral.
latebrosa, Pers. Syn. ii. 318 = Heylandia latebrosa.
leptostachya, *Engelm. ex A. Gray, in Proc. Am. Acad.* xii. (1877) 57.—Am. bor.
lineata, Pers. Syn. ii. 318 = Flemingia lineata.
longifolia, DC. Prod. ii. 349 = capitata.
macrocarpa, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 92.—China.
macrophylla, Bunge, Pl. Monghol.-Chin. 10 = villosa.
macrostyla, Baker, ex Maxim. in Act. Hort. Petrop. ii. (1873) 348 = sericea.
medicaginoides, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 93.—China.
Nuttallii, Darl. Fl. Cestr. ed. I. 420 = Stuvei.
Oldhami, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 48 = Buergeri.
Pallasii, *G. Don, Gen. Syst.* ii. 307.—Sibiria.
paniculata, Royle, ex Maxim. in Act. Hort. Petrop. ii. (1873) 349 = eriocarpa.
parviflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 231.—Burma.
pilosa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 121.—Japon.
pinetorum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 230.—Burma.
polystachya, *Michx. Fl. Bor. Am.* ii. 71. t. 40.—Am. bor.
procumbens, Michx. l. c. 70. t. 39 = repens.
prostrata, Pursh, Fl. Am. Sept. ii. 481 = repens.
repens, *Barton, Prod. Fl. Philad.* ii. 77.—Am. bor.
reticulata, *Pers. Syn.* ii. 318.—Am. bor.
Royleana, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 50 = sericea.
sericea, *Benth. in Miq. Pl. Jungh.* 227.—Reg. Himal.
sericea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 49 = juncea.
sericea, Royle, ex Maxim. in Act. Hort. Petrop. ii. (1873) 349 = stenocarpa.
sessiliflora, Michx. Fl. Bor. Am. ii. 70 = reticulata.
sessiliflora, *Nutt. Gen. Am.* ii. 107.—Am. bor.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 47.—Japon.; Reg. Himal.
speciosa, Royle, ex Maxim. in Act. Hort. Petrop. ii. (1873) 350 = eriocarpa.
stenocarpa, *Maxim. l. c.* 349.—Reg. Himal.
stipulacea, Maxim. Prim. Fl. Amur. 85 = striata.
striata, *Hook. & Arn. Bot. Beech. Voy.* 262.—Japon.
Stuvei, *Nutt. Gen. Am.* ii. 107.—Am. bor.
Swinhoei, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 210 = virgata.
Thomsoni, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 144.—Reg. Himal.
tomentosa, Siebold, ex Maxim. in Act. Hort. Petrop. ii. (1873) 376 = villosa.
trichocarpa, *Pers. Syn.* ii. 318.—Sibiria.
variegata, Cambess. in Jacquem. Voy. Bot. 42 = juncea.
viatorum, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 46.—China.
villosa, *Pers. Syn.* ii. 318.—Am. bor.
violacea, Ell. Sketch, ii. 205 = reticulata.
violacea, *Pers. Syn.* ii. 318.—Am. bor.
virgata, *DC. Prod.* ii. 350.—Japon.
virgata, *Miq. Fl. Ind. Bat.* i. I. 230.—Java.
virgata, Nutt. ex Darl. Fl. Cestr. ed. I. 420 = Stuvei.
virgata, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 121 = cyrtobotrya.

LESPEDEZIA, Spreng. Syst. iii. 202 (1826) = **Lespedeza**, Michx. (Legumin.).

LESQUEREUXIA, Boiss. & Reut. Diagn. Ser. I. xii. 43 (1853) = **Siphonostegia**, Benth. (Scrophul.).
syriaca, Boiss. & Reut. l. c. 44 = S. syriaca.

LESSERTIA, DC. Astrag. 37 (1802). *LEGUMINOSAE*, Benth. & Hook. f. i. 503.
SULITRA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 366.
abbreviata, *E. Mey. Comm. Pl. Afr. Austr.* 118.—Afr. austr.
acuminata, E. Mey. l. c. = physodes.
annua, DC. Astrag. 37 = linearis.
annularis, *Burch. Trav.* i. 304.—Afr. austr.
arcuata, Eckl. & Zeyh. Enum. 248 = annularis.
argentea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 221.—Afr. austr.
astragalina, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 88 = pulchra.
astragalina, Poepp. ex Harv. & Sond. Fl. Cap. ii. 218 = Pappeana.
benguellensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 137.—Afr. trop.
brachypus, *Harv. in Harv. & Sond. Fl. Cap.* ii. 214.—Afr. austr.
brachystachya, *DC. Mém. Lég.* vi. 291.—Afr. austr.
candida, *E. Mey. Comm. Pl. Afr. Austr.* 116.—Afr. austr.
capitata, *E. Mey. l. c.* 118.—Afr. austr.
carnosa, *Eckl. & Zeyh. Enum.* 248.—Afr. austr.
depressa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 219.—Afr. austr.
diffusa, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 327.—Afr. austr.
diffusa, Eckl. & Zeyh. Enum. 250 = excisa.
excisa, *DC. Prod.* ii. 272.—Afr. austr.
excisa, Eckl. & Zeyh. Enum. 250 = tomentosa.
falciformis, *DC. Mém. Lég.* vi. t. 46.—Afr. austr.
falciformis, Eckl. & Zeyh. Enum. 248 = annularis.
flexuosa, *E. Mey. Comm. Pl. Afr. Austr.* 116.—Afr. austr.
fruticosa, *Lindl. Bot. Reg.* t. 970.—Afr. austr.
inflata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 219.—Afr. austr.
lanata, *Harv. l. c.* 224.—Afr. austr.
linearis, *DC. Prod.* ii. 272.—Afr. austr.
macrostachya, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 100.—Afr. austr.
macrostachya, Eckl. & Zeyh. Enum. 249 = flexuosa.
margaritacea, *E. Mey. Comm. Pl. Afr. Austr.* 116.—Afr. austr.
microcarpa, *E. Mey. l. c.* 119.—Afr. austr.
monticola, R. Cunn. ex A. Gray, Bot. U. St. Expl. Exped. i. 411 = Swainsona monticola.
mucronata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 223.—Afr. austr.
obtusata, *DC. Prod.* ii. 272.—Afr. austr.
obtusata, Eckl. & Zeyh. Enum. 247 = stenoloba.
Pappeana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 218.—Afr. austr.
pauciflora, *Harv. l. c.* 216.—Afr. austr.
perennans, *DC. Astrag.* 47.—Afr. austr.
perennans, Eckl. & Zeyh. Enum. 250 = excisa.
physodes, *Eckl. & Zeyh. l. c.* 249.—Afr. austr.
polystachya, *Harv. in Harv. & Sond. Fl. Cap.* ii. 222.—Afr. austr.
procumbens, *DC. Prod.* ii. 273.—Afr. austr.
propinqua, Eckl. & Zeyh. Enum. 250 = linearis.
prostrata, *DC. Prod.* ii. 272.—Afr. austr.
prostrata, Eckl. & Zeyh. Enum. 249 = depressa.
pubescens, *DC. Prod.* ii. 272.—Afr. austr.
pubescens, Eckl. & Zeyh. Enum. 249 = pulchra.
pulchra, *Sims, Bot. Mag.* t. 2064.—Afr. austr.
rigida, DC. Prod. ii. 272 = flexuosa.
rigida, E. Mey. Comm. Pl. Afr. Austr. 115 = fruticosa.
spinescens, *E. Mey. l. c.*—Afr. austr.
stenoloba, *E. Mey. l. c.* 121.—Afr. austr.
subumbellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 217.—Afr. austr.
sulcata, *E. Mey. Comm. Pl. Afr. Austr.* 116.—Afr. austr.
tenuifolia, *E. Mey. l. c.* 117.—Afr. austr.
tomentosa, *DC. Prod.* ii. 272.—Afr. austr.
tomentosa, Eckl. & Zeyh. Enum. 250, partim = lanata.
tumida, Eckl. & Zeyh. l. c. 249 = physodes.
venusta, Eckl. & Zeyh. l. c. 248 = pulchra.
vesicaria, *DC. Prod.* ii. 272.—Afr. austr.
vesicaria, Eckl. & Zeyh. Enum. 250 = argentea.
vesicaria, E. Mey. Comm. Pl. Afr. Austr. 119 = inflata.
villosa, E. Mey. in Linnaea, vii. (1832) 169 = argentea.

LESSINGIA, Cham. in Linnaea, iv. (1829) 203. t. 2. f. 2. *COMPOSITAE*, Benth. & Hook. f. ii. 257.
adenophora, Greene, in Bull. Calif. Acad. i. IV. (1885) 190 = ramulosa.
germanorum, *Cham. in Linnaea*, iv. (1829) 203.—Calif.
glandulifera, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 207.—Calif.
leptoclada, *A. Gray, l. c.* vii. (1868) 351.—Calif.
nana, *A. Gray, in Benth. Pl. Hartw.* 315.—Calif.
nemaclada, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 191.—Calif.
Parryi, *Greene, l. c.*—Calif.
ramulosa, *A. Gray, in Benth. Pl. Hartw.* 314.—Calif.
virgata, *A. Gray, l. c.* 315.—Calif.

LESSONIA, Bert. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 352, in syn.; et in Deless. Ic. Sel. iii. 45. t. 78 (1837) = **Eryngium**, Linn. (Umbellif.).
bupleurifolia, Bert. ex Deless. Ic. Sel. iii. 45 = seq.
bupleuroides, Bert. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 352 = E. bupleuroides.
sarcophyllum, Steud. Nom. ed. II. ii. 34 = E. sarcophyllum.

LESTADIA, Spach, Hist. Vég. Phan. x. 28 (1841) = **Laestadia**, Kunth (Compos.).

LESTIBODEA, Neck. Elem. i. 40 (1790) = **Dimorphotheca**, Moench (Compos.).

LESTIBONDESIA, Reichb. in Moessl. Handb. ed. II. i. p. lii (1827), sphalm. = seq.

LESTIBOUDESIA, Reichb. Consp. 164 (1828) = LESTIBUDESIA, Thou. = **Celosia**, Linn. (Amarant.).

LESTIBUDAEA, Juss. in Dict. Sc. Nat. xxvi. 140 (1823) = **Dimorphotheca**, Moench (Compos.).

LESTIBUDESIA, Thou. Hist. Vég. Iles Austr. Afr. 53. t. 16 (1806) = **Celosia**, Linn. (Amarant.).
altissima, A. Cunn. ex Moq. in DC. Prod. xiii. II. 253 = Deeringia altissima.
arborescens, R. Br. Prod. 414 = Deeringia altissima.
caffra, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 549 (err. typ. 449) = C. virgata?
latifolia, Blume, Bijdr. 541 = C. latifolia.
paniculata, R. Br. Prod. 414 = C. paniculata.
paniculata, Herb. ex Moq. in DC. Prod. xiii. II. 250 = Chamissoa altissima.
philippica, Weinm. in Syll. Ratisb. i. (1824) 118 = C. philippica.
recurva, Moq. in DC. Prod. xiii. II. 247 = Hermbstaedtia elegans.
spicata, Thou. Hist. Vég. Iles Austr. Afr. t. 16 = C. spicata.
trigyna, R. Br. Prod. 414 = C. trigyna.
virgata, R. Br. l. c. = C. virgata.

LETHEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = **Disporum**, Salisb. (Liliac.).

LETHEDON, Spreng. Fl. Hal. Mant. 53 (1807) = **Litsea**?, Lam. (Laurin.).
tannensis, Spreng. l. c.—Ins. Tanna.

LETSOMIA, Reichb. Handb. 302 (1837) = LETTSOMIA, Ruiz & Pav. = **Freziera**, Sw. (Ternstroem.).

LETTSOMIA, Roxb. Hort. Beng. [13] (1814); Fl. Ind. i. 487 (1832). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 870.
adpressa, *Miq. Fl. Ind. Bat.* ii. 591.—Ins. Penang.
aggregata, *Roxb. Hort. Beng.* [13]; *Fl. Ind.* i. 488.—Ind. or.; Burma.
argentea, Roxb. l. c. 489 = Argyreia argentea.
atropurpurea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 194.—Reg. Himal.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 561.—Ins. Banca.
barbata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 195.—Burma.
barbigera, *Benth. & Hook. f. Gen.* ii. 870; *C. B. Clarke, l. c.* 193.—Assam; Burma.
bella, *C. B. Clarke, l. c.* 192.—Ind. or.
Bona-nox, Roxb. Hort. Beng. [13]; Fl. Ind. i. 494 = Rivea hypocrateriformis.
bracteosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 192.—Burma.
capitata, Benth. & Hook. f. Gen. ii. 870 = strigosa.
capitata, Miq. Fl. Ind. Bat. ii. 591 = strigosa.
Chalmersii, *Hance, in Journ. Bot.* xvi. (1878) 230.—China.
Championi, *Benth. & Hook. f. Gen.* ii. 869.—Hongkong.
cuneata, Roxb. Hort. Beng. [13]; Fl. Ind. i. 491 = Argyreia cuneata.
cymosa, Roxb. l. c. 492 = Argyreia cymosa.
elliptica, *Wight, Ic.* t. 1356, in nota.—Ind. or.
festiva, *Benth. & Hook. f. Gen.* ii. 870.—China.
guianensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1092, nomen.—Guian.
hancorniaefolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 196.—Zeylan.
hirsutissima, *C. B. Clarke, l. c.* 193.—Reg. Himal.
hispida, Hook. ex Choisy, Conv. Or. 40 = Argyreia mollis.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 196.—Burma.
Maingayi, *C. B. Clarke, l. c.* 195.—Malaya.
mysorensis, *C. B. Clarke, l. c.* 192.—Ind. or.
nervosa, Hort. Calc. ex C. B. Clarke, l. c. 187 = Argyreia Wallichii.
nervosa, Roxb. Hort. Beng. [13]; Fl. Ind. i. 488 = Argyreia speciosa.
ornata, Roxb. l. c. 496 = Rivea ornata.
peguensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 193.—Burma.
penangiana, *Miq. Fl. Ind. Bat.* ii. 592.—Ins. Penang.
pomacea, Roxb. Hort. Beng. [13]; Fl. Ind. i. 493 = Argyreia pomacea.
rubens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 195.—Assam.
rubicunda, *C. B. Clarke, l. c.*—Malacca.
setosa, *Roxb. Hort. Beng.* [13]; *Fl. Ind.* i. 490.—Ind. or.

LETTSOMIA:—
sikkimensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 194.—Reg. Himal.
speciosa, Roxb. Hort. Beng. [13], nomen = Argyreia speciosa, *Sweet.*
splendens, Roxb. l. c.; Fl. Ind. i. 487 = Argyreia splendens.
strigosa, *Roxb. ll. cc.*—Ind. or.; Malaya.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 560.—Sumatra.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 193.—Reg. Himal.
tomentella, *Miq. Fl. Ind. Bat. Suppl.* 560.—Sumatra.
uniflora, Roxb. Hort. Beng. 84; Fl. Ind. i. 495 = Rivea hypocrateriformis.

LETTSOMIA, Ruiz & Pav. Prod. 77. t. 14 (1794) = **Freziera**, Sw. (Ternstroem.).
candicans, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 125 = F. candicans.
Dombeyana, Choisy, l. c. = F. Dombeyana.
lanata, Ruiz & Pav. Syst. Veg. 135 = F. canescens.
longipes, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 125 = F. longipes.
tomentosa, Ruiz & Pav. Syst. Veg. 134 = F. chrysophylla.

LEUCADENDRON, Berg. in Vet. Akad. Handl. Stockh. (1766) 325, partim; R. Br. in Trans. Linn. Soc. x. (1810) 50. *PROTEACEAE*, Benth. & Hook. f. iii. 169.
CHASME, Salisb. in Knight, Prot. 16 (1809).
CONOCARPUS, Adans. Fam. ii. 284 (1763).
EURYSPERMUM, Salisb. Parad. Lond. t. 75 (1807); et in Knight, Prot. 18 (1809).
GISSONIA, Salisb. l. c. sub t. 67 (1806); et in Knight, l. c. 32 (1809).
LEUCANDRUM, Neck. Elem. i. 106 (1790).
PROTEA, Linn. Gen. ed. II. 38 (1742); ed. V. 41 (1754).
VIONAEA, Neck, Elem. i. 107 (1790).
abietinum, *R. Br. in Trans. Linn. Soc.* x. (1810) 64.—Afr. austr.
abietinum, Drège, ex Meissn. in DC. Prod. xiv. 224 = aemulum.
acuminatum, Buek, ex Meissn. l. c. xiv. 223 = concinnum.
acutum, *Meissn. l. c.* 228.—Afr. austr.
adscendens, *R. Br. in Trans. Linn. Soc.* x. (1810) 61.—Afr. austr.
aemulum, *R. Br. l. c.* 64.—Afr. austr.
angustatum, *R. Br. l. c.* 54.—Afr. austr.
angustatum, Drège, ex Meissn. in DC. Prod. xiv. 215 = imbricatum.
argenteum, *R. Br. in Trans. Linn. Soc.* x. (1810) 52.—Afr. austr.
aurantiacum, *Buek, ex Meissn. in DC. Prod.* xiv. 217.—Afr. austr.
brunioides, *Meissn. l. c.* 215.—Afr. austr.
Buekianum, *Meissn. l. c.* 214.—Afr. austr.
buxifolium, *R. Br. in Trans. Linn. Soc.* x. (1810) 55.—Afr. austr.
callosum, *Hoffmgg. Verz. Pfl.* 72 (Quid?).—Hab.?
canaliculatum, E. Mey. ex Meissn. in DC. Prod. xiv. 215 = imbricatum.
cartilagineum, *R. Br. in Trans. Linn. Soc.* x. (1810) 67.—Afr. austr.
caudatum, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 115.—Hab.?
ciliatum, Buek, ex Meissn. in DC. Prod. xiv. 219 = concolor.
cinereum, *R. Br. in Trans. Linn. Soc.* x. (1810) 57.—Afr. austr.
cinereum, E. Mey. ex Meissn. in DC. Prod. xiv. 217 = nitidum.
comosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 64.—Afr. austr.
concinnum, *R. Br. l. c.* 61.—Afr. austr.
concolor, *R. Br. l. c.* 58.—Afr. austr.
coniferum, *Meissn. in DC. Prod.* xiv. 227.—Afr. austr.
coniferum, Sieber, ex Meissn. l. c. 222 = virgatum.
corymbosum, *Berg. in Vet. Akad. Handl. Stockh.* (1766) 325.—Afr. austr.
crassifolium, *R. Br. in Trans. Linn. Soc.* x. (1810) 66.—Afr. austr.
crinitum, Steud. Nom. ed. II. ii. 399, in syn. = Leucospermum crinitum.
cuspidatum, Klotzsch, ex Meissn. in DC. Prod. xiv. 223 = salignum.
daphnoides, *Meissn. l. c.* 226.—Afr. austr.

LEUCADENDRON:—
decorum, *R. Br. in Trans. Linn. Soc.* x. (1810) 58.—Afr. austr.
decurrens, *R. Br. l. c.* 59.—Afr. austr.
discolor, Buek, ex Meissn. in DC. Prod. xiv. 214 = Buekianum.
divaricatum, Berg. in Vet. Akad. Handl. Stockh. (1766) 324 = Mimetes divaricata.
Dregei, *E. Mey. ex Meissn. in DC. Prod.* xiv. 217.—Afr. austr.
dubium, Buek, ex Meissn. l. c. 215 = buxifolium.
elatum, *Buek, ex Meissn. l. c.* 217.—Afr. austr.
elongatum, Berg. in Vet. Akad. Handl. Stockh. (1766) 327 = Serruria elongata.
ericifolium, *R. Br. in Trans. Linn. Soc.* x. (1810) 66.—Afr. austr.
ericifolium, Drège, ex Meissn. in DC. Prod. xiv. 224 = comosum.
eucalyptifolium, *Buek, ex Meissn. l. c.* 221.—Afr. austr.
flavescens, *Link, Enum. Hort. Berol.* i. 115, nomen.—Hab.?
floridum, *R. Br. in Trans. Linn. Soc.* x. (1810) 63.—Afr. austr.
floridum, Drège, ex Meissn. in DC. Prod. xiv. 222 = lanigerum.
fusciflorum, *R. Br. in Trans. Linn. Soc.* x. (1810) 216.—Afr. austr.
glabrum, *R. Br. l. c.* 60.—Afr. austr.
Globularia, *R. Br. l. c.* 65.—Afr. austr.
grandiflorum, *R. Br. l. c.* 59.—Afr. austr.
grandiflorum, Buek, ex Meissn. in DC. Prod. xiv. 219 = concolor.
heterophyllum, E. Mey. ex Meissn. l. c. 222 = lanigenum.
hirsutum, Hoffmgg. Verz. Pfl. 72 = L. Levisanus.
hirsutum, *Meissn. in DC. Prod.* xiv. 227.—Afr. austr.
humifusum, E. Mey. ex Meissn. l. c. 220 = ovale.
imbricatum, *R. Br. in Trans. Linn. Soc.* x. (1810) 55.—Afr. trop.
inflexum, Klotzsch, ex Meissn. in DC. Prod. xiv. 215 = brunioides.
inflexum, Klotzsch, ex Meissn. l. c. 216 = tortum.
inflexum, *Link, Enum. Hort. Berol.* i. 115 = salignum.
insigne, Buek, ex Meissn. in DC. Prod. xiv. 219 = concolor.
involucratum, *Meissn. l. c.* 228.—Afr. austr.
laeve, *Meissn. l. c.* 227.—Afr. austr.
lanigenum, *Buek, ex Meissn. l. c.* 222.—Afr. austr.
Levisanus, *Berg. in Vet. Akad. Handl. Stockh.* (1766) 324.—Afr. austr.
lineare, Steud. Nom. ed. II. ii. 34 = angustatum.
linifolium, *R. Br. in Trans. Linn. Soc.* x. (1810) 216.—Afr. austr.
marginatum, *Link, Enum. Hort. Berol.* i. 115.—Afr. austr.
Meyerianum, Buek, ex Meissn. in DC. Prod. xiv. 215 = brunioides.
nanum, Berg. in Vet. Akad. Handl. Stockh. (1766) 525 = Protea rosacea.
nitidum, *Buek, ex Meissn. in DC. Prod.* xiv. 217.—Afr. austr.
obtusatum, *Meissn. l. c.* 227.—Afr. austr.
odoratum, *Steud. Nom.* ed. II. ii. 35.—Afr. austr.
oleaefolium, Berg. in Vet. Akad. Handl. Stockh. (1766) 320 = Leucospermum oleaefolium.
ovale, *R. Br. in Trans. Linn. Soc.* x. (1810) 59.—Afr. austr.
pendunculatum, *Meissn. in DC. Prod.* xiv. 216.—Afr. austr.
phylicoides, Berg. in Vet. Akad. Handl. Stockh. (1766) 328 = Serruria phylicoides.
pinifolium, DC. ex Meissn. in DC. Prod. xiv. 241 = Protea rosacea.
platyspermum, *R. Br. in Trans. Linn. Soc.* x. (1810) 63.—Afr. austr.
plumosum, *R. Br. l. c.* 58.—Afr. austr.
polygaloides, Link, Enum. Hort. Berol. i. 115, nomen = imbricatum.
prostratum, *Meissn. in DC. Prod.* xiv. 227.—Afr. austr.
proteoides, E. Mey. ex Meissn. l. c. 217 = aurantiacum.
pruinosum, Mund, ex Meissn. l. c. 216 = tortum.
pubescens, *R. Br. in Trans. Linn. Soc.* x. (1810) 66.—Afr. austr.
pyramidale, *Steud. Nom.* ed. II. ii. 35.—Afr. austr.
ramosissimum, *Buek, ex Meissn. in DC. Prod.* xiv. 221.—Afr. austr.
reticulatum, *Meissn. l. c.* 227.—Afr. austr.

LEUCADENDRON :—
retusum, *R. Br. in Trans. Linn. Soc.* x. (1810) 53.—Afr. austr.
retusum, Drège, ex Meissn. in DC. Prod. xiv. 219 = concolor.
rubricallosum, Buek, ex Meissn. l. c. 222 = lanigenum.
rugosum, *Meissn. l. c.* 227.—Afr. austr.
salignum, Berg. in Vet. Akad. Handl. Stockh. (1766) 323 = adscendens.
salignum, *R. Br. in Trans. Linn. Soc.* x. (1810) 62.—Afr. austr.
scabrum, *R. Br. l. c.* 65.—Afr. austr.
scabrum, Steud. ex Meissn. in DC. Prod. xiv. 225 = ericifolium.
scoparium, E. Mey. ex Meissn. l. c. 305 = Sorocephalus Dregei.
sericeum, *R. Br. in Trans. Linn. Soc.* x. (1810) 65.—Afr. austr.
sessile, *R. Br. l. c.* 54.—Afr. austr.
spathulatum, *R. Br. l. c.*—Afr. austr.
spathulatum, Buek, ex Meissn. in DC. Prod. xiv. 226 = crassifolium.
sphaerocephalum, Berg. in Vet. Akad. Handl. Stockh. (1766) 327 = Serruria Bergii.
spicatum, Berg. l. c. 367 = Nivenia spicata.
squarrosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 58.—Afr. austr.
stellare, Steud. Nom. ed. I. 475 = fusciflorum.
stellatum, Sweet, Hort. Brit. ed. I. 345 = fusciflorum.
strictum, *R. Br. in Trans. Linn. Soc.* x. (1810) 60.—Afr. austr.
tenuifolium, *Meissn. in DC. Prod.* xiv. 227.—Afr. austr.
Thunbergii, Endl. Gen. Suppl. iv. II. 75 = Protea glabra.
thymelaeoides, Berg. in Vet. Akad. Handl. Stockh. (1766) 324 = Mimetes thymelaeoides.
tortum, *R. Br. in Trans. Linn. Soc.* x. (1810) 56.—Afr. austr.
truncatum, *Meissn. in DC. Prod.* xiv. 228.—Afr. austr.
uliginosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 63.—Afr. austr.
undulatum, *Link, Enum. Hort. Berol.* i. 115, nomen.—Afr. austr.?
venosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 59.—Afr. austr.
verticillatum, *Meissn. in DC. Prod.* xiv. 228.—Afr. austr.
virgatum, *R. Br. in Trans. Linn. Soc.* x. (1810) 60.—Afr. austr.
virgatum, Drège, ex Meissn. in DC. Prod. xiv. 222 = adscendens.

LEUCADENDRON, Linn. Syst. ed. II. 18 (1740) = **Protea**, Linn.
acaulon, Linn. Sp. Pl. 92 = P. acaulis.
aureum, Burm. f. Fl. Cap. Prod. 4 (Quid?).—Afr. austr.
bellidifolium, Knight, Prot. 56.—Afr. austr.
calligerum, Knight, l. c. 60 = Leucospermum puberum.
cancellatum, Linn. Sp. Pl. 91 = Aulax pinifolia.
cervinum, Knight, Prot. 55 = Leucospermum conocarpum.
cinaroides, Linn. Sp. Pl. 92 = P. cynaroides, grandiflora.
Conocarpodendron, Linn. l. c. 93 = Leucospermum conocarpum.
cordifolium, Knight, Prot. 54.—Afr. austr.
crassicaule, Knight, l. c. 55 = Leucospermum conocarpum.
criniflorum, Knight, l. c. 58 = Leucospermum crinitum.
cucullatum, Linn. Sp. Pl. 93 = Mimetes cucullata.
cuneiforme, Burm. f. Fl. Cap. Prod. 4.—Afr. austr.
cyanoides, Linn. Sp. Pl. 93 = Serruria cyanoides.
ellipticum, Knight, Prot. 53 = Leucospermum ellipticum.
fallax, Knight, l. c. 52 = Leucospermum lineare.
filamentosum, Burm. f. Fl. Cap. Prod. 4.—Afr. austr.
formosum, Knight, Prot. 54 = Leucospermum medium.
glomeratum, Herb. Linn. ex Meissn. in DC. Prod. xiv. 285 = Serruria furcellata.
glomeratum, Linn. Sp. Pl. ed. II. 134 = Serruria glomerata.
glomiflorum, Knight, Prot. 59.—Afr. austr.
gnaphaliifolium, Knight, l. c. 60.—Afr. austr.
gracile, Knight, l. c. 59.—Afr. austr.
grandiflorum, Salisb. Parad. Lond. t. 116 = Leucospermum grandiflorum.

LEUCADENDRON :—
hirtum, Linn. Pl. Afr. Rar. 8; Amoen. Acad. vi. 83 = Mimetes hirta.
horizontale, Knight, Prot. 53 = Leucospermum tottum.
Hypophyllocarpodendron, Linn. Sp. Pl. 93 = Leucospermum Hypophyllum.
hypophyllum, Knight, Prot. 56 = praec.
Lepidocarpodendron, Linn. Sp. Pl. 91 = P. Lepidocarpodendron.
lineare, Burm. f. Fl. Cap. Prod. 4 = Leucospermum lineare.
parile, Knight, Prot. 57.—Afr. austr.
phyllanthifolium, Knight, l. c. 55 = Leucospermum Zeyheri.
pinifolium, Linn. Mant. i. 36 = Aulax pinifolia.
polifolium, Burm. f. Fl. Cap. Prod. 4.—Afr. austr.
proteoides, Linn. Sp. Pl. 91 = Mimetes purpurea.
puberum, Knight, Prot. 61 = Leucospermum puberum.
pubigerum, Linn. ex Meissn. in DC. Prod. xiv. 229, 259 = Leucospermum puberum.
racemosum, Linn. Sp. Pl. 91 = Spatalla ramulosa.
repens, Linn. l. c. = P. repens.
rodolentum, Knight, Prot. 58 = Leucospermum tomentosum.
royenaefolium, Knight, Prot. 59.—Afr. austr.
rubrum, Burm. f. Fl. Cap. Prod. 4.—Afr. austr.
saxatile, Knight, Prot. 58.—Afr. austr.
scolymocephalum, Linn. Sp. Pl. 92 = P. Scolymus.
Serraria, Burm. ex Meissn. in DC. Prod. xiv. 292 = Serruria phylicoides.
Serraria, Linn. Sp. Pl. 93 = Serruria Burmanni, glomerata.
speciosum, Linn. Mant. i. 36 = P. speciosa.
splendens, Burm. f. Fl. Cap. Prod. 4.—Afr. austr.
tomentosum, Knight, Prot. 57 = Leucospermum tomentosum.
truncatulum, Knight, l. c. 61 = Leucospermum puberum.
xeranthemifolium, Knight, l. c. 60.—Afr. austr.

LEUCADENDRUM, Salisb. Parad. Lond. sub t. 67 (1807) = **Leucospermum**, R. Br. (Proteac.).

LEUCADENIA, Klotzsch, ex Baill. Adansonia, iv. (1864) 338 = **Croton**, Linn. (Euphorb.).
pilosa, Klotzsch, ex Baill. l. c. = C. fuscus.

LEUCADENIUM, Klotzsch, ex Benth. & Hook. f. Gen. iii. 295 (1880) = praec.

LEUCAENA, Benth. in Hook. Journ. Bot. iv. (1842) 416. *LEGUMINOSAE*, Benth. & Hook. f. i. 594.
canescens, *Benth. Pl. Hartw.* 117.—Guatemal.
diversifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 417.—Mexic.
esculenta, *Benth. in Trans. Linn. Soc.* xxx. (1875) 442.—Mexic.
formosa, Griseb. Cat. Pl. Cub. 82 = Lysiloma Sabicu.
Forsteri, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 94.—Ins. Pacif.
glauca, *Benth. in Hook. Journ. Bot.* iv. (1842) 416.—Amphig. trop.
macrophylla, *Benth. Bot. Voy. Sulph.* 90.—Mexic.
pulverulenta, *Benth. in Hook. Journ. Bot.* iv. (1842) 417.—Mexic.
retusa, *Benth. in A. Gray, Pl. Wright.* i. 64.—Texas.
trichodes, *Benth. in Hook. Journ. Bot.* iv. (1842) 417.—Peruv.

LEUCAERIA, DC. in Ann. Mus. Par. xix. (1812) 66 = **Leucheria**, Lag. (Compos.).

LEUCAMPYX, A. Gray, ex Benth. & Hook. f. Gen. ii. 422 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
Newberryi, *A. Gray, ex Porter & Coult. Fl. Colorad.* 77.—Am. bor. occ.

LEUCANDRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 188 = **Tragia**, Plum. (Euphorb.).
betonicaefolia, Klotzsch, l. c. = T. betonicaefolia.

LEUCANDRON, Steud. Nom. ed. II. ii. 35 (1841) = seq.

LEUCANDRUM, Neck. Elem. i. 106 (1790) = **Leucadendron**, Berg. (Proteac.).

LEUCANOTIS, D. Dietr. Syn. Pl. i. 418 (1839) = **Leuconotis**, Jack (Apocyn.).

LEUCANTHA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821) = **Centaurea**, Linn. (Compos.).
cyanifolia, S. F. Gray, l. c. 444 = C. solstitialis.

LEUCANTHEA, Scheele, in Linnaea, xxv. (1852) 258 = **Bouchetia**, DC. (Scrophul.).
Roemeriana, Scheele, l. c. 259 = B. erecta.

LEUCANTHEMUM, [Tourn.] Linn. Syst. ed. I. (1735) = **Chrysanthemum**, Tourn. (Compos.).
affine, Nym. Consp. 371 = C. montanum.
affine, Pomel, Nouv. Mat. Fl. Atl. 58.—Alger.
algidum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) 1. 416 = C. integrifolium.
alpinum, Lam. Fl. Fr. ii. 138 = C. alpinum.
anomalum, DC. Prod. vi. 49 = C. oppositifolium.
arcticum, DC. l. c. 45 = C. arcticum.
arenarium, Pomel, Nouv. Mat. Fl. Atl. 58.—Alger.
armenum, DC. Prod. vii. 297 = C. fruticulosum.
atratum, DC. l. c. vi. 48 = C. atratum.
atratum, Nym. Consp. 371 = C. montanum.
auriculatum, Nym. l. c. = C. montanum.
Barrelieri, Timb. in Bull. Soc. Bot. Fr. xiii. (1866) p. cliii = C. atratum?
Bauhini, Steud. Nom. ed. II. ii. 35 = C. atratum.
Candolleanum, Martr. Fl. Tarn, 356 = C. pallens.
canescens, Pourr. ex Nym. Consp. 370 = C. montanum.
cebennense, DC. Prod. vi. 48 = C. monspeliense.
ceratophylloides, Nym. Syll. 10 = C. atratum.
Chamaemelum, Lam. Fl. Fr. ii. 139 = Matricaria Chamomilla.
commutatum, Timb. & Martr. in Martr. Pl. Crit. Tarn, 29 = C. montanum.
coronopifolium, Vill. Prosp. 32 = C. coronopifolium.
corsicum, DC. Prod. vi. 47 = C. coronopifolium, *Vill.*
corymbosum, Gren. & Godr. Fl. Fr. ii. 145 = C. corymbosum.
crassifolium, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 96.—Hispan.
decipiens, Pomel, Nouv. Mat. Fl. Atl. 59.—Alger.
Delarbrei, Timb. in Lamotte, Prod. Fl. Centr. Fr. ii. 404 = C. Leucanthemum.
discoideum, Bourg. ex Nym. Consp. 370 = C. montanum.
Fontanesii, Boiss. & Reut. Diagn. Ser. II. iii. 26.—Alger.
Gaudini, Dalla Torre, Atl. Alpenfl. 130.—Europ.
glabrum, Boiss. & Reut. Pugill. Pl. Nov. 57 = C. paludosum.
Gmelini, Ledeb. Fl. Ross. ii. 541 = C. arcticum?
gracilicaule, Nym. Consp. 371 = C. montanum.
graminifolium, Lam. Fl. Fr. ii. 137 = C. graminifolium.
Gussonii, Nym. Consp. 371 = C. graminifolium.
Herminii, Nym. l. c. 372 = C. oppositifolium.
heterophyllum, DC. Prod. vi. 47 = C. lanceolatum.
heterophyllum, Schur, ex Nym. Consp. 370 = C. pallens.
integrifolium, DC. Prod. vi. 46 = C. integrifolium.
ircutianum, DC. l. c. 47 = C. ircutianum.
lanceolatum, DC. l. c. = C. lanceolatum.
lanceolatum, Nym. Consp. 371 = C. montanum.
latifolium, DC. Prod. vi. 47 = C. lacustre.
maximum, DC. l. c. 46 = C. maximum.
minimum, Vill. Prosp. 32 = C. alpinum.
montanum, DC. Prod. vi. 48 = C. montanum.
murcicum, J. Gay, ex Willk. in Bot. Zeit. xxii. (1864) 253 = C. glabrum.
nipponicum, Franch. ex Maxim. in Bull. Acad. Pétersb. xvii. (1872) 420.—Japon.
odoratum, Dulac, Fl. Hautes-Pyr. 504 = C. Parthenium.
pallens, DC. Prod. vi. 47 = C. pallens.
palmatum, Lam. Fl. Fr. iii. 138 = C. monspeliense.
Parthenium, Gren. & Godr. Fl. Fr. ii. 145 = C. Parthenium.
pectinatum, DC. Prod. vi. 49 = C. pectinatum.
persicaefolium, Timb. in Rod. & Bot. Agr. ii. 447; ex ej. in Bull. Soc. Sc. Phys. & Nat. Toul. ii. (1874) 460.—Europ.
platylepis, Borb.—Cf. Oestr. Bot. Zeitschr. xxviii. (1878) 259 = C. lanceolatum.
Reboudianum, Pomel, Nouv. Mat. Fl. Atl. 291.—Alger.
rotundifolium, DC. Prod. vi. 46 = C. rotundifolium.
rotundifolium, Schur, Enum. Pl. Transs. 339 = Tanacetum rotundifolium.
sericeum, Nym. Syll. 10 = C. oppositifolium.
setabense, DC. Prod. vi. 49 = C. paludosum.

LEUCANTHEMUM :—
sibiricum, DC. Prod. vi. 46 = C. arcticum.
silvaticum, Willk. & Lange, Prod. Fl. Hisp. ii. 107, sphalm. = montanum.
simplex, Boiss. Fl. Orient. iii. 341 = C. palustre.
sinuatum, DC. Prod. vi. 46 = C. sinuatum.
subglaucum, De Laramb. Soc. Litt. Castres (1861); ex Martr. Fl. Tarn, 356 = C. pallens.
suffruticulosum, Timb. in Bull. Soc. Phys. & Nat. Toul. ii. (1874) 69 = C. montanum.
tomentosum, Gren. & Godr. Fl. Fr. ii. 144 = C. tomentosum.
varians, Martr. Pl. Crit. Tarn, 30; ex Fl. Tarn, 356 = C. monspeliense.
virgatum, Clos, in Bull. Soc. Bot. Fr. xvii. (1870) 185 = C. flosculosum.
vulgare, Lam. Fl. Fr. ii. 137 = C. Leucanthemum.
Weyrichii, Maxim. Prim. Fl. Amur. 156.—Ins. Sachalin.
Zawadskii, Nym. Consp. 372 = C. arcticum.

LEUCAS, Burm. Thes. Zeyl. 140 (1737); R. Br. Prod. 504 (1810). *LABIATAE*, Benth. & Hook. f. ii. 1213.
BLANDINA, Rafin. Fl. Tellur. iii. 88 (1836).
ELBUNIS, Rafin. l. c. (1836).
ENEODON, Rafin. l. c. (1836).
HEMISTEMMA, Reichb. Consp. 116 (1828).
HEMISTOMA, Ehrenb. ex Benth. in Wall. Pl. As. Rar. i. 60 (1830).
HEPTRILIS, Rafin. Fl. Tellur. iii. 89 (1836).
HETREPTA, Rafin. l. c. 88 (1836).
ISODECA, Rafin. l. c. (1836).
LEUCASIA, Rafin. l. c. (1836).
LOXOSTOMA, Benth. ex Steud. Nom. ed. II. ii. 75 (1841).
PHYSOLEUCAS, Jaub. & Spach, Illustr. v. 48. t. 445 (1855 ?).
acrodonta, *Steud. Nom.* ed. II. ii. 36.—Arab.
affinis, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyssin.
angularis, *Benth. in Wall. Pl. As. Rar.* i. 62.—Ind. or.; Malaya.
angustifolia, Wall. l. c. = mollissima.
aspera, *Link, Enum. Hort. Berol.* ii. 113; *Spreng. Syst.* ii. 743.—Ind. or.; Malaya.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 572.—Malaya.
Benthamiana, *Hook. & Arn. Bot. Beech. Voy.* 204.—China.
biflora, Benth. in Wall. Cat. n. 2063, partim = procumbens.
biflora, *R. Br. Prod.* 504.—Ind. or.
calostachys, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 139.—Afr. trop.
capitata, *Desf. in Mém. Mus. Par.* xi. (1824) 8. t. 4.—Ind. or.
Cephalotes, Spreng. Syst ii. 743 = capitata.
chinensis, *R. Br. Prod.* 504.—Ind. or.; China.
ciliata, *Benth. in Wall. Pl. As. Rar.* i. 61.—Ind. or.
ciliata, Hochst. ex Benth. in DC. Prod. xii. 530 = nubica.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 688.—Ind. or.
collina, Dalz. in Hook. Kew Journ. ii. (1850) 338 = lanata.
decemdentata, R. Br. Prod. 504 = flaccida.
decurva, Benth. in Wall. Pl. As. Rar. i. 60 = nutans.
deflexa, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 213.—Afr. trop.
densiflora, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 95.—Afr. trop.
diffusa, *Benth. Lab. Gen. et Sp.* 615.—Ind. or.
dimidiata, Benth. in DC. Prod. xii. 532 = aspera.
dimidiata, *Spreng. Syst.* ii. 743.—Ind. or.
ebracteata, *Peyr. in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 577.—Afr. trop.
eriostoma, *Hook. f. Fl. Brit. Ind.* iv. 686—Ind. or.
flaccida, *R. Br. Prod.* 505.—Ind. or.; Austral.
flaccida, Mor. & Zoll. Verz. Pfl. Zoll. 54 = biflora.
galeopsidea, Hochst. ex Benth. in DC. Prod. xii. 524 = glabrata.
glaberrima, *Jaub. & Spach, Illustr.* iv. 131. t. 387.—Oriens.
glabrata, *R. Br. Prod.* 504.—Arab.
globulifera, *Hassk. Cat. Hort. Bog. Alt.* 133.—Java?
grandis, *Vatke, in Linnaea*, xliii. (1880–82) 96.—Afr. trop.
haematula, Wight & Arn. ex Steud. Nom. ed. II. ii. 36 = seq.
hamatula, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 355 = stelligera.

LEUCAS:—
Hamiltoniana, Benth. in Wall. Pl. As. Rar. i. 62 = montana.
Helferi, *Hook. f. Fl. Brit. Ind.* iv. 681.—Burma.
helianthemifolia, Benth. in Wall. Pl. As. Rar. i. 61 = hirta.
helianthemifolia, *Desf. in Mém. Mus. Par.* xi. (1824) 2. t. 1.—Ind. or.
hirta, *Spreng. Syst.* ii. 743.—Ind. or.
hyssopifolia, *Benth. in Wall. Pl. As. Rar.* i. 60.—Reg. Himal.
indica, *R. Br. Prod.* 504.—Ind. or.
indica, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 95.—Afr. trop.
inflata, *Benth. Lab. Gen. et Sp.* 744.—Nubia; Arab.
inflata, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 36 = acrodonta.
involucrata, Benth. in Wall. Cat. n. 2047 = zeylanica.
javanica, *Benth. Lab. Gen. et Sp.* 611.—Java.
lamiifolia, *Desf. in Mém. Mus Par.* xi. (1824) 4.—Ind. or.
lamiifolia, Zipp. ex Span. in Linnaea, xv. (1841) 333 = chinensis.
lanata, *Benth. in Wall. Pl. As. Rar.* i. 61.—Ind. or.; China.
lanceaefolia, *Desf. in Mém. Mus. Par.* xi. (1824) 5.—Ind. or.
lanciformis, Wall. Cat. n. 2759 = praec.
lavandulaefolia, *Sm. in Rees, Cycl.* xx. n. 2.—Ind. or.; Malaya.
lepistoma, Wight, ex Hook. f. Fl. Brit. Ind. iv. 686 = eriostoma.
leucocephala, *Miq. Fl. Ind. Bat.* ii. 981.—Java.
linifolia, Spreng. Syst. ii. 743 = lavandulaefolia.
longifolia, *Benth. Lab. Gen. et Sp.* 744.—Ind. or.
malayana, *Hance, in Walp. Ann.* iii. 269.—Singapore.
marrubioides, *Desf. in Mém. Mus. Par.* xi. (1824) 6. t. 3. f. 1.—Ind. or.
marrubioides, Wall. Cat. n. 2524 = pubescens.
martinicensis, *R. Br. Prod.* 504.—Amphig. trop.
masaiensis, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 403.—Afr. trop.
melissaeflora, Benth. in Wall. Cat. n. 2062 = seq.
melissaefolia, Benth. in Wall. Pl. As. Rar. i. 62 = flaccida.
membranacea, Benth. in Wall. Cat. n. 2052 = urticaefolia.
microphylla, *Vatke, in Linnaea*, xliii. (1880–82) 97.—Afr. trop.
mollissima, *Wall. Pl. As. Rar.* i. 62.—Ind. or.; China.
moluccoides, *Sm. in Rees, Cycl.* xx. n. 9.—Arab.
montana, *Spreng. Syst.* ii. 742.—Ind. or.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 85.—Afr. austr.
nepetaefolia, *Benth. in Wall. Pl. As. Rar.* i. 62.—Ind. or.
Neufliseana, *Courb. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 145.—Abyssin.
nubica, *Benth. in DC. Prod.* xii. 530.—Nubia.
nutans, Benth. in Wall. Pl. As. Rar. i. 60 = dimidiata.
nutans, *Spreng. Syst.* ii. 743.—Ind. or.
oligocephala, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 213.—Afr. trop.
ovata, *Benth. in Wall. Pl. As. Rar.* i. 61.—Burma.
oxydon, *Miq. Fl. Ind. Bat.* ii. 981.—Java.
parviflora, Benth. in Wall. Pl. As. Rar. i. 62 = flaccida.
paucicrenata, *Vatke, in Linnaea*, xliii. (1880–82) 98.—Afr. trop.
pilosa, *Benth. in Wall. Pl. As. Rar.* i. 62.—Burma.
Plukenetii, Spreng. Syst. ii. 743; Benth. l. c. 60 = aspera.
polycephala, Wall. Cat. n. 2051 = martinicensis.
pratensis, *Vatke, in Linnaea*, xliii. (1880–82) 97.—Afr. trop.
procumbens, *Desf. in Mém. Mus. Par.* xi. (1824) 7. t. 3. f. 2.—Ind. or.
procumbens, Thw. Enum. Pl. Zeyl. 240 = biflora.
pubescens, *Benth. Lab. Gen. et Sp.* 610.—Ind. or.
pubescens, Thw. Enum. Pl. Zeyl. 240 = angularis.
quinquedentata, R. Br. in Salt, Abyss. App. 64, nomen = Lasiocorys abyssinica.
rosmarinifolia, *Benth. in Wall. Pl. As. Rar.* i. 61.—Ind. or.
rufescens, Benth. in Wall. Cat. n. 2053 = hirta.
Schimperi, Hochst. ex A. Br. in Flora, xxiv. (1841) 1. 279 = martinicensis.
Schimperi, Presl, Bot. Bemerk. 101 = acrodonta.

LEUCAS:—
somalensis, *Vatke, in Linnaea*, xl. (1876) 181.—Afr. trop.
stachydiformis, Hochst. in Flora, xxvii. (1844) 100 = Ballota stachydiformis.
stachyoides, Spreng. Syst. ii. 743 = flaccida.
stelligera, *Wall. Pl. As. Rar.* i. 61.—Ind. or.
stricta, *Benth. in Wall. l. c.*—Ind. or.
strigosa, Benth. l. c. 62 = mollissima.
suffruticosa, *Benth. Lab. Gen. et Sp.* 611.—Ind. or.
teres, *Benth. in Wall. Pl. As. Rar.* i. 62.—Burma.
ternifolia, Desf. in Mém. Mus. Par. xi. (1824) 4 = helianthemifolia.
tettensis, *Vatke, in Linnaea*, xl. (1876) 180.—Afr. trop.
trachyphylla, *Jaub. & Spach, Illustr.* iv. 130. t. 386.—Arab.
urticaefolia, *R. Br. Prod.* 504.—Nubia; Arab.; Ind. or.
vestita, *Benth. in Wall. Pl. As. Rar.* i. 61.—Ind. or.
virgata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotr.
Walkeri, Benth. in DC. Prod. xii. 531 = zeylanica.
Wightiana, *Benth. in Wall. Pl. As. Rar.* i. 60.—Ind. or.
zeylanica, Boj. Hort. Maurit. 250 = aspera.
zeylanica, *R. Br. Prod.* 504.—Ind. or.; Malaya.

LEUCASIA, Rafin. Fl. Tellur. iii. 88 (1836) = **Leucas**, Burm. (Labiat.).
zeylanica, Rafin. l. c. = L. zeylanica.

LEUCASTER, Choisy, in Mém. Soc. Phys. Genèv. xii. (1849) 167. *NYCTAGINEAE*, Benth. & Hook. f. iii. 10.
caniflorus, *Choisy, in DC. Prod.* xiii. II. 458.—Bras.

LEUCE, Opiz, Seznam, 59 (1852) = **Populus**, Linn. (Salicin.).
alba, Opiz, l. c. = P. alba.
canescens, Opiz, l. c. = P. alba.

LEUCEA, Presl, Fl. Sicul. i. p. xxx (1826) = **Leuzea**, DC. (Compos.).

LEUCERIA, Lag. Amen. Nat. i. 32 (1811) (Leucheria). *COMPOSITAE*, Benth. & Hook. f. ii. 499.
CASSIOPEA, D. Don, in Trans. Linn. Soc. xvi. (1830) 215.
CHABRAEA, DC. in Ann. Mus. Par. xix. (1812) 65. t. 5.
EIZAGUIRREA, Remy, in C. Gay, Fl. Chil. iii. 401 (1849).
FRAGERIA, Delile, ex Steud. Nom. ed. II. i. 645 (1840).
LASIORRHIZA, Lag. Amen. Nat. i. 32 (1811).
MACLOVIA, DC. ex Steud. Nom. ed. II. i. 338, in syn. (1840).
PTILURUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 218.
abbreviata, *Steud. Nom.* ed. II. ii. 36.—Chili.
acanthoides, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 213.—Chili.
achillaeifolia, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 43.—Patagon.
amoena, *Phil. in Linnaea*, xxviii. (1856) 714.—Chili.
andryaloides, *DC. in Deless. Ic. Sel.* iv. t. 86.—Chili.
Bridgesii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 36.—Chili.
candidissima, *Gill. & D. Don, in Phil. Mag.* (1832) 389.—Chili.
cerberoana, *Remy, in C. Gay, Fl. Chil.* iii. 386.—Chili.
cinerea, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 215.—Chili.
coerulescens, *Remy, in C. Gay, Fl. Chil.* iii. 382.—Chili.
congesta, *Gill. ex D. Don, in Phil. Mag.* (1832) 389.—Chili.
conyzoides, D. Don, ex Baker, in Mart. Fl. Bras. vi. III. 391 = Trixis ochroleuca.
Cummingii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 36.—Chili.
divaricata, D. Don, in Trans. Linn. Soc. xvi. (1830) 214 = paniculata.
echioides, D. Don, in Phil. Mag. (1832) 38 = Trixis brasiliensis.
eriochlaena, Remy, in C. Gay, Fl. Chil. iii. 384.—Chili.
floribunda, *DC. in Deless. Ic. Sel.* iv. t. 88.—Chili.

LEUCERIA :—
foliosa, *Phil. in Linnaea,* xxviii. (1856) 720.—Chili.
Garciana, *Remy, in C. Gay, Fl. Chil.* iii. 381.—Chili.
Gilliesii, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 35 = cinerea.
glandulosa, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 217.—Chili.
glandulosa, *Phil. in Anal. Univ. Chil.* (1872) 743.—Chili.
gossypina, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 43.—Ins. Falkland.
hieracioides, *Cass. in Dict. Sc. Nat.* lv. 392.—Chili.
laciniata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 36.—Chili; Boliv.?
Menana, *Remy, in C. Gay, Fl. Chil.* iii. 387.—Chili.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* I. (1843) 289.—Chili.
multiflora, *Phil. in Anal. Univ. Chil.* (1872) 743.—Chili.
Neaei, *DC. Prod.* vii. 58.—Peruv.; Chili.
oligocephala, *Remy, in C. Gay, Fl. Chil.* iii. 383.—Chili.
paniculata, *Poepp. ex Less. Syn. Comp.* 403.—Chili.
peduncularis, *Remy, in C. Gay, Fl. Chil.* iii. 385.—Chili.
pulchella, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 216.—Chili.
purpurea, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 43.—Reg. Magellan.
rosea, Poepp. ex Less. Syn. Comp. 402 = andryaloides.
runcinata, *Gill. & D. Don, in Phil. Mag.* (1832) 389.—Chili.
scrobiculata, *Gill. & Don, l. c.*—Chili.
senecioides, *Hook. & Arn. Bot. Beech. Voy.* 28.—Chili.
tenuis, *Less. Syn. Comp.* 401.—Chili.
thrincioides, *Griseb. in Goett. Abh.* xix. (1874) 197.—Reg. Argent.
volcanica, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 42.—Chili.

LEUCESTERIA, Meissn. Gen. i. 155 (1838) = **Leycesteria**, Wall. (Caprifol.).

LEUCHAERIA, Less. in Linnaea, v. (1830) 10 = **Leuceria**, Lag. (Compos.).

LEUCHTENBERGIA, [Fisch.] Hook. Bot. Mag. t. 4393 (1848); Salm-Dyck, Cact. (1850) 177. *CACTEAE*, Benth. & Hook. f. i. 848.
Principis, [*Fisch.*] *ll. cc.*—Mexic.

LEUCOBLEPHARIS, Arn. in Mag. Zool. & Bot. ii. (1838) 422 = **Blepharispermum**, Wight (Compos.).
subsessilis, Arn. l. c. = B. subsessile.

LEUCOCARPON, Endl. Gen. 918, in syn. (1841) = seq.

LEUCOCARPUM, A. Rich. Sert. Astrol. 46. t. 18 (1834) = **Denhamia**, Meissn. (Celastr.).
obscurum, A. Rich. l. c. = D. obscurum.
Oleaster, F. Muell, in Trans. Roy. Soc. N. S. Wales, (1881) 15 = D. Oleaster.
pittosporoides, F. Muell, l. c. = D. pittosporoides.

LEUCOCARPUS, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 124 (1830). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 943.
alatus, *D. Don, l. c.*—Am. trop.
fruticosus, Benth. in DC. Prod. x. 336 = Hemichaena fruticosa.
perfoliatus, Benth. l. c. 335 = alatus.

LEUCOCASIA, Schott, in Oestr. Bot. Wochenbl. (1857) 34 = **Colocasia**, Schott (Aroid.).
gigantea, Schott, l. c. = C. indica.

LEUCOCEPHALA, Roxb. Hort. Beng. 68 (1814); Fl. Ind. iii. 612 (1832) = **Eriocaulon**, Linn.
graminifolia, Roxb. ll. cc. 68, 612 = E. quinquangulare.
spathacea, Roxb. ll. cc. 68, 613 = E. sexangulare.

LEUCOCERA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 582 = **Calycera**, Cav.
annua, Turcz. l. c. 583 = C. lanata (in Addend.).

LEUCOCHLAMYS, Poepp. ex Engl. in DC. Monog. Phan. ii. 230 (1879) = **Spathiphyllum**, Schott (Aroid.).
callacea, Poepp. ex Engl. l. c. = S. cannaeforme.

LEUCOCHYLE, Klotzsch, Ind. Sem. Hort. Berol. (1854) App. 1 = **Trichopilia**, Lindl. (Orchid.).
Warscewiczii, Klotzsch, l. c. = T. hymenantha.

LEUCOCNIDE, Miq. Pl. Jungh. i. 36 (1851) = **Debregeasia**, Gaudich. (Urtic.).
affinis, Miq. l. c. 39 = D. dichotoma.
alba, Miq. l. c. 37 = Leucosyke alba.
angustata, Miq. ex Blume, Mus. Bot. Lugd. Bat. ii. 156 = D. velutina.
candidissima, Miq. Pl. Jungh. i. 37 = Leucosyke javanica.
dichotoma, Miq. l. c. 38 = D. dichotoma.
sororia, Miq. l. c. = D. dichotoma.

LEUCOCOCCUS, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 311 = **Pouzolzia**, Gaudich. (Urtic.).
occidentalis, Liebm. l. c. = P. occidentalis.
orientalis, Liebm. l. c. 312, in nota = P. orientalis.

LEUCOCODON, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 4. *RUBIACEAE*, Benth. & Hook. f. ii. 67.
reticulatum, *Gardn. l. c.* 5.—Zeylan.

LEUCOMA, Ehrh. Beitr. iv. 146 (1789) = **Eriophorum**, Linn. (Cyperac.).

LEUCOCORYNE, Lindl. Bot. Reg. t. 1293 (1830). *LILIACEAE*, Benth. & Hook. f. iii. 799.
alliacea, *Lindl. l. c.* sub t. 1293.—Chili.
angustipetala, *C. Gay, Fl. Chil.* vi. 124.—Chili.
Gayi, Baker, in Journ Linn. Soc. xi. (1871) 375 = Tristagma dimorphopetala.
ixioides, *Lindl. Bot. Reg.* sub t. 1293.—Chili.
macropetala, *Phil. in Linnaea*, xxix (1857-58) 74.—Chili.
narcissiflora, Baker, in Journ. Linn. Soc. xi. (1871) 374 = ixioides.
narcissoides, Phil. Fl. Atac. 52 = ixioides.
odorata, Lindl. Bot. Reg. t. 1293 = ixioides.
purpurea, G. Gay, Fl. Chil. vi. 122 = ixioides.

LEUCOCRINUM, Nutt. ex A. Gray, in Ann. Lyc. N. York, iv. (1837) 110. *LILIACEAE*, Benth. & Hook. f. iii. 793.
LEUCRINIS, Rafin. Fl. Tellur. iv. 27 (1836).
montanum, *Nutt. ex A. Gray, l. c.* 110.—Calif.

LEUCOCROTON, Griseb. in Goett. Abh. ix. (1861) 20. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
flavescens, Benth. & Hook. f. Gen. iii. 312 = seq.
flavicans, Muell. Arg. in DC. Prod. xv. II. 757 = Wrightii, revolutus.
revolutus, *Wright, in Sauv. Fl. Cub.* 129.—Cuba.
virens, *Griseb. in Goett. Nachr.* (1865) 175.—Cuba.
Wrightii, *Griseb. in Goett. Abh.* ix. (1861) 21.—Cuba.

LEUCOCYCLUS, Boiss. Diagn. Ser. I. xi. 13 (1849) = **Anacyclus**, Linn. (Compos.).
formosus, Boiss. l. c. 14 = A. formosus.

LEUCODERMIS, Planch. ex Benth. & Hook. f. Gen. i. 357 (1862) = **Ilex**, Tourn.
javanica, Planch. ex Hook. f. Fl. Brit. Ind. i. 605 = I. cymosa.

LEUCODESMIS, Rafin. Fl. Tellur. iv. 19 (1836) = **Haemanthus**, Tourn. (Amaryll.).
pubescens, Rafin. l. c. = H. albiflos.

LEUCODICTYON, Dalz. in Hook. Kew Journ. ii. (1850) 264 = **Galactia**, P. Br. (Legumin.).
malvensis, Walp. Ann. ii. 431 = G. tenuiflora.

LEUCODONIUM, Opiz, Seznam, 59 (1852) = **Cerastium**, Dill. (Caryophyll.).
arvense, Opiz, l. c. = C. arvense.

LEUCOGLOCHIN, Heuff. in Flora, xxvii. (1844) 528 = **Carex**, Linn. (Cyperac.).
pauciflorus, Heuff. l. c. = C. pauciflora.

LEUCOHYLE, Reichb. in Walp. Ann. vi. 679 (1861) sphalm. = LEUCOCHYLE, Klotzsch = **Trichopilia**, Lindl. (Orchid.).

LEUCOIUM, Linn. Gen. ed. I. 102 (1737) = **Leucojum**, Linn. (Amaryll.).

LEUCOIUM, Tourn. ex Adans. Fam. ii. 418 (1763) = **Matthiola**, R. Br. (Crucif.).
incanum, Moench, Meth. 257 = M. incana.
monomotapense, Herb. Burm. ex Sond. in Abh. Geb. Naturw. Hamb. i. (1846) 258 = Heliophila callosa.

LEUCOJUM, Linn. Syst. ed. I (1735). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 720.
ACIS, Salisb. Parad. Lond. sub t. 74 (1807).
ERINOSMA, Herb. Amaryll. 330 (1837).
NIVARIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 421.
RUMINIA, Parl. 'Due Nuov. Gen. Monocot. 3;' ex ej. Fl. Ital. iii. 84 (1858).
aestivum, *Linn. Syst.* ed. X. 975.—Europ.; Oriens.
auctumnale, J. F. Gmel. Syst. 534 = autumnale.
autumnale, Balb. Misc. Alt. 14 = hyemale.
autumnale, Gouan, Hort. Monsp. 163 = aestivum.
autumnale, *Linn. Sp. Pl.* 289.—Reg. Mediterr.
capitulatum, *Lour. Fl. Cochinch.* 199.—Cochinch.
carpathicum, *Steud. Nom.* ed. II. ii. 37; *Salisb. Gen. Pl. Fragm.* 96.—Hungar.
grandiflorum, [DC. in] Red. Lil. t. 217 = trichophyllum.
Hernandezianum, Schult. f. Syst. vii. 784 = seq.
Hernandezii, Cambess. in Mém. Mus. Par. xiv. (1827) 315 = pulchellum.
hyemale, *DC. Fl. Fr. Suppl.* 326 (hiemale).—Europ. austr.
longifolium, *J. Gay, ex Salis, in Flora*, xvi. (1833) 491; *Gren. & Godr. Fl. Fr.* iii. 252.—Ins. Corsic.
nicaeënse, Ardoino, Fl. Alpes-Marit. 371 = hyemale.
pulchellum, *Salisb. Parad. Lond.* t. 74.—Ins. Balear.
roseum, *Mart. Bibl. Phys. Econ.* (1804) 344; *Loisel. Fl. Gall.* ed. I. 190.—Ins. Corsica.
strumosum, Ait. Hort. Kew. ed. I. i. 407 = Hessea filifolia.
tingitanum, *Baker, in Journ. Linn. Soc.* xvi. (1878) 678.—Marocco.
trichophyllum, Reichb. Ic. Pl. Crit. 703 = longifolium.
trichophyllum, *Schousb. Vextr. Marokko*, 154.—Reg. Medit. occ.
vernale, Salisb. Prod. 219 = vernum.
vernum, Gueldenst. Reise, i. 149 = aestivum.
vernum, *Linn. Sp. Pl.* 289.—Europ.

LEUCOLAENA, R. Br. App. Flind. Voy. 557 (1814) = **Xanthosia**, Rudge (Umbellif.).
candida, Benth. in Enum. Pl. Hueg. 55 = X. candida.
Huegelii, Benth. l. c. = X. Huegelii.
montana, Benth. l. c. in nota = X. pilosa.
pannosa, Benth. l. c. in nota = X. pilosa.
peltigera, Hook. Ic. Pl. t. 45 = X. peltigera.
pilosa, Benth. Fl. Austral. iii. 361 in syn. = X. pilosa.
rotundifolia, Benth. in Enum. Pl. Hueg. 55 = X. rotundifolia.
tridentata, R. Br. ex DC. Prod. iv. 75 = X. tridentata.

LEUCOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = **Linum**, Tourn.
suffruticosum, Fourr. l. c. = Linum suffruticosum.
tenuifolium, Fourr. l. c. = Linum tenuifolium.

LEUCOMALLA, Phil. Sert. Mendoc. Alt. 31 (1871) = **Evolvulus**, Linn. (Convolv.).
lanuginosa, Phil. l. c. = E. incanus.

LEUCOMERIS, D. Don, Prod. Fl. Nep. 169 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 490.
celebica, Blume, ex DC. Prod. v. 21 = Vernonia arborea.
decora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 317.—Burma.
glabra, Blume, ex DC. Prod. v. 22 = Vernonia arborea.
javanica, Blume, ex DC. l. c. = Vernonia arborea.
macrophylla, DC. l. c. 25 = spectabilis.
scandens, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126 = Pertya scandens.
spectabilis, *D. Don, Prod. Fl. Nep.* 170.—Nepal.

LEUCOMPHALOS, Benth. in Hook. Ic. Pl. t. 784 (1848). *LEGUMINOSAE*, Benth. & Hook. f. i. 553.
cappariodeus, *Benth. in Hook. Ic. Pl.* t. 784.—Afr. trop.

LEUCOMPHALUS, Benth. in Hook. Niger Fl. 322 (1849) = praec.

LEUCONOË, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323 = **Ranunculus**, Linn.
alpestris, Fourr. l. c. = R. alpestris.
crenatus, Fourr. l. c. = R. crenatus.
magellensis, Fourr. l. c. = R. alpestris?
Trauffelneri, Fourr. l. c. = R. alpestris.

LEUCONOTIS, Jack, in Trans. Linn. Soc. xiv. (1823) 121. t. 4. *APOCYNACEAE*, Benth. & Hook. f. ii. 691.
anceps, *Jack, in Trans. Linn. Soc.* xiv. (1823) 121. t. 4.—Sumatra.
cuspidatus, Blume, Mus. Bot. Lugd. Bat. i. 112 = eugenifolius.
eugenifolius, *A. DC. in DC. Prod.* viii. 331.—Ins. Penang.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 628.—Malacca.
Maingayi, *Dyer, ex Hook. f. l. c.*—Singapore.

EUCOPHAË, Webb & Berth. Phyt. Canar. iii. 99 (1836–50) = **Sideritis**, Linn. (Labiat.).
argosphacelus, Webb & Berth. l. c. 101 = S. argosphacelus.
canariensis, Webb & Berth. l. c. 103 = S. canariensis.
candicans, Webb & Berth. l. c. 100 = S. candicans.
dasygnaphala, Webb & Berth. l. c. 101 = S. dasygnaphala.
macrostachys, Webb & Berth. l. c. 102 = S. macrostachyos.
Massoniana, Webb & Berth. l. c. = S. Massoniana.

LEUCOPHORA, Ehrh. Beitr. iv. 148 (1789) = **Luzula**, DC. (Juncac.).

LEUCOPHOLIS, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 10. *COMPOSITAE*, Benth. & Hook. f. ii. 312.
latifolia, *Benth. in Hook. Ic. Pl.* t. 1115.—Bras.
phylicoides, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 10.—Bras.

LEUCOPHYLLUM, Humb. & Bonpl. Pl. Aequin. ii. 95. t. 109 (1809). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 927.
ambiguum, *Humb. & Bonpl. l. c.*—Mexic.
campanulatum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 254 = ambiguum.
minus, *A. Gray, Bot. Mex. Bound.* 115.—N. Mexic.
texanum, *Benth. in DC. Prod.* x. 344.—Texas.

LEUCOPHYTA, R. Br. in. Trans. Linn. Soc. xii. (1817) 106 = **Calocephalus**, R. Br. (Compos.).
Brownii, Cass. in Dict. Sc. Nat. xxvi. 159 = C. Brownii.

LEUCOPLOCUS, Endl. Ench. 67 (1841) sphalm. = seq.

LEUCOPLOEUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836) = **Hypodiscus**, Nees (Restiac.).
argenteus, Nees, l. c. = H. argenteus.
striatus, Nees, ex Mast. in Journ. Linn. Soc. x. (1868) 261 = H. Neesii.

LEUCOPOA, Griseb. in Ledeb. Fl. Ross. iv. 383 (1853) = **Festuca**, Tourn. (Gramin.).
sibirica, Griseb. l. c. = F. sibirica.

LEUCOPODON, Benth. & Hook. f. Gen. ii. 803 (1873) = seq.

LEUCOPODUM, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 124 = **Chevreulia**, Cass. (Compos.).
campestre, Gardn. l. c. 125 = C. acuminata.

LEUCOPOGON, R. Br. Prod. 541 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 614.
ANACYCLODON, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 49.
PENTAPLELION, Turcz. in Bull. Nat. Mosc. Soc. (1833) II. 194.
PEROA, Pers. Syn. i. 174 (1805).
PEROJOA, Cav. Ic. iv. 29. t. 349 (1797), nomen prius.
PHANERANDRA, Stschegl. in Bull. Soc. Nat. Mosc. (1859) I. 20.

LEUCOPOGON:—
abnormis, *Sond. in Lehm. Pl. Preiss.* i. 325.—Ins. Wagiou.
acicularis, *Benth. Fl. Austral.* iv. 199.—Austral.
acuminatus, Brongn. Voy. Coq. Bot. t. 53 = abnormis.
acuminatus, *R. Br. Prod.* 545.—Austral.
acutifolius, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 18 = unilateralis.
adpressus, Sieber, ex DC. Prod. vii. 747 = muticus.
affinis, R. Br. Prod. 541 = lanceolatus.
albicans, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
Allittii, *F. Muell. Fragm.* iv. 103.—Austral.
alternifolius, *R. Br. Prod.* 543.—Austral.
amplexicaulis, *R. Br. l. c.*—Austral.
angustatus, Benth. in Enum. Pl. Hueg. 77 = revolutus.
apiculatus, *R. Br. Prod.* 542.—Austral.
apiculatus, Sond. in Linnaea, xxvi. (1853–55) 248 = concurvus.
appressus, *R. Br. Prod.* 546.—Austral.
appressus, DC. Prod. vii. 754 = esquamatus.
appressus, Sieber, ex Benth. Fl. Austral. iv. 209 = muticus.
assimilis, Benth. in Enum. Pl. Hueg. 77 = polymorphus.
assimilis, *R. Br. Prod.* 545.—Austral.
astrolomioides, F. Muell. ex Sond. in Linnaea, xxvi. (1853) 249 = rufus.
atherolepis, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 13.—Austral.
attenuatus, *A. Cunn. in Field, N. S. Wales*, 341.—Austral.
australis, *R. Br. Prod.* 541.—Austral.
australis, Sieber, ex DC. Prod. vii. 744 = lanceolatus.
Bellignianus, Raoul, in Ann. Sc. Nat. Sér. II. ii. (1844) 118 = Fraseri.
biflorus, *R. Br. Prod.* 545.—Austral.
Billardieri, *Montr. in Mém. Acad. Lyon*, x. (1860) 226.—N. Caled.
blepharodes, DC. Prod. vii. 753 = Astroloma pallidum.
blepharolepis, *F. Muell. Fragm.* vi. 48.—Austral.
Bossiaea, *F. Muell. l. c.* 47.—Austral.
brachycephalus, DC. Prod. vii. 746 = cucullatus.
bracteolaris, *Benth. Fl. Austral.* iv. 197.—Austral.
brevibarbis, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 14.—N. Zel.
brevicuspis, *Benth. Fl. Austral.* iv. 210.—Austral.
breviflorus, *F. Muell. Fragm.* iv. 102.—Austral.
brevifolius, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 17 = obtusatus.
capitellatus, *DC. Prod.* vii. 747.—Austral.
carinatus, *R. Br. Prod.* 545.—Austral.
carinatus, DC. Prod. vii. 750 = assimilis.
ciliatus, A. Cunn. ex DC. Prod. vii. 746 = collinus.
coelophyllus, A. Cunn. l. c. 753 = Coleanthera coelophylla.
Colensoi, Hook. f. Fl. N. Zel. i. 165 = Cyathodes Colensoi.
collinus, *R. Br. Prod.* 543.—Austral.
compactus, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 13.—Austral.
concinnus, *Benth. Fl. Austral.* iv. 212.—Austral.
concurvus, *F. Muell. Fragm.* iii. 144.—Austral.
confertus, *Benth. Fl. Austral.* iv. 208.—Austral.
conostephioides, *DC. Prod.* vii. 753.—Austral.
cordatus, *Sond. in Lehm. Pl. Preiss.* i. 313.—Austral.
cordifolius, *Lindl. in Mitch. Three Exped.* ii. 122.—Austral.
corifolius, *Endl. Nov. Stirp. Dec.* 15.—Austral.
corynocarpus, *Sond. in Lehm. Pl. Preiss.* i. 322.—Austral.
crassiflorus, *F. Muell. Fragm.* vi. 40.—Austral.
crassifolius, *Sond. in Lehm. Pl. Preiss.* i. 316.—Austral.
cryptanthus, *Benth. Fl. Austral.* iv. 199.—Austral.
cucullatus, *R. Br. Prod.* 545.—Austral.
cucullatus, A. Cunn. ex DC. Prod. vii. 752 = tamariscinus.
cuneifolius, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 18.—Austral.
Cunninghami, R. Br. ex DC. Prod. vii. 745 = lanceolatus.
cuspidatus, *R. Br. Prod.* 545.—Austral.
cuspidatus, Lindl. in Mitch. Journ. Trop. Austral. 225 = Mitchellii.
cymbiformis, *A. Cunn. ex DC. Prod.* vii. 750.—Austral.
Cymbulae, *Labill. Sert. Austro-Caled.* 36. t. 39.—N. Caled.
dammarifolius, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.

LEUCOPOGON:—
dasystylis, Sond. in Lehm. Pl. Preiss. i. 325 = imbricatus.
decussatus, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 11 = elatior.
deformis, *R. Br. Prod.* 546.—Austral.
denudatus, Sieber, ex Spreng. Syst. iv. Cur. Post. 67 = microphyllus.
distans, *R. Br. Prod.* 544.—Austral.
Drummondi, DC. Prod. vii. 745 = australis.
durus, *Benth. Fl. Austral.* iv. 222.—Austral.
elatior, *Sond. in Lehm. Pl. Preiss.* i. 314.—Austral.
elegans, *Sond. l. c.* 318.—Austral.
Epacridis, DC. Prod. vii. 754 = Astroloma divaricatum.
ericoides, *R. Br. Prod.* 543.—Austral.
esquamatus, *R. Br. l. c.* 546.—Austral.
exarrhenus, F. Muell. Fragm. i. 178 = Styphelia pusilliflora.
exolacius, *F. Muell. l. c.* vi. 34.—Austral.
exsertus, F. Muell. l. c. iii. 143 = Styphelia Leucopogon.
fasciculatus, *A. Rich. Bot. Voy. Astrolab.* 215.—N. Zel.
fastigiatus, Sieber, ex Spreng. Syst. iv. Cur. Post. 67 = squamatus.
fimbriatus, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 17.—Austral.
flavescens, *Sond. in Lehm. Pl. Preiss.* i. 322.—Austral.
flexifolius, *R. Br. Prod.* 546.—Austral.
florulentus, *Benth. Fl. Austral.* iv. 194.—Austral.
Forsteri, A. Rich. Bot. Voy. Astrolab. 216 = Cyathodes acerosa.
Fraseri, A. Cunn. ex DC. Prod. vii. 753 = multiflorus.
Fraseri, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 47.—Austral.
fraternus, DC. Prod. vii. 749 = microphyllus.
gibbosus, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 12.—Austral.
Gilbertii, *Stschegl. l. c.* 15.—Austral.
glabellus, *R. Br. Prod.* 544.—Austral.
glacialis, *Lindl. in Mitch. Three Exped.* ii. 175.—Austral.
glaucescens, DC. Prod. vii. 745 = verticillatus.
gnaphalioides, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 14.—Austral.
Gnidium, Gunn, ex Hook. Comp. Bot. Mag. ii. (1836) 41 = Richei?
gracilis, *R. Br. Prod.* 544.—Austral.
gracillimus, *DC. Prod.* vii. 747.—Austral.
grandiusculus, F. Muell. Fragm. vi. 47 = atherolepis.
hirsutus, *Sond. in Lehm. Pl. Preiss.* i. 310.—Austral.
hirtellus, *F. Muell. ex Benth. Fl. Austral.* iv. 218.—Austral.
Hookeri, *Sond. in Linnaea*, xxvi. (1853) 248.—Austral.
Huegelii, *Planch. Hort. Donat.* 46.—Austral.
imbricatus, *R. Br. Prod.* 545.—Austral.
insularis, *A. Cunn. ex DC. Prod.* vii. 754.—Austral.
interruptus, *R. Br. Prod.* 541.—Austral.
javanicus, *De Vriese, in Miq. Pl. Jungh.* 84.—Java.
juniperinus, *R. Br. Prod.* 546.—Austral.
lanceolatus, *R. Br. l. c.* 541.—Austral.
lanceolatus, Sieber, ex DC. Prod. vii. 745 = Richei.
lancifolius, *Hook. f. in Hook. Ic. Pl.* t. 898.—Borneo.
lanigerus, A. Cunn. ex DC. Prod. vii. 749 = glabellus.
lasiophyllus, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 16.—Austral.
lasiostachys, *Stschegl. l. c.* 11.—Austral.
leptanthus, *Benth. Fl. Austral.* iv. 225.—Austral.
leptospermoides, *R. Br. Prod.* 546.—Austral.
linifolius, A. Cunn. ex DC. Prod. vii. 747 = melaleucoides.
lissanthoides, F. Muell. Fragm. iv. 101 = cuneifolius.
longistylis, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
Maccraei, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 106.—Austral.
malayanus, *Jack, in Malay. Misc.* i. (1820) n. v. 20.—Malaya.
margarodes, *R. Br. Prod.* 542.—Austral.
megacarpus, F. Muell. Fragm. iv. 102 = planifolius.
melaleucoides, *A. Cunn. ex DC. Prod.* vii. 750.—Austral.
microphyllus, *R. Br. Prod.* 544.—Austral.
Mitchellii, *Benth. Fl. Austral.* iv. 220.—Austral.
moluccanus, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 419.—Malaya.
mucronatus, DC. Prod. vii. 751 = attenuatus.
multiflorus, *R. Br. Prod.* 542.—Austral.

LEUCOPOGON:—
muticus, *R. Br. Prod.* 543.—Austral.
neo-anglicus, *F. Muell. ex Benth. Fl. Austral.* iv. 223. —Austral.
nervosus, R. Br. Prod. 544 = striatus.
nesophilus, DC. Prod. vii. 752 = Fraseri.
neurophyllus, F. Muell. Fragm. i. 37 = lanceolatus.
oblongifolius, Sond. in Lehm. Pl. Preiss. i. 323 = insularis.
obovatus, R. Br. Prod. 542 = revolutus.
obovatus, *Fawc. in Forbes, Nat. Wand. East. Archip.* 509.—Ins. Timor.
obovatus, Hort. Par. ex DC. Prod. vii. 747 = ericoides.
obtectus, *Benth. Fl. Austral.* iv. 223.—Austral.
obtusatus, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 269 = Hookeri.
obtusatus, *Sond. in Lehm. Pl. Preiss.* i. 313.—Austral.
Oldfieldii, *Benth. Fl. Austral.* iv. 203.—Austral.
opponens, *F. Muell. Fragm.* vi. 48.—Austral.
oppositifolius, *Sond. in Lehm. Pl. Preiss.* i. 316.— Austral.
ovalifolius, *Sond. l. c.* 324.—Austral.
ovatus, Sond. l. c. 319 = carinatus.
Oxycedrus, *Sond. l. c.* 321.—Austral.
ozothamnoides, *F. Muell. ex Benth. Fl. Austral.* iv. 205.—Austral.
Pancheri, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
paniculatus, Sond. in Lehm. Pl. Preiss. i. 306 = australis.
parviflorus, Lindl. Bot. Reg. t. 1560 = Richei.
parvifolius, DC. Prod. vii. 752 = tamariscinus.
parvifolius, Sond. in Lehm. Pl. Preiss. i. 319 = sprengelioides.
pauciflorus, R. Br. Prod. 546 = leptospermoides.
pendulus, Benth. in Enum. Pl. Hueg. 77 = secundiflorus.
pendulus, *R. Br. Prod.* 545.—Austral.
penicillatus, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 12 = distans.
petiolaris, DC. Prod. vii. 753 = Cyathodes ascendens.
phyllostachys, *Benth. Fl. Austral.* iv. 193.—Austral.
pilibundus, A. Cunn. ex DC. Prod. vii. 746 = microphyllus.
pimeleoides, A. Cunn. ex DC. l. c. 744 = lanceolatus.
planifolius, *Sond. in Lehm. Pl. Preiss.* i. 322.—Austral.
pleiospermus, *F. Muell. Fragm.* vi. 41.—Austral.
pleurandroides, *F. Muell. l. c.* iii. 143.—Austral.
plumuliflorus, *F. Muell. l. c.* vi. 29.—Austral.
pluriloculatus, *F. Muell. l. c.* i. 37.—Austral.
pogonocalyx, *F. Muell. ex Benth. Fl. Austral.* iv. 222. —Austral.
polymorphus, *Sond. in Lehm. Pl. Preiss.* i. 309.— Austral.
polystachyus, *R. Br. Prod.* 542.—Austral.
polystachyus, Lodd. Bot. Cab. t. 1436 = Richei.
propinquus, *R. Br. Prod.* 543.—Austral.
psilopus, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 19 = pendulus.
pulchellus, *Sond. in Lehm. Pl. Preiss.* i. 310.—Austral.
pungens, Sond. l. c. 324 = propinquus.
racemulosus, *DC. Prod.* vii. 747.—Austral.
ramulosus, A. Cunn. ex DC. l. c. 753 = attenuatus.
reclinatus, A. Cunn. l. c. 752 = attenuatus.
recurvatus, A. Cunn. l. c. 754 = attenuatus.
reflexus, *R. Br. Prod.* 544.—Austral.
revolutus, *R. Br. l. c.* 542.—Austral.
Richei, *R. Br. l. c.* 541.—Austral.
rigidus, A. Cunn. ex DC. Prod. vii. 753 = conostephioides.
rotundifolius, *R. Br. Prod.* 546.—Austral.
rotundifolius, Sond. in Lehm. Preiss. i. 323 = cordifolius.
rubicundus, *F. Muell. ex Benth. Fl. Austral.* iv. 207.— Austral.
rubicundus, F. Muell. Fragm. iv. 102 = L. Oxycedrus.
rubricaulis, R. Br. Prod. 542 = revolutus.
rudis, F. Muell. Fragm. iv. 106 = assimilis.
rufus, *Lindl. in Mitch. Three Exped.* ii. 179.— Austral.
rupestris, Sond. in Lehm. Pl. Preiss. i. 315 = striatus.
ruscifolius, *R. Br. Prod.* 545.—Austral.
salicifolius, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 68.—N. Caled.
scariosus, Panch. ex Brongn. & Gris, l. c. 6 = albicans.
secundiflorus, Sond. in Lehm. Pl. Preiss. i. 320 = pendulus.

LEUCOPOGON:—
semioppositus, F. Muell. Fragm. vi. 49 = elatior.
setiger, *R. Br. Prod.* 545.—Austral.
Shuttleworthii, Sond. in Lehm. Pl. Preiss. i. 307 = apiculatus.
Sieberi, DC. Prod. vii. 751 = juniperinus.
similis, Sond. in Lehm. Pl. Preiss. i. 321 = biflorus.
sparsus, A. Cunn. ex DC. Prod. vii. 751 = biflorus.
sprengelioides, *Sond. in Lehm. Pl. Preiss.* i. 319.— Austral.
squamatus, Steud. Nom. ed. II. ii. 38 = esquamatus.
squarrosus, *Benth. in Enum. Pl. Hueg.* 77.—Austral.
striatus, *R. Br. Prod.* 544.—Austral.
strictus, *Benth. Fl. Austral.* iv. 219.—Austral.
strongylophyllus, *F. Muell. Fragm.* iv. 101.—Austral.
Stuartii, F. Muell, ex Sond. in Linnaea, xxvi. (1853–55) 249 = Fraseri.
suaveolens, *Hook. f. in Hook. Ic. Pl.* sub t. 898.— Borneo.
subulatus, F. Muell. Fragm. iv. 103 = insularis.
tamariscinus, *R. Br. Prod.* 544.—Austral.
tectus, Sond. in Lehm. Pl. Preiss. i. 318 = carinatus.
tenuis, *DC. Prod.* vii. 744.—Austral.
tetragonus, *Sond. in Lehm. Pl. Preiss.* i. 317.— Austral.
thymifolius, *Lindl. ex Benth. Fl. Austral.* iv. 189.— Austral.
trichocarpus, R. Br. Prod. 543 = ericoides.
triqueter, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 15 = pulchellus.
unilateralis, *Stschegl. l. c.* 19.—Austral.
vaccinioides, Sond. in Lehm. Pl. Preiss. i. 325 = Pentachondra pumila.
vaginans, Sond. l. c. i. 315 = tamariscinus.
variifolius, Sond. l. c. 314 = glabellus.
verticillatus, *R. Br. Prod.* 541.—Austral.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
Vieillardi, Panch. ex Brongn. & Gris, l. c. 66 = L. Cymbulae.
villosus, R. Br. Prod. 542 = revolutus.
villosus, Lindl. ex Benth. Fl. Austral. iv. 190 = thymifolius.
virgatus, *R. Br. Prod.* 543.—Austral.
vitellinus, Sond. in Lehm. Pl. Preiss. i. 309 = assimilis.
Woodsii, *F. Muell. Fragm.* i. 178.—Austral.

LEUCOPSIDIUM, DC. Prod. vi. 43 (1837) = **Aphanostephus**, DC. (Compos.).
arkansanum, DC. l. c. = A. arkansanus.
humile, Benth. Pl. Hartw. 18 = A. humilis.
texanum, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 437 = A. arkansanus.

LEUCOPSIS, Baker, in Mart. Fl. Bras. vi. iii. 5 (1882). *COMPOSITAE.*
calendulacea, *Baker, l. c.* 9.—Bras.; Reg. Argent.
calvata, *Baker, l. c.* 8.—Bras.
diffusa, *Baker, l. c.*—Bras.; Reg. Argent.
gnaphalioides, *Baker, l. c.*—Bras.
Harveyi, *Baker, l. c.* 7, in obs.—Paraguay.
macrocephala, *Baker, l. c.*—Paraguay.
podocomoides, *Baker, l. c.* 6.—Bras.
scaposa, *Baker, l. c.*—Bras.
sericea, *Baker, l. c.* 7.—Am. austr.
Tweediei, *Baker, l. c.* 9.—Bras.

LEUCOPSORA, Rafin. Fl. Tellur. iv. 95 (1836) = **Cephalaria**, Schrad. (Dipsac.).
leucantha, Rafin. l. c. = C. leucantha.

LEUCORAPHIS, T. Anders. in Journ. Linn. Soc. vii. (1864) 21 = LEUCORHAPHIS, Nees = **Brillantaisia**, Beauv. (Acanthac.).

LEUCORCHIS, Blume, Mus. Bot. Lugd. Bat. i. 31 (1849) = **Didymoplexis**, Griff. (Orchid.).
minuta, Blume, Orch. Ind. Arch. 147 = D. pallens.
sylvatica, Blume, Mus. Bot. Lugd. Bat. i. 31 = D. pallens.

LEUCORCHIS, E. Mey. Preuss. Pflanzeng. 50 (1839) = **Habenaria**, Willd. (Orchid.).
albida, E. Mey. ex Schur, Sert. 72. n. 2703; ex ej. Enum. Pl. Transs. 645 = H. albida.
Friwaldskiana, Fuss, Fl. Transs. 625 (= *Gymnadenia Friwaldskiana*).—Transsylv.
lucida, Fuss, l. c. = H. albida?

LEUCORHAPHIS, Nees, in DC. Prod. xi. 97 (1847) = **Brillantaisia**, Beauv. (Acanthac.).
Lamium, Nees, l. c. = B. owariensis.
Vogeliana, Nees, l. c. = B. Vogeliana.

LEUCORRHAPHIS, Walp. Ann. iii. 213 (1852), sphalm. = praec.

LEUCOSCEPTRUM, Sm. Exot. Bot. ii. 113. t. 116 (1806). *LABIATAE* [Benth. & Hook. f. ii. 1221].
canum, *Sm. l. c.*—Reg. Himal.

LEUCOSEDUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 384 = **Sedum**, Tourn. (Crassul.).
album, Fourr. l. c. = S. album.
dasyphyllum, Fourr. l. c. = S. dasyphyllum.
hirsutum, Fourr. l. c. = S. hirsutum.

LEUCOSERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = **Senecio**, Tourn. (Compos.).
maritimus, Fourr. l. c. = S. Cineraria.

LEUCOSERIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 439 = **Malacothrix**, DC. (Compos.).
californica, Nutt. l. c. 440 = M. saxatilis.
saxatilis, Nutt. l. c. = M. saxatilis.
tenuifolia, Nutt. l. c. = M. saxatilis.

LEUCOSIA, Thou. Gen. Nov. Madag. 23 (1806) = **Chailletia**, DC.
Thouarsiana, Roem. & Schult. Syst. v. 324 = C. Leucosia.

LEUCOSIDEA, Eckl. & Zeyh. Enum. 265 (1836). *ROSACEAE*, Benth. & Hook. f. i. 622.
NESTLERA, E. Mey. ex Walp. Ann. iii. 856 (1852).
acaenoides, *Presl, Epim. Bot.* 200.—Afr. austr.
sericea, *Eckl. & Zeyh. Enum.* 265.—Afr. austr.

LEUCOSINAPIS, Spach, Hist. Vég. Phan. vi. 348 (1838) = **Brassica**, Tourn. (Crucif.).
alba, Spach, l. c. 350 = B. alba.
hispida, Spach, l. c. 352 = B. hispida.

LEUCOSMIA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 231. *THYMELAEACEAE*, Benth. & Hook. f. iii. 199.
acuminata, *A. Gray, in Journ. Bot.* iii. (1865) 306.— Ins. Fiji.
Burnettiana, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 231.—Ins. Fiji.
ovata, *Decne. Bot. Voy. Vénus*, 17.—Ins. Fiji.
pubiflora, *A. Gray, in Journ. Bot.* iii. (1865) 306.— Ins. Fiji.

LEUCOSPERMUM, R. Br. in Trans. Linn. Soc. x. (1810) 95. *PROTEACEAE*, Benth. & Hook. f. iii. 170.
DIASTELLA, Salisb. in Knight, Prot. 61 (1809).
LEUCADENDRUM, *Salisb. Parad. Lond.* sub t. 67 (1807).
attenuatum, *R. Br. in Trans. Linn. Soc.* x. (1810) 96. —Afr. austr.
buxifolium, *R. Br. l. c.* 100.—Afr. austr.
candicans, Loud. ex Meissn. in DC. Prod. xiv. 258 = tomentosum.
Conocarpodendron, Buek, Ind. ad DC. Prod. iv. 203 = seq.
conocarpum, *R. Br. in Trans. Linn. Soc.* x. (1810) 99. —Afr. austr.
crinitum, *R. Br. l. c.* 103.—Afr. austr.
crinitum, Klotzsch, ex Meissn. in DC. Prod. xiv. 261 = Mundii.
cryptanthum, Buek, ex Meissn. l. c. 260 = crinitum.
diffusum, *R. Br. in Trans. Linn. Soc.* x. (1810) 104. —Afr. austr.
diffusum, Sieber, ex Meissn. in DC. Prod. xiv. 257 = L. Hypophyllum.
Ecklonii, Buek, ex Meissn. l. c. 258 = tomentosum.
ellipticum, *R. Br. in Trans. Linn. Soc.* x. (1810) 98. —Afr. austr.
formosum, Knight, ex Loud. Encyc. Pl. 82 = Protea formosa.
grandiflorum, *R. Br. in Trans. Linn. Soc.* x. (1810) 100.—Afr. austr.
Gueinzii, *Meissn. in DC. Prod.* xiv. 254.—Afr. austr.

LEUCOSPERMUM:—
Hypophyllum, *R. Br. in Trans. Linn. Soc.* x. (1810) 102.—Afr. austr.
involucrata, *Roem. & Schult. Syst.* iii. 363.—Afr. austr.
lineare, *R. Br. in Trans. Linn. Soc.* x. (1810) 96.—Afr. austr.
marginatum, Spreng. Syst. i. 464 = Leucadendron marginatum.
medium, *R. Br. in Trans. Linn. Soc.* x. (1810) 97.—Afr. austr.
molle, *R. Br. l. c.* 103.—Afr. austr.
molle, Klotzsch, ex Meissn. in DC. Prod. xiv. 260 = crinitum.
Mundii, *Meissn. l. c.* 261.—Afr. austr.
nutans, *R. Br. in Trans. Linn. Soc.* x. (1810) 98.—Afr. austr.
oleaefolium, *R. Br. l. c.* 104.—Afr. austr.
ovatum, *Schult. Mant.* iii. 266.—Afr. austr.
parile, Knight, ex Loud. Encyc. Pl. 82 = Serruria phylicoides.
patulum, R. Br. in Trans. Linn. Soc. x. (1810) 100 = puberum.
pedunculatum, *Klotzsch, ex Krauss, in Flora*, xxviii. (1845) 76.—Afr. austr.
penicillatum, *Buek, in Drège, Zwei Pfl. Docum.* 85, nomen.—Afr. austr.
praemorsum, Buek, ex Meissn. in DC. Prod. xiv. 257 = attenuatum.
puberum, *R. Br. in Trans. Linn. Soc.* x. (1810) 100.—Afr. austr.
purpureum, Mund, ex Meissn. l. c. 261 = Mundii.
reflexum, *Buek, ex Meissn. l. c.* 254.—Afr. austr.
Rochetianum, *A. Rich. in Compt. Rend. Par.* (1851) i. 229.—Abyssin.
spathulatum, *R. Br. in Trans. Linn. Soc.* x. (1810) 101.—Afr. austr.
spathulatum, Drège, ex Meissn. in DC. Prod. xiv. 299 = Nivenia Sceptrum.
tomentosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 101.—Afr. austr.
tottum, *R. Br. l. c.* 97.—Afr. austr.
truncatum, Buek, ex Meissn. in DC. Prod. xiv. 256 = Zeyheri.
Zeyheri, *Meissn. l. c.* 255.—Afr. austr.

LEUCOSPORA, Nutt. in Journ. Acad. Philad. vii. (1834) 87 = **Conobea**, Aubl. (Scrophul.).
multifida, Nutt. l. c. = C. multifida.

LEUCOSTACHYS, Hoffmgg. Preisv. Orch. (1842) 26.—Cf. Linnaea, xvi. (1842) Litt. 234 = **Goodyera**, R. Br. (Orchid.).
procera, Hoffmgg. ll. cc. = G. procera.

LEUCOSTEMMA, Benth. in Wall. Cat. n. 642 (1828); et ex G. Don, Gen. Syst. i. 449 (1831) = **Stellaria**, Linn. (Caryoph.).
angustifolium, Royle, Illustr. Bot. Himal. t. 21 = S. Webbiana.
latifolium, Benth. in Wall. Cat. n. 643; et ex G. Don, Gen. Syst. i. 449 = S. latifolia.
Webbianum, Benth. ll. cc. = S. Webbiana.

LEUCOSTEMMA, D. Don, in Mem. Wern. Soc. v. (1823–24) 540 = **Helichrysum**, Vaill. (Compos.).
lingulatum, D. Don, l. c. 541 = H. vestitum.
vestitum, D. Don, l. c. = H. vestitum.

LEUCOSTOMON, G. Don, Gen. Syst. ii. 481 (1832) = **Lecostemon**, Moç. & Sesse (Rosac.).

LEUCOSYKE, Zoll. & Mor. in Mor. Syst. Verz. Zoll. 76 (1845–46). *URTICACEAE*, Benth. & Hook. f. iii. 391.
Missiessya, Gaudich. Bot. Voy. Bonite, t. 93 (1853?).
alba, *Zoll. & Mor. in Syst. Verz. Ind. Archip.* 100.—Java.
bimensis, Zoll. & Mor. l. c. 100, 103 = alba.
bornensis, *Miq. Fl. Ind. Bat.* i. ii. 267.—Borneo.
candidissima, Wedd. in DC. Prod. xvi. i. 235[26] = javanica.
capitellata, Wedd. l. c. 235[27] = alba.
celebica, *Miq. Fl. Ind. Bat.* i. ii. 267.—Ins. Celebes.
celtidifolia, Miq. l. c. 266 = alba.

LEUCOSPERMUM:—
corymbulosa, *Wedd. in DC. Prod.* xvi. i. 235[30].—Ins. Philipp.
elongata, Miq. Fl. Ind. Bat. i. ii. 266 = alba.
falcata, Miq. l. c. = alba.
hispidissima, *Miq. l. c.* 265.—Ins. Philipp.
javanica, *Zoll. & Mor. Syst. Verz. Zoll.* 76.—Java.
mutabilis, Miq. Fl. Ind. Bat. i. ii. 266 = alba.
ochroneura, *Wedd. in DC. Prod.* xvi. i. 235[29].—Ins. Molucc.
rubiginosa, *Miq. Fl. Ind. Bat.* i. ii. 267.—Borneo.
sumatrana, *Miq. l. c.*—Sumatra.
villosa, Miq. l. c. 266 = alba.

LEUCOTHAMNUS, Lindl. Swan River App. 19 (1839) = **Thomasia**, J. Gay (Stercul.).
montanus, Lindl. l. c. = T. macrocarpa.
polyspermus, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 11 = T. rugosa.

LEUCOTHEA, Moç. & Sesse, ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 419 = **Saurauja**, Willd. (Ternstr.).
serrata, Moç. & Sesse, ex DC. l. c. 420 = S. serrata.
villosa, Moç. & Sesse, ex DC. l. c. = S. villosa.

LEUCOTHOË, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159. *ERICACEAE*, Benth. & Hook. f. ii. 584.
Eubotrys, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 269.
acuminata, *G. Don, Gen. Syst.* iii. 832.—Am. bor.
affinis, *A. Rich. Fl. Cub. Fanerog.* ii. 74.—Cuba.
ambigua, *Meissn. in Mart. Fl. Bras.* vii. 156.—Bras.
axillaris, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 159.—Am. bor.
bahiensis, DC. Prod. vii. 604 = revoluta.
Bojeri, DC. l. c. 603 (Agauriae sp.).—Madag.
bracamorensis, DC. l. c. = Agarista bracamorensis.
breviflora, *Meissn. in Mart. Fl. Bras.* vii. 165.—Bras.
buxifolia, DC. Prod. vii. 603 = Agauria buxifolia.
Catesbaei, *A. Gray, Man. Bot. N. U. St.* ed. II. 252.—Am. bor.
chlorantha, DC. Prod. vii. 604 = Agarista chlorantha.
chlorantha, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 399.—Japon.
ciliata, *DC. Prod.* vii. 605.—Bras.
coccinea; *DC. l. c.*—Bras.
cordifolia, *Meissn. in Mart. Fl. Bras.* vii. 162.—Bras.
coriacea, DC. Prod. vii. 602 = Andromeda nitida.
coriifolia, *DC. l. c.* 605.—Bras.
crassifolia, *DC. l. c.*—Bras.
Davisiae, *Torr. ex A. Gray, in Proc. Am. Acad.* vii. (1868) 400.—Calif.
eriophylla, DC. Prod. vii. 606 = Gaultheria ferruginea.
eucalyptoides, DC. l. c. 605 = Agarista eucalyptoides.
floribunda, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159 = Pieris floribunda.
glandulosa, *A. Rich. Fl. Cub. Fanerog.* ii. 74.—Cuba.
Grayana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 46.—Japon.
Griffithiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 460.—Reg. Himal.
hirta, *DC. Prod.* vii. 606.—Hab.?
hispidula, *Meissn. in Mart. Fl. Bras.* vii. 164.—Bras.
ilicifolia, DC. Prod. vii. 605 = Agarista ilicifolia.
intermedia, *Meissn. in Mart. Fl. Bras.* vii. 163.—Bras.
katagherensis, DC. Prod. vii. 606 = Gaultheria fragrantissima.
Keiskei, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 32.—Japon.
latifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 74.—Cuba.
laxiflora, *Meissn. in Mart. Fl. Bras.* vii. 157.—Bras.
littoralis, DC. Prod. vii. 602 (Agauriae sp.).—Madag.
Mariana, DC. l. c. = Andromeda nitida.
Martii, *Meissn. in Mart. Fl. Bras.* vii. 155.—Bras.
multiflora, DC. Prod. vii. 605 = Agarista multiflora.
myrsinefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 74.—Cuba.
neriifolia, DC. Prod. vii. 605 = Agarista neriifolia.
nummularia, DC. l. c. 603 = Agarista nummularia.
oleifolia, DC. l. c. 605 = Agarista oleifolia.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1852) 132.—Bras.
Pistrix, DC. Prod. vii. 604 = Agarista Pistrix.

LEUCOTHOË:—
pulchella, DC. Prod. vii. 604 = Agarista pulchella.
pulchra, DC. l. c. = Agarista pulchra.
racemosa, *A. Gray, Man. Bot. N. U. St.* ed. II. 252.—Am. bor.
recurva, *A. Gray, l. c.*—Am. bor.
revoluta, *DC. Prod.* vii. 604.—Bras.
rhomboidalis, A. Gray, Syn. Fl. N. Am. ii. i. 33 = Lyonia ferruginea.
salicifolia, DC. Prod. vii. 602 = Agauria salicifolia.
secunda, DC. l. c. 606 = Andromeda Polifolia.
serrata, DC. l. c. = Agarista pulchra.
serrulata, DC. l. c. 604 = Agarista serrulata.
spectabilis, *Meissn. in Mart. Fl. Bras.* vii. 159.—Bras.
spicata, G. Don, Gen. Syst. iii. 832 = racemosa.
spinulosa, G. Don, l. c. = Catesbaei.
subcanescens, *Meissn. in Mart. Fl. Bras.* vii. 163.—Bras.
subrotunda, DC. Prod. vii. 603 = Agarista subrotunda.
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 46.—Japon.
Varnhageniana, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxxiii. (1884) 323.—Bras.

LEUCOXYLON, G. Don, Gen. Syst. ii. 21 (1832) = LEUCOXYLUM, Blume = **Diospyros**, Linn. (Ebenac.).

LEUCOXYLON, Rafin. Sylva Tellur. 77 (1838) = **Tecoma**, Juss. (Bignon.).
acuminata, Rafin. l. c. 78 = T. Leucoxylon.
riparia, Rafin. l. c. 77 = T. Leucoxylon.

LEUCOXYLUM, Blume, Bijdr. 1169 (1826) = **Diospyros**, Linn. (Ebenac.).
buxifolium, Blume, l. c. = D. microphylla.
laurinum, E. Mey. in Drège, Zwei Pfl. Docum. 199, nomen (Quid?).—Afr. austr.

LEUCRINIS, Rafin. Fl. Tellur. iv. 27 (1836) = **Leucocrinum**, Nutt. (Liliac.).

LEUKERIA, Endl. Ench. 249 (1841) = **Leucheria**, Lag. (Compos.).

LEUKOSYKE, Endl. Gen. Suppl. iv. 34 (1847) = **Leucosyke**, Zoll. & Mor. (Urtic.).

LEUNISIA, Phil. in Linnaea, xxxiii. (1864–65) 120. *COMPOSITAE*, Benth. & Hook. f. ii. 501.
laeta, *Phil. l. c.*—Chili.

LEURADIA, Poir. Encyc. Suppl. iii. 346 (1813) = **Lavradia**, Vell. (Violar.).

LEUWENHOEKIA, Bartl. Ord. 49 (1830) = **Levenhookia**, R. Br. (Stylid.).

LEUZEA, DC. Fl. Fr. iv. 109 (1805). *COMPOSITAE*, Benth. & Hook. f. ii. 477.
Rhacoma, Adans. Fam. ii. 117 (1763).
altaica, Fisch. ex Schau. Cat. Sem. Hort. Vrat. (1834) = Centaurea carthamoides.
altaica, Link, Enum. Hort. Berol. ii. 356 = Centaurea salina.
australis, Gaudich. in Freyc. Voy. Bot. 462. t. 92 = Centaurea australis.
carthamoides, DC. Diss. Comp. 49. n. 2 = Centaurea carthamoides.
conifera, *DC. Fl. Fr.* iv. 109.—Reg. Mediterr.
conifera, Spreng. Syst. iii. 382 = Serratula canariensis.
dahurica, Bunge, Enum. Pl. Chin. Bor. 37 = Centaurea monanthos.
exaltata, Cutanda, ex Willk. in Linnaea, xxx. (1859–60) 117 = Centaurea rhaponticoides.
longifolia, *Hoffmgg. & Link, Fl. Port.* ii. 217.—Hispan.
Palmellana, Welw. ex Benth. & Hook. f. Gen. ii. 477 = longifolia.
pinnatifida, *DC. Prod.* vi. 666.—Hab.?
pumila, Steud. Nom. ed. II. ii. 39, sphalm. = seq.
pusilla, Spreng. Syst. iii. 382 = Centaurea pygmaea.
rhaponticoides, Graells (Ramill. 10.) in Mem. Acad. Cienc. Madr. ii. (1859) 468 = Centaurea rhaponticoides.
salina, Spreng. Syst. iii. 382 = Centaurea salina.
sclerophylla, Boiss. & Heldr. Diagn. Ser. I. x. 97 = Centaurea serratuloides.
serratuloides, Fisch. & Mey. ex DC. Prod. vi. 666 = Centaurea serratuloides.

LEVENHOEKIA, Steud. Nom. ed. I. 477 (1821) = seq.

LEVENHOOKIA, R. Br. Prod. 572 (1810). *STYLIDIEAE*, Benth. & Hook. f. ii. 535.
COLEOSTYLIS, Sond. in Lehm. Pl. Preiss. i. 391 (1845).
GYNOCAMPUS, Lesch. ex DC. Prod. vii. 338 (1839).
STIBAS, Comm. ex DC. l. c. (1839).
creberrima, F. Muell. Fragm. iii. 121 = dubia.
dubia, *Sond. in Lehm. Pl. Preiss.* i. 392.—Austral.
leptantha, *Benth. Fl. Austral.* iv. 35.—Austral.
pauciflora, *Benth. in Enum. Pl. Hueg.* 74.—Austral.
Preissii, *F. Muell. Fragm.* iv. 94.—Austral.
pusilla, *R. Br. Prod.* 573.—Austral.
Sonderi, *F. Muell. Fragm.* i. 18.—Austral.
stipitata, *F. Muell. l. c.* iv. 94.—Austral.
stylidioides, F. Muell. l. c. 77 = pauciflora.

LEVIERIA, Becc. Malesia, i. (1877) 192. *MONIMIACEAE*, Benth. & Hook. f. iii. 142.
montana, *Becc. l. c.* 193.—Ins. Amboina.

LEVINA, Adans. Fam. ii. 190 (1763) = **Prasium**, Linn. (Labiat.).

LEVISANUS, Schreb. Gen. i. 149 (1789) = **Staavia**, Dahl (Bruniac.).

LEVISIA, Steud. Nom. ed. II. ii. 39 (1841) = **Lewisia**, Pursh (Portulac.).

LEVISTICUM, Riv. ex Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 916.
argutum, Lindl. in Royle, Illustr. Bot. Himal. 232 = Ligusticum elatum.
grandiflorum, Sond. in Harv. & Sond. Fl. Cap. ii. 553 (Annesorhizae sp.).—Afr. austr.
officinale, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 101. f. 41.—Europ.
paludapifolium, Aschers. Fl. Brandenb. 250 = officinale.
vulgare, Reichb. Fl. Germ. Excurs. 463 = officinale.

LEVRETONIA, Reichb. in Moessl. Handb. ed. II. i. p. LXIII (1827) = LEBRETONIA, Schrank = **Pavonia**, Linn. (Malv.).

LEWISIA, Pursh, Fl. Am. Sept. ii. 368 (1814). *PORTULACEAE*, Benth. & Hook. f. i. 159.
alba, Kellogg, in Proc. Calif. Acad. ii. (1863) 115 = rediviva.
brachycalyx, *Engelm. ex A. Gray, in Proc. Am. Acad.* vii. (1868) 400.—Calif.
brachycarpa, [S. Wats.] in Bot. Calif. i. 79, sphalm. = praec.
rediviva, *Pursh, Fl. Am. Sept.* ii. 368.—Am. bor. occ.

LEXARZA, La Llave, in La Llave, & Lex. Nov. Veg. Desc. fasc. ii. 7 (1825) = **Myrodia**, Sw. (Stercul.).
funebris, La Llave, l. c. = M. funebris.

LEXIPYRETUM, Dulac, Fl. Hautes-Pyr. 449 (1867) = **Gentiana**, Linn.
acaule, campestre, ciliatum, luteum, nivale, Pneumonanthe, punctatum, pyrenaicum, tenellum, vernum, Dulac, l. c. 450, 451 = Gentianae sp. homonymae.

LEYCESTERA, Reichb. Consp. 96 (1828) = seq.

LEYCESTERIA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 181 (1824). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 5.
formosa, *Wall. l. c.* 182.—Reg. Himal.
glaucophylla, *Hook. f. Fl. Brit. Ind.* iii. 16.—Reg. Himal.
gracilis, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 77, in obs. = glaucophylla.

LEYCESTRIA, Endl. Gen. 568 (1838) = praec.

LEYMUS, Hochst. in Flora, xxxi. (1848) 118, in adnot. = **Elymus**, Linn. (Gramin.).
arenarius, Hochst. l. c. = E. arenarius.

LEYSERIA, Neck. Elem. i. 11 (1790) = seq.

LEYSSERA, Linn. Sp. Pl. ed. II. 1249 (1763) (Leysera). *COMPOSITAE*, Benth. & Hook. f. ii. 327.
ASTEROPTERUS, Gaertn. Fruct. ii. 460. t. 173 (1791).
CALLICORNIA, Burm. f. Prod. Fl. Cap. 24 (1768).
LEPTOPHYTUS, Cass. in Bull. Soc. Philom. (1817) 11.
LONGCHAMPIA, Willd. in Ges. Nat. Fr. Berl. Mag. v. (1811) 159.
arctioides, Thunb. Prod. Pl. Cap. 160 = Heterolepis decipiens.
Callicornia, *Linn. Mant.* ii. 286.—Afr. austr.
Callicornia, Thunb. Prod. Pl. Cap. 160 = gnaphalioides.
Callicornia, Willd. ex Less. in DC. Prod. vii. 13 = Printzia aromatica.
capillifolia, *Spreng. Nov. Prov.* 25.—Reg. Mediterr.
caroliniana, *Walt. Fl. Carol.* 211 (Quid ?).—Am. bor.
ciliata, Thunb. Prod. Pl. Cap. 160 = Mairia taxifolia.
discoidea, Spreng. Pugill. ii. 78 = capillifolia.
ericoides, Berg. Desc. Pl. Cap. 294 = Relhania ericoides.
gnaphalodes, *Linn. Pl. Rar. Afr.* 23; *Amoen. Acad.* vi. 104.—Afr. austr.
gnaphaloides, Thunb. Prod. Pl. Cap. 160 = Helipterum gnaphalioides.
incana, *Thunb. l. c.*—Afr. austr.
odorata, Ruiz & Pav. ex DC. Prod. vii. 34 = Onoseris odorata.
ovata, Thunb. Prod. Pl. Cap. 160 = Amellus annuus.
paleacea, Linn. Syst. ed. XII. 560 = Relhania ericoides.
picta, Thunb. Prod. Pl. Cap. 160 = Dicoma radiata.
pilosella, Thunb. l. c. = Aster hispidus, Mairia perezioides.
polifolia, Thunb. l. c. 161 = Printzia Bergii.
squarrosa, Thunb. l. c. 160 = Helipterum gnaphaloides.
tenella, *DC. Prod.* vi. 279.—Afr. austr.
tenuifolia, Salisb. Prod. 202 = gnaphalioides.
tridactyla, E. Mey. ex DC. Prod. v. 215 = Amellus tridactylus.

LHODRA, Endl. Gen. Suppl. ii. 68 (1842) = LODHRA, G. Don = **Symplocos**, Linn. (Styrac.).

LHOTSKYA, Schau. in Linnaea, x. (1835–36) 309. *MYRTACEAE*, Benth. & Hook. f. i. 699.
acutifolia, *Lindl. Swan River App.* 7.—Austral.
brevifolia, *Schau. in Lehm. Pl. Preiss.* i. 103.—Austral.
ciliata, *F. Muell. ex Benth. Fl. Austral.* iii. 54.—Austral.
cuspidata, F. Muell. in Hook. Kew Journ. viii. (1856) 324 = Calythrix achaeta.
ericoides, *Schau. in Linnaea*, x. (1835–36) 309.—Austral.
genethylloides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 16.—Austral.
glaberrima, *F. Muell. Fragm.* i. 13.—Austral.
Harvestiana, *F. Muell. l. c.* xi. 8.—Austral.
hirta, Regel, Gartenfl. (1863) 337. t. 415 = ericoides.
purpurea, *F. Muell. Fragm.* i. 224.—Austral.
scabra, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 324 = ericoides.
violacea, *Lindl. Swan River App.* 7.—Austral.

LHOTZKYA, Endl. in Enum. Pl. Hueg. 46 (1837) = praec.

LIABUM, Adans. Fam. ii. 131 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 435.
ALIBUM, Less. Syn. Comp. 152 (1832).
ANDROMACHIA, Humb. & Bonpl. Pl. Aequin. ii. 104. t. 112 (1809).
CHRYSACTINUM, Wedd. Chlor. And. i. 212. t. 39 (1855).
ERATO, DC. Prod. v. 317 (1836).
KASTNERA, Sch. Bip. in Flora, xxxvi. (1853) 37.
MUNNOZIA, Ruiz & Pav. Prod. Fl. Per. 108. t. 23 (1794).
OLIGACTIS, Cass. in Dict. Sc. Nat. xxxvi. 16 (1825).
PARANEPHELIUS, Poepp. & Endl. Nov. Gen. et Sp. iii. 42. t. 248 (1842).
PRIONOLEPIS, Poepp. & Endl. l. c. 55. t. 261 (1842).
SINCLAIRIA, Hook. & Arn. Bot. Beech. Voy. 433 (1841).
STARKEA, Willd. Sp. Pl. iii. 2216 (1803).
VIVIANIA, Willd. ex DC. Prod. v. 95 (1836).

LIABUM:—
acaule, *DC. Prod.* v. 97.—Ecuador.
amplexicaule, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 43.—Peruv.
Andrieuxii, *Benth. & Hook. f. Gen.* ii. 437.—Mexic.
andromachioides, *Benth. & Hook. f. l. c.* 436.—Mexic.
asclepiadeum, *Sch. Bip. in Linnaea*, xx. (1847) 521.—N. Granat.
auriculatum, *Griseb. in Goett. Abh.* xxiv. (1879) 202.—Reg. Argent.
Bonplandii, *Cass. in Dict. Sc. Nat.* xxvi. 206.—Ecuador.
Brownei, *Cass. l. c.* 203.—Ind. occ.
candidum, *Griseb. in Goett. Abh.* xxiv. (1879) 203.—Reg. Argent.
corymbosum, *Sch. Bip. in Flora*, xxxvi. (1853) 34.—Am. austr.
Deppeanum, *Hemsl. Biol. Centr. Am. Bot.* ii. 232.—Mexic.
discolor, *Benth. & Hook. f. ex Hemsl. l. c.*—Mexic.
ericoides, *Less. in Linnaea*, vi. (1831) 704.—Peruv.
erigeroides, *Benth. Pl. Hartw.* 206.—Mexic.
eriocaulon, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 43. t. 249.—Peruv.
floribundum, Less. in Linnaea, vi. (1831) 702 = Bonplandii.
floribundum, *Sch. Bip. l. c.* xx. (1847) 582.—Am. trop.
glabrum, *Hemsl. Biol. Centr. Am. Bot.* ii. 232.—Mexic.
grandiflorum, *Less, in Linnaea*, vi. (1831) 698.—Ecuador.
hastifolium, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 43.—Peruv.
hieracioides, *DC. Prod.* v. 97.—Peruv.
igniarium, Less. in Linnaea, vi. (1831) 701 = Bonplandii.
Jussieui, *Cass. in Dict. Sc. Nat.* xxvi. 205 (Jussiei).—Am. austr.
lanceolatum, *Sch. Bip. in Flora*, xxxvi. (1853) 34.—Peruv.
Lechleri, Sch. Bip. in Bonplandia, iii. (1855) 236 = Diplostephium Lechleri.
lyratum, *A. Gray, in Proc. Am. Acad.* v. (1861) 115.—Peruv.
megacephalum, *Sch. Bip. in Flora*, xxxvi. (1853) 38.—Venezuela.
melastomoides, *Less. in Linnaea*, vi. (1831) 699.—N. Granat.
nubigenum, *Less. l. c.* 704.—Ecuador.
ovatum, *Ball, in Journ. Linn. Soc.* xxii. (1885) 46.—Peruv.
sagittatum, *Sch. Bip. in Flora*, xxxvi. (1853) 37.—Peruv.
sericolepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 232.—Mexic.
sessiliflorum, *Less. in Linnaea*, vi. (1831) 703.—Peruv.
solidagineum, *Less. l. c.* 700.—Peruv.
trinerve, *Sch. Bip. in Flora*, xxxvi. (1853) 34.—Peruv.
umbellatum, *Sch. Bip. in Journ. Bot.* i. (1863) 236.—Ind. occ.
uniflorum, *Ball, in Journ. Linn. Soc.* xxii. (1885) 46.—Peruv.
uniflorum, *Sch. Bip. in Flora*, xxxvi. (1853) 34.—Am. austr.
venosissimum, *Sch. Bip. in Flora*, xxxvi. (1853) 34.—Peruv.
verbascifolium, *Less. in Linnaea*, vi. (1831) 700.—Ecuador.
volubile, *Less. l. c.* 704.—Ecuador.
Wrightii, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 515.—Ind. occ.

LIATHRIS, C. Muell. in Walp. Ann. v. 162 (1858) = seq.

LIATRIS, Schreb. Gen. 542 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 248.
CALOSTELMA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 184 (1833).
LACIMARIA, Hill, Veg. Syst. iv. 49 (1763).
PSILOSANTHUS, Neck. Elem. i. 69 (1790).
acidota, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1847) 218.—Texas.
amplexicaulis, Rafin. New Fl. Am. iv. 76 (Quid ?).—Am. bor.
aspera, Michx. Fl. Bor. Am. ii. 92 = scariosa.
baicalensis, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 115 = Saussurea pycnocephala.

LIATRIS:—
bellidifolia, Michx. Fl. Bor. Am. ii. 92 = Carphephorus bellidifolius.
borealis, *Nutt. ex Macnab, in Edinb. N. Phil. Journ.* xix. (1835) 60; *Paxt. Mag. Bot.* v. (1838) 27.—Am. bor.
Botrys, Rafin. New Fl. Am. iv. 76 (Quid?).—Am. bor.
Boykini, *Torr. & Gray, Fl. N. Am.* ii. 70.—Am. bor.
brachystachya, Nutt. in Journ. Acad. Philad. vii. (1834) 72 = pycnostachya.
brasiliensis, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 461.—Bras.
Chapmanii, *Torr. & Gray, Fl. N. Am.* ii. 502 (*Suppl.*).—Am. bor.
cordata, Royle, ex C. B. Clarke, Comp. Ind. 247 = Ainsliaea aptera.
corymbosa, Nutt. Gen. Am. ii. 132 = Carphephorus corymbosus.
cylindracea, *Michx. Fl. Bor. Am.* ii. 93.—Am. bor.
cylindrica, Torr. in Ann. Lyc. N. York, ii. (1828) 210 = punctata.
dubia, Barton, Veg. Mat. Med. ii. 223. t. 49 = graminifolia.
elegans, *Willd. Sp. Pl.* iii. 1635.—Am. bor.
flexuosa, D. Thomas, in Am. Journ. Sc. xxxvii. (1839) 338 = cylindracea.
fruticosa, Nutt. l. c. v. (1822) 299 = Garberia fruticosa.
Garberi, *A. Gray, in Proc. Am. Acad.* xv. (1880) 48.—Am. bor.
gracilis, *Pursh, Fl. Am. Sept.* ii. 508.—Am. bor.
graminifolia, *Pursh, l. c.*—Am. bor.
graminifolia, Willd. Sp. Pl. iii. 1636 = cylindracea.
heterophylla, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 503 = scariosa.
hirsutiflora, Kostel. Allg. Med.-Pharm. Fl. ii. 651 = pycnostachya.
hirsutissima, Poir. ex Steud. Nom. ed. II. ii. 40 = pycnostachya.
intermedia, Lindl. Bot. Reg. t. 948 = squarrosa.
laevigata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 285 = tenuifolia.
lanata, Spreng. ex DC. Prod. vi. 540 = Saussurea albescens.
lanceolata, Bertol. Misc. Bot. v. 11. t. 3 = gracilis.
latifolia, D. Don, Prod. Fl. Nep. 169 = Ainsliaea pteropoda.
Linaria, Rafin. Fl. Ludov. 61 = pycnostachya.
lobelioides, Wall. ex C. B. Clarke, Comp. Ind. 246 = Ainsliaea pteropoda.
macrostachya, Michx. Fl. Bor. Am. ii. 91 = spicata.
marginata, *Cass. in Dict. Sc. Nat.* xxvi. 236.—Am. bor.
monocephala, Cass. l. c. 237 = cylindracea, graminifolia.
mucronata, *DC. Prod.* v. 129.—Mexic.
mucronata, Torr. & Gray, Fl. N. Am. ii. 70 = acidota.
odoratissima, Willd. Sp. Pl. iii. 1637 = Trilisia odoratissima.
oppositifolia, Nutt. in Am. Journ. Sc. v. (1822) 299 = Eupatorium ivaefolium.
paniculata, Michx. Fl. Bor. Am. ii. 93; Willd. Sp. Pl. iii. 1637 = Trilisia paniculata.
pauciflora, *Pursh, Fl. Am. Sept.* ii. 510.—Am. bor.
pauciflosculosa, Nutt. in Journ. Acad. Philad. vii. (1834) 71 = gracilis.
picta, Barton, ex DC. Prod. v. 130 = spicata.
pilosa, [Ker-Gawl. in] Bot. Reg. t. 595 = spicata.
pilosa, Willd. Sp. Pl. iii. 1636 = graminifolia.
propinqua, Hook. Bot. Mag. t. 3829 = graminifolia.
pumila, Lodd. Bot. Cab. t. 147 = spicata.
punctata, *Hook. Fl. Bor. Am.* i. 306. t. 105.—Am. bor. occ.
pycnostachya, *Michx. Fl. Bor. Am.* ii. 91.—Am. bor.
radians, Bertol. Misc. Bot. v. 9. t. 1 = elegans.
resinosa, DC. Prod. v. 129 = punctata.
resinosa, Nutt. Gen. Am. ii. 131 = spicata.
scariosa, *Willd. Sp. Pl.* iii. 1635.—Am. bor.
secunda, Ell. Sketch, ii. 278 = pauciflora.
sessiliflora, Bertol. Misc. Bot. v. 10. t. 2. = spicata.
sphaeroides, Michx. Fl. Bor. Am. ii. 92 = scariosa.
spicata, *Willd. Sp. Pl.* iii. 1636.—Am. bor.
squamosa, Nutt. in Journ. Acad. Philad. vii. (1834) 73 = Carphephorus Pseudo-liatris.
squarrosa, *Willd. Sp. Pl.* iii. 1634.—Am. bor.
squarrulosa, Michx. Fl. Bor. Am. ii. 92 = scariosa.
stricta, Macnab, in Edinb. N. Phil. Journ. xix. (1835) 60 = cylindracea.
tenuifolia, *Nutt. Gen. Am.* ii. 131.—Am. bor.
tomentosa, Michx. Fl. Bor. Am. ii. 93 = Carphephorus tomentosus.

LIATRIS:—
trichotoma, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 435 = Leptoclinium trichotomum.
turbinata, Sweet, Hort. Brit. ed. I. 220, nomen = spicata.
umbellata, Bertol. Misc. Bot. v. 13. t. 4 = Vernonia angustifolia.
uniflora, Rafin. New Fl. Am. iv. 76 (Quid?).—Am. bor.
varia, Banks, ex Pursh, Fl. Am. Sept. ii. 508 = scariosa.
virgata, Nutt. in Journ. Acad. Philad. vii. (1834) 72 = graminifolia.
Walteri, Ell. Sketch, ii. 285 = Carphephorus tomentosus.

LIBANOTHAMNUS, Ernst, in Vargasia (1870) 186 = **Espeletia**, Humb. & Bonpl. (Compos.).
neriifolius, Ernst, l. c. = E. neriifolia.

LIBANOTIS, Rafin. Sylva Tellur. 132 (1838) = **Cistus**, Tourn.
populifolius, Rafin. l. c. = C. populifolius.
umbellata, Rafin. l. c. = Helianthemum Libanotis.

LIBANOTIS, Riv. ex Hall. Enum. Helv. ii. 450 (1742) = **Seseli**, Linn. (Umbell.).
alpina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 29 = S. Libanotis.
annua, Crantz, Class. Umb. Emend. 107 = Athamanta cretensis.
annua, [Weber, in] Wigg. Prim. Fl. Holsat. 25 = Selinum sylvestre.
arctica, Rupr. Beitr. Pfl. Russl. ii. 36. t. 2 = *condensata*?
athamanthoides, DC. Prod. iv. 150 = S. athamantoides.
athamantica, Gaertn. Mey. & Scherb. Fl. Wett. i. 427 = S. Libanotis.
athamantoides, Schur, Enum. Pl. Transs. 258 = S. Libanotis.
aurea, Vis. Fl. Dalm. iii. 44 = Athamanta aurea.
bayonnensis, Griseb. Veget. 232, ex Nym. Consp. 295 = Candollei.
buchtormensis, DC. Coll. Mém. v. t. 3. f. 5 = S. buchtormense.
cachroides, DC. Prod. iv. 151 = Phloiodicarpus dahuricus.
Candollei, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 61.—Hispan.
chachroides, Steud. Nom. ed. II. ii. 41, sphalm. = *cachroides*.
condensata, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 724.—Sibiria.
cretensis, Gaertn. Fruct. i. 84. t. 21. f. 1 = S. Libanotis.
cretensis, Scop. Fl. Carn. ed. II. i. 192 = Athamanta cretensis.
crithmoides, Nym. Consp. 395 = S. Libanotis.
daucifolia, Reichb. Fl. Germ. Excurs. 468 = S. Libanotis.
daucoides, Scop. Fl. Carn. ed. II. i. 193 = S. Libanotis.
Daucus-creticus, Crantz, Stirp. Austr. ed. I. fasc. iii. 112; ed. II. 219 = praec.
elata, C. Koch, ex Boiss. Fl. Orient. ii. 967 = S. sibiricum.
eriocarpa, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1843) 195.—Sibir.
fastigiata, Rupr. in Beitr. Pfl. Russ. Reich. ii. 36, in textu = *condensata*?
filifolia, Roehl. ex Steud. Nom. ed. II. ii. 41 = Athamanta Matthioli.
gracilis, Reichb. Fl. Germ. Excurs. 468 = S. Libanotis.
graveolens, G. Don, Gen. Syst. iii. 312 = S. graveolens.
gummifera, Crantz, Class. Umb. Emend. 108 = S. gummiferum.
hirsuta, Lam. Fl. Fr. iii. 428 = Athamanta cretensis.
Hosteana, Schur, Enum. Pl. Transs. 258 = S. Libanotis.
humilis, Schur, in Oestr. Bot. Wochenbl. (1858) 21 = S. Libanotis.
intermedia, Rupr. Beitr. Pfl. Russ. Reich. iv. 53 = S. sibiricum.
italica, Crantz, Stirp. Austr. ed. I. fasc. iii. 117; ed. II. 222 = Athamanta cretensis.
Lehmanniana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 300 = S. Lehmannianum.
macedonica, Crantz, Cl. Umb. Emend. 107 = Athamanta macedonica.
major, Hall. ex Gouan, Illustr. t. 26; fide Pritz. Ic. Ind. 635 = S. Libanotis?
Matthioli, Bertol. Fl. Ital. iii. 320 = Athamanta Matthioli.
montana, Crantz, Stirp. Austr. ed. I. fasc. iii. 117; ed. II. 222; Lam. Fl. Fr. iii. 427 = S. Libanotis.
montana-ferulacea, Lange, Ind. Sem. Hort. Haun. (1855) 25.—Mont. Pyren.

LIBANOTIS:—
nitens, Vis. Fl. Dalm. iii. 43, in syn. = Seseli athamantoides.
nitida, Vis. l. c. t. 28 = praec.
Patriniana, DC. Prod. iv. 150 = S. graveolens.
pubescens, Nym. Consp. 295 = S. Libanotis.
pumila, Crantz, Stirp. Austr. ed. I. fasc. iii. 117 = Athamanta cretensis.
pumila, Schur, Enum. Pl. Transs. 258 = S. Libanotis.
pyrenaica, Bourg. ex Nym. Consp. 295 = S. Libanotis.
rigida, Crantz, Cl. Umb. Emend. 108.—Sicilia.
Riviniana, Scop. Fl. Carn. ed. II. i. 193 = S. Libanotis.
rupestris, Scop. l. c. 192 = Athamanta Matthioli.
seseloides, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 725 = Ligusticum seseloides.
sibirica, Koch, Syn. Fl. Germ. ed. I. 296 = S. Libanotis.
sicula, Bertol. Fl. Ital. iii. 317 = Athamanta sicula.
Stephaniana, DC. Prod. iv. 151.—Sibir.
tenuifolia, DC. l. c. = S. tenuifolium.
Turbith, Crantz, Cl. Umb. Emend. 108 = Athamanta Matthioli.
verticillata, DC. Prod. iv. 151 = Carum graecum.
villosa, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 31.—Sibiria.
vulgaris, DC. Prod. iv. 150 = S. Libanotis.

LIBANOTUS, Stackh. De Liban. 13 (1814) = **Boswellia**, Roxb. (Burser.).
asiaticus, Stackh. l. c. = B. serrata.

LIBANUS, Colebr. in As. Res. ix. (1807) 382. t. 5. f. 1 = **Boswellia**, Roxb. (Burserac.).
thuriferus, Colebr. l. c. = B. serrata.

LIBERTIA, Dum. Comm. Bot. 9 (1822) = **Funkia**, Spreng. (Liliac.).

LIBERTIA, Lej. in Nov. Act. Nat. Cur. xii. (1825) 755. t. 65. = **Bromus**, Linn. (Gramin.).
arduennensis, Lej. l. c. = B. arduennensis.

LIBERTIA, Spreng. Syst. i. 127, 168 (1825). *IRIDEAE*, Benth. & Hook. f. iii. 696.
NEMATOSTIGMA, A. Dietr. Sp. Pl. ii. 509 (1833).
RENEALMIA, R. Br. Prod. Add. (1810).
TAUMASTOS, Rafin. Fl. Tellur. iv. 9 (1836).
caerulescens, *Kunth, in Linnaea*, xix. (1847) 382.—Chili.
chilensis, *Klotzsch, ex Baker, in Journ. Linn. Soc.* xvi. (1877) 153.—Chili.
crassa, R. Grah. in Edinb. N. Phil. Journ. (Oct. 1833) 383 = formosa.
elegans, *Poepp. Fragm. Syn. Pl. Chil.* 1.—Chili.
formosa, *R. Grah. in Edinb. N. Phil. Journ.* (Oct. 1833) 383.—Chili.
graminea, Endl. in Lehm. Pl.-Preiss. ii. 32 = Orthrosanthus gramineus.
grandiflora, Phil. in Bot. Zeit. xiv. (1856) 648 = formosa.
grandiflora, *Sweet, Hort. Brit.* ed. II. 498.—N. Zel.
ixioides, C. Gay, Fl. Chil. vi. 31 = elegans.
ixioides, Klatt, in Mart. Fl. Bras. iii. 530. t. 68. f. 2 = formosa.
ixioides, *Spreng. Syst.* i. 168.—N. Zel.
Laurencii, Hook. f. Fl. Tasman. ii. 34. t. 129 = pulchella.
laxa, Endl. in Lehm. Pl. Preiss. ii. 32 = Orthrosanthus laxus.
macrocarpa, Klatt, in Linnaea, xxxi. (1861–62) 384 = grandiflora.
micrantha, A. Cunn. in Hook. Comp. Bot. Mag. ii. (1836) 375 = pulchella.
orbicularis, Colenso, in Trans. N. Z. Inst. xv. (1883) 329 = ixioides.
paniculata, *Spreng. Syst.* i. 168.—Austral.
pulchella, *Spreng. l. c.* 169.—Austral.?; N. Zel.
restioides, Klatt, in Linnaea, xxxi. (1861–62) 383 = ixioides.
stricta, Endl. in Lehm. Pl. Preiss. ii. 32 = Orthrosanthus multiflorus.
tricocca, *Phil. in Linnaea*, xxix. (1857–58) 63.—Chili.
tricolor, *Lem. Illustr. Hortic.* x. (1863) *Misc.* 35.—N. Zel.

LIBIDIBIA, Schlecht. in Linnaea, v. (1830) 192 = **Caesalpinia**, Linn. (Legum.).
coriaria, Schlecht. l. c. 193 = G. coriaria.
pauciflora, Griseb. Cat. Pl. Cub. 78 = C. pauciflora.
pinnata, Griseb. l. c. 79.—Cuba.

LIBOCEDRUS, Endl. Syn. Conif. 42 (1847). *CONIFERAE*, Benth. & Hook. f. iii. 426.
CALOCEDRUS, Kurz, in Journ. Bot. xi. (1873) 196. t. 133.
HEYDERIA, C. Koch, Dendrol. ii. II. 179 (1873).
austro-caledonicus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xviii. (1871) 140.—N. Caled.
Bidwillii, *Hook. f. Handb. N. Zeal. Fl.* 257.—N. Zel.
chilensis. *Endl. Syn. Conif.* 44.—Chili.
Craigiana, Low, ex Gord. Pinet. Suppl. 102 = Thuya gigantea.
decurrens, *Torr. Pl. Frémont.* 7. t. 3.—Calif.
dolabrata, [Nelson] Senilis, Pinaceae, 65 = Thuya dolabrata.
Doniana, *Endl. Syn. Conif.* 43.—N. Zel.
excelsa, Hort. ex Gord. Pinet. 132 = chilensis.
gigantea, Low, ex Gord. Pinet. Suppl. 102 = Thuya gigantea.
macrolepis, *Benth. & Hook. f. Gen.* iii. 426.—China.
tetragona, *Endl. Syn. Conif.* 44.—Chili.

LIBONIA, C. Koch, ex Linden, Prix-Cour. (1863) n. 18. p. 2 = **Jacobinia**, Moric. (Acanthac.).
floribunda, *C. Koch, l. c.*; *et in Wochenschr.* vi. (1863) 266 = J. pauciflora.

LIBONIA, Lem. Illustr. Hortic. ii. (1855) t. 48 = **Billbergia**, Thunb. (Liliac.).
marmorata, Lem. l. c. = B. marmorata.

LIBONIA, Lem. Jard. Fleur. iii. (1852–53) t. 290 = **Griffinia**, Ker-Gawl. (Liliac.).
bicolor, Lem. l. c. = G. bicolor.

LICANIA, Aubl. Pl. Gui. i. 119. t. 45 (1775). *ROSACEAE*, Benth. & Hook f. i. 606.
HEDYCREA, Schreb. Gen. i. 160 (1789).
angelesia, *Blume, Mélanges Bot.* (1855).—*Cf. Flora*, xli. (1858) 256.—Malaya.
aperta, Benth. in Hook. Journ. Bot. ii. (1840) 218 = Moquilea Turiuva.
arborea, *Seem. Bot. Voy. Herald*, 118. t. 25.—N. Granat.
Benthami, *Hook. f. in Mart. Fl. Bras.* xiv. II. 12.—Bras.
bothynophylla, Mart. in Flora, xxiv. (1841) II. Beibl. 15 = Moquilea bothynophylla.
caracasana, Klotzsch, ex Hook. f. in Mart. Fl. Bras. II. 25 = Moquilea Turiuva.
conduplicata, Klotzsch, in Schomb. Faun. & Fl. Gui. 1198 = Moquilea Turiuva.
coriacea, *Benth. in Hook. Journ. Bot.* ii. (1840) 221.—Guian.
costata, Spruce, ex Hook. f. in Mart. Fl. Bras. xiv. II. 10 = glabra.
crassifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 221.—Guian.
crassivenia, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 14.—Bras.
dealbata, *Hook. f. l. c.* 13.—Bras.
Diamenia, *Blume, Mélanges Bot.* (1855).—*Cf. Flora*, xli. (1858) 256.—Malaya.
divaricata, *Benth. in Hook. Journ. Bot.* ii. (1840) 221.—Guian.
emarginata, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 15.—Bras.
flavicans, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1103, nomen.—Guian.
floribunda, Benth. in Hook. Journ. Bot. ii. (1840) 219 = Moquilea floribunda.
glabra, *Mart. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 10.—Bras.
Glazioviana, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 68.—Bras.
guyanensis, Klotzsch, in Schomb. Faun. & Fl. Gui. 1199, nomen; et ex Hook. f. in Mart. Fl. Bras. xiv. II. 12 = heteromorpha.
habantha, *Mart. ex Hook. f. l. c.* 17.—Bras.
helvola, Spruce, ex Hook. f. l. c. 18 = micrantha.
heteromorpha, *Benth. in Hook. Journ. Bot.* ii. (1840) 221.—Guian.
humilis, Cham. & Schlecht. in Linnaea, i. (1826) 549 = Moquilea humilis.
hypoleuca, *Benth. Bot. Voy. Sulph.* 91. t. 32.—N. Granat.
incana, *Aubl. Pl. Gui.* i. 119. t. 45.—Guian.
incana, Benth. in Hook. Journ. Bot. ii. (1840) 220 = Kunthiana.
intrapetiolaris, *Spreng. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 11.—Bras.
Kunthiana, *Hook. f. l. c.* 16.—Bras.
latifolia, *Benth. ex Hook. f. l. c.* 11.—Bras.
leptostachya, Benth. in Hook. Journ. Bot. ii. (1840) 220 = incana.
leucosepala, *Griseb. in Goett. Abh.* vii. (1857) 198.—Ind. occ.
littoralis, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 67.—Bras.
macrophylla, *Benth. in Hook. Kew Journ.* ii. (1850) 240.—Bras.
macrophylla, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1198, nomen.—Guian.
majuscula, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 306.—Guian.
micrantha, Kunth, ex Hook. f. in Mart. Fl. Bras. xiv. II. 16 = Kunthiana.
micrantha, *Miq. Stirp. Surin. Sel.* 29.—Guian.
microcarpa, *Hook. f. in Mart. Fl. Bras.* xiv. II. 15.—Bras.
mollis, *Benth. in Hook. Journ. Bot.* ii. (1840) 219.—Guian.
myristicoides, Benth. ex Hook. f. in Mart. Fl. Bras. xiv. II. 24 = Moquilea sclerophylla.
nitida, *Hook. f. in Mart. Fl. Bras.* xiv. II. 17.—Bras.
obovata, *Benth. ex Hook. f. l. c.* 11.—Bras.
odorata, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1199, nomen.—Guian.
orbicularis, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 13.—Bras.
pallida, Spruce, ex Hook. f. l. c. 18 = parviflora.
parviflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 221.—Guian.
pendula, Benth. l. c. 218 = Moquilea pendula.
polita, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 17.—Bras.
prismatocarpa, *Spruce, ex Hook. f. l. c.* 19.—Bras.
pubiflora, Benth. in Hook. Journ. Bot. ii. (1840) 219 = Moquilea Turiuva.
pyrifolia, *Griseb. Fl. Brit. W. Ind.* 230.—Ins. Trinit.
rigida, *Benth. in Hook. Journ. Bot.* ii. (1840) 220.—Bras.
robusta, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 306.—Guian.
rufuscens, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1103, nomen.—Guian.
scabra, Benth. ex Hook. f. in Mart. Fl. Bras. xiv. II. 24 = Moquilea sclerophylla.
Schomburgkiana, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1199, nomen.—Guian.
sclerophylla, Mart. ex Hook. f. in Mart. Fl. Bras. xiv. II. 24 = Moquilea sclerophylla.
Sellowiana, Klotzsch, ex Hook. f. l. c. 26 = Moquilea bothynophylla.
spicata, *Hook. f. l. c.* 16.—Bras.
subfalcata, Spruce, ex Hook. f. l. c. 18 = parviflora.
surinamensis, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1198, nomen.—Guian.
triandra, *Mart. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 18.—Bras.
turbinata, Benth. in Hook. Journ. Bot. ii. (1840) 218 = Moquilea turbinata.
Turiuva, Cham. & Schlecht. in Linnaea, ii. (1827) 550 = Moquilea Turiuva.
unceolaris, *Hook. f. in Mart. Fl. Bras.* xiv. II. 15.—Bras.

LICARIA, Aubl. Pl. Gui. 313. t. 121 (1775) 'nomen delend.' Benth. & Hook. f. iii. 150.
guianensis, Aubl. l. c. (= *Dicypellium caryophyllatum*, fide Nees).—Guian.

LICHENORA, Wight, Ic. v. t. 1738 (1852) = **Eria**, Lindl. (Orchid.).
Jerdoniana, Wight, l. c. = E. Lichenora.

LICHINORA, Wight, Ic. v. 18 (1852) = praec.

LICHNIS, Crantz, Inst. i. 194 (1766) = **Lychnis**, Tourn. (Caryoph.).

LICHTENSTEINIA, Cham. & Schlecht. in Linnaea, i. (1826) 394. t. 5. *UMBELLIFERAE*, Benth. & Hook. f. i. 887.
RUTHEA, Bolle, in Verh. Bot. Ver. Brandenb. iii.-iv. (1861–62) 174.

LICHSTENSTEINIA:—
Beiliana, *Eckl. & Zeyh. Enum.* 343.—Afr. austr.
Burchellii, *Hook. f. in Hook. Ic. Pl.* t. 1033.—Ins. St. Helena.
crassijuga, *E. Mey. in Drège, Zwei Pfl. Docum.* 199.—Afr. austr.
inebrians, Eckl. & Zeyh. Enum. 343 = Beiliana.
inebrians, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 548 (= *Glia gummifera*).—Afr. austr.
interrupta, E. Mey. in Drège, Zwei Pfl. Docum. 199 = pyrethrifolia.
lacera, *Cham. & Schlecht. in Linnaea*, i. (1826) 395.—Afr. austr.
latifolia, *Eckl. & Zeyh. Enum.* 343.—Afr. austr.
palmata, DC. Prod. iv. 135 = trifida.
pyrethrifolia, *Cham. & Schlecht. in Linnaea*, i. (1826) 397.—Afr. austr.
runcinata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 543 = trifida.
Sprengeliana, Eckl. & Zeyh. Enum. 344 = pyrethrifolia.
trifida, *Cham. & Schlecht. in Linnaea*, i. (1826) 396.—Afr. austr.
triradiata, E. Mey. in Drège, Zwei Pfl. Docum. 199 = Beiliana.

LICHTENSTEINIA, Wendl. Coll. i. II. 4. t. 39 (1808) = **Loranthus**, Linn.
oleaefolia, Wendl. l. c. 5 = Loranthus oleaefolius.

LICHTENSTEINIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 19. t. 1 = **Ornithoglossum**, Salisb. (Liliac.).
laevigata, Willd. l. c. 20 = O. glaucum.
undulata, Willd. l. c. = O. glaucum.

LICHTERVELDIA, Lem. Illustr. Hortic. ii. (1855) t. 59 = **Odontoglossum**, H. B. & K. (Orchid.).
Lindleyi, Lem. l. c. = O. citrosmum.

LICINIA, Rafin. Fl. Tellur. iii. 57 (1836) = **Anthericum**, Linn. (Liliac.).
canaliculata, Rafin. l. c. = A. canaliculatum.

LICOPERSICUM, Neck. Elem. ii. 61 (1790) = **Lycopersicum**, Mill. (Solan.).

LICOPSIS, Neck. Delic. Gallo-Belg. i. 102 (1768) = **Lycopsis**, Linn. (Borag.).

LICOPUS, Neck. Delic. Gallo-Belg. i. 14 (1768) = **Lycopus**, Tourn. (Labiat.).

LICUALA, Thunb. in Vet. Acad. Nya Handl. (1782) 284; Diss. Nov. Gen. iii. 70. *PALMAE*, Benth. & Hook. f. iii. 928.
PERICYCLA, Blume, Rumphia, ii. 47. t. 94 (1843).
acutifida, *Mart. Hist. Nat. Palm.* iii. 236. t. 135. f. 34.—Malaya.
amplifrons, *Miq. Fl. Ind. Bat. Suppl.* 591.—Sumatra.
aruensis, *Becc. Malesia*, i. (1877) 83.—N. Guin.
australasica, *Hort. ex H. Wendl. in Kerch. Palm.* 249, nomen.—Australas.
bacularia, *Becc. Malesia*, i. (1877) 82.—N. Guin.
Bissula, *Miq. Fl. Ind. Bat.* iii. 57.—Ins. Celebes.
celebica, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 11.—Ins. Celebes.
densiflora, *Hort. ex H. Wendl. in Kerch. Palm.* 249, nomen.—Hab.?
elegans, *Blume, Rumphia*, ii. 42.—Sumatra.
Flabellum, *Mart. Hist. Nat. Palm.* iii. 237.—Ins. Celebes.
glabra, *Griff. ex Mart. l. c.* 238; *et in Calc. Journ. Nat. Hist.* v. (1845) 329.—Malaya.
gracilis, *Blume, in Schult. f. Syst.* vii. 1308.—Java.
grandis, *H. Wendl. in Illustr. Hortic.* xxvii. (1880) t. 412.—Nov. Brit.
horrida, Blume, Rumphia, ii. 39 = spinosa.
insignis, *Becc. Malesia*, i. (1877) 80.—N. Guin.
longipes, *Griff. ex Mart. Hist. Nat. Palm.* iii. 237; *et in Calc. Journ. Nat. Hist.* v. (1845) 330.—Malaya.
Muelleri, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 223.—Austral.
nana, *Blume, Rumphia*, ii. 46. t. 93.—Sumatra.
Oxleyi, H. Wendl. in Kerch. Palm. 249 = paludosa.

LICUALA:—
paludosa, *Griff. ex Mart. Hist. Nat. Palm.* iii. 237; *et in Calc. Journ. Nat. Hist.* v. (1845) 323.—Malaya.
peltata, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 179.—Reg. Himal.; Burma; Ind. or.
penduliflora, [*Zipp. ex*] *Mackl. in Bijdr. Nat. Wet.* v. (1830) 178, nomen; *et ex Blume, Rumphia,* ii. 47.—N. Guin.
pilearia, *Blume, Rumphia,* ii. 42.—Cochinch.
pumila, *Blume, in Schult. f. Syst.* vii. 1302.—Malaya.
ramosa, Blume, l. c. 1303 = spinosa.
rotundifolia, Blume, l. c. 1305 = Livingstona rotundifolia.
Rumphii, *Blume, Rumphia,* ii. 41.—Ins. Molucc.
spectabilis, *Miq. Pl. Jungh.* 163.—Java.
spinosa, Poir. Encyc. Suppl. iii. 430; Blume, Rumphia, ii. 41 = Rumphii.
spinosa, Roxb. Hort. Beng. 25; Fl. Ind. ii. 181 = acutifida.
spinosa, *Thunb. in Vet. Akad. Nya Handl.* (1782) 287.—Malaya.
telifera, *Becc. Malesia,* i. (1877) 81.—N. Guinea.
ternata, *Griff. ex Mart. Hist. Nat. Palm.* iii. 238.—Malaya.
triphylla, Griff. ex Mart. l. c.; et in Calc. Journ. Nat. Hist. v. (1845) 332 = ternata.
Waraguh, Blume, in Schult. f. Syst. vii. 1305 = Rhapis javanica.
Wiru, Mart. Hist. Nat. Palm. iii. 254 = seq.
Wiruh, Blume, Rumphia, ii. 56 = seq.
Wixu, Blume, in Schult. f. Syst. vii. 1305 = Rhapis javanica.

LIDBECKIA, Berg. Desc. Pl. Cap. 306. t. 5. f. 9 (1767). *COMPOSITAE*, Benth. & Hook. f. ii. 423.
LANCISIA, Lam. Illustr. t. 701. f. 2, 3 (1798 ?).
americana, *Spreng. Syst.* iii. 589.—Am. austr.
bipinnata, Thunb. Prod. Pl. Cap. 161 = Lasiospermum radiatum.
capensis, J. F. Gmel. Syst. 227 = pectinata.
lobata, *Thunb. Prod. Pl. Cap.* 161.—Afr. austr.
pectinata, *Berg. Desc. Pl. Cap.* 307. t. 5. f. 9.—Afr. austr.
pruinosa, E. Mey. ex DC. Prod. vi. 82 = Cenia turbinata.
quinqueloba, Cass. in Dict. Sc. Nat. xxvi. 276 = lobata.
turbinata, Thunb. Prod. Pl. Cap. 161 = Cenia turbinata.

LIDBEKIA, Spreng. Gen. ii. 635 (1831) = praec.

LIEBERKUEHNA, Reichb. Consp. 100 (1828) = seq.

LIEBERKUEHNIA, Reichb. Nom. 87 (1841) = seq.

LIEBERKUHNA, Cass. in Dict. Sc. Nat. xxvi. 286 (1823) = **Chaptalia**, Vent. (Compos.).
bracteata, Cass. l. c. 286 = C. piloselloides.
nudipes, Cass. l. c. 287 = C. sinuata.
piloselloides, Steud. Nom. ed. II. ii. 726 = C. piloselloides.
spathulata, DC. Prod. vii. 43 = C. spathulata.

LIEBERKUHNIA, Less. Syn. Comp. 122 (1832) = praec.

LIEBICHIA, Opiz, Seznam, 59 (1852) = **Ribes**, Linn. (Saxifrag.).
alpina, Opiz, l. c. = R. alpinum.

LIEBIGIA, Endl. Gen. 1407 (1841) = **Chirita**, Buch.-Ham. (Gesner.).
bracteosa, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 574 = C. Horsfieldii.
elongata, DC. Prod. ix. 259 = Dichrotrichum ternateum.
speciosa, DC. l. c. = C. Blumii.

LIETZIA, Regel, Gartenfl. (1880) 97. *GESNERACEAE.*
brasiliensis, *Regel & Schmidt, l. c.*—Bras.

LIEVENA, Regel, in Act. Hort. Petrop. vi. (1879) 537 = **Quesnelia**, Gaudich. (Bromel.).
Princeps, Regel, l. c. = Q. rufa.

LIGEA, Poit. ex Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96; et in Arch. Mus. Par. vi. (1852) (Podost. Monog.) 85. tt. 4–7. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 110.
OENONE, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96.
alcicornis, *Tul. in Arch. Mus. Par.* vi. (1852) 94. t. 7. f. 1.—Bras.
corymbosa, *Benth. & Hook. f. Gen.* iii. 111.—Guian.
dissecta, *Benth. & Hook. f. l. c.*—Bras.
flexuosa, *Tul. Podost. Monog.* 88.—Guian.
longifolia, *Tul. l. c.* 86.—Guian.
Richardiana, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96.—Guian.
secundiflora, *Tul. l. c.* 97.—Guian.

LIGERIA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 463 = **Sinningia**, Nees (Gesner.).
andina, Griseb. in Goett. Abh. xix. (1874) 227.—Reg. Argent.
barbata, Hanst. in Mart. Fl. Bras. viii. 390 = S. Carolinae.
caulescens, Decne. in Rev. Hortic. Sér. III. ii. (1848) 463 = S. speciosa.
concinna, Hanst. in Mart. Fl. Bras. viii. 386.—Bras.
digitaliflora, Decne. in Rev. Hortic. Sér. III. ii. (1848) 463 = Gloxinia digitaliflora.
discolor, Decne. ex Hanst. in Mart. Fl. Bras. viii. 388.—Bras.
gesnerifolia, Hanst. l. c. 389 = S. villosa.
gracilis, Hanst. l. c. 390.—Bras.
hirsuta, Decne. ex Hanst. l. c. 386.—Bras.
macrophylla, Hanst. l. c. 387.—Bras.
Maximiliana, Hanst. l. c.—Bras.
Menziesiana, Hanst. l. c. 388 = S. speciosa.
pusilla, Hanst. l. c. 386.—Bras.
speciosa, Decne. in Rev. Hortic. Sér. III. ii. (1848) 464 = S. speciosa.
Teuchleri, Decne. l. c. = *Gloxinia Teuchleri.*
villosa, Hook. ex Hanst. in Mart. Fl. Bras. viii. 420 = S. villosa.

LIGHTFOOTIA, L'Hérit. Sert. Angl. t. 4 (1788). *CAMPANULACEAE*, Benth. & Hook. f. ii. 555.
CONCILIUM, Rafin. Fl. Tellur. ii. 78 (1836).
abyssinica, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 25, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 1.—Abyss.
adpressa, A. DC. Monog. Camp. 110 = Wahlenbergia adpressa.
albanensis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 557.—Afr. austr.
albens, *Spreng. ex A. DC. Monog. Camp.* 110.—Afr. austr.
albicaulis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 556.—Afr. austr.
angustifolia, A. DC. in DC. Prod. vii. 418 = rubens.
annua, *DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 329.—Afr. austr.
anomala, A. DC. in DC. Prod. vii. 418 = Thunbergiana.
arenaria, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 329.—Afr. trop.
axillaris, *Sond. in Harv. & Sond. Fl. Cap.* iii. 558.—Afr. austr.
Buekii, *Sond. l. c.* iii. 560.—Afr. austr.
caledonica, *Sond. l. c.* 559.—Afr. austr.
capillaris, *Buek, in Eckl. & Zeyh. Enum.* 372.—Afr. austr.
ciliata, Sond. in Harv. & Sond. Fl. Cap. iii. 561 = tenella.
ciliata, Spreng. Syst. i. 809 = Prismatocarpus subulatus.
cinerea, Sond. in Harv. & Sond. Fl. Cap. iii. 557 = grisea.
collomioides, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 328.—Afr. trop.
debilis, *A. DC. l. c.* 331.—Afr. trop.
denticulata, Sond. in Harv. & Sond. Fl. Cap. iii. 559 = capillaris.
diffusa, Buek, in Eckl. & Zeyh. Enum. 376 = fasciculata.
divaricata, Buek, l. c. 375 = tenella.
exilis, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 330.—Afr. trop.
fasciculata, *Spreng. ex A. DC. in Monog. Camp.* 112.—Afr. austr.
gracilis, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 330.—Afr. trop.
gracilis, Miq. Fl. Ind. Bat. ii. 567 = Wahlenbergia gracilis.

LIGHTFOOTIA:—
grisea, *Buek, in Eckl. & Zeyh. Enum.* 374.—Afr. austr.
Huttonii, *Sond. in Harv. & Sond. Fl. Cap.* iii. 556.—Afr. austr.
intermedia, Buek, ex Eckl. & Zeyh. Enum. 374 = uitenhagensis.
juncea, *Sond. in Harv. & Sond. Fl. Cap.* iii. 563.—Afr. austr.
lanceolata, Link, Enum. Hort. Berol. i. 217 = oxyococoides?
laricina, Buek, in Eckl. & Zeyh. Enum. 373 = albens, Buekii.
laxiflora, *Sond. in Harv. & Sond. Fl. Cap.* iii. 564.—Afr. austr.
linearis, Eckl. ex A. DC. in DC. Prod. vii. 417 = longifolia.
Loddigesii, A. DC. Monog. Camp. 114 = tenella.
longifolia, *A. DC. l. c.* 107.—Afr. austr.
lycopodioides, A. DC. l. c. 114 = unidentata.
macrostachys, A. DC. l. c. 787 = spicata.
madagascariensis, *A. DC. Monog. Camp.* 116.—Madag.
marginata, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 326.—Afr. trop.
mucronulata, Buek, in Eckl. & Zeyh. Enum. 375 = tenella.
muscosa, Link, Enum. Hort. Berol. i. 217 = oppositifolia.
namaquana, *Sond. in Harv. & Sond. Fl. Cap.* iii. 556.—Afr. austr.
napiformis, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 328.—Afr. trop.
nodosa, Buek, in Eckl. & Zeyh. Enum. 376 = fasciculata.
oppositifolia, *A. DC. Monog. Camp.* 115.—Afr. austr.
oxycoccoides, Drège, ex A. DC. in DC. Prod. vii. 418 = spicata.
oxycoccoides, *L'Hérit. Sert. Angl.* t. 4.—Afr. austr.
oxycoccoides, Spreng. Syst. i. 809, partim = oppositifolia.
paniculata, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 331.—Afr. trop.
paniculata, *Sond. in Harv. & Sond. Fl. Cap.* iii. 560.—Afr. austr.
pubescens, *A. DC. in DC. Prod.* vii. 419.—Afr. austr.
robusta, A. DC. l. c. 420 = Wahlenbergia rigida.
rubens, *Buek, in Eckl. & Zeyh. Enum.* 373.—Afr. austr.
rubioides, Buek, l. c. 377 = ciliata.
rubioides, *A. DC. Monog. Camp.* 116.—Afr. austr.
sessiliflora, Sond. in Harv. & Sand. Fl. Cap. iii. 555 = subulata.
sessiliflora, Spreng. Syst. i. 809 = Prismatocarpa strictus.
sessilifolia, Zeyh. ex A. DC. in DC. Prod. vii. 498 = Cyphia linarioides.
spicata, *Buek, in Eckl. & Zeyh. Enum.* 374.—Afr. austr.
subaphylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 193.—Madag.
subulata, Drège, ex A. DC. in DC. Prod. vii. 418 = spicata.
subulata, *L'Herit. Sert. Angl.* t. 5.—Afr. trop.
tenella, A. DC. Monog. Camp. 111. t. 3. f. B = fasciculata.
tenella, *Lodd. Bot. Cab.* t. 1038.—Afr. austr.
tenerrima, Buek, in Eckl. & Zeyh. Enum. 375 = fasciculata.
tenuifolia, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 327.—Afr. trop.
Thunbergiana, *Buek, in Eckl. & Zeyh. Enum.* 376.—Afr. austr.
thymifolia, *Buek, l. c.* 373.—Afr. austr.
uitenhagensis, *Buek, l. c.*—Afr. austr.
unidentata, *A. DC. Monog. Camp.* 107.—Afr. austr.
Welwitschii, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 332.—Afr. trop.

LIGHTFOOTIA, Schreb. Gen. i. 122 (1789) = **Rondeletia**, Linn. (Rubiac.).

LIGHTFOOTIA, Sw. Prod. Veg. Ind. Occ. 83 (1788) = **Laetia**, Linn. (Bixin.).
indica, Sw. l. c.—Ind. occ.
integrifolia, Vahl, Symb. Bot. iii. 70 = Aphloia mauritiana.
serrata, Sw. Prod. Veg. Ind. Occ. 83 = Aphloia mauritiana.
theaeformis, Vahl, Symb. Bot. iii. 69 = Aphloia mauritiana.

LIGHTIA, Schomb. in Linnaea, xx. (1847) 757. *VOCHYSIACEAE.*
guianensis, *Schomb. l. c.*—Guian.
licanoides, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 122.—Bras.

LIGHTIA, Schomb. in Rep. Brit. Assoc. (1844) II. 71 = **Herrania**, Goudot. (Stercul.).
lemniscata, Schomb. l. c. = H. Mariae.

LIGIA, Fasano, ex Pritz. Ic. Ind. 638 (1855) = LYGIA, Fasano = **Thymelaea**, Tourn.

LIGNONIA, Scop. Introd. 292 (1777) = **Paypayrola**, Aubl. (Violar.).
monodynama, J. F. Gmel. Syst. 373 = P. guianensis.

LIGOSTICON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = **Ligusticum**, Linn. (Umbellif.).
aciphyllum, St. Lag. l. c. = Ligust. Aciphylla.

LIGTU, Adans. Fam. p. (20) (1763) = **Alstroemeria**, Linn. (Amaryll.).

LIGULARIA, Cass. in Bull. Soc. Philom. (1816) 198 = **Senecio**, Tourn. (Compos.).
altaica, DC. Prod. vi. 315 = S. altaicus.
amplexicaulis, DC. l. c. 314 = S. amplexicaulis, *Wall.*
araneosa, DC. l. c. = S. chionopappus.
arnicoides, DC. l. c. = S. arnicoides.
Aucheri, DC. l. c. vii. 300 = S. taraxacifolius.
auriculata, Turcz. ex DC. l. c. vi. 329 = S. dahuricus.
calthaefolia, Maxim. in Bull. Acad. Pétersb. xv. (1871) 374 = S. calthaefolius.
caucasica, G. Don, in Sweet, Hort. Brit. ed III. 382 = S. caucasicus.
clivorum, Maxim. in Bull. Acad. Pétersb. xv. (1871) 374 = S. clivorum.
corymbosa, DC. Prod. vi. 314 = S. amplexicaulis, *Wall.*
euodon, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 188 = S. Ligularia.
Farfugium, C. Koch, in Regel, Gartenfl. (1859) 15 = S. Kaempferi.
Fischeri, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) II. 11 = *speciosa.*
frigida, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 351 = S. arnicoides.
geifolia, DC. Prod. vi. 316 = S. geifolius.
gigantea, Sieb. & Zucc. Fl. Jap. i. 79. t. 36 = S. Kaempferi.
hastata, Less. Syn. Comp. 390 = S. sagittatus.
heterophylla, Rupr. Sert. Tiansch. 53.—Turkestan.
Hodgsoni, Hook. f. Bot. Mag. t. 5417.—Japon.
japonica, Less. Syn. Comp. 390 = S. japonicus.
Kaempferi, Sieb. & Zucc. Fl. Jap. i. 77. t. 35 = S. Kaempferi.
Kareliniana, Stschegl. in Bull. Soc. Nat. Mosc. (1854) I. 176.—As. centr.
lanata, Less. Syn. Comp. 390 = Alciope lanata.
macrophylla, DC. Prod. vi. 316 = S. Ledebourii.
mongolica, DC. l. c. 315 = S. mongolicus.
nana, Decne. in Jacquem. Voy. Bot. 91. t. 99 = Werneria nana.
palmata, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 168 = Cremanthodium palmatum.
persica, Boiss. Diagn. Ser. I. vi. 93 = S. persicus.
pinnatifida, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 167 = Cremanthodium pinnatifidum.
racemosa, DC. Prod. vi. 314 = S. Ligularia.
Reichenbachii, Steud. Nom. ed. II. ii. 42 = S. Ligularia.
renifolia, DC. Prod. vi. 316 = S. renifolius.
reniformis, DC. l. c. 315 = Cremanthodium reniforme.
retusa, DC. l. c. 314 = S. retusus.
robusta, DC. l. c. 316 = S. robustus, *Sch. Bip.*
sibirica, Cass. in Dict. Sc. Nat. xxvi. 402 = S. Ligularia.
speciosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 38 = S. Ligularia?
tabularis, Less. Syn. Comp. 290 = Alciope tabularis.
thyrsoidea, DC. Prod. vi. 315 = S. sibiricus.

LIGULARIA, Duval, Pl. Succ. Hort. Alenç. 11 (1809) = **Saxifraga**, Linn.
sarmentosa, Duval, l. c. = S. sarmentosa.

LIGULARIA, Eckl. & Zeyh. Enum. 69 (1835) = **Pelargonium**, L'Hérit. (Geraniac.).
canescens, Eckl. & Zeyh. l. c. = P. incisum.
dissecta, Eckl. & Zeyh. l. c. = P. dissectum, *Steud.*
hirta, Eckl. & Zeyh. l. c. 70 = P. tenuifolium.
incisa, Eckl. & Zeyh. l. c. 69 = P. incisum.
ionidiflora, Eckl. & Zeyh. Enum. 69 = P. incisum.
ramosissima, Eckl. & Zeyh. l. c. 70 = P. ramosissimum.
tenuifolia, Eckl. & Zeyh. l. c. = P. tenuifolium.
trifida, Eckl. & Zeyh. l. c. 69 = P. tripartitum.
villosa, Eckl. & Zeyh. l. c. = P. villosum.

LIGUSTICUM, Linn. Gen. ed. I. 76 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 911.
CONIOSELINUM, Fisch. in Hoffm. Gen. Umb. 185 (1816).
CORISTOSPERMUM, Bertol. Fl. Ital. iii. 466 (1837).
CYNAPIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 640 (1840).
ENDRESSIA, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 223.
GAYA, Gaud. Fl. Helv. ii. 389 (1828).
HALOSCIAS, Fries, Summ. Veg. Scand. 180 (1846).
HANSENIA, Turcz. in Bull. Soc. Nat. Mosc. xvii. 754 (1844) (Fl. Baic. Dahur. i. 513).
NEOGAYA, Meissn. Gen. 144 (1838).
PACHYPLEURUM, Ledeb. Fl. Alt. i. 296 (1829).
acaule, *Turcz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 386, nomen.—Sibir.
Aciphylla, *Spreng. Umbell. Prod.* 40; & *in Schult. Syst.* vi. 554.—N. Zel.
actaeifolium, Bigel. Fl. Bost. ed. III. 118 = Archangelica Gmelini.
actaeifolium, *Michx. Fl. Bor. Am.* i. 166.—Am. bor.
acutilobum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 203.—Japon.
Ajawain, Roxb. ex Flem. in As. Res. xi. (1810) 171 = Carum copticum.
Ajawain, Spreng. in Schult. Syst. vi. 556 = Ptychotis Ajowan.
alatum, *Boiss. Fl. Orient.* ii. 972.—Reg. Caucas.
alatum, *Spreng. Umbell. Prod.* 40; *Umbell. Spec.* 125.—Reg. Caucas.
alpinum, Spreng. Syst. i. 908 = Carum alpinum.
alsaticum, Link, Enum. Hort. Berol. i. 275 = Peucedanum alsaticum.
Angelica, Stokes, Bot. Mat. Med. ii. 96 = Archangelica officinalis.
angelicoides, Wall. Cat. n. 548 = Pleurospermum angelicoides.
angustilobum, *Phil. in Linnaea*, xxxiii. (1864-65) 95.—Chili.
anisatum, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 89.—N. Zel.
antipodum, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 63.—Ins. Dom. Auckland.
apiifolium, Desf. Tabl. ed. II. 144 = Carum rigidulum.
apiifolium, *A. Gray, in Proc. Am. Acad.* vii. (1868) 347.—Am. bor.
apiifolium, S. Wats. Bot. King's Exped. 125 = filicinum.
apioides, Lam. Encyc. iii. 577 = Carum rigidulum.
apioides, *Phil. in Anal. Univ. Chil.* (1872) 726.—Chili.
aquilegifolium, Willd. Sp. Pl. i. 1425 = Physospermum aquilegifolium.
Archangelica, Geners. Fl. Scep. Elench. n. 252; ex DC. Prod. iv. 244 = Pleurospermum austriacum.
arenarium, Link, Enum. Hort. Berol. i. 276 = Silaus tenuifolius.
argutum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 641 = Arracacia arguta (in Addend.).
aromaticum, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 89.—N. Zel.
athamanthicum, Adams, ex DC. Prod. iv. 151 = Schultzia compacta.
athamantoides, Spreng. Umbell. Prod. 40 = Seseli athamanthoides.
austriacum, Linn. Sp. Pl. 250 = Pleurospermum austriacum.
balearicum, Linn. Mant. ii. 218 = Kundmannia sicula.
barbinode, Michx. Fl. Bor. Am. i. 167 = Thaspium barbinode.
Benthami, Wall. Cat. n. 542 = Pleurospermum Benthami.
biternatum, Stokes, Bot. Mat. Med. ii. 94 = scoticum.
boreale, Salisb. Prod. 164 = scoticum.
Brancionis, Schrank, Baier. Fl. i. 551 = Molopospermum cicutarium.
brevistyle, *Hook. f. Handb. N. Zel. Fl.* 95.—N. Zel.
Brunonis, Wall. Cat. n. 545 = Pleurospermum Brunonis.
Bulbocastanum, Crantz, Cl. Umbell. Emend. 83 = Carum Bulbocastanum.
bulbosum, Boiss. Fl. Orient. ii. 882 = Carum multiflorum.
bulbosum, Pers. Syn. i. 315 = Erigenia bulbosa.

LIGUSTICUM:—
Bunius, Link, Enum. Hort. Berol. i. 276 = Carum Bunius.
candicans, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 348.—Hab.?
Candollei, Wall. Cat. n. 544 = Pleurospermum Candollei.
capense, *DC. Prod.* iv. 159.—Afr. austr.
capillaceum, Lam. Fl. Fr. iii. 454 = Meum athamanticum.
carniolicum, Host, Fl. Austr. i. 378 = Pleurospermum Golaka.
carnulosum, *Hook. f. Handb. N. Zeal. Fl.* 96.—N. Zel.
Carvi, Roth, Tent. Fl. Germ. i. 124 = Carum Carvi.
caucasicum, Willd. ex Schult. Syst. vi. 457 = Eleutherospermum grandifolium.
Cervaria, Spreng. Umbell. Prod. 40 = Peucedanum Cervaria.
cicutaefolium, Vill. Hist. Pl. Dauph. ii. 612 = Carum rigidulum.
cicutarium, Lam. Fl. Fr. iii. 453 = Molopospermum cicutarium.
commutatum, *Regel, Ind. Sem. Hort. Petrop.* (1867) 34.—Hab.?
condensatum, Spreng. Umbell. Prod. 40 = Seseli Libanotis.
coniifolium, DC. Prod. iv. 158 = Selinum tenuifolium.
cornubiense, Linn. Cent. Pl. ii. 13; Amoen. Acad. iv. 310 = Physospermum aquilegifolium.
orsicum, *J. Gay, in Ann. Sc. Nat.* Sér. I. xxvi. (1832) 222.—Corsica.
Cuminum, Crantz, Cl. Umbell. Emend. 82 = Cuminum Cyminum.
cuneifolium, *Guss. Pl. Rar.* 130. t. 26.—Ital.
cynapifolium, *Viv. ex DC. Prod.* iv. 158.—Corsica.
cyprium, Spreng. Umbell. Sp. 125 = Carum multiflorum.
decussatum, Moench, Meth. 81 = Peucedanum carvifolium.
deltoideum, *Cheesm. in Trans. N. Z. Inst.* xiv. (1882) 299.—N. Zel.
dentatum, Wall. Cat. n. 547 = Pleurospermum dentatum.
Dieffenbachii, Hook. f. Handb. N. Zeal. Fl. 729 = Angelica Dieffenbachii.
diffusum, Roxb. ex Sm. in Rees, Cycl. xxi. n. 11; et in Roxb. Hort. Beng. 21; Fl. Ind. ii. 92 = Seseli indicum.
discolor, *Ledeb. Fl. Alt.* i. 321.—Sibir. altaic.
divaricatilobum, *Clos, in C. Gay, Fl. Chil.* iii. 131.—Chili.
divaricatum, Ledeb. Ind. Sem. Hort. Dorpat. (1824) 5 = Silaus longifolius.
dubium, *H. B. & K. Nov. Gen. et Sp.* v. 19.—Mexic.
dubium, *Spreng. Fl. Hal. Mant.* 37.—Hab.?
dumetorum, *Phil. in Linnaea*, xxxiii. (1864-65) 96.—Chili.
elatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 698.—Reg. Himal.
Enysii, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 548.—N. Zel.
ferulaceum, *All. Fl. Pedem.* ii. 13. t. 60. f. 1.—Europ.
ferulaceum, Lam. Fl. Fr. iii. 453 = pyrenaicum.
ferulaceum, Lapeyr. Hist. Abr. Pl. Pyr. 155 = Seseli Libanotis.
filicinum, *S. Wats. in Proc. Am. Acad.* xi. (1876) 140.—Am. bor. occ.
filifolium, *Hook. f. Handb. N. Zeal. Fl.* 95.—N. Zel.
Fischeri, D. Dietr. Syn. Pl. ii. 960 = Selinum Benthami.
Fischeri, Link, Enum. Hort. Berol. i. 276 = Silaus longifolius.
Foeniculum, Roth, Tent. Fl. Germ. i. 124 = Foeniculum vulgare.
Fontanesii, Spreng. in Schult. Syst. vi. 565 (= *Cnidium Fontanesii*).—Numidia.
garganicum, Tenore, Prod. Fl. Nap. p. xix = Laserpitium Siler.
Gingidium, Forst. f. Prod. 22 = Angelica Gingidium.
Gmelini, Cham. & Schlecht. in Linnaea, i. (1826) 391 = Selinum Benthami.
Gmelini, Vill. Prosp. 24; Hist. Pl. Dauph. ii. 610. t. 13 = Pleurospermum austriacum.
Govanianum, Wall. Cat. n. 546; et ex DC. Prod. iv. 246 = Pleurospermum Govanianum.
graecum, DC. Prod. iv. 159 = Carum multiflorum.
Haastii, *F. Muell. ex Hook. f. Handb. N. Zeal. Fl.* 95, 730.—N. Zel.
Hacquetii, Guss. Pl. Rar. 132 = Pleurospermum austriacum.

LIGUSTICUM:—
Halleri, Vill. Prosp. 25 (Quid?).—Gallia.
hirsutum, Crantz, Stirp. Austr. ed. I. fasc. iii. 74 = Chaerophyllum hirsutum.
humile, *Phil. in Anal. Univ. Chil.* (1872) 726.—Chili.
imbricatum, *Hook. f. Handb. N. Zeal. Fl.* 97.—N. Zel.
indicum, Wall. Cat. n. 7215 = Seseli indicum.
intermedium, *Hook. f. Handb. N. Zeal. Fl.* 94.—N. Zel.
japonicum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 270.—Japon.
kamtschaticum, Steud. Nom. ed. I. 478 = Pleurospermum kamtschaticum.
Kellerianum, *Vis. in Mem. Ist. Venet.* vi. (1856) 258.—Europ.
Landbeckii, *Phil. in Linnaea*, xxxiii. (1864–65) 95.—Chili.
latifolium, *Hook. f. Handb. N. Zeal. Fl.* 94.—N. Zel.
Levisticum, Linn. Sp. Pl. 250 = Levisticum officinale.
Lobelii, Vill. Prosp. 24 = Carum rigidulum.
longifolium, Willd. Sp. Pl. i. 1428 = Peucedanum ruthenicum.
lucidum, Mill. Gard. Dict. ed. VIII. n. 4 = pyrenaicum.
luteum, Poir. Voy. Barb. ii. 136 = Ferula sulcata.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 95.—N. Zel.
marginatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 698.—Reg. Himal.
Massalongianum, *Vis. in Mem. Ist. Venet.* vi. (1856) 259.—Europ.
meoides, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 671 = Vicatia coniifolia.
Meum, Crantz, Stirp. Austr. ed. I. fasc. iii. 82; ed. II. 199 = Meum athamanticum.
minus, Lam. Fl. Fr. iii. 454 = Selinum Monnieri.
montanum, *Benth. & Hook. f. Gen.* i. 912; *A. Gray, in Proc. Am. Acad.* vii. (1868) 347.—Am. bor. occ.
montanum, Spreng. Syst. i. 907 = Apium montanum.
multifidum, *Sm. in Rees, Cycl.* xxi. n. 8.—Sibiria.
Mutellina, *Crantz, Stirp. Austr.* ed. I. fasc. iii. 81.—Europ.
mutellinoides, Vill. Prosp. 25 = simplex.
nemorosum, Phil. in Linnaea, xxxiii. (1864–65) 94 = Pimpinella nemorosa.
nepalense, *D. Don, Prod. Fl. Nep.* 125.—Nepal.
nodiflorum, Vill. Prosp. 25 = Trochiscanthes nodiflora.
nodosum, Willd. ex Schult. Syst. vi. 570 = Ferula nodosa.
obtusifolium, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 23.—Afr. bor.
obtusiusculum, Wall. Cat. n. 543 = Trachydium obtusiusculum.
Pansil, *Bert. ex DC. Prod.* iv. 669.—Chili.
Pansil, Phil. ex Benth. & Hook. f. Gen. i. 912 = Pimpinella Pansil.
Panul, Clos, in C. Gay, Fl. Chil. iii. 131 = L. Pansil.
peloponense, Falk, Beitr. ii. 147 (Quid?).—Rossia.
peloponesiacum, Scop. Fl. Carn. ed. II. i. 205 = Molopospermum cicutarium.
peloponnense, Lam. Encyc. iii. 576 = Molopospermum cicutarium.
peloponnense, [Retz.] in Murr. Syst. Veg. ed. XIII. 233 = Retzii.
peloponnesiacum, Georgi, Reise, i. 204 = Pleurospermum uralense.
peloponnesiacum, Linn. Sp. Pl. 250 = Thapsia foetida?
peloponnesiacum, Pall. Reise, ii. 124 = discolor.
peregrinum, Linn. Sp. Pl. ed. II. 360 = Carum peregrinum.
peucedanoides, *Presl, ex DC. Prod.* iv. 158 (sp. dub.).—Chili.
Phelandrium, Roth, Tent. Fl. Germ. i. 123 = seq.?
Phellandrium, Crantz, Stirp. Austr. ed. I. fasc. iii. 84; ed. II. 200 = Oenanthe Phellandrium.
piliferum, *Hook. f. Handb. N. Zeal. Fl.* 96.—N. Zel.
pimpinellifolium, *Phil. in Linnaea*, xxxiii. (1864–65) 94.—Chili.
pinnovelum, *Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) 14.—Hab.?
Podagraria, Crantz, Stirp. Austr. ed. I. fasc. iii. 84; ed. II. 200 = Aegopodium Podagraria.
pumilum, Wall. Cat. n. 550; ex DC. Prod. iv. 245 = Pleurospermum pumilum.
pusillum, Pers. Syn. i. 315 = Apium divaricatum.
pyrenaicum, *Gouan, Illustr.* 14. t. 7.—Mont. Pyren.
resinosum, Guss. Fl. Sic. Prod. i. 335 = Ferula nudicaulis.
Retzii, *Vitm. Summa Pl.* ii. 158.—Hab.?
rotundatum, Wall. Cat. n. 549; et ex DC. Prod. iv. 245 = Pleurospermum rotundatum.

LIGUSTICUM:—
saxifragum, Boiss. & Sprun. in Ann. Sc. Nat. Sér. III. i. (1844) 300 = Carum multiflorum.
scopulorum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 347.—Am. bor. occ.
scopulorum, Parry, in Am. Natural. ix. (1875) 271 = filicinum.
scoticum, *Linn. Sp. Pl.* 250 (scothicum.).—Europ.; As. et Am. bor.
scoticum, Lour. Fl. Cochinch. 224 (Quid?).—Cochinch.
Segueri, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 105.—Alp. Europ. austr.
Seguieri, Vill. Prosp. 25 = ferulaceum.
seseloides, *Fisch. & Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 725.—Davuria.
sibiricum, Spreng. Prod. Umbell. 40 = Seseli Libanotis.
silaifolium, Gaud. Fl. Helv. ii. 399 = Carum rigidulum.
Silaus, Vill. Prosp. 25; All. Fl. Pedem. ii. 14 = Silaus pratensis.
simplex, *All. l. c.* ii. 15.—Europ.
simplex, Benth. Cat. Pl. Pyr. 95 = pyrenaicum.
splendens, Lapeyr. Hist. Abr. Pl. Pyr. 146 (= *Wallrothia tenuifolia*).—Mont. Pyren.
Sprengelii, Sieber, ex Spreng. Umbell. Sp. 124 = Pleurospermum Golaka.
striatum, DC. Prod. iv. 158 = Selinum striatum.
tenuifolium, Ram. in DC. Fl. Fr. iv. 309 (= *Wallrothia tenuifolia.*).—Gallia.
tenuifolium, Salisb. Prod. 164 = Physospermum aquilegifolium.
tenuifolium, *S. Wats. in Proc. Am. Acad.* xvi. (1879) 293.—Am. bor. occ.
ternatum, Willd. ex Schult. Syst. vi. 555 = Apium montanum.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 698.—Reg. Himal.
toluccense, H. B. & K. Nov. Gen. et Sp. v. 19. t. 422 = Arracacia toluccensis.
trifoliolatum, *Hook. f. Handb. New Zeal. Fl.* 97.—N. Zel.
tripartitum, *Dum. Fl. Belg.* 79.—Belg.
vaginatum, Spreng. Pugill. ii. 57 = Pleurospermum austriacum.

LIGUSTRIDIUM, Spach, Hist. Vég. Phan. viii. 271 (1839) = **Ligustrum**, Tourn. (Oleac.).
japonicum, Spach, l. c. 272 = Ligustrum japonicum.

LIGUSTRINA, Rupr. Beitr. Pfl. Russ. Reich. xi. 55 (1859) = **Syringa**, Linn. (Oleac.).
amurensis, Rupr. l. c. 72; Dec. Pl. Amur. t. 9 = S. amurensis.

LIGUSTRUM, [Tourn.] Linn. Syst. ed. I (1735).
OLEACEAE, Benth. & Hook. f. ii. 679.
FAULIA, Rafin. Fl. Tellur. ii. 84 (1836).
LIGUSTRIDIUM, Spach, Hist. Vég. Phan. viii. 271 (1839).
PHLYARODOXA, S. Moore, in Journ. Bot. xiii. (1875) 229.
VISIANIA, DC. Prod. viii. 289 (1844).
album, Gueldenst. ex Ledeb. Fl. Ross. iii. 39 = vulgare.
amurense, *Carr. in Rev. Hortic.* (1861) 352.—Reg. Amur.
amurense, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 11 = Lippia ligustrifolia.
angustifolium, Gilib. Exercit. i. 2; Fl. Lituan. i. 2 = vulgare.
australianum, *F. Muell. Fragm.* v. 20.—Austral.
brachystachyum, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 34.—China.
bracteolatum, D. Don, Prod. Fl. Nep. 107 = nepalense.
bracteolatum, Hook. [f.] ex Decne. in Fl. des Serres, xxii. (1877) 10 = robustum.
caeruleum, Descourt. ex Decne. l. c. 11 (Lippiae sp.).—Ind. occ.
californicum, Hort. ex Decne. l. c. 5 = ovalifolium.
Calleryanum, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 35.—China.
Candolleanum, Blume, Mus. Bot. Lugd. Bat. i. 315 = Roxburghii.
Candolleanum, Decne. in Nouv. Arch. Mus. Par. Sér. II. ii. (1879) 28 = neilgherrense.
ceylanicum, Decne. l. c. 30 = Walkeri.
chinense, Carr. in Rev. Hortic. (1863) 88 = Stauntoni.
ciliatum, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 312 = L. Ibota.
compactum, Hook. f. & Thoms. ex Decne. in Fl. des Serres, xxii. (1877) 10 = Roxburghii, *C. B. Clarke.*

LIGUSTRUM:—
confusum, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 24.—Ind. or.
coriaceum, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = Olea Roxburghii.
cotinifolium, Jacques, Man. Pl. iii. 3 = Linociera cotinifolia.
Cumingianum, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 28.—Ind. or.
Decaisnei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 616.—Ind. or.
glabrum, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = japonicum.
glomeratum, *Blume, Mus. Bot. Lugd. Bat.* i. 314.—Java.
Hookeri, Decne. in Fl. des Serres, xxii. (1877) 10 = Wallichii.
Ibota, *Siebold, in Verh. Batav. Gen.* xii. (1830) 36.—Japon.
insulare, Decne. in Nouv. Arch. Mus. Par. Sér. II. ii. (1879) 24 = seq.
insulense, *Hort. ex Decne. in Fl. des Serres*, xxii. (1877) 10.—Hab.?
italicum, Mill. Gard. Dict. ed. VIII. n. 2 = vulgare.
japonicum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 107 = nepalense.
japonicum, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 8 = lucidum.
Kellerianum, Vis. Revis. Pl. Min. Cogn. Hort. Patav. (1855) 7 = japonicum.
kumaonense, Decne. in Nouv. Arch. Mus. Par. Sér. II. ii. (1879) 28 = nepalense.
lanceolatum, Herb. Lamb. ex D. Don, Prod. Fl. Nep. 107 = nepalense.
lancifolium, Carr. ex Decne. in Fl. des Serres, xxii. (1877) 10 = compactum.
latifolium, Vitm. Summa Pl. i. 21 = japonicum.
laurifolium, Roxb. Hort. Beng. 3, nomen = Myxopyrum smilacifolium.
longifolium, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 8, 10 = Massalongianum, compactum.
lucidum, Ait. Hort. Kew. ed. II. i. 19 = Olea Roxburghii.
lucidum, Buch.-Ham. ex Wall. Cat. n. 2816 C = robustum.
lucidum, Hort. ex C. Koch, Dendrol. ii. 1. 274 = ovalifolium.
macrophyllum, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 9 = japonicum.
Massalongianum, *Vis. Revis. Pl. Min. Cogn. Hort. Patav.* (1855) 8.—Reg. Himal.
medium, *Franch. & Sav. Enum. Pl. Jap.* ii. 437.—Japon.
mellosum, *Decne. in Nouv. Arch. Mus. Par.* ii. Sér. II. (1879) 22.—Tibet.
micranthum, *Zucc. in Abh. Akad. Muench.* iv. III. (1846) 168.—Japon.
microphyllum, *Bedd. Forest. Man. Bot.* 154.—Ind. or.
microphyllum, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 616 = Decaisnei.
molliculum, *Hance, in Journ. Bot.* xx. (1882) 291.—China.
multiflorum, Hort. ex Jacques, in Rev. Hortic. (1863) 339 = Lippia ligustrifolia.
Myrsinites, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 33.—Ind. or.
myrtifolium, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 8 = Massalongianum.
neilgherrense, Decne. in Nouv. Arch. Mus. Par. Sér. II. ii. (1879) 32 = Decaisnei.
neilgherrense, *Wight, Ic.* t. 1243.—Ind. or.
nepalense, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = robustum.
nepalense, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 151.—Reg. Himal.
oblongifolium, Hort. Panorm. ex Decne. in Fl. des Serres, xxii. (1877) 10 = Roxburghii.
obovatum, Decne. in Arch. Nouv. Mus. Par. Sér. II. ii. (1879) 22 = neilgherrense.
obtusifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 168 = L. Ibota.
obtusiusculum, *Blume, Mus. Bot. Lugd. Bat.* i. 314.—Java.
ovalifolium, *Hassk. Cat. Hort. Bog. Alt.* 119.—Japon.
ovalifolium, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = japonicum.

LIGUSTRUM:—
paniculatum, Blume, Mus. Bot. Lugd. Bat. i. 314 = glomeratum.
parviflorum, Vis. in Atti Ist. Venet. Sc. Ser. III. iv. (1858–59) 137 = nepalense.
Perrottetii, *A. DC. in DC. Prod.* viii. 294.—Ind. or.
pubescens, Wall. Cat. n. 2841; Pl. As. Rar. iii. 44 = robustum.
pubinerve, *Blume, Mus. Bot. Lugd. Bat.* i. 314.—Ins. Philipp.
punctatum, Griff. Notul. iv. 741 = robustum.
quadriloculare, Blanco, Fl. Filip. ed. I. 10 = Clerodendron Navesianum.
Quihoui, *Carr. in Rev. Hortic.* (1869) 377.—China.
racemosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = obtusiusculum.
retitulatum, Blume, Mus. Bot. Lugd. Bat. i. 313 = ovalifolium.
robustum, Bedd. For. Man. 154 = Roxburghii.
robustum, *Blume, Mus. Bot. Lugd. Bat.* i. 313.—Ind. or.
robustum, Thw. Enum. Pl. Zeyl. 188 = Walkeri.
rosmarinifolium, Hort. ex Gard. Chron. (1881) II. 149 = Massalongianum.
Roxburghii, Blume, Mus. Bot. Lugd. Bat. i. 315 = lucidum.
Roxburghii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 615.—Ind. or.
Roxburghii, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 6 = L. Ibota.
salicifolium, *Carr. in Rev. Hortic.* (1866) 380.—Hab.?
sempervirens, Fl. Corc. in Ionios Anthol. i. 439 = vulgare.
Sieboldi, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = japonicum.
Simonii, Carr. ex Decne. l. c. = Roxburghii.
sinense, Hort. ex Decne. l. c. 6 = L. Ibota.
sinense, *Lour. Fl. Cochinch.* 19.—China.
spicatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 107 = nepalense.
spicatum, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 8 = Massalongianum.
spicatum, Jacques, in Rev. Hortic. (1863) 339 = Lippia ligustrifolia.
Stauntoni, *DC. Prod.* viii. 294.—China.
Stauntoni, Hort. ex Decne. in Fl. des Serres, xxii. (1877) 10 = insulense.
syringaeflorum, Hort. ex Decne. l. c. 9 = japonicum.
syringaefolium, Hort. ex Decne. l. c. = japonicum.
thibeticum, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 21.—Tibet.
Tschonoskii, *Decne. l. c.* 18.—Japon.
undulatum, *Blume, Mus. Bot. Lugd. Bat.* i. 314.—N. Guinea.
Uva-ursi, Decne. in Nouv. Arch. Mus. Par. Sér. II. ii. (1879) 34 = L. Myrsinites.
vestitum, Wall. Cat. n. 6304 = nepalense.
villosum, May, in Rev. Hortic. (1874) 299; (1875) 460 = sinense.
vulgare, *Linn. Sp. Pl.* 7.—Europ.
vulgare, Thunb. Fl. Jap. 17 = ovalifolium.
Walkeri, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 27.—Zeylan.
Wallichii, Blume, Mus. Bot. Lugd. Bat. i. 315 = nepalense.
Wallichii, *Vis. in Att. Ist. Venet.* Ser. III. iv. (1858–59) 138.—Reg. Himal.

LIJNDENIA, Zoll. & Mor. Syst. Verz. Mor. 10 (1845) = **Memecylon**, Linn. (Melastom.).
dioica, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 282 = seq.
laurina, Zoll. & Mor. Syst. Verz. Zoll. 9.—Java.

LILAC, Tourn. ex Adans. Fam. ii. 223 (1763) = **Syringa**, Linn. (Oleac.).
cordato-folia, Gilib. Exercit. i. 1 = S. vulgaris.
media, Dum.-Cours. Bot. Cult. ed. II. ii. 574 = S. vulgaris.
minor, Moench, Meth. 431 = S. persica.
perpensa, Lam. Encyc. iii. 513 = Forsythia suspensa.
persica, Lam. l. c. = S. persica.
sterilis, Lavy, Etat Gén. Vég. Origin. 12 = S. persica.
suaveolens, Gilib. Fl. Lituan. i. 1 = S. vulgaris.
varina, Dum.-Cours. Bot. Cult. v. 124; ed. II. ii. 574 = S. chinensis.
vulgaris, Lam. Fl. Fr. ii. 305 = S. vulgaris.

LILAEA, Humb. & Bonpl. Pl. Aequin. i. 221. t. 63 (1808). *NAIADACEAE*, Benth. & Hook. f. iii. 1013.
HETEROSTYLIS, Hook. Fl. Bor. Am. ii. 171. t. 185 (1840).
subulata, *Humb. & Bonpl. Pl. Aequin.* i. 22 . 63.—N. Granat.

LILAVIA, Rafin. Fl. Tellur. iv. 85 (1836) = **Alstroemeria**, Linn. (Amaryll.).
psittacina, Rafin. l. c. = A. pulchella

LILENIA, Bert. in Merc. Chil. (1829); ex Bull. Férussac, xx. (1830) 108, nomen = **Azara**, Ruiz & Pav. (Bixin.).

LILIACUM, Renault, Fl. Dép. Orne, 100 (1800) = **Syringa**, Linn. (Oleac.).
album, Renault, l. c. = S. vulgaris.
laciniatum, Renault, l. c. = S. persica.
persicum, Renault, l. c. = S. persica.
rothomagense, Renault, l. c. = S. chinensis.
vulgare, Renault, l. c. = S. vulgaris.

LILIAGO, [Caesalp.] Linn. Class. Pl. 22 (1747); Presl, Bot. Bemerk. 104 (1844); Salisb. Gen. Pl. Fragm. 70 (1866) = **Anthericum**, Linn. (Liliac.).
abyssinica, Presl, l. c. = Chlorophytum tetraphyllum.
baetica, Presl, l. c. = A. Liliago.
ciliata, Presl, l. c. = A. Humboldtii.
eccremorhiza, Presl, l. c. = A. eccremorrhizum.
flavescens, Presl, l. c. = A. flavescens.
glauca, Presl, l. c. = A. glaucum.
indica, Presl, l. c. = Chlorophytum attenuatum.
japonica, Presl, l. c. = A. japonicum.
latifolia, Presl, l. c. = A. glaucum.
nepalensis, Presl, l. c. = Chlorophytum nepalense.
nivea, Presl, l. c. = Chlorophytum tuberosum.
ramosa, Presl, l. c. = A. ramosum.
scilloides, Presl, l. c. = A. scilloides.
tuberosa, Presl, l. c. = Chlorophytum tuberosum.
vulgaris, Presl, l. c. = A. Liliago.

LILIASTRUM, [Tourn.] Link, Handb. i. 173 (1829) = **Paradisea**, Mazz. (Liliac.).
album, Link, l. c. = P. Liliastrum.

LILIORHIZA, Kellogg, in Proc. Calif. Acad. ii. (1863) 46 = **Fritillaria**, Linn. (Liliac.).
lanceolata, Kellogg, l. c. = F. liliacea.

LILIUM, Tourn. ex Linn. Gen. ed. I. 91 (1737).
LILIACEAE, Benth. & Hook. f. iii. 816.
AMBLIRION, Rafin. in Am. Monthly Mag. (1818) 265.
CARDIOCRINUM, Lindl. Veg. Kingd. 205 (1847).
LIRIUM, Linn. Syst. ed. I (1735); J. G. Gmel. Fl. Sibir. i. 41 (1747).
MARTAGON, Rupp. Fl. Jen. ed. Hall. 148 (1745).
NOTHOLIRION, Wall. ex Voigt, Hort. Suburb. Calc. 654 (1845).
abchasicum, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 230 = longiflorum.
affine, Schult. f. Syst. vii. 400 = Fritillaria camtschatcensis.
albanicum, Griseb. Spicil. Fl. Rumel. ii. 385 = chalcedonicum.
albiflorum, Hook. Bot. Mag. t. 3785 = speciosum.
alpinum, *Kit. in Linnaea*, xxxii. (1863) 335 (Quid?).—Hungar.
alternans, Siebold, ex Duch. in Journ. Soc. Hort. Par. Sér. II. iv. (1870) 215, 473 = elegans.
altissimum, Hort. ex Duch. l. c. 478 = testaceum.
andinum, Nutt. in Fraser, Cat. (1813); et in Gen. Am. i. 221 = philadelphicum.
angustifolium, Mill. Gard. Dict. ed. VIII. n. 6 = pomponium.
atrosanguineum, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 629 = bulbiferum.
atrosanguineum, *Siebold, ex Loud. Hort. Brit. Suppl.* ii. 645.—Japon.
aurantiacum, Dum.-Cours. Bot. Cult. ed. II. ii. 1972 = croceum.
aurantiacum, Paxt. Mag. Bot. vi. (1839) 127 = bulbiferum.
aurantianum, Siebold, ex Loud. Hort. Brit. Suppl. ii. 645 = bulbiferum.
aurantium, Haw. ex Loud. l. c. = croceum.
auratum, *Lindl. in Gard. Chron.* (1862) 644.—Japon.

LILIUM:—
autumnale, Lodd. Bot. Cab. t. 355 = superbum.
avenaceum, Fisch. ex Regel, Gartenfl. (1865) 290. t. 485 = maculatum.
Bartramii, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1874) 232 = Washingtonianum.
Batisua, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 52 = Wallichianum.
Belladonna, *Hort. Leichtl. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 232.—Hab.?
biligulatum, Hort. ex Baker, in Journ. Hort. Soc. N. S. iv. (1877) 43 = bulbiferum.
Bloomerianum, Kellogg, in Proc. Calif. Acad. iii. (1863) 160 = Humboldtii.
Boisseri, Orphan. ex Nym. Consp. 721 = L. Martagon.
Bolanderi, *S. Wats. in Proc. Am. Acad.* xx. (1885) 377.—Calif.
Broussarti, Morr. in Van Houtte & Morr. Hortic. Belg. i. (1834) 371. t. 23 = speciosum.
Brownii, Poit. in Rev. Hortic. Sér. II. ii. (1843–44) 496 = japonicum.
bulbiferum, DC. Fl. Fr. iii. 202 = croceum.
bulbiferum, *Linn. Sp. Pl.* 302.—Europ. austr.
bulbiferum, Thunb. Fl. Jap. 134 = lancifolium.
bulbiferum, Thunb. in Trans. Linn. Soc. ii. (1794) 333 = elegans.
Buschianum, Lodd. Bot. Cab. t. 1628 = bulbiferum.
byzantinum, Hort. ex Duch. in Journ. Soc. Hort. Par. Sér. II. v. (1871) 280 = chalcedonicum.
byzantinum, Steud. Nom. ed. II. ii. 44 = candidum.
californicum, [Dombrain,] in Floral Mag. (1872), textu sub t. 33 = pardalinum.
californicum, Hort. ex Duch. in Journ. Soc. Hort. Par. Sér. II. iv. (1870) 216; v. (1871) 98 = Humboldtii.
callosum, *Sieb. & Zucc. Fl. Jap.* i. 86. t. 14.—Japon.
camschatcense, Linn. Sp. Pl. 303 = Fritillaria camtschatcensis.
camschatcense, Lour. Fl. Cochinch. 207 = elegans.
canadense, *Linn. Sp. Pl.* 303.—Am. bor.
canadense, Thunb. Fl. Jap. 135 = medeoloides?
candidum, *Linn. Sp. Pl.* 302.—Europ. austr.; Syria.
candidum, Thunb. Fl. Jap. 133 = longiflorum.
carniolicum, *Bernh. ex Mert. & Koch, Deutschl. Fl.* ii. 536; *Syn. Fl. Germ.* ed. I. 708.—Europ. austr.
carniolicum, Heldr. ex Freyn, in Flora, lxiii. (1880) 28 = Heldreichii.
carolinianum, Bosc, ex Lam. Encyc. iii. 535 = Catesbaei.
carolinianum, Michx. Fl. Bor. Am. i. 197 = superbum.
Catanii, Baker, in Journ. Hort. Soc. N. S. iv. (1877) 45 = L. Martagon.
Catesbaei, Hort. Bouch. ex Kunth, Enum. Pl. iv. 264 = dauricum.
Catesbaei, *Walt. Fl. Carol.* 123.—Am. bor.
Cattaniae, Vis. in Mem. Ist. Venet. xvi. (1871) 64 = L. Martagon.
chalcedonicum, *Linn. Sp. Pl.* 302.—Carniol.
chalcedonicum, Linn. l. c. ed. II. 434, partim = carniolicum.
colchicum, Hort. ex Planch. Fl. des Serres, Sér. I. v. (1849) 507, 509 = monadelphum.
columbianum, *Hort.—Cf. Gard. Chron.* (1871) 1257; *et Hanson, ex Baker, in Journ. Linn. Soc.* xiv. (1874) 243.—Am. bor. occ.
concolor, *Salisb. Parad. Lond.* t. 47.—China.
cordifolium, D. Don, Prod. Fl. Nep. 52 = giganteum.
cordifolium, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332.—Japon.
coridion, Siebold & De Vriese, Tuinbouw Fl. ii. (1855) 341 = concolor.
coruscans, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 240 = elegans.
croceum, *Chaix, in Vill. Hist. Pl. Dauph.* i. 322.—Europ. austr.
dalmaticum, Vis. in Mem. Ist. Venet. xvi. (1871) 64 = L. Martagon.
dauricum, *Ker-Gawl. in Bot. Mag.* sub t. 1210.—Dahuria.
Davidi, *Duch. in Elwes, Monog. Lil.* t. 24.—China.
Dexteri, Hovey, ex Duch. in Journ. Soc. Hort. Par. Sér. II. iv. (1870) 480 = auratum.
elatum, Salisb. Prod. 237 = bulbiferum.
elegans, *Thunb. in Mém. Acad. Pétersb.* iii. (1811) 203. t. 3. f. 2.—Japon.
excelsum, Hort. ex Seem. in Harting. Parad. Vindob. 2 = testaceum.
eximium, Court. ex Spae, Mém. Gen. Lis, 14 = longiflorum.

LILIUM:—
eximium, Hort. ex Kunth, Enum. Pl. iv. 259 = speciosum.
flavum, Lam. Fl. Fr. iii. 283 = pyrenaicum.
formosissimum, *Leichtl. ex Duch. in Journ. Soc. Hort. Par.* Sér. II. iv. (1870) 216, nomen.—Hab. ?
formosum, *Lem. Illustr. Hortic.* xii. (1865) t. 459.—Japon.
Fortunei, *Lindl. in Gard. Chron.* (1862) 212.—Japon.
fulgens, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 645 = atrosanguineum.
fulgens, Morr. ex Spae, Mém. Gen. Lis, 19 = bulbiferum.
giganteum, Wall. Tent. Fl. Nep. 21. t. 12 = cordifolium.
glabrum, Spreng. Syst. ii. 62 = L. Martagon.
Glehni, *F. Schmidt, Reis. Amurl.* 187.—Japon.
gracile, Ebel, Zwölf Tage auf Montenegro, 8. t. 1 = chalcedonicum.
Grayi, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 256.—Am. bor. occ.
haematochroum, Lem. Illustr. Hortic. xiv. (1867) t. 503 = bulbiferum.
Handsoni, *Elwes, Monog. Lil.* i. t. 34.—Mandshuria.
Hansoni, *Leichtl. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 245.—Japon.
Harrisii, Carr. in Rev. Hortic. (1883) 211 = longiflorum.
Heldreichii, *Freyn, in Flora*, lxiii. (1880) 28.—Graecia.
hirsutum, Mill. Gard. Dict. ed. VIII. n. 10 = L. Martagon.
Hookeri, *Baker, in Gard. Chron.* (1871) 201.—Reg. Himal.
Humboldtii, *Roezl & Leichtl. ex Duch. in Journ. Soc. Hort. Par.* Sér. II. v. (1871) 43.—Calif.
humile, Mill. Gard. Dict. ed. VIII. n. 3 = bulbiferum.
Isabellinum, Kunze, in Bot. Zeit. i. (1843) 609 = testaceum.
Jama-Juri, Siebold & De Vriesè, Tuinbouw Fl. i. (1855) 320. t. 11 = longiflorum.
Jankae, Kern. in Oestr. Bot. Zeitschr. xxvii. (1877) 402 = chalcedonicum.
japonicum, D. Don, Prod. Fl. Nep. 40. t. 29 = Wallichianum.
japonicum, *Thunb. Fl. Jap.* 133.—Japon.
Jeffersoni, *Duch. in Journ. Soc. Hort. Par.* Sér. II. iv. (1870) 216, nomen.—Hab. ?
kamtschatkaense, Hoffmgg. Verz. Pfl. 73 = Fritillaria camschatcensis.
Krameri, Hort. Teutschel, ex Hook. f. Bot. Mag. t. 6058 = japonicum.
Kramerianum, Hort. ex Flor. Mag. (1874) t. 105 = praec.
lancifolium, Hort. Bouch. ex Kunth, Enum. Pl. iv. 364 = bulbiferum.
lancifolium, Hort. ex Guill. Arch. Bot. ii. (1833) 271 = speciosum.
lancifolium, *Thunb. in Trans. Linn. Soc.* ii. (1794) 333.—Japon.
lateritium, Hort. ex Baker, in Journ. Hort. Soc. N. S. iv. (1877) 43 = bulbiferum.
latifolium, Link, Enum. Hort. Berol. i. 321 = bulbiferum.
Ledebourii, Boiss. Fl. Orient. v. 175 = monadelphum.
Leichtlinii, *Hook. f. Bot. Mag.* t. 5673.—Japon.
Leopoldi, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 233 = tigrinum.
lilacinum, *Leichtl. ex Duch. in Journ. Soc. Hort. Par.* Sér. II. iv. (1870) 216, nomen.—Hab. ?
linifolium, Hornem. Hort. Hafn. i. 326 = tenuifolium.
Lishmanni, T. Moore, Florist (1872), 260; (1873) 16 = tigrinum.
Loddigesianum, Schult. f. Syst. vii. 416 = monadelphum.
longiflorum, Hort. ex Steud. Nom. ed. II. ii. 440 = eximium.
longiflorum, *Thunb. in Trans. Linn. Soc.* ii. (1794) 333.—Japon.; China.
longiflorum, Wall. Tent. Fl. Nep. 40. t. 29 = Wallichianum.
longifolium, Griff. Ic. Pl. Asiat. t. 277 = roseum.
lucidum, Kellogg, in Proc. Calif. Acad. vi. (1874) 144 = columbianum.
maculatum, *Thunb. in Trans. Linn. Soc.* ii. (1794) 334.—Japon.
Manglesii ×, [Mast. in] Gard. Chron. (1881) II. 152.
maritimum, *Kellogg, in Proc. Calif. Acad.* vi. (1874) 140.—Calif.

LILIUM:—
Martagon, Ledeb. Fl. Ross. iv. 149, partim = avenaceum.
Martagon, *Linn. Sp. Pl.* 303.—Europ. austr.
Martagon, Walt. Fl. Carol. 123 = superbum.
martagonum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Martagon.
Maximowiczii, Regel, Ind. Sem. Hort. Petrop. 1866 Suppl. (1867) 26 = Leichtlinii.
medeoloides, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 415.—Japon.
Metzii, Steud. ex Duch. in Journ. Soc. Hort. Par. Sér. II. iv. (1870) 556 = neilgherrense.
Michauxianum, Schult. f. Syst. vii. 258 = superbum.
Michauxii, Poir. Encyc. iii. 457 = superbum.
Milleri, Schult. Obs. Bot. 67 = L. Martagon.
miniatum, Salisb. Prod. 237 = chalcedonicum.
monadelphum, *Bieb. Fl. Taur. Cauc.* i. 267.—Reg. Caucas.
nanum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 53 = nepalense.
neilgerricum, Hort. Veitch, ex Lem. Illustr. Hortic. x. (1863) t. 353 = seq.
neilgherrense, *Wight, Ic.* t. 2031.—Ind. or.
nepalense, *D. Don, in Mem. Wern. Soc.* iii. (1820) 412.—Reg. Himal.
nigrum, Siebold, in Verh. Batav. Gen. xii. (1830) 16 = Fritillaria camtschatcensis ?
nitidum, *Hort. Bull, ex Baker, in Gard. Chron.* (1880) II. 198.—Calif.
noricum, Bernh. ex Steud. Nom. ed. II. ii. 44 = chalcedonicum.
ochroleucum, Wall. ex Baker, in Journ. Linn. Soc. xiv. (1872) 231 = nepalense.
odorum, Planch. Fl. des Serres, Sér. I. ix. (1853–54) 53. t. 876 = japonicum.
oxypetalum, *Baker, in Journ. Linn. Soc.* xiv. (1874) 234.—Reg. Himal.
Packmanni, T. Moore, in Gard. Chron. (1875) II. 366, sphalm. = *Parkmanni*.
pardalinum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 12.—Calif.
pardinum, Moore, in Flor. & Pom. (1868) 121, in textu = bulbiferum.
Parkmanni ×, Sargent, ex T. Moore, in Gard. Chron. (1875) II. 366, 494, 525.
Parryi, *S. Wats. in Proc. Davenp. Acad.* ii. 188. tt. 5, 6.—Calif.
partheneion, Siebold & De Vriese, in Tuinbouw Fl. ii. (1855) 341 = concolor.
parviflorum, Hort. ex Flor. Mag. N. S. (1874) t. 136 = canadense.
parvum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 179. t. 52.—Calif.
penduliflorum, [DC. in] Red. Lil. t. 105 = canadense.
pendulum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 19 = Crinum asiaticum.
pendulum, Spae, Mém. Gen. Lis, 28 = canadense.
pensylvanicum, Ker-Gawl. in Bot. Mag. t. 872 = bulbiferum.
peregrinum, Hort. Germ. ex Lem. in Fl. des Serres, Sér. I. i. (1845) 92 = testaceum.
peregrinum, Mill. Gard. Dict. ed VIII. n. 2 = candidum.
perenne, Gueldenst. Reise durch Russl. ii. 65, sphalm. = Linum perenne.
persicum, *Fisch. ex Steud. Nom.* ed. I. 480, nomen.—Persia.
philadelphicum, *Linn. Sp. Pl.* ed. II. 435.—Am. bor.
philadelphicum, Thunb. Fl. Jap. 135 = elegans.
philippinense, *Baker, in Gard. Chron.* (1873) 1141.—Ins. Philipp.
pictum, Hort. Siebold, ex Baker, in Journ. Hort. Soc. N. S. iv. (1877) 43 = bulbiferum.
pinifolium, Leichtl. ex Duch. in Journ. Soc. Hort. Par. Sér. II. iv. (1870) 217 nomen (Quid ?).—Hab. ?
polyphyllum, *D. Don, in Royle, Illustr. Bot. Himal.* 388.—Reg. Himal.
pomponicum, Thunb. Fl. Jap. 134 = callosum.
pomponium, [DC. in] Red. Lil. t. 7 = chalcedonicum.
pomponium, *Linn. Sp. Pl.* 302.—Sibir.
pomponium, Lour. Fl. Cochinch. 207 = tigrinum.
ponticum, *C. Koch, in Linnaea*, xxii. (1849) 234.—As. Min.
praecox, Lodd. ex Steud. Nom. ed. II. ii. 44 (Quid ?).—Hab. ?
Pseudo-tigrinum, Carr. in Rev. Hortic. (1867) 411 = Leichtlinii.

LILIUM:—
puberulum, Hort. Leichtl. ex Baker, in Journ. Linn. Soc. xiv. (1874) 242 = pardalinum.
puberulum, *Torr. ex Duch. in Journ. Hort. Soc. Par.* Sér. II. iv. (1870) 217; v. (1871) 54.—Am. bor.
pubescens, Bernh. ex Hornem. Hort. Hafn. ii. 962 = bulbiferum.
pudicum, Pursh, Fl. Am. Sept. i. 228. t. 8 = Fritillaria pudica.
pulchellum, *Fisch. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* vi. 56.—Dahuria.
pulchrum, Salisb. Prod. 237 = canadense.
pumilum, [DC. in] Red. Lil. t. 378 = tenuifolium.
pumilum, Hort. Bouch. ex Kunth, Enum. Pl. iv. 265 = croceum.
punctatum, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 276, 277 = speciosum.
punctatum, Jacquem. ex Duch. in Journ. Hort. Soc. Par. Sér. II. iv. (1870) 545 = polyphyllum.
puniceum, Siebold & De Vriese, in Ann. Hort. Pays-Bas (1861) 23 = tenuifolium.
purpureo-croceum, Bosc, ex Steud. Nom. ed. I. 480 = bulbiferum.
purpureum, Regel, Cat. Pl. Hort. Aksakov, 83 (Quid ?).—Hab. ?
pygmaeum, *Leichtl. ex Duch. in Journ. Hort. Soc. Par.* Sér. II. iv. (1870) 218, nomen.—Hab. ?
pyrenaicum, Baumg. Enum. Stirp. Transs. i. 301 = chalcedonicum.
pyrenaicum, *Gouan, Illustr.* 25.—Mont. Pyren.
pyrenaicum, Ledeb. Fl. Ross. iv. 151 = Ledebourii.
quadrifoliatum, E. Mey. in Presl, Rel. Haenk. i. 126 = Fritillaria camtschatcensis.
recurvum, *Hort. Belv. ex Steud. Nom.* ed. I. 480, nomen.—Hab. ?
Robinsonianum, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 244 = Humboldtii.
Roezli, Regel, Gartenfl. (1870) 321. t. 667 = pardalinum.
roseum, *Wall. Cat.* n. 5077.—Ind. or.
rubescens, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 256.—Calif.
rubrum, Lam. Fl. Fr. iii. 283 = pomponium.
sanguineum, Hort. ex Lindl. Bot. Reg. (1846) t. 50 = bulbiferum.
sarniense, Hort. Angl. ex Curt. Bot. Mag. t. 294 = Nerine sarniense.
Sayii, Nutt. ex Duch. in Journ. Hort. Soc. Par. Sér. II. v. (1871) 273 = columbianum.
scabrum, Moench, Meth. 305 = bulbiferum.
sibiricum, Willd. Enum. Hort. Berol. Suppl. 17 = bulbiferum.
Sieboldi, *Leichtl. ex Duch. in Journ. Hort. Soc. Par.* Sér. II. iv. (1870) 218, nomen.—Japon.
sinense, Hort. ex Steud. Nom. ed. II. ii. 44 = tigrinum.
sinicum, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 115. t. 193 = concolor.
speciosissimum, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 645 = longiflorum.
speciosum, Andr. Bot. Rep. t. 586 = tigrinum.
speciosum, Siebold, in Verh. Batav. Gen. xii. (1830) 15 = longiflorum.
speciosum, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332.—Japon.
spectabile, Link, Enum. Hort. Berol. i. 321 = dauricum.
spectabile, Salisb. Ic. Stirp. Rar. 9 = Catesbaei.
splendens, Baker, in Journ. Hort. Soc. N. S. iv. (1877) 41 = tigrinum.
stenophyllum, Baker, in Journ. Linn. Soc. xiv. (1874) 251 = tenuifolium.
striatum, Baker, in Journ. Hort. Soc. N. S. iv. (1877) 41 = candidum.
superbum, *Linn. Sp. Pl.* ed. II. 434.—Am. bor.
superbum, Thunb. Fl. Jap. 134 = speciosum.
Szovitsianum, Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 58 = monadelphum.
Takesima, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 230 = longiflorum.
tenuifolium, *Fisch. Hort. Gorenk.* ed. II. (1812) 8; *Schrank, Pl. Rar. Hort. Monac.* t. 91.—Davuria.
testaceum, *Lindl. Bot. Reg.* (1842) *Misc.* 51.—Japon.
Thomsonianum, Lindl. l. c. (1845) t. 1 = roseum.
Thunbergianum, Schult. Syst. vii. 415 = elegans.
tigrinum, *Ker-Gawl. in Bot. Mag.* t. 1237.—China; Japon.
triceps, Klotzsch, in Bot. Ergeb. Waldem. Reise, 53 = nepalense.

LILIUM:—
tricolor, Leichtl. ex Duch. in Journ. Hort. Soc. Par. Sér. II. iv. (1870) 219, nomen.—Hab.?
tubiflorum, Wight, Ic. t. 2033 = neilgherrense.
umbellatum, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 239 = dauricum.
umbellatum, Pursh, Fl. Am. Sept. i. 229 = philadelphicum.
venustum, Hort. Berol. (1841); ex Kunth, Enum. Pl. iv. 265 = bulbiferum.
versicolor, Salisb. Prod. 237 = L. Martagon.
verticillatum, Gilib. Exercit. ii. 466 = L. Martagon.
verticillatum, Willd. ex Kunth, Enum. Pl. iv. 263 = philadelphicum.
vestale, Mast. in Gard. Chron. (1872) 1522 = speciosum.
Wallichianum, *Schult. f. Syst.* vii. 1689.—Reg. Himal.
Wallichianum, Wight, Ic. t. 2035 = neilgherrense.
wansharicum, Hort. ex Duch. in Journ. Hortic. Soc. Par. Sér. II. v. (1871) 49 = philadelphicum.
Washingtonianum, [Dombrain,] in Floral Mag. (1872) t. 33 = pardalinum.
Washingtonianum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 13.—Calif.
Wilsoni, Hort. ex T. Moore, in Flor. et Pomol. (1868) 192 = bulbiferum.
Wittei, Suring. in C. Koch, Wochenschr. x. (1867) 294 = auratum.

LILIUM-CONVALLIUM, Tourn. ex Moench, Meth. 636 (1794) = **Convallaria**, Linn. (Liliac.).
majale, Moench, l. c. = C. majalis.

LILLIUM, Hill, Hort. Kew. 354 (1768) = **Lilium**, Tourn.

LIMACIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 383 (1818) = **Xylosma**, Forst. f. (Bixin.).
coriacea, Steud. Nom. ed. I. 480 = X. nitidum.
laurifolia, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 384 = X. nitidum.

LIMACIA, Lour. Fl. Cochinch. 620 (1790). *MENISPERMACEAE*, Benth. & Hook. f. i. 36.
HYPSERPA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40.
Amherstiana, Miers, Contrib. iii. 112 = triandra.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 83.—Borneo.
cerasifera, *Becc. Malesia*, i. (1877) 150.—Borneo.
cuspidata, *Hook. f. & Thoms. Fl. Ind.* i. 189.—Ind. or.; Malaya.
distincta, Miers, Contrib. iii. 111 = velutina.
inornata, Miers, l. c. = velutina.
longifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 367 = Cocculus longifolius.
microphylla, *Miq. Fl. Ind. Bat. Suppl.* 386.—Sumatra.
oblonga, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 43.—Malaya.
scandens, *Lour. Fl. Cochinch.* 621.—Cochinch.
sumatrana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 398.—Sumatra.
triandra, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 43.—Burma; Malaya.
velutina, *Miers, l. c.*—Malaya.
Wallichiana, Miers, Contrib. iii. 113 = triandra.

LIMATODES, Lindl. Gen. et Sp. Orch. 252 (1833) = seq.

LIMATODIS, Blume, Bijdr. 375 (1825) = **Phaius**, Lour. (Orchid.).
gracilis, Lindl. Fol. Orch. Limat. 1 = Calanthe gracilis.
grata, Miq. Fl. Ind. Bat. iii. 672 = P. gratus.
labrosa, Reichb. f. in Gard. Chron. (1879) i. 202 = Calanthe labrosa.
mishmensis, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 36 = P. mishmensis.
pauciflora, Blume, Bijdr. 375. t. 62 = P. pauciflorus.
punctata, Lindl. Fol. Orch. Limat. 2 = P. pauciflorus.
rosea, Lindl. in Lindl. & Paxt. Flow. Gard. iii. (1852–53) 35. t. 81 = Calanthe rosea.

LIMBARDA, Adans. Fam. ii. 125 (1763) = **Inula**, Linn. (Compos.).
arabica, Rafin. Fl. Tellur. ii. 48 = Pulicaria arabica.
crithmifolia, Rafin. l. c. = I. crithmifolia.
crithmoides, Dum. Fl. Belg. 68 = I. crithmoides.

LIMBARDA:—
japonica, Rafin. Fl. Tellur. ii. 48 = Inula japonica.
pinifolia, Rafin. l. c. = Senecio pinifolius.
tricuspis, Cass. in Dict. Sc. Nat. xxvi. 438 = I. crithmoides.

LIMBORCHIA, Scop. Introd. 139 (1777) = **Coutoubea**, Aubl. (Gentian.).

LIMEUM, Linn. Syst. ed. X. 995 (1759). *FICOIDEAE*, Benth. & Hook. f. i. 859.
ACANTHOCARPEA, Klotzsch, in Peters, Reise Mossamb. Bot. 137. t. 24 (1861).
DICARPAEA, Presl, Symb. Bot. i. 37. t. 26 (1830).
GAUDINIA, J. Gay, in Bull. Férussac, xviii. (1829) 412.
LINSCOTIA, Adans. Fam. ii. 269 (1763).
aethiopicum, *Burm. f. Fl. Cap. Prod.* 11.—Afr. austr.
africanum, *Linn. Syst.* ed. X. 995.—Afr. austr.
africanum, Sieber, ex Moq. in DC. Prod. xiii. ii. 23 = viscosum.
africanum, Zeyh. ex Moq. l. c. 22 = capense.
aphyllum, Linn. f. Suppl. 214 = capense.
argute-carinatum, Wawra, in Sitzb. Acad. Wien, Math.-Nat. xxxviii. (1860) 563 = linifolium.
canescens, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* i. (1836) 342.—Afr. austr.
capense, *Thunb. Prod. Pl. Cap.* 68.—Afr. austr.
fluviale, Eckl. & Zeyh. Enum. 287 = aethiopicum.
glomeratum, *Eckl. & Zeyh. l. c.* 288.—Afr. austr.
humile, Forsk. Fl. Aegypt. Arab. 79 = Andrachne telephioides.
indicum, *Stocks, ex T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 30.—Ind. or.; Afr. trop.
linifolium, *Fenzl, in Ann. Wien. Mus.* i. (1836) 342.—Afr. trop.
litorale, Eckl. & Zeyh. Enum. 287 = africanum.
Meyeri, Fenzl, in Ann. Wien. Mus. i. (1836) 342 = glomeratum.
pauciflorum, *Moq. in DC. Prod.* xiii. ii. 23.—Afr. austr.
telephioides, E. Mey. ex Fenzl, Nov. Stirp. Dec. 87 = capense.
viscosum, *Fenzl, l. c.*—Afr. trop.

LIMIA, Vand. Fl. Lusit. 42. t. 3. f. 21 (1788) = **Vitex**, Linn. (Verbenac.).

LIMNANTHEMUM, S. P. Gmel. in Nov. Comm. Acad. Petrop. xiv. i. 1769 (1770) 527. t. 17. f. 2. *GENTIANEAE*, Benth. & Hook. f. ii. 819.
LIMNANTHUS, Neck. Elem. ii. 27 (1790).
NYMPHOIDES, Tourn. ex Medic. Phil. Bot. i. 35 (1789).
SCHWEYCKERTA, C. C. Gmel. Fl. Bad. i. 447 (1805).
TRACHYSPERMA, Rafin. in Med. Repos. N. York, v. (1808) 352.
VILLARSIA, J. F. Gmel. Syst. 447 (1791).
WALDSCHMIDIA, [Web. in] Wigg. Prim. Fl. Holsat. 19 (1780).
albiflorum, F. Muell. Census, 90 = Villarsia albiflora.
aurantiacum, *Dalz. in Hook. Kew Journ.* ii. (1850) 136.—Ind. or.
biflorum, Thw. Enum. Pl. Zeyl. 205 = aurantiacum.
brevipedicellatum, *Vatke, in Linnaea*, xl. (1876) 220.—Abyssin.
calthifolium, F. Muell. Fragm. ix. 164 = Villarsia calthifolia.
calycinum, Miq. Fl. Ind. Bat. ii. 564 = indicum.
capitatum, F. Muell. Fragm. ix. 164 = Villarsia capitata.
congestiflorum, F. Muell. Census, 90 = Villarsia congestiflora.
crenatum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 17.—Austral.
cristatum, *Griseb. Gen. et Sp. Gent.* 342.—Ind. or.; China.
Ecklonianum, Griseb. l. c. 346 = indicum.
exaltatum, F. Muell. Fragm. ix. 165 = Villarsia reniformis.
exiguum, *F. Muell. Fragm.* i. 40.—Tasman.
exiliflorum, *F. Muell. l. c.* v. 46.—Austral.
Forbesianum, *Griseb. Gen. et Sp. Gent.* 345.—Zeylan.
Fraserianum, Griseb. l. c. 346 = indicum.
geminatum, *Griseb. l. c.*—Austral.
Grayanum, *Griseb. Cat. Pl. Cub.* 181.—Cuba.
Gunnii, Hook. f. Fl. Tasm. ii. 368 = Liparophyllum Gunnii.

LIMNANTHEMUM:—
Humboldtianum, *Griseb. Gen. et Sp. Gent.* 347.—Am. austr.
hydrocharoides, *F. Muell. ex Benth. Fl. Austral.* iv. 380.—Austral.
Hydrophyllum, *Griseb. Gen. et Sp. Gent.* 348.—Cochinch.
indicum, Griseb. l. c. 343 = cristatum.
indicum, *Thw. Enum. Pl. Zeyl.* 205.—Amphig. trop.
Kleinianum, Griseb. Gen. et Sp. Gent. 344 = indicum.
lacunosum, *Griseb. l. c.* 347.—Am. bor.
lasiospermum, F. Muell. Fragm. ix. 165 = Villarsia lasiosperma.
latifolium, F. Muell. l. c. 164 = Villarsia latifolia.
microphyllum, *Griseb. Gen. et Sp. Gent.* 364.—Bras.
minimum, *F. Muell. Fragm.* i. 40.—Austral.
Moonii, Thw. Enum. Pl. Zeyl. 205 = parvifolium.
niloticum, *Kotschy & Peyr. Pl. Tinn.* 28. t. 9 A.—Afr. trop.
nymphoides, *Hoffmgg. & Link, Fl. Port.* i. 344.—Europ.; As. bor.
orbiculatum, Griseb. Gen. et Sp. Gent. 348 = indicum?
parnassifolium, F. Muell. Fragm. ix. 165 = Villarsia parnassifolia.
parvifolium, *Griseb. in DC. Prod.* ix. 141.—Ind. or.
peltatum, *S. P. Gmel. in Nov. Comm. Acad. Petrop.* xiv. 1769 (1770) 527. t. 17.—Europ.
peltatum, Griseb. Gen. et Sp. Gent. 348 = Brasenia Schreberi.
tenellum, Sm. ex C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 450 = parvifolium.
Thunbergianum, Griseb. Gen. et Sp. Gent. 345 = indicum.
trachyspermum, *A. Gray, Man. Bot. N. U. St.* ed. I. 363.—Am. bor.
violifolium, F. Muell. Census, 90 = Villarsia violifolia.
Wightianum, Griseb. Gen. et Sp. Gent. 344 = indicum.

LIMNANTHES, R. Br. in Lond. & Edinb. Phil. Mag. ii. (1833) 70. *GERANIACEAE*, Benth. & Hook. f. i. 274.
alba, *Hartw. ex Benth. Pl. Hartw.* 301.—Calif.
Douglasii, *R. Br. in Lond. & Edinb. Phil. Mag.* ii. (1833) 70.—Am. bor. occ.
rosea, Hartw. ex Benth. Pl. Hartw. 302 = Douglasii.
sulphurea, Loud. Encyc. Pl. 1543 = Douglasii?

LIMNANTHES, Stokes, Bot. Mat. Med. i. 300 (1812) = **Limnanthemum**, S. P. Gmel. (Gentian.).
nymphoides, Stokes, l. c. = Limnanthemum nymphoides.
peltata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 340 = Limnanthemum nymphoides.

LIMNANTHUS, Neck. Elem. ii. 27 (1790) = **Limnanthemum**, S. P. Gmel. (Gentian.).

LIMNANTHUS, Reichb. Handb. 290 (1837) = **Limnanthes**, R. Br. (Geran.).

LIMNAS, Ehrh. Beitr. iv. 146 (1789) = **Malaxis**, Sw. (Orchid.).

LIMNAS, Trin. Fund. Agrost. 116. t. 6 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1118.
arkansana, Trin. ex Steud. Nom. ed. II. ii. 45 = Thurberia arkansana.
Stelleri, *Trin. Fund. Agrost.* 116. t. 6.—Kamtschatk.

LIMNASPIDIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
parmularium, Fourr. l. c. = V. scutellaria.
scutellatum, Fourr. l. c. = V. scutellaria.

LIMNETIS, Rich. in Pers. Syn. i. 72 (1805) = **Spartina**, Schreb. (Gramin.).
cynosuroides, Rich. in Pers. l. c. = S. cynosuroides.
glabra, Nutt. Gen. Am. i. 38 = S. cynosuroides.
juncea, Rich. in Pers. Syn. i. 72 = S. juncea.
polystachya, Rich. l. c. = S. polystachya.
pungens, Rich. l. c. = S. stricta.

LIMNIA, Linn. in Act. Soc. Sc. Upsal. (1746) 130. t. 5 = **Claytonia**, Gronov. (Portul.).
alsinoides, Haw. Syn. Pl. Succ. 11 = C. sibirica.
perfoliata, Haw. l. c. 12 = C. perfoliata.
sibirica, Haw. l. c. 11 = C. sibirica.

LIMNIRION, Opiz, Seznam, 59 (1852) = **Iris**, Tourn.
Pseudacorus, Opiz, l. c. = I. Pseudacorus.
sibiricum, Opiz, l. c. = I. sibirica.

LIMNIRIS, Fuss, Fl. Transs. 636 (1866) = **Iris**, Tourn.
graminea, Fuss, l. c. 637 = I. graminea.
humilis, Fuss, l. c. = I. humilis.
Pseudo-acorus, Fuss, l. c. 636 = I. Pseudacorus.
ruthenica, Fuss, l. c. 637 = I. ruthenica.
sibirica, Fuss, l. c. = I. sibirica.
spuria, Fuss, l. c. = I. spuria.

LIMNOBIUM, Rich. in Mém. Inst. Par. (1811) 32, 66. t. 8. *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 452.
JALAMBICEA, La Llave & Lex. Nov. Veg. Desc. ii. 12 (1825).
HYDROCHARELLA, Spruce, ex Benth. & Hook. f. Gen. iii. 452, in nota (1883).
HYDROMYSTRIA, G. F. W. Mey. Prim. Fl. Esseq. 152 (1818).
TRIANEA, Karst. in Linnaea, xxviii. (1856) 424.
Bosci, *Rich. in Mém. Inst. Par.* (1811) 32. t. 8.—Am. bor.
Sinclairi, *Benth. Bot. Voy. Sulph.* 175.—Mexic.
Spongia, Steud. Nom. ed. II. ii. 45 = Bosci.
stoloniferum, *Griseb. Fl. Brit. W. Ind.* 506.—Am. trop.; Ind. occ.

LIMNOCHARIS, Humb. & Bonpl. Pl. Aequin. i. 116. t. 34 (1808). *ALISMACEAE*, Benth. & Hook. f. iii. 1009.
Commersonii, Spreng. Syst. ii. 634 = Hydrocleys Commersonii.
emarginata, *Humb. & Bonpl. Pl. Aequin.* i. 116. t. 32.—Am. trop.
flava, Buch. in Bremen, Abh. ii. (1871) 2 = emarginata.
Haenkei, *Presl, Rel. Haenk.* ii. 88.—Chili.
Humboldti, Rich. in Mém. Mus. Par. i. (1815) 369. t. 19 = Hydrocleys Commersonii.
Laforesti, Duchass. ex Griseb. in Bonplandia, vi. (1858) 11 = emarginata.
Martii, Micheli, in DC. Monog. Phan. iii. 92 = Hydrocleys Martii.
nymphoides, Micheli, l. c. 91 = Hydrocleys Commersonii.
parviflora, Micheli, l. c. 93 = Hydrocleys parviflora.
Plumieri, Rich. in Mém. Mus. Par. i. (1815) 370. t. 19. f. 2 = emarginata.

LIMNOCHARIS, Kunth, Enum. Pl. ii. 153 (1837) = **Eleocharis**, R. Br. (Cyperac.).

LIMNOCHLOA, Beauv. ex Lestib. Ess. Cyp. 41 (1819) = **Eleocharis**, R. Br. (Cyperac.).
acicularis, Reichb. in Moessl. Handb. ed. II. iii. 1808 = E. acicularis.
acutangula, Nees, in Wight, Contrib. 114 = E. fistulosa.
albibracteata, Nees, ex Kunth, Enum. Pl. ii. 143 = E. albibracteata.
alpina, Drejer, Fl. Excurs. Hafn. 18 = Eriophorum alpinum.
alpina, Reichb. Fl. Germ. Excurs. 140[11] = Scirpus pumilus.
articulata, Nees, in Linnaea, ix. (1834) 294 = E. interstincta.
Baeothryon, Reichb. in Moessl. Handb. ed. II. iii. 1808 = Scirpus pauciflorus.
caduciflora, Turcz. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. (1840) 185 = Zizania aquatica.
caespitosa, Reichb. in Moessl. Handb. ed. II. iii. 1808 = Scirpus caespitosus.
calyptrata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 244 = E. sulcata.
capensis, Krauss, in Flora, xxviii. (1845) 759 = E. limosa.
capensis, Nees, in Linnaea, x. (1835–36) 185 = E. palustris.
constricta, Nees, in Mart. Fl. Bras. ii. 1. 99 = E. geniculata.
costulata, Nees, ex Kunth, Enum. Pl. ii. 142 = E. acicularis.
crassiculmis, Nees, in Mart. Fl. Bras. ii. 1. 99 = E. geniculata.
densa, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 244 = E. geniculata.
LIMNOCHLOA :—
equisetoides, Walp. Ann. iii. 684 = Eleocharis interstincta.
fistulosa, Nees, in Linnaea, ix. (1834) 294 = E. fistulosa.
geniculata, Nees, in Mart. Fl. Bras. ii. 1. 99 = E. geniculata.
limosa, Nees, in Linnaea, ix. (1834) 294 = E. limosa.
media, Nees, in Wight, Contrib. 114 = E. fistulosa.
montevidensis, Nees, in Mart. Fl. Bras. ii. 1. 99 = E. montevidensis.
multicaulis, Reichb. Fl. Germ. Excurs. 140[11] = E. multicaulis.
multiculmis, Reichb. in Moessl. Handb. ed. II. iii. 1808 = praec.
mutata, Nees, in Mart. Fl. Bras. ii. 1. 101 = E. mutata.
obsoleta, Nees, l. c. 100 = E. interstincta.
obtusangula, Nees, in Linnaea, ix. (1834) 294 = E. obtusangula.
obtusetrigona, Lindl. & Nees, in Mart. Fl. Bras. ii. 1. 100 = E. mutata.
parvula, Reichb. in Moessl. Handb. ed. II. iii. 1808 = Scirpus nanus.
pauciflora, Peterm. Fl. Bienitz, 21 = Scirpus pauciflorus.
plantaginea, Nees, in Wight, Contrib. 114 = E. plantaginea.
quadrangulata, Nees, in Linnaea, ix. (1834) 294 = E. mutata.
spiralis, Nees, in Wight, Contrib. 114 = E. spiralis.
tetraquetra, Nees, l. c. 113 = E. tetraquetra.
truncata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 244 = E. montana.
tumida, Nees, in Wight, Contrib. 114 = E. plantaginea.

LIMNOCREPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 104 = **Crepis**, Linn. (Compos.).
paludosa, Fourr. l. c. = C. paludosa.

LIMNOGENNETON, Sch. Bip. in Walp. Rep. vi. 146 (1846–47) = **Sigesbeckia**, Linn. (Compos.).
abyssinicum, Sch. Bip. l. c. 147 = S. orientalis.

LIMNOGETON, Edgew. in Lindl. Veg. Kingd. 210 (1847) = **Aponogeton**, Thunb. (Naiad.).

LIMNONESIS, Klotzsch, in Abh. Akad. Berl. (1853) 24, 351, 352 = **Pistia**, Linn. (Aroid.).
commutata, Klotzsch, l. c. = P. Stratiotes.
Friedrichsthaliana, Klotzsch, l. c. = P. Stratiotes.

LIMNOPEUCE, Adans. Fam. ii. 80 (1763) = **Hippuris**, Linn. (Halorag.).

LIMNOPHILA, R. Br. Prod. 442 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 950.
AMBULIA, Lam. Encyc. i. 128 (1783).
CYBBANTHERA, Buch.-Ham. in D. Don, Prod. Fl. Nep. 87 (1825).
DICEROS, Lour. Fl. Cochinch. 381 (1790).
HONOTTIA, Reichb. Consp. 172 (1825).
HOTTONIA, Burm. Thes. Zeyl. 121. t. 55. f. 1 (1737).
HYDROPITYON, Gaertn. f. Fruct. iii. 19. t. 183 (1805).
LYMNOPHILA, Blume, Bijdr. 748 (1826).
TALA, Blanco, Fl. Filip. ed. I. 484 (1837); ed. II. 338 (1845).
albiflora, *Voigt, Hort. Suburb. Calc.* 502.—Ind. or.
balsamea, *Benth. in DC. Prod.* x. 386.—Burma.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 562.—Malaya.
Benthamiana, Miq. ex Hook. f. Fl. Brit. Ind. iv. 271 = racemosa.
campanuloides, Benth. in Wall. Cat. n. 3908 = Mitrasacme polymorpha.
camphorata, Hook. f. Fl. Brit. Ind. iv. 267 = diffusa.
cana, *Griff. Notul.* iv. 98.—Ind. or.
chamaedrifolia, G. Don, Gen. Syst. iv. 543 = gratissima.
conferta, *Benth. in DC. Prod.* x. 387.—Ind. or.
diffusa, *Benth. l. c.*—Ind. or.; Malaya.
diffusa, G. Don, Gen. Syst. iv. 543 = Ebermeiera thyrsoidea.
elongata, Benth. in Wall. Cat. n. 3903 = gratioloides.
erecta, *Benth. in DC. Prod.* x. 388.—Ind. or.
fragrans, Seem. Fl. Vit. 180 = serrata.
gratioloides, *R. Br. Prod.* 442.—Geront. trop.
gratissima, *Blume, Bijdr.* 749.—As. et Austral. trop.

LIMNOPHILA:—
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 271.—Malacca.
Helferi, *Hook. f. l. c.* 269.—Burma.
heterophylla, *Benth. in Wall. Cat.* n. 905; *Scroph. Ind.* 25.—As. trop.
hirsuta, *Benth. in DC. Prod.* x. 388.—As. et Austral. trop.
hypericifolia, *Benth. l. c.* 386.—Ind. or.
hyssopifolia, *Roth, Nov. Pl. Sp.* 297.—Ind. or.
javanica, *A. DC. in DC. Prod.* x. 594.—Java.
laxa, *Benth. in DC. l. c.* 388.—Burma; Zeylan.
laxa, Benth. l. c., partim = pulcherrima.
Menthastrum, Benth. l. c. 386 = Roxburghii.
micrantha, *Benth. l. c.* 387.—As. trop.
Morgania, F. Muell. Fragm. vi. 104 = Morgania floribunda, pubescens, glabra.
myriophylloides, Roth, Nov. Pl. Sp. 294 = gratioloides.
pinnatifida, *Blume, Bijdr.* 749.—Java.
polyantha, *Kurz, ex Hook. f. Fl. Brit. Ind.* iv. 268.—Burma.
polystachya, *Benth. in Wall. Cat.* n. 3909; *Scroph. Ind.* 26.—Ind. or.
pulcherrima, *Hook. f. Fl. Brit. Ind.* iv. 267.—Ind. or.; Malaya.
punctata, Blume, Bijdr. 750 = gratissima.
punctata, Hook. f. Fl. Brit. Ind. iv. 266, sphalm. = conferta.
pygmaea, *Hook. f. l. c.* 268.—Burma; Malacca.
racemosa, *Benth. in Wall. Cat.* n. 3907; *Scroph. Ind.* 26.—Ind. or.; Malaya.
reflexa, Benth. ll. cc. n. 3904; 25 = heterophylla.
repens, Benth. in DC. Prod. x. 387 = conferta.
Roxburghii, Benth. l. c. 388 = racemosa.
Roxburghii, *G. Don, Gen. Syst.* iv. 543.—As. trop.; Ins. Pacif.
serrata, *Gaudich. in Freyc. Voy. Bot.* 448. t. 57.—Malaya; Ins. Pacif.
sessiliflora, *Blume, Bijdr.* 750.—As. trop.
sessiliflora, Griff. Notul. iv. 99 = Griffithii.
tillaeoides, *Hook. f. Fl. Brit. Ind.* iv. 270.—Assam.
torenioides, Baker, in Journ. Bot. xx. (1882) 221 = Mimulus madagascariensis.
trifida, Spreng. Syst. ii. 802 = gratioloides.
villifera, *Miq. Fl. Ind. Bat.* ii. 682.—Sumatra.
villosa, *Blume, Bijdr.* 750.—Java.

LIMNOPHYLLA, Griff. Notul. iv. 97 (1854) = praec.

LIMNOPHYTON, Miq. Fl. Ind. Bat. iii. 242 (1855). *ALISMACEAE*, Benth. & Hook. f. iii. 1005.
DIPSEUDOCHORION, Buchen. in Flora, xlviii. (1865) 245.
obtusifolium, *Miq. l. c.* 243.—As. et Afr. trop.

LIMNOSERIS, 'Peterm.' ex Linnaea, xiii. (1839) Litt. 23 = LIMONOSERIS, Peterm. = **Crepis**, Linn. (Compos.).

LIMNOSIPANEA, Hook. f. in Hook. Ic. Pl. t. 1050 (1868). *RUBIACEAE*, Benth. & Hook. f. ii. 53.
palustris, *Hook. f. l. c.* sub t. 1050.—Am. trop.
Schomburgkii, *Hook. f. l. c.*—Guiana.
Spruceana, *Hook. f. l. c.* t. 1050.—Bras.

LIMNOSTACHYS, F. Muell. Fragm. i. 24 (1858) = **Monochoria**, Presl (Portul.).
cyanea, F. Muell. l. c. = M. cyanea.

LIMNOXERANTHEMUM, Salzm. ex Steud. Syn. Pl. Cyp. 281 (1855) = **Paepalanthus**, Mart. (Erioc.).
glabrum, Salzm. ex Steud. l. c. = P. gracilis.

LIMODORON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = **Limodorum**, Tourn. (Orchid.).
fasciolum, St. Lag. l. c. = Vanilla Fasciola.

LIMODORUM, [Tourn.]. Linn. Syst. ed. II. 29 (1740); Rich. in Mém. Mus. Par. iv. (1818) 50. *ORCHIDEAE*, Benth. & Hook. f. iii. 618.
abortivum, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 80.—Europ.
altum, Linn. Syst. ed. XII. 594 = Bletia verecunda.
angustifolium, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 168 = Cymbidium longifolium.
aphyllum, Roxb. Pl. Corom. i. 34. t. 41 = Dendrobium Pierardi?

LIMODORUM:—
articulatum, Schum. & Thonn. Beskr. Guin. Pl. 399 = Eulophia articulata.
autumnale, Auct. ex Steud. Nom. ed. II. ii. 46 = Earina mucronata.
autumnale, Walt. Fl. Carol. 221 = Spiranthes cernua.
barbatum, Lam. Encyc. iii. 515 = Calopogon pulchellus.
barbatum, Thunb. Prod. Pl. Cap. 4 = Eulophia barbata.
bicallosum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 30 = Liparis bituberculata?
bicolor, Roxb. Hort. Beng. 63; Fl. Ind. iii. 469 = Eulophia herbacea.
bidens, *Afzel. ex Sw. in Vet. Acad. Handl. Stockh.* (1800) 243, nomen; *Pers. Syn.* ii. 521 (Quid?).—Afr. trop.
bidentatum, *Willd. Sp. Pl.* iv. 124 (Quid?).—Ind. or.
boreale, Sw. in Schrad. Neues Journ. i. (1805) 85 = Calypso borealis.
bracteatum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 466 = Phaius albus.
brasiliense, *Nois. ex Steud. Nom.* ed. II. ii. 46, nomen.—Bras.
callosum, Blume, Bijdr. 374 = Phaius callosus.
canaliculatum, *Aubl. Pl. Gui.* ii. 821 (Quid?).—Guian.
candidum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 470 = Geodorum dilatatum?
capense, Berg. Desc. Pl. Cap. 347 = Eulophia tristis.
carinatum, Willd. Sp. Pl. iv. 124 = Eulophia virens?
ciliatum, Schum. & Thonn. Beskr. Guin. Pl. 400 = Eulophia gracilis.
clavatum, Willd. Sp. Pl. iv. 126 = Saccolabium clavatum.
complanatum, Willd. l. c. = Cleisostoma complanatum.
concolor, Thou. Orch. Iles Afr. t. 45 = Eulophia concolor.
coriaceum, Thunb. ex Sw. in Schrad. Journ. ii. (1799) 234 = Aërides coriaceum.
crispum, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 180 = Phaius crispus.
cristatum, Afzel. ex Sw. in Vet. Acad. Handl. Stockh. (1800) 243, nomen; Pers. Syn. ii. 521 = Eulophia cristata.
cucullatum, Afzel. ex Sw. l. c.; et Pers. l. c. = Polystachya grandiflora.
cyperifolium, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 163 = Cyperorchis elegans.
Dabia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 30 = = Eulophia campestris.
densiflorum, Lam. Encyc. iii. 516 = Geodorum dilatatum.
dipterum, Tausch, ex Steud. Nom. ed. II. ii. 46 = Cryptopus elatus.
diurnum, Jacq. Coll. iv. 107 = Cymbidium diurnum.
dubium, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 125 = Eulophia campestris.
eburneum, Willd. Sp. Pl. iv. 125 = Angraecum eburneum.
emarginatum, *Afzel. ex Sw. in Vet. Acad. Handl. Stockh.* (1800) 243, nomen; *Pers. Syn.* ii. 521 (Quid?).—Afr. trop.
ensatum, Thunb. Fl. Jap. 29 = Cymbidium ensifolium.
epidendroides, Willd. Sp. Pl. iv. 124 = Eulophia virens.
Epipogium, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 80 = Epipogum aphyllum.
falcatum, Thunb. in Trans. Linn. Soc. ii. (1794) 326; Sw. in Schrad. Journ. ii. (1799) 231 = Angraecum falcatum.
Fasciola, Sw. l. c. = Vanilla Fasciola.
filiforme, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 80 = Aëranthes filiformis.
flabellatum, Thou. Orch. Iles Afr. tt. 39, 40 = Cymbidium flabellatum.
flavum, Blume, Bijdr. 375 = Phaius flavus.
flexuosum, *Willd. Sp. Pl.* iv. 128 (Quid?).—Ind. occ.
floridum, Salisb. Prod. 9 = Bletia florida.
Flos-aëris, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 80 = Arachnanthe moschifera.
funale, Sw. l. c. 79 = Dendrophylax funalis.
gentianoides, Sw. Prod. Veg. Ind. Occ. 119 = Pogonia gentianoides.
giganteum, Thunb. Prod. Pl. Cap. 4 (= *Cyrtopera gigantea*).—Afr. austr.
grandiflorum, Aubl. Pl. Gui. ii. 818 = Cymbidium grandiflorum.
hians, Thunb. Prod. Pl. Cap. 3 = Eulophia hians.
hyacinthinum, Donn, Hort. Cantab. ed. IV. 201 = Bletia hyacinthina.

LIMODORUM:—
imbricatum, *Sw. in Schrad. Neues Journ.* i. (1805) 87 (Quid?).—Afr. trop.
Incarvillei, Blume, Bijdr. 374 = Phaius Wallichii.
Incarvillei, Pers. Syn. ii. 520 = Phaius grandifolius.
lanceolatum, Aubl. Pl. Gui. ii. 821 (= *Stenorrhynchus lanceolatus*).—Guian.
latifolium, Thunb. ex Sw. in Schrad. Journ. ii. (1799) 234 = Aërides odoratum.
laxiflorum, *Lam. Encyc.* iii. 516 (Quid?).—Ind. or.
longicornu, Sw. in Schrad. Journ. ii. (1799) 230 = Mystacidium filicorne.
longifolium, Buch.-Ham. ex Lindl. Gen. et Sp. Orch. 163 = Cymbidium giganteum.
longifolium, Roxb. Hort. Beng. 63; Fl. Ind. iii. 468 = Cymbidium cyperifolium.
luridum, Pers. Syn. ii. 521 = Eulophia lurida.
luridum, Sw. in Schrad. Neues Journ. i. (1805) 87 = Angraecum luridum.
luteum, Lam. Encyc. iii. 516 = Chloraea crispa.
maculatum, Reinw. ex Lindl. in Journ. Linn. Soc. iii. (1859) 23 = Nephelaphyllum pulchrum.
monile, Thunb. in Trans. Linn. Soc. ii. (1794) 327 = Dendrobium catenatum.
nutans, Roxb. Pl. Corom, i. 33. t. 40 = Geodorum purpureum.
orchideum, Willd. Sp. Pl. iv. 126 (= *Ceratochilus orchideus*).—Ind. or.
pendulum, Aubl. Pl. Gui. ii. 819 = Dichaea echinocarpa.
Piquichen, Lam. Encyc. iii. 516 = Chloraea virescens.
plantagineum, Thou. Orch. Isles Afr. tt. 41, 42 = Cyrtopodium plantagineum.
plicatum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 465 = Phaius callosus.
praecox, Walt. Fl. Carol. 221 = Spiranthes graminea.
pulchellum, Salisb. Prod. 8 = Calopogon pulchellus.
pulchrum, Thou. Orch. Iles Afr. t. 43 = Eulophia pulchra.
purpureum, Lam. Encyc. iii. 515 = Bletia verecunda.
purpureum, Lour. ex Steud. Nom. ed. II. ii. 46 = Epidendrum atropurpureum.
pusillum, Willd. Sp. Pl. iv. 126 (Sarcochili sp.).—Ind. or.
ramentaceum, Roxb. Hort. Beng. 63; Fl. Ind. iii. 467 = Eulophia campestris.
recurvum, Roxb. Pl. Corom. i. tt. 33, 39 = Geodorum dilatatum.
retusum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 80 = Rhynchostylis retusa.
roseum, D. Don, Prod. Fl. Nep. 30 = Epipogum nutans.
scriptum, Thou. Orch. Iles Afr. tt. 46, 47 = Eulophia scripta.
spathulatum, Willd. Sp. Pl. iv. 125 = Vanda spathulata.
spectabile, Salisb. Prod. 9 = Phaius grandifolius.
sphaerochilos, Spreng. Syst. iv. Cur. Post. 306 = abortivum.
sphaerolabium, Viv. App. Fl. Cors. Prod. 6 = abortivum.
striatum, Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 79 = Calanthe striata.
striatum, Thunb. Fl. Jap. 28 = Bletia hyacinthina.
suaveolens, Herb. Reinw. ex Blume, Rumphia, iv. 49 = Vanda tricolor.
subulatum, Willd. Sp. Pl. iv. 126 = Aërides cylindricum.
Tancarvilleae, Banks, in L'Hérit. Sert. Angl. 28 = seq.
Tankervilliae, Ait. Hort. Kew. ed. I. iii. 302 = Phaius grandifolius.
Tankervilliae, Roxb. Fl. Ind. iii. 466 = Phaius Wallichii.
tetragonum, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 41 = Phaius tetragonus.
trifidum, Michx. Fl. Bor. Am. ii. 159 = Bletia verecunda.
triste, Thunb. Prod. Pl. Cap. 4 = Eulophia tristis.
tuberculosum, Thou. Orch. Iles Afr. t. 31 = Phaius tuberculosus.
tuberosum, Jacq. Coll. iv. 108 = Bletia verecunda.
tuberosum, Linn. Sp. Pl. 950 = Calopogon pulchellus.
unguiculatum, Labill. Sert. Austro-Caled. 19. t. 25 = Calanthe striata.
unifolium, Muhl. ex Nutt. Gen. Am. ii. 195 = Tipularia discolor.
utriculatum, Sw. Prod. Veg. Ind. Occ. 119 = Govenia utriculata.
variegatum, Lam. Encyc. iii. 516 = Eulophia virens.
venosum, Lam. l. c. = Chloraea Pavonii.
ventricosum, Steud. Nom. ed. I. 481 = Calanthe veratrifolia.

LIMODORUM:—
veratrifolium, Willd. Sp. Pl. iv. 122 = Calanthe veratrifolia.
verecundum, Salisb. Prod. 9 = Bletia verecunda.
villosum, Thou. Orch. Iles Afr. t. 32 = Phaius villosus.
virens, Roxb. Pl. Corom. i. 32 = Eulophia virens.

LIMON, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Citrus**, Linn. (Rutac.).
racemosum, Mill. l. c. ed. VIII. n. 3 = C. Medica.
spinosum, Mill. l. c. n. 2 = C. Medica.
vulgare, Mill. l. c. n. 1 = C. Medica.

LIMONAETES, Ehrh. Beitr. iv. 148 (1789) = **Carex**, Dill. (Cyperac.).

LIMONANTHUS, Kunth, Enum. Pl. iv. 194, in adnot. (1843) = LEIMANTHIUM, Willd. = **Melanthium**, Linn. (Liliac.).

LIMONIA, Gaertn. Fruct. i. 278. t. 58. f. 4 (1789) = **Scolopia**, Schreb. (Bixin.).
pusilla, Gaertn. l. c. 279 = S. Gaertneri.

LIMONIA, Linn. Sp. Pl. ed. II. 554 (1763). *RUTACEAE*, Benth. & Hook. f. i. 303.
HESPERETHUSA, M. Roem. Syn. Hesper. 31, 38 (1846).
WINTERLIA, Dennst. Schluess. Hort. Malab. 27 (1818).
acidissima, *Linn. Sp. Pl.* ed. II. 554.—Ind. or.
alata, *Wall. Cat.* n. 6363; *Wight & Arn. Prod.* 92.—Ind. or.
alternans, *Wall. ex Voigt, Hort. Suburb. Calc.* 139.—Burma.
alternifolia, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 64 = alternans.
ambigua, DC. Prod. i. 536 = acidissima.
angulata, *Wight & Arn. Prod.* 91.—Ins. Amboina.
angustifolia, Wall. Cat. n. 6360 = Glycosmis pentaphylla.
arborea, Roxb. Pl. Corom. i. 60. t. 85 = Glycosmis pentaphylla.
aurantiifolia, *M. Roem. Syn. Hesper.* 39.—Guian.
australis, A. Cunn. ex G. Don, in Sweet, Hort. Brit. ed. III. 91 = Citrus australis.
bilocularis, Roxb. Hort. Beng. 32; Fl. Ind. ii. 377 = Atalantia buxifolia.
caudata, Wall. ex Voigt, Hort. Suburb. Calc. 139, nomen; Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 25 = Atalantia caudata.
citrifolia, Moon, Cat. Pl. Ceyl. 34 = Atalantia ceylanica.
citrifolia, Roxb. Hort. Beng. 32; Fl. Ind. ii. 379 = Paramignya citrifolia.
citrifolia, *Salisb. Prod.* 320.—Ind. or.
citrifolia, Willd. Enum. Hort. Berol. 448 = Glycosmis pentaphylla.
cochinchinensis, *Baill. Hist. Pl.* iv. 398.—Cochinch.
corymbosa, *Blanco, Fl. Filip.* ed. II. 251.—Ins. Philipp.
crenulata, Roxb. Pl. Corom. i. 60. t. 86 = acidissima.
curvispina, *Miq. Fl. Ind. Bat. Suppl.* 500.—Sumatra.
diacantha, DC. Prod. i. 536 = Triphasia Aurantiola.
diphylla, *M. Roem. Syn. Hesper.* 39.—Java.
disticha, Blanco, Fl. Filip. ed. I. 356 = Atalantia racemosa.
dubia, *Blume, Bijdr.* 133.—Java.
dulcis, *J. F. Gmel. Syst.* 684.—N. Guinea.
flavicans, *Wall. ex Hook. f. Fl. Brit. Ind.* i. 508.—Burma.
glutinosa, Blanco, Fl. Filip. ed. I. 358 = Aegle decandra.
grandiflora, Wall. Cat. n. 6361 = Paramignya grandiflora.
grata, *Wall. ex Voigt, Hort. Suburb. Calc.* 139, nomen.—Hab.?
horrida, *Blume, Bijdr.* 133.—Java.
Laureola, *Blanco, Fl. Filip.* ed. II. 251.—Ins. Philipp.
Laureola, DC. Prod. i. 536 = Skimmia Laureola.
leptostachya, Jack, ex Hook. f. Fl. Brit. Ind. i. 492 = Zanthoxylum ovalifolium.
linearis, Blanco, Fl. Filip. ed. I. 357 = Atalantia monophylla.
lucida, *Forst. f. Prod.* 33.—N. Hebrid.
madagascariensis, *Lam. Encyc.* iii. 517.—Madag.
malliculensis, Forst. ex Steud. Nom. ed. II. ii. 47 = Murraya exotica.
mauritiana, *Lam. Encyc.* iii. 517.—Ins. Maurit.
microphylla, *Voigt, in Syll. Ratisb.* ii. (1828) 53.—Hab.?

LIMONIA:—
minuta, Forst. f. Prod. 33 = Micromelum pubescens.
missionis, Wall. Cat. n. 6358 = Atalantia missionis, racemosa, Paramignya Griffithii.
mollis, Wall. Cat. n. 6362 = Clausena pentaphylla.
monadelpha, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 217.—Afr. trop.
monophylla, Linn. Mant. 237 = Atalantia monophylla.
monophylla, Lour. Fl. Cochinch. 271 = Atalantia Hindsii.
oblonga, Wall. Cat. n. 6359 = Paramignya Griffithii.
octandra, Herb. Heyne, ex Wall. Cat. n. 6353 = Atalantia monophylla.
oligandra, Dalz. in Hook. Kew Journ. ii. (1850) 258 = Toddalia aculeata.
parviflora, Sims, Bot. Mag. t. 2416 = Glycosmis pentaphylla.
pentagyna, Roxb. Fl. Ind. ii. 382 = Bursera serrata.
pentaphylla, Lam. ex DC. Prod. i. 538 = Clausena pentaphylla.
pentaphylla, Retz. Obs. v. 24 = Glycosmis pentaphylla.
pubescens, *Wall. Cat.* n. 6365.—Burma.
retusa, D. Don, Prod. Fl. Nep. 224 = Triphasia Aurantiola?
scandens, Roxb. Hort. Beng. 32; Fl. Ind. ii. 380 = Luvunga scandens.
spectabilis, *Miq. Fl. Ind. Bat. Suppl.* 500.—Sumatra.
spinosa, Spreng. Syst. iv. Cur. Post. 162 (= *Sclerostylis spinosa*).—Java.
torulosa, Dennst. Schluess. Hort. Malab. 30 = Randia virosa.
trichocarpa, *Hance, in Journ. Bot.* xx. (1882) 258.—China.
trifolia, Burm. f. Fl. Ind. 103. t. 35. f. 1 = Triphasia Aurantiola.
trifoliata, Linn. Mant. ii. 237 = Triphasia Aurantiola.
Winterlia, *Steud. Nom.* ed. II. ii. 47.—Ind. or.

LIMONIASTRUM, Moench, Meth. 423 (1794).
PLUMBAGINEAE, Benth. & Hook. f. ii. 627.
BUBANIA, Girard, in Mém. Acad. Montp. (Sect. Sc.) i. (1848) 182.
articulatum, *Moench, Meth.* 423.—Reg. Mediterr.
Guyonianum, *Dur. ex Boiss. in DC. Prod.* xii. 689.—Alger.
monopetalum, Boiss. l. c. = articulatum.
ouarglense, *Pomel, Nouv. Mat. Fl. Atl.* 126.—Alger.
speciosum, Moench, Meth. Suppl. 149 = Statice speciosa.

LIMONIUM, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Statice**, Linn. (Plumbag.).
africanum, Mill. Gard. Dict. ed. VIII. n. 8 = S. Thouini.
bellidifolium, Dum. Fl. Belg. 27 = S. bellidifolia.
commune, S. F. Gray, Nat. Arr. Brit. Pl. ii. 296, 732 = S. Limonium.
confusum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 141 = S. confusa.
cordatum, Mill. Gard. Dict. ed. VIII. n. 10 = S. cordata.
duriusculum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 141 = S. duriuscula.
echioides, Mill. Gard. Dict. ed. VIII. n. 11 = S. echioides.
fruticosum, Mill. l. c. n. 12 = S. cylindrifolia.
Girardianum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 141 = S. Girardiana.
humile, Mill. Gard. Dict. ed. VIII. n. 4 = S. Limonium.
latifolium, Moench, Meth. 623 = S. tatarica.
minutum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 141 = S. minuta.
monopetalum, Hill, Hort. Kew. 183 = Limoniastrum articulatum.
narbonense, Mill. Gard. Dict. ed. VIII. n. 2 = S. Limonium.
oleaefolium, Mill. l. c. n. 3 = S. oleaefolia.
reticulatum, Mill. l. c. n. 9 = S. reticulata.
siculum, Mill. l. c. n. 7 = Limoniastrum articulatum.
sinuatum, Mill. l. c. n. 6 = S. sinuata.
tataricum, Mill. l. c. n. 5 = S. tatarica.
virgatum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 141 = S. virgata.
vulgare, Mill. Gard. Dict. ed. VIII. n. 1; Hill, Hort. Kew. 183 = S. Limonium.

LIMONOSERIS, Peterm. Fl. Lips. 589 (1838) = **Crepis**, Linn. (Compos.).
biennis, Peterm. l. c. = C. biennis.

LIMOSELLA, Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 958.
DANUBIUNCULUS, Sailer, ex Walp. Ann. i. 534 (1849).
MUTAFINIA, Rafin. Fl. Tellur. iv. 57 (1836).
PLANTAGINELLA, Dill. ex Moench, Meth. 427 (1794).
YGRAMELA, Rafin. Atl. Journ. 199 (1833).
annua, Lindern. Hort. Alsat. 156. t. 5 = aquatica.
aquatica, *Linn. Sp. Pl.* 631.—Reg. temp. bor. et austr.
australis, R. Br. Prod. 443 = aquatica.
borealis, Less. ex Ledeb. Fl. Ross. iii. 226 = aquatica.
calycina, Forsk. Fl. Aegypt. Arab. 112 = Herpestis Monnieria.
capensis, Benth. in DC. Prod. x. 427 = grandiflora.
capensis, *Thunb. Prod. Pl. Cap.* 104.—Afr. austr.
coerulea, Burch. Trav. i. 259 = capensis.
Curdieana, *F. Muell. Fragm.* ix. 166.—Austral.
diandra, Krock. Fl. Siles. ii. 1. 406 = aquatica.
diandra, Linn. Mant. i. 252 = Glossostigma spathulatum.
Drummondi, F. Muell. Fragm. vi. 104 = Glossostigma elatinoides, spathulatum, Drummondii.
grandiflora, *Benth. in DC. Prod.* x. 427.—Afr. austr.
maritima, *Rafin. Atl. Journ.* 199 (Quid?).—Am. bor.
plantaginis-folia, Gilib. Fl. Lituan. i. 122 = aquatica.
subulata, Ives, in 'Med. Phys. N. York, 440;' ex eod. in Am. Journ. Sc. i. (1819) 74 = aquatica.
tenuifolia, Wolf. ex Hoffm. Deutschl. Fl. ed. II. i. 11. 29 = aquatica.

LIMOSTELLA, Ledeb. Fl. Ross. iii. 226 (1849), sphalm. = praec.

LINAGROSTIS, [Tourn.] Michx. ex Scop. Fl. Carn. ed. II. i. 47 (1772) = **Eriophorum**, Linn. (Cyperac.).
alpina, Scop. l. c. 48 = E. alpinum.
paniculata, Lam. Fl. Fr. iii. 555 = E. latifolium.
polystachia, Scop. Fl. Carn. ed. II. i. 48 = E. latifolium.
vaginata, Scop. l. c. 47 = E. vaginatum.

LINANTHUS, Benth. in Bot. Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polemon.).
dichotomus, Benth. l. c. = G. dichotoma.

LINARIA, Tourn. ex Mill. Gard. Dict. ed. VI (1752).
SCROPHULARINEAE, Benth. & Hook. f. ii. 932.
CHAENARRHINUM, Reichb. Consp. 123 (1828).
CYMBALARIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 230.
ELATINE, Rupp. Fl. Jen. ed. Hall. 242 (1742).
KICKXIA, Dum. Fl. Belg. 35 (1827).
TERMONTIS, Rafin. Chl. Aetn. 5 (1813).
Acerbiana, *Boiss. Fl. Orient.* iv. 366.—Aegypt.
acutangula, *Tenore, Syll. Fl. Nap.* 302.—Ital.
acutifolia, *Fisch. ex Sweet, Hort. Brit.* ed. II. 391, nomen.—Oriens.
acutiloba, Fisch. ex Reichb. Pl. Crit. v. 14. t. 424 = vulgaris.
aegyptiaca, *Dum.-Cours. Bot. Cult.* ed. I. ii. 92.—Aegypt.; Syria.
aequitriloba, *Spreng. Syst.* ii. 790.—Corsic.
aeruginea, Losc. & Pard. Ser. Inconf. 76 = tristis.
agglutinans, *Pomel, Nouv. Mat. Fl. Atl.* 96.—Alger.
alba, Moench, Meth. 524 = chalepensis.
albifrons, *Spreng. Syst.* ii. 793.—Ins. Rhodus; Afr. bor.; Oriens.
algarbiensis, Welw. ex Nym. Consp. 538 = hirta.
algarviana, *Chav. Monog. Antirrh.* 142.—Hispan.
almijarensis, Campo & Amo, in Revista, v. (1855) 55 = tristis?
alpina, *Mill. Gard. Dict.* ed. VIII. n. 4.—Europ.
alsinaefolia, Benth. in DC. Prod. x. 269 = Acerbiana.
alsinaefolia, *Spreng. Syst.* ii. 792.—Sardin.; Corsic.
altaica, Fisch. ex Ledeb. Fl. Alt. ii. 448 = odora.
ambigua, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 254 = supina.
amethystea, *Hoffmgg. & Link, Fl. Port.* i. 253. t. 47.—Hispan.; Marocc.
amethystina, Spreng. Syst. ii. 795 = praec.
Amoi, Campo, ex Campo & Amo in Revista, v. (1855) 56 = tristis.
angustifolia, Reichb. Fl. Germ. Excurs. 375 = italica.
anticaria, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 86.—Hispan.

LINARIA:—
anthirrhinoides, Coss. ex Boiss. Diagn. Ser. II. iii. 162 = Cavanillesii.
aparinoides, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 395 = heterophylla.
apennina, Tausch, in Syll. Ratisb. ii. (1828) 247 = pallida?
aphylla, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 396; Steud. Nom. ed. I. 482; Spreng. Syst. ii. 797 = Utricularia capensis.
arenaria, *DC. Ic. Gall. Rar.* i. 5. t. 14.—Gallia.
armeniaca, *Chav. Monog. Antirrh.* 147.—Reg. Caucas.
arrabidensis, Welw. ex Nym. Consp. 539 = tristis.
arvensis, *Desf. Fl. Atlant.* ii. 45.—Reg. Meditrr.
ascalonica, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 165.—Palaestin.
asparagoides, *Schweinf. Beitr. Fl. Aethiop.* 99.—Afr. bor.
atlantica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 90.—Hispan.; Afr. bor.
Aucheri, *Boiss. Diagn.* Ser. I. vii. 44.—Persia.
aurasiaca, *Pomel, Nouv. Mat. Fl. Atl.* 299.—Alger.
Balansae, Boiss. & Reut. Diagn. Ser. II. iii. 163 = genistifolia.
barbata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 396 = Antirrhinum barbatum.
Bauhinii, *H. C. Wats. in Hook. Lond. Journ. Bot.* i. (1842) 76.—Europ.
bellidifolia, Dum.-Cours. Bot. Cult. ed. II. iii. 113 = Anarrhinum bellidifolium.
Besseriana, Reichb. Pl. Crit. v. 20 = macroura.
Biancae, Lojac. Osserv. Lin. Eur. 7, 13 = L. Elatine.
Biebersteinii, Bess. Enum. Pl. Volh. 25 = vulgaris.
Biebersteinii, Schur, Enum. Pl. Transs. 491 = hybrida.
bipartita, *Willd. Enum. Hort. Berol.* 640.—Afr. bor. occ.
bipunctata, Dum.-Cours. Bot. Cult. ed. I. ii. 93 = glauca.
bipunctata, Hort. ex Reichb. Fl. Germ. Excurs. 375 = versicolor.
bipunctata, Nym. Consp. 540 = amethystea.
Boissieri, Walp. Rep. iii. 211 = filifolia.
bombycina, Boiss. & Blanch. Diagn. Ser. II. iii. 161 = L. Elatine.
Bourgaei, Jord. Pugill. Pl. Nov. 129 = origanifolia.
Broussonnetii, *Chav. Monog. Antirrh.* 169.—Marocc.; Lusit.
Brunneri, *Benth. in DC. Prod.* x. 270.—Ins. Cap. Verd.
buriatica, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 98, nomen; xxiv. (1851) II. 302.—Sibir. baical.
cabulica, *Benth. in DC. Prod.* x. 270.—Persia; Ind. bor. occ.
caelesyriaca, Boiss. Diagn. Ser. I. xii. 41 = Aucheri.
caerulea, Hort. ex Chav. Monog. Antirrh. 145 = bipartita.
caesia, *DC. ex Chav. l. c.* 174.—Hispan.
calcarata, Scheele, in Flora, xxvi. (1843) 586 = armeniaca.
calycina, Boiss. & Bal. Diagn. Ser. II. vi. 130 = dalmatica.
canadensis, *Dum.-Cours. Bot. Cult.* ed. I. ii. 96.—Am. bor. et austr.
Candollei, Chav. Monog. Antirrh. 165 = arenaria.
capensis, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 397 = Nemesia linearis.
capilliflora, Steud. Nom. ed. II. ii. 48 = glandulifera?
capillipes, Hochst. ex Benth. in DC. Prod. x. 269 = hastata.
capitellata, Steud. Nom. ed. I. 482 = glauca.
capraria, *Moris & De Not. Fl. Caprar.* 98.—Ins. Caprar.
carnosa, Moench, Meth. 523 = arvensis.
caucasica, Muss. ex Spreng. Syst. ii. 790 = L. Elatine.
caulirhiza, Delile, ex Willk. & Lange, Prod. Fl. Hisp. ii. 559 = graeca.
Cavanillesii, *Chav. Monog. Antirrh.* 117.—Hispan.
chalepensis, *Mill. Gard. Dict.* ed. VIII. n. 12.—Europ. austr.; Oriens.
chinensis, Bunge, ex Benth. in DC. Prod x. 273 = vulgaris.
chloraefolia, Reichb. Pl. Crit. v. 21 = genistifolia.
ciliata, Lange, ex Reichb. l. c. 13 = italica.
circinnata, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 235 = sagittata.
cirrhosa, *Willd. Enum. Hort. Berol.* 689.—Reg. Mediterr.
Clementei, *Haensel. ex Boiss. Elench.* 69.—Hispan.

LINARIA:—
coelesyriaca, Auct. = *caelesyriaca*.
commutata, Bernh. ex Reichb. Fl. Germ. Excurs. 373 = graeca.
commutata, Schur, Enum. Pl. Transs. 488 = L. Elatine.
concolor, Griseb. Spicil. Fl. Rumel. ii. 21 = genistifolia.
confertiflora, *Benth. in DC. Prod.* x. 272.—Mesopot.
coridifolia, Boiss. Fl. Orient. iv. 376 = seq.
corifolia, *Desf. in Ann. Mus. Par.* xi. (1808) 53. t. 3.—As. Min.
crassifolia, DC. Fl. Fr. Suppl. 410, in obs.; Kunze, in Flora, xxix. (1846) 635 = origanifolia.
crenifolia, Hort. Matrit. ex Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 560 = L. Elatine.
cretacea, *Fisch. ex Spreng. Syst.* ii. 791.—Sibiria.
crinita, Mabille, Rech. Pl. Corse, fasc. i. 30 = L. Elatine.
cryptarum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 383.—Cataonia.
cuneata, Pritz. Ic. Ind. ii. 170, sphalm. = Veronica cuneata.
Cymbalaria, *Mill. Gard. Dict.* ed. VIII. n. 17.—Europ.
Cyparissias, Tausch, in Flora, xiv. (1831) 221 = repens.
dalmatica, *Mill. Gard. Dict.* ed. VIII. n. 13.—Dalmat.; Oriens.
damascena, *Boiss. & Gaill. Diagn.* Ser. II. vi. 130.—Syria.
dealbata, Hoffmgg. & Link, Fl. Port. i. 232. t. 34 = lanigera.
decumbens, Moench, Meth. 523 = repens.
delphinioides, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 127 = sapphirina.
dentata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 393 = scariosa.
dentata, Schur, Enum. Pl. Transs. 488 = spuria.
dentata, Steud. Nom. ed. I. 482 = Nemesia monomotapensis.
depauperata, *Leresche, ex Willk. & Lange, Prod. Fl. Hisp.* ii. 569.—Hispan.
Derderiana, Auch. ex Boiss. Fl. Orient. iv. 374 = Michauxii.
dianthifolia, *Donnersm. ex Chav. Monog. Antirrh.* 181, nomen.—Hab.?
dichondraefolia, *Benth. in DC. Prod.* x. 270.—Ins. Cap. Verd.
diffusa, Hoffmgg. & Link, Fl. Port. i. 257. t. 49 = multicaulis.
dissita, *Pomel, Nouv. Mat. Fl. Atl.* 98.—Alger.
dshorochensis, *C. Koch, in Linnaea*, xxii. (1849) 718.—As. Min.
dubia, Dum. Fl. Belg. 34 = supina.
Elatine, *Mill. Gard. Dict.* ed. VIII. n. 16 (Elatina).—Europ.; Oriens.
Elatine, Urv. in Mém. Soc. Linn. Par. i. (1822) 330 = graeca.
elatiniflora, Nym. Consp. 542 = spuria.
elatinoides, *Desf. Fl. Atlant.* ii. 39. t. 132.—Afr. bor.
elatinoides, Schur, Enum. Pl. Transs. 488 = L. Elatine.
elegans, *Cav. Desc.* 338.—Hispan.
elegans, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 604 = bipartita.
elegans, *Munby, in Bull. Soc. Bot. Fr.* xi. (1864) 46.—Alger.
elongata, Dum. Fl. Belg. 34 = vulgaris.
eriocalyx, Boiss. Fl. Orient. iv. 372 = pycnophylla.
exilis, *Coss. & Kralik, in Bull. Soc. Bot. Fr.* iv. (1857) 406.—Afr. bor.
fasciculata, *Otto, ex Sweet, Hort. Brit.* ed. II. 391, nomen.—Hab.?
fastigiata, *Chav. Monog. Antirrh.* 125. t. 7.—Persia.
faucicola, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 200.—Hispan.
filicaulis, *Boiss. ex Leresche & Levier, l. c.*—Hispan.
filifolia, *Lag. Gen. et Sp. Nov.* 19.—Hispan.
filiformis, Moench, Meth. 523 = supina.
filiformis, *Spreng. Fl. Hal. Mant.* 45.—Hab.?
flava, *Desf. Fl. Atlant.* ii. 42. t. 136.—Afr. bor.
flavissima, Boiss. Fl. Orient. iv. 378 = genistifolia.
flexuosa, *Desf. Fl. Atlant.* ii. 47. t. 139.—Afr. bor.
floribunda, *Boiss. Diagn.* Ser. I. xii. 40.—Palaestin.
floridana, *Chapm. Fl. S. U. St.* 290.—Am. bor.
Fontanesiana, *Chav. Monog. Antirrh.* 168.—Hispan.
fragilis, *Rodrig. in Bull. Soc. Bot. Fr.* xxv. (1878) 240.—Ins. Balear.
frutescens, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 400 = Nemesia fruticans.

LINARIA:—
fruticans, Spreng. Syst. ii. 789 = praec.
fruticosa, *Desf. Fl. Atlant.* ii. 39. t. 133.—Afr. bor.
galioides, *Ball, in Journ. Bot.* xiii. (1875) 173.—Marocc.
galioides, Schult. ex Steud. Nom. ed. I. 482 = repens.
gangitis, Duval-Jouve, in Mém. Acad. Montp. ix. (1878) reimp. 5 = origanifolia.
Gebleri, Bess. ex Chav. Monog. Antirrh. 132 = vulgaris.
geminiflora, *F. Schmidt, Reis. Amurl.* 161.—Ins. Sachalin.
genistifolia, Benth. Cat. Pl. Pyr. 67 = vulgaris.
genistifolia, DC. Fl. Fr. iii. 591 = italica.
genistifolia, *Mill. Gard. Dict.* ed. VIII. n. 14.—Europ.; As. occ. et bor.
glabrata, H. B. & K. Nov. Gen. et Sp. ii. 362 = triphylla.
glacialis, *Boiss. Elench.* 70.—Hispan.
glandulifera, *Tenore, Ind. Sem. Hort. Neapol.* (1840); *ex Walp. Rep.* iii. 221.—Hab.?
glareosa, Boiss. & Reut. Pugill. Pl. Nov. (1852) 84 = origanifolia.
glauca, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 15.—Hispan.
glauca, Ehrenb. ex Boiss. Fl. Orient. iv. 382 = albifrons.
glauca, Schur, Enum. Pl. Transs. 491 = vulgaris.
glaucophylla, Hoffmgg. & Link, Fl. Port. i. 250 = supina.
glaucophylla, Schur, Enum. Pl. Transs. 491 = vulgaris.
glutinosa, Benth. in DC. Prod. x. 284 = Tournefortii.
glutinosa, Hoffmgg. & Link, Fl. Port. i. 234 = filifolia.
gracilescens, *Pomel, Nouv. Mat. Fl. Atl.* 97.—Alger.
gracilis, *R. Br. in Salt, Abyss. App.* 64, nomen; *Benth. in DC. Prod.* x. 269.—Abyss.
gracilis, Desf. ex Chav. Monog. Antirrh. 168 = Fontanesiana.
gracilis, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 401 = Pelisseriana.
graeca, *Chav. Monog. Antirr.* 108.—Reg. Mediterr.
granatensis, Willk. in Bot. Zeit. v. (1847) 877 = villosa.
grandiflora, Desf. in Ann. Mus. Par. xi. (1808) 51 = dalmatica.
Griffithii, *Benth. in DC. Prod.* x. 272.—Afghan.
Haelava, *Delile, Fl. Aegypt. Illustr.* 66.—Aegypt.; Arab.
Haenseleri, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 88.—Hispan.
hastata, *R. Br. in Salt, Abyss. App.* 64, nomen; *Benth. in DC. Prod.* x. 269.—Abyss.
hastata, Royle, ex Chav. Monog. Antirrh. 112 = ramosissima.
hederaefolia, Salzm. ex Chav. l. c. 100 = hepaticaefolia.
hederaefolia, Steud. Nom. ed. I. 482 = L. Cymbalaria.
Hegelmaieri, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 101.—Hispan.
hepatica, Bunge, in Ledeb. Fl. Alt. ii. 445 = macroura.
hepaticaefolia, *Steud. Nom.* ed. I. 482; *Spreng. Syst.* ii. 790.—Cors.; Sardin.
heterophylla, *Desf. Fl. Atlant.* ii. 48. t. 140.—Reg. Mediterr.
heterophylla, Steud. Nom. ed. I. 482; Spreng. Syst. ii. 790 = sagittata.
hexandra, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 394 = Vandellia crustacea.
hians, Fenzl, ex Benth. in DC. Prod. x. 275 = genistifolia.
hians, Spreng. Syst. ii. 794 = vulgaris.
hirta, *Moench, Meth. Suppl.* 170.—Hispan.
humilis, Guss. Fl. Sic. Syn. ii. 123 = heterophylla.
Huteri, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 99.—Hispan.
hybrida, *Schur, Enum. Pl. Transs.* 491.—Transsilv.
iconia, *Boiss. & Heldr. Diagn.* Ser. I. xii. 42.—Lycaonia.
ignescens, Kunze, in Flora, xxix. (1846) 692 = Broussonnetii.
incana, *Wall. Pl. As. Rar.* ii. 43.—Reg. Himal.
incarnata, Spreng. Syst. ii. 796 = bipartita.
indecora, *Franch. in Révoil, Çomali*, 55.—Afr. trop.
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 175 = vulgaris.
italica, *Trevir. in Nov. Act. Nat. Cur.* xiii. 1. (1826) 188.—Europ. austr.

LINARIA:—
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 115.—Japon.
juncea, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 17.—Hispan.
juncea, Desf. Fl. Atlant. ii. 43 = spartea.
juncea, Reichb. Pl. Crit. v. 15. t. 425 = odora.
Kocianovichii ×, Aschers.—Cf. Bull. Soc. Bot. Fr. xvi. (1869) Bibl. 50.—Europ.
koesensis ×, Simk. in Termész. Füzetek, ii. (1878) 143.—Hungar.
kokanica, *Regel, Pl. Nov. Fedsch.* 60.—Kokania.
kurdica, *Boiss. & Hohen. in Boiss. Diagn.* Ser. I. iv. 73.—Kurdist.
lagopodioides, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 415 = saxatilis.
Lancerottae, Delile, Ind. Sem. Hort. Monsp. (1836) 26 = sagittata.
lancifolia, Schur, in Verh. Siebenb. Naturw. Ver. (1851) 197 = genistifolia.
Langei, *Nym. Consp*, 541.—Hispan.
lanigera, *Desf. Fl. Atlant.* ii. 38. t. 130.—Reg. Mediterr.
lanigera, Hoffmgg. & Link, Fl. Port. i. 231. t. 34 = spuria.
Lapeyrousiana, Jord. Pugill. Pl. Nov. 129 = origanifolia.
lasiopoda, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxvii. (1878) 389 = L. Elatine.
latifolia, *Desf. Fl. Atlant.* ii. 40. t. 134.—Reg. Medit. occ.
latifolia, *Gueldenst. Reisen*, ii. 178 (Quid?).—Ross. austr.
laxiflora, Coss. ex Willk. & Lange, Prod. Fl. Hisp. ii. 565 = nigricans.
laxiflora, *Desf. Fl. Atlant.* ii. 45. t. 138.—Afr. bor.
lilacina, *Lange, Ind. Sem. Hort. Haun.* (1854) 24.—Hispan.
linarioides, Steud. Nom. ed I. 482 = vulgaris.
lineolata, *Boiss, Diagn.* Ser. I. xii. 42.—Persia.
linifolia, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 415 = vulgaris.
linifolia, Reichb. Pl. Crit. v. 14. t. 422 = genistifolia.
linifolia, Rochel, Pl. Rar. Banat. 60. t. 22 = italica.
linogrisea, Hoffmgg. & Link, Fl. Port. i. 240. t. 11 = bipartita.
litoralis, *Willd. Enum. Hort. Berol.* 641.—Mare Adriat.
Loeselii, Schweigger, in Koenigsb. Archiv Naturw. i. (1812) 228 = odora.
longipes, *Boiss. & Heldr. Diagn.* Ser. I. xii. 40.—As. Min.
lurida, Ball, in Journ. Bot. xiii. (1875) 173 = tristis.
lusitanica, Hoffmgg. & Link, Fl. Port. i. 247. t. 43 = tristis.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = triornithophora.
macedonica, *Griseb. Spicil. Fl. Rumel.* ii. 19.—Macedon.
macilenta, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 252.—Arab.
macrocalyx, *Pomel, Nouv. Mat. Fl. Atl.* 98.—Alger.
macropoda, *Boiss. & Reut. Pugill. Pl. Nov.* 83.—Hispan.
macroura, *Link, Enum. Hort. Berol.* ii. 137; *Reichb. Pl. Crit.* v. 20. t. 432.—Reg. Caucas.
macrura, Schur, Enum. Pl. Transs. 491 = hybrida.
marginata, Desf. Fl. Atlant. ii. 43 = tristis.
maritima, DC. Ic. Pl. Rar. 5. t. 12 = supina.
maritima, Reichb. Fl. Germ. Excurs. 375 = odora.
maroccana, *Hook. f. Bot. Mag.* t. 5983.—Marocc.
melanantha, Boiss. & Reut. Pugill. Pl. Nov. 85 = tristis.
meonantha, Hoffmgg. & Link, Fl. Port. i. 236. t. 37 = spartea.
Michauxii, *Chav. Monog. Antirrh.* 124. t. 7. f. B.—Persia.
micrantha, Hoffmgg. & Link, Fl. Port. i. 258 = parviflora.
microcalyx, *Boiss. Diagn.* Ser. I. iv. 72.—Graecia.
microsepala, *A. Kern. Sched. Fl. Exsicc.* i. 50.—Europ.
minor, Desf. Fl. Atlant. ii. 46 = viscida.
minutiflora, C. A. Mey. Verz. Pfl. Cauc. 109 = albifrons.
modesta, Boiss. Diagn. Ser. I. xii. 43 = albifrons.
monochroma, Boiss. & Heldr. in Boiss. Diagn. Ser. I. xii. 41 = genistifolia.

LINARIA:—
monomotapensis, Desf. ex Spreng. Syst. ii. 797 = Nemesia monomotapensis.
monspessulana, Mill. Gard. Dict. ed. VIII. n. 9 = repens.
Muelleri, Moris, Diagn. Stirp. Sard. Nov. 1 = pilosa.
multicaulis, *Mill. Gard. Dict.* ed. VIII. n. 7.—Reg. Mediterr.
multicaulis, Welw. ex Willk. & Lange, Prod. Fl. Hisp. ii. 567 = Broussonnetii.
multipunctata, Hoffmgg. & Link, Fl. Port. i. 254. t. 48 = Broussonnetii.
Munbyana, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 89.—Reg. Mediterr. occ.
muralis, Salzm. ex Chav. Monog. Antirrh. 93 = villosa.
Muteli, Timb. in Bull. Soc. Phys. & Nat. Toul. vi. (1883–84) 135 = origanifolia.
neglecta, Guss. Pl. Rar. 248 = triphylla.
nervosa, Schur, Enum. Pl. Transs. 490 = genistifolia.
nevadensis, Boiss. & Reut. Pugill. Pl. Nov. (1852) 87 = supina.
nigricans, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* ii. 565.—Hispan.
nivea, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 22.—Hispan.
nummularia, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 581 = villosa.
nurensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 374.—Persia.
oblongifolia, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 86.—Hispan.
ochroleuca ×, Bréb. ex Nym. Consp. 539.—Europ.
ocimifolia, *Spreng. Syst.* ii. 793.—Hispan.
odora, [*Fisch.*] *Cat. Hort. Gorenk.* ed. II. (1812) 25; *Chav. Monog. Antirrh.* 136.—Oriens.
oligantha, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 100.—Hispan.
oligotricha ×, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 393.—Europ.
oppositiflora, *Steud. Nom.* ed. I. 483; *Spreng. Syst.* ii. 792.—Marocc.
orchioides, Herb. Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 563 = sappharina.
orgyalis, Reut. ex Boiss. Fl. Orient. iv. 377 = genistifolia.
origanifolia, *DC. Fl. Fr.* iii 591.—Europ. austr.
Osyris, Link, Handb. i. 537 = chalepensis.
ovata, *Boiss. Fl. Orient.* iv. 369.—Afghan.
pallida, Sprun. ex Boiss. l. c. 378 = Sibthorpiana.
pallida, *Tenore, Fl. Nap.* i. 32. t. 159.—Italia.
paniculata, Peyer, ex Vest, in Flora, xii. (1829) 1. Erg. 30 = italica.
parnassica, Boiss. & Heldr. Diagn. Ser. II. iii. 164 = Sibthorpiana.
parviflora, *Desf. Fl. Atlant.* ii. 44. t. 137.—Hispan.; Afr. bor.
parvula, Duf. ex Willk. & Lange, Prod. Fl. Hisp. ii. 567 = Munbyana.
patens, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 403 = Nemesia patens.
pauciflora, Bonpl. Desc. Pl. Rar. Malm. 48. t. 18 = multicaulis.
pedunculata, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* iv. 417.—Hispan.
Pelisseriana, *Mill. Gard. Dict.* ed. VIII. n. 11.—Europ. austr.; As. Min.
Pellicieriana, Puel, in Bull. Soc. Bot. Fr. viii. (1861) 631 = praec.
peloponnesiaca, Boiss. & Heldr. Diagn. Ser. II. iii. 163 = Sibthorpiana.
pensylvanica, Scheele, in Flora, xxvi. (1843) 586 = vulgaris.
Perezii, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 127 = Tournefortii.
persica, *Chav. Monog. Antirrh.* 173.—Persia.
petraea, *Jord. in Compt.-Rend. Soc. Hallér.* (1854–56) 116.—Europ.
petraea, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) 1. 347 = genistifolia.
pilosa, *DC. Fl. Fr.* iii. 584.—Italia.
platycalyx, *Boiss. Voy. Bot. Espagne*, ii. 459. t. 130.—Hispan.
polyclada, Fenzl, in Tchihat. Asie Min. Bot. ii. 23 = genistifolia.
polygalaefolia, Hoffmgg. & Link, Fl. Port. i. 248 = supina.
polygalaefolia, Steud. Nom. ed. I. 483 = italica.

LINARIA:—
polygonifolia, Benth. in DC. Prod. x. 282 = verticillata.
polygonifolia, *Steud. Nom.* ed. I. 483; *Spreng. Syst.* ii. 791.—Hispan.
praealta, *Boiss. & Bal. Diagn.* Ser. II. iii. 162.—Cilicia.
praecox, Bunge, in Ledeb. Fl. Alt. ii. 446 = odora.
praecox, Hoffmgg. & Link, Fl. Port. i. 234. t. 37 = spartea.
praetermissa, Delastre, in Ann. Sc. Nat. Sér. II. xviii. (1842) 151 = viscida.
Prestandreae, Tin. in Guss. Fl. Sic. Syn. ii. 842 = L. Elatine.
procera, DC. Cat. Hort. Monsp. 121 = repens.
propinqua, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 88.—Europ.
propinqua, R. Br. in Salt, Abyss. App. 64, nomen = gracilis.
prostrata, Boenn. Prod. Fl. Monast. 187 = vulgaris.
Pseudo-laxiflora, *Lojac. in Natural. Sicil.* iv. (1885) 136.—Sicilia.
pterospora, *Boiss. Fl. Orient.* iv. 384.—As. Min.; Persia.
pubescens, Desf. ex Chav. Monog. Antirrh. 98 = litoralis.
pubescens, Presl, Delic. Prag. 74 = pilosa.
pulchella, Hohen. ex Boiss. Fl. Orient. iv. 379 = lineolata.
purpurascens, Bernh. ex Hornem. Hort. Hafn. ii. 573 = purpurea.
purpurea, *Mill. Gard. Dict.* ed. VIII. n. 5.—Europ. austr.
pycnophylla, *Boiss. & Bal. Diagn.* Ser. II. vi. 129.—As. Min.
pyramidalis, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 417; *Steud. Nom.* ed. I. 483 = pyramidata.
pyramidata, Schur, Enum. Pl. Transs. 490 = genistifolia.
pyramidata, *Spreng. Syst.* ii. 796.—Reg. Caucas.
pyrenaica, DC. Fl. Fr. iii. 587 = supina.
quadrifolia, *Hance, in Journ. Bot.* ix. (1871) 133.—China.
racemosa, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 405 = vulgaris.
radicans, Le Gall, ex Gren. & Godr. Fl. Fr. ii. 575 = graeca.
Ramburei, Boiss. ex Benth. in DC. Prod. x. 276 = viscosa.
ramosissima, Boiss. Voy. Bot. Espagne, ii. 457 = filifolia.
ramosissima, *Wall. Pl. As. Rar.* ii. 43. t. 153.—Ind. or.; Burma,
Raveyi, Boiss. Elench. 70 = rubrifolia.
reflexa, *Desf. Fl. Atlant.* ii. 42.—Reg. Mediterr.
repens, *Mill. Gard. Dict.* ed. VIII. n. 6.—Europ.
reticulata, *Desf. Fl. Atlant.* ii. 48.—Afr. bor.; Lusit.
reticulata, Dulac, Fl. Hautes-Pyr. 375 = L. Elatine.
reticulata, Reichb. Pl. Crit. v. 19. t. 431. f. 620 = heterophylla.
reticulata, Webb, It. Hisp. 26 = Clementei.
rhegina, Spreng. Syst. ii. 792 = reflexa.
Rossmaessleri, Willk. in Linnaea, xxx. (1859–60) 118 = tristis?
Roylei, Chav. Monog. Antirrh. 112 = ramosissima.
rubioides, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 473.—Austria.
rubrifolia, *Rob. & Cast. ex DC. Fl. Fr. Suppl.* 410.—Europ. austr.
rudis, Janka, ex Nym. Consp. 537 = vulgaris.
rupestris, C. A. Mey. Verz. Pfl. Cauc. 110 = macroura.
rupestris, Guss. Fl. Sic. Syn. ii. 120 = rubrifolia.
rytidosperma, Boiss. Diagn. Ser. I. iv. 73 = persica.
sabulicola, *Pomel, Nouv. Mat. Fl. Atl.* 98.—Alger.
saetabensis, Auct. = *setabensis*.
sagittata, *Hook. f. Bot. Mag.* t. 6060.—Marocc.; Ins. Canar.
sagittata, Steud. Nom. ed. I. 483 = heterophylla.
Salzmannii, Boiss. Voy. Bot. Espagne, ii. 456. t. 128 = viscosa.
saphiria, Steud. Nom. ed. I. 483 = amethystea.
sapphirina, *Hoffmgg. & Link, Fl. Port.* i. 241. t. 42.—Lusitan.
satureioides, *Boiss. Voy. Bot. Espagne*, ii. 463. t. 133.—Hispan.
saxatilis, Benth. in DC. Prod. x. 284 = Tournefortii.
saxatilis, Boiss. Voy. Bot. Espagne, ii. 462 = verticillata.

LINARIA:—
saxatilis, DC. Fl. Fr. iii. 590; Suppl. 409 = arenaria.
saxatilis, *Hoffmgg. & Link, Fl. Port.* i. 238.—Lusitan.
scabra, Spreng. Syst. ii. 792 = Nemesia chamaedrifolia.
scariosa, *Desf. Fl. Atlant.* ii. 38. t. 131.—Afr. bor.
scoparia, Brouss. ex Spreng. Syst. ii. 789 = spartioides.
segetalis, C. Koch, in Linnaea, xvii. (1843) 286 = armeniaca.
segoviensis, Reut. ex Nym. Consp. 541 = origanifolia.
semiglabra, Salzm. ex Chav. Monog. Antirrh. 116 = hirta.
sepium ×, J. G. Allm. in Proc. Irish Acad. ii. (1844) 404.—Europ.
serpyllifolia, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 39.—Hispan.
setabensis, Leresche, ex Lange, in Prod. Fl. Hisp. ii. 580 (saetabensis) = origanifolia.
Sibthorpiana, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 378.—Graecia.
Sieberi, Reichb. Fl. Germ. Excurs. 374 = L. Elatine.
silenifolia, Fisch. Cat. Hort. Gorenk. 25; Reichb. Pl. Crit. v. 22. t. 437 = genistifolia.
simplex, DC. Fl. Fr. iii. 588 = arvensis.
Smithii, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 376 = dalmatica.
somalensis, *Vatke, in Linnaea*, xliii. (1880–82) 305.—Afr. trop.
spartea, *Hoffmgg. & Link, Fl. Port.* i. 233.—Reg. Mediterr. occ.
spartioides, *Brouss. ex Buch, Beschr. Canar. Ins.* 163.—Ins. Canar.
speciosa, Jacq. f. Eclog. i. 141. t. 95 = bipartita.
speciosa, Tenore, Fl. Nap. v. 40 = vulgaris.
spectabilis, *Pomel, Nouv. Mat. Fl. Atl.* 299.—Alger.
sphaerocarpa, Benth. in DC. Prod. x. 287 = Schweinfurthia sphaerocarpa.
spicata, Kunze, in Flora, xxix. (1846) 645 = viscosa.
spinescens, Spreng. Syst. ii. 789 = aegyptiaca.
spuria, *Mill. Gard. Dict.* ed. VIII. n. 15.—Europ.; Afr. bor.
stenantha, *Franch. in Révoil, Çomali*, 55.—Afr. trop.
Steveni, Nym. Syll. Suppl. 22 = genistifolia.
striata, DC. Fl. Fr. iii. 586 = repens.
striata, Ledeb. Fl. Ross. iii. 210 = lineolata.
stricta, Guss. Pl. Rar. Sic. 250 = heterophylla.
stricta, Hornem. Hort. Hafn. ii. 575 = repens.
subalpina, Steud. Nom. ed. I. 483 = amethystea.
supina, *Desf. Fl. Atlant.* ii. 44.—Reg. Mediterr.
syspirensis, C. Koch, in Linnaea, xxii. (1849) 717 = genistifolia.
tatarica, Fisch. ex Ledeb. Fl. Ross. iii. 207 = macroura.
tenella, *DC. Fl. Fr. Suppl.* 410, in obs.—Hispan.
tenella, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 408; Steud. Nom. ed. I. 483 = Collinsia verna.
tenuis, *Spreng. Syst.* ii. 795.—Afr. bor.
texana, Scheele, in Linnaea, xxi. (1848) 761 = canadensis.
Thuillieri, Mérat, Fl. Par. ed. III. ii. 192 = supina.
thymifolia, *DC. Fl. Fr.* iii. 587.—Gallia.
thymophyllum, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 203 = thymifolia.
tingitana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 84 = heterophylla.
Tournefortii, *Steud. Nom.* ed. I. 483; ed. II. ii. 49.—Hispan.
transsilvanica, Schur, Enum. Pl. Transs. 490 = dalmatica.
transtagana, Spreng. Syst. ii. 792 = pedunculata.
triornithophora, *Willd. Enum. Hort. Berol.* 639; *Hoffmgg. & Link, Fl. Port.* i. 244.—Lusitan.
triphylla, *Mill. Gard. Dict.* ed. VIII. n. 2.—Reg. Mediterr.
triquetra, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 74 = spartea.
tristis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Hispan.
tumens, *Steud. Nom.* ed. II. ii. 53, nomen.—Aegypt.
venosa, Lindl. Bot. Reg. (1841) Misc. 69 = odora.
ventricosa, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 252.—Marocc.
veronicoides, *A. Rich. Tent. Fl. Abyss.* ii. 114.—Abyssin.
versicolor, *Moench, Meth.* 523.—Gall. austr.
verticillata, *Boiss. Voy. Bot. Espagne*, ii. 462. t. 132.—Hispan.
villosa, *DC. Fl. Fr. Suppl.* 410, in obs.—Hispan.; Marocc.
villosa, Mill. Gard. Dict. ed. VIII. n. 10 = hirta.

LINARIA:—
violacea, Haensel. ex Willk. in Bot. Zeit. iv. (1846) 313 = satureioides.
virgata, *Desf. Fl. Atlant.* ii. 41. t. 135.—Afr. bor.
virgatula, Spreng. Syst. ii. 796 = spartea.
viscida, *Moench, Meth.* 524.—Europ.; Afr. bor.
viscosa, *Dum.-Cours. Bot. Cult.* ed. I. ii. 93.—Reg. Mediterr. occ.
vulgaris, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.; Reg. Caucas.
vulgaris, Plan. Ens. Fl. Gallega, 315 = Tournefortii.
Warionis, *Pomel, Nouv. Mat. Fl. Atl.* 96.—Alger.
Webbiana, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 240.—Ins. Cap. Verd.
Webbiana, Vis. Ort. Bot. Torin. 142 = sagittata.
Zelliceriana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126, err. typ. = Pelisseriana.

LINARIOPSIS, Welw. in Trans. Linn. Soc. xxvii. (1869) 53. *PEDALINEAE*, Benth. & Hook. f. ii. 1060.
prostrata, *Welw. l. c.* 54.—Afr. trop.

LINCANIA, G. Don, Gen. Syst. ii. 479 (1832) = **Licania**, Aubl. (Rosac.).

LINCONIA, Linn. Mant. ii. 216 (1767). *BRUNIACEAE*, Benth. & Hook. f. i. 672.
alopecuroides, *Linn. l. c.*—Afr. austr.
capitata, Banks, ex Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 381 = Nebelia affinis.
cuspidata, *Sw. in Ges. Naturf. Fr. Berl. Mag.* v. (1811) 284.—Afr. austr.
peruviana, *Lam. Encyc.* iii. 527.—Peruv.
tamariscina, E. Mey. in Drège, Zwei Pfl. Docum. 199 = Brunia pinifolia.
thymifolia, *Sw. in Ges. Naturf. Fr. Berl. Mag.* iv. (1810) 86. t. 4.—Afr. austr.

LINDACKERA, Sieber, ex Endl. Gen. 893, in syn. (1841) = **Capparis**, Linn.
capparoidea, Sieber, ex Steud. Nom. ed. II. ii. 501 = C. aphylla.

LINDACKERIA, Presl, Rel. Haenk. ii. 89. t. 65 (1831) = **Oncoba**, Forsk. (Bixin.).
latifolia, Benth. in Hook. Kew Journ. iii. (1851) 118 = O. latifolia.
laurina, Presl, Rel. Haenk. ii. 89 = Mayna laurina.
maynensis, Poepp. & Endl. Nov. Gen. et Sp. iii. 63. t. 270 = O. maynensis.
pauciflora, Benth. in Hook. Kew Journ. iii. (1851) 118 = O. pauciflora.
vernicosa, Karst. Fl. Columb. ii. 11. t. 106.—N. Granat.

LINDBLOMIA, Fries, in Lindb. Bot. Notiser (1843) 134 = **Habenaria**, Willd. (Orchid.).

LINDELOFIA, Lehm. in Hamb. Gartenz. vi. (1850) 351. *BORAGINEAE*, Benth. & Hook. f. ii. 848.
ANCHUSOPSIS, Bisch. Del. Sem. Hort. Heidelb. (1852).
anchusoides, Lehm. in Hamb. Gartenz. vi. (1850) 352 = Paracarpum heliocarpum.
Benthami, *Hook. f. Fl. Brit. Ind.* iv. 159.—Reg. Himal.
spectabilis, *Lehm. in Hamb. Gartenz.* vi. (1850) 352 —Reg. Himal.

LINDENBERGIA, Lehm. in Link & Otto, Ic. Pl. Rar. 95. t. 48 (1828). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 948.
BOVEA, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 253.
BRACHYCORYS, Schrad. Ind. Sem. Hort. Gotting. (1830).—Cf. Linnaea, vi. (1831) Litt. 72.
HEMIORCHIS, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 241 (1867).
abyssinica, *Hochst. ex Benth. in DC. Prod.* x. 377.—Sind.; Arab.; Abyssin.
fruticosa, *Benth. in DC. Prod.* x. 377.—Arabia.
grandiflora, *Benth. Scroph. Ind.* 22.—Reg. Himal.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 262.—Reg. Himal.
Hookeri, *C. B. Clarke, ex Hook. f. l. c.* iv. 261.—Reg. Himal.
indica, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 10 = polyantha

LINDENBERGIA:—
macrostachya, *Benth. Scroph. Ind.* 22.—Reg. Himal.; China.
nigrescens, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 10.—Afr. trop.
philippinensis, *Benth. in DC. Prod.* x. 377.—Burma; Malaya.
polyantha, *Royle, ex Benth. Scroph. Ind.* 22.—Ind. or.
ruderalis, Voigt, Hort. Suburb. Calc. 501 = urticaefolia.
scutellarioides, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 241.—Afr. trop.
siamensis, Miq. ex Hook. f. Fl. Brit. Ind. iv. 262 = macrostachya.
siamensis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 411.—Siam.
sinaica, *Benth. Scroph. Ind.* 22.—Arabia.
urticaefolia, *Lehm. Ind. Sem. Hort. Berol.* 1829 (1830) 5, nomen; *et in Link & Otto, Ic. Pl. Rar. Berol.* 95. t. 48.—Ind. or.
virens, *Vatke, in Linnaea*, xliii. (1880–82) 306.—Arab.

LINDENIA, Benth. Pl. Hartw. 84, 351 (1842). *RUBIACEAE*, Benth. & Hook. f. ii. 51.
SIPHONIA, Benth. Pl. Hartw. 84 (1841).
acutiflora, *Benth. in Hook. Ic. Pl.* t. 475.—Mexic.
austro-caledonia, *Brongn. in Bull. Soc. Bot. Fr.* xii. (1865) 407.—N. Caled.
rivalis, *Benth. Pl. Hartw.* 351.—Guatemala.
vitiensis, *Seem. in Bonplandia*, ix. (1861) 256; x. (1862) 33. t. 8.—Ins. Fiji.

LINDENIA, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 357 = **Senkenbergia**, Schau. (Nyctagin.).
gypsophiloides, Mart. & Gal. l. c. = S. annulata.

LINDERA, Adans. Fam. ii. 499 (1763) = **Myrrhis**, Scop. (Umbellif.).

LINDERA, Thunb. Diss. Nov. Gen. iii. 44 (1783); Fl. Jap. 9, 145. t. 21 (1784). *LAURINEAE*, Benth. & Hook. f. iii. 163.
APERULA, Blume, Mus. Bot. Lugd. Bat. i. 365 (1851).
BENZOIN, Nees, Laurin. Expos. 17 (1833).
DAPHNIDIUM, Nees, in Wall. Pl. As. Rar. ii. 61, 63 (1831); et l. c. 19 (1833).
EUOSMUS, Nutt. Gen. Am. i. 258 (1818).
EVELYNA, Rafin. Sylva Tellur. 164 (1838).
OZANTHES, Rafin. l. c. 133 (1838).
POLYADENIA, Nees, Laurin. Expos. 18 (1833).
SASSAFRAS, Presl, Rostl. ii. (1825) 67.
acuminata, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
aestivalis, Blume, Mus. Bot. Lugd. Bat. i. 324 = L. Benzoin.
assamica, *Kurz, For. Fl. Brit. Burma*, ii. 308.—Reg. Himal.; Burma.
Benzoin, *Meissn. in DC. Prod.* xv. 1. 244.—Am. bor.
bootanica, *Meissn. l. c.* 245.—Reg. Himal.
caesia, *Reinw. ex Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.; Java.
glauca, *Blume, Mus. Bot. Lugd. Bat.* i. 325.—Japon.
Griffithii, *Meissn. in DC. Prod.* xv. 1. 245.—Reg. Himal.
heterophylla, *Meissn. l. c.* 246.—Reg. Himal.
Hookeri, *Meissn. l. c.* 245.—Reg. Himal.
hypoglauca, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 71.—Japon.
lancifolia, Trimen, Cat. Pl. Ceyl. 76 = Litsea lancifolia.
Loureiri, *Blume, Mus. Bot. Lugd. Bat.* i. 325.—Cochinch.
melastomacea, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Reg. Himal.
melissaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 324.—Am. bor.
membranacea, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 72.—Japon.
mollis, Oliver, in Journ. Linn. Soc. ix. (1865) 168 = obtusiloba.
Neesiana, *Kurz, For. Fl. Brit. Burma*, ii. 309.—Reg. Himal.
nervosa, *Kurz, l. c.* 308.—Burma.
obtusa, *Franch. & Sav. Enum. Pl. Jap.* ii. 483.—Japon.
obtusiloba, *Blume, Mus. Bot. Lugd. Bat.* i. 325.—Japon.
praecox, *Blume, l. c.* 324.—Japon.
reticulata, *Benth. & Hook. f. Gen.* iii. 163.—Reg. Himal.
sericea, *Blume, Mus. Bot. Lugd. Bat.* i. 324.—Japon.

LINDERA:—
sikkimensis, *Meissn. in DC. Prod.* xv. 1. 245.—Reg. Himal.
strychnifolia, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 182.—Ins. Philipp.
triloba, *Blume, Mus. Bot. Lugd. Bat.* i. 325.—Japon.
triloba, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. 1. 246 = heterophylla.
umbellata, *Thunb. Fl. Jap.* 145. t. 21.—Japon.

LINDERNIA, All. Misc. Taur. iii. (1755) 178. t. 5 = **Vandellia**, Linn. (Scrophul.).
alsinoides, R. Br. Prod. 441 = V. alsinoides.
attenuata, Muhl. Cat. ex Ell. Sketch, i. 17 = Ilysanthes riparia.
capensis, Thunb. Prod. Pl. Cap. 104 = Ilysanthes capensis.
clausa, F. Muell. Fragm. vi. 102 = Bonnaya clausa.
crustacea, F. Muell. Census, 97 = V. crustacea.
dianthera, Sw. Prod. Veg. Ind. Occ. 92 = Herpestis chamaedryoides.
dilatata, Muhl. Cat. ex Ell. Sketch, i. 16 = Ilysanthes riparia.
grandiflora, Nutt. Gen. Am. ii. 43 = Ilysanthes grandiflora.
gratioloides, Poir. ex Steud. Nom. ed. I. 484 = V. erecta.
inundata, Schur, Enum. Pl. Transs. 933, sphalm. = V. erecta.
japonica, Thunb. Fl. Jap. 253 = Mazus rugosus.
Kitaibelii, G. Don, Gen. Syst. iv. 551 = V. erecta.
lobelioides, F. Muell. Census, 97 = Ilysanthes lobelioides.
micrantha, D. Don, Prod. Fl. Nep. 85 = V. angustifolia.
montevidensis, Spreng. Syst. ii. 769 = Petunia parviflora.
monticola, Nutt. Gen. Am. i. 9 = Ilysanthes refracta.
plantaginea, F. Muell. Fragm. vi. 102 = Hemiarrhena plantaginea.
pubescens, F. Muell. Census, 97 = V. pubescens.
pyxidaria, All. Misc. Taur. iii. 178 = V. erecta.
pyxidaria, Pursh, Fl. Am. Sept. ii. 419 = Ilysanthes riparia.
refracta, Ell. Sketch, i. 579 = Ilysanthes refracta.
saxicola, M. A. Curt. in Am. Journ. Sc. Ser. I. xliv. (1843) 83 = Ilysanthes refracta.
scapigera, R. Br. Prod. 441 = V. scapigera.
serrata, F. Muell. Census, 97 = Bonnaya brachiata.
sesamoides, Spreng. Neue Entdeck. i. 261 = Herpestis floribunda.
subulata, R. Br. Prod. 441 = V. subulata.
veronicifolia, F. Muell. Fragm. vi. 101 = Bonnaya veronicaefolia.

LINDHEIMERA, A. Gray & Engelm. in Proc. Am. Acad. i. (1847) 47. *COMPOSITAE*, Benth. & Hook. f. ii. 351.
mexicana, *A. Gray, l. c.* xv. (1880) 34.—Mexic.
texana, *A. Gray, l. c.* i. (1847) 47.—Texas.

LINDLEYA, H. B. & K. Nov. Gen. et Sp. v. t. 480 (1821); Kunth, Malv. 10 (1822) = **Casearia**, Jacq. (Samydac.).
glabra, H. B. & K. l. c. t. 479 = C. javitensis.
mollis, H. B. & K. l. c. t. 480 = C. hirsuta.

LINDLEYA, H. B. & K. Nov. Gen. et Sp. vi. 239. t. 562, bis (1823). *ROSACEAE*, Benth. & Hook. f. i. 615.
mespiloides, *H. B. & K. l. c.*—Mexic.

LINDLEYA, Nees, in Flora, iv. (1821) 299 = **Laplacea**, H. B. & K. (Ternstr.).
fruticosa, Nees, l. c. = Laplacea semiserrata.

LINDNERA, Reichb. Handb. 303 (1837) = **Tilia**, Tourn.
alba, Fuss, Fl. Transs. 131 = T. tomentosa.

LINGOUM, Adans. Fam. ii. 319 (1763) = **Pterocarpus**, Linn. (Legumin.).

LINHAREA, Arruda, ex Steud. Nom. ed. I. 484 (1821) = seq.

LINHARIA, Arruda, ex Koster, Trav. Bras. 493 (1816) = **Umbellularia**? Nutt. (Laurin.).
aromatica, Arruda, ex Koster, l. c.—Bras.
tinctoria, Arruda, ex Koster, l. c.—Bras.

LINKIA, Cav. Ic. iv. 61. t. 389 (1797) = **Persoonia**, Sm. (Proteac.).
laevis, Cav. l. c. = P. lanceolata, salicina.

LINKIA, Pers. Syn. i. 219 (1805) = **Desfontainia**, Ruiz & Pav. (Logan.).
peruviana, Pers. l. c. = D. spinosa.
spinosa, Poir. Encyc. Suppl. iii. 449 = D. spinosa.
splendens, Poir. l. c. = D. spinosa.

LINNAEA, Gronov. in Linn. Gen. ed. I. 188 (1737). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 5.
DICODON, Ehrh. Beitr. iv. 146 (1789).
americana, Forbes, Hort. Woburn. 135 = borealis.
borealis, *Linn. Sp. Pl.* 631.—Europ.; Am. bor.
chinensis, A. Br. & Vatke, in Oestr. Bot. Zeitschr. xxii. (1872) 291 = Abelia chinensis.
floribunda, A. Br. & Vatke, l. c. = Abelia floribunda.
rupestris, A. Br. & Vatke, l. c. = Abelia chinensis.
triflora, A. Br. & Vatke, l. c. = Abelia triflora.
uniflora, A. Br. & Vatke, l. c. = Abelia uniflora.

LINNEA, Neck. Elem. i. 378 (1790) = praec.

LINOCHILUS, Benth. Pl. Hartw. 197 (1845) = **Diplostephium**, H. B. & K. (Compos.).
floribundus, Benth. l. c. 203 = D. floribundum.
jodopappus, Sch. Bip. ex Wedd. Chlor. And. i. 204 = D. Schultzii.
lavandulifolius, Sch. Bip. l. c. 200 = Aster lavandulaceus.
Meyenii, Sch. Bip. l. c. 201 = D. Meyenii.
microphyllus, Sch. Bip. l. c. = D. microphyllum.
ochraceus, Sch. Bip. l. c. 205 = D. floribundum.
phylicoides, Sch. Bip. l. c. = Tetramolopium phylicoides.
rosmarinifolius, Benth. Pl. Hartw. 197 = D. rosmarinifolium.

LINOCIERA, Steud. Nom. ed. II. ii. 50 (1841) = LINOCIRIA, Neck. = **Haloragis**, Forst.

LINOCIERA, Sw. in Schreb. Gen. ii. 784 (1791); Fl. Ind. Occ. i. 49. t. 2 (1797). *OLEACEAE*, Benth. & Hook. f. ii. 678.
BONAMICA, Vell. Fl. Flum. 21 (1825); Ic. i. t. 50 (1827).
CERANTHUS, Schreb. Gen. i. 14 (1789).
CHIONANTHUS, Gaertn. Fruct. i. 189. t. 39 (1788).
FREYERIA, Scop. Introd. 208 (1777).
HAENIANTHUS, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 518.
MAYEPEA, Aubl. Pl. Gui. i. 81. t. 31 (1775), nomen prius.
MINUTIA, Vell. Fl. Flum. 19 (1825); Ic. i. t. 47 (1827).
TESSARANDRA, Miers, ex Lindl. Veg. Kingd. 617 (1847); et in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 179.
THOUINIA, Linn. f. Suppl. 9, 89 (1781).
acuminata, Wall. Cat. n. 2844 = terniflora.
alba, Poit. ex DC. Prod. viii. 296 = compacta.
albidiflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 608.—Zeylan.; Austral.
arborea, *Eichl. in Mart. Fl. Bras.* vi. 1. 308.—Bras.
axilliflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 519.—Ind. occ.
bumelioides, *Griseb. Cat. Pl. Cub.* 169.—Cuba.
cambodiana, *Hance, in Journ. Bot.* xv. (1877) 335.—Cambodia.
compacta, *R. Br. Prod.* 523.—Ind. occ.
cotinifolia, *Vahl, Enum.* i. 46.—Zeylan.
crassifolia, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 64.—Bras.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 185.—Ins. Philipp.
densiflora, *Zoll. & Mor. Syst. Verz. Zoll.* 58.—Java.
dichotoma, Wall. Cat. n. 2825 = purpurea.
effusiflora, F. Muell. Fragm. iv. 83 = macrophylla.
elegans, *Eichl. in Mart. Fl. Bras.* vi. 1. 306.—Bras.
flavicans, Vahl, Enum. i. 47 = Noronhia emarginata.
glomerata, *Pohl, Pl. Bras. Ic.* ii. 98. t. 164.—Bras.
insignis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 610.—Burma.
intermedia, *Wight, Ic.* t. 1245.—Ind. or.
latifolia, *Vahl, Enum.* i. 46.—Ins. S. Doming.
laxiflora, *Blume, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 610.—Malaya.
leprocarpa, *C. B. Clarke, l. c.* 608.—Zeylan.

LINOCIERA:—
ligustrina, *Sw. Fl. Ind. Occ.* i. 50.—Ind. occ.
loranthifolia, *Wall. Cat.* n. 2842.—Burma.
luzonica, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 128.—Ins. Philipp.
macrophylla, *Wall. Cat.* n. 2826.—Reg. Himal.
malabarica, *Wall. l. c.* n. 2828.—Ind. or.
mandioccana, *Eichl. in Mart. Fl. Bras.* vi. 1. 308.—Bras.
micrantha, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 63.—Bras.
minutiflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 610.—Burma.
montana, *DC. Prod.* viii. 297.—Java.
Muelleri, Heurck, Obs. Bot. 124 = Bouea microphylla?
nigra, Poit. ex DC. Prod. viii. 296 = ligustrina.
nilotica, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 106.—Afr. trop.
oblonga, Wall. Cat. n. 2843 = intermedia.
odorata, Jack, in Malay. Misc. ii. (1822) n. VII. 96 = purpurea.
pauciflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 609.—Ins. Penang.
picrophloia, F. Muell. Fragm. iii. 139 = intermedia.
purpurea, *Vahl, Enum.* i. 47.—Zeylan.
ramiflora, Wall. Cat. n. 2824 = macrophylla.
rostrata, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 608 = albidiflora.
rostrata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 404.—Java.
subsessilis, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 198.—Bras.
tenuiflora, Wall. Cat. n. 2827 = macrophylla.
terniflora, *Wall. l. c.* n. 2845.—Burma.
tetrandra, *R. Br. Prod.* 523.—Guian.
Wightii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 608.—Ind. or.

LINOCIRIA, Neck. Elem. iii. 366 (1790) = **Haloragis**, Forst.

LINODENDRON, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 187 = **Lasiadenia**, Benth. (Thymel.).
aronifolium, Griseb. Cat. Pl. Cub. 110.—Cuba.
cubense, Griseb. l. c. = Lasiadenia cubensis.
Lagetta, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 187.—Cuba.
venosum, Griseb. Cat. Pl. Cub. 110.—Cuba.

LINOIDES, Rupp. Fl. Jen. ed. Hall. 91 (1745) = **Radiola**, Dill. (Lineae).

LINOPSIS, Reichb. Handb. 306 (1837) = **Linum**, Linn.
aethiopica, Reichb. l. c. 307 = Linum africanum.
africana, Reichb. l. c. = Linum africanum.
corymbosa, Reichb. l. c. = Linum strictum?
maritima, Reichb. l. c. = Linum maritimum.
quadrifolia, Reichb. l. c. = Linum quadrifolium.

LINOSPADIX, Becc. Malesia, i. (1877) 62. *PALMAE*, Benth. & Hook. f. iii. 903.
arfakianus, *Becc. l. c.*—N. Guin.
flabellatus, *Becc. l. c.* 64.—N. Guin.
multifidus, *Becc. l. c.*—N. Guin.

LINOSPADIX, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 177 = **Bacularia**, F. Muell. (Palm.).
monostachyos, H. Wendl. & Drude, l. c. 199 = B. monostachya.

LINOSPARTON, Adans. Fam. ii. 34 (1763) = **Lygeum**, Loefl. (Gramin.).

LINOSPARTUM, Steud. Nom. ed. II. ii. 50 (1841) = praec.

LINOSTACHYS, Klotzsch, ex Schlecht. in Linnaea, xix. (1847) 235, in nota = **Acalypha**, Linn. (Euphorb.).
padifolia, Schlecht. l. c. = A. Schlechtendahliana.
urticaefolia, Klotzsch, ex Schlecht. l. c., in nota = A. urticaefolia.

LINOSTIGMA, Klotzsch, in Linnaea, x. (1835) 438 = **Viviania**, Cav. (Geran.).
petiolatum, Klotzsch, l. c. 439.—Bras.

LINOSTOMA, Wall. Cat. n. 4203 (1831). *THYMELAEACEAE*, Benth. & Hook. f. iii. 197.
ENKLEIA, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 234.
LOPHOSTOMA, Meissn. in Mart. Fl. Bras. v. 1. 72 (1855).
NECTANDRA, Roxb. Fl. Ind. ii. 425 (1832).
PSILAEA, Miq. Fl. Ind. Bat. Suppl. 355 (1860).
calophylloides, *Meissn. in Mart. Fl. Bras.* v. 1. 72.—Bras.
decandrum, *Wall. Cat.* n. 4203.—Ind. or.; Burma.
ovatum, *Benth. & Hook. f. Gen.* iii. 198.—Venezuela.
pauciflorum, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 234.—Burma; Malaya.
scandens, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 83.—Burma; Malaya.
siamense, Kurz, l. c. 82 = scandens.

LINOSTROPHUM, Schrank, Fl. Prim. Salisb. 163 (1792) = **Camelina**, Crantz (Crucif.).
sativum, Schrank, l. c. = C. sativa.

LINOSTYLIS, Fenzl, ex Sond. in Linnaea, xxiii. (1850) 94 = **Calophanes**, D. Don (Acanth.).
fasciculiflora, Fenzl, ex Sond. l. c. 95.—Afr. trop.
ovata, Sond. l. c. 94 = C. Perrottetii.

LINOSYRIS, Cass. in Dict. Sc. Nat. xxxvii. 460, 476 (1825) = **Aster**, Tourn. (Compos.).
candicans, DC. Prod. v. 352 (= *Varthemia candicans*).—Aegypt.
Capusi, Franch. in Ann. Sc. Nat. Sér. VI. xvi. (1883) 306.—Turkestan.
divaricata, DC. Prod. v. 352 = A. divaricatus?
foliosa, Cass. ex Desf. Cat. Hort. Par. ed. III. 166 = A. Linosyris.
glabrata, DC. Prod. v. 353.—Sibir. alt.
Grimmi, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 615.—Turkestan.
hastata, DC. Prod. v. 352 = Vernonia atriplicifolia.
montana, DC. l. c. (= *Varthemia montana*).—Syria; Arab.
palustris, Nym. Consp. 386 = A. Linosyris.
punctata, DC. Prod. v. 352 = A. acris.
punctata, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 613.—Turkestan..
scoparia, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 378.—Sibir.
tarbagatensis, C. Koch, in Linnaea, xxiii. (1850) 703.—Reg. Caucas.
uniflora, DC. Prod. v. 352.—Aegypt.
villosa, DC. l. c.—Europ. or.; As. bor.
vulgaris, Cass. ex Less. Syn. Comp. 195 = A. Linosyris.

LINOSYRIS, Torr. & Gray, Fl. N. Am. ii. 232 (1840) = **Bigelovia**, DC. (Compos.).
albicaulis, Torr. & Gray, l. c. 234 = B. dracunculoides.
arborescens, A. Gray, in Torr. Bot. Mex. Bound. 79 = B. arborescens.
Bigelovii, A. Gray, in Pacif. Rail. Rep. iv. 98 = B. Bigelovii.
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 354 = B. Bolanderi.
carnosa, A. Gray, Pl. Wright. ii. 80 = Aster carnosus.
ceruminosa, Durand & Hilg. in Pacif. Rail. Rep. v. 9 = B. ceruminosa.
coronopifolia, A. Gray, Pl. Wright. i. 96 = B. coronopifolia.
dentata, Kellogg, in Proc. Calif. Acad. ii. (1863) 16 = B. veneta.
depressa, Torr. in Sitgreave, Rep. 161 = B. depressa.
Drummondii, Torr. & Gray, Fl. N. Am. ii. 233 = B. Drummondii.
graveolens, Torr. & Gray, l. c. 234 = B. dracunculoides.
heterophylla, A. Gray, Pl. Wright. i. 95 = B. Wrightii.
hirtella, A. Gray, l. c. = B. Wrightii.
Howardii, Parry, in Proc. Am. Acad. vi. (1865) 541 = B. Howardi.
humilis, Torr. & Gray, Fl. N. Am. ii. 234 = Aster angustus.
lanceolata, Torr. & Gray, l. c. 233 = B. viscidiflora.
Mandonii, Sch. Bip. in Linnaea, xxxiv. (1865–66) 534, nomen; A. Gray, in Proc. Am. Acad. viii. (1873) 638 = B. hypoleuca.
mexicana, Schlecht. in Linnaea, xiv. (1840) Litt. 128 = B. veneta.
nudata, Bertol. Misc. Bot. vi. 27 = B. nudata.

LINOSYRIS :—
Parryi, A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864) 66 = Bigelovia Parryi.
pluriflora, Torr. & Gray, Fl. N. Am. ii. 233 = B. pluriflora.
pulchella, A. Gray, Pl. Wright. i. 96 = B. pulchella.
pumila, A. Gray, l. c. ii. 80, in nota = B. viscidiflora.
ramulosa, A. Gray, l. c. i. 97; ii. 80 = Baccharis pteronioides.
serrulata, Torr. in Stansb. Exped. Utah, 389 = B. viscidiflora.
sonoriensis, A. Gray, in Proc. Am. Acad. viii. (1870) 291 = B. diffusa.
squamata, A. Gray, l. c. = Lepidospartum squamatum.
teretifolia, Durand & Hilg. in Pacif. Rail. Rep. v. 9 = B. teretifolia.
texana, Torr. & Gray, Fl. N. Am. ii. 232 = Baccharis texana.
viscidiflora, Hook. Lond. Journ. Bot. vi. (1847) 243, partim = B. dracunculoides.
viscidiflora, Torr. & Gray, Fl. N. Am. ii. 234 = B. viscidiflora.
Wrightii, A. Gray, Pl. Wright. i. 95 = B. Wrightii.

LINSCHOTENIA, De Vriese, in Mitch. Journ. Trop. Austral. 345 (1848) = **Dampiera**, R. Br. (Gooden.).
discolor, De Vriese, l. c. 346 = D. Linschotenii.

LINSCHOTTIA, Comm. ex Juss. Gen. 343, in obs. (1789) = **Homalium**, Jacq. (Samyd.).

LINSCOTIA, Adans. Fam. ii. 269 (1763) = **Limeum**, Linn. (Ficoid.).

LINSECOMIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 458 = **Helianthus**, Linn. (Compos.).
glauca, Buckl. l. c. = H. ciliaris.

LINTIBULARIA, Gilib. Fl. Lituan. i. 139 (1781), sphalm. = LENTIBULARIA, Adans. = **Utricularia**, Linn. (Lentib.).

LINUM, Tourn. ex Linn. Syst. ed. I (1735). *LINEAE*, Benth. & Hook. f. i. 242.
ADENOLINUM, Reichb. Handb. 306 (1837).
ALSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349.
CATHARTOLINUM, Reichb. Handb. 306 (1837).
CHRYSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349.
CLIOCOCCA, Bab. in Proc. Linn. Soc. i. (1841) 90; et in Trans. Linn. Soc. xix. (1842) 33. t. 3.
LEUCOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349.
LINOPSIS, Reichb. Handb. 306 (1837).
MEIAPINON, Rafin. Fl. Tellur. iii. 32 (1836).
MESYNIUM, Rafin. l. c. 33 (1836).
NEZERA, Rafin. New Fl. Am. iv. 64 (1836).
NUMISAUREUM, Rafin. Fl. Tellur. iii. 32 (1836).
XANTHOLINUM, Reichb. Handb. 306 (1837).
abyssinicum, Hochst. in Flora, xxiv. (1841) 1. Intell. 30 = gallicum, strictum.
acuminatum, E. Mey. in Drège, Zwei Pfl. Docum. 199 = africanum.
acuminatum, Moench, Meth. 287 = angustifolium, narbonense.
adenophyllum, *A. Gray, in Proc. Am. Acad.* viii. (1873) 624.—Calif.
adustum, *E. Mey. ex Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 492.—Afr. austr.
aethiopicum, Thunb. Prod. Pl. Cap. 57 = africanum.
affine, Panz. ex Schult. Syst. vi. 737 = angustifolium.
africanum, *Linn. Sp. Pl.* 280.—Afr. austr.
agrestre, Brot. Fl. Lusit. i. 481 = angustifolium.
album, *Kotschy, ex Boiss. Diagn.* Ser. I. vi. 27.—Persia.
alpinum, *Linn. Sp. Pl.* ed. II. 1672.—Europ.; Afr. bor.
alpinum, Pall. Tabl. Taur. 49 = austriacum.
alpinum, Schangin, in Pall. N. Nord. Beitr. vi. 47 = perenne.
altaicum, *Fisch. ex Steud. Nom.* ed. II. ii. 51, nomen.—Sibir. altaic.
alterniflorum, Delile, ex Spreng. Syst. iv. Cur. Post. 127 = strictum.
alternum, Pieri, Fl. Corcir. 25 = strictum.
ambiguum, Jord. Cat. Jard. Dijon (1848) 27 = angustifolium.
amurense, *Alef. in Bot. Zeit.* xxv. (1867) 251.—Reg. Amur.

LINUM :—
anatolicum, Boiss. Diagn. Ser. I. i. 56 = hirsutum.
anglicum, Mill. Gard. Dict. ed. VIII. n. 5 = perenne.
angustifolium, DC. Prod. i. 426 = marginale.
angustifolium, *Huds. Fl. Angl.* ed. II. 134.—Europ.; Afr. bor.
angustifolium, Lodd. Bot. Cab. t. 1543 = austriacum.
annotinum, *C. Koch, in Linnaea*, xv. (1841) 713.—Armenia.
annuum, Nees, in Neuwied, Reise Nord-Am. ii. 431 = Berlandieri.
aquilinum, Molina, Sagg. Chil. 150 = Chamissonis, Macraei.
arboreum, *Linn. Sp. Pl.* 279.—Creta.
arboreum, Pall. ex Bieb. Fl. Taur. Cauc. i. 255 = flavum.
aretioides, *Boiss. Diagn.* Ser. I. i. 55.—As. Min.
aristatum, *Engelm. in Wisliz. Tour North Mexico*, 101.—Am. bor. occ.
aroanium, *Boiss. & Orph. Diagn.* Ser. II. i. 96.—Graecia.
arvense, Neck. Delic. Gallo-Belg. i. 159 = usitatissimum.
ascyrifolium, Sims, Bot. Mag. t. 1087 = hirsutum.
asiaticum, *Mart. Hort. Erlang.* 84, nomen.—Hab.?
asperifolium, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 25.—Alger.
Aucheri, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 173 = nervosum.
aureum, DC. Prod. i. 432 = nodiflorum.
aureum, Waldst. & Kit. Pl. Rar. Hung. ii. 193. t. 177 = gallicum.
austriacum, DC. Fl. Fr. Suppl. 615 = alpinum.
austriacum, Lam. Encyc. iii. 521 = diffusum.
austriacum, *Linn. Sp. Pl.* 278.—Europ.; Oriens.
austriacum, Sims, Bot. Mag. t. 1086 = angustifolium, perenne.
Babingtonii, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 179 = selaginoides?
Balansae, *Boiss. Fl. Orient.* i. 855.—As. Min.
barbulatum, Lange, ex Reichb. Fl. Germ. Excurs. 832 = austriacum.
Bartlingii, Eckl. & Zeyh. Enum. 35 = africanum.
bavaricum, Schult. in Flora, xxi. (1838) 643 = alpinum.
Beauharnaisianum, *Spadoni, Lino di Siberia*, t. 1.—Sibir.
Berendieri, vide seq.
Berlandieri, *Hook. Bot. Mag.* t. 3480 (sphalm. Berendieri).—Texas.
betsiliense, *Baker, in Journ. Bot.* xx. (1882) 48.—Madag.
bicolor, *Schousb. ex DC. Prod.* i. 428.—Marocc.
Biebersteini, *Spreng. Gartenz.* iii. 133; *ex Schult. Syst.* vi. 758, nomen.—Reg. Caucas.?
bienne, *Mill. Gard. Dict.* ed. VIII. n. 8.—Istria.
bipunctatum, Bartl. ex Steud. Nom. ed. II. ii. 51 = oligophyllum.
boerhaviaefolium, Hort. ex Steud. l. c. ed. I. 484 = Reinwardtia trigyna.
Bootii, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 475 = sulcatum.
brevifolium, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 30.—Bras.
Breweri, *A. Gray, in Proc. Calif. Acad.* iii. (1863) 102.—Calif.
Broteri, *Hoffmgg. ex Schult. Syst.* vi. 758.—Lusitan.
Bungei, *Boiss. Fl. Orient.* i. 861.—Persia.
caespitosum, Sibth. & Sm. Fl. Graec. Prod. i. 216 = arboreum.
californicum, *Benth. Pl. Hartw.* 299.—Calif.
campanulatum, Bieb. Fl. Taur. Cauc. i. 255 = flavum, tauricum.
campanulatum, *Linn. Sp. Pl.* 280.—Europ.
capitatum, Griseb. ex Planch. in Hook. Lond. Journ. Bot. vii. (1848) 510 = flavum?
capitatum, *Kit. ex Schult. Oesterr. Fl.* ed. II. i. 528, in obs.—Europ. austr.
cariense, *Boiss. Diagn.* Ser. I. v. 86.—Caria.
carneum, *A. St. Hil. Fl. Bras. Mer.* i. 132.—Bras.
carnosulum, *Boiss. Diagn.* Ser. I. viii. 104.—As. Min.
catharticum, *Linn. Sp. Pl.* 281.—Europ.; Oriens.
Chamissonis, *Schiede, in Linnaea*, i. (1826) 69.—Chili.
chilense, *Kostel. Allg. Med.-pharm. Flora*, v. 1912.—Chili.
chironioides, Griseb. in Goett. Abh. vi. (1854) 118 = Macraei.
Cicanobum, Buch.-Ham. in D. Don, Prod. Fl. Nep. 217 = Reinwardtia tetragyna.

LINUM :—
cilicicum, Fenzl, in Tchihat. Asie Min. Bot. i. 141 = tenuifolium.
Clevelandi, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 121.—Calif.
collinum, Guss. Fl. Sic. Syn. ii. 808, in textu = austriacum.
complanatum, Gromow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 159, nomen (Quid?).—Ross.
congestum, *A. Gray, in Proc. Am. Acad.* vi. (1864-65) 521.—Calif.
corymbiferum, *Desf. Fl. Atlant.* i. 279. t. 80.—Afr. bor.
corymbulosum, Reichb. Fl. Germ. Excurs. 834 = strictum.
Coulterianum, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 498.—Mexic.
crepitans, Dum. Fl. Belg. 111 = usitatissimum.
cribrosum, Reichb. Ic. Fl. Germ. vi. 63. t. 330. f. 5158 = angustifolium.
crispum, Steud. Nom. ed. I. 484 = salsoloides.
Cruciata, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 499.—Mexic.
Cumingii, Lodd. Bot. Cab. t. 1996 = Macraei.
cuspidatum, *Rafin. New Fl. Am.* iv. 66 (Quid?).—Am. bor.
darmstadinum, *Alef. in Bot. Zeit.* xxv. (1867) 252 (Quid?).—Europ.
davuricum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 260 = stelleroides.
davuricum, *Schult. Syst.* vi. 752.—Dahuria.
decoloratum, Griseb. Spicil. Fl. Rumel. i. 117 = pubescens.
decumbens, *Desf. Fl. Atlant.* i. 278. t. 78.—Tunis.
decurrens, Kellogg, in Proc. Calif. Acad. iii. (1863) 44. f. 11 = perenne.
depressum, *Hort. ex Schult. Syst.* vi. 758, nomen.—Hab.?
dichotomum, Bout. ex Willk. & Lange, Prod. Fl. Hisp. iii. 546 = tenue.
diffusum, *Schult. Obs. Bot.* 63.—Hab.?
diffusum, Wood, Class-Book (1861) 276 = striatum?
digynum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 334.—Calif.
diversifolium, Gilib. Fl. Lituan. ii. 143 = catharticum.
drymarioides, *Curran, in Bull. Calif. Acad.* i. n. III. (1885) 152.—Calif.
elatum, Salisb. Prod. 177 = perenne.
elegans, *Sprun. ex Boiss. Diagn.* Ser. II. i. 99.—Graecia.
emirnense, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1843) 16.—Madag.
empetrifolium, Schott & Kotschy, in Tchihat. Asie Min. Bot. i. 143 = carnosulum.
erigeroides, *St. Hil. Fl. Bras. Mer.* i. 132.—Reg. Argent.
exiguum, Salisb. Prod. 177 = Radiola linoides.
extraaxillare, *Kit. in Linnaea*, xxxii. (1863) 573.—Bannat.
fasciculare, *Van Geuns, Cat. Turic.* (1802); *ex Schult. Syst.* vi. 758, nomen.—Hab.?
fasciculatum, Mart. ex Steud. Nom. ed. II. ii. 51 = fasciculare.
fastigiatum, *Tausch, in Syll. Ratisb.* ii. (1828) 244.—Europ. austr.
filiforme, *Urban, in Linnaea*, xli. (1877) 643.—Ecuador.
flavum, Habl. Taur. 150; ex Bieb. Fl. Taur. Cauc. i. 255 = campanulatum.
flavum, *Linn. Sp. Pl.* 279.—Europ. austr.; Reg. Caucas.
flexuosum, *Hort. ex Steud. Nom.* ed. II. ii. 51, nomen.—Hab.?
formosum, *Urban, in Mart. Fl. Bras.* xii. II. 460.—Bras.
fruticosum, Mill. Gard. Dict. ed. VIII. n. 11 = suffruticosum.
fruticulosum, Planch. ex Walp. Ann. ii. 128 = thesioides?
gallicum, *Linn. Sp. Pl.* ed. II. 401.—Reg. Mediterr.; Abyssin.
gallicum, Sibth. & Sm. Fl. Graec. iv. 3. t. 303 = strictum.
Galloi, Saroti, ex Parl. Fl. Ital. v. 290 = nodiflorum.
glandulosum, Moench, Meth. 287 = campanulatum.
glaucum, Boiss. & Noé, Diagn. Ser. II. v. 66 = alpinum.
globulariaefolium, Poir. Encyc. Suppl. iii. 445 = arboreum.

LINUM:—
gracile, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 496 = africanum.
gracile, Sm. ex Planch. l. c. 169 = marginale.
grandiflorum, *Desf. Fl. Atlant.* i. 278. t. 78.—Alger.
grandiflorum, Hort. ex Reichb. Pl. Crit. iv. 8 = usitatissimum.
Greggii, *Engelm. ex A. Gray, Pl. Wright.* i. 26.—Mexic.
guatemalense, *Benth. Bot. Voy. Sulph.* 67.—Mexic.
heterophyllum, Moench, Meth. Suppl. 99 = maritimum.
heterosepalum, *Regel, in Act. Hort. Petrop.* ii. (1873) 433.—Turkest.
hirsutum, DC. Fl. Fr. iv. 800 = viscosum.
hirsutum, *Linn. Sp. Pl.* 277.—Europ. austr.; As. Min.
hirsutum, Sibth. & Sm. Fl. Graec. Prod. i. 215 = olympicum.
hirsutum, Sieber, ex Presl, Fl. Sic. 170 = pubescens.
hispanicum, Mill. Gard. Dict. ed. VIII. n. 7 = angustifolium.
Hohenackeri, Boiss. Diagn. Ser. II. i. 97 = angustifolium.
hologynum, *Reichb. Fl. Germ. Excurs.* 833.—Bannat.
hudsonioides, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 186 = multicaule.
humile, Heyne, ex Wall. Cat. n. 1507 = mysorense.
humile, Mill. Gard. Dict. ed. VIII. n. 2 = angustifolium.
hypericifolium, *Presl, Rel. Haenk.* ii. 2.—Mexic.
hypericifolium, Salisb. ex Steud. Nom. ed. I. 484 = viscosum.
iberidifolium, Auch. ex Planch. in Hook. Lond. Journ. Bot. vii. (1848) 515 = elegans.
imbricatum, *Rafin. New Fl. Am.* iv. 66 (Quid?).—Am. bor.
inaequale, Presl, Delic. Prag. 58 = strictum.
junceum, *St. Hil. Fl. Bras. Mer.* i. 134. t. 24.—Bras.
juniperifolium, Eckl. & Zeyh. Enum. 35 = africanum.
Kingii, *S. Wats. in Bot. King's Exped.* 49.—Am. bor. occ.
kirgisicum, *Alef. in Bot. Zeit.* xxv. (1867) 255.—Turkest.
laeve, Reichb. Fl. Germ. Excurs. 833 = narbonense.
laeve, Scop. Fl. Carn. ed. II. i. 230 = alpinum.
lambesanum, *Boiss. & Reut. Diagn.* Ser. II. v. 65.—Alger.
lecheoides, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 332.—Am. bor. occ.
Leonii, Schultz, in Flora, xxi. (1838) 644 = alpinum.
leucanthum, *Boiss. & Sprun. Diagn.* Ser. I. i. 55.—Graecia.
Lewisii, Pursh, Fl. Am. Sept. i. 210 = perenne.
liburnicum, Auct. ex Nym. Consp. 126 = strictum.
liburnicum, Scop. Fl. Carn. ed. II. i. 230 = nodiflorum.
limanense, Lamotte, Prod. Fl. Centr. Fr. i. 157 = austriacum.
littorale, *St. Hil. Fl. Bras. Mer.* i. 133.—Bras.
Loreyi, Jord. Cat. Jard. Dijon (1848) 27 = austriacum.
luteolum, Bieb. Fl. Taur. Cauc. i. 256 = nodiflorum.
Lyallanum, Alef. in Bot. Zeit. xxv. (1867) 251 = perenne.
Macraei, *Benth. in Bot. Reg.* sub t. 1326.—Chili.
Macraei, Hook. Bot. Mag. t. 5474 = Chamissonis.
marginale, *A. Cunn. ex Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 169.—Austral.
marginatum, Poir. Encyc. Suppl. iii. 443 = angustifolium.
marginatum, Reichb. Fl. Germ. Excurs. 832 = austriacum.
marinum, Suffren, Pl. Frivul. 132 = seq.
maritimum, *Linn. Sp. Pl.* 280.—Reg. Mediterr.
mauritanicum, *Pomel, Nouv. Mat. Fl. Atl.* 214.—Alger.
melianthum, Brot. Fl. Lusit. i. 484 = tenue?
mexicanum, *H. B. & K. Nov. Gen. et Sp.* vi. 39.—Mexic.
micranthum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 333.—Calif.
millegranum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 641 = Radiola linoides.
monadelphum, Hort. ex Reichb. Pl. Crit. iv. 8 = usitatissimum.
monadelphum, Mart. Hort. Erlang. 84 = praec.?
monogynum, *Forst. f. Prod.* 23.—N. Zel.
monogynum, Hort. ex Reichb. Fl. Germ. Excurs. 833 = hologynum.

LINUM:—
monopetalum, Steph. Fl. Mosq. n. 214.—Cf. DC. Prod. i. 425 = flavum.
montanum, Lorey & Dur. ex Nym. Consp. 125 = austriacum.
montanum, Schleich. Cat. Pl. Helv. ed. III. 19 = alpinum.
mucronatum, Bertol. Misc. Bot. i. 18 = orientale.
mucronatum, Gilib. Fl. Lituan. ii. 143 = usitatissimum.
Muelleri, *Moris, App. ad Elench. Fl. Sard.* 1.—Sardinia.
multicaule, Hedw. ex Steud. Nom. ed. I. 485 = angustifolium.
multicaule, *Hook. ex Torr. & Gray, Fl. N. Am.* i. 678.—Am. bor. occ.
multiflorum, Lam. Fl. Fr. iii. 70 = Radiola linoides.
Munbyanum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 24.—Alger.
mysorense, *Heyne, ex Wall. Cat.* n. 1507.—Ind. or.
narbonense, Desf. Fl. Atlant. i. 279; Bieb. Fl. Taur. Cauc. i. 254 = angustifolium.
narbonense, Hort. ex Reichb. Pl. Crit. iv. 8 = usitatissimum.
narbonense, *Linn. Sp. Pl.* 278.—Europ. austr.
narbonense, Pall. ex Bieb. Fl. Taur. Cauc. i. 254 = nervosum.
narbonense, Suter, Fl. Helvet. i. 184 = alpinum.
neo-mexicanum, *Greene, in Coult. Bot. Gaz.* vi. (1881) 183.—N. Mexic.
nervosum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 109. t. 105.—Europ. austr.; Oriens.
nodiflorum, *Linn. Sp. Pl.* 280.—Europ. austr.; Oriens.
nutans, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 430.—China.
obtusatum, Boiss. Fl. Orient. i. 865 = alpinum.
oligophyllum, Hook. & Arn. Bot. Beech. Voy. 11 = Macraei.
oligophyllum, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 486 = filiforme.
oligophyllum, *Willd. ex Schult. Syst.* vi. 758.—Am. austr.
olympicum, *Boiss. Diagn.* Ser. I. i. 56.—As. Min.
organense, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 100.—Bras.
orientale, *Boiss. Fl. Orient.* i. 855.—As. Min.
Orizabae, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 482.—Mexic.
Ortegae, *Planch. l. c.* 184.—Hispan.
Pallasianum, Schult. Syst. vi. 758 = tauricum.
pallescens, *Bunge, in Ledeb. Fl. Alt.* i. 438.—Sibir. altaic.
palustre, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 99.—Bras.
pamphylicum, Boiss. & Heldr. ex Planch. in Hook. Lond. Journ. Bot. vii. (1848) 511 = orientale.
paniculatum, Desv. ex Steud. Nom. ed. II. ii. 52 = strictum.
paniculatum, Moench, Meth. Suppl. 98 = narbonense.
paposanum, *Phil. Fl. Atac.* 13.—Chili.
parviflorum, *Medic. Beobacht.* (1783) 145.—Hab.?
perenne, Guss. Fl. Sic. Prod. i. 386 = austriacum.
perenne, Lam. Fl. Fr. iii. 66 = alpinum.
perenne, *Linn. Sp. Pl.* 277.—Reg. bor. temp.
perenne, Vill. Hist. Pl. Dauph. ii. 550 = angustifolium.
persicum, *Boiss. Diagn.* Ser. I. vi. 27.—Persia.
piliferum, Boiss. Fl. Orient. i. 860 = pubescens.
piligerum, Presl, Fl. Sic. i. 170 = pubescens.
Plotzii, Hook. Bot. Mag. sub t. 3480 = Berlandieri.
polygaloides, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 487.—Peruv.
procumbens, Hort. ex Steud. Nom. ed. II. ii. 52; Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 280, nota, t. 965. f. 2 = decumbens.
prostratum, *Domb. ex Lam. Encyc.* iii. 525.—Peruv.
provinciale, Jord. Obs. Pl. Crit. vii. (1849) 17 = austriacum.
pubescens, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 268.—Graecia; As. Min.
pubescens, Willd. ex Schult. Syst. vi. 758 = tauricum.
punctatum, *Presl, Delic. Prag.* 58.—Sicil.
pungens, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 491 = africanum.
pusillum, Pall. Tabl. Taur. 49 = nodiflorum.
pycnophyllum, Boiss. & Heldr. Diagn. Ser. II. i. 97 = alpinum.
pyrenaicum, Pourr. in Mém. Acad. Toul. iii. (1788) 322 = angustifolium.
quadrifolium, *Linn. Sp. Pl.* 281.—Afr. austr.

LINUM:—
quadrifolium, Thunb. Fl. Cap. ed. Schult. 277 = Thunbergii.
racemosum, E. Mey. in Drège, Zwei Pfl. Docum. 199 = Thunbergii.
Radiola, Linn. Sp. Pl. 281 = Radiola linoides.
ramosissimum, *C. Gay, Fl. Chil.* i. 463.—Chili.
ramosissimum, Willk. in Bot. Zeit. v. (1847) 240 = tenue?
reflexum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 387.—Europ. austr.
refractum, *Rafin. New Fl. Am.* iv. 64 (Quid?).—Am. bor.
Reichenbachii, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 495 = Thunbergii.
repens, Buch.-Ham. in D. Don, Prod. Fl. Nep. 217 = Reinwardtia trigyna.
rigidum, *Pursh, Fl. Am. Sept.* i. 210.—Am. bor. occ.
rigidum, Sarato, ex Parl. Fl. Ital. v. 293 = strictum.
rigidum, Torr. & Gray, Fl. N. Am. i. 204 = striatum, sulcatum.
rubrum, Rafin. Caratteri, 74 = decumbens.
rupestre, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 232.—Am. bor. occ.
ruscinonense, Timb. in Bull. Soc. Bot. Fr. vii. (1860) 509 = alpinum.
salsoloides, *Lam. Encyc.* iii. 521.—Gallia.
sansabeanum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 450 = Lechea Drummondii.
sardoum, Muell. ex Moris, Fl. Sard. i. 358 = Muelleri.
sativum, Hasselq. Iter Palaest. 462 = usitatissimum.
saxicola, Jord. Cat. Jard. Dijon (1848), 28 = austriacum.
scabrellum, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 507.—Mexic.
scabridum, Steud. ex Boiss. Fl. Orient. i. 852 = strictum.
scabrum, Kunze, in Flora, xxix. (1846) 654 = tenue.
scariosum, *Scheele, in Flora*, xxvi. (1843) 433.—Europ.
Schiedeanum, *Cham. & Schlecht. in Linnaea*, v. (1830) 234.—Mexic.
scoparium, *Griseb. in Goett. Abh.* xix. (1874) 103.—Bras.
selaginoides, *Lam. Encyc.* iii. 525.—Reg. Argent.
selaginoides, Torr. & Gray, Fl. N. Am. i. 205 = multicaule.
semitrigynum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 217 = Reinwardtia trigyna.
serotinum, *Pall. Reise Suedl. Statth. Russ. Reich.* ii. 80.—Tauria.
serrulatum, Bertol. Fl. Ital. iii. 549 = capitatum.
sessiliflorum, Lam. Encyc. iii. 523 = strictum.
setaceum, *Brot. Fl. Lusit.* i. 484.—Reg. Mediterr. occ.
sibiricum, DC. Prod. i. 427 = perenne.
Sibthorpianum, Margot & Reut. in Mém. Soc. Phys. Genèv. viii. (1839) 283 = pubescens.
siculum, Presl, Fl. Sic. 171 = angustifolium.
simplex, Wood, Class Book (1861) 276; Bot. & Flor. 66 = sulcatum?
Simsii, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 512 = arboreum.
spergulinum, *A. Gray, Proc. Am. Acad.* vii. (1868) 333.—Calif.
spicatum, Lam. Encyc. iii. 523 = strictum.
squamatum, Herb. Unio. It. ex Steud. Nom. ed. II. ii. 52 = seq.
squamulosum, Rudolphi, ex Willd. Enum. Hort. Berol. 338 = austriacum.
squarrosum, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 283.—Alger.
stelleroides, *Planch. in Hook. Journ. Lond. Bot.* vii. (1848) 178.—China.
Stocksianum, Boiss. Diagn. Ser. II. i. 98 = perenne.
striatum, Nutt. Gen. Am. i. 206 = sulcatum.
striatum, *Walt. Fl. Carol.* 118.—Am. bor.
strictissimum, Pall. Reise Suedl. Statth. Russ. Reich. ii. 80 = gallicum?
strictum, *Linn. Sp. Pl.* 279.—Reg. Mediterr.; Oriens.
strictum, Pall. Tabl. Taur. 49 = gallicum.
suaedaefolium, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 168.—Austral.
suffruticosum, DC. Fl. Fr. Suppl. 616 = salsoloides.
suffruticosum, *Linn. Sp. Pl.* 279.—Reg. Mediterr. occ.
suffruticosum, Orteg. ex Planch. in Hook. Lond. Journ. Bot. vii. (1848) 184 = Ortegae.
sulcatum, *Riddel, Cat. Pl. Ohio, Suppl.* 10.—Am. bor.
sulphureum, *Boiss. & Hassk. in Boiss. Fl. Orient.* i. 857.—Syria.

LINUM:—
sylvestre, Scop. Fl. Carn. ed. II. i. 229 = viscosum.
syriacum, *Boiss. & Gaill. ex Boiss. Fl. Orient.* i. 856. —Syria.
tauricum, *Willd. Enum. Hort. Berol.* 339.—Graecia; Reg. Cauc.
tenellum, *Cham. & Schlecht. in Linnaea*, v. (1830) 235.—Mexic.
Teneriffae, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 150.—Teneriffa.
tenue, *Desf. Fl. Atlant.* i. 280. t. 81.—Hispan.; Afr. bor.
tenue, Munby, Fl. Algér. 33 = Munbyanum.
tenuifolium, Asso, Syn. Stirp. Arag. 41 = suffruticosum.
tenuifolium, *Linn. Sp. Pl.* 278.—Europ.; As. Min.
tenuifolium, Schousb. Vextr. Marokk. 149 = setaceum.
tenuifolium, With. Bot. Arr. Brit. Pl. ed. III. ii. 322 = angustifolium.
tetragynum, Colebr. ex Wall. Cat. n. 1506 = Reinwardtia tetragyna.
tetrapetalum, Gilib. Fl. Lituan. ii. 144 = Radiola linoides.
tetraphyllum, Hebenstr. ex Steud. Nom. ed. II. ii. 52 = quadrifolium.
thesioides, *Bartl. in Linnaea*, vii. (1832) 540.—Afr. austr.
Thunbergii, *Eckl. & Zeyh. Enum.* 35.—Afr. austr.
tmoleum, *Boiss. Diagn.* Ser. I. viii. 105.—As. Min.
Tommasinii, Nym. Syll. 218 = austriacum.
toxicum, *Boiss. Fl. Orient.* i. 854.—Syria.
transsilvanicum, Schur, Enum. Pl. Transs. 128 = alpinum.
trigynum, Linn. Sp. Pl. 279 = gallicum.
trigynum, Roxb. in As. Res. vi. (1799) 357 = Reinwardtia trigyna.
trinervium, *Freyn, in Oestr. Bot. Zeitschr.* xxvi. (1876) 228.—Transsilv.
trinervium, Herb. Hohen. ex Hook. f. Fl. Brit. Ind. i. 411 = mysorense.
trinervium, Heyne, ex Roth, Nov. Pl. Sp. 187 = usitatissimum.
trisepalum, Kellogg, in Proc. Calif. Acad. iii. (1863) 42 = Helianthemum scoparium.
ucranicum, *Czernw. ex Gruner, in Bull. Soc. Nat. Mosc.* xli. 1868 (1869) II. 130 in textu.—Ucrania.
uniflorum, *Kit. in Linnaea*, xxxii. (1863) 574.—Hungar.
usitatissimum, Griseb. Spicil. Fl. Rumel. i. 117 (excl. syn.) = angustifolium.
usitatissimum, *Linn. Sp. Pl.* 277.—Europ.; Oriens.
utile, Salisb. Prod. 177 = usitatissimum.
velutinum, *Steud. ex Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 517.—Kurdistan.
venustum, Andr. Bot. Rep. t. 477 = viscosum.
verticillatum, *Linn. Sp. Pl.* 281.—Italia.
violascens, *Bunge, in Ledeb. Fl. Alt.* i. 439.—Sibir. altaic.
virgatum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 379 = corymbiferum.
virgatum, Schousb. Vextr. Marokko, 151 = tenue.
virginianum, *Linn. Sp. Pl.* 279.—Am. bor.
virginianum, Reichb. Hort. Bot. ii. 35 = striatum.
virgultorum, *Boiss. & Heldr. ex Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 172.—As. Min.
viscosum, Habl. Taur. 150; ex Bieb. Fl. Taur. Cauc. i. 253 = hirsutum.
viscosum, *Linn. Sp. Pl.* ed. II. 398.—Europ. austr.

LINZIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26 = **Vernonia**, Schreb. (Compos.).
glabra, Steetz, in Peters, Reise Mossamb. Bot. 353 = V. glabra.
vernonioides, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26 = V. Quartiniana.

LIOYDIA, Neck. Elem. i. 4 (1790) = **Printzia**, Cass. (Compos.).

LIPANDRA, Moq. Chenop. Enum. 19 (1840) = **Chenopodium**, Linn.
atriplicoides, Moq. l. c. = C. polyspermum.

LIPARENA, Poit. ex Leman, in Dict. Sc. Nat. xxvii. 6 (1823) = **Drypetes**, Vahl (Euphorb.).

LIPARENE, Poit. ex Baill. Hist. Pl. v. 248 in syn. (1874) = praec.

LIPARIA, Linn. Mant. ii. 156 (1767). *LEGUMINOSAE*, Benth. & Hook. f. i. 472.
badocana, Blanco, Fl. Filip. ed. I. 597 = Psoralea badocana.
Burchellii, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 443.—Afr. austr.
capitata, Thunb. Prod. Pl. Cap. 124 = Priestleya capitata.
comantha, *Eckl. & Zeyh. Enum.* 164.—Afr. austr.
crassinervia, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 63 = parva?
graminifolia, Linn. Mant. ii. 268 = Priestleya graminifolia.
hirsuta, Moench, Meth. Suppl. 52 = Borbonia trinervia.
hirsuta, Thunb. Prod. Pl. Cap. 124 = Priestleya hirsuta.
hybrida, Hort. ex Steud. Nom. ed. I. 485 = Priestleya hirsuta.
laevigata, Thunb. Prod. Pl. Cap. 123 = Priestleya Thunbergii.
myrtifolia, Thunb. l. c. = Priestleya myrtifolia.
opposita, Linn. Mant. ii. 269 = Rafnia opposita.
parva, *Vog. ex Walp. in Linnaea*, xiii. (1839) 468.—Afr. austr.
sericea, Linn. Mant. ii. 269 = Priestleya sericea.
sphaerica, *Linn. l. c.* 268.—Afr. austr.
tecta, Thunb. Prod. Pl. Cap. 124 = Priestleya tecta.
teres, Thunb. l. c. = Priestleya teres.
tomentosa, Thunb. l. c. = Priestleya tomentosa.
umbellata, Linn. Mant. ii. 269 = Priestleya laevigata.
umbellifera, Thunb. Prod. Pl. Cap. 124 = Priestleya umbellifera.
vestita, Thunb. l. c. = Priestleya vestita.
villosa, Andr. Bot. Rep. t. 382 = Priestleya vestita.
villosa, Linn. Mant. ii. 269 = Priestleya villosa.
villosa, Sieber, ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 446 = Priestleya Thunbergii.
villosa, Thunb. Prod. Pl. Cap. 124 = Priestleya lanceolata.

LIPARIS, Rich. in Mém. Mus. Par. iv. (1818) 43, 52.
ORCHIDEAE, Benth. & Hook. f. iii. 495.
ALISPA, Hoffmgg. [Verz. Orchid. 20 (1842)]; ex Linnaea, xvi. (1842) Litt. 228.
ANDROCHILUS, Liebm. in Bot. Notiser (1844), 101.
CESTICHIS, Thou. Orch. Afr. t. 90 (1822).
DISTICHIS, Thou. ex Lindl. Veg. Kingd. 181 (1847). —Cf. Thou. Orch. Isles Austr. Afr. Tabl. 3me.
DITUILIS, Rafin. Fl. Tellur. iv. 49 (1836).
EMPUSA, Lindl. Bot. Reg. sub n. 825 (1824).
EMPUSARIA, Reichb. Consp. 69 (1828).
EPHIPPIANTHUS, Reichb. f. in F. Schmidt, Reisen Amurl. 180. t. 5 (1868).
GASTROGLOTTIS, Blume, Bijdr. 397 (1825).
IEBINE, Rafin. Fl. Tellur. iv. 39 (1836).
MELAXIS, Steud. Nom. ed. II. ii. 118 (1841), sphalm.
MESOPTERA, Rafin. Fl. Tellur. iv. 49 (1836).
PALIRIS, Dum. Fl. Belg. 134 (1827).
PLATYSTYLIS, Lindl. Gen. et Sp. Orch. 18 (1830).
PSEUDORCHIS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 213 (1821).
STICHORCHIS, Thou. Orch. Isles Austr. Afr. Tabl. 1er. (1822).
STURMIA, Reichb. in Moessl. Handb. ed. II. ii. 1576 (1828); Consp. 69 (1828).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 291.—Abyssin.
acutissima, *Reichb. f. Otia Bot. Hamb.* 37.—Cambod.
affinis, *Lindl. Gen. et Sp. Orch.* 31.—Java.
alata, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 17.—Ind. or.
alata, *Scheidw. in Otto & Dietr. Allg. Gartenz.* x. (1842) 293.—Mexic.
angustifolia, Lindl. Gen. et Sp. Orch. 31 = caespitosa.
arnoglossophylla, *Hemsl. Bot. Centr. Am. Bot.* iii. 212. —Mexic.
atropurpurea, *Lindl. Gen. et Sp. Orch.* 28.—Ind. or.
atropurpurea, Wight, Ic. t. 904 = Wightiana.
auriculata, *Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 203.—Japon.
auriculata, *Reichb. f. in Flora*, lv. (1872) 277.—Ind. or.
aurita, *Ridl. in Forbes, Nat. Wand. East. Archip.* 518.—Ins. Timor.
barbata, *Lindl. Gen. et Sp. Orch.* 27.—Zeylan.
Bernaysii, F. Muell. Fragm. xi. 21 = Microstylis congesta.
bicornis, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 458.—Madag.
bidentata, Griff. Notul. iii. 277 (non Liparidis sp.).—Assam.

LIPARIS:—
biloba, *Wight, Ic.* t. 1633.—Ind. or.
bistriata, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 155.—Burma.
bituberculata, *Lindl. Bot. Reg.* sub t. 882.—Reg. Himal.
bootanensis, *Griff. Notul.* iii. 278.—Reg. Himal.
Boothii, Regel, Ind. Sem. Hort. Petrop. (1864) Suppl. 14 = longipes?
Bowkeri, *Harv. Thes. Cap.* ii. 6.—Afr. austr.
brachyglottis, *Reichb. f. ex Trimen, Cat. Pl. Ceyl.* 87. —Zeylan.
brachystalix, *Reichb. f. in Linnaea*, xli. (1877) 43.—Ecuador.
bracteosa, Lindl. in Wall. Cat. n. 7406 = Dendrochilum bracteosum.
caespitosa, *Lindl. in Bot. Reg.* sub t. 882.—Ins. Mascar.
campestris, *Rodrig. Orch. Nov.* i. 36.—Bras.
campylostalix, *Reichb. f. in Linnaea*, xli. (1877) 45. —Ind. or.
capensis, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 314.—Afr. austr.
chlororantha, *Hance, in Journ. Bot.* xxi. (1883) 231.—China.
Clypeolum, *Lindl. Gen. et Sp. Orch.* 29.—Ins. Tahiti.
coelogynoides, *F. Muell. Fragm.* ii. 71.—Austral.
compressa, *Lindl. Gen. et Sp. Orch.* 32.—Malaya.
condylobulbon, *Reichb. f. in Hamb. Gartenz.* xviii. (1862) 34.—Burma.
connata, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 462.—Madag.
Correana, Spreng. Syst. iii. 740 = Loeselii.
crenulata, *Lindl. Gen. et Sp. Orch.* 30.—Java.
crispifolia, *Reich. f. in Linnaea*, xli. (1877) 43.—Peruv.
cuneilabris, F. Muell. Fragm. iv. 164 = reflexa.
decursiva, Reichb. f. in Gard. Chron. (1884) II. 38 = reflexa.
Dendrochilum, *Reichb. f. in Flora*, lv. (1872) 278.—Ind. or.?
densiflora, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 18 = Microstylis versicolor.
diodon, *Reichb. f. in Linnaea*, xli. (1877) 43.—Reg. Himal.
disepala, *Reichb. f. l. c.* 97.—Caled.
disticha, *Lindl. in Bot. Reg.* sub t. 882.—Malaya.
Eggersii, *Reichb. f, in Ber. Deutsch. Bot. Ges.* iii. (1885) 278.—Ind. occ.
elata, *Lindl. Bot. Reg.* t. 1175.—Ind. occ.; Bras.
elegans, *Lindl. in Wall. Cat.* n. 1943.—Malaya.
elliptica, *Reichb. f. in Walp. Ann.* vi. 218.—Ind. occ.; Mexic.
elliptica, Wight, Ic. t. 1735 = viridiflora.
ferruginea, *Lindl. in Gard. Chron.* (1848) 55.—Malaya.
flaccida, *Reichb. f. in Linnaea*, xli. (1877) 45.—Malaya.
flavescens, *Lindl. in Bot. Reg.* sub t. 882.—Ins. Mascar.
foliosa, Lindl. l. c. t. 882 = reflexa.
formosana, Reichb. f. in Gard. Chron. (1880) I. 394 = bituberculata.
Galeottiana, Hemsl. l. c. (1879) I. 559 = elliptica.
Gerrardii, *Reichb. f. in Flora*, l. (1867) 118.—Afr. austr.
Glossula, *Reichb. f. in Linnaea*, xli. (1877) 44.—Reg. Himal.
grandiflora, *Ridl. in Journ. Bot.* xxii. (1884) 333.—Borneo.
gregaria, Lindl. Gen. et Sp. Orch. 33 = disticha.
grossa, *Reichb. f. in Gard. Chron.* (1883) I. 110.—Burma.
guineensis, *Lindl. Bot. Reg.* t. 1671.—Afr. trop.
Guingangae, *Reichb. f. in Flora*, l. (1867) 103.—Afr. trop.
habenarina, *F. Muell. Fragm.* iv. 131.—Austral.
hawaiensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 207.—Ins. Sandvic.
intermedia, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 17 = Microstylis versicolor, *Lindl.*
jamaicensis, Lindl. ex Griseb. Cat. Pl. Cub. 261 = elliptica.
Jovis-pluvii, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 154.—Burma.
Kappleri, *Reichb. f. in Walp. Ann.* vi. 218.—Guian.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 509.—Japon.
labiata, Spreng. Syst. iii. 741 = Leochilus cochlearis.
latifolia, *Lindl. Gen. et Sp. Orch.* 30.—Java.
liliifolia, *A. Rich. ex Lindl. in Bot. Reg.* sub t. 882. —Am. bor.

LIPARIS:—
Lindeniana, Hemsl. in Gard. Chron. (1879) I. 559 = elliptica.
livida, Royle, Illustr. Bot. Himal. 364, nomen = bituberculata?
Loeselii, *A. Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ.
longa, *Reichb. f. in Linnaea*, xli. (1877) 98.—Hab.?
longicaulis, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 461.—Madag.
longipes, *Lindl. Gen. et Sp. Orch.* 30.—As. trop.; Ins. Pacif.
longipetala, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 459.—Madag.
lutea, *Ridl. l. c.* 458.—Madag.
luteola, *Lindl. in Wall. Cat.* n. 1944; *Gen. et Sp. Orch.* 32.—Reg. Himal.; Burma.
Mannii, *Reichb. f. in Flora*, lv. (1872) 275.—Reg. Himal.
minima, *Lindl. Gen. et Sp. Orch.* 32.—Java.
monophylla, Spreng. Syst. iii. 741 = Epidendrum monophyllum?
montana, *Lindl. Gen. et Sp. Orch.* 29.—Java.
mucronata, Lindl. l. c. 32 = disticha.
nepalensis, *Lindl. in Bot. Reg.* sub t. 882.—Reg. Himal.
nervosa, *Lindl. Gen. et Sp. Orch.* 26.—Japon; China.
nesophila, Reichb. f. Otia Bot. Hamb. 56 = longipes?
neuroglossa, *Reichb. f. Xenia Orch.* iii. 26.—Bolivia.
obcordata, *Vahl, ex King, in Atkins. Gaz. N. W. Ind.* 318, nomen.—Ind. or.
ochracea, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 461.—Madag.
odontostoma, *Reichb. f. in Linnaea*, xli. (1877) 97.—Reg. Himal.
odorata, *Lindl. Gen. et Sp. Orch.* 26.—As. trop. et subtrop.
olivacea, *Lindl. in Wall. Cat.* n. 1942; *Gen. et Sp. Orch.* 27.—Reg. Himal.
olivacea, Wight, Ic. t. 903 = atropurpurea.
orbicularis, Lodd. ex Hemsl. in Gard. Chron. (1881) II. 592 = plantaginea.
ornithorrhynchos, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 460.—Madag.
pachypus, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 155.—Burma.
pallida, *Lindl. Gen. et Sp. Orch.* 31.—Java.
Pappei, *Lindl. ex Harv. Thes. Cap.* ii. 7.—Afr. austr.
paradoxa, Reichb. f. in Walp. Ann. vi. 218 = odorata.
parva, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 462.—Madag.
parviflora, *Lindl. Gen. et Sp. Orch.* 31.—Java.
pendula, Lindl. Bot. Reg. (1838) Misc. 94 = longipes.
plantaginea, *Lindl. Gen. et Sp. Orch.* 29.—Reg. Himal.
plicata, *Franch. & Sav. Enum. Pl. Jap.* ii. 509.—Japon.
polycardia, *Reichb. f. in Flora*, lxviii. (1885) 543.—Afr. austr.
priochilus, Lodd. Bot. Cab. t. 1751 = Microstylis versicolor, *Lindl.*
purpurascens, *Lindl. in Bot. Reg.* sub t. 882.—Ins. Mascar.
ramosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 9. t. 112.—Peruv.
reflexa, *Lindl. in Bot. Reg.* sub t. 882.—Austral.
revoluta, *Hook. & Arn. Bot. Beech. Voy.* 70. t. 16.—Ins. Societ.
Rheedii, *Lindl. Gen. et Sp. Orch.* 26.—Java.
rostrata, *Reichb. f. in Linnaea*, xli. (1877) 44.—Reg. Himal.
rupestris, *Griff. Notul.* iii. 276.—Reg. Himal.
Saundersiana, *Reichb. f. in Gard. Chron.* (1872) 1003.—Jamaica.
selligera, Reichb. f. in Linnaea, xli. (1877) 42 = plantaginea.
serraeformis, Lindl. in Wall. Cat. n. 1946 = Sarcochilus arachnites.
spathulata, Lindl. Bot. Reg. (1840) Misc. 81 = longipes.
stachyurus, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
stenoglossa, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 154.—Burma.
Stricklandiana, *Reichb. f. in Gard. Chron.* (1880) I. 232.—Reg. Himal.?
tradescantifolia, *Lindl. Gen. et Sp. Orch.* 29.—Java.

LIPARIS:—
tricallosa, *Reichb. f. in Gard. Chron.* (1879) I. 684.—Borneo.
tristis, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 579, nomen.—Zeylan.
vestita, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
viridiflora, *Lindl. Gen. et Sp. Orch.* 31.—Ind. or.; Malaya.
viridipurpurea, *Griseb. Cat. Pl. Cub.* 261.—Cuba.
Wageneri, *Reichb. f. in Walp. Ann.* vi. 218.—N. Granat.
Walkeriae, *R. Grah. in Edinb. N. Phil. Journ.* (Jan. 1836) 194.—Ind. or.; Zeylan.
Walkeriae, Wight, Ic. t. 905 = atropurpurea.
Welwitschii, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. trop.
Wendlandi, *Reichb. f. Beitr. Orch. Centr. Am.* 98.—Costa Rica.
Wightiana, *Thw. Enum. Pl. Zeyl.* 295.—Zeylan.
Wightii, Reichb. f. in Walp. Ann. vi. 218 = viridiflora.
zeylanica, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 579, nomen.—Zeylan.

LIPAROPHYLLUM, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 472. *GENTIANEAE*, Benth. & Hook. f. ii. 820.
Gunnii, *Hook. f. l. c.* 473.—Tasmania.

LIPEOCERCIS, Nees, Fl. Afr. Austr. 97 (1841), sphalm. = LEPEOCERCIS, Trin. = **Andropogon**, Linn. (Gramin.).

LIPHONOGLOSSA, Torr. in Bot. Mex. Bound. 124, err. typ. = **Siphonoglossa**, Oerst. (Acanthac.).

LIPOCARPHA, R. Br. in Tuckey, Congo, 459 (1818).
CYPERACEAE, Benth. & Hook. f. iii. 1054.
HYPAELYPTUM, Vahl, Enum. ii. 283 (1806), partim.
TUNGA, Roxb. Fl. Ind. ed. Carey, i. 183, partim (1820).
albiceps, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 163.—Afr. trop.
argentea, *R. Br. in Tuckey, Congo*, 477.—Geront. trop.
atra, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 162.—Afr. trop.
cochleata, Griseb. Cat. Pl. Cub. 241 = Humboldtiana.
filiformis, *Kunth, Enum. Pl.* ii. 267.—Afr. trop.
foliosa, Miq. Fl. Ind. Bat. iii. 332 = Remirea maritima.
glomerata, Nees, in Mart. Fl. Bras. ii. I. 64 = Sellowiana.
gracilis, Nees, in Linnaea, ix. (1834) 287; et in Mart. Fl. Bras. ii. I. 64 = sphacelata, Humboldtiana.
Humboldtiana, *Nees, in Linnaea*, ix. (1834) 287.—Am. trop.
laevigata, Nees, in Wight, Contrib. 92 = argentea.
maculata, Torr. in Ann. Lyc. N. York, iii. (1836) 288 = Humboldtiana.
mexicana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 235 = sphacelata.
microcephala, *Kunth, Enum. Pl.* ii. 268.—As. or.; Austral.
Prieuriana, *Steud. Syn. Pl. Cyp.* 130.—Afr. trop.
pulcherrima, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 162.—Afr. trop.
purpureo-lutea, *Ridl. l. c.* 163.—Afr. trop.
Salzmanniana, Steud. Syn. Pl. Cyp. 129 = Humboldtiana.
Schweinfurthiana, Boeck. in Flora, lxii. (1879) 567 = Prieuriana.
Sellowiana, *Kunth, Enum. Pl.* ii. 267.—Am. austr.
sphacelata, *Kunth, l. c.*—Amphig. trop.
triceps, Nees, in Wight, Contrib. 92 = sphacelata.
Zollingeriana, Boeck. in Flora, xlii. (1859) 100 = microcephala.

LIPOCHAETA, DC. Prod. v. 610 (1836), partim.
COMPOSITAE, Benth. & Hook. f. ii. 372.
APHANOPAPPUS, Endl. Gen. Suppl. ii. 43 (1842).
LIPOTRICHE, Less. in Linnaea, vi. (1831) 408, 510.
MACRAEA, Hook. f. in Proc. Linn. Soc. i. (1845) 278.
MICROCHAETA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 450.
SCHIZOPHYLLUM, Nutt. l. c. 452 (1841).
TRIGONOPTERUM, Steetz, in Vet. Akad. Handl. Stockh. (1853) 161.

LIPOCHAETA:—
amazonica, Poepp. & Endl. Nov. Gen. et Sp. iii. 49. t. 256 = Blainvillea amazonica.
australis, A. Gray, in Proc. Am. Acad. v. (1861) 129 = connata, lobata.
calycosa, *A. Gray, l. c.* 130.—Ins. Sandvic.
cavarum, Walp. Rep. ii. 620, sphalm. = lavarum.
connata, *DC. Prod.* v. 611.—Ins. Sandvic.
costaricensis, Benth. & Hook. f. Gen. ii. 373, sphalm. = Zexmenia costaricensis.
fasciculata, DC. Prod. v. 610 = Zexmenia fasciculata.
goyazensis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 406 = Zexmenia goyazensis.
hastata, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 106.—Calif.
hastulata, *DC. Prod.* v. 611.—Ins. Sandvic.
hetrophylla, A. Gray, in Proc. Am. Acad. v. (1861) 130 = lobata.
integrifolia, *A. Gray, l. c.*—Ins. Sandvic.
Lahainae, Wawra, in Flora, lvi. (1873) 77 = lavarum.
lantanifolia, Schau. in Linnaea, xix. (1847) 729 = Zexmenia lantanifolia.
lavarum, *DC. Prod.* v. 611.—Ins. Sandvic.
lobata, *DC. l. c.*—Ins. Sandvic.
longipes, Benth. & Hook. f. Gen. ii. 373, sphalm. = Zexmenia longipes.
macrocephala, Hook. & Arn. Bot. Beech. Voy. 436 = Zexmenia macrocephala.
micrantha, *A. Gray, in Proc. Am. Acad.* v. (1861) 131.—Ins. Sandvic.
monocephala, DC. Prod. v. 610 = Zexmenia monocephala.
Remyi, *A. Gray, in Proc. Am. Acad.* v. (1861) 131.—Ins. Sandvic.
scaberrima, Benth. in Hook. Journ. Bot. ii. (1840) 43 = Zexmenia scaberrima.
serrata, DC. Prod. v. 611 = Zexmenia serrata.
strigosa, DC. l. c. 610 = Zexmenia strigosa.
subcordata, *A. Gray, in Proc. Am. Acad.* v. (1861) 130.—Ins. Sandvic.
succulenta, *DC. Prod.* v. 611.—Ins. Sandvic.
tagetiflora, G. Don, in Sweet, Hort. Brit. ed. III. 360 = Zexmenia tagetiflora.
tenuifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 131.—Ins. Sandvic.
texana, Torr. & Gray, Fl. N. Am. ii. 357 = Zexmenia hispida.
umbellata, DC. Prod. v. 610 = Zexmenia ceanothifolia.
umbellata, Hook. & Arn. Bot. Beech. Voy. 436 = Zexmenia microcephala.

LIPOCHAETE, Benth. in Hook. Journ. Bot. ii. (1840) 43 = praec.

LIPONEURON, Schott, Nym. & Kotschy, Analect. Bot. 55 (1854) = **Lychnis**, Linn. (Caryoph.).

LIPOPHRAGMA, Schott & Kotschy, ex Boiss. Diagn. Ser. II. v. 42 (1856) = **Moriera**, Boiss. (Crucif.).
eunomioides, Schott & Kotschy, ex Boiss. l. c. (= *Crenularia eunomioides*).—Cilicia.

LIPOPHYLLUM, Miers, in Trans. Linn. Soc. xxi. (1855) 251. t. 26 = **Clusia**, Linn. (Guttif.).
latum, Miers, l. c. 257.—Am. trop.

LIPOSTOMA, D. Don, in Edinb. N. Phil. Journ. (Jan.-Apl. 1830) 168 = **Coccocypselum**, R. Br. (Rubiac.).
campanulaeflora, Steud. Nom. ed. II. ii. 54 = C. campanuliflorum.
capitatum, D. Don, in Edinb. N. Phil. Journ. (Jan.-Apl. 1830) 168 = C. campanuliflorum.
sericeum, D. Don, l. c. 169.—Bras.

LIPOTACTES, Blume, in Schult. f. Syst. vii. 1729 (1830) = **Loranthus**, Linn.
pauciflora, Blume, l. c. = Loranthus pauciflorus.

LIPOTRICHE, R. Br. in Trans. Linn. Soc. xii. (1817) 118 = **Melanthera**, Rohr (Compos.).
Brownei, DC. Prod. v. 544 = M. Brownei.

LIPOTRICHE, Less. in Linnaea, vi. (1831) 408, 510 = **Lipochaeta**, DC. (Compos.).
australis, Less. l. c. 510 = Lipochaeta lobata.
gymnolomoides, Less. l. c. 408 = Perymenium gymnolomoides.

LIPOZYGIS, E. Mey. Comm. Pl. Afr. Austr. 76 (1835) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
argentea, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 80.—Afr. austr.
brachyloba, E. Mey. Comm. Pl. Afr. Austr. 78 = Loton. brachyloba.
calycina, E. Mey. l. c. = Loton. calycina.
carinata, E. Mey. l. c. 80 = Loton. carinata.
corymbosa, E. Mey. l. c. 79 = Loton. corymbosa.
erubescens, E. Mey. l. c. 76 = Loton. pumila.
falcata, E. Mey. l. c. 78 = Loton. falcata.
humifusa, E. Mey. l. c. 77 = Loton. humifusa.
involucrata, E. Mey. l. c. 80 = Loton. involucrata.
Kraussiana, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 79 = Loton. versicolor.
mollis, E. Mey. Comm. Pl. Afr. Austr. 79 = Loton. mollis.
peduncularis, E. Mey. l. c. = Loton. peduncularis.
pentaphylla, E. Mey. l. c. = Loton. pentaphylla.
polycephala, E. Mey. l. c. = Loton. polycephala.
quinata, E. Mey. l. c. 77 = Loton. quinata.
Radula, E. Mey. l. c. = Loton. humifusa.
tenella, E. Mey. l. c. 78 = Loton. tenella.
umbellata, E. Mey. l. c. 76 = Loton. umbellata.
villosa, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 609 = Loton. mollis.

LIPPAYA, Endl. Atakt. 13. t. 13 (1833) = **Dentella**, Forst. (Rubiac.).
telephioides, Endl. l. c. = D. repens.

LIPPIA, Houst. ex Linn. Gen. ed. I. 347 (1737). *VERBENACEAE*, Benth. & Hook. ii. 1142.
ACANTHOLIPPIA, Griseb. in Goett. Abh. xix. (1874) 244.
ALOYSIA, Ort. & Palau, ex L'Hérit. Stirp. i. 21 (1784).
CRYPTOCALYX, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 446.
DIPTEROCALYX, Cham. in Linnaea, vii. (1831) 241.
PANOPE, Rafin. Fl. Tellur. ii. 103 (1836).
PHYLA, Lour. Fl. Cochinch. 66 (1790).
PIARIMULA, Rafin. Fl. Tellur. ii. 102 (1836).
PLATONIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
ZAPANIA, Lam. Illustr. t. 17, textu, n. 42 (1791).
acuminata, *Wright, ex Griseb. Cat. Pl. Cub.* 215.—Cuba.
acutidens, *Mart. & Schau. in DC. Prod.* xi. 590.—Bras.
adoënsis, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 23, nomen; *Schau. l. c.* 578.—Abyssin.
aegyptiaca, *Carr. in Rev. Hortic.* (1872) 316.—Aegypt.
affinis, *Schau. in DC. Prod.* xi. 576.—Bras.
alnifolia, *Schau. l. c.* 588.—Bras.
americana, *Linn. Sp. Pl.* 633.—Mexic.
angustifolia, *Cham. in Linnaea*, vii. (1832) 377.—Bras.
argyrophylla, *Schau. in DC. Prod.* xi. 581.—Hab.?
aristata, *Schau. l. c.*—Bras.
asperifolia, Poepp. ex Cham. in Linnaea, vii. (1832) 215 = geminata.
asperifolia, Reichb. Ic. Exot. Bot. ii. 27. t. 169 = dulcis.
asperifolia, *Rich. Cat. Hort. Med. Par.* 67.—Am. austr.
asperrima, *Cham. in Linnaea*, vii. (1832) 215.—Bras.
attenuata, Mart. in Flora, xxi. (1838) ii. Beibl. 59 = vernonioides.
Berlandieri, *Schau. in DC. Prod.* xi. 575.—Mexic.
Berlandieri, Torr. Bot. Mex. Bound. 127 = graveolens.
Berterii, *Spreng. Syst.* ii. 752.—Bras.
betulaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 264.—Venezuela.
bicolor, Kunth & Bouché, Ind. Sem. Hort. Berol. (1845) 10 = callicarpaefolia.
bicolor, *Mart. & Schau. in DC. Prod.* xi. 587.—Bras.
bracteata, *Hort. ex Rev. Hortic.* (1881) 380.—Hab.?
caffra, *Sond. in Linnaea*, xxiii. (1850) 88.—Afr. austr.
callicarpaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 268.—Mexic.
calocephala, Zucc. Del. Sem. Hort. Monac. (1846) 4 = callicarpaefolia.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 263.—Peruv.
capensis, Spreng. Syst. ii. 751 = asperifolia.
cardiostegia, *Benth. Bot. Voy. Sulph.* 153.—Mexic.

LIPPIA:—
chamaedrifolia, *Steud. Nom.* ed. II. ii. 54.—Bras.
Chamissonis, D. Dietr. Syn. Pl. iii. 598 = Lantana Chamissonis.
chilensis, *Schau. in DC. Prod.* xi. 573.—Chili.
citrata, Willd. ex Cham. in Linnaea, vii. (1832) 214 = geminata.
citriodora, *H. B. & K. Nov. Gen. et Sp.* ii. 269.—Am. austr.
Claussenii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 204.—Bras.
cordata, *Turcz. l. c.* 205.—Bras.
corymbosa, *Cham. in Linnaea*, vii. (1832) 219.—Bras.
cuneifolia, *Steud. Nom.* ed. II. ii. 54.—N. Mexic.
cuneifolia, *Zipp. ex Span. in Linnaea*, xv. (1841) 330, nomen.—Ins. Timor.
cylindrica, Scheele, in Linnaea, xvii. (1843) 351 = Stachytarpheta cayennensis.
cymosa, Sw. Prod. Veg. Ind Occ. 93 (Rubiacea).—Jamaic.
densispicata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 12.—Am. austr.
deserticola, *Phil. in Anal. Univ. Chil.* (1865) ii. 350.—Chili.
dracocephaloides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 205.—Bras.
dulcis, *Trevir. in Nov. Act. Nat. Cur.* xiii. (1826) 187.—Cuba.
Echinus, Spreng. Syst. iv. Cur. Post. 351 = Hyptis Sideritis.
elegans, Cham. in Linnaea, vii. (1832) 225 = origanoides.
elliptica, *Schau. in DC. Prod.* xi. 593.—Bras.
eupatorioidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = seq.
Eupatorium, *Schau. in DC. Prod.* xi. 592.—Bras.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* ii. 265.—Peruv.
filifolia, *Mart. & Schau. in DC. Prod.* xi. 586.—Bras.
filiformis, Schrad. Ind. Sem. Hort. Gotting. (1834); ex Schau. l. c. 585 = canescens.
floribunda, H. B. & K. Nov. Gen. et Sp. ii. 267 = americana.
florida, *Cham. in Linnaea*, vii. (1832) 221.—Bras.
foliolosa, *Phil. Sert. Mendoc. Alt.* 34.—Chili.
fragrans, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 203.—Chili.
Gardneriana, *Schau. in DC. Prod.* xi. 592.—Bras.
geminata, *H. B. & K. Nov. Gen. et Sp.* ii. 266.—Am. bor. et austr.
glabrescens, Meissn. ex Walp. Rep. iv. 56 = hirta.
glandulosa, *Schau. in DC. Prod.* xi. 577.—Bras.
gracilis, *Schau. l. c.* 576.—Bras.
grandiflora, *Mart. & Schau. l. c.* 591.—Bras.
grandifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 167.—Abyss.
grata, *Schau. in DC. Prod.* 576.—Bras.
graveolens, *H. B. & K. Nov. Gen. et Sp.* ii. 267.—Am. bor. et austr.
guayaquilensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 202.—Ecuador.
hastulata, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 407.—Reg. Argent.
havanensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 202.—Cuba.
hederaefolia, *Mart. & Schau. in DC. Prod.* xi. 593.—Bras.
hemisphaerica, *Jacq. Enum. Pl. Carib.* 25; *Select. Am.* 176.—Am. austr.
herbacea, *Mart. ex Schau. in DC. Prod.* xi. 589.—Bras.
hermannioides, *Cham. in Linnaea*, vii. (1832) 219.—Bras.
hieracifolia, *Cham. l. c.* 375.—Bras.
hirsuta, *Linn. f. Suppl.* 288.—N. Granat.
hirta, *Meissn. ex Walp. Rep.* iv. 56; *Schau. in DC. Prod.* xi. 579.—Bras.
hyptoides, Benth. Pl. Hartw. 122 = americana.
integrifolia, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 406.—Reg. Argent.
intermedia, *Cham. in Linnaea*, vii. (1832) 378.—Bras.
iodophylla, *Schau. in DC. Prod.* xi. 588.—Bras.
iresinoides, *Griseb. in Goett. Abh.* xxiv. (1879) 278.—Reg. Argent.
javanica, *Spreng. Syst.* ii. 752.—Java.
juncea, Schau. in DC. Prod. xi. 573 = Baillonia juncea.
kituiensis, *Vatke, in Linnaea*, xliii. (1880–82) 528.—Afr. trop.
lacunosa, *Mart. & Schau. in DC. Prod.* xi. 590.—Bras.

LIPPIA:—
lanata, *Walp. Rep.* iv. 43.—Mexic.
lanceolata, *Michx. Fl. Bor. Am.* ii. 15.—Am. bor.
lantanifolia, *Griseb. in Goett. Abh.* xix. (1874) 242.—Reg. Argent.
lantanifolia, *F. Muell. Fragm.* vi. 151.—Austral.
lasiocalycina, *Cham. in Linnaea*, vii. (1832) 231.—Bras.
ligustrifolia, *Thuret, ex Decne. in Fl. des Serres*, xxii. (1877) 11.—Hab.?
linearis, *H. B. & K. Nov. Gen. et Sp.* ii. 262.—N. Granat.
lupulina, *Cham. in Linnaea*, vii. (1832) 222.—Bras.
lycioides, *Steud. Nom.* ed. II. ii. 54.—Am. bor. et austr.
macrophylla, Cham. in Linnaea, vii. (1832) 213 = Lantana macrophylla.
macrostachya, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 134.—N. Mexic.
marrubiifolia, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxxiii. (1884) 322.—Bras.
Martiana, *Schau. in DC. Prod.* xi. 578.—Bras.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 231.—Bras.
microcephala, *Cham. in Linnaea*, vii. (1832) 220.—Bras.
micromera, *Schau. in DC. Prod.* xi. 587.—Ind. occ.
microphylla, Benth. in Hook. Journ. Bot. ii. (1840) 52 = Schomburgkiana.
microphylla, *Cham. in Linnaea*, vii. (1832) 226.—Bras.
microphylla, Phil. in Anal. Univ. Chil. (1865) ii. 350 = deserticola.
montevidensis, Spreng. Syst. ii. 751 = Lantana Sellowiana.
Moritzii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 204.—Venezuela.
myriocephala, *Schlecht. & Cham. in Linnaea*, v. (1830) 98.—Mexic.
myrtifolia, *Griseb. Cat. Pl. Cub.* 215.—Cuba.
nana, *Schau. in DC. Prod.* xi. 582.—Bras.
nepetacea, *Schau. l. c.* 577.—Bras.
nodiflora, Cham. in Linnaea, vii. (1832) 213 = canescens.
nodiflora, *Michx. Fl. Bor. Am.* ii. 15.—Amphig. trop.
oligophylla, Baker, in Journ. Linn. Soc. xx. (1883) 225 = Acrocephalus villosus.
origanoides, *H. B. & K. Nov. Gen. et Sp.* ii. 267.—Am. trop.
ovata, Linn. Mant. i. 89 = Microdon ovatus.
oxycnemis, *Schau. in DC. Prod.* xi. 589.—Bras.
pallescens, Benth. Pl. Hartw. 245 = Lantana canescens.
panamensis, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 201 = geminata.
pauciserrata, *Turcz. l. c.* 203.—N. Granat.
peruviana, *Turcz. l. c.* 200.—Peruv.
Pohliana, *Schau. in DC. Prod.* xi. 577.—Bras.
polygalaefolia, *Steud. Nom.* ed. II. ii. 54.—Bras.
polystachya, *Griseb. in Goett. Abh.* xix. (1874) 242.—Reg. Argent.
Pseudo-thea, *Schau. in DC. Prod.* xi. 582.—Bras.
pumila, *Cham. in Linnaea*, vii. (1832) 218.—Bras.
purpurea, Dum.-Cours. Bot. Cult. ed. II. vii. 131 = Lantana trifolia.
purpurea, Jacq. f. Eclog. i. 126. t. 85 = Lantana purpurea, *Benth. & Hook. f.*
pyramidata, Crantz, Inst. i. 546 = americana.
queretarensis, H. B. & K. Nov. Gen. et Sp. ii. 264 = lanceolata.
renifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 204.—Bras.
repens, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 446 = canescens.
repens, Spreng. Syst. ii. 752 = nodiflora.
reptans, *H. B. & K. Nov. Gen. et Sp.* ii. 263.—Ind. occ.
rhodocnemis, *Mart. & Schau. in DC. Prod.* xi. 592.—Bras.
Riedeliana, *Schau. l. c.* 582.—Bras.
rigida, *Schau. l. c.* 577.—Bras.
rosmarinifolia, *Andersson. in Vet. Akad. Handl. Stockh.* 1853 (1855) 198.—Ins. Galap.
rotundifolia, *Cham. in Linnaea*, vii. (1832) 230.—Bras.
rubiginosa, Gill. ex Ball, in Journ. Linn. Soc. xxi. (1884) 230 = seriphioides.
rubiginosa, *Schau. in DC. Prod.* xi. 574.—Bras.
salicifolia, *Andersson. in Vet. Akad. Handl. Stockh.* 1853 (1855) 198.—Ins. Galap.
salsa, *Griseb. in Goett. Abh.* xix. (1874) 243.—Reg. Argent.
salsoloides, *Benth. & Hook. f. Gen.* ii. 1143.—Reg. Argent.
salviaefolia, *Cham. in Linnaea*, vii. (1832) 227.—Bras.

LIPPIA:—
sarmentosa, Spreng. Syst. ii. 752 = nodiflora.
satureiaefolia, *Mart. & Schau. in DC. Prod.* xi. 586.—Bras.
scaberrima, *Sond. in Linnaea*, xxiii. (1850) 87.—Afr. austr.
scabra, Hochst. in Flora, xxviii. (1845) 68 = asperifolia.
Schaueriana, *Mart. ex Schau. in DC. Prod.* xi. 590.—Bras.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 23 = Lantana viburnoides.
Schimperi, Walp. Rep. iv. 53 = adoënsis.
Schlimi, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 204.—N. Granat.
Schomburgkiana, *Schau. in DC. Prod.* xi. 577.—Guiana.
scirpea, Phil. in Linnaea, xxxiii. (1864–65) 196 (= *Diostea scirpea*).—Chili.
scorodonioides, *H. B. & K. Nov. Gen. et Sp.* ii. 269.—Ecuador.
sericea, *Cham. in Linnaea*, vii. (1832) 228.—Bras.
seriphioides, *A. Gray, in Proc. Am. Acad.* vi. (1862) 49.—Patagon.
sidoides, *Cham. in Linnaea*, vii. (1832) 224.—Bras.
somalensis, *Vatke, l. c.* xliii. (1880–82) 527.—Afr. trop.
sphacelifolia, Benth. Pl. Hartw. 245 = hirsuta.
stachyoides, *Cham. in Linnaea*, vii. (1832) 227.—Bras.
stoechadifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 265.—Ind. occ.
strigulosa, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 319 = reptans.
substrigosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 202.—Mexic.
thymoides, *Mart. & Schau. in DC. Prod.* xi. 586.—Bras.
trifida, *C. Gay, Fl. Chil.* v. 29.—Chili.
trifida, Phil. Fl. Atac. 40 = deserticola.
triplinervis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 133 = microcephala.
turbinata, *Griseb. in Goett. Abh.* xix. (1874) 243.—Reg. Argent.
turnerifolia, *Cham. in Linnaea*, vii. (1832) 217.—Bras.
ukambensis, *Vatke, l. c.* xliii. (1880–82) 528.—Afr. trop.
umbellata, *Cav. Ic.* ii. 75. t. 194.—Mexic.
uncinuligera, Nees, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 377 = canescens.
urticoides, *Steud. Nom.* ed. II. ii. 51.—Bras.
valerianoides, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 377 (Valerianae sp.).—Peruv.
velutina, *Schau. in DC. Prod.* xi. 576.—Bras.
vernonioides, *Cham. in Linnaea*, vii. (1832) 232.—Bras.
virgata, Steud. Nom. ed. II. ii. 751, sphalm. = urticoides.
Wrightii, *A. Gray, in Am. Journ. Sc.* Ser. II. xvi. (1853) 98, nomen; *Torr. Bot. Mex. Bound.* 126.—Mexic.; Arizona.

LIPUSA, Alef. Landw. Fl. 26 (1866) = **Phaseolus**, Linn. (Legumin.).
formosa, Alef. in Bot. Zeit. xxiv. (1866) 254 = P. formosus.
multiflora, Alef. Landw. Fl. 26 = P. multiflorus.

LIQUIDAMBAR, Linn. Hort. Cliff. 486 (1737). *HAMAMELIDEAE*, Benth. & Hook. f. i. 669.
acerifolia, Maxim. in Bull. Acad. Pétersb. x. (1866) 486 = formosana.
Altingia, Blume, Fl. Jav. Balsam. 8. tt. 1, 2 = seq.
Altingiana, Blume, Bijdr. 527 = Altingia excelsa.
Altingiana, Teijsm. ex A. DC. in DC. Prod. xvi. ii. 158 = Bucklandia populnea.
asplenifolia, Linn. Sp. Pl. ed. II. 1418 = Myrica Comptonia.
barbata, Stokes, Bot. Mat. Med. iv. 332 = styraciflua.
chinensis, Champ. ex Benth, in Hook. Kew Journ. iv. (1852) 164 = Altingia chinensis.
formosana, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 215.—China.
gummifera, Salisb. Prod. 393 = L. styraciflua.
imberbis, Ait. Hort. Kew. ed. I. iii. 365 = orientalis.
macrophylla, *Oerst. Am. Centr.* 16 t. 10, 11.—Am. centr.
Maximowiczii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 200 = formosana.

LIQUIDAMBAR:—
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 2.—As. Min.
peregrina, Linn. Sp. Pl. 999 = Myrica Comptonia.
Rasamala, Blume, Cat. Gew. Buitenz. 6 = Altingia excelsa.
styraciflua, *Linn. Sp. Pl.* 999.—Am. bor.
tricuspis, Miq. Fl. Ind. Bat. i. 1. 1097 = Bucklandia populnea.

LIQUIRITIA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 367 = **Glycyrrhiza**, Linn. (Legumin.).
lepidota, DC. Prod. ii. 248 = G. lepidota.
officinalis, Moench, Meth. 152 = G. glabra.
officinarum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 367 = G. glabra.

LIRIACTIS, Rafin. Fl. Tellur. ii. 97 (1836) = **Tulipa**, Tourn. (Liliac.).
albiflora, Rafin. l. c. = T. stellata.

LIRIAMUS, Rafin. Fl. Tellur. iv. 23 (1836) = **Crinum**, Linn. (Amaryll.).
quinqueflorus, Rafin. l. c. = C. angustifolium.

LIRIANTHE, Spach, Hist. Vég. Phan. vii. 485 (1839) = **Magnolia**, Linn.
grandiflora, Spach, l. c. = M. pterocarpa.

LIRIODENDRON, Linn. Sp. Pl. ed. I. 535 (1753). *MAGNOLIACEAE*, Benth. & Hook. f. i. 19.
TULIPIFERA, Herm. ex Mill. Gard. Dict. ed. VI. (1752).
Coco, Lour. Fl. Cochinch. 347 = Magnolia pumila.
Figo, Lour. l. c. = Michelia fuscata.
grandiflora, Roxb. Hort. Beng. 43; Fl. Ind. ii. 653 = Magnolia pterocarpa.
heterophylla, Hort. ex C. Koch, Dendrol. i. 381 = tulipifera.
indica, Spreng. Syst. ii. 642 = Magnolia pterocarpa.
integrifolia, Hort. ex Steud. Nom. ed. I. 486 = tulipifera.
liliifera, Linn. Sp. Pl. ed. II. 755 = Magnolia pumila.
liliifera, Roxb. Hort. Beng. 43; Fl. Ind. ii. 654 = Talauma Rabaniana.
liliiflora, Steud. Nom. ed. II. ii. 55 = Magnolia obovata.
obtusiloba, Hort. ex C. Koch, Dendrol. i. 381 = tulipifera.
procera, Salisb. Prod. 379 = tulipifera.
truncatifolia, Stokes, Bot. Mat. Med. iii. 233 = tulipifera.
tulipifera, *Linn. Sp. Pl.* 535.—Am. bor.; China?
tulipiflora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = praec.

LIRIODENDRUM, Linn. Coroll. 9. (1737) = praec.

LIRIOPE, Herb. App. 41 (1821) = **Elisena**, Herb. (Amaryll.).
ringens, Herb. l. c. 42 = E. ringens.

LIRIOPE, Lour. Fl. Cochinch. 200 (1790). *HAEMODORACEAE*, Benth. & Hook. f. iii. 678.
OPHIOPOGON, Kunth, Enum. Pl. v. 297 (1850).
graminifolia, Baker, in Journ. Linn. Soc. xiv. (1875) 538; xvii. (1879) 499 = spicata.
spicata, *Lour. Fl. Cochinch.* 201.—China; Cochinch.

LIRIOPE, Salisb. Gen. Pl. Fragm. 74 (1866) = **Reineckia**, Kunth (Liliac.).

LIRIOPOGON, Rafin. Fl. Tellur. ii. 35 (1836) = **Tulipa**, Tourn. (Liliac.).
biflorum, Rafin. l. c. = T. biflora.
Celsianum, Rafin. l. c. = T. australis.
sylvestre, Rafin. l. c. = T. sylvestris.

LIRIOPSIS, Reichb. Consp. 62 (1828) = **Elisena**, Herb. (Amaryll.).
longipetala, M. Roem. Syn. Ensat. 184 = seq.?
ringens, Reichb. ex M. Roem. l. c. 183 = E. ringens.

LIRIOPSIS, Spach, Hist. Vég. Phan. vii. 460 (1839) = **Michelia**, Linn. (Magnol.).
fuscata, Spach, l. c. 461 = M. fuscata.

LIRIOSMA, Poepp. & Endl. Nov. Gen. et Sp. iii. 33. t. 239 (1842). *OLACINEAE*, Benth. & Hook. f. i. 347.
DULACIA, Vell. Fl. Flum. i. t. 78 (1827).
HYPOCARPUS, A. DC. in DC. Prod. viii. 245, 673 (1844).
acuta, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. iv. (1859) 363.—Bras.
adhaerens, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. ii. 26.—Bras.; Peruv.
candida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 33. t. 239.—Am. austr.
Gardneriana, *A. DC. in DC. Prod.* viii. 673.—Bras.
grandiflora, *Engl. in Mart. Fl. Bras.* xii. ii. 25.—Bras.
guianensis, *Engl. l. c.*—Guian.
inopiflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. iv. (1859) 362.—Bras.
macrophylla, *Benth. ex Miers, l. c.* 364.—Bras.
micrantha, Spruce, ex Engl. in Mart. Fl. Bras. xii. ii. 28 = inopiflora.
ovata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. iv. (1859) 363.—Bras.
pallida, *Miers, l. c.* 362.—Bras.
pauciflora, *A. DC. in DC. Prod.* viii. 673.—Bras.
Pohliana, *Engl. in Mart. Fl. Bras.* xii. ii. 33.—Bras.
Spruceana, *Engl. l. c.* 24.—Peruv.
Velloziana, *A. DC. in DC. Prod.* viii. 673.—Bras.

LIRIUM, Linn. Syst. ed. I. (1735); J. G. Gmel. Fl. Sibir. i. 41 (1747) = **Lilium**, Tourn.

LISAEA, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 54 = **Caucalis**, Tourn. (Umbellif.).
grandiflora, Boiss. l. c. = C. grandiflora (in Addend.).
heterocarpa, Boiss. Fl. Orient. ii. 1088 = C. grandiflora (in Addend.).
papyracea, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 55.—As. Min.
syrica, Boiss. l. c.—Syria.

LISIANTHIUS, P. Br. Hist. Jamaic. 157 (1756) = seq.

LISIANTHUS, Linn. Mant. i. 6 (1767). *GENTIANEAE*, Benth. & Hook. f. ii. 813.
HELIA, Mart. Nov. Gen. et Sp. ii. 122. t. 191 (1827).
IRLBACHIA, Mart. l. c. 101. t. 179 (1827).
LEIOTHAMNUS, Griseb. Gen. et Sp. Gent. 205 (1839).
SCHLIMIA, Regel, in Act. Hort. Petrop. iii. ii. (1875) 285.
SYMBOLANTHUS, G. Don, Gen. Syst. iv. 210 (1837).
WALLISIA, Regel, Gartenfl. xxiv. (1875) 230. t. 839.
acuminatus, Desr. in Lam. Encyc. iii. 660 = frigidus.
acutangulus, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 49.—Bras.
acutangulus, Ruiz & Pav. Fl. Per. ii. 14 = tetragonus.
alatus, *Aubl. Pl. Gui.* i. 204.—Guian.
alatus, Willd. ex Griseb. Gen. et Sp. Gent. 185 = trifidus.
albus, *Spruce, ex Prog. in Mart. Fl. Bras.* vi. i. 237.—Bras.
alpestris, *Mart. Nov. Gen. et Sp.* ii. 93. t. 171.—Bras.
amoenus, Miq. in Linnaea, xix. (1847) 139 = uliginosus.
amplissimus, *Mart. Nov. Gen. et Sp.* ii. 96. t. 175.—Bras.
angustifolius, *H. B. & K. Nov. Gen. et Sp.* iii. 181.—Venezuela.
angustifolius, Mart. Nov. Gen. et Sp. ii. 99. t. 178 = tenuifolius.
anomalus, *H. B. & K. Nov. Gen. et Sp.* iii. 184.—N. Granat.; Peruv.
aphyllus, *Vell. Fl. Flum.* 78; ii. t. 80.—Bras.
arboreus, *Spruce, ex Prog. in Mart. Fl. Bras.* vi. i. 240.—Bras.
auriculatus, *Benth. Pl. Hartw.* 226.—N. Granat.
bifidus, *H. B. & K. Nov. Gen. et Sp.* iii. 181.—Venezuela.
breviflorus, Benth. in Hook. Journ. Bot. ii. (1840) 45 = Pagaea breviflora.
brevifolius, *Griseb. Gen. et Sp. Gent.* 187.—Bras.
caerulescens, *Aubl. Pl. Gui.* i. 207.—Am. austr.
callosus, Bert. ex Spreng. Syst. i. 585 = Leianthus exsertus.
calygonus, *Ruiz & Pav. Fl. Per.* ii. 14.—Peruv.
campanulaceus, *Desr. in Lam. Encyc.* iii. 660.—Am. austr.

LISIANTHUS:—
campanuloides, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 201.—Bras.
carinatus, Desr. in Lam. Encyc. iii. 658 = Tachiadenus carinatus.
chelonoides, *Linn. f. Suppl.* 134.—Am. austr.
coerulescens, Auct. = caerulescens.
cordifolius, Linn. Mant. i. 43 = Leianthus longifolius.
corymbosus, Ruiz & Pav. Fl. Per. ii. 14 = glaber.
crassicaulis, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 374.—Mexic.
cuspidatus, Bertol. in Nov. Comm. Bonon. iv. (1840) 408 = Leianthus nigrescens.
daturoides, *Griseb. Gen. et Sp. Gent.* 177.—Peruv.
densiflorus, *Benth. Pl. Hartw.* 227.—N. Granat.
elegans, *Mart. Nov. Gen. et Sp.* ii. 98. t. 177.—Bras.
Elisabethae, *Griseb. in Linnaea*, xxii. (1849) 40.—Guian.
exaltatus, Desr. in Lam. Encyc. iii. 662 = Eustoma silenifolium.
exsertus, Sw. Prod. Veg. Ind. Occ. 40 = Leianthus exsertus.
fistulosus, *Poir. in Lam. Encyc. Suppl.* iii. 564.—Am. austr.
frigidus, *Sw. Prod. Veg. Ind. Occ.* 40.—Ind. occ.
glaber, *Linn. f. Suppl.* 134.—Am. austr.
glandulosus, *A. Rich. Fl. Cub. Fanerog.* ii. 102.—Cuba.
glaucifolius, Jacq. Coll. i. 64 = Eustoma silenifolium.
glaucifolius, Nutt. in Trans. Am. Phil. Soc. v. (1837) 197 = Eustoma Russellianum.
gracilis, Griseb. Gen. et Sp. Gent. 182 = uliginosus.
grandiflorus, *Aubl. Pl. Gui.* i. 205.—Am. austr.
grandiflorus, Willd. ex Griseb. Gen. et Sp. Gent. 182 = uliginosus.
inflatus, Mart. Nov. Gen. et Sp. ii. 95. t. 174 = speciosus.
Kunthii, G. Don, Gen. Syst. iv. 208 = angustifolius.
latifolius, Sw. Prod. Veg. Ind. Occ. 40 = Leianthus latifolius.
leucanthus, Lyall, ex Griseb. Gen. et Sp. Gent. 202 = Tachiadenus longiflorus.
longiflorus, Boj. ex Griseb. l. c. = Tachiadenus longiflorus
longiflorus, Willd. ex Roem. & Schult. Syst. iv. 787 = Escobedia scabrifolia.
longifolius, Linn. Mant. i. 43 = Leianthus longifolius.
loranthoides, *Griseb. Gen. et Sp. Gent.* 179.—Peruv.
luteus, *Rafin. Atl. Journ.* 146 (Quid?).—Am. bor.
macranthus, *Benth. Pl. Hartw.* 144.—Ecuador.
macrophyllus, Cham. & Schlecht. in Linnaea, i. (1826) 200 = obtusifolius.
macrophyllus, *H. B. & K. Nov. Gen. et Sp.* iii. 183.—N. Granat.
madagascariensis, Spreng. Syst. i. 586 = Tachiadenus trinervis.
Martii, *Griseb. Gen. et Sp. Gent.* 187.—Bras.
Mathewsii, *Griseb. l. c.* 361.—Peruv.
nerioides, *Griseb. in Linnaea*, xxii. (1849) 39.—N. Granat.
nigrescens, Cham. & Schlecht. l. c. vi. (1831) 388 = Leianthus nigrescens.
oblongifolius, *Griseb. Gen. et Sp. Gent.* 187.—Bras.
obtusifolius, *Griseb. l. c.* 175.—Bras.
Oerstedii, Griseb. in Kjoeb. Vidensk. Meddel. (1853) 54 = alatus.
Ophiorriza, Vell. Fl. Flum. 77; ii. t. 78 = Metternichia principis.
ovalis, *Ruiz & Pav. Fl. Per.* ii. 13.—Peruv.
ovatifolius, Vell. Fl. Flum. 78; ii. t. 79 = alpestris?
paniculatus, *Spreng. Syst.* iv. *Cur. Post.* 340.—Bras.
parvifolius, Desr. in Lam. Encyc. iii. 660 = caerulescens.
pedunculatus, Schlecht. ex Spreng. Syst. iv. Cur. Post. 59, 339 = elegans.
pendulus, *Mart. Nov. Gen. et Sp.* ii. 94. t. 172.—Bras.
pratensis, *H. B. & K. Nov. Gen. et Sp.* iii. 180.—Venezuela.
Princeps, *Lindl. in Journ. Hort. Soc.* iv. (1849) 261.—N. Granat.
pulcher, *Hook. Bot. Mag.* t. 4424.—N. Granat.
pulcherrimus, *Mart. Nov. Gen. et Sp.* ii. 97.—Bras.
pumilus, *Benth. in Hook. Kew Journ.* vi. (1854) 200.—Bras.
purpurascens, *Aubl. Pl. Gui.* i. 201.—Am. austr.
purpurascens, H. B. & K. Nov. Gen. et Sp. iii. 182 = uliginosus.
racemosus, Willd. ex Griseb. Gen. et Sp. Gent. 186 = bifidus.
ramosissimus, *Benth. in Hook. Kew Journ.* vi. (1854) 200.—Bras.
recurvus, Benth. in Hook. Kew Journ. vi. (1854) 201 = Pagaea recurva.
revolutus, Poepp. ex Griseb. in DC. Prod. ix. 77 = glaber.
revolutus, *Ruiz & Pav. Fl. Per.* ii. 14.—Peruv.
Russellianus, Hook. Bot. Mag. t. 3626 = Eustoma Russellianum.
saponarioides, Cham. & Schlecht. in Linnaea, vi. (1831) 389 = Leianthus saponarioides.
scandens, Spreng. Syst. i. 587 = Coutoubea volubilis.
Schomburgkii, *Griseb. in DC. Prod.* ix. 75.—Guian.
sempervirens, Mill. ex Steud. Nom. ed. I. 486 = Gelsemium sempervirens.
spathulatus, *H. B. & K. Nov. Gen. et Sp.* iii. 181.—N. Granat.
speciosus, *Cham. & Schlecht. in Linnaea*, i. (1826) 198.—Am. austr.
splendens, *Hook. Lond. Journ. Bot.* vi. (1847) 264.—N. Granat.
Spruceanus, *Benth. in Hook. Kew Journ.* vi. (1854) 202.—Bras.
subcordatus, Benth. l. c. 200 = Pagaea subcordata.
tenuifolius, *Spreng. Syst.* iv. *Cur. Post.* 339.—Bras.
tetragono-alatus, Steud. Nom. ed. II. ii. 55 = acut-angulus.
tetragonus, *Benth. Pl. Hartw.* 68.—Mexic.
thamnoides, *Griseb. Fl. Brit. W. Ind.* 424.—Ind. occ.
trifidus, *H. B. & K. Nov. Gen. et Sp.* iii. 182.—N. Granat.
trinervis, Desr. in Lam. Encyc. iii. 659 = Tachiadenus trinervis.
tubiflorus, Thou. ex Roem. & Schult. Syst. iv. 788 = Tachiadenus tubiflorus.
uliginosus, *Griseb. Gen. et Sp. Gent.* 181.—Am. austr.
umbellatus, Sw. Prod. Veg. Ind. Occ. 40 = Leianthus umbellatus.
vasculosus, *Griseb. in Linnaea*, xxii. (1849) 37.—Venezuela.
virgatus, Boj. ex Griseb. in DC. Prod. ix. 79, 82 = Tachiadenus gracilis.
virgatus, *Prog. in Mart. Fl. Bras.* vi. 1. 239.—Bras.
viridiflorus, *Mart. Nov. Gen. et Sp.* ii. 94.—Bras.
viscosus, *Ruiz & Pav. Fl. Per.* ii. 14.—Peruv.
volubilis, Salisb. Prod. 137 = Gelsemium sempervirens.
zeylanicus, Spreng. Syst. i. 586 = Exacum Wightianum.

LISIMACHIA, Neck. Delic. Gallo-Belg. i. 108 (1768) = **Lysimachia**, Linn. (Primul.).

LISIONOTUS, Reichb. Handb. 199 (1837) = **Lysionotus**, D. Don (Gesnerac.).

LISSANTHE, R. Br. Prod. 540 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 613.
acerosa, Spreng. Syst. i. 660 = Cyathodes acerosa.
ciliata, R. Br. Prod. 541 = Brachyloma ciliatum.
Cunninghamii, DC. Prod. vii. 743 = Brachyloma daphnoides.
daphnoides, R. Br. Prod. 541 = Brachyloma daphnoides.
depressa, F. Muell. Fragm. i. 36 = Brachyloma depressum.
divaricata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 269 = Cyathodes divaricata.
intermedia, A. Cunn. ex DC. Prod. vii. 742 = strigosa.
montana, *R. Br. Prod.* 540.—Austral.
mucronata, *DC. Prod.* vii. 743.—Austral.
Oxycedrus, Spreng. Syst. i. 660 = Cyathodes acerosa.
parvifolia, Spreng. l. c. = Cyathodes parvifolia.
propinqua, A. Cunn. ex DC. Prod. vii. 742 = Leucopogon juniperinus.
rigida, Benth. in Enum. Pl. Hueg. 76 = strigosa.
sapida, *R. Br. Prod.* 540.—Austral.
stellata, Knowles & Westc. Flor. Cab. iii. (1840) 79 = Brachyloma daphnoides.
strigosa, *R. Br. Prod.* 540.—Austral.
strigosa, Sieber, ex DC. Prod. vii. 751 = Leucopogon juniperinus.
subulata, R. Br. Prod. 540 = strigosa.
ulicina, A. Cunn. ex DC. Prod. vii. 743 = strigosa.
verticillata, Lindl. Swan River App. 25 = Leucopogon verticillatus.

LISSERA, Steud. Nom. ed. II. i. 669, in syn. (1840) = LISTERA, Adans. = **Genista**, Linn. (Legum.).
anglica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = G. anglica.
germanica, Fourr. l. c. 358 = G. germanica.
hispanica, Fourr. l. c. = G. hispanica.

LISSOCARPA, Benth. in Benth. & Hook. f. Gen. ii. 671 (1876). *STYRACEAE*, Benth. & Hook. f. l. c.

LISSOCHILOS, Bartl. Ord. 57 (1830) = seq.

LISSOCHILUS, R. Br. in Lindl. Bot. Reg. t. 573 (err. typ. 578) (1821); Coll. Bot. t. 31 (1822). *ORCHIDEAE*, Benth. & Hook. f. iii. 536.
HYPODEMATIUM, A. Rich. Tent. Fl. Abyss. ii. 286. t. 83 (1851).
aequalis, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 204.—Afr. austr.
Alexandri, *Reichb. f. in Flora*, lxv. (1882) 532.—Afr. trop.
amazonicus, *Rodrig. Orch. Nov.* i. 89.—Bras.
angoanensis, Reichb. f. in Flora, xlviii. (1865) 188 = Galeandra anjoanensis.
angolensis, *Reichb. f. Otia Bot. Hamb.* 64.—Afr. trop.
antennisepalus, *Reichb. f. in Flora*, lxv. (1882) 533.—Afr. trop.
arabicus, *Lindl. Gen. et Sp. Orch.* 192.—Arab.
arenarius, *Lindl. in Journ. Linn. Soc.* vi. (1862) 133.—Afr. trop.
aurantiacus, *Reichb. f. in Flora*, xlviii. (1865) 187.—Afr. trop.
Bouliawongo, Reichb. f. l. c. 188 = Galeandra Bouliawongo.
Buchanani, *Reichb. f. Otia Bot. Hamb.* 64.—Afr. trop.
calopterus, *Reichb. f. in Bonplandia*, iii. (1855) 218.—Afr. trop.
caricifolius, *Reichb. f. in Linnaea*, xli. (1877) 74.—Afr. trop.
clitellifer, *Reichb. f. l. c.* xx. (1847) 687.—Afr. austr.
dilectus, *Reichb. f. Otia Bot. Hamb.* 62.—Afr. trop.
Ephippium, *Reichb. f. in Flora*, lxv. (1882) 533 (Ephippilum).—Afr. trop.
fallax, *Reichb. f. Otia Bot. Hamb.* 115.—Afr. trop.
Friderici, *Reichb. f. in Flora*, l. (1867) 113.—Afr. trop.
giganteus, *Welw. ex Reichb. f. l. c.* xlviii. (1865) 187.—Afr. trop.
graniticus, *Reichb. f. Otia Bot. Hamb.* 75.—Abyssin.
Grantii, *Reichb. f. in Trans. Linn. Soc.* xxix. (1875) 153.—Afr. trop.
heteroglossus, *Reichb. f. Otia Bot. Hamb.* 115.—Afr. trop.
Heudelotii, *Reichb. f. l. c.* 63.—Afr trop.
Horsfallii, *Batem. in Hook. Bot. Mag.* t. 5486.—Afr. trop.
Krebsii, *Reichb. f. in Linnaea*, xx. (1847) 685.—Afr. austr.
Lindleyanus, *Reichb. f. Otia Bot. Hamb.* 65.—Afr. trop.
Livingstonianus, *Reichb. f. l. c.* 114.—Afr. trop.
longifolius, *Benth. in Hook. Niger Fl.* 530.—Afr. trop.
luteus, *Sweet, Hort. Brit.* ed. I. 386, nomen.—Afr. austr.
macranthus, *Lindl. Gen. et Sp. Orch.* 191.—Afr. trop.
madagascariensis, *Kraentz. in Bremen, Abh.* vii. (1882) 256.—Madag.
malanganus, *Reichb. f. in Flora*, lxv. (1882) 533.—Afr. trop.
malangensis, *Reichb. f. Otia Bot. Hamb.* 114.—Afr. trop.
Mechowii, *Reichb. f. l. c.* 75.—Afr. trop.
microceras, *Reichb. f. l. c.* 115.—Afr. trop.
Oliverianus, *Reichb. f. in Trans. Linn. Soc.* xxix. (1875) 154.—Afr. trop.
orthoplectrus, *Reichb. f. Otia Bot. Hamb.* 63.—Afr. trop.
Paivaeanus, *Reichb. f. in Flora*, xlviii. (1865) 187.—Afr. trop.
paludicolus, *Reichb. f. Otia Bot. Hamb.* 63.—Afr. trop.
parviflorus, Lindl. Gen. et Sp. Orch. 191 = Eulophia streptopetala.
platypetalus, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 204.—Afr. austr.
platypterus, *Reichb. f. in Flora*, lxv. (1882) 533.—Afr. trop.
porphyroglossus, *Reichb. f. Otia Bot. Hamb.* 61.—Afr. trop.
purpuratus, *Lindl. in Journ. Linn. Soc.* vi. (1862) 133.—Afr. trop.
pyrophilus, *Reichb. f. Otia Bot. Hamb.* 65.—Afr. trop.
Renschianus, *Reichb. f. l. c.* 75.—Afr. trop.
Roscheri, *Reichb. f. l. c.* 62.—Afr. trop.
roseus, *Lindl. Bot. Reg.* (1843) *Misc.* 25.—Afr. trop.
Rueppelii, *Reichb. f in Linnaea*, xx. (1847) 686.—Abyssin.?
Rutenbergianus, *Kraentz. in Bremen, Abh.* vii. (1882) 257.—Madag.

LISSOCHILUS:—
Sandersoni, *Reichb. f. Otia Bot. Hamb.* 62.—Afr. austr.
Schweinfurthii, *Reichb. f. l. c.* 60.—Afr. trop.
speciosus, *R. Br. ex Lindl. Coll. Bot.* 31.—Afr. austr.
streptopetalus, Lindl. Gen. et Sp. Orch. 191 = Eulophia streptopetala.
stylites, *Reichb. f. Otia Bot. Hamb.* 61.—Afr. trop.
Wakefieldi, *Reichb. f. & S. Moore, in Journ. Bot.* xvi. (1878) 136.—Afr. trop.
Welwitschii, *Reichb. f. in Flora*, xlviii. (1865) 188.—Afr. trop.

LISSOSCHILUS, Hornsch. in Syll. Ratisb. i. (1824) 125 = praec.

LISTERA, Adans. Fam. ii. 321 (1763).—Cf. LISSERA, Steud. = **Genista**, Tourn. (Legum.).

LISTERA, R. Br. in Ait. Hort. Kew. ed. II. v. 201 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 595.
DIPHRYLLUM, Rafin. Med. Repos. N. York, v. (1808) 356.
DISTOMAEA, Spenn. Fl. Friburg. i. 245 (1825).
POLLINIRHIZA, Dulac, Fl. Hautes-Pyr. 120 (1867).
australis, *Lindl. Gen. et Sp. Orch.* 456.—Am. bor.
Banksiana, Lindl. l. c. 455 = Eschscholziana.
convallarioides, Ell. Sketch, ii. 494 = australis.
convallarioides, *Nutt. Gen. Am.* ii. 191.—Am. bor.
cordata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 201.—Reg. bor. temp.
Eschscholziana, *Cham. & Schlecht. in Linnaea*, iii. (1828) 33.—As. bor.
japonica, *Blume, Orch. Archip. Ind.* 136.—Japon.
micrantha, *Lindl. in Journ. Linn. Soc.* i. (1857) 176.—Reg. Himal.
Nidus-avis, Hook. Fl. Lond. t. 58 = Neottia Nidus-avis.
ovata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 201.—Europ.; As. bor.
pinetorum, *Lindl. in Journ. Linn. Soc.* i. (1857) 175.—Reg. Himal.
puberula, *Maxim. in Bull. Acad. Pétersb.* xxix. (1884) 204.—China.
pubescens, Ell. Sketch, ii. 494 = Ponthieva glandulosa.
reniformis, D. Don, Prod. Fl. Nep. 28 (= *Neottia reniformis*).—Ind. or.
tenuis, *Lindl. in Journ. Linn. Soc.* i. (1857) 176.—Reg. Himal.

LISTERIA, Neck. Elem. i. 206 (1790) = **Oldenlandia**, Linn. (Rubiac.).

LISTERIA, Spreng. Anleit. ii. 1. 293 (1817) = **Listera**, R. Br. (Orchid.).

LISTIA, E. Mey. Comm. Pl. Afr. Austr. 80 (1835). *LEGUMINOSAE*, Benth. & Hook. f. i. 476.
heterophylla, *E. Mey. l. c.* 81.—Afr. austr.

LISTROSTACHYS, Reichb. f. in Bot. Zeit. x. (1852) 930 = **Angraecum**, Bory (Orchid.).
anthomaniaca, Reichb. f. in Linnaea, xli. (1877) 75.—Afr. trop.
Aphrodite, Balf. f. & S. Moore, in Baker, Fl. Maurit. 354.—Ins. Rodrig.
arcuata, Reichb. f. in Walp. Ann. vi. 907 = A. arcuatum.
ashantensis, Reichb. f. in Walp. l. c. 908 = A. ashantense.
capitata, Reichb. f. in Flora, xlviii. (1865) 190 = A. capitatum.
caudata, Reichb. f. in Walp. Ann. vi. 907 = A. caudatum.
Cephalotes, Reichb. f. in Gard. Chron. (1872) 1687.—Afr. trop.
Chailluana, Reichb. f. in Flora, lxviii. (1885) 381 = A. Chailluanum.
dactyloceras, Reichb. f. l. c. xlviii. (1865) 190.—Afr. trop.
fragrantissima, Reichb. f. l. c. 189.—Afr. trop.
Jenischiana, Reichb. f. in Bot. Zeit. x. (1852) 930 = A. pertusum.
monodon, Reichb. f. in Walp. Ann. vi. 908 = A. monodon.
Monteirae, Reichb. f. in Linnaea, xli. (1877) 76.—Afr. trop.
odoratissima, Reichb. f. in Gard. Chron. (1880) 1. 806, in obs. = A. odoratissimum.

LISTROSTACHYS:—
Papagayi, Reichb. f. in Flora, xlviii. (1865) 189.—Afr. trop.
parviflora, S. Moore, in Baker, Fl. Maurit. 355 = Angraecum parviflorum.
pellucida, Reichb. f. in Walp. Ann. vi. 908 = A. pellucidum.
pertusa, Reichb. f. in Bot. Zeit. x. (1852) 930 = A. pertusum.
Pescatoriana, S. Moore, in Baker, Fl. Maurit. 354 = A. Pescatorianum.
polystachys, Reichb. f. in Walp. Ann. vi. 909 = Aeonia polystachya.
ringens, Reichb. f. in Gard. Chron. (1878) II. 266.—Afr. trop.
Sedeni, Reichb. f. l. c. (1878) I. 138 = A. arcuatum.
subulata, Reichb. f. in Walp. Ann. vi. 909 = A. subulatum.
tridens, Reichb. f. in Flora, xlviii. (1865) 190 = A. tridens.
vesicata, Reichb. f. l. c. = A. vesicatum.
Welwitschii, Reichb. f. l. c.—Afr. trop.

LISYANTHUS, Aubl. Pl. Gui. i. 201 (1775) = **Lisianthus**, Linn. (Gentian.).

LITA, Schreb. Gen. ii. 795 (1791) = **Voyria**, Aubl. (Gentian.).
aphylla, F. G. Dietr. Vollst. Lexik. Gaertn. v. 531 = V. aphylla.
breviflora, Spreng. Syst. i. 581 = V. breviflora.
coerulea, Willd. Sp. Pl. i. 1071 = V. coerulea.
rosea, Willd. l. c. = V. rosea.
spathacea, Spreng. Syst. i. 581 = V. spathacea.

LITANTHES, Lindl. Veg. Kingd. 205 (1847) = seq.

LITANTHUS, Harv. in Hook. Lond. Journ. Bot. iii. (1844) 314. t. 9. *LILIACEAE*, Benth. & Hook. f. iii. 808.
pusillus, *Harv. l. c.* 315. t. 9.—Afr. austr.

LITCHI, Sonner. Voy. Ind. iii. 255 (1782) = **Nephelium**, Linn. (Sapind.).
chinensis, Sonner. l. c. = N. Litchi.
sinensis, J. F. Gmel. Syst. 635 = N. Litchi.

LITHACHNE, Beauv. Agrost. 135. t. 24 (1812) = **Olyra**, Linn. (Gramin.).
axillaris, Beauv. l. c. 166 = O. pauciflora.
pauciflora, Beauv. l. c. 135 = O. pauciflora.

LITHACNE, Poir. in Dict. Sc. Nat. xxvii. 60 (1823) = praec.

LITHAGROSTIS, Gaertn. Fruct. i. 7. t. 1 (1788) = **Coix**, Linn. (Gramin.).
Lachryma-Jobi, Moench, Meth. 209 = seq.
Lacryma-Jobi, Gaertn. Fruct. i. 7. t. 1. f. 10 = C. Lacryma-Jobi.

LITHANTHUS, Pfeiff. Nom. ii. 136 (1874) = **Litanthus**, Harv. (Liliac.).

LITHOBIUM, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. II. Bot. (1835) 140. t. 8. f. 2. *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
PETROBIUM, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. II. Bot. t. 4 (1838).
cordatum, *Bong. l. c.* 140.—Bras.

LITHOCARPOS, Targ.-Tozz. ex Steud. Nom. ed. II. ii. 56 (1841) = **Cocos**, Linn. (Palm.).
cocciformis, Targ.-Tozz. ex Steud. l. c. = C. lapidea.

LITHOCARPUS, Blume, Bijdr. 526 (1825); Fl. Jav. Cupul. 34. t. 20 (1828) = **Quercus**, Tourn. (Cupul.).
angustifolia, Miq. Fl. Ind. Bat. i. 1. 865 (nom. delend.).—Sumatra.
javensis, Blume, Bijdr. 526 = Q. javensis.
scutigera, Oudem. in Versl. en Med. Kon. Akad. Wetensch. xii. (1861) 207 = Q. costata.

LITHOCARPUS, Targ.-Tozz. ex Steud. Nom. ed. II. i. 170, in syn. (1840) = LITHOCARPOS, Targ.-Tozz. = **Cocos**, Linn. (Palm.).

LITHOCNIDES, Rafin. Fl. Tellur. iii. 48 (1836) = **Rousselia**, Gaudich. (Urtic.).
lappulacea, Rafin. l. c. = R. lappulacea.

LITHODORA, Griseb. Spicil. Fl. Rumel. ii. 85 (1844) = **Lithospermum**, Tourn. (Boragin.).
calabra, Griseb. l. c. = Lithosp. calabrum.
fruticosa, Griseb. l. c. = Lithosp. fruticosum.
graminifolia, Griseb. l. c. = Lithosp. graminifolium.
hispidula, Griseb. l. c. = Lithosp. hispidulum.
oleifolia, Griseb. l. c. = Lithosp. oleaefolium.
prostrata, Griseb. l. c. = Lithosp. fruticosum.

LITHOFRAGMA, Nutt. in Journ. Acad. Philad. vii. (1834) 26 = LITHOPHRAGMA, Torr. & Gray = **Tellima**, R. Br. (Saxif.).

LITHOPHILA, Sw. Prod. Veg. Ind. Occ. 14 (1788); Fl. Ind. Occ. i. 47. t. 1 (1797) = **Alternanthera**, Forsk. (Amarant.).
japonica, J. F. Gmel. Syst. 1577, (Index) lapsu = Globba japonica.
muscoides, Sw. Prod. Veg. Ind. Occ. 14; Fl. Ind. Occ. i. 47 = A. muscoides.

LITHOPHRAGMA, Torr. & Gray, Fl. N. Am. i. 583 (1840) = **Tellima**, R. Br. (Saxifr.).
affinis, A. Gray, in Proc. Am. Acad. vi. (1864–65) 534 = T. affinis.
Bolanderii, A. Gray, l. c. 535 = T. Bolanderi.
Cymbalaria, Torr. & Gray, Fl. N. Am. i. 585 = T. Cymbalaria.
glabra, Nutt. ex Torr. & Gray, l. c. i. 584 = T. tenella.
heterophylla, Torr. & Gray, l. c. = T. heterophylla.
micrantha, Nutt. ex Torr. & Gray, l. c. = T. parviflora.
nudicaulis, Nutt. ex Torr. & Gray, l. c. 587 = Mitella trifida.
parviflora, Nutt. ex Torr. & Gray, l. c. 584 = T. parviflora.
parvifolia, Walp. Rep. iv. 523 (Index) = praec.
tenella, Nutt. ex Torr. & Gray, Fl. N. Am. i. 584 = T. tenella.

LITHOPLIS, Rafin. Sylva Tellur. 32 (1838) = **Rhamnus**, Tourn.
saxatilis, Rafin. l. c. = R. saxatilis.

LITHOSANTHES, 'Blume,' ex Flora, viii. (1825) 129 = **Litosanthes**, Blume (Rubiac.).

LITHOSCIADIUM, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 730 = **Selinum**, Linn. (Umbell.).
multicaule, Turcz. l. c. 731 (= *Cnidium multicaule*).—Sibir. baical.

LITHOSPERMUM, [Tourn.] Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 860.
AEGONYCHON, S. F. Gray, Nat. Arr. Brit. Pl. ii. 354 (1821).
BATSCHIA, J. F. Gmel. Syst. 315 (1791).
BUGLOSSOIDES, [Riv.] Moench, Meth. 418 (1794).
CYPHORIMA, Rafin. in Am. Monthly Mag. (1819) 191, 357; Catal. 13 (1824).
GYMNOLEIMA, Decne. in Jacquem. Voy. Bot. 122 (1844).
LITHODORA, Griseb. Spicil. Fl. Rumel. ii. 85 (1844).
MARGAROSPERMUM, Spach, Hist. Vég. Phan. ix. 31 (1840); Decne. in Jacquem. Voy. Bot. 122 (1844).
PENTALOPHUS, A. DC. in DC. Prod. x. 86 (1846).
RHYTISPERMUM, Link. Handb. i. 579 (1829).
aegyptiacum, Mikan, ex Schrank, in Denkschr. Akad. Muench. 1813 (1814) 73 = tenuiflorum.
affine, *DC. Prod.* x. 78.—Afr. austr.
aggregatum, *Ruiz & Pav. Fl. Per.* ii. 4.—Peruv.
alpinum, Willd. ex Roem. & Schult. Syst. iv. 742 = Krynitzkia linifolia.
ambiguum, *R. Br. in Salt, Abyss. App.* 62, nomen.—Abyssin.
angustifolium, Forsk. Fl. Aegypt. Arab. 39 = callosum.
angustifolium, *Michx. Fl. Bor. Am.* i. 130.—Am. bor.
apulum, Bert. in Merc. Chil. 620; ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 127 = Amsinkia angustifolia.
apulum, Eichw. Casp. Cauc. 6 = tenuiflorum.
apulum, Frivald. ex Boiss. Fl. Orient. iv. 229 = Alkanna primuliflora.
apulum, *Vahl, Symb. Bot.* ii. 33.—Europ.; Reg. Mediterr.
Arnebia, Delile, Fl. Aegypt. Illustr. 55 = Arnebia tinctoria.
arvense, Bové, ex DC. Prod. x. 74 = incrassatum.
arvense, Ledeb. Fl. Alt. i. 174 = tenuiflorum.
arvense, *Linn. Sp. Pl.* 132.—Europ.; Oriens.

LITHOSPERMUM:—
arvense, Thunb. Fl. Jap. 81 = erythrorhizon.
bejariense, DC. Prod. x. 79 = hirtum.
bicolor, Bertol. Misc. Bot. i. 14 = arvense.
breviflorum, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 252 = angustifolium.
caeruleum, Gueldenst. Reisen, ii. 25 = purpurocaeruleum.
calabrum, *Tenore, Fl. Nap.* iii. 177; *Syll. Fl. Neap.* 80.—Italia.
californicum, *A. Gray, in Proc. Am. Acad.* x. (1874) 51.—Calif.
callosum, *Vahl, Symb. Bot.* i. 14.—Am. bor.; Peruv.
calycinum, Moris, in Mem. Acc. Torin. xxxvii. (1834) 98 = Amsinckia angustifolia.
calycinum, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 30.—Reg. Cauc.
canescens, *Lehm. Asperif.* 305.—Am. bor. occ.
canescens, Torr. in Pacif. Rail. Rep. iv. 124 = californicum.
carolinianum, Lam. Tabl. Encyc. i. 397 = Onosmodium molle.
Chamaecistus, Pourr. ex Willd. & Lange, Prod. Fl. Hisp. ii. 499 = fruticosum.
chilense, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 127. t. 40.—Chili.
chinensis, *Hook. & Arn. Bot. Beech. Voy.* 202.—China.
ciliatum, Forsk. Fl. Aegypt. Arab. 39; Lehm. Asperif. 314 = callosum.
ciliatum, *Rafin. New Fl. Am.* iv. 18 (Quid?).—Am. bor.
cinereum, *DC. Prod.* x. 73.—Afr. austr.
circumscissum, Hook. & Arn. Bot. Beech. Voy. 370 = Krynitzkia circumscissa.
clandestinum, Trev. ex DC. Prod. x. 58, 129 = Krynitzkia clandestina.
cobrense, *Greene, in Coult. Bot. Gaz.* vi. (1881) 157.—Mexic.
collinum, Salisb. Prod. 113 = fruticosum.
commutatum, Bianca, in Atti Gioen. Catan. Ser. II. i. (1844) 292 = Alkanna tinctoria.
confertiflorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 95.—Japon.
consobrinum, *Pomel, Nouv. Mat. Fl. Atl.* 296.—Alger.
conspicuum, Spreng. Syst. i. 544 = canescens.
cornutum, Ledeb. Fl. Alt. i. 175 = Arnebia cornuta.
Corsonianum, *G. Don, in Loud. Gard. Mag.* (1842) 371.—Ins. Timor.
corymbosum, Lehm. Pugill. ii. 27 = Mertensia paniculata.
crassifolium, *Lehm. Asperif.* 324.—As. Min.
cuneifolium, *Pers. Syn.* i. 158.—Peruv.
cuspidatum, Rafin. New Fl. Am. iv. 18 (Quid?).—Am. bor.
cyanochroum, Boiss. Diagn. Ser. I. vii. 33 = Macrotomia perennis.
davuricum, Lehm. Asperif. 296 = Mertensia dahurica.
decumbens, Torr. in Ann. Lyc. N. York, ii. (1828) 225 = hirtum, pilosum.
decumbens, Vent. Jard. Cels, 37 = Arnebia cornuta.
deflexum, Schur, Enum. Pl. Transs. 934, Index = Echinospermum deflexum.
densiflorum, Ledeb. ex Nordm. in Bull. Sc. Pétersb. ii. (1837) 312 = Macrotomia Cephalotes.
denticulatum, Lehm. Asperif. ii. 249 = Mertensia sibirica.
dichotomum, Ruiz & Pav. Fl. Per. ii. 5 = Coldenia dichotoma.
dictyocarpum, DC. ex A. DC. in DC. Prod. x. 85, 101 = Alkanna hirsutissima.
diffusum, Lag. in Varied. Cienc. ii. iv. (1805) 39 = fruticosum.
digynum, Forsk. Fl. Aegypt. Arab. 40 = Heliotropium luteum.
discolor, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 337.—Mexic.
dispermum, Linn. Sp. Pl. ed. II. 191 = Rochelia stellulata.
distichum, *Orteg. Hort. Matr. Dec.* 8.—Cuba.
divaricatum, Sieber, ex Spreng. Syst. i. 543 = Echiochilon fruticosum.
diversifolium, *DC. Prod.* x. 77.—Afr. austr.
Drummondii, Lehm. Pugill. ii. 26 = Mertensia sibirica.
echioides, Benth. in Royle, Illustr. Bot. Himal. 305 = Mertensia echioides.
elongatum, Decne. in Jacquem. Voy. Bot. 120 = Mertensia elongata.
ericetorum, Salzm. ex DC. Prod. x. 81 = fruticosum.

LITHOSPERMUM:—
erythrorhizon, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 149.—Japon.
euchromon, Royle, Illustr. Bot. Himal. 305 = Macrotomia perennis.
flavescens, Steud. Nom. ed. II. ii. 56 = Alkanna lutea.
flexuosum, *Lehm. Asperif.* ii. 333.—Afr. austr.
floridanum, *Rafin. New Fl. Am.* iv. 18 (Quid?).—Florida.
frondosum, Salisb. Prod. 113 = Alkanna lutea.
frutescens, *Pall. Reise*, i. 188.—Ross. med.
fruticosum, *Linn. Sp. Pl.* 133.—Europ. austr.
fruticosum, Sibth. & Sm. Fl. Graec. ii. 52. t. 161 = rosmarinifolium.
fruticosum, Vitm. Sagg. 45, 47 = graminifolium.
fulvidulum, Bertol. Misc. Bot. i. 13 = tenuiflorum.
Gasparrinii, Heldr. ex Guss. Fl. Sic. Syn. i. 217 = incrassatum.
Gastoni, *Benth. in DC. Prod.* x. 83.—Mont. Pyren.
glabrum, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 227.—Arizona.
gracile, Rafin. New Fl. Am. iv. 17 (Quid?).—Am. bor.
graminifolium, Roem. & Schult. Syst. iv. 46 = rosmarinifolium.
graminifolium, *Viv. Fl. Ital. Fragm.* i. 3. t. 5.—Italia.
helianthemoides, Shuttl. ex DC. Prod. x. 85, 78 = angustifolium?
heliotropoides, Forsk. Fl. Aegypt. Arab. 39 = Heliotropium supinum.
hirsutissimum, Bertol. Misc. Bot. i. 13 = Alkanna hirsutissima.
hirsutum, E. Mey. ex DC. Prod. x. 77 = diversifolium.
hirtum, *Lehm. Asperif.* 304.—Am. bor.
hispidissimum, Lehm. Ic. Nov. Stirp. t. 39 = Arnebia hispidissima.
hispidulum, *Sibth. & Sm. Fl. Graec.* ii. 53. t. 162.—As. Min.
hispidum, Forsk. Fl. Aegypt. Arab. 39 = Heliotropium undulatum.
hispidum, *Ruiz & Pav. Fl. Per.* ii. 5.—Peruv.
humifusum, Spreng. Syst. i. 543 = Heliotropium humifusum.
hypocrateriforme, Hornem. Hort. Hafn. Suppl. 19 = latifolium?
hypocrateriforme, *Sch. Schult. Pl. Nov. Hort. Tubing.* (1825) 23.—Hab.?
incanum, Forst. f. Prod. 12 = Heliotropium anomalum.
incanum, Ruiz & Pav. Fl. Per. ii. 4 = cuneifolium.
incisum, Lehm. Asperif. ii. 303 = angustifolium.
incrassatum, *Guss. Ind. Sem. Hort. Boccad.* (1826) 6; *Fl. Sic. Prod.* i. 211.—Reg. Mediterr.
incurvum, Thib. ex Stend. Nom. ed. I. 487 = Rochelia stellulata.
inornatum, *DC. Prod.* x. 73.—Afr. austr.
japonicum, *A. Gray, in Bot. Perry, Jap. Exped.* ii. 316.—Japon.
javanicum, Spreng. Syst. i. 547.—Java.
kamczaticum, Turcz. in Bull. Soc. Nat. Mosc. (1840) 75 = Mertensia paniculata.
Kotschyi, Boiss. & Hohen. Diagn. Ser. I. iv. 49 = Sericostoma Kotschyi.
latifolium, Forsk. Fl. Aegypt. Arab. 39 = callosum.
latifolium, *Michx. Fl. Bor. Am.* i. 131.—Am. bor.
Ledebourii, Steud. Nom. ed. II. ii. 56 = Rochelia leiocarpa.
Lehmanni, Tineo, in Guss. Fl. Sic. Syn. ii. 791 = Alkanna tinctoria.
Leithneri, Heldr. ex Boiss. Fl. Orient. iv. 216 = incrassatum.
leucophlaeum, Schweinf. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 152 = Heliotropium rariflorum.
lignosum, Schweinf. ex Vatke, in Oestr. Bot. Zeitschr. (1875) 167 = Heliotropium lignosum.
linearifolium, Goldie, in Edinb. Phil. Journ. (Apr. 1822) 322 = angustifolium.
lineatum, *Lehm. Asperif.* 312.—Hab.?
linifolium, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 338.—Mexic.
longiflorum, Moç. & Sesse, ex A. DC. in DC. Prod. x. 68 = Macromeria longiflora.
longiflorum, Salisb. Prod. 113 = purpurocaeruleum.
longiflorum, Spreng. Syst. i. 554 = angustifolium.
longifolium, Willd. ex Roem. & Schult. Syst. iv. 742 = Onosmodium strigosum.
lutescens, N. Coleman, Cat. Pl. Gr. Rapids Michig. (1874) 29 = latifolium.

LITHOSPERMUM:—
lycopsoides, Lehm. Pugill. ii. 28 = Amsinckia lycopsoides.
majus, Gilib. Fl. Lituan. i. 21 = officinale.
mandanense, Spreng. Syst. i. 554 = angustifolium.
marginatum, Lehm. in Hook. Fl. Bor. Am. ii. 86 = Mertensia oblongifolia.
marginatum, Spreng. Syst. i. 547 = Mertensia lanceolata.
maritimum, Lehm. Asperif. 291 = Mertensia maritima.
matamorense, *DC. Prod.* x. 76.—Texas.
medium, Chev. Fl. Env. Paris, ii. 489 = arvense.
mesopotamicum, DC. Prod. x. 75 = apulum.
micranthum, Viv. Fl. Lib. Spec. 10. t. 1. f. 4 = Arnebia cornuta.
microspermum, *Boiss. Diagn.* Ser. II. iii. 135.—Afr. bor. occ.
minimum, *Moris, Sard. Elench.* ii. 7.—Italia.
minus, Gilib. Fl. Lituan. i. 21 = arvense.
molle, Muhl. Cat. 19 = Onosmodium molle.
moltkioides, Decne. in Jacquem. Voy. Bot. 122 = Mertensia moltkioides.
multiflorum, *S. Wats. in Bot. King's Exped.* 238.—Am. bor. occ.
Murasaki, Siebold, in Verh. Batav. Gen. xii. (1830) 32 = erythrorhizon.
muricatum, Ruiz & Pav. Fl. Per. ii. 4 = Eritrichium muricatum.
myosotoides, Lehm. Asperif. 319 = Plagiobothrys tinctorius.
niveum, Poir. Encyc. Suppl. ii. 836 = callosum.
obtusum, *Vis. in Flora*, xx. (1837) 442.—Aegypt.
ochroleucum, Stokes, Bot. Mat. Med. i. 270 = officinale.
officinale, *Linn. Sp. Pl.* 132.—Europ.; Oriens; As. bor.
oleaefolium, *Lapeyr. Hist. Abr. Pl. Pyr. Suppl.* 28.—Mont. Pyren.
orientale, Georgi, Beschr. Russ. Reich. III. iv. 751 = Nonnea lutea.
orientale, Kotschy, ex A. DC. in DC. Prod. x. 86, 98 = Alkanna Kotschyana.
orientale, Linn. Syst. ed. XII. 145 = Alkanna orientalis.
orientale, Loisel. Fl. Gall. ed. II. i. 149 = Alkanna lutea.
ovalifolium, Decne. in Jacquem. Voy. Bot. 119 = Mertensia racemosa.
Pallasii, Ledeb. Fl. Alt. i. 176 = Mertensia sibirica.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 122.—Mexic.
paniculatum, Lehm. Asperif. ii. 289 = Mertensia paniculata.
papillosum, *Thunb. Prod. Pl. Cap.* 34.—Afr. austr.
pentagonum, Pers. Syn. i. 158 = Arnebia cornuta.
permixtum, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 344 = incrassatum.
peruvianum, *DC. Prod.* x. 77.—Peruv.
petraeum, A. DC. l. c. 82 = Moltkia petraea.
pilosum, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 256 = multiflorum.
pilosum, *Nutt. in Journ. Acad. Sc. Philad.* vii. (1834) 43.—Am. bor. occ.
plebeium, Cham. in Linnaea, ix. (1829) 446 = Krynitzkia plebeia.
prostratum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 462 = matamorense.
prostratum, Loisel. Fl. Gall. i. 105. t. 4 = fruticosum.
pulchrum, Lehm. Asperif. 290 = Mertensia pulmonarioides.
pumilum, *Lehm. l. c.* 319.—Am. bor.
purpurascens, Gueldenst. Reisen, i. 189 = purpurocaeruleum.
purpureo-coeruleum, Desf. ex DC. Prod. x. 83 = Gastoni.
purpureo-coeruleum, Thore, Chlor. Landes, 51 = fruticosum.
purpureum, Gueldenst. Reisen, i. 421 = seq.
purpurocaeruleum, *Linn. Sp. Pl.* 132.—Europ.; Oriens.
ramosum, Willd. ex Lehm. Asperif. 328 = Krynitzkia ramosa.
repens, Stokes, Bot. Mat. Med. i. 271 = purpurocaeruleum.
retortum, Pall. Reise, iii. 718 = Rochelia stellulata.
rivulare, Turcz. in Bull. Soc. Nat. Mosc. (1840) 74 = Mertensia rivularis.
Rochelii, Frivald. in Flora, xviii. (1835) 336 = arvense.
rosmarinifolium, Reichb. Fl. Germ. Excurs. 337 = Moltkia petraea.
rosmarinifolium, *Tenore, Fl. Nap.* iii. 178. t. 114.—Italia; Graecia.
ruderale, Dougl. ex Lehm. Pugill. ii. 28 = pilosum.

LITHOSPERMUM :—
rugosum, DC. Prod. x. 83 = Moltkia coerulea.
rugosum, Lucé, Topog. Nachr. Oesel, 43 = arvense.
scabrum, *Thunb. Prod. Pl. Cap.* 34.—Afr. austr.
Schreberianum, Spreng. Syst. i. 546 = Mertensia paniculata.
secundiflorum, Decne. in Jacquem. Voy. Bot. 120. t. 125 = Mertensia echioides.
sericeum, Lehm. Asperif. 306 = canescens.
serrulatum, Turcz. in Bull. Soc. Nat. Mosc. (1840) 246 = Mertensia serrulata.
setosum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 31 = Macrotomia echioides.
sibiricum, Lehm. Asperif. 293 = Mertensia sibirica.
Sibthorpianum, *Griseb. Spicil. Fl. Rumel.* ii. 86.—Oriens.
simplicissimum, Lehm. Asperif. 295 = Mertensia simplicissima.
spathulatum, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 337.—Mexic.
Splitgerberi, *Guss. Fl. Sic. Syn.* i. 217.—Sicil.
strictum, *Lehm. Asperif.* 303.—Am. austr.
strigosum, Bieb. Fl. Taur. Cauc. iii. 121 = apulum.
strigosum, Rafin. New Fl. Am. iv. 18 (Quid?).—Am. bor.
stylosum, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1840) 249 = Mertensia stylosa.
suffruticosum, A. Kern. Sched. Fl. Exsicc. i. 52 = graminifolium.
tenellum, Nutt. in Trans. Am. Phil. Soc. v. (1837) 188 = Heliotropium tenellum.
tenellum, Rafin. in Am. Monthly Mag. (1819) 195 = Myosotis stricta.
tenuiflorum, Herb. ex Nym. Consp. 519 = minimum.
tenuiflorum, *Linn. f. Suppl.* 130.—Oriens; Ind. or.
tenuiflorum, Sibth. & Sm. Fl. Graec. t. 159 = Sibthorpianum.
tetrastigma, Lam. Encyc. iii. 30 = Arnebia tinctoria.
tinctorium, Auch. ex DC. Prod. x. 86, 101 = Alkanna megacarpa.
tinctorium, Bertol. Fl. Ital. ii. 282 = Alkanna scardica.
tinctorium, Linn. Sp. Pl. 132 = Alkanna tinctoria.
tinctorium, Ruiz & Pav. Fl. Per. ii. 4 = Plagiobothrys tinctorius.
tinctorium, Vahl, Symb. Bot. ii. 33. t. 28 = Arnebia tinctoria.
tingens, *Roem. & Schult. Syst.* iv. 44, 778.—Chili.
Torreyi, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 44 = pilosum.
tubatum, Bertol. Misc. Bot. i. 14 = Arnebia cornuta.
tuberosum, *Rugel, ex A. DC. in DC. Prod.* x. 76.—Am. bor.
vesicarium, Loisel. Fl. Gall. ed. II. i. 149 = Nonnea nigricans.
vestitum, Wall. Cat. n. 941 = Arnebia hispidissima.
villosulum, Lehm. Asperif. 288 = Mertensia villosula.
villosum, G. Don, in Sweet, Hort. Brit. ed. III. 490; et ex DC. Prod. x. 861 = Arnebia hispidissima.
villosum, *Spreng. Syst.* i. 546.—Europ. austr.
violaceum, Lam. Fl. Fr. ii. 271 = purpurocaeruleum.
virginianum, Linn. Sp. Pl. 132 = Onosmodium virginianum.
virginicum, Billb. in Linneska Samfund. i. 1832 (1833) 16 = Mertensia pulmonarioides.
virginicum, Linn. Syst. ed. XII. 145 = *virginianum*.
viride, *Greene, in Coult. Bot. Gaz.* vi. (1881) 158.—N. Mexic.
viridiflorum, Roxb. Hort. Beng. [13]; Fl. Ind. i. 455 = Tournefortia sarmentosa.
Webbii, Coss. & Dur. in Bull. Soc. Bot. Fr. xxii. (1875) 64, nomen = microspermum.
Zollingeri, A. DC. in DC. Prod. x. 587.—Java.

LITHOTHAMNUS, Zipp. ex Span. in Linnaea, xv. (1841) 334 = **Ehretia**, Linn. (Borag.).
buxioides, Zipp. ex Span. l. c. = E. buxifolia.

LITHOXYLON, Endl. Gen. 1122 (1840) = **Actephila**, Blume (Euphorb.).
grandifolium, Muell. Arg. in Linnaea, xxxiv. (1865–66) 65 = A. grandifolia.
Lindleyi, Steud. Nom. ed. II. ii. 57.—Ins. Tahiti.
nitidum, Muell. Arg. in DC. Prod. xv. II. 232 = *Lindleyi*.

LITHRAEA, Miers, Trav. Chili, ii. 529 (1826), partim. *ANACARDIACEAE*, [Benth. & Hook. f. i. 418.]
Aroeirinha, *March. ex Warm. in Kjoeb. Vidensk. Meddel.* (1873) 61.—Bras.; Chili.

LITHRAEA :—
australiensis, Engl. in Engl. Bot. Jahrb. i. (1881) 422 = brasiliensis.
brasiliensis, *March. Anacard. Rev.* 183.—Am. trop.
caustica, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 175 = venenosa.
crenata, *Phil. in Linnaea*, xxxiii. (1864–65) 39.—Chili.
Gilliesii, Griseb. in Goett. Abh. xix. (1874) 114 = L. Aroeirinha.
laurina, Walp. Rep. i. 551 = Rhus laurina.
Molle, C. Gay, Fl. Chil. ii. 45 = Schinus latifolius.
Molle, Griseb. ex Engl. in DC. Monog. Phan. iv. 349 (non Anacard.).—Am. austr.
molleoides, Engl. in Mart. Fl. Bras. xii. II. 394. t. 83 = L. Aroeirinha.
montana, *Phil. in Linnaea*, xxxiii. (1864–65) 37.—Chili.
venenosa, *Miers, Trav. Chil.* ii. 529.—Am. austr.
verrucosa, Miers, ex Engl. in DC. Monog. Phan. iv. 247 = brasiliensis.

LITHREA, Hook. Bot. Misc. iii. (1833) 175, sphalm. = praec.

LITHRUM, Huds. Fl. Angl. ed. I. 179, 180 (1762) = **Lythrum**, Linn.

LITOGYNE, Harv. Thes. Cap. ii. 35. t. 155 (1863) = **Epaltes**, Cass. (Compos.).
glabra, Harv. l. c. = E. gariepina.
scabra, Harv. l. c. 36 = E. alata.

LITONIA, Pritz. Ic. Ind. ii. 171 (1855) = **Littonia**, Hook. (Liliac.).

LITOPHILA, Sw. Prod. Veg. Ind. Occ. 14 (1788) = LITHOPHILA, Sw. = **Alternanthera**, Forsk. (Amarant.).

LITORELLA, Aschers. Fl. Brandenb. 544 (1864) = **Littorella**, Berg. (Plantag.).

LITOSANTHES, Blume, Cat. Gew. Buitenz. 21 (1823); et Bijdr. 994. *RUBIACEAE*, Benth. & Hook. f. ii. 131.
biflora, *Blume, ll. cc.* 22, 994.—Java.

LITREA, Phil. in Linnaea, xxxiii. (1864–65) 38 = seq.

LITRIA, G. Don, in Sweet, Hort. Brit. ed. III. 144 (1839) = **Lithraea**, Miers (Anacard.).

LITSAEA, Pers. Syn. ii. 4 (1807) = seq.

LITSEA, Lam. Encyc. iii. 574 (1789). *LAURINEAE*, Benth. & Hook. f. iii. 161.
BERRYA, Klein, ex Willd. Sp. Pl. ii. 840, in syn. (1800).
BRYANTEA, Rafin. Sylva Tellur. 165 (1838).
CYLICODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 61 (1831).
DARWINIA, Dennst. Schluess. Hort. Malab. 31 (1818).
DECAPENTA, Rafin. Sylva Tellur. 165 (1838).
DIPLIATHUS, Rafin. l. c. 135 (1838).
FIWA, J. F. Gmel. Syst. 745 (1791).
GLABRARIA, Linn. Mant. ii. 156 (1771); Miq. Fl. Ind. Bat. i. I. 940 (1857).
HECKERIA, Rafin. Sylva Tellur. 165 (1838).
HEXANTHUS, Lour. Fl. Cochinch. 195 (1790).
LEPIDADENIA, Nees, in Edinb. N. Phil. Journ. xv. (Apr.–Oct. 1833) 379; Laurin. Expos. 19 (1833).
PIPALIA, Stokes, Bot. Mat. Med. iv. 456 (1812).
SEBIFERA, Lour. Fl. Cochinch. 637 (1790).
TETRADENIA, Nees, in Wall. Pl. As. Rar. ii. 61 (1831).
TETRANTHERA, Jacq. Hort. Schoenb. i. 59. t. 113 (1797).
TOMEX, Thunb. Nov. Gen. 65 (1783).
acerina, *Blume, Mus. Bot. Lugd. Bat.* i. 350.—Java.
aciculata, *Blume, l. c.* 347.—Japon.
amara, *Blume, Bijdr.* 563.—Malaya.
ambigua, *Meissn. in DC. Prod.* xv. I. 226.—Zeylan.
angulata, *Blume, Bijdr.* 563.—Java.
angulata, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 181 = Perrottetii.

LITSEA :—
angustifolia, Blume, Bijdr. 566 = Actinodaphne areolata.
angustifolia, Kurz, For. Fl. Brit. Burma, ii. 305 = Actinodaphne angustifolia.
apetala, R. Br. ex Meissn. in DC. Prod. xv. I. 227 = L. Sebifera.
apetala, Pers. Syn. ii. 4 = polyantha.
Baueri, *Endl. Ic.* iv. t. 44.—Austral.
Bindoniana, *F. Muell. Fragm.* v. 167.—Austral.
brachystachya, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 180.—Ins. Philipp.
californica, *Benth. & Hook. f. Gen.* iii. 161.—Calif.
caloneura, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
cassiaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 348.—Java.
celebica, Blume, l. c. 350 = Actinodaphne celebica.
Cervantesii, H. B. & K. Nov. Gen. et Sp. ii. 168 = glaucescens.
ceylanica, C. & T. Nees, Cinnam. Disp. i. (1823) t. 5 = zeylanica.
chinensis, Blume, Bijdr. 565 = Actinodaphne chinensis.
chinensis, Heyne, ex Nees, in Wall. Pl. As. Rar. ii. 65 = ligustrina.
chinensis, Lam. Encyc. iii. 574 = L. Sebifera.
chrysocoma, *Blume, Mus. Bot. Lugd. Bat.* i. 348.—Java.
cinnamomea, *Blume, l. c.* 349.—Ins. Molucc.
citrata, *Blume, Bijdr.* 565.—Reg. Himal.; Malaya.
citrifolia, Juss. in Ann. Mus. Par. vi. (1805) 211 = L. Sebifera?
concolor, Kurz, For. Fl. Brit. Burma, ii. 304 = Actinodaphne concolor.
consimilis, Nees, Syst. Laurin. 628 = zeylanica.
cordifolia, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
Cubeba, *Pers. Syn.* ii. 4.—Cochinch.; China.
Cuipala, Nees, Syst. Laurin. 638 = lanuginosa.
dealbata, *Nees, l. c.* 630.—Austral.
densifolia, *Miq. Fl. Ind. Bat.* i. I. 973.—Java.
discolor, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 382 = timoriana.
diversifolia, *Blume, Bijdr.* 564.—Java.
elliptica, *Blume, l. c.* 563.—Java.
elongata, *Benth. & Hook. f. Gen.* iii. 163, in nota sub Lindera.—Ind. or.
ferruginea, *Blume, Bijdr.* 561.—Malaya; Java.
foliosa, Nees, Syst. Laurin. 622 = zeylanica.
foliosa, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 207 = aciculata.
fruticosa, Span. ex Blume, Mus. Bot. Lugd. Bat. i. 374 = L. Sebifera?
fulva, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
furfuracea, Nees, Syst. Laurin. 625 = zeylanica.
fuscata, *Thw. Enum. Pl. Zeyl.* 258.—Zeylan.
Gardneri, *Meissn. in DC. Prod.* xv. I. 227; *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
geminata, Blume, Bijdr. 564 = L. Sebifera.
geniculata, *Benth. & Hook. f. Gen.* iii. 162; *Nichols. Dict. Gard.* ii. 287.—Am. bor.
glaberrima, *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
Glabraria, Juss. in Ann. Mus. Par. vi. (1805) 212; Pers. Syn. ii. 4 = L. Sebifera.
glauca, *Siebold, in Verh. Batav. Gen.* xii. (1830) 24.—Japon.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* ii. 168.—Mexic.
glaucescens, Spreng. ex Nees, Syst. Laurin. 145 = Persea pyrifolia.
glomerata, Blume, Bijdr. 566 = Actinodaphne glomerata.
hexantha, *Juss. in Ann. Mus. Par.* vi. (1805) 212.—Cochinch.
hexantha, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 378 = polyantha.
japonica, *Mirb. Hist. Nat. Pl.* ed. II. xi. 150; *Juss. l. c.*—Japon.
javanica, *Blume, Mus. Bot. Lugd. Bat.* i. 348.—Java.
khasyana, *Meissn. in DC. Prod.* xv. I. 227.—Reg. Himal.
laeta, *Benth. & Hook. f. Gen.* iii. 161; *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
laevis, Juss. in Ann. Mus. Par. vi. (1805) 212 = L. Sebifera.
lancifolia, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.

LITSEA:—
lanuginosa, *Nees, Syst. Laurin.* 634.—Ind. or.
latifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 349.—Sumatra.
latifolia, Miq. Fl. Ind. Bat. i. 1. 973, partim = cinnamomea.
leiophylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 103.—Burma.
ligustrina, *Trimen, Cat. Pl. Ceyl.* 76.—Ind. or.
ligustrina, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 180 = L. Sebifera.
litoralis, Villar, l. c. = zeylanica.
longifolia, *Benth. & Hook. f. Gen.* iii. 161.—Zeylan.
lucida, *Blume, Bijdr.* 562.—Java.
luzonica, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
macrocarpa, Blume, Bijdr. 565 (= *Nothaphoebe cuneata*).—Malaya.
macrophylla, Blume, l. c. 567 = Actinodaphne macrophylla.
macrophylla, *Kurz, For. Fl. Brit. Burma*, ii. 305.—Burma; Penang.
magnifica, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
minuta, *Reinw. ex Meissn. in DC. Prod.* xv. 1. 194, 228.—Malaya.
monopetala, Pers. Syn. ii. 4 = polyantha.
multiflora, Blume, Bijdr. 564 = L. Sebifera.
Myrrha, Mirb. Hist. Nat. Pl. ed. II. xi. 150 = trinervia.
Neesiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 76.—Mexic.
nemoralis, *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
Noronhae, *Blume, Bijdr.* 561.—Malaya.
oblonga, *Meissn. in DC. Prod.* xv. I. 228.—Reg. Himal.
oblonga, Nees, Syst. Laurin. 678 = zeylanica.
obovata, Nees, l. c. 636 = Actinodaphne obovata.
obtusa, *Blume, Bijdr.* 564.—Java.
obtusata, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 181.—Ins. Philipp.
orbicularis, *Thw. Enum. Pl. Zeyl.* 258.—Zeylan.
ovalifolia, *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
Perrottetii, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 180.—Ins. Philipp.
piperita, Mirb. Hist. Nat. Pl. ed. II. xi. 150; Juss. ex Pers. Syn. ii. 4 = L. Cubeba.
platyphylla, Pers. Syn. ii. 4 = L. Sebifera.
plumbea, *Blume, Mus. Bot. Lugd. Bat.* i. 350.—Java.
polita, *Blume, l. c.* 348.—Java.
polyantha, *Juss. in Ann. Mus. Par.* vi. (1805) 211.—Ind. or.; Malaya.
procera, Blume, Bijdr. 560 = Actinodaphne procera.
pruinosa, Blume, l. c. = Actinodaphne procera.
pruinosa, Nees, Syst. Laurin. 636 = Cinnamomum Parthenoxylon.
Pseudoculitlawan, *Nees, l. c.* 632.—Hab. ?
puberula, Miq. Fl. Ind. Bat. i. 1. 974 = chrysocoma, plumbea.
pulchella, Meissn. in DC. Prod. xv. 1. 224 = zeylanica.
Reinwardtii, Blume, ex Meissn. l. c. 197, 228 = angulata.
resinosa, *Blume, Bijdr.* 562.—Malaya.
robusta, *Blume, l. c.* 560.—Java.
rubra, *Blume, l. c.* 561.—Java.
rubra, Miq. ex Meissn. in DC. Prod. xv. 1. 228 = Noronhae.
rufa, *Nees, Syst. Laurin.* 631.—Austral.
rugosa, *Kurz, in Flora*, lv. (1872) 172.—Penang.
Rumphii, *Villar, ex Blanco, Fl. Philipp.* ed. III. *Nov. App.* 180.—Ins. Philipp.
scrobiculata, Meissn. in DC. Prod. xv. 1. 223 = zeylanica.
Sebifera, Blume, Bijdr. 560 = L. Tetranthera.
Sebifera, *Pers. Syn.* ii. 4.—Ind. or.; Malaya.
sphaerocarpa, Blume, Bijdr. 559 = Actinodaphne sphaerocarpa.
striolata, Blume, Mus. Bot. Lugd. Bat. i. 347 = zeylanica.
tenuiramis, *Miq. Pl. Jungh.* 185 (= *Daphnidium bibracteatum*).—Java.
Tetranthera, *Mirb. Hist. Nat. Pl.* ed. II. xi. 150; *Pers. Syn.* ii. 4.—Malaya.
Thunbergii, Siebold, in Verh. Batav. Gen. xii. (1830) 24 = japonica.
timoriana, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 350.—Ins. Timor.
tomentosa, *Blume, Bijdr.* 566.—Java.
tomentosa, *Heyne, ex Wall. Cat.* sub n. 2550.—Ind. or.

LITSEA:—
trinervia, Buerg. ex Blume, Mus. Bot. Lugd. Bat. i. 352 = Lindera strychnifolia.
trinervia, *Juss. in Ann. Mus. Par.* vi. (1805) 212.—Cochinch.
trinervia, Juss. in Dict. Sc. Nat. xxvii. 79 = zeylanica.
triplinervia, *Blume, Bijdr.* 559.—Java.
umbrosa, *Nees, Syst. Laurin.* 628.—Reg. Himal.
undulata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 374 = L. Sebifera?
verticillata, *Hance, in Journ. Bot.* xxi. (1883) 356.—China.
villosa, *Blume, Mus. Bot. Lugd. Bat.* i. 349.—Amboina.
villosa, Spreng. ex Meissn. in DC. Prod. xv. 1. 148, 228 = Nectandra villosa.
Walkeri, *Trimen, Cat. Pl. Ceyl.* 76.—Zeylan.
Wightiana, *Benth. & Hook. f. Gen.* iii. 162.—Ind. or.
zeylanica, Blume, Bijdr. 559 = javanica.
zeylanica, *C. & T. Nees, Cinnam. Disp.* i. (1823) 58. t. 5.—As. et Austral. trop.
zeylanica, Thw. Enum. Pl. Zeyl. 257, ex parte = ambigua.

LITTAEA, Tagliabue, in Bibl. Ital. i. (1816) 106 = **Agave**, Linn. (Amaryll.).
geminiflora, Tagliabue, l. c. = A. geminiflora,
gracilis, Versch. Cat. n. v. (1864) = A. atrovirens?

LITTANELLA, Roth, Tent. Fl. Germ. i. 405 (1788), err. typ. = **Littorella**, Berg. (Plantag.).

LITTONIA, Hook. Bot. Mag. t. 4723 (1853). *LILIACEAE*, Benth. & Hook. f. iii. 831.
minor, *Deflers, in Bull. Soc. Bot. Fr.* xxxii. (1885) 353.—Arab.
modesta, *Hook. Bot. Mag.* t. 4723.—Afr. austr.
Revoili, *Franch. in Revoil, Çomali*, 66.—Afr. trop.
Welwitschii, *Benth. & Hook. f. Gen.* iii. 831.—Afr. trop.

LITTORELLA, Berg. in Vet. Acad. Handl. Stockh. xxix. (1768) 341. *PLANTAGINEAE*, Benth. & Hook. f. ii. 1224.
LITTANELLA, Roth, Tent. Fl. Germ. i. 405 (1788), err. typ.
australis, *Griseb. in Lechl. Berb. Am. Austr.* 53.—Chili.
flexuosa, Rafin. in New Fl. Am. iv. 12 (Quid?).—Am. bor.
juncea, *Berg. in Vet. Acad. Handl. Stockh.* xxix. (1768) 341.—Europ.
lacustris, Linn. Mant. ii. 295 = juncea.
uniflora, Aschers. Fl. Brandenb. 544 = juncea.

LIVISTONA, R. Rr. Prod. 267 (1810). *PALMAE*, Benth. & Hook. f. iii. 929.
SARIBUS, Blume, Rumphia, ii. 48. t. 95, 96 (1836).
altissima, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 150.—Ins. Sund.
australis, *Mart. Hist. Nat. Palm.* iii. 242.—Austral.
Bissula, *Mart. l. c.* 319.—Ins. Celebes.
chinensis, *R. Br. Prod.* 268.—China.
cochin-chinensis, *Blume, Rumphia*, ii. 49.—Cochinch.
Diepenhorstii, Hassk. in Bonplandia, vi. (1858) 180 = Pholidocarpus Ihur.
filifera, Hort. ex H. Wendl. in Kerch. Palm. 250 = inermis.
Gaudichaudi, Mart. Hist. Nat. Palm. iii. 242 = Pritchardia Gaudichaudii.
Hasseltii, *Hassk. ex H. Wendl. in Kerch. Palm.* 250.—Hab.?
Hoogendorpii, *Hort. Bog. ex Teijsm. & Binn. Cat. Hort. Bogor.* 71; *André, Illustr. Hortic.* xxi. (1874) 108, 121. t. 174.—Java.
humilis, *R. Br. Prod.* 268.—Austral.
inermis, *R. Br. l. c.*—Austral.
Jenkinsiana, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 334.—Reg. Himal.
Jenkinsii, Griff. ex Mart. Hist. Nat. Palm. iii. 242 = praec.
Leichhardtii, F. Muell. Fragm. v. 49; viii. 221 = humilis.
Mariae, F. Muell. l. c. 283 = humilis?
Martiana, Mart. Hist. Nat. Palm. iii. 242 = seq.
Martii, Gaudich. Bot. Voy. Bonite, t. 58, 59 = Pritchardia Martii.
mauritiana, Wall. ex Voigt, Hort. Suburb. Calc. 641, nomen = chinensis.

LIVISTOMA:—
moluccana, *Hort. ex H. Wendl. in Kerch. Palm.* 250.—Malaya.
Mueller, *Hort. l. c.*—Austral.
olivaeformis, *Mart. Hist. Nat. Palm.* iii. 319.—Java.
papuana, *Becc. Malesia*, i. (1877) 84.—N. Guin.
Ramsayi, F. Muell. Fragm. viii. 221 = inermis.
rotundifolia, *Mart. Hist. Nat. Palm.* iii. 241. t. 102.—Java.
speciosa, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 204.—Burma.
spectabilis, Griff. in Calc. Journ. Nat. Hist. v. (1845) 336 = rotundifolia.
subglobosa, *Mart. Hist. Nat. Palm.* iii. 319.—Java.

LIZERON, Rafin. Fl. Tellur. iv. 70 (1836) = **Convolvulus**, Tourn.
arvensis, Rafin. l. c. = C. arvensis.
chinensis, Rafin. l. c. = C. arvensis.
persicus, Rafin. l. c. = C. persicus.
tricolor, Rafin. l. c. = C. tricolor.
verticillatus, Rafin. l. c. = Jacquemontia hirsuta.

LLAGUNOA, Ruiz & Pav. Prod. 126. t. 28 (1794). *SAPINDACEAE*, Benth. & Hook. f. i. 409.
AMIROLA, Pers. Syn. ii. 565 (1807).
ORBIGNYA, Bert. in Merc. Chil. 737 (1829).—Cf. Am. Journ. Sc. Ser. I. xxiii. (1833) 259.
glandulosa, *G. Don, Gen. Syst.* i. 673.—Chili.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 131.—Ecuador.
nitida, *Ruiz & Pav. Syst. Veg.* 252.—Peruv.
nitida, Steud. Nom. ed. II. ii. 58 = prunifolia.
prunifolia, *H. B. & K. Nov. Gon. et Sp.* v. 131.—Ecuador.

LLANOSIA, Blanco, Fl. Filip. ed. II. 319 (1845) = **Ternstroemia**, Linn.
Toquian, Blanco, l. c. = T. Toquian.

LLAVEA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 95. *CELASTRINEAE*, Benth. & Hook. f. i. 370.
integrifolia, *Hemsl. Diagn. Pl. Nov.* 6.—Mexic.
viscosa, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 96.—Mexic.

LLERASIA, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 37 = **Vernonia**, Schreb. (Compos.).
Lindeni, Triana, l. c.—N. Granat.

LLOYDIA, Benth. & Hook. f. Gen. ii. 333 (1873) = LIOYDIA, Neck. = **Printzia**, Cass. (Compos.).

LLOYDIA, Salisb. in Trans. Hort. Soc. i. (1812) 328. *LILIACEAE*, Benth. & Hook. f. iii. 820.
CRONYXIUM, Rafin. Fl. Tellur. ii. 28 (1836).
HEMIERIUM, Rafin. l. c. 27 (1836).
NECTAROBOTHRIUM, Ledeb. Fl. Alt. ii. 36 (1830).
RHABDOCRINUM, Reichb. Consp. 65 (1828).
alpina, *Salisb. in Trans. Hort. Soc.* i. (1812) 328.—Europ.
cretica, Boiss. ex Baker, in Journ. Linn. Soc. xiv. (1874) 301 = graeca.
graeca, *Endl. Gen.* 140; *Kunth, Enum. Pl.* iv. 245.—Graec.; As. Min.
himalensis, Royle, Illustr. Bot. Himal. i. 388. t. 93. f. 2 = alpina.
kunawurensis, Royle, l. c. 3 = Gagea persica.
libanotica, Hochst. in Lorent, Wanderungen, 326 = rubroviridis.
longiscapa, Hook. Ic. Pl. t. 834 = alpina.
rubroviridis, *Baker, in Journ. Linn. Soc.* xiv. (1874) 301.—Syria.
serotina, Sweet, Hort. Brit. ed. II. 527 = alpina.
sicula, Huet, ex Baker, in Journ. Linn. Soc. xiv. (1874) 300 = alpina.
striata, Sweet, Hort. Brit. ed. II. 527 = alpina.
triflora, *Baker, in Journ. Linn. Soc.* xiv. (1874) 300.—Sibir. or.
trinervia, Coss. in Bull. Soc. Bot. Fr. xii. (1865) 279; 285 = alpina.

LLOYIDIA, Steud. Nom. ed. II. ii. 58 (1841) sphalm. = praec.

LOASA, Adans. Fam. ii. 501 (1763); Juss. Gen. 322 (1782). *LOASEAE*, Benth. & Hook. f. i. 804.
HUIDOBRIA, C. Gay, Fl. Chil. ii. 438. t. 26 (1846).
ILLAIREA, Lenne & Koch, in Verh. Berl. Gartenb. Ver. N. Reihe, i. (1853) 397.

LOASA:—
Ochetocarpus, Meyen, Reise, i. 310 (1834).
Ortiga, Neck. Elem. ii. 400 (1790).
acanthifolia, *Lam. Encyc.* iii. 579.—Chili.
acanthifolia, Lindl. Bot. Reg. t. 785 = Placei.
acerifolia, *Domb. ex A. Juss. in Ann. Mus. Par.* v. (1804) 24. t. 1. f. 2.—Chili.
acuminata, *Wedd. Chlor. And.* ii. 220.—Ecuador.
alba, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 192. —Chili.
Aldunatea, *C. Gay, Fl. Chil.* ii. 442.—Chili.
ambrosiaefolia, Juss. in Ann. Mus. Par. v. (1804) 26. t. 4. f. 1 = hispida.
argemonoides, *Juss. l. c.*—N. Granat.
Arnottiana, *C. Gay, Fl. Chil.* ii. 442.—Chili.
atriplicifolia, *Presl, Symb. Bot.* i. 61.—Am. austr.
aurantiaca, Hort. ex Nichols. Dict. Gard. ii. 289 = lateritia.
Barneoudii, *C. Gay, Fl. Chil.* ii. 452.—Chili.
Bergii, *Hieron. in Bull. Acad. Nac. Córdova*, iii. (1879) 349.—Reg. Argent.
bicolor, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 361.—Am. trop.
Blacei, Ruschen. in Am. Journ. Sc. xxiii. (1833) 94, err. typ. = Placei.
bryoniaefolia, *Schrad. ex DC. Prod.* iii. 341.—Chili.
calycina, *Benth. Pl. Hartw.* 132.—Ecuador.
canarinoides, *Nichols. Dict. Gard.* ii. 289.—Am. centr.
carduifolia, Ball, in Journ. Linn. Soc. xxii. (1885) 39, in syn. = Blumenbachia carduifolia.
chelidonifolia, *Benth. Bot. Voy. Sulph.* 101.—Peruv.
chenopodiifolia, *Lam. Encyc.* iii. 550.—Peruv.
chuquitensis, Meyen, Reise, i. 483 = Blumenbachia chuquitensis.
cirsiifolia, Kunze, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 338 = sclareaefolia.
coccinea, Hort. ex Loud. Encyc. Pl. Suppl. i. 1246 = lateritia.
contorta, Lam. Encyc. iii. 579 = Blumenbachia contorta.
coronata, Gill. ex Arn. in Edinb. Journ. Geogr. Sc. iii. (1831) 274 = Blumenbachia coronata.
decorticans, *Steud. Nom.* ed. II. ii. 58.—Chili.
dissecta, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 330.—Chili.
elongata, *Hook. & Arn. l. c.* 329.—Chili.
filicifolia, *Poepp. Fragm. Syn. Pl. Chil.* 23.—Chili.
floribunda, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 329.—Chili.
furcata, *Phil. in Anal. Univ. Chil.* (1865) II. 328.—Chili.
grandiflora, Lam. Encyc. iii. 580 = Blumenbachia contorta.
helicteroides, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 339 = Blumenbachia scandens.
heptamera, *Wedd. Chlor. And.* ii. 218.—Bolivia.
Herbertii, Paxt. ex Loud. Hort. Brit. Suppl. iii. 579 = *Caiophora Herberti*.
heterophylla, *Bert. in Merc. Chil.* (1829); *ex Am. Journ. Sc.* xxiii. (1833) 94 = seq.?
heterophylla, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 328.—Chili.
hibiscifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 138.—Reg. Argent.
hispida, *Linn. Syst.* ed. XII. 364.—Peruv.
illapelina, *Phil. in Anal. Univ. Chil.* (1863) ii. 380.—Chili.
incana, *R. Grah. in Edinb. N. Phil. Journ.* (Oct.–Dec. 1830) 169.—Peruv.
insons, Poepp. Fragm. Syn. Pl. Chil. 24 = Meyeniana.
intricata, *C. Gay, Fl. Chil.* ii. 455.—Chili.
lateritia, *Gill. ex Arn. in Edinb. Journ. Geogr. Sc.* iii. (1831) 275.—Chili.
Lechleriana, Miq. in Lechl. Berb. Am. Austr. 55 = acerifolia.
leucantha, *Bert. ex Steud. Nom.* ed. II, ii. 58 (Quid?). —Chili.
longiseta, *Phil. in Anal. Univ. Chil.* (1856) II. 347.—Chili.
loxensis, *H. B. & K. Nov. Gen. et Sp.* vi. 116.—Peruv.
magnifica, *Warsz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292.—Bolivia; Peruv.
malesherbioides, *Phil. in Linnaea*, xxxiii. (1864–65) 74.—Chili.
Martini, *Phil. in Anal. Univ. Chil.* (1872) 716.—Chili.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 338.—Chili.
micrantha, *Poepp. Fragm. Syn. Pl. Chil.* 24.—Chili.

LOASA:—
multifida, *C. Gay, Fl. Chil.* ii. 457.—Chili.
muralis, Griseb. in Goett. Abh. xix. (1874) 151 = Blumenbachia insignis.
nitida, Hook. Bot. Mag. f. 2372 = acanthifolia.
nitida, *Lam. Encyc.* iii. 581.—Peruv.; Chili.
Ochagaviae, *Phil. in Linnaea*, xxviii. (1856) 641.—Chili.
pallida, *Gill. ex Arn. in Edinb. Journ. Georgr. Sc.* iii. (1831) 274.—Chili.
palmata, Spreng. Syst. ii. 601 = Blumenbachia insignis.
papaverifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 118.—N. Granat.
parviflora, *Schrad. Pl. Rar. Bras. ex DC. Prod.* iii. 342.—Bras.
patula, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1827) 174 = Blumenbachia insignis.
pedicularifolia, Meyen, Reise, i 310 = volubilis.
peduncularis, *Presl, Symb. Bot.* ii. 41.—Peruv.
Pentlandii, Paxt. Mag. Bot. ix. (1842) 7 (= *Caiophora Pentlandii*).—Peruv.
picta, *Hook. Bot. Mag.* t. 4428.—Peruv.
pilosa, Meyen, Reise, i. 470 = Blumenbachia coronata.
pinnatifida, Gill. ex Arn. in Edinb. Journ. Geogr. Sc. iii. (1831) 275 = filicifolia.
Placei, Bert.—Cf. Am. Journ. Sc. xxiii. (1833) 94 = nitida.
Placei, *Lindl. in Trans. Hort. Soc.* vi. (1826) 95.—Chili.
prostrata, *Gill. ex Arn. in Edinb. Journ. Geogr. Sc.* iii. (1831) 274.—Chili.
punicea, *Phil. in Anal. Univ. Chil.* (1884), reimpr. 4.—Chili.
ranunculifolia, *Humb. & Bonpl. Pl. Aequin.* i. 50. t. 14.—Peruv.
Remyi, *C. Gay, Fl. Chil.* ii. 462.—Chili.
rhoeadifolia, *Schlecht. in Linnaea*, xiv. (1840) 382.—Mexic.
rosulata, *Wedd. Chlor. And.* ii. 219.—Peruv.
rudis, *Benth. Pl. Hartw.* 75.—Ecuador.
Ruiziana, *G. Don, Gen. Syst.* iii. 64.—Peruv.
rupestris, *Gardn. in Hook. Ic. Pl.* t. 663.—Bras.
sagittata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 328.—Chili.
scandens, Meyen, Reise, i. 310 = Blumenbachia scandens.
Schlimiana, *Planch. & Linden, in Fl. des Serres*, x. (1854–55) 89.—Venezuela.
sclareaefolia, *Juss. in Ann. Mus. Par.* v. (1804) 25. t. 1. f. 1.—Chili.
Simoni, *Phil. in Linnaea*, xxx. (1859–60) 188.—Chili.
solanifolia, *C. Gay, Fl. Chil.* ii. 449.—Chili.
stenocarpa, Poepp. Fragm. Syn. Pl. Chil. 23 = Grammatocarpus volubilis.
striata, Meyen, Reise, i. 310 = Grammatocarpus volubilis.
tricolor, *Lindl. Bot. Reg.* t. 667.—Chili.
tricolor, *Weinm. in Syll. Ratisb.* ii. (1828) 21.—Chili.
triloba, *Juss. in Ann. Mus. Par.* v. (1804) 24. t. 1. f. 3.—Peruv.
triphylla, *Juss. l. c.* 27. t. 5. f. 2.—Peruv.
urens, Jacq. Obs. Bot. ii. 15 = hispida.
urens, Vell. Fl. Flum. 225. t. 98 = Blumenbachia latifolia.
Urmenetae, *Phil. in Anal. Univ. Chil.* (1872) 714.—Chili.
urticaefolia, *Bert. ex Steud. Nom.* ed. II. ii. 58 (Quid?). —Chili.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1872) 715.—Chili.
volubilis, Hort. ex G. Don, in Loud. Hort. Brit. 321 = Grammotocarpus volubilis.
volubilis, *Juss. in Ann. Mus. Par.* v. (1804) 26. t. 5. f. 1.—Chili.
vulcanica, *André, in Illustr. Hortic.* xxv. (1878) t. 302. —N. Granat.
Wallisii, Hort. ex Regel, Gartenfl. (1878) 357 = vulcanica.
xanthiifolia, *Juss. in Ann. Mus. Par.* v. (1804) 26. t. 2. f. 1.—Peruv.

LOBADIUM, Rafin. in Am. Monthly Mag. (1819) 357; et in Journ. Phys. lxxxix. (1819) 98 = **Rhus**, Linn. (Anacard.).
amentaceum, Rafin. in Am. Monthly Mag. (1819) 358 = R. aromatica.

LOBADIUM:—
aromaticum, Steud. Nom. ed. I. 689, in syn. = Rhus aromatica.
suaveolens, Rafin. in Journ. Phys. lxxxix. (1819) 98 = R. aromatica.
trifoliatum, Rafin. Cat. 14 = R. aromatica.

LOBAKE, Rafin. Fl. Tellur. iv. 81 (1836) = **Ipomoea**, Linn. (Convolv.).
guyanensis, Rafin. l. c. = I. guyanensis.

LOBARIA, Haw. Enum. Saxifr. 20 (1821) = **Saxifraga**, Tourn.
cernua, Haw. l. c. = S. cernua.
cuneifolia, Haw. l. c. 19 = S. cuneifolia.
hederacea, Haw. l. c. 18 = S. hederacea.
orientalis, Haw. l. c. = S. Cymbalaria.
paradoxa, Haw. l. c. 18, 20 = S. paradoxa.
reticulata, Haw. l. c. = S. Cymbalaria.
rivularis, Haw. l. c. 19 = S. rivularis.
sibirica, Haw. l. c. 21 = S. sibirica.

LOBBIA, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 144. t. 3 = **Thottea**, Rottb. (Aristol.).
dependens, Planch. l. c. = T. dependens.

LOBELIA, Adans. Fam. ii. 157 (1763) = **Scaevola**, Linn. (Goodeniac.).

LOBELIA, Plum. ex Linn. Gen. ed. I. 267 (1737).
CAMPANULACEAE, Benth. & Hook. f. ii. 551.
Dobrowskya, Presl, Prod. Monog. Lobel. 8 (1836).
Dortmanna, Linn. Syst. ed. I (1735).
Grammatotheca, Presl, Prod. Monog. Lobel. 48 (1836).
Holostigma, G. Don, Gen. Syst. iii. 716 (1834).
Holostigmateia, Reichb. Nom. 102 (1841).
Isolobus, A. DC. in DC. Prod. vii. 352 (1839).
Juchia, Neck. Elem. i. 132 (1790).
Mecoschistum, Dulac, Fl. Hautes-Pyr. 459 (1867).
Mezleria, Presl, Prod. Monog. Lobel. 7. (1836).
Monopsis, Salisb. in Trans. Hort. Soc. ii. (1817) 37. t. 2.
Parastranthus, G. Don, Gen. Syst. iii. 716 (1834).
Pratia, G. Don, Gen. Syst. iii. 699 (1834), partim.
Rapuntium, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768).
Rhynchopetalum, Fresen. in Flora, xxi. (1838) II. 603.
Stooria, Neck. Elem. i. 132 (1790).
Trimeris, Presl, Prod. Monog. Lobel. 46 (1836).
Tupa, G. Don, Gen. Syst. iii. 700 (1834).
Tylomium, Presl, Prod. Monog. Lobel. 31 (1836).
Ymnostema, Neck. Elem. i. 132 (1790).
acuminata, *Sw. Prod. Veg. Ind. Occ.* 115.—Ind. occ.
acutangula, *A. DC. in DC. Prod.* vii. 369.—Afr. austr.
acutidens, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 204.—Afr. trop.
adenophora, Roem. & Schult. Syst. v. Index 615 = Lysipoma aretioides.
adscendens, De Vriese, in Lehm. Pl. Preiss. i. 395 = tenuior.
affinis, Mirb. Hist. Nat. Pl. ed. II. [1805] xiv. 237 = Siphocampylus campanulatus.
affinis, *Wall. Cat.* n. 1311.—As. trop.
alata, Hayne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii, 425 = zeylanica.
alata, Labill. Nov. Holl. Pl. i. 51. t. 72 = anceps.
alata, Leperr. & Bout. ex A. DC. in DC. Prod. vii. 368 = madagascariensis.
algoënsis, A. DC. in DC. l. c. 369 = L. Erinus.
alpina, *Vell. Fl. Flum.* viii. t. 154.—Bras.
alsinoides, *Lam. Encyc.* iii. 588.—Afr. austr.
alyssifolia, Salisb. Ic. 19 = pubescens.
ambigua, Cham. in Linnaea, viii. (1833) 221 = Rollandia ambigua.
americana, Willd. ex A. DC. in DC. Prod. vii. 387, 507 = Scaevola Lobelia.
amoena, *Michx. Fl. Bor. Am.* ii. 152.—Am. bor.
amplexicaulis, De Vriese, in Lehm. Pl. Preiss. i. 397 = Bergiana.
amygdalina, *Willd. ex Roem. & Schult. Syst.* v. 57.—Am. austr.
anceps, *Linn. f. Suppl.* 395.—Reg. temp. austr.
andina, *Benth. Pl. Hartw.* 213.—N. Granat.
Andropogon, Cav. Ic. vi. 10 = Centropogon Andropogon.

LOBELIA :—
androsacea, Willd. ex Roem. & Schult. Syst. v. 41 = Lysipoma acaulis.
angulata, Forst. f. Prod. 58 = Pratia angulata.
angulato-dentata, Hook. & Arn. Bot. Beech. Voy. 301 = laxiflora.
angustifolia, Benth. in Enum. Pl. Hueg. 74 = anceps.
angustifolia, Cham. in Linnaea, viii. (1833) 219.—(= *Delissea angustifolia*).—Ins. Sandvic.
antisyphilitica, *Hayne, Arzn. Gew.* xiii. t. 9.—Am. bor.
aphylla, *Nutt. in Am. Journ. Sc.* v. (1822) 297 (Quid?).—Am. bor.
aphylla, Spreng. ex Presl, Prod. Monog. Lobel. 52 (sp. dub.).—Afr. austr.
appendiculata, *A. DC. in DC. Prod.* vii. 376.—Am. bor.
aquatica, *Cham. in Linnaea*, viii. (1833) 211.—Bras.
arabidea, Steud. Nom. ed II. ii. 59 = Laurentia arabidea.
arabidioides, Hook. & Arn. Bot. Beech. Voy. 301. t. 66 = Heterotoma arabidioides.
arborea, Forst. f. Prod. 58 = Sclerotheca arborea.
arenaria, Salzm. ex A. DC. in DC. Prod. vii. 387, 421 = Cephalostigma Perrottetii.
arenarioides, A. DC. l. c. 367 = trigona.
areolata, Rich. ex Juss. in Ann. Mus. Par. xviii. (1811) 3 = stricta.
arguta, Hort. ex A. DC. in DC. Prod. vii. 370 = hederacea.
arguta, Lindl. in Bot. Reg. t. 973 = salicifolia.
aromatica, Moon, Cat. Pl. Ceyl. 14 = excelsa.
ascendens, *De Vriese, in Lehm. Pl. Preiss.* i. 395.—Austral.
asclepiadea, Willd. ex Roem. & Schult. Syst. v. 57 = Siphocampylus ferrugineus.
aspera, *Spreng. Neue Entdeck.* i. 222.—Afr. austr.
asperulata, Klotzsch, in Peters, Reise Mossamb. Bot. 300 = fervens.
asplenifolia, Hort. ex A. DC. in DC. Prod. vii. 387, 412 = Isotoma axillaris.
assurgens, *Linn. Syst.* ed. X. 1237.—Ind. occ.
attenuata, Presl, Prod. Monog. Lobel. 34 = Siphocampylus attenuatus.
ayavensis, Willd. ex Spreng. Syst. i. 709 = Siphocampylus umbellatus.
barbata, Cav. Ic. vi. 12 = Centropogon barbatus (in Addend.).
begonifolia, Wall. in As. Res. xiii. (1820) 377 = Pratia begonifolia.
bellidiflora, *Linn. f. Suppl.* 396.—Afr. austr.
bellidifolia, Thunb. Prod. Pl. Cap. 40 = praec.
Benthami, *F. Muell. Census*, 85.—Austral.
Bergiana, *Cham. in Linnaea*, viii. (1833) 217.—Afr. austr.; Austral.
Berlandieri, *A. DC. in DC. Prod.* vii. 367.—Mexic.
Berlandieri, Torr. Bot. Mex. Bound. 107 = Cliffortiana.
Berteriana, Spreng. Syst. i. 712 = Centropogon Berterianus.
beslerioides, H. B. & K. Nov. Gen. et Sp. iii. 306 = Centropogon beslerioides.
betulaefolia, Cham. in Linnaea, viii. (1833) 204 = Siphocampylus betulaefolius.
bicolor, Boj. Hort. Maurit. 193 = serpens.
bicolor, Sims, Bot. Mag. t. 514 = L. Erinus.
bidentata, Donn, Hort. Cantab. ed. IV. 41 = tenella.
bifida, Thunb. Prod. Pl. Cap. 40 = Laurentia Dregeana.
biserrata, Cav. Ic. vi. 10 = Siphocampylus biserratus.
biserrata, Ruiz & Pav. ex A. DC. in DC. Prod. vii. 397 = Siphocampylus laciniatus.
Bivonae, Tineo, Cat. Pl. Hort. Panorm. (1827) 279 = Laurentia tenella.
Boivinii, *Sond. in Harv. & Sond. Fl. Cap.* iii. 546.—Afr. austr.
Bonplandiana, Roem. & Schult. Syst. v. 57 = Centropogon Bonplandianus.
Boykinii, *Torr. & Gray, ex A. DC. in DC. Prod.* vii. 374.—Am. bor.
bracteolata, *A. DC. l. c.* 371.—Hab.?
bracteolata, *Vatke, in Linnaea*, xxxviii. (1874) 721.—Venezuela.
Brandtii, Hort. ex A. DC. in DC. Prod. vii. 393 = salicifolia.
brevifolia, *Nutt. ex A. DC. l. c.* 377.—Am. bor.
Breynii, *Lam. Encyc.* iii. 588.—Afr. austr.
Breynii, Spreng. ex Eckl. & Zeyh. Enum. 397 = L. Erinus.

LOBELIA :—
Bridgesii, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 278.—Chili.
Broussonetia, Bory, Ess. Isles Fortun. 330 = Wahlenbergia lobelioides.
Browniana, Roem. & Schult. Syst. v. 71 = gibbosa.
bryoides, Willd. l. c. 41 = Arenaria dicranoides.
bulbosa, Linn. Sp. Pl. 933 = Cyphia bulbosa.
Burmanniana, *D. Dietr. Syn. Pl.* i. 728.—Afr. austr.
caerulea, Hook. Bot. Mag. t. 2701 = coronopifolia.
caespitosa, Blume, Bijdr. 729 = radicans.
calcarata, Bertol. Fl. Guatim. 9 = Heterotoma lobelioides.
calycina, Cham. in Linnaea, viii. (1833) 222 = Rollandia calycina.
campanulata, Cav. Ic. vi. 11 = Siphocampylus campanulatus.
campanulata, *Lam. Encyc.* iii. 588.—Afr. austr.
campanuloides, Thunb. in Trans. Linn. Soc. ii. (1794) 331 = radicans.
camporum, *Pohl, Bras. Pl. Ic.* ii. 100.—Bras.
camtschatica, Pall. ex Lamb. in Trans. Linn. Soc. x. (1811) 260 = sessilifolia.
cana, Presl, Prod. Monog. Lobel. 37 = Siphocampylus macropodus.
canariensis, Hort. Par. ex A. DC. in DC. Prod. vii. 388 = Wahlenbergia lobelioides.
Canbyi, *A. Gray, Man. Bot. N. U. St.* ed. V. 284.—Am. bor.
candida, Hort. ex Link, Enum. Hort. Berol. i. 219 = L. Erinus.
canescens, Presl, Prod. Monog. Lobel. 38 = Siphocampylus canescens.
Caoutchouc, H. B. & K. Nov. Gen. et Sp. iii. 304 = Siphocampylus Caoutchouc.
Caoutschouk, Steud. Nom. ed. II. ii. 59 = praec.
capillifolia, *A. DC. in DC. Prod.* vii. 362.—Afr. austr.
capitata, Burm. f. Fl. Cap. Prod. 29. (err. typ. 25) = triquetra.
Cardamines, Thunb. Prod. Pl. Cap. 39 = Cyphia Cardamines.
cardinalis, *Linn. Sp. Pl.* 930.—Am. bor.
cardiophylla, Presl, Prod. Monog. Lobel. 37 = Siphocampylus corymbiferus.
carnosula, Hook. & Arn. Bot. Beech. Voy. 362 = Laurentia carnosula.
Cautschuk, Willd. ex Roem. & Schult. Syst. v. 58 = Siphocampylus Caoutchouc.
Cavanillesiana, Roem. & Schult. Syst. v. 43 = laxiflora.
Cavanillesii, Mart. Ausw. Pfl. Muench. 12. t. 9 = praec.
ceratophylla, A. Dietr. Syn. Pl. i. 728 = tomentosa.
chamaedryfolia, *A. DC. in DC. Prod.* vii. 364.—Afr. austr.
Chamaepitys, Lam. Encyc. iii. 590 = tomentosa.
Chamaepitys, Spreng. Syst. i. 714 = aspera.
Chamissoniana, Presl, Prod. Monog. Lobel. 36 = Siphocampylus Westinianus.
Cheiranthus, Linn. Sp. Pl. 933 = Manulea Cheiranthus.
chenopodiifolia, Wall. Cat. n. 1312 = Cliffortiana.
chinensis, *Lour. Fl. Cochinch.* 514.—China.
chireensis. *A. Rich. Tent. Fl. Abyss.* ii. 6.—Abyssin.
ciliata, De Vriese, in Lehm. Pl. Preiss. i. 397 = tenuior.
ciliata, Spreng. f. ex Presl, Prod. Monog. Lobel. 52 (Prismatocarpi sp.).—Hab.?
cinerea, *Thunb. Prod. Pl. Cap.* 40 = Lightfootia grisea.
circaeoides, *A. DC. ex DC. Prod.* vii. 379.—Mexic.
cirsiifolia, *Lam. Encyc.* iii. 584.—Ins. Cap. Virid.
cladlomesa, Rafin. New Fl. Am. ii. 17 (Quid?).—Am. bor.
Claytoniana, Michx. Fl. Bor. Am. ii. 153 = spicata.
Clermontiana, Gaudich. in Freyc. Voy. Bot. 459 = Clermontia persicifolia.
Cliffortiana, *Linn. Sp. Pl.* 931.—Am. trop.
Cliffortiana, Michx. Fl. Bor. Am. ii. 152 = praec.?
coccinea, Stokes, Bot. Mat. Med. i. 343 = cardinalis.
coelestis, Nutt. ex Loud. Hort. Brit. 592 = fenestralis.
coerulea, Auct. = *caerulea*.
collina, *H. B. & K. Nov. Gen. et Sp.* iii. 312.—Peruv.
colorata, Sweet, Brit. Flow. Gard. Ser. II. t. 180 = amoena.
colorata, *Wall. Pl. As. Rar.* ii. 42.—Reg. Himal.
Columnae, Linn. f. Suppl. 393 = Siphocampylus Columnae.
columnaris, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 14.—Afr. trop.
commutata, Roem. & Schult. Syst. v. 36 = neglecta.
commutata, Roem. & Schult. l. c. 73 = pauciflora.

LOBELIA :—
comosa, Cav. Ic. vi. 9. t. 512. f. 2 = Siphocampylus comosus.
comosa, Linn. Sp. Pl. ed. II. 1323 = triquetra.
concolor, R. Br. Prod. 563 = Pratia erecta.
concolor, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1842), reimp. 15.—Mexic.
conglobata, *Lam. Encyc.* iii. 585.—Ins. Martinic.
conglomerata, Sieber, ex Steud. Nom. ed. II. ii. 59 = praec.
convolvulacea, Cham. in Linnaea, viii. (1833) 205 = Siphocampylus convolvulaceus.
cordata, Willd. ex Roem. & Schult. Syst. v. 308 = Siphocampylus volubilis.
cordifolia, Hook. & Arn. Bot. Beech. Voy. 301 = Heterotoma tenella.
cordigera, *Cav. Ic.* vi. 14. t. 523.—Chili.
cornuta, Linn. Sp. Pl. 930 = Centropogon surinamensis.
coronopifolia, *Linn. l. c.* 933.—Afr. austr.
corymbifera, Presl, Prod. Monog. Lobel. 37 = Siphocampylus corymbiferus.
corymbosa, Berg. Desc. Pl. Cap. 344 = triquetra.
corymbosa, *R. Grah. in Edinb. N. Phil. Journ.* (Jul.-Oct. 1826) 385.—Afr. austr.
crassiuscula, Michx. Fl. Bor. Am. ii. 152 = glandulosa.
crenata, Presl, Prod. Monog. Lobel. 37 = Siphocampylus crenatifolius.
crenata, Thunb. Prod. Pl. Cap. 39 = Cyphia crenata.
cretica, Herb. Juss. ex A. DC. in DC. Prod. vii. 410 = Laurentia tenella.
crispa, Gaudich. in Freyc. Voy. Bot. 459; D. Dietr. Syn. Pl. i. 728 = Rollandia crispa.
crispa, R. Grah. in Edinb. N. Phil. Journ. (Apr.-July 1826) 173 = fenestralis.
cuneifolia, *Link & Otto, Ic. Pl. Sel. Berol.* t. 39.—Afr. austr.
cuneiformis, Labill. Nov. Holl. Pl. i. 51. t. 73 = anceps.
Cymbalaria, *Griseb. in Goett. Abh.* xix. (1874) 200.—Reg. Argent.
Cyphia, J. F. Gmel. Syst. 357 = Cyphia bulbosa.
Cyphia, Thunb. Prod. Pl. Cap. 39 = Cyphia digitata.
cyphioides, *Harv. Thes. Cap.* ii. 40.—Afr. austr.
Davidi, *Franch. Pl. Davidian.* i. (1884) 191.—China.
debilis, Eckl. ex A. DC. in DC. Prod. vii. 370 = triquetra.
debilis, *Linn. f. Suppl.* 395.—Afr. austr.
decipiens, *Sond. in Harv. & Sond. Fl. Cap.* iii. 540.—Afr. austr.
Deckenii, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 466.—Afr. trop.
decumbens, Juss. ex A. DC. in DC. Prod. vii. 388 = Siphocampylus decumbens.
decumbens, *Rich. ex Roem. & Schult. Syst.* v. 67.—Ind. occ.
decumbens, Sims, Bot. Mag. t. 2277 = anceps.
decurrens, *Cav. Ic.* vi. 13. t. 521.—Chili.
decurrens, Roth, Nov. Pl. Sp. 145 = zeylanica.
decurrens, Willd. ex Spreng. Syst. i. 718 = madagascariensis.
Delisseana, Gaudich. in Freyc. Voy. Bot. 437 (= *Delissea acuminata*).—Ins. Sandvic.
densiflora, *Rennie, Mag. Gard. & Bot.* i. (1833) t. 67; *ex Paxt. Mag. Bot.* xvi. (1849) 66.—Hab.?
dentata, *Cav. Ic.* vi. 14. t. 522.—Austral.
dentata, Sieber, ex A. DC. in DC. Prod. vii. 364 = gracilis.
dentata, Torr. ex A. DC. l. c. 377 = brevifolia.
dentata, Willd. ex Roem. & Schult. Syst. v. 57 = Centropogon Bonplandianus.
dependens, Ruiz & Pav. ex A. DC. in DC. Prod. vii. 388, 399 = Siphocampylus dependens.
depressa, *Linn. f. Suppl.* 395.—Afr. austr.
depressa, Thunb. ex Harv. & Sond. Fl. Cap. iii. 551 = Laurentia repens.
dichotoma, Miq. Fl. Ind. Bat. ii. 576 = zeylanica.
digitata, Thunb. 'in Act. Gorenk. v. (1811) 62.'—Cf. Fl. Cap. ii. 50 = Cyphia digitata.
dioica, *R. Br. Prod.* 565.—Austral.
discolor, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Hort. Berol. i. t. 2 = subnuda.
divaricata, *Hook. & Arn. Bot. Beech. Voy.* 301. t. 67.—Mexic.
divaricata, Seem. Bot. Voy. Herald, 316 = Heterotoma intermedia.
diversifolia, Cham. in Linnaea, viii. (1833) 209 = Wahlenbergia linarioides.

LOBELIA :—
diversifolia, *Roem. & Schult. Syst.* v. 68.—Am. austr.
Dobrowskiana, D. Dietr. Syn. Pl. i. 732 = tenella.
Dombeyana, Presl, Prod. Monog. Lobel. 39 = Siphocampylus Dombeyanus.
domingensis, *A. DC. in DC. Prod.* vii. 359.—Ins. S. Doming.
dopatroides, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 77 = Griffithii.
Dortmanna, *Linn. Sp. Pl.* 929.—Europ.; Am. bor.
Draba, Willd. ex Roem. & Schult. Syst. v. 67 = trigona.
dracunculoides, Willd. l. c. 56 = persicaefolia.
Dregeana, *A. DC. in DC. Prod.* vii. 371.—Afr. austr.
dubia, *De Vriese, in Lehm. Pl. Preiss.* ii. 242.—Austral.
Eckloniana, *D. Dietr. Syn. Pl.* i. 736.—Afr. austr.
edulis, Presl, Prod. Monog. Lobel. 48 = Centropogon surinamensis.
Ehrenbergii, *Vatke, in Linnaea*, xxxviii. (1874) 719.—Mexic.
elegans, De Vriese, in Lehm. Pl. Preiss. i. 396 = Isotoma pusilla.
elliptica, *Willd. ex Roem. & Schult. Syst.* v. 57.—Am. austr.
ensiformis, *Vell. Fl. Flum.* viii. t. 155.—Bras.
erecta, De Vriese, in Lehm. Pl. Preiss. i. 395 = anceps.
erecta, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 28.—Reg. Himal.
ericetorum, *A. DC. in DC. Prod.* vii. 784.—Afr. austr.
ericoides, *Presl, ex D. Dietr. Syn. Pl.* i. 733.—Afr. austr.
erina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Erinus.
erinifolia, Salisb. Prod. 129 = Laurentia erinoides.
erinoides, Linn. Sp. Pl. 932 = Laurentia erinoides.
erinoides, Sickm. ex A. DC. in DC. Prod. vii. 408 = repens.
erinoides, Thunb. Prod. Pl. Cap. 40 = radicans.
Erinus, *Linn. Sp. Pl.* 932.—Afr. austr.
Erinus, Thunb. Fl. Cap. 179 = radicans.
euphrasioides, *Steud. Nom.* ed. II. ii. 60.—Afr. austr.
eurostos, *Voigt, Hort. Suburb. Calc.* 367.—Ind. or.
exaltata, *Pohl, Pl. Bras. Ic.* ii. 101. t. 166 = Haynaldia exaltata.
excelsa, Bonpl. Desc. Jard. Malm. 112 = salicifolia.
excelsa, *Leschen. in Roxb. Fl. Ind. ed. Carey*, ii. 114.—Ind. or.
exigua, Pers. Syn. ii. 212 = campanulata.
exilis, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 7.—Afr. trop.
falcata, Rafin. New Fl. Am. ii. 18 = Kalmii?
Farsetia, Vand. Fasc. Pl. 19 = urens.
fastigiata, *H. B. & K. Nov. Gen. et Sp.* iii. 313.—Ecuador.
Feayana, *A. Gray, in Proc. Am. Acad.* xii. (1877) 60.—Am. bor.
fenestralis, *Cav. Ic.* vi. 8. t. 512. f. 1.—Mexic.
ferruginea, Linn. f. Suppl. 394 = Siphocampylus ferrugineus.
fervens, *Thunb. Fl. Cap.* ii. 46.—Afr. austr.
Feuillei, Hort. ex Nichols. Dict. Gard. ii. 291 = L. Tupa.
filiformis, Lam. Encyc. iii. 588 = serpens.
filiformis, Sieber, ex A. DC. in DC. Prod. vii. 372 = Cliffortiana.
fissa, Willd. ex Roem. & Schult. Syst. v. 57 = laxiflora.
fistulosa, Rafin. New Fl. Am. ii. 18 (Quid?).—Am. bor.
fistulosa, *Vell. Fl. Flum.* viii. t. 157.—Bras.
flaccida, A. DC. in DC. Prod. vii. 360 = L. Erinus.
flava, *D. Dietr. Syn. Pl.* i. 734.—Afr. austr.
Fleuryana, Gaudich. in Freyc. Voy. Bot. 456 = Isotoma axillaris.
flexuosa, *A. DC. in DC. Prod.* vii. 379.—Mexic.
floribunda, Bonpl. ex A. DC. l. c. 383 = laxiflora.
floridana, Chapm. in Coult. Bot. Gaz. iii. (1878) 9 = paludosa.
fluminensis, *Vell. Fl. Flum.* viii. t. 159.—Bras.
fluviatilis, R. Br. Prod. 563 = Isotoma fluviatilis.
foetida, Willd. ex H. B. & K. Nov. Gen. et Sp. iii. 304 = Siphocampylus foetidus.
foliosa, H. B. & K. l. c. 310 = decurrens.
formosa, Roth, in Roem. & Schult. Syst. v. 50 = fulgens.
frutescens, Mill. Gard. Dict. ed. VIII. n. 1 = Scaevola Koenigii.
fruticosa, *Linn. f. Suppl.* 395.—Afr. austr.
fulgens, *Willd. Hort. Berol.* t. 85.—Mexic.

LOBELIA :—
fusca, Presl, Prod. Monog. Lobel. 40 = Siphocampylus fuscus.
Gardneriana, *Kanitz, in Mart. Fl. Bras.* vi. IV. 138.—Bras.
Gasparrini, Tineo, Cat. Pl. Panorm. (1827) 279 = Laurentia Michelii.
Gattingeri, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 221.—Am. bor.
Gaudichaudii, *A. DC. in DC. Prod.* vii. 384.—Ins. Sandvic.
gelida, F. Muell. Fragm. iv. 183 = Pratia gelida.
genistoides, *A. DC. in DC. Prod.* vii. 358.—Afr. austr.
Ghiesbreghtii, Decne. in Rev. Hortic. Sér. III. ii. (1848) 341 (Centropogonis sp.?).—Mexic.
Ghiesbreghtii, *Lem. Illustr. Hortic.* i. (1854) t. 34.—Mexic.; Bras.
Gibberoa, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 465.—Afr. trop.
gibbosa, *Labill. Nov. Holl. Pl.* i. 50. t. 71.—Austral.
gigantea, Cav. Ic. vi. 9 = Siphocampylus giganteus.
gigantea, Sims, Bot. Mag. t. 1325 = salicifolia.
glabrata, H. B. & K. Nov. Gen. et Sp. iii. 307 = Siphocampylus glabratus.
glabrifolia, E. Mey. ex Presl, Prod. Monog. Lobel. 12 = leptocarpa.
glandulosa, Dennis, ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 646 = syphilitica.
glandulosa, Engelm. & A. Gray, Pl. Lindh. i. 14 = puberula.
glandulosa, *Walt. Fl. Carol.* 218.—Am. bor.
glandulosa, Willd. in Roem. & Schult. Syst. v. 41 = Lysipoma aretioides.
globosa, *R. Br. ex G. Don, in Loud. Hort. Brit.* 78, nomen.—Hab.?
goodenioides, Willd. Hort. Berol. t. 30 = spicata.
gracilis, *Andr. Bot. Rep.* t. 340.—Afr. austr.; Austral.
gracilis, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 206.—Ins. Galapag.
gracilis, Nutt. Gen. Am. ii. 77 = Nuttallii.
gracilis, Salisb. Prod. 129 = Laurentia Michelii.
graminea, *Lam. Encyc.* iii. 538.—Peruv.
grandidentata, Schlecht. in Linnaea, ix. (1834) 262 = Siphocampylus grandidentatus.
grandiflora, Gaudich. in Freyc. Voy. Bot. 459 = Clermontia grandiflora.
grandis, Linn. f. Suppl. 394 = Centropogon grandis.
gratioloides, Roxb. ex Presl, Prod. Monog. Lobel. 50 = trigona.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 28.—Burma.
Grimesiana, Gaudich. in Freyc. Voy. Bot. 458 = Cyanea Grimesiana.
gruina, *Cav. Ic.* vi. 8. t. 511.—Mexic.
Gulparae, Buch.-Ham. ex Wall. Cat. n. 1311 = affinis.
Haenkeana, *A. DC. in DC. Prod.* vii. 382.—Ins. Philipp.
Hartlaubi, *Buchen. in Bremen, Abh.* vii. (1881) 201.—Madag.
Hartwegi, *Benth. ex A. DC. in DC. Prod.* vii. 361.—Mexic.
hederacea, Cham. in Linnaea, viii. (1833) 212 = Pratia hederacea.
hederacea, *Hort. Kew. ex A. DC. in DC. Prod.* vii. 370.—Hab.?
heteromalla, Schrad. ex A. DC. l. c. 370 = triquetra.
heterophylla, Hook. Bot. Mag. t. 3784 = tenuior.
heterophylla, *Labill. Nov. Holl. Pl.* i. 52. t. 74.—Austral.
heterophylla, Willd. ex Spreng. Syst. i. 736 = Wahlenbergia linarioides.
Heyniana, Roem. & Schult. Syst. v. 50 = zeylanica.
Heyniana, Spreng. Syst. i. 717 = domingensis.
hirsuta, *Linn. Sp. Pl.* 932.—Afr. austr.
hirta, Cav. Ic. vi. 13 = Centropogon hirtus.
hirta, *Linn. Sp. Pl.* 932.—Afr. austr.
hirta, Roxb. Fl. Ind. ii. 113 = affinis.
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 577 = Pratia begonifolia.
hortensis, A. DC. in DC. Prod. vii. 377 = amoena.
Humboldtiana, Presl, Prod. Monog. Lobel. 35 = Siphocampylus Humboldtianus.
Humboldtiana, Roem. & Schult. Syst. v. 68 = Specularia perfoliata.
humilis, Klotzsch, in Peters, Reise Mossamb. Bot. 302 = fervens.
humistrata, *F. Muell. ex Benth. Fl. Austral.* iv. 130.—Austral.

LOBELIA :—
hypocrateriformis, R. Br. Prod. 565 = Isotoma Brownii.
hyssopifolia, *Salzm. ex A. DC. in DC. Prod.* vii. 389.—Chili.
ignea, Paxt. Mag. Bot. vi. (1839) 247 = fulgens.
ignea, *Vell. Fl. Flum.* viii. t. 158.—Bras.
ilicifolia, Ker-Gawl. in Bot. Mag. t. 1896 = purpurascens.
imbricata, Cham. in Linnaea, viii. (1833) 206 = Siphocampylus imbricatus.
incana, *Ruiz & Pav. ex A. DC. in DC. Prod.* vii. 392.—Peruv.
incisa, *D. Dietr. Syn. Pl.* i. 733.—Afr. austr.
incisa, Thunb. Prod. Pl. Cap. 39 = Cyphia incisa.
incisa, Wall. ex Hook. f. & Thoms. Journ. Linn. Soc. ii. (1857) 29 = Cliffortiana.
inconspicua, *A. Rich. Tent. Fl. Abyss.* ii. 8.—Abyss.
incurva, Rafin. New Fl. Am. ii. 18 = Kalmii?
inflata, *Linn. Sp. Pl.* 931.—Am. bor.
integrifolia, *Hort. Nois. ex Presl, Prod. Monog. Lobel.* 50, nomen.—Hab.?
inundata, R. Br. Prod. 563 = Isotoma fluviatilis.
inundata, A. Cunn. ex Walp. Rep. vi. 372 = Pratia erecta.
irasuensis, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 153.—Am. centr.
irrigua, R. Br. Prod. 563 = Pratia irrigua.
Jamesoniana, Presl, Prod. Monog. Lobel. 36 = Siphocampylus Jamesonianus.
japonica, F. G. Dietr. Vollst. Lexik. Gaertn. v. 552 = radicans.
javanica, *Thunb. Fl. Javan.* 9.—Java.
Kalmii, Bart. Fl. Am. Sept. t. 34 = Nuttallii.
Kalmii, *Linn. Sp. Pl.* 930.—Am. bor.
kamtschatica, Pall. ex Spreng. Syst. i. 712 = sessilifolia.
Kohautiana, *Vatke, in Linnaea*, xxxviii. (1874) 719.—Afr. trop.
Kraussii, *R. Grah. in Edinb. N. Phil. Journ.* (Jan.–March 1830) 379.—Ins. S. Doming.
Krebsiana, A. DC. in DC. Prod. vii. 385 = L. Erinus.
Kunthiana, Presl, Prod. Monog. Lobel. 39 = Centropogon Chamissonianus.
laciniata, Lam. Encyc. iii. 584 = Siphocampylus Lamarckii.
laciniata, Presl, Prod. Monog. Lobel. 40 = Siphocampylus laciniatus.
lacustris, Salisb. Prod. 128 = L. Dortmanna.
laevigata, Linn. f. Suppl. 392 = Centropogon laevigatus.
lanceolata, Hook. & Arn. Bot. Beech. Voy. 88 = laxiflora.
laricina, Spreng. f. Tent. Suppl. 9 = Prismatocarpus diffusus.
lasiantha, *A. DC. in. DC. Prod.* vii. 362.—Afr. austr.
Laurentia, Linn. Sp. Pl. 931 = Laurentia Michelii.
Laurentia, Willd. Sp. Pl. i. 948, partim = Laurentia tenella.
lavendulacea, Klotzsch, in Peters, Reise Mossamb. Bot. 302 = triquetra.
laxiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 311.—Mexic.
Lehmanni, De Vriese, in Lehm. Pl. Preiss. i. 394 = Isotoma Brownii.
leptocarpa, *Griessel. in Linnaea*, v. (1830) 419.—Afr. austr.
leptostachys, *A. DC. in DC. Prod.* vii. 376.—Am. bor.
limoselloides, Willd. ex Roem. & Schult. Syst. v. 41 = Lysipoma montioides.
linarioides, *A. DC. in DC. Prod.* vii. 371.—Afr. austr.
linearis, Thunb. ex Harv. & Sond. Fl. Cap. iii. 544 = L. Erinus.
linearis, *Thunb. Prod. Pl. Cap.* 39.—Afr. austr.
linifolia, Willd. ex Roem. & Schult. Syst. v. 57 = collina.
littoralis, *R. Cunn. ex A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 50.—N. Zel.
Lobbiana, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 28.—Reg. Himal.
longepedunculata, De Vriese, in Lehm. Pl. Preiss. i. 394 = tenuior.
longiflora, Linn. Sp. Pl. 930 = Isotoma longiflora.
longifolia, *A. DC. in DC. Prod.* vii. 382.—Ins. Philipp.
longipes, A. DC. l. c. 361 = hirsuta.
longiscapa, *De Vriese, in Lehm. Pl. Preiss.* i. 398.—Austral.
loxensis, *Willd. ex Roem. & Schult. Syst.* v. 72.—Peruv.
Lucaeana, *A. DC. in DC. Prod.* vii. 383.—Chili.
ludoviciana, *A. Gray, in Proc. Am. Acad.* xii. (1877) 60.—Am. bor.
lutea, *Linn. Sp. Pl.* 932.—Afr. austr.

LOBELIA:—
luzoniensis, Pers. ex Steud. Nom. ed. I. 490 = serpens.
lycioides, Cham. in Linnaea, viii. (1833) 207 = Siphocampylus lycioides.
macrocarpa, De Vriese, in Lehm. Pl. Preiss. i. 396 = Bergiana.
macrophylla, Presl, Prod. Monog. Lobel. 39 = Siphocampylus lycioides.
macropoda, Thunb. Pl. Bras. Dec. i. 6; Roem. & Schult. Syst. v. 52 = Siphocampylus macropodus.
macrostachys, *Hook. & Arn. Bot. Beech. Voy.* 88.—Ins. Sandvic.
macrostemon, Presl, Prod. Monog. Lobel. 36 = Siphocampylus macrostemon.
macularis, A. DC. in DC. Prod. vii. 364 = coronopifolia.
maculata, Penny, in Loud. Hort. Brit. Suppl. i. 592 = Pratia angulata.
madagascariensis, *Roem. & Schult. Syst.* v. 67.—Madag.
megapotamica, Spreng. Syst. iv. Cur. Post. 75 = Wahlenbergia linarioides.
Melleri, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 468.—Afr. trop.
membranacea, *R. Br. Prod.* 563.—Austral.
Michauxii, Nutt. Gen. Am. ii. 76 = Cliffortiana?
micrantha, H. B. & K. Nov. Gen. et Sp. iii. 316 = trigona.
microcarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 424.—Burma; Malaya.
microdon, *A. DC. in DC. Prod.* vii. 369.—Afr. austr.
microphylla, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
microsperma, *F. Muell. Fragm.* x. 41.—Austral.
Milleri, *Hort. ex Loud. Encyc. Pl. Suppl.* i. 1164.—Hab.?
Milleri, Steud. Nom. ed. II. ii. 63, in syn. = *Muelleri*.
minima, Sims, Bot. Mag. t. 2077 = Laurentia minuta.
minuta, DC. Fl. Fr. iii. 716 = Laurentia tenella.
minuta, Linn. Sp. Pl. 932 = Laurentia micrantha.
minutiflora, *Kunze, Pugill. Pl. Ined.* i. 10.—Mexic.
mishmica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 426.—Reg. Himal.
modesta, *Wedd. Chlor. And.* ii. 14.—N. Granat.
mollis, *R. Grah. in Edinb. N. Phil. Journ.* (Oct.-Dec. 1829) 185.—Ins. S. Doming.
monanthos, De Vriese, in Lehm. Pl. Preiss. i. 398 = Isotoma scapigera.
montana, Reinw. ex Blume, Bijdr. 728 = Pratia montana.
monticola, *H. B. & K. Nov. Gen. et Sp.* iii. 316.—N. Granat.
mucronata, *Cav. Ic.* vi. 11.—Chili.
mucronata, *Engelm. in Wisliz. Tour N. Mexic.* 108.—Mexic.
Muelleri, Hort. ex Steud. Nom. ed. II. ii. 61 = syphilitica.
multicaule, *Steud. l. c.*—Afr. austr.
multiflora, *Knowles & Westsc. Flor. Cab.* iii. (1840) 126.—Hab.?
Mundtiana, Cham. in Linnaea, viii. (1833) 215 = leptocarpa.
muscoides, *Cham. l. c.*—Afr. austr.
Mutisiana, H. B. & K. Nov. Gen. et Sp. iii. 303 = Siphocampylus Mutisianus.
nana, *H. B. & K. l. c.* 317. t. 272.—Mexic.
natalensis, *A. DC. in DC. Prod.* vii. 369.—Afr. austr.
neglecta, *Roem. & Schult. Syst.* v. 36.—Hab.?
neglecta, *Vatke, in Linnaea*, xxxviii. (1874) 720.—Mexic.
nematophylla, Spreng. ex Eckl. & Zeyh. Enum. 398 = setacea.
neriifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 150.—Ins. Sandvic.
neriifolia, Moris, in Mem. Acc. Torin. xxxviii. (1835) 44 = salicifolia.
nicotianaefolia, *Heyne, in Roth, Nov. Pl. Sp.* 143.—Ind. or.
nitida, Presl, Prod. Monog. Lobel. 37 = Siphocampylus nitidus.
nivea, Rafin. Ann. Nat. (1830) 16 = spicata.
nivea, Willd. ex Roem. & Schult. Syst. v. 58 = Siphocampylus Humboldtianus.
nuda, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 469.—Afr. trop.
nudicaulis, Lam. Encyc. iii. 590 = Cyphia Phyteuma.
nudicaulis, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
Nummularia, Lam. Encyc. iii. 589 = Pratia begonifolia.
nummularioides, *Cham. in Linnaea*, viii. (1833) 209.—Bras.

LOBELIA:—
Nuttallii, *Roem. & Schult. Syst.* v. 39.—Am. bor.
obliqua, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 158 = Pratia bigonifolia.
oblongifolia, Gaudich. in Freyc. Voy. Bot. 459 = Clermontia oblongifolia.
obscura, *Linn. Pl. Surin.* n. 112; *Amoen. Acad.* viii. 262.—Guian.
obtusifolia, Willd. ex Roem. & Schult. Syst. v. 57 = Siphocampylus scandens.
ocimoides, *Kunze, Del. Sem. Hort. Lips.* (1847).—*Cf. Linnaea*, xxiv. (1851) 178.—Mexic.
odorata, R. Grah. in Edinb. N. Phil. Journ. xvi. (1834) 178 = Pratia hederacea.
ophiocephala, De Vriese, in Lehm. Pl. Preiss. i. 397 = Isotoma scapigera.
oppositifolia, Spreng. ex A. DC. in DC. Prod. vii. 390 (Wahlenbergiae sp.).—Afr. austr.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 128 = Haynaldia organensis.
Orizabae, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1842) reimpr. 13.—Mexic.
Ottoniana, *A. DC. in DC. Prod.* vii. 370.—Hab.?
ovalifolia, Hook. & Arn. Bot. Beech. Voy. 301 = laxiflora.
pallida, Muhl. Cat. 22 = spicata.
palmaris, Willd. ex Roem. & Schult. Syst. iv. 56 = monticola.
paludosa, *Nutt. Gen. Am.* ii. 75.—Am. bor.
paniculata, *Linn. Sp. Pl.* 930.—Afr. austr.
paniculata, Rafin. New Fl. Am. ii. 18 = spicata?
parviflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1842) reimpr. 10.—Mexic.
parvifolia, Berg. Desc. Pl. Cap. 345 = Lightfootia oxycoccoides.
parvifolia, *R. Br. Prod.* 564.—Austral.
parvifolia, Rafin. New Fl. Am. ii. 18 (Quid?).—Am. bor.
patula, *Linn. f. Suppl.* 395.—Afr. austr.
paucidentata, A. DC. in DC. Prod. vii. 362 = tomentosa.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* iii. 314.—Mexic.
pectinata, Engelm. in Wisliz. Tour N. Mexic. 108 = fenestralis.
pedicellaris, Presl, Prod. Monog. Lobel. 34 = Siphocampylus longepedunculatus.
pedunculata, R. Br. Prod. 563 = Pratia pedunculata.
pedunculata, Sims, Bot. Mag. t. 2251 = tomentosa.
pendulifolia, Presl, Prod. Monog. Lobel. 35 = Siphocampylus longepedunculatus.
perpusilla, Hook. f. Fl. N. Zel. i. 158 = Pratia perpusilla.
persicaefolia, Cav. Ic. vi. 12. t. 518 = laxiflora.
persicaefolia, *Lam. Encyc.* iii. 585.—Ins. Guadalup.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. 229 = fervens.
phyllostachya, *Engelm. in Wisliz. Tour N. Mexic.* 108.—Mexic.
physaloides, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 50 (= *Colensoa physaloides*).—N. Zel.
Phyteuma, Linn. Sp. Pl. 930 = Cyphia Phyteuma.
phyteumoides, Willd. ex Roem. & Schult. Syst. v. 68 = collina.
pinifolia, *Linn. Sp. Pl.* 929.—Afr. austr.
pinnata, Lam. Encyc. iii. 591 = Cyphia pinnata.
pinnatifida, Cham. in Linnaea, viii. (1833) 220 = Delissea pinnatifida.
platycalyx, F. Muell. Fragm. iv. 183 = Pratia platycalyx.
Plumieri, Burm. f. Fl. Ind. 186 = Scaevola Koenigii.
Plumieri, Linn. Sp. Pl. 929 = Scaevola Lobelia.
Poeppigiana, Kunze, ex A. DC. in DC. Prod. vii. 393 = polyphylla.
polygalaefolia, Willd. ex Roem. & Schult. Syst. v. 56 = rupestris.
polymorpha, Willd. l. c. 68 = serpens.
polyphylla, *Hook. & Arn. Bot. Beech. Voy.* 33.—Chili.
porphyrantha, *Decne. in Rev. Hortic.* Sér. IV. ix. (1860) 118.—Mexic.
pratiana, Gaudich. in Freyc. Voy. Bot. 456 = Hypsela reniformis.
pratioides, *Benth. Fl. Austral.* iv. 131.—Austral.
Preslii, *A. DC. in DC. Prod.* vii. 358.—Afr. austr.
Princeps, Otto & Dietr. Allg. Gartenz. vii. (1839) 298 = fulgens.
procumbens, *Hort. ex A. DC. in DC. Prod.* vii. 387.—Afr. austr.

LOBELIA:—
propinqua, Hort. ex Loud. Hort. Brit. Suppl. ii. 646 = fulgens.
prunifolia, Presl, Prod. Monog. Lobel. 37 = Siphocampylus prunifolius.
psilophylla, Presl, l. c. 36 = Siphocampylus psilophyllus.
pterocaulon, Klotzsch, in Peters, Reise Mossamb. Bot. 299, 302 = fervens.
pteropoda, *A. DC. in DC. Prod.* vii. 364.—Afr. austr.
puberula, *Michx. Fl. Bor. Am.* ii. 152.—Am. bor.
pubescens, Dietr. ex A. DC. in DC. Prod. vii. 355 = leptocarpa.
pubescens, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 498.—Afr. austr.
pubescens, Sieber, ex A. DC. in DC. Prod. vii. 364 = gracilis.
pubescens, Willd. Enum. Hort. Berol. 218 = Laurentia repens.
pulchella, *Vatke, in Linnaea*, xxxviii. (1874) 720.—Mexic.
pulverulenta, Hook. Journ. Bot. i. (1834) 231 = Siphocampylus Jamesonianus.
pulverulenta, Pers. Syn. ii. 212 = Siphocampylus campanulatus.
pumila, Burm. f. Fl. Ind. 186. t. 60. f. 3.—Ind. or.
pumila, Salisb. Prod. 129 = Laurentia minuta.
punicea, Otto & Dietr. Allg. Gartenz. vii. (1839) 299 = fulgens.
purpurascens, *R. Br. Prod.* 563.—Austral.
purpurascens, Wall. Cat. n. 1307 = colorata.
purpurea, Breit. Hort. Breit. (1817) 249; et ex Presl, Prod. Monog. Lobel. 50 (Quid?).—Hab.?
purpurea, *Lindl. Bot. Reg.* t. 1325.—Chili.
purpurea, Ruiz & Pav. ex A. DC. in DC. Prod. vii. 390, 399 = Siphocampylus rosmarinifolius.
purpurea, *Vell. Fl. Flum.* viii. t. 156.—Bras.
pusilla, *A. DC. in DC. Prod.* vii. 379.—Mexic.
pusilla, G. Don, Gen. Syst. iii. 710 = Laurentia micrantha.
pusilla, Poepp. ex A. DC. Prod. vii. 347 (= *Clintonia pusilla*).—Chili.
pygmaea, Thunb. Prod. Pl. Cap. 40 = Laurentia micrantha.
pyramidalis, Hohen. ex Hook. f. in Journ. Linn. Soc. ii. (1857) 29 = rosea.
pyramidalis, *Wall. in As. Res.* xiii. (1820) 376.—Reg. Himal.; China.
quadrangularis, *R. Br. Prod.* 562.—Austral.
racemosa, *Sims, Bot. Mag.* t. 2137.—Ind. occ.
racemosa, *Vest, ex Roem. & Schult. Syst.* v. 58.—Hab.?
radicans, *Thunb. in Trans. Linn. Soc.* ii. (1794) 330.—Reg. Himal.; China.
ramosa, Benth. in Maund, Botanist, ii. t. 93 = tenuior.
ramosa, Hort. ex Burb. in Gard. Chron. (1881) i. 105 = fulgens.
ramosissima, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) reimpr. 11 = Laurentia ramosissima.
rapunculoides, *H. B. & K. Nov. Gen. et Sp.* iii. 512.—Mexic.
recurvifolia, Spreng. Syst. v. 420 (Index) = Simsii.
reflexa, Stokes, Bot. Mat. Med. i. 342 = syphilitica.
Reinwardtiana, A. DC. in DC. Prod. vii. 367 = trigona.
reniformis, *Cham. in Linnaea*, viii. (1833) 210.—Bras.
repanda, T. Martyn, in Mill. Gard. Dict. ed. IX. n. 33 = Pratia angulata.
repens, Thunb. Prod. Pl. Cap. 40 = anceps.
reticulata, *Willd. in Roem. & Schult. Syst.* v. 58.—Am. austr.
retrorsa, *Willd. l. c.* 57.—Am. austr.
rhizophyta, Spreng. Nov. Prov. 25 = anceps.
rhombifolia, *De Vriese, in Lehm. Pl. Preiss.* i. 397.—Austral.
rhynchopetalum, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 465.—Afr. trop.
rhytidosperma, *Benth. Fl. Austral.* iv. 126.—Austral.
rigidula, H. B. & K. Nov. Gen. et Sp. iii. 511 = laxiflora.
robusta, *R. Grah. in Edinb. N. Phil. Journ.* (July-Sept. 1831) 378.—Ins. S. Doming.
robusta, Wall. ex Voigt, Hort. Suburb. Calc. 367 = eurostos.
Rollandia, Gaudich. in Freyc. Voy. Bot. 458 = Rollandia lanceolata.
rosea, *Wall. Pl. As. Rar.* ii. 42. t. 152.—Reg. Himal.; Burma.

LOBELIA:—
rosmarinifolia, Domb. ex Presl, Prod. Monog. Lobel. 36 = Siphocampylus rosmarinifolius.
rotundifolia, *Juss. ex A. DC. in DC. Prod.* vii. 383.—Ins. S. Doming.
Roughii, *Hook. f. Handb. N. Zeal. Fl.* 171.—N. Zel.
ruderalis, Willd. ex Roem. & Schult. Syst. iv. 56 = trigona.
rugosa, *Presl, Prod. Monog. Lobel.* 34.—Peruv.
rugulosa, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1829) 186 = Pratia angulata.
rupestris, *H. B. & K. Nov. Gen. et Sp.* iii. 313.—N. Granat.
rupincola, Bert. ex Hook. & Arn. in Hook. Journ. Bot. i. (1834) 277 = anceps.
salicifolia, Fisch. ex Herd. in Act. Hort. Petrop. i. (1873) 285 = sessilifolia.
salicifolia, *Sweet, Hort. Suburb. Lond.* 37.—Chili.
salicina, Lam. Encyc. iii. 583 = acuminata.
saligna, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 65 = sessilifolia.
salviaefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 69.—Cuba.
Salzmanniana, Presl, Symb. Bot. 31. t. 20 = Laurentia Michelii.
Sartorii, *Vatke, in Linnaea*, xxxviii. (1874) 721.—Mexic.
saxicola, De Vriese, in Lehm. Pl. Preiss. i. 398 = anceps.
scabra, Nees, ex Steud. Nom. ed. II. ii. 62 = tomentosa.
scabra, Presl, Prod. Monog. Lobel. 34 = Siphocampylus longepedunculatus.
scabra, Spreng. Neue Entdeck. i. 272 = aspera.
scabra, *Thunb. in Hoffm. Phytogr. Blaett.* i. (1803) 21.—Afr. austr.
scabripes, *A. DC. in DC. Prod.* vii. 369.—Afr. austr.
scaevolaefolia, *Roxb. ex A. DC. l. c.* 357.—Ins. S. Helen.
scandens, Bert. ex A. DC. l. c. 345 = Centropogon Berterianus.
scandens, H. B. & K. Nov. Gen. et Sp. iii. 309 = Siphocampylus scandens.
scapigera, R. Br. Prod. 565 = Isotoma scapigera.
scapigera, Willd. Enum. Hort. Berol. Suppl. 11 = tenella.
Schiedeana, Heynh. Nom. ii. 367 = fulgens.
Schimperi, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 6.—Abyss.
Schrankii, Sweet, Hort. Brit. ed. II. 323 = anceps.
scoparia, *Buching. ex Krauss, in Flora*, xxvii. (1844) 820.—Afr. austr.
Sebae, A. DC. in DC. Prod. vii. 386 = zeylanica.
secunda, Linn. f. Suppl. 395 = Laurentia repens.
secunda, *Ruiz & Pav. ex A. DC. in DC. Prod.* vii. 390, 392.—Peruv.
secunda, Schrank, Pl. Rar. Hort. Monac. t. 31 = anceps.
Sellowiana, Presl, Prod. Monog. Lobel. 36 = Siphocampylus Westinianus.
senecioides, Hort. ex Gaudich. in Freyc. Voy. Bot. 456; A. Cunn. in Bot. Mag. t. 2702 = Isotoma axillaris.
Senecionis, Spreng. Syst. iv. Cur. Post. 75 = Isotoma axillaris.
senegalensis, *A. DC. in DC. Prod.* vii. 372.—Afr. trop.
serpens, Bout. ex Boj. Hort. Maurit. 193 = filiformis.
serpens, *Lam. Encyc.* iii. 588.—Ins. Mascar.
serrata, *Meyen, Reise*, i. 300.—Chili.
serrulata, Schott, in Roem. & Schult. Syst. v. 54 = urens.
sessilifolia, *Lamb. in Trans. Linn. Soc.* x. (1811) 260.—Kamtchat.
sessilifolia, Willd. ex Roem. & Schult. Syst. v. 68 = Specularia perfoliata.
setacea, Sibth. & Sm. Fl. Graec. Prod. i. 145 = Laurentia tenella.
setacea, *Thunb. in Hoffm. Phyt. Blaett.* i. (1803) 21.—Afr. austr.
setacea, Zeyh. ex Presl, Prod. Monog. Lobel. 52 (Prismatocarpi sp.).—Afr. austr.
simplex, Linn. Mant. ii. 291; Thunb. Prod. Pl. Cap. 39 = debilis.
simplex, Willd. ex A. DC. in DC. Prod. vii. 375 = pauciflora.
simplicicaulis, Benth. in Enum. Pl. Hueg. 74 = rhytidosperma.
simplicicaulis, R. Br. Prod. 564 = gibbosa.
Simsii, *Sweet, Hort. Brit.* ed. I. 247.—Afr. austr.
siphilitica, Linn. Sp. Pl. 931, vide syphilitica.

LOBELIA:—
sonchifolia, *Sw. Fl. Ind. Occ.* iii. 1947.—Ins. S. Doming.
spartioides, *D. Dietr. Syn. Pl.* i. 728.—Afr. austr.
spathulata, Bridges, ex Hook. & Arn. in Hook. Journ. Bot. i. (1834) 277 = Selliera radicans.
speciosa, Sweet, Brit. Flow. Gard. Ser. II. t. 174 = syphilitica.
spectabilis, H. B. & K. Nov. Gen. et Sp. iii. 306 = Centropogon surinamensis.
Speculum, Andr. Bot. Rep. t. 644 = debilis.
sphaerocarpa, Juss. ex A. DC. in DC. Prod. vii. 345, 391 = Centropogon surinamensis.
spicata, *Lam. Encyc.* iii. 587.—Am. bor.
spicata, Moç. ex A. DC. in DC. Prod. vii. 361 = fenestralis.
spicata, *Ruiz & Pav. ex G. Don, Gen. Syst.* iii. 705.—Peruv.
splendens, Willd. Hort. Berol. t. 86 = fulgens.
stellarioides, *Benth. & Hook. f. ex Hemsl. in Oliver, Fl. Trop. Afr.* iii. 470.—Afr. trop.
stenophylla, *Benth. Fl. Austral.* iv. 130.—Austral.
stenotheca, F. Muell. Fragm. ii. 20; iv. 171 = Bergiana.
stimulans, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 157 = pyramidalis.
stipularis, Roth, Nov. Pl. Sp. 144 = trigona.
stricta, R. Br. Prod. 564 = gibbosa.
stricta, De Vriese, in Lehm. Pl. Preiss. i. 396 = anceps.
stricta, Mart. & Gal. in Bull. Acad. Brux. ix. (1862) reimpr. 12 = fenestralis.
stricta, *Sw. Prod. Veg. Ind. Occ.* 117.—Ins. Guadalup.
subcordata, Gaudich. in Freyc. Voy. Bot. 457; D. Dietr. Syn. Pl. i. 728 = Delissea subcordata.
subcuneata, Miq. Fl. Ind. Bat. ii. 574 = affinis.
subincisa, Wall. Cat. n. 1310 = trigona.
subincisa, Wall. l. c. et p. 158 = Raphistemma pulchellum.
submersa, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 50 = Glossostigma elatinoides.
subnuda, *Benth. Pl. Hartw.* 44.—Mexic.
subpubera, *Wedd. Chlor. And.* ii. 14.—Ecuador.
subracemosa, Miq. Fl. Ind. Bat. ii. 576 = trigona.
subtilis, H. B. & K. Nov. Gen. et Sp. iii. 316 = trigona.
subulata, *Benth. Pl. Hartw.* 137.—Ecuador.
subulata, Klotzsch, in Peters, Reise Mossamb. Bot. 299, 302 = fervens.
succulenta, *Blume, Bijdr.* 728.—Java.
superba, Cham. in Linnaea, viii. (1833) 223 = Cyanea superba.
surinamensis, Linn. Sp. Pl. ed. II. 1320 = Centropogon surinamensis.
surrepens, *Hook. f. Fl. Tasm.* i. 237. t. 69 A.—Tasman.
syphilitica, *Linn. Sp. Pl.* 931 (siphilitica).—Afr. austr.
Taccada, Gaertn. Fruct. i. 119. t. 25 = Scaevola Koenigii.
telephioides, A. DC. in DC. Prod. vii. 367 = serpens.
tenella, Bivona, Pl. Sic. Cent. i. 53 = Laurentia tenella.
tenella, *Linn. Mant.* i. 120.—Afr. austr.
tenella, Zeyh. ex A. DC. in DC. Prod. vii. 351 = debilis.
tenera, *H. B. & K. Nov. Gen. et Sp.* iii. 314.—Ecuador.
tenuifolia, D. Dietr. Syn. Pl. i. 731 = linearis.
tenuifolia, Willd. ex Roem. & Schult. Syst. v. 56 = fastigiata.
tenuior, *R. Br. Prod.* 564.—Austral.
terminalis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 424.—Bengal.
tetragona, *Blume, Bijdr.* 729.—Java.
texensis, Rafin. Atl. Journ. 176 = fulgens.
thapsoidea, Schott, in Pohl, Pl. Bras. Ic. ii. 102. t. 167 = Haynaldia thapsoidea.
thapsoides, Cham. in Linnaea, viii. (1833) 209 = Haynaldia organensis.
thermalis, Thunb. Prod. Pl. Cap. 40 = leptocarpa.
Thunbergii, Sweet, Hort. Brit. ed. I. 247 = coronopifolia.
tomentosa, *Linn. f. Suppl.* 394.—Afr. austr.
Trapa, Linn. Sp. Pl. 929 = L. Tupa.
trialata, Buch.-Ham. in D. Don, Prod. Fl. Nep. 157 = trigona.

LOBELIA:—
triangulata, Buch.-Ham. ex Wall. Cat. n. 1312 = Cliffortiana.
triangulata, Roxb. Hort. Beng. 16 = trigona.
trichandra, Wight, Ic. t. 1171 = nicotianaefolia.
trigona, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 506.—Ind. or.
trigona, Thw. Enum. Pl. Zeyl. 169 = zeylanica.
trigonocaulis, Hook. Bot. Mag. t. 5088 = gracilis.
trigonocaulis, *F. Muell. Fragm.* i. 18, 242.—Austral.
trinitensis, *Griseb. Fl. Brit. W. Ind.* 385.—Ind. occ.
triphylla, *Presl, Symb. Bot.* i. 63. t. 40 = Siphocampylus duploserratus.
triquetra, *Linn. Mant.* i. 120.—Afr. austr.
triquetra, Reinw. ex A. DC. in DC. Prod. vii. 367 = Reinwardtiana.
triquetra, Sieber, ex A. DC. l. c. 372 = senegalensis.
trullifolia, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 466.—Afr. trop.
tuberosa, *Burm. f. Fl. Cap. Prod.* 29 (err. typ. 25).—Afr. austr.
Tupa, Donn, ex Steud. Nom. ed. I. 491 = Siphocampylus giganteus.
Tupa, *Linn. Sp. Pl.* ed. II. 1318.—Chili; Peruv.
tyrianthina, *Forbes, Hort. Woburn.* 33.—Peruv.
umbellata, Cham. in Linnaea, viii. (1833) 321 = Centropogon Chamissonianus.
umbellata, H. B. & K. Nov. Gen. et Sp. iii. 304 = Siphocampylus umbellatus.
umbellata, *Vest, ex Roem. & Schult. Syst.* v. 58.—Hab.?
umbrosa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 7 = trigona.
uncinata, De Vriese, in Lehm. Pl. Preiss. i. 396 = anceps.
undulata, Gaudich. in Freyc. Voy. Bot. 437; D. Dietr. Syn. Pl. i. 728 = Delissea undulata.
unidentata, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 356.—Afr. austr.
unidentata, *Hort. Berol. ex Spreng. Nov. Prov.* 26.—Hab.?
uranocoma, Cham. in Linnaea, viii. (1833) 321 = Haynaldia uranocoma.
urens, *Linn. Sp. Pl.* 931.—Europ.
vagans, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 16.—Ins. Mascar.
valdiviana, *Vatke, in Linnaea*, xxxviii. (1874) 725.—Chili.
variifolia, *Sims, Bot. Mag.* t. 1692.—Afr. austr.
velutina, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) reimpr. 10 = Hartwegi.
verbascifolia, Presl, Prod. Monog. Lobel. 38 = Siphocampylus verbascifolius.
verbenaefolia, Salisb. Prod. 129 = urens.
veronicaefolia, *Willd. ex Roem. & Schult. Syst.* v. 56.—Am. austr.
verticillata, Cham. in Linnaea, viii. (1833) 202 = Siphocampylus verticillatus.
verticillata, Kunze, ex Presl, Prod. Monog. Lobel. 52 (Stemodiae sp.).—Chili.
Vestiana, Presl, l. c. 50 = racemosa, *Vest.*
villosula, Presl, l. c. 38 = Siphocampylus villosulus.
virescens, Benth. Pl. Hartw. 77 = Burmeistera virescens.
virgata, Poepp. ex A. DC. in DC. Prod. vii. 391, 398 = Siphocampylus virgatus.
volubilis, Burm. f. Fl. Cap. Prod. 29 (err. typ. 25) = Cyphia volubilis.
volubilis, H. B. & K. Nov. Gen. et Sp. iii. 308 = Siphocampylus volubilis.
Wallichiana, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1857) 29 = pyramidalis.
Wallichii, Steud. Nom. ed. II. ii. 63 = colorata.
Warscewiczii, *Vatke, in Linnaea*, xxxviii. (1874) 718.—Costa Rica.
Westiana, Steud. Nom. ed. I. 491 = seq.
Westiniana, Thunb. Pl. Bras. Dec. i. 5. t. 1. f. 1 = Siphocampylus Westinianus.
Willdenowiana, Presl, Prod. Monog. Lobel. 39 = Siphocampylus umbellatus.
Willdenowiana, Roem. & Schult. Syst. v. 41 = Lysipoma aretioides.
xalapensis, H. B. & K. Nov. Gen. et Sp. iii. 315 = Cliffortiana.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* iii. 539.—Afr. austr.
zeylanica, *Linn. Sp. Pl.* 932.—Ind. or.
zeylanica, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 113 = affinis.

LOBIROTA, Dulac, Fl. Hautes-Pyr. 430 (1867) = **Ramondia**, Rich. (Gesner.).
pyrenaica, Dulac, l. c. = R. pyrenaica.

LOBOCARPUS, Wight & Arn. Prod. 7 (1834) = **Glochidion**, Forst. (Euphorb.).
Candolleanus, Wight & Arn. l. c. = G. neilgherrense.

LOBOMON, Rafin. New Fl. Am. i. 81 (1836) = **Amphicarpaea**, Ell. (Legum.).
acutifolium, montanum, obtusifolium, sarmentosum, Rafin. l. c. 84 = A. monoica.

LOBOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 21 = **Coldenia**, Linn. (Borag.).
tetrandrum, F. Muell. l. c. = C. procumbens.

LOBOPOGON, Schlecht. in Linnaea, xx. (1847) 620 = **Brachyloma**, Sond. (Epacrid.).
ericoides, Schlecht. l. c. = B. ericoides.

LOBOPTERA, Colla, in Mem. Acc. Torin. Ser. II. x. (1849) 221 = **Columnea**, Plum. (Gesner.).
subsessilis, Colla, l. c. = C. Lindleyana.

LOBOSTEMA, Spreng. Gen. i. 126 (1830) = seq.

LOBOSTEMON, Lehm. in Linnaea, v. (1830) 378. t. 5. *BORAGINEAE*, Benth. & Hook. f. ii. 863.
ECHIOPSIS, Reichb. Handb. 192 (1837).
ISORIUM, Rafin. Fl. Tellur. ii. 61 (1836).
OPLEXION, PENTHYSA, TRAXARA, Rafin. l. c. iv. 85, 86 (1836).
acutissimus, *Buek, in Linnaea*, xi. (1837) 139.—Afr. austr.
argenteus, *Buek, l. c.* 133.—Afr. austr.
breviflorus, *DC. Prod.* x. 10.—Afr. austr.
capitatus, *Buek, in Linnaea*, xi. (1837) 143.—Afr. austr.
capitiformis, *DC. Prod.* x. 12.—Afr. austr.
caudatus, *Buek, in Linnaea*, xi. (1837) 147.—Afr. austr.
cephaloides, *DC. Prod.* x. 12.—Afr. austr.
cinereus, *DC. l. c.* 10.—Afr. austr.
curvifolius, *Buek, in Linnaea*, xi. (1837) 137.—Afr. austr.
diversifolius, *Buek, l. c.* 140.—Afr. austr.
Dregei, *DC. Prod.* x. 6.—Afr. austr.
echioides, Lehm. in Linnaea, v. (1830) 378 = Lehmannianus.
Ecklonianus, *Buek, l. c.* xi. (1837) 144.—Afr. austr.
elongatus, *Buek, l. c.* 140.—Afr. austr.
eriostachyus, *Buek, l. c.* 148.—Afr. austr.
fastigiatus, *Buek, l. c.* 141.—Afr. austr.
ferocissimus, *DC. Prod.* x. 7.—Afr. austr.
formosus, *Buek, in Linnaea*, xi. (1837) 132.—Afr. austr.
fruticosus, *Buek, l. c.* 134.—Afr. austr.
glaber, *Buek, l. c.* 137.—Afr. austr.
glaucophyllus, *Buek, l. c.* 138.—Afr. austr.
hispidus, *DC. Prod.* x. 10.—Afr. austr.
laevigatus, *Buek, in Linnaea*, xi. (1837) 129.—Afr. austr.
lasiophyllus, *DC. Prod.* x. 10.—Afr. austr.
latifolius, *Buek, in Linnaea*, xi. (1837) 147.—Afr. austr.
Lehmannianus, *Buek, l. c.* 141.—Afr. austr.
lucidus, *Buek, l. c.* 136.—Afr. austr.
microphyllus, *Buek, l. c.* 142.—Afr. austr.
montanus, *Buek, l. c.* 132.—Afr. austr.
obovatus, *DC. Prod.* x. 10.—Afr. austr.
obtusifolius, *DC. l. c.* 7.—Afr. austr.
oederiaefolius, *DC. l. c.*—Afr. austr.
paniculatus, *Buek, in Linnaea*, xi. (1837) 139.—Afr. austr.
paniculiformis, *DC. Prod.* x. 8.—Afr. austr.
rosmarinifolius, *DC. l. c.* 10.—Afr. austr.
scaber, *DC. l. c.* 6.—Afr. austr.
somalensis, *Franch. in Revoil, Çomali*, 44.—Afr. trop.
sphaerocephalus, *Buek, in Linnaea*, xi. (1837) 143.—Afr. austr.
spicatus, *Buek, l. c.* 145.—Afr. austr.
splendens, *Buek, l. c.* 146.—Afr. austr.
Sprengelianus, *Buek, l. c.* 133.—Afr. austr.
stachydeus, *DC. Prod.* x. 7.—Afr. austr.
strigosus, *Buek, in Linnaea*, xi. (1837) 136.—Afr. austr.
Swartzii, *Buek, l. c.* 137.—Afr. austr.
Thunbergianus, *Buek, l. c.* 135.—Afr. austr.

LOBOSTEMON:—
thymeloideus, *DC. Prod.* x. 11.—Afr. austr.
trichotomus, *DC. l. c.*—Afr. austr.
trigonus, *Buek, in Linnaea*, xi. (1837) 135.—Afr. austr.
verrucosus, *Buek, l. c.* 138.—Afr. austr.
virgatus, *Buek, l. c.* 142.—Afr. austr.
viridi-argenteus, *Buek, l. c.* 144.—Afr. austr.
Wurmbii, *DC. Prod.* x. 11.—Afr. austr.
Zeyheri, *Buek, in Linnaea*, xi. (1837) 134.—Afr. austr.

LOBULARIA, Desv. Journ. Bot. iii. (1814) 172 = **Alyssum**, Tourn. (Crucif.).
canescens, Steud. Nom. ed. II. ii. 64 = A. Fischerianum.
elongata, Steud. l. c. = A. Fischerianum.
halimifolia, Steud. l. c. = A. halimifolium.
intermedia, Webb & Berth. Phyt. Canar. i. 92.—Ins. Canar.
Lagascae, Steud. Nom. ed. II. ii. 64 = A. Lagascae.
libyca, Meisn. Gen. Comm. 11 (libyka) = A. libycum.
longicaulis, Steud. Nom. ed. II. ii. 64 = A. longicaule.
lybica, Webb & Berth. Phyt. Canar. i. 90 = A. libycum.
maritima, Desv. Journ. Bot. iii. (1814) 169 = A. maritimum.
nummularia, Steud. Nom. ed. II. ii. 64 = A. libycum.
pyrenaica, Steud. l. c. = A. pyrenaicum.
rupestris, Steud. l. c. = A. rupestre.
spinosa, Steud. l. c. = A. spinosa.
strigulosa, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 837 = A. strigulosum.
tenuifolia, Steud. Nom. ed. II. ii. 64 = A. tenuifolium.

LOCANDI, Adans. Fam. ii. 449 (1763) = **Samadera**, Gaertn. (Simarub.).

LOCARDI, Steud. Nom. ed. II. ii. 64 (1841), sphalm. = praec.

LOCELLARIA, Welw. Apont. 588 (1858) = **Bauhinia**, Linn. (Legum.).
bauhinioides, Welw. l. c. = B. reticulata.

LOCHEMIA, Arn. in Ann. Sc. Nat. Sér. II. xi. (1839) 172 = **Melochia**, Linn. (Stercul.).
corchorifolia, Arn. l. c. = M. corchorifolia.

LOCHERIA, Neck. Elem. i. 41 (1790) = **Verbesina**, Linn. (Compos.).

LOCHERIA, Regel, in Flora, xxxi. (1848) 251 = **Achimenes**, P. Br. (Gesner.).
heterophylla, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 95 = A. heterophylla.
hirsuta, Regel, in Flora, xxxi. (1848) 252 = A. hirsuta.
ignescens, Klotzsch, ex Hanst. in Linnaea, xxvii. (1854) 713: Regel, Cat. Hort. Turic. (1854) = A. heterophylla.
magnifica, Planch. & Linden, Fl. des Serres, x. (1854–55) t. 1013 = A. magnifica.
multiflora, Regel, in Flora, xxxi. (1848) 252 = A. multiflora.
pedunculata, Regel, l. c. = A. pedunculata.

LOCHIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 409. *PARONYCHIAE*.
bracteata, *Balf. f. l. c.*—Ins. Socotra.

LOCHMOCYDIA, Mart. ex DC. Prod. ix. 177 (1845) = **Cuspidaria**, DC. (Bignon.).
lateriflora, Mart. ex DC. l. c. 179 = C. lateriflora.
multiflora, Mart. ex DC. l. c. = C. multiflora.
populeaster, Mart. ex DC. l. c. 178 = C. pterocarpa.
ramentacea, Mart. ex DC. l. c. = C. ramentacea.

LOCHNERA, Reichb. Consp. 134 (1828) = **Vinca**, Linn. (Apocyn.).
rosea, Reichb. l. c. = V. rosea.

LOCHNERIA, Scop. Introd. 271 (1777) = **Elaeocarpus**, Linn. (Tiliac.).

LOCKHARTIA, Hook. Bot. Mag. t. 2715 (1827). *ORCHIDEAE*, Benth. & Hook. f. iii. 570.
FERNANDEZIA, Lindl. Gen. et Sp. Orch. 207 (1833).
acuta, *Reichb. f. in Bot. Zeit.* x. (1852) 767.—Ins. Trinit.; Venezuela.
amoena, *Endres & Reichb. f. in Gard. Chron.* (1872) 666.—Costa Rica.
elegans, *Hook. Bot. Mag.* t. 2715.—Am. trop.

LOCKHARTIA:—
floribunda, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 179.—Hab.?
goyazensis, *Reichb. f. in Bot. Zeit.* x. (1852) 768.—Bras.
lamellosa, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 300.—Mexic.
ludibunda, *Reichb. f. in Bot. Zeit.* xv. (1857) 159.—Bras.
lunifera, *Reichb. f. l. c.* x. (1852) 767.—Mexic.
micrantha, *Reichb. f. l. c.* 768.—Am. centr.
mirabilis, *Reichb. f. Xenia Orch.* i. 106.—Panama.
obtusa, Regel, Gartenfl. vi. (1857) t. 193 (err. 190) = seq.
obtusifolia, *Regel, in Ann. Sc. Nat.* Sér. II. vi. (1856) 378; *et Gartenfl.* vi. (1857) 131.—N. Granat.
Oerstedii, *Reichb. f. in Bot. Zeit.* x. (1852) 767; *et Xenia Orch.* i. 105. t. 40.—Costa Rica.
pallida, *Reichb. f. in Bonplandia*, ii. (1854) 14; *Xenia Orch.* 102. t. 39.—Panama.
parthenocoma, *Reichb. f. in Bot. Zeit.* x. (1852) 767.—Venezuela.
parthenoglossa, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 300.—Hab.?
platyglossa, *Reichb. f. in Linnaea*, xli. (1877) 106.—N. Granat.
Serra, *Reichb. f. in Otia Bot. Hamb.* 6.—Ecuador.
verrucosa, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 53.—Guatemala.
Weigelti, *Reichb. & Reichb. f. in Bot. Zeit.* x. (1852) 767.—Guiana.

LOCUSTA, Riv. ex Medic. Phil. Bot. i. 55 (1789) = **Valerianella**, Moench (Valer.).
communis, Delarb. Fl. Auv. ed. II. 88 = V. olitoria.
coronata, Delarb. l. c. 89 = V. coronata.
echinata, Delarb. l. c. = V. echinata.
mixta, Delarb. l. c. = V. dentata.

LODDIGESIA, Sims, Bot. Mag. t. 965 (1808). *LEGUMINOSAE*, Benth. & Hook. f. i. 485.
oxalidifolia, *Sims, l. c.*—Afr. austr.

LODDIGGESIA, Reichb. Consp. 153 (1828) = praec.

LODHRA, Guill. in Ann. Sc. Nat. Sér. II. xv. (1841) 161 = **Symplocos**, Jacq. (Styrac.).
acuta, Miers, in Journ. Linn. Soc. xvii. (1879) 301 = S. acuta.
apicalis, Miers, l. c. 302 = S. apicalis.
attenuata, Miers, l. c. 299 = S. spicata.
bractealis, Miers, l. c. 301 = S. bractealis.
caudata, Miers, l. c. 300 = S. caudata.
cordifolia, Miers, l. c. 301 = S. cordifolia.
coronata, Miers, l. c. 302 = S. coronata.
crassifolia, Miers, l. c. = S. japonica.
crataegoides, Decne. in Jacquem. Voy. Bot. 103 = S. crataegoides.
cucullata, Miers, in Journ. Linn. Soc. xvii. (1879) 301 = S. obtusa.
cuneata, Miers, l. c. = S. cuneata.
elegans, Miers, l. c. = S. elegans.
ferruginea, Miers, l. c. 299 = S. ferruginea.
floribunda, Miers, l. c. = S. Samuntia.
foliosa, Miers, l. c. 300 = S. foliosa.
Gardneriana, Miers, l. c. = S. Gardneriana.
grandiflora, Miers, l. c. 299 = S. grandiflora.
Hamiltoniana, Miers, l. c. = S. racemosa.
hirsuta, Miers, l. c. 301 = S. hirsuta.
hispidula, Miers, l. c. = S. hispidula.
javanica, Miers, l. c. 302 = S. ferruginea.
jucunda, Miers, l. c. 301 = S. jucunda.
laeta, Miers, l. c. = S. laeta.
leiostachya, Miers, l. c. 300 = S. leiostachya.
leucantha, Miers, l. c. = S. leucantha.
Lohu, Miers, l. c. 298 = S. spicata.
lucida, Miers, l. c. 299 = S. theaefolia.
macrophylla, Miers, l. c. = S. macrophylla.
marginalis, Miers, l. c. 301 = S. marginalis.
microcarpa, Miers, l. c. 302 = S. laricifolia.
microphylla, Miers, l. c. 300 = S. microphylla.
monantha, Miers, l. c. = S. monantha.
nervosa, Miers, l. c. 299 = S. racemosa.
obovata, Miers, l. c. 301 = S. obtusa.
obtusa, Miers, l. c. 300 = S. obtusa.
oxyphylla, Miers, l. c. = S. oxyphylla.
pedicellata, Miers, l. c. = S. oxyphylla.
polycarpa, Miers, l. c. 301 = S. spicata.
polystachya, Miers, l. c. 300 = S. ferruginea.

LODHRA :—
pulchra, Miers, in Journ. Linn. Soc. xvii. (1879) 300 = Symplocos pulchra.
pyrifolia, Miers, l. c. 299 = S. pyrifolia.
racemosa, Miers, l. c. 298 = S. racemosa.
ramosissima, Miers, l. c. 299 = S. ramosissima.
Ribes, Miers, l. c. 302 = S. Ribes.
rubiginosa, Miers, l. c. 299 = S. rubiginosa.
rufescens, Miers, l. c. 301 = S. versicolor.
Samuntia, Miers, l. c. 298 = S. Samuntia.
spicata, Miers, l. c. = S. spicata.
sulcata, Miers, l. c. 300 = S. sulcata.
Verhuelli, Miers, l. c. 302 = S. ferruginea.
xanthophylla, Miers, l. c. = S. xanthophylla.

LODICULARIA, Beauv. Agrost. 108. t. 21 (1812) = **Hemarthria**, R. Br. (Gramin.).
capensis, Nees, Fl. Afr. Austr. 128 = H. compressa.
fasciculata, Beauv. Agrost. 108 = H. compressa.
fastigiata, Beauv. Agrost. Atlas, 14. t. 21. f. 6.—H. compressa.
Hamiltoniana, Nees, ex Steud. Syn. Pl. Gram. 358 = H. compressa.
laxa, Nees, ex Steud. l. c. = H. compressa.
peruviana, Meyen, Reise, ii. 71 = H. compressa.
protensa, Nees, ex Steud. Syn. Pl. Gram. 359 = H. protensa.

LODOICEA, Comm. ex Jaume St. Hil. Expos. i. 96 (1805). *PALMAE*, Benth. & Hook. f. iii. 939.
callipyge, *Comm. ex Jaume St. Hil. l. c.*—Ins. Seychell.
maldavica, Pers. ex H. Wendl. in Kerch. Palm. 250 = callipyge.
sechellarum, Labill. in Ann. Mus. Par. ix. (1807) 140. t. 13 = callipyge.

LOEFFLINGIA, Neck. Elem. ii. 153 (1790) = seq.

LOEFLINGA, R. Hedw. Gen. 30 (1806) = seq.

LOEFLINGIA, Linn. Sp. Pl. ed. I. 35 (1753). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
caspica, S. G. Gmel. Reise, iii. 310. t. 35. f. 1; et in J. F. Gmel. Syst. 105 = Polycarpon tetraphyllum.
gaditana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 23 = hispanica.
hispanica, Coss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 159 = micrantha.
hispanica, Hort. ex Steud. Nom. ed. II. ii. 64 = Scleranthus annuus.
hispanica, *Linn. Sp. Pl.* 35.—Reg. Mediterr.
indica, Retz. Obs. iv. 8 = Polycarpon Loeflingiae.
micrantha, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 25.—Hispan.
pentandra, *Cav. Ic.* ii. 39.—Hispan.
prostrata, Moench, Meth. 226 = hispanica.
pusilla, *Curran, in Bull. Calif. Acad.* i. n. III. (1885) 152.—Calif.
ramosissima, Weinm. in Flora, iii. (1820) 608 = Pentacaena polycnemoides.
renifolia, Lag. Gen. et Sp. Nov. 2 = Drymaria cordata.
squarrosa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 174.—Am. bor. occ.
texana, Hook. Ic. Pl. t. 275 = squarrosa.

LOESELIA, Linn. Gen. ed. I. 873 (1737). *POLEMONIACEAE*, Benth. & Hook. f. ii. 823.
HOITZIA, Juss. Gen. 136 (1789).
ROYENIA, Houst. ex Linn. Gen. ed. I. 348 (1737).
amplectens, *Benth. in DC. Prod.* ix. 320.—Mexic.
aristata, G. Don, Gen. Syst. iv. 248 = Gilia ciliata.
Cervantesii, G. Don, l. c. = glandulosa.
ciliata, Linn. Sp. Pl. 628 = Gilia ciliata.
coccinea, *G. Don, Gen. Syst.* iv. 247.—Mexic.
coerulea, *G. Don, l. c.* 248.—Mexic.
conglomerata, G. Don, l. c. = glandulosa.
effusa, *A. Gray, in Proc. Am. Acad.* xi. (1876) 86.—Calif.
glandulosa, *G. Don, Gen. Syst.* iv. 248.—Mexic.
Greggii, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 117.—Mexic.
guttata, A. Gray, l. c. xx. (1885) 302 (Giliae sp.).—Calif.
Havardi, A. Gray, l. c. xix. (1883) 87 (Giliae sp.).—Am. bor.
involucrata, *G. Don, Gen. Syst.* iv. 248.—Mexic.
Matthewsii, A. Gray, in Bot. Calif. ii. 466 (Giliae sp.).—Am. bor. occ.
nepetaefolia, G. Don, Gen. Syst. iv. 248 = glandulosa.

LOESELIA :—
pumila, *Walp. Rep.* vi. 527.—Mexic.
ramosissima, *Walp. l. c.* 526.—Mexic.
rupestris, *Benth. in DC. Prod.* ix. 319.—Mexic.
scabra, *Walp. Rep.* vi. 527.—Mexic.
scariosa, *Walp. l. c.*—Mexic.
Schottii, A. Gray, in Bot. Calif. ii. 466 = Gilia Schottii.
tenuifolia, A. Gray, in Proc. Am. Acad. xi. (1876) 86. (Giliae sp.).—Am. bor.

LOETHAINIA, Heynh. Nom. i. 883 (1840), absque char. = **Wiborgia**, Thunb. (Legumin.).
floribunda, Heynh. l. c. ii. 369 = W. obcordata.
obcordata, Heynh. l. c. i. 883 = W. obcordata.
sericea, Heynh. l. c. = W. sericea.

LOEVIGIA, Karst. & Triana, Fl. Neo-Granad. 22; ex Linnaea, xxviii. (1856) 434 = **Monochaetum**, Naud. (Melastom.).
ovalifolia, Karst. in Linnaea, l. c.—Hab. ?
sericea, Karst. l. c. = M. Bonplandia.

LOEZELIA, Adans. Fam. ii. 211 (1763) = **Loeselia**, Linn. (Polemon.).

LOGANIA, R. Br. Prod. 454 (1810). *LOGANIACEAE*, Benth. & Hook. f. ii. 791.
EUOSMA, Andr. Bot. Rep. t. 520 (1811).
alpina, *F. Muell. ex Benth. in Journ. Linn. Soc.* i. (1854) 94.—Austral.
angustifolia, Sieber, ex Spreng. Syst. iv. Cur. Post. 59 = floribunda.
Armstrongii, *J. Buch. in Trans. N. Z. Inst.* xiv. (1882) 347.—N. Zel.
bracteolata, Nees, in Lehm. Pl. Preiss. i. 367 = fasciculata.
buxifolia, *F. Muell. Fragm.* vi. 132.—Austral.
callosa, *F. Muell. l. c.* 134.—Austral.
campanulata, *R. Br. Prod.* 456.—Austral.
capensis, *Eckl. in S. Afr. Quart. Journ.* i. (1830) 371.—Afr. austr.
Centaurium, Nees, in Lehm. Pl. Preiss. ii. 240 = serpyllifolia.
ciliolata, Hook. f. Handb. N. Zeal. Fl. 737 (Veronicae sp.).—N. Zel.
cordifolia, *Hook. in Mitch. Journ. Trop. Austral.* 341.—Austral.
crassifolia, *R. Br. Prod.* 454.—Austral.
depressa, *Hook. f. Fl. N. Zel.* i. 177.—N. Zel.
elliptica, R. Br. Prod. 454 = ovata.
fasciculata, *R. Br. l. c.* 456.—Austral.
floribunda, *R. Br. l. c.*—Austral.
hispidula, Nees, in Lehm. Pl. Preiss. i. 368 = serpyllifolia.
hyssopoides, Nees, l. c. = serpyllifolia.
latifolia, *R. Br. Prod.* 454.—Austral.
ligustrina, *A. Cunn. ex Baxt. in Loud. Hort. Brit. Suppl.* ii. 646, nomen.—N. Zel.
linifolia, *Schlecht. in Linnaea*, xx. (1847) 605.—Austral.
longifolia, *R. Br. Prod.* 454.—Austral.
micrantha, *Benth. in Journ. Linn. Soc.* i. (1857) 94.—Austral.
nuda, *F. Muell. Fragm.* i. 129.—Austral.
ovata, *R. Br. Prod.* 455.—Austral.
paniculata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 12.—Austral.
pusilla, *R. Br. Prod.* 456.—Austral.
revoluta, R. Br. l. c. = floribunda.
serpyllifolia, *R. Br. l. c.*—Austral.
spermacocea, *F. Muell. Fragm.* vi. 134.—Austral.
stenophylla, *F. Muell. l. c.* i. 128.—Austral.
tetragona, Hook. f. Handb. N. Zeal. Fl. 188 (Veronicae sp.).—N. Zel.
vaginalis, F. Muell. Fragm. vi. 132 (sp. aggregat.).—Austral.

LOGANIA, J. F. Gmel. Syst. 422 (1791) = LOGHANIA, Scop. = **Ruyschia**, Jacq. (Ternstroem.).
pentacrina, J. F. Gmel. l. c. = R. Surubaea.

LOGFIA, Cass. in Bull. Soc. Philom. (1819) 143 = **Filago**, Linn. (Compos.).
brevifolia, Cass. in Dict. Sc. Nat. xxvii. 117 = F. arvensis.
dichotoma, Pomel, Nouv. Mat. Fl. Atl. 44 = F. arvensis ?
gallica, Coss. & Germ. in Ann. Sc. Nat. Sér. II. xx. (1843) 291. t. 13. f. A = F. gallica.

LOGFIA :—
lanceolata, Cass. in Dict. Sc. Nat. xxvii. 118 = Filago minima.
montana, Dum. Fl. Belg. 68 = F. arvensis.
minima, Dum. l. c. = F. minima.
spicata, Pomel, Nouv. Mat. Fl. Atl. 43.—Alger.
subulata, Cass. in Dict. Sc. Nat. xxvii. 116 = F. gallica.

LOGHANIA, Scop. Introd. 236 (1777) = **Ruyschia**, Jacq. (Ternstroem.).

LOISELERIA, Reichb. Fl. Germ. Excurs. 416, in nota (1831) = **Rhododendron**, Linn. (Ericac.).
indica, Reichb. l. c. = R. indicum.

LOISELEURIA, Desv. Journ. Bot. iii. (1814) 35. *ERICACEAE*, Benth. & Hook. f. ii. 595.
CHAMAECISTUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 401 (1821).
CHAMAELEDON, Link, Enum. Hort. Berol. i. 210 (1821).
procumbens, *Desf. Journ. Bot.* iii. (1814) 35.—Europ.

LOISELEURIA, Reichb. Nom. 127 (1841), vide LOISELERIA, Reichb.

LOLANARA, Rafin. Fl. Tellur. ii. 34 (1836). An *SAPOTACEA* ?—Cf. Hassk. Neuer Schl. zu Rumph's Herb. Amboin. 55.
odorata, Rafin. l. c.—Ind. or.

LOLIUM, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1202.
ARTHROCHORTUS, Lowe, in Hook. Kew Journ. viii. (1856) 301.
CRAEPALIA, Schrank, Baier. Fl. i. 102, 382 (1789).
CRYPTURUS, Link, in Linnaea, xvii. (1843) 387.
aechicum, Rouv. Monog. 35, sphalm. = *čechicum*.
aegyptiacum, Bell. ex Rouv. l. c. 43 = temulentum.
agreste, Hort. ex Roem. & Schult. Syst. ii. 748 = perenne.
album, Steud. Nom. ed. I. 493 = temulentum.
annuum, Bernh. Sem. Hort. Erf. (1801) = perenne.
annuum, Lam. Fl. Fr. iii. 620 = temulentum.
appenninum, *Brouss. Cat. Hort. Monsp.* (1804) 35, nomen.—Italia.
arenarium, Rouv. Monog. 38 = perenne.
aristatum, Lag. ex Nym. Consp. 445 = multiflorum.
arvense, Schrad. Fl. Germ. 399; Sm. Engl. Fl. i. 175 = remotum.
arvense, With. Arr. Brit. Pl. ed. III. ii. 168 = temulentum.
asperum, Roth, ex Kunth, Enum. Pl. i. 436 = perenne.
Berteronianum, Steud. Syn. Pl. Gram. 340.—Chili.
Boucheanum, Kunth, Rév. Gram. ii. t. 220 = perenne.
brasilianum, *Nees, Agrost. Bras.* 443.—Reg. Argent.
bromoides, Huds. Fl. Angl. ed. II. i. 55 = Festuca Myuros.
canadense, Bernh. ex Rouv. Monog. 27 = perenne.
canadense, *Michx. ex Roem. & Schult. Syst.* ii. 893.—Am. bor.
canariense, *Steud. Syn. Pl. Gram.* 340.—Ins. Canar.
čechicum, Opiz, Oekon.-Teckn. Fl. Boehm. (1836) 379 = multiflorum.
coelorachis, Forst. ex Steud. Nom. ed. II. ii. 64 = Lepturus repens.
complanatum, Schrad. Neues Journ. iv. (1810) 73 = perenne.
compositum, Thuill. Fl. Par. ed. II. 62 = perenne.
compressum, Boiss. & Orph. ex Nym. Consp. 845 = rigidum.
cristatum, Pers. Syn. i. 110 = perenne.
cylindricum, C. Koch, in Linnaea, xxi. (1848) 434 = rigidum.
decipiens, Dum. Obs. Gram. Belg. 98 = temulentum.
distachyum, Linn. Mant. i. 187 = Panicum ciliare.
distachyum, Willd. ex Steud. Nom. ed. II. ii. 65 = Leptochloa monostachya.
durum, C. Koch, in Linnaea, xxi. (1848) 434 = rigidum.
elegans, *Steud. Syn. Pl. Gram.* 341.—Afr. bor.
elongatum, Hort. ex Rouv. Monog. 35 = multiflorum.
felix, Rouv. Monog. 25 = perenne, multiflorum.
Festuca, Rasp. ex Mutel, Fl. Fr. iv. 111 = Festuca elatior.

LOLIUM:—
festucaceum, Link, Handb. i. 819 = perenne.
festucoides, Rasp. ex Mutel, Fl. Fr. iv. 112 = Festuca loliacea.
flagellare, Sprun. ex Boiss. Fl. Orient. v. 680 = rigidum.
Gaudini, *Parl. Fl. Ital.* i. 532, in obs.—Europ.
giganteum, Hort. ex Roem. & Schult. Syst. ii. 750 = maximum.
glumosum, *Plan. Fl. Gallega*, 409.—Hispan.
Gmelini, [Honck.] Verz. aller Gew. Teutschl. 328 = perenne?
gracile, Dum. Obs. Gram. Belg. 99 = perenne?
gracile, Parl. in Webb & Berth. Phyt. Canar. iii. 423 = temulentum.
Gussonei, Nym. Consp. 844 = temulentum.
Halleri, C. C. Gmel. Fl. Bad. i. 271 = perenne.
infelix, Rouv. Monog. 39 = temulentum.
italicum, A. Br. in Flora, xvii. (1834) 259 = multiflorum.
Jechelianum, Opiz, in Lotos, ii. (1852) 229; Seznam, 60 = perenne.
latum, Roth, ex Steud. Nom. ed. I. 492 = perenne.
lepturoides, *Boiss. Diagn.* Ser. I. xiii. 67.—Oriens.
linicolum, A. Br. in Flora, xvii. (1834) 258; Sond. in Koch, Syn. Fl. Germ. ed. II. 957 = perenne.
lucidum, Dum. Obs. Gram. Belg. 98 = temulentum.
macilentum, Delastre, Fl. Dép. Vienne, 498 = rigidum.
maximum, Guss. Fl. Sic. Syn. i. 60 = temulentum.
maximum, Willd. Sp. Pl. i. 462 = praec.?
Marschallii, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) n. III. 103.—Ross.
multiflorum, Gaud. Fl. Helv. i. 354 = Gaudini.
multiflorum, Guss. Fl. Sic. Syn. i. 58 = siculum.
multiflorum, *Lam. Fl. Fr.* iii. 621.—Gallia.
oechicum, Rouv. Monog. 53, sphalm. = *čechicum*.
Osiridis, Fig. & Delile, ex Rouv. l. c. 47. t. 2. f. 2, 3 = perenne?
perenne, Aitch. in Journ. Linn. Soc. xix. (1882) 194 = Agropyron repens.
perenne, *Linn. Sp. Pl.* 83.—Europ.; As. temp.
persicum, *Boiss. & Hohen. Diagn.* Ser. I. xiii. 66.—Oriens.
Phoenice, Rouv. Monog. 36 = seq.
Phoenix, Rouv. l. c. 37 = perenne.
Pseudo-italicum, Schur, Enum. Pl. Transs. 812 = perenne.
Pseudo-linicolum, Gennari, in Nuov. Giorn. Bot. ii. (1870) 97 = perenne.
remotum, Schrank, Baier. Fl. i. 382; Hoffm. Deutschl. Fl. ed. II. i. 63 = perenne.
repens, [Honck.] Verz. aller Gew. Teutschl. 328 = perenne.
rigidum, *Gaud. Agrost. Helv.* i. 334.—Europ.
rigidum, Weiss, ex Nym. Consp. 845 = subulatum.
robustum, Reichb. Fl. Germ. Excurs. Index, 74, nomen; Ic. Fl. Germ. xi. 2. t. 4. f. 1340 = temulentum.
romanum, *Sang. in Att. Accad. Pont. Lincei*, Ser. I. xviii. (1865) 221 (Quid?).—Italia.
rosetlanum, Fig. & Delile, ex Rouv. Monog. 31 = perenne.
scabrum, *J. & C. Presl, Rel. Haenk.* i. 267.—Peruv.; Chili.
siculum, *Parl. Fl. Palerm.* i. 252; *et Giorn. Bot. Ital.* ii. I. (1846) 90.—Sicilia.
Sonderi, Rouv. Monog. 47 = perenne?
speciosum, Stev. ex Bieb. Fl. Taur. Cauc. i. 80 = temulentum.
strictum, Decker, ex Steud. Nom. ed. II. ii. 65 = rigidum.
strictum, Presl, Cyp. & Gram. Sicul. (1820) 49 = perenne.
subulatum, *Vis. Fl. Dalm.* i. 90. t. 3.—Europ. austr.
temulentum, Bert. ex Steud. Nom. ed. II. ii. 65, partim = scabrum.
temulentum, *Linn. Sp. Pl.* 83.—Europ.; As. bor.
temulentum, Thunb. Fl. Jap. 55 = Brachypodium japonicum.
tenue, Bieb. Fl. Taur. Cauc. i. 80 = Marschallii.
tenue, Guss. Fl. Sc. Syn. i. 59 = rigidum.
tenue, Linn. Sp. Pl. ed. II. 122 = perenne.
triticoides, Janka, in Linnaea, xxx. (1859–60) 622 = temulentum.
vulgare, Host, Gram. Austr. i. t. 25 = perenne.

LOMANDRA, Labill. Nov. Holl. Pl. i. 92 (1804) = **Xerotes**, R. Br. (Juncac.) (1810).
angustifolia, Hort. Cels, ex Steud. Nom. ed. II. ii. 65 = seq.?

LOMANDRA:—
latifolia, Cels, Cat. Arb. (1817) 23 = Xerotes longifolia?
longifolia, Labill. Nov. Holl. Pl. i. 92. t. 119 = X. longifolia.
rigida, Labill. l. c. 93. t. 120 = X. rigida.

LOMANODIA, Rafin. Sylva Tellur. 97 (1838) = **Astronia**, Noronha (Melastom.).
glabra, Rafin. l. c. = A. Forsteri.
mucronata, Rafin. l. c. = Miconia holosericea.

LOMANTHERA, Rafin. Sylva Tellur. 96 (1838) = **Tetrazygia**, Rich. (Melastom.).
glandulosa, Rafin. l. c. = T. hispida.

LOMANTHES, Rafin. Sylva Tellur. 92 (1838) = **Phyllanthus**, Linn. (Euphorb.).
latifolia, Rafin. l. c. = P. thymoides.

LOMARESIS, Rafin. Fl. Tellur. iii. 59 (1836) = **Ornithogalum**, Tourn. (Liliac.).
alba, Rafin. l. c. = O. revolutum.

LOMARIA *coriacea*, Kunze, in Linnaea, x. (1836) 506 = Stangeria paradoxa.
eriopus, Kunze, l. c. xiii. (1839) 152; xviii. (1844) 116 = praec.
lagopus, T. Moore, in Hook. Kew Journ. v. (1853) 228 = praec.

LOMASTELMA, Rafin. Sylva Tellur. 107 (1838) = **Eugenia**, Mich. (Myrtac.).
elliptica, Rafin. l. c. = E. venosa?

LOMATIA, R. Br. in Trans. Linn. Soc. x. (1810) 199. *PROTEACEAE*, Benth. & Hook. f. iii. 183.
TRICONDYLUS, Salisb. in Knight, Prot. 121 (1809) nomen prius.
alnifolia, Poepp. ex Meissn. in DC. Prod. xiv. 450 = obliqua.
angustifolia, Schnizl. Ic. ii. t. 113 = longifolia.
chilensis, *C. Gay, Fl. Chil.* v. 310.—Chili.
cymosa, *Soland. ex Seem. Fl. Vit.* 187.—Ins. Tahiti.
dentata, *R. Br. in Trans. Linn. Soc.* x. (1810) 201.—Chili.
ferruginea, *R. Br. l. c.* 200.—China.
Fraseri, R. Br. Prot. Nov. 34 = ilicifolia.
fraxinifolia, *F. Muell. ex Benth. Fl. Australas.* v. 536.—Austral.
ilicifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 200.—Austral.
longifolia, *R. Br. l. c.*—Austral.
Milnei, Hook. Kew Journ. vii. (1855) t. 2 = Stenocarpus Milnei.
obliqua, *R. Br. in Trans. Linn. Soc.* x. (1810) 201.—Am. austr.
opaca, Klotzsch, ex Meissn. in DC. Prod. xiv. 450 = obliqua.
polymorpha, *R. Br. in Trans. Linn. Soc.* x. (1810) 200.—Austral.
silaifolia, *R. Br. l. c.* 199.—Austral.
sinuata, *R. Br. Prot. Nov.* 33.—Austral.
sylvatica, Poepp. ex Meissn. in DC. Prod. xiv. 449 = dentata.
tinctoria, *R. Br. in Trans. Linn. Soc.* x. (1810) 199.—Austral.
trinervis, *Montr. in Mém. Acad. Lyon*, x. (1860) 248.—N. Caled.

LOMATIUM, Rafin. in Journ. Phys. lxxxix. (1819) 101 = **Peucedanum**, Tourn. (Umbell.).
pubescens, Rafin. in Ser. Bull. Bot. (1830) 216 = P. foeniculaceum.
villosum, Rafin. in Journ. Phys. lxxxix. (1819) 101 = P. foeniculaceum.

LOMATOCARUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 59 (1840) = **Selinum**, Linn. (Umbell.).
alpinum, Fisch. & Mey. l. c. = Carum Lomatocarum.

LOMATOGONIUM, A. Br. in Flora, xiii. (1830) 221 = **Pleurogyne**, Eschsch. (Gentian.).
carinthiacum, A. Br. l. c. = P. carinthiaca.
inodora, Pritz. Ic. Ind. 656, sphalm. vide Lonas inodora.
rotatum, Fries, ex Nym. Consp. 500 = P. rotata.
Stellerianum, Kostel. Allg. Med-Pharm. Flora, iii. 1048 = P. carinthiaca.
sulcatum, Reichb. ex Kostel. l. c. = P. rotata.

LOMATOLEPIS, Cass. in Dict. Sc. Nat. xlviii. (1827) 422 = **Launaea**, Cass. (Compos.).
glomerata, Cass. l. c. 423 = Launaea glomerata.
nudicaulis, Cass. l. c. 424 = Launaea nudicaulis.

LOMATOPHYLLUM, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 166. *LILIACEAE*, Benth. & Hook. f. iii. 777.
PHYLLOMA, Ker-Gawl. in Bot. Mag. t. 1585 (1813).
aloiflorum, Nichols. Dict. Gard. ii. 296 = borbonicum.
borbonicum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 166.—Ins. Mascar.
cernuum, Boj. Hort. Maurit. 349 = Dracaena reflexa.
fusciflorum, *Boj. l. c.* 349.—Madag.
Jacquini, Spreng. Syst. ii. 74 = Aloë hexapetala.
macrum, *Salm-Dyck, in Schult. f. Syst.* vii. 1715.—Ins. Mascar.
marginatum, Hoffmgg. Verz. Pfl. 74 = borbonicum.
reflexum, Boj. Hort. Maurit. 349 = Dracaena reflexa.
rufocinctum, *Salm-Dyck, in Schult. f. Syst.* vii. 1715.—'Ind. or.'
tessellatum, *Boj. Hort. Maurit.* 349.—Madag.
umbraculiferum, Boj. l. c. = Dracaena umbraculifera.

LEMATOPODIUM, Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 305 = **Seseli**, Linn. (Umbellif.).
Karelinianum, Turcz. ex Stschegl. in Bull. Soc. Nat. Mosc. xxvii. (1834) I. 167.—As. centr.
Lessingianum, Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 306 = S. Lessingianum.
platyphyllum, Schrenk, l. c.—As. centr.

LOMATOZONA, Baker, in Mart. Fl. Bras. vi. II. 198 (1876). *COMPOSITAE*.
artemisiaefolia, *Baker, l. c.* 199.—Am. trop.

LOMAXETA, Rafin. New Fl. Am. iv. 72 (1836) = **Palafoxia**? Lag. (Compos.).
verrucosa, Rafin. l. c.—Am. bor.

LOMELOSIA, Rafin. Fl. Tellur. iv. 95 (1836) = **Scabiosa**, Tourn. (Dipsac.).
coronopifolia, Rafin. l. c. = S. ceratophylla.
palestina, Rafin. l. c. = S. palaestina.
simplex, Rafin. l. c. = S. stellata.
stellata, Rafin. l. c. = S. stellata.

LOMENIA, Pourr. in Mém. Acad. Toul. iii. (1788) 74. t. 5 = **Watsonia**, Mill. (Irid.).
borbonica, Pourr. l. c. 76.—Ins. Borbon.

LOMERIA, Rafin. Sylva Tellur. 56 (1838) = **Cestrum**, Linn. (Solan.).
purpurea, Rafin. l. c. (Quid?).—Peruv.?

LOMILIS, Rafin. Alsog. Am. 5 (1838) = **Hamamelis**?, Gronov.
ciliata, Rafin. l. c. = H. virginiana?

LOMOPLIS, Rafin. Sylva Tellur. 118 (1838) = **Mimosa**, Linn. (Legumin.).
acantholoba, Rafin. l. c. = M. acantholoba.
Ceratonia, Rafin. l. c. = M. Ceratonia.

LONAS, Adans. Fam. ii. 118 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 417.
inodora, *Gaertn. Fruct.* ii. 396. t. 165.—Reg. Mediterr.
minima, Cass. in Dict. Sc. Nat. xxvii. 167 = inodora.
umbellata, Cass. l. c. = inodora.

LONCHESTIGMA, Dun. in DC. Prod. xiii. I. 476 (1852) = **Jaborosa**, Juss. (Solan.).
bipinnatifidum, Phil. Fl. Atac. 42 = J. caulescens.
caulescens, Dun. in DC. Prod. xiii. 476 = J. caulescens.
crispum, Dun. l. c. 477 = J. crispa.
squarrosum, Dun. l. c. = J. decurrens.

LONCHOCARPUS, H. B. & K. Nov. Gen. et Sp. vi. 383 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 548.
CAPASSA, Klotzsch, in Peters, Reise Mossamb. Bot. 27. t. 5 (1861).
CLOMPANUS, Aubl. Pl. Gui. ii. 773 (1775), nomen prius.
ICTHYOCTONUM, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1884) 440.
NEUROSCAPHA, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 137.

LONCHOCARPUS:—
PHILENOPTERA, Fenzl, in Flora, xxvii. (1844) 312.
SPHINCTOLOBIUM, Vog. in Linnaea, xi. (1837) 417.
albiflorus, *G. Don, Gen. Syst.* ii. 236.—Afr. trop.
americanus, Steud. Nom. ed. II. i. 76, sphalm. = seq.
Amerimnum, DC. Prod. ii. 261 = Dalbergia Amerimnum.
apiciflorus, C. Muell. in Walp. Ann. vii. 796 = spiciflorus.
araripensis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 96.—Bras.
atropurpureus, *Benth. l. c.* 91.—Venezuela.
Barteri, *Benth. l. c.* 99.—Afr. trop.
Blackii, *Benth. Fl. Austral.* ii. 271.—Austral.
Blainii, *Wright, in Sauv. Fl. Cub.* 31.—Cuba.
brachypterus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 100.—Afr. trop.
campestris, *Mart. ex Benth. l. c.* 95.—Bras.
confertiflorus, *Benth. l. c.* 96.—Peruv.
costatus, *Benth. l. c.* 100.—Bras.
cyanescens, *Benth. l. c.* 96.—Afr. trop.
densiflorus, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 433.—Guian.
denudatus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 95.—Bras.
domingensis, DC. Prod. ii. 259 = sericeus.
eriophyllus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 94.—Mexic.
fasciculatus, *Benth. l. c.* 100.—Afr. trop.
Fendleri, *Benth. l. c.* 94.—Venezuela.
filipes, *Benth. l. c.*—Bras.
floribundus, Benth. in Ann. Nat. Hist. Ser. I. iii. (1839) 432 = L. Nicou.
formosianus, DC. Prod. ii. 260 = sericeus.
fruticosus, Span. in Linnaea, xv. (1841) 191 = Tephrosia candida.
glabrescens, *Benth. in Hook. Kew Journ.* ii. (1850) 233.—Bras.
guatemalensis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 87.—Mexic.
guineensis, *D. Dietr. Syn. Pl.* iv. 1052.—Afr. trop.
hedyosmus, *Miq. in Linnaea*, xviii. (1844) 564.—Guian.
heptaphyllus, DC. Prod. ii. 259 = latifolius.
Heudelotianus, Baill. Adansonia, vi. (1866) 222 = Barteri.
hondurensis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 91.—Honduras.
ichthyoctonus, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 440.—Madag.
inconstans, *Vatke, in Linnaea*, xliii. (1880–82) 106.—Madag.
lanceolatus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 92.—Mexic.
latifolius, *H. B. & K. Nov. Gen. et Sp.* vi. 383.—Am. trop.
laxiflorus, *Guill. & Perr. Fl. Seneg. Tent.* i. 226.—Afr. trop.
leonensis, *G. Don, Gen. Syst.* ii. 236.—Afr. trop.
longifolius, Benth. in Mart. Fl. Bras. xv. i. 289 = Derris longifolia.
macrocarpus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 91.—Am. trop.
macrophyllus, H. B. & K. Nov. Gen. et Sp. vi. 383 = sericeus.
macrostachyus, *Hook. f. in Hook. Niger Fl.* 317.—Afr. trop.
maculatus, DC. Prod. ii. 260 = Gliricidia maculata.
mollis, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 95.—Bras.
negrensis, *Benth. l. c.* 98.—Bras.
neuroscapha, *Benth. l. c.* 89.—Bras.
Nicou, *DC. Prod.* ii. 261.—Guian.
nitidulus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 98.—Bras.
nitidus, *Benth. l. c.* 92.—Bras.
obovatus, *Benth. l. c.* 93.—Mexic.
obtusus, *Benth. l. c.* 95.—Bras.
oxycarpus, DC. Prod. ii. 261 = latifolius.
pallescens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 242.—Afr. trop.
parviflorus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 89.—Mexic.
Peckolti, *Wawra, in Flora*, xlvii. (1864) 230.—Bras.
pentaphyllus, H. B. & K. Nov. Gen. et Sp. vi. 383 = latifolius.
phaseolifolius, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 93.—Mexic.

LONCHOCARPUS:—
Philenoptera, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 97 = laxiflorus.
praecox, *Mart. ex Benth. l. c.* 96.—Bras.
Pterocarpus, DC. Prod. ii. 260 = Derris guianensis.
pubescens, *DC. l. c.*—Venezuela.
punctatus, H. B. & K. Nov. Gen. et Sp. vi. 383 = violaceus, *H. B. & K.*
pyxidarius, DC. Prod. ii. 260 = sericeus.
rariflorus, *Mart. ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 98.—Bras.
roseus, *DC. Prod.* ii. 260.—Am. austr.
rubiginosus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 92.—Guiana.
rufescens, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 432.—Guiana.
rugosus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 92.—Mexic.
sepium, *DC. Prod.* ii. 260.—Am. austr.
sericeus, *H. B. & K. Nov. Gen. et Sp.* vi. 383.—Am. trop.
Sophiae, Kotschy & Peyr. Pl. Tinn. 3. t. 2 = laxiflorus.
spiciflorus, *Mart. ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 91.—Bras.
Spruceanus, *Benth. l. c.* 99.—Bras.
subglaucescens, *Mart. ex Benth. l. c.* 96.—Bras.
Swartzii, DC. Prod. ii. 261 = violaceus.
tomentosus, Tul. in Arch. Mus. Par. iv. (1844) 82 = sericeus.
Turpinii, H. B. & K. Nov. Gen. et Sp. vi. 385 = sericeus.
unifoliolatus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 90.—Mexic.
velutinus, *Benth. in Seem. Bot. Voy. Herald*, 111.—Am. centr.
violaceus, *H. B. & K. Nov. Gen. et Sp.* vi. 383, in adnot.—Am. austr.
violaceus, *Oliver, in Trans. Linn. Soc.* xxix. (1872) 63.—Afr. trop.
virgilioides, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 94.—Bras.

LONCHOMERA, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 93 (1872). *ANONACEAE.*
leptopoda, *Hook. f. & Thoms. l. c.* 94.—Malacca.

LONCHOPHORA, Dur. in Duch. Rev. Bot. ii. (1847) 432. *CRUCIFERAE*, Benth. & Hook. f. i. 71.
capiemontana, *Dur. l. c.*—Alger.
Guyoniana, *Dur. Expl. Algér.* t. 72.—Alger.

LONCHOSTEPHUS, Tul. in Arch. Mus. Par. vi. (1852) 198. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 109.
elegans, *Tul. l. c.* 199.—Bras.

LONCHOSTOMA, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 350. t. 10. *BRUNIACEAE*, Benth. & Hook. f. i. 673.
GRAVENHORSTIA, Nees, in Lindl. Nat. Syst. ed. II. 439 (1836).
PTYXOSTOMA, Vahl, in Danske Nat. Selsk. Skriv. vi. (1810) 95, nomen prius.
acutiflorum, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 353. t. 10. f. 1.—Afr. austr.
monostylis, *Sond. in Harv. & Sond. Fl. Cap.* ii. 317.—Afr. austr.
obtusiflorum, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 352. t. 10. f. 2.—Afr. austr.

LONCHOSTYLIS, Torr. in Ann. Lyc. N. York, iii. (1836) 370 = **Rynchospora**, Vahl (Cyperac.).

LONCODILIS, Rafin. Fl. Tellur. ii. 24 (1836) = **Ornithogalum**, Tourn. (Liliac.).
scapigera, Rafin. l. c.—Hab.?

LONCOMELOS, Rafin. Fl. Tellur. ii. 24 (1836) = **Ornithogalum**, Tourn. (Liliac.).
japonicum, Rafin. l. c. = Scilla japonica.
latifolium, Rafin. l. c. = O. latifolium.
narbonense, Rafin. l. c. = O. narbonense.
pyramidale, Rafin. l. c. = O. narbonense.
pyrenaicum, Rafin. l. c. = O. pyrenaicum.
secundum, Rafin. l. c. = Urginea secunda.
suaveolens, Rafin. l. c. = O. suaveolens.

LONCOSTEMON, Rafin. Fl. Tellur. ii. 21 (1836) = **Allium**, Tourn. (Liliac.).
victoriale, Rafin. l. c. = A. victorialis.

LONCOXIS, Rafin. Fl. Tellur. iii. 58 (1836) = **Ornithogalum**, Tourn. (Liliac.).
sulfurea, Rafin. l. c. = O. pyrenaicum.

LONDESIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 40 (1835) = **Chenolea**, Thunb. (Chenop.).
eriantha, Fisch. & Mey. l. c.—Reg. Casp.
eriantha, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 736 = Kirilowia eriantha.

LONGCHAMPIA, Willd. in Ges. Nat. Fr. Berl. Mag. v. (1811) 159 = **Leyssera**, Linn. (Compos.).
capillifolia, Willd. l. c. 159, 160 = L. capillifolia.

LONGETIA, Baill. Adansonia, ii. (1861–62) nomen; et vi. (1866) 352. t. 9. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 279.
buxoides, *Baill. l. c.* ii. (1861–62) 228.—N. Caled.
depauperata, *Baill. l. c.* xi. (1873) 100.—N. Caled.

LONICERA, Adans. Fam. ii. 157 (1763) = **Hamelia**, Jacq. (Rubiac.).

LONICERA, Linn. Gen. ed. I. 57 (1737). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 5.
CAPRIFOLIUM, Tourn. ex Linn. Syst. ed. I (1735).
CHAMAECERASUS, [Tourn.] Medic. Phil. Bot. i. 126 (1789).
COBAEA, Neck. Elem. i. 129 (1790).
DISTEGIA, Rafin. New Fl. Am. iii. 20 (1836).
EUCHYLIA, Dulac, Fl. Hautes-Pyr. 463 (1867).
ISICA, Moench, Meth. 504 (1794).
ISIKA, Adans. Fam. ii. 501 (1763).
ITIA, Molina, Sagg. Chil. ed. II. 281 (1810); ex Roem. & Schult. Syst. v. 259, in obs. (1819).
KANTEMON, Rafin. New Fl. Am. iii. 19 (1836).
NINTOOA, Sweet, Hort. Brit. ed. II. 258 (1830).
PERICLYMENUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 250 (1745); Mill. Gard. Dict. ed. VI. (1752).
PHENIANTHUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83.
XYLOSTEUM, Rupp. Fl. Jen. ed. Hall. 250 (1745).
acuminata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 176.—Reg. Himal.
acutifolia, Rafin. New Fl. Am. iii. 19 (Quid?).—Am. bor.
adenocarpa, Guss. Fl. Sic. Prod. Suppl. 58 = implexa.
affinis, *Hook. & Arn. Bot. Beech. Voy.* 264.—China; Japon.
affinis, Jacquem. Voy. Bot. 77 = asperifolia.
alba, Linn. Sp. Pl. 175 = Chiococca racemosa.
Alberti, *Regel, in Act. Hort. Petrop.* vii. (1880) 550.—Turkest.
albiflora, *Torr. & Gray, Fl. N. Am.* ii. 6.—Am. bor. occ.
alpigena, Gueldenst. Reisen, i. 429 = iberica.
alpigena, *Linn. Sp. Pl.* 174.—Europ.; As. bor.
alpigena, Schangin, in Pall. N. Nord. Beitr. vi. 93 = hispida.
alpigena, Siev. l. c. vii. 274 = microphylla.
alpina, Steud. Nom. ed. II. i. 278, sphalm. = alpigena.
altaica, Pall. Fl. Ross. t. 37 = caerulea.
Altmannii, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 41.—Turkest.
americana, C. Koch, Wochenschr. x. (1867) 279 = grata.
angustata, Steud. Nom. ed. II. ii. 66 = tatarica.
angustifolia, Rafin. New Fl. Am. iii. 19 (Quid?).—Am. bor.
angustifolia, *Wall. Cat.* n. 480; *et in DC. Prod.* iv. 337.—Reg. Himal.
angustifolia, Wender. in Marb. Schr. ii. (1831) 247 = tatarica.
arborea, *Boiss. in Bibl. Univ. Genèv.* (1838); *ex ej. Elench.* 55.—Hispan.
arborea, Boiss. Voy. Bot. Espagne, ii. 277, partim = nummulariaefolia.
asperifolia, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 166.—Reg. Himal.
atrosanguinea, Carr. in Rev. Hortic. (1866) 337 = dimorpha.
Aucherii, *Jaub. & Spach, Illustr.* i. 137. t. 73.—Persia; Arab.
aurea, *Auch. ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 170.—Oriens.
aureo-reticulata, Hort. ex T. Moore, in Flor. & Pom. 1862 (1863) 136 = reticulata.

LONICERA:—
balearica, DC. Fl. Fr. Suppl. 499 = implexa.
bicolor, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 84 (Quid ?).—Reg. Himal.
biflora, *Desf. Fl. Atlant.* i. 184.—Afr. bor.
brachypoda, DC. Prod. iv. 335 = japonica.
bracteata, Royle, Illustr. Bot. Himal. 237 = hispida.
bracteolaris, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 106.—Reg. Caucas.
Brandtii, *Franch. & Sav. Enum. Pl. Jap.* ii. 385.—Japon.
Breweri, A. Gray, in Proc. Am. Acad. vi. (1864–65) 537 = conjugialis.
Brownii, Hort. ex Fl. des Serres, xi. (1856)123 = ciliosa.
Buergeriana, Blume, ex C. Koch, Dendrol. ii. 1. 18 = flexuosa.
bulbalina, Linn. f. Suppl. 146 = Burchellia capensis.
Bungeana, Ledeb. Fl. Ross. ii. 391 = microphylla.
caerulea, Gueldenst. Reisen, i. 423 = orientalis.
caerulea, Kit. in Linnaea, xxxii. (1863) 430 = nigra.
caerulea, *Linn. Sp. Pl.* 174.—Reg. bor. temp.
californica, Torr. & Gray, Fl. N. Am. ii. 7 = hispidula.
campanifolia, G. Don, in Loud. Hort. Brit. ed. I. 471 = nigra.
canadensis, Muhl. ex Roem. & Schult. Syst. v. 260 = ciliata.
canescens, Schousb. Vextr. Marok. 88 = biflora.
caprifolioides, *C. Koch, Ind. Sem. Hort. Berol.* (1871) *App.* 3.—Hab. ?
Caprifolium, *Linn. Sp. Pl.* 173.—Europ.; Reg. Cauc.
Caprifolium, Schousb. Vextr. Marok. 87 = etrusca.
Caprifolium, Tenore, Syll. Fl. Neap. 104 = Stabiana.
caroliniana, Marsh. Arbust. 80 = supervirens.
carpathica, Kit. in Linnaea, xxxii. (1863) 430 = nigra.
caucasica, *Pall. Fl. Ross.* i. 59.—Reg. Caucas.
cerasina, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 41.—Japon.
Chamissoi, *Bunge, in Kirill. Lonic. Russ.* 26; *ex Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 48).—Ins. Sachalin.
chinensis, Wats. Dendrol. t. 117 = japonica.
chlamydophora, C. Koch, in Linnaea, xxiv. (1851) 478 = iberica.
chrysantha, *Turcz. in Bull. Soc. Nat. Mosc.* xi. (1838) 93; xviii. (1845) 304.—Sibir.
ciliata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 463 = ciliosa.
ciliata, *Muhl. Cat.* 22.—Am. bor.
ciliosa, Hook. & Arn. Bot. Beech. Voy. 349 = hispidula.
ciliosa, *Poir. Encyc. Suppl.* v. 612.—Am. bor. occ.
coccinea, Pers. Syn. i. 213 = sempervirens.
cochinchinensis, G. Don, Gen. Syst. iii. 447 = japonica.
coerulea, Auct. vide caerulea.
confusa, *DC. Prod.* iv. 333.—China.
confusa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 269 = japonica.
conjugialis, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 67. f. 15.—Am. bor. occ.
connata, Meerb. Pl. Sel. Ic. t. 11 = sempervirens.
corymbosa, Linn. Sp. Pl. 175 = Loranthus tetrandrus.
cyrenaica, Viv. Fl. Lib. Spec. 12. t. 8. f. 1 = etrusca.
davurica, Siev. in Pall. N. Nord. Beitr. vii. 198 = Viburnum davuricum.
decipiens, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 170.—Reg. Himal.
dentata, Rafin. New Fl. Am. iii. 19 (Quid ?).—Am. bor.
depressa, Royle, Illustr. Bot. Himal. 226 = parvifolia.
Diervilla, Linn. Sp. Pl. 175 = Diervilla trifida.
dimorpha, *Tausch, in Flora*, xxi. (1838) 1. *Beibl.* 80.—Hab. ?
dioica, Linn. Syst. ed. XII. 165 = glauca, *Hill*.
Dioscoridis, Roubieu, ex Nym. Consp. 321 = implexa.
discolor, Lindl. Bot. Reg. (1844) sub t. 33 = orientalis.
diversifolia, Carr. in Rev. Hortic. xxxvii. (1866) 99 = flexuosa.
diversifolia, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 178 = quinquelocularis.
Douglasii, DC. Prod. iv. 332 = hirsuta.
dumetorum, Pers. Syn. i. 213 = L. Xylosteum.
dumosa, A. Gray, Pl. Wright. ii. 66 = albiflora.
edulis, Turcz. in Bull. Soc. Nat. Mosc. xviii. (1845) 306 = caerulea.
Elisae, *Franch. Pl. Davidian.* i. (1884) 152.—China.
elliptica, Rafin. New Fl. Am. iii. 18 (Quid ?).—Am. bor.
elliptica, Royle, Illustr. Bot. Himal. 236 = hypoleuca.
epsomiensis, Hort. ex C. Koch, Dendrol. ii. 1. 12 = glauca, *Hill*.

LONICERA:—
eriensis, Rafin. New Fl. Am. iii. 18 (Quid ?).—Am. bor.
etrusca, Hort. ex Tausch, in Flora, xxi. (1838) 1. Beibl. 80 = dimorpha.
etrusca, Lej. Fl. Spa, ii. 299 = L. Periclymenum.
etrusca, *Santi, Viagg. Mont.* i. 113. t. 1.—Reg. Mediterr.
etrusca, Tenore, Fl. Nap. i. 82 = Stabiana.
Ferdinandi, *Franch. Pl. Davidian.* i. (1884) 151.—China.
Finlaysoniana, Wall. Cat. n. 6302 = japonica.
flammea, Salisb. Prod. 138 = sempervirens.
flava, *Sims, Bot. Mag.* t. 1318.—Am. bor.
flexuosa, Thunb. in Trans. Linn. Soc. ii. (1794) 330 = japonica.
floribunda, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 107. t. 8.—Persia.
florida, *Regel, Cat. Pl. Hort. Aksakov.* 86.—Hab. ?
fragrantissima, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 75.—China.
Fraseri, Poit. in Rev. Hortic. Ser. II. v. (1846) 125 = flava.
germanica, F. G. Dietr. Weim. Fl. 94 = L. Periclymenum.
gibbosa, [Moç. & Sesse,] ex DC. Prod. iv. 336 = involucrata.
gibbosa, *Willd. ex Roem. & Schult. Syst.* v. 257.—Mexic.
glabra, *Boiss. ex Tchihat. Asie Min. Bot.* ii. 193, nomen.—As. Min.
glabrata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 175.—Reg. Himal.
glauca, *Hill, Hort. Kew.* 446.—Am. bor.
glauca, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 166.—Reg. Himal.
glaucophylla, Hook. f. & Thoms. l. c. 165 = Leycesteria glaucophylla.
Glehni, *F. Schmidt, Reis. Amurl.* 143.—Ins. Sachalin.
glutinosa, *Vis. Fl. Dalm.* iii. 18.—Dalmat.
Goldii, Spreng. Syst. i. 758 = hirsuta.
Govaniana, Wall. Cat. n. 481; DC. Prod. iv. 337 = orientalis.
gracilipes, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 85.—Japon.
gracilis, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 77.—Reg. Himal.
grata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 231.—Am. bor.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 173.—Afghan.
hellenica, *Orph. in Boiss. Diagn.* Ser. II. ii. 108.—Graecia.
heterophylla, *Decne. in Jacquem. Voy. Bot.* 80. t. 88.—Reg. Himal.
hetrusca, Host, Fl. Austr. i. 298 = etrusca.
hirsuta, *Eaton, Man. Bot.* ed. II. 307.—Am. bor.
hirtiflora, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 166 = macrantha.
hispanica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 52 = L. Periclymenum.
hispida, *Pall. ex Roem. & Schult. Syst.* v. 258.—Reg. Himal.; Sibir.
hispidula, *Dougl. ex Lindl. Bot. Reg.* t. 1761, in syn.—Am. bor. occ.
humilis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 370.—As. centr.
hypoglauca, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 270 = affinis.
hypoleuca, *Decne. in Jacquem. Voy. Bot.* 81. t. 89.—Reg. Himal.
iberica, *Bieb. Fl. Taur. Cauc.* i. 158.—Reg. Caucas.
implexa, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 231.—Europ. austr.
intermedia, Kellogg, in Proc. Calif. Acad. ii. (1863) 154 = involucrata.
interrupta, Benth. Pl. Hartw. 313 = hispidula.
involucrata, *Banks, ex Spreng. Syst.* i. 759.—Am. bor.
japonica, Andr. Bot. Rep. t. 583 = confusa.
japonica, *Thunb. Fl. Jap.* 89.—Reg. Himal.
japonica, Wall. Cat. 473; in Roxb. Fl. Ind. ed. Carey, ii. 174 = macrantha.
javanica, *DC. Prod.* iv. 334.—Java.
Karelini, *Bunge, in Kirill. Lonic. Russ.* 33.—As. centr.
kurdistana, Schlecht. ex Boiss. Fl. Orient. iii. 7 = nummulariaefolia.
lanceolata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 177 = decipiens.
latifolia, Guss. Fl. Sic. Prod. i. 250 = implexa.

LONICERA:—
Ledebourii, Eschsch. in Mém. Acad. Pétersb. x. (1826) 284 = involucrata.
leiantha, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 188.—Burma.
Leschenaultii, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 270 = affinis.
Leschenaultii, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 178.—Ind. or.
ligustrina, *Wall. in Roxb. l. c.* 179.—Reg. Himal.
linderifolia, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 50.—Japon.
linearis, *Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 168 = spinosa.
longiflora, DC. Prod. iv. 333 = macrantha.
longifolia, Royle, Illustr. Bot. Himal. 236 = praec. ?
Loureiri, Blume, Bijdr. 653 = oxylepis.
Loureiri, *DC. Prod.* iv. 334.—Malaya.
Loureirii, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 172 = acuminata.
Luschani, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 108.—Mesopot.
Maacki, *Herd. in Bull. Soc. Nat. Mosc.* xxxvii. (1864) 204.—Manchur.
macrantha, *Spreng. Syst.* iv. *Cur. Post.* 82.—Reg. Himal.
macrogyne, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 85 (Quid ?).—Reg. Himal.
marilandica, Linn. Sp. Pl. 175 = Spigelia marilandica.
Maximowiczii, *Maxim. Prim. Fl. Amur.* 137.—Reg. Amur.
media, Murr. in Comm. Gotting. vii. (1776) 28. t. 3 = glauca, *Hill*.
micrantha, *Trautv. ex Regel, in Act. Hort. Petrop.* v. (1877) 609.—Turkestan.
microphylla, Hook. Fl. Bor. Am. i. 283 = hispidula.
microphylla, *Willd. ex Roem. & Schult. Syst.* v. 258.—Reg. Himal.; Sibir.
Mociniana, DC. Prod. iv. 336 = involucrata.
mollis, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 45 = etrusca.
mollis, Wall. Cat. n. 6301 = Leschenaultii.
mollissima, Blume, ex Maxim. in Bull. Acad. Pétersb. xxiv. (1878) 37 = affinis.
mongolica, Pall. Reise, iii. 721 = Viburnum davuricum.
Morrowi, A. Gray, in Perry, Jap. Exped. ii. 313 = chrysantha.
multiflora, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 167 = confusa.
Myrtillus, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 168 = parvifolia.
nervosa, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 39.—China.
nigra, *Linn. Sp. Pl.* 173.—Europ.; As. bor.
nigra, Thunb. Fl. Jap. 89 = japonica.
Nummularia, *Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc.* (1839) 156.—Turcoman.
nummulariaefolia, *Jaub. & Spach, Illustr.* i. 133. t. 70.—Oriens.
oblongifolia, *Hook. Fl. Bor. Am.* i. 284. t. 100.—Am. bor.
obovata, *Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 169.—Reg. Himal.
occidentalis, Hook. Fl. Bor. Am. i. 282 = ciliosa.
ochroleuca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = L. Xylosteum.
odora, Salisb. Prod. 138 = L. Periclymenum.
odoratissima, [Lindl. in] Gard. Chron. (1858) 314 = fragrantissima.
Olgae, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 609, adnot.—Turkestan.
orientalis, Bieb. Fl. Taur. Cauc. i. 158 = caucasica.
orientalis, *Lam. Encyc.* i. 731.—Oriens; Reg. Himal.
ovata, *Buch.-Ham. in Wall. Cat.* n. 6300.—Reg. Himal.
ovata, Rafin. New Fl. Am. iii. 18 (Quid ?).—Am. bor.
oxylepis, *Miq. Fl. Ind. Bat.* ii. 125.—Java.
oxyphylla, Edgew. in Trans. Linn. Soc. xx. (1846) 60 = alpigena.
Pallasii, Ledeb. Ind. Sem. Hort. Dorpat. (1821) 20 = caerulea.
pallida, Host, Fl. Austr. i. 298 = L. Caprifolium.
parasitica, Linn. Sp. Pl. 175 = Loranthus lonicerioides.
parviflora, Lam. Encyc. i. 728 = glauca, *Hill*.
parvifolia, *Edgew. in Trans. Linn. Soc.* xx. (1846) 60.—Reg. Himal.
perfoliatum, Edwards, in Phytol. N. S. ii. (1857) 198, sphalm. = L. Caprifolium.

LONICERA :—
Periclymenum, Gouan, Hort. Monsp. 101 = etrusca.
Periclymenum, *Linn. Sp. Pl.* 173.—Europ.
Periclymenum, Lour. Fl. Cochinch. 150 = confusa.
persica, Jaub. & Spach, Illustr. i. 132 = nummulariaefolia.
phyllocarpa, *Maxim. Prim. Fl. Amur.* 138.—Reg. Amur.
Phylomelae, *Carr. in Rev. Hortic.* (1872) 160.—Hab.?
pilosa, Kellogg, in Proc. Calif. Acad. i. (1855) 63 = hispidula.
pilosa, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 47.—Japon.
pilosa, *Willd. ex H. B. & K. Nov. Gen. et Sp.* iii. 427.—Mexic.
plantierensis ×, André, in Illustr. Hortic. xviii. (1871) 233. t. 86.
pubescens, Sweet, Hort. Brit. ed. I. 195 = hirsuta.
punicea, Sims, Bot. Mag. t. 2469 = Symphoricarpus puniceus.
purpurascens, *Walp. Rep.* ii. 449.—Reg. Himal.
pyrenaica, Hort. ex Ledeb. Fl. Ross. ii. 388 = tatarica.
pyrenaica, *Linn. Sp. Pl.* 174.—Mont. Pyren.
pyrenaica, Pall. Reise, ii. 568 = caerulea.
quadrifolia, *Willd. Sp. Pl.* i. 986.—Hab.?
quinquelocularis, *Hardw. in As. Res.* vi. (1799) 351.—Reg. Himal.
racemosa, Auct. ex Steud. Nom. ed. II. ii. 67 = Symphoricarpos racemosus.
Raeseri, Heldr. ex Boiss. Fl. Orient. iii. 5 = etrusca.
ramosissima, *Franch. & Sav. ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 47.—Japon.
repens, Hassk. Cat. Hort. Bog. Alt. 116 = japonica.
reticulata, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 167.—China.
reticulata, Maxim. in Bull. Acad. Pétersb. xxiv. (1878) 40 = venulosa.
reticulata, Rafin. New Fl. Am. iii. 18 (Quid?).—Am. bor.
Roeseri, Heldr. ex Nym. Consp. 321 = etrusca.
rotundifolia, Hort. ex C. C. Gmel. Hort. Carlsr.; ex Steud. Nom. ed. I. 493 (Quid?).—Hab.?
Royleana, Wall. Cat. n. 478 = quinquelocularis.
rubra, Gilib. Fl. Lituan. i. 6 = L. Xylosteum.
rupestris, Rafin. New Fl. Am. iii. 17 (Quid?).—Am. bor.
rupicola, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 168.—Reg. Himal.
Ruprechtiana, *Regel, Ind. Sem. Hort. Petrop.* (1869) 19.—Reg. Amur.
saxatilis, *Fl. Corc. in Ionios Anthol.* v. 210.—Ins. Corcir.
Semenovi, *Regel, in Act. Hort. Petrop.* v. (1877) 608.—Turkestan.
sempervirens, *Linn. Sp. Pl.* 173.—Am. bor.
sericea, Royle, Illustr. Bot. Himal. 236; Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 167 = purpurascens.
sibirica, Hort. ex DC. Prod. iv. 335 = tatarica.
sibirica, *Vest, ex Roem. & Schult. Syst.* v. 259 = alpigena.
sicula, Ucria, in Roem. Arch. i. 1. 68 = Putoria calabrica.
Sieversiana, Bunge, Suppl. Fl. Alt. 18 = microphylla.
speciosa, Wender. Ind. Sem. Hort. Marb. (1825) = sempervirens?
spinosa, *Jacquem. Voy. Bot.* 78.—Reg. Himal.
splendida, *Boiss. Elench.* 54.—Hispan.
Stabiana, *Guss. ex Pasq. in Rendic. Acc. Sc. Nat.* xiii. 1874) 142.—Italia.
Standishii, *Hook. Bot. Mag.* t. 5709.—China.
stipulata, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 165 = Pentapyxis stipulata.
suavis, Salisb. Prod. 138 = L. Caprifolium.
subspicata, Hook. & Arn. Bot. Beech. Voy. 349 = hispidula.
Sullivantii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 76.—Am. bor.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 537.—Sumatra.
Symphoricarpos, Linn. Sp. Pl. 175 = Symphoricarpos vulgaris.
syringantha, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 49.—China.
tangutica, *Maxim. l. c.* 48.—China.
tatarica, *Linn. Sp. Pl.* 173.—Sibiria.
Tatarinowii, *Maxim. Prim. Fl. Amur.* 138.—China.
Telfairii, Hook. & Arn. Bot. Beech. Voy. 190 = confusa.

LONICERA :—
tenuiflora, *Regel & Winkl. in Act. Hort. Petrop.* vi. (1879) 305.—Turkestan.
tomentella, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 167.—Reg. Himal.
translucens, *Carr. in Rev. Hortic.* (1872) 240.—Hab.?
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 39.—Japon.
tubulosa, Benth. Pl. Hartw. 37 = pilosa.
turcomanica, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 156, nomen; *Trautv. in Act. Hort. Petrop.* i. (1870) 274.—Turkestan.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 299.—Turkestan.
utahensis, *S. Wats. in Bot. King's Exped.* 133.—Utah.
variegata, *Regel, Cat. Pl. Hort. Aksakov.* 87.—Hab.?
velutina, DC. Prod. iv. 337 (excl. syn. in parte) = caerulea.
venulosa, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 542.—Japon.
verticillata, *Steud. Nom.* ed. I. 493.—Jamaic.
Vidalii, *Franch. & Sav. Enum. Pl. Jap.* ii. 386.—Japon.
villosa, DC. Prod. iv. 337 = caerulea.
villosa, Muhl. Cat. 22 = hirsuta.
virginiana, Marsh. Arbust. 80 = sempervirens.
viscidula, Boiss. Fl. Orient. iii. 5 = etrusca.
Webbiana, Wall. Cat. n. 476; et ex DC. Prod. iv. 336 = alpigena.
Wightiana, Wall. l. c. sub n. 1520 = ligustrina.
xylosteoides ×, Tausch, in Flora, xxi. (1838) II. 736.
Xylosteum, Georgi, Reise, i. 201; Regel, Fl. Ussur. 80. n. 237 = chrysantha.
Xylosteum, *Linn. Sp. Pl.* 174.—Europ.; As. bor.
Xylosteum, Lour. Fl. Cochinch. 150 = Loureiri.
Xylosteum, Sibth. & Sm. Fl. Graec. iii. 18. t. 223 = nummulariaefolia.
zeylanica, Gaertn. Fruct. i. 137. t. 27 = Loranthus longiflorus.

LONICERA, Plum. ex Gaertn. Fruct. i. 132. t. 27. t. 2 (1788) = **Loranthus**, Linn.

LONTANUS, Benth. & Hook. f. Gen. iii. 939 (1883) = seq.

LONTARUS, Adans. Fam. ii. 25 (1763) = **Borassus**, Linn. (Palm.).
domestica, Gaertn. Fruct. i. 21. t. 8 = B. flabellifera.

LOOSA, Jacq. Obs. Bot. iii. 15 (1768) = **Loasa**, Adans.

LOPADOCALYX, Klotzsch, in Lehm. Pl. Preiss. i. 178 (1844) = **Olax**, Linn.
phyllanthoides, Klotzsch, l. c. = O. Phyllanthi.

LOPANTHUS, Vitm. Summa Veg. ii. 252 (1789) = LOPHANTHUS, Forst. = **Waltheria**, Linn. (Stercul.).

LOPESIA, Juss. in Ann. Mus. Par. iii. (1804) 317 = seq.

LOPEZIA, Cav. Ic. i. 12. t. 18 (1791). *ONAGRARIEAE*, Benth. & Hook. f. i. 791.
JEHLIA, Hort. Germ. ex Planch. in Fl. des Serres, vii. (1851–52) 177.
PISAURA, Bonato, Monog. (1793); ex Endl. Gen. 1194.
albiflora, *Schlecht. in Linnaea*, xxiv. (1851) 696.—Mexic.
annua, Hort. ex DC. Prod. iii. 62 = oppositifolia.
axillaris, Schweigg. in Koenigsb. Arch. Naturw. i. (1812) 236 = coronata.
cordata, *Hornem. Hort. Hafn.* ii. *Add.* 949.—Mexic.
coronata, *Andr. Bot. Rep.* t. 551.—Mexic.
frutescens, Roem. & Schult. Syst. i. 34 = miniata.
fruticosa, Hort. Par. ex Schrank, in Nov. Act. Nat. Cur. ix. (1818) 92; et Pl. Rar. Hort. Monac. t. 20 = miniata.
Galeottii, *Planch. in Fl. des Serres*, vii. (1851–52) 178.—Mexic.
grandiflora, *Zucc. in Flora*, xv. (1832) II. *Beibl.* 101.—Mexic.
haematodes, *Kunze, in Linnaea*, xvii. (1843) 579.—Mexic.

LOPEZIA :—
hirsuta, *Jacq. Coll.* v. 5. t. 15. f. 4.—Mexic.
hispida, Sweet, ex Steud. Nom. ed. II. ii. 68 = praec.?
insignis, *Hemsl. Diagn. Pl. Nov.* 16.—Mexic.
integrifolia, *DC. Prod.* iii. 62.—Mexic.
lineata, *Zuccagn. Pl. Nov. Cogn. Fasc.* ii. 31.—Mexic.
longiflora, *Decne. in Rev. Hortic.* Sér. IV. iii. (1854) 221.—Mexic.
macrantha, *Benth. ex Regel, in Schweiz. Zeitschr. Gartenb.* viii. (1850) 25, cum t. b.—Hab.?
macrophylla, *Benth. Pl. Hartw.* 83.—Guatemala.
mexicana, Jacq. Coll. Suppl. 3 = racemosa.
miniata, *DC. Cat. Hort. Monsp.* (1813) 121.—Mexic.
minima, *Lag. ex Schrank, in Nov. Act. Nat. Cur.* ix. (1818) 92.—Mexic.
minuta, Lag. Gen. et Sp. Nov. 1 = coronata.
oppositifolia, *Lag. l. c.*—Mexic.
paniculata, *Seem. Bot. Voy. Herald*, 120.—Panama.
pubescens, *H. B. & K. Nov. Gen. et Sp.* vi. 96.—Mexic.
pumila, *Bonpl. Desc. Jard. Malm.* 67.—Mexic.
racemosa, *Cav. Ic.* i. 12. t. 18.—Mexic.
thymifolia, Humb. & Bonpl. ex Schult. Mant. i. 50 (Quid?).—Hab.?
trichota, *Schlecht. in Linnaea*, xii. (1838) 273.—Mexic.
trichotoma, Steud. Nom. ed. II. ii. 68 = praec.

LOPHACTIS, Rafin. Atl. Journ. 152 (1832). *'COMPOSITAE'* (Quid?).
uniflora, Rafin. l. c.—Am. bor.

LOPHALIX, Rafin. Sylva Tellur. 70 (1838) = **Alloplectus**, Mart. (Gesner.).
bicolor, Rafin. l. c. = A. cristatus.
coccinea, Rafin. l. c. = A. coccineus.

LOPHANDRA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = **Erica**, Tourn.
blanda, G. Don, Gen. Syst. iii. 810 = E. cornuta.
cubica, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 174 = E. cubica.
seriphiifolia, G. Don, Gen. Syst. iii. 810 = E. seriphiifolia.

LOPHANTHERA, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 328; et in Arch. Mus. Par. iii. (1843) 315 (Monog. 61. t. 6). *MALPIGHIACEAE*, Benth. & Hook. f. i. 255.
Kunthiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 316.—Guiana.
longiflora, C. Muell. in Walp. Ann. vii. 469, sphalm. = seq.
longifolia, Griseb. in Mart. Fl. Bras. xii. i. 25 = Kunthiana.

LOPHANTHERA, Rafin. New Fl. Am. ii. 58 (1836) = **Sopubia**, Buch.-Ham. (Scrophul.).
delphinifolia, Rafin. l. c. = S. delphinifolia.

LOPHANTHUS, Adans. Fam. ii. 194 (1763) = **Nepeta**, Linn. (Labiat.).

LOPHANTHUS, Benth. in Bot. Reg. t. 1282 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1198.
VLECKIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
anisatus, *Benth. in Bot. Reg.* t. 1282.—Am. bor. occ.
chinensis, *Benth. l. c.* sub t. 1282.—As. bor.
multifidus, Benth. l. c. = Nepeta lavandulacea.
nepetoides, *Benth. l. c.*—Am. bor.
obtusifolius, Heynh. Nom. i. 477 = chinensis.
rugosus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 31.—China.
scrophulariaefolius, *Benth. in Bot. Reg.* sub t. 1282.—Am. bor.
tomentosus, *Regel, Pl. Nov. Fedsch.* 63.—Turkestan.
urticifolius, *Benth. in Bot. Reg.* sub t. 1282.—Am. bor.

LOPHANTHUS, Forst. Char. Gen. 27. t. 14 (1776) = **Waltheria**, Linn. (Stercul.).
tomentosus, Forst. l. c. 28 = W. Lophanthus.

LOPHARINA, Neck. Elem. i. 215 (1790) = **Erica**, Tourn.

LOPHATHERUM, Brongn. in Duperr. Voy. Coq. Bot. 49. t. 8 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1191.
ACROËLYTRUM, Steud. in Flora, xxix. (1846) 20.
ALLELOTHECA, Steud. Syn. Pl. Gram. 117 (1855).
annulatum, *Franch. & Sav. Enum. Pl. Jap.* ii. 605.—Japon.
dubium, *Steud. Syn. Pl. Gram.* 300.—Ind. or.
elatum, *Zoll. & Mor. Syst. Verz. Zoll.* 102.—Java.
geminatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 300.—Madag.
gracile, *Brongn. in Duperr. Voy. Coq. Bot.* 50. t. 8.—Ins. Amboina.
humile, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 282.—Japon.
japonicum, Steud. Syn. Pl. Gram. 300 = elatum.
Lehmanni, *Nees, ex Steud. l. c.*—Ind. or.
multiflorum, *Steud. l. c.*—Java.
pilosulum, *Steud. l. c.* 428.—Japon.

LOPHERINA, Juss. in Dict. Sc. Nat. xxvii. 180 (1823) = LOPHARINA, Neck. = **Erica**, Tourn.

LOPHIA, Desv. in Ham. Prod. Pl. Ind. Occ. 47 (1825) = **Alloplectus**, Mart. (Gesner.).
phoenicea, Desv. l. c. = A. cristatus.

LOPHIARIS, Rafin. Fl. Tellur. iv. 40 (1836) = **Oncidium**, Sw. (Orchid.).
fragrans, Rafin. l. c. 41 = O. Lanceanum.

LOPHIOCARPUS, Miq. Illustr. Fl. Arch. Ind. 50 (1871) = **Sagittaria**, Lam. (Alismac.).
calycinus, Micheli, in DC. Monog. Phan. iii. 61 = S. calycina.
cordifolia, Miq. Illustr. Fl. Arch. Ind. 50 = S. cordifolia.
guyanensis, Micheli, in DC. Monog. Phan. iii. 62 = S. guyanensis.
Lappula, Miq. Illustr. Fl. Arch. Ind. 50 = S. Lappula.
Seubertianus, Micheli, in DC. Monog. Phan. iii. 64 = S. Seubertianus.

LOPHIOCARPUS, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 55. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 49.
WALLINIA, Moq. in DC. Prod. xiii. II. 143 (1859).
Burchellii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 50.—Afr. austr.
polystachyus, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 56.—Afr. austr.
tenuissimus, *Hook. f. Ic. Pl.* xv. (1884) 50.—Afr. austr.

LOPHIOLA, Ker-Gawl. in Bot. Mag. t. 1596 (1814). *HAEMODORACEAE*, Benth. & Hook. f. iii. 677.
americana, Wood, Class-Book (1861), 697 = aurea.
aurea, *Ker-Gawl. Bot. Mag.* t. 1596.—Am. bor.

LOPHIOLEPIS, Cass. in Dict. Sc. Nat. xxvii. 180 (1823) = **Cnicus**, Tourn. (Compos.).
araneosa, Cass. l. c. 181 = C. fimbriatus.
calocephala, Cass. l. c. = C. ciliatus.
dubia, Cass. l. c. 183 = C. lanceolatus.
nutans, Cass. l. c. 182 = Carduus cernuus.

LOPHION, Spach, Hist. Vég. Phan. v. 516 (1836) = **Viola**, Linn.
canadense, Spach, l. c. 517 = V. canadensis.

LOPHIRA, Banks, ex Gaertn. f. Fruct. iii. 52. t. 188 (1805). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 193.
africana, Loud. ex G. Don, Gen. Syst. i. 814 = alata.
alata, *Banks, ex Gaertn. f. Fruct.* iii. 52. t. 188. f. 2.—Afr. trop.
simplex, G. Don, Gen. Syst. i. 814 = alata.

LOPHIUM, Steud. Nom. ed. II. ii. 769, in syn. (1841) = LOPHION, Spach = **Viola**, Tourn.

LOPHOBIOS, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
cristata, Rafin. l. c. = E. cristata.
Terracina, Rafin. l. c. = E. Terracina.

LOPHOCACHRYS, Koch, ex DC. Prod. iv. 238 (1830); Bertol. Fl. Ital. iii. 455 (1837) = **Hippomarathrum**, Link (Umbellif.).
echinophora, Bertol. l. c. 456 = H. Bocconi.

LOPHOCARPUS, Link, Diss. Bot. Suerin. (1795) 52 = **Froelichia**, Moench (Amarant.).
procumbens, Link, l. c. = F. interrupta.

LOPHOCARYA, Nutt. ex Moq. in DC. Prod. xiii. II. 113, in syn. (1849) = **Atriplex**, Tourn. (Chenop.).
spinosa, Nutt. ex Moq. l. c. = A. canescens.

LOPHOCHLAENA, Nees, in Ann. Nat. Hist. Ser. I. i. (1838) 283 = **Pleuropogon**, R. Br. (Gramin.).
californica, Nees, l. c. = P. californicum.
refracta, A. Gray, in Proc. Am. Acad. viii. (1872) 409 = P. refractum.

LOPHOCHLOA, Reichb. Fl. Germ. Excurs. 42 (1830) = **Koeleria**, Pers. (Gramin.).
paradoxa, Scheele, in Flora, xxvii. (1844) 61 = K. paradoxa.
phleoides, Reichb. Fl. Germ. Excurs. 42 = K. phleoides.
villosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 184 = K. villosa.

LOPHOCLINIUM, Endl. in Bot. Zeit. i. (1843) 457 = **Podotheca**, Cass. (Compos.).
album, Endl. l. c. = P. fuscescens.
citrinum, Endl. l. c. = P. gnaphalioides.
Manglesii, Endl. l. c. = P. angustifolia.

LOPHOGLOTIS, Rafin. Fl. Tellur. iv. 49 (1836) = **Sophronitis**, Lindl. (Orchid.).

LOPHOGYNE, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 99. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 111.
arculifera, *Tul. & Wedd. l. c.* 100.—Bras.
helicandra, *Tul. l. c.* 99.—Bras.

LOPHOLAENA, DC. Prod. vi. 335 (1837); et in Deless. Ic. Sel. iv. t. 59 (1839). *COMPOSITAE*, Benth. & Hook. f. ii. 441.
Dregeana, *DC. ll. cc.* 336; 26.—Afr. austr.
platyphylla, *Benth. in Hook. Ic. Pl.* t. 1113.—Afr. austr.

LOPHOLEPIS, Decne. in Arch. Mus. Par. i. (1839) 147. *GRAMINEAE*, Benth. & Hook. f. iii. 1123.
HOLBOELLIA, Wall. in Hook. Bot. Misc. ii. (1831) 144. t. 76.
ornithocephala, *Steud. Syn. Pl. Gram.* 112.—Ind. or.

LOPHOLOMA, Cass. in Dict. Sc. Nat. xliv. 37 (1826) = **Centaurea**, Linn. (Compos.).
alpestris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. Scabiosa.
Scabiosa, Cass. in Dict. Sc. Nat. xliv. 37 = C. Scabiosa.

LOPHOPETALUM, Wight ex Arn. in Ann. Nat. Hist. Ser. I. iii. (1839) 150. *CELASTRINEAE*, Benth. & Hook. f. i. 362.
celastroides, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 615.—Burma.
dubium, *M. Laws. l. c.* 616.—Malacca.
filiforme, *M. Laws. l. c.*—Burma.
fimbriatum, *Wight, Illustr.* i. 178.—Ind. or.; Burma.
floribundum, *Wight, l. c.*—Burma.
fuscescens, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 202.—Singapore.
grandiflorum, Wight, ex Arn. in Ann. Nat. Hist. Ser. I. iii. (1839) 151 = Euonymus grandiflorus.
javanum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 598 = fimbriatum.
littorale, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 164.—Burma.
lucidum, Wight, ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 608 = Euonymus grandiflorus.
ovatum, Presl, Bot. Bemerk. 32 = fimbriatum.
pallidum, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 615.—Malacca.
reflexum, *M. Laws. l. c.* 616.—Malacca.
Wallichii, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 299.—Burma.
Wightianum, *Arn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 151.—Ind. or.

LOPHOPHYLLUM, Griff. Notul. iv. 313. t. 491 (1854). *MENISPERMACEAE* [Benth. & Hook. f. i. 38.]
bicristata, *Griff. l. c.*—Ind. or.

LOPHOPHYTUM, Schott & Endl. Melet. i. t. 1. (1832). *BALANOPHOREAE*, Benth. & Hook. f. iii. 238.
ARCHIMEDEA, Leandr. Sacram. in Ann. Sc. Nat. Sér. II. vii. (1837) 32.
LEPIDOPHYTUM, Hook. f. in Lindl. Veg. Kingd. ed. III. 90 (1853), nomen.
bolivianum, *Wedd. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 185. t. 10.—Bolivia.
Leandri, *Eichl. in Mart. Fl. Bras.* iv. II. 59.—Bras.
mirabile, *Schott & Endl. Melet.* i. t. 1.—Bras.
mirabile, Wedd. in Ann. Sc. Nat. Sér. III. xiv. (1850) 185 = Leandri.
Weddelii, *Hook. f. in Trans. Linn. Soc.* xxii. (1856) 30, 49. t. 9.—N. Granat.

LOPHOPTERIS, Griseb. in Linnaea, xiii. (1839) 182 = seq.

LOPHOPTERYS, A. Juss. in Deless. Ic. Sel. iii. 18. t. 29 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 255.
splendens, *A. Juss. l. c.*—Guiana.

LOPHOSCIADIUM, DC. Coll. Mém. v. 57 (1829); Prod. iv. 207 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 905.
Barrelieri, Griseb. Spicil. Fl. Rumel. i. 372 = Ferula monticola.
meifolium, *DC. Coll. Mém.* v. 57.—Banat.
meifolium, C. Koch, in Linnaea, xix. (1847) 38 = Thirkeanum.
millefolium, Steud. Nom. ed. II. i. 68, sphalm. = praec.
olympicum, *C. Koch, in Linnaea*, xix. (1847) 38.—As. Min.
setifolium, C. Koch, l. c. 39 = Ferula pauciflora.
silaifolium, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 326.—Europ. austr.
Thirkeanum, *Boiss. Diagn.* Ser. II. ii. 94.—Anatolia.

LOPHOSPERMUM, D. Don, in Trans. Linn. Soc. xv. (1827) 351 = **Maurandia**, Orteg. (Scrophul.).
atrosanguineum, Zucc. in Abh. Akad. Muench. (1829–30) 305. t. 13 = Rhodochiton volubile.
erubescens, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 68 = M. erubescens.
physalodes, D. Don, in Trans. Linn. Soc. xv. (1827) 353 = Melasma hispidum.
Rhodochiton, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 250 = Rhodochiton volubile.
scandens, D. Don, in Trans. Linn. Soc. xv. (1827) 353.—Mexic.
scandens, Sweet, Brit. Flow. Gard. Ser. II. t. 68 = M. erubescens.

LOPHOSTACHYS, Pohl, Pl. Bras. Ic. ii. 93. t. 161, 162 (1831). *ACANTHACEAE*, Benth. & Hook. f. ii. 1094.
Alvarezia, *Nees, in DC. Prod.* xi. 246.—Peruv.
diandra, *Nees, in Mart. Fl. Bras.* ix. 70.—Bras.
falcata, *Nees, l. c.* 67.—Bras.
floribunda, *Pohl, Pl. Bras. Ic.* ii. 95. t. 162.—Bras.
laxiflora, *Nees, in Mart. Fl. Bras.* ix. 68.—Bras.
montana, *Mart. ex Nees, l. c.* 68.—Bras.
monticola, Mart. ex Nees, in DC. Prod. xi. 416 = Beloperone monticola.
nemoralis, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 69.—Bras.
patula, *Mart. l. c.* 68.—Bras.
semiovata, *Nees, in Mart. l. c.* 69.—Bras.
sessiliflora, *Pohl, Pl. Bras. Ic.* ii. 95, 96. t. 163.—Bras.
villosa, *Pohl, l. c.* 94. t. 161.—Bras.

LOPHOSTEMON, Schott, in Wien. Zeitschr. iii. (1830) 772; ex Endl. in Linnaea, vi. (1831) Litt. 54 = **Tristania**, R. Br. (Myrtac.).
arborescens, Schott, ll. cc. 772; 54 = T. conferta.
depressum, Schott, ll. cc. = T. depressa.

LOPHOSTEPHUS, Harv. Thes. Cap. ii. 9. t. 113 (1863) = **Anisotoma**, Fenzl (Asclep.).
mollis, Harv. l. c. = A. cordifolia.

LOPHOSTOMA, Meissn. in Mart. Fl. Bras. v. I. 72 (1855) = **Linostoma**, Wall. (Thymel.).
calophylloides, Meissn. l. c. = Lin. calophylloides.
ovatum, Meissn. in DC. Prod. xiv. 600 = Lin. ovatum.

LOPHOSTYLIS, Hochst. in Flora, xxv. (1842) 229 = **Securidaca**, Linn. (Polygal.).
angustifolia, Hochst. l. c. 231 = S. longipedunculata.
floribunda, Welw. Apont. 562 = S. Welwitschii.
javanica, Miq. Fl. Ind. Bat. i. II. 128 = S. tavoyana.
oblongifolia, Hochst. in Flora, xxv. (1842) 231 = S. longipedunculata.
pallida, Klotzsch, in Peters, Reise Mossamb. Bot. 115. t. 22 = S. longipedunculata.

LOPHOTAENIA, Griseb. Spicil. Fl. Rumel. i. 377 (1843) = **Malabaila**, Hoffm. (Umbellif.).
aurea, Griseb. l. c. = M. aurea.

LOPHOXERA, Rafin. Fl. Tellur. iii. 42 (1836) = **Celosia**, Linn. (Amarant.).
caudata, Rafin. l. c. = C. caudata.
comosa, Rafin. l. c. = C. cristata.
paniculata, Rafin. l. c. = C. paniculata.
polygonoides, Rafin. l. c. = C. polygonoides.
racemosa, Rafin. l. c. = C. cristata ?

LOPHOZONIA, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 396 = **Fagus**, Tourn. (Cupul.).
heterocarpa, Turcz. l. c. = F. obliqua.

LOPIMIA, Mart. in Nov. Act. Nat. Cur. xi. (1823) t. 96 = **Pavonia**, Cav. (Malvac.).
malacophylla, Mart. l. c. = P. velutina.

LORANTEA, Steud. Nom. ed. I. 494 (1821) = LORENTEA, Orteg. = **Sanvitalia**, Gualt. (Compos.).

LORANTHEA, Steud. Nom. ed. II. ii. 68 (1841) = praec.

LORANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = **Loranthus**, Linn.
corynitum, St. Lag. l. c. = Loranthus Corynitis.
pentagonum, St. Lag. l. c. = Loranthus Pentagonia.
polyrrhizum, St. Lag. l. c. = Loranthus polyrrhizos.
scurrulum, St. Lag. l. c. = Loranthus Scurrula.
spirostylum, St. Lag. l. c. = Loranthus Spirostylis.
stelum, St. Lag. l. c. = Loranthus Stelis.
taguum, St. Lag. l. c. = Loranthus Tagua.

LORANTHUS, Linn. Syst. ed. II. 22 (1740).
LORANTHACEAE, Benth. & Hook. f. iii. 207.
ANTRIBA, Rafin. Sylva Tellur. 126 (1838).
ATKINSONIA, F. Muell. Fragm. v. 34 (1865).
BARATRANTHUS, Miq. Fl. Ind. Bat. i. I. 834 (1855).
DENDROPEMON, Schult. f. Syst. vii. 1729 (1830).
DENDROPHTHOE, Mart. in Flora, xiii. (1830) 109.
ELYTRANTHE, Blume, in Schult. f. Syst. vii. 1730 (1830).
EPICOILA, Rafin. Sylva Tellur. 126 (1838).
ETUBILA, Rafin. l. c. 125 (1838).
GAIADENDRON, G. Don, Gen. Syst. iii. 431 (1834).
GLUTAGO, Comm. ex Poir. in Dict. Sc. Nat. xix. 87 (1821).
HELIXANTHERA, Lour. Fl. Cochinch. 142 (1790).
HYPHIPUS, Rafin. Sylva Tellur. 125 (1838).
ITICANIA, Rafin. l. c. (1838).
LANTHORUS, Presl, Epim. Bot. 256 (1850).
LEPEOSTEGERES, Blume, in Schult. f. Syst. vii. 1731 (1830).
LICHTENSTEINIA, Wendl. Coll. ii. 4. t. 39 (1808).
LIPOTACTES, Schult. f. Syst. vii. 1729 (1830).
LONICERA, Adans. Fam. ii. 157 (1763).
LOXANTHERA, Blume, in Schult. f. Syst. vii. 1612 (1830).
MACROSOLEN, Schult. f. l. c. 1731 (1830).
MEIENA, Rafin. Sylva Tellur. 125 (1838).
MOQUINIA, Spreng. f. Tent. Syst. Suppl. 9 (1828).
NOTANTHERA, G. Don, Gen. Syst. iii. 428 (1834).
ORYCTANTHUS, Eichl. in Mart. Fl. Bras. v. II. 87. tt. 29, 30 (1861).
PASSOWIA, Karst. in Bot. Zeit. x. (1852) 305.
PELTOMESA, Rafin. Sylva Tellur. 126 (1838).
PHOENICANTHEMUM, Schult. f. Syst. vii. 1729 (1830).
PHRYGILANTHUS, Eichl. in Mart. Fl. Bras. v. II. 45 (1868).
PHTHIRUSA, Mart. in Flora, xiii. (1830) 110.
PSITTACANTHUS, Mart. l. c. 106 (1830).
SCURRULA, G. Don, Gen. Syst. iii. 421 (1834).
SPIROSTYLIS, Presl, ex Schult. f. Syst. vii. 163 (1830).
STREPSIMELA, Rafin. Sylva Tell r. 159 (1838).

LORANTHUS:—
STRUTHANTHUS, Mart. in Flora, xiii. (1830) 102.
TAGUARIA, Rafin. Sylva Tellur. 125 (1838).
TAPINANTHUS, Blume, in Schult. f. Syst. vii. 1730 (1830).
TOLYPANTHUS, Blume, l. c. 1731 (1830).
TRISTERIX, Mart. in Flora, xiii. (1830) 108.
Acaciae, *Zucc. in Abh. Akad. Muench.* iii. (1837–43) 249. t. 2. f. 3.—Palaestin. ; Nubia.
acacioides, *A. Cunn. ex Benth. Fl. Austral.* iii. 392.—Austral.
acinarius, *Mart. in Schult. f. Syst.* vii. 130.—Bras.
acuminatus, *Ruiz & Pav. Fl. Peruv.* iii. 49.—Peruv.
acutifolius, *Ruiz & Pav. l. c.* 48.—Peruv.
acutus, Pohl, ex DC. Prod. iv. 287 = podopterus.
Adamsii, *Cheesem. in Trans. N. Z. Inst.* xiii. (1881) 296.—N. Zel.
aduncus, *G. F. W. Mey. Prim. Fl. Esseq.* 149.—Guiana.
affinis, *Mart. in Schult. f. Syst.* vii. 151.—Bras.
albidus, *Blume, Bijdr.* 665.—Burma ; Malaya.
albiflorus, *Hook. Ic. Pl.* t. 683.—N. Granat.
alternatus, Willd. ex Schult. f. Syst. vii. 162 = eugenioides.
alveolatus, *H. B. & K. Nov. Gen. et Sp.* iii. 444.—Ecuador.
alyxifolius, *F. Muell. ex Benth. Fl. Austral.* iii. 391.—Austral.
americanus, Lam. ex Steud. Nom. ed. II. ii. 69 = Plumieri.
americanus, *Linn. Sp. Pl.* 331 ; *Jacq. Select. Am.* 97. t. 67.—Am. trop.
americanus, Sieber, ex Presl, Bot. Bemerk. 75 = martinicensis.
americanus, Sw. Obs. 132 = claviceps.
americanus, *Vell. Fl. Flum.* 138 ; iii. t. 148.—Bras.
amplexicaulis, *H. B. & K. Nov. Gen. et Sp.* iii. 445.—Venezuela.
amplexicaulis, Wall. Cat. n. 520 ; et ex DC. Prod. iv. 306 = longiflorus.
amplexifolius, DC. l. c. 305 = longiflorus.
ampullaceus, *Roxb. Hort. Beng.* 87 ; *Fl. Ind.* i. 189.—Ind. or. ; Malaya.
ampullaceus, Wall. Cat. n. 506 B = capitellatus.
anceps, DC. Prod. iv. 315 = oblongo-ellipticus.
anceps, *Desv. in Ham. Prod. Pl. Ind. Occ.* 33.—Guiana.
anceps, Willd. ex Schult. f. Syst. vii. 131 = conduplicatus.
angustifolius *R. Br. ex Benth. Fl. Austral.* iii. 390.—Austral.
angustifolius, H. B. & K. Nov. Gen. et Sp. iii. 442 = Phoradendron angustifolium.
aphyllus, *Miers, ex DC. Prod.* iv. 307.—Chili.
arboreus, Mutis, ex DC. l. c. iv. 315 = L. Tagua.
Arnottianus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 284.—Sumatra.
Arnottianus, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 361 = lepidophyllus.
artensis, *Montr. in Mem. Acad. Lyon*, x. (1860) 213.—N. Caled.
Atkinsonae, *Benth. in Hook. Ic. Pl.* t. 1319.—Austral.
atropurpureus, *Blume, Bijdr.* 669.—Java.
attenuatus, *Pohl, in DC. Prod.* iv. 315.—Bras.
attenuatus, Willd. ex Schult. f. Syst. vii. 1643 = eugenioides.
aurantiacus, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral. 101 = pendulus.
auriculatus, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 174.—Mexic.
avenis, *Blume, Bijdr.* 663.—Java.
avicularis, Kunth, ex Griseb. Fl. Brit. W. Ind. 749 = occidentalis ?
avicularis, *Mart. ex Schult. f. Syst.* vii. 132.—Bras.
avium, Pohl, ex Eichl. in Mart. Fl. Bras. v. II. 77 = marginatus.
axanthus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 262.—Sumatra.
bahajensis, *Korth. l. c.* 285.—Borneo.
Baroni, *Baker, in Journ. Bot.* xx. (1882) 266.—Madag.
Belvisii, DC. Prod. iv. 303 = lanceolatus.
Berteroi, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 357.—Ins. J. Fernand.
bibracteolatus, *Hance, in Journ. Bot.* xviii. (1880) 301.—China.
bicalyculatus, *Mart. ex Schult. f. Syst.* vii. 128.—Bras.
bicolor, Roxb. Pl. Corom. ii. 19. t. 139 = longiflorus.

LORANTHUS:—
Bidwillii, *Benth. Fl. Austral.* iii. 390.—Austral.
biflorus, Desr. in Lam. Encyc. iii. 600 = L. Scurrula ?
bifurcatus, *Benth. Fl. Austral.* iii. 393.—Austral.
biracemosus, *Jacq. Enum. Pl. Carib.* 35.—Ind. occ.
biternatus, *Hoffmgg. in Schult. f. Syst.* vii. 124.—Bras.
Blancoanus, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 355.—Ins. Philipp.
Blumeanus, Schult. f. Syst. vii. 119 = avenis.
Blumeanus, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 362, partim = obtusatus.
Bojeri, *Baker, Fl. Maurit.* 135.—Ins. Mascar.
bracteatus, DC. Prod. iv. 306 partim = tomentosus.
bracteatus, *Heyne, ex Roxb. Fl. Ind. ed. Carey*, ii. 220.—Ind. or.
bracteatus, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = cucullaris.
bracteatus, Willd. ex DC. Prod. iv. 318 = puracensis.
Brandisianus, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 63.—Burma.
brasiliensis, *Desr. in Lam. Encyc.* iii. 596.—Bras.
brasiliensis, Spreng. ex DC. Prod. iv. 293 = portoricensis.
buddajoides, Walp. Rep. ii. 440, sphalm. = *budleioides*.
buddleioides, Thw. Enum. Pl. Zeyl. 136, partim = cordifolius.
budleioides, Desr. in Lam. Encyc. iii. 600 = L. Scurrula.
Burchellii, *Eckl. & Zeyh. Enum.* 358.—Afr. austr.
buxifolius, *Cham. in Linnaea*, iii. (1828) 207.—Chili.
cactorum, Hook. & Arn. Bot. Beech. Voy. 25 = aphyllus.
caesius, *Spreng. Syst.* ii. 132.—Chili.
calycinus, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyssin.
calyculatus, *DC. Coll. Mém.* vi. t. 10.—Mexic.
Candolleanus, Wight & Arn. Prod. 385 = recurvus.
canescens, *Burch. Trav.* ii. 90 ; *et ex DC. Prod.* iv. 304.—Afr. austr.
cansjeraefolius, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 173.—Costa Rica.
canus, F. Muell. in Hook. Kew Journ. viii. (1856) 145 = L. Quandang.
capitellatus, *Wight & Arn. Prod.* 382.—Ind. or.
cardiphyllus, *Willd. ex Schult. f. Syst.* vii. 162.—Am. austr.
carinatulus, Wall. ex DC. Prod. iv. 296 = ampullaceus.
carnosus, Wall. Cat. n. 552 = pulverulentus.
Casuarinae, Miq. in Lehm. Pl. Preiss. i. 279 = linophyllus.
catocarpus, *Benth. Pl. Hartw.* 190.—N. Granat.
celastroides, *Sieber, ex Schult. f. Syst.* vii. 163.—Austral.
celtidifolius, *Willd. ex Schult. f. l. c.* 161.—Afr. trop.
Chamissonis, Molkenb. in Miq. Pl. Jungh. 116, in nota = robustus.
chinensis, *DC. Coll. Mém.* vi. t. 7.—China.
chloranthus, *Miq. Stirp. Surin. Sel.* 205.—Guiana.
chordocladus, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 173.—Nicaragua.
chrysanthoides, *Korth. in Verh. Batav. Gen.* xvii. (1839) 269.—Borneo.
chrysanthus, *Blume, ex DC. Prod.* iv. 300.—Java.
cinctus, *Mart. ex Schult. f. Syst.* vii. 134.—Bras.
cinnamomeus, Wall. Cat. n. 527 ; et ex DC. Prod. iv. 300 = L. Scurrula.
Citri, *Chatin, Anatom.* 462. t. 85.—Bras.
citricola, *Mart. ex Schult. f. Syst.* vii. 137.—Bras.
clandestinus, *Mart. ex Schult. f. l. c.* 96.—Bras.
clavatus, *Desr. in Lam. Encyc.* iii. 598.—Ins. Mascar.
clavatus, Roxb. Hort. Beng. 87 ; Fl. Ind. ii. 189 = L. Corynitis.
clavatus, Thonn. ex DC. Prod. iv. 303 = Thonningii.
claviceps, *Griseb. Fl. Brit. W. Ind.* 311.—Ind. occ.
clavigerus, Wall. Cat. n. 6873 = longiflorus ?
Cleghornii, *Bedd. in Madr. Journ.* Ser. III. i. (1864) 48.—Ind. or.
clusiaefolius, Willd. ex Eichl. in Mart. Fl. Bras. v. II. 30 = dichotomus.
cneoroides, *Willd. ex Schult. f. Syst.* vii. 162.—Am. austr.
coccineus, Hook. Bot. Misc. i. (1830) t. 58 (non p. 278) = pulcher.
coccineus, *Jack, in Malay. Misc.* i. (1820) n. I. p. 8.—Ind. or. ; Malaya.
coccineus, Reinw. in Blume, Bijdr. 664 = Reinwardtianus.
cochinchinensis, *Lour. Fl. Cochinch.* 195.—Cochinch.

LORANTHUS:—
Colensoi, *Hook. f. in Hook. Ic. Pl.* t. 633.—N. Zel.
concavifolium, Griff. Notul. iv. 615 = L. Scurrula.
concinnus, *Mart. ex Schult. f. Syst.* vii. 150.—Bras.
conduplicatus, *H. B. & K. Nov. Gen. et Sp.* iii. 441.—Venezuela.
confertus, *Mart. ex Schult. f. Syst.* vii. 121.—Bras.
congener, Sieber, ex Schult. f. l. c. 114 = pendulus.
congestus, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyssin.
contractus, Wall. Cat. n. 6864 = pentandrus.
constrictus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 276.—Borneo.
cordatus, *Hoffmgg. ex Schult. f. Syst.* vii. 128.—Bras.
cordifolius, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 222.—Ind. or.
cordifolius, Willd. ex Schult. f. Syst. vii. 162 = cardiphyllus.
coriaceus, *Desr. in Lam. Encyc.* iii. 597.—Malabar.
corymbosus, Desr. l. c. 599 = tetrandrus.
corymbosus, *F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr.* iv. 468.—Peruv.
Corynitis, *Spreng. Syst.* iv. *Cur. Post.* 140.—Ind. or.
crassifolius, *Mart. ex Schult. f. Syst.* vii. 123.—Bras.
crassipes, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 173.—Mexic.
creatophyllus, *Korth. in Verh. Bat. Gen.* xvii. (1839) 282.—Sumatra.
croceus, *E. Mey. in Drège, Zwei Pfl. Docum.* 200, nomen.—Afr. austr.
cubensis, *Griseb. Cat. Pl. Cub.* 119.—Cuba.
cucullaris, *Lam. in Journ. Hist. Nat. Par.* i. (1792) 444. t. 23.—Guian.
cuneatus, *Heyne, in Roth, Nov. Pl. Sp.* 193.—Ind. or.
cuneatus, Wall. Cat. n. 541 = fuscus.
cuneifolius, Reinw. ex Miq. Fl. Ind. Bat. i. 1. 812 = repandus.
cuneifolius, *Ruiz & Pav. Fl. Per.* iii. 46.—Peruv.
Cunninghami, *A. Gray, Bot. U. St. Expl. Exped.* i. 741.—Austral.
cupulatus, *DC. Prod.* iv. 298.—Am. austr.
cupulifer, *H. B. & K. Nov. Gen. et Sp.* iii. 438.—Peruv.
curvatus, *Blume, Bijdr.* 665.—Java.
curviflorus, *Benth. in Hook. Ic. Pl.* xiv. 3. t. 1304.—Abyssin.
cuspidatus, *Mart. ex Schult. f. Syst.* vii. 140.—Bras.
cylindricus, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 213.—Sumatra.
daphnoides, Willd. ex Schult. f. Syst. vii. 158 = microphyllus.
decussatus, *T. Kirk, in Trans. N. Z. Inst.* iii. (1870) 162.—N. Zel.
densiflorus, *Benth. Pl. Hartw.* 62.—Mexic.
Deppeanus, *Cham. & Schlecht. in Linnaea*, v. (1830) 172.—Mexic.
Destructor, *H. B. & K. Nov. Gen. et Sp.* iii. 485.—Ecuador.
dichotomus, *Ruiz & Pav. Fl. Per.* iii. 45.—Am. austr.
dichrous, *Mart. ex Schult. f. Syst.* vii. 122.—Bras.
dictyophlebus, *F. Muell. Rep. Burdek. Exp.* 14.—Austral.
dioicus, Stokes, Bot. Mat. Med. ii. 297 = europaeus.
diplocrater, *Baker, in Journ. Linn. Soc.* xx. (1883) 246.—Madag.
divaricatus, *H. B. & K. Nov. Gen. et Sp.* iii. 433.—Am. austr.
diversifolius, *Benth. Pl. Hartw.* 63.—Mexic.
dodoneaefolius, *DC. Coll. Mém.* vi. t. 9.—Afr. trop.
domingensis, *Desv. in Ham. Prod. Pl. Ind. Occ.* 33.—Ins. S. Doming.
Dregei, *Eckl. & Zeyh. Enum.* 358.—Afr. austr.
elasticus, *Desr. in Lam. Encyc.* iii. 599.—Ind. or.
elatus, *Edgew. in Trans. Linn. Soc.* xx. (1846) 58.—Reg. Himal.
elegans, Cham. & Schlecht. in Linnaea, iii. (1828) 208 = Schlechtendalianus.
elegans, *Mart. ex Schult. f. Syst.* vii. 144.—Bras.
elegans, *Wall. ex DC. Prod.* iv. 304.—Burma.
elegantulus, Steud. Nom. ed. II. ii. 70 = praec.
eleutheropetalus, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 64 = heteranthus.
ellipticus, Korth. in Verh. Batav. Gen. xvii. (1839) 272 (= *Dendrophthoë kramatensis*).—Borneo.
ellipticus, *Ruiz & Pav. Fl. Per.* iii. 47.—Peruv.
emarginatus, *Sw. Prod. Veg. Ind. Occ.* 58.—Ins. S. Doming.

LORANTHUS:—
ensifolius, *Thw. Enum. Pl. Zeyl.* 134.—Zeylan.
epigaeus, *F. Muell. Fragm.* v. 34, in syn.—Austral.
erythrocarpus, *Mart. ex Schult. f. Syst.* vii. 138.—Bras.
erythrostachys, Lam. ex Schult. f. l. c. 103, in obs. = pentapetalus.
Eschscholzianus, *Mart. ex Schult. f. l. c.* 117.—Chili.
eucalyptifolius, *H. B. & K. Nov. Gen. et Sp.* iii. 433.—Venezuela.
eucalyptifolius, Sieber, ex Schult. f. Syst. vii. 163 = celastroides.
eucalyptoides, DC. Prod. iv. 318 = praec.
eugenioides, *H. B. & K. Nov. Gen. et Sp.* iii. 435.—Peruv.
Euphorbiae, Wight, Ic. t. 1063 = elasticus.
europaeus, *Jacq. Enum. Stirp. Vindob.* 55; 230.—Europ. austr.; Oriens.
evenius, Blume, Fl. Jav. Loranth. t. 16 = avenis.
Exocarpi, *Behr, in Linnaea*, xx. (1847) 624.—Austral.
falcatus, Linn. f. Suppl. 211 = longiflorus.
falcifrons, *Mart. ex Schult. f. Syst.* vii. 129.—Bras.
farinaceus, *Griff. Ic. Pl. As.* vi. t. 620.—Ind. or.
farinosus, Desr. in Lam. Encyc. iii. 597 = pentandrus.
fasciculatus, *Blume, Bijdr.* 661.—Java.
ferrugineus, *Roxb. Hort. Beng.* 87; *Fl. Ind. ed. Carey*, ii. 207.—Ind. or.; Malaya.
ferruginosus, Roxb. Fl. Ind. ii. 188 = ferrugineus.
Fieldii, *F. Buch. in Trans. N. Z. Inst.* xvi. (1884) 397.—N. Zel.
filiflorus, *Boj. ex Baker, in Journ. Linn. Soc.* xx. (1883) 246.—Madag.
filifolius, Boj. ex Benth. & Hook. f. Gen. iii. 209 = praec.?
Finlaysonianus, Wall. Cat. n. 6871 = pentandrus.
firmus, Wall. l. c. n. 6874 = Henslowia umbellata.
flagellaris, *Cham. & Schlecht. in Linnaea*, iii. (1828) 213.—Bras.
flavidus, *Hook. f. Fl. N. Zel.* 100. t. 27.—N. Zel.
flavus, Blume, Bijdr. 661 = pentandrus.
flexicaulis, *Mart. ex Schult. f. Syst.* vii. 139.—Bras.
flexistylus, *Miq. in Linnaea*, xviii. (1844) 65.—Guian.
floribundus, Labill. Nov. Holl. Pl. i. 87 = Nuytsia floribunda.
florulentus, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
Fordii, *Hance, in Journ. Bot.* xxiii. (1885) 38.—China.
forficuliflora, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 243 = Faurea forficuliflora.
formosus, *Blume, Bijdr.* 694.—Burma; Malaya.
formosus, Cham. & Schlecht. in Linnaea, iii. (1828) 211 = robustus.
Forsterianus, *Schult. f. Syst.* vii. 114.—Ins. Societ.
Francisianus, Korth. ex Miq. Fl. Ind. Bat. i. 1. 830 = Macrosolen melintangensis.
fulvus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 273.—Borneo.
furcatus, *Mart. ex Schult. f. Syst.* vii. 126.—Bras.
fuscatus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 273.—Sumatra.
fuscoides, Korth. ex Miq. Fl. Ind. Bat. i. 1. 815 = praec.
fuscus, *Blume, Bijdr.* 660.—As. trop.
Gardneri, *Thw. Enum. Pl. Zeyl.* 133.—Zeylan.
Gaudichaudii, *DC. Prod.* iv. 295.—Austral.
gemmiflorus, *Blume, Bijdr.* 665.—Java.
gibbosulus, *A. Rich. Tent. Fl. Abyss.* i. 342.—Abyssin.
glaberrimus, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 170.—Mexic.
glaucocarpus, *Peyr. in Sitzb. Acad. Wien, Math. Nat.* xxxviii. (1860) 571.—Afr. trop.
glaucus, Gill. ex Hook. Bot. Misc. iii. (1833) 357 = Sternbergianus.
glaucus, Pohl, ex DC. Prod. iv. 308 = Pohlii.
glaucus, Ruiz & Pav. Fl. Per. iii. 45 = corymbosus.
glaucus, *Thunb. Prod. Pl. Cap.* 58.—Afr. austr.
globiferus, *A. Rich. Tent. Fl. Abyss.* i. 341.—Abyssin.
globosus, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 187.—Ind. or.; Malaya.
glomeratus, *Mart. ex Schult. f. Syst.* vii. 121.—Bras.
gonocladus, *Baker, in Journ. Linn. Soc.* xx. (1883) 247.—Madag.
goodeniaeflorus, DC. Prod. iv. 306 = cuneatus.
graciliflorus, Wall. ex DC. Prod. iv. 300; et ex Schult. f. Syst. vii. 1785 (Index) = L. Scurrula.
gracilifolius, Roxb. ex Schult. f. l. c. 99 = praec.
Grahami, *Benth. Pl. Hartw.* 62.—Mexic.

LORANTHUS:—
grandibracteus, *F. Muell. Rep. Burdek. Exp.* 14.—Austral.
grandiflorus, *Ruiz & Pav. Fl. Per.* iii. 44.—Peruv.
grandiflorus, *Vell. Fl. Flum.* 138; iii. t. 147.—Bras.
grandifolius, *Mart. ex Schult. f. Syst.* vii. 124.—Bras.
granularis, Willd. ex Schult. f. l. c. 156 = Phoradendron Schottii.
Grewingkii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 106.—Persia.
guadalupensis, *DC. Prod.* iv. 294.—Ins. Guadalup.
guianensis, Klotzsch, in Schomb. Faun. & Fl. Gui. iii. 1161 = cordatus.
Haenkeanus, *Presl, ex Schult. f. Syst.* vii. 113.—Ins. Philipp.
Hartwegi, *Benth. Pl. Hartw.* 62.—Mexic.
heteranthus, *Wall. in DC. Prod.* iv. 306.—Burma; Malaya.
heteromorphus, *A. Rich. Tent. Fl. Abyss.* i. 340.—Abyssin.
heterophyllus, *Ruiz & Pav. Fl. Per.* iii. 48.—Chili.
hexapetalus, Buch.-Ham. ex Schult. f. Syst. vii. 153 = odoratus.
Heyneanus, Schult. f. l. c. 106 = bracteatus.
Heyneanus, Wall. Cat. n. 528 = L. Scurrula.
Heynii, Wall. ex DC. Prod. iv. 300 = L. Scurrula.
hirsutiflorus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 178.—Afr. trop.
Hoffmannseggianus, *Schult. f. Syst.* vii. 113.—Bras.
Hookerianus, *Wight & Arn. Prod.* 381.—Ind. or.
hoyaefolius, *Baker, in Journ. Linn. Soc.* xviii. (1881) 277.—Madag.
hypoglaucus, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 309.—Burma.
ilicifolius, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 259 = Polyosma ilicifolia.
imbricatus, Edgew. in Trans. Linn. Soc. xx. (1846) 59 = longiflorus.
immersus, Benth. Pl. Hartw. 189 = occidentalis.
incanus, *Schum. & Thonn. Beskr. Guin. Pl.* 180.—Afr. austr.
incarnatus, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 213.—Sumatra.
inconspicuus, *Benth. Bot. Voy. Sulph.* 102.—Am. austr.
indicus, Boj. Hort. Maurit. 163 = Bojeri.
indicus, DC. Prod. iv. 305 = signatus.
indicus, Desr. in Lam. Encyc. iii. 601 = longiflorus.
insularum, *A. Gray, Bot. U. St. Expl. Exped.* i. 738.—Ins. Fiji.
intermedius, *Wight, in Calc. Journ. Nat. Hist.* vi. (1846) 361.—Ind. or.
interruptus, *H. B. & K. Nov. Gen. et Sp.* iii. 440.—Mexic.
involucratus, *Roxb. Hort. Beng.* 87; *Fl. Ind.* i. 188.—Reg. Himal.
Jackianus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 279.—Borneo.
Jacquini, DC. Prod. iv. 308 = americanus.
Jacquini, A. Gray, Bot. U. St. Expl. Exped. i. 736 = robustus.
javanicus, *D. Dietr. Syn. Pl.* ii. 1082.—Java.
Junghuhnii, *Molkenb. in Miq. Pl. Jungh.* 113.—Java.
kalahiensis, *Korth. in Verh. Batav. Gen.* xvii. (1839) 271.—Borneo.
Kanneli, *Wight & Arn. Prod.* 387.—Ind. or.
Karwinskyanus, *Schult. f. Syst.* vii. 1641.—Mexic.
Kerberi, *Fourn. in Bull. Soc. Bot. Fr.* xxx. (1883) 185.—Reg. Argent.
Kirkii, *Oliver, in Journ. Linn. Soc.* vii. (1864) 101.—Afr. trop.
Knoblercheri, *Kotschy, in Sitzb. Acad. Wien, Math. Nat.* l. *Abth.* i. (1865) 360.—Afr. trop.
Koenigianus, Agardh, ex Schult. f. Syst. vii. 108 = longiflorus.
Korthalsii, *Molkenb. in Miq. Pl. Jungh.* 110.—Java.
Kraussianus, *Meissn. in Hook. Lond. Journ. Bot.* ii (1843) 539.—Afr. austr.
laetus, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyss.
laevigatus, Wall. ex DC. Prod. iv. 302 = L. Scurrula.
laevigatus, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 1097 = triflorus.
lageniferus, *Wight, Ic.* t. 306.—Ind. or.
lageniflorus, Wight, in Hook. Ic. Pl. tt. 229, 230 = praec.
Lambertianus, *Schult. f. Syst.* vii. 118 (pentapetalus?).—Nepal.
lanceolatus, *Beauv. Fl. Owar.* ii. 8. t. 64.—Afr. trop.

LORANTHUS:—
lanceolatus, Ruiz & Pav. Fl. Per. iii. 47. t. 278 = lancifolius.
lancifolius, *Poir. ex Schult. f. Syst.* vii. 142.—Peruv.
lanosus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 278.—Sumatra.
laurifolius, *H. B. & K. Nov. Gen. et Sp.* iii. 436.—N. Granat.
laurinus, Willd. ex Schult. f. Syst. vii. 162 = praec.
laxiflorus, *Desv. in Ham. Prod. Pl. Ind. Occ.* 33.—Ins. S. Doming.
lenticellatus, *Baker, in Journ. Linn. Soc.* xviii. (1881) 278.—Madag.
lepidobotrys, *Griseb. Fl. Brit. W. Ind.* 311.—Ind. occ.
lepidophyllus, *Walp. Ann.* ii. 730.—Ind. or.
lepidotus, *Blume, Bijdr.* 660.—Java.
lepidotus, Schult. f. Syst. vii. 100 = melanostemon.
leptanthus, Wall. ex DC. Prod. iv. 299 = pulverulentus.
leptolobus, *Benth. in Hook. Niger Fl.* 375.—Afr. trop.
leptostachys, *H. B. & K. Nov. Gen. et Sp.* iii. 440.—N. Granat.
leucosiphon, Griff. Notul. iv. 623 = albidus.
leucostachys, *Molkenb. in Miq. Pl. Jungh.* 114.—Java.
Lichtensteinii, Willd. ex Cham. & Schlecht. in Linnaea, iii. (1828) 209 = oleaefolius.
Liebmanni, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 172.—Mexic.
ligulatus, *Thw. Enum. Pl. Zeyl.* 135.—Zeylan.
ligustrifolius, *Presl, ex Schult. f. Syst.* vii. 135.—Chili; Peruv.
ligustrinus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 219.—Reg. Himal.
ligustrinus, Willd. ex DC. Prod. iv. 314 = ligustrifolius.
Lindenianus, Walp. Ann. i. 363 = Lyndenianus.
linearifolius, *Bert. in 'Merc. Chil.* 1829'; *ex Am. Journ. Sc.* xxiii. (1833) 95.—Chili.
linearifolius, *Hook. in Mitch. Journ. Trop. Austral.* 102.—Austral.
lineatus, Edgew. in Trans. Linn. Soc. xx. (1846) 59 = longiflorus.
linophyllus, *Fenzl, in Enum. Pl. Hueg.* 56.—Austral.
lobeliaeflorus, DC. Prod. iv. 306 = cuneatus.
lonchiphyllus, *Thw. Enum. Pl. Zeyl.* 418.—Zeylan.
longibracteatus, *Desr. in Lam. Encyc.* iii. 599.—Peruv.
longiflorus, *Desr. l. c.* 598.—Malaya; Austral; Ind. or.
longiflorus, Mutis, ex DC. Prod. iv. 312 = Mutisii.
longifolius, Hook. Ic. Pl. t. 880 = pendulus.
longipedunculatus, *Mart. ex Schult. f. Syst.* vii. 141.—Bras.
loniceroides, Koen. ex Schult. f. l. c. 108 = longiflorus.
loniceroides, *Linn. Sp. Pl.* ed. II. 473.—Ind. or.
loxantherus, *DC. Prod.* iv. 316.—Java.
lucarquensis, *H. B. & K. Nov. Gen. et Sp.* iii. 440.—Peruv.
luzonensis, *Presl, ex Schult. f. Syst.* vii. 104.—Ins. Philipp.
Lyndenianus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 578.—Java.
Macklottianus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 264.—Sumatra.
macrantherus, *Hemsl. Biol. Centr. Am. Bot.* iii. 81.—Mexic.
macranthus, *Hook. Ic. Pl.* viii. tt. 743, 744.—Ecuador.
macrophyllus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 280.—Sumatra.
macrophyllus, Zenker, Pl. Ind. 16. t. 16 = loniceroides.
Macrosolen, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 340.—Abyss.
macrostachys, *Korth. in Verh. Batav. Gen.* xvii. (1839) 262.—Borneo.
macrostachys, *Willd. ex Schult. f. Syst.* vii. 162.—Am. austr.
maculatus, *Blume, Bijdr.* 662.—Java.
Magdalenae, *Cham. & Schlecht. in Linnaea*, iii. (1828) 219.—Am. austr.
malelensis, Korth. in Verh. Batav. Gen. xvii. (1839) 268 = repandus.
malifolius, *Presl, ex Schult. f. Syst.* vii. 113.—Ins. Philipp.
Mannii, *Oliver, in Journ. Linn. Soc.* vii. (1864) 101.—Afr. trop.
marginatus, *Desr. in Lam. Encyc.* iii. 596.—Bras.
marginatus, H. B. & K. Nov. Gen. et Sp. iii. 442. t. 299 = orinocensis.
martinicensis, *Presl, ex Schult. f. Syst.* vii. 1640.—Ins. Martinic.
maytenifolius, *A. Gray, Bot. U. St. Expl. Exped.* i. 739. t. 99.—Austral.

LORANTHUS:—
maytenifolius, F. Muell. Rep. Burdek. Exped. 14 = alyxifolius.
medinensis, *Molkenb. in Miq. Pl. Jungh.* 112.—Sumatra.
Melaleucae, Lehm. Pl. Preiss. i. 279 = pendulus.
melanostemon, *Reinw. ex Schult. f. Syst.* vii. 1632.—Java.
melintangensis, *Korth. in Verh. Batav. Gen.* xvii. (1839) 281.—Sumatra.
membranaceus, Willd. ex Schult. f. Syst. vii. 162, 1639 = cupulifer.
memecifolius, Steud. Nom. ed. II. ii. 71 = seq.
memecylifolius, *Wight & Arn. Prod.* 383.—Ind. or.
mexicanus, *Presl, ex Schult. f. Syst.* vii. 129.—Mexic.
Meyeri, *Presl, Bot. Bemerk.* 76.—Afr. austr.
micranthus, *Hook. f. Fl. N. Zel.* 100.—N. Zel.
microcuspis, *Baker, in Journ. Bot.* xx. (1882) 245.—Madag.
microlimbus, *Baker, in Journ. Linn. Soc.* xx. (1883) 246.—Madag.
microphyllus, *H. B. & K. Nov. Gen. et Sp.* iii. 439. t. 300.—Mexic.
microstachys, *Willd. ex Schult. f. Syst.* vii. 162.—Am. bor.
Miqueli, Lehm. Pl. Preiss. i. 280 = pendulus.
miraculosus, Miq. in Lehm. l. c. = pendulus.
Mitchellii, Wall. Cat. n. 6865 = Olax Wightiana?
monophlebius, *Baker, in Journ. Linn. Soc.* xx. (1883) 247.—Madag.
montanus, Macfad. ex Griseb. Fl. Brit. W. Ind. 312 = parvifolius.
montanus, Wight, in Wall. Cat. n. 6866 = cuneatus.
montevidensis, Spreng. Syst. ii. 128 = cuneifolius.
multiflorus, Willd. ex Schult. f. Syst. vii. 162 = suaveolens.
Murrayii, *F. Muell. & Tate, in Trans. R. Soc. S. Austral.* vi. (1883) 109.—Austral.
mutabilis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 61. t. 183.—Chili.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iii. 433.—N. Granat.
myrtifolius, *A. Cunn. ex Benth. Fl. Austral.* iii. 390.—Austral.
natalitius, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1848) 539.—Afr. austr.
neelgherrensis, *Wight & Arn. Prod.* 382.—Ind. or.
nigrans, *Hance, in Journ. Bot.* xix. (1881) 209.—China.
nigritanus, *Hook. f. ex Benth. in Hook. Niger Fl.* 376.—Afr. trop.
nitens, *Mart. ex Schult. f. Syst.* vii. 150.—Bras.
nitidus, *H. B. & K. Nov. Gen. et Sp.* iii. 437.—Ecuador.
nitidus, Willd. ex Eichl. in Mart. Fl. Bras. v. II. 27 = cupulifer.
nodiflorus, *Thw. Enum. Pl. Zeyl.* 134.—Zeylan.
nodosus, *Desr. in Lam. Encyc.* iii. 601.—Peruv.
notothixoides, *Hance, in Journ. Bot.* xxi. (1883) 356.—China.
nutans, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral. 158 = L. Quandang.
nutans, *Spreng. Syst.* ii. 128.—Bras.
oblongifolius, *E. Mey. ex A. Rich. Tent. Fl. Abyss.* i. 342.—Abyss.
oblongifolius, E. Mey. in Drège, Zwei Pfl. Docum. 200 = Meyeri.
oblongo-ellipticus, *Pohl, ex DC. Prod.* iv. 315.—Bras.
oblongus, Schott, ex Pohl, l. c. 287 = Phoradendron rubrum.
obmunitus, *Spreng. Syst.* iv. *Cur. Post.* 139.—Bras.
obovatus, *Benth. Bot. Voy. Sulph.* 103.—Ecuador.
obovatus, Blume, Bijdr. 663 = L. Scurrula.
obovatus, *Griff. Notul.* iv. 622.—Malacca.
obtectus, Wall. in DC. Prod. iv. 299 = L. Scurrula.
obtusatus, *Wall. Cat.* n. 566.—Ind. or.
obtusatus, Zipp. ex Span. in Linnaea, xv. (1841) 208 = rhopalocarpus?
obvallatus, Schult. f. Syst. vii. 162 = puracensis.
occidentalis, *Aubl. Pl. Gui.* 310 = pauciflorus.
occidentalis, *Linn. Syst.* ed. X. 988.—Am. austr.
odontocalyx, *F. Muell. ex Benth. Fl. Austral.* iii. 391.—Austral.
odoratus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 215.—Reg. Himal.
odoriferus, *Vell. Fl. Flum.* 138; iii. t. 146.—Bras.
Oerstedii, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 171.—Nicaragua.

LORANTHUS:—
oleaefolius, *Cham. & Schlecht. in Linnaea*, iii. (1828) 209.—Afr. austr.
oleaefolius, Wall. ex DC. Prod. iv. 297 = globosus.
oleoides, DC. l. c. = globosus.
Oortianus, Korth. in Verh. Batav. Gen. xvii. (1839) 269 = ferrugineus.
orbicularis, *H. B. & K. Nov. Gen. et Sp.* iii. 434.—Ecuador.
oreophilus, *Oliver, ex Hook. f. in Journ. Linn. Soc.* vii. (1864) 196.—Afr. trop.
orinocensis, *Spreng. Syst.* i. 129.—Venezuela.
ovalifolius, *Ruiz & Pav. Fl. Per.* iii. 50.—Peruv.
ovalis, *E. Mey. in Drège, Zwei Pfl. Docum.* 200.—Afr. austr.
ovata, *Pohl, ex DC. Prod.* iv. 315.—Bras.
oxyphyllus, *Pohl, l. c.* 314.—Bras.
pachyphyllus, *Baker, in Journ. Bot.* xx. (1882) 245.—Madag.
pallens, Wall. in DC. Prod. iv. 297 = ampullaceus.
paniculatus, *H. B. & K. Nov. Gen. et Sp.* iii. 442.—Venezuela.
Parkeri, *Baker, in Journ. Linn. Soc.* xx. (1883) 245.—Madag.
parviflorus, Desr. in Lam. Encyc. iii. 595 = uniflorus.
parvifolius, *Sw. Prod. Veg. Ind. Occ.* 58.—Jamaica.
patens, *Mart. ex Schult. f. Syst.* vii. 142.—Bras.
Patrisii, *DC. Prod.* iv. 288.—Guian.
patulus, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 214.—Sumatra.
pauciflorus, *Blanco, Fl. Filip.* ed. I. 235.—Ins. Philipp.
pauciflorus, *Sw. Prod. Veg. Ind. Occ.* 58.—Jamaica.
pedunculatus, *Jacq. Enum. Pl. Carib.* 18.—N. Granat.
pedunculatus, Spreng. ex Steud. Nom. ed. II. ii. 71 = guadalupensis.
pendulus, *Sieber, ex DC. Prod.* iv. 295.—Austral.
Pentagonia, *DC. Coll. Mém.* vi. t. 8.—Afr. trop.
pentandrus, Heyne, ex Schult. f. Syst. vii. 112 = longiflorus.
pentandrus, *Linn. Mant.* i. 63.—Ind. or.; Malaya.
pentandrus, Reinw. ex Schult. f. Syst. vii. 1632 = melanostemon.
pentapetalus, *Roxb. Hort. Beng.* 87; *Fl. Ind.* i. 190.—Ind. or.; Malaya.
pereskiaefolius, Willd. ex Schult. f. Syst. vii. 48 = L. Destructor.
Perrottetii, *DC. Prod.* iv. 292.—Guiana.
philippensis, *Cham. & Schlecht. in Linnaea*, iii. (1828) 204.—Ins. Philipp.
phillyreoides, *H. B. & K. Nov. Gen. et Sp.* iii. 439.—Venezuela.
piperoides, H. B. & K. l. c. 443 = Phoradendron Schottii.
platyphyllus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 341.—Abyss.
Plumieri, *Cham. & Schlecht. in Linnaea*, iii. (1828) 211.—Am. austr.
podopterus, *Cham. & Schlecht. l. c.* 218.—Bras.
Poeppigii, *DC. Prod.* iv. 308.—Chili.
Pohlii, *DC. l. c.*—Bras.
polyanthos, *Mart. ex Schult. f. Syst.* vii. 145.—Bras.
polycarpus, Wall. Cat. n. 540 = pentapetalus.
polychroa, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 241.—N. Zel.
polycryptus, *Didr. in Kjoeb. Vidensk. Meddel.* (1854) 194.—Afr. trop.
polyrrhizos, *Mart. ex Schult. f. Syst.* vii. 139.—Bras.
polystachys, *Ruiz & Pav. Fl. Per.* iii. 50.—Peruv.
polystachyus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 217 = Wallichianus.
portoricensis, *DC. Prod.* iv. 293.—Ins. Porto Rico.
praelongus, *Blume, Bijdr.* 664.—Java.
Preissii, Miq. in Lehm. Pl. Preiss. i. 280 = linifolius.
prunifolius, *E. Mey. in Drège, Zwei Pfl. Docum.* 200.—Afr. austr.
Pseudo-globosus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 279.—Sumatra.
Pseudoperfoliatus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 578.—Java.
psilobotrys, *DC. Prod.* iv. 293.—Ins. S. Doming.
pterygopus, *Mart. ex Schult. f. Syst.* vii. 155.—Bras.
pulcher, Blume, ex Schult. f. l. c. 1649 = loxantherus.
pulcher, *DC. Prod.* iv. 295.—Ins. Penang.
pulverulentus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 221.—Ind. or.; Burma.
pumilus, *Miers, ex Schult. f. Syst.* vii. 1649.—Chili.
punctatus, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 323.—N. Zel.

LORANTHUS:—
punctatus, *Ruiz & Pav. Fl. Per.* iii. 47.—Peruv.
puniceus, *Wall. Cat.* n. 522.—Ins. Penang.
puracensis, *H. B. & K. Nov. Gen. et Sp.* iii. 437.—Ecuador.
purpurascens, Herb. Monac. ex Schult. f. Syst. vii. 110, in obs. = venosus.
pycnanthus, *Benth. Pl. Hartw.* 189.—N. Granat.
pyrifolius, *H. B. & K. Nov. Gen. et Sp.* iii. 441.—N. Granat.
pyrifolius, Wight, ex Wall. Cat. n. 6869 = L. Scurrula.
pyrifolius, Willd. ex Schult. f. Syst. vii. 1647 = leptostachys.
quadrangularis, H. B. & K. Nov. Gen. et Sp. iii. 444 = Phoradendron rubrum?
Quandang, *Lindl. in Mitch. Three Exped.* ii. 69.—Austral.
quercicola, *Schlecht. & Cham. in Linnaea*, v. (1830) 173.—Mexic.
quinquenervis, *Hochst. in Flora*, xxvii. (1844) 432.—Afr. austr.
racemiferus, Wall. in DC. Prod. iv. 296 = coccineus.
radicans, *Cham. & Schlecht. in Linnaea*, iii. (1828) 217.—Bras.
ramiflorus, *Moç. & Sesse, ex DC. Prod.* iv. 308.—Mexic.
recurvus, *Wall. ex DC. l. c.* 299.—Ind. or.
regularis, Steud. ex A. Rich. Tent. Fl. Abyss. i. 339 = rufescens.
Reinwardtianus, *Blume, ex Schult. f. Syst.* vii. 105.—Java.
Reinwardtii, Schult. f. l. c. 1612 = praec.
repandus, *Blume, Fl. Jav. Loranth.* 22. t. 8.—Java.
retroflexus, *Ruiz & Pav. Fl. Per.* iii. 49.—Peruv.
retusus, Cham. & Schlecht. in Linnaea, iii. (1828) 215 = Sellowianus.
retusus, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 212.—Malaya.
rhopalocarpus, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 62 = fuscus.
rhynchanthus, *Benth. Bot. Voy. Sulph.* 102.—Mexic.
rigidulus, Wall. ex G. Don, Gen. Syst. iii. 422 = L. Scurrula.
rigidus, Wall. ex DC. Prod. iv. 298 = pentandrus.
robustus, *Mart. ex Schult. f. Syst.* vii. 125.—Bras.
roseus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 177.—Afr. trop.
rotundatus, *Wall. ex Schult. f. Syst.* vii. 118.—Am. austr.
rotundifolius, *A. St. Hil. in Mém. Mus. Par.* ix. (1822) 328.—Bras.
rotundifolius, Willd. ex Schult. f. Syst. vii. 154 = alveolatus.
Roxburghianus, Blume, ex Schult. f. l. c 1730 = L. Scurrula.
rubens, *Mart. ex Schult. f. l. c.* 144.—Bras.
rubro-viridis, *Baker, in Journ. Linn. Soc.* xx. (1883) 245.—Madag.
rubroviridis, *Oliver, in Hook. Ic. Pl.* t. 1464.—Afr. trop.
rufescens, *DC. Prod.* iv. 303.—Afr. trop.
ruficaulis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 61. t. 183.—Bras.
rufidulus, Wall. in DC. Prod. iv. 302 = L. Scurrula.
rufus, *Mart. ex Schult. f. Syst.* vii. 130.—Bras.
rugulosus, *Heyne, in Roth, Nov. Pl. Sp.* 194.—Ind. or.
sakoembangensis, *Korth. in Verh. Batav. Gen.* xvii. (1839) 285.—Borneo.
salicifolius, *Cham. & Schlecht. in Linnaea*, iii. (1828) 216.—Bras.
salignus, Willd. ex Schult. f. Syst. vii. 155 = Phoradendron angustifolium.
Sampsoni, *Hance, in Journ. Bot.* ix. (1871) 133.—China.
sanguineus, *F. Muell. Fragm.* i. 177.—Austral.
sarcophyllus, *Wall. Cat.* n. 6863.—Ind. or.
sarmentosus, *Ruiz & Pav. Fl. Per.* iii. 49.—Peruv.
Schiedeanus, *Cham. & Schlecht. in Linnaea*, v. (1830) 172.—Mexic.
Schimperi, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 28, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 342.—Abyss.
Schlechtendalianus, *Schult. f. Syst.* vii. 1635.—Afr. austr.
Schultesii, Blume, ex DC. Prod. iv. 300 = ferrugineus.
sclerophyllus, *Thw. Enum. Pl. Zeyl.* 135.—Zeylan.
scoparius, Miq. in Lehm. Pl. Preiss. i. 280 = linophyllus.
Scurrula, *Linn. Sp. Pl.* ed. II. 472.—Ind. or.; Malaya.
sechellensis, *Baker, Fl. Maurit.* 135.—Ins. Seychelles.

LORANTHUS:—
secundus, *Benth. Pl. Hartw.* 189.—N. Granat.
Sellowianus, *DC. Prod.* iv. 312.—Bras.
serrulatus, *Roxb. ex Steud. Nom.* ed. I. 494.—Ind. or.
sessilifolius, *Beauv. Fl. Owar.* ii. 8. t. 23.—Afr. trop.
sessilis, *Jacq. Enum. Pl. Carib.* 18.—N. Granat.
setigerus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 265.—Borneo.
siamensis, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 62.—Siam.
signatus, *F. Muell. ex Benth. Fl. Austral.* iii. 392.—Austral.; Timor.
Smythii, *Schomb. in Proc. Bot. Soc. Lond.* i. (1839) 47.—Guiana.
speciosus, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 473 = oleaefolius.
speciosus, *Pohl, ex DC. Prod.* iv. 309.—Bras.
speciosus, Wall. ex DC. l. c. 295 = pulcher.
sphaerocarpus, Blume, Bijdr. 661 = globosus.
sphaeroideus, DC. Prod. iv. 390 = seq.
sphenoideus, *Blume, Fl. Jav. Loranth.* 23. t. 4.—Java.
spicatus, Herb. Banks, ex Griseb. Fl. Brit. W. Ind. 313 = occidentalis.
spicatus, *Jacq. Enum. Pl. Carib.* 18; *Linn. Mant.* i. 63.—N. Granat.
spicatus, Schreb. ex Schult. f. Syst. vii. 152 = emarginatus.
spicifer, *Villar, in Blanco, Fl. Filipp.* ed. III. *Nov. App.* 183.—Ins. Philipp.
spirostylis, *DC. Prod.* iv. 315.—Am. austr.
staphylinus, *Mart. ex Schult. f. Syst.* vii. 140.—Bras.
Stelis, Forst. f. Prod. 25 = Forsterianus.
Stelis, *Linn. Sp. Pl.* ed. II. 331.—Venezuela.
stenopetalus, *Oliver, in Journ. Linn. Soc.* xv. (1876) 99.—Ins. Celebes.
Sternbergianus, *Schult. f. Syst.* vii. 116.—Chili.
suaveolens, *H. B. & K. Nov. Gen. et Sp.* iii. 435.—Peruv.
subcampestris, *Mart. ex Schult. f. Syst.* vii. 151.—Bras.
subfalcatus, Hook. in Mitch. Journ. Trop. Austral. 224 = L. Exocarpi.
subglobosus, Wall. Cat. n. 538 = globosus.
suborbicularis, *Thw. Enum. Pl. Zeyl.* 134.—Zeylan.
subrotundus, *Pohl, in DC. Prod.* iv. 292.—Bras.
subumbellatus, *Blume, Bijdr.* 662.—Java.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 63.—Guian.
syringaefolius, *Mart. ex Schult. f. Syst.* vii. 142.—Bras.
Tagua, *H. B. & K. Nov. Gen. et Sp.* iii. 436.—N. Granat.
tehuacanensis, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 171.—Mexic.
tenuiflorus, *Hook. f. Fl. Nov. Zel.* 100.—N. Zel.
terniflorus, *Willd. ex Schult. f. Syst.* vii. 161.—Venezuela.
terrestris, Heyne, ex Wall. Cat. n. 509 = Wallichianus.
tetragonus, *Blume, Bijdr.* 663.—Java.
tetragonus, Willd. ex Schult. f. Syst. vii. 131 = paniculatus.
tetrandrus, Heyne, ex Schult. f. l. c. 100 = Wallichianus.
tetrandrus, *Ruiz & Pav. Fl. Per.* iii. 46. t. 275.—Chili.
tetrapetalus, *Linn. f. Suppl.* 211.—N. Zel.
tetraqueter, *Mart. ex Schult. f. Syst.* vii. 140.—Bras.
Theobromae, *Willd. ex Schult. f. l. c.* 132.—Bras.
Thonningii, *Schum. & Thonn. Beskr. Guin. Pl.* 179.—Afr. trop.
thyrsiflorus, *Cham. & Schlecht. in Linnaea*, iii. (1828) 214.—Bras.
tomentosus, Blanco, Fl. Filip. ed. II. 164 = pauciflorus.
tomentosus, *Heyne, ex Roth, Nov. Pl. Sp.* 191.—Ind. or.
tomentosus, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 184 = formosus.
tomentosus, Wight, Ic. t. 378 = bracteatus.
torulosus, H. B. & K. Nov. Gen. et Sp. iii. 443 = Viscum torulosum.
trianthus, *Korth. in Verh. Batav. Gen.* xvii. (1839) 265.—Sumatra.
triflorus, *Span. in Linnaea*, xv. (1841) 208.—Ins. Timor.
trigonus, *Wight & Arn. Prod.* 386.—Ind. or.
tristis, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 579.—Java.

LORANTHUS:—
tubulosus, Willd. ex Schult. f. Syst. vii. 162 = corymbosus.
turbinatus, *DC. Prod.* iv. 305.—Ind. or.
umbellatus, Heyne, in Roth, Nov. Pl. Sp. 192 = loniceroides.
umbellatus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 220 = umbellifer.
umbellifer, Brand. For. Fl. 397 = elatus.
umbellifer, *Schult. f. Syst.* vii. 97.—Reg. Himal.
undulatus, *E. Mey. in Drège, Zwei Pfl. Docum.* 200.—Afr. austr.
uniflorus, *Jacq. Enum. Pl. Carib.* 18; *Linn. Sp. Pl.* ed. II. 331.—Ins. S. Doming.
uniflorus, Spreng. ex Steud. Nom. ed. II. ii. 73 = psilobotrys.
uraguensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 358.—Uruguay.
usuiensis, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 80.—Afr. trop.
Utui, Molina, Sagg. Chile, ed. II. 139 = tetrandrus.
valdivianus, Miq. in Linnaea, xxv. (1852) 653 = heterophyllus.
vegetus, DC. Prod. iv. 313 = venetus.
Vellozianus, Walp. Rep. ii. 443 = dichrous.
venetus, Bert. ex Hemsl. Bot. Voy. Challenger, i. III. 54 = Berteroi.
venetus, *H. B. & K. Nov. Gen. et Sp.* iii. 434.—Mexic.
venosus, *Blume, Bijdr.* 668.—Java.
verticillatus, *Benth. & Hook. f. Gen.* iii. 208.—N. Guinea.
verticillatus, Hoffmgg. ex Schult. f. Syst. vii. 113 = Hoffmanseggianus.
verticillatus, *Ruiz & Pav. Fl. Per.* iii. 47.—Chili.
vestitus, *Miq. in Linnaea*, xviii. (1844) 62.—Guian.
vestitus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 218.—Reg. Himal.
virgatus, *Mart. ex Schult. f. Syst.* vii. 132.—Bras.
viridiflorus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 219 = globosus.
viscifolius, H. B. & K. Nov. Gen. et Sp. iii. 443 = Phoradendron rubrum.
viscifolius, Wight, in Wall. Cat. n. 6868 = cuneatus.
viscoideus, *Poepp. ex Eichl. in Mart. Fl. Bras.* v. II. 49.—Bras.
vitellinus, F. Muell. Rep. Burdek. Exped. 12 = longiflorus.
vitiensis, *Seem. Fl. Vit.* 120.—Ins. Fiji.
vulgaris, *Vell. Fl. Flum.* 137; iii. t. 145.—Bras.
Wallichianus, *Schult. f. Syst.* vii. 100.—Ind. or.
Wightianus, Wall. Cat. n. 6872 = longiflorus.
Yadoriki, *Siebold. ex Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 229.—Japon.
Zeyheri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 576.—Afr. austr.

LORANTUS, Bert. in Merc. Chil. (1829); ex Am. Journ. Sc. xxiii. (1833) 95 = praec.

LOREIA, Rafin. Fl. Tellur. iii. 82 (1836) = **Campanula**, Tourn.
baldensis, Rafin. l. c. = C. ramosissima.

LORENTEA, Lag. Gen. et Sp. Nov. (1816) = **Pectis**, Linn. (Compos.).
affinis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 240 = P. oligocephala.
auricularis, DC. Prod. v. 102 = P. auricularis.
brevipedunculata, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 428 = P. apodocephala.
canescens, DC. Prod. v. 102 = P. canescens.
capillipes, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 70 = P. capillipes.
congesta, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 241 = P. congesta.
cryptocephala, Sch. Bip. in Seem. Bot. Voy. Herald, 309 = seq.
cryptopetala, DC. Prod. v. 101 = P. cryptocephala.
cubensis, A. Rich. Fl. Cub. Fanerog. ii. 86 = P. cubensis.
decumbens, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 241 = P. decumbens.
Haenkeana, DC. Prod. v. 102 = P. Haenkeana.
humifusa, Less. in Linnaea, vi. (1831) 719 = P. humifusa.
Liebmanni, Sch. Bip. ex Hemsl. Biol. Cent. Am. Bot. ii. 226 = P. Liebmanni.
linearis, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 174.—Ins. Galap.

LORENTEA :—
multiflosculosa, DC. Prod. v. 102 = P. multiflosculosa.
oligocephala, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 239 = P. oligocephala.
polycephala, Gardn. l. c. 240 = P. elongata.
prostrata, Lag. Gen. et Sp. Nov. 28 = P. humifusa.
ramosissima, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 240 = P. oligocephala.
satureioides, Less. in Linnaea, v. (1830) 135 ; vi. (1831) 718 = P. satureioides.
sessiliflora, Less. l. c. vi. (1831) 720 = P. sessiliflora.
subsquarrosa, Hook. f. in Trans. Linn. Soc. xx. (1846) 206 = P. subsquarrosa.
tenuifolia, DC. Prod. v. 103 = P. tenuifolia.

LORENTEA, Orteg. Hort. Matr. Dec. 41 (1797) = **Sanvitalia**, Gualt. (Compos.).
atropurpurea, Orteg. l. c. 42. t. 5 = S. procumbens.

LORENTIA, Sweet, Hort. Brit. ed. III. 336 (1839) = praec.

LORENTZIA, Griseb. in Goett. Abh. xix. (1874) 182 = **Pascalia**, Orteg. (Compos.).
pascaloides, Griseb. l. c. = P. glauca.

LORENTZIA, Hieron. in Bol. Acad. Nac. Córdova, iv. (1881) 96. *EUPHORBIACEAE.*
cordobensis, *Hieron. l. c.* 97.—Reg. Argent.
saltenia, *Hieron. l. c.* 99.—Reg. Argent.

LORENZANA, Walp. Ann. iii. 835 (1853) = seq.

LORENZANEA, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 69 = **Meliosma**, Blume (Sabiac.).
dentata, Liebm. l. c.—Mexic.
glabrata, Liebm. l. c. 71.—Costa Rica.
grandifolia, Liebm. l. c. 73.—Mexic.
Ira, Liebm. l. c. 71.—Costa Rica.
vernicosa, Liebm. l. c. 72 = M. vernicosa.

LORETIA, Duval-Jouve, in Rev. Sc. Nat. Sér. II. ii. [1880] 38 = **Festuca**, Linn. (Gramin.).
geniculata, Duval-Jouve, l. c. 42 = F. clavata.
incrassata, Duval-Jouve, l. c. 41 = F. incrassata.
ligustica, Duval-Jouve, l. c. 43 = F. ligustica.
setacea, Duval-Jouve, l. c. 40 = F. Thomasiana.

LOREYA, DC. Prod. iii. 178 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 768.
HETERONEURON, Hook. f. in Benth. & Hook. f. Gen. i. 768 (1867).
OXISMA, Rafin. Sylva Tellur. 94 (1838).
acutifolia, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 142.—Bras.
arborescens, *DC. Prod.* iii. 179.—Guian.
fasiculiflora, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 110 = Henriettella sessilifolia.
flavescens, Naud. l. c. = Henriettella flavescens.
grandiflora, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 143 = Spruceana.
mespiloides, *Miq. in Linnaea*, xviii. (1844) 619.—Guian.
nigricans, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 142.—Bras.
ovata, *O. Berg, ex Triana, l. c.*—Bras.
Spruceana, *Triana, l. c.* 143.—Bras.
trinitensis, Crueg. in Linnaea, xx. (1847) 108 = Henriettella sessilifolia.
umbelluliflora, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 143 = Henriettella umbelluliflora.

LORICARIA, Wedd. Chlor. And. i. 165. t. 27* A, B, C (1856) = **Tafalla**, D. Don (Compos.).
anceps, Sch. Bip. in Bonplandia, viii. (1860) 259.—Peruv.
complanata, Wedd. Chlor. And. i. 167.—N. Granat.
ferruginea, Wedd. l. c. 166 = T. ferruginea.
graveolens, Wedd. l. c. 167.—N. Granat.
stenophylla, Wedd. l. c. 165.—N. Granat.
thyoides, Sch. Bip. in Bonplandia, viii. (1860) 258 = T. thyoides.

LOROGLOSSUM, Rich. in Mém. Mus. Par. iv. (1818) 47 = **Orchis**, Linn. (Orchid.).
anthropophorum, Rich. l. c. 54 = Aceras anthropophora.
brachyglotte, Rich. l. c. = Aceras anthropophora.
hircinum, Rich. l. c. = O. hircina.

LOROPETALUM, R. Br. in Abel, Narr. Journ. Chin. App. 375 (1818). *HAMAMELIDEAE*, Benth. & Hook. f. i. 668.
TETRATHYRIUM, Benth. Fl. Hongk. 132 (1861).
chinense, *Oliver, in Trans. Linn. Soc.* xxiii. (1862) 459.—China ; Reg. Himal.
subcordatum, *Oliver, in Hook. Ic. Pl.* t. 1417.—Hongkong.

LOROSTELMA, Fourn. in Mart. Fl. Bras. vi. IV. 220 (1885). *ASCLEPIADEAE.*
struthianthus, *Fourn. l. c.*—Bras.

LOTEA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384 = **Lotus**, Tourn. (Legumin.).
ornithopodioides, Medic. l. c. = Lotus ornithopodioides.

LOTOASTRAGALUS, Heist. Syst. 9 (1748) (Quid ?).

LOTONONIS, Eckl. & Zeyh. Enum. 176 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 476.
AMPHINOMIA, DC. Prod. ii. 522 (1825).
AULACINTHUS, E. Mey. Comm. Pl. Afr. Austr. 156 (1835).
CAPNITIS, E. Mey. Comm. Pl. Afr. Austr. 81 (1835).
KREBSIA, Eckl. & Zeyh. Enum. 179 (1836).
LEOBORDEA, Delile, in Laborde, Voy. Arab. Pétr. 86 (1833).
LEPTIDIUM, Presl, Bot. Bemerk. 49 (1844) ; in Abh. Boehm. Ges. V. iii. (1845) 479.
LEPTIS, E. Meyer, ex Eckl. & Zeyh. Enum. 174 (1836).
LIPOZYGIS, E. Mey. Comm. Pl. Afr. Austr. 76 (1835).
POLYLOBIUM, Eckl. & Zeyh. Enum. 180 (1836).
TELINA, E. Mey. Comm. Pl. Afr. Austr. 67 (1835).
abyssinica, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat.* li. *Abth.* II. (1865) 263.—Abyssin.
acuminata, *Eckl. & Zeyh. Enum.* 176.—Afr. austr.
acutiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 604.—Afr. austr.
affinis, Eckl. & Zeyh. Enum. 178 = pungens.
angolensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 6.—Afr. trop.
angustifolia, *Steud. Nom.* ed. II. ii. 73.—Afr. austr.
anthylloides, *Harv. in Harv. & Sond. Fl. Cap.* ii. 59.—Afr. austr.
argentea, *Benth. ex Walp. Rep.* v. 459.—Afr. austr.
argentea, *Eckl. & Zeyh. Enum.* 176.—Afr. austr.
azurea, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 600.—Afr. austr.
Bainesii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 6.—Afr. trop.
brachyloba, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 608.—Afr. austr.
bracteata, *Benth. l. c.* 600.—Afr. austr.
Burchellii, *Benth. l. c.* 612.—Afr. austr.
calycina, *Benth. l. c.* 611.—Afr. austr.
Capnitis, *Benth. ex Walp. Rep.* v. 459.—Afr. austr.
carinalis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 60.—Afr. austr.
carinata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 609.—Afr. austr.
carnosa, *Benth. l. c.* 598.—Afr. austr.
clandestina, *Benth. l. c.* 607.—Afr. austr.
corymbosa, *Benth. l. c.* 606.—Afr. austr.
crumaniana, *Burch. ex Benth. l. c.* 612.—Afr. austr.
cytisoides, *Benth. l. c.* 598.—Afr. austr.
debilis, *Benth. l. c.* 604.—Afr. austr.
decidua, Eckl. & Zeyh. Enum. 178 = pungens.
densa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 52.—Afr. austr.
depressa, *Eckl. & Zeyh. Enum.* 178.—Afr. austr.
dichiloides, *Sond. in Linnaea*, xxiii. (1850) 28.—Afr. austr.
dichotoma, Boiss. Fl. Orient. ii. 30 = L. Leobordea.
diffusa, Eckl. & Zeyh. Enum. 177 = Crotalaria humilis.
digitata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 52.—Afr. austr.
divaricata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 599.—Afr. austr.
divaricata, Harv. in Harv. & Sond. Fl. Cap. ii. 613 = calycina.
diversifolia, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 611 = laxa.
eriantha, *Benth. l. c.* 605.—Afr. austr.
excisa, *Steud. Nom.* ed. II. ii. 73.—Afr. austr.
falcata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 608.—Afr. austr.
genistoides, *Benth. l. c.* 607.—As. Min.

LOTONONIS :—
genuflexa, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 598.—Afr. austr.
gracilis, *Benth. l. c.* 597.—Afr. austr.
heterophylla, Eckl. & Zeyh. Enum. 177 = prostrata.
humifusa, *Burch. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 609.—Afr. austr.
involucrata, *Benth. l. c.* 602.—Afr. austr.
lanceolata, *Benth. l. c.* 606.—Afr. austr.
laxa, *Eckl. & Zeyh. Enum.* 177.—Afr. austr.
lenticula, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 611.—Afr. austr.
Leobordea, *Benth. l. c.* 607.—Afr. bor. ; Arab.
lupinifolia, *Benth. l. c.* 607.—Reg. Mediterr.
macrocarpa, *Eckl. & Zeyh. Enum.* 176.—Afr. austr.
maroccana, *Ball, in Journ. Bot.* xi. (1873) 302.—Marocc.
micrantha, Eckl. & Zeyh. Enum. 178 = pumila.
micrantha, *Harv. in Harv. & Sond. Fl. Cap.* ii. 58.—Afr. austr.
microphylla, *Harv. l. c.* 65.—Afr. austr.
mollis, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 609.—Afr. austr.
monophylla, *Harv. Thes. Cap.* i. 39. t. 63.—Afr. austr.
Oxyptera, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 605.—Afr. austr.
pallens, *Benth. l. c.*—Afr. austr.
peduncularis, *Benth. l. c.* 602.—Afr. austr.
pentaphylla, *Benth. l. c.* 605.—Afr. austr.
perplexa, *Eckl. & Zeyh. Enum.* 177, ex parte ; *Benth. l. c.* 604.—Afr. austr.
perplexa, Eckl. & Zeyh. l. c. = Crotalaria humilis.
persica, *Boiss. Fl. Orient.* ii. 30.—Persia ; Arab.
platycarpus, Viv. Pl. Aegypt. Dec. iv. 14 = L. Leobordea.
polycephala, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 605.—Afr. austr.
porrecta, *Benth. l. c.* 606.—Afr. austr.
prostrata, *Benth. l. c.* 600.—Afr. austr.
pumila, *Eckl. & Zeyh. Enum.* 178.—Afr. austr.
pungens, *Eckl. & Zeyh. l. c.* 177.—Afr. austr.
quinata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 608.—Afr. austr.
quinata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 63.—Afr. austr.
Radula, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 609.—Afr. austr.
rigida, *Benth. l. c.* 597.—Afr. austr.
rostrata, Benth. l. c. 604 = micrantha.
secunda, Benth. l. c. 603 = angustifolia.
sericea, *Benth. l. c.* 607.—Afr. austr.
sericophylla, *Benth. l. c.* 599.—Afr. austr.
sessilifolia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 66.—Afr. austr.
sphaerocarpa, Boiss. Fl. Orient. ii. 30, 1130 = persica.
strigosa, Pappe, ex Harv. & Sond. Fl. Cap. ii. 57 = perplexa.
tenella, *Eckl. & Zeyh. Enum.* 178.—Afr. austr.
tenuis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 5.—Afr. trop.
trichopoda, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 603.—Afr. austr.
umbellata, *Benth. l. c.* 602.—Afr. austr.
varia, *Steud. Nom.* ed. II. ii. 73.—Afr. austr.
versicolor, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 610.—Afr. austr.
vexillata, Eckl. & Zeyh. Enum. 176 = prostrata.
viborgioides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 597.—Afr. austr.
villosa, *Steud. Nom.* ed. II. ii. 73.—Afr. austr.
Wrightii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 594.—Afr. austr.

LOTOPHYLLUS, Link, Handb. ii. 156 (1831) = **Argyrolobium**, Eckl. & Zeyh. (1835) (Legumin.).
argenteus, Link, l. c. = A. Linnaeanum.

LOTOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 87 = **Lotus**, Tourn. (Legumin.).

LOTULUS, Rafin. in Am. Monthly Mag. (1818) 268 = seq.

LOTUS, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 490.
ANDACA, Rafin. Fl. Tellur. iii. 82 (1836).
HEINEKENIA, Webb, ex Benth. & Hook. f. Gen. i. 491 (1865).

LOTUS:—
KROKERIA, Moench, Meth. 143 (1794).
LOTEA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384.
LOTULUS, Rafin. in Am. Monthly Mag. (1818) 268.
ORTHOLOTUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 363.
PEDROSIA, Lowe, in Hook. Kew Journ. viii. (1856) 292.
SCANDALIDA, Adans. Fam. ii. 326 (1763).
TETRAGONOLOBUS, Scop. Fl. Carn. ed. II. ii. 87 (1772).
acutus, Wald. & Kit. ex Steud. Nom. ed. I. 495 (Quid?).—Hungar.
aduncus, *Nym. Syll.* 298.—Macedon.
aegeus, *Boiss. Fl. Orient.* ii. 167.—Thracia.
aegyptiacus, Moench, Meth. 151, in syn. = arabicus.
affinis, Bess. Cat. Hort. Crem. 81?; ex DC. Prod. ii. 208 = Dorycnium hirsutum.
albidus, Lodd. Bot. Cab. t. 1063 = australis.
albus, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 202. —Turc.
aleppicus, Boiss. Diagn. Ser. I. ix. 33 = L. Gebelia.
Allionii, Desv. Journ. Bot. iii. (1814) 77; Poir. Encyc. Suppl. iii. 506 = cytisoides.
alopecuroides, Burm. f. Fl. Cap. Prod. 23 = Indigofera coriacea.
alpinus, Schleich. ex Steud. Nom. ed. I. 495; Schur, Enum. Pl. Transs. 160 = corniculatus.
ambiguus, Bess. ex Spreng. Syst. iii. 282 = corniculatus.
americanus, Bisch. Del. Sem. Hort. Heidelb. (1839).—Cf. Linnaea, xiv. (1839) Litt. 132 = Hosackia Purshiana.
americanus, Vell. Fl. Flum. 314; vii. t. 134 (= *Collaea speciosa*).—Peruv.; Bras.
amplexicaulis, *E. Mey. Comm. Pl. Afr. Austr.* 92.—Afr. austr.
ampylocladus, Pritz. Ic. Ind. i. 660, sphalm. = campylocladus.
angivensis, Boj. ex Baker, in Journ. Bot. xx. (1882) 69 Vigna angivensis.
angustifolius, Gouan, Fl. Monsp. 201 = angustissimus.
angustifolius, Gueldenst. Reisen, ii. 65 = corniculatus.
angustifolius, Moç. & Sesse, ex G. Don, Gen. Syst. ii. 200 = Hosackia angustifolia.
angustissimus, *Linn. Sp. Pl.* 774.—Europ.; Oriens; As. bor.
anthylloides, Boiss. & Noé, Diagn. Ser. II. ii. 20 = L. Gebelia.
anthylloides, *Vent. Jard. Malm.* 92. t. 92.—Afr. austr.
approximatus, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 528 = hispidus.
arabicus, Bourg. ex Coss. in Bull. Soc. Bot. Fr. xxvii. (1880) 71 = trigonelloides.
arabicus, *Linn. Mant.* i. 104.—Afr. bor.; Arab.
arabicus, [Soland. in] Russ. Aleppo, ed. II. t. 14 = L. Gebelia.
arboreus, Forst. f. Prod. 52 = Carmichaelia australis.
arenarius, *Brot. Fl. Lusit.* ii. 120.—Reg. Mediterr. occ.
argenteus, Brot. l. c. 118 = Argyrolobium Linnaeanum.
argenteus, Salisb. Prod. 333 = creticus.
argenteus, *Webb & Berth. Phyt. Canar.* ii. 87.—Afr. bor.
argolicus, Link, in Linnaea, ix. (1834) 584 = cytisoides.
aristatus, DC. Cat. Hort. Monsp. 122 = conimbricensis.
arvensis, Pers. in Usteri, Ann. Bot. xiv. (1795) 39; Schkuhr, Handb. ii. 413. t. 211 = corniculatus.
atropurpureus, DC. Cat. Hort. Monsp. 121 = anthylloides.
Aucheri, Boiss. Diagn. Ser. I. ii. 38 = pusillus.
aurantiacus, Boiss. Elench. 37 = arenarius.
aurantius, *Steud. Nom.* ed. II. ii. 74.—Hispan.
australis, *Andr. Bot. Rep.* t. 624.—Austral.
belgradicus, Forsk. Fl. Aegypt. Arab. 215 = Dorycnium latifolium.
Bertholetii, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 160.—Ins. Canar.
biflorus, *Desr. in Lam. Encyc.* iii. 604.—Italia; Sicil.
biflorus, Huet, ex Nym. Consp. 182 = Bivoneus.
Bivoneus, *Guss. Fl. Sic. Syn.* ii. 349.—Sicil.
Bouteloui, Nym. Syll. Suppl. 52 = siliquosus.
brachycarpus, Benth. & Hook. f. ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 225 = Hosackia brachycarpa.
brachycarpus, *Hochst. ex Steud. in Flora*, xxiv. (1841) 1. *Intell.* 32; *et ex A. Rich. Tent. Fl. Abyss.* i. 177. —Abyssin.

LOTUS:—
bracteatus, Wall. Cat. n. 5939 = corniculatus.
Broussonetii, Choisy, ex Ser. in DC. Prod. ii. 211 = Dorycnium Broussonetii.
Brunneri, *Webb, in Hook. Niger Fl.* 119. t. 3.—Ins. Cap. Virid.
campestris, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 181 = tenuifolius.
campylocladus, *Webb & Berth. Phyt. Canar.* ii. 83. —Ins. Canar.
candidus, Mill. Gard. Dict. ed. VIII. n. 7 = Dorycnium hirsutum.
Candolleanus, Sweet, Hort. Brit. ed. I. 477 = australis.
canescens, Kuntze, in Flora, xxix. (1846) 697 = arenarius.
capitellatus, *C. Gay, Fl. Chil.* ii. 80.—Chili.
Carmeli, *Boiss. Diagn.* Ser. I. ix. 34.—Palaestin.
castellanus, *Boiss. & Reut. l. c.*—Hispan.
cernuus, Desr. in Lam. Encyc. ii. 612 = Melolobium cernuum.
cervinus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 340 = edulis.
ciliatus, Amo, Desc. Pl. Nov. 5 = conimbricensis.
ciliatus, Tenore, Hort. Neap. App. 66; & Prod. Fl. Nap. 44 = angustissimus.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 182.—Alger.
coccineus, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 59 = Requieni.
coccineus, Schlecht. in Linnaea, xxi. (1848) 452 = australis.
coccineus, Vell. Fl. Flum. 315; vii. t. 135 (= *Collaea speciosa*).—Bras.
coimbrensis, Willd. Sp. Pl. iii. 1390 = conimbricensis.
coimbricensis, Steud. Nom. ed. II. ii. 74 = praec.
colocensis, *Menyh. Kalocsa Videk. Növenyt.* 67 (Quid?). —Hungar.
commutatus, *Guss. Fl. Sic. Prod.* ii. 545.—Reg. Mediterr. occ.
conimbricensis, *Brot. Fl. Lusit.* ii. 118.—Reg. Mediterr.
conjugatus, Guss. Fl. Sic. Prod. ii. 534 = Requieni.
conjugatus, *Linn. Sp. Pl.* 447.—Reg. Mediterr.
conjugatus, Poir. Voy. Barb. 220 = biflorus.
corniculatus, *Linn. Sp. Pl.* 775.—Geront. temp.; Austral.
coronillaefolius, Guss. Fl. Sic. Prod. ii. 543 = cytisoides.
coronillaefolius, *Webb, in Hook. Niger Fl.* 119.—Ins. Cap. Virid.
Crantzii, *Vis. Fl. Dalm.* iii. 304.—Dalmat.
crassifolius, Pers. Syn. ii. 354 = corniculatus.
creticus, *Linn. Sp. Pl.* 775.—Reg. Mediterr.
creticus, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 423 = commutatus.
creticus, Sibth. Fl. Graec. t. 758 = cytisoides.
creticus, Viv. Fl. Lib. Spec. 46 = argenteus.
cytisoides, *Linn. Sp. Pl.* 776.—Europ. austr.; Afr. bor.
decumbens, Poir. Encyc. Suppl. iii. 508; T. F. Forst. Fl. Tonbr. 86 = tenuifolius.
Delorti, Timb. ex F. Schultz, Arch. Fl. Fr. et Allem. 201 = corniculatus.
depressus, Willd. Enum. Hort. Berol. Suppl. 52 = corniculatus.
dichotomus, Delile, Fl. Aegypt. n. 717; ex Walp. Rep. ii. 838 = Lotononis Leobordea.
diffusus, Schrank, Pl. Rar. Hort. Monac. t. 40; ex DC. Prod. ii. 211 = cytisoides.
diffusus, Sibth. & Sm. Prod. Fl. Graec. ii. 104 = ornithopodioides.
diffusus, Soland. ex Sm. Fl. Brit. ii. 794 = angustissimus.
digitatus, Lam. Fl. Fr. ii. 632 = Dorycnium suffruticosum.
Dioscoridis, All. Fl. Pedem. i. 310. t. 59 = peregrinus.
discolor, *E. Mey. Comm. Pl. Afr. Austr.* 92.—Afr. austr.
dispermus, Desv. Journ. Bot. iii. (1814) 77 = Dorycnium dispermum.
divaricatus, *Boiss. Diagn.* Ser. I. ii. 37.—As. Min.
divaricatus, Soland. ex Buch, in Abh. Acad. Berl. (1816) 198 = hispidus.
dorychnoides, Poir. Encyc. Suppl. iii. 507 = sessilifolius.
Dorycnium, Crantz, Stirp. Austr. ed. II. fasc. v. 402 = Dorycnium herbaceum.
Dorycnium, Linn. Sp. Pl. 778 = Dorycnium suffruticosum.
Dorycnium, Pall. ex Georgi, Beschr. Russ. Reich. III. iv. 1195 = Dorycnium pentaphyllum.
drepanocarpus, *Dur. in Duch. Rev. Bot.* ii. (1846) 438.—Alger.

LOTUS:—
dumetorum, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 423 = arenarius.
edulis, *Linn. Sp. Pl.* 774.—Reg. Mediterr.
Ehrenbergii, Schweinf. ex Boiss. Fl. Orient. ii. 170 = Schimperi.
erectus, Linn. Sp. Pl. 774 = siliquosus.
erectus, Vell. Fl. Flum. 314; vii. t. 136 (= *Collaea speciosa*).—Bras.
eriophthalmus, Webb & Berth. Phyt. Canar. ii. 89 = Dorycnium eriophthalmum.
exstipularis, Steud. Nom. ed. I. 495 = Indigofera sarmentosa.
extipulatus, DC. Prod. ii. 231 in syn. = praec.
filicaulis, *Dur. in Duch. Rev. Bot.* ii. (1846) 438.—Alger.
filiformis, Berg. Desc. Pl. Cap. 227 = Indigofera sarmentosa.
filiformis, Poir. Encyc. Suppl. iii. 504 = hispidus.
flexuosus, *Desr. in Lam. Encyc.* iii. 606.—Hab.?
floridus, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 160.—Ins. Canar.
fluminensis, Vell. Fl. Flum. 313; vii. t. 132 = Clitoria cajanifolia.
Forsteri, Sweet, Hort. Brit. ed. I. 118 = corniculatus.
Fraseri, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 400.—Austral.
fruticosus, Berg. Desc. Pl. Cap. 226 = Indigofera sarmentosa.
fruticulosus, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 57, nomen.—Marocc.
fruticulosus, *Desf. Tabl.* ed. II. 219 (Quid?).—Hab.?
Garcini, *DC. Prod.* ii. 212.—Nubia; Persia; Ind. or.
Gebelia, *Vent. Jard. Cels*, t. 57.—Oriens.
gibbus, Beeke, in Turn. & Dillw. Bot. Guide, ii. 528 = corniculatus.
glaber, Mill. Gard. Dict. ed. VIII. n. 3 = tenuis?
glaberrimus, DC. Cat. Hort. Monsp. 122 = conimbricensis.
glaberrimus, Schur, Enum. Pl. Transs. 160 = angustissimus.
glacialis, Bourg. ex Nym. Consp. 183 = glareosus.
glareosus, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 36.—Hispan.
glaucescens, Presl, Delic. Prag. 47 = cytisoides.
glaucus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 92.—Mader.; Liguria.
glaucus, Sieber, ex Reichb. Fl. Germ. Excurs. 505 = cytisoides.
glinoides, *Delile, in Ann. Sc. Nat.* Sér. II. vii. (1837) 286.—Aegypt.; Arab.
glomeratus, Lam. Fl. Fr. ii. 633 = Dorycnium rectum.
gracilis, Salisb. Prod. 334 = Dorycnium suffruticosum.
gracilis, Waldst. & Kit. Pl. Rar. Hung. iii. 254. t. 229 = angustissimus.
graecus, Linn. Mant. i. 101 = Dorycnium latifolium.
Gussonii, Huet, ex Nym. Consp. 182 = Requieni.
halophilus, Boiss. & Sprun. Diagn. Ser. I. ii. 37 = pusillus.
hemorroidalis, Lam. Fl. Fr. ii. 633 = Dorycnium hirsutum.
hirsutus, Linn. Sp. Pl. 775 = Dorycnium hirsutum.
hispidus, DC. Fl. Fr. iv. 556 = parviflorus.
hispidus, *Desf. Tabl.* ed. I. 190.—Europ. et. Afr. bor. occ.
hispidus, Sprun. & Heldr. ex Nym. Consp. 183 = lamprocarpus.
holosericeus, *Webb & Berth. Phyt. Canar.* ii. 83.—Ins. Canar.
humifusus, Willd. Enum. Hort. Berol. Suppl. 52 = corniculatus.
humilis, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 424 = parviflorus.
incanus, Dougl. ex Hook. Fl. Bor. Am. i. 134 = Hosackia decumbens.
indicus, Desr. in Lam. Encyc. iii. 606 = Rothia trifoliata.
intermedius, Loisel. Fl. Gall. ed. I. 489; Pers. Syn. ii. 354 = Dorycnium hirsutum.
involucratus, Desr. in Lam. Encyc. iii. 611 = Lotononis involucrata.
Jacobaeus, *Linn. Sp. Pl.* 775.—Ins. Cap. Virid.
judaicus, Boiss. Fl. Orient. ii. 165 = cytisoides.
laevigatus, Benth. in Mitch. Journ. Trop. Austral. 62 = australis.
lamprocarpus, *Boiss. Diagn.* Ser. I. ix. 33.—Graecia; Oriens.
lanatus, *C. Sm. in Tuckey. Congo*, 251.—Ins. Cap. Virid.
lancerottensis, *Webb & Berth. Phyt. Canar.* ii. 84.—Ins. Canar.

LOTUS:—
lanuginosus, *Vent. Jard. Malm.* 92.—Arabia.
lateralis, *Presl, Fl. Sicul.* p. xxiii.—Sicilia.
latifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 107 = Dorycnium latifolium.
Levieri, Heldr. in Nuov. Giorn. Bot. vii. (1875) 297 = hispidus.
libanoticus, Boiss. Diagn. Ser. I. ix. 32 = L. Gebelia.
linearis, Walp. in Linnaea, xiii. (1839) 518 = L. Jacobaeus.
longicaulis, Welw. ex Nym. Consp. 183 = lamprocarpus.
longisiliquosus, *Roem. ex Willk. in Linnaea*, xxv. (1852) 22.—Hispan.
Loweanus, *Webb & Berth. Phyt. Canar.* ii. 87.—Ins. Canar.
lugubris, Salisb. Prod. 333 = L. Jacobaeus.
Macraei, Benth. in Trans. Linn. Soc. xvii. (1837) 367 = Hosackia subpinnata.
macranthus, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 24.—Ins. Mader.
macrotrichus, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 250 = divaricatus.
major, *Scop. Fl. Carn.* ed. II. ii. 86.—Europ.
major, Sm. Engl. Bot. t. 2091 = uliginosus.
maritimus, Linn. Sp. Pl. 773 = siliquosus.
maritimus, Vell. Fl. Flum. 314; vii. t. 133 = Phaseolus semierectus.
maroccanus, *Ball, in Journ. Bot.* xi. (1873) 306.—Marocc.
mauritanicus, Linn. Syst. ed. X. 1179 = Indigofera coriacea.
medicaginoides, Retz. Obs. ii. 23 = Trigonella arcuata.
melilotoides, *Webb, in Hook. Niger Fl.* 118.—Ins. Cap. Virid.
Michauxianus, Ser. in DC. Prod. ii. 211 = L. Gebelia.
micranthus, Benth. in Trans. Linn. Soc. xvii. (1837) 367 = Hosackia parviflora.
microcarpus, Brot. Fl. Lusit. ii. 119 = parviflorus.
microphyllus, Hook. Bot. Mag. t. 2808 = Indigofera gracilis, *Spreng.*
minor, Bishop, in Edinb. Phil. Journ. xiv. (1826) 180 = tenuis.
minor, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 531 = praec.?
mollis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 509.—Ins. Socotra; Afr. austr.
mollissimus, S. G. Gmel. ex Ledeb. Fl. Ross. i. 560 = angustissimus.
montanus, A. Rich. Tent. Fl. Abyss. i. 177 = nubicus.
montanus, *Schur, in Verh. Naturf. Ver. Bruenn*, xv. II. (1877) 180.—Europ.
mossamedensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 62.—Afr. trop.
neglecta, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 160.—Ins. Canar.
Nočanus, Boiss. Diagn. Ser. II. ii. 21 = tenuifolius.
nubicus, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 61.—Afr. trop.
nubicus, Stocks, ex Boiss. Fl. Orient. ii. 174 = Garcini.
nummularius, *Reichb. ex Steud. Nom.* ed. II. ii. 74.—Lusitan.
obtusatus, Ser. in DC. Prod. ii. 212 = creticus.
odoratus, Sims, Bot. Mag. t. 1233 = hispidus.
oligoceratos, Desf. Fl. Atlant. ii. 206 = edulis.
oligoceratos, Desr. in Lam. Encyc. iii. 605 = peregrinus.
oligoceratos, Link, ex Nym. Consp. 184 = pusillus.
oligoceratos, Scop. Del. Insub. i. 14 = angustissimus.
ononoides, Desr. in Lam. Encyc. iii. 609 = Ononis aspalathoides, Lotononis involucrata.
ononopsis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 509.—Ins. Socotr.
ornithopodioides, *Linn. Sp. Pl.* 775.—Reg. Mediterr.
ornithopodioides, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 180 = corniculatus.
palustris, Ledeb. Fl. Ross. i. 562 = lamprocarpus.
palustris, Vell. Fl. Flum. 315; vii. t. 137 = Sesbania exasperata.
palustris, Willd. Sp. Pl. iii. 1394 = angustissimus.
parviflorus, *Desf. Fl. Atlant.* ii. 206. t. 211.—Reg. Mediterr.
patens, Presl, Delic. Prag. 48 = cytisoides.
pedunculatus, Cav. Ic. ii. 52. t. 164 = uliginosus.
pentaphyllos, Gilib. Fl. Lituan. ii. 93 = corniculatus.
pentaphyllus, *Link, in Buch, Beschr. Canar. Ins.* 156.—Ins. Canar.
peregrinus, Burm. f. Fl. Ind. 173 = Indigofera trifoliata.
peregrinus, *Linn. Sp. Pl.* 774.—Reg. Mediterr. or.

LOTUS:—
persicus, Burm. f. Fl. Ind. 173 = Ononis aspalathoides.
pilosissimus, Poir. Encyc. Suppl. iii. 504 = hispidus.
pilosissimus, Schur, Enum. Pl. Transs. 160 = corniculatus.
pilosus, Beeke, in Turn. & Dillw. Bot. Guide, ii. 528 = uliginosus.
pilosus, Jord. Pugill. Pl. Nov. (1852) 60 = corniculatus.
pilosus, Medic. Bot. Beobacht. (1783) 226 = hirsutus?
pilosus, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 424 = parviflorus.
pinnatus, Hook. Bot. Mag. t. 2913 = Hosackia bicolor.
pisifolius, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimpr. 24 = uliginosus.
platycarpos, Viv. Pl. Aegypt. Dec. 14 = argenteus.
polyphyllus, Clarke, Trav. iii. 41 = argenteus.
pratensis, Mill. Gard. Dict. ed. VIII. n. 10 = siliquosus.
Preslii, Tenore, Fl. Nap. v. 160 = tenuis.
prostratus, Desf. Fl. Atlant. ii. 206 = cytisoides.
prostratus, Linn. Syst. ed. X. 1179 = Lotononis prostrata.
purpureus, *Webb, in Hook. Niger Fl.* 118. t. 6.—Ins. Cap. Virid.
pusillus, *Medic. Beobacht.* (1783) 226.—Reg. Mediterr. or.
racemosus, Poir. Encyc. Suppl. iii. 508 = Indigofera alopecuroides, coriacea.
rectus, Falk, Beitr. ii. 229 = Dorycnium pentaphyllum.
rectus, Georgi, Reise, i. 227 (Quid?).—Sibir. baical.
rectus, Linn. Sp. Pl. 775 = Dorycnium rectum.
repens, Moç. & Sesse, ex G. Don, Gen. Syst. ii. 200 = Hosackia repens.
Requieni, *Mauri, in Tenore, Viag. Abruz.* 81; *ex Bertol. Fl. Ital.* viii. 214.—Ital.
Requienii, Hort. ex DC. Neuvième Not. Pl. Rar. Genèv. 7 = conjugatus.
riparius, Pers. in Usteri, Ann. Bot. xiv. (1795) 39 = corniculatus.
roseus, Forsk. Fl. Aegypt. Arab. 140 = arabicus.
Salzmanni, Boiss. & Reut. Pugill. Pl. Nov. (1852) 37 = commutatus.
Schimperi, *Steud. Nom.* ed. II. ii. 75.—Aegypt.; Arab.
secundiflorus, Viv. Fl. Lib. Spec. 46. t. 21 = cytisoides.
sericeus, DC. Cat. Hort. Monsp. 122 = Dorycnium hirsutum.
sericeus, Moench, Meth. Suppl. 53 = creticus.
sericeus, Pursh, Fl. Am. Sept. ii. 489 = Hosackia Purshiana.
sessilifolius, *DC. Cat. Hort. Monsp.* 122.—Ins. Teneriff.
siliquosus, *Linn. Syst.* ed. X. 1178.—Reg. Mediterr.; As. Min.
spartioides, *Webb & Berth. Phyt. Canar.* ii. 81.—Ins. Canar.
spectabilis, Choisy, ex Ser. in DC. Prod. ii. 211 = Dorycnium spectabile.
stenodon, Heldr. ex Nym. Consp. 183 = corniculatus.
Stocksii, Boiss. Fl. Orient. ii. 174 = Garcini.
strictus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 32.—As. Min.
suaveolens, *Pers. Syn.* ii. 354.—Gall. austr.
subbiflorus, Lag. in Varied. Cienc. ii. IV. (1805) 213 = hispidus.
subpinnatus, Lag. Gen. et Sp. Nov. 23 = Hosackia subpinnata.
suffruticosus, Burm. f. Fl. Cap. Prod. 22 = Aspalathus suffruticosa.
sulphureus, *Boiss. Diagn.* Ser. I. ii. 35.—As. Min.
symmetricus, Jord. Pugill. Pl. Nov. (1852) 59 = corniculatus.
tauricus, Hort. ex Steud. Nom. ed. II. ii. 75 = corniculatus.
Tchihatchewii, Boiss. Fl. Orient. ii. 165 = corniculatus.
tenuifolius, Burm. f. Fl. Cap. Prod. 22 = Psoralea fascicularis.
tenuifolius, Pollich, ex Steud. Nom. ed. II. ii. 75 = corniculatus.
tenuifolius, *Presl, Delic. Prag.* 46.—Sicil.
tenuifolius, Reichb. Fl. Germ. Excurs. 506 = tenuis.
tenuis, *Waldst. & Kit. ex Willd. Enum. Hort. Berol.* 797.—Europ.; Oriens.
Tetragonolobus, *Linn. Sp. Pl.* 773.—Reg. Mediterr.
tetraphyllus, *Murr. Syst.* ed. XIII. 575.—Ins. Balear.
tetraptera, Stokes, Bot. Mat. Med. iv. 74 = L. Tetragonolobus.
thermalis, Boiss. Diagn. Ser. I. ii. 35 = strictus.
tigrensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 61.—Afr. trop.

LOTUS:—
tingitanus, *Boiss. Diagn.* Ser. I. ii. 36.—Marocc.
tomentosus, DC. Cat. Hort. Monsp. 122, in syn. cum cit. falsa = Dorycnium hirsutum.
tomentosus, *Desr. in Lam. Encyc.* iii. 612.—Hab.?
trichocarpus, Lag. Gen. et Sp. Nov. 23 = conimbricensis.
Trifoliastrum, *Desr. in Lam. Encyc.* iii. 612.—Oriens.
trigonelloides, *Webb & Berth. Phyt. Canar.* ii. 86. t. 65.—Ins. Canar.
tristis, Moench, Meth. Suppl. 53 = L. Jacobaeus.
uliginosus, Hoffm. Deutschl. Fl. ed. II. ii. 100 = corniculatus.
uliginosus, *Schkuhr, Handb.* ii. 412. t. 211.—Europ.; Reg. Mediterr.
unibracteatus, Viv. Fl. Lib. Spec. 48. t. 18. f. 4 = hispidus.
unifoliatus, Benth. in Trans. Linn. Soc. xvii. (1837) 368 = Hosackia Purshiana.
unifultus, Boiss. Diagn. Ser. I. ii. 36 = L. Gebelia.
utricularis, Domb. & Lag. ex DC. Prod. ii. 171 = Hosackia subpinnata.
valde-pilosus, Schur, Enum. Pl. Transs. 160 = corniculatus.
varians, Desv. Journ. Bot. iii. (1814) 77 = creticus.
versicolor, Tineo, Pl. Rar. Sicil. 27 = corniculatus.
villosus, *Burm. f. Fl. Cap. Prod.* 23 (Quid?).—Afr. austr.
villosus, Forsk. Fl. Aegypt. Arab. p. lxxi = peregrinus.
villosus, Thuill. Fl. Par. ed. II. 387 = uliginosus.
vindicatus, Boengh. Cat. Sem. Hort. Monast. (1829); ex Koch, in Roehl. Deutschl. Fl. ed. III. v. 306 = major.
Wiedemanni, *Nym. Consp.* 182.—Arch. Graec.
Wrangelianus, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 41 = Hosackia subpinnata.

LOUCORYNE, Steud. Nom. ed. II. ii. 38 (1841), sphalm. = **Leucocoryne**, Lindl. (Liliac.).

LOUDETIA, Hochst. in Flora, xxiv. (1841) 20; xxiv. (1841) II. 713 = **Trichopteryx**, Nees (Gramin.).
arundinacea, Hochst. ex Steud. Syn. Pl. Gram. 238 = Tristachya arundinacea.
barbata, A. Br. in Flora, xxiv. (1841) II. 714 = Tristachya barbata.
elegans, Hochst. l. c. xxiv. (1841) I. Intell. 20; et ex Steud. Syn. Pl. Gram. 238 = Tristachya elegans.
superba, De Not. Ind. Sem. Hort. Genuens. (1852) 24; et in Ann. Sc. Nat. Sér. III. xix. (1853) 369.—Nubia.

LOUDONIA, Bert. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 193 = **Adesmia**, DC. (Legumin.).
anthylloides, Bert. ex Loud. Encyc. Pl. Suppl. i. 1244 = A. Loudonia.

LOUDONIA, Lindl. Swan River App. 42 (1839). *HALORAGEAE*, Benth. & Hook. f. i. 674.
GLISCHROCARYON, Endl. in Ann. Wien. Mus. ii. (1839) 209.
aurea, *Lindl. Swan River Bot. App.* 42.—Austral.
Behrii, *Schlecht. in Linnaea*, xx. (1847) 648.—Austral.
citrina, F. Muell. l. c. xxv. (1852) 385 = aurea.
flavescens, J. Drumm. in Hook. Lond. Journ. Bot. i. (1842) 396 = aurea.
Roei, *Schlecht. in Linnaea*, xx. (1847) 648.—Austral.

LOUICHEA, L'Hérit. Stirp. Nov. 135. t. 65 (1785) = **Pteranthus**, Forsk. (Illecebrac.).
cervina, L'Hérit. l. c. = P. echinatus.
Pteranthus, L'Hérit. l. c. = P. echinatus.

LOUISIA, Reichb. f. Xenia Orch. 204 (1858) = **Luisia**, Gaudich. (Orchid.).

LOURADIA, Leman, in Dict. Sc. Nat. xxvii. 235 (1823) = **Lavradia**, Vell. (Violar.).

LOUREA, Kunth, in Linnaea, xiii. (1839) 562 = LAUREA, Gaudich. = **Bagassa**, Aubl. (Urtic.).

LOUREA, Neck. Elem. iii. 17 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 522.
campanulata, *Benth. in Miq. Pl. Jungh.* 215.—Burma.
microphylla, Wall. Cat. n. 5672 = obcordata.

LOUREA:—
obcordata, *Desv. Journ. Bot.* i. (1813) 122.—Burma; Malaya; Austral.
paniculata, *Wall. Cat.* n. 5673.—Burma.
reniformis, DC. Prod. ii. 324; Desv. in Mém. Soc. Linn. Par. iv. (1826) 308 = obcordata.
Vespertilionis, *Desv. Journ. Bot.* i. (1813) 122. t. 5. f. 18.—Reg. trop.

LOUREIRA, Cav. Ic. v. 17. t. 429 (1799) = **Jatropha**, Linn. (Euphorb.).
cuneifolia, Cav. l. c. = J. spathulata.
glandulosa, Cav. l. c. 18. t. 430 = J. cordata.
peltata, Desv. Cat. Hort. Par. ed. III. 411 = J. hernandiaefolia.

LOUREIRA, Meissn. Gen. Comm. 53 (1837), 'genus ignotum,' Benth. & Hook. f. i. 322, 418.
TOLUIFERA, Lour. Fl. Cochinch. 362 (1790).
cochinchinensis, *Meissn. l. c.*—Cochinch.

LOUREIRA, Raeusch. Nom. ed. III. 67 (1797) = **Elaeodendron**, Jacq. (Celastr.).
albens, Raeusch. l. c. = E. glaucum.

LOWEA, Lindl. Bot. Reg. t. 1261 (1829) = **Rosa**, Tourn.
berberifolia, Lindl. l. c. = R. berberifolia.

LOWELLIA, A. Gray, in Mem. Am. Ac. N. S. iv. (1849) 89 = **Thymophyllum**, Lag. (Compos.).
aurea, A. Gray, l. c. 91 (= *Hymenatherum aureum*). —N. Mexic.

LOXANTHERA, Blume, in Schult. f. Syst. vii. 1730 (1830) = **Loranthus**, Linn.
corymbosa, Blume, l. c. 1612 = Loranthus tetrandrus.
Eschscholzianus, Blume, l. c. = Loranthus Eschscholzianus.
grandiflorus, Blume, l. c. = Loranthus grandiflorus, *Ruiz & Pav.*
longibracteatus, Blume, l. c. = Loranthus longibracteatus.
speciosa, Blume, l. c. 1730 = Loranthus loxantherus.
spirostylis, G. Don, Gen. Syst. iii. 430 = Loranthus spirostylis.
Sternbergianus, Blume, in Schult. f. Syst. vii. 1612 = Loranthus Sternbergianus.

LOXANTHES, Rafin. Neogenyt. 3 (1825) = **Aphyllon**, Mitch. (Orobanch.).
fasciculata, Rafin. l. c. = A. fasciculatum.

LOXANTHES, Salisb. Gen. Pl. Fragm. 117 (1866) = **Nerine**, Herb. (Amaryll.),
flexuosa, Salisb. l. c. 117 = N. flexuosa.
humilis, Salisb. l. c. = N. humilis.
laticoma, Salisb. l. c. = N. lucida.
marginata, Salisb. l. c. = N. marginata.
pendula, Salisb. l. c. = Cyrtanthus obliquus.
striata, Salisb. l. c. = Brunsvigia striata.
undulata, Salisb. l. c. = N. undulata.

LOXANTHUS, Nees, in Wall. Pl. As. Rar. iii. 76, 322 (1832) = **Phlogacanthus**, Nees (Acanth.).
Gomezii, Nees, in DC. Prod. xi. 322 = Cystacanthus pulcherrimus.
Gomezii, Nees, in Wall. Pl. As. Rar. iii. 99 = P. Wallichii.

LOXIDIUM; Vent. Dec. Nov. Gen. 9 (1808) = **Swainsona**, Salisb. (Legum.).
galegifolium, Vent. ex Poir. in Dict. Sc. Nat. xxvii. 252 = S. galegifolia.

LOXOCARPUS, R. Br. in Benn. Pl. Jav. Rar. 120 (1838) = **Didymocarpus**, Wall. (Gesner.).
alata, R. Br. l. c. = D. incana.
incana, R. Br. l. c. = D. incana.

LOXOCARYA, R. Br. Prod. 249 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1033.
DESMOCLADUS, Nees, in Lehm. Pl. Preiss. ii. 56 (1846).
HAPLOSTIGMA, F. Muell. Fragm. viii. 100 (1873).
Benthami, F. Muell. Census, 124 = Hypolaena Benthamii.
cinerea, *R. Br. Prod.* 249.—Austral.
densa, Benth. Fl. Austral. vii. 241 = Hypolaena densa.

LOXOCARYA:—
fasciculata, Benth. Fl. Austral. vii. 242 = Hypolaena fasciculata.
flexuosa, Benth. l. c. 243 = Hypolaena flexuosa.
pubescens, Benth. l. c. 242 = Hypolaena pubescens.
vestita, Benth. l. c. 241 = Hypolaena vestita.
virgata, Benth. l. c. 242 = Hypolaena Benthamii.

LOXOCOCCUS, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 185. *PALMAE*, Benth. & Hook. f. iii. 888.
rupicola, *H. Wendl. & Drude, l. c.*—Zeylan.

LOXODISCUS, Hook. f. in Hook. Kew Journ. ix. (1857) 200. t. 6. *SAPINDACEAE*, Benth. & Hook. f. i. 397.
coriaceus, *Hook. f. l. c.*—Ins. Pinet.

LOXODON, Cass. in Dict. Sc. Nat. xxvii. 253 (1823) = **Chaptalia**, Vent. (Compos.).
brevipes, Cass. l. c. = C. exscapa.
chilensis, DC. Prod. vii. 44 = C. exscapa.
longipes, Cass. in Dict. Sc. Nat. xxvii. 255 = Chaptalia runcinata.

LOXONIA, Jack, in Trans. Linn. Soc. xiv. (1823) 40. *GESNERACEAE*, Benth. & Hook. f. ii. 1019.
LOXOPHYLLUM, Blume, Bijdr. 750 (1826).
acuminata, *R. Br. in Benn. Pl. Jav. Rar.* 105. t. 25.—Java.
alata, Wall. Cat. n. 809 = Didymocarpus incana.
decurrens, Blume, Bijdr. 776 = Cyrtandromoea decurrens.
discolor, Jack, in Trans. Linn. Soc. xiv. (1823) 40 = acuminata.
hirsuta, Jack, l. c. 41 = acuminata.

LOXOPHYLLUM, Blume, Bijdr. 750 (1826) = **Loxonia**, Jack (Gesnerac.).
racemosum, Blume, l. c. 751 = Loxonia acuminata.

LOXOPTERYGIUM, Hook. f. in Benth. & Hook. f. Gen. i. 419 (1862). *ANACARDIACEAE*, Benth. & Hook. f. l. c.
Gardneri, *Engl. in DC. Monog. Phan.* iv. 460.—Bras.
Grisebachii, *Hiern, ex Griseb. in Goett. Abh.* xxiv. (1879) 95.—Am. austr.
Huasango, *Spruce, ex Engl. in DC. Monog. Phan.* iv. 461.—Ecuador.
Kuasango, Spruce, ex March. Rev. Anacard. 179, nomen = praec.?
Lorentzii, Griseb. in Goett. Abh. xix. (1874) 115 = Quebrachia Lorentzii.
Sagotii, *Hook. f. in Benth. & Hook. f. Gen.* i. 420.—Guiana.

LOXOSPERMUM, Hochst. in Flora, xxix. (1846) 594 = **Trifolium**, Linn. (Legumin.).
calocephalum, Hochst. l. c. 595 = T. calocephalum.
multinerve, Hochst. l. c. = T. multinerve.
Schimperi, Hochst. l. c. 594 = T. Schimperi.

LOXOSTEMON, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129, 147. *CRUCIFERAE*, Benth. & Hook. f. i. 70.
pulchellus, *Hook. f. & Thoms. l. c.* 147.—Reg. Himal.

LOXOSTIGMA, C. B. Clarke, in DC. Monog. Phan. v. 59 (1883). *CYRTANDREAE*.
Griffithii, *C. B. Clarke, l. c.* 60.—Reg. Himal.

LOXOSTYLIS, Spreng. ex Reichb. Ic. Exot. Cent. iii. 3. t. 205 (1827). *ANACARDIACEAE*, Benth. & Hook. f. i. 421.
ANASYLLIS, E. Mey. in Drège, Zwei Pfl. Docum. 163 (1843).
alata, *Spreng. ex Reichb. Ic. Exot. Cent.* iii. 3. t. 205. —Afr. austr.
latifolia, Presl, Bot. Bemerk. 42 = alata.

LOXOTIS, R. Br. in Wall. Pl. As. Rar. iii. 65 (1832) = **Rhynchoglossum**, Blume (Acanthac.).
intermedia, Benth. Scroph. Ind. 57 = R. obliquum.
obliqua, R. Br. in Benn. Pl. Jav. Rar. 102 = R. obliquum.

LOZANIA, Mutis, ex Caldas, in Seman. Nuev. Gran. (1810) III. 20 = **Qualea**, Aubl. (Vochysiac.).

LOZANIA:—
grandiflora, Schult. Mant. i. 109 = Q. grandiflora?
Mutisiana, Schult. l. c. = Q. cryptantha.
nemoralis, DC. Prod. iii. 30 = Q. cryptantha.

LUBINIA, Comm. ex Vent. Jard. Cels, t. 96 (1800) = **Lysimachia**, Tourn. (Primul.).
atropurpurea, Link & Otto, Ic. Pl. Sel. 61. t. 27 = Lys. nutans.
mauritiana, Spreng. Syst. i. 572 = Lys. mauritiana.
pacifica, Seem. Fl. Vit. 147, nomen = Lys. pacifica.
purpurea, Sweet, ex Steud. Nom. ed. II. ii. 75 = Lys. nutans.
spathulata, Vent. Jard. Cels, t. 96 = Lys. mauritiana.

LUCAEA, Kunth, Rév. Gram. ii. 489. t. 159 (1830) = **Arthraxon**, Beauv. (Gramin.).
ciliata, Steud. Syn. Pl. Gram. 414 = A. ciliaris.
gracilis, Kunth, Rév. Gram. ii. 489 = A. ciliaris.
Junghuhnii, Steud. Syn. Pl. Gram. 414 = A. Junghuhnii.
Langsdorffiana, Steud. l. c. 413 = A. ciliaris.
major, Hochst. ex Steud. l. c. 414 = A. ciliaris.
plumbea, Steud. l. c. = A. ciliaris.
Schimperi, Steud. l. c. = A. microphyllus.
typica, Buese, in Miq. Pl. Jungh. 367 = A. Junghuhnii.
violacea, Steud. Syn. Pl. Gram. 414 = A. ciliaris.
Vriesii, Buese, in Miq. Pl. Jungh. 366 = A. ciliaris.

LUCHIA, Steud. Nom. ed. II. ii. 75 (1841) = **Elodea**, Michx. (Hydrochar.).
Berteroniana, Steud. l. c. = E. chilensis.

LUCIANEA, Endl. Gen. 558 (1838) = **Lucinaea**, DC. (Rubiac.).

LUCILIA, Cass. in Bull. Soc. Philom. (1817) 32. *COMPOSITAE* [Benth. & Hook. f. ii. 306].
acutifolia, *Cass. in Dict. Sc. Nat.* xxvii. 64; *Less. in Linnaea*, v. (1830) 363.—Reg. Argent.
acutifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 114 = Jamesoni.
affinis, *Wedd. Chlor. And.* i. 230.—Bolivia.
aretioides, erythractis, hypoleuca, macrocephala, Schultzii, *Wedd. ex Sch. Bip. in Linnaea*, xxxiv. (1865-66) 532, nomina.—Bolivia.
argentea, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 102 = nitens.
chilensis, *Hook. & Arn. l. c.*—Chili.
conoidea, *Wedd. Chlor. And.* i. 154. t. 26. f. C.—Bolivia.
eriophora, *Remy, in C. Gay, Fl. Chil.* iii. 335.—Chili.
facelioides, *Sch. Bip. in Linnaea*, xxxiv. (1865-66) 532, nomen.—Bolivia.
ferruginea, *Baker, in Mart. Fl. Bras.* vi. III. 114.—Bras.
flagelliformis, *Wedd. Chlor. And.* i. 157. t. 26. f. D.—Bolivia.
glomerata, *Baker, in Mart. Fl. Bras.* vi. III. 114.—Bras.
gnaphalioides, Less. in Linnaea, v. (1830) 363 = nitens.
Jamesoni, *Baker, in Mart. Fl. Bras.* vi. III. 113.—Reg. Argent.
linearifolia, *Baker, l. c.* 114.—Bras.
Lundii, *Baker, l. c.* 113.—Bras.
microphylla, Cass. in Dict. Sc. Nat. xxvii. 64 = acutifolia.
nitens, *Less. in Linnaea*, v. (1830) 363.—Bras.
nivalis, *Sch. Bip. l. c.* xxxiv. (1865-66) 532, nomen.—Bolivia.
pedunculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 51.—Chili.
perpusilla, Wedd. Chlor. And. i. t. 26. f. A = Luciliopsis perpusilla.
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 138.—Peruv.
piptolepis, Wedd. Chlor. And. i. t. 26. f. B = Gnaphalium piptolepis.
plumosa, *Wedd. l. c.* 155.—Bolivia.
recurva, *Wedd. l. c.* 156. t. 25 B.—Bolivia.
squarrosa, *Baker, in Mart. Fl. Bras.* vi. III. 114.—Bras.
tomentosa, *Wedd. Chlor. And.* i. 157.—Bolivia.
violacea, *Wedd. l. c.* 155.—Bolivia.

LUCILIOPSIS, Wedd. Chlor. And. i. 159 (1856). *COMPOSITAE*, Benth. & Hook. f. ii. 301.
perpusilla, *Wedd. l. c.* 160 [t. 26. f. A].—Bolivia.

LUCINAEA, DC. Prod. iv. 368 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 67.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 216.—Ins. Philipp.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 187.—Sumatra.
montana, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) 167.—Borneo.
Morindae, *DC. Prod.* iv. 368.—Malaya.
Rheedii, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) 167.—Borneo.

LUCIOLA, Sm. Engl. Fl. ii. 177 (1824) = **Luzula**, DC. (Juncac.).
arcuata, Sm. l. c. 183 = Luzula arcuata.
campestris, Sm. l. c. 181 = Luzula campestris.
congesta, Sm. l. c. = Luzula campestris.
Forsteri, Sm. l. c. 179 = Luzula Forsteri.
pilosa, Sm. l. c. 178 = Luzula vernalis.
spicata, Sm. l. c. 182 = Luzula spicata.
sudetica, Sm. in Johnst. Fl. Berwick, i. 81 = Luzula nigricans.
sylvatica, Sm. Engl. Fl. ii. 180 = Luzula maxima.

LUCULIA, Sweet, Brit. Flow. Gard. t. 145 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 43.
gratissima, *Sweet, Brit. Flow. Gard.* t. 145.—Reg. Himal.
gratissima, Wall. Cat. n. 6117. B = Pinceana.
Pinceana, *Hook. Bot. Mag.* t. 4132.—Reg. Himal.

LUCUMA, Molina, Sagg. Chil. 186 (1782); Juss. Gen. 152 (1789). *SAPOTACEAE*, Benth. & Hook. f. ii. 654.
AMORPHOSPERMUM, F. Muell. Fragm. vii. 112 (1870).
GUAPEA, Endl. Ench. 361, in syn. (1841).
GUAPEBA, Gomez, in Mem. Ac. Lisboa, iii. (1812), Mem. dos Corresp. t. 2.
GUAPEBEIRA, Gomez, l. c. 19 (1812).
LABATIA, Sw. Prod. Veg. Ind. Occ. 32 (1788); Fl. Ind. Occ. i. 263. t. 6 (1797).
NIEMEYERA, F. Muell. Fragm. vii. 114 (1870).
OXYTHECE, Miq. in Mart. Fl. Bras. vii. 105 (1863).
POMETIA, Vell. Fl. Flum. 80 (1825).
PRADOSIA, Liais, Brésil, 614 (1872).
SERSALISIA, R. Br. Prod. 529 (1810), partim.
VITELLARIA, Gaertn. f. Fruct. iii. 131. t. 205 (1805).
Arguacoënsium, *Karst. Fl. Columb.* i. 129. t. 64.—N. Granat.
bifera, *Molina, Sagg. Chil.* 187.—Chili.
biflora, J. F. Gmel. Syst. 1578, sphalm. = praec.
Bonplandia, *H. B. & K. Nov. Gen. et Sp.* iii. 240.—Cuba.
Caimito, *Roem. Schult. & Syst.* iv. 701.—Peruv.
campechiana, *H. B. & K. Nov. Gen. et Sp.* iii. 240.—Mexic.
Capiri, *A. DC. in DC. Prod.* viii. 173.—Mexic.
Casaretti, *A. DC. l. c.* 671.—Bras.
catocladantha, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 202.—Bras.
chrysophyllifolia, Benth. & Hook. f. Gen. ii. 655, sphalm. = seq.
chrysophylloides, *A. DC. in DC. Prod.* viii. 168.—Bras.
curvifolia, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 68.—Bras.
deliciosa, *Planch. & Linden, in Rev. Hortic.* (1870–71) 836.—N. Granat.
dictyoneura, *Benth. & Hook. f. Gen.* ii. 655.—Cuba.
domingensis, *Gaertn. f. Fruct.* iii. 131. t. 204.—Ins. S. Doming.
elongata, Steud. Nom. ed. II. ii. 76 (= *Sapota elongata*).—Hab. ?
ferruginea, *Hook. & Arn. Bot. Beech. Voy.* 302.—Mexic.
galactoxylon, *F. Muell. Census*, 91.—Austral.
Gardneriana, *A. DC. in DC. Prod.* viii. 168.—Bras.
glabrescens, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 72.—Bras.
glaucescens, Mart. l. c. 420 = praec.
glomerata, *Miq. in Mart. l. c.* 81.—Bras.
glycyphloea, *Mart. & Eichl. ex Miq. l. c.* 82.—Bras.
gnaphaloclados, Mart. in Flora, xxi. (1838) II. Beibl. 89 = Sideroxylon Gardnerianum.
gomphiaefolia, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 78.—Bras.
grandiflora, *A. DC. in DC. Prod.* viii. 169.—Bras.

LUCUMA:—
Inophyllum, *Benth. & Hook. f. Gen.* ii. 654.—Bras.
Keule, Molina, Sagg. Chil. 187 = Gomortega nitida.
laevigata, *A. DC. in DC. Prod.* viii. 167.—Bras.
lasiocarpa, *A. DC. l. c.* 166.—Bras.
lateriflora, *Benth. ex Miq. in Mart. Fl. Bras.* vii. 83.—Bras.
laurifolia, *A. DC. in DC. Prod.* viii. 166.—Bras.
leptocarpa, *Benth. & Hook. f. Gen.* ii. 654.—Bras.
littoralis, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 65.—Bras.
longifolia, A. DC. in DC. Prod. viii. 167 = Sellowii.
lucens, *Mart. & Miq. in Mart. Fl. Bras.* vii. 78.—Bras.
mammosa, *Gaertn. f. Fruct.* iii. 129. t. 203.—Ind. or.; Am. austr.
marginata, *Mart. & Eichl. in Mart. Fl. Bras.* vii. 67.—Bras.
multiflora, *A. DC. in DC. Prod.* viii. 168.—Ind. occ.
neriifolia, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 282.—Bras.
nervosa, A. DC. in DC. Prod. viii. 169 = L. Rivicoa.
nitida, *A. DC. l. c.* 166.—Bras.
obovata, *H. B. & K. Nov. Gen. et Sp.* iii. 241.—Peruv.
pallida, *Gaertn. f. Fruct.* iii. 130. t. 204.—Hab. ?
paradoxa, *A. DC. in DC. Prod.* viii. 173.—Hab. ?
parviflora, *Benth. ex Miq. in Mart. Fl. Bras.* vii. 81.—Bras.
pauciflora, *A. DC. in DC. Prod.* vii. 168.—Ind. occ.
peduncularis, *Mart. & Eichl. ex Miq. l. c.* 73.—Bras.
procera, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 57.—Bras.
prunifera, F. Muell. Sel. Pl. Indust. Cult. 142 = Chrysophyllum pruniferum.
psammophila, *A. DC. in DC. Prod.* viii. 167.—Bras.
pulverulenta, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 70.—Bras.
ramiflora, *A. DC. in DC. Prod.* viii. 168.—Bras.
retusa, *Spruce, ex Miq. in Mart. Fl. Bras.* vii. 70.—Bras.
rigida, *Mart. & Eichl. ex Miq. l. c.* 73.—Bras.
Rivicoa, *Gaertn. f. Fruct.* iii. 130.—Bras.
Riviniana, Steud. Nom. ed. II. i. 13, sphalm. = praec.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 241.—Mexic.
Sellowii, *A. DC. in DC. Prod.* viii. 167.—Bras.
sericea; *Benth. & Hook. f. Gen.* ii. 654.—Austral.
serpentaria, *H. B. & K. Nov. Gen. et Sp.* iii. 242.—Cuba.
sphaerocarpa, *A. DC. in DC. Prod.* viii. 169.—Mexic.
spinosa, Molina, Sagg. Chil. 188 = Gourliea chilensis.
splendens, A. DC. in DC. Prod. viii. 171 = valparadisaea.
Temare, *H. B. & K. Nov. Gen. et Sp.* iii. 241.—Venezuela.
tomentosa, Poepp. ex Miq. in Mart. Fl. Bras. vii. 87 (= *Passaveria obovata*).—Bras.
torta, *A. DC. in DC. Prod.* viii. 167.—Bras.
turbinata, *Molina, Sagg. Chil.* 187.—Chili.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 83.—Cuba.
valparadisaea, *Molina, Sagg. Chil.* 187.—Bras.
venosa, *Mart. & Miq. in Mart. Fl. Bras.* vii. 67.—Bras.
venulosa, Spruce, ex Miq. l. c. 52 = Sideroxylon venulosum.
Warmingii, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 201.—Bras.

LUCYA, DC. Prod. iv. 434 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 61.
DUNALIA, Spreng. Pugill. ii. 25 (1815).
tuberosa, *DC. Prod.* iv. 434.—Ind. occ.

LUDDEMANIA, Reichb. f. in Linden, Pescatorea, i. (1860) t. 22 = LUEDDEMANNIA, Reichb. f. = **Cycnoches**, Lindl. (Orchid.).

LUDIA, Comm. ex Juss. Gen. 343 (1789). *BIXINEAE*, Benth. & Hook. f. i. 126.
bivalvis, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 244 = sessiliflora.
foetida, Roxb. Hort. Beng. 38; Fl. Ind. ii. 508 = Flacourtia sumatrana.
heterophylla, Bory, Voy. ii. 116 = Aphloia mauritiana.
heterophylla, Lam. Encyc. iii. 612; Ill. t. 466. f. 1, 2 = sessiliflora.

LUDIA:—
madagascariensis, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 244.—Madag.
mauritanica, J. F. Gmel. Syst. 1579 (Index) = seq.
mauritiana, J. F. Gmel. l. c. 839; Raeusch. Nom. ed. III. 159 = sessiliflora.
myrtifolia, Lam. Encyc. iii. 612 = sessiliflora.
sessiliflora, *Lam. l. c.*—Ins. Mascar.
spinosa, Roxb. Hort. Beng. 38; Fl. Ind. ii. 507 = Scolopia Roxburghii.
tuberculata, Jacq. Hort. Schoenb. i. 59 = sessiliflora.

LUDISIA, A. Rich. in Dict. Clas. Nat. vii. 437 (1825) = **Haemaria**, Lindl. (Orchid.).
argyroneura, Miq. in Journ. Bot. Néerl. i. (1861) 36 = Dossinia marmorata.
discolor, A. Rich. in Dict. Clas. Nat. vii. 437 = H. discolor.
Furetii, Blume, Orch. Archip. Ind. 114.—China.
odorata, Blume, l. c.—Cochinch.

LUDOLFIA, Adans. Fam. ii. 244 (1763) = **Tetragonia**, Linn. (Ficoid.).

LUDOLFIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 320 = **Arundinaria**, Michx. (Gramin.).
falcata, Nees, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 26 = A. falcata.
glaucescens, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 320 = A. glaucescens.
macrosperma, Willd. l. c. = A. macrosperma.
pinifolia, A. Dietr. Sp. Pl. ii. 25 = Chusquea pinifolia.
tecta, A. Dietr. l. c. 24 = A. macrosperma.
verticillata, A. Dietr. l. c. 25 = A. verticillata.

LUDOVIA, Brongn. in Ann. Sc. Nat. Sér. IV. xv. (1861) 361. *CYCLANTHACEAE*, Benth. & Hook. f. iii. 953.
crenifolia, *Drude, in Mart. Fl. Bras.* iii. II. 243.—Bras.
lancaefolia, *Brongn. in Ann. Sc. Nat.* Sér. IV. xv. (1861) 362.—Guiana.

LUDOVIA, Pers. Syn. ii. 576 (1807) = **Carludovica**, Ruiz & Pav. (Cyclanth.).
acuminata, Pers. l. c. = C. acuminata.
angustifolia, Pers. l. c. = Cyclanthus Plumieri.
funifera, Poit. in Mém. Mus. Par. ix. (1822) 28. t. 1 = C. funifera.
latifolia, Pers. Syn. ii. 576 = C. latifolia.
palmata, Pers. l. c. = C. palmata.
subacaulis, Poit. in Mém. Mus. Par. ix. (1822) 28 = C. trigona.
trigona, Pers. Syn. ii. 576 = C. trigona.

LUDOVICIA, Coss. in Bull. Soc. Bot. Fr. iii. (1856) 674 = **Hammatolobium**, Fenzl (Legumin.).
Kremeriana, Coss. l. c. 675 = H. Kremerianum.

LUDWIGHIA, Burm. f. Fl. Ind. 36 (1768) = seq.

LUDWIGIA, Linn. Coroll. Gen. 3 (1737). *ONAGRARIEAE*, Benth. & Hook. f. i. 788.
DANTIA, Petit, ex Linn. Gen. ed. I. 337 (1737).
ISNARDIA, Linn. Gen. ed. I. 337 (1737).
NEMATOPYXIS, Miq. Fl. Ind. Bat. i. I. 600 (1855).
PRIEUREA, DC. Prod. iii. 58 (1828).
QUADRICOSTA, Dulac, Fl. Hautes-Pyr. 329 (1867).
TIPHOGETON, Ehrh. Beitr. iv. 146 (1789).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 274 = prostrata.
alata, *Ell. Sketch*, i. 213.—Am. bor.
alternifolia, Burm. f. Fl. Ind. 36 = erigata.
alternifolia, *Linn. Sp. Pl.* 118.—Am. bor.
alternifolia, Walt. Fl. Carol. 89 = virgata.
angustifolia, Michx. Fl. Bor. Am. i. 88 = linearis.
angustifolia, Vent. ex DC. Prod. iii. 55, in syn. = Jussieua suffruticosa.
apetala, Walt. Fl. Carol. 89 = palustris.
arcuata, *Walt. l. c.*—Am. bor.
aurantiaca, Rafin. in Med. Repos. N. York, v. (1808) 358 = alternifolia.
brachycarpa, DC. Prod. iii. 55, in obs. = Jussieua suffruticosa.
capitata, *Michx. Fl. Bor. Am.* i. 90.—Am. bor.
Cercodia, Steud. Nom. ed. II. ii. 76 = Haloragis alata.
Curtissii, *Chapm. Fl. S. U. St. Suppl.* 621.—Florida.

LUDWIGIA :—
cylindrica, *Ell. Sketch*, i. 213.—Am. bor.
decurrens, Walt. Fl. Carol. 89 = Jussieua decurrens.
diffusa, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 301 = prostrata.
epilobioides, *Maxim. Prim. Fl. Amur.* 104.—Mandshuria.
erigata, *Linn. Mant.* i. 40.—Ind. or.
fluitans, Scheele, in Linnaea, xxi. (1848) 580 = natans.
fruticulosa, Blume, Bijdr. 1133 = prostrata.
glandulosa, Pursh, Fl. Am. Sept. i. 111 = microcarpa.
glandulosa, Walt. Fl. Carol. 88 = cylindrica?
gracilis, *Miq. Fl. Ind. Bat.* i. 1. 629.—Sumatra.
hastata, Spreng. Syst. i. 446 = Ammannia latifolia.
heterophylla, Poir. Encyc. Suppl. iii. 512 = cylindrica.
hirsuta, Desr. in Lam. Encyc. iii. 614 = pilosa.
hirsuta, Pursh, Fl. Am. Sept. i. 110 = hirtella.
hirtella, *Rafin. in Med. Repos. N. York*, v. (1808) 358.—Am. bor.
irrigata, J. F. Gmel. Syst. 1579, sphalm. = erigata.
juncea, Rafin. Autik. Bot. 38 = virgata?
jussiaeoides, *Desr. in Lam. Encyc.* iii. 614.—Afr. trop.
jussiaeoides, Michx. Fl. Bor. Am. i. 89 = Jussieua decurrens.
jussiaeoides, Wall. Cat. n. 6335 = parviflora.
lanceolata, Ell. Sketch, i. 213.—Am. bor.
leucorhiza, *Blume, Bijdr.* 1133.—Java.
linearis, *Walt. Fl. Carol.* 89.—Am. bor.
linifolia, *Poir. Encyc. Suppl.* v. 513.—Am. bor.
lutea, Bosc, ex DC. Prod. iii. 60 = virgata.
lythroides, Blume, Bijdr. 1134 = parviflora.
macrocarpa, Michx. Fl. Bor. Am. i. 89 = alternifolia.
microcarpa, Link, Enum. Hort. Berol. 141 = alternifolia.
microcarpa, *Michx. Fl. Bor. Am.* i. 88.—Am. bor.
mollis, Michx. l. c. 90 = pilosa.
multiflora, Walp. Rep. ii. 75 = parviflora.
natans, *Ell. Sketch*, i. 581.—Texas.
nitida, Michx. Fl. Bor. Am. i. 87 = palustris.
octandra, Banks, ex Kon. in Kon. & Sims, Ann. Bot. i. (1805) 547 = Haloragis tetragyna.
oppositifolia, Linn. Syst. ed. XII. 125 = parviflora.
ovalis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 95.—Japon.
palustris, *Ell. Sketch*, i. 211.—Reg. bor. temp.
parviflora, *Roxb. Hort. Beng.* 11; *Fl. Ind.* i. 419.—Geront. trop.
pedunculosa, Michx. Fl. Bor. Am. i. 38 = arcuata.
perennis, Burm. f. Fl. Ind. 37 = Jussieua suffruticosa.
perennis, *Linn. Sp. Pl.* 119 (sp. dub.).—Ind. or.
permollis, Bart. Fl. Philad. i. 52 = hirtella.
pilosa, Ell. Sketch., i. 216 = hirtella.
pilosa, *Walt. Fl. Carol.* 89.—Am. bor.
polycarpa, *Short & Peter, ex Torr. & Gray, Fl. N. Am.* i. 525.—Am. bor.
prostrata, *Roxb. Hort. Beng.* 11; *Fl. Ind.* i. 420.—As. trop.
pruinosa, Rafin. Autik. Bot. 38 = alternifolia?
ramosa, *Willd. Enum. Hort. Berol.* 166.—Ind. or.
ramosissima, Roth, Catalect. fasc. iii. 24 = virgata.
ramosissima, Walt. Fl. Carol. 89 = alternifolia.
repens, Forst. Cat. Pl. N. Am. 22 = palustris.
repens, *Sw. Prod. Veg. Ind. Occ.* 33.—Reg. trop.
rudis, Walt. Fl. Carol. 89 = pilosa?
salicifolia, Poir. Encyc. Suppl. iii. 512 = alternifolia.
scabriuscula, Kellogg, in Proc. Calif. Acad. vii. 1. (1876) 78 = Ammannia latifolia.
spathulata, *Torr. & Gray, Fl. N. Am.* i. 526.—Florida.
sphaerocarpa, *Ell. Sketch*, i. 214.—Am. bor.
stagnatilis, *Hance, in Walp. Ann.* ii. 531.—China.
subtriflora, Steud. Nom. ed. I. 497 = erigata.
suffruticosa, Walt. Fl. Carol. 90 = capitata.
triflora, Desr. in Lam. Encyc. iii. 615 = erigata.
trifolia, *Burm. f. Fl. Ind.* 37.—Java.
tuberosa, Rafin. Ann. Nat. 15 = virgata?
uniflora, Rafin. in Med. Repos. N. York, v. (1808) 358 = Jussieua decurrens.
virgata, *Michx. Fl. Bor. Am.* i. 89.—Am. bor.
zeylanica, Pers. Syn. i. 145 = parviflora.

LUEDDEMANNIA, Reichb. f. in Bonplandia, ii. (1854) 281. *ORCHIDEAE* [Benth. & Hook. f. iii. 552].
Lehmanni, Reichb. f. in Gard. Chron. (1880) II. 685 = Cycnoches Lehmanni.
Pescatorei, *Reichb. f. in Bonplandia*, ii. (1854) 281.—N. Granat.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 109.—N. Granat.

LUEHEA, F. W. Schmidt, Neue u. Selt. Pfl. 23 (1793) = **Stilbe**, Berg. (Verbenac.).
ericoides, F. W. Schmidt, l. c. = S. ericoides.

LUEHEA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 410. t. 5. *TILIACEAE*, Benth. & Hook. f. i. 235.
ALEGRIA, [Moç. & Sesse] ex DC. Prod. i. 516 (1824).
BROTERA, Vell. Fl. Flum. 322 (1825); vii. t. 163 (1827).
altheaeflora, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 58.—Am. austr.
candicans, *Mart. Nov. Gen. et Sp.* i. 102.—Bras.
candida, *Mart. l. c.*—Mexic.
cymulosa, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 58.—Bras.
densiflora, *A. St. Hil. Pl. Bras. Mer.* i. 294.—Bras.
divaricata, *Mart. Nov. Gen. et Sp.* i. 101.—Bras.
endopogon, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 225.—Venezuela.
ferruginea, *Turcz. l. c.* 224.—Venezuela.
grandiflora, *Mart. Nov. Gen. et Sp.* i. 99.—Bras.
laxiflora, *A. St. Hil. Fl. Bras. Mer.* i. 293.—Bras.
macrophylla, *Pohl, Pl. Bras. Ic.* ii. 127.—Bras.
mexicana, Spach, ex Steud. Nom. ed. II. ii. 77 = candida.
microphylla, *Pohl, Pl. Bras. Ic.* ii. 129.—Bras.
ochrophylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 50.—Bras.
paniculata, *Mart. Nov. Gen. et Sp.* i. 100.—Bras.
platypetala, *A. Rich. Ess. Fl. Cub.* 212.—Mexic.
rufescens, Benth. Bot. Voy. Sulph. 72 = platypetala.
rufescens, *A. St. Hil. Fl. Bras. Mer.* i. 293.—Bras.
rufescens, Seem. Bot. Voy. Herald, 86 = Seemanni.
scabrifolia, *Presl, Epim. Bot.* 212.—Mexic.
Seemannii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 348.—N. Granat.
speciosa, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 410.—Bras.
uniflora, *A. St. Hil. Fl. Bras. Mer.* i. 290.—Bras.
villosa, *Mart. Nov. Gen. et Sp.* i. 102.—Bras.

LUETKEA, Bong. Vég. Sitch. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 130. t. 2 = **Spiraea**, Linn. (Rosac.).
sibbaldioides, Bong. l. c. = S. pectinata.

LUFFA, [Tourn.] Linn. Syst. ed. I (1735); Cav. Ic. i. 7. t. 9 (1791). *CUCURBITACEAE*, Benth. & Hook. f. i. 823.
POPPYA, Neck. Elem. i. 241 (1790).
TREVOUXIA, Scop. Introd. 152 (1777).
TURIA, Forsk. Fl. Aegypt. Arab. 165 (1775).
acutangula, *Roxb. Hort. Beng.* 70; *Fl. Ind.* iii. 713.—Geront. trop.
aegyptiaca, *Mill. Gard. Dict.* ed. VIII.—Geront. trop.
amara, Roxb. Hort. Beng. 70; Fl. Ind. iii. 715 = acutangula.
amara, Wall. Cat. n. 6755 = Momordica tuberosa.
amara, Wall. l. c. n. 6755 B = Momordica Cymbalaria.
arabum, Steud. Nom. ed. I. 497, 536 = aegyptiaca.
Bendaul, Roxb. Hort. Beng. 104; Fl. Ind. iii. 717 = echinata.
Bondel, Buch.-Ham. ex Steud. Nom. ed. II. ii. 77 = praec.
caledonica, Sond. in Harv. & Sond. Fl. Cap. ii. 490 = Peponia caledonica.
Cattu-picinna, Ser. in DC. Prod. iii. 303 = aegyptiaca.
clavata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 714 = aegyptiaca.
cordata, Meissn. in M. Roem. Syn. Pepon. 67 = aegyptiaca.
cordifolia, Blume, Bijdr. 929 = Thladiantha cordifolia.
cylindrica, M. Roem. Syn. Pepon. 63; Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 119 = aegyptiaca.
drastica, Mart. Syst. Mat. Med. Bras. 81 = acutangula.
echinata, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 716.—Ind. or.; Afr. trop.
fluminensis, M. Roem. Syn. Pepon. 64 = acutangula.
foetida, Cav. Ic. i. 7. t. 9 = acutangula.
foetida, [Jacq. f.] Hort. Vindob. ex Schrad. in Linnaea, xii. (1838) 405 = aegyptiaca.
Gosa, Buch.-Ham. in Wall. Cat. n. 6753 = acutangula.
graveolens, *Roxb. Hort. Beng.* 70; *Fl. Ind.* iii. 716.—Ind. or.; Austral.
hederacea, Wall. Cat. n. 6755 = aegyptiaca.

LUFFA :—
Hilapikku, Herb. Madr. ex Wall. Cat. n. 6754 = Momordica tuberosa.
insularum, A. Gray, Bot. U. St. Expl. Exped. i. 644 = aegyptiaca.
Jacquinii, Schrad. in Linnaea, xii. (1838) 405 = aegyptiaca.
Kleinii, Wight & Arn. Prod. 344 = umbellata.
leiocarpa, F. Muell. Fragm. iii. 107 = aegyptiaca.
leucosperma, M. Roem. Syn. Pepon. 63 = aegyptiaca.
longistyla, Edgew. in Journ. As. Soc. Beng. (1852) 172 = echinata.
operculata, Cogn. in Mart. Fl. Bras. vi. IV. 12 = purgans.
Parvala, Buch.-Ham. in Wall. Cat. n. 6758 = aegyptiaca.
pentandra, Roxb. Hort. Beng. 70; Fl. Ind. iii. 712 = aegyptiaca.
Petola, Ser. in DC. Prod. iii. 303 = aegyptiaca.
Plukenetiana, Ser. l. c. 302 = acutangula.
purgans, *Mart. Syst. Mat. Med. Bras.* 81.—Am. trop.
quinquefolia, Seem. Bot. Voy. Herald, 285 = purgans.
racemosa, Roxb. Hort. Beng. 104; Fl. Ind. iii. 715 = aegyptiaca.
saccata, F. Muell. ex Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 124 = graveolens.
Satpatia, Buch.-Ham. in Wall. Cat. n. 6757 = aegyptiaca.
scabra, Schum. & Thonn. Beskr. Guin. Pl. 405 = aegyptiaca.
sphaerica, Sond. in Harv. & Sond. Fl. Cap. ii. 490 = Sphaerosicyos Meyeri.
striata, Schrad. in Linnaea, xi. (1837) Litt. 88 = aegyptiaca.
subangulata, Miq. Fl. Ind. Bat. i. 1. 667 = acutangula.
sylvestris, Miq. l. c. 666 = acutangula.
tuberosa, Roxb. Hort. Beng. 104; Fl. Ind. iii. 717 = Momordica Cymbalaria.
umbellata, *M. Roem. Syn. Pepon.* 63.—Ind. or.
variegata, *Cogn. in Bremen, Abh.* vii. (1882) 250.—Ins. Mascar.
Veitchii, Naud. in Rev. Hortic. (1873) 58 = aegyptiaca.
vittata, Zipp. ex Span. in Linnaea, xv. (1841) 206, nomen = aegyptiaca.

LUGAION, Rafin. Sylva Tellur. 22 (1838) = **Genista** Linn. (Legumin.).
angulatum, Rafin. l. c. = G. parviflora.
aphyllum, Rafin. l. c. = Eremosparton aphyllum.
etnense, Rafin. l. c. = G. aethnensis.
ferox, Rafin. l. c. 23 = G. ferox.
linifolium, Rafin. l. c. 22 = Cytisus linifolius.
multiflorum, Rafin. l. c. = Cytisus albus.
radiatum, Rafin. l. c. = G. radiata.
umbellatum, Rafin. l. c. = G. umbellata.

LUGOA, DC. Prod. vi. 14 (1837) = **Gonospermum**, Less. (Compos.).
revoluta, DC. l. c. = G. revolutum.

LUGONIA, Wedd. Chlor. And. ii. 49. t. 54 (1857). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 758.
andina, *Ball, in Journ. Linn. Soc.* xxii. (1885) 49.—Peruv.
lysimachioides, *Wedd. Chlor. And.* ii. 50.—Peruv.

LUHEA, DC. Prod. i. 517 (1824) = **Luehea**, Willd. (Tiliac.).

LUHEA, A. DC. in DC. Prod. xii. 606, in syn. (1848) = LUEHEA, F. W. Schmidt = **Stilbe**, Berg. (Verbenac.).

LUINA, Benth. in Hook. Ic. Pl. t. 1139 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 438.
hypoleuca, *Benth. l. c.*—Am. bor. occ.

LUISA, Endl. Gen. 199 (1837) = seq.

LUISIA, Gaudich. in Freyc. Voy. Bot. 426. t. 37 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 571.
BIRCHEA, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1838) 66. t. 10.
LOUISIA, Reichb. f. Xenia Orch. 204 (1858).
MESOCLASTES, Lindl. Gen. et Sp. Orch. 44 (1840).
alpina, Lindl. Bot. Reg. (1838) Misc. 56 = Vanda alpina.
antennifera, *Blume, Rumphia*, iv. 50.—Borneo.
bicaudata, *Thw. Enum. Pl. Zeyl.* 302.—Reg. Himal.; Zeylan.; China.

LUISIA:—
Birchea, Blume, Rumphia, iv. 50 = tenuifolia.
brachystachys, *Blume, l. c.*—Reg. Himal.
burmanica, Lindl. Fol. Orch. Luis. 3 = teretifolia.
confusa, *Reichb. f. in Walp. Ann.* vi. 621.—Ins. Amboin.
macrotis, *Reichb. f. in Gard. Chron.* (1869) 1110.—Reg. Himal.
microptera, *Reichb. f. l. c.* (1870) 1503.—Reg. Himal.
occidentalis, Lindl. Fol. Orch. Luis. 4 = teretifolia.
platyglossa, Reichb. f. in Walp. Ann. vi. 622 = teretifolia.
primulina, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 144.—Burma.
Psyche, *Reichb. f. in Bot. Zeit.* xxi. (1863) 98.—Burma.
tenuifolia, *Blume, Rumphia*, iv. 50.—Ind. or.
teres, *Blume, l. c.*—Japon.
teres, Lindl. Fol. Orch. Luis. 1 = Sarcanthus teretifolius.
teretifolia, *Gaudich. in Freyc. Voy. Bot.* 426. t. 37.—As. trop.; N. Caled.
trichorrhiza, *Blume, Rumphia*, iv. 50.—Reg. Himal.
uniflora, Blume, l. c. (Aëridis sp.).—Reg. Himal.
valida, *Reichb. f. in Bonplandia*, x. (1862) 335.—Ins. Philipp.
volucris, *Lindl. Fol. Orch. Luis.* 1.—Reg. Himal.
zeylanica, Lindl. l. c. 3 = teretifolia.
Zollingeri, *Reichb. f. in Walp. Ann.* vi. 622.—Java.

LUITKEA, Auct. ex Steud. Nom. ed. II. ii. 77 (1841) = LUETKEA, Bong. = **Spiraea**, Linn. (Rosac.).

LUMA, A. Gray, Bot. U. St. Expl. Exped. i. 535. t. 66 (1854) = **Myrtus**, Linn.
baeckeoides, Griseb. in Goett. Abh. vi. (1853–55) 120 = Eugenia parvifolia.
Bridgesii, A. Gray, ex Phil. f. Cat. Pl. Vasc. Chil. 76 (= *Myrceugenia Bridgesii*).—Chili.
Cheken, A. Gray, Bot. U. St. Expl. Exped. 536 (= sp. plur.).—Chili.
Chequen, Phil. f. Cat. Pl. Vasc. Chil. 76 = praec.
correaefolia, A. Gray, Bot. U. St. Expl. Exped. 542. t. 66 (= *Myrceugenia correaefolia*).—Chili.
Cruckshanksii, A. Gray, l. c. 540 = Temu Cruickshanksii.
ferruginea, A. Gray, l. c. 542.—Chili.
obtusa, A. Gray, l. c. 541 (= *Myrceugenia obtusa*).—Chili.
ovata, A. Gray, ex Phil. f. Cat. Pl. Vasc. Chil. 76 (= *Myrceugenia ovata*).—Chili.
planipes, A. Gray, Bot. U. St. Expl. Exped. 540, in textu = Eugenia planipes.
stenophylla, A. Gray, l. c. 540.—Chili.
Temu, A. Gray, l. c. 539 = Eugenia divaricata.

LUMANAJA, Blanco, Fl. Filip. ed. I. 821 (1837) = = **Homonoia**, Lour. (Euphorb.).
fluviatilis, Blanco, l. c. = H. riparia.

LUMBRICIDIA, Vell. Fl. Flum. 305; vii. t. 104 (1825) = **Andira**, Lam. (Legumin.).
anthelmia, Vell. l. c. = A. anthelminthica.
legalis, Vell. l. c. 305. t. 105 = A. stipulacea.

LUMNITZERA, Jacq. ex Spreng. Syst. ii. 687 (1825) = **Moschosma**, Reichb. (Labiat.).
caesia, Spreng. Syst. iv. Cur. Post. 223 = M. caesium.
capitata, Spreng. Syst. ii. 687 = Acrocephalus capitatus.
carnosa, Spreng. l. c. iv. Cur. Post. 223 = Ocimum carnosum.
densiflora, Spreng. Syst. ii. 687 = Plectranthus rugosus.
fastigiata, Spreng. l. c. = Salvia plebeia.
moschata, Spreng. l. c. = M. australe.
ocimoides, Jacq. ex Spreng. l. c. = M. polystachyum.
polystachya, Jacq. f. ex Spreng. l. c. = M. polystachyum.
prostrata, Spreng. l. c. = Geniosporum prostratum.
rubicunda, Spreng. l. c. iv. Cur. Post. 223 = Orthosiphon rubicundus.
salvioides, Spreng. l. c. ii. 687 = Ocimum salvioides.
tenuiflora, Spreng. l. c. = Ocimum sanctum.
virgata, Spreng. l. c. iv. Cur. Post. 223 = Orthosiphon rubicundus.

LUMNITZERA, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 186. *COMBRETACEAE*, Benth. & Hook. f. i. 687.
BRUGUIERA, Thou. Gen. Nov. Madag. 21 (1806).
FUNCKIA, Dennst. Schluess. Hort. Malab. 32 (1818).
PETALOMA, Roxb. Fl. Ind. ii. 372 (1832).
POKORNYA, Montr. in Mém. Acad. Lyon, x. (1860) 201.
PROBLASTES, Reinw. in Syll. Ratisb. ii. (1828) 10.
PYRRHANTHUS, Jack, in Malay. Misc. ii. (1822) n. VII. p. 57.
coccinea, *Wight & Arn. Prod.* ii. 316.—As. et Austral. trop.
littorea, Voigt, Hort. Suburb. Calc. 39 = coccinea.
lutea, *Presl, Rep. Bot.* i. 156, adnot.—Ins. Timor.
montana, F. Muell. Fragm. ii. 149 = Macropteranthis montana.
pedicellata, *Presl, Rel. Haenk.* ii. 23.—Ins. Maria.
pentandra, Griff. Notul. iv. 684 = coccinea.
purpurea, Presl, Rep. Bot. i. 155, adnot. = coccinea.
racemosa, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 187.—Geront. trop.
rosea, *Presl, Rep. Bot.* i. 155, adnot.—Ins. Philipp.

LUNANA, Endl. Gen. 1125 (1840), sphalm. = **Lunasia**, Blanco (Rutac.).

LUNANAEA, Endl. Gen. 994, in syn. (1840) = seq.

LUNANEA, DC. Prod. ii. 92 (1825) = **Cola**, Schott (Stercul.).
Bichy, DC. l. c. = C. acuminata.

LUNANIA, Hook. Lond. Journ. Bot. iii. (1844) 317. t. 11. *SAMYDACEAE*, Benth. & Hook. f. i. 797.
cubensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 332.—Cuba.
divaricata, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 90.—Cuba.
dodecandra, *Wright, ex Griseb. Cat. Pl. Cub.* 7.—Cuba.
Grayi, *Griseb. in Goett. Abh.* ix. (1861) 26.—Cuba.
parviflora, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 90.—Peruv.
racemosa, *Hook. Lond. Journ. Bot.* iii. (1844) 317.—Ind. occ.
Sauvalii, *Griseb. Cat. Pl. Cub.* 7.—Cuba.

LUNANIA, Rafin. Med. Fl. iii. 106 (1830) = **Heteranthera**, Ruiz & Pav. (Ponteder.).
uniflora, Rafin. l. c. = H. limosa.

LUNARIA, Tourn. ex Linn. Gen. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 71.
BRACHYPUS, Ledeb. Fl. Ross. i. 133 (1842).
aegyptiaca, Mill. Gard. Dict. ed. VIII. n. 3 = Ricotia cantoniensis.
alpina, Berger. Phyt. iii. 115; ex DC. Syst. ii. 282 = rediviva.
alysina, Steud. Nom. ed. II. ii. 77, sphalm. in syn. = praec.
annua, Jundz. Fl. Lith. 194 = rediviva.
annua, *Linn. Sp. Pl.* 653.—Europ.
argentea, All. Fl. Pedem. i. 245 = Alyssum argenteum.
biennis, Moench, Meth. 261 = annua.
canescens, Willd. Enum. Hort. Berol. 675 = Farsetia clypeata.
cantoniensis, Desv. Journ. Bot. iii. (1814) 174 = Ricotia cantoniensis.
clypeata, All. Fl. Pedem. i. 245 = Farsetia clypeata.
diffusa, Thunb. Prod. Pl. Cap. 107 = Heliophila diffusa.
elongata, Thunb. l. c. = Heliophila Thunbergii.
flagrans, Gilib. Fl. Lituan. ii. 48 = rediviva.
graeca, Willd. Enum. Hort. Berol. 675 = Farsetia lunarioides.
halimifolia, All. Fl. Pedem. i. 245 (*halymifolia*) = Alyssum halimifolium.
inodora, Lam. Fl. Fr. ii. 457 = annua.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 74 = Eutrema Wasabi.
lanceolata, Stokes, Bot. Mat. Med. iii. 442 = rediviva.
libyca, Viv. Pl. Lib. Spec. 34. t. 10. f. 1 = Alyssum libycum.
odorata, Lam. Fl. Fr. ii. 457 = rediviva.
ovalis, Stokes, Bot. Mat. Med. iii. 441 = annua.

LUNARIA:—
parviflora, Delile, Fl. Egypte, 248 = Savignya aegyptiaca.
perennis, C. C. Gmel. El. Bad. iii. 48 = rediviva.
perennis, Mill. Gard. Dict. ed. VIII. n. 4 = Farsetia lunarioides.
pinnata, Thunb. Prod. Pl. Cap. 107 = Heliophila diffusa.
rediviva, *Linn. Sp. Pl.* 653.—Europ.
Ricotia, Desv. Journ. Bot. iii. (1814) 174 = Ricotia cantoniensis.
Ricotia, Gaertn. Fruct. ii. 289 = rediviva.
scabra, Forsk. Fl. Aegypt. Arab. 117 = Farsetia aegyptia.
suffruticosa, Vent. Jard. Cels, 19 = Farsetia suffruticosa.
Tournefortii, Sibth. ex DC. Syst. ii. 288 = Farsetia lunarioides.

LUNASIA, Blanco, Pl. Filip. ed. I. 783 (1837). *RUTACEAE*, Benth. & Hook. f. i. 991.
LUNANA, Endl. Gen. 1125 (1840).
MYTILICOCCUS, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 173.
RABELAISIA, Planch. in Hook. Lond. Journ. Bot. iv. (1845) 519. t. 18.
amara, *Blanco, Fl. Filip.* ed. I. 783.—Ins. Philipp.
costulata, *Miq. Ann. Mus. Lugd. Bat.* iii. 89.—Ins. Sunda.
grandifolia, *Miq. l. c.*—Ins. Sunda.
parvifolia, *Miq. l. c.*—Ins. Sunda.

LUNDIA, DC. in Bibl. Univ. Genèv. xvii. (1838) 127 (Rev. Bign. 11). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1032.
CRATEROTECOMA, Mart. ex DC. Prod. ix. 215 (1845).
PHOENICOCISSUS, Mart. ex Meissn. Gen. Comm. 209 (1840).
acuminata, *Decne. in Rev. Hortic.* Sér. IV. ii. (1853) 221.—Bras.
Chica, Seem. Bot. Voy. Herald, 180 = Bignonia Chica.
cordata, *DC. Prod.* ix. 180.—Bras.
densiflora, *DC. l. c.*—Bras.
erionema, *DC. l. c.*—Bras.
glabra, *DC. l. c.* 180.—Bras.
hebantha, *DC. l. c.* 181.—Bras.
intermedia, DC. l. c. 225, nomen (Quid?).—Bras.
longa, *DC. l. c.* 180.—Bras.
multiflora, Mart. ex DC. l. c. 179 = Cuspidaria multiflora.
nitidula, *DC. l. c.* 181.—Bras.
obliqua, *Sond. in Linnaea*, xxii. (1849) 561.—Bras.
pauciflora, Mart. ex DC. Prod. ix. 179 = Cuspidaria pauciflora.
pubescens, *DC. l. c.* 180.—Bras.
rubicunda, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 196.—Bras.
Schomburgkii, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1158, nomen.—Guian.
Spruceana, *Bur. in Baill. Adansonia*, viii. (1867-68) 279.—Peruv.
umbrosa, *Bur. l. c.* 282.—Bras.; N. Granat.
virginalis, *DC. Prod.* ix. 181.—Bras.

LUNDIA, Puer. ex DC. Prod. ii. 64, in syn. (1825) = **Buchanania**, Roxb. (Anacard.).
mangiferoides, Puer. ex DC. l. c. = B. Lanzan.

LUNDIA, Schum. & Thonn. Beskr. Guin. Pl. 231 (1827) = **Oncoba**, Forsk. (Bixin.).
monacantha, Schum. & Thonn. l. c. = O. monacantha.

LUNGIA, Steud. Nom. ed. II. i. 752, in syn. (1840), sphalm. = LANGIA, Endl. = **Hermbstaedtia**, Reichb. (Amarant.).

LUNTIA, Neck. Elem. ii. 335 (1790) = **Croton**, Linn. (Euphorb.).
sericea, Rafin. Sylva Tellur. 62 = C. matourensis.

LUPINASTER, Buxb. ex Heist. Syst. 9 (1748) = **Trifolium**, Tourn. (Legum.).
africanus, Eckl. & Zeyl. Enum. 223 = T. africanum.
albens, Hort. Gorenk. ex Ledeb. Fl. Ross. i. 552 = T. Lupinaster.
albus, Link, Handb. ii. 174 = T. Lupinaster.

LUPINASTER:—
alpinus, Presl, Symb. Bot. i. 47 = T. alpinum.
amabilis, Presl, l. c. = T. amabile.
Burchellianus, Eckl. & Zeyh. Enum. 222 = T. africanum.
Buxbaumii, Presl, Symb. Bot. i. 47 = T. Buxbaumii.
chilensis, Presl, Rep. Bot. i. 4, in nota = T. chilensis.
circumdatus, Presl, Symb. Bot. i. 50, adnot. = T. circumdatum.
depauperatus, Presl, l. c. = T. depauperatum.
Ecklonianus, Presl, l. c. 47 = T. Burchellianum.
eximius, Presl, l. c. = T. eximium.
hirsutus, Presl, Bot. Bemerk. 52 = T. africanum.
Hookeri, Presl, Rep. Bot. i. 4, in nota = T. polymorphum.
involucratus, Presl, Symb. Bot. i. 47 = T. involucratum.
macrocephalus, Pursh, Fl. Am. Sept. ii. 479 = T. megacephalum.
ochreatus, Presl, Symb. Bot. i. 47 = T. ochreatum.
pentaphyllus, Moench, Meth. Suppl. 50 = T. Lupinaster.
purpurascens, Fisch. ex DC. Prod. ii. 204 = T. Lupinaster.
uniflorus, Presl, Symb. Bot. i. 47 = T. uniflorum.
Wormskioldii, Presl, l. c. = E. involucratum.

LUPINUS, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 480.
affinis, *Agardh, Syn. Gen. Lupin.* 20, partim.—Calif.
africanus, *Lour. Fl. Cochinch.* 429.—Afr. trop.
albescens, Hook. & Arn. in Hook. Bot. Misc. ii. (1833) 201 = multiflorus.
albicaulis, *Dougl. in Hook. Fl. Bor. Am.* i. 165.—Am. bor. occ.
albicaulis, Hook. Lond. Journ. Bot. vi. (1847) 216 = leucophyllus, sericeus.
albicaulis, Torr. ex Stansb. Rep. 386 = holosericeus.
albifrons, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 410 = Chamissonis.
albus, *Linn. Sp. Pl.* 721.—Europ. austr.
alopecuroides, *Desr. in Lam. Encyc.* iii. 626.—Peruv.
Andersoni, *S. Wats. Bot. King's Exp.* 58.—Calif.
andicola, *Gill. in Hook. Bot. Misc.* iii. (1833) 201.—Chili.
angustifolius, Blanco, Fl. Filip. ed. I. 566 = Zornia diphylla.
angustifolius, DC. Fl. Fr. iv. 507 = reticulatus.
angustifolius, *Linn. Sp. Pl.* 721.—Reg. Mediterr.
arboreus, *Sims, Bot. Mag.* t. 682.—Calif.
arbustus, Dougl. ex Lindl. Bot. Reg. 1230 = laxiflorus.
arcticus, *S. Wats. in Proc. Am. Acad.* viii. (1873) 526.—Am. bor. occ.
arenarius, *Gardn. in Hook. Ic. Pl.* t. 511.—Bras.
argenteus, Agardh, Syn. Gen. Lupin. 27 = ornatus.
argenteus, *Pursh, Fl. Am. Sept.* ii. 468.—Am. bor. occ.
argenteus, S. Wats. Bot. King's Exp. 55 = laxiflorus.
argyraeus, *DC. Prod.* ii. 408.—Hort. Teneriff.
aridus, *Dougl. in Lindl. Bot. Reg. t.* 1242.—Am. bor. occ.
aridus, S. Wats. Bot. King's Exp. 57 = Kingii.
arizonicus, *S. Wats. in Proc. Am. Acad.* xii. (1877) 250.—Arizona.
arvensis, *Benth. Pl. Hartw.* 128.—Peruv.
Aschenbornii, *S. Schau. in Linnaea*, xx. (1847) 739.—Mexic.
attenuatus, *Gardn. in Hook. Ic. Pl.* t. 511.—Bras.
aureonitens, *Gill. in Hook. Bot. Misc.* ii. (1833) 201.—Chili.
aureus, *Agardh, Syn. Gen. Lupin.* 8.—Peruv.
barbiger, *S. Wats. in Proc. Am. Acad.* viii. (1873) 528.—Am. bor. occ.
Barkeri, *Lindl. Bot. Reg.* (1839) t. 56.—Mexic.
bicolor, A. Gray, in Proc. Bost. Soc. Nat. Hist. vii. (1859–61) 146 = nanus.
bicolor, Lindl. Bot. Reg. t. 1109 = micranthus.
bilineatus, Benth. Pl. Hartw. 11 = Hartwegii.
bimaculatus, *Desr. in Lam. Encyc.* iii. 626.—Peruv.
bimaculatus, Hook. ex D. Don, in Sweet, Flow. Gard. Ser. II. t. 314 = subcarnosus.
Bivonii, Presl, Fl. Sicul. i. p. xxiv = L. Termis.
Blaschkeanus, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. Suppl. 57 = nootkatensis.
bogotensis, *Benth. Pl. Hartw.* 168.—N. Granat.
bracteolaris, *Desr. in Lam. Encyc.* iii. 622.—Reg. Argent.
brevicaulis, *Griseb. in Goett. Abh.* xxiv. (1879) 98.—Reg. Argent.

LUPINUS:—
brevicaulis, *S. Wats. Bot. King's Exp.* 53.—Am. bor. occ.
Breweri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 334.—Calif.
Bridgesii, A. Gray, ex S. Wats. l. c. viii. (1873) 538 = luteolus.
Burkei, *S. Wats. l. c.* 525.—Am. bor. occ.
caespitosus, *Benth. Pl. Hartw.* 169.—Peruv.
caespitosus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 379.—Am. bor. occ.
calcaratus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 195. f. 60.—Am. bor. occ.
californicus, Hort. ex C. Koch, Wochenschr. i. (1858) 80 = elegans.
campestris, *Cham. & Schlecht. in Linnaea*, v. (1830) 589.—Mexic.
canaliculatus, *Sweet, Brit. Flow. Gard.* Ser. I. iii. t. 283.—Reg. Argent.
canus, *Hemsl. Biol. Centr. Am. Bot.* i. 229.—Mexic.
caudatus, Kellogg, in Proc. Calif. Acad. ii. (1863) 197 = laxiflorus.
cervinus, Kellogg, l. c. 229. t. 73 = affinis.
Chamissonis, *Eschsch. in Mém. Acad. Pétersb.* x. (1826) 288.—Am. bor. occ.
chrysomelas, *Casar. Nov. Stirp. Bras. Dec.* 52.—Bras.
citrinus, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 93.—Calif.
Clarkei, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 1.—Costa Rica.
cochinchinensis, Lour. Fl. Cochinch. 429 = Crotalaria retusa.
comptus, *Mart. ex Benth. in Mart. Fl. Bras.* xv. 1. 14.—Bras.
concinnus, *Agardh, Syn. Gen. Lupin.* 6.—Am. bor. occ.
concinnus, S. Wats. Bibliog. Ind. N. Am. Bot. 237 = lepidus.
confertus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 192.—Calif.
Consentini, Walp. Rep. i. 596 = *Cosentini*.
coriaceus, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 430.—Bras.
Cosentini, Guss. Fl. Sic. Prod. ii. 398 = digitatus.
criticus, *Reichb. f. Ic. Fl. Germ.* xxii. 36.—Dalmatia.
crotalarioides, *Mart. ex Benth. in Mart. Fl. Bras.* xv. 1. 11.—Bras.
Cruckshanksii, *A. Gray, Bot. U. St. Expl. Exped.* i. 393.—Peruv.
Cruckshanksii, *Hook. Bot. Mag.* t. 3056.—Peruv.; Chili.
cytisoides, Agardh, Syn. Gen. Lupin. 18 = rivularis.
Danaus, A. Gray, in Proc. Am. Acad. vii. (1868) 335 = Lyallii.
decumbens, Torr. in Ann. Lyc. N. York, ii. (1828) 191 = argenteus.
decumbens, Torr. in Pacif. Rail. Rep. iv. 81 = laxiflorus?
decurrens, *Gardn. in Hook. Ic. Pl.* t. 521.—Bras.
densiflorus, Agardh, Syn. Gen. Lupin. 3 = microcarpus.
densiflorus, *Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 410.—Calif.
densiflorus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 379 = leucophyllus.
diffusus, *Nutt. Gen. Am.* ii. 93.—Am. bor.
digitatus, *Forsk. Fl. Aegypt. Arab.* 131.—Aegypt.
Douglasii, *Agardh, Syn. Gen. Lupin.* 34.—Calif.
Douglasii, Guss. Fl. Sic. Syn. ii. 852 in syn. = micranthus.
Douglasii, Torr. Bot. Mex. Bound. 57 = Chamissonis.
Ehrenbergii, *Schlecht. in Linnaea*, xii. (1832) 334.—Mexic.
elegans, *H. B. & K. Nov. Gen. et Sp.* vi. 477.—Mexic.
exaltatus, *Zucc. in Abh. Akad. Muench.* i. (1832) 345.—Mexic.
falcifer, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 378.—Am. bor.
flabellaris, *Bertol. Fl. Guatim.* 30.—Guatemala.
flexuosus, *Lindl. ex Agardh, Syn. Gen. Lupin.* 34.—Am. bor. occ.
flexuosus, S. Wats. Bot. King's Exp. 55, 56 = holosericeus, laxiflorus, lepidus, sericeus.
foliolosus, *Benth. Pl. Hartw.* 170.—Peruv.
foliosus, Hook. Lond. Journ. Bot. vi. (1847) 215 = laxiflorus.
foliosus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 377 = albicaulis.
Forskåhlei, Boiss. Diagn. Ser. I. ix. 10 = digitatus.

LUPINUS:—
fruticosus, Hort. ex Steud. Nom. ed. II. ii. 78 = arboreus.
glabellus, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 37.—Mexic.
glabratus, *Agardh, Syn. Gen. Lupin.* 24.—Hab. ?
gracilis, *Agardh, l. c.* 15.—Calif.
gracilis, Durand & Hilg. in Pacif. Rail. Rep. v. 7 = micranthus, sparsiflorus.
gracilis, Kunth, Mimos. 182. t. 52 = sarmentosus.
gracilis, Nutt. in Journ. Acad. Philad. vii. (1834) 115 = Nuttallii.
graecus, *Boiss. & Sprun. Diagn.* Ser. I. ii. 15.—Graecia.
grandifolius, Lindl. ex Agardh, Syn. Gen. Lupin. 18 = polyphyllus.
Grayii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 126.—Calif.
guadalupensis, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 184.—Calif.
guadiloupensis, Hort. ex Steud. Nom. ed. II. ii. 78 = Cyamopsis psoraleoides.
guatimalensis, Hort. l. c. = Cyamopsis psoraleoides.
Gussoneanus, Agardh, Syn. Gen. Lupin. 5 = hirsutus.
Hartwegii, *Lindl. Bot. Reg.* (1839) t. 31.—Mexic.
Havardi, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 369.—Am. bor.
hexaedrus, [Fourn. in] Bull. Soc. Bot. Fr. xiii. (1866) 214, lapsu = Lathyrus hexaedrus.
Hilarianus, *Benth. in Mart. Fl. Bras.* xv. 1. 15.—Bras.
hirsutissimus, *Benth. in Trans. Hort. Soc.* N. S. i. (1835) i. 411.—Calif.
hirsutissimus, A. Gray, in Ives's Rep. 9 = concinnus.
hirsutus, Eichw. Casp. Cauc. 23 = albus.
hirsutus, *Linn. Sp. Pl.* 721.—Reg. Mediterr.
hirsutus, Sieber, ex Guss. Fl. Sic. Prod. ii. 399 = digitatus.
hispanicus, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 17.—Reg. Mediterr.
holosericeus, A. Gray, in Proc. Am. Acad. viii. (1873) 379 = Chamissonis.
holosericeus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 380.—Am. bor. occ.
Hornemanni, *Agardh, Syn. Gen. Lupin.* 22.—Peruv.
humifusus, *Benth. Pl. Hartw.* 169.—N. Granat.
hybridus, Lem. Illustr. Hortic. v. (1858) t. 163 = *insignis*.
incanus, *R. Grah. in Edinb. N. Phil. Journ.* (Jan. 1834) 178.—Am. austr.
insignis ×, Lem. Illustr. Hortic. v. (1858) t. 163.
integrifolius, Desr. in Lam. Encyc. iii. 627 = villosus.
integrifolius, *Linn. Pl. Afr. Rar.* 17; *Amoen. Acad.* vi. 43.—Afr. austr.
interruptus, *Benth. Pl. Hartw.* 168.—N. Granat.
javanicus, *Burm. f. Fl. Ind.* 157 (Quid?).—Java.
Kingii, *S. Wats. in Proc. Am. Acad.* viii. (1873) 534.—Am. bor. occ.
Kunthii, *Agardh, Gen. Syn. Lupin.* 40.—Ecuador.
labiatus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 376 = rivularis.
lacteus, Kellogg, in Proc. Calif. Acad. v. (1875) 37 = microcarpus.
laevigatus, *Benth. in Mart. Fl. Bras.* xv. 1. 13.—Bras.
lanatus, *Benth. l. c.* 16.—Bras.
latifolius, Lindl. ex Agardh, Syn. Gen. Lupin. 18 = rivularis.
laxiflorus, *Dougl. in Lindl. Bot. Reg.* t. 1140.—Am. bor. occ.
laxiflorus, Hook. in Lond. Journ. Bot. vi. (1847) 215 = argenteus.
laxiflorus, Torr. in Pacif. Rail. Rep. iv. 81 = albicaulis.
laxiflorus, S. Wats. Bot. King's Exp. 55 = calcaratus.
laxifolius, A. Gray, in Proc. Am. Acad. viii. (1872) 379 = laxiflorus.
leonensis, *S. Wats. l. c.* xvii. (1881-82) 338.—Mexic.
lepidus, *Dougl. ex Lindl. Bot. Reg.* t. 1149.—Am. bor. occ.
leptocarpus, *Benth. Pl. Hartw.* 11.—Mexic.
leptophyllus, *Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 409.—Calif.
leptophyllus, Cham. & Schlecht. in Linnaea, v. (1830) 589 = Schiedeanus.
leucophyllus, *Dougl. ex Lindl. Bot. Reg.* t. 1124.—Am. bor. occ.
leucopsis, Agardh, Gen. Syn. Lupin. 29 = ornatus.
leucopsis, S. Wats. Bot. King's Exp. 56 = holosericeus.

LUPINUS:—
leucospermus, *Boiss. Diagn.* Ser. I. ix. 8.—Hispan.
Lindleyanus, *Agardh, Syn. Gen. Lupin.* 9.—Peruv.
linearis, *Desr. in Lam. Encyc.* iii. 625.—Reg. Argent.
linifolius, *Roth, Bot. Abh.* 14. t. 5 = reticulatus.
littoralis, *Dougl. ex Lindl. Bot. Reg.* t. 1198.—Am. bor. occ.
littoralis, Torr. Bot. Mex. Bound. 58 = Chamissonis.
lucidus, Benth. ex Loud. Hort. Brit. Suppl. ii. 647 = lepidus?
Ludovicianus, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 184.—Calif.
luteolus, *Kellogg, in Proc. Calif. Acad.* v. (1875) 38.—Calif.
luteus, *Linn. Sp. Pl.* 721.—Reg. Mediterr.
Lyalli, *A. Gray, in Proc. Am. Acad.* vii. (1868) 334.—Am. bor. occ.
macrocarpus, Hook. & Arn. Bot. Beech. Voy. 138 = arboreus.
macrocarpus, Torr. in Pacif. Rail. Rep. iv. 81 = Chamissonis.
macrophyllus, Benth. in Sweet, Brit. Flow. Gard. ii. Ser. II. iv. t. 356 = polyphyllus.
macrorhizos, Georgi, Beschr. Russ. Reich. III. iv. 1162 = nootkatensis.
maderensis, *Seem. Bot. Voy. Herald*, 278. t. 53.—Mexic.
Marschallianus, *Sweet, Brit. Flow. Gard.* Ser. I. ii. t. 139.—Mexic.
meionanthus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 522.—Am. bor. occ.
meionanthus, S. Wats. Bot. King's Exp. 56 = holosericeus.
Menziesii, Agardh, Syn. Gen. Lupin. 2 = densiflorus.
Menziesii, Torr. Bot. Mex. Bound. 57 = microcarpus.
mexicanus, *Cerv. in Lag. Gen. et Sp. Nov.* 22.—Mexic.
mexicanus, A. Gray, Pl. Wright. ii. 49 = Sitgreavesii.
mexicanus, Torr. in Sitgr. Rep. 158 = albicaulis.
micranthus, *Dougl. ex Lindl. Bot. Reg.* t. 1251.—Am. bor. occ.
micranthus, Guss. Fl. Sic. Prod. ii. 400 = hirsutus.
microcarpus, *Sims, Bot. Mag.* t. 2413.—Am. bor. occ.
microphyllus, *Desr. in Lam. Encyc.* iii. 624.—Peruv.
microphyllus, H. B. & K. Nov. Gen. et Sp. vi. 473 = Kunthii.
microphyllus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 372 = gracilis.
Milleri, Agardh, Syn. Gen. Lupin. 43 (Quid?).—Hab.?
minimus, *Dougl. in Hook. Fl. Bor. Am.* i. 163.—Am. bor. occ.
montanus, *H. B. & K. Nov. Gen. et Sp.* vi. 477.—Mexic.
Moritzianus, [Kunth] in Gartenfl. (1852) 10.—Cf. Vatke, Ind. Sem. Hort. Berol. (1876) App. 3 = elegans.
multiflorus, *Desr. in Lam. Encyc.* iii. 624.—Reg. Argent.
mutabilis, Lindl. Bot. Reg. t. 1539 = Cruckshanksii, *Hook.*
mutabilis, *Sweet, Brit. Flow. Gard.* Ser. I. ii. t. 130.—N. Granat.
nanus, *Dougl. ex Benth. in Trans. Hort. Soc.* N. S. i. (1835) 409.—Calif.
nitidissimus, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras.
niveus, *S. Wats. in Proc. Am. Acad.* xi. (1876) 114, 126.—Calif.
nootkatensis, *Donn, Cat. Hort. Cantab.* ed. III. 135; *Sims, Bot. Mag.* t. 1311.—Am. bor. occ.; As. bor. or.
nubigenus, *Kunth, Mimos.* 174. t. 50.—Peruv.
nutcanus, Spreng. Syst. iii. 227 = nootkatensis.
nuthatensis, Cooper, in Pacif. Rail. Rep. xii. 58 = littoralis.
Nuttallii, *S. Wats. in Proc. Am. Acad.* viii. (1873) 526.—Am. bor.
odoratus, Hort. ex DC. Prod. ii. 408 = luteus.
onustus, *S. Wats. in Proc. Am. Acad.* xi. (1876) 127.—Calif.
Orcuttii, *S. Wats. l. c.* xx. (1835) 359.—Calif.
ornatus, *Dougl. ex Lindl. Bot. Reg.* t. 1216.—Am. bor. occ.
ornatus, Newberry, in Pacif. Rail. Rep. vi. 70 = argenteus.
ornatus, Nutt. in Journ. Acad. Philad. vii. (1834) 20 = sericeus.
ornatus, Torr. in Pacif. Rail. Rep. iv. 81 = Chamissonis.
ovalifolius, *Benth. in Mart. Fl. Bras.* xv. i. 11.—Bras.
palaestinus, *Boiss. Diagn.* Ser. I. ix. 9.—Palestin.

LUPINUS:—
Palmeri, *S. Wats. in Proc. Am. Acad.* viii. (1873) 530.—Calif.
palustris, Kellogg, in Proc. Calif. Acad. v. (1875) 16 = microcarpus.
paniculatus, *Desr. in Lam. Encyc.* iii. 624.—Peruv.
parviflorus, *Nutt. ex Hook. & Arn. Bot. Beech. Voy.* 336.—Am. bor. occ.
parvifolius, *Gardn. in Hook. Ic. Pl.* t. 521.—Bras.
perennis, *Linn. Sp. Pl.* 721.—Am. bor.
perennis, Richards. Frankl. Journ. App. 17 = arcticus.
philistaeus, Boiss. Diagn. Ser. I. ix. 9 = reticulatus.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 392.—Peruv.
pilosissimus, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 37.—Mexic.
pilosus, *Murr. Syst.* ed. XIII. 545.—Europ. austr.
pilosus, Walt. Fl. Carol. 180 = villosus.
plattensis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 369.—Am. bor. occ.
plumosus, Dougl. ex Lindl. Bot. Reg. t. 1217 = leucophyllus.
polyphyllus, Ch. Anders. Cat. Pl. Nevad. 119 = Burkei.
polyphyllus, *Lindl. Bot. Reg.* t. 1096.—Calif.
prolifer, Desr. in Lam. Encyc. iii. 622 = L. Termis.
prostratus, Agardh, Syn. Gen. Lupin. 39.—Peruv.
pubescens, *Benth. Pl. Hartw.* 169.—N. Granat.
pulchellus, *Sweet, Brit. Flow. Gard.* Ser. II. t. 67.—Mexic.
pusillus, Ch. Anders. Cat. Pl. Nevad. 119, partim = brevicaulis.
pusillus, *Pursh, Fl. Am. Sept.* ii. 468.—Am. bor. occ.
pusillus, Torr. Bot. Mex. Bound. 57, partim. = concinnus.
ramosissimus, *Benth. Pl. Hartw.* 170.—N. Granat.
recurvatus, *Meyen, Reise*, i. 314.—Chili.
regius, Rudolph, ex Torr. & Gray, Fl. N. Am i. 376 = nootkatensis.
reticulatus, *Desv. in Ann. Sc. Nat.* Sér. II. iii. (1835) 100.—Europ.
rivularis, Agardh, Syn. Gen. Lupin. 24 = arboreus.
rivularis, *Dougl. ex Lindl. Bot. Reg.* t. 1595.—Am. bor. occ.
rivularis, Torr. in Pacif. Rail. Rep. iv. 81 = Chamissonis.
rupestris, *H. B. & K. Nov. Gen. et Sp.* vi. 626.—Peruv.
Sabinianus, Dougl. ex Lindl. Bot. Reg. t. 1435 = seq.
Sabinii, *Dougl. ex Hook. Fl. Bor. Am.* i. 166.—Am. bor. occ.
sarmentosus, *Desr. in Lam. Encyc.* iii. 474.—Peruv.
sativus, Gater. Desc. Pl. Montaub. 126 = albus.
Schiedeanus, *Steud. Nom.* ed. II. ii. 78.—Mexic.
sellulus, Kellogg, in Proc. Calif. Acad. v. (1875) 36 = confertus.
semiverticillatus, Desr. in Lam. Encyc. iii. 623 = varius.
semperflorens, *Hartw. ex Benth. Pl. Hartw.* 128.—Peruv.
sericatus, Kellogg, in Proc. Calif. Acad. vii. (1877) 92 = Chamissonis.
sericeus, Cooper, in Pacif. Rail. Rep. xii. 52 = ornatus.
sericeus, Eschsch. in Mém. Acad. Sc. Pétersb. (1826) 289 = arboreus.
sericeus, Hook. & Arn. Bot. Beech. Voy. 138 = Chamissonis.
sericeus, Nutt. in Journ. Acad. Philad. vii. (1834) 19 = Wyethii.
sericeus, *Pursh, Fl. Am. Sept.* ii. 468.—Am. bor. occ.
sericeus, Torr. & Gray, Fl. N. Am. i. 379, partim = sulphureus?
Sileri, S. Wats. in Proc. Am. Acad. x. (1875) 345 = Kingii.
Sitgreavesii, *S. Wats. l. c.* viii. (1873) 527.—Am. bor. occ.
Smithianus, *Kunth, Mimos.* 177. t. 51.—Ecuador.
sparsiflorus, *Benth. Pl. Hartw.* 303.—Calif.
sparsiflorus, Torr. in Pacif. Rail. Rep. iv. 81, partim = concinnus, hirsutissimus.
sparsiflorus, Torr. l. c. v. 360 = nanus.
stipulatus, *Agardh, Syn. Gen. Lupin.* 23.—Mexic.
Stiveri, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 192. f. 58.—Am. bor. occ.
stoloniferus, *Linn. Cent. Pl.* i. 23 (Quid?).—Sicil.; Arab.
subacaulis, *Griseb. in Goett. Abh.* xxiv. (1879) 98.—Reg. Argent.
subcarnosus, *Hook. Bot. Mag.* t. 3467.—Texas.
subsessilis, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 430.—Bras.
succulentus, Dougl. ex C. Koch, Ind. Sem. Hort. Berol. (1867) App. i. 11 = densiflorus.

LUPINUS:—
sulphureus, *Dougl. in Hook. Fl. Bor. Am.* i. 166.—Am. bor. occ.
sulphureus, S. Wats. Bot. King's Exp. 57 = calcaratus.
sylvaticus, *Hemsl. Biol. Centr. Am. Bot.* i. 231.—Mexic.
sylvestris, Lam. Fl. Fr. ii. 627 = varius.
Tauris, *Benth. Pl. Hartw.* 128.—Peruv.
tenellus, Dougl. ex G. Don, Gen. Syst. ii. 367 = argenteus.
Termis, *Forsk. Fl. Aegypt. Arab.* 131.—Reg. Mediterr. or.
texensis, Hook. Bot. Mag. t. 3492 = subcarnosus.
Thermis, Gasp. in Atti Acc. Nap. vi. (1851) 229. t. 10 = L. Termis.
thermus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 129 = praec.
tomentosus, *DC. Prod.* ii. 409.—Peruv.
tominensis, *Wedd. Chlor. And.* ii. 251.—Bolivia.
Torreyi, A. Gray, ex S. Wats. Bot. King's Exp. 58 = confertus.
trifidus, *Torr. ex S. Wats. in Proc. Am. Acad.* xii. (1877) 250.—Calif.
trifoliatus, Cav. Ic. i. 43 = Cyamopsis psoralioides.
trifoliatus, Rottl. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 223. t. 5 = Crotalaria trifoliastrum.
tristis, Sweet, Hort. Brit. ed. III. 191 = Acacia tristis.
trochophyllus, *Hoffmgg. Preis. Pfl.* viii. *Nachtr.* 45; *ex Walp. Rep.* i. 605.—Am. bor.
truncatus, *Nutt. ex Hook. & Arn. Bot. Beech. Voy.* 336.—Calif.
uncialis, *S. Wats. Bot. King's Exp.* 54. t. 7.—Am. bor. occ.
uncinatus, *Schlecht. in Linnaea*, xii. (1838) 333.—Mexic.
vaginans, *Benth. in Mart. Fl. Bras.* xv. i. 12.—Bras.
vaginatus, *Cham. & Schlecht. in Linnaea*, v. (1830) 590.—Mexic.
variegatus, Poir. Encyc. Suppl. iii. 520 = nootkatensis.
variicolor, Steud. Nom. ed. II. ii. 78 = littoralis.
varius, Gaertn. Fruct. ii. 324. t. 150 f. 4 = albus.
varius, *Linn. Sp. Pl.* 721.—Europ. austr.
varius, Savi, Fl. Pis. ii. 178 = = angustifolius.
varius, Urv. in Mém. Soc. Linn. Par. i. (1822) 342 = pilosus.
velutinus, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 430.—Bras.
venustus, *Hort. ex Vilm. Fl. Pl. Terre*, ed. I. 480.—Calif.
versicolor, Lindl. Bot. Reg. t. 1979 = littoralis.
versicolor, *Sweet, Brit. Flow. Gard.* Ser. II. t. 12.—Mexic.
villosus, *Willd. Sp. Pl.* iii. 1029.—Am. bor.
Wyethii, *S. Wats. in Proc. Am. Acad.* viii. (1873) 525.—Am. bor. occ.

LUPSIA, Neck. Elem. i. 71 (1790) = **Galactites**, Moench (Compos.).

LUPULARIA, Opiz, Seznam, 61 (1852) = **Medicago**, Tourn. (Legumin.).
parviflora, Opiz, l. c. = M. lupulina.

LUPULUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Humulus**, Linn. (Urtic.).
amarus, Gilib. Exercit. ii. 451 = H. Lupulus.
communis, Gaertn. Fruct. i. 358. t. 75 = H. Lupulus.
Humulus, Mill. Gard. Dict. ed. VIII. = H. Lupulus.
scandens, Lam. Fl. Fr. ii. 217 = H. Lupulus.

LUSCADIUM, Endl. Gen. Suppl. v. 94 (1850) = LASCADIUM, Rafin. ('nomen delendum').

LUSEKIA, Opiz, Seznam, 61 (1852) = **Salix**, Linn.
laurina, Opiz, l. c. = S. pentandra.

LUSSACIA, Spreng. Syst. ii. 294 (1825) = **Gaylussacia**, H. B. & K. (Vaccin.).
buxifolia, Spreng. l. c. = G. buxifolia.
hispida, Spreng. l. c. iv. Cur. Post. 160 = G. Pseudo-Gaultheria.
incana, Spreng. l. c. = G. thymelaeoides.
pinifolia, Spreng. l. c. = G. pinifolia.
Rhododendron, Spreng. l. c. = G. Rhododendron.
rugosa, Spreng. l. c. = G. rugosa.
salicifolia, Spreng. l. c. = G. salicifolia.
Vaccinium, Spreng. l. c. = G. brasiliensis.

LUSTRINIA, Rafin. Fl. Tellur. iv. 62 (1836) = **Dianthera**, Gronov. (Acanth.).
geniculata, Rafin. l. c. = D. secunda.

LUSURIAGA, Pers. Syn. i. 373 (1805) = **Luzuriaga**, Ruiz & Pav. (Liliac.).

LUTEOLA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 290 (1745) = **Reseda**, Tourn.
complicata, Webb, Iter Hisp. 67 = R. complicata.
glauca, Webb, Otia Hisp. 19 = R. glauca.
resedoides, Fuss, Fl. Transs. 86 = R. Luteola.
tinctoria, Webb & Berth. Phyt. Canar. i. 106 = R. Luteola.

LUTHERA, Sch. Bip. in Linnaea, x. (1836) 257 = **Krigia**, Schreb. (Compos.).
virginica, Sch. Bip. l. c. = K. amplexicaulis.

LUTKEA, Steud. Nom. ed. II. ii. 78 (1841) sphalm. = LUETKEA, Bong. = **Spiraea**, Linn. (Rosac.).

LUTROSTYLIS, G. Don, Gen. Syst. iv. 391 (1837) = **Ehretia**, Linn. (Boragin.).
inermis, G. Don, l. c. = E. fasciculata.
montevidensis, G. Don, l. c. = Citharoxylum barbinerve.
spinosa, G. Don, l. c. = E. spinosa.

LUUTIA, Neck. Elem. iii. 396 (1790) = **Croton**, Linn. (Euphorb.).

LUVUNGA, Buch.-Ham. in Wall. Cat. n. 6382 (1831). *RUTACEAE*, Benth. & Hook. f. i. 304.
LAVANGA, Meissn. Gen. 46; Comm. 34 (1837).
calophylla, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 69.—Ind. or.
eleutherandra, *Dalz. in Hook. Kew Journ.* ii. (1850) 258.—Ind. or.; Malaya.
Motleyi, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 44.—Borneo.
sarmentosa, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 69 = eleutherandra.
scandens, *Buch.-Ham. in Wight & Arn. Prod.* 90.—Ind. or.; Malaya.
scandens, Thw. Enum. Pl. Zeyl. 47 = eleutherandra.
tavoyana, Lindl. in Wall. Cat. n. 6383 = eleutherandra.

LUXEMBERGIA, A. St. Hil. in Mém. Mus. Par. ix. (1822) 352. *OCHNACEAE*, Benth. & Hook. f. i. 319.
PLECTANTHERA, Mart. Nov. Gen. et Sp. i. 39. t. 36 (1824).
angustifolia, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 596.—Bras.
ciliosa, *Gardn. in Hook. Ic. Pl.* t. 516.—Bras.
corymbosa, A. St. Hil. Pl. Rem. Brés. 335 = polyandra.
nobilis, *Eichl. ex Engl. in Mart. Fl. Bras.* xii. II. 360.—Am. trop.
octandra, *A. St. Hil. in Mém. Mus. Par.* ix. (1822) 352.—Bras.
polyandra, *A. St. Hil. l. c.*—Bras.
speciosa, *A. St. Hil. Pl. Rem. Brés.* 333.—Bras.

LUZIOLA, Juss. Gen. 33 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1115.
ARROZIA, Schrad. ex Kunth, Enum. Pl. i. 11 (1833).
CARYOCHLOA, Trin. Gram. Panic. Diss. ii. 54 (1833?).
alabamensis, *Chapm. Fl. S. U. St.* 584.—Am. bor.
brasiliana, Moric. Pl. Nouv. Am. 94 = peruviana.
caroliniana, Trin. ex Steud. Nom. ed. II. ii. 79 = Hydrochloa carolinensis?
longivalvis, Benth. & Hook. f. Gen. iii. 1116, sphalm. = seq.
longivalvula, *Doell, in Mart. Fl. Bras.* ii. II. 47.—Am. trop.
mexicana, H. B. & K. Nov. Gen. et Sp. i. 199 = peruviana.
micrantha, *Benth. in Journ. Linn. Soc.* xix. (1881) 55.—Bras.
peruviana, *J. F. Gmel. Syst.* 637.—Am. austr.
peruviana, G. F. W. Mey. ex Benth. in Journ. Linn. Soc. xix. (1881) 55 = Spruceana.

LUZIOLA:—
spiciformis, *Anderss. ex Bal. & Poitrass. in Bull. Soc. Hist. Nat. Toul.* xii. (1878) 232.—Paraguay.
Spruceana, *Benth. ex Doell, in Mart. Fl. Bras.* ii. II. 18.—Bras.
striata, *Bal. & Poitrass. in Bull. Soc. Hist. Nat. Toul.* xii. (1878) 231; *et in Bull. Soc. Bot. Fr.* xxxii. (1885) 243.—Paraguay.

LUZOLA, Sang. in Atti Acc. Pont. Lincei, Ser. I. vi. (1855) 609 = seq.

LUZULA, DC. Fl. Fr. iii. 158 (1805). *JUNCACEAE*, Benth. & Hook. f. iii. 868.
GYMNODES, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1809) 172.
JUNCOIDES, Adans. Fam. ii. 47 (1763).
LEUCOPHOBA, Ehrh. Beitr. iv. 148 (1789).
LUCIOLA, Sm. Engl. Fl. ii. 177 (1824).
NEMORINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 172.
abyssinica, Parl. Fl. Ital. ii. 310 = spicata.
africana, *Drège, Zwei Pfl. Docum.* 207, nomen; *et in Steud. Syn. Pl. Cyp.* 294.—Afr. austr.
albida, Bertol. Fl. Ital. iv. 210 = pedemontana.
albida, *DC. Fl. Fr.* iii. 159.—Europ.
Alopecurus, *Desv. Journ. Bot.* i. (1808) 159.—Am. merid.
Alopecurus, H. B. & K. Nov. Gen. et Sp. i. 238 = racemosa, peruviana.
Alopecurus, Poepp. ex Kunth, Enum. Pl. iii. 312 = campestris.
alpigena, Schur, Enum. Pl. Transs. 681 = albida.
alpina, Hoppe, ex Sturm, Deutschl. Fl. (1839) Heft 77 = campestris.
Althii, Herbich, Sel. Pl. Rar. Galic. et Bucov. (1836) 12 = campestris.
altissima, Turcz. ex Ledeb. Fl. Ross. iv. 217 = parviflora.
angustifolia, C. Koch, in Linnaea, xxi. (1848) 615 = campestris.
angustifolia, *Poir. Encyc. Suppl.* iii. 530.—Carolina.
angustifolia, Wender. ex Steud. Nom. ed. II. ii. 79; Garcke, Fl. Deutschl. ed. III. 348 = albida.
antarctica, *Hook. f. Fl. Antarct.* 550; *C. Gay, Fl. Chil.* vi. 139.—Reg. Magell.
arctica, Baer, in Bull. Sc. Acad. Pétersb. iii. (1838) 181 = arcuata.
arctica, Blytt, Norges Fl. i. (1861) 299 = hyperborea.
arcuata, Link, Hort. Berol. ii. 360 = hyperborea.
arcuata, [*Sw.*] *Summ. Syst. Veg. Scand.* 13.—Europ. bor.
arida, *Steud. Syn. Pl. Cyp.* 293 (Quid?).—Labrador.
aurea, Timb. in Bull. Soc. Phys. Sc. Nat. Toul. ii. (1874) 132 (Quid.?).—Gallia.
australasica, Steud. Syn. Pl. Cyp. 294 = campestris.
azorica, H. C. Wats. in Hook. Lond. Journ. Bot. ii. (1843) 408 = parviflora.
Banksiana, E. Mey. in Linnaea, xxii. (1849) 412 = picta.
barbata, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 45 = caricina.
Barrelieri, Pourr. ex Lange, ex Willk. & Lange, Prod. Fl. Hisp. i. 186 = Forsteri.
Berthelotii, Nees, ex Kunth, Enum. Pl. iii. 298 = purpurea.
boliviensis, *Buchen. in Bremen, Abh.* iv. (1874) 128.—Bolivia.
borealis, Fries, Summa Veg. Scand. 219 = Wahlenbergii.
Borreri, Bromf. ex Bab. Man. Bot. ed. III. 333 = vernalis.
brachyphylla, *Phil. in Linnaea*, xxxiii. (1864–65) 268.—Chili.
brevifolia, Desv. Journ. Bot. i. (1808) 152 = lactea.
brunnea, Froeb. ex Steud. Nom. ed. II. ii. 79 = spadicea.
caespitosa, *J. Gay, ex Steud. Syn. Pl. Cyp.* 294.—Hispan.
calabra, Tenore, Fl. Nap. iii. 386 = campestris.
campesiris, Willd. Enum. Hort. Berol. 394, err. typ. = campestris.
campestris, Bong. in Mém. Acad. Pétersb. ii. (1832) 167 = comosa.
campestris, *DC. Fl. Fr.* iii. 161.—Europ.; Am.; Austral.
campestris, Soland. in Phipps, Voy. 201 = hyperborea.

LUZULA:—
canariensis, *Poir. Encyc. Suppl.* iii. 532.—Ins. Canar.
capensis, Link, ex G. Don, in Loud. Hort. Brit. Suppl. iii. 582 = Juncus capensis?
capillaris, Steud. Syn. Pl. Cyp. 293 = comosa.
caricina, *E. Mey. in Linnaea*, xxii. (1849) 418.—Mexic.
Carolinae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 302.—Am. bor.
carpatica, Kit. in Linnaea, xxxii. (1863) 326 = spadicea.
chilensis, [Nees, in] Meyen, Reise, i. 349, nomen; et ex Kunth, Enum. Pl. iii. 312 = campestris.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 293.—N. Zel.
comosa, *E. Mey. Syn. Luz.* 21.—Am. bor. occ.
compacta, Desf. Tabl. ed. II. 25 = campestris?
confusa, *Lindeb. in Bot. Notiser*, 1855 (1856) 9.—Island.
congesta, DC. Fl. Fr. Suppl. 305 = campestris.
conglomerata, Mielich. ex Koch, Syn. Fl. Germ. ed. II. 848 = spicata.
crinita, *Hook. f. Fl. Antarct.* i. 84. t. 48.—Ins. Antarct.
cuprina, Rochel, ex Steud. Nom. ed. II. ii. 79 = albida.
decolor, Webb & Berth. Phyt. Canar. iii. 350. t. 235 = Forsteri.
denticulata, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 46 = gigantea.
Desvauxii, Kunth, Enum. Pl. iii. 304 = glabrata.
divaricata, S. Wats. in Proc. Am. Acad. xiv. (1879) 302 = parviflora.
effusa, *Buchen. Krit. Verz. Junc.* 53, 88.—Reg. Himal.
elegans, *Guthnick, ex Seub. Fl. Azor.* 23.—Ins. Azor.
elegans, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) 532 = purpurea.
erecta, Desv. Journ. Bot. i. (1808) 156 = campestris.
excelsa, *Buchen. in Bremen, Abh.* iv. (1874) 126.—Bolivia.
fastigiata, E. Mey. Syn. Luz. 9 = parviflora.
flavescens, Gaud. Agrost. ii. 239 = Hostii.
Forsteri, *DC. Syn. Pl. Fl. Gall.* (1806) 150.—Europ.; Afr. bor.
Forsteri, Lej. Fl. Spa, i. Suppl. 303 = vernalis.
fuscata, Schur, Reiseber. 132 = albida.
gigantea, *Desv. Journ. Bot.* i. (1808) 145.—Am. austr.
glabrata, Desv. l. c. 143. t. 5. f. 3.—Europ.
glabrata, Fries, Novit. Fl. Suec. Mant. ii. 182 = Wahlenbergii.
glomerata, Mielich. ex Hueben. in Flora, xxii. (1839) 489 = spicata.
gracilis, Welw. in Journ. Bot. x. (1872) 129 = purpurea.
graeca, Guss. Fl. Sic. Syn. ii. Add. 816 = maxima.
graeca, *Kunth, Enum. Pl.* iii. 310.—Graecia; Creta.
havaiiensis, *Buchen. Krit. Verz. Junc.* 54, 85.—Ins. Sandvic.
Hieronymi, *Buchen. & Griseb. in Goett. Abh.* xxiv. (1879) 318.—Reg. Argent.
Hostii, *Desv. Journ. Bot.* i. (1808) 140. t. 6. f. 1.—Europ.
humilis, Buchen. in Bremen, Abh. iv. (1874) 125 = racemosa.
hyperborea, Blytt, [Norges Fl. i. 298?] ex Bab. in Journ. Linn. Soc. x. (1868) 333 = confusa.
hyperborea, *R. Br. in Parry, 1st Voy. App.* 183.—Lappon.; Norweg.
intermedia, Baumg. Enum. Stirp. Transs. iii. 330 = nemorosa.
intermedia, Nocca & Balb. Fl. Ticin. i. 168 = sylvatica.
intermedia, Spenn. Fl. Friburg. i. 177; Schult. f. Syst. vii. 261 = campestris.
interrupta, Desv. Journ. Bot. (1808) i. 163 = racemosa.
involucratus, *Steud. ex Buchen. in Bremen, Abh.* iv. (1874) 121.—Bolivia.
italica, Parl. Fl. Ital. ii. 309 = spicata.
juncoides, Poir. Encyc. Suppl. iii. 535 = Juncus marginatus.
labradorica, *Steud. Syn. Pl. Cyp.* 291.—Labrador.
lactea, Laharpe, in Mém. Soc. Nat. Hist. Par. iii. (1827) 173 = pedemontana.

LUZULA:—
lactea, *E. Mey. Syn. Luz.* 15.—Lusit.; Corsic.; Italia.
laetevirens, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 46 = gigantea.
latifolia, Liebm. l. c. 47 = gigantea.
Leiboldi, *Buchen. in Bremen, Abh.* vi. (1879) 418.—Chili.
longiflora, *Benth. Fl. Austral.* vii. 123.—Austral.
lutea, *DC. Fl. Fr.* iii. 159.—Europ.
macrotricha, Steud. Syn. Pl. Cyp. 294 = spicata.
macusaniensis, *Steud. in Lechl. Berb. Am. Austr.* 56; *et ex Buchen. in Bremen, Abh.* iv. (1874) 181.—Peruv.
maxima, *DC. Fl. Fr.* iii. 160 (1805).—Europ.
maxima, Willd. Enum. Hort. Berol. 393 (1809) = sylvatica.
melanocarpa, Desv. Journ. Bot. i. (1808) 142. t. 5. f. 2 = parviflora.
multiflora, Lej. Fl. Spa, i. 169 = campestris.
neglecta, Kunth, Enum. Pl. iii. 301 = parviflora.
nemorosa, Baumg. Enum. Stirp. Transs. iii. 329 = campestris.
nemorosa, Dulac, Fl. Hautes-Pyr. 100 = Forsteri.
nemorosa, E. Mey. in Linnaea, xxi. (1848) 395 = albida.
nigricans, Desv. Journ. Bot. i. (1808) 158 = campestris.
nigricans, Hoppe, ex Steud. Nom. ed. II. ii. 79 = spicata.
nivalis, Laest. in Vet. Akad. Handl. Stockh. (1822) 334 = hyperborea.
nivea, Baumg. Enum. Stirp. Transs. iii. 328 = albida.
nivea, *DC. Fl. Fr.* iii. 158.—Alp.; Europ. med.
nodulosa, E. Mey. in Linnaea, xxii. (1849) 410 = graeca.
nutans, Duval-Jouve, in Bull. Soc. Bot. Fr. x. (1863) 80 = pediformis.
obtusata, Steud. Syn. Pl. Cyp. 294 = spicata.
Oldfieldii, *Hook. f. Fl. Tasm.* ii. 68.—N. Zel.; Austral.
pallescens, Bess. Enum. Pl. Volh. 15 = campestris.
pallescens, Hoppe, in Flora, ii. (1819) 185 = flavescens.
pallescens, *Schult. f. Syst.* vii. 272, 1665; *Fries, Summa Veg.* 220.—Lappon.
paniculata, *Desv. Journ. Bot.* i. (1808) 147.—Am. merid.
parviflora, *Desv. l. c.* 144.—Reg. bor.
parviflora, Kunth, Enum. Pl. iii. 300 = spadicea.
pauciflora, *Phil. in Linnaea*, xxxiii. (1864–65) 267.—Chili.
pedemontana, *Boiss. & Reut. Pugill. Pl. Nov.* 115.—Europ.
pediformis, Boiss. Voy. Bot. Espagne, ii. 625 = caespitosa.
pediformis, *DC. Fl. Fr.* iii. 162.—Europ.
peruviana, *Desv. Journ. Bot.* i. (1808) 160.—Peruv.
picta, Lesson & A. Rich. in Voy. Astrol. i. 146 = campestris.
pilosa, Baumg. Enum. Stirp. Transs. iii. 331 = campestris.
pilosa, Link, in Buch, Beschr. Canar. Ins. (1825) 140 = Forsteri.
pilosa, Willd. Enum. Hort. Berol. 393 = vernalis.
plumosa, *E. Mey. in Linnaea*, xxii. (1849) 387.—Ind. or.
psilophylla, *Phil. in Linnaea*, xxxiii. (1864–65) 267.—Chili.
pubescens, Schrank, in Flora, i. (1819) 446 = Hostii.
pumila, *Hook. f. Handb. N. Zeal. Fl.* 293.—N. Zel.
purpurea, *Link, in Buch, Beschr. Canar. Ins.* (1825) 140, 179; *E. Mey. in Linnaea*, xxii. (1849) 397.—Ins. Canar.
purpureo-splendens, Seub. Fl. Azor. 23. t. 4. f. 2 = elegans.
racemosa, *Desv. Journ. Bot.* i. (1808) 162.—Am. merid.
rigida, *Phil. in Linnaea*, xxxiii. (1864–65) 267.—Chili.
rubella, Hoppe, ex Gaud. Fl. Helv. ii. 566 = albida.
rufescens, *Fisch. ex E. Mey. in Linnaea*, xxii. (1849) 385.—Sibir.
Seuberti, Lowe, in Hook. Kew Journ. viii. (1856) 300 = elegans.
sicula, Parl. Nuov. Gen. e Spec. 59 = maxima.
Sieberi, Tausch, in Flora, xix. (1836) II. 423 = maxima.
silvatica, Auct. vide *sylvatica*.
spadicea, [Billot,] ex Nym. Consp. 751 = glabrata.

LUZULA:—
spadicea, *DC. Fl. Fr.* iii. 159.—Europ.
spicata, *DC. l. c.* 161.—Europ.; Am. bor.
spicata, Puccin. Syn. Fl. Luc. 198 = campestris.
stenophylla, Steud. Syn. Pl. Cyp. 294 = campestris.
stolonifera, Pourr. ex Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 188 = lactea.
sudetica, DC. Fl. Fr. Suppl. 306 = campestris.
sylvatica, Bichen. in Trans. Linn. Soc. xii. (1818) 331 = maxima.
tenella, Mielich. in Flora, xxii. (1839) 491 = campestris.
tristachya, Desv. Journ. Bot. i. (1808) 159. t. 4 = campestris.
velutina, Lange, in Kjoeb. Vidensk. Meddel. (1881) 93 = lactea.
vernalis, *DC. Fl. Fr.* iii. 160.—Europ.
vernalis, Sebast. & Mauri, Fl. Rom. Prod. 133 = Forsteri.
villosa, Wikstr. in Vet. Akad. Handl. Stockh. (1823) II. 276 = L. Alopecurus.
vulcanica, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 44 = racemosa.
vulgaris, Buchen. in Engl. Bot. Jahrb. vii. (1885) 175 = campestris.
Wahlenbergii, *Rupr. Beitr. Pfl. Russl.* ii. 58.—Europ.

LUZURIAGA, R. Br. Prod. 281 (1810) = **Geitonoplesium**, A. Cunn. (Liliac.).
cymosa, R. Br. l. c. 282 = G. cymosum.
montana, R. Br. l. c. = G. cymosum.

LUZURIAGA, Ruiz & Pav. Fl. Per. iii. 65. t. 298 (1802). *LILIACEAE*, Benth. & Hook. f. iii. 767.
CALLIXENE, Juss. Gen. 41 (1789).
ENARGEA, Banks, ex Gaertn. Fruct. i. 283. t. 59 (1788), nomen prius.
ENARTEA, Steud. Nom. ed. II. i. 553 (1840).
LUSURIAGA, Pers. Syn. i. 373 (1805).
angustifolia, Poir. Encyc. Suppl. iii. 536 = Eustrephus latifolius.
cordata, Bert. in Merc. Chil. 1829; ex Am. Journ. Sc. xxiii. (1833) 96 (Quid?).—Chili.
erecta, *Kunth, Enum. Pl.* v. 280.—Chili.
latifolia, Poir. Encyc. Suppl. iii. 535 = Eustrephus latifolius.
marginata, *Benth. & Hook. f. Gen.* iii. 768.—Chili.
parviflora, *Kunth, Enum. Pl.* v. 281.—N. Zel.
radicans, *Ruiz & Pav. Fl. Per.* iii. 66. t. 298.—Chili.

LYALLIA, Hook. f. Fl. Antarct. ii. 548. t. 122 (1846). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
kerguelensis, *Hook. f. l. c.*—Ins. Kerguelen.

LYCAPSUS, Phil. in Bot. Zeit. xxviii. (1870) 499. t. 8 = **Alomia**, H. B. & K. (Compos.).
tenuifolius, Phil. l. c.—Chili.

LYCASTE, Lindl. Bot. Reg. (1843), Misc. 14. *ORCHIDEAE*, Benth. & Hook. f. iii. 547.
COLAX, Lindl. Bot. Reg. (1843), Misc. 50.
DEPPIA, Rafin. Fl. Tellur. ii. 51 (1836).
PAPHINIA, Lindl. Bot. Reg. (1843), Misc. 14.
aciantha, Reichb. f. in Bonplandia, iii. (1855) 216 = Maxillaria aciantha.
acuminata, Reichb. f. l. c. = Maxillaria acuminata.
aromatica, *Lindl. Bot. Reg.* (1843), *Misc.* 16.—Mexic.
balsamea, *A. Rich. ex Reichb. f. in Walp. Ann.* vi. 602.—Guatemala.
barbifrons, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 383.—Peruv.
Barringtoniae, Hook. f. Bot. Mag. t. 5706 = costata.
Barringtoniae, *Lindl. Bot. Reg.* (1844), *Misc.* 42.—Ind. occ.
brevispatha, Klotzsch, ex Reichb. f. in Walp. Ann. vi. 604 = candida.
candida, *Lindl. in Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 37.—Am. centr.
chrysoptera, *Morr. in Ann. Soc. Bot. Gand*, v. (1849) 7. t. 232.—Mexic.
ciliata, *Lindl. ex Reichb. f. in Bonplandia*, iv. (1856) 324.—Cuba.
cochleata, *Lindl. in Lindl. & Paxt. Flow. Gard.* i. (1850–51) 126.—Mexic.

LYCASTE:—
consobrina, *Reichb. f. in Bot. Zeit.* x. (1852) 669.—Mexic.
costata, *Lindl. Bot. Reg.* (1843), *Misc.* 15.—Peruv.
crinita, *Lindl. l. c.* (1844), *Misc.* 39.—Mexic.
cristata, *Nichols. Dict. Gard.* ii. 304.—Ins. Trinit.
crocea, Linden, Orch. Linden. 21, in nota = fulvescens.
cruenta, *Lindl. Bot. Reg.* (1843), *Misc.* 16.—Guatemala.
Denningiana, *Reichb. f. in Gard. Chron.* (1876) II. 808.—Ecuador.
Denningsiana, *Reichb. f. Otia Bot. Hamb.* 7, 24 (= praec?).—Hab.?
Deppei, *Lindl. Bot. Reg.* (1843), *Misc.* 15.—Mexic.
Dowiana, *Endres & Reichb. f. in Gard. Chron.* (1874) II. 194.—Costa Rica.
fulvescens, *Hook. Bot. Mag.* t. 4193.—N. Granat.
gigantea, *Lindl. Bot. Reg.* (1843), *Misc.* 15; *et in Benth. Pl. Hartw.* 153.—Ecuador.
grandiflora, Beer, Prakt. Orch. 265 = Maxillaria grandiflora.
Harrisoniae, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1468 = Bifrenaria Harrisoniae.
jugosa, *Nichols. Dict. Gard.* ii. 304.—Bras.
lanipes, *Lindl. Bot. Reg.* (1843), *Misc.* 15.—Ecuador.
lasioglossa, *Reichb. f. in Gard. Chron.* (1872) 215.—Guatemala.
leiantha, Beer, Prakt. Orch. 263, 265 = Deppei?
leucantha, *Klotzsch, in Allg. Gartenz.* (1850) 402.—Costa Rica.
leucantha, Lindl. in Lindl. & Paxt. Flow. Gard. ii. (1851–52) 37 = candida.
Linguella, *Reichb. f. in Gard. Chron.* (1871) 738.—Peruv.?
Locusta, *Reichb. f. l. c.* (1879) I. 524.—Peruv.
macrobulbon, *Lindl. in Lindl. & Paxt. Flow. Gard.* i. (1850–51) 126.—N. Granat.
macrophylla, Lindl. Bot. Reg. (1842), Misc. 85, in obs.; (1843), Misc. 14 = plana.
mesochlaena, Reichb. f. & Warsc. in Bonplandia, ii. (1854) 98 = lanipes.
plana, *Lindl. Bot. Reg.* (1842), Misc. 85; (1843) t. 35.—Bolivia.
pleiochroma, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 179.—Hab.?
Puydtii, *Nichols. Dict. Gard.* ii. 304.—Bras.
Reichenbachii, *Gireoud, ex Reichb. f. in Bonplandia*, iv. (1856) 324.—Peruv.
Schilleriana, *Reichb. f. l. c.* iii. (1855) 215.—Hab.?
Skinneri, *Lindl. Bot. Reg.* (1843), *Misc.* 15.—Guatemala.
Smeeana ×, Reichb. f. in Gard. Chron. (1883) II. 198.
sulphurea ×, Reichb. f. l. c. (1882) I. 218.
tetragona, *Lindl. Bot. Reg.* (1843), *Misc.* 49.—Bras.
tricolor, *Reichb. f. in Walp. Ann.* vi. 603.—Guatemala.
tyrianthina, Loud. Hort. Brit. Suppl. iii. 582 = Bifrenaria tyrianthina.
Wittigi, *Reichb. f. in Gard. Chron.* (1878) II. 654.—Bras.
xytriophora, *Linden & Reichb. f. in Saund. Refug. Bot.* ii. t. 131.—Costa Rica.

LYCHNANTHOS, S. Gmel. in Nov. Comm. Acad. Petrop. xiv. (1770) I. 525 = **Cucubalus**, Tourn. (Caryoph.).
scandens, C. C. Gmel. Fl. Bad. ii. 249 = C. baccifer.
volubilis, S. Gmel. in Nov. Comm. Acad. Petrop. xiv. (1770) I. 525. t. 17. f. 1 = C. baccifer.

LYCHNIDEA, Burm. Rar. Afr. Pl. 138 (1737) = **Manulea**, Linn. (Scrophul.).
tomentosa, Moench, Meth. Suppl. 160 = M. tomentosa.

LYCHNIDIA, Pomel, Nouv. Mat. Fl. Atl. 207 (1874) = **Lychnis**, Tourn. (Caryoph.).
Coeli-rosa, Pomel, l. c. = L. Coeli-rosa.

LYCHNIS, [Tourn.] Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 147.
AGROSTEMMA, Linn. Syst. ed. I (1735) (Agrostema); et Gen. ed. I. 135 (1737).
CORONARIA, Linn. Gen. ed. I. 135 (1737).
EUDIANTHE, Reichb. Nom. 206 (1841).
FLOSCUCULI, Opiz, Seznam, 44 (1852).
FLOX, Adans. Fam. ii. p. (22) (1763).
GASTROLYCHNIS, Fenzl, ex Reichb. Nom. 206 (1841).

LYCHNIS:—
Githago, Adans. Fam. ii. 255 (1763).
Hedona, Lour. Fl. Cochinch. 286 (1790).
Liponeuron, Schott, Nym. & Kotschy, Analect. Bot. 55 (1854).
Lychnidia, Pomel, Nouv. Mat. Fl. Atl. 207 (1874).
Melandrium, Roehl. Deutschl. Fl. ed. II. 274 (1812).
Petrocoptis, A. Br. in Flora, xxvi. (1843) 370.
Physocarpon, Neck. Elem. ii. 164 (1790).
Physolychnis, Rupr. in Mém. Acad. Pétersb. Sér. VII. xiv. (1869) n. 4 (Sert. Tianschan.) 41.
Polyschemone, Schott, Nym. & Kotschy, Analect. Bot. 55 (1854).
Pontinia, Fries, in Lindb. Bot. Notiser (1843), 141.
Silenopsis, Willk. in Bot. Zeit. v. (1847) 237.
Steris, Adans. Fam. ii. 255 (1763).
Viscaria, Riv. ex Rupp. Fl. Jen. ed. Hall. 126 (1745).
Wahlbergella, Fries, Summa Veg. Scand. 155 (1846).
acaulis, Scop. Fl. Carn. ed. II. i. 306 = Silene acaulis.
affinis, *F. Vahl, ex Fries, Mant.* iii. 36.—Am. bor.
Agrostemma, Ledeb. Fl. Alt. ii. 184 = L. Githago.
ajanensis, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) ii. 564.—Sibiria.
ajanensis, S. Wats. Bot. King's Exp. 37 = Kingii?
alaschanica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 427.—Mongolia.
alba, *Mill. Gard. Dict.* ed. VIII. n. 4.—Europ.
alpestris, Linn. f. Suppl. 244 = Silene alpestris.
alpina, *Linn. Sp. Pl.* 436.—Reg. bor. et arct.
angustifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 648 = L. Viscaria.
apetala, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 405, partim = montana.
apetala, *Linn. Sp. Pl.* 437.—Reg. bor. et arct.
apetala, Turcz. ex Steud. Nom. ed. II. ii. 81 = pauciflora.
aragonica, Asso, ex Nym. Consp. 86 = pyrenaica.
Armoraria, Scop. Fl. Carn. ed. II. i. 310 = Silene Armeria.
arvensis, Gaertn. Mey. & Scherb, Fl. Wett. ii. 117 = dioica.
aspera, Poir. Encyc. Suppl. iii. 537 = L. Coeli-rosa.
atropurpurea, *Nym. Syll.* 229.—Banat.
baccifera, Scop. Fl. Carn. ed. II. i. 306 = Cucubalus baccifer.
Behen, Scop. l. c. 305 = Silene Cucubalus.
brachypetala, Fisch. Hort. Gorenk. ed. II. (1812) 60, nomen; Hort. Berol. ex Hornem. Hort. Hafn. Suppl. 51 = apetala.
brachypetala, *Schwaeg. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 47 (= praec.?).—Hab.?
Bungeana, *Fisch. ex Lindl. Bot. Reg.* t. 1864.—China.
californica, *S. Wats. in Proc. Am. Acad.* xii. (1877) 248.—Calif.
cancellata, *Jacquem. ex Edgew. & Hook. in f. Fl. Brit. Ind.* i. 226.—Reg. Himal.
cashmeriana, *Royle, ex Benth. in Royle, Illustr. Bot. Himal.* 80.—Reg. Himal.
cerastoides, Scop. Fl. Carn. ed. II. i. 309 = Silene cerastoides.
chalcedonica, *Linn. Sp. Pl.* 436.—Japon.
chilensis, *C. Gay, Fl. Chil.* i. 256.—Chili.
ciliata, Wall. Cat. n. 621 = nutans.
clavata, Moench, Meth. Suppl. 307 = L. Coeli-rosa.
coccinea, Salisb. Prod. 304 = grandiflora.
coccugosantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 71 = L. Flos-cuculi.
coelestis, St. Lag. l. c. 70 = L. Coeli-rosa.
Coeli-rosa, *Desr. in Lam. Encyc.* iii. 644.—Reg. Mediterr.
cognata, Maxim. Prim. Fl. Amur. 55 = fulgens.
conica, Scop. Fl. Carn. ed. II. i. 308 = Silene conica.
coriacea, Moench, Meth. 709 = coronaria.
coronaria, *Desr. in Lam. Encyc.* iii. 643.—Europ. austr.; As. Min.
coronaria, Franch. & Sav. Enum. Pl. Jap. i. 49, in syn. = seq.
coronata, Thunb. Fl. Jap. 187 = grandiflora.
corsica, *Loisel. Notul.* 73.—Corsica; Pyren.
cuneifolia, *Royle, Illustr. Bot. Himal.* 80.—Reg. Himal.
Cyrilli, Richt. ex Reichb. Ic. Fl. Germ. vi. 55 = L. Flos-cuculi.

LYCHNIS:—
decumbens, *Steud. Nom.* ed. II. ii. 80.—Hab.?
Densoni, *Steud. l. c.*—Hab.?
diclinis, *Lag. Gen. et Sp. Nov.* 15.—Hispan.
dioecia, Mill. Gard. Dict. ed. VIII. n. 3, errata = dioica.
dioica, Asso, Syn. Stirp. Arag. 57 = divaricata.
dioica, *Linn. Sp. Pl.* 436.—Europ.; As. bor.
diurna, Sibth. Fl. Oxon. 145 = dioica.
divaricata, *Reichb. Ic. Fl. Germ.* iv. 3.—Sicilia; Hispan.
Dorotheae, *Laest. ex Fries, Mant.* iii. 36; *et ex Nym. Consp.* 87.—Europ.
Drummondi, *S. Wats. Bot. King's Exp.* 37, 432.—Am. bor. occ.
elata, *S. Wats. in Proc. Am. Acad.* xii. (1877) 249.—Am. bor. occ.
eriostemon, Wall. Cat. n. 620 = indica.
Falconeri, Rohr, ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 223 = apetala.
fimbriata, Wall. Cat. n. 619 = indica.
Flos-cuculi, *Linn. Sp. Pl.* 436.—Europ.; As. bor.
Flos-Jovis, *Desr. in Lam. Encyc.* iii. 644.—Europ.
frigida, Schrank, in Denkschr. Bot. Ges. Regensb. i. ii. (1818) 25 = apetala.
fulgens, *Fisch. in Sims, Bot. Mag.* t. 2104; *et ex Spreng. Nov. Prov.* 26.—Sibir.
fulgida, Moench, Meth. 709 = chalcedonica.
Githago, *Scop. Fl. Carn.* ed. II. i. 310.—Europ.
glaberrima, Trevir. in Bot. Zeit. xix. (1861) 205 = praecox.
graminea, Molina, Sagg. Chile, ed. II. 289 = magellanica.
grandiflora, *Jacq. Coll.* i. 149; *Ic. Pl. Rar.* t. 84.—China; Japon.
Haageana, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 486 = Bungeana.
Haageana ×, Lem. Illustr. Hortic. (1859) t. 195.—Japon.
helvetica, G. Don, ex Loud. Hort. Brit. 186 = alpina.
himalayensis, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 223.—Reg. Himal.
inaperta, Scop. Fl. Carn. ed. II, i. 309 = Silene inaperta.
incana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = L. Flos-Jovis.
inclusa, Lehm. in Linnaea, v. (1830) Litt. 46 = apetala.
indica, *Benth. in Royle, Illustr. Bot. Himal.* 81.—Reg. Himal.
inflata, *Wall. Cat.* n. 618.—Reg. Himal.
Kingii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 247.—Am. bor. occ.
laciniata, Lam. Fl. Fr. iii. 51 = L. Flos-cuculi.
laciniata, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 429.—Manchuria.
laciniflora, Stokes, Bot. Mat. Med. ii. 562 = L. Flos-cuculi.
laeta, Ait. Hort. Kew. ed. I. ii. 118 = Silene laeta.
laeta, Moris, Fl. Sard. i. 242 = corsica.
laevis, Poir. Encyc. Suppl. iii. 537 = L. Coeli-rosa.
Lagascae, *Hook. f. Bot. Mag.* t. 5746.—Hispan.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 8 = L. Flos-cuculi.
macrocarpa, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 722.—Reg. Mediterr. occ.
macrorhiza, *Royle, ex Benth. in Royle, Illustr. Bot. Himal.* 80.—Reg. Himal.
madens, Jacquem. ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 223 = macrorhiza.
magellanica, Desr. in Lam. Encyc. iii. 641.—Reg. Magellan.
media, *Hort. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 40.—Hab.?
Miqueliana, *Rohrb. ex Franch. & Sav. Enum. Pl. Jap.* i. 49.—Japon.
montana, *S. Wats. in Proc. Am. Acad.* xii. (1877) 247.—Am. bor. occ.
multicaulis, *Wall. Cat.* n. 622.—Reg. Himal.
neglecta, G. Don, in Loud. Hort. Brit. 186 = L. Viscaria.
nemoralis, *Heuff. ex Reichb. Fl. Germ. Excurs.* 824; *et in Verh. Zool.-Bot. Ges. Wien*, viii. (1858) 73.—Hungar.
nigrescens, *Edgew. in Hook. f. in Fl. Brit. Ind.* i. 223.—Reg. Himal.
nivalis, *Kit. in Schult. Oesterr. Fl.* ed. II. i. 698.—Hungar.
noctiflora, Schreb. Spicil. 31 = Silene noctiflora.
nuda, *S. Wats. Bot. King's Exp.* 37.—Am. bor. occ.
nummularia, Lapeyr. Hist. Abr. Pl. Pyr. 263 = pyrenaica.

LYCHNIS:—
nutans, *Benth. in Royle, Illustr. Bot. Himal.* 80.—Reg. Himal.
nutans, Scop. Fl. Carn. ed. II. i. 310 = Silene nutans.
oculata, Backh. in Hook. Bot. Mag. t. 4075 = L. Coeli-rosa.
officinalis, Scop. Fl. Carn. ed. II. i. 303 = Saponaria officinalis.
orientalis, Scop. l. c. 304 = Saponaria orientalis.
Otites, Scop. l. c. 305 = Silene Otites.
palustris, Brot. Fl. Lusit. ii. 221 = Silene laeta.
Parryii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 248.—Am. bor. occ.
parviflora, Turcz. ex Bess. in Flora, xvii. (1834) i. Beibl. 8 = apetala.
pauciflora, Fisch. ex DC. Prod. i. 386 = apetala.
pauciflora, *Ledeb. in Mém. Acad. Pétersb.* v. (1814) 537.—Sibir. alt.
pilosa, *Edgew. in Trans. Linn. Soc.* xx. (1846) 34.—Reg. Himal.
plumaria, S. F. Gray, Nat. Arr. Brit. Pl. ii. 649 = L. Flos-cuculi.
praecox, *Trevir. in Bot. Zeit.* xix. (1861) 205.—Hispan.
pratensis, Spreng. Fl. Hal. 138 = dioica.
Preslii, *Sekera, in Lotos*, iii. (1853) 134; *et in Oestr. Bot. Wochenbl.* (1854) 197.—Bolivia.
procera, Ledeb. Fl. Ross. i. 298 = Gypsophila paniculata.
pulchra, Willd. ex Steud. Nom. ed. II. ii. 81 = Silene laciniata.
pumila, *Royle, ex Benth. in Royle, Illustr. Bot. Himal.* 80.—Reg. Himal.
Pumilio, Scop. Fl. Carn. ed. II. i. 304 = Silene Pumilio.
pusilla, Link, Enum. Hort. Berol. i. 432 = Silene quadridentata.
pusilla, Poir. Encyc. Suppl. iii. 537 = L. Coeli-rosa.
pyrenaica, *Berger. Fl. Bass. Pyren.* ii. 264.—Mont. Pyren.
quadridentata, Lucé, Fl. Osil. 145 = L. Flos-cuculi.
quadridentata, Murr. Syst. ed. XIII. 362 = Silene quadridentata.
quadrifida, Scop. Fl. Carn. ed. II. i. 307 = Silene alpestris.
rosea, Salisb. Prod. 304 = dioica.
rubra, Patze, Mey. & Elk. Fl. Preuss. 384 = dioica.
saxatilis, *Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 32.—Dahuria.
Saxifraga, Scop. Fl. Carn. ed. II. i. 308 = Silene Saxifraga.
scandens, Schur, Enum. Pl. Transs. 935, Index = Cucubalus baccifer.
segetum, Lam. Fl. Fr. iii. 50 = L. Githago.
Senno, *Sieb. & Zucc. Fl. Jap.* i. 98.—Japon.
sibirica, *Linn. Sp. Pl.* 437.—Sibir.
Sieboldii, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 486 = Bungeana.
Sieboldii, Van Houtte, Fl. des Serres, x. (1854–55) 31 = fulgens.
Siegeri, Steud. Nom. ed. II. ii. 81 = Silene nivalis.
Sigeriana, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 103 = praec.
silvestris, Rafn, Danm. Fl. 790 = dioica.
sordida, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 170.—Sibir. altaic.
speciosa, Carr. in Rev. Hortic. (1870–71) 530 = grandiflora?
splendens, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 524 = L. Viscaria.
stellarioides, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 371.—Japon.
Stewartii, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 224.—Reg. Himal.
suecica, Lodd. Bot. Cab. t. 881 = alpina.
sylvestris, Schk. Handb. i. 403. t. 124 = dioica.
terminalis, *C. Gay, Fl. Chil.* i. 255.—Chili.
thysanodes, *Hook. f. Fl. Antarct.* ii. 246.—Ecuador.
triflora, *R. Br. in Ross, Voy. App.* 142.—Am. arct.
triflora, Hornem. in Fl. Dan. t. 2173 = affinis.
tristis, *Bunge, in Ledeb. Fl. Alt.* ii. 184.—Sibir.
tristis, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 223 = nigrescens.
umbellifera, Lam. Fl. Fr. iii. 52 = L. Flos-Jovis.
uniflora, Ledeb. in Mém. Acad. Pétersb. v. (1814) 536 = tristis.
Vaccaria, Scop. Fl. Carn. ed. II. i. 303 = Saponaria Vaccaria.
variegata, *Desf. in Ann. Mus. Par.* xi. (1808) 442. t. 44.—Creta.

LYCHNIS:—
vespertina, Boiss. Voy. Bot. Espagne, ii. 95 = divaricata.
vespertina, Sibth. Fl. Oxon. 146 = dioica.
Viscaria, *Linn. Sp. Pl.* 436.—Europ.; As. bor.
viscosa, *Scop. Fl. Carn.* ed. II. i. 306.—Europ.
vulnerata, Scop. l. c. 309 = Silene gallica.
Wilfordi, Regel, ex Maxim. in Bull. Acad. Pétersb. xvii. (1872) 178, in textu = fulgens.

LYCHNITIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125 = **Verbascum**, Tourn. (Scrophul.).
alba, Fourr. l. c. = V. Lychnitis.
Boerhavii, Fourr. l. c. = V. Boerhavii.
lutea, Fourr. l. c. = V. Lychnitis.
nigra, Fourr. l. c. = V. nigrum.
pulverulenta, Fourr. l. c. = V. pulverulentum.
sinuata, Fourr. l. c. = V. sinuatum.

LYCHNOCEPHALIOPSIS, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 158 (1873) = **Lychnophora**, Mart. (Compos.).
suffruticulosa, Sch. Bip. ex Baker, l. c. = Lychnophora humillima.

LYCHNOCEPHALUS, Mart. ex DC. Prod. v. 83 (1836) = **Lychnophora**, Mart. (Compos.).
bicolor, Mart. ex DC. l. c. = Eremanthus bicolor.
tomentosus, Mart. ex DC. l. c. = Lychnophora tomentosa.

LYCHNODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 332. *SAPINDACEAE.*
reticulatus, *Radlk. l. c.* 333.—Afr. trop.

LYCHNOPHORA, Mart. in Denkschr. Bot. Ges. Regensb. ii. (1822) 148. tt. 4–10. *COMPOSITAE*, Benth. & Hook. f. ii. 235.
LYCHNOCEPHALIOPSIS, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 158 (1873).
LYCHNOCEPHALUS, Mart. ex DC. Prod. v. 83 (1836).
PIPTOCOMA, Less. in Linnaea, iv. (1829) 315.
affinis, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 233.—Bras.
albertinioides, *Gardn. l. c.* 234.—Bras.
Blanchetii, *Sch. Bip. in Flora*, xxxiii. (1850) 30.—Bras.
brunioides, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 149. t. 4.—Bras.
Candelabrum, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 345.—Bras.
cinerea, Sch. Bip. l. c. 358 = ericoides.
diosmaefolia, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 143 = Piptolepis ericoides.
ericoides, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 151. t. 5.—Bras.
Gardneri, Sch. Bip. in Pollichia, xx–xxi. (1863) 350 = staavioides.
hakeaefolia, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 156.—Bras.
humillima, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 371.—Bras.
lanigera, Pohl, ex Sch. Bip. l. c. 341 = Martiana.
Martiana, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 232.—Bras.
microphylla, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 354.—Bras.
Passerina, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 230 = Haplostephium Passerina.
penninervia, Sch. Bip. in Pollichia, xx–xxi. (1863) 405 = reticulata.
phylicaefolia, *DC. Prod.* v. 79.—Bras.
Pinaster, Mart. in Denkschr. Bot. Ges. Regensb. ii. (1822) 152. t. 6 = ericoides.
Piptocoma, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 155 = trichocarpha.
platyneura, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 357 = ericoides.
Pohlii, *Sch. Bip. l. c.* 353.—Bras.
proteaeformis, Mart. ex DC. Prod. v. 80 = ericoides.
ramosissima, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 232.—Bras.
reticulata, *Gardn. l. c.* 233.—Bras.
Riedelii, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 351.—Bras.
rosmarinifolia, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 155. t. 9.—Bras.
Rosmarinus, Pohl, ex Sch. Bip. in Pollichia, xx–xxi. (1863) 361 = brunioides.

LYCHNOPHORA:—
salicifolia, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 157. t. 10.—Bras.
Sellovii, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 372.—Bras.
staavioides, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 154. t. 8.—Bras.
subulata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 231 = Haplostephium Passerina.
syncephala, Sch. Bip. in Pollichia, xx–xxi. (1863) 404 = reticulata.
tomentosa, *Sch. Bip. l. c.* 369.—Bras.
trichocarpha, *Spreng. Syst.* iv. *Cur. Post.* 298.—Bras.
uniflora, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 347.—Bras.
villosissima, *Mart. in Denkschr. Bot. Ges. Regensb.* ii. (1822) 153. t. 7.—Bras.

LYCHNOPHORIOPSIS, Sch. Bip. in Pollichia, xx–xxi. (1863) 375. *COMPOSITAE*, Benth. & Hook. f. ii. 235.
heterotheca, *Sch. Bip. l. c.* 376.—Bras.

LYCIMNIA, Hance, in Walp. Ann. iii. 30 (1852) = **Melodinus**, Forst. (Apocyn.).
suaveolens, Hance, l. c. = M. suaveolens.

LYCIOPLESIUM, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 330 = **Acnistus**, Schott (Solanac.).
Boissieri, Dun. in DC. Prod. xiii. 493.—Venezuela.
Dombeyanum, Dun. l. c. 492.—Peruv.
fasciculatum, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 336 = A. fasciculatus.
horridum, Miers, l. c. iv. (1845) 331 = Lycium parvifolium.
Meyenianum, Miers, l. c. 332 = Grabowskia Meyeniana.
nitidum, Miers, l. c. 331.—N. Granat.
obovatum, Miers, l. c. 330 = Lycium obovatum.
ovatum, Miers, l. c. 331.—N. Granat.
puberulum, Phil. f. Cat. Pl. Vasc. Chil. 224 = Latua venenosa.
pubiflorum, Griseb. in Goett. Abh. vi. (1854) 128 = Latua venenosa.
pulchellum, Walp. Rep. vi. 618 = Lycium pulchellum.

LYCIOPSIS, Schweinf. Beitr. Fl. Aethiop. 37 (1867) = **Euphorbia**, Linn.
cuneata, Schweinf. l. c. = E. cuneata.

LYCIOPSIS, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 176 = **Fuchsia**, Linn. (Onagrar.).
thymifolia, Spach, l. c. = F. thymifolia.

LYCIOSERISSA, Roem. & Schult. Syst. iv. p. xxiv, 353 (1819) = **Plectronia**, Linn. (Rubiac.).
capensis, Roem. & Schult. l. c. 353 = P. ventosa.

LYCIUM, Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 900.
ASCLEIA, CANTALEA, EVOISTA, OPLUKION, PLICULA, TEREMIS, TRILIENA, Rafin. Fl. Tellur. 53–56 (1838).
JASMINOIDES, Medic. Phil. Bot. i. 134 (1789).
JOHNSONIA, Neck. Elem. ii. 49 (1790).
PANZERIA, J. F. Gmel. Syst. 247 (1791).
abeliaeflorum, Reichb. Ic. Fl. Germ. xx. 10. t. 5 = europaeum.
acnistoides, *Griseb. Cat. Pl. Cub.* 188.—Cuba.
acutifolium, *E. Mey. in Drège, Zwei Pfl. Docum.* 200, nomen; *et ex Dun. in DC. Prod.* xiii. I. 518.—Afr. austr.
afrum, Boiv. ex Dun. l. c. 515 = tenue.
afrum, Boj. Hort. Maurit. 241 = capense.
afrum, Drège, ex Dun. in DC. Prod. xiii. I. 518 = oxycarpum.
afrum, [Ker-Gawl.] Bot. Reg. t. 354 = carnosum.
afrum, *Linn. Sp. Pl.* 191.—Reg. Mediterr.; Afr. austr.
afrum, Mart. ex Dun. in DC. Prod. xiii. I. 524 = mediterraneum.
afrum, Reichb. ex Webb & Berth. Phyt. Canar. iii. 285 = barbarum.
afrum, Thunb. Prod. Fl. Cap. 37; et in Trans. Linn. Soc. ix. (1808) 153 = propinquum.
aggregatum, Ruiz & Pav. Fl. Per. ii. 45 = Acnistus aggregatus.
americanum, *Jacq. Stirp. Am.* 50.—Ins. S. Doming.
Andersonii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 388.—Am. bor. occ.
angustifolium, Mill. Gard. Dict. ed. VIII. n. 8 = tenue.

LYCIUM:—
apiculatum, *Dun. in DC. Prod.* xiii. I. 517.—Afr. austr.
arabicum, Schweinf. ex Boiss. Fl. Orient. iv. 289 = europaeum.
arborescens, Spreng. Syst. i. 701 = Acnistus aggregatus.
arenicolum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 14.—Am. austr.
armatum, Griff. Notul. iv. 95; Ic. Pl. Asiat. t. 415 = ruthenicum.
australe, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 20.—Austral.
austrinum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 13.—Afr. austr.
barbarum, Lam. Encyc. iii. 509 = chinense.
barbarum, *Linn. Sp. Pl.* 192.—Oriens.
barbarum, Linn. l. c. ed. II. 277 = vulgare.
barbarum, Mill. Gard. Dict. ed. VIII. n. 4 = Shawii.
barbatum, Thunb. Prod. Pl. Cap. 37 = Plectronia ventosa.
barbinodum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 138.—Mexic.
Berlandieri, *Dun. in DC. Prod.* xiii. I. 520.—Texas.
Berlandieri, A. Gray, in Proc. Am. Acad. vi. (1866) 47, partim = Andersonii.
boerhaviaefolium, Linn. f. Suppl. 150 = Grabowskia boerhaviaefolia.
brachyanthum, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 426.—Texas.
brevipes, *Benth. Bot. Voy. Sulph.* 40.—Calif.
brevistylum, *Todaro, Nuov. Gen. e Sp.* 74.—Hab.?
californicum, *Nutt. ex A. Gray, Bot. Calif.* i. 542.—Calif.
campanulatum, Drège, ex Dun. in DC. Prod. xiii. I. 523 = carnosum.
canescens, *Griff. Notul.* iv. 95.—Afghan.
canum, *Gill. ex Walp. Rep.* iii. 112, nomen.—Chili.
capense, Mill. Gard. Dict. ed. VIII. n. 7 = tetandrum.
capillare, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 188.—Am. austr.
capsulare, Linn. Cent. Pl. ii. 11; Amoen. Acad. iv. 308 = Hydrolea elegans.
carnosum, Hort. Monsp. ex Dun. in DC. Prod. xiii. I. 520 = Requieni.
carnosum, Hort. Tonelle, ex Dun. l. c. 510 = chinense.
carnosum, *Poir. Encyc. Suppl.* iii. 427.—Afr. austr.
carolinianum, *Walt. Fl. Carol.* 84.—Am. bor.
cestroides, *Schlecht. in Linnaea*, vii. (1832) 70.—Bras.
chilense, *Bert. in Merc. Chil.* (1829) n. xv. 693; *ex Colla, in Mem. Acc. Torin.* xxxviii. (1835) 133.—Chili.
chinense, *Mill. Gard. Dict.* ed. VIII. n. 5.—China.
ciliatum, *Schlecht. in Linnaea*, vii. (1832) 69.—Bras.
cinereum, Krauss, ex Dun. in DC. Prod. xiii. I. 517 = Kraussii.
cinereum, *Thunb. Prod. Pl. Cap.* 57.—Afr. austr.
cochinchinense, *Lour. Fl. Cochinch.* 134.—Cochinch.
confertum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 137.—Mexic.
Cooperi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 388.—Calif.
cordatum, *Mill. Gard. Dict.* ed. VIII. n. 10.—Afr. austr.
cornifolium, H. B. & K. Nov. Gen. et Sp. iii. 54 = Iochroma cornifolia.
crassifolium, Salisb. Prod. 135 = afrum.
depressum, Stocks, in Hook. Kew Journ. iv. (1852) 179 = barbarum.
deserti, *Phil. Fl. Atac.* 43.—Chili.
distichum, Meyen, Reise, i. 448 = Grabowskia disticha.
dumosum, Salisb. Prod. 136 = europaeum.
Dunalianum, Bubani, ex Dun. in DC. Prod. xiii. I. 509 = vulgare.
echinatum, *Dun. l. c.* 515.—Afr. austr.
Edgeworthii, Dun. l. c. 525 = barbarum.
elongatum, *Miers, in Ann. & Mag. Nat. Hist.* Sér. II. xiv. (1854) 136.—Chili.
elongatum, Salisb. Prod. 136 = chinense.
erosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 343.—Reg. Argent.
erythraeum, *Schweinf. ex Vatke, in Linnaea*, xliii. (1880–82) 333.—Afr. trop.
europaeum, Gouan, Hort. Monsp. 111 = flaccidum.
europaeum, Hort. Audib. ex Dun. in DC. Prod. xiii. I. 511 = subglobosum.
europaeum, *Linn. Sp. Pl.* 192.—Reg. Mediterr.; Oriens.
europaeum, Mérat, ex Dun. in DC. Prod. xiii. I. 510 = chinense.
europaeum, Schimp. ex Dun. l. c. 524 = mediterraneum.

LYCIUM:—
exsertum, *A. Gray, in Proc. Am. Acad.* xx. (1885) 305.—Am. bor. occ.
fagosum, Walp. Ann. v. 579, sphalm. = fragosum.
ferocissimum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 187.—Afr. austr.
filifolium, *Gill. ex Miers, l. c.* 336.—Reg. Argent.
flaccidum, *C. Koch, Dendrol.* ii. 1. 347.—Europ.; Afr. bor.
floribundum, *Dun. in DC. Prod.* xiii. 1. 513.—Am. austr.
floribundum, H. B. & K. Nov. Gen. et Sp. iii. 51 = Acnistus arborescens.
floridum, Salisb. Prod. 135 = barbarum.
foetidum, Linn. f. Suppl. 150 = Serissa foetida.
foliosum, Stocks, in Hook. Kew Journ. iv. (1852) 179 = barbarum.
fragosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 132.—Peruv.
Fremonti, *A. Gray, in Proc. Am. Acad.* vi. (1862) 46.—Calif.
fuchsioides, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 603 = Cestrum aurantiacum.
fuchsioides, Humb. & Bonpl. Pl. Aequin. i. 147 = Iochroma fuchsioides.
fuscum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 137.—Reg. Argent.
gelidum, *Wedd. Chlor. And.* ii. 108.—Chili.
gesnerioides, H. B. & K. Nov. Gen. et Sp. iii. 53 = Iochroma gesnerioides.
Gilliesianum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 344.—Reg. Argent.
glaucum, Miers, l. c. 17 = ruthenicum.
glaucum, *Phil. Fl. Atac.* 43.—Chili.
glomeratum, *Sendt. in Mart. Fl. Bras.* x. 154.—Bras.
gracile, Meyen, Reise, i. 380 = chilense.
gracilipes, *A. Gray, in Proc. Am. Acad.* xii. (1877) 81.—Arizona.
grandiflorum, Steud. Nom. ed. II. ii. 82 = seq.
grandifolium, Willd. ex Roem. & Schult. Syst. iv. 698 = Acnistus grandiflorus.
Grevilleanum, *Gill. ex Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 342.—Reg. Argent.
guayaquilense, H. B. & K. Nov. Gen. et Sp. iii. 50 = Acnistus guayaquilensis.
halimifolium, Mill. Gard. Dict. ed. VIII. n. 6 = vulgare.
halophyllum, *Welw. ex Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 11.—Lusitan.
heterophyllum, Murr. in Comm. Gotting. 1783 (1785) 6. t. 2 = Grabowskia boerhaviaefolia.
hirsutum, *Dun. in DC. Prod.* xiii. 1. 521.—Afr. austr.
horridum, H. B. & K. Nov. Gen. et Sp. iii. 52 = parvifolium.
horridum, *Phil. Fl. Atac.* 43.—Chili.
horridum, *Thunb. Prod. Pl. Cap.* 37.—Afr. austr.
humile, *Phil. Fl. Atac.* 43.—Chili.
ignarum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 194.—Am. austr.?
imbricatum, Walp. Rep. iii. 112, sphalm. = *intricatum*.
implexum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 133.—Chili.
indicum, Retz. Obs. ii. 12 = Serissa foetida.
indicum, Wight, Ic. iv. t. 1403 = europaeum.
inerme, Linn. f. Suppl. 150 = Plectronia ventosa.
infaustum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 192.—Am. austr.
intricatum, Boiss. Elench. 66 = europaeum.
italicum, Mill. Gard. Dict. ed. VIII. n. 2 = carnosum.
japonicum, Thunb. Fl. Jap. 93 = Serissa foetida.
Kraussii, *Dun. in DC. Prod.* xiii. 1. 517.—Afr. austr.
lanceolatum, Loisel. in Nouv. Duham. i. 123 = chinense.
leiostemum, *Wedd. Chlor. And.* ii. 108.—Peruv.
longiflorum, *Phil. in Anal. Univ. Chil.* (1862) II. 403.—Chili.
loxense, H. B. & K. Nov. Gen. et Sp. iii. 53 = Iochroma loxensis.
macranthum, Buching. ex Krauss, in Flora, xxvii. (1844) 829 = Rhigozum trichotomum.
macrodon, *A. Gray, in Proc. Am. Acad.* vi. (1862) 46.—Calif.
macrophyllum, Benth. Pl. Hartw. 49 = Acnistus Benthami.
Martii, *Sendt. in Mart. Fl. Bras.* x. 154.—Bras.
mediterraneum, *Dun. in DC. Prod.* xiii. 1. 523.—Reg. Mediterr.; Oriens.
megistocarpum, Dun. l. c. 510 = chinense.

LYCIUM:—
melanopotamicum, *Niederl. in Roca, Exped. Rio-Negro*, ii. 261.—Patag.
Meyenianum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 390 = Grabowskia Meyeniana.
microphyllum, *Loisel. in Nouv. Duham.* i. 112.—Afr. trop.
microphyllum, *Phil. Sert. Mend. Alt.* 39.—Chili.
microphyllum, Willd. ex Roem. & Schult. Syst. iv. 698 = parvifolium.
minutifolium, *Remy, in C. Gay, Fl. Chil.* v. 93.—Chili.
nodosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 139.—Am. austr.
nutans, Poepp. ex DC. 5me Not. Pl. Rar. Jard. Genèv. 24 = chilense.
obliquifolium, Stokes, Bot. Mat. Med. i. 418 = europaeum.
obovatum, Buching. ex Krauss, in Flora, xxvii. (1844) 830 = Ehretia capensis.
obovatum, *Ruiz & Pav. Fl. Per.* ii. 46.—Peruv.
obtusum, *Willd. ex Roem. & Schult. Syst.* iv. 698.—Venezuela.
oreophilum, *Wedd. Chlor. And.* ii. 108.—Peruv.
orientale, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xiv. (1854) 12 = europaeum.
ovale, Willd. ex Dun. in DC. Prod. xiii. 1. 527 = Iochroma cornifolia.
ovatum, Hort. Monsp. ex Dun. l. c. 506 (= *Fregirardia luteiflora*).—Am. austr.
ovatum, Loisel. in Nouv. Duham. i. 117 = chinense.
ovatum, Willd. ex Dun. in DC. Prod. xiii. 1. 527 = Iochroma cornifolia.
oxycarpum, *Dun. l. c.* 518.—Afr. austr.
oxycladum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 15.—Afr. austr.
pallidum, *Miers, l. c.* 131.—N. Mexic.
Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 292.—N. Mexic.
Parishii, *A. Gray, l. c.* xx. (1885) 305.—Calif.
parviflorum, *A. Gray, l. c.* vi. (1862) 48.—Arizona.
parvifolium, *Roem. & Schult. Syst.* iv. 698.—Am. austr.
patagonicum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 340.—Patagon.
pendulinum, *Miers, l. c.* 20.—Afr. austr.
Pendulum, Walp. Ann. vi. 1212 = pendulinum.
persicum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xiv. (1854) 12 = europaeum.
Pringlei, *A. Gray, in Proc. Am. Acad.* xx. (1885) 305.—N. Mexic.
propinquum, *G. Don, Gen. Syst.* iv. 459.—Afr. austr.
pruinosum, *Griseb. in Goett. Abh.* xxiv. (1879) 245.—Reg. Argent.
Prunus-spinosa, *Dun. in DC. Prod.* xiii. 1. 515.—Afr. austr.
puberulum, *A. Gray, in Proc. Am. Acad.* vi. (1862) 46.—N. Mexic.
pubescens, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 339.—Patagon.
pulchellum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 145.—Mexic.
quadrifidum, Moç. & Sesse, ex Dun. in DC. Prod. xiii. 1. 513 = Richii.
quitense, Hook. Ic. Pl. t. 723 = Poecilochroma quitensis.
rachidocladum, *Dun. in DC. Prod.* xiii. 1. 519.—Chili.
Requieni, *Dun. l. c.* 520.—Hab.?
Richii, *A. Gray, in Proc. Am. Acad.* vi. (1862) 46.—Calif.
rigidum, Thunb. Prod. Pl. Cap. 37 = carnosum.
roridum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 15.—Afr. austr.
ruthenicum, *Murr. in Comm. Gotting.* 1779 (1780) 9. t. 2.—Oriens.
saevum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xiv. (1854) 7 = europaeum.
salicifolium, Mill. Gard. Dict. ed. VIII. n. 3 = europaeum.
salsum, Bartr. Trav. ed. II. 57 = carolinianum.
salsum, *Ruiz & Pav. Fl. Per.* ii. 46.—Peruv.
sandwicense, *A. Gray, in Proc. Am. Acad.* vi. (1862) 44.—Ins. Sandvic.
scabrum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 389 = Grabowskia disticha.
Schaffneri, *A. Gray, ex Hemsl. in Biol. Centr. Am. Bot.* ii. 426.—Mexic.

LYCIUM:—
scoparium, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 340.—Reg. Argent.
senticosum, Miers, l. c. 138 = Berlandieri.
serpyllifolium, *Dun. in DC. Prod.* xiii. 1. 509.—Afr. austr.
Shawii, *Roem. & Schult. Syst.* iv. 693.—Afr. austr.
siculum, *Ucria, in Roem. Arch.* i. 1. 68.—Sicilia.
sinense, Auct. = chinense.
sinense, Hort. ex Miers, in Ann. & Mag. Nat. Hist. Ser. II. xiv. (1854) 194 = ignarum.
spathulatum, Mathews, ex Dun. in DC. Prod. xiii. 1. 527, 484 = Dunalia acnistoides.
spathulatum, Ruiz & Pav. Fl. Per. ii. 46 = Acnistus spathulatus.
spinosum, Hasselq. It. Palaest. 76 = europaeum.
spinosum, *Sieber, ex Steud. Nom.* ed. II. ii. 82, nomen.—Hab.?
spinulosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 191.—Reg. Argent.
stenophyllum, *Remy, in C. Gay, Fl. Chil.* v. 93.—Chili.
stolidum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xiv. (1854) 191 = Berlandieri.
subglobosum, *Dun. in DC. Prod.* xiii. 1. 511.—Hab.?
tataricum, Pall. Fl. Ross. i. 78. t. 49 = ruthenicum.
tenue, *Willd. Enum. Hort. Berol.* 245.—Afr. austr.
tenuispinum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 190.—Reg. Argent.
tetrandrum, *Thunb. Prod. Pl. Cap.* 37.—Afr. austr.
Thunbergii, G. Don, Gen. Syst. iv. 459 = Plectronia ventosa.
Torreyi, *A. Gray, in Proc. Am. Acad.* vi. (1866) 47.—Am. bor. occ.
Trewianum, Roem. & Schult. Syst. iv. 693 = chinense.
turbinatum, Loisel. in Nouv. Duham. i. 119 = chinense.
turcomannicum, Fisch. & Mey. ex Ledeb. Fl. Ross. iii. 191 = ruthenicum.
Tweedianum, *Griseb. in Goett. Abh.* xix. (1874) 216.—Reg. Argent.
umbellatum, Ruiz & Pav. Fl. Per. ii. 45 = Acnistus umbellatus.
umbrosum, H. B. & K. Nov. Gen. et Sp. iii. 54 = Iochroma umbrosa.
vimineum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. xiv. (1854) 140.—N. Granat.
vulgare, *Dun. in DC. Prod.* xiii. 1. 509.—Geront. temp.
Wilkesii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 229.—Patagon.

LYCOCTONUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 326 = **Aconitum**, Tourn. (Ranunc.).
sylvaticum, Fourr. l. c. = A. Lycoctonum.

LYCOMORMIUM, Reichb. f. in Bot. Zeit. x. (1852) 833. *ORCHIDEAE*, Benth. & Hook. f. iii. 545.
cerinum, *Benth. in Journ. Linn. Soc.* xviii. (1881) 321.—Am. centr.
serronianum, *Rodrig. Orch. Nov.* i. 102.—Bras.
squalidum, *Reichb. f. in Bot. Zeit.* x. (1852) 833.—N. Granat.

LYCOPERSICON, Tourn. ex Rupp. Fl. Jen. ed. Hall. 50 (1745) = seq.

LYCOPERSICUM, Hill, Veg. Syst. ix. 32 (1765). *SOLANACEAE*, Benth. & Hook. f. ii. 888.
AMATULA, Medic. Bot. Beobacht. 38 (1782); et Malv. 106 (1787).
aethiopicum, Mill. Gard. Dict. ed. VIII. n. 3 = Solanum aethiopicum.
agrimoniaefolium, *Dun. in DC. Prod.* xiii. 1. 24.—Peruv.
agrimoniaefolium, Hort. Berol. ex Dun. l. c. 27 = diadelphum.
atacamense, *Phil. Fl. Atac.* 42.—Chili.
cerasiforme, Dun. Solan. 113 = esculentum.
chilense, *Dun. in DC. Prod.* xiii. 1. 24.—Chili.
commutatum, Roem. & Schult. Syst. iv. 569 = peruvianum.
dentatum, Dun. Syn. Solan. 4 = peruvianum.
diadelphum, *Dun. in DC. Prod.* xiii. 1. 27.—Hab.?
Dulcamara, Medic. Beobacht. (1783) 245 = Solanum Dulcamara.
esculentum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. austr.

LYCOPERSICUM:—
Galeni, Mill. Gard. Dict. ed. VIII. n. 1 = esculentum.
hirsutum, *H. B. & K. Nov. Gen. et Sp.* iii. 18.—Am. austr.
Humboldtii, *Dun. Solan.* 112.—Bras.
inodorum, Juss. ex Steud. Nom. ed. I. 500 = pimpinellifolium.
macrophyllum, *Guss. Enum. Pl. Inar.* 230.—Afr. bor.
peruvianum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Peruv.
philippinarum, *Dun. in DC. Prod.* xiii. 1. 27.—Ins. Philipp.
pimpinellifolium, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. austr.
Pissisi, *Phil. in Anal. Univ. Chil.* (1861) 56; *et in Linnaea*, xxxiii. (1864-65) 205.—Chili.
Pomum-amoris, Moench, Meth. 515 = esculentum.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. austr.
pyriforme, *Dun. Solan.* 112. t. 26.—Am. austr.
racemiforme, Lange, in Bot. Tidsskr. Ser. II. i. (1872) 189 = racemigerum.
racemigerum, *Lange, Ind. Sem. Hort. Haun.* (1865) 26.—Am. bor.
regulare, Dun. Syn. Solan. 3 = peruvianum?
Solanum, Medic. Beobacht. (1783) 245 = esculentum.
Solanum-Lycopersicum, Hill, Hort. Kew. 148 = esculentum.
spurium, *Link, Handb.* i. 566.—Hab.?
tuberosum, Mill. Gard. Dict. ed. VIII. n. 7 = Solanum tuberosum.

LYCOPODIUM *arboreum*, Banks & Soland. ex Carr. Conif. ed. I. 489 = Dacrydium Colensoi.
arboreum, Jungh. in Fror. Notiz. (1842) 184; ex Blume, Rumphia, iii. 221 = Dacrydium elatum.

LYCOPSIS, Linn. Syn. ed. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 856.
BUGLOSSA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 351 (1821).
aegyptiaca, Linn. Sp. Pl. 138 = Anchusa flava.
alpestris, Lehm. Asperif. 260 = Nonnea alpestris.
ampulosus, Gilib. Fl. Lituan. i. 26 = Nonnea pulla.
arvensis, *Linn. Sp. Pl.* 139.—Europ.
axillaris, Rafin. New Fl. Am. iv. 19 (Quid?).—Am. bor.
bullata, Cyrill. Pl. Rar. Neap. Fasc. i. 34 t. 11. f. 3 = variegata.
calycina, Roem. & Schult. Syst. iv. 74 = Nonnea phaneranthera.
capensis, *Burm. f. Prod. Fl. Cap.* 4 (Quid?).—Afr. austr.
caspica, Lehm. Asperif. 256 = Nonnea picta.
ciliata, C. A. Mey. Verz. Pfl. Cauc. 98 = Nonnea pulla.
ciliata, Willd. Sp. Pl. i. 780 = Nonnea lutea.
Colsmanniana, Lehm. Asperif. ii. 265 = Nonnea Colsmanniana.
confertiflora, *Clarke, Trav.* ii. 420.—Palaestin.
crispa, Bertol. Fl. Ital. ii. 337 = Anchusa crispa.
cyrenaica, Spreng. Syst. i. 556 = Alkanna tinctoria.
decurrens, C. A. Mey. Verz. Pfl. Cauc. 98 = Nonnea decurrens.
echioides, Linn. Sp. Pl. ed. II. 199 = Macrotomia echioides.
elongata, Lehm. Asperif. ii. 264. n. 195 = Anchusa undulata.
erecta, Urv. in Mém. Soc. Linn. Par. i. (1822) 278 = orientalis.
flavescens, C. A. Mey. Verz. Pfl. Cauc. 98 = Alkanna lutea.
glomerata, Urv. in Mém. Soc. Linn. Par. i. (1822) 278 = Anchusa aggregata.
hispida, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 43. nomen.—Rossia.
longifolia, Drège, ex A. DC. in DC. Prod. x. 44 = Anchusa Dregei.
lutea, Desr. in Lam. Encyc. iii. 657 = Alkanna lutea.
macrophylla, Desr. l. c. = Anchusa macrophylla.
maroccana, *Willd. ex Roem. & Schult. Syst.* iv. 756 (Quid?).—Marocc.
micrantha, Ledeb. in Pand. Beitr. i. 65 = orientalis.
microphylla, Bertol. Misc. Bot. i. 15 = Nonnea ventricosa.
Milleri, Hornem. Hort. Hafn. i. 181 = Anchusa Milleri.
mollis, Vis. in Atti Riun. Scienz. Torin. 179 = Alkanna orientalis.
nigricans, Desr. in Lam. Encyc. iii. 655 = Nonnea nigricans.
obtusifolia, Willd. Sp. Pl. i. 780 = Nonnea obtusifolia.
orientalis, *Linn. Sp. Pl.* 139.—Europ. austr.; Oriens.
orientalis, Steph. Fl. Mosq. n. 122 = arvensis.
picta, Lehm. Asperif. 255 = Nonnea picta.

LYCOPSIS:—
pulchra, Willd. ex Roem. & Schult. Syst. iv. 756 = Macrotomia echioides.
pulla, Linn. Syst. ed. X. 916 = Nonnea pulla.
rosea, Lehm. Asperif. 261 = Nonnea rosea.
sanguinolenta, Herb. in Journ. Hort. Soc. ii. (1847) 17 = variegata?
saxatilis, Turcz. ex A. DC. in DC. Prod. x. 56, 103 = Onosma saxatilis.
setosa, Lehm. Asperif. 269 = Alkanna lutea.
Sibthorpiana, Lehm. l. c. 258 = Nonnea alba.
taurica, Ledeb. in Pand. Beitr. i. 64 = Nonnea pulla.
taurica, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) 1. 338 = orientalis.
taurica, Willd. ex Ledeb. Fl. Ross. iii. 110 = Nonnea picta.
undulata, Gilib. Fl. Lituan. i. 26 = arvensis.
variegata, *Linn. Sp. Pl.* 138.—Reg. Caucas.
versicolor, C. A. Mey. Verz. Pfl. Cauc. 98 = Nonnea rosea.
vesicaria, Auch. ex A. DC. in DC. Prod. x. 56, 98 = Alkanna tubulosa.
vesicaria, Echeand. ex Willk. & Lange, Prod. Fl. Hisp. ii. 491 = Nonnea alba.
vesicaria, Hornem. Hort. Hafn. i. 181 = Nonnea pulla.
vesicaria, Linn. Sp. Pl. 138 = Nonnea nigricans.
vesicaria, Pall. Voy. ed. Gall. vi. 110 = Craniospermum subvillosum.
virginica, Linn. Sp. Pl. 138 = Myosotis verna.

LYCOPUS, Tourn. ex Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1183.
alboroseus, Gilib. Fl. Lituan. i. 71 = europaeus.
albus, Fl. Corc. in Ionios Anthol. ii. (1834) 446 = europaeus.
americanus, Muhl. Cat. 3; Bart. Fl. Philad. i. 12 = sinuatus.
amethystinus, Stev. in Mém. Soc. Nat. Mosc. v. (1814) 341 = Amethystea caerulea.
angustifolius, Ell. Sketch, i. 26 = sinuatus.
aquaticus, Moench, Meth. 370 = europaeus.
arkansanus, Fresen. in Flora, xxv. (1842) 310, 311 = rubellus.
australis, *R. Br. Prod.* 500.—Austral.
bracteatus, Muhl. ex Link, Jahrb. i. III. 55 = sinuatus.
decrescens, C. Koch, in Linnaea, xxi. (1848) 646 = europaeus.
diantherus, Buch.-Ham. ex Roxb. Hort. Beng. 4; Fl. Ind. i. 144 = Mosla dianthera.
europaeus, *Linn. Sp. Pl.* 21.—Europ.; As. bor.
europaeus, Walt. Fl. Carol. 64 = sinuatus.
exaltatus, Ell. Sketch, i. 26 = sinuatus.
exaltatus, Linn. f. Suppl. 87 = europaeus.
heterophyllus, Rafin. Med. Fl. ii. 28 = sinuatus.
intermedius, Sweet, Hort. Brit. ed. I. 310 = europaeus.
laciniatus, Marz. Vicet. 17; ex Pollini, Fl. Veron. i. 27 = europaeus.
laciniatus, Rouy.—Cf. Bull. Soc. Bot. Fr. xxx (1883), Bibl. 32 = europaeus.
longifolius, Rafin. Med. Fl. ii. 28 = sinuatus.
lucidus, *Turcz. ex Benth. in DC. Prod.* xii. 179.—Sibir.; Am. bor. occ.
macrophyllus, Benth. Lab. Gen. et Sp. 185 = virginicus.
menthifolius, Mabille, Réch. Pl. Corse, fasc. i. 31 = europaeus.
mollis, *Kern. in Oestr. Bot. Zeitschr.* xvi. (1866) 371.—Europ.
niger, Gueldenst. Reisen, ii. 65 = europaeus.
obtusifolius, Benth. in DC. Prod. xii. 178 = lucidus?
obtusifolius, Michx. ex Vahl, Enum. i. 212 = rubellus?
palustris, Burm. f. Fl. Cap. Prod. 1; Lam. Fl. Fr. ii. 430 = europaeus.
parviflorus, *Maxim. Prim. Fl. Amur.* 216.—Reg. Amur.
pauciflorus, Rafin. Med. Fl. ii. 28 = sinuatus.
pensylvanicus, Muhl. ex Rafin. l. c. = sinuatus.
pinnatifidus, Pall. Reise, iii. 655 = europaeus.
praealtus, Nutt. ex Benth. in DC. Prod. xii. 179 = sinuatus.
pumilus, Vahl, Enum. i. 211 = virginicus.
riparius, Salisb. Prod. 72 = europaeus.
rubellus, *Moench, Meth. Suppl.* 146.—Am. bor.
sessilifolius, *A. Gray, in Proc. Am. Acad.* viii. (1870) 285.—Am. bor.
sinuatus, *Ell. Sketch*, i. 26.—Am. bor.
uniflorus, Michx. Fl. Bor. Am. i. 14 = virginicus.
virginicus, *Linn. Sp. Pl.* 21.—Am. bor.
vulgaris, Nutt. Gen. Am. i. 15 = sinuatus.
vulgaris, Pers. Syn. i. 24 = europaeus.

LYCORIS, Herb. App. 20 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 727.
OREXIS, Salisb. Gen. Pl. Fragm. 117 (1866).
PLEURASTIS, Rafin. Fl. Tellur. iv. 12 (1836).
africana, M. Roem. Syn. Ensat. 146 = aurea.
aurea, *Herb. App.* 20.—China.
hyacinthina, Herb. in Bot. Mag. sub t. 2113, p. 5 = Griffinia hyacinthina.
radiata, *Herb. App.* 20.—China.
radiata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 139 = Ungernia trisphaera?
sanguinea, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) 80.—Japon.
Severzowi, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) 1. 435 = Ungernia trisphaera.
squamigera, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) 79.—Japon.
straminea, *Lindl. in Journ. Hort. Soc.* iii. (1848) 76.—China.

LYCORUS, Loud. Hort. Brit. 121, in syn. (1830) = praec.

LYCOSERIS, Cass. in Dict. Sc. Nat. xxxiii. 474 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 495.
DIAZEUXIS, D. Don, in Trans. Linn. Soc. xvi. (1830) 251.
LANGSDORFIA, Willd. ex Less. Syn. Comp. 97 (1832).
bracteata, *Benth. Bot. Voy. Sulph.* 121.—Ecuador.
denticulata, *Less. in Linnaea*, v. (1830) 257.—Am. austr.
grandis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 111.—Costa Rica.
hyssopifolia, Cass. in Dict. Sc. Nat. xxxiii. 474 = Onoseris hyssopifolia.
latifolia, *Benth. Bot. Voy. Sulph.* 121.—Panama.
mexicana, *Cass. in Dict. Sc. Nat.* xxxiii. 474.—N. Granat.
squarrosa, *Benth. Bot. Voy. Sulph.* 121.—Panama.
triplinervia, *Less. in Linnaea*, v. (1830) 257.—N. Granat.

LYCOTIS, Hoffmgg. Verz. Pfl. Nachtr. ii. 99 (1826) = **Arctotis**, Linn. (Compos.).
leucanthemoides, Hoffmgg. l. c. = A. leucanthemoides.

LYCURUS, H. B. & K. Nov. Gen. et Sp. i. 141. t. 45 (1815). *GRAMINEAE*, Benth. & Hook. f. iii. 1145.
PLEOPOGON, Nutt. in Journ. Acad. Phil. Ser. II. i. (1847) 189.
alopecuroides, *Griseb. in Goett. Abh.* xix. (1874) 255.—Reg. Argent.
muticus, Spreng. Syst. iv. Cur. Post. 32 = Elionurus candidus.
phalaroides, *H. B. & K. Nov. Gen. et Sp.* i. 142.—Mexic.
phleoides, *H. B. & K. l. c.* t. 45.—Mexic.

LYDAEA, Molina, Sagg. Chile, ed. II. 300 (1810) = **Kageneckia**, Ruiz & Pav. (Rosac.).
Lyday, Molina, l. c. = K. oblonga.

LYDEA, Molina, Sagg. Chile, ed. II. 164 (1810) = praec.

LYGEUM, Linn. Cent. Pl. i. 4 (1755); Loefl. It. Hisp. 72 (1758). *GRAMINEAE*, Benth. & Hook. f. iii. 1110.
LINOSPARTON, Adans. Fam. ii. 34 (1763).
Spartum, *Loefl. ex Linn. Cent. Pl.* i. 4; *Amoen. Acad.* iv. 264.—Reg. Mediterr. occ.
spathaceum, Lam. Encyc. i. 96 = L. Spartum.
tenax, Salisb. Prod. 26 = L. Spartum.

LYGIA, Fasano, in Atti Acc. Nap. 1787 (1788) 245 (Ligia) = **Thymelaea**, Tourn.
Amani, Kotschy, ex Boiss. Fl. Orient. iv. 1052 = T. Aucheri.
Aucheri, Boiss. l. c. = T. Aucheri.
cilicica, Boiss. l. c. 1053 = T. cilicica.
multicaulis, Schott & Kotschy, ex Boiss. l. c. = T. cilicica.
Passerina, Fasano, in Atti Act. Nap. 1787 (1788) 247.—Cf. C. A. Mey. in Bull. Phys. Math. Acad. Pétersb. i. (1843) 358 = T. arvensis.
pubescens, C. A. Mey. l. c. = T. arvensis.

LYGINIA, R. Br. Prod. 248 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1029.
SCHOENODUM, Labill. Nov. Holl. Pl. ii. 79. t. 29 (1806).
barbata, *R. Br. Prod.* 248.—Austral.
imberbis, R. Br. l. c. = barbata.
montevidensis, *Spreng. Syst.* iv. *Cur. Post.* 26.—Reg. Argent.
symphyonema, F. Muell. Fragm. viii. 79 = barbata.
tenax, Steud. Nom. ed. I. 501 = barbata.

LYGISMA, Hook. f. Ic. Pl. t. 1423 (1883) *ASCLEPIADEAE*.
angustifolia, *Hook. f. l. c.*—Burma.

LYGISTUM, P. Br. Hist. Jamaic. 142 (1756) = **Manettia**, Linn. (Rubiac.).
axillare, Lam. Illustr. i. 286 = M. Lygistum.
spicatum, Lam. l. c. = Gonzalea spicata.

LYGODESMIA, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 311. *COMPOSITAE*, Benth. & Hook. f. ii. 530.
ERYTHREMIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 445.
aphylla, *DC. Prod.* vii. 198.—Am. bor.
exigua, *Benth. & Hook. f. Gen.* ii. 530; *A. Gray, in Proc. Am. Acad.* ix. (1874) 217.—Am. bor. occ.
grandiflora, *Torr. & Gray, Fl. N. Am.* ii. 485.—Am. bor. occ.
juncea, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–Mar. (1829) 311.—Am. bor. occ.
minor, Hook. Fl. Bor. Am. i. 295. t. 103 = Stephanomeria minor.
pumila, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 311 = aphylla.
rostrata, *A. Gray, in Proc. Am. Acad.* ix. (1874) 217.—Am. bor. occ.
spinosa, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 444.—Am. bor. occ.
triquetra, D. Don, in Edinb. N. Phil. Journ. (Jan.–Mar. 1829) 311 = Lactuca triquetra.

LYGODISODEA, Ruiz & Pav. Prod. 32. t. 5 (1794). *RUBIACEAE*, Benth. & Hook. f. ii. 134.
DISODEA, Pers. Syn. i. 210 (1805).
brasiliensis, *Gardn. in Hook. Journ. Bot.* ii. (1840) 25.—Bras.
ciliata, *Bartl. ex DC. Prod.* iv. 470.—Mexic.
foetida, *Ruiz & Pav. Fl. Per.* ii. 48. t. 188.—Peruv.
Lingua, Boj. ex Steud. Nom. ed. II. ii. 83 = Paederia Lingua.
mexicana, *DC. Prod.* iv. 470.—Mexic.

LYGOPLIS, Rafin. Sylva Tellur. 23 (1838) = **Genista**, Linn. (Legumin.).
horridum, Rafin. l. c. = G. horrida.
spinosum, Rafin. l. c. = Calycotome spinosa.
villosum, Rafin. l. c. = Calycotome villosa.

LYGOS, Adans. Fam. ii. 321 (1763) = **Genista**, Linn. (Legumin.).

LYGURUS, D. Dietr. Syn. Pl. i. 134 (1839), sphalm. = **Lycurus**, H. B. & K. (Gramin.).

LYGUSTRUM, Gilib. Fl. Lituan. i. 246 (1781) = **Ligustrum**, Tourn. (Oleac.).

LYMNOPHILA, Blume, Bijdr. 748 (1826) = **Limnophila**, R. Br. (Scrophul.).

LYNCEA, Cham. & Schlecht. in Linnaea, v. (1830) 108. = **Melasma**, Berg. (Scrophul.)
hispida, Cham. & Schlecht. l. c. 109 = M. hispidum.

LYNDENIA, Miq. Fl. Ind. Bat. i. 1. 580 (1855) = LIJNDENIA, Zoll. & Mor. = **Memecylon**, Linn. (Melastom.).

LYONELLA, Rafin. in Am. Monthly Mag. (1818) 266 = **Polygonum**, Linn.

LYONETIA, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 83 (1870), sphalm. = LYONNETIA, Cass. = **Anthemis**, Mich. (Compos.).

LYONETTIA, Endl. Ench. 241 (1841) = praec.

LYONIA, Ell. Sketch, i. 316 (1817) = **Vincetoxicum**, Moench (Asclep.).
maritima, Ell. l. c. = V. palustre.

LYONIA, Nutt. Gen. Am. i. 266 (1818). *ERICACEAE*, Benth. & Hook. f. ii. 587.
acuminata, C. Koch, Dendrol. ii. 1. 123 = Leucothoë acuminata.
arborea, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159 = Oxydendron arboreum.
axillaris, C. Koch, Dendrol. ii. 1. 124 = Leucothoë axillaris.
calyculata, Reichb. Fl. Germ. Excurs. 414 = Cassandra calyculata.
capreaefolia, Wats. Dendrol. t. 127 = paniculata.
Catesbaei, C. Koch, Dendrol. ii. 1. 125 = Leucothoë Catesbaei.
fasciculata, *G. Don, Gen. Syst.* iii. 831.—Jamaic.
ferruginea, *Nutt. Gen. Am.* i. 266.—Am. bor.
frondosa, Nutt. l. c. 267 = paniculata.
glandulosa, *Griseb. Cat. Pl. Cub.* 51.—Cuba.
jamaicensis, *D. Don, in N. Edinb. Phil. Journ.* xvii. (1834) 159, nomen.—Jamaic.
latifolia, *Griseb. Cat. Pl. Cub.* 51.—Ind. occ.
ligustrina, DC. Prod. vii. 599 = paniculata.
lucida, C. Koch, Dendrol. ii. 1. 118 = Pieris nitida.
marginata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159 = Pieris nitida.
Mariana, D. Don, l. c. = Pieris Mariana.
multiflora, Wats. Dendrol. t. 128 = paniculata.
myrtilloides, *Griseb. Cat. Pl. Cub.* 50.—Cuba.
obtusa, *Griseb. l. c.* 51. = Cuba.
octandra, *Griseb. Fl. Brit. W. Ind.* 142.—Jamaic.
paniculata, *Nutt. Gen. Am.* i. 266.—Am. bor.
parabolica, C. Koch, Dendrol. ii. 1. 119 = paniculata.
populifolia, C. Koch, l. c. 123 = Leucothoë acuminata.
pulverulenta, C. Koch, l. c. 117 = Zenobia speciosa.
racemosa, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159 = Leucothoë racemosa.
rhomboidalis, G. Don, Gen. Syst. iii. 831 = ferruginea.
rigida, Nutt. Gen. Am. i. 266 = ferruginea.
rubiginosa, *G. Don, Gen. Syst.* iii. 831.—Ind. occ.
salicifolia, Wats. Dendrol. t. 38 = paniculata.
squamulosa, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 542 = ferruginea.

LYONIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Polygonella**, Michx. (Polygon.).

LYONNETIA, Cass. in Dict. Sc. Nat. xxxiv. 106 (1825) = **Anthemis**, Mich. (Compos.).
abrotanifolia, Less. Syn. Comp. 259 = A. cretica.
abrotanifolia, Webb, Iter Hisp. 37 = A. nobilis.
anthemoides, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 83.—Hispan.
montana, Griseb. ex Boiss. & Reut. Diagn. Pl. Nov. Hisp. 17 = A. montana.
muricata, DC. Prod. vi. 15 = A. muricata.
pectinata, DC. l. c. vii. 295 = A. taygetea, pectinata.
persica, Boiss. Diagn. Ser. I. vi. 85 = A. persica.
pusilla, Cass. in Dict. Sc. Nat. xxxiv. 106 = A. cretica.
rigida, DC. Prod. vi. 14 = A. cretica.
tenuiloba, DC. l. c. 15 = A. montana.

LYONOTHAMNUS, A. Gray, in Proc. Am. Acad. xx. (1885) 291. *ROSACEAE*.
asplenifolius, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 187.—Calif.
floribundus, *A. Gray, in Proc. Am. Acad.* xx. (1885) 292.—Calif.

LYONSIA, R. Br. in Mem. Wern. Soc. i. (1809) 66. *APOCYNACEAE*, Benth. & Hook. f. ii. 711.
celebica, *Oliver, in Journ. Linn. Soc.* xv. (1876) 99.—Ins. Celebes.
cuspidata, Rafin. New Fl. Am. iv. 59 (Quid ?).—Am. bor.
diaphanophlebia, *F. Muell. ex Benth. Fl. Austral.* iv. 322.—Austral.
eucalyptifolia, *F. Muell. l. c.* 323.—Austral.
induplicata, *F. Muell. l. c.* 321.—Austral.
laevis, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 333.—Ins. Fiji.
Langiana, *F. Muell. Fragm.* vi. 128.—Austral.
largiflorens, *F. Muell. ex Benth. Fl. Austral.* iv. 322.—Austral.
latifolia, *Benth. l. c.* 323.—Austral.
Leichhardtii, F. Muell. Fragm. vi. 128 = Parsonsia Leichhardtii.

LYONSIA :—
lilacina, *F. Muell. ex Benth. Fl. Austral.* iv. 321.—Austral.
oblongifolia, *Benth. l. c.* 303.—Austral.
reticulata, *F. Muell. Rep. Burdek. Exped.* 16.—Austral.
scabra, *A. DC. in DC. Prod.* viii. 401.—N. Caled.
straminea, *R. Br. Prod.* 466.—Austral.

LYPERANTHUS, R. Br. Prod. 325 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 611.
FITZGERALDIA, F. Muell. in Wing, South Sc. Record (1882), 56, in obs.
antarcticus, *Hook. f. Fl. Antarct.* ii. 544.—Ins. Dom. Auckl.
Burnettii, F. Muell. Fragm. v. 96 = Burnettia cuneata.
ellipticus, *R. Br. Prod.* 325.—Austral.
Forrestii, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 55.—Austral.
nigricans, *R. Br. Prod.* 325.—Austral.
serratus, Lindl. Gen. et Sp. Orch. 393 = Caladenia serrata.
suaveolens, R. Br. Prod. 325 = Caladenia suaveolens.

LYPERIA, Benth. in Hook. Comp. Bot. Mag. i. (1835) 377. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 945.
URBANIA, Vatke, in Oestr. Bot. Zeitschr. xxv (1875) 10.
amplexicaulis, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 377.—Afr. austr.
argentea, *Benth. l. c.* 379.—Afr. austr.
aspalathoides, *Benth. l. c.* 381.—Afr. austr.
atropurpurea, *Benth. l. c.* 380.—Afr. austr.
Burkeana, *Benth. in DC. Prod.* x. 361.—Afr. austr.
canariensis, *Webb & Berth. Phyt. Canar.* iii. 146.—Ins. Canar.
canescens, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 379.—Afr. austr.
crassicaulis, *Benth. l. c.*—Afr. austr.
crocea, Eckl. ex Benth. in DC. Prod. x. 361 = atropurpurea.
cuneata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 380.—Afr. austr.
diandra, *E. Mey. Ann. Hort. Sem. Regiomont.* (1848) 5.—*Cf. Ann. Sc. Nat.* Sér. III. xi. (1849) 254.—Afr. austr.
filicaulis, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 380.—Afr. austr.
flogiflora, Steud. Nom. ed. II. ii. 84, sphalm. = phlogiflora.
foliolosa, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 380.—Afr. austr.
foliosa, Krauss, in Flora, xxvii. (1844) 835, sphalm. = praec.
fragrans, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 378.—Afr. austr.
fruticosa, *Benth. l. c.* 377.—Afr. austr.
glutinosa, *Benth. l. c.* 378.—Afr. austr.
incisa, *Benth. l. c.* 379.—Afr. austr.
integerrima, *Benth. in DC. Prod.* x. 359.—Afr. austr.
Kraussiana, *Benth. l. c.* 360.—Afr. austr.
macrocarpa, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 378.—Afr. austr.
macrophylla, Walp. Rep. iv. 534, sphalm. = microphylla.
micrantha, *Klotzsch, in Peters, Reise Mossamb. Bot.* 222.—Afr. trop.
microphylla, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 381.—Afr. austr.
mollis, *Benth. l. c.* 380.—Afr. austr.
multifida, *Benth. l. c.*—Afr. austr.
pedicellata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 223.—Afr. trop.
pedunculata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 379.—Afr. austr.
phlogiflora, *Benth. l. c.*—Afr. austr.
pinnatifida, *Benth. l. c.* 380.—Afr. austr.
purpurea, Walp. Rep. iii. 316 = atropurpurea.
racemosa, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 378.—Afr. austr.
simplex, *Benth. l. c.*—Afr. austr.
stricta, *Benth. in DC. Prod.* x. 360.—Afr. austr.
tenuiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 378.—Afr. austr.
tenuifolia, *Bernh. in Flora*, xxvii. (1844) 835.—Afr. austr.
tortuosa, *Benth. in DC. Prod.* x. 362.—Afr. austr.
tristis, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 378.—Afr. austr.
violacea, *Benth. l. c.* 379.—Afr. austr.

LYPERIA, Salisb. Gen. Pl. Fragm. 56 (1866) = **Fritillaria**, Tourn. (Liliac.).

LYPERODENDRON, Willd. ex Meissn. in DC. Prod. xiv. 168, in syn. (1857) = **Coccoloba**, Linn. (Polygon.).
coccoloboides, Willd. ex Meissn. l. c. = C. acuminata.

LYPROLEPIS, Steud. Syn. Pl. Cyp. 130 (1855) = **Kyllinga**, Rottb. (Cyperac.).
denudata, Steud. l. c. = K. peruviana.

LYRAEA, Lindl. Gen. et Sp. Orch. 46 (1830) = **Bulbophyllum**, Thou. (Orchid.).
prismatica, Lindl. l. c. = B. prismaticum.

LYRILOMA, Schlecht. in Bot. Zeit. iii. (1845) 27 sphalm. = **Lysiloma**, Benth. (Legumin.).

LYRIODENDRON, DC. Syst. i. 461 (1818) = **Liriodendron**, Linn. (Magnol.).

LYROCARPA, Hook. & Harv. in Hook. Lond. Journ. Bot. iv. (1845) 76. t. 4. *CRUCIFERAE*, Benth. & Hook. f. i. 93.
Coulteri, *Hook. & Harv. in Hook. Lond. Bot.* iv. (1845) 76.—Calif.
Palmeri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 123.—Am. bor. occ.

LYSANTHE, Salisb. in Knight, Prot. 116 (1809) = **Grevillea**, R. Br. (Proteac.).
cana, Knight, l. c. 117 = G. arenaria.
cytisifolia, Knight, l. c. 119 = G. sericea.
linariaefolia, Knight, l. c. = G. linearis.
podalyriaefolia, Knight, l. c. 117 = G. mucronulata.
riparia, Knight, l. c. 118 = G. linearis?
sericea, Knight, l. c. = G. punicea.
speciosa, Knight, l. c. = G. punicea.
stylosa, Knight, l. c. 117 = G. cinerea (vel G. montana).

LYSIANTHIUS, Adans. Fam. ii. 224 (1763) = **Lisianthus**, Linn. (Gentian.).

LYSIAS, Salisb in Trans. Hort. Soc. i. (1812) 288 = **Habenaria**, Willd. (Orchid.).
bifolia, Salisb. l. c. = H. bifolia.

LYSICARPUS, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 67. *MYRTACEAE*, Benth. & Hook. f. i. 709.
ternifolius, *F. Muell. l. c.* 68.—Austral.

LYSICHITON, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 62, in obs. = seq.

LYSICHITUM, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 62. *AROIDEAE*, Benth. & Hook. f. iii. 994.
ARCTIODRACON, A. Gray, in Mem. Am. Acad. Ser. II. vi. (1858-59) 408.
camtschatcense, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 62.—As. et Am. bor.
camtschaticum, Schott, Gen. Aroid. t. 91 = praec.
japonicum, Schott, ex Miq. Cat. Mus. Bot. Lugd. Bat. 96 = camtschatcense.

LYSIDICE, Hance, in Journ. Bot. v. (1865) 298. *LEGUMINOSAE*.
rhodostegia, *Hance, l. c.* 299.—China.

LYSILOMA, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 82. *LEGUMINOSAE*, Benth. & Hook. f. i. 595.
acapulcensis, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 82.—Mexic.
aurita, *Benth. l. c.*—Mexic.
bahamensis, Benth. l. c. 82 = latisiliqua.
desmostachya, Benth. l. c. 84 = acapulcensis.
divaricata, Benth. l. c. = Schiedeana.
Guachapele, *Benth. in Trans. Linn. Soc.* xxx. (1875) 533.—Panama.
latisiliqua, *Benth. l. c.* 534.—Am. bor.; Ind. occ.
Marchiana, Griseb. Fl. Brit. W. Ind. 223 = Calliandra portoricensis.
microphylla, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 83.—Mexic.
nigricans, Benth. l. c. 84 = Acacia nudiflora.
polyphylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 535.—Am. austr.
rostrata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 84.—Bras.

LYSILOMA:—
Sabicu, *Benth. in Hook. Kew Journ.* vi. (1854) 236.—Cuba.
Schiedeana, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 83.—Mexic.
tergemina, *Benth. in Trans. Linn. Soc.* xxx. (1875) 534.—Mexic.

LYSIMA, Medic. Phil. Bot. ii. 59, 107 (1791) = seq.

LYSIMACHIA, [Tourn.] Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 635.
BORISSA, Rafin. ex Steud. Nom. ed. II. i. 217 (1840).
COXIA, Endl. Gen. 733 (1839).
EPHEMERUM, Reichb. Nom. 124 (1841).
GODINELLA, Lestib. Bot. Belg. ii. 194 (1827).
LEROUXIA, Mérat, Nouv. Fl. Par. 77 (1812).
LISIMA, Medic. Phil. Bot. ii. 59, 107 (1791
LUBINIA, Vent. Jard. Cels, t. 96 (1800).
LYSIMANDRA, Reichb. Nom. 124 (1841).
NAUMBURGIA, Moench, Meth. Suppl. 23 (1802).
NUMMULARIA, Riv. ex Rupp. Fl. Jen. ed. Hall. 18 (1745).
PALLADIA, Moench, Meth. 429 (1794).
THEOPYXIS, Griseb. in Goett. Abh. vi. (1856) 126.
THYRSANTHUS, Schrank, in Denkschr. Akad. Muench. 1813 (1814) 75.
TRIDYNIA, Rafin. ex Steud. Nom. ed. II. ii. 704 (1841).
TRIDYRA, Rafin. ex Steud. l. c. 84, in syn. (1841).
acroadenia, Maxim. in Bull. Acad. Pétersb. xii. (1868) 70 = decurrens.
adoënsis, Hochst. ex Steud. in Flora, xxiv. (1841) 1. Intell. 23, nomen = Anagallis arvensis.
affinis, *Bunge, ex Duby, in DC. Prod.* viii. 67, nomen.—Hab.?
alata, Gardn. ex Thw. Enum. Pl. Zeyl. 172 = ramosa.
Alfredi, *Hance, in Journ. Bot.* xv. (1877) 356.—China.
alpestris, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 299.—Hongkong.
alternifolia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 26.—Reg. Himal.
anagalloides, *Sibth. & Sm. Fl. Graec. Prod.* i. 130.—Graecia.
angustifolia, A. Gray, Man. Bot. ed. I. 283 = Steironema longifolium.
angustifolia, Lam. Tabl. Encyc. i. 440; Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 417 = Steironema heterophyllum.
angustifolia, Michx. Fl. Bor. Am. i. 128 = stricta.
asperulaefolia, *Poir. Encyc. Suppl.* iii. 477.—Am. bor.
atropurpurea, Hook. Exot. Fl. t. 180 = nutans.
atropurpurea, *Linn. Sp. Pl.* 147.—Graecia; As. Min.
atropurpurea, Murr. in Comm. Gotting. (1782) 6. t. 1 = dubia.
azorica, [Hornem.] ex Hook. Bot. Mag. t. 3273 = nemorum.
barystachys, *Bunge, Enum. Pl. Chin. Bor.* 53.—As. or.
barystachys, Klatt, in Abh. Naturf. Ver. Hamb. iv. iv. (1866) 11 = Fortunei.
brachystachys, Carr. in Rev. Hortic. (1881) 90, sphalm. = barystachys.
bulbifera, Curt. Bot. Mag. t. 104 = stricta.
buxifolia, Molina, Sagg. Chile, ed. II. 134 = Anagallis alternifolia.
candida, Lindl. in Journ. Hort. Soc. i. (1846) 301 = obovata.
capitata, Pursh, Fl. Am. Sept. i. 135 = thyrsiflora.
capitellata, Rafin. in Med. Repos. N. York, II. v. (1808) 354 = thyrsiflora.
chenopodioides, *Watt, ex Hook. f. Fl. Brit. Ind.* iii. 503.—Reg. Himal.
chlethroides, Duby, vide clethroides.
Christinae, *Hance, in Journ. Bot.* xi. (1873) 117.—China.
ciliata, Linn. Sp. Pl. 147 = Steironema ciliatum.
ciliata, Walt. Fl. Carol. 92 = Steironema heterophyllum.
Clementsoniana, Wall. Cat. n. 1485 = Leschenaultii.
clethroides, *Duby, in DC. Prod.* viii. 61 (chlethroides).—Japon.
consobrina, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 224 = decurrens.
Cousiniana, *Coss. in Bull. Soc. Bot. Fr.* ix. (1862) 174.—Alger.
cuspidata, *Blume, Bijdr.* 737.—Java.

LYSIMACHIA:—
cuspidata, Klatt, in Abh. Naturf. Ver. Hamb. iv. iv. (1866) 36 = Klattiana.
davurica, Ledeb. in Mém. Acad. Pétersb. v. (1814) 523 = vulgaris.
debilis, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 25 = japonica.
decipiens, Bertol. in Giorn. Ligust. Sc. (Jan. 1827) 9; ex Reichb. Fl. Germ. Excurs. 864 = Steironema heterophyllum.
decurrens, *Forst. f. Prod.* 12.—Reg. Himal.; China; Ins. Pacif.
deltoidea, *Wight, Illustr.* ii. 137. t. 144.—Ind. or.
dubia, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 199.—As. Min.; Persia.
dubia, Willd. Enum. Hort. Berol. 195, in nota = Steironema heterophyllum.
eileta, *Lehm. in Hamb. Gartens.* vi. (1850) 323.—Ind. or.
elata, Salisb. Prod. 120 = vulgaris.
Ephemerum, *Linn. Sp. Pl.* 146.—Europ. austr.
Ephemerum, Thunb. Fl. Jap. 83 = clethroides.
evalvis, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 77.—Reg. Himal.
ferruginea, Edgew. in Trans. Linn. Soc. xx. (1846) 83 = japonica.
floribunda, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 575.—Java.
Foenum-graecum, *Hance, in Journ. Bot.* xv. (1877) 355.—China.
Fortunei, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 68.—China; Japon.
Fraseri, *Duby, in DC. Prod.* viii. 65.—Am. bor.
glandulosa, Edgew. in Trans. Linn. Soc. xx. (1846) 82 = alternifolia.
glandulosa, Klatt, in Abh. Naturf. Ver. Hamb. iv. iv. (1866) 31. t. 17 = chenopodioides.
glauca, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 234 = vulgaris.
glauca, Moench, Meth. 511 = L. Ephemerum.
glaucescens, Wall. ex Hook. f. Fl. Brit. Ind. iii. 502 = obovata.
glaucophylla, *Hook. & Arn. Bot. Beech. Voy.* 306. t. 68.—Mexic.
grammica, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 225.—China.
guestphalica, Weihe, ex Reichb. Fl. Germ. Excurs. 410 = vulgaris.
Herbemonti, Ell. Sketch, i. 232 = asperulaefolia.
heterogenea, *Klatt, in Linnaea*, xxxvii. (1871–73) 501 (heteroganea).—China.
heterophylla, D. Don, Prod. Fl. Nep. 84 = pyramidalis.
heterophylla, Michx. Fl. Bor. Am. i. 127 = Steironema heterophyllum.
Hillebrandi, *Hook. f. ex A. Gray, in Proc. Am. Acad.* v. (1861–62) 328.—Ins. Sandvic.
hirsuta, Michx. Fl. Bor. Am. i. 127 = quadrifolia.
Hochstetteri, Steud. Nom. ed. II. ii. 84 = nemorum.
hybrida, Michx. Fl. Bor. Am. i. 126 = Steironema heterophyllum.
inconspicua, *Miq. in Journ. Bot. Néerl.* i. (1861) 110.—China.
intermedia, Janka, ex Nym. Consp. 600 = dubia.
japonica, *Thunb. Fl. Jap.* 83.—Reg. Himal.; China; Austral.
japonica, Thw. Enum. Pl. Zeyl. 172 = deltoides.
javanica, Blume, Bijdr. 736 = decurrens.
Keiskeana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 120.—Japon.
Klattiana, *Hance, in Journ. Bot.* xvi. (1878) 236.—China.
lanceolata, Pursh, Fl. Am. Sept. ii. 729 = Fraseri.
lanceolata, Walt. Fl. Carol. 92 = Steironema heterophyllum.
Leschenaultii, *Duby, in DC. Prod.* viii. 61.—Ind. or.
leucantha, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 121.—Japon.
linearifolia, *Griff. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 86.—Burma.
linearifloba, Hook. & Arn. Bot. Beech. Voy. 268 = mauritiana.
linifolia, Salisb. Prod. 120 = Asterolinon stellatum.
Linum-stellatum, Linn. Sp. Pl. 148 = Asterolinon stellatum.
lobelioides, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 29.—Reg. Himal.; Java.
longifolia, Pursh, Fl. Am. Sept. i. 135 = Steironema longifolium.

LYSIMACHIA:—
Loomisii, Torr. in Croom, Cat. Pl. New-Bern, 46 = stricta.
lubinioides, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 140 = mauritiana.
lutea, Jiraseck, ex Mert. & Koch, Deutschl. Fl. ii. 130 = vulgaris.
maculata, R. Br. Prod. 428 = japonica.
mauritiana, *Lam. Encyc.* iii. 592.—China.
media, *Willd. ex Roem. & Schult. Syst.* iv. 784.—Sibir.
Metziana, Hohen. ex Hook. f. Fl. Brit. Ind. iii. 505 = deltoides.
minoricensis, *Rodrig. in Bull. Soc. Bot. Fr.* xxv. (1878) 240.—Ins. Balear.
Monnieri, Linn. Cent. Pl. ii. 9 = Herpestis Monnieria.
multiflora, Wall. Cat. n. 1487 = decurrens.
myrtifolia, Molina, Sagg. Chile, ed. II. 134 = Anagallis alternifolia.
nemoralis, Salisb. Prod. 120 = nemorum.
nemorum, Geners. Elench. Scepus. n. 177 ; ex Roem. & Schult. Syst. iv. 128 = L. Nummularia.
nemorum, *Linn. Sp. Pl.* 148.—Europ.
Notoniana, Wall. ex Hook. f. Fl. Brit. Ind. iii. 502 = Leschenaultii.
Nummularia, *Linn. Sp. Pl.* 148.—Europ.
nutans, *Nees, ex Duby, in DC. Prod.* viii. 61.—Afr. austr.
obovata, *Buch.-Ham. ex Wall. Cat.* n. 1488.—Ind. or.; Java; China.
orientalis, Lam. Encyc. iii. 570 = dubia.
Otani, Asso, Syn. Stirp. Arag. 22. t. 2. f. 1 = L. Ephemerum.
pacifica, *F. Muell. Phyt. N. Hebr.* 18.—Ins. Pacif.
paludosa, Baumg. Enum. Stirp. Transs. i. 141 = vulgaris.
paniculata, Gilib. Fl. Lituan. i. 29 = vulgaris.
paridiformis, *Franch. in Bull. Soc. Linn. Par.* i. (1884) 433.—China.
parviflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 196. —Madag.
peduncularis, *Wall. Cat.* n. 1489.—Burma.
pentapetala, Bunge, Enum. Pl. Chin. Bor. 53 = Apochoris pentapetala.
prolifera, *Klatt, in Abh. Naturf. Ver. Hamb.* iv. IV. (1866) 30. t. 16.—Reg. Himal.
punctata, *Linn. Sp. Pl.* 147.—Europ.; As. Min.
punctata, Walt. Fl. Carol. 92 = quadrifolia.
purpurea, Steud. Nom. ed. II. ii. 85 = *atropurpurea*, Hook.
pyramidalis, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 28. —Reg. Himal.
quadriflora, Sims, Bot. Mag. t. 660 = Steironema longifolium.
quadrifolia, *Linn. Sp. Pl.* 147.—Am. bor.
quadrifolia, Mill. Gard. Dict. ed. VIII. n. 10 = punctata.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 16.—Abyssin.
quinquangularis, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 504 = alternifolia.
racemosa, Lam. Encyc. iii. 570; Michx. Fl. Bor. Am. i. 127 = stricta.
radicans, Hook. Comp. Bot. Mag. i. (1835) 177 = Steironema radicans.
ramosa, *Wall. Cat.* n. 1490.—Reg. Himal.; Java.
repens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 299 = L. Nummularia.
repens, Urv. in Mém. Soc. Linn. Par. iv. (1826) 606 = Anagallis alternifolia.
revoluta, Nutt. Gen. Am. i. 122 = Steironema longifolium.
rotundifolia, F. W. Schmidt, Fl. Boëm. ii. 59, in obs. = L. Nummularia.
Ruhmeriana, *Vatke, in Linnaea*, xl. (1876) 204.—Afr. trop.
salicifolia, Mill. Gard. Dict. ed. VIII. n. 6 = L. Ephemerum.
salicifolia, *F. Muell. ex Benth. Fl. Austral.* iv. 269.—Austral.
samolina, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 224 = obovata.
secunda, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 84 = lobelioides.
sedoides, *Lehm. Ind. Sem. Hort. Hamb.* (1856) 7.—*Cf. Linnaea*, xxix. (1857–58) 724.—N. Zel.
serpyllifolia, Poir. Encyc. Suppl. iii. 477 = Asterolinon serpyllifolium.

LYSIMACHIA:—
sessilifolia, Rafin. Atl. Journ. 151 (Quid ?).—Am. bor.
sikokiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 121. —Japon.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 110 = decurrens.
sororia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 120.—Japon.
spathulata, Schouw.—Cf. Linnaea, xxiv. (1851) 160 = mauritiana.
spatulata, Klatt, in Abh. Naturf. Ver. Hamb. iv. IV. (1866) 32 = mauritiana.
stellata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = Asterolinon stellatum.
stricta, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 199.—Am. bor.
suaveolens, Schoenh. ex Garcke, Fl. Deutschl. ed. IV. 272 = L. Nummularia.
subcapitata, Rafin. in Med. Rep. N. York, Ser. III. ii. (1811) 333 (Quid ?).—Am. bor.
tenella, Linn. Sp. Pl. 148 = Anagallis tenella.
tenella, Wall. Cat. n. 1491 = Centunculus tenellus.
tetragona, D. Don, Prod. Fl. Nep. 83 = alternifolia.
thyrsantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = thyrsiflora.
thyrsiflora, Geners. Elench. Scepus, n. 176 ; ex Roem. & Schult. Syst. iv. 120 = vulgaris.
thyrsiflora, *Linn. Sp. Pl.* 147.—Reg. bor.
tomentosa, Presl, Fl. Sic. p. xxxviii = vulgaris.
Trientalis, Klatt, in Linnaea, xxxvii. (1871–73) 499 = Trientalis americana, europaea.
uliginosa, *Blume, Bijdr.* 737.—Java.
umbellata, *Phil. in Linnaea*, xxx. (1859–60) 195.—Chili.
umbrosa, Gardn. ex Thw. Enum. Pl. Zeyl. 172 = deltoides.
verticillaris, Spreng. Fl. Hal. Mant. 36 = punctata.
verticillata, Bieb. Fl. Taur. Cauc. i. 141 = praec.
villosa, F. W. Schmidt, Fl. Boëm. ii. 57 = punctata.
vulgaris, *Linn. Sp. Pl.* 146.—Europ.; As. bor.
vulgaris, Walt. Fl. Carol. 92 = stricta.
vulneraria, *Nois. ex Steud. Nom.* ed. II. ii. 85, nomen. —Hab.?
westphalica, Weihe, in Flora, v. (1822) 411 = vulgaris.
Zavadskii, Wiesn. in Oestr. Bot. Wochenbl. iv. (1854) 257 = nemorum?

LYSIMANDRA, Reichb. Nom. 124 (1841) = **Lysimachia**, Tourn. (Primul.).

LYSIMNIA, Rafin. Fl. Tellur. iv. 43 (1836) = **Brassavola**, R. Br. (Orchid.).
bicolor, Rafin. l. c. = B. cordata.

LYSINEMA, R. Br. Prod. 552 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 615.
JULIETA, Leschen. ex DC. Prod. vii. 764 (1839).
LASYNEMA, Poir. Encyc. Suppl. ii. 553, 555 (1811).
WOOLLSIA, F. Muell. Fragm. viii. 52, 55 (1873).
attenuatum, Link, Enum. Hort. Berol. i. 211 = pungens.
brevilimbatum, F. Muell. Fragm. iii. 142 = lasianthum.
ciliatum, *R. Br. Prod.* 552.—Austral.
conspicuum, *R. Br. l. c.*—Austral.
curvatum, Lindl. Swan River App. 25 = ciliatum.
elegans, *Sond. in Lehm. Pl. Preiss.* i. 327.—Austral.
fimbriatum, *F. Muell. Fragm.* iv. 125.—Austral.
lasianthum, *R. Br. Prod.* 552.—Austral.
ovatum, Sond. in Lehm. Pl. Preiss. i. 329 = ciliatum.
pentapetalum, R. Br. Prod. 552 = ciliatum.
pungens, *R. Br. l. c.*—Austral.
ruscifolium, Sieber, ex Spreng. Syst. iv. Cur. Post. 64 = Epacris purpurascens.
Sieberi, Benth. in Enum. Pl. Hueg. 76 = pungens.
spicatum, Lindl. Swan River App. 25 = ciliatum.
virgatum, DC. Prod. vii. 765 = ciliatum.

LYSINOTUS, Low, Sarawak, 67, sphalm. (1848) = seq.

LYSIONOTHUS, D. Dietr. Syn. Pl. i. 30 (1839) = seq.

LYSIONOTIS, G. Don, Gen. Syst. iv. 644 (1837) = seq.

LYSIONOTUS, D. Don, in Edinb. Phil. Journ. vii. (1822) 85. *GESNERACEAE*, Benth. & Hook. f. ii. 1015.
albida, Blume, Bijdr. 765 = Aeschynanthus purpurascens.
angustifolia, Blume, l. c. = Aeschynanthus angustifolia.
Aucklandii, *Low, Sarawak*, 67.—Borneo.
conferta, *C. B. Clarke, in DC. Monog. Phan.* v. 58.—Assam.
Griffithii, C. B. Clarke, l. c. 60 = Loxostigma Griffithii.
longiflora, Blume, Bijdr. 766 = Aeschynanthus longiflora.
pauciflora, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 534.—Japon.
pulchella, Blume, ex C. B. Clarke, in DC. Monog. Phan. v. 34 = Aeschynanthus Horsfieldii.
serrata, *D. Don, in Edinb. Phil. Journ.* vii. (1822) 86. —Reg. Himal.
ternifolia, Wall. Pl. As. Rar. ii. 20. t. 118 = serrata.

LYSIOSEPALUM, F. Muell. Fragm. i. 142 (1859). *STERCULIACEAE*, Benth. & Hook. f. i. 228.
Barryanum, *F. Muell. l. c.* 143.—Austral.
rugosum, *Benth. Fl. Austral.* i. 267.—Austral.

LYSIOSTYLES, Benth. in Hook. Lond. Journ. Bot. v. (1846) 356. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 878.
scandens, *Benth. l. c.*—Guian.

LYSIPOMA, Spreng. Syst. i. 719 (1325) = seq.

LYSIPOMIA, H. B. & K. Nov. Gen. et Sp. iii. 318 (1818). *CAMPANULACEAE*, Benth. & Hook. f. ii. 549.
acaulis, *H. B. & K. l. c.* 321.—Peruv.
aretioides, *H. B. & K. l. c.*—Peruv.
glanduliferum, Schlecht. in Lechl. Berb. Am. Austr. (1857) 58 ; et ex Wedd. Chlor. And. ii. 11 = Pratia glandulifera.
laciniata, *A. DC. in DC. Prod.* vii. 349.—Bolivia.
montioides, *H. B. & K. Nov. Gen. et Sp.* iii. 320.—Ecuador.
muscoides, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 286. t. 9 A.—N. Granat.
reniformis, *H. B. & K. Nov. Gen. et Sp.* iii. 320.—Peruv.
repens, Phil. f. Cat. Pl. Vasc. Chil. 189, sphalm. = seq.
serpens, Kunze, ex Cham. in Linnaea, viii. (1833) 224 = Selliera radicans.
sphagnophilum, *Griseb. in Lechl. Berb. Am. Austr.* (1857) 58; *et ex Wedd. Chlor. And.* ii. 15.—Peruv.
subulata, *G. Don, Gen. Syst.* iii. 717.—Peruv.

LYSISTEMMA, Steetz, in Peters, Reise Mossamb. Bot. 340 (1863) = **Vernonia**, Schreb. (Compos.).
dendigulense, Steetz, l. c. 341 = V. indica.
divergens, Steetz, l. c. = V. divergens.
multiflorum, Steetz, l. c. 342 = V. divergens.
pectiniforme, Steetz, l. c. 343 = V. pectiniformis.
salignum, Steetz, l. c. = V. saligna.

LYSISTIGMA, Schott, in Bonplandia, x. (1862) 222 = **Taccarum**, Brongn. (Aroid.).
peregrinum, Schott, l. c. 223 = T. cylindricum.

LYSSANTHE, D. Dietr. Syn. Pl. i. 634 (1839) = **Lissanthe**, R. Br. (Epacrid.).

LYSSANTHE, Endl. Gen. 340 (1837) = LYSANTHE, Salisb. = **Grevillea**, R. Br. (Proteac.).

LYTHOSPERMUM, Lucé, Topog. Nachr. Oesel, 43 (1825) = **Lithospermum**, Linn. (Borag.).

LYTHRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 129 = **Lythrum**, Linn.
salicarium, St. Lag. ll. cc. = Lythrum Salicaria.

LYTHROPSIS, Welw. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 305 = **Lythrum**, Linn.
peploides, Welw. ex Koehne, l. c. 311 = Lythrum hispidulum.

LYTHRUM, Linn. Syst. ed. I (1735). *LYTHRARIEAE*, Benth. & Hook. f. i. 779.
ANISOTES, Lindl. Introd. Nat. Syst. ed. II. 101, App. 441 (1836).
EDITELES, Rafin. Sylva Tellur. 103 (1838).
HYSSOPIFOLIA, Opiz, Seznam, 53 (1852).
LYTHROPSIS, Welw. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 305.
MOZULA, Rafin. in Journ. Phys. xci. (1820) 71.
PENTAGLOSSUM, Forst. Fl. Aegypt. Arab. 11 (1775).
PYTHAGOREA, Rafin. in Journ. Phys. lxxxix. (1819) 96.
SALICA, Hill, Hort. Kew. 172[2d] (1768).
SALICARIA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
acinifolium, [*Moç. & Sesse,*] *ex DC. Prod.* iii. 81, in syn. —Mexic.
acuminatum, Willd. Sp. Pl. ii. 866 = virgatum.
acutangulum, Lag. Gen. et Sp. Nov. 16 = Graefferi.
alatum, Hemsl. Biol. Centr. Am. Bot. i. 447, partim = album.
alatum, Presl, Delic. Prag. 55 = Graefferi.
alatum, *Pursh, Fl. Am. Sept.* i. 334.—Am. bor.
albicaule, Bert. in Merc. Chil. (1829) 694.—Cf. Am. Journ. Sc. xxiii. (1833) 250 = album.
album, *H. B. & K. Nov. Gen. et Sp.* vi. 193.—Mexic.
album, [Moç. & Sessé,] ex DC. Prod. iii. 84 = Cuphea jorullensis.
alternifolium, Lorey, Fl. Côte-d'Or, i. 348. t. 2 = L. Salicaria.
altissimum, *Pomel, Nouv. Mat. Fl. Atl.* 155.—Alger.
americanum, *Mill. Gard. Dict.* ed. VIII. n. 9.—Mexic.
anomalum, St. Hil. Fl. Bras. Mer. iii. 130 = Pleurophora anomala.
apetalum, Spreng. Syst. ii. 454 = Ammannia latifolia.
arnhemicum, F. Muell. Fragm. iii. 109 = Nesaea arnhemica.
austriacum, Jacq. Enum. Stirp. Vindob. 80, 243 = virgatum.
axillare, *Vell. Fl. Flum.* 201; v. t. 9.—Bras.
bahiense, Casar. ex Koehne, in Mart. Fl. Bras. xiii. II. 307 = Pleurophora anomala.
bibracteatum, Salzm. ex DC. Prod. iii. 81 = tribracteatum.
biflorum, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 227 = L. Salicaria.
Bocconi, Déségl.—Cf. Bull. Soc. Bot. Fr. xxv. (1878) Bibl. 108 = L. Salicaria.
breviflorum, S. Wats. in Proc. Am. Acad. xii. (1877) 251 = alatum.
calcaratum, Jacq. ex Koehne, in Engl. Bot. Jahrb. ii. (1882) 407 = Cuphea jorullensis.
calcaratum, *Vell. Fl. Flum.* 200; v. t. 7.—Bras.
californicum, Torr. & Gray, Fl. N. Am. i. 482 = alatum.
campestre, Griseb. in Goett. Abh. xix. (1874) 140 = maritimum.
carthagenense, Jacq. Enum. Pl. Carib. 22 = Cuphea Balsamona.
cashmerianum, Royle, Illustr. Bot. Himal. 203 = L. Salicaria.
ciliatum, Sw. Prod. Veg. Ind. Occ. 76 = Cuphea decandra.
cinereum, Griseb. Spicil. Fl. Rumel. i. 110 = L. Salicaria.
cistifolium, L'Hérit. ex DC. Prod. iii. 88 = Cuphea microphylla.
cordifolium, [Moç. & Sesse,] ex DC. l. c. 85 = Cuphea cyanaea.
cordifolium, Sw. Prod. Veg. Ind. Occ. 76 = Cuphea Swartziana.
coronense, Schur, Enum. Pl. Transs. 218 = L. Salicaria.
Cuphea, Linn. f. Suppl. 249 = Cuphea viscosissima.
dibracteatum, Guss. Fl. Sic. Syn. i. 526 = tribracteatum.
diffusum, Sweet, Brit. Flow. Gard. Ser. I. ii. t. 149 = L. Salicaria.
diffusum, *Vell. Fl. Flum.* 201; v. t. 8.—Bras.
dipetalum, Linn. f. Suppl. 250 = Cuphea verticillata.
divaricatum, Colla, in Mem. Acc. Torin. xxxviii. (1835) 3 = Pleurophora pusilla.
divaricatum, Schur, Enum. Pl. Transs. 219 = virgatum.
domingense, Spreng. ex DC. Prod. iii. 86 = Cuphea micrantha.
dubium, Schult. Oesterr. Fl. ed. II. ii. 5 = L. Salicaria.

LYTHRUM:—
elegans, Schur, Enum. Pl. Transs. 219 = virgatum.
filiforme, Perr. ex DC. Prod. iii. 77 = Ammannia senegalensis.
flagellare, Shuttlew. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 321 = ovalifolium.
flavum, Spreng. Syst. ii. 454 = Nesaea salicifolia.
flexicaule, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 116 = nanum.
flexuosum, Lag. Gen. et Sp. Nov. 16 = Graefferi.
floribundum, F. Muell. Fragm. vii. 146 = Nesaea floribunda.
fruticosum, Linn. Syst. ed. X. 1045 = Woodfordia floribunda.
geminiflorum, Bertol. Fl. Ital. v. 16 = thesioides.
glaucescens, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 116 = thesioides.
glaucum, Dulac, Fl. Hautes-Pyr. 327 = L. Hyssopifolia.
gracile, *Benth. Pl. Hartw.* 7.—Mexic.
gracile, DC. Cat. Hort. Monsp. 123 = L. Salicaria.
gracile, Hemsl. Biol. Centr. Am. Bot. i. 447 = acinifolium.
Graefferi, C. Gay, Fl. Chil. ii. 368 = album.
Graefferi, Lodd. Bot. Cab. t. 1338 = L. Hyssopifolia.
Graefferi, *Tenore, Prod. Fl. Nap.* p. lxviii.—Geront. temp. et trop.
grandiflorum, Perr. ex DC. Prod. iii. 80 = Nesaea Candollei.
Gussoni, Presl, Delic. Prag. 55 = Graefferi.
hexagonum, Berchtold, ex Opiz, Boehm. Gew. (1823) 58 = L. Salicaria.
hispanicum, Mill. Gard. Dict. ed. VIII. n. 5 = Cuphea Parsonsia.
hispidulum, *Koehne, in Sitzb. Bot. Ver. Brandenb.* (1880) 29.—Europ.; Afr.
Hunteri, DC. Prod. iii. 83 = Woodfordia floribunda.
Hyssopifolia, Desf. Fl. Atlant. i. 372 = Graefferi.
Hyssopifolia, *Linn. Sp. Pl.* 447.—Cosmop. temp.
hyssopifolium, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 119 = alatum.
hyssopifolium, Thunb. Prod. Pl. Cap. 85 = L. Hyssopifolia.
intermedium, Ledeb. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 92 = L. Salicaria.
junceum, [Soland. in] Russ. Aleppo, ed. II. ii. 253 = Graefferi.
Kennedianum, H. B. & K. Nov. Gen. et Sp. vi. 194 = alatum.
lanceolatum, Ell. Sketch, i. 544 = alatum.
lineare, Bert. ex DC. Prod. iii. 82 = L. Hyssopifolia.
lineare, Hook. & Arn. Bot. Beech. Voy. 343 = alatum.
lineare, *Linn. Sp. Pl.* 447.—Am. bor.
linifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 421 = thesioides.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 3 = virgatum.
lycioides, Willd. ex Steud. Nom. ed. II. ii. 86 = maritimum.
maculatum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 12 = Graefferi.
maritimum, Cham. & Schlecht. in Linnaea, v. (1830) 568 = acinifolium.
maritimum, *H. B. & K. Nov. Gen. et Sp.* vi. 194.—Peruv.
Melanium, Linn. Syst. ed. X. 1045 = Cuphea viscosissima.
micranthum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 422 = nanum.
microphyllum, Kar. & Kir. l. c. 421 = tribracteatum.
monanthum, Link, ex Steud. Nom. ed. I. 502 = Graefferi.
multiflorum, Perr. ex DC. Prod. iii. 77 = Ammannia senegalensis.
myrtifolium, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 189.—Am. bor.
nanum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 422.—Soongaria.
nanum, Nym. Consp. 252 = tribracteatum.
nummulariaefolium, *Loisel. in Desv. Journ. Bot.* ii. (1809) 330.—Corsica.
nummulariaefolium, Pers. Syn. ii. 8 = L. Salicaria.
ovalifolium, *Engelm. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 187.—Texas.
palustre, Salisb. Prod. 365 = L. Salicaria.
Parsonsia, Linn. Syst. ed. X. 1045 = Cuphea Parsonsia.

LYTHRUM:—
parviflorum, Bast. ex Reichb. Fl. Germ. Excurs. 640 = L. Hyssopifolia.
Pemphis, Linn. f. Suppl. 249 = Pemphis acidula.
petiolatum, Linn. Sp. Pl. 446 = Cuphea viscosissima.
Preslii, Guss. Pl. Rar. 188 = Graefferi.
propinquum, Weinm. in Bull. Soc. Nat. Mosc. xxiii. (1850) I. 544 = L. Salicaria.
prostratum, Domb. ex Koehne, in Mart. Fl. Bras. xiii. II. 199 = L. Hyssopifolia.
pubescens, Sweet, Hort. Brit. ed. II. 202 = L. Salicaria.
punctatum, Span. in Linnaea, xv. (1841) 202 = Woodfordia floribunda.
punicifolium, Cham. & Schlecht. in Linnaea, ii. (1827) 356 = Graefferi.
Purshianum, Steud. Nom. ed. II. ii. 86 = L. Salicaria.
quintuplinervium, Neuw. Reise, v. ii; ex Flora, iv. (1821) 302 = Diplusodon quintuplinervius.
racemosum, Linn. f. Suppl. 250 = Cuphea spicata.
ramosum, Perr. ex DC. Prod. iii. 80 = Ammania senegalensis.
rigidulum, Sond. in Linnaea, xxiii. (1850) 41 = Nesaea rigidula.
Robertsii, F. Muell. Fragm. vii. 145 = Nesaea Robertsii.
rotundifolium, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 283.—Abyssin.
sagittifolium, Sond. in Linnaea, xxiii. (1850) 41 = Nesaea sagittifolia.
Salicaria, *Linn. Sp. Pl.* 446.—Reg. bor. temp.; Austral.
Salzmanni, Jord. Obs. Pl. Crit. v. (1847) 42 = tribracteatum.
satureifolium, [Moç. & Sesse,] ex DC. Prod. iii. 82 = alatum.
scabrum ×, Simk. in Herm. Termész. Füzetek, i. (1877) 241.—Europ.
semiamplexans, Ruiz, ex Koehne, in Engl. Bot. Jahrb. I. (1881) 244 = Ammannia senegalensis.
silenoides, *Boiss. & Noé, Diagn.* Ser. II. ii. 55.—Persia; Afghan.
spicatum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 553 = L. Salicaria.
spiciforme, Dulac, Fl. Hautes-Pyr. 327 = L. Salicaria.
strictum, Willd. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 317 = L. Thymifolia.
tenellum, Thunb. Prod. Pl. Cap. 85 = L. Hyssopifolia.
thesioides, *Bieb. Fl. Taur. Cauc.* i. 367.—Reg. Caucas.
Thymifolia, All. Fl. Pedem. ii. 168 = Graefferi.
Thymifolia, Krock. Fl. Siles. ii. 92 = L. Hyssopifolia.
Thymifolia, *Linn. Sp. Pl.* 447—Europ.; Afr. bor.
thymifolium, Boiss. Fl. Orient. ii. 740 = thesioides.
thymifolium, Sibth. & Sm. Fl. Graec. Prod. i. 321 = tribracteatum.
thymophyllum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 129 (thymouphyllum) = L. Thymifolia.
tomentosum, Mill. Gard. Dict. ed. VIII. n. 2 = L. Salicaria.
triantha, Vent. ex DC. Prod. iii. 90 = Nesaea triflora.
tribracteatum, *Salzm. ex Tenore, Ind. Sem. Hort. Neap.* (1830) 13; *ex Spreng. Syst.* iv. *Cur. Post.* 190.—Europ.; As. bor.
triflorum, Linn. f. Suppl. 249 = Nesaea triflora.
truxillense, Steud. Nom. ed. II. ii. 86 = Cuphea aequipetala.
tuxtlense, [Moç. & Sesse,] ex DC. Prod. iii. 88 = Cuphea aequipetala.
uniflorum, Perr. ex DC. l. c. 78 = Ammannia elatinoides.
verticillatum, Linn. Sp. Pl. 447 = Nesaea verticillata, *H. B. & K.*
virgatum, *Linn. l. c.*—Europ.; As. bor.
virgatum, Spreng. ex DC. Prod. iii. 82 = lineare.
virgatum, Walt. Fl. Carol. 120 = alatum?
virginicum, Kenn. ex DC. Prod. iii. 81 = alatum.
virgultosum, Griseb. Cat. Pl. Cub. 106 = lineare.
Vulneraria, Ait. ex Schrank, Pl. Rar. Hort. Monac. t. 27 = alatum.
vulneria, Steud. Nom. ed. II. ii. 86, sphalm. = praec.
Wagneri, *Bunge, ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 64.—Reg. Casp.

LYTRUM, Vill. Fl. Dauph. iii. 818 (1789) = **Lythrum**, Linn.

M

MAACKIA, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 128, 143 = **Cladrastis**, Rafin. (Legumin.).
amurensis, Rupr. l. c. = C. amurensis.

MAASA, Roem. & Schult. Syst. v. 220 (1819), sphalm. = **Maesa**, Forsk. (Myrsin.).

MABA, Forst. Char. Gen. 121. t. 61 (1776). *EBENACEAE*, Benth. & Hook. f. ii. 664.
EBENOXYLUM, Lour. Fl. Cochinch. ii. 613 (1790).
FERREOLA, Koen. ex Roxb. Pl. Corom. i. 35. t. 45. (1795).
HOLOCHILUS, Dalz. in Hook. Kew Journ. iv. (1852) 290.
MACREIGHTIA, A. DC. in DC. Prod. viii. 220 (1844).
RHIPIDOSTIGMA, Hassk. Retzia, 103 (1855).
abyssinica, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 132.—Abyssin.
acapulcensis, *Hiern, l. c.* 128.—Mexic.
acuminata, *Hiern, l. c.* 112.—Zeylan.
albens, *Hiern, l. c.* 126.—Mexic.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 138.—Ins. Andam.
Andersoni, [*Soland. ex*] *Seem. Fl. Vit.* 152.—Ins. Tonga.
angustifolia, Miq. Anal. Bot. Ind. iii. 13 = buxifolia.
Beccarii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 140.—Borneo.
buxifolia, *Pers. Syn.* ii. 606.—As. et Afr. trop.
Cargillia, F. Muell. Fragm. v. 162 = Diospyros Cargillia.
caribaea, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 125.—Ind. occ.
cauliflora, *Hiern, l. c.* 142.—Am. trop.
compacta, *R. Br. Prod.* 528.—Austral.
confertiflora, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 136.—Borneo.
cordata, *Hiern, l. c.* 141.—Borneo.
Cumingiana, A. DC. in DC. Prod. viii. 241 = buxifolia.
cupulosa, F. Muell. Fragm. v. 164 = rufa.
diffusa, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 111.—Madag.
Ebenoxylon, G. Don, Gen. Syst. iv. 43 = elliptica.
Ebenus, Spreng. Syst. ii. 126 = elliptica.
Ebenus, Wight, Ic. iv. t. 1228–9 = buxifolia.
elliptica, *Forst. Char. Gen.* 122.—Ins. Amicor.
elliptica, Seem. Fl. Vit. 152 = sandwicensis.
fasciculosa, *F. Muell. Fragm.* v. 163.—Austral.; N. Caled.
foliosa, *Rich, ex A. Gray, in Proc. Am. Acad.* v. (1862) 326.—Ins. Fiji.
geminata, *R. Br. Prod.* 527.—Austral.
glauca, *Montr. in Mém. Acad. Lyon*, x. (1860) 230.—N. Caled.
Grisebachii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 125.—Cuba.
guineensis, A. DC. in DC. Prod. viii. 241 = buxifolia.
hemicycloides, *F. Muell. ex Benth. Fl. Austral.* iv. 290.—Austral.
hermaphroditica, *Zoll. Syst. Verz. Ind. Arch.* 135.—Java.
Hilairei, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 143.—Bras.
Hillebrandii, *Seem. Fl. Vit.* 151.—Ins. Sandvic.
humilis, *R. Br. Prod.* 527.—Austral.
inconstans, *Griseb. Fl. Brit. W. Ind.* 404.—Ind. occ.
interstans, F. Muell. Fragm. v. 163 = reticulata.
intricata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 126.—Calif.
javanica, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 158.—Java.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 584.—Sumatra.
lancea, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 118.—Afr. trop.
lanceolata, *Hiern, l. c.* 131.—Madag.
lateriflora, *Hiern, in Journ. Linn. Soc.* xx. (1883) 366.—Ins. Fiji.
laurina, *R. Br. Prod.* 527.—Austral.
laxiflora, Benth. Fl. Austral. iv. 290 = fasciculosa.
littorea, R. Br. Prod. 527 = buxifolia.

MABA:—
madagascariensis, A. DC. in DC. Prod. viii. 241 = buxifolia.
Maingayi, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 138.—Malaya.
major, *Forst. f. Pl. Escul.* 54.—Ins. Fiji.
Mannii, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 129.—Afr. trop.
megalocarpa, F. Muell. Fragm. v. 163 = Diospyros maritima.
Mellinoni, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 143.—Am. trop.
merguensis, *Hiern, l. c.* 134.—Burma; Malaya.
micrantha, *Hiern, l. c.* 133.—Ind. or.
Motleyi, *Hiern, l. c.* 139.—Borneo.
Mualala, *Welw. ex Hiern, l. c.* 111.—Afr. trop.
myristicoides, *Hiern, l. c.* 142.—Bras.
myrmecocalyx, *Hiern, l. c.* 139.—Borneo.
myrmecocarpa, *Hiern, l. c.* 141.—Bras.
natalensis, *Harv. Thes. Cap.* ii. 7.—Afr. austr.
neilgerrensis, Wight, Ic. iv. 10 = buxifolia.
nigrescens, *Dalz. in Dalz. & Gibs. Bomb. Fl.* 142.—Ind. or.
oblongifolia, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 112.—Zeylan.
obovata, *R. Br. Prod.* 527.—Austral.
ovalifolia, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 113.—Zeylan.
Pavonii, *Hiern, l. c.* 129.—Am. trop.
pentamera, F. Muell. Fragm. v. 163 = Diospyros pentamera.
punctata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 136.—Borneo.
quadridentata, F. Muell. Fragm. v. 162 = Diospyros mabacea.
quiloënsis, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 132.—Afr. trop.
reticulata, *R. Br. Prod.* 528.—Austral.
revoluta, Vieill. ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 114 = rufa.
rosea, *Montr. in Mém. Acad. Lyon*, x. (1860) 231.—N. Caled.
rufa, *Labill. Sert. Austro-Caled.* 33. t. 36.—Austral.; N. Caled.
ruminata, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 135.—N. Caled.
salicifolia, *Hiern, l. c.* 129.—Am. trop.
samoënsis, *Hiern, in Journ. Bot.* xv. (1877) 99.—Ins. Samoa.
sandwicensis, *A. DC. in DC. Prod.* viii. 242.—Ins. Sandvic.
sericea, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 140.—Bras.
sericocarpa, F. Muell. Fragm. v. 164 = rufa.
seychellarum, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 130.—Ins. Seychelles.
Smeathmanni, A. DC. in DC. Prod. viii. 241 = buxifolia.
sumatrana, *Miq. Pl. Jungh.* 204.—Sumatra.
Teijsmanni, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 137.—Malaya.
vacciniaefolia, Benth. in Hook. Niger Flora, 442 = buxifolia.
Vieillardi, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 124.—N. Caled.

MABEA, Aubl. Pl. Gui. ii. 867. t. 334 (1775). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 331.
angustifolia, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 365.—Bras.
anomala, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 526.—Bras.
biglandulosa, *Baill. Etud. Gén. Euph.* 413, nomen.—Guiana.
brasiliensis, *Muell. Arg. in DC. Prod.* xv. ii. 1151.—Bras.
ferruginea, Benth. in Hook. Kew Journ. vi. (1854) 366 = fistulifera.
fistulifera, Benth. l. c. 365 = angustifolia.
fistulifera, *Mart. Reise Bras.* 479.—Bras.
Gaudichaudiana, *Baill. Adansonia*, iv. (1863–64) 372.—Bras.
lactescens, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 523.—Bras.
maynensis, *Spruce, in Journ. Linn. Soc.* v. (1861) 7; *Muell. Arg. in DC. Prod.* xv. ii. 1150.—Peruv.
montana, *Muell. Arg. l. c.* 1151.—N. Granat.
nitida, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 367.—Bras.

MABEA:—
occidentalis, *Benth. in Hook. Kew Journ.* vi. (1854) 364.—Bras.
pallida, Baill. Adansonia, iv. (1863–64) 371 = nitida.
pallida, *Muell. Arg. in DC. Prod.* xv. ii. 1150.—N. Granat.
paniculata, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 367.—Bras.
paraguensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 527.—Bras.
Piriri, *Aubl. Pl. Gui.* ii. 867. t. 334.—Guian.
Piriri, Benth. Bot. Voy. Sulph. 165 = occidentalis.
Pohliana, *Muell. Arg. in DC. Prod.* xv. ii. 1152.—Bras.
pulcherrima, *Muell. Arg. in Flora*, lv. (1872) 44.—Guatemala.
Riedelii, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 518.—Bras.
rubrinervis, Baill. Adansonia, iv. (1863–64) 371 = nitida.
Schomburgkii, *Benth. in Hook. Kew Journ.* vi. (1854) 365.—Guian.
speciosa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 520.—Bras.
subserrulata, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 366.—Bras.
surinamensis, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1185, nomen.—Guian.
Taquari, *Aubl. Pl. Gui.* ii. 870. t. 334.—Guian.
volubilis, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1098, nomen.—Guian.
Weddelliana, Baill. Adansonia, iv. (1863–64) 371 = nitida.

MABOLA, Rafin. Sylva Tellur. 11 (1838) = **Diospyros**, Linn. (Ebenac.).
edulis, Rafin. l. c. = D. discolor.

MABURNIA, Thou. Gen. Nov. Madag. 4 (1806) = **Burmannia**, Linn.

MACADAMIA, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 72, cum ic. *PROTEACEAE*, Benth. & Hook. f. iii. 178.
ternifolia, *F. Muell. l. c.*—Austral.
verticillata, *F. Muell. ex Benth. Fl. Austral.* v. 407.—Austral.
Youngiana, *F. Muell. ex Benth. l. c.* 406.—Austral.

MACAGLIA, Rich. ex Vahl, in Skrivt. Nat. Selsk. Kiobenh. vi. (1810) 107 = **Aspidosperma**, Mart. & Zucc. (Apocyn.).
alba, Vahl, l. c.—Guian.
tuberculata, Vahl, l. c. 109.—Guian.

MACAHANEA, Aubl. Pl. Gui. Suppl. 6. t. 371 (1775) = **Salacia**, Linn. (Celastr.).
guianensis, Aubl. l. c. (= *Clercia guianensis*).—Guian.

MACAIREA, DC. Prod. iii. 109 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 742.
adenostemon, *DC. Prod.* iii. 109.—Bras.
albiflora, *Cogn. in Mart. Fl. Bras.* xiv. iii. 246.—Bras.
bracteolaris, Triana, in Trans. Linn. Soc. xxviii. (1871) 38 = Marcetia bracteolaris.
calvescens, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1856) 35.—Guian.
ledifolia, *Cogn. in Mart. Fl. Bras.* xiv. iii. 245.—Bras.
Mosenii, *Cogn. l. c.* 242.—Bras.
multinervia, *Benth. in Hook. Journ. Bot.* ii. (1840) 291.—Am. trop.
pachyphylla, *Benth. l. c.* 292.—Guian.
pachyphylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 37 = calvescens.
parvifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 293.—Bras.
Radula, *DC. Prod.* iii. 109.—Bras.
rigida, *Benth. in Hook. Journ. Bot.* ii. (1840) 292.—Guian.
rufescens, *DC. Prod.* iii. 109.—Bras.
sericea, *Cogn. in Mart. Fl. Bras.* xiv. iii. 243.—Bras.
Spruceana, *Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 38.—Bras.
stylosa, *Triana, l. c.*—Bras.
sulcata, *Triana, l. c.*—Peruv.
thyrsiflora, *DC. Prod.* iii. 109.—Bras.

MACANANGA, Reichb. Consp. 195 (1828) = **Macaranga**, Thou. (Euphorb.).

MACANEA, Juss. Gen. 257 (1789) = MACAHANEA, Aubl. = **Salacia**, Linn. (Celastr.).

MACARANGA, Thou. Gen. Nov. Madag. 26 (1806). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 320.
Mappa, A. Juss. Euph. Tent. 44. t. 14 (1824).
Mecostylis, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Indie, xxvii. (1864) 44.
Pachystemon, Blume, Bijdr. 626 (1825).
Panopia, Noronha, ex Thou. Gen. Nov. Madag. 26 (1806).
aleuritoides, *F. Muell. Papuan Pl.* 21.—N. Guin.
alnifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 256.—Madag.
amboinensis, *Muell. Arg. in DC. Prod.* xv. ii. 1002.—Amboina.
andamanica, *Kurz, For. Fl. Brit. Burma*, ii. 389.—Ins. Andam.
angolensis, *Muell. Arg. in DC. Prod.* xv. ii. 994.—Afr. trop.
asterolasia, F. Muell. Fragm. iv. 140 = involucrata.
Bailloniana, *Muell. Arg. in DC. Prod.* xv. ii. 1013.—Ins. Comor.
bancana, *Muell. Arg. l. c.* 990.—Ins. Banca.
Barteri, *Muell. Arg. in Flora*, xlvii. (1864) 535.—Afr. trop.
bicolor, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 199.—Ins. Philipp.
borneensis, Muell. Arg. in DC. Prod. xv. ii. 1003 = trichocarpa.
boutonioides, *Baill. Etud. Gén. Euph.* 432, nomen; *Adansonia*, i. (1860–61) 265.—Ins. Comor.
caladifolia, *Becc. Malesia*, ii. (1884) 46.—Borneo.
Chatiniana, *Muell. Arg. in DC. Prod.* xv. ii. 996.—Hab. ?
ciliata, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 256 = cupularis.
cissifolia, *Muell. Arg. in DC. Prod.* xv. ii. 993.—Sumatra.
conifera, *Muell. Arg. l. c.* 1005.—Sumatra.
cordifolia, Boiv. ex Baill. Etud. Gén. Euph. 432 = Bailloniana.
cordifolia, *Muell. Arg. in DC. Prod.* xv. ii. 1002.—Ins. Molucc.
coriacea, *Muell. Arg. l. c.* 1006.—N. Caled.
cornuta, *Muell. Arg. l. c.* 988.—Java.
corymbosa, *Muell. Arg. l. c.* 1006.—N. Caled.
Cumingii, *Muell. Arg. l. c.* 1005.—Ins. Philipp.
cuneifolia, *Muell. Arg. l. c.* 1004.—Ins. Banca.
cupularis, *Muell. Arg. in Flora*, xlvii. (1864) 466.—Madag.
cuspidata, *Boiv. ex Baill. Etud. Gén. Euph.* 432, nomen; *Adansonia*, i. (1860–61) 260.—Madag.
Dallachyi, *F. Muell. ex Benth. Fl. Austral.* vi. 144.—Austral.
denticulata, *Muell. Arg. in DC. Prod.* xv. ii. 1000.—Reg. Himal.; Malaya.
depressa, *Muell. Arg. l. c.* 989.—Malaya.
Diepenhorstii, *Muell. Arg. l. c.* 998.—Sumatra.
digyna, *Muell. Arg. l. c.* 1007.—Zeylan.
dioica, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 199.—Ins. Pacif.
divergens, *Muell. Arg. in DC. Prod.* xv. ii. 989.—Borneo.
echinocarpa, *Baker, in Journ. Linn. Soc.* xx. (1883) 255.—Madag.
eglandulosa, Baill. Etud. Gén. Euph. 432, nomen = Bailloniana.
flexuosa, Wight, Ic. v. ii. 23 = indica.
gigantea, *Muell. Arg. in DC. Prod.* xv. ii. 995.—Sumatra.
Grayana, *Muell. Arg. l. c.* 1001.—Ins. Pacif.
Griffithiana, *Muell. Arg. l. c.* 993.—Malaya.
gummiflua, Muell. Arg. l. c. 1000 = denticulata.
Harveyana, *Muell. Arg. l. c.* 998.—Ins. Pacif.
Helferi, Muell. Arg. l. c. 1004 = minutiflora.
heterophylla, *Muell. Arg. l. c.* 993.—Afr. occ.
Heudelotii, *Baill. Adansonia*, i. (1860–61) 69.—Afr. trop.
hexandra, *Muell. Arg. in DC. Prod.* xv. ii. 1003.—Malaya.
hispida, *Muell. Arg. l. c.* 990.—Ins. Molucc.
hypoleuca, *Muell. Arg. l. c.* 992.—Malaya.
inamoena, *F. Muell. ex Benth. Fl. Austral.* vi. 145.—Austral.
indica, *Wight, Ic.* t. 1883.—Ind. or.
involucrata, *Baill. Etud. Gén. Euph.* 432, nomen; *Muell. Arg. in DC. Prod.* xv. ii. 1011.—Ins. Molucc. ?; Austral.

MACARANGA:—
javanica, *Muell. Arg. in DC. Prod.* xv. ii. 1004.—Malaya.
leptostachya, Muell. Arg. l. c. 1007 (Cleidionis sp.).—Ins. Pacif.
longestipulata, *Muell. Arg. l. c.* 991.—Ins. Molucc.
macrophylla, *Muell. Arg. l. c.*—Ins. Fiji.
macropoda, *Baker, in Journ. Linn. Soc.* xx. (1883) 257.—Madag.
madagascariensis, *Steud. Nom.* ed. II. ii. 86, nomen.—Madag.
mallotoides, F. Muell. Fragm. iv. 139 = involucrata.
Mappa, *Muell. Arg. in DC. Prod.* xv. ii. 1000.—Malaya.
Maudslayi, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 371, nomen.—Ins. Fiji.
mauritiana, *Boj. Hort. Maurit.* 283; *Baill. Etud. Gén. Euph.* 432. t. 21. f. 8–9.—Ins. Maurit.
megalophylla, *Muell. Arg. in DC. Prod.* xv. ii. 995.—Malaya.
membranacea, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 246.—Burma.
membranacea, *Muell. Arg. in DC. Prod.* xv. ii. 996.—Ins. Fiji.
minutiflora, *Muell. Arg. in Flora*, xlvii. (1864) 466.—Burma.
molliuscula, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 245 = M. Tanarius.
monandra, *Muell. Arg. in Journ. Bot.* ii. (1864) 337.—Afr. trop.
Motleyana, *Muell. Arg. in DC. Prod.* xv. ii. 994.—Borneo.
myriantha, *Muell. Arg. l. c.* 1008.—Amboina.
myriolepida, *Baker, in Journ. Linn. Soc.* xxi. (1885) 442.—Madag.
oblongifolia, *Baill. Etud. Gén. Euph.* 432. t. 21. f. 5–7.—Madag.
obovata, *Boiv. ex Baill. l. c.* nomen; *Adansonia*, i. (1860–61) 263.—Madag.
occidentalis, *Muell. Arg. in DC. Prod.* xv. ii. 994.—Afr. trop.
ovata, Boiv. ex Baill. Etud. Gén. Euph. 432, nomen; Adansonia, i. (1860–61) 266 = boutonioides.
pachyphylla, *Muell. Arg. in DC. Prod.* xv. ii. 999.—Sumatra.
peltata, Boiv. ex Baill. Etud. Gén. Euph. 432, nomen = cuspidata.
peltata, Muell. Arg. in DC. Prod. xv. ii. 1010 = Roxburghii.
populifolia, *Muell. l. c.* 1006.—Sumatra.
pruinosa, *Muell. Arg. l. c.* 992.—Sumatra.
reticulata, Baill. Etud. Gén. Euph. 432 = obovata.
rhizinoides, *Muell. Arg. in DC. Prod.* xv. ii. 1011.—Java.
ribesioides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 442.—Madag.
rottleroides, *Baill. Etud. Gén. Euph.* 432, nomen; *Adansonia*, i. (1860–61) 262.—Madag.
Roxburghii, *Wight, Ic.* v. ii. 23. t. 1949. f. 4.—Ind. or.
rugosa, Muell. Arg. in DC. Prod. xv. ii. 995 = megalophylla.
Sampsoni, *Hance, in Journ. Bot.* ix. (1871) 134.—China.
secunda, *Muell. Arg. in DC. Prod.* xv. ii. 996.—Ins. Fiji.
Seemanni, *Muell. Arg. l. c.* 999.—Ins. Pacif.
sinensis, *Muell. Arg. l. c.* 1001.—China.
sphaerophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 257.—Madag.
spinosa, *Muell. Arg. in Flora*, xlvii. (1864) 466.—Afr. trop.
stipulosa, *Muell. Arg. in DC. Prod.* xv. ii. 1001.—Ins. Pacif.
stricta, *Muell. Arg. l. c.* 1004.—Sumatra.
subdentata, *Benth. Fl. Austral.* vi. 145.—Austral.
subfalcata, *Muell. Arg. in DC. Prod.* xv. ii. 1007.—Java.
sumatrana, *Muell. Arg. l. c.* 1003.—Sumatra.
taitensis, *Muell. Arg. l. c.* 997.—Ins. Pacif.
Tanarius, *Muell. Arg. l. c.*—Malaya.
tenuifolia, *Muell. Arg. l. c.* 990.—Java.
Teysmanni, *Muell. Arg. l. c.* 999.—Sumatra.
Thouarsii, *Baill. Etud. Gén. Euph.* 432, nomen.—Afr. trop.
tomentosa, Wight, Ic. v. ii. 23. t. 1949. f. 1 = Roxburghii.
trichocarpa, *Muell. Arg. in DC. Prod.* xv. ii. 1003.—Malaya.
triloba, *Muell. Arg. l. c.* 989.—Malaya.
Vedeliana, *Muell. Arg. l. c.* 1002.—N. Caled.

MACARANGA:—
velutina, *Muell. Arg. in DC. Prod.* xv. ii. 993.—Sumatra.
Vieillardi, *Muell. Arg. l. c.* 1007.—N. Caled.
Wightiana, Baill. Etud. Gén. Euph. 432 = Roxburghii.

MACARISIA, Thou. Hist. Vég. Afr. 49. t. 14 (1807). *RHIZOPHOREAE*, Benth. & Hook. f. i. 682.
icosandra, Planch. ex Hook. f. Fl. Brit. Ind. i. 416 = Ixonanthes icosandra.
lanceolata, *Baill. in Adansonia*, iii. (1862–63) 20.—Ins. Nossibé.
pyramidata, *Thou. Hist. Vég. Afr.* 49. t. 14.—Madag.

MACARTHURIA, Hueg. ex Endl. in Enum. Pl. Hueg. 11 (1837). *FICOIDEAE*, Benth. & Hook. f. i. 856.
apetala, *Harv. in Hook. Kew Journ.* vii. (1855) 55.—Austral.
australis, *Hueg. ex Endl. in Enum. Pl. Hueg.* 11.—Austral.
foliosa, Preiss. ex Steud. in Lehm. Pl. Preiss. i. 230 = australis.
neo-cambrica, *F. Muell. Fragm.* v. 28.—Austral.

MACBRIDEA, Ell. ex Nutt. Gen. Am. ii. 36 (1818); Sketch, ii. 86 (1824). *LABIATAE*, Benth. & Hook. f. ii. 1204.
alba, *Chapm. Fl. S. U. St.* 325.—Am. bor.
pulchella, Benth. Lab. Gen. et Sp. 505 = pulchra.
pulchra, *Ell. ex Nutt. Gen. Am.* ii. 36; *Sketch*, ii. 86.—Am. bor.

MACCLELLANDIA, Wight, Ic. vi. 14. t. 1996 (1853) = **Pemphis**, Forst. (Lythrar.).
Griffithiana, Wight, l. c. = P. acidula.

MACCOYA, F. Muell. Fragm. i. 127 (1859) = **Rochelia**, Reichb. (Borag.).
plurisepalea, F. Muell. l. c. = R. Maccoya.

MACDONALDIA, Gunn, ex Lindl. Swan River App. 50. t. 9 (1839) = **Thelymitra**, Forst. (Orchid.).
antennifera, Lindl. l. c. = T. antennifera.
concolor, Lindl. l. c. t. 9 = T. flexuosa.
cyanea, Lindl. l. c. 50 = T. cyanea.
Smithiana, Gunn, ex Lindl. l. c. = T. flexuosa.
spiralis, Lindl. l. c. = T. variegata.
variegata, Lindl. l. c. = T. variegata.
venosa, Lindl. l. c. = T. venosa.

MACELLA, C. Koch, in Ind. Sem. Hort. Berol. (1855) App. 13 = **Jaegeria**, H. B. & K. (Compos.).
hirta, C. Koch, l. c. = J. hirta.

MACFADYENA, A. DC. in DC. Prod. ix. 179 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1034.
Dolichandra, Cham. in Linnaea, vii. (1832) 657.
Pharseophora, Miers, in Proc. Hort. Soc. iii. (1863) 185.
Phryganocydia, Mart. ex DC. Prod. ix. 198, etc. (1845).
amanda, *Miers, in Proc. Hort. Soc.* iii. (1863) 200, nomen.—Bras.
amazonica, *Miers, l. c.* nomen.—Bras.
bifolia, *Miers, l. c.*—Ind. occ.
bracteosa, *Benth. & Hook. f. Gen.* ii. 1035.—Guiana.
citrifolia, *Miers, in Proc. Hort. Soc.* iii. (1863) 200.—Bras.
coccinea, *Miers, l. c.*—Bras.
Coito, *Miers, l. c.*—Bras.
corymbosa, *Griseb. in Bonplandia*, vi. (1858) 10.—Ins. Trinit.
diffusa, *Miers, in Proc. Hort. Soc.* iii. (1863) 200, nomen.—Bras.
Dolichandra, *Benth. & Hook. f. Gen.* ii. 1034.—Bras.
Fenzliana, *Miq. Stirp. Surin. Sel.* 125.—Guian.
Fockeana, *Miers, in Proc. Hort. Soc.* iii. (1863) 200.—Guian.
guttiflua, *Miers, l. c.* nomen.—Bras.
hispida, *Seem. in Journ. Bot.* i. (1863) 227.—Bras.
japurensis, *Miers, in Proc. Hort. Soc.* iii. (1863) 200.—Bras.
laurifolia, *Miers, l. c.*—Venezuela.
lepidota, Seem. ex Miers, l. c. = corymbosa.
mollis, *Seem. in Journ. Bot.* i. (1863) 227.—Bras.
obovata, *Miers, in Proc. Hort. Soc.* iii. (1863) 200.—N. Granat.
orinocensis, *Miers, l. c.*—Venezuela.
ovata, Miers, l. c. = corymbosa?
phellosperma, *Hemsl. Biol. Centr. Am. Bot.* ii. 492.—Panama.

MACFADYENA :—
platypoda, *Miers, in Proc. Hort. Soc.* iii. (1863) 200.—Bras.
podopogon, *Griseb. Cat. Pl. Cub.* 195.—Cuba.
stipuloidea, *Miers, in Proc. Hort. Soc.* iii. (1863) 200, nomen.—Bras.
terniflora, *Miers, l. c.*—Bras.
Tweediana, Griseb. Veg. Prov. Entre-Rios, 83 = Bignonia Tweediana.
uncinata, *DC. Prod.* ix. 180.—Panama.

MACGREGORIA, F. Muell. in Caruel, Giorn. Bot. Ital. (1873) 128 ; et Fragm, viii. 160 (1874). *STACKHOUSIEAE.*
racemigera, *F. Muell. ll. cc.* 128 ; 161.—Austral.
racemosa, Benth. in Hook. Ic. Pl. xiii. 23. t. 1230 = racemigera.

MACHADOA, Welw. ex Benth. & Hook. f. Gen. i. 814 (1867). *PASSIFLOREAE*, Benth. & Hook. f. i. 814.
huillensis, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 29. t. 10.—Afr. trop.

MACHAERANTHERA, Nees, Gen. et Sp. Ast. 224 (1833) = **Aster**, Tourn. (Compos.).
canescens, A. Gray, Pl. Wright. i. 89 ; ii. 75 = A. canescens.
grandiflora, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = Xanthisma texanum.
parviflora, A. Gray, Pl. Wright. i. 90 = A. parviflora.
setigera, Nees, in Linnaea, xix. (1847) 722 = A. gymnocephalus.
shastensis, A. Gray, in Proc. Am. Acad. vi. (1864–65) 538 = A. shastensis.
tanacetifolia, Nees, Gen. et Sp. Ast. 225 = A. tanacetifolius.

MACHAERINA, Nees, in Lehm. Pugill. viii. 54 (1844) (Macherina) = **Lepidosperma**, Labill. (Cyperac.).
resinosa, Nees, l. c. = L. resinosum.

MACHAERINA, Vahl, Enum. ii. 238 (1806) = **Cladium**, P. Br. (Cyperac.).
anceps, Boj. Hort. Maurit. 385 = C. iridifolium.
filifolia, Griseb. Cat. Pl. Cub. 242 = C. filifolium.
hirta, Boeck. in Flora, lviii. (1875) 117.—Ins. Samoa.
lavarum, Boj. Hort. Maurit. 385 = C. iridifolium.
restioides, Gaudich. in Freyc. Voy. Bot. 104, 417 (= *Vincentia angustifolia*).—Ins. Sandvic.
restioides, Vahl, Enum. ii. 238.—Ind. occ.

MACHAERIUM, Pers. Syn. ii. 276 (1807). *LEGUMINOSAE*, Benth. & Hook. f. i. 545.
acaciaefolium, Mart. ex Benth. in Mart. Fl. Bras. xv. 1. 236 = angustifolium.
aculeatum, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 398.—Bras.
acuminatum, *H. B. & K. Nov. Gen. et Sp.* vi. 391.—Venezuela.
acutifolium, *Vog. in Linnaea*, xi. (1837) 187.—Bras.
affine, Benth. in Ann. Wien. Mus. ii. (1838) 98 = angustifolium.
Allemani, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 63 ; *et in Mart. Fl. Bras.* xv. 1. 248.—Bras.
amplum, *Benth. in Ann. Wien. Mus.* ii. (1838) 97.—Bras.
angustifolium, Mart. ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 57 = eriocarpum.
angustifolium, *Vog. in Linnaea*, xi. (1837) 193.—Bras.
arboreum, *Vog. l. c.* 182 ; *Benth. in Ann. Wien. Mus.* ii. (1838) 100.—Am. austr.
armatum, Vog. l. c. 191 = angustifolium.
bracteatum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 65.—N. Granat.
brasiliense, *Vog. in Linnaea*, xi. (1837) 185.—Bras.
campestre, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 100 = opacum.
ciliatum, Benth. l. c. 99 = brasiliense.
densicomum, *Mart. ex Benth. l. c.*—Bras.
diadelphum, Vog. in Linnaea, xi. (1837) 182 = ferrugineum.
discolor, *Vog. l. c.* 204.—Bras.
dubium, *Allem. in Trab. Soc. Vell.* [1852] 56.—Bras.
erianthum, *Benth. in Ann. Wien. Mus.* ii. (1838) 99.—Bras.
eriocarpum, *Benth. l. c.* 98.—Bras.
eriostemon, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 68 ; *et in Mart. Fl. Bras.* xv. 1. 254.—Bras.

MACHAERIUM :—
eriostemum, Benth. ex Micheli, in Kjoeb. Vidensk. Meddel. (1875) 99 = praec. ?
ferrugineum, *Pers. Syn.* ii. 276.—Guiana.
fertile, *Griseb. in Goett. Abh.* xix. (1874) 127.—Reg. Argent.
firmum, *Benth. in Ann. Wien. Mus.* ii. (1838) 101.—Bras.
floribundum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 68 ; *et in Mart. Fl. Bras.* xv. 1. 254.—Bras.
Gardneri, *Benth. ll. cc.* 60 ; 242.—Bras.
glabrum, *Vog. in Linnaea*, xi. (1837) 187.—Bras.
Goudoti. *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 59.—N. Granat.
gracile, *Benth. in Ann. Wien. Mus.* ii. (1838) 98.—Bras.
heterophyllum, Presl, Bot. Bemerk. 61 = oblongifolium.
heteropterum, Allem. in Trab. Soc. Vell. [1852] 4 = Tipuana heteroptera.
Humboldtianum, *Vog. in Linnaea*, xi. (1837) 194.—Am. austr.
incorruptibile, *Allem. in Trab. Soc. Vell.* [1852] 56.—Bras.
incorruptibile, Benth. in Ann. Wien. Mus. ii. (1838) 101 = legale.
Kegelii, *Meissn. in Linnaea*, xxi. (1848) 257.—Guiana.
lanatum, *Tul. in Arch. Mus. Par.* iv. (1844) 96.—Bras.
legale, *Benth. in Ann. Wien. Mus.* ii. (1838) 101.—Bras.
leiocarpum, Vog. in Linnaea, xi. (1837) 203 = secundiflorum.
leiophyllum, *Benth. in Ann. Wien. Mus.* ii. (1838) 100.—Guiana.
leucopterum, *Vog. in Linnaea*, xi. (1837) 189.—Bras.
lignosum, Benth. in Mart. Fl. Bras. xv. 1. 253 = ferrugineum.
Lindenianum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 67.—N. Granat.
lineatum, Benth. in Ann. Wien. Mus. ii. (1838) 98 = discolor.
longifolium, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 55 ; *et in Mart. Fl. Bras.* xv. 1. 235.—Bras.
Luschnathianum, Presl, Bot. Bemerk. 61 = brasiliense.
macrocarpum, Benth. in Hook. Journ. Bot. ii. (1840) 67 = Tipuana macrocarpa.
macrophyllum, *Benth. in Ann. Wien. Mus.* ii. (1838) 99.—Bras.
macrophyllum, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 68 = floribundum.
Martii, Tul. in Arch. Mus. Par. iv. (1844) 92 = eriocarpum.
micropterum, *Benth. in Ann. Wien. Mus.* ii. (1838) 101.—Hort. Teneriff.
minutiflorum, Tul. in Arch. Mus. Par. iv. (1844) 94 = stipitatum.
Moritzianum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 58.—Venezuela.
mucronatum, Vog. in Linnaea, xi. (1837) 191 = incorruptibile.
mucronulatum, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 100.—Bras.
muticum, Benth. l. c. = acutifolium.
myrianthum, *Spruce, ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 59 ; *et in Mart. Fl. Bras.* xv. 1. 241.—Bras.
nervosum, Vog. in Linnaea, xi. (1837) 186 = ferrugineum.
nictitans, *Benth. in Ann. Wien. Mus.* ii. (1838) 98.—Bras.
nigrum, *Vog. in Linnaea*, xi. (1837) 188.—Bras.
oblongifolium, *Vog. l. c.* 184.—Bras.
opacum, *Vog. l. c.* 187.—Bras.
paniculatum, Allem. ex Benth. in Mart. Fl. Bras. xv. 1. 246 = stipitatum.
parviflorum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 60 ; *et in Mart. Fl. Bras.* xv. 1. 242.—Bras.
pedicellatum, *Vog. in Linnaea*, xi. (1837) 202.—Bras.
pilosum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 57 ; *et in Mart. Fl. Bras.* xv. 1. 238.—Bras.
polyphyllum, *Benth. in Ann. Wien. Mus.* ii. (1838) 101.—Guian.
Pseudotipe, *Griseb. in Goett. Abh.* xxiv. (1879) 110.—Reg. Argent.
puberulum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 99 = oblongifolium.
punctatum, Pers. Syn. ii. 276 = stipitatum.
pungens, Allem. in Trab. Soc. Vell. [1852] 56 = leucopterum.

MACHAERIUM :—
reticulatum, *Pers. Syn.* ii. 276.—Madag.
retusum, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1884) 436 = Dalbergia retusa.
robiniaefolium, *Vog. in Linnaea*, xi. (1837) 190.—Ins. S. Vincent.
Salzmanni, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 63 ; *et in Mart. Fl. Bras.* xv. 1. 248.—Bras.
Schomburgkii, *Benth. in Hook. Journ. Bot.* ii. (1840) 67.—Guian.
scleroxylon, *Tul. in Arch. Mus. Par.* iv. (1844) 93.—Bras.
scleroxylum, Allem. in Trab. Soc. Vell. [1852] 56 = Allemani.
secundiflorum, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 118.—Bras.
Seemanni, *Benth. ex Seem. Bot. Voy. Herald*, 110.—Panama.
sericiflorum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 341 = Gardneri.
sericiflorum, Vog. in Linnaea, xi. (1837) 192 = aculeatum.
Sieberi, Benth. in Ann. Wien. Mus. ii. (1838) 98 = robiniaefolium.
sordidum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 58 ; *et in Mart. Fl. Bras.* xv. 1. 241.—Bras.
spinosum, *Benth. in Ann. Wien. Mus.* ii. (1838) 99.—Bras.
splendens, *Vog. in Linnaea*, xi. (1837) 192.—Bras.
stipitatum, *Vog. l. c.* 189.—Bras.
surinamense, Miq. Stirp. Surin. Sel. 21 = leiophyllum.
Tipu, Benth. in Hook. Kew Journ. v. (1853) 267 = Tipuana speciosa.
triste, *Vog. in Linnaea*, xi. (1837) 416.—Bras.
uncinatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 98.—Bras.
Vellosianum, *Benth. l. c.*—Bras.
velutinum, Benth. l. c. 100 = nigrum.
verrucosum, Vog. in Linnaea, xi. (1837) 183 = oblongifolium.
vestitum, *Vog. l. c.* 190.—Bras.
villosulum, Mart. in Flora, xx. (1837) ii. Beibl. 118 = oblongifolium.
villosum, *Vog. in Linnaea*, xi. (1837) 189.—Bras.
violaceo-purpureum, Duchass. & Walp. Ann. ii. 437 = angustifolium.
violaceum, *Vog. in Linnaea*, xi. (1837) 186.—Bras.

MACHAEROPHORUS, Schlecht. in Linnaea, xxviii. (1856) 469 = **Mathewsia**, Hook. (Crucif.).
matthioloides, Schlecht. l. c. = Mathewsia matthioloides.

MACHANAEA, Steud. Nom. ed. I. 503 (1821) = MACAHANEA, Aubl. = **Salacia**, Linn. (Celastr.).

MACHAONIA, Humb. & Bonpl. Pl. Aequin. i. 101. t. 29 (1808). *RUBIACEAE*, Benth. & Hook. f. ii. 102.
acuminata, *Humb. & Bonpl. Pl. Aequin.* i. 101.—Ecuador.
brasiliensis, *Cham. & Schlecht. in Linnaea*, iv. (1829) 2.—Bras.
cymosa, *Griseb. Fl. Brit. W. Ind.* 348.—Ind. occ.
fasciculata, *A. Gray, in Proc. Am. Acad.* xix. (1883) 77.—Mexic.
Galeottiana, *Baill. in Bull. Soc. Linn. Par.* i. (1879) 204.—Mexic.
Hahniana, *Baill. l. c.*—Mexic.
Lindeniana, *Baill. l. c.*—Mexic.
microphylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 510.—Ind. occ.
portoricensis, *Baill. in Bull. Soc. Linn. Par.* i. (1879) 204.—Porto Rico.
rotundata, *Griseb. Fl. Brit. W. Ind.* 348.—Ind. occ.
spinosa, *Cham. & Schlecht. in Linnaea*, iv. (1829) 2.—Bras.
velutina, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 134.—Mexic.
veracruzeana, Baill. in Bull. Soc. Linn. Par. i. (1879) 204 = velutina.

MACHARINA, Steud. Syn. Pl. Cyp. 156 (1855) = MACHAERINA, Vahl = **Cladium**, P. Br. (Cyperac.).

MACHARISIA, Spreng. Syst. iii. 66 (1826) = **Macarisia**, Thou. (Rhizoph.).

MACHILUS, Nees, in Wall. Pl. As. Rar. ii. 70 (1831). *LAURINEAE*, Benth. & Hook. f. iii. 156.
bootanica, *Meissn. in DC. Prod.* xv. 1. 42.—Reg. Himal.
cuneata, Hassk. Cat. Hort. Bog. Alt. 88 = Phoebe cuneata.
fruticosa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 101.—Burma.
glaucescens, Wight, Ic. t. 1825 = macrantha.
Grijsii, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 226.—China.
incrassata, Nees, in Wall. Pl. As. Rar. ii. 70 = Dehaasia microcarpa.
indica, Kurz, For. Fl. Brit. Burma, ii. 291 = odoratissima.
japonica, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1843) 302.—Japon.
khasyana, *Meissn. in DC. Prod.* xv. 1. 42.—Reg. Himal.
longifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 331.—Japon.
macrantha, *Nees, in Wall. Pl. As. Rar.* ii. 70.—Ind. or.
macrophylla, *Blume, Mus. Bot. Lugd. Bat.* i. 331.—Java.
macrophylla, Nees, Syst. Laurin. 674 = Phoebe macrophylla, cuneata.
Madang, Miq. in Pl. Jungh. i. 176 = odoratissima.
odoratissima, *Nees, in Wall. Pl. As. Rar.* ii. 70.—Ind. or.; Malaya; China.
oreophila, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 227.—China.
parviflora, *Meissn. in DC. Prod.* xv. 1. 505.—Reg. Himal.
peduncularis, Nees, in Wall. Pl. As. Rar. ii. 70 (= *Haasia peduncularis*).—Ind. or.
pilosa, *Nees, Syst. Laurin.* 176.—Cochinch.
pubescens, *Blume, Mus. Bot. Lugd. Bat.* i. 330.—Nepal.
rimosa, *Blume, l. c.*—Java.
salicina, *Hance, in Journ. Bot.* xxiii. (1885) 327.—China.
sericea, *Blume, Mus. Bot. Lugd. Bat.* i. 330—Reg. Himal.
tavoyana, Meissn. in DC. Prod. xv. 1. 41 (Phoebes sp.).—Burma.
Thunbergii, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1843) 302.—Japon.
velutina, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 198.—Hongkong.

MACHLIS, DC. Prod. vi. 140 (1837) = **Cotula**, Tourn. (Compos.).
hemisphaerica, DC. l. c. = C. hemisphaerica.

MACHURA, Steud. Nom. ed. I. 503 (1821), sphalm. = **Maclura**, Nutt. (Urtic.).

MACIELIA, Vand. Fl. Lusit. 14 (1788) = **Cordia**, Linn. (Borag.).
arborea, Steud. Nom. ed. II. ii. 87 = *racemifera*.
racemifera, Vand. Fl. Lusit. 15.—Bras.

MACINTYRIA, F. Muell. Fragm. v. 8 (1865) = **Xanthophyllum**, Roxb. (Polygal.).
octandra, F. Muell l. c. = X. Macintyrii.

MACKAYA, Arn. in Jard. Mag. Zool. & Bot. ii. (1838) 550 = **Erythropalum**, Blume (Olacin.).
populifolia, Arn. l. c. = E. populifolium.

MACKAYA, Harv. Thes. Cap. i. 8. t. 13 (1859) = **Asystasia**, Blume (Acanthac.).
bella, Harv. l. c.; et in Proc. Dubl. Zool. & Bot. Assoc. i. (1859) 253. t. 26 = A. bella (in Addend.).

MACKENIA, Harv. Gen. S. Afr. Pl. ed. II. 233 (1868) = **Schizoglossum**, E. Mey. (Asclep.).

MACKENZIEA, Nees, in DC. Prod. xi. 308 (1845) = **Strobilanthes**, Blume (Acanthac.).
sessilis, Nees, l. c. = S. sexennis.

MACKINLAYA, F. Muell. Fragm. iv. 119 (1864). *ARALIACEAE*, Benth. & Hook. f. i. 937.
macrosciadea, *F. Muell. l. c.* 120.—Austral.

MACKLEYA, Walp. Rep. i. 109 (1842) = MACLEAYA, R. Br. = **Bocconia**, Linn. (Papav.).

MACKLOTTIA, Korth. in Nederl. Kruidk. Arch. i. (1847) 196 = **Leptospermum**, Forst. (Myrtac.).
amboinensis, Korth. l. c. = L. amboinense.
javanica, Korth. l. c. = L. javanicum.

MACLAYA, Bernh. in Linnaea, viii. (1833) 460 = MACLEAYA, R. Br. = **Bocconia**, Linn. (Papav.).

MACLEANIA, Hook. Ic. Pl. t. 109 (1837). *VACCINIACEAE*, Benth. & Hook. f. ii. 566.
BIRAMIA, Neraud, in Freyc. Voy. Bot. 28 (1826).
TYRIA, Klotzsch, in Linnaea, xxiv. (1851) 21.
angulata, *Hook. Bot. Mag.* t. 3979.—Peruv.
Benthamiana, *Walp. Rep.* vi. 415.—Peruv.
coccinea, *Decne. in Rev. Hortic.* Sér. III. v. (1851) 301.—Mexic.; Peruv.
colorata, Klotzsch, in Linnaea, xxiv. (1851) 68 = cordata.
cordata, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) 312.—Peruv.
cordifolia, *Benth. Pl. Hartw.* 223.—Peruv.
floribunda, Benth. l. c. 141 = Benthamiana.
floribunda, *Hook. Ic. Pl.* t. 109.—Peruv.
insignis, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 531.—Mexic.
longiflora, *Lindl. Bot. Reg.* (1844) t. 25.—Peruv.
macrantha, *Benth. Pl. Hartw.* 223.—Peruv.
ovata, Klotzsch, in Linnaea, xxiv. (1851) 20 = cordata.
pauciflora, *Benth. & Hook. f. Gen.* ii. 567, in textu.—Peruv.
pubiflora, *Benth. Pl. Hartw.* 224.—Reg. Granat.
pulchra, *Hook. f. Bot. Mag.* t. 5465.—N. Granat.
punctata, *Hook. l. c.* t. 4426.—Ecuador.
Salapa, *Benth. & Hook. f. Gen.* ii. 566.—Peruv.
speciosissima, *Hook. Bot. Mag.* t. 5453.—N. Granat.
tenuiflora, Klotzsch, in Linnaea, xxiv. (1851) 19 = longiflora.
tenuifolia, Walp. Rep. vi. 415 = longiflora.

MAC-LEAYA, Benth. & Hook. f. Gen. i. 573 (1865) = MAC-LEAYIA, Montr. = **Cassia**, Tourn. (Legum.).

MACLEAYA, R. Br. in App. Denh. & Clapp. Trav. 218, in adnot. (1826) = **Bocconia**, Linn. (Papav.).
cordata, R. Br. l. c. = B. cordata.
yedoënsis, André, in Rev. Hortic. (1866) 369 = B. cordata.

MAC-LEAYIA, Montr. in Mém. Acad. Lyon, x. (1860) 198 = **Cassia**, Tourn. (Legumin.).
multiflora, Montr. l. c.—N. Caled.

MACLEDIUM, Cass. in Dict. Sc. Nat. xxxiv. 39 (1825) = **Dicoma**, Cass. (Compos.).
Burmanni, Cass. l. c. = D. Burmanni.

MACLELANDIA, Wight, Ic. in tab. 1996 (1853) = MACCLELLANDIA, Wight = **Pemphis**, Forst. (Lythrar.).

MACLEYA, Reichb. Consp. 187 (1828) = MACLEAYA, R. Br. = **Bocconia**, Linn. (Papav.).

MACLURA, Nutt. Gen. Am. ii. 233 (1817). *URTICACEAE*, Benth. & Hook. f. iii. 363.
FUSTICUS, Rafin. New Fl. Am. iii. 43 (1836).
IOXYLON, Rafin. in Am. Monthly Mag. iv. (1818) 195.
SUKAMINEA, Rafin. New Fl. Am. iii. 44 (1836).
TOXYLON, Rafin. in Am. Monthly Mag. iv. (1818) 188. (Cf. 195.)
affinis, Miq. in Mart. Fl. Bras. iv. 1. 155 = Chlorophora tinctoria.
amboinensis, Blume, Mus. Bot. Lugd. Bat. ii. 84 = Cudrania javanensis.
aurantiaca, *Nutt. Gen. Am.* ii. 234.—Am. bor.
brasiliensis, *Endl. Gen. Suppl.* iv. ii. 34.—Bras.
Calcar-galli, Lodd. ex Voigt, Hort. Suburb. Calc. 291, nomen = Cudrania javanensis?
chlorocarpa, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 314 = Chlorophora tinctoria.
excelsa, Bureau, in DC. Prod. xvii. 231 = Chlorophora excelsa (in Addend.).
gerontogaea, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 220 = Cudrania javanensis.
javanica, Blume, Mus. Bot. Lugd. Bat. ii. 83. t. 31 = Cudrania javanensis.
javanica, Miq. in Zoll. Syst. Verz. Ind. Archip. 89 = Malaisia tortuosa.

MACLURA:—
Mora, *Griseb. in Goett. Abh.* xxiv. (1879) 86.—Reg. Argent.
Plumiera, D. Don, ex Steud. Nom. ed. II. ii. 87 = Chlorophora tinctoria.
polyneura, Miq. in Mart. Fl. Bras. iv. 1. 154 = Chlorophora tinctoria.
sempervirens, Tenore, Cat. Hort. Bot. Nap. (1845) 87 = Chlorophora tinctoria.
Sieberi, Blume, Mus. Bot. Lugd. Bat. ii. 83 = Chlorophora tinctoria.
subintegerrima, Miq. in Mart. Fl. Bras. iv. 1. 157 = Chlorophora tinctoria.
timorensis, Blume, Mus. Bot. Lugd. Bat. ii. 84 = Cudrania javanensis.
tinctoria, D. Don, ex Steud. Nom. ed. II. ii. 87 = Chlorophora tinctoria.
tricuspidata, *Carr. in Rev. Hortic.* (1864) 390.—China.
velutina, Blume, Mus. Bot. Lugd. Bat. ii. 82 = Chlorophora tinctoria.
xanthoxylon, Endl. Gen. Suppl. iv. ii. 34 = Chlorophora tinctoria.

MACNABIA, Benth. in DC. Prod. vii. 612 (1839). *ERICACEAE*, Benth. & Hook. f. ii. 589.
NABEA, Lehm. Ind. Sem. Hort. Hamb. (1831); ex Klotzsch, in Linnaea, viii. (1833) 666.
montana, *Benth. in DC. Prod.* vii. 612.—Afr. austr.

MACNEMARAEA, Willem. in Usteri, Ann. Bot. xviii. (1796) 34 (Genus dubium).
opulina, Willem. l. c.—China?

MACODES, Lindl. Gen. et Sp. Orch. 496 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 602.
lanceolata, Reichb. f. Xenia Orch. i. 226 = Hetaeria lanceolata.
marmorata, Reichb. f. l. c. 228. t. 96 = Dossinia marmorata.
Petola, *Lindl. Gen. et Sp. Orch.* 497.—Java.
setaceus, *Boxall, ex Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 249, nomen.—Ins. Philipp.
striatus, *Boxall, l. c.* nomen.—Ins. Philipp.
Veitchii, *Boxall, l. c.*—Ins. Philipp.
xanthophyllus, *Boxall, l. c.*—Ins. Philipp.

MACOUBEA, Aubl. Pl. Gui. Suppl. 17. t. 378 (1775). *INC. SED.*—Cf. Benth. & Hook. f. i. 169.
guianensis, *Aubl. l. c.* 18. t. 378.—Guiana.

MACOUCOUA, Aubl, Pl. Gui. i. 88. t. 34 (1775) = **Ilex**, Tourn.
guianensis, Aubl. l. c. = I. Macoucoua.

MACOWANIA, Oliver, in Hook. Ic. Pl. t. 1062 (1870). *COMPOSITAE*, Benth. & Hook. f. ii. 327.
revoluta, *Oliver, in Hook. Ic. Pl.* t. 1062.—Afr. austr.

MACPHERSONIA, Blume, Rumphia, iii. 156 (1847). *SAPINDACEAE*, Benth. & Hook. f. i. 402.
gracilis, *O. Hoffm. Sert. Madag.* 14.—Madag.
Hildebrandtii, *O. Hoffm. l. c.*—Madag.
madagascariensis, *Blume, Rumphia*, iii. 157.—Madag.
pteridophylla, *Baill. Adansonia*, xi. (1874) 240.—Ins. Nossibé.

MACQUINIA, Steud. Nom. ed. II. ii. 88 (1841) sphalm. = MOQUINIA, Spreng. = **Loranthus**, Linn.

MACRACHAENIUM, Hook f. Fl. Antarct. ii. 321 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 498.
gracile, *Hook. f. Fl. Antarct.* ii. 321.—Patagon.

MACRADENIA, R. Br. in Bot. Reg. t. 612 (1822). *ORCHIDEAE*, Benth. & Hook. f. iii. 586.
RHYNCHADENIA, A. Rich. Fl. Cub. Fanerog. ii. 248. t. 85 (1853).
Brassavolae, *Reichb. f. in Bot. Zeit.* x. (1852) 734.—Guatemala.
delicatula, *Rodrig. Orch. Nov.* ii. 183.—Bras.
loxoglottis, *Focke & Reichb. f. in Walp. Ann.* vi. 697.—Guiana.
lutescens, *R. Br. in Bot. Reg.* t. 612.—Ins. Trinit.
lutescens, Lodd. Bot. Cab. t. 1556 = Epidendrum odoratissimum.
mutica, *Lindl. Bot. Reg.* (1839) *Misc.* 20.—Ins. Trinit.
paraënsis, *Rodrig. Orch. Nov.* i. 139.—Bras.

MACRADENIA:—
polystachya, *Spreng. Syst.* iii. 726.—Ins. Mascar.
Regnellii, *Rodrig. Orch. Nov.* ii. 183.—Bras.
rubescens, *Rodrig. l. c.* i. 139.—Bras.
surinamensis, *Reichb. f. & Wullschl. in Walp. Ann.* vi. 697.—Guian.
triandra, *Lindl. Bot. Reg.* t. 1815.—Guiana; Cuba.

MACRAEA, Hook. f. in Proc. Linn. Soc. i. (1845) 278; et in Trans. Linn. Soc. xx. (1847) 209 = **Lipochaeta**, DC. (Compos.).
lancifolia, Hook. f. in Trans. Linn. Soc. xx. (1847) 210.—Ins. Galap.

MACRAEA, Lindl. in Brand. Journ. (1828) 104 = **Vivianiа**, Cav. (Geran.).
cordata, Meyen, Reise, i. 342 = V. grandifolia.
crenata, Hook. Bot. Misc. i. (1830) 177 = V. crenata.
grandifolia, Lindl. in Brand. Journ. (1828) 105 = V. grandifolia.
parvifolia, Lindl. l. c. = V. parvifolia.
rosea, Lindl. l. c. = V. marifolia.

MACRAEA, Wight, Ic. v. II. 27 (1852) = **Phyllanthus**, Linn. (Euphorb.).
Gardneriana, Wight, l. c. = P. simplex.
myrtifolia, Wight, l. c. = P. myrtifolius.
oblongifolia, Wight, l. c. = P. simplex.
ovalifolia, Wight, l. c. = P. simplex.
Rheedii, Wight, l. c. = P. Macraei.

MACRANDRIA, Meissn. Gen. 160 (1838) = **Hedyotis**, Linn. (Rubiac.).

MACRANTHERA, Torr. ex Benth. in Hook. Comp. Bot. Mag. i. (1835) 203. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 971.
CONRADIA, Nutt. in Journ. Acad. Philad. vii. (1834) 88. t. 12.
TOMILIX, Rafin. New Fl. Am. ii. 72 (1836).
TOXOPUS, Rafin. l. c. 71 (1836).
fuchsioides, *Torr. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 203.—Am. bor.
Lecontii, *Torr. in Ann. Lyc. N. York*, iv. (1837) 80. t. 4.—Am. bor.

MACRANTHUS, Poir. Encyc. Suppl. iii. 569 (1813) = MARCANTHUS, Lour. = **Mucuna**, Adans. (Legumin.).

MACREIGHTIA, A. DC. in DC. Prod. viii. 220 (1844) = **Maba**, Forst. (Ebenac.).
acapulcensis, A. DC. l. c. 221 = Maba acapulcensis.
acuminata, Thw. Enum. Pl. Zeyl. 424 = Maba acuminata.
albens, A. DC. in DC. Prod. viii. 221 = Maba albens.
andamanica, Kurz, in Rep. Veg. Andam. ed. II. i. 42 = Maba andamanica.
buxifolia, Griseb. Cat. Pl. Cub. 169 = Maba Grisebachii.
caribaea, A. DC. in DC. Prod. viii. 221 = Maba caribaea.
conduplicata, A. DC. l. c. = Maba inconstans.
inconstans, A. DC. l. c. = Maba inconstans.
intricata, A. Gray, in Proc. Am. Acad. v. (1861) 163 = Maba intricata.
myristicoidea, Spruce, in Journ. Linn. Soc. v. (1861) 8 = Maba myristicoides.
oblongifolia, Kurz, in Rep. Veg. Andam. ed. I. i. 11 = Maba andamanica.
oblongifolia, Thw. Enum. Pl. Zeyl. 183, 423 = Maba oblongifolia.
obovata, Mart. ex Miq. in Mart. Fl. Bras. vii. 9 = Maba inconstans.
ovalifolia, Thw. Enum. Pl. Zeyl. 424 = Maba ovalifolia.
Pavonii, A. DC. in DC. Prod. viii. 222 = Maba Pavonii.
psidioides, A. DC. l. c. 221 = Maba inconstans.

MACRIA, Spach, Hist. Vég. Phan. ix. 245 (1840) = **Selago**, Linn.

MACRIA, Tenore, in Mem. Soc. Ital. Mod. xxiv. (1848) 366 = **Cordia**, Linn. (Borag.).
brasiliensis, Tenore, ex Pasq. Cat. Hort. Bot. Neap. (1867) 63, in nota = seq.
callipticantha, Tenore, in Mem. Soc. Ital. Mod. xxiv. (1848) 367 = C. rufescens.
calliptychantha, Ces. in Atti Acc. Nap. vi. (1875) n. 11. p. 6 = praec.

MACROBLEPHARUS, Phil. in Linnaea, xxix. (1857–58) 100 = **Eragrostis**, Beauv. (Gramin.).
contractus, Phil. l. c. = E. ciliaris.

MACROCALYX, Miers, ex Lindl. Veg. Kingd. 764 (1847). *RUBIACEAE* (Quid?).

MACROCALYX, Trew, in Act. Nat. Cur. ii. (1761) 332 = **Ellisia**, Linn. (Hydroph.).

MACROCAPNOS, Royle, ex Lindl. Nat. Syst. ed. II. 439 (1836) = **Dicentra**, Bernh. (Papav.).

MACROCARPHUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 376 = **Chaenactis**, DC. (Compos.).
achilleaefolius, Nutt. l. c. = C. Douglasii.
Douglasii, Nutt. l. c. = C. Douglasii.

MACROCENTRUM, Hook. f. in Benth. & Hook. f. Gen. i. 756 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
AULACIDIUM, Rich. ex DC. Prod. iii. 112 (1828).
cristatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 79.—Guian.
droseroides, *Triana, l. c.* 80.—Guian.
fasciculatum, *Hook. f. in Benth. & Hook. f. Gen.* i. 756.—Guian.
parviflora, *Hook. f. l. c.*—Bras.

MACROCENTRUM, Phil. Sert. Mend. Alt. 42 (1871) = **Habenaria**, Willd. (Orchid.).
mendocinum, Phil. l. c. 43.—Chili.

MACROCEPHALUS, Lindl. Veg. Kingd. 712 (1847), sphalm. = MACROCARPHUS, Nutt. = **Chaenactis**, DC. (Compos.).

MACROCERATIDES, Raddi, in Mem. Mod. xviii. Fis. (1820) 392 = **Mucuna**, Adans. (Legumin.).
Pseudostizolobium, Raddi, l. c. = Mucuna Macroceratides.

MACROCERATIUM, Reichb. Consp. 184 (1828) = **Andrzeiowskia**, Reichb. (Crucif.).

MACROCHAETA, Steud. in Zoll. Syst. Verz. Ind. Archip. 60 (1854) = **Pennisetum**, Rich. (Gramin.).
sacchariformis, Steud. l. c. (= *Sericura elegans.*).—Java.

MACROCHAETIUM, Steud. Syn. Pl. Cyp. 159 (1855). *CYPERACEAE*.
Dregei, *Steud. l. c.*—Afr. austr.

MACROCHILUS, Knowles & Westc. Flor. Cab. i. (1837) 93 = **Miltonia**, Lindl. (Orchid.).
Fryanus, Knowles & Westc. l. c. = Miltonia spectabilis.

MACROCHILUS, Presl, Prod. Monog. Lobel. 47 (1836) = **Cyanea**, Gaudich. (Campan.).
superbus, Presl, l. c. = C. superba.

MACROCHITON, M. Roem. Syn. Hesper. 84, 104 (1846) = **Dysoxylon**, Blume (Meliac.).
excelsum, M. Roem. l. c. 104 = D. excelsum.
Forsteri, M. Roem. l. c. = D. Forsteri.
Lessertianum, M. Roem. l. c. = D. Lessertianum.
mollissimum, M. Roem. l. c. = D. mollissimum.

MACROCHLAMYS, Decne. in Rev. Hortic. Sér. III. iii. (1849) 243 = **Alloplectus**, Mart. (Gesner.).
guttatus, Decne. l. c.—Hab.?
involucratus, Decne. l. c. nomen.—Hab.?
Miquelii, Decne. l. c. = A. Patrisii?
Patrisii, Decne. l. c. = A. Patrisii?
speciosus, Decne. l. c. = A. capitatus?

MACROCHLOA, Kunth, Rév. Gram. i. 58 (1815) = **Stipa**, Linn. (Gramin.).
arenaria, Kunth, l. c. 59 = S. arenaria.
gigantea, Hack. Cat. Gram. Port. 16 = S. arenaria.
tenacissima, Kunth, Rév. Gram. i. 59 = S. tenacissima.

MACROCHORDION, De Vriese, in Jaarb. Nederl. Maat. Tuinb. (1853) 14 = **Aechmea**, Ruiz & Pav. (Bromel.).
bromeliaefolium, Beer, Bromel. 146 = Ae. bromeliaefolium.
luteum, Regel & Linden, in Ind. Sem. Hort. Petrop. (1866) 82 = Ae. calyculata.
melananthum, Beer, Bromel. 146 = Ae. bromeliaefolia.
nudiusculum, C. Koch, Wochenschr. vii. (1864) 176 = Ae. calyculata.
Pinelianum, Lem. Illustr. Hortic. ix. (1862) Misc. 62 = Ae. Pineliana.
pulchrum, Beer, Bromel. 147 = Ae. bromeliaefolia.
recurvatum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiv. (1856) 393.—Hab.?
strictum, Beer, Bromel. 147 = Ae. bromeliaefolia.
tinctorium, De Vriese, in Jaarb. Nederl. Maat. Tuinb. (1853) 14 = Ae. bromeliaefolia.

MACROCHORDIUM, Beer, Bromel. 145 (1857) = praec.

MACROCLADUS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 489 = **Orania**, Zipp. (Palmae).
sylvicola, Griff. l. c. 490 = O. Macrocladus.

MACROCLINIDIUM, Maxim. in Bull. Acad. Pétersb. xv. (1871) 375. *COMPOSITAE*, Benth. & Hook. f. ii. 493.
robustum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 376.—Japon.
verticillatum, *Franch. & Sav. Enum. Pl. Jap.* i. 265 in nota.—Japon.

MACROCNEMUM, P. Br. Hist. Jamaic. 165 (1756). *RUBIACEAE*, Benth. & Hook. f. ii. 35.
LASIONEMA, D. Don, in Brewst. Phil. Mag. ii. (1833) 377; et in Trans. Linn. Soc. xvii. (1834) 141.
candidissimum, Vahl, Symb. Bot. ii. 38. t. 30 = Calycophyllum candidissimum.
cinchonoides, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 76.—N. Granat.
coccineum, Vahl, Symb. Bot. ii. 38. t. 29 = Warscewiczia coccinea.
corymbosum, Ruiz & Pav. Fl. Per. ii. 48. t. 189 = Condaminea corymbosa.
cubense, *Griseb. Cat. Pl. Cub.* 124.—Cuba.
exsertum, *Oerst. in Kjoeb. Vedensk. Meddel.* (1852) 45.—Costa Rica.
glabratum, Bartl. ex DC. Prod. iv. 402 = Condaminea glabrata.
glabrescens, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 76.—Costa Rica.
grandiflorum, *Wedd. l. c.*—N. Granat.
Humboldtianum, *Wedd. l. c.*—Peruv.
jamaicense, *Linn. Pl. Jam. Pugill.* 31; *Amoen. Acad.* v. 413.—Jamaic.
longifolium, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 279 = Chimarrhis cymosa.
microcarpum, Ruiz & Pav. Fl. Per. ii. 49 = Condaminea microcarpa.
parviflorum, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 525 (Quid?).—Ins. Molucc.
pastoënse, *Karst. in Linnaea*, xxx. (1859–60) 150.—N. Granat.
pubescens, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 76.—N. Granat.
roseum, *Wedd. l. c.*—Peruv.
speciosum, Jacq. Hort. Schoenb. i. 19 = Mussaenda speciosa.
stipulaceum, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 525.—Ins. Molucc.
strictum, Roxb. ex Roem. & Schult. Syst. v. 6 = Hyptianthera stricta.
stylocarpum, *Karst. in Linnaea*, xxx. (1859–60) 149.—N. Granat.
tetrandrum, Cav. ex Roem. & Schult. Syst. v. 254 = Mussaenda tetrandra.
tetrandrum, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 279.—Bras.
tinctorium, H. B. & K. Nov. Gen. et Sp. iii. 399 = Condaminea tinctoria.
tubulosum, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 279 (Pogonopi sp.).—Bras.
venosum, Ruiz & Pav. Fl. Per. ii. 49 = Condaminea venosa.

MACROCNEUMUM, Vand. Fl. Lusit. et Bras. 14 (1788) = **Remijia**, DC. (Rubiac.).
ferrugineum, Vell. ex Steud. Nom. ed. II. ii. 88 = R. ferruginea.

MACROCOCCULUS, Becc. Malesia, i. (1877) 160. *MENISPERMACEAE.*
pomiferus, *Becc. l. c.* 161.—N. Guin.

MACROCROTON, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 1186 (1848) = **Croton**, Linn. (Euphorb.).
cuneatus, Klotzsch, l. c. = C. cuneatus.
surinamensis, Klotzsch, l. c. = C. surinamensis?

MACROCYMBIUM, Walp. in Flora, xxxvi. (1853) 149 = **Erythrina**, Linn. (Legumin.).
Vogelii, Walp. l. c. = E. senegalensis.

MACRODISCUS, Bur. Monog. Bign. 46. t. 11 (1864) = **Distictis**, Mart. (Bignon.).
rigescens, Bur. l. c. = D. rigescens.

MACROGYNE, Link & Otto, Ic. Pl. Sel. 69. t. 31 (1823) = **Aspidistra**, Ker-Gawl. (Liliac.).
convallariaefolia, Link & Otto, l. c. = A. lurida.

MACROLENES, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 311 = **Marumia**, Blume (Melastom.).
annulata, Naud. l. c. = Marumia annulata.

MACROLEPIS, A. Rich. Sert. Astrol. 25. t. 19 (1833) = **Bulbophyllum**, Thou. (Orchid.).
longiscapa, A. Rich. l. c.—Vanikora.

MACROLINIUM, Rodrig. Orch. Nov. ii. 236 (1881), *ORCHIDEAE.*
roseum, *Rodrig. l. c.* 237.—Bras.

MACROLINUM, Klotzsch, in Linnaea, xii. (1838) 242 = **Simocheilus**, Klotzsch (Ericac.).
ciliatum, Klotzsch, l. c. 243 (= *Syndesmanthus ciliatus*).—Afr. austr.
paucifolium, Klotzsch, l. c. 242 (= *Syndesmanthus paucifolius*).—Afr. austr.

MACROLINUM, Reichb. Handb. 306 (1827) = **Reinwardtia**, Dum. (Lineae).
trigynum, Reichb. l. c. = R. trigyna.

MACROLOBIUM, Schreb. Gen. 30 (1789). *LEGUMINOSAE*, Benth. & Hook. f. i. 579.
ANTHONOTA, Beauv. Fl. Ow. et Ben. i. 70. t. 42 (1805).
KRUEGERIA, Scop. Introd. 314 (1777).
OUTEA, Aubl. Pl. Gui. 28. t. 9 (1775).
VOUAPA, Aubl. l. c. 25. t. 7 (1775).
acaciaefolium, *Benth. in Mart. Fl. Bras.* xv. II. 224.—Am. trop.
bifolium, Pers. Syn. i. 39 = M. Vuapa.
bijugum, Colebr. in Trans. Linn. Soc. xii. (1819) 359. t. 17 = Afzelia bijuga.
canaliculatum, *Spruce, ex Mart. Fl. Bras.* xv. II. 219.—Bras.
chrysostachyum, *Benth. l. c.* 220.—Bras.; Guian.
demonstrans, *Oliver, Fl. Trop. Afr.* ii. 299.—Afr. trop.
discolor, *Benth. in Mart. Fl. Bras.* xv. II. 222.—Bras.
elegans, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 40 = M. Vuapa.
flexuosum, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 223.—Bras.
floridum, *Karst. Fl. Columb.* i. 151. t. 75.—Venezuela.
gracile, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 223.—Bras.
Heudelotii, *Planch. ex Benth. in Trans. Linn. Soc.* xxv. (1865) 308.—Afr. trop.
hymenaeoides, Willd. Sp. Pl. i. 186 = M Vuapa.
latifolium, *Vog. in Linnaea*, xi. (1837) 414.—Bras.
limbatum, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 218.—Bras.
multijugum, *Benth. l. c.* 222.—Am. trop.
Outea, Steud. Nom. ed. I. 503 = M. Utea.
Palisoti, *Benth. in Trans. Linn. Soc.* xxv. (1865) 308.—Afr. trop.
pendulum, *Willd. ex Vog. in Linnaea*, xi. (1837) 412.—Bras.
phaseolocarpum, Steud. Nom. ed. II. ii. 88 = Cynometra racemosa.
pinnatum, Willd. Sp. Pl. i. 186 = M. Utea.
punctatum, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 219.—Bras.
racemigerum, Tul. in Arch. Mus. Par. iv. (1844) 174 = pendulum.
Simira, *J. F. Gmel. Syst.* 93.—Guian.

MACROLOBIUM:—
sphaerocarpum, Willd. Sp. Pl. i. 186 = M. Simira.
stamineum, G. F. W. Mey. Prim. Fl. Esseq. 18 = M. Vuapa.
stipulaceum, *Benth. in Trans. Linn. Soc.* xxv. (1865) 308.—Afr. trop.
suaveolens, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 219.—Bras.
taxifolium, *Spruce, ex Benth. l. c.* 224.—Bras.
Utea, *J. F. Gmel. Syst.* 93.—Guian.
venulosum, *Benth. in Mart. Fl. Bras.* xv. II. 223.—Bras.
Vuapa, *J. F. Gmel. Syst.* 93.—Am. trop.

MACROLOMIA, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 181. t. 24 (1842) = **Scleria**, Berg. (Cyperac.).
bracteata, Schrad. ex Nees, l. c. 182 = S. bracteata.

MACROMERIA, D. Don, in Edinb. N. Phil. Journ. (Jul.-Oct. 1832) 239. *BORAGINEAE*, Benth. & Hook. f. ii. 859.
PHILONOMIA, DC. ex Steud. Nom. ed. II. ii. 88, 320 (1841).
cinerascens, *A. DC. in DC. Prod.* x. 69.—Peruv.
discolor, *Benth. Pl. Hartw.* 49.—Mexic.
exserta, *D. Don, in Edinb. N. Phil. Journ.* (Jul.-Oct. 1832) 239.—Mexic.
hispida, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 339 = longiflora.
longiflora, *D. Don, in Edinb. N. Phil. Journ.* (Jul.-Oct. 1832) 239.—Mexic.
viridiflora, *DC. Prod.* x. 68.—Mexic.

MACROMERUM, Burch. Trav. i. 388 (1824) = **Cadaba**, Forsk. (Cappar.).
junceum, Burch. l. c. = C. juncea.

MACROMISCUS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 507 = **Aeschynomene**, Linn. (Legumin.).
brasiliensis, Turcz. l. c. 508 = A. montevidensis.
glandulosus, Turcz. l. c. xx. (1847) I. 174 = A. rostrata.

MACROMYRTUS, Miq. Fl. Ind. Bat. i. I. 439 (1855) = **Eugenia**, Linn. (Myrtac.).
javanica, Miq. l. c. 440.—Java.

MACRONAX, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Arundinaria**, Michx. (Gramin.).

MACRONEMA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 322 = **Chrysopsis**, Ell. (Compos.).
discoidea, Nutt. l. c. = Aplopappus Macronema.
suffruticosa, Nutt. l. c. = Aplopappus suffruticosus.

MACRONYX, Dalz. in Hook. Kew Journ. ii. (1858) 35 = **Tephrosia**, Pers. (Legumin.).
strigosus, Dalz. l. c. = T. tenuis.
tenuis, Hook. l. c. in nota = T. tenuis.

MACROPANAX, Miq. Fl. Ind. Bat. i. I. 763 (1855). *ARALIACEAE*, Benth. & Hook. f. i. 945.
concinnum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 220.—Java.
cyrtostylum, *Miq. l. c.* 13.—Sumatra.
floribundum, Miq. Fl. Ind. Bat. i. I. 764 = oreophilum.
glomerulatum, Miq. l. c. = Brassaiopsis speciosa.
oreophilum, *Miq. l. c.*—Reg. Himal.; Malaya.
undulatum, *Seem. in Journ. Bot.* ii. (1864) 294.—Reg. Himal.

MACROPETALUM, Burch. ex Decne. in DC. Prod. viii. 626 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 778.
Burchellii, *Decne. in DC. Prod.* viii. 627.—Afr. austr.

MACROPHORA, Rafin. Fl. Tellur. iv. 103 (1836) = **Passiflora**, Linn.
sanguinea, Rafin. l. c. = P. vitifolia.

MACROPHTHALMA, Gasp. Ricerche Caprif. 83. t. 8 (1845) = **Ficus**, Tourn. (Urtic.).
elastica, Gasp. l. c. = F. elastica.

MACROPIDIA, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 57. *HAEMODORACEAE*, Benth. & Hook. f. iii. 677.
fumosa, *J. Drumm. ex Harv. in Hook. Kew Journ.* vii. (1855) 57.—Austral.

MACROPIPER, Miq. Enum. Phyt. 35 (1839) = **Piper**, Linn.
excelsum, Miq. Syst. Pip. 221 = P. excelsum.
latifolium, Miq. l. c. 218 = P. latifolium, guahamense.
latifolium, Miq. in Linnaea, xx. (1847) 130 = P. methysticum.
latum, Presl, Epim. Bot. 223 = P. latum.
methysticum, Hook. & Arn. in Bot. Beech. Voy. 96 = P. methysticum.
methysticum, Miq. Syst. Pip. 217 = P. guahamense.
potamogetonifolium, Miq. l. c. 221 = P. potamogetonifolium.
psittacorum, Miq. l. c. = P. excelsum.
puberulum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 235 = P. Macgillivrayi.
Reinwardtianum, Miq. in Linnaea, xxi. (1848) 481 = P. Reinwardtianum.

MACROPLACIS, Blume, Mus. Bot. Lugd. Bat. i. 7 (1849) = **Kibessia**, DC. (Melastom.).
cordata, Blume, l. c. = K. cordata.

MACROPLATIS, Triana, in Trans. Linn. Soc. xxviii. (1871) 153, sphalm. = praec.

MACROPODIA, Benth. Fl. Austral. vi. 447 (1873), sphalm. = **Macropidia**, J. Drumm. (Haemodor.).

MACROPODIUM, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 108 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 69.
elongatum, Schur, ex Nym. Consp. 46 = Brassica elongata.
laciniatum, Hook. Fl. Bor. Am. i. 43 = Thelypodium laciniatum.
nivale, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 108.—Sibir. altaic.
pterospermum, *F. Schmidt, Reis. Amurl.* 113.—Ins. Sachalin.

MACROPSIDIUM, Blume, Mus. Bot. Lugd. Bat. i. 85 (1849) = **Myrtus**, Tourn.
elegans, Blume, l. c.—Ins. Molucc.
rubrum, Blume, l. c.—Cochinch.

MACROPTERANTHES, F. Muell. Fragm. iii. 91 (1862). *COMBRETACEAE*, Benth. & Hook. f. i. 687.
Fitzalani, *F. Muell. Fragm.* viii. 160.—Austral.
Kekwickii, *F. Muell. l. c.* iii. 151.—Austral.
Leichhardtii, *F. Muell. l. c.* 91.—Austral.
montana, *F. Muell. l. c.* 91.—Austral.

MACRORHAMNUS, Baill. Adansonia, xi. (1875) 273. *RHAMNEAE.*
decipiens, *Baill. l. c.*—Madag.

MACRORHYNCHUS, Less. Syn. Comp. 139 (1832) = **Troximon**, Nutt. (Compos.).
angustifolius, Kellogg, in Proc. Calif. Acad. v. (1873) 47 = T. retrorsum.
aurantiacus, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 40 = T. aurantiacum.
australis, Phil. in Anal. Univ. Chil. (1873) 482.—Chili.
californicus, Torr. & Gray, Fl. N. Am. ii. 492 = T. heterophyllum.
chilensis, Hook. Lond. Journ. Bot. vi. (1847) 256 = T. heterophyllum.
chilensis, Less. Syn. Comp. 139.—Chili.
coronopifolius, Hook. f. Fl. Antarct. t. 112 = *pumilus.*
cynthioides, Hook. Lond. Journ. Bot. vi. (1847) 256 = T. glaucum.
elatus, Torr. & Gray, Fl. N. Am. ii. 492 = T. Nuttallii.
glaucus, Eaton, in Bot. King's Exp. 204 = T. glaucum.
grandiflorus, Eaton, l. c. = T. Nuttallii.
grandiflorus, Torr. & Gray, Fl. N. Am. ii. 492 = T. grandiflorum.
Harfordii, Kellogg, in Proc. Calif. Acad. v. (1873) 47 = T. humile.
heterophyllus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 430 = T. heterophyllum.
humilis, Benth. Pl. Hartw. 320 = T. humile.
laciniatus, Torr. & Gray, Fl. N. Am. ii. 492 = T. laciniatum.
laevigatus, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 41 = *chilensis.*
Lessingii, Hook. & Arn. Bot. Beech. Voy. 145 = Troximon humile.
Poeppigii, DC. Prod. vii. 152.—Chili.

MACRORHYNCHUS:—
pterocarpus, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 41.—Chili.
pumilus, DC. Prod. vii. 152.—Ins. Falkland.
purpureus, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 114 = T. aurantiacum.
retrorsus, Benth. Pl. Hartw. 320 = T. retrorsum.
troximoides, Torr. & Gray, Fl. N. Am. ii. 491 = T. aurantiacum.

MACROSCEPIS, H. B. & K. Nov. Gen. et Sp. iii. 200. t. 233 (1818). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 751.
aurea, *Fourn. in Mart. Fl. Bras.* vi. IV. 301.—Bras.
guianensis, *Klotzsch, in Schomb. Fauna et Fl. Brit. Gui.* 1147, nomen.—Guiana.
longiflora, Spreng. Syst. i. 851 (= *Schubertia longiflora*).—Am. austr.
obovata, *H. B. & K. Nov. Gen. et Sp.* iii. 201.—Mexic.
patens, Spreng. Syst. i. 851 = obovata.
rotata, *Decne. in DC. Prod.* viii. 599.—Mexic.
Selloana, *Fourn. in Mart. Fl. Bras.* vi. IV. 302.—Bras.
Trianae, *Decne. ex Triana, in Compt. Rend. Par.* lxxiv. (1872) 882.—N. Granat.
tristis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 321.—Panama.
urceolata, *Karst. Fl. Columb.* ii. 115. t. 161.—N. Granat.

MACROSELINUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 30 = **Peucedanum**, Tourn. (Umbell.).
latifolium, Schur, l. c. = P. latifolium.

MACROSEMA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = **Astragalus**, Tourn. (Legumin.).
aduncum, Stev. l. c. = A. aduncus.
bicolor, Stev. l. c. = A. bicolor.
flaccidus, Stev. l. c. xxix. (1856) II. 146 = A. flaccidus.
onobrychioides, Stev. l. c. iv. (1832) 266 = A. Cephalotes.
Onobrychis, Stev. l. c. = A. Onobrychis.
succulentum, Stev. l. c. = A. caryocarpus.

MACROSEPALUM, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 25 (1882). *CRASSULACEAE*.
turkestanicum, *Regel & Schmalh. l. c.*—Turkestan.

MACROSIPHON, Hochst. in Flora, xxiv. (1841) 373 = **Rhamphicarpa**, Benth. (Scrophul.).
elongatus, Hochst. l. c. 374 = R. longiflora.
fistulosus, Hochst. l. c. = R. longiflora.

MACROSIPHON, Miq. in Linnaea, xix. (1846) 442 = **Hindsia**, Benth. (Rubiac.).
brasiliensis, Miq. l. c. 443 = H. longiflora.

MACROSIPHONIA, Muell. Arg. in Mart. Fl. Bras. vi. I. 137. t. 42, 43 (1860). *APOCYNACEAE*, Benth. & Hook. f. ii. 726.
Berlandieri, *A. Gray, Syn. Fl. N. Am.* ii. I. 83.—Mexic.
brachysiphon, *A. Gray, l. c.*—Mexic.
guaranitica, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 139.—Bras.
hypoleuca, *Muell. Arg. in Linnaea*, xxx. (1859–60) 452.—Mexic.
lanuginosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 316.—Mexic.
longiflora, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 140.—Bras.
Martii, *Muell. Arg. l. c.* 138.—Bras.
pinifolia, Miers, Apocyn. Am. 131 = verticillata.
prostrata, Miers, l. c. = longiflora.
Velame, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 138.—Bras.
verticillata, *Muell. Arg. l. c.* 140.—Bras.; Reg. Argent.
virescens, *Muell. Arg. l. c.* 139.—Bras.
Wrightii, *A. Gray, Syn. Fl. N. Am.* ii. I. 83.—Am. bor. occ.

MACROSOLEN, Blume, in Schult. f. Syst. vii. 1731 (1830) = **Loranthus**, Linn.
ampullaceus, Blume, l. c. = L. ampullaceus.
clavatus, Blume, l. c. = L. clavatus.
cochinchinensis, Blume, l. c. = L. cochinchinensis.
creatophyllus, Miq. Fl. Ind. Bat. i. I. 832 = L. creatophyllus.
elasticus, Blume, in Schult. f. Syst. vii. 1731 = L. elasticus.
evenius, Miq. Fl. Ind. Bat. i. I. 827 = L. avenis.
formosus, Blume, in Schult. f. Syst. vii. 1731 = L. formosus.
Jackianus, Miq. Fl. Ind. Bat. i. I. 828 = L. Jackianus.

MACROSOLEN:—
macrophyllus, Miq. Fl. Ind. Bat. i. I. 829 = Loranthus macrophyllus.
melintangensis, Miq. l. c. 830 = L. melintangensis.
oleoides, Miq. l. c. 831 = L. globosus.
pallens, Miq. l. c. = L. ampullaceus.
patulus, Blume, in Schult. f. Syst. vii. 1731 = L. patulus.
Pseudo-globosus, Miq. Fl. Ind. Bat. i. I. 829 = L. Pseudo-globosus.
Pseudo-perfoliatus, Miq. l. c. 828 = L. Pseudo-perfoliatus.
Reinwardtianus, Blume, in Schult. f. Syst. vii. 1731 = L. Reinwardtianus.
retusus, Blume, l. c. = L. subumbellatus.
retusus, Miq. Fl. Ind. Bat. i. I. 828 = L. retusus.
rotundatus, Miq. l. c. Suppl. 346.—Ins. Banca.
sphaerocarpus, Blume, in Schult. f. Syst. vii. 1731 = L. globosus.
subumbellatus, Blume, l. c. = L. subumbellatus.
tetragonus, Blume, l. c. = L. tetragonus.
tomentosus, Blume, l. c. = L. tomentosus.
viridiflorus, Blume, l. c. = L. globosus.

MACROSPERMUM, Steud. Nom. ed. II. ii. 88 (1841) sphalm. = MACROSPORUM, DC. = **Sobolewskia**, Bieb. (Crucif.).

MACROSPHYRA, Hook. f. in Benth. & Hook. f. ii. 86 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
longistyla, *Hook. f. l. c.*—Afr. trop.

MACROSPORUM, DC. Syst. ii. 575, in obs. (1821) = **Sobolewskia**, Bieb. (Crucif.).

MACROSTACHYA, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 408, in nota (1851) = **Chloris**, Sw. (Gramin.).

MACROSTEGIA, Nees, in DC. Prod. xi. 218 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
Ruiziana, *Nees, l. c.*—Peruv.

MACROSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 177 = **Pimelea**, Banks (Thymel.).
erubescens, Turcz. l. c. 178 = P. physodes.

MACROSTEMA, Pers. Syn. i. 185 (1805) = **Ipomoea**, Linn. (Convolv.).
vitifolia, Pers. Syn. l. c. = I. peduncularis.

MACROSTIGMA, Hook. Ic. Pl. t. 412 (1842) = **Stylobasium**, Desf. (Rosac.).
australis, Hook. l. c. = S. lineare.

MACROSTIGMA, Kunth, Ind. Sem. Hort. Berol. (1848) 12; in Ann. Sc. Nat. Sér. III. xi. (1849) 220; Enum. Pl. v. 319 (1850) = **Tupistra**, Ker-Gawl. (Liliac.).
tupistroides, Kunth, ll. cc. 12; 221; 319 = T. squalida.

MACROSTOMIUM, Blume, Bijdr. 335. t. 37 (1825) = **Dendrobium**, Sw. (Orchid.).
aloëfolium, Blume, l. c. = D. Serra.

MACROSTOMUM, Benth. & Hook. f. Gen. iii. 499 (1880), sphalm. = praec.

MACROSTYLIS, Bartl. & Wendl. f. Beitr. 191. t. B (1824). *RUTACEAE*, Benth. & Hook. f. i. 289.
barbata, St. Hil. ex Steud. Nom. ed. II. ii. 89 = lanceolata.
barbigera, *Bartl. & Wendl. f. Beitr.* i. 195.—Afr. austr.
cordata, G. Don, Gen. Syst. i. 789 = Agathosma imbricata.
crassifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 440.—Afr. austr.
decipiens, *E. Mey. in Drège, Zwei Pfl. Docum.* 200, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 439.—Afr. austr.
dubia, *Benth. & Hook. f. Gen.* i. 289 (in nota sub Euchaetide).—Afr. austr.
hirta, *E. Mey. ex Drège, Zwei Pfl. Docum.* 200, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 441.—Afr. austr.
lanceolata, *Bartl. & Wendl. f. Beitr.* i. 194.—Afr. austr.
obtusa, Bartl. & Wendl. f. l. c. 197 = squarrosa.
ovata, *Sond. in Harv. & Sond. Fl. Cap.* i. 441.—Afr. austr.
Sieberi, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 439.—Afr. austr.
squarrosa, *Bartl. & Wendl. f. Beitr.* i. 198.—Afr. austr.

MACROSTYLIS:—
tenuis, *E. Mey. in Drège, Zwei Pfl. Docum.* 200, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 440.—Afr. austr.
villosa, Sond. in Harv. & Sond. l. c. 439 = lanceolata.

MACROSTYLIS, Breda, Orch. Kuhl. et Hass. [t. 2] (1827) = **Corymbis**, Thou. (Orchid.).
decumbens, Reichb. f. in Bonplandia, ii. (1854) 11 (= *Chloidia decumbens*).—Bras.
disticha, Breda, Orch. Kuhl. et Hass. [t. 2] = C. corymbosa.
forcipigera, Reichb. f. in Bonplandia, ii. (1854) 96.—Am. trop.
galipanensis, Reichb. f. l. c. 11 = C. galipanensis.
vernalis, Reichb. f. l. c. (= *Chloidia vernalis*).—Jamaic.

MACROSYPHONIA, Duby, in Mém. Soc. Phys. Genèv. (1844) 426 = **Dionysia**, Fenzl (Primul.).
cespitosa, Duby, l. c. 427 = D. caespitosa.

MACROTHYRSUS, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 61 = **Aesculus**, Linn. (Sapind.).
discolor, Spach, l. c. 62 = A. parviflora.
odorata, Rafin. Alsog. Am. 75 = A. parviflora.

MACROTOMIA, DC. in Meissn. Gen. 281; Comm. 190 (1840); Prod. x. 26 (1846). *BORAGINEAE*, Benth. & Hook. f. ii. 862.
AIPYANTHUS, Stev. in Bull. Soc. Nat. Mosc. (1851) I. 599.
LEPTANTHE, Klotzsch, in Bot. Ergeb. Waldem. Reise, 95. t. 63 (1862).
MUNBYA, Boiss. Diagn. Ser. I. xi. 114 (1849).
Benthami, *DC. ex Meissn. Gen. Comm.* 190; *Prod.* x. 27 —Reg. Himal.
Cephalotes, *Boiss. Fl. Orient.* iv. 212.—Graecia; As. Min.
cyanochroa, Boiss. l. c. = perennis.
echioides, *Boiss. l. c.* 211.—Oriens.
endochroma, Hook. f. & Thoms. ex Henders. & Hume, Yarkand, 328 = perennis.
onosmoides, Regel & Smirnow, in Act. Hort. Petrop. v. (1877) 624 = perennis.
perennis, *Boiss. Fl. Orient.* iv. 212, in nota.—Reg. Himal.; Turkestan.

MACROTONICA, Steud. Nom. ed. II. ii. 89 (1841), sphalm. = praec.

MACROTROPIS, DC. Prod. ii. 98 (1825) = **Ormosia**, Jacks. (Legumin.).
bancana, Miq. Fl. Ind. Bat. Suppl. 295 = O. parvifolia.
emarginata, Walp. Rep. ii. 903 = O. emarginata.
foetida, DC. Prod. ii. 98 = Anagyris sinensis.
inodora, DC. l. c. 99 = Anagyris inodora.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 294.—Sumatra.

MACROTRULLION, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 1202 (1848) sine desc. *LEGUMINOSAE* (Quid?).
elegans, Klotzsch, l. c.—Guian.
splendens, Klotzsch, l. c.—Guian.

MACROTRYS, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Cimicifuga**, Linn. (Ranunc.).
actaeoides, Rafin. l. c. = C. racemosa.
racemosa, Sweet, Hort. Brit. ed. I. 9 = C. racemosa.

MACROTYBUS, Dulac, Fl. Hautes-Pyr. 425 (1867) = **Douglasia**, Lindl. (Primul.).
luteus, Dulac, l. c. = D. Vitaliana.

MACROTYS, 'Rafin.' ex DC. Syst. i. 383 (1818) = MACROTRYS, Rafin. = **Cimicifuga**, Linn. (Ranunc.).

MACROZAMIA, Miq. Monog. Cycad. 36. t. 4, 5 (1841). *CYCADACEAE*, Benth. & Hook. f. iii. 445.
CATAKIDOZAMIA, T. Hill, in Gard. Chron. (1865) 1107.
LEPIDOZAMIA, Regel, Gartenfl. vi. (1857) 11.
corallipes, Hook. f. Bot. Mag. t. 5943 = spiralis.
cylindrica, André, Illustr. Hortic. xxiii. (1881) 32 = spiralis.
cylindrica, *C. Moore, in Journ. R. Soc. N. S. W.* xvii. (1884) 119.—Austral.
Denisonii, F. Muell. Fragm. i. 41 = Peroffskyana.

MACROZAMIA:—
Douglasii, *T. Hill, ex F. Muell. Census*, 110, nomen.—Austral.
eriolepis, Brongn. ex Miq. in Verh. Kon. Akad. Wet nsch. II. iii. (1869) 56 = Peroffskyana.
Fawcetti, *C. Moore, in Journ. R. Soc. N. S. W.* xvii. (1884) 120.—Austral.
flexuosa, *C. Moore, l. c.* 121.—Austral.
Fraseri, *Miq. Monog. Cycad.* 37.—Austral.
Gigas, Hort. ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 56 = Peroffskyana.
heteromera, *C. Moore, in Journ. R. Soc. N. S. W.* xvii. (1884) 122.—Austral.
Hopei, *T. Hill, ex C. Moore, l. c.* 116.—Austral.
littoralis, Liebm. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 191 = Dioon edule.
Macdonnelli, F. Muell. Fragm. ii. 179 = Fraseri.
Macleayi, Miq. Arch. Néerl. iii. (1868) 250 = spiralis.
Miquelii, *A. DC. in DC. Prod.* xvi. II. 535.—Austral.
Oldfieldi, Miq. in Versl. Akad. Wet. xv. (1863) 370 = Fraseri.
Pauli-Guilielmi, *Hill & Muell. ex F. Muell. Fragm.* i. 86.—Austral.
pectinata, Liebm. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 191 = Dioon edule.
Peroffskyana, *Miq. in Arch. Néerl.* iii. (1868) 252.—Austral.
Perowskiana, Auct. = praec.
plumosa, A. Mohr, in Rev. Hort. Belg. iii. (1877) 69. f. 7 = M. Pauli-Guilielmi.
Preissii, Lehm. Cat. Hort. Hamb. 1842; ex ej. Pugill. viii. 31 = Fraseri.
secunda, *C. Moore, in Journ. R. Soc. N. S. W.* xvii. (1884) 120.—Austral.
spiralis, *Miq. Monog. Cycad.* 36. t. 4, 5.—Austral.
spiralis, Miq. l. c. 36, partim = Fraseri.
tenuifolia, Hort. ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 53 = M. Pauli-Guilielmi.
tridentata, Regel, Gartenfl. (1876) 229 = Miquelii.

MACUBEA, Jaume St. Hil. Expos. ii. 29 (1805) = **Macoubea**, Aubl. (Inc. sed.).

MACUCUA, J. F. Gmel. Syst. 210, 262 (1791) = MACOUCOUA, Aubl. = **Ilex**, Tourn.

MACUILLAMIA, Rafin. Neogenyt. 2 (1825) = **Herpestis**, Gaertn. f. (Scroph.).

MACUNA, Marcgr. ex Scop. Introd. 309 (1777) = **Mucuna**, Adans. (Legumin.).

MADACARPUS, Wight, Ic. t. 1152 (1846) = **Senecio**, Tourn. (Compos.).
belgaumensis, Wight, l. c. = S. belgaumensis.

MADARACTIS, DC. Prod. vi. 322 (1837) = **Senecio**, Tourn. (Compos.).
arachnoidea, Herb. Calc. ex C. B. Clarke, Comp. Ind. 199 = S. Grahami.
glabra, DC. Prod. vi. 440 = S. saxatilis.
incana, Regel, Ind. Sem. Hort. Turic. (1847) 4; ex Walp. Ann. i. 427 = S. incanus?
lanuginosa, Edgew. in Trans. Linn. Soc. xx. (1846) 75 = S. Edgeworthii.
Metziana, Sch. Bip. ex C. B. Clarke, Comp. Ind. 344 = S. saxatilis.
pinnatifida, DC. Prod. vi. 439 = S. Hohenackeri.
polycephala, DC. l. c. 440 = S. polycephalus.
scabra, DC. l. c. = S. Lessingianus.
variegata, Bartl. Ind. Sem. Hort. Gotting. (1852) 3; ex Urban, Add. Ind. Sem. Hort. Berol. (1880) 35.—Cf. *Madaria variegata*.

MADARAGLOSSA, Hook. Ic. Pl. iv. t. 326 (1841) err. typ. = MADAROGLOSSA, DC. = **Layia**, Hook. & Arn. (Compos.).

MADARIA, DC. in Mém. Soc. Phys. Genèv. vii. II. (1836) 280 = **Madia**, Molina (Compos.).
corymbosa, DC. Prod. v. 692 = Madia elegans.
elegans, DC. l. c. = Madia elegans.
racemosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 386 = Madia sativa.
variegata, Bartl. Ind. Sem. Hort. Gotting. (1852) 3.—Cf. Linnaea, xxvi. (1853) 736.—Bras.

MADARIOPSIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 387 = **Madia**, Molina (Compos.).
chilensis, Nutt. l. c. 388 = Madia sativa.

MADAROGLOSSA, DC. Prod. v. 694 (1836) = **Layia**, Hook. & Arn. (Compos.).
angustifolia, DC. Prod. v. 694 = L. glandulosa.
angustifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 394 = L. platyglossa.
carnosa, Nutt. l. c. 393 = L. carnosa.
Douglasii, Walp. Rep. ii. 632 = L. Calliglossa.
elegans, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 393 = L. elegans.
heterotricha, DC. Prod. v. 694 = L. heterotricha.
hieracioides, DC. l. c. = L. hieracioides.
hirsuta, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 394 = L. platyglossa.

MADAROSPERMA, Benth. in Benth. & Hook. f. Gen. ii. 1241 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
aripecurense, *Fourn. in Mart. Fl. Bras.* vi. IV. 213.—Bras.
confusum, *Fourn. l. c.* 214.—Bras.
Traillianum, *Benth. in Benth. & Hook. f. Gen.* ii. 1241.—Bras.

MADDENIA, Hook. f. & Thoms. in Hook. Kew Journ. vi. (1854) 381. t. 12. *ROSACEAE*, Benth. & Hook. f. i. 610.
himalaica, *Hook. f. & Thoms. in Hook. Kew Journ.* vi. (1854) 381.—Reg. Himal.
pedicellata, *Hook. f. Fl. Brit. Ind.* ii. 318.—Reg. Himal.

MADEA, Soland. ex DC. Prod. v. 301 (1836) = **Boltonia**, L'Hérit. (Compos.).
glauca, Soland. ex DC. l. c. = B. asteroides.

MADHUCA, J. F. Gmel. Syst. 799 (1791) = **Bassia**, Koen. (Sapot.).
indica, J. F. Gmel. l. c. = B. latifolia.

MADIA, Molina, Sagg. Chile, ed. I. 136 (1782). *COMPOSITAE*, Benth. & Hook. f. ii. 393.
AMIDA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 390.
ANISOCARPUS, Nutt. l. c. (1841) 388.
BIOTIA, Cass. in Dict. Sc. Nat. xxxiv. 308 (1825).
HARPAECARPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 389.
MADARIA, DC. in Mém. Soc. Phys. Genèv. vii. II. (1836) 280.
MADARIOPSIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 387.
MADORELLA, Nutt. l. c. (1841).
PYROPSIS, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 22 (1840).
anomala, Greene, in Bull. Calif. Acad. i. III. (1885) 91 (Hemizoniae sp.).—Calif.
Bolanderi, *A. Gray, in Proc. Am. Acad.* viii. (1872) 391.—Calif.
capitata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 386 = sativa.
citriodora, Greene, in Bull. Torrey Bot. Club. ix. (1882) 63 = Hemizonia citriodora.
corymbosa, Lindl. ex W. Baxt. in Lond. Hort. Brit. Suppl. iii. 584 = elegans.
dissitiflora, *Torr. & Gray, Fl. N. Am.* ii. 405.—Am. bor. occ.
elegans, *D. Don, in Lindl. Bot. Reg.* t. 1458.—Am. bor. occ.
filipes, *A. Gray, in Proc. Am. Acad.* viii. (1873) 391.—Am. bor. occ.
glomerata, *Hook. Fl. Bor. Am.* ii. 24.—Am. bor. occ.
hirsuta, Meyen, Reise, i. 311 = sativa.
mellita, J. F. Gmel. Syst. 1249 = sativa.
mellosa, Molina, Sagg. Chile, ed. I. 137 = sativa.
Nuttallii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 391.—Am. bor. occ.
racemosa, Torr. & Gray, Fl. N. Am. ii. 405 = sativa.
radiata, *Kellogg, in Proc. Calif. Acad.* iv. (1870) 190.—Calif.
Rammii, Greene, in Bull. Calif. Acad. i. III. (1885) 90 = yosemitana.
sativa, *Molina, Sagg. Chile*, ed. I. 136.—Am. occ. bor. et austr.
stellata, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 32 = sativa.
viscosa, Cav. Ic. iii. 50 = sativa.
yosemitana, *Parry, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 219.—Calif.

MADIOLA, A. St. Hil. Fl. Bras. Mer. i. t. 43 (1825), sphalm. = **Modiola**, Moench (Malvac.).

MADORELLA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 387 = **Madia**, Molina (Compos.).
dissitiflora, Nutt. l. c. = Madia dissitiflora.
racemosa, Nutt. l. c. = Madia sativa.

MADVIGIA, Liebm. Ind. Sem. Hort. Haun. (1854) 26; et in Ann. Sc. Nat. Sér. IV. ii. (1854) 373 = **Cryptanthus**, Otto & Dietr. (Bromel.).
densiflora, Liebm. ll. cc. 27; 373 = C. undulatus.
humilis, Liebm. ex Antoine, Bromel. 38 = Karatas humilis.

MAECHARANTHERA, Pritz. Ic. Ind. i. 673 (1855), sphalm. = MACHAERANTHERA, Nees = **Aster**, Tourn. (Compos.).

MAELENIA, Dum. in Mém. Acad. Sc. Brux. ix. (1834) 13. t. 10 = **Cattleya**, Lindl. (Orchid.).
paradoxa, Dum. l. c. = C. Forbesii.

MAERANTHUS, Benth. & Hook. f. Gen. i. 533 (1865), sphalm. = MARCANTHUS, Lour. = **Mucuna**, Adans. (Legumin.).

MAERLENSIA, Vell. Fl. Flum. 231 (1825); v. t. 112 (1827) = **Corchorus**, Linn. (Tiliac.).
hirsuta, Vell. ll. cc. = C. hirtus.

MAERUA, Forsk. Fl. Aegypt. Arab. 104 (1775). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
STREBLOCARPUS, Arn. in Ann. Sc. Nat. Sér. II. ii. (1834) 235.
WIEGMANNIA, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 787 (1841).
acuminata, *Oliver, Fl. Trop. Afr.* i. 85.—Afr. trop.
aethiopica, *Oliver, l. c.*—Afr. trop.
angolensis, *DC. Prod.* i. 254.—Afr. trop.
angustifolia, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 29. t. 8 = oblongifolia.
arabica, J. F. Gmel. Syst. 827 = crassifolia.
arenaria, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 171 = ovalifolia.
crassifolia, *Forsk. Fl. Aegypt. Arab.* p. cxiii.—Arab.
Currori, *Hook. f. in Hook. Niger Fl.* 218.—Afr. trop.
Grantii, *Oliver, Fl. Trop. Afr.* i. 84.—Afr. trop.
lucida, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 33 = angolensis.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 466.—Madag.
nervosa, *Oliver, Fl. Trop. Afr.* i. 84.—Afr. trop.
oblongifolia, *A. Rich. Tent. Fl. Abyss.* i. 32. t. 5.—Abyss.
ovalifolia, *Cambess. in Jacquem. Voy. Bot.* 22. t. 24.—Ind. or.
pubescens, *C. Muell. in Peters, Reise Mossamb. Bot.* 166.—Afr. trop.
racemosa, *Vahl, Symb. Bot.* i. 36.—Arab.
retusa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 33 = angolensis.
rigida, R. Br. ex G. Don, Gen. Syst. i. 286 = crassifolia.
scabra, Cambess. in Jacquem. Voy. Bot. 22. t. 23 = ovalifolia.
scandens, *C. Muell. in Peters, Reise Mossamb. Bot.* i. 165.—Afr. trop.
senegalensis, R. Br. ex G. Don, Gen. Syst. i. 286 = crassifolia.
Thomsoni, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 5.—Arab.
triphylla, *A. Rich. Tent. Fl. Abyss.* i. 32. t. 6.—Abyssin.
uniflora, Vahl, Symb. i. 36 = crassifolia.

MAESA, Forsk. Fl. Aegypt. Arab. 66 (1775). *MYRSINEAE*, Benth. & Hook. f. ii. 641.
BAEOBOTRYS, Forst. Char. Gen. t. 11 (1776).
DORAENA, Thunb. Diss. Nov. Gen. iii. (1783) 59.
SIBURATIA, Thou. Gen. Nov. Madag. 12 (1809).
acuminata, A. DC. in DC. Prod. viii. 77 = ramentacea.
acuminata, Wall. Cat. n. 2321 = Gymnospora acuminata.
alnifolia, *Harv. Thes. Cap.* ii. 20.—Afr. austr.
amboinensis, *Scheff. Comm. Myrsin. Arch. Ind.* 29.—Malaya.
andamanica, *Kurz, For. Fl. Brit. Burma*, ii. 575.—Ins. Andaman.

MAESA:—
angustifolia, *A. DC. in DC. Prod.* viii. 81.—Madag.
arabica, *J. F. Gmel. Syst.* 403.—Arab.
argentea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 233.—Reg. Himal.
Baeobotrys, *Roem. & Schult. Syst.* iv. 226.—Java.
Blumei, *D. Don, Gen. Syst.* iv. 22.—Java.
brachybotrya, *Miq. Fl. Ind. Bat.* ii. 1010.—Java.
canarana, Miq. ex Hook. f. Fl. Brit. Ind. iii. 510 = dubia.
Chisia, *D. Don, Prod. Fl. Nep.* 148.—Reg. Himal.
cordifolia, *Miq. Fl. Ind. Bat.* ii. 1008.—Java.
coriacea, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 300 = M. Doraena.
corylifolia, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 330.—Ins. Fiji.
costulata, *Miq. Fl. Ind. Bat.* ii. 1007.—Java.
dependens, *F. Muell. Fragm.* v. 107.—Austral.
dioica, A. DC. in DC. Prod. viii. 79 = M. Chisia.
Doraena, *Blume, ex Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 138.—Japon.
dubia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 235.—Malabar.
emirnensis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 79.—Madag.
floribunda, *Scheff. Comm. Myrsin. Archip. Ind.* 18.—Sumatra.
fulvinervis, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 80 = mollis.
Gaudichaudii, *A. DC. l. c.*—Ins. Philipp.
glabra, A. DC. in DC. Prod. viii. 82 = ramentacea.
grandifolia, *Miq. Fl. Ind. Bat.* ii. 1007.—Java.
haplobotrys, *F. Muell. Fragm.* v. 161.—Austral.
hirtella, *Miq. Pl. Jungh.* 186.—Java.
indica, Hook. f. in Journ. Linn. Soc. vi. (1861) 16 = lanceolata.
indica, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 230, in nota.—Ind. or.; Malaya.
japonica, *Mor. ex Zoll. Syst. Verz. Ind. Archip.* 61.—Japon.
Junghuhniana, *Scheff. Comm. Myrsin. Archip. Ind.* 25.—Sumatra.
Korthalsiana, *Scheff. l. c.* 18.—Sumatra.
laevigata, *Scheff. l. c.* 17.—N. Guin.
lanceolata, G. Don, Gen. Syst. iv. 22 = ramentacea.
lanceolata, *Forsk. Fl. Aegypt. Arab.* p. cvi.—Afr. bor.
lanceolata, Voigt, Hort. Suburb. Calc. 338 = indica.
latifolia, A. DC. in Trans. Linn. Soc. xvii. (1834) 133 = indica.
leptobotrya, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 224.—China.
leucocarpa, *Blume, ex Scheff. Comm. Myrsin. Archip. Ind.* 30.—Malaya.
lutea, *A. DC. in DC. Prod.* viii. 82.—Hab.?
macrocarpa, *Scheff. Comm. Myrsin. Archip. Ind.* 25.—Borneo.
macrophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 234.—Reg. Himal.
macrothyrsa, *Miq. Fl. Ind. Bat. Suppl.* 573.—Sumatra.
membranacea, Blanco, Fl. Filip. ed. II. Suppl. 590 = indica.
membranacea, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 80.—Ins. Philipp.
missionis, A. DC. in Trans. Linn. Soc. xvii. (1834) 134 = ramentacea.
mollis, *A. DC. l. c.*—Burma; Malaya.
mollissima, *A. DC. l. c.*—Burma; Malaya.
mollissima, Kurz, For. Fl. Br. Burma, ii. 100 = mollis.
montana, A. DC. in DC. Prod. viii. 79 = M. Chisia.
muscosa, A. DC. in Trans. Linn. Soc. xvii. (1834) 133 = membranacea.
muscosa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 87.—Burma.
nemoralis, A. DC. in Trans. Linn. Soc. xvii. (1834) 134 = M. Baeobotrys.
novo-guineensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 32.—N. Guin.
ovata, A. DC. in Trans. Linn. Soc. xvii. (1834) 133 = ramentacea.
ovata, Thou. ex Roem. & Schult. Syst. v. 226 = indica.
palustris, *Hochst. in Flora*, xxvii. (1844) 825.—Afr. austr.
paniculata, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 133.—Reg. Himal.; Burma.
parviflora, *Scheff. Comm. Myrsin. Archip. Ind.* 28.—Java.

MAESA:—
permollis, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 66 = mollis.
Perrottetiana, A. DC. in DC. Prod. viii. 80 = indica.
persicaefolia, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 330.—Ins. Fiji.
Pickeringii, *A. Gray, l. c.* 329.—Ins. Fiji.
picta, Hochst. in Flora, xxiv. (1841) I. Intell. 25 = lanceolata.
pisocarpa, *Blume, ex Scheff. Comm. Myrsin. Archip. Ind.* 21.—Ins. Celebes.
polyantha, *Scheff. l. c.* 16.—Borneo.
protracta, *F. Muell. Papuan Pl.* i. 92.—N. Guin.
pubescens, *G. Don, in Loud. Hort. Brit.* 70.—Ind. or.
pulchella, *Fawc. in Forbes, Natur. Wand. East. Archip.* 509.—Ins. Timor.
pyrifolia, *Miq. Fl. Ind. Bat.* ii. 1008.—Reg. Himal.; Java.
ramentacea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 230.—Ind. or.; Malaya.
Reinwardtii, *Blume, ex Scheff. Comm. Myrsin. Archip. Ind.* 27.—Malaya.
rubiginosa, *Blume, ex Scheff. l. c.* 26.—Malaya.
rufescens, A. DC. in DC. Prod. viii. 81 = lanceolata.
rugosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 508.—Reg. Himal.
sinensis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 80.—Hongkong.
subdendata, *A. DC. l. c.*—Cochinch.
sumatrana, Scheff. Comm. Myrsin. Archip. Ind. 15 = ramentacea.
tetrandra, *A. DC. in DC. Prod.* viii. 82.—Ins. Molucc.
tomentosa, D. Don, Prod. Fl. Nep. 148 = macrophylla.
trichophlebia, *Baker, in Journ. Linn. Soc.* xx. (1883) 197.—Madag.
verrucosa, Kurz, For. Fl. Brit. Burma, ii. 98 = andamanica.
verrucosa, *Scheff. Comm. Myrsin. Archip. Ind.* 46.—N. Guin.
virgata, A. DC. in Trans. Linn. Soc. xvii. (1834) 133 = indica.
vitiensis, *Seem. Fl. Vit.* 148.—Ins. Fiji.
Zollingeri, *Scheff. Comm. Myrsin. Archip. Ind.* 22.—Java.

MAESIA, Gaertn. Fruct. i. 344. t. 70 (1788) = **Gomphia**? Schreb. (Ochnac.).
serrata, Gaertn. l. c. = G. angustifolia?

MAESOBOTRYA, Benth. in Hook. Ic. Pl. t. 1296 (1879). *EUPHORBIABEAE*, Benth. & Hook. f. iii. 284.
floribunda, *Benth. in Hook. Ic. Pl.* t. 1296.—Afr. trop.

MAFUREIRA, Bertol. in Mem. Acc. Sc. Bolog. ii. (1850) 269 (Misc. Bot. ix. 6. t. 2) = **Trichilia**, Linn. (Meliac.).
oleifera, Bertol. ll. cc. 270; 6 = T. emetica.

MAGALLANA, Cav. Ic. iv. 50. t. 374 (1797) = **Tropaeolum**, Linn. (Geraniac.).
porifolia, Cav. l. c. 51. t. 374 = T. pentaphyllum.
porrifolia, DC. Prod. i. 684 sphalm. = praec.

MAGALLANA, Comm. ex DC. Syst. i. 443 (1818) = MAGELLANIA, Comm. = **Drimys**, Forst. (Magnol.).

MAGASTACHYA, Beauv. Agrost. 74 (1812) err. typ. = MEGASTACHYA, Beauv. = **Eragrostis**, Beauv. (Gramin.).

MAGELLANA, Poir. in Dict. Sc. Nat. xxviii. 13 (1823) = MAGALLANA, Cav. = **Tropaeolum**, Linn. (Geraniac.).

MAGELLANIA, Comm. ex Lam. Encyc. ii. 330 (1786) = **Drimys**, Forst. (Magnol.).

MAGHANIA, Steud. Nom. ed. II. ii. 89 (1841) = MOGHANIA, Jaume St. Hil. = **Flemingia**, Roxb. (Legumin.).

MAGNOLIA, Linn. Syst. ed. I (1735); Gen. ed. I. 162 (1737). *MAGNOLIACEAE*, Benth. & Hook. f. i. 18.
GWILLIMIA, Rottl. ex DC. Syst. i. 455 (1818).
KOBUS, Kaempf. ex Salisb. Parad. Lond. t. 87 (1807).
LIRIANTHE, Spach, Hist. Vég. Phan. vii. 485 (1839).

MAGNOLIA:—
SPHENOCARPUS (vel SPHAEROCARPUS), Wall. Cat. p. 236, sub n. 975 (1832?).
TULIPASTRUM, Spach, Hist. Vég. Phan. vii. 481 (1839).
YULANIA, Spach, l. c. 462 (1839).
acuminata, *Linn. Syst.* ed. X. 1082.—Am. bor.
alexandrina, Hort. ex Steud. Nom. ed. II. ii. 89 = M. Yulan.
angatensis, Blanco, Fl. Filip. ed. I. 859 = Talauma angatensis.
anonaefolia, Salisb. Parad. Lond. t. 5 = Michelia fuscata.
atropurpurea, Hort. ex Steud. Nom. ed. I. 504 = obovata.
auricularis, Salisb. Parad. Lond. t. 43 = Fraseri.
auriculata, Desr. in Lam. Encyc. iii. 673 = Fraseri.
Bailloni, *Pierre, Fl. For. Cochinch.* t. 2.—Cambodia.
Burchelliana, Hort. ex Steud. Nom. ed. II. ii. 89 = glauca.
Campbelli, *Hook. f. & Thoms. Fl. Ind.* i. 77.—Reg. Himal.
Candollii, Link, Handb. ii. 375 = acuminata.
Champaca, Baill. ex Pierre, Fl. For. Cochinch. t. 3 = Michelia Champaca.
Championi, Benth. Fl. Hongk. 8 = pumila.
citriodora, Hort. ex Steud. Nom. ed. II. ii. 89 = M. Yulan.
Coco, DC. Syst. i. 459 = pumila.
compressa, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 417.—Japon.
conspicua, Salisb. Parad. Lond. t. 38 = M. Yulan.
cordata, *Michx. Fl. Bor. Am.* i. 328.—Am. bor.
cyathiformis, Ruiz, ex C. Koch, Dendrol. i. 376 = M. Yulan.
dealbata, *Zucc. in Abh. Akad. Muench.* ii. (1836) 373. tt. 3, 4.—Mexic.
Decandollii, Savi, in Bibliot. Ital. xvi. (1819) 224 = acuminata.
denudata, Desr. in Lam. Encyc. iii. 675 = obovata.
discolor, Vent. Jard. Malm. 24 = obovata.
Duperreana, *Pierre, Fl. For. Cochinch.* t. 1.—Cambodia.
elliptica, Link, Handb. ii. 375 = grandiflora.
excelsa, Wall. Tent. Fl. Nep. 5. t. 2 = Michelia excelsa.
fasciata, Vent. Jard. Malm. n. 24. adn. 2 = Michelia fuscata.
fatiscens, Rich. ex DC. Prod. i. 82 = Talauma Plumieri.
Figo, *DC. Syst.* i. 460.—China.
fragrans, Rafin. Fl. Ludov. 91 (Quid?).—Am. bor.
fragrans, Salisb. Prod. 379 = glauca.
Fraseri, *Walt. Fl. Carol.* 159.—Am. bor.
frondosa, Salisb. Prod. 379 = M. Umbrella.
fuscata, Andr. Bot. Rep. t. 229 = Michelia fuscata.
glauca, *Linn. Syst.* ed. X. 1082.—Am. bor.
glauca, [Moç. & Sesse,] ex DC. Prod. i. 80 = Talauma mexicana.
glauca, Pierre, Fl. For. Cochinch. sub t. 2 = Talauma glauca.
glauca, Thunb. Fl. Jap. 236 = hypoleuca.
globosa, *Hook. f. & Thoms. Fl. Ind.* i. 77.—Reg. Himal.
Gordoniana, Hort. ex Steud. Nom. ed. II. ii. 90 = glauca.
gracilis, Salisb. Parad. Lond. t. 87 = M. Kobus.
grandiflora, *Linn. Syst.* ed. X. 1082.—Am. bor.
grandiflora, [Moç. & Sesse] ex DC. Prod. i. 80 = Talauma mexicana.
Griffithii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 41.—Reg. Himal.
Halleana, Hort. ex Flor. Mag. N. S. (1878) sub t. 309 = stellata.
Hoonoki, Siebold, in Verh. Batav. Gen. xii. (1830) 50 = hypoleuca.
hypoleuca, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 187.—Japon.
inodora, *DC. Syst.* i. 459.—China.
insignis, *Blume, Fl. Jav. Magnol.* 23.—Ind. or.
insignis, Wall. Tent. Fl. Nap. 5. t. 1 = Manglietia insignis.
Kobus, *DC. Syst.* i. 456.—Japon.
lanceolata, Link, Handb. ii. 375 = grandiflora.
Lennei ×, Van Houtte, Fl. des Serres, xvi. (1865-67) t. 1693-94.
liliflora, Desr. in Lam. Encyc. iii. 675 = obovata.
liliifera, Baill. Hist. Pl. i. 141 = M. Coco.
longifolia, Sweet, Hort. Brit. ed. I. 11 = glauca.
macrophylla, *Michx. Fl. Bor. Am.* i. 328.—Am. bor.
maxima, Lodd. ex G. Don, in Loud. Hort. Brit. 226 = acuminata.

MAGNOLIA:—
meleagrioides, Hort. ex DC. Syst. i. 458 = Michelia fuscata.
mexicana, DC. l. c. 451 = Talauma mexicana.
Michauxiana, Hort. ex DC. l. c. 455 = macrophylla.
mutabilis, *Regel, Cat. Pl. Hort. Aksakov.* 88.—Hab.?
obovata, Ait. ex Link, Handb. ii. 375 = grandiflora.
obovata, *Thunb. in Trans. Linn. Soc.* ii. (1794) 336. —Japon.
odoratissima, Reinw. ex Blume, Fl. Jav. Magnol. 32 = Talauma Candollei.
ovata, Spreng. Syst. iv. Cur. Post. 217 = Talauma ovata.
parviflora, Blume, Bijdr. 9 = Michelia fuscata.
parviflora, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 187.—Japon.
parvifolia, *DC. Syst.* i. 459.—Java.
pensylvanica, Hort. ex DC. l. c. 453 = acuminata.
Plumieri, Sw. Prod. Veg. Ind. Occ. 87 = Talauma Plumieri.
portoricensis, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1880) 233.—Porto Rico.
Precia, Correa, ex Vent. Jard. Malm. 24, in adnot. 2 = M. Yulan.
pterocarpa, *Roxb. Corom. Pl.* iii. 62. t. 266.—Reg. Himal.
pumila, *Andr. Bot. Rep.* t. 226.—Java; China.
punduana, *Wall. Cat.* n. 974.—Reg. Himal.
purpurea, Curt. Bot. Mag. t. 390 = obovata.
pyramidata, Bartr. ex Pursh, Fl. Am. Sept. ii. 382 = Fraseri.
Rumphii, Spreng. Syst. iv. Cur. Post. 217 = Talauma Rumphii.
rustica, Hort. ex DC. Syst. i. 453 = acuminata.
salicifolia, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 418.—Japon.
Schiedeana, *Schlecht. in Bot. Zeit.* xxii. (1864) 144.—Mexic.
Selloi, Spreng. Syst. iv. Cur. Post. 216 = Talauma Sellowiana.
Soulangiana ×, Hort. [ex Thieb.] in Mém. Soc. Linn. Par. v. (1827) p. lxxiii.
speciosa ×, Reichb. Fl. Exot. v. 37. i. 343.
sphenocarpa, Hook. f. & Thoms. Fl. Ind. i. 78, errore speciosa = pterocarpa.
stellata, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 418.—Japon.
Thomsoniana, Hort. ex Steud. Nom. ed. II. ii. 90 = glauca.
tomentosa, Thunb. in Trans. Linn. Soc. ii. (1794) 336 = M. Kobus.
tripetala, Linn. Syst. ed. X. 1082 = M. Umbrella.
umbellata, Hort. ex Steud. Nom. ed. II. ii. 90 = seq.
Umbrella, *Desr. in Lam. Encyc.* iii. 673.—Am. bor.
versicolor, Salisb. Parad. Lond. t. 5 = Michelia fuscata.
virginiana, Linn. Sp. Pl. 535 = acuminata, glauca, M. Umbrella.
Vrieseana, *Baill. ex Pierre, Fl. For. Cochinch.* sub t. 2. —Ins. Celebes.
Yulan, *Desf. Hist. Arb.* ii. 6.—China.

MAGNUSIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 125; et in Abh. Akad. Berl. (1855) 101. t. 9 B = **Begonia**, Linn.
fusca, Klotzsch, in Abh. Akad. Berl. (1855) 102.—Mexic.
maxima, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 125.—Mexic.

MAGONAEA, G. Don, Gen. Syst. i. 674 (1831) = seq.

MAGONIA, A. St. Hil. Pl. Rem. Brés. i. 239 (1824); et in Mém. Mus. Par. xii. (1825) 336. t. 12, 13. *SAPINDACEAE*, Benth. & Hook. f. i. 398.
PHAEOCARPUS, Mart. Nov. Gen. et Sp. i. 61. t. 37, 38 (1824).
glabrata, *A. St. Hil. Pl. Rem. Brés.* i. 241.—Bras.
pubescens, *A. St. Hil. l. c.* 239.—Bras.

MAGONIA, Vell. Fl. Flum. 165 (1825); iv. t. 60 (1827) = **Ruprechtia**, C. A. Mey. (Polygon.).
scandens, Vell. ll. cc. = R. laurifolia.

MAGYDARIS, Koch, ex DC. Coll. Mém. v. 68 (1829); et Prod. iv. 241 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 904.
ERIOCACHRYS, DC. Prod. iv. 241, in syn. (1830).
ambigua, DC. Prod. iv. 241 = panacina.
panacifolia, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 62 = seq.
panacina, *DC. Prod.* iv. 241.—Hispan.; Afr. bor.
tomentosa, *Koch, ex DC. Prod.* iv. 241.—Sicil.; Afr. bor.

MAHAGONI, Adans. Fam. ii. 343 (1763) = **Swietenia**, Jacq. (Meliac.).

MAHARANGA, DC. Prod. x. 71 (1846) = **Onosma**, Linn. (Borag.).
bicolor, DC. l. c. = O. bicolor.
Emodi, DC. l. c. = O. Emodi.
Wallichiana, DC. l. c. = O. Wallichianum.

MAHERNIA, Linn. Mant. i. 8 (1767). *STERCULIACEAE*, Benth. & Hook. f. i. 223.
abyssinica, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 29, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 75.—Afr. austr.
angustifolia, Harv. in Harv. & Sond. Fl. Cap. i. 219, in syn. (Hermanniae sp.).—Afr. austr.
anthemifolia, *Harv. l. c.* 213.—Afr. austr.
betonicaefolia, *Harv. l. c.* 218.—Afr. austr.
bicrenata, *E. Mey. in Drège, Zwei Pfl. Docum.* 200, nomen.—Afr. austr.
bipinnata, DC. Prod. i. 496 = pinnata.
biserrata, *Cav. Diss.* vi. 326.—Afr. austr.
biserrata, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 211 = pulchella.
borraginiflora, F. Muell. Fragm. ix. 42, in obs. = Hermannia boraginiflora.
brachycarpa, Harv. in Harv. & Sond. Fl. Cap. i. 212 = sisymbrifolia.
Burchellii, Sweet, Hort. Brit. ed. I. 57 = grandiflora.
chrysantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 219.—Afr. austr.
coccocarpa, *Eckl. & Zeyh. Enum.* 50.—Afr. austr.
cordata, E. Mey. in Drège, Zwei Pfl. Docum. 200 = betonicaefolia.
dentata, Harv. in Harv. & Sond. Fl. Cap. i. 216 = abyssinica.
diffusa, *Jacq. Hort. Schoenb.* ii. t. 201.—Afr. austr.
diffusa, E. Mey. ex Harv. & Sond. Fl. Cap. i. 211 = pulchella.
dryadiphylla, Eckl. & Zeyh. Enum. 51 = Hermannia dryadifolia.
Elliottiana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 589. —Afr. austr.
erodioides, *Burch. ex DC. Prod.* i. 496.—Afr. austr.
exappendiculata, *Mast. in Oliver, Fl. Trop. Afr.* i. 234. —Afr. trop.
fragrans, *Reichb. Ic. Pl. Cult.* i. t. 65.—Afr. austr.
geranioides, Harv. in Harv. & Sond. Fl. Cap. i. 213 = marginata.
Gilesii, F. Muell. Fragm. ix. 42, in obs. = Hermannia Gilesii.
glabrata, *Cav. Diss.* vi. 326. t. 200. f. 1.—Afr. austr.
glandulosa, Presl, Bot. Bemerk. 22 = pulchella.
gracilis, *Harv. in Harv. & Sond. Fl. Cap.* i. 215.—Afr. austr.
grandiflora, *Burch. Trav.* i. 295.—Afr. austr.
grandistipula, *Buching. ex Hochst. in Flora*, xxvii. (1844) 294.—Afr. austr.
heterophylla, *Cav. Diss.* vi. 324. t. 178. f. 1.—Afr. austr.
hilaris, Eckl. & Zeyh. Enum. 49 = Hermannia trifurca.
humifusa, *Eckl. & Zeyh. l. c.* 51.—Afr. austr.
humilis, *E. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 218.—Afr. austr.
incana, Eckl. & Zeyh. Enum. 49 = Hermannia trifurca.
incisa, Eckl. & Zeyh. l. c. 51 = vesicaria.
incisa, *Jacq. Hort. Schoenb.* i. t. 54.—Afr. austr.
jacobeaefolia, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 220 = pinnata.
lacera, E. Mey. in Drège, Zwei Pfl. Docum. 200 = scabra.
linearis, *Harv. in Harv. & Sond. Fl. Cap.* i. 217.—Afr. austr.
linnaeoides, *Burch. ex DC. Prod.* i. 497.—Afr. austr.
linnaeoides, E. Mey. in Drège, Zwei Pfl. Docum. 200 = nana.
marginata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 221.—Afr. austr.
modesta, *Planch. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 292.—Afr. austr.
multicaulis, *E. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 218.—Afr. austr.
multifida, *E. Mey. in Drège, Zwei Pfl. Docum.* 200.—Afr. austr.
myrrhifolia, Eckl. & Zeyh. Enum. 52 = pilosula.
namaquensis, E. Mey. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 219 = heterophylla.

MAHERNIA:—
nana, *Eckl. & Zeyh. Enum.* 50.—Afr. austr.
oblongifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 219. —Afr. austr.
odorata, Eckl. & Zeyh. Enum. 52 = Hermannia Presliana.
ovalis, Harv. in Harv. & Sond. Fl. Cap. i. 215 = saccifera.
ovata, *E. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 221.—Afr. austr.
oxalidiflora, *Burch. Trav.* i. 295.—Afr. austr.
oxalidiflora, Eckl. & Zeyh. Enum. 49 = grandiflora.
parviflora, *Eckl. & Zeyh. l. c.* 50.—Afr. austr.
pauciflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 220.—Afr. austr.
pilosula, *Harv. in Harv. & Sond. Fl. Cap.* i. 212.—Afr. austr.
pinnata, Eckl. & Zeyh. ex Harv. & Sond. l. c. 203 = Hermannia tenuifolia.
pinnata, *Linn. Syst.* ed. XII. 227.—Afr. austr.
pulchella, *Cav. Diss.* vi. 177. f. 3.—Afr. austr.
Quartiniana, Hochst. ex Walp. Ann. iii. 833 = abyssinica.
resedaefolia, *Burch. ex DC. Prod.* i. 496.—Afr. austr.
rotundata, E. Mey. in Drège, Zwei Pfl. Docum. 192 = linnaeoides.
rutila, *Jacq. Hort. Schoenb.* t. 263.—Afr. austr.
saccifera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 218.—Afr. austr.
salviaefolia, *Turcz. l. c.* 220.—Afr. austr.
scabra, *Eckl. & Zeyh. Enum.* 50.—Afr. austr.
scoparia, Eckl. & Zeyh. l. c. 51 = Hermannia scoparia.
sessilifolia, *DC. Prod.* i. 498.—Afr. austr.
sisymbriifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 221.—Afr. austr.
spinosa, Burch. Trav. i. 279 = Hermannia spinosa.
stellulata, *Harv. in Harv. & Sond. Fl. Cap.* i. 216.—Afr. austr.
stricta, E. Mey. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 222 = Hermannia stricta.
tomentosa, Harv. in Harv. & Sond. Fl. Cap. i. 219 (Hermanniae sp.).—Afr. austr.
tomentosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 219.—Afr. austr.
tomentosa, Wendl. f. in Bartl. & Wendl. f. Beitr. ii. 28 = saccifera.
vernicata, *Burch. Trav.* i. 278.—Afr. austr.
vernicata, Eckl. & Zeyh. Enum. 50 = saccifera.
veronicaefolia, *Eckl. & Zeyh. l. c.*—Afr. austr.
verticillata, *Linn. Mant.* i. 59.—Afr. austr.
vesicaria, *DC. Prod.* i. 497.—Afr. austr.
vestita, *Harv. in Harv. & Sond. Fl. Cap.* i. 218.—Afr. austr.
violacea, *Burch. ex DC. Prod.* i. 497.—Afr. austr.

MAHOMETA, DC. Prod. v. 457, in syn. (1836) = **Monarrhenus**, Cass. (Compos.).

MAHONIA, Nutt. Gen. Am. i. 211 (1818) = **Berberis**, Linn.
acanthifolia, Wall. ex G. Don, Gen. Syst. i. 118 = B. nepalensis.
Aquifolium, Nutt. Gen. Am. i. 212 = B. Aquifolium.
Beali, Pynaert, in Rev. Hortic. Belge, i. (1875) 156. f. 41 = B. Bealei.
diversifolia, Sweet, Brit. Flow. Gard. Ser. II. t. 94 = B. Aquifolium.
fascicularis, DC. Syst. ii. 19 = B. Aquifolium, pinnata.
glumacea, DC. l. c. 21 = B. nervosa.
ilicina, Schlecht. in Linnaea, x. (1835–36) 236 = B. ilicina.
intermedia, Dupuis, in Rev. Hortic. Sér. IV. ix. (1860) 22 = B. nepalensis?
japonica, DC. Syst. ii. 22 = B. nepalensis.
japonica, Thunb. Fl. Jap. 77; Ic. Jap. t. 22 = B. japonica.
Knightii, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 20 = B. Bealei.
napaulensis, DC. Syst. ii. 21 = B. nepalensis.
nervosa, Nutt. Gen. Am. i. 212 = B. nervosa.
paniculata, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 36.—Costa Rica.
repens, G. Don, Gen. Syst. i. 118 = B. repens.
tenuifolia, Loud. ex Steud. Nom. ed. II. i. 197, in syn. = B. tenuifolia.
trifolia, Cham. & Schlecht. in Linnaea, v. (1830) 211. = B. ilicina.
trifurca, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1346 = B. trifurca.

MAHUREA, Aubl. Pl. Gui. i. 558. t. 222 (1775). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 188.
BONNETIA, Schreb. Gen. 363 (1789).
casiquiarensis, *Benth. in Journ. Linn. Soc.* v. (1861) 64.—Bras.
exstipulata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 365.—Guian.
linguiformis, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 340.—Guian.
palustris, *Aubl. Pl. Gui.* i. 558. t. 222.—Guian.
speciosa, *Choisy, in DC. Prod.* i. 558.—Hab.?

MAIA, Salisb. Gen. Pl. Fragm. 64 (1866) = **Maianthemum**, Weber (Liliac.).
bifolia, Salisb. l. c. = Maianthemum Convallaria.

MAIANTHEMUM, [Weber, in] Wigg. Prim. Fl. Holsat. 14 (1780). *LILIACEAE*, Benth. & Hook. f. iii. 770.
BIFOLIUM, Gaertn. Mey. & Scherb. Fl. Wett. i. 209 (1799).
MAIA, Salisb. Gen. Pl. Fragm. 64 (1866).
SCIOPHILA, Wibel, Prim. Fl. Werth. 147 (1799).
STYRANDRA, Rafin. Am. Monthly Mag. (1818) 266.
UNIFOLIUM, Hall. Enum. Stirp. Helv. i. 288 (1742).
Bifolium, [DC.] in Red. Lil. iv. t. 216. f. 2 = M. Convallaria.
canadense, Desf. in Ann. Mus. Par. ix. (1807) 54 = M. Convallaria.
Convallaria, [Weber, in] Wigg. Prim. Fl. Holsat. 15. —Reg. bor. temp.
cordifolium, Moench, Meth. 638 = M. Convallaria.
racemosum, Link, Enum. Hort. Berol. i. 343 = Smilacina racemosa.
stellatum, Link, l. c. = Smilacina stellata.
umbellatum, Link, l. c. = Clintonia umbellata.

MAIETA, Aubl. Pl. Gui. i. 443. t. 176 (1775). *MELASTOMACEAE*, Benth. & Hook. f. i. 767.
HOSANGIA, Neck. Elem. i. 225 (1790).
angustifolia, Baill. Hist. Pl. vii. 35 = Heterotrichum angustifolium.
annulata, Vent. Choix, 32 = Marumia annulata.
argentea, Vent. l. c. 34, in adnot. = Miconia squamulosa.
dependens, Baill. Hist. Pl. vii. 35 = Clidemia dependens.
dispar, Miq. in Linnaea, xviii. (1844) 277 = guianensis.
guianensis, *Aubl. Pl. Gui.* i. 443. t. 176.—Guian.
heterophylla, *DC. Prod.* iii. 166.—Peruv.
hypophysca, DC. l. c. = guianensis.
macrophylla, Poepp. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 148 = Blakea hirsuta.
pilosa, Baill. Hist. Pl. vii. 19 (Clidemiae sp.).—Am. trop.
Poeppigii, *Mart. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 141.—Peruv.
rubra, Baill. Hist. Pl. vii. 35 = Clidemia rubra.
scalpta, Vent. Choix, 33 = Ossaea scalpta.

MAILELOU, Adans. Fam. ii. 200 (1763) = **Vitex**, Tourn. (Verben.).

MAILLARDIA, Frapp. & Duch. in Maill. Réunion, i. 148 (1863). *URTICACEAE*, Benth. & Hook. f. iii. 360.
borbonica, *Duch. in Maill. l. c.*—Ins. Borbon.
lancifolia, Frapp. ex Duch. l. c. = borbonica.

MAILLEA, Parl. Pl. Nov. 31 (1842). *GRAMINEAE*, Benth. & Hook. f. iii. 1146.
crypsoides, *Boiss. Fl. Orient.* v. 479.—Graecia; Sardin.
Urvillei, *Parl. Pl. Nov.* 31.—Graecia.

MAINEA, Vell. Fl. Flum. 275 (1825); vii. t. 8 (1827) = **Trigonia**, Aubl. (Vochys.).
racemosa, Vell. ll. cc. = T. crotonoides.

MAINGAYA, Oliver, in Trans. Linn. Soc. xxviii. (1873) 517. *HAMAMELIDEAE*.
malayana, *Oliver, l. c.*—Malaya.

MAIREANA, Moq. Chenop. Enum. 95 (1840); et in Ann. Sc. Nat. Sér. II. xv. (1841) 97. t. 13 = **Kochia**, Roth (Chenop.).
stelligera, F. Muell. Fragm. i. 139 = K. stelligera.
tomentosa, Moq. Chenop. Enum. 97 = K. villosa.

MAIRERIA, Scop. Introd. 192 (1777) = **Maripa**, Aubl. (Convolv.).

MAIRIA, Nees, Gen. et Sp. Ast. 247 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 266.
ZYRPHELIS, Cass. in Ann. Sc. Nat. Sér. I. xvii. (1829) 420.
Burchelli, *DC. Prod.* v. 218.—Afr. austr.
corymbosa, *Harv. in Harv. & Sond. Pl. Cap.* iii. 66.—Afr. austr.
crenata, *Nees, Gen. & Sp. Ast.* 248.—Afr. austr.
Ecklonis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 66.—Afr. austr.
foliosa, *Harv. l. c.*—Afr. austr.
hirsuta, *DC. Prod.* v. 217.—Afr. austr.
lasiocarpa, *DC. l. c.* 218.—Afr. austr.
lasiocarpa, Drège, ex Harv. & Sond. Fl. Cap. iii. 65 = taxifolia.
microcephala, *DC. Prod.* v. 218.—Afr. austr.
perezioides, *Nees, Gen. & Sp. Ast.* 249.—Afr. austr.
pinifolia, Sch. Bip. ex Walp. Rep. ii. 955 = taxifolia.
taxifolia, *DC. Prod.* v. 218.—Afr. austr.

MAIRRANIA, Neck. Elem. i. 219 (1791) = **Arctostaphylos**, Adans. (Ericac.).
alpina, Desv. Journ. Bot. i. (1813) 37 = A. alpina.
Uva-ursi, Desv. l. c. = A. Uva-ursi.

MAIS, Adans. Fam. ii. 39 (1763) = **Zea**, Mich. (Gramin.).

MAIZILLA, Schlecht. in Bot. Zeit. viii. (1850) 601 = **Paspalum**, Linn. (Gramin.).
stolonifera, Schlecht. l. c. 605 = P. stoloniferum.

MAJA, Klotzsch, in Schomb. Fauna et Fl. Guin. 1191 (1848) = **Cuphea**, P. Br. (Lythrar.).
hypericoides, Klotzsch, l. c. = C. mimuloides.

MAJA, Wedd. Chlor. And. i. 228. t. 27 (1856) = **Pterygopappus**, Hook. f. (Compos.).
compacta, Wedd. l. c. 229. t. 27 = P. Laurencii.

MAJIDEA, J. Kirk, ex Oliver, in Hook. Ic. Pl. t. 1097 (1871). *SAPINDACEAE*.
zanguebarica, *J. Kirk, l. c.*—Afr. trop.

MAJORANA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 235 (1745) = **Origanum**, Tourn. (Labiat.).
aegyptiaca, Kostel. Allg. Med.-Pharm. Fl. iii. 770 = O. Maru.
crassa, Moench, Meth. 406 = O. Majorana.
crassifolia, Benth. Lab. Gen. et Sp. 339 = O. Maru.
Dictamnus, Kostel. Allg. Med.-Pharm. Fl. iii. 770 = O. Dictamnus.
fragrans, Rafin. Fl. Tellur. iii. 86 = O. Majorana.
hortensis, Moench, Meth. 406 = O. Majorana.
mexicana, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 191 = O. Majorana?
microphylla, Benth. Lab. Gen. et Sp. 338 = O. microphyllum.
neglecta, Walp. Rep. iii. 697 = O. hirtum.
nervosa, Benth. Lab. Gen. et Sp. 339 = O. Maru?
Onites, Benth. l. c. = O. Onites.
Oreja, Walp. Rep. iii. 697 = O. Orega.
ovalifolia, Stokes, Bot. Mat. Med. iii. 350 = O. Majorana.
ovatifolia, Stokes, l. c. 351 = O. Majorana.
paniculata, Spenn. in T. Nees, Gen. Pl. Germ. Gamop. ii. n. 15 = O. paniculatum.
scutellifolia, Stokes, Bot. Mat. Med. iii. 349 = O. Maru.
sipylea, Kostel. Allg. Med. Pharm. Fl. iii. 770 = O. sipyleum.
suffruticosa, Rafin. Fl. Tellur. iii. 86 = O. Majorana?
syriaca, Kostel. Allg. Med. Pharm. Fl. iii. 769 = O. Maru.
tenuifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 381 = O. Majorana.
tenuifolia, Rafin. Fl. Tellur. iii. 86 = O. Majorana?
tomentosa, Stokes, Bot. Mat. Med. iii. 348 = O. Dictamnus.
turbinata, Walp. Rep. iii. 697 = O. turbinatum.
uncinata, Stokes, Bot. Mat. Med. iii. 353 = O. Majorana.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 381 = O. Majorana.

MALABAILA, Hoffm. Gen. Umb. 125 (1814). *UMBELLIFERAE*, Benth. & Hook. f. i. 923.
ANALYRIUM, E. Mey. ex Baill. Hist. Pl. vii. 206 (1879).
LEIOTULUS, Ehrenb. in Linnaea, iv. (1829) 399.
LOPHOTAENIA, Griseb. Spicil. Fl. Rumel. i. 377 (1843).
abyssinica, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 338.—Abyssin.
atropurpurea, *Vatke, in Linnaea*, xl. (1876) 189.—Abyssin.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 336.—Assyr.
aurea, *Boiss. Fl. Orient.* ii. 1053.—Graecia.
brachytaenia, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 338.—Persia.
carvifolia, *Boiss. & Bal. Diagn.* Ser. II. vi. 85.—As. Min.
dasyantha, Fisch. & Mey. ex C. Koch, in Linnaea, xvi. (1842) 359 = pimpinellifolia.
echinulata, Boiss. & Bal. Diagn. Ser. II. v. 100 = M. Sekakul.
graveolens, *Hoffm. Gen. Umb.* 126.—Rossia; Reg. Caucas.
hispidula, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 1056. —As. Min.
involucrata, *Boiss. & Sprun. in Ann. Sc. Nat.* Sér. III. i. (1844) 337.—Graecia.
Kotschyi, *Boiss. Diagn.* Ser. I. x. 42.—Persia.
lasiocarpa, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 337.—Kurdist.
numidica, *Coss. in Bull. Soc. Bot. Fr.* ix. (1862) 297.—Afr. bor.
obtusifolia, *Boiss. Fl. Orient.* ii. 1054.—As. Min.
Opoponax, Baill. Hist. Pl. vii. 186 = Opoponax Chironium.
orientalis, Benth. & Hook. f. Gen. i. 923 = Opoponax hispidus.
pastinacaefolia, *Boiss. & Bal. Diagn.* Ser. II. iv. 84.—Cappadoc.
pimpinellifolia, *Hoffm. Gen. Umb.* 126.—Reg. Caucas.
pimpinellifolia, Kotschy, ex Boiss. Fl. Orient. ii. 1058 = Kotschyi.
platyptera, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 336 = M. Sekakul.
pumila, *Boiss. Fl. Orient.* ii. 1058.—Aegypt.
rectistyla, Boiss. & Sprun. in Ann. Sc. Nat. Sér. III. i. (1844) 334 = aurea.
Sekakul, *Boiss. Fl. Orient.* ii. 1057.—As. Min.; Syria.
suaveolens, *Coss. in Bull. Soc. Bot. Fr.* xix. (1872) 82. —Aegypt.
sulcata, *Boiss. Fl. Orient.* ii. 1055.—Reg. Caucas.

MALABAILA, Tausch, in Flora, xvii. (1834) 356 = **Pleurospermum**, Hoffm. (Umbell.).
Golaka, Walp. Rep. ii. 425 = P. Golaka.
Hacquetii, Tausch, in Flora. xvii. (1834) 356 = P. Golaka.

MALABATHRIS, Rafin. Sylva Tellur. 97 (1838) = **Otanthera**, Blume (Melastom.).
cyanoides, Rafin. l. c. = O. moluccana.
nigra, Rafin. l. c. = praec.?

MALACARYA, Rafin. in Am. Monthly Mag. (1819) 190 = **Thalia**, Linn. (Scitam.).
dealbata, Rafin. l. c. = T. dealbata.

MALACHADENIA, Lindl. Bot. Reg. (1839) Misc. 67 = **Bulbophyllum**, Thou. (Orchid.).
clavata, Lindl. l. c.; et Hook. Bot. Mag. t. 4334.—Bras.

MALACHE, B. Vog. in Trew, Pl. Select. 50. t. 90 (1772) = **Pavonia**, Cav. (Malvac.).
scabra, B. Vog. l. c. = P. racemosa?

MALACHIUM, Fries, Fl. Hall. 77 (1817) = **Stellaria**, Linn. (Caryoph.).
aquaticum, Fries, l. c. = S. aquatica.
calycinum, Willk. in Bot. Zeit. v. (1847) 239 = Arenaria calycina.
coeruleum, Jaub. & Spach, Illustr. iii. 41 = Cerastium caeruleum.
erectum, Gren. in Acad. Sc. Besanç. Séances Publ. (1839) 186 = Cerastium quaternellum.
manticum, Reichb. Fl. Germ. Excurs. 795 = Cerastium manticum.
octandrum, Gren. in Acad. Sc. Besanç. Séances Pulb. (1839) 186 = Cerastium quaternellum.

MALACHOCHAETE, 'Nees,' ex Benth. & Hook. f. Gen. iii. 1051 (1883) = MALACOCHAETE, Nees = **Scirpus**, Tourn. (Cyperac.).

MALACHODENDRON, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 216 = **Stuartia**, Linn. (Ternstroem.).
corchoroides, J. F. Gmel. Syst. 1069 ; Raeusch. Nom. ed. III. 201 = Sida spinosa.
monogynum, Dum.-Cours. Bot. Cult. ed. II. v. 106 = S. Malachodendron.
ovatum, Cav. Diss. v. 302 = S. pentagyna.
pentagynum, Dum.-Cours. Bot. Cult. ed. II. v. 107 = S. pentagyna.

MALACHRA, Linn. Mant. i. 13 (1767). *MALVACEAE*, Benth. & Hook. f. i. 205.
alceaefolia, Jacq. Coll. ii. 350 ; Ic. Pl. Rar. iii. t. 549 = capitata.
Berterii, *Spreng. Syst.* iii. 94.—Am. austr.
bracteata, *Cav. Diss.* ii. t. 34. f. 2.—Am. austr.
capitata, *Linn. Syst.* ed. XII. 458.—Cosmop. trop.
ciliata, Poir. Encyc. Suppl. iii. 578 = urens.
conglomerata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 205 = capitata.
cordata, *Poir. Encyc. Suppl.* iii. 578.—Portorico.
digitata, *Presl, Rel. Haenk.* ii. 126.—Mexic.
fasciata, *Jacq. Coll.* ii. 352 ; *Ic. Pl. Rar.* iii. 11. t. 548.—Venezuela.
Gaudichaudiana, A. St. Hil. Fl. Bras. Mer. i. 218 = radiata.
helodes, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 61.—Bras.
heptaphylla, *Fisch. in Hornem. Hort. Hafn. Suppl.* 78.—Bras.
hispida, *Guill. & Perr. Fl. Seneg. Tent.* i. 47.—Afr. trop.
horrida, *Miq. Fl. Ind. Bat.* i. II. 149.—Ins. Timor.
humilis, *Benth. Bot. Voy. Sulph.* 70.—Ecuador.
Kegeliana, *Garcke, in Linnaea*, xxii. (1849) 52.—Bras.
linearilobа, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 206.—Ins. Philipp.
mexicana, Schrad. Cat. Gen. Hort. Gotting. (1830).—Cf. Linnaea, vi. (1831) Litt. 73 = capitata.
officinalis, *Klotzsch, in Bot. Zeit.* iv. (1846) 107.—N. Granat.
ovata, Presl, Rel. Haenk. ii. 125 = Pavonia sessiliflora.
palmata, *Moench, Meth.* 615.—Ind. occ.
plumosa, *Desr. in Lam. Encyc.* iii. 686.—Bras.
radiata, *Linn. Syst.* ed. XII. 458.—Am. trop.
rosea, *Hoffmgg. Verz. Pfl. Nachtr.* II. 154.—Hab. ?
rotundifolia, Schrank, Pl. Rar. Hort. Monac. t. 56 = capitata.
rudis, *Benth. Pl. Hartw.* 164.—Am. austr.
triloba, [Desf.] Hort. Par. ex Poir. Encyc. Suppl. iii. 578 = palmata.
trinervis, Presl, Rel. Haenk. ii. 126 = Pavonia sessiliflora.
Urena, *DC. Prod.* i. 441.—China.
urens, *Poit. in Schrad. Neues Journ.* ii. (1807) 293.—Ind. occ.
urticifolia, *Presl, Rel. Haenk.* ii. 126.—Mexic.
velutina, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 182.—N. Granat.
viminea, *Fisch. ex Steud. Nom.* ed. II. ii. 91.—Hab. ?

MALACION, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 183 = MALACHIUM, Fries, = **Stellaria**, Linn. (Caryoph.).

MALACMAEA, Griseb. in Linnaea, xiii. (1839) 248 = **Bunchosia**, Rich. & Juss. (Malpig.).
fluminensis, Griseb. l. c. = B. fluminensis.

MALACOCARPUS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 78. (1843) = **Peganum**, Linn. (Rutac.).
crithmifolius, Fisch. & Mey. l. c. 79 = P. crithmifolium.

MALACOCARPUS, Salm-Dyck, Cact. Hort. Dyck. ed. II. 24, 141 (1850) = **Echinocactus**, Link & Otto (Cact.).
aciculatus, Salm-Dyck, l. c. 25 = E. aciculatus.
acuatus, Salm-Dyck, l. c. = E. acuatus.
corynodes, Salm-Dyck, l. c. 25, 141 = E. corynodes.
Courantii, Salm-Dyck, l. c. 25, 142 = E. Courantii.
erinaceus, Lem. ex Foerst. Handb. Cact. ed. II. i. 455 = E. erinaceus.
Martini, Labour. ex Foerst. l. c. 454.—Hab. ?
polyacanthus, Salm-Dyck, Cact. Hort. Dyck. ed. II. 25 = E. polyacanthus.
Sellowianus, Salm-Dyck, l. c. 25, 142 = E. Sellowii.

MALACOCEPHALUS, Tausch, in Flora, xi. (1828) II. 481 = **Centaurea**, Linn. (Compos.).
altaicus, Tausch, l. c. = C. salina.
pinnatifidus, Tausch, l. c. 482 = Leuzea pinnatifida.

MALACOCHAETE, Nees, in Linnaea, ix. (1834) 292 = **Scirpus**, Tourn. (Cyperac.).
assimilis, Cesati, in Atti Soc. Sc. Nap. v. (1873) n. VII. 17 = S. riparius.
chilensis, Nees & Meyen, ex Boeck. in Linnaea, xxxvi. (1869–70) 717 = S. riparius.
corymbosa. Arn. ex Boeck. l. c. 706 = S. corymbosus.
Lehmanni, Nees, ex Boeck. l. c. 716 = S. littoralis.
littoralis, Nees, l. c. ix. (1834) 292 = S. littoralis.
oligostachya, Phil. l. c. xxix. (1857–58) 78 = S. riparius.
pectinata, Nees, l. c. ix. (1834) 292 = S. littoralis.
Pterolepis, Nees, l. c. ; et x. (1836) 184 = S. Pterolepis.
riparia, Nees & Meyen, l. c. ; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 90 = S. riparius.
sanguinolenta, Nees & Meyen, ex Boeck. in Linnaea, xxxvi. (1869–70) 718 = S. riparius.
Tatora, Meyen, Reise, i. 483, 491, nomen ; & in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 90 = S. Tatora.

MALACOIDES, Tourn. ex Adans. Fam. ii. 400 (1763) = **Malope**, Linn. (Malvac.).

MALACOMERIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 435 = **Malacothrix**, DC. (Compos.).
incanus, Nutt. l. c. = Malacothrix incana.

MALACOTHRIX, DC. Prod. vii. 192 (1838).
COMPOSITAE, Benth. & Hook. f. ii. 518.
LEPTOSERIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 438.
LEUCOSERIS, Nutt. l. c. (1841) 439.
MALACOMERIS, Nutt. l. c. (1841) 435.
altissima, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 195.—Calif.
californica, *DC. Prod.* vii. 192.—Calif.
Clevelandi, *A. Gray, in Bot. Calif.* i. 433.—Calif.
commutata, Torr. & Gray, Fl. N. Am. ii. 487 = saxatilis.
Coulteri, *Harv. & Gray, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 113.—Calif.
crepoides, A. Gray, in Pacif. Rail. Rep. xii. II. 53 = Crepis virens.
Fendleri, *A. Gray, Pl. Wright.* ii. 104.—Mexic.
glabrata, *A. Gray, Syn. Fl. N. Am.* i. II. 422.—Am. bor occ.
incana, *Torr. & Gray, Fl. N. Am.* ii. 486.—Calif.
insularis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 194.—Calif.
obtusa, *Benth. Pl. Hartw.* 321.—Calif.
parviflora, Benth. l. c. = obtusa.
platyphylla, A. Gray, in Proc. Am. Acad. ix. (1874) 214 = Atrichoseris platyphylla.
saxatilis, *Torr. & Gray, Fl. N. Am.* ii. 486.—Calif.
sonchoides, *Torr. & Gray, l. c.*—Am. bor. occ.
sonchoides, Torr. in Stansb. Exped. Utah, 392 = Torreyi.
tenuifolia, Torr. & Gray, Fl. N. Am. ii. 487 = saxatilis.
Torreyi, *A. Gray, in Proc. Am. Acad.* ix. (1874) 213.—Am. bor. occ.
Xanti, *A. Gray, l. c.*—Calif.

MALACOXYLUM, Jacq. Fragm. 31. t. 35 A (1800 ?) = **Zanthoxylum** ? Linn. (Rutac.).
pinnatum, Jacq. l. c.—Hab. ?

MALAISIA, Blanco, Fl. Filip. ed. I. 789 (1837) ; ed. II. 543 (1845). *URTICACEAE*, Benth. & Hook. f. iii. 360.
CATURUS, Lour. Fl. Cochinch. 612 (1790).
CEPHALOTROPHIS, Blume, Mus. Bot. Lugd. Bat. ii. 75, t. 27 (1853).
DUMARTROYA, Gaudich. Bot. Voy. Bonite, t. 97 (1844–52).
aculeata, Blume, Mus. Bot. Lugd. Bat. ii. 76 = tortuosa.
acuminata, Planch. in Ann. Sc. Nat. Sér. IV. iii. (1855) 294 = tortuosa.
Cunninghamii, Planch. l. c. = tortuosa.
puberula, *Bur. in DC. Prod.* xvii. 222.—Sumatra.
scandens, Planch. in Ann. Sc. Nat. Sér. IV. iii. (1855) 293 = tortuosa.
spinosa, Blume, Mus. Bot. Lugd. Bat. ii. 76 = tortuosa ?
tortuosa, *Blanco, Fl. Filip.* ed. I. 789.—Malaya.
viridescens, Planch. in Ann. Sc. Nat. Sér. IV. iii. (1855) 293 = tortuosa.

MALANEA, Aubl. Pl. Gui. i. 106. t. 41 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 103.
CUNNINGHAMIA, Schreb. Gen. 789 (1789).
angustifolia, Bartl. in Schomb. Fauna & Fl. Brit. Gui. 947, nomen (Quid. ?).—Guian.
bahiensis, *Muell. Arg. in Flora*, lviii. (1875) 453.—Bras.
bifurcata, Desr. in Lam. Encyc. iii. 688 = Antirrhoea bifurcata.
citrifolia, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1854) 203.—Ind. occ.
dichotoma, *A. Rich. Fl. Cub. Fanerog.* ii. 22.—Cuba.
dioica, Thou. ex DC. Prod. iv. 460, in obs. = Antirrhoea verticillata.
erecta, *Seem. Bot. Voy. Herald*, 136.—Panama.
evenosa, *Muell. Arg. in Flora*, lviii. (1875) 453.—Bras.
forsteronioides, *Muell. Arg. l. c.*—Bras.
gabrielensis, *Muell. l. c.*—Bras.
glabra, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 203.—Guian.
glabrescens, Bartl. in Schomb. Fauna & Fl. Brit. Gui. 947 = sarmentosa.
lucida, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 203.—Ind. occ.
macrophylla, *Bartl. in Schomb. Faun. & Fl. Brit. Gui.* 947 ; *Griseb. Fl. Brit. W. Ind.* 337.—Ind. occ.
Martiana, *Muell. Arg. in Flora*, lviii. (1875) 453.—Bras.
nitida, Desr. in Lam. Encyc. iii. 688 (*Stenostomum nitidum*).—Ind. occ.
panurensis, *Muell. Arg. in Flora*, lviii. (1875) 453.—Bras.
parviflora, Muell. Arg. l. c. = Chomelia parviflora.
ribesioides, Muell. Arg. l. c. = Chomelia ribesioides.
sarmentosa, *Aubl. Pl. Gui.* i. 106. t. 41.—Ind. occ. ; Guian.
spicata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 457.—Bras.
verticillata, Lam. Encyc. iii. 688 = Danais fragrans.
verticillata, Sieber, ex DC. Prod. iv. 361 = Danais rotundifolia.

MALAPARIUS, Miq. Fl. Ind. Bat. i. I. 1082 (1855) = **Pterocarpus** ? Linn. (Legumin.).
flavus, Miq. l. c. = P. indicus ?

MALASMA, Scop. Introd. 178 (1777), sphalm. = **Melasma**, Berg. (Scrophul.).

MALASPINAEA, Presl, Reliq. Haenk. ii. 68. t. 61 (1830) = **Aegiceras**, Gaertn. (Myrsin.).
laurifolia, Presl, l. c. = A. majus.

MALAXIS, Soland. ex Sw. Prod. Veg. Ind. Occ. 119 (1778) ; et in Vet. Acad. Nya Handl. Stockh. xxi. (1800) 233, partim. *ORCHIDEAE*, Benth. & Hook. f. iii. 493
LIMNAS, Ehrh. Beitr. iv. 146 (1789).
acuminata, D. Don, Prod. Fl. Nep. 29 = Microstylis Wallichii.
affinis, Blume, Bijdr. 392 = Liparis affinis.
anceps, Reichb. f. in Walp. Ann. vi. 214 = Oberonia anceps.
angustifolia, Blume, Bijdr. 393 = Liparis caespitosa.
angustifolia, Reichb. f. in Walp. Ann. vi. 213 = Oberonia angustifolia.
anthropophora, Reichb. f. l. c. 215 = Oberonia anthropophora.
aporophylla, Reichb. f. l. c. 214 = Oberonia aporophylla.
atropurpurea, Blume, Bijdr. 390 (= *Platystylis atropurpurea*).—Java.
autumnalis, Spreng. Pugill. ii. 81 = Earina mucronata.
bicornis, Reichb. f. in Walp. Ann. vi. 211 = Oberonia bicornis.
brachystachya, Reichb. f. l. c. = Oberonia brachystachys.
brasiliensis, *Spreng. Syst.* iii. 740.—Bras.
brevifolia, Reichb. f. in Walp. Ann. vi. 215 = Oberonia brevifolia.
Brunoniana, Reichb. f. l. c. 209 = Oberonia Brunoniana.
caespitosa, Thou. Orch. Iles Austr. Afr. t. 90 = Liparis caespitosa.
caudata, Willd. Sp. Pl. iv. 93 = Brassia caudata.
caulescens, Reichb. f. in Walp. Ann. vi. 215 = Oberonia caulescens.
cernua, Willd. Sp. Pl. iv. 93 = Geodorum dilatatum.
cimicina, Reichb. f. in Walp. Ann. vi. 216 = Oberonia Griffithiana.
cochleariaefolia, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 = Microstylis ichthyorrhyncha.
compressa, Blume, Bijdr. 390 = Liparis compressa.
cordifolia, Sm. in Rees, Cycl. xxii. n. 12 = Liparis nepalensis.
Correanaa, Bart. Fl. Philad. Prod. 86 = Liparis Loeselii.
crenulata, Blume, Bijdr. 393 = Liparis crenulata.

MALAXIS:—
cylindrica, Reichb. f. in Walp. Ann. vi. 208 = Oberonia cylindrica.
decurrens, Blume, Bijdr. 390. t. 54 (= *Platystylis decurrens*).—Java.
demissa, Reichb. f. in Walp. Ann. vi. 211 = Oberonia demissa.
denticulata, Reichb. f. l. c. 208 = Oberonia iridifolia.
diphyllos, Cham. & Schlecht. in Linnaea, iii. (1828) 34 = Microstylis monophylla.
disticha, Thou. Orch. Iles Austr. Afr. t. 89 = Liparis disticha.
elliptica, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 (sp. dub.).—Mexic.
ensiformis, Sm. in Rees, Cycl. xxii. n. 14 = Oberonia ensiformis.
equitans, Blume, Bijdr. 395 = Oberonia spathulata.
flavescens, Thou. Orch. Iles Austr. Afr. t. 25 = Liparis flavescens.
forcipata, Reichb. f. in Walp. Ann. vi. 208 = Oberonia forcipata.
Galeottiana, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 = Liparis elliptica.
glandulosa, Reichb. f. in Walp. Ann. vi. 215 = Oberonia glandulosa.
Griffithiana, Reichb. f. l. c. 208 = Oberonia Griffithiana.
heliophila, *Reichb. f. Otia Bot. Hamb.* 56.—Ins. Fiji et Samoa.
ichthiorhynca, A. Rich. & Gal. in Ann. Sc. Nat Sér. III. iii. (1845) 18 = Microstylis ichthyorrhyncha.
imbricata, Blume, Bijdr. 395 = Oberonia imbricata.
iridifolia, Reichb. f. in Walp. Ann. vi. 208 = Oberonia iridifolia.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 257.—Japon.
Jenkinsiana, Reichb. f. in Walp. Ann. vi. 211 = Oberonia Jenkinsiana.
lancifolia, Sm. in Rees, Cycl. xxii. n. 7 = Liparis odorata.
latifolia, Blume, Bijdr. 393 = Liparis latifolia.
latifolia, Sm. in Rees, Cycl. xxii. n. 3 = Microstylis congesta.
liliifolia, Sw. in Vet. Akad. Nya Handl. Stockh. xxi. (1800) 235 = Liparis liliifolia.
Lindeniana, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 = Liparis elliptica.
Lindleyana, Reichb. f. in Walp. Ann. vi. 210 = Oberonia Lindleyana.
Lobbiana, Reichb. f. l. c. 216 = Oberonia Lobbiana.
Loeselii, Sw. in Vet. Akad. Nya Handl. Stockh. xxi. (1800) 235 = Liparis Loeselii.
longibracteata, Reichb. f. in Walp. Ann. vi. 209 = Oberonia longibracteata.
longifolia, Bart. Fl. Philad. ii. 142 = Liparis Loeselii.
lunata, Blume, Bijdr. 394 = Oberonia lunata.
maianthemifolia, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 = Microstylis fastigiata.
majanthemifolia, Cham. & Schlecht. in Linnaea, vi. (1831) 59 = Microstylis majanthemifolia.
microphylla, Blume, Bijdr. 395 = Oberonia microphylla.
miniata, Reichb. f. in Walp. Ann. vi. 215 = Oberonia miniata.
minima, Blume, Bijdr. 391 = Liparis minima.
monophyllos, Sw. in Vet. Akad. Nya Handl. Stockh. xxi. (1800) 234 = Microstylis monophylla.
monstruosa, Blume, Bijdr. 395 = Oberonia monstruosa.
montana, Blume, Bijdr. 388 = Liparis montana.
monticola, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 (sp. dub.).—Mexic.
mucronata, Blume, Bijdr. 391 = Liparis disticha.
myodes, Bernh. Syst. Verz. Erf. 315 = Ophrys muscifera.
Myosurus, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 154 = Oberonia Myosurus.
myriantha, Reichb. f. in Walp. Ann. vi. 213 = Oberonia myriantha.
nervosa, Sw. in Vet. Akad. Nya Handl. Stockh. xxi. (1800) 235 = Liparis nervosa.
Nidus-avis, Bernh. Syst. Verz. Erf. 314 = Neottia Nidus-avis.
nutans, Willd. Sp. Pl. iv. 93 = Geodorum purpureum.
obcordata, Reichb. f. in Walp. Ann. vi. 216 = Oberonia obcordata.
oblongifolia, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 (sp. dub.).—Mexic.
odorata, Willd. Sp. Pl. iv. 91 = Liparis odorata.
ophioglossoides, Muhl. ex Willd. Sp. Pl. iv. 90 = Microstylis ophioglossoides.

MALAXIS:—
ovata, Bernh. Syst. Verz. Erf. 315 = Listera ovata.
pachystachys, *Reichb. f. in Walp. Ann.* vi. 214.—Ins. Pacif.
pallida, *Blume, Bijdr.* 391.—Java.
palmicola, F. Muell. Fragm. vii. 30 = Oberonia palmicola.
paludosa, *Sw. in Vet. Akad. Nya Handl. Stockh.* xxi. (1800) 235. t. 3.—Europ.; As. bor.
Parthoni, Morr. in Bull. Acad. Roy. Brux. (1839) 485 = Microstylis histionantha.
parviflora, Blume, Bijdr. 392 = Liparis parviflora.
platycaulon, Reichb. f. in Walp. Ann. vi. 209 = Oberonia platycaulon.
plicata, Roxb. Hort. Beng. 63; Fl. Ind. iii. 456 = Microstylis congesta.
Pumilio, *Reichb. f. in Walp. Ann.* vi. 216.—Java.
purpurascens, Thou. Orch. Iles Austr. Afr. tt. 26, 27 = Liparis purpurascens.
pyrulifera, Reichb. f. in Walp. Ann. vi. 211 = Oberonia pyrulifera.
Querci, De Vriese, Ic. 238; ex Lindl. Folia Orch. Oberonia, 8.—Hab. ?
recurva, Reichb. f. in Walp. Ann. vi. 212 = Oberonia recurva.
Rheedii, Blume, Bijdr. 389 = Liparis Rheedii.
Rheedii, Heyne, ex Wall. Cat. sub n. 1939 = Microstylis versicolor, *Lindl.*
Rheedii, Sw. in Vet. Acad. Nya Handl. Stockh. xxi. (1800) 235 = Microstylis Rheedii.
rufilabris, Reichb. f. in Walp. Ann. vi. 213 = Oberonia rufilabris.
Scyllae, Reichb. f. l. c. 213 = Oberonia Scyllae.
setifera, Reichb. f. l. c. 210 = Oberonia recurva.
sikkimensis, Reichb. f. l. c. 212 = Oberonia myriantha.
similis, Blume, Bijdr. 394 = Oberonia similis.
spathulata, Reichb. f. in Walp. Ann. vi. 215 = Oberonia spathulata.
spicata, Sw. Prod. Veg. Ind. Occ. 118 = Microstylis spicata.
spiralis, Reichb. f. in Walp. Ann. vi. 215 = Oberonia spiralis.
subulata, Labill. Nov. Holl. Pl. ii. 62. t. 212 = Cryptostylis longifolia.
tahitensis, Reichb. f. in Walp. Ann. vi. 208 = Oberonia tahitensis.
tenuis, Reichb. f. l. c. 211 = Oberonia tenuis.
Thisbe, Reichb. f. l. c. 213 = Oberonia Thisbe.
thlaspiformis, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 18 (sp. dub.).—Mexic.
Titania, *Reichb. f. in Walp. Ann.* vi. 216.—Ins. Norfolk.
tradescantifolia, Blume, Bijdr. 389 = Liparis tradescantifolia.
truncata, Reichb. f. in Walp. Ann. vi. 210 = Oberonia truncata.
umbellata, Steud. Nom. ed. II. ii. 144, in syn. = seq.
umbelliflora, Sw. Prod. Veg. Ind. Occ. 119 = Microstylis umbellulata.
umbellulata, Sw. Fl. Ind. Occ. iii. 1995 = praec.
unifolia, Michx. Fl. Bor. Am. ii. 157 = Microstylis ophioglossoides.
verticillata, Reichb. f. in Walp. Ann. vi. 210 = Oberonia verticillata.
Vieillardi, *Reichb. f. in Linnaea*, xli. (1877) 98.—N. Caled.
viridiflora, Blume, Bijdr. 392 = Liparis viridiflora.
Wightiana, Reichb. f. in Walp. Ann. vi. 212 = Oberonia Wightiana.

MALBRANCIA, Neck. Elem. ii. 366 (1790) = **Rourea**, Aubl. (Connar.).

MALCHOMIA, Sang. in Atti Acc. Pont. Lincei, Ser. I. xv. (1862) 36, 185 = seq.

MALCOLMIA, Spreng. Anleit. ii. II. 716 (1818) = seq.

MALCOMIA, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 121 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 77.
EREMOBIUM, Boiss. Fl. Orient. i. 156 (1867).
KLUKIA, Andrz. ex DC. Syst. ii. 459 (1821).
STRIGOSELLA, Boiss. Diagn. Ser. II. iv. 22 (1859).
WILCKIA, Scop. Introd. 317 (1777).
aculeolata, *Boiss. Fl. Orient.* i. 226.—Arab.; Persia; Beluch.

MALCOMIA:—
aegyptiaca, *Spreng. Syst.* ii. 898.—Aegypt.
africana, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 121.—Europ. austr.; Oriens.
alyssoides, DC. Syst. ii. 444 = littorea.
angulifolia, Boiss. & Orph. Diagn. Ser. II. v. 19 = cymbalaria.
arenaria, *DC. Syst.* ii. 442.—Afr. bor.
bicolor, *Boiss. & Heldr. Diagn.* Ser. I. vi. 10.—Graecia.
biloba, *Pomel, Nouv. Mat. Fl. Atl.* 229.—Alger.
binervis, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 71 = nana.
brevipes, *Boiss. Fl. Orient.* i. 226.—Persia; Turkest.
Broussonetii, DC. Syst. ii. 445 = littorea.
Bungei, *Boiss. Fl. Orient.* i. 226.—Afghan.; Turkest.
cabulica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 156.—Afghan.
chia, *DC. Syst.* ii. 440.—Reg. Mediterr. or.
circinata, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 155 = Bungei.
circinnata, *Boiss. Fl. Orient.* i. 227.—Turkest.
confusa, Boiss. Fl. Orient. i. 221 = nana.
conringioides, Boiss. l. c. 230 = exacoides.
contortuplicata, *Boiss. l. c.* 224.—Reg. Caucas.
crenulata, *Boiss. l. c.* 229.—Syria.
cymbalaria, *Heldr. & Sart. ex Boiss. l. c.* 227.—Graecia.
divaricata, Fisch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 33 = africana.
erosa, *DC. Syst.* ii. 446.—Baeotia.
exacoides, *Spreng. Syst.* ii. 899.—Syria.
flexuosa, *Sibth. & Sm. Fl. Graec.* t. 634.—Graecia.
graeca, *Boiss. & Sprun. Diagn.* Ser. I. i. 71.—Graecia.
incrassata, DC. Syst. ii. 441 = flexuosa.
intermedia, C. A. Mey. Verz. Pfl. Cauc. 186 = africana.
lacera, *DC. Syst.* ii. 445.—Hispan.; Marocc.
laxa, *DC. l. c.* 440.—Sibir.
Ledebourii, *Boiss. Fl. Orient.* i. 224.—Reg. Caucas.
littorea, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 121.—Reg. Mediterr. occ.
lyrata, Bory & Chaub. Fl. Pélop. 43, partim; ex Boiss. Diagn. Ser. I. vi. 11 = graeca.
lyrata, *DC. Syst.* ii. 443.—Ins. Cyprus.
maritima, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 121.—Reg. Mediterr.
Meyeri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 71.—Reg. Caucas.
mongolica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 422.—Mongolia.
multicaulis, *Pomel, Nouv. Mat. Fl. Atl.* 371.—Alger.
nana, *Boiss. Fl. Orient.* i. 222.—Reg. Mediterr. et Casp.
Orsiniana, *Tenore, Fl. Nap.* v. 67.—Europ.
parviflora, *DC. Syst.* ii. 442.—Europ. austr.
patula, *DC. l. c.* 444.—Hispan.
pulchella, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 70.—Syria.
pygmaea, *Boiss. Fl. Orient.* i. 222.—Aegypt; Persia.
runcinata, *C. A. Mey. Verz. Pfl. Cauc.* 186.—Reg. Casp.; Persia.
runcinata, Spreng. ex Steud. Nom. ed. I. 402, in syn. = Hesperis matronalis.
scorpioides, *Boiss. Fl. Orient.* i. 225.—Persia; Turkest.
stenopetala, Bernh. ex Ledeb. Fl. Ross. i. 170 = africana.
stricta, Cambess. in Jacquem. Voy. Bot. 16 = Sisymbrium strictum, *Hook. f. & Thoms.*
strigosa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 70.—Persia; Afghan.
taraxacifolia, DC. Syst. ii. 439 = runcinata.
torulosa, *Boiss. Fl. Orient.* i. 225.—Afr. bor.; Oriens.
trichocarpa, Boiss. ex Baker, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 21 = africana.
triloba, *Spreng. Syst.* ii. 899.—Reg. Mediterr.
veluchensis, Boiss. & Heldr. Diagn. Ser. II. vi. 10 = bicolor.
versicolor, *Pomel, Nouv. Mat. Fl. Atl.* 228.—Alger.

MALESHERBIA, Ruiz & Pav. Prod. 45 (1794). *PASSIFLOREAE*, Benth. & Hook. f. i. 809.
coerulea, Steud. Nom. ed. II. ii. 92 = Gynopleura caerulea.
coronata, D. Don, in Edinb. N. Phil. Journ. (1831–32) 112 = Gynopleura coronata.
deserticola, *Phil. Fl. Atac.* 18.—Chili.
dilatata, Phil. f. Cat. Pl. Vasc. Chil. 81, sphalm., vide *humilis*.

MALESHERBIA:—
fasciculata, D. Don, in Edinb. N. Phil. Journ. (1831–32) 206 = Gynopleura fasciculata.
humilis, D. Don, l. c. 111 = Gynopleura humilis.
lactea, *Phil. Fl. Atac.* 18.—Chili.
linearifolia, Poir. Encyc. Suppl. iii. 581 = Gynopleura linearifolia.
Lirana, *C. Gay, Fl. Chil.* ii. 422.—Chili.
oblongifolia, *Phil. in Anal. Univ. Chil.* (1865) II. 347.—Chili.
ovata, *Phil. Fl. Atac.* 18.—Chili.
paniculata, D. Don, in Edinb. N. Phil. Journ. (1827) 321 = Gynopleura paniculata.
propinqua, *C. Gay, Fl. Chil.* ii. 423.—Chili.
rugosa, *C. Gay, l. c.*—Chili.
solanoides, Meyen, Reise, i. 402 = Gynopleura solanoides.
subalpina, Poepp. ex M. Roem. Syn. Pepon. 210 = Gynopleura linearifolia.
tenuifolia, *D. Don, in Edin. N. Phil. Journ.* (Apr.–Oct. 1832) 244.—Peruv.
thyrsiflora, *Ruiz & Pav. Prod.* 45; *Fl. Per.* iii. 30.—Peruv.; Chili.
thyrsoidea, Bert. ex Steud. Nom. ed. II. ii. 92 = Gynopleura fasciculata.

MALICOPE, Vitm. Summa Veg. ii. 442 (1789) sphalm. = **Melicope**, Forst. (Rutac.).

MALIDRA, Rafin. Sylva Tellur. 107 (1838) = **Eugenia**, Mich. (Myrtac.).
aquea, Rafin. l. c. = E. aquea.

MALIGA, Rafin. Fl. Tellur. ii. 19 (1836) = **Allium**, Tourn. (Liliac.).
fastigiata, Rafin. l. c.—Hab.?
gracilis, Rafin. l. c. = Nothoscordum fragrans.
laxa, Rafin. l. c. = A. angulosum.

MALINVAUDIA, Fourn. in Mart. Fl. Bras. vi. IV. 312 (1885). *ASCLEPIADEAE.*
capillacea, *Fourn. l. c.* 313.—Bras.

MALLEA, A. Juss. in Mém. Mus. Par. xix. (1830) 221 = **Cipadessa**, Blume (Meliac.).
integerrima, Wall. ex Voigt, Hort. Suburb. Calc. 134 = C. fruticosa.
Rothii, A. Juss. in Mém. Mus. Par. xix. (1830) 221 = C. fruticosa.
subscandens, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 422 = C. fruticosa.

MALLINGTONIA, Willd. Sp. Pl. iii. 382 (1800) sphalm. = **Millingtonia**, Linn. f. (Bignon.).

MALLOCOCCA, Forst. Char. Gen. 77. t. 39 (1776) = **Grewia**, Linn. (Tiliac.).
crenata, Blanco, Fl. Filip. ed. I. 442 = G. umbellata.
crenata, Forst. Char. Gen. 78 = G. Mallococca.
parva, Blanco, Fl. Filip. ed. I. 443 = G. laevigata.

MALLOGONUM, Reichb. Handb. 236 (1837) = **Psammotropha**, Eckl. (Ficoid.).

MALLOPHORA, Endl. in Ann. Wien. Mus. ii. (1838) 206. *VERBENACEAE*, Benth. & Hook. f. ii. 1139.
LACHNOCEPHALUS, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 36.
corymbosa, *Endl. in Ann. Wien. Mus.* ii. (1838) 207.—Austral.
globifera, *Endl. l. c.* 206.—Austral.

MALLOSTOMA, Karst. Fl. Columb. ii. 9. t. 105 (1862). *RUBIACEAE*, Benth. & Hook. f. ii. 60.
ARCYTOPHYLLUM, [Willd.] Schult. Mant. iii. 5 (1827).
PSEUDORHACHICALLIS, Karst. Fl. Columb. ii. 10 (1862).
acerosum, Hemsl. Biol. Centr. Am. Bot. ii. 31 = Houstonia acerosa.
cachirense, *Karst. Fl. Columb.* ii. 9. t. 105.—N. Granat.
caracasanum, *Benth. & Hook. f. Gen.* ii. 60.—N. Granat.
confertum, *Benth. & Hook. f. l. c.*—Peruv.
filiforme, *Benth. & Hook. f. l. c.*—Peruv.
hyssopifolium, *Benth. & Hook. f. l. c.* 61.—Chili.
juniperifolium, *Benth. & Hook. f. l. c.*—Peruv.
laricifolium, *Benth. & Hook. f. l. c.*—Chili.
nitidum, *Benth. & Hook. f. l. c.* 60.—N. Granat.
setosum, *Benth. & Hook. f. . c.* 61.—Peruv.
thymifolium, *Benth. & Hook. f. l. c.*—Peruv.

MALLOTOPUS, Franch. & Sav. Enum. Pl. Jap. ii. 394 (1879). *COMPOSITAE.*
japonicus, *Franch. & Sav. l. c.*—Japon.

MALLOTUS, Lour. Fl. Cochinch. 635 (1790). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 319.
ADISCA, Blume, Bijdr. 609 (1825).
AXENFELDIA, Baill. Etud. Gén. Euph. 419 (1858).
BLUMEODENDRON, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 245.
BOUTONIA, Boj. Hort. Maurit. 282 (1837).
CORDEMOYA, Baill. Adansonia, i. (1860–61) 255.
DIPLOCHLAMYS, Muell. Arg. in Flora, xlvii. (1864) 539.
ECHINOCROTON, F. Muell. Fragm. i. 31 (1858).
ECHINUS, Lour. Fl. Cochinch. 633 (1790).
HANCEA, Seem. Bot. Voy. Herald, 409. t. 96 (1857)
LASIPANA, Rafin. Sylva Tellur. 21 (1858).
MELANOLEPIS, Reichb. & Zoll. in Verh. Natuurk. Ver. Nederl. Ind. i. (1856) 22; et in Linnaea, xxviii. (1856) 324.
PLAGIANTHERA, Reichb. & Zoll. ll. cc. 19; 321 (1856).
ROTTLERA, Roxb. Pl. Corom. ii. 36. t. 168 (1798).
STYLANTHUS, Reichb. & Zoll. in Linnaea, xxviii. (1856) 312.
acuminatus, *Muell. Arg. in. Linnaea*, xxxiv. (1865–66) 117.—Burma; Malaya.
albus, *Muell. Arg. l. c.* 188.—Reg. Himal.
amentiformis, Muell. Arg. in Flora, xlvii. (1864) 468 = floribundus.
angulatus, *Muell. Arg. in DC. Prod.* xv. II. 958.—Sumatra.
angustifolius, *Benth. Fl. Austral.* vi. 141.—Austral.
apelta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 189.—China.
atrovirens, *Muell. Arg. l. c.* 195.—Ind. or.
aureo-punctatus, Muell. Arg. in DC. Prod. xv. II. 973 = Lawii.
Baillonianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 186.—Madag.
barbatus, *Muell. Arg. l. c.* 184.—Ind. or.; Burma; Malaya.
Blumeanus, *Muell. Arg. l. c.* 195.—Java.
borneensis, *Muell. Arg. in DC. Prod.* xv. II. 980.—Borneo.
calcosus, *Muell. Arg. l. c.* 958.—Ins. Amboina.
capensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 189.—Afr. austr.
chinensis, Lour. Fl. Cochinch. 635 = cochinchinensis
claoxyloides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 192.—Afr. austr.
cochinchinensis, *Lour. Fl. Cochinch.* 635.—Burma; China; Malaya.
contubernalis, *Hance, in Journ. Bot.* xx. (1882) 293.—China.
Cumingii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 195.—Ins. Philipp.
Dallachyi, F. Muell. Fragm. vi. 184 = Macaranga Dallachyi.
decipiens, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 194.—Burma.
diadenus, *Muell. Arg. in DC. Prod.* xv. II. 959.—Sumatra.
discolor, *F. Muell. ex Benth. Fl. Austral.* vi. 143.—Austral.
dispar, *Muell. Arg. in DC. Prod.* xv. II. 971.—Java.
distans, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 194.—Ind. or.
eriocarpoides, Muell. Arg. l. c. 185 = Wallichianus.
eriocarpus, *Muell. Arg. l. c.*—Zeylan.
ferrugineus, *Muell. Arg. in DC. Prod.* xv. II. 982.—Ind. or.
floribundus, *Muell. Arg. in Flora*, xlvii. (1864) 469.—Burma; Malaya.
Furetianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 190.—China.
fuscescens, *Muell. Arg. l. c.* 195.—Zeylan.
glaberrimus, *Muell. Arg. l. c.* 192.—Java.
Helferi, *Muell. Arg. l. c.* 190.—Burma; Malaya.
Hookerianus, *Muell. Arg. l. c.* 193.—Ins. Hongkong.
inamoenus, F. Muell. ex Benth. Fl. Austral. vi. 146 = Macaranga inamoena.
insignis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 193.—Singapore.
integrifolius, *Muell. Arg. l. c.* 186.—Ins. Borbon.
japonicus, Muell. Arg. l. c. 189 = Rottlera japonica.

MALLOTUS:—
Korthalsii, *Muell. Arg. in DC. Prod.* xv. II. 976.—Borneo.
Lambertianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 190.—Ins. Molucc.
lappaceus, Muell. Arg. in DC. Prod. xv. II. 957 = Coelodiscus lappaceus.
Lawii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 192.—Ind. or.
leucocalyx, *Muell. Arg. in DC. Prod.* xv. II. 970.—Ins. Philipp.
longifolius, *Muell. Arg. l. c.* 967.—Sumatra.
longipes, Muell. Arg. in Linnaea, xxxiv. (1865–66) 193 = Coelodiscus hirsutulus.
macrostachyus, *Muell. Arg. in DC. Prod.* xv. II. 963.—Malaya.
Melleri, *Muell. Arg. in Flora*, xlvii. (1864) 468.—Afr. trop.
micrantha, Muell. Arg. in Linnaea, xxxiv. (1865–66) 191 = rhamnifolius.
moluccanus, *Muell. Arg. l. c.* 185.—Ins. Philipp.
Moritzianus, *Muell. Arg. in DC. Prod.* xv. II. 971.—Java.
muricatus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 191.—Ind. or.
nepalensis, *Muell. Arg. l. c.* 188.—Reg. Himal.
nesophilus, *Muell. Arg. l. c.* 196.—Austral.
nitidus, Muell. Arg. in DC. Prod. xv. II. 979 = Cleidion nitidum.
oblongifolius, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 192.—Java.
oppositifolius, *Muell. Arg. l. c.* 194.—Afr. trop.
oreophilus, Muell. Arg. l. c. 188 = nepalensis.
paniculatus, Muell. Arg. l. c. 189 = cochinchinensis.
peltatus, *Muell. Arg. l. c.* 187.—Ind. or.
penangensis, *Muell. Arg. l. c.* 186.—Malaya.
philippinensis, *Muell. Arg. l. c.* 196.—As. et Austral. trop.
polyadenos, *F. Muell. Fragm.* vi. 184.—Austral.
Porterianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 115.—Ins. Penang.
pycnostachys, F. Muell. Fragm. iv. 138 = ricinoides.
repandus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 197.—As. trop.; N. Caled.
rhamnifolius, *Muell. Arg. l. c.* 196.—Ind. or.
ricinoides, *Muell. Arg. l. c.* 187.—Burma; China; Malaya.
Roxburghianus, *Muell. Arg. l. c.* 186.—Reg. Himal.; Burma.
rufidulus, *Muell. Arg. in DC. Prod.* xv. II. 970.—Sumatra.
scabrifolius, Muell. Arg. l. c. 699, sphalm. = repandus.
scandens, *Muell. Arg. l. c.* 982.—Ins. Timor.
sphaerocarpus, *Muell. Arg. l. c.* 976.—Sumatra,
stenanthus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 191.—Ind. or.
stipulosus, *Muell. Arg. l. c.* 192.—Java.
stylaris, *Muell. Arg. in DC. Prod.* xv. II. 973.—Borneo.
subpeltatus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 189.—Malaya.
subulatus, *Muell. Arg. l. c.* 192.—Afr. trop.
tetracoccus, Kurz, in Journ. As. Soc. Beng. xli. (1873) II. 245 = albus.
tiliaefolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 190.—Java.
Tokbrai, *Muell. Arg. in DC. Prod.* xv. II. 956.—Java.
Wallichianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 196.—Burma.
zeylanicus, Muell. Arg. l. c. 195 = rhamnifolius.
Zippellii, F. Muell. Fragm. iv. 139 = ricinoides.
Zollingeri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 193.—Java.

MALNAREGAM, Adans. Fam. ii. 344 (1763) = **Atalantia**, Correa (Rutac.).

MALNEREGA, Rafin. Sylva Tellur. 143 (1838) = **Atalantia**, Correa (Rutac.).
malabarica, Rafin. l. c. = A. monophylla.

MALOCCHIA, Savi, in Nuov. Giorn. Pisa, viii. (1824) 113 = **Canavalia**, Adans. (Legumin.).
ensiformis, Savi, l. c. = C. ensiformis.
gladiata, Savi, l. c. 116 = C. ensiformis.

MALOCOPSIS, Walp. Ann. i. 104 (1848) = MALVEOPSIS, Presl = **Sphaeralcea**, A. St. Hil. (Malvac.).

MALOPE, Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 200.
MALACOIDES, Tourn. ex Adans. Fam. ii. 400 (1763).
althaeoides, Moris, in Reichb. Fl. Germ. Excurs. 873 = malacoides.
asterotricha, *Pomel, Nouv. Mat. Fl. Atl.* 346.—Alger.
ciliata, F. G. Dietr. in Neue Bot. Gart. Journ. i. 126 = trifida.
grandiflora, F. G. Dietr. Vollst. Lexik. Gaertn. II. Nachtr. v. 403; C. A. Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 75 = trifida.
hispida, *Boiss. Diagn.* Ser. II. i. 100.—Alger.
indica, Wight, Ic. iii. IV. 1, textu lapsu = Melicope indica.
laevigata, *Pomel, Nouv. Mat. Fl. Atl.* 347.—Alger.
lutea, *Rafin. Atl. Journ.* 147.—Hab. ?
malacoides, *Linn. Sp. Pl.* 692.—Europ. austr.; As. Min.
malacoides, Walt. Fl. Carol. 176 = Malvastrum tricuspidatum ?
multiflora, *Triguer. in Cav. Diss.* ii. 85.—Hispan.
parviflora, L'Hérit. Stirp. Nov. 105. t. 50 = Palavia malvaefolia.
rotundifolia, *C. C. Gmel. Hort. Carlsr. ex Steud. Nom.* ed. I. 505.—Hab. ?
stellipilis, *Boiss. & Reut. Diagn.* Ser. II. i. 101.—Hispan.
stipulacea, Cav. in Anal. Cienc. Nat. iii. (1801) 74 = malacoides.
trifida, *Cav. Diss.* ii. 85. t. 27. f. 2.—Hispan.; Afr. bor.
tripartita, *Boiss. & Reut. Diagn.* Ser. II. i. 101.—Alger.

MALORTIEA, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 25; 145. *PALMAE*, Benth. & Hook. f. iii. 906.
gracilis, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 26; 146.—Costa Rica.
intermedia, *H. Wendl. in Kerch. Palm.* 250, nomen.—Hab. ?
lacerata, *Hort. ex H. Wendl. l. c.*—Hab. ?
latisecta, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 146.—Guatemala.
simplex, *H. Wendl. in Bot. Zeit.* xvii. (1859) 5.—Costa Rica.
speciosa, *Hort. ex H. Wendl. in Kerch. Palm.* 250, nomen.—Hab. ?

MALOSMA, Engl. in DC. Monog. Phan. iv. 393 (1883) = **Rhus**, Tourn. (Anacard.).
laurina, Engl. l. c. = R. laurina.

MALOUETIA, A. DC. in DC. Prod. viii. 378 (1844) *APOCYNACEAE*, Benth. & Hook. f. ii. 703.
ROBBIA, A. DC. in DC. Prod. viii. 444 (1844).
albiflora, *Miq. Stirp. Surin. Sel.* 161.—Guian.
amplexicaulis, *Spruce, ex Muell. Arg. in Mart. Fl. Bras.* vi. I. 91.—Bras.
arborea, Miers, Apocyn. S. Am. 89 = Secondatia arborea.
asiatica, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 163.—Japon.
cestroides, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 94.—Bras.
cubana, *A. DC. in DC. Prod.* viii. 379.—Cuba.
flavescens, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 95.—Bras.
furfuracea, *Benth. ex Muell. Arg. l. c.* 93.—Bras.
glandulifera, Miers, Apocyn. S. Am. 90 = M. Tamaquarina.
gracilis, *A. DC. in DC. Prod.* viii. 380.—Guiana.
guianensis, Klotzsch, in Schomb. Faun. et Fl. Gui. 952, nomen; Miers, Apocyn. S. Am. 87 = M. Tamaquarina.
Heudeloti, *A. DC. in DC. Prod.* viii. 380.—Afr. trop.
jasminoides, *A. DC. l. c.* 379.—Venezuela; N. Granat.
lactiflua, *Miers, Apocyn. S. Am.* 88.—Guian.
lanceolata, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 93.—Bras.
Martii, *Muell. Arg. l. c.* 94.—Bras.
nitida, *Spruce, ex Muell. Arg. l. c.*—Bras.
obtusiloba, *A. DC. in DC. Prod.* viii. 379.—Guian.
odorata, *A. DC. l. c.*—Guian.
panamensis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 185.—N. Granat.
puberula, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 952, nomen.—Guiana.
retroflexa, *Muell. Arg. in Linnaea*, xxx. (1859–60) 408.—Ins. S. Vincent.

MALOUETIA:—
riparia, *A. DC. in DC. Prod.* viii. 380.—N. Granat.
Schomburgkiana, *Muell. Arg. in Linnaea*, xxx. (1859–60) 409.—Guian.
sessilis, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 96.—Bras.
Tamaquarina, *A. DC. in DC. Prod.* viii. 378.—Bras.
tetrastachya, Miers, Apocyn. S. Am. 92 = Tabernaemontana tetrastachya.
virescens, *Spruce, ex Muell. Arg. in Mart. Fl. Bras.* vi. I. 92.—Bras.

MALPIGHIA, Plum. ex Linn. Syst. ed. I (1735). *MALPIGHIACEAE*, Benth. & Hook. f. i. 251.
RUDOLPHIA, Medic. Malv. iii. (1787).
alternifolia, Schum. & Thonn. Beskr. Guin. Pl. 222 = Acridocarpus Smeathmanni.
altissima, Aubl. Pl. Gui. i. 455 = Byrsonima altissima.
altissima, Jacq. Obs. Bot. i. 40 = Byrsonima spicata.
ambigua, Salzm. ex A. Juss. in Arch. Mus. Par. iii. (1843) 277 = Byrsonima Salzmanniana.
angustifolia, *Linn. Sp. Pl.* ed. II. 610.—Ind. occ.
aquifolia, *Linn. l. c.* ed. I. 426.—Am. austr.
argentea, Jacq. Fragm. 57 = Bunchosia argentea.
armeniaca, Cav. Diss. 410 = Bunchosia armeniaca.
aureifolia, C. Muell. in Walp. Ann. vii. 467, sphalm. = variifolia.
bannisterioides, *Linn. Pl. Surin.* 55; *Amoen. Acad.* viii. 256.—Guian.
Berteriana, *Spreng. Syst.* ii. 383.—Ins. Guadalupe.
bicolor, Mart. ex Juss. Monogr. Malpig. 26 = Byrsonima verbascifolia.
biflora, Poir. in Lam. Encyc. iv. 326 = punicifolia.
biglandulosa, Poir. Encyc. Suppl. iv. 7 = Galphimia glandulosa.
brachystachya, Spreng. Syst. ii. 383 = Byrsonima brachystachya.
campechiensis, Poir. in Lam. Encyc. iv. 333 = incana.
canescens, [Soland in] Ait. Hort. Kew. ed. II. ii. 105 = Bunchosia canescens.
Caucanthus, Poir. Encyc. Suppl. iv. 6 = Caucanthus arabicus.
cinerea, Poir. Encyc. Suppl. iv. 7 = Byrsonima cinerea.
Cnide, *Spreng. Neue Entdeck.* iii. 51.—Ins. S. Doming.
coccigera, *Linn. Sp. Pl.* ed. II. 426.—Ind. occ.
Coccigrya, Linn. l. c. 611 = praec.
coccolobaefolia, Spreng. Syst. ii. 384 = Byrsonima coccolobaefolia.
coriacea, Sw. Prod. Veg. Ind. Occ. 74 = Byrsonima spicata.
cotinifolia, Spreng. Syst. ii. 384 = Byrsonima cotinifolia.
crassifolia, Linn. Sp. Pl. 426 = Byrsonima crassifolia.
crassifolia, Vell. Fl. Flum. 193; iv. t. 170 = Byrsonima verbascifolia.
cubensis, H. B. & K. Nov. Gen. et Sp. v. 145 = angustifolia.
cuneata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 390.—Cuba.
densa, Poir. Encyc. Suppl. iv. 7 = Byrsonima densa.
diphylla, Jacq. Enum. Pl. Carib. 21 = Byrsonima diphylla.
dubia, Cav. Diss. viii. 413. t. 242 = Heteropteris laurifolia.
elegans, G. F. W. Mey. Prim. Fl. Esseq. 178 = Byrsonima elegans.
elliptica, *Presl, Bot. Bemerk.* 25.—Ins. Trinit.
emarginata, [*Moç. & Sesse,*] *ex DC. Prod.* i. 578.—Mexic.
faginea, *Sw. Prod. Veg. Ind. Occ.* 74.—Ind. occ.
fallax, Salisb. Prod. 318 = glabra.
fasciculata, *Vell. Fl. Flum.* 194; iv. t. 174.—Bras.
fruticosa, Vell. l. c. 193; iv. t. 169 = Heteropteris saligna.
fucata, *Ker-Gawl. in Bot. Reg.* t. 189.—Jamaic.
Galeottiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 261.—Mexic.
glabra, *Linn. Sp. Pl.* 425.—Am. trop.
glandulifera, Jacq. Coll. iv. 207 = Bunchosia glandulifera.
glandulosa, Cav. Diss. viii. 411. t. 239. f. 2 = Bunchosia glandulosa.
glandulosa, Jacq. Ic. Pl. Rar. t. 469 = Bunchosia glandulifera.
glauca, Poir. Encyc. Suppl. iv. 7 = Galphimia glauca.
Gnidae, G. Don, Gen. Syst. i. 634, sphalm. = M. Cnide.
grandiflora, *Jacq. Enum. Pl. Carib.* 21.—Ins. Martiniq.
grandifolia, J. F. Gmel. Syst. 724, sphalm. = praec.
guadalupensis, Spreng. Syst. ii. 383 = Byrsonima spicata.

MALPIGHIA:—
heterantha, *Wight, Illustr.* i. 138. t. 49.—Ind. or.
heterophylla, *Griseb. in Linnaea*, xxii. (1849) 2.—Mexic.
hirsuta, Poir. Encyc. Suppl. iv. 6 = Galphimia hirsuta.
hirsuta, Vell. Fl. Flum. 193; iv. t. 171 = Byrsonima pachyphylla.
ilicifolia, Mill. Gard. Dict. ed. VIII. n. 8 = aquifolia.
incana, *Mill. l. c.* n. 3.—Honduras.
incana, Salisb. Prod. 318 = Bunchosia canescens.
infestissima, Rich. ex Walp. Rep. v. 151 = urens.
japurensis, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 305 = Byrsonima japurensis.
laevigata, Poir. in Lam. Encyc. iv. 332 = Byrsonima laevigata.
lanceolata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 537 = Bunchosia glandulosa.
lanceolata, Griseb. in Goett. Abh. vii. (1857) 185 = urens.
lanceolata, C. Muell. in Walp. Ann. vii. 467 = cuneata.
lanceolata, Poir. in Lam. Encyc. iv. 332 = Byrsonima crassifolia.
laurifolia, Spreng. Syst. ii. 385 = Byrsonima laurifolia.
linearis, Jacq. Enum. Pl. Carib. 21 = angustifolia.
Loddigesii, *Regel, Ind. Sem. Hort. Petrop.* (1857) 54.—Ind. occ.
lucida, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 20 = Byrsonima vacciniaefolia.
lucida, Pav. ex Moric. Pl. Nouv. Am. 109 = peruviana.
lucida, Sw. Prod. Veg. Ind. Occ. 74 = Byrsonima lucida.
macrophylla, Desf. Cat. Hort. Par. ed. III. 232 = fucata.
macrophylla, Pers. Syn. i. 506 = Byrsonima nervosa.
macrophylla, *Willd. Enum. Hort. Berol. Suppl.* 37.—Bras.
malifolia, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 21 = Banisteria crotonifolia.
maritima, Vell. Fl. Flum. 194; iv. t. 173 = Bunchosia fluminensis.
martinicensis, Jacq. Enum. Pl. Carib. 21 = urens.
media, [R. Br. in] Ait. Hort. Kew. ed. II. iii. 103 = Bunchosia media.
mexicana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 267.—Mexic.
mitis, Reichb. ex Griseb. Fl. Brit. W. Ind. 108 = Mammea humilis.
mollis, Poir. Encyc. Suppl. iv. 6 = Byrsonima mollis.
montana, Spreng. Syst. ii. 385 = Byrsonima crassifolia.
Moureila, Aubl. Pl. Gui. i. 459 = Byrsonima crassifolia.
myrtifolia, *Desf. Cat. Hort. Par.* ed. III. 406.—Ind. occ.
Neumanniana, *A. Juss. in Rev. Hortic.* Sér. II. v. (1846) 103.—Hab. ?
nitida, Cav. Diss. viii. t. 239 = Bunchosia polystachya.
nitida, *Crantz, Inst.* ii. 478.—Venezuela.
nitida, Jacq. Enum. Pl. Carib. 21 = Bunchosia nitida.
nitida, Mill. Gard. Dict. ed. VIII. n. 5 = glabra.
nitida, Sw. Obs. Bot. 180 = Bunchosia Swartziana.
oblongifolia, *Griseb. Ind. Sem. Hort. Gotting.* (1878) 8.—Hab. ?
obovata, *H. B. & K. Nov. Gen. et Sp.* v. 146.—N. Granat.
odorata, Jacq. Enum. Pl. Carib. 21 = Bunchosia odorata.
oxycocca, *Griseb. Fl. Brit. W. Ind.* 117.—Ind. occ.
pallida, Poir. Encyc. Suppl. iv. 7 = Byrsonima pallida.
paniculata, Mill. Gard. Dict. ed. VIII. n. 6 = Triopteris ovata.
parvifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 267.—Mexic.
peruviana, *Moric. Pl. Nouv. Am.* 109.—Peruv.
phlomoides, Spreng. Syst. ii. 385 = Tetrapteris rotundifolia.
piauhiensis, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 273 = Byrsonima sericea.
platyphylla, Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1827) 66 = Bunchosia polystachya.
polystachya, Andr. Bot. Rep. t. 604 = Bunchosia polystachya.
polytricha, A. Juss. in Arch. Mus. Par. iii. (1843) 258 (*polytrhica*) = setosa.
pruinosa, Sieber, ex A. Juss. l. c. 302 = Byrsonima coccolobaefolia.
pruinosa, Spreng. Syst. ii. 384 = Byrsonima angustifolia.
pulchra, [Moç. & Sesse,] ex DC. Prod. i. 580 = Byrsonima pulchra.
punicifolia, *Linn. Sp. Pl.* ed. II. 609.—Am. trop.
reticulata, Poir. Encyc. Suppl. iv. 8 = Heteropteris platyptera.
reticulata, Spreng. ex A. Juss. in Arch. Mus. Par. iii. (1843) 295 = Byrsonima spicata.

MALPIGHIA:—
retusa, *Benth. Bot. Voy. Sulph.* 74.—Ecuador.
rhopalaefolia, Spreng. Syst. ii. 384 = Byrsonima crassifolia.
rufa, Poir. in Lam. Encyc. iv. 332 = Byrsonima crassifolia.
rutilans, Salzm. ex Griseb. in Mart. Fl. Bras. xii. 1. 11 = Byrsonima sericea.
saccharina, *G. Don, Gen. Syst.* i. 635.—Afr. trop.
Semeruco, *A. Juss. in Arch. Mus. Par.* iii. (1843) 266. —Venezuela.
setosa, *Spreng. Neue Entdeck.* iii. 50.—Ind. occ.
singularis, Vell. Fl. Flum. 194; iv. t. 172 = Camarea affinis.
spicata, Cav. Diss. viii. 409 = Byrsonima spicata.
terminaliaefolia, Mart. ex Griseb. in Mart. Fl. Bras. xii. 1. 16 = Byrsonima vacciniifolia.
ternifolia, H. B. & K. Nov. Gen. et Sp. v. 146.—N. Granat.
tomentosa, *Pav. ex Moric. Pl. Nouv. Am.* 111. t. 68.—Peruv.
tuberculata, Jacq. Hort. Schoenb. i. 54 = Bunchosia tuberculata.
undulata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 266. —Mexic.
uniflora, Tussac, Fl. Ant. iii. t. 27 = punicifolia.
urens, *Linn. Sp. Pl.* 426.—Ind. occ.
urens, Macfad. Fl. Jamaic. 146 = oxycocca.
variifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 581.—Am. trop.?
verbascifolia, Linn. Sp. Pl. 426 = Byrsonima verbascifolia.
volubilis, Sims, Bot. Mag. t. 809 = Hiraea Simsiana.

MALPIGIA, P. Br. Hist. Jamaic. 229 (1756) = praec.

MALTEBRUNIA, Kunth, Rév. Gram. i. 183. t. 3 (1830) = **Potamophila**, R. Br. (Gramin.).
leersioides, Kunth, l. c. = P. leersioides.
prehensilis, Nees, Fl. Afr. Austr. i. 194 = P. prehensilis.

MALTEBURNIA, Steud. Nom. ed. II. ii. 93 (1841) sphalm. = praec.

MALTHEWSIA, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 93, 105 (1841) sphalm. = **Mathewsia**, Hook. & Arn. (Crucif.).

MALULUCBAN, Blanco, Fl. Filip. ed. I. 188 (1837) = **Champereia**, Griff. (Santal.).

MALUS, Tourn. ex Linn. Gen. ed. I. 145, in syn. (1737); Mill. Gard. Dict. ed. VI (1752) = **Pyrus**, Tourn. (Rosac.).
acerba, Mérat, Fl. Par. i. 187 = P. Malus.
angustifolia, Michx. Fl. Bor. Am. i. 292 = P. angustifolia.
apiosa, Hort. ex Steud. Nom. ed. II. ii. 93 = P. Malus.
astracanica, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 426 = P. Malus.
baccata, Borckh. Handb. Forstbot. ii. 1280; Moench, Meth. 682 = P. baccata.
baccata, Loisel. in Duham. ed. Nov. vi. t. 43. f. 2 = P. cerasifera.
betulifolia, Wenzig, in Eichl. Jahrb. Berl. ii. (1883) 292 = P. betulaefolia.
cerasifera, Spach, Hist. Vég. Phan. ii. 152 = P. baccata.
communis, Poir. Encyc. v. 560 = P. Malus.
coronaria, Mill. Gard. Dict. ed. VIII. n. 2 = P. coronaria.
csikloviensis, Kit. in Rochel. Pl. Banat. Rar. 26, nomen. (Quid?).—Banat.
dasyphylla, Borckh. Handb. Forstbot. ii. 1269 = P. Malus.
dioica, Moench, Meth. Suppl. 289; Borckh. l. c. 1276 = P. Malus.
diversifolia, M. Roem. Syn. Rosifl. 215 = P. rivularis.
domestica, Baumg. Enum. Stirp. Transs. ii. 43 = P. Malus.
domestica, Borckh. Handb. Forstbot. ii. 1272 = P. Sorbus?
floribunda, Siebold, ex Van Houtte, Fl. des Serres, xv. (1862–65) t. 1585 = P. spectabilis.
Fontanesiana, Spach, Hist. Vég. Phan. ii. 150 = P. Malus.
frutescens, Medic. Geschich. 78 = P. Malus.

MALUS:—
heterophylla, Spach, Hist. Vég. Phan. ii. 138 = Pyrus heterophylla, *Steud.*
hybrida, Desf. Hist. Arbr. ii. 141 = P. cerasifera.
japonica, Andr. Bot. Rep. t. 462 = P. japonica.
Kaido, Hort. ex C. Koch, Dendrol. i. 210 = P. spectabilis.
longipes, Wenzig, in Eichl. Jahrb. Berl. ii. (1883) 292 = P. longipes.
microcarpa, Rafin. ex DC. Prod. ii. 632 = Amelanchier canadensis.
nivalis, Desf. ex Steud. Nom. ed. II. ii. 93 = P. elaeagnifolia.
paradisiaca, Medic. Geschich. 78 = P. Malus.
Pashia, Wenzig, in Eichl. Jahrb. Berl. ii. (1883) 292 = P. Pashia.
praecox, Borckh. Handb. Forstbot. ii. 1271 = P. Malus.
prunifolia, Borkh. l. c. 1278 = P. prunifolia.
prunifolia, Steud. Nom. ed. II. ii. 93 = P. baccata.
pumila, Mill. Gard. Dict. ed. VIII. n. 3 = P. Malus.
quinqueflora, M. Roem. Syn. Rosifl. 216 = P. quinqueflora.
rivularis, M. Roem. l. c. 215 = P. rivularis.
rossica, Medic. Geschich. 78 = P. Malus.
rubicunda, M. Roem. Syn. Rosifl. 216 = P. rubicunda.
sativa, Dum.-Cours. Bot. Cult. ed. II. v. 426 = P. Malus.
sempervirens, Desf. Hist. Arbr. ii. 141 = P. angustifolia.
sibirica, Borckh. in Roem. Arch. i. iii. 89 = P. baccata.
Sieversii, M. Roem. Syn. Rosifl. 216 = P. Sieversii.
sinensis, Dum.-Cours. Bot. Cult. ed. II. v. 429 = P. spectabilis.
Sorbus, Borckh. in Roem. Arch. i. iii. 89 = P. Sorbus.
spectabilis, Borckh. Handb. Forstbot. ii. 1279 = P. spectabilis.
sphaerocarpa, M. Roem. Syn. Rosifl. 214 = P. sphaerocarpa.
suaveolens, M. Roem. l. c. 215 = P. suaveolens.
subcordata, M. Roem. l. c. 192 = P. rivularis.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 1 = P. Malus.
tatarica, Hort. ex C. Koch, Dendrol. i. 204 = P. Malus.
tomentosa, Dum.-Cours. Bot. Cult. ed. II. v. 438 = P. tomentosa.
Toringo, Hort. ex C. Koch, Dendrol. i. 210 = P. spectabilis.
Torringo, Carr. in Rev. Hortic. (1871) 451; (1872) 210 = P. Torringo.
umbellata, Gilib. Fl. Lituan. ii. 235 = P. Malus.
vulgaris, Pall. Reise, iii. 653 = P. communis?

MALVA, [Tourn.] Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 200.
BISMALVA, Medic. Malv. 39 (1787).
abulensis, Cav. Diss. 76 = M. Alcea.
abutiloides, Linn. Sp. Pl. ed. II. 971 = Sphaeralcea abutiloides.
abutiloides, Pursh, Fl. Am. Sept. ii. 454 = Kosteletzkya virginica?
abyssinica, *A. Br. in Flora*, xxiv. (1841) 1. 279.—Abyssin.
acaulis, Cav. Diss. ii. 82. t. 35. f. 2 = Malvastrum acaule.
acerifolia, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera acerifolia.
acerifolia, Nutt. ex Walp. Rep. i. 296 = Sphaeracea rivularis.
acutiloba, C. Koch, in Linnaea, xix. (1847) 58; Martr. Fl. Tarn, 120 = sylvestris.
adulterina, Wallr. Beitr. Fl. Hercyn. 241 (Quid?).—Europ.
aegyptia, *Linn. Sp. Pl.* 690.—Reg. Mediterr.; Oriens.
aegyptiaca, Steud. Nom. ed. II. ii. 93 = praec.
africana, Mill. ex Steud. l. c. ed. I. 506 = capensis.
albens, E. Mey. ex Harv. & Sond. Fl. Cap. i. 163 = Malvastrum albens.
albiflora, Presl, Delic. Prag. 29 = sylvestris.
albulensis, Steud. Nom. ed. I. 506, sphalm. = *abulensis*.
Alcea, *Linn. Sp. Pl.* 689.—Europ.
Alcea, Plan. Ens. Fl. Gallega, 135 = Tournefortiana.
alceoides, Tenore, Fl. Neap. Prod. Suppl. i. 62 (1815); Fl. Nap. ii. 109 = M. Alcea.
alchemillaefolia, Wall. Cat. n. 1884 D = verticillata.
althaeoides, *Cav. Ic.* ii. 30. t. 135.—Reg. Mediterr.
althaeoides, Sibth. & Sm. Fl. Graec. vii. t. 664 = cretica.
ambigua, *Guss. Fl. Sic. Prod.* ii. 331.—Europ. austr.
americana, Linn. Sp. Pl. 687 = Malvastrum tricuspidatum.

MALVA:—
americana, A. Rich. Ess. Fl. Cub. 115 = Malvastrum spicatum.
amoena, Drège, ex Harv. & Sond. Fl. Cap. i. 161 = Malvastrum grossulariaefolium.
amoena, Sims, Bot. Mag. t. 1998 = Malvastrum calycinum.
angustifolia, Cav. Diss. ii. 64. t. 29. f. 3 = Sphaeralcea angustifolia.
anomala, Eckl. & Zeyh. Enum. 290; et ex Harv. & Sond. Fl. Cap. i. 165 = Sphaeralcea elegans.
anomala, Link & Otto, Ic. Pl. Select. i. 51 = Lavatera tripartita.
antillarum, Zuccagni, in Roem. Collect. 147 = Malvastrum tricuspidatum.
arborea, A. St. Hil. Fl. Bras. Mer. i. 215. t. 134. f. 4 = Hibiscus arboreus.
arborea, Webb & Berth. Phyt. Canar. i. 30 = Lavatera arborea.
arvensis, Presl, Delic. Prag. 28 = nicaeënsis.
asperrima, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 164 = Malvastrum dissectum.
asperrima, Jacq. Hort. Schoenb. ii. 7 = Malvastrum asperrimum.
asterocarpa, Steud. Nom. ed. II. ii. 94; et in Flora, xxxix. (1856) 436 (Malvastri sp.).—Chili.
astrolasia, Zipp. ex Span. in Linnaea, xv. (1841) 168 = Malvastrum spicatum.
atropurpurea, *Noé, ex Alef. in Oestr. Bot. Zeitschr.* xii. (1862) 261.—Hab.?
aurantiaca, Scheele, in Linnaea, xxi. (1848) 469 = Malvastrum Wrightii.
balsamica, Jacq. Misc. ii. 321 = Malvastrum capense.
Behriana, Schlecht. in Linnaea, xx. (1847) 633 = Lavatera plebeia.
Belloa, C. Gay, Fl. Chil. i. 304. t. 7 = Malvastrum Belloa.
Berteroniana, *Steud. Nom.* ed. II. ii. 94; *et in Flora*, xxxix. (1856) 437.—Chili.
betulina, Desr. in Lam. Encyc. iii. 754 = Malvastrum spicatum.
betuloides, Schrad. in Linnaea, xi. (1837) 88 = Malvastrum ruderale.
biflora, Desr. in Lam. Encyc. iii. 747 = Malvastrum capense.
Bismalva, Bernh. ex Lej. Rev. Fl. Spa, 147 = M. Alcea.
Bivoniana, *Presl, Fl. Sicil.* i. 176.—Sicilia.
Blumeana, Steud. Nom. ed. II. ii. 94 = Malvastrum spicatum.
bonariensis, Cav. Diss. ii. 69. t. 22. f. 1 = Sphaeralcea bonariensis.
borbonica, Willd. Enum. Hort. Berol. 728 = Malvastrum tricuspidatum.
borealis, *Wallm. in Liljebl. Svensk. Fl.* ed. III. 374. —Europ.; As. bor.
Boryana, *DC. Prod.* i. 435.—Ins. Borbon.
brachystachya, F. Muell. in Linnaea, xxv. (1852) 378 = Malvastrum spicatum.
bracteata, Reichb. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 186 = borealis.
brasiliensis, *Desr. in Lam. Encyc.* iii. 744.—Bras.
brevifolia, Gilib. Fl. Lituan. i. 64 = crispa.
brevipes, *Phil. in Anal. Univ. Chil.* (1870) ii. 163.—Chili.
bryoniaefolia, Drège, ex Harv. & Sond. Fl. Cap. i. 161 = Malvastrum asperrimum.
bryonifolia, Drège, l. c. = Malvastrum grossulariaefolium.
bryonifolia, Linn. Sp. Pl. 688 = Malvastrum bryonifolium?
californica, Presl, Rel. Haenk. ii. 121 = Sida hederacea.
calycina, Cav. Diss. ii. 81. t. 22. f. 4 = Malvastrum calycinum.
campanulata, Paxt. Mag. Bot. ix. (1843) 173 = Malvastrum campanulatum.
campanuloides, *Lodd. Bot. Cab.* t. 1670.—Am. bor.
cannabina, Serres, in Bull. Soc. Bot. Fr. iii. (1856) 276 = M. Alcea.
capensis, Linn. Sp. Pl. 688; Cav. Diss. ii. 70. t. 18. f. 3 = Malvastrum capense.
capitata, Cav. Diss. v. 280. t. 137. f. 1 = Malvastrum capitatum.
caroliniana, Linn. Sp. Pl. 688 = Modiola multifida.
carpinifolia, Desr. in Lam. Encyc. iii. 754 = Malvastrum tricuspidatum.
caschmireana, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera cachemiriana.

MALVA:—
chilensis, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1857) 1.—Chili.
chinensis, Mill. Gard. Dict. ed. VIII. n. 6 = verticillata.
ciliata, Wall. Cat. n. 1878 = sylvestris.
circinnata, Vis. Prod. Fl. Cors. App. 6 = nicaeënsis.
cisplatina, Steud. Nom. ed. II. ii. 94 = Sphaeralcea cisplatina.
coccinea, Nutt. Gen. Am. ii. 81 = Malvastrum coccineum.
cognata, *Steud. in Flora*, xxxix. (1856) 425.—Chili.
Colmeiroi, Willk. in Linnaea, xxx. (1859–60) 98 = Tournefortiana.
Columnae, Tenore, ex Steud. Nom. ed. II. ii. 94 (Quid?).—Ital.
concinna, *Knowles & Westc. Flor. Cab.* i. (1837) 79.—Am. austr.
corchorifolia, Desr. in Lam. Encyc. iii. 743 = Malvastrum Rugelii.
cordistipula, *Steud. Nom.* ed. II. ii. 94.—Chili.
coromandelina, Linn. Sp. Pl. 687 = Malvastrum tricuspidatum.
coronata, *Pomel, Nouv. Mat. Fl. Atl.* 346.—Alger.
corymbosa, E. Mey. ex Harv. & Sond. Fl. Cap. i. 161 = Malvastrum asperrimum?
costata, Presl, Rel. Haenk. ii. 123 = Malvastrum peruvianum?
Creeana, R. Grah. in Hook. Bot. Mag. t. 3698 = Malvastrum coccineum, *vel* Sphaeralcea pedata.
crenata, Kit. in Sadl. Fl. Pest. ed. I. ii. 161 = borealis.
cretica, *Cav. Diss.* v. 280. t. 138. f. 2.—Ins. Creta.
cretica, Tenore, Fl. Nap. ii. 106 = althaeoides.
crispa, *Linn. Syst.* ed. X. 1147.—Europ.
cuneifolia, Cav. Diss. ii. 64. t. 20. f. 1 = hispanica.
curassavica, Desr. in Lam. Encyc. iii. 754 = Malvastrum tricuspidatum.
cymbalariaefolia, DC. Prod. i. 431 = seq.
cymbalarifolia, Desr. in Lam. Encyc. iii. 753 = Sida Sherardiana.
Cyrilli, Vis. Fl. Dalm. iii. 207 = Lavatera thuringiaca.
debilis, Salisb. Prod. 380 = Malvastrum capense.
decumbens, Host, Fl. Austr. ii. 298 = M. Alcea.
decumbens, Willd. Enum. Hort. Berol. 731 = Modiola multifida.
deflexa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 186 = Malvastrum grossulariaefolium.
Dethardingii, Link, ex Dethard. Fl. Megalop. 54 = moschata.
digitata, Torr. & Gray, Fl. N. Am. i. 246 = Callirhoë digitata.
Dilleniana, Eckl. & Zeyh. Enum. 36 = Malvastrum virgatum.
diphylla, Moench, Meth. 611 = aegyptia.
divaricata, Andr. Bot. Rep. t. 182 = Malvastrum divaricatum.
divaricata, Hort. ex Steud. Nom. ed. II. ii. 94 = Malvastrum tridactylites.
divaricata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 165 (Sphaeralceae sp.).—Afr. austr.
domingensis, Spreng. ex DC. Prod. i. 431 = Malvastrum tricuspidatum.
Durieui, Spach, in Linnaea, xxiv. (1851) 233 = Lavatera mauritanica.
echinata, Presl, Rel. Haenk. ii. 122 (Malvastri sp.).—Peruv.
elata, *Pomel, Nouv. Mat. Fl. Atl.* 345.—Alger.
elata, Salisb. Prod. 381 = sylvestris.
elegans, Cav. Diss. ii. 59. t. 16. f. 1 = Sphaeralcea elegans.
equina, Wallr. Beitr. Fl. Hercyn. 241 = sylvestris.
erecta, Gilib. Fl. Lituan. i. 64; Presl, Delic. Prag. 30 = sylvestris.
eriocalyx, *Steud. Nom.* ed. II. ii. 94.—Chili.
eriocarpa, DC. Prod. i. 436 = Modiola multifida.
erodiifolia, Presl, Rel. Haenk. ii. 122 (Malvastri sp.).—Chili.
europaea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = Malvastrum tricuspidatum.
exasperata, Moench, Meth. 610 = Malvastrum peruvianum.
excelsa, Presl, Delic. Prag. 27 = nicaeënsis.
excisa, Reichb. Ic. Fl. Germ. v. t. 170 = M. Alcea.
fasciculata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 225 = Malvastrum Thurberi.
fastigiata, Cav. Diss. ii. 75. t. 23. f. 2 = M. Alcea.
fastuosa, Salisb. Prod. 381 = Lavatera arborea.
flava, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera flava.

MALVA:—
flexuosa, Gill. ex Hook. Bot. Misc. iii. (1833) 151 = Sphaeralcea obtusiloba.
flexuosa, Hornem. Hort. Hafn. ii. 655 = parviflora.
floribunda, *Steud. Nom.* ed. II. ii. 94.—Mexic.
florida, Salisb. Prod. 381 = Althaea rosea.
fluminensis, Vell. Fl. Flum. 287; vii. t. 48 = Malvastrum spicatum.
fragrans, Jacq. Hort. Vindob. iii. 20. t. 35 = Malvastrum capense.
fruticans, Dorn. ex Reichb. Ic. Fl. Germ. v. 18 = rotundifolia.
fruticosa, Berg. Desc. Pl. Cap. 181 = Malvastrum tridactylites.
gangetica, Linn. Sp. Pl. ed. II. 967 = Malvastrum tricuspidatum.
geraniifolia, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 125 = moschata.
geranioides, Cham. & Schlecht. in Linnaea, v. (1830) 226 = Malvastrum geranioides.
geranioides, Gill. ex Hook. Bot. Misc. iii. (1833) 152 = Modiola geranioides.
Gilliesii, *Phil. in Anal. Univ. Chil.* (1870) II. 16.—Chili.
Gilliesii, Steud. Nom. ed. II. ii. 94 = Modiola geranioides.
glabra, Desr. in Lam. Encyc. iii. 752 = sylvestris.
glabrifolia, Tenore, ex Steud. Nom. ed. II. ii. 94 (Quid?).—Hab.?
glomerata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 151 = Malvastrum glomeratum.
glomerata, Hort. ex G. Don, in Loud. Hort. Brit. 286 = verticillata.
glutinosa, Desr. in Lam. Encyc. iii. 747 = Malvastrum calycinum, grossulariaefolium.
grandiflora, *C. Morr. in Ann. Soc. Bot. Gand*, iii. (1847) 453. t. 167.—Am. bor.
grossulariaefolia, Cav. Diss. ii. 71 = Malvastrum grossulariaefolium.
grossulariaefolia, Drège, ex Presl, Bot. Bemerk. 18 = hispida.
grossulariaefolia, E. Mey. ex Harv. & Sond. El. Cap. i. 163 = Malvastrum albens.
gymnocarpa, *Pomel, Nouv. Mat. Fl. Atl.* 345.—Alger.
Haenkeana, Presl, Reliq. Haenk. ii. 123 = Malvastrum peruvianum?
hederacea, Dougl. in Hook. Fl. Bor. Am. i. 107 = Sida hederacea.
hederaefolia, Vis. Fl. Dalm. iii. 205 = Lavatera cretica.
Henningii, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 133 = borealis.
Hermanniae, *Bubani, ex Timb. in Bull. Soc. Dauph.* [ii?] 39; *ex eod. in Mém. Acad. Toul.* Sér. VIII. i. II. (1877) 191.—Hab.?
hibiscoides, *Desr. in Lam. Encyc.* iii. 748.—Ins. Borbon.
hirsuta, Aubl. Pl. Gui. ii. 704 = Sida linifolia.
hirsuta, Tenore, Prod. Fl. Nap. 40 = cretica.
hirsuta, Viv. Fl. Cors. Diagn. 12; Presl, Fl. Sic. 175 = sylvestris.
hispanica, Asso, Syn. Pl. Arag. 90 = stipulacea.
hispanica, *Linn. Sp. Pl.* 689.—Reg. Mediterr. occ.
hispida, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera Olbia.
hispida, Moench, Meth. 609 = Malvastrum tricuspidatum.
hispida, *Presl, Bot. Bemerk.* 18.—Afr. austr.
Hornschuchiana, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 303 (Sphaeralceae sp.).—Peruv.
horrida, *Span. in Linnaea*, xv. (1841) 168.—Ins. Timor.
hortensis, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 886 = Althaea rosea.
Houghtonii, Torr. & Gray, Fl. N. Am. i. 225 = Callirhoë triangulata.
humifusa, Henning, in Mém. Soc. Nat. Mosc. vi. (1823) 76 = borealis.
humifusa, Salisb. Prod. 380 = Sida Sherardiana.
humilis, Gill. in Hook. Bot. Misc. iii. (1833) 150 = Malvastrum humile.
hybrida, Čelak. in Arch. Naturf. Landesd. Boehm. iii. (1875) 515 = rotundifolia.
hybrida, Hort. ex Steud. Nom. ed II. ii. 94 = verticillata.
incana, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 419.—Hab.?
incana, *Presl, Rel. Haenk.* ii. 121.—Chili.
intermedia, Bor. Fl. Centr. Fr. ed. II. 98 = M. Alcea.

MALVA:—
involucrata, Torr. & Gray, Fl. N. Am. i. 226 = Callirhoë involucrata.
italica, Pollini, Hort. & Prov. Veron. Pl. 17; ex ej. Fl. Veron. ii. 438 = M. Alcea.
Julii, Spreng. Syst. iii. 89 (= *Sphaeroma Julii*).—Afr. austr.
juvenalis, Lespinasse, in Bull. Soc. Bot. Fr. vi. (1859) 647, in nota = Modiola erecta.
laciniata, Desr. in Lam. Encyc. iii. 750 = moschata.
lactea, [Soland. in] Ait. Hort. Kew. ed. I. ii. 448 = Malvastrum vitifolium.
laeta, Phil. in Anal. Univ. Chil. (1862), 442 = *tenuifolia*, Hook. & Arn.
laevis, Moench, Meth. 610 = Malvastrum limense.
Lagascae, Hort. ex DC. Prod. i. 430 = Malvastrum scabrum.
Lagascae, *Lás. & Turb. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 416.—Hispan.
Lamottei, Jord. ex Nym. Consp. 129 = moschata.
lasiocarpa, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 45 = Malvastrum lasiocarpum.
lateritia, Hook. Bot. Mag. t. 3846 = Malvastrum lateritium.
lateritia, *Regel, in Bot. Zeit.* ix. (1851) 891.—Hab.?
Lecontii, Buckl. in Am. Journ. Sc. xlv. (1843) 176 = Pavonia hastata.
leprosa, Orteg. Hort. Matr. Dec. 95. (Sidae sp.).—Cuba.
libyca, *Pomel, Nouv. Mat. Fl. Atl.* 344.—Alger.
limensis, Linn. Cent. Pl. ii. 27 = Malvastrum limense.
Lindheimeriana, Scheele, in Linnaea, xxi. (1848) 470 = Malvastrum tricuspidatum.
lineariloba, Young, Fl. Texas, 180; ex S. Wats. Bibliog. Ind. N. Am. Bot. 133 = Callirhoë involucrata.
litoralis, Dethard. ex Reichb. Fl. Germ. Excurs. 771 = rotundifolia.
lobata, Cav. Diss. ii. 76 = M. Alcea.
lucida, Lindl. Bot. Reg. (1839) Misc. 82 = sylvestris.
lutea, *Roem. Hort. Turic. ex Steud. Nom.* ed. II. ii. 94.—Hab.?
luzonica, Blanco, Fl. Filip. ed. II. 385 = Malvastrum tricuspidatum.
macrocalyx, Presl, Bot. Bemerk. 19 = Sphaeralcea elegans.
macropoda, *Steud. in Flora*, xxxix. (1856) 425.—Chili.
macrostachya, Presl, Rel. Haenk. ii. 120 = Malvastrum macrostachyum.
malachroides, Hook. & Arn. Bot. Beech. Voy. 326 = Sidalcea malachroides.
malwensis, Edgew. in Hook. Journ. Bot. ii. (1840) 284 = Althaea Ludwigii.
mamillosa, Lloyd, ex Nym. Consp. 128 = Lavatera cretica.
mareotica, Delile, ex DC. Prod. i. 433 = parviflora.
maritima, Lam. Fl. Fr. iii. 140 = Tournefortiana.
maritima, Salisb. Prod. 381 = Althaea officinalis.
Martensiana, *E. Morr. in Belg. Hortic.* xiv. (1864) 165.—Austral.
Mathewsii, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 563 = Malvastrum peruvianum.
mauritanica, Spreng. Syst. iii. 90 = seq.
mauritiana, Linn. Sp. Pl. 689 = sylvestris.
mauritiana, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 581 = Lavatera cretica.
mendocina, Phil. Sert. Mendoc. Alt. 5 = Sphaeralcea mendocina.
mexicana, S. Schau. in Linnaea, xx. (1847) 724 = Malvastrum mexicanum.
micans, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera micans.
microcalyx, Hort. ex C. A. Mey. in Ind. Sem. Hort. Petrop. xi. 75 = parviflora.
microcarpa, Desf. Tabl. ed. I. 144; et ex DC. Prod. i. 433 = parviflora.
microphylla, E. Mey. ex Harv. & Sond. Fl. Cap. i. 163 = Malvastrum divaricatum.
miniata, Cav. Ic. iii. 40 = Sphaeralcea miniata.
miniata, *Jacq. Fragm.* 83.—Hab.?
minoricensis, *Rodrig. in Anal. Soc. Esp. Hist. Nat.* iii. (1874) 13.—Ins. Balear.
mohileviensis, *Down. in Bull. Soc. Nat. Mosc.* xxxiv. (1861) i. 177.—Rossia.
Monroana, Otto & Dietr. Allg. Gartenz. v. (1837) 104 = Malvastrum Munroanum.
montana, Forsk. Fl. Aegypt. Arab. 124 = nicaeënsis.
Moreniana, Pollini, Fl. Veron. ii. 437, in syn. = M. Alcea.

MALVA:—
Morenii, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 34 = M. Alcea.
Morisonii, Pollini, ex Schur, Enum. Pl. Transs. 130 = M. Alcea.
moschata, Blanco, Fl. Filip. ed. I. 51 = Pelargonium odoratissimum.
moschata, *Linn. Sp. Pl.* 690.—Europ.
Mulleri, Mull. ex Parl. Fl. Ital. v. 79 = Lavatera triloba.
Munroana, Dougl. ex Lindl. Bot. Reg. t. 1306 = Malvastrum Munroanum.
neglecta, Wallr. in Syll. Ratisb. i. (1824) 140 = rotundifolia.
neilgherrensis, Wight, Ic. t. 950 = verticillata.
nepalensis, Hort. ex Steud. Nom. ed. II. ii. 95 = verticillata.
nicaeënsis, *All. Fl. Pedem.* ii. 40.—Reg. Mediterr.
nodosa, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 39; *et in Oken, Isis*, xxi. (1828) 991.—Thracia.
nubigena, Wedd. Chlor. And. ii. 276 (= Malvastri sp.).—Bolivia.
nuttalloides, Croom, in Am. Journ. Sc. xxvi. (1834) 313 = Callirhoë Papaver.
obliqua, Nutt. ex A. Gray, in Mem. Am. Ac. N. S. iv. (1849) 23 = Sida hederacea.
obtusa, Moench, Meth. 610 = sylvestris.
obtusa, Torr. & Gray, Fl. N. Am. i. 225 = borealis.
obtusifolia, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 303.—Chili.
obtusiloba, Hook. Bot. Mag. t. 2787 = Sphaeralcea obtusiloba.
odorata, [Westc.] in Maund, Botanist, v. t. 218 = Malvastrum capense.
officinalis, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 885 = Althaea officinalis.
Olbia, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera Olbia.
operculata, Cav. Diss. ii. 65. t. 35. f. 1 (= Malvastri sp.).—Peruv.
orbiculata, *Pomel, Nouv. Mat. Fl. Atl.* 344.—Alger.
Oriastrum, Wedd. Chlor. And. ii. 277 (= Malvastri sp.).—Bolivia.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 3.—Oriens.
Orsiniana, Tenore, Fl. Nap. v. 85 = moschata.
ovata, Cav. Diss. ii. 81. t. 20. f. 2 = Malvastrum spicatum.
ovata, Dulac, Fl. Hautes-Pyr. 232 = M. Alcea.
oxyacanthoides, Eckl. & Zeyh. ex Harv. et Sond. Fl. Cap. i. 163 = Malvastrum divaricatum.
oxyacanthoides, Hornem. Hort. Hafn. ii. 654 = Malvastrum tridactylites.
oxyloba, *Boiss, Diagn.* Ser. I. viii. 109.—Palaestina.
pallida, Salisb. Prod. 381 = Lavateria Olbia.
Papaver, Cav. Diss. ii. 64. t. 15. f. 3 = Callirhoë Papaver.
papaveriformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = praec.
parnassiaefolia, Wedd. Chlor. And. ii. 275 = Malvastrum parnassiaefolium.
parviflora, Huds. Fl. Angl. ed. I. 268 = borealis.
parviflora, *Linn. Diss. Dem. Pl. Nov.*; *Amoen. Acad.* iii. 416.—Europ.
patagonica, *Niederl. in Roca, Exped. Rio-Negro*, ii. 196.—Patagon.
pedata, Torr. & Gray, Fl. N. Am. i. 227 = Callirhoë alcaeoides, digitata.
peduncularis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 150.—Chili.
pedunculata, *Rafin. Caratteri*, 76.—Sicil.
perpusilla, Nutt. ex A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 22 = Malvastrum angustum?
peruviana, Linn. Sp. Pl. 688 = Malvastrum peruvianum.
phoenicea, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera phoenicea.
pinnatipartita, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 45.—Bras.
plebeja, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 325 = sylvestris.
plicata, Nutt. in Torr. & Gray, Fl. N. Am. i. 227 = Sida hederacea.
plicata, *Thunb. Prod. Pl. Cap.* 119.—Afr. austr.
plumosa, Presl, Rel. Haenk. ii. 124 = Malvastrum plumosum.
polyantha, *Steud. Nom.* ed. II. ii. 95.—Chili.
polymorpha, Guss. Fl. Sic. Syn. ii. 226 = sylvestris.
polystachya, Cav. Diss. v. 280. t. 638. f. 3 = Malvastrum spicatum.
praemorsa, Hoffmgg. Verz. Pfl. 76 = Malvastrum tridactylites.

MALVA:—
Preissiana, Miq. in Lehm. Pl. Preiss. i. 238 = Lavatera plebeia.
propinqua, Salisb. Prod. 380 = Malvastrum grossulariaefolium.
prostrata, Cav. Diss. ii. 59. t. 16. f. 3 = Modiola multifida.
prostrata, Gilib. Fl. Lituan. i. 63 = rotundifolia.
prostrata, Phil. in Anal. Univ. Chil. (1870) ii. 163 = Sphaeralcea bonariensis.
Pseudo-borealis, Schur, Enum. Pl. Transs. 130 = borealis.
Pseudo-lavatera, Webb & Berth. Phyt. Canar. i. 29 = Lavatera cretica.
Pseudo-olbia, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = L. Olbia.
ptarmicaefolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 46.—Bras.
pulchella, Bernh. Sel. Sem. Hort. Erfurt. (1832) n. 8 = verticillata.
pulchra, *Boj. Hort. Maurit.* 25.—Hab.?
pumila, *Retz. Obs.* iii. 37.—Hab.?
punctata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera punctata.
Purdiaei, A. Gray, in Wedd. Chlor. And. ii. 275 = Malvastrum Purdiaei.
Purdiaei, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 154.—N. Granat.
purpurata, Lindl. Bot. Reg. t. 1362 = Malvastrum campanulatum.
purpurea, C. Gay, Fl. Chil. iii. 305 = praec.
pusilla, Sm. Engl. Bot. iv. t. 241 = borealis.
pygmaea, Wedd. Chlor. And. ii. 274 = Malvastrum pygmaeum.
racemosa, E. Mey. ex Harv. & Sond. Fl. Cap. i. 164 = Malvastrum racemosum.
racemosa, Presl, Delic. Prag. 29 = sylvestris.
recta, Opiz, Boehm. Gew. (1823) 83 = sylvestris.
reflexa, Andr. Bot. Rep. t. 135 = Malvastrum tridactylites.
reflexa, *Wendl. Bot. Beobacht.* 53; *Hort. Herrenh.* 6. tab. 4.—Hab.?
repens, Gueldenst. Reisen, i. 52 = borealis.
retusa, Cav. Diss. ii. 93. t. 21. f. 1 = Malvastrum trilobum.
retusa, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 160 = Malvastrum calycinum.
rhizantha, Wedd. Chlor. And. ii. 276 = Malvastrum rhizanthum.
ribifolia, Schlecht. in Linnaea, xi. (1837) 351 = Malvastrum ribifolium.
ribifolia, Viv. Fl. Cors. Prod. App. 85 = ambigua.
Richii, Wedd. Chlor. And. ii. 276 = Malvastrum Richii.
rivularis, Dougl. ex Hook. Fl. Bor. Am. i. 107 = Sphaeralcea rivularis.
Rodigii, *Presl, Delic. Prag.* 28.—Europ.
rosea, [Moç. & Sesse,] ex DC. Prod. i. 435 = Sphaeralcea umbellata.
rosea, Roxb. ex Wight & Arn. Prod. 45 = borealis.
rotundifolia, Cav. Diss. ii. 79. t. 26. f. 3 = sylvestris.
rotundifolia, Gorter, Fl. Ingr. 110 = borealis.
rotundifolia, *Linn. Sp. Pl.* 688.—Europ.; As. bor.
ruderalis, Blume, Bijdr. 64 = Malvastrum tricuspidatum.
ruderalis, Salisb. Prod. 381 = sylvestris.
rugosa, Desr. in Lam. Encyc. iii. 747 (Malvastri sp.).—Afr. austr.
rugosa, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 165 = Sphaeralcea elegans.
scaberrima, *Lag. ex Steud. Nom.* ed. II. ii. 95 (Quid?).—Hispan.
salvitellensis, Briganti, Stirp. Rar. Pempt. i. 1. t. 1; ex Tenore, Syll. 337 = rotundifolia.
scabra, Cav. Diss. v. 281. t. 138. f. 1 = Malvastrum scabrum.
scabrosa, Linn. Cent. Pl. ii. 27; Amoen. Acad. iv. 325 = Malvastrum capense.
scandens, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
Schiedeana, Steud. Nom. ed. II. ii. 95 = Malvastrum ribifolium.
scoparia, Jacq. Coll. i. 59 = Malvastrum Rugelii.
scoparia, L'Hérit. Srirp. Nov. 53; Domb. ex Cav. Diss. 65. t. 21. f. 4 = Malvastrum scoparium.
scorpioides, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 562 = Malvastrum peruvianum.
setigera, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 886 = Althaea hirsuta.
setosa, Moench, Meth. Suppl. 199 = nicaeënsis.
Sherardiana, Linn. Sp. Pl. ed. II. 1675 = Sida Sherardiana.

MALVA:—
simpliciuscula, *Steud. in Flora*, xxxix. (1856) 426.—Chili.
sinensis, Cav. Diss. ii. 77. t. 25. f. 4 = sylvestris.
spicata, Linn. Syst. ed. X. 1146 = Malvastrum spicatum.
spithamea, Cav. Diss. ii. 61. t. 18. f. 3 = hispanica.
stellata, D. Dietr. Syn. Pl. iv. 816 = Sphaeralcea angustifolia.
stellata, Thunb. Prod. Pl. Cap. 119 = Malvastrum asperrimum.
stenophylla, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 156.—Hab.?
stipulacea, *Cav. Diss.* ii. 62. t. 15. f. 2.—Hispan.
striata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 165 = Sphaeralcea elegans.
stricta, Eckl. & Zeyh. ex Harv. & Sond. l. c. 162 = Malvastrum tridactylites.
stricta, Jacq. Hort. Schoenb. iii. t. 294 = Malvastrum strictum.
subhastata, Cav. Diss. ii. 72. t. 21. f. 3 = Malvastrum tricuspidatum.
sublobata, Desr. in Lam. Encyc. iii. 754 = Malvastrum spicatum.
subtriflora, Lag. Gen. et Sp. Nov. 21 = Malvastrum subtriflorum.
sulphurea, Gill. in Hook. Bot. Misc. iii. (1833) 149 = Sida sulphurea.
sylvestris, *Linn. Sp. Pl.* 689.—Europ.; As. temp.
tenella, Cav. Ic. v. 14. t. 422. f. 3 = Malvastrum tenellum.
tenuicalyx, *Tenore, Ind. Sem. Hort. Neap.* (1857).—Hab.?
tenuifolia, Desr. in Lam. Encyc. iii. 750 = Tournefortiana.
tenuifolia, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 150 (Malvastri sp.).—Chili.
tenuifolia, Savi, Due Cent. 122 = moschata.
Thirkeana, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera thuringiaca.
thuringiaca, *Vis. Fl. Dalm.* iii. 207.—Europ.
timoriensis, DC. Prod. i. 430; Blume, Bijdr. 64 = Malvastrum spicatum.
tomentella, Presl, Fl. Sicul. 174 = sylvestris.
tomentosa, *Linn. Sp. Pl.* 687.—Malaya.
Tournefortiana, Asso, ex Losc. & Pard. Ser. Imperf. ed. II. 75 = stipulacea.
Tournefortiana, *Linn. Cent. Pl.* i. 21; *Amoen. Acad.* iv. 283.—Europ.
Tournefortii, Timb. in Bull. Soc. Dauph. [ii.] 39; ex eod. in Mém. Acad. Toul. Sér. VIII. i. ii. (1877) 191 = praec.?
Toxicodendrum, Hort. ex Steud. Nom. ed. II. ii. 95 = Malvastrum tridactylites.
trachelifolia, *Link, Enum. Hort. Berol.* ii. 209.—Hab.?
triangulata, Leavenw. in Am. Journ. Sc. vii. (1824) 62 = Callirhoë triangulata.
tricuspidata, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 210 = Malvastrum tricuspidatum.
tridactylites, Cav. Diss. ii. 77. t. 21. f. 2 = Malvastrum tridactylites.
trifida, *Cav. l. c.* v. 280. t. 137. f. 2.—Hispan.
triloba, Cav. in Anal. Cienc. Nat. vi. (1803) 338 = praec.?
triloba, Thunb. Prod. Pl. Cap. 118 = Malvastrum trilobum.
trimestris, Salisb. Prod. 381 = Lavatera trimestris.
trionoides, DC. Prod. i. 433 = parviflora.
truncata, *Opiz, Boehm. Gew.* (1823) 83, nomen.—Europ.
tuberculata, *Link, ex Steud. Nom.* ed. II. ii. 95.—Hab.?
ulmifolia, Balb. ex DC. Prod. i. 430 = Malvastrum tricuspidatum.
umbellata, Bert. in Merc. Chil. [1829?] 695; ex C. Gay, Fl. Chil. i. 289 = Lavatera arborea.
umbellata, Cav. Ic. i. 64. t. 95 = Sphaeralcea vitifolia.
unguiculata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = Lavatera unguiculata.
urticaefolia, H. B. & K. Nov. Gen. et Sp. v. 276 = Modiola multifida.
velutina, *J. A. Schmidt, Beitr. Fl. Ins. Cap. Verd.* 282.—Ins. Cap. Virid.
veneta, Hort. ex DC. Prod. i. 432 = sylvestris.
venosa, Thunb. Prod. Pl. Cap. 119 = Sphaeralcea elegans?
verticillata, *Linn. Sp. Pl.* 689.—Europ.; As. et Afr. bor.

MALVA:—
villosa, *Thunb. Prod. Pl. Cap.* 118.—Afr. austr.
violacea, Phil. Sert. Mendoc. Alt. 4 = Malvastrum violaceum.
virgata, Cav. Diss. ii. 70. t. 18. f. 2 = Malvastrum virgatum.
virgata, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 163 = Malvastrum albens.
virgata, *Murr. in Comm. Gotting.* ii. 1779 (1780) n. 1. 20. t. 6.—Afr. austr.
viscosa, Salisb. Prod. 381 = Malvastrum capense.
vitifolia, Cav. Ic. i. t. 30 = Malvastrum vitifolium.
vitifolia, *Cervant. ex Steud. Nom.* ed. II. ii. 95 (Quid?).—Mexic.
vulgaris, Fries, Novit. Fl. Suec. 219 = rotundifolia.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 638 = sylvestris.
waltherifolia, Link, Enum. Hort. Berol. ii. 209 = tomentosa.
Willkommiana, Scheele, in Linnaea, xxi. (1848) 570 = Lavatera cretica.
zebrina, Hort. Turic.—Cf. Linnaea, xxiv. (1851) 175 = sylvestris.

MALVASTRUM, A. Gray, in Mém. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 21. *MALVACEAE*, Benth. & Hook. f. i. 201.
NOTOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1863) 1. 567.
acaule, *A. Gray, Bot. U. St. Expl. Exped.* i. 150.—Am. austr.
albens, *Harv. in Harv. & Sond. Fl. Cap.* i. 163.—Afr. austr.
americanum, Torr. Bot. Mex. Bound. 38 = tricuspidatum.
amoenum, Garcke, in Bonplandia, v. (1857) 296 = calycinum.
angustifolium, Hemsl. Biol. Centr. Am. Bot. i. 99 = Sphaeralcea angustifolia.
angustum, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 22.—Am. bor.
anthemidifolium, *A. Gray, Bot. U. St. Expl. Exped.* i. 152.—Bolivia.
aretioides, *A. Gray, l. c.* 153.—Peruv.
asperrimum, *Garcke, in Bonplandia*, v. (1857) 296; *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 161.—Afr. austr.
aurantiacum, Walp. Ann. ii. 153 = Wrightii.
Belloa, *A. Gray, in Bot. U. St. Expl. Exped.* i. 150, in adnot.—Chili.
borussicum, *Wedd. Chlor. And.* ii. 278.—Peruv.
bryonifolium, *Garcke, in Bonplandia*, v. (1857) 296; *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 161.—Afr. austr.
calycinum, *Garcke, l. c.*; *A. Gray & Harv. l. c.* 160.—Afr. austr.
campanulatum, *Nichols. Dict. Gard.* ii. 319.—Chili.
capense, *Garcke, in Bonplandia*, v. (1857) 296; *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 160.—Afr. austr.
capitatum, *Griseb. in Goett. Abh.* xix. (1874) 90.—Reg. Argent.
carpinifolium, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 22 = tricuspidatum.
Castelnaeanum, *Wedd. Chlor. And.* ii. 283—Peruv.
Cavanillesii, *A. Gray, Bot. U. St. Expl. Exped.* i. 156.—Peruv.
coccineum, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 24.—Am. bor. occ.
compactum, *A. Gray, Bot. U. St. Expl. Exped.* i. 152.—Chili.
coromandelianum, Regel, Ind. Sem. Hort. Petrop. (1861) 51 = seq.
coromandelinum, Garcke, in Bonplandia, v. (1857) 297 = tricuspidatum.
Coulteri, S. Wats. in Proc. Am. Acad. xi. (1876) 125 (Sphaeralceae sp.).—Calif.
densiflorum, *S. Wats. l. c.* xvii. (1881–82) 368.—Calif.
dissectum, *Harv. in Harv. & Sond. Fl. Cap.* i. 164.—Afr. austr.
divaricatum, *Garcke, in Bonplandia*, v. (1857) 296; *A. Gray & Harv. in Harv. & Sond. l. c.* 163.—Afr. austr.
exile, *A. Gray, in Ives, Rep. Colorado River, Bot.* 8.—Am. bor. occ.
flabellatum, *Wedd. Chlor. And.* ii. 281.—Bolivia.
foliosum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 356.—Calif.

MALVASTRUM:—
fragrans, A. Gray & Harv. in Harv. & Sond.. Fl. Cap. i. 159 = capense.
Fremontii, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 21 = Sphaeralcea Lindheimeri.
geranioides, *Hemsl. Biol. Centr. Am. Bot.* i. 99.—Mexic.
Gilliesii, *Baker, in Gard. Chron.* (1885) II. 166.—Parana.
glomeratum, *Griseb. in Goett. Abh.* xxiv. (1879) 42.—Reg. Argent.
grossulariaefolium, *Garcke, in Bonplandia*, v. (1857) 294; *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 162.—Afr. austr.
grossulariaefolium, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 21 = Sphaeralcea pedata.
heterophyllum, Griseb. in Goett. Abh. xxiv. (1879) 43 (Sphaeralceae sp.).—Reg. Argent.
humile, *A. Gray, Bot. U. St. Expl. Exped.* i. 150.—Chili.
lasiocarpum, *Griseb. in Goett. Abh.* xxiv. (1879) 43.—Reg. Argent.
lateritium, *Nichols. Dict. Gard.* ii. 319.—Uruguay.
leptophyllum, *A. Gray, Pl. Wright.* i. 17.—N. Mexic.
limense, *Ball, in Journ. Linn. Soc.* xxii. (1885) 32.—Peruv.
Lindheimerianum, *Walp. Ann.* ii. 753.—Texas.
linearifolium, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 449 = Sida ciliaris.
linoides, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 336.—Reg. Argent.
lobulatum, Wedd. Chlor. And. ii. 275 = parnassifolium.
longirostre, *Wedd. l. c.* 281.—Bolivia.
Macleani, *A. Gray, Bot. U. St. Expl. Exped.* i. 152.—Peruv.
macrostachyum, *Hemsl. Biol. Centr. Am. Bot.* i. 99.—Mexic.
Mandonianum, *Wedd. Chlor. And.* ii. 282.—Bolivia.
marrubioides, *Durand & Hilg. in Pacif. Rail. Rep.* v. (1855) 6.—Calif.
mexicanum, *Hemsl. Biol. Centr. Am. Bot.* i. 99.—Mexic.
Munroanum, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 21 = Sphaeralcea Munroana.
Orbignyanum, *Wedd. Chlor. And.* ii. 279.—Bolivia.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 250.—Calif.
parnassiaefolium, *A. Gray, Bot. U. St. Expl. Exped.* i. 150, adnot.—Ecuador.
pedatifidum, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 160 = Sphaeralcea Fendleri.
pedicullariaefolium, *A. Gray, Bot. U. St. Expl. Exped.* i. 152.—Peruv.
peruvianum, *A. Gray, l. c.* 146.—Mexic.
phyllanthos, *A. Gray, l. c.* 152.—Ecuador; Peruv.
pichinchense, *A. Gray, l. c.*—Ecuador.
pinnatum, *A. Gray, l. c.* 154.—Peruv.
plumosum, *A. Gray, l. c.* 147.—Peruv.
procumbens, *Harv. in Harv. & Sond. Fl. Cap.* i. 164.—Afr. austr.
prostratum, *Hieron, in Bol. Acad. Nac. Córdova*, iv. (1881) 15.—Reg. Argent.
Purdiaei, *A. Gray, Bot. U. St. Expl. Exped.* i. 150.—N. Granat.
pygmaeum, *A. Gray, l. c.* adnot.—Bolivia.
pygmaeum, Griseb. in Goett. Abh. xxiv. (1879) 43 = tenellum.
racemosum, *Harv. in Harv. & Sond. Fl. Cap.* i. 164.—Afr. austr.
rhizanthum, *A. Gray, Bot. U. St. Expl. Exped.* i. 148.—Peruv.
ribifolium, *Hemsl. Biol. Centr. Am. Bot.* i. 100.—Mexic.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 149.—Peruv.
roseum, Hemsl. Biol. Centr. Am. Bot. i. 100 = Sphaeralcea umbellata.
rotundifolium, *A. Gray, in Proc. Am. Acad.* vii. (1868) 333.—Calif.
ruderale, Hance, in Walp. Ann. iii. 830 = tricuspidatum.
Rugelii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 367.—Am. bor.; Ind. occ.
scabrum, *A. Gray, Bot. U. St. Expl. Exped.* i. 147.—Peruv.
scoparium, *A. Gray, l. c.* in textu.—Am. trop.

MALVASTRUM:—
setosum, *Harv. in Harv. & Sond. Fl. Cap.* i. 163.—Afr. austr.
spicatum, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 22.—Reg. trop.
splendidum, *Kellogg, in Proc. Calif. Acad.* i. (1855) 65.—Calif.
stenopetalum, *A. Gray, Bot. U. St. Expl. Exped.* i. 155.—Peruv.
strictum, *A. Gray, & Harv. in Harv. & Sond. Fl. Cap.* i. 160.—Afr. austr.
subtriflorum, *Hemsl. Biol. Centr. Am. Bot.* i. 100.—Mexic.
sulphureum, Griseb. in Goett. Abh. xxiv. (1879) 43 = Sida sulphurea.
tenellum, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 15.—Reg. Argent.
Thurberi, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 307.—Calif.
tricuspidatum, *A. Gray, Pl. Wright.* i. 16.—Cosmop. trop.
tridactylites, *Garcke, in Bonplandia*, v. (1857) 296; *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 162.—Afr. austr.
trilobum, *Garcke, in Bot. Zeit.* xxii. (1864) 11.—Afr. austr.
ulophyllum, *A. Gray, Bot. U. St. Expl. Exped.* i. 150.—Peruv.
violaceum, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 15.—Reg. Argent.
virgatum, *A. Gray & Harv. in Harv. & Sond. Fl. Cap.* i. 162.—Afr. austr.
vitifolium, *Hemsl. Biol. Centr. Am. Bot.* i. 100.—Mexic.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 21.—Texas.

MALVAVISCUS, Dill. ex Adans. Fam. ii. 399 (1763). *MALVACEAE*, Benth. & Hook. f. i. 206.
ACHANIA, Sw. Prod. Fl. Ind. Occ. 102 (1788).
acapulcensis, *H. B. & K. Nov. Gen. et Sp.* v. 286.—Mexic.
acerifolius, *Presl, Rel. Haenk.* ii. 135.—Mexic.
arboreus, *Cav. Diss.* iii. t. 48. f. 1.—Am. austr.
Balbisii, *DC. Prod.* i. 445.—Bras.
brevipes, *Benth. Bot. Voy. Sulph.* 68.—Mexic.
candidus, [*Moç. & Sesse,*] *ex DC. Prod.* i. 445.—Mexic.
ciliatus, *DC. l. c.*—Ins. Tobago.
coccineus, Medic. Malv. 49 = Hibiscus coccineus.
concinnus, *H. B. & K. Nov. Gen. et Sp.* v. 286.—Peruv.
concinnus, Schlecht. in Linnaea, xi. (1837) 361 = sepium.
cordatus, *DC. Prod.* i. 445.—Bras.
cordifolius, Moench, Meth. Suppl. 208 = arboreus.
cuspidatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 190.—Venezuela.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 230.—Texas.
flavidus, [*Moç. & Sesse,*] *ex DC. Prod.* i. 446.—Mexic.
floridanus, Nutt. in Journ. Acad. Philad. vii. (1834) 89 = Hibiscus Bancroftianus.
fragilis, Bory, ex DC. Prod. i. 446 = Hibiscus liliiflorus.
glabrescens, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 168 = oligotrichus.
grandiflorus, *H. B. & K. Nov. Gen. et Sp.* v. 286.—Mexic.
leucocarpus, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 169.—N. Granat.
longifolius, *Garcke, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 321.—Peruv.
longifolius, Spach, Hist. Vég. Phan. iii. 370 = Pavonia longifolia.
mollis, *DC. Prod.* i. 445.—Mexic.
montanus, Mart. ex Garcke, in Eichl. Jahrb. Berl. i. (1881) 222 = Pavonia montana.
multiflorus, Spach, Hist. Vég. Phan. iii. 370 = Pavonia multiflora.
oligotrichus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 190.—N. Granat.
penduliflorus, [Moç. & Sesse,] ex DC. Prod. i. 445 = Hibiscus Bancroftianus?
pentacarpus, [*Moç. & Sesse,*] *l. c.*—Mexic.
pilosus, DC. Prod. l. c. = Hibiscus Bancroftianus.

MALVAVISCUS:—
pilosus, Macfad. Fl. Jamaic. i. 64 (excl. syn. Sw.) = arboreus.
pleurantherus, [*Moç. & Sesse,*] *ex DC. Prod.* i. 446.—Mexic.
pleurogonus, [*Moç. & Sesse,*] *l. c.*—Mexic.
Poeppigii, G. Don, Gen. Syst. i. 475 = Hibiscus Poeppigii.
populifolius, *Presl, Rel. Haenk.* ii. 135.—Mexic.
populneus, Gaertn. Fruct. ii. 253 = Thespesia populnea.
pulvinatus, *A. Rich. Ess. Fl. Cub.* 133.—Cuba.
puniceus, Bory, ex DC. Prod. i. 446 = Hibiscus liliiflorus.
Sagraeanus, A. Rich. Ess. Fl. Cub. 131 = arboreus.
sepium, *Schlecht. in Linnaea*, xi. (1837) 361.—Mexic.
spathulatus, *Garcke, in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 321.—Mexic.
velutinus, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 168.—N. Granat.

MALVELLA, Jaub. & Spach, Illustr. v. 47, t. 444 (1853-57) = **Sida**, Linn. (Malvac.).
cymbalariaefolia, Boiss. Fl. Orient. i. 835 = S. Sherardiana.
Sherardiana, Jaub. & Spach, Illustr. v. 46. t. 444 = S. Sherardiana.

MALVEOPSIS, Presl, Bot. Bemerk. 19 (1844) = **Sphaeralcea**, A. St. Hil. (Malvac.).
anomala, Presl, l. c. = S. elegans.

MALVINDA, Dill. ex Medic. Malv. 23 (1787) = **Sida**, Linn. (Malvac.).
alba, Medic. Malv. 24 = S. spinosa.
alnifolia, Medic. l. c. 23 = S. spinosa.
angustifolia, Medic. l. c. 24 = S. angustifolia.
carpinifolia, Medic. l. c. = S. acuta.
cordifolia, Medic. l. c. = S. cordifolia.
palmata, Moench, Meth. Suppl. 203 = S. ricinoides.
spinosa, Medic. Malv. 23 = S. spinosa.
umbellata, Medic. l. c. 24 = Abutilon umbellatum.

MALYA, Opiz, Seznam, 62 (1852) = **Ventenata**, Koel. (Gramin.).
tenuis, Opiz, l. c. 63 = V. avenacea.

MAMBOGA, Blanco, Fl. Filip. ed. I. 140 (1837) = **Mitragyna**, Korth. (Rubiac.).
capitata, Blanco, l. c.—Ins. Philipp.

MAMILLARIA, Reichb. in Moessl. Handb. ed. II. i. p. 1 (1827) = **Mammillaria**, Haw. (Cact.).

MAMMEA, Linn. Gen. ed. I. 344 (1737). *GUTTIFERAE*, Benth. & Hook. f. i. 176.
POTAMOCHARIS, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 296.
africana, G. Don, Gen. Syst. i. 619 = Ochrocarpus africanus.
americana, *Linn. Sp. Pl.* 512.—Am. trop.
angustifolia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 246.—Madag.
asiatica, Linn. Sp. Pl. 512 = Barringtonia Butonica.
birmannica, T. Anders. in Journ. Linn. Soc. ix. (1867) 261 = Ochrocarpus siamensis.
emarginata, *Moç. & Sesse, ex DC. Prod.* i. 561.—Mexic.
eugenioides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 245.—Madag.
excelsa, Planch. & Triana, l. c. 244 = Ochrocarpus ovalifolius.
humilis, *Vahl, Eclog.* ii. 40.—Am. trop.
lateriflora, Griseb. in Goett. Abh. vii. (1857) 184 = humilis.
longifolia, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 240 = Ochrocarpus longifolius.
sessiliflora, *Planch. & Triana, l. c.* 245.—Madag.
siamensis, T. Anders. in Journ. Linn. Soc. ix. (1867) 261 = Ochrocarpus siamensis.

MAMMILARIA, Torr. & Gray, Fl. N. Am. i. 553 (1840) sphalm. = seq.

MAMMILLARIA, Haw. Syn. Pl. Succ. 177 (1812). *CACTEAE*, Benth. & Hook. f. i. 847.
ANHALONIUM, Lem. Cact. Nov. Gen. et Sp. 1 (1841).
ARIOCARPUS, Scheidw. in Bull. Acad. Brux. v. (1838) 491.
CORYPHANTA, Lem. Cact. 32 (1868).
STROMATOCACTUS, Karw. ex Foerst. Handb. Cact. ed. II. i. 229 (1885).

MAMMILLARIA:—
acanthophlegma, *Lehm. Del. Sem. Hamb.* (1832); *Pfeiff. Enum. Diagn. Cact.* 26.—Mexic.
acanthostephes, *Lehm. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 228.—Mexic.
acicularis, *Lem. Cact. Nov. Gen. et Sp.* (1839) 34.—Mexic.
aciculata, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = discolor.
aciculata, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 29.—Mexic.
actinoplea, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 266.—Mexic.
aduncа, Scheidw. ex Foerst. Handb. Cact. ed. I. 222 = uncinata.
aeruginosa, *Scheidw. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 338.—Mexic.
affinis, *DC. Coll. Mém.* viii. *Cact.* 11. t. 6.—Mexic.
aggregata, Engelm. in Emory, Mil. Reconnois. 156 = Cereus phoeniceus.
albida, *Haage, ex Pfeiff. Enum. Diagn. Cact.* 28.—Mexic.
albiseta, *Hort. ex Foerst. Handb. Cact.* ed. II. i. 354.—Mexic. ?
aloides, Monv. Cat. (1846); ex Labour. Cact. 153 = prismatica.
alpina, *Mart. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 7, 79.—Mexic.
amabilis, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 326.—Mexic.
ambigua, *G. Don, ex Loud. Hort. Brit.* 194.—Chili.
amoena, *Hopfr. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 12, 99.—Hab. ?
anancistria, *Lem. Cact. Gen. et Sp. Nov.* (1839) 39.—Hab. ?
ancistracantha, Lem. l. c. 36 = raphidacantha.
ancistrata, Schelh. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 8 = ancistroides.
ancistria, *Walp. Rep.* ii. 296.—Hab. ?
ancistrina, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 10 = ancistroides.
ancistroides, Lehm. Del. Sem. Hort. Hamb. (1832) = glochidiata.
ancistroides, *Lem. Cact. Gen. et Sp. Nov.* (1839) 38.—Hab. ?
Andreae, Pfeiff. ex Salm-Dyck, Cact. Hort. Dyck. (1845) 7 = rhodantha.
anguinea, *Otto, ex Salm-Dyck, l. c.* ed. II. 101.—Mexic.
angularis, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830) 430, nomen; et ex Pfeiff. Enum. Diagn. Cact. 12 = compressa.
anisacantha, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 11; ed. II. 17 = polyëdra.
applanata, Engelm. ex Salm-Dyck, l. c. ed. II. 109 = Heyderi.
areolosa, *Hemsl. Biol. Centr. Am. Bot.* i. 503.—Mexic.
argentea, *Fennel, in Otto & Dietr. Allg. Gartenz.* xv. (1847) 65.—Mexic.
arietina, Lem. Cact. Aliq. Nov. Desc. 10 = magnimamma.
arizonica, *Engelm. in S. Wats. Pl. Wheeler*, 9.—Am. bor. occ.
Asterias, *Cels, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 19, 129.—Hab. ?
atrata, Hort. Mack. ex Hook. Bot. Mag. sub t. 3642 = rhodantha.
atrorubra, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 327.—Mexic.
atrosanguinea, *Ehrenb. l. c.* 270.—Mexic.
aulacantha, *DC. Rév. Cact.* 113.—Mexic.
aulacothele, *Lem. Cact. Aliq. Nov. Desc.* 8.—Mexic.
aurata, Hort. ex Pfeiff. Enum. Diagn. Cact. 31 = rhodantha.
aurea, Hort. ex Foerst. Handb. Cact. ed. I. 200 = Odieriana.
aurea, Pfeiff. ex Foerst. l. c. 198 = rhodantha.
aureiceps, *Lem. Cact. Aliq. Nov. Desc.* 8.—Mexic.
auricoma, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 308.—Mexic.
aurorea, *A. Dietr. l. c.* xvii. (1849) 363.—Mexic.
autumnalis, *A. Dietr. l. c.* xvi. (1848) 287.—Hab. ?
badispina, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 159.—Hab. ?
barbata, *Engelm. in Wisliz. Tour North Mexic.* 105.—Mexic
Barlowii, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 46.—Mexic.
bellatula, *Foerst. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 51.—Bras.

MAMMILLARIA:—
Beneckei, Ehrenb. in Bot. Zeit. ii. (1844) 833 = Goodridgii.
Bergenii, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 326.—Mexic.
Bergii, *Miq. Comm. Phyt.* iii. 104.—Mexic.
bicolor, *Lehm. Del. Sem. Hort. Hamb.* (1830) 7.—Mexic.
bifurca, *A. Dietr. in Allg. Gartenz.* xviii. (1850) 188.—Mexic.
biglandulosa, *Pfeiff. in Otto & Dietr. l. c.* vi. (1838) 274.—Mexic.
bihamata, *Pfeiff. l. c.*—Mexic.
binops, *Haage, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab. ?
bocasana, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 94.—Texas.
Bockii, *Foerst. l. c.* xv. (1847) 50.—Mexic.
brevimamma, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 34.—Mexic.
breviseta, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvi. (1849) 251.—Mexic.
Brongniartii, Hort. Par. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 9, 86 = microthele.
bumamma, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 243 = elephantidea.
caesia, Ehrenb. l. c. 251 = spinosissima.
caespititia, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 7 = multiceps.
caespitosa, A. Gray, Struct. Bot. 421. f. 838 = missouriensis.
caespitosa, Hort. ex Pfeiff. Enum. Diagn. Cact. 6 = echinata.
calcarata, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 195.—Texas.
candida, Scheidw. in Bull. Acad. Brux. v. (1838) 496 = sphaerotricha.
canescens, Hort. ex Pfeiff. Enum. Diagn. Cact. 28, 33 = discolor, grandiflora.
canescens, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 89.—Mexic.
Cantera, *Haage, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab. ?
Caput-Medusae, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830), 430, nomen; et ex Pfeiff. Enum. Diagn. Cact. 22 = Sempervivi.
caracassana, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 14; 107.—Venezuela.
carnea, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 19.—Mexic.
castaneoides, Hort. ex Salm-Dyck, Cact. Hort. Dyck. 5 = Wegeneri.
castaneoides, *Lem. ex Labour. Cact.* 37.—Mexic.
catafracta, Mart. ex Salm-Dyck, Hort. Dyck. 155 = affinis.
cataphracta, Mart. ex Pfeiff. Enum. Diagn. Cact. 11 = affinis.
caudata, *G. Don, ex Loud. Hort. Brit.* 194.—Chili.
Celsiana, *Lem. Cact. Nov. Gen. et Sp.* (1839) 41.—Mexic.
centricirrha, *Lem. l. c.* 42.—Mexic.
centrispina, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 258.—Mexic.
cephalophora, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 137.—Mexic.
ceratocentra, *Berg, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 130.—Mexic.
ceratophora, Lehm. l. c. (1835) 228 = magnamamma.
cespititia, *DC. in Mém. Mus. Par.* xvii. (1828) 112.—Mexic.
chlorantha, *Engelm. in Wheeler, Rep. Bot.* vi. 127.—Mexic.
chrysacantha, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 430, nomen; *et Pfeiff. Enum. Diagn. Cact.* 28.—Mexic.
circumtexta, *Mart. Hort. Reg. Monac.* (1829) 127, nomen.—Mexic.
cirrhifera, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. 1. (1832) 234.—Mexic.
cirrifera, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 18 = compressa.
cirrosa, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 94.—Mexic.
citrina, *Scheidw. ex Foerst. Handb. Cact.* ed. I. 518 = Mexic. ?
Clava, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 282.—Mexic.

MAMMILLARIA:—
clavata, Scheidw. in Bull. Acad. Brux. v. (1838) 494 = raphidacantha.
Closiana, *Roumeg. in Bull. Soc. Bot. Fr.* ii. (1855) 372.—Mexic.
coccinea, *G. Don, in Loud. Hort. Brit.* 194.—Chili.
columbiana, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 12, 99.—N. Granat.
columnaris, Mart. in Nov. Act. Nat. Cur. xvi. 1. (1832) 330 = polythele.
communis, Steud. Nom. ed. II. i. 245, in syn. = Melocactus communis.
compacta, *Engelm. in Wisliz. Tour North Mexic.* 105.—N. Mexic.
compacta, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 9 = microthele.
compressa, *DC. in Mém. Mus. Par.* xvii. (1828) 112.—Mexic.
confinis, Haage, Cat. Cact. 1886; ex Walp. Rep. ii. 289 = albida.
conica, *Haw. Pl. Succ. Suppl.* 71.—Am. austr.
coniflora, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = discolor.
conimamma, *A. Linke, in Koch, Allg. Gartenz.* xxv. (1857) 239.—Mexic.
conoidea, *DC. in Mém. Mus. Par.* xvii. (1828) 112.—Mexic.
conopsea, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 12 = Neumanniana.
conopsea, *Scheidw. in Bull. Acad. Brux.* v. (1838) 496; vi. (1839) 92.—Mexic.
contacta, *Wendl. ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
convoluta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = M. Helicteres.
corioides, *Bosch, ex Sweet, Hort. Brit.* ed. III. 281.—Am. austr.
cornifera, *DC. in Mém. Mus. Par.* xvii. (1828) 112.—Mexic.
corollaria, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 294.—Mexic.
coronaria, *Haw. Rev. Pl. Succ.* 69.—Mexic.
coronata, Scheidw. in Otto & Dietr. Allg. Gartenz. viii. (1840) 338 = crebrispina.
coryphides, *Forb. ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
crassispina, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 406.—Mexic.
crebrispina, *DC. in Mém. Mus. Par.* xvii. (1828) 111.—Mexic.
criniformis, DC. in Mém. Cact. 8. t. 4 = glochidiata.
crinigera, *Otto, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
crinita, *DC. in Mém. Mus. Par.* xvii. (1828) 112.—Mexic.
crocidata, *Lem. Cact. Aliq. Nov. Desc.* 9.—Mexic.
crucigera, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 9, 87 = formosa.
crucigera, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. 1. (1832) 340.—Mexic.
cubensis, Zucc. ex Labour. Cact. 59 = crucigera.
cuneiflora, *Hitchen, ex Forbes, Hort. Tour Germ.* 1 nomen.—Mexic.
curvata, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 15 = exsudans.
curvispina, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 7 = discolor.
curvispina, *Otto, in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 204.—Mexic.
cylindracea, DC. in Mém. Mus. Par. xvii. (1828) 111 = eriacantha.
cylindrica, *Hitchen, ex Forbes, Hort. Tour Germ.* 149, nomen.—Am. austr.
cylindrica, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830) 429, nomen = eriacantha.
dactylothele, Labour. Monog. Cact. 146 = macromeris.
daedalea, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 6 = bicolor.
daemonoceras, Lem. ex Foerst. Handb. Cact. ed. I. 251 = cornifera.
daimonoceras, Lem. Cact. Aliq. Nov. Desc. 5 = scolymoides.
dasyacantha, *Engelm. Syn. Cact.* 268.—Mexic.
dealbata, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 309.—Mexic.
decipiens, *Scheidw. in Bull. Acad. Brux.* v. (1838) 496.—Mexic.

MAMMILLARIA:—
declivis, A. Dietr. in Allg. Gartenz. xviii. (1850) 235 = Heyderi.
decora, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 159.—Mexic.
deficiens, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 7 = decipiens.
deficum, Hort. ex Foerst. Handb. Cact. ed. I. 185 = decipiens.
deflexispina, Lem. Cact. Aliq. Nov. Desc. 6 = gladiata.
densa, Link & Otto, Ic. Pl. Rar. t. 35 = echinata.
depressa, DC. Cat. Hort. Monsp. ex DC. Mém. Cact. 28 = discolor.
depressa, *Scheidw. in Bull. Acad. Brux.* v. (1838) 449.—Mexic.
deserti, *Engelm. in S. Wats. Bot. Calif.* ii. 449.—Calif.
diacantha, Haage, ex Steud. Nom. ed. II. ii. 96 = Haageana.
diacantha, Lem. Cact. Aliq. Nov. Desc. 2 = Sempervivi.
diacentra, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 91.—Hab.?
Diadema, *Muehlenpf. in Otto & Dietr. l. c.* xiii. (1845) 346.—Mexic.
diaphanacantha, Lem. Cact. Aliq. Nov. Desc. 39 = conoidea.
dichotoma, *Hort. ex Sweet, Hort. Brit.* ed. III. 281, nomen.—Hab.?
digitalis, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 267 = Haynii.
disciformis, DC. in Mém. Mus. Par. xvii. (1828) 114 = Echinocactus turbiniformis.
discolor, *Haw. Syn. Pl. Succ.* 177.—Am. austr.
divaricata, *A. Dietr. in Allg. Gartenz.* xvi. (1848) 210.—Mexic.?
divergens, *DC. in Mém. Mus. Par.* xvii. (1828) 113.—Am. austr.
dolichacantha, Lem. ex Foerst. Handb. Cact. ed. I. 213 = dolichocentra.
dolichocentra, *Lem. Cact. Aliq. Nov. Desc.* 3.—Mexic.
Dyckiana, Zucc. ex Pfeiff. Enum. Diagn. Cact. 26 = acanthophlegma.
eborina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 309.—Mexic.
eburnea, Miq. in Linnaea, xii. (1838) 14 = bicolor.
echinaria, DC. in Mém. Mus. Par. xvii. (1828) 110 = echinata.
echinata, *DC. Mém. Cact.* 3.—Mexic.
echinocactoides, Pfeiff. in Otto & Dietr. Allg. Gartenz. viii. (1840) 281 = conoidea.
Echinops, Scheidw. ex Foerst. Handb. Cact. ed. I. 209 = oothele.
Echinus, *Engelm. Syn. Cact.* 11.—Texas.
Ehrenbergii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 274.—Mexic.
elegans, *DC. in Mém. Mus. Par.* xvii. (1828) 111.—Mexic.
elephantidens, *Lem. Cact. Aliq. Nov. Desc.* 1.—Mexic.
elongata, *DC. in Mém. Mus. Par.* xvii. (1828) 109.—Mexic.
elongata, *Hemsl. in Biol. Cent. Am. Bot.* i. 509.—Mexic.
Emundtsiana, *Hort. ex Foerst. Handb. Cact.* ed. II. i. 341.—Hab.?
enneacantha, *Otto, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
erecta, *Lem. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 370; *Lem. Cact. Aliq. Nov. Desc.* 3.—Mexic.
erectacantha, *Foerst. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 50.—Mexic.
eriacantha, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 32.—Mexic.
eriacantha, *Hort. ex Sweet, Hort. Brit.* ed. III. 281, nomen.—Mexic.
eriantha, Hort. ex Pfeiff. Enum. Diagn. Cact. 32 = eriacantha, *Hort. Berol.*
erinacea, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 18.—Mexic.
erinacea, Wendl. f. Cat. Herrenh. (1835) = rhodantha.
euchlora, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 266.—Mexic.
Eugenia, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 7 = rutila.
evanescens, Hort. Belg. ex Foerst. Handb. Cact. ed. I. 243 = erecta.

MAMMILLARIA:—
eximia, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 309.—Mexic.
exsudans, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 15.—Mexic.
falcata, *Aut. ex Foerst. Handb. Cact.* ed. II. i. 345.—Mexic.
farinosa, Hort. ex Fennel, in Otto & Dietr. Allg. Gartenz. xv. (1847) 66 = livida.
fasciculata, *Engelm. in Emory, Mil. Reconnois.* 156. f. 2.—Am. bor. occ.
Fellneri, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvi. (1849) 261.—Mexic.
Fennelii, Hopper, in Otto & Dietr. l. c. xi. (1843) 3 = zephyranthoides.
Fischeri, *Pfeiff. in Otto & Dietr. l. c.* iv. (1836) 257.—Mexic.
fissurata, *Engelm. Syn. Cact.* 14.—Mexic.
flava, *Otto & Dietr. Allg. Gartenz.* xvii. (1849) 261.—Mexic.
flavescens, *DC. Prod.* iii. 459.—Am. trop.
flavescens, *Pfeiff. Enum. Diagn. Cact.* 10.—Mexic.
flavescens, Zucc. ex Foerst. Handb. Cact. ed. II. i. 348 = Karwinskiana.
flaviceps, Scheidw. ex Labour. Cact. 42 = crassispina.
flavicoma, *Hort. ex Foerst. Handb. Cact.* ed. II. i. 298.—Hab.?
flavovirens, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 16, 117.—Hab.?
floccigera, Otto, ex Salm-Dyck, l. c. 11 = crassispina.
floribunda, *Hook. Bot. Mag.* t. 3647.—Chili.
Foersteri, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 371.—Mexic.
formosa, *Gal. ex Scheidw. in Bull. Acad. Brux.* v. (1838) 497.—Mexic.
formosa, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xii. (1844) 186.—Mexic.
foveolata, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 372.—Mexic.
fuliginosa, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 10, 93.—Venezuela?
fulvispina, *Haw. in Phil. Mag.* (1829) 106; (1830) 109.—Bras.; Mexic.
Funkii, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 43.—Mexic.
fuscata, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1820) 429, nomen; *et ex Pfeiff. Enum. Diagn. Cact.* 28.—Mexic.
Galeotti, Scheidw. ex Foerst. Handb. Cact. ed. I. 213 = dolichocentra.
Gebweileriana, *Haage, Cat. ex Foerst. Handb. Cact.* ed. II. i. 358.—Hab.?
geminata, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 42.—Mexic.
geminispina, DC. in Mém. Mus. Par. xvii. (1828) 30. t. 3 = acanthophlegma.
geminispina, Haw. in Till. Phil. Mag. lxiii. (1824) 42 = bicolor.
gibbosa, *Salm-Dyck, Hort. Dyck.* 343.—Hab.?
gigantothele, Hort. ex Foerst. Handb. Cact. ed. I. 183 = longimamma.
glabrata, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 109.—Mexic.
gladiata, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. 1. (1832) 336.—Mexic.
glanduligera, *Otto & Dietr. Allg. Gartenz.* xvi. (1848) 298.—Mexic.
glauca, *Dietr. ex A. Linke, l. c.* 330.—Mexic.
globosa, *A. Linke, in Koch, Allg. Gartenz.* (1857) 240.—Mexic.
glochidiata, *Mart. Hort. Reg. Monac.* (1829) 12, nomen; *et in Nov. Act. Nat. Cur.* xvi. 1. (1832) 337. t. 23.—Mexic.
glomerata, *DC. Prod.* iii. 459.—Ins. S. Doming.
Goodridgii, *Scheer, ex Salm-Dyck, Cact. Hort. Dyck* ed. II. 10, 91.—Calif.
gracilis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 275.—Mexic.
Grahami, *Engelm. Syn. Cact.* 262.—Am. bor. occ.
grandicornis, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 372.—Mexic.
grandiflora, Hort. ex Salm-Dyck, Cact. Hort. Dyck. 4 = vetula.
grandiflora, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 430, nomen; *et ex Pfeiff. Enum. Diagn. Cact.* 38.—Mexic.

MAMMILLARIA:—

grandiflora, Scheidw. ex Foerst. Handb. Cact. ed. I. 189 = macrantha.
grandis, *Hitchen, ex Forbes, Hort. Tour Germ.* 149, nomen.—Mexic.
granulata, *Meinsh. in C. Koch & Fint. Wochenschr.* (1858) 264.—Mexic.
grisea, *Gal. ex Foerst. Handb. Cact.* ed. I. 219.—Mexic.?
grisea, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 110.—Mexic.
Guilleminiana, Lem. Cact. Gen. et Sp. Nov. (1839) 48 = decipiens.
gummifera, *Engelm. in Wisliz. Tour North Mexic.* 105.—Mexic.
Haageana, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 257.—Mexic.
haematactina, *Ehrenb. in Otto & Dietr. l. c.* xvi. (1848) 266.—Mexic.
hamata, *Lehm. Del. Sem. Hort. Hamb.* (1832); *et ex Pfeiff. Enum. Cact.* 34.—Mexic.
Haselaffii, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 303 = seq.
Hasseloffii, Ehrenb. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 8 = senilis.
Haynii, *Ehrenb. in Bot. Zeit.* ii. (1844) 833.—Mexic.
Heideri, Auct. vide Heyderi.
Heinei, *Ehrenb. in Bot. Zeit.* ii. (1844) 833.—Mexic.
Helicteres, *DC. in Mém. Mus. Par.* xvii. (1828) 31. t. 5.—Mexic.
hemisphaerica, Engelm. in Bost. Journ. Nat. Hist. vi. (1850) 198 = Heyderi.
Hepatica, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 267 = Seegerii.
Herrmanni, *Ehrenb. l. c.* xvii. (1849) 303.—Mexic.
heteracantha, Mart. Hort. Reg. Monac. (1829) 127, nomen; *et ex Pfeiff. Enum. Diagn. Cact.* 13 = loricata.
heteracentra, *Otto, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Mexic.?
heteromorpha, *Scheer, in Seem. Bot. Voy. Herald*, 289.—Mexic.
hexacantha, *Salm-Dyck, Hort. Dyck.* 344.—Mexic.
hexacentra, Otto, ex Foerst. Handb. Cact. ed. I. 183 = longimamma.
Heyderi, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 20.—Mexic.
Hoffmannseggii, *Hoffmgg. Preiss-Verz.* ed. VII. (1833) 23.—Hab.?
Hopferiana, *A. Linke, in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 329.—Mexic.?
horripila, Lem. Cact. Aliq. Nov. Desc. 7 = Echinocactus horripilus.
Humboldtii, *Ehrenb. in Linnaea*, xiv. (1842) 378.—Mexic.
hybrida, Hort. ex Pfeiff. Enum. Diagn. Cact. 31 = rhodantha.
Hystrix, *Mart. ex Pfeiff. l. c.* 21.—Mexic.
imbricata, Wegen. in Otto & Dietr. Allg. Gartenz. xii. (1844) 66 = Stuberi.
impexicoma, Lem. Cact. Aliq. Nov. Desc. 5 = cornifera.
inclinis, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 9.—Hab.?
inconspicua, Scheidw. in Bull. Acad. Brux. v. (1838) 495 = conoidea.
incurva, *Scheidw. l. c.* vi. (1839) 92.—Mexic.
intertexta, *DC. in Mém. Mus. Par.* xvii. (1828) 110.—Mexic.
intricata, *Otto, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
inuncta, Hoffmgg. Preiss-Verz. ed. VII. (1833) 23 = rhodantha.
irregularis, *DC. in Mém. Mus. Par.* xvii. (1828) 111.—Mexic.
isabellina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 309.—Mexic.
jalappensis, Hort. ex Pfeiff. Enum. Diagn. Cact. 17 = subpolyedra.
jucunda, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 250.—Mexic.
Karstenii, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 95.—Reg. Argent.
Karwinskiana, *Mart. in Nov. Act. Nat. Cur.* xvi. 1. (1832) 335. t. 22.—Mexic.
kewensis, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 112.—Hab.?
Kleinii, *Regel, Ind. Sem. Hort. Petrop.* (1860) 47.—Mexic.

MAMMILLARIA:—

Klugei, Ehrenb. in Bot. Zeit. ii. (1844) 834 = elegans.
Krameri, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 347.—Mexic.
Kunthii, *Ehrenb. in Bot. Zeit.* ii. (1844) 835.—Mexic.
lactescens, *Meinsh. in Koch & Fint. Wochenschr.* (1859) 117.—Mexic.
laeta, Meinsh. ex Foerst. Handb. Cact. ed. II. i. 246 = alpina.
lamprochaeta, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 82.
lanifera, *Haw. in Till. Phil. Mag.* lxiii. (1824) 41.—Mexic.
lanifera, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 12, 98.—Mexic.
lasiacantha, *Engelm. Syn. Cact.* 5.—Texas.
latimamma, DC. in Mém. Mus. Par. xvii. (1828) 114 = pycnacantha.
Lehmanni, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830) 430, nomen; et ex Pfeiff. Enum. Cact. 15 = recurva.
Lehmannii, *Fennel, in Otto & Dietr. Allg. Gartenz.* xv. (1847) 66.—Hab.?
leona, *Poselg. l. c.* xxi. (1853) 94.—Mexic.?
leucacantha, *DC. in Mém. Mus. Par.* xvii. (1828) 113.—Mexic.
leucantha, DC. ex Steud. Nom. ed. II. ii. 97 = recurva.
leucocarpa, Scheidw. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 17 = maschalacantha.
leucocentra, *Berg, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 130.—Mexic.
leucocephala, Hort. Par. ex Pfeiff. Enum. Diagn. Cact. 180 = acanthophlegma.
leucodasys, *Salm-Dyck, ex Scheer, in Seem. Bot. Voy. Herald*, 286.—Mexic.
leucodictia, *Link, ex Otto & Dietr. Allg. Gartenz.* xvi. (1848) 330.—Mexic.
leucotricha, *Scheidw. l. c.* viii. (1840) 338.—Mexic.
Lindheimeri, Engelm. ex Hemsl. Biol. Centr. Am. Bot. i. 525 = texensis.
Linkeana, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 308.—Mexic.
Linkei, Ehrenb. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 8 = senilis.
livida, *Fennel, in Otto & Dietr. Allg. Gartenz.* xv. (1847) 66.—Mexic.
longimamma, *DC. in Mém. Mus. Par.* xvii. (1828) 113.—Mexic.
longiseta, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 346.—Mexic.
longispina, Reichb. Suppl. Terscheck, Cact. Verz. 1 = dolichocentra.
loricata, Hort. ex Foerst. Handb. Cact. ed. I. 214 = polythele.
loricata, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et ex Pfeiff. Enum. Diagn. Cact.* 13.—Mexic.
Ludwigii, *Ehrenb. in Linnaea*, xiv. (1840) 376.—Mexic.
lutescens, *Hort. ex Forbes, Hort. Tour Germ.* 149, nomen.—Hab.?
macracantha, *DC. in Mém. Mus. Par.* xvii. (1828) 113.—Mexic.
macrantha, *Salm-Dyck, ex Foerst. Handb. Cact.* ed. I. 189 (an praec. ?).—Hab.?
macromeris, *Engelm. in Wisliz. Tour North Mexic.* 97.—Mexic.
macrothele, Hort. Berol. ex Pfeiff. in Linnaea, xii. (1838) 140 = Martiana.
macrothele, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 24.—Mexic.
Maelenii, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 14.—Hab.?
magnimamma, *Haw. in Till. Phil. Mag.* lxiii. (1824) 41.—Mexic.
Martiana, *Pfeiff. in Linnaea*, xii. (1838) 140.—Mexic.
maschalacantha, *Monv. Cat.* 77; *ex Labour. Cact.* 106.—Mexic.
megacantha, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 123.—Mexic.
meiacantha, *Engelm. in Pacif. Rail. Rep.* iv. 27.—Texas.
Meissneri, Ehrenb. in Bot. Zeit. ii. (1844) 834 = acanthophlegma.
melaleuca, *Karw. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 108.—Mexic.
melanacantha, *Hort. ex Foerst. Handb. Cact.* ed. II. i. 386.—Hab.?
melanocentra, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 18.—Mexic.

MAMMILLARIA:—

meschalacantha, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 10 = maschalacantha.
micans, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 330.—Mexic.
micracantha, *Miq. in Linnaea*, xii. (1838) 16.—Mexic.
micracantha, Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 9 = caracassana.
micrantha, Hort. ex Foerst. Handb. Cact. ed. II. i. 335 = caracassana.
microcarpa, Engelm. in Emory, Mil. Reconnois. 156. f. 3 = Grahami?
microceras, *Lem. Cact. Aliq. Nov. Desc.* 6.—Mexic.
micromeris, *Engelm. Syn. Cact.* 260.—Texas.
microthele, Mart. Hort. Reg. Monac. (1829) 127, nomen (Quid?).—Mexic.
microthele, Monv. ex Foerst. Handb. Cact. ed. II. i. 335 = caracassana.
microthele, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 11.—Mexic.?
minima, *Reichb. in Terscheck, Suppl. Cact. Verz.* 1; *et ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 100.—Mexic.
Miqueliana, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
mirabilis, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 251.—Mexic.
missouriensis, Scheer, in Seem. Bot. Voy. Herald, 287 = vivipara.
missouriensis, *Sweet, Hort. Brit.* ed. I. 171.—Am. bor. occ.
mitis, *Mill. Dict. ex DC. Prod.* iii. 460.—Am. austr.
monancistria, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 7 = Wildiana.
monocentra, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 90.—Hab.?
mucronata, *Ehrenb. l. c.* xvii. (1849) 294.—Mexic.
Muehlenpfordtii, *Foerst. l. c.* xv. (1847) 49.—Mexic.
multiceps, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 7, 81.—Mexic.
multiradiata, *Mart. Hort. Reg. Monac.* (1829) 127, nomen.—Mexic.
multiseta, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 242.—Mexic.
mutabilis, *Scheidw. l. c.* ix. (1841) 43.—Mexic.
Mystax, Hort. ex Pfeiff. Enum. Diagn. Cact. 18 = villifera.
Mystax, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. 1. (1832) 332. t. 21.—Mexic.
neglecta, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = rhodantha.
Neumanniana, *Lem. Cact. Nov. Gen. et Sp.* 53; *ex Walp. Rep.* ii. 297.—Mexic.
nigra, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 287.—Mexic.
nigricans, *Fennel, l. c.* xv. (1847) 66.—Hab.?
nigricans, Pfeiff. Abbild. 2. t. 23 = phaeacantha.
nitens, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = discolor.
nitens, *Otto, ex A. Linke, in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 331.—Mexic.?
nitida, Scheidw. l. c. ix. (1841) 42 = cespititia.
nivea, Wendl. f. Cat. Hort. Herrenh. (1835) = bicolor.
nivosa, Hort. ex Forbes, Hort. Tour Germ. 149, nomen = seq.?
nivosa, *Link, ex Pfeiff. Enum. Diagn. Cact.* 11.—Ins. Tortola.
nobilis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 282.—Mexic.
nuda, *DC. Prod.* iii. 460.—Mexic.
Nuttallii, Engelm. in A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 49 = missouriensis.
obconella, Scheidw. in Bull. Acad. Brux. vi. (1839) 91 = dolichocentra.
obliqua, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 250.—Mexic.
obscura, Scheidw. ex Foerst. Handb. Cact. ed. I. 213 = dolichocentra.
obvallata, *Otto, in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 308.—Mexic.
octacantha, *DC. in Mém. Mus. Par.* xvii. (1828) 113.—Mexic.
Odieriana, *Lem. Cact. Nov. Gen. et Sp.* (1839) 46.—Mexic.
olivacea, Hort. ex Pfeiff. Enum. Diagn. Cact. 180 = tentaculata.

MAMMILLARIA:—

olorina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 326.—Mexic.
oothele, *Lem. Cact. Nov. Gen. et Sp.* (1839) 37.—Hab.?
Ottonis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 274.—Mexic.
ovimamma, *Lem. Cact. Nov. Gen. et Sp.* (1839) 49.—Hab.?
pachytele, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 17.—Mexic.
pachythele, Salm-Dyck, Cact. Hort. Dyck. ed. II. 17 = Krameri.
pallescens, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 41.—Mexic.
Palmeri, *Jacobi, l. c.* xxiv. (1856) 82.—Hab.?
papyracantha, Engelm. in Mem. Am. Acad. N. S. iv. (1849) 49 = Echinocactus papyracanthus.
Parkinsonii, *Ehrenb. in Linnaea*, xiv. (1840) 375.—Mexic.
Parmentieri, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830) 429 = flavescens.
parvimamma, *Haw. Suppl. Pl. Succ.* 72.—Ind. occ.
parvissima, Karw. ex Foerst. Handb. Cact. ed. II. i. 260 = multiceps.
Pazzani, *Steeber, in Bot. Zeit.* v. (1847) 491.—Mexic.
Peacocki, Hort. Angl. ex Foerst. Handb. Cact. ed. II. i. 286 = dealbata.
pectinata, *Engelm. Syn. Cact.* 10.—Texas.
pentacantha, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 406.—Mexic.
Perote, Hort. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. iv. (1836) 257 = Haageana.
perpusilla, Meinsh. ex Foerst. Handb. Cact. ed. II. i. 260 = multiceps.
persicina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 250.—Mexic.
Pfeifferi, Booth, ex Scheidw. in Bull. Acad. Brux. vi. (1839) 93 = rhodantha.
Pfeifferiana, *De Vriese, in Tijdschr.* vi. (1839) 51. t. 1. f. 2.—Mexic.
phaeacantha, *Lem. Cact. Nov. Gen. et Sp.* (1839) 47.—Mexic.
phaeotricha, *Monv. Cact.* (1846), *ex Labour. Cact.* 39.—Mexic.
phellosperma, *Engelm. Syn. Cact.* 6.—Calif.
phymatothele, *Berg, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 129.—Mexic.
picta, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 27.—Mexic.
picturata, *Labour. in Rev. Hortic.* Sér. IV. iv. (1855) 28.—Chili.
Plaschnickii, *Pfeiff. Enum. Diagn. Cact.* 24.—Mexic.
plecostigma, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 27.—Mexic.
pleiocephala, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 47.—Mexic.
plinthimorpha, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 92.—Mexic.
polyacantha, Ehrenb. l. c. xvi. (1848) 265 = spinosissima.
polyactina, Ehrenb. l. c. 260 = spinosissima.
polycentra, *Berg, l. c.* viii. (1840) 130.—Mexic.
polycephala, *Muehlenpf. l. c.* xiii. (1845) 347.—Mexic.
polychlora, Scheidw. ex Foerst. Handb. Cact. ed. I. 205 = crebrispina.
polyëdra, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 326. t. 18.—Mexic.
polygona, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 17. 120.—Mexic.
polygona, Zucc. ex Pfeiff. Enum. Diagn. Cact. 17 = subpolyëdra.
polymorpha, *Scheer, ex Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiv. (1846) 373.—Hab.?
polythele, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 328. t. 19.—Mexic.
polytricha, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* x. (1842) 289.—Hab.?
pomacea, *Ehrenb. l. c.* xvi. (1848) 267.—Mexic.
porphyracantha, *Jacobi, l. c.* xxiv. (1856) 81.—Hab.?
Poselgeriana, *Haage, Cat. ex Foerst. Handb. Cact.* ed. II. i. 269.—Mexic.
potosiana, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 92.—Mexic.

MAMMILLARIA:—

Pottsii, *Scheer, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 104.—Texas.
Praelii, Muehlenpf. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 372 = viridis.
pretiosa, *Ehrenb. l. c.* xvii. (1849) 250.—Mexic.
prismatica, *Hemsl. Biol. Centr. Am. Bot.* i. 517.—Mexic.
procera, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 241.—Mexic.
prolifera, *Haw. Syn. Pl. Succ.* 177.—Am. austr.
prolifera, Hort. ex Pfeiff. Enum. Diagn. Cact. 9 = parvimamma.
prolifera, Pfeiff. ex Foerst. Handb. Cact. ed. II. i. 307 = discolor.
pruinosa, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 261 = Seegerii.
Pseudo-mamillaris, Salm-Dyck, Cact. Hort. Dyck. ed. I. 7; II. 11 = discolor.
pugionacantha, *Foest. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 50.—Mexic.
pulchella, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 10, 94.—Mexic.
pulchella, Otto, ex A. Linke, in Otto & Dietr. Allg. Gartenz. xvi. (1848) 331 = discolor.
pulcherrima, *Ehrenb. l. c.* xvii. (1849) 249.—Mexic.
pulcherrima, *Hort. ex Forbes, Hort. Tour Germ.* 149, nomen.—Hab.?
pulchra, *Haw. in Bot. Reg.* t. 1329.—Mexic.
pulvilligera, Monv. Cat. ex Foerst. Handb. Cact. ed. II. i. 231 = elongata.
punctata, *Labour. ex Foerst. l. c.* 293.—Hab.?
purpurascens, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 260.—Mexic.
purpurea, *Ehrenb. l. c.* 270.—Mexic.
pusilla, *Sweet, Hort. Brit.* ed. I. 171.—'Am.'
pycnacantha, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 325.—Mexic.
pyramidalis, Link & Otto, in Verh. Ver. Befoerd. Gartenb. vi. (1830) 429, nomen = rhodantha.
pyrrhacantha, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
pyrrhocentra, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 10, 93 = fulvispina.
pyrrhocephala, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 42.—Mexic.
pyrrhochracantha, *Lem. Cact. Nov. Gen. et Sp.* (1839) 51.—Mexic.
quadrata, *G. Don, ex Loud. Hort. Brit.* 194.—Chili.
quadrispina, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 329.—Mexic.
radians, *DC. in Mém. Mus. Par.* xvii. (1828) 111.—Mexic.
radians, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 20 = cornifera.
radiosa, Engelm. in Bost. Journ. Nat. Hist. vi. (1850) 196 = vivipara.
Radula, Scheidw. ex Foerst. Handb. Cact. ed. I. 208 = phaeacantha.
raphidacantha, *Lem. Cact. Nov. Gen. et Sp.* (1839) 34.—Mexic.
recta, Miq. ex Labour. Cact. 63 = acanthophlegma.
recurva, *Lehm. ex Pfeiff. Enum. Diagn. Cact.* 15.—Mexic.
recurvata, *Engelm. in Trans. Acad. St. Louis*, ii. (1860–63) 202.—N. Mexic.
recurvispina, *De Vriese, in Tijdschr.* vi. (1839) 53. t. 1. f. 3.—Mexic.
recurvispina, Engelm. Syn. Cact. 10 = recurvata.
regia, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii (1849) 269.—Mexic.
retusa, *Hort. Belg. ex Pfeiff. in Otto & Dietr. l. c.* v. (1837) 369.—Mexic.
retusa, Scheidw. ex Foerst. Handb. Cact. ed. I. 248 = sulco-lanata.
rhaphidacantha, *Lem. Cact. Nov. Gen. et Sp.* (1839) 34.—Mexic.
rhodacantha, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 11. 96.—Hab.?
rhodantha, *Link & Otto, Ic. Select.* t. 26.—Mexic.
rhodeocentra, *Lem. Cact. Nov. Gen. et Sp.* (1839) 52.—Mexic.
robusta, Otto, ex Foerst. Handb. Cact. ed. I. 207; et Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = crassispina.
robustispina, *Schott, ex Engelm. Cact. U. St. Bound.* 10.—Mexic.

MAMMILLARIA:—

rosea, *Gal. ex Foerst. Handb. Cact.* ed. I. 191.—Mexic.?
rosea, *Scheidw. in Hortic. Belg.* (1838) 118. t. 5 (err. 7).—Mexic.
ruficeps, Lem. Cact. Nov. Gen. et Sp. (1839) 37 = rhodantha.
rufidula, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 295.—Mexic.
rufocrocea, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 102.—Mexic.
Ruschiana, *Regel, Ind. Sem. Hort. Turic.* (1850) 4, in adnot.—*Cf. Linnaea*, xxiv. (1851) 231.—Mexic.
russea, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 347.—Mexic.
rutila, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 29.—Mexic.
Salm-Dyckiana, *Scheer, ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 394.—Mexic.
Salmiana, *Fennel, ex Foerst. Handb. Cact.* ed. I. 254, nomen.—Hab.?
sanguinea, *Haage, ex Regel, in Act. Hort. Petrop.* viii. (1883) 276; *Gartenfl.* (1883) 66.—Mexic.
saxatilis, *Scheer, in Seem. Bot. Voy. Herald*, 286.—Mexic.
scepontocentra, *Lem. Cact. Nov. Gen. et Sp.* (1839) 45.—Mexic.
Schaeferi, *Fennel, in Otto & Dietr. Allg. Gartenz.* xv. (1847) 66.—Mexic.
Scheerii, Muehlenpf. l. c. xiii. (1845) 346 = conoidea.
Scheerii, *Muehlenpf. l. c.* xv. (1847) 97.—Bras.
Scheidweileriana, *Otto, in Otto & Dietr. l. c.* ix. (1841) 179.—Mexic.
Schelhasii, *Pfeiff. l. c.* vi. (1838) 274.—Mexic.
Schiedeana, Hort. ex Pfeiff. Enum. Diagn. Cact. 14 = magnimamma.
Schiedeana, *Schlecht. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 249.—Mexic.
Schlechtendalii, *Ehrenb. in Linnaea*, xiv. (1840) 377.—Mexic.
Schmerwitzii, *Haage, ex Foerst. Handb. Cact.* ed. II. i. 270.—Mexic.
Schmidtii, *Sencke, ex Foerst. l. c.* 376.—Hab.?
scolymoides, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 44.—Mexic.
Seegerii, *Ehrenb. l. c.* xvii. (1849) 261.—Mexic.
Seemannii, *Scheer, in Seem. Bot. Voy. Herald*, 288.—Mexic.
Seidelii, *Terscheck, Suppl. Cact. Verz.* 1.—Mexic.
Seitziana, Hort. ex Foerst. Handb. Cact. ed. I. 223 = Karwinskiana.
Seitziana, *Mart. in Pfeiff. Enum. Diagn. Cact.* 18.—Mexic.
Seitziana, Miq. in Linnaea, xii. (1838) 10 = Bergii.
Sempervivi, *DC. in Mém. Mus. Par.* xvii. (1828) 114.—Mexic.
senilis, *Hort. Angl. ex Foerst. Handb. Cact.* ed. I. 189 (an seq. ?).—Hab.?
senilis, *Lodd. ex Scheer, in Seem. Bot. Voy. Herald*, 286.—Mexic.
Senkii, Foerst. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 16 = mutabilis.
sericata, Lem. Cact. Nov. Gen. et Sp. (1839) 44 = magnimamma.
setosa, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 379.—Mexic.
Severini, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 46.—Mexic.
similis, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 246 = missouriensis.
simplex, *Haw. Syn. Pl. Succ.* 177.—Am. austr.
simplex, Torr. & Gray, Fl. N. Am. i. 553 = missouriensis.
solitaria, *G. Don, ex Loud. Hort. Brit.* 194.—Chili.
sororia, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 28.—Mexic.
speciosa, *De Vriese, in Tijdschr.* vi. (1839) 52.—Mexic.
speciosa, *G. Don, ex Steud. Nom.* ed. II. ii. 97.—Chili.
spectabilis, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 15 = kewensis.
spectabilis, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1846) 346.—Mexic.
sphacelata, *Mart. Hort. Reg. Monac.* (1829) 127, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 339.—Mexic.
sphaerica, *Dietr. ex Engelm. Cact. Mex. Bound.* 9.—Texas; Mexic.

MAMMILLARIA :—
sphaerotricha, *Lem. Cact. Nov. Gen. et Sp.* (1839) 33. —Mexic.
spinaurea, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 394.—Mexic.
Spinii, Colla, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = discolor.
spinosa, *G. Don, ex Loud. Hort. Brit.* 194.—Chili.
spinosissima, *Lem. Cact. Aliq. Nov. Desc.* 4.—Mexic.
splendens, Ehrenb. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 242 = acanthophlegma.
squarrosa, *Meinsh. in C. Koch & Fint. Wochenschr.* (1859) 116.—Mexic.
staurotypa, Hort. Belg. ex Foerst. Handb. Cact. ed. I. 221 = Sempervivi.
Stella-aurata, *Mart. Hort. Reg. Monac.* (1829) 128, nomen ; *et in Zucc. Pl. Nov. Fasc.* iii. 101.—Mexic.
stellaris, Haw. Syn. Pl. Succ. 72 = pusilla.
stellata, *Haw. in Till. Phil. Mag.* vii. (1830) 114.—Am. austr.
stellata, *Sweet, Hort. Brit.* ed. I. 171.—Am. austr.
stenocephala, Scheidw. ex Labour. Cact. 50 = dolichocentra.
stenocephala, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 43.—Mexic.
Stephani, *Hort. Vind. ex Foerst. Hand. Cact.* ed. I. 254, nomen.—Hab. ?
stipitata, Scheidw. in Bull. Acad. Brux. v. (1838) 495 = raphidacantha.
straminea, Haw. Suppl. Pl. Succ. 71 = flavescens, *DC.*
strobiliformis, Engelm. in Wisliz. Tour North Mexic. 113 = conoidea.
strobiliformis, Muehlenpf. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 19 = calcarata.
strobiliformis, *Scheer, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 104.—Mexic.
Stueberii, *Foerst. Handb. Cact.* ed. I. 517.—Mexic.
suaveolens, *Hort. ex Foerst. l. c.* ed. II. i. 297.—Mexic.
subangularis, *DC. in Mém. Mus. Par.* xvii. (1828) 112. —Mexic.
subcirrhifera, Hort. ex Foerst. Handb. Cact. ed. I. 234 = subangularis.
subcrocea, *DC. in Mém. Mus. Par.* xvii. (1828) 110.—Mexic.
subcurvata, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xii. (1844) 232.—Mexic.
subechinata, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 101. —Mexic.
subpolyëdra, *Salm-Dyck, l. c.* 343.—Mexic.
subtetragona, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 169.—Mexic.
subulata, *Muehlenpf. l. c.* xiii. (1845) 347.—Chili.
subulifera, *Ehrenb. l. c.* xvii. (1849) 242.—Mexic.
sulcata, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 246.—Texas.
sulcata, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 78.—Mexic.
sulcimamma, Pfeiff. ex Salm-Dyck, l. c. 19 = Plaschnickii.
sulco-glandulifera, *Jacobi, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 92.—Hab. ?
sulco-lanata, *Lem. Cact. Aliq. Nov. Desc.* 2.—Mexic.
sulphurea, Foerst. Handb. Cact. ed. I. 200 ; et ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 11 = rhodantha.
supertexta, Hort. ex Pfeiff. Enum. Diagn. Cact. 25 = elegans.
supertexta, *Mart. Hort. Reg. Monac.* (1829) 128, nomen; *et ex Pfeiff. l. c.*—Mexic.
tecta, *Miq. in Linnaea*, xii. (1838) 12.—Mexic.
tentaculata, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 29.—Mexic.
tentaculata, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 430, nomen ; *et ex Pfeiff. l. c.*—Mexic.
tenuis, *DC. in Mém. Mus. Par.* xvii. (1828) 110.—Mexic.
tetracantha, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 18.—Mexic.
tetracentra, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 112.—Mexic.
tetracentra, *Otto, ex Foerst. Handb. Cact.* ed. I. 214.—Hab. ?
tetrancistra, Engelm. in Am. Journ. Sc. Ser. II. xiv. (1852) 337 (1852), partim = phellosperma.
texana, Young, Fl. Texas. 279 = pusilla.
texensis, Labour. Cact. 89 = Heyderi.
thelocamptos, *Lehm. in Linnaea*, xiii. (1839) *Litt.* 101. —Mexic.
Toaldoae, Lehm. ex Walp. Rep. ii. 289 = bicolor.

MAMMILLARIA :—
tomentosa, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 262.—Mexic.
tortolensis, Hort. ex Pfeiff. Enum. Diagn. Cact. 11 = nivosa.
triacantha, *DC. in Mém. Mus. Par.* xvii. (1828) 113. —Mexic.
tuberculata, Engelm. Syn. Cact. 12 = strobiliformis.
turbinata, *Hook. Bot. Mag.* t. 3984.—Mexic.
uberiformis, *Zucc. in Pfeiff. Enum. Diagn. Cact.* 23.—Mexic.
Uhdeana, Salm-Dyck, Cact. Hort. Dyck. ed. II. 83 = spinosissima.
umbrina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 287.—Mexic.
uncinata, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 34.—Mexic.
variamamma, *Ehrenb. ex Otto & Dietr. Allg. Gartenz.* xvii. (1849) 242.—Mexic.
versicolor, *Scheidw. in Bull. Acad. Brux.* v. (1838) 494.—Mexic.
vetula, *Mart. Hort. Reg. Monac.* (1829) 128, nomen; *et in Nov. Act. Nat. Cur.* xvi. i. (1832) 338. t. 24.—Mexic.
villifera, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 18.—Mexic.
villosa, *Fennel, ex Foerst. Handb. Cact.* ed. I. 255, nomen.—Hab. ?
virens, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 43.—Mexic.
viridis, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 116.—Mexic.
viridula, Ehrenb. in Otto & Dietr. Allg. Gartenz. (1848) 267 = Haynii.
vivipara, *Haw. Suppl. Pl. Succ.* 72.—Am. bor.
vulpina, *Ehrenb. in Otto & Dietr. Allg. Gartenz.* xvii. (1849) 308.—Mexic.
Webbiana, *Lem. Cact. Nov. Gen. et Sp.* (1832) 45.—Mexic.
Wegeneri, *Ehrenb. in Bot. Zeit.* i. (1843) 738.—Mexic.
Wildiana, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 37.—Mexic.
Wildii, *Hort. Berol. ex A. Dietr. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 137.—Hab. ?
Winklerii, *Foerst. in Otto & Dietr. l. c.* xv. (1847) 50. —Mexic.
woburnensis, *Scheer, in Hook. Lond. Journ. Bot.* iv. (1845) 136.—Guatemala.
Wrightii, *Engelm. in Pacif. Rail. Rep.* iv. 27.—N. Mexic.
xanthotricha, *Scheidw. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 338.—Mexic.
Zegschwitzii, *Terscheck, Suppl. Cact. Verz.* 1.—Hab. ?
zephyranthoides. *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 41.—Mexic.
Zepnickii, *Ehrenb. in Bot. Zeit.* ii. (1844) 835.—Mexic.
Zuccariniana, Hort. ex Foerst. Handb. Cact. ed. I. 238 = macrantha.
Zuccariniana, *Mart. in Nov. Act. Nat. Cur.* xvi. i. (1832) 331. t. 20.—Mexic.

MAMPATA, 'Adans.' ex Steud. Nom. ed. II. ii. 98 (1841) = **Parinarium**, Juss. (Rosac.).

MANABAEA, R. Hedw. Gen. 368 (1806) = seq.

MANABEA, Aubl. Pl. Gui. i. 61. t. 23, 25 (1775) = **Aegiphila**, Jacq. (Verben.).
arborescens, Aubl. l. c. 64 = A. arborescens.
laevis, Aubl. l. c. 66 = A. laevis, *Willd.*
villosa, Aubl. l. c. 62 = A. villosa.

MANAELIA, 'Bowdich,' Madeira, ed. Germ. 381 (182–?), vide MANOELIA, Bowdich.

MANAGA, Aubl. Pl. Gui. Suppl. 2 (1775) (Quid ?).
guianensis, Aubl. l. c.—Guiana.

MANÇANILLA, Plum. ex Adans. Fam. ii. 354 (1763) = seq.

MANCINELLA, Tussac, Fl. Antill. iii. 21. t. 5 (1824) = **Hippomane**, Linn. (Euphorb.).
venenata, Tussac, l. c. = H. Mancinella.

MANCOA, Rafin. Fl. Tellur. iii. 56 (1836) = **Mohlana**, Mart. (Phytolac.).
secunda, Rafin. l. c. = Mohlana secunda.

MANCOA, Wedd. Chlor. And. i. t. 86 D (1857). *CRUCIFERAE*, Benth. & Hook. f. i. 86.
hispida, *Wedd. Chlor. And.* i. t. 86 D ; *et in Ann. Sc. Nat.* Sér. V. i. (1864) 285.—Boliv.
laevis, *Wedd. l. c.* 285.—Boliv.

MANDARUS, Rafin. Sylva Tellur. 122 (1838) = **Bauhinia**, Linn. (Legum.).
acuminata, Rafin. l. c. = B. acuminata.
divaricata, Rafin. l. c. = B. divaricata.
Pes-caprae, Rafin. l. c. = B. Pes-caprae.
rotundifolia, Rafin. l. c. = B. subrotundifolia.
Utimuta, Rafin. l. c. = B. velutina.

MANDELORNA, Steud. Syn. Pl. Gram. 359 (1855) = **Andropogon**, Linn. (Gramin.).
insignis, Steud. l. c. = A. squarrosus.

MANDEVILLA, Lindl. Bot. Reg. (1840) t. 7. *APOCYNACEAE*, Benth. & Hook. f. ii. 726.
AMBLYANTHERA, Muell. Arg. in Mart. Fl. Bras. vi. i. 141. t. 44, 45 (1860).
Andrieuxii, *Hemsl. Biol. Centr. Am. Bot.* ii. 316.—Mexic.
convolvulacea, *Hemsl. l. c.*—Mexic.
foliosa, *Hemsl. l. c.*—Mexic.
hispida, *Hemsl. l. c.*—Mexic.
Karwinskii, *Hemsl. l. c.*—Mexic.
oaxacana, *Hemsl. l. c.*—Mexic.
palustris, *Hemsl. l. c.* 317.—Costa Rica.
suaveolens, *Lindl. Bot. Reg.* (1840) t. 7.—Reg. Argent.
subcarnosa, *Benth. & Hook. f. Gen.* ii. 727.—Guian.
Vanheurckii, *Benth. & Hook. f. l. c.*—Bras.
veraguensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 317.—N. Granat.

MANDIOCA, Link, Handb. ii. 436 (1831) = **Manihot**, Tourn. (Euphorb.).
Aipi, Pohl, ex Link, l. c. = Manihot Aipi.
dulcis, Parodi, in Anal. Soc. Cient. Argent. iv. (1877) 127 = Manihot utilissima.
utilissima, Pohl, ex Link, Handb. ii. 436 = Manihot utilissima.

MANDIROLA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 468 = **Achimenes**, P. Br. (Gesner.).
ichthyostoma, Seem. ex Hanst. in Mart. Fl. Bras. viii. 348 = A. ichthyostoma.
lanata, Planch. & Linden, in Illustr. Hortic. iii. (1856) t. 80 = A. lanata.
multiflora, Decne. in Rev. Hortic. Sér. III. ii. (1848) 468 = A. multiflora.
Naegelia ×, Morr. Belg. Hortic. v. (1855) 294.
Seemannii, Lem. Illustr. Hort. iii. (1856) sub t. 80 = A. Scheerii.

MANDONIA, Hassk. in Flora, liv. (1871) 260 = **Tradescantia**, Linn. (Commelin.).
boliviana, Hassk. l. c. = T. ambigua.

MANDONIA, Sch. Bip. in Linnaea, xxxiii. (1664–65) 757 = **Hieracium**, Linn. (Compos.).
Pilosella, Sch. Bip. l. c. 758 = H. Pilosella.

MANDONIA, Wedd. in Bull. Soc. Bot. Fr. xi. (1864) 50. t. 1 = **Tridax**, Linn. (Compos.).
boliviensis, Wedd. l. c. 51.—Bolivia.

MANDRAGORA, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 125. *SOLANACEAE*, Benth. & Hook. f. ii. 900.
acaulis, Gaertn. Fruct. ii. 236 = officinarum.
autumnalis, *Bertol. Elench.* 6 ; *Spreng. Syst.* i. 699.—Reg. Mediterr.
caulescens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 242.—Reg. Himal.
microcarpa, Bertol. in Nov. Comm. Bonon. ii. (1836) 391. t. 25 = autumnalis.
neglecta, G. Don, ex Loud. Hort. Brit. 71 = officinarum ?
officinalis, Mill. Gard. Dict. ed. VIII. = officinarum.
officinalis, Moris, Stirp. Sard. Elench. fasc. i. 33 = autumnalis.
officinarum, Bertol. Comm. Mandr. 10. t. 2 = autumnalis.
officinarum, *Linn. Sp. Pl.* 181.—Reg. Mediterr.
praecox, Sweet, Brit. Flow. Gard. t. 198 = officinarum ?
vernalis, Bertol. in Giorn. Arcad. xxi. (1824) 191 [bis] = officinarum.

MANDUYTA, 'Comm.' ex Steud. Nom. ed. II. ii. 98 (1841) = **Samadera**, Gaertn. (Simarub.).

MANETTIA, Adans. Fam. ii. 242 (1763) = **Mesembryanthemum**, Dill. (Ficoid.).

MANETTIA, Crantz, Inst. i. 580 (1766) = **Selago**, Linn.
coccinea, Crantz, l. c. 581 = S. spuria.
corymbosa, Crantz, l. c. 580 = S. corymbosa.
dubia, Crantz, l. c. 581 = Agathelpis angustifolia.
lychnoidea, Crantz, l. c. = Lyperia fragrans.
pinastra, Crantz, l. c. 580 = Stilbe pinastra.
rapunculoides, Crantz, l. c. = S. spuria.
spuria, Crantz, l. c. 581 = S. spuria.
tomentosa, Crantz, l. c. = Manulea tomentosa.

MANETTIA, Mutis, ex Linn. Mant. ii. 558 (1771). *RUBIACEAE*, Benth. & Hook. f. ii. 37.
CONOTRICHIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 277. t. 24. f. 1.
ENDOLASIA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 579.
GUAGNEBINA, Vell. Fl. Flum. 45 (1825); Ic. i. tt. 115–121 (1827).
LYGISTUM, P. Br. Hist. Jamaic. 142. t. 3 (1756).
NACIBEA, Aubl. Pl. Gui. i. 95. t. 37 (1775).
VANESSA, Rafin. Fl. Tellur. iii. 57 (1836).
acutiflora, Roem. & Schult. Syst. ii. 203 = seq.
acutifolia, *Ruiz & Pav. Fl. Per.* i. 58. t. 89.—Peruv.
albiflora, *Schott, in DC. Prod.* iv. 364.—Bras.
asperula, Benth. in Linnaea, xxiii. (1850) 444 = cordifolia.
attenuata, Nees & Mart. in Nov. Act. Nat. Cur. xii. i. (1824) 14 = cordifolia.
auratifolia, *Silva Manso, Enum. Subst. Bras.* 25.—Bras.
barbata, *Oerst. in Kjoeb. Vedensk. Meddel.* (1852) 47.—Costa Rica.
bicolor, Paxt. Mag. Bot. x. (1843) 27 = luteo-rubra.
calycosa, *Griseb. Fl. Brit. W. Ind.* 330.—Ind. occ.
ciliata, *Cham. & Schlecht. in Linnaea*, iv. (1829) 176.—Bras.
coccinea, Griseb. Fl. Brit. W. Ind. 329 = uniflora.
coccinea, Willd. Sp. Pl. i. 625 = M. Lygistum.
confertiflora, Benth. in Linnaea, xxiii. (1850) 443 = cordifolia.
cordifolia, *Mart. Spec. Mat. Med. Bras.* i. 19. t. 7.—Bras.
cuspidata, Bert. ex Spreng. Syst. i. 415 = uniflora.
filicaulis, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 281 = luteo-rubra.
fimbriata, *Cham. & Schlecht. in Linnaea*, iv. (1829) 173.—Bras.
Gaertneri, *DC. Prod.* iv. 364.—Hab. ?
glabra, Cham. & Schlecht. in Linnaea, iv. (1829), 159 = cordifolia.
glabra, Steud. Nom. ed. II. ii. 98 = Gaertneri.
glandulosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 24.—Chili.
gracilis, *Cham. & Schlecht. in Linnaea*, iv. (1829) 169.—Bras.
gracilis, Steud. Nom. ed. II. ii. 98 = Gaertneri.
grandiflora, Miq. in Linnaea, xxiii. (1850) 803 = cordifolia.
havanensis, H. B. & K. Nov. Gen. et Sp. iii. 388 = uniflora.
hirtella, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 238.—Mexic.
hispida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 24. t. 228.—Peruv.
lanceolata, Vahl, Symb. Bot. i. 12 = Mussaenda luteola.
leianthiflora, Griseb. in Goett. Abh. xix. (1874) 159 = cordifolia.
luteo-rubra, *Benth. in Linnaea*, xxiii. (1850) 445.—Bras.
lygista, *St. Lag. in Ann. Soc. Bot. Lyon*, vii. (1880) 130 = M. Lygistum.
lygistoides, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 505 = M. Lygistum.
Lygistum, Spreng. Syst. i. 415 = uniflora.
Lygistum, *Sw. Prod. Veg. Ind. Occ.* 37.—Am. trop.
micans, Poepp. & Endl. Nov. Gen. et Sp. iii. 24 = cordifolia.
miniata, Lem. in Fl. des Serres, iv. (1848) t. 317 = cordifolia.
multiflora, Cham. in Linnaea, ix. (1834) 243 = albiflora.

MANETTIA :—
mutabilis, Pers. Syn. i. 134 = racemosa.
panamensis, Duchass. & Walp. in Linnnaea, xxiii. (1850) 733 = uniflora.
paniculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 24.—Peruv.
picta, Willd. Sp. Pl. i. 625 = M. Lygistum.
Pseudodiodia, *Cham. & Schlecht. in Linnaea*, iv. (1829) 174.—Bras.
pubescens, Cham. & Schlecht. l. c. 170 = luteo-rubra.
racemosa, *Ruiz & Pav. Fl. Per.* i. 58. t. 89.—Peruv.
reclinata, *Linn. Mant.* ii. 553.—Mexic.
rosea, Pohl, ex DC. Prod. iv. 364 = fimbriata.
secundiflora, *DC. l. c.*—Ins. S. Doming.
serrata, *Spreng. in Schult. Mant.* iii. 147.—Ins. S. Doming.
splendens, *Regel, Cat. Pl. Hort. Aksakov.* 89.—Bras.
tenuiflora, Willd. ex Steud. Nom. ed. II. ii. 98 = uniflora.
tenuifolia, Schult. Mant. iii. 145 = uniflora.
umbellata, *Ruiz & Pav. Fl. Per.* i. 58.—Peruv.
uniflora, *H. B. & K. Nov. Gen. et Sp.* iii. 387.—Am. trop.
villosa, Cham. & Schlecht. in Linnaea, iv. (1829) 172 = pubescens.
zimapanica, *Hemsl. Diagn. Pl. Nov.* 30.—Mexic.

MANFREDA, Salisb. Gen. Pl. Fragm. 78 (1866) = **Agave**, Linn. (Amaryll.).
virginica, Salisb. l. c. = A. virginica.

MANGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = **Mangifera**, Linn. &c. (Anacard.).
acida, Noronha, l. c. = Bouea burmanica.
deleteria, Noronha, ex Hassk. in Hoev. & De Vriese, Tijdschr. xi. (1844) 224 = Gluta Benghas.

MANGAS, Adans. Fam. ii. 345 (1763) = **Mangifera**, Linn. (Anacard.).

MANGHAS, Burm. Thes. Zeyl. 150 (1737) = **Cerbera**, Linn. (Apocyn.).

MANGIFERA, Linn. Fl. Zeyl. 211 (1747). *ANACARDIACEAE*, Benth. & Hook. f. i. 420.
MANGAS, Adans. Fam. ii. 345 (1763).
africana, *Oliver, Fl. Trop. Afr.* i. 443.—Afr. trop.
altissima, *Blanco, Fl. Filip.* ed. I. 181.—Ins. Philipp.
Amba, Forsk. Fl. Aegypt. Arab. 205 = indica.
anisodora, Blanco, Fl. Filip. ed. II. 129 = fragrans.
axillaris, Desr. in Lam. Encyc. iii. 697 = Buchanania Lanzan.
caesia, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 441.—Malaya.
caloneura, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 66.—Burma.
cochinchinensis, *Engl. in DC. Monog. Phan.* iv. 205.—Cochinch.
domestica, Gaertn. Fruct. ii. 95. t. 100 = indica.
foetida, Blume, Bijdr. 1153 = caesia.
foetida, *Lour. Fl. Cochinch.* 160.—Malaya.
fragrans, *Maingay, in Hook. f. Fl. Brit. Ind.* ii. 18 = Malaya.
gabonensis, Aubry-Lecomte, ex O'Rorke, in Journ. Pharm. et Chim. Sér. III. xxxi. (1857) 275 = Irvingia Barteri.
Gandaria, *Roxb. Hort. Beng.* 18, nomen.—Ins. Molucc.
Gedebi, *Miq. Fl. Ind. Bat. Suppl.* 522.—Sumatra.
gladiata, Boj. Hort. Maurit. 73 = indica.
glauca, *Blume, Bijdr.* 1158.—Java.
glauca, Rottb. in Nye Saml. Danske Vidensk. Selsk. Skr. ii. (1783) 534. t. 4. f. 1 = Elaeodendron glaucum.
glauca, Span. in Linnaea, xv. (1841) 188 = timoriensis.
gracilipes, *Hook. f. Fl. Brit. Ind.* ii. 16.—Malacca.
Griffithii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 168.—Malacca.
Horsfieldii, Miq. Fl. Ind. Bat. i. ii. 632 = foetida.
indica, Blume, Bijdr. 1157 = foetida.
indica, *Linn. Sp. Pl.* 200.—Ind. or.; Malaya.
indica, Thw. Enum. Pl. Zeyl. 75 = zeylanica.
indica, Wall. Cat. n. 8487 I = sylvatica.
Kemanga, Blume, Mus. Bot. Lugd. Bat. i. 202 = caesia.
Kukula, Blume, l, c. 192 = indica.
lagenifera, Griff. Notul. iv. 414; Ic. Pl. Asiat. t. 567.—Malacca.

MANGIFERA :—
Langang, *Miq. Fl. Ind. Bat. Suppl.* 521.—Sumatra.
laurina, *Blume, Mus. Bot. Lugd. Bat.* i. 195.—Malaya.
laxiflora, *Desr. in Lam. Encyc.* iii. 97.—Ins. Maurit.
Leschenaultii, March. Rev. Anacard. 189 = foetida.
Linnaei, Korth. ex Hassk. Cat. Hort. Bog. Alt. 245 = indica.
longipes, *Griff. Notul.* iv. 419.—Malaya.
macrocarpa, *Blume, Bijdr.* 1158.—Java.
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 17.—Malacca.
maritima, Lechaume, in Rev. Hortic. (1870–71) 369 = indica.
membranacea, *Blume, Mus. Bot. Lugd. Bat.* i. 195.—Ins. Molucc.
microphylla, *Griff. ex Hook. f. Fl. Brit. Ind.* ii. 17.—Malacca.
minor, *Blume, Mus. Bot. Lugd. Bat.* i. 198.—Ins. Molucc.
montana, Heyne, ex Wight & Arn. Prod. 170 = indica.
mucronulata, *Blume, Mus. Bot. Lugd. Bat.* i. 201.—N. Guian.
oblongifolia, *Hook. f. Fl. Brit. Ind.* ii. 16.—Malacca.
odorata, *Griff. Notul.* iv. 417.—Malaya.
oppositifolia, Roxb. Hort. Beng. 18; Fl. Ind. i. 640 = Bouea burmanica.
Parish, Miq. Fl. Ind. Bat. i. ii. 630 = longipes.
pentandra, *Hook. f. Fl. Brit. Ind.* ii. 14.—Malacca.
pinnata, Desr. in Lam. Encyc. iii. 697 = Sorindeia madagascariensis.
pinnata, Linn. f. Suppl. 156 = Spondias Mangifera.
polycarpa, *Griff. Notul.* iv. 416. t. 167. f. 2.—Malacca.
quadrifida, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 440.—Malaya.
racemosa, Boj. Hort. Maurit. 73 = indica.
racemosa, Lam. Illustr. ii. 113 = Holigarna Arnottiana ?
rigida, *Blume, Mus. Bot. Lugd. Bat.* i. 200.—Sumatra.
rostrata, Blanco, Fl. Filip. ed II. 129 = sylvatica.
rubra, Boj. Hort. Maurit. 76 = indica.
sativa, Roem. & Schult. Syst. i. 37 in obs. = indica.
sclerophylla, *Hook. f. Fl. Brit. Ind.* ii. 15.—Malacca.
spathulaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 200.—Borneo.
similis, *Blume, l. c.*—Malaya.
sumatrana, Miq. Fl. Ind. Bat. i. ii. 630 = longipes.
superba, *Hook. f. Fl. Brit. Ind.* ii. 19.—Malacca.
sylvatica, *Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 644.—Reg. Himal.
Taipon, *Ham. ex Miq. Fl. Ind. Bat.* i. ii. 631.—N. Guinea.
timorensis, *Blume, Mus. Bot. Lugd. Bat.* i. 199.—Ins. Timor.
viridis, Boj. Hort. Maurit. 73 = indica.
zeylanica, *Hook. f. Fl. Brit. Ind.* ii. 16.—Zeylan.

MANGIUM, Rumpf, ex Scop. Introd. 218 (1777) = **Rhizophora**, Linn.

MANGLE, Adans. Fam. ii. 445 (1763) = **Rhizophora**, Linn.

MANGLES, Plum. ex DC. Prod. iii. 32 (1828) = praec.

MANGLESIA, Endl. Nov. Stirp. Dec. 25 (1839) = **Grevillea**, R. Br. (Proteac.).
cuneata, Endl. l. c. = G. glabrata.
glabrata, Lindl. Swan River App. 37 = G. glabrata.
Lawrenceana, Hort. ex Meissn. in DC. Prod. xiv. 392 = G. vestita.
tridentifera, Endl. Nov. Stirp. Dec. 25 = G. tridentifera.
vestita, Endl. l. c. 26 = G. vestita.

MANGLESIA, Lindl. Swan River App. t. 3 (1839) = **Beaufortia**, R. Br. (Myrtac.).
purpurea, Lindl. l. c. = B. purpurea.

MANGLIETIA, Blume, in Verh. Batav. Gen. ix. (1823) 149; et Bijdr. 8 (1825). *MAGNOLIACEAE*, Benth. & Hook. f. i. 19.
Candollei, Wall. Cat. n. 6497 = Talauma mutabilis.
Caveana, *Hook. f. & Thom. Fl. Ind.* i. 76.—Reg. Himal.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 72.—Ins. Celebes.
glauca, *Blume, in Verh. Batav. Gen.* ix. (1823) 150; *et Bijdr.* 8.—Java.
insignis, *Blume, Fl. Jav. Magnol.* 23.—Reg. Himal.

MANGLIETIA:—
Macklottii, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 97.—Sumatra.
Oortii, Korth. l. c. = Talauma glauca.
Sebassa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 71.—Sumatra.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 367.—Sumatra.

MANGLILLA, Juss. Gen. 151 (1789) = **Myrsine**, Linn.
brasiliensis, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 86 = Myrsine brasiliensis.
canariensis, Roem. & Schult. Syst. iv. 505 = Myrsine canariensis.
Caporoca, A. St. Hil. ex Miq. in Mart. Fl. Bras. x. 309 = Myrsine Gardneriana.
dentata, Roem. & Schult. Syst. iv. 506 = Myrsine dentata.
dependens, Roem. & Schult. l. c. = Myrsine dependens.
ferruginea, Roem. & Schult. l. c. = Myrsine ferruginea.
Jussieui, Pers. Syn. i. 237 = Myrsine Manglilla.
latifolia, Roem. & Schult. Syst. iv. 505 = Myrsine latifolia.
melanophleos, Pers. Syn. i. 287 = Myrsine melanophleos.
Milleriana, Pers. l. c. = Myrsine mitis.
oblonga, Roem. & Schult. Syst. iv. 506 = Myrsine Manglilla.
pellucida, Roem. & Schult. l. c. = Myrsine pellucida.
peruviana, J. F. Gmel. Syst. 398 = Myrsine Manglilla.
racemosa, Roem. & Schult. Syst. iv. 803 = Myrsine racemosa.
venosissima, Roem. & Schult. l. c. 506 = Myrsine venosissima.
venulosa, Roem. & Schult. l. c. 505 = Myrsine melanophleos.

MANGONIA, Schott, in Oestr. Bot. Wochenbl. (1857) 77. *AROIODEAE*, Benth. & Hook. f. iii. 988.
Tweedieana, *Schott, in Oestr. Bot. Wochenbl.* (1857) 77.—Bras.

MANGOSTANA, Rumph. ex Gaertn. Fruct. ii. 105. t. 105. (1791) = **Garcinia**, Linn. (Guttif.).
Cambogia, Gaertn. l. c. 106 = G. Cambogia.
Garcinia, Gaertn. l. c. 105 = G. Mangostana.
Morella, Gaertn. l. c. 106 = G. Morella.

MANICARIA, Gaertn. Fruct. ii. 468. t. 176 (1791). *PALMAE*, Benth. & Hook. f. iii. 919.
PILOPHORA, Jacq. Fragm. 32. t. 35 (1800).
Plukenetii, Griseb. & H. Wendl. in Griseb. Fl. Brit. W. Ind. 518 = saccifera.
saccifera, *Gaertn. Fruct.* ii. 468. t. 176.—Guian.

MANIHOT, Tourn. ex Adans. Fam. ii. 356 (1763). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 306.
JANIPHA, H. B. K. Nov. Gen. et Sp. ii. 106 (1817).
MANDIOCA, Link, Handb. ii. 436 (1831).
acuminatissima, *Muell. in Arg. Mart. Fl. Bras.* xi. II. 455.—Bras.
aesculifolia, *Pohl, Pl. Bras. Ic.* i. 55.—Mexic.
Aipi, Pohl, l. c. 29 = palmata.
amaroleitensis, *Baill. Adansonia*, iv. (1863–64) 281.—Bras.
angustifolia, Pohl, ex Steud. Nom. ed. II. i. 799; ii. 99 = *Jatropha angustifolia* (nom. delend.).
angustifrons, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 461.—Bras.
angustiloba, *Muell. Arg. in DC. Prod.* xv. II. 1073.—Peruv.
anisophylla, *Muell. Arg. in Journ. Bot.* xii. (1874) 230.—Reg. Argent.
anomala, *Pohl, Pl. Bras. Ic.* i. 27. t. 21.—Bras.
arcuata, Pohl, l. c. 42 = violacea.
attenuata, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 443.—Bras.
Aypi, Spruce, in Journ. Linn. Soc. v. (1861) 10 = palmata.
brachyloba, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 451.—Bras.
Burchelli, *Muell. Arg. l. c.* 457.—Bras.
cajaniformis, Pohl, Pl. Bras. Ic. i. 45 = tripartita.
caricaefolia, *Pohl, l. c.* 40.—Bras.
carthaginensis, *Muell. Arg. in DC. Prod.* xv. II. 1073.—Am. trop.
cecropiaefolia, Pohl, Pl. Bras. Ic. i. 49 = violacea.
cleomaefolia, Pohl, l. c. 51 = tripartita.

MANIHOT:—
coerulea, Steud. Nom. ed. II. i. 799; ii. 99, sphalm. = seq.
coerulescens, *Pohl, Pl. Bras. Ic.* i. 56.—Bras.
conulifera, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 474.—Bras.
crotalariaeformis, *Pohl, Pl. Bras. Ic.* i. 24. t. 18.—Bras.
cujabensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 207.—Bras.
Curcas, Crantz, Inst. i. 167 = Jatropha Curcas.
dalechampiaeformis, Pohl, Pl. Bras. Ic. i. 51 = tripartita.
diffusa, Pohl, l. c. 55 = palmata.
digitata, Sweet, Hort. Brit. ed. II. 458 = *Jatropha Fischeri* (nom. delend.).
digitiformis, *Pohl, Pl. Bras. Ic.* i. 36.—Bras.
divergens, Pohl, l. c. 41 = violacea.
diversifolia, Sweet, Hort. Brit. ed. II. 458, nomen = *Jatropha diversifolia* (nom. delend.).
edule, A. Rich. Fl. Cub. Fanerog. ii. 208 = utilissima.
elegans, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 485.—Bras.
esculenta, Crantz, Inst. i. 167 = utilissima.
flabellifolia, Pohl, Pl. Bras. Ic. i. 35. t. 25 = palmata.
foetida, *Pohl, l. c.* 55.—Mexic.
Glaziovii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 446.—Bras.
gossypifolia, Crantz, Inst. i. 167 = Jatropha gossypifolia.
gracilis, *Pohl, Pl. Bras. Ic.* i. 23. t. 16.—Bras.
Grahami, Hook. Ic. Pl. t. 530 = palmata.
grandiflora, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 471.—Bras.
guyanensis, *Klotzsch, in Schomb. Faun. et Fl. Gui.* 1186.—Guian.
hemitrichandra, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 454.—Bras.
herbacea, Crantz, Inst. i. 167 = Jatropha urens.
heterophylla, *Pohl, Pl. Bras. Ic.* i. 39. t. 31.—Bras.
Hilariana, *Baill. Adansonia*, iv. (1863–64) 282.—Bras.
humilis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 448.—Bras.
inflata, *Muell. Arg. l. c.* 450.—Bras.
intercedens, *Muell. Arg. l. c.* 483.—Bras.
jacobinensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 205.—Bras.
Janipha, Pohl, Pl. Bras. Ic. i. 55 = carthaginensis.
janiphoides, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 480.—Bras.
laciniosa, *Pohl, Pl. Bras. Ic.* i. 54.—Bras.
lagoënsis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 475.—Bras.
Langsdorffii, *Muell. Arg. l. c.* 455.—Bras.
linearifolia, *Muell. Arg. in Flora*, lv. (1872) 43.—Peruv.
Loeflingii, Muell. Arg. in Mart. Fl. Bras. xi. II. 460 = palmata.
longepetiolata, *Pohl, Pl. Bras. Ic.* i. 25. t. 19.—Bras.
Loureirii, *Pohl, l. c.* 55.—Cochinch.
melanobasis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 206.—Guian.
microcarpa, *Muell. Arg. in Flora*, lv. (1872) 42.—Am. centr.
moluccana, Crantz, Inst. i. 167 = Aleurites triloba.
multifida, Crantz, l. c. = Jatropha multifida.
nana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 448.—Bras.
occidentalis, *Muell. Arg. l. c.* 468.—Bras.
orbicularis, *Pohl, Pl. Bras. Ic.* i. 20. t. 12.—Bras.
palmata, *Muell. Arg. in DC. Prod.* xv. II. 1062.—Bras.
paraënsis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 470.—Bras.
pardina, *Muell. Arg. l. c.* 484.—Bras.
paviaefolia, *Pohl, Pl. Bras. Ic.* i. 52.—Bras.
Pavoniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 205.—Peruv.
pedicellaris, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 453.—Bras.
peltata, *Pohl, Pl. Bras. Ic.* i. 18. t. 13.—Bras.
pentaphylla, *Pohl, l. c.* 53.—Bras.
peruviana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 206.—Peruv.
pilosa, *Pohl, Pl. Bras. Ic.* i. 55.—Bras.
Pohliana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 464.—Bras.
Pohlii, *Wawra, in Flora*, xlvii. (1864) 252.—Bras.

MANIHOT:—
porrecta, Pohl, Pl. Bras. Ic. i. 46 = tripartita.
procumbens, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 206.—Bras.
pronifolia, *Pohl, Pl. Bras. Ic.* i. 24. t. 17.—Bras.
pruinosa, *Pohl, l. c.* 28. t. 22.—Bras.
pubescens, *Pohl, l. c.* 50. t. 43.—Bras.
purpureo-costata, *Pohl, l. c.* 19.—Bras.
pusilla, *Pohl, l. c.* 36.—Bras.
quinquefolia, *Pohl, l. c.* 56.—Bras.
quinqueformis, Steud. Nom. ed. II. i. 800, in syn. = praec.?
quinqueloba, *Pohl, Pl. Bras. Ic.* i. 21.—Bras.
reniformis, *Pohl, l. c.* 56.—Bras.
rhomboidea, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 205.—Mexic.
Riedeliana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 443.—Bras.
rigidula, *Muell. Arg. l. c.* 474.—Bras.
sagittato-partita, *Pohl, Pl. Bras. Ic.* i. 22. t. 15.—Bras.
salicifolia, *Pohl, l. c.* 18.—Bras.
sinuata, *Pohl, l. c.*—Bras.
sparsifolia, *Pohl, l. c.* 26.—Bras.
speciosa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 470.—Bras.
spinosissima, Steud. Nom. ed. II. ii. 99 = Jatropha vitifolia.
stricta, *Baill. Adansonia*, iv. (1863–64) 282.—Bras.
subquinqueloba, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 446.—Bras.
tenella, *Muell. Arg. l. c.* 484.—Bras.
tenerrima, Pohl, Pl. Bras. Ic. i. 39. t. 30 = gracilis.
tenuifolia, Pohl, l. c. 38. t. 29 = gracilis.
tomentella, Pohl, l. c. 45 = tripartita.
tomentosa, *Pohl, l. c.* 50.—Bras.
tripartita, *Muell. Arg. in DC. Prod.* xv. II. 1068.—Bras.
triphylla, *Pohl, Pl. Bras. Ic.* i. 37.—Bras.
tristis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 449.—Bras.
Tweedieana, *Muell. Arg. l. c.* 450.—Bras.
urens, Crantz, Inst. i. 167 = Jatropha urens.
utilissima, *Pohl, Pl. Bras. Ic.* i. 32. t. 24.—Bras.
varians, *Pohl, l. c.* 53.—Bras.
violacea, *Pohl, l. c.* 43.—Bras.
Warmingii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 481.—Bras.
Weddelliana, *Baill. Adansonia*, iv. (1863–64) 281.—Bras.

MANILKARA, Adans. Fam. ii. 166 (1763) = **Mimusops**, Linn. (Sapot.).

MANILTOA, Scheff. in Ann. Jard. Buitenz. i. (1876) 20. *LEGUMINOSAE*.
grandiflora, *Scheff. l. c.*—N. Guin.

MANISURIS, Linn. Mant. ii. 164 (1767) = **Rottboellia**, Linn. f. (Gramin.).
Myurus, Linn. l. c. 300 = R. Myurus.

MANISURIS, Sw. Prod. Veg. Ind. Occ. 25 (1788), partim; Fl. Ind. Occ. i. 186 (1797). *GRAMINEAE*, Benth. & Hook. f. iii. 1130.
granularis, *Sw. ll. cc.*—Reg. trop. et subtrop.
polystachya, Beauv. Fl. Owar. i. 24. t. 14 = granularis.

MANITIA, Giseke, Prael. Linn. 209 (1792) = **Globba**, Linn. (Scitam.).
aurea, Giseke, l. c. 251 = G. versicolor.

MANLILIA, Salisb. Gen. Pl. Fragm. 17 (1866) = **Polyxena**, Kunth (Liliac.).
pygmaea, Salisb. l. c. = P. pygmaea.

MANNA, D. Don, Prod. Fl. Nep. 246 (1825) = **Alhagi**, Tourn. (Legum.).
caspica, D. Don, l. c. 247 = A. camelorum.
hebraica, D. Don, l. c. = A. maurorum.
nepalensis, D. Don, l. c. = A. maurorum.

MANNAPHORUS, Rafin. in Am. Monthly Mag. (1818) 175 = **Fraxinus**, Tourn. (Oleac.).

MANNARIA, Heist. Syst. 12 (1748) = **Fraxinus**? Tourn. (Oleac.).

MANNIA, Hook. f. in Benth. & Hook. f. Gen. i. 309 (1862). *SIMARUBEAE*, Benth. & Hook. f. l. c.
africana, *Hook. f. l. c.*—Afr. trop.

MANNIELLA, Reichb. f. Otia Bot. Hamb. 109 (1881). *ORCHIDEAE*, Benth. & Hook. f. iii. 605.
Gustavi, *Reichb. f. l. c.*—Afr. trop.

MANNIOPHYTON, Muell. Arg. in Flora. xlvii. (1864) 530. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 297.
africanus, *Muell. Arg. in Flora*, xlvii. (1864) 531.—Afr. trop.
fulvum, *Muell. Arg. in Journ. Bot.* ii. (1864) 332.—Afr. trop.

MANNOPAPPUS, Sch. Bip. in Flora, xxvii. (1844) 677 = **Helichrysum**, Vaill. (Compos.).
anomalus, Sch. Bip. l. c. = H. anomalum.

MANOELIA, Bowdich, Madeira, 246 (1825) = (Gen. dub. an Primulac. ?).
pallida, Bowdich, l. c.—Ins. Madeira.

MANOELLIA, Reichb. Consp. 212* (1828) = praec.

MANOTES, Soland. ex Planch. in Linnaea, xxiii. (1850) 438. *CONNARACEAE*, Benth. & Hook. f. i. 433.
expansa, *Soland. ex Planch. in Linnaea*, xxiii. (1850) 439.—Afr. trop.
Griffoniana, *Baill. Adansoniana*, vii. (1866–67) 244.—Afr. trop.
longiflora, *Baker, in Oliver, Fl. Trop. Afr.* i. 460.—Afr. trop.
Palisotii, Planch. in Linnaea, xxiii. (1850) 439 = Rourea Palisotii.

MANOTHRIX, Miers, Apocyn. S. Am. 28 (1878). *APOCYNEAE*.
nodosa, *Miers, l. c.* 30.—Bras.
valida, *Miers, l. c.* 29.—Bras.

MANSANA, J. F. Gmel. Syst. 580 (1791) = **Zizyphus**, Adans. (Rhamnac.).
arborea, J. F. Gmel. l. c. = Z. Jujuba.

MANSOA, DC. in Bibl. Univ. Genèv. xvii. (1838) 128 [Rev. Bign. 12] = **Cuspidaria**, DC. (Bignon.).
hirsuta, DC. Prod. ix. 182.—Bras.
laevis, DC. l. c.—Bras.

MANTEIA, Rafin. Sylva Tellur. 161 (1838) = **Rubus**, Tourn. (Rosac.).
acaulis, Rafin. l. c. = R. arcticus.
stellatus, Rafin. l. c. = R. stellatus.

MANTISALCA, Cass. in Bull. Soc. Philom. (1818) 142 = **Centaurea**, Linn. Compos.).
elegans, Cass. in Dict. Sc. Nat. xxix. 81 = C. salmantica.

MANTISIA, Sims, Bot. Mag. t. 1320 (1810). *SCITAMINEAE*, Benth. & Hook. f. iii. 639.
saltatoria, *Sims, Bot. Mag.* t. 1320.—Ind. or.
spathulata, *Schult. Mant.* i. 49.—Ind. or.

MANTODDA, Adans. Fam. ii. 508 (1763) = **Aeschynomene**, Linn. (Legumin.).

MANUELIA, Pritz. Ic. Ind. ii. 180 (1855) sphalm. = seq.

MANULEA, Linn. Mant. i. 12 (1767). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 946.
LYCHNIDEA, Burm. Rar. Afr. Pl. 138 (1737).
NEMIA, Berg. Desc. Pl. Cap. 160 (1767).
aequipetala, Sternb. ex Steud. Nom. ed. I. 508 = Chaenostoma foetidum.
aethiopica, Thunb. Prod. Pl. Cap. 101 = Polycarena aurea.
alternifolia, Desf. ex Pers. Syn. ii. 148 = Chaenostoma foetidum.
altissima, *Linn. f. Suppl.* 285.—Afr. austr.
androsacea, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 381.—Afr. austr.
angustifolia, Link & Otto, Ic. Pl. Sel. 47. t. 20 = rubra.
antirrhinoides, *Linn. f. Suppl.* 286.—Afr. austr.
argentea, Linn. f. l. c. = Lyperia argentea.
aurantiaca, *Jarosz, Pl. Nov. Cap.* 18.—Afr. austr.
bellidifolia, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
capensis, Thunb. Prod. Pl. Cap. 101 = Polycarena capensis.
capillaris, Linn. f. Suppl. 285 = Polycarena capillaris.
capitata, Linn. f. l. c. 286 = Phyllopodium capitatum.
Cephalotes, *Thunb. Prod. Pl. Cap.* 101.—Afr. austr.
Cheiranthus, *Linn. Mant.* i. 88.—Afr. austr.
coerulea, Linn. f. l. c. 285 = Sphenandra viscosa.
cordata, Thunb. Prod. Pl. Cap. 102 = Chaenostoma cordatum.
corymbosa, *Linn. f. Suppl.* 286.—Afr. austr.
crassifolia, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
crystallina, Weinm. in Syll. Ratisb. i. (1824) 226 = Lyperia violacea.
cuneifolia, Linn. f. Suppl. 285 = Phyllopodium cuneifolium.
densiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
divaricata, Thunb. Prod. Pl. Cap. 101 = Zaluzianskya divaricata.
exaltata, *Banks, ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 384.—Afr. austr.
foetida, Pers. Syn. ii. 148 = Chaenostoma foetidum.
gariepina, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 384.—Afr. austr.
heterophylla, Linn. f. Suppl. 285 = Phyllopodium heterophyllum.
hirta, Gaertn. Fruct. i. 258; Desr. in Lam. Encyc. iii. 707 = rubra.
hirta, *Thunb. Prod. Pl. Cap.* 101.—Afr. austr.
hispida, Thunb. l. c. 102 = Chaenostoma hispidum.
incana, *Thunb. l. c.* 101.—Afr. austr.
indiana, Lour. Fl. Cochinch. 386 = Adenosma capitatum ?
indica, Willd. in Lour. l. c. ed. II. ii. 469 = pr ec.
integrifolia, Linn. f. Suppl. 285 = Chaenostoma integrifolium.
juncea, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
leiostachys, *Benth. l. c.* 383.—Afr. austr.
lichnidea, Desr. in Lam. Encyc. iii. 707 = Lyperia fragrans.
linifolia, Thunb. Prod. Pl. Cap. 100 = Chaenostoma linifolium.
longiflora, *Jarosz, Pl. Nov. Cap.* 18.—Afr. austr.
longifolia, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
lychnoides, Steud. Nom. ed. I. 508, vide *lichnidea*.
microphylla, Linn. f. Suppl. 285 = Lyperia microphylla.
natalensis, Bernh. in Flora, xxvii. (1844) 835 = parviflora.
nervosa, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 381.—Afr. austr.
obovata, *Benth. l. c.* 383.—Afr. austr.
oppositiflora, Vent. Jard. Malm. t. 15 = Chaenostoma hispidum.
oppositifolia, Steud. Nom. ed. II. ii. 99 = praec.
paniculata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 383.—Afr. austr.
parviflora, *Benth. l. c.* 383.—Afr. austr.
pedunculata, Pers. Syn. ii. 184 = Lyperia pedunculata.
pinnatifida, Linn. f. Suppl. 286 = Lyperia pinnatifida.
plantaginea, Thunb. Prod. Pl. Cap. 101 = seq.
Plantaginis, Linn. f. Suppl. 286 = Polycarena plantaginea.
pusilla, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 384.—Afr. austr.
revoluta, Thunb. Prod. Pl. Cap. 100 = Chaenostoma revolutum.
rhynchantha, Link, Enum. Hort. Berol. ii. 142 = M. Cheiranthus.
rigida, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 382.—Afr. austr.
rotata, Desr. in Lam. Encyc. iii. 706 = Sphenandra viscosa.
rubra, *Linn. f. Suppl.* 286.—Afr. austr.
satureioides, Desr. in Lam. Encyc. iii. 705 = Chaenostoma revolutum ?
scabra, *Wendl. ex Steud. Nom.* ed. II. 99, nomen.—Afr. austr.
silenioides, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 381.—Afr. austr.
Sopubia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 88 = Sopubia trifida.
stellata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 384.—Afr. austr.
Thunbergii, *G. Don, Gen. Syst.* iv. 596.—Afr. austr.
thysiflora, *Linn. f. Suppl.* 286.—Afr. austr.
tomentosa, Benth. in DC. Prod. x. 367, sphalm. = Thunbergii.
tomentosa, *Murr. Syst.* ed. XIII. 476.—Afr. austr.
turrita, *Banks, ex Benth. in Hook. Comp. Bot. Mag.* i. (1835) 383.—Afr. austr.
uncinata, *Desr. in Lam. Encyc.* iii. 706.—Afr. austr.
villosa, Pers. Syn. ii. 148 = Polycarena capensis.
violacea, Link, Enum. Hort. Berol. ii. 142 = Lyperia violacea.
virgata, *Thunb. Prod. Pl. Cap.* 101.—Afr. austr.
viscosa, Willd. Enum. Hort. Berol. 653 = Sphenandra viscosa.

MANUNGALA, Blanco, Fl. Filip. ed. I. 306 (1837) = **Samadera**, Gaertn. (Simarub.).
pendula, Blanco, l. c. = S. indica.

MAOUTIA, Montr. in Mém. Acad. Lyon, x. (1860) 241 = **Oxera**, Labill. (Verbenac.).
neriifolia, Montr. l. c.—N. Caled.

MAOUTIA, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 193. *URTICACEAE*, Benth. & Hook. f. iii. 391.
LECANOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 193. t. 12 (1856).
ambigua, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 483.—Ins. Amboina.
aspera, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 194 = odontophylla.
australis, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 480.—Ins. Pacif.
diversifolia, Wedd. in DC. Prod. xvi. 1. 235^{32} = rugosa.
odontophylla, *Miq. Fl. Ind. Bat.* i. II. 273.—Java.
platistigma, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 194.—Ins. Philipp.
Puya, *Wedd. l. c.* 193.—Reg. Himal.; Burma.
reticulata, *Wedd. in DC. Prod.* xvi. 1. 235^{31}.—Ins. Philipp.
rigida, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 194 = Boehmeria rigida.
rugosa, *Wedd. l. c.*—Java.
setosa, *Wedd. l. c.*—Ins. Philipp.
Zollingeriana, Wedd. l. c. = rugosa.

MAPA, Vell. Fl. Flum. 59 (1825); i. t. 153 = **Petiveria**, Linn. (Phytolacc.).
graveolens, Vell. ll. cc. = P. alliacea.

MAPANIA, Aubl. Pl. Gui. i. 47. t. 17 (1775). *CYPERACEAE*, Benth. & Hook. f. iii. 1055.
CEPHALOSCIRPUS, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) II. 83.
HALOSTEMMA, Benth. & Hook. f. Gen. iii. 1056 (1883).
LEPISTACHYA, Zipp. ex Miq. Illustr. Fl. Ind. Arch. 61 (1871).
PANDANOPHYLLUM, Hassk. Cat. Hort. Bog. Alt. 297 (1844).
africana, *Boeck. in Linnaea*, xxxvii. (1871–73) 137.—Afr. trop.
bancana, Benth. & Hook. f. Gen. iii. 1055 = Thoracostachyum bancanum.
humilis, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 309.—Malaya.
hypolytroides, F. Muell. ex Benth. Fl. Austral. vii. 341 (Thoracostachyi sp.).—Austral.
immersa, *Trimen, Cat. Pl. Zeyl.* 102.—Zeylan.
lucida, N. E. Br. in Illustr. Hortic. xxxii. (1885) 77. t. 557 = humilis.
palustris, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 309.—Ins. Philipp.
pycnocephala, *Benth. in Journ. Linn. Soc.* xv. (1877) 512 (lapsu *pycnostachya*).—Panama.
sumatrana, *Benth. & Hook. f. Gen.* iii. 1055.—Malaya.
sylvatica, *Aubl. Pl. Gui.* i. 47. t. 17.—Guian.; Panama.
sylvatica, Pursh, Fl. Am. Sept. i. 47 = Carex Fraseri.
sylvestris, Vahl, Enum. ii. 391 = sylvatica.
zeylanica, *Trimen, Cat. Pl. Zeyl.* 102.—Ind. or.; Borneo.

MAPIRA, Adans. Fam. ii. 39 (1763) = **Olyra**, Linn. (Gramin.).

MAPOURIA, Aubl. Pl. Gui. i. 175. t. 67 (1775) = **Psychotria**, Linn. (Rubiac.).
aemulans, Muell. Arg. in Flora, lix. (1876) 496, 497.—Bras.
alba, Muell. Arg. l. c. 458 = P. alba.
apocynacea, Muell. Arg. l. c. 457, 464.—Bras.
australis, Muell. Arg. l. c. 459, 465.—Bras.
borjensis, Muell. Arg. l. c. 459, 466 = P. borjensis.
brachypoda, Muell. Arg. in Mart. Fl. Bras. vi. v. 422.—Bras.
brevicollis, Muell. Arg. in Flora, lix. (1876) 459, 465 = P. brevicollis.
Burchelliana, Muell. Arg. l. c. = P. Burchelliana.
caldasica, Muell. Arg. l. c.—Bras.
capillacea, Muell. Arg. in Mart. Fl. Bras. vi. v. 405.—Bras.
capituliflora, Muell. Arg. in Flora, lix. (1876) 495, 497.—Bras.
castellana, Muell. Arg. l. c. 458, 464 = P. castellana.
catharinensis, Muell. Arg. l. c. 459, 465.—Bras.
cephalantha, Muell. Arg. l. c. 495, 497.—Bras.
chaenotricha, Muell. Arg. l. c. 496 = P. chaenotricha.
chionantha, Muell. Arg. in Mart. Fl. Bras. vi. v. 387 = P. Luschnathii.
cinchonoides, Muell. Arg. l. c. 388.—Bras.
colarensis, Muell. Arg. in Flora, lix. (1876) 459, 465.—Bras.
compaginata, Muell. Arg. l. c. 457, 464.—Bras.
cordata, Muell. Arg. in Mart. Fl. Bras. vi. v. 426.—Bras.
corymbifera, Muell. Arg. in Flora, lix. (1876) 458, 464.—Bras.
coussareoides, Muell. Arg. l. c. 460, 466.—Bras.
crassa, Muell. Arg. in Mart. Fl. Bras. vi. v. 389.—Bras.
cuparis, Muell. Arg. in Flora, lix. (1876) 459, 465.—Bras.
cymuligera, Muell. Arg. l. c. 496, 497.—Bras.
depauperata, Muell. Arg. in Mart. Fl. Bras. vi. v. 466.—Bras.
ferruginea, Muell. Arg. in Flora, lix. (1876) 457, 464.—Bras.
formosa, Muell. Arg. l. c. 459 = P. formosa.
graciliflora, Oerst. Am. Centr. 17. t. 14. f. 6 = P. graciliflora.
guianensis, Aubl. Pl. Gui. i. 175 = P. nitida.
herbacea, Muell. Arg. in Mart. Fl. Bras. vi. v. 427 = P. herbacea.
insignis, Muell. Arg. l. c. 424.—Bras.
japurensis, Muell. Arg. l. c. 412 = P. japurensis.
laevifolia, Muell. Arg. l. c. 416.—Bras.
Langsdorffiana, Muell. Arg. l. c. 395 = P. Langsdorffiana.
lurida, Muell. Arg. l. c. 405.—Bras.
Luschnathii, Muell. Arg. in Flora, lix. (1876) 457 = P. Luschnathii.
macrocarpa, Muell. Arg. in Mart. Fl. Bras. vi. v. 425 = P. herbacea.
mandiocana, Muell. Arg. in Flora, lix. (1876) 496, 498 = P. mandiocana.
Martiana, Muell. Arg. l. c. 458, 464 = P. Martiana.
microcarpa, Muell. Arg. l. c. 496, 497 = P. microcarpa.
miradorensis, Oerst. Am. Centr. 17. t. 14. f. 9 = P. miradorensis.
myriantha, Muell. Arg. in Mart. Fl. Bras. vi. v. 400 = P. myriantha.
Nettoana, Muell. Arg. in Flora, lix. (1876) 458, 464.—Bras.
niveo-barbata, Muell. Arg. in Mart. Fl. Bras. vi. v. 401.—Bras.
obovata, Oerst. Am. Centr. 17. t. 14. f. 3, 4 = P. obovata.
panurensis, Muell. Arg. in Flora, lix. (1876) 460, 466.—Bras.
papantlensis, Oerst. Am. Centr. 17. t. 14. f. 5 = P. papantlensis.
parvifolia, Oerst. l. c. = P. parvifolia.
peraffinis, Muell. Arg. in Flora, lix. (1876) 496, 497.—Bras.
podocephala, Muell. Arg. l. c. 460, 466.—Bras.
Pohliana, Muell. Arg. l. c. 457, 464.—Bras.
ptychophylla, Muell. Arg. l. c. 458, 464.—Bras.
puberulina, Muell. Arg. l. c. 459, 465.—Bras.
Rabeniana, Muell. Arg. l. c. 458, 464 = P. Rabeniana.
remota, Muell. Arg. l. c. 459 = P. remota.
Riedeliana, Muell. Arg. l. c. 496 = P. Riedeliana.
salicifolia, Oerst. Am. Centr. 17. t. 14. f. 2 = P. salicifolia, *Hemsl.*

MAPOURIA:—
sambucina, Muell. Arg. in Flora, lix. (1876) 458 = Psychotria sambucina.
Schlechtendaliana, Muell. Arg. l. c. 457 = P. elliptica.
sclerocalyx, Muell. Arg. l. c. 496, 497.—Bras.
sessiliflora, Muell. Arg. in Mart. Fl. Bras. vi. v. 416 = P. sessiliflora.
Simira, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 174 = P. parviflora, *Willd.*
sphaerocarpus, Muell. Arg. in Mart. Fl. Bras. vi. v. 418.—Bras.
Spruceana, Muell. Arg. in Flora, lix. (1876) 496, 497.—Bras.
stipulata, Oerst. Am. Centr. 17. t. 14. f. 7 = P. stipulata.
subsessilis, Muell. Arg. in Flora, lix. (1876) 460 = P. subsessilis?
subspathulata, Muell. Arg. l. c. 459, 465.—Bras.
tenuis, Muell. Arg. in Mart. Fl. Bras. vi. v. 425.—Bras.
tomentosa, Oerst. Am. Centr. 17. t. 14. f. 8 = P. tomentosa, *Hemsl.*
triadica, Muell. Arg. in Mart. Fl. Bras. vi. v. 423.—Bras.
tricephala, Muell. Arg. l. c.—Bras.
trichogyne, Muell. Arg. l. c. 426.—Bras.
tristis, Muell. Arg. in Flora, lix. (1876) 458, 465.—Bras.
umbelluligera, Muell. Arg. l. c. 496, 497.—Bras.
umbrosa, Muell. Arg. l. c. 459, 466.—Bras.
vasivensis, Muell. Arg. l. c. 460, 466.—Bras.
Velhana, Muell. Arg. l. c. 457, 464.—Bras.
vestita, Muell. Arg. in Mart. Fl. Bras. vi. v. 421 = P. vestita.
xanthophylla, Muell. Arg. in Flora, lix. (1876) 496, 497 = P. xanthophylla.
xanthophylloides, Muell. Arg. l. c.—Bras.

MAPPA, A. Juss. Euph. Tent. 44. t. 14 (1824) = **Macaranga**, Thou. (Euphorb.).
acuminatissima, Zipp. ex Span. in Linnaea, xv. (1841) 349, nomen = Mallotus subpeltatus.
acutifolia, Zoll. & Mor. Syst. Verz. Zoll. 17 = Mallotus acuminatus.
amboinensis, Muell. Arg. in Linnaea, xxxiv. (1865-66) 197 = Mac. amboinensis.
angolensis, Muell. Arg. in Journ. Bot. ii. (1864) 337 = Mac. angolensis.
bancana, Miq. Fl. Ind. Bat. Suppl. 458 = Mac. javanica.
borneensis, Muell. Arg. in Flora, xlvii. (1864) 468 = Mac. trichocarpa.
capensis, Baill. Etud. Gén. Euph. 430 = Mallotus capensis.
Chatiniana, Baill. Adansonia, i. (1860-61) 349 = Mac. Chatiniana.
cissifolia, Zoll. in Linnaea, xxix. (1857-58) 466 = Mac. cissifolia.
cochinchinensis, Spreng. Syst. iii. 878 = Mallotus cochinchinensis.
conifera, Zoll. in Linnaea, xxix. (1857-58) 466 = Mac. conifera.
coriacea, Muell. Arg. l. c. xxxiv. (1865-66) 198 = Mac. coriacea.
corymbosa, Muell. Arg. l. c. 199 = Mac. corymbosa.
costulata, Miq. Fl. Ind. Bat. Suppl. 459 = Baccaurea costulata.
Cumingii, Muell. Arg. in DC. Prod. xv. II. 1005 = Mac. Cumingii.
cuneifolia, Zoll. in Linnaea, xxix. (1857-58) 467 = Mac. cuneifolia.
denticulata, Blume, Bijdr. 625 = Mac. denticulata.
Diepenhorstii, Miq. Fl. Ind. Bat. Suppl. 457 = Mac. Diepenhorstii.
digyna, Muell. Arg. in Linnaea, xxxiv. (1865-66) 198 = Mac. digyna.
fastuosa, Linden, in Belg. Hortic. xv. (1865) 100 = Homalanthus fastuosus.
floribunda, Zoll. & Mor. Syst. Verz. Zoll. 17 = Mallotus floribundus.
gigantea, Zoll. in Linnaea, xxix. (1857-58) 465 = Mac. gigantea.
glabra, A. Juss. Euph. Tent. 114. t. 14 = Mac. Tanarius.
gummiflua, Miq. Fl. Ind. Bat. Suppl. 458 = Mac. denticulata.
Harveyana, Muell. Arg. in Flora, xlvii. (1864) 467 = Mac. Harveyana.
heterophylla, Muell. Arg. in Journ. Bot. ii. (1864) 336 = Mac. heterophylla.
hispida, Blume, Bijdr. 624 = Mac. hispida.

MAPPA:—
hypoleuca, Reichb. & Zoll. in Verh. Natuurk. Ver. Ned. Ind. Bat. i. (1856) 30; et in Linnaea, xxviii. (1856) 309 = Mac. hypoleuca.
jamaicensis, Spreng. Syst. iii. 878 = Mettenia globosa.
javanica, Blume, Bijdr. 625 = Mac. javanica.
leptostachya, Muell. Arg. in Linnaea, xxxiv. (1865-66) 198 (Cleidionis sp.).—Ins. Pacif.
macrophylla, A. Gray, ex Seem. in Bonplandia, ix. (1861) 258 = Mac. macrophylla.
macrophylla, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvi. (1864) 44 = Mac. gigantea.
mallotoides, F. Muell. Fragm. iv. 139 = Mac. involucrata.
megalophylla, Muell. Arg. in Flora, xlvii. (1864) 467 = Mac. megalophylla.
moluccana, Spreng. Syst. iii. 878 = Mac. Mappa.
moluccana, Wight, Ic. t. 816 = Mac. Tanarius.
Motleyana, Muell. Arg. in Flora, xlvii. (1864) 467 = Mac. Motleyana.
occidentalis, Muell. Arg. l. c. = Mac. occidentalis.
paniculata, Wall. Cat. n. 7810 = Mac. denticulata.
pellata, Wight, Ic. t. 817 = Mac. Roxburghii.
populifolia, Muell. Arg. in Linnaea, xxxiv. (1865-66) 198 = Mac. populifolia.
pruinosa, Miq. Fl. Ind. Bat. Suppl. 457 = Mac. pruinosa.
pubescens, Brongn. ex Baill. Etud. Gén. Euph. t. 20. f. 5-7 (Quid?).—Hab.?
rhynchophylla, Miq. Fl. Ind. Bat. i. II. 403 = Mallotus subpeltatus.
rugosa, Muell. Arg. in Linnaea, xxxiv. (1865-66) 197 = Mac. megalophylla.
scandens, Panch. ex Baill. Adansonia, ii. (1861-62) 223 = Mallotus repandus.
Seemanni, Muell. Arg. in Flora, xlvii. (1864) 468 = Mac. Seemanni.
sinensis, Muell. Arg. in DC. Prod. xv. II. 1001 = Mac. sinensis.
stricta, Reichb. & Zoll. in Verh. Nat. Ver. Ned. Ind. Bat. i. (1856) 31; et in Linnaea, xxviii. (1856) 310 = Mac. stricta.
taitensis, Muell. Arg. in Linnaea, xxxiv. (1865-66) 197 = Mac. taitensis.
tanaria, Spreng. Syst. iii. 878 = seq.
Tanarius, Blume, Bijdr. 624 = Mac. Tanarius.
tannensis, Spreng. Syst. iii. 878 = Mac. dioica.
Teysmanni, Zoll. in Linnaea, xxix. (1857-58) 465 = Mac. Teysmanni.
tomentosa, Blume, Bijdr. 624 = Mac. Tanarius.
trichocarpa, Zoll. in Linnaea, xxviii. (1856) 307 = Mac. trichocarpa.
triloba, Muell. Arg. in Flora, xlvii. (1864) 466 = Mac. Griffithiana.
truncata, Muell. Arg. in Linnaea, xxxiv. (1865-66) 198 = Mac. denticulata.
Vedeliana, Muell. Arg. l. c. = Mac. Vedeliana.
velutina, Reichb. & Zoll. l. c. xxix. (1857-58) 468 = Mac. velutina.
Vieillardi, Muell. Arg. l. c. xxxiv. (1865-66) 199 = Mac. Vieillardi.
Wallichii, Muell. Arg. in DC. Prod. xv. II. 1000 = Mac. denticulata.
Zippelii, Zoll. & Mor. Syst. Verz. Zoll. 17 = Mallotus ricinoides.
Zollingeriana, Miq. Fl. Ind. Bat. Suppl. 457 = Mac. trichocarpa.

MAPPIA, Habl. ex Ledeb. Fl. Ross. ii. 403 (1844) = **Crucianella**, Linn. (Rubiac.).
fasciculata, Habl. l. c. = C. glauca.

MAPPIA, Heist. ex Adans. Fam. ii. 193 (1763) = **Cunila**, Linn. (Labiat.).

MAPPIA, Jacq. Hort. Schoenb. i. 22. t. 47 (1797). *OLACINEAE*, Benth. & Hook. f. i. 351.
LERETIA, Vell. Fl. Flum. iii. t. 2 (1827).
affinis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. ix. (1852) 395.—Jamaic.
ampla, *Engl. in Mart. Fl. Bras.* xii. II. 51.—Bras.
angustifolia, *Griseb. Cat. Pl. Cub.* 119.—Cuba.
Championiana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 397 = oblonga.
cordata, *Engl. in Mart. Fl. Bras.* xii. II. 50.—Bras.
foetida, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. ix. (1852) 395.—Ind. or.
Gardneriana, Miers, l. c. 396 = ovata.
montana, *Miers, l. c.* 398.—Java.

MAPPIA :—
nitida, *Engl. in Mart. Fl. Bras.* xii. II. 51.—Bras.
oblonga, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. ix. (1852) 396.—Ind. or.
ovata, *Miers, l. c.*—Ind. or.
Poeppigiana, *Baill. Adansonia*, xi. (1874) 175.—Bras.
racemosa, *Jacq. Hort. Schoenb.* i. 22.—Jamaic.
senegalensis, *Baill. Hist. Pl.* v. 277.—Afr. trop.
tomentosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. ix. (1852) 397.—Ind. or.
Wightiana, Miers, l. c. = ovata.

MAPPIA, Schreb. Gen. 806 (1791) = **Doliocarpus**, Roland. (Dillen.).

MAPROUNEA, Aubl. Pl. Gui. ii. 895. t. 342 (1775).
EUPHORBIACEAE, Benth. & Hook. f. iii. 333.
AEGOPRICUM, Linn. Pl. Surinam. 15 (1775) ; et in Amoen. Acad. viii. 263.
africana, *Muell. Arg. in DC. Prod.* xv. II. 1191.—Afr. occ.
brasiliensis, *A. St. Hil. Pl. Us. Brés.* t. 65.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 895.—Bras. ; Guian.

MAPROUNIA, Ham. Prod. Pl. Ind. Occ. 53 (1825) = praec.

MAPRUNEA, J. F. Gmel. Syst. 272 (1791) = praec.

MAPURIA, J. F. Gmel. Syst. 371 (1791) = MAPOURIA, Aubl. = **Psychotria**, Linn. (Rubiac.).

MAQUIRA, Aubl. Pl. Gui. Suppl. 36. t. 389 (1775) = **Helicostylis**? Tréc. (Urtic.).
granatensis, Baill. Adansonia, xi. (1875) 292.—N. Granat.
grandifolia, Baill. ex Benth. & Hook. f. Gen. iii. 371 = Olmedia grandifolia.
guianensis, Aubl. Pl. Gui. Suppl. 36.—Guian.
laurifolia, Bur. ex Benth. & Hook. f. Gen. iii. 373 = Perebea laurifolia.

MARAH, Kellogg, in Proc. Calif. Acad. ii. (1863) 18 ; et i. ed. II. (1873) 37 = **Echinocystis**, Torr. & Gray (Cucurb.).
minima, Kellogg, l. c. ii. (1863) 18 = E. minima.
muricatus, Kellogg, l. c. i. ed. II. (1873) 37 = E. Marah.

MARALIA, Thou. Gen. Nov. Madag. 13 (1806) = **Panax**, Linn. (Araliac.).
madagascariensis, DC. Prod. iv. 255 = P. Maralia.

MARAMA, Rafin. Fl. Tellur. iv. 62 (1836) = **Graptophyllum**, Nees (Acanth.).
picta, Rafin. l. c. = G. hortense.

MARANTA, Plum. ex Linn. Gen. ed. I. 331 (1737).
SCITAMINEAE, Benth. & Hook. f. iii. 649.
PSYDARANTA, Neck. Elem. iii. 145 (1790).
affinis, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 57 = Porteana.
Allouia, Aubl. Pl. Gui. i. 3 = Calathea Allouia.
amabilis, *Linden, Cat.* n. 22 (1869) 7.—Bras.
angustifolia, Sims, Bot. Mag. t. 2398 = Stromanthe Tonckat.
argyraea, Hort. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 142 = Calathea argyraea.
argyrophylla, *Linden, ex C. Koch, in Berl. Gartenz.* (1857) 248.—Hab. ?
Arouma, Aubl. Pl. Gui. i. 3 = Ischnosiphon Arouma.
arundinacea, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 108 = Thalia geniculata.
arundinacea, Blanco, Fl. Filip. ed. I. 7 = Clinogyne dichotoma.
arundinacea, *Linn. Sp. Pl.* 2.—Am. austr.
aurantiaca, *Hort. ex C. Koch, in Berl. Gartenz.* (1857) 149, nomen.—Hab. ?
bambusacea, D. Dietr. Syn. Pl. i. 7 = Ischnosiphon bambusacea.
Baraquinii, Lem. Illustr. Hortic. xv. (1868) t. 542 = Calathea Baraquini (in Addend.).
bella.—Cf. Gard. Chron. (1875) I. 822 = Kegeljani.
bicolor, *Ker-Gawl. in Bot. Reg.* t. 786.—Bras.
bicolor, Vell. Fl. Flum. 3 ; i. t. 7 = Calathea zebrina.
Blumei, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 52 = Clinogyne Blumei (in Addend.).

MARANTA :—
borussica, Linden, Cat. n. 13 (1858) 4 ; et ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 125 = Calathea fasciata.
brachystachys, Benth. in Hook. Niger Fl. 531 = Phrynium brachystachyum.
Cachibou, Jacq. Fragm. 52 = Calathea discolor.
caespitosa, A. Dietr. Sp. Pl. i. 31 = Phrynium spicatum, *Roxb.*
capitata, Ruiz & Pav. Fl. Per. i. 3. t. 5 = Calathea capitata.
Casupito, Jacq. Fragm. 51 = Calathea discolor.
Casupo, Jacq. l. c. = Calathea discolor.
chimboracensis, Linden, in Illustr. Hortic. xvii. (1870) t. 6 = Calathea chimboracensis.
Chouca, *Hort. ex C. Koch, in Berl. Gartenz.* (1857) 149, nomen.—Hab. ?
cinerea, *Hort. ex Denester, in Bull. Soc. Linn. Brux.* i. (1892) 59.—Peruv.
clavata, Vell. Fl. Flum. 3 ; i. t. 9 = Calathea grandifolia.
coccinea, *Hort. ex C. Koch, in Berl. Gartenz.* (1857) 149, nomen.—Hab. ?
comosa, Linn. f. Suppl. 80 = Calathea comosa.
composita, Hort. ex Koern. in Regel, Gartenfl. (1858) 83 = Riedelianum.
compressa, A. Dietr. Sp. Pl. i. 22 = Ctenanthe Luschnathiana.
concinna, Hort. Bull, ex Regel, in Gartenfl. (1877) 35 = Calathea leopardina.
cordata, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 39.—Bras.
cristata, Nees & Mart. in Nov. Act. Nat. Cur. xi. I. (1823) 25 = bicolor.
cuspidata, Rosc. Scitam. t. 31 = Clinogyne filipes ?
cuyabensis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 68.—Bras.
cylindrica, A. Dietr. Sp. Pl. i. 27 = Calathea grandifolia.
dealbata, A. Dietr. l. c. 24 = Thalia dealbata.
depressa, *E. Morr. in Belg. Hortic.* xxx. (1880) 97. t. 6.—Bras.
dichotoma, Poepp. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 45 = Ruiziana.
dichotoma, Wall. Cat. n. 6614 = Clinogyne dichotoma.
discolor, Hort. ex Koern. iu Bull. Soc. Nat. Mosc. xxxv. (1862) I. 136 = Calathea angustifolia.
discolor, Hort. ex Koern. l. c. = Calathea varians.
disticha, Buchoz, Ic. Coll. t. 156 = Calathea lutea.
divaricata, *Rosc. Scitam.* t. 27.—Bras.
dubia, Roem. & Schult. Syst. Add. i. 70 = Ischnosiphon spicatus.
eburnea, *Linden & André, in Illustr. Hortic.* xx. (1873) 171.—N. Granat.
eximia, L. Mathieu, ex Koern. in Regel, Gartenfl. (1858) 87 = Calathea eximia.
eximia, *Regel, l. c.* (1878) 103.—Hab. ?
fasciata, Linden, ex C. Koch, in Berl. Gartenz. (1857) 248 = Calathea fasciata.
flavescens, A. Dietr. Sp. Pl. i. 25 = Calathea flavescens.
flexuosa, G. Don, ex Benth. in Hook. Niger Fl. 531 = Phrynium flexuosum.
flexuosa, *Lestib. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 217.—Hab. ?
flexuosa, Presl, Rel. Haenk. i. 107 = Thalia geniculata.
foliosa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 47.—Bras.
Friedrichsthaliana, *Koern. l. c.* 42.—Am. centr.
fruticosa, Stokes, Bot. Mat. Med. i. 5 = Stromanthe Tonckat.
furcata, Nees & Mart. in Nov. Act. Nat. Cur. xi. I. (1823) 24 = gibba.
Galanga, Linn. Sp. Pl. ed. II. 3 = Alpinia Galanga.
geniculata, Lam. Tabl. Encyc. i. 9 = Thalia geniculata.
gibba, *Sm. in Rees, Cyclop.* xxii. n. 2.—Ins. Barbados.
glabra, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 63 = Ctenanthe glabra.
glumacea, *Van Houtte, ex Koern. l. c.* 62.—Am. trop.
gracilis, Rudge, Pl. Guian. Rar. 8. t. 3 = Ischnosiphon gracilis.
graminifolia, Ruiz & Pav. ex Steud. Nom. ed. II. ii. 100 = Stromanthe Tonckat.
grandiflora, A. Dietr. Sp. Pl. i. 30 = Calathea flavescens.
grandifolia, A. Dietr. l. c. 26 = Calathea grandifolia.
grandis, Miq. Fl. Ind. Bat. Suppl. 616 = Clinogyne grandis.
Herderiana, *Regel, Ind. Sem. Hort. Petrop.* (1867) 28. —Hab. ?
hexantha, *D. Dietr. Syn. Pl.* i. 6.—Bras.

MARANTA :—
hieroglyphica, *Linden & André, in Belg. Hortic.* xxiii. (1873) 111.—N. Granat.
Hjalmarssoni, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 54.—Honduras.
Hoyeti, *Jacob-Makoy, ex Gard. Chron.* (1878) I. 440. —Bras.
humilis, *Aubl. Pl. Gui.* i. 4.—Guian.
humilis, Vell. Fl. Flum. 3 ; i. t. 6 = Calathea violacea.
illustris, *Linden, Illustr. Hortic.* xiv. (1867) t. 515. —Hab. ?
imbricata, A. Dietr. Sp. Pl. i. 30 = Phrynium imbricatum.
indica, Tussac, Fl. Antill. i. 183 = arundinacea.
inscripta, *Hort.—Cf. Gard. Chron.* (1876) I. 735.—Hab. ?
insignis, *Hort. ex C. Koch, in Berl. Gartenz.* (1857) 149, nomen.—Hab. ?
Jacquini, Presl, Rel. Haenk. i. 107 = Ischnosiphon polyphyllus.
Jacquini, Roem. & Schult. Syst. i. 558 = Stromanthe lutea.
Jagoriana, Hort. ex Regel, Gartenfl. (1879) 297 = Calathea Jagoriana.
juncea, Lam. Encyc. ii. 589 = Ischnosiphon Arouna.
juncea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = Stromanthe lutea.
Kegeljani, *E. Morr. in Belg. Hortic.* xxv. (1875) 135, 216.—Hab. ?
Kerchovei, Hort.—Cf. Gard. Chron. (1878) I. 440 = seq.
Kerchoviana, E. Morr. in Belg. Hortic. (1879) 77. t. 5 = leuconeura.
Klotzschiana, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 50.—Bras.
lachnocaula, *Miq. in Linnaea*, xviii. (1844) 755.—Guiana.
lateralis, Ruiz & Pav. Fl. Per. i. 4 = Calathea lateralis.
latifolia, *Regel, ex Horan. Prod. Scitam.* 10, nomen.—Hab. ?
laxa, D. Dietr. Syn. Pl. i. 7 = Ischnosiphon laxus.
Legrelleana, Linden, in Belg. Hortic. xvii. (1867) 104 = Calathea Legrelliana.
leopardina, Hort.—Cf. Gard. Chron. (1875) I. 523 ; et Van Geert, in Rev. Hortic. Belg. vi. (1880) 13 = Calathea leopardina.
leptostachya, *Regel & Koern. in Ind. Sem. Hort. Petrop.* (1857) 32.—Bras.
leucocephala, D. Dietr. Syn. Pl. i. 7 = Ischnosiphon leucophoeus.
leuconeura, *E. Morr. Belg. Hortic.* xxiv. (1874) 323 ; *et* xxv. (1875) 172.—Bras.
Lindeni, Hort. ex E. Morr. l. c. xx. (1870) 1 = Calathea Lindeniana.
longibracteata, A. Dietr. Sp. Pl. i. 26 = Calathea longibracteata.
longifolia, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 426 = Calathea longifolia.
lucida, *Regel, ex Horan. Prod. Scitam.* 10, nomen.—Hab. ?
Luschnathiana, Regel & Koern. in Ind. Sem. Hort. Petrop. (1857) 31 = Ctenanthe Luschnathiana.
lutea, Aubl. Pl. Gui. i. 4 = Calathea discolor.
lutea, Jacq. Coll. iv. 117 = Stromanthe lutea.
macilenta, A. Dietr. Sp. Pl. i. 27 = Calathea macilenta.
maculata, Pav. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 116 = Calathea Pavonii.
majestica, Linden, in Belg. Hortic. xv. (1865) 103 = Calathea ornata.
malaccensis, Burm. f. Fl. Ind. 2 = Alpinia malaccensis.
Massangeana, *E. Morr. Belg. Hortic.* (1875) 178. t. 10. —Bras.
Mazellii, *Hort.—Cf. Gard. Chron.* (1872) 7.—Peruv.
mediopicta, E. Morr. in Belg. Hortic. xxv. (1875) 135 = Calathea mediopicta.
metallica, Hort. ex Koern. in Regel, Gartenfl. (1858) 88 = Calathea metallica.
mischanta, *Vell. Fl. Flum.* 4 ; i. t. 12.—Bras.
monophylla, Vell. l. c. t. 11 = Calathea monophylla.
Moritziana, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 66.—Am. trop.
Morreni, *Jacob-Makoy.—Cf. Gard. Chron.* (1878) I. 440.—Bras.
Myrosma, A. Dietr. Sp. Pl. i. 22 = Calathea Myrosma.
nana, Poepp. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) 72 = Calathea micans.
niveiflora, *A. Dietr. Sp. Pl.* i. 23.—Ins. Trinit.
noctiflora, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1858) 31.—Hab. ?

MARANTA:—
obliqua, Rudge, Pl. Guian. Rar. 8. t. 2 = Ischnosiphon obliquus.
obliqua, Seubert, ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 85 = Ischnosiphon laxus.
oblongifolia, *Regel, Gartenfl.* (1878) 101.—Bras.
olivaris, Hort. Veitch.—Cf. Gard. Chron. (1872) 1589 = Calathea Makoyana.
ornata, Linden, in Fl. des Serres, iv. (1848). t. 413 = Calathea ornata.
ovata, A Dietr. Sp. Pl. i. 29 = Calathea ovata.
Parkeri, A. Dietr. l. c. 25 = Ischnosiphon Parkeri.
parviflora, A. Dietr. l. c. 30 = Phrynium parviflorum.
petiolata, Rudge, Pl. Guian. Rar. 25. t. 37 = Ischnosiphon Arouma.
phrynioides, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 40.—Bras.
Pierardi, *Jacob-Makoy.—Cf. Gard. Chron.* (1878) i. 440.—Bras.
pilosa, Humb. ex Link, Jahrb. i. iii. 20 = Stromanthe Tonckat.
pilosa, Schau. in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) 426 = Ctenanthe pilosa.
pinnato-picta, Hort.—Cf. Gard. Chron. (1875) i. 822 = Calathea applicata.
Placentaria, A. Dietr. Sp. Pl. i. 30 = Phrynium capitatum.
Pohliana, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 44 = divaricata.
polyphylla, D. Dietr. Syn. Pl. i. 6 = Ischnosiphon polyphyllus.
Porteana, Horan. Prod. Scitam. 10; Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 57 = Stromanthe Porteana.
prasina, Hort.—Cf. Gard. Chron. (1875) i. 822 = Calathea medio-picta.
Princeps, Linden, ex Regel, Gartenfl. (1879) 302 = Calathea Princeps.
prolifera, *Vell. Fl. Flum.* 4; i. t. 10.—Bras.
protracta, *Miq. in Linnaea*, xviii. (1844) 71.—Guian.
pruinosa, *Hort. Bull, ex Regel, Gartenfl.* (1878) 104. —Am. trop.
pubescens, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 917 = divaricata.
pulchella, Linden, ex C. Koch, in Berl. Allg. Gartenz. (1857) 243 = Calathea zebrina.
pulchella, E. Morr. in Belg. Hortic. xxv. (1875) 272. f. 6 = Calathea pulchella.
pulverulenta, *Hort. ex Koern. in Regel, Gartenfl.* (1858) 83, nomen.—Hab.?
pumila, Vell. Fl. Flum. 3; i. t. 8 = Calathea pumila.
purpurascens, Link, ex A. Dietr. Sp. Pl. i. 20 = divaricata.
pygmaea, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 61.—Bras.
racemosa, A. Dietr. Sp. Pl. i. 21 = Stromanthe Tonckat.
ramosissima, Wall. Pl. As. Rar. iii. 51. t. 286 = Clinogyne dichotoma.
regalis, Hort. Angl. ex Lem. Illustr. Hortic. ii. (1855) t. 74 = Calathea ornata.
Riedeliana, *Koern. in Regel, Gartenfl.* (1858) 80.—Bras.
roseo-picta, Hort.—Cf. Gard. Chron. (1865) 365; Linden, Illustr. Hortic. xiv. (1867) t. 508 = Calathea roseo-picta.
rotundifolia, A. Dietr. Syn. Pl. i. 6 = Ischnosiphon rotundifolius.
rotundifolia, Hort. ex Koern. in Regel, Gartenfl. vii. (1858) 83 = Calathea rotundifolia.
Ruiziana, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 45 (Ruicziana).—Peruv.
sanguinea, Hort. ex Planch. in Fl. des Serres, viii. (1852–53) 97 = Stromanthe sanguinea.
secunda, R. Grah. in Edinb. Phil. Journ. xiii. (1825) 357 = Myrosma setosum.
Seemanni, *Hort. ex Gard. Chron.* (1872) 323. f. 113.—Hab.?
Selloi, *Horan. Prod. Scitam.* 10.—Hab.?
Selloi, Hort. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 64 = Ctenanthe Luschnathiana.
semperflorens, Horan. Prod. Scitam. 10 = Calathea Allouia.
sericea, *Hort. ex Koern. in Regel, Gartenfl.* (1858) 83, nomen.—Hab.?
setosa, A. Dietr. Sp. Pl. i. 23 = Myrosma setosum.
smaragdina, Linden, ex Regel, Gartenfl. (1878) 104 = Calathea smaragdina.
speciosa, Bull. Cat. (1884) 14 = leptostachya.

MARANTA:—
spectabilis, Koern. in Regel, Gartenfl. (1858) 79 = Stromanthe sanguinea.
spicata, Aubl. Pl. Gui. i. 4 = Ischnosiphon spicatus.
spicata, Thw. Enum. Pl. Zeyl. 320, excl. syn. = Phrynium zeylanicum.
splendida, *Hort.—Cf. Gard. Chron.* (1864) 414.—Hab.?
splendida, *Hort. Versch. ex Lem. Illustr. Hortic.* xiii. (1866) tt. 467–8.—Bras.
striata, *Veitch.—Cf. Gard. Chron.* (1864) 671; *Nichols. Dict. Gard.* ii. 327.—Ins. Philipp.
strigosa, *Horan. Prod. Scitam.* 10.—Bras.
strobilifera, Miq. in Linnaea, xxii. (1849) 79 = Calathea strobilifera.
surinamensis, Miq. in Linnaea, xviii. (1844) 603 = Ischnosiphon surinamensis.
sylvatica, Rosc. in Trans. Linn. Soc. viii. (1807) 340, nomen; et Sm. in Rees, Cyclop. xxii. n. 3 = arundinacea.
Tonchat, Blume, Enum. Pl. Jav. 36 = Clinogyne dichotoma.
Tonchat, Aubl. Pl. Gui. i. 3 = Stromanthe Tonckat.
truncata, Link, ex A. Dietr. Sp. Pl. i. 26 = Calathea orbiculata.
tuberosa, Vell. Fl. Flum. 4; i. t. 13 = Calathea Arribida.
tubispatha, Hort. ex Denester, in Bull. Soc. Linn. Brux. i. (1872) 58 = Calathea tubispatha.
undulata, Linden & André, Cat. n. 87 (1871) p. 4 = Calathea undulata.
unilateralis, D. Dietr. Syn. Pl. i. 6 = Myrosma australe.
Van-den-Heckei, Hort. Versch. ex Lem. Illustr. Hortic. x. (1863) t. 380 = Calathea Vandenheckei.
variegata, Hort. ex Koern. in Regel, Gartenfl. (1858) 88 = Calathea variegata.
variegata, *Lodd. ex G. Don, in Loud. Hort. Brit.* 1, nomen.—Am. austr.
Veitchiana, Van Houtte, in Fl. des Serres, Sér. II vi. (1865–67) 89 = Calathea Veitchiana.
Veitchii, *Hort.—Cf. Gard. Chron.* (1864) 414, *et* (1870) 924. t. 170.—Hab.?
villosa, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 49 = Stromanthe Tonckat.
violacea, A. Dietr. Sp. Pl. i. 25 = Calathea violacea.
virgata, A. Dietr. l. c. 21 = Phrynium virgatum.
virgata, Wall. Cat. n. 6616 = Clinogyne virgata.
virginalis, *Linden, ex Lem. Illustr. Hortic.* xv. (1868) t. 550.—Bras.
vittata, Hort. ex Koern. in Regel, Gartenfl. (1858) 88 = Calathea vittata.
Wallisi, *Linden, Belg. Hortic.* xvii. (1867) 105.—Peruv.
Warscewiczii, Hort. ex Planch. in Fl. des Serres, ix (1853–54) 209. t. 939 = Calathea Warscewiczii.
Warszewiczii, Lem. Illustr. Hortic. i. (1854) t. 7 = praec.
Wiotii, E. Morr. Belg. Hortic. xxv. (1875) 273 f. 7 = Calathea Wiotiana.
zebrina, Sims, Bot. Mag. t. 1926 = Calathea zebrina.

MARANTHES, Blume, Bijdr. 89 (1825) = **Parinarium**, Juss. (Rosac.).
corymbosa, Blume, l. c.—Java.
multiflora, Korth. Verh. Nat. Gesch. Bot. 259. t. 70 = P. Griffithianum.

MARANTHUS, Reichb. Consp. 204 (1828) = praec.

MARANTOCHLOA, Brongn. ex Gris, in Bull. Soc. Bot. Fr. vii. (1860) 321. *SCITAMINEAE.*
comorensis, *Brongn. ex Gris, l. c.*—Ins. Comor.

MARANTOPSIS, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 97 = **Stromanthe**, Sond. (Scitam.).
lutea, Koern. l. c. = S. lutea.
Schottiana, Koern. l. c. 99 = S. Schottiana.

MARARA, Karst. in Linnaea, xxviii. (1856) 389 = **Martinezia**, Ruiz & Pav. (Palm.).
aculeata, Karst. ex H. Wendl. in Kerch. Palm. 250 = M. Aiphanes.
bicuspidata, Karst. in Linnaea, xxviii. (1856) 390 = M. elegans.
caryotaefolia, Karst. ex H. Wendl. in Kerch. Palm. 250 = M. caryotaefolia.
erinacea, Karst. in Linnaea, xxviii. (1856) 391 (= *Aiphanes erinacea*).—N. Granat.

MARASMODES, DC. Prod. vi. 136 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 432.
ADENOSOLEN, DC. Prod. vi. 136 (1837).
BRACHYMERIS, DC. l. c. 76 (1837).
BRACHYSTYLIS, E. Mey. ex DC. l. c. (1837).
OLIGODORELLA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 187.
adenoxelon, *Harv. in Harv. & Sond. Fl. Cap.* iii. 175. —Afr. austr.
oligocephalus, *DC. Prod.* vi. 136.—Afr. austr.
polycephalus, *DC. l. c.*—Afr. austr.

MARATHRUM, Humb. & Bonpl. Pl. Aequin. i. 39. t. 11 (1808). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 110.
cubanum, *Wright, in Sauv. Fl. Cub.* 150.—Cuba.
flexuosum, *Liebm. in Forhandl. Skand. Naturf. Kiöbenh.* (1849) 511.—Mexic.
foeniculaceum, *Humb. & Bonpl. Pl. Aequin.* i. 40.—Mexic.; N. Granat.
foeniculaceum, Seem. Bot. Voy. Herald, 197 = oxycarpum.
lacunosum, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 169 = Lophogyne helicandra.
oxycarpum, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 93.—Am. centr.
pauciflorum, *Tul. l. c.* 94.—Guian.
ricciaeforme, *Liebm. in Forhandl. Skand. Naturf. Kiöbenh.* (1849) 512.—Mexic.
Schiedeanum, *Cham. in Linnaea*, xi. (1837) 504. t. 6. —Mexic.
squamosum, *Wedd. in DC. Prod.* xvii. 54.—Bras.
tenue, *Liebm. in Forhandl. Skand. Naturf. Kiöbenh.* (1849) 511.—Mexic.
utile, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 95.—N. Granat.

MARATHRUM, Link, Handb. i. 348 (1829) = **Seseli**, Linn. (Umbellif.).
elatum, Link, l. c. = S. elatum.
glaucum, Link, l. c. = S. montanum.
montanum, Link, l. c. = S. montanum.
tortuosum, Link, l. c. = S. tortuosum.

MARATHRUM, Rafin. in Journ. Phys. lxxxix. (1819) 101 = **Musenium**, Nutt. (Umbell.).
divaricatum, Rafin. l. c. = Mus. divaricatum.

MARCANILLA, Steud. Nom. ed. II. i. 769, in syn. (1840) sphalm. = MANÇANILLA, Adans. = **Hippomane**, Linn. (Euphorb.).

MARCANTHUS, Lour. Fl. Cochinch. 460 (1790) = **Mucuna**, Adans. (Legumin.).
cochinchinensis, Lour. l. c.—Cochinch.
guadalupensis, Spreng. Syst. iii. 248.—Ins. Guadalup.

MARCELIA, Cass. in Dict. Sc. Nat. xxxiv. 107 (1825) = **Anthemis**, Linn. (Compos.).
aurea, Cass. l. c. = A. aurea.

MARCELLIA, Mart. ex Choisy, in Mém. Soc. Phys. Genèv. x. (1844) 443 = **Ipomoea**, Linn. (Convolv.).
villosa, Choisy, l. c. = I. villosa, *Meissn.*

MARCETIA, DC. Prod. iii. 124 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 744.
acerosa, *DC. l. c.* 125.—Bras.
andicola, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 44 = cordigera.
bracteolaris, *Cogn. in Mart. Fl. Bras.* xiv. iii. 441.—Bras.
caespitosa, A. St. Hil. Voy. Distr. Diam. i. 375 = Fritzschia erecta.
canescens, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 44.—Bras.
carinata, *Naud. l. c.* 47.—Bras.
cinerea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 50. —Bras.
cordigera, *DC. Prod.* iii. 124.—Bras.
decussata, *DC. l. c.*—Bras.
decussata, Maund, Bot. v. t. 223 = cordigera.
denudata, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 46.—Bras.
disticha, *Cogn. in Mart. Fl. Bras.* xiv. iii. 447.—Bras.
ericoides, Cogn. l. c. 456 = carinata.

MARCETIA:—
excoriata, Cham. in Linnaea, ix. (1834) 401 = cordigera.
excoriata, *DC. Prod.* iii. 124.—Bras.
excoriata, Miq. in Linnaea, xxii. (1849) 538 = Comolia lanceaeflora.
fastigiata, *Cogn. in Mart. Fl. Bras.* xiv. III. 452.—Bras.
Gardneri, *Cogn. l. c.* 448.—Bras.
glandulosa, DC. Prod. iii. 124 = taxifolia.
Glazioviana, *Cogn. in Mart. Fl. Bras.* xiv. III. 449.—Bras.
gracillima, *Cogn. l. c.* 455.—Bras.
hirsuta, *Cogn. l. c* 451.—Bras.
juniperina, *DC. Prod.* iii. 125.—Bras.
latifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 44.—Bras.
pubescens, DC. Prod. iii. 124 = taxifolia.
Sertularia, DC. l. c. 125 = Comolia Sertularia.
tamariscina, *DC. l. c.*—Bras.
taxifolia, *DC. l. c.* 124.—Bras.
taxifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 50 = cordigera.
tenuifolia, *DC. Prod.* iii. 125.—Bras.
tetrasticha, *Cogn. in Mart. Fl. Bras.* xiv. III. 457.—Bras.

MARCGRAAVIA, Griseb. in Abh. Goett. vii. (1854) 184 = seq.

MARCGRAFIA, Gled. Syst. 231 (1764) = seq.

MARCGRAVIA, Plum. ex Linn. Gen. ed. I. 363 (1737). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 181.
JATROPS, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 293.
acuminata, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 37 = coriacea.
affinis, *Hemsl. Diagn. Pl. Nov.* 3.—Costa Rica.
angustifolia, Pav. ex Wittm. in Mart. Fl. Bras. xii. I. 224 = polyantha.
cacabifera, Ruiz & Pav. ex G. Don, Gen. Syst. i. 625 = Norantea cacabifera.
caudata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 362.—N. Granat.
comosa, Presl, Bot. Bemerk. 23 = myriostigma?
cordachida, Ruiz & Pav. ex G. Don, Gen. Syst. i. 625 = Norantea cordachida.
coriacea, *Vahl, Eclog. Am.* ii. 39.—Am. trop.
crenata, *Poepp. ex Wittm. in Mart. Fl. Bras.* xii. I. 223.—Peruv.
cuneifolia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 335 = Norantea cuneifolia.
cuspidata, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 370 = coriacea.
cyrtogastra, Delp. in Atti Soc. Ital. Sc. Nat. Milan. xii. (1869) 183, 212 = umbellata.
cyrtonota, Delp. l. c. = coriacea.
dubia, H. B. & K. Nov. Gen. et Sp. vii. 217 = umbellata.
Eichleriana, *Wittm. in Mart. Fl. Bras.* xii. I. 230.—Bras.
flagellaris, Poepp. ex Wittm. l. c. 223 = rectiflora.
gracilis, *Sagot, in Ann. Sc. Nat.* Sér. VI. xi. (1881) 170.—Guian.
macrocalyptra, Delp. in Atti. Soc. Ital. Sc. Nat. Milan. xii. (1869) 183, 211 = rectiflora.
macrocarpa, Pav. ex Wittm. in Mart. Fl. Bras. xii. I. 223 = oblongifolia.
macrocarpa, Ruiz & Pav. ex G. Don, Gen. Syst. i. 625 = rectiflora.
micrantha, Pav. ex Wittm. in Mart. Fl. Bras. xii. I. 249 = Ruyschia sphaeradenia.
monogyna, Pav. ex Delp. in Nuov. Giorn. Bot. i. (1869) 282 = Norantea cacabifera.
myriostigma, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 369.—N. Granat.
nepenthoides, *Seem. in Journ. Bot.* viii. (1870) 245.—Nicaragua.
nervosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 363.—N. Granat.
oblongifolia, *Pav. ex Wittm. in Mart. Fl. Bras.* xii. I. 223.—Peruv.
oblongifolia, Pav. ex Wittm. l. c. 224 = polyantha.
obovata, Ruiz & Pav. ex G. Don, Gen. Syst. i. 625 = Norantea obovata.

MARCGRAVIA:—
octandra, Pav. ex Wittm. in Mart. Fl. Bras. xii. I. 241 = Norantea cacabifera.
oligandra, *Wright, ex Griseb. Cat. Pl. Cub.* 39.—Cuba.
paradoxa, Bull. Cat. (1872) 7; et in Gard. Chron. (1877) II. 13. f. 1, 2 = Monstera tenuis.
parviflora, *Rich. ex Wittm. in Mart. Fl. Bras.* xii. I. 227.—Am. trop.
pedunculosa, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 371 = parviflora.
pentandra, Pav. ex Delp. in Nuov. Giorn. Bot. i. (1869) 282 = Norantea cacabifera.
pentandra, Pav. ex Wittm. in Mart. Fl. Bras. xii. I. 240 = Norantea anomala.
pentandra, Pav. ex Wittm. l. c. 249 = Ruyschia sphaeradenia.
picta, *Willd. in Ges. Nat. Fr. Berl. Mag.* ii. (1808) 172.—Bras.
polyantha, *Delp. in Atti Soc. Ital. Sc. Milan.* xii. (1869) 182, 210.—Bras.; Mexic.
rectiflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 364.—Peruv.
spiciflora, Juss. in Ann. Mus. Par. xiv. (1809) 402. t. 25 = Norantea Jussiaei.
Trianae, *Baill. Adansonia*, x. (1872) 244.—Venezuela.
Trinitatis, Presl, Bot. Bemerk. 122 = umbellata.
umbellata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 167 = oligandra.
umbellata, Jacq. Select. Am. 156, partim = rectiflora.
umbellata, *Linn. Sp. Pl.* 503.—Am. trop.
umbellata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Hornemannia martinicensis.
Wrightii, Seem. in Journ. Bot. viii. (1870) 246 = oligandra.

MARCIELIA, Steud. Nom. ed. II. ii. 101 (1841) = MACIELIA, Vand. = **Cordia**, Linn. (Borag.).
Vandellii, Steud. l. c. (= *Macielia racemifera*).—Bras.

MARCKEA, Rich. in Dict. Class. x. 168 (1826) = **Markea**, Rich. (Solan.).

MARCORELLA, Neck. Elem. ii. 122 (1790) = **Colubrina**, Rich. (Rhamnac.).
Colubrina, Rafin. Sylva Tellur. 31 = C. ferruginosa.
cubensis, Rafin. l. c. = Ceanothus reclinatus?

MARCUCCIA, Becc. in Nuov. Giorn. Bot. iii. (1871) 181. *ANONACEAE*.
grandiflora, *Becc. l. c.* 183.—Borneo.

MARENGA, Endl. Gen. 223, in syn. (1837) = MAROGNA, Salisb. = **Amomum**, Linn. (Scitam.).

MARENTERIA, Thou. Gen. Nov. Madag. 18 (1806) = **Uvaria**, Linn. (Anonac.).

MARESIA, Pomel, Nouv. Mat. Fl. Atl. 228 (1874) = **Sisymbrium**, Tourn. (Crucif.).
binervis, Pomel, l. c. = Malcomia nana.
malcomioides, Pomel, l. c. = S. malcolmioides.

MAREYA, Baill. Adansonia, i. (1860–61) 73. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
leonensis, Baill. Hist. Pl. v. 213 = spicata.
micrantha, Muell. Arg. in DC. Prod. xv. II. 792 = spicata.
spicata, *Baill. Adansonia*, i. (1860–61) 74.—Afr. trop.

MARGACOLA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457 = **Trichocoronis**, A. Gray (Compos.).
parvula, Buckl. l. c. = T. Wrightii.

MARGARANTHUS, Schlecht. Ind. Sem. Hort. Hal. (1838).—Cf. Linnaea, xiii. (1839) Litt. 99. *SOLANACEAE*, Benth. & Hook. f. ii. 893.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 91.—Arizona.
solanaceus, *Schlecht. Ind. Sem. Hort. Hal.* (1838).—*Cf. Linnaea*, xiii. (1839) *Litt.* 99.—Texas.
tenuis, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 250 = solanacens.

MARGARETTA, Oliver, in Trans. Linn. Soc. xxix. (1875) 111. t. 76. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 759.
rosea, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 111.—Afr. trop.

MARGARIPES, DC. ex Steud. Nom. ed. II. ii. 101 (1841) = **Anaphalis**, DC. (Compos.).

MARGARIS, DC. Prod. iv. 483 (1830) = **Symphoricarpos**, Dill. (Caprifol.).
barbigera, DC. l. c. = S. microphyllus.
nudiflora, DC. l. c.—Mexic.

MARGARIS, Griseb. Cat. Pl. Cub. 134 = **Margaritopsis**, Wright (Rubiac.).
nudiflora, Griseb. l. c. = Margaritopsis acuifolia.

MARGARITA, Gaud. Fl. Helv. v. 335 (1829) = **Aster**, Tourn. (Compos.).
Bellidiastrum, Gaud. l. c. 336 = A. Bellidiastrum.

MARGARITARIA, Linn. f. Suppl. 66 (1781), pro parte = **Phyllanthus**, Linn. (Euphorb.).
adelioides, Rich. ex Baill. Etud. Gén. Euph. 618, nomen = P. antillanus.
nobilis, Linn. f. Suppl. 428, quoad plantam ♀ (♂ est Combretacea) = P. antillanus.

MARGARITARIA, Opiz, Seznam, 63 (1852) = **Anaphalis**, DC. (Compos.).
ramosa, Opiz, l. c. = A. margaritacea.
simplex, Opiz, l. c. = Antennaria dioica.

MARGARITOPSIS, Wright, in Sauv. Fl. Cub. 68 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 133.
MARGARIS, Griseb. Cat. Pl. Cub. 134 (1866).
acuifolia, *Wright, in Sauv. Fl. Cub.* 69.—Cuba.

MARGAROCARPUS, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 203 = **Pouzolzia**, Gaudich. (Urtic.).
asper, Wedd. l. c. 204 = P. aspera.
Boiviniauus, Wedd. l. c. = P. Boiviniana.
cinerascens, Buek, Ind. ad DC. Prod. iv. 215 = P. cinerascens.
conglobatus, Buek, l. c. = P. conglobata.
exasperatus, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 204 = P. viminea.
heterophyllus, Wedd. l. c. = Boehmeria Blumei.
laevis, Wedd. l. c. = P. virgata.
obliquus, Wedd. l. c. = P. obliqua.
ovatus, Wedd. l. c. = P. ovata.
Poeppigianus, Wedd. l. c. = P. discolor.
procridioides, Wedd. l. c. = P. procridioides.
Schimperianus, Wedd. l. c. 205 = P. abyssinica.
vimineus, Wedd. l. c. 203 = P. viminea.

MARGAROSPERMUM, Spach, Hist. Vég. Phan. ix. 31 (1840); Decne. in Jacquem. Voy. Bot. 122 (1844) = **Lithospermum**, Linn. (Borag.).
apulum, Decne. l. c. = L. apulum.
arvense, Decne. l. c. = L. arvense.
fruticosum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 122 = L. fruticosum.
latifolium, Decne. in Jacquem. Voy. Bot. 122 = L. latifolium.
officinale, Decne. l. c. = L. officinale.
purpureo-caeruleum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 122 = L. purpurocaeruleum.

MARGGRAVIA, Willd. in Ges. Nat. Fr. Berl. Mag. ii. (1808) 172 = **Marcgravia**, Plum. (Ternstr.).

MARGOTIA, Boiss. Elench. 52 (1838) = **Elaeoselinum**, Koch (Umbell.).
gummifera, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 25 = laserpitioides.
laserpitioides, Boiss. Elench. 52.—Hispan.

MARGYRICARPUS, Ruiz & Pav. Prod. 7. t. 33 (1794). *ROSACEAE*, Benth. & Hook. f. i. 623.
TETRAGLOCHIN, Kunze, ex Poepp. Fragm. Synops. 26 (1833).
alatus, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 305.—Chili.
Clarazii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 217.—Patagon.
imberbis, *Presl, Epim. Bot.* 200.—Peruv.

MARGYRICARPUS:—
laevis, Willd, in Ges. Nat. Fr. Neue Schr. iii. (1801) 437 = setosus.
linearis, Steud. Nom. ed. II. ii. 101 = alatus.
microphyllus, *Niederl. in Roca, Exped. Rio-Negro*, ii. 215.—Am. austr.
prolifer, Steud. Nom. ed. II. ii. 101 = alatus.
setosus, *Ruiz & Pav. Fl. Per.* i. 28. t. 8. f. d.—Am. austr.

MARIA-ANTONIA, Parl. Mar. Ant. Nov. Gen. 3 (1844) = **Crotalaria**, Linn. (Legumin.).
orientalis, Parl. l. c. 5 = C. microcarpa.

MARIALVA, Vand. Fl. Lusit. 37 (1788) = **Tovomita**, Aubl. (Guttif.).
amazonica, Poepp. & Endl. Nov. Gen. et Sp. iii. 13. t. 212 = T. amazonica.
brasiliensis, Mart. Nov. Gen. et Sp. ii. 83 = T. brasiliensis.
calyptrata, Steud. Nom. ed. II. ii. 101 = T. calyptrata.
elliptica, Stahl, Estud. Fl. Puerto-Rico, Foll. ii. 126.—Porto Rico.
fructipendula, Choisy, in DC. Prod. i. 560 = T. fructipendula.
guianensis, Choisy, l. c. = T. guianensis.
macrophylla, Poepp. & Endl. Nov. Gen. et Sp. iii. 14 = T. macrophylla.
microcarpa, Poepp. & Endl. l. c. = T. microcarpa.
paniculata, Steud. Nom. ed. II. ii. 101 = T. paniculata.
uniflora, Choisy, in DC. Prod. i. 560 = T. uniflora.

MARIALVAEA, Mart. Nov. Gen. et Sp. ii. 82 (1827) = praec.

MARIALVEA, Spreng. Anleit. ii. II. 783 (1818) = praec.

MARIANA, Hill, Veg. Syst. iv. 19 (1762) = **Silybum**, Gaertn. (1788) (Compos.).
lactea, Hill, Herb. Brit. i. 75 = S. Marianum.
Mariana, Hill, Hort. Kew. 61 = S. Marianum.

MARIANTHEMUM, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 34 = **Campanula**, Linn.
aggregatum, Schrank, in Flora, vii. (1824) II. Beil. 54 = C. glomerata.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = C. alpina.
barbatum, Schur, l. c. = C. barbata.
betonicaefolium, Schrank, in Flora, vii. (1824) II. Beil. 54 = C. sarmatica.
divergens, Schur, Enum. Pl. Transs. 433 = C. sibirica.
Grossekii, Schur, l. c. 936 (Index) = C. Grossekii.
lingulatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = C. lingulata.
longifolium, Schrank, in Flora, vii. (1824) II. Beil. 54 = C. speciosa.
Medium, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = C. Medium.
pannonicum, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 34 = C. sibirica.
sibiricum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = C. sibirica.
spathulatum, Schur, l. c. = C. sibirica.

MARIANTHUS, Hueg. in Enum. Pl. Hueg. 8 (1837). *PITTOSPOREAE*, Benth. & Hook. f. i. 132.
CALOPETALON, Harv. in Hook. Kew Journ. vii. (1855) 52.
ONCOSPORUM, Putterl. Syn. Pittosp. 21 (1839).
RHYTIDOSPORUM, F. Muell. in Hook. f. Fl. Tasm. i. 39 (1855).
angustifolius, Putterl. in Lehm. Pl. Preiss. i. 200 = Billardiera Lehmanniana.
bignoniaceus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 6.—Austral.
candidus, *Hueg. in Enum. Pl. Hueg.* 8.—Austral.
coelestis, Putterl. Syn. Pittosp. 23 = Billardiera variifolia.
coerulea-punctatus, *Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Rar.* 28. t. 12.—Austral.
cuneatus, *C. Muell. in Walp. Ann.* vii. 238.—Austral.
Drummondianus, *Benth. Fl. Austral.* i. 119.—Austral.
erubescens, *Putterl. in Endl. Stirp. Nov. Dec.* 60.—Austral.
floribundus, *Putterl. l. c.* 61.—Austral.

MARIANTHUS:—
granulatus, *Benth. Fl. Austral.* i. 118.—Austral.
laxiflorus, *Benth. l. c.* 119.—Austral.
lineatus, *F. Muell. Fragm.* i. 217.—Austral.
microphyllus, *Benth. Fl. Austral.* i. 117.—Austral.
parviflorus, *F. Muell. Fragm.* ii. 144.—Austral.
pictus, *Lindl. Swan River App.* 22.—Austral.
procumbens, *Benth. Fl. Austral.* i. 117.—Austral.
purpureus, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 364 = erubescens.
rhytidosporeus, F. Muell. Fragm. ii. 145 = microphyllus.
ringens, *F. Muell. l. c.* i. 218.—Austral.
tenuis, *Benth. Fl. Austral.* i. 119.—Austral.
venustus, Putterl. Syn. Pittosp. 23 = Billardiera variifolia.
villosus, *Benth. Fl. Austral.* i. 117.—Austral.

MARICA, Ker-Gawl. in Bot. Mag. t. 654 (1803). *IRIDEAE*, Benth. & Hook. f. iii. 689.
CYPELLA, Klatt, in Linnaea, xxxi. (1861–62) 538.
GALATHEA, Liebm. Ind. Sem. Hort. Haun. (1855) 26.
acorifolia, Ker-Gawl. Irid. Gen. 24, excl. syn. = Sisyrinchium convolutum.
acorifolia, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 111.—Mexic.
anceps, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 174 = Sisyrinchium angustifolium.
aphylla, Ker-Gawl. in Bot. Reg. sub t. 229 = Bobartea aphylla.
Bermudiana, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 174 = Sisyrinchium Bermudiana.
bicolor, *Regel, Cat. Pl. Hort. Aksakov.* 90, nomen.—Hab.?
bogotensis, Ker-Gawl. Irid. Gen. 23 = Sisyrinchium bogotense.
brachypus, *Baker, in Journ. Linn. Soc.* xvi. (1877) 150.—Ind. occ.
bulbosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 374.—Bras.
caerulea, *Ker-Gawl. in Bot. Reg.* t. 713.—Bras.
californica, Ker-Gawl. in Bot. Mag. t. 983 = Sisyrinchium californicum.
coelestina, Ker-Gawl. Irid. Gen. 19 = Nemastylis coelestina.
coelestis, *Lehm. Ind. Sem. Hort. Hamb.* (1828) 17.—Bras.
coerulea, Auct. = caerulea.
convoluta, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 174 = Sisyrinchium convolutum.
filiformis, Ker-Gawl. in Bot. Reg. sub t. 229 = Bobartia filiformis.
foliosa, Willd. ex Steud. Nom. ed. I. 509 = Tigridia violacea.
fugax, Willd. Enum. Hort. Berol. Suppl. 4 = Geissorrhiza quadrangula.
gladiata, Ker-Gawl. in Bot. Reg. t. 229 = Bobartea gladiata.
glauca, *Baker, in Journ. Linn. Soc.* xvi. (1877) 149.—Bras.
gracilis, *Herb. in Bot. Mag.* t. 3713.—Am. trop.
Herbertiana, Ker-Gawl. Irid. Gen. 19 = Cypella Herberti.
humilis, Klatt, in Mart. Fl. Bras. iii. 1. 521, sphalm. = gracilis.
humilis, *Lodd. Bot. Cab.* t. 1081.—Bras.
iridifolia, Ker-Gawl. in Bot. Mag. t. 646 = Sisyrinchium iridifolium.
longifolia, *Link & Otto, Ic. Pl. Sel.* 123. t. 58.—Bras.
lutea, *Herb. in Bot. Mag.* t. 3809.—Bras.
martinicensis, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 245 = Trimeza lurida.
micrantha, Ker-Gawl. Irid. Gen. 22 = Sisyrinchium micranthum.
mucronata, Ker-Gawl. l. c. = Sisyrinchium angustifolium.
Northiana, *Ker-Gawl. in Bot. Mag.* t. 654.—Am. trop.
palmifolia, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 246 = Sisyrinchium palmifolium.
pantherina, Salisb. in Trans. Hort. Soc. i. (1812) 308 = M. Northiana.
plicata, Ker-Gawl. in Bot. Mag. t. 655 = Eleutherine plicata.
pusilla, Ker-Gawl. Irid. Gen. 24 = Sisyrinchium pusillum.
Sabini, Lindl. in Trans. Hort. Soc. vi. (1826) 75. t. 1 = caerulea.

MARICA:—
Sabini, Lodd. Bot. Cab. t. 1164 = Northiana.
Sabiniana, Herb. in Bot. Mag. sub t. 3809 = caerulea?
Sabiniana, Lodd. ex Voigt, Hort. Suburb. Calc. 602 = Northiana.
semiaperta, Lodd. Bot. Cab. t. 685 = Trimeza lurida.
spathacea, Ker-Gawl. in Bot. Reg. sub t. 229 = Bobartia indica.
splendens, *Regel, Cat. Pl. Hort. Aksakov.* 90, nomen.—Hab.?
striata, Ker-Gawl. in Bot. Mag. t. 701 = Sisyrinchium striatum.
tenuifolia, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 174. t. 275 = Sisyrinchium tenuifolium.
Tigridia, Lehm. in Ind. Sem. Hort. Hamb. (1828) 17 = Tigridia Pavonia?
tinctoria, Ker-Gawl. Irid. Gen. 23 = Sisyrinchium tinctorium.
variegata, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 112.—Mexic.
Warmingii, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 375.—Bras.
zebrina, *Hoffmgg. Verz. Pfl.* 76, nomen.—Bras.?

MARICA, Schreb. Gen. 37 (1789) = **Cipura**, Aubl. (Irid.).
Cipura, J. F. Gmel. Syst. 118 = C. paludosa.
graminea, Roem. & Schult. Syst. i 450 = C. paludosa.
humilis, Roem. & Schult. l. c. = C. paludosa.
paludosa, Willd. Sp. Pl. i. 246 = C. paludosa.

MARIERA, Walp. Rep. i. 180 (1842), sphalm. = **Moriera**, Boiss. (Crucif.).

MARIGNIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 351 = **Protium**, Burm. f. (Burser.).
acutifolia, Boj. Hort. Maurit. 83 = P. obtusifolium.
acutifolia, DC. Prod. ii. 79 = Canarium rostratum?
madagascariensis, March. ex Engl. in DC. Monog. Phan. iv. 92 = P. madagascariense.
nitida, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 475 = Glycosmis pentaphylla.
obtusifolia, DC. Prod. ii. 79 = P. obtusifolium.

MARILA, Sw. Prod. Veg. Ind. Occ. 84 (1788). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 189.
MONOPORINA, J. Presl, Rostlin, ii. 277 (1825).
SCYPHAEA, C. Presl, Scyph. Nov. Pl. Gen. (1829); Symb. i. 7. t. 4 (1830).
alternifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 271.—N. Granat.
cespedesiana, *Triana & Planch. l. c.* 270.—N. Granat.
dissitiflora, *Wright, in Sauv. Fl. Cub.* 10.—Cuba.
grandiflora, *Griseb. Fl. Brit. W. Ind.* 111.—Ind. occ.
macrophylla, *Benth. Bot. Voy. Sulph.* 72.—Panama.
nitida, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 64.—Bras.
racemosa, *Sw. Prod. Veg. Ind. Occ.* 84.—Ind. occ.
tomentosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 15.—Peruv.

MARINA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 103. *LEGUMINOSAE*, Benth. & Hook. f. i. 492.
gracilis, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 103.—Mexic.

MARIPA, Aubl. Pl. Gui. i. 230. t. 91 (1775). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 869.
MAIRERIA, Scop. Introd. 192 (1777).
MOUROUCOA, Aubl. Pl. Gui. i. 141. t. 54 (1775).
axilliflora, *Mart. ex Meissn. in Mart. Fl. Bras.* vii. 208.—Bras.
cayennensis, *Meissn. l. c.*—Guian.
cordifolia, *Klotzsch, ex Meissn. l. c.* 209.—Guian.
densiflora, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 351.—Bras.
erecta, *G. F. W. Mey. Prim. Fl. Esseq.* 115.—Guiana.
glabra, *Choisy, in DC. Prod.* ix. 327.—Guian.
nicaraguensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 382.—Nicaragua.
panamensis, *Hemsl. l. c.*—Panama.
passifloroides, *Spruce, ex Meissn. in Mart. Fl. Bras.* vii. 207.—Bras.
scandens, *Aubl. Pl. Gui.* i. 230. t. 91.—Guian.
spectabilis, Choisy, in DC. Prod. ix. 327 (= *Prevostea spectabilis*).—Bras.
villosa, *Spreng. Syst.* i. 645.—Bras.

MARISCUS, Gaertn. Fruct. i. 11 (1788); Vahl. Enum. ii. 372 (1806). *CYPERACEAE*, [Benth. & Hook. f. iii. 1044.]

ADUPLA, Bosc. in Jaume St. Hil. Expos. Fam. Nat. i. 65 (1805).

BORABORA, Steud. Syn. Pl. Cyp. 71 (1855).

DICLIDIUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 51 (1842).

EPIPHYSTIS, Trin. Fund. Agrost. 97 (1820).

acicularis, Moench, Meth. 350 = Eleocharis acicularis.
aggregatus, Willd. Enum. Hort. Berol. 70 = flavus.
alatus, Schult. Mant. ii. 144 = elatus.
albescens, *Gaudich. in Freyc. Voy. Bot.* 415.—Geront. trop.
albus, Gilib. Exercit. ii. 512 = Rynchospora alba.
alopecuroides, *Steud. Syn. Pl. Cyp.* 65.—Bras.
alternifolius, Vahl, Enum. ii. 376 = umbellatus.
ambiguus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 231.—Mexic.
aphyllus, Vahl, Enum. ii. 373 = Kyllinga peruviana.
appendiculatus, *Brongn. in Duperr. Voy. Coq. Bot.* 178. t. 34.—Ins. S. Trinit.; Fernando Noronha; Ascension.
asper, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 229.—Mexic.
atrosanguineus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 490 = bulbocaulis.
atrosanguineus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 228.—Mexic.
aureus, Bertol. Misc. Bot. viii. 29. t. 3. f. 2 = cylindricus.
badius, Willd. ex Kunth, Enum. Pl. ii. 123 = paniceus.
Beyrichii, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 48.—Bras.
biglumis, Gaertn. Fruct. i. 12. t. 2. f. 8 = Sieberianus.
bracheilema, Steud. Syn. Pl. Cyp. 317 = ovularis.
brachystachys, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 179.—Ins. Galapag.
bracteosus, *Schlecht. in Linnaea*, xxv. (1852) 164.—Bras.
bulbocaulis, *Hochst. in Flora*, xxvii. (1844) 102.—Abyssin.
bulbocaulis, A. Rich. Tent. Fl. Abyss. ii. 490 = plateilema.
bulbosus, Steud. Syn. Pl. Cyp. 62 = bulbocaulis.
capensis, *Schrad. Anal. Cyp.* 13.—Afr. austr.
capillaris, *Vahl, Enum.* ii. 372.—Ind. occ.
capitatus, Steud. Syn. Pl. Cyp. 60 = Remirea maritima.
cariciformis, *F. Muell. Fragm.* v. 92.—Austral.
castaneus, *Kunze, ex Kunth, Enum. Pl.* ii. 91.—Chili.
Cephalotes, Poepp. ex Kunth, l. c. 23 = Cyperus humilis.
coloratus, *Nees, in Linnaea*, ix. (1834) 286.—Afr. trop.
concinnus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 47 = Sieberianus.
conferto-spicatus, Steud. Syn. Pl. Cyp. 64 = flavus.
confertus, H. B. & K. Nov. Gen. et Sp. i. 213 = flavus.
conflexus, Link, Enum. Hort. Berol. i. 47 = Jacquinii.
conicus, *R. Br. Prod.* 218.—Austral.
coriaceus, G. F. W. Mey. Prim. Pl. Esseq. 38 = rufus.
cornutus, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 151.—Ins. Galapag.
corymbosus, Boeck. in Flora, xli. (1858) 409 = Sieberianus.
cylindricus, *Ell. Sketch*, i. 74.—Am. bor.
cylindristachyus, Steud. Syn. Pl. Cyp. 65 = umbellatus.
cyperiformis, Torr. Cat. Pl. N. York, 14 = glomeratus.
cyperinus, Nees, in Wight, Contrib. 90 = paniceus, Sieberianus.
cyperinus, *Vahl, Enum.* ii. 377.—As. or.; Polynesia.
cyperoides, A. Dietr. Sp. Pl. ii. 348 = Courtoisia cyperoides.
decompositus, *R. Br. Prod.* 218.—Austral.
dilutus, Nees, in Wight, Contrib. 90 = microcephalus.
Dregeanus, *Kunth, Enum. Pl.* ii. 120.—Afr.; As.; Austral.
Drummondii, Steud. Syn. Pl. Cyp. 316 = ovularis.
echinatus, *Ell. Sketch*, i. 75.—Am. bor.
elatus, Kunth, Enum. Pl. ii. 116 = flavus.
elatus, *Vahl, Enum.* ii. 377.—Am. trop.
filiformis, *H. B. & K. Nov. Gen. et Sp.* i. 213.—Am. trop.
filiformis, Roem. & Schult. Syst. ii. 242, partim; Schult. Mant. i. 140 = havanensis.
filiformis, Spreng. Syst. i. 234 = Swartzii.
flabelliformis, *H. B. & K. Nov. Gen. et Sp.* i. 215.—Reg. trop.

MARISCUS:—

flavus, Seem. Fl. Vit. 319 = laevigatus.
flavus, *Vahl, Enum.* ii. 374.—Am. bor. & austr.
foliosissimus, *Steud. Syn. Pl. Cyp.* 65.—Afr. trop.
giganteus, Boeck. in Flora, xlii. (1859) 443 = microcephalus.
glandulosus, Boj. Hort. Maurit. 382 = rufus.
glomeratus, *Barton, Fl. Philad.* 18.—Am. bor.
gracilis, *Vahl, Enum.* ii. 273.—Ind. occ.
Haenkei, *J. & C. Presl, Rel. Haenk.* i. 181.—Calif.; Mexic.
havanensis, *H. B. & K. Nov. Gen. et Sp.* i. 212.—Ind. occ.
Hochstetteri, Walp. Ann. iii. 678 = bulbocaulis.
huarmensis, *H. B. & K. Nov. Gen. et Sp.* i. 214.—Peruv.
humilis, *Zeyh. ex Schrad. Anal. Cyp.* 10.—Afr. austr.
irroratus, Nees, in Hook. Kew Journ. vi. (1854) 28 = Dregeanus.
Jacquinii, *H. B. & K. Nov. Gen. et Sp.* i. 216, in obs.—Am. trop.
Karwinskianus, Kunth, Enum. Pl. ii. 123 = Mutisii.
Kraussii, Hochst. in Flora, xxviii. (1845) 756 = Dregeanus.
Kunthianus, *Gaudich. in Freyc. Voy. Bot.* 415.—Ins. Sandvic.
kyllingiaeformis, Boeck. in Flora, xlii. (1859) 443 = Dregeanus.
kyllingioides, Steud. Syn. Pl. Cyp. 60 = Kyllinga brevifolia.
laevigatus, *Hook. & Arn. Bot. Beech. Voy.* 72.—Ins. Pacif.
laevigatus, Roem. & Schult. Syst. ii. 243 = flavus.
laevis, *R. Br. Prod.* 218.—Austral.
laevis, H. B. & K. Nov. Gen. et Sp. i. 214 = flavus.
latifolius, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 50 = elatus.
leptochloides, Steud. Syn. Pl. Cyp. 317 = Cyperus radiatus.
littoralis, G. F. W. Mey. Prim. Fl. Esseq. 37 = flavus.
longiradiatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 232 = Jacquinii.
longus, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 44.—Bras.
macer, *Kunth, Enum. Pl.* ii. 121.—Afr. austr.
macrocarpus, *Kunth, l. c.* 120.—Afr. austr.
macrophyllus, *Brongn. in Duperr. Voy. Coq. Bot.* 179. t. 33.—Ins. Societ.
maculatus, Roem. & Schult. Syst. ii. 243 = Lipocarpha Humboldtiana.
maritimus, *Miq. Fl. Ind. Bat. Suppl.* 600.—Ins. Banca.
mexicanus, Steud. Syn. Pl. Cyp. 64 = Mutisii.
mexicanus, Willd. ex Link, Jahrb. i. III. 90 = flavus.
Meyenianus, *Nees, in Mart. Fl. Bras.* ii. 1. 49.—Bras.
micans, Nees, l. c. 50 = Mutisii.
microcephalus, J. & C. Presl, Rel. Haenk. i. 182.—Ind. or.; China; Malaya.
mucronatus, Gaertn. Fruct. i. 11. t. 2. f. 5 (Rynchosporae sp.).—Hab.?
mucronatus, Presl, Cyp. & Gram. Sicul. 12 = Cyperus aegyptiacus.
Mutisii, Hook. f. in Trans. Linn. Soc. xx. (1847) 178 = cornutus, brachystachys.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* i. 216. t. 66.—Am. trop.
neglectus, Schult. Mant. ii. 144 = cylindricus.
nossibeensis, *Steud. Syn. Pl. Cyp.* 63.—Afr. trop.
oaxacensis, Steud. l. c. 65 = Mutisii.
oligostachys, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 48.—Bras.
ovularis, Kunth, Enum. Pl. ii. 121, partim = echinatus.
ovularis, Torr. Fl. U. St. i. 58 = cylindricus.
ovularis, *Vahl, Enum.* ii. 374.—Am. bor.
pachystachyus, *Steud. Syn. Pl. Cyp.* 65.—Hab.?
pallens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 230 = Sieberianus.
palustris, *Schrad. in Schult. Mant.* ii. 141.—Bras.
paniceus, *Vahl, Enum.* ii. 373.—Ind. or.; Maurit.
parviflorus, Nees, in Flora, xi. (1828) 329 = Cyperus Luzulae.
patulus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 47 = Jacquinii.
philippensis, *Steud. Syn. Pl. Cyp.* 66.—Ins. Philipp.
phleoides, *Nees, in Linnaea*, ix. (1834) 286.—Ins. Sandvic.

MARISCUS:—

pictus, *Nees, in Wight, Contrib.* 90.—Ind. or.
piluliferus, Bertol. f. in Mem. Acc. Sc. Bolog. v. (1854) 466 = Rynchospora cyperoides.
plateilema, *Steud. in Flora*, xxv. (1842) 596.—Abyss.
Poeppigianus, Kunth, Enum. Pl. ii. 123 = Mutisii.
Pohlianus, *Nees, in Mart. Fl. Bras.* ii. 1. 50.—Bras.
polycephalus, Link, Hort. Berol. i. 324 = Cyperus Luzulae.
polycephalus, A. Rich. Sert. Astrolab. p. lii = appendiculatus.
polyphyllus, *H. B. & K. Nov. Gen. et Sp.* i. 217.—Peruv.
polyphyllus, Steud. in Flora, xxv. (1842) 596 = Sieberianus.
polystachyus, Brongn. ex A. Rich. Sert. Astrolab. p. liv = appendiculatus.
procerus, *A. Rich. Tent. Fl. Abyss.* ii. 489.—Abyssin.
procerus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 46 = elatus.
pubescens, J. & C. Presl, Rel. Haenk. i. 181 = retrofractus.
Pullu, Steud. Syn. Pl. Cyp. 66 = paniceus.
pungens, Steud. l. c. 60 = Remirea maritima.
pycnostachyus, *H. B. & K. Nov. Gen. et Sp.* i. 215. t. 65.—Am. centr.
radiatus, *Hochst. in Flora*, xxviii. (1845) 757.—Afr. austr.
retroflexus, Willd. ex Kunth, Enum. Pl. ii. 121 = ovularis.
retrofractus, Gaertn. Fruct. i. 12. t. 2. f. 5 = uniflorus.?
retrofractus, *Vahl, Enum.* ii. 375.—Am. bor.
rhaphiostachys, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 232.—Nicaragua.
Richardi, Steud. Syn. Pl. Cyp. 66 = procerus.
rigidus, Spreng. Syst. iv. Cur. Post. 29 = flavus.
riparius, *Schrad. in Goett. Gel. Anz.* (1821) III. 2067.—Afr. austr.
rivularis, Walp. Ann. iii. 674, sphalm. = ovularis.
rufus, *H. B. & K. Nov. Gen. et Sp.* i. 216. t. 67.—Afr. et Am. trop.
scaber, *R. Br. Prod.* 218.—Austral.
Schaffnerianus, *Schlecht. in Linnaea*, xxvi. (1853) 379.—Mexic.
Schimperi, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 20, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 491.—Abyss.
serratus, Gilib. Exercit. ii. 512 = Cladium germanicum.
setaceus, Moench, Meth. 350 = Scirpus setaceus.
Sieberianus, *Nees, in Linnaea*, ix. (1834) 286.—Reg. trop.
sphaericus, Willd. ex Kunth, Enum. Pl. ii. 121 = ovularis.
sphaerocephalus, Steud. Syn. Pl. Cyp. 316 = ovularis.
subulatus, *Nees, in Linnaea*, ix. (1834) 286, nomen.—Hab.?
sundaicus, Miq. Fl. Ind. Bat. iii. 289 = cyperinus.
Swartzii, *A. Dietr. Sp. Pl.* ii. 343.—Ind. occ.
sylvestris, H. B. & K. Nov. Gen. et Sp. i. 377 = flavus.
tenuifolius, *Schrad. ex Nees, in Mart. Fl. Bras.* ii. 1. 46.—Ind. or.
Thunbergii, Schrad. Anal. Cyp. 13 = riparius.
toluccensis, Steud. Syn. Pl. Cyp. 65 = flavus.
Tovari, *H. B. & K. Nov. Gen. et Sp.* i. 215.—Venezuela.
tribrachiatus, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 230.—Costa Rica.
uitenhagensis, Steud. Syn. Pl. Cyp. 317 = capensis.
uliginosus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 46 = Meyenianus.
umbellatus, A. Gray, in Perry, Exped. iii. 322 = Sieberianus.
umbellatus, Horne, Fiji, 264 = cyperinus.
umbellatus, J. & C. Presl, Rel. Haenk. i. 181 = philippensis.
umbellatus, Pursh, Fl. Am. Sept. i. 59 = cylindricus, glomeratus.
umbellatus, *Vahl, Enum.* ii. 376.—Afric.
umbellatus, Zoll. Syst. Verz. Ind. Archip. 63, partim = Rynchospora Wallichiana.
uniflorus, *Steud. Syn. Pl. Cyp.* 64.—Texas.
viridis, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 217 = Schimperi.
Wallichianus, Kunth, Ennm. Pl. ii. 117 = paniceus.

MARJORANA, G. Don, Gen. Syst. iv. 766 (1837) = MAJORANA, Tourn. = **Origanum**, Linn. (Labiat.).

MARKEA, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107. *SOLANACEAE*, Benth. & Hook. f. ii. 903.
LAMARKEA, Pers. Syn. i. 218 (1805).
coccinea, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
longiflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 186.—Ind. occ.
neurantha, *Hemsl. Biol. Centr. Am. Bot.* ii. 429.—Costa Rica.

MARKHAMIA, Seem. Journ. Bot. i. (1863) 226 = **Dolichandrone**, Fenzl (Bignon.).
stipulata, Seem. l. c. = D. stipulata.

MARLEA, Roxb. Hort. Beng. [28] (1814); Pl. Corom. iii. 80. t. 283 (1819). *CORNACEAE*, Benth. & Hook. f. i. 949.
DIACICARPIUM, Blume, Bijdr. 657 (1825).
PAUTSAUVIA, Juss. in Mém. Mus. Par. iii. (1817) 443.
PSEUDALANGIUM, F. Muell. Fragm. ii. 84 (1860).
RHYTIDANDRA, A. Gray, Bot. U. St. Expl. Exped. i. 302. t. 28 (1854).
STYLIDIUM, Lour. Fl. Cochinch. 220 (1790).
STYLIS, Poir. Encyc. Suppl. v. 260 (1817).
affinis, Decne. in Jacquem. Voy. Bot. 74. t. 83 = begoniifolia.
barbata, *R. Br. in Wall. Cat.* n. 7129.—Reg. Himal.
begonifolia, *Roxb. Hort. Beng.* [28]; *Pl. Corom.* iii. 80.—Ind. or.; China; Japon.
Bussyana, *Baill. Adansonia*, x. (1872) 183.—N. Caled.
ebenacea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 742.—Malacca.
Griffithii, C. B. Clarke, l. c. = unilocularis.
macrophylla, Sieb. & Zucc. in Abh. Akad. Muench, iv. II. (1843) 135 = platanifolia.
nobilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 743.—Malaya.
platanifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 134.—China; Japan.
tomentosa, Endl. ex Hassk. in Flora, xxvii. (1844) 605 = begonifolia.
unilocularis, *Griff. Notul.* iv. 679.—Malaya.
villosa, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 61.—Java.
virgata, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 157.—Java.
vitiensis, *Benth. Fl. Austral.* iii. 386.—Austral.; Ins. Pacif.

MARLIERA, D. Dietr. Syn. Pl. iii. 86 (1843) = seq.

MARLIEREA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 373. t. 156 (1829). *MYRTACEAE*, Benth. & Hook. f. i. 716.
EUGENIOPSIS, Berg, in Linnaea, xxvii. (1854) 80.
RUBACHIA, Berg, l. c. 11 (1854).
acuminata, *Berg, l. c.* 13.—Guiana.
capitata, *Berg, in Mart. Fl. Bras.* xiv. I. 33.—Bras.
dimorpha, *Berg, l. c.* 538.—Bras.
elliptica, *Griseb. Fl. Brit. W. Ind.* 233.—Ind. occ.
estrellensis, *Berg, in Mart. Fl. Bras.* xiv. I. 34.—Bras.
excoriata, *Mart. in Flora*, xx. (1837) II. *Beibl.* 88.—Bras.
glabra, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 374.—Bras.
glomerata, *Berg, in Linnaea*, xxvii. (1854) 14.—Guiana.
grandifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 37.—Bras.
Langsdorffii, *Berg, l. c.* 536.—Bras.
obscura, *Berg, l. c.* 36.—Bras.
obtusa, *Berg, in Linnaea*, xxvii. (1854) 15.—Guian.
parviflora, *Berg, in Mart. Fl. Bras.* xiv. I. 537.—Bras.
parvifolia, *Berg, l. c.* 33.—Bras.
Regeliana, *Berg, l. c.* 537.—Bras.
resupinata, *Berg, l. c.* 37.—Bras.
Richardiana, *Berg, in Linnaea*, xxx. (1859–60) 650.—Guian.
rufa, *Berg, in Mart. Fl. Bras.* xiv. I. 37.—Bras.
Schomburgkiana, *Berg, in Linnaea*, xxix. (1857–58) 209.—Hab. ?
Schottiana, *Berg, in Mart. Fl. Bras.* xiv. I. 36.—Bras.
sessiliflora, *Berg, l. c.* 34.—Bras.
spathulata, *Berg, l. c.* 32.—Bras.
Spruceana, *Berg, l. c.* 34.—Bras.
strigipes, *Berg, l. c.* 32.—Bras.
suaveolens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 374.—Bras.

MARLIEREA:—
suffruticosa, *Berg, in Linnaea*, xxvii. (1854) 16.—Guian.
tomentosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 373.—Bras.
tovarensis, *Berg, in Linnaea*, xxvii. (1854) 13.—Venezuela.
uaupensis, *Berg, in Mart. Fl. Bras.* xiv. I. 516.—Bras.
umbraticola, *Berg, l. c.* 35.—Venezuela.
verticillaris, *Berg, l. c.* 538.—Bras.

MARLIERIA, Benth. & Hook. f. Gen. i. 716 (1865) err. typ. = praec.

MARMORITES, Benth. in Hook. Bot. Misc. iii. (1833) 377 = **Nepeta**, Linn. (Labiat.).
rotundifolia, Benth. l. c. = N. floccosa.

MAROGNA, Salisb. in Trans. Hort. Soc. i. (1812) 283 = **Amomum**, Linn. (Scitam.).
paludosa, Salisb. l. c. = A. angustifolium.

MAROPSIS, Pomel, Nouv. Mat. Fl. Atl. 121 (1874) = **Marrubium**, Tourn. (Labiat.).
deserti, Pomel, l. c. = Marrubium deserti.

MAROTTIA, Rafin. Sylva Tellur. 155 (1838) = **Hydnocarpus**, Gaertn. (Bixin.).
oleosa, Rafin. l. c. = H. Wightiana.

MARQUARTIA, Hassk. in Flora, xxv. (1842) II. Beibl. 14 = **Pandanus**, Linn.
Billardieri, Walp. Ann. i. 1084, lapsu, loco *Marquisia* = Coprosma Billardieri.
globosa, Hassk. in Flora, xxv. (1842) II. Beibl. 14 = P. utilis.
leucacantha, Hassk. Cat. Hort. Bog. Alt. 61 = P. odoratissimus.

MARQUARTIA, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 35 = **Millettia**, Wight & Arn. (Legumin.).
tomentosa, Vog. l. c. = Millettia nitida.

MARQUISIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 192 = **Coprosma**, Forst. (Rubiac.).
Billardieri, A. Rich. l. c. = C. Billardieri.

MARRHUBIUM, Delarb. Fl. Auv. ed. II. 153 (1800) = MARRUBIUM, Tourn. (Labiat.).

MARRUBIASTRUM, Tourn. ex Moench, Meth. 391 (1794) = **Sideritis**, Tourn. (Labiat.).
elegans, Moench, l. c. = S. lanata.
tomentosum, Moench, l. c. = S. canariensis.

MARRUBIUM, Tourn. ex Linn. Syst. ed. I (1735) = *LABIATAE*, Benth. & Hook. f. ii. 1206.
ANISODONTIUM, Reichb. Handb. 188 (1837).
KRASCHNIKOWIA, Turcz. ex Ledeb. Fl. Ross. iii. 402, in syn. (1849).
LAGOPSIS, Bunge, ex Benth. Lab. Gen. et Sp. 586 (1834).
MAROPSIS, Pomel, Nouv. Mat. Fl. Atl. 121 (1874).
PADOTA, Adans. Fam. ii. 505 (1763).
acetabulosum, Duf. ex Willk. & Lange, Prod. Fl. Hisp. ii. 446 = Ballota hirsuta.
acetabulosum, Linn. Sp. Pl. 584 = Ballota acetabulosa.
adfine, Spreng. Pugill. ii. 68 = seq.
affine, Hornem. Hort. Hafn. 556 = leonuroides.
affine, Host, Fl. Austr. ii. 172 = peregrinum.
africanum, Linn. Sp. Pl. 583 = Ballota africana.
album, Boiss. et Bal. Diagn. Ser. II. iv. 52 = parviflorum.
album, Gilib. Fl. Lituan. i. 87 = vulgare.
alyssoides, *Pomel, Nouv. Mat. Fl. Atl.* 120.—Alger.
Alysson, *Linn. Sp. Pl.* 582.—Reg. Mediterr.
angustifolium, Moench, Meth. 399 = peregrinum.
anisodon, C. Koch, in Linnaea, xxi. (1848) 696 = vulgare.
apulum, Tenore, Prod. Fl. Nap. 34 = vulgare.
aquaticum, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 367 = Mentha aquatica.
astracanicum, Bieb. Fl. Taur. Cauc. ii. 52 = leonuroides.
astracanicum, *Jacq. Ic. Pl. Rar.* i. 11. t. 109.—As. Min.; Pers.
ballotoides, *Boiss. & Bal. Diagn.* Ser. II. iv. 53.—As. Min.

MARRUBIUM:—
Bourgaei, *Boiss. Fl. Orient.* iv. 698.—As. Min.
brachyodon, Boiss. Diagn. Ser. I. vii. 58 = Kotschyi.
candidissimum, Frivald. ex Boiss. l. c. xii. 75 = Friwaldskyanum.
candidissimum, C. Koch, in Linnaea, xxi. (1848) 696 = parviflorum.
candidissimum, *Linn. Sp. Pl.* 583.—Eur. austr.; As. Min.
candidissimum, C. A. Mey. ex Boiss. Fl. Orient. iv. 700 = catariaefolium.
candidissimum, Orsini, ex Tenore, Syll. 292 = peregrinum.
catariaefolium, *Desr. in Lam. Encyc.* iii. 717.—Reg. Cauc.; As. Min.
catariaefolium, Reichb. Pl. Crit. iv. 10. t. 313 = candidissimum.
cephalanthum, *Boiss. & Noé, Diagn.* Ser. II. iv. 50.—As. Min.
cinerascens, Fisch. ex Steud. Nom. ed. I. 510, nomen (Quid ?).—Hab. ?
cinereum, Desr. in Lam. Encyc. iii. 719 = Ballota hirsuta.
cinereum, Hort. ex Benth. Lab. Gen. et Sp. 595 = Ballota mollissima.
circinnatum, Desr. in Lam. Encyc. iii. 717 = velutinum.
coerulescens, Desf. Cat. Hort. Par. ed. III. 395 = propinquum ?
condensatum, *Boiss. Diagn.* Ser. II. iv. 51; *Fl. Orient.* iv. 697.—Mesopot.
crassidens, *Boiss. Diagn.* Ser. I. v. 35.—Persia.
creticum, Mill. Gard. Dict. ed. VIII. n. 3 = peregrinum.
creticum, Roth, Tent. Fl. Germ. i. 257 = pannonicum.
crispum, Desr. in Lam. Encyc. iii. 719 = Ballota hirsuta.
crispum, Linn. Sp. Pl. ed. II. 1674 = Ballota africana.
crispum, Sieber, ex Boiss. Fl. Orient. iv. 773 = Ballota undulata.
cuneatum, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 255.—Syria; Mesopot.
cylleneum, Boiss. et Heldr. Diagn. Ser. II. iv. 51 = velutinum.
depauperatum, *Boiss. et Bal. l. c.* 49.—Cappadoc.
deserti, *Noé, ex Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 472, 490.—Alger.
echinatum, *Ball, in Journ. Bot.* xiii. (1878) 175.—Marocco.
eriostachyum, *Benth. Lab. Gen. et Sp.* 586.—Sibir.
faucidens, *Boiss. & Bal. Diagn.* Ser. II. iv. 50.—As. Min.
flavum, *Walp. Rep.* iii. 856.—Mont. Alatau.
flexuosum, Moench, Meth. 399 = candidissimum.
Friwaldskyanum, *Boiss. Diagn.* Ser. I. xii. 74.—Rumel.
gamodon, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 48.—Persia.
germanicum, Schrank, ex Steud. Nom. ed. I. 510 = vulgare.
globosum, *Montbr. et Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 53.—As. Min.
hamatum, H. B. et K. Nov. Gen. et Sp. ii. 310 = vulgare.
Hermonis, Boiss. Diagn. Ser. I. xii. 74 = libanoticum.
heterodon, *Boiss. & Bal. l. c.* II. iv. 52.—As. Min.
hirsutum, Willd. Sp. Pl. iii. 113 = Ballota hirsuta.
hispanicum, Desf. Cat. Hort. Par. ed. III. 102 = Ballota hirsuta.
hispanicum, Linn. Sp. Pl. 583 = Ballota rupestris.
humile, Desf. Cat. Hort. Par. ed. III. 395 = Ballota saxatilis.
incanum, Desr. in Lam. Encyc. iii. 76 = candidissimum.
incisum, *Benth. Lab. Gen. et Sp.* 586.—Sibir.
indicum, Blanco, Fl. Filip. ed. I. 477 = Hyptis suaveolens.
indicum, Burm. f. Fl. Ind. 127 = Anisomeles ovata.
Kotschyi, *Boiss. et Hohen. Diagn.* Ser. I. v. 33.—Kurdist.
lanatum, *Benth. Lab. Gen. et Sp.* 587.—Sibir. altaic.
laurifolium, Hort. Par. ex Steud. Nom. ed. I. 510 = Ballota africana.
leonuroides, *Desr. in Lam. Encyc.* iii. 715.—Reg. Caucas.
libanoticum, *Boiss. Diagn.* Ser. I. xii. 73.—Syria.
lutescens, *Boiss. et Heldr. l. c.* v. 34.—Graecia.
Malcolmianum, Dalz. in Hook. Kew Journ. iv. (1852) 109 = Micromeria capitellata.
Marrubiastrum, Ssowinski, ex Trautv. in Act. Hort. Petrop. ix. (1884) 103 = Leonurus Marrubiastrum.

MARRUBIUM:—
micranthum, *Boiss. et Heldr. Diagn.* Ser. I. xii. 78.—As. Min.
microphyllum, Desr. Lam. Encyc. iii. 720 = Otostegia Schimperi.
mollissimum, D. Don, Prod. Fl. Nep. 112 = Leucas lanata.
nigrum, Crantz, Stirp. Austr. ed. II. fasc. iv. 272 = Ballota nigra.
Nočanum, Boiss. Fl. Orient. iv. 696 = Kotschyi.
odoratissimum, Pourr. ex Steud. Nom. ed. I. 510 = candidissimum.
orientale, Spreng. Syst. ii. 740 (= *Ballota orientalis*).—As. Min.
pallidum, Salisb. Prod. 84 = Ballota Pseudo-dictamnus.
paniculatum, Desr. in Lam. Encyc. iii. 716 = peregrinum.
pannonicum, *Reichb. Fl. Germ. Excurs.* 325.—Reg. Caucas.
parnassicum, Reut. ex Nym. Consp. 582 = velutinum.
parviflorum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 33.—As. Min.; Persia.
pauciflorum, Wallr. Sched. Crit. 295 = peregrinum.
peregrinum, Asso, Syn. Stirp. Arrag. 76 = supinum.
peregrinum, *Linn. Sp. Pl.* 582.—Europ.; As. Min.
peregrinum, C. A. Mey. Verz. Pfl. Cauc. 96 = parviflorum.
peregrinum, Reichb. Pl. Crit. iv. 10. t. 312 = candidissimum.
peregrinum, Spreng. Fl. Hal. 172. t. 6 = pannonicum.
persicum, *C. A. Mey. Verz. Pfl. Cauc.* 95.—Reg. Cauc.
Pestalozzae, Boiss. Diagn. Ser. II. iv. 53 = praecox.
plicatum, Forsk. Fl. Aegypt. Arab. 213 = M. Alysson.
plumosum, C. A. Mey. Verz. Pfl. Cauc. 96 = leonuroides.
polyodon, *Boiss. Diagn.* Ser. I. xii. 75.—Persia.
praecox, Janka, in Oestr. Bot. Zeitschr. xxv. (1875) 63 = peregrinum.
procerum, *Bunge, Lab. Pers.* 66.—Persia.
propinquum, Benth. in DC. Prod. xii. 450 = vulgare.
propinquum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 33.—Reg. Caucas.; Pers.
Pseudo-alyssum, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 34 = micranthum.
Pseudo-dictamnus, Linn. Sp. 583 = Ballota Pseudo-dictamnus.
Pseudo-dictamnus, Sibth. et Sm. Fl. Graec. vi. 50. t. 562 = Ballota acetabulosa.
purpureum, *Bunge, Lab. Pers.* 67.—Persia.
radiatum, Delile, ex Benth. Lab. Gen. et Sp. 591 = cuneatum.
remotum, Janka, in Oestr. Bot. Zeitschr. xxv. (1875) 62 = peregrinum.
remotum, Kit. ex Schult. Oesterr. Fl. ed. II. ii. 161 = pannonicum.
rotundifolium, *Boiss. Diagn.* Ser. I. v. 33.—As. Min.
rubrum, Roth, Nov. Pl. Sp. 259 = pannonicum.
rugosum, [Soland. in] Russ. Aleppo, ed. II. ii. 255; Desf. Cat. Hort. Par. ed. III. 395 = Ballota saxatilis.
rupestre, Bivona, Stirp. Rar. Sicil. ii. [5]. t. 1 = Ballota rupestris.
saxatile, Rafin. Préc. Découv. 38 = Ballota saxatilis.
scrophulariaefolium, Hort. ex Benth. Lab. Gen. et Sp. 595 = Ballota mollissima.
scrophulariaefolium, Lag. ex Benth. l. c. 597 = Ballota rupestris.
sericeum, Boiss. Elench. 77 = supinum.
setaceum, Desr. in Lam. Encyc. iii. 717 = candidissimum.
suffruticosum, Mill. Gard. Dict. ed. VIII. n. 9 = Ballota acetabulosa.
supinum, *Linn. Sp. Pl.* 583.—Europ. austr.
supinum, Scop. Fl. Carn. ed. II. i. 414 = candidissimum.
thessalum, *Boiss. & Heldr. Diagn.* Ser. II. iv. 50.—Graecia.
Thouini, Schult. ex Weinm. in Syll. Ratisb. ii. (1828) 23 = Ballota africana.
trachyticum, *Boiss. Fl. Orient.* iv. 698.—Anatol.
uncinatum, Hornem. Hort. Hafn. 968 = candidissimum.
undulatum, Sieber, ex Fresen. in Mus. Senckenb. i. (1834) 92 = Ballota undulata.
Vaillantii, *Coss. & Germ. in Ann. Sc. Nat.* Sér. II. xx. (1843) 293.—Europ.
velutinum, *Sibth. & Sm. Fl. Graec. Prod.* i. 412.—Graecia.
vulgare, *Linn. Sp. Pl.* 583.—Europ.; As.; Afr. bor.

MARSANA, Sonner. Voy. Ind. Or. iii. 282 (1782) = **Murraya**, Linn. (Rutac.).
buxifolia, Sonner. l. c. = M. exotica.

MARSCHALLIA, Bartl. Ord. 142 (1830) = **Marshallia**, Schreb. (Compos.).

MARSDENIA, R. Br. in Mem. Wern. Soc. i. (1809) 28. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 772.
BAXTERA, Reichb. Consp. 131 (1828).
CHLOROCHLAMYS, Miq. in Ann. Mus. Lugd. Bat. iv. (1848) 142.
CIONURA, Griseb. Spicil. Fl. Rumel. ii. 69 (1844).
HARRISONIA, Hook. Bot. Mag. t. 2699 (1826).
KOELREUTERIA, Medic. Beobacht. 1782 (1783) 22.
LEICHHARDTIA, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 81 (1849).
RUEHSSIA, Karst. in Verh. Ver. Gartenb. Preuss. xix. (1849) II. 304.
SICYOCARPUS, Boj. Hort. Maurit. 214 (1837).
SPHINCTOSTOMA, Benth. ex Benth. & Hook. f. Gen. ii. 772 (1876).
TETRAGONOCARPUS, Hassk. in Flora, xx. (1857) 99.
affinis, A. Rich. Fl. Cub. Fanerog. ii. 100 = clausa.
agglomerata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 276.—Ins. S. Doming.
Akkar, Blanco, Fl. Filip. ed. I. 118 = tinctoria.
angustifolia, Wight, Contrib. 40 = Lygisma angustifolia.
araujacea, *F. Muell. Fragm.* vi. 135.—Austral.
Billardieri, *Decne. in DC. Prod.* viii. 615.—N. Caled.
brasiliensis, *Decne. l. c.* 617.—Bras.
Brunoniana, *Wight & Arn. in Wight, Contrib.* 40.—Ind. or.
Burchellii, *Fourn. in Mart. Fl. Bras.* vi. IV. 322.—Bras.
Calesiana, *Wight, Contrib.* 41.—Reg. Himal.
campanulata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 521.—Cuba.
cinerascens, *R. Br. in Mem. Wern. Soc.* i. (1809) 30.—Austral.
clausa, *R. Br. l. c.*—Jamaic.
coronata, *Benth. Fl. Austral.* iv. 341.—Austral.
Coulteri, *Hemsl. Biol. Centr. Am. Bot.* ii. 336.—Mexic.
crassipes, *Hemsl. l. c.* 337.—Panama.
cubensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 322.—Cuba.
Cundurango, *Nichols. Dict. Gard.* ii. 329.—N. Granat.
Cundurango, Reichb. f. in Bot. Zeit. xxx. (1872) 552 = Reichenbachii.
cymulosa, *Benth. Fl. Austral.* iv. 338.—Austral.
elliptica, *Decne. in DC. Prod.* viii. 616.—Ind. occ.
erecta, *R. Br. in Mem. Wern. Soc.* i. (1809) 31.—Syria.
eriocarpa, *Hook. f. Fl. Brit. Ind.* iv. 35.—Reg. Himal.
flavescens, *A. Cunn. in Bot. Mag.* t. 3289.—Austral.
Fraseri, *Benth. Fl. Austral.* iv. 339.—Austral.
fusca, *Wright, ex Griseb. Cat. Pl. Cub.* 178.—Cuba.
glabriflora, *Benth. in Hook. Niger Fl.* 455.—Afr. trop.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 36.—Reg. Himal.
Hamiltonii, *Wight, Contrib.* 41.—Ind. or.
Hilariana, *Fourn. in Mart. Fl. Bras.* vi. IV. 322.—Bras.
hemiptera, *Reichb. f. Beitr. Syst. Pfl.* 73.—Austral.
Hullsii, *F. Muell. ex Benth. Fl. Austral.* iv. 338.—Austral.
Jenkinsii, *Hook. f. Fl. Brit. Ind.* iv. 36.—Reg. Himal.
lachnostoma, *Benth. Fl. Hongk.* 226.—Ins. Hongkong.
Leichhardtiana, *F. Muell. Fragm.* v. 160.—Austral.
leonensis, *Benth. in Hook. Niger Fl.* 455.—Afr. trop.
leptophylla, *F. Muell. ex Benth. Fl. Austral.* iv. 340.—Austral.
linearis, *Decne. in DC. Prod.* viii. 617.—Ins. S. Doming.
longiflora, *A. Rich. Fl. Cub. Fanerog.* ii. 100.—Cuba.
longiloba, *Benth. Fl. Austral.* iv. 340.—Austral.
loniceroides, *Fourn. in Mart. Fl. Bras.* vi. IV. 323.—Bras.
lucida, *Edgew. ex Madden, in Journ. As. Soc. Beng.* xvii. I. (1848) 370.—Reg. Himal.
lucida, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 37 = Griffithii.
macrophylla, *Fourn. in Mart. Fl. Bras.* vi. IV. 321.—Bras.
maculata, *Hook. Bot. Mag.* t. 4299.—Am. trop.
mexicana, *Decne. in DC. Prod.* viii. 617.—Mexic.
microlepis, *Benth. Fl. Austral.* iv. 342.—Austral.

MARSDENIA:—
mollissima, *Fourn. in Mart. Fl. Bras.* vi. IV. 322.—Bras.
monostachya, Wall. ex Hook. f. Fl. Brit. Ind. iv. 34 = tinctoria.
nitida, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 275.—Ins. S. Doming.
obovata, Turcz. in Bull. Soc. Nat. Mosc. xxv. II. (1852) 321 = clausa.
ovata, *Fourn. in Mart. Fl. Bras.* vi. IV. 321, in obs.—Guiana.
parasita, *Blanco, Fl. Filip.* ed. I. 120.—Ins. Philipp.
pauciflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 321.—Cuba.
picta, *Decne. in DC. Prod.* viii. 615.—Ins. S. Doming.
propinqua, *Hemsl. Biol. Centr. Am. Bot.* ii. 337.—Mexic.
Reichenbachii, *Triana, in Bull. Soc. Bot. Fr.* xx. (1873) 36.—Ecuador.
rhyncholepis, *F. Muell. Fragm.* xi. 78.—Austral.
robusta, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 79.—Ins. Socotr.
rostrata, *R. Br. in Mem. Wern. Soc.* i. (1809) 31.—Austral.
Rothii, *G. Don, Gen. Syst.* iv. 131.—Ind. or.
rotundifolia, *Decne. in DC. Prod.* viii. 615.—Ind. or.
Roylei, *Wight, Contrib.* 40.—Reg. Himal.
rubro-fusca, *Fourn. in Mart. Fl. Bras.* vi. IV. 321.—Bras.
satureiaefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 100.—Cuba.
Schimperi, *Decne. in DC. Prod.* viii. 616.—Abyssin.
stenophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) I. 260.—Ins. S. Doming.
suaveolens, *R. Br. in Mem. Wern. Soc.* i. (1809) 30.—Austral.
syringaefolia, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 275.—Malaya.
tagudinia, *Blanco, Fl. Filip.* ed. I. 121.—Ins. Philipp.
tenacissima, *Wight & Arn. in Wight, Contrib.* 41.—Ind. or.; Malaya.
tenuissima, Steud. Nom. ed. II. ii. 103, sphalm. = praec.
thyrsiflora, *Hook. f. Fl. Brit. Ind.* iv. 37.—Ind. or.
tinctoria, *R. Br. in Mem. Wern. Soc.* i. (1809) 30.—Afr. trop.
tinctoria, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 37 = thyrsiflora.
tomentosa, *C. Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 172.—Japon.
tubulosa, *F. Muell. Fragm.* ix. 71.—Austral.
umbellata, *Griseb. Cat. Pl. Cub.* 179.—Cuba.
urceolata, *Decne. in DC. Prod.* viii. 617.—Cochinch.
velutina, *R. Br. in Mem. Wern. Soc.* i. (1809) 29.—Austral.
verrucosa, *Decne. in DC. Prod.* viii. 615.—Madag.
villosa, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 126 = Tylophora villosa.
vinciflora, *Griseb. Cat. Pl. Cub.* 179.—Cuba.
viridiflora, *R. Br. in Mem. Wern. Soc.* i. (1809) 30.—Austral.
Warmingii, *Fourn. in Mart. Fl. Bras.* vi. IV. 323.—Bras.
zimapanica, *Hemsl. Biol. Centr. Am. Bot.* ii. 338.—Mexic.

MARSEA, Adans. Fam. ii. 122 (1763) = **Conyza**, Linn. (Compos.).

MASEMA, Dulac, Fl. Hautes-Pyr. 476 (1867) = **Valerianella**, Tourn. (Valer.).
auriculatum, Dulac, l. c. 477 = V. rimosa.
carinatum, Dulac, l. c. = V. carinata.
dentatum, Dulac, l. c. = V. dentata.
olitorium, Dulac, l. c. 476 = V. olitoria.

MARSESINA, Rafin. Sylva Tellur. 112 (1838) = **Capparis**, Tourn.

MARSHALLIA, J. F. Gmel. Syst. 836 (1791) = **Homalium**, Jacq. (Samyd.).
Lagunac, J. F. Gmel. l. c.—Hab.?

MARSHALLIA, Schreb. Gen. ii. 810 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 392.
PERSOONIA, Michx. Fl. Bor. Am. ii. 104. t. 43 (1803).
PHYTEUMOPSIS, Juss. ex Poir. Encyc. Suppl. iv. 405 (1816).

MARSHALLIA:—
THEROLEPTA, Rafin. Neogenyt. 3 (1825).
TRATTENIKIA, Pers. Syn. ii. 403 (1807).
aliena, Spreng. Syst. iii. 446 = Heterothalamus brunioides.
angustifolia, *Pursh, Fl. Am. Sept.* ii. 520.—Am. bor.
australis, Spreng. ex DC. Prod. v. 672 = Calea Berteriana.
bulbosa, *Rafin. Fl. Ludov.* 61.—Louisian.
caespitosa, *Nutt. ex DC. Prod.* v. 680.—Am. bor. occ.
dentata, *Bertol. Misc. Bot.* xv. 22.—Am. bor.
lanceolata, *Pursh, Fl. Am. Sept.* ii. 519.—Am. bor.
latifolia, *Pursh, l. c.*—Am. bor.
pumila, *Rafin. New Fl. Am.* iv. 78 (Quid?).—Am. bor.
Schreberi, Tratt. Arch. Gew. i. 108 = latifolia.
spiralis, *Rafin. New Fl. Am.* iv. 77 (Quid?).—Am. bor.
tenuifolia, *Rafin. l. c.* (Quid?).—Am. bor.

MARSIPPOSPERMUM, Desv. Journ. Bot. i. (1808) 328. t. 12 = **Rostkovia**, Desv. (Juncac.).
calyculatum, Desv. l. c. = R. grandiflora.
gracile, Buchen. in Bremen, Abh. vi. (1879) 374 = R. gracile, *Hook. f.*
grandiflorum, Hook. Ic. Pl. t. 533 = R. grandiflora.

MARSSONIA, Karst. Fl. Columb. i. 97. t. 48 (1859?) = **Napeanthus**, Gardn. (Gesnerac.).
primulina, Karst. l. c. = N. primulinus.

MARSUPIANTHES, Reichb. Consp. 116 (1828) = seq.

MARSYPIANTHES, Mart. ex Benth. Lab. Gen. et Sp. 64 (1833). *LABIATAE*, Benth. & Hook. f. ii. 1178.
foliolosa, *Benth. in DC. Prod.* xii. 85.—Bras.
glomerata, *Presl, Bot. Bemerk.* 101.—Hab.?
hyptoides, *Mart. ex Benth. Lab. Gen. et Sp.* 64.—Bras.
montana, *Benth. in DC. Prod.* xii. 85.—Bras.
secundiflora, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 185 = hyptoides.
sessiliflora, Presl, Bot. Bemerk. 101 = hyptoides.
squarrosa, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1149, nomen.—Guian.
viscosa, *Klotzsch, l. c.* 1148, nomen.—Guian.

MARSYPIANTHUS, Bartl. Ord. 183 (1830) = praec.

MARSYPOCARPUS, Neck. Elem. iii. 91 (1790) = **Capsella**, Medic. (1794) (Crucif.).

MARSYPOPETALUM, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 342; et in Flora, liii. (1870) 243. *ANONACEAE*.
ceratosanthes, *Scheff. ll. cc.* 343; 244.—Java.

MARSYROCARPUS, Steud. Nom. ed. I. 511 (1821) = MARSYPOCARPUS, Neck.

MARTAGON, Rupp. Fl. Jen. ed. Hall. 148 (1745) = **Lilium**, Tourn.
montanum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 158 = L. Martagon.
sylvaticum, Opiz, Seznam, 63 = L. Martagon.

MARTENSIA, Giseke, Prael. Linn. 199, 207 (1792) = **Alpinia**, Linn. (Scitam.).
aquatica, Raeusch. Nom. ed. III. 2 = A. aquatica.

MARTHA, Fritz Muell. in Bot. Zeit. xxiv. (1866) 129. *RUBIACEAE* (an **Posoqueria**, Aubl.?).
fragrans, *Fritz Muell. l. c.*—Bras.

MARTIA, Benth. in Hook. Journ. Bot. ii. (1840) 146. *LEGUMINOSAE*, Benth. & Hook. f. i. 571.
MARTIUSIA, Benth. in Hook. Journ. Bot. ii. (1840) 84.
excelsa, *Benth. l. c.* 84 (Martiusia).—Guian.
parvifolia, *Benth. l. c.* 103 (Martiusia).—Bras.

MARTIA, Lacerda, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 261 (1862) = **Brunfelsia**, Plum. (Solanac.).
opifera, Lacerda, ex J. A. Schmidt, l. c. = B. Hopeana.

MARTIA, Leandr. Sacr. in Denkschr. Akad. Muench. vii. (1821) 238. t. 12 = **Clitoria**, Linn. (Legumin.).
brasiliensis, Zucc. ex Steud. Nom. ed. II. ii. 104 = C. glycinoides.
mexicana, Zucc. in Abh. Akad. Muench. i. (1832) 339. (= *Neurocarpum mexicanum*).—Mexic.
physodes, Leand. Sacr. in Denkschr. Akad. Muench. vii. (1821) 233. t. 12 = C. glycinoides.

MARTIA, Spreng. Anleit. ii. II. 788 (1818) = **Hypericum**, Tourn.
campanulata, Spreng. Syst. iii. 333 = H. virginicum.
dichotoma, Spreng. l. c. = H. dichotomum, *Lam.*
petiolata, Spreng. l. c. = H. paludosum.
polyandra, Spreng. l. c. = H. aegyptiacum.
virginicum, Spreng. Anleit. ii. II. 788 = H. virginicum.

MARTINERIA, Pfeiff. Nom. ii. 236 (1874) = MARTINIERIA, Vell. = **Kielmeyera**, Mart. (Ternstr.).

MARTINEZIA, Ruiz & Pav. Prod. 148. t. 32 (1794). *PALMAE*, Benth. & Hook. f. iii. 944.
AIPHANES, Willd. in Mém. Acad. Berl. (1804) 32.
MARARA, Karst. in Linnaea, xxviii. (1856) 389.
aculeata, Klotzsch, in Linnaea, xx. (1847) 455 = M. Aiphanes.
Aiphanes, *Mart. Palmet. Orbign.* 77.—Venezuela.
antioquensis, *Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
caryotaefolia, *H. B. & K. Nov. Gen. et Sp.* i. 305. t. 699.—Am. austr.
ciliata, Ruiz & Pav. Syst. Veg. 295 = Bactris ciliata.
corallina, *Mart. Hist. Nat. Palm.* iii. 284.—Ins. Martinique.
disticha, *Linden, Cat.* n. 93 (1875) 32, cum ic.; *Wallis, ex Regel, Gartenfl.* (1879) 163. t. 977. f. 1.—N. Granat.
elegans, *Linden & H. Wendl. in Linnaea*, xxviii. (1856) 351.—N. Granat.
ensiformis, Ruiz & Pav. Syst. Veg. 297 = Euterpe ensiformis.
erosa, *Linden, Cat.* n. 87 (1871) 5.—Ind. occ.
granatensis, *Hort. ex H. Wendl. in Kerch. Palm.* 251, nomen.—N. Granat.
interrupta, *Ruiz & Pav. Syst. Veg.* 296.—Peruv.
lanceolata, Ruiz & Pav. l. c. 297 = Chamaedorea gracilis.
leucophoeus, *Hort.—Cf. Gard. Chron.* (1875) I. 589.—N. Granat.
Lindeniana, *H. Wendl. in Linnaea*, xxviii. (1856) 349.—N. Granat.
linearis, Ruiz & Pav. Syst. Veg. 297 = Chamaedorea linearis.
minor, *Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
Roezlii, *Hort.—Cf. Gard. Chron.* (1876) I. 735; (1878) I. 439.—Am. trop.
truncata, *Brongn. in Mart. Palmet. Orbign.* 75.—Bolivia.

MARTINIERA, Guill. in Deless. Ic. Sel. iii. 23. t. 40 (1837) = **Wendtia**, Meyen (Geran.).
potentilloides, Guill. l. c. = W. gracilis.

MARTINIERIA, Vell. Fl. Flum. 232 (1825); v. t. 114. (1827) = **Kielmeyera**, Mart. (Ternstr.).
arborea, Vell. ll. cc. = K. coriacea.

MARTINIERIA, Walp. Ann. i. 141 (1848) = MARTINIERA, Guill. = **Wendtia**, Meyen (Geran.).

MARTINSIA, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii.] 1853 (1854) 378 = **Boreava**, Jaub. & Spach (Crucif.).
glastifolia, Godr. l. c. = B. orientalis.

MARTIUSA, Benth. & Hook. f. Gen. i. 571 (1865) = seq.

MARTIUSIA, Benth. in Hook. Journ. Bot. ii. (1840) 84 = **Martia**, Benth. (Legumin.).
excelsa, Benth. l. c. = Martia excelsa.
parvifolia, Benth. l. c. 103 = Martia parvifolia.

MARTIUSIA, Schult. Mant. i. 69 (1822) = **Clitoria**, Linn. (Legumin.).
physalodes, Schult. l. c. 226 = C. glycinoides.

MARTRASIA, Lag. Amen. Nat. i. 36 (1811) = **Jungia**, Linn. f. (Compos.).
auriculata, Lag. l. c. = J. auriculata.
axillaris, Lag. ex Steud. Nom. ed. II. ii. 104 = J. axillaris.
crenata, Lag. Amen. Nat. i. 36 = J. crenata.
pubescens, Lag. l. c. = J. floribunda, *Less.*
sericea, Lag. l. c. = J. sericea.

MARTYNIA, Houst. ex Linn. Syst. ed. I (1735). *PEDALINEAE*, Benth. & Hook. f. ii. 1055.
BASONCA, Rafin. Fl. Tellur. iv. 68 (1836).
CARPOCERAS, A. Rich. Elem. Bot. (1846) 706.
DISTERIA, Rafin. Fl. Tellur. iv. 68 (1836).
HOLOREGMIA, Nees, in Flora, iv. (1821) 300.
MYRTINIA, Nees, in DC. Prod. xi. 174 (1847).
PROBOSCIDEA, Schmid, Ic. 49. t. 12, 13 (1747).
SESAMUM, Adans. Fam. ii. 213 (1763).
VATKEA, Hildeb. & O. Hoffm. in Verh. Bot. Ver. Brandenb. xxii. (1880) 45, 78.
alternifolia, Lam. Encyc. ii. 112 = proboscidea.
altheaefolia, *Benth. Bot. Voy. Sulph.* 37.—Am. bor. occ.
angulosa, Lam. Encyc. ii. 112 = diandra.
annua, Linn. Sp. Pl. 618 = proboscidea, Craniolaria annua.
arenaria, Engelm. in Wisliz. Tour North Mexic. 100 = altheaefolia.
Botterii, *Hemsl. Biol. Centr. Am. Bot.* ii. 499.—Mexic.
capensis, Glox. Obs. 13 = Rogeria longiflora.
caulescens, Desf. ex Steud. Nom. ed. II. ii. 104 = Sinningia speciosa.
Craniolaria, Glox. Obs. 14 = Craniolaria annua.
diandra, *Glox. l. c.* t. 1.—Bras.; Mexic.
fallax, Kunze, in Linnaea, xviii. (1844) 164 = Craniolaria fallax.
fragrans, *Lindl. Bot. Reg.* (1840) *Misc.* 85.—Mexic.
fruticosa, Glox. Obs. 15 = Pentarhaphia Craniolaria.
hirsuta, Desf. ex Steud. Nom. ed. II. ii. 104 (= *Ligeria hirsuta*).—Bras.
integrifolia, Steud. l. c. = Craniolaria integrifolia.
lanceolata, Macrae, ex Nees, in DC. Prod. xi. 174, 182 = Strobilanthes lanceolatus.
lanceolata, Moon, Cat. Pl. Ceyl. 45 = Chirita Moonii, Strobilanthes lanceolatus.
longiflora, Linn. Syst. ed. XII. 412 = Rogeria longiflora.
louisiana, Mill. Gard. Dict. ed. VIII. n. 3 = seq.
louisianica, Mill. l. c. (in Err.) = proboscidea.
lutea, *Lindl. in Bot. Reg.* t. 934.—Bras.
montevidensis, *Cham. in Linnaea*, vii. (1832) 724.—Bras.
pallida, Salisb. Prod. 103 = proboscidia.
perennis, Linn. Sp. Pl. 618 = Gloxinia maculata.
proboscidea, *Glox. Obs.* 14.—Am. bor.
spathacea, Lam. Encyc. ii. 112 = Craniolaria annua.
spathacea, *Spreng. Syst.* iv. *Cur. Post.* 238.—Bras.
speciosa, Loisel. Herb. Amat. t. 225 = Sinningia speciosa.
triloba, *Cham. & Schlecht. in Linnaea*, v. (1830) 121.—Mexic.
violacea, Engelm. in Wisliz. Tour North Mexic. 101 = fragrans.
zanguebarica, Lour. Fl. Cochinch. ii. 386 = Pretrea zanguebarica.

MARUBIUM, Roth, Tent. Fl. Germ. ii. II. 34 (1793) = **Marrubium**, Tourn. (Labiat.).

MARUMIA, Blume, in Flora, xiv. (1831) 503. *MELASTOMACEAE*, Benth. & Hook. f. i. 757.
MACROLENES, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 311.
affinis, Korth. Verh. Nat. Gesch. Bot. 241. t. 60 = nemorosa.
annulata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 82.—Java.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 355.—Ins. Banca.
bipulvinata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 82.—Java.
echinulata, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 280 = zeylanica.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. I. 536.—Java.
Jackii, Korth. Verh. Nat. Gesch. Bot. 243 = stellulata.
Korthalsiana, *Miq. Fl. Ind. Bat. Suppl.* 318.—Sumatra.
leprosa, *Korth. Verh. Nat. Gesch. Bot.* 241.—Borneo.
muscosa, *Blume, in Flora*, xiv. (1831) 504.—Java.
nemorosa, *Blume, l. c.* 505.—Malaya.
oligantha, Naud. in Ann. Sc. Nat. Sér. xv. (1851) 281 = reticulata.

MARUMIA:—
pachygyna, *Korth. Verh. Nat. Gesch. Bot.* 242.—Borneo.
reticulata, *Blume, Mus. Bot. Lugd. Bat.* i. 34.—Sumatra.
rhodocarpa, Triana, in Trans. Linn. Soc. xxviii. (1871) 176 = zeylanica.
stellulata, *Blume, in Flora*, xiv. (1831) 505.—Sumatra.
stellulata, Korth. Verh. Nat. Gesch. Bot. 243 = reticulata.
vulcanica, Korth. l. c. = stellulata.
zeylanica, *Blume, in Flora*, xiv. (1831) 505.—Zeylan.

MARUMIA, Reinw. ex Blume, Cat. Gew. Buitenz. 79 (1828) = **Saurauja**, Willd. (Ternstr.).
angustifolia, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 50 = S. lanceolata.
cauliflora, Reinw. ex Blume, Cat. Gew. Buitenz. 79 = S. cauliflora.
dioica, Reinw. ex Miq. Fl. Ind. Bat. i. II. 484 = S. tristyla.
macrophylla, Reinw. ex Blume, Cat. Gew. Buitenz. 79. —Hab.?
montana, Reinw. ex Blume, l. c. 80 = S. hirsuta.
muricata, Reinw. ex Blume, l. c. = S. muricata.
pentapetala, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 48 = S. euryolepis.

MARUPA, Miers, in Journ. Bot. xi. (1873) 258. *SIMARUBEAE*.
ODINA, Netto, in Ann. Sc. Nat. Sér. V. v. (1866) 85.
Francoana, *Miers, in Journ. Bot.* xi. (1873) 260.—Bras.
paraënsis, *Miers, in Journ. Linn. Soc.* xvii. (1878) 151.—Bras.

MARUTA, Cass. in Bull. Soc. Philom. (1818) 167 = **Anthemis**, Mich. (Compos.).
Cotula, DC. Prod. vi. 13 = A. Cotula.
foetida, S. F. Gray, Nat. Arr. Brit. Pl. ii. 456 = A. Cotula.
fuscata, DC. Prod. vi. 14 = A. fuscata.
hemisphaerica, Hook. f. Fl. Brit. Ind. iii. 316, sphalm. = Cotula hemisphaerica.
lithuanica, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 53 = Matricaria confusa.
Micheliana, Fisch. Mey. & Avé-Lall. l. c. xi. 76 = A. maritima.
microcephala, Schrenk, in Fisch. Mey. & Avé-Lall. l. c. x. 53; et xi. 77.—Soongar.
mixta, Moris, Fl. Sard. ii. 416 = Ormenis mixta.
vulgaris, Bluff & Fingerh. Comp. Fl. Germ. ii. 392 = A. Cotula.

MASCAGNIA, Bert. ex Colla, Hort. Ripul. 86 (1824) = **Hiraea**, Jacq. (Malpig.).
ambigua, Griseb. in Mart. Fl. Bras. xii. I. 90 = H. ambigua.
americana, Bert. ex DC. Prod. i. 585 = H. oblongifolia.
anisopetala, Griseb. in Mart. Fl. Bras. xii. I. 95 = H. anisopetala.
argentea, Griseb. l. c. 90 = H. argentea.
biglandulosa, Griseb. l. c. 91 = Heladena biglandulosa.
bracteosa, Griseb. l. c. 97.—Bras.
brevifolia, Griseb. in Goett. Abh. xxiv. (1879) 67.—Reg. Argent.
bunchosioides, Griseb. in Mart. Fl. Bras. xii. I. 92 = Heladena bunchosioides.
chlorocarpa, Griseb. l. c. 93 = H. chlorocarpa.
cordifolia, Griseb. l. c. 95 = H. cordifolia.
coriacea, Griseb. l. c. 92.—Bras.
cynanchifolia, Griseb. l. c. 95.—Bras.
Doniana, Griseb. l. c. 94.—Bras.
dumetorum, Griseb. in Goett. Abh. xxiv. (1879) 67.—Reg. Argent.
elegans, Griseb. in Mart. Fl. Bras. xii. I. 95 = H. elegans.
exotropica, Griseb. l. c. 93.—Bras.
fluminensis, Griseb. l. c. 94 = H. fluminensis.
leucanthele, Griseb. l. c. 96.—Bras.
microphylla, Griseb. l. c. 98 = H. microphylla.
oblongifolia, Bert. ex DC. Prod. i. 585 = H. oblongifolia.
ovatifolia, Griseb. Fl. Brit. W. Ind. 121 = H. ovatifolia.
psilophylla, Griseb. in Mart. Fl. Bras. xii. I. 94 = H. psilophylla.
pubiflora, Griseb. l. c. 91 = H. pubiflora.

MASCAGNIA:—
rigida, Griseb. in Mart. Fl. Bras. xii. I. 92 = Hiraea rigida.
rubra, Griseb. l. c. 90.—Bras.
salicifolia, Griseb. l. c. = H. salicifolia.
sepium, Griseb. l. c. 96 = H. sepium.
septentrionalis, Griseb. l. c. 93, in obs. = H. septentrionalis.
Simsiana, Griseb. Fl. Brit. W. Ind. 121 = H. Simsiana.
sinemariensis, Griseb. in Mart. Fl. Bras. xii. I. 93, in obs. = H. Simsiana.

MASCARENHASIA, A. DC. in DC. Prod. viii. 487 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 721.
angustifolia, *A. DC. in DC. Prod.* viii. 488.—Ins. Nossibé.
arborescens, *A. DC. l. c.*—Madag.
brevituba, *Vatke, in Bremen, Abh.* ix. (1885) 125.—Ins. Nossibé.
Curnowiana, *Hort.—Cf. Gard. Chron.* (1884) II. 146.—Madag.
Havetii, *A. DC. in DC. Prod.* viii. 488.—Madag.
lanceolata, *A. DC. l. c.*—Ins. Nossibé.
lisianthiflora, *A. DC. l. c.* 487.—Madag.
macrocalyx, *Baker, in Journ. Bot.* xx. (1882) 219.—Madag.
Rutenbergiana, *Vatke, in Bremen, Abh.* ix (1885) 124.—Ins. Mascar.

MASCHALANTHE, Blume, Fl. Jav. Praef. p. vi (1828) = **Urophyllum**, Wall. (Rubiac.).

MASCHALANTHUS, Nutt. in Trans. Am. Phil. Soc. v. (1837) 175 = **Andrachne**, Linn. (Euphorb.).
obovatus, Nutt. l. c. = A. Roemeriana.
polygonoides, Nutt. l. c. = A. Roemeriana.

MASDEVALLIA, Ruiz & Pav. Prod. 122. t. 27 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 492.
abbreviata, *Reichb. f. in Gard. Chron.* (1878) II. 106.—Peruv.?
acrochordonia, *Reichb. f. Xenia Orch.* ii. 213.—Ecuador.
aequiloba, *Regel, Gartenfl.* ix. (1860) 82. t. 285.—Peruv.
affinis, *Lindl. Orch. Linden.* 5.—N. Granat.
amabilis, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 116.—Peruv.
amanda, *Reichb. f. & Warsc. l. c.* 115.—N. Granat.
amethystina, *Reichb. f. Otia Bot. Hamb.* 14.—Reg. Andin.
anachaeta, *Reichb. f. l. c.* 17.—Ecuador.
anchorifera, *Reichb. f. in Gard. Chron.* (1884) I. 577.—Costa Rica.
angulata, *Reichb. f. Otia Bot. Hamb.* 15.—Ecuador.
aristata, *Rodrig. Orch. Nov.* i. 30.—Bras.
Arminii, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 283.—N. Granat.
atropurpurea, Reichb. f. in Walp. Ann. vi. 1216 (Index) = auropurpurea.
attenuata, *Reichb. f. in Gard. Chron.* (1871) 834.—Costa Rica.
aurantiaca, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 357.—Bras.
auriculigera, *Reichb. f. Otia Bot. Hamb.* 93.—Bras.
auropurpurea, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 115.—N. Granat.
aviceps, *Reichb. f. Otia Bot. Hamb.* 93.—Bras.
Backhousiana, *Reichb. f. in Gard. Chron.* (1879) I. 716.—N. Granat.
Barlaeana, *Reichb. f. in Gard. Chron.* (1876) I. 170.—Peruv.
bella, *Reichb. f. l. c.* (1878) I. 725.—N. Granat.
Benedicti, *Reichb. f. Xenia Orch.* ii. 197.—N. Granat.
bicolor, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 6.—Peruv.
biflora, E. Morr. in Fl. des Serres, Sér. II. ix. (1873) 128 = bicolor.
Bonplandi, *Reichb. f. in Bonplandia*, iii. (1855) 69.—Hab.?
brevis, *Reichb. f. in Gard. Chron.* (1883) II. 588.—Guian.
Bruchmulleri, *E. Morr. in Fl. des Serres*, Sér. II. ix. (1873) 128.—Hab.?
Buccinator, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 115.—N. Granat.

MASDEVALLIA:—
caloptera, *Reichb. f. in Gard. Chron.* (1874) I. 338.—Peruv.
calura, *Reichb. f. l. c.* (1883) II. 230.—Hab.?
campyloglossa, *Reichb. f. l. c.* (1878) II. 588.—Hab.?
candida, Klotzsch & Karst. ex Reichb. f. in Bonplandia, ii. (1854) 23 = tovarensis.
Carderi, *Reichb. f. in Gard. Chron.* (1883) I. 784.—Hab.?
caudata, *Lindl. Gen. et Sp. Orch.* 193.—N. Granat.
cayennensis, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 242.—Guian.
Chelsoni ×, Reichb. f. in Gard. Chron. (1880) I. 554.
Chestertoni, *Reichb. f. in Gard. Chron.* (1883) I. 532.—N. Granat.
Chimaera, André, in Illustr. Hortic. xx. (1873) 25. t. 117–118 = nycterina.
Chimaera, *Reichb. f. in Gard. Chron.* (1872) 463.—Am. austr.
chontalensis, *Reichb. f. Otia Bot. Hamb.* 17.—Nicaragua.
cinnamomea, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Peruv.
civilis, *Reichb. f. l. c.* ii. (1854) 115.—Peruv.
coccinea, *Linden, ex Lindl. Orch. Linden.* 5.—N. Granat.
Colibri, Burbidge, in Florist & Pomol. (1873) 3 = M. Ephippium.
constricta, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 6.—Peruv.
coriacea, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 257.—N. Granat.
corniculata, *Reichb. f. in Gard. Chron.* (1878) I. 72.—N. Granat.
cucullata, *Lindl. Orch. Linden.* 4.—N. Granat.
cuprea, *Lindl. Bot. Reg.* (1843) *Misc.* 81.—Guian.
cupularis, *Reichb. f. Beitr. Orch. Centr. Am.* 93.—Costa Rica.
curtipes, *Rodrig. Orch. Nov.* ii. 95.—Bras.
Davisii, *Reichb. f. in Gard. Chron.* (1874) II. 710.—Am. austr.
Dayana, *Reichb. f. in Gard. Chron.* (1880) II. 295.—N. Granat.
Denisoni, *D'Ombr. in Fl. Mag.* N. S. t. 79.—Hab.?
Echidna, *Reichb. f. in Bonplandia*, iii. (1855) 69.—N. Granat.
Eduardi, *Reichb. f. in Gard. Chron.* (1880) II. 778.—N. Granat.
elephanticeps, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 116.—N. Granat.
ellipes, *Reichb. f. in Linnaea*, xli. (1877) 11.—N. Granat.
ensata, *Reichb. f. l. c.* xxii. (1849) 818.—N. Granat.
Ephippium, *Reichb. f. in Bot. Zeit.* (1873) 390.—N. Granat.
erinacea, *Reichb. f. in Linnaea*, xli. (1877) 11.—N. Granat.
erythrochaete, *Reichb. f. in Gard. Chron.* (1882) II. 392.—Am. centr.
Estradae, *Reichb. f. l. c.* (1874) II. 435.—N. Granat.
expansa, *Reichb. f. in Otia Bot. Hamb.* 16.—N. Granat.
Falcago, *Reichb. f. l. c.*—N. Granat.
fasciata, *Reichb. f. in Gard. Chron.* (1881) I. 202.—N. Granat.
fenestrata, *Linden, Illustr. Hortic.* xxix. (1882) 67, nomen.—N. Granat.
fenestrata, *Lindl. ex Hook. Bot. Mag.* t. 4164.—Jamaic.
flaveola, *Reichb. f. in Gard. Chron.* (1884) I. 638.—Costa Rica.
floribunda, *Lindl. Bot. Reg.* (1843) *Misc.* 72.—Mexic.
Fraseri ×, Reichb. f. in Gard. Chron. (1882) I. 143.
Gairiana ×, Reichb. f. l. c. (1884) II. 38.
Galeottiana, Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 17 = floribunda.
Gargantua, *Reichb. f. in Gard. Chron.* (1876) II. 516.—N. Granat.
Gaskelliana, *Reichb. f. l. c.* (1883) II. 294.—Hab.?
gemmata, *Reichb. f. l. c.*—Hab.?
gibberosa, *Reichb. f. l. c.* (1876) I. 8.—N. Granat.
gracilenta, *Reichb. f. l. c.* (1875) 98.—Costa Rica.
guayanensis, *Lindl. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 673.—Guian.
Gustavi, *Reichb. f. in Gard. Chron.* (1875) I. 461.—Hab.?
guttulata, *Reichb. f. in Linnaea*, xli. (1877) 118.—Ecuador?

MASDEVALLIA:—
Harryana, Reichb. f. in Gard. Chron. (1871) 1421 = coccinea.
heteroptera, *Reichb. f. l. c.* (1875) I. 590.—N. Granat.
hians, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 283.—N. Granat.
hieroglyphica, *Reichb. f. in Gard. Chron.* (1882) II. 230.—N. Granat.
Houtteana, *Reichb. f. l. c.* (1874) II. 98.—N. Granat.
hymenantha, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Peruv.
Hypodiscus, *Reichb. f. in Gard. Chron.* (1878) II. 234.—Hab.?
ignea, *Reichb. f. l. c.* (1871) 1482.—N. Granat.
inequalis, *Reichb. f. l. c.* (1874) I. 372.—N. Granat.
inflata, *Reichb. f. l. c.* (1881) II. 716.—Hab.?
infracta, *Lindl. Gen. et Sp. Orch.* 193.—Bras.
ionocharis, *Reichb. f. in Gard. Chron.* (1875) II. 388.—Peruv.
Klabocthorum, *Reichb. f. l. c.* (1876) I. 720.—Am. austr.
laevis, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 257.—N. Granat.
Lansbergii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 317.—Venezuela.
lata, *Reichb. f. in Gard. Chron.* (1877) I. 653.—Am. centr.
Lehmanni, *Reichb. f. l. c.* (1877) II. 38.—Ecuador.
leontoglossa, *Reichb. f. in Bonplandia*, iii. (1855) 69.—N. Granat.
lepida, *Reichb. f. l. c.*—N. Granat.
Lindeni, André, Illustr. Hortic. xvii. (1870) 226. t. 42 = coccinea.
Lindeniana, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 17.—Mexic.
Livingstoneana, *Roezl & Reichb. f. in Gard. Chron.* (1874) II. 322.—Panama.
longicaudata, Lem. Illustr. Hortic. xv. (1868) Misc. 109 = infracta.
ludibunda, *Reichb. f. in Gard. Chron.* (1882) I. 179.—N. Granat.
macrodactyla, *Reichb. f. l. c.* (1872) 571.—N. Granat.
macroglossa, *Reichb. f. in Otia Bot. Hamb.* 15.—Venezuela.
macrura, *Reichb. f. in Gard. Chron.* (1874) I. 240.—N. Granat.
maculata, *Klotzsch & Karst. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 330.—Venezuela.
marginella, *Reichb. f. in Gard. Chron.* (1883) II. 38.—Hab.?
Mastodon, *Reichb. f. in Bonplandia*, iii. (1855) 69.—N. Granat.
melanopus, *Reichb. f. in Gard. Chron.* (1874) I. 338.—Peruv.
melanoxantha, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 283.—N. Granat.
Meleagris, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 257.—N. Granat.
militaris, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 115.—N. Granat.
minuta, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 396.—Guiana.
Molossus, *Reichb. f. in Linnaea*, xli. (1877) 10.—N. Granat.
Mooreana, *Reichb. f. in Gard. Chron.* (1884) I. 408.—Hab.?
muscosa, *Reichb. f. l. c.* (1875) I. 460.—N. Granat.
myriostigma, E. Morr. in Fl. des Serres, Sér. II. ix. (1869–70) 131, 132 = floribunda.
nidifica, *Reichb. f. Otia Bot. Hamb.* 18.—Ecuador.
Normanni, Hort.—Cf. Gard. Chron. (1881) II. 230 = Reichenbachiana.
nycterina, *Reichb. f. l. c.* (1873) 1238.—N. Granat.
ochracea, Burbidge, l. c. (1871) 1421 = coriacea.
ochthodes, *Reichb. f. in Bonplandia*, iii. (1855) 70.—N. Granat.
ophioglossa, *Reichb. f. Otia Bot. Hamb.* 17.—Ecuador.
pachyantha, *Reichb. f. in Gard. Chron.* (1884) I. 174.—N. Granat.
pachyura, *Reichb. f. l. c.* (1874) II. 322.—Hab.?
Paivaeana, *Reichb. f. Xenia Orch.* iii. 25.—Bolivia.
pardina, *Reichb. f. Otia Bot. Hamb.* 15.—N. Granat.
Parlatoreana, *Reichb. f. in Gard. Chron.* (1879) I. 172.—Peruv.
Peristeria, *Reichb. f. l. c.* (1874) I. 500.—N. Granat.
picturata, *Reichb. f. Otia Bot. Hamb.* 16.—Venezuela.

MASDEVALLIA:—
platyglossa, *Reichb. f. in Gard. Chron.* (1882) II. 552.—Hab.?
polyantha, *Lindl. Orch. Linden.* 6.—N. Granat.
polysticta, Hook. f. Bot. Mag. t. 6258 = melanopus.
polysticta, *Reichb. f. in Gard. Chron.* (1874) I. 338.—Peruv.
porcelliceps, *Reichb. f. l. c.* (1883) I. 10.—Hab.?
psittacina, *Reichb. f. l. c.* (1876) I. 817.—N. Granat.?
pulvinaris, *Reichb. f. l. c.* (1880) I. 200.—Hab.?
pumila, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 6.—Peruv.
racemosa, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 256.—N. Granat.
radiosa, *Reichb. f. in Gard. Chron.* (1877) I. 684.—N. Granat.
Reichenbachiana, *Endres, l. c.* (1875) II. 257.—Costa Rica.
riograndensis, *Hort.—Cf. Gard. Chron.* (1871) 1421, nomen.—Hab.?
Roezlii, *Reichb. f. Xenia Orch.* ii. 196.—N. Granat.
rosea, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 257.—N. Granat.
rufolutea, Lindl. in Gard. Chron. (1853) 292, nomen = civilis.
Saltatrix, *Reichb. f. in Linnaea*, xli. (1877) 10.—N. Granat.
Sceptrum, *Reichb. f. in Bonplandia*, ii. (1854) 283.—N. Granat.
Schlimii, *Linden, ex Lindl. Orch. Linden.* 5.—Venezuela.
senilis, *Reichb. f. in Gard. Chron.* (1885) II. 489.—Hab.?
severa, *Reichb. f. l. c.* (1875) I. 170.—N. Granat.
Shuttleworthii, *Reichb. f. l. c.*—N. Granat.
simula, *Reichb. f. l. c.* 8.—N. Granat.
Spectrum, *Reichb. f. l. c.* 429.—Hab.?
splendida, *Reichb. f. l. c.* (1878) I. 493.—Am. austr.
Sprucei, *Reichb. f. Otia Bot. Hamb.* 17.—Bras.
surinamensis, Focke, in Tijdschr. Nat. Wetensch. iv. (1851) 62 = minuta.
swertiaefolia, *Reichb. f. in Gard. Chron.* (1880) II. 390.—N. Granat.
torta, *Reichb. f. l. c.* (1883) I. 110.—Hab.?
tovarensis, *Reichb. f. in Linnaea*, xxii. (1849) 818.—N. Granat.
triangularis, *Lindl. Orch. Linden.* 5.—N. Granat.
triaristella, *Reichb. f. in Gard. Chron.* (1876) II. 226.—Costa Rica.
trichaete, *Reichb. f. l. c.* (1883) II. 360.—Hab.?
tricolor, *Reichb. f. l. c.* (1882) II. 102.—N. Granat.
tricolor, *Reichb. f. in Linnaea*, xxii. (1849) 818.—Venezuela.
tridactylites, *Reichb. f. in Gard. Chron.* (1883) I. 784.—Hab.?
tridens, *Reichb. f. Otia Bot. Hamb.* 13.—Ecuador.
tridentata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 357.—Bras.
Triglochin, *Reichb. f. in Gard. Chron.* (1877) II. 648.—Ecuador.
triquetra, *Scheidw. in Otto & Dietr. Allg. Gartenz.* vii. (1839) 146.—Hab.?
Trochilus, *Linden & André, ex Gard. Chron.* (1873) 711; *Illustr. Hortic.* xxi. (1874) 136. t. 180.—N. Granat.
Troglodytes, *E. Morr. Belg. Hortic.* xxvii. (1877) 97. t. 5.—N. Granat.
Tubeana, *Reichb. f. in Gard. Chron.* (1878) I. 234.—Ecuador.
tubulosa, *Lindl. Orch. Linden.* 4.—Venezuela.
uncifera, *Reichb. f. Otia Bot. Hamb.* 16.—Ecuador.
uniflora, *Ruiz & Pav. Syst. Veg.* 288.—Ecuador.
urostachya, *Reichb. f. in Gard. Chron.* (1882) I. 765.—Hab.?
Veitchiana, *Reichb. f. in Gard. Chron.* (1868) 814; *et ex Hook. f. Bot. Mag.* t. 5739.—Peruv.
velifera, *Reichb. f. in Gard. Chron.* (1874) I. 406, nomen; (1874) II. 98.—Hab.?
velutina, *Reichb. f. l. c.* (1875) II. 420.—N. Granat.
ventricularia, *Reichb. f. Otia Bot. Hamb.* 14.—Ecuador.
venusta, *Hort.—Cf. Gard. Chron.* (1871) 1421, nomen.—Hab.?
verrucosa, *Reichb. f. in Linnaea*, xxii. (1849) 819.—Venezuela.
Vespertilio, *Reichb. f. in Bot. Zeit.* xxxi. (1873) 390.—N. Granat.

MASDEVALLIA:—
Wageneriana, *Linden, in Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 74.—Venezuela.
Wallisii, *Reichb. f. in Gard. Chron.* (1875) II. 258.—N. Granat.
Winniana, *Reichb. f. l. c.* (1881) II. 198.—Hab.?
xanthina, *Reichb. f. l. c.* (1880) I. 681.—Hab.?
xanthodactyla, *Reichb. f. l. c.* (1877) II. 552.—Hab.?
xipheres, *Reichb. f. in Linnaea*, xli. (1877) 12.—N. Granat.
xylina, *Reichb. f. in Bot. Zeit.* xxxi. (1873) 390.—N. Granat.

MASSANGEA, E. Morr. in Belg. Hortic. (1877) 59, 199. t. 8 = **Caraguata**, Lindl. (Bromel.).
hieroglyphica, Carr. in Rev. Hortic. (1878) 175.—Bras.
Lindeni, André, in Illustr. Hortic. xxv. (1878) 55. t. 309.—Peruv.
musaica, E. Morr. Belg. Hortic. xxvii. (1877) 59, 199. tt. 8, 9 = C. musaica.
tigrina, Hort.—Cf. Gard. Chron. (1883) I. 510.—Bras.

MASSANGIA, Benth. & Hook. f. Gen. iii. 668 (1883), err. typ. = praec.

MASSOIA, Becc. in D'Albertis, New Guinea, ii. 398 (1880). *LAURINEAE.*
aromatica, *Becc. l. c.*—N. Guin.

MASSONIA, Thunb. ex Linn. f. Suppl. 27, 193 (1781). *LILIACEAE*, Benth. & Hook. f. iii. 806. PODOCALLIS, Salisb. Gen. Pl. Fragm. 17 (1866).
angustifolia, [Delile, in] Red. Lil. t. 392 = Zeyheri.
angustifolia, *Linn. f. Suppl.* 193.—Afr. austr.
aurea, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 699 = Daubenya aurea.
Bowkeri, *Baker, in Journ. Linn. Soc.* xi. (1871) 390.—Afr. austr.
brachypus, *Baker, in Journ. Bot.* xii. (1874) 368.—Afr. austr.
breviscapa, *Lehm. Delect. Sem. Hort. Hamb.* (1838) 8.—Afr. austr.
Burchellii, *Baker, in Journ. Linn. Soc.* xi. (1871) 393.—Afr. austr.
calvata, *Baker, in Journ. Bot.* xvi. (1878) 321.—Afr. austr.
candida, *Burch. ex Kunth, Enum. Pl.* iv. 297, in syn.—Afr. austr.
comata, *Burch. ex Baker, in Journ. Linn. Soc.* xi. (1871) 392.—Afr. austr.
cordata, *Jacq. Hort. Schoenb.* iv. 30. t. 459.—Afr. austr.
coronata, Jacq. l. c. t. 460 = sanguinea.
corymbosa, Ker-Gawl. in Bot. Mag. t. 991 = Hyacinthus corymbosus.
depressa, Houtt. Handleid. xii. 424 = latifolia.
echinata, *Linn. f. Suppl.* 193.—Afr. austr.
ensifolia, *Ker-Gawl. in Bot. Mag.* t. 554.—Afr. austr.
grandiflora, Lindl. in Bot. Reg. t. 958 = obovata.
grandifolia, Ker-Gawl. in Bot. Mag. sub t. 991 = latifolia.
hirsuta, *Link & Otto, Ic. Pl. Rar.* i. t. 1.—Afr. austr.
Huttoni, *Baker, in Journ. Linn. Soc.* xi. (1871) 390.—Afr. austr.
jasminiflora, *Burch. ex Baker, l. c.*—Afr. austr.
lanceaefolia, *Jacq. Hort. Schoenb.* iv. 29. t. 456.—Afr. austr.
lanceolata, Thunb. Nov. Gen. 40 = angustifolia.
lanceolata, Zeyh. ex Kunth, Enum. Pl. iv. 298 = Zeyheri.
latifolia, Ker-Gawl. in Bot. Mag. t. 848 = sanguinea.
latifolia, *Linn. f. Suppl.* 193.—Afr. austr.
linearis, Drège, ex Kunth, Enum. Pl. iv. 293 = Hyacinthus corymbosus.
longifolia, *Jacq. Hort. Schoenb.* iv. 29. t. 457.—Afr. austr.
lutea, Hort. ex Lindl. Bot Reg. t. 1813 = Daubenya aurea.
marginata, *Willd. ex Kunth, Enum. Pl.* iv. 299.—Afr. austr.
muricata, *Ker-Gawl. in Bot. Mag.* t. 559.—Afr. austr.
nervosa, *Hornem. Hort. Hafn. Suppl.* 39.—Afr. austr.
nodicarpa, *Gaertn. f. Fruct.* iii. 13.—Afr. austr.

MASSONIA:—
obovata, *Jacq. Hort. Schoenb.* iv. 29. t. 458.—Afr. austr.
odorata, *Hook. f. Bot. Mag.* t. 5891.—Afr. austr.
orientalis, *Baker, in Journ. Bot.* xvi. (1878) 321.—Afr. austr.
ovalis, Drège, ex Kunth, Enum. Pl. iv. 295 = ensifolia.
pauciflora, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 210.—Afr. austr.
Pinsilla, Mass. ex Salisb. Gen. Pl. Fragm. 17 = hirsuta.
pusilla, Baker, in Journ. Linn. Soc. xi. (1870) 389, sphalm. = hirsuta.
pusilla, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 991 = undulata.
pustulata, *Jacq. Coll.* iv. 177.—Afr. austr.
pygmaea, *Schlecht. ex Kunth, Enum. Pl.* iv. 298.—Afr. austr.
rugulosa, Licht. ex Kunth, l. c. 299 = marginata.
sanguinea, *Jacq. Hort. Schoenb.* iv. 31.—Afr. austr.
scabra, Thunb. Prod. Pl. Cap. 188 = pustulata.
Schlechtendalii, *Baker, in Journ. Bot.* xii. (1874) 5.—Afr. austr.
setulosa, *Baker, in Journ. Linn. Soc.* xi. (1871) 389.—Afr. austr.
tenella, *Soland. ex Baker, l. c.*—Afr. austr.
undulata, Thunb. Nov. Gen. 41 = Lachenalia pusilla.
uniflora, *Banks, ex Ker-Gawl. in Bot. Mag.* sub t. 991, nomen; *et ex Baker, in Journ. Linn. Soc.* xi. (1870) 393.—Afr. austr.
versicolor, *Baker, in Journ. Bot.* xiv. (1876) 186.—Afr. austr.
violacea, Andr. Bot. Rep. t. 46 = ensifolia.
Zeyheri, *Kunth, Enum. Pl.* iv. 298.—Afr. austr.

MASSOUNIA, Thunb. Fl. Cap. ii. 275 (1818) = **Massonia**, Thunb. (Liliac.).

MASSOVIA, Benth. & Hook. f. Gen. iii. 998 (1883) = seq.

MASSOWIA, C. Koch, in Bot. Zeit. x. (1852) 277 = **Spathiphyllum**, Schott (Aroid.).
cannaefolia, C. Koch, l. c. 278 = S. cannaefolium.
cannaeformis, C. Koch, in Bonplandia, iv. (1856) 11 = S. Humboldtii.
commutata, Ender, Ind. Aroid. 52 = S. commutatum.
Friedrichsthalii, C. Koch, in Bonplandia, iv. (1856) 11 = S. Friedrichsthalii.
Gardneri, C. Koch, l. c. = S. Gardneri.
heliconiaefolia, C. Koch, l. c. = S. heliconiaefolium.
lanceaefolia, C. Koch, l. c. = S. cochlearispathum.
lanceolata, C. Koch, l. c. = S. Friedrichsthalii.
Lechleriana, C. Koch, ex Ender, Ind. Aroid. 53 = S. Lechlerianum.
marginata, C. Koch, l. c., nomen.—N. Granat.

MASSULA, Dulac, Fl. Hautes-Pyr. 47 (1867) = **Typha**, Tourn.
angustifolia, Dulac, l. c. = T. angustifolia.
latifolia, Dulac, l. c. = T. latifolia.

MASTACANTHUS, Endl. Gen. 638 (1838) = **Caryopteris**, Bunge (Verben.).
Barbula, Steud. Nom. ed. II. ii. 105 = C. Mastacanthus.
sinensis, Endl. ex Walp. Rep. iv. 3 = C. Mastacanthus.

MASTERSIA, Benth. in Trans. Linn. Soc. xxv. (1865) 300. t. 34. *LEGUMINOSAE*, Benth. & Hook. f. i. 535.
assamica, *Benth. in Trans. Linn. Soc.* xxv. (1865) 300.—Reg. Himal.
cleistocarpa, Baker, in Hook. f. Fl. Brit. Ind. ii. 195 = assamica.

MASTICHINA, Adans. Fam. ii. 194 (1763) = **Thymus**, Tourn. (Labiat.).

MASTICHODENDRON, R. Hedw. Gen. 116 (1806) = **Sideroxylon**, Linn. (Sapot.).

MASTIGLOSCLERIA, Nees, in Mart. Fl. Bras. ii. 1. 177 (1842) = **Scleria**, Berg. (Cyperac.).
reflexa, Nees, l. c. = S. reflexa.

MASTIGOPHORUS, Cass. in Dict. Sc. Nat. xxxiv. 222 (1825) = **Nassauvia**, Comm. (Compos.).
Gaudichaudii, Cass. l. c. 223 = N. Gaudichaudii.

MASTIXIA, Blume, Bijdr. 654 (1825). *CORNACEAE*, Benth. & Hook. f. i. 950.
BURSINOPETALUM, Wight, Ic. t. 956 (1847).
MASTYXIA, Spach, Hist. Vég. Phan. viii. 88 (1839).

MASTIXIA:—
acuminatissima, *Blume, Mus. Bot. Lugd. Bat.* i. 258.—Sumatra.
arborea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 745.—Ind. or.
bracteata, *C. B. Clarke, l. c.* 746.—Malacca.
caesia, *Blume, Mus. Bot. Lugd. Bat.* i. 258.—Java.
cuneata, *Blume, l. c.* 257.—Java.
cuspidata, Blume, l. c. 256 = pentandra.
Gardneriana, Baill. Adansonia, iii. (1862-63) 83 = arborea.
heterophylla, *Blume, Mus. Bot. Lugd. Bat.* i. 257.—Sumatra.
Junghuhniana, *Miq. Fl. Ind. Bat.* i. 1. 772.—Malacca; Java.
Kimanilla, *Blume, Mus. Bot. Lugd. Bat.* i. 258.—Java.
lanceolata, Baill. Adansonia, iii. (1862-63) 83 = tetrandra.
laxa, *Blume, Mus. Bot. Lugd. Bat.* i. 257.—Java.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 746.—Singapore.
pentandra, *Blume, Bijdr.* 654.—Ind. or.; Java.
rostrata, *Blume, Mus. Bot. Lugd. Bat.* i. 258.—Java.
tetrandra, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 745.—Zeylan.
Thwaitesii, Baill. Adansonia, iii. (1862-63) 83 = arborea.
trichotoma, *Blume, Bijdr.* 655.—Java.
trichotoma, Blume, Mus. Bot. Lugd. Bat. i. 258 = laxa.

MASTOSTIGMA, Stocks, in Hook. Ic. Pl. t. 863 (1852) = **Glossonema**, Decne. (Apocyn.).
varians, Stocks, in Hook. l. c. = G. varians.

MASTOSUKE, Rafin. Sylva Tellur. 59 (1838) = **Ficus**, Tourn. (Urtic.).
rubiginosa, Rafin. l. c. = F. rubiginosa.

MASTRUCIUM, Cass. in Dict. Sc. Nat. xxxv. 173 (1825) = **Serratula**, Linn. (Compos.).
coronata, Cass. l. c. = S. coronata.
pinnatifidum, DC. Prod. vi. 667 = S. coronata.

MASTRUTIUM, Endl. Ench. 247 (1841) = praec.

MASTYXIA, Spach, Hist. Vég. Phan. viii. 88 (1839) = **Mastixia**, Blume (Cornac.).

MASUS, G. Don, Gen. Syst. iv. 542 (1837) = **Mazus**, Lour. (Scrophul.).

MATAIBA, R. Hedw. Gen. 264 (1806) = MATAYBA, Aubl. = **Ratonia**, DC. (Sapind.).

MATALEA, A. Gray, Syn. Fl. N. Am. ii. 1. 96, in syn. (1878), sphalm. = **Matelea**, Aubl. (Asclep.).

MATAMORIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 8 (1824) = **Elephantopus**, Linn. (Compos.).
spicata, La Llave, l. c. = E. spicatus.

MATAXA, Spreng. Syst. iv. Cur. Post. 297, 303 (1827) = **Lasiospermum**, Lag. (Compos.).
capensis, Spreng. l. c. 303 = L. radiatum.

MATAYBA, Aubl. Pl. Gui. i. 331. t. 128 (1775), male desc. = **Ratonia**, DC. (1824) (Sapind.).
apetala, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 624 = R. apetala.
arborescens, Radlk. l. c. 625.—Guian.
camptoneura, Radlk. l. c. 629.—Guian.
discolor, Radlk. l. c. 630.—Bras.
domingensis, Radlk. l. c. 624 = R. domingensis.
elaeagnoides, Radlk. l. c. 635.—Reg. Argent.
elegans, Radlk. l. c. 629.—N. Granat.
floribunda, Radlk. l. c. 626.—Mexic.
glaberrima, Radlk. l. c. 628.—Panama.
grandis, Radlk. l. c. 635.—Bras.
guianensis, Aubl. Pl. Gui. i. 331.—Guian.
heterophylla, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 634.—Bras.
inelegans, Radlk. l. c. 627.—Bras.
juglandifolia, Radlk. l. c. 635.—Bras.
laevigata, Radlk. l. c. 628 = Cupania laevigata.
longipes, Radlk. l. c. 626.—Venezuela.
macrostylis, Radlk. l. c. 625.—Guian.
marginata, Radlk. l. c. 634.—Bras.
mexicana, Radlk. l. c. 624.—Mexic.

MATAYBA:—
mollis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 634.—Bras.
opaca, Radlk. l. c. 628 = Cupania laevigata.
Patrisiana, DC. Prod. i. 609.—Guian.
paucijuga, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 628.—Bras.
peruviana, Radlk. l. c. 627.—Peruv.
punctata, Radlk. l. c. 634.—Bras.
purgans, Radlk. l. c. 629.—Bras.
robusta, Radlk. l. c. 627.—Bras.
scrobiculata, Radlk. l. c.—N. Granat.
Spruceana, Radlk. l. c.—Bras.
sylvatica, Radlk. l. c. 631.—Bras.
tovarensis, Radlk. l. c. 626.—Venezuela.
Voua-rana, DC. Prod. i. 609.—Guian.

MATEATIA, Vell. Fl. Flum. ix. t. 95 (1827) = **Sterculia**, Linn.
cariosa, Steud. Nom. ed. II. ii. 105, sphalm. = seq.
curiosa, Vell. Fl. Flum. ix. t. 95 = S. carthaginensis.

MATELEA, Aubl. Pl. Gui. i. 277. t. 109 (1775). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 766.
HOSTEA, Willd. Sp. Pl. i. 1274 (1797).
laevis, Nutt. in Am. Journ. Sc. v. (1822) 291 = Asclepias perennis.
latifolia, *Aubl. Pl. Gui.* i. 278. t. 109. f. 2.—Guian.
linearis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 321 = Nephradenia linearis.
palustris, *Aubl. Pl. Gui.* i. 278. t. 109. f. 1.—Guian.

MATELLA, Bartl. Ord. 202 (1830) = praec.

MATHAEA, Vell. Fl. Flum. 22 (1825) = **Schwenkia**, Linn. (Solanac.).
apostolica, Vell. l. c.—Bras.

MATHEA, Vell. Fl. Flum. i. t. 51 (1827) = praec

MATHEWSIA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 140. t. 96. *CRUCIFERAE*, Benth. & Hook. f. i. 81.
MACHAEROPHORUS, Schlecht. in Linnaea, xxviii. (1856) 469.
foliosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 140. t. 94.—Chili.
incana, *Phil. Fl. Atac.* 8.—Chili.
linearifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 550.—Chili.
matthioloides, *C. Muell. in Walp. Ann.* vii. 144.—Peruv.

MATHIEUA, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 337 = **Eucharis**, Planch. (Amaryll.).
galanthoides, Klotzsch, l. c. = E. galanthoides.

MATHIOLA, R. Br. in Ait. Hort. Kew. ed. II. iv. 119 (1812) = **Matthiola**, R. Br. (Crucif.).

MATHIOLA, DC. Syst. ii. 162, in nota (1821) = MATTHIOLA, Plum. = **Guettarda**, Linn. (Rubiac.).

MATHURINA, Balf. f. in Journ. Linn. Soc. xv. (1876) 159. *TURNERACEAE*.
penduliflora, *Balf. f. l. c.* 160.—Ins. Mascar.

MATISIA, Humb. & Bonpl. Pl. Aequin. i. 9. t. 2 (1805). *MALVACEAE*, Benth. & Hook. f. i. 211.
alchornaefolia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 327.—N. Granat.
Castaño, *Triana & Karst. Fl. Neo-Granad.* 24; *ex iisd. l. c.* 326.—N. Granat.
cordata, *Humb. & Bonpl. Pl. Aequin.* i. 9. t. 2.—Am. austr.
Cornu-copiae, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 326.—N. Granat.
glandifera, *Planch. & Triana, l. c.* 327.—N. Granat.
oblongifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 35.—Peruv.

MATOUREA, Aubl. Pl. Gui. ii. 641. t. 259 (1775) = **Stemodia**, Linn. (Scrophul.).
dubia, G. Don, Gen. Syst. iv. 547 = Herpestis dubia.
guianensis, Aubl. Pl. Gui. ii. 642 = S. foliosa.
nigrescens, Benth. in Hook. Comp. Bot. Mag. i. (1835) 173 = Herpestis nigrescens.
pratensis, Aubl. Pl. Gui. ii. 642 = S. pusilla.

MATRELLA, Pers. Syn. i. 73 (1805) = **Zoysia**, Willd. (Gramin.).
juncea, Pers. l. c. = Z. pungens.

MATRICARIA, Hall. ex Scop. Fl. Carn. ed. II. ii. 147 (1772) = **Chrysanthemum**, Tourn. (Compos.).

MATRICARIA, [Tourn.] Linn. Gen. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 427.
Akylopsis, Lehm. Ind. Sem. Hort. Hamb. (1850) 3.
Chamaemelum, Vis. Fl. Dalm. ii. 84 (1847).
Chamomilla, [Hall.] S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 (1821).
Courrantia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276. t. 89 ('1835–60').
Dibothrospermum, Knaf, in Flora, xxix. (1846) 298.
Gastrostylum, Sch. Bip. ex Webb & Berth. Phyt. Canar. ii. 277 ('1835–60').
Gastrosulum, Sch. Bip. Tanacet. 29 (1844).
Lepidanthus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 396.
Lepidotheca, Nutt. in Trans. Am. Phil. Soc. vii. (1841) [454].
Rhytidospermum, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 277 ('1835–60').
Sphaeroclinium, Sch. Bip. Tanacet. 20 (1844).
Trallesia, Zumag. Fl. Pedem. i. 361 (1849).
Tripleurospermum, Sch. Bip. Tanacet. 31 (1844).
absinthioides, Desr. in Lam. Encyc. iii. 733 = Chrysanthemum millefolium.
achillaeifolia, Poir. Encyc. Suppl. v. 729 = Chrysanthemum achilleaefolium.
acutiloba, *Harv. in Harv. & Sond. Fl. Cap.* iii. 166. —Afr. austr.
africana, *Berg. Desc. Pl. Cap.* 296.—Afr. austr.
albida, *Fenzl, ex Harv. in Harv. & Sond. Fl. Cap.* iii. 166.—Afr. austr.
alpina, Desr. in Lam. Encyc. iii. 730 = Chrysanthemum alpinum.
americana, Mill. Gard. Dict. ed. VIII. n. 5 = Aster nemoralis.
anethifolia, Poir. Encyc. Suppl. iii. 605 = Chrysanthemum anethifolium.
anomala, Poir. l. c. 608 = Chrysanthemum anomalum.
apetala, Hort. Erf. ex DC. Prod. vi. 63 = Chrysanthemum Balsamita.
arctica, Desr. in Lam. Encyc. iii. 735 = Chrysanthemum arcticum.
argentea, Linn. Sp. Pl. 891 = Chrysanthemum argenteum.
arvensis, Baill. Hist. Pl. viii. 310 = Anthemis arvensis.
arvensis, Nym. Syll. 11 = glabra.
asteroides, Linn. Mant. i. 116 = Boltonia asteroides.
atrata, Schrank, Baier. Fl. ii. 406 = Chrysanthemum atratum.
aurea, Sch. Bip. in Bonplandia, viii. (1860) 369 = Cotula aurea.
auriculata, *Benth. & Hook. f. Gen.* ii. 428; *Ball, in Journ. Linn. Soc.* xvi. (1878) 512.—Palaest.; Arab.
Balsamita, Desr. in Lam. Encyc. iii. 733 = Chrysanthemum Balsamita.
Bayeri, Kanitz, in Magyar Tudom. Ertekez. (Comm. Scient. Hung.) (1862) 321 = M. Chamomilla.
bipinnata, Desr. in Lam. Encyc. iii. 736 = Chrysanthemum bipinnatum.
bipinnata, Spreng. Syst. iii. 582 = Lasiospermum radiatum.
Bocconi, Poir. Encyc. Suppl. iii. 607 = Chrysanthemum Bocconi.
brachyglossa, *Benth. & Hook. f. Gen.* ii. 428.—Hab.?
Broussonetii, Poir. Encyc. Suppl. iii. 608 = Chrysanthemum Broussonetii.
Burchellii, *Benth. & Hook. f. Gen.* ii. 428.—Afr. austr.
cantoniensis, Lour. Fl. Cochinch. 498 = A. indica.
capensis, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 681 = Anthemis parthenoides.
capensis, Linn. Mant. i. 115 = africana.
carinata, Poir. Encyc. Suppl. iii. 602 = Chrysanthemum carinatum.
carnea, Poir. l. c. v. 728 = Chrysanthemum coccineum.
carnosa, Desr. in Lam. Encyc. iii. 732 = Chrysanthemum paludosum.
caucasica, Poir. Encyc. Suppl. iii. 604 = Chrysanthemum caucasicum.
ceratophylloides, Poir. l. c. 601 = Chrysanthemum atratum.

MATRICARIA:—
Chamaemilla, Hill, Veg. Syst. ii. 98 = M. Chamomilla.
Chamomilla, Blanco, Fl. Filip. ed. I. 631 = Chrysanthemum indicum.
Chamomilla, *Linn. Sp. Pl.* 891.—Europ.
chia, Baill. Hist. Pl. viii. 310 = Anthemis chia.
coccinea, Poir. Encyc. Suppl. iii. 602 = Chrysanthemum coccineum.
confusa, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* x. 53.—Hab.?
conoclinia, *Nym. Consp.* 374.—Byzant.
coronaria, Desr. in Lam. Encyc. iii. 737 = Chrysanthemum coronarium.
coronata, J. Gay, ex Koch, Syn. Fl. Germ. ed. II. 416 = M. Chamomilla.
coronopifolia, Poir. Encyc. Suppl. iii. 609 = Chrysanthemum coronopifolium, *Masf.*
corymbifera, *DC. Prod.* vii. 297.—As. Min.; Persia.
corymbosa, Desr. in Lam. Encyc. iii. 734 = Chrysanthemum corymbosum.
Cotula, Baill. Hist. Pl. viii. 310 = Anthemis Cotula.
Courrantiana, DC. Prod. vi. 52 = M. Chamomilla.
crithmifolia, Poir. Encyc. Suppl. iii. 609 = Chrysanthemum anethifolium.
daucifolia, Poir. l. c. 608 = Chrysanthemum daucifolium.
decipiens, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 14.—As. Min.; Persia.
deflexa, Gilib. Fl. Lituan. i. 221 = M. Chamomilla.
densa, Poir. Encyc. Suppl. iii. 606 = Chrysanthemum densum.
dichotoma, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 167. —Afr. austr.
disciformis, DC. Prod. vi. 51 = corymbifera.
discoidea, *DC. l. c.* 50.—As. et Am. bor.
elegans, Nym. Consp. 374 = inodora.
feniculacea, Poir. Encyc. Suppl. iii. 605 = Chrysanthemum anethifolium.
florida, Salisb. Prod. 203 = Chrysanthemum Parthenium.
frutescens, Desr. in Lam. Encyc. iii. 730 = Chrysanthemum frutescens.
fuscata, *Poir. Encyc. Suppl.* iii. 602; *Ball, in Journ. Linn. Soc.* xvi. (1878) 512.—Afr. bor.
glabra, *Ball, l. c.* 511.—Marocc.
glabrata, *Poir. Encyc. Suppl.* iii. 609; *DC. Prod.* vi. 51.—Afr. austr.
glastifolia, Hill, Hort. Kew. 19 = Boltonia glastifolia.
glauca, Poir. Encyc. Suppl. iii. 603 = Chrysanthemum multifidum.
globifera, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 165. —Afr. austr.
graminifolia, Desr. in Lam. Encyc. iii. 732 = Chrysanthemum graminifolium.
grandiflora, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 166.—Afr. austr.
grandiflora, C. Koch, in Linnaea, xxiv. (1851) 336 = Chrysanthemum caucasicum.
grandiflora, Poir. Encyc. Suppl. iii. 609 = Chrysanthemum grandiflorum.
grandis, Desr. in Lam. Encyc. iii. 738 (= *Plagius grandiflorus*).—Afr. bor.
Halleri, Poir. Encyc. Suppl. iii. 603 = Chrysanthemum coronopifolium.
heterocarpa, *C. Koch, in Linnaea*, xxiv. (1851) 333.—As. Min.
heterophylla, Poir. Encyc. Suppl. iii. 600 = Chrysanthemum heterophyllum.
hirta, *Poir. l. c.* 609; *DC. Prod.* vi. 51.—Afr. austr.
hispida, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 194.—Afr. trop.
incana, Poir. Encyc. Suppl. iii. 609 = Pentzia virgata.
inciso-serrata, Poir. l. c. 604 = Chrysanthemum uliginosum.
indica, Mill. Gard. Dict. ed. VIII. n. 3 = Chrysanthemum indicum.
inodora, Lam. Fl. Fr. ii. 136 = Chrysanthemum corymbosum.
inodora, *Linn. Fl. Suec.* ii. 765.—Europ.; Oriens.
italica, Desr. in Lam. Encyc. iii. 736 = Chrysanthemum italicum.
japonica, Desr. l. c. 738 = Chrysanthemum indicum.
Kochianum, Sch. Bip. Tanacet. 26 = M. Chamomilla.
lacustris, Poir. Encyc. Suppl. iii. 608 = Chrysanthemum lacustre.
lamellata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 335.—Turkestan.

MATRICARIA:—
lasiocarpa, *Boiss. Fl. Orient.* iii. 324.—Beluchist.
latifolium, Gilib. Fl. Lituan. i. 220 = Chrysanthemum Parthenium.
laxa, *Tratt.* (*Arch.* t. 200; *ex Pritz. Ic. Ind.* 681) *ex Steud. Nom.* ed. I. 512.—Hab.?
Leucanthemum, Scop. Fl. Carn. ed. II. ii. 147 = Chrysanthemum Leucanthemum.
macrophylla, Poir. Encyc. Suppl. iii. 603 = Chrysanthemum macrophyllum.
maritima, Fries, ex Nym. Consp. 374 = inodora.
maritima, *Linn. Sp. Pl.* 891.—Europ.
maroccana, *Ball, in Journ. Bot.* xi. (1873) 367.—Morocc.
maxima, Poir. Encyc. Suppl. iii. 600 = Chrysanthemum maximum.
microcephala, *C. Koch, in Linnaea*, xxiv. (1851) 336. —Reg. Caucas.
millefoliata, Desr. in Lam. Encyc. iii. 736 = Chrysanthemum millefoliatum.
minima, Desr. l. c. 731 = Chrysanthemum alpinum, pulverulentum.
modesta, Boiss. ex Boiss. Fl. Orient. iii. 326 = praecox.
monspeliensis, Desr. in Lam. Encyc. iii. 733 = Chrysanthemum monspeliense, arcticum.
montana, Desr. in Lam. Encyc. iii. 732 = Chrysanthemum montanum.
multicaulis, Poir. Encyc. Suppl. iii. 606 = Chrysanthemum multicaule.
multifida, Baill. Hist. Pl. viii. 310 = Cotula villosa.
multiflora, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 166. —Afr. austr.
Myconis, Desr. in Lam. Encyc. iii. 736 = Chrysanthemum Myconis.
nigellaefolia, *DC. Prod.* vi. 50.—Afr. austr.
nigrescens, Baill. Hist. Pl. viii. 310 = Anthemis austriaca.
nobilis, Baill. l. c. 60, 309 = Anthemis nobilis.
obliqua, Dulac, Fl. Hautes-Pyr. 505 = M. Chamomilla.
odorata, Lam. Fl. Fr. ii. 136 = Chrysanthemum Parthenium.
oleracea, Buch.-Ham. in Wall. Cat. n. 3229 = Chrysanthemum coronarium.
oreades, *Boiss. Diagn.* Ser. I. vi. 87.—Syria.
orientalis, Poir. Encyc. Suppl. iii. 607 = Anthemis Biebersteiniana.
pallida, Poir. l. c. 607 = Chrysanthemum pulverulentum, Anthemis mixta.
paludosa, Poir. l. c. 600 = Chrysanthemum paludosum.
partenifolium, Poir. l. c. 608 = Chrysanthemum praealtum.
Parthenium, Linn. Sp. Pl. 890 = Chrysanthemum Parthenium.
parthenoides, Desf. Cat. Hort. Par. ed. III. 169, 400 = Anthemis parthenoides.
parviflora, Poir. Encyc. Suppl. iii. 608 = praecox.
patens, Gilib. Fl. Lituan. i. 220 = M. Chamomilla.
pectinata, Desr. in Lam. Encyc. iii. 735 = Chrysanthemum pectinatum.
perforata, Mérat, Fl. Par. ed. I. 332 = inodora.
pilosa, Poir. Encyc. Suppl. iii. 602 = Chrysanthemum tanacetifolium.
pinnatifida, Desr. in Lam. Encyc. iii. 729 = Chrysanthemum pinnatifidum.
praealta, Poir. Encyc. Suppl. iii. 602 = Chrysanthemum praealtum.
praecox, *Poir. l. c.* v. 729; *DC. Prod.* vi. 52.—As. Min.; Persia.
procumbens, Poir. l. c. iii. 604 = Chrysanthemum indicum.
prostrata, Sw. Prod. Veg. Ind. Occ. 114 = Egletes domingensis.
ptarmicaefolia, Poir. Encyc. Suppl. iii. 603 = Achillea Ptarmica.
pubescens, Sch. Bip. in Bonplandia, viii. (1860) 369 = Cotula pubescens.
pubesceus, Ball, in Journ. Linn. Soc. xvi. (1878) 512, sphalm. = praec.
pulverulenta, Poir. Encyc. Suppl. iii. 605 = Chrysanthemum praealtum
pumila, Nym. Syll. 12 = inodora.
punctata, Desr. in Lam. Encyc. iii. 732 = Chrysanthemum palustre.
pusilla, Willd. Enum. Hort. Berol. 907 = M. Chamomilla.
pyrethrifolia, Poir. Encyc. Suppl. iii. 605 = praecox.
pyrethroides, DC. Prod. vi. 52 = M. Chamomilla.

MATRICARIA:—
Pyrethrum, Baill. Hist. Pl. viii. 61, 310 = Anacyclus Pyrethrum.
radicans, Poir. Encyc. Suppl. iii. 608 = Chrysanthemum radicans.
recutita, Linn. Sp. Pl. 891 = M. Chamomilla.
rigida, Desr. in Lam. Encyc. iii. 737 = Chrysanthemum flosculosum.
rosea, Baill. Hist. Pl. viii. 310 = Anthemis rosea.
rosella, *Nym. Consp.* 374.—Graecia.
rotundifolia, Poir. Encyc. Suppl. iii. 608 = Chrysanthemum rotundifolium.
salina, Nym. Consp. 374 = inodora.
salina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 40 = M. Chamomilla.
segetum, Schrank, Baier. Fl. ii. 406 = Chrysanthemum segetum.
sericea, Poir. Encyc. Suppl. v. 729 = Chrysanthemum argenteum.
serotina, Desr. in Lam. Encyc. iii. 730 = Chrysanthemum serotinum.
suaveolens, Koch, Syn. Fl. Germ. ed. II. 416, 1026 = inodora.
suaveolens, Linn. Fl. Suec. 297 = M. Chamomilla.
tanacetifolia, Desr. in Lam. Encyc. iii. 733 = Chrysanthemum multifidum.
tanacetoides, Fisch. & Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vii. 52 = discoidea.
tenella, *DC. Prod.* vi. 51.—Afr. austr.
tenuifolia, Poir. Encyc. Suppl. iii. 609 = Chrysanthemum tenuifolium.
tenuifolia, Salisb. Prod. 208 = M. Chamomilla.
tinctoria, Baill. Hist. Pl. viii. 310 = Anthemis tinctoria.
trichophylla, *Boiss. Diagn.* Ser. I. vi. 88.—As. Min.
tridentata, *Ball, in Journ. Linn. Soc.* xvi. (1878) 512. —Marocc.
trifurcata, Poir. Encyc. Suppl. iii. 606 = Chrysanthemum trifurcatum.
uliginosa, Poir. l. c. 608 = Chrysanthemum uliginosum.
umbrosa, Poir. l. c. 606 = Chrysanthemum umbrosum.
uniglandulosa, C. Koch, in Linnaea, xxiv. (1851) 333 = trichophylla.
virgata, Desr. in Lam. Encyc. iii. 737 = Chrysanthemum flosculosum.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 = M. Parthenium.

MATRICARIOIDES, Spach, Hist. Vég. Phan. x. 24 (1841) = **Tanacetum**, Tourn. (Compos.).

MATTHAEA, Blume, Mus. Bot. Lugd. Bat. ii. 89. t. 10. 1852 [1856]. *MONIMIACEAE*, Benth. & Hook. f. iii. 141.
sancta, *Blume, Mus. Bot. Lugd. Bat.* ii. 90.—Malaya.

MATTHEWSIA, Reichb. Handb. 260 (1837) = **Mathewsia**, Hook. & Arn. (Crucif.).

MATTHIOLA, R. Br. in Ait. Hort. Kew. ed. II. iv. 119 (1812) (Mathiola). *CRUCIFERAE*, Benth. & Hook. f. i. 67.
LEUCOIUM, Tourn. ex Adans. Fam. ii. 418 (1763).
MICROSTIGMA, Trautv. Pl. Imag. 34. t. 35 (1845).
TRICERAS, Andrz. ex Reichb. Consp. 185 (1828).
acaulis, DC. Syst. ii. 168 = humilis.
albicaulis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 46.—Persia.
anchusoides, Webb, in Parl. Giorn. Bot. Ital. ii. II. (1847) 213 = aspera.
angulosa, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 52 = oxyceras.
annua, Sweet, Hort. Suburb. Lond. 147 = incana.
anoplia, *Pomel, Nouv. Mat. Fl. Atl.* 231.—Alger.
arabica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 49.—Arabia.
arborescens, *Sternb. in Syll. Ratisb.* i. (1824) 188.—Creta.
aspera, *Boiss. Diagn.* Ser. I. viii. 16.—Palaestin.
bicornis, *DC. Syst.* ii. 177.—Graecia; As. Min.
biennis, Nym. Consp. 974 (Index) = praec.
capiomontiana, *Pomel, Nouv. Mat. Fl. Atl.* 374.—Alger.
chenopodifolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 33.—Reg. Casp.
chorassanica, *Bunge, ex Boiss. Fl. Orient.* i. 151.—Persia.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 230.—Alger.
coronopifolia, DC. Syst. ii. 173 = tristis.

MATTHIOLA:—
crassifolia, *Boiss. & Gaill. in Boiss. Fl. Orient.* i. 149. —Syria.
crucigera, *DC. Syst.* ii. 177.—Sicil.
damascena, *Boiss. Diagn.* Ser. I. viii. 16.—Syria.
deflexa, *Bunge, Del. Sem. Hort. Dorpat.* (1839) 8.—*Cf. Linnaea*, xiv. (1840) 120.—Sibir. altaic.
dimorpha, Hort. ex Bernh. Sel. Sem. Hort. Erf. (1835). —Cf. Linnaea, xi. (1837) Litt. 85 = Diptychocarpus strictus.
dumulosa, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 14.—Persia.
elliptica, *R. Br. in Salt, Abyss. App.* 65, nomen.—Abyssin.
euboea, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 48 = damascena.
farinosa, *Bunge, ex Boiss. Fl. Orient.* i. 150.—Persia.
fasciculata, *DC. Syst.* ii. 714.—Sicil.
fenestralis, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 419.—Ins. Creta.
Fischeri, Bernh. Sel. Sem. Hort. Erf. (1835).—Cf. Linnaea, xi. (1837) Litt. 85 = Diptychocarpus strictus.
flavida, *Boiss. Diagn.* Ser. I. vi. 9.—Persia.
fragrans, *Bunge, Del. Sem. Hort. Dorpat.* (1839) 8.—Ross. mer.
glabra, DC. Syst. ii. 165 = incana.
glabrata, DC. Prod. i. 133 = praec.
glandulosa, *Pomel, Nouv. Mat. Fl. Atl.* 373.—Alger.
glandulosa, *Vis. Fl. Dalm.* iii. 124.—Dalmat.
graeca, Sweet, Hort. Suburb. Lond. 147 = incana.
hesperoides, Less. ex Ledeb. Fl. Ross. i. 110 = chenopodifolia.
humilis, *DC. Syst.* ii. 177.—Aegypt.
incana, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 119.—Reg. Mediterr.
Kralikii, *Pomel, Nouv. Mat. Fl. Atl.* 374.—Alger.
lacera, *Spreng. Syst.* ii. 897, 899.—Hispan.
linearis, Delile, in Laborde, Voy. Arab. Pétr. 85 = Malcomia aegyptiaca.
livida, *DC. Syst.* ii. 174.—Aegypt.; Arab.
livida, Kral. ex Pomel, Nouv. Mat. Fl. Atl. 374 = Kralikii.
lonchophora, *Pomel, l. c.*—Alger.
longipetala, DC. Syst. ii. 174 = oxyceras.
lunata, *DC. l. c.* 176.—Hispan.
macropetala, Spreng. Syst. ii. 897 = oxyceras.
maderensis, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 29.—Ins. Mader.
maroccana, *Coss. in Bull. Soc. Bot. Fr.* xxvii. (1880) 68.—Marocc.
montana, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 47.—As. Min.
montana, *Pomel, Nouv. Mat. Fl. Atl.* 230.—Alger.
nana, *Spreng. Syst.* ii. 897.—Sibir.
nudicaulis, *Trautv. in Act. Hort. Petrop.* i. (1871) 51. —As. centr.
obovata, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 136.—Turkestan.
odoratissima, Boiss. & Buhse, ex Boiss. Fl. Orient. i. 150, partim = ovatifolia.
odoratissima, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 120. —Reg. Caucas.
odoratissima, Claus, in Goebel, Reise, ii. 255 = fragrans.
ovatifolia, *Boiss. Diagn.* Ser. I. vi. 9.—As. Min.; Persia.
oxyceras, *DC. Syst.* ii. 173.—Syria; Persia.
oxyceratia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = praec.
oyensis, Menier & Viaud, in Bull. Soc. Bot. Fr. xxiv. (1877) 203 = sinuata.
parviflora, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 121.—Reg. Mediterr. occ.
patens, *Presl, Delic. Prag.* 17.—Sicil.
persica, DC. Syst. ii. 168 = Anchonium Tournefortii.
Phlox, *Didr. Ind. Sem. Hort. Haun.* (1865) 27.—Alger.
prostrata, *Hochst. & Steud. ex Steud. Nom.* ed. II. ii. 106.—Arabia.
pumila, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
Pumilio, DC. Syst. ii. 177 = bicornis.
revoluta, *Bunge, ex Boiss. Fl. Orient.* i. 151.—Persia.
robusta, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 137.—Reg. Casp.
rostrata, Haberle, ex Bernh. Sel. Sem. Hort. Erf. (1835).—Cf. Linnaea, xi. (1837) Litt. 85 = Diptychocarpus strictus.

MATTHIOLA:—
runcinata, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) 1. 254.—As. temp.
rupestris, Bernh. ex Steud. Nom. ed. II. ii. 106 = saxatilis.
rupestris, DC. Syst. ii. 714 = incana.
saxatilis, *Bernh. ex Steud. Nom.* ed. II. ii. 106.—Lusitan.
simplicicaulis, Sweet, Hort. Brit. ed. I. 17 = incana.
sinuata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 120.—Reg. Mediterr.
stelligera, *Sond. in Linnaea*, xxii. (1849) 1.—Afr austr.
stenopetala, *Bernh. ex Steud. Nom.* ed. II. ii. 106.—Sibir. altaic.
stenopetala, Coss. in Bull. Soc. Bot. Fr. xxvii. (1880) 68 = maroccana.
stenopetala, *Pomel, Nouv. Mat. Fl. Atl.* 372.—Alger.
Stoddarti, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 138.—Turkestan.
stylosa, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 106 = Farsetia Hamiltonii.
tatarica, *DC. Syst.* ii. 170.—Rossia; Turkest.
Telum, *Pomel, Nouv. Mat. Fl. Atl.* 372.—Alger.
tenella, DC. Syst. ii. 169 = oxyceras.
thessala, *Boiss. & Orph. in Boiss. Fl. Orient.* i. 153.—Thessalia.
tomentosa, Bélang. Voy. Indes Or., Ic. absque num. = ovatifolia.
torulosa, *DC. Syst.* ii. 169.—Afr. austr.
tricuspidata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 120. —Reg. Mediterr.
tristis, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 120.—Reg. Mediterr.
tristis, Decne. ex Boiss. Fl. Orient. i. 152 = arabica.
undulata, Tineo, Cat. Pl. Hort. Panorm. (1827) 279 = incana.
valesiaca, J. Gay, ex Nym. Consp. 30 = varia.
varia, *DC. Syst.* ii. 171.—Ital.; Graec.
varia, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 807 = Erysimum linifolium.
vulgaris, Spach, Hist. Vég. Phan. vi. 403 = incana, fenestralis.

MATTHIOLA, Plum. ex Linn. Gen. ed. I. 49 (1737) = **Guettarda**, Linn. (Rubiac.).
scabra, Linn. Sp. Pl. ed. I. 1192 = G. Matthiola.

MATTHISONIA, Lindl. Veg. Kingd. 795 (1847) = seq.

MATTHISSONIA, Raddi, in Mem. Mod. xviii. Fis. (1820) 390 = **Schwenkia**, Linn. (Solan.).
paniculata, Raddi, l. c.—Bras.

MATTIA, Schult. Obs. Bot. 30, 32 (1809) = **Rindera**, Pall. (Borag.).
albida, Wettstein, ex Stapf, in Denkschr. Acad. Wien, l. (1885) 32.—Persia.
angustifolia, G. Don, Gen. Syst. iv. 310 = R. angustifolia.
Aucheri, DC. & A. DC. in Prod. x. 169 = Paracaryum Aucheri.
brachyantha, *Boiss. Diagn.* Ser. I. xi. 127.—As. Min.; Persia.
bracteata, *Orph. ex Boiss. Fl. Orient.* iv. 274.—Graecia.
Bungei, Boiss. l. c.—Persia.
caespitosa, A. DC. in DC. Prod. x. 168 = R. caespitosa.
canescens, A. DC. l. c. = R. canescens.
Columnae, G. Don, Gen. Syst. iv. 310 = Cynoglossum Columnae.
corymbiformis, DC. & A. DC. Prod. x. 169 = Paracaryum corymbiforme.
cristata, G. Don, Gen. Syst. iv. 310 = Paracaryum cristatum.
emarginata, Roem. & Schult. Syst. iv. 83, 765 = Paracaryum angustifolium.
eriantha, Ledeb. Fl. Ross. iii. 173 = R. pubescens.
glastifolia, G. Don, Gen. Syst. iv. 310 = Paracaryum glastifolium.
graeca, A. DC. in DC. Prod. x. 168 = R. graeca.
gymnandra, Coss. in Bull. Soc. Bot. Fr. iii. (1856) 708.—Alger.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 94 = Paracaryum himalayense.
incana, Ledeb. Fl. Ross. iii. 173 = Paracaryum incanum.
laevigata, Schult. Obs. Bot. 31 = R. tetraspis.

MATTIA:—
lanata, Schult. Obs. Bot. 31 (= *Cyphomattia lanata*).—As. Min.; Persia.
lateriflorum, G. Don, Gen. Syst. iv. 310 = Pectocarya lateriflora.
leptophylla, DC. & A. DC. Prod. x. 170 = Paracaryum leptophyllum.
magellensis, G. Don, Gen. Syst. iv. 310 = Cynoglossum magellense.
pilosa, G. Don, l. c. = Pectocarya lateriflora.
punctata, A. DC. in DC. Prod. x. 167 = Rindera pubescens.
Schlumbergeri, Boiss. Fl. Orient. iv. 274.—Syria.
Schmidtii, Heldr. in Ann. Sc. Nat. Sér. IV. xiii. (1860) 382 = R. graeca.
staminea, Roem. & Schult. Syst. iv. 82, 764 = Solenanthus Tournefortii.
umbellata, C. Koch, in Linnaea, xvii. (1843) 302 = R. pubescens.
umbellata, Schult. Obs. Bot. 32.—Europ. austr.

MATTIOLA, Sang. in Att. Acc. Pont. Lincei, Ser. I. xv. (1862) 36 = **Matthiola**, R. Br. (Crucif.).

MATTUSCHKAEA, Schreb. Gen. 788 (1791) = **Perama**, Aubl. (Rubiac.).
galioides, H. B. & K. Nov. Gen. et Sp. ii. 272 = P. galioides.
hirsuta, Vahl, Symb. Bot. iii. 11 = P. hirsuta.
hispida, H. B. & K. Nov. Gen. et Sp. ii. 271 = P. hispida.
incana, Spreng. Syst. i. 418.—Bras.
Perama, Schult. Mant. iii. 47 = P. hirsuta.

MATTUSCHKEA, Batsch, Tab. 193 (1802) = praec.

MATTUSCHKIA, J. F. Gmel. Syst. 589 (1791) = **Saururus**, Plum. (Piperac.).
aquatica, J. F. Gmel. l. c. = S. cernuus.

MATTUSKEA, Rafin. Princ. Somiol. 30 (1814) = MATTUSCHKAEA, Schreb. = **Perama**, Aubl. (Rubiac.).

MATURNA, Rafin. Fl. Tellur. ii. 99 (1836) = **Gomesa**, R. Br. (Orchid.).
suaveolens, Rafin. l. c. = G. foliosa.

MAUCHARTIA, Neck, Elem. i. 172 (1790) = **Apium**, Tourn. (Umbellif.).

MAUDUITA, Comm. ex DC. Prod. i. 592 (1824) = **Samadera**, Gaertn. (Simarub.).

MAUDUYTA, Comm. ex Endl. Gen. 1144 (1841) = praec.

MAUHLIA, Dahl, Obs. Syst. Linn. 25 (1787) = **Agapanthus**, L'Hérit. (Liliac.).
africana, Dahl, l. c. 26 = A. umbellatus.
ensifolia, Thunb. Nov. Gen. 111 = Massonia ensifolia.
latifolia, Thunb. Mus. Upsal. ex Schult. f. Syst. vii. 999 (Quid?).—Hab.?
linearis, Thunb. Nov. Gen. 111 = A. umbellatus.
tortilis, Thunb. Mus. Upsal. ex Schult. f. Syst. vii. 999 (Quid?).—Hab.?
umbellata, Thunb. Hort. Upsal. ex Schult. f. l. c. 997 = A. umbellatus.

MAUKSCHIA, Heuff. in Flora, xxvii. (1844) 527 = **Carex**, Dill. (Cyperac.).
laevis, Heuff. l. c. = C. dioica.
scabra, Heuff. l. c. 528 = C. Davalliana.

MAUNDIA, F. Muell. Fragm. i. 22 (1858) = **Triglochin**, Linn. (Naiad.).
triglochinoides, F. Muell. l. c. 23 = T. Maundii.

MAUNEIA, Thou. Nov. Gen. Madag. 6 (1806). *INC. SED.*
madagascariensis, *Steud. Nom.* ed. II. ii. 107.—Madag.

MAURANDIA, Orteg. Hort. Matr. Dec. 21 (1797). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 935.
LOPHOSPERMUM, D. Don, in Trans. Linn. Soc. xv. (1827) 359.
REICHARDIA, Roth, Catal. ii. 64 (1800).
USTERIA, Cav. Ic. ii. 15. t. 116 (1793).
antirrhiniflora, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 659 = Antirrhinum maurandioides.

MAURANDIA:—
atrosanguinea, Nichols. Dict. Gard. ii. 337 = Rhodochiton volubile.
Barclaiana, *Lindl. Bot. Reg.* t. 1108.—Mexic.
Barclayana, Auct. = praec.
erecta, *Hemsl. in Gard. Chron.* (1882) i. 22.—Mexic.
erubescens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 377.—Mexic.
uncea, Benth. Bot. Voy. Sulph. 41 = Antirrhinum junceum.
personata, Lag. Gen. et Sp. Nov. 19 = Antirrhinum maurandioides.
scandens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 377.—Mexic.
scandens, Pers. Syn. ii. 160 = semperflorens.
semperflorens, *Jacq. Hort. Schoenb.* iii. 20.—Mexic.
stricta, Hook. & Arn. Bot. Beech. Voy. 375 = Antirrhinum strictum.
Wislizeni, *Engelm. in Torr. Bot. Mex. Bound.* 111.—Mexic.

MAURANDYA, G. Don, Gen. Syst. iv. 532 (1837) = praec.

MAURIA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 338. *ANACARDIACEAE*, Benth. & Hook. f. i. 426.
aurantiodora, *Engl. in DC. Monog. Phan.* iv. 326.—Peruv.
Biringo, *Tul. in Ann. Sc. Nat.* Sér. III. vi. (1846) 365.—N. Granat.
cyclocarpa, Mart. Herb. Fl. Bras. n. 1271 = Tapirira guianensis.
decandra, Karst. ex Engl. in DC. Monog. Phan. iv. 327 = heterophylla.
ferruginea, *Tul. in Ann. Sc. Nat.* Sér. III. vi. (1846) 366.—N. Granat.
guianensis, Brongn. ex Engl. in DC. Monog. Phan. iv. 301 = Sorindeia madagascariensis.
heterophylla, *H. B. & K. Nov. Gen. et Sp.* vii. 11. t. 606.—Peruv.
juglandifolia, Benth. ex Engl. in DC. Monog. Phan. iv. 244 = Spondias lutea.
multiflora, *Mart. ex Benth. in Hook. Kew Journ.* iv. (1852) 14.—Bras.
ovalifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 467.—N. Granat.
puberula, *Tul. in Ann. Sc. Nat.* Sér. III. vi. (1846) 363.—N. Granat.
Schickendantzii, Hieron. & Lor. ex Engl. in DC. Monog. Phan. iv. 344 = Schinus latifolius.
Seemanni, Planch. & Oerst. ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xiv. (1872) 292 = puberula.
simplicifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 11. t. 605.—Chili.
suaveolens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 77.—Peruv.; Guian.
subbijuga, Mart. Herb. Fl. Bras. n. 1275 = Tapirira guianensis.

MAURITIA, Linn. f. Suppl. 70 (1781). *PALMAE*, Benth. & Hook. f. iii. 937.
LEPIDOCOCCUS, H. Wendl. & Drude, in Kerch. Palm. 249 (1878).
OROPHOMA, Drude, in Mart. Fl. Bras. iii. II. 294 (1882).
aculeata, *H. B. & K. Nov. Gen. et Sp.* i. 311.—Am. austr.
aculeata, Mart. Hist. Nat. Palm. ii. 47. t. 39. f. 3, 4 et t. 44 = M. Martiana.
armata, *Mart. l. c.* 45. tt. 41–43.—Bras.
Carana, *Wallace, ex Archer, in Hook. Kew Journ.* vii. (1855) 218.—Am. trop.
casiquiarensis, Spruce, in Journ. Linn. Soc. xi. (1869) 173 = Lepidocaryum tenue.
flexuosa, *Linn. f. Suppl.* 454.—Am. austr.
gracilis, *Mart. ex Spruce, in Journ. Linn. Soc.* xi. (1869) 169.—Am. trop.
gracilis, Wallace, Palm-trees Amaz. 57, t. 20 = aculeata.
guainiensis, Spruce, in Journ. Linn. Soc. xi. (1869) 174 = Lepidocaryum guianiense.
limnophylla, Barb.-Rodr. Enum. Palm. Nov. 18 = aculeata.
Martiana, *Spruce, in Journ. Linn. Soc.* xi. (1869) 171.—Am. trop.
Piritu, *Linden, in Illustr. Hortic.* xxviii. (1881) 32.—Bras.
pumila, *Wallace, Palm-trees Amaz.* 59. t. 21.—Am. trop.

MAURITIA:—
quadripartita, *Spruce, in Journ. Linn. Soc.* xi. (1869) 172.—Am. trop.
Sagus, *Schult. f. Syst.* vii. 1321.—Guiana.
setigera, *Griseb. & H. Wendl. ex Griseb. Fl. Brit. W. Ind.* 515.—Ind. occ.
subinermis, *Spruce, in Journ. Linn. Soc.* xi. (1869) 171.—Bras.
tenuis, *Mart. ex Spruce, l. c.* 169.—Am. trop.
vinifera, *Mart. Hist. Nat. Palm.* ii. 42. tt. 38, 39. f. 1, 2.—Bras.

MAUROCENA, Linn. Syst. ed. II. 21 (1740) = seq.

MAUROCENIA, Linn. Gen. ed. I. 85 (1737) = **Cassine**, Linn. (Celastr.).
americana, Mill. Gard. Dict. ed. VIII. n. 4.—Jamaic.
capensis, Sond. in Harv. & Sond. Fl. Cap. i. 465 = C. Maurocenia.
Cerasus, Mill. Gard. Dict. ed. VIII. n. 3.—Afr. austr.
Frangula, Mill. l. c. n. 1 = C. Maurocenia.
Frangularia, Pers. Syn. i. 327, in obs. = praec.
Phylliraea, Mill. Gard. Dict. ed. VIII. n. 2 = C. capensis.

MAUROCENNA, Linn. Phil. Bot. 29 (1751) = praec.

MAVIA, Bertol. f. in Mem. Acc. Sc. Bolog. ii. (1850) 570. t. 39 = **Erythrophleum**, Afzel. (Legumin.).
judicia, Walp. Ann. ii. 461 = seq.
judicialis, Bertol. f. in Mem. Acc. Sc. Bolog. ii. (1850) 570. t. 39 = E. guineense.

MAXILLARIA, Ruiz & Pav. Prod. 116. t. 25 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 555.
DICRYPTA, Lindl. Gen. et Sp. Orch. 44, 152 (1830).
HETEROTAXIS, Lindl. Bot. Reg. t. 1028 (1826).
MENADENA, Rafin. Fl. Tellur. ii. 98 (1836).
ONKERIPUS, Rafin. l. c. iv. 42 (1836).
PENTULOPS, Rafin. l. c. (1836).
PSITTACOGLOSSUM, La Llave & Lex. Nov. Veg. Desc. ii. Orch. 29 (1825).
acervata, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Costa Rica.
aciantha, *Reichb. f. in Bot. Zeit.* x. (1852) 858.—Guatemala.
acicularis, *Herb. ex Lindl. Bot. Reg.* sub t. 1986.—Bras.
acuminata, *Lindl. in Benth. Pl. Hartw.* 155.—Peruv.
acutifolia, Lindl. Bot. Reg. (1839) Misc. 92 = rufescens.
acutipetala, *Hook. Bot. Mag.* t. 3966.—Am. centr.
affinis, Hort. Petrop. ex Reichb. f. in Walp. Ann. vi. 511 = *hyacinthina*.
aggregata, Lindl. Gen. et Sp. Orch. 151 = Ornithidium aggregatum.
alata, *Ruiz & Pav. Syst. Veg.* 223.—Peruv.
alba, *Lindl. Gen. et Sp. Orch.* 143.—Am. austr.
albata, *Lindl. Orch. Linden.* 19.—N. Granat.
alpestris, *Lindl. in Benth. Pl. Hartw.* 154.—Ecuador.
anatamorum, *Reichb. f. in Bot. Zeit.* x. (1852) 935.—Am. austr.
angustifolia, Hook. Ic. Pl. t. 348 = variabilis.
Arachnites, *Reichb. f. in Gard. Chron.* (1880) i. 394.—N. Granat.
Arbuscula, *Reichb. f. in Bonplandia*, iv. (1856) 213.—Peruv.
argyrophylla, *Poepp. & Endl. Nov. Gen. et Sp.* i. 36.—Peruv.
aromatica, R. Grah. in Edinb. N. Phil. Journ. (Apr.–Jul. 1826) 173; et in Hook. Exot. Fl. t. 219 = Lycaste aromatica.
articulata, Klotzsch, Ind. Sem. Hort. Berol. (1853) 11 = rufescens.
atrata, *Reichb. f. Beitr. Orch. Centr. Am.* 31. t. 6.—Guatemala.
atropurpurea, Lodd. Bot. Cab. t. 1877 = Bifrenaria atropurpurea.
aurantiaca, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 25 = Bifrenaria aurantiaca?
aurea, Hort. ex Reichb. f. in Bonplandia, iii. (1855) 216 = porrecta.
aureofulva, Hook. Bot. Mag. t. 3629 = Bifrenaria aureofulva.
balsamea, Beer, Prakt. Orch. 264 = Lycaste balsamea.
barbata, Knowles & Westc. Phyt. 7; ex Reichb. f. in Walp. Ann. vi. 549 = Bifrenaria vitellina.
Barkerii, *Lindl. ex Beer, Prakt. Orch.* 266.—Hab.?

MAXILLARIA:—
Barringtoniae, Hort. ex Reichb. f. Xenia Orch. i. 224 = Bifrenaria Harrisoniae.
Barringtoniae, Lodd. Bot. Cab. t. 1824 = Lycaste Barringtoniae.
Batemanni, *Poepp. & Endl. Nov. Gen. et Sp.* i. 38. t. 65.—Peruv.
bicolor, *Ruiz & Pav. Syst. Veg.* 224.—Peruv.
Boothii, Lindl. Bot. Reg. (1838) Misc. 52 = Epidendrum auritum.
brachyglossa, *Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 24.—Mexic.
brachypus, Reichb. f. in Bot. Zeit. x. (1852) 734 = Xylobium brachypus.
bractescens, Lindl. Bot. Reg. (1842) Misc. 84 (Xylobii sp.).—Peruv.
brevifolia, *Reichb. f. in Walp. Ann.* vi. 540.—Ecuador.
breviscapa, *Poepp. & Endl. Nov. Gen. et Sp.* i. 36.—Peruv.
brevispatha, Klotzsch, in Otto & Dietr. Allg. Gartenz. xix. (1851) 217 = Lycaste candida.
Brocklehurstiana, Lindl. Bot. Reg. (1841) Misc. 10 = Houlletia Brocklehurstii.
brunnea, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 281.—N. Granat.
caespitifica, *Reichb. f. in Linnaea*, xli. (1877) 73.—Costa Rica.
callichroma, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Venezuela.
caloglossa, *Reichb. f. in Gard. Chron.* (1878) II. 654.—N. Granat.
Camaridii, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 547.—Costa Rica.
candida, *G. Lodd. ex Lindl. Bot. Reg.* (1841) *Misc.* 23.—Bras.
carinata, *Rodrig. Orch. Nov.* ii. 206.—Bras.
carinulata, *Reichb. f. in Linnaea*, xli. (1877) 6.—N. Granat.
cassapensis, *Reichb. f. in Walp. Ann.* vi. 539.—Peruv.
Cepula, *Reichb. f. in Bonplandia*, iii. (1855) 216, 226.—Bras.
cerifera, *Rodrig. Orch. Nov.* i. 118.—Bras.
chlorantha, *Lindl. Bot. Reg.* sub t. 1986.—Guian.
christobalensis, *Reichb. f. in Bot. Zeit.* x. (1852) 857.—Peruv.
chrysantha, *Rodrig. Orch. Nov.* i. 115.—Bras.
chrysoptera, Beer, Prakt. Orch. 264 = Lycaste chrysoptera.
ciliata, Ruiz & Pav. Syst. Veg. 220 = Lycaste Barringtoniae.
cochleata, Beer, Prakt. Orch. 264 = Lycaste cochleata.
coelia, Reichb. f. & Warsc. in Bonplandia, ii. (1854) 97 (Xylobii sp.).—N. Granat.
Colleyi, Batem. ex Lindl. Bot. Reg. (1838) Misc. 87 = Batemannia Colleyi.
colorata, *Reichb. f. in Walp. Ann.* vi. 523.—Peruv.
concava, Lindl. Bot. Reg. (1844) Misc. 4 = Xylobium concavum.
consanguinea, *Klotzsch, in Ann. Sc. Nat.* Sér. III. xix. (1853) 376.—Bras.
coriacea, *Rodrig. Orch. Nov.* ii. 199.—Bras.
corrugata, Lindl. Bot. Reg. (1844) Misc. 14 (Xylobii sp.).—Venezuela.
costata, Lindl. l. c. (1838) Misc. 93 = Lycaste costata.
crassifolia, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Bras.
crinita, Beer, Prakt. Orch. 264 = Lycaste crinita.
cristata, Lindl. Bot. Reg. t. 1811 = Lycaste cristata.
crocea, *Lindl. l. c.* t. 1749.—Bras.
crocea, Poepp. & Endl. Nov. Gen. et Sp. 37 = Poeppigiana.
cruenta, Lindl. Bot. Reg. (1842) t. 13 = Lycaste cruenta.
ctenostachys, *Reichb. f. in Gard. Chron.* (1870) 39.—Costa Rica.
cucullata, *Lindl. Bot. Reg.* (1840) t. 12.—Mexic.
cuneiformis, *Ruiz & Pav. Syst. Veg.* 223.—Peruv.
curtipes, *Hook. Ic. Pl.* t. 384.—Mexic.
cyanea, Beer, Prakt. Orch. 264 = Warrea cyanea.
cylindrobulba, *Regel, Gartenfl.* (1858) 341.—Mexic.
decolor, Lindl. Bot. Reg. t. 1549 (Xylobii sp.).—Jamaic.
deflexa, *Klotzsch, ex A. Br. Klotzsch & Bouché, Ind. Sem. Hort. Berol.* (1852) *App.* 1.—*Cf. Ann. Sc. Nat.* Sér. III. xix. (1853) 375.—Bras.
densa, Lindl. Bot. Reg. t. 1804 = Ornithidium densum.
densifolia, Reichb. f. in Bonplandia, ii. (1854) 16 = guareimensis.
Deppei, Lodd. Bot. Cab. t. 1612 = Lycaste Deppei.

MAXILLARIA:—
Desvauxiana, *Reichb. f. in Bonplandia*, iii. (1854) 67.—Guian.
discolor, *Reichb. in Walp. Ann.* vi. 529.—Guian.
divitiflora, *Reichb. f. Otia Bot. Hamb.* 7.—Ecuador.
eburnea, *Lindl. Sert. Orch.* t. 40. n. 2.—Bras.
echinophyta, *Rodrig. Orch. Nov.* i. 122.—Bras.
elatior, *Reichb. f. in Walp. Ann.* vi. 532.—Guatemala.
elongata, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 69. f. 264 = Xylobium elongatum.
exigua, Regel, Ind. Sem. Hort. Petrop. (1855) 20; et Gartenfl. (1856) 118 = Ornithidium vestitum.
Ferdinandiana, *Rodrig. Orch. Nov.* ii. 204.—Bras.
fimbriata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 38. t. 52. f. 2.—Peruv.
flagellifera, Lindl. ex Reichb. f. in Walp. Ann. vi. 551 = Scuticaria Steelii.
flavoviridis, *Rodrig. Orch. Nov.* i. 119.—Bras.
floribunda, *Lindl. Orch. Linden.* 19.—Peruv.
foveata, Lindl. Bot. Reg. (1839) Misc. 2 (Xylobii sp.).—Guian.
fractiflexa, *Reichb. f. Otia Bot. Hamb.* 8.—Ecuador.
Friedrichsthalii, *Reichb. f. in Bot. Zeit.* x. (1852) 858.—Guatemala.
fulvescens, Beer, Prakt. Orch. 265 = Lycaste fulvescens.
funerea, *Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 188.—Bras.
furfuracea, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xii. (1844) 67.—Bras.
fuscata, Klotzsch, ex Reichb. f. in Walp. Ann. Bot. vi. 520 = picta.
fuscata, Reichb. f. l. c. 525 = rufescens.
galeata, Lindl. in Lodd. Bot. Cab. t. 1645 = Gongora galeata.
galeata, *Scheidw. in Otto & Dietr. Allg. Gartenz.* x. (1842) 309.—Mexic.
Galeottiana, *Regel, Ind. Sem. Hort. Petrop.* (1856) 19.—*Cf. Ann. Sc. Nat.* Sér. IV. vi. (1856) 375.—Bras.
Galeottiana, *Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 24.—Mexic.
gigantea, Beer, Prakt. Orch. 265 = Lycaste gigantea.
glauca, *Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 586.—Guian.
glomerata, *Gal. Cact. et Orch. Brux.* 6.—Mexic.
glumibracteum, Hemsl. Biol. Centr. Am. Bot. iii. 266, sphalm. = Epidendrum glumibracteum.
Goeringii, *Reichb. f. in Bot. Zeit.* iii. (1845) 334.—Japon.
gracilis, *Lodd. Bot. Cab.* t. 1837.—Bras.
graminea, Lindl. Bot. Reg. xxi. (1836) sub t. 1802 (Aganosmatis sp.).—Guian.
graminifolia, *Reichb. f. in Walp. Ann.* vi. 538.—Paraguay.
grandiflora, *Lindl. Gen. et Sp. Orch.* 147.—N. Granat.; Peruv.
grandis, *Reichb. f. in Linnaea*, xli. (1877) 29.—Ecuador.
guareimensis, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Venezuela.
guttata, Hort. ex Reichb. f. in Bot. Zeit. x. (1852) 672 = Zygopetalum xanthinum.
Harrisoniae, Lindl. in Bot. Reg. t. 897 = Bifrenaria Harrisoniae.
hastata, *Ruiz & Pav. Syst. Veg.* 222.—Peruv.
hastulata, *Lindl. in Benth. Pl. Hartw.* 155.—Ecuador.
Heismanniana, *Rodrig. Orch. Nov.* ii. 201 (*Cf. Index* p. 8).—Bras.
hematoglossa, Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 24 = rhombea.
Henchmanni, Hook. Bot. Mag. t. 3614 = variabilis.
heraldica, *Reichb. f. in Bot. Zeit.* x. (1852) 763.—Guatemala.
heteroclita, *Poepp. & Endl. Nov. Gen. et Sp.* 37. t. 63.—Peruv.
Heynderycxii, E. Morr. in Ann. Soc. Bot. Gand, i. (1845) 97. t. 9 = Lycaste gigantea.
hirtilabia, *Lindl. in Journ. Hort. Soc.* viii. (1853) 132.—N. Granat.
histiologorum, *Reichb. f. in Bot. Zeit.* x. (1852) 936.—Bras.
Houtteana, *Reichb. f. in Hamb. Gartenz.* xiv. (1858) 212; *et in Regel, Gartenfl.* vii. (1858) 286.—Guatemala; Venezuela.
Houtteii, *Beer, Prakt. Orch.* 265.—Hab.?
hyacinthina, Reichb. f. in Linnaea, xxii. (1849) 855 (Xylobii sp.).—Venezuela.
hypocrita, Reichb. f. in Hamb. Gartenz. xvi. (1860) 15 (Xylobii sp.).—Hab.?

MAXILLARIA:—
imbricata, *Rodrig. Orch. Nov.* i. 120.—Bras.
inaudita, *Reichb. f. Beitr. Orch. Centr. Am.* 76.—Costa Rica.
infausta, *Reichb. f. in Bonplandia*, iii. (1855) 216.—Peruv.
inflexa, Griseb. Fl. Brit. W. Ind. 626 = Ornithidium inflexum.
iridifolia, *Reichb. f. in Bonplandia*, ii. (1854) 16.—N. Granat.
irrorata, *Reichb. f. in Gard. Chron.* (1883) II. 102.—Reg. Andin.
jugosa, Lindl. Bot. Reg. (1841) Misc. 51 = Lycaste jugosa.
Kalbreyeri, *Reichb. f. in Gard. Chron.* (1885) I. 239.—N. Granat.
Kegelii, *Reichb. f. in Linnaea*, xli. (1877) 127.—Guian.
Kreysigii, Hoffmgg. ex Regel, Gartenfl. v. (1856) 328 = picta.
Kreysigii, *Reichb. f. ex Beer, Prakt. Orch.* 265.—Hab.?
laevilabris, *Lindl. in Benth. Pl. Hartw.* 155.—Peruv.
lancifolia, *Reichb. f. in Walp. Ann.* vi. 539.—Venezuela.
latifolia, *Lindl. Gen. et Sp. Orch.* 149.—Reg. Andin.
Lehmanni, *Reichb. f. Otia Bot. Hamb.* 8.—Ecuador.
leiantha, Beer, Prakt. Orch. 265 = Lycaste Deppei?
lentiginosa, Lindl. Bot. Reg. (1839) Misc. 16 = Zygopetalum lentiginosum.
leontoglossa, Reichb. f. in Bonplandia, iii. (1855) 67 (Xylobii sp.).—N. Granat.
lepidota, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 383.—N. Granat.
leptosepala, Hook. Bot. Mag. t. 4434 = setigera.
leucaimata, *Rodrig. Orch. Nov.* ii. 198.—Bras.
leucantha, Klotzsch, in Otto & Dietr. Allg. Gartenz. xviii. (1850) 402 = Lycaste leucantha.
leucocheile, Hoffmgg. Verz. Orch.—Cf. Bot. Zeit. i. (1843) 833 = picta.
ligulata, *Ruiz & Pav. Syst. Veg.* 222.—Peruv.
liliacea, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 12 = Govenia liliacea.
liliacea, *Rodrig. Orch. Nov.* ii. 201.—Bras.
Lindeniana, *Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 24.—Mexic.
linguaeformis, *Regel, in Ind. Sem. Hort. Petrop.* (1856) 20; *et Gartenfl.* (1856) 328.—Hab.?
longibracteata, Reichb. f. in Walp. Ann. vi. 540 = Camaridium longibracteatum.
longifolia, Lindl. Gen. et Sp. Orch. 150 = Xylobium longifolium.
longipes, *Lindl. in Benth. Pl. Hartw.* 154.—Peruv.; Ecuador.
longipetala, *Rodrig. Orch. Nov.* i. 120.—Bras.
longipetala, *Ruiz & Pav. Syst. Veg.* 220.—Peruv.
longissima, *Lindl. Orch. Linden.* 20.—Venezuela.
lorifolia, *Reichb. f. in Bonplandia*, ii. (1854) 92.—Venezuela.
luteo-alba, *Lindl. Orch. Linden.* 20.—N. Granat.
luteo-grandiflora, *Hort. ex Flor. Mag.* x. (1871) t. 599.—Hab.?
luteola, Beer, Prakt. Orch. 266 = Polystachya luteola.
luteo-rubra, *Reichb. f. in Walp. Ann.* vi. 539.—Peruv.; N. Granat.
lutescens, Scheidw. in Otto & Dietr. Allg. Gartenz. vii. (1839) 145 = Camaridium lutescens.
lyncea, Lindl. Gen. et Sp. Orch. 151 = Stanhopea tigrina.
Lyonii, *Lindl. Bot. Reg.* (1845) *Misc.* 17.—Mexic.
Macleei, *Batem. ex Lindl. l. c.* (1840) *Misc.* 70.—Guatemala.
macrobulbon, Hook. Bot. Mag. t. 4228 = Lycaste macrobulbon.
macrophylla, Poepp. & Endl. Nov. Gen. et Sp. i. 37. t. 64 = Lycaste macrophylla.
macrura, *Reichb. f. in Linnaea*, xli. (1877) 28.—Venezuela.
maculata, *Lindl. Gen. et Sp. Orch.* 149.—Ecuador.
madida, Lindl. Bot. Reg. (1838) Misc. 44 = H. Cepula.
marginata, *Fenzl, in Fl. des Serres*, Sér. I. x. (1854–55) 112; *Reichb. f. in Bonplandia*, iii. (1855) 216.—Bras.
Mathewsii, *Lindl. in Benth. Pl. Hartw.* 155.—Peruv.
Matthewsii, *Reichb. f. in Walp. Ann.* vi. 539.—Peruv.
Meirax, *Reichb. f. & Warm. in Otia Bot. Hamb.* 88.—Venezuela.
Meleagris, *Lindl. Bot. Reg.* (1844) *Misc.* 3.—Venezuel.

MAXILLARIA:—
melina, *Lindl. Orch. Linden.* 20.—Venezuela.
meridensis, *Lindl. l. c.* 19.—Venezuela.
miliacea, *Reichb. f. Xenia Orch.* iii. 22.—Bolivia.
monoceros, Klotzsch, ex Reichb. f. in Walp. Ann. vi. 520 = picta.
multiflora, *Rodrig. Orch. Nov.* i. 117.—Bras.
muscicola, *Reichb. f. Otia Bot. Hamb.* 7.—Ecuador.
mutabilis, Hort. ex Reichb. f. in Bonplandia, iii. (1855) 216 = porrecta.
nana, Hook. Ic. Pl. t. 315 = uncata.
nanegalensis, *Reichb. f. in Linnaea*, xli. (1877) 34.—Ecuador.
nasuta, *Reichb. f. Beitr. Orch. Centr. Am.* 104.—Costa Rica.
neophylla, *Reichb. f. in Gard. Chron.* (1878) ii. 588.—N. Granat.
Neowiedii, *Reichb. f. in Linnaea*, xli. (1877) 29.—Bras.
nigrescens, *Lindl. Orch. Linden.* 20.—Venezuela.
nitidula, *Reichb. f. in Linnaea*, xli. (1877) 29.—Ecuador.
notylioglossa, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Venezuela.
nutans, *Lindl. in Benth. Pl. Hartw.* 154.—Peruv.
obscura, Linden & Reichb. f. Beitr. Orch. Centr. Am. 31 = cucullata.
ochroleuca, *Lodd. ex Lindl. Gen. et Sp. Orch.* 143.—Bras.
ornata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 257.—Peruv.
ornithoglossa, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 586.—Mexic.
pallidiflora, Hook. Bot. Mag. t. 2806 (Xylobii sp.).—Ins. S. Vincent.
palmifolia, Lindl. Gen. et Sp. Orch. 148 (Xylobii sp.).—Jamaica.
pamplonensis, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 281.—N. Granat.
paniculata, *Ruiz & Pav. Syst. Veg.* 223.—Peruv.
pantherina, *Reichb. f. in Bonplandia*, iii. (1855) 237.—Reg. Andin.
paranaënsis, *Rodrig. Orch. Nov.* ii. 205.—Bras.
Parkerii, *Hook. Bot. Mag.* t. 2729.—Guian.
Parthoni, *Dum. in Hortic. Belge*, ii. (1834) 245.—Bras.
parvula, Hook. Exot. Fl. t. 217 = Bifrenaria parvula.
pastensis, *Reichb. f. in Bonplandia*, iii. (1855) 239.—N. Granat.
pauciflora, *Rodrig. Orch. Nov.* i. 116.—Bras.
penduliflora, Fenzl, in Fl. des Serres, Sér. I. x. (1854–55) 110 = gracilis ?
pentura, *Lindl. Orch. Linden.* 20.—Venezuela.
pertusa, Reichb. f. in Gard. Chron. (1878) i. 169 = lepidota.
phoenicanthera, *Rodrig. Orch. Nov.* ii. 200.—Bras.
phyllomega, Steud. Nom. ed. II. ii. 107 = Lycaste macrophylla.
picta, *Hook. Bot. Mag.* t. 3154.—Bras.
placanthera, Hook. l. c. t. 3173 = *viridis*.
platypetala, *Ruiz & Pav. Syst. Veg.* 221.—Peruv.
plebeja, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 57.—Bras.
Poeppigiana, *Steud. Nom.* ed. II. ii. 107.—Peruv.
polyantha, *Rodrig. Orch. Nov.* ii. 202.—Bras.
polypetala, Steud. Nom. ed. I. 261; ed. II. i. 490, in syn. = platypetala.
polyphylla, *Reichb. f. in Walp. Ann.* vi. 539.—Ecuador.
ponerantha, *Reichb. f. in Bonplandia*, ii. (1854) 17.—Venezuela.
porphyrostele, *Reichb. f. in Gard. Chron.* (1878) i. 978.—Bras.
porrecta, *Lindl. Bot. Reg.* (1838) *Misc.* 192.—Guian.
praestans, *Reichb. f. in Gard. Chron.* (1885) i. 566.—Guatemala.
praetexta, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Venezuela.
proboscidea, *Reichb. f. l. c.*—Venezuela.
procurrens, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 383.—N. Granat.
prolifera, *Ruiz & Pav. Syst. Veg.* 226.—Peruv.
psittacina, *Batem. ex Steud. Nom.* ed. II. ii. 107 (Quid ?).—Mexic.
pterocarpa, *Rodrig. Orch. Nov.* ii. 204.—Bras.
pubigera, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 106 = Bifrenaria Harrisoniae.
pulla, *Linden & Reichb. f. in Bonplandia*, ii. (1845) 280.—N. Granat.

MAXILLARIA:—
pumila, *Hook. Bot. Mag.* t. 3613.—Guian.
punctata, *Lodd. Bot. Cab.* t. 1914.—Bras.
puncto-striata, *Reichb. f. in Linnaea*, xli. (1877) 28.—Mexic.
punctulata, Klotzsch, in Otto & Dietr. Allg. Gartenz. xix. (1851) 249 = marginata.
purpurascens, Knowl. & Westc. in Phytolog. 8; ex Reichb. f. in Walp. Ann. vi. 547 = Bifrenaria parvula.
purpurata, *Reichb. f. in Walp. Ann.* vi. 538.—Venezuela.
purpurea, Beer, Prakt. Orch. 266 = Polystachya purpurea.
Queirogana, *Rodrig. Orch. Nov.* i. 116.—Bras.
racemosa, Hook. Bot. Mag. t. 2789 = Bifrenaria racemosa.
ramosa, *Ruiz & Pav. Syst. Veg.* 226.—Peruv.
rebellis, Reichb. f. in Fl. des Serres, Sér. I. ix. (1853–54) 102; et in Bonplandia, iii. (1855) 217 (Xylobii sp.).—Hab. ?
Reichenheimiana, *Reichb. f. in Gard. Chron.* (1871) 16, 78.—Costa Rica.
revoluta, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 186 = variabilis.
rhombea, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 12.—Mexic.
rigida, *Rodrig. Orch. Nov.* ii. 206.—Bras.
ringens, *Reichb. f. in Walp. Ann.* vi. 523.—Mexic.
robusta, *Rodrig. Orch. Nov.* ii. 197.—Bras.
Roezli, *Reichb. f. ex Linden, Illustr. Hortic.* xxix. (1882) 68, nomen.—N. Granat.
Rollinsonii, Steud. Nom. ed. II. ii. 108 = seq.
Rollissonii, Lindl. Bot. Reg. sub t. 1986 = Zygopetalum Rollissonii.
rubro-fusca, Klotzsch, Ind. Sem. Hort. Berol. (1853) 12 = nigrescens.
rufescens, *Lindl. Bot. Reg.* (1836) t. 1848.—Ins. Trinit.
rugosa, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xi. (1843) 101.—Bras.
rupestris, *Rodrig. Orch. Nov.* ii. 199.—Bras.
saxatilis, *Reichb. f. in Bonplandia*, iii. (1855) 216.—Peruv.
scabrilinguis, Lindl. Bot. Reg. (1844), Misc. 71 (Xylobii sp.).—Ecuador.
serotina, *Regnell & Rodrig. in Rodrig. Orch. Nov.* ii. 203.—Bras.
sessilis, *Lindl. in Benth. Pl. Hartw.* 155.—Ecuador.
setigera, *Lindl. in Bot. Reg.* (1845), *Misc.* 80.—Venezuela.
sinuosa, *Batem. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 587.—Guian.
Skinneri, Batem. ex Lindl. in Bot. Reg. (1840) Misc. 48; et (1842) Misc. 10 = Lycaste Skinneri.
spathacea, Lindl. Gen. & Sp. Orch. 151 = Bifrenaria Harrisoniae.
speciosa, *Reichb. f. in Gard. Chron.* (1876) ii. 197.—N. Granat.
spilotantha, *Reichb. f. in Bonplandia*, ii. (1854) 17.—Venezuela.
splendens, *Poepp. & Endl. Nov. Gen. et Sp.* i. 38. t. 66.—Peruv.
squalens, Hook. Bot. Mag. t. 2955 = Xylobium squalens.
squamata, *Rodrig. Orch. Nov.* i. 118.—Bras.
stachyobiorum, Reichb. f. in Bot. Zeit. x. (1852) 735 = Xylobium stachyobiorum.
stapelioides, Lindl. Gen. & Sp. Orch. 146 = Zygopetalum stapelioides.
Steelii, Hook. Bot. Mag. t. 3573 = Scuticaria Steelii.
stenobulbum, Klotzsch, Ind. Sem. Hort. Berol. (1853) 11 (Xylobii sp.).—Venezuela.
stenopetala, Knowl. & Westc. Flor. Cab. ii. (1838) 112 (Bifrenariae sp.).—Bras.
stenophylla, *Reichb. f. in Bonplandia*, ii. (1854) 17.—Venezuela.
subulata, *Lindl. Gen. & Sp. Orch.* 147.—Bras.
sulfurina, Lem. in Fl. des Serres, Sér. I. iv. (1848) t. 330 b (Xylobii sp.).—Guatemala.
superba, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 13 = Govenia superba.
superflua, *Reichb. f. in Bonplandia*, iv. (1856) 323.—Hab. ?
supina, Poepp. & Endl. Nov. Gen. et Sp. i. 39. t. 67 (Xylobii sp.).—Peruv.
surinamensis, Focke, ex Reichb. f. in Walp. Ann. vi. 492 (Ornithidii sp.).—Peruv.
tenuifolia, *Lindl. Bot. Reg.* sub t. 1986.—Mexic.
tetragona, Lindl. l. c. t. 1428 = Lycaste tetragona.

MAXILLARIA:—
triangularis, Lindl. Bot. Reg. (1845) Misc. 9 = elatior.
tricolor, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 186 = Lycaste tricolor.
tricolor, Lindl. in Hook. Lond. Journ. Bot. i. (1842) 189 = marginata.
tricolor, *Ruiz & Pav. Syst. Veg.* 224.—Peruv.
triloris, *E. Morr. in Belg. Hortic.* xx. (1870) 240. t. 13.—Venezuela ?
triphylla, *Ruiz & Pav. Syst. Veg.* 225.—Peruv.
truxillensis, Reichb. f. in Bonplandia, ii. (1854) 17 (Xylobii sp.).—Mexic.
Turneri, *Nichols. Dict. Gard.* i. 339.—Hab. ?
tyrianthina, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 582 = Bifrenaria tyrianthina.
uncata, *Lindl. Bot. Reg.* sub t. 1986.—Guiana.
undatiflora, *Ruiz & Pav. Syst. Veg.* 225.—Peruv.
undulata, *Ruiz & Pav. l. c.* 221.—Peruv.
vaginalis, *Reichb. f. Beitr. Orch. Centr. Am.* 77.—Costa Rica.
variabilis, *Batem. ex Lindl. Bot. Reg.* sub t. 1986.—Mexic.
varicosa, *Reichb. f. in Gard. Chron.* (1883) ii. 392.—Bolivia.
variegata, *Ruiz & Pav. Syst. Veg.* 222.—Peruv.
venusta, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 280.—N. Granat.
vernicosa, *Rodrig. Orch. Nov.* i. 121.—Bras.
violaceopunctata, *Reichb. f. in Bonplandia*, iii. (1855) 216.—Guian.
virginalis, *Scheidw. in Bull. Acad. Brux.* ix. 1. (1842) 25.—Mexic.
virguncula, *Reichb. f. in Bonplandia*, ii. (1854) 16.—Venezuela.
viridis, Lindl. Bot. Reg. t. 1510 (an Lycastes sp. ?).—Bras.
vitellina, Lindl. l. c. (1838) Misc. 62 (Bifrenariae sp.).—Bras.
vitelliniflora, *Rodrig. Orch. Nov.* i. 121.—Bras.
Wageneri, Reichb. f. in Regel, Gartenfl. (1856) 118 = Bifrenaria Wageneri.
Warreana, Lodd. ex Lindl. Gen. & Sp. Orch. 148 (Warreae sp.?).—Bras.
xanthina, Lindl. Bot. Reg. (1839) sub t. 17 = Zygopetalum xanthinum.

MAXIMILIANA, Mart. Hist. Nat. Palm. ii. 131 (1824). *PALMAE*, Benth. & Hook. f. iii. 946.
attaleoides, Barb. Rodr. Enum. Palm. Nov. 41 = Attalea spectabilis.
caribaea, *Griseb. & H. Wendl. ex Griseb. Fl. Brit. W. Ind.* 522.—Ind. occ.
crassispatha, *Mart. Palmet. Orbign.* 110.—Ins. S. Doming.
elegans, Karst. in Linnaea, xxviii. (1856) 271 = M. Maripa.
Inajai, Spruce, in Journ. Linn. Soc. xi. (1869) 163 = Cocos Inajai.
insignis, *Mart. Hist. Nat. Palm.* ii. 133. t. 94.—Bras.
Jagua, *Seem. ex Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
Maripa, *Drude, in Mart. Fl. Bras.* iii. ii. 452.—Guian.
Martiana, Karst. in Linnaea, xxviii. (1856) 273 = regia.
panamensis, *Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
Princeps, Mart. Palmet. Orbign. t. 4 = Scheelea Princeps.
regia, *Mart. Hist. Nat. Palm.* ii. 132. tt. 91–93.—Bras.
regia, Wallace, Palm-trees Amaz. 121. t. 47 = M. Maripa.
spiralis, *Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—Bras.
tetrasticha, *Drude, in Mart. Fl. Bras.* iii. ii. 455.—Bras.
venatorum, *H. Wendl. in Kerch. Palm.* 251.—Peruv. ?

MAXIMILIANA, Mart. in Flora, ii. (1819) 451 = **Cochlospermum**, Kunth (Bixin.).
regia, Mart. l. c. 452 = C. insigne.

MAXIMOVITZIA, Benth. & Hook. f. Gen. i. 19 (1862) = MAXIMOWICZIA, Rupr. = **Schisandra**, Michx.

MAXIMOWICZIA, Cogn. in DC. Monog. Phan. iii. 726 (1881). *CUCURBITACEAE*.
Lindheimeri, *Cogn. l. c.* 727.—Texas.
tripartita, *Cogn. l. c.* 728.—Texas.

MAXIMOWICZIA, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 142, 124 = **Schisandra**, Michx. (Magnol.).
amurensis, Rupr. l. c. 124 = S. chinensis.
chinensis, Rupr. in Maxim. Prim. Fl. Amur. 31 = S. chinensis.
japonica, C. Koch, Dendrol. i. 386 = S. chinensis.

MAXWELLIA, Baill. Adansonia, x. (1871) 100. *STERCULIACEAE.*
lepidota, *Baill. l. c.*—N. Caled.

MAYACA, Aubl. Pl. Gui. i. 42. t. 15 (1775). *MAYACEAE*, Benth. & Hook. f. iii. 843.
Biaslia, Vand. Fl. Lusit. et Bras. Spec. 4. t. 1. f. 2 (1788).
Coletia, Vell. Fl. Flum. 32 (1825).
Syena, Schreb. Gen. 39 (1789).
Aubletii, *Michx. Fl. Bor. Am.* i. 26.—Am. bor.
Aubletii, Schott & Endl. Melet. 24. t. 3 = fluviatilis.
Endlicheri, *Poepp. ex Seub. in Mart. Fl. Bras.* iii. 1. 230.—Peruv.
fluviatilis, *Aubl. Pl. Gui.* i. 42.—Guian.
fluviatilis, Ell. Sketch, i. 50 = Aubletii.
Kunthii, *Seub. in Mart. Fl. Bras.* iii. 1. 228.—Bras.
lagoënsis, *Warm. in Kjoeb. Vidensk. Meddel.* (1867) 42.—Bras.
longipes, *Mart. ex Seub. in Mart. Fl. Bras.* iii. 1. 229.—Bras.
Michauxii, Schott & Endl. Melet. 24 = Aubletii.
Sellowiana, *Kunth, Enum. Pl.* iv. 32.—Bras.
Vandellii, Kunth, l. c. = Kunthii.
Vandellii, *Schott & Endl. Melet.* 24. t. 3.—Bras.
Wrightii, *Griseb. Cat. Pl. Cub.* 224.—Cuba.

MAYANTHEMUM, DC. Fl. Fr. iii. 107 (1805) = **Maianthemum**, Weber (Liliac.).

MAYCOCKIA, A. DC. in DC. Prod. viii. 324 (1844) = **Condylocarpon**, Desf. (Apocyn.).
Rauwolfiae, A. DC. l. c. = C. Rauwolfiae.

MAYEPEA, Aubl. Pl. Gui. i. 81. t. 31 (1775) = **Linociera**, Sw. (Oleac.).
axillaris, F. Muell. Census, 92 = L. albidiflora?
guianensis, Aubl. Pl. Gui. i. 81. t. 31 = L. tetrandra.
picrophloia, F. Muell. Census, 92 = L. intermedia.
quadristaminea, F. Muell. Fragm. x. 89.—Austral.
ramiflora, F. Muell. Census, 92 = L. macrophylla.

MAYETA, Juss. Gen. 330 (1789) = **Maieta**, Aubl. (Melastom.).

MAYNA, Aubl. Pl. Gui. ii. 921. t. 352 (1775) = **Oncoba**, Forsk. (Bixin.)
brasiliensis, Raddi, in Mem. Mod. xviii. Fis. (1820) 402.—Bras.
denticulata, Benth. in Hook. Journ. Bot. iv. (1842) 115 = Dendrostylis denticulata.
echinata, Spruce, ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 82.—Peruv.
latifolia, Benth. l. c. 80 = O. latifolia.
laurina, Benth. l. c. 81.—Mexic.
laxiflora, Benth. in Hook. Journ. Bot. iv. (1842) 114 = O. maynensis.
longifolia, Poepp. & Endl. Nov. Gen. et Sp. iii. 64. t. 271 = Carpotroche longifolia.
odorata, Aubl. Pl. Gui. ii. 921. t. 352 = Dendrostylis suaveolens?
ovata, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 81 = O. ovata.
paludosa, Benth. in Hook. Journ. Bot. iv. (1842) 114 = O. maynensis.
pauciflora, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 81 = O. pauciflora.
sericea, Spreng. Neue Entdeck. ii. 153.—Bras.
vernicosa, Karst. Fl. Columb. ii. 11. t. 106.—N. Granat.

MAYODENDRON, Kurz, For. Fl. Brit. Burma, ii. 232 (1877). *BIGNONIACEAE.*
igneum, *Kurz, l. c.* 233.—Burma.

MAYS, Tourn. ex Gaertn. Fruct. i. 6. t. 1 (1788) = **Zea**, Linn. (Gramin.).
americana, Baumg. Enum. Stirp. Transs. iii. 281 = Z. Mays.
vulgaris, Ser. Mél. Bot. i. 182 = Z. Mays.
Zea, Gaertn. Fruct. i. 6. t. 1. f. 9 = Z. Mays.

MAYTENUS, Molina, Sagg. Chile, ed. I. 177 (1782); *CELASTRINEAE*, Benth. & Hook. f. i. 364.
Boaria, DC. Prod. viii. 299 (1844).
Euthalis, Banks & Soland. ex Hook. f. Fl. Antarct. 254 (1847).
Haenkea, Ruiz & Pav. Prod. 36. t. 6 (1794).
Monteverdia, A. Rich. Ess. Fl. Cub. 346 (1845).
Tricerma, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 97.
acanthophylla, *Reiss. in Mart. Fl. Bras.* xi. 1. 6.—Bras.
acerifolia, Walp. Rep. ii. 827, sphalm. = aquifolium.
acutangula, *Reiss. in Mart. Fl. Bras.* xi. 1. 4.—Bras.
alaternoides, *Reiss. l. c.* 15.—Bras.
amazonica, *Mart. ex Reiss. l. c.* 25.—Bras.
amygdalina, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 451.—Guian.
andina, Phil. in Linnaea, xxviii. (1856) 617 = magellanica.
Aquifolium, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 91.—Bras.
ardisiaefolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 26.—Bras.
attenuata, *Reiss. l. c.* 23.—Bras.
basidentata, *Reiss. l. c.* 7.—Bras.
Boaria, *Molina, Sagg. Chile*, ed. I. 177.—Chili.
brasiliensis, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 87.—Bras.
buxifolia, *Griseb. Cat. Pl. Cub.* 53.—Cuba.
buxifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 367.—N. Granat.
capitellata, *Presl, Bot. Bemerk.* 35.—Peruv.
cassineformis, *Reiss. in Mart. Fl. Bras.* xi. 1. 15.—Bras.
castaneaeformis, *Reiss. l. c.* 7.—Bras.
catingarum, *Reiss. l. c.* 26.—Bras.
cestrifolia, *Reiss. l. c.* 24.—Bras.
chilensis, DC. Prod. ii. 9 = M. Boaria.
cochlearifolia, *Griseb. Cat. Pl. Cub.* 53.—Cuba.
communis, *Reiss. in Mart. Fl. Bras.* xi. 1. 22.—Bras.
commutata, *Reiss. l. c.* 21.—Bras.
comocladiaeformis, *Reiss. l. c.* 8.—Bras.
crenulata, *Presl, Bot. Bemerk.* 35.—Chili.
cuneifolia, *Griseb. Cat. Pl. Cub.* 64.—Cuba.
dasyclados, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 89.—Bras.
diospyroides, *Reiss. in Mart. Fl. Bras.* xi. 1. 17.—Bras.
distichophylla, *Mart. ex Reiss. l. c.*—Bras.
ebenifolia, *Reiss. l. c.* 18.—Bras.
elaeodendroides, *Griseb. Cat. Pl. Cub.* 54.—Cuba.
erythroxylon, *Reiss. in Mart. Fl. Bras.* xi. 1. 21.—Bras.
evonymoides, *Reiss. l. c.* 11.—Bras.
ficiformis, *Reiss. l. c.* 20.—Guian.
floribunda, *Reiss. l. c.* 16.—Bras.
glaucescens, *Reiss. l. c.* 10.—Bras.
gonoclada, Griseb. Fl. Brit. W. Ind. 145 = tetragona.
gonoclada, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 89.—Bras.
grandiflora, *Reiss. in Mart. Fl. Bras.* xi. 1. 23.—Bras.
guianensis, *Klotzsch, in Schomb. Fauna & Fl. Brit. Gui.* 1097.—Guian.
horrida, *Reiss. in Mart. Fl. Bras.* vi. 1. 5.—Bras.
ilicifolia, *Mart. ex Reiss. l. c.* 8.—Bras.
imbricata, *Mart. ex Reiss. l. c.* 16.—Bras.
impressa, *Reiss. l. c.* 29.—Bras.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxxii. (1882) 38.—Bras.
Karstenii, *Reiss. in Mart. Fl. Bras.* xi. 1. 19.—Venezuela.
laevis, *Reiss. l. c.* 27.—Bras.
lagoënsis, *Warm. in Kjoeb. Vidensk. Meddel.* (1880) 365.—Bras.
laxiflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 367.—N. Granat.
ligustrina, *Reiss. in Mart. Fl. Bras.* xi. 1. 25.—Bras.
lineatus, *Wright, ex Griseb. Cat. Pl. Cub.* 54.—Cuba.
macrodonta, *Reiss. in Mart. Fl. Bras.* xi. 1. 6.—Bras.
macrophylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 95.—Bras.
magellanica, *Hook. f. Fl. Antarct.* 254.—Reg. Magell.
multiflora, *Reiss. in Mart. Fl. Bras.* xi. 1. 24.—Bras.
myricoides, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 366.—N. Granat.
myrsinoides, *Reiss. in Mart. Fl. Bras.* xi. 1. 21.—Guian.

MAYTENUS:—
nitida, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 91.—Bras.
oblongata, *Reiss. in Mart. Fl. Bras.* xi. 1. 25.—Guian.
obovata, *Hook. f. in Trans. Linn. Soc.* xx. (1837) 230.—Ins. Galap.
obtusifolia, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 88.—Bras.
octogona, *DC. Prod.* ii. 9.—Peruv.
opaca, *Reiss. in Mart. Fl. Bras.* xi. 1. 28.—Bras.
oxyodonta, *Reiss. l. c.* 5.—Bras.
patens, *Reiss. l. c.* 13.—Bras.
pendulina, Steud. Nom. ed. II. ii. 108 = M. Boaria.
phyllanthoides, *Benth. Bot. Voy. Sulph.* 54.—Mexic.
pruinosa, *Reiss. in Mart. Fl. Bras.* xi. 1. 20.—Guian.
prunifolia, *Presl, Bot. Bemerk.* 35.—Peruv.
Pseudocasearia, *Reiss. in Mart. Fl. Bras.* xi. 1. 11.—Bras.
pulviniflora, *Reiss. l. c.* 14.—Bras.
Pyraster, *Reiss. l. c.* 13.—Bras.
racemosa, *Reiss. l. c.* 30.—Bras.
repanda, *Reiss. l. c.* 29.—Bras.
repanda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 450.—Mexic.
rhamnoides, *Reiss. in Mart. Fl. Bras.* xi. 1. 13.—Bras.
Riedeliana, *Regel, Ind. Sem. Hort. Petrop.* (1856) 36.—Bras.
rigida, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 90.—Bras.
robusta, *Reiss. in Mart. Fl. Bras.* xi. 1. 15.—Bras.
salicifolia, *Reiss. l. c.* 10.—Bras.
samydaeformis, *Reiss. l. c.* 11.—Bras.
sapotiformis, *Reiss. l. c.* 27.—Guian.
subalata, *Reiss. l. c.* 12.—Bras.
tetragona, *Griseb. Fl. Brit. W. Ind.* 709.—Ins. Trinit.; Guian.
tovarensis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 383.—Venezuela.
trichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 451.—Mexic.
truncata, *Reiss. in Mart. Fl. Bras.* xi. 1. 5.—Bras.
uliginosa, *H. B. & K. Nov. Gen. et Sp.* vii. 65.—Ecuador.
uncinata, *G. Don, Gen. Syst.* ii. 11.—Chili.
verticillata, *DC. Prod.* ii. 10.—Peruv.
vincentina, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 600.—Ins. S. Vincent.
viscifolia, *Griseb. in Goett. Abh.* xix. (1874) 110.—Reg. Argent.
Vitis-idaea, *Griseb. l. c.*—Peruv.

MAYZEA, Rafin. Med. Fl. ii. 241 (1830) = **Zea**, Linn. (Gramin.).
cerealis, Rafin. l. c. = Z. Mays.
vestita, Rafin. l. c. = praec.?

MAZEUTOXERON, Labill. Voy. ii. 11 (1798) = **Correa**, Andr. (Rutac.).
reflexum, Labill. l. c. 66. t. 19 = C. speciosa.
rufum, Labill. l. c. 12. t. 17 = C. alba.

MAZINNA, Spach, Hist. Vég. Phan. ii. 487 (1834) = MOZINNA, Ort. = **Jatropha**, Linn. (Euphorb.).

MAZUS, Lour. Fl. Cochinch. 385 (1790). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 947.
Hornemannia, Willd. Enum. Hort. Berol. 653 (1809).
bicolor, Benth. in Wall. Cat. n. 3913 = rugosus.
caducifer, *Hance, in Journ. Bot.* xx. (1882) 292.—China.
dentatus, *Wall. Cat.* n. 3914.—Reg. Himal.
laevifolius, *Blume, Bijdr.* 753.—Java.
pinnatus, Wall. Cat. n. 3915 = Sibthorpia pinnata.
Pumilio, *R. Br. Prod.* 439.—Austral.; N. Zel.
rugosus, *Lour. Fl. Cochinch.* 385.—Reg. Himal.; Malaya; China.
stachydifolius, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 438.—China.
surculosus, *D. Don, Prod. Fl. Nep.* 86.—Reg. Himal.
vandellioides, Hance, in Walp. Ann. iii. 198 = rugosus.
villosus, Hemsl. in Journ. Bot. xiv. (1876) 209 = stachydifolius.

MEADIA, Catesby, ex Mill. Gard. Dict. ed. VI. App. 175 (1752) = **Dodecatheon**, Linn. (Primul.).
dodecathea, Crantz, Inst. ii. 309 = seq.
Dodecatheon, Mill. Gard. Dict. ed. VIII. = D. Meadia.

MEBORA, Steud. Nom. ed. II. ii. 108 (1841) = seq.

MEBOREA, Aubl. Pl. Gui. ii. 825. t. 323 (1775) = **Phyllanthus**, Linn. (Euphorb.).
guianensis, Aubl. l. c. = P. attenuatus.

MECARDONIA, Ruiz & Pav. Prod. 95 (1794) = **Herpestis**, Gaertn. (Scrophul.).
ovata, Ruiz & Pav. Syst. Veg. 164.—Peruv.
pusilla, Mart. Nov. Gen. et Sp. iii. 16. t. 28. = H. serpylloides.

MECKELIA, Mart. ex A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 326, in syn. = **Spachea**, A. Juss. (Malpig.).
multiflora, Mart. ex A. Juss. l. c. = S. tricarpa.

MECLATIS, Spach, Hist. Vég. Phan. vii. 272 (1839) = **Clematis**, Tourn. (Ranunc.).
orientalis, Spach, l. c. 274 = C. orientalis.
sibirica, Spach, l. c. 273 = C. Flammula.

MECOMISCHUS, Coss. ex Benth. & Hook. f. Gen. ii. 418 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c. FRADINIA, Pomel, Nouv. Mat. Fl. Atl. 51 (1874).
Geslini, *Coss. ex Benth. & Hook. f. Gen.* ii. 418.—Algeria.
pedunculatus, Coss. ex Benth. & Hook. f. l. c. 419 = Anthemis peduncularis.

MECONELLA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 64 (1838) = **Platystigma**, Benth. (Papav.).
californica, Torr. in Frém. Rep. 312 = P. californicum.
oregana, Nutt. ex Torr. & Gray, Fl. N. Am. i. 64 = P. oreganum.

MECONOPSIS, Vig. Diss. 20, 48 (1814); ex DC. Syst. ii. 86. *PAPAVERACEAE*, Benth & Hook. f. i. 52.
aculeata, *Royle, Illustr. Bot. Himal.* 67.—Reg. Himal.
cambrica, *Vig. Diss.* 48. f. 3.—Europ.
crassifolia, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 408 = heterophylla.
diphylla, DC. Syst. ii. 88 = Stylophorum diphyllum.
Guilelmi-Waldemarii, Klotzsch, in Bot. Ergeb. Waldem. Reise, 129 = aculeata.
heterophylla, *Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 408.—Calif.
horridula, *Hook f. & Thoms. Fl. Ind.* i. 252.—Reg. Himal.
nepalensis, *DC. Prod.* i. 121.—Reg. Himal.
petiolata, DC. Syst. ii. 87 = Stylophorum diphyllum.
quintuplinervia, *Regel, Gartenfl.* (1876) 291.—Mandshuria.
racemosa, *Maxim. in Bull. Acad. Pétersb.* xxviii. (1877) 310.—China.
robusta, *Hook. f. & Thoms. Fl. Ind.* i. 253.—Reg. Himal.
simplicifolia, *G. Don, Gen. Syst.* i. 135.—Reg. Himal.
Wallichii, *Hook. Bot. Mag.* t. 4668.—Reg. Himal.

MECONOSTIGMA, Schott, Melet. 20 (1832) = **Philodendron**, Schott (Aroid.).
speciosum, Schott, l. c. = P. speciosum.

MECOPUS, Benn. Pl. Jav. Rar. 154. t. 32 (1838). *LEGUMINOSAE*, Benth. & Hook. f. i. 521.
nidulans, *Benn. Pl. Jav. Rar.* 154.—Java.

MECOSA, Blume, Bijdr. 403. t. 1 (1825) = **Habenaria**, Willd. (Orchid.).
angustata, Blume, l. c. 404.—Java.
dilatata, Blume, l. c. = H. dilatata.

MECOSCHISTUM, Dulac, Fl. Hautes-Pyr. 459 (1867) = **Lobelia**, Linn. (Campan.).
urens, Dulac, l. c. = L. urens.

MECOSTYLIS, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 44 = **Macaranga**, Thou. (Euphorb.).
acalyphoides, Kurz, ex Teijsm. & Binn. l. c.—Ins. Molucc.

MECRANIUM, Hook. f. in Benth. & Hook. f. Gen. i. 767 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
amygdalinum, *Hook. f. l. c.*—Ind. occ.

MECRANIUM:—
birimosum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Ins. S. Doming.
haemanthum, *Triana, l. c.*—Cuba.
integrifolium, *Triana, l. c.* 139.—Ind. occ.
multiflorum, *Triana, l. c.*—Ins. S. Doming.
purpurascens, *Hook. f. in Benth. & Hook. f. Gen.* i. 767.—Ins. Jamaic.
racemosum, *Wright, in Sauv. Fl. Cub.* 46.—Cuba.
virgatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Jamaic.

MEDEA, Klotzsch, in Wiegm. Arch. vii. (1841) 198 = **Croton**, Linn. (Euphorb.).
erecta, D. Dietr. Syn. Pl. v. 331 = seq.
hirta, Klotsch, in Wiegm. Arch. vii. (1841) 198 = C. timandroides.
timandroides, Didr. in Kjoeb. Vidensk. Meddel. (1857) 141 = C. timandroides.

MEDEMIA, Wuert. ex H. Wendl. in Bot. Zeit. xxxix. (1881) 89-90. *PALMAE.*
abiadensis, *H. Wendl. l. c.* 93.—Afr. trop.
Argun, Wuert. ex H. Wendl. l. c. = Hyphaena Argun.

MEDEOLA, Gronov. ex Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 833. GYROMIA, Nutt. Gen. Am. i. 238 (1818).
aculeata, Linn. Sp. Pl. 339 = Jacquinia ruscifolia.
angustifolia, [Delile, in] Red. Lil. t. 393 = Geitonoplesium cymosum.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = Asparagus medeoloides.
asparagoides, Linn. Sp. Pl. 339 = Asparagus medeoloides.
latifolia, Salisb. Prod. 252 = Asparagus medeoloides.
triphylla, Thib. ex Baker, in Journ. Linn. Soc. xiv. (1875) 607 = Asparagus crispus.
verticillifolia, Stokes, Bot. Mat. Med. ii. 325 = virginica.
virginica, *Linn. Sp. Pl.* 339.—Am. bor.

MEDICA, Cothen. Disp. Veg. 7 (1790) = **Tourretia**, Juss. (Bignon.).

MEDICA, Tourn. ex Linn. Syst. ed. I (1735) = **Medicago**, Tourn. (Legumin.).
arabica, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386 = Medicago maculata.
arborea, Mill. Gard. Dict. ed. VIII. n. 7; Medic. Beobacht. (1783) 231 = Medicago arborea.
bimuricata, Moench, Meth. 117 = *Medicago polymorpha.*
ciliaris, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386 = Medicago ciliaris.
coronata, Medic. l. c. = Medicago coronata.
cretica, Mill. Gard. Dict. ed. VIII. n. 6 = Medicago cretica.
echinata, Lam. Fl. Fr. ii. 587 = Medicago orbicularis.
falcata, Mill. Gard. Dict. ed. VIII. n. 2 = Medicago falcata.
hispanica, Mill. l. c. n. 4 = Hymenocarpos circinnatus.
incana, Moench, Meth. 117 = Medicago muricata.
inermis, Lam. Fl. Fr. ii. 586 = Medicago orbicularis, scutellata, turbinata.
intertexta, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386 = Medicago ciliaris.
italica, Mill. Gard. Dict. ed. VIII. n. 5 = Medicago obscura.
lupulina, Scop. Fl. Carn. ed. II. ii. 88 = Medicago lupulina.
marina, Lam. Fl. Fr. ii. 584 = Medicago marina.
media, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 359 = Medicago sativa.
orbicularis, Medic. in Vorles. Churf. Phys. Ges. ii. (1787) 386 = Medicago orbicularis.
polymorpha, Scop. Fl. Carn. ed. II. ii. 89.—Cf. *Medicago polymorpha.*
radiata, Mill. Gard. Dict. ed. VIII. n. 3 = Trigonella radiata.
sativa, Lam. Fl. Fr. ii. 584 = Medicago sativa.
tomentosa, Moench, Meth. Suppl. 38 = Medicago marina.
tornata, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386 = Medicago tornata.
turbinata, Medic. l. c. = Medicago turbinata.

MEDICAGO, Tourn. ex Linn. Gen. ed. I. 225 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 487.
DIPLOPRION, Viv. Fl. Lib. Spec. 48. t. 19 (1824).
LUPULARIA, Opiz, Seznam, 61 (1852).
MEDICA, Tourn. ex Linn. Syst. ed. I (1735).
MEDICULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386.
RADIATA, Medic. Phil. Bot. i. 208 (1789).
SPIROCARPUS, Opiz, Seznam, 93 (1852).
aculeata, Gaertn. Fruct. ii. 349 = hispida.
aculeata, Reichb. Fl. Germ. Excurs. 503 = truncatula.
aculeata, Willd. Sp. Pl. iii. 1410 = turbinata.
aegagropila, Desv. Journ. Bot. iii. (1814) 76 = hispida.
agrestis, *Tenore, Fl. Nap. Prod.* 45; *et Cat.* 59; *Prod. Suppl.* ii. p. lxxi.—Europ. austr.; Syria.
ambigua, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 147 = orbicularis.
amethystina, *C. L. Gmel. ex Steud. Nom.* ed. I. 513.—Hab. ?
angulata, Láng, Enum. 2; ex Neilr. Aufz. Ung. Gefaesspfl. 331 = sativa.
annularis, Bess. Prim. Fl. Galic. ii. 127 = glutinosa.
apennina, *J. Woods, Tourist's Fl.* 84.—Mont. Apenn.
apiculata, Bast. Fl. Maine-et-Loire, 280 = tuberculata.
apiculata, Willd. Sp. Pl. iii. 1414 = denticulata.
applanata, Willd. Enum. Hort. Berol. Suppl. 52 = orbicularis.
arabica, All. Fl. Pedem. i. 315 = maculata.
arborea, *Aubl. Pl. Gui.* ii. 779 (Quid ?).—Guian.
arborea, *Linn. Sp. Pl.* 778.—Italia; Graecia.
arborescens, Presl, Fl. Sicul. p. xx = arborea.
arcuata, *Trautv. in Bull. Sc. Acad. Pétersb.* viii. (1841) 272.—Reg. Caucas.
arenaria, Tenore, Cat. Hort. Neap. (1819) 59 = littoralis.
Aschersoniana, *Urban, in Verh. Bot. Ver. Brandenb.* xv. (1873) 77.—Afr. bor.
astroites, Bertol. Fl. Ital. viii. 272 = M. Helix.
astroites, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella astroites.
aurantiaca, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 425.—Hab. ?
banofcensis, Kit. in Linnaea, xxxii. (1863) 615 = rigidula.
Berteriana, Moris, in Ann. Stor. Nat. iv. (1830) 59 = tuberculata.
Berteroana, Nym. Consp. 169 = praec.
Biancae, Todaro, ex Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 60 = orbicularis.
bimuricata, Retz. ex Steud. Nom. ed. I. 513.—Cf. *polymorpha.*
Blancheana, *Boiss. Diagn*, Ser. II. v. 75.—Syria.
Bonarotiana, *Archang. in Nuov. Giorn. Bot.* viii. (1876) 5.—Italia.
brachyacantha, Kern. in Oestr. Bot. Zeitschr. xviii. (1868) 386 = minima.
brachycarpa, Fisch. ex Bieb. Fl. Taur. Canar. iii. 517 = Trigonella brachycarpa.
brachyloba, Fisch. ex Moris, in Mem. Acc. Torin. xxxvi. (1833) 190 = praec.
Braunii, Gren. & Godr. Fl. Fr. i. 393 = littoralis.
breviflora, Gilib. Fl. Lituan. ii. 95 = lupulina.
caerulea, Less. ex Ledeb. Fl. Ross. i. 526 = sativa.
Calcar, Lowe, in Hook. Kew Journ. viii. (1856) 291 = obscura.
canariensis, Benth. in Webb & Berth. Phyt. Canar. iii. 2. ii. t. 56 = obscura.
cancellata, *Bieb. Fl. Taur. Cauc.* ii. 226.—Reg. Casp.
cancellata, Tenore, Prod. p. xliv = Tenoreana.
canescens, R. Grah. in Wall. Cat. n. 5946 = denticulata.
canescens, *Menyh. Kalocsa Vidék. Növényt.* 61 (Quid?).—Hungar.
carstiensis, *Wulf. in Jacq. Coll.* i. 86; *Jacq. Ic. Pl. Rar.* i. 13. t. 156.—Carinth.
caspica, Jacq. ex Spreng. Nov. Prov. 27 = denticulata.
catalonica, Schrank, Pl. Rar. Hort. Monac. t. 28 = tuberculata.
catalonica, Urban, in Ind. Sem. Hort. Berol. (1872) 4 = hispida.
cibiniensis, Lerchen. ex Schur, Enum. Pl. Transs. 151 = sativa.
ciliaris, Balb. ex DC. Fl. Fr. Suppl. 548 = tuberculata.
ciliaris, *Krock, Fl. Siles.* ii. II. 244.—Reg. Mediterr.; Oriens.
ciliaris, Lucé, Fl. Osil. 253 = lupulina.
ciliaris, Savi, Due Cent. Fl. Etrusc. 148 = denticulata.

MEDICAGO:—
cinerascens, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 316 = rigidula.
circinnata, Gaertn. Fruct. ii. 348 = Hymenocarpos nummularius.
circinnata, Linn. Sp. Pl. 778 = Hymenocarpos circinnatus.
clypeata, Lindl. Bot. Reg. (1839) Misc. 60 = orbicularis.
cochleata, *Pall. Reise in Suedl. Statth. Russ. Reich.* ii. 336.—Reg. Caucas.
coerulea, Less. ex Nym. Consp. 166 = sativa.
commutata, Todaro, Ind. Sem. Hort. Panorm. (1867) 23, 31 = tornata.
confinis, Koch, Syn. Fl. Germ. ed. I. 164 = tuberculata.
conica, Schk. Handb. ii. 420 = hispida.
connivens, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella striata.
constricta, Dur. in Act. Linn. Soc. Bord. xxix. (1873) 15 = globosa.
contorta, Gilib. Fl. Lituan. ii. 94 = sativa.
cordata, Desr. in Lam. Encyc. iii. 636 = maculata.
corniculata, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella corniculata.
coronata, *Desr. in Lam. Encyc.* iii. 634.—Reg. Mediterr.; Oriens.
coronata, Gaertn. Fruct. ii. t. 155. f. 7 = tuberculata.
coronata, Pall. Ind. Taur.; ex Bieb. Fl. Taur. Cauc. ii. 227 = maculata.
corrugata, Dur. in Duch. Rev. Bot. i. (1845) 365 = obscura.
corymbifera, W. L. E. Schmidt, ex Schlecht. in Linnaea, iv. (1829) 74 = lupulina.
crassispina, Vis. in Flora, xii. (1829) I. Erg. 20 = truncatula.
cretacea, *Bieb. Fl. Taur. Cauc.* ii. 223.—Tauria.
cretica, Steud. Nom. ed. I. 512 (Quid?).—Ins. Creta.
crinita, Presl. Fl. Sicul. i. p. xx = M. Echinus.
cuneata, *J. Woods, Tourist's Fl.* 84.—Italia.
Cupianiana, Guss. Fl. Sic. Syn. ii. 362 = lupulina.
cuspidata, Tausch. in Flora, xiv. (1831) 211 = saxatilis.
cylindracea, DC. Cat. Hort. Monsp. 123 = littoralis.
daghestanica, *Rupr. ex Boiss. Fl. Orient.* ii. 95.—Reg. Caucas.
Decaisneana, *Todaro, Ind. Sem. Hort. Panorm.* (1866) 35.—Hab.?
Decandollii, *Tineo, ex Guss. Fl. Sic. Syn.* ii. 369.—Sicilia.
declinata, Kit. in Linnaea, xxxii. (1863) 613 = prostrata.
denticulata, Boiss. Fl. Orient. ii. 102 = hispida.
denticulata, *Willd. Sp. Pl.* iii. 1414.—Geront. bor. temp.
depressa, Jord. Cat. Jard. Dijon (1848), 28 = rigidula.
dicarpa, *Trautv. in Act. Hort. Petrop.* i. (1870) 24.—Rossia.
diffusa, Poir. Encyc. Suppl. iii. 524 = laciniata.
diffusa, Steud. Nom. ed. I. 513 = prostrata.
disciformis, *DC. Cat. Hort. Monsp.* 124.—Europ. austr.; As. Min.
distans, Poir. Encyc. Suppl. iii. 326 = hispida.
doliata, Carmign. in Giorn. Sc. Acc. Ital. i. (1818) 48; ex Bertol. Fl. Ital. viii. 292 = turbinata.
echinata, Bouch. Abbev. 56; ex DC. Fl. Fr. iv. 548 = hispida.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = M. Echinus.
echinata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 261.—Syria.
echinata, Steud. Nom. ed. II. ii. 109 = orbicularis.
Echinus, *DC. Fl. Fr.* iv. 546.—Reg. Mediterr.
elegans, Jacq. ex Willd. Sp. Pl. iii. 1408 = rugosa.
elegans, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 384 = truncatula, rigidula.
esculenta, Rottl. ex Willd. Enum. Hort. Berol. 799 = Trigonella corniculata.
falcata, Lam. Encyc. iii. 639 = sativa.
falcata, *Linn. Sp. Pl.* 779.—Geront. bor. temp.
Fischeriana, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella Fischeriana.
fissa, Trautv. l. c. 272 = Trigonella polycerata.
flexuosa, Tenore, Prod. Fl. Nap. p. xlv = denticulata.
fructicans, Hornem. Cat. Hort. Hafn. 24; ex Willd. Enum. Hort. Berol. 801 = cretacea.
galilaea, *Boiss. Diagn.* Ser. I. ix. 10.—Palestin.
Gerardi, Waldst. & Kit. ex Willd. Sp. Pl. iii. 1415 = rigidula.
germana, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 315 = rigidula.

MEDICAGO:—
glabella, *Presl, Delic. Prag.* 44.—Sicilia.
glabra, *Aubl. Pl. Gui.* ii. 778 (Quid?).—Guian.
glandulosa, Koch, Syn. Fl. Germ. ed. II 176 = falcata.
globosa, *Presl, Delic. Prag.* 45.—Sicilia.
globulifera, Desf. Tabl. ed. I. 190; Poir. Encyc. Suppl. iii. 526 = sphaerocarpa.
globulosa, *Desv. Journ. Bot.* iii. (1814) 77.—Afr. bor.
glomerata, Balb. Elencho, 93 = glutinosa.
glomerata, Fisch. ex Moris, in Mem. Acc. Torin. xxxvi. (1833) 190 = Trigonella brachycarpa.
glutinosa, *Bieb. Casp.* 191.—Reg. Caucas.
gracilis, *Bivona, in A. Bivona, Nuov. Piant. Ined.* 10; *Tod. Ind. Sem. Hort. Panorm.* (1858) 26.—Hab.?
gracillima, Tineo, ex Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 74 = hispida.
graeca, Hornem. Hort. Hafn. ii. 728 = minima.
granatensis, *Willd. Enum. Hort. Berol.* 803.—Hispan.
hamata, *Presl, Delic. Prag.* 45.—Sicilia.
hamosa, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella hamosa.
Helix, *Willd. Sp. Pl.* iii. 1409.—Reg. Mediterr.
heterocarpa, *Spach, ex Dur. Expl. Sc. Algérie*, t. 89.—Alger.
heterocarpa, Urban, in Ind. Sem. Hort. Berol. (1872) 3 = hispida.
hirsuta, All. Fl. Pedem. i. 315 = minima.
hirsuta, Schk. Handb. ii. 420 = denticulata.
hirsuta, Thuill. Fl. Par. ed. II. 390 = rigidula.
hirsuta, Vill. Hist. Pl. Dauph. iii. 403 = hispida.
hirsutissima, Cyrill. ex Steud. Nom. ed. II. ii. 109 = minima.
hispanica, Steud. l. c. = Hymenocarpos circinnatus.
hispida, *Gaertn. Fruct.* ii. 349.—Europ. austr.
Hornemanniana, *Ser. ex DC. Prod.* ii. 177.—Marocc.
hybrida, *Boiss. Diagn.* Ser. I. ix. 12, nomen.—Hab.?
hybrida, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Pourretii.
Hystrix, Tenore, Prod. Fl. Nap. 45 = denticulata.
inermis, Steud. Nom. ed. I. 513 = orbicularis, scutellata, tornata, turbinata.
inermis, Urban, in Ind. Sem. Hort. Berol. (1872) 4 = hispida.
intermedia, DC. Prod. ii. 174 = obscura.
intermedia, Schult. Obs. Bot. 160 = falcata.
intertexta, All. Fl. Pedem. i. 315 = M. Echinus.
intertexta, *Mill. Gard. Dict.* ed. VIII. n. 4.—Europ. austr.
intertexta, Willd. Sp. Pl. iii. 1411 = ciliaris.
italica, Steud. Nom. ed. II. ii. 109 = obscura.
laciniata, *Mill. Gard. Dict.* ed. VIII. n. 5.—Reg. Mediterr.
laevis, Desf. Fl. Atlant. ii. 213 = M. Helix.
Langeana, *Tod. Ind. Sem. Hort. Panorm.* (1867) 23, 31.—Europ.
lappacea, Desr. in Lam. Encyc. iii. 637 = denticulata.
laxiflora, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 172 = falcata.
leiocarpa, Benth. Cat. Pl. Pyr. 100 = suffruticosa.
lenticularis, Desr. in Lam. Encyc. iii. 630 = obscura.
Lessingii, *Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc.* (1839) 150, nomen.—Turcoman.
libyca, *Spreng. Syst.* iii. 289.—Libya.
lineata, Tausch, in Flora, xiv. (1831) 210 = minima.
littoralis, *Rhode, ex Hornem. Hort. Hafn.* ii. *Suppl.* 85.—Reg. Mediterr.
lunata, Boiss. Diagn. Ser. I. Index, 94 = Trigonella lunata.
lunata, Reichb. Fl. Germ. Excurs. 501 = Trigonella radiata.
lupulina, *Linn. Sp. Pl.* 779.—Geront. bor. temp.
macrocarpa, Boiss. Fl. Orient. ii. 96 = papillosa.
macrocarpa, Moris, Fl. Sard. i. 446. t. 45 = truncatula.
macrocarpa, Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 56 = falcata.
maculata, Hook. & Arn. Bot. Beech. Voy. 330 = denticulata.
maculata, *Willd. Sp. Pl.* iii. 1412.—Europ.
marginata, Willd. Enum. Hort. Berol. 802 = orbicularis.
marina, *Linn. Sp. Pl.* 779.—Reg. Mediterr.
marina, Pall. Reise, iii. 590 = rigidula.
maritima, Pall. ex Georgi, Beschr. Russ. Reich. III. iv. 1197 = rigidula.
media, Pers. Syn. ii. 356 = sativa.
mexicana, *Sweet, Hort. Brit.* ed. II. 138, nomen.—Mexic.

MEDICAGO:—
Meyeri, *Gruner, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 416.—Reg. Caucas.
microcarpa, Urban, in Ind. Sem. Hort. Berol. (1875) App. = hispida.
microcarpa, Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 57 = falcata.
microcarpa, Urban, l. c. 58 = papillosa.
microdon, Ehrenb. in Cat. Sem. Hort. Berol. (1827) = denticulata.
minima, *Linn. Fl. Ang.* 21; *Amoen. Acad.* iv. 105.—Europ.; Oriens?
mitis, Delile, ex Nym. Consp. 168 = agrestis.
mitis, Willd. ex Urban, in Ind. Sem. Hort. Berol. (1872) 4 = hispida.
mniocarpa, Wallr. ex Ser. in DC. Prod. ii. 172 = lupulina.
mollissima, Roth, Catalect. Bot. iii. 74 = minima.
monantha, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 272 = Trigonella polycerata.
monspeliaca, Trautv. l. c. = Trigonella monspeliaca.
Morisiana, Jord. Pugill. 53 = rigidula.
Murex, Willd. Sp. Pl. iii. 1410 = truncatula.
muricata, *All. Fl. Pedem.* i. 316.—Europ.
muricata, Benth. Cat. Pl. Pyr. 102 = turbinata.
muricata, Desr. in Lam. Encyc. iii. 634 = denticulata.
muricata, Griseb. Spicil. Fl. Rumel. i. 17 = sphaerocarpa.
muricata, Salzm, ex Ball, in Journ. Linn. Soc. xvi. (1878) 416 = ciliaris.
muricoleptis, DC. Prod. ii. 179 = Decandollii.
muricoleptis, *Tineo, Pl. Sic. Rar. Pugill.* i. 18.—Sicilia.
muricoleptis, Urban, in Ind. Sem. Hort. Berol. (1872) 4 = hispida.
muricula, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = truncatula.
neglecta, Guss. Fl. Sic. Prod. ii. 575 = turbinata.
neurocarpos, Kunze, ex Urban, in Ind. Sem. Hort. Berol. (1872) 3 = hispida.
nigra, Krock. Fl. Siles. ii. II. 244 = denticulata.
Noccae, Balb. in Brugnat. Giorn. Fis. viii. (1815) 65. t. 2 = carstiensis.
Noëana, *Boiss. Diagn.* Ser. II. ii. 10.—As. Min.; Syria.
nummularia, DC. Cat. Hort. Monsp. 124 = Hymenocarpos nummularius.
nummularia, Stev. in Bess. Cat. Hort. Crem. (1816) 85 = cretacea.
obscura, *Retz. Obs.* i. 24. t. 1.—Gall. austr.
obscura, Vis. Fl. Dalm. iii. 283 = hispida.
olivaeformis, Guss. Pl. Rar. 390 = turbinata.
orbicularis, *All. Fl. Pedem.* i. 314.—Reg. Mediterr.; Abyss.
ornithopodioides, Trautv. in Bull. Acad. Sc. Pétersb. viii. (1841) 271 = Trigonella ornithopodioides.
orthoceras, *Trautv. in Act. Hort. Petrop.* v. (1877) 421.—Reg. Caucas.
ovalis, *Urban, in Verh. Bot. Ver. Brandenb.* xv. (1873) 81.—Hispan.
ovata, Hort. Erf. ex Steud. Nom. ed. I. 514; Carmign. in Giorn. Acad. Ital. (1810); ex Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 73 = truncatula.
oxalioides, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 174 = maculata.
Paillerii, Phil. ex Nym. Consp. 166 = suffruticosa.
panormitana, Tineo, ex Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 62 = M. Echinus.
papillosa, *Boiss. Diagn.* Ser. I. ii. 23.—As. Min.
paradoxa, *Spreng. ex Steud. Nom.* ed. I. 514 (Quid?).—Hab.?
pauciflora, *Ledeb. Fl. Ross.* i. 526.—Reg. Casp.
pedunculata, *Ehrenb. ex Sweet, Hort. Brit.* ed. II. 134.—Aegypt.
pennata, *Schwaegr. ex Steud. Nom.* ed. I. 514 (Quid?).—Europ.
pentacycla, DC. Cat. Hort. Monsp. 124 = denticulata.
persica, Stev. ex Fisch. Cat. Hort. Gorenk. ed II. (1812) 69 = denticulata.
pilosa, Fisch. ex Moris, in Mem. Acc. Torin. xxxvi. (1833) 190 = Trigonella brachycarpa.
pilosa, *Schleich. Cat. Pl. Helvet.* ed. III. 46 (Quid?).—Helvet.
pinnatifida, Steud. Nom. ed. I. 514 = granatensis.
Pironae, *Vis. Ind. Sem. Hort. Patav.* (1855).—*Cf. Linnaea*, xxviii. (1856) 365.—Europ. austr.
plagiospira, Dur. in Duch. Rev. Bot. i. (1845) 366 = Soleirolii.

MEDICAGO:—
platycarpa, *Trautv. in Bull. Acad. Sc. Pétersb.* viii. (1841) 271.—Soongaria.
platycarpos, *Ledeb. Fl. Ross.* i. 523.—Sibiria.
plumbea, Bertol. Fl. Ital. viii. 267 = M. Helix.
polyantha, Dulac, Fl. Hautes-Pyr. 278 = denticulata.
polycarpa, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 175 (Quid ?).—Europ.
polycarpa, Willd. Enum. Hort. Berol. Suppl. 52 = hispida.
polycerata, Sauv. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 272 = Trigonella polycerata.
polycycla, Moris, in Mem. Acc. Torin. xxxviii. (1835) p. xxxi = denticulata.
polymorpha, Linn. Sp. Pl. 779 = orbicularis (etc.).
polymorpha, Roxb. Hort. Beng. 58 ; Fl. Ind. iii. 390 = denticulata.
pontificalis, *Gennari, in Nuov. Giorn. Bot. Ital.* ii. (1870) 132.—Ins. Caprera.
Pourretii, *Noul. Fl. Bass. Sous-Pyr.* 151.—Europ.
praecox, *DC. Cat. Hort. Monsp.* 123.—Europ. austr.
procumbens, Bess. Prim. Fl. Galic. ii. 127 = falcata.
prostrata, *Jacq. Hort. Vindob.* i. 39. t. 89.—Italia.
Pseudo-Gerardi, Nym. Consp. 168 = agrestis.
Pseudo-Gerardi, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 174 = rigidula.
psilocarpa, *Trautv. in Bull. Sc. Acad. Pétersb.* viii. (1841) 272.—Hab. ?
pubescens, DC. Cat. Hort. Monsp. 124 = tuberculata.
pubescens, *Guss. Pl. Rar.* 313.—Europ. austr.
pubescens, Hornem. Hort. Hafn. ii. 726 = Hornemanniana, truncatula.
pugionata, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 272 = Trigonella spinosa.
pulchella, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimpr. 25 = minima.
pulchella, *Tod. Ind. Sem. Hort. Panorm.* (1868) 36.—Hab. ?
pungens, Trautv. in Bull. Acad. Sc. Pétersb. viii. (1841) 271 = Trigonella maritima.
purpurea, *Gueldenst. Reise*, i. 195.—Rossia.
pusilla, Viv. Pl. Aegypt. Arab. Dec. 14 = littoralis.
radiata, Linn. Sp. Pl. 778 = Trigonella radiata.
recta, Desf. Fl. Atlant. ii. 212 = minima.
recta, *Kit. in Linnaea*, xxxii. (1863) 613 (Quid ?).—Hungar.
reniformis, Dulac, Fl. Hautes-Pyr. 278 = lupulina.
reticulata, Benth. Cat. Pl. Pyr. 101 = hispida.
rigida, Boiss. Fl. Orient. ii. 100 = rigidula.
rigidula, Bourg. ex Nym. Consp. 167 = M. Helix.
rigidula, *Desr. in Lam. Encyc.* iii. 634.—Europ.
rigidula, Hoffm. ex Steud. Nom. ed. II. ii. 109 = minima.
rigidula, Willd. Sp. Pl. iii. 1417 = truncatula.
rotata, *Boiss. Diagn.* Ser. I. ii. 24.—Syria.
rugosa, *Desr. in Lam. Encyc.* iii. 632.—Europ.
rugosa, Urv. in Mém. Soc. Linn. Par. i. (1822) 354 = tuberculata.
rupestris, *Bieb. Fl. Taur. Cauc.* ii. 225.—Tauria.
ruthenica, *Trautv. in Bull. Sc. Acad. Pétersb.* viii. (1841) 271 ; *Ledeb. Fl. Ross.* i. 523.—Sibiria.
sardoa, Moris, Stirp. Sard. Elench. 15 = denticulata.
sativa, *Linn. Sp. Pl.* 778.—Europ. ; Oriens.
sativa, Urban, in Verh. Bot. Ver. Brandenb. xv. (1873) 56 = falcata.
saxatilis, *Bieb. Fl. Taur. Cauc.* ii. 225.—Tauria.
Schimperiana, Hochst. ex Boiss. Fl. Orient. ii. 104 = laciniata.
scutellaris, Reichb. Ic. Fl. Germ. t. 2114 = seq.
scutellata, *Mill. Gard. Dict.* ed. VIII. n. 2.—Reg. Mediterr.
scutellata, Pall. Ind. Taur. ; ex Bieb. Fl. Taur. Cauc. ii. 226 = orbicularis.
secundiflora, *Dur. in Duch. Rev. Bot.* i. (1845) 365.—Alger. ; Hispan. ?
secundiramea, Willk. & Lange, Prod. Fl. Hisp. iii. 417 = praec.
sibirica, *DC. Prod.* ii. 173.—Sibiria.
sicula, Tod. Ind. Sem. Hort. Panorm. (1868) 27 = sphaerocarpa.
Soleirolii, *Duby, Bot. Gall.* 124.—Italia.
Sorrentini, Tineo, in Tod. Ind. Sem. Hort. Panorm. (1859).—Cf. Linnaea, xxx. (1859–60) 759 = truncatula.
sphaerocarpa, *Bertol. Amoen. Ital.* 91.—Reg. Mediterr.
sphaerocarpa, Costa, Fl. Catal. 60 = turbinata.
spinulosa, DC. Fl. Fr. v. 569 = tuberculata.
stellata, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 272 = Trigonella stellata.

MEDICAGO:—
stipularis, Wallr. Beitr. Fl. Hercyn. 260 ; et in Linnaea, xiv. (1840) 630 = lupulina.
striata, *Bast. in Desv. Journ. Bot.* iii. (1814) 19.—Italia.
stricta, Wallr. ex Urban, in Ind. Sem. Hort. Berol. (1872) 3 = hispida.
strumaria, Hort. Erf. ex Steud. Nom. ed. I. 514 (Quid ?).—Europ. austr.
subfalcata, Schur, Enum. Pl. Transs. 151 = sativa.
subinermis, Bertol. Fl. Ital. viii. 290 = littoralis.
suffruticosa, *Ram. ex DC. Fl. Fr.* iv. 541.—Reg. Mediterr. occ.
sylvestris, Fries, Mant. iii. 92 = falcata.
Tenoreana, *Ser. in DC. Prod.* ii. 180.—Italia.
tentaculata, Willd. Sp. Pl. iii. 1413 = truncatula.
tenuifolia, Presl, Fl. Sicul. i. p. xx = prostrata.
tenuis, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = Trigonella striata.
terebella, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
Terebellum, Willd. Sp. Pl. iii. 1416 = hispida.
tetracycla, Presl, Fl. Sicul. i. p. xx = littoralis.
Timeroyi, Jord. Cat. Jard. Dijon (1848), 29 = truncatula.
tomentosa, Dulac, Fl. Hautes-Pyr. 278 = suffruticosa.
tornata, Desr. in Lam. Encyc. iii. 633 = turbinata.
tornata, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ. austr.
tornata, Webb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 412 = M. Helix.
tortuosa, *Schwaegr. ex Steud. Nom.* ed. I. 514.—Hab. ?
tribuloides, Desr. in Lam. Encyc. iii. 635 = trunculata.
tricycla, DC. Cat. Hort. Monsp. 125 = striata.
truncatula, *Gaertn. Fruct.* ii. 150.—Europ.
truncatulata, Gren. & Godr. Fl. Fr. i. 395, sphalm. = praec.
tuberculata, Urban, in Ind. Sem. Hort. Berol. (1872) App. 3 = hispida.
tuberculata, *Willd. Sp. Pl.* iii. 1410.—Reg. Mediterr.
turbinata, Carmign. in Giorn. Pis. xii. (1810) I. 13 ; ex DC. Fl. Fr. Suppl. 567 = tuberculata.
turbinata, *Willd. Sp. Pl.* iii. 1409.—Reg. Mediterr.
umbellata, Hort. ex DC. Prod. ii. 178 = carstiensis.
uncinata, Auct. ex Nym. Consp. 169 = denticulata.
uncinata, Urban, in Ind. Sem. Hort. Berol. (1872) 4 = hispida.
uncinata, Willd. Sp. Pl. iii. 1417 = truncatula.
varia, Martyn, Fl. Rust. iii. 87 = sativa.
Verloti, Song. & Perrier, in Billotia, i. (1864) 74 = rigidula.
vermicularis, Cesati, in Friedr. Reise, 274 = coronata.
villosa, Brot. Fl. Lusit. ii. 116 = minima.
villosa, DC. Fl. Fr. iv. 545 = rigidula.
villosula, Baumg. Enum. Stirp. Transs. ii. 382 = truncatula.
virginica, Linn. Sp. Pl. 778 = Lespedeza capitata, reticulata.
viscida, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 173 = minima.
vulgaris, Alef. Landw. Fl. 75 = sativa ?
Willdenowii, Mérat, Fl. Par. 296 = lupulina.

MEDICASIA, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 219 (1870), sphalm. = MEDICUSIA, Moench = **Picris**, Linn. (Compos.).

MEDICIA, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 324 = **Gelsemium**, Juss. (Logan.).
elegans, Gardn. & Champ. l. c. 325 = G. elegans.

MEDICOSMA, Hook. f. in Benth. & Hook. f. Gen. i. 296 (1862). *RUTACEAE*, Benth. & Hook. f. l. c.
Cunninghamii, *Hook. f. l. c.* 297.—Austral.

MEDICULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 386 = **Medicago**, Tourn. (Compos.).
lupulina, Medic. l. c. = Medicago lupulina.

MEDICUSIA, Moench, Meth. 536 (1794) = **Picris**, Linn. (Compos.).
aspera, Moench, l. c. 537 = P. Sprengeriana.
lappacea, Reichb. Fl. Germ. Excurs. 254 = P. pauciflora.
Sprengeriana, Reichb. l. c. = P. Sprengeriana.

MEDINILLA, Gaudich. in Freyc. Voy. Bot. 484. t. 106 (1826). *MELASTOMACEAE*, Benth. & Hook. f. i. 759.
DACTYLIOTA, Blume, Mus. Bot. Lugd. Bat. i. 21 (1849).
DIPLOGENEA, Lindl. in Brand. Quart. Journ. (1828) 122.
ERPETINA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 299.
GALLARIA, Schrank, ex Endl. Gen. 1217 (1841).
HYPENANTHE, Blume, Mus. Bot. Lugd. Bat. i. 21 (1849).
TRIPLECTRUM, D. Don, in Wight & Arn. Prod. 324 (1834).
alpestris, *Blume, in Flora*, xiv. (1831) 514.—Sumatra.
alternifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 19.—Sumatra.
amabilis, Dyer, in Gard. Chron. (1874) I. 373 = Teysmanni.
amoena, *Seem. Fl. Vit.* 88.—Ins. Fiji.
amplexicaulis, *Blume, Mus. Bot. Lugd. Bat.* i. 20.—Sumatra.
astronioides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 88.—Ins. Philipp.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 548.—Ind. or.
bornensis, *Blume, Mus. Bot. Lugd. Bat.* i. 20.—Borneo.
bracteata, *Blume, in Flora*, xiv. (1831) 519.—Ins. Molucc.
bracteata, Hort. Veitch, in Fl. des Serres, Sér. I. vi. (1850–51) t. 572 = magnifica.
celebica, *Blume, Mus. Bot. Lugd. Bat.* i. 18.—Ins. Celebes.
crassifolia, *Blume, in Flora*, xiv. (1831) 511.—Java.
crassifolia, Mor. Syst. Verz. Arch. Ind. 11 = succulenta.
crassifolia, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 288 = Naudinii.
crassifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 86, partim = Hasseltii.
crassinervia, *Blume, in Flora*, xiv. (1831) 510.—Ins. Banda.
crispata, *Blume, l. c.* 517.—Ins. Molucc.
Cummingii, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 292.—Ins. Philipp.
Curtisii, *Hort. Veitch, in Gard. Chron.* (1888) II. 621.—Sumatra.
cuspidata, *Blume, Mus. Bot. Lugd. Bat.* i. 18.—Borneo.
divaricata, *Baker, in Journ. Linn. Soc.* xx. (1883) 149.—Madag.
Erpetina, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87.—Malaya.
erythrophylla, Lindl. Bot. Reg. (1838) Misc. 85 = rubicunda.
eximia, *Blume, in Flora*, xiv. (1831) 515.—Sumatra.
eximia, Siebold, ex Planch. in Fl. des Serres, Sér. I. v. (1849) 482 = Sieboldiana.
farinosa, Hort. Groenew, ex Regel, Ind. Sem. Hort. Petrop. (1865) 52 = venosa.
fasciculata, *Baker, in Journ. Linn. Soc.* xx. (1883) 148.—Madag.
fuchsioides, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 12.—Zeylan.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 549.—Reg. Himal.
Halogeton, *S. Moore, in Journ. Bot.* xviii. (1880) 3.—Ins. Admiralt.
Hasseltii, *Blume, in Flora*, xiv. (1831) 513.—Malaya.
heterophylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 598. t. 75.—Ins. Fiji.
himalayana, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 88.—Reg. Himal.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. I. 545.—Java.
humilis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 427.—Ins. Celebes.
hypericifolia, *Blume, in Flora*, xiv. (1831) 513.—Java.
inaequalis, *Miq. Fl. Ind. Bat.* i. I. 542.—Java.
intermedia, *Blume, in Flora*, xiv. (1831) 516.—Java.
javanensis, *Blume, l. c.* 515.—Java.
javanensis, Korth. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 88 = bornensis.
javanica, Triana, l. c. t. 7. f. 94 a = javanensis.
javensis, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 20 = bornensis.
Korthalsii, *Blume, l. c.* 18.—Sumatra.
Kuhlii, *Blume, in Flora*, xiv. (1831) 514.—Java.

MEDINILLA:—
Lagunae, *Vidal, Cat. Pl. Pr. Manila*, 31.—Ins. Philipp.
lanceolata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 344.—Madag.
laurifolia, *Blume, in Flora*, xiv. (1831) 511.—Java.
leptophylla, *Baker, in Journ. Linn. Soc.* xxi. (1884) 343.—Madag.
lophoclada, *Baker, l. c.* 344.—Madag.
loranthoides, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 290.—Ins. Borbon.
luzonensis, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 88.—Ins. Philipp.
macrocarpa, *Blume, in Flora*, xiv. (1831) 510.—Madag.
macrocarpa, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 547 = crassinervia.
macrophylla, *Blume, Mus. Bot. Lugd. Bat.* i. 19.—Sumatra.
maculata, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 13.—Zeylan.
magnifica, *Lindl. in Paxt. Flow. Gard.* i. (1850) 55. t. 12.—Ins. Philipp.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 549.—Malacca.
malabarica, *Bedd. Ic. Pl. Ind. Or.* i. 33.—Ind. or.
Mannii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 460; *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87.—Afr. trop.
merguiensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 549.—Burma.
Motleyi, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 89.—Borneo.
muricata, *Blume, Mus. Bot. Lugd. Bat.* i. 20.—Sumatra.
myrtiformis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 86.—Ins. Philipp.
Naudinii, *Triana, l. c.*—Java.
occidentalis, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 295.—Madag.
papillosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 148.—Madag.
papuana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 24.—N. Guin.
parviflora, Triana, in Trans. Linn. Soc. xxviii. (1871) 89 = parvifolia, *Seem.*
parvifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 149.—Madag.
parvifolia, *Seem. Fl. Vit.* 89.—Ins. Fiji.
pauciflora, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 88.—Reg. Himal.
polyantha, Korth. in Verh. Nat. Gesch. Bot. 245. t. 61 = quadrifolia.
pterocaula, *Blume, in Flora*, xiv. (1831) 509.—Java.
quadrifolia, *Blume, l. c.*—Java.
radicans, Bedd. Ic. Pl. Ind. Or. t. 184 = Beddomei.
radicans, *Blume, in Flora*, xiv. (1831) 509.—Java.
rhodochlaena, *A. Gray, Bot. U. St. Expl. Exped.* i. 600.—Ins. Fiji.
rosea, *Gaudich. in Freyc. Voy. Bot.* 484. t. 106.—Malaya.
rubicunda, *Blume, in Flora*, xiv. (1831) 512.—Malaya.
salicifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 21.—Java.
setigera, *Miq. Fl. Ind. Bat.* i. 1. 550.—Sumatra.
Sieboldiana, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 126, sub. t. 572.—Ins. Molucc.
Sieboldtiana, Planch. l. c. v. (1849) t. 482 = praec.
speciosa, *Blume, in Flora*, xiv. (1831) 515.—Java.
speciosa, Hook. Bot. Mag. t. 4321 = Cummingii.
succulenta, *Blume, in Flora*, xiv. (1831) 513.—Java.
succulenta, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 18 = Korthalsii.
superba, Teijsm. & Binn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 87 = magnifica.
ternatensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 217.—Ins. Ternate.
ternifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 88.—Ins. Philipp.
Teysmanni, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 217.—Ins. Celebes.
venosa, *Blume, in Flora*, xiv. (1831) 518.—Malaya.
verrucosa, *Blume, l. c.* 514.—Java.
viscoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 87.—Madag.
Walkeri, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 11 = Kendrickia Walkeri.
Waterhousei, *Seem. Fl. Vit.* 89.—Ins. Fiji.

MEDIUM, [Tourn.] Fisch. ex A. DC. Monog. Camp. 213 (1830) = **Campanula**, Tourn.
alliariaefolium, Spach, Hist. Vég. Phan. ix. 557 = C. alliariaefolia.
barbatum, Spach, l. c. 555 = C. barbata.
grandiflorum, Spach, l. c. 553 = C. Medium.
gummiferum, Spach, l. c. 556 = C. sarmatica.
saxifragum, Fisch. ex A. DC. Monog. Camp. 228, 229 = C. Saxifraga, ciliata.
sayanense, Fisch. ex A. DC. l. c. 229 = C. pilosa.
sibiricum, Spach, Hist. Vég. Phan. ix. 554 = C. sibirica.

MEDORA, Kunth, Enum. Pl. v. 155 (1850) = **Smilacina**, Desf. (Liliac.).
divaricata, Kunth, l. c. 156 = S. fusca.

MEDUSA, Lour. Fl. Cochinch. 406 (1790) = **Alsodeia**, Thou. (Violar.).
anguifera, Lour. l. c. = A. echinocarpa.

MEDUSAEA, Reichb. Nom. 193 (1841) = MEDUSEA, Haw. = **Euphorbia**, Linn.

MEDUSAGYNE, Baker, Fl. Maurit. 16 (1877). *TERNSTROEMIACEAE.*
oppositifolia, *Baker, l. c.* 17; *et in Hook. Ic. Pl.* t. 1252.—Ins. Mascar.

MEDUSANTHERA, Seem. in Journ. Bot. ii. (1864) 74 = **Lasianthera**, Beauv. (Olacin.).
vitiensis, Seem. l. c.—Ins. Fiji.

MEDUSEA, Haw. Syn. Pl. Succ. 133 (1812) = **Euphorbia**, Linn.
Fructus-pini, Haw. l. c. 134 = E. Caput-Medusae.
globosa, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251 = E. globosa.
hamata, Klotzsch & Garcke, l. c. = E. hamata.
major, Haw. Syn. Pl. Succ. 134 = E. Caput-Medusae.
patula, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251 = E. anacantha.
procumbens, Haw. Syn. Pl. Succ. 134 = E. procumbens.
tessellata, Haw. l. c. 135 = E. Caput-Medusae.
tridentata, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251 = E. anacantha.
tuberculata, Klotzsch & Garcke, l. c. = E. tuberculata.

MEDUSULA, Pers. Syn. ii. 215 (1807) = MEDUSA, Lour. = **Alsodeia**, Thou. (Violar.).

MEDYPHYLLA, Opiz, in Lotos, viii. (1858) 65 = **Astragalus**, Tourn. (Legumin.).
rotundifolia, Opiz, l. c. = A. nummularioides.

MEERBURGIA, Moench, Meth. Suppl. 116 (1802) = **Pollichia**, Soland. (Illecebr.).
glomerata, Moench, l. c. = P. campestris.

MEESIA, Gaertn. Fruct. i. 344 (1788) = **Gomphia**, Schreb. (Ochnac.).
serrata, Gaertn. l. c. = G. angustifolia.

MEGABOTRYA, Hance, in Walp. Ann. ii. 259 (1851) = **Evodia**, Forst. (Rutac.).
meliaefolia, Hance, l. c. = E. meliaefolia.

MEGACARPAEA, DC. Syst. ii. 417 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 91.
angulata, *DC. Syst.* ii. 418.—Sibir.
bifida, *Benth. in Hook. Kew Journ.* vii. (1855) 357.—Reg. Himal.
gigantea, *Regel, in Bull. Soc. Nat. Mosc.* xlii. (1870) i. 263.—Turkestan.
laciniata, *DC. Syst.* ii. 417.—Reg. Casp.
polyandra, *Benth. in Hook. Kew Journ.* vii. (1855) 353. t. 7.—Reg. Himal.

MEGACARPHA, Hochst. in Flora, xxvii. (1844) 551 = **Oxyanthus**, DC. (Rubiac.).
pyriformis, Hochst. l. c. = O. natalensis.

MEGACARYON, Boiss. Pl. Or. Nov. Dec. i. 7 (1875). *BORAGINEAE*, Benth. & Hook. f. ii. 863.
armenum, *Boiss. Pl. Or. Nov. Dec.* i. 7.—Armenia.
orientale, Boiss. Fl. Orient. iv. 204 = armenum.

MEGACLINIUM, Lindl. Bot. Reg. t. 989 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 505.
Bufo, *Lindl. Bot. Reg.* (1841) *Misc.* 42.—Afr. trop.
colubrinum, *Reichb. f. in Bonplandia*, iii. (1855) 223.—Hab. ?
falcatum, *Lindl. Bot. Reg.* t. 989.—Afr. trop.
flaccidum, Hook. Cent. Orch. t. 3 = maximum.
maximum, *Lindl. Gen. et Sp. Orch.* 47.—Afr. trop.
melanorrhachis, *Reichb. f. in Gard. Chron.* (1875) ii. 162.—Afr. trop.
Melleri, *Hook. f. Bot. Mag.* sub t. 5936, nomen; *Reichb. f. in Flora*, lxi. (1878) 78.—Afr. trop.
oxypterum, *Lindl. Bot. Reg.* (1839) *Misc.* 14.—Afr. trop.
purpuratum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 128.—Afr. trop.
Sandersoni, *Oliver, in Bot. Mag.* sub t. 5936, nomen; *Reichb. f. in Flora*, lxi. (1878) 78.—Afr. austr.
tentaculigerum, *Reichb. f. in Flora*, lxi. (1878) 78, in textu.—Afr. trop.
velutinum, *Lindl. Bot. Reg.* (1847) t. 32.—Afr. trop

MEGADENDRON, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1875) 109 = **Barringtonia**, Forst. (Myrtac.).
ambiguum, Miers, l. c. 110.—Ins. Timor.
macrocarpum, Miers, l. c, 109 = B. macrocarpa.
pallidum, Miers, l. c. 110.—Java.

MEGADENUS, Rafin. Neogenyt. 4 (1825) = **Scirpus**, Tourn. (Cyperac.).
capitatus, Rafin. l. c. = Eleocharis capitata.
palustris, Rafin. l. c. = S. palustris.
tuberculatus, Rafin. l. c. = Eleocharis tuberculosa.

MEGALACHNE, Steud. in Flora, xxxiii. (1850) 229, nomen; et Syn. Pl. Gram. 237 (1855). *GRAMINEAE.*
Berteroniana, *Steud. ll. cc.*—Ins. J. Fernand.

MEGALACHNE, Thw. Enum. Pl. Zeyl. 372 (1864) = **Eriachne**, R. Br. (Gramin.).
zeylanica, Thw. l. c. = E. triseta.

MEGALOTHECA, F. Muell. Fragm. viii. 98 (1873) = **Restio**, Linn.
stricta, F. Muell. l. c. 99 = R. Megalotheca.

MEGALOTROPIS, Griff. Notul. iv. 441 (1854) = **Butea**, Roxb. (Legumin.).
buteiformis, 'Voigt,' ex Griff. l. c. = B. minor.

MEGAPLEILIS, Rafin. Fl. Tellur. ii. 57 (1836) = **Gesneria**, Mart.
tuberosa, Rafin. l. c. = G. bulbosa.

MEGAPTERIUM, Spach, Hist. Vég. Phan. iv. 363 (1835) = **Oenothera**, Linn. (Onagr.).
missouriense, Spach, l. c. 364 = Oe. missouriensis.
Nuttallianum, Spach, l. c. = Oe. missouriensis.

MEGARRHIZA, Torr. & Gray, in Pacif. Rail. Rep. vi. 74, nomen (1857); xii. ii. 61 (1860) = **Echinocystis**, Torr. (Cucurb.).
californica, Torr. l. c. vi. 74, nomen = E. fabacea.
californica, S. Wats. in Bot. Calif. i. 241, partim. = E. macrocarpa.
gilensis, Greene, in Bull. Torrey Bot. Club, viii. (1881) 97 = E. gilensis.
guadalupensis, S. Wats. in Proc. Am. Acad. xi. (1876) 115, 138 = E. guadalupensis.
Marah, S. Wats. l. c. 138 = E. Marah.
muricata, S. Wats. l. c. 139 = E. Watsonii.
oregona, Torr. in Pacif. Rail. Rep. xii. ii. 61 = E. oregana.

MEGASEA, Haw. Enum. Saxifr. 6 (1821) = **Saxifraga**, Tourn.
ciliata, Haw. l. c. 7 = S. ligulata.
cordifolia, Haw. l. c. = S. crassifolia.
crassifolia, Haw. l. c. 6 = S. crassifolia.
media, Haw. l. c. 7 = S. crassifolia.

MEGASTACHYA, Beauv. Agrost. 74. t. 15. f. 5 (1812) = **Eragrostis**, Beauv. (Gramin.).
abyssinica, Hochst. ex Steud. Syn. Pl. Gram. 203 = Enderopogon macrostachya.
acutiflora, Roem. & Schult. Syst. ii. 588 = E. acutiflora.

MEGASTACHYA :—
amabilis, Beauv. Agrost. 74 = Eragrostis amabilis.
amoena, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 572 = E. amoena.
aturensis, Roem. & Schult. Syst. ii. 587 = E. aturensis.
Berteroniana, Schult. Mant. ii. 330 = E. Berteroniana.
bipennata, Beauv. Agrost. 74 = E. cynosuroides.
Boryana, Boj. Hort. Maurit. 369 = E. ciliaris.
Boryana, Roem. & Schult. Syst. ii. 592 = E. Boryana.
brasiliensis, Schult. Mant. ii. 620 = E. elegans.
breviflora, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 576 = E. reptans ?
brizoides, Beauv. Agrost. 74 = E. brizoides, *Nees*.
canadensis, Roem. & Schult. Syst. ii. 593 = Briza canadensis.
ciliaris, Beauv. Agrost. 74 = E. ciliaris, *Link*.
ciliata, Schult. Mant. ii. 328 = E. ciliata.
condensata, J. & C. Presl, Rel. Haenk. i. 284 = Distichlis condensata.
corymbifera, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 576 = E. reptans ?
cuspidata, Schult. Mant. ii. 329 = E. Vahlii.
cylindrica, Schult. l. c. 328 = E. ciliata.
elongata, Beauv. Agrost. 74 = E. elongata.
Eragrostis, Beauv. l. c = E. major.
fasciculata, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 576 = E. reptans ?
glomerata, Schult. Mant. ii. 327 = E. conferta.
Gouini, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 576 = E. reptans ?
hypnoides, Beauv. Agrost. 74 = E. reptans.
infirma, Roem. & Schult. Syst. ii. 585 = E. infirma.
leersioides, Presl, Cyp. et Gram. Sic. 44 = Poa leersioides.
maxima, Boj. ex Baker, in Journ. Linn. Soc. xxi. (1885) 454 = E. maxima.
maypurensis, Roem. & Schult. Syst. ii. 588 = E. maypurensis.
Montufari, Roem. & Schult. Syst. ii. 586 = E. Montufari.
mucronata, Beauv. Agrost. 74 = E. mucronata.
multiflora, Beauv. l. c. = E. major.
nigricans, Roem. & Schult. Syst. ii. 586 = E. nigricans.
oblonga, Beauv. Agrost. 74 = E. major.
obtusa, Schult. Mant. ii. 326 = Poa obtusa.
Olmedi, Roem. & Schult. Syst. ii. 586 = E. Olmedi.
oxylepis, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 574 = E. oxylepis.
panamensis, Fourn. ex Hemsl. l. c. = E. maypurensis.
panicoides, J. & C. Presl, Rel. Haenk. i. 283 = E. panicoides.
papposa, Roem. & Schult. Syst. ii. 585 = E. papposa.
pastoënsis, Roem. & Schult. l. c. 587 = E. pastoënsis.
patula, Roem. & Schult. l. c. 585 = E. patula.
polymorpha, Beauv. Agrost. 74 = E. Brownei.
prostrata, Roem. & Schult. Syst. ii. 590 = Distichlis prostrata.
pulchella, Roem. & Schult. Syst. ii. 591 = Festuca rigida.
purpurascens, Schult. Mant. ii. 326 = Briza purpurascens.
reptans, Beauv. Agrost. 74 = E. reptans.
rigida, Roem. & Schult. Syst. ii. 591 = Festuca rigida.
riparia, Roem. & Schult. l. c. 593 = E. ciliaris, *Link*.
rupestris, Roem. & Schult. l. c. 592 = E. ciliata.
simpliciflora, J. & C. Presl, Rel. Haenk. i. 283 = Poa simpliciflora.
spectabilis, Roem. & Schult. Syst. ii. 589 = E. pectinacea.
Swainsonii, Raddi, Agrost. Bras. 52 = E. Vahlii.
tenax, Roem. & Schult. Syst. ii. 587 = E. tenax.
tenella, Boj. Hort. Maurit. 369 = E. plumosa.
thalassica, Roem. & Schult. Syst. ii. 590 = Poa thalassica.
triticea, Presl, Cyp. et Gram. Sic. 45 = E. triticea.
uninervia, J. & C. Presl, Rel. Haenk. i. 279 = E. uninervia.
unioloides, Presl, Cyp. et Gram. Sic. 44 = Demazeria sicula.

MEGASTEGIA, G. Don, Gen. Syst. ii. 468 (1832) = **Brongniartia**, H. B. & K. (Legumin.).
speciosa, G. Don, l. c. = B. thermoides.

MEGASTIGMA, Hook. f. in Benth. & Hook. f. Gen. i. 299 (1862). *RUTACEAE*, Benth. & Hook. f. l. c.
Galeottii, *Baill. Adansonia*, x. (1871–72) 331.—Mexic.
Skinneri, *Hook. f. in Benth. & Hook. f. Gen.* i. 299.—Guatemala.

MEGASTOMA, Coss. & Dur. ex Benth. & Hook. f. Gen. ii. 851 (1876) = **Eritrichium**, Schrad. (Borag.).

MEGISTA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 115 = **Physalis**, Linn. (Solanac.).
maxima, Fourr. l. c. = P. Alkakengi.

MEGOPIZA, Rafin. Fl. Tellur. iv. 110 (1836) = **Utricularia**, Linn. (Lentibul.).
fornicata, Rafin. l. c. = U. gibba.
integra, Rafin. l. c. = U. biflora.
longirostris, Rafin. l. c. = U. fibrosa.
macrorhiza, Rafin. l. c. = U. vulgaris.

MEGOTIGEA, Rafin. Fl. Tellur. iii. 64 (1836) = **Helicodiceros**, Schott (1853) (Aroid.).
crinita, Rafin. l. c. = H. crinitus.

MEGOTRYS, Rafin. in Am. Monthly Mag. (1818) 267 = **Cimicifuga**, Linn. (Ranunc.).
serpentaria, Rafin. l. c. = C. racemosa.

MEGYATHUS, Rafin. Fl. Tellur. iii. 93 (1836) = **Salvia**, Tourn. (Labiat.).
Acetabulum, Rafin. l. c. = S. acetabulosa.

MEHENBETHENE, Besl. ex Gaertn. Fruct. ii. 98 (1788); Walp. Ann. ii. 291 (1851) = **Canarium**, Linn. (Burser.).

MEIALISA, Rafin. Sylva Tellur. 63 (1838) = **Adriana**, Gaudich. (Euphorb.).
australis, Rafin. l. c. 64 = A. quadripartita.

MEIAPINON, Rafin. Fl. Tellur. iii. 32 (1836) = **Linum**, Tourn.
saginoides, Rafin. l. c. 33 = L. selaginoides ?

MEIBOMIA, Heist. ex Adans. Fam. ii. 509 (1763) = **Desmodium**, Desv. (Legumin.).

MEIEMIANTHERA, Rafin. Sylva Tellur. 25 (1838) = **Cytisus**, Linn. (Legumin.).
eolica, Rafin. l. c = C. aeolicus.

MEIENA, Rafin. Sylva Tellur. 125 (1838) = **Loranthus**, Linn.
axillaris, Rafin. l. c. = L. pentandrus.

MEINECKIA, Baill. Etud. Gén. Euph. 586 (1858) = **Securinega**, Comm. (Euphorb.).
phyllanthoides, Baill. l. c. 587 = S. phyllanthoides.

MEIOGYNE, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 12 (1865–66) = **Unona**, Linn. f. (Anonac.).
virgata, Miq. l. c.—Malaya.

MEIONECTES, R. Br. in Flind. Voy. App. ii. 550 (1814). *HALORAGEAE*, Benth. & Hook. f. i. 675.
Brownii, *Hook. f. in Ic. Pl.* t. 306.—Austral.
Preissii, Nees, in Lehm. Pl. Preiss. ii. 224 = Brownii.

MEIONULA, Rafin. Fl. Tellur. iv. 108 (1836) = **Utricularia**, Linn. (Lentibul.).
parviflora, Rafin. l. c. = U. lateriflora.

MEIOPERIS, Rafin. Fl. Tellur. iv. 103 (1836) = **Passiflora**, Linn.
angustifolia, Rafin. l. c. = P. suberosa.
hederacea, Rafin. l. c. = P. suberosa.
minima, Rafin. l. c. = P. suberosa *vel* serrulata.
multiflora, Rafin. l. c. = P. multiflora.
pallida, Rafin. l. c. = P. suberosa.
pannosa, Rafin. l. c. = P. pannosa.
peltata, Rafin. l. c. = P. suberosa.
suberosa, Rafin. l. c. = P. suberosa.

MEIOSPERMA, Rafin. Fl. Tellur. iv. 64 (1836) = **Justicia**, Linn. (Acanth.).
debilis, Rafin. l. c. = J. debilis.

MEIRACYLLIUM, Reichb. f. Xenia Orch. i. 12. t. 6. (1854). *ORCHIDEAE*, Benth. & Hook. f. iii. 493.
Gemma, *Reichb. f. in Gard. Chron.* (1869) 988; *Xenia Orch.* iii. 209.—Mexic.
trinasutum, *Reichb. f. Xenia Orch.* i. 12; *et in Bonplandia*, iv. (1856) 214.—Am. austr.
Wendlandi, *Reichb. f. Orch. Centr. Am.* 73.—Guatemala.

MEISNERIA, DC. Prod. iii. 114 (1828) = **Siphanthera**, Pohl (Melastom.).
arenaria, DC. l. c. = S. arenaria.
cordata, Triana, in Trans. Linn. Soc. xxviii. (1871) 32 = S. cordata.
cordifolia, Benth. in Hook. Journ. Bot. ii. (1840) 299 = S. microlicioides.
microlicioides, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 204 = S. microlicioides.
paludosa, DC. Prod. iii. 114 = S. paludosa.
subtilis, Triana, in Trans. Linn. Soc. xxviii. (1871) 32 = S. subtilis.
tenera, Triana, l. c. = S. tenera, Hostmannii.

MEISSARRHENA, R. Br. in Salt, Abyss. App. 63 (1841) = **Anticharis**, Endl. (Scrophul.).
tomentosa, R. Br. l. c. = A. arabica.

MEISTERA, Giseke, Prael. Linn. 199, 205 (1792) = **Amomum**, Linn. (Scitam.).
Amomum, Raeusch. Nom. ed. III. i. (Quid ?).—Ind. or.

MEISTERIA, Scop. Introd. 124 (1777) = **Poraqueiba**, Aubl. (Olacin.).
anonyma, J. F. Gmel. Syst. 391 = P. guianensis.

MEISTERIA, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 127 = **Enkianthus**, Lour. (Ericac.).
cernua, Sieb. & Zucc. l. c.—Japon.

MEIZOTROPIS, Voigt, Hort. Suburb. Calc. 239 (1845) = **Butea**, Koen. (Legumin.).
buteaeformis, Voigt, l. c. = B. minor.

MELACHNE, Schrad. ex Schult. f. Syst. vii. II. p. lxxxv (1830) = **Gahnia**, Forst. (Cyperac.).
Sieberi, Schrad. ex Schult. f. l. c. = G. tristis.

MELADENDRON, Molina, Sagg. Chile, ed. II. 143 (1810) (Solanac.) (Quid ?).
chilensis, Molina, l. c.—Chili.

MELADENDRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 64, 130 = **Melaleuca**, Linn. (Myrtac.).
leucocladum, St. Lag. l. c. = Melaleuca Leucadendron.

MELADENIA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 576 = **Psoralea**, Linn. (Legumin.).
densiflora, Turcz. l. c. = P. badocana.

MELAENACRANIS, Roem. & Schult. Syst. ii. 6, 236 (1817) = MELANCRANIS, Vahl = **Ficinia**, Schrad. (Cyperac.).

MELALEMA, Hook. f. Fl. Antarct. ii. 311 (1846). *COMPOSITAE*, Benth. & Hook. f. ii. 443.
humifusa, *Hook. f. Fl. Antarct.* ii. 311.—Reg. Magell.

MELALEUCA, Linn. Mant. i. 14 (1767). *MYRTACEAE*, Benth. & Hook. f. i. 705.
ASTEROMYRTUS, Schau. in Linnaea, xvii. (1843) 242.
GYMNAGATHIS, Schau. l. c. (1843) 243.
MELADENDRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 64, 130.
MELANOLEUCE, St. Lag. l. c. viii. (1881) 203.
abietina, Sm. in Rees, Cycl. xxiii. n. 15 = cuticularis.
acacoides, *F. Muell. Fragm.* iii. 116.—Austral.
acerosa, *Schau. in Lehm. Pl. Preiss.* i. 137.—Austral.
acuminata, *F. Muell. Fragm.* i. 15.—Austral.
adnata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 343.—Austral.
aestuosa, *Forst. f. Prod.* 38.—Ins. Tahiti.
affinis, *Regel & Koern. in Ind. Sem. Hort. Petrop.* (1857) 56.—Austral.
alba, Hort. ex Steud. Nom. ed. I. 515 = seq.?
albiflora, Raeusch. Nom. ed. III. 142 = Metrosideros albiflora.
alsophila, *A. Cunn. ex Benth. Fl. Austral.* iii. 137.—Austral.
angulata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 342.—Austral.
angustifolia, Blume, Mus. Bot. Lugd. Bat. i. 83 = M. Leucadendron.
angustifolia, *Gaertn. Fruct.* i. 172. t. 35.—Austral.
apodocephala, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 340.—Austral.
arachnoidea, Raeusch. Nom. ed. III. 143 = Leptospermum arachnoideum.

MELALEUCA:—
armillaris, *Sm. in Trans. Linn. Soc.* iii. (1797) 277.—Austral.
armillaris, Wendl. ex Steud. Nom. ed. I. 515 = taxifolia.
aspalathoides, *Schau. in Lehm. Pl. Preiss.* i. 140.—Austral.
axillaris, Hort. ex Steud. Nom. ed. II. ii. 111 = ericifolia.
Baxteri, *Benth. Fl. Austral.* iii. 138.—Austral.
basicephala, *Benth. l. c.* 133.—Austral.
bicolor, Poir. Encyc. Suppl. iii. 624 = Memecylon umbellatum.
bisulcata, F. Muell. Fragm. iii. 118 = acerosa.
blaeriaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 165.—Austral.
brachyphylla, Schau. in Otto & Dietr. Allg. Gartenz. iii. (1835) 162 = microphylla.
brachystachya, *F. Muell. Fragm.* iii. 119.—Austral.
bracteata, F. Muell. l. c. i. 15 = genistifolia.
bracteosa, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 340.—Austral.
brevifolia, F. Muell. Fragm. i. 116 = microphylla.
brevifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 342.—Austral.
Cajaputi, Roxb. Hort. Beng. 59; Fl. Ind. iii. 394 = M. Leucadendron.
Caja-putti, Hort. ex DC. Prod. iii. 215 = squarrosa.
callistemonea, Lindl. Swan River App. 8 = lateritia.
calothamnoides, *F. Muell. Fragm.* iii. 114.—Austral.
calycina, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 416.—Austral.
calycina, Steud. Nom. ed. I. 515 = thymifolia.
canaliculata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 342.—Austral.
canescens, Otto, Hort. Berol. 37; ex Link & Otto, Pl. Sel. Hort. Berol. 81. t. 37 = incana.
cardiophylla, *F. Muell. Fragm.* i. 225.—Austral.
carinata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 344.—Austral.
chlorantha, Bonpl. Desc. Pl. Rar. Malm. 22. t. 8 = diosmifolia.
ciliata, Forst. f. Prod. 38 = Metrosideros ciliata.
ciliosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxv. (1862) II. 326.—Austral.
citrina, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 341.—Austral.
commutata, Miq. Anal. Bot. Ind. i. 14 = M. Leucadendron.
concinna, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 339.—Austral.
concreta, *F. Muell. Fragm.* iii. 118.—Austral.
conferta, *Benth. Fl. Austral.* iii. 142.—Austral.
conferta, Steud. Nom. ed. I. 515 = Tristania conferta.
cordata, *Benth. Fl. Austral.* iii. 149.—Austral.
cordata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 339 (an praec. ?).—Austral.
coronata, Andr. Bot. Rep. t. 278 = thymifolia.
costata, Raeusch. Nom. ed. III. 142 = Angophora lanceolata.
crassifolia, *Benth. Fl. Austral.* iii. 145.—Austral.
cucullata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 343.—Austral.
Cumingiana, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 164 = M. Leucadendron.
cuneata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 339.—Austral.
Cunninghami, Schau. ex Walp. Rep. ii. 927 = M. Leucadendron.
cupressina, F. Muell. Fragm. iii. 114 = thyoides.
curvifolia, Schlecht. in Linnaea, xx. (1847) 654 = Preissiana.
cuspidata, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 327 = Conothamnus trinervis.
cuticularis, *Labill. Nov. Holl. Pl.* ii. 30. t. 171.—Austral.
cylindrica, R. Br. ex Benth. Fl. Austral. iii. 146 = armillaris.
cymbifolia, *Benth. l. c.* 148.—Austral.
cyrtodonta, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 24 = Beaufortia cyrtodonta.
decurrens, Wall. Cat. n. 3649 = Tristania merguensis.
decussata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 415.—Austral.
deltoidea, *Benth. Fl. Austral.* iii. 161.—Austral.
densa, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 416—Austral.
densa, Colla, Hort. Ripul. 87 = pulchella.

MELALEUCA:—
depauperata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 343.—Austral.
diffusa, Forst. f. Prod. 37 = Metrosideros diffusa.
diosmatifolia, Dum.-Cours. Bot. Cult. ed. II. v. 373 = ericifolia.
diosmifolia, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 410.—Austral.
discolor, Reichb. Ic. Bot. Exot. ii. 5. t. 113 = thymifolia.
dissitiflora, *F. Muell. Fragm.* iii. 153.—Austral.
divaricata, Donn, ex Steud. Nom. ed. I. 515 (Quid ?).—Hab. ?
divaricata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 344.—Austral.
Drummondii, Schau. in Lehm. Pl. Preiss. i. 138 = uncinata.
elachophylla, *F. Muell. Fragm.* iii. 120.—Austral.
elegans, Hornsch. in Otto & Dietr. Allg. Gartenz. ii. (1834) 329 = decussata?
eleuterostachya, *F. Muell. Fragm.* iii. 117.—Austral.
elliptica, *Labill. Nov. Holl. Pl.* ii. 31. t. 173.—Austral.
empetrifolia, Reichb. Ic. Bot. Exot. ii. 1. t. 102 = Beaufortia micrantha.
Endlicheriana, Schau. in Lehm. Pl. Preiss. i. 134 = seriata.
epacridioides, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 165 = densa.
eremaea, F. Muell. Fragm. iii. 114 = trichophylla.
eriantha, *Benth. Fl. Austral.* iii. 153.—Austral.
ericaefolia, Andr. Bot. Rep. t. 175 = armillaris.
ericifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 276.—Austral.
eriocephala, Sieber, in Spreng. Syst. iii. 836 = Kunzea capitata.
erubescens, Otto, Hort. Berol. 37 = ericifolia.
erucaeformis, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 344.—Austral.
eugeniifolia, Wall. Cat. n. 3648 = Tristania merguensis.
exarata, *F. Muell. Fragm.* iii. 114.—Austral.
fascicularis, Labill. Nov. Holl. Pl. ii. 29. t. 170 = Astartea fascicularis.
fasciculiflora, *Benth. Fl. Austral.* iii. 137.—Austral.
ferrea, *A. Cunn. ex A. Gray, Bot. U. St. Expl. Exped.* i. 566.—Austral.
filifolia, *F. Muell. Fragm.* iii. 119.—Austral.
fimbriata, Donn, Hort. Cantab. ed. VI. 207 = thymifolia.
florida, Forst. f. Prod. 37 = Metrosideros diffusa.
foliolosa, *A. Cunn. ex Benth. Fl. Austral.* iii. 161.—Austral.
foliosa, Dum.-Cours. Bot. Cult. ed. II. v. 373 = diosmifolia.
Fraseri, Hook. Bot. Mag. t. 3210 = striata.
fulgens, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 415.—Austral.
genistifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 277.—Austral.
gibbosa, *Labill. Nov. Holl. Pl.* ii. 30. t. 172.—Austral.
glaberrima, *F. Muell. Fragm.* iii. 119.—Austral.
globifera, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 410.—Austral.
glomerata, *F. Muell. Rep. Babbage Exp.* 8.—Austral.
gnidiaefolia, Vent. Jard. Malm. 4 = thymifolia.
gnidioides, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. ii. (1864) 139.—N. Caled.
grandiflora, Blanco, Fl. Filip. ed. I. 615 = Bombax malabaricum.
grandis, *Fisch. ex Steud. Nom.* ed. II. ii. 112.—Austral.
gummifera, Steud. Nom. ed. I. 515 = Eucalyptus resinifera.
Gunniana, Schau. in Walp. Rep. ii. 928 = ericifolia.
hakeacea, F. Muell. Fragm. iii. 117 = teretifolia.
hakeoides, *F. Muell. ex Benth. Fl. Austral.* iii. 151.—Austral.
halmaturorum, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 122 = pustulata.
hamata, Field. & Gardn. Sert. Pl. t. 74 = uncinata.
hamulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 165.—Austral.
heliophila, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 120 = ericifolia.
hispida, *Wendl. ex DC. Prod.* iii. 215.—Austral.
holosericea, *Schau. in Lehm. Pl. Preiss.* i. 139.—Austral.
Huegelii, *Endl. in Enum. Pl. Hueg.* 48.—Austral.
hypericifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 279.—Austral.
hypochondriaca, Dehnh. Revista Napolit. i. III. 177; ex Walp. Rep. ii. 163 = incana.

MELALEUCA:—
imbricata, *Hort. ex Link, Enum. Hort. Berol.* ii. 272.—Austral.
incana, *R. Br. in Bot. Reg.* t. 410.—Austral.
juniperina, Sieber, ex Reichb. Ic. Exot. Bot. ii. 4. t. 112 = nodosa.
juniperoides, DC. Prod. iii. 213 = nodosa.
lanata, Nois. ex Steud. Nom. ed. I. 515 = incana.
lanceolata, R. Br. ex Benth. Fl. Austral. iii. 143 = M. Leucadendron.
lanceolata, Otto, Hort. Berol. 36 = genistifolia.
lancifolia, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 164 = M. Leucadendron.
lanigera, Wendl. ex DC. Prod, iii. 215 = incana?
lasiandra, *F. Muell. Fragm.* iii. 115.—Austral.
lateralis, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 339.—Austral.
laterifolia, *Benth. Fl. Austral.* iii. 136.—Austral.
lateritia, *Otto & Dietr. Allg. Gartenz.* ii. (1834) 257.—Austral.
latifolia, Raeusch. Nom. ed. III. 142 = M. Leucadendron.
laurina, Sm. in Trans. Linn. Soc. iii. (1797) 275 = Tristania laurina.
laxiflora, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 341.—Austral.
Lehmannii, Schau. in Lehm. Pl. Preiss. i. 142 = viminea.
leiocarpa, *F. Muell. Fragm.* x. 55.—Austral.
leiopyxis, *F. Muell. ex Benth. Fl. Austral.* iii. 160.—Austral.
leptoclada, *Benth. l. c.* 132.—Austral.
leptospermoides, Schau. in Lehm. Pl. Preiss. i. 139 = ciliosa.
Leucadendron, *Linn. Mant.* i. 105.—Austral.
linariifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 278.—Austral.
linearis, Wendl. & Schrad. Sert. Hannov. 19. t. 11 = Callistemon linearis.
linophylla, *F. Muell. Fragm.* iii. 115.—Austral.
longicoma, *Benth. Fl. Austral.* iii. 129.—Austral.
lucida, Forst. f. Prod. 38 = Metrosideros lucida.
lucida, Linn. f. Suppl. 342 = Metrosideros diffusa.
lucidula, Hort. ex DC. Prod. iii. 213 = thymoides?
macronychia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 340.—Austral.
Manglesii, Schau. in Lehm. Pl. Preiss. i. 135 = scabra.
megacephala, *F. Muell. Fragm.* iii. 117.—Austral.
micromera, *Schau. in Lehm. Pl. Preiss.* i. 146.—Austral.
microphylla, *Sm. in Rees, Cycl.* xxiii. n. 9.—Austral.
mimosoides, A. Cunn. ex Schau. in Walp. Rep. ii. 927 = M. Leucadendron.
minor, Sm. in Rees, Cycl. xxiii. n. 2 = M. Leucadendron.
minutifolia, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 45.—Austral.
muricata, Hort. ex DC. Prod. iii. 213 = scabra.
myrtifolia, Hort. ex Steud. Nom. ed. I. 515 = hypericifolia.
myrtifolia, Vent. Jard. Malm. 47 = squarrosa.
nematophylla, F. Muell. Fragm. iii. 119 = filifolia.
neriifolia, Sims, Bot. Mag. t. 1058 = Tristania neriifolia.
nervosa, Hort. ex Link, Enum. Hort. Berol. ii. 25 = Leptospermum flavescens.
nesophila, *F. Muell. Fragm.* iii. 113.—Austral.
nodosa, Link, Enum. Hort. Berol. ii. 273 = ericifolia.
nodosa, *Sm. in Trans. Linn. Soc.* iii. (1797) 276.—Austral.
nummularia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 341.—Austral.
obliqua, Hort. ex Steud. Nom. ed. I. 515 = stypheliioides.
Oldfieldii, *F. Muell. Fragm.* iii. 118.—Austral.
oligantha, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 123 = decussata.
ornata, Schau. in Lehm. Pl. Preiss. i. 135 = seriata.
Ottonis, Schau. in Otto & Dietr. Allg. Gartenz. iii. (1835) 167 = squamea.
ovatifolia, Poir. Encyc. Suppl. iii. 624 = Memecylon angulatum.
paludosa, R. Br. in Ait. Hort. Kew. ed. II. iv. 410 = Callistemon speciosus.
paludosa, Schlecht. in Linnaea, xx. (1847) 653 = Callistemon salignus.
parviceps, Lindl. Swan River App. 8 = scabra.
parviflora, Lindl. l. c. = Preissiana.
parviflora, Link, Enum. Hort. Berol. ii. 272, sphalm. = *parvifolia*.

MELALEUCA:—
parviflora, Reichb. Ic. Bot. Exot. t. 31 = decussata.
parvifolia, Otto, Hort. Berol. 37 = thymifolia.
pauciflora, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 166.—Austral.
pauperiflora, *F. Muell. Fragm.* iii. 116.—Austral.
pendulina, *Lodd. ex G. Don, in Loud. Hort. Brit.* 319.—Austral.
pentagona, *Labill. Nov. Holl. Pl.* ii. 27. t. 166.—Austral.
perforata, Forst. f. Prod. 37 = Metrosideros perforata.
pinifolia, *Hort. ex Colla, Hort. Ripul. App.* ii. 352.—Austral.
pinifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 166 (an praec. ?).—Austral.
pithyoides, F. Muell. ex Benth. Fl. Austral. iii. 121 = Callistemon salignus.
polycephala, *Benth. Fl. Austral.* iii. 152.—Austral.
polygaloides, *Schau. in Lehm. Pl. Preiss.* i. 142.—Austral.
polygonoides, Hoffmgg. Verz. Pfl. 78 = genistifolia.
Preissiana, *Schau. in Lehm. Pl. Preiss.* i. 143.—Austral.
propinqua, Schau. in Otto & Dietr. Allg. Gartenz. iii. (1835) 166 = densa.
pubescens, Schau. in Walp. Rep. ii. 928 = Preissiana.
pulchella, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 414.—Austral.
pumila, Otto, ex DC. Prod. iii. 214 = decussata.
pungens, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. ii. (1864) 139.—N. Caled.
pungens, *Schau. in Lehm. Pl. Preiss.* i. 138.—Austral.
pustulata, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 476.—Austral.
Radula, *Lindl. Swan River App.* 8.—Austral.
Regelii, Planch. Hort. Donat. 88. App. [227] = Beaufortia micrantha ?
rhaphiophylla, *Schau. in Lehm. Pl. Preiss.* i. 143.—Austral.
rigida, *Roxb. Hort. Beng.* [100]; *Fl. Ind.* iii. 399.—Ins. Mollucc.
rigidifolia, *Turcz. in Bull. Phys.-Math. Akad. Pétersb.* x. (1852) 342.—Austral.
salicifolia, Andr. Bot. Rep. t. 485 = Tristania neriifolia.
saligna, Schau. in Walp. Rep. ii. 927; Blume, Mus. Bot. Lugd. Bat. i. 66 = M. Leucadendron.
scabra, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 414.—Austral.
scandens, Raeusch. Nom. ed. III. 142 = Metrosideros florida.
scoparia, Forst. f. Prod. 37 = Leptospermum scoparium.
scoparia, *Wendl. Sert. Hannov.* 25. t. 15.—N. Zel.
semiteres, Schau. in Lehm. Pl. Preiss. i. 143 = uncinata.
seriata, *Lindl. Swan River App.* 8.—Austral.
serpyllifolia, Dum.-Cours. Bot. Cult. ed. II. v. 374 = pulchella.
serpyllifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 339.—Austral.
Sieberi, Schau. in Walp. Rep. ii. 928 = M. Leucadendron.
sparsiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 167.—Austral.
spathulata, *Schau. in Lehm. Pl. Preiss.* i. 134.—Austral.
spectabilis, Raeusch. Nom. ed. III. 142 = Metrosideros florida.
spinosa, Lindl. Swan River App. 8 = thymoides.
splendens, Lee, ex Kerner, Hort. t. 639 = fulgens.
sprengelioides, DC. Prod. iii. 215 = Beaufortia Dampieri.
squamea, *Labill. Nov. Holl. Pl.* ii. 28. t. 158.—Austral.
squarrosa, *Donn, Hort. Cantab.* ed. II. 101; *Sm. in Trans. Linn. Soc.* vi. (1802) 300.—Austral.
striata, *Labill. Nov. Holl. Pl.* ii. 26. t. 165.—Austral.
striata, Walp. Rep. ii. 162, sphalm. = seriata.
stricta, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 375 = linariifolia.
styphelioides, *Sm. in Trans. Linn. Soc.* iii. (1797) 275.—Austral.
suaveolens, Soland. ex Gaertn. Fruct. i. 173. t. 35 = Tristania suaveolens.
subfalcata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 341.—Austral.
subtrigona, *Schau. in Lehm. Pl. Preiss.* i. 139.—Austral.
symphyocarpa, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 44.—Austral.
tamariscina, *Hook. in Mitch. Journ. Trop. Austral.* 262.—Austral.

MELALEUCA:—
taxifolia, *Schlecht. ex Spreng. Syst.* iii. 336.—Austral.
tenella, *Benth. Fl. Austral.* iii. 160.—Austral.
tenuifolia, DC. Prod. iii. 213 = nodosa.
tenuifolia, Wendl. Bot. Beobacht. 50 = Leptospermum scoparium.
tenuissima, Tausch, in Flora, xix. (1836) 411 = microphylla.
teretifolia, *Endl. in Enum. Pl. Hueg.* 49.—Austral.
ternifolia, *F. Muell. ex Miq. in Nederl. Kruidk. Arch.* iv. (1859) 123.—Austral.
tetragona, Lodd. ex Otto, Hort. Berol. 37; DC. Prod. iii. 215 = decussata.
Thea, Wendl. Sert. Hannov. 24. t. 13 = Leptospermum flavescens.
thymifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 278.—Austral.
thymoides, *Labill. Nov. Holl. Pl.* ii. 27. t. 167.—Austral.
thyoides, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 167.—Austral.
tomentosa, Colla, Hort. Ripul. 87. t. 37 = incana.
trichophylla, *Lindl. Swan River App.* 8.—Austral.
trichostachya, Lindl. in Mitch. Journ. Trop. Austral. 277 = linariifolia.
trinervia, White, Journ. Voy. N. S. Wales, App. 229. t. 24 = Leptospermum flavescens *vel* pubescens.
tuberculata, Schau. in Lehm. Pl. Preiss. i. 139 = subtrigona.
uncinata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 414.—Austral.
undulata, *Benth. Fl. Austral.* iii. 135.—Austral.
uniflora, Hort. ex Steud. Nom. ed. II. ii. 112 (Quid ?).—Austral.
urceolaris, *F. Muell. ex Benth. Fl. Austral.* iii. 154.—Austral.
villosa, *Linn. f. Suppl.* 342.—Ins. Tahiti.
viminea, *Lindl. Swan River App.* 8.—Austral.
violacea, *Schau. in Lehm. Pl. Preiss.* i. 146.—Austral.
virgata, Linn. f. Suppl. 343 = Baeckea virgata.
viridiflora, Soland. ex Gaertn. Fruct. i. 175. t. 35 = M. Leucadendron.
Waeberi, Reichb. ex Schau. in Otto & Dietr. Allg. Gartenz. iii. (1835) 167 = seriata.
Wilsonii, *F. Muell. Fragm.* ii. 124. t. 15.—Austral.

MELALEUCON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 64 = praec.

MELAMPIRUM, Neck. Delic. Gallo-Belg. i. 264 (1768) = **Melampyrum**, Tourn. (Scrophul.).

MELAMPODIUM, Linn. Coroll. Gen. 17 (1737).
COMPOSITAE, Benth. & Hook. f. ii. 348.
ALCINA, Cav. Ic. i. 10. t. 15 (1791).
CAMUTIA, Bonato, ex Steud. Nom. ed. II. i. 273 (1840).
CARGILLA, Adans. Fam. ii. 130 (1763).
DYSODIUM, Rich. in Pers. Syn. ii. 489 (1807).
PRONACRON, Cass. in Dict. Sc. Nat. xliii. 370 (1826).
UNXIA, Linn. f. Suppl. 56 (1781).
ZARABELLIA, Cass. in Dict. Sc. Nat. lix. 240 (1829).
achillaeoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 145.—Mexic.
americanum, *Linn. Sp. Pl.* 921.—Mexic.
angustifolium, *DC. Prod.* v. 519.—Péruv.
australe, Loefl. It. Hisp. 268 = Acanthospermum brasilum.
Baranquillae, *Spreng. Syst.* iii. 619.—N. Granat.
Berterianum, *Spreng. l. c.*—Ind. occ.
camphoratum, *Benth. & Hook. f. Gen.* ii. 349; *Baker, in Mart. Fl. Bras.* vi. III. 161.—Bras.
cinereum, *DC. Prod.* v. 518.—Am. bor. occ.
connatum, [*Moç. & Sesse,*] *ex DC. Prod.* v. 521.—Mexic.
coronopifolium, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 145.—Mexic.
cupulatum, *A. Gray, in Proc. Am. Acad.* viii. (1870) 291.—Mexic.
diffusum, *Cass. in Dict. Sc. Nat.* lix. 238.—Ins. Philipp.
digynum, *Benth. & Hook. f. Gen.* ii. 349.—Panama.
divaricatum, DC. Prod. v. 520 = paludosum.
Dombeyanum, *DC. l. c.* 521.—Peruv.
flaccidum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 86.—Nicaragua.

MELAMPODIUM:—
gracile, *Less. in Linnaea*, vi. (1831) 407.—Mexic.
heterophyllum, *Lag. Gen. et Sp. Nov.* 33.—Mexic.
Hidalgoa, *DC. Prod.* v. 521.—Mexic.
hirsutum, Benth. & Hook. f. Gen. ii. 349 = camphoratum.
hispidum, *H. B. & K. Nov. Gen. et Sp.* iv. 273. t. 399.—Arizona.
humile, Sw. Prod. Veg. Ind. Occ. 114 = Acanthospermum humile.
Kunthianum, *DC. Prod.* v. 519.—Mexic.
lanceolatum, *DC. l. c.*—Am. austr.
leucanthum, Torr. & Gray, Fl. N. Am. ii. 271 = cinereum.
linearilobium, *DC. Prod.* v. 518.—Mexic.
longicorne, *A. Gray, in Mem. Am. Acad.* N. S. v.(1855) 321.—Arizona.
longifolium, Brouss. ex Willd. Enum. Hort. Berol. 934 = seq. ?
longifolium, *Cerv. ex Cav. in Anal. Cienc. Nat.* vi. (1803) 333.—Mexic.
manillense, Less. in Linnaea, v. (1830) 155. t. 2 = diffusum.
microcephalum, *Less. l. c.* ix. (1834) 268.—Mexic.
montanum, *Benth. Pl. Hartw.* 64.—Mexic.
oblongifolium, *DC. Prod.* v. 519.—Mexic.
ovatifolium, Reichb. Ic. Exot. t. 42 = paludosum.
paludosum, *H. B. & K. Nov. Gen. et Sp.* iv. 273.—Am. austr.
paniculatum, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 287.—Bras.
perfoliatum, *H. B. & K. Nov. Gen. et Sp.* iv. 274.—Mexic.
pumilum, *Benth. Pl. Hartw.* 64.—Mexic.
ramosissimum, DC. Prod. v. 518 = cinereum.
rhomboideum, *DC. l. c.* 520.—Am. austr.
ruderale, Sw. Fl. Ind. Occ. iii. 1372 = Eleutheranthera ovata.
sericeum, H. B. & K. Nov. Gen. et Sp. iv. t. 398 = Kunthianum.
sericeum, *Lag. Elench. Hort. Madr.* (1805); *et Gen. et Sp. Nov.* 32.—Mexic.
tenellum, *Hook. & Arn. Bot. Beech. Voy.* 299.—Mexic.
ternatum, DC. Prod. v. 521 = Hidalgoa ternata.

MELAMPYRUM, [Tourn.] Linn. Syst. ed. I (1735).
SCROPHULARINEAE, Benth. & Hook. f. ii. 979.
albicans, Porc. in Magyar Növényt. Lapok, ix. (1885) 130 = arvense.
alpestre, Pers. Syn. ii. 151 = sylvaticum.
americanum, *Michx. Fl. Am. Bor.* ii. 16.—Am. bor.
angustissimum, *G. Beck, in Verh. Zool. Bot. Ges. Wien*, xxxii. (1883) 187.—Austria.
argyrocomum, Fisch. ex Steud. Nom. ed. II. ii. 113 = arvense.
arvense, *Linn. Sp. Pl.* 605.—Europ.; Reg. Caucas.
barbatum, Ledeb. Fl. Ross. iii. 305 = caucasicum.
barbatum, Schur, in Verh. Siebenb. Ver. Naturf. x. (1859) 61 = arvense.
barbatum, *Waldst. & Kit. Pl. Rar. Hung.* i. 89. t. 86.—Europ. austr.
bihariense, *A. Kern. Sched. Fl. Exsicc.* i. 35.—Austria.
bohemicum, *A. Kern. l. c.*—Europ.
brachiatum, Schwein. in Long, Exped. Winnep. ii. 391 = americanum.
caerulescens, Gilib. Fl. Lituan. i. 131 = nemorosum.
caeruleum, Gueldenst. Reisen, i. 424 = nemorosum.
carpaticum, *Schult. Obs. Bot.* 116.—Carpath.
catalaunicum, *Freyn, in Flora*, lxvii. (1884) 680.—Hispan.
caucasicum, *Bunge, Pl. Abich.* 16.—Reg. Caucas.
ciliare, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 122 = roseum.
ciliatum, Boiss. & Heldr. Diagn. Ser. II. iii. 176 = barbatum.
commutatum, Tausch, ex Koch, Syn. Fl. Germ. ed. II. 621 = pratense.
cretaceum, *Czern. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 79, nomen.—Rossia.
cristatum, Habl. ex Steud. Nom. ed. II. ii. 113 = arvense.
cristatum, *Linn. Sp. Pl.* 605.—Europ.; As. bor.
dentatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 56 = sylvaticum.
digitatum, Schur, Enum. Pl. Transs. 506 = pratense.
elatius, Reut. ex Boiss. Fl. Orient. iv. 481 = arvense.
grandiflorum, *A. Kern. Sched. Fl. Exsicc.* ii. 114.—Austria.

MELAMPYRUM:—
hastatum, Gilib. Fl. Lituan. i. 131 = pratense.
heracleoticum, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 482 = pratense.
hians, Gilib. Exercit. i. 132 = sylvaticum.
hyans, Gilib. l. c. 131 = praec.
hybridum, Wolfn. in Oestr. Bot. Wochenschr. (1857) 232 = arvense.
indicum, *Hook. f. & Thoms. in Hook. Kew Journ.* ix. (1857) 246. t. 8.—Reg. Himal.
jedoënse, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 122 = roseum.
laciniatum, *Kosh. & Zinger, in Bull. Soc. Nat. Mosc.* lvi. (1881) i. 313.—Rossia.
laricetorum, *Kern. ex Dalla Torre, Atl. Alpenfl.* 175. —Europ. alp.
latifolium, Muhl. Cat.; ex Nutt. Gen. Am. ii. 58 = americanum.
laxum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 123.—Japon.
lineare, Lam. Encyc. iv. 22 = americanum.
lineare, Saut. ex Nym. Consp. 556 = pratense.
montanum, Johnst. Fl. Berwick, i. 136 = pratense.
moravicum, *H. Braun, in Oestr. Bot. Zeitschr.* xxxiv. (1884) 422.—Austr.
nemorosum, Baumg. Enum. Stirp. Transs. ii. 198 = bihariense.
nemorosum, *Linn. Sp. Pl.* 605.—Europ.; Reg. Caucas.
pratense, *Linn. l. c.*—Europ.; As. bor.
pratense, Tausch, ex Koch, Syn. Fl. Germ. ed. II. 621 = sylvaticum.
Pseudo-arvense, Schur, Enum. Pl. Transs. 506 = barbatum.
Pseudo-barbatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 56 = arvense.
Pseudo-sylvaticum, Schur, Enum. Pl. Transs. 506 = pratense.
purpurescens, Gilib. Fl. Lituan. i. 130 = arvense.
roseum, *Maxim. Prim. Fl. Amur.* 210.—Manchuria.
saxosum, Baumg. Enum. Stirp. Transs. ii. 199 = sylvaticum.
stenophyllum, *Boiss. Fl. Orient.* iv. 482.—As. Min.
stenotaton, Wiesb. in Oestr. Bot. Zeitschr. xxxiv. (1884) 269 = bihariense.
subalpinum, Kern. l. c. xiii. (1863) 363 = bihariense.
sylvaticum, Hook. Fl. Bor. Am. ii. 106 = americanum.
sylvaticum, C. Koch, in Linnaea, xxiii. (1850) 679 = stenophyllum.
sylvaticum, *Linn. Sp. Pl.* 605.—Europ.; As. bor.
transsilvanicum, Schur, Enum. Pl. Transs. 507 = sylvaticum.
violaceum, Lam. Fl. Fr. ii. 356 = nemorosum.
vulgatum, Pers. Syn. ii. 151 = pratense.

MELANACRANIS, Reichb. Consp. 56 (1828) = MELANCRANIS, Vahl = **Ficinia**, Schrad. (Cyperac.).

MELANANTHERA, Michx. Fl. Bor. Am. ii. 106 (1803) = **Melanthera**, Rohr (Compos.).

MELANANTHUS, Walp. in Bot. Zeit. viii. (1850) 788. *SOLANACEAE* [Benth. & Hook. f. ii. 1137]. MICROSCHWENKIA, Benth. ex Hemsl. Biol. Centr. Am. Bot. ii. 438 (1882).
dipyrenoides, *Walp. in Bot. Zeit.* viii. (1850) 788.—Bras.

MELANCHRYSUM, Cass. in Bull. Soc. Philom. (1817) 12 = **Gazania**, Gaertn. (Compos.).
Pavonium, Cass. ex DC. Prod. vi. 510 = G. Pavonia.
rigens, Cass. in Dict. Sc. Nat. xxix. 443 = G. rigens.
spinulosum, Cass. l. c. = G. pinnata.

MELANCIUM, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 175. *CUCURBITACEAE*, Benth. & Hook. f. i. 833.
campestre, *Naud. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 175.—Bras.

MELANCRANIS, Vahl, Enum. ii. 239 (1806) = **Ficinia**, Schrad. (Cyperac.).
gracilis, Nees, in Linnaea, ix. (1834) 287 = F. bracteata.
nigrescens, Schrad. in Goett. Gel. Anz. (1821) iii. 2072 = F. bracteata.
pallida, Nees, ex Kunth, Enum. Pl. ii. 257 = F. setiformis.
radiata, Kunth, Enum. Pl. ii. 260 = F. brevifolia.
radiata, Vahl, Enum. ii. 239 (Gen. ?).—Afr. austr.

MELANCRANIS:—
rigidula, Nees, in Linnaea, x. (1836) 141 = Ficinia bracteata.
rubiginosa, Spreng. Syst. i. 236 = Cladium glomeratum.
scariosa, Vahl, Enum. ii. 239 = F. bracteata.
tenuis, Eckl. ex Kunth, Enum. Pl. ii. 265.—Afr. austr.

MELANDRIUM, Roehl. Deutschl. Fl. ed. II. 274 (1812) = **Lychnis**, Linn. (Caryoph.).
affine, J. Vahl, in Fl. Dan. xiv. 5 = L. affinis.
ajanense, Rohrb. in Linnaea, xxxvi. (1869–70) 230 = L. ajanensis.
album, Garcke, Fl. Deutschl. ed. IV. 55 = L. alba.
Allamani, Rohrb. Monog. Silene, 231 = Silene laciniata.
andicolum, Rohrb. in Linnaea, xxxvi. (1869–70) 221 = Saponaria andicola.
angustiflorum, Walp. Ann. i. 93 = L. affinis.
apetalum, Fenzl, in Ledeb. Fl. Ross. i. 326 = L. apetala.
apricum, Rohrb. Monog. Silene, 231 = Silene aprica.
auriculatum, Rohrb. l. c. = Silene auriculata.
Baldwini, Rohrb. l. c. 231, 232 = Silene Baldwini.
bellidioides, Rohrb. in Linnaea, xxxvi. (1869–70) 247 = Silene bellidioides.
Bolanderi, Rohrb. Monog. Silene, 232 = Silene Hookeri.
brachypetalum, Fenzl, in Ledeb. Fl. Ross. i. 326 = L. brachypetala.
cabulicum, Boiss. Diagn. Ser. II. i. 79.—Afghan.
caffrum, Rohrb. Monog. Silene, 232 = Silene caffra.
californicum, Rohrb. in Linnaea, xxxvi. (1869–70) 252 = Silene californica.
capense, Rohrb. Monog. Silene, 232 = Silene capensis.
cashmerianum, Walp. Rep. i. 280 = L. cachemiriana.
caspum, Gruner, in Bull. Soc. Nat. Mosc. x. (1867) ii. 407.—Reg. Casp.
chilense, Rohrb. in Linnaea, xxxvi. (1869–70) 222 = L. chilensis.
cucubaloides, Fenzl, ex Rohrb. l. c. 223.—Chili; Reg. Argent.
cuneifolium, Walp. Rep. i. 280 = L. cuneifolia.
dicline, Willk. Ic. Desc. Pl. Hisp. 27. t. 14 = L. diclinis.
dioicum, Coss. & Germ. Fl. Par. ed. I. 28; ed. II. 37 = Silene conoidea.
diurnum, Fries, in Bot. Notiser (1842), 170 = L. sylvestris.
divaricatum, Fenzl, in Ledeb. Fl. Ross. i. 328 = L. divaricata.
dubium, Hampe, ex Garcke, Fl. N. M. Deutschl. ed. VIII. 66.—Europ.
Echegarayi, Hieron. in Bol. Acad. Nac. Córdova, iv. (1881) 11.—Reg. Argent.
Ecklonianum, Rohrb. Monog. Silene, 232 = Silene Eckloniana.
Elisabethae, Rohrb. l. c. = Silene Elisabethae.
eriocalycinum, Boiss. Diagn. Ser. II. i. 78 = L. divaricata.
Falconeri, Rohrb. in Linnaea, xxxvi. (1869–70) 220 = L. apetala.
fimbriatum, Walp. Rep. i. 280 = L. indica.
firmum, Rohrb. Monog. Silene, 232, 233 = Silene aprica.
Flos-cuculi, Roehl. Deutschl. Fl. ed. II. 274 = Lychnis Flos-cuculi.
Greggii, Rohrb. in Linnaea, xxxvi. (1869–70) 256 = Silene laciniata.
Griffithii, Rohrb. Monog. Silene, 232 = Silene Griffithii.
Hookeri, Rohrb. l. c. 233 = Silene Hookeri.
illinoënse, Rohrb. in Linnaea, xxxvi. (1869–70) 250 = Silene regia.
indicum, Walp. Rep. i. 280 = L. indica.
inflatum, Walp. l. c. = L. inflata.
intermedium, Schur, Enum. Pl. Transs. 108 = L. sylvestris.
intrusum, Rohrb. Monog. Silene, 233 = Silene colorata.
involucratum, Rohrb. in Linnaea, xxxvi. (1869–70) 216 = L. affinis.
laciniatum, Rohrb. Monog. Silene, 233, 234 = Silene laciniata.
lanuginosum, Rohrb. l. c. 231 = Silene auriculata.
lanuginosum, Rohrb. l. c. 233 = Silene lanuginosa.
macrocarpum, Willk. Ic. Desc. Pl. Hisp. i. 28 = L. divaricata.
macrorhizum, Rohrb. in Linnaea, xxxvi. (1869–70) 228 = L. macrorhiza.
magellanicum, Fenzl, ex Rohrb. l. c. 224 = L. magellanica.
Mandonii, Rohrb. l. c. 222.—Boliv.

MELANDRIUM:—
multicaule, Walp. Rep. i. 280 = Lychnis multicaulis.
nemorale, A. Br. in Flora, xxvi. (1843) 371 = L. nemoralis.
nivale, Nym. Syll. 230 = L. nivalis.
noctiflorum, Fries, Bot. Notiser (1843), 143 = Silene noctiflora.
nutans, Walp. Rep. i. 280 = L. nutans.
Oldhamianum, Rohrb. Monog. Silene, 233 = Silene aprica.
Olgae, Maxim. in Bull. Acad. Pétersb. xv. (1871) 225 = Silene Olgae, *Rohrb.*
ornatum, Aschers. ex Rohrb. Monog. Silene, 231, 232, 233 = Silene ornata.
ovalifolium, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 15.—Turkestan.
pennsilvanicum, Rohrb. Monog. Silene, 232, 233, 235 = Silene pensylvanica.
pilosum, Edgew. in Trans. Linn. Soc. xx. (1846) 34 = L. nutans.
pratense, Roehl. Deutschl. Fl. ed. II. 274 = L. sylvestris.
Preslii, Nym. Syll. Suppl. 41 = L. sylvestre.
pulchrum, Rohrb. Monog. Silene, 232, 234 = Silene californica, laciniata.
pumilum, Walp. Rep. i. 280 = L. pumila.
purpureum, Rupr. Fl. Ingric. i. 163 = L. sylvestris.
regium, A. Br. in Flora, xxvi. (1843) 372 = Silene regia.
Requieni, Rohrb. Monog. Silene, 232, 234, 235 = Silene Requienii.
rotundifolium, Rohrb. l. c. 234 = Silene rotundifolia.
rubrum, Garcke, Fl. Deutschl. ed. IV. 55 = L. sylvestris.
saxatile, A. Br. in Flora, xxvi. (1843) 371 = L. saxatilis.
setabense, J. Gay, ex Willk. & Lange, Prod. Fl. Hisp. iii. 641 = L. diclinis.
sibiricum, A. Br. in Flora, xxvi. (1843) 371 = L. sibirica.
songaricum, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 14 = L. apetala.
sordidum, Rohrb. in Linnaea, xxxvi. (1869–70) 233 = L. sordida.
stenophyllum, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 130 = L. sylvestris.
sylvestre, Roehl. Deutschl. Fl. ed. II. i. 274 = L. sylvestris.
thysanodes, A. Gray, Bot. U. St. Expl. Exped. i. 114 = L. thysanodes.
triflorum, J. Vahl, ex Liebm. in Fl. Dan. xiv. 5 = L. triflora.
triste, Fenzl, in Ledeb. Fl. Ross. i. 326 = L. tristis.
undulatum, Rohrb. in Linnaea, xxxvi. (1869–70) 245 = Silene undulata.
Vahlii, Walp. Ann. i. 93.—Ross. bor.
vespertinum, Fries, in Bot. Notiser (1842), 170 = L. alba.
virginicum, A. Br. in Flora, xxvi. (1843) 372 = Silene virginica.
viscosum, Čelak. in Lotos, xviii. (1868) 118 = Silene viscosa.
Wrightii, Rohrb. Monog. Silene, 235 = Silene Wrightii.
Zawadskii, A. Br. in Flora, xxvi. (1843), 387 = Silene Zawadskii.

MELANDRUM, Blytt, Norges Fl. iii. 1068 (1876) = praec.

MELANDRYUM, Reichb. Handb. 298 (1837) = praec.

MELANEA, Pers. Syn. i. 131 (1805) = **Malanea**, Aubl. (Rubiac.).

MELANENTHERA, Link, Enum. Hort. Berol. ii. 309 (1823) = **Melanthera**, Rohr (Compos.).

MELANIUM, P. Br. Hist. Jamaic. 215 (1756) = **Cuphea**, P. Br. (Lythrar.).
alliaceum, Spreng. Syst. ii. 454 = C. viscosissima.
fruticosum, Spreng. l. c. 455 = Pemphis acidula.
hirtum, Spreng. l. c. 454 = C. micrantha.
rupestre, Zipp. in Alg. Konst. en Letter-Bode, i. (1829) 298.—Java.
scabrum, Spreng. Syst. ii. 455 = C. Parsonsia.

MELANIUM, Rich. ex DC. Prod. iii. 135 (1828) = **Arthrostema**, Ruiz & Pav. (Melastom.).

MELANOCARPUM, Hook. f. in Benth. & Hook. f. Gen. iii. 24 (1880) = **Pleuropetalum**, Hook. f. (Amarant.).
Sprucei, Hook. f. l. c. = P. costaricense.

MELANOCARYA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) II. 453 = **Euonymus**, Linn. (Celastr.).
alata, Turcz. l. c. 454 = E. Thunbergianus.

MELANOCENCHRIS, Nees, in Proc. Linn. Soc. i. (1841) 94. *GRAMINEAE*, Benth. & Hook. f. iii. 1169.
PTILONEILEMA, Steud. Syn. Pl. Gram. 201 (1855).
Jacquemontii, *Jaub. & Spach, Illustr.* iv. 36.—Ind. or.
Perrottetii, *Jaub. & Spach, l. c.* 38.—Ind. or.
plumosa, *Hochst. in Flora*, xxxviii. (1855) 273.—Abyssin.
Rothiana, *Nees, in Proc. Linn. Soc.* i. (1841) 95.—Ind. or.
Royleana, *Nees, l. c.*—Ind. or.

MELANOCHYLA, Hook. f. Fl. Brit. Ind. ii. 38 (1876). *ANACARDIACEAE.*
angustifolia, *Hook. f. l. c.* 39.—Malacca.
auriculata, *Hook. f. l. c.*—Malacca.
Maingayi, *Hook. f. l. c.*—Malacca.
tomentosa, Engl. in DC. Monog. Phan. iv. 450, lapsu = Rhus retusa.
tomentosa, *Hook. f. Fl. Brit. Ind.* ii. 38.—Malacca.

MELANOCOCCA, Blume, Mus. Bot. Lugd. Bat. i. 236 (1850). *RUTACEAE*, Benth. & Hook. f. i. 298.
tomentosa, *Blume, Mus. Bot. Lugd. Bat.* i. 236.—N. Guinea.

MELANODENDRON, DC. Prod. v. 279 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 278.
integrifolium, *DC. Prod.* v. 280.—Ins. S. Helena.

MELANOLEPIS, Reichb. & Zoll. in Verh. Natuurk. Ver. Ned. Ind. Bat. i. (1856) 22; et in Linnaea, xxviii. (1856) 324 = **Mallotus**, Lour. (Euphorb.).
angulata, Miq. Fl. Ind. Bat. Suppl. 455 = Mallotus angulatus.
calcosa, Miq. l. c. i. II. 399 = Mallotus calcosus.
diadena, Miq. l. c. Suppl. 455 = Mallotus diadena.
multiglandulosa, Reichb. & Zoll. in Linnaea, xxviii. (1856) 324 = Mallotus moluccanus.

MELANOLEUCE, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 203 = **Melaleuca**, Linn. (Myrtac.).

MELANOLOMA, Cass. in Dict. Sc. Nat. xxix. 472 (1823) = **Centaurea**, Linn. (Compos.).
excelsior, Cass. l. c. 473 = C. pullata.
Fontanesii, Cass. l. c. l. 252 = C. involucrata.
humilis, Cass. l. c. xxix. 473 = C. pullata.
pullata, Cass. l. c. = C. pullata.
Tchihatchewii, Boiss. Fl. Orient. iii. 704 = C. Tchihatcheffii.

MELANOPHYLLA, Baker, in Journ. Linn. Soc. xxi. (1884) 352. *CORNACEAE.*
alnifolia, *Baker, l. c.*—Madag.
aucubaefolia, *Baker, l. c.* 353.—Madag.

MELANOPSIDIUM, Colla, Hort. Ripul. 88. t. 35 (1824) = **Billiottia**, DC. (Rubiac.).
nigrum, Colla, l. c. = B. psychotrioides.

MELANOPSIDIUM, Poit. ex DC. Prod. iv. 443 (1830) = **Alibertia**, Rich. (Rubiac.).

MELANORRHOEA, Wall. Pl. As. Rar. i. 9. tt. 11, 12 (1829). *ANACARDIACEAE*, Benth. & Hook. f. i. 421.
Beccarii, *Engl. Bot. Jahrb.* i. (1881) 45.—Borneo.
glabra, *Wall. Pl. As. Rar.* iii. 50. t. 283.—Burma.
laccifera, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 538.—Cochinch.
macrocarpa, *Engl. in DC. Monog. Phan.* iv. 236.—Borneo.
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 25.—Malacca.
usitata, *Wall. Pl. As. Rar.* i. 9. tt. 11, 12.—Burma.
usitatissima, Steud. Nom. ed. II. ii. 114 = praec.
Wallichii, *Hook. f. Fl. Brit. Ind.* ii. 25.—Malaya.

MELANOSELINUM, Hoffm. Gen. Umb. ed. I. 156 (1814) = **Thapsia**, Linn. (Umbell.).
decipiens, Hoffm. l. c.—Ins. Madera.
Seguieri, Sweet, Hort. Brit. 190 = Ligusticum ferulaceum.

MELANOSERIS, Decne. in Jacquem. Voy. Bot. 101. t. 109 (1844) = **Lactuca**, Tourn. (Compos.).
bracteata, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 409 = L. bracteata.
cyanea, Edgew. in Trans. Linn. Soc. xx. (1846) 81 = L. hastata.
hastata, Edgew. l. c. = L. hastata.
Lessertiana, Decne. in Jacquem. Voy. Bot. 102 = L. Lessertiana.
lyrata, Decne. l. c. 101. t. 109 = L. Lessertiana.
paniculata, Edgew. in Trans. Linn. Soc. xx. (1846) 80 = L. hastata.
rapunculoides, Edgew. l. c. 81 = L. rapunculoides.
saxatilis, Edgew. l. c. 79 = L. macrorhiza.

MELANOSINAPIS, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 944 (1829) = **Brassica**, Linn. (Crucif.).
communis, Spenn. l. c. 945 = B. nigra.

MELANOSTICTA, DC. Mém. Lég. 474. t. 69 (1825) = **Hoffmanseggia**, Cav. (Legumin.).
Burchelli, DC. l. c. = H. Burchellii.
Sandersoni, Harv. in Harv. & Sond. Fl. Cap. iii.. 270.—Afr. austr.

MELANOTIS, Neck. Elem. i. 281 (1790) (Quid?).

MELANOXYLON, Schott, in Spreng. Syst. iv. Cur. Post. 406 (1827). *LEGUMINOSAE*, Benth. & Hook. f. i. 563.
PERITTIUM, Vog. in Linnaea, xi. (1837) 408.
Brauna, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 406.—Bras.

MELANTHERA, Rohr, in Skrift. Nat. Selsk. Kioebenh. ii. (1792) 213. *COMPOSITAE*, Benth. & Hook. f. ii. 377.
LIPOTRICHE, R. Br. in Trans. Linn. Soc. xii. (1817) 118.
MELANANTHERA, Michx. Fl. Bor. Am. ii. 106 (1803).
PSATHUROCHAETA, DC. Prod. v. 609 (1836).
TRIGONOTHECA, Sch. Bip. in Flora, xxvii. (1844) 672, in syn.
WUERSCHMITTIA, Sch. Bip. l. c. xxiv. (1841) I. Intell. 27.
abyssinica, *Benth. & Hook. f. ex Vatke, in Linnaea*, xxxix. (1875) 496.—Afr. trop.
angustifolia, A. Rich. Fl. Cub. Fanerog. ii. 54 = lanceolata.
aspera, Steud. Nom. ed. II. ii. 113 = deltoidea.
Brownei, *Sch. Bip. in Flora*, xxvii. (1844) 673.—Afr. trop.
corymbosa, *Spreng. Neue Entdeck.* ii. 135; iii. 40.—Ind. occ.
deltoidea, *Michx. Fl. Bor. Am.* ii. 106.—Am. trop.
glandulosa, Wall. ex DC. Prod. v. 547 = Wedelia biflora.
hastata, *Michx. Fl. Bor. Am.* ii. 106.—Am. trop.
lanceolata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 88.—Am. trop.
Linnaei, H. B. & K. Nov. Gen. et Sp. iv. 199 = deltoidea.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 418.—Madag.
microphylla, Steetz, in Seem. Bot. Voy. Herald, 156 = lanceolata.
orientalis, Spreng. ex DC. Prod. v. 483 = Caesulia axillaris.
oxylepis, *DC. l. c.* 545.—Ecuador.
panduriformis, Cass. in Dict. Sc. Nat. xxix. 485 = hastata.
pungens, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 382.—Afr. trop.
trilobata, Cass. in Dict. Sc. Nat. xxix. 485 = hastata.
urticaefolia, Cass. l. c. 484 = deltoidea.

MELANTHESA, Blume, Bijdr. 590 (1825) = **Breynia**, Forst. (Euphorb.).
acuminata, Muell. Arg. in Linnaea, xxxii. (1863) 74 = B. acuminata.
anceps, Miq. Fl. Ind. Bat. i. II. 371 = Phyllanthus simplex.
canescens, Zipp. ex Span. in Linnaea, xv. (1841) 347, nomen.—Ins. Timor.
cernua, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 483 = B. cernua.

MELANTHESA:—
chinensis, Blume, Bijdr. 592 (= *Melanthesopsis fruticosa*).—China.
glaucescens, Miq. in Journ. Bot. Néerl. i. (1861) 97.—China.
microphylla, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 49 = Breynia microphylla.
neo-caledonica, Baill. Adansonia, ii. (1861–62) 240 = B. disticha.
obliqua, Wight, Ic. t. 1898 (= *Melanthesopsis patens*).—Ind. or.
ovalifolia, Kostel. Allg. Med.-Pharm. Flora, v. 1772 = B. rhamnoides.
racemosa, Blume, Bijdr. 592 = B. racemosa.
reclinata, Muell. Arg. in Linnaea, xxxii. (1863) 74 = B. racemosa.
retusa, Kostel. Allg. Med.-Pharm. Flora, v. 1771 (= *Melanthesopsis patens*).—Ind. or.
rhamnoides, Blume, Bijdr. 591 = B. rhamnoides.
rhamnoides, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 483 = B. oblongifolia.
rubra, Blume, Bijdr. 591 = B. rubra.
rupestris, Miq. Fl. Ind. Bat. i. II. 371 = Phyllanthus simplex?
turbinata, Wight, Ic. t. 1897 (= *Melanthesopsis patens*).—Ind. or.
virgata, Blume, Bijdr. 592 = B. virgata.

MELANTHESIOPSIS, Benth. & Hook. f. Gen. iii. 277 (1880) = seq.

MELANTHESOPSIS, Muell. Arg. in Linnaea, xxxii. (1863) 74 = **Breynia**, Forst. (Euphorb.).
fruticosa, Muell. Arg. in DC. Prod. xv. II. 437.—China.
lucens, Muell. Arg. in Linnaea, xxxii. (1863) 75 = praec.
patens, Muell. Arg. in DC. Prod. xv. II. 437.—Ind. or.
variabilis, Muell. Arg. in Linnaea, xxxii. (1863) 75.—Ind. or.

MELANTHIUM, Clayt. ex Linn. Gen. ed. II. 524 (1742). *LILIACEAE*, Benth. & Hook. f. iii. 834.
EVONYXIS, Rafin. Fl. Tellur. ii. 29 (1836).
LEIMANTHEMUM, Ritg. in Marb. Schrift. ii. (1830) 130.
LEIMANTHIUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 24.
LIMONANTHUS, Kunth, Enum. Pl. iv. 194, in adnot. (1843).
aethiopicum, Desr. in Lam. Encyc. iv. 29 = Baeometra columellaris.
album, Thunb. Diss. Melanth. 8 = Veratrum album.
angustifolium, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 23 = Androcymbium punctatum.
aspericaule, *Poir. Encyc. Suppl.* iii. 628.—Am. bor.
biglandulosum, Bertol. in Mem. Acc. Sc. Bolog. ii. (1850) 316 = virginicum.
blandum, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 694 = Androcymbium leucanthum.
bracteolare, Desr. in Lam. Encyc. iv. 25 = Veratrum album.
Brownii, Schlecht. in Linnaea, i. (1826) 86 = Anguillaria dioica.
capense, Linn. Sp. Pl. ed. II. 483 = Androcymbium leucanthum.
caricinum, Roth, Nov. Pl. Sp. 199 = Iphigenia indica.
cochinchinense, Lour. Fl. Cochinch. 216.—Cochinch.
densum, Desr. in Lam. Encyc. iv. 26 = Chamaelirium carolinianum.
dioicum, Walt. Fl. Carol. 126 = Chamaelirium carolinianum.
eucomoides, Jacq. Coll. Suppl. 104 = Androcymbium eucomoides.
flavum, Sm. in Rees, Cycl. xxiii. n. 7 = Baeometra columellaris.
glaucum, Nutt. Gen. Am. i. 232 = Zigadenus glaucus.
gramineum, Cav. in Anal. Cienc. Nat. iii. (1801) 50 = Androcymbium punctatum.
gramineum, Heyne, ex Kunth, Enum. Pl. iv. 214 = Iphigenia indica.
graminifolium, Steud. Nom. ed. II. ii. 114 = Androcymbium punctatum.
guineense, Welw. ex Baker, in Journ. Linn. Soc. xvii. (1879) 451 = Iphigenia guineensis.

MELANTHIUM:—
Hultgreenii, [Soland. ex] Thunb. in Mus. Ac. Ups. App. xxvi. 47 = Zygadenus glaucus.
hyacinthium, Heyne, ex Roth, Nov. Pl. Sp. 195 = Scilla indica.
hybridum, Ell. Sketch, i. 419 = virginicum?
hybridum, Nutt. Gen. Am. i. 232 = Veratrum parviflorum.
hybridum, Walt. Fl. Carol. 125 = virginicum.
indicum, Linn. Mant. ii. 226 = Iphigenia indica.
inustum, [Soland.] Ker-Gawl. in Bot. Mag. sub t. 694 = Wurmbea capensis.
japonicum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 22 = Chionographis japonicum.
laetum, Kinn. ex Kunth, Enum. Pl. iv. 195 = virginicum.
laetum, [Soland. in] Ait. Hort. Kew. ed. I. i. 488 = Zygadenus muscitoxicus.
latifolium, Desr. in Lam. Encyc. iv. 25 = virginicum.
longiflorum, Ker-Gawl. in Bot. Mag. sub t. 694 = Wurmbea capensis.
lucidum, Ker-Gawl. l. c. = seq.
luridum, Banks, ex Ker-Gawl. l. c. t. 994 = Wurmbea campanulata.
luteum, Thunb. Fl. Jap. 152 = Chionographis japonica.
luteum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 22 = Chamaelirium carolinianum.
marginatum, Desr. in Lam. Encyc. iv. 29 = Wurmbea capensis.
massoniaefolium, Andr. Bot. Rep. t. 368 = Whiteheadia latifolia.
monoicum, Pursh, Fl. Am. Sept. i. 241 = virginicum.
monoicum, Walt. Fl. Carol. 125 = Veratrum parviflorum.
monopetalum, Linn. f. Suppl. 213 = Wurmbea capensis.
muscaetoxicum, Walt. Fl. Carol. 125 = Zygadenus muscaetoxicus.
myoctonum, J. F. Gmel. Syst. i. 587 = Zygadenus muscaetoxicus.
nigrum, Thunb. Diss. Melanth. 8 = Veratrum nigrum.
nudum, Heyne, in Wall. Herb. (errore *indum*) in Wall. Cat. n. 5062 D = Urginea indica.
Nuttallii, D. Dietr. Syn. Pl. ii. 1206 = Zygadenus Nuttallii.
parviflorum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 276.—Am. bor.
phalangioides, Desr. in Lam. Encyc. iv. 27 = Zygadenus muscaetoxicus.
polygamum, Desr. l. c. 25 = virginicum.
pumilum, Forst. in Comm. Goett. ix. (1789) 30 = Astelia pumila.
pumilum, Ker-Gawl. in Bot. Mag. sub t. 694 = Wurmbea capensis.
punctatum, Cav. in Anal. Cienc. Nat. iii. (1801) 49. t. 26. f. 1 = Androcymbium punctatum.
punctatum, *Linn. Pl. Afr. Rar.* 10; *in Amoen. Acad.* vi. 87 (Quid?).—Afr.
racemosum, Michx. Fl. Bor. Am. ii. 251 = virginicum.
racemosum, Roth, Nov. Pl. Sp. 199 = Iphigenia indica.
racemosum, Walt. Fl. Carol. 125 = Tofieldia pubens.
remotum, [Soland.] Ker-Gawl. in Bot. Mag. sub t. 694 = Wurmbea capensis.
revolutum, [Soland.] Ker-Gawl. l. c. = Wurmbea capensis.
Sabadilla, Thunb. Diss. Melanth. 4 = Veratrum Sabadilla.
sibiricum, *Linn. Sp. Pl.* 339.—Sibir.
spicatum, Burm. f. Fl. Cap. Prod. 11 = Wurmbea capensis.
spicatum, Walt. Fl. Carol. 125 = Xerophyllum setifolium.
stigmosum, Ker-Gawl. in Bot. Mag. sub t. 694 = Wurmbea stigmosa.
striatum, Hill, Veg. Syst. xvi. 56 (Quid?).—Am. bor.
tenue, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 223.—Afr. trop.
uniflorum, Jacq. Coll. iv. 100 = Baeometra columellaris.
virens, Thunb. Diss. Melanth. 4 = Veratrum album.
virescens, Willd. ex Kunth, Enum. Pl. iv. 193 = Zygadenus mexicanus.
virginicum, *Linn. Sp. Pl.* 339.—Am. bor.
virginicum, Walt. Fl. Carol. 125 = Zygadenus glaberrimus.
viride, Linn. f. Suppl. 213 = Ornithoglossum glaucum.
Wurmbeum, Thunb. Prod. Pl. Cap. 67 = Wurmbea capensis.

MELANTHIUM, Kunth, Enum. Pl. iv. 155 (1843) = **Dipidax**, Salisb. (Liliac.).
Bergii, Schlecht. in Linnaea, i. (1826) 84 = D. ciliata.
blandum, Soland. ex Baker, in Journ. Linn. Soc. xvii. (1879) 447 = D. ciliata.
caeruleum, Eckl. ex Schult. f. Syst. vii. 1547 = D. ciliata.
capense, Licht. ex Schult. f. l. c. 1546 = D. ciliata.
ciliatum, Linn. f. Suppl. 213 = D. ciliata.
Garnotianum, Kunth, Enum. Pl. iv. 157 = D. ciliata.
gracile, Desr. in Lam. Encyc. iv. 29 = D. ciliata.
junceum, Jacq. Coll. Suppl. 105 = D. rosea.
luridum, Soland. ex Baker, in Journ. Linn. Soc. xvii. (1879) 447 = D. ciliata.
marginatum, Schlecht. in Linnaea, i. (1826) 84 = D. ciliata.
punctatum, Mill. Gard. Dict. ed. VIII. n. 3 = D. ciliata.
rubicundum, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 22 = D. ciliata.
Schlechtendalianum, Schult. f. Syst. vii. 1546 = D. ciliata.
secundum, Desr. in Lam. Encyc. iv. 28 = D. ciliata.
triquetrum, Linn. f. Suppl. 213 = D. rosae.
virescens, Kunth, Enum. Pl. iv. 193 = D. ciliata.

MELANTHIUM, Medic. Phil. Bot. i. 96 (1789) = **Nigella**, Tourn. (Ranunc.).
damascenum, Medic. l. c. = N. damascena.

MELARGYRA, Rafin. Fl. Tellur. iii. 81 (1836) = **Spergularia**, C. & J. Presl (Caryoph.).
canadensis, Rafin. l. c. = S. media, salina.
media, Rafin. l. c. = S. media.
purpurea, Rafin. l. c. = S. capillacea.
rosea, Rafin. l. c. = S. rubra?
rubra, Rafin. l. c. = S. rubra.

MELARHIZA, Kellogg, in Proc. Calif. Acad. i. ed. II. 37 (1873) = **Wyethia**, Nutt. (Compos.).
inuloides, Kellogg, l. c. = W. helenioides.

MELASANTHUS, Pohl, Pl. Bras. Ic. i. 75. tt. 60–65 (1827) = **Stachytarpheta**, Vahl (Verben.).
glaucus, Pohl, l. c. 81. t. 65 = S. glauca.
integrifolius, Pohl, l. c. 80. t. 64 = S. glauca.
longespicatus, Pohl, l. c. 77. t. 61 = S. Chamissonis.
rhomboidalis, Pohl, l. c. 78. t. 62 = S. rhomboidalis.
triphyllus, Pohl, l. c. 79. t. 63 = S. rhomboidalis.
villosus, Pohl, l. c. 76. t. 60 = S. villosa.

MELASCUS, Rafin. Fl. Tellur. iv. 80 (1836) = **Ipomoea**, Linn. (Convolv.).
latifolius, Rafin. l. c. = I. Bona-nox.

MELASMA, Berg. Desc. Pl. Cap. 162. t. 3. f. 4 (1767).
SCROPHULARINEAE, Benth. & Hook. f. ii. 966.
GASTROMERIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 75 (1830).
LYNCEA, Cham. & Schlecht. in Linnaea, v. (1830) 108.
NIGRINA, Linn. Mant. i. 5 (1767).
hispidum, Benth. in Hook. Comp. Bot. Mag. i. (1835) 202.—Mexic.
ovatum, *E. Mey. in Drège, Zwei Pfl. Docum.* 201, nomen; *et ex Benth. in DC. Prod.* x. 338.—Afr. austr.
rhinanthoides, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 202.—Bras.
scabrum, *Berg. Desc. Pl. Cap.* 162.—Afr. austr.
Zeyheri, Hook. Ic. Pl. t. 255 = Graderia scabra.

MELASPHAERULA, Ker-Gawl. in Bot. Mag. t. 615 (1805). *IRIDEAE*, Benth. & Hook. f. iii. 707.
AGLAEA, Pers. Syn. i. 46 (1805).
DIASIA, DC. in Bull. Soc. Philom. iii. (1803) 151.
graminea, *Ker-Gawl. in Bot. Mag.* t. 615.—Afr. austr.
intermedia, Sweet, Hort. Brit. ed. I. 399 = graminea.
iridifolia, Sweet, l. c. ed. II. 502 = graminea.
parviflora, Lodd. Bot. Cab. t. 1444 = graminea.

MELASTOMA, Burm. ex Linn. Gen. ed. I. 127 (1737).
MELASTOMACEAE, Benth. & Hook. f. i. 746.
ACINOLIS, ALOSEMIS, ANTHERYTA, ANTISOLA, AULIPHAS, DANCERA, EXODICLIS, GYNOMPHIS, OCTELLA, OCTONUM, Rafin. Sylva Tellur. 93–98 (1838).
BENKARA, Adans. Fam. ii. 85 (1763).

MELASTOMA:—
Acinodendron, Linn. Sp. Pl. 389 = Miconia Acinodendron, Sagraea guadalupensis.
aculeatum, Presl, in Oken, Isis, xxi. (1828) 272 = Charianthus ciliatus.
acuminatum, Desr. in Lam. Encyc. iv. 54 = Miconia semicrenata.
acuminatum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Miconia laevigata.
acuminatum, Vahl, Eclog. Am. i. 47 = Tetrazygia discolor.
acutiflorum, Schrank & Mart. ex DC. Prod. iii. 154 = Leandra scabra.
acutipetalum, Rich. in Humb. & Bonpl. Mélast. t. 38 = Ossaea scalpta.
acutum, Mill. Gard. Dict. ed. VIII. n. 13 = Tetrazygia discolor.
adpressum, Benth. in Wall. Cat. n. 4081 = malabathricum.
adscendens, Sw. Fl. Ind. Occ. ii. 772 = Adelobotrys adscendens.
aeleagnoides, Sw. Prod. Veg. Ind. Occ. 72 = Tetrazygia elaeagnoides.
affine, D. Don, in Mem. Wern. Soc. iv. (1823) 288 = malabathricum.
Afzelianum, D. Don, l. c. 291 = Osbeckia multiflora.
aggregatum, Mart. & Schrank, ex DC. Prod. iii. 156 = Ossaea confertiflora.
agreste, Aubl. Pl. Gui. i. 425. t. 166 = Oxymeris agrestis.
agreste, Schrank & Mart. ex DC. Prod. iii. 160 = Miconia astrotricha.
alatum, Aubl. Pl. Gui. i. 410 = Miconia alata.
albicans, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 136 = Miconia cinerascens.
albicans, Sw. Prod. Veg. Ind. Occ. 70 = Miconia holosericea.
albiflorum, G. Don, Gen. Syst. ii. 764 = Tristemma albiflorum.
album, Reinw. ex Blume, in Flora, xiv. (1831) 498 = Dissochaeta gracilis.
alpestre, Jack, in Trans. Linn. Soc. xiv. (1823) 20 = Medinilla alpestris.
alpinum, Sw. Prod. Veg. Ind. Occ. 71 = Charianthus coccineus.
ambiguum, Humb. & Bonpl. Mélast. t. 25 = Miconia ambigua.
amplexicaule, Pav. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 138 = Miconia dipsacea.
amygdalinum, Desr. in Lam. Encyc. iv. 35 = Mecranium amygdalinum.
amygdaloides, Schrank & Mart. ex DC. Prod. iii. 156 = Ossaea amygdaloides.
angustifolium, Sw. Prod. Veg. Ind. Occ. 71 = Tetrazygia angustifolia.
Anhanga, Vell. Fl. Flum. 182; iv. t. 133 = Clidemia hirta.
anisotrichon, Schlecht. in Linnaea, xiii. (1839) 427 = Miconia Berghesiana.
annulatum, Poir. ex Steud. Nom. ed. II. i. 778, in syn. = Marumia annulata.
annulatum, Wall. Cat. n. 4056 = Anplectrum annulatum.
annuum, Schrank, ex DC. Prod. iii. 115 = Aciotis annua.
anoplanthum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 277 = malabathricum.
antennatum, Cham. in Linnaea, ix. (1834) 376 = Huberia ovalifolia.
aplostachyum, Humb. & Bonpl. Mélast. t. 1 = Miconia aplostachya.
apterum, Mart. & Schrank, ex DC. Prod. iii. 161 = Oxymeris aptera.
aquaticum, Aubl. Pl. Gui. i. 430. t. 169 = Nepsera aquatica.
arborescens, Aubl. Pl. Gui. i. 429 = Loreya arborescens.
arborescens, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Blakea pulverulenta.
arboreum, Schlecht. in Linnaea, xiii. (1839) 424 = Conostegia arborea.
arboreum, Vell. Fl. Flum. 182; iv. t. 131 = Miconia calvescens.
argenteum, Desr. in Lam. Encyc. iv. 45 = Tibouchina holosericea.
argenteum, Poir. ex Steud. Nom. ed. I. 516 = Miconia squamulosa.
argenteum, Sw. Prod. Veg. Ind. Occ. 70 = Miconia argentea.
argyratum, Presl, in Oken, Isis, xxi. (1828) 272 = Tetrazygia discolor.

MELASTOMA:—
argyrophyllum, Schrank & Mart. ex DC. Prod. iii. 181 = Miconia argyrophylla.
aristatum, Mart. ex DC. l. c. 157 = Clidemia hirta.
aromaticum, Vahl, Eclog. Am. i. 41 = Tibouchina aspera.
articulatum, Desr. in Lam. Encyc. iv. 56 = Dichaetanthera articulata.
articulatum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 285 = malabathricum.
arvense, Vell. Fl. Flum. 180; iv. t. 126 (= Tremblyae sp.).—Bras.
aspergillare, Humb. & Bonpl. Mélast. t. 9 = Miconia aspergillaris.
asperiusculum, Rich. ex DC. Prod. iii. 155 = Oxymeris asperiuscula.
aspersum, Presl, in Oken, Isis, xxi. (1828) 272 = Miconia laevigata.
asperum, *Blume, Bijdr.* 1076.—Java.
asperum, Linn. Sp. Pl. 389 = Osbeckia aspera.
asperum, Meerb. ex Walp. Rep. ii. 136 = Osbeckia Royeni.
asperum, Mill. Gard. Dict. ed. VIII. n. 17 = Conostegia rufescens.
asperum, Pav. ex DC. Prod. iii. 152 (= *Pleroma mexicanum*).—Mexic.
Astrolasion, Spreng. Neue Entdeck. iii. 62 = Miconia rubiginosa.
Aubletianum, Steud. Nom. ed. II. ii. 115 = Miconia prasina.
auriculatum, Mart. ex DC. Prod. iii. 185 = Miconia auriculata.
azureum, Blume, Bijdr. 1079 = Kibessia azurea.
azureum, Zipp. ex Blume, in Flora, xiv. (1831) 525 = Pternandra caerulescens.
Balbisianum, Ser. ex DC. Prod. iii. 177 = Miconia macrophylla.
bananale, Schrank, ex DC. l. c. 158 = Clidemia blepharodes.
Banksii, Cunn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 59 = malabathricum.
barbatum, Benth. in Wall. Cat. n. 4082 = Anplectrum barbatum.
Baumeanum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 289.—Ins. Philipp.
Berterianum, Spreng. ex DC. Prod. iii. 197 = Charianthus glaberrimus.
Berteroanum, Ser. l. c. 173; Spreng. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 99 = Heterotrichum angustifolium.
bibracteatum, Blume, Bijdr. 1071 = Creochiton bibracteata.
bicolor, Mill. Gard. Dict. ed. VIII. n. 6 = Miconia bicolor.
biglomeratum, Humb. & Bonpl. Mélast. t. 15 = Miconia biglomerata.
bipartitum, Pers. Syn. i. 472 = Miconia triplinervis.
biseptenum, Schrank & Mart. ex DC. Prod. iii. 164 = Oxymeris dichotoma.
biserratum, Schrank & Mart. l. c. 158 = Clidemia biserrata.
bivalve, Aubl. Pl. Gui. i. 404 = Acisanthera brevifolia.
blepharodes, Mart. ex DC. Prod. iii. 158 = Clidemia blepharodes.
Boryanum, *Korth. Verh. Nat. Gesch. Bot.* 232.—Borneo.
brachiatum, Vahl, Eclog. Am. iii. 24 = Miconia laevigata.
brachiatum, Vell. Fl. Flum. 180; iv. t. 125 = Tibouchina Lhotzkyana.
brachyodon, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 292 = polyanthum.
brachyotum, Triana, in Trans. Linn. Soc. xxviii. (1871) 151 = Tibouchina Lhotzkyana.
bracteatum, Jack, in Trans. Linn. Soc. xiv. (1846) 9 = Dissochaeta bracteata.
bracteatum, Mart. & Schrank, ex DC. Prod. iii. 176 = Miconia holosericea.
bracteatum, Wall. Cat. n. 4044, partim = Dissochaeta annulata.
bracteolatum, Humb. & Bonpl. Mélast. 62 t. 20 = Miconia bracteolata.
brasiliense, Spreng. Syst. ii. 297 = Miconia brasiliensis.
brunneum, Mart. ex DC. Prod. iii. 184 = Miconia brunnea.
brunneum, Vahl, Eclog. Am. iii. 19. t. 23 = Henriettea succosa.
bubalinum, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 319 = Miconia bubalina.

MELASTOMA:—
bullosum, Schrank & Mart. ex DC. Prod. iii. 158 = Clidemia bullosa.
bullosum, Spreng. Neue Entdeck. iii. 172 = Clidemia biserrata.
Cacatin, Aubl. Pl. Gui. i. 437 = Bellucia Cacatin.
caelatum, Humb. & Bonpl. Mélast. t. 6 = Miconia caelata.
calvescens, Schrank & Mart. ex DC. Prod. iii. 185 = Miconia calvescens.
calycinum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 485 = candidum.
calycopteron, Rich. in Bonpl. Mélast. t. 42 = Calycogonium stellatum.
calyptratum, Schrank & Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 109 = Miconia Chamissois.
calyptratum, Sieber, ex Triana, l. c. 102 = Miconia semicrenata.
calyptratum, Vahl, Eclog. Am. i. 40 = Conostegia calyptrata.
candidum, *D. Don, in Mem. Wern. Soc.* iv. (1823) 288.—China.
capillare, Pav. ex DC. Prod. iii. 157 = Ossaea leptopus.
capillare, Sw. Prod. Veg. Ind. Occ. 71 = Sagraea capillaris.
capitatum, G. Don, Gen. Syst. ii. 764 = Dissotis capitata.
capitatum, Vahl, Eclog. Am. i. 45 = Tristemma capitatum.
capitellatum, Humb. & Bonpl. Mélast. t. 3 = Clidemia dependens.
carassanum, Schrank & Mart. ex DC. Prod. iii. 162 = Oxymeris carassana.
cardinale, Spreng. Syst. ii. 298 = Tibouchina cardinalis.
carneum, Zipp. ex Blume, in Flora, xiv. (1831) 511 = Medinilla laurifolia.
caudatum, Humb. & Bonpl. Mélast. 13. t. 7 = Miconia caudata.
caudatum, Schrank & Mart. ex DC. Prod. iii. 186 = Miconia caudigera.
celebicum, *Blume, in Flora*, xiv. (1831) 487.—Ins. Celebes.
ceramense, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 276.—Ins. Ceram.
ceramicarpum, Schrank & Mart. ex DC. Prod. iii. 160 = Miconia spondylantha.
cereaceum, Pickering, ex A. Gray, Bot. U. St. Expl. Exped. i. 581, in textu (Astroniae sp. ?).—Ins. Samoa.
cernuum, Benth. in Wall. Cat. n. 4054 = Anplectrum glaucum.
cernuum, Bonpl. ex Loud. Hort. Brit. 174 = Clidemia cernua.
cernuum, Roxb. Hort. Beng. 33; Fl. Ind. ii. 404 = Oxyspora cernua.
chaetodon, Schrank & Mart. ex DC. Prod. iii. 117 = Spennera chaetodon.
Christophoriana, Ham. Prod. Pl. Ind. Occ. 37 = Tetrazygia discolor.
chrysophyllum, Desr. in Lam. Encyc. iv. 50 = Rousseauxia chrysophylla.
chrysophyllum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Miconia fulva.
chulese, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 222 = Osbeckia chulesis.
ciliatum, Desr. in Lam. Encyc. iv. 49 (= *Pleroma strigosum*).—Am. trop.
ciliatum, Pav. ex D. Don, Mem. Wern. Soc. v. (1822) 309 = Calophysa ciliata.
ciliatum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Miconia ciliata.
ciliatum, Schrank & Mart. ex DC. Prod. iii. 180 = Miconia pileata.
cinchonaefolia, Mart. l. c. 184 = Miconia cinchonaefolia.
cinereum, Vahl, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 103 = Miconia pallens.
cinnamomifolium, Jacq. Coll. ii. 110 = Miconia cinnamomifolia.
clavatum, Pers. Syn. i. 472 = Tibouchina holosericea.
coccineum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Miconia spondylantha.
coccineum, Vahl, Eclog. Am. i. 48 = Charianthus purpureus.
coccineum, Vell. Fl. Flum. 183; iv. t. 136 = Clidemia blepharodes.
coeleste, Pav. ex DC. Prod. iii. 192 = Miconia coelestis.
coeruleum, Hort. ex Reichb. Hort. Bot. iii. 18 = Tibouchina viminea.
coeruleum, Pav. ex DC. Prod. iii. 192 = Miconia caerulea.

MELASTOMA:—
columneaefolia, Schrank, ex DC. Prod. iii. 170 = Clidemia rubra.
compressum, Vahl, Eclog. Am. iii. 23. t. 17 = Miconia Fothergilla.
confertum, Vell. Fl. Flum. 181; iv. t. 127 = Acanthella Sprucei.
conglomeratum, Rich. ex DC. Prod. iii. 156 = Clidemia conglomerata.
conglomeratum, Schrank & Mart. l. c. 163 = Oxymeris Nianga.
constrictum, Blume, Bijdr. 1072 = Pachycentria constricta.
cordatum, Schrank & Mart. ex DC. Prod. iii. 107 = Rhynchanthera cordata.
cordatum, Vell. Fl. Flum. 178; iv. t. 114 = Aciotis dysophylla.
cordifolium, Hort. ex Link, Enum. Hort. Berol. i. 397 (Quid ?).—Bras.?
cordifolium, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 405.—Ind. or.
coriaceum, Juss. ex DC. Prod. iii. 190 = Miconia Thomasiana.
coriaceum, Sw. Prod. Veg. Ind. Occ. 70 = Miconia coriacea.
cornifolium, Desr. in Lam. Encyc. iv. 51 = Miconia cinnamomifolia.
cornoides, Schlecht. ex Cham. in Linnaea, v. (1830) 563 = Oxymeris cornioides.
coronatum, Humb. & Bonpl. Mélast. t. 56 = Miconia coronata.
corymbosum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109; Vahl, Eclog. Am. iii. 25 = Charianthus glaberrimus.
corymbosum, Sims, in Bot. Mag. t. 904 = Amphiblemma cymosum.
crassifolium, Reinw. ex Blume, Bijdr. 1075 = Medinilla crassifolia.
crenatum, Pav. ex DC. Prod. iii. 165 = Oxymeris crenata.
crenatum, Vahl, Eclog. Am. i. 41 = Clidemia hirta.
crinitum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 280 = sanguineum.
crinitum, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 402 = Osbeckia stellata.
crinitum, *Vahl, Eclog. Am.* iii. 28.—Ins. Martinica.
crispatum, Linn. Sp. Pl. ed. II. 560 = Medinilla crispata.
croceum, Desr. in Lam. Encyc. iv. 55 = Miconia crocea.
croceum, Spreng. Syst. ii. 302 = Miconia racemosa.
crotonifolium, Desr. in Lam. Encyc. iv. 43 = Tetrazygia crotonifolia.
cryptodon, Schrank & Mart. ex DC. Prod. iii. 157 = Clidemia hirta.
cucullatum, Pav. l. c. 176 = Calyptrella cucullata.
cupreum, Sm. in Rees, Cycl. xxiii. n. 14, 15 = Miconia rubiginosa.
curvum, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 406.—Ind. or.
cuspidatum, Willd. ex Cham. in Linnaea, ix. (1834) 458 = Miconia laevigata.
cyanocarpum, Blume, Bijdr. 1073 = Anplectrum cyanocarpum.
cyanoides, Sm. in Rees, Cycl. xxiii. n. 56, 57 = Otanthera moluccana.
cymbifolium, Rich. in Bonpl. Mélast. t. 26 = Miconia coriacea.
cymosum, Vent. ex DC. Prod. iii. 147 = Amphiblemma cymosum.
cymosum, Wendl. ex Spreng. Syst. ii. 299 = Heterotrichum patens.
decemdentatum, Kostel. Allg. Med. Pharm. Fl. iv. 1511 (Osbeckiae sp. ?).—Ind. or.
decemfidum, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 406 = sanguineum.
decumbens, Beauv. Fl. Owar. i. 69. t. 41 = Dissotis decumbens.
decussatum, Vahl, Eclog. Am. i. 42 = Miconia ciliata.
densifrons, Schrank & Mart. ex DC. Prod. iii. 194 = Miconia ligustroides.
dentatum, Pav. ex DC. l. c. 158 = Clidemia dentata.
denticulatum, Bonpl. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 214 = Miconia denticulata.
denticulatum, Labill. Sert. Austro-Caled. i. 65. t. 64 = malabathricum.
depauperatum, Rich. ex DC. Prod. iii. 161 = Leandra agrestis.
dependens, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 307 = Clidemia dependens.

MELASTOMA :—
detergibile, Schrank & Mart. ex DC. Prod. iii. 181 = Miconia albicans.
diaphanum, Blume, Bijdr. 1075 = Medinilla crassifolia.
dichotomum, Desr. in Lam. Encyc. iv. 41 = Rhynchanthera dichotoma.
dichotomum, Pav. ex DC. Prod. iii. 160 = Oxymeris dichotoma.
diffusum, Balb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Ossaea microphylla.
diffusum, Desr. in Lam. Encyc. iv. 38 = Acisanthera recurva.
diffusum, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 291 (= *Pleroma diffusum*).—Porto Rico.
diffusum, Reinw. ex Blume, in Flora, xiv. (1831) 493 = Dissochaeta inappendiculata.
dimorphum, Schrank & Mart. ex DC. Prod. iii. 193 = Miconia milleflora.
discolor, Bert. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 99 = Charianthus coccineus.
discolor, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 131 = Oxymeris cinnamomea.
discolor, Linn. Syst. ed. X. 1022 = Tetrazygia discolor.
divaricatum, Willd. Sp. Pl. ii. 596 = Anplectrum divaricatum.
diversifolium, Humb. & Bonpl. Mélast. t. 59 = Octopleura diversifolia.
Dodandianum, Ham. Prod. Pl. Ind. Occ. 37 = Charianthus ciliatus.
dodecandrum, Blanco, Fl. Filip. ed. II. 258 = imbricatum.
dodecandrum, Desr. in Lam. Encyc. iv. 46 = Miconia Fothergilla.
dodecandrum, Lour. Fl. Cochinch. 274 = repens.
dodecandrum, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 406.—Ins. Molucc.
dubium, Schrank & Mart. ex DC. Prod. iii. 154 = Oxymeris dubia.
ebracteatum, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 637 = Miconia semicrenata.
echinatum, Reinw. ex Blume, in Flora, xiv. (1831) 524 = Kibessia azurea.
ecostatum, [Dryand. in] Ait. Hort. Kew. ed. II. iii. 46 = Miconia ecostata.
egense, Mart. ex DC. Prod. iii. 165 = Myriaspora egensis.
elaeagnoides, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Tetrazygia discolor.
elaeagnoides, Sw. Fl. Ind. Occ. ii. 815 = Tetrazygia elaeagnoides.
elatum, Sw. Prod. Veg. Ind. Occ. 70 = Miconia elata.
elegans, Aubl. Pl. Gui. i. 427. t. 167 = Clidemia hirta.
ellipticum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 291 = malabathricum?
elongatum, D. Don, in Mem. Wern. Soc. iv. (1822) 290 = Dissotis grandiflora.
elongatum, Vahl, Eclog. Am. ii. 16 = Miconia semicrenata.
enanum, Vell. Fl. Flum. 179; iv. t. 117 = Bertolonia Leuzeana.
Epibaterium, Mart. ex DC. Prod. iii. 157 = Octopleura Epibaterium.
epidendrum, Reinw. ex Blume, in Flora, xiv. (1831) 515 = Medinilla javanensis.
erectum, *Jack, in Trans. Linn. Soc.* xiv. (1823) 5.—Malaya.
erostratum, Mart. & Schrank, ex DC. Prod. iii. 160 = Oxymeris erostrata.
erythrophyllum, Wall. Cat. n. 4085 = Medinilla rubicunda.
exiguum, Jack, in Trans. Linn. Soc. xiv. (1823) 10 = Allomorphia exigua.
eximium, Blume, Bijdr. 1072 = Medinilla speciosa.
eximium, Jack, in Trans. Linn. Soc. xiv. (1823) 17 = Medinilla eximia.
eximium, Siebold, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 87 = Medinilla Sieboldiana.
expansum, Vahl, Eclog. Am. iii. 21 = Miconia laevigata.
extinctorium, Humb. & Bonpl. Mélast. t. 57 = Conostegia extinctoria.
fallax, Blume, Bijdr. 1068 = Dissochaeta intermedia.
fallax, Jack, in Trans. Linn. Soc. xiv. (1823) 13 = Omphalopus fallax.
fallax, Wall. Cat. n. 4050 = Dissochaeta celebica.
farinulentum, *Sw. ex Wikstr. in Vet. Acad. Handl. Stockh.* (1827) 65.—Ins. Guadalup.

MELASTOMA :—
fasciculare, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 288 = imbricatum.
fasciculare, Sw. Prod. Veg. Ind. Occ. 71 = Henriettella fascicularis.
fasciculare, Willd. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 146 = Ossaea scalpta.
fastigiatum, Mart. ex Triana, l. c. 116 = Miconia stenostachya.
favosum, Desr. in Lam. Encyc. iv. 47 = Miconia favosa.
ferruginatum, DC. Prod. iii. 181 = seq.
ferrugineum, Desr. in Lam. Encyc. iv. 42 = Miconia ferruginea.
ferrugineum, Reinw. ex Blume, in Flora, xiv. (1831) 505 = Marumia muscosa.
ferrugineum, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 402 = Dissochaeta intermedia.
Finlaysonii, Wall. Cat. n. 4041 = Dissotis Findlaysonii.
flavescens, Aubl. Pl. Gui. i. 423 = Henriettella flavescens.
flavum, Vell. Fl. Flum. 182; iv. t. 135 = Oxymeris longibarbis?
floribundum, Humb. & Bonpl. Mélast. t. 53 = Miconia floribunda.
floribundum, Sieber, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 71 = Graffenrieda latifolia.
Fontanesii, Spreng. Syst. ii. 299 = Tibouchina granulosa.
Fothergilla, Desr. in Lam. Encyc. iv. 45; Humb. & Bonpl. Mélast. t. 32 = Miconia Fothergilla.
fothergilloides, Schrank & Mart. ex DC. Prod. iii. 108 = Rhynchanthera grandiflora.
foveolatum, Schrank & Mart. l. c. 164 = Oxymeris foveolata.
fragile, DC. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 136 = Clidemia biserrata.
fragile, Linn. f. Suppl. 236 = Miconia eriopoda.
fragile, Rich. ex DC. Prod. iii. 115 = Aciotis fragilis.
fulvum, Rich. in Humb. & Bonpl. Mélast. t. 11 = Miconia fulva.
furcatum, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 406.—Ins. Molucc.
furfuraceum, Vahl, Eclog. Am. iii. 13. t. 22 = Miconia furfuracea.
Gaudichaudianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 278 = sanguineum.
geniculatum, Roxb. Fl. Ind. ii. 403 (Osbeckiae sp.?).—Ind. or.
glaberrimum, Schlecht. in Linnaea, xiii. (1839) 427 = Miconia glaberrima.
glabratum, Sw. Prod. Veg. Ind. Occ. 71 = Calycogonium glabratum.
glabrescens, Schrank & Mart. ex DC. Prod. iii. 187 = Miconia staminea.
glabrum, Forst. f. Prod. 34 = Astronia Forsteri, pileolata.
glabrum, Vell. Fl. Flum. 180: iv. t. 121 = Henriettella umbelluliflora.
glandulosum, Sw. Fl. Ind. Occ. ii. 799 = Heterotrichium octandrum.
glaucocarpum, Schlecht. in Linnaea, xiii. (1839) 421 = Miconia oligotricha.
glaucum, Griff. Ic. Pl. Asiat. t. 637 = Dissochaete gracilis.
glaucum, Jack, in Trans. Linn. Soc. xiv. (1823) 15 = Anplectrum divaricatum.
glaucum, Wall. Cat. n. 4055 = Anplectrum glaucum.
globuliferum, Walp. Rep. ii. 133 = seq.
globuliflorum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Miconia Sieberi.
glomeratum, Klein, ex Wight & Arn. Prod. 323 = Osbeckia octandra.
glomeratum, Vahl, ex DC. Prod. iii. 168 = Ossaea scalpta.
glutinosum, Linn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 23 = Bucquetia glutinosa.
gracile, Jack, l. c. xiv. (1823) 14 = Dissochaeta gracilis.
grandiflorum, Aubl. Pl. Gui. i. 414. t. 160 = Rhynchanthera grandiflora.
grandiflorum, Vell. Fl. Flum. 178; iv. t. 109 = Tibouchina estrellensis.
grandifolium, Hort. Chels. ex Loud. Hort. Brit. 174 = Miconia grandifolia.
grandifolium, Schrank, ex DC. Prod. iii. 107 = Rhynchanthera Haenkeana.
granulosum, Desr. in Lam. Encyc. iv. 44 = Tibouchina granulosa.
granulosum, Humb. & Bonpl. Mélast. t. 12 = Miconia granulosa.

MELASTOMA :—
granulosum, Willd. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 118 = Miconia cinerascens.
Grossularia, Crantz, Inst. ii. 195 = seq.
grossularioides, Linn. Sp. Pl. 390 = Bellucia Aubletii.
grossum, Linn. f. Suppl. 236 = Tibouchina grossa.
guayaquilense, Humb. & Bonpl. Mélast. t. 49 = Miconia guayaquilensis.
guianense, Poir. ex Steud. Nom. ed. I. 517 = Maieta guianensis.
Haenkeanum, Mart. ex DC. Prod. iii. 107 = Rhynchanthera Haenkeana.
haplostachyum, Spreng. Syst. ii. 297 = Miconia aplostachya.
herpeticum, Schrank & Mart. in DC. Prod. iii. 181 = Miconia herpetica.
heterobase, Mart. ex DC. l. c. 164 = Oxymeris heterabasis.
heterocarpum, Schrank & Mart. ex DC. l. c. 155 = Oxymeris rufescens.
heteromallum, D. Don, in Bot. Reg. t. 644 = Tibouchina heteromalla.
heteroneuron, Schrank & Mart. ex DC. Prod. iii. 201 = Sagraea heteroneura.
heterophyllum, Desr. in Lam. Encyc. iv. 34 = Calophysa heterophylla.
heterostegium, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 286 = malabathricum.
hirsutum, Sw. Prod. Veg. Ind. Occ. 72 = Ossaea hirsuta.
hirsutum, Vell. Fl. Flum. 179; iv. t. 119 = Leandra melastomoides.
hirtellum, Sw. Prod. Veg. Ind. Occ. 72. = Ossaea hirtella.
hirtum, Chapel. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 57 = Antherotoma Naudinii.
hirtum, Desr. in Lam. Encyc. iv. 43 = Heterotrichum angustifolium.
hirtum, Linn. Sp. Pl. 390 partim = Clidemia hirta.
hirtum, Mill. Gard. Dict. ed. VIII. n. 3 = Conostegia speciosa.
hirtum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Sagraea umbrosa.
hispidum, Mart. ex DC. Prod. iii. 164 = Oxymeris intermedia.
hispidum, Sw. Prod. Veg. Ind. Occ. 72 = Tetrazygia hispida.
holosericeum, Linn. Sp. Pl. 390 = Miconia holosericea.
holosericeum, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 316 = Conostegia xalapensis.
holosericeum, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 116 = Miconia stenostachya.
holosericeum, Sw. Obs. Bot. 176 = Tibouchina holosericea.
holosericeum, Vahl, Eclog. Am. ii. 42 = Miconia albicans.
holosericeum, Vell. Fl. Flum. 179; iv. t. 118 = Leandra scabra.
Hombronianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 278 = polyanthum.
homostegium, Naud. l. c. 286 = obovatum.
Houtteanum, *Naud. l. c.* 291.—Burma.
humile, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 222 = Osbeckia nepalensis.
hymenonervium, *Raddi, in Mem. Soc. Ital. Moden.* xx. (1829) 135.—Bras.
hypericifolium, Blume, Bijdr. 1070 = Medinilla hypericifolia.
hypophyscon, Schrank & Mart. ex DC. Prod. iii. 166 = Maieta guianensis.
ibaguense, Humb. & Bonpl. Mélast. i. 105. t. 45 = Miconia auriculata.
icosandrum, Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1827) 64. (Quid?).—Ins. Guadelup.
idorense, Steud. Nom. ed. II. ii. 116, in syn. sphalm. = *tidorense*.
imbricatum, *Wall. Cat.* n. 4047.—Ind. or.; Malaya.
impetiginosum, Mart. ex DC. Prod. iii. 183 = Miconia rufescens.
impetiolare, Sw. Prod. Veg. Ind. Occ. 70 = Miconia impetiolaris.
impuberum, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 404 = Allomorphia exigua.
inaequalifolium, Schrank & Mart. ex DC. Prod. iii. 164 = Oxymeris inaequalifolia.
inundatum, Steud. Nom. ed. II. ii. 116 = Acisanthera inundata.
involucratum, Desr. in Lam. Encyc. iv. 35 = Meriania obtusifolia.

MELASTOMA :—

involucratum, D. Don, in Mem. Wern. Soc. iv. (1823) 290 = Tristemma involucratum.
involucratum, Mart. & Schrank, ex DC. Prod. iii. 154 = Leandra involucrata.
involucratum, Rich. ex DC. l. c. 163 = Clidemia involucrata.
Jackianum, Korth. Verh. Nat. Gesch. Bot. 227 = malabathricum.
januarense, Schrank & Mart. ex DC. Prod. iii. 156 = Ossaea marginata.
japurense, Schrank & Mart. l. c. 159 = Clidemia japurensis.
javanense, Blume, Bijdr. 1078 = Medinilla javanensis.
lacerum, Humb. & Bonpl. Mélast. t. 5 = Miconia lacera.
laevigatum, Aubl. Pl. Gui. i. 412 = Miconia prasina.
laevigatum, Linn. Sp. Pl. 390 = Miconia laevigata.
laevigatum, Mart. & Schrank, ex DC. Prod. iii. 187 = Tococa subciliata.
laevigatum, Sieber, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 97 = Conostegia montana.
Lambertianum, Ser. ex DC. Prod. iii. 165 = Oxymeris crenata.
lanatum, Hort. Chels. ex G. Don, in Loud. Hort. Brit. 174 = Tococa lanata.
lanatum, Lamb. ex DC. Prod. iii. 162 = Miconia lanata.
lanceolatum, Desr. in Lam. Encyc. iv. 43 = Miconia lanceolata.
laniflorum, D. Don, in Mem. Wern. Soc. iv. (1823) 292 = Trembleya laniflora.
lanuginosum, *Blume, Bijdr.* 1077.—Java.
lappaceum, Desr. in Lam. Encyc. iv. 42 = Heterotrichum patens.
lappaceum, Rich. ex DC. Prod. iii. 162 = Miconia lappacea.
lasiopetalum, Mart. & Schrank, ex DC. l. c. 177 = Miconia macrophylla.
lateriflorum, Vahl, Eclog. Am. i. 48 = Henriettella lateriflora.
latifolium, Desr. in Lam. Encyc. iv. 31 = Sagraea latifolia.
laurifolium, Benth. in Wall. Cat. n. 4084 = Medinilla Hasseltii.
laurifolium, Blume, Bijdr. 1069 = Medinilla laurifolia.
laurinum, Ser. ex DC. Prod. iii. 167 = Huberia laurina.
laxiflorum, Schlecht. in Linnaea, xiii. (1839) 426 = Clidemia laxiflora.
laxum, Desr. in Lam. Encyc. iv. 41 (= *Pleroma laxum*).—Peruv.
laxum, Rich. ex DC. Prod. iii. 116 = Aciotis laxa.
ledifolium, Desr. in Lam. Encyc. iv. 48 = Brachyotum ledifolium.
leonense, Lodd. ex G. Don, in Loud. Hort. Brit. 173 = Amphiblemma cymosum.
lepidotum, Mart. & Schrank, ex DC. Prod. iii. 180 = Miconia lepidota.
leprosum, Blume, Bijdr. 1068 = Dissochaeta leprosa.
leucandrum, Schrank, ex DC. Prod. iii. 184 = Miconia cinchonaefolia.
leucocarpum, Mart. & Schrank, ex DC. l. c. 182 = Miconia leucocarpa.
Lhotzkyi, Cogn. in Mart. Fl. Bras. xiv. III. 50, in syn. = Microlicia insignis.
ligustrinum, Mart. & Schrank, ex DC. Prod. iii. 194 = Miconia ligustroides.
ligustrinum, Sm. in Rees, Cycl. xxiii. n. 27–28 = Miconia rhamnoidea.
Lima, Desr. in Lam. Encyc. iv. 47 = Ossaea Lima.
lineare, Reinw. ex Blume, in Flora, xiv. (1831) 474 = Osbeckia linearis.
lineatum, Schlecht. in Linnaea, xiii. (1839) 423 = Miconia auriculata.
litoreum, Wall. Cat. n. 4087 = Ochthocharis javanica.
longibarbe, Schrank & Mart. ex DC. Prod. iii. 163 = Oxymeris longibarbis.
longiflorum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 290 = asperum.
longifolium, Aubl. Pl. Gui. i. 432 = Miconia longifolia.
longifolium, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 293.—Singapore.
loxense, Humb. & Bonpl. Mélast. t. 43 = Miconia loxensis.
lutescens, Humb. & Bonpl. l. c. t. 17 = Miconia lutescens.
lutescens, Vahl, Eclog. Am. ii. 17 = Miconia cinnamomifolia.
luteum, Ruiz & Pav. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Maieta Poeppigii.

MELASTOMA :—

macrocarpum, D. Don, in Mem. Wern. Soc. iv. (1823) 289 = candidum.
macrocarpum, Hook. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 59 = malabathricum.
macrocarpum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 281 = sanguineum.
macrophyllum, Desr. in Lam. Encyc. iv. 44 = Miconia impetiolaris.
macrophyllum, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 319 = Miconia platyhedra.
macrophyllum, Vell. Fl. Flum. 180; iv. t. 122 = Miconia macrophylla.
madagascariense, D. Don, in Mem. Wern. Soc. iv. (1823) 290 = Dichaetanthera madagascariensis.
Maieta, Desr. in Lam. Encyc. iv. 34 = Maieta guianensis.
Majeta, Schrank & Mart. ex DC. Prod. iii. 166 = Maieta hypophysca.
malabathricum, Jack, in Trans. Linn. Soc. xiv. (1823) 4 = polyanthum.
malabathricum, *Linn. Sp. Pl.* 390.—Ind. or.; Malaya.
malabathricum, Sims, Bot. Mag. t. 529 = candidum.
malabatricum, Blanco, Fl. Filip. ed. I. 367 = sanguineum.
mandiocanum, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 132 = Miconia calvescens.
marginatum, Desr. in Lam. Encyc. iv. 32 = Ossaea marginata.
marianum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 276.—Ins. Marian.
Martianum, Schrank, ex DC. Prod. iii. 186 = Miconia Martiusiana.
martinicense, Presl, in Oken, Isis, xxi. (1828) 272 = Charianthus glaberrimus.
Martiusianum, DC. Prod. iii. 186 = Miconia Martiusiana.
Matuba, Spreng. Neue Entdeck. iii. 61 = Miconia trichotoma.
Maximilianum, Mart. ex DC. Prod. iii. 186 = Miconia Maximiliana.
megalophyllum, Mart. l. c. 183 = Miconia tomentosa.
melanophyllum, Spreng. Neue Entdeck. ii. 171 = Sagraea melanophyllum.
mellinum, Mart. & Schrank, ex DC. Prod. iii. 182 = Miconia mellina.
mexicanum, D. Don, in Mem. Wern. Soc. iv. (1823) 290 (= *Pleroma mexicanum*, Triana).—Mexic.
mexicanum, Hook. & Arn. Bot. Beech. Voy. 290 = Heeria subtriplinervia.
mexicanum, Humb. & Bonpl. Mélast. t. 56 = Miconia mexicana.
micranthum, Bert. ex DC. Prod. iii. 169 = Ossaea acuminata.
micranthum, Sw. Prod. Veg. Ind. Occ. 71 = Octopleura micrantha.
microphyllum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 293.—Ind. or.
microphyllum, Sw. Prod. Veg. Ind. Occ. 72 = Ossaea microphylla, *Triana*.
milleflorum, Schrank & Mart. ex DC. Prod. iii. 193 = Miconia milleflora.
minutiflorum, Humb. & Bonpl. Mélast. t. 22 = Miconia milleflora.
Molkenboerii, *Miq. Fl. Ind. Bat.* i. 1. 513. t. 8. f. B.—Java.
molle, *Wall. Cat. n.* 4046.—Singapor.
moluccanum, Blume, Bijdr. 1078 = Otanthera moluccana.
montanum, Spreng. Syst. ii. 296 = Conostegia parviflora.
montanum, Sw. Prod. Veg. Ind. Occ. 69 = Conostegia montana.
mucronatum, Desr. in Lam. Encyc. iv. 46 = Miconia holosericea.
multiflorum, Desr. l. c. 35 = Mecranium multiflorum.
muscosum, Blume, Bijdr. 1070 = Marumia muscosa.
mutabile, Rich. ex DC. Prod. iii. 162 = Miconia mutabilis.
mutabile, Vell. Fl. Flum. 181; iv. t. 130 = Tibouchina mutabilis.
Mutisii, Humb. & Bonpl. Mélast. t. 136 = Centronia Mutisii.
myrcioides, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Conostegia montana.
myricoides, Desr. in Lam. Encyc. iv. 51 = Miconia Sieberi.
Nabotan, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 262 = *M. Noötan.*
napalense, Lodd. Bot. Cab. t. 707 = normale.

MELASTOMA :—

nemorosum, Jack, in Trans. Linn. Soc. xiv. (1823) 8 = Marumia nemorosa.
nervosum, Sm. in Rees, Cycl. xxiii. n. 31 = Miconia spondylantha.
nicotianaefolium, Desr. in Lam. Encyc. iv. 53 = Miconia laevigata.
nigricans, Vahl, Eclog. Am. iii. 26 = Miconia trichotoma.
nitens, Schrank & Mart. ex DC. Prod. iii. 194 = Miconia ligustroides.
nitidum, Korth. Verh. Nat. Gesch. Bot. 234 = sanguineum.
nitidum, Pav. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 116 = Miconia albicans.
nitidum, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 586.—Java.
niveum, Desr. in Lam. Encyc. iv. 42 = Heterotrichum patens.
Nobotan, Blume, Mus. Bot. Lugd. Bat. i. 54 = candidum.
nodosum, Desr. in Lam. Encyc. iv. 55 = Charianthus ciliatus.
nodosum, Zipp. ex Blume, in Flora, xiv. (1831) 510 = Medinilla macrocarpa.
normale, *D. Don, Prod. Fl. Nep.* 220.—Reg. Himal.
Novae-Hollandiae, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 290 = malabathricum.
obovatum, Pav. ex DC. Prod. iii. 192 = Miconia myrtiformis.
obscurum, Humb. & Bonpl. Mélast. t. 52 = Miconia obscura.
obvolutum, Jack, in Trans. Linn. Soc. xiv. (1823) 3 = malabathricum.
octandrum, Linn. ex Sm. in Rees, Cyclop. xxiii. n. 67 = repens.
octandrum, Linn. Sp. Pl. 391 = Osbeckia octandra.
octonum, Humb. & Bonpl. Mélast. t. 4 = Heterotrichum octonum.
oliganthum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 292 = polyanthum.
ornatum, Sw. Prod. Veg. Ind. Occ. 69 (= *Pleroma strigosum.*).—Am. austr.
osbeckioides, Ker-Gawl. in Bot. Mag. t. 2235 = Osbeckia octandra.
oxypetalum, Mart. & Schrank, ex DC. Prod. iii. 190 = Oxymeris quinquedentata.
oxypetalum, Spreng. Syst. ii. 303 = Ossaea scalpta.
oxyphyllum, Benth. in Wall. Cat. n. 4083 = Ochthocharis paniculata.
paleaceum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 279.—Cochinch.
pallens, Spreng. Neue. Entdeck. iii. 62 = Miconia pallens.
pallidum, Jack, in Trans. Linn. Soc. xiv. (1823) 12 = Dissochaeta pallida.
paniculatum, Mart. & Schrank, ex DC. Prod. iii. 194 = Miconia paniculata.
paniculatum, Mill. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 95 = Oxymeris secundiflora.
papillosum, Desr. in Lam. Encyc. iv. 48 = Miconia papillosa.
paradoxum, Mart. ex DC. Prod. iii. 202 = Miconia paradoxa.
parviflorum, Aubl. Pl. Gui. i. 433. t. 171 = Miconia prasina.
patens, Spreng. ex DC. Prod. iii. 177 = Miconia platyhedra.
patens, Sw. Prod. Veg. Ind. Occ. 68 = Heterotrichum patens.
paucidens, Schrank & Mart. ex DC. Prod. iii. 186 = Miconia paucidens.
pauciflorum, Desr. in Lam. Encyc. iv. 39 = Clidemia hirta.
paucifolium, Bonpl. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 143 = Henriettella Goudotiana.
paulense, Schrank, ex DC. Prod. iii. 165 = Myriaspora egensis.
paulinum, Schrank & Mart. ex DC. l. c. 153 = Leandra paulina.
pelagicum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 279 = normale.
pellucens, Rich. ex DC. Prod. iii. 115 = Aciotis purpurascens.
pendulifolium, Humb. & Bonpl. Mélast. t. 35 = Miconia laevigata.
pendulifolium, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Miconia prasina.
pendulum, Salzm. ex Griseb. Fl. Brit. W. Ind. 257 = Miconia minutiflora.
penicillatum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 280.—Ins. Philipp.

MELASTOMA:—
petandrum, *Vell. Fl. Flum.* 181; iv. t. 128.—Bras.
petiolare, Schlecht. in Linnaea, v. (1830) 562 = Clidemia petiolaris.
petiolare, Wall. Cat. n. 4053 = Anplectrum pallens.
petiolatum, *Mill. Gard. Dict.* ed. VIII. n. 9.—Hab. ?
petiolatum, Rich. ex DC. Prod. iii. 157 = Sagraea petiolata.
pexiricum, Vell. Fl. Flum. 182; iv. t. 134 = Clidemia rubra.
Phyllopus, Schrank & Mart. ex DC. Prod. iii. 178 = Henriettea Martii.
physiphorum, Vahl, Eclog. Am. i. 45 = Tococa guianensis.
pictum, *Rudolphi, in Schrad. Neues Journ.* ii. (1807) 293.—Ins. S. Doming.
pictum, Vahl, Eclog. Am. ii. 15 = Miconia cinnamomifolia.
pileatum, Schrank & Mart. ex DC. Prod. iii. 180 = Miconia pileata.
pilosum, Pav. ex DC. l. c. 161 = Calophysa pilosa.
pilosum, Spreng. Syst. ii. 304 = Sagraea domingensis.
pilosum, Sw. Prod. Veg. Ind. Occ. 72 = Sagraea pilosa.
piluliferum, Vahl, Eclog. Am. iii. 15 = Miconia laevigata.
plantaginifolium, Mill. Gard. Dict. ed. VIII. n. 1 = Miconia splendens.
pleurocarpum, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 795 = Miconia Acinodendron.
Plukenetii, Schrank & Mart. ex DC. Prod. iii. 181 = Miconia ferruginata.
plumosum, Desr. in Lam. Encyc. iv. 31 = Sagraea plumosa.
plumosum, D. Don, in Mem. Wern. Soc. iv. (1823) 290 = Dissotis rotundifolia.
polyanthum, Benth. in Wall. Cat. n. 4051 = Anplectrum divaricatum.
polyanthum, *Blume, in Flora*, xiv. (1831) 480.—Malaya; Austral.
polyanthum, Korth. ex Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 288 = pusillum, *Blume*.
polygamum, Bonpl. ex Naud. l. c. xvi. (1851) 195 = Miconia rubens.
porphyraeum, Zipp. ex Blume, in Flora, xiv. (1831) 487 = sanguineum.
portoricense, Spreng. Neue Entdeck. iii. 61 = Adelobotrys adscendens, Miconia laevigata.
prasinum, Sw. Prod. Veg. Ind. Occ. 69 = Miconia prasina.
procerum, Mart. & Schrank, ex DC. Prod. iii. 187 = Tococa subciliata.
procerum, Sw. Prod. Veg. Ind. Occ. 68 = Conostegia procera.
procumbens, Vell. Fl. Flum. 180; iv. t. 124 (= Blakeae sp. ?).—Bras.
prostratum, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 220 = Dissotis prostrata.
pudibundum, Blume, Bijdr. 1071 = Creochiton pudibunda.
pulchellum, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 403 = Osbeckia rostrata.
pulcherrimum, *Korth. Verh. Nat. Gesch. Bot.* 231.—Malaya.
pulchrum, Vell. Fl. Flum. 178; iv. t. 110 = Miconia prasina.
pulverulentum, Blume, Bijdr. 1072 = Pogonathera reflexa.
pulverulentum, Jack, in Trans. Linn. Soc. xiv. (1823) 19 = Pogonathera pulverulenta.
pulverulentum, Rich. ex DC. Prod. iii. 155 = Oxymeris pulverulenta.
pumilum, Vell. Fl. Flum. 179; iv. t. 116 = Trembleya phlogiformis.
punctatum, Desr. in Lam. Encyc. iv. 50 = Miconia punctata.
punctatum, Korth. Verh. Nat. Gesch. Bot. 229 = asperum.
punctulatum, Sw. ex Wikst. in Vet. Acad. Handl. Stockh. (1827) 65 (sp. dub.).—Ins. Guadelup.
purpurascens, Aubl. Pl. Gui. i. 402 = Aciotis purpurascens.
purpurascens, Domb. ex DC. Prod. iii. 158 = Clidemia dentata.
purpurascens, Schrank & Mart. ex DC. l. c. 161 = Oxymeris purpurascens.
purpurascens, Sw. Prod. Veg. Ind. Occ. 71 = Mecranium purpurascens.

MELASTOMA:—
purpurascens, Vahl, ex DC. Prod. iii. 180 = Miconia ciliata.
purpureum, Pav. ex DC. l. c. 158 = Clidemia dentata.
purpureum, Willd. Sp. Pl. ii. 590 = Aciotis purpurascens.
pusilliflorum, Mart. & Schrank, ex DC. Prod. iii. 194 = Miconia Gardneriana.
pusillum, Blume, Mus. Bot. Lugd. Bat. i. 54.—Borneo.
pusillum, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 292 (Quid ?).—Guian.
pustulatum, Schrank & Mart. ex DC. Prod. iii. 159 = Clidemia umbonata.
pyramidale, Desr. in Lam. Encyc. iv. 53 = Miconia laevigata.
pyramidale, Mart. & Schrank, ex DC. Prod. iii. 194 = Miconia latecrenata.
quadrangulare, Sieber, ex Steud. Nom. ed. II. ii. 117 = Conostegia quadrangularis.
quadrangulare, Sw. Prod. Veg. Ind. Occ. 69 = Miconia quadrangularis.
quadrifolium, Blume, Bijdr. 1069 = Medinilla quadrifolia.
quinquedentatum, Steud. Nom. ed. II. ii. 117 = Oxymeris quinquedentata.
quinquenerve, Vell. Fl. Flum. 179; iv. t. 120 = Henriettea succosa.
quinquenervium, Mill. Gard. Dict. ed. VIII. n. 15 = Octopleura quinquenervia.
quinquenervium, Salzm. ex Griseb. Fl. Brit. W. Ind. 297 = Miconia prasina.
quinquenode, Schrank & Mart. ex DC. Prod. iii. 191 = Oxymeris quinquenodis.
racemiferum, Schrank & Mart. l. c. 155 = Miconia racemifera.
racemosum, Aubl. Pl. Gui. i. 406. t. 156 = Miconia racemosa.
radicans, Blume, Bijdr. 1069 = Medinilla radicans.
Radula, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 159.—Bras.
ramiflorum, Macfad. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 143 = Henriettella Macfadyenii.
ramiflorum, Spreng. Syst. ii. 295 = Henriettella fasciculatis.
ramiflorum, Sw. Prod. Veg. Ind. Occ. 69 = Henriettea ramiflora.
rariflorum, Humb. & Bonpl. Mélast. t. 50 = Sagraea rariflora.
reclinatum, Humb. & Bonpl. l. c. t. 48 = Miconia reclinata.
reflexum, Reinw. ex Blume, in Flora, xiv. (1831) 521 = Pogonathera reflexa.
Reinwardtianum, Blume, Bijdr. 1069 = Dissochaeta inappendiculata.
repandulum, Schrank & Mart. ex DC. Prod. iii. 194 = Miconia ligustroides.
repens, *Desr. in Lam. Encyc.* iv. 54.—China.
reptans, Schrank & Mart. ex DC. Prod. iii. 133 = Tibouchina clinopodifolia.
reticulata, Vell. Fl. Flum. 178; iv. t. 108 = Miconia staminea.
reversa, Ser. ex DC. Prod. iii. 161 = Oxymeris reversa.
revoluta, Dietr. ex Steud. Nom. ed. II. ii. 117 = Miconia granulosa.
rhaphistigma, Schrank, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 135 = Clidemia hirta.
rhexioides, Vell. Fl. Flum. 181; iv. t. 123 = Comolia denudata.
rhodocarpum, Wall. Cat. n. 4045 = zeylanica.
rhodopogon, Schrank & Mart. ex DC. Prod. iii. 163 = Oxymeris rhodopogon.
rigidum, Sw. Prod. Veg. Ind. Occ. 69 = Miconia rigida.
roseum, Poir. Encyc. Suppl. iii. 644 = repens.
rosmarinifolium, Pav. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 127 = Miconia salicifolia.
rostratum, *Bertol. Fl. Guatimal.* 17.—Guatemal.
rostratum, Blume, Bijdr. 1074 = Anplectrum rostratum.
rostratum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 221 = Osbeckia rostrata.
rotundifolium, Jack, in Trans. Linn. Soc. xiv. (1823) 11 = Phyllagathis rotundifolia.
Royenii, Blume, in Flora, xiv. (1831) 483 = polyanthum.
rubens, Sw. Prod. Veg. Ind. Occ. 71 = Miconia rubens.
rubicundum, Jack, in Trans. Linn. Soc. xiv. (1823) 18 = Medinilla rubicunda, Pogonanthera pulverulenta.
rubicundum, Wall. Cat. n. 4086 = Pogonanthera reflexa.
rubiginosum, Humb. & Bonpl. Mélast. t. 47 = Miconia rubiginosa.

MELASTOMA:—
rubiginosum, Wall. Cat. n. 4052, partim = Dissochaeta intermedia.
rubricaule, Schrank, ex DC. Prod. iii. 116 = Aciotis rubricaulis.
rubrolimbata, Link & Otto, Ic. Pl. Sel. t. 41 = Otanthera rubro-limbata.
rubrum, Aubl. Pl. Gui. i. 416 = Clidemia rubra.
rufescens, Aubl. Pl. Gui. i. 408 = Miconia rufescens.
rufidulum, Rich. ex DC. Prod. iii. 155 = Oxymeris rufescens.
rugosum, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 292 (Quid ?).—Guian.
rugosum, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 525 = Oxyspora paniculata.
rusticum, Forst. ex DC. Prod. iii. 157 = Clidemia hirta.
salicifolium, Bonpl. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 234 = Miconia salicifolia.
salicinum, *Ser. in DC. Prod.* iii. 199.—Bras.
sanguineum, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1822) 305 = Tococa sanguinea.
sanguineum, *Sims, Bot. Mag.* t. 2241.—Malaya; China.
sanguinolentum, Naud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 127 = Miconia Pavoniana.
saxatile, Dennst. Schluess. Hort. Malab. 30 = Osbeckia virgata.
saxatile, Vell. Fl. Flum. 181; iv. t. 129 = Rhynchanthera dichotoma.
scabrosum, Bert. ex DC. Prod. iii. 168 = Calycogonium stellatum.
scabrosum, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 130 = Conostegia purpurea.
scabrosum, Linn. Syst. ed. X. 1022 = Ossaea scabrosa.
scalptum, Poir. ex Steud. Nom. ed. II. ii. 117 = Ossaea scalpta.
scandens, Aubl. Pl. Gui. i. 435 = Adelobotrys scandens.
scandens, Sw. Prod. Veg. Ind. Occ. 69 = Adelobotrys adscendens.
scorpioides, *Schlecht. in Linnaea*, v. (1830) 564.—Mexic.
Sechellarum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 290 = malabathricum.
secundiflora, Steud. Nom. ed. II. ii. 117 = Clidemia secundiflora.
secundum, Mart. & Schrank, ex DC. Prod. iii. 164 = Oxymeris secundiflora.
secundum, Pav. ex DC. l. c = Oxymeris secunda.
semicrenatum, Rich. in Humb. & Bonpl. Mélast. t. 31 = Miconia semicrenata.
sepiarium, Mart. ex DC. Prod. iii. 185 = Miconia prasina.
septemnervium, *Jacq. Enum. Pl. Carib.* 20; *Stirp. Am. Select.* 131 (Quid ?).—Ins. Martinic.
septemnervium, *Lour. Fl. Cochinch.* 273 (sp. dub.).—Cochinch.
sericeum, *Vell. Fl. Flum.* 179; iv. t. 115 (Miconiae sp.?).—Bras.
serrulatum, Pav. ex DC. Prod. iii. 192 = Miconia galactantha.
serrulatum, Rich. ex DC. l. c. 177 = Miconia platyhedra.
serrulatum, Schlecht. in Linnaea, xiii. (1839) 425 = Clidemia serrulata.
serrulatum, Spreng. Syst. ii. 300 = Miconia Acinodendron.
sessile, Schum. & Thonn. Beskr. Guin. Pl. 219 = Tristemma incompletum.
sessile, Spreng. Syst. ii. 303 = Comolia sessilis.
sessiliflorum, Mart. ex DC. Prod. iii. 180 = Miconia ciliata.
sessiliflorum, Spreng. ex DC. l. c. 157 = Clidemia erythropogon.
sessiliflorum, Vahl, Eclog. Am. i. 49 = Clidemia rubra.
sessilifolium, Linn. Syst. ed. X. 1022 = Miconia Plukenetii, Henriettella sessilifolia.
setigerum, *Blume, Bijdr.* 1077.—Java.
setigerum, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 22 = Medinilla setigera.
setinode, Humb. & Bonpl. Mélast. t. 12 = Miconia setinodis.
smilacifolium, Wall. Cat. n. 4057 = Astronia smilacifolia.
solaniflorum, Schrank, ex DC. Prod. iii. 164 = Oxymeris heterobasis.
sparsiflorum, Rich. ex DC. l. c. 169 = Henriettella lateriflora.

MELASTOMA:—
speciosum, Reinw. ex Blume, in Flora, xiv. (1831) 516 = Medinilla speciosa.
spicatum, Aubl. Pl. Gui. i. 423 = Clidemia dependens.
spicatum, Pav. ex DC. Prod. iii. 160 = Miconia spondylantha.
splendens, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Charianthus glaberrimus.
splendens, Sw. Prod. Veg. Ind. Occ. 70 = Miconia splendens.
spondylanthum, Mart. ex DC. Prod. iii. 184 = Miconia spondylantha.
squamosum, Buch.-Ham. in Wall. Cat. n. 4061 = Osbeckia nepalensis.
squamulosum, Sm. in Rees, Cycl. xxiii. n. 14–15 = Miconia squamulosa.
stamineum, Desr. in Lam. Encyc. iv. 53 = Miconia staminea.
stellatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 221 = Osbeckia stellata.
stellatum, Vahl, ex DC. Prod. iii. 168 = Calycogonium stellatum.
stellulatum, Jack, in Trans. Linn. Soc. xiv. (1823) 6 = Marumia stellulata.
stenobotrys, Rich. in Humb. & Bonpl. Mélast. t. 30 = Miconia quadrangularis.
stenopetalum, Mart. & Schrank, ex DC. Prod. iii. 163 = Oxymeris dichotoma.
stenostachyum, Mart. & Schrank, l. c. 181 = Miconia stenostachya.
stipulare, Blume, Bijdr. 1073 = Anplectrum stipulare.
strangulatum, Raddi, in Mem. Soc. Ital. Modena, xx. (1829) 139 = Miconia platyhedra.
striatum, Vahl, Eclog. Am. iii. 14 = Miconia semicrenata, *Naud.*
striatum, Willd. ex Cham. in Linnaea, ix. (1834) 458 = Miconia laevigata.
strigillosum, Sw. Prod. Veg. Ind. Occ. 71 = Clidemia strigillosa.
strigosum, Linn. f. Suppl. 236 = Brachyotum strigosum.
strigosum, Rich. ex DC. Prod. iii. 161 = Oxymeris anisophylla.
strigosum, Spreng. Neue Entdeck. ii. 171 (= *Pleroma strigosum*).—Am. trop.
suaveolens, Raddi, in Mem. Soc. Ital. Modena, xx. (1829) 133 = Miconia pusilliflora.
subferrugineum, Naud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 127 = Miconia Pavonium.
subrepandum, Vell. Fl. Flum. 178; iv. t. 111 = Miconia pusilliflora?
subtriplinervium, Link & Otto, Ic. Pl. Rar. 47. t. 24 = Heeria subtriplinervia.
succosum, Aubl. Pl. Gui. i. 418 = Henriettea succosa.
succulentum, Blume, Bijdr. 1070 = Medinilla succulenta.
superbum, Bonpl. ex D. Don, in Mem. Wern. Soc. iv. (1822) 317 = Conostegia superba.
Swartzianum, Rich. in Humb. & Bonpl. Mélast. t. 33 = Miconia Fothergilla.
sylvaticum, *Blume, Bijdr.* 1077.—Java.
sylvaticum, Schlecht. in Linnaea, xiii. (1839) 422 = Miconia laevigata.
sylvestre, Mart. & Schrank, ex DC. Prod. iii. 154 = Oxymeris sylvestris.
taitense, DC. l. c. 144 = denticulatum.
Tamonea, Blanco, Fl. Filip. ed. I. 367 = imbricatum.
Tamonea, Sw. Prod. Veg. Ind. Occ. 70 = Miconia Fothergilla.
Taniboa, J. F. Gmel. Syst. 689 = Terminalia Tanibouca.
Taquari, Vell. Fl. Flum. 182; iv. t. 132 = Miconia latecrenata.
tetrandrum, Spreng. Syst. ii. 305 = Charianthus coccineus.
tetrandrum, Sw. Prod. Veg. Ind. Occ. 72 = Miconia tetrandra.
Teysmanni, *Miq. Fl. Ind. Bat. Suppl.* 316.—Sumatra.
theaefolium, G. Don, Gen. Syst. ii. 764 = Tristemma theaefolium.
theaeforme, Schrank & Mart. ex DC. Prod. iii. 185 = Miconia Maximiliana.
theaezans, Humb. & Bonpl. Mélast. t. 9 = Miconia milleflora.
thyrsoideum, Vahl, Eclog. Am. iii. 13 = Miconia globuliflora.
Tibouchina, Desr. in Lam. Encyc. iv. 49 = Tibouchina aspera.
tidorense, Blume, in Flora, xiv. (1831) 482 = polyanthum.
tiliaefolium, Schrank & Mart. ex DC. Prod. iii. 158 = Clidemia tiliaefolia.

MELASTOMA:—
tinctorium, Steud. Nom. ed. II. ii. 118 = Miconia fulva.
Tococa, Desr. in Lam. Encyc. iv. 39 = Tococa guianensis.
tomentosum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Miconia tomentosa.
tondanense, Blume, Mus. Bot. Lugd. Bat. i. 54 = polyanthum.
trachycaulon, *Miq. Fl. Ind. Bat. Suppl.* 316.—Sumatra.
trichotomum, Desr. in Lam. Encyc. iv. 52 = Miconia trichotoma.
trichotomum, Willd. ex DC. Prod. iii. 136 = Pterolepis trichotoma.
triflorum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 293.—Ind. or.
triflorum, Vahl, Eclog. Am. i. 46 = Henriettella triflora.
trinerve, Vell. Fl. Flum. 178; iv. t. 113 = Miconia paniculata.
trinervium, Salzm. ex Griseb. Fl. Brit. W. Ind. 257 = Miconia prasina.
trinervium, Sw. Prod. Veg. Ind. Occ. 69 = Miconia trinervia.
trineurum, Vell. Fl. Flum. 178; iv. t. 112 = Miconia pyrifolia.
triplinervium, G. Don, Gen. Syst. ii. 803 = Heeria subtriplinervia.
triste, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1822) 307 = Clidemia strigillosa.
triste, Rich. ex DC. Prod. iii. 162 = Miconia bracteata.
trivalve, Aubl. Pl. Gui. i. 406. t. 155 = Acisanthera trivalvis.
truncatum, Mart. & Schrank, ex DC. Prod. iii. 106 = Miconia caryophyllacea.
tunicatum, Humb. & Bonpl. Mélast. t. 60 = Miconia tunicata.
uliginosum, Bonpl. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1849) 144 = Aciotis uliginosa.
umbellatum, Mart. & Schrank, ex DC. Prod. iii. 154 = Leandra umbellata.
umbellatum, Mill. Gard. Dict. ed. VIII. n. 10 = Heterotrichum patens.
umbilicatum, Bertol. Fl. Guatimal. 16 (sp. dub.).—Guatemal.
umbonatum, Mart. & Schrank, ex DC. Prod. iii. 158 = Clidemia umbonata.
umbrosum, Spreng. Syst. ii. 304 = Sagraea latifolia.
umbrosum, Sw. Prod. Veg. Ind. Occ. 72 = Sagraea umbrosa.
urceolatum, Mart. & Schrank, ex DC. Prod. iii. 158 = Clidemia neglecta.
vaccinioides, Humb. & Bonpl. Melast. t. 8 = Miconia vaccinioides.
vacillans, Blume, Bijdr. 1074 = Dissochaeta inappendiculata.
vagans, Roxb. Hort. Beng. 33; Fl. Ind. ii. 404 = Oxyspora vagans.
varingiaefolium, Blume, Bijdr. 1071 = Pachycentria varingiaefolia.
velutinum, Mart. & Schrank, ex DC. Prod. iii. 182 = Miconia fallax.
velutinum, Seem. Fl. Vit. 90 = normale.
velutinum, Willd. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 116 = Miconia holosericea.
velutinum, Willd. Sp. Pl. ii. 584 = Tibouchina holosericea.
venosum, Blume, Bijdr. 1075 = Medinilla venosa.
verrucosum, Blume, l. c. 1078 = Medinilla verrucosa.
verticillatum, Mill. Gard. Dict. ed. VIII. n. 12 = Clidemia rubra.
verticillatum, Rich. ex DC. Prod. iii. 155 = Oxymeris rufescens.
verticillatum, Vahl, Eclog. Am. i. 47 = Ossaea verticillata.
vesiculosum, Sessé, ex DC. Prod. iii. 166 = Calophysa vesiculosa.
villosum, Aubl. Pl. Gui. i. 428. t. 168 = Desmoscelis villosa.
villosum, *Sims, Bot. Mag.* t. 2630.—Cochinch.
viminale, Jack, in Trans. Linn. Soc. xiv. (1823) 16 = Anplectrum viminale.
violaceum, Rich. ex DC. Prod. iii. 162 = Miconia spondylantha.
virescens, Vahl, Eclog. Am. iii. 18 = Miconia virescens.
virgatum, Sw. Prod. Veg. Ind. Occ. 72 = Mecranium virgatum.
virusana, Comm. ex Juss. Gen. 329 = Tristemma virusanum.

MELASTOMA:—
viscosum, Bonpl. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 146 = Aciotis viscosa.
vitiense, Naud. l. c. xiii. (1849) 275 = denticulatum.
volubile, Bonpl. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 35 = Arthrostema volubile.
volubile, Schrank, ex DC. Prod. iii. 157 = Octopleura Epibaterium.
Wallichii, DC. Prod. iii. 146 = normale.
xalapense, Humb. & Bonpl. Mélast. t. 54 = Conostegia xalapensis.
Xantholasia, Schrank & Mart. ex DC. Prod. iii. 163 = Oxymeris Xantholasia.
zeylanicum, Pers. Syn. i. 472 = Osbeckia octandra.
Zollingeri, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 277 = nitidum.

MELASTOMASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 296. t. 5 = **Dissotis**, Benth. (Melastom.).
erectum, Naud. l. c. = D. capitata.

MELAXIS, Steud. Nom. ed. II. i. 118 (1841), sphalm. = MALAXIS, Sm. = **Liparis**, Rich. (Orchid.).
lancifolia, Steud. l. c. = L. paradoxa.

MELEAGRINEX, Arruda, ex Koster, Trav. Braz. 496 (1816) (Quid?).
pernambucana, Arruda, l. c.—Bras.
pernambucea, Arruda, ex Steud. Nom. ed. II. ii. 118 = praec.

MELFONA, Rafin. Sylva Tellur. 103 (1838) = **Cuphea**, P. Br. (Lythrac.).
purpurea, Rafin. l. c. = C. viscosissima.

MELHANIA, Forsk. Fl. Aegypt. Arab. 64 (1775).
STERCULIACEAE, Benth. & Hook. f. i. 222.
BROTERA, Cav. Ic. v. 19. t. 433 (1799).
CARDIOSTEGIA, Presl, Epim. Bot. 249 (1850).
PENTAGLOTTIS, Wall. Cat. n. 1156 (1831).
SPRENGELIA, Schult. Obs. Bot. 134 (1804).
VIALIA, Vis. Ind. Sem. Hort. Patav. (1840).—Cf. Linnaea, xv. (1841) Litt. 103.
abutiloides, Aitch. Cat. Pl. Punjab, 22, partim = tomentosa.
abutiloides, Arn. ex Wight, Ic. t. 23 = Hamiltoniana.
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 76.—Sind; Afr. trop.
acuminata, *Mast. in Oliver, Fl. Trop. Afr.* i. 231.—Afr. trop.
bracteosa, Boiss. Fl. Orient. i. 841 = Denhamii.
Burchellii, *DC. Prod.* i. 499.—Afr. austr.
cannabina, *Wight, ex Mast. in Hook. f. Fl. Brit. Ind.* i. 372.—Ind. or.
chrysantha, E. Mey. in Drège, Zwei Pfl. Docum. 201, nomen = Mahernia chrysantha.
corchoriflora, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 501.—Madag.
cyclophylla, *Hochst. ex Mast. in Oliver, Fl. Trop. Afr.* i. 230.—Abyssin.
damarana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 590.—Afr. austr.
decanthera, DC. Prod. i. 499 = Trochetia decanthera.
Denhamii, *R. Br. in Denh. & Clapp. Trav. App.* 232.—Sind; Afr. trop.
didyma, *Eckl. & Zeyh. Enum.* 52.—Afr. austr.
Erythroxylon, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 146.—Ins. S. Helen.
ferruginea, *A. Rich. Tent. Fl. Abyss.* i. 76.—Abyssin.
Forbesii, *Planch. ex Mast. in Oliver, Fl. Trop. Afr.* i. 231.—Afr. trop.
futteyporensis, *Munro, ex Mast. in Hook. f. Fl. Brit. Ind.* i. 373.—Ind. bor. occ.
Hamiltoniana, Munro, Hort. Agrens. 9 = futteyporensis.
Hamiltoniana, *Wall. Pl. As. Rar.* i. 69.—Ind. or.; Burma.
incana, *Heyne, in Wall. Cat.* n. 1200.—Ind. or.; Austral.
Kotschyi, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = Denhamii.
laurifolia, Boj. in Ann. Sc. Nat. Sér. II. xviii. (1842) 192 = Trochetia laurifolia.
Leprieurii, Webb, in Hook. Niger Fl. 113. tt. 4, 5 = abyssinica.
leucantha, E. Mey. in Drège, Zwei Pfl. Docum. 201, nomen = didyma.
linearifolia, *Sond. in Linnaea*, xxiii. (1850) 18.—Afr. austr.

MELHANIA:—
macrophylla, *Vis. Sem. Hort. Patav.* (1855).—*Cf. Linnaea*, xxviii. (1856) 366, in obs.—Hab.?
Melanoxylon, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 146.—Ins. S. Helen.
muricata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 503.—Ins. Socotra.
oblongata, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 232 = abyssinica.
oblongifolia, F. Muell. Fragm. i. 69 = incana.
ovata, Spreng. Syst. iii. 32 = abyssinica.
prostrata, *Burch. Trav.* ii. 263.—Afr. austr.
reniformis, *Hochst. & Steud. ex Steud. Nom.* ed. II. ii. 118.—Arab.
rotundata, *Hochst. ex Mast. in Oliver, Fl. Trop. Afr.* i. 230.—Abyssin.
rupestris, *Wight, in Hook. Journ. Bot.* iii. (1841) 163.—Ind. or.
sidaeflora, Span. in Linnaea, xv. (1841) 174 = incana.
sidaefolia, Stocks, ex Boiss. Fl. Orient. i. 841 = Denhamii.
Steudneri, *Schweinf. in Verh. Zool. Bot. Ges. Wien*, xviii. (1868) 672.—Afr. trop.
tomentosa, *Stocks, ex Mast. in Hook. f. Fl. Brit. Ind.* i. 373.—Ind. bor. occ.
velutina, *Forsk. Fl. Aegypt. Arab.* 64.—Arab.

MELIA, Linn. Gen. ed. I. 127 (1737). *MELIACEAE*, Benth. & Hook. f. i. 332.
AZADIRACHTA, A. Juss. in Mém. Mus. Par. xix. (1830) 220.
AZEDARAC, Adans. Fam. ii. 342 (1763).
aethiopica, Welw. Apont. 584 = dubia.
angustifolia, Schum. & Thonn. Beskr. Guin. Pl. 214 = M. Azedarach.
argentea, Buch.-Ham. ex Wall. Cat. p. 214. n. 1254 C = dubia.
arguta, *DC. Prod.* i. 622.—Ins. Molucc.
australasica, Juss. in Mém. Mus. Par. xix. (1830) 257 = dubia.
australis, Sweet, Hort. Brit. ed. II. 85 = M. Azedarach.
Azadirachta, *Linn. Sp. Pl.* 385.—Ind. or.
Azedarach, Blanco, Fl. Filip. ed. I. 345 = dubia.
Azedarach, *Linn. Sp. Pl.* 384.—Reg. Himal. (alibi cult.).
baccata, Heyne, ex Wall. Cat. sub n. 1256 = seq.
baccifera, Roth, Nov. Pl. Sp. 215 = Cipadessa fruticosa.
Bambolo, Welw. Apont. 561 = dubia.
birmanica, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 183.—Burma.
Bukayun, Royle, Illustr. Bot. Himal. 144 = M. Azedarach.
Candollei, *Juss. in Mém. Mus. Par.* xix. (1830) 258.—Ins. Timor.
chinensis, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 23 = M. Toosendan.
cochinchinensis, M. Roem. Syn. Hesper. 95 = M. Azedarach.
Commelini, Medic. ex Steud. Nom. ed. II. ii. 118 = M. Azedarach.
composita, Benth. Fl. Austral. i. 380 = M. Azedarach.
composita, DC. Prod. i. 622 = M. Candollei.
composita, Willd. Sp. Pl. ii. 559 = dubia.
dubia, *Cav. Diss.* vii. 364.—As. Afr. & Austral. trop.
dubia, Hiern, in Hook. f. Fl. Brit. Ind. i. 545, partim = birmanica.
elegans, *Seem. Fl. Vit.* 36.—Ins. Fiji.
excelsa, *Jack, in Malay. Misc.* i. (1820) n. 1. 12.—Malaya.
flaccida, Zipp. ex Span. in Linnaea, xv. (1841) 182 = Candollei.
floribunda, *Carr. in Rev. Hortic.* (1872) 470.—Hab.?
florida, Salisb. Prod. 317 = M. Azedarach.
fraxinifolia, Salisb. l. c. = M. Azadirachta.
Guara, Jacq. Enum. Pl. Carib. 220 = Guarea trichilioides.
guineensis, G. Don, in Loud. Hort. Brit. 168 = M. Azedarach.
Iloilo, Blanco, Fl. Filip. ed. II. 241 = Aglaia argentea.
indica, Brand. For. Fl. Ind. 67 = M. Azadirachta.
integerrima, *Buch.-Ham. in Trans. Linn. Soc.* xvii. (1835) 231.—Ind. or.
japonica, G. Don, Gen. Syst. i. 680 = M. Azedarach.
japonica, Hassk. Cat. Hort. Bogor. Alt. 219 = M. Azadirachta.
javanica, M. Roem. Syn. Hesper. 96 = praec.
Koetjape, Burm. f. Fl. Ind. 101 = Sandoricum indicum.
montana, Herb. Madr. ex Wall. Cat. p. 214. 1256 D = Cipadessa fruticosa.

MELIA:—
neilgherrica, Walp. Rep. iv. 551, sphalm. = Munronia neilgherrica.
orientalis, M. Roem. Syn. Hesper. 95 = M. Azedarach.
parviflora, Moon, Cat. Pl. Ceyl. 35 = M. Azadirachta.
pendula, Reinw. ex C. DC. in DC. Monog. Phan. i. 529 = Chisocheton patens.
penduliflora, Wall. Cat. n. 1255 = Chisocheton penduliflorus.
pinnata, Stokes, Bot. Mat. Med. ii. 482 = M. Azadirachta.
pumila, Moon, Cat. Pl. Ceyl. 35 = Munronia pumila.
robusta, Roxb. Hort. Beng. 33; Fl. Ind. ii. 397 = dubia.
sambucina, Blume, Bijdr. 162 = M. Azedarach.
sempervirens, Sw. Prod Veg. Ind. Occ. 67 = M. Azedarach.
superba, Roxb. Hort. Beng. 33; Fl. Ind. ii. 396 = dubia.
tomentosa, Kurz, Rep. Veg. Andam. ed. I. 4 = Chickrassia tabularis.
tomentosa, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 394.—Penang.
Toosendan, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 159.—Japon.

MELIANTHUS, Linn. Syst. ed. I (1735). *SAPINDACEAE*, Benth. & Hook. f. i. 411,
DIPLERISMA, Planch. in Trans. Linn. Soc. xx. (1851) 416. f. 15–20.
comosus, *Vahl, Symb. Bot.* iii. 86.—Afr. austr.
Dregeanus, *Sond. in Harv. & Sond. Fl. Cap.* i. 368.—Afr. austr.
himalayanus, Wall. Cat. n. 1190 = major.
major, *Linn. Sp. Pl.* 639.—Afr. austr.
minor, *Linn. l. c.*—Afr. austr.
pectinatus, *Harv. in Harv. & Sond. Fl. Cap.* i. *Add. sub praef.* 21*—Afr. austr.
sibiricus, *Pall. ex Georgi, Beschr. Russ. Reich.* III. iv. 1114 (Quid?).—Sibir.
Trimenianus, Hook. f. in Journ. Bot. xi. (1873) 353 = pectinatus.

MELICA, Franch. & Sav. Enum. Pl. Jap. i. 368, in syn. (1875) sphalm. = **Melissa**, Tourn. (Labiat.).

MELICA, Linn. Fl. Lappon. 23 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1189.
CHONDRACHYRUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836).
DALUCUM, Adans. Fam. ii. 323 (1763).
acuminata, *Boland. in Proc. Calif. Acad.* iv. (1870) 104.—Am. bor. occ.
alata, *Nees, ex Steud. Syn. Pl. Gram.* 290.—Chili.
altissima, *Linn. Sp. Pl.* 66.—Europ. austr.; As. bor.
altissima, Sobol. Fl. Petrop. 21 = Glyceria aquatica.
altissima, Walt. Fl. Carol. 78 = diffusa.
amethystina, Pourr. in Mém. Acad. Toul. iii. (1788) 322 = Bauhini.
ampla, *Phil. in Linnaea*, xxxiii. (1864–65) 292.—Chili.
angustifolia, Boiss. & Blanch. Diagn. Ser. II. iv. 131 = minuta.
aquatica, Weber, ex Beauv. Agrost. 167 = Glyceria aquatica.
argentata, *E. Desv. in C. Gay, Fl. Chil.* vi. 374.—Chili.
argentea, C. Muell. in Walp. Ann. vi. 1024, sphalm. = praec.
aristata, *Thurb. ex Boland. in Proc. Calif. Acad.* iv. (1870) 103.—Calif.
armena, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 585 = minuta.
armena, Boiss. & Kotschy, l. c. 590 = Cupani.
arrecta, Kunze, in Flora, xxix. (1846) 740 = minuta.
arundinacea, Raeusch. Nom. ed. III. 24 = caerulea.
aspera, Desf. Fl. Atlant. i. 71 = minuta.
asperrima, Hort. ex Roem. & Schult. Syst. ii. 530 = exasperata.
aurantiaca, *Desr. in Lam. Encyc.* iv. 70.—Reg. Argent.
australis, Coss. ex Parl. Fl. Ital. i. 306 = minuta.
baetica, Bout. ex Willk. & Lange, Prod. Fl. Hisp. i. 84 = minuta.
Balansae, Boiss. & Bal. Diagn. Ser. II. iv. 132 = ciliata.
barbata, Moon, Cat. Pl. Ceyl. 8 = Chloris barbata.
Bauhini, *All. Auct. Pedem.* 43.—Reg. Mediterr.
Bauhini, Guss. Fl. Sic. Prod. i. 66 = Cupani.
Bauhini, Tenore, Syll. Fl. Neap. 37 = ciliata.

MELICA:—
Bossieri, Reut. ex Boiss. Fl. Orient. v. 589 = ciliata.
brasiliana, Ard. Animad. ii. 17 = papilionacea.
breviflora, Boiss. Diagn. Ser. I. vii. 124 = Cupani.
bromoides, *A. Gray, in Proc. Am. Acad.* viii. (1873) 409, in nota.—Calif.
bulbosa, *Geyer, ex Hook. Kew Journ. Bot.* viii. (1856) 19, nomen.—Am. bor. occ.
bulbosa, Porter & Coulter, Fl. Colorado, 149 = spectabilis.
bulbosa, Thurb. in Bot. Calif. ii. 304 = californica.
caerulea, Linn. Mant. ii. 325 = Molinia caerulea.
caespitosa, Schur, Sert. Fl. Transs. 86; ex eod. in Oestr. Bot. Zeitschr. ix. (1859) 329 = ciliata.
caffrorum, *Schrad. in Goett. Gel. Anz.* (1821) III. 2073.—Afr. austr.
californica, *Scribner, in Proc. Acad. Philad.* (1885) 46.—Am. bor. occ.
calycina, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Poa calycina.
capensis, Thunb. Prod. Pl. Cap. 21 = Ehrharta melicoides.
capillaris, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 243.—Syria.
cappadocica, Boiss. Fl. Orient. v. 589 = ciliata.
caricina, Urv. in Mém. Soc. Linn. Par. i. (1822) 263 = minuta.
chilensis, *J. & C. Presl, Rel. Haenk.* i. 270.—Chili.
ciliata, *Linn. Sp. Pl.* 66.—Europ.; Afr. bor.
cimicina, Salisb. Prod. 20 = Panicum cimicinum.
coerulea, Auct. = *caerulea*.
colorata, Dulac, Fl. Hautes-Pyr. 80 = ciliata.
colpodioides, Nees, in Ann. Nat. Hist. i. (1838) 283 = imperfecta.
Commersoni, *Nees, ex Steud. Syn. Pl. Gram.* 290.—Chili.
commutata, *Steud. l. c.*—Chili.
cretica, Boiss. & Heldr. Diagn. Ser. I. xiii. 54 = ciliata.
cristata, Trin. Fund. Agrost. 163 = ciliata.
Cupani, *Guss. Suppl. Fl. Sic. Prod.* 17.—Reg. Mediterr.; Oriens; Himal.
cynosuroides, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 42 = Enteropogon melicoides.
cyrenaica, Viv. Fl. Lib. Spec. 4. t. 25. f. 1 = minuta.
decumbens, *Thunb. Prod. Pl. Cap.* 21.—Afr. austr.
decumbens, Weber, Spicil. Fl. Gotting. 3 = Triodia decumbens.
dendroides, *Lehm. Pugill.* iii. 39.—Afr. austr.
diandra, Roxb. Hort. Beng. 8; Fl. Ind. i. 327 = Centotheca lappacea?
diffusa, *Pursh, Fl. Am. Sept.* i. 77.—Am. bor.
digitata, Roxb. Hort. Beng. 82; Fl. Ind. i. 326 = Chloris digitata.
divaricata, Meig. & Wenig. Syst. Verz. Rheins, 16 = Molinia caerulea.
effusa, Salisb. Prod. 20 = Milium effusum.
eligulata, Boiss. Diagn. Ser. I. xiii. 56 = Cupani.
exaltata, *Phil. in Anal. Univ. Chil.* (1873) 571.—Chili.
exasperata, *Roem. & Schult. Syst.* ii. 530.—Hab.?
expansa, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Falx, Linn. f. Suppl. 109 = Harpechloa capensis.
festucoides, Licht. ex Roem. & Schult. Syst. ii. 530 = Ehrharta panicea.
festucoides, *Willd. ex Steud. Nom.* ed. II. ii. 119, nomen.—Afr. austr.
filiculmis, *E. Desv. in C. Gay, Fl. Chil.* vi. 378.—Chili.
Fischeri, Spreng. Syst. iv. Cur. Post. 32 = Graphephorum Fischeri.
flava, *Nees, ex Steud. Syn. Pl. Gram.* 290.—Chili.
fluitans, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Glyceria fluitans.
frutescens, *Scribner, in Proc. Acad. Philad.* (1885) 45.—Calif.
fugax, *Boland. in Proc. Calif. Acad.* iv. (1870) 104.—Calif.
geniculata, Thunb. Prod. Pl. Cap. 21 = Ehrharta calycina.
Geyeri, Munro, ex Boland. in Proc. Calif. Acad. iv. (1870) 103 = bromoides.
Geyeri, Thurb. in Bot. U. St. Expl. Exped. ii. 492 = fugax.
gigantea, Thurb. Prod. Pl. Cap. 21 = Ehrharta gigantea.
glabra, Michx. Fl. Bor. Am. i. 62, partim = diffusa.
glabra, Pursh, Fl. Am. Sept. i. 77 = mutica.
glabra, *Steud. Nom.* ed. II. ii. 119; *et Syn. Pl. Gram.* 290.—Chili.

MELICA :—
glauca, F. Schultz, in Flora, xlv. (1862) 462 = ciliata.
glaucescens, Steud. Syn. Pl. Gram. 289 = Cupani.
Gmelini, Roth, Tent. Fl. Germ. ii. 1. 104 = Koeleria cristata.
Gmelini, *Turc. ex Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 368.—Sibir.
gracilis, Aitch. & Hemsl. in Journ. Linn. Soc. xix. (1882) 192 = secunda.
Hallii, *Vasey, in Coult. Bot. Gaz.* vi. (1881) 296.—Am. bor. occ.
Harfordii, *Boland. in Proc. Calif. Acad.* iv. (1870) 102.—Calif.
hirsuta, Koel. Desc. Gram. 144 = Koeleria cristata.
hirta, *Phil. in Linnaea*, xxxiii. (1864–65) 292.—Chili.
Hohenackeri, Boiss. Diagn. Ser. I. xiii. 54 = Cupani.
humilis, Boiss. Voy. Bot. Espagne, ii. 662 = Cupani.
hyalina, *Doell, in Mart. Fl. Bras.* ii. II. 127.—Bras.
imperfecta, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Calif.
imperforata, Nees, in Hook. & Arn. Bot. Beech. Voy. 403 = praec.
inaequiglumis, Boiss. Diagn. Ser. I. vii. 124 = Cupani.
interrupta, Reichb.? ex Schur, in Oestr. Bot. Zeitschr. ix. (1859) 329 = ciliata.
Jacquemontii, Decne. in Jacquem. Voy. Bot. 174. t. 175 = Cupani.
Kotschyi, Hochst. ex Steud. Syn. Pl. Gram. 289 = Cupani.
lanata, Steud. l. c. = Cupani.
lanuginosa, *Desf. Tabl.* ed. I. 15, nomen.—Hab.?
lappacea, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Centotheca lappacea.
latifolia, Roxb. Hort. Beng. 8; Fl. Ind. i. 328 = Thysanolaena Agrostis.
laxiflora, Boiss. & Blanch. ex Boiss. Fl. Orient. v. 589 = ciliata.
laxiflora, *Cav. Ic.* v. 48. t. 473. f. 2.—Chili.
litoralis, *Phil. in Anal. Univ. Chil.* (1873) 572.—Chili.
littoralis, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Aeluropus laevis.
lobata, Schur, Enum. Pl. Transs. 765 = ciliata.
Lobelii, Vill. Fl. Delph. 6; Hist. Pl. Dauph. ii. 89 = uniflora.
longiflora, *Steud. Nom.* ed. II. ii. 119; *Syn. Pl. Gram.* 290.—Chili.
macra, *Nees, Agrost. Bras.* 486.—Reg. Argent.
magellanica, Désr. in Lam. Encyc. iv. 72 = Hierochloa redolens.
Magnolii, Gren. & Godr. Fl. Fr. iii. 550 = ciliata.
major, Sibth. & Sm. Fl. Graec. Prod. i. 51 = minuta.
micrantha, Boiss. & Hohen. Diagn. Ser. I. xiii. 53 = ciliata.
minor, *Hack. ex Boiss. Fl. Orient.* v. 586.—Reg. Caucas.
minuta, *Linn. Mant.* i. 32.—Europ. austr.; Persia.
montana, Cav. Desc. 44 = minuta.
montana, Huds. Fl. Angl. ed. II. 37 = nutans.
Muehlenbergiana, Schult. Mant. ii. 294 = mutica.
Munroi, Boiss. Fl. Orient. v. 587 = secunda.
mutica, *Walt. Fl. Carol.* 78.—Am. bor.
nebrodensis, Parl. Giorn. Bot. Ital. ii. 1. (1844) 198 = ciliata.
nervosa, *Moon, Cat. Pl. Ceyl.* 8, nomen.—Ceylon.
nitens, Nutt. ex Scribner, in Proc. Acad. Philad. (1885) 44 = diffusa.
nodosa, Pill. & Mitterp. It. Posegan. 143 = Diplachne serotina.
nodosa, Turcz. ex Ledeb. Fl. Ross. iv. 398 = virgata.
nutans, Cav. Ic. ii. 58. t. 175. f. 2 = minuta.
nutans, Lam. Tabl. Encyc. i. 179 = uniflora.
nutans, *Linn. Sp. Pl.* 66.—Europ.; As. bor.
Onoei, *Franch. & Sav. Enum. Pl. Jap.* ii. 603.—Japon.
ovalis, *Nees, Fl. Afr. Austr.* i. 417.—Afr. austr.
pallida, *H. B. & K. Nov. Gen. et Sp.* i. 164. t. 698.—Ecuador.
panicoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 188 = imperfecta.
pannosa, Boiss. Diagn. Ser. I. xiii. 55 = Cupani.
papilionacea, *Linn. Mant.* i. 31.—Bras.; Mexic.
penicillaris, *Boiss. & Bal. Diagn.* Ser. II. iv. 132.—Syria.
persica, Kunth, Rév. Gram. i. 351 = Cupani.
picta, *C. Koch, in Linnaea*, xxi. (1848) 395.—Reg. Caucas.
poaeoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 188 = imperfecta.
poaeoides, Torr. in Pacif. Rail. Rep. iv. 157 = californica.

MELICA :—
poecilantha, *E. Desv. in C. Gay, Fl. Chil.* vi. 379.—Chili.
Porteri, *Scribner, in Proc. Acad. Philad.* (1885) 44; *et ex Coult. Man. Bot. Rocky Mount.* 419.—Am. bor. occ.
provincialis, Clar. ex Steud. Nom. ed. II. ii. 119 = ciliata.
pubescens, Desv. Opusc. 65 = ciliata.
pyramidalis, Lam. Fl. Fr. iii. 585 = minuta.
racemosa, Muhl. Desc. Gram. 88 = mutica.
racemosa, *Thunb. Prod. Pl. Cap.* 21.—Afr. austr.
Radula, *Franch. Pl. Davidian.* i. (1884) 336.—China.
ramosa, Thunb. Prod. Pl. Cap. 21 = Ehrharta ramosa.
ramosa, Vill. Hist. Pl. Dauph. ii. 91 = minuta.
rariflora, Schreb. Beschr. Graes. ii. 157 = mutica.
rectiflora, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 56.—Ins. Creta.
refracta, Roxb. Hort. Beng. 8; Fl. Ind. i. 327 = Centotheca lappacea.
reptans, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Aeluropus laevis?
rigida, *Cav. Ic.* v. 47. t. 473. f. 1.—Reg. Argent.
rigida, Wibel, Prim. Fl. Werth. 117 = Triodia decumbens.
Sabinii, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 443 = Pleuropogon Sabini.
sarmentosa, *Nees, Agrost. Bras.* 485.—Bras.
saxatilis, Sibth. & Sm. Fl. Graec. Prod. i. 51 = minuta.
scabra, *H. B. & K. Nov. Gen. et Sp.* i. 164.—Ecuador.
scabra, Nutt. in Trans. Am. Phil. Soc. v. (1837) 148 = diffusa.
scabrosa, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1838) 59.—China.
Schimperi, Presl, Bot. Bemerk. 120, nomen = Cupani.
secunda, *Regel, in Act. Hort. Petrop.* vii. (1880) 629.—Afghan.; Turkestan.
setacea, Pers. Syn. i. 78 (err. typ. 73) = Bauhini.
sibirica, Lam. Tabl. Encyc. i. 179 = altissima.
sinaica, De Not. ex Steud. Syn. Pl. Gram. 427 = Cupani.
speciosa, Muhl. in Trans. Phil. Soc. iii. (1793) 161 = mutica.
spectabilis, Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 42 = papilionacea.
spectabilis, *Scribner, in Proc. Acad. Philad.* (1885) 45.—Am. bor. occ.
spinosa, Munro, ex Boiss. Fl. Orient. v. 587 = secunda.
stricta, *Boland. in Proc. Calif. Acad.* iii. (1863) 4; *et* iv. (1870) 104.—Am. bor. occ.
stricta, Brandegee, in Bull. Geol. & Geog. Surv. Territ. ii. n. III. (1876) 244 = Porteri.
subenervis, Boiss. Fl. Orient. v. 586 = minor.
subulata, Scribner, in Proc. Acad. Philad. (1885) 47 = acuminata.
sylvatica, Link, ex Steud. Nom. ed. I. 520 = Molinia caerulea.
taurica, C. Koch, in Linnaea, xxi. (1848) 395 = ciliata.
Tinei, Lojac. in Natur. Sic. i. (1882) 9 = nebrodensis.
Torreyana, *Scribner, in Proc. Acad. Philad.* (1885) 43.—Calif.
trachyantha, Boiss. Diagn. Ser. I. vii. 125 = Cupani.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 86 = ciliata.
triflora, *Bigel. in New Engl. Med. Journ.* v. (1816) 334.—Am. bor.
typhina, Bor. in Mém. Soc. Acad. Maine-et-Loire, i. (1857) 91 = ciliata.
uniflora, *Retz. Obs.* i. 10.—Europ.; Oriens.
valdiviana, *Phil. in Anal. Univ. Chil.* (1873) 572.—Chili.
variabilis, Schur, Enum. Pl. Transs. 936 (Index) = Molinia caerulea.
vestita, Boiss. Diagn. Ser. I. vii. 125 = Cupani.
violacea, *Cav. Ic.* v. 47.—Chili.
virgata, *Turcz. ex Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 369.—Sibir.
viridiflora, Czern. Consp. Fl. Ucran. (1859); ex Nym. Consp. 818 = picta.
viridis, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 590 = Cupani.

MELICHO, Salisb. Gen. Pl. Fragm. 130 (1866) = **Haemanthus**, Linn. (Amaryll.).
amarylloides, Salisb. l. c. = H. amarylloides.
carneus, Salisb. l. c. = H. carneus.
humilis, Salisb. l. c. = H. humilis.
lanceaefolia, Salisb. l. c. = H. lanceaefolius.
Pumilio, Salisb. l. c. = H. Pumilio.
sanguineus, Salisb. l. c. = H. sanguineus.

MELICHRUS, R. Br. Prod. 539 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 612.
VINTENATIA, Cav. Ic. iv. 28. t. 349. f. 1 (1797) partim.
adpressus, A. Cunn. ex DC. Prod. vii. 740 = urceolatus.
erubescens, A. Cunn. l. c. = urceolatus.
medius, A. Cunn. in Field, N. S. Wales, 341 = urceolatus.
rotatus, *R. Br. Prod.* 539.—Austral.
squarrosus, Sond. in Lehm. Pl. Preiss. i. 304 = Andersonia depressa?
urceolatus, *R. Br. Prod.* 539.—Austral.

MELICLIS, Rafin. Fl. Tellur. ii. 99 (1836) = **Coryanthes**, Hook. (Orchid.).
speciosa, Rafin. l. c. 100 = C. speciosa.

MELICOCCA, Linn. Sp. Pl. ed. II. 495 (1763). *SAPINDACEAE*, Benth. & Hook. f. i. 401.
CASIMIRA, Scop. Introd. 233 (1777).
amoena, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 139.—Java.
apetala, Poir. Encyc. Suppl. iii. 224 = Doratoxylon diversifolium.
australis, *Steud. Nom.* ed. II. ii. 120 (sp. dub.).—Austral.
bijuga, *Linn. Sp. Pl.* ed. II. 495.—Am. trop.
carpoodea, Juss. in Mém. Mus. Par. iii. (1817) 187. t. 4 = bijuga.
dentata, *Juss. l. c.*—Ins. Maurit.
diphylla, Boj. Hort. Maurit. 56 = seq.?
diversifolia, Juss. in Mém. Mus. Par. iii. (1817) 187. t. 7 = Doratoxylon diversifolium.
geniculata, *Spreng. Syst.* ii. 220.—Bras.
lepidopetala, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 344.—Bolivia.
olivaeformis, H. B. & K. Nov. Gen. et Sp. v. 130 = Talisia olivaeformis.
paniculata, Juss. in Mém. Mus. Par. iii. (1817) 187. t. 5 = Hypelate paniculata.
pubescens, DC. Prod. i. 615 = Schleichera trijuga.
trijuga, Juss. in Mém. Mus. Par. iii. (1817) 187. t. 8 = Schleichera trijuga.
triptera, Blanco, Fl. Filip. ed. II. 203 = Tristira triptera.

MELICOCCUS, P. Br. Hist. Jamaic. 210 (1756) = **Melicocca**, Linn. (Sapind.).
bijugatus, Jacq. Enum. Pl. Carib. 19 = Melicocca bijuga.

MELICOPE, Forst. Char. Gen. 55 (1776). *RUTACEAE*, Benth. & Hook. f. i. 295.
ASTORGANTHUS, Endl. Cat. Hort. Vindob. ii. 196 (1843).
BOUCHARDATIA, Baill. Adansonia, vii. (1867) 350.
ENTOGANUM, Banks, ex Gaertn. Fruct. i. 331. t. 68 (1788).
MALICOPE, Vitm. Summa Veg. ii. 442 (1789).
PELEA, A. Gray, in Bot. U. St. Expl. Exped. 339. tt. 35, 38 (1854).
acronychioides, F. Muell. Fragm. iv. 117 = Evodia acronychioides.
auriculata, *Nadeaud, Enum. Pl. Tahiti*, 76.—Ins. Tahiti.
australasica, *F. Muell. Fragm.* ii. 102.—Austral.
barbigera, *A. Gray, Bot. U. St. Expl. Exped.* i. 351. t. 39.—Ins. Sandvic.
cinerea, *A. Gray, l. c.* 350. t. 39.—Ins. Sandvic.
contermina, *C. Moore & Muell. ex F. Muell. Fragm.* vii. 144.—Ins. Dom. Howe.
elliptica, *A. Gray, Bot. U. St. Expl. Exped.* i. 353.—Ins. Sandvic.
erythrococca, *Benth. Fl. Austral.* i. 360.—Austral.
grandifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 354.—Ins. Sandvic.
Helferi, *Hook. f. Fl. Brit. Ind.* i. 492.—Burma.
indica, *Wight, Ic.* t. 1051.—Ind. or.
leguminosa, *Nadeaud, Enum. Pl. Tahiti*, 76.—Ins. Tahiti.
Mantellii, *J. Buch. in Trans. N. Z. Inst.* iii. (1871) 212.—N. Zel.
neurococca, *Benth. Fl. Austral.* i. 360.—Austral.
Pagetii, F. Muell. Fragm. v. 178 = Pagetia medicinalis.
simplex, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 315.—N. Zel.
spathulata, *A. Gray, Bot. U. St. Expl. Exped.* i. 352.—Ins. Sandvic.

MELICOPE:—
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 75.—Ins. Tahiti.
ternata, *Forst. Char. Gen.* 56.—N. Zel.
tetrandra, Blanco, Fl. Filip. ed. I. 293 = Evodia triphylla.
tetrandra, Roxb. Hort. Beng. 88; Fl. Ind. ii. 257 = Tetractomia Roxburghii.

MELICOPSIDIUM, Baill. Adansonia, xi. (1874) 243 = **Cossinia**, Comm. (Sapind.).
trifoliatum, Baill. l. c. = C. trifoliata.

MELICYTUS, Forst. Char. Gen. 123. t. 62 (1776). *VIOLARIEAE*, Benth. & Hook. f. i. 119.
TRACHYTIS, Soland. ex Gaertn. Fruct. i. 205 (1788).
collinus, *Colenso, Excurs. N. Island*, 85, nomen.—N. Zel.
lanceolatus, *Hook. f. Fl. N. Zel.* i. 18. t. 8.—N. Zel.
macrophyllus, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 256.—N. Zel.
micranthus, *Hook. f. Fl. N. Zel.* i. 18.—N. Zel.
Oleaster, Lindl. in Mitch. Journ. Trop. Austral. 383 = Denhamia Oleaster.
ramiflorus, *Forst. Char. Gen.* 124.—N. Zel.
umbellatus, Gaertn. Fruct. i. 206. t. 44. f. 1 = ramiflorus.

MELIDISCUS, Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Cappar.).
gigantea, Rafin. l. c. = C. gigantea.

MELIDORA, Noronha, ex Salisb. in Trans. Hort. Soc. ii. (1817) 156 = **Enkianthus**, Lour. (Ericac.).
pellucida, Noronha, ex Salisb. l. c. = E. quinqueflorus.

MELILOBUS, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) App. 215 = **Gleditschia**, Linn. (Legumin.).
heterophyla, Rafin. Sylva Tellur. 121 = G. triacanthos.

MELILOTA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 382 = **Melilotus**, Tourn. (Legumin.).
messanensis, Medic. l. c. = Melilotus messanensis.

MELILOTHUS, Hornem. Hort. Hafn. Suppl. 83 (1819), sphalm. = seq.

MELILOTUS, Tourn. ex Hall. Enum. Stirp. Helv. ii. 587 (1742). *LEGUMINOSAE*, Benth. & Hook. f. i. 487.
BRACHYLOBUS, Dulac, Fl. Hautes-Pyr. 279 (1867).
MELILOTA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 382.
abyssinica, *Baker, in Oliver, Fl. Trop. Afr.* ii. 53.—Afr. trop.
abyssinica, Hochst. ex Boiss. Fl. Orient. ii. 107 = elegans.
agraria, Desf. Fl. Atlant. ii. 193 = Trifolium procumbens.
alba, *Desr. in Lam. Encyc.* iv. 63.—Europ.; Asia.
alba, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 382 = praec. ?
altissima, Thuill. Fl. Par. ed. II. 378 = officinalis.
altissima, Wall. Cat. n. 5942 = alba.
angulata, Hort. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 270 = vulgaris.
anomala, Ledeb. ex Nordm. l. c. ii. (1837) 313 = Trifolium filiforme.
arborea, Cast. ex DC. Prod. ii. 187 = alba.
arguta, Reichb. Fl. Germ. Excurs. 499 = alba.
armena, Boiss. Diagn. Ser. I. ix. 11 = officinalis.
arvensis, Wallr. Sched. Crit. 391 = officinalis.
Baumetti, Ser. in DC. Prod. ii. 189, sphalm. = *Boumetti*.
Berardi, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 382.—Europ.
Besseriana, Ser. in DC. Prod. ii. 188 = gracilis.
bicolor, *Boiss. & Bal. Diagn.* Ser. II. vi. 46.—Phrygia.
Bonplandi, *Tenore, Ind. Sem. Hort. Neap.* (1833) 14.—Hab.?
Boumetti, Hornem. Hort. Hafn. Suppl. 84 = indica.
brachyloba, Fisch. ex Ser. in DC. Prod. iii. 172 = Trigonella brachycarpa.
brachystachya, *Willd. ex Spreng. Syst.* iii. 207.—Am. austr.
brachystachys, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 219.—Turkestan.
Bungeana, *Boiss. Fl. Orient.* ii. 108.—Persia.
caerulea, Desr. in Lam. Encyc. iv. 62 = Trigonella caerulea.
caspia, *Gruner, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 420, in textu.—Reg. Casp.
citrina, Duval, ex Steud. Nom. ed. I. 520 = officinalis.
coerulea, Auct. vide *caerulea*.
collina, Guss. Suppl. Fl. Sic. Prod. 230 = elegans.
compacta, Salzm. ex Guss. Fl. Sic. Prod. ii. 485 = sulcata.
connata, Bernh. ex Reichb. Fl. Germ. Excurs. 500 = Trigonella caerulea.
cretica, Chaub. Nouv. Fl. Pélop. 50 = Trigonella graeca.
cretica, Desr. in Lam. Encyc. iv. 64 = Trigonella cretica.
dentata, *Pers. Syn.* ii. 348.—Europ.; As. bor.
diffusa, Koch, ex DC. Fl. Fr. Suppl. 564 = officinalis.
diffusa, *Trevir. Del. Sem. Hort. Bonn.* (1832).—*Cf. Flora*, xvi. (1833) 123.—Chili.
Dorycnium, Schrank, Baier. Fl. ii. 280 = Lotus Dorycnium.
elegans, *Salzm. ex Ser. in DC. Prod.* ii. 198.—Reg. Mediterr.
Emodi, R. Grah. in Wall. Cat. n. 5941 = Trigonella Emodi.
exaltata, *Bianca, in Atti Acc. Gioen. Sc. Nat.* Ser. II. xiii. (1857) 211.—Sicilia.
expansa, Hort. ex Reichb. Fl. Germ. Excurs. 498 = officinalis.
flava, Pall. Reise, iii. 654 = officinalis.
gigantea, Hort. ex Trautv. in Bull. Acad. Sc. Pétersb. viii. (1841) 270 = vulgaris.
gigantea, *Wender. in Flora*, ix. (1826) 357.—Hab.?
glaucescens, Godr. ex Trautv. in Bull. Acad. Sc. Pétersb. viii. (1841) 271 = gracilis.
globulosa, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 138, in textu = gracilis.
glomerata, Fisch. ex Ser. in DC. Prod. ii. 172 = Trigonella brachycarpa.
gracilis, *DC. Fl. Fr. Suppl.* 565.—Reg. Mediterr.; As. Min.
graveolens, Bunge, Enum. Pl. Chin. Bor. 16 = suaveolens.
hamosa, Link, Enum. Hort. Berol. ii. 260 = Trigonella spicata.
heterophylla, *Schlecht. in Linnaea*, xxi. (1848) 571.—Dalmat.
imbricata, Bess. ex Ser. in DC. Prod. ii. 188 = gracilis.
indica, *All. Fl. Pedem.* i. 308.—Europ.; As. bor.
indica, Desr. in Lam. Encyc. iv. 65 = mauritanica.
infesta, *Guss. Fl. Sic. Prod.* ii. 486.—Reg. Mediterr.
integerrima, Stev. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = ruthenica.
intermedia, Boiss. Voy. Bot. Espagne, ii. 167 = sulcata.
intricata, Nym. Consp. 152 = gracilis.
italica, *Lam. Fl. Fr.* ii. 594.—Reg. Mediterr.
italica, A. Rich. Tent. Fl. Abyss. i. 166 = abyssinica.
Kochiana, DC. Fl. Fr. Suppl. 564 = officinalis.
Kochiana, Willd. Enum. Hort. Berol. 790 = dentata.
laevis, Moench, Meth. 110 = sulcata.
laxa, Stev. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = ruthenica.
laxiflora, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 39; et ex Heuff. in Verh. Zool. Bot. Ges. Wien, viii. (1858) 88 = Trigonella caerulea.
leiosperma, *Pomel, Nouv. Mat. Fl. Atl.* 179.—Alger.
leucantha, Koch, ex DC. Fl. Fr. Suppl. 564 = alba.
linearifolia, Nocca, ex Steud. Nom. ed. I. 520 = dentata.
linearis, Cav. ex Pers. Syn. ii. 348 = dentata ?
Lippoldiana, *Lowe, in Hook. Kew Journ.* viii. (1856) 292.—Ins. Madera.
longepedicellata, Rosbach, in Verh. Naturh. Ver. Rheinl. (1875) Sitzb. 292 = officinalis.
longifolia, Tenore, Prod. Fl. Nap. 43 = sulcata.
lupulina, Lam. Fl. Fr. ii. 593 = Trifolium agrarium, etc.
lupulina, Trautv. in Bull. Sci. Acad. Pétersb. viii. (1841) 271 = Medicago lupulina.
lutea, Gueldenst. Reisen, ii. 255 = officinalis.
lutescens, Gilib. Fl. Lituan. ii. 84 = officinalis ?
macrorhiza, Bess. Enum. Pl. Volhyn. 30 = pallida.
macrorrhiza, Pers. Syn. ii. 348 = officinalis.
macrosperma, C. Koch, ex Boiss. Fl. Orient. ii. 109 = officinalis.
macrostachys, *Pomel, Nouv. Mat. Fl. Atl.* 180.—Alger.
mauritanica, *Willd. Enum. Hort. Berol.* 789.—Gall.; Afr. bor.
maxima, Le Grand, Stat. Bot. du Forez, 101 = officinalis.
medicaginoides, Zumag. Fl. Pedem. ii. 214 = Medicago lupulina.
melanosperma, Bess. ex Ser. in DC. Prod. ii. 186 = alba.
messanensis, *All. Fl. Pedem.* i. 309.—Reg. Mediterr.
micrantha, Willd. ex Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 133 = ruthenica.
microcarpa, Balb. ex Bernh. Hort. Erf. (1840).—Cf. Linnaea, xv. (1841), Lit. 90 = Trifolium micranthum.
microcarpa, C. A. Mey. ex Boiss. Fl. Orient. ii. 108 = gracilis.
minima, Roth, Nov. Pl. Sp. 361 = indica.
montana, Gaud. Fl. Helv. iv. 607 = officinalis.
neapolitana, Tenore, ex Guss. Fl. Sic. Prod. ii. 482 = gracilis.
nebrodensis, Jord. ex Nym. Consp. 172 = officinalis.
neglecta, *Tenore, Fl. Nap.* v. 135.—Ital.
occidentalis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 321 = indica.
officinalis, *Lam. Fl. Fr.* ii. 594; *Desr. in Lam. Encyc.* iv. 63.—Europ.; As. bor.
olympica, Hort. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 270 = dentata.
ornithopodioides, Desr. in Lam. Encyc. iv. 67 = Trigonella ornithopodioides.
pallida, *Bess. Ind. Sem. Hort. Cremen.* (1823); *et ex Ser. in DC. Prod.* ii. 188.—Rossia.
paluster, Menyh. in Oestr. Bot. Zeitschr. xxvii. (1877), 231, 262 = seq.
palustris, Schult. Oesterr. Fl. ed. II. ii. 346 = officinalis.
parviflora, Desf. Fl. Atlant. ii. 192 = indica.
permixta, Jord. Pugill. Pl. Nov. 55 = indica.
Pertusiana, *Gennari, in Mem. Acc. Torin.* II. xvii. (1858) 460.—Ital.
Petitpierreana, Willd. Enum. Hort. Berol. 790 = officinalis.
physocarpa, *Pomel, Nouv. Mat. Fl. Atl.* 180.—Alger.
plicata, Stev. ex Ser. in DC. Prod. ii. 188 = gracilis.
polonica, Pall. Reise, iii. 537 = ruthenica.
polonica, *Pers. Syn.* ii. 348.—Europ.
procumbens, Bess. Enum. Pl. Volh. 30 = Trigonella caerulea.
procumbens, Hort. Prag. ex Ser. in DC. Prod. ii. 186 = dentata.
psoraloides, Nutt. Gen. Am. ii. 104 = Psoralea melilotoides.
rariflora, Eichw. Pl. Casp. Cauc. 7 = ruthenica.
reticulata, *Pomel, Nouv. Mat. Fl. Atl.* 322.—Alger.
rigida, *Pomel, l. c.* 321.—Alger.
Robini, *Rafin. Fl. Ludov.* 102 (Quid ?).—Am. bor.
rotundifolia, Tenore, Prod. Fl. Nap. 43 = italica.
rugosa, Gilib. Fl. Lituan. ii. 83 = officinalis.
rugosa, Moench, Meth. 111 = italica.
rugulosa, Hort. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 270 = vulgaris.
rugulosa, Willd. Enum. Hort. Berol. 789 = indica.
ruthenica, *Ser. in DC. Prod.* ii. 186.—Rossia.
segetalis, Ser. l. c. 187 = sulcata ?
sibirica, Poir. Encyc. Suppl. iii. 647 = Medicago sibirica.
sicula, *Vitm. Summa Pl.* iv. 326.—Sicilia.
speciosa, *Dur. in Duch. Rev. Bot.* i. (1845) 365.—Alger.
striata, Moench, Meth. 111 = messanensis.
stricta, Hort. ex Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 270 = vulgaris.
suaveolens, *Ledeb. Ind. Sem. Hort. Dorp. Suppl.* (1824) 5.—Dahuria.
sulcata, *Desf. Fl. Atlant.* ii. 193.—Reg. Mediterr.
taurica, Ser. in DC. Prod. ii. 188 = gracilis.
tenella, Wallr. Beitr. Fl. Hercyn. 247 = alba.
Tommasinii, Jord. Pugill. Pl. Nov. 55 = indica.
uncinata, Poir. Encyc. Suppl. iii. 648 = Trigonella spicata.
virescens, Jord. Cat. Gr. Grenoble (1853) 11; et in F. Schultz, Fl. Fr. et Allem. 316 = officinalis ?
vulgaris, Willd. Enum. Hort. Berol. 790 = alba.
wolgica, Poir. Encyc. Suppl. iii. 648 = ruthenica.

MELINIA, Decne. in DC. Prod. viii. 588 (1844).
ASCLEPIADEAE, Benth. & Hook. f. ii. 757.
BRACHYLEPIS, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 290.
angustifolia, *A. Gray, in Proc. Am. Acad.* xii. (1877) 70.—Arizona.
bicornuta, *Griseb. in Goett. Abh.* xxiv. (1879) 228.—Reg. Argent.
Candolleana, *Decne. in DC. Prod.* viii. 589.—Reg. Argent.
Commersoniana, *Decne. l. c.*—Bras.
Glaziovi, *Fourn. in Mart. Fl. Bras.* vi. IV. 289.—Bras.
Hilariana, *Fourn.* l. c. 290.—Bras.

MELINIS, Beauv. Agrost. 54. t. 11. f. 4 (1812).
GRAMINEAE, Benth. & Hook. f. iii. 1119.
SUARDIA, Schrank, Pl. Rar. Hort. Monac. t. 58 (1819).
TRISTEGIS, Nees, Hor. Phys. Berol. 47, 54 (1820).
minutiflora, *Beauv. Agrost.* 54. t. 11. f. 4.—Bras.

MELINONIA, Brongn. in E. Morr. Cat. Brom. Jard. Liège (1873), 11 = **Pitcairnia**, L'Hérit. (Bromel.).
incarnata, Brongn. l. c. = Guiana.
rubiginosa, Brogn. l. c.—Guiana.

MELINOSPERMUM, Walp. in Linnaea, xiii. (1839) 527 = **Dichilus**, DC. (Legumin.).
pumilum, Walp. l. c. = D. gracilis.

MELINUM, Link, Handb. i. 96 (1829) = **Zizania**, Linn. (Gramin.).
palustre, Link, l. c. = Z. aquatica.

MELINUM, Medic. in Staatsw. Vorles. Churpf. Phys. Oek. Ges. i. (1791) 200 = **Salvia**, Tourn. (Labiat.).

MELIOBLASTIS, C. Muell. in Bot. Zeit. iv. (1846) 158, sphalm. = MYRIOBLASTUS, Wall. = **Cryptocoryne**, Fisch. (Aroid.).

MELIOCARPUS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 84 = **Prangos**, Lindl. (Umbell.).
alatus, Boiss. l. c. 85 (= *Colladonia alata*).—Syria.
anatolicus, Boiss. l. c. 84 (= *Colladonia anatolica*).—As. Min.
anisopterus, Boiss. l. c. 85 (= *Colladonia anisoptera*).—Palestin.
cilicicus, Boiss. et Bal. Diagn. Ser. II. v. 102 (= *Colladonia cilicica*).—Cilicia.
microcarpus, Boiss. Diagn. Ser. I. x. 53.—Syria.
pedunculatus, Boiss. l. c. 52 (= *Colladonia crenata*).—Syria.

MELIOPSIS, Reichb. Nom. 135 (1841) = **Fraxinus**, Tourn. (Oleac.).

MELIOSMA, Blume, Cat. Gew. Buitenz. 10 (1823).
SABIACEAE, Benth. & Hook. f. i. 414.
KINGSBOROUGHIA, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 67.
LORENZANEA, Liebm. l. c. 69 (1850).
MILLINGTONIA, Roxb. Fl. Ind. ed. Carey, i. 102 (1820).
OLIGOSTEMON, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 447.
WELLINGTONIA, Meissn. Gen. Comm. 207, in nota (1840).
acuminata, Hook. f. Fl. Brit. Ind. ii. 4, 774 = pungens.
alba, *Walp. Rep.* ii. 816.—Mexic.
angulata, Blume, Rumphia, iii. 197 = simplicifolia.
Arnottiana, *Walp. Rep.* i. 423.—Ind. or.
confusa, *Blume, Rumphia*, iii. 200.—Sumatra.
congesta, Walp. Rep. iv. 552 = pungens.
cuspidata, *Blume, Rumphia*, iii. 202, in nota.—Borneo.
dilleniaefolia, *Walp. Rep.* i. 423.—Reg. Himal.
elliptica, *Hook. f. Fl. Brit. Ind.* ii. 5.—Malaya.
ferruginea, *Blume, Cat. Gew. Buitenz.* 32.—Java; Borneo.
ferruginea, Sieb. et Zucc. ex Hook. f. Fl. Brit. Ind. ii. 4 = pungens.

MELIOSMA:—
floribunda, *Blume, Rumphia*, iii. 200.—Java.
fruticosa, *Blume, l. c.* 198.—Java.
glauca, *Blume, l. c.* 200 = Java; Sumatra.
hirsuta, *Blume, l. c.*—Borneo.
lanceolata, *Blume, Cat. Gew. Buitenz.* 32.—Malaya.
lancifolia, *Hook. f. Fl. Brit. Ind.* ii. 5.—Malacca.
laurina, *Blume, Rumphia*, iii. 198.—Borneo.
lepidota, *Blume, l. c.* 199.—Sumatra.
myriantha, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 153.—China; Japon.
nitida, *Blume, Cat. Gew. Buitenz.* 32.—Malacca; Sumatra.
Oldhami, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 94; *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 63.—Japon.
oppositifolia, *Griseb. Cat. Pl. Cub.* 46.—Cuba.
petiolaris, *Miq. Fl. Ind. Bat. Suppl.* 519.—Sumatra.
pinnata, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 64.—Reg. Himal.; Japon.
polyptera, *Miq. Fl. Ind. Bat. Suppl.* 520.—Sumatra.
pungens, *Walp. Rep.* i. 423.—Reg. Himal.; Japon.
rhoifolia, Maxim. in Bull. Acad. Pétersb. xii. (1868) 63 = pinnata?
rigida, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 153 = pungens.
sambucina, *Miq. Illustr. Fl. Arch. Ind.* 74.—Java.
simplicifolia, *Walp. Rep.* i. 423.—As. trop.
squamulata, *Hance, in Journ. Bot.* xiv. (1876) 364.—Hongkong.
sumatrana, Walp. Ann. i. 135; Miq. Fl. Ind. Bat. i. II. 617 = nitida.
tenuiflora, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 93.—Japon.
tenuis, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 63.—Japon.
vernicosa, *Planch. ex Griseb. Cat. Pl. Cub.* 47, in obs.—Costa Rica.
Wallichii, *Planch. ex Hook. f. Fl. Brit. Ind.* ii. 6.—Reg. Himal.; Korea.
Wightii, *Planch. ex Brand. For. Fl.* 116.—Ind. or.

MELIPHLEA, Zucc. in Abh. Akad. Muench. ii. (1847) 359 = **Sphaeralcea**, A. St. Hil. (Malvac.).
vitifolia, Zucc. l. c. 360 = S. vitifolia.

MELISITUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 383 = **Trigonella**, Linn. (Legumin.).
creticus, Medic. l. c. = T. cretica.

MELISSA, Tourn. ex Linn. Gen. ed. I. 172 (1737).
LABIATAE, Benth. & Hook. f. ii. 1191.
MELICA, Franch. & Sav. Enum. Pl. Jap. i. 368, in syn. (1875), sphalm.
MUTELIA, Gren. ex Mutel, Fl. Fr. iii. 21 (1836).
abyssinica, Hochst. in Flora, xxiv. (1841) I. Intell. 23 = Micromeria abyssinica.
acinoides, Nym. Syll. 101 = Calamintha suaveolens.
Acinos, Benth. Lab. Gen. et Sp. 389 = Calamintha Acinos.
aetheos, Hort. ex Benth. l. c. 387 = Calamintha Nepeta.
alba, Waldst. & Kit. Pl. Rar. Hung. iii. 227 = Micromeria rupestris.
alpina, Benth. Lab. Gen. et Sp. 390 = Calamintha alpina.
alpina, Boiss. Voy. Bot. Espagne, ii. 497 = Calamintha alpina.
altissima, Sibth. & Sm. Fl. Graec. Prod. i. 423 = officinalis.
arborescens, *Desf. Tabl.* ed. I. 62, nomen.—Hab.?
arundana, Boiss. Voy. Bot. Espagne, ii. 498. t. 145 = Calamintha arundana.
asiatica, *G. Don, Gen. Syst.* iv. 784.—Cochinch.
Calamintha, Linn. Sp. Pl. 593 = Calamintha officinalis.
candidissima, Munby, Fl. Algér. 61 = Calamintha candidissima.
caroliniana, Benth. Lab. Gen. et Sp. 388 = Calamintha caroliniana.
Clinopodium, Benth. l. c. 392 = Calamintha Clinopodium.
coccinea, Spreng. Syst. ii. 224 = Calamintha coccinea.
cordifolia, Pers. Syn. ii. 132 = officinalis.
corsica, Hort. ex Benth. Lab. Gen. et Sp. 732 = officinalis.
cretica, Lam. Encyc. iv. 79 = Micromeria marifolia.
cretica, Linn. Sp. Pl. 593 = Calamintha cretica.

MELISSA:—
cretica, [Soland. in] Russ. Aleppo, ed. II. ii. 256 = Calamintha incana.
debilis, Benth. Lab. Gen. et Sp. 391 = Calamintha debilis.
flava, *Benth. in Wall. Pl. As. Rar.* i. 65.—Reg. Himal.
foliosa, Opiz, ex Reichb. Fl. Germ. Excurs. 329 = officinalis.
fruticosa, Linn. Sp. Pl. 593 = Micromeria marifolia.
fruticosa, Willd. ex Steud. Nom. ed. II. ii. 121 = Micromeria rupestris.
glandulosa, Benth. Lab. Gen. et Sp. 387 = Calamintha glandulosa.
glomerata, Stokes, Bot. Mat. Med. iii. 366 = Calamintha Clinopodium.
granatensis, Nym. Syll. 101 = Calamintha alpina.
grandiflora, Linn. Sp. Pl. 592 = Calamintha grandiflora.
graveolens, Benth. Lab. Gen. et Sp. 390 = Calamintha graveolens.
graveolens, Host, Fl. Austr. ii. 128 = officinalis.
hirsuta, *Blume, Bijdr.* 830.—Java.
hirsuta, Hornem. Hort. Hafn. ii. 562 = officinalis.
hortensis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 (Quid?).—Java.
humilis, G. Don, Gen. Syst. iv. 784 (= *Clinopodium humile*).—Am. bor.
incana, Benth. Lab. Gen. et Sp. 386 = Calamintha incana.
inodora, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 127.—Hab.?
intermedia, Baumg. Enum. Stirp. Transs. ii. 184 = Calamintha officinalis.
longicaulis, Wall. ex Benth. Lab. Gen. et Sp. 395 = Calamintha longicaulis.
longifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = hirsuta.
macrostema, Moç. & Sesse, ex Benth. Lab. Gen. et Sp. 395 = Calamintha macrostema.
majoranaefolia, Mill. Gard. Dict. ed. VIII. n. 7 = Calamintha patavina.
marifolia, Griseb. Spicil. Fl. Rumel. ii. 125 = Micromeria serpyllifolia.
marifolia, Pers. Syn. ii. 132 = Micromeria marifolia.
marifolia, Willd. ex Steud. Nom. ed. II. ii. 122 = Gardoquia discolor.
maritima, Nym. Syll. 101 = Calamintha maritima.
maxima, Ard. Animad. ii. 28 = Perilla ocimoides.
microphylla, Benth. Lab. Gen. et Sp. 390 = Calamintha corsica.
montana, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = Coleus galeatus.
nepalensis, Benth. in Wall. Pl. As. Rar. i. 66 = Mosla dianthera.
Nepeta, Linn. Sp. Pl. 593 = Calamintha Nepeta.
nova, Portenschl. ex Steud. Nom. ed. II. ii. 122 = Micromeria Pulegium.
obtusifolia, Pers. Syn. ii. 132 = Calamintha Nepeta.
occidentalis, Rafin. ex Benth. Lab. Gen. et Sp. 732 = officinalis.
officinalis, *Linn. Sp. Pl.* 592.—Reg. Mediterr.; Oriens.
origanifolia, Benth. Lab. Gen. et Sp. 392 = Calamintha origanifolia.
parviflora, *Benth. in Wall. Pl. As. Rar.* i. 65.—Reg. Himal.; Malaya.
parviflora, Salisb. Prod. 86 = Calamintha Nepeta.
patavina, Benth. Lab. Gen. et Sp. 389 = Calamintha patavina.
perennis, Forsk. Fl. Aegypt. Arab. 108 = Salvia aegyptiaca.
polyanthos, *Lodd. ex G. Don, in Loud. Hort. Brit.* 239, nomen.—Hab.?
pulegioides, Linn. Sp. Pl. 593 = Hedeoma pulegioides.
Pulegium, Griseb. Spicil. Fl. Rumel. ii. 125 = Mentha Pulegium.
Pulegium, Rochel, Pl. Banat. Rar. 62 = Micromeria Pulegium.
purpurascens, Benth. Lab. Gen. et Sp. 391 = Calamintha graveolens.
pyrenaica, Scop. Ann. II. Hist. Nat. (1769) 55 = Horminum pyrenaicum.
repens, Benth. Lab. Gen. et Sp. 392 = Calamintha umbrosa.
romana, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
rotundifolia, Benth. Lab. Gen. et Sp. 391 = Calamintha patavina.
rotundifolia, Soland. ex Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimp. 14 = Calamintha officinalis.

MELISSA:—
rugosa, *Lour. Fl. Cochinch.* 368.—Cochinchin.
rupestris, Salisb. Prod. 86 = Micromeria marifolia.
serpyllifolia, Nym. Suppl. Syll. 20 = Micromeria serpyllifolia.
serpyllifolia, Willd. ex Benth. in Linnaea, xi. (1837) 341 = Micromeria marifolia.
stricta, *Kitt. Taschenb.* ed. II. 376.—Europ.
suaveolens, Nym. Suppl. Syll. 20 = Calamintha suaveolens.
subnuda, Waldst. & Kit. Pl. Rar. Hung. iii. 291 = Micromeria Pulegium.
sylvestris, Lam. Fl. Fr. ii. 401 = Melittis Melissophyllum.
taurica, Hort. ex Benth. Lab. Gen. et Sp. 393 = officinalis.
teucriifolia, Willd. ex Ledeb. Fl. Ross. iii. 379 = Nepeta teucriifolia.
teucrioides, Boiss. Diagn. Ser. I. vii. 49 = Nepeta lasiocephala.
thymifolia, Benth. Lab. Gen. et Sp. 386 = Calamintha thymifolia.
umbrosa, Auct. Germ. ex Steud. Nom. ed. II. ii. 122 = M. Calamintha.
umbrosa, Bieb. Fl. Taur. Cauc. ii. 63 = Calamintha umbrosa.
veronicaefolia, *Pers. Syn.* ii. 132.—Ins. Bahama.
vulgaris, Trevis. Prosp. Fl. Eugan. 26 = Calamintha Clinopodium.

MELISSITUS, Medic. Phil. Bot. i. 209 (1789) = **Trigonella**, Linn. (Legumin.).
dentata, Moench, Meth. 111 = T. cretica.

MELISSOPHYLLON, Adans. Fam. ii. 189 (1763) = **Melittis**, Linn. (Labiat.).

MELISSOPSIS, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 193 (1876) = **Ageratum**, Linn. (Compos.).
oppositifolia, Sch. Bip. ex Baker, l. c. 195 = A. melissaefolium.

MELISTAURUM, Forst. Char. Gen. 143. t. 72 (1776) = **Casearia**, Jacq. (Samyd.).
arboreum, Van Hasselt, ex Blume, Mus. Bot. Lugd. Bat. i. 254 = C. flavovirens.
distichum, Forst. Char. Gen. 144 = C. Melistaurum.

MELITIS, Gled. in Mém. Berl. Acad. v. (1749) 129 = seq.

MELITTIS, Linn. Gen. ed. I. 172 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1205.
MELISSOPHYLLON, Adans. Fam. ii. 189 (1763).
albida, Guss. Fl. Sic. Prod. ii. 140 = M. Melissophyllum.
caroliniana, Spreng. Syst. ii. 700 = Macbridea pulchra.
grandiflora, Sm. Engl. Bot. t. 636 = M. Melissophyllum.
japonica, Thunb. in Trans. Linn. Soc. ii. (1794) 338 = Rehmannia chinensis.
melissaefolium, Gilib. Fl. Lituan. i. 76; Salisb. Prod. 87 = seq.
Melissophyllum, *Linn. Sp. Pl.* 597.—Europ.
Melissophyllum, Thunb. Fl. Jap. 248 = Rehmannia chinensis.

MELLA, Vand. Fl. Lusit. et Bras. Spec. 43. f. 53 (1788) = **Herpestis**, Gaertn. f. (Scrophul.).

MELLERA, S. Moore, in Journ. Bot. xvii. (1879) 225. *ACANTHACEAE*.
lobulata, *S. Moore, l. c.*—Afr. trop.

MELLIGO, Rafin. Fl. Tellur. iii. 92 (1836) = **Salvia**, Tourn. (Labiat.).
amena, Rafin. l. c. = S. lamiifolia.

MELLISSIA, Hook. f. in Hook. Ic. Pl. t. 1021 (1867). *SOLANACEAE*, Benth. & Hook. f. ii. 891.
begonifolia, *Hook. f. Ic. Pl.* t. 1021.—Ins. S. Helena.

MELLITIS, Scop. Fl. Carn. ed. II. i. 421 (1772) = **Melittis**, Linn. (Labiat.).

MELLOA, Bur. in Adansonia, viii. (1868) 379. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1035.
populifolia, *Bur. ex Benth. & Hook. f. Gen.* ii. 1035.—Bras.

MELLOBIUM, A. Juss. in Orb. Dict. vii. 269 (1849) = **Melolobium**, Eckl. & Zeyh. (Legumin.).

MELLOCA, Lindl. in Gard. Chron. (1847) 685 = **Ullucus**, Lozano (Chenop.).
peruviana, Moq. in DC. Prod. xiii. II. 225 = U. tuberosa.
tuberosa, Lindl. in Gard. Chron. (1847) 685 = U. tuberosa.

MELLOLOBIUM, Reichb. Nom. 151 (1841) = **Melolobium**, Eckl. & Zeyh. (Legumin.).

MELO [Tourn.] Linn. Syst. ed. I (1735) = **Cucumis**, Tourn. (Cucurb.).
Chate, Sageret, ex M. Roem. Syn. Pepon. 74 = C. Melo.
flexuosus, Sageret, l. c. 75 = C. Melo.
persicus, Sageret, l. c. 71 = C. Melo.
sativus, Sageret, l. c. 68 = C. Melo.
vulgaris, Cogn. in DC. Monog. Phan. iii. 484, sphalm. = C. Melo.

MELOCACTUS, [Tourn.] Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) 417. tt. 11, 12. *CACTEAE*, Benth. & Hook. f. i. 847.
acuatus, Link & Otto, l. c. t. 23 = Echinocactus acuatus.
ambiguus, Hort. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. i. (1833) 364 = Cereus leucanthus.
amoenus, *Hoffmgg. Preiss. Verz.* ed. VII. (1833) 22; *et ex Pfeiff. Enum. Diagn. Cact.* 43.—N. Granat.
atrosanguineus, *Hort. Berol. ex Pfeiff. l. c.* 44.—Ins. S. Thomas.
atrovirens, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 279, nomen.—Hab. ?
Besleri, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) 420. t. 21 = placentiformis.
Besleri-affinis, Hort. ex Pfeiff. Enum. Diagn. Cact. 60 = Echinocactus spiralis.
bradypus, Lehm. ex Steud. Nom. ed. II. ii. 122 = Cereus senilis.
Brongniartii, *Lem. Cact. Aliq. Nov. Desc.* 12.—Hab. ?
caesius, *H. L. Wendl. ex Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 184.—N. Granat.
cephalonoplus, *Lem. ex Foerst. Handb. Cact.* ed. I. 279, nomen.—Hab. ?
cephalophora, Salm-Dyck, Cact. Hort. Dyck. ed. II. 137 = mammillariaeformis.
Columna-Trajani, Pfeiff. Enum. Diagn. Cact. 46 = Cereus Columna-Trajani.
communis, *Link & Otto, in Verh. Preuss. Ver. Gartenb.* iii. (1827) 417. t. 11.—Ind. occ.
coronatus, *Cels, ex Foerst. Handb. Cact.* ed. I. 279, nomen.—Hab. ?
crassicostatus, Lem. Cact. Aliq. Nov. Desc. 13 = obtusipetalus.
crassispinus, Salm-Dyck, in Otto & Dietr. Allg. Gartenz. viii. (1840) 10 = Lemairei.
curvispinus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 46.—Mexic.
Delessertianus, *Lem. ex Labour. Monog. Cact.* 19.—Mexic.
depressus, *Hook. Bot. Mag.* t. 3691.—Bras.
depressus, Salm-Dyck, ex DC. Prod. iii. 463 = Cereus leucanthus.
dichroacanthus, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 147.—Ins. S. Thomas.
elegans, Hort. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 244 = Cereus leucanthus.
Ellemetii, *Miq. in Nederl. Kruidk. Arch.* iv. (1859) 336.—Bras.
excavatus, *Hort. ex G. Don, in Sweet, Hort. Brit.* ed. III. 282, nomen.—Mexic.
ferox, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 519; *et ex Labour. Monog. Cact.* 16.—Mexic.
Gardenerianus, Booth, ex Foerst. l. c. 277 = depressus.
Gilliesii, Hort. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 364; et ex Pfeiff. Enum. Diagn. Cact. 50 (= *Echinopsis formosus*).—Chili.
gladiatus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 17 = Echinocactus gladiatus.
goniodacanthus, *Lem. Cact. Aliq. Nov. Desc.* 11.—Hab. ?
Grengelii, Hort. ex Forbes, Hort. Tour Germ. 151, nomen = communis.
griseus, *H. L. Wendl. ex Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 185.—N. Granat.
havannensis. *Miq. l. c.* 144.—Cuba.
Hookerianus, *Forbes, ex Foerst. Handb. Cact.* ed. I. 279, nomen.—Hab. ?

MELOCACTUS:—
Hystrix, *Parment. ex Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 138.—Hab. ?
ingens, Karw. ex Pfeiff. Enum. Diagn. Cact. 54 = Echinocactus ingens.
intricatus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 24 = Echinocactus intricatus.
Lamarckii, G. Don, Gen. Syst. iii. 160 = communis.
Langsdorfii, DC. Prod. iii. 461 = Echinocactus Langsdorfi.
latispinus, Hort. ex Pfeiff. Enum. Diagn. Cact. 56 = Echinocactus cornigerus.
Lehmanni, *Miq. in Linnaea*, xi. (1837) 642.—Venezuela.
Lemairei, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 192.—Ins. S. Doming.
Leopoldi, *Hort.—Cf. Gard. Chron.* (1876) I. 603.—Hab. ?
leucaster, *Hoffmgg. Preiss. Verz.* ed. VII. (1833) 22.—Hab. ?
macracanthoides, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 173.—Ins. S. Thomas.
macrocanthus, *Link & Otto, in Verh. Preuss. Ver. Gartenb.* iii. (1827) 418. t. 12; *Salm-Dyck, Hort. Dyck.* 344.—Ind. occ.
macrocanthus, Miq. in Linnaea, xi. (1837) 157 = macracanthoides.
mammillariaeformis, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* iv. (1836) 148.—Mexic.
melocactoides, *DC. Prod.* iii. 461.—Bras.
meonacanthus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 15 = Echinocactus meonacanthus.
microcephalus, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 157.—Ins. Curaçao.
Miquelii, *Lehm. Del. Sem. Hort. Hamb.* (1838); *ex Miq. l. c.* 149.—Ins. S. Crucis.
monoclonos, *DC. ex Steud. Nom.* ed. II. ii. 122.—Ind. occ.; Guian.
Monvillianus, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 133.—Hab. ?
obtusipetalus, *Lem. Cact. Aliq. Nov. Desc.* 11.—N. Granat.
octogonus, *Hort. ex Forbes, Hort. Tour Germ.* 151, nomen.—Mexic.
Oreas, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 192.—Bras.
orthacanthus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 18 = Echinocactus orthacanthus.
Parthoni, Hort. ex Miq. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 190 = violaceus.
parvispinus, DC. Prod. iii. 463, in syn. = Echinocactus parvispinus.
pentacentrus, *Lem. Cact. Nov. Gen. et Sp.* 108.—Bras.
placentiformis, *DC. Prod.* iii. 460.—Bras.
platyacanthus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 14 = Echinocactus platyacanthus.
poliacanthus, Link & Otto, l. c. t. 16 = Echinocactus polyacanthus.
prolifer, Hort. ex Otto & Dietr. Allg. Gartenz. i. (1833) 364 = Echinocactus robustus.
pycnacanthus, Cels, ex Miq. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 200 b = Lemairei.
pyramidalis, *Link & Otto, in Verh. Preuss. Ver. Gartenb.* iii. (1827) 419. t. 25.—Am. trop.
recurvus, Link & Otto, l. c. t. 20 = Echinocactus recurvus.
rubens, Hort. ex Miq. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 145 = amoenus.
rubens, *Pfeiff. Enum. Diagn. Cact.* 43.—Ind. occ.
rufispinus, *Bertol. in Nov. Comm. Bonon.* iii. (1839) 322. t. 26.—Hab. ?
Salmianus, *Link & Otto, in Verh. Preuss. Ver. Gartenb.* iii. (1827) 423. t. 13; *et ex Salm-Dyck, Hort. Dyck.* 345.—Ins. Curaçao.
Schlumbergerianus, *Lem. Illustr. Hortic.* viii. (1861) *Misc.* 32.—Ins. S. Thomas.
Sellowi, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 22; et DC. Prod. iii. 461 = Echinocactus Sellowii.
Spatangus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 45.—Ins. Curaçao.
tenuispinus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 19 = Echinocactus tenuispinus.

MELOCACTUS:—
tephracanthus, Link & Otto, in Verh. Preuss. Ver. Gartenb. iii. (1827) t. 16 = Echinocactus tephracanthus.
tuberculatus, Link & Otto, l. c. t. 26 = Echinocactus tuberculatus.
violaceus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 313.—Bras.
viridescens, Nutt. ex Teschem. in Journ. Bost. Soc. Nat. Hist. v. (1845) 293 = Echinocactus viridescens.
Wendlandii, *Miq. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 146.—Ins. S. Thomas.
xanthacanthus, *Miq. l. c.* 169.—Ins. S. Thomas.
Zuccarinii, *Miq. in Linnaea*, xi. (1837) 645.—Ins. Curaçao.

MELOCALAMUS, Benth. in Journ. Linn. Soc. xix. (1881) 134. *GRAMINEAE*, Benth. & Hook. f. iii. 1212.
compactiflorus, *Benth. in Journ. Linn. Soc.* xix. (1881) 134.—Burma.

MELOCANNA, Trin. in Spreng. Neue Entdeck. ii. 43 (1821). *GRAMINEAE*, Benth. & Hook. f. iii. 1214.
IRULIA, Bedd. Forest. Man. Bot. 235 (1873).
bambusoides, *Trin. in Spreng. Neue Entdeck.* ii. 43.—Ind. or.
excelsa, Roep. ex Trin. Clav. Agrost. 105, 397 = Bambusa maxima.
gracilis, *Kurz, ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 133.—Malaya.
humilis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 251.—Burma.
humilis, *Roep. ex Trin. Clav. Agrost.* 105.—Malaya.
Kurzii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 134.—Ins. Andaman.
Rheedii, Steud. Syn. Pl. Gram. 332 = bambusoides.
travancorica, *F. Muell. Sel. Pl. Indust. Cult.* 129.—Ind. or.
virgata, Munro, in Trans. Linn. Soc. xxvi. (1868) 133 = Cephalostachyum virgatum.
Zollingeri, *Kurz, ex Munro, l. c.* 134.—Java.

MELOCHIA, Dill. ex Linn. Syst. ed. I (1735). *STERCULIACEAE*, Benth. & Hook. f. i. 223.
ALEURODENDRON, Reinw. in Syll. Ratisb. ii. (1828) 12.
ALTHERIA, Thou. Gen. Nov. Madag. 19 (1806).
ANAMORPHA, Karst. & Triana, in Linnaea, xviii. (1844) 443.
ANTIPHYLA, Rafin. New Fl. Am. iv. 16 (1836).
GLOSSOSPERMUM, Wall. Cat. n. 1153 (1831).
LOCHEMIA, Arn. in Ann. Sc. Nat. Sér. II. xi. (1839) 172.
MELUCHIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 217.
MOLUCHIA, Medic. Malv. 10 (1787).
MOUGEOTIA, H. B. & K. Nov. Gen. et Sp. v. 326. tt. 483, 484 (1821).
PHYSOCODON, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 212.
POLYCHLAENA, G. Don, Gen. Syst. i. 488 (1831).
RIDDELIA, Rafin. New Fl. Am. iv. 15 (1836).
RIEDLEA, Vent. in Mém. Inst. Sc. Phys. i. (1807) 2.
VISENIA, Houtt. Handleid. viii. 308 (1777).
acutangula, *Span. in Linnaea*, xv. (1841) 174.—Ins. Timor.
affinis, Wall. Cat. n. 1198 = corchorifolia.
anomala, *Griseb. in Goett. Abh.* xix. (1874) 93.—Reg. Argent.
arborea, *Blanco, Fl. Filip.* ed. I. 524.—Ins. Philipp.
arenosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 127.—Bras.
aristata, *A. Gray, Bot. U. St. Expl. Exped.* i. 198.—Ins. Samoa.
Berterianum, Balb. ex DC. Prod. i. 492 = Waltheria glabra.
betonicaefolia, *A. St. Hil. Fl. Bras. Mer.* i. 165.—Bras.
borbonica, *Cav. Diss.* vi. 321.—Ins. Borbon.
Burmanni, Zoll. & Mor. Syst. Verz. Zoll. 27 = Corchorus javanicus.
capitata, Boj. Hort. Maurit. 38 = corchorifolia.
caracassana, *Jacq. Coll.* ii. 369.—Venezuela.
carpinifolia, *Wendl. Bot. Beobacht.* 52.—Ind. occ.
Chamaedrys, *A. St. Hil. Fl. Bras. Mer.* i. 161.—Bras.
cinerea, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 35.—Bras.
Clinopodium, A. St. Hil. & Naud. l. c. 36 = hirsuta, *Cav.*
concatenata, Linn. Sp. Pl. 675 = corchorifolia.

MELOCHIA:—
concinna, *Miq. in Linnaea*, xxii. (1849) 467.—Guian.
corchorifolia, *Linn. Sp. Pl.* 675.—Cosmop. trop.
corchorifolia, Wall. Cat. n. 1196 H = Waltheria indica.
cordata, *Burm. f. Fl. Ind.* 143.—Ind. or.
cordiformis, *A. St. Hil. Fl. Bras. Mer.* i. 166.—Bras.
corymbosa, Meissn. ex Steud. Nom. ed. II. ii. 123 = Physodium corymbosum.
crenata, Bert. ex Spreng. Syst. iii. 29 = portoricensis.
crenata, Vahl, Symb. Bot. iii. 86 = tomentosa.
densiflora, *Miq. in Linnaea*, xxii. (1849) 800.—Bras.
depressa, *Linn. Sp. Pl.* 674.—Cuba.
depressa, Mill. Gard. Dict. ed. VIII. n. 3 = peruviana.
diffusa, Bert. ex Spreng. Syst. iii. 30 = tomentosa.
domingensis, Jacq. Select. Am. 194 = pyramidata.
erecta, *Burm. f. Fl. Ind.* 143.—Java.
fasciculata, Benth. in Hook. Journ. Bot. iv. (1842) 127 = parvifolia.
fasciculata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = corchorifolia.
frutescens, Jacq. Obs. Bot. ii. 22 = tomentosa.
globifera, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 343.—N. Granat.
gracilis, A. St. Hil. & Naud. l. c. Sér. II. xviii. (1842) 35 = venosa.
graminifolia, *A. St. Hil. Fl. Bras. Mer.* i. 160.—Bras.
hermannioides, *A. St. Hil. l. c.* 163.—Bras.
hirsuta, *Cav. Diss.* vi. 323.—Mexic.
hirsuta, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 570.—Peruv.
hispida, *Hook. & Arn. Bot. Beech. Voy.* 60.—Ins. Societ.
hypoleuca, *Miq. in Linnaea*, xxii. (1849) 468.—Guian.
inflata, Benth. Bot. Voy. Sulph. 71 = lupulina.
interrupta, *Hemsl. Biol. Centr. Am. Bot.* i. 131.—Mexic.
jamaicensis, Balb. ex DC. Prod. i. 492 = venosa.
Jurgensenii, *Hemsl. Biol. Centr. Am. Bot.* i. 131.—Mexic.
kerriaefolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 341.—N. Granat.
lanata, *A. St. Hil. Fl. Bras. Mer.* i. 167.—Bras.
lanceolata, *Benth. in Hook. Journ. Bot.* iv. (1842) 128.—Bras.
lilacina, *A. St. Hil. Fl. Bras. Mer.* i. 162.—Bras.
lupulina, *Sw. Prod. Veg. Ind. Occ.* 97.—Am. trop.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 324.—Mexic.
manducata, *Wright, in Sauv. Fl. Cub.* 14.—Cuba.
melissaefolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 129.—Guian.
mollis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 343.—N. Granat.
mollissima, *Desf. Tabl.* ed. I. 148, nomen; *Pers. Syn.* ii. 218.—Am. austr.
monoica, Baill. Hist. Pl. iv. 74 = Dicarpidium monoicum.
nepetoides, *A. St. Hil. Fl. Bras. Mer.* i. 165.—Bras.
nitidula, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 35 = venosa.
nodiflora, *Sw. Prod. Veg. Ind. Occ.* 97.—Am. trop.
oblonga, *Benth. in Hook. Journ. Bot.* iv. (1842) 128.—Guian.
odorata, *Linn. f. Suppl.* 302; *Forst. f. Prod.* 47.—Ins. Tanna.
ovata, Desf. Tabl. ed. II. 172 = Melhania abyssinica.
parviflora, Loud. Hort. Brit. 269 = seq.?
parvifolia, *H. B. & K. Nov. Gen. et Sp.* v. 325.—Venezuela.
pauciflora, Wall. Cat. n. 1199 = corchorifolia.
peruviana, *Desr. in Lam. Encyc.* iv. 83.—Peruv.
plicata, *Presl, Rel. Haenk.* ii. 145.—Mexic.
polystachya, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 341 = venosa.
portoricensis, *Spreng. Syst.* iii. 29.—Ins. Portorico.
pulverulenta, *Miers, in Proc. Roy. Hort. Soc.* iv. (1864) 185.—N. Granat.
pyramidata, *Linn. Sp. Pl.* 674.—Am. calid.
rhodocalyx, *C. Koch & Bouché, in Ind. Sem. Hort. Berol.* (1854) 14.—Hab.?
scordiifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 207.—N. Granat.
scutellarioides, *Hemsl. Biol. Centr. Am. Bot.* i. 131.—Mexic.
sericea, A. St. Hil. Fl. Bras. Mer. i. 160 = venosa.
serrata, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 36 = hirsuta, *Cav.*
simplex, *A. St. Hil. Fl. Bras. Mer.* i. 164.—Bras.
splendens, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 36.—Bras.

MELOCHIA:—
supina, Linn. Sp. Pl. 675 = corchorifolia.
tenella, *Hemsl. Biol. Centr. Am. Bot.* i. 132.—Mexic.
tomentella, *Hemsl. l. c.*—Mexic.
tomentosa, *Linn. Syst.* ed. X. 1140.—Am. trop.
truncata, Wall. Cat. n. 1196 K = Sida Schimperiana.
truncata, Willd. Sp. Pl. iii. 601 = corchorifolia.
Turpiniana, H. B. & K. Nov. Gen. et Sp. v. 323 = tomentosa.
ulmarioides, *A. St. Hil. Fl. Bras. Mer.* i. 159.—Bras.
ulmifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 129.—Am. trop.
ulmifolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 36.—Bras.
varronifolia, *Desf. Cat. Hort. Par.* ed. III. 252, nomen.—Hab.?
velutina, Wall. ex Bedd. For. Man. Bot. 35. t. 5. f. 3 = arborea.
venosa, *Sw. Prod. Veg. Ind. Occ.* 97.—Bras.
vestita, Benth. in Hook. Journ. Bot. iv. (1842) 130 = hirsuta, *Cav.*
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 193 = odorata.

MELOCHIA, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 296, absque char. (Quid?).
candelaris, Rottb. l. c.—Guian.

MELODINUS, Forst. Char. Gen. 37. t. 19 (1776). *APOCYNACEAE*, Benth. & Hook. f. ii. 694.
BICORONA, A. DC. in DC. Prod. viii. 330 (1844).
ECHALTIUM, Wight, Ic. t. 472 (1841).
LYCIMNIA, Hance, in Walp. Ann. iii. 20 (1852).
ONCINUS, Lour. Fl. Cochinch. 123 (1790).
acutiflorus, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 71.—Austral.
Baueri, *Endl. Prod. Fl. Norf.* 57.—Ins. Norfolk.
chilocarpoides, F. Muell. Fragm. vi. 118 = Chilocarpus australis.
Cumingii, *A. DC. in DC. Prod.* viii. 330.—Ins. Philipp.; China.
eugenifolius, Wall. Cat. n. 1616 = Leuconotis eugenifolius.
fasciculatus, *Blume, Mus. Bot. Lugd. Bat.* i. 155.—Borneo.
Forbesii, *Fawc. in Forbes, Nat. Wand. East. Archip.* 510.—Ins. Timor.
fusiformis, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 332.—Ins. Hongkong.
Guilfoylei, *F. Muell. Fragm.* vi. 118.—Austral.
khasianus, *Hook. f. Fl. Brit. Ind.* iii. 629.—Reg. Himal.
laetus, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 332 = monogynus.
laevigatus, *Blume, Mus. Bot. Lugd. Bat.* i. 155.—Malaya.
laxiflorus, *Blume, l. c.*—Java.
micranthus, *Hook. f. Fl. Brit. Ind.* iii. 629.—Malacca.
monogynus, *Roxb. Hort. Beng.* 20; *Fl. Ind.* ii. 56.—Reg. Himal.; Malaya; China.
orientalis, *Blume, Bijdr.* 1026.—Malaya.
parvifolius, *Lodd. Cat.*; *ex Sweet, Hort. Brit.* ed. II. 356, nomen.—Ind. or.
philippensis, *A. DC. in DC. Prod.* viii. 330.—Ins. Philipp.
philliraeoides, *Labill. Sert. Austro-Caled.* 29. t. 33.—N. Caled.
scandens, *Forst. Char. Gen.* 38.—N. Caled.
scandens, Hassk. Cat. Hort. Bogor. Alt. 120 = laxiflorus.
scandens, Seem. Fl. Vit. 155 = vitiensis.
suaveolens, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 333.—Ins. Hongkong.
terminalis, *Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
undulatus, *Lodd. Cat.*; *ex Sweet, Hort. Brit.* ed. II. 356, nomen.—Ind. or.
vitiensis, *Rolfe, in Journ. Bot.* xxi. (1883) 201.—Ins. Fiji.

MELODORUM, Hook. f. & Thoms. Fl. Ind. i. 115 (1855). *ANONACEAE*, Benth. & Hook. f. i. 28.
FISSISTIGMA, Griff. Notul. ii. 706 (1854).
MITRELLA, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 38 (1865).
PYRAMIDANTHE, Miq. l. c. 39 (1865).
africanum, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 477.—Afr. trop.

MELODORUM:—
bancanum, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 343; *et in Flora*, liii. (1870) 244.—Malaya.
Beccarii, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 24.—N. Guin.
bicolor, *Hook. f. & Thoms. Fl. Ind.* i. 119.—Reg. Himal.; Burma.
borneense, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 36.—Borneo.
cinerascens, *Miq. l. c.* 37.—Sumatra.
clavipes, *Hance, in Journ. Bot.* xv. (1877) 328.—China.
cylindricum, *Maing. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 80.—Malacca.
elegans, *Hook. f. & Thoms. Fl. Ind.* i. 122.—Malaya.
fulgens, *Hook. f. & Thoms. l. c.* 120.—Malaya.
glaucescens, *Hance, in Journ. Bot.* xix. (1881) 112.—China.
Griffithii, *Hook. f. & Thoms. Fl. Ind.* i. 120.—Burma.
hypoglaucum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 36.—Sumatra.
Kentii, *Hook. f. & Thoms. Fl. Ind.* i. 116.—Java.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 37.—Borneo.
lanuginosum, *Hook. f. & Thoms. Fl. Ind.* i. 117.—Malaya.
latifolium, *Hook. f. & Thoms. l. c.*—Malaya.
Lefevrii, *Baill. Adansonia*, x. (1871) 108.—Cochinch.
Leichhardtii, *Benth. Fl. Austral.* i. 52.—Austral.
Maccreai, *F. Muell. Fragm.* vi. 176.—Austral.
macranthum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 291.—Burma.
Maingayi, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 80.—Penang.
manubriatum, *Hook. f. & Thoms. Fl. Ind.* i. 118.—Malaya.
mollissimum, Miq. Fl. Ind. Bat. Suppl. 374 = latifolium.
parviflorum, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 344; *et in Flora*, liii. (1870) 244.—Malaya.
pisocarpum, *Hook. f. & Thoms. Fl. Ind.* i. 123.—Malacca.
polyanthum, *Hook. f. & Thoms. l. c.* 121.—Reg. Himal.
polycarpum, Benth. in Trans. Linn. Soc. xxiii. (1862) 477 = Xylopia polycarpa.
prismaticum, *Hook. f. & Thoms. Fl. Ind.* i. 121.—Malaya.
punctulatum, *Baill. Adansonia*, x. (1871) 107.—N. Caled.
rubiginosum, *Hook. f. & Thoms. Fl. Ind.* i. 116.—Reg. Himal.; Burma.
rufinervum, *Hook. f. & Thoms. l. c.*—Reg. Himal.
sphaerocarpum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 38.—Java; Sumatra.
Uhrii, *F. Muell. Fragm.* vi. 2.—Austral.
verrucosum, *Hook. f. & Thoms. Fl. Ind.* i. 119.—Reg. Himal.
Wallichii, *Hook. f. & Thoms. l. c.* 118.—Reg. Himal.
Zippelii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 38.—Sumatra.

MELODORUM, Lour. Fl. Cochinch. 351 (1790) = **Unona**, Linn. f. (Anonac.).
arboreum, Lour. l. c. = U. sylvatica.
fruticosum, Lour. l. c. = U. dumetorum.

MELOLOBIUM, Eckl. & Zeyh. Enum. 188 (1836).
LEGUMINOSAE, Benth. & Hook. f. i. 478.
SPHINGIUM, E. Mey. Comm. Pl. Afr. Austr. 65 (1835).
adenoides, *Eckl. & Zeyh. Enum.* 190.—Afr. austr.
alpinum, *Eckl. & Zeyh. l. c.*—Afr. austr.
calycinum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 350.—Afr. austr.
canaliculatum, *Benth. l. c.* 353.—Afr. austr.
candicans, *Eckl. & Zeyh. Enum.* 189.—Afr. austr.
canescens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 351.—Afr. austr.
cernuum, *Eckl. & Zeyh. Enum.* 190.—Afr. austr.
collinum, *Eckl. & Zeyh. l. c.* 189.—Afr. austr.
decumbens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 352.—Afr. austr.
exudans, Harv. in Harv. & Sond. Fl. Cap. ii. 80.—Afr. austr.
humile, *Eckl. & Zeyh. Enum.* 190.—Afr. austr.
microphyllum, *Eckl. & Zeyh. l. c.* 189.—Afr. austr.
obcordatum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 81.—Afr. austr.
parviflorum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 351.—Afr. austr.
spicatum, Eckl. & Zeyh. Enum. 190 = cernuum.
squarrosum, Eckl. & Zeyh. l. c. 189 = candicans.
stipulatum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 81.—Afr. austr.
viscidulum, *Steud. Nom.* ed. II. ii. 123.—Afr. austr.

MELOMPHIS, Rafin. Fl. Tellur. ii. 21 (1836) = **Ornithogalum**, Tourn. (Liliac.).
arabica, Rafin. l. c. 22 = O. arabicum.
patens, Rafin. l. c. iii. 59 = O. arabicum.
peruviana, Rafin. l. c. ii. 22 = Scilla peruviana.
sicula, Rafin. l. c. = Scilla peruviana.

MELONEURA, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentib.).
purpurea, Rafin. l. c. = U. striatula.

MELONGENA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Solanum**, Tourn.
incurva, Mill. l. c. ed. VIII. n. 3 = S. Melongena.
ovata, Mill. l. c. n. 1 = S. Melongena.
spinosa, Mill. l. c. n. 4 = S. Melongena.
teres, Mill. l. c. n. 2 = S. Melongena.

MELORIMA, Rafin. Fl. Tellur. iv. 53 (1836) = **Fritillaria**, Linn. (Liliac.).
cirrhosa, Rafin. l. c. = F. cirrhosa.

MELOSMON, Rafin. Fl. Tellur. iii. 85 (1836) = **Teucrium**, Tourn. (Labiat.).
bicolor, Rafin. l. c. = T. bicolor.
campanulatum, Rafin. l. c. = T. campanulatum.
orientale, Rafin. l. c. = T. orientale.
rotundifolium, Rafin. l. c. = T. pyrenaicum?

MELOSPERMA, Benth. in DC. Prod. x. 374 (1846).
SCROPHULARINEAE, Benth. & Hook. f. ii. 946.
andicola, *Benth. in DC. Prod.* x. 374.—Chili.
angustifolia, *Phil. in Anal. Univ. Chil.* (1873) 525.—Chili.
glabra, *Phil. in Linnaea*, xxix. (1857–58) 26.—Chili.

MELOTHRIA, Linn. Coroll. Gen. 1 (1737). *CUCURBITACEAE*, Benth. & Hook. f. i. 830.
AECHMANDRA, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274.
ARKEZOSTIS, Rafin. New Fl. Am. iv. 100 (1836).
DICLIDOSTIGMA, Kunze, in Linnaea, xvii. (1843) 576.
HARLANDIA, Hance, in Walp. Ann. ii. 648 (1852).
JUCHIA, M. Roem. Syn. Pepon. 11, 48 (1846).
KARIVIA, Arn. in Hook. Journ. Bot. iii. (1841) 275.
LANDERSIA, Macfad. Fl. Jamaic. ii. 142 (1837).
MELOTHRIX, M. Laws. in Hook. f. Fl. Brit. Ind. i. 662 (1875).
MELOTRIA, P. Br. Hist. Jamaic. 124 (1756).
MUKIA, Arn. in Hook. Journ. Bot. iii. (1841) 276.
PILOGYNE, Schrad. Ind. Sem. Hort. Gotting. (1835).—Cf. Linnaea, xx. (1837) Litt. 89.
SOLENA, Lour. Fl. Cochinch. 514 (1790).
ZEHNERIA, Endl. Prod. Fl. Norf. 69 (1833).
amplexicaulis, *Cogn. in DC. Monog. Phan.* iii. 621.—Ind. or.
angolensis, *Cogn. l. c.* 606.—Afr. trop.
angustiloba, *Cogn. l. c.* 579.—Mexic.
Baueriana, *F. Muell. Fragm.* vi. 188.—Ins. Norfolk.
bicirrhosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 627.—Burma.
Candolleana, *Cogn. in DC. Monog. Phan.* iii. 578.—Paraguay.
capillacea, Cogn. l. c. 600 = triangularis.
celebica, *Cogn. l. c.* 625.—Ins. Celebes.
chlorocarpa, Engelm. ex S. Wats. Bibliog. Ind. N. Am. Bot. 395 = pendula.
cordata, *Cogn. in DC. Monog. Phan.* iii. 613.—Afr. austr.
cordifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 563.—Afr. trop.
cucumerina, *Naud. in Ann. Sc. Nat.* Sér. V. vi. (1866) 10.—China.
Cucumis, *Vell. Fl. Flum.* 29; i. t. 70.—Bras.
Cunninghamii, *F. Muell. ex Benth. Fl. Austral.* iii. 320.—Austral.
deltoidea, *Benth. in Hook. Niger Fl.* 368.—Afr. trop.
deltoidea, Thw. Enum. Pl. Zeyl. 124 = zeylanica.
edulis, Rafin. New Fl. Am. iv. 15 = pendula.
emirnensis, *Baker, in Journ. Linn. Soc.* xxi. (1884) 346.—Madag.
fluminensis, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 173.—Bras.
foetida, Desr. in Lam. Encyc. iv. 87 (fetida) = Kedrostis foetidissima.
grandifolia, *Torr. & Gray, Fl. N. Am.* i. 541.—Am. bor.
Grayana, *Cogn. in DC. Monog. Phan.* iii. 591.—Ins. Tahiti.
guadalupensis, Cogn. l. c. 580 = pervaga.
hederacea, *Cogn. l. c.* 611.—Afr. austr.
heterophylla, *Cogn. l. c.* 618.—As. trop.
hirsuta, *Cogn. in Mart. Fl. Bras.* vi. IV. 28. t. 5.—Bras.
Hookeri, *Cogn. in DC. Monog. Phan.* iii. 588.—Peruv.
indica, *Lour. Fl. Cochinch.* 35.—Cochinch.
japonica, *Maxim. ex Cogn. in DC. Monog. Phan.* iii. 599.—Japon.
javanica, *Cogn. l. c.* 625.—Java.
leiosperma, *Cogn. l. c.* 622.—Ind. or.
leucocarpa, Cogn. l. c. 601 = odorata.
longepedunculata, *Cogn. l. c.* 612.—Abyssin.
lucida, *Cogn. l. c.* 606.—Madag.
maderaspatana, *Cogn. l. c.* 623.—As. et Afr. trop.
Mannii, *Cogn. l. c.* 617.—Afr. trop.
marginata, Cogn. l. c. 593 = Rumphiana.
micrantha, F. Muell. ex Cogn. l. c. 603 = Muelleri.
microcarpa, Griseb. Cat. Pl. Cub. 112 = fluminensis.
microcarpa, Shuttl. ex Cogn. in DC. Monog. Phan. iii 587 = pendula.
microsperma, *Cogn. l. c.* 611.—Afr. trop.
minutiflora, *Cogn. l. c.*—Afr. trop.
monoica, Schrad. ex Steud. Nom. ed. II. ii. 123 = pendula.
mucronata, *Cogn. in DC. Monog. Phan.* iii. 608.—Ind. or.; Malaya.
Muelleri, *Benth. Fl. Austral.* iii. 320.—Austral.
nigra, Rafin. Ann. Nat. (1820) 15 = pendula.
obtusiloba, *Cogn. in DC. Monog. Phan.* iii. 616.—Afr. austr.
obtusiloba, Spruce, ex Cogn. l. c. 585 = fluminensis.
odorata, *Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 626.—Ind. or.; Malaya.
pendula, Brewer & S. Wats. Bot. Calif. i. 240 = Echinocystis Bigelovii.
pendula, *Linn. Sp. Pl.* 35.—Am. bor.; Ind. occ.
pendula, G. F. W. Mey. Prim. Fl. Esseq. 279 = fluminensis.
pendula, Sieber, ex Cogn. in DC. Monog. Phan. iii. 436 = Momordica Charantia.
Peneyana, *Cogn. l. c.* 592.—Abyssin.
pentaphylla, *Naud. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 196.—N. Caled.
perpusilla, *Cogn. in DC. Monog. Phan.* iii. 607.—Ind. or.; Malaya.
pervaga, *Griseb. Fl. Brit. W. Ind.* 289.—Ind. occ.
prunifera, Poepp. & Endl. Nov. Gen. et Sp. ii. 55. t. 174 = Selysia prunifera.
punctata, *Cogn. in DC. Monog. Phan.* iii. 615.—Afr. austr.
punctata, Rafin. New Fl. Am. iv. 15 = pendula.
punctatissima, *Cogn. in Mart. Fl. Bras.* vi. IV. 29.—Bras.
quinqueloba, Eckl. & Zeyh. ex M. Roem. Syn. Pepon. 99 = Coccinia quinqueloba.
racemosa, *Cogn. in DC. Monog. Phan.* iii. 617.—Afr. trop.
Rauwenhoffii, *Cogn. l. c.* 597.—Java.
Regelii, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 35 = japonica.
repanda, Rafin. New Fl. Am. iv. 14 = pendula.
Rumphiana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 25.—Java.
Rutenbergiana, *Cogn. in Bremen, Abh.* vii. (1882) 251.—Madag.
samoënsis, A. Gray, Bot. U. St. Expl. Exped. i. 641 = Baueriana.
scabra, *Naud. in Ann. Sc. Nat.* Sér. V. vi. (1866) 10.—Mexic.
scrobiculata, *Cogn. in DC. Monog. Phan.* iii. 605.—Abyssin.
Thwaitesii, Schweinf. Reliq. Kotschy. 44. t. 29 = tridactyla, zeylanica.
tomentosa, *Cogn. in DC. Monog. Phan.* iii. 614.—Abyssin.

MELOTHRIA:—
triangularis, *Benth. in Hook. Niger Fl.* 367.—Afr. trop.
tridactyla, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 562.—Afr. trop.
trifoliolata, F. Muell. Fragm. v. 181 = Alsomitra Hookeri.
trilobata, *Cogn. in Mart. Fl. Bras.* vi. IV. 26.—Bras.
uliginosa, *Cogn. l. c.*—Bras.
velutina, *Cogn. in DC. Monog. Phan.* iii. 613.—Afr. austr.
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 626.—Burma.
Warmingii, *Cogn. in Mart. Fl. Bras.* vi. IV. 27.—Bras.
zeylanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 626.—Zeylan.
zeylanica, Koen. ex Wight & Arn. Prod. 129 = Vitis pedata.

MELOTHRIX, M. Laws. in Hook. f. Fl. Brit. Ind. i. 662, (1875), err. typ. = **Melothria**, Linn. (Cucurb.).

MELOTRIA, P. Br. Hist. Jamaic. 124 (1756) = praec.

MELUCHIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 217 = **Melochia**, Dill. (Stercul.).

MELVILLA, A. Anders. in Journ. Arts & Sc. xxv. (1807) 207; et ex Lindl. Bot. Reg. t. 852 (1824) = **Cuphea**, P. Br. (Lythrar.).
speciosa, A. Anders. l. c. = C. Melvilla.

MEMAECYLUM, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) App. 213 = **Epigaea**, Linn. (Ericac.).

MEMECYLON, Linn. Diss. Dass. 16 (1747); Amoen. Acad. i. 399 (1749). *MELASTOMACEAE*, Benth. & Hook. f. i. 773.
LIJNDENIA, Zoll. & Mor. Syst. Verz. Mor. 10 (1846).
SCUTULA, Lour. Fl. Cochinch. 235 (1790).
SPATHANDRA, Guill. & Perr. Fl. Seneg. Tent. i. 313. t. 71 (1832).
VALIKAHA, Adans. Fam. ii. 84 (1763).
acuminatissimum, *Blume, Mus. Bot. Lugd. Bat.* i. 360.—Sumatra.
acuminatum, *Sm. in Rees, Cycl.* xxiii. n. 4.—Malacca.
acuminatum, Wall. Cat. n. 4105 = (Gen. ?).—Ind. or.
Afzelii, R. Br. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 156 = polyanthemos.
Afzelii, *G. Don, Gen. Syst.* ii. 655.—Afr. trop.
amabile, *Bedd. Forest. Man. Bot.* 115.—Ind. or.
ambiguum, Blume, Mus. Bot. Lugd. Bat. i. 359 = grande.
Amherstianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 557.—Burma.
amplexicaule, *Roxb. Hort. Beng.* [88]; *Fl. Ind.* ii. 260.—Ind. or.; Penang.
angulatum, *Reichb. ex DC. Prod.* iii. 5.—Ins. Maurit.
angustifolium, *Wight, Illustr.* i. 215.—Ind. or.
appendiculatum, *Blume, Mus. Bot. Lugd. Bat.* i. 361.—Borneo.
Arnottianum, *Wight, ex Thw. Enum. Pl. Zeyl.* 113.—Zeylan.
australe, F. Muell. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 159 = pauciflorum.
Barteri, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 462.—Afr. trop.
blakeoides, *G. Don, Gen. Syst.* ii. 655 (blackioides).—Afr. trop.
buxifolium, *Blume, Mus. Bot. Lugd. Bat.* i. 363.—Madag.
caeruleum, Hook. f. in Oliver, Fl. Trop. Afr. ii. 461 = M. Spathandra.
caeruleum, *Jack, in Malay. Misc.* i. (1820) v. 26.—Malaya.
calderense, A. Gray, Bot. U. St. Expl. Exped. i. 574. t. 71 = paniculatum.
Calleryanum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 275 = Cumingianum.
caloneuron, *Miq. Fl. Ind. Bat. Suppl.* 321.—Ins. Banca.
campanulatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 563.—Malaya.
campongum, C. B. Clarke, l. c. 554, err. typ., vide lampongum.
capense, *Eckl. & Zeyh. Enum.* 274.—Afr. austr.
capitellatum, Blume, Bijdr. 1094 = myrsinoides.

MEMECYLON:—
capitellatum, Heyne, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 158 = edule.
capitellatum, *Linn. Sp. Pl.* 349.—Zeylan.
capitellatum, Sm. in Rees, Cyclop. xxiii. n. 1 = sphaerocarpum.
capitellatum, Span. in Linnaea, xv. (1841) 203 = pauciflorum.
capitellatum, Willd. Sp. Pl. iv. 347 = floribundum.
celastrinum, Kurz, For. Fl. Brit. Burma, i. 515 = grande.
cerasiforme, *Kurz, l. c.* 516.—Chittagong.
cinnamomioides, *G. Don, Gen. Syst.* ii. 655.—Afr. trop.
clausiflorum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 274 = Cumingianum.
coeruleum, Auct. vide caeruleum.
confine, Blume, Mus. Bot. Lugd. Bat. i. 357 = glomeratum.
confusum, Blume, l. c. 362 = sphaerocarpum.
cordatum, *Desr. in Lam. Encyc.* iv. 89.—Ins. Maurit.
cordatum, Gardn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 158 = amplexicaule.
cordatum, Wall. Cat. n. 4100 = amplexicaule, caeruleum.
costatum, *Miq. Anal. Bot. Ind.* i. 29.—Malaya.
crassinerve, *Blume, Mus. Bot. Lugd. Bat.* i. 363.—Madag.
Cumingianum, *Presl, Epim. Bot.* 209.—Ins. Philipp.
Cumingii, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 273 (Cummingii) = Preslianum.
cuneatum, Thw. Enum. Pl. Zeyl. 112 = edule.
deccanense, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 560.—Ind. or.
depressum, Benth. in Wall. Cat. n. 4101 = amplexicaule.
diversifolium, *Presl, Epim. Bot.* 208.—Ins. Philipp.
dolichophyllum, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 266.—Madag.
Donianum, *Planch. ex Benth. in Hook. Niger Fl.* 357.—Afr. trop.
dumosum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 270 = buxifolium.
edule, *Roxb. Pl. Corom.* i. 59. t. 82.—Ind. or.; Malaya.
Elaeagni, *Blume, Mus. Bot. Lugd. Bat.* i. 356.—Ins. Seychell.
elegans, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 307.—Ins. Andam.
elegantulum, *Thw. Enum. Pl. Zeyl.* 112.—Zeylan.
ellipticum, *Thw. l. c.*—Zeylan.
eurhynchum, *Miq. Fl. Ind. Bat. Suppl.* 322.—Sumatra.
excelsum, *Blume, Bijdr.* 1094.—Java.
fasciculare, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 263, 282.—Afr. trop.
ferreum, Blume, Bijdr. 1095 = edule.
floribundum, Benth. in Wall. Cat. n. 4113 = Eugenia balsamea.
floribundum, *Blume, Mus. Bot. Lugd. Bat.* i. 361.—Java.
fuscescens, *Thw. Enum. Pl. Zeyl.* 111.—Zeylan.
garcinioides, *Blume, Mus. Bot. Lugd. Bat.* i. 358.—Sumatra.
Gardneri, *Thw. Enum. Pl. Zeyl.* 113.—Zeylan.
globiferum, Wall. Cat. n. 4108 = edule.
glomeratum, *Blume, Mus. Bot. Lugd. Bat.* i. 356.—Sumatra.
gracile, *Bedd. Forest. Man. Bot.* 115.—Ind. or.
grande, Blume, Bijdr. 1095 = costatum.
grande, Heyne, in Wall. Cat. n. 4102 = Heyneanum.
grande, *Retz. Obs.* iv. 26.—Zeylan.; Malaya.
grande, Sm. in Rees, Cyclop. xxiii. n. 6 = caeruleum.
grande, Wall. Cat. n. 4103, partim = edule.
grandifolium, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 266.—Madag.
Griffithianum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 82 = grande.
Griffithianum, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 274.—Burma.
Harveyi, *Seem. Fl. Vit.* 85.—Ins. Fiji.
hepaticum, *Blume, Mus. Bot. Lugd. Bat.* i. 357.—Borneo.
heteropleurum, *Blume, l. c.* 362.—Sumatra.
Heudelotii, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 265 = fasciculare.
Heyneanum, *Benth. in Wall. Cat.* n. 4102.—Ind. or.
Heyneanum, Wight & Arn. Prod. 319 = deccanense.
Hookeri, *Thw. Enum. Pl. Zeyl.* 113.—Zeylan.

MEMECYLON:—
Horsfieldii, Miq. Fl. Ind. Bat. i. I. 572 = grande.
intermedium, *Blume, Mus. Bot. Lugd. Bat.* i. 358.—Burma; Malaya.
jambosioides, Wight, Illustr. i. 215 = Heyneanum.
laevigatum, *Blume, Mus. Bot. Lugd. Bat.* i. 358.—Burma; Malaya.
lampongum, *Miq. Fl. Ind. Bat. Suppl.* 321.—Sumatra.
lanceolatum, Blanco, Fl. Filip. ed. I. 301 = Presli-anum.
latifolium, Boj. Hort. Maurit. 133 = cordatum.
laurifolium, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 277 = floribundum.
laurinum, *Blume, Mus. Bot. Lugd. Bat.* i. 359.—Malaya.
laxiflorum, DC. Prod. iii. 6 = sphaerocarpum.
laxiflorum, Wall. Cat. n. 4472 = grande.
leucanthum, Thw. Enum. Pl. Zeyl. 110 = edule.
ligustrifolium, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 117.—Ins. Hongkong.
ligustrinum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 271 = sphaerocarpum.
lilacinum, Zoll. & Mor. Syst. Verz. Zoll. 9 = myrsinoides.
longicuspe, *Baker, in Journ. Linn. Soc.* xx. (1883) 150.—Madag.
Loureirii, Triana, in Trans. Linn. Soc. xxviii. (1871) 156 = *Lurerii*.
lucidum, Presl, Epim. Bot. 209 = edule.
Lurerii, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 282 = umbellatum.
luteolum, *Miq. Fl. Ind. Bat. Suppl.* 323.—Sumatra.
lutescens, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 269.—Madag.
lutescens, Presl, Epim. Bot. 208 = caeruleum.
macrocarpum, Thw. Enum. Pl. Zeyl. 110 = ovoideum.
macrophyllum, *Thw. l. c.* 111.—Zeylan.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 557.—Malacca.
malabaricum, *Kostel. Allg. Med.-Pharm. Flora*, iv. 1517.—Ind. or.
manillanum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 276 = caeruleum.
Mannii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 461.—Afr. trop.
marginatum, *Blume, Mus. Bot. Lugd. Bat.* i. 363.—Malaya.
melastomoides, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 265.—Madag.
membranifolium, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 462.—Afr. trop.
micranthum, Blume, Mus. Bot. Lugd. Bat. i. 360 = oleaefolium, *Blume*.
microstomum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 557.—Malacca.
minutiflorum, *Miq. Fl. Ind. Bat. Suppl.* 323.—Sumatra.
myricoides, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 269.—Madag.
myrsinoides, *Blume, Mus. Bot. Lugd. Bat.* i. 356.—Malaya.
myrtifolium, Wall. Cat. n. 4111 = edule.
myrtiforme, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 270.—Madag.
Myrtilli, Blume, Mus. Bot. Lugd. Bat. i. 357 = laevigatum.
nigrescens, *Hook. & Arn. Bot. Beech. Voy.* 186.—China.
nigrescens, Miq. Fl. Ind. Bat. i. I. 578 = glomeratum.
nudum, Blume, Mus. Bot. Lugd. Bat. i. 361 = appendiculatum.
obovatum, Neraud, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 156 = sphaerocarpum.
obtusum, Wall. Cat. n. 4110 = edule.
oleaefolium, *Baker, in Journ. Linn. Soc.* xxi. (1884) 343.—Madag.
oleaefolium, *Blume, Mus. Bot. Lugd. Bat.* i. 359.—Sumatra.
oligoneurum, *Blume, l. c.* 353.—Java.
orbiculare, *Thw. Enum. Pl. Zeyl.* 112.—Zeylan.
ovatum, Sm. in Rees, Cyclop. xxiii. n. 3 = edule.
ovoideum, *Thw. Enum. Pl. Zeyl.* 112.—Zeylan.
pachyderme, Wall. Cat. n. 4104 = laevigatum.
paniculatum, *Jack, in Malay. Misc.* ii. (1822) n. VII. 62.—Malaya.

MEMECYLON:—
paniculatum, Miq. Fl. Ind. Bat. i. 1. 572 = acuminatissimum.
parviflorum, Blanco, Fl. Filip. ed. I. 300 = edule.
parvifolium, Thw. Enum. Pl. Zeyl. 113 = varians.
pauciflorum, *Blume, Mus. Bot. Lugd. Bat.* i. 356.—Ind. or.; Malaya; Austral.
pauciflorum, Wall. Cat. n. 414 = Canthium confertum.
Pervilleanum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 272 = M. Elaeagni.
phyllanthifolium, *Thw. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 159, nomen.—Zeylan.
Pierrei, *Hance, in Journ. Bot.* xv. (1877) 334.—Cambodia.
plebejum, *Kurz, For. Fl. Brit. Burma*, i. 513.—Burma.
polyanthemos, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 463.—Afr. trop.
prasinum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 275 = edule.
Preslianum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 157.—Ins. Philipp.
procerum, *Thw. Enum. Pl. Zeyl.* 415.—Zeylan.
prunifolium, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 268 = angulatum.
Pseudo-nigrescens, Blume, Mus. Bot. Lugd. Bat. i. 357 = glomeratum.
pulchrum, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 307 = caeruleum.
punctatum, Presl, Bot. Bemerk. 67 = edule.
pyrifolium, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 277 = edule.
pyrifolium, Presl, Epim. Bot. 210 = Cumingianum.
pyriforme, Wight, in Wall. Cat. n. 4106 (non Melastomac.).—Ind. or.
ramiflorum, Boj. Hort. Maurit. 132 = sphaerocarpum.
ramiflorum, Desr. in Lam. Encyc. iv. 88 = edule.
revolutum, *Thw. Enum. Pl. Zeyl.* 111.—Zeylan.
rhamnoideum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 271 = sphaerocarpum.
rhinophyllum, *Thw. Enum. Pl. Zeyl.* 110.—Zeylan.
roboreum, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 268.—Madag.
rostratum, *Thw. Enum. Pl. Zeyl.* 111.—Zeylan.
Royenii, Blume, Mus. Bot. Lugd. Bat. i. 360 = grande.
rubro-caeruleum, Thw. Enum. Pl. Zeyl. 415 = edule.
scutellatum, Hook. & Arn. Bot. Beech. Voy. 186 = ligustrifolium.
scutellatum, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 282 = edule.
sessile, Benth. in Wall. Cat. n. 4112 = amplexicaule, edule.
sparsiflorum, *Boj. Hort. Maurit.* 132 (Quid?).—Hab.?
Spathandra, *Blume, Mus. Bot. Lugd. Bat.* i. 361.—Afr. trop.
sphaerocarpum, *DC. Prod.* iii. 6.—Ins. Maurit.
strumosum, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 267.—Madag.
subquadrangulare, DC. Prod. iii. 6 = edule.
subtrinervium, *Miq. Fl. Ind. Bat. Suppl.* 322.—Sumatra.
sylvaticum, Thw. Enum. Pl. Zeyl. 110 = laevigatum.
terminale, *Dalz. in Hook. Kew Journ.* iii. (1851) 121.—Ind. or.
Thomsoni, Thw. Enum. Pl. Zeyl. 416 = varians.
Thouarsianum, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 267.—Madag.
tinctorium, Blume, Bijdr. 1094 = floribundum.
tinctorium, Koen. ex Willd. Sp. Pl. ii. 347 = edule.
tinctorium, Sieber, ex Blume, Mus. Bot. Lugd. Bat. i. 362 = sphaerocarpum.
trinerve, *DC. Prod.* iii. 5.—Ins. Maurit.
trinerve, Hassk. Cat. Hort. Bog. Alt. 259 = oligoneurum.
ulopterum, *DC. Prod.* iii. 5.—Madag.
umbellatum, Benth. Fl. Austral. iii. 293 = pauciflorum.
umbellatum, Blume, Bijdr. 1094 = intermedium.
umbellatum, Burm. f. Fl. Ind. 87 = edule.
umbellatum, Gaertn. Fruct. ii. 207 = Samara laeta.
umbellatum, *Kostel. Allg. Med.-Pharm. Flora*, iv. 1517.—Cochinch.
umbellatum, Presl, Epim. Bot. 208 = Preslianum.
varians, *Thw. Enum. Pl. Zeyl.* 112.—Zeylan.
virescens, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 463.—Afr. trop.

MEMECYLON:—
vitiense, *A. Gray, Bot. U. St. Expl. Exped.* i. 573.—Ins. Fiji.
Vogelii, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 263.—Afr. trop.
Vosmaerianum, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 356 = laevigatum.
Walkeri, Hook. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 564 = edule.
Wightianum, Triana, in Trans. Linn. Soc. xxviii. (1871) 158 = deccanense.
Wightii, *Thw. Enum. Pl. Zeyl.* 113.—Ind. or.; Zeylan.

MEMORA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 185 = **Adenocalymma**, Mart. (Bignon.).
alba, Miers, l. c. = Macfadyena bracteosa.
albiflora, Miers, l. c. = Bignonia albiflora.
Candolleana, Miers, l. c. = Spathodea Candolleana.
crassifolia, Miers, l. c.—Bras.
diffusa, Miers, l. c.—Venezuela.
fluminensis, Miers, l. c. = Bignonia fluminensis.
laserpitiifolia, Miers, l. c. = Bignonia laserpitiifolia.
nobilis, Miers, l. c.—Guian.
nodosa, Miers, l. c. = Bignonia nodosa.
patula, Miers, l. c. iv. (1864) 161.—Venezuela.
pedunculata, Miers, l. c. iii. (1863) 185 = Bignonia pedunculata.
perianthomega, Miers, l. c. = Bignonia perianthomega.
pinnatifolia, Miers, l. c.—Guian.
Schomburgkii, Miers, l. c. = Spathodea Schomburgkii.
tetraquetras, Miers, l. c. = Bignonia tetraquetra.
vaga, Miers, l. c.—Bras.

MEMORIALIS, Buch.-Ham. in Wall. Cat. sub n. 4598 et n. 4601 (1831) = **Pouzolzia**, Gaudich. (Urtic.).
ambigua, Wedd. in Arch. Mus. Par. viii. (1855–56) 430 = P. Wightii.
aquatica, Wedd. l. c. 418 = P. pentandra.
bracteata, Wedd. l. c. 421 = P. caudata.
caudata, Wedd. l. c. 425 = P. caudata.
ciliaris, Buch.-Ham. in Wall. Cat. n. 4598 = P. pentandra.
concinna, Wedd. in DC. Prod. xvi. 1. 235^{4}. = P. Wightii.
Dalzellii, Wedd. in Arch. Mus. Par. viii. (1855–56) 431 = P. integrifolia.
hirta, Wedd. in DC. Prod. xvi. 1. 235^{6} = P. Bennettiana.
hispida, Buch.-Ham. ex Wall. Cat. sub n. 4601 = P. Bennettiana.
integrifolia, Wedd. in DC. Prod. xvi. 1. 235^{8} = P. integrifolia.
lythroides, F. Muell. Fragm. v. 194 = P. quinquenervis.
mysorensis, Wedd. in Arch. Mus. Par. viii. (1855–56) 428 = P. Bennettiana.
nilghirensis, Wedd. l. c. 420 = P. Wightii.
parvifolia, Wedd. l. c. 432 = P. parvifolia.
pentandra, Wedd. in DC. Prod. xvi. 1. 235^{2} = P. pentandra.
quinquenervis, Buch.-Ham. in Wall. Cat. sub n. 4601 = P. quinquenervis.
scabra, Wedd. in Arch. Mus. Par. viii. (1855–56) 423 = P. scabra.
ternata, Wedd. l. c. 422 = P. Wightii.
triandra, Wedd. in DC. Prod. xvi. 1. 235^{9} = P. parvifolia.

MEMYCYLON, Griff. Notul. iv. 673 (1854), sphalm. = **Memecylon**, Linn. (Melastom.).

MENADENA, Rafin. Fl. Tellur. ii. 98 (1836) = **Maxillaria**, Ruiz & Pav. (Orchid.).
Parkeri, Rafin. l. c. 99 = M. Parkerii.

MENADENIUM, Rafin. Fl. Tellur. iv. 45 (1836) = **Zygopetalum**, Hook. (Orchid.).
rostratum, Rafin. l. c. = Z. rostratum.

MENAIS, Loefl. It. Hisp. 306 (1758). Gen. dub.—Cf. Benth. & Hook. f. ii. 838.
blanda, Blanco, Fl. Filip. ed. I. 139; ex DC. Prod. ix. 512 = seq.
mollis, Blanco, l. c. = Ehretia acuminata.
topiaria, Linn. Sp. Pl. ed. II. 251.—Am. austr.

MENALIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 = **Wallichia**? Roxb. (Palm.).

MENANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 43, 56 = **Menyanthes**, Tourn. (Gentian.).

MENARDA, Comm. ex A. Juss. Euph. Tent. 23. t. 6 (1824) = **Phyllanthus**, Linn. (Euphorb.).
capillaris, Baill. Adansonia, i. (1860–61) 85 = P. capillaris.
cryptophila, A. Juss. Euph. Tent. 109 = P. cryptophilus.
Goudotiana, Baill. Adansonia, ii. (1861–62) 62 = P. Goudotianus.
linifolia, Baill. l. c. i. (1860–61) 84 = P. pentandrus.
nummularifolia, Baill. Etud. Gén. Euph. 609 = P. nummulariaefolius.
pulchella, Baill. l. c. = P. multiflorus.

MENDEVILLA, Poit. in Rev. Hortic. Sér. II. iv. (1845–46) 166, sphalm. = **Mandevilla**, Lindl. (Apocyn.).

MENDEZIA, DC. Prod. v. 532 (1836) = **Spilanthes**, Linn. (Compos.).
bicolor, DC. l. c. 533 = S. bicolor & Zinnia bicolor.

MENDOCINA, Walp. Ann. iii. 209 (1852–53) sphalm. = seq.

MENDONCIA, Vell. ex Vand. Fl. Lusit. 43, t. 3. f. 22 (1788). *ACANTHACEAE*, Benth. & Hook. f. ii. 1072.
ENGELIA, Karst. ex Nees, in DC. Prod. xi. 721 (1847).
MENDOZIA, Ruiz et Pav. Prod. 89, t. 17 (1794).
albida, Vell. Fl. Flum. 263; vi. t. 85 = puberula.
angustifolia, Poepp. ex Nees, in DC. Prod. xi. 52 = hirsuta.
aspera, *Nees, l. c.* 51.—Peruv.
coccinea, *Vell. Fl. Flum.* 263; vi. t. 86.—Bras.
costaricana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 113.—Costa Rica.
glabra, *Nees, in DC. Prod.* xi. 52.—Peruv.
hirsuta, *Nees, l. c.*—Peruv.
Hoffmannseggiana, *Nees, l. c.* 50.—Bras.
madagascariensis, *Radlk. in Bremen, Abh.* viii. (1884) 467.—Madag.
Meyeniana, *Nees, in DC. Prod.* xi. 51.—Bras.
multiflora, *Nees, l. c.* 52.—Bras.
Perrottetiana, *Nees, l. c.* 53.—Guian.
pilosa, *Nees, l. c.* 50.—Bras.
puberula, *Nees, l. c.* 53.—Bras.
pubescens, Nees, l. c. = puberula.
racemosa, *Nees, l. c.* 54.—Peruv.
Riedeliana, *Nees, l. c.* 51.—Bras.
rotundifolia, Nees, l. c. = aspera.
Schomburgkiana, *Nees, l. c.* 53.—Guiana.
Schottiana, *Nees, l. c.* 51.—Bras.
Sellowiana, *Nees, l. c.* 53.—Bras.
speciosa, *Nees, l. c.* 54.—Peruv.
Splitgerberiana, *De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 353.—Guiana.
squamuligera, *Nees, in DC. Prod.* xi. 50.—Guiana.
strigosa, Poepp. ex Nees, l. c. 52 = multiflora.
tomentosa, *Poepp. ex Nees, l. c.*—Bras.
Velloziana, *Nees, l. c.*—Bras.

MENDONI, Adans. Fam. ii. 48, 576 (1763) = **Gloriosa**, Linn. (Liliac.).

MENDOZIA, Ruiz et Pav. Prod. 89, t. 17 (1794) = **Mendoncia**, Vell. (Acanth.).
spera, Ruiz & Pav. Syst. Veg. 158 = Mendoncia aspera.
glabra, Poepp. & Endl. Nov. Gen. et Sp. iii. 10 = Mendoncia glabra.
granulosa, Poepp. ex Nees, in Mart. Fl. Bras. ix. 11 = Mendoncia pilosa.
hirsuta, Poepp. & Endl. Nov. Gen. et Sp. iii. 10 = Mendoncia hirsuta.
Hoffmannseggiana, Nees, in Mart. Fl. Bras. ix. 9 = Mendoncia Hoffmannseggiana.
lancifolia, Presl, Bot. Bemerk. 147.—Bras.
Meyeniana, Nees, in Mart. Fl. Bras. ix. 11 = Mendoncia Meyeniana.
multiflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 10 = Mendoncia multiflora.
pilosa, Mart. Nov. Gen. et Sp. iii. 21 = Mendoncia pilosa.
puberula, Mart. l. c. 24. t. 211 = Mendoncia puberula.

MENDOZIA:—
pubescens, Hoffmgg. ex Nees, in Mart. Fl. Bras. ix. 9 = Mendoncia Hoffmannseggiana.
pubescens, Poepp. & Endl. Nov. Gen. et Sp. iii. 11 = Mendoncia puberula.
racemosa, Ruiz & Pav. Syst. Veg. 158 = Mendoncia racemosa.
Riedeliana, Nees, in Mart. Fl. Bras. ix. 12 = Mendoncia Riedeliana.
rotundifolia, Poepp. & Endl. Nov. Gen. et Sp. iii. 10 = Mendoncia aspera.
Schottiana, Pohl, ex Nees, in Mart. Fl. Bras. ix. 12 = Mendoncia Schottiana.
Selloviana, Nees, l. c. 10 = Mendoncia Sellowiana.
strigosa, Poepp. ex Nees, l. c. 12 = Mendoncia multiflora.
tomentosa, Poepp. l. c. = Mendoncia tomentosa.
Velloziana, Mart. Nov. Gen. et Sp. iii. 22 = Mendoncia Velloziana.

MENEGHINIA, Endl. Gen. 648 (1839) = **Arnebia**, Forsk. (Borag.).

MENEGHINIA, Vis. in Atti 8va Riun. Scienz. Ital. (1846) 560 = **Niphaea**, Lindl. (Gesnerac.).
alba, Vis. l. c. = N. oblonga.

MENEPHORA, Rafin. Fl. Tellur. iv. 46 (1836) = **Cypripedium**, Linn. (Orchid.).
bicolor, Rafin. l. c. = C. purpuratum?

MENESTORIA, DC. Prod. iv. 390 (1830). Gen. delendum.—Cf. Benth. & Hook. f. ii. 29.
Hameliae, DC. l. c. (= Adenosacmes sp.).—Nepal.
Mussaendae, DC. l. c. (= Mussaendae sp.).—Nepal.
rigida, DC. l. c. (= Randiae sp.).—Nepal.
Tocoyenae, DC. l. c. = Mussaenda Roxburghii?

MENESTRATA, Vell. Fl. Flum. 199 (1825); v. t. 2 (1827) = **Persea**? Gaertn. (Laurin.).
racemosa, Vell. ll. cc. (= *Oreodaphne crassiramea*).—Bras.

MENETHO, Rafin. Fl. Tellur. ii. 93 (1836) = **Frankenia**, Linn.
sedifolius, Rafin. l. c. = F. tetrapetala.

MENGEA, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 405 (Pl. Meyen.) = **Amaranthus**, Linn.
californica, Moq. in DC. Prod. xiii. II. 270 = A. californicus.
peruviana, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 406.—Peruv.
tenuifolia, Moq. in DC. Prod. xiii. II. 271 = A. tenuifolius.

MENIANTHES, Neck. Delic. Gallo-Belg. i. 105 (1768) = **Menyanthes**, Tourn. (Gentian.).

MENIANTHUS, Gouan, Hort. Monsp. 89 (1762) = praec.

MENICHEA, Sonner. Voy. N. Guin. 133 (1776) = **Barringtonia**, Forst. (Myrtac.).
rosata, Sonner. l. c.—Afr. trop.

MENICOSTA, D. Dietr. Syn. Pl. ii. 923 (1839) = MENISCOSTA, Blume = **Sabia**, Colebr.

MENINIA, Fua, ex Hook. f. Bot. Mag. t. 6043 (1873) = **Cystacanthus**, T. Anders. (Acanth.).
turgida, Fua, ex Hook. f. l. c. = C. turgidus.

MENIOCUS, Desv. Journ. Bot. iii. (1814) 173 = **Alyssum**, Tourn. (Crucif.).
aureus, Fenzl. Pugill. Pl. Nov. Syr. 13 = A. aureum.
australasicus, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 297 = A. linifolium.
Ehrenbergii, Presl, Bot. Bemerk. 9 = A. libycum.
filifolius, Jaub. & Spach, Illustr. i. 107 = A. meniocoides.
grandiflorus, Jaub. & Spach, l. c. 105 = A. aureum.
hirsutus, Boiss. & Huet, Diagn. Ser. II. v. 32 = A. Huetii.
linifolius, DC. Syst. ii. 325 = A. linifolium.
serpyllifolius, Desv. Journ. Bot. iii. (1814) 173 = A. linifolium.
stylaris, Boiss. Diagn. Ser. II. vi. 16 = A. stylare.

MENISCOSTA, Blume, Bijdr. 28 (1825) = **Sabia**, Colebr.
javanica, Blume, l. c. 29 = S. Meniscosta.
scandens, Spreng. Syst. iv. Cur. Post. 114 = S. Meniscosta.

MENISPERMUM, [Tourn.] Linn. Syst. ed. I (1735).
MENISPERMACEAE, Benth. & Hook. f. i. 37.
TRILOPHUS, Fisch. Hort. Gorenk. ed. II. 56 (1812).
Abuta, Lam. Encyc. iv. 100 = Abuta rufescens.
acuminatum, Lam. l. c. 101 = Tiliacora racemosa.
acutum, *Thunb. Fl. Jap.* 193.—Japon.
amarum, Willd. Sp. Pl. iv. 826 = Aristolochia glaucescens.
angulatum, Moench, Meth. 277 = canadense.
bantamense, Spreng. Syst. iv. Cur. Post. 143 = Cocculus bantamensis.
brachystachyon, Spreng. Syst. ii. 157 = Cocculus brachystachyus.
Calumba, Stokes, Bot. Mat. Med. iv. 700 = Jateorhiza Columba.
canadense, *Linn. Sp. Pl.* 340.—Am. bor.
canadense, Pall. Reise, ii. 687 = dauricum.
capense, Thunb. Prod. Pl. Cap. 85 = Cissampelos torulosa.
carinianum, Mill. Gard. Dict. ed. VIII. n. 3, sphalm. = *carolinum*.
carolinianum, Hill, Veg. Syst. xvi. 27 = Cocculus carolinus.
carolinum, Linn. Sp. Pl. 340 = Cocculus carolinus.
Chondrodendron, Spreng. Syst. ii. 155 = Cocculus Chondodendrum.
cocculiferum, Stokes, Bot. Mat. Med. iii. 49 = Anamirta paniculata.
Cocculus, Linn. Sp. Pl. 340 = Anamirta paniculata.
Cocculus, Willd. Sp. Pl. iv. 825 = Cocculus glaucus, Pachygone ovata.
Columba, Roxb. Hort. Beng. 72; Fl. Ind. iii. 807 = Jateorhiza Columba.
cordifolium, Willd. Sp. Pl. iv. 826 = Tinospora cordifolia.
coriaceum, Spreng. Syst. iv. Cur. Post. 143 = Cocculus coriaceus.
corymbosum, Spreng. l. c. = Cocculus corymbosus.
Cotoneaster, Spreng. l. c. ii. 156 = Cocculus Cotoneaster.
crispum, Linn. Sp. Pl. ed. II. 1468 = Tinospora crispa.
dauricum, *DC. Syst.* i. 540.—Dahuria.
dioicum, *Griff. Notul.* iv. 315.—Burma.
diversifolium, Spreng. Syst. ii. 155 = Cocculus diversifolius.
domingense, Spreng. l. c. 157 = Pachygone domingensis.
dubium, Muhl. ex Rafin. Fl. Ludov. 138, nomen = Calycocarpum Lyonii?
edule, Vahl, Symb. Bot. i. 80 = Cocculus Cebatha.
ellipticum, Poir. Encyc. Suppl. iii. 657 = Cocculus Leaeba.
Epibaterium, Spreng. Syst. ii. 156 = Cocculus Epibaterium.
fenestratum, Gaertn. Fruct. i. 219. t. 46. f. 5 = Coscinium fenestratum.
flavescens, Lam. Encyc. iv. 98 = Cocculus flavescens.
flavum, Linn. Amoen. Acad. iv. 128 = Cocculus flavescens.
glabrum, Burm. f. Fl. Ind. 316 [recte 216] = Tinospora cordifolia.
glabrum, Koen. ex Willd. Sp. Pl. iv. 797 = Tiliacora racemosa.
glaucescens, Spreng. Syst. iv. Cur. Post. 143 = Cocculus glaucescens.
glaucum, Lam. Encyc. iv. 100 = Cocculus glaucus.
gomphioides, Spreng. Syst. ii. 157 = Cocculus gomphioides.
hastatum, Lam. Encyc. iv. 98 = Cocculus villosus.
heteroclitum, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 817 = Anamirta paniculata.
hexagynum, Roxb. ll. cc. 72; 816 = Cocculus hexagynus.
hirsutum, Linn. Sp. Pl. 341 = Cocculus villosus.
japonicum, Roxb. ex DC. Syst. i. 516 = Stephania rotunda.
japonicum, Thunb. Fl. Jap. 193 = Cocculus japonicus.
lacunosum, Lam. Encyc. iv. 98 = Anamirta paniculata.
lanuginosum, Spreng. Syst. iv. Cur. Post. 143 = Cocculus lanuginosus.
laurifolium, Roxb. Hort. Beng. 72; Fl. Ind. iii. 815 = Cocculus laurifolius.
Leaeba, Delile, Fl. Egypt. 284 = Cocculus Leaeba.
leptostachyon, Spreng. Syst. ii. 157 = Cocculus leptostachyus.

MENISPERMUM:—
Limacia, Spreng. Syst. ii. 155 = Cocculus Limacia.
Lyoni, Pursh, Fl. Am. Sept. ii. 371 = Calycocarpum Lyonii.
malabaricum, Lam. Encyc. iv. 96 = Tinospora malabarica.
megaspermum, Roxb. Hort. Beng. 72 = Cocculus megaspermus.
milleflorum, Spreng. Syst. ii. 157 = Cocculus milleflorus.
monadelphum, Roxb. ex Wight & Arn. Prod. 446 = Anamirta paniculata.
Myosotis, Linn. Syst. ed. X. 992 = seq.
myosotoides, Linn. Sp. Pl. 341 = Cocculus villosus.
myosuroides, Hill, Veg. Syst. xvi. 27 = praec.?
oblongifolium, Spreng. Syst. ii. 155 = Cocculus oblongifolius.
orbiculatum, Linn. Sp. Pl. 341 = Cissampelos Pareira.
orbiculatum, Thunb. Fl. Jap. 194 = Cocculus Thunbergii.
ovalifolium, Pers. Syn. ii. 628 = Cocculus Thunbergii.
palmatum, Lam. Encyc. iv. 99 = Jateorhiza Columba.
peltatum, Forst. ex DC. Syst. i. 517 = Cocculus Forsteri.
peltatum, Lam. Encyc. iv. 96 = Cyclea peltata.
polycarpon, Roxb. Hort. Beng. 72; Fl. Ind. iii. 816 = Tiliacora racemosa.
populifolium, Spreng. Syst. ii. 156 = Anamirta paniculata.
radiatum, Lam. Encyc. iv. 100 = Tiliacora racemosa.
reniforme, Spreng. Syst. ii. 156 = Cocculus Nephroia.
rimosum, Blanco, Fl. Filip. ed. I. 810 = Tinospora crispa.
rimosum, Spreng. Syst. iv. Cur. Post. 143 = Cocculus rimosus.
rotundifolium, Spreng. Syst. ii. 156 = Cocculus rotundifolius.
Roxburghii, Spreng. l. c. 155 = Stephania rotunda.
sagittatum, Spreng. l. c. = Muehlenbeckia sagittifolia.
Schimperi, Hochst. ex Steud. Nom. ed. II. ii. 124 = Stephania abyssinica.
smilacinum, DC. Syst. i. 541 = canadense.
tamoides, Spreng. Syst. ii. 156 = Chondodendron tamoides.
tinctorium, Spreng. l. c. = Fibraurea tinctoria.
tomentosum, Roxb. Hort. Beng. 72; Fl. Ind. iii. 813 = Tinospora tomentosa.
triandrum, Roxb. ll. cc. 72; 816 = Limacia triandra.
triflorum, Spreng. Syst. ii. 157 = Cocculus triflorus.
trilobum, Thunb. Fl. Jap. 194 = Cocculus trilobus.
tuberculatum, Lam. Encyc. iv. 96 = Tinospora crispa.
verrucosum, Roxb. ex Flem. in As. Res. xi. (1810) 171 = Tinospora crispa.
villosum, Lam. Encyc. iv. 97 = Cocculus villosus.
villosum, Roxb. Hort. Beng. 72; Fl. Ind. iii. 812 = Pericampylus incanus.
virginicum, Linn. Sp. Pl. 340 = Cocculus carolinus.

MENKEA, Lehm. Ind. Sem. Hort. Hamb. (1843) 8.
CRUCIFERAE, Benth. & Hook. f. i. 83.
australis, *Lehm. Ind. Sem. Hort. Hamb.* (1843) 8.—Austral.
australis, F. Muell. Fragm. ii. 142 = draboides.
draboides, *Hook. f. ex Benth. Fl. Austral.* i. 80.—Austral.
procumbens, F. Muell. Fragm. ii. 142 = australis.
sphaerocarpa, *F. Muell. l. c.* viii. 223.—Austral.

MENNICHEA, Steud. Nom. ed. II. ii. 121 (1841) = MENICHEA, Sonner. = **Barringtonia**, Forst. (Myrtac.).

MENOCERAS, Lindl. Veg. Kingd. 695 (1847) = **Velleia**, Sm. (Gooden.).

MENODORA, Humb. & Bonpl. Pl. Aequin. ii. 98. t. 110 (1809). *OLEACEAE*, Benth. & Hook. f. ii. 674.
BOLIVARIA, Cham. & Schlecht. in Linnaea, i. (1826) 207. t. 4.
CALYPTROSPERMUM, A. Dietr. Sp. Pl. i. 226 (1831).
africana, *Hook. Ic. Pl.* t. 586.—Afr. austr.
Coulteri, *A. Gray, in Am. Journ. Sc.* Ser. II. xiv. (1852) 44.—Mexic.
decemfida, *A. Gray, l. c.*—Chili.
helianthemoides, *Humb. & Bonpl. Pl. Aequin.* ii. 98. t. 110.—Mexic.
heterophylla, *Moric. ex DC. Prod.* viii. 316.—Texas.
integrifolia, *Steud. Nom.* ed. II. ii. 124.—Reg. Argent.
juncea, *Harv. Gen. S. Afr. Pl.* ed. II. 220.—Afr. austr.

MENODORA:—
linoides, *Phil. in Linnaea*, xxxiii. (1864-65) 173.—Chili.
longiflora, *A. Gray, in Am. Journ. Sc.* Ser. II. xiv. (1852) 45.—Texas.
mexicana, *A. Gray, l. c.*—Mexic.
pinnatifida, *Mart. Del. Sem. Hort. Monac.* (1852); *ex Ann. Sc. Nat.* Sér. III. xix. (1853) 366.—Texas.
pubens, *A. Gray, in Am. Journ. Sc.* Ser. II. xiv. (1852) 45.—N. Mexic.
robusta, *A. Gray, l. c.* 43.—Patagon.
scabra, *A. Gray, l. c.* 44.—N. Mexic.
scoparia, *Engelm. ex A. Gray, in Bot. Calif.* i. 471.—Am. bor. occ.
spinescens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 388.—Am. bor. occ.
trifida, *Steud. Nom.* ed. II. ii. 124.—Bras.

MENOMPHALUS, Pomel, Nouv. Mat. Fl. Atl. 32 (1874) = **Centaurea**, Linn. (Compos.).
acaulis, Pomel, l. c. 33 = C. acaulis.
amourensis, Pomel, l. c. = C. amourensis.
Choulettiana, Pomel, l. c. = C. Choulettiana.
Cineraria, Pomel, l. c. = C. Cineraria.
nanus, Pomel, l. c. = C. nana.
omphalothrix, Pomel, l. c. = C. omphalodes.
pubescens, Pomel, l. c. = C. incana.
rupicola, Pomel, l. c. = C. rupicola.
Saharae, Pomel, l. c. = C. Saharae.
Scabiosa, Pomel, l. c. = C. Scabiosa.

MENONVILLEA, R. Br. ex DC. Syst. ii. 419 (1821).
CRUCIFERAE, Benth. & Hook. f. i. 90.
CYMATOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 309.
DISPELTOPHORUS, Lehm. Ind. Sem. Hort. Hamb. (1836.)
angustifolia, Presl, Bot. Bemerk. 9 = filifolia.
aptera, *Phil. Fl. Atac.* 9.—Chili.
chilensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 551.—Chili.
filifolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 34.—Chili.
flexuosa, *Phil. in Anal. Univ. Chil.* (1872) 674.—Chili.
Gayi, *Phil. in Linnaea*, xxviii. (1856) 610.—Chili.
linearis, Bert. ex Presl, Bot. Bemerk. 9 = filifolia.
linearis, *DC. Syst.* ii. 420.—Peruv.; Chili.
orbiculata, *Phil. Fl. Atac.* 9.—Chili.
parviflora, *Phil. l. c.* 8.—Chili.
parvifolia, Phil. f. Cat. Pl. Vasc. Chil. 16, sphalm. = praec.
pinnatifida, *Barn. in C. Gay, Fl. Chil.* i. 182.—Chili.
robustula, *Steud. in Flora*, xxviii. (1856) 411.—Chili.
trifida, *Phil. in Linnaea*, xxviii. (1856) 610.—Chili.
trifida, Steud. Nom. ed. II. ii. 125 = filifolia.
virgata, *Phil. in Linnaea*, xxviii. (1856) 671.—Chili.

MENOPHYLA, Rafin. Fl. Tellur. iii. 45 (1836) = **Rumex**, Linn. (Polygon.).
Lunaria, Rafin. l. c. = R. Lunaria.
polygama, Rafin. l. c. = R. Lunaria.

MENOTRICHE, Steetz, in Peters, Reise Mossamb. Bot. 472 (1862) = **Wedelia**, Jacq. (Compos.).
strigosa, Steetz, l. c. 475 = W. Menotricha.

MENTHA, [Tourn.] Linn. Syst. ed. I (1735).
LABIATAE, Benth. & Hook. f. ii. 1182.
AUDIBERTIA, Benth. in Bot. Reg. sub n. 1282 (1829).
MENTHELLA, Pérard, in Bull. Soc. Bot. Fr. xvii. (1870) 205.
PULEGIUM, Ray, ex Mill. Gard. Dict. ed. VI. (1752).
acuta, Opiz, Seznam, 64 = aquatica.
acutata, Opiz, in Lotos, iii. (1853) 209 = aquatica.
acutifolia, Rabenhorst, Fl. Lusat. 160 = gentilis.
acutifolia, Sm. in Trans. Linn. Soc. v. (1800) 203 = aquatica.
acuti-serrata, Opiz, Natural. x. (1825) 194 = aquatica.
adspersa, Moench, Meth. 379 = aquatica.
adulterina ×, Borb.—Cf. Oestr. Bot. Zeitschr. xxxiii. (1883) 119.—Europ.
affinis, Bor. Fl. Centr. Fr. ed. III. ii. 509 = aquatica.
affinis, Opiz, Nom. Bot. 74 (Quid?).—Europ.
Agardhiana, Fries, Nov. Fl. Suec. ed. II. 184 = gentilis.

MENTHA:—
Agardhiana, Hartm. Skand. Fl. ed. V. 43 = sativa.
agrestris, Sole, Menth. Brit. 33. t. 14 = arvensis.
alata, Koch, Syn. Fl. Germ. ed. I. 552, sphalm. = *elata*.
Allionii, Bor. Fl. Centr. Fr. ed. III. ii. 515 = arvensis.
alopecuroides, Hull, Brit. Fl. i. 126 = sylvestris.
alpigena, *A. Kern. Sched. Fl. Exsicc.* ii. 121.—Austria.
amaurophylla, Timb. in Bull. Soc. Bot. Fr. vii. (1860) 257 = rotundifolia.
ambigua, Guss. Enum. Pl. Inar. 253 = sylvestris.
ambigua, Personnat, in Act. Congr. Bot. Par. (1867) 161 = rotundifolia.
angustifolia, *Host, Fl. Austr.* ii. 148.—Austria.
angustifolia, Schreb. ex Schweigg. & Koerte, Fl. Erlang. ii. 8 = arvensis.
anomala, Hérib. in Bull. Soc. Bot. Fr. xxvii. (1880) 169 = arvensis?
aquatica, *Linn. Sp. Pl.* 576.—Europ.; As. et Afr. bor.
aquatica, O. F. Muell. in Fl. Dan. t. 673 = Ballota nigra.
arguta, Opiz, Natural. viii. (1824) 69 = arvensis.
aromatica, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 197 = aquatica.
aromatica, Salisb. Prod. 80 = M. Pulegium.
arvensis, *Linn. Sp. Pl.* 577.—Europ.; As. bor.
atrata, Schur, Enum. Pl. Transs. 516 = viridis.
atrovirens, Host, Fl. Austr. ii. 152 = arvensis.
augusta, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 198 = aqnatica.
auneticensis, Opiz, Seznam, 64 = aquatica, arvense?
auricularia, Blanco, Fl. Filip. ed. II. 329 = Pogostemon Cablin.
auricularia, Linn. Mant. i. 81 = Dysophylla auricularia.
aurita, Weihe, ex Fresen. in Syll. Ratisb. ii. (1828) 226 = aquatica.
australis, *R. Br. Prod.* 505.—Austral.
austriaca, Jacq. Fl. Austr. v. 14 = arvensis.
Avellini, Tod. ex Bertol. Fl. Ital. vi. 96 = aquatica.
Ayassei, Malinv. in Bull. Soc. Bot. Fr. xxiv. (1877) 234 = aquatica.
badensis, Fellm. ex Ledeb. Fl. Ross. iii. 340 = arvensis.
badensis, C. C. Gmel. Fl. Bad. ii. 604 = arvensis.
baikalensis, Georgi, Beschr. Russ. Reich. III. v. 1083 = Elsholtzia cristata.
ballotaefolia, Opiz, Natural. iv. (1823) [21] = aquatica.
balsamea, Reichb. Fl. Germ. Excurs. 309 = viridis.
balsamea, Willd. Enum. Hort. Berol. 608 = piperita, suavis.
Bandeliana, Opiz, Nom. Bot. 70 (Quid?).—Europ.
barbata, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 211 = arvensis.
Barkhausiana, Opiz, ex Steud. Nom. ed. II. ii. 126, nomen (Quid?).—Europ.
belgica, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 324.—Belgium.
bellojocensis, *Gillot, in Ann. Soc. Bot. Lyon*, viii. (1881) 26.—Gallia.
beneschina, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 206.—Europ.
Billotiana, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 319.—Gallia.
biserrata, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 202 = aquatica.
blanda, DC. Pl. Rar. Hort. Genev. 27. t. 8 = Elsholtzia incisa.
blanda, Wall. ex Benth. Lab. Gen. et Sp. 162 = Elsholtzia blanda.
Boenninghausiana, Opiz, Nom. Bot. 64 (Quid?).—Europ.
borealis, Michx. Fl. Bor. Am. ii. 2 = canadensis.
brachiata, Weihe, ex Fresen. in Syll. Ratisb. ii. (1828) 226 = aquatica.
Brachtii, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 196.—Europ.
brachystachya ×, Borb. in Oestr. Bot. Zeitschr. xxx. (1880) 19.—Europ.
bracteolata, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 211 = arvensis.
Braunii, *Oborny, in Verh. Naturf. Ver. Bruenn*, xxii. II. (1884) 370.—Europ.
brevispicata, Lehm. Ind. Sem. Hort. Hamb. (1828) 16; et in Linnaea, v. (1830) 380 = viridis.
Brideliana, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 333.—Gallia.
Brittingeri, Opiz, Natural. xi. (1826) 300 = sylvestris.
bugulaefolia, Weihe, ex Fresen. in Syll. Ratisb. ii. (1828) 226 = aquatica.
Burckhardtiana, Opiz, Natural. xi. (1826) 301 = rotundifolia.

MENTHA:—
Cablin, Blanco, Fl. Filip. ed. I. 473 = Pogostemon Cablin.
caerulescens, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 216 = sylvestris.
calliantha, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 36.—Persia.
campestris, Schur, Enum. Pl. Transs. 937 (Index) = arvensis.
canadensis, *Linn. Sp. Pl.* 577.—Am. bor.; China.
canariensis, Lam. Encyc. iv. 113 = Bystropogon origanifolius.
canariensis, Linn. Sp. Pl. 578 = Bystropogon canariensis.
candicans, Mill. Gard. Dict. ed. VIII. n. 3; Crantz, Stirp. Austr. ed. II. fasc. iv. (1769) 330 = sylvestris.
canescens, Roth, Catalect. fasc. ii. 46 = sylvestris.
cantalica, Hérib. in Bull. Soc. Bot. Fr. xxvii. (1880) 167 = aquatica.
capensis, Burch. Trav. i. 255 = sylvestris.
capensis, Thunb. Prod. Pl. Cap. 95 = salicina.
capitata, Gilib. Fl. Lituan. i. 72 = aquatica.
cardiaca, *Baker, in Journ. Bot.* iii. (1865) 245.—Europ.
carinthiaca, Host, Fl. Austr. ii. 149 = arvensis.
carniolica, Host, l. c. 146 = arvensis.
cechobrodensis, Opiz, in Lotos, iii. (1853) 210 = sativa.
Cephalotes, Heyne, ex Steud. Nom. ed. II. ii. 126 = Acrocephalus capitatus.
cervina, Linn. Sp. Pl. 578 = Preslia cervina.
chalepensis, Mill. Gard. Dict. ed. VIII. n. 10 = sylvestris.
ciliata, Opiz, ex Fresen. in Syll. Ratisb. ii. (1828) 240 = gentilis.
cinerascens, *Timb. ex Déségl.—Cf. Bull. Soc. Bot. Fr.* xxv. (1878) *Rev. Bibl.* 109.—Europ.
cinerea, *Holuby, in Oestr. Bot. Zeitschr.* xxvi. (1876) 149.—Europ.
cinerea, Opiz, Seznam, 64 = gentilis.
cinerea, Rafin. New Fl. Am. iv. 94 (Quid?).—Am. bor.
citrata, Ehrh. Beitr. vii. 150 = aquatica.
clinopodiifolia, *Krock. Fl. Siles. Suppl.* ii. 144.—Europ.
coccinea, *Roxb. ex Steud. Nom.* ed. II. ii. 126, nomen.—Ind. or.
coerulea, Opiz, Natural. iv. (1823) [21] = arvensis.
concolor, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 35.—Persia.
cordata, Jan, ex Nym. Consp. 596 = aquatica.
cordifolia, Opiz, ex Fresen. in Syll. Ratisb. ii. (1828) 232 = viridis.
corymbosa, Lam. Encyc. iv. 114 = Bystropogon canariensis.
crenata, Becker, Fl. Franc. i. 225 = arvensis.
Crépiniana, Durand, in Bull. Soc. Bot. Belg. xvi. (1877) 115 = gentilis.
cretica, Hort. ex Benth. Lab. Gen. et Sp. 170 = sylvestris.
crispa, Linn. Sp. Pl. 576 = aquatica.
crispa, Roth, Tent. Fl. Germ. i. 248 = viridis.
crispa, Tenore, Fl. Nap. ii. 29 = sylvestris.
crispata, Schrad. ex Willd. Enum. Hort. Berol. 608 = viridis.
crispula, Wender. in Flora, xii. (1829) I. Erg. 27 = aquatica.
cristata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 115 = Elsholtzia cristata.
croatica, Weld. in Flora, xiii. (1830) 218 = Micromeria rupestris.
cruciata, Opiz, ex Steud. Nom. ed. II. ii. 126 = gentilis.
Cunninghamii, *Benth. in DC. Prod.* xii. 174.—N. Zel.
cuspidata, Opiz, ex Steud. Nom. ed. II. ii. 126, nomen (Quid?).—Europ.
dahurica, *Fisch. ex Benth. Lab. Gen. et Sp.* 181.—Sibir.
dalmatica, Tausch, in Syll. Ratisb. ii. (1828) 249 = arvensis.
Decloetiana, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 215 = sylvestris.
deflexa, Dum. Fl. Belg. 49 = arvensis.
densiflora, Opiz, Seznam, 65 = arvensis.
dentata, Moench, Meth. 380; Willd. Sp. Pl. iii. 80 = gentilis.
Deseglisei, Malinv. in Bull. Soc. Bot. Fr. xxviii. (1881) 369 = aquatica.
diemenica, Spreng. Syst. ii. 724 = gracilis.
diffusa, Lej. Rev. Fl. Spa, 117 = arvensis.
discolor, Opiz, Seznam, 64, nomen (Quid?).—Europ.

MENTHA:—
dissimilis, *Déségl. in Bull. Soc. Sc. Angers* (1882), 201. —Europ.
divaricata, Host, Fl. Austr. ii. 150 = arvensis.
divaricata, Lag. ex Benth. Lab. Gen. et Sp. 175 = suavis.
diversifolia, Dum. Fl. Belg. 49 (Quid?).—Europ.
domestica, Tausch, ex Nym. Consp. 595 = sylvestris.
Dossiniana, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 323.—Belgium.
dubia, Chaix, ex Vill. Hist. Pl. Dauph. ii. 358 = aquatica.
dubia, Schleich. ex Suter, Fl. Helv. ii. 10 = arvensis.
dulcissima, Dum. Fl. Belg. 48 = sylvestris.
dumetorum, Schult. Obs. Bot. 108 = aquatica, sylvestris.
Dumortieri, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 334.—Europ.
egerensis, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 213.—Europ.
Ehrhartiana, Lej. & Court. Comp. Fl. Belg. ii. 232 = arvensis.
Eisenstadiana, Opiz, ex Tenore, Syll. Fl. Neap. 281 = sylvestris.
Eisensteiniana, Opiz, Natural. xi. (1826) 301 = sylvestris.
elata, Host, Fl. Austr. ii. 145 = arvensis.
elegans, Lej. Rev. Fl. Spa, 117 = arvensis.
elliptica, Lej. l. c. = arvensis.
elongata, Tenore, ex Steud. Nom. ed. II. ii. 126 = rotundifolia.
emarginata, Reichb. Fl. Germ. Excurs. 309 = sylvestris.
eriantha, C. Koch, in Linnaea, xxi. (1848) 650 = aquatica.
exigua, Linn. Cent. Pl. ii. 20; Amoen. Acad. iv. 318 = M. Pulegium.
exigua, Lucé, Fl. Osil. 205 = arvensis.
fallax, *Déségl. in Ann. Soc. Bot. Lyon*, viii. (1881) 224. —Europ.
farinosa, Rafin. Fl. Ludov. 42 (Quid?).—Am. bor.
Favrati, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 331.—Europ.
Fieberiana, Opiz, ex Steud. Nom. ed. II. ii. 126, nomen (Quid?).—Europ.
filina, Wallr. Beitr. Fl. Hercyn. 225; et in Linnaea, xiv. (1840) 595 = sylvestris.
flagellifera, Schur, Enum. Pl. Transs. 518 = arvensis.
foetens, Wall. ex Benth. Lab. Gen. et Sp. 162 = Elsholtzia incisa.
foetida, Burm. f. Fl. Ind. 126 = Dysophylla auricularia.
foliiconia, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 210.—Europ.
foliosa, Opiz, Natural. xi. (1826) 437 = sylvestris.
fontana, Weihe, ex Strail, in Kicksia, n. 252; et Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 37 = arvensis.
fragrans, Presl, Delic. Prag. 78 = rotundifolia.
fruticosa, Hilsenb. & Boj. ex Benth. Lab. Gen. et Sp. 165 = Tetradenia fruticosa.
fruticosa, Roxb. ex D. Don, Prod. Fl. Nep. 115; et Fl. Ind. iii. 6 = Pogostemon plectranthoides.
galeobdolifolia, Opiz, Boehm. Gew. (1823) 72, nomen (Quid?).—Europ.
galeopsifolia, *Opiz, ex Déségl. in Bull. Sc. Angers* (1882), 206.—Europ.
genevensis, *Déségl. & Durand, in Bull. Soc. Belg.* xvii. (1878) 315.—Europ.
Genevierii, *Déségl. & Durand, l. c.* 316.—Gallia.
gentilis, *Forsk. Fl. Aegypt. Arab.* 213 (Quid?).—Arab.
gentilis, Georgi, Beschr. Russ. Reich. III. v. 1082 = arvensis.
gentilis, *Linn. Sp. Pl.* 577.—Europ.
gentilis, Sm. in Trans. Linn. Soc. v. (1800) 208 = sativa.
germanica, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 312.—Europ.
gibraltarica, Willd. Enum. Hort. Berol. 611 = M. Pulegium.
Gillotii, Déségl. & Durand, in Bull. Soc. Bot. Belg. xvii. (1878) 324 = rotundifolia.
Giusliana, *Opiz, ex Déségl. l. c.* xxi. II. (1882) 115.—Europ.
glabra, Colla, Herb. Pedem. t. 84; ex Benth. in DC. Prod. xii. 171 = aquatica.
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = viridis.
glabra, Sole, ex Benth. Lab. Gen. et Sp. 179 = arvensis.
glabrata, Vahl, Symb. Bot. iii. 75 = piperita.
glomerata, Stokes, Bot. Mat. Med. iii. 324 = aquatica.
glomerulata, Jan, ex Benth. Lab. Gen. et Sp. 715 = suavis.

MENTHA:—
gracilis, *R. Br. Prod.* 505.—Austral.
gracilis, Pérard, in Bull. Soc. Bot. Fr. xvii. (1870) 337 = rotundifolia.
gracilis, Sole, Menth. Brit. 37. t. 16 = arvensis.
grandiflora, *Benth. in Mitch. Journ. Trop. Austral.* 362.—Austral.
grandifolia, *Hegetschw. Fl. Schw.* 581.—Helvet.
grata, Host, Fl. Austr. ii. 152 = arvensis.
gratissima, [Web. in] Wigg. Prim. Fl. Holsat. 43 = sylvestris.
graveolens, Opiz, Seznam, 65 = arvensis.
graveolens, *Presl, Fl. Sic.* p. xxxvi.—Sicilia.
Hakka, Siebold, in Verh. Batav. Gen. xii. (1830) 32 = arvensis.
Halleri, Dierb. Fl. Heidelb. ii. 169 = rotundifolia.
Halleri, C. C. Gmel. Fl. Bad. ii. 594 = sylvestris.
hamadanensis, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 35.—Persia.
Haynaldiana ×, Borb. in Oestr. Bot. Zeitschr. xxx. (1880) 19.—Europ.
hercynica, Roehl. Deutschl. Fl. ed. II. 328; ex Koch, Syn. Fl. Germ. ed. I. 550 = viridis.
Hillebrandtii, Ortm. ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 29 = arvensis.
hircina, Hull, Brit. Fl. i. 127 = piperita.
hirsuta, Huds. Fl. Angl. ed. I. 223 = aquatica.
hirta, Cald. Fl. Fav. Tent. 178 = aquatica.
hirta, Willd. Enum. Hort. Berol. 608 = sylvestris.
hispidula, Bor. Fl. Centr. Fr. ed. III. ii. 507 = sylvestris.
hispidula, Poepp. ex Benth. in DC. Prod. xii. 169 = piperita.
Hoferiana, Opiz, ex Steud. Nom. ed. II. 126, nomen (Quid?).—Europ.
hortensis, Opiz, ex Fresen. in Syll. Ratisb. ii. (1828) 227 = viridis.
hortensis, Tausch, ex Koch, Syn. Fl. Germ. ed. II. 635 = gentilis.
Hostii, Bor. Fl. Centr. Fr. ed. III. ii. 512 = arvensis.
Huguenini, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 330.—Gallia.
hungarica, *Borb. in Oestr. Bot. Zeitschr.* xxx. (1880) 19. —Hungar.
hybrida, Aresch. Skånes Fl. ed. II.; e in Bot. Notiser (1881), 158 = aquatica.
hybrida, Schleich. Cat. Pl. Helv. ed. IV. 23 = sylvestris.
hyptiformis, Lam. Encyc. iv. 110 = Hyptis verticillata.
imbricata, Roxb. Hort. Beng. 44 (Elsholtziae sp. ?).—Ind. or.
inarimensis, Guss. Enum. Pl. Inar. 255 = viridis.
incana, Willd. Enum. Hort. Berol. 609 = sylvestris.
incisa, Humnicki, Cat. Pl. Luxueil. 48 = Lamium purpureum?
incisa, Wall. ex Benth. Lab. Gen. et Sp. 162 = Elsholtzia incisa.
insularis, Req. ex Gren. & Godr. Fl. Fr. ii. 649 = rotundifolia.
integra, Buch.-Ham. ex Benth. in Wall. Pl. As. Rar. i. 31 = Pogostemon parviflorum.
intermedia, Host, Fl. Austr. ii. 141 = aquatica.
intermedia, *Krock. Fl. Siles. Suppl.* ii. 143.—Europ.
intermedia, Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 20 = arvensis.
intermedia, Opiz, Natural. xi. (1826) 436 = dissimilis.
javanica, Blume, Bijdr. 826 = arvensis.
javanica, Span. in Linnaea, xv. (1841) 322 = Hyptis brevipes.
Juliaca, Benth. in DC. Prod. xii. 165, sphalm. = *niliaca*.
kahirina, Forsk. Fl. Aegypt. Arab. 213 = piperita.
Korteana, Boenn. ex Steud. Nom. ed. II. ii. 126 = sylvestris.
laciniosa, Schur, Enum. Pl. Transs. 515 = viridis.
laevigata, Willd. Enum. Hort. Berol. 609 = viridis.
laevis, Opiz, Nom. Bot. 73 (Quid?).—Europ.
Laggeri, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 322.—Europ.
Lamarckii, Tenore, Fl. Nap. Prod. App. V. 19 = sylvestris.
lamiifolia, Host, Fl. Austr. ii. 144 = arvensis.
lamiifolia, Tenore, Fl. Nap. v. 8 = suavis.
lanceolata, Benth. Lab. Gen. et Sp. 181 = arvensis.
lanceolata, Opiz, Boehm. Gew. (1823) 72, nomen (Quid?).—Europ.
Langii, Geig. ex T. Nees, Duesseld. Samml. Suppl. [ii. 25. t. 1] = piperita.
Langii, Geig. Pharm. Bot. i. 471 = suavis.
lapponica, Wahlenb. Fl. Lapp. 161. t. 10 = arvensis.

MENTHA:—
lata, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 211 = arvensis.
latifolia, Host, Fl. Austr. ii. 145 = arvensis.
latifolia, Nolte, ex Hornem. in Fl. Dan. t. 2057 = aquatica.
latissima, Schur, Enum. Pl. Transs. 519 = arvensis.
lavandulacea, Willd. Enum. Hort. Berol. 609 = sylvestris.
lavandulaefolia, Pers. Syn. ii. 119 = sylvestris.
laxa, Host, Fl. Austr. ii. 151 = arvensis.
laxiflora, *Benth. in DC. Prod.* xii. 174.—Austral.
Lejeuneana, Opiz, Nom. 69 = seq.?
Lejeunii, Opiz, ex Reichb. Fl. Germ. Excurs. 309 = viridis.
Lereschii, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 332.—Gallia.
libenensis, Opiz, in Lotos, iii. (1853) 208 (Quid?).—Europ.
Linnaei, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 311.—Europ.
Lloydii, Bor. Fl. Centr. Fr. ed. III. ii. 507 = aquatica.
longifolia, Host, Fl. Austr. ii. 144 = arvensis.
longifolia, Huds. Fl. Angl. ed. I. 221 = sylvestris.
longistachya ×, Timb.—Cf. Bull. Soc. Bot. Fr. xxv. (1878) Rev. Bibl. 109.
Lucandiana, *Pérard, in Bull. Soc. Bot. Fr.* xvii. (1870) 335.—Europ.
macrantha, [Fisch.] Hort. Gorenk. ed. II. 23, nomen = *micrantha*?
macrophylla, Opiz, Natural. ix. (1825) 131 (Quid?).—Europ.
macrostachya, Tenore, Prod. Fl. Nap. p. xxxiii = rotundifolia.
maculata, Host, Fl. Austr. ii. 143 = arvensis.
malabarica, Heyne, ex Benth. in Wall. Pl. As. Rar. i. 30 = Dysophylla stellata.
marisensis, *Simk. in Termész. Füzetek*, ix. (1885) 25.—Europ.
Marrubiastrum, F. Schultz, ex Nym. Consp. 597 = arvensis.
Masneriana, Opiz, Natural. n. ix. (1825) 131 (Quid?). —Europ.
Maximiliana, F. Schultz, in Flora, xxxvii. (1854) 472 = aquatica.
meduanensis, Déségl. & Durand, in Bull. Soc. Bot. Belg. xvii. (1878) 310 = rotundifolia.
melissaefolia, Host, Fl. Austr. ii. 144 = arvensis.
mexicana, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 189 = rotundifolia.
Michelii, Tenore, ex Reichb. Pl. Crit. x. 27 = viridis.
micrantha, Fisch ex Benth. in DC. Prod. xii. 175 = M. Pulegium.
microphylla, C. Koch, in Linnaea, xxi. (1848) 648 = sylvestris.
minor, Opiz, ex Déségl. in Ann. Soc. Bot. Lyon, viii. (1881) 225; et in Bull. Soc. Sc. Angers (1882), 218 = arvensis.
mollicoma, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 216 = sylvestris.
mollis, Benth. Lab. Gen. et Sp. 182 = Bystropogon mollis.
mollissima, Borckh. ex Gaertn. Mey. & Scherb. Fl. Wett. ii. 348 = sylvestris.
montana, Host, Fl. Austr. ii. 145 = arvensis.
montana, Lowe, ex Benth. in DC. Prod. xii. 175 = M. Pulegium.
monticola, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 329.—Europ.
Morenii, *Déségl. & Dur. l. c.* 318.—Europ.
mosana, Lej. & Court. Comp. Fl. Belg. ii. 232 = arvensis.
motolensis ×, Opiz, in Lotos, iii. (1853) 208.—Europ.
mucronulata, Opiz, Natural. n. ix. (1825) 131 = rotundifolia.
Muelleriana, Sch. Bip. in Flora, xxxvii. (1854) 543 = arvensis.
multifida, Stokes, Bot. Mat. Med. iii. 318 = Preslia cervina.
multiflora, Host, Fl. Austr. ii. 147 = arvensis.
myosuroides, Roth, Nov. Pl. Sp. 257 = Dysophylla myosuroides.
napolitana, Tenore, Fl. Nap. t. 242; ex Benth. in DC. Prod. xii. 169 = piperita.
Neesiana, Opiz, ex Steud. Nom. ed. II. ii. 126 = arvensis.
neglecta, Tenore, Cat. Pl. Hort. Neap. (1819) App. ii. 82; Fl. Nap. ii. 379. t. 157. f. 2 = rotundifolia.

MENTHA:—
nemorosa, Host, Fl. Austr. ii. 144 = arvensis.
nemorosa, Willd. Sp. Pl. iii. 75 = sylvestris.
nemorum, *Bor. Fl. Centr. Fr.* ed. III. ii. 511.—Europ.
nepetifolia, Lej. ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 15 = sylvestris.
nepetoides, Lej. Rev. Fl. Spa, 116 = sylvestris.
nigrescens, C. Koch, in Linnaea, xxi. (1848) 648 = sylvestris.
nigrescens, Weihe, ex Fresen. in Syll. Ratisb. ii. (1828) 226 = aquatica.
nigricans, Mill. Gard. Dict. ed. VIII. n. 12 = aquatica.
niliaca, Jacq. Hort. Vindob. iii. 46 = sylvestris.
niliacea, Vahl, Symb. Bot. iii. 75 = viridis.
nitida, Host, Fl. Austr. ii. 143 = arvensis.
nivea, *Hegetschw. Fl. Schw.* 580.—Europ.
nobilis, Weihe, ex Fingerh. in Flora, xviii. (1835) II. 406 = arvensis.
Noëana, Boiss. Fl. Orient. iv. 543 = sylvestris.
Nouletiana, Timb. in Bull. Soc. Bot. Fr. vii. (1860) 259 = viridis.
numidica, *Poir. Encyc. Suppl.* iii. 662.—Afr. bor.
nummularia, Schreb. in Schweigg. & Koerte, Fl. Erlang. ii. 7 = arvensis.
nusleensis, Opiz, in Lotos, iii. (1853) 207 (Quid?).—Europ.
oblonga, Opiz, in Flora, vii. (1824) II. 525 = sylvestris.
oblongifolia, Opiz, ex Steud. Nom. ed. II. ii. 127, nomen (Quid?).—Europ.
obtusata, Opiz, Natural. x. (1825) 194 = arvensis.
obtuseserrata, Opiz, ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 29 = aquatica.
obtusifolia, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 200 = aquatica.
ocymiodora, Opiz, Natural. iv. (1823) [21–22] = viridis.
ocymoides, Host, Fl. Austr. ii. 148 = arvensis.
ocymoides, Lam. Encyc. iv. 103 = Geniosporum prostratum.
odora, Salisb. Prod. 80 = piperita.
odorata, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 211 = arvensis.
odorata, Sole, Menth. Brit. 21. t. 9 = aquatica.
officinalis, Hull, Brit. Fl. i. 127 = p perita.
Opitziana, Opiz, Nom. 72 (Quid?).—Europ.
origanifolia, J. F. Gmel. Syst. 902 = Bystropogon origanifolius.
origanifolia, Host, Fl. Austr. ii. 142 = arvensis.
origanoides, Lej. ex Fingerh. in Flora, xviii. (1835) II. 393 = aquatica.
origanoides, Maxim. ex Trautv. in Act. Hort. Petrop. ix. (1884) 87 = dahurica?
origanoides, Tenore, Fl. Nap. Prod. App. v. 19; Ind. Sem. Hort. Neap. (1825) 11. n. 16 = aquatica.
Ortmanniana, Opiz, Natural. xi. (1826) 437 = aquatica.
ovalifolia, Opiz, l. c. viii. (1824) 70 = arvensis.
ovata, Cav. Ic. iv. 36 = Elsholtzia cristata.
ovata, Schur, Enum. Pl. Transs. 519 = arvensis.
palatina, F. Schultz, Arch. Fl. 315 = arvensis.
pallida, Nees, ex Steud. Nom. ed. I. 523 = sylvestris.
palmata, Fisch. ex Steud. l. c. ed. II. ii. 127, nomen (Quid?).—Hab.?
paludosa, Nees, ex Bluff. & Fingerh. Comp. Fl. Germ. ed. I. ii. 15 = arvensis.
paludosa, Sole, Menth. Brit. t. 22 = aquatica.
palustris, Mill. Gard. Dict. ed. VIII. n. 11 = aquatica.
palustris, Moench, Meth. 380 = sativa.
paniculata, Roxb. Hort. Beng. 44; Fl. Ind. iii. 4 = Elsholtzia incisa.
parietariaefolia, Steud. Nom. ed. II. ii. 127 = arvensis.
parviflora, Schultz, Prod. Fl. Starg. Suppl. i. 33 = arvensis.
parvifolia, Opiz, Natural. ix. (1825) 132 = arvensis.
parvifolia, *Schur, Enum. Pl. Transs.* 514.—Europ.
pascuicola, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 321.—Europ.
Patrini, Lepech. in Nov. Act. Acad. Petrop. i. (1783) 336. t. 8 = Elsholtzia cristata.
Pauliana, F. Schultz, in Pollichia, xi. (1853) 40 = arvensis.
peckaënsis, Opiz, Natural. x. (1825) 195 (Quid?).—Europ.
pedunculariis, Bor. Fl. Centr. Fr. ed. III. ii. 510 = aquatica.
pedunculata, Pers. Syn. ii. 119 = aquatica.
perilloides, Hort. ex [Spreng.] Bot. Gart. Halle, 48 = Elsholtzia cristata.

MENTHA:—
perilloides, Lam. Encyc. iv. 112 = Perilla ocimoides.
perilloides, Linn. Syst. ed. XII. 736 = Hyptis pectinata.
persica, Hort. ex Benth. Lab. Gen. et Sp. 170 = sylvestris.
pilosa, Spreng. ex Wallr. Ann. Bot. 75 = arvensis.
Pimentum, Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 13 = piperita.
piperata, Stokes, Bot. Mat. Med. iii. 315 = piperita.
Piperella, Opiz, ex Lej. & Court. Comp. Fl. Belg. ii. 226 = viridis.
piperita, *Linn. Sp. Pl.* 576.—Europ.; As. et Afr. bor.
plicata, Lej. & Court. Comp. Fl. Belg. 231, in syn. = sativa.
plicata, Nees, ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 22 = praec.?
plicata, Opiz, Natural. viii. (1824) 70 = arvensis.
plumosa, Linn. f. Suppl. 273 = Bystropogon origanifolius.
polymorpha, Host, Fl. Austr. ii. 152 = arvensis.
polystachya, Wall. ex Steud. Nom. ed. II. ii. 127 = Elsholtzia polystachya.
postelbergensis, Opiz, Seznam, 65 = gentilis.
prachinensis, Opiz, l. c. = sativa.
praecox, Sole, Menth. Brit. 31. t. 13 = arvensis.
pratensis, Sole, l. c. 39. t. 17 = gentilis.
praticola, Opiz, Seznam, 65 = arvensis.
premysliana, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 213.—Europ.
probabilis, Schur, Enum. Pl. Transs. 518 = aquatica.
procumbens, Thuill. Pl. Par. ed. II. 288 = arvensis.
prostrata, Host, Fl. Austr. ii. 151 = arvensis.
pubescens, Willd. Enum. Hort. Berol. 608 = sylvestris.
pudina, Buch.-Ham. ex Benth. in Wall. Pl. As. Rar. i. 29 = viridis.
pulchella, Host, Fl. Austr. ii. 148 = arvensis.
pulegioides, Dum. Fl. Belg. 49 = M. Pulegium.
Pulegium, *Linn. Sp. Pl.* 577.—Europ.; As. et Afr. bor.
Pulegium, Lucé, Fl. Osil. 24 = arvensis.
pumila, R. Grah. in Edinb. N. Phil. Journ. (Jan.-Mar. 1828) 393 = Dysophylla crassicaulis.
pumila, Host, Fl. Austr. ii. 150 = arvensis.
punctata, Moench, Meth. Suppl. 146 = Preslia cervina.
purpurascens, Host, ex Steud. Nom. ed. II. ii. 127 = reversa.
purpurea, Host, Fl. Austr. ii. 141 = aquatica.
pyramidalis, Tenore, Prod. Fl. Nap. 34 = aquatica.
quadrifolia, Buch.-Ham. ex Benth. in Wall. Pl. As. Rar. i. 30 = Dysophylla crassicaulis.
quadrifolia, D. Don, Prod. Fl. Nep. 113 = Dysophylla cruciata.
quadrifolia, Roxb. Hort. Beng. 44; Fl. Ind. iii. 4 = Dysophylla quadrifolia.
quaternifolia, Heyne, ex Roth, Nov. Pl. Sp. 256 = Dysophylla quadrifolia.
ranina, Opiz, in Lotos, iii. (1853) 207 = aquatica.
ratisbonensis, Opiz, in Flora, vii. (1824) II. 525 = sylvestris.
recta, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 331.—Europ.
reflexifolia, Opiz, Natural. viii. (1824) 71; Flora, vii. (1824) II. 525 = sylvestris.
Requieni, *Benth. Lab. Gen. et Sp.* 182.—Ins. Corsica.
resinosa, Opiz, Natural. x. (1825) 195 = gentilis.
reticulosa, *Hance, in Walp. Ann.* iii. 247.—Hongkong.
reversa, *Reichb. ex Steud. Nom.* ed. II. ii. 127.—Europ.
riparia, Lej. ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 17 = aquatica?
riparia, Schreb. in Schweigg. & Koerte, Fl. Erlang. ii. 6 = sativa.
Ripartii, Déségl. & Durand, in Bull. Soc. Bot. Belg. xvii. (1878) 313 = rotundifolia.
rivalis, Sole, Menth. Brit. 45. t. 20 = arvensis.
rivularis, *Hegetschw. Fl. Schw.* 582.—Helvet.
Rodriguezii, *Malinv. in Marès, Cat. Pl. Vasc. Balear.* 335, in nota.—Ins. Balear.
Rosani, Tenore, Fl. Nap. App. v. 18; et Syll. 81 = sylvestris.
Rothii, Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 18 = arvensis.
rotundata, Opiz, in Flora, vii. (1824) II. 525 = arvensis.
rotundifolia, *Huds. Fl. Angl.* ed. I. 221.—Europ.; As. et Afr. bor.
rotundifolia, Sole, Menth. Brit. 9. t. 4 = sylvestris.
Royleana, Wall. Cat. n. 1537; Benth. in Wall. Pl. As. Rar. i. 29 = sylvestris.

MENTHA:—
rubra, Huds. Fl. Angl. ed. II. 252; Sole, Menth. Brit. 41 = gentilis.
rubra, *Mill. Gard. Dict.* ed. VIII. n. 9.—Hab.?
rubra, Sm. in Trans. Linn. Soc. v. (1800) 205 = sativa.
rubricaulis, Opiz, in Lotos, iii. (1853) 208 (Quid?).—Europ.
rubrohirta, Lej. & Court. Comp. Fl. Belg. ii. 230 = arvensis.
Rudaeana, Opiz, Nom. 28 (Quid?).—Europ.
rugosa, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 638 = Dysophylla rugosa.
rugosa, Lam. Fl. Fr. ii. 420 = rotundifolia.
salebrosa, Bor. Fl. Centr. Fr. ed. III. ii. 511 = arvensis.
salicina, *Burch. ex Benth. Lab. Gen. et Sp.* 170.—Afr. austr.
sapida, Tausch, ex Reichb. Fl. Germ. Excurs. 310 = sylvestris.
sativa, *Linn. Sp. Pl.* ed. II. 805.—Europ.
sativa, Roxb. Hort. Beng. 44; Fl. Ind. iii. 6 = arvensis.
sativa, Sm. in Trans. Linn. Soc. v. (1800) 199 = aquatica.
satureioides, *R. Br. Prod.* 505.—Austral.
Schleicheri, Opiz, Nom. 30 (Quid?).—Europ.
Schultzii, Boutigny, ex F. Schultz, Arch. Fl. 283 = rotundifolia.
Scordiastrum, F. Schultz, ex Malinv. in Soc. Dauph. Ech. Pl. iv. (1877) 120 = arvensis.
scotophylla, *Timb. in Mém. Acad. Toul.* Sér. V. iv. (1860) 526.—Europ.
Scribae, F. Schultz, in Flora, lvi. (1873) 237 = arvensis.
scrophulariaefolia, Lej. & Court. Comp. Fl. Belg. ii. 230 = arvensis.
secunda, Roxb. Hort. Beng. 44 = Pogostemon plectranthoides.
segetalis, Opiz, in Lotos, iii. (1853) 211 (Quid?).—Europ.
Sehlmeyeri, Opiz, Nom. 63 (Quid?).—Europ.
semi-integra, *Opiz, ex Déségl. in Bull. Soc. Bot. Belg.* xxi. II. (1882) 109.—Europ.
seriata, *A. Kern. Sched. Fl. Exsicc.* ii. 121.—Dalmat.
serotina, Host, Fl. Austr. ii. 143 = arvensis.
serotina, Tenore, Prod. Fl. Nap. App. iv. 27 = sylvestris.
serpyllifolia, *Benth. in DC. Prod.* xii. 174.—Tasman.
serrata, Pérard, in Bull. Soc. Bot. Fr. xvii. (1870) 337 = rotundifolia.
serrulata, *Opiz, ex Déségl. in Bull. Soc. Bot. Belg.* xxi. II. (1882) 105.—Europ.
Sieberi, C. Koch, in Linnaea, xxi. (1848) 649 = sylvestris.
silesiaca, Opiz, Natural. ix. (1825) 132 (Quid?).—Europ.
similis, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 314.—Europ.
simplex, Host, Fl. Austr. ii. 148 = M. Pulegium.
Skofitziana ×, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 385.—Europ.
slichoviensis, Opiz, Seznam, 65 = arvensis.
Speckmoseriana, Opiz, Natural. x. (1825) 196 = arvensis.
spicata, Crantz, Stirp. Austr. ed. II. fasc. iv. 328 = viridis.
spicata, Linn. Sp. Pl. 576 = sylvestris.
Sprengeliana, Opiz, ex Steud. Nom. ed. II. ii. 127, nomen (Quid?).—Europ.
stachyoides, Host, Fl. Austr. ii. 146 = arvensis.
statenicensis, Opiz, in Lotos, iii. (1853) 211 = sativa.
stellata, Buch.-Ham. ex Roxb. Hort. Beng. [95]; Fl. Ind. iii. 5 = Dysophylla stellata.
stellata, Lour. Fl. Cochinch. ii. 361 = Dysophylla verticillata.
stolonifera, *Krock. Fl. Siles. Suppl.* ii. 146.—Europ.
stolonifera, Opiz, Seznam, 64 = aquatica.
Strailii, Durand, in Bull. Soc. Bot. Belg. xiv. (1875) 307 = aquatica.
stricta, Becker, ex Steud. Nom. ed. II. ii. 127 = aquatica.
stricta, Stokes, Bot. Mat. Med. iii. 322 = sativa.
strobilifera, Wall. ex Benth. in Wall. Pl. As. Rar. i 30 = Elsholtzia strobilifera.
stylosa, Opiz, in Lotos, iii. (1853) 184 (Quid?).—Europ.
suaveolens, Ehrh. Beitr. vii. 149 = rotundifolia.
suaveolens, Host, Fl. Austr. ii. 146 = arvensis.
suaveolens, Steph. ex Ledeb. Fl. Ross. iii. 340 = dahurica.

MENTHA:—
suavis, *Guss. Pl. Rar.* 387. t. 66.—Europ. austr.
suavis, Hoffm. ex Sm. in Rees, Cycl. xxiii. sub n. 6 = sylvestris.
suavissima, Lej. ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 5 = sylvestris.
subcordata, Colla, ex Lamotte, Prod. Fl. Centr. Fr. 593 = arvensis.
subglabra, Opiz, in Lotos, iii. (1853) 209 (Quid?).—Europ.
subinodora, Schur, Enum. Pl. Transs. 518 = arvensis.
subspicata, Weihe, ex Fresen. in Syll. Ratisb. ii. (1828) 226 = aquatica, arvensis.
sudetica, *Opiz, ex Déségl. in Bull. Soc. Sc. Angers* (1882), 207.—Europ.
sylvatica, Host, Fl. Austr. ii. 149 = arvensis.
sylvestris, *Linn. Sp. Pl.* ed. II. 804.—Reg. bor. temp.
sylvestris, Sole, Menth. Brit. 7. t. 3 = rotundifolia.
Tauschiana, Opiz, Boehm. Gew. (1823) 71, nomen (Quid?).—Europ.
Tenorii, Pérard, in Bull. Soc. Bot. Fr. xvii. (1870) 335 = sylvestris.
tenuiflora, Opiz, Natural. xi. (1826) 438 = viridis?
tenuifolia, Host, Fl. Austr. ii. 147 = arvensis.
tenuifolia, Opiz, ex Reichb. Fl. Germ. Excurs. 309 = viridis.
tenuis, Frank, ex Benth. in DC. Prod. xii. 169 = piperita.
tenuis, Michx. Fl. Bor. Am. ii. 2 = viridis.
terebinthinacea, Willd. ex Steud. Nom. ed. II. ii. 128 = canadensis.
Thurmannii, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 320.—Europ.
Timija, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 65, nomen.—Marocc.
Tinantiana, Lej. ex Malinv. in Bull. Soc. Linn. Normand. Sér. III. iii. (1879) 29 = aquatica?
Todari, *Lojac. in Natural. Sicil.* iii. (1884) 46.—Sicil.
tomentella, Hoffmgg. & Link, Fl. Port. i. 73 = M. Pulegium.
tomentosa, Sm. in Rees, Cyclop. xxiii. n. 23 = M. Pulegium.
tomentosa, Urv. in Mém. Soc. Linn. Par. i. (1822) 323 = sylvestris.
tortuosa, Host, Fl. Austr. ii. 142 = arvensis.
transmota, *Déségl. & Durand, in Bull. Soc. Bot. Belg.* xvii. (1878) 319.—Europ.
transsilvanica, Schur, Enum. Pl. Transs. 513 = sylvestris.
tristis, Opiz, Natural. x. (1825) 196 (Quid?).—Europ.
uliginosa, Salisb. Prod. 80 = sylvestris.
umbrosa, Opiz, Boehm. Gew. (1823) 72; Weihe, ex Fingerh. in Flora, xviii. (1835) 393 = aquatica.
undulata, Hort. Canal. ex Opiz, Natural. xi. (1826) 437 (Quid?).—Europ.
undulata, Willd. Enum. Hort. Berol. 608 = sylvestris.
urticaefolia, Tenore, Fl. Nap. t. 157; Ind. Sem. Hort. Neap. (1825) 11. n. 17 = aquatica.
varians, Host, Fl. Austr. ii. 150 = arvensis.
variegata, Sole, Menth. Brit. 43. t. 19 = arvensis.
vegeta, Baker, in Journ. Bot. iii. (1865) 245 = cardiaca.
velutina, Lej. Rev. Fl. Spa, 115; Bab. Man. Brit. Bot. ed. VI. 263, in syn. = sylvestris.
velutina, Wall. ex Steud. Nom. ed. II. ii. 128 = Dysophylla verticillata.
venulosa, Heyne, ex Wall. Cat. sub n. 1547 = Dysophylla myosuroides.
veronicaefolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 114 = Dysophylla crassicaulis.
veronicaeformis, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 217 = sylvestris.
verticillata, D. Don, Prod. Fl. Nep. 114 = Dysophylla crassicaulis.
verticillata, Gaertn. Mey. & Scherb. Fl. Wett. ii. 352 = sativa.
verticillata, Linn. Syst. ed. X. 1099 = arvensis.
verticillata, Roxb. Hort. Beng. 44; Fl. Ind. iii. 5 = Dysophylla verticillata.
viennensis, Opiz, Seznam, 64 = aquatica.
villosa, Becker, Fl. Frankf. i. 223 = arvensis.
villosa, Heyne, ex Benth. in Wall. Pl. As. Rar. i. 31 = Pogostemon rotundatum.
villosa, Huds. Fl. Angl. ed. II. 250 = sylvestris.
villoso-nervata, Opiz, Nom. 60 (Quid?).—Europ.

MENTHA:—
virgata, Salisb. ex Steud. Nom. ed. I. 523, 524 = arvensis.
viridis, *Linn. Pl.* ed. II. 804.—Europ.; As. et Afr. bor.
viridula, Host, Fl. Austr. ii. 147 = arvensis.
Wagneriana, Opiz, Natural. x. (1825) 196 (Quid?).—Europ.
Walteriana, Opiz, l. c. ix. (1825) 132 (Quid?).—Europ.
Weidenhofferi, Opiz, Nom. 32 = aquatica.
Weiheana, Opiz, Natural. viii. (1824) 71 = aquatica.
Weinerniana, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 216 = sylvestris.
weissenburgensis, Schultz, ex Nym. Consp. 596 = aquatica.
Weitenweberi, Opiz, ex Steud. Nom. ed. II. ii. 128; et ex Déségl. in Ann. Soc. Bot. Lyon, viii. (1881) 228 (Quid?).—Europ.
Welwitschii, *Rouy.—Cf. Bull. Soc. Bot. Fr.* xxx. (1883) *Bibl.* 32.—Europ.
Wierzbickiana, Opiz, in Flora, vii. (1824) II. 525 = sylvestris.
Wigmanniana, Opiz, ex Steud. Nom. ed. II. ii. 128, nomen (Quid?).—Europ.
Willdenowii, Déségl. & Durand, in Bull. Soc. Bot. Belg. xvii. (1878) 317 = rotundifolia.
Winkleriana, Opiz, ex Steud. Nom. ed. II. ii. 128; et ex Déségl. in Ann. Soc. Bot. Lyon, viii. (1881) 228 (Quid?).—Europ.
Wirtgeniana, F. Schultz, in Pollichia, xii. (1854) 41 = sativa.
Wohlwerthiana, F. Schultz, l. c. 37 = arvensis.
Wondračekii, Opiz, ex Déségl. in Bull. Soc. Sc. Angers (1882), 217 = sylvestris.
zatecensis, *Opiz, l. c.* 211.—Europ.
zeylanica, Burm. ex Steud. Nom. ed. II. ii. 128 = Geniosporum prostratum.

MENTHELLA, Pérard, in Bull. Soc. Bot. Fr. xvii. (1870) 205 = **Mentha**, Tourn. (Labiat.).
Requienii, Pérard, l. c. 206 = Mentha Requieni.

MENTZELIA, Plum. ex Linn. Gen. ed. I. 338 (1737).
LOASEAE, Benth. & Hook. f. i. 804.
ACROLASIA, Presl, Rel. Haenk. ii. 39. t. 55 (1831).
BARTONIA, Sims, Bot. Mag. t. 1487 (1804).
CHRYSOSTOMA, Lilja, in Linnaea, xv. (1841) 263.
CREOLOBUS, Lilja, Flora Oefver Sver. Odl. Vext. 67 (1839).
EUCNIDE, Zucc. Del. Sem. Monac.—Cf. Linnaea, xviii. (1844) 508; et in Abh. Akad. Muench. iv. 1. (1845) t. 1.
MENZELIA, Linn. Hort. Cliff. 492 (1737).
MICROSPERMA, Hook. Ic. Pl. t. 234 (1839).
TRACHYPHYTUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 533 (1840).
adhaerens, *Benth. Bot. Voy. Sulph.* 15.—Calif.
albicaulis, *Dougl. ex Hook. Fl. Bor. Am.* i. 222.—Am. bor. occ.
albicaulis, A. Gray, Pl. Wright. i. 74 = dispersa.
albicaulis, Torr. Bot. Mex. Bound. 67, partim = aspera.
aspera, Cav. Ic. i. 51. t. 70 = hispida.
aspera, *Linn. Sp. Pl.* 516.—Am. bor. occ.
aurea, Nutt. Gen. Am. i. 300 = oligosperma.
Bartonia, Steud. Nom. ed. II. ii. 128 = Lindleyi.
Brandegei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 367.—Am. bor. occ.
chilensis, *C. Gay, Fl. Chil.* ii. 431.—Chili.
chrysantha, Engelm. in Brandeg. Fl. S. W. Color. 237 = pumila.
congesta, *Torr. & Gray, Fl. N. Am.* i. 534.—Am. bor. occ.
cordata, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 33.—Calif.
crocea, Kellogg, l. c. vii. (1876) 110 = Lindleyi.
dispersa, *S. Wats. in Proc. Am. Acad.* xi. (1876) 115, 137.—Am. bor. occ.
floridana, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 532.—Am. bor.
gracilenta, *Torr. & Gray, l. c.* 534.—Am. bor. occ.
grandiflora, *Ruiz & Pav. ex G. Don, Gen. Syst.* iii. 66.—Peruv.
gronoviaefolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* x. 54.—Mexic.
Haenkii, *A. Gray, Pl. Wright.* i. 74, in nota.—Chili.
hirsutissima, *S. Wats. in Proc. Am. Acad.* xii. (1877) 252.—Calif.

MENTZELIA:—
hirta, *Pav. ex G. Don, Gen. Syst.* iii. 66.—Mexic.
hispida, *Willd. Sp. Pl.* ii. 1176.—Mexic.
involucrata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 367.—Calif.
laevicaulis, *Torr. & Gray, Fl. N. Am.* i. 535.—Am. bor. occ.
Lindleyi, *Torr. & Gray, l. c.* 533.—Calif.
lobata, *Walp. Rep.* ii. 224; *Hemsl. Biol. Centr. Am. Bot.* i. 472.—Mexic.
micrantha, *Torr. & Gray, Fl. N. Am.* i. 535.—Calif.
multiflora, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 48.—Am. bor.
nuda, *Torr. & Gray, Fl. N. Am.* i. 535.—Am. bor.
oligosperma, *Nutt. ex Sims, Bot. Mag.* t. 1760.—Louisiana.
ornata, *Torr. & Gray, Fl. N. Am.* i. 534.—Am. bor.
ornata, Torr. in Stansb. Exped. Utah, 387 = laevicaulis.
pectinata, Kellogg, in Proc. Calif. Acad. iii. (1863) 40 = gracilenta.
pedicellata, Presl, Epim. Bot. 246 = stipitata.
petiolata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 455 = oligosperma.
pumila, *Torr. & Gray, Fl. N. Am.* i. 535.—Am. bor. occ.
rhombifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 533 = oligosperma.
scabra, *H. B. & K. Nov. Gen. et Sp.* vi. 120.—N. Granat.
stipitata, [Moç. & Sesse,] ex DC. Prod. iii. 343 = hispida.
stipitata, *Presl, Rel. Haenk.* ii. 42.—Mexic.
strigosa, *H. B. & K. Nov. Gen. et Sp.* vi. 120.—Mexic.
Torreyi, *A. Gray, in Proc. Am. Acad.* x. (1874) 72.—Am. bor. occ.
tricuspis, *A. Gray, in Am. Natural.* ix. (1875) 271.—Calif.
urens, *Parry, ex A. Gray, in Proc. Am. Acad.* x. (1874) 71.—Utah.
urens, *Vell. Fl. Flum.* v. t. 97.—Bras.
Veatchiana, Kellogg, in Proc. Calif. Acad. ii. (1863) 99 = albicaulis.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 48.—Texas.
Wrightii, Porter, Fl. Color. 47 = pumila.

MENYANTES, Zumag. Fl. Pedem. i. 275 (1849) = seq.

MENYANTHES, [Tourn.] Linn. Syst. ed. I (1735).
GENTIANEAE, Benth. & Hook. f. ii. 819.
americana, Sweet, Hort. Brit. ed. I. 282 = trifoliata.
biflora, Moon, Cat. Pl. Ceyl. 13 = Limnanthemum aurantiacum.
brachysperma, Griseb. in DC. Prod. ix. 137, sphalm. = Limnanthemum trachyspermum.
brasilica, Vell. Fl. Flum. 75; ii. t. 70 = Limnanthemum Humboldtianum.
campestris, Macrae & Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 132, partim = Limnanthemum Forbesianum.
campestris, Moon, Cat. Pl. Ceyl. 13 = Limnanthemum parvifolium.
capensis, Thunb. Prod. Pl. Cap. 34 = Villarsia ovata.
Crista-galli, *Menz. ex Hook. Bot. Misc.* i. (1830) 45. t. 34.—Am. bor. occ.
cristata, Roxb. Pl. Corom. ii. 3. t. 105 = Limnanthemum cristatum.
exaltata, Sims, Bot. Mag. t. 1029 = Villarsia parnassifolia.
Hydrophyllum, Lour. Fl. Cochinch. 105 = Limnanthemum Hydrophyllum.
indica, Aubl. Pl. Gui. i. 118 = Limnanthemum Humboldtianum.
indica, Bory, Voy. i. 236 = Limnanthemum cristatum.
indica, Linn. Sp. Pl. 145 = Limnanthemum indicum.
latifolia, Rafin. New Fl. Am. ii. 20 = trifoliata?
macrophylla, Roth, ex Roem. & Schult. Syst. iv. 180 = Limnanthemum Kleinianum.
meridionalis, Willd. ex Griseb. Gen. et Sp. Gent. 347 = Limanthemum Humboldtianum.
natans, Lam. Fl. Fr. ii. 203 = Limnanthemum nymphoides.
nymphoides, Linn. Sp. Pl. 145 = Limnanthemum nymphoides.
nymphoides, Thunb. Fl. Jap. 82 = Brasenia Schreberi.

MENYANTHES:—
ovata, Linn. f. Suppl. 133 = Villarsia ovata.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 340 = trifoliata.
paradoxa, Fries, Nov. Fl. Suec. ed. II. 71 = trifoliata.
peltata, Thunb. in Nov. Act. Soc. Sc. Upsal. vii. (1815) 142 = Brasenia Schreberi.
petioliflora, Stokes, Bot. Mat. Med. i. 299 = Limnanthemum indicum.
pumila, Dougl. ex Griseb. Gen. et Sp. Gent. 338 = Hesperochiron californicus.
punctata, Muhl. ex Griseb. l. c. 348 = Limnanthemum lacunosum.
sarmentosa, Sims, Bot. Mag. t. 1328 = Villarsia reniformis.
sinarica, Buch.-Ham. ex Steud. Nom. ed. II. ii. 128 = Limnanthemum cristatum.
trachyspermà, Michx. Fl. Bor. Am. i. 126 = Limnanthemum trachyspermum.
tridentata, Rafin. New Fl. Am. ii. 20 = trifoliata?
trifoliata, *Linn. Sp. Pl.* 145.—Reg. bor. temp.
Trifolium, Neck. Delic. Gallo-Belg. i. 105 = trifoliata.
verna, Rafin. Med. Fl. ii. 33 = trifoliata.

MENZELIA, Linn. Hort. Cliff. 492 (1737) = **Mentzelia**, Plum. (Loasac.).

MENZIESIA, Sm. Pl. Ic. Ined. t. 56 (1791). *ERICACEAE*, Benth. & Hook. f. ii. 602.
CANDOLLEA, Baumg. Cat. Hort. Bollw. (1810).
aleutica, Spreng. Syst. ii. 202 = Bryanthus aleuticus.
Bruckenthalii, Baumg. Enum. Stirp. Transs. i. 333 = Bruckenthalia spiculifolia.
bryantha, Sw. in Trans. Linn. Soc. x. (1811) 377. t. 30. f. b = Bryanthus Gmelini.
caerulea, Sw. l. c.; Wahlenb. Fl. Lappon. 105 = Bryanthus taxifolius.
ciliicalyx, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. ix. 10.—Japon.
Daboëci, DC. Fl. Fr. iii. 674 = Daboëcia polifolia.
empetriformis, Pursh, Fl. Am. Sept. i. 264 = Cassiope Stelleriana.
empetriformis, Sw. in Trans. Linn. Soc. x. (1811) 380 = Bryanthus empetriformis.
ferruginea, A. Gray, in Proc. Am. Acad. viii. (1872) 393 = glabella.
ferruginea, *Sm. Pl. Ic. Ined.* t. 56.—Am. bor. occ.
glabella, *A. Gray, Syn. Fl. N. Am.* ii. 1. 39.—Am. bor. occ.
glanduliflora, Hook. Fl. Bor. Am. ii. 40 = Bryanthus glanduliflorus.
globularis, Hook. l. c. 41 = glabella.
globularis, *Salisb. Parad. Lond.* t. 44.—Am. bor.
Grahamii, Hook. Fl. Bor. Am. ii. 40 = Bryanthus empetriformis.
intermedia, Hook. l. c. = Bryanthus empetriformis.
minor, Hoffmgg. Verz. Pfl. 79 = Daboëcia polifolia.
multiflora, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. ix. 11.—Japon.
pentandra, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 431.—Japon.
phylicifolia, Fisch. ex Hook. Fl. Bor. Am. ii. 40 = Bryanthus aleuticus.
pilosa, Juss. in Ann. Mus. Par. i. (1802) 56 = globularis.
polifolia, Juss. l. c. 55. t. 4. f. 1 = Daboëcia polifolia.
purpurea, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 431.—Japon.
Smithii, Michx. Fl. Bor. Am. i. 235 = globularis.
Stelleriana, Fisch. ex Hook. Fl. Bor. Am. ii. 37 = Cassiope Stelleriana.
triflora, Bert. ex Spreng. Syst. ii. 202 = Lyonia octandra.
urceolaris, Salisb. Parad. Lond. sub t. 44 = ferruginea.

MEON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = **Meum**, Tourn. (Umbell.).
mutellinum, St. Lag. l. c. = Meum Mutellina.

MEOSCHIUM, Beauv. Agrost. 111. t. 21 (1812) = **Ischaemum**, Linn. (Gramin.).
aristatum, Beauv. Agrost. 111 = I. aristatum.
Arnottianum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 198 = I. rugosum.
australe, Beauv. ex Roem. & Schult. Syst. ii. 790 = I. australe.
barbatum, Beauv. Agrost. 111 = I. aristatum.
ciliare, Beauv. l. c. = I. ciliare.
elatum, Nees, ex Steud. Syn. Pl. Gram. 376 = I. aristatum.

MEOSCHIUM:—
elegans, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 199 = Ischaemum rugosum.
Griffithii, Nees & Arn. in Ann. Nat. Hist. Ser. I. i. (1838) 284 = I. rugosum.
lodiculare, Nees, in Hook. & Arn. Bot. Beech. Voy. 246 = I. aristatum.
Meyenianum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 197 = I. aristatum.
monostachys, Wight & Arn. ex Steud. Syn. Pl. Gram. 369 = I. aristatum.
Neesianum, Arn. ex Steud. l. c. 375 = I. aristatum.
Royleanum, Nees, ex Steud. l. c. = I. rugosum.
rugosum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 200 = I. rugosum.
semisagittatum, Schult. Mant. ii. 435 = I. semisagittatum.
triticeum, Beauv. ex Roem. & Schult. Syst. ii. 790 = I. triticeum.
villosum, Beauv. l. c. = I. australe.
Wightianum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 199 = I. rugosum.

MEPHITIDIA, Reinw. ex Blume, Cat. Gew. Buitenz. 51 (1823) = **Lasianthus**, Jack (Rubiac.).
acuminata, Walp. Ann. ii. 762 = L. acuminatus.
apiocarpa, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 221 = L. apiocarpus.
attenuata, DC. Prod. iv. 452 = L. attenuatus.
Blumeana, Walp. Ann. ii. 761 = L. Blumeanus.
bracteata, Walp. l. c. 759 = L. cyanocarpus.
capitata, DC. Prod. iv. 453 = L. capitatus.
capitulata, Walp. Ann. ii. 762 = L. capitulatus.
chinensis, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 196 = L. chinensis.
chrysoneura, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 222 = L. chrysoneurus.
ciliata, Walp. Ann. ii. 761 = L. ciliatus.
constricta, Walp. l. c. 762 = L. constrictus.
cyanocarpa, DC. Prod. iv. 452 = L. cyanocarpus.
dichotoma, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 224.—Java.
dichotoma, Walp. Ann. ii. 761 = L. dichotomus.
dubia, Walp. l. c. 763 = Psychotria Wightiana.
elliptica, Walp. l. c. 761 = L. ellipticus.
foetens, Walp. l. c. 763 = Saprosma ceylanicum.
Gardneri, Thw. Enum. Pl. Zeyl. 145 = L. Gardneri.
Griffithii, Walp. Ann. ii. 760 = L. Griffithii.
hexandra, Blume, Cat. Gew. Buitenz. 51 = L. hexandrus.
inaequalis, DC. Prod. iv. 453 = L. cyanocarpus.
inaequalis, Zoll. ex Miq. Fl. Ind. Bat. ii. 321 = L. densifolius.
inodora, DC. Prod. iv. 453 = L. inodorus.
Jackiana, Walp. Ann. ii. 760 = L. Jackianus.
laevigata, DC. Prod. iv. 453 = L. laevigatus.
lanceolata, Hassk. Cat. Hort. Bog. Alt. 111.—Hab.?
latifolia, Blume, ex DC. Prod. iv. 453 = L. latifolius.
longifolia, Walp. Ann. ii. 762 = L. longifolius.
lucida, DC. Prod. iv. 453 = L. lucidus.
Moonii, Walp. Ann. ii. 760 = L. Moonii.
mucronulata, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 223 = L. mucronulatus.
obscura, DC. Prod. iv. 453 = L. obscurus.
obscura, Zoll. ex Miq. Fl. Ind. Bat. ii. 318 = L. tomentosus.
oligantha, Thw. Enum. Pl. Zeyl. 146 = L. oliganthus.
ovata, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 221 = L. ovatus?
parvifolia, Walp. Ann. ii. 762 = L. parvifolius.
pauciflora, Walp. l. c. 763 = L. pauciflorus.
pilosa, Walp. l. c. 760 = L. pilosus.
platyphylla, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 222 = L. platyphyllus.
protracta, Thw. Enum. Pl. Zeyl. 146 = L. strigosus.
pterosperma, Walp. Ann. ii. 762 = L. pterospermus.
purpurea, DC. Prod. iv. 454 = L. purpureus.
reticulata, DC. l. c. = L. reticulatus.
retosa, Walp. Ann. ii. 760 = L. retosus.
rhinocerotis, Hort. Bogor. ex Miq. Fl. Ind. Bat. ii. 317 = L. obscurus.
rhinozerotis, DC. Prod. iv. 453 = L. rhinozerotis.
rhinozerotis, Kurz, in. Andam. For. Rep. Append. A. 40 = L. cyanocarpus.
rhinophylla, Thw. Enum. Pl. Zeyl. 145 = L. rhinophyllus.
rostrata, Walp. Ann. ii. 761 = L. ciliatus.
Roxburghii, Wight & Arn. Prod. 390 = L. cyanocarpus.
rufa, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 224 = L. rufus.

MEPHITIDIA:—
stercoraria, DC. Prod. iv. 454 = Lasianthus stercorarius.
stipularis, DC. l. c. 453 = L. stipularis.
strigosa, Walp. Ann. ii. 762 = L. strigosus.
sylvestris, DC. Prod. iv. 454 = L. sylvestris.
tomentosa, DC. l. c. 453 = L. tomentosus.
tomentosa, Thw. Enum. Pl. Zeyl. 145 = L. Thwaitesii.
tubiflora, DC. Prod. iv. 454 = Hamiltonia suaveolens.
varians, Thw. Enum. Pl. Zeyl. 146 = L. Walkerianus, strigosus, varians.
venosa, DC. Prod. iv. 453 = L. venosus.
venosa, Zoll. ex Miq. Fl. Ind. Bat. ii. 325 = L. reticulatus.
venulosa, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 508, in syn. = L. venulosus.
Walkeriana, Walp. Ann. ii. 762 = L. Walkerianus.
Wallichii, Wight & Arn. Prod. 390 = L. Wallichii.

MERATHREPTA, Rafin. in Ser. Bull. Bot. i. (1830) 221 = **Danthonia**, DC. (Gramin.).
spicata, Rafin. l. c. = D. spicata.

MERATIA, Cass. in Dict. Sc. Nat. xxx. 65 (1824) = **Elvira**, Cass. (Compos.).
Sprengelii, Cass. l. c. 66 = E. Martyni.

MERATIA, A. DC. in DC. Prod. x. 104 (1846) = **Moritzia**, DC. (Borag.).
Lindenii, A. DC. l. c.—Venezuela.

MERATIA, Loisel. Herb. Gen. Amat. iii. t. 173 (1819) = **Chimonanthus**, Lindl. (Calycanth.).
fragrans, Loisel. l. c. = C. fragrans.

MERCADOA, Naves, in Blanco, Fl. Philipp. ed. III. t. 463 (1880?) = **Sumbavia**, Baill. (Euphorb.).
mandalojonensis, Naves, l. c. = S. rottleroides.

MERCIERA, A. DC. Monog. Camp. 369. t. 5 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 560.
brevifolia, *A. DC. Monog. Camp.* 371.—Afr. austr.
Eckloniana, Buek, in Eckl. & Zeyh. Enum. 387 = tenuifolia.
heteromorpha, Buek, l. c. (Rubiacea).—Afr. austr.
heterophylla, Steud. Nom. ed. II. ii. 129, sphalm. = praec.
leptoloba, *A. DC. Monog. Camp.* 371. t. 5.—Afr. austr.
tenuifolia, *A. DC. l. c.* 370.—Afr. austr.

MERCKIA, Fisch. ex Cham. & Schlecht. in Linnaea, i. (1826) 59 = **Arenaria**, Rupp. (Caryoph.).
peploides, G. Don, Gen. Syst. i. 441 = A. peploides.
physodes, Fisch. ex Cham. & Schlecht. in Linnaea, i. (1826) 59 = A. physodes.

MERCKLINIA, Regel, Ind. Sem. Hort. Petrop. (1856) 25; et Gartenfl. (1857) 155 = **Hakea**, Schrad. (Proteac.).
petrophiloides, Regel, Gartenfl. (1857) 156 = H. lissocarpa.
rosea, Regel, Ind. Sem. Hort. Petrop. (1856) 25; Gartenfl. (1857) 155 = H. lissocarpa.

MERCURIALIS, [Tourn.] Linn. Syst. ed. I (1735). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 309.
SYNEMA, Dulac, Fl. Hautes-Pyr. 154 (1867).
afra, Linn. Mant. ii. 298 = Hydrocotyle villosa.
alpina, Schur, Enum. Pl. Transs. 600 = perennis.
alternifolia, Hochst. ex Baill. Etud. Gén. Euph. 490 = Acalypha Hochstetteriana.
alternifolia, Lam. Encyc. iv. 120 = Claoxylon Mercurialis.
ambigua, Linn. Sp. Pl. ed. II. 1465 = annua.
androgyna, Steud. Nom. ed. I. 524 = Leidesia capensis.
angustifolia, Baill. Adansonia, vi. (1866) 322 = Claoxylon angustifolium.
annua, *Linn. Sp. Pl.* 1035.—Europ.; Afr. bor.
annua, Thunb. Fl. Cap. ed. Schult. 387 = Leidesia capensis.
annua, Uspenski, ex Ledeb. Fl. Ross. iii. 580 = perennis.
australis, Baill. Adansonia, vi. (1866) 322 = Claoxylon australe.
bupleuroides, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 557 = Adenocline pauciflora.
caffra, Meissn. in Hook. l. c. 558 = Adenocline acuta.
capensis, Sond. in Linnaea, xxiii. (1850) 112 = Leidesia capensis.

MERCURIALIS :—
capensis, Spreng. ex Sond. in Linnaea, xxiii. (1850) 111 = Leidesia obtusa.
ciliata, Presl, Delic. Prag. 56 = annua.
corsica, *Coss. & Kral. Notes Crit.* 63.—Ins. Corsica.
Cynocrambe, Scop. Fl. Carn. ed. II. ii. 266 = perennis.
Dregeana, Buchinger, ex Krauss, in Flora, xxviii. (1845) 84 = Adenocline acuta.
elliptica, Duby, Bot. Gall. ed. II. i. 417 = corsica.
elliptica, *Lam. Encyc.* iv. 119.—Reg. Mediterr. occ.
Huetii, Hanry, in Billotia, i. (1864) 21 = annua.
indica, Lour. Fl. Cochinch. 628 (Claoxyli sp.?).—Cochinchin.
Ladanum, Hartm. in Anderss. Bot. Notiser (1849), 67 = annua.
leiocarpa, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* 37.—Japon.
livida, Portenschl. ex Baumg. Enum. Stirp. Transs. iii. 344 = perennis.
longifolia, Host, Fl. Austr. ii. 666 = perennis.
longifolia, Lam. Encyc. iv. 119 = annua.
nemoralis, Salisb. Prod. 390 = perennis.
ovata, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. (1815) 170. t. 4 = perennis.
pauciflora, Baill. Adansonia, iii. (1862–63) 159 = Adenocline pauciflora.
perennis, *Linn. Sp. Pl.* 1037.—Europ.; As. occ.
procumbens, Linn. l. c. 1036 (= *Paradenocline procumbens*).—Afr. austr.
pumila, Sond. in Linnaea, xxiii. (1850) 112 = Tragia triandra.
sericea, Salisb. Prod. 390 = tomentosa.
serrata, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 557 = Adenocline pauciflora.
subcordata, Buchinger, ex Krauss, in Flora, xxviii. (1845) 84 = Adenocline acuta.
sylvatica, Hoppe, in Flora, i. (1818) 472 = perennis.
tenella, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 556 (err. typ. 456) = Adenocline pauciflora.
tenerifolia, Baill. Adansonia, vi. (1866) 323 = Claoxylon tenerifolium.
tomentosa, *Linn. Sp. Pl.* 1035.—Europ. austr.
triandra, E. Mey. in Linnaea, iv. (1829) 237 = Tragia triandra.
tricocca, Eckl. & Zeyh. ex Krauss, in Flora, xxviii. (1845) 85 = seq.?
tricocca, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 111 = Leidesia capensis, obtusa.
violaefolia, Kunze, Cat. Sem. Hort. Lips. (1846).—Cf. Linnaea, xxiv. (1851) 162 (= *Paradenocline procumbens*).—Afr. austr.
Zeyheri, Kunze, in Linnaea, xx. (1847) 54 = Adenocline pauciflora.

MERCURIASTRUM, Heist. Syst. 11 (1748) (Quid?).

MERENDERA, Ram. in Bull. Soc. Philom. (1798) 178. t. 12. *LILIACEAE*, Benth. & Hook. f. iii. 822.
GEOPHILA, Berger. Fl. Bass. Pyr. ii. 184 (1803).
MINDERERA, Ram. ex Schrad. Journ. 1800, II. (1801) 467.
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 337.—Abyssin.
Aitchisoni, Hook. f. Bot. Mag. t. 6012 = persica.
attica, *Boiss. & Sprun. Diagn.* Ser. I. v. 67.—Graecia.
bulbocodioides, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 27 = M. Bulbocodium.
Bulbocodium, Balb. Misc. ii. 14 = Bulbocodium ruthenicum.
Bulbocodium, *Ram. in Bull. Soc. Philom.* (1798) 178. t. 12. f. 2.—Hispan.
caucasica, *Bieb. Fl. Taur. Cauc.* i. 293; *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 27.—Reg. Caucas.; Persia.
caucasica, Sprun. ex Nym. Consp. 744 = attica.
Eichleri, *Boiss. Fl. Orient.* v. 168.—Reg. Caucas.
filifolia, *Cambess. in Mém. Mus. Par.* xiv. (1827) 319.—Afr. bor.
hastulata, *Baker, in Journ. Linn. Soc.* xvii. (1879) 439.—Rumelia.
hissarica, *Regel, in Act. Hort. Petrop.* viii. (1884) 645.—Turkestan.
linifolia, Munby, Fl. Algér. 34 = filifolia.
longispatha, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 337 = abyssinica.
montana, Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 193 = M. Bulbocodium.
nivalis, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 18.—Persia.

MERENDERA :—
persica, *Boiss. Diagn.* Ser. I. xiii. 37.—Persia; Afghan.
pyrenaica, Auct. ex Steud. Nom. ed. II. ii. 130 = M. Bulbocodium.
quadrifolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 19.—Persia.
Raddeana, *Regel, in Act. Hort. Petrop.* vii. (1880) 544.—Reg. Caucas.
robusta, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 515.—Turkest.
Schimperiana, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 337 = abyssinica.
sobolifera, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 34.—As. Min.; Persia.

MERETRICIA, Neraud, in Freyc. Voy. Bot. 28 (1826), sine desc. (Quid?). *RUBIACEAE*.

MERIANA, Trev. Pl. Sel. Ehret. 11. t. 40 (1754) = **Watsonia**, Mill. (Irid.).

MERIANA, Vell. Fl. Flum. 128 (1825); iii. t. 109 (1827) = **Evolvulus**, Linn. (Convolv.).
procumbens, Vell. ll. cc. = E. pusillus.

MERIANA, Vent. in Mém. Inst. Par. (1807) 11 = **Meriania**, Sw. (Melastom.).

MERIANDRA, Benth. in Bot. Reg. sub t. 1282 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1194.
abyssinica, F. Muell. Sel. Pl. Indust. Cult. 130 = benghalensis.
benghalensis, *Benth. in Bot. Reg.* sub t. 1282; & *in Wall. Pl. As. Rar.* i. 29.—Abyss. (Ind. or. cult.)
strobilifera, *Benth. ll. cc.* 1282; 29.—Reg. Himal.

MERIANIA, Sw. Fl. Ind. Occ. ii. 823. t. 15 (1800). *MELASTOMACEAE*, Benth. & Hook. f. i. 749.
ADELBERTIA, Meissn. Gen. 114 (1837).
DAVYA, DC. Prod. iii. 105 (1828).
NOTOCENTRUM, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 131.
PACHYMERIA, Benth. Pl. Hartw. 130 (1844).
SCHWERINIA, Karst. Ausw. Gew. Venez. i. 12 (1848).
WRIGHTEA, Tussac, Fl. Antill. i. sub t. 78 (1808).
adenostemon, Schrank, ex DC. Prod. iii. 131 = Tibouchina adenostemon.
aemula, Schrank & Mart. ex DC. l. c. 128 = Tibouchina aemula.
arborea. *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—N. Granat.
barbinervis, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 129.—Venezuela.
bifrons, Naud. l. c. 127 = purpurea.
bullifera, Griseb. Fl. Brit. W. Ind. 265 = purpurea.
calophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—Bras.
calyptrata, *Triana, l. c.* 66.—Bras.
ciliaris, Vent. Choix, t. 34 (= *Pleroma ciliare*).—N. Granat.
Claussenii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Bras.
coccinea, Sw. ex Loud. Hort. Brit. 174, nomen = leucantha.
cordata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 65.—N. Granat.
dependens, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 124, 129 = Axinaea dependens.
fuscescens, Baill. Hist. Pl. vii. 20 = Adelobotrys fuscescens.
glabra, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Bras.
grandidens, *Triana, l. c.* 65.—N. Granat.
involucrata, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 125 = obtusifolia.
Karstenii, *Naud. l. c.* 128.—Venezuela.
leucantha, *Sw. Fl. Ind. Occ.* ii. 826.—Jamaica.
longipes, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Bras.
macrantha, Linden, ex Planch. in Fl. des Serres, viii. (1852–53) t. 767 = Karstenii.
macrophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—N. Granat.
majalis, Benth. Pl. Hartw. 180 = speciosa.
nana, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 126 = leucantha.

MERIANIA :—
nobilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—N. Granat.
obtusifolia, *DC. Prod.* iii. 101.—Ins. S. Doming.
paniculata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Bras.
parviflora, D. Don, in Mem. Wern. Soc. iv. (1823) 323 = Trembleya parviflora.
prunifolia, *D. Don, l. c.*—Peruv.
purpurea, *Sw. Fl. Ind. Occ.* ii. 829.—Jamaica.
quintuplinervis, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 129.—N. Granat.
Radula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 66.—Peruv.
rigida, *Triana, l. c.*—Ecuador.
rosea, Tuss. Fl. Antill. i. 78. t. 6 = leucantha.
sclerophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—Guiana.
speciosa, *Naud. in Ann. Sc. Nat.* Sér. III. xxviii. (1852) 128.—N. Granat.
speciosa, Naud. l. c., partim = grandidens, Karstenii.
splendens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—N. Granat.
tetraquetra, *Triana, l. c.* 66.—Peruv.
truncata, Schrank & Mart. ex DC. Prod. iii. 106 = Miconia caryophyllaea.
tuberculata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—N. Granat.
umbellata, *Karst. Fl. Columb.* i. 49. t. 24.—N. Granat.
urceolata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 67.—Am. trop.
violacea, Schrank, ex DC. Prod. iii. 108 = Rhynchanthera limosa.

MERICARPAEA, Boiss. Diagn. Ser. I. iii. 51 (1843) *RUBIACEAE*, Benth. & Hook. f. ii. 149.
vaillantioides, *Boiss. Diagn.* Ser. I. iii. 52.—Assyria.

MERIDA, Neck. Elem. ii. 382 (1790) = **Portulaca**, Tourn.

MERIDIANA, Hill, Veg. Syst. ii. 121** (1761) = **Gazania**, Gaertn. (Compos.).
tesselata, Hill, Hort. Kew. 26 = G. rigens.

MERIDIANA, Linn. f. Suppl. 248 (1781) = **Portulaca**, Tourn.
axilliflora, Schrank, in Bot. Zeit. Regensb. iii. (1804) 354 = P. axilliflora.
elliptica, Poir. in Dict. Sc. Nat. xxx. 112 = P. Meridiana.
quadrifida, Poir. l. c. 111 = P. Meridiana.

MERIMEA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 160. t. 107 (1829) = **Bergia**, Linn. (Elatin.).
arenarioides, Cambess. l. c. = B. arenarioides.
texana, Hook. Ic. Pl. t. 278 = B. texana.

MERIOLIX, Rafin. in Am. Monthly Mag. (1819) 192 = **Oenothera**, Linn. (Onagr.).
Berlandieri, Walp. Rep. ii. 79 = Oe. serrulata.
serrulata, Rafin. in Am. Monthly Mag. (1819) 192 = Oe. serrulata.

MERIONE, Salisb. Gen. Pl. Fragm. 12 (1866) = **Dioscorea**, Linn.
peltata, Salisb. l. c., nomen = D. peltata?
villosa, Salisb. l. c. = D. villosa.

MERISACHNE, Steud. in Flora, xxxiii. (1850) 228, nomen; Syn. Pl. Gram. 117 (1855) = **Triodia**, R. Br. (Gramin.).
Drummondii, Steud. ll. cc.—Texas.

MERISTA, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 47 = **Myrsine**, Linn.
laevigata, Banks & Soland. ex A. Cunn. l. c. = Myrsine Urvillei.

MERISTOSTIGMA, A. Dietr. Sp. Pl. ii. 593 (1833) = **Lapeyrousia**, Pourr. (Irid.).
aculeatum, Steud. Nom. ed. II. ii. 130 = L. compressa.
anceps, A. Dietr. Sp. Pl. ii. 596 = L. compressa.
azureum, Steud. Nom. ed. II. ii. 130 = L. corymbosa.
bracteatum, A. Dietr. Sp. Pl. ii. 597 = L. fissifolia.
corymbosum, A. Dietr. l. c. 594 = L. corymbosa.
Fabricii, A. Dietr. l. c. 597 = L. Fabricii.
falcatum, A. Dietr. l. c. 594 = L. falcata.
fissifolium, A. Dietr. l. c. 595 = L. fissifolia.
heterophyllum, A. Dietr. l. c. = L. fasciculata.

MERISTOSTIGMA :—
junceum, Steud. Nom. ed. II. ii. 130 = Lapeyrousia juncea.
laxum, A. Dietr. Sp. Pl. ii. 597.—Afr. austr.
manuleaeflorum, Steud. Nom. ed. II. ii. 130 = L. micrantha.
silenoides, A. Dietr. Sp. Pl. ii. 596 = L. silenoides.
Sparmanni, Steud. Nom. ed. II. ii. 130 = Freesia refracta.

MERISTOSTYLIS, Klotzsch, in Peters, Reise Mossamb. Bot. 267 (1861). *GENTIANEAE.*
brachycalyx, *Klotzsch, l. c.* 270.—Afr. trop.
grandiflora, *Klotzsch, l. c.* 268.—Afr. trop.
macrocalyx, *Klotzsch, l. c.* 269.—Afr. trop.

MERISTOTROPIS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 95 (1843) = **Glycyrrhiza**, Tourn. (Legumin.).
triphylla, Fisch. & Mey. l. c. 96 = G. triphylla.

MERIZADENIA, Miers, Apocyn. S. Am. 78 (1878) = **Tabernaemontana**, Plum. (Apocyn.).
amplifolia, Miers, l. c. 79 = T. macrophylla, *Poir.*
arcuata, Miers, l. c. = T. arcuata.
Sananho, Miers, l. c. 78 = T. Sananho.

MERKIA, Reichb. Handb. 298 (1837) = MERCKIA, Fisch. = **Arenaria**, Rupp. (Caryoph.).

MERKUSIA, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 150 = **Scaevola**, Linn. (Gooden.).
aemula, De Vriese, in Verh. Haarl. x. (1854) 74 = S. aemula.
anchusaefolia, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 165 = S. anchusaefolia.
angulata, De Vriese, l. c. 167 = S. angulata.
attenuata, De Vriese, l. c. 162 = S. attenuata.
caespitosa, De Vriese, l. c. 164 = S. globulifera.
crassifolia, De Vriese, l. c. 151 = S. crassifolia.
cuneiformis, De Vriese, l. c. 157 = S. cuneiformis.
depauperata, De Vriese, in Verh. Haarl. x. (1854) 74 = S. depauperata.
depressa, De Vriese, l. c. 70 = S. depressa.
fastigiata, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 152 = S. nitida.
glandulifera, De Vriese, l. c. 166 = S. glandulifera.
globulifera, De Vriese, l. c. 156 = S. globulifera.
hispida, De Vriese, l. c. 162 = S. hispida.
Hookeri, De Vriese, l. c. 159 = S. Hookeri.
humifusa, De Vriese, in Verh. Haarl. x. (1854) 70 = S. humifusa.
humilis, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 161 = S. humilis.
linearis, De Vriese, l. 166 = S. linearis.
longifolia, De Vriese, in Verh. Haarl. x. (1854) 69 = S. longifolia.
lyratifolia, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 161 = S. lyratifolia.
macrostachya, De Vriese, l. c. 154 = S. macrostachya.
microcarpa, De Vriese, l. c. 157 = S. microcarpa.
molluginea, De Vriese, in Verh. Haarl. x. (1854) 71 = S. humifusa.
multiflora, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 152 = S. nitida.
myrtifolia, De Vriese, in Verh. Haarl. x. (1854) 72 = S. Groeneri.
nitida, De Vriese, l. c. 73 = S. nitida.
ovalifolia, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 153 = S. ovalifolia.
pallida, De Vriese, l. c. 159 = S. microcarpa.
paludosa, De Vriese, l. c. 167 = S. paludosa.
revoluta, De Vriese, l. c. 164 = S. revoluta.
sinuata, De Vriese, l. c. 160 = S. aemula.
suaveolens, De Vriese, l. c. 163 = S. suaveolens.
thesioides, De Vriese, l. c. 155 = S. thesioides.

MEROPE, M. Roem. Syn. Hesper. 44 (1846) = **Atalantia**, Correa (Rutac.).
spinosa, M. Roem. l. c. (= *Sclerostylis spinosa*).—Java.

MEROPE, Wedd. Chlor. And. i. 160. t. 24 (1856) = **Gnaphalium**, Linn. (Compos.).
aretioides, Wedd. l. c. 164 = Antennaria aretioides.
argentea, Wedd. l. c. 163.—Boliv.
caespititia, Wedd. l. c. 164.—Boliv.
erythractis, Wedd. l. c. 162 = G. erythractis.

MEROPE :—
Kunthiana, Wedd. Chlor. And. i. 161. t. 24 (1856) = G. depressum ?
piptolepis, Wedd. l. c. 162 = G. piptolepis.
Schultzii, Wedd. l. c. 163 = G. evacoides.
virescens, Wedd. l. c.—Boliv.

MEROPHRAGMA, Dulac, Fl. Hautes-Pyr. 365 (1867) = **Telephium**, Tourn. (Ficoid.).
terrestre, Dulac, l. c. = T. Imperati.

MEROSTACHYS, Spreng. Syst. i. 132 (1825). *GRAMINEAE*, Benth. & Hook. f. iii. 1209.
brevispica, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 49.—Peruv.
Burchellii, *Munro, l. c.* 51.—Bras.
capitata, Hook. Ic. Pl. tt. 273, 274 = Athroostachys capitata.
cirrhosa, Nees, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 48 = speciosa.
Clausseni, *Munro, l. c.*—Bras.
Fischeriana, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 215.—Bras.
fistulosa, *Doell, l. c.* 209.—Bras.
Kunthii, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 127. t. 10. f. 30.—Bras.
Neesii, *Rupr. l. c.*—Bras.
petiolata, *Doell, in Mart. Fl. Bras.* ii. III. 216.—Bras.
racemiflora, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 586 = Arthrostylidium racemiflorum.
Riedeliana, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 213.—Bras.
Sellovii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 51.—Paraguay.
sparsiflora, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 127. t. 10. f. 32.—Bras.
speciosa, Kunth, Rév. Gram. i. 139 = Kunthii.
speciosa, Nees, Agrost. Bras. 527 = Neesii.
speciosa, *Spreng. Syst.* 249.—Bras.
ternata, *Nees, Agrost. Bras.* 529.—Bras.

MERREMIA, Dennst. Schluess. Hort. Malab. 34 (1818) = **Convolvulus**, Tourn.
convolvulacea, Dennst. l. c. = C. flavus.

MERTENSIA, H. B. & K. Nov. Gen. et Sp. ii. 80. t. 103 (1817) = **Celtis**, Tourn. (Urtic.).
aculeata, Miq. in Mart. Fl. Bras. iv. I. 173, in syn. = C. aculeata.
brasiliensis, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 339 = C. brasiliensis.
citrifolia, H. B. & K. Nov. Gen. et Sp. ii. 32 = C. pubescens.
commutata, Hemsl. Biol. Centr. Am. Bot. iii. 138, in syn. = C. aculeata.
Goudotii, Planch. in DC. Prod. xvii. 187, in syn. = C. pubescens.
Iguanea, Schult. Syst. vi. 312 = C. aculeata.
integrifolia, Schult. l. c. 313 = C. mississipiensis.
laevigata, H. B. & K. Nov. Gen. et Sp. ii. 31. t. 103 = C. aculeata.
pubescens, Schult. Syst. vi. 312 = C. pubescens.
rhamnoides, Schult. l. c. 313 = C. aculeata.
senegalensis, Perr. ex Blume, Mus. Bot. Lugd. Bat. ii 69 = C. mississipiensis.
Tala, Wedd. ex Planch. in DC. Prod. xvii. 191 = C. Tala.
texana, Tréc. ex Planch. l. c. = C. Tala.
zizyphoides, H. B. & K. Nov. Gen. et Sp. ii. 31 = C. aculeata.

MERTENSIA, Roth, Catalect. i. 34 (1797). *BORAGINEAE*, Benth. & Hook. f. ii. 857.
CASSELIA, Dum. Comm. Bot. 21 (1822).
HIPPOGLOSSUM, Hartm. ex Lilja, in Linnaea, xvii. (1843) 111.
OREOCHARIS, Lindl. Veg. Kingd. 656 (1847).
STEENHAMMERA, Reichb. Fl. Germ. Excurs. 337 (1831).
PLATYNEMA, Schrad. Ind. Sem. Hort. Gotting. (1835).—Cf. Linnaea, xi. (1837) Litt. 89.
PNEUMARIA, Hill, Veg. Syst. vii. 40 (1764), nomen prius.
WINKLERIA, Reichb. Nom. 236 (1841).
alpina, *G. Don, Gen. Syst.* iv. 372.—Am. bor. occ.
alpina, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862). 340, partim = lanceolata.
brevistyla, S. Wats. in Bot. King's Exp. 239 = alpina.
ciliata, G. Don, Gen. Syst. iv. 372 = sibirica.

MERTENSIA :—
corymbosa, G. Don, Gen. Syst. iv. 319 = paniculata.
dahurica, *G. Don, l. c.* 318.—Dahuria.
denticulata, G. Don, l. c. 319 = sibirica.
Drummondii, G. Don, l. c. = sibirica.
dschagastanica, *Regel, in Act. Hort. Petrop.* vi. (1879) 340.—Turkestan.
echioides, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 170.—Reg. Himal.
elliptica, *Ledeb. Fl. Ross.* iii. 135.—Kamschatka.
elongata, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 170.—Reg. Himal.
Fendleri, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 339 = lanceolata.
kamczatica, DC. Prod. x. 91 = paniculata.
lanceolata, *DC. l. c.* 88.—Am. bor. occ.
longistyla, *Ledeb. Fl. Ross.* iii. 135.—Ins. Karaginsk.
marginata, G. Don, Gen. Syst. iv. 319 = lanceolata.
maritima, *S. F. Gray, Nat. Arr. Brit. Pl.* ii. 354.—Reg. bor. temp. et arct.
moltkioides, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 170.—Reg. Himal.
oblongifolia, *G. Don, Gen. Syst.* iv. 372.—Am. bor. occ.
Pallasii, G. Don, l. c. 319 = sibirica.
paniculata, *G. Don, l. c.* 318.—Am. et As. bor.
parviflora, G. Don, l. c. 320 = maritima.
pilosa, DC. Prod. x. 90 = paniculata.
primuloides, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 170.—Reg. Himal.
pubescens, DC. Prod. x. 90 = paniculata.
pulmonarioides, *Roth, Catalect.* i. 34.—Am. bor.
racemosa, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 171.—Reg. Himal.
rivularis, *DC. Prod.* x. 90.—Sibiria.
Sendtneri, Janka, in Oestr. Bot. Zeitschr. ix. (1859) 314 = Zwackhia aurea.
serbica, *Janka, l. c.*—Europ.
serrulata, *DC. Prod.* x. 89.—Sibir. baical.
sibirica, *G. Don, Gen. Syst.* iv. 319.—Sibir. or.; Am. bor. occ.
sibirica, Torr. in Bot. U. St. Expl. Exp. ii. 412 = paniculata.
simplicissima, *G. Don, Gen. Syst.* iv. 319.—Sibir. or.
stomatechoides, Kellogg, in Proc. Calif. Acad. ii. (1863) 148 = sibirica.
stylosa, *DC. Prod.* x. 91.—Sibir. baical.
symphytoides, Fisch. ex Herd. in Act. Hort. Petrop. i. (1873) 511 = paniculata.
tibetica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 171.—Reg. Himal.
villosula, *G. Don, Gen. Syst.* iv. 319.—Mont. Carpath.
virginica, Link, Handb. i. 580 = pumonarioides.

MERYTA, Forst, Char. Gen. 119. t. 60 (1776) *ARALIACEAE*, Benth. & Hook. f. i. 940.
BOTRYODENDRUM, Endl. Prod. Fl. Norf. 62 (1833).
NEARA, Soland. ex Seem. Fl. Vit. 118 (1865).
agglomerata, *Poir. Tabl. Encyc.* iii. 399. t. 803.—Hab. ?
angustifolia, *Seem. in Bonplandia*, x. (1862) 295.—Ins. Norfolk.
Balansae, *Baill. Adansonia*, xii. (1878) 156.—N. Caled.
coriacea, *Baill. l. c.* 153.—N. Caled.
Denhamii, *Seem. in Bonplandia*, x. (1862) 295.—N. Caled.
lanceolata, *Forst. Char. Gen.* 120.—Ins. Pacif.
latifolia, *Seem. in Bonplandia*, x. (1862) 295.—Ins. Norfolk.
macrocarpa, *Baill. Adansonia*, xii. (1878) 155.—N. Caled.
macrocephala, *Baill. l. c.* 158.—N. Caled.
macrophylla, *Seem. in Bonplandia*, x. (1862) 294.—Ins. Samoa.
microcarpa, *Baill. Adansonia*, xii. (1878) 156.—N. Caled.
oxylaena, *Baill. l. c.* 157.—N. Caled.
pachycarpa, *Baill. l. c.*—N. Caled.
pallens, *Baill. l. c.*—N. Caled.
schizolaena, *Baill. l. c.*—N. Caled.
Sinclairii, *Seem. in Bonplandia*, x. (1862) 295.—N. Zel.
sonchifolia, *Linden & André, Illustr. Hortic.* xxvi. (1879) 42. t. 340.—N. Caled.

MESADENIA, Rafin. in Loud. Gard. Mag. viii. (1832) 247 = **Senecio**, Tourn. (Compos.).
atriplicifolia, Rafin. New Fl. Am. iv. 79 = S. atriplicifolius.

MESADENIA:—
dentata, Rafin. New Fl. Am. iv. 79 (Quid?).—Am. bor.
lanceolata, Rafin. l. c. = Senecio Boscianus.
oblonga, Rafin. l. c. 80 (Quid?).—Am. bor.
ovata, Rafin. l. c. 79 = S. Walteri?
paniculata, Rafin. l. c. = S. Nuttallii?
plantaginea, Rafin. l. c. (Quid?).—Am. bor.
pteranthes, Rafin. l. c. = S. Nuttalli?
pulverulenta, Rafin. l. c. (Quid?).—Am. bor.
reniformis, Rafin. l. c. = S. Muehlenbergii.
rotundifolia, Rafin. l. c. = Vernonia cinerea.

MESANCHUM, Dulac, Fl. Hautes-Pyr. 320 (1867) = **Tillaea**, Linn. (Crassul.).
connatum, Dulac, l. c. = T. muscosa.

MESANDRINIA, Rafin. Neogenyt. 3 (1825) = **Jatropha**?, Linn. (Euphorb.).

MESANTHEMUM, Koern. in Linnaea, xxvii. (1854) 572. *ERIOCAULEAE*, Benth. & Hook. f. iii. 1021.
platyphyllum, *Baker, in Journ. Linn. Soc.* xx. (1883) 278.—Madag.
Prescottianum, *Koern. in Mart. Fl. Bras.* iii. 1. 472.—Bras.
pubescens, *Koern. in Linnaea*, xxvii. (1854) 575.—Madag.
radicans, *Koern. l. c.* 573.—Afr. trop.
Rutenbergianum, *Koern. in Bremen, Abh.* vii. (1880) 34.—Afr. trop.

MESANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836) = **Cannomois**, Beauv. (Restiac.).
macrocarpus, Nees, l. c. = C. virgata.
Ricinus, Nees, ex Mast. in Journ. Linn. Soc. x. (1869) 236 = C. scirpoides.

MESECHITES, Muell. Arg. in Mart. Fl. Bras. vi. 1. 150. t. 46 (1860) = **Echites**, Linn. (Apocyn.).
Andrieuxii, Miers, Apocyn. S. Am. 235 = Mandevilla Andrieuxii.
angustata, Miers, l. c. 231 = Macrosiphonia longiflora.
angustifolia, Miers, l. c. 230 = E. angustifolia.
Brownei, Miers, l. c. 232 = E. torulosa.
dichotoma, Miers, l. c. 233 = E. dichotoma, *H. B. & K.*
guayaquilensis, Miers, l. c. 233 = E. guayaquilensis (in Addend.).
guianensis, Miers, l. c. 235 = E. guianensis.
hastata, Miers, l. c. 233.—Cuba.
hirtella, Miers, l. c. 234 = Prestonia hirsuta.
hirtellula, Miers, l. c. = Mandevilla oaxacana.
japurensis, Muell. Arg. in Mart. Fl. Bras. vi. 1. 152 = E. japurense.
jasminiflora, Miers, Apocyn. S. Am. 235 = E. jasminiflora.
lanceolata, Miers, l. c. 230 = Parsonia lanceolata.
linearifolia, Miers, l. c. = E. linearifolia.
myrtifolia, Muell. Arg. in Linnaea, xxx. (1859–60) 445 = E. rosea.
oaxacana, Miers, Apocyn. S. Am. 234 = Mandevilla oaxacana.
repens, Miers, l. c. 229 = E. repens.
rosea, Miers, l. c. 232 = E. rosea.
subcarnosa, Miers, l. c. = E. subcarnosa.
sulphurea, Muell. Arg. in Mart. Fl. Bras. vi. 1. 151 = E. sulphurea.
torulosa, Miers, Apocyn. S. Am. 229 = E. torulosa.
trifida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 151 = E. trifida.

MESEMBRIANTHEMUM, Spreng. Nachr. i. Bot. Gart. Halle (1801), 28 = seq.

MESEMBRYANTHEMUM, Dill. ex Linn. Syst. ed. I (1735). *FICOIDEAE*, Benth. & Hook. f. i. 853.
ABRYANTHEMUM, Neck. Elem. ii. 82 (1790).
FICOIDES, Rupp. Fl. Jen. ed. Hall. 164 (1745).
FICULA, Heist. Syst. 8 (1748).
GASOUL, Adans. Fam. ii. 243 (1763).
GYNICIDIA, Neck. Elem. ii. 81 (1790).
HYMENOGYNE, Haw. Rev. Pl. Succ. 192 (1821).
MANETTIA, Adans. Fam. ii. 242 (1763).
MESEMBRYANTHUS, Neck. Elem. ii. 81 (1790).
MESEMBRYUM, Adans. Fam. ii. 242 (1763).
NYCTERANTHUS, Neck. Elem. ii. 82 (1790).
VOSSIA, Adans. Fam. ii. 243 (1763).

MESEMBRYANTHEMUM:—
abbreviatum, *Haw. in Phil. Mag.* (Dec. 1826) 330.—Afr. austr.
abyssinicum, *Regel, Gartenfl.* (1873) 299.—Abyssin.
acaule, Linn. Syst. ed. X. 1059 = bellidiflorum.
acinaciforme, DC. Pl. Grass. t. 89 = lacerum.
acinaciforme, *Linn. Sp. Pl.* 485.—Afr. austr.
acuminatum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 426.—Afr. austr.
acutangulum, *Haw. l. c.* 424.—Afr. austr.
acutum, *Haw. Misc. Nat.* 26.—Afr. austr.
adscendens, *Haw. Syn. Pl. Succ.* 220.—Afr. austr.
aduncum, Eckl. & Zeyh. Enum. 316 = micranthum.
aduncum, *Haw. Misc. Nat.* 87.—Afr. austr.
aduncum, Jacq. Fragm. 43 = Schollii.
aduncum, Willd. Enum. Hort. Berol. 534 = curvifolium.
aequilaterale, *Haw. Misc. Nat.* 77.—Austral.; Am. austr.
aggregatum, Haw. Obs. Mesemb. 419 = grossum, *Soland.*
agninum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 127.—Afr. austr.
Aitonis, *Jacq. Hort. Vindob.* iii. 8.—Afr. austr.
albicaule, *Haw. in Till. Phil. Mag.* lxviii. (1826) 331.—Afr. austr.
albidum, *Linn. Sp. Pl.* ed. II. 699.—Afr. austr.
albinatum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 126.—Afr. austr.
albipunctatum, Haw. l. c. = albinatum.
album, Haw. ex Steud. Nom. ed. II. ii. 131 = sessiliflorum.
aloides, *Haw. Suppl. Pl. Succ.* 88; *Burch. Trav.* ii. 332.—Afr. austr.
alsinifolium, *Hort. ex G. Don, in Loud. Hort. Brit.* 480, nomen.—Austral.?
amoenum, *Salm-Dyck, ex DC. Prod.* iii. 466.—Afr. austr.
ampullaceum, *Cat. Hort. Liverp.* (1808) 203; *Steud. Nom.* ed. II. ii. 131.—Hab.?
anatomicum, *Haw. Misc. Nat.* 50.—Afr. austr.
anceps, *Haw. Syn. Pl. Succ.* 289.—Afr. austr.
angulatum, *Thunb. Prod. Pl. Cap.* 91.—Afr. austr.
angustum, *Haw. Obs. Mesemb.* 176.—Afr. austr.
anomalum, Willd. Enum. Hort. Berol. 531 = umbellatum.
apetalum, *Linn. f. Suppl.* 258.—Afr. austr.
aphyllum, Hort. ex DC. Prod. iii. 446 = junceum.
arboriforme, *Burch. Trav.* i. 343.—Afr. austr.
aridum, Moench, Meth. Suppl. 192 = tortuosum.
aristulatum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 424.—Afr. austr.
articulatum, *Jacq. ex Steud. Nom.* ed. I. 525; ed. II. ii. 131 (Quid?).—Afr. austr.
articulatum, Thunb. in Nov. Act. Nat. Cur. viii. (1791), App. 10 = junceum, granulicaule.
asperulum, *Salm-Dyck, Monog.* iv. § 50. f. 2.—Afr. austr.
asperum, *Haw. Rev. Pl. Succ.* 145.—Afr. austr.
attenuatum, *Haw. l. c.* 188.—Afr. austr.
aurantiacum, *Haw. Misc. Nat.* 84.—Afr. austr.
aurantium, Willd. Enum. Hort. Berol. 537 = aurantiacum.
auratum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 449.—Afr. austr.
aureum, *Linn. Syst. Nat.* ed. X. 1060.—Afr. austr.
aureum, Thunb. in Nov. Act. Nat. Cur. viii. (1791), App. 16 = auratum.
australe, *Soland. ex Forst. f. Prod.* 90; *et in Ait. Hort. Kew.* ed. I. ii. 187.—Austral.; Ins. Pacif.
barbatum, *Linn. Sp. Pl.* 482.—Afr. austr.
bellidiflorum, *Linn. l. c.*—Afr. austr.
bellidifolium, Linn. Syst. ed. XII. 345 = praec.
bellidiforme, Burm. f. Fl. Cap. Prod. 15 = praec.?
bibracteatum, Haw. Syn. Pl. Succ. 215 = rostratum.
bicolor, *Linn. Sp. Pl.* 485.—Afr. austr.
bicolorum, Linn. l. c. ed. II. 695 = praec.
bicorne, *Sond. in Harv. & Sond. Fl. Cap.* ii. 434.—Afr. austr.
bidentatum, *Haw. Suppl. Pl. Succ.* 89.—Afr. austr.
bifidum, *Haw. Misc. Nat.* 29.—Afr. austr.
bifolium, *Breit. ex Hoffmgg. Verz. Pfl.* 79.—Hab.?
bigibberatum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 329.—Afr. austr.
blandum, *Haw. Suppl. Pl. Succ.* 95.—Afr. austr.
Boerhavii, Eckl. & Zeyh. Enum. 318 = hispidum.
Bolusii, *Hook. f. Bot. Mag.* t. 6664.—Afr. austr.
brachiatum, DC. Pl. Grass. t. 129 = nitidum, *Haw.*

MESEMBRYANTHEMUM:—
brachiatum, Eckl. & Zeyh. Enum. 318 = subincanum.
brachiatum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 191 = decussatum.
brachyphyllum, *Welw. in Journ. Bot.* xi. (1873) 290. t. 136.—Lusitan.
bracteatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 185.—Afr. austr.
bracteatum, Willd. Sp. Pl. ii. 1050 = stipulaceum.
brevicaule, *Haw. Suppl. Pl. Succ.* 91.—Afr. austr.
brevifolium, Eckl. & Zeyh. Enum. 318 = obliquum, *Haw.*
brevifolium, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 188.—Afr. austr.
bulbosum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 428.—Afr. austr.
Burchelli, Hort. Genev. ex A. DC. 7me Not. Pl. Rar. Jard. Genèv. 29 = blandum.
Burmanni, *Haw. Rev. Pl. Succ.* 106.—Afr. austr.
caducum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 179.—Afr. austr.
calamiforme, *Linn. Sp. Pl.* 481.—Afr. austr.
calendulaceum, Eckl. & Zeyh. Enum. 323 = sabulosum.
calendulaceum, Haw. Misc. Nat. 47 = helianthoides.
calendulaceum, Haw. Rev. Pl. Succ. 161 = pomeridianum.
calycinum, Eckl. & Zeyh. Enum. 319 = longispinulum.
calycinum, *Haw. Rev. Pl. Succ.* 187.—Afr. austr.
campestre, *Burch. Trav.* i. 259.—Afr. austr.
canaliculatum, *Haw. Obs. Mesemb.* 218.—Afr. austr.
candens, *Haw. Rev. Pl. Succ.* 186.—Afr. austr.
Candollii, Eckl. & Zeyh. Enum. 323 = pomeridianum.
Candollii, *Haw. Rev. Pl. Succ.* 160.—Afr. austr.
canescens, Haw. l. c. 135 = pulchellum.
caninum, *Lam. Encyc.* ii. 487; *Haw. Obs. Mesemb.* 159.—Afr. austr.
canum, *Haw. l. c.* 158.—Afr. austr.
canum, Salm-Dyck, Obs. 20 = carinans.
capillare, Thunb. in Nov. Act. Nat. Cur. viii. (1791), App. 13 = brevifolium.
capitatum, *Haw. Misc. Nat.* 41.—Afr. austr.
carinans, *Haw. Rev. Pl. Succ.* 90.—Afr. austr.
carinatum, Vent. Jard. Malm. 109 = lacerum.
carneum, Haw. Obs. Mesemb. 206 = grossum, *Soland.*
caspicum, S. G. Gmel. Iter, iv. 48; ex Ledeb. Fl. Ross. i. 490 = Peganum crithmifolium.
caulescens, *Mill. Gard. Dict.* ed. VIII. n. 12; *Haw. Obs. Mesemb.* 367.—Afr. austr.
cauliculatum, Haw. Suppl. Pl. Succ. 90 = loreum, *Linn.*
ceratophyllum, Willd. Enum. Hort. Berol. Suppl. 36 = curvifolium.
chilense, *Molina, Sagg. Chile*, ed. II. 133.—Chili.
ciliatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 179.—Afr. austr.
ciliatum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791), *App.* 11.—Afr. austr.
clandestinum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 129.—Afr. austr.
clavatum, Haw. Obs. Mesemb. 478, nomen = pyropeum.
clavatum, *Jacq. Hort. Schoenb.* i. 56. t. 108.—Afr. austr.
clavellatum, Haw. Misc. Nat. 79 = australe.
claviforme, DC. Prod. iii. 448 = pyropeum.
coccineum, Eckl. & Zeyh. Enum. 316 = bicolor.
coccineum, *Haw. Obs. Mesemb.* 247.—Afr. austr.
collinum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 443.—Afr. austr.
compactum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 191.—Afr. austr.
compressum, *Haw. Obs. Mesemb.* 326.—Afr. austr.
compressum, Haw. l. c. 416, nomen = virgatum.
concavum, *Haw. Rev. Pl. Succ.* 168.—Afr. austr.
confertum, Haw. Syn. Pl. Succ. 301 = incurvum, roseum.
congestum, *Salm-Dyck, Monog.* iii. § 23. f. 2.—Afr. austr.
conspicuum, *Haw. Syn. Pl. Succ.* 240.—Afr. austr.
conspicuum, *Spreng. Nov. Prov.* 27 (an praec.?).—Hab.?
Cooperi, *Hook. f. Bot. Mag.* t. 6312.—Afr. austr.
copticum, Jacq. Hort. Vindob. iii. t. 6 = apetalum.
copticum, Linn. Sp. Pl. ed. II. 688 = Aizoon hispanicum.
coralliflorum, *Salm-Dyck, Monog.* iv. § 40. f. 2.—Afr. austr.
coralliforme, *Vest, in Syll. Ratisb.* ii. (1828) 224.—Hab.?

MESEMBRYANTHEMUM:—
corallinum, Haw. Rev. Pl. Succ. 154 = coralliflorum.
corallinum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 12.—Afr. austr.
coralloides, *DC. Prod.* iii. 451.—Afr. austr.
cordifolium, *Linn. f. Suppl.* 260.—Afr. austr.
coriarium, *Burch. Trav.* i. 243.—Afr. austr.
corniculatum, Eckl. & Zeyh. Enum. 309 = Thunbergii.
corniculatum, Haw. Obs. Mesemb. 411, nomen = loreum, *Linn.*
corniculatum, *Linn. Sp. Pl.* ed. II. 676.—Afr. austr.
cornutum, *Salm. ex Spreng. Syst.* ii. 519.—Afr. austr.
coruscans, *Haw. Suppl. Pl. Succ.* 90.—Afr. austr.
crassicaule, *Haw. in Till. Phil. Mag.* lxiv. (1824) 425.—Afr. austr.
crassifolium, Eckl. & Zeyh. Enum. 310 = Thunbergii.
crassifolium, *Linn. Sp. Pl.* 484.—Afr. austr.
crassulinum, *Spreng. Syst.* iv. *Cur. Post.* 196; *DC. Prod.* iii. 445.—Afr. austr.
crassuloides, Haw. Rev. Pl. Succ. 170 = crassulinum.
criniflorum, *Linn. f. Suppl.* 259; *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
crispatum, Haw. Obs. Mesemb. 478, nomen = crispum.
crispum, *Haw. Misc. Bot.* 44.—Afr. austr.
croceum, *Jacq. Fragm.* 17. t. 11. f. 2.—Afr. austr.
cruciatum, *Haw. Obs. Mesemb.* 173.—Afr. austr.
cryptanthum, *Hook. f. in Hook. Ic. Pl.* t. 1034.—Ins. S. Helen.
crystallinum, *Linn. Sp. Pl.* 480.—Graecia; Ins. Canar.; Afr. austr.; Calif.
crystallophanes, Eckl. & Zeyh. Enum. 322 = angulatum.
cultratum, *Salm-Dyck, Obs.* (1820) 7.—Afr. austr.
cuneifolium, Jacq. Coll. ii. 319 = criniflorum.
curtum, *Haw. Syn. Pl. Succ.* 334.—Afr. austr.
curviflorum, *Haw. Rev. Pl. Succ.* 147.—Afr. austr.
curvifolium, DC. Prod. iii. 436 = praec.
curvifolium, *Haw. Misc. Nat.* 88.—Afr. austr.
cylindricum, *Haw. Obs. Mesemb.* 411.—Afr. austr.
cymbifolium, Eckl. & Zeyh. Enum. 312 = Dregeanum.
cymbifolium, *Haw. in Till. Phil. Mag.* lxiv. (1824) 424.—Afr. austr.
cymbiforme, Haw. Obs. Mesemb. 264 = sessile.
dactylinum, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 582.—Afr. trop.
debile, Haw. in Till. Phil. Mag. lxviii. (1826) 330 = reptans.
decipiens, Haw. Rev. Pl. Succ. 110 = Thunbergii.
decumbens, *Mill. ex Weston, Cat.* 162; *ex Haw. Obs. Mesemb.* 352.—Afr. austr.
decussatum, *Thunb. Prod. Pl. Cap.* 88.—Afr. austr.
deflexum, Salm-Dyck, Obs. 27 = elegans.
deflexum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 187.—Afr. austr.
defoliatum, Haw. Misc. Nat. 64 = clavatum.
deltoides, *Linn. Sp. Pl.* 482.—Afr. austr.
demissum, Willd. Enum. Hort. Berol. Suppl. 36 = australe.
densum, *Haw. Obs. Mesemb.* 302.—Afr. austr.
dentatum, *Kern. Hort.* t. 26; *ex Pritz. Ic. Ind.* 706 (Quid?).—Hab.?
denticulatum, *Haw. Obs. Mesemb.* 149.—Afr. austr.
denticulatum-canum, *Haw. l. c.*—Afr. austr.
depressum, *Haw. Misc. Nat.* 33.—Afr. austr.
depressum, Sims, Bot. Mag. t. 1866 = longum.
difforme, *Haw. Syn. Pl. Succ.* 225.—Afr. austr.
difforme, *Linn. Sp. Pl.* 487.—Afr. austr.
difforme, Thunb. Fl. Cap. ed Schult. 423 = denticulatum, namaquense.
digitatum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 181 = seq.
digitiforme, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 6.—Afr. austr.
dilatatum, *Haw. Misc. Nat.* 68.—Afr. austr.
dimidiatum, *Haw. Syn. Pl. Succ.* 233; *Rev. Pl. Succ.* 119.—Afr. austr.
diminutum, Haw. Misc. Nat. 26 = loreum, *Linn.*
dimorphum, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 582.—Afr. trop.
diversifolium, *Haw. Misc. Nat.* 38.—Afr. austr.
diversiphyllum, Haw. Obs. Mesemb. 228 = diversifolium.
dolabriforme, *Linn. Sp. Pl.* 487.—Afr. austr.
dolabriformoides, Haw. Obs. Mesemb. 168 = praec.
Dregeanum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 414.—Afr. austr.
dubium, Haw. Misc. Nat. 39 = Thunbergii.
dubium, Salm-Dyck, Cat. Hort. Dyck. 22 = procumbens.

MESEMBRYANTHEMUM:—
dunense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 411.—Afr. austr.
echinatum, *Lam. Encyc.* ii. 478; [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 194.—Afr. austr.
Ecklonis, *Salm-Dyck, Monog.* iv. § 49. f. 5.—Afr. austr.
edentulum, Haw. Rev. Pl. Succ. 125 = uncinatum.
edule, *Linn. Syst.* ed. X. 1060.—Afr. austr.
effusum, Salm-Dyck, ex Steud. Nom. ed. II. ii. 132 = geminiflorum.
elegans, *Jacq. Hort. Schoenb.* iv. 18.—Afr. austr.
elongatum, Eckl. & Zeyh. Enum. 321 = angulatum.
elongatum, *Haw. Obs. Mesemb.* 236.—Afr. austr
emarcidum, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 9 = anatomicum.
emarginatoides, Haw. Obs. Mesemb. 343 = elegans.
emarginatum, Eckl. & Zeyh. Enum. 317 = bicolor, linguiforme, *Linn.*
emarginatum, *Linn. Sp. Pl.* ed. II. 692.—Afr. austr.
equilateratum, *Haw. Obs. Mesemb.* 390.—Afr. austr.
erectum, *Haw. Misc. Nat.* 53.—Afr. austr.
erectum, *Hort. ex Link, Enum. Hort. Berol.* ii. 41 (an praec.?).—Afr. austr.
erigeriflorum, Jacq. Hort. Schoenb. iv. 39 = brevifolium.
erminium, *Haw. in Till. Phil. Mag.* lxviii. (1826) 126.—Afr. austr.
expansum, DC. Pl. Grass. t. 47 = pallens.
expansum, *Linn. Syst.* ed. X. 1059.—Afr. austr.
falcatum, Lam. Encyc. ii. 482 = glomeratum.
falcatum, *Linn. Sp. Pl.* 484.—Afr. austr.
falcatum, Murr. Syst. ed. XIII. 391 = acinaciforme.
falciforme, *Haw. Syn. Pl. Succ.* 299.—Afr. austr.
fasciculatum, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 11 = tetragonum.
fastigiatum, *Haw. Obs. Mesemb.* 210.—Afr. austr.
fastigiatum, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 7 = papuliferum.
felinum, *Lam. Encyc.* ii. 486; *Haw. Obs. Mesemb.* 161.—Afr. austr.
fibulaeforme, Haw. Syn. Pl. Succ. 204 = seq.
fibuliforme, *Haw. Misc. Nat.* 22.—Afr. austr.
ficiforme, *Haw. Rev. Pl. Succ.* 83.—Afr. austr.
filamentosum, DC. Pl. Grass. t. 60 = mutabile.
filamentosum, Eckl. & Zeyh. Enum. 311 = sarmentosum.
filamentosum, *Linn. Syst.* ed. X. 1060.—Afr. austr.
filicaule, *Haw. Misc. Nat.* 87.—Afr. austr.
filiforme, *Thunb. Prod. Pl. Cap.* 89.—Afr. austr.
fimbriatum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 393.—Afr. austr.
fissoides, Haw. Obs. Mesemb. 135 = obtusum.
fissum, *Haw. l. c.* 134.—Afr. austr.
flaccidum, *Jacq. Hort. Vindob.* t. 475.—Afr. austr.
flavum, *Haw. Rev. Pl. Succ.* 183.—Afr. austr.
flexifolium, Eckl. & Zeyh. Enum. 316 = micranthum.
flexifolium, Haw. Suppl. Pl. Succ. 98 = curvifolium.
flexile, Haw. Rev. Pl. Succ. 148 = polyanthon.
flexuosum, Eckl. & Zeyh. Enum. 315 = elegans.
flexuosum, *Haw. Misc. Nat.* 61.—Afr. austr.
floribundum, *Haw. l. c.* 100.—Afr. austr.
foliosum, Eckl. & Zeyh. Enum. 314 = multiflorum.
foliosum, Haw. Misc. Nat. 97 = tumidulum.
forficatum, Jacq. Hort. Vindob. i. 9. t. 26 = mutabile.
forficatum, *Linn. Syst.* ed. X. 1060.—Afr. austr.
formosum, *Haw. Rev. Pl. Succ.* 145.—Afr. austr.
Forskählei, *Hochst. ex Boiss. Fl. Orient.* ii. 765.—Mare rubr.
fragrans, *Salm-Dyck, Obs.* (1820) 8.—Afr. austr.
fulvum, Eckl. & Zeyh. Enum. 315 = simile.
fulvum, *Haw. Rev. Pl. Succ.* 180.—Afr. austr.
furfureum, Haw. in Phil. Mag. N. S. x. (1831) 421 = floribundum.
fuscum, Hort. ex Steud. Nom. ed. I. 526; ed. II. ii. 132 (Quid?).—Afr. austr.
fusiforme, Haw. Misc. Nat. 41 = elongatum.
geminatum, *Haw. l. c.* 92.—Afr. austr.
geminatum, Jacq. Fragment. 43 = geminiflorum.
geminiflorum, Eckl. & Zeyh. Enum. 310 = testaceum.
geminiflorum, *Haw. Rev. Pl. Succ.* 114; *Lodd. Bot. Cab.* t. 1160.—Afr. austr.
geniculiflorum, Forsk. Fl. Aegypt. Arab. 98 = Forskählei.
geniculiflorum, *Linn. Sp. Pl.* 481.—Afr. austr.
gibbosum, *Haw. Obs. Mesemb.* 137.—Afr. austr.
glabrum, Haw. ex Steud. Nom. ed. I. 526; ed. II. ii. 132 = pomeridianum.

MESEMBRYANTHEMUM:—
glabrum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 193.—Afr. austr.
glaciale, Haw. Suppl. Pl. Succ. 92 = crystallinum.
gladiatum, Jacq. Hort. Schoenb. iv. 19 = lacerum.
glaucescens, Haw. Syn. Pl. Succ. 236 = aequilaterale.
glaucinum, Eckl. & Zeyh. Enum. 311 = congestum.
glaucinum, Haw. Suppl. Pl. Succ. 97 = mutabile.
glaucoides, Haw. Obs. Mesemb. 330 = aurantiacum.
glaucum, Eckl. & Zeyh. Enum. 312 = croceum.
glaucum, *Linn. Sp. Pl.* 486.—Afr. austr.
glomeratum. *Linn. Syst.* ed. X. 1060.—Afr. austr.
gracile, Eckl. & Zeyh. Enum. 312 = bracteatum, patulum.
gracile, *Haw. Rev. Pl. Succ.* 144.—Afr. austr.
gramineum, Haw. Misc. Nat. 55 = pyropeum.
grandiflorum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 328.—Afr. austr.
graniforme, *Haw. Misc. Nat.* 82.—Afr. austr.
granulicaule, *Haw. in Till. Phil. Mag.* lxiv. (1824) 427.—Afr. austr.
grossum, *Haw. Misc. Nat.* 56.—Afr. austr.
grossum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 191.—Afr. austr.
guianense, *Klotzsch, in Schomb. Gui.* ii. 300; *ex Walp. Ann.* i. 326.—Guian.
hamatum, Willd. ex DC. Prod. iii. 432 = vaginatum.
Haworthii, *Donn, Cat. Hort. Cantab.* ed. II. 66.—Afr. austr.
Haworthii, Eckl. & Zeyh. Enum. 314 = Zeyheri.
helianthoides, DC. Pl. Grass. t. 135 = Candollii.
helianthoides, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 193.—Afr. austr.
Hermanni, *Haw. Obs. Mesemb.* 429.—Afr. austr.
heteropetalum, Eckl. & Zeyh. Enum. 311 = congestum.
heteropetalum, *Haw. Misc. Nat.* 67.—Afr. austr.
heterophyllum, Andr. Bot. Rep. t. 540 = angustum, praepingue.
heterophyllum, Haw. Obs. Mesemb. 420 = angustum.
hexaphyllum, Haw. Rev. Pl. Succ. 165 = Aizoon sarmentosum.
hirsutum, *Haw. Obs. Mesemb.* 297.—Afr. austr.
hirtellum, Eckl. & Zeyh. Enum. 319 = striatum.
hirtellum, *Haw. Obs. Mesemb.* 284.—Afr. austr.
hirtellum, Salm-Dyck, Obs. 24 = candens.
hirtum, *N. E. Br. in Gard. Chron.* (1878) II. 139.—Afr. austr.
hispidifolium, Steud. Nom. ed. II. ii. 133 = striatum.
hispidum, *Linn. Sp. Pl.* 482.—Afr. austr.
hispifolium, Haw. Suppl. Pl. Succ. 94 = striatum.
horizontale, Haw. Syn. Pl. Succ. 261 = clavatum.
humifusum, *Lee, ex Haw. Obs. Mesemb.* 420 (an seq.?)—Hab.?
humifusum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 179.—Afr. austr.
humile, *Haw. Misc. Nat.* 80.—Afr. austr.
hybridum, *Haw. Syn. Pl. Succ.* 218.—Afr. austr.
imbricans, Haw. Suppl. Pl. Succ. 94 = polyanthon.
imbricatum, Eckl. & Zeyh. Enum. 314 = curtum.
imbricatum, Haw. Obs. Mesemb. 317 = multiflorum.
inaequale, *Haw. Syn. Pl. Succ.* 266.—Afr. austr.
inclaudens, *Haw. l. c.* 295.—Afr. austr.
incomptum, *Haw. Suppl. Pl. Succ.* 96.—Afr. austr.
inconspicuum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 128.—Afr. austr.
incurvum, Eckl. & Zeyh. Enum. 313 = elegans.
incurvum, *Haw. Misc. Nat.* 69.—Afr. austr.
incurvum, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 416 = patulum.
inflexum, *Haw. Suppl. Pl. Succ.* 98.—Afr. austr.
insititium, Willd. Enum. Hort. Berol. 536 = croceum.
intonsum, Eckl. & Zeyh. Enum. 320 = bulbosum.
intonsum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 62.—Afr. austr.
introrsum, Hook. f. Bot. Mag. t. 6057, sphalm. = praec.
junceum, *Haw. Misc. Nat.* 59.—Afr. austr.
lacerum, Eckl. & Zeyh. Enum. 311 = anceps.
lacerum, *Haw. Obs. Mesemb.* 383.—Afr. austr.
laeve, Haw. Misc. Nat. 64 = coralliflorum.
laeve, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 187.—Afr. austr.
laeve, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 16 = Thunbergii.
laevigatum, Haw. Misc. Nat. 76 = acinaciforme.
lanceolatum, *Haw. l. c.* 45.—Afr. austr.
lanceum, Eckl. & Zeyh. Enum. 322 = Ecklonis.
lanceum, *Thunb. Prod. Pl. Cap.* 89.—Afr. austr.
lateriflorum, DC. Pl. Grass. t. 164 = brevifolium.

MESEMBRYANTHEMUM:—
latum, *Haw. Obs. Mesemb.* 186.—Afr. austr.
laurifolium, *Haw. l. c.* 478, nomen.—Afr. austr.
laxum, *Willd. Enum. Hort. Berol.* 536.—Afr. austr.
Lehmanni, *Eckl. & Zeyh. Enum.* 310.—Afr. austr.
lepidum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 130. —Afr. austr.
leptaleum, Haw. Rev. Pl. Succ. 140 = elegans.
limpidum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 183 = criniflorum, pyropeum.
lineare, Thunb. Prod. Pl. Cap. 88 = pyropeum.
lineolatum, *Haw. Rev. Pl. Succ.* 130.—Afr. austr.
lingueforme, Haw. Obs. Mesemb. 182 = obliquum, *Willd.*
linguiforme, DC. Pl. Grass. t. 71 = longum.
linguiforme, *Linn. Sp. Pl.* 488.—Afr. austr.
linguiforme, Linn. l. c., partim = latum, obliquum *Willd.*, longum.
longispinulum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 426.—Afr. austr.
longistylum, *DC. Pl. Grass.* t. 147.—Afr. austr.
longistylum, Eckl. & Zeyh. Enum. 315 = elegans.
longum, *Haw. Obs. Mesemb.* 177.—Afr. austr.
loratum, Haw. Rev. Pl. Succ. 168 = pallens.
loreum, *Haw. Misc. Nat.* 38.—Afr. austr.
loreum, *Linn. Sp. Pl.* 486.—Afr. austr.
lucidum, Haw. Rev. Pl. Succ. 95 = linguiforme.
lucidum, *Mill. Gard. Dict.* ed. VIII. n. 43.—Afr. austr.
lunatum, *Willd. Enum. Hort. Berol.* 538.—Afr. austr.
lupinum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 111.— Afr. austr.
luteolum, *Haw. l. c.* lxviii. (1826) 129.—Afr. austr.
luteo-viride, *Haw. Syn. Pl. Succ.* 226.—Afr. austr.
luteum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 128.— Afr. austr.
macrorhizum, Haw. l. c. 332 = megarhizum.
macrorrhizum, *DC. Prod.* iii. 425.—Ias. Borbon.
maculatum, *Haw. Obs. Mesemb.* 270.—Afr. austr.
magnipunctatum, *Haw. Rev. Pl. Succ.* 86.—Afr. austr.
marginatum, *Haw. Obs. Mesemb.* 412.—Afr. austr.
maximum, *Haw. l. c.* 402.—Afr. austr.
medium, *Haw. Suppl. Pl. Succ.* 88.—Afr. austr.
megarhizum, *G. Don, Gen. Syst.* iii. 145.—Afr. austr.
micans, *Linn. Sp. Pl.* 485.—Afr. austr.
micans, Thunb. Fl. Cap. 426 = collinum.
micranthum, Eckl. & Zeyh. Enum. 316 = bicorne.
micranthum, *Haw. Syn. Pl. Succ.* 257.—Afr. austr.
micropetalum, Pers. Syn. ii. 43 = apetalum.
microphyllum, *Haw. Obs. Mesemb.* 417; *Misc. Nat.* 73.—Afr. austr.
Milleri, Willd. Enum. Hort. Berol. Suppl. 37 = lacerum.
minimum, *Haw. Obs. Mesemb.* 126.—Afr. austr.
minutum, *Haw. Misc. Nat.* 21.—Afr. austr.
molle, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 192.— Afr. austr.
moniliforme, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 7.—Afr. austr.
monticolum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 428. —Afr. austr.
mucosum, *Haw. Obs. Mesemb.* 429.—Afr. austr.
mucronatum, *Haw. Misc. Nat.* 73.—Afr. austr.
mucroniferum, *Haw. in Till. Phil. Mag.* lxii. (1823) 381.—Afr. austr.
mucroniflorum, Harv. & Sond. Fl. Cap. ii. 615 = praec.
multiceps, *Salm-Dyck, Monog.* iii. § 6. f. 2.—Afr. austr.
multiflorum, Eckl. & Zeyh. Enum. 314 = Schollii.
multiflorum, *Haw. Obs. Mesemb.* 318.—Afr. austr.
multipunctatum, *Salm-Dyck, Hort. Dyck.* 357.—Afr. austr.
multiradiatum, Jacq. Fragm. 44 = roseum.
muricatum, *Haw. Misc. Nat.* 75.—Afr. austr.
murinum, *Haw. Obs. Mesemb.* 165.—Afr. austr.
musculinum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 327.—Afr. austr.
mustelinum, *Salm-Dyck, Obs.* 9.—Afr. austr.
mutabile, *Haw. Obs. Mesemb.* 377.—Afr. austr.
namaquense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 399. —Afr. austr.
nigrescens, Benth. Fl. Austral. iii. 324, in syn. = aequilaterale.
nitidum, *Haw. Obs. Mesemb.* 412; *Misc. Nat.* 57.— Afr. austr.
nitidum, *Salm. ex Spreng. Nov. Prov.* 27.—Hab. ?
nobile, *Haw. in Till. Phil. Mag.* lxii. (1823) 381.— Afr. austr.

MESEMBRYANTHEMUM:—
noctiflorum, *Linn. Sp. Pl.* 481.—Afr. austr.
nodiflorum, *Linn. l. c.* 480.—Reg. Mediterr.; Persia.
notabile, *Link, Enum. Hort. Berol.* ii. 44.—Afr. austr.
nuciforme, Haw. Obs. Mesemb. 129 = minutum.
obconellum, *Haw. Misc. Nat.* 21.—Afr. austr.
obcordellum, *Haw. l. c.*—Afr. austr.
obliquum, Eckl. & Zeyh. Enum. 318 = flavum.
obliquum, *Haw. Rev. Pl. Succ.* 183.—Afr. austr.
obliquum, *Willd. Sp. Pl.* ii. 1027.—Afr. austr.
obsubulatum, *Haw. Misc. Nat.* 26.—Afr. austr.
obtusum, *Haw. l. c.* 25.—Afr. austr.
octophyllum, Haw. Rev. Pl. Succ. 85 = testiculatum.
oligandrum, *Kunze, Del. Sem. Hort. Lips.* (1845).—*Cf. Linnaea*, xix. (1847) 405.—Afr. austr.
orbiculare, Page, Prod. Southamp. Bot. Gard. 169 = fibuliforme, minimum, minutum, obconellum, obcordellum, truncatellum.
ovatum, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 8 = angulatum.
pallens, Jacq. Hort. Schoenb. iii. t. 279 = longistylum.
pallens, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 182.— Afr. austr.
pallescens, Haw. Rev. Pl. Succ. 174 = longistylum.
papuliferum, *DC. Prod.* iii. 448.—Afr. austr.
papulosum, Eckl. & Zeyh. Enum. 322 = puberulum.
papulosum, *Linn. f. Suppl.* 259.—Afr. austr.
parviflorum, *Haw. Misc. Nat.* 95.—Afr. austr.
parviflorum, Jacq. Hort. Schoenb. iii. 15 = micranthum.
parvifolium, Eckl. & Zeyh. Enum. 318 = brevifolium.
parvifolium, *Haw. Rev. Pl. Succ.* 184.—Afr. austr.
parvifolium, Lam. Encyc. ii. 482 = violaceum.
patens, Willd. Enum. Hort. Berol. Suppl. 37 = multiflorum.
patulum, *Haw. Syn. Pl. Succ.* 334.—Afr. austr.
perfoliatum, *Mill. Gard. Dict.* ed. VIII. n. 19; *Haw. Obs. Mesemb.* 307.—Afr. austr.
perpusillum, *Haw. Rev. Pl. Succ.* 82.—Afr. austr.
perviride, *Haw. Obs. Mesemb.* 136.—Afr. austr.
Petiveri, Haw. l. c. 478 = humile.
pilifolium, Haw. ex Link, Enum. Hort. Berol. ii. 48 = striatum, hirtellum.
pilosum, Haw. Misc. Nat. 47 = helianthoides.
pinnatifidum, *Linn. f. Suppl.* 260.—Afr. austr.; Chili.
pinnatum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) App. 15.—Afr. austr.
pisiforme, *Haw. Misc. Nat.* 23.—Afr. austr.
polyanthon, *Haw. Syn. Pl. Succ.* 270.—Afr. austr.
polyanthum, Eckl. & Zeyh. Enum. 317 = glomeratum.
polyphyllum, Haw. Rev. Pl. Succ. 141 = violaceum.
pomeridianum, *Linn. Sp. Pl.* ed. II. 698.—Afr. austr.
praecox, F. Muell. in Linnaea, xxv. (1852) 384 = aequilaterale.
praepingue, *Haw. Obs. Mesemb.* 179.—Afr. austr.
procumbens, *Haw. Rev. Pl. Succ.* 111.—Afr. austr.
productum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 425. —Afr. austr.
protractum, *Link, Enum. Hort. Berol.* ii. 41.—Afr. austr.
pruinosum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 17.—Afr. austr.
puberulum, *Haw. in Till. Phil. Mag.* N. S. x. (1831) 421.—Afr. austr.
pubescens, *Haw. Obs. Mesemb.* 138.—Afr. austr.
pugioniforme, DC. Pl. Grass. t. 72 = elongatum.
pugioniforme, Haw. Obs. Mesemb. 390 = capitatum.
pugioniforme, *Linn. Sp. Pl.* 488.—Afr. austr.
pulchellum, *Willd. Enum. Hort. Berol.* 537; *Haw. Misc. Nat.* 72.—Afr. austr.
pulverulentum, Eckl. & Zeyh. Enum. 318, ex parte = parvifolium.
pulverulentum, *Haw. Obs. Mesemb.* 265.—Afr. austr.
pulverulentum, Willd. Enum. Hort. Berol. 533 = mucroniferum.
punctatum, *Haw. Obs. Mesemb.* 411; *Rev. Pl. Succ.* 107.—Afr. austr.
puniceum, Jacq. Hort. Schoenb. iv. 20 = violaceum.
purpurascens, *Salm-Dyck, ex Link, Enum. Hort. Berol.* ii. 45, nomen; *Obs.* (1822) 28.—Afr. austr.
purpureo-album, *Haw. in Till. Phil. Mag.* lxviii. (1826) 329.—Afr. austr.
purpureo-croceum, Haw. Obs. Mesemb. 257 = croceum.
purpureum, *Fisch. ex Steud. Nom.* ed. I. 527; ed. II. ii. 134 (Quid ?).—Hab. ?
pusiforme, Sond. in Harv. & Sond. Fl. Cap. ii. 450, sphalm. = pisiforme.
pustulatum, *Haw. Suppl. Pl. Succ.* 88.—Afr. austr.
pygmaeum, *Haw. l. c.* 98.—Afr. austr.

MESEMBRYANTHEMUM:—
pyropeum, *Haw. Suppl. Pl. Succ.* 99.—Afr. austr.
quadrifidum, *Haw. Misc. Nat.* 28.—Afr. austr.
radiatum, *Haw. Obs. Mesemb.* 322.—Afr. austr.
radicans, *DC. Pl. Grass.* t. 171.—Afr. austr.
ramulosum, *Haw. Misc. Nat.* 29.—Afr. austr.
rapaceum, Eckl. & Zeyh. Enum. 320 = grossum, *Haw.*
rapaceum, *Jacq. Fragm.* 43. t. 52. f. 1.—Afr. austr.
rectum, *Haw. Rev. Pl. Succ.* 170.—Afr. austr.
recurvum, Haw. Suppl. Pl. Succ. 90 = Schollii.
recurvum, *Moench, Meth. Suppl.* 190.—Afr. austr.
reflexum, Haw. Obs. Mesemb. 213 = longistylum.
relaxatum, *Willd. Enum. Hort. Berol. Suppl.* 36.— Afr. austr.
repens, Salisb. Prod. 347 = reptans.
repens, Salm. ex Spreng. Syst. ii. 517 = canaliculatum.
reptans, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 185.— Afr. austr.
retroflexum, Haw. Misc. Nat. 71 = elegans.
rhomboideum, *Salm-Dyck, Monog.* iii. § 6. f. 5.—Afr. austr.
rigidicaule, Eckl. & Zeyh. Enum. 310 = unidens.
rigidicaule, *Haw. Rev. Pl. Succ.* 116.—Afr. austr.
rigidum, *Haw. Misc. Nat.* 95.—Afr. austr.
ringens, Linn. Sp. Pl. 487 = caninum, felinum.
ringens, Thunb. ex Harv. & Sond. Fl. Cap. ii. 397 = tigrinum.
robustum, *Haw. Misc. Nat.* 28.—Afr. austr.
roseum, *Willd. Enum. Hort. Berol.* 535.—Afr. austr.
Rossi, Haw. Rev. Pl. Succ. 120 = aequilaterale.
rostellatum, DC. Prod. iii. 430 = seq.
Rostellum, *Haw. Rev. Pl. Succ.* 123.—Afr. austr.
rostratoides, Haw. Obs. Mesemb. 154 = ramulosum, acutum.
rostratum, *Linn. Sp. Pl.* 486.—Afr. austr.
rubricaule, *Willd. Enum. Hort. Berol.* 536; *Haw. Misc. Nat.* 77.—Afr. austr.
rubrocinctum, Eckl. & Zeyh. Enum. 311 = acinaciforme.
rubrocinctum, *Haw. Syn. Pl. Succ.* 234.—Afr. trop.
rufescens, Haw. l. c. 221 = obliquum, *Willd.*
sabulosum, Eckl. & Zeyh. Enum. 321 = papulosum.
sabulosum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 17.—Afr. austr.
Salmii, *Haw. Suppl. Pl. Succ.* 89.—Afr. austr.
salmoneum, Eckl. & Zeyh. Enum. 320 = nitidum.
salmoneum, *Haw. Rev. Pl. Succ.* 76.—Afr. austr.
sarmentosum, *Haw. Syn. Pl. Succ.* 238.—Afr. austr.; Austral.
scabrum, Linn. Sp. Pl. 483 = linguiforme.
scabrum, Thunb. ex Sond. in Harv. & Sond. Fl. Cap. ii. 416 = radiatum.
scalpratum, Haw. Obs. Mesemb. 187 = linguiforme.
scapigerum, Eckl. & Zeyh. Enum. 309 = rhomboideum.
scapigerum, *Haw. in Till. Phil. Mag.* lxiv. (1824) 423. —Afr. austr.
Schlichtianum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 435.—Afr. austr.
Schollii, *Salm-Dyck, Obs.* (1820) 10.—Afr. austr.
semicylindricum, *Haw. Obs. Mesemb.* 238.—Afr. austr.
semidentatum, *Haw. Suppl. Pl. Succ.* 95.—Afr. austr.?
semidentatum, *Salm-Dyck, Obs.* (1820) 9.—Afr. austr.
serratifolium, Hort. ex Steud. Nom. ed. I. 527; ed. II. ii. 134 = serratum.
serratulum, Nichols. Dict. Gard. ii. 359, sphalm. = serrulatum.
serratum, *Linn. Sp. Pl.* 485.—Afr. austr.
serrulatum, *Haw. Misc. Nat.* 77.—Afr. austr.
sessile, *Thunb. in Nov. Act. Cur.* viii. (1791), *App.* 14. —Afr. austr.
sessiliflorum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 193.—Afr. austr.
setosum, Moench, Meth. Suppl. 193 = echinatum.
setuliferum, *N. E. Br. in Gard. Chron.* (1876) II. 8.— Afr. austr.
simile, Haw. Rev. Pl. Succ. 115 = sarmentosum.
simile, *Sond. in Harv. & Sond. Fl. Cap.* ii. 435.—Afr. austr.
sordidum, DC. ex Spreng. Syst. ii. 518 = spinuliferum.
spathulatum, Thunb. in Nov. Act. Nat. Cur. viii. (1791), App. 5 = criniflorum.
spatulatum, Haw. Misc. Nat. 46 = praec.
speciosum, *Haw. Obs. Mesemb.* 267.—Afr. austr.
spectabile, Eckl. & Zeyh. Enum. 312 = Thunbergii.
spectabile, *Haw. Obs. Mesemb.* 385.—Afr. austr.
spiniforme, *Haw. l. c.* 240.—Afr. austr.

MESEMBRYANTHEMUM:—
spinosum, *Linn. Sp. Pl.* 483.—Afr. austr.
spinuliferum, *Haw. Obs. Mesemb.* 206.—Afr. austr.
spinuliforme, Harv. & Sond. Fl. Cap. ii. 616 = praec.
splendens, Eckl. & Zeyh. Enum. 315 = aduncum.
splendens, *Linn. Sp. Pl.* 486.—Afr. austr.
stellatum, Haw. Obs. Mesemb. 416; Misc. Nat. 91 = gracile.
stellatum, *Mill. Gard. Dict.* ed. VIII. n. 14.—Afr. austr.
stelligerum, Haw. in Till. Phil. Mag. lxiv. (1824) 61 = barbatum.
stenum, *Haw. l. c.* x. (1831) 420.—Afr. austr.
stipulaceum, *Linn. Sp. Pl.* 484.—Afr. austr.
stramineum, Haw. Obs. Mesemb. 252 = clavatum, fulvum.
stramineum, Willd. Enum. Hort. Berol. 533 = tricolorum.
striatum, *Haw. Obs. Mesemb.* 280.—Afr. austr.
strictum, Eckl. & Zeyh. Enum. 312 = Dregeanum.
strictum, *Haw. Misc. Nat.* 82.—Afr. austr.
strumosum, *Haw. Rev. Pl. Succ.* 190.—Afr. austr.
suaveolens, *Jacq. Fragm.* 66.—Afr. austr.
subaduncum, Haw. Suppl. Pl. Succ. 96 = viride.
subalatoides, Steud. Nom. ed. II. ii. 134, sphalm. = *subulatoides*.
subalatum, Haw. Misc. Nat. 76 = acinaciforme.
subcompressum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 131.—Afr. austr.
subglobosum, Haw. Misc. Nat. 62 = brevifolium.
subhispidum, *Haw. in Till. Phil. Mag.* N. S. x. (1831) 421.—Afr. austr.
subincanum, *Haw. l. c.* lxiv. (1824) 427.—Afr. austr.
subrectum, Steud. Nom. ed. II. ii. 134 = surrectum.
subrostratum, Willd. Enum. Hort. Berol. 529 = acutum.
subulatoides, Haw. Obs. Mesemb. 141 = acutum.
subulatum, Mill. Gard. Dict. ed. VIII. n. 10 = bellidiflorum.
sulcatum, *Haw. Rev. Pl. Succ.* 173.—Afr. austr.
surrectum, *Haw. l. c.* 101.—Afr. austr.
Sutherlandii, *Hook. f. Bot. Mag.* t. 6299.—Afr. austr.
taurinum, *Haw. Syn. Pl. Succ.* 224.—Afr. austr.
tegens, *F. Muell. Fragm.* v. 157.—Austral.
tenellum, *Haw. Obs. Mesemb.* 315.—Afr. austr.
tenue, Eckl. & Zeyh. Enum. 316 = bicorne.
tenue, Haw. Rev. Pl. Succ. 175 = micranthum.
tenuiflorum, Jacq. Fragm. 44. t. 32. f. 3 = viridiflorum.
tenuifolium, Eckl. & Zeyh. Enum. 316 = coccineum.
tenuifolium, *Linn. Sp. Pl.* 484.—Afr. austr.
tenuifolium, Thunb.—Cf. Sond. in Harv. & Sond. Fl. Cap. ii. 429, 437, 439, 440 = productum, bicolorum, glomeratum, elegans.
teretifolium, *Haw. Syn. Pl. Succ.* 210.—Afr. austr.
teretiusculum, *Haw. Obs. Mesemb.* 410.—Afr. austr.
ternifolium, Thunb. ex Harv. & Sond. Fl. Cap. ii. 415 = gracile.
testaceum, Eckl. & Zeyh. Enum. 317 = subincanum.
testaceum, *Haw. Suppl. Pl. Succ.* 97.—Afr. austr.
testiculare, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 181.—Afr. austr.
testiculare, Thunb. in Nov. Act. Nat. Cur. viii. (1791) App. 6 = testiculatum.
testiculatum, *Jacq. Fragm.* 20. t. 12.—Afr. austr.
tetragonum, Eckl. & Zeyh. Enum. 315 = productum.
tetragonum, *Moench, Meth. Suppl.* 191.—Hab. ?
tetragonum, *Thunb. Prod. Pl. Cap.* 91.—Afr. austr.
tetrasepalum, *Regel, in Act. Hort. Petrop.* ii. (1873) 434.—Hab. ?
Theris, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 126 (Quid ?).—Malaya.
Thunbergii, Eckl. & Zeyh. Enum. 315, partim = elegans.
Thunbergii, *Haw. Misc. Nat.* 86.—Afr. austr.
tigrinum, *Haw. Obs. Mesemb.* 164.—Afr. austr.
torquatum, Haw. Rev. Pl. Succ. 187 = floribundum.
tortuosum, DC. Pl. Grass. t. 94 = expansum.
tortuosum, Eckl. & Zeyh. Enum. 322 = anatomicum.
tortuosum, *Linn. Sp. Pl.* 487.—Afr. austr.
trichotomum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 14.—Afr. austr.
tricolor, Jacq. Hort. Schoenb. iv. 20 = mutabile.
tricolor, Steud. Nom. ed. II. ii. 135 = tricolorum.
tricolor, Willd. Hort. Berol. i. 22 = pyropeum.
tricolorum, *Haw. Obs. Mesemb.* 233.—Afr. austr.
Tripolium, *Linn. Sp. Pl.* 481.—Afr. austr.
truncatulum, *Haw. Misc. Nat.* 22.—Afr. austr.

MESEMBRYANTHEMUM:—
truncatum, *Thunb. in Nov. Act. Nat. Cur.* viii. (1791) *App.* 5.—Afr. austr.
tuberculatum, DC. ex Pers. Syn. ii. 44 = striatum.
tuberculatum, Eckl. & Zeyh. Enum. 319 = floribundum.
tuberculatum, Mill. Gard. Dict. ed. VIII. n. 32 = rostratum.
tuberosum, *Linn. Sp. Pl.* 484.—Afr. austr.
tumidulum, *Haw. Syn. Pl. Succ.* 286.—Afr. austr.
turbinatum, *Jacq. Hort. Vindob.* t. 476.—Afr. austr.
turbiniforme, Haw. Rev. Pl. Succ. 84 = truncatum.
umbellatum, *Linn. Sp. Pl.* 481.—Afr. austr.
umbellatum, Thunb. ex Harv. & Sond. Fl. Cap. ii. 441 = tuberosum.
umbelliflorum, *Jacq. Hort. Schoenb.* iv. 39.—Afr. austr.
uncatum, *Salm-Dyck, Monog.* iii. § 8. f. 6.—Afr. austr.
uncinatum, Eckl. & Zeyh. Enum. 313 = tenellum.
uncinatum, *Linn. Sp. Pl.* 483.—Afr. austr.
uncinellum, *Salm-Dyck, ex Haw. Rev. Pl. Succ.* 125.—Afr. austr.
unidens, *Haw. in Till. Phil. Mag.* lxviii. (1826) 331.—Afr. austr.
uvaeforme, *Haw. Rev. Pl. Succ.* 84.—Afr. austr.
vaginatum, *Lam. Encyc.* ii. 478; *Haw. Misc. Nat.* 95.—Afr. austr.
validum, *Haw. in Till. Phil. Mag.* lxviii. (1826) 329.—Afr. austr.
variabile, *Haw. Misc. Nat.* 85.—Afr. austr.
varians, Haw. l. c. 51 = tortuosum.
verruculoides, *Sond. in Harv. & Sond. Fl. Cap.* ii. 428.—Afr. austr.
verruculatum, *Linn. Sp. Pl.* 486.—Afr. austr.
verruculatum, Thunb. ex Harv. & Sond. Fl. Cap. ii. 417 = aurantiacum, verruculoides.
versicolor, Eckl. & Zeyh. Enum. 317 = elegans.
versicolor, *Haw. Misc. Nat.* 70.—Afr. austr.
villosum, Eckl. & Zeyh. Enum. 332 = Ecklonis.
villosum, *Linn. Sp. Pl.* 483.—Afr. austr.
violaceum, *DC. Pl. Grass.* t. 84.—Afr. austr.
violaceum, Eckl. & Zeyh. Enum. 317 = polyanthum.
virens, *Haw. Rev. Pl. Succ.* 121.—Afr. austr.
virescens, Haw. Syn. Pl. Succ. 236 = aequilaterale.
virgatum, *Haw. Misc. Nat.* 88.—Afr. austr.
viride, *Haw. Obs. Mesemb.* 314.—Afr. austr.
viridiflorum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 196.—Afr. austr.
vittatum, *N. E. Br. in Gard. Chron.* (1876) II. 772.—Afr. austr.
Volckameri, Haw. Obs. Mesemb. 426 = lanceolatum.
vulpinum, Haw. Syn. Pl. Succ. 217 = caninum.
Zeyheri, *Salm-Dyck, Monog.* iv. § 40. f. 5.—Afr. austr.

MESEMBRYANTHES, Stokes, Bot. Mat. Med. iii. 127 (1812) = praec.

MESEMBRYANTHUS, Neck. Elem. ii. 81 (1790) = praec.

MESEMBRYUM, Adans. Fam. ii. 242 (1763) = praec.

MESICERA, Rafin. Neogenyt. 4. (1825) = **Habenaria**, Willd. (Orchid.).
Michauxii, Rafin. Fl. Tellur. ii. 39 = H. Michauxii.
quinqueseta, Rafin. Neogenyt. 4 = H. Michauxii.
repens, Rafin. l. c. = H. repens.

MESOCENTRON, Cass. in Dict. Sc. Nat. xliv. 38 (1826) = **Centaurea**, Linn. (Compos.).
eriophorum, Cass. l. c. = C. eriophora.

MESOCHLOA, Rafin. Fl. Tellur. iv. 11 (1836) = **Zephyranthes**, Herb. (Amaryll.).
canaliculata, Rafin. l. c. = Z. Mesochloa.

MESOCLASTES, Lindl. Gen. et Sp. Orch. 44 (1830) = **Luisia**, Gaudich. (Orchid.).
brachystachys, Lindl. l. c.; et in Wall. Cat. n. 1994 = L. brachystachys.
uniflora, Lindl. in Wall. Cat. n. 1993 (= *L. uniflora*).—Reg. Himal.

MESODACTYLIS, Wall. Pl. As. Rar. i. 74, in obs. (1830) = **Apostasia**, Blume (Orchid.).
deflexa, Wall. l. c. = A. Wallichii.
odorata, Endl. Gen. 221 = A. odorata.
Wallichii, Endl. l. c. = A. Wallichii.

MESODACTYLUS, Endl. Gen. 221 (1837) = praec.

MESODETRA, Rafin. Fl. Ludov. 141 (1817) = **Helenium**, Linn. (Compos.).
alata, Rafin. l. c. = H. quadridentatum.

MESODISCUS, Rafin. New Fl. Am. iv. 19 (1836) = **Cryptotaenia**, DC. (Umbell.).
proliferus, Rafin. l. c. 20 = C. canadensis.
simplex, Rafin. l. c. = C. canadensis.

MESOGRAMMA, DC. Prod. vi. 304 (1837) = **Senecio**, Tourn. (Compos.).
apiifolium, DC. l. c.—Afr. austr.

MESOLIGUS, Rafin. Fl. Tellur. ii. 44 (1836) = **Aster**, Tourn. (Compos.).
subulatus, Rafin. l. c. = A. subulatus.

MESOMELAENA, Nees, in Lehm. Pl. Preiss. ii. 88 (1846). *CYPERACEAE*, Benth. & Hook. f. iii. 1061.
GYMNOSCHOENUS, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 47.
PTILANTHELIUM, Steud. Syn. Pl. Cyp. 166 (1855).
anceps, *Benth. Fl. Austral.* vii. 380.—Austral.
bicapitata, Steud. Syn. Pl. Cyp. 163 = Caustis dioica.
deusta, *Benth. Fl. Austral.* vii. 379.—Austral.
Preissii, Nees, in Lehm. Pl. Preiss. ii. 88 = stygia.
sphaerocephala, *Benth. Fl. Austral.* vii. 380.—Austral.
stygia, *Nees, in Lehm. Pl. Preiss.* ii. 89.—Austral.
tetragona, *F. Muell. Fragm.* ix. 36.—Austral.

MESONA, Blume, Bijdr. 838 (1826). *LABIATAE*, Benth. & Hook. f. ii. 1172.
chinensis, *Benth. Fl. Hongk.* 274.—China.
palustris, *Blume, Bijdr.* 838.—Java.
Wallichiana, *Benth. in DC. Prod.* xii. 46.—Reg. Himal.

MESONEURIS, A. Gray, in Proc. Am. Acad. viii. (1873) 661 = **Senecio**, Tourn. (Compos.).
bipinnatifida, A. Gray, l. c. = S. cervariaefolius.

MESOPTERA, Hook. f. in Benth. & Hook. f. Gen. ii. 130 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Maingayi, *Hook. f. l. c.* 131.—Malacca.

MESOPTERA, Rafin. Fl. Tellur. iv. 49 (1836) = **Liparis**, Rich. (Orchid.).

MESOSETUM, Steud. in Flora, xxxiii. (1850) 228, nomen; Syn. Pl. Gram. 118 (1855) = **Panicum**, Linn. (Gramin.).
cayennense, Steud. ll. cc.—Guian.

MESOSPHAERUM, P. Br. Hist. Jamaic. 257 (1756) = **Hyptis**, Jacq. (Labiat.).

MESOSPINIDIUM, Reichb. f. in Bot. Zeit. x. (1852) 929 = **Odontoglossum**, H. B. & K. (Orchid.).
aurantiacum, Reichb. f. in Walp. Ann. vi. 857 = Ada aurantiaca.
Bowmani, Reichb. f. in Gard. Chron. (1869) 1182 (Bowmanni).—N. Granat.
cinnabarinum, Reichb. f. in Walp. Ann. vi. 858 = Brassia cinnabarina.
densiflorum, Reichb. f. in Gard. Chron. (1872) 393 = Cochlioda densiflora.
incantans, Reichb. f. l. c. (1880) I. 586.—N. Granat.
jucundum, Reichb. f. l. c. (1876) II. 580.—Bras.
radiatum, Reichb. f. in Walp. Ann. vi. 856 = Abola radiata.
roseum, Reichb. f. in Gard. Chron. (1872) 393; et Otia Bot. Hamb. 5, 23 = Cochlioda rosea (in Addend.).
sanguineum, Reichb. f. in Walp. Ann. vi. 858 = Cochlioda sanguinea (in Addend.).
vulcanicum, Reichb. f. in Gard. Chron. (1872) 393 (= Cochliodae sp.).—Peruv.
Wallisii, Reichb. f. in Linnaea, xli. (1877) 106.—Peruv.
Warscewiczii, Reichb. f. in Bot. Zeit. x. (1852) 929 = O. Warszewiczii.

MESOTRICHA, Stschegl. in Bull. Soc. Nat. Mosc. (1859) I. 9 = **Astroloma**, R. Br. (Epacrid.).
discolor, Stschegl. l. c. = A. longiflorum.
longiflora, Stschegl. l. c. = A. longiflorum.

MESPILODAPHNE, Nees, in Linnaea, viii. (1833) 45 = **Ocotea**, Aubl. (Laurin.).
acuminata, Baill. in Bull. Soc. Linn. Par. i. (1885) 448.—Madag.
attenuata, Meissn. in DC. Prod. xv. 1. 108.—Bras.
aurantiodora, Meissn. l. c. 101 = Nectandra aurantiodora.
baccans, Klotzsch & Karst. ex Meissn. l. c. 107.—N. Granat.
bahiensis, Meissn. l. c. 108.—Bras.
balanocarpa, Nees, in Linnaea, xxi. (1848) 495.—Peruv.
Bernieri, Baill. in Bull. Soc. Linn. Par. i. (1885) 453.—Madag.
borbonica, Meissn. in DC. Prod. xv. 1. 105.—Ins. Borbon.
bullata, Meissn. l. c. 511 = O. bullata.
ceanothifolia, Nees, Syst. Laurin. 236.—Bras.
complicata, Meissn. in DC. Prod. xv. 1. 103.—Bras.
cordata, Meissn. l. c. 101.—Bras.
corymbosa, Meissn. l. c. 98.—Bras.
cupularis, Meissn. l. c. 104.—Ins. Mascar.
cymosa, Meissn. l. c. 105 = *cupularis*.
fasciculata, Meissn. l. c. 108 = O. firmula.
foetens, Meissn. l. c. 511 = O. foetens.
Gardneri, Meissn. l. c. 99.—Bras.
glauca, Meissn. l. c. 97.—Bras.
heteromorpha, Meissn. l. c. 105 = *cupularis*.
indecora, Meissn. l. c. 102 = O. indecora.
lanata, Meissn. l. c.—Bras.
laxiflora, Meissn. l. c. 107.—Bras.
leucophloea, Nees, in Linnaea, viii. (1833) 45 = O. indecora.
Lindleyana, Meissn. in DC. Prod. xv. 1. 105 = *cupularis*.
madagascariensis, Meissn. l. c. 105.—Madag.
marginata, Meissn. l. c. 106 = *cupularis*.
mauritiana, Meissn. l. c. 104 = *cupularis*.
Maximilianea, Meissn. l. c. 108 = *prolifera*.
Meyendorffiana, Meissn. l. c. 99.—Bras.
microneura, Meissn. l. c. 98 = *prolifera*.
montana, Meissn. l. c. 98.—Bras.
moschata, Meissn. l. c. 106.—Peruv.; Bras.
nitida, Meissn. l. c.—Peruv.
nitidula, Meissn. l. c. 510 = Nectandra nitidula.
notata, Meissn. l. c. 98.—Bras.
novo-granatensis, Meissn. l. c. 107 = Nectandra concinna.
nummularia, Meissn. l. c. 101.—Bras.
nutans, Meissn. l. c. 98.—Bras.
oblonga, Meissn. l. c. 107.—Guian.
opifera, Meissn. l. c. 510 = O. opifera.
organensis, l. c. 97.—Bras.
parviflora, Meissn. l. c. 109.—Bras.
patens, Meissn. in Kjoeb. Vidensk. Meddel. (1870) 134.—Bras.
Persooniana, Meissn. in DC. Prod. xv. 1. 105.—Ins. Mascar.
petiolaris, Meissn. l. c. 99.—Bras.
phillyraeoides, Meissn. l. c. 100.—Bras.
Pohlii, Meissn. l. c. 97.—Bras.
pretiosa, Nees, in Linnaea, viii. (1833) 45 = O. pretiosa.
prolifera, Nees, l. c. 46.—Bras.
pulchella, Meissn. in DC. Prod. xv. 1. 99 = O. pulchella.
pyriformis, Nees, ex Meissn. l. c. 108.—Guian.
Riedelii, Meissn. l. c. 511.—Bras.
rigens, Meissn. l. c. 99, in obs. ad n. 11 = O. tristis.
Sassafras, Meissn. l. c. 102.—Bras.
Sprucei, Meissn. l. c. 97.—Bras.
trichophlebia, Baill. in Bull. Soc. Linn. Par. i. (1885) 448.—Madag.
tristis, Meissn. in DC. Prod. xv. 1. 100 = O. tristis.
vaccinioides, Meissn. l. c. = O. pulchella.
vaginans, Meissn. l. c. 104.—Venezuela.

MESPILOPHORA, Neck. Elem. ii. 73 (1790) = **Pyrus**, Tourn. (Rosac.).

MESPILUS, [Tourn.] Linn. Syst. ed. I (1735) = **Pyrus**, Tourn. etc. (Rosac.).
acerifolia, Poir. in Lam. Encyc. iv. 442 = Crataegus cordata.
acuminata, Lodd. Bot. Cab. t. 919 = Cotoneaster acuminata.
aestivalis, Walt. Fl. Carol. 148 = Crataegus aestivalis.
affinis, D. Don, Prod. Fl. Nep. 238 = Cotoneaster bacillaris.

MESPILUS:—
Amelanchier, Linn. Sp. Pl. 478 = Amelanchier vulgaris.
Amelanchier, Walt. Fl. Carol. 148 = Amelanchier canadensis.
anthyllidifolia, Spreng. Syst. ii. 508 = Osteomeles anthyllidifolia.
apiifolia, Medic. Gesch. Bot. 82 = Crataegus Oxyacantha.
apiifolia, Poir. Encyc. Suppl. iv. 68 = Crataegus apiifolia.
arborea, Michx. f. Hist. Arbr. Am. Sept. iii. 68. t. 11 = Amelanchier canadensis.
arborescens, C. Koch, Wochenschr. v. (1862) 380 = Crataegus arborescens.
arbutifolia, Link, Enum. Hort. Berol. ii. 236 = Photinia arbutifolia.
arbutifolia, Linn. Sp. Pl. 478 = Pyrus arbutifolia.
Aria, Scop. Fl. Carn. i. 345 = Pyrus Aria.
Aronia, Spach, Hist. Vég. Phan. ii. 69 = Crataegus Azarolus.
Aronia, Willd. Enum. Hort. Berol. Suppl. 35 = Crataegus orientalis.
atrofusca, C. Koch, Wochenschr. v. (1862) 400 = Crataegus pentagyna.
Aucuparia, [Web. in] Wigg. Prim. Fl. Holsat. 38; All. Fl. Pedem. ii. 142 = Pyrus Aucuparia.
axillaris, Pers. Syn. ii. 39 = Crataegus parvifolia.
Azarolus, All. Fl. Pedem. ii. 141 = Crataegus Azarolus.
Azarolus, Marsh, Arbust. 89 = Crataegus spathulata.
badiata, Bosc, Encyc. Agric. vii. 592 = Crataegus Crus-galli.
bengalensis, Roxb. Hort. Beng. 38; Fl. Ind. ii. 510 = Eriobotrya bengalensis.
berberifolia, C. Koch, Wochenschr. v. (1862) 383 = Crataegus berberifolia.
Bosciana, Spach, Hist. Vég. Phan. ii. 58 = Crataegus Crus-galli.
Calpodendron, Ehrh. Beitr. ii. 67 = Crataegus tomentosa.
calycina, Stokes, Bot. Mat. Med. iii. 107 = Pyrus germanica.
canadensis, Linn. Sp. Pl. 478 = Amelanchier canadensis, Pyrus arbutifolia.
caroliniana, Poir. in Lam. Encyc. iv. 442 = Crataegus flava.
caroliniana, Wendl. f. in Flora, vi. (1823) 701 = = Crataegus tomentosa.
caucasica, C. Koch, Wochenschr. v. (1862) 405 = Crataegus Oxyacantha.
Celsiana, Dum.-Cours. Bot. Cult. Suppl. 286 = Crataegus nigra.
Chamaemespilus, Linn. Sp. Pl. 479 = Pyrus Chamaemespilus.
chinensis, Blume, Bijdr. 1102 = Crataegus chinensis.
coccinea, Marsh. Arbust. 87 = Crataegus coccinea.
coccinea, Waldst. & Kit. Pl. Rar. Hung. iii. 284 = Cotoneaster tomentosa.
communis, Gueldenst. Reisen, i. 419 = Pyrus germanica.
constantinopolitana, Hort. ex Poir. Encyc. Suppl. iv. 69 = Crataegus coccinea, C. Azarolus.
corallina, Desf. Tabl. ed. I. 174 = Crataegus cordata.
cordata, Willd. Enum. Hort. Berol. 523 = Crataegus cordata.
cordifolia, Mill. Gard. Dict. ed. VIII. n. 4, errata = Crataegus cordata.
cornifolia, Muench. Hausv. v. 145 = Crataegus tomentosa.
Cosansasi, C. Koch, Wochenschr. v. (1862) 396 = Crataegus sanguinea.
Cotoneaster, Linn. Sp. Pl. 479 = Cotoneaster integerrima.
Crataegus, Borckh. Beschr. Hess. Hoelz. 190; ex Roem. Arch. i. III. 86 = Crataegus Oxyacantha.
crenulata, D. Don, Prod. Fl. Nep. 238 = Crataegus crenulata.
Crus-galli, Marsh. Arbust. 88; Poir. in Lam. Encyc. iv. 441 = Crataegus Crus-galli.
Cuila, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 238 = Eriobotrya elliptica.
cuneata, C. Koch, Wochenschr. v. (1862) 388 = Crataegus cuneata.
cuneifolia, Ehrh. Beitr. iii. 21 = Crataegus tomentosa.
cuneifolia, Moench, Meth. 684 = Craetaegus Crus-galli.
cuneiformis, Marsh. Arbust. 88 = Crataegus Crus-galli.
denticulata, Spreng. Syst. ii. 505 = Cotoneaster denticulata.
depressa, C. Koch, Wochenschr. v. (1862) 365 = Crataegus depressa.

MESPILUS:—
digyna, S. F. Gray, Nat. Arr. Brit. Pl. ii. 565 = Crataegus Oxyacantha.
dissecta, Dum.-Cours. Bot. Cult. ed. II. v. 454 = Crataegus Oxyacantha.
diversifolia, Poir. Encyc. Suppl. iv. 73 = Crataegus Oxyacantha.
domestica, All. Fl. Pedem. ii. 142 = Pyrus Sorbus.
domestica, Gater. Pl. Montaub. 92; et ex DC. Fl. Fr. iv. 434 = Pyrus germanica.
elegans, Poir. in Lam. Encyc. iv. 439 = Crataegus Oxyacantha.
elliptica, Guimp. Otto & Hayne, Fremd. Holz. t. 144 = Crataegus Crus-galli.
elliptica, Lam. Encyc. iv. 447 = Crataegus flava.
eriocarpa, DC. Fl. Fr. Suppl. 532 = Cotoneaster tomentosa.
eriocarpa, Desf. Cat. Hort. Par. ed. III. 409 = Cotoneaster integerrima.
ferruginea, Poir. Encyc. Suppl. iv. 73; Spreng. Syst. ii. 506 = Eriobotrya cordata.
fissa, Poir. l. c. 72 = Crataegus Oxyacantha.
flabellata, Bosc, Encyc. Agric. vii. 591 = Crataegus coccinea.
flava, Willd. Enum. Hort. Berol. 523 = Crataegus flava.
flavescens, Bosc, Encyc. Agric. vii. 591 = Crataegus flavescens.
flexispina, C. Koch, Dendrol. i. 139 = Crataegus flava.
flexispina, Moench, Verz. 62 = Crataegus parvifolia.
flexuosa, Bosc, Encyc. Agric. vii. 591 = Crataegus tomentosa.
flexuosa, Poir. Encyc. Suppl. iv. 73 = Crataegus parvifolia.
florentina, Bertol. Amoen. Ital. 29 = Pyrus crataegifolia.
Fontanesiana, Spach, in Ann. Sc. Nat. Sér. II. iii. (1835) 105 = Crataegus Crus-galli.
germanica, Linn. Sp. Pl. 478 = Pyrus germanica.
glabra, Poir. in Lam. Encyc. iv. 446 = Photinia serrulata, *Lindl.*
glabrata, Spreng. Syst. ii. 506 = Osteomeles glabrata.
glandulosa, Willd. Enum. Hort. Berol. 523 = Crataegus coccinea.
grandiflora, Sm. Exot. Bot. i. 33 = Crataegus grandiflora.
heterophylla, Poir. Encyc. Suppl. iv. 68 = Crataegus Azarolus.
heterophylla, Ruiz & Pav. Fl. Per. iv. t. 425 = Eriobotrya heterophylla.
hyemalis, Walt. Fl. Carol. 148 = Crataegus flava.
indica, Poir. Encyc. Suppl. iv. 73 = Raphiolepis indica.
Insegnae, Tineo, ex Guss. Fl. Sic. Syn. ii. 830 = Crataegus Insegnae.
intermedia, Poir. Encyc. Suppl. iv. 68 = Pyrus intermedia.
japonica, Thunb. Fl. Jap. 206 = Eriobotrya japonica.
laciniata, Guss. Fl. Sic. Prod. i. 565 = Crataegus laciniata.
laciniata, Walt. Fl. Carol. 147 = Crataegus parvifolia.
laevigata, Poir. in Lam. Encyc. iv. 439 = Crataegus Oxyacantha.
laevis, Poir. in Lam. Encyc. iv. 445 = Photinia laevis.
lanuginosa, Ruiz & Pav. Fl. Per. iv. t. 425 = Osteomeles latifolia.
latifolia, Hort. ex Poir. in Lam. Encyc. iv. 444 = Crataegus tomentosa.
leucophloeos, C. Koch, Dendrol. i. 136 = Crataegus tomentosa.
linearis, Desf. Hist. Arbr. ii. 156 = Crataegus Crus-galli.
lobata, Poir. Encyc. Suppl. iv. 71 = Crataegus tomentosa.
Loddigesiana, Spach, in Ann. Sc. Nat. Sér. II. iii. (1835) 105 = Crataegus mexicana.
Loureiri, Kostel, Allg. Med. Pharm. Fl. iv. 1479 = Crataegus Pyracantha.
lucida, Ehrh. Beitr. iv. 17 = Crataegus Crus-galli.
lutea, Poir. Encyc. Suppl. iv. 72 = Crataegus tomentosa.
macracantha, Wenzig, in Linnaea, xxxviii. (1874) 137 = Crataegus sanguinea.
macrocarpa, DC. ex E. Morr. Belg. Hortic. vi. (1856) 256. t. 63 = Pyrus germanica.
maroccana, Poir. Encyc. Suppl. iv. 74 = Crataegus Oxyacantha.
maura, Poir. l. c. 73 = Crataegus Oxyacantha.
maxima, Dum.-Cours. Bot. Cult. ed. II. v. 451 = Crataegus Crus-galli.
melanocarpa, Bess. Cat. Hort. Krzem. (1814) 88; Poir. Encyc. Suppl. iv. 68 = Crataegus pentagyna.

MESPILUS :—
melanocarpa, Fisch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 48 = Cotoneaster integerrima.
mexicana, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. iii. t. 300; C. Koch, Wochenschr. v. (1862) 363 = Crataegus mexicana.
Michauxii, Hornem. Hort. Hafn. 455 = Crataegus flava.
microphylla, C. Koch, Wochenschr. v. (1862) 405 = Crataegus Oxyacantha.
monogyna, All. Fl. Pedem. ii. 141 = Crataegus Oxyacantha.
montana, Hort. ex M. Roem. Syn. Rosifl. 161 = Pyrus arbutifolia.
myrtifolia, C. Koch, Wochenschr. v. (1862) 365 = Crataegus myrtifolia.
nana, Dum.-Cours. Bot. Cult. ed. II. vii. 286 = Crataegus Crus-galli.
nebrodensis, Bertol. Fl. Ital. vii. 629 = Cotoneaster nebrodensis.
nigra, Willd. Enum. Hort. Berol. 524 = Crataegus Oxyacantha.
nivea, Marsh. Arbust. 90 = Amelanchier canadensis.
obovata, Bosc, ex Steud. Nom. ed. I. 528, sphalm. = *odorata*.
obtusifolia, Herb. Juss. ex Poir. Encyc. Suppl. iv. 73 = Osteomeles glabrata.
odorata, Wendl. f. in Flora, vi. (1823) 700 = Crataegus tomentosa.
odoratissima, Andr. Bot. Rep. t. 590 = Crataegus tanacetifolia.
Oliveriana, Bosc, Encyc. Agric. vii. 591; Dum.-Cours. Bot. Cult. ed. II. v. 454 = Crataegus pentagyna.
Oliveriana, Poir. Encyc. Suppl. iv. 72 = Crataegus Oxyacantha.
orientalis, Habl. Taur. 108; ex Bieb. Fl. Taur. Cauc. i. 387 = Crataegus tanacetifolia.
orientalis, Mill. Gard. Dict. ed. VIII. n. 9 = Cotoneaster tomentosa.
orientalis, Poir. Encyc. Suppl. iv. 72 = Crataegus orientalis.
ovalifolia, Hornem. Hort. Hafn. Suppl. 52 = Crataegus Crus-galli.
oxyacantha, Crantz, Stirp. Austr. ed. I. fasc. ii. 39; ed. II. 82* = Crataegus Oxyacantha.
oxyacanthoides, DC. Fl. Fr. iv. 433 = Crataegus Oxyacantha.
parvifolia, Willd. Enum. Hort. Berol. 523 = Crataegus parvifolia.
pauciflora, Poir. in Lam. Encyc. iv. 441 = Crataegus Pyracantha.
pectinata, Dum.-Cours. Bot. Cult. ed. II. v. 454 = Crataegus pectinata.
pentagyna, Spreng. Syst. ii. 507 = Crataegus pentagyna.
Persoonii, Spreng. l. c. 506 = Osteomeles glabrata.
peruviana, C. Koch, Wochenschr. v. (1862) 365 = Crataegus peruviana (in Addend.).
phaenopyrum, Ehrh. Beitr. ii. 67 = Crataegus cordata.
pinnata, Dum.-Cours. ex Steud. Nom. ed. I. 529 = Crataegus tanacetifolia.
pinnatifida, C. Koch, in Wochenschr. v. (1862) 398 = Crataegus pentagyna.
pirifolia, Desf. Hist. Arbr. ii. 155 = Crataegus tomentosa.
Poiretiana, Sweet, Hort. Brit. ed. II. 176 = Crataegus Crus-galli.
polyacantha, Guss. Fl. Sic. Prod. Suppl. 154 = Crataegus Oxyacantha.
populifolia, Poir. in Lam. Encyc. iv. 447 = Crataegus coccinea.
pruinosa, Wendl. f. in Flora, vi. (1823) 701 = Crataegus tomentosa.
prunellifolia, Bosc, ex Poir. Encyc. Suppl. iv. 72 = Crataegus Crus-galli.
prunifolia, Marsh. Arbust. 90; Poir. in Lam. Encyc. iv. 443 = Crataegus Crus-galli.
pubescens, H. B. & K. Nov. Gen. et Sp. vi. 213. t. 555 = Crataegus pubescens.
pubescens, Presl, Delic. Prag. 53 = Crataegus tanacetifolia.
pubescens, Wendl. f. in Flora, vi. (1823) 700 = Crataegus coccinea.
pumila, Schmidt, ex Steud. Nom. ed. II. ii. 136 = Pyrus arbutifolia.
punctata, Willd. Enum. Hort. Berol. 524 = Crataegus tomentosa.
purpurea, Poir. Encyc. Suppl. iv. 73 = Crataegus sanguinea.

MESPILUS :—
pycnoloba, C. Koch, Wochenschr. v. (1862) 390 = Crataegus pycnoloba.
pygmaea, Baumg. Enum. Stirp. Transs. ii. 40 = Amelanchier vulgaris.
Pyracantha, Linn. Sp. Pl. 478 = Crataegus Pyracantha.
pyracanthifolia, Sweet, Hort. Brit. ed. II. 175 = Crataegus Crus-galli.
pyrifolia, Willd. Enum. Hort. Berol. 523 = Crataegus grandiflora.
pyriformis, Hort. ex M. Roem. Syn. Rosifl. 156 = Crataegus coccinea.
quinqueloba, Bosc, Encyc. Agric. vii. 591.—Hab.?
quitensis, C. Koch, Wochenschr. v. (1862) 364 = Crataegus rivularis.
racemiflora, Desf. Cat. Hort. Par. ed. III. 409 = Crata gus Crus-galli.
rivularis, C. Koch, Wochenschr. v. (1862) 372 = Crataegus rivularis.
rotundifolia, Ehrh. Beitr. iii. 28 = Crataegus coccinea.
rubra, Stokes, Bot. Mat. Med. iii. 110 = Raphiolepis indica.
salicifolia, Hort. ex C. Koch, Dendrol. i. 144 = Crataegus Crus-galli.
sanguinea, Spach, Hist. Vég. Phan. ii. 62 = Crataegus sanguinea.
Sieboldi, Blume, Bijdr. 1102 = Raphiolepis japonica.
sinaica, C. Koch, Wochenschr. v. (1862) 391 = Crataegus sinaica.
sinensis, Poir. Encyc. Suppl. iv. 70 = Raphiolepis indica.
Smithii, DC. Prod. ii. 633 = Crataegus grandiflora.
sorbifolia, Hort. ex Poir. Encyc. Suppl. iv. 72 = Pyrus spuria.
spathulata, Poir. l. c. 68 = Crataegus spathulata.
spectabilis, Hort. ex C. Koch, Hort. Dendrol. 182 = Pyrus spectabilis.
spiralis, Blume, Bijdr. 1102 = Rhaphiolepis indica.
stipulacea, Desf. ex Steud. Nom. ed. II. i. 434 = Crataegus mexicana.
stipulosa, H. B. & K. Nov. Gen. et Sp. vi. 213 = Crataegus stipulosa.
subspinosa, Vent. ex DC. Prod. ii. 626 = Crataegus subspinosa.
succulenta, Wenzig, in Linnaea, xxxviii. (1874) 136 = Crataegus sanguinea.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 1 = Pyrus germanica.
tanacetifolia, Poir. in Lam. Encyc. iv. 440 = Crataegus tanacetifolia.
tiliaefolia, C. Koch, Wochenschr. v. (1862) 375 = Crataegus subvillosa.
tinctoria, D. Don, Prod. Fl. Nep. 238 = Eriobotrya bengalensis.
tomentosa, Ait. Hort. Kew. ed. I. ii. 174 = Cotoneaster tomentosa.
tomentosa, Hohen. Enum. Pl. Elisabethpol. 235 = Cotoneaster nummularia.
tomentosa, Poir. in Lam. Encyc. iv. 440 = Crataegus parvifolia.
torminalis, [Web. in] Wigg. Prim. Fl. Holsat. 38; All. Fl. Pedem. ii. 141 = Pyrus torminalis.
Trewiana, Tausch, in Flora, xxi. (1838) 716 = Crataegus Crus-galli.
trifoliata, Bosc, Encyc. Agric. vii. 591 = Crataegus trifoliata.
triloba, Poir. Voy. Barb. ii. 171 = Crataegus Oxyacantha.
turbinata, Spreng. Syst. ii. 506 = Crataegus flava.
uniflora, C. Koch, Wochenschr. v. (1862) 383 = Crataegus parvifolia.
unilateralis, Poir. Encyc. Suppl. iv. 73 = Crataegus parvifolia.
villosa, Poir. l. c. 445 = Photinia villosa.
virginiana, Mill. Gard. Dict. ed. VIII. n. 11 = Crataegus flava?
viridis, Sweet, Hort. Brit. ed. I. 134 = Crataegus coccinea.
vulgaris, Reichb. Fl. Germ. Excurs. 630 = Pyrus germanica.
Watsoniana, Spach, Hist. Vég. Phan. ii. 57 = Crataegus Crus-galli.
Wendlandii, Opiz, in Flora, xvii. (1834) 590 = Crataegus coccinea.
xalapensis, C. Koch, Wochenschr. v. (1862) 364.—Mexic.
xanthocarpa, Linn. f. Suppl. 254; Ehrh. Beitr. ii. 67 = Crataegus parvifolia.

MESSANTHEMUM, Pritz. Ic. Ind. ii. 185 (1866) sphalm. = **Mesanthemum**, Koern. (Eriocaul.).

MESSERSCHMIDIA, Murr. Syst. ed. XIII. 161 (1774) = seq.

MESSERSCHMIDTIA, G. Don, Gen. Syst. iv. 370 (1837) = seq.

MESSERSMIDIA, Linn. Hort. Upsal. 36 (1748) = **Tournefortia**, Linn. (Borag.).
angustifolia, Lam. Tabl. Encyc. ii. 415 = T. fruticosa, *Ker-Gawl.*
Argunia, Gaertn. Fruct. ii. 130. t. 109. f. 3 = T. sibirica.
Arguzia, Linn. Mant. i. 42 = T. sibirica.
Blanchetii, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 195 = T. Blanchetii.
cancellata, Asso, Syn. Pl. Arag. n. 162. t. 1. f. 1 = Rochelia stellulata.
candida, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 334 = T. candida.
candidula, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 202.—Bras.
chrysantha, Mart. & Gall. in Bull. Acad. Brux. xi. II. (1844) 334 = T. chrysantha.
corymbosa, G. Don, Gen. Syst. iv. 371 = T. foetidissima.
elegans, G. Don, l. c. = T. elegans.
ferruginea, G. Don, l. c. = T. scandens.
floribunda, G. Don, l. c. 370 = T. floribunda.
floribunda, Salisb. Prod. 112 = T. fruticosa, *Ker-Gawl.*
fruticosa, Linn. f. Suppl. 132 = T. fruticosa, *Ker-Gawl.*
hirsutissima, Roem. & Schult. Syst. iv. 541 = T. hirsutissima.
hispida, Benth. in Royle, Illustr. Bot. Himal. 306 = Heliotropium zeylanicum.
incana, G. F. W. Mey. Prim. Fl. Esseq. 92 = T. sessilifolia.
laurifolia, Roem. & Schult. Syst. iv. 543 = T. laurifolia.
longifolia, Roem. & Schult. l. c. = T. longifolia.
macroloba, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 196 = T. macroloba.
Martii, Miers, l. c. 194 = T. Martii.
membranacea, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 181 = T. scandens.
microphylla, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 198 = T. volubilis.
Milleri, Roem. & Schult. Syst. iv. 544 = T. scandens.
minor, G. Don, Gen. Syst. iv. 372 = T. incana.
minuta, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 198 = T. incana.
montana, Roem. & Schult. Syst. iv. 544 = T. montana.
organensis, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 194.—Bras.
paniculata, G. Don, Gen. Syst. iv. 370 = T. paniculata.
piliochros, G. Don, l. c. 371 = T. poliochros.
Pohlii, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 203 = T. Pohlii.
polystachya, Roem. & Schult. Syst. iv. 542 = = T. polystachya.
psilostachya, G. Don, Gen. Syst. iv. 370 = T. psilostachya.
punctata, Spreng. Neue Entdeck. iii. 28 = T. psilostachya.
ramiflora, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 197.—Bras.
rosmarinifolia, Roem. & Schult. Syst. iv. 844 = T. sibirica.
salicifolia, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 181 = T. salicifolia.
Salzmanni, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 201 = T. Salzmanni.
scandens, Roem. & Schult. Syst. iv. 542 = T. scandens.
sericea, Roem. & Schult. l. c. = T. tomentosa?
sibirica, Linn. Mant. ii. 334 = T. sibirica.
spigelliflora, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 200 = T. spigeliaeflora.
subsessilis, G. Don, Gen. Syst. iv. 370 = T. subsessilis.
subulata, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 532 = T. Gardneri.
syringaefolia, Roem. & Schult. Syst. iv. 542 = T. laurifolia.
tomentosa, G. Don, Gen. Syst. iv. 371 = T. tomentosa.
umbellata, G. Don, l. c. = T. umbellata.
valga, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 198.—Bras.
Vauthieri, Miers, l. c. 196 = T. Vauthieri.
velutina, G. Don, Gen. Syst. iv. 370 = T. velutina.
vicinia, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 197.—Bras.
villosa, Miers, l. c. 202 = T. villosa.
volubilis, Roem. & Schult. Syst. iv. 841 = T. volubilis.

MESTOTES, Soland. ex DC. Prod. ii. 57 (1825) = **Chailletia**, DC.

MESUA, Linn. Syst. ed. I (1735). *GUTTIFERAE*, Benth. & Hook. f. i. 176.
NAGASSARI, Adans. Fam. ii. 449 (1763).
NAGATAMPO, Adans. l. c. 444 (1763).
NAGHAS, Mirb. ex Steud. Nom. ed. II. ii. 136, 180 (1841).
bracteata, *Spreng. Syst.* iii. 127.—Cochinch.
coromandelina, Wight, Illustr. 127 = ferrea.
ferrea, *Linn. Sp. Pl.* 515.—Ind. or.
glabra, *Forsk. Fl. Aegypt. Arab.* 100.—Arab.
lepidota, *T. Anders. in Hook. f. Fl Brit. Ind.* i. 278.—Malaya.
nagana, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 4 = ferrea.
nervosa, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 307 = Kayea nervosa.
pedunculata, Wight, Illustr. i. 127 = ferrea.
pulchella, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 307 = ferrea.
Roxburghii, Wight, Illustr. i. 127 = ferrea.
salicina, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 302 = ferrea.
sclerophylla, Thw. Enum. Pl. Zeyl. 407 = ferrea.
singaporiana, Wall. Cat. n. 4836 = Kayea racemosa.
speciosa, Choisy, in DC. Prod. i. 562 = ferrea.
Thwaitesii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 305.—Zeylan.
Walkeriana, Planch. & Triana, l. c. 301 = ferrea.

MESYNIUM, Rafin. Fl. Tellur. iii. 33 (1836) = **Linum**, Tourn.
africanum, Rafin. l. c. = L. africanum.
chilense, Rafin. l. c. nomen = L. chilense?
mexicanum, Rafin. l. c. = L. mexicanum.
texense, Rafin. l. c. = L. Berlandieri.

METABASIS, DC. Prod. vii. 97, 307 (1838) = **Hypochoeris**, Linn. (Compos.).
aethnensis, DC. l. c. 307 = H. serioloides.
cretensis, DC. l. c. = H. cretensis.
hymettia, DC. l. c. 97 = H. cretensis.
neapolitanus, Nym. Consp. 471 = H. radicata.

METABOLOS, Blume, Bijdr. 990 (1826) = **Hedyotis**, Linn. (Rubiac.).
angustifolius, DC. Prod. iv. 436 = H. angustifolia.
caeruleus, Blume, Bijdr. 992 = H. vestita.
ciliaris, Blume, ex Miq. Fl. Ind. Bat. ii. 182 = H. Auricularia.
ferrugineus, DC. Prod. iv. 436.—Ins. Philipp.
laevigatus, DC. l. c. = H. laevigata.
latifolius, Blume, Bijdr. 991 = H. latifolia.
lineatus, DC. Prod. iv. 435 = H. vestita.
macrophyllus, Zoll. & Mor. Syst. Verz. Zoll. 63 = H. iodoneura.
prostratus, Blume, Bijdr. 991 = H. prostrata.
radicans, DC. Prod. iv. 435 = H. radicans.
rigidus, Blume, Bijdr. 992 = H. rigida.
rugosus, Blume, l. c. 991 = H. rugosa.
venosus, Blume, l. c. = H. Auricularia.

METABOLUS, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 237 = praec.

METACHILUM, Lindl. Gen. et Sp. Orch. 74 (1830) = **Appendicula**, Blume (Orchid.).
cyathiforme, Lindl. l. c. = A. Lewisii.

METAGNANTHUS, Endl. Gen. 438 (1838) = **Athanasia**, Linn. (Compos.).
parviflorus, Heynh. Nom. ii. 402 = A. parviflora, *Thunb.*

METAGNATHUS, Benth. & Hook. f. Gen. ii. 417 (1878), sphalm. = praec.

METAGONIA, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 263 = **Vaccinium**, Linn.
acuminata, Nutt. l. c. 266 = V. acuminatum.
alaternoides, Nutt. l. c. 265 = V. alaternoides, *H. B. & K.*
calycina, Nutt. l. c. 264 = V. calycinum.
cerea, Nutt. l. c. = V. cereum.
corymbodendron, Nutt. l. c. = V. corymbodendron.
crenulata, Nutt. l. c. 266 = V. crenulatum.

METAGONIA:—
didymantha, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 265 = Vaccinium didymanthum.
empetrifolia, Nutt. l. c. = V. empetrifolia.
marginata, Nutt. l. c. 266 = V. marginatum.
meridionalis, Nutt. l. c. 264 = V. meridionale.
myrtifolia, Nutt. l. c. 265 = V. crassifolium.
ovata, Nutt. l. c. 264 = V. ovatum.
penaeoides, Nutt. l. c. 265 = V. penaeoides.
penduliflora, Nutt. l. c. 264 = V. penduliflorum.
prostrata, Nutt. l. c. 265.—Peruv.
Pyxothamnus, Nutt. l. c. 264 = V. ovatum.
Schlechtendalii, Nutt. l. c. 266 = V. Schlechtendalii.
terniflora, Nutt. l. c. 265 = V. terniflorum.
villosa, Nutt. l. c. 266 = V. villosum.

METALASIA, R. Br. in Trans. Linn. Soc. xii. (1817). *COMPOSITAE*, Benth. & Hook. f. ii. 325.
ENDOLEUCA, Cass. in Bull. Soc. Philom. (1819) 47; et in Dict. Sc. Nat. xiv. 474 (1819).
ERYTHROPOGON, DC. Prod. vi. 254 (1837).
TULOCLINIA, Rafin. Fl. Tellur. iv. 119 (1836).
aduncа, *Less. Syn. Comp.* 341.—Afr. austr.
aristata, DC. Prod. vi. 250 = muricata.
aurea, *D. Don, in Mem. Wern. Soc.* v. (1826) 558.—Afr. austr.
capitata, *Less. Syn. Comp.* 337.—Afr. austr.
Cephalotes, DC. Prod. vi. 252 = concinna.
Cephalotes, *Less. Syn. Comp.* 338.—Afr. austr.
compacta, *Zeyh. ex Sch. Bip. in Flora*, xxvii. (1844) 693.—Afr. austr.
concinna, *Harv. in Harv. & Sond. Fl. Cap.* iii. 271.—Afr. austr.
cymbaefolia, *Harv. l. c.* 267.—Afr. austr.
cymosa, Cass. in Dict. Sc. Nat. xxx. 223 = muricata.
depressa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 267.—Afr. austr.
distans, *DC. Prod.* vi. 254.—Afr. austr.
divergens, *D. Don, in Mem. Wern. Soc.* v. (1826) 557.—Afr. austr.
divergens, Eckl. ex Harv. & Sond. Fl. Cap. iii. 267 = depressa.
Dregeana, *DC. Prod.* vi. 251.—Afr. austr.
Eckloniana, DC. l. c. 252 = fasciculata.
ericoides, Sieber, ex DC. Prod. vi. 253 = capitata.
erubescens, *DC. l. c.* 254.—Afr. austr.
fasciculata, *D. Don, in Mem. Wern. Soc.* v. (1826) 557.—Afr. austr.
fastigiata, D. Don, l. c. = muricata.
glomerata, DC. Prod. vi. 252 = M. Cephalotes.
hispida, D. Don, in Mem. Wern. Soc. v. (1826) 558 = Elytropappus spinellosus.
imbricata, Harv. in Harv. & Sond. Fl. Cap. iii. 266 = uniflora.
intermedia, DC. Prod. vi. 253 = erubescens.
lanceolata, DC. l. c. 251 = pulcherrima.
Lichtensteinii, *Less. Syn. Comp.* 335.—Afr. austr.
mucronata, R. Br. in Trans. Linn. Soc. xii. (1817) 124 = Helichrysum mucronatum.
muraltiaefolia, DC. Prod. vi. 253 = M. Cephalotes.
muricata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 124; *D. Don, in Mem. Wern. Soc.* v. (1826) 557.—Afr. austr.
nitidula, *Harv. in Harv. & Sond. Fl. Cap.* iii. 268.—Afr. austr.
octoflora, *DC. Prod.* vi. 250.—Afr. austr.
phylicoides, D. Don, in Mem. Wern. Soc. v. (1826) 559 = muricata.
polyanthos, D. Don, l. c. 558 = muricata.
pulcherrima, Eckl. ex Harv. & Sond. Fl. Cap. iii. 280 = Stoebe gomphrenoides.
pulcherrima, *Less. Syn. Comp.* 340.—Afr. austr.
pungens, D. Don, in Mem. Wern. Soc. v. (1826) 558 = muricata.
quinqueflora, *DC. Prod.* vi. 254.—Afr. austr.
rosea, DC. l. c. 252 = M. Cephalotes.
rosmarinifolia, Sieber, ex DC. l. c. 166 = Helichrysum diosmaefolium.
seriphiifolia, *DC. Prod.* vi. 254.—Afr. austr.
seriphioides, R. Br. in Trans. Linn. Soc. xii. (1817) 124 = Trichogyne seriphioides.
stricta, *Less. Syn. Comp.* 336.—Afr. austr.
tenuifolia, *DC. Prod.* vi. 253.—Afr. austr.
umbellata, Cass. in Dict. Sc. Nat. xxx. 223 = Lichtensteinii.
umbellata, *D. Don, in Mem. Wern. Soc.* v. (1826) 556.—Afr. austr.
uniflora, *D. Don, l. c.* 559—Afr. austr.

METALEPIS, Griseb. Cat. Pl. Cub. 179 (1866). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 766.
cubensis, *Griseb. l. c.* 180.—Cuba.

METANARTHECIUM, Maxim. in Bull. Acad. Pétersb. xi. (1867) 438. *LILIACEAE*, Benth. & Hook. f. iii. 825.
foliatum, *Maxim. in Trautv. Regel, Maxim. et Winkl. Decas Pl. Nov.* 10.—Japon.
luteo-viride, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 438.—Japon.

METAPLEXIS, R. Br. in Mem. Wern. Soc. i. (1809) 48. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 765.
UROSTELMA, Bunge, Enum. Pl. Chin. Bor. 44 (1835).
chinensis, *Decne. in DC. Prod.* viii. 511.—China.
fimbriata, *Spreng. Syst.* i. 854.—Venezuela.
mucronata, Spreng. Neue Entdeck. i. 269 (Gomphocarpi sp.).—Afr. austr.
rostellata, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 253.—Manchuria.
Stauntoni, *Schult. Syst.* vi. 111.—China.

METASTELMA, R. Br. in Mem. Wern. Soc. i. (1809) 52. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 755.
ACROCORYNE, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 316.
IRMISCHIA, Schlecht. in Linnaea, xix. (1847) 738.
acuminatum, *Decne. in DC. Prod.* viii. 514.—Bras.
albiflorum, *Griseb. Fl. Brit. W. Ind.* 417.—Ind. occ.
angustifolium, Torr. in Bot. Mex. Bound. 159 = Melinia angustifolia.
angustifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 315.—Mexic.
aphyllum, Decne. in DC. Prod. viii. 515 (= *Amphistelma aphyllum*).—Bras.
arizonicum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85.—Arizona.
bahamense, *Griseb. Cat. Pl. Cub.* 174.—Ind. occ.
barbigerum, *Scheele, in Linnaea*, xxi. (1848) 760.—Mexic.
Berterianum, *Decne. in DC. Prod.* viii. 515.—N. Granat.
Blodgetti, *A. Gray, in Proc. Am. Acad.* xii. (1877) 73.—Florida.
brachystephanum, *Griseb. Cat. Pl. Cub.* 174.—Cuba.
bracteolatum, *Fourn. in Mart. Fl. Bras.* vi. iv. 209.—Bras.
calcaratum, Decne. in DC. Prod. viii. 513 = Zygostelma calcaratum.
californicum, *Benth. Bot. Voy. Sulph.* 33. t. 18.—Calif.
campanulatum, *Decne. in DC. Prod.* viii. 514.—Guiana.
cordatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 253.—Bras.
costatum, D. Dietr. Syn. Pl. ii. 908 = Parapodium costatum.
cubense, *Decne. in DC. Prod.* viii. 515.—Ind. occ.
cubense, Griseb. Fl. Brit. W. Ind. 417 = bahamense.
cuspidatum, *Decne. in DC. Prod.* viii. 516.—Bras.
Decaisneana, *Benth. & Hook. f. Gen.* ii. 756, in nota sub Tassadia.—Guian.
diffusum, *Decne. in DC. Prod.* viii. 516.—Bras.
filiforme, Wright, in Sauv. Fl. Cub. 120 = Vincetoxicum scoparium.
Fraseri, Decne. in DC. Prod. viii. 513 = albiflorum?
fusculum, *Wright, in Sauv. Fl. Cub.* 120.—Cuba.
gracile, *Decne. in DC. Prod.* viii. 515.—Ins. S. Doming.
grandiflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 314.—Venezuela.
guianensis, *Klotzsch, in Schomb. Fauna & Fl. Brit. Gui.* 1083, 1147, nomen.—Guian.
hamatum, *Griseb. Cat. Pl. Cub.* 173.—Cuba.
hernandifolium, Decne. in DC. Prod. viii. 516 = Jobinia hernandifolia.
Hilarianum, *Fourn. in Mart. Fl. Bras.* vi. iv. 209.—Bras.
hirsutum, *Klotzsch, in Linnae*, xiv. (1840) 290.—Bras.
latipes, Decne. in DC. Prod. viii. 513 = Cyathostelma latipes.
lineare, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 292.—Porto Rico.
linearifolium, A. Rich. Fl. Cub. Fanerog. ii. 96 = (*Amphistelma filiforme*).—Cuba.
longicaule, *Fourn. in Mart. Fl. Bras.* vi. iv. 210.—Bras.
Martianum, Decne. in DC. Prod. viii. 513 = Petalostelma Martiana.

METASTELMA:—
mucronatum, *Decne. in DC. Prod.* viii. 514.—N. Granat.
myrtifolium, Decne. l. c. = Stelmation myrtifolium.
obscurum, *Fourn. in Mart. Fl. Bras.* vi. IV. 208.—Bras.
odoratum, *Decne. in DC. Prod.* viii. 513.—Bras.
organense, *Fourn. in Mart. Fl. Bras.* vi. IV. 209.—Bras.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 115.—Texas.
Paralias, *Decne. in DC. Prod.* viii. 514.—Ind. occ.
parviflorum, *R. Br. in Mem. Wern. Soc.* i. (1809) 52.—Am. trop.
parviflorum, Chapm. Fl. S. U. St. 367 = Blodgetti.
parviflorum, Schlecht. in Linnaea, vi. (1831) 731 = Schlechtendalii.
parvifolium, *Klotzsch, in Schomb. Fauna & Fl. Brit. Gui.* 1147, nomen.—Guian.
pedunculare, *Decne. in DC. Prod.* viii. 514.—Bras.
penicillatum, *Griseb. Cat. Pl. Cub.* 174.—Cuba.
purpurascens, *Benth. Pl. Hartw.* 215.—Ecuador.
Riedelii, *Fourn. in Mart. Fl. Bras.* vi. IV. 209.—Bras.
rotundifolium, *Decne. in DC. Prod.* viii. 514.—Bras.
rugosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 314.—N. Granat.
Schlechtendalii, Chapm. Fl. S. U. St. 366 = bahamense.
Schlechtendalii, *Decne. in DC. Prod.* viii. 513.—Am. trop.
sessilifolium, *Fourn. in Mart. Fl. Bras.* vi. IV. 207.—Bras.
stenolobum, *Decne. in DC. Prod.* viii. 515.—Guian.
Steudelianum, *Miq. in Linnaea*, xviii. (1844) 753.—Guian.
suaveolens, *Schlecht. in Linnaea*, xxvi. (1853–55) 667.—Austral.
suberosum, *Griseb. Fl. Brit. W. Ind.* 417.—Am. trop.
tomentosum, *Decne. in DC. Prod.* viii. 514.—Bras.
trifurcatum, *Griseb. in Goett. Abh.* xxiv. (1879) 227.—Reg. Argent.
tubatum, *Griseb. l. c.*—Reg. Argent.
uncinatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 211.—Bras.
urceolatum, *Fourn. l. c.* 210.—Bras.
venosum, *Decne. in DC. Prod.* viii. 514.—Bras.
virgatum, Decne. l. c. 515 (= *Orthosia virgata*).—Bras.

METAXANTHUS, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 280 = seq.

METAZANTHUS, Meyen, Reise, i. 356 (1834) = **Senecio**, Tourn. (Compos.).
cacalioides, Meyen, l. c.—Chili.
grandiflorus, Meyen, l. c.—Chili.

METEORINA, Cass. in Bull. Soc. Philom. (1818) 167 = **Dimorphotheca**, Vaill. (Compos.).
crassipes, Cass. in Dict. Sc. Nat. xxx. 322 = D. pluvialis.
gracilipes, Cass. l. c. 320 = D. pluvialis.
gracilis, Less. ex Steud. Nom. ed. II. i. 509; ii. 137 = D. pluvialis.
lyrata, Cass. in Dict. Sc. Nat. xxx. 321 = D. pluvialis.

METEORUS, Lour. Fl. Cochinch. 410 (1790) = **Barringtonia**, Forst. (Myrtac.).
coccineus, Lour. l. c. = B. excelsa.

METERANA, Rafin. Sylva Tellur. 65 (1838) = **Caperonia**, A. St. Hil. (Euphorb.).
arborea, Rafin. l. c. 66.—Hab.?
castaneaefolia, Rafin. l. c. = C. castaneaefolia.
dimidiata, Rafin. l. c. = Acalypha australis.
palustris, Rafin. l. c. = C. palustris.

METHONICA, Tourn. ex Crantz, Inst. i. 474 (1766); Juss. Gen. 48 (1789) = **Gloriosa**, Linn. (Liliac.).
abyssinica, Walp. Ann. iii. 630 = G. abyssinica.
Doniana, Kunth, Enum. Pl. iv. 277 = G. superba.
Gloriosa, Salisb. in Trans. Hort. Soc. i. (1812) 331 = G. simplex.
grandiflora, Hook. Bot. Mag. t. 5216 = G. simplex.
Leopoldi, Van Houtte, ex Lem. in Fl. des Serres, ii. (1846) t. 163–4 = G. simplex.
Petersiana, Klotzsch, ex Garcke, in Peters, Reise Mossamb. Bot. 519. t. 55 = G. simplex.
Plantii, Hort. ex Planch. in Fl. des Serres, ix. (1853–54) t. 865 = G. simplex.

METHONICA:—
platyphylla, Klotzsch, ex Garcke, in Peters, Reise Mossamb. Bot. 520. t. 55 = Gloriosa simplex.
superba, Crantz, Inst. i. 474; Kunth, Enum. Pl. iv. 276 = G. superba.
virescens, Kunth, l. c. 277 = G. simplex.

METHORIUM, Schott & Endl. Melet. 29. t. 5 (1832) = **Helicteres**, Pluk. (Stercul.).
canum, Schott & Endl. l. c. = H. cana.
dentatum, F. Muell. Census, 15 = H. dentata.
integrifolium, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 40 = H. cana.
semiglabrum, F. Muell. Fragm. v. 43 = H. semiglabra.
spicatum, F. Muell. Census, 15.—Austral.

METHYSCOPHYLLUM, Eckl. & Zeyh. Enum. 152 (1836) = **Catha**, Forsk. (Celastr.).
glaucum, Eckl. & Zeyh. l. c.—Afr. austr.

METHYSTICUM, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
esculentum, Rafin. l. c. = P. Methysticum.

METOPIUM, P. Br. Hist. Jamaic. 177 (1756) = **Rhus**, Tourn. (Anacard.).
Linnaei, Engl. in DC. Monog. Phan. iv. 367 = R. Metopium.
venosum, Engl. l. c. = R. venosa.

METROCYNIA, Thou. Gen. Nov. Madag. 22 (1806) = **Cynometra**, Linn. (Legumin.).
Commersonii, DC. Prod. ii. 507.—Madag.

METRODOREA, A. St. Hil. Fl. Bras. Mer. i. 81. t. 16 (1825) = **Esenbeckia**, Kunth (Rutac.).
atropurpurea, Fisch. ex Lem. in Fl. des Serres, iv. (1848) t. 337 = *nigra*.
nigra, A. St. Hil. Fl. Bras. Mer. i. 81.—Bras.
pubescens, A. St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 139.—Bras.
stipularis, Mart. in Flora, xx. (1837) II. Beibl. 124.—Bras.

METROSIDEROS, Banks, ex Gaertn. Fruct. i. 170. t. 34 (1788). *MYRTACEAE*, Benth. & Hook. f. i. 710.
NANI, Adans. Fam. ii. 88 (1763), nomen prius.
NANIA, Miq. Fl. Ind. Bat. i. I. 399 (1855).
STENOSPERMUM, Sweet, Hort. Brit. ed. II. 209 (1830).
abietina, Hoffmgg. Verz. Pfl. 80 = Kunzea corifolia.
acuminata, Decne. Bot. Voy. Vénus, 29 = villosa.
acuminata, *Dum.-Cours. Bot. Cult.* ed. II. v. 381 (Quid?).—Austral.
albida, Sieber, ex DC. Prod. iii. 212 = Melaleuca Leucadendron.
albiflora, *Soland. ex Gaertn. Fruct.* i. 172. t. 34. f. 11.—N. Zel.
angustifolia, Dum.-Cours. Bot. Cult. ed. II. v. 378 = Callistemon lanceolatus.
angustifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 270.—Afr. austr.
anomala, Vent. Jard. Malm. 5 = Angophora cordifolia.
apocynifolia, Salisb. Prod. 351 = Angophora lanceolata.
armillaris, Cav. Ic. iv. 19 = Melaleuca ericifolia.
armillaris, Soland. ex Gaertn. Fruct. i. 171. t. 34 = Melaleuca armillaris.
aromatica, Salisb. Prod. 351 = Eucalyptus piperita.
aspera, *Dum.-Cours. Bot. Cult.* ed. II. v. 381.—Austral.
aspera, Hoffmgg. Verz. Pfl. 80 = Callistemon rigidus.
australis, R. Br. ex Bonpl. Desc. Pl. Rar. Malm. 102, in textu = Callistemon salignus.
buxifolia, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 111 = scandens.
buxifolia, *Dum.-Cours. Bot. Cult.* ed. II. v. 379.—Austral.
calicina, Cav. Ic. iv. 20 = Melaleuca thymifolia.
calyculatus, Sieber, ex DC. Prod. iii. 223 = Callistemon linearis.
canaliculata, Hort. Angl. ex Dum.-Cours. Bot. Cult. ed. II. v. 381 (Quid?).—Austral.
capitata, Sm. in Trans. Linn. Soc. iii. (1797) 273 = Kunzea capitata.
chrysantha, F. Muell. Fragm. iv. 159 = Xanthostemon chrysanthus.
ciliaris, Dum.-Cours. Bot. Cult. ed. II. v. 378 = seq.
ciliata, *Sm. in Trans. Linn. Soc.* iii. (1797) 271.—N. Caled.

METROSIDEROS:—
citrina, Curt. Bot. Mag. t. 260 = Callistemon lanceolatus.
Colensoi, *Hook. f. Fl. N. Zel.* i. 68.—N. Zel.
collina, A. Gray, Bot. U. St. Expl. Exped. i. 558. t. 68 = villosa.
comosa, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii. 478.—Ins. Molucc.
connata, Desf. Tabl. ed. I. 171 = Eugenia Ventenatii.
cordifolia, Pers. Syn. ii. 25 = Angophora cordifolia.
coriacea, Salisb. Prod. 352; Poir. Encyc. Suppl. iii. 685 = Melaleuca Leucadendron.
corifolia, Vent. Jard. Malm. 46. t. 46 = Kunzea corifolia.
costata, Gaertn. Fruct. i. 171 = Angophora lanceolata.
crassifolia, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 379 = Callistemon speciosus.
decora, Salisb. Prod. 352 = Melaleuca genistifolia.
decurrens, Hort. Angl. ex Steud. Nom. ed. II. ii. 137 (Quid?).—Austral.
diffusa, A. Cunn. in Ann. Nat. Hist. iii. (1839) 114 = albiflora.
diffusa, Hook. & Arn. Bot. Beech. Voy. 63 = villosa.
diffusa, *Sm. in Trans. Linn. Soc.* iii. (1794) 268.—N. Zel.
ericifolia, Sm. in Rees, Cycl. xxiii. n. 16 = Kunzea ericifolia.
eucalyptoides, *F. Muell. Fragm.* i. 243.—Austral.
excelsa, *Soland. ex Gaertn. Fruct.* i. 172. t. 34. f. 8 (= Leptospermi sp.?).—Hab.?
falcata, Dehnh. Revist. Napolit. i. III. 172; ex Walp. Rep. ii. 165 = seq.?
falcata, *Dum.-Cours. Bot. Cult.* ed. II. v. 378.—Austral.
fasciculata, Dum.-Cours. l. c. 381 = aspera?
flexuosa, Willd. Enum. Hort. Berol. 514 = Agonis flexuosa.
floribunda, Sm. in Trans. Linn. Soc. iii. (1797) 267 = Angophora intermedia.
floribunda, Vent. Jard. Malm. t. 75 = Eugenia Ventenatii.
florida, Hook. Bot. Mag. t. 4471 = robusta.
florida, *Sm. in Trans. Linn. Soc.* iii. (1797) 269.—N. Zel.
fulgens, Soland. ex Gaertn. Fruct. i. 172 = Angophora lanceolata.
glandulosa, Desf. Cat. Hort. Par. ed. III. 407 = Callistemon rigidus.
glauca, Bonpl. Desc. Pl. Rar. Malm. 86. t. 34 = Callistemon speciosus.
glauca, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 380 = praec.?
glomeruliflora, A. Gray, Bot. U. St. Expl. Exped. i. 556, in syn. = Syncarpia laurifolia.
glomulifera, Sm. in Trans. Linn. Soc. iii. (1797) 269 = Syncarpia laurifolia.
gracilis, Salisb. Prod. 352 = Melaleuca thymifolia.
gummifera, Gaertn. Fruct. i. 170. t. 34. f. 1 = Eucalyptus corymbosa.
hirsuta, Andr. Bot. Rep. t. 281 = Angophora cordifolia.
hispida, Sm. in Trans. Linn. Soc. iii. (1797) 267 = Angophora cordifolia.
Homeana, Turcz. ex Hook. f. Handb. N. Zeal. Fl. 71 = diffusa, *Sm.*
hybrida, Otto, ex DC. Prod. iii. 224 = Callistemon hybridus.
hypericifolia, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 114.—N. Zel.
hypericifolia, Salisb. Prod. 351 = Melaleuca hypericifolia.
hyssopifolia, Cav. Ic. iv. 20 = Melaleuca linariifolia.
juniperina, Reichb. ex Spreng. Syst. iv. Cur. Post. 194 = Melaleuca nodosa.
juniperoides, Reichb. ex DC. Prod. iii. 213 = Melaleuca nodosa.
lanceolata, Pers. Syn. ii. 25 = Angophora lanceolata.
lanceolata, Sm. in Trans. Linn. Soc. iii. (1797) 272 = Callistemon lanceolatus.
latifolia, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 379 = Callistemon lanceolatus.
laurifolia, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 300.—N. Caled.
laurifolia, Desf. Tabl. ed. I. 171, nomen = Eugenia Smithii.
laurifolia, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 376 = Eugenia Ventenatii.
leptopetala, *F. Muell. Docum. Intercol. Exhib.* 30.—Austral.

METROSIDEROS:—
linearifolia, Link, Enum. Hort. Berol. ii. 26 = Callistemon rigidus.
linearis, Sm. in Trans. Linn. Soc. iii. (1797) 271 = Callistemon linearis.
linearis, Willd. Enum. Hort. Berol. 513 = Callistemon rigidus.
linifolia, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 277 = Callistemon rigidus.
longifolia, Dum.-Cours. l. c. v. 378 = Callistemon lanceolatus.
lophantha, Vent. Jard. Cels, t. 69 = Callistemon lanceolatus.
lucida, Linn. f. ex Hook. f. Handb. N. Zeal. Fl. 787 = diffusa, *Sm.*
lucida, *A. Rich. Fl. N. Zel.* i. 333.—N. Zel.
lutea, A. Gray, Bot. U. St. Expl. Exped. i. 560. t. 69. = villosa.
macrophylla, *Lam. Tabl. Encyc.* ii. 538.—Madag.
macropunctata, Dum.-Cours. Bot. Cult. ed. II. vii. 277 = Callistemon rigidus.
macropus, *Hook. & Arn. Bot. Beech. Voy.* 83.—Ins. Sandvic.
marginata, Cav. Ic. iv. 18 = Callistemon lanceolatus.
myrtifolia, Dum.-Cours. Bot. Cult. ed. II. v. 379 = scandens?
myrtifolia, Hoffmgg. Verz. Pfl. 80 = Callistemon lanceolatus.
myrtifolia, Soland. ex Gaertn. Fruct. i. 172. t. 34. f. 12 = diffusa.
nana, C. C. Gmel. Hort. Carlsr. ex DC. Prod. iii. 226 (Quid?).—Hab.?
nervulosa, *C. Moore & F. Muell. Fragm.* viii. 15.—Ins. Dom. Howe.
nitida, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 182.—N. Caled.
nodosa, Soland. ex Gaertn. Fruct. i. 172. t. 34 = Melaleuca nodosa.
obovata, Hook. & Arn. Bot. Beech. Voy. 63 = villosa.
operculata, *Labill. Sert. Austro-Caled.* 61. t. 60.—N. Caled.
pallida, Bonpl. Desc. Pl. Rar. Malm. 101. t. 41 = Callistemon salignus.
paradoxa, F. Muell. Fragm. i. 243 = Xanthostemon paradoxus.
Parkinsonii, *J. Buch. in Trans. N. Z. Inst.* xv. (1883) 339.—N. Zel.
pauciflora, Endl. in Enum. Pl. Hueg. 50 = Eremaea pilosa.
pendens, *Colenso, in Trans. N. Z. Inst.* xii. (1880) 360.—N. Zel.
perforata, A. Rich. Ess. Fl. Nouv. Zél. 334 = scandens.
pictipetala, Blanco, Fl. Filip. ed. II. 295 = vera.
pinifolia, Wendl. Coll. i. 53. t. 16 = Callistemon pinifolius.
Pleurocalyptus, Baill. Hist. Pl. vi. 337 = Pleurocalyptus Deplanchei.
pliniaefolia, Desf. Tabl. ed. I. 171, nomen (Quid?).—Austral.
polymorpha, Gaudich. in Freyc. Voy. Bot. 482. t. 85 = villosa.
procera, Salisb. Prod. 351 = Syncarpia laurifolia.
propinqua, Endl. in Enum. Pl. Hueg. 50 = Kunzea ericifolia.
propinqua, Salisb. Prod. 351 = Syncarpia laurifolia.
pubescens, *Baill. Hist. Pl.* vi. 337.—N. Caled.
pubescens, Hort. Angl. ex Dum.-Cours. Bot. Cult. ed. II. vii. 277 (Quid?).—Austral.
pungens, Reichb. in Spreng. Syst. iv. Cur. Post. 194 = Melaleuca nodosa.
quinquenervia, Cav. Ic. iv. 19 = Melaleuca Leucadendron.
rigida, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 381 = Callistemon rigidus.
rigidifolia, Hoffmgg. Verz. Pfl. 80 = Callistemon rigidus.
robusta, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 112.—N. Zel.
rubra, *Baill. Hist. Pl.* vi. 337.—N. Caled.
rugosa, A. Gray, Bot. U. St. Expl. Exped. i. 561. t. 69. = villosa.
rugulosa, Sieber, ex Benth. Fl. Austral. iii. 120 = Callistemon lanceolatus.
rugulosa, Willd. Enum. Hort. Berol. Suppl. 31, nomen; et ex Link, Enum. Hort. Berol. ii. 27 = Callistemon rigidus.
salicifolia, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 114 = Olea montana, Santalum Cunninghamii.

METROSIDEROS:—
salicifolia, Soland. ex Gaertn. Fruct. i. 171. t. 34. f. 3 (Eucalypti sp.?).—Hab.?
saligna, Sm. in Trans. Linn. Soc. iii. (1797) 272 = Callistemon salignus.
scabra, Colla, Hort. Ripnl. 91 = Callistemon rigidus.
scabra, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 381 = acuminata, *Dum.-Cours.*
scandens, *Soland. ex Gaertn. Fruct.* i. 172. t. 34. f. 10.—N. Zel.
scariosa, *Wendl. ex Hornem. Hort. Hafn. Suppl.* 139.—Hab.?
semperflorens, Lodd. Bot. Cab. t. 523 = Callistemon lanceolatus.
sororia, Endl. in Enum. Pl. Hueg. 49 = Kunzea recurva?
speciosa, Sims, Bot. Mag. t. 1761 = Callistemon speciosus.
spectabilis, Soland. ex Gaertn. Fruct. i. 172. t. 34. f. 9 = villosa.
splendens, DC. Prod. iii. 222, in syn. = Angophora lanceolata.
stipularis, *Hook. f. Fl. Antarct.* 275.—Chili.
suberosa, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii. 478.—Ins. Molucc.
subsimilis, *Colenso, in Trans. N. Z. Inst.* xii. (1880) 361.—N. Zel.
taitensis, Decne. Bot. Voy. Vénus, 30 = villosa.
ternifolia, *F. Muell. Docum. Intercol. Exhib.* 30.—Austral.
tetrapetala, *F. Muell. Fragm.* vii. 41.—Austral.
tomentosa, *A. Rich. Ess. Fl. Nouv. Zél.* 336.—N. Zel.
trinervia, *Hort. Angl. ex Steud. Nom.* ed. II. ii. 138 (Quid?).—Austral.
tubiflora, *Breit. Hort. Breit.* 282 (Quid?).—Austral.
umbellata, Cav. Ic. iv. 20. t. 337 = lucida.
vera, *Lindl. Coll. Bot.* t. 18.—Amboina; Java.
vesiculata, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 327.—N. Zel.
villosa, *Sm. in Trans. Linn. Soc.* iii. (1797) 268.—Ins. Tahiti.
viminalis, Soland. ex Gaertn. Fruct. i. 171. t. 34. f. 4 = Callistemon rigidus.
viridiflora, Sims, Bot. Mag. t. 2602 = Callistemon salignus.

METROXYLON, Rottb. in Nye Saml. Dansk. Vidensk. Selsk. Skrift. ii. (1783) 525. t. 1. *PALMAE*, Benth. & Hook. f. iii. 935.
Coelococcus, H. Wendl. in Bonplandia, x. (1862) 199.
Sagus, Rumph. ex Gaertn. Fruct. i. 27 (1788).
elatum, Hort. ex Scheff. in Ann. Jard. Buitenz. i. (1876) 162 = Heterospathe elata.
elatum, Mart. Hist. Nat. Palm. iii. 216. t. 102 = Pigafetta elata.
filare, Mart. l. c. 215 = Pigafetta filaris.
hermaphroditum, *Hassk. in Hoev. & De Vriese Tijdschr.* ix. (1842) 175.—Malaya.
inerme, Mart. Hist. Nat. Palm. iii. 215 = M. Sagu.
laeve, *Mart. l. c.* 214.—Siam.
longispinum, *Mart. l. c.* 215.—Amboina.
micracanthum, *Mart. l. c.*—Ins. Ceram.
microcarpum, Kunth, Enum. Pl. iii. 215 = Pigafetta filaris.
microspermum, Kunth, l. c. = Pigafetta filaris.
Rumphii, *Mart. Hist. Nat. Palm.* 213.—Malaya.
Sago, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 193. t. 4 = seq.
Sagu, *Rottb. in Nye Samml. Dansk. Vidensk. Selsk. Skrift.* ii. (1783) 527.—Ins. Molucc.
Sagus, Spreng. Syst. ii. 139 = praec.
sylvestre, *Mart. Hist. Nat. Palm.* iii. 215.—Amboina.
vitiense, *Benth. & Hook. f. Gen.* iii. 935.—Ins. Fiji.

METROXYLON, Spreng. Gen. 283 (1830) = **Raphia**, Beauv. (Palm.).
Ruffia, Spreng. Syst. ii. 139 = R. Ruffia.
taedigerum, Spreng. l. c. = R. taedigera.
textile, Welw. Apont. 584 = R. textilis.
viniferum, Spreng. Syst. ii. 139 = R. vinifera.

METTENIA, Griseb. Fl. Brit. W. Ind. 43 (1859). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 324.
globosa, *Griseb. Fl. Brit. W. Ind.* 43.—Ind. occ.

METTENIUSA, Karst. Fl. Columb. i. 79. t. 39 (1859?). *OLACINEAE*?
edulis, *Karst. l. c.*—N. Granat.

METTERNICHIA, Mikan, Del. Fl. et Faun. Bras. (1825) cum ic. *SOLANACEAE*, Benth. & Hook. f. ii. 905.
affinis, *Presl, Bot. Bemerk.* 102.—Bras.
megalandra, *Dun. in DC. Prod.* xiii. 1. 595.—N. Granat.
Princeps, Miers, in Hook. Lond. Journ. Bot. v. (1846) 145 = seq.
principis, *Mikan, Del. Fl. & Faun. Bras.* 3. t. 1.—Bras.
spectabilis, Regel, Cat. Pl. Hort. Aksakov. 93 = praec.?

MEUM, [Tourn.] Adans. Fam. ii. 97 (1763). *UMBELLIFERAE*, Benth. & Hook. f. i. 911.
adonidifolium, J. Gay.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 575, 576 = Ligusticum Mutellina.
alatum, Baill. Hist. Pl. vii. 107 = Ligusticum alatum.
anethifolium, Gaertn. Mey. & Scherb. Fl. Wett. iii. 11. 10 = athamanticum.
athamanticum, *Jacq. Fl. Austr.* iv. 2.—Europ.
atlanticum, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 59, nomen; *Ball, in Journ. Linn. Soc.* xvi. (1878) 474.—Marocc.
Bunius, Jacq. ex G. Don, in Loud. Hort. Brit. 104 = Carum Bunius.
capillaceum, Dulac, Fl. Hautes-Pyr. 354 = athamanticum.
carvifolium, Bertol. Fl. Ital. iii. 313 = Carum flexuosum.
corsicum, Bertol. l. c. 615 = Ligusticum corsicum.
Dalechampii, Tenore, Prod. Fl. Neap. 19 = Trinia vulgaris.
diffusum, Baill. Hist. Pl. vii. 191 = Seseli indicum.
fatuum, Pers. Syn. i. 319 = Carum verticillatum.
Foeniculum, Spreng. in Schult. Syst. vi. 433 = Foeniculum vulgare.
gayoides, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 32.—Kokan.
graecum, Boiss. & Heldr. ex Boiss. Fl. Orient. ii. 881 = Carum graecum.
heterophyllum, Moench, Meth. 86 = Carum Bunius.
inundatum, Spreng. Umbellif. Sp. 115 = Apium inundatum.
Mutellina, Gaertn. Fruct. i. 106. t. 23 = Ligusticum Mutellina.
nevadense, Boiss. Voy. Bot. Espagne, ii. 732 = athamanticum.
nodiflorum, Baill. Hist. Pl. vii. 191 = Apium nodiflorum.
nudicaule, Moench, Meth. 86 = Seseli montanum.
nudicaule, Trevir. Symb. Phytol. i. 31 = Peucedanum redivivum.
nutans, Moench, Meth. 86 = Carum rigidulum.
piperatum, Tenore, Fl. Nap. iii. 291 = seq.
piperinum, Colla, Herb. Pedem. iii. 46 = seq.
piperitum, Schult. Syst. vi. 435 = Foeniculum piperitum.
pyrenaicum, J. Gay, ex DC. Prod. iv. 162 = Ligusticum pyrenaicum.
resinosum, Baill. Hist. Pl. vii. 191 = Ferula nudicaulis.
rigidulum, Bertol. Fl. Ital. iii. 314 = Carum rigidulum.
scoticum, Baill. Hist. Pl. vii. 191 = Ligusticum scoticum.
segetum, Guss. Fl. Sic. Prod. i. 346 = Carum Ridolfia.
sibiricum, Spreng. Pugill. ii. 56 = Silaus tenuifolius.
siifolium, Presl, ex Steud. Nom. ed. II. ii. 138 (Quid?).—Sicilia.
Silaus, Baill. Hist. Pl. vii. 191 = Silaus flavescens.
tenuifolium, Duby, Bot. Gall. 230 = Seseli tenuifolium.
venosum, Baill. Hist. Pl. vii. 191 = Selinum sylvestre.

MEYENIA, Nees, in Wall. Pl. As. Rar. iii. 78 (1832). **Thunbergia**, Retz. (Acanthac.).
erecta, Benth. in Hook. Niger Fl. 476 = T. erecta.
Hawtayneana, Nees, in Wall. Pl. As. Rar. iii. 78 = T. Hawtayneana.
longiflora, Benth. in Hook. Niger Fl. 476 = T. fragrans.
Vogeliana, Benth. in Bot. Mag. t. 5389 = T. Vogeliana.

MEYENIA, Schlecht. in Linnaea, viii. (1833) 251 = **Cestrum**, Linn. (Solanac.).
corymbosa, Schlecht. l. c. 252 = C. Endlicheri.
cyanea, Heynh. Nom. ii. 404 = Iochroma tubulosa.
fasciculata, Schlecht. in Linnaea, viii. (1833) 251 = C. fasciculatum.
purpurea, Heynh. Nom. ii. 404 = C. elegans.
tomentosa, Heynh. l. c. = C. Benthami.

MEYERA, Adans. Fam. ii. 257 (1763) = **Holosteum**, Dill. (Caryoph.).

MEYERA, Schreb. Gen. 570 (1791) = **Enhydra**, Lour. (Compos.).
capitata, Spreng. Syst. iii. 601 = Wedelia biflora.
fluctuans, Spreng. l. c. 602 = E. fluctuans.
guineensis, Spreng. l. c. = E. fluctuans.
Heloncha, Buch.-Ham. ex Wall. Cat. n. 3195 = E. fluctuans.
maritima, H. B. & K. Nov. Gen. et Sp. iv. 269 = E. maritima.
orientalis, D. Don, Prod. Fl. Nep. 180 = Caesulia axillaris.
sessilis, Sw. Fl. Ind. Occ. iii. 1364 = E. sessilis.

MEYERIA, DC. Prod. v. 670 (1836) = **Calea**, Linn. (Compos.).
angustifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 414 = C. Gardneriana.
Candolleana, Gardn. l. c. = C. Candolleana.
elongata, Gardn. l. c. 415 = C. elongata.
hispida, DC. Prod. v. 671 = C. hispida.
hypericifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 413 = C. hypericifolia.
longifolia, DC. Prod. v. 671 = C. longifolia.
microphylla, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 413 = C. microphylla.
myrtifolia, DC. Prod. v. 670 = C. myrtifolia.
parvifolia, DC. l. c. = C. parvifolia.
teucriifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 412 = C. teucriifolia.

MEYNA, Roxb. ex Link, Jahrb. i. III. 32 (1820) = **Vangueria**, Juss. (Rubiac.).
spinosa, Roxb. ex Link, l. c. = V. spinosa.

MEYNIA, Schult. Mant. i. 67 (1822) = praec.

MEZEREUM, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 358 = **Daphne**, Linn. (Thymel.).
officinarum, C. A. Mey. l. c. = D. Mezereum.

MEZIERA, Baker, Fl. Maurit. 129 (1877), sphalm. = seq.

MEZIEREA, Gaudich. Bot. Voy. Bonite, t. 32 (1844–52) = **Begonia**, Linn.
Calleryana, Herb. Mus. Par. ex A. DC. in DC. Prod. xv. 1. 408 = B. aptera, *Roxb.*
Griffithiana, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 144 = B. episcopalis.
nepalensis, A. DC. l. c. = B. gigantea.
salaziensis, Gaudich. Bot. Voy. Bonite, t. 32 = B. aptera, *Roxb.*

MEZLERIA, Presl, Prod. Monog. Lobel. 7 (1836) = **Lobelia**, Linn. (Campan.).
depressa, Presl, l. c.; Sond. in Harv. & Sond. Fl. Cap. iii. 533 = L. depressa.
Dregeana, Sond. l. c. = L. Dregeana.
filicaulis, Presl, Prod. Monog. Lobel. 7 = L. depressa?
humifusa, A. DC. in DC. Prod. vii. 351 = L. alsinoides?
valdiviana, Phil. in Bot. Zeit. xxii. (1864) 217 = L. valdiviana.

MEZONEURUM, Desf. in Mém. Mus. Par. iv. (1818) 245. tt. 10, 11. *LEGUMINOSAE*, Benth. & Hook. f. i. 565.
angolense, *Welw. ex Oliver*, *Fl. Trop. Afr.* ii. 261.—Afr. trop.
Benthamianum, *Baill. Adansonia*, vi. (1866) 196.—Afr. trop.
brachycarpum, *Benth. Fl. Austral.* ii. 278.—Austral.
cucullatum, *Wight & Arn. Prod.* 283.—Ind. or.; Malaya.
enneaphyllum, *Wight & Arn. l. c.*—Ind. or.; Malaya.
glabrum, *Desf. in Mém. Mus. Par.* iv. (1818) 245. t. 10.—Burma; Malaya.
grande, Heyne, ex Wall. Cat. n. 5830 = cucullatum.
grande, Miq. Fl. Ind. Bat. Suppl. 291 = Acrocarpus grandis.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 338.—Afr. trop.
hymenocarpum, *Wight & Arn. Prod.* 283.—Ind. or.
intermedium, *Zipp. ex Span. in Linnaea*, xv. (1841) 200, nomen.—Ins. Timor.
macrophyllum, Benth. ex Miq. Fl. Ind. Bat. i. 1. 104 = cucullatum.
procumbens, Blanco, Fl. Filip. ed. II. 235 = glabrum.
pubescens, *Desf. in Mém. Mus. Par.* iv. (1818) 245. t. 11.—Burma; Malaya.

MEZONEURUM:—
Scortechinii, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 73.—Austral.
sulfureum, *Miq. Fl. Ind. Bat.* i. 1. 105.—Java.
sumatranum, *Wight & Arn. Prod.* 283.—Malaya.
Welwitschianum, *Oliver, Fl. Trop. Afr.* ii. 261.—Afr. trop.

MEZZETTIA, Becc. in Nuov. Giorn. Bot. iii. (1871) 187. *ANONACEAE.*
parviflora, *Becc. l. c.* 188.—Borneo.
umbellata, *Becc. l. c.* 187.—Borneo.

MIAGRUM, Crantz, Stirp. Austr. ed. I. 4 (1762) = **Myagrum**, Linn. (Crucif.).

MIBORA, Adans. Fam. ii. 495 (1763). *GRAMINEAE*, Benth. & Hook. f. iii. 1147.
CHAMAGROSTIS, Borkh. ex Wibel, Prim. Fl. Werth. 126 (1799).
KNAPPIA, Sm. Engl. Bot. t. 1127 (1803).
STURMIA, Hoppe, in Sturm, Deutschl. Fl. Heft vii. (1799).
Desvauxii, Lange, in Kjoeb. Vidensk. Meddel. (1860) 24 = verna.
minima, Beauv. Agrost. 30; Desv. Fl. Anj. 46 = verna.
verna, *Beauv. l. c.* 167; *S. F. Gray, Nat. Arr. Brit. Pl.* ii. 155.—Europ.; Afr. bor.; As. Min.

MICADANIA, R. Br. in Denh. & Clapp. Trav. App. 239 (1827) = **Butyrospermum**, Kotschy (Sapot.).

MICALIA, Rafin. Fl. Tellur. ii. 104 (1836) = **Escobedia**, Ruiz & Pav. (Scrophul.).
grandiflora, Rafin. l. c. 105 = E. scabrifolia.

MICAMBE, Adans. Fam. ii. 407 (1763) = **Cleome**, Linn. (Capparid.).
ornithopodioides, Medic. Phil. Bot. i. 43 = C. ornithopodioides.
violacea, Medic. l. c. = C. graveolens.

MICANIA, D. Dietr. Syn. Pl. iv. 1377 (1847) = **Mikania**, Willd. (Compos.).

MICHAUXIA, L'Hérit. Monog. [(1788)]; ex Ait. Hort. Kew. ed. I. ii. 8 (1789). *CAMPANULACEAE*, Benth. & Hook. f. ii. 561.
MINDIUM, Adans. Fam. ii. 134 (1763).
campanuloides, *L'Hérit. ex Ait. Hort. Kew.* ed. I. ii. 8.—As. Min.; Syria.
columnaris, Boiss. Diagn. Ser. II. iii. 117 = Tchihatcheffii.
decandra, Fisch. ex DC. Prod. vii. 457 = laevigata.
dodecandra, Fisch. ex Steud. Nom. ed. II. ii. 138 = laevigata.
laevigata, *Vent. Jard. Cels*, 81. t. 81.—Persia.
nova, J. F. Gmel. Syst. 618 = campanuloides.
nuda, *A. DC. in DC. Prod.* vii. 457.—Mesopot.
stenophylla, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 891.—Persia.
strigosa, Pers. Syn. i. 418 = campanuloides.
Tchihatcheffii, *Fisch. & Heldr. in Ann. Sc. Nat.* Sér. IV. i. (1854) 32.—As. Min.
thyrsoides, *Boiss. & Heldr. Diagn.* Ser. I. xi. 61.—As. Min.

MICHAUXIA, Neck. Elem. i. 12 (1790) = **Relhania**, L'Hérit. (Compos.).

MICHAUXIA, Raeusch. Nom. ed. III. 282 (1797) (Quid?).
spicata, Raeusch. l. c.—Hab.?

MICHAUXIA, Salisb. Prod. 386 (1796) = **Gordonia**, Ellis (Ternstroem.).
sessilis, Salisb. l. c. = G. pubescens.

MICHELARIA, Dum. Obs. Gram. Belg. 77 (1823) = **Bromus**, Linn. (Gramin.).
arduennensis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 59 = B. arduennensis.
bromoides, Dum. Obs. Gram. Belg. 77 = B. arduennensis.
eburonensis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 60 = B. racemosus.
hirsuta, Davr. Nécr. Lièg. (1854) 36 = B. racemosus.
villosa, Dum. ex Thielens, in Mém. Soc. Hainault, Sér. III. v. (1870) 200 = Agropyron repens?
villosa, Strail, in Bull. Acad. Brux. xxii. II. (1855) 516 = B. racemosus.

MICHELIA, Linn. Gen. ed. I. 119 (1737). *MAGNOLIACEAE*, Benth. & Hook. f. i. 19.
CHAMPACA, Adans. Fam. ii. 365 (1763).
LIRIOPSIS, Spach, Hist. Vég. Phan. vii. 460 (1839).
alba, *DC. Syst.* i. 449 (gen. dub.).—Java.
aurantiaca, Wall. Pl. As. Rar. ii. 39 = M. Champaca.
Blumei, Steud. Nom. ed. II. ii. 139 = M. Champaca.
caerulea, *DC. Syst.* i. 449 (sp. dub.).—Java.
Cathcartii, *Hook. f. & Thoms. Fl. Ind.* i. 79.—Reg. Himal.
Champaca, *Linn. Sp. Pl.* 536.—Malaya.
coerulea, Steud. Nom. ed. I. 533 = caerulea.
Doltsopa, Buch.-Ham. ex DC. Syst. i. 448 = M. Champaca.
ecicatrisata, *Miq. Fl. Ind. Bat. Suppl.* 368.—Sumatra.
evonymoides, Burm. f. Fl. Ind. 124 = M. Champaca.
excelsa, *Blume, Fl. Jav. Magnol.* 9, in obs.—Reg. Himal.
Figo, *Spreng. Syst.* ii. 643.—China.
fuscata, *Blume, Fl. Jav. Magnol.* 8, in textu.—China.
glauca, Wight, Illustr. i. 14 = nilagirica.
gracilis, Kostel. Allg. Med. Pharm. Fl. v. 1701 = Magnolia Kobus.
Kisopa, *Buch.-Ham. ex DC. Syst.* i. 448.—Reg. Himal.
lactea, Buch.-Ham. ex Wall. Cat. n. 6491 = oblonga.
lanuginosa, *Wall. Tent. Fl. Nep.* 8. t. 5.—Reg. Himal.
longifolia, Blume, Bijdr. 7 = M. Champaca.
macrophylla, D. Don, Prod. Fl. Nep. 226 = Magnolia pterocarpa.
montana, *Blume, Bijdr.* 7.—Reg. Himal.; Malaya.
nilagirica, *Zenker, Pl. Ind.* 21. t. 20.—Ind. or.
oblonga, *Wall. Cat.* n. 972.—Reg. Himal.
ovalifolia, Wight, Illustr. i. 13 = nilagirica.
parvifolia, *Blume, Fl. Jav. Magnol.* 8, in textu.—Java.
parvifolia, *DC. Syst.* i. 449 (sp. dub.).—Ins. Ternate.
pubinervia, Blume, Fl. Jav. Magnol. 14 = M. Champaca.
pulneyensis, Wight, Illustr. i. 14 = nilagirica.
punduana, *Hook. f. & Thoms. Fl. Ind.* i. 81.—Reg. Himal.
Rheedii, Wight, Illustr. i. 14. t. 5. f. 6 = M. Champaca.
rufinervis, DC. Syst. i. 449 = M. Champaca.
sericea, Pers. Syn. ii. 94 = M. Champaca.
suaveolens, Pers. l. c. = M. Champaca.
Tsiampaca, *Linn. Mant.* i. 78.—Amboina.
velutina, Blume, Fl. Jav. Magnol. 17 = M. Tsiampaca.
velutina, DC. Prod. i. 79 = lanuginosa.
Walkeri, Wight, Illustr. i. 13 = nilagirica.
Yulan, Kostel. Allg. Med. Pharm. Fl. v. 1700 = Magnolia conspicua.

MICHIEA, F. Muell. Fragm. iv. 95. t. 27 (1864) = **Coleanthera**, Stschegl. (Epacrid.).
symphyanthera, F. Muell. l. c. 96 = C. myrtoides.

MICHOXIA, Vell. Fl. Flum. 227 (1825); v. t. 103 (1827). Gen. dub.—Cf. Benth. & Hook. f. i. 981.
calyptrata, Vell. Fl. Flum. 227; v. t. 103.—Bras.

MICONIA, Ruiz & Pav. Prod. 60 (1794). *MELASTOMACEAE*, Benth. & Hook. f. i. 763.
ANTISOLA, Rafin. Sylva Tellur. 97 (1838).
AUGUSTINEA, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 145.
AULIPHAS, Rafin. Sylva Tellur. 95 (1838).
CATACHAENIA, Griseb. in Goett. Abh. x. (1861) 54.
CHAENOPLEURA, Rich. ex DC. Prod. iii. 197 (1828).
CHILOPORUS, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 57.
CHITONIA, D. Don, in Mem. Wern. Soc. iv. (1823) 317.
CLASTILIX, Rafin. Sylva Tellur. 99 (1838).
CREMANIUM, D. Don, in Mem. Wern. Soc. iv. (1823) 310.
CYANOPHYLLUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 324.
CYATHANTHERA, Pohl, Pl. Bras. Ic. ii. 130 (1831).
DECARAPHE, Miq. Phytog. ii. 84 (1838–40).
DIPLOCHITA, DC. Prod. iii. 176 (1828).
EURYCHAENIA, Griseb. Fl. Brit. W. Ind. 258 1860); et in Goett. Abh. ix. (1861) 46.
FOLOMFIS, Rafin. Sylva Tellur. 97 (1838).
FOTHERGILLA, Aubl. Pl. Gui. i. 440. t. 175 (1775).
GALLASIA, Mart. ex DC. Prod. iii. 182 (1828).
GLOSSOCENTRUM, Crueg. in Linnaea, xx. (1847) 111.
HARTIGIA, Miq. in Linnaea, xviii. (1844) 284.

MICONIA :—
Hypoxanthus, Rich. ex DC. Prod. iii. 180 (1828).
Jucunda, Cham. in Linnaea, ix. (1834) 456.
Leonicenia, Scop. Introd. 212 (1777).
Octomeris. Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 378.
Pholomphis, Rafin. Sylva Tellur. 97 (1838).
Pleurochaenia, Griseb. in Goett. Abh. ix. (1861) 55.
Pogonorhynchus, Crueg. in Linnaea, xx. (1847) 107.
Pomereula, Domb. ex DC. Prod. iii, 190 (1828).
Schizanthera, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 322.
Sericola, Rafin. Sylva Tellur. 93 (1838).
Soltmannia, [Klotzsch] ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 132.
Synoptera, Rafin. Sylva Tellur. 98 (1838).
Tamonea, Aubl. Pl. Gui. i. 441 (1775).
Truncaria, DC. Prod. iii. 106 (1828).
Ziegera, Rafin. Sylva Tellur. 94 (1838).
Zulatia, Neck. Elem. ii. 117 (1790).
abortiva, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 232 = Sieberi.
acalephoides, *Naud. l. c.* 207.—N. Granat.
Acinodendron, *Sweet*, *Hort. Brit.* ed. I. 159; *D. Don*, *in Lond. Hort. Brit.* 174.—Am. austr.
acuminata, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 244 = axilliflora.
acuminifera, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 105.—N. Granat.
adenophora, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 182 = aurea.
aegrotans, Naud. l. c. 211 = arbutifolia.
aeruginosa, Naud. l. c. 135 = desmantha.
affinis, *DC. Prod.* iii. 187.—Guian.
affinis, Macfad. ex Griseb. Fl. Brit. W. Ind. 257 = laevigata.
agrestis, Baill. Hist. Pl. vii. 35 = Leandra agrestis.
agyna, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 232 = caerulea.
alata, *DC. Prod.* iii. 184.—Am. trop.
albicans, Steud. Nom. ed. II. ii. 139 = holosericea.
albo-rufescens, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 160.—Bras.
alternans, *Naud. l. c.* 161 in adnot.—Guian.
alypifolia, *Naud. l. c.* 215.—Peruv.
amazonica, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 103.—Bras.
ambigua, *DC. Prod.* iii. 189.—Venezuela.
amblyandra, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 206.—N. Granat.
amoena, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 115.—Bras.
ampla, *Triana*, *l. c.* 101.—Ins. Trinit.
amplexicaulis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 136.—Peruv.
amygdalina, Naud. l. c. 141 = fasciculata.
anceps, Naud. l. c. 150 = scorpioides.
ancistrophora, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 103.—Cuba.
andina, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 236.—Venezuela.
androsaemifolia, *Griseb. Cat. Pl. Cub.* 100.—Cuba.
angustata, *D. Don*, *ex Steud. Nom.* ed. II. ii. 139.—Ins. Trinit.
angustiflora, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 246.—Jamaic.
angustifolia, Griseb. in Goett. Abh. vii. (1857) 217 = Tetrazygia angustifolia.
anisophylla, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
anisotricha, Triana, l. c. 102 = Berghesiana.
annulata, *Triana*, *l. c.* 121.—N. Granat.
aplostachya, *DC. Prod.* iii. 183.—Am. austr.
aponeura, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 106.—N. Granat.
appendiculata, *Triana*, *l. c.* 131.—N. Granat.
arborea, Pav. ex Triana, l. c. 108 = calvescens.
arbutifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 210.—Venezuela.
argentea, *DC. Prod.* iii. 182.—Ind. occ.
argyrophylla, *DC. l. c.* 181.—Bras.
argyrophylla, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 147 = stenostachya.
Aschenborniana, Naud. l. c. 246 = Oxymeris cornioides.

MICONIA :—
asclepiadea, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 121.—N. Granat.
aspergillaris, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 214.—Ecuador.
asperrima, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 124.—N. Granat.
astrolasia, DC. Prod. iii. 190 = rubiginosa.
astrotricha, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 113.—Bras.
atrata, *Wawra*, *in Oestr. Bot. Zeitschr.* xiii. (1883) 89.—Bras.
attenuata, DC. Prod. iii. 186 = prasina.
Aubletiana, Steud. Nom. ed. II. ii. 139 = prasina.
aulocalyx, *Mart. ex Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 115.—Peruv.
aurea, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Am. trop.
auriculata, *DC. Prod.* iii. 185.—Bras.
axilliflora, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 179.—Guiana.
barbigera, DC. Prod. iii. 179 = racemosa.
barbinervis, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 113.—Am. austr.
Benthamiana, *Triana*, *l. c.* 102.—Am. austr.
berbiceana, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 174 = M. Acinodendron.
Berghesiana, *Naud. l. c.* 246.—Mexic.
Bergii, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 129 = Conostegia xalapensis.
Bergii, Wawra, Bot. Ergeb. Maxim. Bras. 19. t. 4 = atrata.
Beurlingii, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 107.—Ins. Trinit.
bicolor, Triana, l. c. 103 (Tetrazygiae sp.).—Ind. occ.
bifrons, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 178 = lepidota.
biglomerata, *DC. Prod.* iii. 183.—Bras.
Blanchetiana, Naud. in Ann. Sc. Nat. Sér. III. xxi. (1851) 202 = hypoleuca.
brachyanthera, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 129.—Peruv.
brachybotrya, *Triana*, *l. c.* 111.—Venezuela.
brachycalyx, *Triana*, *l. c.* 102.—N. Granat.
brachypoda, DC. Prod. iii. 180 = racemosa.
brachystyla, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 230 = glaberrima.
bracteata, *Macfad. Fl. Jamaic.* ii. 82.—Jamaic.
bracteata, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 111.—Am. austr.
bracteolaris, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 150 = eriodonta.
bracteolata, *DC. Prod.* iii. 182.—Peruv.
brasiliensis, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 118.—Bras.
brevipes, *Benth. in Hook. Journ. Bot.* ii. (1840) 313.—Guiana.
brunnea, *DC. Prod.* iii. 184.—Bras.
bubalina, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Peruv.
budlejoides, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 118.—Bras.
bullata, *Triana*, *l. c.* 130.—Peruv.
Burchellii, *Triana*, *l. c.* 116.—Bras.
buxifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 208.—N. Granat.
caelata, *DC. Prod.* iii. 183.—Peruv.
caerulea, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Peruv.
calophylla, Crueg. in Linnaea, xx. (1847) 109 = ciliata.
calophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 129 = myrtiformis.
calvescens, *DC. Prod.* iii. 185.—Bras.
calyptrata, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 109 = Chamissois.
campestris, *Triana*, *l. c.* 113.—Guian.
Candolleana, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 244 = cinnamomifolia.
Candolleana, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 117.—Bras.
caryophyllacea, *Triana*, *l. c.* 106.—Bras.
castaneaefolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 132.—Bras.
cataractae, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 123.—N. Granat.
caudata, *DC. Prod.* iii. 187.—N. Granat.
caudata, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 184, partim = Doneana.

MICONIA :—
caudigera, *DC. Prod.* iii. 186.—Bras.
ceanothina, DC. l. c. 189 = laevigata.
cecidophora, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 166.—Peruv.
centrodesma, *Naud. l. c.* 164.—N. Granat.
centrophora, *Naud. l. c.* 209.—Guian.
cernua, *Naud. l. c.* 205.—N. Granat.
chaenopleuroides, Naud. l. c. 245 = latecrenata.
chaetodon, Naud. l. c. 134 = desmantha.
Chamissois, *Naud. l. c.* 179.—Bolivia.
chartacea, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 119.—Bras.
chionophila, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 236.—Peruv.
chionophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 130 = chionophila.
chlorophylla, Wright, in Sauv. Fl. Cub. 47 = delicatula.
chrysandra, Griseb. Cat. Pl. Cub. 99 = Graffenrieda chrysandra.
chrysoneura, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 111.—Mexic.
chrysopetala, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 233 = coelestis.
ciliaris, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 105.—Peruv.
ciliata, Benth. in Hook. Journ. Bot. ii. (1840) 311 = racemosa.
ciliata, *DC. Prod.* iii. 179.—Guian.
cinchonaefolia, *DC. l. c.* 184.—Bras.
cinerascens, *Miq. in Linnaea*, xxii. (1849) 543.—Bras.
cinnamomea, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 131 = Oxymeris cinnamomea.
cinnamomifolia, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 168 = Candolleana.
cinnamomifolia, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 101.—Ind. occ.
circumsecta, Macfad. Fl. Jamaic. ii. 77 = racemosa.
clathrantha, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 129.—Peruv.
Claussenii, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 143 = pepericarpa.
clidemioides, Steud. in Flora, xxvii. (1844) 724 = M. Acinodendron.
coelestis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Peruv.
coerulea, Auct., vide caerulea.
collina, DC. Prod. iii. 185 = prasina.
compressa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 128.—Bras.
congestiflora, DC. Prod. iii. 180 = ciliata.
conostegioides, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 127.—Bras.
corallina, *Spring*, *ex Mart. in Flora*, xx. (1837) II. *Beibl.* 77 = mexicana.
cordata, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 114.—Am. trop.
coriacea, *DC. Prod.* iii. 189.—Ins. Guadalup.
cornifolia, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 126 = cinnamomifolia.
coronata, *DC. Prod.* iii. 187.—Ecuador.
crassifolia, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 125.—Peruv.
crassipes, *Triana*, *l. c.* 116.—Peruv.
cremophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 228.—Peruv.
crinita, *Naud. l. c.* 194.—N. Granat.
crispula, Spruce, ex Naud. l. c. 172 = prasina.
cristulata, Naud. l. c. 177 = prasina.
crocea, *Naud. l. c.* 245.—Peruv.
Cruegeri, Triana, in Trans. Linn. Soc. xxviii. (1871) 131 = ciliata.
Cruegeriana, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 245 = hypoleuca.
cruenta, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
cubensis, *Wright*, *in Sauv. Fl. Cub.* 48.—Cuba.
cuneata, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 113.—Peruv.
cuspidata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 163.—Bras.
cyanocarpa, *Naud. l. c.* 221.—Bolivia.
Cyathanthera, *Triana*, *in Trans. Linn. Soc.* xxviii. (1871) 127.—Bras.
decussata, D. Don, ex Loud. Hort. Brit. 174 = ciliata.
delicatula, *A. Rich. Ess. Fl. Cub.* 566.—Cuba.
densiflora, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 245 = pusilliflora.

MICONIA:—
denticulata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 214.—N. Granat.
depauperata, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 346.—Bras.
depauperata, Triana, in Trans. Linn. Soc. xxviii. (1871) 117, partim = paniculata.
desmantha, *Benth. Pl. Hartw.* 181.—N. Granat.
detergibilis, DC. Prod. iii. 181 = holosericea.
detergibilis, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 149 = herpetica.
difficilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
dimorpha, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 245 = milleflora.
dipsacea, *Naud. l. c.* 138.—Peruv.
discolor, *DC. Prod.* iii. 114.—Bras.
discolor, Macfad. Fl. Jamaic. ii. 85 = punctata.
dispar, *Benth. in Hook. Kew Journ.* ii. (1850) 241.—Bras.
dispar, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 112 = eriodonta.
divaricata, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 345.—Bras.
divergens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 127.—N. Granat.
dolichopoda, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 217.—Venezuela.
dolichorrhyncha, *Naud. l. c.* 166.—Peruv.
Donaeana, *Naud. l. c.* 244.—Peruv.
dura, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 131.—Bras.
ecostata, *Sweet, Hort. Brit.* ed. I. 159 (sp. dub.).—Jamaica.
elaeagnoides, *Griseb. in Goett. Abh.* vii. (1857) 217.—Ind. occ.
Elaeodendron, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Bras.
elaeoides, *Naud. l. c.* 223.—N. Granat.
elata, *DC. Prod.* iii. 182.—Jamaica.
elliptica, Macfad. Fl. Jamaic. ii. 88 = obovalis.
emarginata, Ruiz & Pav. Syst. i. 106 = Graffenrieda emarginata.
erioclada, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 111.—Peruv.
eriodonta, *DC. Prod.* iii. 185.—Guian.
eriophora, Miq. Comm. Phyt. 86 = auriculata.
eriopoda, Benth. in Hook. Journ. Bot. ii. (1840) 311 = M. Acinodendron.
erythrantha, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 213.—Mexic.
erythropila, Steud. in Flora, xxvii. (1844) 723 = spondylantha.
eugenioides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 109.—Bras.
eurychaenioides, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 185 = elata.
excelsa, A. St. Hil. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 126 = inconspicua.
fallax, *DC. Prod.* iii. 181.—Bras.
fasciculata, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 583.—Bras.
fasciculata, Miq. in Linnaea, xxii. (1849) 543 = chartacea.
favosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 190.—Ins. S. Doming.
ferruginata, *DC. Prod.* iii. 181.—Bras.
ferruginea, *DC. l. c.* 182.—Ins. S. Doming.
flammea, *Casar. Nov. Stirp. Bras. Dec.* 85.—Bras.
Fleischeriana, Steud. in Flora, xxvii. (1844) 723 = prasina.
floribunda, *DC. Prod.* iii. 188.—Peruv.
florida, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Guian.
Fockeana, Miq. Comm. Phyt. ii. 88. t. 11. f. A = spondylantha.
foliosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 106.—N. Granat.
Fothergilla, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 119.—Mexic.
fragilis, *Naud. l. c.* 171.—Guian.
fulva, *DC. Prod.* iii. 180.—Guian.
furfuracea, *Griseb. in Goett. Abh.* ix. (1861) 54, in obs.—Ins. S. Doming.
galactantha, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Peruv.
Gardneriana, *Naud. l. c* 245.—Bras.

MICONIA:—
glaberrima, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Mexic.
glabra, *Macfad. Fl. Jamaic.* ii. 85.—Jamaic.
glandulosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Jamaic.
glaucescens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 115.—Peruv.
glaucocarpa, *Steud. Nom.* ed. II. ii. 140.—Mexic.
globulifera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 139.—Mexic.
globuliflora, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 129 = Sieberi.
glomerata, *Triana, l. c.* 110.—Peruv.
glossocentra, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 243 = minutiflora.
gonioclada, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 128.—Ecuador.
goniostigma, *Triana, l. c.* 124.—N. Granat.
gossypina, *Triana, l. c.* 123.—N. Granat.
Goudotii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 155.—N. Granat.
gracilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 107.—N. Granat.
grandifolia, *D. Don, in Loud. Hort. Brit.* 174.—Ins. Trinit.
granulosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 218.—N. Granat.
gratissima, *Benth. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 101.—Bras.
Grisebachii, *Triana, l. c.* 130.—Cuba.
guayaquilensis, *D. Don, in Mem. Wern. Soc.* iv. (1823) 316.—Ecuador.
haematostemon, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 193.—N. Granat.
haplobotrya, Naud. l. c. 131 = triplinervis.
haplostachya, C. Muell. in Walp. Ann. iv. 711 = aplostachya.
haplostachya, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 130 = macrostachya.
hartigioides, Naud. l. c. 245 = milleflora.
havanensis, DC. Prod. iii. 188 = laevigata.
heliotropoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 116.—Bras.
hemenostigma, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 230.—Mexic.
herpetica, *DC. Prod.* iii. 181.—Bras.
heterochroa, Miq. in Linnaea, xviii. (1844) 621 = holosericea.
heteromera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 124.—Bras.
hexaptera, Naud. l. c. 129 = triplinervis.
hirta, Baill. Hist. Pl. vii. 35 = Clidemia hirta.
hispidissima, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 191.—Peruv.
holosericea, *DC. Prod.* iii. 181.—Ind. occ.
holosericea, Triana, in Trans. Linn. Soc. xxviii. (1871) 101 = mucronata.
Hookeriana, *Triana, l. c.* 125.—Hab.?
hyemalis, *A. St. Hil. & Naud. ex Naud. in Ann. Sc Nat.* Sér. III. xvi. (1851) 142.—Bras.
hygrophila, *Naud. l. c.* 220.—Bolivia.
hymenanthera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 130.—Ecuador.
hypargyrea, Miq. in Linnaea, xviii. (1844) 622 = stenostachya.
hyperprasina, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 186.—Mexic.
hypoleuca, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 119.—Am. trop.
ibaguensis, Schlecht. in Linnaea, v. (1830) 562; ex Triana, l. c. 109 = Maximiliana.
ibaguensis, Triana, l. c. 110 = auriculata.
ibaguensis, *Wright, in Sauv. Fl. Cub.* 48.—Cuba.
igniaria, *Bonpl. ex Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 197.—Am. trop.
impetiginosa, DC. Prod. iii. 183 = rufescens.
impetiolaris, *D. Don, in Mem. Wern. Soc.* iv. (1823) 316.—Ind. occ.
inaequalifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 110.—Bras.
inaequidens, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Bras.
inaperta, *Naud. l. c.* 210.—Venezuela.
inconspicua, *Miq. in Linnaea*, xxii. (1849) 542.—Bras.
intricata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 105.—N. Granat.

MICONIA:—
iodopila, Steud. in Flora, xxvii. (1844) 725 = spondylantha.
ioneura, *Griseb. in Goett. Abh.* xxiv. (1879) 129.—Reg. Argent.
Jucunda, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 101.—Bras.
Kapplerii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 162.—Guian.
Klotzschii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 118.—Bras.
lacera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 152.—Am. trop.
laevigata, *DC. Prod.* iii. 188.—Ind. occ.
laevigata, Griseb. Fl. Brit. W. Ind. 257, partim = virescens.
Lambertiana, DC. Prod. iii. 185 = longifolia.
lamprarrhena, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
lamprophylla, *Triana, l. c.* 108.—N. Granat.
lanata, *Triana, l. c.* 113.—Am. trop.
lanceolata, *DC. Prod.* iii. 190.—Ins. S. Doming.
lanuginosa, *Ruiz & Pav. Syst. Veg.* i. 105.—Peruv.
lappacea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 111.—Guian.
latecrenata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 239.—Bras.
latifolia, *Naud. l. c.* 244.—Peruv.
lauriformis, *Naud. l. c.* 189.—N. Granat.
laurina, *Naud. l. c.* 245.—Peruv.
Lechleri, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 124.—Peruv.
ledifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 224.—Ecuador.
lepidota, *DC. Prod.* iii. 180.—Bras.
lepidota, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 162 = Kapplerii.
Leschenaultiana, Sagot, ex Naud. l. c. 325 = spondylantha.
leucocarpa, *DC. Prod.* iii. 182.—Bras.
leucocephala, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 244 = macrophylla.
Lhotzkyana, Triana, in Trans. Linn. Soc. xxviii. (1871) 151 = staminea.
ligustrina, Triana, l. c. 128 = rhamnoidea.
ligustroides, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 167.—Bras.
lilacina, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 129.—Peruv.
Lindenii, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 135 = desmantha.
lineata, Schlecht. in Linnaea, xiii. (1839) 423; ex Triana, in Trans. Linn. Soc. xxviii. (1871) 110 = auriculata.
livida, *Triana, l. c.* 121.—Peruv.
lonchophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 176.—Venezuela.
longifolia, *DC. Prod.* iii. 184.—Guian.
longifolia, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 157 = fulva.
longispicata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 117.—Bras.
longistyla, Seem. Bot. Voy. Herald, 123 = argentea.
longistyla, Steud. in Flora, xxvii. (1844) 724 = argyrophylla.
loreyoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 121.—N. Granat.
loxensis, *DC. Prod.* iii. 186.—Ecuador.
lucida, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 196.—Venezuela.
lutescens, *DC. Prod.* iii. 189.—Peruv.
macrantha, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 123.—N. Granat.
macrophylla, Macfad. Fl. Jamaic. ii. 84 = impetiolaris.
macrophylla, Steud. in Flora, xxvii. (1844) 723 = prasina.
macrophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 103 = platyhedra.
macrostachya, *DC. Prod.* iii. 190.—Bras.
macrostoma, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 235 = nitida.
macrothyrsa, *Benth. in Hook. Journ. Bot.* ii. (1840) 312.—Guian.
Magdalenae, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 115.—N. Granat.
magnifica, *Triana, l. c.* 131.—Mexic.
marginata, *Triana, l. c.* 110.—Bras.
Martiusiana, *DC. Prod.* iii. 186.—Bras.

MICONIA:—
Matthaei, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 176.—Peruv.
Maximilianea, *DC. Prod.* iii. 186.—Bras.
media, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Peruv.
melanodendron, Naud. l. c. 162 = minutiflora.
Melinonis, *Naud. l. c.* 125.—Guian.
mellina, *DC. Prod.* iii. 182.—Bras.
membranacea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 115.—Peruv.
meridensis, *Triana, l. c.* 102.—Venezuela.
metallica, *Triana, l. c.* 106.—Venezuela.
mexicana, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Mexic.
microbotrya, Naud. l. c. 195 = rubens.
microcarpa, *DC. Prod.* iii. 189.—Bras.
microcarpa, Macfad. ex Griseb. Fl. Brit. W. Ind. 257 = laevigata.
microcarpa, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 159 = cecidophora.
microcarpa, Triana, in Trans. Linn. Soc. xxviii. (1871) 111 = Klotzschii.
microphylla, Steud. ex Hemsl. Biol. Centr. Am. Bot. i. 428 = prasina.
milleflora, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 237.—N. Granat.
minutiflora, *DC. Prod.* iii. 189.—Venezuela.
miocarpa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 237.—N. Granat.
Miqueliana, Naud. l. c. 245 = M. Acinodendron.
mollicula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 113.—N. Granat.
mollis, *Triana, l. c.* 113.—Bras.
molybdea, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 185.—Bolivia.
montana, Crueg. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 116 = holosericea.
mucronata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 120.—Guian.
multispicata, *Naud. l. c.* 131.—Jamaic.
muricata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 102.—Peruv.
mutabilis, *Triana, l. c.* 112.—Guian.
myriantha, Benth. in Hook. Journ. Bot. ii. (1840) 314 = minutiflora.
myricoides, Triana, in Trans. Linn. Soc. xxviii. (1871) 129 = Sieberi.
myriocarpa, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 231 = glaberrima.
myrtiformis, *Naud. l. c.* 244.—Peruv.
myrtillifolia, *Naud. l. c.* 198.—Venezuela.
myura, Naud. l. c. 129 = Sancti-Philippi.
Naudiniana, Miq. Stirp. Surin. Sel. 50 = lepidota.
nemoralis, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 178 = pteropoda.
neriifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 127.—Peruv.
nervosa, Triana, l. c. 111 = spondylantha.
neurocarpa, *Triana, l. c.* 108.—N. Granat.
nicotianaefolia, DC. Prod. iii. 188 = laevigata.
nitens, Benth. in Hook. Journ. Bot. ii. (1840) 313 = Tococa Weddellii.
nitida, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Peruv.
notabilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 103.—N. Granat.
novemnervia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 216.—Venezuela.
obovalis, *Naud. l. c.* 183.—Jamaica.
obovata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 123.—N. Granat.
obscura, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Peruv.
obtusa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 103.—Cuba.
ochracea, *Triana, l. c.* 127.—N. Granat.
Olfersiana, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 123 = jucunda.
oligotricha, *Naud. l. c.* 245.—Mexic.
orcheotoma, *Naud. l. c.* 227.—N. Granat.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 345.—Bras.
ossaeiformis, Naud. in Ann. Sc. Nat. Sér. VIII. xvi. (1851) 212 = Sagraea ossaeaeformis.
otodesma, Naud. l. c. 205 = squamulosa.
pallens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 103.—Ind. occ.

MICONIA:—
paludosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Bras.
palustris, Macfad. Fl. Jamaic. ii. 89 = prasina.
pandurata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 108.—Bras.
paniculata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 245.—Bras.
papillosa, *Naud. l. c.* 216.—N. Granat.
paradoxa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 121.—Bras.
parviflora, Macfad. ex Griseb. Fl. Brit. W. Ind. 257 = laevigata.
parviflora, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 245 = pusilliflora.
pastoënsis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 126.—N. Granat.
paucidens, *DC. Prod.* iii. 186.—Bras.
pauciflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 115.—Bras.
pauciglandulosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 183.—Peruv.
paulensis, *Naud. l. c.* 187.—Bras.
pauperula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 102.—N. Granat.
Pavoniana, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 180.—Peruv.
pendulifolia, DC. Prod. iii. 187 = laevigata.
Pentlandii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 182.—Bolivia.
pepericarpa, *DC. Prod.* iii. 182.—Bras.
pernettifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 127.—N. Granat.
persicaefolia, Naud. & A. St. Hil. ex Triana, l. c. 119 = fasciculata.
phaeophylla, *Triana, l. c.* 113.—Bras.
phaeotricha, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 193.—Mexic.
pichinchensis, *Benth. Pl. Hartw.* 182.—Ecuador.
pileata, *DC. Prod.* iii. 180.—Bras.
pinetorum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 229.—Mexic.
piperifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 109.—N. Granat.
planinervia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 160.—Venezuela.
platyhedra, *Naud. l. c.* 244.—Am. trop.
plebeia, Naud. l. c. 170 = brevipes.
plethorica, *Naud. l. c.* 227.—N. Granat.?
pleurocarpa, Naud. l. c. 245 = M. Acinodendron.
plicata, Steud. Nom. ed. II. 115, sphalm. = pileata.
Plukenetii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 116.—Guian.
plumifera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 124.—Peruv.
Poeppigii, *Triana, l. c.* 107.—Peruv.
polyandra, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 346.—Bras.
polygama, Triana, in Trans. Linn. Soc. xxviii. (1871) 181 = rubens.
polyneura, *Triana, l. c.* 128.—N. Granat.
praecox, *Wright, in Sauv. Fl. Cub.* 47.—Cuba.
prasina, *DC. Prod.* iii. 188.—Ind. occ.
prasina, Triana, in Trans. Linn. Soc. xxviii. (1871) 109, partim = pteropoda.
procera, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 245 = argentea.
proteoides, A. St. Hil. & Naud. l. c. 141 = cinerascens.
psychrophila, *Naud. l. c.* 220.—N. Granat.
pterocaulon, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 114.—N. Granat.
pterophora, Miq. in Linnaea, xviii. (1844) 622 = alata.
pteropoda, *Benth. in Hook. Journ. Bot.* ii. (1840) 314.—Bras.
pteropoda, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 161 = alternans.
pubipetala, Miq. Stirp. Surin. Sel. 50 = verticilliflora.
pulverulenta, DC. Prod. iii. 190 = M. Tiri.
pulverulenta, Hook. Bot. Mag. t. 5411 = Hookeriana.
pulverulenta, *Ruiz & Pav. Syst. Veg.* i. 104.—Peruv.
punctata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1883) 268.—Porto Rico.
punctata, *D. Don, in Mem. Wern. Soc.* iv. (1823) 316.—Ind. occ.
punctulata, *Macfad. Fl. Jamaic.* ii. 84.—Jamaic.
purpurascens, DC. Prod. iii. 179 = Aciotis purpurascens.

MICONIA:—
pusilliflora, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 130 = Beurlingii.
pusilliflora, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 171 = depauperata.
pusilliflora, Triana, in Trans. Linn. Soc. xxviii. (1871) 120 = Gardneriana.
pustulata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 208.—N. Granat.
pyramidalis, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1883) 268.—Porto Rico.
pyramidalis, DC. Prod. iii. 188 = laevigata.
pyrifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 164.—Bras.
quadrangularis, *Naud. l. c.* 197.—Ind. occ.
quadrangularis, Triana, in Trans. Linn. Soc. xxviii. (1871) 130 = cubensis.
quadrifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 223.—Guiana.
quitensis, Benth. Pl. Hartw. 182 = papillosa.
racemifera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 119.—Bras.
racemosa, *DC. Prod.* iii. 179.—Guiana; Porto Rico.
racemosa, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 153, pro parte = ciliata.
radulaefolia, *Naud. l. c.* 243.—Bras.
Rangeliana, Wright, ex Griseb. Cat. Pl. Cub. 99 = Tetrazygia laxiflora.
reclinata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 209.—Peruv.
reducens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 106.—N. Granat.
Renggeri, Steud. in Flora, xxvii. (1844) 723 = fallax.
repando-crenata, Steud. l. c. 724 = prasina.
resima, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 226.—N. Granat.
reticulata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 123.—N. Granat.
revoluta, Benth. in Hook. Journ. Bot. ii. (1840) 313 = prasina.
revoluta, *Miq. in Linnaea*, xxii. (1849) 544.—Bras.
rhamnoidea, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 226.—N. Granat.
rhytidophylla, *Naud. l. c.* 139.—Guian.
Riedelii, *Naud. l. c.* 125.—Bras.
rigens, *Naud. l. c.* 225.—N. Granat.
rigida, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 130.—Jamaic.
rimalis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 238.—Bras.
riparia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 117.—Bras.
Rivoeriae, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 170 = trichotoma.
rostrata, *Bertol. Fl. Guatim.* 17.—Guatemala.
rotundifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 235.—Peruv.
rubens, *Naud. l. c.* 169.—Ind. occ.
rubescens, D. Don, in Loud. Hort. Brit. 174, nomen = rufescens?
rubiginosa, *DC. Prod.* iii. 183.—Ecuador.
rubricans, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 121.—N. Granat.
rufa, *Triana, l. c.* 130.—Cuba.
rufescens, *DC. Prod.* iii. 180.—Guian.
rufescens, Macfad. ex Griseb. Fl. Brit. W. Ind. 256 = holosericea.
rufipila, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 104.—Bras.
rugifolia, *Triana, l. c.* 125.—Peruv.
rugosa, *Triana, l. c.* 106.—Bras.
Ruizii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 192.—Peruv.
salicifolia, *Naud. l. c.* 234.—N. Granat.
Sancti-Philippi, *Naud. l. c.* 121.—Venezuela.
sanguinea, Triana, in Trans. Linn. Soc. xxviii. (1871) 125 = hispidissima.
scandens, Ruiz & Pav. Fl. Per. iv. t. 395 = Adelobotrys adscendens.
Schlimii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 102.—N. Granat.
Schomburgkii, Benth. in Hook. Journ. Bot. ii. (1840) 312 = lepidota.
sclerophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 119.—Peruv.?
scorpioides, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Am. trop.

MICONIA:—
Sellowiana, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 206.—Bras.
semicrenata, *D. Don, ex Loud. Hort. Brit.* 174, nomen.—Ins. Guadalup.
semicrenata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 126.—Ind. occ.
sepiaria, DC. Prod. iii. 185 = prasina.
serialis, *DC. l. c.* 182.—Bras.
serrulata, Macfad. ex Griseb. Fl. Brit. W. Ind. 258 = racemosa.
serrulata, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 118 = macrophylla.
serrulata, Triana, in Trans. Linn. Soc. xxviii. (1871) 129 = galactantha.
sessiliflora, C. Muell. in Walp. Ann. vi. 1225, sphalm. = seq.
sessilifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 181.—Bolivia.
setinodis, *DC. Prod.* iii. 179.—N. Granat.
Sieberi, *DC. Prod.* iii. 189.—Ins. Trinit.
simplex, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 104.—Ecuador.
smaragdina, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 186.—N. Granat.
Soltmanniana, Naud. l. c. 132 = corallina.
sordida, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
speciosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Bras.
spennerostachya, *Naud. l. c.* 187.—Peruv.
spicata, Macfad. ex Griseb. Fl. Brit. W. Ind. 257 = triplinervis.
spicellata, *Bonpl. ex Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 134.—Bras.
spinulosa, *Naud. l. c.* 195.—Venezuela.
spireaefolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 122.—N. Granat.
splendens, Griseb. Fl. Brit. W. Ind. 256 = obovalis, trichotoma.
splendens, *Griseb. in Goett. Abh.* ix. (1861) 47.—Ind. occ.
splendens, Wright, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 109 = prasina.
spondylantha, *DC. Prod.* iii. 184.—Bras.
Sprucei, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 108.—Bras.
squamulosa, *Triana, l. c.* 128.—N. Granat.
staminea, *DC. Prod.* iii. 187.—Bras.
staminea, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 123, partim = jucunda.
staphidioides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 102.—Bolivia.
stellifera, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 142 = barbinervis.
stelligera, Naud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 113 = cuneata.
stenobotrys, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 240 = quadrangularis.
stenopetala, Griseb. Fl. Brit. W. Ind. 711 = Platycentrum clidemioides.
stenostachya, *DC. Prod.* iii. 181.—Mexic.
stenourea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 104.—N. Granat.
stictophylla, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 167 = spicellata.
stipularis, *Naud. l. c.* 194.—N. Granat.
strangulata, Naud. l. c. 117 = coronata.
subciliata, DC. Prod. iii. 187 = Tococa subciliata.
subcrustulata, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 130 = Conostegia purpurea.
subnodosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 106.—N. Granat.
sylvatica, Schlecht. in Linnaea, xiii. (1839) 422; ex Triana, l. c. 107 = laevigata.
symphyandra, *Triana, l. c.* 126.—N. Granat.
symplocoidea, *Triana, l. c.* 108.—N. Granat.
tentaculifera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 241.—Bras.
tenuifolia, *D. Don, ex Loud. Hort. Brit.* 174, nomen.—Am. austr.
tenuis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 120.—N. Granat.
Terera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 196.—Peruv.
ternatifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 118.—Peruv.
tetrandra, *D. Don, in Loud. Hort. Brit.* 174; *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Ind. occ.
tetrastoma, *Naud. l. c.* 236.—Cuba.

MICONIA:—
Teysmanniana, Regel, Ind. Sem. Hort. Petrop. (1866) 105; et Gartenfl. (1867) t. 537 = staminea.
theaeforme, Schrank & Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 109 = Maximilianea.
Thomasiana, *DC. Prod.* iii. 189.—Ins. S. Thomas.
thyrsiflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 130.—Peruv.
thyrsoidea, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 244.—Peruv.
tiliaefolia, *Naud. l. c.* 151.—Bolivia.
tinctoria, Naud. l. c. 229 = media.
tinifolia, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 130 = andina.
tinifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 225.—Venezuela.
Tiri, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 127.—Peruv.
Titanea, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 203 = molybdea.
tococoides, Naud. l. c. 127 = mexicana.
tomentosa, *D. Don, in Mem. Wern. Soc.* iv. (1823) 316.—Guian.
trichodes, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 112.—Guian.
trichotoma, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1883) 268 = prasina.
trichotoma, *DC. Prod.* iii. 188.—Ins. Martinic.
trinervia, *D. Don, ex Loud. Hort. Brit.* 174.—Jamaic.
trinervis, Griseb. Fl. Brit. W. Ind. 257 = triplinervis.
Trinitatis, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 246 = minutiflora.
triplinervis, *Ruiz & Pav. Syst. Veg.* i. 105.—Peruv.
triseriata, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 145 = ferruginata.
tristis, *Spring, ex Mart. in Flora*, xx. (1837) II. *Beibl.* 76.—Bras.
truncata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 104.—Bras.
tuberculata, *Triana, l. c.* 102.—Am. trop.
tunicata, Griseb. ex Triana, l. c. 115 = glaucescens.
tunicata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 243.—Peruv. ?
ulmarioides, Naud. l. c. 211 = arbutifolia.
umbilicata, Triana, in Trans. Linn. Soc. xxviii. (1871) 131 = Melastoma umbilicata.
umbrifera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 116.—Peruv.
undata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 107.—Bras.
urophylla, *DC. Prod.* iii. 186.—Bras.
urticoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 123.—N. Granat.
uvifera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 222.—Bolivia.
vaccinioides, *Naud. l. c.* 233.—Bras.
Valtherii, *Naud. l. c.* 155.—Bras.
velutina, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 113.—N. Granat.
venustula, Benth. ex Triana, l. c. 118 = brasiliensis.
vernicosa, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 191 = Thomasiana.
verruculosa, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 120 = racemosa.
versicolor, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1851) 189.—N. Granat.
verticillata, *Macfad. Fl. Jamaic.* ii. 78.—Jamaic.
verticillata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 117.—Bras.
verticilliflora, *Steud. ex Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 122.—N. Guin.
villosissima, Crueg. in Linnaea, xx. (1847) 109 = lacera.
virescens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 107.—Ind. occ.
vismioides, *Triana, l. c.* 124.—Bras.
vulcanica, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 213.—Ins. Guadalup.
Weddellii, *Naud. l. c.* 144.—Bras.
Willdenowii, *Klotzsch, ex Naud. l. c.* 199.—Bras.
Wrightii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 103.—Cuba.
Wydleriana, DC. Mém. Mélast. 77 = impetiolaris.

MICONIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 341 = **Tetrazygia**, Rich. (Melastom.).
Lambertianum, Naud. l. c. (= *Miconia bicolor*).—Jamaic.

MICRACTIS, DC. Prod. v. 619 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 363.
Bojeri, *DC. Prod.* v. 620.—Madag.

MICRADENIA, Miers, Apocyn. S. Am. 158 (1878) = **Diplademia**, A. DC. (Apocyn.).
acuminata, Miers, l. c. 162 = D. acuminata.
atroviolacea, Miers, l. c. 159 = D. atroviolacea.
crassinoda, Miers, l. c. 158 = D. crassinoda.
fragrans, Miers, l. c. 162 = D. fragrans.
hirsutula, Miers, l. c. 160.—Bras.
Martiana, Miers, l. c. 161 = D. Martiana.
Moricandiana, Miers, l. c. 162 = D. Moricandiana.
nodulosa, Miers, l. c. 159 = D. Martiana.
Riedelii, Miers, l. c. 160 = D. Riedelii.
Sellowii, Miers, l. c. 161 = D. Sellowii.
splendens, Miers, l. c. 163 = D. splendens.
urophylla, Miers, l. c. 161 = D. urophylla.

MICRAEA, Miers, Trav. Chili, ii. 529 (1826). *GEN. DUB.*—Cf. Benth. & Hook. f. ii. 803.
pulchella, Miers, Trav. Chili, ii. 529, nomen.—Chili.

MICRAIRA, F. Muell. Fragm. v. 208 (1866). *GRAMINEAE*, Benth. & Hook. f. iii. 1156.
subulifolia, *F. Muell. Fragm.* v. 208.—Austral.

MICRAMPELIS, Rafin. in Med. Repos. N. York, v. (1808) 350 = **Sicyos**, Linn. (Cucurb.).
echinata, Rafin. l. c. = S. angulata.

MICRANDRA, Benth. in Hook. Kew Journ. vi. (1854) 371. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 289.
POGONOPHYLLUM, Didr. in Kjoeb. Vidensk. Meddel. (1857) 144.
bracteosa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 290.—Bras.
Cunuri, Baill. ex Muell. Arg. in DC. Prod. xv. II. 1123 = Cunuria Spruceana.
elata, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 142.—Bras.
major, Baill. Adansonia, iv. (1863-64) 286 = siphonioides.
minor, Benth. in Hook. Kew Journ. vi. (1854) 372 = siphonioides.
siphonioides, *Benth. l. c.* 371.—Bras.

MICRANDRA, R. Br. in Benn. Pl. Jav. Rar. 237 (1844) = **Hevea**, Aubl. (Euphorb.).
ternata, R. Br. l. c. = H. discolor.

MICRANGELIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 388 = **Selinum**, Linn. (Umbell.).
pyrenaea, Fourr. l. c. = S. pyrenaeum.

MICRANTHEA, A. Juss. Tent. Euph. 24 (1824) = **Micrantheum**, Desf. (Euphorb.).
hexandra, Walp. Ann. i. 630, sphalm. = Micrantheum hexandrum.

MICRANTHELLA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 347 = **Tibouchina**, Aubl. (Melastom.).
Candollei, Naud. l. c. 352 (= *Pleroma exappendiculatum*).—Peruv.
capitata, Naud. l. c. 351 (= *Pleroma capitatum*).—Bolivia.
ciliaris, Naud. l. c. 348 (= *Pleroma ciliare*).—N. Granat.
citrina, Naud. l. c. 350 (= *Pleroma citrinum*).—Bolivia.
clinopodifolia, Naud. l. c. 348 = T. clinopodifolia.
confertiflora, Naud. l. c. 349 (= *Pleroma confertiflorum*).—Peruv.
Gayana, Naud. l. c. 350 (= *Pleroma Gayanum*).—N. Granat.
havanensis, Naud. l. c. 352 = T. longifolia.
hispida, Naud. l. c. 349 = T. longifolia.
lanceolata, Naud. l. c. 348 = T. longifolia.
latifolia, Naud. l. c. 351 (= *Pleroma latifolium*).—Bolivia.
longifolia, Naud. l. c. 352 = T. longifolia.
Orbignyana, Naud. l. c. 350 (= *Pleroma Orbignyanum*).—Bolivia.
rosea, Naud. l. c. 349 = T. longifolia.
stenopetala, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 46 (= *Pleroma stenopetalum*).—Am. austr.

MICRANTHEMUM, Michx. Fl. Bor. Am. i. 10. t. 2 (1803). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 956.
AMPHIOLANTHUS, Griseb, Cat. Pl. Cub. 186 (1866).
GLOBIFERA, J. F. Gmel. Syst. 32 (1791), nomen prius.
HEMIANTHUS, Nutt. in Journ. Acad. Philad. i. (1817) 119. t. 6.
PINARDA, Vell. Fl. Flum. 23 (1825).
callitrichoides, *Wright, in Sauv. Fl. Cub.* 101.—Cuba.
ciliolatum, *Wright, l. c.* 102.—Cuba.
elatinoides, *Wright, l. c.* 101.—Cuba.
emarginatum, Ell. Sketch, 18 = orbiculatum.
indicum, Hook. f. & Thoms. in Hook. Kew Journ. ix. (1857) 245 = Bythophyton indicum.
micrantha, Wood, Class-Book (1861), 525 = Nuttallii?
Nuttallii, *A. Gray, Man. Bot. N. U. St.* ed. V. 331.—Am. bor.
orbiculatum, *Michx. Fl. Bor. Am.* i. 10. t. 2.—Am. bor. et austr.
pilosum, *Ernst, in Flora*, lvii. (1874) 215.—Venezuela.
reflexum, *Wright, in Sauv. Fl. Cub.* 103.—Cuba.
rotundatum, *Griseb. Cat. Pl. Cub.* 184.—Cuba.
tetrandum, *Wright, in Sauv. Fl. Cub.* 102.—Cuba.
Tweedii, Benth. in DC. Prod. x. 423 = orbiculatum.

MICRANTHERA, Choisy, in Mém. Soc. Hist. Nat. Par. i. (1823) 224. tt. 11, 12 = **Tovomita**, Aubl. (Guttif.).
clusiaefolia, DC. Prod. i. 560 = seq.
clusioides, Choisy, in Mém. Soc. Hist. Nat. Par. i. (1823) 224. tt. 11, 12 = T. clusiaefolia.

MICRANTHES, Bertol. in Mem. Acc. Sc. Bolog. ix. (1858) 172; (Misc. Bot. xix. 8. t. 4) = **Hoslundia**, Vahl (Labiat.).
menthoides, Bertol. ll. cc. = H. decumbens.

MICRANTHES, Haw. Syn. Pl. Succ. 320 (1812) = **Saxifraga**, Tourn.
hieracifolia, Haw. Rev. Saxif. Suppl. i. 4 = S. pensylvanica.
hirta, Haw. Enum. Saxif. 46 = S. pensylvanica.
pensylvanica, Haw. l. c. 45 = S. pensylvanica.
semipubescens, Haw. Syn. Pl. Succ. App. 321 = S. pensylvanica.

MICRANTHEUM, Desf. in Mém. Mus. Par. iv. (1818) 253. t. 14. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 262.
CALETIA, Baill. Etud. Gén. Euph. 553. t. 26 (1858).
MICRANTHEA, A. Juss. Tent. Euph. 24 (1824).
boroniaceum, F. Muell. Fragm. i. 32 = ericoides.
ericoides, *Desf. in Mém. Mus. Par.* iv. (1818) 253. t. 14.—Austral.
hexandrum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 283.—Austral.
inversum, Panch. ex Baill. Adansonia, ii. (1861–62) 237 = Phyllanthus Fagueti.
triandrum, Hook. in Mitch. Journ. Trop. Austral. 342 = Phyllanthus Mitchelli.

MICRANTHEUM, Presl, Symb. Bot. i. 47 (1830) = **Trifolium**, Linn. (Legumin.).
glomeratum, Presl, l. c. 48 = T. glomeratum.
suffocatum, Presl, l. c. 48 = T. suffocatum.

MICRANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = seq.

MICRANTHUS, Eckl. Top. Verz. 43 (1827). *IRIDEAE*, Benth. & Hook. f. iii. 706.
alopecuroideus, *Eckl. Top. Verz.* 43.—Afr. austr.
fistulosus, Eckl. l. c. = alopecuroideus.
plantagineus, Eckl. l. c. = alopecuroideus.
spicatus, Heynh. Nom. ii. 405 = Watsonia punctata.
triticeus, Heynh. l. c. = Watsonia punctata.

MICRANTHUS, Loud. Hort. Brit. i. 314 (1830) = MARCANTHUS, Lour. = **Mucuna**, Adans. (Legumin.).

MICRANTHUS, Wendl. Bot. Beobacht. 38, 39 (1798) = **Phaylopsis**, Willd. (Acanthac.).
oppositifolia, Wendl. l. c. 39 = P. parviflora.
serpyllifolius, Roth, Nov. Pl. Sp. 282 = P. parviflora.

MICRARGERIA, Benth. in DC. Prod. x. 509 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 971.
GERARDIANELLA, Klotzsch, in Peters, Reise Mossamb. Bot. 229. t. 36 (1861).
Wightii, *Benth. in DC. Prod.* x. 509.—Ind. or.

MICRASTER, Harv. Gen. S. Afr. Pl. ed. II. 242 (1868) = **Brachystelma**, R. Br. (Asclep.).
pulchellus, Harv. l. c.—Afr. austr.

MICRAUCHENIA, Froel. in DC. Prod. vii. 239 (1838) = **Lactuca**, Tourn. (Compos.).
hispida, Froel. l. c. = L. Dubyaea.

MICRECHITES, Miq. Fl. Ind. Bat. ii. 456 (1856). *APOCYNACEAE*, Benth. & Hook. f. ii. 714.
elliptica, *Hook. f. Fl. Brit. Ind.* iii. 671.—Reg. Himal.
polyantha, *Miq. Fl. Ind. Bat.* ii. 457.—Java.
Schrieckii, *Vidal, Phan. Cuming. Philipp.* 26, 126.—Ins. Philipp.

MICRELIUM, Forsk. Fl. Aegypt. Arab. 152 (1775) = **Eclipta**, Linn. (Compos.).
asteroides, Forsk. l. c. 153 = E. erecta.
Tolax, Forsk. l. c. = E. erecta.

MICROBLEPHARIS, M. Roem. Syn. Pepon. 133, 200 (1846) = **Modecca**, Lam. (Passifl.).
acuminata, M. Roem. l. c. 200 = Modecca acuminata.
aliena, M. Roem. l. c. 202 (Asclepiad.).—Ind. or.
cordifolia, M. Roem. l. c. = Modecca cordifolia.
heterophylla, M. Roem. l. c. 200 = Modecca heterophylla.
macrophylla, M. Roem. l. c. 202 = Modecca macrophylla.
obtusa, M. Roem. l. c. 200 = Modecca obtusa.
Wightiana, M. Roem. l. c. = Modecca Wightiana.

MICROCACHRYS, Hook. f. in Hook. Lond. Journ. Bot. iv. (1845) 149. *CONIFERAE*, Benth. & Hook. f. iii. 433.
tetragona, *Hook. f. in Hook. Lond. Journ. Bot.* iv. (1845) 150.—Tasmania.

MICROCAELIA, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 285 (1851) = **Angraecum**, Thou. (Orchid.).
taeniophyllum, Hochst. ex A. Rich. l. c. = A. Guyonianum.

MICROCALA, Hoffmgg. & Link, Fl. Port. i. 359 (1806). *GENTIANEAE*, Benth. & Hook. f. ii. 806.
FRANQUEVILLIA, Salisb. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 338 (1821).
ISCHALEON, Ehrh. Beitr. iv. 147 (1789).
filiformis, Colm. Cat. Pl. Cataluña, 108 = Cicendia pusilla.
filiformis, *Hoffmgg. & Link, Fl. Port.* i. 359.—Europ.
pulchella, G. Don, Gen. Syst. iv. 213 = Erythraea ramosissima?
pusilla, G. Don, l. c. = Cicendia pusilla.
quadrangularis, *Griseb. in DC. Prod.* ix. 63.—Am. bor. et austr.

MICROCALIA, A. Rich. Fl. Nouv. Zél. 230. t. 30 (1832) = **Lagenophora**, Cass. (Compos.).
australis, A. Rich. l. c. 231 = L. Forsteri.

MICROCARPAEA, R. Br. Prod. 435 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 957.
alterniflora, Blume, Bijdr. 744 = muscosa.
americana, Spreng. Syst. i. 42 = Herpestis chamaedryoides.
cochlearifolia, Sm. in Rees, Cycl. xxiii. n. 2 = Peplidium humifusum.
cuneifolia, Hook. ex Wall. Cat. n. 3196 = Glossostigma spathulatum.
diandra, Griff. Notul. iv. 101 = Peplidium humifusum?
montevidensis, Spreng. Syst. i. 42 = Scoparia flava.
muscosa, *R. Br. Prod.* 436.—As. et Austral. trop.
spathulata, Hook. Bot. Misc. ii. (1831) 101 = Glossostigma spathulatum.

MICROCASIA, Becc. in Bull. Soc. Tosc. Ortic. iv. (1879) 180. *AROIDEAE*, Benth. & Hook. f. iii. 986.
elliptica, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 299.—Borneo.
pygmaea, *Becc. l. c.* 180, 299.—Borneo.

MICROCERASUS, M. Roem. Syn. Rosifl. 93 (1847) = **Prunus**, Tourn. (Rosac.).
aspera, M. Roem. l. c. 96 = P. aspera.
glandulosa, M. Roem. l. c. 95 = P. japonica.
humilis, M. Roem. l. c. 94 = P. prostrata.

MICROCERASUS:—
incana, M. Roem. Syn. Rosifl. 94 = Prunus prostrata.
incisa, M. Roem. l. c. 95 = P. incisa.
japonica, M. Roem. l. c. = P. japonica.
orientalis, M. Roem. l. c. 96 = P. microcarpa.
pectinata, M. Roem. l. c. 94 = P. Puddum.
Phoshia, M. Roem. l. c. 93 = P. Puddum.
prostrata, M. Roem. l. c. = P. prostrata.
sinensis, M. Roem. l. c. 95 = P. japonica.

MICROCHAETA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 450 = **Lipochaeta**, DC. (Compos.).
cavarum, Walp. Rep. ii. 620, lapsu = L. lavarum.
connata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 452 = L. connata.
hastulata, Walp. Rep. ii. 620 = L. hastulata.
integrifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 451 = L. integrifolia.
lanceolata, Nutt. l. c. = L. connata.
lavarum, Nutt. l. c. = L. lavarum.
lobata, Nutt. l. c. = L. hastulata.
procumbens, Nutt. l. c. 452 = Chrysogonum procumbens.
strigulosa, Nutt. l. c. = Wedelia aristata.
succulenta, Nutt. l. c. 451 = L. succulenta.

MICHROCHAETA, Reichb. Nom. Ind. 192 (1841) = **Rynchospora**, Vahl (Cyperac.).

MICROCHAETE, Benth. Pl. Hartw. 209 (1845) = **Senecio**, Tourn. (Compos.).
corymbosa, Benth. l. c. 196.—N. Granat.
glabrata, Benth. l. c. 210.—N. Granat.
humilis, Sch. Bip. in Flora. xxxviii. (1855) 121.—Reg. Magellan.
pulchella, Benth. Pl. Hartw. 210 = S. pulchellus.
teretifolia, Benth. l. c. 209 = S. teretifolius?
trichopus, Benth. l. c. 210.—N. Granat.
vaccinioides, Benth. l. c. = S. vaccinioides.

MICROCHARIS, Benth. in Trans. Linn. Soc. xxv. (1865) 297. t. 33. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
angolensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 162.—Afr. trop.
latifolia, *Benth. in Trans. Linn. Soc.* xxv. (1865) 298.—Afr. trop.
tenella, *Benth. l. c.* 297.—Afr. trop.

MICROCHILUS, Presl, Rel. Haenk. i. 94 (1827) = **Physurus**, Rich. (Orchid.).
bifalcis, D. Dietr. Syn. Pl. v. 166 = P. bifalcis.
Blumei, D. Dietr. l. c. = P. Blumei.
debilis, D. Dietr. l. c. = P. debilis.
densiflorus, D. Dietr. l. c. = P. densiflorus.
foliosus, D. Dietr. l. c. = P. foliosus.
hirtellus, D. Dietr. l. c. = P. hirtellus.
major, Presl, Rel. Haenk. i. 94 = P. Preslei.
minor, Presl, l. c. = P. minor.
ovatus, D. Dietr. Syn. Pl. v. 166 = P. ovatus.
pauciflorus, D. Dietr. l. c. = P. pauciflorus.
pictus, D. Dietr. l. c. = P. pictus.
plantagineus, D. Dietr. l. c. = P. plantagineus.
querceticola, D. Dietr. l. c. 167 = P. querceticola.
repens, D. Dietr. l. c. 166 = P. repens.
roseus, D. Dietr. l. c. = P. roseus.

MICROCHLAENA, Wight & Arn. Prod. 70 (1834) = **Eriolaena**, DC. (Stercul.).
flavescens, Garcke, ex Mast. in Hook. f. Fl. Brit. Ind. i. 371 = E. quinquelocularis.
quinqueloculare, Wight & Arn. Prod. 71 = E. quinquelocularis.
spectabilis, Endl. ex Walp. Rep. i. 351 = E. spectabilis.

MICROCHLOA, R. Br. Prod. 208 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1163.
abyssinica, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 20, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 404.—Abyss.
caffra, *Nees, Fl. Afr. Austr.* i. 246.—Afr. austr.
capensis, Nees, ex Steud. Nom. ed. II. ii. 141 = caffra.
elongata, *R. Br. in Wall. Cat.* n. 3807.—Ind. or.
Kunthii, *Desv. Opusc.* 75.—Am. trop.
obtusiflora, *Benth. in Hook. Niger Fl.* 565.—Afr. trop.
setacea, *R. Br. Prod.* i. 208.—Reg. trop.

MICROCLISIA, Benth. in Benth. & Hook. f. Gen. i. Pars 1ma, Add. 435 (1862) = **Pleogyne**, Miers (Menisp.).
australis, Benth. Fl. Austral. i. 59 = P. Cunninghami.

MICROCNEMUM, Ung.-Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 280, 272. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 66.
fastigiatum, *Ung.-Sternb. in Atti Congr. Bot. Firenze*, 1874 (1876), 280.—Hispan.

MICROCOCCA, Benth. in Hook. Niger Fl. 503 (1849). *EUPHORBIACEAE*, Benth. & Hook. f. ii. 309.
Mercurialis, *Benth. in Hook. Niger Fl.* 503.—As. et Afr. trop.

MICROCOCOS, Phil. in Bot. Zeit. xvii. (1859) 362 = **Jubaea**, H. B. & K. (Palm.).
chilensis, Phil. l. c.; et in Anal. Univ. Chil. (1859) 651 = J. spectabilis.

MICROCODON, A. DC. Monog. Camp. 127. t. 19 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 556.
brevibracteatum, Buek, ex Eckl. et Zeyh. Enum. 377 = glomeratum.
Candollianum, Buek, l. c. = glomeratum, sparsiflorum.
depressum, *A. DC. in DC. Prod.* vii. 422.—Afr. austr.
glomeratum, *A. DC. Monog. Camp.* 127. t. 19.—Afr. austr.
hispidulum, Sond. in Harv. & Sond. Fl. Cap. iii. 565 = depressum.
lineare, *Buek, ex Eckl. & Zeyh. Enum.* 378.—Afr. austr.
longebracteatum, Buek, l. c. 378 = Treichelia longebracteata.
pygmaeum, Buek, l. c. 377 = glomeratum.
sparsiflorum, *A. DC. Monog. Camp.* 128.—Afr. austr.

MICROCOECIA, Hook. f. in Proc. Linn. Soc. i. (1845) 278; et in Trans. Linn. Soc. xx. (1847) 209 = **Elvira**, Cass. (Compos.).
repens, Hook. f. in Trans. Linn. Soc. l. c.—Ins. Galap.

MICROCOELIA, Lindl. Gen. et Sp. Orch. 60 (1830) = **Mystacidium**, Lindl. (Orchid.).
exilis, Lindl. l. c. 61.—Madag.

MICROCORYS, R. Br. Prod. 502 (1810). *LABIATAE*, Benth. & Hook. f. ii. 1218.
ANISANDRA, Bartl. in Lehm. Pl. Preiss. i. 361 (1845).
barbata, *R. Br. Prod.* 502.—Austral.
brevidens, Benth. in DC. Prod. xii. 569 = glabra.
capitata, *Benth. l. c.* 568.—Austral.
ericifolia, *Benth. l. c.* 569.—Austral.
exserta, *Benth. Fl. Austral.* v. 124.—Austral.
glabra, *Benth. in DC. Prod.* xii. 569.—Austral.
lenticularis, *F. Muell. Fragm.* vi. 113.—Austral.
loganiacea, *F. Muell. l. c.*—Austral.
longiflora, *F. Muell. l. c.*—Austral.
longifolia, *Benth. in DC. Prod.* xii. 568.—Austral.
Macredieana, *F. Muell. Fragm.* viii. 231.—Austral.
obovata, *Benth. in DC. Prod.* xii. 569.—Austral.
parvifolia, Benth. l. c. = ericifolia.
pimeloides, *F. Muell. Fragm.* i. 156.—Austral.
purpurea, *R. Br. Prod.* 502.—Austral.
selaginoides, Bartl. in Lehm. Pl. Preiss. i. 363 = virgata.
subcanescens, *Benth. Fl. Austral.* v. 123.—Austral.
tenuifolia, *Benth. l. c.* 122.—Austral.
virgata, *R. Br. Prod.* 502.—Austral.

MICROCOS, Burm. ex Linn. Diss. Dass. 20 (1747); Amoen. Acad. i. 130 (1749) = **Grewia**, Linn. (Tiliac.).
begoniaefolia, G. Don, Gen. Syst. i. 551 = G. begonifolia.
glabra, Jack, in Malay. Misc. i. (1820) n. 1. 14 = G. Microcos.
involucrata, G. Don, Gen. Syst. i. 551 = G. involucrata.
lateriflora, Linn. Sp. Pl. 514 = G. tiliaefolia.
Mala, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 549 = G. Microcos.
paniculata, Linn. Sp. Pl. 514 = G. Microcos.
scabra, Sm. in Rees, Cycl. xxiii. n. 3 = G. scabra.
Stauntoniana, G. Don, Gen. Syst. i. 551 = G. Microcos.
tomentosa, Sm. in Rees, Cycl. xxiii. n. 2 = G. paniculata.

MICROCYBE, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 166. *RUTACEAE*, Benth. & Hook. f. i. 293.
albiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 167.—Austral.
multiflora, *Turcz. l. c.* 166.—Austral.
pauciflora, *Turcz. l. c.* 167.—Austral.

MICROCYCAS, A. DC. in DC. Prod. xvi. ii. 538 (1868). *CYCADACEAE*, Benth. & Hook. f. iii. 447.
calocoma, *A. DC. in DC. Prod.* xvi. ii. 538.—Cuba.

MICRODERIS, DC. Prod. vii. 127 (1838) = **Picris**, Linn. (Compos.).
filii, Hochst. ex Seub. Fl. Azor. 34. t. 13 = P. filii.
rigens, DC. Prod. vii. 128 = P. rigens.
umbellata, Hochst. ex Seub. Fl. Azor. 33. t. 12. f. 2, 2a.—Ins. Azor.

MICRODESMIS, Hook. f. in Hook. Ic. Pl. t. 758 (1844). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 287.
PENTABRACHION, Muell. Arg. in Flora, xlvii. (1864) 532.
caseariaefolia, *Planch. ex Hook. Ic. Pl.* sub t. 758.—Burma; Malaya.
puberula, *Hook. f. in Hook. l. c.* t. 758.—Afr. trop.

MICRODON, Choisy, in Mém. Soc. Phys. Genèv. ii. (1823) ii. 97 (Mém. Selag. 27). *SELAGINEAE*, Benth. & Hook. f. ii. 1129.
DALEA, Gaertn. Fruct. i. 235. t. 51 (1788).
cylindricus, *E. Mey. Comm. Pl. Afr. Austr.* 253.—Afr. austr.
linearis, Choisy, in DC. Prod. xii. 23 = ovatus.
Linkii, *Walp. Rep.* iv. 169.—Afr. austr.
lucidus, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1823) ii. 97.—Afr. austr.
orbicularis, *Choisy, in DC. Prod.* xii. 22.—Afr. austr.
ovatus, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1823) ii. 97.—Afr. austr.

MICRODONTA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 369 = **Heterospermum**, Cav. (Compos.).
nana, Nutt. l. c. 370.—Peruv.

MICROELUS, Wight & Arn. in Edinb. N. Phil. Journ. [xiv.] (Apr. 1833) 298 = **Bischofia**, Blume (Euphorb.).
Roeperianus, Wight & Arn. l. c. = B. javanica.

MICROGENETES, A. DC. in DC. Prod. ix. 292 (1845) = **Phacelia**, Juss. (Hydroph.).
Cumingii, A. DC. l. c. 293 = P. Cumingii.

MICROGLOSSA, DC. Prod. v. 320 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 282.
AMPHIRHAPIS, DC. Prod. vii. 279 (1838).
FRIVALDIA, Endl. Gen. 384 (1838).
HOMOSTYLIUM, Nees, in Linnaea, xviii. (1844) 513.
albescens, *C. B. Clarke, Comp. Ind.* 59.—Reg. Himal.
altissima, DC. Prod. v. 320 = Nidorella altissima.
angolensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 309.—Afr. trop.
cabulica, C. B. Clarke, Comp. Ind. 57 = albescens.
densiflora, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 200.—Afr. trop.
Griffithii, C. B. Clarke, Comp. Ind. 58 = albescens.
hispida, DC. Prod. v. 321 = Nidorella hispida.
mespiloides, *Benth. & Hook. f. Gen.* ii. 282.—Afr. austr.
mikanioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 182.—Madag.
petiolaris, *DC. Prod.* v. 321.—Afr. trop.
psiadioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 182.—Madag.
sessilifolia, *DC. Prod.* v. 320.—Madag.
volubilis, *DC. l. c.*—As. trop.
zeylanica, *Benth. & Hook. f. Gen.* ii. 282.—Ind. or.

MICROGYNE, Cass. in Dict. Sc. Nat. l. 493 (1827) = **Eriocephalus**, Linn. (Compos.).

MICROGYNE, Less. Syn. Comp. 190 (1832) = **Vittadinia**, Rich. (Compos.).
trifurcata, Less. l. c. = V. trifurcata.

MICROGYNOECIUM, Hook. f. in Benth. & Hook. f. Gen. iii. 56 (1880). *CHENOPODIACEAE*, Benth. & Hook. f. l. c.
tibeticum, *Hook. f. in Herb. Kew.*—Reg. Himal.

MICROJAMBOSA, Blume, Mus. Bot. Lugd. Bat. i. 117 (1849) = **Eugenia**, Linn. (Myrtac.).
besukiensis, Hassk. ex Miq. Fl. Ind. Bat. Suppl. 311 (= *Jambosa buxifolia*).—Ins. Banca.
conferta, Blume, Mus. Bot. Lugd. Bat. i. 118.—Borneo.
cuspidata, Blume, l. c. 119.—Java.
ferruginea, Blume, l. c. 118, in obs. = E. Thumra.
reticulata, Blume, l. c. = E. mangifolia.
splendens, Blume, l. c. 119.—Java.
trifida, Blume, l. c. 118.—Sumatra.

MICROKENTIA, H. Wendl. ex Benth. & Hook. f. Gen. iii. 895 (1883). *PALMAE*, Benth. & Hook. f. l. c.
Billardieri, *Benth. & Hook. f. l. c.*—N. Caled.
Deplanchei, *Benth. & Hook. f. l. c.*—N. Caled.
eriostachys, *Benth. & Hook. f. l. c.*—N. Caled.
gracilis, *Benth. & Hook. f. l. c.*—N. Caled.
Pancheri, *Benth. & Hook. f. l. c.*—N. Caled.
surculosa, *Benth. & Hook. f. l. c.*—N. Caled.

MICROLAENA, R. Br. Prod. 210 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1137.
DIPLAX, Soland. ex Benn. Pl. Jav. Rar. 11 (1838).
avenacea, *Hook. f. Handb. N. Zeal. Fl.* 320.—N. Zel.
distichophylla, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 296 = Tetrarrhena distichophylla.
Gunnii, Hook. f. Fl. Tasm. ii. 105 = stipoides.
polynoda, *Hook. f. Handb. N. Zeal. Fl.* 320.—N. Zel.
stipoides, *R. Br. Prod.* 210.—Austral.; N. Zel.
tasmanica, *Hook. f. ex Benth. Fl. Austral.* vii. 552.—Tasman.

MICROLAENA, Wall. Cat. n. 1173 (1832) = **Eriolaena**, DC. (Stercul.).
spectabilis, Wall. l. c. = E. spectabilis.

MICROLECANE, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 384 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
abyssinica, *Benth. & Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 386.—Abyss.

MICROLEPIDIUM, F. Muell. in Linnaea, xxv. (1852) 371 = **Capsella**, Moench (Crucif.).
fenestratum, F. Muell. ex Rupr. in Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 234 = C. procumbens.
pilosulum, F. Muell. in Linnaea, xxv. (1852) 371 = C. pilosula.

MICROLEPIS, Eichw. Casp. Cauc. 2 (1831) = **Anabasis**, Linn. (Chenop.).
salsa, Eichw. l. c. (= *Brachylepis salsa*).—Reg. Casp.

MICROLEPIS, Miq. Comm. Phytog. 71 (1839–40). *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
Mosenii, Cogn. in Mart. Fl. Bras. xiv. iii. 235 = quaternifolia, *Miq.*
oleaefolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 36. t. 2. f. 25.—Bras.
quaternifolia, *Cogn. in Mart. Fl. Bras.* xiv. iii. 236.—Bras.
quaternifolia, *Miq. in Linnaea*, xxii. (1849) 541.—Bras.
quaternifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 36 = Trianaei.
Trianaei, *Cogn. in Mart. Fl. Bras.* xiv. iii. 238.—Bras.

MICROLEPIS, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 157, 167 (1842) = **Cryptangium**, Schrad. et **Lagenocarpus**, Nees (Cyperac.).
bahiensis, Schrad. l. c. = L. Martii.
elatior, Schrad. l. c. = L. rigidus.
Minarum-Generalium, Schrad. l. c. = C. Minarum.
rigidifolia, Schrad. l. c. = L. topazinus.

MICROLICIA, D. Don, in Mem. Wern. Soc. iv. (1823) 301. *MELASTOMACEAE*, Benth. & Hook. f. i. 737.
IARAVAEA, Scop. Introd. 215 (1777).
acuminata, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 179 = Chaetostoma acuminatum.
agrestis, *Cogn. in Mart. Fl. Bras.* xiv. iii. 105.—Bras.
alsinefolia, DC. Prod. iii. 117 = Acisanthera alsinaefolia.

MICROLICIA:—
alternans, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 26 = serpyllifolia.
amplexicaulis, *Cogn. in Mart. Fl. Bras.* xiv. III. 82.—Bras.
arenariaefolia, Cham. in Linnaea, ix. (1834) 396 = multicaulis.
arenariaefolia, *DC. Prod.* iii. 120.—Peruv.
aristata, DC. Mém. Mélast. 30; et Prod. iii. 484 = setosa.
australis, Baill. Hist. Pl. vii. 9. f. 12 = Lavoisiera australis.
avicularis, *Mart. ex Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 176.—Bras.
baccharoides, *Mart. ex Naud. l. c.* 174.—Bras.
baccharoides, Naud. l. c. xii. (1849) 245 = myrtifolia.
balsamifera, *Mart. Nov. Gen. et Sp.* iii. 127, in obs.—Bras.
balsamifera, Triana, in Trans. Linn. Soc. xxviii. (1871) 27, partim = lavoisieraeformis.
Benthamiana, *Triana, l. c.* (Bentamiana).—Guian.
bivalvis, DC. Prod. iii. 117 = Acisanthera brevifolia.
Blanchetiana, Cogn. in Mart. Fl. Bras. xiv. III. 79 = lavoisieraeformis.
brevifolia, DC. Prod. iii. 117 = Acisanthera trivalvis.
Burchelliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 88.—Bras.
canastrensis, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 174.—Bras.
cardiophora, *Naud. l. c.* xii. (1849) 247.—Bras.
carnosula, *Naud. l. c.* 242.—Bras.
castrata, Naud. l. c. 230 = Chaetostoma castratum.
Chamaepitys, Naud. l. c. iii. (1845) 185 = juniperina.
Chamissois, *Naud. l. c.* xii. (1849) 263.—Bras.
chloracea, Naud. l. c. 252 = insignis.
ciliosa, Naud. l. c. iii. (1845) 179 = serrulata.
cinerea, *Cogn. in Mart. Fl. Bras.* xiv. III. 86.—Bras.
Clausseniana, *Cogn. l. c.* 54.—Bras.
confertiflora, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 176.—Bras.
cordata, *Cham. in Linnaea*, ix. (1834) 390.—Bras.
cordata, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 249 = confertiflora.
crenulata, *Mart. Nov. Gen. et Sp.* iii. 106. t. 251.—Bras.
cryptandra, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 175.—Bras.
cuneata, *Naud. l. c.* 180.—Bras.
cupressina, *D. Don, in Mem. Wern. Soc.* iv. (1823) 302.—Bras.
cuspidifolia, *Mart. Herb. Fl. Bras.* n. 937, nomen; *et ex Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 186.—Bras.
decipiens, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 232.—Bras.
decussata, *Naud. l. c.* iii. (1845) 175.—Bras.
denudata, *Cogn. in Mart. Fl. Bras.* xiv. III. 51.—Bras.
depauperata, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 253.—Bras.
doryphylla, *Naud. l. c.* iii. (1845) 181.—Bras.
doryphylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 27 = parvifolia.
elegans, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 187.—Bras.
ericoides, *D. Don, in Mem. Wern. Soc.* iv. (1823) 302.—Bras.
ericoides, Mart. Herb. Fl. Bras. n. 915 = Martiana.
euphorbioides, *Mart. Nov. Gen. et Sp.* iii. 107. t. 152.—Bras.
fasciculata, *Mart. Herb. Fl. Bras.* n. 936.—Bras.
ferruginea, *Scheele, in Linnaea*, xvii. (1843) 338.—Bras.
formosa, *Cham. in Linnaea*, ix. (1834) 393.—Bras.
Francavillana, *Cogn. in Mart. Fl. Bras.* xiv. III. 48.—Bras.
fulva, Cham. in Linnaea, ix. (1834) 391 = serpyllifolia.
Gardneri, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 240 = vestita.
glabra, *DC. Prod.* iii. 120.—Bras.
glossophylla, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 186 = formosa.
glutinosa, *Naud. l. c.* 180.—Bras.
grandiflora, Baill. Hist. Pl. vii. 34 = Lavoisiera grandiflora.
graveolens, *DC. Prod.* iii. 119.—Bras.

MICROLICIA:—
helvola, Triana, in Trans. Linn. Soc. xxviii. (1871) 25 = hypericoides.
heterophylla, Klotzsch, in Schomb. Faun. et Fl. Brit. Gui. 1016, 1100 = Rhynchanthera serrulata.
Hilariana, Miq. in Linnaea, xxii. (1849) 537 = serpyllifolia.
Hilariana, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 178.—Bras.
hirsuta, Triana, in Trans. Linn. Soc. xxviii. (1871) 25, in syn. = euphorbioides.
hirsutissima, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 182.—Bras.
hirto-ferruginea, *Naud. l. c.* 176.—Bras.
hispidula, *Naud. l. c.* 182.—Bras.
holosericea, *Naud. l. c.*—Bras.
humilis, *Naud. l. c.* 179.—Bras.
hypericoides, *Cham. in Linnaea*, ix. (1834) 389.—Bras.
imbricata, Cham. l. c. 395 = ericoides.
inquinans, *Naud. in. Ann. Sc. Nat.* Sér. III. iii. (1845) 172.—Bras.
insignis, *Cham. in Linnaea*, ix. (1834) 388.—Bras.
insignis, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 183 = linifolia.
inundata, DC. Prod. iii. 117 = Acisanthera inundata.
isophylla, *DC. l. c.* 120.—Bras.
jungermannioides, *DC. l. c.* 119.—Bras.
jungermannioides, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 173 = cordata.
juniperina, *A. St. Hil. Voy. Distr. Diam.* i. 373.—Bras.
lanceaeflora, DC. Prod. iii. 118 = Comolia lanceaeflora.
laniflora, Baill. Adansonia, xii. (1877) 95 = Trembleya laniflora.
lavoisieraeformis, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 232.—Bras.
leucantha, *Naud. l. c.* 239.—Bras.
Lhotzkyana, *Steud. Nom.* ed. II. ii. 142.—Bras.
limnobios, DC. Prod. iii. 117 = Acisanthera limnobios.
linifolia, *Cham. in Linnaea*, ix. (1834) 395.—Bras.
linophylla, D. Don, in Mem. Wern. Soc. iv. (1823) 301 = setosa.
loricata, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 188.—Bras.
macrophylla, *Naud. l. c.* 173.—Bras.
marifolia, D. Don, in Mem. Wern. Soc. iv. (1823) 303 = cordata.
marifolia, Herb. Berol. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 26 = serpyllifolia.
Martialis, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 248 = serpyllifolia.
Martiana, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 28.—Bras.
Maximowicziana, *Cogn. in Mart. Fl. Bras.* xiv. III. 94.—Bras.
Melastoma, Lhotzky & Manso, ex Walp. Rep. ii. 125 = insignis.
microphylla, *Cogn. in Mart. Fl. Bras.* xiv. III. 55.—Bras.
minutiflora, *Cogn. l. c.* 60.—Bras.
multicaulis, *Mart. Herb. Fl. Bras.* n. 932, nomen.—Bras.
myrtifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 245.—Bras.
myrtoidea, Benth. in Hook. Journ. Bot. ii. (1840) 294 = Benthamiana.
myrtoidea, *Cham. in Linnaea*, ix. (1834) 393.—Bras.
neglecta, *Cogn. in Mart. Fl. Bras.* xiv. III. 118.—Bras.
oblongifolia, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 177 = euphorbioides.
obtusa, *Naud. l. c.* 173.—Bras.
occidentalis, *Naud. l. c.* xii. (1849) 235.—Bras.
oligantha, *Naud. l. c.* 241.—Bras.
oxyanthera, Naud. l. c. 230 = Chaetostoma oxyantherum.
pallida, *Cogn. in Mast. Fl. Bras.* xiv. III. 49.—Bras.
parvifolia, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 177.—Bras.
passerina, Naud. l. c. xii. (1849) 244 = myrtifolia.
peruviana, *Cogn. in Mart. Fl. Bras.* xiv. III. 83.—Peruv.
pilosissima, *Cogn. l. c.* 116.—Bras.
polystemma, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 179.—Bras.
Pseudoscoparia, *Cogn. in Mart. Fl. Bras.* xiv. III. 60.—Bras.

MICROLICIA:—
pulchella, *Cham. in Linnaea*, ix. (1834) 392.—Bras.
pulchella, Mart. Herb. Fl. Bras. n. 916 = isophylla.
pulchella, Mart. l. c. n. 947 = serpyllifolia.
punctatissima, DC. Prod. iii. 118 = Acisanthera punctatissima.
pusilla, *Cogn. in Mart. Fl. Bras.* xiv. III. 53.—Bras.
recurva, DC. Prod. iii. 118 = Acisanthera recurva.
Regeliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 92.—Bras.
Reichardtiana, *Cogn. l. c.* 66.—Bras.
resinosa, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 27 = lavoisieraeformis.
resinosa, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 172. t. 7 = sincorensis.
Riedeliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 80.—Bras.
scoparia, Cham. in Linnaea, ix. (1834) 394 = M. Pseudoscoparia.
scoparia, *DC. Prod.* iii. 120.—Bras.
selaginea, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 188.—Bras.
Sellowiana, Naud. l. c. xii. (1849) 257 = setosa.
Seringeana, DC. Prod. iii. 119 = graveolens.
serpyllifolia, *D. Don, in Mem. Wern. Soc.* iv. (1823) 302.—Bras.
serrulata, *Cham. in Linnaea*, ix. (1834) 390.—Bras.
setosa, *DC. Prod.* iii. 120.—Bras.
setosa, Mart. Nov. Gen. et Sp. iii. t. 253. f. 1 = subsetosa.
sincorensis, *Mart. l. c.* 127 in obs.—Bras.
stenocladon, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 188 = cupressina.
stricta, *Cogn. in Mart. Fl. Bras.* xiv. III. 93.—Bras.
subsetosa, *DC. Prod.* iii. 119.—Bras.
subuliformis, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 185 = ericoides.
taxifolia, *Naud. l. c.* 183.—Bras.
ternata, *Cogn. in Mart. Fl. Bras.* xiv. III. 70.—Bras.
tetrasticha, *Cogn. l. c.* 80.—Bras.
tomentella, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 174.—Bras.
trembleyaeformis, *Naud. l. c.* 172.—Bras.
trichocalycina, *DC. Prod.* iii. 120.—Bras.
uncata, *Cogn. in Mart. Fl. Bras.* xiv. III. 64.—Bras.
variabilis, Mart. ex DC. Prod. iii. 118 = Acisanthera variabilis.
variolosa, DC. l. c. 119 = oligantha, euphorbioides, fasciculata.
variolosa, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 181 = oligantha.
venusta, Mart. Herb. Fl. Bras. n. 938 = cordata.
venusta, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 177 = serpyllifolia.
vestita, *DC. Prod.* iii. 119.—Bras.
viminalis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 28.—Bras.
violacea, DC. Prod. iii. 118 = Comolia violacea.
violacea, Mart. Herb. Fl. Bras. n. 950 = Comolia lanceaeflora.
virgata, *Cogn. in Mart. Fl. Bras.* xiv. III. 44.—Bras.
virgulata, DC. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 26 = serpyllifolia.
viscidula, Cham. in Linnaea, ix. (1834) 387 = Comolia sessilis.
Warmingiana, *Cogn. in Mart. Fl. Bras.* xiv. III. 47.—Bras.
Weddellii, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 251.—Peruv.

MICROLOBIUM, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 104 = **Apoplanesia**, Presl (Legumin.).
glandulosum, Liebm. 105 = A. paniculata.

MICROLOBIUS, Presl, Bot. Bemerk. 66 (1844); in Abh. Boehm. Ges. V. iii. (1845) 496. *LEGUMINOSAE*?—Cf. Benth. & Hook. f. i. 465.
mimosoides, *Presl, Bot. Bemerk.* 67.—Mexic.

MICROLOMA, R. Br. in Mem. Wern. Soc. i. (1809) 53. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 747.
angustifolium, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 64 = Leptadenia Spartum.
calycinum, *E. Mey. Comm. Pl. Afr. Austr.* 223.—Afr. austr.
glabratum, *E. Mey. l. c.* 222.—Afr. austr.
incanum, *Decne. in DC. Prod.* viii. 511.—Afr. austr.

MICROLOMA :—
lineare, *R. Br. in Mem. Wern. Soc.* i. (1809) 53.—Afr. austr.
pyrotechnicum, Spreng. Syst. i. 855 = Leptadenia Spartum.
sagittatum, *R. Br. in Mem. Wern. Soc.* i. (1809) 53.—Afr. austr.

MICROLONCHUS, Cass. in Dict. Sc. Nat. xliv. 38 (1826) = **Centaurea**, Linn. (Compos.).
albispinus, Bunge, Del. Sem. Hort. Dorp. (1843) 8; et in Linnaea, xviii. (1844) 155.—Persia; Beluchist.
cichoraceus, C. Koch, in Linnaea, xxiv. (1851) 416 = C. salmantica.
Clusii, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 166 = C. salmantica.
Delestrii, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 164.—Afr. bor.
Delilianus, Spach, l. c. 165 = C. salmantica.
divaricatus, DC. Prod. vi. 562 = Volutarella divaricata.
Duriaei, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 166.—Reg. Mediterr.
elatus, Spach, l. c. 168 = C. salmantica?
gracilis, Pomel, Nouv. Mat. Fl. Atl. 280.—Alger.
iserniánus, J. Gay & Webb, ex Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 466 = C. Janerii.
leptolonchus, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 168.—Afr. bor.
minimus, Boiss. Fl. Orient. iii. 701.—Persia.
oligochaetus, Boiss. l. c.—As. Min.; Persia.
pappósus, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 167.—Hab.?
persicus, DC. Prod. vi. 563 = C. Candolleana.
Reboudii, Pomel, Nouv. Mat. Fl. Atl. 279, in obs.—Alger.
salmanticus, DC. Prod. vi. 563 = C. salmantica.
spinulosus, Rouy, in Bull. Soc. Bot. Fr. xxxi. (1884) 277, nomen.—Hispan.
strictus, Pomel, Nouv. Mat. Fl. Atl. 279.—Alger.
tenellus, Spach, in Ann. Sc. Nat. Sér. III. iv. (1845) 165 = C. salmantica.
valdemorensis, Cut. Fl. Madrid, 420 = *Duriaei*.
yserniánus, J. Gay & Webb, ex Graells, Ind. Pl. Nov. 7, vide *iserniánus*.

MICROLOPHIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = **Polygala**, Tourn.
exile, Fourr. l. c. = P. exilis.

MICROLOPHUS, Cass. in Dict. Sc. Nat. xliv. 37 (1826) = **Centaurea**, Linn. (Compos.).
alatus, Cass. l. c. liv. 491 = C. Behen.

MICROMELES, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 168 = **Pyrus**, Tourn. (Rosac.).
castaneifolia, Decne. l. c. 169 = P. castaneifolia.
Griffithii, Decne. l. c. 170 = P. Griffithii.
khasiana, Decne. l. c. 169 = P. khasiana.
rhamnoides, Decne. l. c. = P. rhamnoides.
verrucosa, Decne. l. c. = P. cuspidata.

MICROMELUM, Blume, Bijdr. 137 (1825). *RUTACEAE*, Benth. & Hook. f. i. 303.
ceylanicum, Wight, Illustr. i. 109, in obs. = pubescens?
coriaceum, *Seem. Fl. Vit.* 32.—N. Caled.
diversifolium, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.—Malaya.
glabrescens, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 212 = pubescens.
glabrescens, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 36 = tephrocarpum.
hirsutum, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 40.—Burma; Malaya.
integerrimum, Wight & Arn. Prod. 94 = pubescens?
minutum, Wight & Arn. l. c. 448; Seem. Mission to Viti, 434 = pubescens.
molle, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 380.—Ins. Philipp.
monophyllum, Wight, Illustr. i. t. 42 = Paramignya monophylla.
octandrum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 578.—Cochinch.
pubescens, *Blume, Bijdr.* 138.—As. trop.; Ins. Pacif.
tephrocarpum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 379.—Ins. Philipp.
timoriense, Zipp. ex Span. in Linnaea, xv. (1841) 178 = pubescens.

MICROMERIA, Benth. in Bot. Reg. sub t. 1282 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1188.
APOZIA, Willd. ex Benth. in Linnaea, xi. (1837) 328.
CUSPIDOCARPUS, Spenn. in T. Nees, Gen. Fl. Germ. Gamop. ii. n. 18 (1843).
MICRONEMA, Schott, in Oestr. Bot. Wochenbl. (1857) 95.
PIPERELLA, Presl, Fl. Sic. p. xxxvii (1826).
SABBATIA, Moench, Meth. 386 (1794).
TENDANA, Reichb. f. Ic. Fl. Germ. xviii. 39. t. 1271 (1858).
XENOPOMA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399.
ZYGIS, Desv. in Ham. Prod. Pl. Ind. Occ. 46 (1825).
abyssinica, *Benth. in DC. Prod.* xii. 224.—Arab.; Abyssin.
affinis, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 274 = Mentha serpyllifolia.
alternipilosa, C. Koch, in Linnaea, xix. (1847) 25 = Satureia alternipilosa.
approximata, *Reichb. Fl. Germ. Excurs.* 859.—Europ. austr.
arkansana, Benth. Lab. Gen. et Sp. 730 = Calamintha Nuttallii.
australis, Benth. l. c. 380 = Mentha australis.
barbata, Boiss. & Kotschy, Diagn. Ser. II. iv. 14 = serpyllifolia, *Boiss.*
barbata, Fisch. & Mey. Ind. Sem. Hort. Petrop. viii. 67 = Douglasii.
Barceloi, Willk. in Oestr. Bot. Zeitschr. xxv. (1875) 111 = approximata.
Benthami, *Webb & Berth. Phyt. Canar.* iii. 77. t. 159.—Ins. Canar.
biflora, *Benth. Lab. Gen. et Sp.* 378.—Ind. or.; Arab.; Afr. trop.
boliviana, *Benth. l. c.* 731.—Bolivia.
bonariensis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* x. 56.—Reg. Argent.
Bourgaeana, *Webb, ex C. Bolle, in Bonplandia*, viii. (1860) 283.—Ins. Canar.
bracteolata, *Benth. Lab. Gen. et Sp.* 371.—Am. bor.
Brownei, *Benth. l. c.* 372.—Am. bor. et austr.
Buchii, Webb & Berth. Phyt. Canar. t. 161 = Linkii.
byzantina, Walp. Ann. iii. 251 = Calamintha Nepeta.
Calverti, Boiss. Fl. Orient. iv. 571 = elliptica.
canescens, *Benth. Lab. Gen. et Sp.* 376.—Sicil.
canescens, Walp. Ann. iii. 251 = Calamintha Nepeta.
capitellata, *Benth. in DC. Prod.* xii. 218.—Ind. or.
congesta, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 575.—Cataonia.
cordata, *Moris, Diagn. Stirp. Sard. Nov.* 2; *ex ej. Fl. Sard.* iii. 285.—Sardin.
cosentina, Terrac. in Nuov. Giorn. Bot. v. (1873) 227 = graeca.
cremnophila, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 570.—Graecia.
cristata, *Griseb. Spicil. Fl. Rumel.* ii. 122.—Graecia; As. Min.; Syria.
croatica, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 93.—Croatia.
Cunninghamii, Benth. Lab. Gen. et Sp. 730 = Mentha Cunninghamii.
cymuligera, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 569.—Cataonia.
cypria, Kotschy, in Unger, Cypern, 270 = graeca.
dalmatica, *Benth. in DC. Prod.* xii. 225.—Thracia.
dalmatica, Fenzl, Cat. Sem. Hort. Vindob. (1851); in Mem. Ist. Venet. vi. (1856) 247 = Mentha Pulegium.
Darwinii, *Benth. in DC. Prod.* xii. 222.—Patagon.
debilis, *Pomel, Nouv. Mat. Fl. Atl.* 122.—Alger.
densiflora, *Benth. Lab. Gen. et Sp.* 375.—Ins. Canar.
Douglasii, *Benth. l. c.* 372 (Douglassii).—Am. bor. occ.
elliptica, *C. Koch, in Linnaea*, xxi. (1848) 669.—Armenia.
eugenioides, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 413.—Reg. Argent.
Fenzlii, Regel, Ind. Sem. Hort. Petrop. (1866) 93 = Mentha Pulegium.
filicaulis, *Schott & Kotschy, in Tchihat. Asie Min. Bot.* ii. 131, nomen.—As. Min.
filiformis, *Benth. Lab. Gen. et Sp.* 378.—Reg. Mediterr.
flagellaris, *Baker, in Journ. Linn. Soc.* xx. (1883) 232.—Madag.
Fontanesii, *Pomel, Nouv. Mat. Fl. Atl.* 123.—Alger.
Forbesii, *Benth. Lab. Gen. et Sp.* 376.—Ins. Cap. Virid.

MICROMERIA :—
Forskåhlei, *Benth. Lab. Gen. et Sp.* 379.—Arab.
Gilliesii, *Benth. l. c.* 381.—Chili.
glabella, Benth. l. c. 371 = Calamintha glabella.
gracilis, Benth. l. c. 380 = Mentha gracilis.
graeca, *Benth. l. c.* 373.—Graecia; As. Min.
helianthemifolia, *Webb & Berth. Phyt. Canar.* iii. 79.—Ins. Canar.
herpyllomorpha, *Webb & Berth. l. c.* 72. t. 155.—Ins. Canar.
hirsuta, Nym. Consp. 590 = Juliana.
hirsuta, Mazziar. ex Nym. l. c. = nervosa.
hispida, *Boiss. & Heldr. Diagn.* Ser. I. xii. 48.—Ins. Creta.
Hydaspidis, *Falc. ex Benth. in DC. Prod.* xii. 224.—Reg. Himal.
hyssopifolia, Webb & Berth. Phyt. Cauar. iii. 72 = varia.
inodora, *Benth. Lab. Gen. et Sp.* 375.—Afr. bor.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 106.—Japon.
Juliana, *Benth. Lab. Gen. et Sp.* 373.—Reg. Mediterr. or.
julianoides, *Webb & Berth. Phyt. Canar.* iii. 78. t. 160.—Ins. Canar.
lachnophylla, *Webb & Berth. l. c.* 73. t. 156.—Ins. Canar.
lanata, *Benth. Lab. Gen. et Sp.* 731.—Ins. Canar.
lanata, Benth. l. c. 374 = Benthami.
lasiophylla, Webb & Berth. Phyt. Canar. iii. 74 = varia.
lepida, *Webb & Berth. l. c.* 74. t. 157.—Ins. Canar.
libanotica, *Boiss. Diagn.* Ser. I. xii. 50.—Syria.
Linkii, *Webb & Berth. Phyt. Canar.* iii. 79.—Ins. Canar.
longiflora, Tod. ex Nym. Consp. 590 = graeca.
longifolia, *Scheele, in Linnaea*, xxii. (1849) 593.—Dalmatia.
lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 94.—Lycia.
macrosiphon, *Coss. in Bull. Soc. Bot. Fr.* xxvii. (1880) 72.—Marocc.
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 244.—Madag.
Malcolmiana, Benth. ex Hook. f. Fl. Brit. Ind. iv. 650 = capitellata.
marifolia, *Benth. Lab. Gen. et Sp.* 382.—Europ. austr.
marifolia, Benth. l. c., partim = serpyllifolia.
microphylla, *Benth. l. c.* 377.—Reg. Mediterr.
mollis, *Benth. in DC. Prod.* xii. 225.—Mesopot.
montana, Reichb. Fl. Germ. Excurs. 311 = Satureia montana.
myrtifolia, Boiss. & Hohen. Diagn. Ser. I. v. 19 = M. Juliana.
nervosa, *Benth. Lab. Gen. et Sp.* 376.—Reg. Mediterr.
nubigena, *Benth. l. c.* 381.—Peruv.
nummularifolia, *Boiss. Diagn.* Ser. I. xii. 49.—Syria.
Nuttallii, Torr. & Gray, ex Benth. in DC. Prod. xii. 230 = Calamintha Nuttallii.
obovata, *Benth. Lab. Gen. et Sp.* 381.—Ind. occ.
odora, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 413.—Reg. Argent.
origanifolia, Benth. in DC. Prod. xii. 214 = Calamintha origanifolia.
origanifolia, Boiss. Fl. Orient. iv. 575 = dalmatica.
ovata, Beck. ex Hook. f. Fl. Brit. Ind. iv. 650 = biflora.
ovata, *Benth. Lab. Gen. et Sp.* 377.—Abyssin.
parviflora, *Reichb. Fl. Germ. Excurs.* 859.—Dalmatia.
Perezii, *C. Bolle, in Bonplandia*, viii. (1860) 282.—Ins. Canar.
perforata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 106.—Japon.
persica, *Boiss. Diagn.* Ser. I. vii. 48.—Persia.
Piperella, *Benth. Lab. Gen. et Sp.* 379.—Europ. austr.
Piperella, Panč. ex Nym. Consp. 591 = croatica?
plumosa, Buek, Cand. Ind. iii. 297; Heldr. ex Nym. Consp. 590 = nervosa.
polioides, *Webb & Berth. Phyt. Canar.* iii. 76. t. 158.—Ins. Canar.
Preauxii, Webb & Berth. l. c. 75 = varia.
pubescens, Boiss. & Kotschy, ex Boiss. Fl. Orient. iv. 571 = elliptica.
pulchella, *Wedd. Chlor. And.* ii. 151.—Chili.
Pulegium, *Benth. Lab. Gen. et Sp.* 382.—Banat.
punctata, *Benth. l. c.* 378.—Abyssin.
purpurea, *A. Gray, in Bot. Calif.* i. 595.—Calif.
pygmaea, Reichb. Fl. Germ. Excurs. 311 = Satureia montana.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 190.—Abyssin.

MICROMERIA:—
Reinholdi, Heldr. ex Boiss. Fl. Orient. iv. 570 = M. Juliana.
repens, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 274 = Mentha serpyllifolia.
Rodriguezii, Freyn & Janka, in Oestr. Bot. Zeitschr. xxiv. (1874) 16 = nervosa.
rupestris, *Benth. in DC. Prod.* xii. 225.—Europ. austr.
Rutenbergiana, *Vatke, in Bremen, Abh.* ix. (1885) 135.—Madag.
satureioides, Benth. Lab. Gen. et Sp. 380 = Mentha satureioides.
Schimperi, *Vatke, in Linnaea*, xxxvii. (1871–73) 326.—Abyssin.
serpyllifolia, *Boiss. Diagn.* Ser. II. iv. 13.—As. Min.; Syria.
serpyllifolia, *Schlecht. in Linnaea*, xxii. (1849) 593.—Dalmatia; Creta.
serpyllimorpha, Benth. in DC. Prod. xii. 217 = herpyllomorpha.
sessilis, Benth. Fl. Austral. v. 84, sphalm. = Mentha serpyllifolia.
sicula, Nym. Consp. 590 = graeca.
sinaica, *Benth. Lab. Gen. et Sp.* 380.—Arab.
sphaciotica, Boiss. & Heldr. ex Benth. in DC. Prod. xii. 220 = microphylla.
sphaerophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 232.—Madag.
spicata, Reichb. Fl. Germ. Excurs. 311 = Satureia cuneifolia.
spicigera, C. Koch, in Linnaea, xvii. (1843) 295 = Satureia alternipilosa.
spicigera, Walp. Ann. iii. 251 = serpyllifolia.
staminea, Boiss. & Hohen. Diagn. Ser. I. v. 19 = Calamintha staminea.
stolonifera, *Benth. Lab. Gen. et Sp.* 371.—Mexic.
subcordata, *Vis. ex Benth. l. c.* 379.—Dalmatia.
Teneriffae, *Benth. Lab. Gen. et Sp.* 378.—Ins. Teneriffa.
tenuifolia, Benth. in DC. Prod. xii. 215; Terrac. in Nuov. Giorn. Bot. v. (1873) 226 = graeca.
tenuifolia, Reichb. Fl. Germ. Excurs. 311 = parviflora.
tenuis, Benth. Lab. Gen. et Sp. 731 = Benthami.
tenuis, Benth. l. c. 374 = lanata.
tenuis, *Link, in Webb & Berth. Phyt. Canar.* iii. 75.—Ins. Canar.
terebinthinacea, Webb & Berth. l. c. 80 = Teneriffae.
teydensis, *C. Bolle, in Bonplandia*, viii. (1860) 282.—Ins. Canar.
Thymbra, Kostel. Allg. Med.-Pharm. Flora, iii. 763 = Satureia Thymbra.
thymoides, *De Not. Prosp. Fl. Ligust.* 54.—Italia.
thymoides, Webb & Berth. Phyt. Canar. iii. 71 = varia.
Tragothymus, Webb & Berth. l. c. iii. 73 = varia.
unguentaria, *Schweinf. Beitr. Fl. Aethiop.* 124.—Abyssin.
varia, *Benth. Lab. Gen. et Sp.* 374.—Ins. Canar.
variegata, Reichb. Fl. Germ. Excurs. 859 = Satureia montana.
xalapensis, Benth. Lab. Gen. et Sp. 372 = Brownei.
xylorrhiza, Boiss. & Heldr. Diagn. Ser. I. xii. 49 = cristata.

MICROMYRTUS, Benth. in Benth. & Hook. f. Gen. i. 700 (1865). *MYRTACEAE*, Benth. & Hook. f. l. c.
Drummondii, *Benth. Fl. Austral.* iii. 64.—Austral.
elobata, *Benth. l. c.*—Austral.
imbricata, *Benth. l. c.*—Austral.
leptocalyx, *Benth. l. c.* 65.—Austral.
microphylla, *Benth. l. c.*—Austral.
minutiflora, *Benth. l. c.*—Austral.
racemosa, *Benth. l. c.* 64.—Austral.

MICRONEMA, Schott, in Oestr. Bot. Wochenbl. (1857) 95 = **Micromeria**, Benth. (Labiat.).
Piperella, Schott, l. c. = M. Piperella.

MICRONOMA, H. Wendl. ex Benth. & Hook. f. Gen. iii. 882 (1883). *PALMARUM*, genus dubium.

MICRONYCHIA, Oliver, in Hook. Ic. Pl. xiv. 27. t. 1337 (1881). *ANACARDIACEAE.*
madagascariensis, *Oliver, l. c.*—Madag.

MICROPEPLIS, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 474 (Rel. Lehm. 295) = **Halogeton**, C. A. Mey. (Chenop.).
arachnoidea, Bunge, l. c. = H. arachnoideus.

MICROPERA, Dalz. in Hook. Kew Journ. iii. (1851) 282 = **Saccolabium**, Blume (Orchid.).
maculata, Dalz. l. c. (= *Sarcochilus maculatus*).—Ind. or.
viridiflora, Dalz. l. c. = S. viridiflorum.

MICROPERA, Lindl. Bot. Reg. sub t. 1522 (1832) = **Sarcochilus**, R. Br. (Orchid.).
Banksii, Heynh. Nom. i. 523, nomen.—N. Zel.
media, Heynh. Nom. i. 523, nomen.—Austral.
pallida, Lindl. in Bot. Reg. sub t. 1522.—Ind. or.
pallida, Lindl. in Wall. Cat. n. 7321, partim = Sarcanthus secundus.
parviflora, Heynh. Nom. i. 523, nomen = Sarcochilus parviflorus.

MICROPETALON, Pers. Syn. i. 509 (1805) = **Stellaria**, Linn. (Caryoph.).
gramineum, Pers. l. c. = S. longifolia.
lanceolatum, Pers. l. c. = S. borealis.
lanuginosum, Pers. l. c. = Arenaria lanuginosa.

MICROPEUCE, Gord. Pinet. Suppl. 13 (1862) = **Tsuga**, Carr. (Conif.).
Brunoniana, Carr. Conif. ed. II. 247 = T. Brunoniana.
Sieboldii, Gord. Pinet. Suppl. 13 = T. Sieboldii.

MICROPHOENIX, Naud. ex Carr. in Rev. Hortic. (1885) 513. *PALMAE* (hybrida).
decipiens ×, Naud. ex Carr. l. c.
Sahuti ×, Carr. l. c.

MICROPHYES, Phil. Fl. Atac. 20. t. 1 F (1868). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 154.
lanuginosa, *Phil. in Linnaea*, xxx. (1859–60) 209.—Chili.
litoralis, *Phil. Fl. Atac.* 20.—Chili.

MICROPHYSA, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 99. *MELASTOMACEAE*, Benth. & Hook. f. i. 767.
quadrialata, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 99.—Peruv.
rotundifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 141.—Bras.

MICROPHYSA, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 115 = **Galium**, Linn. (Rubiac.).
galioides, Schrenk, l. c. 116.—Soongaria.

MICROPHYSCA, Benth. & Hook. f. Gen. i. 767 (1867) (sphalm.) = praec.

MICROPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = **Trifolium**, Linn. (Legumin.).
glomeratum, Fourr. l. c. = T. glomeratum.
suffocatum, Fourr. l. c. = T. suffocatum.

MICROPIPER, Miq. in Bull. Sc. Phys. Nat. Néerl. (1839) 447 = **Peperomia**, Ruiz & Pav. (Piperac.).
alatum, Miq. Comm. Phyt. 52, in obs. = P. alata.
bilineatum, Miq. l. c. 59. t. 8 = P. bilineata.
blandum, Miq. l. c. 54 = P. blanda?
Blattarum, Miq. l. c. 52, in obs. = Piper Blattarum.
Blumei, Miq. l. c. 53. t. 8. f. G = P. polystachya.
candidum, Miq. l. c. 58. t. 8. = P. candida.
convexum, Miq. l. c. 60. t. 8 = P. convexa.
exiguum, Miq. l. c. 55. t. 9 = P. exigua.
laevifolium, Miq. l. c. 56. t. 9 = P. laevifolia.
Langsdorffii, Miq. l. c. 52. t. 8. f. F = P. Langsdorffii.
macrostachyum, Miq. l. c., in obs. = P. macrostachya.
melanostigma, Miq. l. c. 51 = P. melanostigma.
nervoso-venosum, Miq. l. c. 57. t. 8 = P. nervoso-venosa.
pellucidum, Miq. l. c. 54 = P. pellucida.
pusillum, Miq. l. c. 62. t. 5 = P. reflexa.
recurvatum, Miq. l. c. 61. t. 9 = P. recurvata.
subrotundum, Miq. l. c. 59. t. 8 = P. Reinwardtiana.
tenellum, Klotzsch, ex Miq. Fl. Ind. Bat. i. II. 432 = P. exigua.
tomentosum, Miq. Comm. Phyt. 63. t. 9 = P. tomentosa.
trinerve, Miq. l. c. 52, in obs. = P. trinervis.

MICROPLEURA, Lag. in Ocios Esp. Emigr. iv. (1825) 347, adnot. (Obs. Aparasol. 15). *UMBELLIFERAE*, Benth. & Hook. f. i. 873.
renifolia, *Lag. in Ocios Esp. Emigr.* iv. (1825) 347, adnot.—Ins. Chiloe.

MICROPODIUM, Reichb. Consp. 185 (1828) = **Brassica**, Tourn. (Crucif.).

MICROPOGON, 'Spreng.' ex Pfeiff. Nom. ii. 310 (1874), cum cit. falsa; an **Microchloa**, R. Br. (Gramin.)?

MICROPSIS, DC. Prod. v. 459 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 298.
LASIOPHYTON, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 44.
nana, *DC. Prod.* v. 460.—Chili.

MICROPTELEA, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 358 = **Ulmus**, Linn. (Urtic.).
parvifolia, Spach, l. c. 359 = U. parvifolia.

MICROPTERYX, Walp. in Linnaea, xxiii. (1850) 739 = **Erythrina**, Linn. (Legumin.).
Crista-galli, Walp. l. c. 740 = E. Crista-galli.
fasciculata, Walp. Ann. ii. 425 = E. Crista-galli.
laurifolia, Walp. l. c. 425; et in Otto & Dietr. Allg. Gartenz. xix. (1851) 50 = E. Crista-galli.
Poeppigiana, Walp. in Linnaea, xx. iii. (1850) 740 = E. Micropteryx.
reticulata, Walp. l. c. 741 = E. reticulata.
stricta, Walp. l. c. 740 = E. stricta.
suberosa, Walp. l. c. = E. suberosa.
sublobata, Walp. l. c. = E. suberosa.

MICROPUS, Linn. Gen. ed. I. 359 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 297.
ANCISTROCARPHUS, A. Gray, in Proc. Am. Acad. vii. (1868) 355.
GNAPHALODES, Adans. Fam. ii. 118 (1763).
amphibolus, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 214.—Calif.
angustifolius, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 339 = californicus.
bombycinus, *Lag. Gen. et Sp. Nov.* 32.—Reg. Mediterr.
brevissimus, Walp. Rep. ii. 600 = Psilocarphus brevissimus.
californicus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 42.—Am. bor. occ.
caulescens, Walp. Ann. ii. 842 = Psilocarphus caulescens.
conyzaeus, Dubois, Méth. ed. II. 386 = erectus.
discolor, Pers. Syn. ii. 423 = bombycinus.
erectus, *Linn. Sp. Pl.* ed. I. *Add. post Ind.*—Europ.
erectus, Tenore, ex Nym. Consp. 395 = bombycinus.
exiguus, Urv. in Mém. Soc. Linn. Par. i. (1822) 365 = Evax exigua.
globiferus, Bert. in Merc. Chil. 1829; ex Am. Journ. Sc. xxiii. (1833) 254 (= Psilocarphi sp.).—Chili.
Grayanus, Hemsl. Biol. Centr. Am. Bot. ii. 135 = Stylocline micropoides.
heterophyllus, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 178.—Calif.
longifolius, *Boiss. & Reut. in Boiss. Fl. Orient.* iii. 241.—Oriens.
mareoticus, Spreng. Syst. iii. 499 = Filago mareotica.
minimus, DC. Prod. v. 461 = Evax multicaulis.
multicaulis, *Dubois, Méth.* ed. II. 386.—Europ.
oregonus, Walp. Rep. ii. 600 = Psilocarphus oregonus.
pygmaeus, Desf. Fl. Atlant. ii. 307 = Evax umbellata.
supinus, *Linn. Sp. Pl.* 927.—Reg. Mediterr.
tenellus, Walp. Rep. ii. 600 = Psilocarphus tenellus.

MICROPYRUM, Link, in Linnaea, xvii. (1843) 397 = **Festuca**, Linn. (Gramin.).
tenellum, Link, l. c. 398 = F. Poa.

MICROPYXIS, Duby, in DC. Prod. viii. 71 (1844) = **Centunculus**, Dill. (Primul.).
elongata, Duby, l. c.—Bolivia.
ovalis, Phil. in Anal. Univ. Chil. (1856) 161 = C. pentandrus.
pumila, Duby, in DC. Prod. viii. 72 = C. pentandrus.
rubricaulis, Duby, l. c.—Madag.
tenella, Wight, Ic. t. 1585 = C. pentandrus.

MICRORHAMNUS, A. Gray, Pl. Wright. i. 33 (1852). *RHAMNEAE*, Benth. & Hook. f. i. 376.
RHAMNELLA, Miq. Ann. Mus. Bot. Lugd. Bat. iii 30 (1867).
ericoides, *A. Gray, Pl. Wright.* i. 34.—Mexic.
franguloides, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. n. XI. (1866) 4.—Japon.

MICRORHINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 127 = **Linaria**, Tourn. (Scrophul.).
minus, Fourr. l. c. = L. viscida.

MICRORHYNCHUS, Less. Syn. Comp. 139 (1832) = **Launaea**, Cass. (Compos.).
acaulis, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 208 = Crepis acaulis.
altissimus, Steud. Nom. ed. II. ii. 143, nomen.—Arab.
arabicus, Jaub. & Spach, Illustr. iii. 107. t. 277 = L. nudicaulis.
asplenifolius, DC. Prod. vii. 181 = L. aspleniifolia.
bellidifolius, DC. l. c. = L. bellidifolia.
Brunneri, Walp. Ann. ii. 976 = Sonchus Brunneri.
chondrilloides, C. B. Clarke, Comp. Ind. 277 = L. chondrilloides.
chondrilloides, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 485, nomen = Prenanthes chondrilloides.
Dregeanus, DC. Prod. vii. 181 = L. bellidifolia.
fallax, Jaub. & Spach, Illustr. iii. 106. t. 276 = L. nudicaulis.
glaber, Wight, Ic. t. 1145 = Crepis acaulis.
glomeratus, Jaub. & Spach, Illustr. iii. 105 = L. glomerata.
nudicaulis, Less. Syn. Comp. 139 = L. nudicaulis.
octophyllus, Hochst. ex Oliver, Fl. Trop. Afr. iii. 452 = Lactuca goraeënsis.
patens, DC. Prod. vii. 181 = Lactuca Heyneana.
pentaphyllus, Hochst. ex Oliver, Fl. Trop. Afr. iii. 451 = Lactuca taraxacifolia.
picridioides, Walp. Ann. i. 463 (= *Rhabdotheca picridioides*).—Ins. Cap. Virid.
sarmentosus, DC. Prod. vii. 181 = L. pinnatifida.
secundus, C. B. Clarke, Comp. Ind. 276 = L. secunda.
spinosus, Benth. & Hook. f. Gen. ii. 529 = L. spinosa.
surinamensis, Miq. in Versl. Medel. Kon. Nederl. Inst. 1842 (1843) 204.—Guian.

MICRORYNCHUS, Sch. Bip. in Linnaea, xv. (1841) 725 = praec.

MICROSACCUS, Blume, Bijdr. 367 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 582.
javensis, *Blume, Bijdr.* 367.—Java.

MICROSCHWENKIA, Benth. ex Hemsl. Biol. Centr. Am. Bot. ii. 438 (1882) = **Melananthus**, Walp. (Solanac.).
guatemalensis, Benth. ex Hemsl. l. c. = Mel. dipyrenoides.

MICROSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 141. *UMBELLIFERAE*, Benth. & Hook. f. i. 892.
CYMINUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 142, in syn.
tenuifolium, *Boiss. l. c.* 142.—As. Min.

MICROSCIADIUM, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 468 [bis] = **Azorella**, Lam. (Umbell.).
cuneifolium, F. Muell. in Hook. Kew Journ. vii. (1855) 378 = A. cuneifolia.
Saxifraga, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 468 [bis] = A. Saxifraga.

MICROSECHIUM, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 25. *CUCURBITACEAE*, Benth. & Hook. f. i. 838.
guatemalense, Hemsl. Diagn. Pl. Nov. 16 = ruderale.
Helleri, Cogn. in DC. Monog. Phan. iii. 910 = ruderale.
palmatum, *Cogn. l. c.* 911.—Mexic.
ruderale, *Naud. in Ann. Sc. Nat.* Sér. V. vi. (1866) 25.—Mexic.

MICROSELINUM, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 407, nomen. *UMBELLIFERAE*.
Besseri, Andrz. l. c. (Quid ?).—Rossia.

MICROSEMMA, Labill. Sert. Austro-Caled. 58. t. 57 (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 187.
MICROSTEMMA, Reichb. Consp. 208 (1828).
Balansae, *Baill. Adansonia*, xi. (1876), 367.—N. Caled.
cernua, *Baill. l. c.*—N. Caled.
salicifolia, *Labill. Sert. Austro-Caled.* 58. t. 57.—N. Caled.

MICROSEPALA, Miq. Fl. Ind. Bat. Suppl. 444 (1860) **Baccaurea**, Lour. (Euphorb.).
acuminata, Miq. l. c. = B. acuminata.

MICROSERIS, D. Don, in Phil. Mag. xi. (Apr. 1832) 388. *COMPOSITAE*, Benth. & Hook. f. ii. 506.
BELLARDIA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 40. t. 34.
CALAIS, DC. Prod. vii. 85 (1838).
FICHTEA, Sch. Bip. in Linnaea, x. (1836) 255.
GALASIA, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 310.
LEPIDONEMA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 31 (1835).
MYCROSERIS, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 250.
PHYLLOPAPPUS, Walp. in Linnaea, xiv. (1840) 507.
PTILOPHORA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 112.
SCORZONELLA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 426.
UROPAPPUS, Nutt. l. c. (1841) 424.
acuminata, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 88.—Calif.
aphantocarpha, *Sch. Bip. in Pollichia*, xxii-xxiv. (1866) 308.—Calif.
attenuata, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 111.—Calif.
Bigelowii, *Sch. Bip. in Pollichia*, xxii–xxiv. (1866) 308.—Calif.
Bolanderi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 64.—Am. bor. occ.
borealis, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 310 = Leontodon borealis.
cuspidata, Sch. Bip. l. c. 309 = Troximon cuspidatum.
cyclocarpha, Sch. Bip. l. c. 308 = Douglasii.
Douglasii, *Sch. Bip. l. c.*—Calif.
elegans, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. II. 419.—Calif.
eriocarpha, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 308 = Douglasii.
Forsteri, *Hook. f. Fl. N. Zel.* 151.—Austral. ; N. Zel.
Howellii, *A. Gray, in Proc. Am. Acad.* xx. (1855) 300.—Am. bor. occ.
laciniata, *Sch. Bip. in Pollichia*, xxii–xxiv. (1866) 309.—Am. bor. occ.
lanceolata, Sch. Bip. l. c. 310 = Forsteri.
leptosepala, *A. Gray, in Proc. Am. Acad.* ix. (1874) 209 ; xix. (1883) 64.—Am. bor. occ.
leptosepala, A. Gray, l. c. 209, partim = Bolanderi.
Lindleyi, *A. Gray, l. c.* 210.—Calif.
linearifolia, *Sch. Bip. in Pollichia*, xxii–xxiv. (1866) 308.—Am. bor. occ.
macrochaeta, *Sch. Bip. l. c.* 309.—Am. bor. occ.
major, *Sch. Bip. l. c.*—Am. bor. occ.
nutans, *Sch. Bip. l. c.*—Am. bor. occ.
Parryi, *Sch. Bip. l. c.*—Calif.
platycarpha, *Sch. Bip. l. c.* 308.—Calif.
procera, *A. Gray, in Proc. Am. Acad.* xix. (1883) 64.—Calif.
pygmaea, *D. Don, in Phil. Mag.* (Apr. 1832) 388.—Chili.
scapigera, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 310 = Forsteri.
sylvatica, *Sch. Bip. l. c.* 309.—Calif.
tenella, Sch. Bip. l. c. 308 = aphantocarpha.
troximoides, *A. Gray, in Proc. Am. Acad.* ix. (1874) 211.—Am. bor. occ.

MICROSPERMA, Hook. Ic. Pl. t. 234 (1839) = **Mentzelia**, Linn. (Loaseae).
bartonioides, Walp. Rep. v. 777 = Ment. gronoviaefolia.
grandiflora, Groenl. in Rev. Hortic. (1861) 349.—Mexic.
lobata, Hook. Ic. Pl. t. 234 = Ment. lobata.
rudis, S. Schau. in Linnaea, xx. (1847) 721 = Ment. lobata.

MICROSPERMUM, Lag. Gen. et Sp. Nov. 25 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 407.
MIRADORIA, Sch. Bip. ex Benth. & Hook. f. l. c. (1873).
debile, *Benth. Pl. Hartw.* 64.—Mexic.
nummulariaefolium, *Lag. Gen. et Sp. Nov.* 25.—Mexic.

MICROSPLENIUM, Hook. f. in Benth. & Hook. f. Gen. ii. 4 (1873). *CAPRIFOLIACEAE*, Benth. & Hook. f. l. c.
Coulteri, *Hook. f. l. c.*—Mexic.

MICROSTACHYS, A. Juss. Euph. Tent. 48. t. 15 (1824) = **Sebastiania**, Spreng. (Euphorb.).
bicornis, A. Juss. l. c. 49. t. 15 = S. corniculata.
Chamaelea, Juss. l. c. = S. Chamaelea.
corniculata, Juss. l. c. = S. corniculata.
crotonoides, Klotzsch, ex Benth. in Hook. Kew Journ. vi. (1854) 325 = S. corniculata.
daphnoides, Muell. Arg. in Linnaea, xxxii. (1863) 91 = S. daphnoides.
fruticulosa, Karst. Fl. Columb. ii. 33 = S. corniculata.
guianensis, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 45 = S. corniculata.
hastata, Klotzsch, ex Baill. Adansonia, v. (1864–65) 324, nomen = S. ditassoides.
marginata, Klotzsch, ex Muell. Arg. in Linnaea, xxxii. (1863) 90 = S. marginata.
Mercurialis, Dalz. & Gibs. Bomb. Fl. 227 = Claoxylon Mercurialis.
micrantha, Benth. in Hook. Kew Journ. vi. (1854) 326 = S. corniculata.
polymorpha, Muell. Arg. in Linnaea, xxxii. (1863) 91 = S. corniculata.
ramosissima, A. St. Hil. Pl. Rém. Brés. 242 = S. brasiliensis.
serrulata, Muell. Arg. in Linnaea, xxxii. (1863) 90 = S. serrulata.
sessilifolia, Muell. Arg. l. c. 95 = S. ditassoides.
stipulacea, Klotzsch, ex Baill. Adansonia, v. (1864–65) 325 = S. stipulacea.
Vahlii, A. Rich. Fl. Cub. Fanerog. ii. 202 = S. corniculata.
virgata, Muell. Arg. in Linnaea, xxxii. (1863) 94 = S. virgata.

MICROSTEGIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836) = **Pollinia**, Trin. (Gramin.).
Willdenovianum, Nees, l. c. = P. imberbis.

MICROSTEIRA, Baker, in Journ. Linn. Soc. xx. (1883) 111. *MALPIGHIACEAE*.
Curtisii, *Baker, l. c.*—Madag.

MICROSTEMMA, R. Br. in Mem. Wern. Soc. i. (1809) 25. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 778.
glabriflorum, *F. Muell. Fragm.* i. 58.—Austral.
tuberosum, *R. Br. Prod.* 459.—Austral.

MICROSTEMMA, Reichb. Consp. 208 (1828) = **Microsemma**, Labill. (Ternstroem.).

MICROSTEMON, Engl. Bot. Jahrb. i. (1881) 376. *ANACARDIACEAE*.
velutina, *Engl. l. c.*; *et in DC. Monog. Phan.* iv. 294.—Malacca.

MICROSTEPHIUM, Less. in Linnaea, vi. (1831) 92 = **Cryptostemma**, R. Br. (Compos.).
niveum, Less. l. c. = C. niveum.

MICROSTIGMA, Trautv. Pl. Imag. 36. t. 25 (1844) = **Matthiola**, R. Br. (Crucif.).
Bungei, Trautv. l. c. = M. deflexa.

MICROSTYLIS, Nutt. Gen. Am. ii. 196 (1818). *ORCHIDEAE*, Benth. & Hook. f. iii. 494.
ACHROANTHES, Rafin. in Desv. Journ. Bot. ii. (1809) 17.
CREPIDIUM, Blume, Bijdr. 387 (1825).
DIENIA, Lindl. in Bot. Reg. sub t. 825 (1824).
PEDILEA, Lindl. Orch. Scel. 27 (1826).
PTEROCHILUS, Hook. & Arn. Bot. Beech. Voy. 71. t. 17 (1832).
atropurpurea, Miq. Fl. Ind. Bat. iii. 625 (Liparidis sp.).—Java.
bella, Reichb. f. in Illustr. Hortic. xxxii. (1885) t. 581 = Rheedii, *Lindl.*
Bernaysii, F. Muell. Fragm. xi. 21 = congesta.
biaurita, *Lindl. in Wall. Cat.* n. 1941.—Reg. Himal.
biloba, Lindl. l. c. n. 1940 = Wallichii.
brachypoda, A. Gray, in Ann. Lyc. N. York, iii. (1828–36) 228 = monophylla.
brachystachys, *Reichb. f. in Linnaea*, xxii. (1849) 834.—Mexic.
calophylla, *Reichb. f. in Gard. Chron.* (1879) II. 718.—Malaya.
cardiophylla, *Reichb. f. in Flora*, lxviii. (1885) 543.—Ins. Comoro.

MICROSTYLIS:—
carinata, *Reichb. f. in Walp. Ann.* vi. 207.—Ins. Philipp.
caulescens, *Lindl. Bot. Reg.* (1841) sub t. 1.—Peruv.
chlorophrys, *Reichb. f. in Gard. Chron.* (1881) I. 266.—Borneo.
cochleariaefolia, Reichb. f. in Linnaea, xxii. (1849) 834 = ichthyorrhyncha.
commelynifolia, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, i. (1844) 402.—Java.
congesta, *Reichb. f. in Walp. Ann.* vi. 206.—As. et Austral. trop.
cordata, *Reichb. f. l. c.* 207.—Mexic.
corymbosa, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 195.—Arizona.
crispata, *Reichb. f. in Walp. Ann.* vi. 207.—Mexic.
crispifolia, *Reichb. f. Beitr. Orch. Centr. Am.* 100.—Costa Rica.
cylindrostachya, *Reichb. f. in Walp. Ann.* vi. 207.—Reg. Himal.
decurrens, Miq. Fl. Ind. Bat. iii. 625 (= *Platystylis decurrens*).—Java.
diphyllos, Lindl. Gen. et Sp. Orch. 19 = monophyllos.
discolor, *Lindl. l. c.* 20.—Zeylan.
disepala, *Reichb. f. in Linnaea*, xxvi. (1853–55) 142.—Venezuela.
Ehrenbergii, *Reichb. f. l. c.* xxii. (1849) 835.—Mexic.
excavata, *Lindl. Bot. Reg.* (1838), *Misc.* 51.—Mexic.
fastigiata, *Reichb. f. in Linnaea*, xxii. (1849) 834.—Mexic.
flavescens, Lindl. Gen. et Sp. Orch. 20 = congesta.
floridana, *Chapman, Fl. S. U. St.* 454.—Florida.
fusca, Reichb. f. in Walp. Ann. vi. 207 = congesta.
Godefroyi, *Reichb. f. Otia Bot. Hamb.* 37.—Cambodia.
hastilabia, *Reichb. f. Beitr. Orch. Centr. Am.* 101.—Am. centr.; Bras.
histionantha, *Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol.* 11. t. 5.—Nicaragua.
ichthyorrhyncha, *Reichb. f. Beitr. Orch. Centr. Am.* 99.—Mexic.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 203.—Japon.
Josephiana, *Reichb. f. in Hook. Bot. Mag.* t. 6325.—Reg. Himal.
lagotis, *Reichb. f. Beitr. Orch. Centr. Am.* 101.—Costa Rica.
lancifolia, *Thw. Enum. Pl. Zeyl.* 297.—Zeylan.
Lowi, *E. Morr. in Belg. Hortic.* xxxiv. (1884) 281. t. 14. f. 3.—Borneo.
luteola, Wight, Ic. t. 1632 = versicolor.
macrostachya, *Lindl. Gen. et Orch.* 21.—Mexic.
maianthemifolia, *Reichb. f. in Linnaea*, xxii. (1849) 834.—Mexic.
metallica, *Reichb. f. in Gard. Chron.* (1879) II. 750.—Borneo.
monophyllos, *Lindl. Gen. et Sp. Orch.* 19.—Reg. bor. temp.
montana, *Engelm. ex Rothr. Rep. Bot. Wheeler, Surv.* vi. 264.—Arizona.
myurus, *Reichb. f. in Walp. Ann.* vi. 207.—Mexic.
odorata, *Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 230.—Ins. Philipp.
ophioglossoides, Lindl. Bot. Reg. t. 1290 = fastigiata.
ophioglossoides, *Nutt. Gen. Am.* ii. 196.—Am. bor.
Parthonis, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol. 12, in textu = histionantha.
plantaginea, Steud. Nom. ed. II. ii. 144 = Rheedii, *Lindl.*
platychila, *Reichb. f. in Seem. Fl. Vit.* 302.—Ins. Fiji.
pubescens, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 662.—Bras.
purpurea, *Lindl. Gen. et Sp. Orch.* 20.—Zeylan.
purpurea, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 195.—Arizona.
Rheedii, *Lindl. Gen. et Sp. Orch.* 21.—Malaya; Ins. Pacif.
Rheedii, *Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 138 (non alior.).—Burma.
Rheedii, Wight, Ic. t. 902 = versicolor, *Lindl.*
rupestris, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 8.—Peruv.
simillima, *Reichb. f. Beitr. Orch. Centr. Am.* 101.—Costa Rica.
spicata, *Lindl. Gen. et Sp. Orch.* 19.—Jamaic.
stelidostachya, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. trop.
taurina, *Reichb. f. in Linnaea*, xli. (1877) 97.—N. Caled.

MICROSTYLIS:—
tipuloides, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 256.—Costa Rica.
trilobulata, Kurz, Rep. Veg. Andam. App. B. 19, nomen = congesta.
umbellata, Steud. Nom. ed. II. ii. 144 = seq.
umbellulata, *Lindl. Gen. et Sp. Orch.* 19.—Ind. occ.
Ventilabrum, *Reichb. f. in Gard. Chron.* (1881) II. 717.—Ins. Sunda.
ventricosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 9.—Peruv.
versicolor, *Lindl. Gen. et Sp. Orch.* 21.—Ind. or.
versicolor, *Wight, Ic.* t. 901.—Ind. or.
Wallichii, *Lindl. in Wall. Cat.* n. 1938.—Reg. Himal.
Warmingii, *Reichb. f. Otia Bot. Hamb.* 94.—Bras.

MICROSYPHUS, Presl, Bot. Bemerk. 91 (1844) = **Striga**, Lour. (Scrophul.).
parviflorus, Presl, l. c. = S. orobanchoides.

MICROTEA, Sw. Prod. Veg. Ind. Occ. 53. *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 82 (1880).
ANCISTROCARPUS, H. B. & K. Nov. Gen. et Sp. ii. 186. t. 122 (1817).
APHANANTHE, Link, Enum. Hort. Berol. i. 383 (1821).
CERATOCOCCA, Willd. ex Roem. & Schult. Syst. vi. p. lxx (1820).
POTAMOPHILA, Schrank, Pl. Rar. Hort. Monac. ii. t. 63 (1819).
SCHOLLERA, Rohr, in Skrivt. Naturhist. Selsk. Kjobenh. ii. (1792) 210.
celosioides, Moq. in DC. Prod. xiii. II. 18 = paniculata.
debilis, *Sw. Prod. Veg. Ind. Occ.* 53.—Ind. occ.
glochidiata, *Moq. in DC. Prod.* xiii. II. 18.—Bras.
lanceolata, Delile, Hort. Monsp. (1827); ex Moq. l. c. 17 = maypurensis.
maypurensis, *G. Don, in Loud. Hort. Brit.* 98.—Am. austr.
ovata, Delile, Hort. Monsp. (1827); ex Moq. in DC. Prod. xiii. II. 17 = debilis.
paniculata, *Moq. l. c.* 18.—Bras.
portoricensis, *Urban, in Ber. Deutsch. Bot. Ges.* iii. (1885) 324.—Ind. occ.
scabrida, *Urban, l. c.* 325.—Ind. occ.
Sprengelii, *Moq. in DC. Prod.* xiii. II. 19.—Bras.
tenuifolia, *Moq. l. c.* 18.—Bras.

MICROTHOUAREIA, Steud. Nom. ed. I. 531 (1821) = seq.

MICROTHUAREIA, Thou. Gen. Nov. Madag. 3 (1806) = **Thouarea**, Pers. (Gramin.).

MICROTINUS, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 293 = **Viburnum**, Linn. (Caprifol.).
odoratissimus, Oerst. l. c. 294 = V. odoratissimum.
Simonsii, Oerst. l. c. = V. Simonsii.

MICROTIS, R. Br. Prod. 320 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 609.
alba, *R. Br. Prod.* 321.—Austral.
arenaria, Lindl. Gen. et Sp. Orch. 396 = porrifolia.
atrata, *Lindl. Swan River App.* 54.—Austral.
Banksii, Hook. Bot. Mag. sub t. 3377 = porrifolia.
Benthamiana, Reichb. f. Beitr. Syst. Pfl. 24 = parviflora.
Brownii, Reichb. f. l. c. = media.
frutetorum, Schlecht. in Linnaea, xx. (1847) 568 = porrifolia.
javanica, *Reichb. f. in Bonplandia*, v. (1857) 36.—Java.
longifolia, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 247.—N. Zel.
media, *R. Br. Prod.* 321.—Austral.
minutiflora, F. Muell. Fragm. i. 90 = atrata.
pallida, *Heynh. Nom.* i. 523, nomen (Quid?).—Ind. or.
parviflora, *R. Br. Prod.* 321.—Austral.
porrifolia, *R. Br. l. c.* 320.—Austral.; N. Zel.
pulchella, *R. Br. l. c.* 321.—Austral.
pulchella, Lindl. Gen. et Sp. Orch. 395 = porrifolia.
rara, R. Br. Prod. 321 = porrifolia.
unifolia, Reichb. f. Beitr. Syst. Pfl. 62 = porrifolia.
viridis, F. Muell. ex Benth. Fl. Austral. vi. 347, in obs. = media, parviflora, porrifolia.

MICROTREMA, Klotzsch, in Linnaea, xii. (1838) 499 = **Erica**, Tourn.
opulentum, Klotzsch, l. c. = E. opulenta.

MICROTRICHIA, DC. Prod. v. 366 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 260.
Perrottetii, *DC. Prod.* v. 366.—Afr. trop.

MICROTROPIA, Reichb. Nom. 236 (1841) = seq.

MICROTROPIS, E. Mey. Comm. Pl. Afr. Austr. 65 (1835) = **Euchlora**, Eckl. & Zeyh. (Legumin.).
hirsuta, E. Mey. l. c. = E. serpens.

MICROTROPIS, Wall. Cat. pp. 152, 250 (1831); et ex Meissn. Gen. 68; Comm. 49 (1837). *CELASTRINEAE*, Benth. & Hook. f. i. 361.
PARACELASTRUS, Miq. Fl. Ind. Bat. i. II. 590 (1859).
bivalvis, *Wall. Cat.* n. 4340.—Malaya.
coriacea, Wall. l. c. 4338 = Gynotroches axillaris.
densiflora, *Wight, Ic.* t. 1052.—Ind. or.
discolor, *Wall. Cat.* n. 4337.—Reg. Himal.; Burma.
garcinifolia, Wall. ex Wight, Ic. t. 761 = discolor.
latifolia, *Gardn. ex Bedd. in Trans. Linn. Soc.* xxv. (1865) 213, nomen.—Ind. or.
latifolia, *Wight, ex M. Laws. in Hook. f. Fl. Brit. Ind.* i. 613.—Ind. or.
longifolia, Wall. Cat. n. 4339 = latifolia, Salacia flavescens.
microcarpa, *Wight, Ic.* t. 975.—Ind. or.
ovalifolia, *Wight, l. c.* t. 976.—Ind. or.
ramiflora, *Wight, l. c.* t. 977.—Ind. or.
Wallichiana, *Wight, ex Thw. Enum. Pl. Zeyl.* 71.—Zeylan.

MICROULA, Benth. in Benth. & Hook. f. Gen. ii. 853 (1876). *BORAGINEAE*, Benth. & Hook. f. l. c.
Benthami, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 167.—Reg. Himal.
tangutica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 500.—China.

MIDA, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 376 = **Fusanus**, R. Br. (Santal.).
eucalyptoides, A. Cunn. l. c. (= *Santalum Cunninghamii*).—N. Zel.
myrtifolia, A. Cunn. l. c. 377 (= *Santalum Cunninghamii*).—N. Zel.
salicifolia, A. Cunn. l. c. 376 (= *Santalum Cunninghamii*).—N. Zel.
undulata, Colenso, Excurs. N. Island, 84 (= *Santalum Cunninghamii*).—N. Zel.

MIDDENDORFIA, Trautv. in Mém. Sav. Etr. Pétersb. iv. (1842) 489 = **Peplis**, Linn. (Lythrar.).
borysthenica, Trautv. l. c. 490 = Lythrum nummulariifolium.
hamulosa, Trautv. l. c. 491 = P. erecta.

MIEGIA, Neck. Elem. i. 49 (1790) = **Hieracium**, Tourn. (Compos.).

MIEGIA, Pers. Syn. i. 101 (1805) = **Arundinaria**, Michx. (Gramin.).
arundinacea, Torr. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 15 = A. macrosperma.
gigantea, Nutt. Gen. Am. i. 39 = A. macrosperma.
macrosperma, Pers. Syn. i. 102 = A. macrosperma.
maritima, Willd. Sp. Pl. i. 311 = A. macrosperma.
pumila, Nutt. ex Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 112 = A. macrosperma.

MIEGIA, Schreb. Gen. 786 (1791) = **Remirea**, Aubl. (Gramin.). Cyperaceae

MIERIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 9 (1825) = **Schkuhria**, Roth (Compos.).
virgata, La Llave, l. c. = S. virgata.

MIERSIA, Lindl. in Miers, Trav. Chil. ii. 529 (1826), nomen; et in Bot. Reg. sub t. 992 (1826). *LILIACEAE*, Benth. & Hook. f. iii. 804.
chilensis, *Lindl. in Miers, Trav. Chil.* ii, 529; *et in Bot. Reg.* sub t. 992.—Chili.
cornuta, Phil. in Anal. Univ. Chil. (1873) 548 = chilensis.
major, Kunth, Enum. Pl. iv. 486 = chilensis.
minor, Kunth, l. c. 487 = myodes.
myodes, *Bert. in Merc. Chil.* 700.—*Cf Am. Journ. Sc.* xxiii. (1833) 254.—Chili.

MIKANIA, Neck. Elem. ii. 217 (1790) = **Perebea**, Aubl. (Urtic.).

MIKANIA, F. W. Schmidt, Samml. Phys. Aufs. i. 272 (1795) = **Lactuca**, Linn. (Compos.).
floridana, F. W. Schmidt, l. c. = L. cretica.

MIKANIA, Willd. Sp. Pl. iii. 1742 (1803). *COMPOSITAE*, Benth. & Hook. f. ii. 246.
CARELIA, Cav. in Anal. Cienc. Nat. vi. (1802) 317.
CORYNANTHELIUM, Kunze, in Linnaea, xx. (1847) 19.
MORONOA, Hort. ex Kunze, l. c. (1847).
MORRENIA, Hort. ex Kunze, l. c. (1847).
WILLUGBAEYA, Neck. Elem. i. 82 (1790).
acuminata, *DC. Prod.* vii. 270.—Bras.
affinis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 120 = lanuginosa.
alata, DC. Prod. v. 197 = scandens.
amara, Bert. ex DC. Prod. v. 199 = cordifolia.
amara, *Willd. Sp. Pl.* iii. 1744.—Am. austr.
amazonica, *Baker, in Mart. Fl. Bras.* vi. II. 266.—Bras.
ambigua, DC. Prod. v. 187 = penstemonoides.
Andrachne, DC. l. c. 192 = laevis.
angularis, *Humb. & Bonpl. Pl. Aequin.* ii. 87. t. 106.—N. Granat.
angulata, *La Llave, in Mosaico Mexicano*, ii. 299; *ex Hook. Kew Journ.* v. (1853) 78.—Mexic.
angulosa, Rafin. Fl. Ludov. 62 = scandens.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 138 = Eupatorium angustifolium.
antennaria, Sch. Bip. in Linnaea, xxx. (1859–60) 182 = microdonta.
apiifolia, *DC. Prod.* v. 202.—Bras.
arborea, H. B. & K. Nov. Gen. et Sp. iv. 139 = Eupatorium dendroides.
arborea, Roxb. in Beats. Tracts St. Helen. App. 313 = Senecio prenanthiflora.
Argyreiae, *DC. Prod.* v. 193.—Bras.
argyropappa, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 268.—Bras.
argyrostigma, Miq. in Linnaea, xvii. (1843) 69 = amara.
armigera, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 52. t. 260.—Peruv.
aromatica, *Oerst. in Overs. Dansk. Vidensk. Selsk. Forh.* (1863) 10.—Bras.
artemisioides, Cass. in Dict. Sc. Nat. lv. 130 = Eupatorium foeniculaceum.
asparagoides, Licht. ex Less. Syn. Comp. 394 = Euryops asparagoides.
aspera, *Miq. in Linnaea*, xvii. (1843) 68.—Guian.
asperifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 256 = phaeoclados.
atriplicifolia, Sch. Bip. ex Miq. Stirp. Surin. Sel. 189 = scandens.
attenuata, *DC. Prod.* v. 195.—Peruv.
auricularis, Griseb. in Goett. Abh. xix. (1874) 170 = scandens.
auriculata, Willd. Sp. Pl. iii. 1745 = Senecio deltoideus.
Badieri, *DC. Prod.* v. 194.—Ins. Guadalup.
Banisteriae, *DC. l. c.* 193.—Bras.
batataefolia, DC. l. c. 197 = scandens.
Berteriana, Spreng. Syst. iii. 423 = Eupatorium guadalupense.
biformis, *DC. Prod.* v. 202.—Bras.
biformis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 254 = cordifolia.
bogotensis, *Benth. Pl. Hartw.* 201.—N. Granat.
boliviensis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
brachiata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 53.—Peruv.
brachypoda, DC. Prod. v. 201 = officinalis.
bracteosa, *DC. l. c.* 194.—Bras.
brevipetiolata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 230 = myriocephala.
buddleiaefolia, *DC. Prod.* v. 192.—Bras.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. II. 232.—Bras.
callineura, *Sch. Bip. ex Baker, l. c.* 258.—Bras.
campanulata, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 489.—Bras.
Candolleana, *Gardn. l. c.* 484.—Bras.
capensis, *DC. Prod.* v. 198.—Afr. austr.
carnosula, DC. l. c. 197 = cordifolia.
caudata, *Benth. Pl. Hartw.* 201.—N. Granat.
Charua, *Griseb. in Goett. Abh.* xxiv. (1879) 174.—Reg. Argent.
chenopodifolia, Willd. Sp. Pl. iii. 1745 = scandens.
chenopodiodes, Wall. Cat. n. 3175 = Senecio chenopodifolius.
chlorolepis, *Baker, in Mart. Fl. Bras.* vi. II. 247.—Bras.
cinerea, Sch. Bip. ex Baker, l. c. 236 = sericea.

MIKANIA:—
cissampelina, *DC. Prod.* v. 195.—Ins. S. Doming.
Clausseni, Sch. Bip. in Linnaea, xxx. (1859–60) 182 = nummularia.
clavellata, DC. Prod. v. 192 = Ophryosporus Freyreissii.
clematidea, Wall. Cat. n. 3173 = Eupatorium Reevesii.
coarctata, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 488.—Bras.
cocaënsis, Gardn. l. c. 487 = cordifolia.
complanata, Gardn. l. c. 485 = obovata.
conferta, *Gardn. l. c.* iv. (1845) 119.—Bras.
confertissima, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 242.—Bras.
congesta, DC. Prod. v. 197 = scandens.
consanguinea, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 448 = psilostachya.
convolvulacea, DC. Prod. v. 199 = cordifolia.
cordifolia, *Willd. Sp. Pl.* iii. 1746.—Am. mer.
coriacea, *La Llave, in Mosaico Mexicano*, ii. 299; *ex Hook. Kew Journ.* v. (1853) 77.—Mexic.
cornifolia, G. Don, ex Baker, in Mart. Fl. Bras. vi. II. 237 = amara.
corydalifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 512.—Ind. occ.
corymbulosa, Benth. Pl. Hartw. 211 = Kanimia corymbulosa.
crispiflora, *Wright, in Sauv. Fl. Cub.* 78.—Cuba.
Cumingii, Sch. Bip. in Linnaea, xxxiv. (1865–66) 535, nomen = Ophryosporus Cumingii.
cuneata, Sch. Bip. in Bonplandia, iv. (1856) 54, nomen = amara.
cynanchifolia, Hook. & Arn. ex Baker, in Mart. Fl. Bras. vi. II. 249 = scandens.
decora, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 53.—Peruv.
deflexa, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 482 = Lundiana.
deltoides, *Poepp. ex Spreng. Syst.* iii. 423.—Cuba.
dentata, *Spreng. l. c.* 422.—Bras.
denticulata, Willd. Sp. Pl. iii. 1744 = scandens.
dioscoreaefolia, *DC. Prod.* v. 198.—Madag.
discolor, *Baker, in Mart. Fl. Bras.* vi. II. 234.—Bras.
divaricata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 488 = microcephala.
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 53.—Bras.
diversifolia, DC. Prod. v. 201 = biformis.
elliptica, *DC. l. c.* vii. 270.—Bras.
ericoides, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 372 = Brickellia pinifolia.
erioclada, *DC. Prod.* v. 192.—Bras.
eriophora, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 103.—Bras.
erithalina, DC. Prod. v. 193 = Kanimia erythralina.
estrellenis, *Baker, in Mart. Fl. Bras.* vi. II. 231.—Bras.
fastuosa, *Lem. in Fl. des Serres*, iii. (1847) sub t. 262. *Misc.* n. 42.—Hab.?
Fendleri, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 324.—Panama.
fimbriata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 119 = cordifolia.
firmula, *Baker, in Mart. Fl. Bras.* vi. II. 266.—Bras.
flavescens, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 481 = nummularia.
floribunda, *DC. Prod.* v. 198.—Madag.
Fockeana, Miq. in Linnaea, xvii. (1843) 68 = psilostachya.
fragrans, Tenore, ex Lem. in Fl. des Serres, iii. (1847) sub t. 261. Misc. n. 42 = fastuosa.
fulva, *Baker, in Mart. Fl. Bras.* vi. II. 222.—Bras.
Gabrieli, *Baker, l. c.* 235.—Bras.
glastifolia, Mart. ex Baker, l. c. 321 = Eupatorium dendroides.
glauca, *Mart. Reise Bras.* 401, nomen; *et ex Baker, l. c.* 224.—Bras.
Glaziovii, *Baker, l. c.* 251.—Bras.
glechomaefolia, Sch. Bip. ex Baker, l. c. 249 = scandens.
glomerata, *Spreng. Syst.* iii. 421.—Bras.
glumiflora, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 369 = Kanimia oblongifolia.
gonoclada, DC. Prod. v. 199 = cordifolia.
gracilis, *Sch. Bip. ex Miq. Stirp. Surin. Sel.* 187.—Guiana.
grandifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 246 = vitifolia.
Guaco, Humb. & Bonpl. Pl. Aequin. ii. 84. t. 105 = amara.

MIKANIA:—
Haenkeana, *DC. Prod.* v. 196.—Venezuela.
hastata, *Willd. Sp. Pl.* iii. 1742.—Ind. occ.
hastifolia, *Baker, in Mart. Fl. Bras.* vi. II. 270.—Bras.
hederaefolia, DC. Prod. v. 202 = glomerata.
hederaefolia, Sch. Bip. in Linnaea, xxx. (1859–60) 182 = triangularis.
hemisphaerica, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 254.—Bras.
hirsutissima, *DC. Prod.* v. 200.—Bras.
hispida, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 490.—Bras.
Hookeriana, *DC. Prod.* v. 195.—Guiana.
Hostmani, Miq. Stirp. Surin. Sel. 189 = cordifolia.
Houstonis, *Willd. Sp. Pl.* iii. 1742.—Ind. occ.
Huaco, Rieux, ex Spreng. Syst. iii. 422 = amara.
Ildefonsiana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 484 = Argyreiae.
Imrayana, *Griseb. Fl. Brit. W. Ind.* 363.—Ind. occ.
involucrata, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 243 = bracteosa.
itambana, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 481 (stambana).—Bras.
laevigata, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 241.—Bras.
laevis, *DC. Prod.* v. 194.—Bras.
lagoënsis, *Baker, in Mart. Fl. Bras.* vi. II. 220.—Bras.
lanigera, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 479 = leiolaena.
lanuginosa, *DC. Prod.* v. 201.—Bras.
Lasiandrae, *DC. l. c.* 189.—Bras.
latifolia, *Sm. in Rees, Cycl.* xxiii. n. 8.—Ins. S. Lucia.
laurifolia, *Willd. Sp. Pl.* iii. 1746.—N. Granat.
laxa, *DC. Prod.* v. 200.—Peruv.
Lechleri, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
leiolaena, *DC. Prod.* v. 188.—Bras.
leiostachya, *Benth. Pl. Hartw.* 201.—N. Granat.
leptotricha, *Baker, in Mart. Fl. Bras.* vi. II. 234.—Bras.
Liervalii, *Hérincq, in Hortic. Franç.* (1866) 18.—N. Granat.
ligustrifolia, *DC. Prod.* v. 191.—Bras.
Lima, Schlecht. ex Mart. in Flora, xxiv. (1841) II. Beibl. 110 = psilostachya.
Lindbergii, *Baker, in Mart. Fl. Bras.* vi. II. 232.—Bras.
Lindleyana, *DC. Prod.* v. 195.—Guian.
linearifolia, *DC. l. c.* 187.—Bras.
longicaulis, Wall. Cat. sub n. 3073 = Eupatorium longicaule.
longipes, *Baker, in Mart. Fl. Bras.* vi. II. 233.—Bras.
loxensis, *H. B. & K. Nov. Gen. et Sp.* iv. 136.—Am. austr.
Lundiana, *DC. Prod.* vii. 270.—Bras.
Mandonii, Sch. Bip. in Linnaea, xxxiv. (1865–66) 536 = Ophryosporus piquerioides.
Manettiae, DC. Prod. v. 191 = stipulacea.
maritima, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 489 = stipulacea.
Martiana, Gardn. l. c. 484 = hirsutissima.
melissaefolia, Willd. Sp. Pl. iii. 1747 (= *Nothites melissaefolia*).—Peruv.
mendocina, *Phil. Sert. Mendoc. Alt.* 20.—Reg. Argent.
menispermea, DC. Prod. v. 200 = cordifolia.
micrantha, H. B. & K. Nov. Gen. et Sp. iv. 134 = scandens.
microcephala, *DC. Prod.* v. 200.—Bras.
microdonta, *DC. l. c.*—Bras.
microlepis, *Baker, in Mart. Fl. Bras.* vi. II. 259.—Bras.
microphylla, *Sch. Bip. ex Baker, l. c.* 219.—Bras.
microptera, DC. Prod. v. 196 = scandens.
mollis, H. B. & K. Nov. Gen. et Sp. iv. 135 = cordifolia.
montevidensis, Spreng. ex Baker, in Mart. Fl. Bras. vi. II. 222 = ternifolia.
mossambicensis, *Steetz, in Peters, Reise Mossamb. Bot.* 377.—Afr. trop.
multinervia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 59.—Ecuador.
myricaefolia, *DC. Prod.* v. 188.—Madag.
myriocephala, *DC. l. c.* 191.—Bras.
natalensis, *DC. l. c.* 198.—Afr. austr.
neurocaula, *DC. l. c.* 188.—Bras.
nigricans, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 486.—Bras.

MIKANIA:—
nitidula, *Baker, in Mart. Fl. Bras.* vi. II. 243.—Bras.
nodosa, *Spreng. Syst.* iii. 423.—Cuba.
nodulosa, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 235.—Bras.
nummularia, *DC. Prod.* v. 188.—Bras.
oblongifolia, DC. l. c. = Kanimia oblongifolia.
obovata, *DC. l. c.* 192.—Bras.
obscura, Spreng. ex Baker, in Mart. Fl. Bras. vi. II. 362 = Eupatorium betonicaeforme.
obtusata, *DC. Prod.* v. 192.—Bras.
odorata, Lehm. in Hamb. Gartenz. v. (1849) 369 = fastuosa.
officinalis, *Mart. Reise Bras.* i. 283.—Bras.
opifera, Mart. l. c. 279 = cordifolia.
orinocensis, H. B. & K. Nov. Gen. et Sp. iv. 134 = scandens.
ovalis, *Griseb. Fl. Brit. W. Ind.* 363.—Ind. occ.
oxylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 220.—Bras.
oxyota, DC. Prod. v. 198 = scandens.
pachylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 239.—Bras.
paniculata, *DC. Prod.* v. 194.—Bras.
pannosa, *Baker, in Mart. Fl. Bras.* vi. II. 261.—Guian.
papillosa, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 325.—Ins. S. Doming.
Parkeriana, DC. Prod. v. 199 = scandens.
parvifolia, *Baker, in Mart. Fl. Bras.* vi. II. 226.—Bras.
penstemonoides, *DC. Prod.* v. 189.—Bras.
periplocifolia, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 243 = scandens.
pernambucensis, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 490.—Bras.
phaeoclados, *Mart. ex Baker, in Mart. Fl. Bras.* vi. II. 256.—Bras.
phyllopodon, *Griseb. in Goett. Abh.* xix. (1874) 170.—Reg. Argent.
pilosa, *Baker, in Mart. Fl. Bras.* vi. II. 234.—Bras.
pinnatiloba, *DC. Prod.* v. 187.—Bras.
platyphylla, DC. l. c. 195 = Hookeriana.
Poeppigii, Spreng. Syst. iii. 422 = scandens.
Pohliana, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 232.—Bras.
polybotrys, G. Don, ex Baker, l. c. 266 = psilostachya.
polystachya, DC. Prod. v. 190 = psilostachya.
populifolia, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 485.—Bras.
premnifolia, *Gardn. l. c.* 480.—Bras.
psilostachya, *DC. Prod.* v. 190.—Peruv.
psychotrioides, DC. l. c. 193 = obovata.
pterocaula, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 103.—Mexic.
pteropoda, *DC. Prod.* v. 191.—Bras.
puberula, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 482 = Selloi.
pubescens, Nutt. Gen. Am. ii. 136 = scandens.
punctata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 118 = pteropoda.
purpurascens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 371 = Kanimia purpurascens.
pyrifolia, *DC. Prod.* v. 195.—Madag.
racemosa, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 270 = ligustrifolia.
racemulosa, Benth. in Hook. Journ. Bot. ii. (1840) 41 = psilostachya.
racemulosa, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 325.—Venezuela.
ramosissima, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 483.—Bras.
ranunculifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 45.—Cuba.
Regnellii, Sch. Bip. in Linnaea, xxii. (1849) 572; et xxx. (1859–60) 182 = sessilifolia.
repanda, *La Llave, in Mosaico Mexicano*, ii. 299; *ex Hook. Kew Journ.* v. (1853) 78.—Mexic.
resinosa, Spreng. Neue Entdeck. ii. 134 = Eupatorium sordescens.
reticulata, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 480.—Bras.
reticulosa, *Wright, in Sauv. Fl. Cub.* 78.—Cuba.
retifolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 242.—Bras.
rotunda, *Griseb. Fl. Brit. W. Ind.* 364.—Ind. occ.
rubiginosa, Sm. in Rees, Cyclop. xxiii. n. 12 = cordifolia.

MIKANIA:—
rufa, *Benth. Pl. Hartw.* 201.—N. Granat.
rufescens, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 238.—Bras.
Ruiziana, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 53.—Peruv.
salviaefolia, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 487.—Bras.
salzmanniaefolia, *DC. Prod.* v. 193.—Bras.
sarcodes, *Baker, in Mart. Fl. Bras.* vi. II. 269.—Bras.
satureiaefolia, Willd. Sp. Pl. iii. 1747 = Stevia satureiaefolia.
scabra, DC. Prod. v. 190 = psilostachya.
scabrida, *Baker, in Mart. Fl. Bras.* vi. II. 258.—Bras.
scandens, *Willd. Sp. Pl.* iii. 1743.—Am. calid.
scansoria, DC. Prod. v. 202 = glomerata.
Selloi, *Spreng. Syst.* iii. 421.—Bras.
senecioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 223 = ternifolia.
sepiaria, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 488 = hirsutissima.
sericea, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 243.—Bras.
serratifolia, H. B. & K. Nov. Gen. et Sp. iv. 138 = Eupatorium serratifolium.
serratifolia, Sieber, ex DC. Prod. v. 181 = Eupatorium iresinoides.
serrulata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 230 = myriocephala.
sessilifolia, *DC. Prod.* v. 188.—Bras.
setigera, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 259.—Bras.
Sieberiana, DC. Prod. v. 196 = scandens.
smilacina, *DC. l. c.* 192.—Bras.
sophiaefolia, Spreng. Syst. iii. 423 = Eupatorium sophioides.
speciosa, *DC. Prod.* v. 196.—Peruv.
Spiraeastrum, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 243 = obtusata.
Sprucei, *Baker, l. c.* 264.—Bras.
stambana, Gardn. vide itambana.
stenocephala, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 249 = scandens.
stipitata, Sch. Bip. ex Miq. Stirp. Surin. Sel. 191 = amara.
stipulacea, *Willd. Sp. Pl.* iii. 1745.—Bras.
strigosa, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 119 = Lasiandrae.
strobilifera, *Gardn. l. c.* v. (1846) 479.—Bras.
stylosa, Gardn. l. c. 486 = microdonta.
suaveolens, H. B. & K. Nov. Gen. et Sp. iv. 135 = cordifolia.
subcordata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 118 = Argyreiae.
subcrenata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 243.—Reg. Argent.
subcymosa, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 448 = scandens.
subverticillata, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 224.—Bras.
surinamensis, Miq. Stirp. Surin. Sel. 188 = cordifolia.
Swartziana, *Griseb. Fl. Brit. W. Ind.* 363.—Ind. occ.
Tafallae, Spreng. Syst. iii. 422 = seq.
Tafallana, H. B. & K. Nov. Gen. et Sp. iv. 137 = amara.
tamoides, DC. Prod. v. 197 = scandens.
tenax, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
tenuiflora, Griseb. in Goett. Abh. xxiv. (1879) 174 = Eupatorium patens.
ternifolia, *DC. Prod.* v. 188.—Bras.
testudinaria, *DC. l. c.* 197.—Bras.
teucriifolia, Spreng. Neue Entdeck. ii. 134 = Eupatorium Gaudichaudianum.
thapsoides, DC. Prod. v. 189 = Kanimia thapsoides.
thunbergiaefolia, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 448 = cordifolia.
thunbergioides, *DC. Prod.* v. 198.—Madag.
thyrsoidea, *Baker, in Mart. Fl. Bras.* vi. II. 267.—Bras.
Tlalixcoyan, *La Llave, in Mosaico Mexicano*, ii. 299; *ex Hook. Kew Journ.* v. (1853) 79.—Mexic.
tomentosa, Willd. Sp. Pl. iii. 1744 = Senecio penicillatus.
triangularis, *Baker, in Mart. Fl. Bras.* vi. II. 246.—Bras.

MIKANIA:—
trichophila, *DC. Prod.* v. 200.—Bras.
trifolia, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
trinervis, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 244 = laevis.
trinitaria, *DC. Prod.* v. 194.—Ins. Trinit.
triphylla, *Spreng. ex Baker, in Mart. Fl. Bras.* vi. II. 263.—Bras.
umbellifera, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 119 = scandens.
ursina, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 261 = hirsutissima.
urticaefolia, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 244 = cordifolia.
variabilis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 486 = scandens.
variabilis, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 257.—Peruv.
Vauthieriana, *Baker, in Mart. Fl. Bras.* vi. II. 269.—Bras.
verrucosa, Spreng. Syst. iii. 423 = Eupatorium celtidifolium.
verticillata, *Sch. Bip. in Bonplandia*, ix. (1861) 175.—Bras.
viminaria, Steud. Nom. ed. II. ii. 145, sphalm. = seq.
viminea, *DC. Prod.* v. 187.—Bras.
violacea, *Regel, Cat. Pl. Hort. Aksakov.* 93, nomen.—Hab.?
viscosa, Spreng. Neue Entdeck. i. 277 = Symphyopappus viscosus.
vismiaefolia, *DC. Prod.* v. 189.—Bras.
vitifolia, *DC. l. c.* 202.—Bras.
volubilis, Willd. Sp. Pl. iii. 1743 = scandens.
Warmingii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 242.—Bras.

MILDEA, Griseb. Cat. Pl. Cub. 63 (1866) = **Verhuellia**, Miq. (Piperac.).
elegans, Griseb. l. c. = P. elegans.
hydrocotylifolia, Griseb. l. c. = P. cordifolia.

MILDEA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 88 (1867) = **Paranephelium**, Miq. (Sapind.).
gibbosa, Miq. l. c. 89 = P. gibbosa.
xestophylla, Miq. l. c. = P. xestophylla.

MILHANIA, Neck. Elem. iii. 24 (1790) = **Ipomoea**, Linn. (Convolv.).

MILHANIA, Rafin. Fl. Tellur. iv. 71 (1836) = **Calystegia**, R. Br. (Convolv.).
bracteata, Rafin. l. c. = Ipomoea setifera?
Catesbiana, Rafin. l. c. = C. Catesbeiana.
involucrata, Rafin. l. c. = Ipomoea pileata?
mutabilis, Rafin. l. c. = C. sepium?
pentaphyla, Rafin. l. c. = Ipomoea pentaphylla.
riparia, Rafin. l. c. = C. riparia.
sepium, Rafin. l. c. = C. sepium.
spithamea, Rafin. l. c. = C. spithamaea.
villosa, Rafin. l. c. = C. villosa.

MILIARIUM, Moench, Meth. 204 (1794) = **Milium**, Linn. (Gramin.).
effusum, Moench, l. c. = Milium effusum.

MILIUM, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 143.
MILIARIUM, Moench, Meth. 204 (1794).
adscendens, *Roxb. Hort. Beng.* 8, nomen.—Nepal.
alepense, Cav. Desc. Index, 619, vide *halepense*.
alternans, *Bubani, in Nuov. Giorn. Bot. Ital.* v. (1873) 317.—Hispan.
amphicarpon, Pursh, Fl. Am. Sept. i. 62 = Amphicarpum Purshii.
arundinaceum, Koen. ex Steud. Syn. Pl. Gram. 70 = Panicum sparsum, *Nees*.
arundinaceum, Sibth. & Sm. Fl. Graec. Prod. i. 45; Fl. Graec. i. 50. t. 66 (= *Piptatherum miliaceum*).—Europ. austr.
attenuatum, Moench, Meth. 204 = Panicum miliaceum.
capense, Linn. Mant. ii. 185 = Danthonia papillosa.
capillaceum, Koen. ex Steud. Nom. ed. II. ii. 145 = Paspalum pedicellare.
capillare, Moench, Meth. 203 = Panicum capillare.
capillare, Roth, Nov. Pl. Sp. 39 = Paspalum brevifolium.

MILIUM:—
capillare, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 271 = Panicum commelinaefolium.
ciliatum, Moench, Meth. Suppl. 66 = Panicum dichotomum.
ciliatum, Muhl. Desc. Gram. 77 = Amphicarpum Purshii.
cimicinum, Linn. Mant. ii. 184 = Panicum cimicinum.
clandestinum, Moench, Meth. 204 = Panicum clandestinum.
coerulescens, Desf. Fl. Atlant. i. 66 = Oryzopsis coerulescens.
colonum, Moench, Meth. 202 = Panicum colonum.
coloratum, Moench, l. c. 203 = Panicum coloratum.
comosum, Poir. Encyc. Suppl. i. 257 (= *Piptatherum miliaceum*).—Europ. austr.
compactum, Cav. Desc. 307 = Sorghum vulgare.
compressum, Poir. Encyc. Suppl. i. 259 = Agrostis compressa, *Poir.*
compressum, Sw. Prod. Veg. Ind. Occ. 24 = Paspalum platycaule.
confertum, Guss. Fl. Sic. Syn. i. 131 = vernale.
confertum, *Linn. Sp. Pl.* 61.—Europ.
confertum, Mill. Gard. Dict. ed VIII. n. 4 = effusum.
Crus-galli, Moench, Meth. 202 = Panicum Crus-galli.
cuspidatum, Spreng. Syst. i. 251 = Oryzopsis cuspidata.
Dactylon, Moench, Meth. Suppl. 67 = Cynodon Dactylon.
digitatum, Sw. Prod. Veg. Ind. Occ. 24 = Panicum sanguinale.
distichum, Muhl. Desc. Gram. 78 = Paspalum distichum.
effusum, *Linn. Sp. Pl.* 61.—Reg. bor. temp.
effusum, Lour. Fl. Cochinch. 49 = Panicum miliare.
elegans, Steud. Nom. ed. II. ii. 146 (= *Piptatherum elegans*).—Hab. ?
esculentum, Moench, Meth. 203 = Panicum miliaceum.
filiforme, Lag. Gen. et Sp. Nov. 2 = Paspalum velutinum.
filiforme, Roxb. Hort. Beng. 8; Fl. Ind. i. 314 = Paspalum brevifolium.
frutescens, *Link Enum. Hort. Berol.* i. 72.—Oriens.
frutescens, Sieber, ex Kunth, Enum. Pl. i. 177 (= *Piptatherum miliaceum*).—Europ. austr.; Sibir.
gaditanum, Hort. ex Steud. Nom. ed. II. ii. 146 (= *Urachne parviflora*).—Europ. austr.
gallecicum, Roem. & Schult. Syst. ii. 320 = Aira Lagascae.
globosum, Thunb. Fl. Jap. 49 = Isachne australis.
halepense, Cav. Desc. 306 = Sorghum halepense.
hirsutum, Beauv. Agrost. 13 = Panicum leucopheum.
holciforme, Spreng. Syst. i. 251 = Oryzopis holciformis.
juncoides, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1883) 126.—Reg. Argent.
lanatum, Roem. & Schult. Syst. ii. 322 = Anthaenantia lanata.
latifolium, Cav. Ic. iii. 37 = Paspalum racemosum.
latifolium, Moench, Meth. 204 = Panicum latifolium.
lendigerum, Delile, ex Boiss. Fl. Orient. v. 519 = Triplachne nitens.
lendigerum, Linn. Sp. Pl. ed. II. 91 = Gastridium australe.
maritimum, Koen. ex Steud. Nom. ed. II. ii. 146 = Zoysia pungens.
maritimum, Pall. Tabl. Taur. 45 = Aeluropus laevis.
maritimum, Roxas Clem. Ensay. Vid. 285 = Agrostis alba.
Matrella, Poir. ex Steud. Nom. ed. II. ii. 146 = Zoysia pungens.
membranaceum, Moench, Meth. 203 = Panicum sanguinale.
mexicanum, Spreng. Syst. i. 251 = Oryzopsis fimbriata.
microspermum, Hornem. Hort. Hafn. Suppl. 8 (= *Piptatherum miliaceum*).—Ins. Canar.?
microspermum, *Lag. Gen. et Sp. Nov.* 2.—Mexic.
minimum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 56 = Gastridium australe.
minutiflorum, Trin. in Mem. Acad. Pétersb. Sér. VI. iii. (1835) 121 = Panicum Perrottetii.
molle, *Kunth, Rév. Gram.* i. 28.—Bras.
Montianum, Parl. Fl. Ital. i. 156 = vernale.
Muehlenbergianum, *Schult. Mant.* ii. 178.—Am. bor.
multiflorum, Cav. Desc. 36 (= *Piptatherum miliaceum*).—Europ. austr.; Sibir.
muticum, Hort. Berol. ex Spreng. Nov. Prov. 20 = Gastridium muticum.
natans, Spreng. Syst. i. 250 = Luziola peruviana.

MILIUM:—
nemorosum, Moench, Meth. Suppl. 67 = Ichnanthus nemorosus.
nigricans, *Ruiz & Pav. Fl. Per.* i. 47.—Peruv.
nigrum, Lam. Fl. Fr. iii. 568 = Oryzopsis paradoxa.
orinoccensis, Willd. ex Steud. Nom. ed. II. ii. 146 = Panicum aturense.
paniceum, Host, Fl. Austr. i. 94 = Polypogon monspeliensis.
paniceum, Sw. Prod. Veg. Ind. Occ. 24 = Panicum filiforme.
Panicum, Mill. Gard. Dict. ed. VIII. n. 1 = Panicum miliaceum.
paradoxum, All. Fl. Pedem. ii. 236 = Oryzopsis coerulescens.
paradoxum, Linn. Syst. ed. X. 872; Scop. Fl. Carn. ed. II. i. 58 = Oryzopsis paradoxa.
paspaloides, Ell. Sketch, i. 103 = Paspalum distichum.
pauciflorum, Hort. Matrit. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 11 (= *Piptatherum miliaceum*).—Europ. austr.
phleoides, *Sweet, Hort. Brit.* ed. II. 555, nomen.—Ind. or.
poaeforme, Balansa, in Bull. Soc. Bot. Fr. v. (1858) 170 = trichopodum.
polystachyon, Spreng. Syst. i. 251 = Eriochloa polystachya.
pulcher, *Roxb. Hort. Beng.* 8, nomen.—China.
punctatum, Linn. Syst. ed. X. 872 = Eriochloa polystachya.
pungens, Torr. ex Spreng. Neue Entdeck. ii. 102 = Oryzopsis canadensis.
purpureum, Lapeyr. Hist. Abr. Pl. Pyr. 33 = Oryzopsis coerulescens.
racemosum, Sm. in Rees, Cyclop. xxiii. n. 15 = Oryzopsis melanocarpa.
ramosum, Retz. Obs. vi. 22 = Eriochloa annulata.
rigidifolium, Roem. & Schult. Syst. ii. 319 = Panicum rigidifolium.
rigidum, Poir. Encyc. Suppl. i. 257 = Panicum rigidifolium.
roseum, *Spreng. Syst.* i. 251.—Ins. Mascar.
sanguinale, Roxb. Hort. Beng. 8; Fl. Ind. i. 315 = Panicum pedicellatum.
scabrum, Guss. Fl. Sic. Prod. i. 54 = Gastridium muticum.
scabrum, Rich. in Merlet, Herbor. Maine-et-Loire (1809), 131 = effusum.
Schmidtianum, C. Koch, in Linnaea, xxi. (1848) 438 = effusum.
Scopolianum, Willem. ex Steud. Nom. ed. II. ii. 146 = Oryzopsis paradoxa.
setaceum, Koen. ex Steud. l. c. = Paspalum granulare.
Sorghum, Cav. Desc. 306 = Sorghum vulgare.
sparsum, Mill. Gard. Dict. ed. VIII. n. 2 = Panicum ambitiosum.
striatum, Buch.-Ham. ex Wall. Cat. n. 8747 A = Paspalum pedicellatum.
strictum, Willd. ex Steud. Nom. ed. II. ii. 146 = Trachypogon polymorphus.
tenellum, Cav. Ic. iii. 37 = Airopsis globosa.
Thomasii, Duby, Bot. Gall. i. 505 (= *Piptatherum miliaceum*).—Europ. austr.; Sibir.
tomentosum, Koen. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 220 = Panicum subeglume.
transsilvanicum, Schur, Enum. Pl. Transs. 741 = effusum.
trichopodium, *Boiss. Diagn.* Ser. I. xiii. 45.—As. Min.; Syria.
undulatifolium, Moench, Meth. 202 = Oplismenus undulatifolius.
velutinum, DC. Cat. Hort. Monsp. 126 = Paspalum velutinum.
vernale, *Bieb. Fl. Taur. Cauc.* i. 53.—Graecia; Reg. Caucas.
vernale, Van Hall, Syn. 32 (1821); Kunth, Enum. Pl. i. 66 = effusum.
verticillatum, *Boiss. & Bal. Diagn.* Ser. II. iv. 126.—Phrygia.
villosum, Sw. Prod. Veg. Ind. Occ. 24 = Panicum leucophaeum.

MILIUM, [Tourn.] Adans. Fam. ii. 34 (1763) = **Panicum**, Linn. (Gramin.).

MILIUSA, Leschen. ex A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 213. *ANONACEAE*, Benth. & Hook. f. i. 28.
HYALOSTEMMA, Wall. Cat. n. 6434 (1831).

MILIUSA:—
Bailloni, *Pierre, Fl. For. Cochinch.* t. 38.—Anam.; Ins. Philipp.
Brahei, F. Muell. Fragm. viii. 159 = Saccopetalum Brahei.
campanulata, *Pierre, Fl. For. Cochinch.* t. 41.—Cambodia.
fusca, *Pierre, l. c.* 42.—Cambodia.
Horsfieldii, Baill. ex Pierre, l. c. sub t. 38 = Saccopetalum Horsfieldii.
indica, *Leschen. ex A. DC. Mém. Anon.* 37.—Ind. or.
macrocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 150.—Reg. Himal.
macropoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 40.—Borneo.
mollis, *Pierre, Fl. For. Cochinch.* t. 40.—Cambodia.
montana, Gardn. ex Hook. f. & Thoms. Fl. Ind. i. 148 = indica.
nilagirica, *Bedd. Ic. Pl. Ind. Or.* i. 18.—Ind. or.
Roxburghiana, *Hook. f. & Thoms. Fl. Ind.* i. 150.—Reg. Himal.; Burma.
sclerocarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 291.—Burma.
tristis, *Kurz, l. c.* xliii. (1874) II. 58, 181.—Burma.
velutina, *Hook. f. & Thoms. Fl. Ind.* i. 151.—Ind. or.
Wallichiana, Hook. f. & Thoms. l. c. 149 = Roxburghiana.
Wightiana, *Hook. f. & Thoms. l. c.*—Ind. or.
zeylanica, *Gardn. ex Hook. f. & Thoms. l. c.*—Zeylan.

MILLA, Cav. Ic. ii. 76. t. 196 (1793). *LILIACEAE*, Benth. & Hook. f. iii. 799.
ASKOLAME, Rafin. Fl. Tellur. ii. 11 (1836).
GYRENIA, Knowles & Westc. ex Loud. Hort. Brit. Suppl. ii. 639 (1839).
IPHEION, Rafin. Fl. Tellur. ii. 12 (1836).
MILLEA, Willd. Sp. Pl. ii. 62 (1799).
TULOPHOS, Rafin. Fl. Tellur. iii. 71 (1836).
andicola, *Baker, in Journ. Linn. Soc.* xi. (1871) 381.—Peruv.; Boliv.
aurea, *Baker, l. c.* 386.—Reg. Argent.
biflora, *Cav. Ic.* ii. 76. t. 196.—Arizona; Mexic.
bivalvis, *Baker, in Journ. Linn. Soc.* xi. (1871) 386.—Chili.
bonariensis, Gill. ex Baker, l. c. 382 = uniflora.
borbonica, Baker, Fl. Maurit. 373 = Nothoscordum borbonicum.
brevipes, *Baker, in Journ. Linn. Soc.* xi. (1871) 386.—Chili.
capitata, Baker, l. c. 381 = Brodiaea capitata.
coerulea, Scheele, in Linnaea, xxv. (1852) 260 = Androstephium violaceum.
crocea, Baker, in Journ. Linn. Soc. xi. (1871) 384 = Brodiaea crocea.
grandiflora, Baker, l. c. 380 = Brodiaea Douglasii.
hirtella, *Baker, l. c.* 385.—Reg. Argent.
hyacinthina, Baker, l. c. = Brodiaea lactea.
ixioides, Baker, l. c. 383 = Brodiaea ixioides.
laxa, Baker, l. c. 384 = Brodiaea laxa.
Leichtlinii, *Baker, in Gard. Chron.* (1875) I. 234.—Reg. Andin.
macrostemon, *Baker, in Journ. Linn. Soc.* xi. (1871) 381.—Reg. Argent.
maritima, S. Wats. in Bot. King's Exped. 354 = Muilla maritima.
nivalis, *Baker, in Journ. Linn. Soc.* xi. (1871) 383.—Chili.
patagonica, *Baker, l. c.* 382.—Patagon.
pedunculatis, Baker, l. c. 384 = Brodiaea peduncularis.
Poeppigiana, *Baker, l. c.* 383.—Chili.
porrifolia, *Baker, l. c.* 386.—Chili.
Sellowiana, *Baker, l. c.* 383.—Reg. Argent.
sessiliflora, *Baker, l. c.* 382.—Chili.
setacea, *Baker, l. c.* 385.—Tucuman.
subbiflora, *Baker, l. c.*—Chili.
Tweediana, *Griseb. in Goett. Abh.* xxiv. (1879) 318.—Reg. Argent.
uniflora, *R. Grah. in Edinb. N. Phil. Journ.* (Jan. 1833) 174.—Reg. Argent.
violacea, *Baker, in Journ. Bot.* xii. (1874) 5.—Chili.
vittata, *Griseb. in Goett. Abh.* xxiv. (1879) 318.—Reg. Argent.

MILLA, Vand. Fl. Lusit. et Bras. Spec. 43. t. 3. f. 23 (1788) = **Herpestis**, Gaertn. (Scrophul.).

MILLANIA, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 1852 [1856] 128 = **Pemphis**, Forst. (Lythrar.).
rupestris, Zipp. ex Blume, l. c. = P. acidula.

MILLEA, Willd. Sp. Pl. ii. 62 (1799) = **Milla**, Cav. (Liliac.).

MILLEFOLIUM, Tourn. ex Adans. Fam. ii. 128 (1763) = **Achillea**, Linn. (Compos.).
compactum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 91 = A. compacta.
nobile, Fourr. l. c. = A. nobilis.
odoratum, Fourr. l. c. = A. odorata.
officinale, Gueldenst. ex Ledeb. Fl. Ross. ii. 532 = A. Millefolium.
setaceum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 91 = A. setacea.
tomentosum, Fourr. l. c. = A. tomentosa.
vulgare, Gueldenst. ex Ledeb. Fl. Ross. ii. 532 = A. Millefolium.

MILLEGRANA, Adans. Fam. ii. 269 (1763) = **Radiola**, Dill. (Lineae).

MILLEGRANA, Juss. ex Turp. in Ann. Mus. Par. vii. (1806) 220 = **Cypselea**, Turp. (Ficoid.).
Surian, Steud. Nom. ed. II. ii. 146, sphalm. = C. humifusa.

MILLERA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.

MILLERIA, Houst. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 344.
alba, Hort. ex Pers. Syn. ii. 489 = Baltimora recta.
angustifolia, Cav. Ic. iii. 12. t. 223 = Flaveria angustifolia.
biflora, Linn. Sp. Pl. 919 = Elvira Martyni.
Contrahierba, Lam. Encyc. iv. 183 = seq.
Contrayerba, Cav. Ic. i. 2 = Flaveria Contrayerba.
dichotoma, Cav. l. c. 58 = quinqueflora.
glandulosa, DC. Prod. v. 503 = quinqueflora.
maculata, Mill. Gard. Dict. ed. VIII. n. 2 = quinqueflora.
quinqueflora, *Linn. Sp. Pl.* 919.—Mexic.
trifolia, *Mill. Gard. Dict.* ed. VIII. n. 4.—Mexic.

MILLETIA, Meissn. Gen. 95 (1837) = seq.

MILLETTIA, Wight & Arn. Prod. 263 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 498.
CALLERYA, Endl. Gen. Suppl. iii. 104 (1843).
FORNASINIA, Bertol. Misc. Bot. viii. 18. t. 1 (1849).
MARQUARTIA, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 35. t. 1.
OTOSEMA, Benth. in Miq. Pl. Jungh. 248 (1852).
PADBRUGGEA, Miq. Fl. Ind. Bat. i. 1. 150 (1855).
aboënsis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 130.—Afr. trop.
amygdalina, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 296.—Madag.
antsahalanbensis, *Baill. l. c.* 395.—Madag.
argentea, *Miq. Fl. Ind. Bat.* i. 1. 156.—Java.
atropurpurea, *Benth. in Miq. Pl. Jungh.* 249.—Burma; Malaya.
auriculata, *Baker, ex Brand. For. Fl. Ind.* 138.—Reg. Himal.
australis, *Benth. in Miq. Pl. Jungh.* 250.—Ins. Norfolk.
Blackii, F. Muell. Fragm. ii. 123 = Lonchoçarpus Blackii.
Bojeri, *Vatke, in Bremen, Abh.* vii. (1882) 246.—Madag.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 69.—Burma.
caerulea, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 107.—Burma; Malaya.
caffra, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 99.—Afr. austr.
cana, *Benth. in Miq. Pl. Jungh.* 250.—Burma.
caudata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 109.—Reg. Himal.
Championi, *Benth. in Hook. Kew Journ.* iv. (1852) 74.—Ins. Hongkong.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 395.—Madag.
chinensis, Benth. in Miq. Pl. Jungh. 249 = Wistaria chinensis.
cinerea, *Benth. l. c.*—Reg. Himal.; Burma.
cognata, *Hance, in Journ. Bot.* xviii. (1880) 260.—China.
drastica, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 128.—Afr. trop.

MILLETTIA:—
elliptica, Wight & Arn. Prod. 263 = Derris elliptica.
eriantha, *Benth. in Miq. Pl. Jungh.* 250.—Malacca.
extensa, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 109.—Burma.
ferruginea, *Hochst. in Flora*, xxix. (1846) 597, in syn.; *Baker, in Oliver, Fl. Trop. Afr.* ii. 130.—Afr. trop.
fruticosa, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 109.—Ind. or.
glaucescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 67.—Burma.
gracilis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 129.—Afr. trop.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 395.—Madag.
Griffoniana, Baill. Adansonia, vi. (1866) 222 = Thonningii.
heterophylla, Blume, ex Miq. Fl. Ind. Bat. i. 1. 154 = sericea.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1859) 386.—Japon.
leiogyna, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 67.—Burma.
lenneoides, *Vatke, in Linnaea*, xliii. (1880–82) 335.—Madag.
leucantha, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 68 = pendula.
leucantha, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 223.—Afr. trop.
luzonensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 456.—Ins. Philipp.
macrophylla, *Benth. in Hook. Ic. Pl.* t. 788–9.—Afr. austr.
macrophylla, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 276 = auriculata.
madagascariensis, *Vatke, in Linnaea*, xliii. (1880–82) 101.—Ins. Nossibé.
Maingayi, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 110.—Singapore.
Mannii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 127.—Afr. trop.
megasperma, *Benth. Fl. Austral.* ii. 211.—Austral.
monticola, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 67.—Burma.
nitida, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 484.—China.
nudiflora, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 130.—Afr. trop.
obtusa, Blume, ex Miq. Fl. Ind. Bat. i. 1. 154 = sericea.
ovalifolia, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 68.—Burma.
pachycarpa, *Benth. in Miq. Pl. Jungh.* 250.—Reg. Himal.; Malaya.
pallida, Dalz. in Journ. Linn. Soc. xiii. (1873) 187 = racemosa.
paniculata, *Miq. Fl. Ind. Bat. Suppl.* 301.—Sumatra.
pendula, *Benth. in Miq. Pl. Jungh.* 250.—Burma.
pirifolia, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 215.—Afr. trop.
piscidia, *Wight & Arn. Prod.* 263.—Reg. Himal.
pubinervis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 68.—Burma.
pulchra, *Kurz, l. c.* 69.—Reg. Himal.; Burma.
racemosa, *Benth. in Miq. Pl. Jungh.* 249.—Ind. or.
reticulata, *Benth. l. c.*—China.
rhodantha, *Baill. Adansonia*, vi. (1866) 223.—Afr. trop.
rostrata, *Miq. Fl. Ind. Bat.* i. 1. 155.—Java.
rubiginosa, *Wight & Arn. Prod.* 263.—Ind. or.
sericea, *Wight & Arn. l. c.*—Burma; Malaya.
sericosema, *Hance, in Journ. Bot.* xx. (1882) 259.—China.
speciosa, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 73.—China.
speciosa, *Welw. Apont.* 585. n. 23.—Afr. trop.
splendens, *Wight & Arn. Prod.* 263.—Ind. or.
splendidissima, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 156.—Borneo.
subcordata, Blume, l. c. 154 = sericea.
Sutherlandi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 211.—Afr. austr.
tetraptera, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 69.—Burma.
Thonningii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 128.—Afr. trop.
thyrsiflora, Benth. in Miq. Pl. Jungh. 249, in adnot. = Derris thyrsiflora.

MILLETTIA:—
turgida, *Miq. Fl. Ind. Bat. Suppl.* 300.—Sumatra.
urophylla, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 129.—Afr. trop.
versicolor, *Welw. l. c.*—Afr. trop.
xylocarpa, *Miq. Fl. Ind. Bat.* i. 1. 157.—Java.
Zollingeriana, *Miq. l. c.* 156.—Java.

MILLIGANIA, Hook. f. in Hook. Ic. Pl. t. 299 (1840) = **Gunnera**, Linn.
cordifolia, Hook. f. l. c. = G. cordifolia.

MILLIGANIA, Hook. f. in Hook. Kew Journ. v. (1853) 296. t. 9. *LILIACEAE*, Benth. & Hook. f. iii. 781.
densiflora, *Hook. f. in Hook. Kew Journ.* v. (1853) 298.—Tasmania.
Johnstoni, *F. Muell. ex Benth. Fl. Austral.* vii. 26.—Tasmania.
longifolia, *Hook. f. in Hook. Kew Journ.* v. (1853) 297.—Tasmania.
stylosa, *F. Muell. ex Benth. Fl. Austral.* vii. 27.—Tasmania.

MILLINA, Cass. in Dict. Sc. Nat. xxxi. 89 (1824) = **Leontodon**, Linn. (Compos.).
arabica, Boiss. Diagn. Ser. I. xi. 38 = L. arabicus.
chicoracea, Pomel, Nouv. Mat. Fl. Atl. 270 = L. chicoraceus.
hispidula, Boiss. Diagn. Ser. I. xi. 38 = L. hispidulus.
hyoseroides, DC. Prod. vii. 110 = L. creticus.
leontodontoides, Cass. in Dict. Sc. Nat. xxxi. 90 = L. fasciculatus.
persica, Boiss. Diagn. Ser. I. vii. 3 = L. hispidulus.

MILLINGTONIA, Linn. f. Suppl. 45, 291 (1781). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1040.
NEVRILIS, Rafin. Sylva Tellur. 138 (1838).
Clematis, *Schrad. in Goett. Gel. Anz.* i. (1821) 712.—Bras.
dubiosa, Span. in Linnaea, xv. (1841) 326 = hortensis.
hortensis, *Linn. f. Suppl.* 291.—Burma.
pinnata, Blanco, Fl. Filip. 501 = Stereosperma pinnatum.
quadripinnata, Blanco, l. c. = Stereosperma quadripinnatum.

MILLINGTONIA, Roxb. ex Wight & Arn. Prod. 241, in syn. (1834) = **Flemingia**, Roxb. (Legumin.).
congesta, Roxb. l. c. = F. congesta.
nana, Roxb. l. c. 242 = F. congesta.
semialata, Roxb. l. c. 241 = F. congesta.
stricta, Roxb. l. c. = F. stricta.

MILLINGTONIA, Roxb. Fl. Ind. ed. Carey, i. 102 (1820) = **Meliosma**, Blume (Sabiac.).
acuminata, Royle, Illustr. Bot. Himal. 139 = Mel. pungens.
alba, Schlecht. in Linnaea, xvi. (1842) 395 = Mel. alba.
Arnottiana, Wight, Illustr. i. 144. t. 53 = Mel. Arnottiana.
congesta, Wight & Arn. in Edinb. N. Phil. Journ. xv. (1833) 179 = Mel. pungens.
dilleniifolia, Wall. ex Wight & Arn. l. c. = Mel. dilleniaefolia.
ferruginea, Schult. Mant. i. Addit. ii. 250 = Mel. ferruginea.
integrifolia, Wall. Cat. n. 8114 = Mel. pungens, simplicifolia.
lanceolata, Schult. Mant. i. Addit. ii. 250 = Mel. lanceolata.
nitida, Schult. l. c. = Mel. nitida.
pinnata, Korth. ex Blume, Rumphia, iii. 200 = Mel. confusa.
pinnata, Roxb. Hort. Beng. 3; Fl. Ind. i. 104 = Mel. pinnata.
pungens, Wall. ex Wight & Arn. in Edinb. N. Phil. Journ. xv. (1833) 178 = Mel. pungens.
sambucina, Jungh. in Hoev. & De Vriese, Tijdschr. viii. (1841) 365 = Mel. glauca.
simplicifolia, Roxb. Hort. Beng. 3; Pl. Corom. iii. 50 = Mel. simplicifolia.
simplicifolia, Wall. Cat. n. 8114 A = Mel. Wightii.
sumatrana, Jack, in Malay. Misc. ii. (1822) n. VII. 30 = Mel. nitida.

MILLOTIA, Cass. in Ann. Sc. Nat. Sér. I. xvii. (1829) 416. *COMPOSITAE*, Benth. & Hook. f. ii. 315.
glabra, Steetz, in Lehm. Pl. Preiss. i. 458 = tenuifolia.
Greevesii, *F. Muell. Fragm.* iii. 18. t. 19.—Austral.
Kempei, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 2.—Austral.
myosotidifolia, Steetz, in Lehm. Pl. Preiss. i. 457 = tenuifolia.
robusta, Steetz, l. c. 458 = tenuifolia.
tenuifolia, *Cass. in Ann. Sc. Nat.* Sér. I. xvii. (1829) 417.—Austral.

MILNEA, Rafin. Sylva Tellur. 167 (1838) = **Ardisia**, Sw. (Myrsin.).

MILNEA, Roxb. Hort. Beng. 18 (1814); Fl. Ind. i. 637 (1832) = **Aglaia**, Lour. (Meliac.).
apiocarpa, Thw. Enum. Pl. Zeyl. 60 = A. apiocarpa.
argentea, Reinw. Cat. Gew. Buitenz. 71 = A. argentea.
Blumei, Teijsm. & Binn. Cat. Hort. Bog. 211 = A. Diepenhorstii.
dulcis, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 253.—Ins. Celebes.
edulis, Roxb. Hort. Beng. 18; Fl. Ind. i. 637 = A. edulis.
lancifolia, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 165 = Hearnia lancifolia.
montana, Jack, in Trans. Linn. Soc. xiv. (1823) 118.—Sumatra.
montana, Miq. Fl. Ind. Bat. Suppl. 197 = A. oligophylla.
montana, Teijsm. & Binn. Cat. Hort. Bog. 211 = A. subgrisea.
racemosa, M. Roem. Syn. Hesper. 98 = A. edulis.
Roxburghiana, Wight & Arn. Prod. 119 = A. Roxburghiana.
Sulingi, Teijsm. & Binn. Cat. Hort. Bog. 211 = A. Sulingi.
undulata, Wall. ex C. DC. in DC. Monog. Phan. i. 605 = A. undulata.

MILTIANTHUS, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 197 (Rel. Lehm. 58). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 266.
portulacoides, *Bunge, in Abh. Naturf. Ver. Riga*, i. (1847) 197 (*Rel. Lehm.* 58).—Afghan.

MILTITZIA, A. DC. in DC. Prod. ix. 296 (1845). **Emmenanthe**, Benth. (Hydroph.).
lutea, A. DC. l. c. = E. lutea.

MILTONIA, Lindl. Bot. Reg. sub t. 1976 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 563.
GYNIZODON, Rafin. Fl. Tellur. iv. 40 (1836).
MACROCHILUS, Knowles & Westc. Flor. Cab. t. 45 (1837).
anceps, *Lindl. Fol. Orch. Milt.* 3.—Bras.
bicolor, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 589, nomen.—Bras.
Bluntii ×, Reichb. f. in Gard. Chron. (1879) II. 489.—Bras.
candida, *Lindl. Bot. Reg.* (1838) *Misc.* 25.—Bras.
cereola, *Lem. Illustr. Hortic.* xii. (1865) t. 446.—Bras.
Clowesii, *Lindl. Sert. Orch.* t. 34.—Bras.
cuneata, *Lindl. Bot. Reg.* (1844) *Misc.* 26.—Bras.
cyrtochiloides, *Rodrig. Orch. Nov.* i. 98.—Bras.
Endresii, *Nichols. Dict. Gard.* ii. 368.—Am. centr.
festiva, *Nichols. l. c.*—Hab.?
flava, *Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 589, nomen.—Bras.
flavescens, *Lindl. Sert. Orch.* sub t. 48.—Bras.
Karwinskii, Lindl. in Journ. Hort. Soc. iv. (1849) 83 = Odontoglossum Karwinskii.
Lamarckeana ×, Reichb. f. in Gard. Chron. (1885) I. 530.—Bras.
Moreliana, Hort. ex Lindl. Fol. Orch. Milt. 1 = spectabilis.
odorata, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 589, nomen.—Bras.
Phalaenopsis, *Nichols. Gard. Dict.* ii. 369.—N. Granat.
pulchella, Hort. ex Linden, Pescatorea, i. (1860) sub t. 44 = M. Phalaenopsis.
Regnelli, *Reichb. f. in Linnaea*, xxii. (1849) 851.—Bras.
Roezlii, *Nichols. Dict. Gard.* ii. 369.—N. Granat.
rosea, Versch. ex Lem. Illustr. Hortic. xiv. (1867) t. 524 = spectabilis.

MILTONIA:—
Rosina, *Rodrig. Orch. Nov.* i. 99.—Bras.
Russelliana, *Lindl. Sert. Orch.* sub t. 48.—Bras.
speciosa, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 129.—Bras.
spectabilis, *Lindl. Bot. Reg.* sub t. 1976.—Bras.
stellata, Lindl. Sert. Orch. sub t. 48 = flavescens.
vexillaria, *Nichols. Dict. Gard.* ii. 369.—N. Granat.
Warneri, Nichols. l. c. = spectabilis.
Warszewiczii, *Reichb. f. in Xenia Orch.* i. 132.—Bras.

MILTUS, Lour. Fl. Cochinch. 302 (1790) = **Gisekia**, Linn. (Ficoid.).
africana, Lour. l. c. = G. Miltus.

MIMAECYLON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 78 = **Memecylon**, Linn. (Melastom.).

MIMELA, Phil. in Anal. Univ. Chil. (1865) II. 336. *COMPOSITAE*.
pedicilariaefolia, *Phil. l. c.*—Chili.

MIMETANTHE, Greene, in Bull. Calif. Acad. i. IV. (1885) 181 = **Mimulus**, Linn. (Scrophul.).
pilosa, Greene, l. c. = Mimulus exilis.

MIMETES, Salisb. Parad. Lond. sub n. 67 (1807). *PROTEACEAE*, Benth. & Hook. f. iii. 171.
DEASTELLA, Loud. Hort. Brit. 38, in syn. (1830).
OROTHAMNUS, Pappe, ex Hook. Bot. Mag. t. 4357 (1848).
argentea, *Knight, Prot.* 67.—Afr. austr.
capitulata, *R. Br. in Trans. Linn. Soc.* x. (1810) 106.—Afr. austr.
capitulata, Sieber, ex Schult. Mant. iii. 268 = hirta.
coriacea, *Steud. Nom.* ed. II. ii. 147; *Meissn. in DC. Prod.* xiv. 266.—Afr. austr.
cucullata, R. Br. in Trans. Linn. Soc. x. (1810) 107 = lyrigera.
decapitata, *Meissn. in DC. Prod.* xiv. 262.—Afr. austr.
dichotoma, *Roem. & Schult. Syst.* iii. 383.—Afr. austr.
divaricata, *R. Br. in Trans. Linn. Soc.* x. (1810) 111.—Afr. austr.
fimbriaefolia, *Knight, Prot.* 65.—Afr. austr.
floccosa, *Knight, l. c.*—Afr. austr.
Hartogii, *R. Br. in Trans. Linn. Soc.* x. (1810) 108.—Afr. austr.
Hibbertii, R. Br. l. c. = splendida.
hirta, *Knight, Prot.* 66; *R. Br. l. c.* 105.—Afr. austr.
homomalla, Reichb. f. ex Meissn. in DC. Prod. xiv. 266 = purpurea.
intermedia, *Buek, ex Meissn. l. c.* 265.—Afr. austr.
Ludwigii, Steud. ex Meissn. l. c. 263 = lyrigera.
lyrigera, *Knight, Prot.* 65.—Afr. austr.
Massoni, *R. Br. in Trans. Linn. Soc.* x. (1810) 109.—Afr. austr.
myrtifolia, *R. Br. l. c.* 110.—Afr. austr.
nitens, *Roem. & Schult. Syst.* iii. 384.—Afr. austr.
palustris, *Knight, Prot.* 66.—Afr. austr.
parviflora, Klotzsch, ex Krauss, in Flora, xxviii. (1845) 77 = divaricata.
pauciflora, *R. Br. in Trans. Linn. Soc.* x. (1810) 106.—Afr. austr.
purpurea, *R. Br. l. c.* 111.—Afr. austr.
spectabilis, Steud. Nom. ed. I. 532 = Protea latifolia.
splendida, *Knight, Prot.* 66.—Afr. austr.
thymelaeoides, *R. Br. in Trans. Linn. Soc.* x. (1810) 109.—Afr. austr.
vacciniifolia, *Sweet, Hort. Brit.* ed. I. 347.—Afr. austr.
Zeyheri, *Meissn. in DC. Prod.* xiv. 264.—Afr. austr.

MIMOSA, Linn. Gen. ed. I. 155 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 593.
EBURNAX, Rafin. New Fl. Am. i. 42 (1836).
LOMOPLIS, Rafin. Sylva Tellur. 118 (1838).
SENSITIVA, Rafin. l. c. 119 (1838).
abietina, Poir. Encyc. Suppl. i. 60 = Acacia linifolia.
abstergens, Roxb. ex Spreng. Syst. ii. 207 = Acacia concinna, *DC.*
acacioides, Benth. in Hook. Journ. Bot. ii. (1840) 132 = Piptadenia peregrina.
acanthocarpa, *Poir. Encyc. Suppl.* v. 529; *Benth. l. c.* iv. (1842) 409.—Mexic.
acantholoba, *Poir. l. c.* i. 83.—Am. trop.
acerba, *Benth. in Hook. Journ. Bot.* iv. (1842) 378.—Bras.
acicularis, Poir. Encyc. Suppl. v. 529 = Acacia Farnesiana.

MIMOSA:—
Acle, Blanco, Fl. Filip. ed. I. 738 = Xylia dolabriformis.
aculeata, *Mill. Gard. Dict.* ed. VIII. add.—Hab.?
aculeaticarpa, Orteg. Hort. Matr. Dec. 134 = acanthocarpa.
acutiflora, Benth. in Hook. Journ. Bot. iv. (1842) 397 = somnians.
acutistipula, *Benth. l. c.* 391.—Bras.
adenanthera, Hort. ex Steud. Nom. ed. II. i. 3 = Acacia adenanthera (in Addend.).
adenanthera, Roxb. Hort. Beng. 40; Fl. Ind. ii. 554 = Neptunia plena.
adenantheroides, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 88.—Am. trop.
adenocarpa, *Benth. in Hook. Journ. Bot.* iv. (1842) 399.—Bras.
adenotricha, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 91.—Bras.
adhaerens, *H. B. & K. Nov. Gen. et Sp.* vi. 249.—Am. trop.
adhaerens, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 529 = Acacia paniculata.
adhaerens, Mart. in Flora, xx. (1837) II. Beibl. 122 = Acacia adhaerens.
adianthifolia, Schum. & Thonn. Beskr. Pl. Guin. 322 = Albizzia fastigiata.
adpressa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 202.—Am. austr.
adstringens, Schum. & Thonn. Beskr. Guin. Pl. 327 = Acacia arabica.
adversa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 439; *et in Mart. Fl. Bras.* xv. II. 384.—Am. trop.
aemula, *Spreng. Syst.* ii. 206.—Bras.
Aeschynomenes, Benth. Bot. Voy. Sulph. 89 = camporum.
aggregata, *Pers. Syn.* ii. 263.—Ind. or.
agrestis, Sieber, ex Spreng. Syst. ii. 206 = Prosopis Stephaniana.
alata, Poir. Encyc. Suppl. v. 530 = Acacia alata.
alba, Roxb. ex Rottl. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 208 = Acacia leucophloea.
alba, Sw. Prod. Veg. Ind. Occ. 85 = Inga alba.
alba, Vahl, Eclog. Am. iii. 31 = Inga Bourgoni.
albida, *Humb. & Bonpl. in Willd. Sp. Pl.* iv. 1030.—Am. trop.
albida, Roxb. ex Wall. Cat. sub n. 5227 = Acacia Suma.
Algarrobo, Azara, Voy. ii. 483 = Prosopis juliflora.
alternans, Vahl, ex Benth. in Trans. Linn. Soc. xxx. (1875) 548 = Calliandra alternans.
altissima, Roxb. Hort. Beng. 41 = Pithecolobium altissimum?
amara, Roxb. Pl. Corom. ii. 13. t. 122 = Albizzia amara.
ambigua, Salisb. Prod. 325 = Acacia suaveolens.
andina, *Benth. in Trans. Linn. Soc.* xxx. (1875) 427.—Reg. Andin.
angulata, Poir. Encyc. Suppl. v. 530 = Acacia discolor.
angulata, *Spreng. Neue Entdeck.* ii. 158.—Bras.
angusta, *Benth. in Hook. Journ. Bot.* iv. (1842) 362.—Bras.
angustifolia, Jacq. Hort. Schoenb. iii. 74. t. 39 = Acacia suaveolens.
angustisiliqua, Lam. Encyc. i. 10 = Desmanthus virgatus.
angustissima, Mill. Gard. Dict. ed. VIII. n. 19 = Acacia filicina.
annularis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 419; *et in Mart. Fl. Bras.* xv. II. 362.—Bras.
Antillarum, Poir. Encyc. Suppl. i. 80 = Pithecolobium parviflorum.
antrorsa, Benth. in Hook. Journ. Bot. iv. (1842) 403 = adversa.
apodocarpa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 415; *et in Mart. Fl. Bras.* xv. II. 359.—Am. trop.
appressa, Benth. in Mart. Fl. Bras. xv. II. 347 = adpressa.
aquatica, Pers. Syn. ii. 263 = Neptunia oleracea.
arabica, Lam. Encyc. i. 19 = Acacia arabica.
arborea, Forsk. Fl. Aegypt. Arab. 177; Thunb. Fl. Jap. 229 = Albizzia Julibrissin.
arborea, Linn. Sp. Pl. 519 = Acacia villosa, Pithecolobium filicifolium.
arcuata, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 310.—Mexic.
arenosa, *Poir. Encyc. Suppl.* i. 66.—Am. trop.
armata, Poir. l. c. v. 530 = Acacia armata.
armata, Rottl. ex Spreng. Syst. ii. 206 = hamata.
arvensis, Sieber, ex Steud. Nom. ed. I. 533 = Prosopis Stephaniana?

MIMOSA:—
Asak, Forsk. Fl. Aegypt. Arab. 176 = Acacia hamulosa?
asperata, Blanco, Fl. Filip. ed. I. 732 = pudica.
asperata, *Linn. Syst.* ed. X. 1312.—Ind. occ.
asperrima, *Benth. in Hook. Journ. Bot.* iv. (1842) 377.—Am. trop.
atakta, Steud. in Flora, xxvi. (1843) 758 = Pithecolobium trapezifolium.
atomaria, Poir. Encyc. Suppl. i. 81 = Acacia macracantha.
auriculata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 430.—Am. trop.
aurivillus, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 52.—Bras.
axillaris, *Benth. in Hook. Journ. Bot.* iv. (1842) 382.—Bras.
Azak, G. Don, Gen. Syst. ii. 412 = Acacia hamulosa!
bahamensis, *Benth. in Hook. Journ. Bot.* iv. (1842) 408.—Am. trop.
Balansae, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 52.—Paraguay.
balsamica, Molina, Sagg. Chile, 165 = Larrea nitida.
Barbadetimam, Vell. Fl. Flum. xi. t. 7 = Stryphnodendron Barbatimam.
barbigera, *Benth. in Hook. Journ. Bot.* iv. (1842) 371.—Bras.
Barclayana, *Sweet, Hort. Brit.* ed. I. 100, nomen.—Madag.
bauhiniaefolia, *Karst. Fl. Columb.* ii. 65. t. 133.—N. Granat.
bauhiniaefolia, *Salisb. Prod.* 324.—Hab.?
Bellatrix, Hoffmgg. Verz. Nachtr. II. 162 = asperata.
Berlandieri, *A. Gray, ex Torr. in Bot. Mex. Bound.* 61.—Texas.
biceps, Poir. Encyc. Suppl. i. 75 = Leucaena glauca.
bicolor, Bacle, ex DC. Prod. ii. 445; Schum. & Thonn. Beskr. Pl. Guin. 326 = Dichrostachys nutans.
bicolor, Hilsenb. & Boj. ex Benth. in Hook. Journ. Bot. iv. (1842) 353 = Dichrostachys tenuifolia.
biflora, Poir. Encyc. Suppl. v. 530 = Acacia biflora.
bifurca, *Benth. in Hook. Journ. Bot.* iv. (1842) 407.—Am. trop.
bigemina, Linn. Sp. Pl. 517 = Pithecolobium bigeminum.
biglobosa, Jacq. Select. Am. 267 = Parkia africana.
biglobosa, Roxb. Hort. Beng. 40; Fl. Ind. ii. 551 = Parkia Roxburghii.
bijuga, *Benth. in Hook. Journ. Bot.* iv. (1842) 398.—Bras.
bijuga, Vell. Fl. Flum. xi. t. 32 = Acacia Miersii.
binervia, Wendl. Bot. Beobacht. 56 = Acacia glaucescens.
bipinnata, Aubl. Pl. Gui. ii. 946 = Entada polystachya.
biuncifera, *Benth. Pl. Hartw.* 12.—Mexic.
Blanchetii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 433; *et in Mart. Fl. Bras.* xv. II. 372.—Am. trop.
Blancoana, *Llanos, in Mem. Acad. Cienc. Mad.* ii. (1859) 503 (Quid?).—Ins. Philipp.
boliviana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 396.—Bolivia.
Bonplandii, *Benth. in Hook. Journ. Bot.* iv. (1842) 387.—Reg. Argent.
borealis, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 39.—Texas.
botrycephala, Vent. Jard. Cels, 1 = Acacia discolor.
Bourgoni, Aubl. Pl. Gui. ii. 941 = Inga Bourgoni.
brachyacantha, Poir. Encyc. Suppl. v. 529 = acanthocarpa.
brachycarpa, *Benth. in Hook. Journ. Bot.* iv. (1842) 410.—Bras.
brachyloba, Muhl. ex Steud. Nom. ed. II. ii. 147 = Desmanthus brachylobus.
brachystachys, DC. Prod. ii. 437, in syn. = Pithecolobium macrostachyum.
bracteolaris, *Benth. in Hook. Journ. Bot.* iv. (1842) 408.—Bras.
brasiliensis, *Spreng. Syst.* ii. 207.—Bras.
brevifolia, *Humb. ex Spreng. l. c.* 205.—Am. austr.
brevipes, *Benth. in Hook. Journ. Bot.* iv. (1842) 381.—Bras.
brevipinna, *Benth. in Trans. Linn. Soc.* xxx. (1875) 432; *et in Mart. Fl. Bras.* xv. II. 371.—Bras.
brevissima, Benth. in Mart. l. c. 518 (Ind.) sphalm. = praec.
Brownei, Poir. Encyc. Suppl. v. 530 = Acacia juniperina.
Burchellii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 432; *et in Mart. Fl. Bras.* xv. II. 370.—Am. trop.

MIMOSA:—
Cabrera, *Karst. Fl. Columb.* ii. 63. t. 132.—N. Granat.
caduca, *Poir. Encyc. Suppl.* i. 83.—Am. trop.
caesalpiniaefolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 392.—Bras.
caesia, Burm. f. Fl. Ind. 224 = Acacia Burmanniana.
caesia, Linn. Sp. Pl. 522 = Acacia Intsia.
caesia, Russ. ex Wall. Cat. n. 5264 C = Acacia planifrons.
caffra, Thunb. Prod. Pl. Cap. 92 = Acacia caffra.
calcarea, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 453 = Prosopis cinerascens.
calistachya, Presl, Bot. Bemerk. 65 = millefoliata.
callosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 376.—Am. trop.
calocephala, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 53.—Bras.
Calodendron, *Mart. Herb. Fl. Bras.* n. 1087; *Benth. in Hook. Journ. Bot.* iv. (1842) 389.—Bras.
Calothamnus, *Mart. l. c.* n. 1089; *Benth. l. c.* 388.—Bras.
calycina, *Benth. l. c.* 377.—Bras.
cameleopardalis, *Page, Prod. Southamp. Bot. Gard.* 207, nomen.—Afr. austr.
campeachiana, Mill. Gard. Dict. ed. VIII. n. 20 = Acacia sphaerocephala.
camporum, *Benth. in Hook. Journ. Bot.* ii. (1840) 130.—Guian.
canescens, Willd. Sp. Pl. iv. 1038 = asperata.
capensis, Burm. f. Fl. Cap. Prod. 31 (err. typ. 27) = Acacia horrida?
capillaris, *Dum.-Cours. Bot. Cult.* ed. II. vii. 296.—Hab.?
capillipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 432; *et in Mart. Fl. Bras.* xv. II. 371.—Bras.
capitata, Poir. Encyc. Suppl. v. 528 = Inga capitata.
caracasana, Jacq. Coll. iv. 142 = Calliandra caracasana.
carinata, *Griseb. in Goett. Abh.* xxiv. (1879) 120.—Reg. Argent.
caripensis, Poir. Encyc. Suppl. i. 39 = Calliandra tergemina.
Carisquis, *Blanco, Fl. Filip.* ed. I. 734.—Ins. Philipp.
Cascabelillo, *Colla, Hort. Ripul.* 91.—Hab.?
cassioides, Poir. Encyc. Suppl. v. 529 = Acacia cassioides.
casta, *Linn. Sp. Pl.* 518.—Am. trop.
Catechu, Linn. f. Suppl. 439 = Acacia Catechu.
catechuoides, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 562 = Acacia Catechu.
caudata, Vahl, Eclog. Am. iii. 35 = Entada polystachya.
cauliflora, Poir. Encyc. Suppl. i. 45 = Pithecolobium cauliflorum.
Cavalum, *Roxb. Hort. Beng.* 41 (Quid?).—Ind. or.
Cavaty-tooma, Roxb. ex DC. Prod. ii. 462 = Acacia tomentosa.
Cavenia, Molina, Sagg. Chile, 354 = Acacia Farnesiana.
Ceratonia, *Linn. Sp. Pl.* 523.—Am. trop.
chiliantha, G. F. W. Mey. Prim. Fl. Esseq. 163 = Entada polystachya.
chionacantha, Rafin. Fl. Ludov. 99 (Quid?).—Am. bor.
chrysantha, Vahl, Eclog. Am. iii. 38 = Acacia chrysantha.
chrysastra, *Mart. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 410.—Am. trop.
chrysostachys, Hilsenb. & Boj. ex Benth. in Hook. Journ. Bot. iv. (1842) 343 = Piptadenia chrysostachys.
Chundra, Roxb. & Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 207 = Acacia Sundra.
ciliata, *Spreng. Syst.* ii. 205.—Am. austr.
ciliata, Willd. Enum. Hort. Berol. 1048 = asperata.
cineraria, Linn. Sp. Pl. 517 = Prosopis spicigera.
cinerea, Linn. l. c. 520 = Dichrostachys cinerea.
cinerea, *Vell. Fl. Flum.* xi. t. 35.—Bras.
circinalis, Benth. in Trans. Linn. Soc. xxx. (1875) 635, in syn. = Inga alternifolia.
circinalis, Cav. Ic. vi. 41, in nota = Prosopis strombulifera.
circinalis, Linn. Sp. Pl. 517 = Pithecolobium circinale.
circinnatus, Sesse & Moç. ex G. Don, Gen. Syst. ii. 319 = Inga alternifolia.
Clausseni, *Benth. in Hook. Journ. Bot.* iv. (1842) 405.—Bras.
cochlearis, Labill. Nov. Holl. Pl. ii. 15. t. 234 = Acacia cochlearis.

MIMOSA:—
cochleata, Poir. Encyc. Suppl. i. 45 = Pithecolobium cochleatum.
cochliacantha, Poir. l. c. 78 = Acacia cochliacantha.
cochliocarpos, Gomez, in Mem. Acad. Lisboa, iii. (1812); Mem. Corresp. 34 = Pithecolobium Auaremotemo.
colubrina, Vell. Fl. Flum. xi. t. 16 = Piptadenia colubrina.
comosa, Sw. Prod. Veg. Ind. Occ. 85 = Calliandra comosa.
concinna, *Benth. in Trans. Linn. Soc.* xxx. (1875) 404; *et in Mart. Fl. Bras.* xv. II. 337.—Bras.
concinna, Willd. Sp. Pl. iv. 1039 = Acacia concinna, *DC.*
concordiana, Roxb. Fl. Ind. ii. 556 = Pithecolobium umbellatum.
conferta, *Benth. in Trans. Linn. Soc.* xxx. (1875) 402; *et in Mart. Fl. Bras.* xv. II. 331.—Bras.
contortisiliqua, Vell. Fl. Flum. xi. t. 25 = Enterolobium Timboüva.
contortuplicata, Zuccagni, in Roem. Collect. 160 = Desmanthus brachylobus.
coorgensis, Heyne, ex Wall. Cat. n. 5277 = Xylia dolabriformis.
corcondiana, Roxb. Hort. Beng. 40 = Pithecolobium umbellatum.
cordistipula, *Benth. in Hook. Journ. Bot.* iv. (1842) 411.—Bras.
coriacea, Pers. Syn. ii. 262 = Inga sertulifera.
coriacea, Poir. Encyc. Suppl. i. 39 = Calliandra emarginata.
coriacea, Sieber, ex Presl, Symb. Bot. i. 65 = Inga martinicensis.
coriaria, Blanco, Fl. Filip. ed. I. 734 = Albizzia procera.
corniculata, Lour. Fl. Cochinch. ii. 651 = Albizzia Millettii, Sophora japonica.
cornigera, Linn. Sp. Pl. 520 = Acacia sphaerocephala, latronum.
cornigera, Linn. f. Suppl. 438 = Acacia latronum.
cornuta, J. F. Gmel. Syst. 504 (Quid?).—Afr. trop.
coronillaefolia, Pers. Syn. ii. 263 = Acacia coronillaefolia.
coruscans, Poir. Encyc. Suppl. i. 43 = Inga coruscans.
coruscans, Sieber, ex Presl, Bot. Bemerk. 66 = Inga laurina.
corymbosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = Pithecolobium corymbosum.
costaricensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 423.—Guatemala.
crinita, Pers. Syn. ii. 261 = pilosa.
crocea, Vell. Fl. Flum. xi. t. 17 = Swartzia crocea.
cruenta, *Benth. in Hook. Journ. Bot.* iv. (1842) 407.—Bras.
cumana, Poir. Encyc. Suppl. i. 65 = Prosopis juliflora.
cyclocarpa, Jacq. Fragm. 30 = Enterolobium cyclocarpum.
cylindracea, *Benth. in Hook. Journ. Bot.* iv. (1842) 389.—Bras.
cylindrica, Vell. Fl. Flum. xi. t. 9 = Inga cylindrica.
Dalea, Poir. Encyc. Suppl. v. 530 = Dichrostachys cinerea.
daleoides, *Benth. in Hook. Journ. Bot.* iv. (1842) 389.—Bras.
dasyphylla, *Baker, in Journ. Linn. Soc.* xxi. (1884) 338.—Madag.
dealbata, Page, Prod. Southamp. Bot. Gard. 148 = Acacia dealbata.
debilis, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1029.—Am. austr.
decipiens, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 366 = Acacia decipiens.
decurrens, *Boj. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 420.—Afr. trop.
decurrens, Boj. Hort. Maurit. 113 = Barclayana.
decurrens, Wendl. Bot. Beobacht. 57 = Acacia decurrens.
deltoides, *Nois. ex Steud. Nom.* ed. II. ii. 148 (Quid?).—Hab.?
densa, *Benth. in Hook. Journ. Bot.* iv. (1842) 405.—Bras.
depauperata, *Benth. Pl. Hartw.* 13.—Mexic.
depressa, Poir. Encyc. Suppl. i. 58 = Desmanthus depressus.
desmodioides, *Benth. in Trans. Linn. Soc.* xxx. (1875) 392; *et in Mart. Fl. Bras.* xv. II. 309.—Bras.
detinens, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 89.—Reg. Argent.

MIMOSA :—
Diadema, Vell. Fl. Flum. xi. t. 45 = Inga Diadema.
dichotoma, Vell. l. c. t. 42 = Inga dichotoma.
didyma, *Vell. l. c.* t. 46.—Bras.
diffusa, *Benth. in Hook. Journ. Bot.* iv. (1842) 373.—Bras.
digitata, *Benth. l. c.* 368.—Bras.
dimidiata, *Benth. l. c.* 378.—Bras.
diplacantha, *Benth. in Trans. Linn. Soc.* xxx. (1875) 424.—Ind. occ.
diplotricha, Wright, in Sauv. Fl. Cub. 34 = invisa.
diptera, Poir. Encyc. Suppl. v. 529 = Prosopis juliflora.
discolor, Andr. Bot. Rep. t. 235 = Acacia discolor.
discolor, *Benth. in Hook. Journ. Bot.* iv. (1842) 391.—Bras.
discolor, Poir. Encyc. Suppl. i. 46 = Pithecolobium adiantifolium.
dispersa, Benth. in Hook. Journ. Bot. iv. (1842) 369 = hirsuta.
distachya, *Cav. Ic.* iii. 48. t. 295.—Mexic.
distachya, Vent. Jard. Cels, 20 = Albizzia lophantha.
distans, *Benth. in Hook. Journ. Bot.* iv. (1842) 382.—Bras.
divaricata, Jacq. Hort. Schoenb. iii. 76 = Lysiloma Schiedeana.
divergens, Poir. Encyc. Suppl. i. 59 = Dichrostachys nutans.
diversifolia, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 57.—Paraguay.
Djiringa, Roxb. Hort. Beng. [93] = Pithecolobium lobatum.
dodonaeifolia, Pers. Syn. ii. 261 = Acacia dodonaeifolia.
dolens, *Vell. Fl. Flum.* xi. t. 34.—Bras.
domingensis, *Benth. in Hook. Journ. Bot.* iv. (1842) 409.—Ins. St. Doming.
dominiciana, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 424 = casta.
dormiens, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1035.—Am. austr.
dulcis, Roxb. Pl. Corom. i. 67. t. 99 = Pithecolobium dulce.
dulcis, Vell. Fl. Flum. xi. t. 4 = Inga affinis.
dumetorum, A. St. Hill. in Mém. Mus. Par. ix. (1822) 316 = Acacia dumetorum.
dumosa, Roxb. Hort. Beng. 40; Fl. Ind. ii. 559 = Acacia modesta.
dysocarpa, *Benth. in A. Gray, Pl. Wright.* i. 62.—Am. bor. occ.
eburnea, Hort. Par. ex Benth. in Trans. Linn. Soc. xxx. (1875) 507 = Acacia horrida.
eburnea, Linn. f. Suppl. 437 = Acacia eburnea.
eburnea, Roxb. Pl. Corom. ii. 54. t. 199 = Acacia planifrons.
echinocarpa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 398; *et in Mart. Fl. Bras.* xv. II. 320.—Bras.
echinocaula, *Benth. ll. cc.* 421; 365.—Bras.
edulis, Poir. Encyc. Suppl. v. 529 = Acacia Farnesiana.
elata, Roxb. Hort. Beng. 40; Fl. Ind. ii. 546 = Albizzia procera.
elegans, Andr. Bot. Rep. t. 563 = Albizzia lophantha.
elliptica, *Benth. in Hook. Journ. Bot.* iv. (1842) 400.—Bras.
elongata, *Benth. l. c.* 369.—Bras.
emarginata, Poir. Encyc. Suppl. i. 39 = Calliandra emarginata.
emirnensis, *Benth. in Hook. Journ. Bot.* iv. (1842) 396.—Madag.
Emoryana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 426.—Am. bor. occ.
Endymionis, Mart. in Flora, xxi. (1838) II. Beibl. 51 = pudica.
ensifolia, Sm. ex Pers. Syn. ii. 261, in syn. = Acacia longifolia.
Entada, Linn. Sp. Pl. 518 = Entada polystachya.
ephedroides, *Benth. in Hook. Journ. Bot.* iv. (1842) 413.—Chili.
eriantha, Poir. Encyc. Suppl. v. 530 = Acacia eriantha.
erinacea, Benth. in Hook. Journ. Bot. iv. (1842) 383 = pogocephala.
eriocarpa, *Benth. l. c.* 387.—Bras.
eriocaulis, *Benth. l. c.* 383.—Bras.
eriophylla, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 87.—Bras.
Ervendbergii, *A. Gray, in Proc. Am. Acad.* v. (1862) 178.—Mexic.
esculenta, Pav. ex Benth. in Trans. Linn. Soc. xxx. (1875) 443 = Leucaena esculenta.

MIMOSA :—
expansa, Pav. ex Benth. in Trans. Linn. Soc. xxx. (1875) 549 = Calliandra expansa.
extensa, *Benth. in Hook. Journ. Bot.* iv. (1842) 393.—Bras.
extranea, *Benth. in Trans. Linn. Soc.* xxx. (1875) 433.—Ins. S. Doming.
fagaracantha, *Griseb. Cat. Pl. Cub.* 81.—Cuba.
fagifolia, Jacq. Select. Am. 264 = Inga laurina.
fagifolia, Linn. Sp. Pl. 516 = Inga fagifolia.
falcata, Dum.-Cours. Bot. Cult. ed. I. 420; Desf. Tabl. ed. I. 180 = Acacia longifolia.
falcata, Poir. Encyc. Suppl. i. 61 = Acacia falcata.
falcipinna, *Benth. in Hook. Journ. Bot.* iv. (1842) 371.—Bras.
farcta, [Soland. in] Russ. Aleppo, ed. II. ii. 266 = Prosopis Stephaniana.
farinosa, *Griseb. in Goett. Abh.* xix. (1874) 134.—Reg. Argent.
Farnesiana, Linn. Sp. Pl. 521 = Acacia Farnesiana.
fasciculata, Benth. in Hook. Journ. Bot. iv. (1842) 383 = microcarpa.
fasciculata, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 88.—Bras.
fasciculata, Poir. Encyc. Suppl. i. 46 = Calliandra surinamensis.
fastuosa, Jacq. Fragm. 15. t. 10 = Inga fastuosa.
fera, Lour. Fl. Cochinch. 652 (= Gleditschiae sp.).—China.
ferruginea, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 208 = Acacia pennata.
ferruginea, Roxb. Hort. Beng. 41; Fl. Ind. ii. 561 = Acacia ferruginea.
fervida, Mart. in Flora, xxi. (1838) II. Beibl. 57 = obtusifolia.
filicifolia, Lam. Encyc. i. 13 = Pithecolobium filicifolium.
filicina, Poir. Encyc. Suppl. i. 47 = Parkia filicina.
filicoides, Cav. Ic. i. 55 = Acacia filicina.
filiformis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 432; *et in Mart. Fl. Bras.* xv. II. 371.—Bras.
filipes, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 52.—Bras.
flagellaris, *Benth. in Hook. Journ. Bot.* iv. (1842) 372.—Bras.
flagelliformis, Vell. Fl. Flum. xi. t. 27 = Inga flagelliformis.
flava, *Forsk. Fl. Aegypt. Arab.* 176.—Arab.
flavescens, Splitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 110 = camporum.
flaviseta, Benth. in Hook. Lond. Journ. Bot. v. (1846) 90 = camporum.
flexuosa, *Benth. in A. Gray, Pl. Wright.* i. 62.—Am. bor. occ.
flexuosa, Poir. Encyc. Suppl. i. 79 = Acacia macracantha.
flexuosa, Rottl. ex Wight & Arn. Prod. 275 = Albizzia latifolia.
floribunda, Benth. in Hook. Journ. Bot. ii. (1840) 230 = sensitiva.
floribunda, Vent. Choix, 13 = Acacia longifolia.
floribunda, *Willd. Sp. Pl.* iv. 1031.—Am. trop.
fluminensis, Vell. Fl. Flum. xi. t. 38 = Acacia plumosa.
foetida, Jacq. Hort. Schoenb. iii. 73 = Piptadenia foetida.
foliolosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 406.—Bras.
fragrans, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 182.—Texas.
franciscana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 431; *et in Mart. Fl. Bras.* xv. II. 369.—Bras.
fraxinea, Poir. Encyc. Suppl. i. 44 = Inga alba.
frondosa, Heyne, ex Wall. Cat. n. 5227 = Acacia Suma.
frondosa, Klein, ex Poir. Encyc. Suppl. i. 76 = Leucaena glauca.
frondosa, Koen. ex Wall. Cat. n. 5265 C = Albizzia Lebbek.
fruticosa, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 549.—China.
fruticosa, Vell. Fl. Flum. xi. t. 6 = Piptadenia laxa.
furcata, Desf. Tabl. ed. I. 180; Pers. Syn. ii. 263 = Prosopis juliflora.
furfuracea, *Benth. in Hook. Journ. Bot.* iv. (1842) 389.—Bras.
Galeottii, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 92.—Mexic.
Gardneri, *Benth. in Hook. Journ. Bot.* iv. (1842) 405.—Bras.

MIMOSA :—
geminata, Koen. ex Wight & Arn. Prod. 269 = Pithecolobium geminatum.
geminata, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 427.—Mexic.
Gilliesii, *Benth. in Hook. Journ. Bot.* iv. (1842) 413.—Am. austr.
giraffae, Poir. Encyc. Suppl. v. 529 = Acacia giraffae.
glaberrima, Schum. & Thonn. Beskr. Guin. Pl. 321 = Albizzia glaberrima.
glabra, *Benth. in Hook. Journ. Bot.* iv. (1842) 386.—Bras.
glandulosa, Forst. f. Prod. 92 = Leucaena Forsteri.
glandulosa, Michx. Fl. Bor. Am. ii. 254 = Desmanthus brachylobus.
glandulosa, *C. Sm. in Tuckey, Congo*, 249, nomen.—Ins. Cap. Virid.
glandulosa, Vahl, Eclog. Am. iii. 38 = Inga glandulosa.
glauca, Koen. ex Roxb. Hort. Beng. 41 = Albizzia amara.
glauca, Linn. Sp. Pl. 520 = Leucaena glauca.
glaucescens, *Benth. in Hook. Journ. Bot.* iv. (1842) 362.—Bras.
glaucescens, Poir. Encyc. Suppl. i. 61 = Acacia glaucescens.
Glazioui, *Benth. in Trans. Linn. Soc.* xxx. (1875) 397.—Bras.
Glaziovii, Benth. in Mart. Fl. Bras. xv. II. 316 = praec.
Glockeri, Meissn. ex Benth. in Trans. Linn. Soc. xxx. (1875) 397 = polydactyla.
glomerata, *Forsk. Fl. Aegypt. Arab.* 177.—Arab.
glomerata, Vell. Fl. Flum. xi. t. 43 = Pithecolobium glomeratum.
gonoclada, *Benth. in Trans. Linn. Soc.* xxx. (1875) 427.—Bolivia.
goyazensis, *Benth. in Hook. Journ. Bot.* iv. (1842) 380.—Bras.
gracilis, *Benth. l. c.* 411.—Bras.
Grahami, *A. Gray, Pl. Wright.* ii. 52.—N. Mexic.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 357.—Madag.
grandiflora, Herb. Madr. ex Wight & Arn. Prod. 279 = Parkia biglandulosa.
grandiflora, L'Hérit. Sert. Angl. 30 = Calliandra grandiflora.
grandiflora, Soland. ex Benth. Fl. Austral. ii. 424 = Pithecolobium grandiflorum.
grandisiliqua, Vell. Fl. Flum. xi. t. 37 = Acacia paniculata.
grata, Poir. Encyc. Suppl. v. 529 = Piptadenia macrocarpa.
guadalupensis, Pers. Syn. ii. 262 = Pithecolobium Unguis-cati.
guatemalensis, *Benth. Bot. Voy. Sulph.* 89.—Guatemal.
guayaquilensis, *Steud. Nom.* ed. I. 533; ed. II. ii. 148.—Ecuador.
guianensis, Aubl. Pl. Gui. ii. 938 = Stryphnodendron guianense.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 323 = Calliandra portoricensis.
gummifera, Brouss. ex Poir. Encyc. Suppl. i. 64 = Acacia gummifera.
gummifera, Forsk. Fl. Aegypt. Arab. p. cxxiv = Acacia spirocarpa.
gymnoloma, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 85.—Bras.
Habbas, Delile, Fl. Aeg. Illustr. 79 = asperata.
Haematoxylon, Poir. Encyc. Suppl. v. 529 = Acacia Haematoxylon.
hamata, *Willd. Sp. Pl.* iv. 1033.—Ind. or.
hebecarpa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 87.—Bras.
hebeclada, Poir. Encyc. Suppl. v. 529 = Acacia hebeclada.
Helvilleana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 357.—Madag.
herbacea, Roxb. Hort. Beng. 41 (Quid ?).—Am.
heterocarpa, *Presl, Bot. Bemerk.* 64.—Bras.
heterophylla, Lam. Encyc. i. 14 = Acacia heterophylla.
heterophylla, Roxb. Hort. Beng. 40; Fl. Ind. ii. 545 = Pithecolobium angulatum.
hexaphylla, Salzm. ex Benth. in Mart. Fl. Bras. xv. II. 317 = polydactyla.
hirsuta, Moç. & Sesse, ex G. Don, Gen. Syst. ii. 383 = pudica.
hirsuta, *Spreng. Syst.* ii. 204.—Reg. Argent.
hirsutissima, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 55.—Bras.
hirta, *Vell. Fl. Flum.* xi. t. 31.—Bras.
hispida, Willd. Sp. Pl. iv. 1037 = asperata.

MIMOSA:—
hispidula, H. B. & K. Nov. Gen. et Sp. vi. 252 = pudica.
hispidula, Sm. Bot. N. Holl. 59. t. 16 = Acacia hispidula.
honesta, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 57.—Bras.
horrida, Linn. Sp. Pl. 521 = Acacia horrida.
horrida, Russ. ex Wall. Cat. n. 5261 E = Acacia leucophloea.
horrida, Sm. in Rees, Cyclop. xxiii. in obs. = Acacia planifrons.
horridula, Michx. Fl. Bor. Am. ii. 254 = Schrankia uncinata.
hostilis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 415; *et in Mart. Fl. Bras.* xv. II. 359.—Bras.
Hostmanni, Benth. in Hook. Lond. Journ. Bot. v. (1846) 84 = debilis.
Houstoni, L'Hérit. Sert. Angl. 30 = seq.
Houstoniana, Mill. Gard. Dict. ed. VIII. n. 16 = Calliandra Houstoni.
humifusa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 406; *et in Mart. Fl. Bras.* xv. II. 342.—Bras.
humilis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1037 = dormiens.
hymenaeodes, Rich. ex Pers. Syn. ii. 262 = Calliandra hymenaeoides.
hymeneaefolia, Poir. Encyc. Suppl. i. 38 = Pithecolobium hymeneaefolium.
hypoglauca, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 53.—Bras.
ignava, H. B. & K. Nov. Gen. et Sp. vi. 259 = Lysiloma rostrata.
illinoënsis, Michx. Fl. Bor. Am. ii. 254 = Desmanthus brachylobus.
imbricata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1842) 378.—Bras.
implexa, *Benth. l. c.* 411.—Bras.
inaequalis, Poir. Encyc. Suppl. i. 44 = Pithecolobium inaequale.
inamoena, *Benth. in Trans. Linn. Soc.* xxx. (1875) 424; *et in Mart. Fl. Bras.* xv. II. 364.—Bras.
incana, *Benth. in Hook. Journ. Bot.* iv. (1842) 387.—Bras.
incurialis, Vell. Fl. Flum. xi. t. 24 = Pithecolobium incuriale.
indica, Pers. Syn. ii. 266 = Acacia Farnesiana.
Inga, Linn. Sp. Pl. 516 = Inga vera.
Inga, Vell. Fl. Flum. xi. t. 3 (Ynga) = Inga edulis.
ingoides, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = Inga ingoides.
insidiosa, Benth. in Hook. Journ. Bot. iv. (1842) 365 = trachycephala.
insidiosa, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 54.—Bras.
intermedia, Kunth, Mimos. 16. t. 6 = dormiens.
interrupta, *Benth. in Hook. Journ. Bot.* iv. (1842) 392.—Bras.
intricata, *Benth. l. c.* 407.—Bras.
Intsia, Linn. Sp. Pl. 522 = Acacia Intsia.
Intsia, Walt. Fl. Carol. 252 = Schrankia uncinata.
Intsia, Wight, ex Wall. Cat. n. 5250 A = Acacia pinnata?
invisa, *Mart. in Flora*, xx. (1837) II. *Beibl.* 121.—Bras.
involucrata, *Benth. in Hook. Journ. Bot.* iv. (1842) 386.—Bras.
irritabilis, *Dum.-Cours. Bot. Cult.* ed. II. vi. 15.—Hab.?
irritabilis, Presl, Bot. Bemerk. 64 = pudica.
jiramenensis, Karst. Fl. Columb. ii. 59. t. 130 = Velloziana.
Jiringa, Jack, in Malay. Misc. i. (1820) I. n. 14 = Pithecolobium lobatum.
juglandifolia, Poir. Encyc. Suppl. i. 44 = Inga juglandifolia.
Julibrissin, Scop. Delic. Insub. i. 18 = Acacia Julibrissin.
juliflora, Sw. Prod. Veg. Ind. Occ. 85 = Prosopis juliflora.
juniperina, Vent. Jard. Malm. 64 = Acacia juniperina.
Jupunba, Poir. Encyc. Suppl. i. 70 = Pithecolobium trapezifolium.
Kaeringa, Baker, in Hook. f. Fl. Brit. Ind. ii. 305, sphalm. = *M. Koeringa*.
Kalkora, Roxb. Hort. Beng. 40; Fl. Ind. ii. 547 = Albizzia Julibrissin.
Kantuffa, Poir. Encyc. Suppl. v. 529; DC. Prod. ii. 431 = Pterolobium lacerans.
kermesina, A. Dietr. in Otto & Dietr. Allg. Gartenz. iii. (1835) 209 = lucidula.

MIMOSA:—
Kleinii, Poir. Encyc. Suppl. i. 82 = Acacia tomentosa.
Koeringa, Roxb. Hort. Beng. 40; Fl. Ind. ii. 543 = Pithecolobium lobatum.
lactiflua, *Delile, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 393.—Am. trop.
lacustris, Humb. & Bonpl. Pl. Aequin. i. 55. t. 16 = Neptunia oleracea.
laevigata, *Benth. in Linnaea*, xxii. (1849) 530.—Bras.
laevigata, Benth. in Trans. Linn. Soc. xxx. (1875) 378, in syn. = *levigata*.
lanata, *Benth. in Hook. Journ. Bot.* iv. (1842) 379.—Bras.
lanceolata, *Jacq. Fragm.* 44.—Hab.?
lanceolata, Poir. Encyc. Suppl. i. 37 = Pithecolobium ligustrinum.
lasiocarpa, *Benth. in Hook. Journ. Bot.* iv. (1842) 399.—Bras.
lasiocephala, *Benth. l. c.* 372.—Bras.
lasiophylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 414; *et in Mart. Fl. Bras.* xv. II. 357.—Bras.
latifolia, Linn. Syst. ed. X. 1312 = Pithecolobium latifolium.
latisiliqua, Linn. Sp Pl. 519 = Lysiloma latisiliqua.
latispinosa, *Lam. Encyc.* i. 22.—Madag.
latronum, Linn. f. Suppl. 438 = Acacia latronum.
laurina, Sw. Prod. Veg. Ind. Occ. 85 = Inga laurina.
laxa, Poir. Encyc. Suppl. i. 72 = Calliandra laxa.
laxa, Vell. Fl. Flum. xi. t. 23 (Swartziae sp.).—Bras.
laxiflora, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 93.—Mexic.
Lebbeck, Linn. Sp. Pl. 516 = Albizzia latifolia.
Lebbek, Blanco, Fl. Filip. ed. I. 733 = Albizzia retusa.
Lebbek, Forsk. Fl. Aegypt. Arab. 177 = Albizzia Lebbek?
leiocarpa, Benth. in Hook. Journ. Bot. iv. (1842) 391 = malacocentra.
leiocarpa, *DC. Prod.* ii. 429.—Am. trop.
leiocephala, *Benth. in Hook. Journ. Bot.* iv. (1842) 404.—Bras.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 76.—Arizona.
lenis, Vell. Fl. Flum. xi. t. 29 (Acaciae sp.).—Bras.
lentiscifolia, Pers. Syn. ii. 263 = Acacia lentiscifolia.
leptantha, *Benth. in Hook. Journ. Bot.* iv. (1842) 411.—Bras.
leptocaulis, *Benth. l. c.* 398.—Bras.
leptophylla, Brouss. Elench. Hort. Monsp. (1805) 38, nomen; Cav. ex Lag. Gen. et Sp. Nov. 16 = Inga leptophylla.
leptophylla, Poir. Encyc. Suppl. v. 529 = Acacia Farnesiana.
leptorhachis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 434; *et in Mart. Fl. Bras.* xv. II. 376.—Bras.
leucacantha, Jacq. Hort. Schoenb. iii. 75. t. 393 = Acacia horrida.
leucaenoides, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 89.—Mexic.
leucocephala, Lam. Encyc. i. 12 = Leucaena glauca.
leucophloea, Roxb. Pl. Corom. ii. 27. t. 150 = Acacia leucophloea.
levigata, Poir. Encyc. Suppl. i. 65 = Prosopis juliflora.
ligustrina, Jacq. Fragm. 29 = Pithecolobium ligustrinum.
ligustrina, Vahl, Eclog. Am. iii. 34 = Pithecolobium Vahlianum.
Lindheimeri, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 181.—Texas.
linearis, Wendl. Bot. Beobacht. 56 = Acacia linifolia.
linifolia, Vent. Jard. Cels, 2 = Acacia linifolia.
litigiosa, Mart. in Flora, xxi. (1838) II. Beibl. 58 = sensitiva.
longifolia, Andr. Bot. Rep. t. 207 = Acacia longifolia.
longifolia, Poir. Encyc. Suppl. i. 59 = Pithecolobium Vahlianum.
longipes, *Benth. in Hook. Journ. Bot.* iv. (1842) 370.—Bras.
longipinna, Benth. l. c. 366 = polycarpa.
longisiliqua, Lam. Encyc. i. 21.—Cf. invisa, asperata.
longisiliqua, *Vell. Fl. Flum.* xi. t. 15.—Bras.
longissima, Nois. ex Steud. Nom. ed. I. 533 (Quid?).—Hab.?
lophantha, Pers. Syn. ii. 264 = Albizzia lophantha.
Lorentzii, *Griseb. in Goett. Abh.* xix. (1874) 134.—Reg. Argent.

MIMOSA:—
lucida, Roxb. Hort. Beng. 40; Fl. Ind. ii. 544 = Albizzia lucida.
lucida, Vahl, Eclog. Am. iii. 31. t. 24 = Inga pilosiuscula.
lucidula, *Benth. in Hook. Journ. Bot.* iv. (1842) 407.—Bras.
Lundiana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 395; *et in Mart. Fl. Bras.* xv. II. 313.—Bras.
lupulina, *Benth. in Hook. Journ. Bot.* iv. (1842) 399.—Bras.
lusoria, Vell. Fl. Flum. xi. t. 14 = Pithecolobium lusorium.
lutea, Mill. Gard. Dict. ed. VIII. n. 17 = Acacia macracantha.
lycopodioides, Desf. Tabl. ed. I. 182; Pers. Syn. ii. 263 = Neptunia plena.
macracantha, Poir. Encyc. Suppl. i. 78 = Acacia macracantha.
macrocalyx, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 54.—Paraguay.
macrocephala, *Benth. in Trans. Linn. Soc.* xxx. (1875) 398; *et in Mart. Fl. Bras.* xv. II. 320.—Bras.
macroloba, Poir. Encyc. Suppl. i. 66 = Pentaclethra filamentosa.
macrophylla, Poir. l. c. 42 = Inga macrophylla.
macrostachya, Poir. l. c. 61 = Acacia longifolia.
macrostachys, Vahl, Eclog. Am. iii. 34. t. 26 = Pithecolobium macrostachyum.
madagascariensis, G. Don, Gen. Syst. ii. 387, sphalm. = *mascarenensis*.
malacocentra, *Mart. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 415; *et in Mart. Fl. Bras.* xv. II. 360.—Bras.
malacophylla, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 182.—Texas.
malitiosa, Mart. in Flora, xxi. (1838) II. Beibl. 58 = sensitiva.
mangensis, Jacq. Enum. Pl. Carib. 34 = Pithecolobium parvifolium.
Mangium, Forst. f. Prod. 75 = Acacia laurifolia.
Mansii, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 50.—Bras.
marginata, Lam. Encyc. i. 12 = Albizzia odoratissima.
marginata, *Lindl. Bot. Reg.* (1838) *Misc.* 82.—Bras.
Martiusiana, Steud. Nom. ed. II. ii. 148 = Acacia adhaerens.
mascarenensis, Spreng. Syst. ii. 207 = latispinosa.
mauroceana, Desf. Tabl. ed. I. 181 = Acacia mauroceana.
megalodena, Poir. Encyc. Suppl. v. 530 = Acacia pennata.
melanocarpa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 437; *et in Mart. Fl. Bras.* xv. II. 381.—Bras.
melanoxylum, Poir. Encyc. Suppl. v. 530 = Acacia melanoxylon.
mellifera, Vahl, Symb. Bot. ii. 103 = Acacia mellifera.
membranulacea, Blanco, Fl. Filip. ed. I. 739 = Pterolobium indicum.
meticulosa, Benth. in Hook. Journ. Bot. iv. (1842) 381 = neuroloma.
meticulosa, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 55.—Bras.
miamensis, Roxb. Hort. Beng. 41, nomen (Quid?).—Afr. austr.
micracantha, *Benth. in Hook. Journ. Bot.* ii. (1840) 131.—Bras.
micrantha, Vahl, ex Walp. Rep. v. 582 = Prosopis Stephaniana.
microcantha, Poir. Encyc. Suppl. v. 530 = Pithecolobium parvifolium.
microcarpa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 403; *et in Mart. Fl. Bras.* xv. II. 333.—Bras.
microcephala, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1039.—Am. trop.
microphylla, Poir. Encyc. Suppl. i. 36 = Pithecolobium Unguis-cati.
microphylla, Roxb. Hort. Beng. 40; Fl. Ind. ii. 549 = Albizzia myriophylla.
microphylla, Sm. ex Steud. Nom. ed. I. 533 = Schrankia uncinata.
micropteris, *Benth. in Hook. Journ. Bot.* iv. (1842) 384.—Bras.
millefoliata, *Scheele, in Linnaea*, xvii. (1843) 337.—Bras.
misera, *Benth. in Hook. Journ. Bot.* iv. (1842) 411.—Bras.

MIMOSA :—
modesta, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 55.—Bras.
mollis, *Benth. in Hook. Journ. Bot.* iv. (1842) 408.—Mexic.
mollissima, Hort. ex Willd. Enum. Hort. Berol. 1053 = Acacia decurrens.
mollissima, Poir. Encyc. Suppl. i. 46 = Calliandra mollissima.
monacantha, Poir. l. c. v. 529 = Acacia monacantha.
monadelpha, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 544 = Pithecolobium bigeminum.
monancistra, *Benth. Pl. Hartw.* 12.—Mexic.
monilifera, Bertol. Fl. Guatim. 41 = Pithecolobium Bertolonii.
Monjollo, Vell. Fl. Flum. xi. t. 26 = Enterolobium Mongollo.
montana, Hassk. Cat. Hort. Bog. Alt. 290 = Pithecolobium moniliferum.
montana, *Kunth, Mimos.* 31. t. 10.—Peruv.
mucronulata, *Hort. ex Dum.-Cours. Bot. Cult.* ed. II. vii. 296.—Hab.?
multiglandulosa, *Humb. ex Spreng. Syst.* ii. 205.—Am. austr.
multipinna, *Benth. in Trans. Linn. Soc.* xxx. (1875) 440; *et in Mart. Fl. Bras.* xv. II. 388.—Bras.
multiplex, *Benth. in Hook. Journ. Bot.* iv. (1842) 373.—Bras.
muricata, Linn. Syst. ed. X. 1311 = Acacia muricata.
Musa, Vell. Fl. Flum. xi. t. 19 (Swartziae sp.).—Bras.
mutabilis, Roxb. Hort. Beng. 41; Fl. Ind. ii. 564 = rubricaulis.
myriacantha, *Baker, in Journ. Linn. Soc.* xxi. (1884) 339.—Madag.
myriadena, *Benth. in Trans. Linn. Soc.* xxx. (1875) 408; *et in Mart. Fl. Bras.* xv. II. 344.—Bras.
myriophylla, *Bong. ex Benth. in Hook. Journ. Bot.* iv. (1842) 384.—Bras.
myrtifolia, Sm. in Trans. Linn. Soc. i. (1791) 252 = Acacia myrtifolia.
natans, Linn. f. Suppl. 439 = Neptunia triquetra.
natans, Vahl, Symb. Bot. iii. 102 = Neptunia oleracea.
Nellyrenza, Heyne, in Wall. Cat. n. 5240 = Albizzia amara.
Nemu, Poir. Encyc. Suppl. i. 69 = Albizzia Julibrissin.
nepalensis, *Hoffmgg. Verz. Pfl. Nachtr.* II. 163.—Ind. or.
nepuria, Blume, ex Miq. Fl. Ind. Bat. i. I. 38 = Pithecolobium umbellatum.
nervosa, *Bong. ex Benth. in Hook. Journ. Bot.* iv. (1842) 370.—Bras.
neuroloma, *Benth. in Trans. Linn. Soc.* xxx. (1875) 406; *et in Mart. Fl. Bras.* xv. II. 341.—Bras.
nigricans, Labill. Nov. Holl. Pl. ii. 88. t. 238 = Acacia nigricans.
nigricans, Vahl, Eclog. Am. iii. 37. t. 29 = Acacia nudiflora.
nilotica, Linn. Sp. Pl. 521 = Acacia arabica.
nilotica, Thunb. Prod. Pl. Cap. 92 = Acacia horrida.
Niopo, Poir. Encyc. Suppl. i. 48 = Piptadenia peregrina.
nissobiensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 420.—Madag.
nitens, *Benth. in Hook. Journ. Bot.* iv. (1842) 404.—Bras.
nitida, Poir. Encyc. Suppl. i. 41 = Inga nitida.
nitida, Vahl, Symb. Bot. ii. 103 = Pithecolobium umbellatum.
nobilis, Poir. Encyc. Suppl. v. 528 = Inga nobilis.
nodosa, Linn. Sp. Pl. 516 = Cassia bacillaris.
notata, Steud. in Flora, xxvi. (1843) 758 = debilis.
nuda, *Benth. in Hook. Journ. Bot.* iv. (1842) 362.—Bras.
nudiflora, Rich. ex Poir. Encyc. Suppl. i. 65 = Acacia nudiflora.
nutans, Pers. Syn. ii. 266 = Dichrostachys nutans.
nutans, Vell. Fl. Flum. xi. t. 44 = Inga nutans.
obliqua, Lam. in Journ. Hist. Nat. Par. i. (1792) 89. t. 5 = Acacia suaveolens.
obliqua, Wendl. Bot. Beobacht. 57 = Acacia falcata.
oblonga, *Benth. in Hook. Journ. Bot.* iv. (1842) 365.—Bras.
obovata, *Benth. l. c.* 394.—Bras.
obovata, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 561 = Acacia modesta.
obtusa, Poir. Encyc. Suppl. i. 82 = Acacia macracantha.
obtusifolia, Poir. l. c. 46 = Calliandra purpurea.
obtusifolia, *Willd. Enum. Hort. Berol.* 1047.—Bras.

MIMOSA :—
octandra, Roxb. Pl. Corom. ii. 55. t. 200 = rubicaulis.
odoratissima, Linn. f. Suppl. 437 = Albizzia odoratissima.
Oerfota, Forsk. Fl. Aegypt. Arab. 177 = Acacia nubica?
oligacantha, DC. Prod. ii. 429 = distachya.
oligophylla, *Micheli, in Mem. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 56.—Paraguay.
ophthalmocentra, *Mart. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 415; *et in Mart. Fl. Bras.* xv. II. 359.—Bras.
Orfora, Steud. Nom. ed. I. 534; ed. II. ii. 149, sphalm. = *M. Oerfota*.
orthacantha, *Benth. in Hook. Journ. Bot.* iv. (1842) 365.—Bras.
orthocarpa, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 437; *et in Mart. Fl. Bras.* xv. II. 380.—Bras.
Ouyrarema, Aubl. Pl. Gui. ii. 946 = Acacia Ouyrarema.
Ouyratama, Vitm. Summa Pl. vi. 52 = Pithecolobium filicifolium.
Pacay, *Aubl. Pl. Gui.* ii. 946.—Guiana.
pachycarpa, *Benth. in Hook. Journ. Bot.* iv. (1842) 406.—Bras.
Pacoba, Vell. Fl. Flum. xi. t. 20 = Swartzia Flemmingii.
pallida, Poir. Encyc. Suppl. i. 65 = Prosopis juliflora.
palpitans, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1036 = somnians.
paludosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 400.—Bras.
paniculata, *Benth. l. c.* ii. (1840) 131.—Guiana.
paniculata, Benth. l. c. iv. (1842) 362 = rixosa.
paniculata, Poir. Encyc. Suppl. i. 74 = Acacia paniculata.
paniculata, Wendl. Bot. Beobacht. 57 = Acacia discolor.
paniculata, West, Beskr. Ste-Croix, 312 = Acacia riparia.
papposa, *Benth. in Hook. Journ. Bot.* iv. (1842) 375.—Bras.
paradoxa, Poir. Encyc. Suppl. v. 529 = Acacia armata.
Parae, Poir. l. c. i. 44 = Inga heterophylla.
Paraguariae, *Micheli, in Mem. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 53.—Paraguay.
paratyensis, Vell. Fl. Flum. xi. t. 41 = Acacia pedicellata.
parvifolia, Poir. Encyc. Suppl. i. 74 = Piptadenia peregrina.
parvifolia, Sw. Fl. Ind. Occ. ii. 984 = Pithecolobium parvifolium.
parvipinna, *Benth. in Hook. Journ. Bot.* iv. (1842) 384.—Bras.
patula, Poir. Encyc. Suppl. v. 529 = Acacia patula.
paucifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 412.—Bras.
pauciseta, Benth. in Trans. Linn. Soc. xxx. (1875) 391, in syn. = rixosa.
paupera, *Benth. in Hook. Journ. Bot.* iv. (1842) 372.—Bras.
pectinata, *Kunth, Mimos.* 5.—Ecuador.
pectinata, Poir. Encyc. Suppl. i. 48 = Parkia pectinata.
peduncularis, *Bong. ex Benth. in Hook. Journ. Bot.* iv. (1842) 388.—Bras.
pedunculata, Poir. Encyc. Suppl. i. 81 = Acacia Farnesiana.
pedunculata, Roxb. Hort. Beng. 40; Fl. Ind. ii. 551 = Parkia biglandulosa.
pedunculosa, *Micheli, in Mem. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 55.—Paraguay.
pellacantha, Meyen, Reise, i. 445 (pellocantha) = Acacia macracantha.
pellita, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1037 = asperata.
pendula, Poir. Encyc. Suppl. i. 47 = Parkia pendula.
pennata, Linn. Sp. Pl. 522 = Acacia pennata.
pennata, Roxb. ex Wight & Arn. Prod. 278 = Acacia caesia?
pennata, Russ. ex Wall. Cat. n. 5289 = rubicaulis.
pentagona, Schum. & Thonn. Beskr. Guin. Pl. 324 = Acacia pennata.
peregrina, Blanco, Fl. Filip. ed. I. 737 = Parkia Roxburghii.
peregrina, Linn. Sp. Pl. 520 = Piptadenia peregrina.
pernambucana, Linn. l. c. 519 = Desmanthus depressus.

MIMOSA :—
pernambucana, Mill. Gard. Dict. ed. VIII. n. 3 = Desmanthus virgatus.
peruviana, Poir. Encyc. Suppl. v. 529 = Acacia peruviana.
petiolaris, *Benth. in Hook. Journ. Bot.* iv. (1842) 376.—Bras.
philicoides, Cav. ex Steud. Nom. ed. II. i. 4, in syn.; ii. 149 = Lysiloma Schiedeana.
phyllocantha, Pers. Syn. ii. 267 = latispinosa.
phyllodinea, *Benth. in Hook. Journ. Bot.* iv. (1842) 412.—Bras.
pigra, Linn. Cent. Pl. i. 13; Amoen. Acad. iv. 274 = asperata.
piliflora, Sw. Fl. Ind. Occ. ii. 986 = Prosopis juliflora.
pilosa, *Lour. Fl. Cochinch.* 650.—China.
pilosiuscula, DC. Prod. ii. 432, in syn. = seq.
pilosula, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = Inga pilosiuscula.
pilulifera, Benth. in Hook. Journ. Bot. iv. (1842) 386 = incana.
pinifolia, Page, Prod. Southamp. Bot. Gard. 148 = Acacia linifolia.
pistaciaefolia, *Willd. Sp. Pl.* iv. 1028.—Am. trop.
pithecolobioides, *Benth. in Trans. Linn. Soc.* xxx. (1875) 413; *et in Mart. Fl. Bras.* xv. II. 357.—Bras.
plana, Vell. Fl. Flum. xi. t. 10 = Inga cordistipula.
plana, Vell. l. c. t. 28 = Acacia riparia.
planifrons, Koen. ex Wight & Arn. Prod. 276 = Acacia planifrons.
platycarpa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 417.—Guatemala.
platyloma, Benth. in Hook. Journ. Bot. iv. (1842) 406 = densa.
platyphylla, *Benth. l. c.* v. (1846) 85.—Bras.
plena, Linn. Sp. Pl. 519 = Neptunia plena.
podocarpa, Benth. in Hook. Journ. Bot. iv. (1842) 397 = somnians.
pogocephala, *Benth. l. c.* 382.—Bras.
pogonoclada, *Benth. in Trans. Linn. Soc.* xxx. (1875) 399; *et in Mart. Fl. Bras.* xv. II. 321.—Bras.
Pohlii, Benth. in Hook. Journ. Bot. iv. (1842) 375 = calocephala.
polyacantha, Willd. Sp. Pl. iv. 1034 = asperata.
polyancistra, *Benth. in Trans. Linn. Soc.* xxx. (1875) 422.—Mexic.
polyantha, *Benth. in Hook. Journ. Bot.* iv. (1842) 410.—Mexic.
polycarpa, *Kunth, Mimos.* 8. t. 3.—Peruv.
polycephala, *Benth. in Hook. Journ. Bot.* iv. (1842) 374.—Bras.
polydactyla, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1033.—Guian.
polyphylla, Poir. Encyc. Suppl. v. 530 = Acacia polyphylla.
polystachia, Linn. Sp. Pl. 520 = Entada polystachya.
portoricensis, Jacq. Coll. iv. 143 = Calliandra portoricensis.
procera, Roxb. Pl. Corom. ii. 12. t. 121 = Albizzia procera.
procumbens, *Boj. Hort. Maurit.* 113, nomen.—Madag.
procumbens, Schum. & Thonn. Beskr. Guin. Pl. 324 = asperata.
procurrens, *Benth. in Hook. Journ. Bot.* iv. (1842) 371.—Bras.
prostrata, Hort. Berol. ex Schleid. & Vog. in Nov. Act. Nat. Cur. xix. II. (1842) 70, nomen = marginata.
prostrata, Lam. Encyc. i. 10 = Neptunia oleracea.
prostrata, *Regel, in Act. Hort. Petrop.* iii. (1875) 117.—Hab.?
Psoralea, *Benth. in Trans. Linn. Soc.* xxx. (1875) 421.—Madag.
ptericina, Poir. Encyc. Suppl. i. 74 = Acacia filicina.
pteridifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 391.—Bras.
pterocarpa, Lam. Encyc. i. 13 = Gagnebina tamariscina.
puberula, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 88.—Mexic.
pubescens, Vent. Jard. Malm. 21 = Acacia pubescens, *R. Br.*
pubifera, Poir. Encyc. Suppl. i. 47 = Pithecolobium Saman.
pubigera, Hort. Par. ex Poir. Encyc. Suppl. i. 71 = Acacia pubescens, *R. Br.*

MIMOSA:—
pudibunda, Willd. Sp. Pl. iv. 1032 = pudica.
pudica, *Linn. Sp. Pl.* 518.—Bras.
pudica, Mill. Gard. Dict. ed. VIII. n. 4 = aculeata.
pulchella, Poir. Encyc. Suppl. v. 530 = Acacia pulchella.
pulchella, Roxb. Hort. Beng. 40; Fl. Ind. ii. 548 = Albizzia amara.
pulcherrima, Poir. Encyc. Suppl. i. 66 = Stryphnodendron floribundum.
pulchra, Dum.-Cours. Bot. Cult. ed. I. 420 = Acacia decurrens.
pulchra, Vell. Fl. Flum. xi. t. 18 = Swartzia Langsdorffii.
pumila, *Schlecht. in Linnaea*, xii. (1838) 557.—Mexic.
punctata, Linn. Syst. ed. X. 1311 = Neptunia plena.
punctulata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 408; *et in Mart. Fl. Bras.* xv. II. 345.—Bras.
pungens, Poir. Encyc. Suppl. i. 36 = Pithecolobium dulce.
purpurea, Linn. Sp. Pl. 517 = Calliandra purpurea.
pusilla, *Benth. Bot. Voy. Sulph.* 90.—Ecuador.
pycnocoma, Benth. in Hook. Journ. Bot. iv. (1842) 405 = Clausseni.
quadrangularis, Poir. Encyc. Suppl. i. 72 = Calliandra tetragona.
quadrangularis, *Vell. Fl. Flum.* xi. t. 36.—Bras.
quadrijuga, Salzm. ex Benth. in Hook. Journ. Bot. iv. (1842) 398 = somnians.
quadrivalvis, Linn. Sp. Pl. 522 = Schrankia aculeata.
quassiaefolia, Poir. Encyc. Suppl. i. 41 = Inga nitida.
quitensis, *Benth. in Hook. Journ. Bot.* iv. (1842) 408.—Ecuador.
racemosa, Schlecht. in Linnaea, xii. (1838) 557 = floribunda.
Radula, *Benth. in Hook. Journ. Bot.* iv. (1842) 377.—Bras.
ramosissima, *Benth. in Trans. Linn. Soc.* xxx. (1875) 394; *et in Mart. Fl. Bras.* xv. II. 312.—Bras.
ramulosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 385.—Bras.
Regnellii, *Benth. in Linnaea*, xxii. (1849) 529.—Bras.
remota, Benth. in Hook. Lond. Journ. Bot. v. (1846) 88 = distachya.
reptans, *Benth. in Hook. Journ. Bot.* iv. (1842) 372.—Bras.
reticulata, Herb. Wight, ex Wall. Cat. n. 5286 = Pithecolobium geminatum.
reticulata, Linn. Mant. i. 129 = Acacia horrida.
retortunium, Hort. Par. ex Lam. Encyc. i. 15 = Prosopis strombulifera.
retrorsa, Benth. in Hook. Journ. Bot. iv. (1842) 364 = insidiosa.
retusa, Jacq. Enum. Pl. Carib. 34 = Acacia riparia.
reversa, Sieber, ex Steud. Nom. ed. II. ii. 149 = Drepanocarpus lunatus.
revoluta, *Benth. in Hook. Journ. Bot.* iv. (1842) 409.—Am. trop.
rhodacantha, Pers. Syn. ii. 266 = Acacia rhodacantha.
rhodostachya, *Benth. in Trans. Linn. Soc.* xxx. (1875) 407; *et in Mart. Fl. Bras.* xv. II. 343.—Bras.
rhoifolia, Poir. Encyc. Suppl. v. 528 = Inga rhoifolia.
rhombifolia, Pers. Syn. ii. 263 = Acacia rhombifolia.
Riedeli, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 91.—Bras.
rigescens, Benth. in Hook. Journ. Bot. iv. (1842) 376 = meticulosa.
rigida, *Benth. l. c.* 377.—Bras.
rixosa, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 57.—Bras.
Rocae, *Lor. & Niederl. in Roca, Exped. Rio-Negro*, ii. 213.—Patagon.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 456 = Schrankia platycarpa.
rosea, Vahl, Eclog. Am. iii. 33. t. 25 = Pithecolobium Unguis-cati.
rostrata, Poir. Encyc. Suppl. i. 66 = Lysiloma rostrata.
Rottleri, Spreng. Syst. ii. 206 = rubicaulis.
rotundata, Pav. ex Benth. in Trans. Linn. Soc. xxx. (1875) 521 = Acacia rotundata.
rubicaulis, *Lam. Encyc.* i. 20.—Ind. or.
rubiginosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = Inga rubiginosa.
rudis, *Benth. in Hook. Journ. Bot.* iv. (1842) 377.—Bras.
rufescens, *Benth. in Trans. Linn. Soc.* xxx. (1875) 418; *et in Mart. Fl. Bras.* xv. II. 362.—Bras.
rufipila, *Benth. in Hook. Journ. Bot.* iv. (1842) 368.—Bras.

MIMOSA:—
rugata, Lam. Encyc. i. 20 = Acacia concinna, *DC.*
rupestris, *Benth. in Hook. Journ. Bot.* iv. (1842) 385.—Bras.
rutaefolia, Nois. ex Steud. Nom. ed. I. 534; ed. II. ii. 149 = Acacia rutaefolia.
rutaeolens, *Page, Prod. Southamp. Bot. Gard.* 148, nomen (Quid?).—Austral.
Sagotiana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 394; *et in Mart. Fl. Bras.* xv. II. 311.—Am. trop.
saligna, Labill. Nov. Holl. Pl. ii. 86. t. 235 = Acacia saligna.
salinarum, Rohr, ex Benth. in Trans. Linn. Soc. xxx. (1875) 501 = Acacia tortuosa.
salinarum, Vahl, Eclog. Am. iii. 35 = Prosopis juliflora.
Saman, Jacq. Fragm. 15. t. 9 = Pithecolobium Saman.
sanguinea, Bruce, Trav. ed. II. vii. 148; J. F. Gmel. Syst. 803 = Dichrostachys nutans.
sapindoides, Poir. Encyc. Suppl. i. 40 = Inga sapindoides.
saponaria, Lour. Fl. Cochinch. 653 = Albizzia saponaria.
saponaria, Roxb. ex Wight & Arn. Prod. 278 = Acacia concinna, *DC.*
sarmentosa, Desf. Tabl. ed. I. 181; Pers. Syn. ii. 266 = Acacia caesia.
Sassa, Poir. Encyc. Suppl. i. 49 = Albizzia fastigiata.
scabrella, *Benth. in Hook. Journ. Bot.* iv. (1842) 387.—Bras.
scandens, Linn. Sp. Pl. ed. II. 1501 = Entada scandens.
Schomburgkii, *Benth. in Hook. Journ. Bot.* ii. (1840) 133.—Guiana.
schrankioides, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 86.—Guiana.
Scolopendra, *Nois. ex Steud. Nom.* ed. I. 534; ed. II. ii. 149, nomen (Quid?).—Hab. ?
scorpioides, Forsk. Fl. Aegypt. Arab. p. lxxvii = Acacia Farnesiana.
scorpioides, Linn. Sp. Pl. 521 = Acacia arabica.
scutifera, Blanco, Fl. Filip. ed. I. 735 = Pithecolobium lobatum.
Seeressa, Steud. Nom. ed. II. ii. 149 = *M. Sirissa.*
Sejal, Forsk. Fl. Aegypt. Arab. 177 = Acacia hadiensis.
Selloi, *Benth. in Trans. Linn. Soc.* xxx. (1875) 436; *et in Mart. Fl. Bras.* xv. II. 379.—Bras.
semialata, Vell. Fl. Flum. xi. t. 5 = Inga marginata.
semicordata, *Roxb. Hort. Beng.* 40, nomen.—Ind. or.
semispinosa, *Linn. Sp. Pl.* 522.—N. Granat.
semispinosa, Spreng. Syst. ii. 206.—Cf. caduca.
Senegal, Linn. Sp. Pl. 521 = Acacia Senegal.
senegalensis, Forsk. Fl. Aegypt. Arab. 176 = seq. ?
senegalensis, Lam. Encyc. i. 19 = Acacia Senegal.
sensibilis, *Griseb. in Goett. Abh.* xxiv. (1879) 119.—Reg. Argent.
sensitiva, *Linn. Sp. Pl.* 518.—Am. trop.
sensitiva, Lodd. Bot. Cab. t. 249 = albida.
sepiaria, *Benth. in Hook. Journ. Bot.* iv. (1842) 395.—Bras.
sericantha, *Benth. l. c.* 392.—Bras.
sericea, Poir. Encyc. Suppl. i. 42 = Inga punctata.
sessilis, Vell. Fl. Flum. xi. t. 21 = Inga sessilis.
setipes, *Benth. in Hook. Journ. Bot.* iv. (1842) 376.—Bras.
setistipula, *Benth. l. c.* 379.—Bras.
setosa, *Benth. l. c.* 404.—Bras.
Seyal, DC. Prod. ii. 472, in syn. = *M. Sejal.*
Sicaria, Hoffmgg. Verz. Pfl. Nachtr. i. 221 = asperata.
simplicifolia, Linn. f. Suppl. 436 = Acacia laurifolia.
sinemariensis, Aubl. Pl. Gui. ii. 945 = Inga angustifolia; Feuillei.
sinuata, *Lour. Fl. Cochinch.* 653.—Cochinch.
Sirissa, Roxb. Hort. Beng. 40; Fl. Ind. ii. 544 = Albizzia Lebbek.
Skinneri, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 85.—Guatemala.
skleroxyla, Poir. Encyc. Suppl. v. 530 = Acacia scleroxyla.
Smithiana, Roxb. Hort. Beng. 40; Fl. Ind. ii. 550 = Albizzia stipulata.
somnians, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1036.—N. Granat.
somniculosa, H. B. & K. Nov. Gen. et Sp. vi. 257 = somnians.
Sophorae, Labill. Nov. Holl. Pl. ii. 87 = Acacia longifolia.

MIMOSA:—
soratensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 427.—Bolivia.
sordida, *Benth. in Hook. Journ. Bot.* iv. (1842) 389.—Bras.
sparsa, *Benth. l. c.* 385.—Bras.
spartioides, Vahl, ex DC. Prod. ii. 456 = Calliandra haematomma.
speciosa, Jacq. Ic. Pl. Rar. i. 19. t. 198; Coll i. 47 = Albizzia Lebbek.
speciosa, Thunb. in Trans. Linn. Soc. ii. (1794) 336 = Albizzia Julibrissin.
spectabilis, Vahl, in Skrift. Nat. Selsk. Kjoeb. ii. (1792) i. 219 = Inga spectabilis.
spicata, E. Mey. Comm. Pl. Afr. Austr. 164 = Entada natalensis.
spiciflora, *Karst. Fl. Columb.* ii. 61. t. 131.—Venezuela.
spinaefolia, Poir. Encyc. Suppl. v. 528 = Pithecolobium circinale.
spinisiliqua, Klein, ex Poir. Encyc. Suppl. i. 54; Rottl. ex Spreng. Syst. ii. 206 = rubicaulis.
spinosissima, Poir. Encyc. Suppl. i. 78 = Acacia Catechu *vel* A. Suma.
spiriflora, C. Muell. in Walp. Ann. vii. 829, sphalm. = spiciflora.
splendens, Poir. Encyc. Suppl. i. 43 = Inga splendens.
Sprengelii, DC. Prod. ii. 430 = ciliata.
Spruceana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 418; *et in Mart. Fl. Bras.* xv. II. 362.—Bras.
spuria, Poir. Encyc. Suppl. i. 40 = Inga spuria.
staminea, Thunb. Pl. Bras. Dec. ii. 22 = Calliandra virgata.
stellata, Forsk. Fl. Aegypt. Arab. 177 = Acacia stellata.
stellata, Lour. Fl. Cochinch. 651 = Acacia taxifolia.
stenoptera, *Benth. in Trans. Linn. Soc.* xxx. (1875) 395.—Reg. Argent.
Stephaniana, Bieb. Casp. 205; ex ej. Fl. Taur. Cauc. ii. 449 = Prosopis Stephaniana.
stipulacea, Roxb. Fl. Ind. ii. 549 = Albizzia stipulata.
stipularis, *Bong. ex Benth. in Hook. Journ. Bot.* iv. (1842) 380.—Bras.
stipulata, Roxb. Hort. Beng. 40, nomen = Albizzia stipulata.
stolonifera, Perr. ex DC. Prod. ii. 444 = Neptunia oleracea.
striata, Poir. Encyc. Suppl. i. 83 = Acacia striata.
striato-stipula, Steud. in Flora, xxvi. (1843) 758 = pudica.
stricta, Andr. Bot. Rep. t. 53 = Acacia stricta.
stricta, Dum.-Cours. Bot. Cult. ed. 1. 420 = Albizzia lophantha.
strigillosa, *Torr. & Gray, Fl. N. Am.* i. 399.—Am. bor. occ.
strigosa, Pers. Syn. ii. 263 = Acacia strigulosa.
strigosa, Willd. Sp. Pl. iv. 1030 = floribunda.
strombulifera, Lam. Encyc. i. 15 = Prosopis strombulifera.
suaveolens, Salisb. Prod. 325 = Acacia Farnesiana.
suaveolens, Sm. in Trans. Linn. Soc. i. (1791) 253 = Acacia suaveolens.
subenervis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 432; *et in Mart. Fl. Bras.* xv. II. 370.—Bras.
subinermis, Benth. in Hook. Journ. Bot. iv. (1842) 385 = rupestris.
subnervosa, Pav. ex Benth. in Trans. Linn. Soc. xxx. (1875) 541 = Calliandra subnervosa.
subsericea, *Benth. in Hook. Journ. Bot.* iv. (1842) 380.—Bras.
subvestita, *Benth. l. c.* 379.—Bras.
sulcata, Poir. Encyc. Suppl. v. 530 = Acacia sulcata.
Suma, Roxb. Hort. Beng. 41; Fl. Ind. ii. 563 = Acacia Suma.
Sundra, Roxb. ll. cc. 41; 562 = Acacia Sundra.
tamarindifolia, Linn. Sp. Pl. 523 = Acacia tamarindifolia.
tamariscifolia, Russ. Herb. ex Steud. Nom. ed. II. ii. 149 = Acacia concinna, *DC.*
tamariscifolia, Steud. l. c. sphalm. = *tamarindifolia.*
tamariscina, Heyne, ex Wall. Cat. sub n. 5227 = Acacia Suma.
tamariscina, Lam. Encyc. i. 13 = Gagnebina tamariscina.
taxifolia, *Bong. ex Benth. in Hook. Journ. Bot.* iv. (1842) 373.—Bras.
taxifolia, Pers. Syn. ii. 266 = Parkia africana.
taxifolia, Poir. Encyc. Suppl. i. 59 = Acacia taxifolia.

MIMOSA:—
tenuiflora, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 92.—Mexic.
tenuiflora, Poir. Encyc. Suppl. i. 82 = M. Cabrera.
tenuifolia, Blanco, Fl. Filip. ed. I. 739 = Acacia concinna, *DC.*
tenuifolia, Linn. Sp. Pl. 523 = Acacia paniculata.
tenuifolia, Roxb. Hort. Beng. 41 = Acacia Intsia.
tenuis, Vell. Fl. Flum. xi. t. 11 = Inga nutans.
tergemina, Linn. Sp. Pl. 517 = Calliandra tergemina.
terminalis, Salisb. Prod. 325 = Acacia elata.
terminalis, Vell. Fl. Flum. xi. t. 30 = Pithecolobium polycephalum.
ternata, Pers. Syn. ii. 261 = Acacia taxifolia.
tetragona, Poir. Encyc. Suppl. i. 56 = Schrankia hamata.
tetrandra, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1032 = pudica.
tetraphylla, Vell. Fl. Flum. xi. t. 8 = Inga fagifolia.
thyrsoidea, *Griseb. Fl. Brit. W. Ind.* 219.—Jamaica.
thyrsoidea, Poir. Encyc. Suppl. v. 528 = Inga alba.
tomentosa, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1033.—Venezuela.
tomentosa, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 208 = Acacia tomentosa.
torta, Roxb. Hort. Beng. 41; Fl. Ind. ii. 566 = Acacia pennata.
tortilis, Forsk. Fl. Aegypt. Arab. 176 = Acacia spirocarpa *vel* tortilis.
tortuosa, Burm. f. Fl. Ind. 224 = Acacia Burmanniana.
tortuosa, Linn. Syst. ed. X. 1312 = Acacia tortuosa.
tovarensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 407.—Venezuela.
trachycarpa, *Benth. in Hook. Journ. Bot.* iv. (1842) 407.—Bras.
trachycephala, *Benth. in Trans. Linn. Soc.* xxx. (1875) 395; *et in Mart. Fl. Bras.* xv. II. 314.—Bras.
trapezifolia, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 546 = Pithecolobium Clypearia.
trapezifolia, Vahl, Eclog. Am. iii. 36 = Pithecolobium trapezifolium.
tremula, *Benth. in Hook. Journ. Bot.* iv. (1842) 380.—Bras.
Trianae, *Benth. in Trans. Linn. Soc.* xxx. (1875) 412.—N. Granat.
tricephala, *Cham. & Schlecht. in Linnaea*, v. (1830) 591.—Mexic.
trichocephala, *Benth. in Trans. Linn. Soc.* xxx. (1875) 394; *et in Mart. Fl. Bras.* xv. II. 312.—Bras.
trichodes, Jacq. Hort. Schoenb. iii. 76. t. 394 = Leucaena trichodes.
triflora, Pav. ex Benth. in Trans. Linn. Soc. xxx. (1875) 591 = Pithecolobium triflorum.
trijuga, *Benth. in Hook. Journ. Bot.* iv. (1842) 398.—Bras.
trinervis, Desf. Tabl. ed. I. 181 = Acacia trinervis.
triphylla, Vell. Fl. Flum. xi. t. 22 = Swartzia elegans?
triquetra, Vahl, Symb. Bot. iii. 102 = Neptunia triquetra.
trisperma, Vell. Fl. Flum. xi. t. 40 = Piptadenia trisperma.
ulicifolia, Salisb. Prod. 324; Wendl. Coll. i. 25. t. 6 = Acacia juniperina.
ulicina, Benth. Fl. Austral. ii. 332, in syn. sphalm. = *ulicifolia*.
umbellata, Vahl, Symb. Bot. ii. 103 = Pithecolobium umbellatum.
umbellata, Vell. Fl. Flum. xi. t. 12 = Inga affinis.
umbellifera, Vahl, Eclog. Am. iii. 30 = Inga umbellifera.
umbrifera, Banks, ex Wight & Arn. Prod. 276 = Acacia planifrons.
uncinella, Poir. Encyc. Suppl. i. 80 = acanthocarpa.
Unguis-cati, Blanco, Fl. Filip. ed. I. 731 = Pithecolobium dulce.
Unguis-cati, Forsk. Fl. Aegypt. Arab. 176 = Acacia mellifera.
Unguis-cati, Linn. Sp. Pl. 517 = Pithecolobium Unguis-cati.
unijuga, Duchass. & Walp. in Linnaea, xxiii. (1850) 744 = pudica.
uraguensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 202.—Uraguay.
ursina, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 56.—Bras.
vaga, Aubl. Pl. Gui. ii. 945 = Pithecolobium trapezifolium.
vaga, Linn. Sp. Pl. 519 = Acacia vaga.

MIMOSA:—
vaga, Vell. Fl. Flum. xi. t. 13 = Pithecolobium Auaremotemo.
varia, Perr. ex DC. Prod. ii. 445 = Dichrostachys nutans.
Velloziana, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 9.—Bras.
velutina, Poir. Encyc. Suppl. i. 42 = Inga velutina.
venusta, Poir. l. c. v. 529 = Calliandra portoricensis.
venustula, *Dum.-Cours. Bot. Cult.* ed. II. vii. 297 (Quid?).—Austral.
verecunda, *Benth. in Hook. Journ. Bot.* iv. (1842) 368.—Bras.
vernicosa, *Bong. ex Benth. l. c.* 404.—Bras.
verrucosa, *Benth. l. c.* 390.—Bras.
verticillata, L'Hérit. Sert. Angl. 30 = Acacia verticillata.
vestita, *Benth. in Hook. Journ. Bot.* iv. (1842) 369.—Bras.
vialis, Benth. in Trans. Linn. Soc. xxx. (1875) 434 = somnians.
villosa, Sw. Prod. Veg. Ind. Occ. 85 = Acacia villosa.
violacea, *Bolle, in Peters, Reise Mossamb. Bot.* 8.—Afr. trop.
violacea, Bonpl. ex Tenore, Ind. Sem. Hort. Neap. (1842) 12. n. 6 = rixosa.
virescens, Poir. Encyc. Suppl. v. 530 = Acacia virescens.
virgata, Linn. Sp. Pl. 519 = Desmanthus virgatus.
virginalis, *Arruda, in Koster, Trav. Bras.* 500.—Bras.
virgultosa, Vahl, Eclog. Am. iii. 32 = Inga virgultosa.
viscida, *Willd. Enum. Hort. Berol.* 1048.—Bras.
viva, *Linn. Sp. Pl.* 517.—Jamaica.
viva, Vell. Flum. xi. t. 33 = Velloziana.
Warmingii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 413; *et in Mart. Fl. Bras.* xv. II. 356.—Bras.
Weddelliana, *Benth. ll. cc.* 438; 383.—Bras.
Willdenowii, Poir. Encyc. Suppl. i. 50 = floribunda.
Woony, Heyne, ex Wall. Cat. n. 5226 = Acacia ferruginea.
Wrightii, *A. Gray, Pl. Wright.* ii. 52.—N. Mexic.
xanthocentra, Mart. in Flora, xxi. (1838) II. Beibl. 50 = polycarpa.
xantholasia, Benth. in Hook. Lond. Journ. Bot. v. (1846) 88 = arenosa.
Xanti, *A. Gray, in Proc. Am. Acad.* v. (1861) 157.—Calif.
xylocarpa, Roxb. Pl. Corom. i. 68. t. 100 = Xylia dolabriformis.
Ynga, Vell. vide *M. Inga*, Vell.
zygophylla, *Benth. in A. Gray, Pl. Wright.* i. 61.—N. Mexic.

MIMULOPSIS, Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 677. *ACANTHACEAE*, Benth. & Hook. f. ii. 1080.
diffusa, *Baker, in Journ. Linn. Soc.* xx. (1883) 219.—Madag.
lanceolata, *Baker, l. c.* 220.—Madag.
Solmsii, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 677.—Afr. trop.
speciosa, *Baker, in Journ. Linn. Soc.* xviii. (1881) 274.—Madag.

MIMULUS, Linn. in Act. Soc. Sc. Upsal. (1741) 82. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 946.
CYNORRHYNCHIUM, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 207.
DIPLACUS, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 137.
ERYTHRANTHE, Spach. Hist. Vég. Phan. ix. 312 (1840).
EUNANUS, Benth. in DC. Prod. x. 374 (1846).
MIMETANTHE, Greene, in Bull. Calif. Acad. i. IV. (1885) 181.
MONAVIA, Adans. Fam. ii. 211 (1763).
UVEDALIA, R. Br. Prod. 440 (1810).
acutidens, Greene, in Bull. Calif. Acad. i. III. (1885) 117 = inconspicuus.
alatus, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 361.—Am. bor.
Alecterolophus, Scop. Fl. Garn. ed. II. i. 435 = Rhinanthus Crista-galli.
alsinoides, *Dougl. ex Benth. Scoph. Ind.* 29.—Am. bor. occ.
andicolus, *H. B. & K. Nov. Gen. et Sp.* ii. 370.—Peruv.

MIMULUS:—
androsaceus, Curran, ex Greene, in Bull. Calif. Acad. i. III. (1885) 121 = Palmeri.
angustifolius, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 119.—Abyss.
assamicus, Griff. in Madr. Journ. Sc. iv. (1836) 375 = nepalensis.
atropurpureus, *Kellogg, in Proc. Calif. Acad.* i. (1873) 59.—Calif.
aurantiacus, Curt. Bot. Mag. t. 354 = glutinosus.
aurantiacus, *Renjifo, in Renjifo & Phil. Desc. Alg. Pl. Nuev.* (1884) 7.—Chili.
barbatus, *Greene, in Bull. Calif. Acad.* i. I. (1884) 9.—Calif.
bicolor, *Hartw. ex Benth. Pl. Hartw.* 328.—Calif.
Bigelovii, *A. Gray, Bot. Calif.* i. 564; *et in Proc. Am. Acad.* xi. (1876) 96.—Am. bor. occ.
Bolanderi, *A. Gray, l. c.* vii. (1868) 381.—Calif.
brevipes, *Benth. Scroph. Ind.* 28.—Calif.
brevipes, A. Gray, in Pacif. Rail. Rep. iv. 120 = Bolanderi.
Bridgesii, *Clos, in C. Gay, Fl. Chil.* 141.—Chili.
cardinalis, *Dougl. ex Benth. Scroph. Ind.* 28.—Am. bor. occ.
Colensoi, *T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 179.—N. Zel.
Crista-galli, Scop. Fl. Carn. ed. II. i. 434 = Rhinanthus Crista-galli.
cupreus, Regel, Gartenfl. (1864) 3 = luteus.
debilis, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 62 = Uvedaliae.
dentatus, A. Gray, in Pacif. Rail. Rep. iv. 120 = nudatus.
dentatus, *Nutt. ex Benth. in DC. Prod.* x. 373.—Am. bor. occ.
depressus, *Phil. Fl. Atac.* 45.—Chili.
Douglasii, *A. Gray, in Bot. Calif.* i. 563; *et in Proc. Am. Acad.* xi. (1876) 95.—Calif.
Eisenii, Kellogg, in Proc. Calif. Acad. vii. I. (1876) 89 = laciniatus.
exiguus, *A. Gray, in Proc. Am. Acad.* xx. (1885) 307.—Calif.
exilis, *Durand & Hilg. in Pacif. Rail. Rep.* v. 12.—Calif.
floribundus, *Dougl. in Lindl. Bot. Reg.* t. 1125.—Calif.
Fremonti, *A. Gray, in Bot. Calif.* i. 565; *et in Proc. Am. Acad.* xxi. (1876) 96.—Calif.
geniculatus, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 280.—Calif.
Geyeri, Torr. in Nicoll. Rep. Hydrog. Upper Missis. (1843) 157 = glabratus.
glabratus, A. Gray, Bot. Mex. Bound. 116 = Jamesii.
glabratus, *H. B. & K. Nov. Gen. et Sp.* ii. 370.—Mexic.
glandulosus, *Lehm. Del. Sem. Hort. Hamb.* (1830) 7.—*Cf. Linnaea*, vi. (1831) *Litt.* 11.—Am. bor.
glaucescens, *Greene, in Bull. Calif. Acad.* i. (1885) 113.—Calif.
glutinosus, *Wendl. Bot. Beobacht.* 51.—Calif.
gracilis, *R. Br. Prod.* 439.—Austral.
guttatus, [Fisch.] Hort. Gorenk. ed. II. (1812) 25; DC. Cat. Hort. Monsp. 127 = luteus.
Hallii, Greene, in Bull. Calif. Acad. i. III. (1885) 113 = glabratus.
Harrisonianus ×, Hort.—Cf. Gard. Chron. (1879) II. 86.
hirsutus, Blume, Bijdr. 756 = Torenia flava.
Hodsoni ×, Turner, in Loud. Gard. Mag. xiii. (1837) 333.
inconspicuus, *A. Gray, in Pacif. Rail. Rep.* iv. 120.—Calif.
inodorus, Greene, in Bull. Calif. Acad. i. III. (1885) 119 = moschatus.
Jamesii, *Torr. & Gray, ex Benth. in DC. Prod.* x. 371.—Am. bor. occ.
javanicus, Blume, Cat. Gew. Buitenz. 83 = Vandellia crustacea.
Kingii, *Phil. in Anal. Univ. Chil.* (1873) 528.—Chili.
laciniatus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 98.—Calif.
lanatus, *Pursh, ex Loud. Hort. Brit.* 251 (Quid?).—Am. bor.
latifolius, *A. Gray, in Proc. Am. Acad.* xi. (1876) 95.—Calif.
leptaleus, *A. Gray, l. c.* 96.—Calif.
Lewisii, *Pursh, Fl. Am. Sept.* ii. 427. t. 20.—Am. bor. occ.
linearis, Benth. Scroph. Ind. 27 = glutinosus.

MIMULUS:—
longipes, *Phil. in Linnaea*, xxix. (1857–58) 28.—Chili.
luteus, *Linn. Sp. Pl.* ed. II. 884.—Am. bor. et austr.
lyratus, Benth. Scroph. Ind. 28 = luteus.
Maclaineanus, Paxt. Fl. Mag. ix. (1842) t. 147 = seq.
Maclainii, Fergus. ex Hook. Bot. Mag. t. 3924 = cardinalis.
madagascariensis, *Benth. in DC. Prod.* x. 369.—Madag.
madrensis, Seem. Bot. Voy. Herald, 322. t. 58 = glabratus.
mephiticus, *Greene, in Bull. Calif. Acad.* i. 1. (1884) 9.—Calif.
microphyllus, Benth. in DC. Prod. x. 371 = luteus.
mohavensis, *Lemmon, in Coult. Bot. Gaz.* ix. (1884) 142.—Calif.
moniliformis, *Greene, in Bull. Calif. Acad.* i. 1. (1884) 10.—Calif.
montioides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 380.—Am. bor. occ.
moschatus, *Dougl. in Lindl. Bot. Reg.* t. 1118.—Am. bor. occ.
nanus, *Hook. & Arn. Bot. Beech. Voy.* 379.—Am. bor. occ.
nanus, *Phil. Fl. Atac.* 45.—Chili.
nasutus, *Greene, in Bull. Calif. Acad.* i. III. (1885) 112.—Am. bor. occ.
nepalensis, *Benth. in Wall. Cat.* n. 3917.—Reg Himal.; As. bor. occ.
nudatus, *Curran, ex Greene, in Bull. Calif. Acad.* i. III. (1885) 114.—Calif.
nummularis, C. Gay, Fl. Chil. Atl. t. 57 = luteus.
ocellatus, Bert. ex Steud. Nom. ed. II. ii. 150 = luteus.
orbicularis, *Wall. Cat.* n. 3919.—Burma.
Orizabae, *Benth. in DC. Prod.* x. 372.—Mexic.
pallidus, Salisb. Prod. 100 = ringens.
Palmeri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 82.—Calif.
Parishii, *Greene, in Bull. Calif. Acad.* i. III. (1885) 108.—Calif.
Parryi, *A. Gray, in Proc. Am. Acad.* xi. (1876) 197.—Am. bor. occ.
parviflorus, Lindl. Bot. Reg. t. 874 = pilosiusculus.
peduncularis, *Dougl. ex Benth. Scroph. Ind.* 29.—Am. bor. occ.
perfoliatus, H. B. & K. Nov. Gen. et Sp. ii. 371 = Leucocarpus alatus.
Pilingi, Regel, Gartenfl. (1869) t. 631, sphalm. = *Tilingii*.
pilosiusculus, *H. B. & K. Nov. Gen. et Sp.* ii. 370.—Peruv.
pilosus, S. Wats. in Bot. King's Exp. 225 = exilis.
Pissisi, *Phil. in Anal. Univ. Chil.* (1861) 57; *et in Linnaea*, xxxiii. (1864–65) 212.—Chili.
pluriflorus, Hook. & Arn. Bot. Beech. Voy. 378 = nanus.
Prattenii, Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 98 = bicolor.
primuloides, *Benth. Scroph. Ind.* 29.—Am. bor. or.
propinquus, Lindl. Bot. Reg. i. 1330 = glabratus.
prostratus, *Benth. in DC. Prod.* x. 373.—Austral.
pteropus, Rafin. Fl. Ludov. 44 (Quid?).—Am. bor.
pubescens, *Benth. in DC. Prod.* x. 372.—Mexic.
Pulsiferae, *A. Gray, in Proc. Am. Acad.* xi. (1876) 98.—Calif.
punctatus, Miers, ex Bert. in Merc. Chil. (1829) 700.—Cf. Am. Journ. Sc. xxiii. (1833) 254 = luteus.
puniceus, *Steud. Nom.* ed. II. ii. 150.—Calif.
pusillus, Benth. in DC. Prod. x. 369 = gracilis.
radicans, *Hook. f. Fl. N. Zel.* i. 188.—N. Zel.
Rattani, *A. Gray, in Proc. Am. Acad.* xx. (1885) 307.—Calif.
reniformis, Engelm. ex Benth. in DC. Prod. x. 371 = Jamesii.
repens, *R. Br. Prod.* 432.—Austral.; N. Zel.
ringens, *Linn. Sp. Pl.* 634.—Am. bor.
rivularis, Nutt. in Journ. Acad. Philad. vii. (1834) 47 = luteus.
roseus, Dougl. ex Lindl. Bot. Reg. t. 1591 = Lewisii.
rubellus, *A. Gray, in Torr. Bot. Mex. Bound.* 116.—Am. bor. occ.
Scouleri, *Hook. Fl. Bor. Am.* ii. 100.—Am. bor. occ.
sessilifolius, *Maxim. in Bull. Acad. Pétersb.* xx. (1874) 436.—Japon.
Smithii, Lindl. Bot. Reg. t. 1674 = luteus.
splendens, Hort. ex Steud. Nom. ed. I. 534, 535 = luteus.

MIMULUS:—
stachydifolius, *Maxim. in Bull. Acad. Pétersb.* xx. (1874) 438.—As. or.
strictus, Benth. in Wall. Cat. n. 3918 = gracilis.
sylvaticus, *Phil. in Linnaea*, xxx. (1859–60) 197.—Chili.
tenellus, Bunge, Enum. Pl. Chin. Bor. 49 = nepalensis.
tenellus, Nutt. ex A. Gray, in Proc. Am. Acad. xi. (1876) 98 = luteus.
Tilingii, Regel. Gartenfl. (1869) 321 = luteus.
Torreyi, *A. Gray, in Proc. Am. Acad.* xi. (1876) 197.—Calif.
tricolor, *Hartw. ex Lindl. in Journ. Hort. Soc.* iv. (1849) 222.—Calif.
Uvedaliae, *Benth. in DC. Prod.* x. 369.—Austral.
variegatus, Lodd. Bot. Cab. t. 1872 = luteus.
violaceus, Blanco, Fl. Filip. ed. II. 357 = Vandellia mollis?
viscosus, Moench, Meth. Suppl. 168 = glutinosus.
Wilsoni, D. Don, ex Steud. Nom. ed. II. ii. 150 = glutinosus.

MIMUSOPS, Linn. Diss. Dass. 14 (1747); Amoen. Acad. i. 397 (1749). *SAPOTACEAE*, Benth. & Hook. f. ii. 661.
ELENGI, Adans. Fam. ii. 166 (1763).
MANILKARA, Adans. l. c. (1763).
STISSERIA, Scop. Introd. 199 (1777).
SYNARRHENA, Fisch. & Mey. in Bull. Sc. Acad. Pétersb. viii. (1841) 255.
acuminata, *Blume, Bijdr.* 672.—Java.
angustifolia, *Boj. Hort. Maurit.* 198.—Ins. Mascar.
attenuata, Wall. Cat. n. 4150 = Sideroxylon ferrugineum.
Balata, Crueg. ex Griseb. Fl. Brit. W. Ind. 400 = globosa.
Balata, Gaertn. f. Fruct. iii. 133 = M. Kauki.
Balota, Blume, Bijdr. 673 = praec.
bidentata, A. DC. in DC. Prod. viii. 204 = globosa.
Bojeri, *A. DC. l. c.* 205.—Ins. Maurit.
Browniana, Benth. Fl. Austral. iv. 285 = M. Kauki.
caffra, *E. Mey. ex A. DC. in DC. Prod.* viii. 203.—Afr. austr.
calyptranthifolia, *Span. ex Hook. Comp. Bot. Mag.* i. (1835) 348, nomen.—Ins. Timor.
coriacea, *Miq. in Mart. Fl. Bras.* vii. 44.—Bras.
cotinifolia, A. Cunn. ex A. DC. in DC. Prod. viii. 176, 207 = Hormogyne cotinifolia.
cuneifolia, *Baker, in Oliver, Fl. Trop. Afr.* iii. 506.—Afr. trop.
cyanocarpa, A. Cunn. ex Sweet, Hort. Brit. ed. III. 451, nomen = Browniana?
dissecta, R. Br. Prod. 531, in obs. = M. Kauki.
dissecta, Buch.-Ham. ex A. DC. in DC. Prod. viii. 205 = hexandra.
elata, *Allem. ex Miq. in Mart. Fl. Bras.* vii. 42.—Bras.
Elengi, Boj. Hort. Maurit. 198 = M. Kauki?
Elengi, *Linn. Sp. Pl.* 349.—Ind. or.; Malaya.
Elengi, Sieber, ex A. DC. in DC. Prod viii. 201, 207 = Labramia Bojeri.
Elengi, Wight, in Wall. Cat. n. 4146 = Sapota elengoides.
erythroxyla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
erythroxylon, *Boj. Hort. Maurit.* 198.—Ins. Maurit.
erythroxylon, Llanos, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 126 = parvifolia, *R. Br.*
floribunda, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 3.—Bras.
fruticosa, *Boj. Hort. Maurit.* 198.—Afr. trop.
globosa, *Gaertn. f. Fruct.* iii. 132.—Am. trop.
hexandra, Bout. ex Boj. Hort. Maurit. 198 = Labramia Bojeri.
hexandra, *Roxb. Pl. Corom.* i. 16. t. 15.—Ind. or.
hexandra, Wall. Cat. n. 4148 = Achras Sapota.
Hookeri, A. DC. in DC. Prod. viii. 204 = M. Kauki.
Horneana, Hartog, in Journ. Bot. xvii. (1879) 358, nomen = Northea seychellana.
Imbricaria, Blume, Bijdr. 675 = Labramia Bojeri.
Imbricaria, Bout. ex A. DC. in DC. Prod. viii. 200 = Imbricaria petiolaris.
Imbricaria, Willd. Sp. Pl. ii. 326 = Imbricaria maxima.
indica, A. DC. in DC. Prod. viii. 205 = hexandra.
indica, Kurz, Andam. Rep. 42 = littoralis.
Jainiqui, *Wright, ex Griseb. Cat. Pl. Cub.* 164.—Cuba.
javensis, *Burck, in Ann. Jard. Buitenz.* v. (1885) 63.—Java.

MIMUSOPS:—
Kauki, Boj. Hort. Maurit. 198 = Bojeri.
Kauki, *Linn. Sp. Pl.* 349.—Burma; Malaya; Austral.
Kauki, Wall. Cat. n. 4149 A, C = hexandra.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* iii. 507.—Afr. trop.
Kummel, *Bruce, ex A. DC. in DC. Prod.* viii. 203.—Abyssin.
lacera, *Baker, in Oliver, Fl. Trop. Afr.* iii. 507.—Afr. trop.
laevis, *Wall. Cat.* n. 4152.—Burma.
littoralis, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 138.—Ins. Nicobar.
longifolia, *A. DC. in DC. Prod.* viii. 205.—Bras.
lucida, *Poir. Encyc. Suppl.* iii. 705.—Ind. or.
lucida, Wall. Cat. n. 4147 = Payena lucida.
Manilkara, G. Don, Gen. Syst. iv. 35 = M. Kauki.
Mochisia, *Baker, in Oliver, Fl. Trop. Afr.* iii. 506.—Afr. trop.
moluccana, Kostel. Allg. Med.-Pharm. Fl. iii. 1104 = Maba elliptica.
multinervis, *Baker, in Oliver, Fl. Afr. trop.* iii. 506.—Afr. trop.
Nattarium, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 30.—Ins. Maurit.
obovata, *Sond. in Linnaea*, xxiii. (1850) 75.—Afr. austr.
obtusifolia, Lam. Encyc. iv. 186 = M. Kauki?
obtusifolia, Wall. Cat. n. 4473 = Labourdonnaisia calophylloides.
parvifolia, *R. Br. Prod.* 531.—Austral.
parvifolia, *Kurz, For. Fl. Brit. Burma*, ii. 124.—Burma.
racemosa, *Jacq. Coll. Suppl.* 108.—Ins. Maurit.
retusa, Willem. in Usteri, Ann. Bot. xviii. (1796) 30 = Imbricaria maxima.
revoluta, *Hochst. in Flora*, xxvii. (1844) 825.—Afr. austr.
Roxburghiana, *Wight, Ic.* t. 1588.—Ind. or.
rufula, *Miq. in Mart. Fl. Bras.* vii. 44.—Bras.
Salzmanni, *A. DC. in DC. Prod.* viii. 205.—Bras.
Schimperi, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 22.—Abyssin.
Sieberi, *A. DC. in DC. Prod.* viii. 204.—Am. bor.; Ind. occ.
subsericea, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 3.—Bras.
surinamensis, *Miq. in Mart. Fl. Bras.* vii. 43.—Guian.
Talosan, Blanco, Fl. Filip. ed. I. 284 = Helicteres spicata.
timorensis, *Burck, in Ann. Jard. Buitenz.* v. (1885) 63.—Ins. Timor.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 74.—Afr. austr.

MINA, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 3 (1824) = **Ipomoea**, Linn. (Convolv.).
lobata, Cerv. l. c. = I. versicolor.

MINAEA, Lojac. in Nuov. Giorn. Bot. xiii. (1881) 305 = **Thlaspi**, Tourn. (Crucif.).
Prolongoi, Lojac. l. c. 306 = T. Prolongi.
Saviana, Lojac. l. c. = Bivonaea Saviana.

MINDERERA, Ram. ex Schrad. Journ. 1800. II. (1801) 467 = **Merendera**? Ram. (Liliac.).

MINDIUM, Adans. Fam. ii. 134 (1763); Juss. Gen. 164 (1789) = **Michauxia**, L'Hérit. (Campan.).
canariense, Rafin. Fl. Tellur. ii. 79 = Canarina campanulata.
Rhazis, Juss. Gen. 164 = Mich. campanuloides.
spicatum, J. F. Gmel. Syst. 618 = Mich. campanuloides.

MINGUARTIA, Miers, in Journ. Linn. Soc. xvii. (1879) 338, sphalm. = **Minquartia**, Aubl. (Bignoniac.).

MINKELERSIA, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 200. *LEGUMINOSAE*, Benth. & Hook. f. i. 539.
biflora, *Hemsl. Diagn. Pl. Nov.* 48.—Mexic.
galactioides, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 200.—Mexic.

MINQUARTIA, Aubl. Pl. Gui. Suppl. 4 (1775). *BIGNONIACEAE*.
guianensis, *Aubl. l. c.*—Guian.

MINTHA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 78 = seq.

MINTHE, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = **Mentha**, Tourn. (Labiat.).
pulegia, St. Lag. l. c. = Mentha Pulegium.

MINTOSTACHYS, Spach, Hist. Vég. Phan. ix. 164 (1840) = **Bystropogon**, L'Hérit. (Labiat.).

MINUARTIA, Linn. Sp. Pl. ed. I. 89 (1753) = **Arenaria**, Rupp. (Caryoph.).
campestris, DC. Prod. i. 380 = A. montana.
campestris, Linn. Sp. Pl. 89 (= *Alsine campestris*).—Hispan.
dichotoma, Echeand. Fl. Caesaraug.; ex Willd. & Lange, Prod. Fl. Hisp. iii. 608 = A. montana.
dichotoma, Linn. Sp. Pl. 89 (= *Alsine dichotoma*).—Hispan.
fasciculata, Reichb. Ic. Fl. Germ. v. 28. t. 206.—A. globulosa.
fastigiata, Reichb. l. c. 28 = A. globulosa.
montana, Cav. ex Nym. Consp. 120 (= *Alsine campestris*).—Hispan.
montana, Linn. Sp. Pl. 90 = A. montana.
Petteri, Reichb. ex Nym. Consp. 118 = A. globulosa.
ramosissima, Schur, Enum. Pl. Transs. 112 = A. ramosissima.
rostrata, Reichb. Ic. Fl. Germ. v. 28 = A. rostrata.
tenuifolia, Nees, ex Mart. Hort. Erlang. 44 = A. mucronata.

MINURIA, DC. Prod. v. 298 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 267.
ELACHOTHAMNUS, DC. Prod. v. 398 (1836).
KIPPISTIA, F. Muell. Rep. Babb. Exped. 12 (1858).
THEROGERON, DC. Prod. v. 283 (1836).
asteroidea, Sond. in Linnaea, xxv. (1852) 467 = leptophylla.
Candollei, *F. Muell. Fragm.* ix. 119.—Austral.
Cunninghami, *Benth. Fl. Austral.* iii. 498.—Austral.
denticulata, *Benth. l. c.* 499.—Austral.
integerrima, *Benth. l. c.*—Austral.
leptophylla, *DC. Prod.* v. 298.—Austral.
suaedifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 499.—Austral.
tenuissima, DC. Prod. v. 298 = leptophylla.

MINUROTHAMNUS, DC. Prod. vii. 286 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 331.
phagnaloides, *DC. l. c.*—Afr. austr.

MINUTALIA, Fenzl, in Flora, xxvii. (1844) 312, nomen = **Antidesma**, Linn. (Euphorb.).
tomentosa, Fenzl, l. c. = A. venosum.

MINUTIA, Vell. Fl. Flum. 19 (1825); i. t. 47 (1827) = **Linociera**, Sw. (Oleac.).
trichotoma, Vell. ll. cc. = L. mandioccana?

MINYRANTHES, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 180 = **Sigesbeckia**, Linn. (Compos.).
heterophylla, Turcz. l. c. 181 = S. orientalis.

MINYTHODES, Phil. ex Benth. & Hook. f. Gen. ii. 496 (1873) = **Chaetanthera**, Ruiz & Pav. (Compos.).

MIOCARPUS, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 146 = **Acisanthera**, P. Br. (Melastom.).
paludosa, Naud. l. c. 147 = A. limnobios.

MIONANDRA, Griseb. in Goett. Abh. xix. (1874) 101. *MALPIGHIACEAE*.
argentea, *Griseb. l. c.* 102.—Reg. Argent.
camareoides, *Griseb. l. c.*—Reg. Argent.

MIOPTRILA, Rafin. Am. Man. Mulb. 37 (1839) = **Zanthoxylum**, Linn. (Rutac.).
odorata, Rafin. l. c. = Z. americanum.

MIQUELIA, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 177 = **Garnotia**, Brongn. (Gramin.).
barbulata, Nees, l. c. = G. stricta.
courtallensis, Arn. & Nees, l. c.—Ind. or.
Emodi, Arn. & Nees, l. c.—Reg. Himal.

MIQUELIA, Blume, in Bull. Sc. Phys. et Nat. Néerl. i. (1838) 94 = **Staüranthera**, Benth. (Gesnerac.).
azurea, Blume, Rumphia, iv. 34 = S. ecalcarata.
caerulea, Blume, in Bull. Sc. Phys. et Nat. Néerl. i. (1838) 94 = S. ecalcarata.

MIQUELIA, Meissn. Gen. 152 (1838). *OLACINEAE*, Benth. & Hook. f. i. 354.
JENKINSIA, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 231. t. 12.
assamica, *Mast. in Hook. f. Fl. Brit. Ind.* i. 593.—Reg. Himal.
cancellata, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 201.—Malacca.
celebica, *Blume, Rumphia*, vi. 371.—Ins. Celebes.
Cumingii, *Baill. Adansonia*, x. (1871–72) 278.—Ins. Philipp.
dentata, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 38 = Kleinii?
gibba, *Baill. Adansonia*, x. (1871–72) 278.—Ind. or.
Kleinii, Mast. in Hook. f. Fl. Brit. Ind. i. 593 = assamica.
Kleinii, *Meissn. Gen. Comm.* 109.—Ind. or.

MIRA, Colenso, Excurs. N. Island, 84 (1844) sphalm. = MIDA, R. Cunn. = **Fusanus**, R. Br. (Santal.).

MIRABELLIA, Bert. ex Baill. Etud. Gén. Euph. 436 (1858) = **Dysopsis**, Baill. (Euphorb.).
glechomoides, Bert. ex Baill. l. c. = D. Gayana.

MIRABILIS, Riv. ex Linn. Syst. ed. I (1735). *NYCTAGINEAE*, Benth. & Hook. f. iii. 3.
NYCTAGO, Juss. Gen. 90 (1789).
QUAMOCLIDION, Choisy, in DC. Prod. xiii. ii. 429 (1849).
aggregata, Cav. Ic. v. 22 = Oxybaphus aggregatus.
ambigua, Trautv. Del. Sem. Hort. Kiov. (1840) 6.—Cf. Linnaea, xv. (1841) Litt. 97 = M. Jalapa.
californica, A. Gray, in Torr. Bot. Mex. Bound. 173 = Oxybaphus californicus.
coccinea, *Benth. & Hook. f. Gen.* iii. 3.—Am. bor. occ.
cordata, Brouss. ex Roem. & Schult. Syst. iv. 2 = hybrida.
corymbosa, Cav. Ic. iv. 55. t. 379 = Oxybaphus glabrifolius.
dichotoma, Gater. Pl. Montaub. 46 = M. Jalapa.
dichotoma, *Linn. Syst.* ed. X. 931.—Mexic.
divaricata, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 17.—Ins. Madera.
Froebelii, *Greene, in Bull. Calif. Acad.* i. iii. (1885) 124.—Calif.
Greenei, *S. Wats. in Proc. Am. Acad.* xii. (1877) 253.—Calif.
hybrida, *Lepel. in Ann. Mus. Par.* viii. (1806) 481.—Mexic.; N. Granat.
Jalapa, *Linn. Sp. Pl.* 177.—Am. trop.
Jalappa, Thunb. Fl. Jap. 91 = praec.
longiflora, Blanco, Fl. Filip. ed. I. 77 = M. Jalapa.
longiflora, *Linn. in Vet. Akad. Handl. Stockh.* (1755) 176. t. 6. f. 1.—Mexic.
multiflora, *A. Gray, in Torr. Bot. Mex. Bound.* 173.—Am. bor. occ.
odorata, Hort.—Cf. Linnaea, xii. (1838) Litt. 75 = longiflora.
odorata, Linn. Cent. Pl. i. 7; Amoen. Acad. iv. 267 = dichotoma.
oxybaphoides, *A. Gray, in Torr. Bot. Mex. Bound.* 173.—Am. bor. occ.
parviflora, Targ.-Tozz. ex Steud. Nom. ed. I. 535 = dichotoma.
pedunculata, Stokes, Bot. Mat. Med. i. 311 = M. Jalapa
planiflora, Trautv. in Bull. Sc. Acad. Pétersb. vi. (1840) 215 = M. Jalapa.
procera, Bertol. in Nov. Comm. Bonon. iii. (1839) 15 = M. Jalapa.
pubescens, Zipp. ex Span. in Linnaea, xv. (1841) 342 = M. Jalapa.
suaveolens, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 141 = M. Jalapa.
suaveolens, H. B. & K. Nov. Gen. et Sp. ii. 213 = longiflora.
tenuiloba, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 375.—Calif.
triflora, *Benth. Pl. Hartw.* 23.—Mexic.
undulata, Steud. Nom. ed. II. i. 796 = dichotoma.
uniflora, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 34.—Bras.
viscosa, Cav. Ic. i. 13 = Oxybaphus viscosus.
Xalapa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = M. Jalapa.

MIRADORIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 407 (1873) = **Microspermum**, Lag. (Compos.).
Sartorii, Sch. Bip. l. c. = Microspermum debile.

MIRASOLIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 367 (1873) = **Gymnolomia**, H. B. & K. (Compos.).
calva, Benth. & Hook. f. l. c. 368 = G. calva.
diversifolia, Hemsl. Biol. Centr. Am. Bot. ii. 168 = Tithonia diversifolia.
scaberrima, Benth. & Hook. f. Gen. ii. 368 = G. platylepis.

MIRBELIA, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 511. *LEGUMINOSAE*, Benth. & Hook. f. i. 468.
DICHOSEMA, Benth. in Enum. Pl. Hueg. 35 (1837).
OXYCLADIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 20.
angustifolia, R. Grah. in Edinb. N. Phil. Journ. xxvi. (1838) 197 = reticulata.
aotoides, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 53.—Austral.
aphylla, F. Muell. Fragm. iv. 11 = daviesioides.
aspera, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 281 = ovata.
Baxteri, Lindl. Bot. Reg. t. 1434 = Oxylobium scandens.
cordata, Pritz, Ic. Ind. i. 718 = Daviesia cordata.
daviesioides, *Benth. Fl. Austral.* ii. 37.—Austral.
dilatata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 21.—Austral.
floribunda, *Benth. in Lindl. Swan River App.* 12.—Austral.
grandiflora, *Ait. ex Hook. Bot. Mag.* t. 2771.—Austral.
ilicifolia, *Hort. ex Regel, Cat. Pl. Hort. Aksakov.* 94, nomen.—Austral.
Meissneri, Hook. Bot. Mag. t. 4419 = dilatata.
microphylla, *Benth. Fl. Austral.* ii. 37.—Austral.
multicaulis, *Benth. l. c.*—Austral.
ovata, *Meissn. in Lehm. Pl. Preiss.* i. 77.—Austral.
oxyclada, *F. Muell. Fragm.* iv. 12.—Austral.
oxylobioides, *F. Muell. l. c.* ii. 154.—Austral.
pulchella, Meissn. in Lehm. Pl. Preiss. ii. 221 = floribunda.
pungens, *A. Cunn. ex G. Don, Gen. Syst.* ii. 126.—Austral.
racemosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 282.—Austral.
reticulata, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 511.—Austral.
rubiaefolia, G. Don, Gen. Syst. ii. 126 = reticulata.
speciosa, *Sieber, ex DC. Prod.* ii. 115.—Austral.
speciosa, Sweet, Fl. Austral. t. 34 = floribunda.
spinosa, *Benth. Fl. Austral.* ii. 36.—Austral.
subcordata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 282.—Austral.

MIRICA, Nocca, in Usteri, Ann. Bot. v. (1793) 17 = **Myrica**, Linn.

MIRMECODIA, Gaudich. in Freyc. Voy. Bot. 472 (1826) = **Myrmecodia**, Jack (Rubiac.).

MIROBALANUS, Steud. Nom. ed. II. ii. 151 (1841) = MYROBALANUS, Gaertn. = **Terminalia**, Linn. (Combret.).

MIROXILUM, Blanco, Fl. Filip. ed. I. 813 (1837) = MYROXYLON, Forst. = **Xylosma**, Forst. (Bixin.).

MIROXYLON, Scop. Introd. 267 (1777) = praec.

MIROXYLUM, Blanco, Fl. Filip. ed. I. 812 (1837) = praec.

MISANDRA, Comm. ex Juss. Gen. 405 (1789) = **Gunnera**, Linn. (Halorag.).
lobata, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 193 = G. lobata.
magellanica, J. F. Gmel. Syst. 76 = G. magellanica.
prorepens, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 193 = G. prorepens.

MISANDRA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 102 (1819) = **Tillandsia**, Linn. (Bromel.).
juncea, F. G. Dietr. l. c. 103 = F. setacea.
strobilantha, F. G. Dietr. l. c. 104 = T. strobilantha.

MISANDROPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 192 = **Gunnera**, Linn. (Halorag.).
monoica, Oerst. l. c. = G. monoica.

MISANORA, Urv. in Mém. Soc. Linn. Par. iv. (1826) 621, sphalm. = MISANDRA, Comm. = **Gunnera**, Linn. (Halorag.).

MISANTECA, Cham. & Schlecht. in Linnaea, vi. (1831) 367. *LAURINEAE*, Benth. & Hook. f. iii. 155.
SYMPHYSODAPHNE, A. Rich. Fl. Cub. Fanerog. ii. 190. t. 67 (1853).
anacardioides, *Benth. in Hook. Ic. Pl.* t. 1259.—Bras.
capitata, *Cham. & Schlecht. in Linnaea*, vi. (1831) 367.—Mexic.
crassiramea, *Benth. in Hook. Ic. Pl.* xiii. 47, in obs.—Bras.
cubensis, *Benth. l. c.*—Cuba.

MISCANTHUS, Anderss. in Oefvers. Vet. Akad. Stockh. (1855) 165. *GRAMINEAE*, Benth. & Hook. f. iii. 1125.
ECCOILOPUS, Steud. Syn. Pl. Gram. 123 (1855).
EULALIA, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 332, partim.
capensis, Anderss. in Oefvers. Vet. Akad. Stockh. (1855) 165 = Erianthus capensis.
cotulifer, *Benth. in Journ. Linn. Soc.* xix. (1881) 65.—Japon.
fuscus, Benth. l. c. = Saccharum fuscum.
japonicus, *Anderss. in Oefvers. Vet. Akad. Stockh.* (1855) 166.—As. or.; Polynes.
luzonensis, Anderss. l. c. = japonicus.
purpurascens, *Anderss. l. c.* 167.—Japon.
saccharifer, *Benth. in Journ. Linn. Soc.* xix. (1881) 65 (sphalm. sacchariflorus).—As. or.; Japon.
sinensis, *Anderss. in Oefvers. Vet. Akad. Stockh.* (1855) 166.—China; Japon.

MISCHANTHUS, Coss. in Bull. Soc. Bot. Fr. iv. (1857) 14 = **Mecomischus**, Coss. (Compos.).

MISCHANTHUS, Ind. Kew. supra, i. 912, in syn., sphalm. = **Miscanthus**, Anderss. (Gramin.).

MISCHOCARPUS, Blume, Bijdr. 238 (1825); Rumphia, iii. 166 (1847) = **Ratonia**, DC. (Sapind.).
anodontus, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 647 = R. anodonta.
exangulatus, Radlk. l. c. = R. exangulata.
fuscescens, Blume, Rumphia, iii. 169 = Cupania Helferi.
grandissimus, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 647 = R. grandissima.
lachnocarpus, Radlk. l. c. = R. lachnocarpa.
pentapetalus, Radlk. l. c. 646 = Cupania pentapetala.
pyriformis, Radlk. l. c. 647 = R. pyriformis.
sumatranus, Blume, Rumphia, iii. 168 = Cupania sumatrana.
sundaicus, Blume, Bijdr. 238 = Cupania Lessertiana.

MISCHODON, Thw. in Hook. Kew Journ. vi. (1854) 299. t. 10. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 280.
zeylanicus, *Thw. l. c.* 300.—Zeylan.

MISCHOPHLOEUS, Scheff. in Ann. Jard. Buitenz. ii. (1876) 115. *PALMAE*, Benth. & Hook. f. iii. 883.
paniculata, *Scheff. l. c.* 152.—Ins. Ternate.

MISCHOSPORA, Boeck. in Flora, xliii. (1860) 113 = **Fimbristylis**, Vahl. (Cyperac.).
efoliata, Boeck. l. c. = F. tetragona.

MISCODENDRUM, Steud. Nom. ed. II. ii. 151 (1841) = **Myzodendron**, Banks (Santal.).

MISCOLOBIUM, Vog. in Linnaea, xi. (1837) 200 = **Dalbergia**, Linn. (Legumin.).
densiflorum, Benth. in Ann. Wien. Mus. ii. (1838) 104 = D. densiflora.
divaricatum, Benth. l. c. = D. densiflora.
glandulosum, Benth. in Mart. Fl. Bras. xv. i. 223 = D. glandulosa.
glaucescens, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 103 = D. glaucescens.
lateriflorum, Benth. in Mart. Fl. Bras. xv. i. 225 = D. lateriflora.
nigrum, Allem. in Trab. Soc. Vell. 66 = D. nigra.
nigrum, Mart. Herb. Fl. Bras. n. 1126 = D. Miscolobium.
parviflorum, Benth. in Mart. Fl. Bras. xv. i. 242 = Machaerium parviflorum.

MISCOLOBIUM:—
polyphyllum, Benth. in Ann. Wien. Mus. ii. (1838) 103 = Dalbergia foliolosa.
scandens, Benth. in Mart. Fl. Bras. xv. i. 226 = D. gracilis.
Spruceanum, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 35 = D. Spruceana.
villosum, Benth. in Ann. Wien. Mus. ii. (1838) 104 = villosa.
violaceum, Vog. in Linnaea, xi. (1837) 201 = D. Miscolobium.

MISCOPETALUM, Haw. Syn. Pl. Succ. 323 (1812) = **Saxifraga**, Linn.
cernuum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 530 = S. cernua.
rotundifolium, Haw. Syn. Pl. Succ. 323 = S. rotundifolius.

MISIPUS, Rafin. Sylva Tellur. 60 (1838) = **Elaeocarpus**, Linn. (Tiliac.).
serratus, Rafin. l. c. = E. serratus.

MISODENDRON, G. Don, Gen. Syst. iii. 408 (1834) = seq.

MISODENDRUM, DC. Coll. Mém. vi. 14. tt. 11, 12 (1830); et Prod. iv. 285 (1830) = **Myzodendron**, Banks. (Santal.).

MISSIESSIA, Benth. & Hook. f. Gen. iii. 391 (1880) = seq.

MISSIESSYA, Gaudich. Bot. Voy. Bonite, t. 93 (1853?) = **Leucosyke**, Zoll. (Urtic.).
alba, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 195 = L. alba.
aspera, Wedd. l. c. = L. alba.
bimensis, Wedd. l. c. = L. alba.
bornensis, Blume, Mus. Bot. Lugd. Bat. ii. 164 = L. bornensis.
candidissima, Blume, l. c. 165 = L. javanica.
celebica, Blume, l. c. 165 = L. celebica.
celtidifolia, Gaudich. Bot. Voy. Bonite, t. 93 = L. alba.
corymbulosa, Wedd. in Arch. Mus. Par. viii. (1855–56) 476 = L. corymbulosa.
elongata, Blume, Mus. Bot. Lugd. Bat. ii. 163 = L. alba.
fagifolia, Gaudich. Bot. Voy. Bonite, t. 93 = L. alba.
falcata, Blume, Mus. Bot. Lugd. Bat. ii. 164 = L. alba.
hispidissima, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 195 = L. hispidissima.
hypoleuca, Wedd. l. c. = Debregeasia hypoleuca.
javanica, Wedd. l. c. = L. javanica.
mutabilis, Blume, Mus. Bot. Lugd. Bat. ii. 163 = L. alba.
ochroneura, Blume, l. c. 165 = L. ochroneura.
parvifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 195 = Debregeasia edulis.
rubiginosa, Blume, Mus. Bot. Lugd. Bat. ii. 164 = L. rubiginosa.
sumatrana, Blume, l. c. 165 = L. sumatrana.
velutina, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 195 = Debregeasia velutina.
villosa, Blume, Mus. Bot. Lugd. Bat. ii. 163 = L. alba.
Wallichii, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 195 = Debregeasia Wallichiana.

MISTRALIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 147 = **Daphne**, Linn. (Thymel.).
Gnidium, Fourr. l. c. = D. Gnidium.

MISTYLLUS, Presl, Symb. Bot. i. 49 (1830) = **Trifolium**, Tourn. (Legumin.).
spumosus, Presl, l. c. = T. spumosum.
turgidus, Presl, l. c. = T. versiculosum.

MITCHELLA, Linn. Diss. Chen. 24 (1751); Amoen. Acad. iii. 16 (1756). *RUBIACEAE*, Benth. & Hook. f. ii. 137.
DISPERMA, J. F. Gmel. Syst. 892 (1791).
GEOHERPUM, Willd. ex Schult. Mant. iii. 18, 350 (1827).
ovata, *DC. Prod.* iv. 452.—Ecuador.
pilosa, Benth. Pl. Hartw. 194 = Corynula pilosa.
repens, *Linn. Sp. Pl.* 111.—Am. bor.; Japon.
undulata, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 175 = repens.

MITCHERLICHIA, Klotzsch, in Monatsb. Berl. Acad. (1854) 124 = MITSCHERLICHIA, Klotzsch = **Begonia**, Linn.
albo-coccinea, Klotzsch, l. c. = B. albo-coccinea.

MITELLA, Tourn. ex Linn. Syst. ed. I (1735). *SAXIFRAGEAE*, Benth. & Hook. f. i. 638.
DRUMMONDIA, DC. Prod. iv. 49 (1830).
MITELLOPSIS, Meissn. Gen. 136 (1838).
OZOMELIS, Rafin. Fl. Tellur. ii. 73 (1836).
PECTIANTIA, Rafin. l. c. 72 (1836).
acerifolia, Schlecht. ex Ledeb. Fl. Ross. ii. 228 = Tellima grandiflora.
Breweri, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 533.—Calif.
caulescens, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 586.—Am. bor. occ.
cordifolia, Lam. Encyc. iv. 196 = nuda.
diphylla, *Linn. Sp. Pl.* 406.—Am. bor.
grandiflora, Pursh, Fl. Am. Sept. i. 314 = Tellima grandiflora.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 96; *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 431.—Japon.
nuda, *Linn. Sp. Pl.* 406.—Sibir.; Am. bor.
parviflora, D. Dietr. Syn. Pl. ii. 1539 = Tellima parviflora.
pentandra, *Hook. Bot. Mag.* t. 2933.—Am. bor. occ.
prostrata, Michx. Fl. Bor. Am. i. 270 = nuda.
reniformis, Lam. Encyc. iv. 196 = nuda.
trifida, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.–June, 1829) 185.—Am. bor. occ.

MITELLOPSIS, Meissn. Gen. 136 (1830) = **Mitella**, Tourn. (Saxifr.).
Drummondia, Meissn. Gen. Comm. 100 = Mitella pentandra.
Hookeri, Meissn. l. c. = Mitella trifida.
pentandra, Walp. Rep. ii. 370 = Mitella pentandra.
trifida, Walp. l. c. = Mitella trifida.

MITESIA, Rafin. New Fl. Am. iv. 47 (1836); Fl. Tellur. iii. 15 (1836) = **Polygonum**, Tourn.
albiflora, Rafin. Fl. Tellur. iii. 15.—Hab.?
divergens, Rafin. New Fl. Am. iv. 48; Fl. Tellur. iii. 15.—Am. bor.
floridana, Rafin. New Fl. Am. 47.—Am. bor.
hirsuta, Rafin. l. c. 48; Fl. Tellur. iii. 15 = P. hirsutum.
ligularis, Rafin. New Fl. Am. iv. 47.—Am. bor.
montana, Rafin. Fl. Tellur. iii. 15.—Hab.?
punctata, Rafin. l. c. = P. punctatum.

MITHRACARPUS, Reichb. Consp. 94 (1828) = **Mitracarpum**, Zucc. (Rubiac.).

MITHRIDATEA, Comm. ex Schreb. Gen. 783 (1791) = **Tambourissa**, Sonner. (Monim.).
amplifolia, Boj. Hort. Maurit. 290 = T. amplifolia.
cymosa, Willd. ex Steud. Nom. ed. II. ii. 151, nomen; Tul. in Arch. Mus. Par. viii. (1855–56) 306 (Rubiacea).—Hab.?
quadrifida, Willd. Sp. Pl. i. 27 = T. quadrifida.
Tambourissa, Boj. Hort. Maurit. 290 = T. quadrifida.

MITHRIDATIUM, Adans. Fam. ii. 48 (1763) = **Erythronium**, Linn. (Liliac.).

MITINA, Adans. Fam. ii. 116 (1763) = **Carlina**, Linn. (Compos.).
corymbosa, Cass. in Dict. Sc. Nat. xlvii. 508 = C. corymbosa.
lanata, Cass. l. c. 507 = C. lanata.
sulphurea, Cass. l. c. 508 = C. racemosa.

MITODENDRON, Walp. Ann. i. 360 (1849), sphalm. = **Myzodendron**, Banks (Santal.).

MITOLEPIS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 78. *ASCLEPIADEAE*.
intricata, *Balf. f. l. c.*—Ins. Socotra.

MITOPETALUM, Blume, Praef. Fl. Jav. 8 (1828) = **Tainia**, Blume (Orchid.).
angustifolium, Blume, Mus. Bot. Lugd. Bat. ii. 185 = T. angustifolia.
bicorne, Blume, l. c. = T. bicornis.
fimbriatum, Miq. Fl. Ind. Bat. iii. 674 = T. fimbriata.

MITOPETALUM :—
latifolium, Blume, Mus. Bot. Lugd. Bat. ii. 185 = Tainia latifolia.
parviflorum, Blume, Orch. Archip. Ind. 160, t. 50. f. 2 = trinerve.
plicatum, Blume, Mus. Bot. Lugd. Bat. ii. 185.—Java.
rubescens, Blume, Orch. Archip. Ind. 159. t. 51 = *plicatum*.
speciosum, Blume, Praef. Fl. Jav. 8 = T. speciosa.
trinerve, Blume, Mus. Bot. Lugd. Bat. ii. 185 = T. trinervis.
trinervium, Blume, Orch. Archip. Ind. 160 = praec.
Wightii, Blume, Mus. Bot. Lugd. Bat. ii. 185 = T. bicornis.

MITOSTAX, Rafin. Sylva Tellur. 120 (1838) = **Prosopis**, Linn. (Legumin.).
pallida, Rafin. l. c. = P. juliflora.

MITOSTEMMA, Mast. in Journ. Bot. xxi. (1883) 33. *PASSIFLOREAE*.
Glaziovii, *Mast. l. c.* 34.—Bras.
Jenmanii, *Mast. l. c.*—Guian.

MITOSTIGMA, Blume, Mus. Bot. Lugd. Bat. ii. 189 (1856) = **Habenaria**, Willd. (Orchid.).
gracile, Blume, l. c. 190 (= *Gymnadenia gracilis*).—Japon.

MITOSTIGMA, Decne. in DC. Prod. viii. 507 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
affine, *Griseb. in Goett. Abh.* xxiv. (1879) 225.—Reg. Argent.
cionophorum, *Griseb. l. c.* 226.—Reg. Argent.
latiflorum, *Griseb. l. c.*—Reg. Argent.
niveum, *Griseb. l. c.*—Reg. Argent.
rhynchophorum, *Griseb. l. c.*—Reg. Argent.
tomentosum, *Decne. in DC. Prod.* viii. 507.—Reg. Argent.

MITOSTYLIS, Rafin. Sylva Tellur. 114 (1838) = **Cleome**, Linn. (Cappar.).
procumbens, Rafin. l. c. = C. procumbens.

MITOZUS, Miers, Apocyn. S. Am. 217 (1878) = **Echites**, R. Br. (Apocyn.).
Blanchetii, Miers, l. c. 219 = E. Blanchetii.
brachystachys, Miers, l. c. 222 = E. brachystachya.
brevipes, Miers, l. c. 223 = E. brevipes.
concinnus, Miers, l. c.—Bras.
cuyabensis, Miers, l. c. = E. cuyabensis.
discolor, Miers, l. c. 224 = E. discolor.
exilis, Miers, l. c. 218 = E. funiformis.
funiformis, Miers, l. c. 219 = E. funiformis.
gracilipes, Miers, l. c. 220 = E. gracilipes.
guanabaricus, Miers, l. c. 218 = E. Velami.
jamaicensis, Miers, l. c. 225 = E. jamaicensis.
leptophyllus, Miers, l. c. 220 = E. leptophylla.
mexicanus, Miers, l. c. 225 = Prestonia mexicana.
microphyllus, Miers, l. c. 219 = E. microphylla.
rugosus, Miers, l. c. 222 = E. rugosa.
scabridulus, Miers, l. c. 224.—Bras.
symphytocarpus, Miers, l. c. 222 = E. symphytocarpa.
tenellus, Miers, l. c. 220 = Odontadenia angustifolia.
tenuicaulis, Miers, l. c. 221 = E. tenuicaulis.
versicolor, Miers, l. c. = E. versicolor.

MITRA, Houst. ex Linn. Gen. ed. I. 377 (1737) = **Mitreola**, Linn. (Logan.).

MITRACARPIUM, Benth. in Hook. Journ. Bot. iii. (1841) 238 = seq.

MITRACARPUM, Zucc. in Schult. Mant. iii. 210, in obs. (1827). *RUBIACEAE*, Benth. & Hook. f. ii, 146.
CRUSEA, Cham. ex DC. Prod. iv. 571 (1830).
MITHRACARPUS, Reichb. Consp. 94 (1828).
SCHIZANGIUM, Bartl. ex DC. Prod. iv. 571 (1830).
STAUROSPERMUM, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 73 (1827).
ampliatum, Hochst. in Flora, xxv. (1842) I. Beibl. 135, nomen ; et ex A. Rich. Tent. Fl. Abyss. i. 349 = Spermacoce ampliata.
breviflorum, *A. Gray, Pl. Wright.* ii. 68.—Arizona.
crassifolium, *A. Rich. Fl. Cub. Fanerog.* ii. 31.—Cuba.
cuspidatum, DC. Prod. iv. 572 = Sellowianum.
diffusum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 363.—Am. austr.

MITRACARPUM :—
diodioides, *A. Rich. Fl. Cub. Fanerog.* ii. 31.—Cuba.
discolor, Miq. in Linnaea, xviii. (1844) 616 = Humboldtianum.
Dregeanum, *E. Mey. in Drège, Zwei Pfl. Docum.* 202, nomen ; *et ex Sond. in Harv. & Sond. Fl. Cap.* iii. 25.—Afr. austr.
Fischeri, *Cham. & Schlecht. in Linnaea*, iii. (1828) 363.—Jamaic.
hirtum, DC. Prod. iv. 572 = villosum.
Humboldtianum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 358.—Am. austr.
laeteviride, *Wright, in Sauv. Fl. Cub.* 73.—Cuba.
Lhotzkianum, *Cham. in Linnaea*, ix. (1834) 219.—Bras.
lineare, *Benth. Bot. Voy. Sulph.* 20.—Calif.
linearifolium, *A. Rich. Fl. Cub. Fanerog.* ii. 31.—Cuba.
neglectum, DC. Prod. iv. 573 = Sellowianum.
pallidum, *Hook. & Arn. Bot. Beech. Voy.* 430.—Mexic.
Peladilla, *Griseb. in Goett. Abh.* xxiv. (1879) 158.—Reg. Argent.
pilosum, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 151. t. 14. f. 4 = villosum ?
puberulum, Benth. in Hook. Journ. Bot. iii. (1841) 238 = villosum.
rude, Benth. l. c. = Humboldtianum.
Sagraeanum, *DC. Prod.* iv. 572.—Cuba.
Salzmannianum, DC. l. c. 571 = Humboldtianum.
scabrellum, *Benth. in Hook. Journ. Bot.* iii. (1841) 238.—Bras.
scabrum, *Zucc. in Schult. Mant.* iii. 210.—Afr. trop.
Schizangium, *DC. Prod.* iv. 572.—Mexic.
Sellowianum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 361.—Bras.
senegalense, DC. Prod. iv. 572 = scabrum.
sphaerostigma, Hochst. in Flora, xxiv. (1841) I. Intell. 27 = Spermacoce ampliata.
squarrosum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 363.—Cuba.
stylosum, *Cham. & Schlecht. l. c.*—Ins. Philipp.
Torresianum, Cham. & Schlecht. l. c. 360 = villosum.
verticillatum, Vatke, in Linnaea, xl. (1876) 196 = scabrum.
villosum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 363.—Jamaic.
virgatum, *Cham. & Schlecht. l. c.*—Bras.

MITRACARPUS, A. Rich. Tent. Fl. Abyss. i. 348 (1847) = **Mitracarpum**, Zucc. (Rubiac.).

MITRACME, Schult. Mant. iii. 67 (1827) = **Mitrasacme**, Labill. (Logan.).

MITRAGYNA, Korth. Obs. Naucl. Ind. 19 (1839). *RUBIACEAE* [Benth. & Hook. f. ii. 31].
MAMBOGA, Blanco, Fl. Filip. ed. I. 140 (1837).
STEPHEGYNE, Korth. Verh. Nat. Gesch. Bot. 160 (1840 ?).
africana, *Korth. Obs. Naucl. Ind.* 19.—Afr. trop.
macrophylla, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 41.—Afr. trop.
parvifolia, *Korth. Obs. Naucl. Ind.* 19.—Reg. Himal.
speciosa, *Korth. l. c.*—Borneo.

MITRAGYNE, R. Br. Prod. 452 (1810) = **Mitrasacme**, Labill. (Logan.).

MITRANTHES, Berg, in Linnaea, xxvii. (1854) 316 = **Calyptranthes**, Sw. (Myrtac.).
Browniana, Berg, in Mart. Fl. Bras. xiv. I. 355.—Bras.
eugenioides, Berg, l. c. = C. eugenioides.
Gardneriana, Berg, l. c. 354.—Bras.
Langsdorffii, Berg, l. c. 595.—Bras.
Ottonis, Berg, in Linnaea, xxvii. (1854) 316 = C. Ottonis.
ovalifolia, Berg, in Mart. Fl. Bras. xiv. I. 356.—Bras.
Riedliana, Berg, l. c. 595.—Bras.
Sartoriana, Berg, in Linnaea, xxix. (1857-58) 248.—Mexic.

MITRANTHUS, Hochst. in Flora, xxvii. (1844) 103 = **Vandellia**, Linn. (Scrophul.).
latifolius, Hochst. l. c. = V. sessiliflora.

MITRARIA, Cav. in Anal. Cienc. Nat. iii. (1801) 230. t. 31. *GESNERACEAE*, Benth. & Hook. f. ii. 1012.
DIPLOCALYX, Presl, Bot. Bemerk. 146 (1844).
coccinea, *Cav. in Anal. Cienc. Nat.* iii. (1801) 231.—Ins. Chiloe.
pallida, *Hanst. in Linnaea*, xxxiv. (1865-66) 419.—Mexic.
tomentella, *Hanst. l. c.* 420.—Mexic.

MITRARIA, J. F. Gmel. Syst. 799 (1791) = **Barringtonia**, Forst. (Myrtac.).
Commersonia, J. F. Gmel. l. c. = B. Butonica.

MITRASACME, Labill. Nov. Holl. Pl. i. 35. t. 49 (1804), *LOGANIACEAE*, Benth. & Hook. f. ii. 790.
MITRACME, Schult. Mant. iii. 67 (1827).
MITRAGYNE, R. Br. Prod. 452 (1810).
alsinoides, *R. Br. Prod.* 453.—As. or. ; Austral.
ambigua, *R. Br. l. c.* 454.—Austral.
Archeri, *Hook. f. Fl. Tasman.* ii. 368.—Tasmania.
canescens, R. Br. Prod. 453 = polymorpha.
capillaris, Wall. in Roxb. Fl. Ind. ed. Carey, i. 420 = polymorpha.
Cheesemanii, J. Buch. in Trans. N. Z. Inst. xiv. (1882) 348. t. 29 (Veronicae sp.).—N. Zel.
chinensis, Griseb. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 51 = nudicaulis.
cinerascens, R. Br. Prod. 452 = polymorpha.
connata, *R. Br. l. c.* 454.—Austral.
constricta, F. Muell. Fragm. i. 131 = connata.
crystallina, Griff. Notul. iv. 87 = alsinoides.
Cunninghamii, *Benth. Fl. Austral.* iv. 357.—Austral.
diffusa, Benth. in Journ. Linn. Soc. i. (1857) 93 = paludosa.
distylis, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 20.—Austral.
divergens, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 272 = paradoxa.
elata, *R. Br. Prod.* 453.—Austral.
elata, F. Muell. Fragm. i. 132 = longiflora.
exserta, *F. Muell. l. c.* 131.—Austral.
gentianea, *F. Muell. l. c.* 130.—Austral.
hirsuta, Presl, Bot. Bemerk. 104 = polymorpha
Hookeri, J. Buch. in Trans. N. Z. Inst. xiv. (1882) 348. t. 29 (Veronicae sp.).—N. Zel.
indica, Wight, Ic. Pl. t. 1601 = alsinoides.
laevis, *Benth. in Journ. Linn. Soc.* i. (1857) 93.—Austral.
laricifolia, *R. Br. Prod.* 453.—Austral.
longiflora, *F. Muell. ex Benth. Fl. Austral.* iv. 353.—Austral.
lutea, *F. Muell. Fragm.* i. 133.—Austral.
malaccensis, Wight, Ic. t. 1601 = polymorpha.
montana, *Hook. f. ex Benth. in Journ. Linn. Soc.* i. (1857) 93.—Austral.
multicaulis, *R. Br. Prod.* 453.—Austral.
nuda, Nees, in Lehm. Pl. Preiss. ii. 239 = paradoxa.
nudicaulis, *Reinw. ex Blume, Bijdr.* 849.—As. or. ; Austral.
paludosa, *R. Br. Prod.* 453.—Austral.
paradoxa, *R. Br. l. c.* 454.—Austral.
perpusilla, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 275 = serpyllifolia.
Petriei, J. Buch. in Trans. N. Z. Inst. xiv. (1882) 349. t. 30. f. 1 (Veronicae sp.).—N. Zel.
phascoides, *R. Br. Prod.* 454.—Austral.
pilosa, *Labill. Nov. Holl. Pl.* i. 36. t. 49.—Austral.
pilosula, F. Muell. Fragm. i. 134 = paludosa.
polymorpha, *R. Br. Prod.* 452.—As. or. ; Austral.
prolifera, *R. Br. l. c.* 453.—Austral.
prolifera, Sieber, ex Schult. Mant. iii. 67 = polymorpha.
pusilla, Dalz. in Hook. Kew Journ. ii. (1850) 136 = alsinoides.
pygmaea, *R. Br. Prod.* 453.—Austral.
ramosa, R. Br. l. c. = multicaulis.
serpyllifolia, *R. Br. l. c.* 454.—Austral.
setosa, *Hance, in Journ. Bot.* xv. (1877) 335.—Cambodia.
Sieberi, A. DC. in DC. Prod. ix. 10 = polymorpha.
squarrosa, R. Br. Prod. 452 = polymorpha.
stellata, *R. Br. l. c.* 454.—Austral.
subvolubilis, F. Muell. Fragm. i. 133 = laevis.
tenuiflora, *Benth. Fl. Austral.* iv. 354.—Austral.
trinervis, Span. in Linnaea, xv. (1841) 335 = polymorpha.
villosa, Pritz. Ic. Ind. i. 719, sphalm. = pilosa.

MITRASTIGMA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 20 = **Plectronia**, Linn. (Rubiac.).
lucidum, Harv. l. c. (= *Canthium obovatum*).—Afr. austr.

MITRELLA, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 38 (1865-66) = **Melodorum**, Dun. (Anonac.).
Kentii, Miq. l. c. 39.—Sumatra; Borneo.

MITREOLA, Linn. Gen. ed. I. 377 (1737). *LOGANIACEAE*, Benth. & Hook. f. ii. 790.
MITRA, Houst. ex Linn. Gen. ed. I. 377 (1737).
PAROPHIORRHIZA, C. B. Clarke, ex Hook. f. Fl. Brit. Ind. iii. 85 (1880).
inconspicua, Zoll. & Mor. Syst. Verz. Zoll. 55 = oldenlandioides.
oldenlandioides, *Wall. Cat.* n. 4350.—As. or.; Austral.
ophiorhizoides, Rich. in Mém. Soc. Hist. Nat. Par. i. (1823) 63. t. 3 = petiolata, sessilifolia.
paniculata, A. DC. in DC. Prod. ix. 9 = petiolata.
paniculata, Wall. Cat. n. 4349 = oldenlandioides.
pedicellata, *Benth. in Journ. Linn. Soc.* i. (1857) 91.—Reg. Himal.
petiolata, *Torr. & Gray, Fl. N. Am.* ii. 45.—Am. bor. et trop.
sessilifolia, *Torr. & Gray, l. c.*—Am. bor.
Swartzii, G. Don, Gen. Syst. iv. 171 = petiolata.

MITREPHORA, Hook. f. & Thoms. Fl. Ind. i. 112 (1855). *ANONACEAE*, Benth. & Hook. f. i. 26.
PSEUDUVARIA, Miq. Fl. Ind. Bat. i. II. 32 (1859).
aperta, *Teijsm. & Binn. Cat. Hort. Bog.* 175, nomen.—Java.
Bousigoniana, *Pierre, Fl. For. Cochinch.* t. 36.—Cochinch.
celebica, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 389.—Ins. Celebes.
Diepenhorstii, *Teijsm. & Binn. Cat. Hort. Bog.* 175, nomen.—Sumatra.
diversifolia, *Miq. Fl. Ind. Bat.* i. II. 32.—Ins. Timor.
Edwardsii, *Pierre, Fl. For. Cochinch.* t. 35.—Cambodia.
excelsa, *Hook. f. & Thoms. Fl. Ind.* i. 114.—Penang.
glabra, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 20.—Borneo.
grandiflora, *Bedd. Fl. Sylv.* i. t. 75.—Ind. or.
Heyneana, *Thw. Enum. Pl. Zeyl.* 8.—Ind. or.
humilis, *Miq. Fl. Ind. Bat.* i. II. 31.—Java.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 28.—Borneo.
longipetala, *Miq. l. c.*—Borneo.
macrantha, *Hassk. Retzia*, i. 235.—Java.
Maingayi, *Hook. f. & Thoms. in Hook. Fl. Brit. Ind.* i. 77.—Malacca.
micrantha, *Teijsm. & Binn. Cat. Hort. Bog.* 176, nomen.—Java.
multiflora, *Miq. Fl. Ind. Bat.* i. II. 32.—Java.
obtusa, *Hook. f. & Thoms. Fl. Ind.* i. 113.—Malaya.
polypyrena, *Miq. Fl. Ind. Bat.* i. II. 32.—Java.
reticulata, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 77.—Burma; Java.
subaequalis, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 20.—N. Guin.
Teysmanni, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 12; *et in Flora*, lii. (1870) 302.—Ins. Banca.
Thorelii, *Pierre, Fl. For. Cochinch.* t. 37.—Cochinch.
tomentosa, *Hook. f. & Thoms. Fl. Ind.* i. 113.—Reg. Himal.
vandaeflora, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 57, 181.—Burma.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 27.—Ins. Amboina.

MITRIOSTIGMA, Hochst. in Flora, xxv. (1842) 235. *RUBIACEAE*, Benth. & Hook. f. ii. 90.
axillare, *Hochst. l. c.* 236.—Afr. austr.
Barteri, *Hook. f. in Benth. & Hook. f. Gen.* ii. 90, nomen; *et ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 111.—Afr. trop.
subpunctatum, *Hiern, l. c.*—Afr. trop.

MITROCARPUM, Hook. Bot. Misc. iii. (1833) 360, sphalm. = **Mitracarpum**, Zucc.

MITROPHORA, Neck. Elem. i. 123 (1790) = **Fedia**, Gaertn. (Valer.).
Cornucopia, Rafin. Chl. Aetn. 5 = F. Cornucopiae.

MITROSICYOS, Maxim. Prim. Fl. Amur. 112 (1859); et in Ann. Sc. Nat. Sér. IV. xiii. (1860) 95 = **Actinostemma**, Griff. (Cucurb.).
lobatus, Maxim. Prim. Fl. Amur. 112. t. 7 = A. lobatum.
paniculatus, Maxim. l. c. 113, in nota = A. paniculatum.
racemosa, Maxim. l. c. 112, in nota = A. racemosum.

MITROSPORA, Nees, in Linnaea, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.).
cephalophora, Nees, in Mart. Fl. Bras. ii. I. 133 = R. cephalophora.
ciliata, Nees, l. c. = R. Neesii.
pallida, Nees, l. c. = R. pallida.
polyphylla, Nees, in Linnaea, ix. (1834) 295 = R. polyphylla.
recurvata, Nees, in Mart. Fl. Bras. ii. I. 132 = R. recurvata.
Wageneriana, Schlecht. in Linnaea, xxvi. (1853) 129.—N. Granat.

MITSA, Chapel. ex Benth. Lab. Gen. et Sp. 52 (1832) = **Coleus**, Lour. (Labiat.).
maculata, Chapel. ex Benth. l. c. = C. Bojeri.

MITSCHERLICHA, Klotzsch, in Abh. Akad. Berl. (1855) 73. t. 6 C. = **Begonia**, Tourn.
albo-coccinea, Klotzsch, l. c. = B. albo-coccinea.
coriacea, Klotzsch, l. c. 74 = B. peltata.
Grahamiana, Hassk. Hort. Bog. Desc. 334 = B. albo-coccinea.
Junghuhniana, Miq. Fl. Ind. Bat. i. I. 696 = B. Junghuhniana.
Lobbii, Hassk. Hort. Bog. Desc. 331 = B. Lobbii.
repens, Miq. Fl. Bat. Ind. Suppl. 333 = B. mollis.
rubro-setulosa, Hassk. Hort. Bog. Desc. 336 = B. rubro-setulosa.
stictopoda, Miq. Fl. Ind. Bat. i. I. 1092 = B. stictopoda.
trichopoda, Miq. l. c. Suppl. 333 = B. trichopoda.

MITSCHERLICHIA, Kunth, in Abh. Akad. Berl. (1831) 209 = **Neea**, Ruiz & Pav. (Nyctag.).
spectabilis, Kunth, l. c. = N. Mitscherlichia.

MITTELLA, Linn. Diss. Chen. 29 (1751) = **Mitella**, Linn. (Saxifr.).

MNASIUM, Schreb. Gen. 214 (1789) = **Rapatea**, Aubl.
paludosum, Poepp. ex Koern. in Linnaea, xxxvii. (1871–73) 449 = Cephalostemon gracilis.
paludosum, Willd. Sp. Pl. ii. 22 = R. paludosa.
sphaerocephalum, Rudge, Pl. Gui. 12 = R. sphaerocephala.
unilaterale, Rudge, l. c, 12. t. 11 = R. unilateralis.

MNASIUM, Stackh. Extr. Bruce's Trav. Abyss. 18. t. 2 (1815) = **Musa**, Linn. (Scitam.).
Theophrasti, Pritz. Ic. Ind. 719 = M. Ensete.

MNASSEA, Vell. Fl. Flum. 188 (1825) (Quid?).
pinnata, Vell. l. c. 188; iv. t. 147.—Bras.

MNEMION, Spach, Hist. Vég. Phan. v. 510 (1836) = **Viola**, Tourn.
agreste, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 341 = V. tricolor.
alpestre, Fourr. l. c. 342 = V. tricolor.
calcaratum, Spach, Hist. Vég. Phan. v. 512 = V. calcarata.
Carioti, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 342.—Gallia.
cheiranthifolium, Webb & Berth. Phyt. Canar. i. 113 = V. cheiranthifolia.
contemptum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 341 = V. tricolor.
cornutum, Spach, Hist. Vég. Phan. v. 512 = V. cornuta.
elegans, Spach, l. c. 513 = V. tricolor.
gracilescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 342 = V. tricolor.
grandiflorum, Spach, Hist. Vég. Phan. v. 514 = V. altaica.
hirsutum, Spach, l. c. = V. rothomagensis.
Lagascae, Webb, Iter Hisp. 68 = V. nevadensis.
nemausense, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 341 = V. tricolor.
obtusifolium, Fourr. l. c.—Gallia.

MNEMION:—
palmense, Webb & Berth. Phyt. Canar. i. 112 = Viola palmensis.
Sagoti, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 342 = V. tricolor.
segetale, Fourr. l. c. 341 = V. tricolor.
sudeticum, Fourr. l. c. = V. tricolor.
tenellum, Webb, Iter Hisp. 68 = V. tricolor.
tricolor, Spach, Hist. Vég. Phan. v. 515 = V. tricolor.
variatum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 341 = V. tricolor.

MNEMOSILLA, Forsk. Fl. Aegypt. Arab. 122 (1775) = **Hypecoum**, Linn. (Papav.).
aegyptiaca, Forsk. l. c. = H. imberbe.

MNESITEON, Rafin. Fl. Ludov. 67 (1817) = **Baldwinia**, Nutt. (Compos.).
album, Rafin. l. c. 68 = B. multiflora.
luteum, Rafin. l. c. = B. multiflora?

MNESITHEA, Kunth, Rév. Gram. i. 153 (1830); et Enum. Pl. i. 465 (1833) = **Ophiurus**, R. Br. (Gramin.).
laevis, Kunth, Rév. Gram. i. 154 = O. perforatus.

MNESITHEON, Spreng. Gen. ii. 635 (1831) = MNESITEON, Rafin. = **Baldwinia**, Nutt. (Compos.).

MNIANTHUS, Walp. Ann. iii. 443 (1853) = **Terniola**, Tul. (Podost.).
longipes, Walp. l. c. = T. longipes.
pulchellus, Walp. l. c. = T. pulchella.
zeylanicus, Walp. l. c. = T. zeylanica.

MNIARUM, Forst. Char. Gen. i. t. 1 (1776) = **Scleranthus**, Linn. (Illecebr.).
biflorum, Forst. l. c. 2. t. 1 = S. biflorus.
fasciculatum, R. Br. Prod. 412 = S. biflorus.
pedunculatum, Labill. Nov. Holl. Pl. i. 8. t. 2 = S. biflorus.
singuliflorum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 13 = S. mniaroides.

MNIODES, A. Gray, ex Benth. & Hook. f. Gen. ii. 301 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
andina, *A. Gray, in Herb. Kew.*—Peruv.

MNIOPSIS, Mart. Nov. Gen. et Sp. i. 3. t. 1 (1822). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 113.
CRENIAS, Spreng. Syst. iv. Cur. Post. 247 (1827).
Glazioviana, *Warm. in Vidensk. Selsk. Skr.* VI. ii. (1881) 34.—Bras.
Hookeriana, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 105 = Podostemon Hookerianus.
Johnsonii, Wight, Ic. t. 1919 = Podostemon Johnsonii.
scaturiginum, *Mart. Nov. Gen. et Sp.* i. 3. t. 1.—Bras.
selaginoides, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 54 = Podostemon selaginoides.
Weddeliana, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 105.—Java.

MOACURRA, Roxb. Hort. Beng. 21 (1814); Fl. Ind. ii. 69 (1832) = **Chailletia**, DC.
gelonioides, Roxb. ll. cc. = C. gelonioides.

MOCANERA, Blanco, Fl. Filip. ed. I. 446 (1837) = **Dipterocarpus**, Gaertn. f. et **Anisoptera**, Korth. (Dipteroc.).
grandiflora, Blanco, l. c. 451 = Dipterocarpus grandiflorus.
Guiso, Blanco, l. c. 449 = Anisoptera Guiso.
Malaanonan, Blanco, l. c. 858 = Shorea Malaanonan.
Mangachapoi, Blanco, l. c. 450 = Shorea Mangachapoi.
Mayapis, Blanco, l. c. 449 = Dipterocarpus Mayapis.
plagata, Blanco, l. c. 447 = Anisoptera plagata.
polysperma, Blanco, l. c. 448 = Dipterocarpus polysperma.
thurifera, Blanco, l. c. 446 = Anisoptera thurifera.
verniciflua, Blanco, l. c. 450 = Dipterocarpus vernicifluus.

MOCANERA, Juss. Gen. 318 (1789) = **Visnea**, Linn. f. (Ternstroem.).
canariensis, Heynh. Nom. i. 884, nomen = V. Mocanera.

MOCINIA, DC. Prod. vii. 26 (1838) = **Stifftia**, Mikan (Compos.).
mutisioides, DC. l. c. = S. chrysantha.

MOCINNA, Benth. Pl. Hartw. 9 (1839), sphalm. = MOZINNA, Orteg. = **Jatropha**, Linn. (Euphorb.).

MOCINNA, Cerv. ex La Llave, in La Naturaleza, vii. (1885) Apénd. 70 = **Carica**, Linn. (Passifl.).
heterophila, Cerv. l. c. = C. heterophylla.

MOCINNA, Lag. Gen. et Sp. Nov. 31 (1816) = **Calea**, Linn. (Compos.).
brachiata, Lag. l. c. = C. brachiata.
brasiliensis, Spreng. Neue Entdeck. ii. 139 = C. pinnatifida.
serrata, Lag. Gen. et Sp. Nov. 31 = C. axillaris.

MOCQUINIA, Steud. Nom. ed. II. ii. 152 (1841), sphalm. = MOQUINIA, Spreng. = **Loranthus**, Linn.

MODANTHOS, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 11 = **Modiola**, Moench (Malvac.).
caroliniana, Alef. l. c. 12 = Modiola multifida.
decumbens, Alef. l. c. = Modiola multifida.
eriocarpa, Alef. l. c. = Modiola multifida.
geranioides, Alef. l. c. = Modiola geranioides.
prostrata, Alef. l. c. = Modiola multifida.
reptans, Alef. l. c. = Modiola reptans.
urticifolia, Alef. l. c. = Modiola multifida.

MODECCA, Lam. Encyc. iv. 208 (1797). *PASSIFLOREAE*, Benth. & Hook. f. i. 813.
BLEPHARANTHES, Sm. Gramm. Bot. 188 (1821).
CLEMANTHUS, Klotzsch, in Peters, Reise Mossamb. Bot. 143 (1861).
ERYTHROCARPUS, M. Roem. Syn. Pepon. 204 (1846).
KOLBIA, Beauv. Fl. Owar. ii. 91. t. 120 (1807).
MICROBLEPHARIS, M. Roem. Syn. Pepon. 133, 200 (1846).
PASCHANTHUS, Burch. Trav. i. 543 (1822).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 297. —Afr. trop.
aculeata, *Oliver, ex Hook. Ic. Pl.* t. 1317.—Afr. trop.
acuminata, *Blume, Bijdr.* 940.—Java.
aliena, Wall. Cat. n. 6766 (= Asclepiad.).—Ind. or.
antongilliana, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 51.—Madag.
apiculata, *Mast. in Hook. f. Fl. Brit. Ind.* ii. 603.—Burma.
australis, *R. Br. ex DC. Prod.* iii. 337.—Austral.
bracteata, Lam. Encyc. iv. 210 = Trichosanthes pubera.
cardiocarpa, *Hassk. Cat. Hort. Bog. Alt.* 187 (kardiocarpa).—Java.
cardiophylla, *Mast. in Hook. f. Fl. Brit. Ind.* ii. 602.—Reg. Himal.
cissampeloides, Planch. in Hook. Niger Fl. 365 = Ophiocaulon cissampeloides.
coccinea, Blanco, Fl. Filip. ed. II. 453 = heterophylla.
cordifolia, *Blume, Bijdr*, 939.—Ins. Andam.; Java.
cynanchifolia, Benth. in Hook. Niger Fl. 366 = Ophiocaulon cynanchifolius.
densiflora, *Baker, in Journ. Bot.* xx. (1882) 112.—Madag.
digitata, *Harv. Thes. Cap.* i. 8.—Afr. austr.
diversifolia, Schum. & Thonn. Beskr. Guin. Pl. 435 = lobata.
diversifolia, Wall. Cat. n. 6763 = Wightiana.
dubia, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 135.—Sumatra.
furfuracea, *Wall. Cat.* n. 1235.—Burma.
gummifera, Harv. in Harv. & Sond. Fl. Cap. ii. 500 = Ophiocaulon gummifer.
hastata, *Harv. Thes. Cap.* ii. 43.—Afr. austr.
heterophylla, *Blume, Bijdr.* 940.—Java.
heterophylla, Kurz, Andam. Rep. Append. A. 39 = cordifolia.
integrifolia, Lam. Encyc. iv. 209 = palmata.
integrifolia, *Wall. Cat.* n. 6765.—Hab.?
kardiocarpa, Hassk., vide cardiocarpa.
Kirkii, *Mast. in Oliver, Fl. Trop. Afr.* ii. 515.—Afr. trop.
lobata, *Jacq. Fragm.* 82. t. 131.—Afr. trop.
macrophylla, *Blume, Bijdr.* 939.—Java.
Mannii, *Mast. in Oliver, Fl. Trop. Afr.* ii. 516.—Afr. trop.
nicobarica, *Kurz, in Journ. Bot.* xiii. (1875) 326.—Malaya; Ins. Nicobar.

MODECCA:—
oblonga, *Hassk. Cat. Hort. Bog. Alt.* 187.—Java.
obtusa, *Blume, Bijdr.* 939.—Java.
palmata, *Lam. Encyc.* iv. 209.—Ind. or.
parviflora, Blanco, Fl. Filip. ed. I. 453 = cardiophylla.
parviflora, G. Don, Gen. Syst. iii. 59 = Ophiocaulon cynanchifolius?
Paschanthus, *Harv. in Harv. & Sond. Fl. Cap.* ii. 500. —Afr. austr.
peltata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 345.—Madag.
populifolia, *Blume, Rumphia*, i. 168.—Malaya.
quintuplinervia, *Miq. Fl. Ind. Bat.* i. 1. 1093.—Sumatra.
refracta, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 52. —Madag.
saponaria, Blanco, Fl. Filip. ed. II. 453 = trilobata.
senensis, *Mast. in Oliver, Fl. Trop. Afr.* ii. 513.—Afr. trop.
septemloba, E. Mey. in Drège, Zwei Pfl. Docum. 202 = Ceratiosicyos Ecklonii.
singaporeana, *Mast. in Hook. f. Fl. Brit. Ind.* ii. 601. —Singapore.
stricta, *Mast. in Oliver, Fl. Trop. Afr.* ii. 515.—Afr. trop.
tamnifolia, Planch. ex Mast. l. c. 517 = lobata.
tenuifolia, Planch. ex Benth. in Hook. Niger Fl. 366 sphalm. = lobata.
trilobata, Blanco, Fl. Filip. ed. II. 452 = palmata.
trilobata, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 132.—Ind. or.
trisecta, *Mast. in Oliver, Fl. Trop. Afr.* ii. 514.—Afr. trop.
tuberosa, Roxb. Hort. Beng. 49; Fl. Ind. iii. 134 = palmata.
Welwitschii, *Mast. in Oliver, Fl. Trop. Afr.* ii. 513.—Afr. trop.
Wightiana, *Wall. Cat.* n. 6764.—Ind. or.

MODECCOPSIS, Griff. in Proc. Linn. Soc. i. (1843) 171 = **Erythropalum**, Blume (Olacin.).
vaga, Griff. Notul. iv. 633 = E. vagum.

MODECOPSIS, Griff. Notul. iv. 633. t. 628 (1854) = praec.

MODESTA, Rafin. Fl. Tellur. iv. 75 (1836) = **Ipomoea**, Linn. (Convolv.).
congesta, Rafin. l. c. 76 = I. digitata, *Linn.*
coriacea, Rafin. l. c. = Rivea hypocrateriformis.
insignis, Rafin. l. c. = I. insignis.
macrorhiza, Rafin. l. c. = I. macrorrhiza.
mutabilis, Rafin. l. c. = I. acuminata.
paniculata, Rafin. l. c. 75 = I. digitata, *Linn.*
setosa, Rafin. l. c. 76 = I. setosa.
tuberculata, Rafin. l. c. = I. dasysperma.

MODIOLA, Moench, Meth. 619 (1794). *MALVACEAE*, Benth. & Hook. f. i. 205.
HAYNEA, Reichb. Consp. 202 (1828).
MODANTHOS, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 11.
caroliniana, G. Don, Gen. Syst. i. 466 = multifida.
decumbens, G. Don, l. c. = multifida.
erecta, *Lespinasse, in Bull. Soc. Bot. Fr.* vi. (1859) 647.—Am. austr.?
eriocarpa, G. Don, Gen. Syst. i. 466 = multifida.
fissistipula, Presl, Bot. Bemerk. 19 = multifida.
geranioides, *Walp. Rep.* i. 296.—Chili.
malvifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 45.—Reg. Argent.
multifida, *Moench, Meth.* 620.—Am. bor. occ. & austr.
prostrata, A. St. Hil. Fl. Bras. Mer. i. 211 = multifida.
reptans, *A. St. Hil. l. c.* 212.—Bras.
urticifolia, G. Don, Gen. Syst. i. 466 = multifida.

MOEHNIA, Neck. Elem. i. 9 (1790) = **Gazania**, Gaertn. (Compos.).

MOEHRINGIA, Linn. Syst. ed. II. 22 (1740) = **Arenaria**, Rupp. (Caryoph.).
bavarica, Dalla Torre, Atl. Alpenfl. 78 = A. bavarica, *Linn.*
bavarica, Gren. in Mém. Soc. Emul. Doubs, i. (1841) Bot. 37 = A. sperguloides, *Mutel* (in Addend.).
dasyphylla, Bruno, ex Balb. in Mém. Acad. Turin, Sc. Phys. i. [1804] 391 = A. bavarica, *Linn.*
diversifolia, Dolliner, ex Koch, in Flora, xxii. (1839) 2.—Styria.

MOEHRINGIA:—
erecta, Martr. Fl. Tarn, 106.—Europ.
flaccida, Schloss. & Vukot. Fl. Croat. 352 = Arenaria sperguloides, *Mutel* (in Addend.).
glabrescens, Reichb. ex Nym. Consp. 113 = A. Wulfenii.
glauca, Leybold, in Flora, xxxvi. (1853) 586 = *glaucovirens.*
glaucescens, Neilr. Veget. Croat. (1868) 199, in obs. = A. bavarica, *Linn.*
glaucovirens, Bertol. Fl. Ital. vi. 626 = A. bavarica, *Linn.*
Grisebachii, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 194.—Bulgar.
heterophylla, Dolliner, ex Reichb. Ic. Fl. Germ. v. 31. —Europ.
intermedia, Loisel. Fl. Gall. ed. II. i. 282 = A. bavarica, *Linn.*
intricata, Willk. Enum. Pl. Nov. Rar. Hisp. Austr. (1852) 14 = A. intricata.
Jankae, Griseb. ex Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 195.—Bulgar.
lateriflora, Fenzl, Verbreit. Alsin. tab. ad p. 18, 38 = A. lateriflora.
macrophylla, Fenzl, l. c. = A. macrophylla.
muscosa, Linn. Sp. Pl. 359 = A. sperguloides, *Mutel* (in Addend.).
muscosa, Lucé, Fl. Osil. 125 = Sagina procumbens.
nemorosa, Fenzl, ex Hemsl. Biol. Centr. Am. Bot. i. 69 = A. lanuginosa.
nodosa, [Clairv.] Man. Herb. 150 = Spergula nodosa.
obtusa, Reichb. ex Nym. Consp. 113 = A. polygonoides.
papulosa, Bertol. Fl. Ital. iv. 363.—Ital.
pendula, Fenzl, Verbreit. Alsin. tab. ad p. 46 = A. pendula.
pentandra, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 230 = A. trinervia.
platysperma, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 373.—Japon.
polygonoides, Mert. & Koch, Deutschl. Fl. iii. 272 = A. polygonoides.
Ponae, Fenzl, Verbreit. Alsin. tab. ad p. 46 = A. bavarica, *Linn.*
radiolata, Panč. ex Nym. Consp. 113 (Quid?).—Europ.
sedifolia, Willd. in Ges. Naturf. Fr. Berl. Mag. viii. (1818) 101 = A. bavarica, *Linn.*
sedoides, [Clairv.] Man. Herb. 150 = A. Cherleria.
sedoides, Cumino, ex Balb. in Mém. Acad. Turin, Sc. Phys. i. [1804] 391 = A. bavarica, *Linn.*
sperguloides, Nym. Consp. 113 = A. sperguloides, *Mutel* (in Addend.).
sphagnoides, Reichb. Ic. Fl. Germ. v. 32.—Austria.
stellarioides, Coss. in Bull. Soc. Bot. Fr. ix. (1862) 170. —Alger.
stenopetala, Dalla Torre, Atl. Alpenfl. 78 = A. trinervia.
stricta, Sibth. & Sm. Fl. Graec. Prod. i. 264 = Bufonia brachyphylla.
subulata, [Clairv.] Man. Herb. 150.—Europ.
thesiifolia, Dalla Torre, Atl. Alpenfl. 78 = A. thesiifolia.
thesiifolia, Reichb. Ic. Fl. Germ. v. 33, in obs. = A. polygonoides.
Thomasiana, J. Gay, ex Bertol. Fl. Ital. vi. 625 = A. Villarsii.
Tommasinii, Marches, in Bull. Adr. Sc. Nat. Trieste, v. (1880) 327 = *papulosa.*
trinervia, [Clairv.] Man. Herb. 150 = A. trinervia.
umbrosa, Fenzl, Verbreit. Alsin. 38 = A. umbrosa.
umbrosa, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 13 = A. macrophylla.
villosa, Fenzl, Verbreit. Alsin. tab. ad p. 46 = A. Wulfenii.

MOELLERIA, Scop. Introd. 335 (1777) = **Casearia**, Jacq. (Samyd.).

MOENCHIA, Ehrh. Beitr. ii. 177 (1788) = **Cerastium**, Dill. (Caryoph.).
cerastoides, Spreng. Syst. i. 497 = C. tetrandrum.
coerulea, Boiss. Diagn. Ser. I. viii. 104 = C. caeruleum.
erecta, Gaertn. Mey. & Scherb. Fl. Wett. i. 219 = C. quaternellum.
filiforme, Risso, Fl. Nice, 71.—Gallia.
glauca, Pers. Syn. i. 153 = C. quaternellum.
graeca, Boiss. & Heldr. Diagn. Ser. II. i. 91.—Graecia.
mantica, Bartl. Cat. Sem. Hort. Gotting. ex Koch, Syn. Fl. Germ. ed. II. 131 = C. manticum.
octandra, J. Gay, in Perrym. Pl. Phan. Fréjus, 55 = C. quaternellum.
quaternella, Ehrh. Beitr. ii. 180 = C. quaternellum.

MOENCHIA, Medic. in Act. Acad. Palat. vi. Phys. (1790) 493 = **Allium**, Tourn. (Liliac.).
narcissiflora, Medic. l. c. = A. senescens.
obliqua, Medic. l. c. = A. obliquum.

MOENCHIA, Neck. Elem. ii. 166 (1790) = **Cucubalus**, Tourn. (Caryoph.).

MOENCHIA, Roth, Tent. Fl. Germ. i. 273 (1788) = **Alyssum**, Tourn. (Crucif.).
aizoides, Roth, l. c. = Draba aizoides.
arvensis, Bernh. ex Hoffm. Fl. Germ. Comp. (1804) i. 32 = Camelina sativa.
campestris, Roth, Tent. Fl. Germ. i. 274 = A. calycinum.
incana, Roth, l. c. 273 = A. incanum.
podolicum, Bess. Enum. Pl. Volh. 26 = A. podolicum.
sativa, Roth, Tent. Fl. Germ. i. 274 = Camelina sativa.
saxatilis, Schrank, Fl. Monac. iv. t. 372 = A. montanum.

MOENCHIA, Wender. ex Steud. Nom. ed. II. ii. 153 (1841) = **Paspalum**, Linn. (Gramin.).
speciosa, Wender. ex Steud. l. c. = P. saccharoides.

MOERENHOUTIA, Blume, Orch. Archip. Ind. 99. t. 28, 42 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 604.
plantaginea, *Blume, l. c.* 100.—Ins. Tahiti.

MOERHINGIA, Linn. Phil. Bot. 32 (1751) = MOEHRENGIA, Linn. = **Arenaria**, Rupp. (Caryoph.).

MOERKENSTEINIA, Opiz, Seznam, 66 (1852) = **Senecio**, Tourn. (Compos.).
sylvatica, Opiz, l. c. = S. sylvaticus.

MOERORIS, Rafin. Sylva Tellur. 91 (1838) = **Phyllanthus**, Linn. (Euphorb.).
stipulata, Rafin. l. c. = P. Niruri.

MOESA, Blanco, Fl. Filip. ed. II. Suppl. 590 (1845) = MAESA, Forsk. (Myrsin.).

MOESSLERA, Reichb. in Moessl. Handb. ed. II. i. p. l. (1827); et Consp. 160 (1828) = **Tittmannia**, Brongn. (Bruniac.).
lateriflora, Eckl. & Zeyh. Enum. 142 = T. lateriflora.

MOGHAMIA, Steud. Nom. ed. II. ii. 153 (1841) = seq.

MOGHANIA, Jaume St. Hil. in Desv. Journ. Bot. i. (1813) 61 = **Flemingia**, Roxb. (Legumin.).
strobilifera, Jaume St. Hil. l. c. 62 = F. strobilifera.

MOGIPHANES, Mart. Nov. Gen. et Sp. ii. 29. t. 129 (1826). *AMARANTACEAE*, Benth. & Hook. f. iii. 37.
aquatica, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 316; *Contrib.* 84.—Paraguay.
brasiliensis, *Mart. Nov. Gen. et Sp.* ii. 34. t. 133.—Bras.
diffusa, Mart. l. c. 36 = ramosissima.
Dunaliana, Griseb. in Goett. Abh. xxiv. (1879) 35 = Gomphrena Dunaliana.
emarginata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 317; *Contrib.* 86.—Paraguay.
fistulosa, *Parodi, ll. cc.* 317; 85.—Paraguay.
flavescens, *Mart. Nov. Gen. et Sp.* ii. 36.—N. Granat.
glauca, *Griseb. in Goett. Abh.* xxiv. (1879) 35.—Reg. Argent.
hirtula, *Mart. Nov. Gen. et Sp.* ii. 30. t. 129.—Bras.
Jacquini, Schrad. Ind. Sem. Hort. Gotting. (1834) 4 = brasiliensis.
multicaulis, Mart. Nov. Gen. et Sp. ii. 32. t. 131 = straminea.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 318; *Contrib.* 87.—Paraguay.
philoxeroides, *Parodi, ll. cc.* 316; 85 (philorexoides).—Paraguay.
pilosa, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 348 = Gomphrena pilosa.
ramosissima, *Mart. Nov. Gen. et Sp.* ii. 31. t. 130.—Bras.
straminea, *Mart. l. c.* 35. t. 135.—Am. trop.
villosa, *Mart. l. c.* 33. t. 132.—Bras.
virgata, Schrad. Ind. Sem. Hort. Gotting. (1834); ex Moq. in DC. Prod. xiii. II. 382 = straminea?

MOGORI, Adans. Fam. ii. 223 (1763) = seq.

MOGORIUM, Juss. Gen. 106 (1789) = **Jasminum**, Tourn. (Oleac.).
aculeatum, Blanco, Fl. Filip. ed. I. 9 = J. marianum.
acuminatum, Lam. Tabl. Encyc. i. 23 = J. trinerve.
dichotomum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 105 = J. glandulosum.
dichotomum, Poir. Encyc. Suppl. iii. 713 = J. dichotomum.
elongatum, Lam. Tabl. Encyc. i. 23 = J. elongatum.
gimea, Zuccagni, in Roem. Collect. 114 = J. Sambac.
goaense, Zucc. l. c. = J. Sambac.
hirsutum, Poir. Encyc. Suppl. iii. 713 = J. pubescens.
lanceolatum, Poir. l. c. = J. lanceolatum.
multiflorum, Lam. Encyc. iv. 211 = J. undulatum.
myrtifolium, Lam. Tabl. Encyc. i. 23 = J. glaucum.
pubescens, Lam. Encyc. iv. 213 = J. pubescens.
Sambac, Lam. Tabl. Encyc. i. 23; et Encyc. iv. 210 = J. Sambac.
scandens, Lam. Encyc. iv. 214 = J. scandens.
sessiliflorum, Poir. Encyc. Suppl. iii. 713 = J. sessiliflorum.
triflorum, Lam. Tabl. Encyc. i. 23 = J. angustifolium.
trifoliatum, Lam. l. c. 24 = J. auriculatum.
undulatum, Lam. l. c. 23; et Encyc. iv. 212 = J. Sambac.
vimineum, Lam. Encyc. iv. 213 = J. angustifolium.

MOHAVEA, A. Gray, in Pacif. Rail. Rep. iv. 122 (1856). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 934.
viscida, *A. Gray, l. c.*—Calif.

MOHERINGIA, Zumag. Fl. Pedem. ii. 166 (1864) = MOEHRINGIA, Linn. = **Arenaria**, Rupp. (Caryoph.).

MOHLANA, Mart. Nov. Gen. et Sp. iii. 170. t. 290 (1829). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 81.
HILLERIA, Vell. Fl. Flum. 47 (1825); Ic. i. t 122 (1827).
MANCOA, Rafin. Fl. Tellur. iii. 56 (1836).
guineensis, *Moq. in DC. Prod.* xiii. II. 15.—Afr. trop.
latifolia, *Moq. l. c.* 16.—Madag.; Venez.?
nemoralis, *Mart. Nov. Gen. et Sp.* iii. 171.—Bras.
secunda, *Mart. l. c.* 172.—Peruv.; N. Granat.

MOKOF, Adans. Fam. ii. 50 (1763) = **Cleyera**, Thunb. (Ternstroem.).

MOLDAVICA, [Tourn.] Linn. Syst. ed. I (1735) = **Dracocephalum**, Linn. (Labiat.).
elata, Moench, Meth. 410 = Nepeta macrantha.
punctata, Moench, l. c. = D. Moldavica.
setosa, Stokes, Bot. Mat. Med. iii. 367 = D. Moldavica.
sibirica, Steud. Nom. ed. I. 285, in syn. = Nepeta macrantha.
suaveolens, Gilib. Fl. Lituan. i. 79 = D. Moldavica.

MOLDENHAUERA, Spreng. Syst. i. 373 (1825) = **Pyrenacantha**, Wight (Olacin.).
scandens, Spreng. l. c. 489 = P. scandens.

MOLDENHAWERA, Schrad. in Goett. Gel. Anz. i. (1821) 718. *LEGUMINOSAE*, Benth. & Hook. f. i. 569.
DOLICHONEMA, Nees, in Flora, iv. (1821) 303.
Blanchetiana, *Tul. in Arch. Mus. Par.* iv. (1844) 158.—Bras.
cuprea, *Pohl, Pl. Bras. Ic.* ii. 91. t. 160.—Bras.
emarginata, Moric. ex Benth. in Mart. Fl. Bras. xv. II. 77 = cuprea.
floribunda, *Schrad. in Goett. Gel. Anz.* i. (1821) 718.—Bras.
lucida, Benth. in Mart. Fl. Bras. xv. II. 77 = cuprea.
mollis, *Benth. l. c.*, in obs.—Venezuela.
speciosa, Allem. in Trab. Soc. Vell. (1852) 56 = floribunda.

MOLDENHAWERIA, Steud. Nom. ed. II. ii. 153 (1841) = praec.

MOLINA, Cav. Diss. 435. t. 263 (1790) = **Hiptage**, Gaertn. (Malpig.).
racemosa, Cav. l. c. = H. Madablota.

MOLINA, C. Gay, Fl. Chil. v. 345. t. 61 (1833) = **Dysopsis**, Baill. (Euphorb.).
chilensis, C. Gay, l. c. 346 = D. Gayana.

MOLINA, Giseke, Prael. Linn. 368 (1792) = MOLINAEA, Comm. = **Cupania**, Plum. (Sapind.).

MOLINA, Ruiz & Pav. Prod. 111. t. 24 (1794) = **Baccharis**, Linn. (Compos.).
articulata, Less. in Linnaea, vi. (1831) 140 = B. articulata.
caespitosa, Ruiz & Pav. Syst. Veg. 203 = B. caespitosa.
concava, Ruiz & Pav. l. c. 206 = B. concava.
conyzoides, Less. in Linnaea, vi. (1831) 139 = B. conyzoides.
corymbosa, Ruiz & Pav. Syst. Veg. 210 = B. corymbosa, *Pers.*
crispa, Less. in Linnaea, vi. (1831) 141 = B. genistelloides.
cylindrica, Less. l. c. 144 = B. genistelloides.
dependens, Ruiz & Pav. Syst. Veg. 202 = B. dependens.
emarginata, Ruiz & Pav. l. c. = B. emarginata.
ferruginea, Ruiz & Pav. l. c. 211 = B. ferruginea.
glutinosa, Alam. ex DC. Prod. v. 312 = Gymnosperma corymbosum.
incana, Ruiz & Pav. Syst. Veg. 211 = B. thyoides.
juncea, Less. in Linnaea, vi. (1831) 145 = B. juncifformis.
latifolia, Ruiz & Pav. Syst. Veg. 208 = B. latifolia.
linearis, Less. in Linnaea, vi. (1831) 139, 505 = B. serrulata.
linearis, Ruiz & Pav. Syst. 205 = B. Pingraea?
microcephala, Less. in Linnaea, vi. (1831) 142 = B. microcephala.
milleflora, Less. l. c. 143 = B. genistelloides.
nitida, Ruiz & Pav. Syst. Veg. 204 = B. nitida.
oblongifolia, Ruiz & Pav. l. c. 203 = B. oblongifolia.
obovata, Ruiz & Pav. l. c. 206 = B. obovata.
parviflora, Ruiz & Pav. l. c. 209 = B. marginalis.
pentaptera, Less. in Linnaea, vi. (1831) 505 = B. pentaptera.
phyteumoides, Less. l. c. 146 = B. phyteumoides.
prostrata, Ruiz & Pav. Syst. Veg. 204 = B. prostrata.
racemosa, Ruiz & Pav. l. c. 209 = B. racemosa.
reticulata, Ruiz & Pav. l. c. 212 = B. genistelloides.
rhexioides, Less. in Linnaea, vi. (1831) 406 = B. rhexioides.
sagittalis, Less. l. c. 144 = B. sagittalis.
salicifolia, Ruiz & Pav. Syst. Veg. 210 = B. salicifolia.
scabra, Ruiz & Pav. l. c. = B. scabra.
scandens, Ruiz & Pav. l. c. 205 = B. scandens, *Pers.*
scoparia, Less. in Linnaea, vi. (1831) 147, in obs. = B. scoparia.
sessilifolia, Less. l. c. 140 = B. sessilifolia.
striata, Ruiz & Pav. Syst. Veg. 207 = B. Fevillei.
tomentosa, Ruiz & Pav. l. c. 203 = B. tomentosa.
trimera, Less. in Linnaea, vi. (1831) 141 = B. genistelloides.
uniflora, Ruiz & Pav. Syst. Veg. 208 = B. uniflora.
venosa, Ruiz & Pav. l. c. 212 = B. venosa.
viscosa, Ruiz & Pav. l. c. 207 = B. glutinosa.

MOLINAEA, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, xx. (1830) 109; et Am. Journ. Soc. xx. (1831) 152 = **Jubaea**, H. B. & K. (Palm.).
microcos, Bert. ll. cc. = J. spectabilis.

MOLINAEA, Comm. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 364 = **Retanilla**, Brongn. (Rhamn.).

MOLINAEA, Comm. ex Juss. Gen. 248 (1789) = **Cupania**, Plum. (Sapind.).
alternifolia, Willd. Sp. Pl. ii. 329 = C. alternifolia.
arborea, Blanco, Fl. Filip. ed. I. 292 = Hemigyrosa canescens.
brevipes, Radlk. in Sitzb. Math-Phys. Acad. Muench. ix. (1879) 651.—Madag.
canescens, Roxb. Pl. Corom. i. 43. t. 60 = Hemigyrosa canescens.
cupanioides, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 650 = C. laevis.
laevis, Willd. Sp. Pl. ii. 329 = C. laevis.
macrantha, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 650.—Ins. Maurit.
petiolaris, Radlk. l. c. 651.—Madag.
retusa, Radlk. l. c. 650.—Madag.
Tolambitou, Radlk. l. c. 651 = C. Tolambitou.

MOLINERIA, Colla, Hort. Ripul. App. ii. 333. t. 18 (1826) = **Curculigo**, Gaertn. (Amaryll.).
capitulata, Herb. Amaryll. 84 = C. recurvata.
crassifolia, Baker, in Journ. Linn. Soc. xvii. (1878) 121.—Reg. Himal.
Finlaysoniana, Baker, l. c. = C. Finlaysoniana.
gracilis, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 177 = C. gracilis.
latifolia, Herb. ex Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 232 = C. latifolia.
longiflora, Kurz, l. c.—Ins. Banca.
plicata, Colla, Hort. Rip. App. ii. 333 = C. recurvata.
plicata, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 176 = C. latifolia.
recurvata, Herb. Amaryll. 84 = C. recurvata.
rhizophylla, Baker, in Journ. Linn. Soc. xvii. (1878) 121.—Ins. Seychelles.
sulcata, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 175 = C. recurvata.
sumatrana, Herb. Amaryll. 84 = C. latifolia.
villosa, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 176 = C. latifolia.

MOLINERIA, Parl. Fl. Ital. i. 236 (1848) = **Aira**, Linn. (Gramin.).
laevis, Hack. Cat. Gram. Portug. 17 = *lendigera*.
lendigera, Hack. in Oestr. Bot. Zeitschr. xxvii. (1877) 120.—Lusit.; Hisp.
minuta, Parl. Fl. Ital. i. 237 = A. minuta.
palustris, Tod. ex Nym. Consp. 807 = A. minuta.

MOLINIA, Schrank, Baier. Fl. i. 100. 334 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1185.
AMBLYTES, Dulac, Fl. Hautes-Pyr. 80 (1867).
ENODIUM, Gaud. Agrost. Helv. i. 145 (1811).
MONILIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 110 (1821).
altissima, Link, Hort. Berol. i. 197 = caerulea.
aquatica, Wibel, Prim. Fl. Werth. 116 = Catabrosa aquatica.
arundinacea, Schrank, Baier. Fl. i. 336 = caerulea.
caerulea, *Moench, Meth.* 183.—Europ.; As. bor.
capillaris, Hartm. Handb. Skand. Fl. ed. I. 56 = Glyceria distans.
conferta, *Hartm. l. c.* ed. IV. 33.—Europ.
distans, Hartm. l. c. ed. I. 56 = Glyceria distans.
fluitans, Hartm. l. c. = Glyceria fluitans.
litoralis, Host, Fl. Austr. i. 118 = caerulea.
maritima, Hartm. Handb. Skand. Fl. ed. I. 55 = Glyceria maritima.
maxima, Hartm. l. c. 56 = Glyceria aquatica.
Olgae, *Regel, in Act. Hort. Petrop.* vii. (1880) 625.—Kokania.
pendulina, *Hartm. Handb. Skand. Fl.* ed. IV. 34.—Europ.
plicata, Hartm. l. c. = Glyceria fluitans.
remota, *Hartm. l. c.* 33.—Europ.
rivulorum, *Pomel, Nouv. Mat. Fl. Atl.* 388.—Alger.
serotina, Mert. & Koch, Deutschl. Fl. i. 585 = Diplachne serotina.
squarrosa, *Trin. ex Ledeb. Fl. Alt.* i. 105.—Mont. Altai.
sylvatica, Link, Hort. Berol. i. 197, in obs. = caerulea.
varia, Schrank, Baier. Fl. i. 334 = caerulea.
variabilis, Wibel, Prim. Fl. Werth. 115 = caerulea.

MOLINIERA, Ball, in Journ. Linn. Soc. xvi. (1878) 718, sphalm. = MOLINERIA, Parl. = **Aira**, Linn. (Gramin.).

MOLIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = **Allium**, Tourn. (Liliac.).
roseum, Fourr. l. c. = A. roseum.

MOLKENBOERIA, De Vriese, in Verh. Haarl. x. (1854) 38 = **Scaevola**, Linn. (Gooden.).
macrophylla, De Vriese, l. c. 44 = S. macrophylla.
membranacea, De Vriese, l. c. 40 = S. pilosa.
microphylla, De Vriese, l. c. 44. t. 9. f. 1–10 = S. microphylla.
pilosa, De Vriese, l. c. 39 = S. pilosa.
platyphylla, De Vriese, l. c. 43. t. 6 = S. platyphylla.
semiamplexicaulis, De Vriese, l. c. 41. t. 5 = S. pilosa.
striata, De Vriese, l. c. 42 = S. striata.

MOLLE, [Tourn.] Adans. Fam. ii. 341 (1763) = **Schinus**, Linn. (Anacard.).

MOLLIA, J. F. Gmel. Syst. 420 (1791) = **Baeckea**, Linn. (Myrtac.).
imbricata, J. F. Gmel. l. c. = B. crenulata.

MOLLIA, Mart. Nov. Gen. et Sp. i. 96. t. 60 (1824). *TILIACEAE*, Benth. & Hook. f. i. 236.
SCHLECHTENDALIA, Spreng. Syst. iv. Cur. Post. 295 (1827).
glabrescens, *Benth. in Hook. Journ. Bot.* iv. (1842) 131.—Guiana.
gracilis, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 59.—Bras.
lepidota, *Spruce, ex Benth. l. c.*—Bras.
longifolia, *Spruce, ex Benth. l. c.*—Bras.
speciosa, *Mart. Nov. Gen. et Sp.* i. 97. t. 60.—Bras.
tomentosa, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 60.—Bras.

MOLLIA, Willd. Hort. Berol. 11. t. 11 (1806) = **Polycarpaea**, Lam. (Caryoph.).
alsinifolia, Spreng. Syst. i. 795 = Polycarpon alsinifolium.
aristata, Ait. Hort. Kew. ed. II. ii. 62 = P. corymbosa.
corymbosa, Willd. Hort. Berol. 11 = P. corymbosa.
diffusa, Willd. l. c. = P. Teneriffae.
fragilis, Spreng. Syst. i. 795 = P. fragilis.
gnaphalodes, Spreng. l. c. = P. gnaphalodes.
latifolia, Willd. Enum. Hort. Berol. 269 = P. latifolia.
minuartoides, Spreng. Mant. Fl. Hal. i. 37 = P. minuartoides.
polycarpoides, Link, Handb. ii. 422 = Polycarpon peploides.
Polycarpon, Spreng. Nov. Prov. 28 = Polycarpon peploides.
rosea, Hochst. ex Boiss. Fl. Orient. i. 737 = Polycarpon succulentum.
spadicea, Willd. Hort. Berol. 11 = P. corymbosa.
stellata, Willd. l. c. = P. stellata.
tenuifolia, Willd. l. c. = P. tenuifolia.

MOLLINEDIA, Ruiz & Pav. Prod. 83 (1794). *MONIMIACEAE*, Benth. & Hook. f. iii. 140.
TETRATOME, Poepp. & Endl. Nov. Gen. et Sp. ii. 46 (1838).
acuminata, *F. Muell. Fragm.* v. 155.—Austral.
boliviensis, *A. DC. in Journ. Bot.* iii. (1865) 220.—Bolivia.
brasiliensis, *Schott, ex Tul. in Arch. Mus. Par.* viii. (1855–56) 377.—Bras.
campanulacea, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 42.—N. Granat.
cinerea, *Tul. in Arch. Mus. Par.* viii. (1855–56) 380.—Bras.
clavigera, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 44.—Bras.
coriacea, Baill. Hist. Pl. i. 304 = Kibara Blumei.
cuspidata, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 320 = nitida.
elegans, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 44.—Bras.
elliptica, A. DC. in DC. Prod. xvi. II. 664 = nitida.
floribunda, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 41.—Bras.
gracilis, *Tul. l. c.*—Bras.
Huegeliana, *Tul. l. c.* 45.—Austral.
ibaguensis, *Tul. l. c.* 41.—N. Granat.
lanceolata, *Ruiz & Pav. Syst. Veg.* 143.—Peruv.
latifolia, *Tul. in Arch. Mus. Par.* viii. (1855–56) 402.—Bras.
laurina, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 43.—Guiana; Ins. Trinit.
laxiflora, F. Muell. Census, 3 = Kibara laxiflora.
lepidota, Tul. in Arch. Mus. Par. viii. (1855–56) 403 = Citriosma Poeppigii.
ligustrina, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 44.—Bras.
longifolia, *Tul. l. c.* 41.—Bras.
longipes, F. Muell. Census, 3 = Kibara longipes.
loxocarya, *Benth. Fl. Austral.* v. 287.—Austral.
macrantha, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 42.—N. Granat.
macrophylla, Tul. l. c. 45 = Kibara macrophylla.
multiflora, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 317 = floribunda.
myrtoidea, Baill. in Bull. Soc. Linn. Par. i. (1882) 342 = Ephippiandra myrtoidea.

MOLLINEDIA:—
nigrescens, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 42.—Mexic.
nitida, *Tul. l. c.* 43.—Bras.
ovata, *Ruiz & Pav. Syst. Veg.* 143.—Peruv.
pellucens, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 43.—Bras.
polycarpa, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 318 = laurina.
pubescens, F. Muell. Census, 3 = Kibara pubescens.
racemosa, *Tul. in Arch. Mus. Par.* viii. (1855–56) 382.—Peruv.
repanda, *Ruiz & Pav. Syst. Veg.* 142.—Peruv.
sancta, Baill. Hist. Pl. i. 306 = Matthaea sancta.
Selloi, *A. DC. in DC. Prod.* xvi. II. 666.—Bras.
sericiflora, *A. DC. in Journ. Bot.* iii. (1865) 220.—Bras.
tomentosa, *Tul. in Arch. Mus. Par.* viii. (1855–56) 402.—Peruv.
triflora, *Tul. l. c.* 394.—Bras.
umbellata, *Tul. l. c.* 397.—Bras.
utriculata, *Mart. ex Tul. in Mart. Fl. Bras.* iv. 1. 319.—Bras.
viridiflora, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 43.—Mexic.
Wardellii, *F. Muell. Fragm.* v. 155.—Austral.
Widgrenii, *A. DC. in Journ. Bot.* iii. (1865) 220.—Bras.

MOLLOYA, Meissn. in Hook. Kew Journ. vii. (1855) 382 **Grevillea**, R. Br. (Proteac.).
cynanchicarpa, Meissn. in DC. Prod. xiv. 348 = G. cynanchicarpa.

MOLLUGO, Linn. Gen. ed. I. 336 (1737). *FICOIDEAE*, Benth. & Hook. f. i. 857.
DOOSERA, Roxb. ex Wight & Arn. Prod. 362 (1834).
GLINUS, Loefl. ex Linn. Sp. Pl. ed. I. 463 (1753).
LAMPETIA, Rafin. Fl. Tellur. iii. 34 (1836).
NEMALLOSIS, Rafin. l. c. (1836).
PAULO-WILHELMIA, Hochst. in Flora, xxvii. (1844) 17.
PHYSA, Thou. Gen. Nov. Madag. 20 (1806).
PLENCHIA, Rafin. Specch. i. 194 (1814).
ROLOFA, Adans. Fam. ii. 256 (1763).
TRIGASTROTHECA, F. Muell. in Hook. Kew Journ. ix. (1857) 16.
arenaria, H. B. & K. Nov. Gen. et Sp. vi. 20 = verticillata.
axillaris, Schlecht. ex Rohrb. in Mart. Fl. Bras. xiv. II. 242 = verticillata.
Bainesii, *Oliver, Fl. Trop. Afr.* ii. 590.—Afr. trop.
bellidifolia, Ser. in DC. Prod. i. 391 = nudicaulis.
Berteriana, Ser. l. c. = verticillata.
Cerviana, *Ser. l. c.* 392.—Geront. trop. et subtrop.
cordifolia, Ser. l. c. = Polycarpon tetraphyllum.
denticulata, Guill. & Perr. Fl. Seneg. Tent. i. 45 = oppositifolia.
dichotoma, Schrank, Pl. Rar. Hort. Monac. t. 64 = verticillata.
dichotoma, Ser. in DC. Prod. i. 392 = Pharnaceum dichotomum.
diffusa, Willd. ex Steud. Nom. ed. II. ii. 154 = verticillata.
digyna, *Montr. in Mém. Acad. Lyon*, x. (1860) 179.—N. Caled.
disticha, Ser. in DC. Prod. i. 392 = racemosa.
enneandra, *Wright, in Sauv. Fl. Cub.* 7.—Cuba.
erecta, *Burm. f. Fl. Ind.* 32.—Zeylan.
flavescens, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 226.—Ins. Galapag.
fluvialis, Steud. Nom. ed. II. ii. 154 = Pharnaceum dichotomum.
fragilis. *Wawra, in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 565.—Afr. trop.
glinoides, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 171 = radiata.
glinoides, A. Rich. Tent. Fl. Abyss. i. 48 = oppositifolia.
Glinus, A. Rich. l. c. = hirta.
glomerata, Ser. in DC. Prod. i. 392 = Adenogramma galioides.
gracillima, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 226 = verticillata.
hirta, *Thunb. Prod. Pl. Cap.* 24.—Reg. trop. et subtrop.
Hoffmannseggiana, Ser. in DC. Prod. i. 393 = verticillata.

MOLLUGO:—
juncea, Fenzl, in Ann. Wien. Mus. i. (1838) 378 = verticillata.
lanata, Steud. Nom. ed. II. ii. 154 = Pharnaceum lanatum.
Lichtensteiniana, Ser. in DC. Prod. i. 393 = Adenogramma diffusa.
linearis, Ser. l. c. 392 = Pharnaceum lineare.
Linkii, Ser. l. c. = pentaphylla.
lotoides, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 662, 776 = hirta.
marginata, Ser. in DC. Prod. i. 392 = Psammotropha parvifolia.
maritima, Ser. l. c. 393 = Sesuvium pentandrum.
microphylla, Ser. l. c. 392 = Pharnaceum microphyllum.
mucronata, Ser. l. c. 393 = Psammotropha androsacea.
multiflora, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20; Ser. in DC. Prod. i. 392 = pentaphylla.
novo-hollandica, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 14 = oppositifolia.
nudicaulis, *Lam. Encyc.* iv. 234.—Reg. trop. et subtrop.
oppositifolia, *Linn. Sp. Pl.* 89.—Geront. trop.
orygioides, *F. Muell. ex Benth. Fl. Austral.* iii. 333. —Austral.
paniculata, Burm. f. Fl. Ind. 32 = pentaphylla, oppositifolia.
parviflora, Ser. ex DC. Prod. i. 391 = oppositifolia.
patens, Steud. Nom. ed. II. ii. 154 = Pharnaceum dichotomum.
Pauli-Wilhelmi, *Schweinf. Beitr. Fl. Aethiop.* 267, nomen.—Afr. trop.
pentaphylla, *Linn. Sp. Pl.* 89.—As. trop.; Polynes.
Pluckenetiana, Steud. Nom. ed. II. ii. 154 = triphylla.
pumila, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = Bergia repens.
quadrangularis, Ser. in DC. Prod. i. 392 = Psammotropha quadrangularis.
racemosa, *Lam. Tabl. Encyc.* i. 218.—Ind. or.
radiata, *Ruiz & Pav. Fl. Per.* i. 48.—Chili.
reflexa, Steud. Nom. ed. II. ii. 154 = Pharnaceum reflexum.
rigida, Steud. l. c. = Adenogramma rigida.
Schrankii, Ser. in DC. Prod. i. 391 = verticillata.
serpyllifolia, Ser. l. c. = Pharnaceum serpyllifolium.
serrulata, Sond. in Linnaea. xxiii. (1850) 15 = oppositifolia.
sinaica, *Steud. Nom.* ed. II. ii. 154.—Arab.
spathulata, Poepp. ex Fenzl, in Ann. Wien. Mus. i. (1838) 359 = oppositifolia.
Spergula, Linn. Syst. ed. X. 881 = oppositifolia.
spergulaefolia, Willd. ex Steud. Nom. ed. II. ii. 154 = verticillata.
sperguloides, *Ser. ex DC. Prod.* i. 392.—Ind. or.
stricta, Linn. Sp. Pl. ed. II. 131 = pentaphylla.
subserrata, Blanco, Fl. Filip. ed. I. 51 = pentaphylla.
teretifolia, Ser. in DC. Prod. i. 393 = Pharnaceum teretifolium.
tetraphylla, Linn. Sp. Pl. 89 = Polycarpon tetraphyllum.
trifolia, Schult. Mant. ii. 471 = pentaphylla.
trigastrotheca, *F. Muell. Pl. Vict.* i. 201.—Austral.
trigona, Steud. Nom. ed. II. ii. 154 = Pharnaceum trigonum.
triphylla, *Burm. f. Fl. Ind.* 32.—Ind. or.
triphylla, Lour. Fl. Cochinch. 62; Link, Enum. Hort. Berol. i. 108 = pentaphylla.
triphylla, Schrank, ex Steud. Nom. ed. II. ii. 154 = verticillata.
umbellata, Ser. in DC. Prod. i. 393 = M. Cerviana.
verticillata, *Linn. Sp. Pl.* 89.—Am. et Afr. trop.
verticillata, Roxb. Hort. Beng. 9; Fl. Ind. i. 360 = oppositifolia.

MOLOPANTHERA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 580. *RUBIACEAE*, Benth. & Hook. f. ii. 41.
Burchellii, *Hook. f. in Benth. & Hook. f. Gen.* ii. 42.—Bras.
paniculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 581.—Bras.

MOLOPOSPERMUM, Koch, in Nov. Act. Nat. Cur. xii. (1824) 108. *UMBELLIFERAE*, Benth. & Hook. f. i. 882.
cicutarium, *DC. Prod.* iv. 230.—Europ.
peloponnesiacum, Koch, in Nov. Act. Nat. Cur. xii. (1824) 108 = cicutarium.

MOLOSPERMUM, Steud. Nom. ed. II. ii. 155 (1841) = praec.

MOLPADIA, Cass. in Bull. Soc. Philom. (1818) 166 = **Buphthalmum**, Linn. (Compos.).
suaveolens, Cass. in Dict. Sc. Nat. xxxii. 401 = B. speciosum.

MOLTKIA, Lehm. in Neue Schr. Naturf. Ges. Halle, iii. II. (1817) 3. *BORAGINEAE*, Benth. & Hook. f. ii. 861.
anatolica, Boiss. Diagn. Ser. I. xi. 114 = coerulea.
angustifolia, *DC. Prod.* x. 72.—Assyria.
aurea, *Boiss. Diagn.* Ser. I. iv. 49.—As. Min.
coerulea, *Lehm. in Neue Schr. Naturf. Ges. Halle*, iii. II. (1817) 6.—As. Min.; Persia; Syria.
cyrenaica, Spreng. Syst. i. 548 = Nonnea phaneranthera.
graminifolia, Nym. Consp. 518 = Lithospermum graminifolium.
libanotica, Zucc. in Abh. Akad. Muench. iii. (1837-43) 246. t. 3. f. 2 = Solenanthus Tournefortii.
parviflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 171.—Reg. Himal.
petraea, *Boiss. Fl. Orient.* iv. 221.—Europ. or.
punctata, Lehm. in Neue Schr. Naturf. Ges. Halle, iii. II. (1817) 5 = coerulea.
Sendtneri, Boiss. Diagn. Ser. II. iii. 138 = Zwackhia aurea.

MOLUBDA, Rafin. Sylva Tellur. 123 (1838) = **Plumbago**, Linn.
scandens, Rafin. l. c. = P. scandens.

MOLUCCA, [Tourn.] Linn. Syst. ed. I (1735) = **Moluccella**, Linn. (Labiat.).
levis, Moench, Meth. 404 = Moluccella laevis.
spinosa, Moench, l. c. = Moluccella spinosa.

MOLUCCELLA, Linn. Gen. ed. I. 176 (1737).
LABIATAE, Benth. & Hook. f. ii. 1211.
CHASMONIA, Presl, Fl. Sic. p. xxxvii (1826).
MOLUCCA, [Tourn.] Linn. Syst. ed. I (1735).
acutiloba, Ledeb. in Eichw. Pl. Nov. Casp. Cauc. 39. t. 35 = Lagochilus acutilobus.
armata, Sieber, ex Benth. Lab. Gen. et Sp. 640 = spinosa.
diacanthophylla, Pall. in Nov. Act. Acad. Petrop. x. (1797) 380. t. 11 = Lagochilus diacanthophyllus.
frutescens, Linn. Sp. Pl. 587 = Ballota spinosa.
frutescens, Sibth. & Sm. Fl. Graec. vi. 55. t. 568 = Ballota integrifolia.
fruticosa, Forsk. Fl. Aegypt. Arab. p. xxviii = Ballota Pseudodictamnus.
grandiflora, Ledeb. Fl. Alt. ii. 418 = Lagochilus Bungei.
grandiflora, Steph. ex Willd. Sp. Pl. iii. 130 = Lagochilus diacanthophyllus.
integrifolia, R. Br. in Salt, Abyss. App. p. lxiv = Otostegia integrifolia.
laevis, *Linn. Sp. Pl.* 587.—As. occ.
laniflora, Willd. ex Benth. in Linnaea, xi. (1837) 339 = Marrubium lanatum.
lanigera, Bieb. ex Poir. Encyc. Suppl. iii. 722 = Eremostachys laciniata.
Marrubiastrum, Steph. in Mém. Soc. Nat. Mosc. ii. (1809) 8 = Marrubium lanatum.
microphylla, Delile, in Laborde, Voy. Arab. Pétr. 83 = Otostegia Schimperi.
mongholica, Turcz. ex Ledeb. Fl. Ross. iii. 402 = Marrubium eriostachyum.
persica, Burm. f. Fl. Ind. 128 = Otostegia microphylla.
phlomoides, Wight, ex Wall. Cat. sub n. 2527 = Leucas chinensis.
quadrangula, Sievers, in Pall. N. Nord. Beitr. vii. 275 = Eremostachys phlomoides.
repanda, R. Br. in Salt, Abyss. App. p. lxiv = Otostegia repanda.
scariosa, R. Br. l. c. = Otostegia scariosa.
sinaitica, Ehrenb. ex Boiss. Fl. Orient. iv. 777 = Otostegia scariosa.
spinosa, Falk, Beitr. ii. 208 = Eremostachys tuberosa.
spinosa, *Linn. Sp. Pl.* 587.—Europ. austr.; Syria.
tuberosa, Pall. Reise, iii. 738 = Eremostachys tuberosa.
tuberosa, Sievers, in Pall. N. Nord. Beitr. vii. 331 = Eremostachys moluccelloides.

MOLUCELLA, Juss. Gen. 115 (1789) = praec.

MOLUCHIA, Medic. Malv. 10 (1787) = **Melochia**, Dill. (Stercul.).
domingensis, Steud. Nom. ed. II. ii. 155, sphalm. = Melochia pyramidata.
frutescens, Medic. Malv. 10 = Melochia tomentosa.
fruticosa, Medic. l. c. = Melochia pyramidata.
herbacea, Medic. l. c. = Melochia pyramidata.

MOLY, [Rupp.] Moench, Meth. 286 (1794) = **Allium**, (Liliac.).
latifolium, S. F. Gray, Nat. Arr. Brit. Pl. ii. 180 = A. ursinum.
odorum, Moench, Meth. Suppl. 98 = A. odorum.
speciosum, Moench, Meth. 286 = A. nigrum.

MOLYZA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
Moly, Salisb. l. c. 92 = A. Moly.

MOMISIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 122 (1819) = **Celtis**, Tourn. (Urtic.).
aculeata, Klotzsch, in Linnaea, xx. (1847) 539 = C. aculeata.
alnifolia, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 193 = C. alnifolia.
brevifolia, Klotzsch, in Linnaea, xx. (1847) 538 = C. Tala.
Chichape, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 193 = C. Tala.
Clausseniana, Wedd. l. c. 194 = C. Clausseniana.
crenata, Wedd. l. c. 195 = C. crenata.
dichotoma, Klotzsch, in Linnaea, xx. (1847) 539 = C. dichotoma.
Ehrenbergiana, Klotzsch, l. c. 538 = C. aculeata.
ferruginea, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 194 = C. ferruginea.
flexuosa, Wedd. l. c. 195 = C. boliviensis?
integrifolia, Wedd. l. c. 196 = C. Selloviana.
laevigata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 123 = C. aculeata.
Lamarckii, Blume, Mus. Bot. Lugd. Bat. ii. 69 = C. mississipiensis.
lancifolia, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 196 = C. lancifolia.
membranacea, Wedd. l. c. = C. membranacea.
pallida, Planch. in DC. Prod. xvii. 191, in syn. = C. Tala.
pubescens, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 123 = C. pubescens.
spinosissima, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 195 = C. spinosissima.
Tala, Planch. in DC. Prod. xvii. 191, in syn. = C. Tala.
tarijensis, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 194 = C. glycycarpa.
triflora, Klotzsch, in Linnaea, xx. (1847) 537 = C. Tala.
zizyphoides, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 124 = C. zizyphoides.

MOMORDICA, [Tourn.] Linn. Syst. ed. I (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 825.
AMORDICA, Neck. Elem. i. 241 (1790).
MURICIA, Lour. Fl. Cochinch. 596 (1790).
NEVROSPERMA, Rafin. in Am. Monthly Mag. (1818) 40.
POPPYA, Neck. Elem. i. 241 (1790).
ZUCCA, Comm. ex Juss. Gen. 398, in obs. (1789).
aculeata, *Poir. Encyc. Suppl.* iii. 723.—Hab.?
acutangula, Steud. Nom. ed. I. 536 = aculeata.
adoënsis, Hochst. in Flora, xxiv. (1841) i. Intell. 22; et ex A. Rich. Tent. Fl. Abyss. i. 293 = Coccinia adoënsis.
angulata, Savi, ex Steud. Nom. ed. II. ii. 155, nomen = M. Elaterium.
anigosantha, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 536. —Afr. trop.
anthelmintica, Schum. & Thonn. Beskr. Guin. Pl. 423 = M. Charantia.
aspera, Lam. Fl. Fr. ii. 191 = Ecballium Elaterium.
auriculata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 (Quid?).—Java.
Balsamina, Descourt. Fl. Antill. iii. 62 = M. Charantia.
Balsamina, *Linn. Sp. Pl.* 1009.—Amphig. trop.
Balsamina, Wall. Cat. n. 6741 A, B = dioica.
bicolor, Blume, Bijdr. 928 = Coccinia indica.
brachybotrys, Poepp. & Endl. Nov. Gen. et Sp. ii. 54 = Cyclanthera brachybotrys.

MOMORDICA:—
calcarata, Colebr. ex Wall. Cat. n. 6740 = Thladiantha calcarata.
cardiospermoides, Klotzsch, in Peters, Reise Mossamb. Bot. 150 = clematidea.
carinata, Vell. Fl. Flum. x. t. 97 = Luffa aegyptiaca.
Casea, Buch.-Ham. ex Wall. Cat. n. 6740 = Thladiantha calcarata.
Charantia, Descourt. Fl. Antill. viii. 355 = M. Balsamina.
Charantia, *Linn. Sp. Pl.* 1009.—Amphig. trop.
cirrhiflora, Wight, ex Steud. Nom. ed. II. ii. 155 = Modecca Wightiana.
cissoides, *Planch. ex Benth. in Hook. Niger Fl.* 370.—Afr. trop.
clematidea, *Sond. in Harv. & Sond. Fl. Cap.* ii. 491.—Afr. austr.
cochinchinensis, *Spreng. Syst.* iii. 14.—As. trop.
cordatifolia, Godoy Torres, in Patriota, iii. 71; ex Cogn. in Mart. Fl. Bras. vi. IV. 87 = Cayaponia Martiana.
cordifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 492 = foetida.
corymbifera, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 539.—Afr. trop.
Covel, Dennst. Schluess. Hort. Malab. 34 = Coccinia indica.
crinocarpa, Fenzl, ex M. Roem. Syn. Pepon. 59, sphalm. = *echinocarpa*.
cucullata, Hook. f. in Oliver, Fl. Trop. Afr. ii. 538 = Schimperiana.
cylindrica, Blanco, Fl. Filip. ed. I. 769 = M. Charantia.
cylindrica, Linn. Sp. Pl. 1009 = Luffa cylindrica.
Cymbalaria, *Fenzl, ex Naud. in Ann. Sc. Nat.* Sér. IV. xii. (1859) 134.—Afr. trop.
dasycarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 291 = Cucumis dipsaceus.
denticulata, *Miq. Fl. Ind. Bat.* i. 1. 1090.—Sumatra; Borneo.
denudata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 618.—Zeylan.
dictyosperma, *Griseb. in Lorentz, Veg. Nord Entre-Rios*, 117, nomen.—Reg. Argent.
dioica, *Roxb. ex Willd. Sp. Pl.* iv. 605.—Ind. or.
dioica, Wall. Cat. n. 6750 A, F = cochinchinensis.
echinata, Muhl. in Trans. Am. Phil. Soc. iii. (1793)180, nomen; et ex Willd. Sp. Pl. iv. 605 = Echinocystis lobata.
echinocarpa, Fenzl, ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 531 = Luffa echinata.
ecirrhata, Stokes, Bot. Mat. Med. iv. 465 = Ecballium Elaterium.
elastica, Salisb. Prod. 158 = Ecballium Elaterium.
Elaterium, Linn. Sp. Pl. 1010 = Ecballium Elaterium.
elegans, Salisb. Prod. 158 = M. Charantia.
erinocarpa, Fenzl, ex Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 135 = Luffa echinata.
foetida, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 426.—Afr. trop.
Gabonii, *Cogn. in DC. Monog. Phan.* iii. 450.—Afr. trop.
garipensis, E. Mey. in Drège, Zwei Pfl. Docum. 202 = M. Balsamina.
garriepensis, Arn. in Hook. Journ. Bot. iii. (1841) 277 = praec.
glauca, Mor. ex Cogn. in DC. Monog. Phan. iii. 836 = Cyclanthera glauca.
guttata, Planch. ex Benth. in Hook. Niger Fl. 371 = cissoides.
Hamiltoniana, Wall. Cat. n. 6748 = dioica.
heterophylla, Wight, ex M. Roem. Syn. Pepon. 203 = Modecca Wightiana.
Heyneana, Wall. Cat. n. 6744 = dioica.
hispida, Dennst. Schluess. Hort. Malab. 34 = dioica.
Huberii, Todaro, in Giorn. Ist. Agric. Sicil. (1863) 165 = involucrata.
humilis, Wall. Cat. n. 6747 = M. Charantia.
Hystrix, Gill. in Hook. Bot. Misc. iii. (1833) 324 = Cyclanthera Hystrix.
involucrata, *E. Mey. in Drège, Zwei Pfl. Docum.* 202, nomen; *et ex Sond. in Harv. & Sond. Fl. Cap.* ii. 491.—Afr. austr.
jagorana, C. Koch, Wochenschr. i. (1858) Nachr. 44 = M. Charantia.
Lambertiana, Ser. in DC. Prod. iii. 311 = Cucumis trigonus.
lanata, Thunb. Prod. Pl. Cap. 13 = Citrullus vulgaris.

MOMORDICA:—
Luffa, Linn. Sp. Pl. 1009 = Luffa aegyptiaca.
Luffa, Vell. Fl. Flum. x. t. 93 = Luffa acutangula.
macropetala, Mart. in Hook. Lond. Journ. Bot. v. (1846) 504 = M. Charantia.
macropoda, Poepp. & Endl. Nov. Gen. et Sp. ii. 54 = Cyclanthera macropoda.
maculata, Planch. ex Benth. in Hook. Niger Fl. 370 = cissoides.
Mannii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 539.—Afr. trop.
missionis, Wall. Cat. n. 6739 = dioica.
mixta, Roxb. Hort. Beng. 70; Fl. Ind. iii. 709 = cochinchinensis.
monadelpha, Roxb. ll. cc. 70; 708 = Coccinia indica.
Morkorra, A. Rich. Tent. Fl. Abyss. i. 292. t. 53 = foetida.
multiflora, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 540.—Afr. trop.
muricata, Vell. Fl. Flum. x. t. 91 = Echinocystis muricata.
muricata, Willd. Sp. Pl. iv. 602 = M. Charantia.
Officinarum-elaterium, Crantz, Inst. i. 171 = M. Elaterium.
operculata, Blanco, Fl. Filip. ed. I. 770 = Luffa aegyptiaca.
operculata, Linn. Syst. ed. X. 1278 = Luffa operculata.
operculata, Vell. Fl. Flum. x. t. 92 = M. Charantia.
ovata, *Cogn. in DC. Monog. Phan.* iii. 446.—Ins. Philipp.
Paina, Buch.-Ham. ex Wall. Cat. n. 6742 = Actinostemma tenerum.
palmata, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 222 = Anguria Makoyana.
palmata, E. Mey. in Drège, Zwei Pfl. Docum. 202 = Coccinia palmata.
papillosa, Peckolt, ex Rosenth. Syn. Pl. Diaphor. 678 = M. Charantia.
pedata, Linn. Sp. Pl. 1009 = Cyclanthera pedata.
pedisecta, Ser. in DC. Prod. iii. 319, in syn. = Cyclanthera pedata.
pterocarpa, *Hochst. in Flora*, xxix. (1841) 1. *Intell.* 22, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 292.—Abyss.
purgans, Mart. Reise Bras. ii. 546 = Luffa operculata.
quinquefida, Hook. & Arn. Bot. Beech. Voy. 424 = Luffa operculata.
quinqueloba, E. Mey. in Drège, Zwei Pfl. Docum. 202 = Coccinia quinqueloba.
quinquelobata, Vell. Fl. Flum. x. t. 95 = Cyclanthera quinquelobata.
racemiflora, *Cogn. in DC. Monog. Phan.* iii. 448.—Sumatra.
racemosa, Steud. Nom. ed. II. ii. 155 = Echinocystis muricata.
renigera, Wall. Cat. n. 6743 = dioica.
reticulata, Salisb. Prod. 158 = Luffa aegyptiaca.
rotunda, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = Melothria perpusilla.
Roxburghiana, G. Don, Gen. Syst. iii. 35 = M. Charantia.
sativa, Roxb. ex Wight & Arn. Prod. 341 = Cucumis Melo.
Schimperiana, *Naud. in Ann. Sc. Nat.* Sér. V. v. (1866) 23.—Afr. trop.
senegalensis, Lam. Encyc. iv. 239 = M. Charantia.
sicyoides, Ser. in DC. Prod. iii. 312 = dioica.
sphaeroidea, Blanco, Fl. Filip. ed. I. 771 = ovata?
spicata, Linn. ex Sm. in Rees, Cycl. xxiii. n. 13 = Gymnopetalum cochinchinense.
subangulata, Blume, Bijdr. 928 = dioica.
surculata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 = Gymnopetalum cochinchinense.
Suringarii, *Cogn. in DC. Monog. Phan.* iii. 434.—Malaya.
trifolia, Linn. Syst. ed. X. 1278 = seq.
trifoliata, Linn. Sp. Pl. ed. II. 1434 = Trichosanthes trifoliata.
trifoliolata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 537.—Afr. trop.
trilobata, Wight, ex Steud. Nom. ed. II. ii. 155 = dioica.
tuberosa, Cogn. in DC. Monog. Phan. iii. 454 = M. Cymbalaria.
tuberosa, Dennst. Schluess. Hort. Malab. 34 = dioica.
tubiflora, Roxb. Hort. Beng. 70; Fl. Ind. iii. 711 = Gymnopetalum cochinchinense.
tubiflora, Wall. Cat. n. 6749 = Luffa acutangula.
tubulosa, Buch.-Ham. ex Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 136 = Zehneria umbellata.

MOMORDICA:—
umbellata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 710 = Melothria heterophylla.
verticillata, Vell. Fl. Flum. x. t. 96 = Wilbrandia verticillata.
Vogelii, Planch. ex Benth. in Hook. Niger Fl. 369 = foetida.
Wallichii, M. Roem. Syn. Pepon. 58 = dioica.
Welwitschii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 538.—Afr. trop.
zeylanica, Mill. Gard. Dict. ed. VIII. n. 3 = M. Charantia.

MONACANTHUS, G. Don, in Sweet, Hort. Brit. ed. III. 644 (1839) = MONACHANTHUS, Lindl. = **Catasetum**, Rich. (Orchid.).

MONACATHER, Benth. in Journ. Linn. Soc. xxi. (1881) 99 = MONACHATHER, Steud. = **Danthonia**, DC. (Gramin.).

MONACHANTHUS, Lindl. in Bot. Reg. sub t. 1538 (1832) = **Catasetum**, Rich. (Orchid.).
Bushnani, Hook. Bot. Mag. t. 3832 = C. discolor.
discolor, Lindl. Bot. Reg. t. 1735 = C. discolor.
fimbriatus, Gardn. ex Hook. Bot. Mag. t. 3708 = C. discolor.
longifolius, Lindl. in Bot. Mag. t. 3819 = C. longifolium.
roseo-albus, Hook. l. c. t. 3796 = C. discolor.
viridis, Lindl. in Bot. Reg. sub t. 1538 = C. trifidum.
viridis, Lindl. l. c. t. 1752 = C. macrocarpum.
viridis, Schomb. in Trans. Linn. Soc. xvii. (1837) 551, partim = C. barbatum.

MONACHATHER, Steud. Syn. Pl. Gram. 247 (1855) = **Danthonia**, DC. (Gramin.).
paradoxus, Steud. l. c. = D. bipartita.

MONACHNE, Beauv. Agrost. 49. t. 10 (1812) = **Eriochloa**, H. B. & K. (Gramin.).
racemosa, Beauv. l. c. 168. t. 10. f. 10 = Panicum reptans.
rufa, Bertol. Misc. Bot. xi. 10 = Panicum reptans?
unilateralis, Beauv. Agrost. 49. t. 10. f. 9 = E. polystachya.
unilateralis, Roem. & Schult. Syst. ii. 468 = Panicum leucophaeum.

MONACHOCHLAMYS, Baker, in Journ. Linn. Soc. xx. (1883) 217. *ACANTHACEAE*.
flagellaris, *Baker, l. c.*—Madag.

MONACHYRON, Parl. in Hook. Niger Fl. 190 (1849). *GRAMINEAE*, Benth. & Hook. f. iii. 1158.
villosum, *Parl. l. c.* 191.—Ins. Cap. Virid.

MONACTINEIRMA, Bory, in Ann. Sc. Gén. Phys. ii. (1819) 138 = **Passiflora**, Linn.
angustifolia, Bory. l. c. = P. suberosa.
coriacea, Bory, l. c. = P. coriacea.
hederacea, Bory, l. c. = P. suberosa.
mexicana, Bory, l. c. = P. mexicana.
minima, Bory, l. c. = P. suberosa.
peltata, Bory, l. c. = P. suberosa.
suberosa, Bory, l. c. = P. suberosa.

MONACTIS, H. B. & K. Nov. Gen. et Sp. iv. 286. t. 403 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 358.
dubia, H. B. & K. l. c. 287 = Astemma dubium.
Flaveriae, DC. Prod. v. 546 = flaverioides.
flaverioides, *H. B. & K. Nov. Gen. et Sp.* iv. 287. t. 403.—Bras.

MONADELPHANTHUS, Karst. Fl. Columb. i. 67. t. 33 (1859?). *RUBIACEAE*, Benth. & Hook. f. ii. 38.
floridus, *Karst. l. c.*—N. Granat.

MONADENIA, Lindl. Gen. et Sp. Orch. 356 (1838) = **Disa**, Berg (Orchid.).
brevicornis, Lindl. l. c. 357.—Afr. austr.
comosa, Reichb. f. in Linnaea, xx. (1847) 687 = D. affinis.
densiflora, Lindl. Gen. et Sp. Orch. 357.—Afr. austr.
inflata, Sond. in Linnaea, xix. (1847) 102 = D. cernua.
lancifolia, Sond. l. c. 100 = *ophrydea*.
leptostachys, Sond. l. c. 101 = D. rufescens.

MONADENIA:—
macrocera, Lindl. Gen. et Sp. Orch. 358 = Disa rufescens.
macrostachya, Lindl. l. c. 357.—Afr. austr.
micrantha, Lindl. l. c.—Afr. austr.
multiflora, Sond. in Linnaea, xix. (1847) 101.—Afr. austr.
ophrydea, Lindl. Gen. et Sp. Orch. 358.—Afr. austr.
physodes, Reichb. f. in Flora, lxvi. (1883) 461 = D. physodes.
prasinata, Lindl. Gen. et Sp. Orch. 358 = D. cernua.
rufescens, Lindl. l. c. 356 (excl. syn.) = D. affinis.
rufescens, Reichb. f. in Flora, lxvi. (1883) 464 = D. rufescens.

MONADENUS, Salisb. Gen. Pl. Fragm. 51 (1866) = **Zygadenus**, Michx. (Liliac.).

MONANDRAIRA, E. Desv. in C. Gay, Fl. Chil. vi. 341. t. 79 (1881) = **Deschampsia**, Beauv. (Gramin.).
Berteroana, E. Desv. l. c. 343 = D. Berteroana.
glauca, E. Desv. l. c. 342.—Chili.
patula, Phil. in Anal. Univ. Chil. (1873) 565.—Chili.

MONANTHEMUM, Griseb. Fl. Brit. W. Ind. 354 (1861) = **Piptocarpha**, R. Br. (Compos.).
Cruegerii, Griseb. l. c. = P. Cruegeri.

MONANTHES, Haw. Rev. Pl. Succ. 68 (1821). *CRASSULACEAE*, Benth. & Hook. f. i. 660.
PETROPHYES, Webb & Berth. Phyt. Canar. i. 201. t. 36 (1836–45).
atlantica, *Ball, in Journ. Bot.* xi. (1873) 333.—Marocc.
brachycaulon, *Lowe, Fl. Salv. Tent.* 12.—Ins. Salv.
muralis, Hook. f. in Bot. Mag. t. 5988 = atlantica.
polyphylla, Haw. Rev. Pl. Succ. 68.—Ins. Madera.
spathulata, Cree, ex Steud. Nom. ed. II. ii. 155 = Sempervivum micranthes.

MONANTHOCHLOË, Engelm. in Trans. Acad. Sc. St. Louis i. (1859), 436. tt. 13, 14. *GRAMINEAE*, Benth. & Hook. f. iii. 1180.
littoralis, *Engelm. l. c.*—Texas.

MONARDA, Linn. Gen. ed. I. 6 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1197.
CHEILYCTIS, Benth. Lab. Gen. et Sp. 726 (1836).
CORYANTHUS, Benth. l. c. (1836).
affinis, Link, Enum. Hort. Berol. i. 9 = fistulosa.
albiflora, *Morr. in Belg. Hortic.* i. (1851) 40.—Am. bor.
allophylla, Michx. Fl. Bor. Am. i. 16 = fistulosa.
altissima, Willd. Enum. Hort. Berol. 33 = clinopodia.
amplexicaulis, Fisch. ex Morr. in Belg. Hortic. i. (1851) 41 = Bradburiana.
aristata, Hook. Bot. Mag. t. 3526 = clinopodioides.
aristata, Nutt. in Trans. Am. Phil. Soc. v. (1837) 186 = citriodora.
barbata, Wender. Ind. Sem. Hort. Marb. (1828); et in Flora, xii. (1829) i. Erg. 27 = fistulosa.
Beckii, *Eaton, Man. Bot.* ed. VI. 229.—Am. bor.
Bradburiana, *Beck, in Am. Journ. Sc.* x. (1826) 260.—Am. bor.
ciliata, Linn. Sp. Pl. 23 = Blephilia ciliata.
ciliata, Michx. Fl. Bor. Am. i. 16 = Blephilia hirsuta.
citriodora, *Cerv. ex Lag. Gen. et Sp. Nov.* 2.—Am. bor. occ.; Mexic.
clinopodia, *Linn. Sp. Pl.* 22.—Am. bor.
clinopodioides, *A. Gray, Syn. Fl. N. Am.* ii. i. 375.—Texas.
coccinea, Michx. Fl. Bor. Am. i. 16 = didyma.
coerulea, Hort. ex Benth. Lab. Gen. et Sp. 317 = fistulosa.
commutata, Wender. Ind. Sem. Hort. Marb. (1828); et in Flora, xii. (1829) i. Erg. 28 = fistulosa.
contorta, *Morr. in Belg. Hortic.* i. (1851) 37. t. 6. f. a.—Am. bor.
cristata, Hort. ex Benth. Lab. Gen. et Sp. 317 = fistulosa.
didyma, *Linn. Sp. Pl.* 22.—Am. bor.
didynama, Stokes, Bot. Mat. Med. i. 44 = praec.
dubia, Hort. ex Benth. Lab. Gen. et Sp. 317 = fistulosa.
elongata, A. Dietr. Sp. Pl. i. 356, in syn. = fistulosa.
fistulosa, Hook. Bot. Mag. t. 3310 = Bradburiana.
fistulosa, Curt. l. c. t. 145 = didyma.
fistulosa, *Linn. Sp. Pl.* 22.—Am. bor.
glabra, Lam. Encyc. iv. 256 = clinopodia.

MONARDA:—
gracilis, Pursh, Fl. Am. Sept. i. 17 = Pycnanthemum montanum?
hirsuta, Pursh, l. c. 19 = Blephilia hirsuta.
hybrida, Wender. Ind. Sem. Hort. Marb. (1828); et in Flora, xii. (1829) i. Erg. 28 = fistulosa.
involucrata, Wender. ll. cc. = fistulosa.
Kalmiana, Pursh, Fl. Am. Sept. i. 17 = didyma.
lilacina, Wender. Ind. Sem. Hort. Marb. (1828); et in Flora, xii. (1329) i. Erg. 28 = fistulosa.
Lindheimeri, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 228 = fistulosa.
longifolia, Lam. Encyc. iv. 255 = fistulosa.
lutea, Michx. Fl. Bor. Am. i. 16 = punctata.
media, Willd. Enum. Hort. Berol. 32 = fistulosa.
menthaefolia, R. Grah. in Edinb. N. Phil. Journ. (Jul.–Oct. 1829) 347 = fistulosa.
mollis, Linn. Amoen. Acad. iii. 399 = fistulosa.
oblongata, [Soland. in] Ait. Hort. Kew. ed. I. i. 36 = fistulosa.
oswegoënsis, Bart. Prod. Fl. Pensylv. i. 34; ex Benth. Lab. Gen. et Sp. 726 = didyma.
pectinata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847), 82.—N. Mexic.
penicillata, A. Gray, in Proc. Am. Acad. viii. (1821) 369, sphalm. = praec.
pratensis, *Rafin. Med. Fl.* ii. 37.—Am. bor.
punctata, *Linn. Sp. Pl.* 22.—Am. bor.
purpurascens, Dum.-Cours. Bot. Cult. ed. II. iii. 4 = fistulosa.
purpurascens, Wender. Ind. Sem. Hort. Marb. (1828); et in Flora, xii. (1829) i. Erg. 28 = didyma.
purpurea, Lam. Encyc. iv. 256 = didyma.
purpurea, Pursh, Fl. Am. Sept. i. 17 = fistulosa.
rigida, *Rafin. Med. Fl.* ii. 37.—Am. bor.
rugosa, [Soland. in] Ait. Hort. Kew. ed. I. i. 36 = clinopodia.
Russeliana, *Nutt. in Journ. Trav. Arkans. Terr.* 141, nomen; *et in Trans. Am. Phil. Soc.* v. (1837) 185.—Am. bor. occ.
scabra, Beck, in Am. Journ. Sc. x. (1826) 260 = fistulosa.
undulata, Tausch, ex Reichb. Hort. Bot. ii. 31. t. 181 = fistulosa.
varians, Bart. Prod. Fl. Pensylv. i. 36; ex Benth. Lab. Gen. et Sp. 726 = fistulosa.
villosa, Martens, in Bull. Acad. Brux. viii. (1841) 66 = Bradburiana.
violacea, Desf. Tabl. ed. II. 66 = fistulosa.
virgata, *Rafin. Med. Fl.* ii. 37.—Am. bor.
virginalis, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 709 = fistulosa.
zeylanica, Burm. f. Fl. Ind. 12 = Anisomeles ovata.

MONARDELLA, Benth. Lab. Gen. et Sp. 331 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1185.
Breweri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 386.—Calif.
candicans, *Benth. Pl. Hartw.* 330.—Calif.
caroliniana, Benth. Lab. Gen. et Sp. 332 = Monarda clinopodia.
Douglasii, *Benth. l. c.*—Calif.
glabra, Nutt. ex Benth. in DC. Prod. xii. 190 = odoratissima.
hypoleuca, *A. Gray, Syn. Fl. N. Am.* ii. i. 356.—Calif.
lanceolata, *A. Gray, in Proc. Am. Acad.* xi. (1876) 102.—Calif.
leucocephala, *A. Gray, l. c.* vii. (1868) 385.—Calif.
linoides, *A. Gray, l. c.* xi. (1876) 101.—Calif.
macrantha, *A. Gray, l. c.* 100.—Calif.
montana, Benth. Lab. Gen. et Sp. 331 = Pycnanthemum montanum.
nana, *A. Gray, in Proc. Am. Acad.* xi. (1876) 101.—Calif.
odoratissima, *Benth. Lab. Gen. et Sp.* 332.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 82.—Calif.
Pringlei, *A. Gray, l. c.* xix. (1883) 96.—Calif.
Sheltoni, Torr. ex Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 99 = villosa.
tenuiflora, S. Wats. ex A. Gray, in Proc. Am. Acad. xvii. (1881–82) 230 = macrantha.
thymifolia, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 211.—Calif.
undulata, *Benth. Lab. Gen. et Sp.* 332.—Calif.
villosa, *Benth. Bot. Voy. Sulph.* 42. t. 21.—Calif.

MONARRHENUS, Cass. in Bull. Soc. Philom. (1817) 31. *COMPOSITAE*, Benth. & Hook. f. ii. 295.
MAHOMETA, DC. Prod. v. 457, in syn. (1836).
pinifolius, *Cass. in Dict. Sc. Nat.* xxxii. 434.—Ins. Borbon.
rufescens, *DC. Prod.* v. 457.—Ins. Maurit.
salicifolius, *Cass. in Dict. Sc. Nat.* xxxii. 434.—Ins. Maurit.

MONATHERA, Rafin. in Am. Monthly Mag. (1819) 190 = **Ctenium**, Panz. (Gramin.).

MONAVIA, Adans. Fam. ii. 211 (1763) = **Mimulus**, Linn. (Scrophul.).

MONBIN, Plum. ex Adans. Fam. ii. 341 (1763) = **Spondias**, Linn. (Anacard.).

MONDO, Adans. Fam. ii. 496 (1763) = **Carex**, Dill. (Cyperac.).

MONECHMA, Hochst. in Flora, xxiv. (1841) 374 = **Justicia**, Linn. (Acanth.).
affine, Hochst. l. c. xxvi. (1843) 76 = J. debilis.
angustifolium, Nees, in DC. Prod. xi. 411 = J. debilis.
bracteatum, Hochst. in Flora, xxiv. (1841) 375 = J. debilis.
ciliatum, Hochst. ex Nees, in DC. Prod. xi. 411 = J. violacea.
debile, Nees, l. c. = J. debilis.
hispidum, Hochst. in Flora, xxiv. (1841) 375 = Schwabea ciliaris.
Pilosella, Nees, in DC. Prod. xi. 412 = Siphonoglossa Pilosella.
violaceum, Nees, l. c. 411 = J. violacea.

MONELLA, Herb. App. 29 (1821) = **Cyrtanthus**, Ait. (Amaryll.).
angustifolia, Herb. l. c. = C. angustifolius.
collina, Salisb. Gen. Pl. Fragm. 139 = C. collinus.
glauca, Herb. App. 29 = C. collinus.
ochroleuca, Herb. l. c. = C. lutescens.
odora, Herb. l. c. = C. odorus.
pallida, Herb. ex Kunth, Enum. Pl. v. 536 = C. pallidus.
spiralis, Herb. App. 29 = C. spiralis.
ventricosa, Herb. l. c. = C. angustifolius.

MONENCYANTHES, A. Gray, in Hook. Kew Journ. iv. (1852) 229 = **Helipterum**, DC. (Compos.).
gnaphaloides, A. Gray, l. c. 230 = H. moschatum.

MONENTELES, Labill. Sert. Austro-Caled. 42. tt. 43, 44 (1825) = **Pterocaulon**, Ell. (Compos.).
glandulosus, F. Muell. ex Benth. Fl. Austral. iii. 523 = P. glandulosus.
globiferus, DC. Prod. v. 455 = P. sphacelatus.
intermedius, DC. l. c. = P. sphacelatus.
Pterocaulon, DC. l. c. = P. Bojeri.
redolens, DC. l. c.—N. Caled.
serrulatus, Montr. in Mém. Acad. Lyon, x. (1860) 225.—N. Caled.
sphacelatus, Labill. Sert. Austro-Caled. 43. t. 44 = P. sphacelatus.
sphaeranthoides, DC. Prod. v. 556 = P. sphaeranthoides.
spicatus, Labill. Sert. Austro-Caled. 43. t. 43 = P. cylindrostachynum.
tomentosus, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 122, nomen.—Malaya.
verbascifolius, F. Muell. ex Benth. Fl. Austral. iii. 523 = P. verbascifolius.

MONERMA, Beauv. Agrost. 116 (1812) = **Psilurus**, Trin. (Gramin.).
cylindrica, Coss. & Dur. Expl. Alg. 214 = Lepturus cylindricus.
Gangitis, Roem. & Schult. Syst. ii. 800 = Ctenium carolinianum.
monandra, Beauv. Agrost. 117 = Psilurus nardoides.
ramosa, Schult. Mant. ii. 441 = Lepturus ramosus.
repens, Beauv. Agrost. 117 = Lepturus repens.
simplex, Gaudich. in Freyc. Voy. Bot. 412 = Lepturus cylindricus?
stricta, Presl, ex Steud. Nom. ed. II. 156 = Lolium perenne.
subulata, Beauv. Agrost. 117 = Lepturus cylindricus.

MONESES, Salisb. ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 403 (1821). *ERICACEAE*, Benth. & Hook. f. ii. 603.
BRYOPHTHALMUM, E. Mey. Preuss. Pflanzeng. 101 (1839).
brevicaulis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 49 ; x. (1859) 139 = grandiflora.
grandiflora, *S. F. Gray, Nat. Arr. Brit. Pl.* ii. 403.—Reg. bor. et arct.
reticulata, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 271 = grandiflora.
uniflora, A. Gray, Man. Bot. N. U. St. ed. I. 273 = grandiflora.
verticillata, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 170 = grandiflora.

MONESIS, Walp. Rep. ii. 734 (1843), err. typ. = praec.

MONETIA, L'Hérit. Stirp. Nov. 1. t. 1 (1784) = **Azima**, Linn. (Salvad.).
angustifolia, Boj. ex A. DC. in DC. Prod. xvii. 29 = A. angustifolia.
barlerioides, L'Hérit. Stirp. Nov. 1. t. 1 = A. tetracantha.
Brunoniana, Wall. Cat. n. 7493 = A. sarmentosa.
diacantha, Willd. Sp. Pl. i. 670 = Plectronia parviflora.
laxa, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 191 = A. sarmentosa.
paleacea, Spreng. Syst. i. 442.—Afr. austr.
sarmentosa, Baill. Adansonia, ix. (1870) 289. t. 10 = A. sarmentosa.
tetracantha, Salisb. Prod. 65 = A. tetracantha.
triphylla, Dennst. Schluess. Hort. Malab. 31 = Plectronia didyma.

MONGESIA, Miers, in Journ. Linn. Soc. xvii. (1879) 285, sphalm. = seq.

MONGEZIA, Vell. Fl. Flum. 229 (1825) = **Symplocos**, Linn. (Styrac.).
glabra, Vell. l. c. ; v. t. 106 = S. laxiflora.
glabrata, Steud. Nom. ed. II. ii. 157, sphalm. = praec.
pilosa, Vell. Fl. Flum. 229 ; v. t. 105 = S. pubescens.

MONGORIUM, Desf. Fl. Atlant. i. 5 (1798) = MOGORIUM, Juss. = **Jasminum**, Tourn. (Oleac.).

MONGUIA, Chapel. ex Baill. Adansonia, i. (1860–61) 147 = **Croton**, Linn. (Euphorb.).
cordifolia, Chapel. ex Baill. l. c. 158 = C. Mongue.
lanceolata, Chapel. ex Baill. l. c. 147 = C. Chrysodaphne.

MONIERA, B. Juss. ex P. Br. Hist. Jamaic. 269 (1756) = **Herpestis**, Gaertn. (Scrophul.).

MONIERA, Loefl. It. Hisp. 197, et MONIERIA, Loefl. l. c. 176 (1758) = **Monnieria**, Linn. (Rutac.).

MONILIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 110 (1821) = **Molinia**, Schrank (Gramin.).

MONILIFERA, Adans. Fam. ii. 127 (1763) = **Osteospermum**, Linn. (Compos.).

MONILISTUS, Rafin. Alsog. Am. 42 (1838) = **Populus**, Tourn. (Salic.).
monilifera, Rafin. l. c. = P. canadensis.

MONIMIA, Thou. Hist. Vég. Isles de Fr. 35. t. 9 (1804). *MONIMIACEAE*, Benth. & Hook. f. iii. 139.
citrina, *Tul. in Ann. Sc. Nat.* Sér. IV. iii. (1855) 32.—Ins. Borbon.
Lastelliana, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 342.—Madag.
ovalifolia, *Thou. Hist. Vég. Isles de Fr.* t. 9. f. 1.—Ins. Mascar.
rotundifolia, *Thou. l. c.* f. 2.—Ins. Mascar.

MONINA, Pers. Syn. ii. 270 (1807) = **Monnina**, Ruiz & Pav. (Polygal.).

MONIPSIS, Rafin. Fl. Tellur. iii. 85 (1836) = **Teucrium**, Tourn. (Labiat.).
orchidea, Rafin. l. c. = T. bicolor.

MONIZIA, Lowe, in Hook. Kew Journ. viii. (1856) 295 = **Thapsia**, Linn. (Umbell.).
edulis, Lowe, l. c.—Ins. Mader.

MONNELLA, Salisb. Gen. Pl. Fragm. 139 (1866) = MONELLA, Herb. = **Cyrtanthus**, Ait. (Amaryll.).

MONNERIA, Spreng. Syst. iii. 157 (1826) = **Monnieria**, Linn. (Rutac.).

MONNIERA, B. Juss. ex P. Br. Hist. Jamaic. 269 (1756) (Moniera) = **Herpestis**, Gaertn. f. (Scrophul.).
africana, Pers. Syn. ii. 166 = H. Monnieria.
amplexicaulis, Michx. Fl. Bor. Am. ii. 22 = H. amplexicaulis
Brownei, Pers. Syn. ii. 166 = H. Monnieria.
cuneifolia, Michx. Fl. Bor. Am. ii. 22 = H. Monnieria.
graveolens, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 308 = H. depressa.
pedunculosa, Pers. Syn. ii. 166 = H. Monnieria.
rotundifolia, Michx. Fl. Bor. Am. ii. 22 = H. rotundifolia.

MONNIERA, Juss. Gen. 421 (1789) = seq.

MONNIERIA, Linn. Syst. ed. X. 1153 (1759). *RUTACEAE*, Benth. & Hook. f. i. 286.
AUBLETIA, Rich. in Pers. Syn. ii. 638 (1807).
ERTELA, Adans. Fam. ii. 358 (1763).
MONIERA, Loefl. It. Hisp. 197 (1758).
bahiensis, *Engl. in Mart. Fl. Bras.* xii. II. 129.—Bras.
semiserrata, *Schrank, in Syll. Ratisb.* ii. (1828) 63 (Quid ?).—Bras.
subserrata, Mart. ex Schrank, l. c. = praec.
trifolia, *Linn. Syst.* ed. X. 1153.—Am. austr.

MONNINA, Ruiz & Pav. Syst. Veg. 169 (1798). *POLYGALEAE*, Benth. & Hook. f. i. 139.
HEBEANDRA, Bonpl. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 40.
aestuans, *DC. Prod.* i. 338.—N. Granat.
angustata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 140.—N. Granat.
angustifolia, *DC. Prod.* i. 340.—Peruv.
angustifolia, Schlecht. in Linnaea, xiv. (1840) 380 = Schlechtendaliana.
bifurcata, [*Moç. & Sesse*] *ex DC. Prod.* i. 339.—Mexic.
brachystachya, *Griseb. in Goett. Abh.* xix. (1874) 75.—Reg. Argent.
caerulea, *G. Don, Gen. Syst.* i. 367.—Mexic.
calophylla, Poepp. & Endl. Nov. Gen. et Sp. iii. 66 = Securidaca Corytholobium.
cardiocarpa, *A. St. Hil. Fl. Bras. Mer.* ii. 60. t. 93.—Bras.
cestrifolia, *H. B. & K. Nov. Gen. et Sp.* v. 413. t. 502.—Am. austr.
ciliolata, [*Moç. & Sesse*] *ex DC. Prod.* i. 340.—Mexic.
ciliolosa, G. Don, Gen. Syst. i. 368 = praec.
cladostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 357.—Venezuela.
conferta, *Ruiz & Pav. Syst. Veg.* 173.—Peruv.
cordata, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 192.—Bras.
crassifolia, *H. B. & K. Nov. Gen. et Sp.* v. 411.—Ecuador.
crassinervia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 143.—N. Granat.
crotalariaefolia, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 590 = seq.
crotalarioides, *DC. Prod.* i. 339.—Am. austr.
cuneata, Herb. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 56 = insignis.
cuneata, *A. St. Hil. Fl. Bras. Mer.* ii. 62. t. 95.—Bras.
cuspidata, *Benth. Pl. Hartw.* 162.—Ecuador.
densa, *Planch. & Linden, ex Wedd. Chlor. And.* ii. 268.—N. Granat.
Deppei, *G. Don, Gen. Syst.* i. 367.—Mexic.
dictyocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 23.—Reg. Argent.
diversifolia, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 58 = cuneata.
elongata, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 137.—N. Granat.
emarginata, *A. St. Hil. Fl. Bras. Mer.* ii. 63.—Bras.
epilobioides, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 58 = Richardiana.
euonymoides, *Schlecht. in Linnaea*, xiv. (1840) 380.—Mexic.
exaltata, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 59.—Bras.
excelsa, Spreng. Syst. iv. Cur. Post. 265 = Xanthophyllum excelsum.
fastigiata, *DC. Prod.* i. 338.—N. Granat.
floribunda, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 139.—N. Granat.

MONNINA :—
herbacea, *DC. Prod.* i. 340.—Peruv.
Hilairiana, Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 192 = cuneata.
insignis, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 56.—Bras.
lanceolata, *DC. Prod.* i. 339.—Peruv.
lancifolia, *G. Don, Gen. Syst.* i. 367.—Bras.
latifolia, *DC. Prod.* i. 338.—N. Granat.
leptostachya, *Benth. Pl. Hartw.* 125.—Peruv.
ligustrifolia, *H. B. & K. Nov. Gen. et Sp.* v. 417.—Ecuador.
linearifolia, *Ruiz & Pav. Syst. Veg.* 173.—Chili.
longifolia, *DC. Prod.* i. 339.—Bras.
longifolia, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 57 = stenophylla.
longifolia, Spreng. Syst. iv. Cur. Post. 265 = Xanthophyllum longifolium.
loxensis, *Benth. Pl. Hartw.* 125.—Am. austr.
luteola, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 57 = Tristaniana.
macrophylla, Steud. Nom. ed. II. ii. 157 = Xanthophyllum longifolium.
macrostachya, *Ruiz & Pav. Syst. Veg.* 173.—Peruv.
marginata, *Presl, Rel. Haenk.* ii. 102.—Peruv.
Martiana, *Klotzsch, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 59.—Bras.
membranacea, *Miq. Stirp. Surin. Sel.* 30.—Guiana.
meridensis, *Planch. & Linden, ex Wedd. Chlor. And.* ii. 268.—Venezuela.
mexicana, *G. Don, Gen. Syst.* i. 367.—Mexic.
mollis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 139.—N. Granat.
myrtilloides, *DC. Prod.* i. 339.—Peruv.
nemorosa, *H. B. & K. Nov. Gen. et Sp.* v. 416. t. 504.—Ecuador.
nervosa, Steud. Nom. ed. II. i. 724, in syn. = praec.
obscura, *G. Don, Gen. Syst.* i. 367.—Mexic.
obtusifolia, *H. B. & K. Nov. Gen. et Sp.* v. 411.—N. Granat.
Olfersiana, Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 192 = Richardiana.
pallida, *Spreng. Syst.* iii. 175.—Bras.
paniculata, *Benth. Pl. Hartw.* 126.—Peruv.
parviflora, *H. B. & K. Nov. Gen. et Sp.* v. 419.—N. Granat.
phytolaccaefolia, *H. B. & K. l. c.* 413.—Am. austr.
pilosa, *H. B. & K. l. c.* 419.—Ecuador.
polystachya, *Ruiz & Pav. Syst. Veg.* 171.—Peruv.
pterocarpa, *Ruiz & Pav. l. c.* 174.—Peruv.
pubescens, *H. B. & K. Nov. Gen. et Sp.* v. 418.—Venezuela.
resedoides, *A. St. Hil. Fl. Bras. Mer.* ii. 61. t. 94.—Bras.
retusa, *Presl, Rel. Haenk.* ii. 102.—Chili.
revoluta, *H. B. & K. Nov. Gen. et Sp.* v. 412.—Peruv.
Richardiana, *A. St. Hil. Fl. Bras. Mer.* ii. 66.—Bras.
rupestris, *H. B. & K. Nov. Gen. et Sp.* v. 415.—N. Granat.
salicifolia, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 58 = Richardiana.
salicifolia, *Ruiz & Pav. Syst. Veg.* 172.—Peruv.
Schlechtendaliana, *D. Dietr. Syn. Pl.* iv. 912.—Mexic.
Selloi, *Spreng. Syst.* iii. 175.—Bras.
semiserrata, Steud. Nom. ed. II. ii. 157, sphalm. = Monnieria semiserrata.
solandraefolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 138.—N. Granat.
speciosa, *Triana & Planch. l. c.* 144.—N. Granat.
stenophylla, *A. St. Hil. Fl. Bras. Mer.* ii. 64.—Bras.
stenophylla, Steud. Nom. ed. II. ii. 157 = Schlechtendaliana.
subscandens, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 143.—N. Granat.
sylvatica, *Cham. & Schlecht. in Linnaea*, v. (1830) 231.—Mexic.
Tristaniana, *A. St. Hil. Fl. Bras. Mer.* ii. 65.—Bras.
tuberosa, *G. Don, Gen. Syst.* i. 367.—Bras.
vitellina, Spreng. Syst. iv. Cur. Post. 265 = Xanthophyllum vitellinum.
Wrightii, *A. Gray, Pl. Wright*, ii. 31.—N. Mexic.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* v. 414.—Mexic.

MONNURIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) t. 18, sphalm. = **Monnieria**, Linn. (Rutac.).

MONOBOTHRIUM, Hochst. in Flora, xxvii. (1844) 27 = **Swertia**, Linn. (Gentian.).
Schimperi, Hochst. l. c. 28 = S. Schimperi.

MONOCALLIS, Salisb. Gen. Pl. Fragm. 27 (1866) = **Scilla**, Linn. (Liliac.).
monophylla, Salisb. l. c. 28 = S. monophyllos.

MONOCARPIA, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 12 (1865-66). *ANONACEAE*, Benth. & Hook. f. i. 956.
Blancoi, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 6.—Ins. Philipp.
cuneura, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 12.—Borneo.

MONOCARYUM, R. Br. in Denh. & Clapp. Trav. App. 243 (1826) = **Colchicum**, Tourn. (Liliac.).
fasciculare, Schult. f. Syst. vii. 1135 = C. montanum.

MONOCERA, Ell. Sketch, i. 176 (1817) = **Ctenium**, Panz. (Gramin.).
aromatica, Ell. l. c. i. 177 = C. carolinianum.

MONOCERA, Jack, in Malay. Misc. i. (1820) n. v. 42 = **Elaeocarpus**, Burm. (Tiliac.).
aristata, Jack, l. c. 43, in obs. = E. aristatus.
atrovirens, Miq. Fl. Ind. Bat. i. II. 213 = E. atrovirens.
bilocularis, Jack, in Malay. Misc. i. (1820) n. v. 43, in obs. = F. tuberculatus.
ceylanica, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 323 = E. ceylanicus.
cuneifolia, Miq. Fl. Ind. Bat. Suppl. 409.—Sumatra.
cyanea, Hassk. Cat. Hort. Bog. Alt. 208 = E. cyaneus.
cyrtostachya, Miq. Fl. Ind. Bat. Suppl. 408.—Sumatra.
densiflora, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 236, in obs. = E. ellipticus?
elliptica, Turcz. l. c. = E. Varunua.
ferruginea, Jack, in Malay. Misc. i. (1820) n. v. 44; Wight, Ic. t. 205 = E. ferrugineus.
glandulifera, Hook. ex Wight, Illustr. i. 84 = E. glandulifer.
grandiflora, Hook. Bot. Mag. t. 4680 = E. grandiflorus.
Griffithii, C. Muell. Annot. Elaeocarp. 12 = E. paniculatus.
Griffithii, Wight, Illustr. i. 84 = E. Griffithii.
holopetala, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 496 = E. Griffithii.
integra, C. Muell. Annot. Elaeocarp. 12 = E. integer.
isotricha, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 494 = E. isotrichus.
lanceolata, Hassk. Cat. Hort. Bog. Alt. 208 = E. lanceolatus.
leucobotrya, Miq. Fl. Ind. Bat. Suppl. 409 = E. paniculatus.
luzonica, Miq. Fl. Ind. Bat. i. II. 213 = E. Monocera.
macrocera, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 494 = E. Monocera.
multiflora, Turcz. l. c. 495 = E. multiflorus.
Munroii, Wight, Illustr. i. 84 = E. Munroii.
obtusa, Hassk. Cat. Hort. Bog. Alt. 208 = E. obtusus.
odontopetala, Miq. Fl. Bat. Suppl. 409.—Sumatra.
palembanica, Miq. l. c. 408.—Sumatra.
paniculata, Wall. ex A. Gray, Bot. U. St. Expl. Exp. i. 203 = E. Griffithii.
petiolaris, Steud. Nom. ed. II. ii. 157, sphalm. = seq.
petiolata, Jack, in Malay. Misc. i. (1820) n. v. 43 = E. integer.
prunifolia, C. Muell. Annot. Elaeocarp. 15 = E. prunifolius.
robusta, Miq. Fl. Ind. Bat. i. II. 212.—Java.
Roxburghii, Wight, Ic. t. 63 = E. aristatus.
rugosa, Wight, Ic. t. 61 = E. rugosus.
serrata, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 495, in obs. = E. serratus.
subintegerrima, Miq. ex Mast. in Hook. f. Fl. Brit. Ind. i. 405 = E. tuberculatus.
tricanthera, Griff. Notul. iv. 518 = E. Griffithii.
tuberculata, Wight & Arn. Prod. 83 = E. tuberculatus.
Walkerii, Wight, Illustr. i. 84 = E. ceylanicus.

MONOCERAS, Auct. ex Steud. Nom. ed. II. ii. 159 (1841) = praec.

MONOCERAS, Steud. Nom. ed. I. 537 (1821) = **Velleia**, Sm. (Gooden.).

MONOCHAETE, Doell, in Mart. Fl. Bras. ii. III. 78. t. 22 (1878). *GRAMINEAE*, Benth. & Hook. f. iii. 1167.
fastigiata, *Doell, l. c.* 79.—Bras.

MONOCHAETUM, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 48. t. 2. *MELASTOMACEAE*, Benth. & Hook. f. i. 747.
EPHYNES, Rafin. Sylva Tellur. 101 (1838).
GRISCHOWIA, Karst. Ausw. Neuer Gew. Venez. 15. t. 5 (1848).
LOEVIGIA, Karst. & Triana, Fl. Neo-Granad. 22; ex Linnaea, xxviii. (1856) 434.
alpestre, *Naud. in Ann. Sc. Nat.* Sér. III. iv. (1845) 50.—Mexic.
Bonplandii, *Naud. l. c.* 51; *et* xiv. (1850) 165.—Am. trop.
brachyurum, *Naud. l. c.* xiv. 162.—N. Granat.
bracteolatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 64.—Panama.
calcaratum, Triana, l. c. 63 = Candolleanum.
Candolleanum, *Naud. in Ann. Sc. Nat.* Sér. III. iv. (1845) 51.—Mexic.
Deppeanum, *Naud. l. c.* (1850) 165.—Mexic.
dicranantherum, *Naud. l. c.* iv. (1845) 52.—Peruv.
dicranantherum, Triana, ex Hook. f. Bot. Mag. t. 5506 = Hartwegianum.
discolor, *Karst. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 63.—Venezuela.
ensiferum, Hook. Bot. Mag. t. 5132 = alpestre.
ensiferum, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 50 = Candolleanum.
floribundum, Naud. l. c. xiv. (1850) 165 = rivulare.
fuscescens, Naud. l. c. iv. (1845) 50 = lineatum.
glanduliferum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 62.—N. Granat.
Hartwegianum, *Naud. in Ann. Sc. Nat.* Sér. III. iv. (1845) 51.—N. Granat.
hirtum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 64.—Venezuela.
Humboldtianum, Seem. Bot. Voy. Herald, 122 = bracteolatum.
Humboldtianum, *Walp. Rep.* v. 702.—Venezuela.
latifolium, *Naud. in Ann. Sc. Nat.* Sér. III. iv. (1845) 49.—Venezuela.
Lindenianum, *Naud. l. c.* xiv. (1850) 158.—N. Granat.
lineatum, *Naud. l. c.* iv. (1845) 52.—Am. trop.
meridense, *Naud. l. c.* xiv. (1850) 165.—Venezuela; N. Granat.
multiflorum, *Naud. l. c.* iv. (1845) 52.—N. Granat.
myrtoideum, *Naud. l. c.* 51.—Mexic.; N. Granat.
myrtoideum, Seem. Bot. Voy. Herald, 284 = Candolleanum.
Naudinianum, Neum. in Rev. Hortic. (1861) 211 = alpestre.
oliganthum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 159 = Candolleanum.
osbeckioides, Naud. l. c. 165 = Humboldtianum.
pauciflorum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 63.—Ecuador.
plumosum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 160 = pulchrum.
polyneurum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 62.—N. Granat.
pulchellum, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 161 = pulchrum.
pulchrum, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 101. f. 6.—Mexic.
quadrangulare, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 63.—N. Granat.
rivulare, *Naud. in Ann. Sc. Nat.* Sér. III. iv. (1845) 50.—Am. centr.
sericeum, Naud. ex Linden, in Belg. Hortic. viii. (1858) 169 = Bonplandii.
stellulatum, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 160.—N. Granat.
tenellum, *Naud. l. c.* 159.—Guatemala.
triplinerve, Naud. l. c. iv. (1845) 51 = Deppeanum.
umbellatum, Naud. l. c. 49 = Humboldtianum.

MONOCHASMA, Maxim. ex Franch. & Sav. Enum. Pl. Jap. ii. 458 (1876). *SCROPHULARINEAE.*
Savateri. *Franch. ex Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) I. 39.—China.
Sheareri, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 458.—Japon.

MONOCHILA, Spach, Hist. Vég. Phan. ix. 227 (1840) = **Goodenia**, Sm.

MONOCHILON, Dulac, Fl. Hautes-Pyr. 405 (1867) = **Teucrium**, Tourn. (Labiat.).
bipinnatifidus, Dulac, l. c. = T. Botrys.
cordifolius, Dulac, l. c. 406 = T. Scorodonia.
palustris, Dulac, l. c. 405 = T. Scordium.
reptans, Dulac, l. c. 406 = T. pyrenaicum.
rubellus, Dulac, l. c. = T. Chamaedrys.
tomentosus, Dulac, l. c. 405 = T. Polium.

MONOCHILUS, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 34 (1835). *VERBENACEAE*, Benth. & Hook. f. ii. 1147.
gloxinifolius, *Fisch. & Mey. l. c.*—Bras.

MONOCHILUS, Wall. ex Lindl. Gen. et Sp. Orch. 486 (1840) = **Zeuxine**, Lindl. (Orchid.).
affinis, Lindl. l. c. 487.—Burma.
affinis, Wight, Ic. t. 1728 = Z. longilabris.
Boryi, Reichb. f. in Linnaea, xli. (1877) 60.—Ins. Borbon.
clandestinus, Miq. Fl. Ind. Bat. iii. 724 = Z. clandestina.
flabellatus, Wight, Ic. t. 1727 = Cheirostylis flabellata.
flavus, Wall. ex Lindl. Gen. et Sp. Orch. 487 = Z. flava.
galeatus, Lindl. in Journ. Linn. Soc. i. (1857) 187 = Z. goodyeroides.
goodyeroides, Lindl. l. c. = Z. goodyeroides.
gracilis, Lindl. ex Miq. Fl. Ind. Bat. iii. 723 = Z. gracilis.
gymnochiloides, Ridl. in Journ. Linn. Soc. xxi. (1885) 499 = Cheirostylis gymnochiloides.
lepidus, Reichb. f. Otia Bot. Hamb. 110.—Afr. trop.
longilabris, Lindl. Gen. et Sp. Orch. 487 = Z. longilabris.
nervosus, Wall. ex Lindl. l. c. = Z. nervosa.
parviflorus, Miq. Fl. Ind. Bat. iii. 724.—Java.
plantagineus, Reichb. f. Otia Bot. Hamb. 52.—Ins. Samoa.
regius, Lindl. Gen. et Sp. Orch. 487 = Z. regia.
stenophyllus, Reichb. f. Otia Bot. Hamb. 52.—Ins. Samoa.
tetrapterus, Reichb. f. l. c. 110.—Afr. trop.
Vieillardi, Reichb. f. in Linnaea, xli. (1877) 60.—N. Caled.
Zollingeri, Reichb. f. in Bonplandia, v. (1857) 35.—Java.

MONOCHLAENA, Cass. in Dict. Sc. Nat. l. 496 (1827) = **Eriocephalus**, Linn. (Compos.).
racemosa, Cass. l. c. = E. umbellulatus.

MONOCHORIA, Presl, Rel. Haenk. i. 127 (1827). *PONTEDERIACEAE*, Benth. & Hook. f. iii. 839.
GOMPHIMA, Rafin. Fl. Tellur. ii. 10 (1836).
LIMNOSTACHYS, F. Muell. Fragm. i. 24 (1858).
cyanea, *F. Muell. Fragm.* viii. 44.—Austral.
dilatata, Kunth, Enum. Pl. iv. 134 = hastaefolia.
dubia, Miq. Fl. Ind. Bat. iii. 549 = vaginalis.
hastaefolia, *Presl, Rel. Haenk.* i. 128.—As. trop.
hastata, Solms, in DC. Monog. Phan. iv. 523, sphalm. = hastaefolia.
Junghuhniana, Hassk. in Flora, xxxv. (1852) 115 = vaginalis.
Korsakowii, Regel & Maack, in Mém. Acad. Pétersb. Sér. VII. iv. (1861) n. IV. 155 = vaginalis.
linearis, Miq. Fl. Ind. Bat. iii. 549 = vaginalis.
Loureirii, Kunth, Enum. Pl. iv. 135 = vaginalis.
ovata, *Kunth, l. c.* 665.—China.
pauciflora, Kunth, l. c. 135 = vaginalis.
plantaginea, Kunth, l. c. = vaginalis.
sagittata, Kunth, l. c. 134 = hastaefolia.
vaginalis, *Presl, Rel. Haenk.* i. 128, in obs.—As. et Afr. trop.

MONOCOCCUS, F. Muell. Fragm. i. 46 (1858). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 83.
echinophorus, *F. Muell. l. c.* 47.—Austral.

MONOCODON, Salisb. Gen. Pl. Fragm. 56 (1866) = **Fritillaria**, Tourn. (Liliac.).

MONOCOSMIA, Fenzl, Nov. Stirp. Dec. 84 (1839). *PORTULACEAE*, Benth. & Hook. f. i. 158.
corrigioloides, *Fenzl, l. c.* 93.—Chili.
verrucosa, *Phil. in Linnaea*, xxxiii. (1864-65) 79.—Chili.

MONOCYCLIS, Wall. ex Voigt, Hort. Suburb. Calc. 135 (1845) = **Walsura**, Roxb. (Meliac.).
robusta, Wall. ex Voigt, l. c. = W. robusta.

MONOCYSTIS, Lindl. Introd. Nat. Syst. ed. II. 445 (1836) = **Alpinia**, Linn. (Scitam.).
abnormis, Lindl. l. c. (= *Hellenia abnormis*).—China.

MONODORA, Dun. Monog. Anon. 79 (1817). *ANONACEAE*, Benth. & Hook. f. i. 26.
angolensis, *Welw. Apont.* 587.—Afr. trop.
brevipes, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 475.—Afr. trop.
Grandidieri, *Baill. Adansonia*, viii. (1867-68) 301.—Afr. trop.
grandiflora, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 474.—Afr. trop.
madagascariensis, *Baill. Adansonia*, viii. (1867-68) 299.—Madag.
microcarpa, Dun. Monog. Anon. 80 = Diospyros Cargillia.
Myristica, Blanco, Fl. Filip. ed. II. 300 = Monocarpia Blancoi.
Myristica, *Dun. Monog. Anon.* 80.—Jamaica.
stenopetala, *Oliver, Fl. Trop. Afr.* i. 39.—Afr. trop.
tenuifolia, *Benth. in Journ. Linn. Soc.* v. (1861) 72.—Afr. trop.

MONODYNAMIS, J. F. Gmel. Syst. 10 (1791) = **Usteria**, Willd. (Logan.).
guineensis, Poir. Encyc. Suppl. iii. 732 = U. guineensis.
Iserti, J. F. Gmel. Syst. 10 = U. guineensis.

MONODYNAMUS, Pohl, Pl. Bras. Ic. ii. 67. t. 144 (1831) = **Anacardium**, Rottl.
humilis, Pohl, l. c. 68. t. 144 = A. nanum.

MONOËSTES, Salisb. Gen. Pl. Fragm. 21 (1866) = **Lachenalia**, Jacq. (Liliac.).
unifolia, Salisb. l. c. = L. unifolia.

MONOGRAPHIDIUM, Presl, Epim. Bot. 202 (1849) = **Cliffortia**, Linn. (Rosac.).
obcordatum, Presl, l. c. = C. obcordata.

MONOGYNELLA, Des Moul. Etudes Org. Cusc. 65 (1853) = **Cuscuta**, Linn. (Convolv.).
Blancheana, Des Moul. ex Engelm. in Trans. Acad. St. Louis, i. (1859) 515 = C. monogyna.
Vahliana, Des Moul. Etudes Org. Cusc. 65 = C. monogyna.

MONOLENA, Triana, in Benth. & Hook. f. Gen. i. 756 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
cordifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 80.—N. Granat.
coriacea, *Triana, l. c.*—N. Granat.
primulaeflora, *Hook. f. Bot. Mag.* t. 5818.—Peruv.; N. Granat.
Sprucei, Triana, in Trans. Linn. Soc. xxviii. (1871) 80 = primulaeflora.

MONOLEPIS, Schrad. Ind. Sem. Hort. Gotting. (1830) 4. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 50.
asiatica, *Fisch. & Mey. in Bull. Phys. Math. Acad. Pétersb.* ii. (1844) 131.—Sibir.
chenopodioides, Moq. in DC. Prod. xiii. II. 85 = trifida.
pusilla, *Torr. ex S. Wats. in Bot. King's Exp.* 289.—Am. bor. occ.
spathulata, *A. Gray, in Proc. Am. Acad.* vii. (1868) 389.—Calif.
trifida, *Schrad. Ind. Sem. Hort. Gotting.* (1830) 4.—Sibir.

MONOLIX, Rafin. Cat. 14 (1824) = **Callirhoë**, Nutt. (Malvac.).
digitata, Rafin. l. c. = C. digitata.

MONOLOPHUS, Wall. Pl. As. Rar. i. 24. t. 24 (1830) = **Kaempferia**, Linn. (Scitam.).
coenobialis, Hance, in Journ. Bot. viii. (1870) 75.—China.
elegans, Wall. Pl. As. Rar. i. 24 = K. elegans.
linearis, Wall. in Roxb. Fl. Ind. ed. Carey, i. 20 = K. linearis, secunda.
philippianus, A. Dietr. Allg. Gartenz. xvii. (1849) 266.—'Ind. or.'
scaposus, Dalz. in Hook. Kew Journ. ii. (1850) 143 = K. scaposa.
secunda, Wall. in Roxb. Fl. Ind. ed. Carey, i. 19 = K. secunda.

MONOLOPIA, DC. Prod. vi. 74 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 400.
SPIRIDANTHES, Fenzl, in Endl. Gen. Suppl. ii. 105 (1842).
bahiaefolia, *Benth. Pl. Hartw.* 317.—Calif.
californica, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 80 = Lasthenia glabrata.
glabrata, Fisch. & Mey. l. c. = Lasthenia glabrata.
gracilens, *A. Gray, in Proc. Am. Acad.* xix. (1883) 20.—Calif.
Heermannii, Durand, in Journ. Acad. Sc. Nat. Philad. Ser. II. iii. (1855) 93 = bahiaefolia.
lanceolata, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 175.—Calif.
major, *DC. Prod.* vi. 74.—Calif.
minor, *DC. l. c.*—Calif.

MONOMERIA, Lindl. Gen. et Sp. Orch. 61 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 506.
barbata, *Lindl. in Wall. Cat.* n. 1978; *ex ej. Gen. et Sp. Orch.* 61.—Burma.
Crabro, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 143 = barbata.

MONOMESIA, Rafin. Fl. Tellur. iv. 87 (1836) = **Coldenia**, Linn. (Borag.).

MONOON, Miq. Ann. Mus. Lugd. Bat. ii. 15 (1865) = **Polyalthia**, Blume (Anonac.).
biglandulosum, Miq. l. c. 19 = P. biglandulosa.
canangioides, Miq. l. c. 18.—Sumatra.
chloraxanthum, Miq. l. c. 17.—N. Guin.
chryseum, Miq. l. c. 15.—Sumatra.
costigerum, Miq. l. c. 18.—Sumatra.
glaucum, Miq. l. c. 19.—Java.
hirtum, Miq. l. c. 16.—N. Guin.
incertum, Miq. l. c. 19.—Java.
lateriflorum, Miq. l. c. = P. lateriflora.
littorale, Miq. l. c.—Java.
longipes, Miq. l. c. 15.—Java.
macropodium, Miq. l. c. 17.—N. Guin.
oligocarpum, Miq. l. c. 18.—Sumatra.
pallidum, Miq. l. c. 19.—Java.
sublanceolatum, Miq. l. c. 18.—Borneo.
submitratum, Miq. l. c. 16.—Borneo.
sumatranum, Miq. l. c. 19 = P. sumatrana.
Teysmannii, Miq. l. c.—Sumatra.

MONOPANAX, Regel, Gartenfl. xviii. (1869) 35. t. 606. = **Oreopanax**, Decne. & Planch. (Araliac.).
Ghiesbreghtii, Regel, l. c. = O. xalapense.

MONOPHALACRUS, Cass. in Dict. Sc. Nat. liii. 235 (1828) = **Tessaria**, Ruiz & Pav. (Compos.).

MONOPHYLLAEA, R. Br. in Benn. Pl. Jav. Rar. 121 (1838). *GESNERACEAE*, Benth. & Hook. f. ii. 1017.
Beccarii, *C. B. Clarke, in DC. Monog. Phan.* v. 183.—Borneo.
glauca, *C. B. Clarke, l. c.* 182.—Borneo.
hirtella, *Miq. Fl. Ind. Bat. Suppl.* 564.—Sumatra.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 115, 121.—Java.
Lowei, *C. B. Clarke, in DC. Monog. Phan.* v. 183.—Borneo.
pygmaea, *C. B. Clarke, l. c.*—Java.

MONOPHYLLON, Delarb. Fl. Auv. ed. II. 615 (1800) = **Maianthemum**, Web. (Liliac.).
Lobelii, Delarb. l. c. = Maianth. Convallaria.

MONOPLECTRA, Rafin. Fl. Ludov. 106 (1817) = **Sesbania**, Pers. (Legumin.).

MONOPLOCA, Bunge, in Lehm. Pl. Preiss. i. 259 (1845) **Lepidium**, Linn. (Crucif.).
leptopetala, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 35 = leptopetalum.
linifolia, Bunge, in Lehm. Pl. Preiss. i. 260 = L. rotundum.
phlebopetala, F. Muell. in Linnaea, xxv. (1852) 369 = L. phlebopetalum.
rotunda, Bunge, in Lehm. Pl. Preiss. i. 260 = L. rotundum.

MONOPOGON, J. & C. Presl, Rel. Haenk. i. 324. t. 44 (1830) = **Tristachya**, Nees (Gramin.).
avenaceus, J. & C. Presl, l. c. = T. leiostachya.

MONOPORANDRA, Thw. in Hook. Kew Journ. vi. (1854) 69. t. 2. *DIPTEROCARPEAE*, Benth. & Hook. f. i. 194.
cordifolia, *Thw. l. c.* 70.—Zeylan.
elegans, *Thw. l. c.* 69. t. 2. f. B.—Zeylan.
lancifolia, *Thw. Enum. Pl. Zeyl.* 39.—Zeylan.

MONOPORINA, J. Presl, Rostl. ii. 277 (1825) = **Marila**, Sw. (Cappar.).
oenotheroides, J. Presl, l. c. = M. racemosa.

MONOPORUS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91. t. 7 = **Ardisia**, Sw. (Myrsin.)
paludosus, A. DC. l. c.—Madag.

MONOPSIS, Salisb. in Trans. Hort. Soc. ii. (1817) 37 = **Lobelia**, Linn. (Campan.).
aspera, Urb. in Eichl. Bot. Jahrb. i. (1881) 274 = L. aspera.
campanulata, Sond. in Harv. & Sond. Fl. Cap. iii. 534 = L. campanulata.
conspicua, Salisb. in Trans. Hort. Soc. ii. (1817) 37 = L. debilis.
corymbosa, Presl, Prod. Monog. Lobel. 11 = L. corymbosa.
debilis, Presl, l. c. = L. debilis.
dioica, Presl, l. c. = L. dioica.
gracilis, A. DC. in DC. Prod. vii. 352 = L. campanulata.
inconspicua, Salisb. in Trans. Hort. Soc. ii. (1817) 40 = L. anceps.
litigiosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. vii. 53.—Afr. austr.
lutea, Urb. in Eichl. Bot. Jahrb. i. (1881) 276 = L. lutea.
scabra, Urb. l. c. 274 = L. scabra.
Schimperiana, Urb. l. c. 275 = L. stellarioides.
Speculum, A. DC. in DC. Prod. vii. 351 = L. debilis.
stellarioides, Urb. in Eichl. Bot. Jahrb. i. (1881) 275 = L. stellarioides.
tenella, Urb. l. c. 273 = L. tenella.
variifolia, Urb. l. c. 277 = L. variifolia.

MONOPTERA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 253 (1836-50) = **Chrysanthemum**, Linn. (Compos.).
filifolia, Sch. Bip. l. c. = C. dissectum?

MONOPTERIS, Klotzsch, in Schomb. Faun. & Fl. Gui. 1181, nomen. *SAPINDACEAE*.
guianensis, Klotzsch, l. c.—Guiana.

MONOPTERYX, Spruce, ex Benth. in Mart. Fl. Bras. xv. 1. 307. t. 122 (1862). *LEGUMINOSAE*, Benth. & Hook. f. i. 552.
angustifolia, *Spruce, ex Benth. l. c.* 307.—Bras.
Uauca, *Spruce, ex Benth. l. c.* 308.—Bras.

MONOPTILON, Torr. & Gray, in Proc. Bost. Soc. Nat. Hist. i. (1844) 210; et ex A. Gray, in Bost. Journ. Nat. Hist. v. (1847) 106. t. 13. *COMPOSITAE*, Benth. & Hook. f. ii. 267.
bellidiforme, *Torr. & Gray, ex A. Gray, in Bost. Journ. Nat. Hist.* v. (1847) 106.—Am. bor. occ.

MONOPYLE, Moritz, ex Benth. & Hook. f. Gen. ii. 997 (1876). *GESNERACEAE*, Benth. & Hook. f. l. c.
leucantha, *Moritz, ex Benth. in Hook. Ic. Pl.* xii. 87.—Venezuela.
macrocarpa, *Benth. l. c.* 85. t. 1198.—Costa Rica.
macrophylla, *Benth. l. c.* 86.—N. Granat.
paniculata, *Benth. l. c.*—Ecuador.
racemosa, *Benth. l. c.* 87.—N. Granat.
subsessilis, *Benth. l. c.* 86.—Peruv.

MONORCHIS, Ehrh. Beitr. iv. 147 (1789) = **Herminium**, Linn. (Orchid.).

MONOSIS, DC. Prod. v. 77 (1836) = **Vernonia**, Schreb. (Compos.).
brasiliensis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 230.—Bras.
elaeagnoides, Kunth, Ind. Sem. Hort. Berol. (1848) 14 = V. cinerea?
foliosa, Benth. Pl. Hartw. 19 = V. foliosa, *Sch. Bip.*

MONOSIS :—
insularum, A. Gray, in Proc. Am. Acad. v. (1861) 115.—Ins. Fiji.
salicifolia, DC. Prod. v. 77 = Vernonia salicifolia.
tarchonanthifolius, DC. l. c. = V. tarchonanthifolia.
tomentosa, DC. l. c. = V. Monosis.
Wightiana, DC. in Wight, Contrib. 1 = V. Monosis.

MONOSOMA, Griff. Notul. iv. 502 (1854) = **Carapa**, Aubl. (Meliac.).
littorata, Griff. l. c. = C. moluccensis.

MONOSPORA, Hochst. in Flora, xxiv. (1841) 660 = **Trimeria**, Harv. (Bixin.).
grandifolia, Hochst. l. c. 661.—Afr. austr.
rotundifolia, Hochst. l. c. = T. alnifolia.

MONOSTEMMA, Turcz. in Bull. Soc. Nat. Mosc. (1848) i. 255 = **Sarcostemma**, R. Br. (Asclep.).
aphyllum, Turcz. l. c. = S. aphyllum.
tetrapterum, Turcz. l. c. = S. tetrapterum.

MONOSTERIA, Rafin. Fl. Tellur. iii. 49 (1836) = **Gentiana**, Tourn.
dichotoma, Rafin. l. c. = G. tenella.

MONOSTICHE, Koern. in Regel, Gartenfl. (1858) 76, 88 = **Calathea**, G. F. W. Mey. (Scitam.).
colorata, Koern. l. c. 88 = C. colorata.
Danielli, Horan. Prod. Scitam. 13.—Afr. trop.
Hookeri, Koern. in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 360. t. 6. f. 4 = C. colorata.

MONOSTYLIS, Tul. in Arch. Mus. Par. vi. (1852) (Podost. Monog. 201) = **Apinagia**, Tul. (Podost.).
capillacea, Tul. l. c. 202.—Bras.

MONOTASSA, Salisb. Gen. Pl. Fragm. 36 (1866) = **Urginea**, Steinh. (Liliac.).
secunda, Salisb. l. c. 37 = U. secunda.

MONOTAXIS, Brongn. in Duperr. Voy. Coq. Bot. 223. t. 49 (1829). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 264.
HIPPOCREPANDRA, Muell. Arg. in Linnaea, xxxiv. (1865–66) 61.
REISSIPA, Steud. ex Klotzsch, in Lehm. Pl. Preiss. ii. 230 (1847).
bracteata, Nees, ex Klotzsch, in Lehm. Pl. Preiss. ii. 230 = grandiflora.
cuneifolia, Klotzsch, l. c. 229 = occidentalis.
ericoides, Klotzsch, l. c. i. 177 = grandiflora.
gracilis, *Baill. Adansonia*, vi. (1865–66) 293.—Austral.
grandiflora, *Endl. in Enum. Pl. Hueg.* 19.—Austral.
linifolia, *Brongn. in Duperr. Voy. Coq. Bot.* 223. t. 49.—Austral.
lurida, *Benth. Fl. Austral.* vi. 80.—Austral.
luteiflora, *F. Muell. Fragm.* x. 51.—Austral.
macrophylla, *Benth. Fl. Austral.* vi. 79.—Austral.
megacarpa, *F. Muell. Fragm.* iv. 143.—Austral.
Neesiana, Baill. Adansonia, vi. (1865–66) 293 = grandiflora.
occidentalis, *Endl. in Enum. Pl. Hueg.* 19.—Austral.
Oldfieldii, Baill. Adansonia, vi. (1865–66) 293 = lurida.
porantheroides, F. Muell. ex Benth. Fl. Austral. vi. 80 = occidentalis.
simplex, Brongn. ex Steud. Nom. ed. II. ii. 158 = linifolia ?
tridentata, Endl. Atakt. 8. t. 8 = linifolia.

MONOTELES, Rafin. Sylva Tellur. 122 (1838) = **Bauhinia**, Linn. (Legumin.).
paradoxa, Rafin. l. c. (Quid ?).—Hab. ?

MONOTES, A. DC. in DC. Prod. xvi. II. 623 (1868). *DIPTEROCARPEAE*.
africana, *A. DC. l. c.* 624.—Afr. trop.

MONOTHECA, A. DC. in DC. Prod. viii. 152 (1844) = **Reptonia**, A. DC. (Myrsin.).
mascatensis, A. DC. l. c. = R. buxifolia.

MONOTHECIUM, Hochst. in Flora, xxiv. (1841) 374. *ACANTHACEAE*, Benth. & Hook. f. ii. 1104.
ANTHOCOMETES, Nees, in DC. Prod. xi. 311 (1847).
aristatum, *T. Anders. in Thw. Enum. Pl. Zeyl.* 234.—Ind. or.
glandulosum, *Hochst. in Flora*, xxiv. (1841) 374.—Abyssin.

MONOTHRIX, Torr. in Stansb. Utah Exped. 389. t. 7 (1852) = **Laphamia**, A. Gray (Compos.).
Stansburiana, Torr. l. c. 390 = L. Stansburii.

MONOTHYLACEUM, G. Don, Gen. Syst. iv. 116 (1837) = **Hoodia**, Sweet (Asclep.).
Gordoni, G. Don, l. c. = H. Gordoni.

MONOTHYLACIUM, Benth. & Hook. f. Gen. ii. 783 (1876), sphalm. = praec.

MONOTOCA, R. Br. Prod. 546 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 615.
albens, R. Br. l. c. 547 = elliptica.
elliptica, *R. Br. l. c.* 546.—Austral.
empetrifolia, *R. Br. l. c.* 547.—Austral.
ledifolia, *A. Cunn. ex DC. Prod.* vii. 756.—Austral.
lineata, *R. Br. Prod.* 547.—Austral.
oligarrhenoides, *F. Muell. Fragm.* ix. 47.—Austral.
patens, A. Cunn. ex DC. Prod. vii. 756 = scoparia.
propinqua, A. Cunn. l. c. = scoparia.
Pumilio, Spreng. Syst. i. 654 = Needhamia Pumilio.
scoparia, *R. Br. Prod.* 547.—Austral.
tamariscina, *F. Muell. Fragm.* vi. 79.—Austral.

MONOTREMA, Koern. in Linnaea, xxxvii. (1871–72) 475 = **Stegolepis**, Klotzsch (Rapat.).
aemulans, Koern. l. c. 477 = S. aemulans.
flavum, Koern. l. c. 478.—Venezuela.

MONOTRIS, Lindl. Bot. Reg. sub t. 1701 (1834) **Holothrix**, Rich. (Orchid.).
secunda, Lindl. l. c. = H. Monotris.

MONOTROPA, Linn. Syst. ed. I (1735). *MONOTROPEAE*, Benth. & Hook. f. ii. 607.
HYPOPITHYS, Adans. Fam. ii. 443 (1763).
MONOTROPION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130.
abietina, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 342.—Europ.
coccinea, Zucc. in Flora, xv. (1832) II. Beibl. 100 = uniflora.
epirrhizium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = Hypopitys multiflora.
fimbriata, *A. Gray, in Proc. Am. Acad.* viii. (1873) 629.—Calif.
flagrans, Gilib. Fl. Lituan. ii. 191 = Hypopitys multiflora.
glabra, Bernh. ex Reichb. Fl. Germ. Excurs. 411 = Hypopitys multiflora.
hirsuta, Hornem. ex Willk. & Lange, Prod. Fl. Hisp. ii. 339 = Hypopitys multiflora.
humilis, D. Don, Prod. Fl. Nep. 151 = uniflora.
hypophega, Wallr. Sched. Crit. 191 = Hypopitys multiflora.
hypopithys, Linn. Sp. Pl. 387 = Hypopitys multiflora.
Hypopitys, Walt. Fl. Carol. 136 = praec.
hypoxya, Spreng. Syst. ii. 317 = Hypopitys multiflora.
japonica, *Franch. & Sav. Enum. Pl. Jap.* i. 297.—Japon.
lanuginosa, Michx. Fl. Bor. Am. ii. 266 = Hypopitys multiflora.
Morisoni, Pers. Syn. i. 469 = uniflora.
Morisoniana, Michx. Fl. Bor. Am. i. 266 = praec.
nepalensis, Wall. Cat. n. 7252 = Hypopitys multiflora.
squamiformis, Dulac, Fl. Hautes-Pyr. 421 = Hypopitys multiflora.
uniflora, *Linn. Sp. Pl.* 387.—Reg. Himal.; Japon.: Am. bor.

MONOTROPION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = praec.

MONOTROPSIS, Schwein. ex Ell. Sketch, i. 478 (1817) = **Schweinitzia**, Ell. (Monotrop.).
odorata, Schwein. ex Ell. l. c. = S. odorata.

MONOXORA, Wight, Illustr. ii. 12. t. 97 (1841) = **Rhodamnia**, Jack (Myrtac.).
latifolia, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 218 = R. latifolia.
rubescens, Benth. l. c. 219 = R. trinervia.
spectabilis, Wight, Illustr. ii. 12 = R. trinervia.

MONROA, Torr. in Pacif. Rail. Rep. iv. 158 (1856) = **Munroa**, 'Torr.' (Gramin.).

MONSONIA, Linn. Mant. i. 14 (1767). *GERANIACEAE*, Benth. & Hook. f. i. 271.
OLOPETALUM, Klotzsch, in Linnaea, x. (1835) 428.
angustifolia, E. Mey. in Drège, Zwei Pfl. Docum. 203 = biflora.
apiculata, E. Mey. l. c. (Quid ?).—Afr. austr.
arabica, Steud. Nom. ed. II. ii. 158 = ovata.
asiatica, Vicary, in Journ. As. Soc. Beng. xvi. (1847) II. 1161 = senegalensis.
attenuata, *Harv. & Sond. Fl. Cap.* i. 155.—Afr. austr.
biflora, *DC. Prod.* i. 138.—Afr. austr.
Burkeana, *Planch. ex Harv. & Sond. Fl. Cap.* i. 155.—Afr. austr.
Burmanni, DC. Prod. i. 638 = Sarcocaulon Burmanni.
chumbalensis, Wight, Ic. iii. iv. 5. t. 1047 = senegalensis.
Dregeana, *Presl, Bot. Bemerk.* 26.—Afr. austr.
emarginata, L'Hérit. Geran. t. 41 = ovata.
Filia, Hort. ex Willd. Enum. Hort. Berol. 717 = pilosa.
Filia, Linn. f. Suppl. 341 = lobata.
grandiflora, Burm. f. Fl. Cap. Prod. 23 = spinosa.
heliotropoides, *Boiss. Fl. Orient.* i. 897.—Egypt.; As. occ.
Heritieri, Steud. Nom. ed. II. ii. 158 = *L'Heritieri*.
hispida, Boiss. Diagn. Ser. I. viii. 120 = heliotropoides.
incisa, Dum.-Cours. Bot. Cult. ed. II. v. 49 = lobata.
Lawiana, Stocks, ex Wight, in Calc. Journ. Nat. Hist. vii. (1847) 19 = senegalensis.
L'Heritieri, DC. Prod. i. 638 = Sarcocaulon L'Heritieri.
lobata, *Mont. in Gothob. Wet. Samk. Handb.* i. *Wet. Afd.* II. (1780) 1. t. 1.—Afr. austr.
macilenta, E. Mey. in Drège, Zwei Pfl. Docum. 203 = Sarcocaulon Patersonii.
mallica, Edgew. in Journ. Linn. Soc. vi. (1862) 200 = heliotropioides.
marginata, Klotzsch, in Linnaea, x. (1835–36) 429, sphalm. = *emarginata*.
mossamedensis, *Welw. in Oliver, Fl. Trop. Afr.* i. 290.—Afr. trop.
multifida, E. Mey. in Drège, Zwei Pfl. Docum. 203 (Quid ?).—Afr. austr.
nivea, *Webb, Fragm. Fl. Aethiop.* 59.—Afr. bor.
obcordata, E. Mey. in Drège, Zwei Pfl. Docum. 203 = Sarcochilon Patersonii.
ovata, *Cav. Diss.* iv. 193. t. 113. f. 1.—Afr. bor. et austr.
Patersonii, DC. Prod. i. 639 = Sarcocaulon Patersonii.
pilosa, *Willd. Enum. Hort. Berol.* 717.—Afr. austr.
praemorsa, E. Mey. in Drège, Zwei Pfl. Docum. 203 = ovata.
senegalensis, *Guill. & Perr. Fl. Seneg. Tent.* i. 131.—Afr. trop. et austr.; As. trop. occ.
speciosa, Drège, ex Presl, Bot. Bemerk. 26 = Dregeana.
speciosa, *Linn. Mant.* i. 105.—Afr. austr.
speciosa, Sweet, Geran. t. 77 = pilosa.
spinosa, L'Hérit. Geran. t. 42 = Sarochilon L'Heritieri.
tenuifolia, Hort. ex Steud. Nom. ed. II. ii. 158 = pilosa.
tenuifolia, *Willd. Sp. Pl.* iii. 717.—Afr. austr.
umbellata, *Harv. in Harv. & Sond. Fl. Cap.* i. 255.—Afr. austr.

MONSSONIA, Linn. Mant. i. 105 (1767) = praec.

MONSTERA, Adans. Fam. ii. 470 (1763). *AROIDEAE*, Benth. & Hook. f. iii. 991.
ALLOSCHEMONE, Schott, Prod. Aroid. 358 (1860).
FORNELIA, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179, sphalm.
SERANGIUM, W. Wood, ex Salisb. Gen. Pl. Fragm. 5 (1866).
TORNELIA, Gutierez. ex Schlecht. in Linnaea, xxvi. (1853–55) 382.
acuminata, *C. Koch, Ind. Sem. Hort. Berol.* (1855) *App.* 4.—Am. trop.

MONSTERA:—
Adansonii, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52; Melet. i. 21 = pertusa.
amomifolia, Poepp. & Endl. Nov. Gen. et Sp. iii. 88 = Stenospermatium amomifolium.
Blanchetii, Schott, Prod. Aroid. 367 = pertusa.
Borsigiana, C. Koch, Wochenschr. v. (1862) 275 = deliciosa?
cannaefolia, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52; Melet. i. 21 = Spathiphyllum cannaeforme.
caudata, Schott, ll. cc. 1028; 53 = Epipremnum mirabile.
crassifolia, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 40 = pertusa.
cuspidata, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 540 = Philodendron propinquum.
decursiva, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52 = Rhaphidophora decursiva.
deliciosa, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 19.—Am. trop.
dilacerata, *C. Koch, Ind. Sem. Hort. Berol.* (1855) *App.* 5.—Costa Rica.
dimidiata, Hort. ex Schott, Prod. Aroid. 372 = acuminata.
egregia, *Schott, in Journ. Bot.* ii. (1864) 53.—Hab.?
expilata, Schott, Prod. Aroid. 367 = obliqua.
fenestrata, Schott, in Oestr. Bot. Wochenbl. v. (1855) 290 = pertusa.
Friedrichsthalii, *Schott, l. c.* iv. (1854) 65.—Guatemala.
Gaudichaudii, Schott, Prod. Aroid. 368 = pertusa.
gigantea, C. Koch, ex Ender, Ind. Aroid. 74 = Epipremnum giganteum.
gigantea, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52 = Epipremnum giganteum.
glauca, C. Koch, ex Ender, Ind. Aroid. 54 = Rhaphidophora glauca.
gracilis, *Engl. in DC. Monog. Phan.* ii. 258.—N. Granat.
Holtoniana, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 40 = pertusa.
Imrayana, Schott, l. c. = pertusa.
Jacquini, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 66 = pertusa.
Karwinskyi, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 99.—Mexic.
Klotzschiana, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 419 = pertusa.
lanceaefolia, Schott, Prod. Aroid. 368 = pertusa.
Lechleriana, Schott, l. c. 366 = pertusa.
Lennea, C. Koch, in Bot. Zeit. x. (1852) 277; Ind. Sem. Hort. Berol. (1853) 15 = deliciosa.
lingulata, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52 = Rhaphidophora lingulata?
macrophylla, Schott, Prod. Aroid. 362 = pertusa.
Maximiliani, *Engl. in Mart. Fl. Bras.* iii. II. 113.—Bras.
microstachys, Schott, in Oestr. Bot. Wochenbl. v. (1855) 274 = obliqua.
Milleriana, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 386 = pertusa.
modesta, Schott, Gen. Aroid. 75 = pertusa.
multijuga, C. Koch, ex Ender, Ind. Aroid. 73 = Rhaphidophora decursiva.
obliqua, *Miq. in Linnaea*, xviii. (1844) 79.—Bras.; Guiana.
oblongifolia, Schott, Prod. Aroid. 369 = pertusa.
occidentalis, *C. Koch, ex Ender, Ind. Aroid.* 4.—Bras.
officinalis, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52 = Scindapsus officinalis.
ovata, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 387 = pertusa.
Parkeriana, Schott, l. c. ix. (1859) 41 = dilacerata.
Peepla, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 52 = Rhaphidophora Peepla.
pertusa, *Schott, in Wien. Zeitschr.* iv. (1830) 1028; *ex Endl. in Linnaea*, vi. (1831) Litt. 52.—Am. trop.
pinnata, Schott, ll. cc. 1028; 53 = Epipremnum mirabile.
pinnatifida, Schott, in Wien. Zeitschr. iv. (1830) 1028; ex Endl. in Linnaea, vi. (1831) Litt. 53 = pertusa.
Poeppigii, Schott, Prod. Aroid. 365 = pertusa.
protensa, Schott, ex Engl. in DC. Monog. Phan. ii. 263 = pertusa.
punctulata, *Schott, ex Engl. l. c.* 259.—Mexic.
Seemanni, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 40 = pertusa?
Spruceana, *Engl. in Mart. Fl. Bras.* iii. II. 115.—Bras.
subpinnata, *Engl. in DC. Monog. Phan.* ii. 267.—Peruv.
surinamensis, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 387 = Rhaphidophora pertusa.
tenuis, C. Koch, Ind. Sem. Hort. Berol. (1855) 4 = acuminata.
trijuga, C. Koch, ex Ender, Ind. Aroid. 73 = Rhaphidophora decursiva.
Velloziana, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 66 = pertusa.

MONTABEA, Roem. & Schult. Syst. iv 527 (1819), sphalm. = **Moutabea**, Aubl. (Polygal.).

MONTAGNAEA, DC. Prod. v. 564 (1836) = **Montañoa**, Cerv. (Compos.).
arborescens, DC. l. c. 565 = Montañoa arborescens.
atriplicifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 304 = Montañoa arborescens.
clematidea, Walp. in Linnaea, xiv. (1840) 308 = Montañoa clematidea.
ensifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 304 = Gymnolomia ensifolia.
excelsa, Ernst, in Vargasia, i. (1870) 186.—Venezuela.
floribunda, DC. Prod. v. 564 = Montañoa floribunda.
frutescens, DC. l. c. 565 = Montañoa frutescens.
grandiflora, DC. l. c. = Montañoa grandiflora.
heracleifolia, Brongn. in Rev. Hortic. (1863) 370 = Montañoa bipinnatifolia.
heterophylla, Walp. Rep. ii. 612 = Montañoa tomentosa.
hibiscifolia, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 89 = Montañoa hibiscifolia.
Karvinskii, DC. Prod. v. 565 = Montañoa Karwinski.
ovalifolia, DC. l. c. 566 = Montañoa ovalifolia.
purpurea, Brongn. in Rev. Hortic. (1863) 370, in obs. = Montañoa purpurea.
speciosa, DC. Prod. v. 565 = Montañoa speciosa.
tomentosa, DC. l. c. 564 = Montañoa tomentosa.

MONTALBANIA, Neck, Elem. i. 273 (1790) = **Clerodendron**, Linn. (Verben.).

MONTAÑOA, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 11 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 364.
ERIOCARPHA, Cass. in Dict. Sc. Nat. lix. 236 (1829).
ERIOCOMA, H. B. & K. Nov. Gen. et Sp. iv. 267. t. 396 (1820).
MONTAGNAEA, DC. Prod. v. 564 (1836).
PRIESTLEYA, Moç. & Sesse, ex DC. Prod. v. 564 (1836).
UHDEA, Kunth, Ind. Sem. Hort. Berol. (1847).—Cf. Ann. Sc. Nat. Sér. III. ix. (1848) 316.
arborescens, *C. Koch, Wochenschr.* vii. (1864) 406.—Mexic.
Aschenbornii, *Sch. Bip. ex C. Koch, l. c.* 407.—Mexic.
atriplicifolia, C. Koch, l. c. = arborescens.
bipinnatifida, *C. Koch, l. c.*—Mexic.
clematidea, *Hemsl. Biol. Centr. Am. Bot.* ii. 165.—Mexic.
crenata, *Sch. Bip. ex C. Koch, Wochenschr.* vii. (1864) 407.—Mexic.
elegans, *C. Koch, l. c.* 408.—Mexic.
floribunda, *C. Koch, l. c.* 406.—Mexic.
frutescens, *Hemsl. Biol. Centr. Am. Bot.* ii. 165.—Mexic.
gracilis, *Sch. Bip. ex C. Koch, Wochenschr.* vii. (1864) 407.—Mexic.
grandiflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 165.—Mexic.
grandiflora, *Hemsl. l. c.* (sp. diversa).—Mexic.
heracleifolia, Brongn. in Rev. Hortic. Sér. IV. v. (1857) 544 = bipinnatifida.

MONTAÑOA:—
heterophylla, Hemsl. Biol. Centr. Am. Bot. ii. 166, in syn. = tomentosa.
hibiscifolia, *C. Koch, Wochenschr.* vii. (1864) 407.—Guatemala.
Karwinski, *C. Koch, l. c.* —Mexic.
microcephala, *Sch. Bip. ex C. Koch, l. c.* 406.—Mexic.
mollissima, *Brongn. in Rev. Hortic.* Sér. IV. vi. (1857) 543.—Mexic.
Olivae, *Sch. Bip. ex C. Koch, Wochenschr.* vii. (1864) 406.—Mexic.
Orbignyana, *Klatt, Comp. Francav.* 8.—N. Mexic.
ovalifolia, *C. Koch, Wochenschr.* vii. (1864) 407.—N. Granat.
purpurea, *Brongn. in Rev. Hortic.* Sér. IV. vi. (1857) 544, nomen.—Hab.?
pyramidata, *Sch. Bip. ex C. Koch, Wochenschr.* vii. (1864) 408.—Mexic.
quadrangularis, *Sch. Bip. ex C. Koch, l. c.* 407.—Venezuela.
speciosa, *C. Koch, l. c.* 408.—Mexic.
ternifolia, *Sch. Bip. ex C. Koch, l. c.* 406.—Mexic.
Thomasii, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 328.—Mexic.
tomentosa, *Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc.* ii. 11.—Mexic.
triloba, *Sch. Bip. ex C. Koch, Wochenschr.* vii. (1864) 406.—Mexic.
uncinata, *Sch. Bip. ex C. Koch, l. c.*—Mexic.
xanthiifolia, *Sch. Bip. ex C. Koch, l. c.*—Mexic.

MONTBRETIA, DC. in Bull. Soc. Philom. iii. (1803) 151 = **Tritonia**, Ker-Gawl. (Irid.).
abyssinica, Hochst. in Flora, xxiv. (1841) I. Intell. 21 = Lapeyrousia abyssinica.
aequimaculata, Heynh. Nom. ii. 417.—Hab.?
anigozanthaeflora, Heynh. l. c. = T. anigozanthiflora.
capensis, Voigt, Hort. Suburb. Calc. 611 = T. rosea.
concolor, Voigt, l. c. = T. concolor.
crispa, Voigt, l. c. 610 = T. crispa.
crocata, Voigt, l. c. 611 = T. crocata.
deusta, Voigt, l. c. = T. deusta.
dubia, Baker, in Journ. Linn. Soc. xvi. (1877) 169 = T. dubia.
fenestrata, Voigt, Hort. Suburb. Calc. 611 = T. fenestrata.
flava, Heynh. Nom. ii. 417; Klatt, in Linnaea, xxxii. (1863) 753 = T. flava.
fucata, Heynh. l. c. = Antholyza fucata.
gallabatensis, Schweinf. ex Baker, in Journ. Linn. Soc. xvi. (1877) 155 = Lapeyrousia abyssinica.
kamisbergensis, Baker, l. c. 169 = T. kamisbergensis.
lacerata, Baker, l. c. 168 = T. crispa.
lancea, Endl. ex Heynh. Nom. ii. 417 = T. squalida.
laxifolia, Klatt, in Linnaea, xxxii. (1863) 754 = T. laxifolia.
lineata, Voigt, Hort. Suburb. Calc. 610 = T. lineata.
longiflora, Voigt, l. c. = Ixia paniculata.
miniata, Voigt, l. c. 611 = T. miniata.
odorata, Heynh. Nom. ii. 418 = Freesia refracta.
pallida, Baker, in Journ. Linn. Soc. xvi. (1877) 168 = Ixia paniculata.
pauciflora, Baker, Journ. Bot. xiv. (1876) 336 = T. pauciflora.
pectinata, Heynh. Nom. ii. 418 = T. crispa.
Pottsii, Baker, in Gard. Chron. (1877) II. 424 = T. Pottsii.
purpurea, Heynh. Nom. ii. 418 = T. crocata.
refracta, Endl. ex Heynh. l. c. = Freesia refracta.
Rocheana, Heynh. l. c. = T. Rochensis.
rosea, Voigt, Hort. Suburb. Calc. 610; Baker, in Journ. Linn. Soc. xvi. (1877) 169 = T. rosea.
sanguinea, Heynh. Nom. ii. 418 = T. crocata.
securigera, [DC. in] Red. Lil. t. 53 = T. securigera.
squalida, Voigt, Hort. Suburb. Calc. 611 = T. squalida.
striata, Baker, in Journ. Bot. xiv. (1876) 337 = T. Bakeri.
striata, Voigt, Hort. Suburb. Calc. 610 = Gladiolus striatus.
strictifolia, Klatt, in Linnaea, xxxii. (1863) 753 = T. strictifolia.
tenuiflora, Voigt, Hort. Suburb. Calc. 611 = Ixia paniculata.
viridis, Voigt, l. c. 610 = T. viridis.
xanthospila, Heynh. Nom. ii. 418 = Freesia xanthospila.

MONTELIA, A. Gray, Man. Bot. N. U. St. ed. II. 369 (1856) = **Acnida**, Linn. (Amarant.).
tamariscina, A. Gray, l. c. 370 = A. tamariscina.

MONTEVERDIA, A. Rich. Ess. Fl. Cub. 346 (1845) = **Maytenus**, Feuill. (Celastr.).
buxifolia, A. Rich. l. c. 347 = Maytenus buxifolia.

MONTEZUMA, [Moç. & Sesse,] ex DC. Prod. i. 477 (1824). *MALVACEAE*, Benth. & Hook. f. i. 212.
speciosissima, [*Moç. & Sesse,*] *ex DC. l. c.*—Mexic.

MONTIA, Houst. ex Linn. Hort. Cliff. 211 (1737) = **Heliocarpus**, Linn. (Tiliac.).

MONTIA, Mich. ex Linn. Syst. ed. I (1735). *PORTULACEAE*, Benth. & Hook. f. i. 159.
LATERIFISSUM, Dulac, Fl. Hautes-Pyr. 366 (1867).
LEPTRINA, Rafin. in Am. Monthly Mag. (1819) 192.
LEPTRINIA, Schult. Mant. ii. 38 (1824).
alsine-facie, Gilib. Fl. Lituan. i. 122 = fontana.
arvensis, Wallr. Beitr. Fl. Hercyn. 177 ; et in Linnaea, xiv. (1840) 547 = fontana.
decumbens, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = fontana.
erecta, Steud. Nom. ed. I. 538 = fontana.
fontana, *Linn. Sp. Pl.* 87.—Reg. temp. bor. et austr.
gibba, *Griseb. in Goett. Abh.* vi. (1854) 117.—Chili.
Howellii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 191.—Am. bor.
lamprosperma, Cham. in Linnaea, vi. (1831) 565 = fontana.
linearifolia, Urv. in Mém. Soc. Linn. Par. iv. (1826) 619 = fontana.
major, Steud. Nom. ed. II. ii. 159 = fontana.
minor, C. C. Gmel. Fl. Bad. i. 301 = fontana.
pentandra, Willd. ex Cham. in Linnaea, vi. (1831) 565 = fontana.
repens, C. C. Gmel. Fl. Bad. i. 301 = fontana.
rivularis, C. C. Gmel. l. c. = fontana.
tenella, Froel. ex Steud. Nom. ed. II. ii. 159 = fontana.
terrestris, Dum. Fl. Belg. 86 = fontana.
verna, Neck. Delic. Gallo-Belg. i. 78 = fontana.

MONTINIA, Thunb. in Phys. Sållsk. Handl. i. (1776) 108. *ONAGRARIEAE*, Benth. & Hook. f. i. 794.
acris, Linn. f. Suppl. 427 = caryophyllacea.
caryophyllacea, *Thunb. in Phys. Sållsk. Handl.* i. (1776) 109.—Afr. austr.
frutescens, Harv. & Sond. Fl. Cap. ii. 308, in syn. = seq.
fruticosa, Gaertn. Fruct. i. 170 = caryophyllacea.

MONTIRA, Aubl. Pl. Gui. ii. 637. t. 257 (1775) = **Spigelia**, Linn. (Logan.).
guianensis, Aubl. l. c.—Guian.

MONTOLIVAEA, Reichb. f. in Otia Bot. Hamb. 107 (1881) = **Habenaria**, Willd. (Orchid.).
elegans, Reichb. f. l. c. = H. elegans.

MONTRICHARDIA, Crueg. in Bot. Zeit. xii. (1854) 25. *AROIDEAE*, Benth. & Hook. f. iii. 982.
aculeatum, *Crueg. l. c.*; *Schott, Syn. Aroid.* 72.—Bras.
arborea, Schott, Syn. Aroid. 72 = aculeatum.
arborescens, Schott, Araceen Betreff. 4 = aculeatum.
Fendleri, *Schott, Gen. Aroid.* t. 49.—Panama.
linifera, *Schott, Araceen Betreff.* 5.—Bras.
Splitgerberi, *Schott, in Bonplandia*, vii. (1859) 29.—Guian.

MONTROUZERIA, Benth. & Hook. f. Gen. i. 173 (1862), sphalm. = seq.

MONTROUZIERA, Panch. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 292. *GUTTIFERAE*, Benth. & Hook. f. i. 173.
cauliflora, *Planch. & Triana, l. c.* 294.—N. Caled.
Gabriellae, *Baill. Adansonia*, xi. (1873–76) 366.—N. Caled.
robusta, *Vieill. ex Panch. & Sebert, Not. Bois N. Caléd.* 220.—N. Caled.
sphaeroidea, *Panch. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 293.—N. Caled.
spheraeflora, *Panch. in Panch. & Sebert, Not. Bois N. Caléd.* 220.—N. Caled.
verticillata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 293.—N. Caled.

MONTTEA, C. Gay, Fl. Chil. iv. 416. t. 51 (1849). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 946.
OXYCLADUS, Miers, in Proc. Linn. Soc. ii. (1852) 156 ; et in Trans. Linn. Soc. xxi. (1853) 146. t. 18.
chilensis, *J. Gay, Fl. Chil.* iv. 416.—Chili.
Schickendantzii, *Griseb. in Goett. Abh.* xxiv. (1879) 240.—Reg. Argent.

MONUSTES, Rafin. Fl. Tellur. ii. 87 (1836) = **Spiranthes**, Rich. (Orchid.).
australis, Rafin. l. c. = S. australis.

MOONIA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 348 = **Chrysogonum**, Linn. (Compos.).
Arnottiana, Wight, Ic. t. 1105 = C. Arnottianum.
ecliptoides, Benth. Fl. Austral. iii. 540 = C. ecliptoides.
heterophylla, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 349 = C. heterophyllum.
procumbens, Benth. Fl. Austral. iii. 540 = C. procumbens.
trichodesmoides, Benth. l. c. = C. trichodesmoides.

MOORCROFTIA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 431 (Conv. Or. 49. t. 4) = **Lettsomia**, Roxb. (Convolv.).
adpressa, Choisy, l. c. = L. adpressa.
pinangiana, Choisy, l. c. 432 = L. penangiana.

MOORIA, Montr. in Mém. Acad. Lyon, x. (1860) 207 = **Cloezia**, Brongn. (Myrtac.).
artensis, Montr. l. c. = C. floribunda ?

MOPHIGANES, Steud. Nom. ed. II. i. 699, in syn. (1840), sphalm. = **Mogiphanes**, Mart. (Amarant.).

MOQUILEA, Aubl. Pl. Gui. i. 521. t. 208 (1775). *ROSACEAE*, Benth. & Hook. f. i. 606.
DAHURONIA, Scop. Introd. 219 (1777).
Aubletiana, *Blume, Mus. Bot. Lugd. Bat.* ii. 92.—Guian.
bothynophylla, *Mart. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 26.—Bras.
bracteosa, Walp. Rep. ii. 7 = Couepia bracteosa.
canomensis, Mart. Nov. Gen. et Sp. ii. 80. t. 166 = Couepia canomensis.
chrysocalyx, Poepp. & Endl. Nov. Gen. et Sp. iii. 75. t. 286 = Couepia chrysocalyx.
comosa, Walp. Rep. ii. 6 = Couepia comosa.
Couepia, Steud. Nom. ed. II. ii. 159 = Couepia guianensis.
Eliti, Walp. Rep. ii. 6, sphalm. = *M. Uiti*.
floribunda, *Benth. in Hook. Journ. Bot.* ii. (1840) 219.—Bras.
Gardneri, *Hook. f. in Mart. Fl. Bras.* xiv. II. 23.—Bras.
glandulosa, *Walp. Ann.* ii. 463.—Guian.
grandiflora, Mart. & Zucc. in Abh. Akad. Muench. i. (1832) 388 = Couepia grandiflora.
guianensis, *Aubl. Pl. Gui.* i. 521. t. 208.—Guian.
humilis, *Hook. f. in Mart. Fl. Bras.* xiv. II. 26.—Bras.
hypoleuca, *Miq. in Linnaea*, xix. (1847) 438.—Mexic.
Kunthiana, Mart. & Zucc. in Abh. Akad. Muench. i. (1832) 390 = Couepia Kunthiana.
licaniaeflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 308.—Guian.
longistyla, *Hook. f. in Mart. Fl. Bras.* xiv. II. 24.—Bras.
minutiflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 308.—Guian.
multiflora, Walp. Rep. ii. 7 = Couepia multiflora.
organensis, *Miers, in Journ. Linn. Soc.* xvii. (1879) 374.—Bras.
pallida, *Hook. f. in Mart. Fl. Bras.* xiv. II. 25.—Bras.
paraënsis, *Mart. & Zucc. in Abh. Akad. Muench.* i. (1832) 390.—Bras.
Parilla, Steud. Nom. ed. II. ii. 159 = Couepia Parillo.
parviflora, *Blume, Mus. Bot. Lugd. Bat.* ii. 92.—Guian.
pendula, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 22.—Bras.
platypus, *Hemsl. Diagn. Pl. Nov.* 9.—Nicaragua.
Salzmanni, *Hook. f. in Mart. Fl. Bras.* xiv. II. 21.—Bras.
sclerophylla, *Mart. ex Hook. f. l. c.* 23.—Bras.
Sprucei, *Hook. f. l. c.* 22.—Bras.
Steudeliana, *Walp. Ann.* ii. 463.—Guian.
tomentosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Guian.

MOQUILEA :—
turbinata, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 23.—Bras.
Turiuva, *Hook. f. l. c.* 25.—Bras.
Uiti, Mart. & Zucc. in Abh. Akad. Muench. i. (1832) 390 = Couepia Uiti.
utilis, *Hook. f. in Mart. Fl. Bras.* xiv. II. 24.—Bras.

MOQUINIA, DC. Prod. vii. 22 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 490.
SPADONIA, Less. Syn. Comp. 99 (1832), nomen prius.
Bojeri, DC. Prod. vii. 23 (al. gen.).—Afr. trop.
cinerea, DC. l. c. = polymorpha.
cinerea, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 463 = Gardneri.
congesta, *Gardn. l. c.* 459.—Bras.
cratensis, Gardn. l. c. 458.—Bras.
curviflora, *Griseb. in Goett. Abh.* xxiv. (1879) 211.—Reg. Argent.
desertorum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 459 = velutina.
eriosematoides, Walp. Rep. ii. 679 = Inula Cappa.
flavescens, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 458.—Bras.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 348.—Bras.
hypoleuca, DC. Prod. vii. 23 = Gochnatia hypoleuca.
lucida, *Baker, in Mart. Fl. Bras.* vi. III. 347.—Bras.
oligocephala, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 457.—Bras.
paniculata, *DC. Prod.* vii. 23.—Bras.
polycephala, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 458 = velutina.
polymorpha, *DC. Coll. Mém.* ix. t. 13.—Bras.
racemosa, *DC. Prod.* vii. 23.—Bras.
tomentosa, Bong. Nov. Comp. Bras. 43. t. 9 = paniculata.
tomentosa, Schlecht. ex Mart. in Flora, xxiv. (1841) II. Beibl. 109 = velutina.
velutina, *Bong. Nov. Comp. Bras.* 41. t. 8.—Bras.

MOQUINIA, Spreng. f. Tent. Syst. Suppl. 9 (1828) = **Loranthus**, Linn.
rubra, Spreng. f. l. c. = L. glaucus.

MORA, Schomb. ex Benth. in Trans. Linn. Soc. xviii. (1839) 210. t. 16 = **Dimorphandra**, Schott (Legumin.).
conjugata, Splitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 109.—Guian.
excelsa, Benth. in Trans. Linn. Soc. xviii. (1839) 210 = D. Mora.

MORAEA, Mill. ex Linn. Sp. Pl. ed. II. 59 (1762). *IRIDEAE*, Benth. & Hook. f. iii. 688.
DIETES, Salisb. in Trans. Hort. Soc. i. (1812) 307.
FRENCHENIA, Eckl. Top. Verz. 14 (1827).
HELIXYRA, Salisb. in Trans. Hort. Soc. i. (1812) 305.
HYMENOSTIGMA, Hochst. in Flora, xxvii. (1844) 24.
IRIDOPSIS, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 270.
NARON, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 419.
PHAIANTHES, Rafin. Fl. Tellur. iv. 30 (1836).
VIEUSSEUXIA, Delar. Diss. 31. t. 5 (1766).
acorifolia, H. B. K. Nov. Gen. et Sp. i. 322 = Orthrosanthus chimboracensis.
africana, Thunb. Diss. Moraea, 7 = Aristaea cyanea.
alata, Vahl, Enum. i. 154 = Sisyrinchium palmifolium.
andongensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 271.—Afr. trop.
angusta, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 240.—Afr. austr.
aphylla, Linn. f. Suppl. 99 ; Thunb. Diss. Moraea, 9 = Bobartia aphylla.
Aristea, Lam. Illustr. i. 114 = Aristea cyanea.
aurantiaca, *A. Dietr. Sp. Pl.* ii. 485.—Afr. austr.
aurantiaca, *Eckl. Top. Verz.* 15.—Afr. austr.
barbigera, Salisb. in Trans. Hort. Soc. i. (1812) 306 = pilosa.
Bermudiana, Thunb. Diss. Moraea, 7 = Sisyrinchum Bermudiana.
bicolor, *Steud. Nom.* ed. II. ii. 159.—Afr. austr.
bituminosa, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 240 = viscaria.
brasiliensis, Link & Otto, ex Steud. Nom. ed. II. ii. 159 (Quid ?).—Bras.
bulbifera, Jacq. Hort. Schoenb. ii. 38 = Sparaxis bulbifera.

MORAEA:—
Burchellii, *Baker, in Journ. Linn. Soc.* xvi. (1877) 132, nomen.—Afr. austr.
caerulea, Thunb. Diss. Moraea, 12. t. 2. f. 2 = Aristea capitata.
Candelabrum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 271.—Afr. trop.
Candolleana, *Spreng. Syst.* i. 164.—Afr. austr.
catenulata, Lindl. Bot. Reg. t. 1074 = iridioides.
chimboracensis, H. B. & K. Nov. Gen. et Sp. i. 322 = Orthrosanthus chimboracensis.
chinensis, Thunb. Fl. Jap. 34 = Pardanthus chinensis.
ciliata, Drège, ex Baker, in Journ. Linn. Soc. xvi. (1877) 129 = papilionacea.
ciliata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 241.—Afr. austr.
coerulea, Auct. = *caerulea*.
collina, Thunb. Diss. Moraea, 11 = Homeria collina.
corniculata, Lam. Illustr. i. 115 = Aristea umbellata.
crispa, *Ker-Gawl. in Bot. Mag.* t. 1284.—Afr. austr.
crispa, Thunb. Diss. Moraea, 13 = undulata.
cultriformis, Schott, ex Klatt, in Mart. Fl. Bras. iii. 1. 521 = Cypella gracilis.
decussata, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 367 = crispa.
diandra, Vahl, Enum. ii. 154 = Diplarrhena Moraea.
dichotoma, Thunb. Prod. Pl. Cap. 186 = Aristea dichotoma.
diversifolia, *Baker, in Journ. Linn. Soc.* xvi. (1877) 130.—Abyss.
edulis, *Ker-Gawl. in Bot. Mag.* t. 613.—Afr. austr.
elegans, Jacq. Hort. Schoenb. i. 6 = Homeria elegans.
exaltata, Steud. Nom. ed. II. ii. 160 = Homeria collina.
falcifolia, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 366.—Afr. austr.
fasciculata, *Klatt, l. c.*—Afr. austr.
Ferrariola, Jacq. Coll. iv. 141 = Ferraria Ferrariola.
filiformis, Linn. f. Suppl. 100 = Bobartia filiformis.
fimbriata, *Klatt, in Linnaea*, xxxiv. (1865–66) 561.—Afr. austr.
fimbriata, Loisel. in Delaun. Herb. Amat. vi. 412 = Iris japonica.
flaccida, Steud. Nom. ed. II. ii. 160 = Homeria collina.
flexuosa, Linn. f. Suppl. 100 = Hexaglottis longifolia.
foliosa, H. B. & K. Nov. Gen. et Sp. i. 322 = Nemastylis multiflora.
fugax, Jacq. Hort. Vindob. iii. 14. t. 20 = edulis.
fugax, *Steud. Nom.* ed. II. ii. 160.—Afr. austr.
fugax, Tenore, Syll. Fl. Neap. 26 = Iris Sisyrinchium.
galaxioides, *Baker, in Journ. Linn. Soc.* xvi. (1877) 130, nomen.—Afr. austr.
Gawleri, Spreng. Syst. v. (Index) 462 = crispa.
gigantea, *Klatt, in Linnaea*, xxxv. (1867–68) 381.—Afr. austr.
gladiata, Thunb. Diss. Moraea, 8 = Bobartia gladiata.
gladioloides, H. B. & K. Nov. Gen. et Sp. i. 322 = Orthosanthus chimboracensis.
glaucopis, *Drapiez, Encyc.* iv. (1836) cum ic. col.—Afr. austr.
glutinosa, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 271.—Afr. trop.
gracilis, Baker, l. c. 272 = edulis.
gracilis, *A. Dietr. Sp. Pl.* ii. 478.—Afr. austr.
grandiflora, Eckl. Top. Verz. 14 = Homeria collina.
grandiflora, Pers. Syn. i. 49 = Sisyrinchium grandiflorum.
guttata, Stokes, Bot. Comment. i. 229 = Pardanthus chinensis.
hantamensis, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 365 = ciliata.
Herberti, Lindl. in Bot. Reg. t. 949 = Cypella Herberti.
hirsuta, Ker-Gawl. Irid. Gen. 43; A. Dietr. Sp. Pl. ii. 478 = papilonacea.
humilis, *Eckl. Top. Verz.* 15.—Afr. austr.
iridioides, *Linn. Mant.* i. 28.—Afr. austr.
irioides, Gaertn. Fruct. i. 40 = praec.
iriopetala, *Linn. f. Suppl.* 100.—Afr. austr.
iriopetala, Willd. Sp. Pl. i. 244 = tristis, Homeria collina.
ixioides, Thunb. Diss. Moraea, 8 = Libertia ixioides.
juncea, *Linn. Sp. Pl.* ed. II. 59.—Afr. austr.
linearis, H. B. & K. Nov. Gen. et Sp. i. 321 = Herbertia lineata.
lineata, Steud. Nom. ed. II. ii. 160 = Homeria lineata.
longiflora, *Ker-Gawl. in Bot. Mag.* t. 712.—Afr. austr.
longifolia, Pers. Syn. i. 49 = Hexaglottis longifolia.

MORAEA:—
longifolia, Sweet, Hort. Brit. ed. II. 497 = edulis.
longispatha, Klatt, in Linnaea, xxxiv. (1865–66) 360 = spathacea.
lugens, Linn. f. Suppl. 99 = Aristea melaleuca.
lurida, *Ker-Gawl. in Bot. Reg.* t. 312.—Afr. austr.
magellanica, Willd. Sp. Pl. i. 241 = Tapeinia magellanica.
melaleuca, Thunb. Prod. Pl. Cap. 10 = Aristea melaleuca.
miniata, Andr. Bot Rep. t. 404 = Homeria collina.
minor, *Eckl. Top. Verz.* 15.—Afr. austr.
minuta, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 241 = ciliata.
mira, *Klatt, in Trans. S. Afr. Phil. Soc.* iii. (1885) 202.—Afr. austr.
multiflora, *Eckl. Top. Verz.* 14, nomen.—Afr. austr.
Northiana, Schneevogt, Ic. tt. 41, 42; Andr. Bot. Rep. t. 255 = Marica Northiana.
ochroleuca, *Drapiez, Encycl.* iv. (1836).—Afr. austr.
odora, Salisb. Parad. Lond. t. 10 = edulis.
ovata, Thunb. Prod. Pl. Cap. 186 = Lapeyrousia fissifolia.
palmifolia, Jacq. Ic. Pl. Rar. ii. t. 227 = Eleutherina plicata.
palmifolia, Thunb. Diss. Moraea, 8 = Sisyrinchium palmifolium.
papilionacea, *Ker-Gawl. in Bot. Mag.* t. 750.—Afr. austr.
pauciflora, *Eckl. Top. Verz.* 14, nomen.—Afr. austr.
Pavonia, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 240.—Afr. austr.
Pavonia, Thunb. Diss. Moraea, 14 = Tigridia Pavonia.
pilosa, Wendl. Bot. Beobacht. 42 = ciliata.
plicata, Sw. Fl. Ind. Occ. i. 82 = Eleutherine plicata.
plumaria, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 240 = iriopetala.
polyanthos, *Thunb. Diss. Moraea*, 11.—Afr. austr.
polyphylla, Baker, in Journ. Linn. Soc. xvi. (1877) 130, nomen. = fasciculata.
polystachya, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 240.—Afr. austr.
porrifolia, Steud. Nom. ed. II. ii. 160 = Homeria lineata.
pusilla, Thunb. Diss. Moraea, 7 = Aristea pusilla.
racemosa, Drège, ex Baker, in Journ. Linn. Soc. xvi. (1877) 131 = ramosa.
ramosa, Endl. in Harting. Parad. Vindob. t. 7 = Gladiolus carneus.
ramosa, *Ker-Gawl. in Bot. Mag.* t. 771.—Afr. austr.
repens, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 175 = Aristea repens.
Robinsoniana, *C. Moore & F. Muell. Fragm.* vii. 153.—Ins. Dom. Howe.
sertula, Jacq. Hort. Schoenb. i. 6. t. 11 = Sisyrinchium striatum.
setacea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 240.—Afr. austr.
setacea, Steud. Nom. ed. II. ii. 160 = praec.
setifolia, *Eckl. Top. Verz.* 14, nomen.—Afr. austr.
similis, *Eckl. l. c.* 15.—Afr. austr.
Sisyrinchium, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 241 = Iris Sisyrinchium.
sordescens, Jacq. Coll. Suppl. 29 = tristis.
spathacea, *Ker-Gawl. in Bot. Mag.* sub t. 1103.—Afr. austr.
spathacea, Thunb. Diss. Moraea, 9 = Bobartia indica.
spicata, Ker-Gawl. in Bot. Mag. t. 1283 = Homeria elegans.
spiralis, Linn. f. Suppl. 99; Thunb. Diss. Moraea, 6 = Aristea spiralis.
spithamaea, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 271.—Afr. trop.
Tenoreana, Sweet, Brit. Flow. Gard. t. 110 = Iris Sisyrinchium.
tenuis, *Ker-Gawl. in Bot. Mag.* t. 1047.—Afr. austr.
textilis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 270.—Afr. trop.
Tigridia, Baker, in Journ. Linn. Soc. xvi. (1877) 136 = Tigridia Pavonia.
tricolor, Andr. Bot. Rep. t. 83 = ciliata.
tricuspis, *Ker-Gawl. in Bot. Mag.* t. 696.—Afr. austr.
tripetala, *Ker-Gawl. l. c.* t. 702.—Afr. austr.
tristis, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 241.—Afr. austr.
umbellata, Thunb. Diss. Moraea, 13 = Aristea umbellata.
undulata, *Ker-Gawl. Gen. Irid.* 43.—Afr. austr.

MORAEA:—
undulata, Thunb. Diss. Moraea, 14 = Ferraria undulata.
unguicularis, *Lam. Encyc.* iv. 275.—Afr. austr.
unguiculata, Ker-Gawl. in Bot. Mag. t. 593 = praec.
vaginata, [DC. in] Red. Lil. i. t. 56 = Marica Northiana.
vegeta, Herb. Linn. ex Baker, in Journ. Linn. Soc. xvi. (1877) 130 = iriopetala.
vegeta, Jacq. Ic. Pl. Rar. t. 224 = edulis.
vegeta, Linn. Sp. Pl. 59 = tristis.
vegeta, Mill. Gard. Dict. ed. VIII. n. 1 = iridioides.
villosa, Ker-Gawl. ex Reichb. Fl. Exot. i. 22. t. 19. f. 1 = M. Pavonia.
virgata, Jacq. Coll. iii. 194 = Hexaglottis virgata.
viscaria, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 240.—Afr. austr.
Welwitschii, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 270.—Afr. trop.
zambesiaca, *Baker, in Journ. Linn. Soc.* xvi. (1877) 130, nomen.—Afr. trop.
Zeyheri, *Lehm. in Ann. Sc. Nat.* Sér. II. vi. (1836) 107.—Afr. austr.

MORANDA, Scop. Introd. 289 (1777) = **Pentapetes**, Linn. (Stercul.).

MORELIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 232. *RUBIACEAE* [Benth. & Hook. f. ii. 88.]
senegalensis, *A. Rich. l. c.*—Afr. trop.

MORELLA, Lour. Fl. Cochinch. 548 (1790) = **Myrica**, Linn.
rubra, Lour. l. c. = M. Nagi.

MORELOSIA, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 1 (1824) = **Bourreria**, P. Br. (Borag.).
Huanita, Lex. l. c. = B. Huanita.

MORELOTIA, Gaudich. in Freyc. Voy. Bot. 416. t. 28 (1826) = **Gahnia**, Forst. (Cyperac.).
gahniaeformis, Gaudich. l. c. = G. Gaudichaudii.

MORENIA, Ruiz & Pav. Prod. 150. t. 32 (1794) = **Chamaedorea**, Willd. (Palm.).
Chonta, Phil. in Bot. Zeit. xiv. (1856) 648; et in Anal. Univ. Chil. (1856) 168 = Juania australis.
corallina, Karst. in Linnaea, xxviii. (1856) 274.—N. Granat.
corallocarpa, Hort. ex H. Wendl. Ind. Palm. 29.—N. Granat.?
Ernesti-Augusti, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 3 = C. Ernesti-Augusti.
fragrans, Ruiz & Pav. Syst. Veg. 299 = C. fragrans.
integrifolia, Trail, in Journ. Bot. xiv. (1876) 331.—Bras.
Lindeniana, H. Wendl. in Bot. Zeit. xvii. (1859) 17 = C. Lindeniana.
oblongata, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 3 = C. Sartorii.
pauciflora, Drude, in Mart. Fl. Bras. iii. II. 526.—Bras.
Poeppigiana, Mart. Hist. Nat. Palm. iii. 161. tt. 140, 141.—Peruv.

MORENOA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 5 (1824) = **Ipomoea**, Linn. (Convolv.).
globosa, La Llave, l. c. = I. peduncularis.
grandiflora, La Llave, l. c. 17 = I. Funis.
patula, La Llave, ex Steud. Nom. ed. II. ii. 160 (Quid?).—Mexic.

MORETTIA, DC. Syst. ii. 426 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 70.
NECTOUXIA, DC. Syst. ii. 149 (1821).
asperrima, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 60 = philaeana.
canescens, *Boiss. Diagn.* Ser. I. viii. 17.—Arab.
parviflora, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 60.—Arab.
philaea, Spreng. Syst. ii. 883 = philaeana.
philaeana, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 60 = canescens.
philaeana, *DC. Syst.* ii. 426.—Arab.; Nubia.
Revoili, *Franch. in Revoil, Çomali*, 10.—Afr. trop.

MORGAGNIA, Bubani, in Nouv. Ann. Sc. Nat. Bologn. ix. (1848) 92 = **Simethis**, Kunth (Liliac.).
bicolor, Bubani, l. c. 94 = S. bicolor.

MORGANIA, R. Br. Prod. 441 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 950.
aspera, Spreng. Syst. ii. 803 = Vandellia crustacea.
caerulea, Poir. Encyc. Suppl. iv. 1 = glabra.
callitrichoides, *Spreng. Syst.* ii. 803.—Venezuela.
floribunda, *Benth. in Mitch. Journ. Trop. Austral.* 384.—Austral.
glabra, *R. Br. Prod.* 441.—Austral.
hyssopioides, Spreng. Syst. ii. 803 = Ilysanthes hyssopioides.
juncea, Spreng. l. c. = Dopatrium junceum.
lucida, Spreng. l. c. 802 = Vandellia crustacea.
parviflora, *Benth. Fl. Austral.* iv. 489.—Austral.
pubescens, *R. Br. Prod.* 441.—Austral.
rotundifolia, Spreng. Syst. ii. 803 = Ilysanthes rotundifolia.
strigosa, Spreng. l. c. = Gratiola strigosa.

MORICANDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 173 = seq.

MORICANDIA, DC. Syst. ii. 626 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 85.
CRANTZIA, Lag. ex DC. Syst. ii. 509, 627, in syn. (1821).
DOUEPEA, Cambess. in Jacquem. Voy. Bot. 18. t. 18 (1837).
ORYCHOPHRAGMUS, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 81.
OUDNEYA, R. Br. in Denh. & Clapp. Trav. App. 20 (1826).
alyssifolia, *Pomel, Nouv. Mat. Fl. Atl.* 366.—Alger.
arvensis, *DC. Syst.* ii. 626.—Europ.; Reg. Mediterr.
baetica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 8.—Hispan.
clavata, *Boiss. & Reut. Diagn.* Ser. II. v. 25.—Alger.; Arab.
crassifolia, J. Gay, ex Boiss. Fl. Orient. i. 389 = Diplotaxis Sieberi.
dumosa, *Boiss. Diagn.* Ser. I. viii. 25.—Arab.
foetida, *Bourg. ex Coss. Notes Crit.* 143.—Hispan.
hesperidiflora, DC. Syst. ii. 627 = Diplotaxis Sieberi.
longirostris, *Pomel, Nouv. Mat. Fl. Atl.* 367.—Alger.
pallida, *Pomel, l. c.* 365.—Alger.
papillosa, *Steud. Nom.* ed. II. ii. 161, nomen.—Arab.
patula, *Pomel, Nouv. Mat. Fl. Atl.* 225.—Alger.
Ramburii, *Webb, Iter Hisp.* 73.—Hispan.
sinaica, *Boiss. Fl. Orient.* i. 386.—Arab.; Pers.; Beluchist.
sonchifolia, *Hook. Bot. Mag.* t. 6243.—China.
spinosa, *Pomel, Nouv. Mat. Fl. Atl.* 365.—Alger.
suffruticosa, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 395.—Alger.
teretifolia, *DC. Syst.* ii. 628.—Afr. bor.
tortuosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 172.—Ind. bor. occ.

MORIERA, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 380. *CRUCIFERAE*, Benth. & Hook. f. i. 94.
CRENULARIA, Boiss. l. c. (1842) 181.
LIPOPHRAGMA, Schott & Kotschy, ex Boiss. Diagn. Ser. II. v. 42 (1856).
cabulica, Boiss. Diagn. Ser. II. i. 41 = spinosa.
spinosa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 182.—Persia.
umbellata, *Boiss. Diagn.* Ser. I. vi. 16.—Persia.
virgata, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 183 = Aethionema elongatum.

MORIERINA, Vieill. in Bull. Soc. Linn. Norm. ix. (1865) 344. *RUBIACEAE*, Benth. & Hook. f. ii. 47.
montana, *Vieill. l. c.*—N. Caled.
propinqua, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870-71) 401.—N. Caled.

MORILANDIA, Neck. Elem. ii. 99 (1790) = **Cliffortia**, Linn. (Rosac.).

MORINA, Tourn. ex Linn. Syst. ed. I. (1735). *DIPSACEAE*, Benth. & Hook. f. ii. 158.
ASAPHES, Spreng. Syst. iv. Cur. Post. 222 (1827).
Aucheri, Jaub. & Spach, Illustr. v. 32 = persica.
betonicoides, *Benth. in Hook. Ic. Pl.* t. 1171.—Reg. Himal.

MORINA :—
breviflora, Edgew. in Trans. Linn. Soc. xx. (1846) 62 = Coulteriana.
Coulteriana, *Royle, Illustr. Bot. Himal.* 245.—Reg. Himal.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 9.—China.
elegans, Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 67 = longifolia.
graeca, Jaub. & Spach, Illustr. v. 32 = persica.
kokanica, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) I. 14.—Turkestan.
Lehmanniana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 321.—Turkestan.
longifolia, *Wall. Cat.* n. 426.—Reg. Himal.
nana, Wall. l. c. n. 424 = nepalensis.
nepalensis, *D. Don, Prod. Fl. Nep.* 161.—Reg. Himal.
orientalis, Mill. Gard. Dict. ed. VIII = persica.
parviflora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 373.—Soongar.
persica, *Linn. Sp. Pl.* 28.—Oriens; Reg. Himal.
polyphylla, *Wall. Cat.* n. 425.—Reg. Himal.
subinermis, *Boiss. Diagn.* Ser. II. vi. 94.—Bithynia.
Tournefortii, Jaub. & Spach, Illustr. v. 30 = persica.
verticillata, Moench, Meth. Suppl. 186 = persica.
Wallichiana, Royle, Illustr. Bot. Himal. 245. t. 55. f. 1 = persica.

MORINDA, Linn. Gen. ed. I. 57 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 117.
GUTTENBERGIA, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1846) 2.
ROJOC, Adans. Fam. ii. 146 (1763).
SPHAEROPHORA, Blume, Mus. Bot. Lugd. Bat. i. 179, f. 36 (1850).
acutifolia, *F. Muell. Fragm.* ix. 179.—Austral.
angustifolia, Roth, Nov. Pl. Sp. 147 = tinctoria.
angustifolia, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 547.—Ind. or.
artensis, *Montr. in Mém. Acad. Lyon*, x. (1860) 219.—N. Caled.
aspera, Wight & Arn. Prod. 420 = tinctoria.
axillaris, Poir. in Lam. Encyc. iv. 315 = Saldinia Pseudomorinda.
Billardieri, *Baill. Adansonia*, xii. (1879) 231.—N. Caled.
bracteata, *Hort. Angl. ex Link, Enum. Hort. Berol.* i. 225 (an seq.?).—Hab.?
bracteata, Roxb. Hort. Beng. 15; Fl. Ind. i. 544 = citrifolia.
bucidaefolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 41.—Ins. Fiji.
celebica, *Miq. Fl. Ind. Bat.* ii. 246.—Ins. Celebes.
Chachuca, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 536 = tinctoria.
choriophylla, *Baill. Adansonia*, xii. (1879) 246.—N. Caled.
chrysorrhiza, *DC. Prod.* iv. 450.—Afr. trop.
citrifolia, Hunter, in As. Res. iv. (1795) 37; Bedd. Fl. Sylv. t. 220 = tinctoria.
citrifolia, *Linn. Sp. Pl.* 176.—As. et Austral. trop.; Ins. Pacif.
cochinchinensis, *DC. Prod.* iv. 449.—Cochinch.
Coreia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 537 = tinctoria.
Cumingiana, Vidal, Phan. Cuming. Philipp. 184 = Lucinaea Cumingiana.
exserta, Roxb. Hort. Beng. 15; Fl. Ind. i. 545 = tinctoria.
fasciculata, *Benth. Pl. Hartw.* 123.—Guatemala.
ferruginea, *A. Rich. Fl. Cub. Fanerog.* ii. 19.—Cuba.
Forsteri, *Seem. Fl. Vit.* 129.—Ins. Fiji.
gemella, *Miq. Fl. Ind. Bat.* ii. 247.—N. Guin.
geminata, DC. Prod. iv. 447 = citrifolia.
glomerata, *Miq. Fl. Ind. Bat.* ii. 247.—N. Guin.
Grayi, *Seem. Fl. Vit.* 130.—Ins. Fiji.
Jackiana, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 227.—Malaya.
jasminoides, *A. Cunn. in Hook. Bot. Mag.* t. 3351.—Austral.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 214.—Sumatra.
lanceolata, Wall. Cat. n. 8435 = persicaefolia.
Lastelliana, *Baill. Adansonia*, xii. (1879) 231.—Madag.
laxa, *Bartl. ex DC. Prod.* iv. 449.—Mexic.
leiantha, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 313 = tinctoria.

MORINDA :—
ligulata, Blanco, Fl. Filip. ed. II. 105 = citrifolia.
littoralis, Blanco, l. c. 109 = citrifolia.
longiflora, *G. Don, Gen. Syst.* iii. 545.—Afr. trop.
lucida, Benth. in Hook. Niger Fl. 406 = citrifolia.
lucida, *A. Gray, in Proc. Am. Acad.* iv. (1860) 41.—Ins. Fiji.
macrophylla, Desf. Cat. Hort. Par. ed. III. 404 = citrifolia.
microcephala, *Bartl. ex DC. Prod.* iv. 449.—Ins. Philipp.
microphylla, D. Dietr. Syn. Pl. i. 784 = microcephala.
mollis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 41.—Ins. Fiji.
Mudia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 536 = tinctoria.
multiflora, Roxb. Hort. Beng. 15; Fl. Ind. i. 546 = tinctoria.
muscosa, Jacq. Enum. Pl. Carib. 16 = Cephaëlis muscosa.
myrtifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 41.—Ins. Fiji.
Naudia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 156. err. typ. = *Mudia*.
nodosa, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 537 = tinctoria.
Padavara, Juss. Gen. 206.—Cf. Poir. Encyc. Suppl. iv. 3, in obs. = umbellata.
palmetorum, *DC. Prod.* iv. 448.—Afr. trop.
panamensis, *Seem. Bot. Voy. Herald*, 136.—Panama.
parvifolia, *Bartl. ex DC. Prod.* iv. 449.—Ins. Philipp.
peduncularis, *H. B. & K. Nov. Gen. et Sp.* iii. 380.—Venezuela.
persicaefolia, *Buch.-Ham. in Trans. Linn. Soc.* xiii. (1822) 535.—Burma; Malaya.
phylliraeoides, *Labill. Sert. Austro-Caled.* 49. t. 49.—N. Caled.
piperiformis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 135.—Ins. Timor.
polyneura, *Miq. Fl. Ind. Bat.* ii. 244.—Java.
polysperma, Jack, in Malay. Misc. i. (1820) n. v. 14 = Lucinaea Morinda.
pubescens, *Sm. in Rees, Cycl.* xxiv. n. 3.—Ins. Maurit.
quadrangularis, G. Don, Gen. Syst. iii. 545 = citrifolia.
reticulata, Benth. Fl. Austral. iii. 424 = Coelospermum decipiens.
retusa, *Poir. in Lam. Encyc.* iv. 316.—Madag.
rigida, *Miq. Fl. Ind. Bat.* ii. 246.—Malaya.
Rojoc, Lour. Fl. Cochinch. i. 174 = umbellata.
Royoc, Blanco, Fl. Filip. ed. I. 148 = tinctoria.
Royoc, Linn. Sp. Pl. 176 = umbellata.
sarmentosa, *Blume, Bijdr.* 1006.—Java.
scandens, Roxb. Hort. Beng. 86; Fl. Ind. i. 548 = umbellata.
speciosa, Wall. Cat. n. 8436 = Rennellia speciosa.
squarrosa, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 535 = angustifolia.
stenophylla, Spreng. Syst. i. 749 = tinctoria.
sumatrana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 213.—Sumatra.
tetrandra, Jack, in Malay. Misc. i. (1820) n. v. 13 = umbellata.
Teysmanniana, Miq. Fl. Ind. Bat. ii. 244 = tinctoria.
tinctoria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = citrifolia.
tinctoria, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 543.—Ind. or.; Malay.
tomentosa, Heyne, in Roth, Nov. Pl. Sp. 147 = tinctoria.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* iii. 380.—N. Granat.
umbellata, Labill. ex Baill. Adansonia, xii. (1879) 232 = Billardieri.
umbellata, *Linn. Sp. Pl.* 176.—As. et Austral. trop.
umbellata, Lour. Fl. Cochinch. 173 = cochinchinensis.
Vieillardi, *Baill. Adansonia*, xii. (1879) 230.—N. Caled.
villosa, *Hook. f. Fl. Brit. Ind.* iii. 158.—Reg. Himal.
Wallichii, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 313.—Burma.
Zollingeriana, Miq. Fl. Ind. Bat. ii. 243 = tinctoria.

MORINDOPSIS, Hook. f. in Benth. & Hook. f. Gen. ii. 93 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
capillaris, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 189.—Burma.

MORINGA, Burm. Thes. Zeyl. 162. t. 75 (1737).
MORINGEAE, Benth. & Hook. f. i. 430.
ALANDINA, Neck. Elem. ii. 456 (1790).
ANOMA, Lour. Fl. Cochinch. 278 (1790).
HYPERANTHERA, Forsk. Fl. Aegypt. Arab. 67 (1775).
aptera, *Gaertn. Fruct.* ii. 315.—Nubia; Arabia.
arabica, Pers. Syn. i. 460 = aptera.
concanensis, *Nimmo, in J. Grah. Cat. Bomb. Pl.* 43.—Ind. or.
edulis, Medic. Theod. 48 = pterygosperma?
erecta, Salisb. Prod. 326 = pterygosperma.
Nux-eben, Desf. Cat. Hort. Par. ed. III. 303 = pterygosperma.
octogona, Stokes, Bot. Mat. Med. ii. 466 = pterygosperma.
oleifera, Lam. Encyc. i. 398 = pterygosperma.
parvifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 20 = pterygosperma.
polygona, DC. Prod. ii. 478 = pterygosperma.
pterygosperma, *Gaertn. Fruct.* ii. 314.—Ind. bor. occ. (alibi cult.).
robusta, Boj. Hort. Maurit. 117 = pterygosperma?
zeylanica, Pers. Syn. i. 461 = pterygosperma.

MORISIA, J. Gay, in Colla, Hort. Rip. App. iv. 50 (1829). *CRUCIFERAE*, Benth. & Hook. f. i. 100.
hypogaea, *J. Gay, l. c.*—Corsic.; Sardin.

MORISIA, Nees, in Edinb. N. Phil. Journ. (Oct. 1834) 265 = **Rynchospora**, Vahl (Cyperac.).
capitata, Nees, in Linnaea, ix. (1834) 295 = R. Wallichiana.
Wallichii, Nees, in Edinb. N. Phil. Journ. (Oct. 1834) 265; et in Wight, Contrib. 115 = R. Wallichiana.

MORISONA, Plum. ex Linn. Gen. ed. I. 364 (1737) = seq.

MORISONIA, Linn. Sp. Pl. ed. I. 503 (1753). *CAPPARIDEAE*, Benth. & Hook. f. i. 107.
americana, *Linn. l. c.*; *Jacq. Am. Select.* t. 97.—Am. austr.
asiatica, Stocks, in Proc. Linn. Soc. i. (1848) 367, err. typ. = Monsonia senegalensis.
flexuosa, Linn. Pl. Jam. Pugill. 14; Amoen. Acad. v. 398 = Capparis Cynophallophora.
Imrayi, *Griseb. Fl. Brit. W. Ind.* 19.—Ins. Dominic.
Lawiana, Stocks, in Proc. Linn. Soc. i. (1848) 367, err. typ. = Monsonia senegalensis.
multiflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 87.—N. Granat.

MORITZIA, DC. ex Meissn. Gen. 280 (1840). *BORAGINEAE*, Benth. & Hook. f. ii. 859.
MERATIA, A. DC. in DC. Prod. x. 104 (1846).
ciliata, *DC. Prod.* x. 56.—Bras.
dasyantha, *Fresen. in Mart. Fl. Bras.* viii. 63.—Bras.
Sellowiana, *Fresen. l. c.*—Bras.

MORITZIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 279 (1873) = **Podocoma**, Cass. (Compos.).
glandulosa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 14, in nota = P. glandulosa.

MORMODES, Lindl. Introd. Nat. Syst. ed. II. 446 (1836). *ORCHIDEAE*, Benth. & Hook. f. iii. 552.
CYCLOSIA, Klotzsch, in Otto & Dietr. Allg. Gartenz. vi. (1838) 305.
aromatica, *Lindl. in Bot. Reg.* (1841) *Misc.* 76; (1843) t. 56.—Mexic.
atropurpurea, Hook. Bot. Mag. t. 4577 = Hookeri.
atropurpurea, *Lindl. Introd. Nat. Syst.* ed. II. 446.—Ind. or.
barbata, Lindl. & Paxt. Flow. Gard. ii. (1851-52) 57 = Hookeri.
brachystachya, Klotzsch, Ind. Sem. Hort. Berol. (1852) 13 = M. Buccinator.
Buccinator, *Lindl. Bot. Reg.* (1840) *Misc.* 9.—Mexic.
Cartoni, *Hook. Bot. Mag.* t. 4214.—Bras.
Colossus, *Reichb. f. in Bot. Zeit.* x. (1852) 636.—Am. centr.
convoluta, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 98, sub t. 93.—Bras.
Dayana, *Reichb. f. in Gard. Chron.* (1885) II. 552.—Mexic.

MORMODES:—
flavida, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 113 = M. Buccinator.
fractiflexa, *Reichb. f. in Gard. Chron.* (1872) 141.—Costa Rica.
Greenii, Hook. f. Bot. Mag. t. 5802 = M. Uncia.
Histrio, *Linden & Reichb. f. in Otto, Hamb. Gartenz.* xv. (1859) 54.—Mexic.
holochrysa, Reichb. f. Xenia Orch. ii. 172, in obs. t. 170 = variabilis.
Hookeri, *Lem. Jard. Fleur.* i. (1851) *Misc.* 116; ii. (1852) t. 118.—Panama.
ignea, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 97, t. 93.—N. Granat.
incisa, Reichb. f. in Gard. Chron. (1879) II. 582, lapsu = M. Uncia.
lentiginosa, Hook. Bot. Mag. t. 4455 = M. Buccinator.
leucochila, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 289 = M. Buccinator.
lineata, *Batem. ex Lindl. Bot. Reg.* (1841) *Misc.* 52; (1843) t. 43.—Guatemala.
luxata, *Lindl. l. c.* (1842) *Misc.* 60; (1843) t. 33.—Mexic.
macrantha, Lindl. & Paxt. Flow. Gard. iii. (1852-53) 98, sub t. 93 = M. Colossus.
marmorea, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 290 = M. Buccinator.
Ocanae, *Linden & Reichb. f. in Walp. Ann.* vi. 581 (Ocannae).—N. Granat.
pardina, *Batem. Orch. Mex. & Guatem.* t. 14.—Mexic.
rosea, *Rodrig. Orch. Nov.* i. 131.—Bras.
sinuata, *Reichb. f. & Warm. in Otia Bot. Hamb.* 90.—Bras.
Skinneri, *Reichb. f. in Gard. Chron.* (1869) 50.—Guatemala.
speciosa, *Linden, ex Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 174.—N. Granat.
Tibicen, *Reichb. f. in Gard. Chron.* (1870) 1085.—N. Granat.?
tigrina, *Rodrig. Orch. Nov.* i. 131.—Bras.
Uncia, *Reichb. f. in Gard. Chron.* (1869) 892.—Mexic.
variabilis, *Reichb. f. l. c.* 1038.—Am. austr.
vitellina, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 290 = M. Buccinator.
Wageneriana, Klotzsch, l. c. 147 = M. Buccinator.
Warszewiczii, Klotzsch, l. c. xxii. (1854) 65 = lineata.
Wendlandi, *Reichb. f. in Walp. Ann.* vi. 581.—Am. centr.
Williamsii, Hort. ex Nichols. Dict. Gard. ii. 385 = luxata.

MORMOLYCE, Fenzl, in Denkschr. Acad. Wien, Math. Nat. i. (1850) 253 (Nov. Gen. et Sp. 1. t. 2). *ORCHIDEAE*, Benth. & Hook. f. iii. 554.
lineolata, *Fenzl, ll. cc.*—Mexic.

MORMORAPHIS, Jack, ex Wall. Cat. n. 4931 (1831) = **Arthrophyllum**, Blume (Araliac.).
sumatrana, Jack, ex Wall. l. c. = A. diversifolium.

MORNA, Lindl. Bot. Reg. t. 1941 (1837); et (1838) t. 9 = **Waitzia**, Wendl. (Compos.).
nitida, Lindl. l. c. t. 1941 = W. corymbosa.
nivea, Lindl. l. c. (1838) t. 9 = W. nivea.

MOROCARPUS, Adans. Fam. ii. 261 (1763) = **Chenopodium**, Tourn.
capitatus, Scop. Fl. Carn. ed. II. i. 6 = C. capitatum.
foliosus, Moench, Meth. 342 = C. capitatum.

MOROCARPUS, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 218 (Fl. Jap. Fam. Nat. ii. 94) = **Debregeasia**, Gaudich. (Urticac.).
affinis, Blume, Mus. Bot. Lugd. Bat. ii. 157.—Java.
angustatus, Blume, ex Zoll. Syst. Verz. Ind. Archip. 101 = D. velutina.
dichotomus, Blume, Mus. Bot. Lugd. Bat. ii. 157 = D. dichotoma.
edulis, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 218 = D. edulis.
japonicus, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 105 = D. edulis?
leucophyllus, Blume, Mus. Bot. Lugd. Bat. ii. 155 = D. Wallichiana.
longifolius, Blume, l. c. 156 = D. velutina.

MOROCARPUS:—
lutescens, Blume, ex Zoll. Syst. Verz. Ind. Archip. 101.—Java.
microcephalus, Benth. in Hook. Kew Journ. vi. (1854) 74 = Villebrunea frutescens.
salicifolius, Blume, Mus. Bot. Lugd. Bat. ii. 157 = D. hypoleuca.
sororius, Miq. Fl. Ind. Bat. i. II. 272.—Java.
velutinus, Blume, Mus. Bot. Lugd. Bat. ii. 156 = D. velutina.
virens, Blume, l. c.—Sumatra.
Wallichianus, Kurz, Fl. For. Brit. Burma, ii. 428; Miq. Ann. Mus. Bot. Lugd. Bat. iv. 306 = D. Wallichiana.
Wallichianus, Thw. Enum. Pl. Zeyl. 262.—Zeylan.

MOROEA, Franch. & Sav. Enum. Pl. Jap. i. 43 (1879), err. typ. = **Moraea**, Linn. (Irid.).

MORONOA, Hort. ex Kunze, in Linnaea, xx. (1847) 19 = **Mikania**, Willd. (Compos.).

MORONOBEA, Aubl. Pl. Gui. ii. 788. t. 313 (1775), partim. *GUTTIFERAE*, Benth. & Hook. f. i. 174.
BLAKSTONIA, Scop. Introd. 276 (1777).
LEUCONOCARPUS, Spruce, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 295 (in Addend.).
PENTADESMOS, Spruce, ex Planch. & Triana, l. c. 295 (1860).
PICCIA, Neck. Elem. ii. 436 (1790).
coccinea, Aubl. Pl. Gui. ii. 788. t. 313 = grandiflora, Symphonia globulifera.
esculenta, Arruda, in Koster, Trav. Braz. 490 = *coccinea*.
exserens, Endl. ex Walp. Rep. i. 394 = Symphonia globulifera.
globulifera, Schlecht. in Linnaea, viii. (1833) 190 = Symphonia globulifera.
grandiflora, *Choisy, in Mém. Soc. Hist. Nat. Par.* i. (1823) 230.—Guian.
montana, Schlecht. in Linnaea, viii. (1833) 189 = grandiflora.
riparia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 296.—Bras.

MOROPHORUM, Neck. Elem. iii. 255 (1790) = **Morus**, Tourn. (Urtic.).

MORPHAEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid?).

MORPHIXIA, Ker-Gawl. Gen. Irid. 105 (1827) = **Ixia**, Linn. (Irid.).
angustifolia, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 384 = I. angustifolia.
aulica, Ker-Gawl. Gen. Irid. 107 = I. scariosa.
capillaris, Ker-Gawl. l. c. 106 = I. capillaris.
columellaris, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 384 = I. columellaris.
Cooperi, Baker, in Journ. Bot. xiv. (1876) 237.—Afr. austr.
curta, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 384 = I. monadelpha.
grandiflora, Klatt, l. c. 383 = I. monadelpha.
incarnata, Ker-Gawl. Gen. Irid. 107 = I. scariosa.
juncifolia, Baker, in Journ. Bot. xiv. (1876) 238 (Watsoniae sp.).—Afr. austr.
lancea, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 383.—Afr. austr.
latifolia, Klatt, l. c. 385 = I. monadelpha.
linearis, Ker-Gawl. Gen. Irid. 106 = I. linearis.
monadelpha, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 384 = I. monadelpha.
nervosa, Baker, in Journ. Bot. xiv. (1876) 237.—Afr. austr.
odorata, Baker, in Journ. Linn. Soc. xvi. (1877) 97 = I. odorata.
paniculata, Baker, Fl. Maurit. 365 = I. paniculata.
purpurea, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 383 = I. monadelpha.
trichorhiza, Baker, in Journ. Bot. xiv. (1876) 237.—Afr. austr.
versicolor, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 385 = I. monadelpha.

MORRENIA, Hort. ex Kunze, in Linnaea, xx. (1847) 19 = **Mikania**, Willd. (Compos.).
odorata, Hort. ex Kunze, l. c. (= *Corynanthelium Moronoa*).—Bras.

MORRENIA, Lindl. Bot. Reg. (1838) Misc. 69. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 761.
brachystephana, *Griseb. in Goett. Abh.* xix. (1874) 205. —Reg. Argent.
odorata, *Lindl. Bot. Reg.* (1838) *Misc.* 71.—Reg. Argent.

MORSTDORFFIA, Steud. Nom. ed. II. ii. 161 (1841) = **Chirita**, Buch.-Ham. (Gentian.).
elongata, Steud. l. c. = Dichrotrichum ternateum.
speciosa, Steud. l. c. = C. Horsfieldii.

MORTONIA, A. Gray, Pl. Wright. i. 34. t. 4 (1852); ii. 28; (in Smithson. Contrib. iii). *CELASTRINEAE*, Benth. & Hook. f. i. 368.
effusa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 453.—Mexic.
Greggii, *A. Gray, Pl. Wright.* i. 35, in adnot.—N. Mexic.
Palmeri, *Hemsl. Diagn. Pl. Nov.* 24.—Mexic.
scabrella, *A. Gray, Pl. Wright.* ii. 28.—N. Mexic.
sempervirens, *A. Gray, l. c.* 35.—Texas.

MORUS, [Tourn.] Linn. Syst. ed. I (1735). *URTICACEAE*, Benth. & Hook. f. iii. 364.
MOROPHORUM, Neck. Elem. iii. 255 (1790).
acidosa, Griff. Notul. iv. 388 = indica?
Alaisia, Deless. ex Moretti, in Giorn. Ist. Lomb. Sc. i. (1841) 182 = laevigata.
alba, Bur. in DC. Prod. xvii. 243, partim = indica.
alba, *Linn. Sp. Pl.* 986.—As. temp.
alpina, Rafin. Man. Am. Mulb. 91, nomen (Quid?). —Hab.?
altissima, *Miq. Fl. Ind. Bat. Suppl.* 415.—Sumatra.
Ampalis, Poir. in Lam. Encyc. iv. 380 = Ampalis madagascariensis.
atrata, Rafin. Man. Am. Mulb. 27, nomen = nigra?
atropurpurea, Roxb. Hort. Beng. 67; Fl. Ind. iii. 595 = alba.
australis, Poir. in Lam. Encyc. iv. 380 = Broussonetia papyrifera.
bifaria, Wall. ex Bur. in DC. Prod. xvii. 248 = Ampalis madagascariensis.
bifaria, Wall. ex Voigt, Hort. Suburb. Calc. 284, partim = Sloetia Sideroxylon.
brasiliensis, Pohl, ex Bur. in DC. Prod. xvii. 230 = Chlorophora tinctoria.
Brunoniana, Endl. Atakta, t. 32 = Pseudomorus Brunoniana.
bullata, Balb. ex Loud. Arboret. iii. 1348 = alba.
byzantina, Sieber, ex Steud. Nom. ed. II. ii. 162 = alba.
Calcar-galli, A. Cunn. ex Loud. Arboret. iii. 1361 = Cudrania javanensis.
canadensis, Poir. in Lam. Encyc. iv. 380 = rubra.
caroliniana, Hort. ex Moretti, in Giorn. Ist. Lomb. Sc. i. (1841) 181 = rubra.
celtidifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 33.—Ecuador.
chinensis, Lodd. Cat. (1836); ex Loud. Arboret. iii. 1350 = alba.
constantinopolitana, Hort. ex Poir. in Lam. Encyc. iv. 381 = alba.
corylifolia, H. B. & K. Nov. Gen. et Sp. ii. 33 = celtidifolia.
cretica, Rafin. Man. Am. Mulb. 23 = nigra?
cucullata, Bonaf. Sul Gels. Philipp. 7 = alba.
Cudranus, Buch.-Ham. ex Wall. Cat. n. 4643 = Cudrania javanensis.
cuspidata, Wall. l. c. 4646 = indica.
dioica, Crantz, Inst. i. 149 = Chlorophora tinctoria.
excelsa, Welw. in Trans. Linn. Soc. xxvii. (1869) 69 = Chlorophora excelsa (in Addend.).
expansa, Hort. ex Rafin. Am. Man. Mulb. 26, nomen = latifolia?
furcata, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
Guzziola, Hort. ex Steud. l. c. = alba.
heterophylla, Loud. Arboret. iii. i. 1361 = alba.
hispanica, Hort. ex Loud. l. c. 1349 = alba.
indica, *Linn. Sp. Pl.* 986.—As. trop.
indica, Macé? ex Ser. Desc. Mur. 226 = alba.
indica, Ser. l. c. 229 = Streblus asper.
insignis, *Bur. in DC. Prod.* xvii. 247.—N. Granat.
insularis, *Spreng. Pugill.* i. 60.—Ins. Pacif.
intermedia, Perr. in Guill. Arch. Bot. i. (1833) 234 = alba.
italica, Poir. ex Lam. Encyc. iv. 377 = alba.
japonica, Audib. ex Ser. Desc. Mur. 226 = alba.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 27 (an praec.?).—Japon.

MORUS:—
javanica, Blume, Bijdr. 488 = Malaisia tortuosa.
Kaempferi, Ser. Desc. Mur. 228 = Broussonetia Kazinoki.
laciniata, Audib. ex Loisel. in Dict. Sc. Nat. xxxiii. 345 = alba.
laciniata, Mill. Gard. Dict. ed. VIII. n. 2 = nigra.
laciniata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = indica.
laciniata, Poir. in Lam. Encyc. iv. 381 = Ruizia variabilis.
laevigata, *Wall. Cat.* n. 4649.—Ind. or.
latifolia, Hort. ex Spach, Hist. Vég. Phan. xi. 45 = alba.
latifolia, Poir. in Lam. Encyc. iv. 381 = indica.
leucophylla, *Miq. Fl. Ind. Bat. Suppl.* 415.—Sumatra.
longifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = Villebrunea sylvatica.
lucida, Hort. ex Loud. Arboret. iii. 1350 = alba.
luzonica, Blanco, Fl. Filip. ed. I. 703 = Allaeanthus luzonicus.
macrophylla, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
macroura, *Miq. Pl. Jungh.* 42.—Java.
Mariettii, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
mauritiana, Jacq. Coll. iii. 206; Fragm. 9 = Ampalis madagascariensis.
membranacea, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
mexicana, Benth. Pl. Hartw. 71 = celtidifolia.
microphylla, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 8 = rubra.
missouriensis, Audib. ex Moretti, in Giorn. Ist. Lomb. Sc. i. (1841) 181 = rubra.
Moretti, Audib. ex Bur. in DC. Prod. xvii. 239 = alba.
Morettiana, Lodd. Cat. (1836); ex Loud. Arboret. iii. 1349 = alba.
multicaulis, Perr. in Mém. Soc. Linn. Par. iii. (1825) 129 = alba.
multicaulis, Rafin. New Fl. Am. iii. 48 (Quid?).—Am. bor.
nana, Audib. ex Loisel. in Dict. Sc. Nat. xxxiii. 345 = alba.
nervosa, Hort. in Bon. Jard. (1836); ex Loud. Arboret. iii. 1349 = alba.
nigra, *Linn. Sp. Pl.* 986.—As. temp.
nigra, Thunb. Fl. Jap. 71 = Broussonetia Kaempferi.
nitida, Willem. in Usteri, Ann. Bot. xviii. (1796) 56 = Ampalis madagascariensis.
norwegica, Tillands, ex Steud. Nom. ed. I. 539; ed. II. ii. 162 = Rubus Chamaemorus.
pabularia, Decne. in Jacquem. Voy. Bot. 149. t. 151 = serrata.
paniculata, Roxb. Hort. Beng. 67; Fl. Ind. iii. 599 = Pipturus velutinus.
papyrifera, Linn. Sp. Pl. 986 = Broussonetia papyrifera.
parvifolia, Rafin. New Fl. Am. iii. 47 (Quid?).—Am. bor.
patavina, Hort. ex Spach, Hist. Vég. Phan. xi. 43 = alba.
pendulina, Ferd. Bauer, in Endl. Prod. Fl. Norf. 40 = Pseudomorus Brunoniana.
pensylvanica, Nois. ex Loud. Hort. Brit. 378 = rubra.
petiolaris, Rafin. Am. Man. Mulb. 24 = nigra?
Plumiera, Steud. Nom. ed. II. ii. 87, 162 = Chlorophora tinctoria.
pumila, Balb. Cat. Hort. Bot. Taur. (1813) 52 = alba.
pyramidalis, Hort. ex Rafin. Man. Am. Mulb. 27 (Quid?).—Hab.?
reticulata, Rafin. l. c. 28 = rubra?
rigida, Hassk. Cat. Hort. Bog. Alt. 74 = Ampalis madagascariensis.
riparia, Rafin. New Fl. Am. iii. 46 (Quid?).—Am. bor.
romana, Lodd. ex Spach, Hist. Vég. Phan. xi. 45 = alba.
rubra, *Linn. Sp. Pl.* 986.—Am. bor.
rubra, Lour. Fl. Cochinch. 555 = alba.
scabra, Moretti, Bibl. Agr. ed. I; ex eod. in Giorn. Ist. Lomb. Sc. i. (1841) 180 = nigra.
scabra, Willd. Enum. Hort. Berol. 967 = rubra.
scandens, Wall. Cat. n. 4652 = Malaisia tortuosa.
serotina, Mart. ex Bur. in DC. Prod. xvii. 239 = alba.
serrata, *Roxb. Hort. Beng.* [103]; *Fl. Ind.* iii. 596.—Reg. Himal.
serrata, Wall. Cat. n. 4648 B = alba.
siciliana, Mill. Gard. Dict. ed. VIII. n. 2 in Erratis = nigra.

MORUS:—
sinensis, Hort. ex Loud. Hort. Brit. 378 = alba.
spinosa, Herb. Mus. Par. ex Gaudich. in Freyc. Voy. Bot. 508, in nota = Chlorophora tinctoria.
stylosa, Ser. Desc. Mur. 225 = alba.
subalba, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
Tataiba, Vell. Fl. Flum. x. t. 21 = Chlorophora tinctoria.
tatarica, Linn. Sp. Pl. 986 = alba.
tatarica, Mill. Gard. Dict. ed. VIII. n. 7 = Streblus asper.
Tatayiba, Parodi, in Anal. Soc. Cient. Argent. v. (1878) 91; Contrib. 39 = Chlorophora tinctoria?
tinctoria, Blanco, Fl. Filip. ed. I. 704 = Cudrania javanensis.
tinctoria, Linn. Sp. Pl. 986 = Chlorophora tinctoria.
Tokwa, Hort. ex C. Koch, Dendrol. ii. i. 447 = alba.
tomentosa, Rafin. Fl. Ludov. 113 = rubra.
tortuosa, Audib. ex Moretti, in Giorn. Ist. Lomb. Sc. i. (1841) 181 = alba.
Venassaini, Hort. ex Steud. Nom. ed. II. ii. 162 = alba.
venosa, Delile, ex Spach, Hist. Vég. Phan. xi. 43 = alba.
vicorum, Jacquem. Voy. Bot. 149 = serrata.
viridis, Buch.-Ham. ex Wall. Cat. n. 4649 = laevigata.
Xanthoxylon, Linn. Syst. ed. X. 1267; Mill. Gard. Dict. ed. VIII. n. 8 = Chlorophora tinctoria.

MORYSIA, Cass. in Dict. Sc. Nat. xxxiii. 59 (1824) = **Athanasia**, Linn. (Compos.).
acerosa, DC. Prod. vi. 92 = A. acerosa.
aspera, Less. Syn. Comp. 263 = A. aspera.
dentata, DC. Prod. vi. 91 = A. dentata.
diversifolia, Cass. in Dict. Sc. Nat. xxxiii. 59 = A. dentata.
fasciculata, Less. Syn. Comp. 262 = A. fasciculata.
juncea, DC. Prod. vi. 91 = A. juncea.
linearifoloba, DC. l. c. 92 = A. pectinata.
longifolia, Less. Syn. Comp. 262 = A. longifolia.
microcephala, DC. Prod. vi. 91 = A. linifolia.
pachyphylla, Sch. Bip. in Flora, xxvii. (1844) 676 = A. dentata.
pauciflora, DC. Prod. vi. 91 = A. linifolia.
pinnata, DC. l. c. = A. pinnata.
spathulata, DC. l. c. = A. spathulata.
velutina, DC. l. c. = A. tomentosa.

MOSCARIA, Pers. Syn. ii. 379 (1807) = **Moscharia**, Ruiz & Pav. (Compos.).

MOSCATELLA, Adans. Fam. ii. 243 (1763) = **Adoxa**, Linn. (Caprifol.).

MOSCHAREA, Salisb. Gen. Pl. Fragm. 25 (1866) = **Muscari**, Tourn. (Liliac.).
Muscari, Salisb. l. c. = Muscari moschatum.

MOSCHARIA, Forsk. Fl. Aegypt. Arab. 158 (1775) = **Ajuga**, Linn. (Labiat.).
arabica, Vitm. Summa Pl. v. 278 = A. Iva.
asperifolia, Forsk. Fl. Aegypt. Arab. p. lxxiv = A. Iva.
Forskâhlii, J. F. Gmel. Syst. 275 = A. Iva.

MOSCHARIA, Ruiz & Pav. Prod. Fl. Per. 103 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 503.
GASTROCARPHA, D. Don, in Trans. Linn. Soc. xvi. (1830) 231.
MOSCHIFERA, Molina, Sagg. Chile, ed. II. 294 (1810).
MOSIGIA, Spreng. Syst. iii. 366, 661 (1826).
pinnatifida, *Ruiz & Pav. Syst. Veg.* i. 186.—Chili.

MOSCHATELLA, Scop. Fl. Carn. ed. II. i. 281 (1772) = **Adoxa**, Linn. (Caprifol.).
Adoxa, Scop. l. c. = A. Moschatellina.

MOSCHATELLINA, [Tourn.] Hall. Enum. Stirp. Helv. i. 412 (1742) = **Adoxa**, Linn. (Caprifol.).
Adoxa, All. Fl. Pedem. i. 134 = A. Moschatellina.
fumariaefolia, Gilib. Fl. Lituan. i. 61 = A. Moschatellina.
tetragona, Moench, Meth. 478 = A. Moschatellina.

MOSCHIFERA, Molina, Sagg. Chile, ed. II. 294 (1810) = **Moscharia**, Ruiz & Pav. (Compos.).
pinnatifida, Molina, l. c. = Moscharia pinnatifida.

MOSCHKOWITZIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 127; et in Abh. Akad. Berl. (1855) 76 t. 8 A = **Begonia**, Linn.
fagopyroides, Klotzsch, ll. cc. 127; 77 = B. fagopyroides.
Wagneria, Klotzsch, ex A. DC. in DC. Prod. xvi. 366 = seq.
Wageneriana, Klotzsch, in Abh. Akad. Berl. (1855) 77 = B. dominicalis.

MOSCHOPSIS, Phil. in Anal. Univ. Chil. (1865) II. 335. *CALYCEREAE*.
Leyboldi, *Phil. l. c.*—Chili.

MOSCHOSMA, Reichb. Consp. 171 (1828) in adnot. *LABIATAE*, Benth. & Hook. f. ii. 1173.
LEHMANNIA, Jacq. ex Steud. Nom. ed. II. ii. 21 (1841); Jacq. f. Eclog. ii. t. 108 (1844).
LUMNITZERA, Jacq. ex Spreng. Syst. ii. 687 (1825).
australe, *Benth. Lab. Gen. et Sp.* 708.—Austral.
Brownii, Heynh. Nom. i. 532, nomen; ii. 421 = australe.
caesium, *Steud. Nom.* ed. II. ii. 163 (sp. dub.).—Hab.?
dimidiatum, Benth. Lab. Gen. et Sp. 24 = polystachyum.
multiflorum, *Benth. in DC. Prod.* xii. 49.—Abyssin.
myriostachyum, *Benth. & Hook. f. Gen.* ii. 1173.—Afr. trop.
ocymoides, Benth. in Wall. Pl. As. Rar. ii. 13 = Mosla ocymoides.
polycladon, Reichb. ex Steud. Nom. ed. II. ii. 162 = Mosla ocymoides.
polystachyum, *Benth. in Wall. Pl. As. Rar.* ii. 13.—Geront. trop.
riparium, *Hochst. in Flora*, xxviii. (1845) 67.—Afr. austr.
tenuiflorum, Heynh. Nom. i. 532; ii. 421 = polystachyum.

MOSCHOXYLON, Meissn. Gen. 49; Comm. 36 (1837) = seq.

MOSCHOXYLUM, A. Juss. in Mém. Mus. Par. xix. (1830) 238 (Mém. Mel. 86. t. 8) = **Trichilia**, P. Br. (Meliac.).
affine, Juss. l. c. 239 = T. Catigua.
Catigna, C. DC. in DC. Monog. Phan. i. 689, sphalm. = seq.
Catigua, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. Catigua.
Cipo, A. Juss. l. c. = T. Cipo.
cuneatum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 413 = T. glabra.
elegans, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. elegans.
emarginatum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 413 = T. emarginata.
multiflorum, Karst. Fl. Columb. i. 71. t. 136 = T. verrucosa.
odoratum, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. odoratum.
pentandrum, Poepp. & Endl. Nov. Gen. et Sp. iii. 39 = T. Tocacheana.
Pleeanum, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. Pleeana.
propinquum, Miq. Stirp. Surin. Sel. 74 = T. propinqua.
Pseudostipulare, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. Pseudostipularis.
Richardianum, A. Juss. l. c. 238 = T. Richardiana.
surinamensis, Miq. Stirp. Surin. Sel. 73 = T. surinamensis.
Swartzii, A. Juss. in Mém. Mus. Par. xix. (1830) 239 = T. moschata.
trachyanthum, Griseb, in Mem. Am. Acad. N. S. viii. (1861) 169 = T. trachyantha.
veraguense, Seem. Bot. Voy. Herald, 93 = T. glabra.

MOSIGIA, Spreng. Syst. iii. 366, 661 (1826) = **Moscharia**, Ruiz & Pav. (Compos.).
pinnatifida, Spreng. l. c. 661 = Moscharia pinnatifida.

MOSINA, Adans. Fam. ii. 272 (1763) = **Ortegia**, Loefl. (Caryoph.).

MOSKERION, Rafin. Fl. Tellur. iv. 21 (1836) = **Narcissus**, Tourn. (Amaryll.).
hispanicum, Rafin. l. c. = N. Pseudo-narcissus.
moschatum, Rafin. l. c. = N. Pseudo-narcissus.

MOSLA, Buch.-Ham. ex Benth. in Wall. Pl. As. Rar. i. 66, in syn. (1830). *LABIATAE*, Benth. & Hook. f. ii. 1182.
ORTHODON, Benth. & Oliver, in Journ. Linn. Soc. ix. (1865) 167.
chinensis, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 177.—China.
dianthera, Maxim. l. c. xx. (1865) 457 = ocymoides.
formosana, *Maxim. l. c.* 459.—Ins. Formosa.
grosseserrata, *Maxim. l. c.* 458.—Japon.
japonica, *Maxim. l. c.* 461.—Japon.
lanceolata, *Maxim, l. c.* 459.—Japon.
ocymoides, *Buch.-Ham. ex Benth. in Wall. Pl. As. Rar.* i. 66, in syn.—Ind. or.; China.
punctata, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 460.—Japon.

MOSTUEA, Didr. in Kjoeb. Vidensk. Meddel. (1853) 86. *LOGANIACEAE*, Benth. & Hook. f. ii. 789.
LEPTOCLADUS, Oliver, in Journ. Linn. Soc. viii. (1864) 160. t. 12.
Brunonis, *Didr. in Kjoeb. Vidensk. Meddel.* (1853) 87.—Afr. trop.
madagascarica, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 245.—Madag.
Pervilleana, *Baill. l. c.* 246.—Am. et Afr. trop.
surinamensis, *Benth. in Hook. Ic. Pl.* xii. 83, t. 1196.—Guian.
Thomsoni, *Benth. l. c.* sub t. 1196.—Afr. trop.

MOTANDRA, A. DC. in DC. Prod. viii. 423 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 715.
guineensis, *A. DC. l. c.*—Afr. trop.

MOTHERWELLIA, F. Muell. Fragm. vii. 107 (1870) = **Aralia**, Tourn.
haplosciadea, F. Muell. l. c. = A. Motherwelli.

MOUFFETTA, Neck. Elem. i. 124 (1790) = **Patrinia**, Juss. (Valer.).

MOUGEOTIA, H. B. & K. Nov. Gen. et Sp. v. 326. tt. 483, 484 (1821) = **Melochia**, Linn. (Stercul.).
caracasana, H. B. & K. l. c. 329 = Mel. caracasana.
corchorifolia, H. B. & K. l. c. 326, adnot. = Mel. corchorifolia.
hirsuta, H. B. & K. l. c. 331 = Mel. hirsuta, *Cav.*
inflata, H. B. & K. l. c. 330. t. 484 = Mel. lupulina.
mollis, H. B. & K. l. c. 328 = Mel. mollis.
nodiflora, H. B. & K. l. c. 330 = Mel. nodiflora.
polystachya, H. B. & K. l. c. 328. t. 483 = Mel. venosa.

MOULINSIA, Cambess. in Mém. Mus. Par. xviii. (1829) 27. t. 2 = **Erioglossum**, Blume (Sapind.).
cupanioides, Cambess. l. c. = E. edule.
rubiginosa, G. Don, Gen. Syst. i. 667 = E. edule.

MOULINSIA, Rafin. in Ser. Bull. Bot. i. (1830) 221 = **Aristida**, Linn. (Gramin.).
lanosa, Rafin. l. c. = A. lanosa.

MOULLAVA, Adans. Fam. ii. 318 (1763). *GEN. INCERT. SED.*
ALMELOVEENIA, Dennst. Schluess. Hort. Malab. 32 (1816).

MOURERA, Aubl. Pl. Gui. i. 582. t. 233 (1775). *PODOSTOMACEAE*, Benth. & Hook. f. iii. 109.
LACIS, Schreb. Gen. 366 (1789), partim.
STENGELIA, Neck. Elem. ii. 258 (1790).
alcicornis, Benth. ex Tul. in Arch. Mus. Par. vi. (1852) 94 = Ligea alcicornis.
aspera, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 93.—Bras.
fluviatilis, *Aubl. Pl. Gui.* i. 582. t. 233.—Am. trop.
Weddelliana, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 93.—Bras.

MOURICOU, Adans. Fam. ii. 326 (1763) = **Erythrina**, Linn. (Legumin.).

MOURIRI, Aubl. Pl. Gui. i. 452 (1775) = **Mouriria**, Juss. (Melastom.).
guianensis, Aubl. l. c. 583 = Mouriria guianensis.

MOURIRIA, Juss. Gen. 320 (1789). *MELASTOMACEAE*, Benth. & Hook. f. i. 772.
BOCKIA, Scop. Introd. 10 (1777).

MOURIRIA:—
GUILDINGIA, Hook. Bot. Misc. i. (1830) 122. t. 30.
MOURIRI, Aubl. Pl. Gui. i. 452 (1775).
MURIRI, J. F. Gmel. Syst. 693 (1791).
OLISBEA, DC. Prod. iii. 31 (1828).
PETALOMA, Sw. Prod. Veg. Ind. Occ. 4, 73 (1788); Fl. Ind. Occ. ii. 831 (1800).
abnormis, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 286.—Guian.
acuta, Griseb. Cat. Pl. Cub. 92 = parvifolia.
acutiflora, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 284.—Guian.
alternifolia, Roxb. ex Steud. Nom. ed. II. ii. 162 (Quid?).—Ind. or.
angustifolia, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 155.—Bras.
Apiranga, *Spruce, ex Triana, l. c.* 154.—Bras.
arborea, *Gardn. in Hook. Ic. Pl.* t. 515.—Bras.
brevipes, *Gardn. in Hook. Journ. Bot.* ii. (1840) 24.—Bras.
calva, Gardn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 153 = elliptica.
cauliflora, *DC. Prod.* iii. 7.—Bras.
crassifolia, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 336.—Guian.
domingensis, *Spach, Hist. Vég. Phan.* iv. 276.—Ind. occ.
elliptica, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 59.—Bras.
emarginata, *Griseb. Cat. Pl. Cub.* 92.—Cuba.
eugeniaefolia, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 154.—Bras.
Gardneri, *Triana, l. c.* 155.—Bras.
grandiflora, *DC. Prod.* iii. 8.—Bras.
graveolens, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 153 = vernicosa.
guianensis, *Poir. in Lam. Encyc.* ix. 334.—Guian.
guianensis, Miq. Stirp. Surin. Sel. 43 = acutiflora.
guianensis, Sagot, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 155 = Sagotiana.
lanceolata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 183.—Cuba.
macrophylla, *Willd. ex Cham. & Schlecht. in Linnaea*, x. (1835-36) 218.—Bras.
mexicana, [*Moç. & Sesse*], *ex DC. Prod.* iii. 8.—Mexic.
myrtifolia, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 154.—Bras.
myrtilloides, *Poir. in Dict. Sc. Nat.* xxxiii. 163.—Ind. occ.
parvifolia, *Benth. Bot. Voy. Sulph.* 97. t. 36.—N. Granat.
parvifolia, Gardn. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 155 = Gardneri.
pauciflora, Spruce, ex Triana, l. c. 154 = acutiflora.
polyantha, Miq. in Linnaea, xviii. (1844) 290 = guianensis.
Princeps, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 283.—Guian.
Pusa, *Gardn. in Hook. Journ. Bot.* ii. (1840) 23.—Bras.
rhizophoraefolia, *Gardn. in Hook. Ic. Pl.* sub t. 515; *Triana, in Trans. Linn. Soc.* xxviii. (1871) 153.—Ins. Martinic.
rivularis, Gardn. ex Triana, l. c. 154 = cauliflora.
Sagotiana, Triana, l. c. 155.—Guian.
Sideroxylon, *Sagot, ex Triana, l. c.*—Guian.
spathulata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 183.—Cuba.
subumbellata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 154.—Bras.
Valenzuelana, *A. Rich. Ess. Fl. Cub.* 575.—Cuba.
vernicosa, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 285.—Guian.
Weddellii, Naud. l. c. 286 = cauliflora.

MOUROUCOA, Aubl. Pl. Gui. i. 141. t. 54 (1775) = **Maripa**, Aubl. (Convolv.).
violacea, Aubl. l. c. 142 = Maripa passifloroides.

MOUSSONIA, Regel, in Flora, xxxi. (1848) 248 = **Isoloma**, Decne. (Gesnerac.).
costaricensis, Klotzsch, ex Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 107 = I. costaricense.
Deppeana, Klotzsch, ex Hanst. in Linnaea, xxxiv. (1865-66) 284 = I. Deppeanum.
elegans, Decne. ex Planch. in Fl. des Serres, v. (1849) t. 489 = I. elegans.
elongata, Regel, in Flora, xxxi. (1848) 248 = I. Deppeanum.

MOUSSONIA :—
formosa, Van Houtte, in Regel, Gartenfl. iii. (1854) 310. t. 101 = Isoloma elegans.
melastoma, Hanst. in Linnaea, xxxiv. (1865-66) 288.—Peruv.
papillosa, Oerst. ex Hanst. l. c.—Mexic.
purpurea, Hanst. l. c. 289 = I. purpureum.

MOUTABEA, Aubl. Pl. Gui. ii. 679. t. 274 (1775). *POLYGALEAE*, Benth. & Hook. f. i. 141.
ACOSTA, Ruiz & Pav. Prod. i. t. 1 (1794).
CRYPTOSTOMUM, Schreb. Gen. 137 (1789).
Acostae, Roem. & Schult. Syst. iv. 527 = aculeata.
aculeata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 62.—Am. trop.
dibotrya, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 14.—Bras.
excoriata, *Mart. ex Miq. l. c.* 16.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 680. t. 274.—Guian.
longifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 62. t. 186.—Bras.

MOUTAN, Reichb. in Moessl. Handb. ed. II. i. p. lx (1827) = **Paeonia**, Linn. (Ranunc.).
officinalis, Lindl. & Paxt. Flow. Gard. i. (1850–51) 162 = P. officinalis.

MOUTOUCHI, Aubl. Pl. Gui. ii. 748. t. 299 (1775) = **Pterocarpus**, Linn. (Legumin.).
crispata, Benth. in Ann. Wien. Mus. ii. (1838) 94 = P. Draco.
Draco, Benth. l. c. = P. Draco.
suberosa, Aubl. Pl. Gui. ii. 748 = P. Draco.

MOUTOUCHIA, Benth. in Ann. Wien. Mus. ii. (1838) 94 = praec.

MOYA, Griseb. in Goett. Abh. xix. (1874) 111. *CELASTRINEAE*.
ferox, *Griseb. l. c.* xxiv. (1879) 83.—Reg. Argent.
spinosa, *Griseb. l. c.* xix. (1874) 111.—Reg. Argent.
scutioides, *Griseb. l. c.* xxiv. (1879) 84.—Reg. Argent.

MOZAMBE, Rafin. Sylva Tellur. 112 (1838) = **Cadaba**, Forsk. (Cappar.).
levigata, Rafin. l. c. = C. indica.

MOZINNA, Orteg. Hort. Matr. Dec. 104. t. 13 (1797) = **Jatropha**, Linn. (Euphorb.).
canescens, Benth. Bot. Voy. Sulph. 52. t. 25 = J. canescens.
cardiophylla, Torr. in Bot. Mex. Bound. 198 = J. cardiophylla.
cinerea, Orteg. Hort. Matr. Dec. 108 = J. cinerea.
cordata, Orteg. l. c. 107 = J. cordata.
peltata, Steud. Nom. ed. II. ii. 163 = J. hernandiaefolia.
spathulata, Orteg. Hort. Matr. Dec. 105. t. 13 = J. spathulata.

MOZULA, Rafin. in Journ. Phys. xci. (1820) 71 = **Lythrum**, Linn.

MUCKIA, Hassk. Pl. Jav. Rar. 184 (1848) = MUKIA, Arn. = **Melothria**, Linn. (Cucurb.).

MUCRONEA, Benth. in Trans. Linn. Soc. xvii. (1836) 419. t. 20 = **Chorizanthe**, R. Br. (Polygon.).
californica, Benth. l. c. = C. californica.

MUCUNA, Adans. Fam. ii. 325 (1763). *LEGUMINOSAE*, Benth. & Hook. f. i. 533.
CACUVALLUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 399.
CARPOPOGON, Roxb. Hort. Beng. 54 (1814); Fl. Ind. iii. 283 (1832).
CITTA, Lour. Fl. Cochinch. 456 (1790).
HORNERA, Neck. Elem. iii. 43 (1790).
LABRADIA, Swediaur, Arzneymitt. i. 164 (1801).
MACRANTHUS, Poir. Encyc. Suppl. iii. 569 (1813).
MARCANTHUS, Lour. Fl. Cochinch. 460 (1790).
MACROCERATIDES, Raddi, in Mem. Mod. xviii. Fis. (1820) 392.
MACUNA, Marcgr. ex Scop. Introd. 309 (1777).
NEGRETIA, Ruiz & Pav. Prod. 98. t. 21 (1794).
PILLERA, Endl. Prod. Fl. Norf. 91 (1833).
STIZOLOBIUM, P. Br. Hist. Jamaic. 290 (1756).
ZOOPHTHALMUM, P. Br. l. c. 295. t. 31 (1756).
acuminata, *R. Grah. in Wall. Cat.* n. 5621.—Ind. or.

MUCUNA :—
Albertisi, *F. Muell. Papuan Pl.* i. 64.—N. Guin.
altissima, Boj. ex Benth. in Ann. Wien. Mus. ii. (1838) 133 = Dioclea violacea.
altissima, *DC. Prod.* ii. 405.—Ins. Martinic.
altissima, Hook. & Arn. Bot. Beech. Voy. 81; fide A. Gray, Bot. U. St. Expl. Exped. i. 446 = Strongylogon ruber.
anguina, Wall. Pl. As. Rar. iii. 19 = monosperma.
anguinea, Sweet, Hort. Brit. ed. I. 482 = praec.
atropurpurea, *DC. Prod.* ii. 406.—Ind. or.
Balfouriana, *A. Murr. in Trans. Bot. Soc. Edinb.* vi. (1860) 289.—Afr. trop.
Bennetti, *F. Muell. Papuan Pl.* i. 63.—N. Guin.
Bernieriana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 382.—Madag.
biplicata, *Teijsm. & Binn. Cat. Hort. Bog.* 261, nomen.—Borneo.
bracteata, *DC. Prod.* ii. 406.—Reg. Himal.; Burma.
capitata, *Sweet, Hort. Brit.* ed. I. 482; *Wight & Arn. Prod.* 255.—Reg. Himal.; Malaya.
Championi, *Benth. in Hook. Kew Journ.* iv. (1852) 49.—China.
comorensis, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 262.—Ins. Johan.
comosa, *DC. Prod.* ii. 405.—Guiana.
coriacea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 187.—Afr. trop.
corymbosa, R. Grah. in Wall. Cat. n. 5622 = monosperma.
cristata, Buch.-Ham. ex Wall. Cat. n. 5623 = monosperma.
cylindrosperma, Welw. ex Baker, in Oliver, Fl. Trop. Afr. ii. 186 = Physostigma cylindrospermum.
elliptica, *DC. Prod.* ii. 405.—Peruv.
flagellipes, *Vog. ex Benth. in Hook. Niger Fl.* 307.—Afr. trop.
gigantea, *DC. Prod.* ii. 405.—Ind. or.; Malaya.
gracilis, R. Grah. in Wall. Cat. n. 5626 = Cochlianthus gracilis.
hirsuta, *Wight & Arn. Prod.* 254.—Ind. or.
horrida, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 381.—Madag.
imbricata, *DC. Prod.* ii. 406.—Reg. Himal.
inflexa, *DC. l. c.* 405.—Peruv.
macrobotrys, *Hance, in Walp. Ann.* ii. 422.—China.
macrocarpa, *Wall. Cat.* n. 5618; et Pl. As. Rar. i. 41. t. 47.—Reg. Himal.
Macroceratides, *DC. Prod.* ii. 406.—Bras.
macrophylla, *Miq. Fl. Ind. Bat.* i. 1. 213.—Java.
melanocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 215.—Afr. trop.
mitis, *DC. Prod.* ii. 405.—Peruv.
mollis, *DC. l. c.* 405.—N. Granat.
mollissima, *Teijsm. & Binn. Cat. Hort. Bog.* 261, nomen.—Ins. Molucc.
monosperma, *DC. Prod.* ii. 406.—Ind. or.
monosperma, Wall. Cat. n. 5623 C, partim = imbricata.
Mutisiana, *DC. Prod.* ii. 406.—N. Granat.
nigra, Buch.-Ham. in Wall. Cat. n. 5617 = nivea.
nigricans, *Steud. Nom.* ed. II. ii. 163.—Cochinch.
nivea, *DC. Prod.* ii. 406.—Bengal; Burma.
nova-guineensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 18.—N. Guin.
paniculata, *Baker, in Journ. Linn. Soc.* xx. (1883) 132.—Madag.
platycarpa, *DC. Prod.* ii. 406.—Peruv.
platyphylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 448.—Ins. Pacif.
pruriens, *DC. Prod.* ii. 405.—Amphig. trop.
prurita, Hook. Bot. Mag. t. 4945 = imbricata.
prurita, Hook. Bot. Misc. ii. (1831) 348 = pruriens.
psittacina, *Miers, in Proc. Roy. Hort. Soc.* iv. (1864) 160.—N. Granat.
quadrialata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 186.—Afr. trop.
recta, Buch.-Ham. in Wall. Cat. n. 5625 = Vigna dolichoides.
rostrata, *Benth. in Mart. Fl. Bras.* xv. 1. 171.—Bras.
rutilans, [Moç. & Sesse], ex DC. Prod. ii. 404 = Canavalia obtusifolia.
spuria, *Steud. Nom.* ed. II. ii. 164.—Guiana.
stans, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 187.—Afr. trop.
umbellata, Salzm. ex Benth. in Mart. Fl. Bras. xv. 1. 169 = urens.
urens, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 399; *DC. Prod.* ii. 405.—Am. austr.

MUCUNA :—
utilis, Wall. ex Wight, Ic. ii. t. 80 = pruriens.
velutina, Hassk. in Flora, xxv. (1842) II. Beibl. 77 = capitata.
venenosa, *A. Murr. in Trans. Bot. Soc. Edinb.* vi. (1860) 289.—Afr. trop.
virgata, Desv. ex Steud. Nom. ed. II. ii. 164 = Dolichos virgatus.
volubilis, *Steud. l. c.*—Ind. or.

MUEHLENBECKIA, Meissn. Gen. 316; Comm. 227 (1840.) *POLYGONACEAE*, Benth. & Hook. f. iii. 101.
CALACINUM, Rafin. Fl. Tellur. ii. 33 (1836).
CONOBEA, Bert. ex Steud. Nom. ed. II. i. 404 (1840).
KARKINETRON, Rafin. Fl. Tellur. iii. 11 (1836).
SARCOGONUM, G. Don, in Sweet, Hort. Brit. ed. III. 577 (1839).
adpressa, *Meissn. Gen. Comm.* 227.—Austral.
australis, *Meissn. l. c.*—N. Zel.
axillaris, *Walp. Ann.* i. 552.—Austral.; N. Zel.
Benthamii, Endl. Gen. Suppl. iv. 51 = tamnifolia.
chilensis, *Meissn. in DC. Prod.* xiy. 148.—Chili.
complexa, *Meissn. Gen. Comm.* 227.—N. Zel.
Cunninghami, *F. Muell. Fragm.* v. 91.—Austral.
ephedroides, *Hook. f. Fl. N. Zel.* 211.—N. Zel.
flexuosa, Meissn. in Lehm. Pl. Preiss. i. 624 = adpressa.
florulenta, Meissn. in DC. Prod. xiv. 146 = Cunninghami.
gracillima, *Meissn. l. c.* 145.—Austral.
Gunnii, Walp. Ann. i. 552 = adpressa.
hastifolia, F. Muell. ex Meissn. in DC. Prod. xiv. 146 = adpressa.
leptobotrys, *Meissn. l. c.* 149.—Peruv.
parvifolia, Meissn. in Linnaea, xxvi. (1853–55) 362 = axillaris.
peruviana, *Meissn. in DC. Prod.* xiv. 150.—Peruv.
platyclados, *Meissn. in Bot. Zeit.* xxii. (1865) 313.—Ins. Salomon.
polybotrya, *Meissn. in Lehm. Pl. Preiss.* i. 623.—Austral.
polygonoides, *F. Muell. Fragm.* v. 73.—Austral.
quadrangulata, Endl. Gen. Suppl. iv. 51 = tamnifolia.
rhytocarya, *F. Muell. Fragm.* v. 92.—Austral.
rupestris, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 256.—Bolivia.
sagittaefolia, Remy, in C. Gay, Fl. Chil. v. 274 = chilensis.
sagittifolia, *Meissn. Gen. Comm.* 227.—Am. austr.
stenophylla, F. Muell. Fragm. i. 138 = polygonoides.
tamnifolia, *Meissn. Gen. Comm.* 227.—Mexic.
tiliaefolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 255.—Bolivia.
varians, *Meissn. in Bot. Zeit.* x. (1852) 347.—Hab.?
volcanica, Endl. Gen. Suppl. iv. 51 = seq.
vulcanica, *Meissn. in DC. Prod.* xiv. 148.—Ecuador.

MUEHLENBERGIA, Schreb. Gen. 44 (1789) (Muhlenbergia). *GRAMINEAE*, Benth. & Hook. f. iii. 1143.
CALYCODON, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 186.
CLOMENA, Beauv. Agrost. 28. t. 7. f. 10 (1812).
DACTYLOGRAMMA, Link, Hort. Berol. ii. 248 (1827).
DILEPYRUM, Michx. Fl. Bor. Am. i. 40 (1803).
DIPLACHYRIUM, Nees, in Flora. xi. (1828) 303.
PODOSAEMUM, Kunth, in Mém. Mus. Par. ii. (1815) 72.
TOSAGRIS, Beauv. Agrost. 29. t. 8. f. 3 (1812).
TRICHOCHLOA, Beauv. l. c. t. 8. f. 2 (1812).
VASEYA, Thurb. in Proc. Acad. Nat. Sc. Philad. (1863) 79.
acutifolia, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
adspersa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 291.—Peruv.
affinis, *Trin. l. c.* 301.—Mexic.
alpestris, *Trin. Gram. Unifl.* 194.—Sibir.
ambigua, *Torr. in Nicollet, Rep. Hydrog. Basin Missis.* (1843) 164.—Am. bor.
anemagrostoides, *Trin. ex Steud. Nom.* ed. II. ii. 164 (Quid?).—'Amer.'
angustata, *Kunth, Enum. Pl.* i. 202.—Peruv.
anomala, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 539.—Mexic.
arachnoides, Trin. ex Steud. Nom. ed. II. i. 365; ii. 164 = Cinna arundinacea.
arenicola, Buckl. in Prod. Acad. Sc. Philad. 1862 (1863) 91 = gracillima.
aristata, *Pers. Syn.* i. 73.—Mexic.

MUEHLENBERGIA:—
aspera, Trin. Gram. Unifl. 193 = Sporobolus asper.
aspericaulis, *Nees, ex Steud. Nom.* ed. II. ii. 164.—Hab.?
baicalensis, *Trin. ex Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 21.—Sibir. baical.
Berlandieri, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 299.—Mexic.
Berteroniana, *Kunth, Rév. Gram.* i. 64.—Ins. S. Doming.
Beyrichiana, Kunth, l. c. 63 = Perielema brasilianum.
Botteri, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 539.—Mexic.
Bourgaei, *Fourn. ex Hemsl. l. c.*—Mexic.
Brachyelytrum, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 306 = aristata.
brachyphylla, *Nees, in Linnaea*, xix. (1847) 690.—Mexic.
brasiliensis, *Steud. Syn. Pl. Gram.* 177.—Bras.
breviseta, *Griseb. ex Hemsl. Biol. Centr. Am. Bot.* iii. 539.—Mexic.
caespitosa, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 18.—Am. bor.
calamagrostidea, *Kunth, Rév. Gram.* i. 63.—Mexic.
calycina, Trin. Gram. Unifl. 193 = glomerata.
capillaris, *Trin. l. c.* 191.—Am. bor.; Mexic.
ciliata, *Trin. l. c.* 193.—Mexic.
Cinna, Trin. l. c. 191 = Cinna arundinacea.
ciurea, Benth. Fl. Austral. vii. 575, err. typ. = Dichelachne sciurea.
clandestina, Trin. l. c. 190 = Sporobolus asper.
Clomena, *Trin. l. c.* 194.—Mexic.; Peruv.
comata, *Benth. ex Vasey, in Rep. Dep. Agric.* (1881–82) 247. t. 5; *et ex Coult. Man. Rocky Mt. Bot.* 409.—Am. bor. occ.
composita, Trin. ex Steud. Nom. ed. II. i. 40; ii. 164 = Agrostis composita.
crinita, Trin. Gram. Unifl. 193 = Dichelachne crinita.
debilis, *Trin. l. c.*—Calif.; Mexic.
depauperata, *Scribn. in Coult. Bot. Gaz.* ix. (1884) 187.—Arizona.
diffusa, Willd. Sp. Pl. i. 320 = Schreberi.
distichophylla, *Kunth, Enum. Pl.* i. 202.—Mexic.
distichophylla, Munro, ex Vasey, Cat. Gram. U. St. 45, nomen = Epicampes distichophylla.
divaricata, Hort. ex DC. Cat. Hort. Monsp. 98 = Chaeturus fasciculatus.
diversiglumis, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 298 = Trinii.
dubia, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 540.—Mexic.
elegans, *Trin. Gram. Unifl.* 194.—Ecuador.
erecta, *J. & C. Presl, Rel. Haenk.* i. 231.—Mexic.
erecta, Schreb. Graes. ii. 139. t. 50 = aristata.
exilis, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 540.—Mexic.
expansa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 299.—Mexic.
fasciculata, Trin. Gram. Unifl. 192 = rigida.
filipes, *M. A. Curt. in Am. Journ. Sc.* Ser. I. xliv. (1843) 83 = capillaris.
foliosa, Trin. Gram. Unifl. 190 = mexicana.
geniculata, *Nees, ex Steud. Syn. Pl. Gram.* 178.—Reg. Himal.
glabrata, *Trin. Gram. Unifl.* 194.—Mexic.
glauca, *Nees, in Linnaea*, xix. (1847) 689.—Mexic.
glomerata, *Trin. Gram. Unifl.* 191. t. 5. f. 10.—Am. bor.
gracilis, Rupr. in Bull. Acad. Brux. ix. (1842) 229 = quadridentata.
gracilis, *Trin. Gram. Unifl.* 193.—Calif.; Mexic.
gracillima, *Torr. in Pacif. Rail. Rep.* iv. 155.—Am. bor. occ.
gymnostyla, *Nees, Del. Sem. Hort. Vratisl.* (1850) 3.—Hab.?
hirtula, *Steud. Syn. Pl. Gram.* 180.—Texas.
Huegelii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 293.—Reg. Himal.
implicata, *Trin. Gram. Unifl.* 193.—Mexic.
interrupta, Steud. Syn. Pl. Gram. 177 = Apera interrupta.
japonica, *Steud. l. c.* 422.—Japon.
lateriflora, Trin. Gram. Unifl. 193 = mexicana.
Liebmannii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 541.—Mexic.
longiseta, Benth. Pl. Hartw. 28 = calamagrostidea.
macrostachya, *Boj. Hort. Maurit.* 367.—Madag.
Mertensii, *Trin. ex Steud. Nom.* ed. II. ii. 164, nomen.—Hab.?
mexicana, *Trin. Gram. Unifl.* 189.—Am. bor.

MUEHLENBERGIA:—
microsperma, Trin. Gram. Unifl. 193 = debilis.
mollicoma, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 414 = Dichelachne crinita.
monticola, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 91 = sylvatica.
mucronata, *Trin. Gram. Unifl.* 194.—Mexic.
multiflora, Beauv. Agrost. 168 (Index) = Schreberi.
nana, Benth. Pl. Hartw. 262 = M. Clomena.
nardifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 294.—Reg. Argent.
pauciflora, *Buckl. in Proc. Acad. Sc. Philad.* 1862 (1863) 91.—Texas.
pendula, Bong. Veget. Sitch. 172 = Cinna arundinacea.
peruviana, Steud. Nom. ed. II. i. 41 = M. Clomena.
phragmitoides, *Griseb. in Goett. Abh.* xix. (1874) 255.—Reg. Argent.
Polypogon, Trin. Gram. Unifl. 193 = capillaris.
pungens, *Thurb. in Proc. Acad. Sc. Philad.* 1863 (1864) 78.—Am. bor. occ.
purpurea, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 186 = debilis.
pusilla, Steud. Syn. Pl. Gram. 177 = M. Clomena.
quadridentata, *Trin. Gram. Unifl.* 194.—Mexic.
rara, Trin. l. c. 193 = Agrostis rara.
rariflora, Hook. f. Fl. Antarct. 371. t. 131 = Stipa rariflora.
rigida, *Trin. Gram. Unifl.* 194.—Mexic.
rupestris, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
scabra, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 183 = Aristida scabra.
scabra, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 174.—Mexic.
Schaffneri, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 542.—Mexic.
Schreberi, *J. F. Gmel. Syst.* 171.—Am. bor.
sciurea, Trin. Gram. Unifl. 193 = Dichelachne sciurea.
setarioides, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 542.—Mexic.
setifolia, *Coult. Bot. Gaz.* vii. (1882) 92.—Am. bor.
setosa, Trin. Gram. Unifl. 193 = glomerata.
Sieberiana, Trin. ex Hook. f. Handb. N. Zeal. Fl. 326 = Dichelachne sciurea.
simplex, Kunth, Rév. Gram. i. 64 = Polypogon maritimus?
sobolifera, *Trin. Gram. Unifl.* 189.—Am. bor.
spicata, *Kunth, Rév. Gram.* i. 64.—Bras.
Spica-venti, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 285 = Apera Spica-venti.
spiciformis, *Trin. l. c.* 288.—Mexic.
Sprengelii, Trin. Gram. Unifl. 189 = tenuifolia.
stipoides, *Trin. l. c.* 194.—Mexic.; Ecuador.
stricta, Kunth, Enum. Pl. i. 202 = capillaris.
subalpina, *Vasey, Cat. Gram. U. St.* 40.—Am. bor.
subtilis, *Nees, in Linnaea*, xix. (1847) 689.—Mexic.
sylvatica, *Torr. & Gray, ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. ii. (1845) 292.—Am. bor.
tenella, Rupr. in Bull. Acad. Brux. ix. (1842) 230 = exilis.
tenella, *Trin. Gram. Unifl.* 193.—Mexic.
tenuifolia, *Trin. l. c.*—Mexic.
tenuis, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. (1842) 183 = Aristida tenuis.
tenuissima, *Kunth, Enum. Pl.* i. 198.—Am. centr.
texana, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 91 = Sporobolus ramulosus.
texana, *Thurb. ex Coult. Man. Rocky Mt. Bot.* 410.—Am. bor. occ.
trichodes, *Steud. Syn. Pl. Gram.* 177.—Am. austr.
trichopodes, Chapm. Fl. S. U. St. 553 = affinis.
Trinii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 543.—Mexic.
virescens, *Trin. Gram. Unifl.* 193.—Mexic.
viridissima, *Nees, ex Steud. Syn. Pl. Gram.* 178.—Indor.
viridissima, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. ii. (1845) 293 = sylvatica.
Willdenowii, *Trin. Gram. Unifl.* 188. t. 5. f. 3.—Am. bor.
Wrightii, *Vasey, ex Coult. Man. Rocky Mt. Bot.* 409.—Am. bor. occ.

MUELLERA, Linn. f. Suppl. 53 (1781). *LEGUMINOSAE*, Benth. & Hook. f. i. 550.
COUBLANDIA, Aubl. Pl. Gui. ii. 937. t. 356 (1775).
CYANOBOTRYS, Zucc. in Abh. Akad. Muench. iv. ii. (1845) 28 (Pl. Nov. Fasc. v. 30. t. 5).
MULLERA, Juss. Gen. 352 (1789).

MUELLERA:—
mexicana, *Benth. in Journ. Linn. Soc.* iv. (1860) *Suppl.* 117.—Mexic.
moniliformis, *Linn. f. Suppl.* 53.—Nicaragua.
verrucosa, Herb. Mus. Par. ex DC. Prod. ii. 353 = Bremontiera Ammoxylon.
verrucosa, Pers. Syn. ii. 311 = Ormocarpum verrucosum.

MUELLERARGIA, Cogn. in DC. Monog. Phan. iii. 630 (1881). *CUCURBITACEAE.*
timorensis, *Cogn. l. c.*—Ins. Timor.

MUENCHAUSIA, Scop. Introd. 220 (1777) = seq.

MUENCHHAUSIA, Murr. Syst. ed. XIII. 581 (1774) = MUNCHAUSIA, Linn. = **Lagerstroemia**, Linn. (Lythrar.).

MUENTERIA, Seem. Journ. Bot. iii. (1865) 329. tt. 35, 36 = **Dolichandrone**, Fenzl (Bignon.).
lutea, Seem. l. c. viii. (1870) 211 = D. lutea.
puberula, Seem. l. c. 339.—Afr. trop.
stenocarpa, Seem. l. c. iii. (1865) 329.—Afr. trop.
tomentosa, Seem. l. c. 330 = D. tomentosa.
zanzibarica, Seem. l. c. viii. (1870) 339.—Afr. trop.

MUENTERIA, Walp. Rep. v. 398 (1846) = **Picraena**, Lindl. (Simarub.).
excelsa, Walp. l. c. = P. excelsa.

MUHLENBERGIA, Schreb. Gen. 44 (1789) = **Muehlenbergia**, Schreb. (Gramin.).

MUILLA, S. Wats. in Proc. Am. Acad. xiv. (1879) 235. *LILIACEAE*, Benth. & Hook. f. iii. 801.
maritima, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 235.—Calif.

MUKIA, Arn. in Hook. Journ. Bot. iii. (1841) 276 = **Melothria**, Linn. (Cucurb.).
althaeoides, M. Roem. Syn. Pepon. 47 = Mel. maderaspatana.
leiosperma, Wight, in Ann. & Mag. Nat. Hist. Ser. I. viii. (1842) 268 = Mel. leiosperma.
maderaspatana, M. Roem. Syn. Pepon. 47 = Mel. maderaspatana.
micrantha, F. Muell. Fragm. ii. 180 = Mel. Muelleri.
Rottleri, M. Roem. Syn. Pepon. 47 = Mel. maderaspatana.
scabrella, Arn. in Hook. Journ. Bot. iii. (1841) 276 = Mel. maderaspatana.

MULDERA, Miq. in Bull. Néerl. (1839) 448; et Comm. Phyt. 34 (1840) = **Piper**, Linn.
baccata, Miq. Syst. Pip. 341 = P. baccatum.
cordata, Miq. Pl. Jungh. 17 = P. ungaramense.
diandra, Thw. Enum. Pl. Zeyl. 428 = P. trineuron.
firma, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 140 = P. firmum.
galeata, Miq. in Hook. Lond. Journ. Bot. v. (1846) 557 = P. galeatum.
multinervis, Miq. l. c. 558 = P. nigrum.
recurva, Miq. Syst. Pip. 342 = P. recurvum.
recurva, Zoll. Syst. Verz. Arch. Ind. 84 = P. baccatum.
trichostachya, Miq. in Hook. Lond. Journ. Bot. v. (1846) 556 = P. trichostachyon.
Wightiana, Miq. l. c. 557 = P. galeatum, nigrum.

MULGEDIUM, Cass. in Dict. Sc. Nat. xxxiii. 296 (1824) = **Lactuca**, Tourn. (Compos.).
abietinum, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 802.—As. Min.
acuminatum, DC. Prod. vii. 250 = L. acuminata.
albanum, DC. l. c. 249 = L. racemosa.
alpinum, Less. Syn. Comp. 142 = L. alpina.
armeniacum, Boiss. Fl. Orient. iii. 800 = L. racemosa.
aureum, Sch. Bip. ex Vis. in Mem. Ist. Venet. xv. (1870) 7 = L. sonchifolia.
Aurora, Sch. Bip. ex Nym. Consp. 433 = L. tatarica.
azureum, DC. Prod. vii. 248.—Sibir. altaic.
Bourgaei, Boiss. ex Boiss. Fl. Orient. iii. 801.—As. Min.
cacaliaefolium, DC. Prod. vii. 250.—Reg. Caucas.
crassicaule, Trautv. in Bull. Soc. Nat. Mosc. xxxix. (1866) i. 390.—As. centr.
cyaneum, DC. Prod. vii. 249 = L. hastata.
decipiens, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 266 = L. decipiens.

MULGEDIUM :—
divaricatum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 442.—Am. bor.
djimilense, Boiss. Fl. Orient. iii. 800 = seq.
dshimilense, C. Koch, in Linnaea, xxiii. (1850) 669 = Lactuca racemosa.
dubium, C. Koch, l. c. 670.—As. Min.
fissifolium, Steud. Nom. ed. II. ii. 165.—Sibir.
floridanum, DC. Prod. vii. 249 = L. floridana.
glabrum, C. Koch, in Linnaea, xxiii. (1850) 668 = L. macrophylla.
grande, C. Koch, l. c. xvii. (1843) 279 = L. macrophylla.
hastatum, Less. Syn. Comp. 142 = Prenanthes alata.
heterophyllum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 441 = L. pulchella.
hispidum, DC. Prod. vii. 250.—Sibiria.
integrifolium, Cass. in Dict. Sc. Nat. xxxiii. 298 = L. canadensis.
kamtschaticum, Ledeb. in Denkschr. Bot. Ges. Regensb. iii. (1841) 65.—Kamtschat.
laevigatum, DC. Prod. vii. 249 = L. macrorrhiza.
lazicum, Boiss. & Bal. in Boiss. Fl. Orient. iii. 801 = *cacaliaefolium*.
Lessertianum, DC. Prod. vii. 251 = L. Lessertiana.
leucophaeum, DC. l. c. 250 = alpina.
lyratum, Cass. in Dict. Sc. Nat. xxxiii. 297 = L. floridana.
macranthum, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 267 = L. macrantha.
macrophyllum, DC. Prod. vii. 248 = L. macrophylla.
macrorhizum, Royle, Illustr. Bot. Himal. t. 61. f. 1 = L. macrorhiza.
multiflorum, DC. Prod. vii. 249 = L. alpina.
neilgherrense, Wight, Ic. t. 1144 = L. hastata.
Pančičii, Vis. in Mem. Inst. Venet. ix. (1860) 173.—Europ. austr.
petiolatum, C Koch, in Linnaea, xvii. (1843) 279 = *cacaliaefolium*.
Plumieri, DC. Prod. vii. 248 = L. Plumieri.
ponticum, Boiss. in Ann. Sc. Nat. Sér. II. iv. (1854) 248. t. 2 = *cacaliaefolium*.
prenanthoides, DC. Prod. vii. 248.—Reg. Caucas.
pulchellum, G. Don, in Sweet, Hort. Brit. ed. III. 418 = L. pulchella.
rapunculoides, DC. Prod. vii. 249 = L. rapunculoides.
robustum, DC. l. c. 250 = L. hastata.
rostratum, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 125 = L. graciliflora.
runcinatum, Cass. in Dict. Sc. Nat. xxxiii. 296 = L. tatarica.
sagittatum, Royle, Illustr. Bot. Himal. 252. t. 61. f. 2 = L. longifolia.
salicifolium, C. Koch, in Linnaea, xxiii. (1850) 669.—Armenia.
sibiricum, Less. Syn. Comp. 142 = L. sibirica.
sonchifolium, Vis. & Panč. in Mem. Ist. Venet. xv. (1870) 5 = L. tatarica.
tataricum, DC. Prod. vii. 248 = L. tatarica.
thianschanicum, Regel & Schmalh. in Act. Hort. Petrop. vi. (1879) 329.—As. centr.

MULINUM, Pers. Syn. i. 309 (1805). *UMBELLIFERAE*, Benth. & Hook. f. i. 877.
acaule, Pers. Syn. i. 309 = Laretia acaulis.
albovaginatum, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 328.—Chili.
angulatum, DC. Prod. iv. 80 (= *Pozoa coriacea*).—Chili.
axilliflorum, *Griseb. in Goett. Abh.* xix. (1874) 154.—Reg. Argent.
crassifolium, *Phil. Fl. Atac.* 24.—Chili.
cryptanthum, *Clos, in C. Gay, Fl. Chil.* iii. 90.—Chili.
cuneatum, *Hook. & Arn. Bot. Beech. Voy.* 26.—Chili.
Dipterygia, DC. Prod. iv. 80 = Asteriscium polycephalum.
Echegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 27.—Reg. Argent.
Echinus, DC. Prod. iv. 79 = ulicinum.
hirsutum, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 346 = Bowlesia multiradiata.
integrifolium, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 28.—Reg. Argent.
isatidicarpum, DC. Prod. iv. 80 = Asteriscium isatidicarpum.
laxum, *Phil. in Linnaea*, xxxiii. (1864-65) 90.—Chili.

MULINUM:—
leoninum, *Lor. in Roca, Exped. Rio-Negro*, ii. 222.—Patagon.
microphyllum, *Pers. Syn.* i. 309.—Am. austr.
proliferum, *Pers. l. c.*—Am. austr.
saniculaefolium, Spreng. Umbell. Prod. 15 = Diposis saniculaefolia.
spinosum, *Pers. Syn.* i. 309.—Chili.
triacanthum, *Griseb. in Goett. Abh.* xix. (1874) 154.—Reg. Argent.
ulicinum, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 328. t. 64.—Chili.

MULLERA, Juss. Gen. 352 (1789) = **Muellera**, Linn. f. (Legumin.).

MUNBYA, Boiss. Diagn. Ser. I. xi. 114 (1849 = **Macrotomia**, DC. (Borag.).
Cephalotes, Boiss. l. c. 116 = Macrotomia Cephalotes.
conglobata, Boiss. l. c. = Macrotomia Cephalotes.
cyanochroa, Boiss. l. c. 115 = Macrotomia perennis.
densiflora, Boiss. l. c. 116.—Songaria.
perennis, Boiss. l. c. 115 = Macrotomia perennis.

MUNBYA, Pomel, Mat. Fl. Atl. 8 (1860). *LEGUMINOSAE*.
LAMOTTEA, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.
plicata, *Pomel, Mat. Fl. Atl.* 9.—Alger.
polystachya, *Pomel, l. c.*—Alger.

MUNCHAUSIA, Linn. Mant. ii. 153 (1771) = **Lagerstroemia**, Linn. (Lythrar.).
ovata, Jaume St. Hil. Expos. ii. 176 = L. hirsuta.
speciosa, Linn. Mant. ii. 243; et ex Muench. Hausv. i. 357 = L. Flos-reginae.

MUNCHHAUSIA, Murr. Prod. Gotting. Praef. (1770) = praec.

MUNCHUSIA, Heist. ex Rafin. Sylva Tellur. 115 (1838) = **Hibiscus**, Linn. (Malvac.).
tomentosa, Rafin. l. c. = H. clypeatus.

MUNDTIA, H. B. & K. Nov. Gen. et Sp. v. 392, in adnot. (1821) (Mundia); Harv. in Harv. & Sond. Fl. Cap. i. 95 (1859-60). *POLYGALEAE*, Benth. & Hook. f. i. 137.
NYLANDTIA, Dum. Comm. Bot. 31 (1822).
VASCOA, DC. Prod. i. 337, in obs. (1824).
albiflora, Eckl. & Zeyh. Enum. 29 = spinosa.
brasiliensis, A. St. Hil. Fl. Bras. Mer. ii. 58. t. 92 = Acanthocladus brasiliensis.
Candolleana, Walp. Rep. ii. 769 = Muraltia De-Candollei.
desertorum, Eckl. & Zeyh. Enum. 30 = Muraltia juniperifolia.
glauca, Eckl. & Zeyh. l. c. = spinosa.
heterophylla, Walp. Rep. ii. 769 = Muraltia ononidifolia.
Kraussiana, Walp. l. c. = Muraltia Kraussiana.
montana, Eckl. & Zeyh. Enum. 30 = spinosa.
scoparia, Eckl. & Zeyh. l. c. = spinosa.
spinosa, *DC. Prod.* i. 338.—Afr. austr.
tabularis, Eckl. & Zeyh. Enum. 29 = spinosa.

MUNDUBI, Adans. Fam. ii. 323 (1763) = **Arachis**, Linn. (Legumin.).

MUNDULEA, Benth. in Miq. Pl. Jungh. ii. 248 (1852). *LEGUMINOSAE*, Benth. & Hook. f. i. 497.
ambatoana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 390.—Madag.
ambongoënsis, *Baill. l. c.* 391.—Madag.
antanossarum, *Baill. l. c.* 390.—Madag.
densicoma, *Baill. l. c.* 391.—Madag.
Grandidieri, *Baill. l. c.* 390.—Madag.
Hookeri, *Baill. l. c.* 389.—Madag.
Millettii, *Benth. in Miq. Pl. Jungh.* 249.—China.
pauciflora, *Baker, in Journ. Bot.* xx. (1882) 68.—Madag.
pulchra, Benth. in Miq. Pl. Jungh. 248 = Millettia pulchra.
revoluta, *Baker, in Journ. Linn. Soc.* xx. (1883) 129.—Madag.
Richardiana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 389.—Madag.

MUNDULEA:—
suberosa, *Benth. in Miq. Pl. Jungh.* 248.—Afr. trop.
Telfairii, *Baker, in Journ. Bot.* xx. (1882) 69.—Madag.
tullearensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 390.—Madag.

MUNGOS, Adans. Fam. ii. 225 (1763) = **Ophiorrhiza**, Linn. (Rubiac.).

MUNNICKIA, Reichb. Consp. 85 (1828) = **Bragantia**, Lour. (Aristol.).

MUNNICKSIA, Dennst. Schluess. Hort. Malab. 27 (1818) = **Hydnocarpus**, Gaertn. (Bixin.).
laurifolia, Dennst. l. c. = H. Wightiana.

MUNNOZIA, Ruiz & Pav. Prod. Fl. Per. 108. t. 23 (1794) = **Liabum**, Adans. (Compos.).
corymbosa, Ruiz & Pav. Syst. Veg. 195 = L. corymbosum.
lanceolata, Ruiz & Pav. l. c. 196 = L. lanceolatum.
sagittata, Wedd. Chlor. And. i. 211 = L. sagittatum.
Senecionidis, Benth. Pl. Hartw. 134.—Ecuador.
trinervis, Ruiz & Pav. Syst. Veg. 195 = L. trinerve.
venosissima, Ruiz & Pav. l. c. = L. venosissimum.

MUNROA, Torr. in Pacif. Rail. Rep. iv. 158 (1856) (Monroa). *GRAMINEAE*, Benth. & Hook. f. iii. 1180.
MONROA, Torr. l. c. (1856), errore, et auct. plur.
argentina, *Griseb. in Goett. Abh.* xxiv. (1879) 300.—Reg. Argent.
mendocina, *Phil. Sert. Mend. Alt.* 52.—Chili.
squarrosa, *Torr. in Pacif. Rail. Rep.* iv. 158.—N. Mexic.

MUNRONIA, Wight, Illustr. i. 147. t. 54 (1839). *MELIACEAE*, Benth. & Hook. f. i. 331.
javanica, *Benn. Pl. Jav. Rar.* i. 176.—Java.
neilgherrica, Wight, Illustr. i. 147. t. 54 = Wallichii.
pumila, *Wight, Ic.* i. t. 91.—Ind. or.
timoriensis, *Baill. Adansonia*, xi. (1874) 266.—Ins. Timor.
Wallichii, *Wight, Illustr.* i. 117.—Ind. or.

MUNROZIA, Steud. Nom. ed. II. ii. 166 (1841), sphalm. = MUNNOZIA, Ruiz & Pav. = **Liabum**, Adans. (Compos.).

MUNTINGIA, Plum. ex Linn. Syst. ed. I (1735). *TILIACEAE*, Benth. & Hook. f. i. 236.
Calabura, *Linn. Sp. Pl.* 509.—Ind. occ.
glabra, *Spreng. Syst.* ii. 592.—Bras.
rosea, *Karst. Fl. Columb.* ii. 55. t. 128.—N. Granat.

MUNYCHIA, Cass. in Dict. Sc. Nat. xxxvii. 483 (1825) = **Felicia**, Cass. (Compos.).
brachyglossa, Cass. l. c. = Aster Cymbalariae.
Cymbalariae, Nees, Gen. & Sp. Ast. 218 = Aster Cymbalariae.
cymbalarioides, DC. Prod. v. 223 = Aster Candollii.
hirsuta, DC. l. c. 222 = Aster hirsutus.

MURALTA, Adans. Fam. ii. 460 (1763) = **Clematis**, Tourn. (Ranunc.).

MURALTIA, Neck. Elem. iii. 62 (1790). *POLYGALEAE*, Benth. & Hook. f. i. 137.
HEISTERIA, Linn. Gen. ed. I. 350 (1737), nomen prius.
acerosa, *Harv. in Harv. & Sond. Fl. Cap.* i. *Add.* 20*.—Afr. austr.
acicularis, *Harv. l. c.* 102.—Afr. austr.
acipetala, *Harv. l. c.* 101.—Afr. austr.
affinis, Harv. l. c. 102 = longicuspis.
alopecuroides, *DC. Prod.* i. 335.—Afr. austr.
alopecuroides, Eckl. & Zeyh. Enum. 26 = squarrosa.
angulosa, *Spreng. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 356.—Afr. austr.
anthospermifolia, *Eckl. & Zeyh. Enum.* 26.—Afr. austr.
aspalatha, DC. Prod. i. 336 = phylicoides.
asparagifolia, *Eckl. & Zeyh. Enum.* 28.—Afr. austr.
aspera, Lehm. ex Eckl. & Zeyh. l. c. 25 = thymifolia.

MURALTIA:—
Beiliana, *Harv. in Harv. & Sond. Fl. Cap.* i. 111.—Afr. austr.
brevicornu, *DC. Prod.* i. 337.—Afr. austr.
brevicornu, E. Mey. in Drège, Zwei Pfl. Docum. 203 = striata.
brevifolia, DC. Prod. i. 335 = thymifolia.
brevifolia, Eckl. & Zeyh. Enum. 29 = obovata.
Burchellii, Eckl. & Zeyh. l. c. 27 = alopecuroides.
Burmanni, *DC. Prod.* i. 337.—Afr. austr.
Burmanni, Eckl. & Zeyh. Enum. 25 = satureioides.
calycina, *Harv. in Harv. & Sond. Fl. Cap.* i. 101.—Afr. austr.
Candolleana, Meissn. in Hook. Lond. Journ. Bot. i. (1842) 471 = De-Candollei.
Candollei, Harv. in Harv. & Sond. Fl. Cap. i. 10 = De-Candollei.
carnosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 203, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 111.—Afr. austr.
ciliaris, *DC. Prod.* i. 336.—Afr. austr.
ciliaris, Eckl. & Zeyh. Enum. 27 = incompta.
ciliais, Sieber, ex Harv. & Sond. Fl. Cap. i. 106 = phylicoides.
cliffortiaefolia, *Eckl. & Zeyh. Enum.* 25.—Afr. austr.
condensata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 354.—Afr. austr.
conferta, DC. Prod. i. 335 = satureioides.
conferta, Hochst. ex Harv. & Sond. Fl. Cap. i. 108 = lancifolia.
conferta, E. Mey. in Drège, Zwei Pfl. Docum. 203 = M. Heisteria.
crassifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 112.—Afr. austr.
De-Candollei, *Eckl. & Zeyh. Enum.* 27.—Afr. austr.
depressa, *Burch. ex DC. Prod.* i. 336.—Afr. austr.
depressa, E. Mey. in Drège, Zwei Pfl. Docum. 203 = mixta.
diffusa, Burch. ex DC. Prod. i. 336 = stipulacea.
divaricata, *Eckl. & Zeyh. Enum.* 28.—Afr. austr.
Dregei, *Harv. in Harv. & Sond. Fl. Cap.* i. 108.—Afr. austr.
Dregei, Presl, Bot. Bemerk. 17 = squarrosa.
dumosa, *DC. Prod.* i. 337.—Afr. austr.
ericaefolia, *DC. l. c.* 336.—Afr. austr.
ericaefolia, Eckl. & Zeyh. Enum. 28 = divaricata.
ericaefolia, Pappe, ex Harv. & Sond. Fl. Cap. i. 101 = satureioides.
ericifolia, E. Mey. in Drège, Zwei Pfl. Docum. 203 = satureioides.
fasciculata, *DC. Prod.* i. 337.—Afr. austr.
fasciculata, Eckl. & Zeyh. Enum. 27 = ciliaris.
filiformis, *DC. Prod.* i. 337.—Afr. austr.
hamata, E. Mey. in Drège, Zwei Pfl. Docum. 203 = macrocarpa.
Heisteria, *DC. Prod.* i. 335.—Afr. austr.
heterophylla, E. Mey. in Drège, Zwei Pfl. Docum. 203 = ononidifolia.
humilis, *DC. Prod.* i. 337.—Afr. austr.
incompta, *E. Mey. in Drège, Zwei Pfl. Docum.* 203, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 107.—Afr. austr.
juniperifolia, *DC. Prod.* i. 336.—Afr. austr.
juniperifolia, Eckl. & Zeyh. Enum. 26 = ericaefolia.
Kraussiana, *Meissn. in Hook. Lond. Journ. Bot.* i. (1842) 473.—Afr. austr.
lancifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 108.—Afr. austr.
laricifolia, *Eckl. & Zeyh. Enum.* 27.—Afr. austr.
laxa, DC. Prod. i. 337.—Afr. austr.
laxa, Eckl. & Zeyh. Enum. 29 = divaricata.
laxa, E. Mey. in Drège, Zwei Pfl. Docum. 203 = Dregei.
leptorhiza, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 356.—Afr. austr.
linifolia, Harv. in Harv. & Sond. Fl. Cap. i. 111 = angulosa.
linophylla, Burch. ex DC. Prod. i. 336 = filiformis.
longicuspis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 355.—Afr. austr.
lycopodioides, *Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 182.—Afr. austr.
macrocarpa, *Lehm. ex Eckl. & Zeyh. Enum.* 26.—Afr. austr.
macroceras, *Burch. ex DC. Prod.* i. 336.—Afr. austr.
macroceratia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = praec.
macropetala, *Harv. in Harv. & Sond. Fl. Cap.* i. 100.—Afr. austr.

MURALTIA:—
marifolia, Eckl. & Zeyh. Enum. 28 = serpylloides.
metalasiaefolia, Eckl. & Zeyh. l. c. 26 = rubeacea.
micrantha, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* ii. (1834) 122.—Afr. austr.
micrantha, Eckl. & Zeyh. Enum. 28 = stipulacea.
mixta, *DC. Prod.* i. 336.—Afr. austr.
mixta, E. Mey. in Drège, Zwei Pfl. Docum. 203, partim = longicuspis.
obovata, *DC. Prod.* i. 337.—Afr. austr.
obovata, Eckl. & Zeyh. Enum. 29 = origanoides.
ononidifolia, *Eckl. & Zeyh. l. c.* 25.—Afr. austr.
origanoides, *Presl, Bot. Bemerk.* 16.—Afr. austr.
Pappeana, *Harv. in Harv. & Sond. Fl. Cap.* i. 106.—Afr. austr.
parviflora, *DC. Prod.* i. 337.—Am.
pauciflora, *DC. l. c.*—Afr. austr.
phylicoides, *DC. l. c.*—Afr. austr.
pilosa, *DC. l. c.*—Afr. austr.
Poiretii, *DC. l. c.*—Afr. austr.
pubescens, *DC. l. c.* 336.—Afr. austr.
reticulata, Harv. in Harv. & Sond. Fl. Cap. i. 109 = origanoides.
rigida, *E. Mey. in Drège, Zwei Pfl. Docum.* 203, nomen; *et ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 357.—Afr. austr.
rubeacea, *Eckl. & Zeyh. Enum.* 26.—Afr. austr.
rubicunda, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 354 = dumosa.
rupestris, Pappe, ex Harv. & Sond. Fl. Cap. i. 107 = pubescens.
ruscifolia, Eckl. & Zeyh. Enum. 25 = squarrosa.
satureioides, *Burch. ex DC. Prod.* i. 336.—Afr. austr.
serpyllifolia, G. Don, Gen. Syst. i. 364 = seq.
serpylloides, *DC. Prod.* i. 335.—Afr. austr.
serpylloides, Eckl. & Zeyh. Enum. 25 = thymifolia.
sprengelioides, Burch. ex DC. Prod. i. 336 = juniperifolia.
sprengelioides, Eckl. & Zeyh. Enum. 26 = ericaefolia.
sprengelioides, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 357 = Beiliana.
squarrosa, *DC. Prod.* i. 335.—Afr. austr.
squarrosa, Eckl. & Zeyh. Enum. 27 = alopecuroides.
squarrosa, Sieber, ex Presl, Bot. Bemerk. 16 = stipulacea.
stipulacea, Andr. Bot. Rep. t. 363 = macropetala.
stipulacea, *Burch. ex DC. Prod.* i. 336.—Afr. austr.
stipulacea, Eckl. & Zeyh. Enum. 25 = dumosa.
striata, *DC. Prod.* i. 337.—Afr. austr.
striata, Eckl. & Zeyh. Enum. 29 = angulosa.
sulcata, Harv. in Harv. & Sond. Fl. Cap. i. 113 = leptorhiza.
tenuifolia, *DC. Prod.* i. 336.—Afr. austr.
tenuifolia, Meissn. in Hook. Lond. Journ. Bot. i. (1842) 474 = lancifolia.
Thunbergii, Eckl. & Zeyh. Enum. 26 = phylicoides.
thymifolia, *DC. Prod.* i. 337.—Afr. austr.
trinervia, *DC. l. c.* 335.—Afr. austr.
viminea, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 356 = striata.
virgata, Burch. ex DC. Prod. i. 336 = filiformis.
virgata, Eckl. & Zeyh. Enum. 27 = alopecuroides.
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 353 (err. typ. Zeygeri).—Afr. austr.

MURDANNIA, Royle, Illustr. Bot. Himal. 403. t. 95 (1839) = **Aneilema**, R. Br. (Commel.).
scapiflora, Royle, l. c. = A. scapiflorum.
tuberosa, C. B. Clarke, in DC. Monog. Phan. iii. 201, sphalm. = praec.

MURERA, Jaume St. Hil. Expos. ii. 376 (1805) = **Mourera**, Aubl. (Podostem.).

MURETIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 143. *UMBELLIFERAE*, Benth. & Hook. f. i. 895.
amplifolia, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 859.—Persia.
aurea, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1884) 143.—As. Min.
lutea, Boiss. Fl. Orient. ii. 858 = tanaicensis.
tanaicensis, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 143.—Reg. Casp.

MURICARIA, Desv. Journ. Bot. iii. (1814) 159. t. 25. f. 2. *CRUCIFERAE*, Benth. & Hook. f. i. 99.
CORVINA, Steud. Nom. ed. II. i. 256 (1840) sphalm.
KREMERIA, Coss. & Dur. in Bull. Soc. Bot. Fr. iii. (1856) 671.

MURICARIA:—
corydylocarpa, *C. Muell. in Walp. Ann.* vii. 177.—Alger.
prostrata, *Desv. Journ. Bot.* iii. (1814) 159. t. 25. f. 2.—Alger.

MURICIA, Lour. Fl. Cochinch. 596 (1790) = **Momordica**, Linn. (Cucurb.).
cochinchinensis, Lour. l. c. = Momord. cochinchinensis.

MURIEA, Hartog, in Journ. Bot. xvi. (1878) 145 = **Labourdonnaisia**, Boj. (Sapot.).
albescens, Hartog. l. c. 145.—Cf. 72 = L. albescens.
discolor, Hartog. l. c. 145.—Cf. 72 = L. discolor.

MURIRI, J. F. Gmel. Syst. 693 (1791) = **Mouriria**, Juss. (Melastom.).

MUROCOA, Jaume St. Hil. Expos. i. 307 (1805) = MOUROUCOA, Aubl. = **Maripa**, Sw. (Ternstr.).

MURRAEA, Koen. ex Linn. Mant. ii. 554 (1771) = seq.

MURRAYA, Koen. ex Linn. Mant. ii. 554 (1771) (Murraea); Murr. Syst. ed. XIII. 331 (1774). *RUTACEAE*, Benth. & Hook. f. i. 304.
BERGERA, Koen. in Linn. Mant. ii. 555 (1771).
CHALCAS, Linn. Mant. i. 68 (1767).
MARSANA, Sonner. Voy. Ind. Or. iii. 282 (1782).
NIMBO, Dennst. Schluess. Hort. Malab. 30 (1818).
SICKLERA, M. Roem. Syn. Pepon. i. 49 (1846).
amoena, Salisb. Prod. 320 = exotica.
brevifolia, Thw. Enum. Pl. Zeyl. 45 = exotica.
Burmanni, Spreng. Syst. ii. 315 = Clausena excavata.
cerasiformis, Blanco, Fl. Filip. ed. I. 363 = Glycosmis pentaphylla.
crenulata, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 29.—Ins. Philipp.; Austral.
elongata, *A. DC. ex Hook. f. Fl. Brit. Ind.* i. 503.—Burma.
exotica, Blanco, Fl. Filip. ed. II. 255 = Glycosmis pentaphylla.
exotica, *Linn. Mant.* ii. 563.—As. et Austral. trop.; Polynes.
exotica, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 48 = Aglaia odorata.
foetidissima, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 41.—Java.
Glenieii, Thw. ex Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 29 = exotica.
heptaphylla, *Span. in Linnaea*, xv. (1841) 178.—Ins. Timor.
japonensis, Raeusch. Nom. ed. III. 119 = exotica.
Koenigii, *Spreng. Syst.* ii. 315.—Ind. or.
lobata, *Blanco, Fl. Filip.* ed. I. 363.—Ins. Philipp.
longifolia, *Blume, Bijdr.* 137.—Java.
odorata, Blanco, Fl. Filip. ed. II. 256 = Feronia elephantum.
paniculata, Jack, in Malay. Misc. i. (1820) n. v. 31 = exotica.
sumatrana, Roxb. Hort. Beng. 32; Fl. Ind. ii. 375 = exotica.

MURRINEA, Rafin. Sylva Tellur. 104 (1838) = **Baeckea**, Linn. (Myrtac.).

MURRITHIA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 576 = **Pimpinella**, Riv. (Umbell.).
cordata, Zoll. & Mor. l. c.—Java.

MURRYA, Griff. Notul. iv. 494 (1854), sphalm. = **Murraya**, Linn. (Rutac.).

MURTEKIAS, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
Myrsinites, Rafin. l. c. = E. Myrsinites.

MURUCOA, J. F. Gmel. Syst. 344 (1791) = MOUROUCOA, Aubl. = **Maripa**, Sw. (Ternstr.).

MURUCUJA, Tourn. ex Medic. Malv. 97 (1787) = **Passiflora**, Linn.
adiantifolia, Sweet, Hort. Brit. ed. I. 355 = P. glabra.
aurantia, Pers. Syn. ii. 222 = P. aurantia.
Baueri, Lindl. Coll. Bot. t. 36 = P. Baueriana.

MURUCUJA :—
Herbertiana, Sweet, Hort. Brit. ed. I. 355 = Passiflora Herbertiana.
lunata, Medic. Malv. 97 = P. lunata, *Willd.*
mollissima, Spreng. Syst. iii. 43 = Tacsonia mollissima.
ocellata, Pers. Syn. ii. 222 = P. Murucuja.
orbiculata, Pers. l. c. = P. Murucuja.
pedunculariis, Spreng. Syst. iii. 43 = Tacsonia pedunculariis.
perfoliata, Spreng. l. c. = P. perfoliata.
speciosa, Spreng. l. c. = Tacsonia mixta.
viridiflora, Spreng. l. c. = P. viridiflora.

MUSA, Linn. Mus. Cliff. 2 (1736); Gen. ed. I. 315 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 655.
ENSETE, Bruce, 'Trav. v. App. 36' ex Horan. Prod. Scit. 40 (1862).
KARKANDELA, Rafin. Sylva Tellur. 106 (1838).
MNASIUM, Stackh. Extr. Bruce's Trav. Abyss. 18. t. 2 (1815).
Abaca, Perr. in Mém. Soc. Linn. Par. iii. (1825) 130 = textilis ?
acuminata, *Colla, Mem. Gen. Musa*, 66.—Ins. Amboin.
alphurica, Miq. Fl. Ind. Bat. iii. 589 = sapientum.
amboinensis, Miq. l. c. 588 = textilis.
Bacoba, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 302 (Quid ?).—Guiana.
Balbisiana, Colla, Mem. Gen. Musa, 56 = sapientum.
Banksiana, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 164 = seq.
Banksii, *F. Muell. Fragm.* iv. 132.—Austral.
Basjoo, *Siebold, in Verh. Batav. Gen.* xii. (1830) 18. —Japon.
Berteri, Colla, Mem. Gen. Musa, 57 = sapientum.
Berteroniana, Steud. Nom. ed. II. ii. 167, sphalm. = praec.
Bihai, Linn. Sp. Pl. 1043 = Heliconia Bihai.
Carolinae, Sterler, Hort. Nymph. 109 = rosacea.
Cavendishii, Lambert, ex Paxt. Mag. Bot. iii. (1837) 51, 52 = chinensis.
Chapara, Perr. in Mém. Soc. Linn. Par. iii. (1825) 131 = sapientum.
chinensis, *Sweet, Hort. Brit.* ed. II. 596.—China.
Cliffortiana, Linn. Sp. Pl. 1043, in syn. = sapientum.
coccinea, *Andr. Bot. Rep.* t. 47.—China.
corniculata, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 166 = acuminata.
corniculata, *Lour. Fl. Cochinch.* 644.—Cochinch.
Dacca, Hort. Berol. ex Horan. Prod. Scitam. 41 = sapientum.
dasycarpa, Kurz, in Journ. Agr. Hort. Soc. Ind. xiv. (1865) 301, nomen = velutina.
discolor, *Hort. ex Horan. Prod. Scitam.* 41.—Hab. ?
Ensete, *J. F. Gmel. Syst.* 567.—Afr. trop.
Fehi, *Bert. ex Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 45.—Tahiti; N. Caled.
Fei, Bert. ex Nadeaud. Pl. Tahiti, 39 = praec.
Fitzalani, *F. Muell. Fragm.* ix. 188.—Austral.
glauca, *Roxb. Hort. Beng.* 19; *Fl. Ind.* i. 669.—Burma.
Hillii, *F. Muell. Fragm.* ix. 169.—Austral.
humilis, Aubl. Pl. Gui. ii. 931 = Heliconia humilis.
humilis, Perr. in Mém. Soc. Linn. Par. iii. (1825) 131 = chinensis ?
Jackeyi, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 163 = Hillii ?
Livingstoniana, *J. Kirk, in Journ. Linn. Soc.* ix. (1867) 128.—Afr. trop.
maculata, *Jacq. Hort. Schoenb.* iv. 23. t. 446.—Ins. Maurit.
mensaria, Moench, Meth. 647 = sapientum.
mindanensis, Miq. Fl. Ind. Bat iii. 588 = sapientum.
nana, *Lour. Fl. Cochinch.* 644.—Cochinch.
nepalensis, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 492. —Reg. Himal.
nigra, Perr. in Mém. Soc. Linn. Par. iii. (1825) 131 (Quid ?).—Ins. Philipp.
odorata, Lour. Fl. Cochinch. 644 = sapientum.
oleracea, Vieill. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 46 = sapientum.
ornata, Roxb. Hort. Beng. 19; Fl. Ind. i. 666 = rosacea.
paradisiaca, Linn. Sp. Pl. 1043 = sapientum.
rosacea, *Jacq. Hort. Schoenb.* iv. 22.—As. trop.
rubra, Hort. ex Steud. Nom. ed. II. ii. 167 = rosacea.
Rumphiana, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 164 = acuminata.

MUSA :—
salaccensis, *Zoll. Syst. Verz. Ind. Archip.* 74.—Java.
sanguinea, *Hook. f. Bot. Mag.* t. 5975.—Reg. Himal.
sapientum, *Linn. Syst.* ed. X. 1303.—As. trop.
Seemanni, F. Muell. Fragm. ix. 190 = M. Fehi ?
seminifera, Lour. Fl. Cochinch. 644 = sapientum.
sikkimensis, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 164 = sapientum.
simiarum, *Kurz, l. c.* xiv. (1867) 297.—Malaya.
speciosa, Tenore, Ind. Sem. Hort. Neap. (1829) 16 = rosacea.
sumatrana, *Becc. in Illustr. Hortic.* xxvii. (1880) 37. t. 375.—Sumatra.
superba, *Roxb. Hort. Beng.* 19; *Fl. Ind.* i. 667.—Ind. or.
sylvestris, Colla, Mem. Gen. Musa, 58 = textilis.
textilis, *Née, in Anal. Cienc. Nat.* iv. (1801) 123.—Malaya.
Troglodytarum, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 163, partim = M. Fehi.
Troglodytarum, Linn. Sp. Pl. ed. II. 1477 = sapientum.
Uranoscopus, Colla, Mem. Gen. Musa, 59 = M. Troglodytarum.
Uranoscopus, Lour. Fl. Cochinch. 645 = coccinea.
Uranoscopus, Seem. Fl. Vit. 290 = M. Fehi.
velutina, *H. Wendl. & Drude, in Gartenfl.* (1875) 65. —Reg. Himal.
ventricosa, *Welw. Apont.* 587. n. 45.—Afr. trop.
vittata, W. Ackm. ex Rodigas, in Fl. des Serres, xv. (1862–65) 25 = sapientum.
zebrina, Van Houtte, ex Planch. l. c. x. (1854-55) 223. t. 1061 = sapientum.

MUSANGA, C. Sm. ex R. Br. App. Tuckey, Congo, 453 (1818). *URTICACEAE*, Benth. & Hook. f. iii. 379.
Smithii, *R. Br. in Benn. Pl. Jav. Rar.* 49.—Afr. trop.

MUSCARI, Tourn. ex Mill. Gard. Dict. ed. VI (1752). *LILIACEAE*, Benth. & Hook. f. iii. 811.
BOTRYANTHUS, Kunth, Enum. Pl. iv. 310 (1843).
BOTRYCOMUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160.
BOTRYPHILE, Salisb. Gen. Pl. Fragm. 25 (1866).
COMUS, Salisb. l. c. 24 (1866).
CZEKELIA, Schur, in Oestr. Bot. Wochenschr. vi. (1856) 237.
ETHEIRANTHUS, Kostel. Ind. Hort. Prag. 56 (1844).
LEOPOLDIA, Parl. Fl. Palerm. i. 435 (1845).
MOSCHARIA, Salisb. Gen. Pl. Fragm. 25 (1866).
acutifolium, *Boiss. Fl. Orient.* v. 300.—Reg. Caucas.
acutilobum, Bertol. Misc. Bot. i. 21 = commutatum.
aestivale, *Baker, in Bot. Mag.* t. 6269.—Oriens.
albovirens, Nym. Syll. Suppl. 65 = commutatum.
alpinum, *J. Gay, ex Baker, in Journ. Linn. Soc.* xi. (1871) 413.—As. Min.
ambrosiacum, Moench, Meth. 633 = moschatum.
Argaei, *Hort.—Cf. Gard. Chron.* (1883) II. 394 (Quid?). —Graecia ?
armeniacum, *Leichtlin, ex Baker, in Gard. Chron.* (1878) I. 798.—As. Min.
atlanticum, Boiss. & Reut. Pugill. Pl. Nov. 114 = racemosum.
Aucheri, *Baker, in Journ. Linn. Soc.* xi. (1871) 418. —As. Min.
autumnale, Guss. Ind. Sem. Hort. Boccad. (1825); ex ej. Fl. Sic. Prod. i. 427 = parviflorum.
azureum, Fenzl, Del. Sem. Hort. Vindob. (1858) 4; et in Ann. Sc. Nat. Sér. IV. xii. (1859) 165 = Hyacinthus ciliatus.
bicolor, *Boiss. Fl. Orient.* v. 294.—Aegypt.
bootanensis, Griff. Ic. Pl. Asiat. t. 280 = racemosum.
botryoides, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.; Oriens.
botryoides, Steud. Nom. ed. II. ii. 167 = pedunculare.
Bourgaei, *Baker, in Journ. Linn. Soc.* xi. (1871) 416. —As. Min.
Calandrinianum, Nym. Suppl. Syll. 65 = comosum.
caucasicum, *Baker, in Journ. Linn. Soc.* xi. (1871) 414.—Reg. Caucas.
ciliatum, Ker-Gawl. in Bot. Reg. t. 394 = Hyacinthus ciliatus.
ciliatum, Kotschy, ex Baker, in Journ. Linn. Soc. xi. (1871) 429 = Hyacinthus sessiliflorus.
cilicicum, Kotschy, Reise Taur. 379 = parviflorum.
Clusianum, C. Koch, in Linnaea, xxii. (1849) 253 = comosum.
commutatum, *Guss. Pl. Rar.* 145.—Sicil.

MUSCARI :—
comosum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Reg. Mediterr.; Oriens.
compactum, *Baker, in Gard. Chron.* (1878) I. 799.—Europ.
compactum, Jord. ex Nym. Consp. 734 = racemosum.
concinnum, *Baker, in Gard. Chron.* (1878) I. 799.—Hab. ?
conicum, *Baker, l. c.*—Hab. ?
constrictum, *Tausch, in Flora*, xxiv. (1841) 234.—Corsica.
Courtilleri, Bor. Fl. Centr. Fr. ed. III. ii. 621 = comosum.
crassifolium, *Boj. Hort. Maurit.* 347, nomen.—Madag.
Cupanianum, *Tarant. & Gerb. Cat. Pl. Calat.* 18 (1845); *et ex Guss. Fl. Sic. Syn.* ii. 814.—Sicil.
curtum, *Boiss. Fl. Orient.* v. 292.—Graecia.
dilutum, *Baker, in Gard. Chron.* (1878) I. 798.—Hab. ?
discolor, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 300.—Mesopotam.
Elwesii, *Baker, in Gard. Chron.* (1878) I. 798.—Caria.
filifolium, Wahlenb. in Bergg. Resor, ii. Bihang, 60; et in Oken, Isis, xxi. (1828) 1001 = parviflorum.
fuliginosum, *Freyn, in Flora*, lxviii. (1885) 8.—Tauria.
glaucum, Lindl. Bot. Reg. t. 1085 = Scilla Hohenackeri.
graecum, *Boiss. Fl. Orient.* v. 291.—Graecia.
granatense, *Freyn, in Flora*, lxviii. (1885) 7.—Hispan.
grandifolium, *Baker, in Saund. Refug. Bot.* t. 173.—Hab. ?
Gussonei, Tod. ex Nym. Consp. 734 = racemosum.
Gussonii, *Nym. Syll. Suppl.* 65; *Baker, in Journ. Linn. Soc.* xi. (1871) 415.—Sicil.
Heldreichii, *Boiss. Diagn.* Ser. II. iv. 109.—Graecia.
Holzmanni, *Hirc, in Oestr. Bot. Zeitschr.* xxxiii. (1883) 358.—Graecia.
hymenophorum, Heldr. ex Boiss. Diagn. Ser. II. iv. 109 = Heldreichii.
Kerneri, Marches. in Boll. Soc. Adr. Sc. Trieste, vii. (1882) 266 = botryoides.
Lafarinae, Tineo, ex Parl. Fl. Ital. ii. 503 = commutatum.
latifolium, *J. Kirk, in Trans. Bot. Soc. Edinb.* vi. (1860) 30.—Phrygia.
laxum, *Freyn, in Flora*, lxviii. (1885) 21.—Persia.
Lelievrii, Bor. Fl. Centr. Fr. ed. III. ii. 621 = botryoides.
Letourneuxii, *Boiss. Fl. Orient.* v. 299.—Aegypt.
leucophaeum, C. Koch, in Linnaea, xxii. (1849) 254 = Hyacinthus leucophaeus.
lingulatum, *Baker, in Journ. Bot.* xii. (1874) 6.—As. Min.
longipes, *Boiss. Diagn.* Ser. I. xiii. 36.—Palaest.
longissimum, Medic, in Usteri, Ann. Bot. ii. (1791) 15 = comosum.
luteum, Tod. ex Nym. Consp. 733 = moschatum.
macrocarpum, Sweet, Brit. Flow. Gard. Ser. I. iii. t. 210 = moschatum.
maritimum, *Desf. Fl. Atlant.* i. 308.—Reg. Mediterr.
maritimum, Guss. Fl. Sic. Prod. i. 426 = Gussonii.
micranthum, *Baker, in Gard. Chron.* (1878) I. 799.—Hab. ?
monstrosum, Mill. Gard. Dict. ed. VIII. n. 4 = comosum.
montanum, C. Koch, in Linnaea, xxii. (1849) 253 = Hyacinthus flexuosus.
Mordoanum, *Heldr. in Oestr. Bot. Zeitschr.* xxviii. (1878) 52.—Ins. Corcyra.
moschatum, *Willd. Enum. Hort. Berol.* 378.—As. Min.
Muscarimi, Medic. in Usteri, Ann. Bot. ii. (1791) 15 = moschatum.
neglectum, *Guss. ex Tenore, Fl. Neap. Syll. App.* v. 13.—Reg. Mediterr.
Neumayeri, *Boiss. Fl. Orient.* v. 293.—Graecia.
nivale, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 13. —Persia.
nutans, Bieb. ex Steud. Nom. ed. II. ii. 167 = Scilla festalis.
orchioides, Mill. Gard. Dict. ed. VIII. n. 5 = Lachenalia orchioides.
pallens, Boiss. Fl. Orient. v. 305, sphalm. in syn. = Hyacinthus leucophaeus.
pallens, *Fisch. Ind. Pl. Hort. Gorenk.* (1812) 9; *Link, Enum. Hort. Berol.* i. 331.—Reg. Caucas.
pallens, Hohen. ex Baker, in Journ. Linn. Soc. xi. (1871) 414 = caucasicum.

MUSCARI:—
paniculatum, Medic. in Usteri, Ann. Bot. ii. (1791) 16 = comosum.
paradoxum, *C. Koch, in Linnaea*, xxii. (1849) 253.—Armenia.
parviflorum, *Desf. Fl. Atlant.* i. 309.—Reg. Mediterr.
pauperulum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 78.—Lycia.
pedunculare, Haw. ex Steud. Nom. ed. II. ii. 167 = botryoides.
pendulum, *Trautv. in Act. Hort. Petrop.* ii. (1873) 484.—Armenia.
pharmacusanum, *Boiss. Fl. Orient.* v. 291.—Graecia.
Pinardi, *Boiss. Diagn.* Ser. I. vii. 110.—As. Min.; Syria.
polyanthum, *Boiss. Fl. Orient.* v. 297.—As. Min.
polyphyllum, Schur, Enum. Pl. Transs. 676 = botryoides.
pulchellum, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 109 = racemosum.
pycnanthum, C. Koch, in Linnaea, xxii. (1849) 255 = Hyacinthus pycnanthus.
pyramidale, *Tausch, in Flora*, xxiv. (1841) 235.—Corsica.
racemosum, *Mill. Gard. Dict.* ed. VIII. n. 3.—Reg. Mediterr. et Caucas.
Sartorianum, *Boiss. Fl. Orient.* v. 292.—Graecia.
Schliemanni, *Freyn & Aschers. in Flora*, lxviii. (1885) 6.—As. Min.
sessiliflorum, Spreng. Syst. ii. 66 = Hyacinthus sessiliflorus.
speciosum, Marches. in Bull. Soc. Adr. Trieste, viii. (1882) 267 = neglectum.
stenanthum, *Freyn, in Flora*, lxviii. (1885) 4.—Afr. bor.
Strangwaisii, Kotschy, ex Baker, in Journ. Linn. Soc. xi. (1871) 417 = commutatum.
Strangwaysii, Tenore, Ind. Sem. Hort. Neap. (1837) 5 (Stangwaysii); (1839) 11. n. 9 = racemosum.
suaveolens, Fisch. ex Steud. Nom. ed. II. ii. 168, nomen = moschatum?
Szovitsianum, *Baker, in Gard. Chron.* (1878) i. 799.—Persia; Reg. Caucas.
Szovitsianum, Rupr. ex Boiss. Fl. Orient. v. 295 = racemosum.
tenuiflorum, Tausch, in Flora, xxiv. (1841) 324 = comosum.
theraeum, *Boiss. Fl. Orient.* v. 290.—Ins. Santorin.
transsilvanicum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 76 = botryoides.
trojanum, *Boiss. Fl. Orient.* v. 290.—Agr. Trojan.
tubiflorum, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 84 = comosum.
Weissii, *Freyn, in Oestr. Bot. Zeitschr.* xxviii. (1878) 87.—Archip. Graec.
Wilhelmsii, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 84 = Scilla Hohenackeri.

MUSCARIA, Haw. Saxifr. Enum. 36 (1821) = **Saxifraga**, Tourn.
acaulis, Haw. l. c. 38 = S. muscoides.
atropurpurea, Haw. l. c. = S. muscoides.
basaltica, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386, nomen = S. hypnoides.
caespitosa, Haw. Saxifr. Enum. 37 = S. muscoides.
crocea, Haw. l. c. 38 = S. muscoides.
helviensis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386, nomen = S. hypnoides.
hypnoides, Jord. & Fourr. l. c. = S. hypnoides.
indivisa, Jord. & Fourr. l. c. = S. hypnoides.
laeta, Jord. & Fourr. l. c. = S. hypnoides.
magellanica, Haw. Saxifr. Enum. 38 = S. magellanica.
muscoides, Haw. l. c. 37 = S. muscoides.
parvula, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386, nomen = S. hypnoides.
rhodanica, Jord. & Fourr. l. c. = S. hypnoides.
stenopetala, Haw. Saxifr. Enum. 39 = S. aphylla.
vivariensis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386, nomen = S. hypnoides.
vulcanorum, Jord. & Fourr. l. c. = S. hypnoides.

MUSCIPULA, Rupp. Fl. Jen. ed. Hall. 125 (1745) = **Silene**, Linn. (Caryoph.).
australis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. Muscipula.
portensis, Fourr. l. c. = S. portensis.

MUSENIUM, Nutt. in Torr. & Gray, Fl. N. Am. i. 642 (1840). *UMBELLIFERAE*, Benth. & Hook. f. i. 884.
MARATHRUM, Rafin. in Journ. Phys. lxxxix. (1819) 101.
MUSINEON, Rafin. in Journ. Phys. (1820); ex DC. Prod. iv. 146 (1830).
angustifolium, Nutt. in Torr. & Gray, Fl. N. Am. i. 642 = divaricatum.
divaricatum, *Nutt. l. c.*—Am. bor.
Greenei, A. Gray, in Proc. Am. Acad. viii. (1872) 387 = Carum Hallii.
Hookeri, Nutt. ex Torr. & Gray, Fl. N. Am. i. 642 = divaricatum.
tenuifolium, Hook. Lond. Journ. Bot. vi. (1847) 237 = Peucedanum graveolens, *S. Wats.*
tenuifolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 642.—Am. bor. occ.
trachyspermum, Nutt. l. c. = divaricatum.

MUSINEON, Rafin. in Journ. Phys. (1820); ex DC. Prod. iv. 146 (1830) = praec.

MUSSAENDA, Burm. ex Linn. Diss. Dass. 10 (1747); Amoen. Acad. i. 394 (1749). *RUBIACEAE*, Benth. & Hook. f. ii. 64.
BELILLA, Adans. Fam. ii. 159 (1763).
LANDIA, Comm. ex Juss. Gen. 201, in obs. (1789).
SPALLANZANIA, DC. Prod. iv. 406 (1830).
acuminata, *Blume, Bijdr.* 986.—Java.
acutiflora, *Bartl. ex DC. Prod.* iv. 370 (Quid?).—Mexic.
acutifolia, Steud. Nom. ed. II. ii. 168, sphalm. = praec.
aegyptiaca, Poir. in Lam. Encyc. iv. 394 = luteola.
Afzelii, *G. Don, Gen. Syst.* iii. 490.—Afr. trop.
angustifolia, Wall. ex G. Don, l. c. 491 = parva.
anisophylla, *Vidal, Phan. Cuming. Philipp.* 178.—Ins. Philipp.
arcuata, *Poir. in Lam. Encyc.* iv. 392.—Afr. trop.; Ins. Mascar.
Belilla, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 203 = frondosa.
calycina, Wall. Cat. n. 6253 = macrophylla.
candida, Poir. in Lam. Encyc. iv. 394 = Calycophyllum candidissimum.
candidissima, Roem. & Schult. Syst. v. 251 = Calycophyllum candidissimum.
capsulifera, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 836.—Ins. Socotr.
chinensis, *Lour. Fl. Cochinch.* 152.—Cochinch.
citrifolia, *Lam. ex Poir. in Lam. Encyc.* iv. 393.—Madag.
coccinea, Poir. l. c. 394 = Warszewiczia coccinea.
cordifolia, *Wall. Cat.* n. 6260.—Ins. Penang.
coriacea, Spreng. Neue Entdeck. ii. 145 = Rudgea eriantha.
corymbosa, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 556.—Ind. or.
corymbosa, Wall. Cat. n. 6252, partim = Roxburghii.
cubensis, *Spreng. Syst.* iv. *Cur. Post.* 73.—Cuba.
cuneifolia, D. Don, Prod. Fl. Nep. 139 = Hymenopogon parasiticus.
dasyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 187.—Ins. Ternate.
discolor, Thonn. ex DC. Prod. iv. 372 = elegans.
discolor, *Thou. ex Roem. & Schult. Syst.* v. 254.—Hab.?
Dovinia, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 203 = frondosa.
echitioides, *Willd. ex Roem. & Schult. Syst.* v. 254.—Hab.?
elegans, *Schum. & Thonn. Beskr. Guin. Pl.* 117.—Afr. trop.
eriantha, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 246.—Madag.
erosa, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 193.—Hongkong.
erythrophylla, *Schum. & Thonn. Beskr. Guin. Pl.* 116.—Afr. trop.
flavescens, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 203 = frondosa.
formosa, Jacq. Enum. Pl. Carib. 16 = Randia Mussaendae.
formosa, Linn. Mant. i. 45 = frondosa.
Forsteniana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 188.—Amboina.
frondosa, Blanco, Fl. Filip. ed. I. 167 = grandiflora, *Rolfe*.

MUSSAENDA:—
frondosa, *Linn. Sp. Pl.* 177.—Ind. or.; Malaya.
frondosa, Lour. Fl. Cochinch. 151 = pubescens.
frondosa, Roxb. Hort. Beng. 15; Fl. Ind. i. 557 = Roxburghii.
frondosa, Wall. Cat. 6250 A = macrophylla, Roxburghii.
fruticosa, Linn. Syst. ed. XII. 168 = frondosa.
fulgens, R. Br. ex Tedlie, in Bowdich, Miss. 374 = erythrophylla.
fuscopilosa, *Baker, in Journ. Linn. Soc.* xxi. (1885) 410.—Madag.
glabra, *Hook. & Arn. Bot. Beech. Voy.* 264.—China.
glabra, *Vahl, Symb. Bot.* iii. 38.—Reg. Himal.; Malaya.
glabra, Wall. Cat. n. 6251 = Wallichii.
glomerulata, Lam. ex Poir. in Lam. Encyc. iv. 393 = Rondeletia glomerulata.
grandiflora, *Benth. in Hook. Niger Fl.* 392.—Afr. trop.
grandiflora, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 311.—Ins Philipp.
Griffithii, *Wight, ex Hook. f. Fl. Brit. Ind.* iii. 88.—Burma.
heinsioides, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 70.—Afr. trop.
hirsutula, *Miq. in Journ. Bot. Néerl.* i. (1861) 109.—China.
hispida, D. Don, Prod. Fl. Nep. 139 = macrophylla.
holosericea, Sm. in Rees, Cycl. xxiv. n. 6 = M. Landia.
Humboldtiana, Steud. Nom. ed. I. 540 = Randia Humboldtiana.
hymenopogonoides, *Baker, in Journ. Bot.* xx. (1882) 138.—Madag.
incana, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 229.—Reg. Himal.
ingrata, Wall. ex Hook. f. Fl. Brit. Ind. iii. 89 = frondosa.
Isertiana, *DC. Prod.* iv. 371.—Afr. trop.
Jelinekii, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 135 = Wallichii.
Keenani, *Hook. f. Fl. Brit. Ind.* iii. 87.—Reg. Himal.
lanceolata, Poir. in Lam. Encyc. iv. 393 = Chasalia divaricata.
lanceolata, Spreng. Syst. i. 705 = luteola.
Landia, *Poir. in Lam. Encyc.* iv. 392.—Ins. Maurit.
latifolia, Poir. Encyc. Suppl. iv. 36 = M. Landia.
longifolia, *Lam. ex Poir. in Lam. Encyc.* iv. 393.—Madag.
longifolia, Wall. Cat. n. 6258 = Wallichii.
Luculia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 139 = Luculia gratissima.
luteola, *Delile, Cent. Pl. Méroé*, 65.—Arab.; Nubia.
luteola, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 357 = Pentas carnea.
macrophylla, Kurz, For. Fl. Brit. Burma, ii. 57 = frondosa.
macrophylla, Schum. & Vahl, in Schum. & Thonn. Beskr. Guin. Pl. 118 = Isertiana.
macrophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 228.—Reg. Himal.; Malaya.
macropoda, *Baker, in Journ. Linn. Soc.* xxi. (1885) 410.—Madag.
nitida, H. B. & K. Nov. Gen. et Sp. iii. 410 = Randia nitida.
odorifera, Vell. Fl. Flum. 94; ii. t. 144 (Randiae sp.?).—Bras.
parva, *Wall. Cat.* n. 6261.—Burma.
parviflora, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 110.—Japon.
pavettaefolia, *Kurz, For. Fl. Brit. Burma*, ii. 57.—Burma.
penangensis, Miq. Fl. Ind. Bat. ii. 214 = glabra.
philippica, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 245.—Ins. Philipp.
platyphylla, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 70.—Afr. trop.
polita, *Hiern, l. c.* 67.—Afr. trop.
pubescens, Buch-Ham. in Wall. Cat. n. 6257 = Roxburghii, incana.
pubescens, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 372.—China.
pubescens, H. B. & K. Nov. Gen. et Sp. iii. 410 = Randia Humboldtiana.
racemosa, Sieber, ex DC. Prod. iv. 392 = Bertiera Zaluzania.
Reinwardtiana, *Miq. Fl. Ind. Bat.* ii. 211.—Ins. Celebes.
repens, Wall. Cat. n. 6263 = Ophiorrhiza calcarata.

MUSSAENDA:—
Roxburghii, *Hook. f. Fl. Brit. Ind.* iii. 87.—Reg. Himal.
rubra, Steud. Nom. ed. II. ii. 168, sphalm. = rufa, zanzibarica.
rufa, Boj. in Ann. Sc. Nat. Sér. II. iv. (1835) 264 = zanzibarica.
rufa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 246.—Afr. trop.
rufinervia, *Miq. Fl. Ind. Bat.* ii. 212.—Sumatra.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 27 = Pentas Schimperiana.
sericea, *Blume, Bijdr.* 986.—Java.
setulosa, Klotzsch, in Ber. Verh. Berl. Acad. (1853) 499 = glabra.
speciosa, *Poir. Encyc. Suppl.* iv. 37.—Venezuela.
spinosa, Jacq. Enum. Pl. Carib. 16 = Randia armata.
splendida, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 36. t. 13.—Afr. trop.
Stadmanni, Michx. ex DC. Prod. iv. 372 = M. Landia.
stenocarpa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 68.—Afr. trop.
sumatrensis, *Heyne, ex Roth, Nov. Pl. Sp.* 152.—Sumatra.
tenuiflora, *Benth. in Hook. Niger Fl.* 392.—Afr. trop.
tetracantha, Cav. Ic. v. 20 = Randia tetracantha.
tetrandra, *Roem. & Schult. Syst.* v. 254.—Hab.?
Teysmanniana, *Miq. Fl. Ind. Bat* ii. 213.—Ins. Bali.
theifera, Pierre, ex Carr. in Rev. Hortic. (1883) 93 = uniflora.
Thouarsiana, *Baill. Adansonia*, xii. (1879) 295.—Madag.
tomentosa, Reinw. ex Miq. Fl. Ind. Bat. ii. 211 = Reinwardtiana.
tomentosa, Wight, ex Hook. f. Fl. Brit. Ind. iii. 90 = frondosa.
tomentosa, *Wight, in Wall. Cat.* n. 6265.—Ind. or.
trichophlebia, *Baker, in Journ. Linn. Soc.* xx. (1883) 166.—Madag.
tubulosa, *D. Dietr. Syn. Pl.* i. 724.—Bras.
uniflora, *Wall. Cat.* n. 6264.—Burma.
variolosa, *Wall. l. c.* n. 6259.—Burma.
vestita, *Baker, in Journ. Linn. Soc.* xx. (1883) 166.—Madag.
villosa, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 90 = frondosa.
villosa, *Wall. Cat.* n. 6254.—Penang.
Wallichii, *G. Don, Gen. Syst.* iii. 490.—Burma.
zanzibarica, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 264.—Afr. trop.
Zollingeriana, *Klotzsch, in Ber. Verh. Berl. Acad.* (1853) 500.—Java.

MUSSAENDOPSIS, Baill. Adansonia, xii. (1879) 282. *RUBIACEAE.*
Beccariana, *Baill. l. c.*—Borneo.

MUSSCHIA, Dum. Comm. Bot. 28 (1822). *CAMPANULACEAE*, Benth. & Hook. f. ii. 560.
BENAUREA, Rafin. Fl. Tellur. ii. 77 (1836).
CHRYSANGIA, Link, Handb. i. 632 (1829).
aurea, *Dum. Comm. Bot.* 28.—Ins. Madera.
Wollastoni, *Lowe, in Hook. Kew Journ.* viii. (1856) 298.—Ins. Madera.

MUSSINIA, Willd. Sp. Pl. iii. 2263 (1803) = **Gazania**, Gaertn. (Compos.).
incisa, Willd. l. c. 2264 = G. pinnata.
linearis, Willd. l. c. 2263 = G. subulata.
Othonna, Willd. l. c. 2264 = G. Othonnites.
pinnata, Willd. l. c. 2265 = G. pinnata.
pusilla, Licht. ex Less. Syn. Comp. 50 = G. Lichtensteinii.
speciosa, Willd. Sp. Pl. iii. 2264 = G. pinnata.
uniflora, Willd. l. c. 2263 = G. uniflora.

MUSTELIA, Cav. ex Steud. Nom. ed. II. i. 361, in syn. (1840); ii. 168 = **Chusquea**, Kunth (Gramin.).
arundinacea, Cav. ex Steud. ll. cc. = C. scandens.

MUSTELIA, Spreng. Nachr. i. Bot. Gart. Halle, 28 (1801); et in Trans. Linn. Soc. vi. (1802) 152. t. 13 = **Stevia**, Cav. (Compos.).
Eupatoria, Spreng. ll. cc. = S. Eupatoria.

MUSTERON, Rafin. Fl. Tellur. ii. 50 (1836) = **Conyza**, Linn. (Compos.).
bellidifolium, Rafin. l. c. = Vernonia cinerea.
glabrum, Rafin. l. c. = Vernonia longifolia.
pulchellum, Rafin. l. c. = C. sophiaefolia.
uniflorum, Rafin. l. c. = C. uniflora.

MUTABEA, J. F. Gmel. Syst. 1009 (1791) = **Moutabea**, Aubl. (Polygal.).

MUTAFINIA, Rafin. Atl. Journ. 199 (1833) = **Limosella**, Linn. (Scrophul.).
capensis, Rafin. Fl. Tellur. iv. 57 = L. capensis.
silesiaca, Rafin. l. c. = L. aquatica.

MUTARDA, Bernh. Syst. Verz. Erf. 184 (1800) = **Brassica**, Tourn. (Crucif.).
nigra, Bernh. l. c. 197 = B. nigra.

MUTELIA, Gren. ex Mutel, Fl. Fr. iii. 21 (1836) = **Melissa**, Tourn. (Labiat.).
officinalis, Gren. ex Mutel, l. c. Suppl. 87 = Melissa officinalis.

MUTISIA, Linn. f. Suppl. 57 (1781). *COMPOSITAE*, Benth. & Hook. f. ii. 485.
APLOPHYLLUM, Cass. in Dict. Sc. Nat. xxxiii. 463, 472 (1824).
GUARIRUMA, Cass. l. c. 463, 472 (1824).
acerosa, *Poepp. ex Less. Syn. Comp.* 107.—Chili.
acuminata, *Ruiz & Pav. Syst. Veg.* 192.—Chili.
arachnoidea, Mart. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 264 = speciosa.
araucana, *Phil. in Anal. Univ. Chil.* (1872) 737.—Chili.
auriculata, Poepp. ex Less. Syn. Comp. 105 = spinosa.
Berteriana, *Poepp. & Endl. Nov. Gen. et Sp.* i. 19.—Chili.
Berterii, *DC. Prod.* vii. 8.—Chili.
Bipontini, *Mandon, ex Sch.-Bip. in Linnaea*, xxxiv. (1865–66) 527.—Bolivia.
bracteata, Remy, in C. Gay, Fl. Chil. iii. 268 = latifolia.
breviflora, *Phil. in Anal. Univ. Chil.* (1884) reimpr. 4; *et in Gartenfl.* (1884) 227.—Ins. J. Fernand.
Bridgesii, *Poepp. & Endl. Nov. Gen. et Sp.* i. 17.—Chili.
campanulata, *Less. in Linnaea*, v. (1830) 269.—Bras.
cana, *Poepp. & Endl. Nov. Gen. et Sp.* i. 17.—Chili.
Candolleana, Gardn. in Field. Sert. Pl. t. 45–46 = viciaefolia.
caryophyllata, *Vell. Fl. Flum.* viii. t. 103.—Bras.
Clematis, *Linn. f. Suppl.* 373.—Peruv.
cneorifolia, Decne. ex Wedd. Chlor. And. i. 22 = Orbignyana.
coccinea, *A. St. Hil. Voy. Distr. Diam.* i. 386.—Bras.
dealbata, DC. Prod. vii. 6 = coccinea.
decurrens, *Cav. Ic.* 65. t. 497.—Chili.
Gayana, *Remy, in C. Gay, Fl. Chil.* iii. 268.—Chili.
glauca, *Phil. in Linnaea*, xxxiii. (1864–65) 110.—Chili.
gracilis, *Meyen, Reise*, i. 342.—Chili.
grandiflora, *Humb. & Bonpl. Pl. Aequin.* i. 177. t. 50.—N. Granat.
grossedentata, *Phil. in Linnaea*, xxviii. (1856) 710.—Chili.
hastata, Bert. ex DC. Prod. vii. 7 = spinosa.
hastata, *Cav. Ic.* v. 64. t. 494.—Chili.
heliantha, Poepp. ex Less. Syn. Comp. 107 = decurrens.
hirsuta, Meyen, Reise, i. 451 = viciaefolia.
homoeantha, *Wedd. Chlor. And.* i. 21.—Bolivia.
Hookeri, *Meyen, Reise*, i. 348.—Chili.
hypoleuca, Mart. & Zucc. ex Baker, in Mart. Fl. Bras. vi. III. 366 = coccinea.
ilicifolia, *Cav. Ic.* v. 63.—Chili.
inflexa, Cav. l. c. 65 = subulata.
involucrata, *Phil. in Linnaea*, xxviii. (1856) 710.—Chili.
lanata, *Ruiz & Pav. Syst. Veg.* 192.—Chili.
lanigera, *Wedd. Chlor. And.* i. 16.—Bolivia.
latifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 270.—Chili.
Lechleri, Sch. Bip. in Lechl. Berb. Am. Austr. 54 = retusa.
ledifolia, *Decne. ex Wedd. Chlor. And.* i. 20.—Bolivia.
linariaefolia, *Remy, in C. Gay, Fl. Chil.* iii. 271.—Chili.
linearifolia, *Cav. Ic.* v. 66. t. 500.—Chili.
linearifolia, Hook. Bot. Misc. i. (1830) 11. t. 8 = Hookeri.
linifolia, *Hook. Bot. Misc.* i. (1830) 12. t. 9.—Chili.
macrophylla, *Phil. in Linnaea*, xxxiii. (1864–65) 108.—Chili.
Mathewsii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 107, in nota.—Peruv.

MUTISIA:—
Meyeniana, *Remy, ex Phil. in Anal. Univ. Chil.* (1861) i. 62.—Chili.
microphylla, Willd. ex DC. Prod. vii. 6 = speciosa.
neriifolia, Decne. ex Wedd. Chlor. And. i. 22 = Orbignyana.
obovata, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 351 = Stifftia Martiana.
oligodon, *Poepp. & Endl. Nov. Gen. et Sp.* i. 18. t. 29.—Chili.
Orbignyana, *Wedd. Chlor. And.* i. 22.—Peruv.
parviflora, Remy, in C. Gay, Fl. Chil. iii. 274 = acerosa.
patagonica, *Phil. in Linnaea*, xxviii. (1856) 711.—Chili.
peduncularis, *Cav. Ic.* v. 62. t. 691.—Peruv.
petiolulata, Decne. ex Wedd. Chlor. And. i. 15 = viciaefolia.
pichinchensis, *Karst. Fl. Columb.* i. 93. t. 46.—Ecuador.
polyphylla, *Phil. in Linnaea*, xxxiii. (1864–65) 110.—Chili.
reticulata, *Phil. l. c.* 108.—Chili.
retrorsa, *Cav. Ic.* v. 65. t. 498.—Patagon.
retusa, *Remy, in C. Gay, Fl. Chil.* iii. 269.—Chili.
rosea, *Poepp. ex Less. Syn. Comp.* 106.—Chili.
runcinata, Willd. Sp. Pl. iii. 2069 = retrorsa.
sagittata, Willd. l. c. 2070 = hastata.
sanguinea, *Meyen, Reise*, i. 343.—Chili.
sinuata, *Cav. Ic.* v. 66. t. 499.—Chili.
sinuata, Willd. ex Less. in Linnaea, v. (1830) 271 = taraxacifolia.
speciosa, *Ait. ex Hook. Bot. Mag.* t. 2705.—Ecuador.
spectabilis, *Phil. in Linnaea*, xxxiii. (1864–65) 107.—Chili.
spinosa, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 107 = ilicifolia.
spinosa, *Ruiz & Pav. Syst. Veg.* 193.—Chili.
splendens, *Renjifo, in Renjifo & Phil. Desc. Alg. Pl. Nuev.* (1884) 5.—Chili.
subspinosa, *Cav. Ic.* v. 64. t. 495.—Peruv.; Chili.
subulata, *Ruiz & Pav. Syst. Veg.* 193.—Chili.
taraxacifolia, *Less. in Linnaea*, v. (1830) 271.—Chili.
tridens, *Poepp. ex Less. Syn. Comp.* 105.—Chili.
truncata, D. Don, in Trans. Linn. Soc. xvi. (1830) 269 = ilicifolia.
ulicina, D. Don, in Phil. Mag. xi. (1832) 391 = acerosa.
versicolor, *Phil. in Anal. Univ. Chil.* (1884), reimpr. 5; *et in Gartenfl.* (1884) 228.—Ins. J. Fernand.
viciaefolia, *Cav. Ic.* v. 62. t. 490.—Chili.
vitellina, *Roezl, in Belg. Hortic.* (1874) 39.—Peruv.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 109.—Chili.

MUTUCHI, J. F. Gmel. Syst. 1085 (1791) = MOUTOUCHI, Aubl. = **Pterocarpus**, Linn. (Legumin.).

MUXIRIA, Welw. Apont. 573 (1858) = **Eriosema**, DC. (Legum.).
utilis, Welw. l. c. = E. Muxiria.

MUZA, Stokes, Bot. Mat. Med. i. 472 (1812) = **Musa**, Linn. (Scitam.).

MYAGRUM, [Tourn.] Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 96.
BRICOUR, Adans. Fam. ii. 423 (1763).
DELTOCARPUS, L'Hérit. ex DC. Syst. ii. 573 (1821).
aegyptium, Linn. Sp. Pl. 641 = Rapistrum aegyptiacum.
alpinum, Lapeyr. Hist. Abr. Pl. Pyr. i. 362 = Cochlearia auriculata.
Alyssum, Mill. Gard. Dict. ed. VIII. n. 2 = Camelina sativa.
amphibium, Loisel. Fl. Gall. ed. II. ii. 50 = Nasturtium amphibium.
amplexicaule, Moench, Meth. 221 = perfoliatum.
aquaticum, Lam. Fl. Fr. ii. 483 = Nasturtium amphibium.
arborescens, Jacq. Coll. i. 39 = Crambe strigosa.
argenteum, Pursh, Fl. Am. Sept. ii. 434 = Vesicaria ludoviciana.
armeniacum, Steud. Nom. ed. I. 541 = Camelina sylvestris.
asperum, Poir. Encyc. Suppl. ii. 47 = Bunias Erucago.
auriculatum, DC. Fl. Fr. vi. 597 = Cochlearia saxatilis.
austriacum, Jacq. Obs. Bot. i. 43 = Nasturtium austriacum.

MYAGRUM:—
balearicum, Lam. Encyc. i. 571 = Succowia balearica.
Bauhini, C. C. Gmel. Fl. Bad. iii. 7 = Camelina foetida.
biarticulatum, Crantz, Stirp. Austr. ed. I. fasc. i. 4; ed. II. 7 = Rapistrum perenne.
bursifolium, Thuill. Fl. Par. ed. II. i. 319 = Calepina Corvini.
chloraefolium, Willd. Sp. Pl. iii. 407 = Aethionema cordatum.
clavatum, Lam. Fl. Fr. ii. 482 = Bunias Erucago.
clavatum, Poir. Encyc. Suppl. ii. 47 = Rapistrum rugosum.
cornutum, Lam. Encyc. i. 571 = Pugionium cornutum.
Crantzii, Vitm. Summa Pl. iv. 9 = Nasturtium austriacum.
dentatum, Willd. Phyt. 9 = Camelina sativa.
erucaefolium, Vill. Fl. Delph. 66; Hist. Pl. Dauph. iii. 279 = Calepina Corvini.
Erucago, Lam. Encyc. i. 571 = Bunias Erucago.
erucoides, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 757 = Calepina Corvini.
glabrum, Gilib. Fl. Lituan. ii. 49 = Camelina sativa.
grandiflorum, Banks, ex DC. Syst. ii. 699 = Brachycarpaea varians.
hispanicum, Linn. Sp. Pl. 640 = Rapistrum Linnaeanum.
iberioides, Brot. Fl. Lusit. i. 562 = Calepina Corvini.
irregulare, Asso, Syn. Stirp. Arag. 82 = Calepina Corvini.
littorale, Scop. Fl. Carn. ed. II. ii. 12 = perfoliatum.
monospermum, Forsk. Fl. Aegypt. Arab. p. xxix = Rapistrum rugosum.
montanum, Berger. Phyt. iii. 140; ex DC. Syst. ii. 360 = Cochlearia auriculata.
natans, Patrin, ex DC. Syst. ii. 198 = Cochlearia amphibia.
orientale, Linn. Sp. Pl. 640 = Rapistrum orientale.
orientale, Sieber, ex Steud. Nom. ed. II. ii. 169 = Bunias Erucago.
palustre, Lam. Encyc. i. 572 = Nasturtium palustre.
paniculatum, Linn. Sp. Pl. 641 = Neslia paniculata.
perenne, Linn. l. c. 640 = Rapistrum perenne.
perenne, Scop. Fl. Carn. ed. II. ii. 10 = Rapistrum rugosum.
perenne, Stev. ex DC. Syst. ii. 434 = Rapistrum costatum.
perfoliatum, *Linn. Sp. Pl.* 640.—Europ.; Oriens.
pinnatifidum, Ehrh. ex DC. Syst. ii. 516 = Camelina sativa.
pinnatum, [Soland. in] Russ. Aleppo, ed. II. ii. 256 = Brassica adpressa?
procumbens, Pourr. in Mém. Acad. Toul. iii. (1788) 322 (Quid?).—Gallia.
prostratum, Poir. Encyc. Suppl. ii. 48 = Muricaria prostrata.
pumilum, Lam. Encyc. i. 572 = Nasturtium palustre.
pyrenaicum, Lam. l. c. 571 = Nasturtium pyrenaicum.
rigidum, Pall. Reise, iii. 741 = Euclidium syriacum.
rostratum, Scop. Fl. Carn. ed. II. ii. 11 = Euclidium syriacum.
rugosum, Bieb. Fl. Taur. Cauc. ii. 86 = Rapistrum costatum.
rugosum, Gilib. Fl. Lituan. ii. 49 = Neslia paniculata.
rugosum, Linn. Sp. Pl. 640 = Rapistrum rugosum.
rugosum, Vill. Prosp. 37 = Calepina Corvini.
sativum, Aubry, Morb. An. x. 28; ex DC. Syst. ii. 378 = Thlaspi alliaceum.
sativum, Cav. Ic. i. 47. t. 66 = Camelina sylvestris.
sativum, Linn. Sp. Pl. 641 = Camelina sativa.
saxatile, Linn. Syst. ed. X. 1126 = Cochlearia saxatilis.
sphaerocarpum, Jacq. Obs. Bot. ii. 20. t. 41 = Crambe hispanica.
spinosum, Lam. Encyc. i. 571 = Zilla myagroides.
stylosum, Gochn. ex DC. Syst. ii. 432 = Rapistrum rugosum.
syriacum, Lam. Encyc. i. 570 = Euclidium syriacum.
taraxacifolium, Lam. l. c. = Bunias orientalis.
tataricum, Poir. Encyc. Suppl. ii. 48 = Euclidium tataricum.
utriculatum, Berger. ex DC. Syst. ii. 296 = Vesicaria utriculata.
venosum, Pers. Syn. ii. 183 = Rapistrum rugosum.
verrucosum, Lam. Encyc. i. 570 = Ochthodium aegyptiacum.

MYANTHE, Salisb. Gen. Pl. Fragm. 34 (1866) = **Ornithogalum**, Tourn. (Liliac.).
arabica, Salisb. l. c. = O. arabicum.

MYANTHUS, Lindl. Bot. Reg. sub t. 1538 (1832) = **Catasetum**, Rich. (Orchid.).
barbatus, Lindl. l. c. t. 1778 = C. barbatum.
callosus, Beer, Prakt. Orch. 272 = C. callosum.
cernuus, Lindl. in Bot. Reg. sub t. 1538 = C. trifidum.
deltoideus, Lindl. l. c. t. 1896 = C. deltoideum.
fimbriatus, C. Morr. in Ann. Soc. Bot. Gand, iv. (1848) 453. t. 231 = C. fimbriatum.
grandiflorus, Beer, Prakt. Orch. 272 = C. callosum.
Lansbergii, Reinw. & De Vriese.—Cf. Linnaea, xxiv. (1851) 170 = C. Lansbergii.
sanguineus, Linden, Cat. n. 6 (1851) 7; et ex Lem. Jard. Fleur. ii. (1852) Misc. 134 = C. sanguineum.
spinosus, Hook. Bot. Mag. t. 3802 = C. barbatum.
Warczewicii, Beer, Prakt. Orch. 272 = C. Warczewitzii.

MYARIS, Presl, Bot. Bemerk. 40 (1844); in Abh. Boehm. Ges. V. iii. (1845) 470, 553 = **Clausena**, Burm. (Rutac.).
inaequalis, Presl, l. c. = C. inaequalis.

MYCARANTHES, Blume, Bijdr. 352. t. 57 (1825) = **Eria**, Lindl. (Orchid.).
latifolia, Blume, l. c. 353 = E. latifolia, *Reichb. f.*
lobata, Blume, l. c. 352 = E. lobata.
obliqua, Lindl. Bot. Reg. (1840) Misc. 77 = E. obliqua.
oblitterata, Blume, Bijdr. 353 = E. oblitterata.
stricta, Lindl. in Wall. Cat. n. 1970 = E. stricta.

MYCARANTHUS, Benth. & Hook. f. Gen. iii. 510 (1883), sphalm. = praec.

MYCELIS, Cass. in Dict. Sc. Nat. xxxiii. 483 (1824) = **Lactuca**, Tourn. (Compos.).
ambigua, Cass. l. c. xlviii. 426 = L. deltoidea.
angulosa, Cass. l. c. xxxiii. 484 = L. muralis.
muralis, Dum. Fl. Belg. 60; Reichb. Fl. Germ. Excurs. 272 = L. muralis.

MYCETANTHE, Reichb. Nom. 61 (1841) = **Brugmansia**, Blume (Cytinac.).

MYCETIA, Reinw. in Syll. Ratisb. ii. (1828) 9 = **Adenosacme**, R. Br. (Rubiac.).
cauliflora, Reinw. l. c. = A. longifolia.

MYCONIA, Lapeyr. Hist. Abr. Pl. Pyr. 115 (1813) = **Ramondia**, Rich. (Scrophul.).
borraginea, Lapeyr. l. c. = R. pyrenaica.
pyrenaica, Auct. ex Steud. Nom. ed. II. ii. 169 = R. pyrenaica.

MYCONIA, Neck. Elem. i. 22 (1790) = **Chrysanthemum**, Tourn. (Compos.).
chrysanthemoides, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 247 = C. Myconis.
Chrysanthemum, Sch. Bip. l. c. = C. Myconis.
macrotus, Walp. Ann. i. 421 = C. macrotum.
multicaulis, Walp. l. c. = C. multicaule.

MYCOSTYLIS, Rafin. Sylva Tellur. 19 (1838) = **Sonneratia**, Linn. f. (Lythrar.).

MYCROPUS, Gouan, Fl. Monsp. 339 (1765) = **Micropus**, Linn. (Compos.).

MYCROSERIS, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 250 = **Microseris**, D. Don (Compos.).

MYCTANTHES, Rafin. New Fl. Am. ii. 22 (1836) = **Aster**, Tourn. (Compos.).
axillaris, Rafin. l. c. = Solidago nemoralis?
latifolia, Rafin. l. c. = A. macrophyllus?
punicea, Rafin. l. c. = A. puniceus.

MYGALURUS, Link, Enum. Hort. Berol. i. 92 (1821) = **Festuca**, Linn. (Gramin.).
alopecuroides, Link, l. c. = F. Alopecuros.
bromoides, Link, l. c. = F. bromoides.
caudatus, Link, l. c. = F. Myuros.
delicatulus, Link, l. c. = F. delicatula.
geniculatus, Link. l. c. = F. clavata.
uniglumis, Link, l. c. = F. bromoides.

MYGINDA, Jacq. Enum. Pl. Carib. 1 (1760). *CELASTRINEAE*, Benth. & Hook. f. i. 366.
CROSSOPETALUM, P. Br. Hist. Jamaic. 145 (1756).
RHACOMA, Linn. Syst. ed. X. 896 (1759).
Aquifolia, Griseb. Cat. Pl. Cub. 55 = ilicifolia.
brasiliensis, Spreng. Syst. i. 496 (= *Mespilodaphne glauca*).—Bras.
Bredemeyeri, *Schult. Mant.* iii. 349.—Portorico.
coccinea, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 604 = M. Uragoga.
Congonha, Miers, in Ann. & Mag. Nat. Hist. Ser. III. ix. (1862) 113, sphalm. = *M. Gongonha.*
Crossopetalum, Schult. Mant. iii. 346, sphalm. = *M. Glossopetalum.*
disticha, *Hook. f. Fl. Antarct.* 254.—Reg. Magellan.
Glossopetalum, Raeusch. Nom. ed. III. 43 = M. Rhacoma.
Gongonha, DC. Prod. ii. 13 = Villaresia mucronata.
ilicifolia, *Poir. in Lam. Encyc.* iv. 396.—Am. bor.; Ind. occ.
integrifolia, *Poir. l. c.*—Ind. occ.
latifolia, *Sw. Prod. Veg. Ind. Occ.* 39.—Am. bor.; Ind. occ.
microphylla, *DC. Prod.* ii. 12.—Ins. S. Doming.
myricoides, Willd. ex Schult. Mant. iii. 349 = Ilex myricoides.
myrsinoides, *H. B. & K. Nov. Gen. et Sp.* vii. 68. t. 620.—Peruv.
myrtifolia, Nutt. Gen. Am. i. 109 = Pachistima Myrsinites.
pallens, *Sm. in Rees, Cycl.* xxiv. n. 4.—Am. bor.; Ind. occ.
pentandra, Willd. ex Schult. Mant. iii. 349 = Bredemeyeri.
pungens, *Wright, in Sauv. Fl. Cub.* 22.—Cuba.
Rhacoma, *Sw. Prod. Veg. Ind. Occ.* 39.—Am. bor.; Ind. occ.
rotundata, *Lam. Tabl. Encyc.* i. 308.—Ins. Portoric.
scoparia, *Hook. & Arn. Bot. Beech. Voy.* 283.—Mexic.
theoides, *Benth. Pl. Hartw.* 127.—Peruv.
Uragoga, *Jacq. Enum. Pl. Carib.* 12; *Select. Am.* 24. t. 16.—Am. trop. et bor.

MYGINDUS, Hook. & Arn. Bot. Beech. Voy. 283 (1838?), sphalm. = praec.

MYLACHNE, Steud. Nom. ed. II. i. 561, in syn. (1840); ii. 170 = seq.

MYLANCHE, Wallr. Orob. Gen. 75 (1825) = **Epifagus**, Nutt. (Orob.).
virginiana, Wallr. l. c. 76 = E. americanus.

MYLINUM, Gaud. Fl. Helv. ii. 17, 344 (1828) = **Selinum**, Linn. (Umbell.).
Carvifolia, Gaud. l. c. 344 = S. Carvifolia.
commutatum, Gaud. l. c. 329, in obs. = Peucedanum alsaticum?

MYLOCARYUM, Willd. Enum. Hort. Berol. 454 (1809) = **Cliftonia**, Banks (Cyrill.).
ligustrinum, Willd. l. c. = C. nitida.

MYLOSPHORA, Neck. Elem. ii. 280 (1790) = **Singana**, Aubl. (Inc. sed.).

MYOBROMA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = **Astragalus**, Tourn. (Legumin.).
albana, Stev. l. c. 268 = A. nummularius.
buchtormensis, Stev. l. c. = A. buchtormensis.
cahyrica, Stev. l. c. = A. kahiricus.
caprina, Stev. l. c. = A. caprinus.
diffusa, Stev. l. c. = A. dolichophyllus.
exscapa, Stev. l. c. = A. exscapus.
fabacea, Stev. l. c. = A. fabaceus.
galactites, Stev. l. c. = A. galactites.
geminiflora, Stev. l. c. 269 = A. geminiflorus.
Henningii, Stev. l. c. xxix. (1856) ii. 151 = A. utriger.
humilis, Stev. l. c. iv. (1832) 268 = A. humilis.
lanigera, Stev. l. c. = A. lanigerus.
lanuginosa, Stev. l. c. xxix. (1856) ii. 150 = A. lanuginosus, *Kar. & Kir.*
lasiopetala, Stev. l. c. = A. lasiopetalus.
lignosa, Stev. l. c. iv. (1832) 268 = A. declinatus.
longiflora, Stev. l. c. = A. longiflorus.
megalantha, Stev. l. c. = A. megalanthus.
mollis, Stev. l. c. = A. mollis.

MYOBROMA :—
nummularia, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = Astragalus nummularius.
ovina, Stev. l. c. = A. ovinus.
pubiflora, Stev. l. c. = A. pubiflorus.
rupifraga, Stev. l. c. = A. rupifragus.
Schanginiana, Stev. l. c. xxix. (1856) II. 150 = A. Schanginianus.
testiculata, Stev. l. c. iv. (1832) 268 = A. testiculatus.
triphylla, Stev. l. c. 269 = A. triphyllus.
uniflora, Stev. l. c. = A. uniflorus.
utrigera, Stev. l. c. 268 = A. utriger.

MYODA, Lindl. in Wall. Cat. n. 7390 (1832); et Bot. Reg. sub t. 1618 (1833) = **Haemaria**, Lindl. (Orchid.).
rufescens, Lindl. in Wall. l. c. = H. discolor.

MYODIUM, Salisb. in Trans. Hort. Soc. i. (1812) 289 = **Ophrys**, Linn. (Orchid.).
araniferum, Salisb. l. c. = O. aranifera.

MYODOCARPUS, Brongn. & Gris, in Bull. Soc. Bot. Fr. viii. (1861) 123. *ARALIACEAE*, Benth. & Hook. f. i. 935.
fraxinifolius, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 270.—N. Caled.
pinnatus, *Brongn. & Gris, l. c.* viii. (1861) 123.—N. Caled.
simplicifolius, *Brongn. & Gris, l. c.*—N. Caled.
Vieillardi, *Brongn. & Gris, l. c.* xii. (1865) 270.—N. Caled.

MYOGALUM, Link, Handb. i. 163 (1829) = **Ornithogalum**, Tourn. (Liliac.).
affine, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1855) App. 10 = O. nutans.
Boucheanum, Kunth, Enum. Pl. iv. 348 = O nutans.
nutans, Link, Handb. i. 164 = O. nutans.
prasandrum, Walp. Ann. i. 858 = O nutans.
Thirkeanum, C. Koch, in Linnaea, xix. (1847) 11 = O. nutans.

MYONIMA, Comm. ex Juss. Gen. 206 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 115.
NESCIDIA, A. Rich. in Mém Soc. Hist. Nat. Par. v. (1834) 192.
Borboniae, Raeusch. Nom. ed. III. 37 = obovata.
grandiflora, *Willd. ex Schult. Mant.* iii. 49.—Hab. ?
heterophylla, Boj. Hort. Maurit. 169 = myrtifolia.
lanceolata, Willd. Sp. Pl. i. 614 = myrtifolia.
latifolia, Boj. Hort. Maurit. 169 = multiflora.
multiflora, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 212.—Ins. Maurit.
myrtifolia, *Lam. Tabl. Encyc.* i. 288.—Ins. Mascar.
oblongifolia, Willd. ex Schult. Mant. iii. 49 = multiflora ?
obovata, *Lam. Tabl. Encyc.* i. 288.—Ins. Mascar.
umbellata, Bartl. ex DC. Prod. iv. 463 = Plectronia umbellata.
umbellata, Hook. & Arn. Bot. Beech. Voy. 86 (= *Canthium odoratum*).—Ins. Sandvic.

MYOPORDON, Boiss. Diagn. Ser. I. vi. 107 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 474.
Aucheri, *Boiss. & Diagn.* Ser. I. vi. 108.—Persia.
persicum, *Boiss. l. c.* 107.—Persia.

MYOPORUM, Banks & Soland. ex Forst. f. Prod. 44 (1786). *MYOPORINEAE*, Benth. & Hook. f. ii. 1124.
ANDREUSIA, Vent. Jard. Malm. t. 108 (1807).
BERTOLONIA, Spin. Cat. Jard. St. Sebast. (1809) 24.
DISOON, A. DC. in DC. Prod. xi. 703 (1847).
PENTACOELIUM, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 151.
POGONIA, Andr. Bot. Rep. tt. 212, 283 (1801).
POLYCOELIUM, A. DC. in DC. Prod. xi. 705 (1847).
acuminatum, *R. Br. Prod.* 515.—Austral.
adscendens, R. Br. l. c. 516 = serratum.
apiculatum, DC. Prod. xi. 707 = serratum.
Bateae, *F. Muell. in Proc. Linn. Soc. N. S. W.* vi. (1881) 792.—Austral.
Beckeri, *F. Muell. ex Benth. Fl. Austral.* v. 7.—Austral.
bontioides, *A. Gray, in Proc. Am. Acad.* vi. (1862) 52.—Japon.
brevifolium, Bartl. in Lehm. Pl. Preiss. i. 350 = Pholidia brevifolia.

MYOPORUM:—
brevipes, *Benth. Fl. Austral.* v. 6.—Austral.
caprarioides, Benth. in Enum. Pl. Hueg. 77 = serratum.
chinensis, *A. Gray, in Proc. Am. Acad.* vi. (1862) 52.—China.
crassifolium, *Forst. f. Prod.* 44.—Ins. Pacif.
crystallinum, *Hort. ex Kunze, in Linnaea*, xx. (1847) 31.—N. Zel. ?
Cunninghamii, Benth. in Enum. Pl. Hueg. 78 = acuminatum.
cuspidatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 226.—N. Caled.
cyanantherum, A. Cunn. ex A. DC. in DC. Prod. xi. 708 = acuminatum.
Dampieri, A. Cunn. ex A. DC. l. c. = acuminatum.
debile, *R. Br. Prod.* 516.—Austral.
deserti, *A. Cunn. ex Benth. in Enum. Pl. Hueg.* 78.—Austral.
diffusum, R. Br. Prod. 515 = debile.
dulce, Benth. in Mitch. Journ. Trop. Austral. 384 = deserti.
ellipticum, R. Br. Prod. 515 = acuminatum.
euphrasioides, Hook. & Arn. Bot. Beech. Voy. 67 = Nesogenes euphrasioides.
floribundum, *A. Cunn. ex Benth. Enum. Pl. Hueg.* 78.—Austral.
glandulosum, DC. Prod. xi. 709 = serratum.
gracile, Bartl. in Lehm. Pl. Preiss. i. 350 = serratum.
humile, R. Br. Prod. 516 = parvifolium.
insulare, R. Br. l. c. = serratum.
laetum, *Forst. f. Prod.* 44.—N. Zel.
laxiflorum, *Benth. Fl. Austral.* v. 6.—Austral.
mauritianum, *A. DC. in DC. Prod.* xi. 711.—Ins. Mascar.
montanum, R. Br. Prod. 515 = acuminatum.
mucronulatum, *DC. Prod.* xi. 706.—Austral.
obscurum, *Endl. Prod. Fl. Norf.* 54.—Ins. Norfolk.
ophioglossoides, Steud. Nom. ed. II. ii. 170, sphalm. = Pogonia ophioglossoides.
oppositifolium, *R. Br. Prod.* 516.—Austral.
parvifolium, *R. Br. l. c.*—Austral.
patens, A. Cunn. ex A. DC. in DC. Prod. xi. 708 = deserti.
platycarpum, *R. Br. Prod.* 516.—Austral.
pubescens, *Forst. f. Prod.* 44.—N. Zel.
punctulatum, *Schlecht. in Linnaea*, xx. (1847) 613.—Austral.
rugulosum, F. Muell. l. c. xxv. (1852) 427 = deserti.
salsoloides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 226.—Austral.
sandwicense, *A. Gray, in Proc. Am. Acad.* vi. (1862) 52.—Ins. Sandvic.
serratum, *R. Br. Prod.* 516.—Austral.
strictum, A. Cunn. ex A. DC. in DC. Prod. xi. 708 = deserti.
subserratum, Nees, in Lehm. Pl. Preiss. i. 350 = serratum.
tasmanicum, A. DC. in DC. Prod. xi. 709 = serratum.
tenuifolium, *Forst. f. Prod.* 44.—Austral.; N. Caled.
tuberculatum, R. Br. Prod. 516 = serratum.
verrucosum, *Poir. Encyc. Suppl.* iv. 42.—Austral.
viscosum, R. Br. Prod. 516 = serratum.

MYOPSIA, Presl, Prod. Monog. Lobel. 8 (1836) = **Heterotoma**, Zucc. (Campan.).
mexicana, Presl, l. c. = H. lobelioides.

MYOPSIS, Benth. & Hook. f. Gen. ii. 553 (1876), sphalm. = praec.

MYOSANTHUS, [Dum. Anal. Fam. 49 (1829)]; Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 348 = **Stellaria**, Linn. (Caryoph.).
aquaticus, Fourr. l. c. = S. aquatica.

MYOSCHILOS, Ruiz & Pav. Prod. 41. t. 34 (1794). *SANTALACEAE*, Benth. & Hook. f. iii. 226.
oblongum, *Ruiz & Pav. Syst. Veg.* 73.—Chili.

MYOSERIS, Link, Enum. Hort. Berol. ii. 291 (1822) = **Pterotheca**, Cass. (Compos.).
urpurea, Link, l. c. = P. bifida.

MYOSOTIDIUM, Hook. Bot. Mag. t. 5137 (1859). *BORAGINEAE*, Benth. & Hook. f. ii. 849.
nobile, *Hook. l. c.*—Ins. Chatham.

MYOSOTIS, [Tourn.] Dill. ex Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 858.
ECHIOIDES, Moench, Meth. 416 (1794).
EXARRHENA, R. Br. Prod. 495 (1810).
SCORPIOIDES, Gilib. Fl. Lituan. i. 20 (1781).
SCORPIURUS, Hall. Enum. Stirp. Helv. ii. 519 (1742).
STROPHIOSTOMA, Turcz. in Bull. Soc. Nat. Mosc. (1840) 258.
abyssinica, *Boiss. & Reut. Diagn.* Ser. I. xi. 122.—Abyssin.
adpressa, Stokes, Bot. Mat. Med. i. 269 = palustre.
adulterina, Lebel, Obs. Pl. Manche, 17 = intermedia.
alba, Colla, in Mem. Acc. Torin. xxxviii. (1835) 128. t. 41 = Plagiobothrys rufescens.
albida, H. B. & K. Nov. Gen. et Sp. iii. 91 = Krynitzkia ramosa.
albida, T. Kirk, in Trans. N. Z. Inst. xvii. (1885) 224 = antarctica.
albiflora, *Banks & Soland. ex Hook. f. Fl. Antarct.* ii. 329.—Reg. Magellan.
alpestris, F. W. Schmidt, Fl. Boëm. iii. 26 = sylvatica.
alpina, Lapeyr. Hist. Abr. Pl. Pyr. 85 = sylvatica.
amoena, *Boiss. Fl. Orient.* iv. 241.—Reg. Caucas.
amoena, Schott & Kotschy, in Oestr. Bot. Wochenbl. vii. (1857) 206 = sylvatica.
anchusiformis, Griff. Notul. iv. 202 = Bothriospermum tenellum ?
annua, Moench, Enum. Pl. Hass. 81 = arvensis.
antarctica, *Hook. f. Fl. Antarct.* i. 57.—N. Zel.
aprica, Opiz, Seznam, 67 = collina.
apula, Linn. Sp. Pl. 131 = Lithospermum apulum.
arenaria, Schrad. in Schultz, Prod. Fl. Starg. Suppl. i. 12 = stricta.
aretioides, Cham. in Linnaea, iv. (1829) 443 = Eritrichium nanum.
arvensis, Hill, Veg. Syst. vii. 55 = versicolor.
arvensis, *Lam. Fl. Fr.* ii. 213; *Willd. Fl. Berol. Prod.* 77.—Europ.; As.; Am. bor.
arvensis, Link, Enum. Hort. Berol. i. 164 = collina.
arvensis, Reichb. Fl. Germ. Excurs. 340 = stricta.
arvensis, Roth, Abh. ii. 20 = intermedia.
aspera, Lamotte, Prod. Fl. Centr. Fr. ii. 538 = palustris.
aspera, Schur, Enum. Pl. Transs. 476 = arvensis.
australis, *R. Br. Prod.* 415.—Austral.; N. Zel.
azorica, *H. C. Wats. ex Hook. Bot. Mag.* t. 4122.—Ins. Azor.
baeotica, Reut. ex Boiss. Pl. Or. Nov. Dec. ii. 9 = idaea.
Balbisiana, Jord. Pugill. Pl. Nov. 128 = versicolor.
barbata, Bieb. Fl. Taur. Cauc. i. 121 = Echinospermum barbatum.
Barrelieri, Bess. ex Steud. Nom. ed. I. 543 = Anchusa Barrelieri.
borbonica, Lam. Tabl. Encyc. i. 396 = Cynoglossum borbonicum.
brachypoda, *Gren. in Mém. Soc. Emul. Doubs*, Sér. III. ii. (1858) 425.—Europ. ?
bracteata, A. Br. ex A. Rich. Tent. Fl. Abyss. ii. 89, in obs. = collina.
cadmea, Boiss. Diagn. Ser. I. xi. 122 = idaea.
caespitosa, Auct. vide cespitosa.
caespitosa, DC. Prod. x. 135, partim = laxa.
caespitosa, Tenore, ex Nym. Syll. 519 = sylvatica.
californica, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 42 = Krynitzkia californica.
campestris, Godet, ex DC. Prod. x. 108, in nota = sylvatica.
capitata, *Hook. f. Fl. Antarct.* i. 56.—N. Zel.; Ins. Dom. Auckl.
cespitosa, *Schultz, Prod. Fl. Starg. Suppl.* i. 11.—Reg. bor.
chinensis, *DC. Prod.* x. 106.—China.
Chorisiana, Cham. in Linnaea, iv. (1829) 444 = Krynitzkia Chorisiana.
Chorisiana, Lehm. in Hook. Fl. Bor. Am. ii. 83 = Krynitzkia Scouleri.
chrysantha, Welw. ex Nym. Consp. 520 = versicolor.
ciliata, Rudolph, in Mém. Acad. Pétersb. (1803–6) 352. t. 11 = Eritrichium pectinatum.
ciliata, Turcz. ex Ledeb. Fl. Ross. iii. 151 = Eritrichium sericeum.
clavata, Ledeb. ex Spreng. Syst. i. 553 = Trigonotis peduncularis.
collina, Bab. in Journ. Linn. Soc. xi. (1871) 319 = arvensis.
collina, *Hoffm. Deutschl. Fl.* ed. I. 61.—Europ.
commutata, Roem. & Schult. Syst. iv. 102 = caespitosa.

MYOSOTIS :—

compressa, Turcz. ex Ledeb. Fl. Ross. iii. 154 (= *Anoplocaryum compressum*).—Sibir. baical.
concinna, *Cheesem. in Trans. N. Z. Inst.* xvii. (1885) 235.—N. Zel.
condylophora, Blume, ex Ledeb. Fl. Ross. iii. 153 = Trigonotis peduncularis.
coronaria, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 350 = palustris.
corymbosa, *Ruiz & Pav. Fl. Per.* ii. 5.—Chili.
cretica, Boiss. & Heldr. Diagn. Ser. I. xi. 121 = pusilla.
cretica, Unger, ex Nym. Consp. 519 = idaea.
cyanea, Boiss. & Heldr. ex Orph. in Bull. Congr. Bot. St. Pétersb. 1869 (1870) 129 = sylvatica.
cymosa, *Nutt. ex Hook. Kew Journ.* iii. (1851) 294.—Am. bor.
cynoglossoides, Lam. Tabl. Encyc. i. 396 = Echinospermum cynoglossoides.
davurica. Pall. ex Roem. & Schult. Syst. iv. 774 = Krynitzkia obovata.
debilis, *Pomel, Nouv. Mat. Fl. Atl.* 298.—Alger.
decumbens, Host, Fl. Austr. i. 228 = sylvatica.
deflexa, Wahlenb. in Vet. Acad. Handl. Stockh. xxxi. (1810) 113. f. 4 = Echinospermum deflexum.
densiflora, C. Koch, in Linnaea, xix. (1847) 20 = sylvatica.
dentata, *Jensch. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 47, nomen.—Reg. Transcauc.
diffusa, Dougl. ex Lehm. Pugill. ii. 23 = Echinospermum diffusum.
discolor, Pers. in Murr. Syst. ed. XV. 190, in obs. = versicolor.
dissitiflora, *Baker, in Gard. Chron.* (1868) 599.—Helvet.
drabaefolia, Turcz. ex Steud. Nom. ed. II. ii. 170 = Eritrichium villosum.
dubia, Arrond. Cat. Pl. Morb. 70 = versicolor.
Dumortieri, Thielens, in Bull. Soc. Bot. Belg. vii. (1868) 86 = palustris.
echinata, Thunb. Fl. Cap. ii. 352 = Echinospermum cynoglossoides.
echinophora, Pall. Reise, iii. 717 = Echinospermum minimum.
echinophora, Wahlenb. in Vet. Acad. Handl. Stockh. xxxi. 114 = Echinospermum Redowskii.
echinosperma, *Fl. Corc. in Ionios Anthol.* iv. 960.—Ins. Corcyr.
elongata, *Strobl, in Flora*, lxvii. (1884) 635.—Sicil.
exscapa, Steud. Nom. ed. I. 543 = Eritrichium nanum.
fallacina, Jord. in Bor. Fl. Centr. Fr. ed. III. ii. 463 = versicolor.
filiformis, Schleich. Cat. Pl. Helv. ed. IV. 23 = collina.
flaccida, Dougl. ex Lehm. Pugill. ii. 22 = Krynitzkia oxycarya.
foliosa, *Lehm. Asperif.* 99.—Am. austr.
Forsteri, *Lehm. l. c.* 95.—N. Zel.
fruticosa, Haenke, in Abh. Boehm. Ges. Wiss. ii. (1786) 46, nomen = Lithospermum purpurocaeruleum.
fruticosa, Linn. Mant. ii. 201 = Heliotropium paniculatum.
fulva, Hook. & Arn. Bot. Beech. Voy. 38 = Plagiobothrys rufescens.
fulva, Hook. & Arn. l. c. 369 = Plagiobothrys nothofulvus.
geniculata, Schur, Enum. Pl. Transs. 474 = palustris.
glomerata, Nutt. Gen. Am. i. 112 = Krynitzkia glomerata.
gracilis, Ruiz & Pav. Fl. Per. ii. 5 = Pectocarya linearis.
gracillima, Losc. Ser. Inconf. Pl. Arag. 72 = collina.
graminifolia, *DC. Prod.* x. 110.—Afr. austr.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 90. t. 99.—Ecuador,
granulosa, *Ruiz & Pav. Fl. Per.* ii. 5.—Peruv.
Hacquetii, D. Dietr. Syn. Pl. i. 604 = Eritrichium nanum.
hamosa, Stokes, Bot. Mat. Med. i. 269 = arvensis.
Hectori, Hook. f. Handb. N. Zeal. Fl. 193 = pulvinaris.
heterodoxa, *Pomel, Nouv. Mat. Fl. Atl.* 298.—Alger.
heteropoda, Trautv. in Act. Hort. Petrop. ii. (1873) 479 = intermedia.
hispida, C. Koch, in Linnaea, xvii. (1843) 303 = stricta.
hispida, Schlecht. in Ges. Naturf. Fr. Berl. Mag. viii. (1814) 230 = collina.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 174.—Reg. Himal.
hortensis, Decne. in Deless. Ic. Sel. v. 42. t. 99 = Myosotidium nobile.

MYOSOTIS :—

humilis, *Ruiz & Pav. Fl. Per.* ii. 5.—Peruv.
idaea, *Boiss. & Heldr. Diagn.* Ser. I. xi. 121.—Graecia ; As. Min.
incana, Turcz. in Bull. Soc. Nat. Mosc. (1838) 97 ; et xxiii. (1850) i. 515 = Eritrichium pectinatum.
incrassata, Guss. Fl. Sic. Syn. i. 214 = pusilla.
inflexa, Engelm. in Am. Journ. Sc. Ser. I. xlvi. (1844) 98 = verna.
intermedia, Lechart. ex A. DC. in DC. Prod. x. 109 = versicolor.
intermedia, *Link, ex Schultz, Prod. Fl. Starg. Suppl.* i. 12 ; *et Link, Enum. Hort. Berol.* i. 164.—Europ. ; As. bor. ; Oriens.
involucrata, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 253.—Reg. Caucas.
javanica, Sw. ex Lehm. in Neue Schr. Naturf. Ges. Halle, iii. 11. (1817) 21 = Echinospermum javanicum.
Kiesenwetteri, Heldr. ex Boiss. Fl. Orient. iv. 236 = idaea.
lactea, Boenn. Prod. Fl. Monast. 321 (Index) = sylvatica.
laeta, *Cheesem. in Trans. N. Z. Inst.* xvii. (1885) 236.—N. Zel.
Lappula, Bouton, ex DC. Prod. x. 152 = Cynoglossum Rochelia.
Lappula, Linn. Sp. Pl. 131 = Echinospermum Lappula.
latifolia, Boiss. & Kotschy, ex Boiss. Pl. Or. Nov. Dec. ii. 9 = platyphylla.
latifolia, Poir. Encyc. Suppl. iv. 45 = sylvatica.
laxa, Hort. ex Steud. Nom. ed. II. ii. 171 = sparsiflora.
laxa, *Lehm. Asperif.* 83.—Reg. bor.
laxiflora, Reichb. in Sturm, Deutschl. Fl. Heft 42 = palustris.
Lebelii, Gren. & Godr. Fl. Fr. ii. 532 = intermedia.
leucophaea, Dougl. ex Lehm. Pugill. ii. 22 = Krynitzkia leucophaea.
linearis, Colla, in Mem. Acc. Torin. xxxviii. (1835) 129. t. 42. f. 2 = Krynitzkia linearis.
lingulata, Lehm. Asperif. 110 ; Roem. & Schult. Syst. iv. 780, in obs. = cespitosa.
lithospermifolia, Hornem. Hort. Hafn. i. 173 = sylvatica.
litoralis, *Stev. ex Fisch. Cat. Hort. Gorenk.* (1812) 27.—Graecia ; As. Min.
littoralis, Heldr. ex DC. Prod. x. 106, in nota = pusilla.
longiflora, *Boeber, ex Georgi, Beschr. Russ. Reich.* III. iv. 750 (Quid ?).—Rossia.
longiflora, Wall. Cat. n. 929 = Mertensia racemosa.
lutea, Lam. Fl. Fr. ii. 282 = Lithospermum apulum.
lutea, Pers. Syn. i. 156 = versicolor.
Lyallii, *Hook. f. Fl. N. Zel.* i. 202.—N. Zel.
macrophylla, Adam, in Web. & Mohr, Beitr. i. 46 ; Bieb. Fl. Taur. Cauc. i. 119 = Anchusa myosotidiflora.
macrosperma, Engelm. in Am. Journ. Sc. Ser. I. xlvi. (1844) 98 = verna.
marginata, Bieb. Fl. Taur. Cauc. i. 120 = Echinospermum marginatum.
maritima, *Hochst. ex Seub. Fl. Azor.* 37.—Ins. Azor.
maritima, Ledeb. Fl. Ross. iii. 144 = cespitosa.
micrantha, Guss. Fl. Sic. Prod. i. 207 = sicula.
micrantha, Pall. ex Lehm. in Neue Schr. Naturf. Ges. Halle, iii. 11. (1817) 24 = stricta.
microcarpa, Wall. Cat. n. 928 = Trigonotis microcarpa.
microphylla, DC. Prod. x. 115, sphalm. = *macrophylla*.
microsepala, Schott & Kotschy, ex Boiss. Pl. Or. Nov. Dec. ii. 9 = olympica.
minutiflora, Boiss. & Reut. Pugill. Pl. Nov. 80 = stricta.
modesta, Schott & Kotschy, ex Boiss. Pl. Or. Nov. Dec. ii. 9 = olympica.
moltkioides, Benth. in Royle, Illustr. Bot. Himal. 305 = Mertensia moltkioides.
montana, Bess. Fl. Galic. i. 142 = sylvatica.
montana, Pourr. ex Steud. Nom. ed. I. 543 = palustris.
monticola, *Wender. ex Steud. l. c.* ed. II. ii. 171, nomen.—Hab. ?
multicaulis, Turcz. in Bull. Soc. Nat. Mosc. (1838) 97 = Eritrichium sericeum, jenisseense.
multicaulis, Wall. Cat. n. 7020 = Trigonotis multicaulis.
multiflora, Mérat, Rev. Fl. Par. 204 = cespitosa.
muricata, Hook. & Arn. Bot. Beech. Voy. 369 = Krynitzkia muriculata.
nana, Miégev. in Bull. Soc. Bot. Fr. x. (1863) 28 = intermedia.

MYOSOTIS :—

nana, Sibth. & Sm. Fl. Graec. Prod. i. 112 = sylvatica.
nana, Vill. Prosp. 21 ; Hist. Pl. Dauph. ii. 459 = Eritrichium nanum.
nemorosa, Bess. Enum. Pl. Volh. 52 = palustris.
nemorosa, Bunge, ex A. DC. in DC. Prod. x. 105 = cespitosa.
nemorosa, Martr. Fl. Tarn. 492 = sylvatica ?
oblongata, Link, in Buch, Beschr. Canar. Ins. 144 = sylvatica.
obovata, Ledeb. Fl. Alt. i. 190 = Krynitzkia obovata.
obtusa, Waldst. & Kit. Pl. Rar. Hung. i. 103 = Anchusa Barrelieri.
odorata, Poir. Encyc. Suppl. iv. 44 = sylvatica.
oenesia, Heldr. ex Boiss. Fl. Orient. iv. 237 = pusilla.
olympica, *Boiss. Diagn.* Ser. I. iv. 50.—As. Min.
oraria, *Dum. in Bull. Soc. Bot. Belg.* vii. (1868) 350.—Europ.
orientalis, Schenk, Pl. Sp. Aegypt. 26 = Anchusa neglecta.
ovalifolia, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 9 = Trigonotis ovalifolia.
pallens, Wall. Cat. n. 7019 = sylvatica.
palustris, Benth. in Royle, Illustr. Bot. Himal. 305 = cespitosa.
palustris, *Lam. Fl. Fr.* ii. 283 ; *With. Bot. Arr. Brit. Pl.* ed. III. ii. 225.—Europ. ; As. et Am. bor.
pauciflora, Ledeb. in Mém. Acad. Pétersb. v. (1812) 517 = Eritrichium rupestre.
pauciflora, Vitm. Summa Pl. i. 381 = Eritrichium nanum.
pectinata, Pall. Reise, iii. 717 = Eritrichium pectinatum.
pectinata, Turcz. ex Ledeb. Fl. Ross. iii. 155 = Echinospermum deflexum.
pectinata, Turcz. in Bull. Soc. Nat. Mosc. xxiii. (1850) i. 513 = Eritrichium villosum.
peduncularis, Trev. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 147 = Trigonotis peduncularis.
pedunculata, Roem. & Schult. Syst. iv. 104 = Trigonotis peduncularis.
perennis, Moench, Hass. 81 = palustris.
perpusilla, *Pomel, Nouv. Mat. Fl. Atl.* 297.—Alger.
petiolata, *Hook. f. Fl. N. Zel.* i. 202.—N. Zel.
phoenicea, Ehrenb. ex Boiss. Fl. Orient. iv. 157 = Anchusa neglecta.
platyphylla, *Boiss. Pl. Or. Nov. Dec.* ii. 9 ; *et Fl. Orient.* iv. 237.—Armenia.
pontica, C. Koch, in Linnaea, xxii. (1849) 642 = sylvatica.
procumbens, Colla, in Mem. Acc. Torin. xxxviii. (1835) 130 = Plagiobothrys procumbens.
propinqua, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 163.—As. Min. ; Persia.
Pseudo-sylvatica, Schur, Enum. Pl. Transs. 475 = arvensis.
pulvinaris, *Hook. f. Handb. N. Zeal. Fl.* 193.—N. Zel.
pusilla, *Loisel. in Desv. Journ. Bot.* ii. (1809) 260. t. 8. f. 2.—Reg. Mediterr.
pusilla, Orph. ex Nym. Consp. 519 = litoralis.
pygmaea, Bertol. Misc. Bot. i. 13 = collina.
pygmaea, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 334.—N. Zel.
pyrenaica, Pourr. in Mém. Acad. Toul. iii. (1788) 322 = sylvatica.
racemosa, Royle, Illustr. Bot. Himal. 305 = Mertensia racemosa.
radicans, Turcz. in Bull. Soc. Nat. Mosc. (1840) 258 = Trigonotis radicans.
ramosissima, Rochel, ex Schult. Oesterr. Fl. ed. II. i. 366, in obs. = collina.
Redowskii, Hornem. Hort. Hafn. i. 174 = Echinospermum Redowskii.
refracta, *Boiss. Voy. Bot. Espagne*, ii. 433. t. 125.—Hispan. ; Graecia ; Syria ; Turkest.
Rehsteineri, Wartm. ex Reut. in Compt.-Rend. Soc. Hallér. (1854–56) 115 = cespitosa.
repens, G. Don, Gen. Syst. iv. 344 = palustris.
rigida, *Pomel, Nouv. Mat. Fl. Atl.* 297.—Alger.
robusta, *D. Don, Prod. Fl. Nep.* 101.—Nepal.
rotundifolia, Wall. Cat. n. 930 = Trigonotis rotundifolia.
rupestris, Pall. Reise, iii. 716 = Eritrichium rupestre.
rupestris, Willd. ex Ledeb. Fl. Ross. iii. 151 = Eritrichium sericeum.
rupicola, Mert. in Linnaea, v. (1830) 62 = Eritrichium nanum.
rupicola, Sm. Engl. Bot. t. 2559 = sylvatica.

MYOSOTIS:—
saxatilis, Pall. Tabl. Taur. 47 = Echinospermum barbatum.
saxosa, *Hook. f. Fl. N. Zel.* i. 202.—N. Zel.
scorpioides, Krock. Fl. Siles. i. 275 = sparsiflora.
scorpioides, Linn. Sp. Pl. 131 = arvensis, palustris, sylvatica.
Scorpiurus, Reichard, Fl. Moeno-Franç. i. 31 = *scorpioides*, Linn.
Scouleri, Hook. & Arn. Bot. Beech. Voy. 370, in obs. = Krynitzkia Scouleri.
semiamplexicaulis, *DC. Prod.* x. 110.—Afr. austr.
sepium, Bess. in Cat. Hort. Cracov. (1808) = sparsiflora.
sericea, Lehm. Asperif. 98 = Eritrichium sericeum.
sericea, *Nutt. ex Hook. Kew Journ.* iii. (1851) 295.—Am. bor.
sicula, *Guss. Fl. Sic. Syn.* i. 214.—Reg. Mediterr.
Soleirolii, Godr. in Gren. & Godr. Fl. Fr. ii. 534 = sylvatica.
sparsiflora, Kirschl. ex Mutel, Fl. Fr. ii. 428, 446 = cespitosa.
sparsiflora, *Mikan, in Hoppe, Taschenb.* (1807) 74.—Europ.; Oriens; As. bor.
spathulata, Poir. Encyc. Suppl. iv. 45 = australis.
spathulata, A. Rich. Ess. Fl. N. Zél. 198 = Forsteri.
spatulata, *Forst. f. Prod.* 12.—N. Zel.
speciosa, *Pomel, Nouv. Mat. Fl. Atl.* 296.—Alger.
speluncicola, Schott, ex Boiss. Fl. Orient. iv. 240 = stricta.
spinocarpos, Vahl, Symb. Bot. ii. 32 = Echinospermum Vahlianum.
squarrosa, Bieb. Fl. Taur. Cauc. i. 120 = Echinospermum patulum.
squarrosa, Retz. Obs. ii. 9 = Echinospermum Lappula.
staminea, Lehm. Pl. Preiss. i. 348 = australis.
stolonifera, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 351 = cespitosa.
stricta, Friedrichs. ex Nym. Consp. 519 = idaea.
stricta, A. Gray, Man. Bot. N. U. St. ed. I. 338 = verna.
stricta, Lehm. ex Unger, Reise Griechenl. (1862) 127 = littoralis.
stricta, *Link, ex Roem. & Schult. Syst.* iv. 104; *et Enum. Hort. Berol.* i. 164.—Europ.; Oriens; As. bor.
strigosa, Schlecht. in Ges. Naturf. Fr. Berl. Mag. viii. (1818) 233; Willd. ex Roem. & Schult. Syst. iv. 783 = Cynoglossum borbonicum.
strigulosa, Reichb. in Sturm, Deutschl. Fl. Heft 42 = palustris.
suaveolens, *Poir. Encyc. Suppl.* iv. 44.—Austral.
suaveolens, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 176 = sylvatica.
subvillosa, *Ledeb. ex Steud. Nom.* ed. II. ii. 171, nomen.—Sibir.
suffruticosa, Torr. in Ann. Lyc. N. York, ii. (1828) 225 = Krynitzkia Jamesii.
sylvatica, *Hoffm. Deutschl. Fl.* ed. I. 61.—Reg. bor. temp.; Oriens.
sylvatica, H. Mart. Fl. Mosq. 32 = intermedia.
tenella, C. A. Mey. ex Ledeb. Fl. Ross. iii. 148 = sparsiflora.
tenella, Nutt. in Hook. Kew Journ. iii. (1851) 295 = Plagiobothrys tenellus.
tenuiflora, Viv. Fl. Lib. Spec. 9. t. 1. f. 2 = Lithospermum tenuiflorum.
terglovensis, Hacq. Pl. Alp. Carn. 12. t. 2. f. 6 = Eritrichium nanum.
texana, Hook. Kew Journ. iii. (1851) 295 = Krynitzkia texana.
Traillii, T. Kirk, in Trans. N. Z. Inst. xvi. (1884) 373 = antarctica.
transsylvanica, Porc. in Magyar, Növenyt. Lapok, ix. (1885) 130 = sylvatica.
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 194.—N. Zel.
ucrainica, *Czern. in Bull. Soc. Nat. Mosc.* xviii. (1845) II. 133, in adnot.—Ross.
uliginosa, Schrad. ex Mert. & Koch, Deutschl. Fl. ii. 42 = cespitosa.
umbrata, Auct. ex Mart. & Koch, l. c. 47 = intermedia.
umbrosa, Nym. Consp. 520 = intermedia.
umbrosa, Schleich. ex Reichb. Fl. Germ. Excurs. 341 = sylvatica.
uncata, *Boiss. & Bal. Diagn.* Ser. II. iii. 139.—Cilicia.
ungulata, *G. Don, ex Loud. Hort. Brit. Suppl.* iii. 591, nomen—Sibiria.

MYOSOTIS:—
uniflora, *Hook. f. Handb. N. Zeal. Fl.* 192.—N. Zel.
variabilis, Angel. ex Maly, in Flora, xxiii. (1840) 607 = sylvatica.
verna, *Nutt. Gen. Am.* ii. *Addend.*—Am. bor.
verna, Opiz, in Flora, v. (1822) 267 = stricta.
vernalis, Willd. ex Roem. & Schult. Syst. iv. 778, in obs. = litoralis.
versicolor, Lehm. in Hook. Fl. Bor. Am. ii. 81 = verna.
versicolor, *Sm. Engl. Bot.* sub t. 2558.—Europ.
villosa, Cham. in Linnaea, iv. (1829) 442 = Eritrichium nanum.
villosa, Ledeb. in Mém. Acad. Pétersb. v. (1812) 516 = Eritrichium villosum.
virginiana, Linn. Sp. Pl. 131 = Echinospermum virginicum.
Welwitschii, *Boiss. & Reut. Diagn.* Ser. II. iii. 138.—Hispan.
zeylanica, Sw. ex Lehm. in Neue Schr. Naturf. Ges. Halle, iii. II. (1817) 20 = Echinospermum denticulatum.

MYOSOTIS, Tourn. ex Moench, Meth. 224 (1794) = **Cerastium**, Dill. (Caryoph.).
connata, Moench, l. c. = C. perfoliatum.
dichotoma, Moench, l. c. 225 = C. dichotomum.
lanata, Moench, l. c. Suppl. 308 = C. tomentosum.
linearis, Moench, Meth. 225 = C. arvense.
repens, Moench, l. c. Suppl. 308 = C. adscendens.
semidecandra, Moench, Meth. 225 = C. semidecandrum.
vulgaris, Moench, l. c. 224 = C. glomeratum.

MYOSOTON, Moench, Meth. 225 (1794) = **Stellaria**, Linn. (Caryoph.).
aquaticum, Moench, l. c. = S. aquatica.

MYOSPYRUM, Lindl. Veg. Kingd. ed. III. 617 (1853), sphalm. = MYXOPYRUM, Blume = **Glycosmis**, Correa (Rutac.).

MYOSTEMMA, Salisb. Gen. Pl. Fragm. 135 (1866) = **Hippeastrum**, Herb. (Amaryll.).
advena, Salisb. l. c. = H. advena.

MYOSTOMA, Miers, in Trans. Linn. Soc. xxv. (1866) 474. t. 57 = **Thismia**, Griff. (Burmann.).
hyalinum, Miers, l. c.—Bras.

MYOSURANDRA, Baill. Adansonia, ix. (1870) 325. *HAMAMELIDEAE.*
moschata, *Baill. l. c.* 329.—Madag.

MYOSUROS, Adans. Fam. ii. 459 (1763) = seq.

MYOSURUS, Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
alopecuroides, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 278.—Calif.
apetalus, C. Gay, Fl. Chil. i. 31 = aristatus.
aristatus, *Benth. in Hook. Lond. Journ. Bot.* vi. (1847) 459 [bis].—N. Zel.; Am. austr.
australis, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 6 = minimus.
Cauda-muris, Neck. Delic. Gallo-Belg. i. 161 = minimus.
cupulatus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Calif.
europaea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 723 = minimus.
minimus, *Linn. Sp. Pl.* 284.—Europ.; Am. bor.
scaposus, Gilib. Exercit. i. 376 = minimus.
sessilis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Am. bor. occ.
Shortii, Rafin. in Am. Journ. Sc. Ser. I. i. (1819) 379 = minimus.

MYOTOCA, Griseb. in Abh. Goett. vi. (1856) 129 = **Collomia**, Nutt. (Polemon.).

MYOXANTHUS, Poepp. & Endl. Nov. Gen. et Sp. i. 50. t. 88 (1835) = **Pleurothallis**, R. Br. (Orchid.).
monophyllus, Poepp. & Endl. l. c. = P. Raymondi.

MYRACRODRUON, Allem. in Trab. Comm. Sc. Expl. Ceará; Secç. Bot. 3. tt. 1, 2 (1862) = **Astronium**, Jacq. (Anacard.).
concinnus, Engl. in Mart. Fl. Bras. xii. II. 401 = A. concinnum.

MYRACRODRUON:—
macrocalyx, Engl. in Mart. Fl. Bras. xii. II. 402 = Astronium macrocalyx.
Urundeuva, Allem. in Trab. Comm. Sc. Expl. Ceará; Secç. Bot. 3. tt. 1, 2 = A. Urundeuva.

MYRCEUGENIA, Berg, in Linnaea, xxvii. (1854) 131 = **Myrtus**, Linn.
Bridgesii, Berg, l. c. xxx. (1859–60) 671 = Eugenia Bridgesii.
camphorata, Berg, l. c. xxvii. (1854) 134 = Myrtus camphorata.
chilensis, Berg, l. c. 132 = Myrtus reticulata, *Poepp.*
correaefolia, Berg, l. c. xxx. (1859–60) 670.—Chili.
exsucca, Berg, l. c. 671 = Eugenia exsucca.
Lechleriana, Berg, l. c. xxvii. (1854) 133 = Eugenia exsucca.
Luma, Berg, l. c. xxx. (1859–60) 671 = Myrtus fernandeziana.
montevidiensis, Berg, in Mart. Fl. Bras. xiv. I. 211.—Reg. Argent.
myrcoides, Berg, in Linnaea, xxvii. (1854) 134 = Eugenia myrcioides.
myrtoides, Berg, in Mart. Fl. Bras. xiv. I. 211.—Reg. Argent.
obtusa, Berg, in Linnaea, xxx. (1859–60) 669 = Eugenia Raran.
ovata, Berg, l. c. 670 = Eugenia ovata.
Pitra, Berg, l. c. 671 = Eugenia Pitra.
planipes, Berg, in Bot. Zeit. xvi. (1858) 250 = Eugenia planipes.
Sellowiana, Berg, in Mart. Fl. Bras. xiv. I. 212.—Bras.
stenophylla, Berg, in Linnaea, xxx. (1859–60) 670 = Eugenia stenophylla.

MYRCIA, DC. ex Guill. in Dict. Class. Hist. Nat. xi. 378, 401 (1826). *MYRTACEAE*, Benth. & Hook. f. i. 716.
AGUAVA, Rafin. Sylva Tellur. 107 (1838).
AULOMYRCIA, Berg, in Linnaea, xxvii. (1854) 35.
CALYCAMPE, Berg, l. c. 129 (1854).
CALYPTROMYRCIA, Berg, l. c. 34 (1854).
CERQUIERIA, Berg, l. c. 5 (1854).
GOMIDESIA, Berg, l. c. 6 (1854).
GOMIDEZIA, Benth. & Hook. f. Gen. i. 716 (1865).
acris, DC. Prod. iii. 243 = Pimenta acris.
acuminata, *DC. l. c.* 256.—N. Granat.
acuminatissima, *Berg, in Mart. Fl. Bras.* xiv. I. 167.—Bras.
acutata, *Berg, l. c.* 175.—Bras.
acutiloba, *Berg, l. c.* 189.—Bras.
affinis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 307.—Bras.
alagoënsis, *Berg, in Mart. Fl. Bras.* xiv. I. 165.—Bras.
albotomentosa, *DC. Prod.* iii. 254.—Bras.
alternifolia, *Miq. in Linnaea*, xxii. (1849) 534.—Bras.
amazonica, *DC. Prod.* iii. 250.—Bras.
ambigua, *DC. l. c.* 252.—Bras.
amplexicaulis, *Hook. f. Bot. Mag.* t. 5790.—Bras.
anacardiaefolia, *Gardn. in Hook. Lond. Journ. Bot.* (1843) 354.—Bras.
anceps, *Berg, in Mart. Fl. Bras.* xiv. I. 186.—Bras.
andromedoides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 323.—Bras.
angusta, *G. Don, Gen. Syst.* ii. 845.—Bras.
anomala, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 328.—Bras.
arborescens, *Berg, in Mart. Fl. Bras.* xiv. I. 200.—Bras.
aromatica, Schlecht. in Linnaea, xiii. (1839) 415 = Calyptranthes Schiedeana, Schlechtendaliana.
Australasiae, F. Muell. Rep. Burdek. Exp. 7 = Memecylon pauciflorum.
Ayresiana, *Berg, in Linnaea*, xxvii. (1854) 100.—Venezuel.
Balbisiana, *DC. Prod.* iii. 243.—Ins. Guadalup.
banisteriaefolia, *DC. l. c.* 246.—Bras.
barrensis, *Berg, in Mart. Fl. Bras.* xiv. I. 187.—Bras.
bella, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 322.—Bras.
Berberis, *DC. Prod.* iii. 254.—Bras.
Berberis, Schau. in Linnaea, xxi. (1848) 273 = Kegeliana.
Bergiana, *Berg, in Mart. Fl. Bras.* xiv. I. 194.—Bras.

MYRCIA:—
Billardiana, *DC. Prod.* iii. 255.—Mexic.
bracteata, *DC. l. c.* 245.—Am. austr.
bracteolaris, *DC. l. c.*—Guian.
Brandami, *Berg, in Mart. Fl. Bras.* xiv. i. 164.—Bras.
Bredemeyeriana, *Berg, in Linnaea,* xxvii. (1854) 85.—Venezuela.
Browniana, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 354.—Bras.
brunea, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 306.—Bras.
bullata, *Berg, in Mart. Fl. Bras.* xiv. i. 153.—Bras.
buxifolia, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 101.—Bras.
Camaraeana, *DC. Prod.* iii. 251.—Bras.
Cambessediana, *Berg, in Mart. Fl. Bras.* xiv. i. 202.—Bras.
campestris, *DC. Prod.* iii. 247.—Bras.
cancellata, *Berg, in Mart. Fl. Bras.* xiv. i. 154.—Bras.
canescens, *Berg, l. c.* 206.—Bras.
capitata, *Berg, l. c.* 154.—Bras.
capivarhyensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 328.—Bras.
caracasana, *Klotzsch, ex Berg, in Linnaea,* xxvii. (1854) 92.—Venezuela.
cardiaca, *Berg, in Mart. Fl. Bras.* xiv. i. 204.—Bras.
cardiophylla, *Reichardt, in Verh. Zool.-Bot. Ges. Wien,* xxxiii. (1884) 324.—Bras.
carnea, *DC. Prod.* iii. 245.—Guiana.
cassinioides, *DC. l. c.* 249.—Bras.
Catharinae, *Berg, in Mart. Fl. Bras.* xiv. i. 176.—Bras.
chilensis, *Berg, in Linnaea,* xxvii. (1854) 99.—Chili.
ciarensis, *Berg, in Mart. Fl. Bras.* xiv. i. 178.—Bras.
clusiaefolia, *DC. Prod.* iii. 255.—Am. austr.
clusiaefolia, Miq. in Linnaea, xxii. (1849) 533 = Calyptranthes clusiaefolia.
coccolobaefolia, *DC. Prod.* iii. 255.—N. Granat.
Comete, Steud. Nom. ed. II. ii. 177 in syn. sphalm. = M. Coumeta.
communis, *Berg, in Mart. Fl. Bras.* xiv. i. 183.—Bras.
complicata, *DC. Prod.* iii. 255.—Am. austr.
corcovadensis, *Berg, in Mart. Fl. Bras.* xiv. i. 177.—Bras.
cordata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 330.—Bras.
cordiaefolia, *DC. Prod.* iii. 248.—Bras.
cordifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 205.—Bras.
coriacea, *DC. Prod.* iii. 243.—Ind. occ.
corymbosa, *DC. l. c.* 252.—Bras.
costa-ricensis, *Berg, in Linnaea,* xxvii. (1854) 104.—Costa Rica.
costata, *DC. Prod.* iii. 252.—Bras.
Coumeta, *DC. l. c.* 245.—Guiana.
coumetoides, *Berg, in Linnaea,* xxvii. (1854) 102.—Ind. occ.
crassicaulis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 311.—Bras.
crassinervia, *DC. Prod.* iii. 245.—Guiana.
cucullata, *Berg, in Linnaea,* xxvii. (1854) 97.—N. Granat.
cujabensis, *Berg, in Mart. Fl. Bras.* xiv. i. 161.—Bras.
curatellaefolia, *DC. Prod.* iii. 253.—Bras.
daphnoides, *DC. l. c.* 246.—Bras.
dasyblasta, *Berg, in Mart. Fl. Bras.* xiv. i. 207.—Bras.
dealbata, *DC. Prod.* iii. 254.—Bras.
debilis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 326.—Bras.
decorticans, *DC. Prod.* iii. 252.—Bras.
deflexa, *DC. l. c.* 244.—Ins. S. Doming.
delicatula, Steud. Nom. ed. II. ii. 177, in syn. sphalm. = Eugenia delicatula.
detergens, *Miq. in Linnaea,* xxii. (1849) 795.—Bras.
discolor, *Berg, l. c.* xxvii. (1854) 111.—Costa Rica.
divaricata, *DC. Prod.* iii. 243.—Ind. occ.
divergens, *DC. l. c.* 245.—Guiana.
Doniana, *Berg, in Mart. Fl. Bras.* xiv. i. 518.—Bras.
Duchassaingiana, Berg, in Linnaea, xxvii. (1854) 88 = ferruginea, *DC.*
duriuscula, *Mart. ex DC. Prod.* iii. 253.—Bras.
elaeodendra, *DC. l. c.* iii. 250.—Bras.
elegans, *DC. l. c.* 251.—Bras.
elliptica, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 352.—Bras.
ellipticifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 312.—Bras.

MYRCIA:—
elongata, *Berg, in Mart. Fl. Bras.* xiv. i. 159.—Bras.
eriocalyx, *DC. Prod.* iii. 247.—Bras.
eriopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = seq.
Eriopus, *DC. Prod.* iii. 255.—Bras.
Eriopus, Mart. ex DC. l. c. 257 = Calyptranthes eriopoda.
erythroxylon, *Berg, in Mart. Fl. Bras.* xiv. i. 173.—Bras.
eugenioides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 302.—Bras.
eximia, *DC. Prod.* iii. 248 —Bras.
exsucca, *DC. l. c.* 247.—Bras.
fallax, *DC. l. c.* 244.—Guiana.
fascicularis, *Berg, in Linnaea,* xxvii. (1854) 87.—Peruv.
Felisberti, Berg, in Mart. Fl. Bras. xiv. i. 562 = involucrata.
fenestrata, *DC. Prod.* iii. 251.—Bras.
Fenzliana, *Berg, in Mart. Fl. Bras.* xiv. i. 196.—Bras.
ferruginea, *DC. Prod.* iii. 245.—Am. trop.
ferruginea, *G. Don, Gen. Syst.* ii. 845.—Bras.
floribunda, *Miq. in Linnaea,* xxii. (1849) 534.—Bras.
florida, *Lem. Jard. Fleur.* iii. (1852–53) *Misc.* 100.—Bras.
formosiana, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 299 = M. Hilariana.
formosiana, *DC. Prod.* iii. 255.—Bras.
friburgensis, *Berg, in Mart. Fl. Bras.* xiv. i. 161.—Bras.
Gardneriana, *Berg, l. c.* 184.—Bras.
garopabensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 324.—Bras.
gemmiflora, *Berg, in Mart. Fl. Bras.* xiv. i. 190.—Bras.
Gestasiana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 303.—Bras.
glauca, *Cambess. l. c.* 318.—Bras.
Gollmeriana, *Berg, in Linnaea,* xxix. (1857–58) 221.—Venezuela.
goyazensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 305.—Bras.
graciliflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 185.—Guian.
gracilis, *Berg, in Mart. Fl. Bras.* xiv. i. 174.—Bras.; N. Granat.
grandifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 298.—Bras.
guajavaefolia, *Berg, in Mart. Fl. Bras.* xiv. i. 160.—Bras.
Guavira, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 182; *Contrib.* 142.—Paraguay.
Guavira-mi, *Parodi, l. c.* 140; 141.—Paraguay.
guianensis, *DC. Prod.* iii. 245.—Guiana.
Hayneana, *DC. l. c.* 246.—Bras.
hebepetala, *DC. l. c.*—Bras.
hiemalis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 332.—Bras.
Hilariana, *Berg, in Mart. Fl. Bras.* xiv. i. 199.—Bras.
hirsuta, *Berg, l. c.* 151.—Bras.
hirtiflora, *DC. Prod.* iii. 249.—Bras.
hispida, *Berg, in Mart. Fl. Bras.* xiv. i. 152.—Bras.
Hoffmannseggii, *Berg, l. c.* 157.—Bras.
huanocensis, *Berg, in Linnaea,* xxxi. (1861–62) 250.—Peruv.
Humboldtiana, *DC. Prod.* iii. 256.—Venezuela.
hypericoides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 317.—Bras.
hypoleuca, *Spring, ex Mart. in Flora,* xx. (1837) II. *Beibl.* 80.—Bras.
imbricata, Gardn. in Field. Sert. Pl. 75 = scutulifera.
Imperatoris-Maximiliani, *Wawra, Bot. Ergeb. Maxim. Bras.* 11.—Bras.
impressa, *Berg, in Mart. Fl. Bras.* xiv. i. 184.—Bras.
insularis, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 536.—Bras.
involucrata, *Berg, in Mart. Fl. Bras.* xiv. i. 156.—Bras.
itajuruensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 307.—Bras.
itambensis, *Berg, in Mart. Fl. Bras.* xiv. i. 190.—Bras.
Jaboticaba, *Baill. Hist. Pl.* vi. 345.—Bras.
Kegeliana, *Berg, in Linnaea,* xxvii. (1854) 99.—Guiana.

MYRCIA:—
Klotzschiana, *Berg, in Mart. Fl. Bras.* xiv. i. 169.—Bras.
Kunthiana, *Steud. Nom.* ed. II. ii. 172.—Venezuel.
Lacerdaeana, *Berg, in Mart. Fl. Bras.* xiv. i. 193.—Bras.
laevigata, *Berg, l. c.* 175.—Bras.
laevis, *Berg, in Linnaea,* xxxi. (1861–62) 252.—Guiana.
laevis, *G. Don, Gen. Syst.* ii. 845.—Bras.
lanceolata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 329.—Bras.
Langsdorffii, *Berg, in Mart. Fl. Bras.* xiv. i. 562.—Bras.
lanuginosa, *Berg, l. c.* 205.—Bras.
Laruotteana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 311.—Bras.
lasiantha, *DC. Prod.* iii. 254.—Bras.
lasiopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
lasiopus, *DC. Prod.* iii. 253.—Bras.
latifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 170.—Bras.
lauriflora, *DC. Prod.* iii. 252.—Bras.
laurifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 301.—Bras.
laxiflora, *Cambess. l. c.* 319.—Bras.
Lechleriana, *Miq. in Linnaea,* xxv. (1852) 651.—Chili.
leptoclada, *DC. Prod.* iii. 244.—Ins. S. Doming.
leucadendra, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
leucadendron, *DC. Prod.* iii. 251.—Bras.
leucophlaea, *DC. l. c.* 247.—Bras.
Lindeniana, *Wright, in Sauv. Fl. Cub.* 40.—Cuba.
linearifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 334.—Bras.
Linkiana, *DC. Prod.* iii. 248.—Bras.
littoralis, *DC. l. c.* 249.—Bras.
longifolia, *DC. l. c.* 256.—Venezuela.
Luschnathiana, *Berg, in Mart. Fl. Bras.* xiv. i. 194.—Bras.
lutescens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 301.—Bras.
macrocarpa, *DC. Prod.* iii. 249.—Bras.
macrochlamys, *DC. l. c.* 247.—Bras.
macrophylla, *DC. l. c.* 248.—Bras.
magnoliaefolia, *DC. l. c.*—Bras.
Mansoniana, *Berg, in Mart. Fl. Bras.* xiv. i. 163.—Bras.
maragnana, *DC. Prod.* iii. 249.—Bras.
marginata, *Berg, in Mart. Fl. Bras.* xiv. i. 565.—Bras.
Martiana, *Berg, l. c.* 159.—Bras.
Maximiliana, *Berg, in Linnaea,* xxvii. (1854) 89.—Bras.
melanoclada, *Berg, l. c.* 113.—Costa Rica.
melastomoides, *DC. Prod.* iii. 256.—Ins. Tobago.
micrantha, *Berg, in Mart. Fl. Bras.* xiv. i 169.—Bras.
microcarpa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 324.—Bras.
microphylla, *Berg, in Mart. Fl. Bras.* xiv. i. 566.—Bras.
Mikaniana, *Berg, l. c.* 182.—Bras.
Mini, *Sweet, Hort. Brit.* ed. II. 211.—Guiana.
minutiflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 185.—Guiana.
mollis, *DC. Prod.* iii. 256.—N. Granat.
montana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 325.—Bras.
mugiensis, *Cambess. l. c.* 327.—Bras.
multiflora, *DC. Prod.* iii. 244.—Guiana.
myoporina, *DC. l. c.* 246.—Bras.
myrtillifolia, *DC.* l. c. 250.—Bras.
Nandu-apysa, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 183; *Contrib.* 143.—Paraguay.
Neesiana, *DC. Prod.* iii. 249.—Bras.
negrensis, *Berg, in Mart. Fl. Bras.* xiv. i. 187.—Bras.
nigrescens, *DC. Prod.* iii. 246.—Bras.
nitens, *Berg, in Mart. Fl. Bras.* xiv. i. 178.—Bras.
nitida, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 309.—Bras.
nivea, *Cambess. l. c.* 332.—Bras.
nobilis, *Berg, in Mart. Fl. Bras.* xiv. i. 195.—Bras.
oblongata, *DC. Prod.* iii. 251.—Bras.
obtusa, *Schau. in Linnaea,* xxi. (1848) 273.—Guiana.
ochracea, *Berg, in Mart. Fl. Bras.* xiv. i. 190.—Bras.
ochroides, *Berg, l. c.* 208.—Bras.

MYRCIA:—
Oerstediana, *Berg, in Linnaea*, xxvii. (1854) 112.—Mexic.
Oitchi, *Berg, in Mart. Fl. Bras.* xiv. i. 563.—Bras.
oleaefolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 313.—Bras.
oligantha, *Berg, in Mart. Fl. Bras.* xiv. i. 184.—Bras.
oocarpa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 298.—Bras.
opaca, *Berg, in Mart. Fl. Bras.* xiv. i. 177.—Bras.
ovata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 319.—Bras.
Paivae, *Berg, in Mart. Fl. Bras.* xiv. i. 179.—Bras.
pallens, *DC. Prod.* iii. 252.—Bras.
palustris, *DC. l. c.* 246.—Bras.
pauciflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 321.—Bras.
pellucida, *Berg, in Mart. Fl. Bras.* xiv. i. 173.—Bras.
perforata, *Berg, l. c.* 197.—Bras.
pernambucensis, *Berg, l. c.* 194.—Bras.
pertusa, *DC. Prod.* iii. 251.
phaeoclada, *Berg, in Mart. Fl. Bras.* xiv. i. 167.—Bras.
piauhiensis, *Berg, l. c.* 196.—Bras.
pilosa, *DC. Prod.* iii. 253.—Bras.
pimentoides, DC. Prod. iii. 243 = Eugenia acris.
Pinaster, Mart. ex Berg, in Linnaea, xxvii. (1854) 338 = Eugenia myriophylla.
pinifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 333.—Bras.
platyclada, *DC. Prod.* iii. 244.—Guiana.
plicato-costata, *Berg, in Linnaea*, xxvii. (1854) 114.—Costa Rica.
Poeppigiana, *Berg, in Mart. Fl. Bras.* xiv. i. 157.—Bras.
Pohliana, *Berg, l. c.* 199.—Bras.
polyantha, *DC. Prod.* iii. 252.—Bras.
polyantha, DC. l. c. 255 = Kunthiana.
prunifolia, *DC. l. c.* 253.—Bras.
Pseudo-Mini, *DC. l. c.* 252.—Bras.
puberula, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 316.—Bras.
pubescens, *DC. Prod.* iii. 247.—Bras.
pubescens, *G. Don, Gen. Syst.* ii. 845.—Bras.
pubiflora, *DC. Prod.* iii. 249.—Bras.
pubipetala, *Miq. in Linnaea*, xix. (1847) 441.—Bras.
punctata, *DC. Prod.* iii. 243.—Ins. S. Crucis.
pyramidata, *Berg, in Mart. Fl. Bras.* xiv. i. 193.—Bras.
quitarensis, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 184.—Guian.
racemulosa, *DC. Prod.* iii. 254.—Bras.
ramulosa, *DC. l. c.* 250.—Bras.
recurvata, *Berg, in Mart. Fl. Bras.* xiv. i. 166.—Bras.
Regeliana, *Berg, l. c.* 562.—Bras.
Regnelliana, *Berg, l. c.* 174.—Bras.
reticulata, *Berg, in Linnaea*, xxvii. (1854) 101.—Ins. Martinic.
reticulata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 304.—Bras.
reticulosa, *Miq. in Linnaea*, xxii. (1849) 794.—Bras.
retorta, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 322.—Bras.
Riedeliana, *Berg, in Mart. Fl. Bras.* xiv. i. 565.—Bras.
rimosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 333.—Bras.
riparia, *Berg, in Mart. Fl. Bras.* xiv. i. 179.—Bras.
rostrata, *DC. Prod.* iii. 255.—Bras.
rubella, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 317.—Bras.
rubiginosa, *Cambess. l. c.* 300.—Bras.
rufidula, *Schlecht. in Linnaea*, xiii. (1839) 416.—Mexic.
rufipes, *DC. Prod.* iii. 247.—Bras.
rufula, *Miq. in Linnaea*, xix. (1847) 440.—Bras.
salicifolia, *DC. Prod.* iii. 246.—Bras.
Salzmanni, *Berg, in Mart. Fl. Bras.* xix. i. 207.—Bras.
Sartoriana, *Berg, in Linnaea*, xxix. (1857–58) 220.—Mexic.
saxicola, *Berg, l. c.* xxvii. (1854) 92.—Peruv.
Schaueriana, *Berg, in Mart. Fl. Bras.* xiv. i. 163.—Bras.
Schomburgkiana, *Berg, in Linnaea*, xxvii. (1854) 110.—Guiana.

MYRCIA:—
Schottiana, *Berg, in Mart. Fl. Bras.* xiv. i. 188.—Bras.
Schrankiana, *DC. Prod.* iii. 247.—Bras.
Schuechiana, *Berg, in Mart. Fl. Bras.* xiv. i. 181 = Bras.
scrobiculata, *Berg, in Linnaea*, xxx. (1859–60) 668.—Bras.
scutulifera, *DC. Prod.* iii. 254.—Bras.
Sellowiana, *Berg, in Mart. Fl. Bras.* xiv. i. 197.—Bras.
sepiaria, *DC. Prod.* iii. 249.—Bras.
sericea, *Berg, in Linnaea*, xxvii. (1854) 114.—Peruv.; Chili.
sericea, *G. Don, Gen. Syst.* ii. 844.—Bras.
sericiflora, *Berg, in Mart. Fl. Bras.* xiv. i. 178.—Bras.
sororia, *DC. Prod.* iii. 243.—Ins. Trinit.
spectabilis, *DC. l. c.* 248.—Bras.
sphaerocarpa, *DC. l. c.* 251.—Bras.
Spixiana, *DC. l. c.*—Bras.
splendens, *DC. l. c.* 244.—Ind. occ.
Sprengeliana, *Berg, in Linnaea*, xxvii. (1854) 109.—Ins. S. Thomas.
Spruceana, *Berg, in Mart. Fl. Bras.* xiv. i. 165.—Bras.
Stewartiana, *Berg, in Linnaea*, xxix. (1857–58) 219.—Am. trop.
stigmatosa, *Berg, in Mart. Fl. Bras.* xiv. i. 564.—Bras.
strigipes, Mart. in Flora, xxiv. (1841) ii. Beibl. 108 = Marlierea strigipes.
suaveolens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 315.—Bras.
subalpestris, *DC. Prod.* iii. 250.—Bras.
subcordata, *DC. l. c.* 253.—Bras.
subsericea, *A. Gray, Bot. U. St. Expl. Exped.* i. 533.—Bras.
subsessilis, *Berg, in Linnaea*, xxxi. (1861–62) 251.—Am. trop.
suffruticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 189.—Bras.
sulcata, *Berg, in Linnaea*, xxx. (1859–60) 667.—Guiana.
superba, *Berg, in Mart. Fl. Bras.* xiv. i. 198.—Bras.
surinamensis, *Miq. in Linnaea*, xxii. (1849) 170.—Guiana.
sylvatica, *DC. Prod.* iii. 244.—Guiana.
terebinthacea, Poepp. ex Berg, in Mart. Fl. Bras. xiv. i. 155 = lanceolata.
thomasiana, *DC. Prod.* iii. 244.—Ins. S. Thomas.
thyrsoïdea, *Berg, in Mart. Fl. Bras.* xiv. i. 192.—Bras.
tingens, *Berg, l. c.* 564.—Bras.
tomentosa, *DC. Prod.* iii. 245.—Guiana.
torta, *DC. l. c.* 250.—Bras.
tovarensis, *Berg, in Linnaea*, xxvii. (1854) 118.—Venezuela.
triantha, *DC. Prod.* iii. 256.—Hab. ?
triflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 327.—Bras.
uberavensis, *Berg, in Mart. Fl. Bras.* xiv. i. 568.—Bras.
umbellulifera, *DC. Prod.* iii. 255.—Venezuela.
undulata, *Berg, in Mart. Fl. Bras.* xiv. i. 185.—Bras.
Valenzuelana, Griseb. Cat. Pl. Cub. 86 = Eugenia Valenzuelana.
variabilis, *DC. Prod.* iii. 254.—Bras.
Vauthiereana, *Berg, in Mart. Fl. Bras.* xiv. i. 154.—Bras.
Velloziana, *Berg, l. c.* 171.—Bras.
velutina, *Berg, l. c.* 182.—Bras.
venezuelensis, *Berg, in Linnaea*, xxvii. (1854) 96.—Venezuela.
venulosa, *DC. Prod.* iii. 250.—Bras.
vernicosa, *DC. l. c.* 256.—Am. austr.
verticillaris, *Berg, in Mart. Fl. Bras.* xiv. i. 206.—Bras.
vestita, *DC. Prod.* iii. 248.—Bras.
virgata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 320.—Bras.
xylopioides, *DC. Prod.* iii. 256.—N. Granat.
ypanemensis, *Berg, in Mart. Fl. Bras.* xiv. i. 186.—Bras.

MYRCIA, Soland. ex Lindl. Coll. Bot. sub t. 19 (1821–25) = **Pimenta**, Lindl. (Myrtac.).

MYRCIANTHES, Berg, in Linnaea, xxvii. (1854) 315 = **Myrtus**, Linn.
apiculata, Berg, in Mart. Fl. Bras. xiv. i. 352.—Reg. Argent.
brunnea, Berg, in Linnaea, xxvii. (1854) 315 = Myrtus ovalis, *Spreng.*
cisplatensis, Berg, l. c. = Eugenia cisplatensis.
edulis, Berg, in Mart. Fl. Bras. xiv. i. 353 = Eugenia edulis, *Benth. & Hook. f.*

MYRCIARIA, Berg, in Linnaea, xxvii. (1854) 320 = **Eugenia**, Linn. (Myrtac.).
amazonica, Berg, in Mart. Fl. Bras. xiv. i. 374.—Bras.
axillaris, Berg, l. c. 373.—Bras.
bipennis, Berg, in Linnaea, xxxi. (1861–62) 259.—Bras.
cauliflora, Berg, in Mart. Fl. Bras. xiv. i. 361 = E. edulis.
chartacea, Berg, l. c. 371.—Bras.
ciliolata, Berg, l. c. 366 = E. ciliolata.
cordata, Berg, in Linnaea, xxvii. (1854) 337.—Guiana.
cuspidata, Berg, in Mart. Fl. Bras. xiv. i. 367 = E. tenella, *Miq.*
delicatula, Berg, l. c. 362 = E. delicatula.
deserti, Berg, l. c. 359 = E. deserti.
disticha, Berg, l. c. 366.—Bras.
divaricata, Berg, in Linnaea, xxvii. (1854) 334 = E. divaricata, *Benth.*
Ehrenbergiana, Berg, l. c. 321.—Guiana.
ferruginea, Berg, in Mart. Fl. Bras. xiv. i. 597.—Bras.
floribunda, Berg, in Linnaea, xxvii. (1854) 330 = E. floribunda.
glomerata, Berg, in Mart. Fl. Bras. xiv. i. 365.—Bras.
Gnapurium, Berg, in Linnaea, xxvii. (1854) 325 = E. Gnapurium.
Hagendorffii, Berg, in Mart. Fl. Bras. xiv. i. 360.—Bras.
herbacea, Berg, l. c. 369.—Bras.
involucrata, Berg, l. c. 375.—Bras.
Jaboticaba, Berg, l. c. 361 = Myrcia Jaboticaba.
lanceolata, Berg, l. c. 363.—Bras.
leucadendron, Berg, l. c. 364.—Bras.
leucophloea, Berg, l. c. 370.—Bras.
linearifolia, Berg, l. c. 362.—Bras.
lituatinervia, Berg, in Linnaea, xxvii. (1854) 322.—Guiana.
longipes, Berg, in Mart. Fl. Bras. xiv. i. 599.—Bras.
maragnanensis, Berg, l. c. 372.—Bras.
maranhensis, Berg, l. c. Index 652 = praec.
marowynensis, Berg, in Linnaea, xxvii. (1854) 335 = E. marowynensis.
Maximiliana, Berg, in Mart. Fl. Bras. xiv. i. 370 = E. Maximiliana, *DC.*
micrantha, Berg, l. c. 360.—Bras.
minensis, Berg, l. c. 369.—Bras.
myriophylla, Berg, l. c. 375 = E. myriophylla.
nitida, Berg, in Linnaea, xxvii. (1854) 324 = E. nitida, *Benth.*
obscura, Berg, in Mart. Fl. Bras. xiv. i. 374.—Bras.
pallida, Berg, l. c. 368.—Bras.
paraënsis, Berg, l. c. 364.—Bras.
perforata, Berg, l. c. 598.—Bras.
phillyraeoides, Berg, in Linnaea, xxvii. (1854) 326.—Am. austr.
plicato-costata, Berg, in Mart. Fl. Bras. xiv. i. 366 = E. edulis, *Vell.*
polyantha, Berg, in Linnaea, xxvii. (1864) 322 = E. polyantha, *Miq.* (in Addend.).
prasina, Berg, in Mart. Fl. Bras. xiv. i. 373.—Bras.
protracta, Berg, in Linnaea, xxvii. (1854) 330 = E. floribunda.
pumila, Berg, in Mart. Fl. Bras. xiv. i. 366 = E. pumila.
quitarensis, Berg, in Linnaea, xxvii. (1854) 323 = E. quitarensis.
ramiflora, Berg, l. c. 334 = E. coffeaefolia.
Riedeliana, Berg, in Mart. Fl. Bras. xiv. i. 598.—Bras.
rivularis, Berg, l. c. 360 = E. rivularis, *Cambess.*
rubiginosa, Berg, l. c. 359 (= *Eugeniopsis rubiginosa*).—Bras.
Salzmanni, Berg, in Linnaea, xxvii. (1854) 331 = E. Salzmanni.
Schaueriana, Berg, l. c. 323 = E. Schaueriana.
Schuechiana, Berg, in Mart. Fl. Bras. xiv. i. 373.—Bras.
Sellowiana, Berg, l. c. 374.—Bras.

MYRCIARIA :—
sericea, Berg, in Mart. Fl. Bras. xiv. 1. 375.—Bras.
splendens, Berg, l. c. 371.—Bras.
Spruceana, Berg, l. c. 365.—Bras.
stirpiflora, Berg, in Linnaea, xxx. (1859–60) 702 = Eugenia trunciflora, *Rich.* (in Addend.).
strigipes, Berg, in Mart. Fl. Bras. xiv. 1. 364.—Bras.
tenella, Berg, l. c. 368 = E. tenella, *DC.*
tenuiramis, Berg, l. c. 371.—Bras.
tolypantha, Berg, l. c. 372.—Bras.
Trinitatis, Berg, in Linnaea, xxvii. (1854) 336.—Ins. Trinit.
trunciflora, Berg, in Mart. Fl. Bras. xiv. 1. 361.—Bras.
uliginosa, Berg, in Linnaea, xxvii. (1854) 329.—Guiana.
undulata, Berg, in Mart. Fl. Bras. xiv. 1. 368.—Bras.
verticillata, Berg, in Linnaea, xxvii. (1854) 332.—Guiana.
vismeaefolia, Berg, l. c. 336 = E. vismeaefolia.

MYRIACHAETA, Zoll. & Mor. Syst. Verz. Zoll. 101 (1845–46) = **Thysanolaena**, Nees (Gramin.).
arundinacea, Zoll. & Mor. l. c. = T. Agrostis.
glauca, Mor. ex Steud. Syn. Pl. Gram. 404.—Malaya.

MYRIACTIS, Less. in Linnaea, vi. (1831) 127. *COMPOSITAE*, Benth. & Hook. f. ii. 263.
BOTRYADENIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 30 (1835).
carnosa, Wall. Cat. n. 3219 = nepalensis.
glutinosa, *Schlecht. in Linnaea*, xxv. (1852) 207.—Ind. or.
Gmelini, DC. Prod. v. 309 = nepalensis.
gracilis, Edgew. in Trans. Linn. Soc. xx. (1846) 66 = nepalensis.
javanica, DC. Prod. v. 308 = Wightii.
nepalensis, *Less. in Linnaea*, vi. (1831) 128.—Reg. Himal. ; As. centr.
oleosa, Edgew. in Trans. Linn. Soc. xx. (1846) 65 = nepalensis.
pilosa, *Blume, ex DC. Prod.* v. 308.—Java.
Wallichii, *Less. in Linnaea*, vi. (1831) 129.—Reg. Himal.
Wightii, *DC. in Wight, Contrib.* 10.—Ind. or.

MYRIADENUS, Cass. in Bull. Soc. Philom. (1817) 138 = **Inula**, Linn. (Compos.).
glutinosus, Cass. l. c. = I. viscosa.

MYRIADENUS, Desv. Journ. Bot. i. (1813) 121. t. 4. f. 11 = **Zornia**, J. F. Gmel. (Legumin.).
echinocarpus, Moric. ex Meissn. Gen. Comm. 64 = Z. echinocarpa.
tetraphyllus, Desv. Journ. Bot. i. (1813) 121 ; et DC. Prod. ii. 316 = Z. myriadena.

MYRIANDRA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 364 = **Hypericum**, Linn.
brachyphylla, Spach, l. c. 365 = H. fasciculatum.
Brathydis, Spach, Hist. Vég. Phan. v. 436 = H. fasciculatum.
galioides, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 365 = H. galioides.
glauca, Spach, l. c. = H. myrtifolium.
ledifolia, Spach, l. c. = H. nudiflorum.
Michauxii, Spach, l. c. = H. galioides.
nitida, Spach, l. c. = H. fasciculatum.
nudiflora, Spach, l. c. = H. nudiflorum.
prolifica, Spach, l. c. = H. prolificum.
spathulata, Spach, l. c. = H. prolificum.

MYRIANTHEA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 65 = seq.

MYRIANTHEIA, Thou. Gen. Nov. Madag. 21 (1806) = **Homalium**, Jacq. (Samyd.).
laxiflora, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 66.—Madag.
oppositifolia, Tul. l. c. 65.—Madag.

MYRIANTHUS, Beauv. Fl. Owar. i. 16 (1805). *URTICACEAE*, Benth. & Hook. f. iii. 379.
DICRANOSTACHYS, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 85. t. 1.
arboreus, *Beauv. Fl. Owar.* i. 16.—Afr. bor.

MYRIASPORA, DC. Prod. iii. 165 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 769.
decipiens, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 101.—Am. trop.
egensis, *DC. Prod.* iii. 165.—Bras.
integrifolia, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 145 = decipiens.
paulensis, DC. Prod. iii. 165 = egensis.
surinamensis, Steud. in Flora, xxvii. (1844) 722 = egensis.

MYRICA, Linn. Syst. ed. I (1735). *MYRICACEAE*, Benth. & Hook. f. iii. 400.
CEROPHORA, Rafin. Alsog. Am. 11 (1838).
COMPTONIA, Banks, ex Gaertn. Fruct. ii. 58. t. 90 (1791).
FAYA, Webb & Berth. Phyt. Canar. iv. 272. t. 216 (1836–50)
FAYANA, Rafin. Alsog. Am. 12 (1838).
GALE, Tourn. ex Adans. Fam. ii. 345 (1763).
MORELLA, Lour. Fl. Cochinch. 548 (1790).
NAGEIA, Gaertn. Fruct. i. 191 (1788).
PIMECARIA, Rafin. Alsog. Am. 64 (1838).
adenophora, *Hance, in Journ. Bot.* xxi. (1883) 357.—China.
aethiopica, *Linn. Mant.* ii. 298.—Afr. austr.
alaternoides, Crantz, Inst. i. 182 = cordifolia.
altera, *C. DC. in DC. Prod.* xvi. II. 595.—Ind. occ.
arabica, Willd. Sp. Pl. iv. 749, in syn. = Myrsine bottensis.
arguta, *H. B. & K. Nov. Gen. et Sp.* ii. 17. t. 98.—N. Granat.
asplenifolia, *Linn. Sp. Pl.* 1024 ; *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. bor.
australasica, *F. Muell. Pl. Fitzalan*, 7.—Austral.
banksiaefolia, Wendl. Coll. Pl. i. 70 = aethiopica.
Bojeriana, *Baker, in Journ. Linn. Soc.* xx. (1883) 267.—Madag.
brabantica, S. F. Gray, Nat. Arr. Brit. Pl. ii. 249 = M. Gale.
brevifolia, *E. Mey. ex C. DC. in DC. Prod.* xvi. II. 150.—Afr. austr.
Burmani, *E. Mey. l. c.* 149.—Afr. austr.
californica, *Cham. & Schlecht. in Linnaea*, vi. (1831) 335.—Calif.
capensis, Hort. ex Steud. Nom. ed. II. ii. 173 = aethiopica.
caracasana, *H. B. & K. Nov. Gen. et Sp.* ii. 18.—Venezuela.
carolinensis, Mill. Gard. Dict. ed. VIII. n. 3 = cerifera.
cerifera, *Linn. Sp. Pl.* 1024.—Am. bor.
Comptonia, C. DC. in DC. Prod. xvi. II. 151 = asplenifolia.
conifera, Burm. f. Fl. Cap. Prod. 31 (err. typ. 27) = aethiopica.
cordifolia, *Linn. Sp. Pl.* 1025.—Afr. austr.
domingana, *C. DC. in DC. Prod.* xvi. II. 154.—Ins. S. Doming.
esculenta, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 56 = M. Nagi.
Farquhariana, Wall. Tent. Fl. Nep. 60 = M. Nagi.
Faya, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 397.—Ins. Canar. et Azor.
florida, *Regel, Cat. Pl. Hort. Aksakov.* 96, nomen.—Hab. ?
frondosa, Salisb. Prod. 396 = M. Faya.
fuscata, Rafin. Alsog. Am. 9 (Quid ?).—Am. bor.
Gale, *Linn. Sp. Pl.* 1024.—Reg. temp. bor.
Hartwegi, *S. Wats. in Proc. Am. Acad.* x. (1875) 350.—Calif.
heterophyla, Rafin. Alsog. Am. 9 (Quid ?).—Am. bor.
hirsuta, Mill. Gard. Dict. ed. VIII. n. 6 = quercifolia.
humilis, *Cham. & Schlecht. in Linnaea*, vi. (1831) 535.—Afr. austr.
ilicifolia, *Burm. f. Fl. Cap. Prod.* 31 (err. typ. 27).—Afr. austr.
inodora, *Bartr. Trav.* ed. II. (1794) 403.—Florida.
integrifolia, Roxb. Hort. Beng. 71 ; Fl. Ind. iii. 765 = M. Nagi.
interrupta, *Benth. Pl. Hartw.* 251.—N. Granat.
javanica, *Blume, Bijdr.* 517.—Java.
Kraussiana, *Buching. in Flora*, xxviii. (1845) 89.—Afr. austr.
laciniata, Willd. Enum. Hort. Berol. 1012 = quercifolia.
Laureola, *Tréc. ex C. DC. in DC. Prod.* xvi. II. 154.—Am. bor.
Lindeniana, *C. DC. l. c.* 150.—Mexic.
linearis, *C. DC. l. c.* 154.—Hab. ?

MYRICA :—
Lobbii, *Teysm. & Binn. ex Miq. Fl. Ind. Bat.* i. 1. 872.—Java.
longifolia, *Teysm. & Binn. ex C. DC. in DC. Prod.* xvi. II. 152.—Java.
luzonica, *Vidal, Sin. Pl. Leñ. Filip. Atl.* p. xl. t. 90. f. B.—Ins. Philipp.
macrocarpa, H. B. & K. Nov. Gen. et Sp. ii. 16 = arguta.
macrophylla, Mirb. in Mém. Mus. Par. xiv. (1827) 472. t. 27 = javanica.
mexicana, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1011.—Mexic.
microcarpa, *Benth. Pl. Hartw.* 266.—Ind. occ.
missionis, Wall. Cat. n. 7297 = M. Nagi.
montana, Vahl, Symb. Bot. ii. 99 = Myrsine bottensis.
Nagi, *Thunb. Fl. Jap.* 76.—As. trop. et subtrop.
natalensis, *C. DC. in DC. Prod.* xvi. II. 148.—Afr. austr.
obovata, *Cham. ex C. DC. l. c.* 150.—Florida.
octandra, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 56.—Nepal.
ovata, *Wendl. f. in Bartl. & Wendl. f. Beitr.* ii. 3.—Afr. austr.
palustris, Lam. Fl. Fr. ii. 236 = M. Gale.
parvifolia, *Benth. Pl. Hartw.* 251.—N. Granat.
Pavonis, *C. DC. in DC. Prod.* xvi. II. 151.—Peruv.
pensylvanica, Hort. Reg. ex Lam. Encyc. ii. 592 = cerifera.
phillyreaefolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 267.—Madag.
polycarpa, *H. B. & K. Nov. Gen. et Sp.* ii. 18—N. Granat.
polycarpa, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 134 = xalapensis.
pubescens, *Willd. Sp. Pl.* iv. 746.—N. Granat.
punctata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 177 = cerifera.
pusilla, Rafin. Alsog. Am. 10 (Quid ?).—Am. bor.
quercifolia, Linn. Sp. Pl. 1025.—Afr. austr.
rotundifolia, Salisb. Prod. 396 = cordifolia.
rubra, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 230 = M. Nagi.
salicifolia, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 267 = Bojeriana.
salicifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 277.—Abyssin.
sapida, Wall. Tent. Fl. Nep. 59. t. 45 = M. Nagi.
segregata, Jacq. Coll. ii. 273 = Forestiera porulosa.
serrata, *Lam. Encyc.* ii. 592 = aethiopica.
sessilifolia, Rafin. Alsog. Am. 10 (Quid ?).—Am.. bor.
spathulata, *Mirb. in Mém. Mus. Par.* xiv. (1827) 474. t. 28. f. 1.—Madag.
trifoliata, Hort. ex Turp. in Ann. Mus. Par. v. (1804) 447 = Rhus aromatica.
trifoliata, Linn. Pl. Rar. Afr. 28 ; Amoen. Acad. vi. 112 = Toddalia lanceolata.
trifoliolata, Hort. ex DC. Prod. ii. 73 = Rhus aromatica.
undulata, Rafin. Alsog. Am. 10 (Quid ?).—Am. bor.
verrucosa, Rafin. l. c. 9 (Quid ?).—Am. bor.
Vidaliana, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 316.—Ins. Philipp.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* ii. 16.—Mexic.
xalapensis, Poepp. ex Griseb. Fl. Brit. W. Ind. 177 = microcarpa.
Zeyheri, *C. DC. in DC. Prod.* xvi. II. 149.—Afr. austr.

MYRICARIA, Desv. in Ann. Sc. Nat. Sér. I. iv. (1825) 348. *TAMARISCINEAE*, Benth. & Hook. f. i. 161.
alopecuroides, Schrenk, Enum. Pl. Nov. 65 = germanica.
armena, *Boiss. Diagn.* Ser. II. 58. ii.—Armenia.
bracteata, Royle, Illustr. Bot. Himal. 214. t. 44. fig. 2 = germanica.
brevifolia, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 70.—Mongolia.
dahurica, DC. Prod. iii. 98 = seq.
davurica, *Ehrenb. in Linnaea*, ii. (1827) 278.—Sibir.
elegans, *Royle, Illustr. Bot. Himal.* 214.—Reg. Himal.
germanica, *Desv. in Ann. Sc. Nat.* Sér. I. iv. (1825) 349.—Europ. ; Oriens ; As. bor.
herbacea, Desv. l. c. 350 = germanica ?
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 120 = germanica.

MYRICARIA:—
linearifolia, Desv. in Ann. Sc. Nat. Sér. I. iv. (1825) 349 = longifolia.
longifolia, *Ehrenb. in Linnaea*, ii. (1827) 279; *DC. Prod.* iii. 97.—Sibir. baical.
macrostachya, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 423 = germanica.
platyphylla, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 425.—Mongolia.
prostrata, Hook. f. & Thoms. ex Benth. & Hook. f. Gen. i. 161 = germanica.
squamosa, Desv. in Ann. Sc. Nat. Sér. I. iv. (1825) 350 = germanica.
vaginata, Desv. l. c. = Tamarix ericoides.

MYRICE, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = praec.

MYRINIA, Lilja, Fl. Sverig. Suppl. i. 25 (1840) = **Fuchsia**, Linn. (Onagr.).
microphylla, Lilja, l. c. = F. microphylla.

MYRIOBLASTUS, Wall. ex Griff. in Proc. Linn. Soc. i. (1845) 264 = **Cryptocoryne**, Fisch. (Aroid.).

MYRIOCARPA, Benth. Bot. Voy. Sulph. 168. t. 55 (1844). *URTICACEAE*, Benth. & Hook. f. iii. 391.
bifurca, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 307.—Nicaragua.
colipensis, Liebm. l. c. 306 = longipes.
cordifolia, *Liebm. l. c.*—Mexic.
densiflora, *Benth. Bot. Voy. Sulph.* 169.—Peruv.
Dombeyana, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 232 = densiflora.
humilis, Wedd. l. c. = stipitata.
inaequilatera, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 307.—Costa Rica.
longipes, *Liebm. l. c.* 306.—Costa Rica; Mexic.
macrophylla, *Bartl.—Cf. Ind. Sem. Hort. Gotting.* (1871) 5.—Hab.?
polystachya, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 231 = stipitata.
stipitata, *Benth. Bot. Voy. Sulph.* 168. t. 55.—Am. trop.

MYRIOCEPHALUS, Benth. in Enum. Pl. Hueg. 61 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 318.
ANTHEIDOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 173.
ELACHOPAPPUS, F. Muell. Fragm. iii. 157 (1863).
GILBERTA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 192.
HIRNELLIA, Cass. in Bull. Soc. Philom. (1820) 57, nomen prius.
HYALOLEPIS, DC. Prod. vi. 149 (1837).
LAMPROCHLAENA, F. Muell. Fragm. iii. 157 (1863).
POLYCALYMMA, F. Muell. & Sond. in Linnaea, xxv. (1852) 494.
appendiculatus, *Benth. in Enum. Pl. Hueg.* 61.—Austral.
cotuloides, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 73 = Gnephosis brevifolia.
Decazesii, F. Muell. Fragm. xi. 72 = Decazesia hecatocephala.
gracilis, *Benth. Fl. Austral.* ii. 559.—Austral.
Guerinae, *F. Muell. Fragm.* viii. 169.—Austral.
helichrysoides, *A. Gray, in Hook. Kew Journ.* iii. (1851) 175.—Austral.
nudus, *A. Gray, l. c.* 174.—Austral.
rhizocephalus, *Benth. Fl. Austral.* iii. 557.—Austral.
Rudallii, *F. Muell. Fragm.* iii. 157; *Benth. l. c.* 559.—Austral.
Stuartii, *Benth. l. c.* 560.—Austral.
suffruticosus, *Benth. l. c.* 559.—Austral.
villosissimus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 74 = Gnephosis eriocephala.

MYRIOGOMPHUS, Didr. in Kjoeb. Vidensk. Meddel. (1857) 142 = **Croton**, Linn. (Euphorb.).
fuscus, Didr. l. c. 143 = C. fuscus.

MYRIOGYNE, Less. in Linnaea, vi. (1831) 219 = **Centipeda**, Lour. (Compos.).
Cunninghami, DC. Prod. vi. 139 = C. Cunninghami.
elatinoides, Less. in Linnaea, vi. (1831) 219.—Chili.
latifolia, Hassk. Cat. Hort Bog. Alt. 102 = Dichrocephala latifolia.
minuta, Less. in Linnaea, vi. (1831) 219 = C. obicularis.
racemosa, Hook. in Mitch. Journ. Trop. Austral. 353 = C. racemosa.

MYRIONEURON, R. Br. in Wall. Cat. n. 6225 (1832). *RUBIACEAE*, Benth. & Hook. f. ii. 69.
BERLIERA, Buch.-Ham. ex Wall. Cat. sub n. 6225 (1832).
angustifolium, *Hook. f. Fl. Brit. Ind.* iii. 97.—Reg. Himal.
Clarkei, *Hook. f. l. c.* 96.—Reg. Himal.
hirsutum, *Kurz, For. Fl. Brit. Burma*, ii. 55.—Burma.
nutans, *Wall. Cat.* n. 6225.—Reg. Himal.

MYRIOPHILLUM, Neck. Delic. Gallo-Belg. ii. 386 (1768) = seq.

MYRIOPHYLLUM, Ponted. ex Linn. Syst. ed. I (1735). *HALORAGEAE*, Benth. & Hook. f. i. 676.
BURSHIA, Rafin. in Med. Repos. N. York, v. (1808) 357.
ENYDRIA, Vell. Fl. Flum. 56 (1825); i. t. 150 (1827).
HYLAS, Bigel. ex DC. Prod. iii. 69, in obs. (1828).
PELONASTES, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 474.
PENTAPTERIS, Hall. Enum. Stirp. Helv. i. 201 (1742)
POTAMOGETON, Walt. Fl. Carol. 90 (1788).
PTILOPHYLLUM, Reichb. Nom. 169 (1841).
PURSHIA, Rafin. in Am. Monthly Mag. (1819) 191, sub n. 33.
SPHONDYLASTRUM, Reichb. Nom. 169 (1841).
alterniflorum, *DC. Fl. Fr. Suppl.* 529.—Europ.; Groenl.
ambiguum, *Nutt. Gen. Am.* ii. 212.—Am. bor.
amphibium, *Labill. Nov. Holl. Pl.* ii. 70. t. 220.—Tasmania.
axilliflorum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 340.—Madag.
brasiliense, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 252.—Bras.
capillaceum, Torr. Compend. 355 = ambiguum.
chuquitense, *Meyen, Reise*, i. 491, nomen.—Peruv.
denudatum, La Pilaye, ex Steud. Nom. ed. II. ii. 174 (Quid?).—Terra nova.
dicoccum, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 41.—Austral.
Drummondii, *Benth. Fl. Austral.* ii. 490.—Austral.
elatinoides, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 105.—Austral.; N. Zel.; Am. austr.
filiforme, *Benth. Fl. Austral.* ii. 489.—Austral.
fulvescens, Bertol. Misc. Bot. xvi. 23 = Proserpinaca pectinata.
gracile, *Benth. Fl. Austral.* ii. 489.—Austral.
heterophyllum, *Michx. Fl. Bor. Am.* ii. 191.—Am. bor.
hippuroides, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 530.—Am. bor. occ.
indicum, Griff. Notul. iv. 687 = tuberculatum.
indicum, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 432 = intermedium, tuberculatum.
indicum, *Willd. Sp. Pl.* iv. 407.—Ind. or.
integrifolium, *Hook. f. Fl. Tasman.* i. 123. t. 23.—Austral.
intermedium, *DC. Prod.* iii. 69.—Ind. or.; Malaya; Austral.; N. Zel.
intermedium, Wight, Illustr. ii. t. 102, partim = tuberculatum.
latifolium, *F. Muell. Fragm.* ii. 87.—Austral.
laxum, *Shuttl. ex Chapm. Fl. S. U. St.* 143.—Am. bor.
limosum, Hectot, ex DC. Fl. Fr. Suppl. 530 = verticillatum.
limosum, Nutt. Gen. Am. ii. 212 = ambiguum.
lineare, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 433 = intermedium.
montanum, Martr. Pl. Crit. Tarn. 254 = alterniflorum.
Muelleri, *Sond. in Linnaea*, xxviii. (1856) 233.—Austral.
nudum, La Pilaye, ex DC. Prod. iii. 69 = tenellum.
pectinatum, DC. Fl. Fr. Suppl. 529 = spicatum, verticillatum.
pedunculatum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 474.—Austral.
procumbens, Bigel. Fl. Bost. ed. II. 346 = ambiguum.
propinquum, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 30 = elatinoides.
proserpinacoides, *Gill. ex Hook. & Arn. in Bot. Misc.* iii. (1833) 313.—Am. austr.
pusillum, *Blume, Mus. Bot. Lugd. Bat.* i. 111.—Ins. Banca.
quitense, *H. B. & K. Nov. Gen. & Sp.* vi. 89.—Ecuador.

MYRIOPHYLLUM:—
robustum, *Hook. f. Handb. N. Zeal. Fl.* 67.—N. Zel.
scabratum, Cham. in Linnaea, iv. (1829) 506 = hippuroides.
scabratum, *Michx. Fl. Bor. Am.* ii. 190.—Am. bor.
setaceum, La Pilaye, in Mém. Soc. Linn. Par. iv. (1826) 504, 535 (Quid?).—Terra nova.
siculum, Guss. Fl. Sic. Syn. ii. 599 = verticillatum.
sparsiflorum, *Wright, in Sauv. Fl. Cub.* 37.—Cuba.
spicatum, S. G. Gmel. Reise, ii. 198. t. 33 = verticillatum.
spicatum, Linn. Sp. Pl. 992.—Reg. bor. temp.
tenellum, *Bigel. Fl. Bost.* ed. II. 346.—Am. bor.
ternatum, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 106.—Chili; Patagon.
tetrandrum, Roxb. Fl. Ind. ed. Carey, i. 470 = indicum.
titikakense, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 352.—Peruv.
trachycarpum, *F. Muell. Fragm.* ii. 87.—Austral.
tuberculatum, *Roxb. Hort. Beng.* 12; *Fl. Ind.* i. 451.—Ind. or.
ussuriense, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 182.—Mandschur.; Japon.
variaefolium, Hook. f. Ic. Pl. t. 289 = intermedium.
verrucosum, *Lindl. in Mitch. Journ. Trop. Austral.* 384.—Austral.
verticillatum, *Linn. Sp. Pl.* 992.—Reg. temp. bor. & austr.
viridescens, Gill. ex Hook. & Arn. in Bot. Misc. iii. (1833) 314 = ternatum.
Wrightii, A. Gray, in Hall, Pl. Texas, 9, sphalm. = Mentzelia Wrightii.

MYRIOPTERON, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 385. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 744.
JENKINSIA, Wall. ex Voigt, Hort. Suburb. Calc. 544 (1845).
VICARYA, Wall. ex Voigt, l. c. (1845).
Horsfieldii, Hook. f. Fl. Brit. Ind. iv. 11, sphalm. = seq.
paniculatum, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 385.—Assam; Burma; Malaya.

MYRIOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 554 = **Abatia**, Ruiz & Pav. (Samyd.).
spicata, Turcz. l. c. 555.—Peruv.

MYRIPNOIS, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 112; Enum. Pl. Chin. Bor. 38. *COMPOSITAE*, Benth. & Hook. f. ii. 494.
dioica, *Bunge, Enum. Pl. Chin. Bor.* 38.—China.
uniflora, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 495.—China.

MYRISTICA, Linn. Gen. ed. II. 524 (1742). *MYRISTICEAE*, Benth. & Hook. f. iii. 136.
COMACUM, Adans. Fam. ii. 345 (1763).
HORSFIELDIA, Willd. Sp. Pl. iv. 872 (1805).
KNEMA, Lour. Fl. Cochinch. 604 (1790).
PYRRHOSA, Endl. Gen. 830 (1839).
SEBOPHORA, Neck. Elem. ii. 188 (1790).
VIROLA, Aubl. Pl. Gui. 904. t. 345 (1775).
acuminata, *Lam. in Mém. Acad. Sc. Par.* 1788 (1791), 164.—Madag.
americana, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 302 = fragrans?
amygdalina, J. Grah. Cat. Pl. Bomb. 175 = attenuata.
amygdalina, Kurz, For. Fl. Br. Burma, ii. 283 = glabra.
amygdalina, Thw. ex A. DC. in DC. Prod. xiv. 190 = laurifolia.
amygdalina, *Wall. Pl. As. Rar.* i. 79. t. 90.—Ind. or.
angustifolia, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 847 = missionis? *vel* glauca?
ardisiaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 31.—Ins. Philipp.
aromatica, Sw. Prod. Veg. Ind. Occ. 96; Lam. in Mém. Acad. Sc. Par. 1788 (1791) 155 = fragrans.
aruana, *Blume, Rumphia*, i. 191.—Ins. Molucc.
attenuata, *Wall. Cat.* n. 6791.—Ind. or.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 383.—Ins. Banca.
Bicuhyba, *Schott, in Spreng. Syst.* iv. *App.* 409.—Bras.
bracteata, *A. DC. in DC. Prod.* xiv. 192.—Malaya.
caesia, Zipp. ex Span. in Linnaea, xv. (1841) 346 = glauca.

MYRISTICA:—
calocarpa, Miq. Fl. Ind. Bat. i. II. 71 = elliptica.
calophylla, *Spruce, in Journ. Linn. Soc.* v. (1861) 4. —Bras.
canariformis, *Blume, Rumphia*, i. 191.—Ins. Amboin.
capitellata, *Poepp. ex A. DC. in DC. Prod.* xiv. 697.— Bras.
carinata, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 2.—Bras.
castaneaefolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 32. —Ins. Fiji.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 47.—Ins. Celebes.
ceylanica, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 29.—Zeylan.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 455.—Madag.
cimicifera, R. Br. Prod. 400 = insipida.
cinerea, *Poir. Encyc. Suppl.* iv. 35.—Ins. Molucc.
Commersoni, *Blume, Rumphia*, i. 181.—Hab.?
cordifolia, Mart. ex A. DC. in DC. Prod. xiv. 195 = sebifera.
cordifolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 46 = fatua.
corticosa, Bedd. Fl. Sylvat. t. 271 = attenuata.
corticosa, Hook. f. & Thoms. Fl. Ind. i. 158 = missionis, erratica, glauca.
costulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 48.— Ins. Celebes.
crassifolia, *Hook. f. & Thoms. Fl. Ind.* i. 160.— Malacca.
Cumaru, Poepp. ex Meissn. in Mart. Fl. Bras. v. I. 122 = M. Otoba.
cuspidata, *Benth. in Hook. Kew Journ.* v. (1853) 5.— Bras.
dactyloides, Gaertn. Fruct. i. 195 = fatua.
dactyloides, Wall. Cat. n. 6786 = malabarica.
debilis, *Spruce, ex A. DC. in DC. Prod.* xiv. 697.— Bras.
diospyrifolia, A. DC. in Ann. Sc. Nat. Sér. IV. iv. (1855) 29 = laurifolia.
diversifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 205. —Ins. Amboin.
elliptica, *Wall. Cat.* n. 6798.—Malaya.
elongata, *Benth. in Hook. Kew Journ.* v. (1853) 5.— Bras.
erratica, *Hook. f. & Thoms. Fl. Ind.* i. 156.—Reg. Himal.
eugeniaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 29.—Penang.
exaltata, Wall. Cat. n. 6804 = M. Irya, amygdalina.
Farquhariana, *Wall. l. c.* n. 6795.—Malaya.
fatua, Blume, Rumphia, i. t. 59 = Litsea obtusa.
fatua, *Houtt. Handleid.* iii. 337.—Ins. Molucc.
fatua, Sw. Prod. Veg. Ind. Occ. 96 = surinamensis.
ferruginea, Wall. Cat. n. 6803 = M. Horsfieldia.
Finlaysoniana, Wall. l. c. n. 6793 = Melodorum fulgens.
floribunda, Wall. l. c. n. 6805 = glabra.
fragrans, *Houtt. Handleid.* iii. 333.—Ins. Molucc.
furfuracea, *Hook. f. & Thoms. Fl. Ind.* i. 159.— Malaya.
Gardneri, *A. DC. in DC. Prod.* xiv. 197.—Bras.
geminata, *Miq. Fl. Ind. Bat. Suppl.* 385.—Burma; Malaya.
gibbosa. *Hook. f. & Thoms. Fl. Ind.* i. 158.—Reg. Himal.; Malaya.
glabra, *Blume, Bijdr.* 576.—Malaya.
glauca, *Blume, l. c.*—Burma; Malaya.
glauca, Zoll. ex A. DC. in DC. Prod. xiv. 206 = intermedia.
glaucescens, *Hook. f. & Thoms. Fl. Ind.* i. 157.— Malaya.
glaucescens, Wall. Cat. n. 6790 = Tetranthera venulosa.
globularia, *Blume, Rumphia*, i. 191. t. 64. f. 2.— Malaya.
globularia, *Hook. f. & Thoms. Fl. Ind.* i. 160 (non alior.).—Malaya.
globularia, Lam. in Mém. Acad. Sc. Par. 1788 (1791), 162 = microcarpa.
glomerata, Miq. Pl. Jungh. 171 = M. Horsfieldia.
gordoniaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 21, 30.—Peruv.
gracilis, *A. DC. l. c.* 30.—Bras.
grandifolia, *A. DC. in DC. Prod.* xiv. 194.—Ins. Fiji.
grandis, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 157.—Borneo.

MYRISTICA:—
guatemalensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 66.— Guatemala.
guatteriaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 30.—Ins. Philipp.
heterophylla, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 178.—Ins. Philipp.
Heyneana, *Wall. Cat.* n. 6789 (sp. dub.).—Ind. or.
Hookeriana, *Wall. l. c.* n. 6802 A.—Malaya.
Hookeriana, Wall. l. c. n. 6802 B (sp. indeterm.).— Malaya.
Horsfieldia, *Blume, Bijdr.* 577.—Zeylan.
Horsfieldii, Blume, Rumphia, i. 192 = praec.
Horsfieldii, Wall. Cat. n. 6806 = Wallichii.
Hostmanni, *Benth. in Hook. Kew Journ.* v. (1853) 7. —Guian.
hypargyraea, *A. Gray, Bot. U. St. Expl. Exped.* i. 33.— Ins. Navigat.
hypoleuca, *Spruce, in Journ. Linn. Soc.* v. (1861) 4. —Bras.
hypostícta, *Miq. Fl. Ind. Bat.* i. II. 55.—Java.
iners, *Blume, Bijdr.* 575.—Java.
insipida, *R. Br. Prod.* 400.—Austral.
integra, Wall. Cat. n. 6799 = glabra?
integrifolia, Steud. Nom. ed. II. ii. 174, sphalm. = *Myrica integrifolia.*
intermedia, *Blume, Rumphia*, i. 187.—Malaya.
inutilis, *Rich, ex A. Gray, Bot. U. St. Expl. Exped.* i. 34.—Ins. Samoa.
Iriagdehi, Steud. Nom. ed. II. ii. 174, sphalm. = *Iryaghedhi.*
Irya, *Gaertn. Fruct.* i. 195. t. 41.—Malaya; Zeylan.
Iryaghedhi, Gaertn. l. c. 196 = M. Horsfieldia.
iteophylla, *Miq. Fl. Ind. Bat.* i. II. 59.—Sumatra.
javanica, Blume, Bijdr. 576 = M. Irya.
Knema, Steud. Nom. ed. II. ii. 174 = furfuracea?
Kombo, *Baill. Adansonia*, ix. (1868) 79.—Afr. trop.
laevigata, *Blume, Rumphia*, i. 191. t. 64. f. 3.—Java.
lanceolata, Wall. Cat. n. 6794 = missionis.
lancifolia, *Poir. Encyc. Suppl.* iv. 35.—Ins. Molucc.
laurifolia, *Hook. f. & Thoms. Fl. Ind.* i. 163.— Zeylan.
laurifolia, Spruce, ex A. DC. in DC. Prod. xiv. 199 = Sprucei.
laurina, *Blume, Rumphia*, i. 189. t. 61.—Malaya.
Lemanniana, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 31.—Malacca.
lepidota, *Blume, Rumphia*, i. 183. t. 57.—N. Guinea.
leucoxyla, *Miq. Fl. Ind. Bat. Suppl.* 385.—Sumatra.
linifolia, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 847 = erratica *vel* longifolia?
litoralis, *Miq. Fl. Ind. Bat.* i. II. 57.—Java.
longifolia, Hook. f. & Thoms. Fl. Ind. i. 156 = furfuracea.
longifolia, *Wall. Cat.* n. 6801.—Reg. Himal.; Burma.
luzonica, Blanco, Fl. Filip. ed. I. 664 = bracteata.
macrocarpa, *Blume, Rumphia*, i. 185.—Ins. Philipp.
macrocoma, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 207. —Ins. Molucc.
macrophylla, A. Gray, Bot. U. St. Expl. Exped. i. 33 = grandifolia.
macrophylla, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 846 (sp. dub.).—Ins. Molucc.?
macrophylla, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 6.—Bras.
macrothyrsa, *Miq. Pl. Jungh.* 172.—Sumatra.
madagascariensis, *Lam. in Mém. Acad. Sc. Par.* 1788 (1791) 163.—Madag.
madagascariensis, Vent. ex A. DC. in DC. Prod. xiv. 192 = bracteata.
magnifica, *Bedd. Fl. Sylv.* ii. t. 268.—Travancor.
malabarica, *Lam. in Mém. Acad. Sc. Par.* 1788 (1791) 162.—Ind. or.
Mandaharan, *Miq. Fl. Ind. Bat. Suppl.* 384.—Sumatra.
Mannii, *Benth. in Hook. Ic. Pl.* t. 1262.—Afr. trop.
mascula, *Reinw. ex De Vriese, Pl. Ind. Bat. Or.* 93.— Malaya.
membranacea, *Poepp. ex A. DC. in DC. Prod.* xiv. 196. —Bras.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* iii. 67.— Mexic.
micrantha, *Wall. Cat.* n. 6807.—Siam.
microcarpa, *Willd. Sp. Pl.* iv. 871, partim.—Ins. Molucc.; Cochinch.
microcephala, *Benth. in Hook. Ic. Pl.* t. 1261.—Ins. Fernando Po.
missionis, *Wall. Cat.* n. 6788.—Burma; Malaya.

MYRISTICA:—
Mocoa, *Poepp. ex A. DC. in DC. Prod.* xiv. 195.—Bras.
mollissima, *Poepp. l. c.* 696.—Bras.
montana, *Roxb. Hort. Beng.* [105]; *Fl. Ind.* iii. 846. —Malaya.
montana, Wall. Cat. n. 6792 = iners.
morindaefolia, Blume, Rumphia, i. 186 = fatua.
moschata, Thunb. in Vet. Akad. Handl. Stockh. (1782) 49 = fragrans.
myrmecophila, *Becc. Malesia*, ii. (1884) 37.—N. Guin.
nesophila, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 206.— Malaya.
Niohue, *Baill. Adansonia*, ix. (1868) 79.—Afr. trop.
notha, Wall. Cat. n. 6787 = malabarica.
obtusifolia, Wall. l. c. n. 6808 (Laurinea?).—Singapore.
officinalis, Benth. ex A. DC. in DC. Prod. xiv. 197 = Gardneri.
officinalis, Linn. f. Suppl. 265 = fragrans.
officinalis, *Mart. Reise Bras.* i. 364, 543.—Bras.
orinocensis, H. B. & K. Nov. Gen. et Sp. vii. 191 (Perae sp.).—Venezuela.
Otoba, *Humb. & Bonpl. Pl. Aequin.* ii. 78. t. 103.— N. Granat.
palembanica, Miq. Fl. Ind. Bat. Suppl. 384 = glaucescens.
panamensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 67.— Panama.
paniculata, A. DC. in Ann. Sc. Nat. Sér. IV. iv. (1855) 31 = Farquhariana.
papuana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 46. N. Guin.
parviflora, *Roxb. Fl. Ind.* iii. 847 (Quid?).—Ins. Molucc.
pavonis, *A. DC. in DC. Prod.* xiv. 697.—Bras.
peltata, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 846.—Ins. Molucc.
peruviana, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 30.—Peruv.
philippensis, *Lam. in Mém. Acad. Sc. Par.* 1788 (1791) 161.—Ins. Philipp.
pinnaeformis, *Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 49.—N. Guin.
platysperma, *Spruce, ex A. DC. in DC. Prod.* xiv. 695. —Bras.
pulchra, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 51.— Borneo.
punctata, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 6.—Bras.
punicifolia, Roxb. Hort. Beng. [105], nomen = parviflora?
Radja, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 206.—Ins. Molucc.
rufula, Mart. ex A. DC. in DC. Prod. xiv. 696 = cuspidata.
rugulosa, *Spruce, in Journ. Linn. Soc.* v. (1861) 4.— Bras.
Sagotiana, *Benth. in Hook. Ic. Pl.* t. 1260.—Guian.
salicifolia, Willd. Sp. Pl. iv. 871 = sylvestris.
sapida, Steud. Nom. ed. II. ii. 175, sphalm. = Myrica Nagi.
sebifera, Seem. Bot. Voy. Herald, 75 = panamensis.
sebifera, *Sw. Prod. Veg. Ind. Occ.* 96.—Am. trop.
sesquipedalis, Wall. Cat. n. 6809 = Actinodaphne sesquipedalis.
sessilis, *A. DC. in DC. Prod.* xiv. 198.—Bras.
simiarum, *A. DC. in Ann. Sc. Nat.* Sér. IV. iv. (1855) 29.—Ins. Philipp.
spadicea, Blume, Bijdr. 577 = fatua.
Spanogheana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 47. —Ins. Timor.
sphaerocarpa, Wall. Pl. As. Rar. i. 79. t. 89 = M. Irya.
spicata, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 847.—Ins. Molucc.
Sprucei, *A. DC. in DC. Prod.* xiv. 199.—Bras.
subalulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 47. —N. Granat.
subcordata, Blume, Rumphia, i. 186 = fatua.
subglobosa, *Miq. Fl. Ind. Bat. Suppl.* 383.—Sumatra.
sublanceolata, Miq. l. c. i. II. 58 = iners.
subsessilis, *Benth. in Hook. Kew Journ.* v. (1853) 4.— Bras.
subtilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 50.—N. Guinea.
succedanea, *Blume, Rumphia*, i. 186.—Malaya.
sumatrana, Blume, l. c. 187 = glauca.
superba, *Hook. f. & Thoms. Fl. Ind.* i. 162.—Malaya.
surinamensis, *Roland. ex Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 281.—Ind. occ.

MYRISTICA:—
sycocarpa, Miq. Fl. Ind. Bat. i. II. 68 = elliptica.
sylvestris, *Houtt. Handleid.* iii. 340.—Ins. Molucc.
sylvestris, Sieber, ex A. DC. in DC. Prod. xiv. 192 = bracteata.
Teysmanni, *Miq. Fl. Ind. Bat.* i. II. 57.—Java.
theiodora, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 6.—Bras.
tingens, *Blume, Rumphia*, i. 190.—Ins. Amboin.
tomentosa, Blume, Bijdr. 577 = laurina.
tomentosa, J. Grah. Cat. Pl. Bomb. 175 = malabarica.
tomentosa, *Hook. f. & Thoms. Fl. Ind.* i. 161.—Malaya.
tomentosa, Thunb. in Vet. Akad. Handl. Stockh. (1782) 49 = fatua.
tomentosa, Thw. ex A. DC. in Ann. Sc. Nat. Sér. IV. iv. (1855) 29 = laurifolia.
tubiflora, *Blume, Rumphia*, i. 182. t. 56.—N. Guin.
uapensis, *Spruce, ex A. DC. in DC. Prod.* xiv. 696.—Bras.
uviformis, *Lam. in Mém. Acad. Sc. Par.* 1788 (1791) 166.—Ins. Molucc.
valida, *Miq. Fl. Ind. Bat.* i. II. 67.—Sumatra.
venosa, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.; Peruv.
Virola, Raeusch. Nom. ed. III. 292 = sebifera.
Vouri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 454.—Madag.
Vrieseana, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 49 = M. Irya.
Wallichii, *Hook. f. & Thoms. Fl. Ind.* i. 161.—Malaya.
zeylanica, Thw. Enum. Pl. Zeyl. 11, 399 = laurifolia.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 50.—N. Guin.

MYRMECHIS, Blume, Orch. Archip. Ind. 76. t. 21 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 601.
glabra, *Blume, l. c.* 78.—Java.
gracilis, *Blume, l. c.* 77.—Java.

MYRMECIA, Schreb. Gen. 74 (1789) = **Tachia**, Aubl. (Gentian.).
calygona, Steud. Nom. ed. II. ii. 175 = T. calygona.
scandens, Willd. Sp. Pl. i. 623 = T. guianensis.
Tachia, J. F. Gmel. Syst. 247 = T. guianensis.

MYRMECODIA, Jack, in Trans. Linn. Soc. xiv. (1823) 122. *RUBIACEAE*, Benth. & Hook. f. ii. 132.
alata, *Becc. Malesia*, ii. (1884) 106.—N. Guin.
Albertisii, *Becc. l. c.* 112.—N. Guin.
amboinensis, Becc. l. c. 97 = Rumphii?
Antonii, *Becc. l. c.* 116.—Austral.
armata, DC. Prod. iv. 450 = tuberosa.
aruensis, *Becc. Malesia*, ii. (1884) 108.—Ins. Aru.
bullosa, *Becc. l, c.*—N. Guin.
echinata, Antoine, in Oestr. Bot. Zeitschr. xxxii. (1882) 347 = Antonii.
echinata, *Gaudich. in Freyc. Voy. Bot.* 472.—Ins. Rawak et Aru.
echinata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 257 = menadensis.
echinata, Miq. Fl. Ind. Bat. ii. 310 = tuberosa.
echinata, F. Muell. Papuan Pl. 90 = Muelleri.
erinacea, *Becc. Malesia*, ii. (1884) 105.—N. Guin.
glabra, *Britt. in Proc. Linn. Soc.* (1879–80) p. liii. (Hydnophyti sp. ?).—Borneo.
goramensis, *Becc. Malesia*, ii. (1884) 118.—Ins. Molucc.
hispida, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 91, sphalm. = echinata.
imberbis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 42.—Ins. Fiji.
inermis, Benth. & Hook. f. Gen. ii. 132, in nota = imberbis.
inermis, DC. Prod. iv. 450 = tuberosa.
inermis, Gaudich. in Freyc. Voy. Bot. 472. t. 95 = Hydnophytum Gaudichaudii.
jobiensis, *Becc. Malesia*, ii. (1884) 111.—Ins. Jobi.
kandariensis, *Becc. l. c.* 100.—Ins. Celebes.
menadensis, *Becc. l. c.* (1885) 176.—Ins. Celebes.
Muelleri, *Becc. l. c.* (1884) 102.—N. Guin.
oninensis, *Becc. l. c.* 110.—N. Guin.
platyrea, *Becc. l. c.* 115.—N. Guin.
pulvinata, *Becc. l. c.* 103.—N. Guin.
Rumphii, *Becc. l. c.* 117.—Malaya.
salomonensis, *Becc. l. c.* (1885) 175.—Ins. Salomon.

MYRMECODIA:—
selebica, Becc. in Nuov. Giorn. Bot. Ital. vi. (1874) 195 = Myrmephytum selebicum.
tuberosa, *Jack, in Trans. Linn. Soc.* xiv. (1823) 123.—Malaya.
vitiensis, Seem. in Bonplandia, ix. (1861) 256; x. (1862) 36 = Hydnophytum longiflorum.

MYRMECYLON, Hook. & Arn. Bot. Beech. Voy. 186 (1833) sphalm. = **Memecylon**, Linn. (Melastom.).

MYRMEDOMA, Becc. Malesia, ii. (1884) 94. t. 10. *RUBIACEAE*.
arfakiana, *Becc. l. c.*—N. Guin.

MYRMEPHYTUM, Becc. Malesia, ii. (1884) 92. tt. 8, 9. *RUBIACEAE*.
selebicum, *Becc. l. c.* 93.—Ins. Celebes.

MYRMIDONE, Mart. Nov. Gen. et Sp. iii. 149. t. 279 (1829). *MELASTOMACEAE*, Benth. & Hook. f. i. 767.
macrosperma, *Mart. l. c.* 149.—Bras.
rotundifolia, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Microphysa rotundifolia.

MYROBALANUS, Gaertn. Fruct. ii. 90. t. 97 (1791) = **Terminalia**, Linn. (Combret.).
angustifolia, Kostel. Allg. Med.-pharm. Fl. iv. 1497 = T. angustifolia.
Badamia, Poir. Encyc. Suppl. iii. 707 = T. Badamia.
Bellirica, Gaertn. Fruct. ii. 90. t. 97 = T. Bellerica.
Chebula, Gaertn. l. c. 91 = T. Chebula.
citrina, Gaertn. l. c. = T. citrina.
Fatraea, Poir. Encyc. Suppl. iii. 708 = T. Fatraea.
gangetica, Kostel. Allg. Med.-Pharm. Fl. iv. 1497 = T. Chebula.
indica, Buch.-Ham. ex Steud. Nom. ed. II. ii. 151 = Scleropyrum Wallichianum.
rhomboidea, Poir. Encyc. Suppl. iii. 708 = T. rhomboidea.
Terminalia, Poir. l. c. 707 = T. Badamia.

MYROBROMA, Salisb. Parad. Lond. t. 82 (1807) = **Vanilla**, Sw. (Orchid.).
fragrans, Salisb. l. c. = V. planifolia.

MYROCARPUS, Allem. Diss. (1847). *LEGUMINOSAE*, Benth. & Hook. f. i. 559.
fastigiatus, *Allem. Diss.* (1847).—Bras.
frondosus, *Allem. l. c.* (1848).—Bras.

MYRODENDRON, Schreb. Gen. 358 (1789) = **Humiria**, Aubl.
amplexicaule, Willd. Sp. Pl. ii. 1171 = H. balsamifera.
balsamiferum, J. F. Gmel. Syst. 817 = H. balsamifera.
crassifolium, Spreng. Syst. iv. Cur. Post. 345 = H. crassifolia.
floribundum, Spreng. l. c. = H. floribunda.
subvaginale, Mart. ex Urban, in Mart. Fl. Bras. xii. II. 441 = H. balsamifera.

MYRODIA, Sw. Prod. Veg. Ind. Occ. 102 (1788). *STERCULIACEAE*, Benth. & Hook. f. i. 219.
LEXARZA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 7 (1825).
angustifolia, Mart. in Flora, xxii. (1839) I. Beibl. 17 = Quararibea turbinata.
breviflora, Benth. in Journ. Linn. Soc. vi. (1862) 115 = Scleronema Spruceana.
Cacao, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 324.—N. Granat.
floribunda, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 212 (Quararibeae sp.).—Bras.
funebris, *Benth. in Journ. Linn. Soc.* vi. (1862) 115.—Mexic.
longiflora, Sw. Prod. Veg. Ind. Occ. 103 = Quararibea guianensis.
macrophylla, *Klotzsch, in Schomb. Fauna & Fl. Brit. Gui.* 1091, nomen.—Guiana.
Martii, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 212 = Quararibea turbinata.
multiflora, *Steud. in Flora*, xxvi. (1843) 755.—Guiana.
ovata, Moç. & Sesse, ex DC. Prod. i. 447 = Quararibea turbinata.
parviflora, Benth. in Journ. Linn. Soc. vi. (1862) 109 = Scleronema Spruceana.

MYRODIA:—
penduliflora, A. St. Hil. Fl. Bras. Mer. i. 269. t. 53 (Quararibeae sp.).—Bras.
scabra, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 19.—Bras.
trinervia, *Mart. l. c.* 17.—Bras.
turbinata, Sw. Prod. Veg. Ind. Occ. 102 = Quararibea turbinata.
verticillaris, [*Moç. & Sesse,*] *ex DC. Prod.* i. 477.—Mexic.

MYROSMA, Linn. f. Suppl. 8, 80 (1781). *SCITAMINEAE*, Benth. & Hook. f. iii. 651.
THALIANTHUS, Klotzsch, in Rich. Schomb. Reise Brit. Gui. iii. (1848) 1125.
cannaefolium, *Linn. f. Suppl.* 80.—Guiana.
cannaeforme, Willd. Sp. Pl. i. 13 = praec.
comosum, *Spreng. Syst.* i. 9.—Guiana.
madagascariense, *Benth. in Benth. & Hook. f. Gen.* iii. 651.—Madag.
setosum, *Benth. l. c.*—Bras.
unilaterale, Benth. & Hook. f. l. c.—Peruv.

MYROSMODES, Reichb. f. Xenia Orch. i. 19. t. 8 (1854) = **Altensteinia**, H. B. & K. (Orchid.).
nubigenum, Reichb. f. l. c. = A. nubigena.

MYROSPERMUM, Jacq. Enum. Pl. Carib. 4 (1760). *LEGUMINOSAE*, Benth. & Hook. f. i. 558.
CALUSIA, Bert. ex Steud. Nom. ed. II. i. 262 (1840).
balsamiferum, Ruiz & Pav. Ic. Ined. t. 373.—Peruv.?
emarginatum, Klotzsch, in Bonplandia, v. (1857) 277 = frutescens.
Erythroxylum, Allem. in Rev. Braz. i. (1857) 51 = Myroxylon peruiferum.
frutescens, *Jacq. Enum. Pl. Carib.* 20.—Am. austr.
pedicellatum, Lam. Encyc. iv. 191 = Myroxylon peruiferum.
Pereirae, Royle, Man. Mat. Med. ed. II. 414 = Myroxylon Pereirae.
peruiferum, DC. Prod. ii. 95 = Myroxylon peruiferum.
pubescens, DC. l. c. = Myroxylon pubescens.
punctatum, Walp. Rep. i. 805 = Myroxylon toluiferum.
robinifolium, Warsz. ex Klotzsch, in Bonplandia, v. (1857) 276 = Myroxylon robiniaefolium.
secundum, *Klotzsch, l. c.* 277.—Venezuela.
toluiferum, DC. Prod. ii. 95 = Myroxylon toluiferum.

MYROTHAMNUS, Welw. Apont. 578 (1858). *HAMAMELIDEAE*, Benth. & Hook. f. i. 1005.
flabellifolia, *Welw. l. c.*—Afr. trop.
moschata, Baill. Adansonia, ix. (1870) 330 = Myosurandra moschata.

MYROXYLON, Forst. Char. Gen. 125 (1776) = **Xylosma**, Forst. (Bixin.).
orbiculatum, Forst. l. c. 126 = X. orbiculatum.
suaveolens, Forst. l. c. = X. suaveolens.

MYROXYLON, Linn. f. Suppl. 34 (1781). *LEGUMINOSAE*, Benth. & Hook. f. i. 558.
TOLUIFERA, Linn. Gen. ed. II. 182 (1742).
abruptifolium, Stokes, Bot. Mat. Med. ii. 471 = peruiferum.
dicline, *Blanco, Fl. Filip.* ed. I. 813.—Ins. Philipp.
frutescens, Willd. Sp. Pl. ii. 546 = Myrospermum frutescens.
Hanburyanum, *Klotzsch, in Bonplandia*, v. (1857) 275.—Ecuador.
pedicellatum, *Klotzsch, l. c.* 276.—Peruv.
Pereirae, *Klotzsch, l. c.* 274.—Am. centr.
peruiferum, *Linn. f. Suppl.* 233.—Ecuador; Peruv.; Bras.
pubescens, *H. B. & K. Nov. Gen. et Sp.* vi. 374.—N. Granat.
punctatum, Klotzsch, in Hayne, Arzneigew. xiv. t. 12 = toluiferum.
robiniaefolium, *Klotzsch, in Bonplandia*, v. (1857) 276.—Ecuador.
toluiferum, *H. B. & K. Nov. Gen. et Sp.* vi. 375.—Am. trop.
trialatum, Stokes, Bot. Mat. Med. ed. II. 473 = Cinchona officinalis.

MYROXYLUM, Schreb. Gen. i. 281 (1798) = praec.

MYRRHA, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 217. *UMBELLIFERAE* (Quid?).

MYRRHIDIUM, Eckl. & Zeyh. Enum. 71 (1835) = **Pelargonium**, L'Hérit. (Geran.).
anemonefolium, Eckl. & Zeyh. l. c. 71 = P. myrrhifolium.
artemisiaefolium, Eckl. & Zeyh. l. c. 72 = P. artemisiaefolium.
athamanthoides, Eckl. & Zeyh. l. c. = P. athamanthoides.
betonium, Eckl. & Zeyh. l. c. 71 = P. myrrhifolium.
bullatum, Eckl. & Zeyh. l. c. = P. myrrhifolium.
candicans, Eckl. & Zeyh. l. c. = P. candicans.
coriandrifolium, Eckl. & Zeyh. l. c. 72 = P. myrrhifolium.
fruticosum, Eckl. & Zeyh. l. c. = P. fruticosum.
lacerum, Eckl. & Zeyh. l. c. = P. myrrhifolium.
longicaule, Eckl. & Zeyh. l. c. 71 = P. longicaule.
multicaule, Eckl. & Zeyh. l. c. = P. multicaule.
senecioides, Eckl. & Zeyh. l. c. 72 = P. senecioides.
triangulare, Eckl. & Zeyh. l. c. 71 = P. triangulare.
urbanum, Eckl. & Zeyh. l. c. = P. urbanum.

MYRRHINA, Rupr. in Mém. Acad. Pétersb. Sér. VII. xv. n. II. (1869) (Fl. Cauc.) 277 = **Erodium**, L'Hérit. (Geran.).
absinthoides, Rupr. Fl. Cauc. 277, in textu = E. absinthoides.
alpina, Rupr. l. c. 278, in textu = E. alpinum.
anthemoides, Rupr. l. c. 277, in textu = E. anthemidifolium.
fumarioides, Rupr. l. c. 277 = E. absinthoides.
gunibensis, Rupr. l. c. 278 = E. absinthoides.
inodora, Rupr. l. c. 277 = E. cicutarium.

MYRRHINIUM, Schott, in Spreng. Syst. iv. Cur. Post. 404 (1827). *MYRTACEAE*, Benth. & Hook. f. i. 717.
FELICIANEA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 375. t. 157 (1829).
TETRASTEMON, Hook. Bot. Misc. iii. (1833) 317.
atropurpureum, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 404.—Bras.
peruvianum, *Berg, in Linnaea*, xxvii. (1854) 438.—Peruv.
rubriflorum, *Berg, in Mart. Fl. Bras.* xiv. 1. 466.—Bras.
sarcopetalum, *Lem. in Rev. Hortic.* Sér. IV. vii. (1858) 444.—Bras.

MYRRHIS, [Tourn.] Linn. Syst. ed. I (1735), partim; Scop. Fl. Carn. ed. II. i. 207 (1772). *UMBELLIFERAE*, Benth. & Hook. f. i. 897.
GLYCOSMA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 639 (1840).
LINDERA, Adans, Fam. ii. 499 (1763).
alpina, Steud. Nom. ed. I. 545 = Anthriscus sylvestris.
ammoides, Spreng. in Schult. Syst. vi. 517 = Conopodium denudatum.
andicola, H. B. & K. Nov. Gen. et Sp. v. 13. t. 419 = Oreomyrrhis andicola.
andina, Hemsl. Biol. Centr. Am. Bot. i. 567, sphalm. = praec.
angelicaefolia, Schult. Syst. vi. 509 = Chaerophyllum angelicaefolium.
arborescens, Spreng. Umbellif. Prod. 29 = Aralia spinosa?
aristata, Spreng. Spec. Umbellif. 133 = Osmorhiza Claytoni.
aromatica, Spreng. in Schult. Syst. vi. 509 = Chaerophyllum aromaticum.
aurea, All. Fl. Pedem. ii. 29 = Chaerophyllum aureum.
australis, All. l. c. = Scandix australis.
Berterii, D. Dietr. Syn. Pl. ii. 984 = Osmorhiza Berteri.
bifida, Spreng. Umbellif. Prod. 29 = Chaerophyllum procumbens.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 346.—Calif.
brevipedunculata, *Hoffm. Gen. Umbell.* ed. I. 30.—Reg. Caucas.
brevistylis, D. Dietr. Syn. Pl. ii. 984 = Osmorhiza brevistylis.
bulbosa, All. Fl. Pedem. ii. 29 = Anthriscus alpestris.
bulbosa, Spreng. Umbellif. Prod. 29 = Chaerophyllum bulbosum.
Bunium, Spreng. Spec. Umbellif. 131 = Conopodium denudatum.
canadensis, Gaertn. Fruct. i. 109 = Cryptotaenia canadensis.

MYRRHIS:—
capensis, Spreng. Spec. Umbellif. 32 = Annesorhiza capensis.
capilliflora, Guss. Fl. Sic. Prod. i. 351 = Conopodium capillifolium.
chaerophyllea, Lam. Fl. Fr. iii. 442 = Anthriscus vulgaris.
chaerophylloides, *Hance, in Journ. Bot.* xvi. (1878) 108.—China.
Cicutaria, Spreng. Umbellif. Prod. 29 = Chaerophyllum hirsutum.
Cicutaria, Tenore, Fl. Nap. iii. 320, partim = Chaerophyllum calabricum.
clavata, Spreng. Syst. i. 903 = Chaerophyllum macrospermum.
Claytoni, D. Dietr. Syn. Pl. ii. 984 = Osmorrhiza longistylis.
Claytoni, Michx. Fl. Bor. Am. i. 170 = Osmorrhiza brevistylis.
colorata, Spreng. in Schult. Syst. vi. 515 = Chaerophyllum coloratum.
cynapioides, Guss. Rar. 127. t. 24 = Carum cylindricum?
gilanica, Schult. Syst. vi. 520 = Chaerophyllum macrospermum.
gracilis, Spreng. Syst. iv. Cur. Post. 120 = Conopodium Cyminum.
hirsuta, All. Fl. Pedem. ii. 28 = Chaerophyllum hirsutum.
humilis, Schult. Syst. vi. 519 = Chaerophyllum humile.
iberica, Hoffm. Gen. Umbellif. ed. I. 30 = brevipedunculata.
longifolia, Spreng. ex DC. Prod. iv. 232 = Chaerophyllum millefolium.
longistylis, Torr. Fl. U. St. i. 310 = Osmorrhiza longistylis.
maculata, Sweet, Hort. Brit. ed. I. 189 = Chaerophyllum aureum.
magellensis, Bertol. Fl. Ital. iii. 213 = Chaerophyllum magellense.
millefolia, Spreng. in Schult. Syst. vi. 519 = Chaerophyllum millefolium.
nodosa, Lam. Fl. Fr. iii. 441 = Chaerophyllum nodosum.
occidentalis, *Benth. & Hook. f. Gen.* i. 897.—Am. bor.
odorata, *Scop. Fl. Carn.* ed. II. i. 207.—Europ. austr.; Reg. Caucas.
Pecten-Veneris, All. Fl. Pedem. ii. 29 = Scandix Pecten-Veneris.
procumbens, Spreng. Pugill. ii. 56 = Chaerophyllum procumbens.
pyrenaica, Spreng. in Schult. Syst. vi. 518 = Conopodium denudatum.
rosea, Spreng. l. c. 519 = Chaerophyllum roseum.
scabra, Spreng. Syst. i. 902 = Caucalis Anthriscus.
sulcata, Lag. Gen. et Sp. Nov. 13 = odorata.
sylvestris, Spreng. Umbellif. Prod. 29 = Anthriscus sylvestris.
temula, All. Fl. Pedem. ii. 29 = Chaerophyllum temulum.
tenerrima, Presl, Delic. Prag. 131 = Conopodium denudatum.
tenuifolia, Schult. Syst. vi. 520 = Chaerophyllum tenuifolium.
tenuifolia, Spreng. Nov. Prov. 28 = praec.?
ternata, Moench, Meth. 101 = Cryptotaenia canadensis.
torquata, Schult. Syst. vi. 514 = Anthriscus alpestris.
tuberosa, Jundz. Opis. Rosl. w Litw. 118, sphalm. = Chaerophyllum bulbosum?

MYRSINE, Linn. Syst. ed. I (1735); Gen. ed. I. 54 (1737). *MYRSINEAE*, Benth. & Hook. f. ii. 642.
AGERIA, Adans. Fam. ii. 166 (1763).
ATHRUPHYLLUM, Lour. Fl. Cochinch. 120 (1790).
CABALLERIA, Ruiz & Pav. Prod. 141. t. 30 (1794).
DUHAMELIA, Domb. ex Lam. Encyc. i. 245 (1783).
FIALARIS, Rafin. Sylva Tellur. 166 (1838).
HEBERDENIA, Banks, ex A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 79.
HEURLINIA, Rafin. Sylva Tellur. 166 (1838).
MANGLILLA, Juss. Gen. 151 (1789).
MERISTA, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 47.
NIARA, Dennst. Schluess. Hort. Malab. 31 (1818).
PLEIOMERIS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 87.
PLOTIA, Adans. Fam. ii. 226 (1763).

MYRSINE:—
RAPANEA, Aubl. Pl. Gui. i. 121. t. 46 (1775).
ROEMERIA, Thunb. Nov. Gen. ix. 130 (1798), pro parte.
SAMARA, Sw. Prod. Veg. Ind. Occ. 32 (1788); Fl. Ind. Occ. i. 261 (1797).
SCLEROXYLUM, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 57; et Enum. Hort. Berol. 249 (1809).
SUTTONIA, A. Rich. Fl. Nov. Zél. 349. t. 38 (1832).
achradifolia, *F. Muell. Fragm.* vi. 164.—Austral.
acuminata, Royle, Illustr. Bot. Himal. 265 = semiserrata.
acuta, Salisb. Prod. 141 = africana.
affinis, *A. DC. in DC. Prod.* viii. 96.—Malaya.
africana, *Linn. Sp. Pl.* 196.—Reg. Himal.; Afr. austr. et trop.; Ins. Azor.
angustifolia, D. Dietr. Syn. Pl. i. 619 = Embelia angustifolia.
ardisioides, H. B. & K. Nov. Gen. et Sp. iii. 249 = Ardisia decipiens.
ardisioides, *Hook. & Arn. Bot. Beech. Voy.* 197.—China.
Athyrophyllum, *R. Br. Prod.* 533; *Roem. & Schult. Syst.* iv. 509.—Cochinch.
aurantiaca, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 300 = Embelia viridiflora.
avenis, A. DC. in Trans. Linn. Soc. xvii. (1834) 107 = capitellata.
baccata, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 86.—Porto Rico.
bahiensis, Casar. Nov. Stirp. Bras. Dec. 53 = parvifolia.
Barthesia, Roem. & Schult. Syst. iv. 507 = Ardisia insularis.
Berterii, A. DC. in Trans. Linn. Soc. xvii. (1834) 109 = laeta.
bifaria, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 296 = africana.
borneensis, *Scheff. Comm. Myrsin.* 50.—Borneo.
bottensis, *A. DC. in DC. Prod.* viii. 93.—Abyssin.
Brackenridgei, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 330.—Ins. Fiji.
brasiliensis, *A. DC. in DC. Prod.* viii. 103.—Bras.
buxifolia, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 225.—China.
campanulata, *F. Muell. Fragm.* vi. 235.—Austral.
canariensis, *Spreng. Syst.* i. 663.—Ins. Canar.
capitellata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 295.—As. trop.; Malaya.
Capororoca, Casar. Nov. Stirp. Bras. Dec. 54 = laeta.
chathamica, *F. Muell. Veg. Chath. Isl.* 38.—Ins. Chatham.
ciliata, H. B. & K. Nov. Gen. et Sp. iii. 248 = dependens.
cochin-chinensis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 84.—Cochinch.
collina, *Nadeaud, Enum. Pl. Tahiti*, 61.—Ins. Tahiti.
cordata, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 422.—Malaya.
coriacea, *R. Br. Prod.* 533.—Am. trop.
coriacea, *Nadeaud, Enum. Pl. Tahiti*, 61.—Ins. Tahiti.
coriacea, Sieber, ex A. DC. in Trans. Linn. Soc. xvii. (1834) 108 = laeta.
crassa, D. Dietr. Syn. Pl. i. 619 = Ardisia insularis.
crassifolia, *R. Br. Prod.* 534.—Austral.
cubana, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 86 = coriacea, *R. Br.*
Daphnites, Mart. in Flora, xxiv. (1841) II. Beibl. 17 = umbellata.
densiflora, *Scheff. Comm. Myrsin.* 50.—Malaya.
dentata, *Spreng. Syst.* i. 664.—Peruv.
dependens, *Spreng. l. c.*—Peruv.
divaricata, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 47.—N. Zel.
emarginella, *Miq. in Mart. Fl. Bras.* x. 312.—Bras.
erythroxyloides, *Benth. Bot. Voy. Sulph.* 123.—Am. trop.
excelsa, D. Don, Prod. Fl. Nep. 147 = capitellata.
excelsa, Link, Enum. Hort. Berol. i. 196 = Ardisia excelsa.
falcata, *Nadeaud, Enum. Pl. Tahiti*, 61.—Ins. Tahiti.
ferruginea, *Spreng. Syst.* i. 664.—Peruv.
flocculosa, Mart. in Flora, xxiv. (1841) II. Beibl. 17 = laeta.
floribunda, *R. Br. Prod.* 533.—Am. trop.
floridana, A. DC. in Trans. Linn. Soc. xvii. (1834) 107 = floribunda.
fragilis, Casar. Nov. Stirp. Bras. Dec. 53 = Gardneriana.

MYRSINE:—
Gardneriana, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 86.—Bras.
Gaudichaudii, *A. DC. l. c.* 85.—Ins. Sandvic.
Gilliana, *Sond. in Linnæa*, xxiii. (1850) 76.—Afr. austr.
glabra, Gaertn. Fruct. i. 282 = africana.
glauca, Casar. Nov. Stirp. Bras. Dec. 54 = umbellata.
Glazioviana, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 17.—Bras.
glomeriflora, Mart. in Flora, xxiv. (1841) II. Beibl. 19 = umbellata.
Grisebachii, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 365.—Reg. Argent.
Hasseltii, *Blume, ex Scheff. Comm. Myrsin.* 49.—Java.
Heberdenia, Roem. & Schult. Syst. iv. 538 = Ardisia excelsa.
Hookeriana, Steud. Nom. ed. II. ii. 176 = ardisioides.
Icacorea, Roem. & Schult. Syst. iv. 508 = Ardisia acuminata.
ilicifolia, *Lodd. ex Steud. Nom.* ed. II. ii. 176.—Austral.
insularis, D. Dietr. Syn. Pl. i. 619 = Ardisia insularis.
Kellau, Hochst. in Flora, xxiv. (1841) I. Intell. 25 = Euclea Kellau.
khasyana, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 21 = semiserrata.
Korthalsii, *Miq. Pl. Jungh.* 189.—Java.
laeta, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 112.—Am. trop.
lanceolata, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 192.—N. Caled.
lanceolata, Wall. Cat. n. 2297 = capitellata.
lancifolia, *Mart. Herb. Fl. Bras.* n. 1193.—Bras.
latifolia, *Spreng. Syst.* i. 664.—Peruv.
laurifolia, *Casar. Nov. Stirp. Bras. Dec.* 55.—Bras.
lepidocarpon, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 512 = capitellata.
Lessertiana, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 85.—Ins. Sandvic.
leuconeura, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 18.—Bras.
linearis, Poir. Encyc. Suppl. iii. 709 = M. Athyrophyllum.
longifolia, *Nadeaud, Enum. Pl. Tahiti*, 61.—Ins. Tahiti.
lucida, Wall. Cat. n. 2298 = capitellata.
Lundiana, *A. DC. in DC. Prod.* viii. 114.—Bras.?
madagascariensis, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 107.—Madag.
Manglilla, *R. Br. Prod.* 533; *Roem. & Schult. Syst.* iv. 509.—Peruv.
marginata, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 283.—Bras.
maritima, Casar. Nov. Stirp. Bras. Dec. 56 = floribunda.
Martiana, A. DC. in DC. Prod. viii. 101 = umbellata.
melanophleos, *R. Br. Prod.* 533.—Afr. austr.
micrantha, D. Dietr. Syn. Pl. i. 619 = Embelia micrantha.
mitis, *Spreng. Syst.* i. 663.—Afr. austr.
montana, *Hook. f. Handb. N. Zeal. Fl.* 184.—N. Zel.
monticola, Mart. in Flora, xxiv. (1841) II. Beibl. 18 = umbellata.
multiflora, Herb. Vindob. ex Miq. in Mart. Fl. Bras. x. 310 = leuconeura.
myricaefolia, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 330.—Ins. Pacif.
myricoides, Schlecht. in Linnaea, viii. (1833) 525 = laeta.
myrtifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 112.—Am. austr.
Myrtillus, Hook. Ic. Pl. t. 825 = Embelia Myrtillus.
myrtoides, *Hook. l. c.* t. 87.—N. Granat.
neriifolia, Casar. Nov. Stirp. Bras. Dec. 55 = umbellata, lancifolia.
neriifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 137 = capitellata.
nummularia, *Hook. f. Handb. N. Zeal. Fl.* 184.—N. Zel.
oblonga, Spreng. Syst. i. 664 = M. Manglilla.
ovalifolia, D. Dietr. Syn. Pl. i. 612 (= *Badula ovalifolia*).—Ins. Maurit.
ovalifolia, *Miq. in Mart. Fl. Bras.* x. 313.—Bras.
ovalis, *Nadeaud, Enum. Pl. Tahiti*, 61.—Ins. Tahiti.
pachysandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 297 = Pimelandra Wallichii.

MYRSINE:—
parvifolia, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 87.—Bras.
paulensis, A. DC. l. c. 86 = lancifolia.
pellucida, *Spreng. Syst.* i. 664.—Peruv.
pellucido-punctata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 133.—Costa Rica.
penduliflora, [Moç. & Sesse, ex] A. DC. in Trans. Linn. Soc. xvii. (1834) 110.—Mexic.
pentandra, *R. Br. Prod.* 533.—Afr. austr.
philippensis, A. DC. in DC. Prod. viii. 94 = capitellata.
platystigma, *F. Muell. Fragm.* viii. 48.—Austral.
popayanensis, *H. B. & K. Nov. Gen. et Sp.* iii. 249.—N. Granat.
porosa, F. Muell. Fragm. v. 108 = crassifolia.
Porteriana, Wall. Cat. n. 6525 = capitellata.
Potama, D. Don, Prod. Fl. Nep. 146 = africana.
querimbensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 185.—Afr. trop.
racemosa, *Steud. Nom.* ed. II. ii. 176.—Am. austr.
Rapanea, Roem. & Schult. Syst. iv. 509 = floribunda.
rawacensis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 85.—Ins. Rawak.
retusa, [Soland. in] Ait. Hort. Kew. ed. I. 271 = africana.
Richardiana, Endl. in Ann. Wien. Mus. i. (1836) 171 = Urvillei.
Roraimae, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1086, nomen.—Guian.
rotundifolia, Lam. Encyc. iv. 194 = africana.
rufescens, A. DC. in Trans. Linn. Soc. xvii. (1834) 109 = laeta.
ruminata, *E. Mey. in Drège, Zwei Pfl. Docum.* 204, nomen; *et ex A. DC. in DC. Prod.* viii. 104.—Afr. austr.
salicifolia, A. DC. in Trans. Linn. Soc. xvii. (1834) 108 = laeta.
salicina, *Heward, in Hook. Lond. Journ. Bot.* i. (1842) 283, in nota.—N. Zel.
saligna, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 112.—Am. austr.
Samara, Roem. & Schult. Syst. iv. 511 = pentandra.
sandwicensis, *DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 85.—Ins. Sandvic.
scabra, Gaertn. Fruct. i. 282 = africana.
Schomburgkiana, *Miq. in Mart. Fl. Bras.* x. 315.—Guiana.
semiserrata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 293.—Reg. Himal.; Burma.
serrata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 134.—Costa Rica.
sessilis, D. Don, Prod. Fl. Nep. 146 = semiserrata.
Sieberi, D. Dietr. Syn. Pl. i. 619 = Ardisia guadaloupensis.
simensis, *Hochst. ex A. DC. in DC. Prod.* viii. 97.—Afr. trop.
spicata, *H. B. & K. Nov. Gen. et Sp.* iii. 250.—Am. austr.
subsessilis, F. Muell. Fragm. iv. 81 = crassifolia.
subspinosa, D. Don, Prod. Fl. Nep. 147 = semiserrata.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 574.—Sumatra.
taitensis, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 330.—Ins. Tahiti.
tetrandra, Willd. ex Roem. & Schult. Syst. iv. 803 = spicata.
theaefolia, Wall. Cat. n. 6391 = Ilex theaefolia.
tomentosa, *Presl, Rel. Haenk.* ii. 63.—Peruv.
Trinitatis, A. DC. in Trans. Linn. Soc. xvii. (1834) 108 = laeta.
umbellata, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 18.—Bras.
umbellulata, A. DC. in Trans. Linn. Soc. xvii. (1834) 135 = capitellata.
umbrosa, Mart. ex Miq. in Mart. Fl. Bras. x. 308 = floribunda.
undulata, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 47 = Urvillei.
undulata, Phytopol. ex Schlecht. in DC. Prod. xiv. 615 = Olearia Forsteri.
undulata, Wall. Cat. n. 2301 = Embelia Nagushia.
urceolata, *R. Br. Prod.* 534.—Austral.
Urvillei, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 105.—N. Zel.
variabilis, *R. Br. Prod.* 534.—Austral.
venosa, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 86 = leuconeura.
venosissima, *Spreng. Syst.* i. 664.—Peruv.

MYRSINE:—
venulosa, Spreng. *Syst.* i. 663 = melanophleos.
verticillata, *Presl, Rel. Haenk.* ii. 64.—Peruv.
villosissima, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 18.—Bras.
Wightiana, Wall. Cat. n. 2300 = capitellata.

MYRSIPHYLLUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 25 = **Asparagus**, Tourn. (Liliac.).
angustifolium, Willd. l. c. = A. medeoloides.
asparagoides, Willd. l. c. = A. medeoloides.
erectum, Schlecht. ex Kunth, Enum. Pl. v. 109 = A. erectus.
falciforme, Kunth, l. c. 107 = A. medeoloides.
gramineum, Kunth, l. c. 108 = A. Krausii.
Krausianum, Kunth, l. c. 107 = A. Krausii.
striatum, Schlecht. ex Kunth, l. c. 110 = A. striatus.
undulatum, Schlecht. ex Kunth, l. c. 109 = A. undulatus.

MYRSSIPHYLLA, Rafin. Sylva Tellur. 148 (1838) = **Pychotria**, Linn. (Rubiac.).
rigida, Rafin. l. c. = P. myrtiphylla.

MYRSTIPHYLLUM, P. Br. Hist. Jamaic. 152 (1756) = **Psychotria**, Linn. (Rubiac.).

MYRTELLA, F. Muell. Papuan Pl. i. 105 (1877 ?). *MYRTACEAE.*
Beccarii, *F. Muell. l. c.* 106.—N. Guin.
hirsutula, *F. Muell. l. c.*—N. Guin.

MYRTEOLA, Berg, in Linnaea, xxvii. (1854) 393 = **Myrtus**, Tourn.
Barneoudii, Berg, l. c. xxix. (1857–58) 252.—Chili.
bullata, Berg, l. c. xxvii. (1854) 394 = Myrtus bullata, *Poepp.*
humifusa, Berg, l. c. xxix. (1857–58) 252.—Chili.
microphylla, Berg, l. c. xxvii. (1854) 393 = Myrtus microphylla, *Humb. & Bonpl.*
myrsinoides, Berg, l. c. 396 = Myrtus myrsinoides.
nummularia, Berg, l. c. = Myrtus nummularia.
oxycoccoides, Berg, l. c. = Myrtus oxycoccoides.
repens, Berg, in Bot. Zeit. xvi. (1858) 250 = Myrtus repens.
vaccinioides, Berg, in Linnaea, xxvii. (1854) 395 = Myrtus vaccinioides.

MYRTHUS, Scop. Introd. 218 (1777) = **Myrtus**, Tourn.

MYRTILLOIDES, Banks & Soland. ex Hook. Ic. Pl. t. 639 (1844) = **Fagus**, Tourn. (Cupul.).
cinerascens, Banks & Soland. ex Hook. l. c. = F. Solandri.

MYRTILLUS, Gilib. Fl. Lituan. i. 4 (1781) = **Vaccinium**, Linn.
niger, Gilib. l. c. = V. Myrtillus.
sylvaticus, Drej. Fl. Excurs. Hafn. 147 = V. Myrtillus.
uliginosus, Drej. l. c. = V. uliginosum.

MYRTINIA, Nees, in DC. Prod. xi. (1847), sphalm. = **Martynia**, Linn. (Pedal.).

MYRTOBIUM, Miq. in Linnaea, xxv. (1852) 652 = **Lepidoceras**, Hook. f. (Loranth.).
microphyllum, Miq. l. c. = L. Kingii.

MYRTOPHYLLUM, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 604 = **Azara**, Ruiz & Pav. (Bixin.).
chilense, Turcz. l. c. 605 = A. macrophylla.

MYRTOPSIS, O. Hoffm. in Linnaea, xliii. (1880–82) 133. *MYRTACEAE.*
malangensis, *O. Hoffm. l. c.* 134.—Afr. trop.

MYRTUS, [Tourn.] Linn. Syst. ed. I (1735). *MYRTACEAE*, Benth. & Hook. f. i. 714.
ANAMOMIS, Griseb. Fl. Brit. W. Ind. 240 (1860).
BLEPHAROCALYX, Berg, in Linnaea, xxvii. (1854) 412.
CALOMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 76, in obs. (1849).
CLUACENA, Rafin. Sylva Tellur. 104 (1838).
EVANESCA, Rafin. l. c. 105 (1838).
HETEROMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 76, in obs. (1849).

MYRTUS:—
Leantria, Soland. ex Forst. in Comm. Gotting. ix. (1789) 45.
Luma, A. Gray, Bot. U. St. Expl. Exped. i. 535. t. 66 (1854).
Macropsidium, Blume, Mus. Bot. Lugd. Bat. i. 85 (1849).
Myrceugenia, Berg, in Linnaea, xxvii. (1854) 131.
Myrcianthes, Berg, l. c. 315 (1854).
Myrteola, Berg, l. c. 393 (1854).
Pseudocaryophyllus, Berg, l. c. 415.
Temu, Berg, in Linnaea, xxx. (1859–60) 710.
Temus, Molina, Sagg. Chil. 178 (1782).
Ugni, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 579.
acerosa, *Berg, in Linnaea*, xxix. (1857–58) 253.—Am. austr.
acetosans, Spreng. Syst. i. 488 = Eugenia acetosans.
Acka, Juss. ex DC. Prod. iii. 277 = Eugenia Acka.
acmenioides, *F. Muell. Fragm.* i. 77.—Austral.
acris, Sw. Prod. Veg. Ind. Occ. 79 = Pimenta acris.
acuminata, H. B. & K. Nov. Gen. et Sp. vi. 141 = Myrcia acuminata.
acuminata, Mill. ex Steud. Nom. ed. I. 546, nomen = communis.
acuminatissima, Blume, Bijdr. 1088 = Eugenia acuminatissima.
acuta, Mill. Gard. Dict. ed. VIII. n. 3 = communis.
acutata, *Berg, in Mart. Fl. Bras.* xiv. i. 415.—Bras.
aequalis, *Vell. Fl. Flum.* 217; v. t. 76.—Bras.
aeruginea, Forsyth, ex DC. Prod. iii. 283 = Eugenia aeruginea.
aeruginosa, *Griseb. in Goett. Abh.* xxiv. (1879) 127.—Reg. Argent.
aeruginosa, Steud. Nom. ed. II. ii. 177 = *aeruginea*.
affinis, *Sweet, Hort. Brit.* ed. I. 158, nomen.—China?
aggregata, Vell. Fl. Flum. 215; v. t. 65 (= *Phyllocalyx cerasiflorum*).—Bras.
alata, Zipp. ex Span. in Linnaea, xv. (1841) 204 = Gustavia globosa.
alaternoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 177.—N. Caled.
alba, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = Melaleuca Leucadendron.
albida, H. B. & K. Nov. Gen. et Sp. vi. 143 = Eugenia albida.
albida, Vell. Fl. Flum. 216; v. t. 72 = Eugenia discrossa.
albo-tomentosa, Mart. ex DC. Prod. iii. 254 = Myrica albotomentosa.
alpigena, Mart. l. c. 265 = Eugenia alpigena.
alpina, *Sw. Prod. Veg. Ind. Occ.* 77 = Eugenia alpina.
amazonica, Mart. ex DC. Prod. iii. 250 = Myrcia amazonica.
ambrosiaca, Mor. ex Berg, in Linnaea, xxvii. (1854) 351 = Psidiopsis Moritziana.
anceps, Spreng. Neue Entdeck. ii. 170; Syst. ii. 487 = Myrcia anceps.
androsaemoides, Linn. Sp. Pl. 472 = Eugenia androsaemoides.
androsaemoides, Poir. Encyc. iv. 409 = Myonima multiflora.
angustifolia, Linn. Mant. i. 74 = Metrosideros angustifolia.
angustifolia, Rafin. Sylva Tellur. 105 = communis.
angustifolia, *Parodi, in Anal. Soc. Cienc. Argent.* vii. (1879) 186; *Contrib.* 146.—Paraguay.
angustifolia, Spreng. Syst. ii. 479 = Eugenia angustifolia.
Arayan, *H. B. & K. Nov. Gen. et Sp.* vi. 133.—Peruv.
aromatica, Poir. in Lam. Encyc. iv. 410 = seq.?
aromatica, Salisb. Prod. 354 = Pimenta vulgaris.
Aubletii, Spreng. Syst. ii. 486 = Myrcia tomentosa.
auriculata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 98, 104 (= *Jambosa ovalifolia, auriculata*).—N. Guin.
australis, Spreng. Syst. ii. 482 = Eugenia myrtifolia, *Sims*.
axillaris, Poir. in Lam. Encyc. iv. 412 = Eugenia Poiretii.
axillaris, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia monticola.
baetica, Hoffmgg. Verz. Pfl. 82, vide *boetica*.
baladensis, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 177.—N. Caled.
balsamica, Spreng. Syst. ii. 481 = Eugenia balsamica.
banisteriaefolia, Mart. ex DC. Prod. iii. 246 = Myrcia banisteriaefolia.

MYRTUS:—
Barneoudii, *Berg, in Linnaea*, xxvii. (1854) 398.—Chili.
baruensis, Spreng. Syst. ii. 483 = Eugenia baruensis.
Becklerii, *F. Muell. Fragm.* ii. 85.—Austral.
belgica, Mill. Gard. Dict. ed. VIII. n. 2 = communis.
Berberis, Mart. ex DC. Prod. iii. 254 = Myrcia Berberis.
Berlandiereana, *Berg, in Linnaea*, xxvii. (1854) 403.—Mexic.
Berteriana, Spreng. Syst. ii. 482 = Eugenia Berteriana.
Berteroi, Phil. in Bot. Zeit. xiv. (1856) 644 = Selkirkii.
bicolor, H. B. & K. Nov. Gen. et Sp. vi. t. 540 = Eugenia discolor.
Bidevillii, *Benth. Fl. Austral.* iii. 275.—Austral.
biflora, Linn. Syst. ed. X. 1056 = Eugenia biflora.
biflora, Willd. ex Berg, in Linnaea, xxvii. (1854) 174 = Eugenia commutata.
Billardiana, H. B. & K. Nov. Gen. et Sp. vi. 139 = Myrcia Billardiana.
Blanchetiana, *Berg, in Mart. Fl. Bras.* xiv. i. 418.—Bras.
boetica, Mill. Gard. Dict. ed. VIII. n. 4 = communis.
borbonica, Spreng. Syst. ii. 481 = Eugenia coriacea.
brachyphylla, Kunze, ex Berg, in Linnaea, xxvii. (1854) 144 = Eugenia leptospermoides.
brachystemon, *DC. Prod.* iii. 240.—Ind. occ.
bracteata, Mart. ex DC. Prod. iii. 245 = Myrcia bracteata.
bracteata, Willd. Sp. Pl. ii. 969 = Eugenia bracteata.
bracteolaris, Poir. in Lam. Encyc. iv. 411 = Myrcia bracteolaris.
bracteolaris, Steud. Nom. ed. II. ii. 177 = Eugenia brasiliensis.
brasiliana, Linn. Sp. Pl. 701 = Eugenia Michelii.
bullata, *Poepp. ex Berg, in Linnaea*, xxvii. (1854) 394, in syn.—Chili.
bullata, Salisb. Prod. 354 = Calyptranthes bullata.
bullata, *Soland. ex A. Cunn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 115.—N. Zel.
buxifolia, Poir. in Lam. Encyc. iv. 413 = Eugenia coriacea.
buxifolia, Rafin. Sylva Tellur. 104 = communis.
buxifolia, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia buxifolia.
Caledoniae, Spreng. Syst. iv. Cur. Post. 193 (= *Caryophyllus ellipticus*).—N. Caled.
calophylla, *H. B. & K. Nov. Gen. et Sp.* vi. 133.—Venezuela.
camaraeana, Mart. ex DC. Prod. iii. 251 = Myrcia camaraeana.
campestris, Mart. ex DC. l. c. 274 = Eugenia campestris.
camphorata, *Baill. Hist. Pl.* vi. 340.—Chili.
cana, Mart. ex DC. Prod. iii. 264 = Eugenia cana.
Candolleana, Mart. ex DC. l. c. 281 = Eugenia Candolleana.
Candollii, *Barn. in C. Gay, Fl. Chil.* ii. 382.—Chili.
canescens, Lour. Fl. Cochinch. 311 = Rhodomyrtus tomentosa.
capensis, *Burm. f. Fl. Cap. Prod.* 14.—Afr. austr.
capensis, Harv. Gen. S. Afr. Pl. 99 = Eugenia capensis.
capparidifolia, Mart. ex DC. Prod. iii. 281 = Eugenia capparidifolia.
carnea, G. F. Mey. Prim. Fl. Esseq. 191 = Myrcia carnea.
caryophyllata, Jacq. Obs. Bot. ii. 1 = Pimenta acris.
caryophyllata, Linn. Sp. Pl. 472 = Eugenia caryophyllaea.
caryophyllata, Vell. Fl. Flum. 216; v. t. 70 = M. Pseudo-caryophyllus.
Caryophyllus, Blume, ex Steud. Nom. ed. II. ii. 177, nomen (Quid?)—Java.
Caryophyllus, Spreng. Syst. ii. 485 = Eugenia caryophyllaea.
casearioides, H. B. & K. Nov. Gen. et Sp. vi. 145 = Eugenia casearioides.
cassinoides, Salzm. ex Berg, in Mart. Fl. Bras. xiv. i. 286 = Eugenia cassinoides, *Berg*.
cassinoides, Spreng. Syst. ii. 481 = Eugenia cassinoides, *Lam*.
cassinoides, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 124 (= *Jossinia litoralis*).—N. Guinea.
caudata, *Wall. Cat.* n. 3631.—Singapur.
cauliflora, Blume, Bijdr. 1086 = Eugenia polycephala.
cauliflora, Mart. Reise Bras. i. 285 = Eugenia edulis, *Vell*.
cayennensis, Spreng. Syst. ii. 480 = Eugenia citrifolia.
cerasiformis, Blume, Bijdr. 1087 = Eugenia cerasiformis.

MYRTUS:—
cerasina, Vahl, Symb. Bot. ii. 57 = Eugenia ligustrina, *Willd*.
Cheken, Spreng. Syst. ii. 485 = Eugenia Chequen.
Chequen, Berg, in Linnaea, xxvii. (1854) 146 = praec.
chinensis, Lour. Fl. Cochinch. 313 = Symplocos sinica.
chrysocarpa, Poepp. ex Berg, in Linnaea, xxvii. (1854) 169 = Eugenia chrysocarpa.
Chrysophyllum, Spreng. Syst. ii. 482 = Eugenia ferruginea.
Chytraculia, Linn. Syst. ed. X. 1056 = Calyptranthus Chytraculia.
ciliaris, Mor. ex Berg, in Linnaea, xxvii. (1854) 246 = Eugenia ternifolia.
citrifolia, Poir. Encyc. iv. 410 (*cytrifolia*) = Pimenta officinalis.
citrifolia, Willd. ex Berg, in Linnaea, xxvii. (1854) 381 = Calycolpus calophyllus.
clinocarpa, Mart. ex DC. Prod. iii. 267 = Eugenia clinocarpa.
clusiaefolia, H. B. & K. Nov. Gen. et Sp. vi. 138 = Myrcia clusiaefolia.
coarensis, Mart. ex DC. Prod. iii. 267 = Eugenia coarensis.
coccolobaefolia, H. B. & K. Nov. Gen. et Sp. vi. 139 = Myrcia coccolobaefolia.
collina, Mart. ex DC. Prod. iii. 281 = Eugenia collina.
Comete, Spreng. Syst. ii. 488 = Myrcia Coumeta.
Commersonii, Spreng. l. c. 479 = Eugenia cotinifolia.
communis, Blanco, Fl. Filip. ed. I. 422 = Decaspermum Blancoi.
communis, *Linn. Sp. Pl.* 471.—Europ. austr.; Oriens.
complicata, H. B. & K. Nov. Gen. et Sp. vi. 140 = Myrcia complicata.
compressa, H. B. & K. l. c. 135 = Eugenia compressa.
coquimbensis, *Barn. in C. Gay, Fl. Chil.* ii. 382.—Chili.
cordata, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia cordata, *DC*.
cordiaefolia, Mart. ex DC. Prod. iii. 248 = Myrcia cordiaefolia.
coriacea, Sieber, ex DC. Prod. iii. 259 = Eugenia glomerata.
coriacea, Vahl, Symb. Bot. ii. 59 = Myrcia coriacea.
coriandriformis, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 74 = Nelitris Coriandri.
coromandelina, Koen. ex Roxb. Fl. Ind. ii. 491 = Eugenia bracteata.
corticosa, Spreng. Syst. ii. 488 = Eugenia Jambolana.
costata, Mart. ex DC. Prod. iii. 252 = Myrcia costata.
cotinifolia, Burm. ex Steud. Nom. ed. II. ii. 177, nomen (Quid?).—Am. austr.
cotinifolia, Poir. in Lam. Encyc. iv. 410 = Myrcia coriacea.
cotinifolia, Spreng. Syst. ii. 481 = Eugenia cotinifolia.
crassifolia, Mart. ex DC. Prod. iii. 266 = Eugenia crassifolia, *Miq*.
crenulata, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia crenulata, *Willd*.
Cumini, Linn. Sp. Pl. 471 = Eugenia Jambolana.
cuneata, Heyne, ex Wall. Cat. sub n. 3625 = Eugenia Jossinia.
curatellaefolia, Mart. ex DC. Prod. iii. 253 = Myrcia curatellaefolia.
cuspidata, *Berg, in Mart. Fl. Bras.* xiv. i. 415.—Bras.
cuspidata, Willd. ex Berg, in Linnaea, xxvii. (1854) 283 = Eugenia intermedia.
cuspidatifolia, Mart. ex DC. Prod. iii. 279 = Eugenia cuspidifolia.
cuspidifolia, Mart. ex DC. l. c. = Eugenia sphaerosperma?
cymiflora, F. Muell. Fragm. v. 12 = Rhodomyrtus cymiflora.
cymosa, Blume, Bijdr. 1086 = Eugenia polyantha.
cymosa, Korth. ex Miq. Fl. Ind. Bat. i. i. 445 = Eugenia pamatensis.
cymosa, Spreng. Syst. ii. 486 = Eugenia cymosa.
cytrifolia, Poir. in Lam. Encyc. iv. 410 = *citrifolia*.
daphnoides, Spreng. Syst. ii. 480 = Eugenia ruscifolia.
Darwinii, *Barn. in C. Gay, Fl. Chil.* ii. 383.—Chili.
dealbata, Mart. ex DC. Prod. iii. 254 = Myrcia dealbata.
decorticans, Mart. ex DC. Prod. iii. 253 = Myrcia decorticans.
decussata, Mart. ex DC. l. c. 235 = Psidium decussatum.
decussata, Vell. Fl. Flum. 27; v. t. 73 = Eugenia tenifolia.

MYRTUS:—
deflexa, H. B. & K. Nov. Gen. et Sp. vi. 142 = Myrcia Humboldtiana.
deflexa, Steud. Nom. ed. II. ii. 177 = Myrcia deflexa.
delicatula, Mart. ex DC. Prod. iii. 273 = Eugenia delicatula.
densa, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 69 = Calyptranthes densa.
densiflora, Blume, Bijdr. 1087 = Eugenia densiflora.
dichotoma, Poir. Encyc. Suppl. iv. 53 = Eugenia dichotoma.
dichotoma, Salisb. Prod. 354 = Calyptranthes Syzygium.
dichroa, Kunze, ex Berg, in Linnaea, xxvii. (1854) 254 = Eugenia stenophylla.
dioica, *Linn. Syst.* ed. X. 1056,—'Am.'
dioica, Sieber, ex DC. Prod. iii. 280 = Eugenia Trinitatis.
dioica, Spreng. Syst. ii. 486 = Myrcia Schaueriana.
discolor, H. B. & K. Nov. Gen. et Sp. vi. 134 = Eugenia discolor.
discolor, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 315 = Eugenia Hoffmannseggii.
disticha, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia disticha.
divaricata, Ham. Prod. Pl. Ind. Occ. 45 = Eugenia divaricata, *Berg*.
dives, Kunze, ex Berg, in Linnaea, xxvii. (1854) 147 = Eugenia Chequen.
Dombeyi, Spreng. Syst. ii. 485 = Eugenia brasiliensis.
Duarti, Mart. ex DC. Prod. iii. 247 = Myrcia exsucca.
dumetorum, Poir. Encyc. Suppl. iv. 52 = Eugenia dumetorum.
dumosa, L'Hérit. ex DC. Prod. iii. 272 = Eugenia L'Heritieriana.
dumosa, Spreng. Syst. ii. 483 = Eugenia salicifolia.
dumosa, Vahl, Symb. Bot. ii. 57 = Eugenia dumosa.
dysenterica, Mart. Reise Bras. ii. 571 = Eugenia dysenterica.
egensis, Mart. ex DC. Prod. iii. 281 = Eugenia egensis.
Ehrenbergii, *Berg, in Linnaea*, xxvii. (1854) 404.—Mexic.
elachantha, F. Muell. Fragm. iv. 56 = Decaspermum paniculatum.
elegans, *DC. Prod.* iii. 240.—Bras.
elegans, Mart. ex DC. l. c. 251 = Myrcia elegans.
elegantula, Kunze, ex Berg, in Linnaea, xxvii. (1854) 155 = Eugenia distoma.
elliptica, Spreng. Syst. ii. 483 = Eugenia cotinifolia.
elliptica, *Wright, in Sauv. Fl. Cub.* 39.—Cuba.
emarginata, H. B. & K. Nov. Gen. et Sp. vi. 142 = Eugenia emarginata, *DC.*
emarginata, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 177.—N. Caled.
emarginata, Ruiz, ex Berg, in Linnaea, xxvii. (1854) 274 = Eugenia andina.
eriopus, Mart. ex DC. Prod. iii. 255 = Myrcia eriopus.
erythrocarpa, H. B. & K. Nov. Gen. et Sp. vi. 148 = Eugenia erythrocarpa.
erythroxyloides, *H. B. & K. l. c.* 149.—Venezuel.
eugeniaeflora, Kunth, ex Berg, in Linnaea, xxvii. (1854) 139 (= *Acca domingensis*).—Ind. occ.
excelsa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 293.—Bras.
exsucca, Domb. ex DC. Prod. iii. 278 = Eugenia exsucca.
exsucca, Mart. ex DC. l. c. 247 = Myrcia exsucca.
fascicularis, *DC. l. c.* 240.—Guian.
fasciculata, *Vell. Fl. Flum.* 210; v. t. 69.—Bras.
Felisberti, Mart. ex DC. Prod. iii. 266 = Eugenia Felisberti.
fenestrata, Mart. ex DC. l. c. 251 = Myrcia fenestrata.
fernandeziana, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 316.—Ins. J. Fernand.
ferruginea, Spreng. Syst. ii. 487 = Myrcia ferruginea, *DC.*
fimbriata, H. B. & K. Nov. Gen. et Sp. vi. 137 = Eugenia fimbriata.
firma, Mart. ex DC. Prod. iii. 282 = Eugenia capparidifolia?
firma, Spreng. Syst. ii. 487 = Myrcia nobilis.
flava, Mart. ex DC. Prod. iii. 272 = Eugenia flavescens.
floribunda, Spreng. Syst. ii. 484 = Eugenia floribunda.
florida, Mart. ex DC. Prod. iii. 283 = Eugenia florida.
foetida, Spreng. Syst. ii. 481 = Eugenia foetida.
foliosa, H. B. & K. Nov. Gen. et Sp. vi. 134 = Eugenia foliosa.
fragrans, Sw. Prod. Veg. Ind. Occ. 79 = Eugenia fragrans.

MYRTUS:—
fragrantissima, *F. Muell. ex Benth. Fl. Austral.* iii. 277.—Austral.
frigida, Willd. ex Berg, in Linnaea, xxvii. (1854) 413 = salicifolia.
fruticulosa, Mart. ex DC. Prod. iii. 270 = Eugenia fruticulosa.
fruticulosa, Poepp. ex Berg, in Linnaea, xxvii. (1854) 244 = Eugenia crenulata, *Willd.*
fulva, *Spreng. Syst.* ii. 487.—Bras.
Gayana, *Berg, in Linnaea*, xxvii. (1854) 399.—Chili.
glabra, Vell. Fl. Flum. 215; v. t. 67 = Eugenia badia.
glabrata, Blume, Bijdr. 1088 = Eugenia ramosissima.
glabrata, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia glabrata.
globosa, Korth. in Nederl. Kruidk. Arch. i. (1848) 197 = Rhodamnia globosa.
glomerata, Spreng. Syst. ii. 483 = Eugenia glomerata.
Goetheana, Mart. ex DC. Prod. iii. 240 = Calycolpus Goetheanus.
gonoclada, *F. Muell. ex Benth. Fl. Austral.* iii. 275.—Austral.
grammica, *Spreng. Syst.* ii. 480.—Bras.
grandifolia, *Berg, in Mart. Fl. Bras.* xiv. i. 419.—Bras.
Greggii, Nees & Mart. in Nov. Act. Nat. Cur. xii. i. (1824) 51 = Myrcia Maximiliana.
Gregii, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia Greggii.
Grumixama, Vell. Fl. Flum. 216; v. t. 69 = Eugenia brasiliensis.
Guayabillo, *Wright, in Sauv. Fl. Cub.* 39.—Cuba.
guayaquilensis, H. B. & K. Nov. Gen. et Sp. vi. 147 = Eugenia guayaquilensis.
Gudilla, Colla, in Mem. Acc. Torin. xxxvii. (1834) 66 = Eugenia stenophylla.
guianensis, Ham. Prod. Pl. Ind. Occ. 45; Spreng. Syst. ii. 482 = Myrcia guayanensis.
Hayneana, Mart. ex DC. Prod. iii. 246 = Myrcia Hayneana.
Heynei, Spreng. Syst. ii. 482 = Eugenia bracteata.
Hilariana, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 275 = Eugenia Hilariana.
Hillii, *Benth. Fl. Austral.* iii. 275.—Austral.
hirta, Willd. ex Berg, in Mart. Fl. Bras. xiv. i. 156 = Myrcia lanceolata.
horizontalis, Vent. Jard. Malm. 60 = Eugenia disticha.
hortensis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 20 = communis?
hypericifolia, Blume, Bijdr. 1082 (= *Jambosa hypericifolia*).—Java.
hypericifolia, Salisb. Prod. 354 (= *Syzygium hypericifolium*).—Ins. Borbon.
idaea, Willd. ex Berg, in Linnaea, xxvii. (1854) 395 = vaccinioides.
incana, *Berg, in Mart. Fl. Bras.* xiv. i. 416.—Bras.
indica, Walp. Rep. ii. 172 = Eugenia Jossinia.
inocarpus, Mart. ex DC. Prod. iii. 264 = Eugenia inocarpa.
inundata, Mart. ex DC. l. c. 280 = Eugenia inundata.
italica, Mill. Gard. Dict. ed. VIII. n. 5 = communis.
Jaboticaba, Vell. Fl. Flum. 214; v. t. 62 = Myrcia Jaboticaba.
Jacquiniana, *Berg, in Linnaea*, xxvii. (1854) 406.—Hab.?
Jambos, H. B. & K. Nov. Gen. et Sp. vi. 144 = Eugenia Jambos.
javanica, Blume, Bijdr. 1084 = Eugenia aquea.
javanica, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 96 = Eugenia javanica.
javanica, Spreng. Syst. ii. 484 = Eugenia bifaria.
Kalah, Miq. Anal. Bot. Ind. i. 23 = Eugenia kalahiensis.
Kochiana, Mart. ex DC. Prod. iii. 268 = Eugenia Kochiana.
Krausei, *Phil. in Linnaea*, xxviii. (1856) 638.—Chili.
Kuma, *Barn. in C. Gay, Fl. Chil.* ii. 384.—Chili.
Kunthiana, Mart. ex DC. Prod. iii. 265 = Eugenia Kunthiana.
laevigata, Mart. ex DC. l. c. 284 = Eugenia laevigata.
laevis, *Thunb. Fl. Jap.* 198.—Japon.
Lambertiana, Mart. ex DC. Prod. iii. 271 = Eugenia bahiensis.
lancea, Spreng. Syst. ii. 482 = Eugenia lancea.
lanceolata, Juss. ex Jaume St. Hil. in Duham. Arb. ed. Nov. i. 208 = Eugenia Dombeyana.
lanceolata, Rafin. Sylva Tellur. 105 = communis.
lasiantha, Mart. ex DC. Prod. iii. 254 = Myrcia lasiantha.

MYRTUS:—
lasioclada, *F. Muell. Fragm.* ix. 148.—Austral.
lasiopus, Steud. Nom. ed. II. ii. 178 = Myrcia lasiopus.
lateriflora, Mart. ex DC. Prod. iii. 273 = Eugenia dicrossa.
latifolia, Forsyth, ex DC. l. c. 245 = Myrcia crassinervia.
latifolia, Heyne, ex Roth, Nov. Pl. Sp. 232 = Eugenia bracteata.
latifolia, Rafin. Sylva Tellur. 104 = communis.
latifolia, Spreng. Syst. ii. 481 = Eugenia latifolia.
lauriflora, Mart. ex DC. Prod. iii. 252 = Myrcia lauriflora.
laurifolia, Hort. Par. ex DC. l. c. 243 = Myrcia divaricata.
laurina, Retz. Obs. iv. 26 = Symplocos spicata.
laxa, Mart. ex DC. Prod. iii. 263 = Eugenia laxa.
leriocarpa, *Wright, in Sauv. Fl. Cub.* 38.—Cuba.
Leucadendron, Linn. Syst. ed. X. 1056 = Melaleuca Leucadendron.
leucomyrtillus, *Griseb. in Goett. Abh.* vi. (1854) 120.—Chili.
leucomyrtus, *Griseb. ex Phil. in Linnaea*, xxviii. (1856) 637 (an praec.?).—Chili.
leucophlaea, Mart. ex DC. Prod. iii. 247 = Myrcia leucophlaea.
leucoxyla, *Ortega, Hort. Matr. Dec.* 129.—N. Granat.
leucoxylon, Korth. ex Miq. Anal. Bot. Ind. i. 26 = Eugenia leucoxylon.
ligustrina, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia ligustrina, *Willd.*
limbata, H. B. & K. Nov. Gen. et Sp. vi. 136 = Eugenia limbata.
limbosa, Ruiz, ex Berg, in Linnaea, xxvii. (1854) 295 = Eugenia limbosa.
Lindleyana, H. B. & K. Nov. Gen. et Sp. vi. 138. t. 543 = Eugenia Lindleyana.
Lindleyana, Mart. ex DC. Prod. iii. 269 = Eugenia Martiusiana.
linearis, *J. F. Gmel. Syst.* 793 (Quid?).—Am. trop.
lineata, Blume, Bijdr. 1087 = Eugenia lineata, *Duthie.*
lineata, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia lineata, *DC.*
linifolia, Spreng. Syst. ii. 479 = Eugenia Sprengelii.
Linkiana, Mart. ex DC. Prod. iii. 248 = Myrcia Linkiana.
littoralis, Mart. ex DC. l. c. 249 = Myrcia littoralis.
littoralis, Roxb. ex Wight & Arn. Prod. 331 = Eugenia bracteata.
littoralis, Salisb. Prod. 353 = communis.
littoralis, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 124 = Jossinia litoralis.
longifolia, H. B. & K. Nov. Gen. et Sp. vi. 258 = Myrcia longifolia.
longifolia, Mart. ex DC. Prod. iii. 269 = Eugenia longifolia.
longifolia, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 75 = Decaspermum paniculatum.
Loureiri, Spreng. Syst. ii. 488 = Eugenia nervosa.
lucida, *Linn. Syst.* ed. X. 1056.—Guian.
lucida, Steud. Nom. ed. II. ii. 178 = Eugenia cotinifolia.
lugens, Poepp. ex Berg, in Linnaea, xxvii. (1854) 300 = Eugenia lugens.
Luma, *Barn. in C. Gay, Fl. Chil.* ii. 384.—Chili.
Luma, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 317 = fernandeziana.
Luma, Molina, Sagg. Chile, 173 = Eugenia apiculata.
Luma, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 333 = Eugenia Chequen.
Luma, Spreng. ex Berg, in Linnaea, xxvii. (1854) 262 = Eugenia Cruckshanksii.
lurida, *Spreng. Syst.* ii. 480.—Reg. Argent.
lusitanica, Willd. ex Steud. Nom. ed. I. 546 = communis.
macrochlamys, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 419.—Bras.
macrophylla, Ehrh. Beitr. vii. 50, nomen. = communis?
macrophylla, Spreng. Syst. ii. 483 = Eugenia malaccensis.
magnoliaefolia, Blume, Bijdr. 1088 (= *Syzygium magnoliaefolium*).—Java.
Makal, J. F. Gmel. Syst. 792 (= *Syzygium Makul*).—Zeylan.
malaccensis, Spreng. Syst. ii. 484 = Eugenia malaccensis.
malpighioides, H. B. & K. Nov. Gen. et Sp. vi. 146 = Eugenia malpighioides.
maragnana, Mart. ex DC. Prod. iii. 249 = Myrcia maragnana.

MYRTUS:—
marginata, Spreng. Syst. ii. 438 = Eugenia marginata.
maritima, H. B. & K. Nov. Gen. et Sp. vi. 146 = Eugenia acapulcensis.
maritima, Mart. ex DC. Prod. iii. 271 = Eugenia maritima, *DC.*
maxima, *Molina, Sagg. Chile*, 173.—Chili.
media, Hoffmgg. Verz. Pfl. 82 = communis.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 193.—Bras.
melastomaefolia, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 102 (= *Fambosa melastomaefolia.*).—Malaya.
melastomoides, F. Muell. Fragm. i. 76 = Rhodamnia trinervia.
Meli, *Phil. in Linnaea*, xxviii. (1856) 638.—Chili.
mespiloides, Spreng. Syst. ii. 481 = Eugenia cotinifolia.
micrantha, H. B. & K. Nov. Gen. et Sp. vi. 144 = Eugenia micrantha, *Bertol.*
micrantha, Nees & Mart. in Nov. Act. Nat. Cur. xii. i. (1824) 51 = Eugenia Maximiliana, *DC.*
microphylla, Ehrh. Beitr. vii. 50, nomen = communis?
microphylla, Humb. & Bonpl. Pl. Aequin. i. 19. t. 4.—Peruv.
Mikaniana, Mart. ex DC. Prod. iii. 283 = Eugenia Mikaniana.
Mini, Spreng. Syst. ii. 486 = Myrcia Mini.
minima, Mill. Gard. Dict. ed. VIII. n. 7 = communis.
modesta, Mart. ex DC. Prod. iii. 279 = Eugenia modesta.
Molinae, *Barn. in C. Gay, Fl. Chil.* ii. 381.—Chili.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 141 = Myrcia mollis.
montana, *Benth. Pl. Hartw.* 61.—Mexic.
monticola, Sw. Prod. Veg. Ind. Occ. 78 = Eugenia monticola.
monticola, Vahl, ex DC. Prod. iii. 280 = Eugenia ludibunda.
mucronata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 294.—Bras.
mucronata, Pers. Syn. ii. 30 = communis.
Mulleri, Korth. in Nederl. Kruidk. Arch. i. (1848) 197 = Rhodamnia trinervia.
multiflora, *Juss. ex Jaume St. Hil. in Duham. Arb.* ed. Nov. i. 208.—Chili.
multiflora, Spreng. Syst. ii. 485 = Myrcia multiflora.
muricata, Mart. ex DC. Prod. iii. 283 = Eugenia muricata.
myricoides, *H. B. & K. Nov. Gen. et Sp.* vi. 131. t. 539.—Peruv.
myrobalana, Mart. ex DC. Prod. iii. 278 = Eugenia myrobalana.
Myrsinites, Mart. ex DC. l. c. 236 = Psidium Myrsinites.
myrsinoides, *H. B. & K. Nov. Gen. et Sp.* vi. 132.—Peruv.
myrtillifolia, Mart. ex DC. Prod. iii. 250 = Myrcia myrtillifolia.
Neesiana, Mart. ex DC. l. c. 249 = Myrcia Neesiana.
nemoralis, Mart. ex DC. l. c. 267 = Eugenia nemoralis.
nigra, Mart. ex DC. l. c. 268 = Eugenia nigra.
nigrescens, Mart. ex DC. l. c. 246 = Myrcia nigrescens.
nitens, Poir. Encyc. Suppl. iv. 51 = Eugenia nitens.
nitida, Vell. Fl. Flum. 216; v. t. 68 = Eugenia Candolleana.
nivea, *Berg, in Mart. Fl. Bras.* xiv. i. 414.—Reg. Argent.
nummularia, Barn. in C. Gay, Fl. Chil. ii. 379 = Barneoudii.
nummularia, *Poir. in Lam. Encyc.* iv. 407.—Reg. Magell.
obcordata, *Hook. f. Fl. N. Zeal.* i. 71.—N. Zel.
oblongata, Mart. ex DC. Prod. iii. 251 = Myrcia oblongata.
obovata, Korth. ex Miq. Fl. Ind. Bat. i. i. 453 = Eugenia Muelleri.
obovata, Spreng. Syst. ii. 486 = Eugenia obovata, *Poir.*
obscura, Lindl. Bot. Reg. t. 1044 = Eugenia obscura.
obtusa, Juss. ex DC. Prod. iii. 266 = Eugenia obtusa.
obtusissima, Blume, Bijdr. 1086 = Eugenia javanica.
Oerstedeana, *Berg, in Linnaea*, xxvii. (1851) 405.—Ins. S. Crucis.
Oerstedi, *Hemsl. Biol. Centr. Am. Bot.* i. 407.—Costa Rica.
oleaefolia, H. B. & K. Nov. Gen. et Sp. vi. 147 = Eugenia oleaefolia.
Oleaster, Mart. ex DC. Prod. iii. 282 = Eugenia Pseudocaryophyllus.

MYRTUS:—
oligoneura, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 354 = Memecylon oligoneurum.
orbiculata, Spreng. Syst. ii. 480 = Eugenia cotinifolia.
orinocensis, Bonpl. ex Berg, in Linnaea, xxvii. (1854) 110 = Myrcia polyantha.
ovalifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 381.—Bras.
ovalis, *Berg, in Mart. Fl. Bras.* xiv. i. 417.—Uruguay.
ovalis, Fl. Per. ex Berg, in Linnaea, xxvii. (1854) 156 = Eugenia ovalis.
ovalis, *Spreng. Syst.* ii. 479.—Bras.
owariensis, Steud. Nom. ed. II. ii. 178 = Eugenia owariensis.
oxycoccoides, *Benth. Pl. Hartw.* 174.—N. Granat.
pallens, Vahl, Symb. Bot. ii. 57 = Eugenia pallens.
palustris, Mart. ex DC. Prod. iii. 246 = Myrcia palustris.
paniculata, J. F. Gmel. Syst. 792 = Eugenia paniculata.
parviflora, Spreng. Syst. ii. 486 = Eugenia parviflora.
parvifolia, Juss. ex DC. Prod. iii. 266 = Eugenia parvifolia.
Patrisii, Spreng. Syst. iii. 480 = Eugenia Patrisii.
patula, Mart. ex DC. Prod. iii. 284 = Eugenia patula.
pauciflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 296.—Reg. Argent.
Pavonii, *Poepp. ex Berg, in Linnaea*, xxvii. (1854) 38.—Peruv.; Chili.
pedunculata, *Hook. f. in Hook. Ic. Pl.* t. 629.—N. Zel.
pendula, Blume, Bijdr. 1085 = Eugenia pendula.
phillyraeoides, *Willd. ex Berg, in Linnaea*, xxvii. (1854) 326 (= *Myrciaria phillyraeoides*).—Am. austr.
phlebophylla, Kunze, ex Berg, l. c. 155 = Eugenia distoma.
phylicoides, *Benth. Pl. Hartw.* 131.—Peruv.
pia, Mart. ex DC. Prod. iii. 263 = Eugenia pia.
pilosa, Mart. ex DC. l. c. 253 = Myrcia pilosa.
Pimenta, Linn. Sp. Pl. 472 = Pimenta vulgaris.
pimentoides, *Lindl. ex Steud. Nom.* ed. II. ii. 178, nomen.—Ind. occ.
pistaciaefolia, Mart. ex DC. Prod. iii. 270 = Eugenia pistaciaefolia.
Pohliana, Mart. ex DC. l. c. 264 = Eugenia Pohliana.
Poiretii, Spreng. Syst. ii. 483 = Eugenia Poiretii.
polyantha, H. B. & K. Nov. Gen. et Sp. vi. 140 = Myrcia Humboldtiana.
polyantha, Mart. ex DC. Prod. iii. 252 = Myrcia polyantha.
procera, Sw. Prod. Veg. Ind. Occ. 77 = Eugenia procera.
prunifolia, Mart. ex DC. Prod. iii. 253 = Myrcia prunifolia.
Pseudocaryophyllus, Gomez, in Mem. Ac. Lisboa, iii. (1812) Mem. Corresp. 92 = Eugenia Pseudocaryophyllus.
Pseudomini, Mart. ex Steud. Nom. ed. II. ii. 178 = Myrcia Pseudomini.
Pseudopsidium, Spreng. Syst. ii. 480 = Eugenia Pseudopsidium.
psidioides, Desv. in Ham. Prod. Pl. Ind. Occ. 44 = Eugenia Desvauxiana.
pubescens, *Berg, in Mart. Fl. Bras.* xiv. i. 415.—Reg. Argent.
pubescens, H. B. & K. Nov. Gen. et Sp. vi. 143 = Eugenia pubescens.
pulchella, *Regel, Ind. Sem. Hort. Petrop.* (1857) 56.—Hab.?
punctata, Mart. ex DC. Prod. iii. 246 = Myrcia myoporina.
punctata, Spreng. Syst. ii. 482 = Myrcia punctata.
punicaefolia, H. B. & K. Nov. Gen. et Sp. vi. 149 = Eugenia punicaefolia.
pyriformis, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 216; *Contrib.* 151.—Paraguay.
quadrangularis, Buch.-Ham. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 486 = Eugenia grata.
quadriflora, Vell. Fl. Flum. 217; v. t. 77 = Eugenia lanceolata.
quadripartita, Van Royen, ex Blume, Mus. Bot. Lugd. Bat. i. 87 = Eugenia bracteata.
quadrisperma, Vell. Fl. Flum. 215; v. t. 63 = Eugenia Mikaniana.
racemosa, Mart. ex DC. Prod. iii. 281 = Eugenia racemosa.
racemosa, Vell. Fl. Flum. 213; v. t. 60 (= *Eugeniopsis cannaefolia*).—Bras.
racemulosa, *Benth. Fl. Austral.* iii. 276.—Austral.
racemulosa, Mart. ex DC. Prod. iii. 254 = Myrcia racemulosa.

MYRTUS:—
Ralphii, *Hook. f. Fl. N. Zel.* ii. 329.—N. Zel.
ramiflora, Benth. ex Berg, in Linnaea, xxvii. (1854) 335 = Eugenia coffeaefolia.
ramiflora, *Vahl, in West, Bidr. Ste.-Croix*, 290.—Ins. S. Crucis.
Raran, Colla, in Mem. Acc. Torin. xxxvii. (1834) 66 = Eugenia Raran.
Reinwardtiana, Blume, Bijdr. 1082 = Eugenia Reinwardtiana.
Reloncavi, Barn. ex Phil. f. Cat. Pl. Vasc. Chil. 79 = M. Luma, *Barn.*
repens, *Phil. in Bot. Zeit.* xv. (1857) 400.—Chili.
resupinata, Vell. Fl. Flum. 214; v. t. 61 = Marlierea resupinata.
reticulata, *Kunze, ex C. Gay, Fl. Chil.* ii. 380.—Chili.
reticulata, *Poepp. ex Berg, in Linnaea*, xxvii. (1854) 132, in syn.—Chili.
revoluta, *Schau. in Linnaea*, xxi. (1848) 272.—Guian.
rhopaloides, H. B. & K. Nov. Gen. et Sp. vi. 137 = Eugenia rhopaloides.
rhytisperma, *F. Muell. Fragm.* i. 77.—Austral.
rigida, *Berg, in Mart. Fl. Bras.* xiv. i. 417.—Bras.
rigida, J. F. Gmel. Syst. 792 = Calyptranthes rigida.
rigida, Mart. ex DC. Prod. iii. 273 = Eugenia rigida.
rigida, Sw. ex DC. l. c. 265 = Eugenia alpina.
riparia, Mart. ex DC. l. c. 283 = Eugenia riparia.
romana, Hoffmgg. Verz. Pfl. 82 = communis.
rosmarinifolia, Pers. Syn. ii. 30 = Eugenia rosmarinifolia.
rosmarinifolia, Spreng. Syst. ii. 479 = Eugenia angustifolia.
rostrata, Mart. ex DC. Prod. iii. 255 = Myrcia rostrata.
rufa, Colla, in Mem. Acc. Torin. xxxvii. (1834) 66 = Eugenia ferruginea.
rufa, Vell. Fl. Flum. 217: v. t. 75 = Marlierea rufa.
rufescens, Spreng. Syst. ii. 487 = Eugenia rufescens.
rufidula, Bonpl. ex Berg, in Linnaea, xxvii. (1854) 123 = Myrcia mollis.
rufipes, Mart. ex DC. Prod. iii. 247 = Myrcia rufipes.
rufo-punctata, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 176.—N. Caled.
Ruizii, Poepp. ex Berg, in Linnaea, xxvii. (1854) 99 = Myrcia chilensis.
ruscifolia, Rottl. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 486 = Eugenia linearis, *Duthie.*
ruscifolia, Wight, ex Wall. Cat. n. 3623 = Eugenia uniflora, *Linn.*
ruscifolia, Willd. Sp. Pl. ii. 970 = Eugenia bracteata.
Sagraea, *Berg, in Linnaea*, xxx. (1859–60) 710.—Cuba.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 136. t. 541.—N. Granat.
salicifolia, Mart. ex DC. Prod. iii. 246 = Myrcia salicifolia.
salicifolia, Willd. ex Berg, in Linnaea, xxvii. (1854) 374 = Psidium crenatum.
saligna, Burm. f. Fl. Ind. 116 = Melaleuca Leucadendron.
salutaris, *H. B. & K. Nov. Gen. et Sp.* vi. 132.—Venezuela.
samarangensis, Blume, Bijdr. 1084 = Eugenia javanica.
sancta, Mart. ex DC. Prod. iii. 267 = Eugenia sancta.
saxicola, Poepp. ex Berg, in Linnaea, xxvii. (1854) 92 = Myrcia saxicola.
scabra, *Arruda, in Koster, Trav. Braz.* 491.—Bras.
Schomburgkii, *Klotzsch, in Schomb. Faun. et Fl. Brit. Gui.* 1196, nomen.—Guian.
Schrankiana, Mart. ex DC. Prod. iii. 247 = Myrcia Schrankiana.
scutulifera, Mart. ex DC. l. c. 254 = Myrcia scutulifera.
Selkirkii, *Hemsl. Bot. Voy. Challeng.* i. III. 36.—Ins. J. Fernand.
Selloi, *Spreng. Syst.* ii. 482.—Bras.
Sellowiana, *Berg, in Mart. Fl. Bras.* xiv. i. 413.—Bras.
Sellowiana, Mart. ex DC. Prod. iii. 264 = Eugenia Sellowiana.
sepiaria, Mart. ex DC. l. c. 249 = Myrcia sepiaria.
sericea, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 295.—Bras.
seringapatamensis, Herb. Madr. ex Wall. Cat. n. 3623 = Eugenia uniflora, *Linn.*
serrata, Koen. ex Steud. Nom. ed. I. 547 = Symplocos spicata.

MYRTUS:—
serratifolia, *Griseb. in Goett. Abh.* xix. (1874) 139.—Reg. Argent.
sessiliflora, Spreng. Syst. ii. 479 = Eugenia sessiliflora.
sessilifolia, Mart. ex DC. Prod. iii. 263 = Eugenia sessilifolia.
sessilis, Korth. ex Miq. Anal. Bot. Ind. i. 25 = Eugenia Korthalsiana.
Shepherdi, *F. Muell. Fragm.* ix. 148.—Austral.
silvestris, Vell. Fl. Flum. 215; v. t. 64 = Myrcia anceps.
sinemariensis, Spreng. Syst. ii. 483 = Eugenia sinemariensis.
smilacifolia, Wall. Cat. n. 3629 = Rhodamnia trinervia.
Smithii, Spreng. Syst. ii. 487 = Eugenia Smithii.
Sonneratii, Spreng. l. c. 485 = Eugenia Wightiana.
sparsiflora, Mart. ex DC. Prod. iii. 263 = Eugenia sparsiflora.
sparsifolia, *Berg, in Linnaea*, xxvii. (1854) 402.—Ins. S. Crucis.
spectabilis, Blume, Bijdr. 1083 (Rhodomyrti sp.).—Malaya.
spectabilis, Mart. ex DC. Prod. iii. 248 = Myrcia spectabilis.
spicata, Buch.-Ham. ex Wall. Cat. n. 3621 = Eugenia bracteata.
spiciflora, Spreng. Syst. iv. Cur. Post. 193 = Eugenia Sprengelii.
splendens, Sw. Prod. Veg. Ind. Occ. 79 = Myrcia splendens.
sponsalis, Hort. ex Steud. Nom. ed. II. ii. 179 = communis.
stictopetala, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 293 = Eugenia stictopetala.
stigmatosa, Mart. ex DC. Prod. iii. 258 = Eugenia stigmatosa.
stipularis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 316 = Tepualia stipularis.
stipularis, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 88 = Eugenia stipularis.
Stoupii, Spreng. Syst. ii. 484 = Myrcia bracteolaris.
suaveolens, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 220; *Contrib.* 155.—Paraguay.
subalpestris, Mart. ex DC. Prod. iii. 250 = Myrcia subalpestris.
subamplexicaulis, Mart. ex DC. l. c. 277 = Eugenia subamplexicaulis.
subrubens, Blanco, Fl. Filip. ed. II. 294 = Eugenia cymosa.
suffruticosa, *Berg, in Mart. Fl. Bras.* xiv. i. 418.—Uruguay.
sylvatica, G. F. W. Mey. Prim. Fl. Esseq. 191 = Myrcia sylvatica.
sylvestris, Auct. = *silvestris*.
Syzygium, Linn. Amoen. Acad. v. 398 = *M. Zuzygium*.
Tabasco, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 559 = Pimenta officinalis.
tarentina, Mill. Gard. Dict. ed. VIII. n. 6; Bertol. Fl. Ital. v. 120 = communis.
tenella, Mart. ex DC. Prod. iii. 272 = Eugenia tenella.
tenuifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 280.—Austral.
thelocarpa, Mart. ex Berg, in Mart. Fl. Bras. xiv. i. 138 = Myrcia torta.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 204 = Eugenia javanica.
tinifolia, Spreng. Syst. ii. 481 = Eugenia cotinifolia.
tomentosa, [Soland. ex] Ait. Hort. Kew. ed. I. ii. 159 = Rhodomyrtus tomentosa.
torta. Mart. ex DC. Prod. iii. 250, 252 = Myrcia pallens, torta.
Tozerii, F. Muell. Fragm. ii. 86. t. 13 = Rhodomyrtus psidioides.
triflora, Herb. Madr. ex Wall. Cat. n. 3623 = Eugenia uniflora, *Linn.*
triflora, Jacq. Enum. Pl. Carib. 23 = Eugenia triflora.
triflora, *Spreng. Syst.* ii. 482.—Reg. Argent.
trinervia, Lour. Fl. Cochinch. 312 = Eugenia dumetorum.
trinervia, Sm. in Trans. Linn. Soc. iii. (1797) 280 = Rhodamnia trinervia.
trineura, F. Muell. Fragm. iv. 117 = Rhodomyrtus trineura.
tripinnata, Blanco, Fl. Filip. ed. I. 421 = Eugenia cymosa.

MYRTUS:—
trunciflora, Schlecht. & Cham. in Linnaea, v. (1830) 561 = Eugenia trunciflora.
tuberculata, H. B. & K. Nov. Gen. et Sp. vi. 148 = Eugenia tuberculata.
Ugni, Bert. in Ann. Sc. Nat. Sér. I. xxi. (1830) 347 = Selkirkii.
Ugni, *Molina, Sagg. Chile*, 161.—Chili.
uliginosa, *Miq. in Linnaea*, xxv. (1852) 652.—Chili.
umbellata, *Desv. in Ham. Prod. Pl. Fl. Ind. Occ.* 45.—Guian.
umbellata, Mart. ex DC. Prod. iii. 269 = Eugenia micropora.
umbellulifera, H. B. & K. Nov. Gen. et Sp. vi. 135 = Myrcia umbellulifera.
umbilicata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 296.—Bras.
umbraticola, *H. B. & K. Nov. Gen. et Sp.* vii. 258.—Venezuela.
undulata, Spreng. Syst. ii. 482 = Eugenia undulata.
Ungui, Steud. Nom. ed. II. ii. 179, sphalm. = E. Ugni.
vaccinioides, *H. B. & K. Nov. Gen. et Sp.* vi. 130.—Peruv.
vaccinioides, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 176.—N. Caled.
Vahlii, Spreng. Syst. ii. 486 = Eugenia Vahlii.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 688.—Chili.
variabilis, Mart. ex DC. Prod. iii. 254 = Myrcia variabilis.
variegata, Blume, Bijdr. 1082 = Eugenia minima.
venosa, Spreng. Syst. ii. 485 = Eugenia venosa.
venulosa, Mart. ex DC. Prod. iii. 250 = Myrcia venulosa.
verrucosa, *Berg, in Linnaea*, xxvii. (1854) 405.—Hab.?
verrucosula, Mart. ex DC. Prod. iii. 285 = Eugenia verrucosula.
verticillata, Salzm. ex Berg, in Linnaea, xxvii. (1854) 331 (= *Myrciaria splendens*).—Bras.
verticillata, Vell. Fl. Flum. 215; v. t. 66 = Eugenia Riedeliana.
vestita, Mart. ex DC. Prod. iii. 248 = Myrcia vestita.
vestita, *Spreng. Syst. iv. Cur. Post.* 193.—Bras.
vetula, Mart. ex DC. Prod. iii. 271 = Eugenia vetula.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 177.—N. Caled.
villosa, Spreng. Syst. ii. 487 = Monimia rotundifolia.
violacea, Spreng. l. c. 488 = Eugenia violacea.
virgata, Mor. ex Berg, in Linnaea, xxvii. (1854) 391 = myricoides.
virgultosa, Mart. ex DC. Prod. iii. 250 = Myrcia ramulosa.
virgultosa, Sw. Prod. Veg. Ind. Occ. 79 = Eugenia virgultosa.
viridis, Vell. Fl. Flum. 217; v. t. 74 = Eugenia viridis.
Vulcani, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 197 (Decaspermi sp. ?).—Sumatra.
Willdenowii, Spreng. Syst. ii. 480 = Eugenia portoricensis, uniflora, *Linn.*
xalapensis, H. B. & K. Nov. Gen. et Sp. vi. 145 = Eugenia xalapensis.
xylapioides, H. B. & K. l. c. vii. 257. t. 657 = Myrcia xylopioides.
Zeyheri, Harv. Gen. S. Afr. Pl. 99 = Eugenia Zeyheri.
zeylanica, Linn. Sp. Pl. 472 = Eugenia zeylanica.
Zuzygium, Linn. Syst. ed. X. 1056 = Calyptranthes Suzygium.

MYSCOLUS, Cass. in Bull. Soc. Philom. (1818) 33 = **Scolymus**, Tourn. (Compos.).
grandiflorus, D. Dietr. Syn. Pl. iv. 1333 = S. grandiflorus.
hispanicus, D. Dietr. l. c. = S. hispanicus.
megacephalus, Cass. in Dict. Sc. Nat. xxxiv. 84 = S. grandiflorus.
microcephalus, Cass. l. c. 85 = S. hispanicus.

MYSICARPUS, Webb, in Hook. Niger Fl. 122 (1849) lapsu = **Alysicarpus**, Neck. (Legumin.).

MYSOTIS, Hill, Veg. Syst. vii. 55 (1764), sphalm. = **Myosotis**, Rupp. (Borag.).

MYSTACIDIUM, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 205. *ORCHIDEAE*, Benth. & Hook. f. iii. 584.
AËRANTHUS, Reichb. f. in Walp. Ann. vi. 899 (1861).
GUSSONEA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. 67 (1828).
GUSSONIA, Spreng. Gen. ii. 664 (1831).
MICROCOELIA, Lindl. Gen. et Sp. Orch. 60 (1830).
caulescens, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 488.—Madag.
filicorne, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. austr.
gracile, *Harv. Thes. Cap.* ii. 48.—Afr. austr.
graminifolium, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 490.—Madag.
inapertum, *Ridl. l. c.* 489.—Madag.
ochraceum, *Ridl. l. c.* 488.—Madag.
pusillum, *Harv. Thes. Cap.* ii. 47.—Afr. austr.
tenellum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 489.—Madag.
zeylanicum, *Trimen, Cat. Pl. Ceyl.* 90.—Zeylan.

MYSTACINUS, Rafin. Sylva Tellur. 30 (1838) = **Helinus**, E. Mey. (1840) (Rhamn.).
cirrhiferus, Rafin. l. c. = H. Mystacinus.

MYSTROPETALON, Harv. Gen. S. Afr. Pl. 418 (1838); et in Ann. Nat. Hist. Ser. I. ii. (1839) 385. t. 19, 20. *BALANOPHOREAE*, Benth. & Hook. f. iii. 234.
BLEPHAROCHLAMYS, Presl, Epim. Bot. 245 (1850).
Polemanni, *Harv. Gen. S. Afr. Pl.* 418.—Afr. austr.
Thomii, *Harv. l. c.* 419.—Afr. austr.

MYSTROXYLON, Eckl. & Zeyh. Enum. 125 (1835) = **Elaeodendron**, Jacq. (Tiliac.).
athranthum, Eckl. & Zeyh. l. c. 126 = E. athranthum.
Burkeanum, Sond. in Harv. & Sond. Fl. Cap. i. 470.—Afr. austr.
confertiflorum, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 106 = E. athranthum.
eucleaeforme, Eckl. & Zeyh. Enum. 125.—Afr. austr.
filiforme, Eckl. & Zeyh. l. c.—Afr. austr.
Kuba, Eckl. & Zeyh. l. c. 127.—Afr. austr.
macrocarpum, Sond. in Harv. & Sond. Fl. Cap. i. 471.—Afr. austr.
oligocarpum, Eckl. & Zeyh. Enum. 125.—Afr. austr.
pubescens, Eckl. & Zeyh. l. c. 126.—Afr. austr.
reticulatum, D. Dietr. Syn. Pl. i. 817.—Afr. austr.
sessiliflorum, Eckl. & Zeyh. Enum. 126.—Afr. austr.
sphaerophyllum, Eckl. & Zeyh. l. c. = E. sphaerophyllum.
spilocarpum, Eckl. & Zeyh. l. c.—Afr. austr.

MYSTYLLUS, Presl, ex Ann. Sc. Nat. Sér. II. i. (1834) 365 = MISTYLLUS, Presl = **Trifolium**, Tourn. (Legumin.).

MYTILICOCCUS, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 173 = **Lunasia**, Blanco (Rutac.).
costulatus, Miq. Fl. Ind. Bat. i. II. 388 = L. amara.
grandifolius, Miq. l. c. = L. amara.
quercifolius, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 173 = L. amara.

MYXOPYRUM, Blume, Bijdr. 683 (1825). *OLEACEAE*, Benth. & Hook. f. ii. 680.
CHRONDROSPERMUM, Wall. Cat. n. 2837 (1832).
coriaceum, *Blume, Mus. Bot. Lugd. Bat.* i. 320.—Borneo.
nervosum, *Blume, Bijdr.* 683.—Malaya.
smilacifolium, *Blume, Mus. Bot. Lugd. Bat.* i. 320.—Ind. or.; Malaya.

MYXOSPERMUM, M. Roem. Syn. Hesper. 40 (1846) = **Glycosmis**, Correa (Rutac.).
chylocarpum, M. Roem. l. c. = G. pentaphylla.

MYZODENDRON, [Banks] Soland. ex Forst. in Comm. Gotting ix. (1789) 45 (Myzodendrum). *SANTALACEAE*, Benth. & Hook. f. iii. 229.
angulatum, *Phil. in Anal. Univ. Chil.* (1865) II. 315.—Chili.
brachystachyum, *DC. Coll. Mém.* vi. t. 12. f. 1.—Reg. Magellan.
imbricatum, *Poepp. & Endl. Nov. Gen. et Sp.* i. t. 3.—Chili.

MYZODENDRON:—
lineare, Poepp. & Endl. Nov. Gen. et Sp. i. 1. t. 1 = linearifolium.
linearifolium, *DC. Prod.* iv. 671.—Chili.
macrolepis, *Phil. in Anal. Univ. Chil.* (1865) II. 316.—Chili.
microphyllum, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 357 = linearifolium.
oblongifolium, *DC. Prod.* iv. 671.—Chili.
planifolium, Banks & Soland. ex Hook. f. Fl. Antarct. 297 = brachystachyum.
punctulatum, [*Banks*] *Soland. ex Forst. f. in Comm. Gotting.* ix. (1789) 45, nomen; *et ex DC. Prod.* iv. 286.—Reg. Magellan.
quadriflorum, *DC. Coll. Mém.* vi. t. 12. f. 2.—Reg. Magellan.

MYZODENDRUM, Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 45 = praec.

MYZORRHIZA, Phil. in Linnaea, xxix. (1857–58) 36 = **Aphyllon**, Mitch. (Oroban.).
chilensis, Phil. l. c.—Chili.

N

NABALUS, Cass. in Dict. Sc. Nat. xxxiv. 94 (1825) = **Prenanthes**, Vaill. (Compos.).
acerifolius, Maxim. in Bull. Acad. Pétersb. xv. (1871) 376.—Japon.
alatus, Hook. Fl. Bor. Am. i. 294. t. 102 = P. alata.
albus, Hook. l. c. 294 = P. alba.
altissimus, Hook. l. c. 294 = P. altissima.
asper, Torr. & Gray, Fl. N. Am. ii. 483 = P. aspera.
Boottii, DC. Prod. vii. 241 = P. Boottii.
cordatus, Hook. Fl. Bor. Am. i. 294 = P. altissima.
crepidineus, DC. Prod. ii. 242 = P. crepidinea.
deltoideus, DC. l. c. = P. altissima.
Fraseri, DC. l. c. 241 = P. serpentaria.
glaucus, DC. l. c. = P. glauca.
illinoënsis, DC. l. c. 242 = P. aspera.
integrifolius, Cass. in Dict. Sc. Nat. xxxiv. 96 = P. serpentaria, alba.
nanus, DC. Prod. vii. 241 = P. serpentaria.
nipponicus, Franch. & Sav. Enum. Pl. Jap. ii. 420.—Japon.
ochroleucus, Maxim. in Bull. Acad. Pétersb. xv. (1871) 376.—Manchuria.
quercifolius, Bertol. Misc. Bot. iv. 11.—Am. bor.
racemosus, Hook. Fl. Bor. Am. i. 294 = P. racemosa.
repens, Ledeb. Fl. Ross. ii. 840 = Lactuca repens.
roanensis, Chickering, in Coult. Bot. Gaz. v. (1880) 155 = P. serpentaria.
serpentarius, Hook. Fl. Bor. Am. i. 294 = P. serpentaria.
suavis, DC. Prod. vii. 241 = P. alba.
trifoliatus, Cass. in Dict. Sc. Nat. xxxiv. 95 = P. alba.
trilobatus, Cass. l. c. = P. serpentaria.
virgatus, DC. Prod. vii. 242 = P. virgata.

NABEA, Lehm. Ind. Sem. Hort. Hamb. (1831); ex Klotzsch, in Linnaea, viii. (1833) 666 = **Macnabia**, Benth. (Ericac.).
montana, Lehm. l. c. = M. montana.

NABLONIUM, Cass. in Dict. Sc. Nat. xxxiv. 101 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 341.
calyceroides, *Cass. l. c.*—Austral.

NACIBAEA, Poir. Encyc. Suppl. iv. 55 (1816) = seq.

NACIBEA, Aubl. Pl. Gui. i. 95. t. 37 = **Manettia**, Mutis (Rubiac.).
acutiflora, Poir. Encyc. Suppl. iv. 55 = M. acutifolia.
alba, Aubl. Pl. Gui. i. 96. t. 37. f. 1 = M. Lygistum.
angustifolia, Vahl, ex DC. Prod. iv. 364 = M. secundiflora.
coccinea, Aubl. Pl. Gui. i. 96 = M. Lygistum.

NACIBEA:—
glabra, Gaertn. f. Fruct. iii. 102. t. 197. f. 5 = Manettia Gaertneri.
mutabilis, Poir. Encyc. Suppl. iv. 55 = M. racemosa.
racemosa, Poir. in Dict. Sc. Nat. xxxiv. 110 = M. racemosa.
reclinata, Poir. in Lam. Encyc. iv. 416 = M. reclinata.
umbellata, Poir. Encyc. Suppl. iv. 55 = M. umbellata.
uniflora, Poir. in Dict. Sc. Nat. xxxiv. 110 = M. uniflora.

NAEGELIA, Lindl. Veg. Kingd. 560 (1847) = **Cotoneaster**, Medic. (Rosac.).

NAEGELIA, Regel, in Flora, xxxi. (1848) 249. *GESNERACEAE*, Benth. & Hook. f. ii. 999.
amabilis, Decne. in Fl. des Serres, xvi. (1865) 115 = multiflora.
cinnabarina, *Linden, Suppl. Cat. Pl. Exot. Jard. Brux.* (1856), cum fig.; *ex Hook. Bot. Mag.* t. 5036.—Mexic.
fulgida, *Ortgies, in Regel, Gartenfl.* (1867) 97.—Mexic.
Geroltiana, *Regel, in Flora*, xxxi. (1848) 249.—Mexic.
hyacinthina ×, Carr. in Rev. Hortic. (1877) 29.
multiflora, *Hook. Bot. Mag.* t. 5083.—Mexic.
punctata, *Hemsl. Biol. Centr. Am. Bot.* ii. 476.—Mexic.
secunda, *Oerst. ex Hanst. in Linnaea*, xxix. (1857–58) 510.—Mexic.
zebrina, *Regel, in Flora*, xxxi. (1848) 249.—Mexic.

NAEGELIA, Zoll. & Mor. Syst. Verz. Zoll. 20 (1845–46) = **Gouania**, Linn. (Rhamn.).
dubia, Zoll. & Mor. l. c. = G. leptostachya.
tomentosa, Hassk. in Flora, xxxv. (1852) 114 = G. javanica.

NAEMATOSPERMUM, Steud. Nom. ed. II. ii. 180 (1841) = NEMATOSPERMUM, Rich. = **Lacistema**, Sw.

NAGASSARI, Adans. Fam. ii. 449 (1763) = seq.

NAGATAMPO, Adans. Fam. ii. 444 (1763) = **Mesua**, Linn. (Guttif.).

NAGEIA, Gaertn. Fruct. i. 191 (1788), partim = **Myrica**, Linn.
arabica, Willd. Sp. Pl. iv. 749 = M. arabica.
japonica, Gaertn. Fruct. i. 191. t. 39 = M. Nagi.

NAGEIA, Gaertn. Fruct. i. 191. t. 39 (1788) partim = **Podocarpus**, L'Hérit. (Conif.).
alpina, F. Muell. in Papers Roy. Soc. Tasm. (1879) 23 = P. alpina.
amara, F. Muell. Sel. Pl. Indust. Cult. 138 = P. amara.
andina, F. Muell. l. c. = P. andina.
Blumei, Gord. Pinet. 135 = P. Blumei.
bracteata, F. Muell. Sel. Pl. Indust. Cult. 138 = P. macrophylla.
chilina, F. Muell. l. c. = P. chilina.
coriacea, F. Muell. l. c. = P. coriacea.
cupressina, F. Muell. Phyt. New Hebr. 20 = P. cupressina.
cuspidata, Gord. Pinet. 136 = P. cuspidata.
dacrydioides, F. Muell. Sel. Pl. Indust. Cult. 138 = P. dacrydioides.
Drouyniana, F. Muell. Census, 109 = P. Drouyniana.
elata, F. Muell. Sel. Pl. Indust. Cult. 138 = P. elata.
elongata, F. Muell. l. c. = P. elongata.
falcata, Carr. in Rev. Hortic. (1869) 370 = P. falcata.
ferruginea, F. Muell. Sel. Pl. Indust. Cult. 139 = P. ferruginea.
grandifolia, Gord. Pinet. 137 = P. grandifolia.
Lamberti, F. Muell. Sel. Pl. Indust. Cult. 139 = P. Lamberti.
latifolia, Gord. Pinet. 138 = P. latifolia?
macrophylla, F. Muell. Ausw. Aussertrop. Pfl. 245 = P. macrophylla.
minor, Carr. Conif. ed. II. 641 = P. minor.
nubigena, F. Muell. Sel. Pl. Indust. Cult. 139 = P. nubigena.
ovata, Gord. Pinet. Suppl. 42 = P. ovata.
Purdieana, F. Muell. Sel. Pl. Indust. Cult. 139 = P. Purdieana.
Rumphii, F. Muell. Papuan Pl. 92 = P. Rumphii.
spicata, F. Muell. Sel. Pl. Indust. Cult. 139 = P. spicata.

NAGEIA:—
spinulosa, F. Muell. Census, 109 = P. spinulosa.
Thunbergi, F. Muell. Sel. Pl. Indust. Cult. 139 = P. Thunbergii.
Totara, F. Muell. l. c. = P. Totara.

NAGEIA, Roxb. Hort. Beng. 71 (1814); Fl. Ind. iii. 766 (1832) = **Putranjiva**, Wall. (Euphorb.).
Putranjiva, Roxb. ll. cc. 71; 766 = P. Roxburghii.

NAGELIA, Lindl. in Bot. Reg. (1845) Misc. 40 = **Cotoneaster**, Medic. (Rosac.).
denticulata, Lindl. l. c. = C. denticulata.

NAGHAS, Mirb. ex Steud. Nom. ed. II. ii. 180 (1841) = **Mesua**, Linn. (Guttif.).

NAHUSIA, Schneev. Ic. 21 (1793) = **Fuchsia**, Linn. (Onagr.).
coccinea, Schneev. l. c. = F. coccinea.

NAIAS, Juss. Gen. 19 (1789) = seq.

NAJAS, Linn. Syst. ed. I (1735). *NAIADACEAE*, Benth. & Hook. f. iii. 1018.
CAULINIA, Willd. in Mém. Acad. Sc. Berl. 87 (1798).
FLUVIALIS, Mich. ex Adans. Fam. ii. 472 (1763).
ITTNERA, C. C. Gmel. Fl. Bad. iii. t. 4 (1808).
alagnensis, Masé, ex Nym. Consp. 685 = fragilis.
alagnensis, Pollini, Fl. Veron. iii. 49 = graminea.
ancistrocarpa, *A. Br. ex Magnus, Beitr. Najas*, p. vii.—Japon.
arguta, *H. B. & K. Nov. Gen. et Sp.* i. 371.—Am. austr.
australis, Bory, ex Cham. in Linnaea, iv. (1829) 501 = minor.
canadensis, Michx. Fl. Am. Bor. ii. 220 = flexilis.
conferta, *A. Br. in Sitzb. Ges. Naturf. Fr. Berl.* (1868) 17.—Hab.?
dichotoma, Roxb. Hort. Beng. 71; Fl. Ind. iii. 749 = minor.
falciculata, *A. Br. in Journ. Bot.* ii. (1864) 278.—Ind. or.; Ins. Philipp.
flexilis, *Rostk. & Schmidt, Fl. Sedin.* 382.—Europ.; Aegypt.
fluviatilis, Poir. in Lam. Encyc. iv. 416 = major.
foveolata, *A. Br. ex Magnus, Beitr. Najas*, p. vii.—Java.
fragilis, *Delile, Fl. Aeg. Illustr.* 75.—Aegypt.
fragilis, Rostk. & Schmidt, Fl. Sedin. 382 = minor.
fucoides, Griff. Notul. iii. 182 = major.
gracillima, A. Br. ex Magnus, Beitr. Najas, 23 = minor.
graminea, *Delile, Fl. Egypt.* 282.—Geront. temp. & trop.
guadalupensis, *Magnus, Beitr. Najas*, p. viii.—Ind. occ.
heteromorpha, *Griff. ex Voigt, Hort. Suburb. Calc.* 694, nomen.—Ind. or.
horrida, A. Br. ex Magnus, Beitr. Najas, 46 (= *Caulinia pectinata*).—Italia.
indica, Cham. in Linnaea, iv. (1829) 501 = minor.
intermedia, Gorski, in Eichw. Skizze, 126 = major.
latifolia, *A. Br. in Journ. Bot.* ii. (1864) 276.—Venezuela.
Leichhardtii, *Magnus, Beitr. Najas*, 50.—Austral.
lobata, Blanco, Fl. Filip. ed. II. 459 = minor.
major, *All. Fl. Pedem.* ii. 221.—Geront. temp. et trop.
marina, Linn. Sp. Pl. 1015 = major, minor.
maritima, Pall. Reise, i. 369 = major.
microdon, A. Br. in Sitzb. Ges. Naturf. Fr. Berl. (1868) 17 = guadalupensis.
microphylla, Reichb. Fl. Germ. Excurs. 843 = graminea, minor.
minor, *All. Fl. Pedem.* ii. 221.—Geront. temp. et trop.
monosperma, Willd. Sp. Pl. iv. 331 = major.
muricata, *Delile, Fl. Egypt.* 281.—Aegypt.
muricata, Thuill. Fl. Par. ed. II. 509 = major.
obvoluta, *Blanco, Fl. Filip.* ed. II. 460.—Ins. Philipp.
palustris, Blanco, l. c. ed. I. 660 = tenuifolia.
podostemon, *Magnus, Beitr. Najas*, p. vii.—Bras.
rigida, *Griff. Notul.* iii. 181 (sp. dub.).—Ind. or.
seminuda, Griff. l. c. 184 = graminea.
serristipula, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 72.—Japon.
serristipula, Nocca & Balb. Fl. Ticin. t. 15 = graminea.
spinosa, Buch.-Ham. in Wall. Cat. n. 5182 = major.
subulata, Thuill. Fl. Par. ed. II. 500 = minor.

NAJAS:—
tenera, *Schrad. in Goett. Gel. Anz.* i. (1821) 715.—Bras.
tenuifolia, Aschers. in Att. Soc. Ital. Sc. Nat. Milan. x. (1867) 267, 268 = graminea.
tenuifolia, *R. Br. Prod.* 345.—Austral.
tenuis, *A. Br. ex Magnus, Beitr. Najas*, p. vii. (sp. dub.).—Ind. or.
ternata, Roxb. ex Griff. Notul. iii. 183; Pl. As. t. 252 = minor.
tetrasperma, Willd. Sp. Pl. iv. 331 = major.
Wrightiana, *A. Br. in Sitzb. Ges. Naturf. Fr. Berl.* (1868) 17.—Ind. occ.

NALAGU, Adans. Fam. ii. 445 (1763) = **Aralia**, Linn.

NAMA, Linn. Diss. Dass. ii. (1747); Amoen. Acad. i. 394 (1749). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 831.
Berlandieri, A. Gray, in Proc. Am. Acad. viii. (1870) 282 = undulatum.
biflorum, *Choisy, in DC. Prod.* x. 183.—Mexic.
convolvuloides, Willd. ex Schult. Syst. vi. 189 = Evolvulus alsinoides.
Coulteri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 283.—Am. bor. occ.
demissum, *A. Gray, l. c.*—Am. bor. occ.
depressum, *Lemmon, ex A. Gray, l. c.* xx. (1885) 304.—Calif.
dichotomum, *Choisy, in DC. Prod.* x. 182.—Peruv.; Mexic.
echioides, Griseb. in Goett. Abh. xix. (1874) 230 = = undulatum.
erectum, *Roxb. Hort. Beng.* 84, nomen.—Ind. or.
evolvuloides, Willd. ex Choisy. in DC. Prod. ix. 447 = Evolvulus alsinoides.
Havardi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 304.—Texas.
hirsutum, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 277.—Mexic.
hispidum, *A. Gray, in Proc. Am. Acad.* v. (1861) 339.—Texas; Mexic.
jamaicense, *Linn. Syst.* ed. X. 950.—Ind. occ.
latifolium, *A. Gray, in Proc. Am. Acad.* viii. (1870) 284.—Mexic.
Lobbii, *A. Gray, l. c.* vi. (1864) 37.—Am. bor. occ.
longiflorum, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 114 = sericeum.
origanifolium, *H. B. & K. Nov. Gen. et Sp.* iii. 130. t. 218.—Am. bor. occ.; Mexic.
Palmeri, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 361.—Mexic.
Parryi, *A. Gray, in Bot. Calif.* i. 621.—Am. bor. occ.
pusillum, *Lemmon, ex A. Gray, in Proc. Am. Acad.* xx. (1885) 304.—Calif.
racemosum, Kellogg, in Proc. Calif. Acad. v. (1873) 51 = Phacelia namatoides.
Rothrockii, *A. Gray, in Bot. Calif.* i. 621.—Calif.
rupicolum, *Bonpl. ex Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 114.—Mexic.; Peruv.
sandwicense, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 338.—Ins. Sandvic.
Schaffneri, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 361.—Mexic.
sericeum, *Willd. ex Schult. Syst.* vi. 189.—Mexic.
serpylloides, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 363.—Mexic.
stenocarpum, *A. Gray, in Proc. Am. Acad.* x. (1875) 331.—Calif.
stenophyllum, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 361.—Mexic.; Texas.
strictum, *Phil. Fl. Atac.* 37.—Chili.
subincanum, Willd. ex Schult. Syst. vi. 189 = origanifolium.
subpetiolare, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 365.—Mexic.
systylum, *A. Gray, in Proc. Am. Acad.* v. (1861) 37.—Calif.
tetrandrum, Pav. ex Steud. Nom. ed. II. ii. 180 = dichotomum.
undulatum, A. Gray, in Proc. Am. Acad. viii. (1870) 282 = stenocarpum.
undulatum, *H. B. & K. Nov. Gen. et Sp.* iii. 130.—Mexic.
zeylanicum, Linn. Sp. Pl. 226 = Hydrolea zeylanica.

NANANTHEA, DC. Prod. vi. 45 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 428.
perpusilla, *DC. l. c.*—Corsica; Sardin.

NANDINA, Thunb. Nov. Gen. i. 14 (1781). *BERBERIDEAE*, Benth. & Hook. f. i. 44.
domestica, *Thunb. l. c.*; *Fl. Jap.* 147.—Japon.; China.
Tsermonanten, Hassk. Cat. Hort. Bog. Alt. 180 (= praec. ?).—Japon.

NANGHA, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1830) 178, nomen = **Artocarpus** ?, Forst. (Urtic.).
amplifolia, Zipp. l. c. = A. integrifolia ?

NANI, Adans. Fam. ii. 88 (1763) = seq.

NANIA, Miq. Fl. Ind. Bat. i. 1. 399 (1855) = **Metrosideros**, Banks (Myrtac.).
vera, Miq. l. c. 400.—Ins. Amboina.

NANNOGLOTTIS, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 480. *COMPOSITAE*.
carpesioides, *Maxim. l. c.* 481.—China.

NANNORRHOPS, H. Wendl. in Bot. Zeit. xxxvii. (1879) 147. *PALMAE*, Benth. & Hook. f. iii. 923.
Ritchieana, *H. Wendl. l. c.* 148.—Ind. bor. occ.

NANOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 154. t. 17. 1855 (1856). *URTICACEAE*, Benth. & Hook. f. iii. 381.
japonica, *Blume, l. c.* 155. t. 17.—Japon.; Corea.
lobata, *Wedd. in DC. Prod.* xvi. 69.—Ins. Loochoo.

NANODEA, Banks, ex Gaertn. f. Fruct. iii. 251. t. 225 (1805). *SANTALACEAE*, Benth. & Hook. f. iii. 226.
muscosa, *Gaertn. f. l. c.*—Reg. Magellan.

NANODES, Lindl. Bot. Reg. t. 1541 (1832) = **Epidendrum**, Linn. (Orchid.).
discolor, Lindl. l. c.—Bras.
Medusae, Reichb. f. in Fl. des Serres, xvii. (1867–68) 78.—Am. trop.

NANOLIRION, Benth. in Benth. & Hook. f. Gen. iii. 793 (1883). *LILIACEAE*, Benth. & Hook. f. l. c.
capense, *Benth. l. c.*—Afr. austr.

NANOPETALUM, Hassk. in Versl. Kon. Akad. Wetensch. iv. (1855) 140 = **Cleistanthus**, Benth. (Euphorb.).
myrianthum, Hassk. l. c. 141 = C. myrianthus.

NANOPHYTON, Less. in Linnaea, ix. (1834) 197. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 73.
caspicum, *Less. l. c.*—Reg. Casp.
erinaceum, Bunge, Anabas. Rev. 51 = caspicum.
juniperinum, C. A. Mey. in. Bull. Phys-Math. Acad. Pétersb. ii. (1844) 133 = caspicum.
macranthum, Fisch. & Mey. l. c. 134 = caspicum.

NANOPHYTUM, Endl. Gen. 299 (1837) = **Nanophyton**, Less. (Chenop.).

NANOTHAMNUS, T. Thoms. in Journ. Linn. Soc. ix. (1867) 342. t. 8. *COMPOSITAE*, Benth. & Hook. f. ii. 293.
sericeus, *T. Thoms. l. c.*—Ind. or.

NANSIATUM, Miq. Fl. Ind. Bat. i. II. 797 (1855), sphalm. = **Natsiatum**, Buch.-Ham. (Olacin.).

NAPAEA, Linn. Syst. ed. VI. Add. (1748). *MALVACEAE*, Benth. & Hook. f. i. 201.
SCHIZOICA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 249.
crispa, Moench, Meth. Suppl. 207 = Abutilon crispum.
dioecia, Hill, Veg. Syst. xv. 50 = seq.
dioica, *Linn. Sp. Pl.* 686.—Am. bor.
discolor, Alef. in Bot. Zeit. xxv. (1867) 220 = Plagianthus sidoides.
divaricata, Alef. l. c. = Plagianthus divaricatus.
hermaphrodita, Linn. Sp. Pl. 686 = dioica.
incurva, Moench, Meth. Suppl. 206 = Abutilon hirtum.
laevis, Linn. Mant. ii. 435 = dioica.
latifolia, Blanco, Fl. Filip. ed. II. 387 = Malachra capitata.

NAPAEA:—
lobata, Moench, Meth. 621 = dioica.
palmata, Moench, l. c. = dioica.
pulchella, Alef. in Bot. Zeit. xxv. (1867) 220 = Plagianthus pulchellus.
rhombifolia, Moench, Meth. 621 = Sida rhombifolia.
scabra, Blanco, Fl. Filip. ed. I. 553 = Malachra horrida.
scabra, Linn. Mant. ii. 435 = dioica.

NAPEA, Crantz, Inst. i. 200 (1766) = praec.

NAPEANTHUS, Gardn. in Hook. Lond. Journ. ii. (1843) 13. *GESNERACEAE*, Benth. & Hook. f. ii. 1018.
MARSSONIA, Karst. Fl. Columb. i. 97. t. 48 (1859 ?).
brasiliensis, *Gardn. in Hook. Lond. Journ.* ii. (1843) 13.—Bras.
primulinus, *Benth. & Hook. f. Gen.* ii. 1018.—N. Granat.
subacaulis, *Benth. & Hook. f. l. c.*—Ins. Trinit.

NAPELLUS, Rupp. Fl. Jen. ed. Hall. 275 (1745) = **Aconitum**, Linn. (Ranunc.).
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 327 = A. Napellus.

NAPIMOGA, Aubl. Pl. Gui. i. 592 (1775) = **Homalium**, Jacq. (Samyd.).
guianensis, Aubl. l. c. = H. Napimoga.

NAPOEA, Hill, Veg. Syst. xv. 50 (1769) = **Napaea**, Linn. (Malv.).

NAPOLEONA, Beauv. Fl. Owar. ii. 29. t. 7 (1805). *MYRTACEAE*, Benth. & Hook. f. i. 723.
BELVISIA, Desv. Journ. Bot. iv. (1814) 130.
angolensis, *Welw. Apont.* 586. n. 25.—Afr. trop.
cuspidata, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 10 = imperialis.
Heudelotii, *A. Juss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 227.—Afr. trop.
imperialis, *Beauv. Fl. Owar.* ii. 32. t. 78.—Afr. trop.
imperialis, Hook. Bot. Mag. t. 4387 = Whitfieldii.
Mannii, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 11 = imperialis.
Vogelii, Hook. & Planch. in Hook. Ic. Pl. t. 799–800 = imperialis.
Whitfieldii, Decne. in Rev. Hortic. (1853) 301. t. 16 = imperialis.
Whitfieldii, *Van Houtte, ex Lem. in Fl. des Serres*, i. (1843) 2.—Afr. trop.

NAPUS, [Tourn.] Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 939 (1829) = **Brassica**, Linn. (Crucif.).
Agriasinapis, Schimp. & Spenn. l. c. 944 = B. Sinapis.
campestris, Schimp. & Spenn. l. c. 941, in obs. = B. campestris.
Leucosinapsis, Schimp. & Spenn. l. c. 943 = B. alba.
oleifera, Schimp. & Spenn. l. c. 940 = B. campestris.
oleracea, Schimp. & Spenn. l. c. 939 = B. oleracea.
Rapa, Schimp. & Spenn. l. c. 941 = B. campestris.
Villarsii, Schimp. & Spenn. l. c. 942 = B. Cheiranthus.

NARAVEL, Adans. Fam. ii. 460 (1763) = seq.

NARAVELIA, DC. Syst. i. 167 (1818). *RANUNCULACEAE*, Benth. & Hook. f. i. 4.
dasyoneura, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 208.—Borneo.
Finlaysoniana, Wall. Cat. n. 4686 = laurifolia.
laurifolia, *Wall. l. c.* n. 4685.—Malaya.
pilulifera, Hance, in Journ. Bot. vi. (1868) 111 = zeylanica.
zeylanica, *DC. Syst.* i. 167.—Ind. or.; Malaya.

NARBALIA, Rafin. New Fl. Am. iv. 84 (1838) = **Prenanthes**, Vaill. (Compos.).
latifolia, Rafin. l. c. = P. Serpentaria ?

NARCISSOIDES, Heist. Syst. 5 (1748) (Quid ?).

NARCISSOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = **Narcissus**, Tourn. (Amaryll.).

NARCISSULUS, Heist. Syst. 5 (1748) (Quid ?).

NARCISSUS, [Tourn.] Linn. Syst. ed. I (1735).
AMARYLLIDEAE, Benth. & Hook. f. iii. 718.
AJAX, CORBULARIA, GANYMEDES, HERMIONE, PHILOGYNE, QUELTIA, Salisb. in Trans. Hort. Soc. i. (1812) 351–357.
ARGENOPE, CHIONE, CYDERIS, PANZA, PATROCLES, PLATEANA, PRASITELES, TITYRUS, VENIERA, Salisb. Gen. Pl. Fragm. 99–102 (1866).
ASSARACUS, Haw. Monog. Narciss. 4 (1831).
ASSARCUS, Kunth, Enum. Pl. v. 717 (1850).
AURELIA, J. Gay, in Ann. Sc. Nat. Sér. IV. x. (1858) 95.
AUTOGENES, Rafin. Fl. Tellur. iv. 20 (1836).
CALATHINUS, Rafin. l. c. 21 (1836).
CHLORASTER, Haw. in Phil. Mag. lxiii. (1824) 103.
CODIAMINUM, Rafin. Fl. Tellur. iv. 21 (1836).
DIOMEDES, Haw. in Phil. Mag. lxii. (1823) 440.
HELENA, Haw. Monog. Narciss. 13 (1831).
ILLUS, Haw. l. c. 4 (1831).
JONQUILLA, Haw. l. c. 7 (1831).
MOSKERION, Rafin. Fl. Tellur. iv. 21 (1836).
OILEUS, Haw. Monog. Narciss. 4 (1831).
SCHISANTHES, Haw. Suppl. Pl. Succ. 128 (1819).
STEPHANOPHORIUM, Dulac, Fl. Hautes-Pyr. 133 (1867).
TROS, Haw. Monog. Narciss. 5 (1831).
abscissus, Schult. f. Syst. vii. 941 = N. Pseudo-Narcissus.
aequilimbus, Nym. Syll. 365 = serotinus.
albicans, Spreng. Syst. ii. 45 = N. Pseudo-Narcissus.
albidus, Heyne, Nom. i. 539 = N. Pseudo-Narcissus.
albidus, Schult. f. Syst. vii. 949 = incomparabilis.
albus, Haw. in Trans. Linn. Soc. v. (1800) 243 = moschatus.
albus, Mill. Gard. Dict. ed. VIII. n. 5 = biflorus.
albus, Spach, Hist. Vég. Phan. xii. 440 = incomparabilis.
algirus, *Pomel, Nouv. Mat. Fl. Atl.* 384.—Alger.
Amancaes, Ruiz & Pav. Fl. Per. iii. 53. t. 281 = Hymenocallis Amancaes.
ambrosiacus, *Risso, Fl. Nice*, 476.—Gall. austr.
amplus, Salisb. Prod. 224 = incomparabilis.
anceps, DC. Cat. Hort. Monsp. 127; et in Red. Lil. viii. sub t. 486 = papyraceus.
Andersoni, Sabine, ex M. Roem. Syn. Ensat. 198 = N. Pseudo-Narcissus.
angustifolius, Curt. Bot. Mag. t. 193 = poëticus.
angustifolius, Hort. ex M. Roem. Syn. Ensat. 233 = gracilis.
angustifolius, *Willd. Enum. Hort. Berol.* 351.—Hab.?
annuus, Bull. Férussac, xiii. (1828) 328, sphalm. = aureus.
aperticorona, Steud. Nom. ed. II. ii. 181 = N. Tazetta.
apodanthus, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 25 = juncifolius.
arcuatus, Heynh. Nom. i. 539 = cernuus.
Aschersonii, Bolle, in Atti Soc. Ital. Sc. Nat. viii. (1865) 90 = N. Tazetta.
Assoanus, Dufour, ex Schult. f. Syst. vii. 962 = odorus.
Assoi, Dufour, ex Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 481 = juncifolius.
auranticoronus, Spach, Hist. Vég. Phan. xii. 448 = N. Tazetta.
aurantius, Schult. f. Syst. vii. 948 = incomparabilis.
aureo-tinctus ×, T. Moore & Ayres, Gard. Mag. Bot. iii. (1851) 289.
aureus, [Delile,] Red. Lil. viii. sub t. 486 = N. Bulbocodium.
aureus, *Loisel. Nouv. Not.* 13.—Reg. Mediterr.
autumnalis, Link, in Linnaea, ix. (1834) 569 = elegans.
Barlae, Parl. Fl. Ital. iii. 129 = N. Tazetta.
bericus, Turra, ex Marzari, Elenco (1802) 27; et ex Sacc. in Atti Ist. Venet. Ser. III. ix. (1863–64) 486 = incomparabilis.
Bernardi, DC. ex Hénon, in Mém. Acad. Lyon, iv. (1854) 5 = Macleayi.
Bertolonii, Parl. Fl. Ital. iii. 132 = N. Tazetta.
Biancae, Tod. in Ind. Sem. Hort. Panorm. (1857) 43 = N. Tazetta.
Bicchianus, Parl. Fl. Ital. iii. 156 = N. Tazetta.
bicolor, Auct. ex Schult. f. Syst. vii. 944 = lorifolius.
bicolor, Linn. Sp. Pl. ed. II. 415 = N. Pseudo-Narcissus.
bicrenatus, Hort. ex Nichols. Dict. Gard. ii. 411 = N. Tazetta.
biflorus, *Curt. Bot. Mag.* t. 197.—Europ.
biflorus, Gren. & Godr. Fl. Fr. iii. 256 = N. Pseudo-Narcissus.
biflorus, Schur, Enum. Pl. Transs. 658 = radiiflorus.
bifrons, Ker-Gawl. in Bot. Mag. t. 1186 = N. Tazetta.

NARCISSUS:—
biscrenatus, *Haw. ex Spach, Hist. Vég. Phan.* xii. 444. —Europ. austr.
breviflorus, Spach, Hist. Vég. Phan. xii. 448 = N. Tazetta.
breviflos, Steud. Nom. ed. II. ii. 181 = N. Pseudo-Narcissus.
Broussonetii, *Lag. Gen. et Sp. Nov.* 13.—Marocc.
Bulbocodium, *Linn. Sp. Pl.* 289.—Hispan.; Lusitan.
byzantinus, Hort. ex Boiss. Fl. Orient. v. 151 = N. Tazetta.
byzantinus, *Turra, ex Sacc. in Atti Ist. Venet.* Ser. III. ix. (1863–64) 487.—Oriens.
calathinus, Colm. ex Willk. & Lange, Prod. Fl. Hisp. i. 153 = triandrus.
calathinus, Delarb. Fl. Auv. i. 138 = heminalis.
calathinus, Linn. Sp. Pl. ed. II. 415 = odorus.
calathinus, Loisel. Fl. Gall. ed. II. i. 235 = capax.
cambricus, Steud. Nom. ed. II. ii. 181 = N. Pseudo-Narcissus.
campanulatus, Link, Enum. Hort. Berol. i. 311, in obs. = triandrus.
Campernelli, Hort. ex Haw. Monog. Narciss. 6 = odorus.
canaliculatus, Guss. Enum. Pl. Inar. 329. t. 15 = N. Tazetta.
canariensis, Burbidge, Narciss. 58. t. 48 = N. Tazetta.
candicans, Haw. ex Willk. & Lange, Prod. Fl. Hisp. i. 151 = N. Bulbocodium.
candidissimus, [DC. in] Red. Lil. t. 188 = N. Pseudo-Narcissus.
cantabricus, [DC.] l. c. viii. sub t. 486 = N. Bulbocodium.
capax, *Schult. f. Syst.* vii. 950.—Gallia.
cerinus, Schult. f. l. c. 972 = N. Tazetta.
cernuus, Bourg. ex Nym. Consp. 710 = triandrus.
cernuus, Roth, Catalect. fasc. i. 43 = N. Pseudo-Narcissus.
cernuus, *Salisb. Prod.* 223.—Gallia; Hispan.
Chloraster, *Schult. f. Syst.* vii. 981.—Afr. bor.
chrysanthus, DC. Fl. Fr. Suppl. 323 = N. Tazetta.
citrinus, Link, Handb. i. 202; Schult. f. Syst. vii. 968 = N. Tazetta.
Clusii, Dun. in Mém. Acad. Montp. (Sect. Sc.) i. (1847) 9 = N. Bulbocodium.
commutatus, Parl. Fl. Ital. iii. 152, in obs. = N. Tazetta.
compressus, *Haw. in Trans. Linn. Soc.* v. (1800) 245. —Europ. austr.
concolor, Link, Handb. i. 201; Schult. f. Syst. vii. 949 = triandrus.
conspicuus, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 326 = N. Bulbocodium.
conspicuus, Salisb. Prod. 224 = odorus.
constantinopolitanus, Hort. ex Boiss. Fl. Orient. v. 151 = N. Tazetta.
Coornei, [*DC. in*] *Red. Lil.* viii. sub t. 486.—Hab.?
corcyrensis, Nym. Syll. 365 = N. Tazetta.
corifolius, Steud. Nom. ed. II. ii. 181, sphalm. = lorifolius.
coronatus, [*DC. in*] *Red. Lil.* viii. sub t. 486.—Hab.?
cothurnalis, Salisb. Prod. 225 = biflorus.
cothurnalis, Williams, ex Steud. Nom. ed. II. ii. 181 = odorus.
crenulatus, Haw. in Trans. Linn. Soc. v. (1800) 245 = papyraceus.
crispicoronus, Spach, Hist. Vég. Phan. xii. 448 = N. Tazetta.
croceo-cinctus, *Spach, l. c.* 453.—Hab.?
cuneiflorus, Link, Handb. i. 204 = N. Pseudo-Narcissus.
Cupanianus, Grech-Delic. Fl. Melit. 36 (1853) = serotinus.
Cupanianus, Guss. Fl. Sic. Syn. i. 382 = elegans.
cupularis, Bertol. in Schult. f. Syst. vii. 970 = N. Tazetta.
cupularis, Puccin. ex Parl. Fl. Ital. iii. 133 = Puccinellii.
Curtisii, Spach, Hist. Vég. Phan. xii. 441 = odorus.
cyclamineus, [DC. in] Red. Lil. viii. sub t. 486 = N. Pseudo-Narcissus.
Cypri, Haw. Monog. Narciss. 10; Sweet, Brit. Flow. Gard. Ser. II. t. 92 = N. Tazetta.
decorus, Spach, Hist. Vég. Phan. xii. 449 = N. Tazetta.
deficiens, Herb. in Lindl. Bot. Reg. (1847) t. 22 = serotinus.
deflexicaulis, Spach, Hist. Vég. Phan. xii. 445 = N. Tazetta.
dianthos, Haw. Monog. Narciss. 14 = biflorus.
dubius, Gouan, Illustr. 22 = N. Tazetta.

NARCISSUS:—
elatior, Haw. in Trans. Linn. Soc. v. (1800) 244 = odorus.
elatus, Guss. in Bertol. Fl. Ital. vii. 627 = N. Tazetta.
elegans, *Spach, Hist. Vég. Phan.* xii. 452.—Reg. Mediterr.
etruscus, Parl. Fl. Ital. iii. 146 = N. Tazetta.
exiguus, Salisb. Prod. 220 = N. Pseudo-Narcissus.
eystettensis, Hort. ex Gard. Chron. (1884) i. 484 = N. Pseudo-Narcissus.
festalis, Salisb. Prod. 220 = N. Pseudo-Narcissus.
fistulosus, Schult. f. Syst. vii. 972 = N. Tazetta.
flaveolus, Spach, Hist. Vég. Phan. xii. 445 = N. Tazetta.
flavus, Lag. Gen. et Sp. Nov. 13 = N. Jonquilla.
flexiflorus, Spach, Hist. Vég. Phan. xii. 447 = N. Tazetta.
floribundus, Schult. f. Syst. vii. 968 = N. Tazetta.
Flos-lactis, Steud. Nom. ed. II. ii. 181 = N. Tazetta?
gaditanus, Boiss. & Reut. Diagn. Ser. II. iv. 96 = juncifolius.
galanthifolius, *Schult. f. Syst.* vii. 950.—Mont. Pyren.
Gennarii, Parl. Fl. Ital. iii. 130 = N. Tazetta.
Gigas, Steud. Nom. ed. II. ii. 181 = N. Bulbocodium.
glaucifolius, Pourr. in Mém. Acad. Toul. iii. (1788) 322 = N. Tazetta.
glaucus, Hornem. Hort. Hafn. i. 315 = propinquus, N. Pseudo-Narcissus.
Gouani, Roth, in Roem. Arch. i. ii. 39 = incomparabilis.
gracilis, *Sabine, in Bot. Reg.* t. 816.—Europ.
Graellsii, Webb, ex Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 471; Ind. Pl. Nov. 8 = N. Bulbocodium.
grandicrenatus, Parl. Fl. Ital. iii. 154 = N. Tazetta.
grandiflorus, Haw. Suppl. Pl. Succ. 142, in syn. = N. Trewianus.
grandiflorus, Salisb. Prod. 221 = N. Pseudo-Narcissus.
Haworthii, G. Don, in Loud. Hort. Brit. 116 = triandrus.
heminalis, *Schult. f. Syst.* vii. 960.—Gall. austr.
herminicus, Link, ex Willk. & Lange, Prod. Fl. Hisp. i. 152 = juncifolius.
hispanicus, Gouan, Illustr. 23 = N. Pseudo-Narcissus.
hispanicus, Willd. ex Willk. & Lange, Prod. Fl. Hisp. i. 152 = juncifolius.
homochroos, Schult. f. Syst. vii. 955 = triandrus.
Horsfeldii, Hort. ex Burbidge, Narcissus, 30 = N. Pseudo-Narcissus.
incomparabilis, *Mill. Gard. Dict.* ed. VIII. n. 3.— Europ. austr.
inflatus, Haw. in Trans. Linn. Soc. v. (1800) 243 = N. Bulbocodium.
infundibularis, Salisb. in Loud. Hort. Brit. 116 = seq.?
Infundibulum, Poir. Encyc. iv. 427 = heminalis, odorus.
integer, Spach, Hist. Vég. Phan. xii. 443 = viridiflorus.
interjectus, Schult. f. Syst. vii. 960 = odorus.
intermedius, Loisel. Fl. Gall. i. 237. t. 7 = N. Tazetta.
italicus, Ker-Gawl. in Bot. Mag. t. 1188 = N. Tazetta.
jasmineus, Schult. f. Syst. vii. 1734 = N. Tazetta.
Jonquilla, *Linn. Sp. Pl.* 290.—Reg. Mediterr.
Jonquilla, Pourr. ex Nym. Consp. 712 = juncifolius.
jonquilloides, Willd. ex Schult. f. Syst. vii. 968 = triandrus.
jonquilloides, Willk. in Bot. Zeit. xviii. (1860) 103 = N. Jonquilla.
juncifolius, *Req. ex Lag. Gen. et Sp. Nov.* 13; *et Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 406.—Gallia.
juncifolius, Salisb. Prod. 223 = N. Jonquilla.
lacticolor, Steud. Nom. ed. II. ii. 182 = N. Tazetta.
laetus, Salisb. Prod. 224 = odorus.
latifolius, Schult. f. Syst. vii. 972 = N. Tazetta.
Leedsii ×, T. Moore, in Moore & Ayres, Gard. Mag. Bot. iii. (1851) 169.
leucojifolius, Auct. ex Schult. f. Syst. vii. 968 = teretifolius.
lobatus, Poir. Encyc. iv. 427 = odorus.
lobularis, Schult. f. Syst. vii. 1732 = N. Pseudo-Narcissus.
lobulatus, Haw. Misc. Nat. 178 = N. Bulbocodium.
longiflorus, Willd. Enum. Hort. Berol. 351 = N. Tazetta?
longipetalus, Schleich. ex Steud. Nom. ed. II. ii. 182 = poëticus.
lorifolius, Schult. f. Syst. vii. 944 = N. Pseudo-Narcissus.
Luna, Schult. f. l. c. 974 = N. Tazetta.

NARCISSUS:—
Macleayi, *Lindl. in Bot. Reg.* sub t. 762 (Macleaii). —Mont. Pyren.
majalis, Curt. Bot. Mag. sub t. 193 = poëticus.
major, Curt. l. c. t. 51 = N. Pseudo-Narcissus.
Mastersianus ×, Hort.—Cf. Gard. Chron. (1882) I. 472.
maximus, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 286 = N. Pseudo-Narcissus.
medio-luteus, Mill. Gard. Dict. ed. VIII. n. 4 = biflorus.
medius, M. Roem. Syn. Ensat. 235 = M. Jonquilla.
megacodium, Dur. ex Munby, Fl. Alger. 34 = N. Bulbocodium.
mignon, Kunth, Enum. Pl. v. 730 in syn. sphalm. = *mion*.
minimus, Kunth, l. c. 895 (Index) = N. Pseudo-Narcissus.
minor, Linn. Sp. Pl. ed. II. 415 = N. Pseudo-Narcissus.
minutiflorus, Willk. in Bot. Zeit. xviii. (1860) 104 = juncifolius.
mion, Schult. f. Syst. vii. 961 = heminalis.
monophyllus, T. Moore, in Florist (1870) 84 = N. Bulbocodium.
montanus, Ker-Gawl. in Bot. Reg. t. 123 = poculiformis.
moschatus, *Linn. Sp. Pl.* ed. II. 415.—Hispan.; Gall.
multiflorus, Lam. Fl. Fr. iii. 391 = N. Tazetta.
muticus, J. Gay, ex Baker, in Flor. Mag. N. S. (1876) t. 224 = N. Pseudo-Narcissus.
nanus, Steud. Nom. ed. II. ii. 182 = N. Pseudo-Narcissus.
neglectus, Tenore, Ind. Sem. Hort. Neap. (1837) 5 = N. Tazetta.
Nelsoni ×, Hort. ex Vilm. Fl. Pl. Terre, Suppl. 122.
nivalis, Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 473 = N. Bulbocodium.
niveus, Loisel. in Desv. Journ. Bot. ii. (1809) 278 = papyraceus.
niveus, Panizz. ex Parl. Fl. Ital. iii. 128 = N. Tazetta.
nobilis, *Schult. f. Syst.* vii. 939.—Europ. austr.
nutans, *Haw. Misc. Nat.* 179.—Europ. austr.
obesus, Salisb. Prod. 222 = N. Bulbocodium.
obliquus, Guss. in Bertol. Fl. Ital. vii. 627 = N. Tazetta.
obliquus, Tausch, in Flora, xxiv. (1841) 237 = poëticus?
oblitteratus, Willd. ex Schult. f. Syst. vii. 981 = N. Broussonetii.
obscissus, Loud. Hort. Brit. 117, sphalm. = *abscissus*.
obsoletus, Spach, Hist. Vég. Phan. xii. 452 = serotinus.
obsoletus, Steud. Nom. ed. II. ii. 182 = elegans.
obvallaris, Salisb. Prod. 221 = N. Pseudo-Narcissus.
ochroleucus, Loisel. Not. 163 = N. Tazetta.
odorus, *Linn. Cent. Pl.* ii. 14; *Sp. Pl.* ed. II. 416.—Europ. austr.
orientalis, Linn. Mant. i. 38 = N. Tazetta.
orientalis, Tenore, Fl. Neap. i. 143 = N. Tazetta, biflorus.
ornatus, Haw. Monog. Narciss. 14 = poëticus.
oxypetalus, Boiss. Voy. Bot. Espagne, ii. 606 = elegans.
pachybolbus, Dur. in Duch. Rev. Bot. ii. (1846) 425 = N. Tazetta.
pallidulus, Graells, in Mem. Acad. Cienc. Madr. ii. (1854) 478 = triandrus.
pallidus, Graells, ex Nym. Consp. 713 = N. Bulbocodium.
pallidus, Poir. in Lam. Encyc. iv. 424 = N. Tazetta.
Panizzianus, Parl. Fl. Ital. iii. 128 = N. Tazetta.
papyraceus, *Ker-Gawl. in Bot. Mag.* t. 947.—Reg. Medit. occ.
papyratius, Spach, Hist. Vég. Phan. xii. 451 = N. Tazetta.
Parkinsoni, Steud. Nom. ed. II. ii. 182; Spach, l. c. 438 = Macleayi.
parvicorona, M. Roem. Syn. Ensat. 235 = N. Jonquilla?
parvulus, Sweet, Hort. Brit. ed. II. 515 = juncifolius.
patellaris, Salisb. Prod. 225 = poëticus.
patulus, Loisel. in Desv. Journ. Bot. ii. (1809) 276 = N. Tazetta.
perluteus, Spach, Hist. Vég. Phan. xii. 445 = N. Tazetta.
planicorona, Spach, l. c. 453 = gracilis.
poculiformis, *Salisb. Prod.* 224.—Mont. Pyren.
poëtarum, Haw. Monog. Narciss. 14 = poëticus.
poëticus, Huds. Fl. Angl. 122; ed. II. 141 = biflorus.
poëticus, *Linn. Sp. Pl.* 289.—Europ.
poëticus, Sibth. & Sm. Fl. Graec. Prod. 758 = radiiflorus.
polyanthos, Loisel. in Desv. Journ. Bot. ii. (1809) 277 = N. Tazetta.
praecox, Tenore, Cat. Hort. Neap. (1819) 39 = N. Tazetta.

NARCISSUS:—
primulinus, Haw. Suppl. Pl. Succ. 140, in syn. = N. Tazetta.
propinquus, *Salisb. Prod.* 221.—Europ. austr.
Pseudo-Jonquilla, Desf. Tabl. ed. II. 55 = juncifolius.
Pseudo-Narcissus, *Linn. Sp. Pl.* 289.—Europ.
Pseudo-poëticus, Boutigny, ex Nym. Consp. 710 = poëticus.
Puccinellii, *Parl. Fl. Ital.* iii. 133.—Ital.
pulchellus, *Salisb. Prod.* 223.—Europ. austr.
pumilus, [*Delile, in*] *Red. Lil.* vii. t. 409.—Hab.?
pumilus, Salisb. Prod. 220 = N. Pseudo-Narcissus.
purpureo-cinctus, Spach, Hist. Vég. Phan. xii. 453 = poëticus.
pusillus, G. Don, in Loud. Hort. Brit. 116 = juncifolius.
pyrenaicus, Pers. Syn. i. 350 = triandrus.
Queltia, Sweet, Hort. Brit. ed. I. 407 = incomparabilis.
radians, Lapeyr. Hist. Abr. Pl. Pyren. 177 = N. Pseudo-Narcissus.
radiatus, [Delile, in] Red. Lil. viii. t. 459 = N. Tazetta.
radiiflorus, *Salisb. Prod.* 225.—Europ. austr.
recurvifolius, Lodd. ex Steud. Nom. ed. II. ii. 183 = seq.?
recurvus, Haw. Syn. Pl. Succ. 331 = poëticus.
reflexus, Brot. Fl. Lusit. i. 550 = triandrus.
reflexus, Loisel. Not. 165 = odorus?
regulosus, Sweet, Hort. Brit. ed. III. 682, sphalm. = *rugulosus*.
remopolensis, Panizzi, in Parl. Giorn. Bot. Ital. ii. 1. (1847) 3 = N. Tazetta.
Renaudi, Bavoux, in Mém. Soc. Emul. Doubs, Sér. II. iv. (1854) 114 = N. Pseudo-Narcissus.
Requienii, M. Roem. Syn. Ensat. 236 = juncifolius.
Ricasolianus, Parl. Fl. Ital. iii. 147 = N. Tazetta.
rotularis, *Spach, Hist. Vég. Phan.* xii. 444.—Hab.?
rugilobus, Steud. Nom. ed. II. ii. 183 = N. Pseudo-Narcissus.
rugulosus, Link, Handb. i. 201; Schult. f. Syst. vii. 956 = odorus.
rupicola, Dufour, ex Schult. f. l. c. 958 = juncifolius.
Sabini, Lindl. in Bot. Reg. t. 762 = Macleayi.
semipartitus, Link, Handb. i. 200; Schult. f. Syst. vii. 949 = odorus.
seriorflorens, Schur, Enum. Pl. Transs. 657 = radiiflorus.
serotinus, *Linn. Sp. Pl.* 290.—Reg. Mediterr.
serotinus, Salzm. ex Schousb. Vextr. Marok. 155 = elegans.
serratus, *Haw. Misc. Nat.* 179.—Hab.?
sexlobatus, Haw. ex Schult. f. Syst. vii. 1723 = N. Tazetta?
Sibthorpii, Haw. in Trans. Linn. Soc. v. (1800) 24 = N. Pseudo-Narcissus.
siculus, Parl. Fl. Ital. iii. 149 = N. Tazetta.
similis, Steud. Nom. ed. II. ii. 183; M. Roem. Syn. Ensat. 235 = N. Jonquilla.
solaris, Spach, Hist. Vég. Phan. xii. 446 = N. Tazetta.
spathulatus, Haw. Monog. Narciss. 14 = poëticus.
spiralis, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 30 = serotinus.
spiralis, Parl. Fl. Ital. iii. 152 = N. Tazetta.
Spofforthiae ×, Herb. in Bot. Reg. (1843) t. 38. ff. 3, 4.
spurius, Schult. f. Syst. vii. 939 = N. Pseudo-Narcissus.
stellaris, Haw. Monog. Narciss. 15 = poëticus.
stellatus, DC. Fl. Fr. Suppl. 323 = papyraceus.
stellatus, Sprun. ex Nym. Consp. 710 = N. Tazetta.
stelliflorus, Schur, in Oestr. Bot. Zeitschr. xix. (1869) 205 = poëticus.
stramineus, Spach, Hist. Vég. Phan. xii. 451 = N. Tazetta.
striatellus, Spach, l. c. 438 = triandrus.
subalbidus, Loisel. Not. 163 = N. Tazetta.
sulcicaulis, Spach, Hist. Vég. Phan. xii. 449 = N. Tazetta.
sulphureus, Sweet, Hort. Brit. ed. I. 407 = incomparabilis.
sylvestris, Lam. Fl. Fr. iii. 390 = N. Pseudo-Narcissus.
syriacus, Boiss. & Gaill. Diagn. Ser. II. iv. 96 = N. Tazetta.
Tazetta, Ker-Gawl. in Bot. Mag. t. 925 = aureus.
Tazetta, *Linn. Sp. Pl.* 290.—Reg. Mediterr.
Tazetto-poëticus, Gren. & Godr. Fl. Fr. iii. 257 = biflorus.
tazettus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = N. Tazetta.
Telamonius, Link, Handb. i. 204; Schult. f. Syst. vii. 11. 940 = N. Pseudo-Narcissus.
Tenorii, Parl. Fl. Ital. iii. 137 = N. Tazetta.
tenuiflorus, Schult. f. Syst. vii. 1733 = N. Tazetta.

NARCISSUS:—
tenuifolius, Salisb. Prod. 222 = N. Bulbocodium.
tenuior, Curt. Bot. Mag. t. 379; Haw. in Trans. Linn. Soc. v. (1800) 245 = gracilis.
tenuis, Agardh, ex Schult. f. Syst. vii. 964 = praec.?
tereticaulis, *Haw. in Trans. Linn. Soc.* v. (1800) 245. —Hispan.
Timbali, Gautier, Herb. Comm. Agr. Narb. 18; ex Timb. in Bull. Soc. Sc. Phys. & Nat. Toul. iii. (1875-76) 143 = N. Tazetta.
Tinei, Tod. ex Parl. Fl. Ital. iii. 146, in textu = N. Tazetta.
tortuosus, Haw. Misc. Nat. 179 = N. Pseudo-Narcissus.
Trewianus, *Ker-Gawl. in Bot. Mag.* sub t. 1301.—Oriens.
triandrus, Curt. Bot. Mag. t. 48 = cernuus.
triandrus, *Linn. Sp. Pl.* ed. II. 416.—Mont. Pyren.
tridymus ×, Burbidge, in Gard. Chron. (1881) II. 603.
triflorus, Haw. Monog. Narciss. 14 = biflorus.
trilobus, Linn. Sp. Pl. ed. II. 417 = odorus.
tripartitus, Hornem. Hort. Hafn. i. 316 = N. Tazetta.
tripedalis, Lodd. ex Schult. f. Syst. vii. 987 = poëticus.
tripodalis, Salisb. ex Herb. Amaryll. 317 = praec.
tubaeflorus, Salisb. Prod. 221 = N. Pseudo-Narcissus.
turgidus, Salisb. l. c. 222 = N. Bulbocodium.
unicolor, Tenore, Fl. Neap. i. 144. t. 26 = N. Tazetta.
unicolor, Vent. Jard. Malm. t. 26 = papyraceus.
uniflorus, Hort. ex Schult. f. Syst. vii. 964 = gracilis.
varians, Guss. ex Parl. Fl. Ital. iii. 141 = N. Tazetta.
verbanensis, Herb. Amaryll. 318. t. 37 = poëticus.
vergellensis, Parl. Fl. Ital. iii. 148 = N. Tazetta.
viridiflorus, *Schousb. Vextr. Marok.* 157. t. 2.—Reg. Mediterr. occ.
Webbii, Parl. Fl. Ital. iii. 123, in textu = N. Jonquilla.

NARDA, Vell. Fl. Flum. 108 (1825); iii. t. 24 (1827) = **Strychnos**, Linn. (Logan.).
spinosa, Vell. ll. cc. = S. brasiliensis.

NARDINA, Murr. Syst. ed. XIV. 344 (1784), sphalm. = **Nandina**, Thunb. (Berberid.).

NARDOPHYLLUM, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 44. *COMPOSITAE*, Benth. & Hook. f. ii. 257.
ANACTINIA, Remy, in C. Gay, Fl. Chil. iv. 8 (1849).
DOLICHOGYNE, DC. Prod. vii. 257 (1838).
chiliotrichoides, *A. Gray, in Proc. Am. Acad.* v. (1862) 123.—Chili.
Darwini, *A. Gray, l. c.*—Patagon.
genistoides, *A. Gray, l. c.* viii. (1873) 636.—Chili.
humile, *A. Gray, l. c.* v. (1862) 123.—Patagon.
Kingii, *A. Gray, l. c.*—Patagon.
obtusifolium, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 144.—Patagon.
revolutum, *Hook. & Arn. l. c.*—Chili.

NARDOSMIA, Cass. in Dict. Sc. Nat. xxxiv. 186 (1825) = **Petasites**, Tourn. (Compos.).
angulosa, Cass. l. c. 188 = P. frigidus.
Billingsiana, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) I. 370 = *glacialis*.
corymbosa, Hook. Fl. Bor. Am. i. 307 = P. frigidus.
denticulata, Cass. in Dict. Sc. Nat. xxxiv. 187 = P. fragrans.
fragrans, Reichb. Fl. Germ. Excurs. 289 = P. fragrans.
frigida, Hook. Fl. Bor. Am. i. 307 = P. frigidus.
glacialis, Ledeb. Fl. Ross. ii. 466.—Sibir. arct.
Gmelini, DC. Prod. vii. 271 = Gerbera Anandria.
Hookeriana, Nutt. in Trans. Ann. Phil. Soc. N. S. vii. (1841) 288 = P. palmatus?
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 181 = P. japonicus.
laevigata, DC. Prod. v. 205 = P. laevigatus.
palmata, Hook. Fl. Bor. Am. i. 308 = P. palmatus.
racemosa, Pasq. Cat. Orto Nap. 112 = P. fragrans.
sagittata, Hook. Fl. Bor. Am. i. 307 = P. sagittatus.
saxatilis, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 138 = P. frigidus?
speciosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 289 = P. palmata?
spuria, Less. in Linnaea, ix. (1834) 156 = P. spurius.
straminea, Cass. in Dict. Sc. Nat. xxxiv. 189 = P. laevigatus.

NARDOSTACHYS, DC. Coll. Mém. vii. 4. tt. 1, 2 (1830). *VALERIANEAE*, Benth. & Hook. f. ii. 153.
grandiflora, DC. l. c. t. 2 = N. Jatamansi.
Jatamansi, *DC. l. c.*—Reg. Himal.

NARDURUS, Reichb. Fl. Germ. Excurs. 19, in obs. (1830); et Nom. 39 (1841) = **Festuca**, Tourn. (Gramin.).
Lachenalii, Godr. Fl. Lorr. ed. I. iii. 187 = F. Poa.
montanus, Boiss. & Reut. Pugill. Pl. Nov. 129 = F. montana, *Steud.*
orientalis, Boiss. Diagn. Ser. I. vii. 127 = F. aleppica.
patens, Hack. Cat. Gram. Portug. 31 = *F. loliacea.*
persicus, Boiss. & Buhse, in Mém. Soc. Nat. Mosc. xii. (1860) 225 = Poa persica.
Poa, Boiss. Voy. Bot. Espagne, ii. 667 = F. Poa.
Salzmanni, Boiss. l. c. = F. Salzmanni.
tenellus, Fries.—Cf. Linnaea, xxix. (1857–58) 730; Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 131 = F. montana? *Steud.*
tenuiculmis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 189 = F. Poa?
tenuiflorus, Boiss. Voy. Bot. Espagne, ii. 667 = F. tenuiflora.
unilateralis, Fries.—Cf. Linnaea, xxix. (1857–58) 730; Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 131 = F. tenuiflora.

NARDUS, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1205.
SPARTUM *vel* SPARTIUM, Heist. Syst. 12 (1748).
aristata, Linn. Sp. Pl. ed. II. 78 = Psilurus nardoides.
articulata, Linn. l. c. ed. I. 53 = Lepturus incurvatus.
ciliaris, Linn. l. c. = Perotis latifolia?
ciliata, Lam. Tabl. Encyc. i. 152 = praec.
dactyloides, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 269 = Panicum lanatum.
Gangitis, Linn. Sp. Pl. 53 = Ctenium carolinianum.
incurva, Gouan, Hort. Monsp. 33 = Psilurus nardoides.
indica, Linn. f. Suppl. 105 = Microchloa setacea.
monandra, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 438 = Psilurus nardoides.
scorpioides, Lam. Tabl. Encyc. i. 153 = Ctenium carolinianum.
stricta, *Linn. Sp. Pl.* 53.—Europ.; As. bor.
Thomaea, Linn. f. Suppl. 105 = Rottboellia Thomaea.

NAREGA, Rafin. Sylva Tellur. 98 (1838) = **Randia**, Linn. (Rubiac.).
Coduva, Rafin. l. c. = R. dumetorum.

NAREGAMIA, Wight & Arn. Prod. 116 (1834). *MELIACEAE*, Benth. & Hook. f. i. 331.
alata, *Wight & Arn. l. c.* 117.—Ind. or.
dentata, Miq. ex Hook. f. Fl. Brit. Ind. i. 542 = alata.

NARGEDIA, Bedd. Fl. Sylv. t. 328 (1873?). *RUBIACEAE*, Benth. & Hook. f. ii. 1228.
macrocarpa, *Bedd. Forest. Man. Bot.* 134/2. t. 328.—Ind. or.

NARICA, Rafin. Fl. Tellur. ii. 87 (1836) = **Spiranthes**, Rich. (Orchid.).
moschata, Rafin. l. c. = S. picta.

NARINGI, Adans. Fam. ii. 341 (1763) = **Limonia**, Linn. (Rutac.).

NARKETIS, Rafin. Fl. Tellur. iii. 26 (1836) = **Pleurogyne**, Eschsch. (Gentian.).
carinthiaca, Rafin. l. c. = P. carinthiaca.
hyperborea, Rafin. l. c. = praec.?
japonica, Rafin. l. c., nomen = P. rotata.
rotata, Rafin. l. c. = P. rotata.
sulcata, Rafin. l. c. = P. rotata.

NARON, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 419 = **Moraea**, Mill. (Irid.).
iridioideum, Moench, Meth. 627 = M. iridioides.
orientale, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 419 = M. iridioides.

NARTHECIUM, Moehr. in Act. Acad. Nat. Cur. vi. (1742) 384. t. 5. *LILIACEAE*, Benth. & Hook. f. iii. 825.
ABAMA, Adans. Fam. ii. 47 (1763).
alpinum, Michx. Fl. Bor. Am. i. 209, in obs. = Tofieldia calyculata.
americanum, Ker-Gawl. in Bot. Mag. t. 1505 = ossifragum.
anthericoides, Hoppe, ex Mert. & Koch, Deutschl. Fl. ii. 559 = ossifragum.

NARTHECIUM:—
asiaticum, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 438.—Japon.
boreale, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1805) 124 = Tofieldia palustris.
californicum, *Baker, in Journ. Linn. Soc.* xv. (1876) 351.—Calif.
calyculatum, Lam. Fl. Fr. iii. 298; All. Fl. Pedem. ii. 165 = Tofieldia calyculata.
falcatum, Poir. Encyc. Suppl. iv. 61 = Tofieldia falcata.
falcatum, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1805) 126 = Tofieldia glutinosa.
flavescens, Wahlenb. Veg. Helv. 68, in syn. = Tofieldia calyculata.
glutinosum, Michx. Fl. Bor. Am. i. 210 = Tofieldia glutinosa.
iridifolium, Vill. Hist. Pl. Dauph. ii. 225 = Tofieldia calyculata.
ossifragum, *Huds. Fl. Angl.* ed. I. 128; ed. II. 145.—Europ.; As. Min.; Am. bor.
pubens, Michx. Fl. Bor. Am. i. 209 = Tofieldia pubens.
pusillum, Michx. l. c. = Tofieldia palustris.
scabrum, *Rafin. Préc. Découv.* 45; *et in Desv. Journ. Bot.* iv. (1814) 273.—Am. bor.

NARTHEX, Falc. in Proc. Linn. Soc. i. 309 (1846); et in Trans. Linn. Soc. xx. (1846) 285 = **Ferula**, Tourn. (Umbell.).
Asafoetida, Falc. ll. cc. = F. Narthex.
Silphium, Oerst. in Overs. Danske Vidensk. Selsk. Forh. (1869) 21.—Afr. bor.

NARUKILA, Adans. Fam. ii. 54 (1763) = **Pontederia**, Linn.

NARUM, Adans. Fam. ii. 365 (1763) = **Ubaria**, Linn. (Anonac.).

NARVALINA, Cass. in Dict. Sc. Nat. xxxviii. 17 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 388.
NEEDHAMIA, Cass. in Dict. Sc. Nat. xxxiv. 335 (1825).
domingensis, *Cass. in Dict. Sc. Nat.* xxxviii. 17.—Ins. S. Doming.

NARVELIA, Link, Enum. Hort. Berol. ii. 89 (1822), sphalm. = **Naravelia**, DC. (Ranunc.).

NASMYTHIA, Huds. Fl. Angl. ed. II. ii. 414 (1778) = **Eriocaulon**, Linn.
articulata, Huds. l. c. 415 = E. septangulare.
septangularis, Mart. in Nov. Act. Nat. Cur. xviii. 1. (1835) t. 2 = E. septangulare.

NASONIA, Lindl. in Benth. Pl. Hartw. 150 (1844) = **Centropetalum**, Lindl. (Orchid.).
cinnabarina, Linden & Reichb. f. in Gard. Chron. (1867) 544 = *punctata.*
Hartwegii, Reichb. f. Xenia Orch. i. 97 = *punctata.*
isnanthera, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 99.—Peruv.
Myrtillus, Reichb. f. l. c. iii. (1855) 239 = C. Myrtillus.
punctata, Lindl. in Benth. Pl. Hartw. 150.—Peruv.
sanguinea, Lindl. Orch. Linden. 18.—N. Granat.

NASSAUVIA, Comm. ex Juss. Gen. 175 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 502.
ACANTHOPHYLLUM, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 37.
ANAGYRUM, Steud. Nom. ed. II. i. 84 (1840).
CALOPAPPUS, Meyen, Reise, i. 315 (1834).
CALOPTILIUM, Lag. Amen. Nat. i. 34 (1811).
MASTIGOPHORUS, Cass. in Dict. Sc. Nat. xxxiv. 222 (1825).
NASSOVIA, Batsch, Tab. 251 (1802).
PANARGYRUS, Lag. Amen. Nat. i. 33 (1811).
PENTANTHUS, Less. Syn. Comp. 397 (1832).
PIPTOSTEMMA, Spach, Hist. Vég. Phan. x. 34 (1841).
PORTALESIA, Meyen, Reise, i. 316 (1834).
SPHAEROCEPHALUS, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 67.
STRONGYLOMA, DC. Prod. vii. 52 (1838).
TRIACHNE, Cass. in Bull. Soc. Philom. (1818) 11.
TRIANTHUS, Hook. f. Fl. Antarct. ii. 320 (1846).
abbreviata, *Benth. & Hook. f. Gen.* ii. 503.—Patagon.
acerosa, *Wedd. Chlor. And.* i.51.—Chili.

NASSAUVIA:—
aculeata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 13. t. 20.—Chili.
axillaris, *D. Don, in Phil. Mag.* xi. (1832) 390.—Chili.
capillata, D. Don, in Phil. Mag. xi. (1832) 390, nomen = Triptilion capillatum.
cespitosa, *Wedd. Chlor. And.* i. 48.—Chili.
Commersonii, Cass. in Dict. Sc. Nat. xxxviii. 457 = magellanica.
cordifolia, D. Don, in Phil. Mag. xi. (1832) 390, nomen = Triptilion cordifolium.
Cumingii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 37.—Chili.
dentata, *Griseb. in Goett. Abh.* vi. (1854) 125.—Chili.
diffusa, D. Don, in Phil. Mag. xi. (1832) 390, nomen = Triptilion diffusum.
digitata, *Wedd. Chlor. And.* i. 48.—Chili.
Durvillei, Cass. in Dict. Sc. Nat. xxxviii. 456 = serpens.
Gaudichaudii, *Cass. l. c.* 457.—Reg. Magellan.
glabrata, *Phil. in Anal. Univ. Chil.* (1872) 741.—Chili.
glomerata, *Wedd. Chlor. And.* i. 53.—Chili.
glomerulosa, *D. Don, in Phil. Mag.* xi. (1832) 390, nomen.—Chili.
Lagascae, *Benth. & Hook. f. Gen.* ii. 503.—Chili.
lycopodioides, *Phil. in Linnaea*, xxviii. (1856) 714.—Chili.
macracantha, DC. Prod. vii. 49 = Cumingii.
magellanica, *J. F. Gmel. Syst.* 1281.—Reg. Magell.
multiflora, *Meyen, Reise*, i. 356.—Chili.
nivalis, *Poepp. & Endl. Nov. Gen. et Sp.* i. 13. t. 22.—Chili.
oligocephala, *Wedd. Chlor. And.* i. 53.—Chili; Reg. Argent.
pinnigera, *D. Don, in Phil. Mag.* xi. (1832) 390; *Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 37.—Chili.
planifolia, *Wedd. Chlor. And.* i. 48.—Chili.
pumila, *Poepp. & Endl. Nov. Gen. et Sp.* i. 13. t. 21.—Chili.
pungens, Phil. in Anal. Univ. Chil. (1872) 741 = pyramidalis.
pygmaea, *Hook. f. Fl. Antarct.* ii. 319.—Reg. Magellan.
pyramidalis, *Meyen, Reise*, i. 356.—Chili.
ramosissima, *DC. Prod.* vii. 49.—Chili.
Remyana, *Wedd. Chlor. And.* i. 51. t. 12 a.—Chili.
revoluta, *D. Don, in Phil. Mag.* xi. (1832) 390; *Gill. ex Hook. Comp. Bot. Mag.* i. (1835) 37.—Chili.
rosulata, *Ball, in Journ. Linn. Soc.* xxi. (1884) 225.—Patagon.
sericea, *Poepp. & Endl. Nov. Gen. et Sp.* i. 13.—Chili.
serpens, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 609. Ins. Falkland.
spicata, *Remy, in C. Gay, Fl. Chil.* iii. 343.—Chili.
spinosa, D. Don, in Phil. Mag. xi. (1832) 390, nomen = Triptilion spinosum.
sprengelioides, *DC. Coll. Mém.* ix. 35. t. 14; *et Prod.* iii. 49.—Chili.
suaveolens, D. Don, in Phil. Mag. xi. (1832) 389 = Cumingii.
suaveolens, Willd. Sp. Pl. iii. 2396 = magellanica.

NASSAVIA, Spreng. Gen. ii. 624 (1831) = praec.

NASSAVIA, Vell. Fl. Flum. 141 (1825); iii. tt. 155, 156 (1827) = **Schmidelia**, Linn. (Sapind.).
axillaris, Vell. ll. cc. 142; t. 155.—Bras.
terminalis, Vell. ll. cc. 141; t. 156.—Bras.

NASSAWIA, Lag. Amen. Nat. i. 34 (1811) = **Nassauvia**, Comm. (Compos.).

NASSELLA, E. Desv. in C. Gay, Fl. Chil. vi. 263. t. 75. f. 1 (1853) = **Oryzopsis**, Michx. (Gramin.).
caespitosa, Griseb. in Goett. Abh. xix. (1874) 258.—Reg. Argent.
chilensis, E. Desv. in C. Gay, Fl. Chil. vi. 267.—Chili.
exserta, Phil. in Linnaea, xxxiii. (1864–65) 278.—Chili.
humilis, Phil. l. c.—Chili.
juncea, Phil. l. c. 277.—Chili.
Landbecki, Phil. l. c. 279.—Chili.

NASSELLA:—
laxiflora, Phil. in Linnaea, xxxiii. (1864–65) 279.—Chili.
major, E. Desv. in C. Gay, Fl. Chil. vi. 265.—Chili.
melanocarpa, Phil. in Anal. Univ. Chil. (1873) 558.—Chili.
planifolia, Phil. l. c.—Chili.
pubiflora, E. Desv. in C. Gay, Fl. Chil. vi. 264.—Chili.
pungens, E. Desv. l. c. 258.—Chili.
ramosa, E. Desv. l. c. 266.—Chili.
Sadae, Phil. in Linnaea, xxxiii. (1864–65) 277.—Chili.

NASSOVIA, Batsch, Tab. 251 (1802) = **Nassauvia**, Comm. (Compos.).

NASTANTHUS, Miers, ex Lindl. Veg. Kingd. 701 (1847), nomen; et in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 184 = **Boopis**, Juss. (Calycer.).
agglomeratus, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 186.—Chili.
compactus, Miers, l. c. 189 = B. compacta.
Gayanus, Miers, l. c. = B. Gayana.
Gilliesii, Miers, l. c. 188.—Chili.
laciniatus, Miers, l. c. 187 = B. laciniata.
pinnatifidus, Miers, l. c.—Chili.
scapigerus, Miers, l. c. 188 = B. scapigera.
spathulatus, Miers, l. c. 189 = B. spathulata.
ventosus, Miers, l. c. 188 = Calycera ventosa.

NASTURTIOIDES, Medic. Pflanzengatt. i. 81 (1792); Moench, Meth. 270 (1794) = **Lepidium**, Tourn. (Crucif.).
inconspicuum, Medic. l. c. = L. ruderale.

NASTURTIOLUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 692 (1821) = **Hutchinsia**, R. Br. (Crucif.).
montanum, S. F. Gray, l. c. = H. petraea.

NASTURTIOLUM, Medic. Pflanzengatt. i. 82 (1792) = **Senebiera**, DC. (1799) (Crucif.).
castratum, Medic. l. c. = S. pinnatifida.
pinnatum, Moench, Meth. Suppl. 71 = S. pinnatifida.

NASTURTIOPSIS, Boiss. Fl. Orient. i. 237 (1867) = **Nasturtium**, Linn. (Crucif.).
arabica, Boiss. l. c.—Arabia.

NASTURTIUM, Linn. Syst. ed. I (1735); R. Br. in Ait. Hort. Kew. ed. II. iv. 109 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 68.
Brachiolobos, All. Fl. Pedem. i. 278 (1785).
Clandestinaria, Spach, Hist. Vég. Phan. vi. 427 (1838).
Leiolobrum, Reichb. Consp. 184 (1828).
Nasturtiopsis, Boiss. Fl. Orient. i. 237 (1867).
Radicula, Dill. ex Moench, Meth. 262 (1794).
Roripa, Scop. Fl. Carn. ed. I. 520 (1760).
Tetracellion, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 39.
Tetrapoma, Turcz. ex Fisch. & Mey. l. c. (1835).
acaule, *Welw. Apont.* 589.—Afr. trop.
album, Spreng. Syst. ii. 883 = Smelowskia cinerea.
alpinum, Crantz, Crucif. t. i. 80 = Hutchinsia alpina.
amphibium, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 110.—Reg. bor. temp.
anceps, Auct. ex Nym. Consp. 3 = sylvestre.
anceps, *DC. Prod.* i. 137.—Europ.
anethifolium, *Phil. Fl. Atac.* 7.—Chili.
apetalum, DC. Syst. ii. 200 = indicum.
aquaticum, Wahlenb. in Svensk Bot. t. 624 = officinale.
aquaticum, Wallr. Sched. Crit. 371 = amphibium.
arabiforme, *DC. Syst.* ii. 220.—Am. austr.
Armoracia, Fries, Fl. Scan. 65 = Cochlearia Armoracia.
armoracioides, *Tausch, in Flora*, xxiii. (1840) 707.—Europ.
arvense, Laest. ex Nym. Consp. 36 = palustre.
Aschersonianum, *Janka.—Cf. Just, Jahresber.* iv. (1876) 1061.—Europ.
asperum, Boiss. Voy. Bot. Espagne, ii. 28 = Sisymbrium Boissieri.
asperum, Coss. Notes Crit. 27 = Sisymbrium asperum.
astyloides, Knaf, ex Opiz, Seznam, 68 = sylvestre.
astylon, Reichb. Fl. Germ. Excurs. 684 = austriacum.

NASTURTIUM:—
atlanticum, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocc.
atrovirens, DC. Syst. ii. 201 = indicum.
aureum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 17.—Syria.
austriacum, *Crantz, Stirp. Austr.* ed. I. 15.—Europ. or.; As. Min.
barbaraeoides, Tausch, in Flora, xxiii. (1840) 713 = sylvestre.
barbareaefolium, *Baker, Fl. Maurit.* 7.—Ins. Maurit.
benghalense, DC. Syst. ii. 198 = indicum.
benghalense, Hance, in Journ. Linn. Soc. xiii. (1873) 99 = microspermum.
Boissieri, Coss. Notes Crit. ii. 26 = Sisymbrium Boissieri.
Boissieri, Noé, ex Boiss. Fl. Orient. i. 181 = thracicum.
bonariense, *DC. Syst.* ii. 193.—Reg. Argent.
brachycarpum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 8.—Rossia; Sibir.
Brachylobus, C. Muell. in Walp. Ann. vii. 100 = austriacum.
brachypus, *Webb, Fragm. Fl. Aethiop.* 13.—Afr. trop.
brachystylum, *Wallr. Beitr. Fl. Hercyn.* 236.—Europ.
brevipes, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 154.—Cuba.
brevistylon, Schur, Enum. Pl. Transs. 39 = sylvestre.
Bursa-pastoris, Roth, Tent. Fl. Germ. i. 281 = Capsella Bursa-pastoris.
bursifolium, *DC. Syst.* ii. 194.—Kamtchat.
caledonicum, *Sond. in Linnaea*, xxiii. (1850) 2.—Afr. austr.
Camelinae, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 34 = austriacum.
cantoniense, Hance, in Journ. Bot. iii. (1865) 378 = Cochlearia globosa.
ceratophyllum, *DC. Syst.* ii. 193.—Mauritan.
cernuum, Nutt. in Torr. & Gray, Fl. N. Am. i. 74 = curvisiliqua.
chilense, Hort. ex Steud. Nom. ed. II. ii. 185 = erectum.
clandestinum, *Spreng. Nov. Prov.* 29.—Bras.
commutatum, Opiz, Seznam, 68 = amphibium.
coronopifolium, *DC. Syst.* ii. 194.—Afr. bor.
crispum, Medic. Pflanzengatt. 81 (*cryspum*) = Lepidium sativum.
cryptanthum, *A. Rich. Tent. Fl. Abyss.* i. 15.—Abyssin.
curvisiliqua, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 73.—Am. bor. occ.
densiflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 226.—As. centr.
dentaculatum, Walp. Rep. ii. 755, sphalm. = *tentaculatum*.
deserticola, *Phil. in Anal. Univ. Chil.* (1865) II. 339.—Chili.
diandrum, Moench, Meth. 270 = Lepidium Iberis.
diffusum, DC. Prod. i. 139 = indicum.
divaricatum, *Wall. Cat.* n. 4776.—Ind. or.
diversifolium, Wall. l. c. n. 4775 = palustre.
Draba, Crantz, Crucif. 91 = Lepidium Draba.
ellipticum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 59 (= *Chrysochamela elliptica*).—As. Min.
erectum, *Trevir. ex Sweet, Hort. Brit.* ed. III. 26.—Chili.
erythrospermum, Boiss. & Huet, Diagn. Sér. II. v. 19, in obs. = sylvestre.
flaccidum, *Barn. in C. Gay, Fl. Chil.* i. 119.—Chili.
flexicaule, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 58 = Cochlearia velutina.
flexuosum, Seub. Fl. Azor. 43 = officinale?
fluviatile, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 21.—Afr. austr.
fontanum, Aschers. Fl. Brandenb. 32 = officinale.
gangeticum, Buch.-Ham. in Wall. Cat. n. 4777 = indicum.
glaucophyllum, DC. Syst. ii. 195 = Cochlearia glaucophylla.
globosum, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 35 = Cochlearia globosa.
Halleri, Andrz. ex DC. Syst. ii. 233 = Arabis Halleri.
heterophyllum, Blume, Bijdr. 50 = indicum.
heterophyllum, D. Don, Prod. Fl. Nep. 202 = palustre.
hispanicum, *Boiss. & Reut. Diagn.* Ser. I. viii. 18.—Hispan.
hispidum, DC. Syst. ii. 201 = palustre.
humifusum, *Guill. & Perr. Fl. Seneg. Tent.* i. 19.—Afr. trop.

NASTURTIUM:—
hybridum, Nym. Consp. 35, in syn. = austriacum.
Iberis, Gaertn. Mey. & Scherb. Fl. Wett. ii. 432 = Lepidium Iberis.
Impatiens, *Cham. & Schlecht. in Linnaea*, v. (1830) 212.—Mexic.
indicum, *DC. Syst.* ii. 199.—Ind. or.; China; Malaya.
Kerneri, Menyh. ex Janka, in Termés. Füzetek, vii. (1883) 117, in syn. = brachycarpum.
lacustre, *A. Gray, Gen. Pl. N. Am.* i. 132.—Am. bor.
limosum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 74.—Am. bor.
linifolium, Nutt. in Journ. Acad. Philad. vii. (1834) 12 = Sisymbrium linifolium.
lippizense, *DC. Syst.* ii. 195.—Europ. austr. or.
lyratum, Coop. in Pacif. Rail. Rep. xii. 51 = palustre.
lyratum, Nutt. in Torr. & Gray, Fl. N. Am. i. 73 = curvisiliqua.
macrocarpum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 18.—Syria.
macrorhizum, Steud. in Flora, xxxix. (1856) 409 = flaccidum.
madagascariense, *DC. Syst.* ii. 192.—Madag.
madagascariense, Wight & Arn. Prod. 19 = indicum.
Menziesii, Spreng. Syst. ii. 883 = Sisymbrium canescens.
mexicanum, *Moç. Sesse & Cerv. ex DC. Syst.* ii. 193.—Mexic.
micranthum, *DC. Prod.* i. 137.—Ind. or.
micropetalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 41 = tanacetifolium.
microphyllum, Reichb. Ic. Fl. Germ. i. 15. t. 50. f. 4360 = officinale.
microspermum, *DC. Syst.* ii. 199.—China.
Mirbellii, Spreng. ex Steud. Nom. ed. II. ii. 185 = erectum.
montanum, *Wall. Cat.* n. 4778, partim.—Reg. Himal.; Java; China.
Morisonii, Tausch, in Flora, xxiii. (1840) 715 = sylvestre.
mucronulatum, Lacroix, in Bull. Soc. Bot. Fr. vi. (1859) 551 = sylvestre.
multifidum, Spreng. Syst. ii. 883 = Sisymbrium canescens.
Munbyanum, *Boiss. & Reut. Diagn.* Ser. II. v. 19.—Alger.
myriophyllum, Spreng. Syst. ii. 883 = Sisymbrium myriophyllum.
nanum, *Wedd. in Ann. Sc. Nat.* Sér. IV. v. (1864) 290.—Bolivia.
natans, *DC. Syst.* ii. 198.—Sibir.
natans, Hook. Fl. Bor. Am. i. 39 = lacustre.
nebrodense, Rafin. in Desv. Journ. iv. (1814) 270 = Lepidium nebrodense.
niloticum, *Boiss. Diagn.* Ser. I. viii. 19.—Nubia.
Noëanum, Boiss. l. c. Ser. II. i. 21 (= *Chrysochamela Noëana*).—Armenia.
obliquum, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nêerl. Indie*, ii. (1845) 580.—Java.
obtusum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 74.—Am. bor.
obtusum, Torr. in Pacif. Rail. Rep. iv. 65 = curvisiliqua.
officinale, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 111.—Reg. bor. temp.
Orizabae, *Cham. & Schlecht. in Linnaea*, v. (1830) 212.—Mexic.
ovirense, Andrz. ex DC. Syst. ii. 234 = Arabis Halleri.
palustre, *DC. Syst.* ii. 191.—Reg. bor. et austr.
paniculatum, Crantz, Stirp. Austr. ed. I. fasc. i. 15; ed. II. 15 = Neslia paniculata.
petraeum, Crantz, Crucif. 80. t. 2 = Hutchinsia petraea.
perfoliatum, Bess. Enum. Pl. Volh. 26 = Lepidium perfoliatum.
plebejum, *Polak. in Linnaea*, xli. (1858) 546.—Costa Rica.
polymorphum, Nutt. in Torr. & Gray, Fl. N. Am. i. 74 = curvisiliqua.
portoricense, *Spreng. Syst.* ii. 882.—Ind. occ.
procurrens, Andrz. ex DC. Syst. ii. 229 = Arabis procurrens.
proliferum, Heuff. in Flora, xxxvi. (1853) 624 = brachycarpum.
pumilum, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 119.—Bras.
pumilum, Nutt. in Journ. Acad. Philad. vii. (1834) 12 = Sisymbrium linifolium.
pumilum, *Pomel, Nouv. Mat. Fl. Atl.* 376.—Alger.
pyrenaicum, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 110.—Europ. austr.

NASTURTIUM:—
pyrenaicum, Reut. ex Boiss. Diagn. Ser. I. viii. 19 = hispanicum.
radicans, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 247.—Chili.
Reichenbachii, Knaf, ex Opiz, Seznam, 68 = sylvestre.
riparium, Gremli, Excursions-Fl. Schweiz, ed. I, in nota, 80 = sylvestre.
riparium, Wallr. Sched. Crit. 373 = amphibium.
rivulare, Reichb. Ic. Fl. Germ. vi. (1827) 14. t. 517 = sylvestre.
rotundifolium, Rafin. Fl. Ludov. 86 (Quid?).—Louisian.
ruderale, Scop. Fl. Carn. ed. II. ii. 13 = Lepidium ruderale.
sagittatum, Auch. ex Fourn. Thèse Crucif. 124 = Sisymbrium pumilum.
sagittatum, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 111 = Alyssopsis deflexa.
sativum, Moench, Meth. 270 = Lepidium sativum.
saxatile, Crantz, Stirp. Austr. ed. I. Fasc. i. 14; ed. II. 14 = Cochlearia saxatilis.
scapigerum, Auch. ex Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 166 = Arabis sulphurea.
semipinnatifidum, Hook. Journ. Bot. i. (1834) 246 = palustre.
sessiliflorum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 73.—Am. bor. occ.
siifolium, Reichb. Fl. Germ. Excurs. 683 = officinale.
sikokianum, *Franch. & Sav. Enum. Pl. Jap.* ii. 277.—Japon.
sinense, *DC. Syst.* ii. 699.—China.
sinuatum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 73.—Am. bor. occ.
sparsum, D. Don, Prod. Fl. Nep. 202 = micranthum.
sphaerocarpum, *A. Gray, in Mem. Am. Ac.* N. S. iv. (1849) 6.—N. Mexic.
stenocarpum, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 21 = anceps.
sublyratum, Franch. & Sav. Enum. Pl. Jap. ii. 281 = montanum.
supinum, Beurl. ex Nym. Consp. 44 = Sisymbrium supinum.
sylvestre, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 110.—Reg. bor. temp.
tanacetifolium, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 190.—Am. bor.
tentaculatum, Wallr. in Linnaea, xiv. (1840) 607 = amphibium.
tenue, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 71.—Japon.
terrestre, R. Br. in Ait. Hort. Kew. ed. II. iv. 110 = palustre.
terrestre, Tausch, in Flora, xxiii. (1840) 711 = anceps?
Thaliana, Andrz. ex DC. Syst. ii. 226 = Sisymbrium Thalianum.
thracicum, *Griseb. Spicil. Fl. Rumel.* i. 258.—Banat.; Graecia.
trachycarpum, *A. Gray, in Bull. Geol. Geogr. Surv.* ii. 233.—Am. bor. occ.
tuberosum, Rafin. Fl. Ludov. 84 = Cardamine rhomboidea?
Turczaninowii, Czern. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 372, in textu = anceps?
turfosum, *Kunze, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 247.—Chili.
verrucarium, Garsault, Desc. Pl. t. 402; ex DC. Syst. ii. 525 = Senebiera Coronopus.
Walteri, Wood, Class-Book (1861), 228 = tanacetifolium.
Wulfenianum, Host, Fl. Austr. ii. 259 = lippizense.

NASTUS, Juss. Gen. 34 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1210.
STEMMATOSPERMUM, Beauv. Agrost. 144. t. 25 (1812).
arundinaceus, Sm. in Rees, Cycl. xxiv. n. 1 = Bambusa arundinacea.
baccifer, Roxb. ex Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 442 = Melocanna bambusoides.
barbatus, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 132.—Bras.
borbonicus, *J. F. Gmel. Syst.* ii. 580.—Ins. Borbon.
bruneus, *Desv. Opusc.* 107 (err. typ. 211).—Bras.
capitatus, Kunth, Rév. Gram. i. 325 (= *Beesha capitata*).—Ins. Borbon.
Chusque, H. B. & K. Nov. Gen. et Sp. i. 201 = Chusquea scandens.
Chusquea, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 442 = praec.
Guadua, Spreng. Syst. ii. 113 = Guadua angustifolia.
humilus, Hassk. Cat. Hort. Bog. Alt. 19 = Dinochloa Tjankorreh.
latifolius, Spreng. Syst. ii. 113 = Guadua latifolia.
macrospermus, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 442, 458. t. 8. f. 1 = Arundinaria macrosperma.
paniculatus, Sm. in Rees, Cycl. xxiv. n. 4 = borbonicus.
plicatilis, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1830) 178, nomen (Quid?).—N. Guin.
prolifer, *Desv. Opusc.* 107 (err. typ. 211).—Bras.
Quila, Schult. f. Syst. vii. 1361 = Chusquea Quila.
strictus, Sm. in Rees, Cycl. xxiv. n. 2 = Dendrocalamus strictus.
tessellatus, Nees, Fl. Afr. Austr. i. 463 = Arundinaria tessellata.
Thouarsii, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 442; Spreng. Syst. ii. 113 = Bambusa vulgaris.
Tjankorreh, Schult. f. Syst. vii. 1358 = Dinochloa Tjankorreh.
verticillatus, Sm. in Rees, Cycl. xxiv. n. 3 = Gigantochloa verticillata.
viviparus, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 458. t. 8. f. 3 = Bambusa vulgaris.

NATALANTHE, Sond. in Linnaea, xxiii. (1850) 52 = **Tricalysia**, A. Rich. (Rubiac.).
floribunda, Sond. l. c. (= *Bunburya capensis*).—Afr. austr.

NATALIA, Hochst. in Flora, xxiv. (1841) II. 663 = **Bersama**, Fresen. (Sapind.).
lucens, Hochst. l. c. 664.—Afr. austr.
paullinioides, Planch. in Hook. Ic. Pl. t. 780 = B. paullinioides.

NATHUSIA, Hochst. in Flora, xxiv. (1841) I. Intell. 25; II. 671 = **Schrebera**, Roxb. (Oleac.).
alata, Hochst. ll. cc. 25; 672 = S. alata.

NATRIX, Moench, Meth. 158 (1794) = **Ononis**, Linn. (Legumin.).
crispa, Moench, Meth. Suppl. 54 = O. crispa.
fruticosa, Moench, Meth. 158 = O. fruticosa.
pinguis, Moench, l. c. = O. Natrix.
rotundifolia, Moench, l. c. = O. rotundifolia.

NATSIATOPSIS, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 201. *OLACINEAE*.
thunbergiaefolia, *Kurz, l. c.*—Burma.

NATSIATUM, Buch.-Ham. in Wall. Cat. n. 4252 (1831). *OLACINEAE*, Benth. & Hook. f. i. 355.
NANSIATUM, Miq. Fl. Ind. Bat. i. 1. 797 (1855).
gamosepalum, Griff. Notul. iv. 330 = Lophophyllum bicristatum.
herpeticum, *Buch.-Ham. in Wall. Cat.* n. 4252.—Reg. Himal.; Burma.
oppositifolium, Planch. in Hook. Lond. Journ. Bot. v. (1846) 247 = Iodes ovalis.

NAUCHEA, Descourt. in Mém. Soc. Linn. Par. iv. (1826) 7 = **Clitoria**, Linn. (Legumin.).
bracteata, Dupuis, ex Descourt. l. c. 11 = C. Ternatea.
brasiliana, Descourt. l. c. 9 = Centrosema brasilianum.
clypeata, Pritz. Ic. Ind. i. 738 = C. glycinoides.
falcata, Descourt. in Mém. Soc. Linn. Par. iv. (1826) 9 = Periandra coccinea.
heterophylla, Descourt. l. c. 8 = C. heterophylla.
laurifolia, Ledru, ex Descourt. l. c. 11 = Centrosema vexillatum.
Mariana, Descourt. l. c. 9 = C. Mariana.
multiflora, Descourt. l. c. 10 = Vilmorinia multiflora.
Plumieri, Pers. ex Descourt. l. c. 12 = Centrosema Plumieri.
polyphylla, Ledru, ex Descourt. l. c. 10 = Barbieria polyphylla.
pudica, Descourt. l. c. 12 = C. pudica.
rubiginosa, Descourt. l. c. = C. glycinoides.
Ternatea, Descourt. l. c. 8 = C. Ternatea.
virginiana, Descourt. l. c. 9 = Centrosema virginianum.

NAUCLEA, Linn. Sp. Pl. ed. II. 243 (1762). *RUBIACEAE*, Benth. & Hook. f. ii. 31.
PLATANOCARPUM, Korth. Obs. Naucl. 18 (1839), partim.
acida, Hunter, in Trans. Linn. Soc. ix. (1808) 223 = Uncaria acida.
aculeata, H. B. & K. Nov. Gen. et Sp. iii. 382 = Uncaria tomentosa.
aculeata, Willd. Sp. Pl. i. 929 = Uncaria guianensis.
Adina, Blanco, Fl. Filip. ed. II. 102 (= *Stephegyne diversifolia*).—Ins. Philipp.
Adina, Sm. in Rees, Cycl. xxiv. n. 13 = Adina globiflora.
adinoides, Lindl. in Bot. Reg. sub t. 895 = Adina globiflora.
affinis, *Miq. Fl. Ind. Bat.* ii. 137.—Ins. Banca.
africana, Willd. Sp. Pl. i. 929 = Mitragyna africana.
appendiculata, Walp. Rep. ii. 943 = Uncaria appendiculata.
aralioides, Miq. Fl. Ind. Bat. ii. 344, in textu = Adina polycephala.
attenuata, Walp. Rep. ii. 513 = Uncaria attenuata.
Bartlingii, *DC. Prod.* iv. 344.—Ins. Philipp.
Blancoi, *Vidal, Phan. Cuming. Philipp.* 175.—Ins. Philipp.
bracteosa, Welw. Syn. Explicat. 48. n. 130 = Mitragyna macrophylla.
Brunonis, Wall. Cat. n. 6097 (= *Stephegyne diversifolia*).—Ind. or.
Cadamba, Roxb. Hort. Beng. 14; Fl. Ind. i. 512 = Anthocephalus morindaefolius.
callophylla, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 182 = purpurascens.
calycina, *Bartl. ex DC. Prod.* iv. 346.—Ins. Philipp.
canescens, *Bartl. l. c.*—Ins. Philipp.
canescens, Walp. Rep. ii. 513 = Uncaria canescens.
capitellata, Voigt, Hort. Suburb. Calc. 375 = Adina polycephala.
Cinchonae, *DC. Prod.* iv. 345.—Am. austr.
citrifolia, Poir. in Lam. Encyc. iv. 435 = Anthocephalus morindaefolius.
coadunata, Roxb. ex Sm. in Rees, Cycl. xxiv. n. 6 = Sarcocephalus cordatus.
cordata, *Blume, Bijdr.* 1011.—Java.
cordata, Roxb. Hort. Beng. 14; Fl. Ind. i. 509 = Sarcocephalus cordatus.
cordifolia, Roxb. Pl. Corom. i. 40. t. 53 = Adina cordifolia.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 176.—Ins. Philipp.
cyclophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 181.—Ins. Molucc.
cyrtopoda, *Miq. Fl. Ind. Bat.* ii. 342.—Sumatra.
dasyoneura, Walp. Rep. ii. 513 = Uncaria dasyoneura.
digitata, Blanco, Fl. Filip. ed. II. 102 = Heptapleurum Cephalotes.
diversifolia, Wall. Cat. n. 609 (= *Stephegyne diversifolia*).—Ind. or.
elliptica, Dalz. & Gibs. Bomb. Fl. 118 = missionis.
elliptica, Walp. Rep. ii. 512 = Uncaria dasyoneura.
excelsa, *Blume, Bijdr.* 1009.—Java.
fagifolia, *Teijsm. & Binn. Cat. Hort. Bog.* 117.—Ins. Amboina.
ferrea, Blume, Bijdr. 1014 = Uncaria ferrea.
ferruginea, Blume, l. c. 1013 = Uncaria pedicellata.
Forsteri, *Seem. Fl. Vit.* 121, in obs.—Polynes.
Gambir, Hunter, in Trans. Linn. Soc. ix. (1808) 218. t. 22 = Uncaria Gambier.
glaberrima, *Bartl. ex DC. Prod.* iv. 344.—Ins. Philipp.
glaberrima, Blanco, Fl. Filip. ed. II. 100 = Sarcocephalus subditus.
glabra, Blanco, Fl. Filip. ed. II. 101 = Anthocephalus morindaefolius.
glabra, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 512.—Ins. Molucc.
glabrata, Blume, Bijdr. 1012 = Uncaria glabrata.
glandulifera, *Span. in Linnaea*, xv. (1841) 315.—Ins. Timor.
glandulosa, Blanco, Fl. Filip. ed. I. 143 = Anthocephalus morindaefolius.
gracilis, *Vidal, Phan. Cuming. Philipp.* 176.—Ins. Philipp.
grandifolia, DC. Prod. iv. 345 = Sarcocephalus cordatus.
grandifolia, Spreng. Syst. iv. Cur. Post. 81 = Uncaria macrophylla.
guianensis, Poir. in Lam. Encyc. iv. 436; D. Dietr. Syn. Pl. i. 792 = Uncaria guianensis.
Haenkeana, *Steud. Nom.* ed. II. ii. 186.—Ins. Philipp.
Hallii, Walp. Rep. ii. 512 = Uncaria pedicellata.
imbricata, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 182 = purpurascens.
inermis, Baill. in Bull. Soc. Linn. Par. i. (1879) 201 = Mitragyna africana.

NAUCLEA:—
insignis, D. Dietr. Syn. Pl. i. 791 = Uncaria sclerophylla.
Korthalsii, Steud. Nom. ed. II. ii. 186 = Mitragyna speciosa.
laevigata, Walp. Rep. ii. 512 = Uncaria laevigata.
lanceolata, Blume, Bijdr. 1009 = purpurea.
lanosa, Poir. Encyc. Suppl. iv. 64 = Uncaria lanosa.
latifolia, Blanco, Fl. Filip. ed. I. 144 = obtusa.
latifolia, Sm. in Rees, Cycl. xxiv. n. 5 = Sarcocephalus esculentus.
longiflora, Poir. Encyc. Suppl. iv. 63 = Uncaria acida.
lutea, Blanco, Fl. Filip. ed. I. 141 = Sarcocephalus subditus.
luzonensis, D. Dietr. Syn. Pl. i. 791 = Haenkeana.
luzoniensis, Blanco, Fl. Filip. ed. II. 102 = Mitragyna speciosa.
macrophylla, Blume, Bijdr. 1010 = Sarcocephalus cordatus.
macrophylla, Perr. & Lepr. ex DC. Prod. iv. 346 = Mitragyna macrophylla.
macrophylla, *Roxb. Hort. Beng.* 14; *Fl. Ind.* i. 511.—Ins. Amboin.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 27.—Malaya.
microcephala, Delile, in Caill. Pl. Afr. 67. n. 54 = Adina microcephala.
microcephala, Wall. ex Voigt, Hort. Suburb. Calc. 375; et ex Hook. f. Fl. Brit. Ind. iii. 25 = Adina polycephala.
missionis, *Wall. Cat.* n. 6099.—Ind. or.
mollis, Bartl. ex DC. Prod. iv. 344 = Bartlingii.
mollis, *Blume, Bijdr.* 1010.—Java.
moluccana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 183.—Ins. Molucc.
morindaefolia, Blume, Bijdr. 1011 = Anthocephalus morindaefolius.
nemorosa, Walp. Rep. ii. 512 = Uncaria nemorosa.
oblongata, Miq. ex Hook. f. Fl. Brit. Ind. iii. 27 = missionis.
obtusa, *Blume, Bijdr.* 1009.—Java.
orientalis, Forst. f. Prod. 15 = Forsteri.
orientalis, Gaertn. Fruct. i. 151. t. 30 = Mitragyna parvifolia.
orientalis, Herb. Madr. ex Wight & Arn. Prod. 392 = missionis.
orientalis, *Linn. Sp. Pl.* ed. II. 243.—Ind. or.
orientalis, *Wilid. Sp. Pl.* i. 928.—Ind. or.; Java.
ovalifolia, Roxb. Hort. Beng. 14; Fl. Ind. i. 516 = Adina sessilifolia?
ovalifolia, Spreng. Syst. iv. Cur. Post. 80 = Uncaria ovalifolia.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* 538.—Sumatra.
pallida, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 182 = purpurascens.
parviflora, Pers. Syn. i. 202 = Mitragyna parvifolia.
parvifolia, Roxb. Pl. Corom. i. 40. t. 52 = Mitragyna parvifolia.
parvifolia, Wall. Cat. n. 6093 B = Sarcocephalus cordatus.
parvifolia, Willd. Sp. Pl. i. 929 = Mitagyna parvifolia.
pedicellata, Blume, Bijdr. 1012 = Uncaria pedicellata.
peduncularis, Thw. Enum. Pl. Zeyl. 137 = zeylanica.
peduncularis, *Wall. Cat.* n. 6091.—Malaya.
pilosa, Blume, Bijdr. 1013 = Uncaria pilosa.
pilulifera, Baill. ex Franch. Pl. Davidian. i. 154 = Adina globiflora.
platanocarpa, Hook. f. in Hook. Ic. Pl. t. 787 = Mitragyna africana.
polycephala, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 289 = Cinchonae.
polycephala, Wall. Cat. n. 6100 = Adina polycephala.
purpurascens, *Korth. Verh. Nat. Gesch. Bot.* 157.—Malaya.
purpurea, Herb. Reinw. ex Miq. Fl. Ind. Bat. ii. 343 = glabra.
purpurea, *Roxb. Pl. Corom.* i. 41. t. 54.—Ind. or.; China.
purpurea, Wall. Cat. n. 6090 D = undulata.
racemosa, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 178 = Adina racemosa.
rhynchophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 108.—Japon.
rotundifolia, Bartl. ex DC. Prod. iv. 346 = Haenkeana.
rotundifolia, Roxb. Hort. Beng. 86; Fl. Ind. i. 516 (= *Stephegyne diversifolia*).—Ind. or.
Roxburghiana, Walp. Rep. ii. 513 = Uncaria Roxburghiana.

NAUCLEA:—
Roxburghii, G. Don, Gen. Syst. iii. 469 = Sarcocephalus cordatus.
sambucina, T. Winterb. Account Sierra Leone, ii. 45 = Sarcocephalus esculentus.
scandens, Roxb. ex Hook. f. Fl. Brit. Ind. iii. 30 = Uncaria sessilifructus.
scandens, Sm. in Rees, Cycl. xxiv. n. 9 = Uncaria pilosa.
sclerophylla, Hunter, in Trans. Linn. Soc. ix. (1808) 223 = Uncaria sclerophylla.
sericea, *Span. in Linnaea*, xv. (1841) 314.—Ins. Timor.
sericea, Wall. Cat. n. 6095 = Adina sessilifolia.
sessilifolia, Roxb. Hort. Beng. 86; Fl. Ind. i. 515 = Adina sessilifolia.
sessilifructus, D. Dietr. Syn. Pl. i. 792 = Uncaria sessilifructus.
sessilis, Spreng. Syst. iv. Cur. Post. 81 = Uncaria sessilifructus.
setigera, Blume, Bijdr. 1013 = Uncaria lanosa.
setiloba, Walp. Rep. ii. 943 = Uncaria setiloba.
silhetana, D. Dietr. Syn. Pl. i. 791 = Uncaria macrophylla.
speciosa, Miq. Fl. Ind. Bat. ii. 140 = Mitragyna speciosa.
speciosa, Walp. Rep. ii. 512 = Uncaria sclerophylla.
stellata, *Wall. Cat.* n. 6102.—Ind. or.
sterculiaefolia, Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 289 = Adina cordifolia.
stipulacea, G. Don, Gen. Syst. iii. 469 = Mitragyna macrophylla.
stipulacea, Roxb. ex Hook. f. Fl. Brit. Ind. iii. 22 = Sarcocephalus cordatus.
stipulata, Benth. & Hook. f. Gen. ii. 31 = seq.
stipulosa, DC. Prod. iv. 346 = Mitragyna macrophylla.
strigosa, *Korth. Verh. Nat. Gesch. Bot.* 157.—Malaya.
subdita, Steud. Nom. ed. II. ii. 186 = Sarcocephalus subditus.
surinamensis, *Miq. in Linnaea*, xix. (1847) 129.—Guian.
synkorynes, *Korth. Verh. Nat. Gesch. Bot.* 158.—Borneo.
tetrandra, Roxb. Hort. Beng. 14; Fl. Ind. i. 516 = Cephalanthus naucleoides.
tomentosa, Willd. in Roem. & Schult. Syst. v. 221 = Uncaria tomentosa.
trichotoma, Zoll. & Mor. Syst. Verz. Zoll. 61 = Adina polycephala.
tubulosa, Arn. ex Thw. Enum. Pl. Zeyl. 137 = (*Stephegyne tubulosa*).—Ind. or.
Uncaria, D. Dietr. Syn. Pl. i. 791 = Uncaria ovalifolia.
undulata, *Roxb. Hort. Beng.* 14; *Fl. Ind.* i. 508.—Ins. Molucc.
undulata, Wall. Cat. n. 6094 = Sarcocephalus cordatus.
verticillata, *Baill. Adansonia*, xii. (1879) 284.—Madag.
vestita, Zipp. ex Span. in Linnaea, xv. (1841) 314 = sericea.
Wallichiana, R. Br. in Wall. Cat. n. 6098 = Sarcocephalus cordatus.
Wallichiana, Spreng. Syst. iv. Cur. Post. 81 = Uncaria lanosa.
Wallichii, Walp. Rep. ii. 513 = Uncaria Wallichii.
zeylanica, *Hook. f. Fl. Brit. Ind.* iii. 26.—Zeylan.

NAUCLEOPSIS, Miq. in Mart. Fl. Bras. iv. 1. 120. t. 35 (1853) = **Noyera**, Tréc. (Urtic.).
macrophylla, Miq. l. c.—Am. trop.

NAUCOREPHES, Rafin. Fl. Tellur. ii. 34 (1836) = **Coccoloba**, Linn. (Polygon.).
pubescens, Rafin. l. c. = C. coronata.
punctata, Rafin. l. c. = C. coronata.

NAUDINIA, Decne. ex Triana, in Seem. Fl. Vit. 86 (1866) = **Astronia**, Blume (Melastom.).
Forsteri, Decne. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 152 = A. Forsteri.
glabra, Decne. ex Seem. Fl. Vit. 86 = A. pileolata.
ovalifolia, Decne. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 152 = A. ovalifolia.
pileolata, Decne. ex Triana, l. c. = A. pileolata.
Umbella, Decne. ex Triana, l. c. = A. Forsteri.

NAUDINIA, Planch. & Linden, in Ann. Sc. Nat. Sér. III. xix. (1853) 79. *RUTACEAE*, Benth. & Hook. f. i. 285.
amabilis, *Planch. & Linden, l. c.*—N. Granat.

NAUDINIA, A. Rich. Ess. Fl. Cub. 561 (1845) = **Tetrazygia**, Rich. (Melastom.).
argyrophylla, A. Rich. l. c. 562 (= *Miconia bicolor*).—Cuba.
chrysophylla, A. Rich. l. c. 563 (= *Miconia pallens*).—Cuba.

NAUENBURGIA, Willd. Sp. Pl. iii. 2393 (1803) = **Flaveria**, Juss. (Compos.).
trinervata, Willd. l. c. = F. repanda.

NAUENIA, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 193 = **Lacaena**, Lindl. (Orchid.).
spectabilis, Klotzsch, l. c. = L. spectabilis.

NAUMBURGIA, Moench, Meth. Suppl. 23 (1802) = **Lysimachia**, Tourn. (Primul.).
guttata, Moench, l. c. = L. thyrsiflora.
thyrsiflora, Reichb. Fl. Germ. Excurs. 410 = L. thyrsiflora.

NAUPLIUS, Cass. in Dict. Sc. Nat. xxiii. 566 (1822) = **Odontospermum**, Neck. (Compos.).
aquaticus, Cass. l. c. xxxiv. 273 = O. aquaticum.
intermedius, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 236 = O. intermedium.
maritimus, Cass. in Dict. Sc. Nat. xxxiv. 274 = O. maritimum.
sericeus, Cass. ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 234 = O. sericeum.
stenophyllus, Webb, l. c. 237 = O. stenophyllum.

NAUTEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Tectona**, Linn. f. (Verbenac.).

NAUTICALYX, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1404 (1855) = seq.

NAUTILOCALYX, Linden, Cat. [1851, p. 12] ex Hanst. in Linnaea, xxvi. (1853) 207 = **Episcia**, Mart. (Gesnerac.).
hastatus, Linden, ex Hanst. l. c. = E. bractescens.
panamensis, Seem. Bot. Voy. Herald, 250 = Achimenes panamensis.

NAUTONIA, Decne. in DC. Prod. viii. 509 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 747.
nummularia, *Decne. l. c.* 510.—Bras.

NAVAEA, Webb & Berth. Phyt. Canar. i. 32. t. 1 C (1836) = **Lavatera**, Linn. (Malvac.).
phoenicea, Webb & Berth. l. c. 34 = L. phoenicea.

NAVARETTIA, Hedw. f. Gen. 107 (1806) = seq.

NAVARRETIA, Ruiz & Pav. Prod. 20 (1794) = **Gilia**, Ruiz & Pav. (Polem.).
atractyloides, Hook. & Arn. Bot. Beech. Voy. 368 = G. atractyloides.
cotulaefolia, Hook. & Arn. l. c. = G. cotulaefolia.
heterophylla, Benth. in DC. Prod. ix. 309 = Collomia heterophylla.
intertexta, Hook. Fl. Bor. Am. ii. 75 = G. intertexta.
involucrata, Ruiz & Pav. Fl. Per. ii. 8 = G. Navarettia.
leucocephala, Benth. Pl. Hartw. 324 = G. leucocephala.
minima, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 160 = G. minima.
pubescens, Hook. & Arn. Bot. Beech. Voy. 368 = G. cotulaefolia.
pungens, Hook. Fl. Bor. Am. ii. 75 = G. squarrosa.
pusilla, Hort. ex Steud. Nom. ed. II. i. 683, in syn.; ii. 186 = G. Navarettia.
Schottii, Torr. in Bot. Mex. Bound. 145 = G. setosissima.
setosissima, Torr. & Gray, ex Torr. in Ives, Rep. Colorado River Bot. 22 = G. setosissima.
squarrosa, Hook. & Arn. Bot. Beech. Voy. 368 = G. squarrosa.
viscidula, Benth. Pl. Hartw. 324 = G. viscidula.

NAVENIA, Klotzsch, ex Benth. & Hook. f. Gen. iii. 547 (1883) = NAUENIA, Klotzsch = **Lacaena**, Lindl. (Orchid.).

NAVIA, Schult. f. Syst. vii. 65, 1195 (1830) = **Dyckia**, Schult. (Bromel.).
acaulis, Mart. ex Schult. f. l. c. 1196.—Bras.
brevifolia, Griseb. in Goett. Abh. xxiv. (1879) 332.—Reg. Argent.
caulescens, Mart. ex Schult. f. Syst. vii. 1195.—Bras.

NAVICULARIA, Heist. ex Adans. Fam. ii. 188 (1763) = **Sideritis**, Tourn. (Labiat.).
Heisteri, Fabr. ex Benth. in DC. Prod. xii. 440 = S. syriaca.

NAVICULARIA, Raddi, Agrost. Bras. 38 (1823) = **Ichnanthus**, Beauv. (Gramin.).
glabra, Raddi, l. c. 39 = I. Martianus, I. Cyanotis.
hirta, Raddi, l. c. = I. Planotis.
lanata, Raddi, l. c. 40. t. 1. f. 5 = I. leiocarpus ?

NAVIDURA, Alef. in Bonplandia, ix. (1861) 147 = **Lathyrus**, Tourn. (Legumin.).
tingitana, Alef. l. c. = L. tingitanus.

NAYAS, Neck. Elem. iii. 279 (1790) = **Najas**, Linn.

NAZIA, Adans. Fam. ii. 31 (1763) = **Tragus**, Hall. (Gramin.).

NEACTELIS, Rafin. New Fl. Am. iv. 69 (1836). *COMPOSITAE* (Quid ?).
strigosa, Rafin. l. c.—Am. bor.

NEAEA, Juss. in Ann. Mus. Par. ii. (1803) 275 = **Neea**, Ruiz & Pav. (Nyctag.).

NEAERA, Salisb. Gen. Fl. Fragm. 109 (1866) = **Stenomesson**, Herb. (Amaryll.).
crocea, Salisb. l. c. = S. croceum.

NEARA, Soland. ex Seem. Fl. Vit. 118 (1865) = **Meryta**, Forst. (Araliac.).
longifolia, Soland. l. c. = M. lanceolata.

NEBELIA, Neck. Elem. i. 113 (1790). *BRUNIACEAE*.
BERARDIA, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 380. t. 37.
DIBERARA, Baill. in Bull. Soc. Linn. Par. i. (1881) 279.
affinis, *Sweet, Hort. Brit.* ed. II. 116.—Afr. austr.
paleacea, *Sweet, l. c.*—Afr. austr.
phylicoides, Sweet, l. c. = Raspalia phylicoides.

NEBRA, Noronha, ex Choisy, in DC. Prod. xiii. II. 447 (1849) = **Neea**, Ruiz & Pav. (Nyctag.).
rudifolia, Noronha, ex Choisy, l. c. 448 = Neea floribunda.

NEBROPSIS, Rafin. Alsog. Am. 68 (1838) = **Aesculus**, Linn. (Sapind.).
alba, Rafin. l. c. 69 = Ae. alba.
glabra, Rafin. l. c. = Ae. glabra.
muricata, Rafin. l. c. 68 = Ae. muricata.
neglecta, Rafin. l. c. 70 = Ae. flava.
ochroleuca, Rafin. l. c. 68 = Ae. ochroleuca.
pallida, Rafin. l. c. 69 = Ae. glabra.
verrucosa, Rafin. l. c. = Ae. verrucosa.

NECALISTIS, Rafin. Sylva Tellur. 58 (1838) = **Ficus**, Tourn. (Urtic.).
aspera, Rafin. l. c. = F. hirta.
turbinata, Rafin. l. c. = F. turbinata.

NECHAMANDRA, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 78 = **Lagarosiphon**, Harv. (Hydroch.).
alternifolia, Thw. Enum. Pl. Zeyl. 332 = L. Roxburghii.
Roxburghii, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 78 = L. Roxburghii.

NECKERIA, J. F. Gmel. Syst. 3, 16 (1791) = **Pollichia**, Soland. (Illecebr.).
campestris, J. F. Gmel. l. c. = P. campestris.

NECKERIA, Scop. Introd. 313 (1777) = **Corydalis**, Vent. (Papav.).
capnoides, Neck. Elem. iii. 60 = C. lutea.
lutea, Neck. l. c. = C. lutea.
sempervirens, Neck. l. c. = C. glauca.

NECKIA, Korth. in Nederl. Kruidk. Arch. i. (1848) 358. *VIOLARIEAE*, Benth. & Hook. f. i. 120.
humilis, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 158.—Borneo.
lancifolia, *Hook. f. l. c.*—Borneo.
serrata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 358.—Sumatra.

NECTALISMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 156 = **Elisma**, Buchen. (Alismac.).
natans, Fourr. l. c. = E. natans.

NECTANDRA, Berg. Desc. Pl. Cap. 131 (1767) = **Gnidia**, Linn. (Thymel.).
laevigata, Berg. l. c. 134 = G. oppositifolia.
sericea, Berg. l. c. 131 = G. sericea.
tetrandra, Berg. l. c. 133 = Struthiola erecta.

NECTANDRA, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 279. *LAURINEAE*, Benth. & Hook. f. i. iii. 159.
DAMBURNEYA, Rafin. Sylva Tellur. 136 (1838).
POMATIUM, Nees, ex Meissn. in Mart. Fl. Bras. v. II. 252 (1866).
POROSTEMA, Schreb. Gen. 517 (1791), partim.
abortiens, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 281 = Ocotea guianensis.
acutangula, *Miq. in Linnaea*, xxii. (1849) 806.—Bras.
acutissima, Nees, ex Meissn. in DC. Prod. xv. I. 147 = rigida.
amara, Meissn. l. c. 158 = puberula.
amara, Nees, ex Meissn. l. c. 156 = cymbarum.
amazonum, *Nees, Syst. Laurin.* 282.—Bras.
ambigua, *Meissn. in DC. Prod.* xv. I. 158.—Guian.
angustifolia, Miq. ex Meissn. l. c. = puberula.
angustifolia, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 48.—Bras.
anonyma, Steud. Nom. ed. II. ii. 187 = Willdenoviana.
antillana, *Meissn. in DC. Prod.* xv. I. 153.—Ind. occ.
Arnottiana, *Nees, Syst. Laurin.* 289.—Peruv.
aurantiodora, *Benth. & Hook. f. Gen.* iii. 158.—Bras.
barcellensis, *Meissn. in DC. Prod.* xv. I. 155.—Bras.
berchemiaefolia, *Meissn. l. c.* 154.—N. Granat.
bijuga, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 281 = Ocotea guianensis.
Boniato, *A. Rich. Fl. Cub. Fanerog.* ii. 188.—Cuba.
Bredemeyeriana, Nees, in Linnaea, xxi. (1848) 505 = turbacensis, Willdenoviana.
canescens, Meissn. in DC. Prod. xv. I. 147 = amazonum.
canescens, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 47.—Bras.
caudata, Nees & Mart. ex Meissn. in DC. Prod. xv. I. 85 = Acrodiclidium guianense.
chrysothyrsus, Benth. & Hook. f. Gen. iii. 159 = Arnottiana.
Cigua, A. Rich. Fl. Cub. Fanerog. ii. 187 = Willdenoviana.
cinnamomoides, *Nees, Syst. Laurin.* 307.—N. Granat.
cissiflora, *Nees, l. c.* 296.—Peruv.
concinna, *Nees, l. c.* 322.—Am. trop.
coriacea, Griseb. Fl. Brit. W. Ind. 281 = Willdenoviana.
coriacea, Meissn. in DC. Prod. xv. I. 163 = patens.
crassifolia, *Benth. Pl. Hartw.* 253.—Peruv.
crassipes, *Meissn. in DC. Prod.* xv. I. 150.—Peruv.
cuneata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 188 (Ocoteae sp.).—Cuba.
cuspidata, *Nees & Mart. ex Nees, Syst. Laurin.* 330.—Bras.
cymbarum, *Nees, l. c.* 305.—Venezuela.
discolor, *Nees, Syst. Laurin.* 286.—N. Granat.
effusa, Meissn. in DC. Prod. xv. I. 158 = concinna.
exaltata, *Griseb. Cat. Pl. Cub.* 111.—Cuba.
exaltata, Griseb. Fl. Brit. W. Ind. 281, partim = sanguinea.
filiformis, Rafin. Sylva Tellur. 134 (Quid ?).—Guian. ?
floribunda, *Nees, Syst. Laurin.* 339.—Jamaic.
furcata, *Nees, in Linnaea*, xxi. (1848) 501.—Peruv.
Gardneri, *Meissn. in DC. Prod.* xv. I. 155.—Bras.
glabrescens, *Benth. Bot. Voy. Sulph.* 161.—Mexic.
glauca, *Warm. in Kjoeb. Vidensk. Meddel.* (1870) 142.—Bras.
grandiflora, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 49.—Bras.
grandis, Klotzsch & Karst. ex Nees, l. c. xxi. (1848) 505 = villosa.
guianensis, *Meissn. in DC. Prod.* xv. I. 160.—Guian.

NECTANDRA :—
japurensis, *Nees & Mart. ex Nees, Syst. Laurin.* 335.—Bras.
lanceolata, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 47.—Bras.
Laurel, *Klotzsch, ex Nees, l. c.* xxi. (1848) 505.—N. Granat.
Laurel-hu, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 267 ; *Contrib.* 73.—Reg. Argent.
Laurel-y, *Parodi, ll. cc.* 268 ; 74.—Reg. Argent.
leucantha, Griseb. Fl. Brit. W. Ind. 282 = antillana.
leucantha, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 48.—Bras.
leucantha, Nees, l. c. xxi. (1848) 507, partim = ambigua.
leucothyrsus, *Meissn. in DC. Prod.* xv. I. 160.—Bras.
limbata, *Nees, in Linnaea*, xxi. (1848) 509.—Peruv.
limbosa, Nees, ex Meissn. in DC. Prod. xv. I. 156 = limbata.
longifolia, *Nees, in Linnaea*, xxi. (1848) 502.—Peruv.
longifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 188.—Cuba.
lucida, *Nees, in Linnaea*, viii. (1833) 47.—Bras.
macrophylla, *Nees, Syst. Laurin.* 292.—N. Granat.
magnoliaefolia, *Meissn. in DC. Prod.* xv. I. 154.—Bras.
magnoliaefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 188.—Cuba.
Matthewsii, *Meissn. in DC. Prod.* xv. I. 159.—Peruv.
membranacea, *Griseb. Fl. Brit. W. Ind.* 282.—Ind. occ.
microcarpa, *Meissn. in DC. Prod.* xv. I. 162.—Peruv.
mollis, Nees, Syst. Laurin. 287 = villosa.
Moritziana, *Klotzsch, ex Nees, in Linnaea*, xxi. (1848) 503.—N. Granat.
myriantha, *Meissn. in DC. Prod.* xv. I. 163.—Am. trop.
Neesiana, Miq. in Linnaea, xviii. (1844) 745 (= *Oreodaphne fallax*).—Guian.
nitidula, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 48.—Bras.
oblongifolia, Nees, ex Meissn. in DC. Prod. xv. I. 152 = grandiflora.
oppositifolia, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 47 = rigida.
Oreadum, Mart. ex Meissn. in DC. Prod. xv. I. 148 = lanceolata.
pallida, Miq. Stirp. Surin. Sel. 196 = vaga.
pallida, *Nees, in Linnaea*, xxi. (1848) 510.—Guian.
panurensis, *Meissn. in DC. Prod.* xv. I. 157.—Bras.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 266 ; *Contrib.* 72.—Paraguay.
patens, *Griseb. Fl. Brit. W. Ind.* 281.—Ind. occ.
paterifera, *Nees, Syst. Laurin.* 308.—Bras.
pendula, Nees, ex Meissn. in DC. Prod. xv. I. 137 = Ocotea Sieberi.
Pisi, *Miq. Stirp. Surin. Sel.* 199.—Guian.
polita, *Nees & Mart. ex Nees, Syst. Laurin.* 325.—Bras.
polyphylla, Nees, l. c. 332 = Acrodiclidium guianense.
porphyria, *Griseb. in Goett. Abh.* xix. (1874) 144.—Reg. Argent.
psammophila, *Nees & Mart. ex Nees, Syst. Laurin.* 303.—Bras.
puberula, *Nees, l. c.* 332.—Bras.
Puchury-major, *Nees & Mart. ex Nees, l. c.* 328.—Bras.
Puchury-minor, *Nees & Mart. ex Nees, l. c.* 336.—Bras.
pulverulenta, *Nees, l. c.* 283.—Bras.
punctata, *Klotzsch, ex Nees, in Linnaea*, xxi. (1848) 509.—Peruv.
pyrifolia, Nees, ex Meissn. in DC. Prod. xv. I. 166 = squarrosa.
racemifera, *Meissn. in Mart. Fl. Bras.* v. II. 310.—Bras.
rectinervia, *Meissn. in DC. Prod.* xv. I. 158.—Venezuela.
Regnellii, *Meissn. in Mart. Fl. Bras.* v. II. 310.—Am. trop.
rhynchophylla, *Meissn. in DC. Prod.* xv. I. 155.—Guian.
Riedelii, *Meissn. l. c.* 161.—Bras.
rigida, *Nees, Syst. Laurin.* 284.—N. Granat.
Rodiei, Schomb. ex Meissn. in DC. Prod. xv. I. 155 = seq.
Rodioei, *Hook. Lond. Journ. Bot.* iii. (1844) 626.—Guian.

NECTANDRA :—
salicifolia, Nees, in Linnaea, xxi. (1848) 506 (Ocoteae sp.).—Mexic.
salicifolia, Nees, Syst. Laurin. 302 = sanguinea.
saligna, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 48.—Bras.
sanguinea, Griseb. Fl. Brit. W. Ind. 281 = Willdenoviana.
sanguinea, Nees, ex Meissn. in DC. Prod. xv. 1. 164 = polita.
sanguinea, *Roland. ex Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 279.—Am. austr.
sarcocalyx, Nees, Syst. Laurin. 338 = nitidula.
Schomburgkii, Meissn. in DC. Prod. xv. 1. 151 = lucida.
Schottii, Meissn. l. c. 147 = rigida.
spicata, Meissn. l. c. 151 = leucantha.
squarrosa, Nees, Syst. Laurin. 326 = Ocotea squarrosa.
staminea, *Griseb. Fl. Brit. W. Ind.* 282.—Ind. occ.
stenophylla, Meissn. in DC. Prod. xv. 1. 162 = Ocotea angustifolia.
striata, *Nees, Syst. Laurin.* 309.—Mexic.
tovarensis, Klotzsch & Karst. ex Nees, in Linnaea, xxi. (1848) 505 = N. Laurel.
turbacensis, *Nees, Syst. Laurin.* 316.—N. Granat.
urophylla, *Meissn. in DC. Prod.* xv. 1. 157.—Bras.
vaga, *Meissn. l. c.* 153.—Guiana.
venulosa, *Meissn. l. c.*—Bras.
viburnoides, *Meissn. l. c.* 162.—Am. trop.
villosa, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 47.—Bras.
virgultosa, Nees, Syst. Laurin. 304 = Ocotea virgultosa.
Warmingii, *Meissn. ex Warm. in Kjoeb. Vidensk. Meddel.* (1870) 141.—Bras.
Weddellii, *Meissn. in DC. Prod.* xv. 1. 148.—Bras.
Willdenoviana, *Nees, Syst. Laurin.* 321. n. 29.—Ind. occ.
Willdenoviana, Nees, l. c. 290. n. 9 = turbacensis.
xanthocalyx, Nees, l. c. 312 = Ocotea xanthocalyx.

NECTANDRA, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 425 (1832) = **Linostoma**, Wall. (Thymel.).
decandra, Roxb. ll. cc. = L. decandrum.

NECTAROBOTHRIUM, Ledeb. Fl. Alt. ii. 36 (1830) = **Lloydia**, Salisb. (Liliac.).
Redowskianum, Cham. in Linnaea, vi. (1831) 585 = L. alpina.
striatum, Ledeb. Fl. Alt. ii. 36 = L. alpina.

NECTAROSCILLA, Parl. Nuov. Gen. e Spec. 26 (1854) = **Scilla**, Linn. (Liliac.).
hyacinthoides, Parl. l. c. 27 = S. hyacinthoides.

NECTAROSCORDUM, Lindl. Bot. Reg. t. 1912 (1836) = **Allium**, Tourn. (Liliac.).
bulgaricum, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 242 = A. Dioscoridis.
siculum, Lindl. Bot. Reg. t. 1912 = A. Dioscoridis.

NECTOLIS, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
ambigua, Rafin. l. c. = S. nigra.
hippophaëfolia, Rafin. l. c. = S. multiformis.
membranacea, Rafin. l. c. = S. purpurea.
vitellina, Rafin. l. c. = S. alba.

NECTOPIX, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
caprea, Rafin. l. c. = S. caprea.
rhamnifolia, Rafin. l. c. = S. rhamnifolia.
sphacelata, Rafin. l. c. = S. grandifolia.

NECTOUXIA, DC. Syst. ii. 149 (1821) = **Morettia**, DC. (Crucif.).

NECTOUXIA, H. B. & K. Nov. Gen. et Sp. iii. 10. t. 193 (1818). *SOLANACEAE*, Benth. & Hook. f. ii. 899.
NETOUXIA, G. Don, Gen. Syst. iv. 454 (1837).
bella, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 138 = formosa.
formosa, *H. B. & K. Nov. Gen. et Sp.* iii. 10. t. 193. —Mexic.

NECTRIS, Schreb. Gen. i. 237 (1789) = **Cabomba**, Aubl. (Nymph.).
aquatica, J. F. Gmel. Syst. 580; Willd. Sp. Pl. ii. 248 = C. aquatica.
aquatica, Nutt. Gen. Am. i. 229 = C. caroliniana.
caroliniana, Steud. Nom. ed. II. ii. 188 = C. caroliniana.
furcata, Leandr. ex Schult. f. Syst. vii. 1379 = C. furcata.
peltata, Pursh, Fl. Am. Sept. i. 238 = C. caroliniana.
pinnata, Pursh, l. c. 239 = C. caroliniana.

NECTUSION, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
reticulata, Rafin. l. c. = S. reticulata.

NEEA, Ruiz & Pav. Prod. 52. t. 9. (1794). *NYCTAGINEAE*, Benth. & Hook. f. i. 9.
MITSCHERLICHIA, Kunth, in Abh. Berl. Akad. (1831) 209.
NEBRA, Noronha, ex Choisy, in DC. Prod. xiii. II. 447 (1849).
NEEANIA, Rafin. Princ. Somiol. 30 (1814).
altissima, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 46.—Bras.
anisophylla, *Ernst, in Flora*, lvii. (1874) 215.—Venezuela.
compressa, *J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 366.—Bras.
constricta, *Spruce, ex J. A. Schmidt, l. c.* 368.—Bras.
divaricata, Poepp. & Endl. Nov. Gen. et Sp. ii. 45 = oppositifolia.
floribunda, *Poepp. & Endl. l. c.* 46.—Peruv.
hirsuta, *Poepp. & Endl. l. c.* 45. t. 161.—Peruv.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 71.—Jamaic.
lanceolata, *Choisy, in DC. Prod.* xiii. II. 449.—Bras.
laxa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 45. t. 162.—Peruv.
macrophylla, *Poepp. & Endl. l. c.* 46.—Peruv.
Mitscherlichia, *Steud. Nom.* ed. II. ii. 152.—Hab.?
mollis, *Spruce, ex J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 367.—Bras.
oppositifolia, *Ruiz & Pav. Syst. Veg.* 91; *Fl. Per.* iv. t. 329.—Peruv.
ovalifolia, *Spruce, ex J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 368.—Bras.
parviflora, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 46.—Peruv.
pubescens, *Poepp. & Endl. l. c.* 45.—Bras.
rosea, *Mart. ex J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 366.—Bras.
rufula, *Mart. ex J. A. Schmidt, l. c.* 368.—Bras.
spectabilis, Choisy, in DC. Prod. xiii. II. 448 = N. Mitscherlichia.
theifera, *Oerst. in Overs. Danske Vidensk. Selsk. Forh.* (1863) 9.—Bras.
verticillata, *Ruiz & Pav. Syst. Veg.* 90; *Fl. Per.* iv. t. 328.—Peruv.

NEEAEA, Poepp. & Endl. Nov. Gen. et Sp. ii. 45 (1838) = praec.

NEEANIA, Rafin. Princ. Somiol. 30 (1814) = praec.

NEEDHAMIA, R. Br. Prod. 549 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 613.
Pumilio, *R. Br. l. c.*—Austral.

NEEDHAMIA, Cass. in Dict. Sc. Nat. xxxiv. 335 (1825) = **Narvalina**, Cass. (Compos.).
domingensis, Cass. l. c. = Narvalina domingensis.

NEEDHAMIA, Scop. Introd. 310 (1777) = **Tephrosia**, Pers. (Legumin.).

NEERIJA, Roxb. Hort. Beng. 86; Fl. Ind. i. 646 (1832) = **Elaeodendron**, Jacq. (Tiliac.).
dichotoma, Roxb. ll. cc. = E. glaucum.

NEESIA, Blume, in Nov. Act. Nat. Cur. xvii. 1. (1835) 83. *MALVACEAE*, Benth. & Hook. f. i. 213.
COTYLEPHORA, Meissn. Gen. Comm. 28 (1837).
ESENBECKIA, Blume, Bijdr. 118 (1825).
altissima, *Blume, in Nov. Act. Nat. Cur.* xvii. (1835) 83. t. 6.—Java.
Griffithii, Planch. ex Mast. in Hook. f. Fl. Brit. Ind. i. 352 = Durio Oxleyanus.
strigosa, *Mast. in Journ. Linn. Soc.* xiv. (1875) 504. —Malaya.
synandra, *Mast. in Hook. f. Fl. Brit. Ind.* i. 352.—Ins. Penang.

NEESIA, Mart. ex Meissn. in Mart. Fl. Bras. v. 1. 67 (1855) = **Funifera**, Leandr. (Thymel.).
daphnoides, Mart. ex Meissn. l. c. = F. utilis.

NEESIA, Spreng. Anleit. ii. II. 547 (1818) = **Diotis**, Desf. (Compos.).
Lupinaster, Spreng. ex Steud. Nom. ed. II. ii. 188 = Trifolium Lupinaster.
maritima, Spreng. ex Steud. l. c. ed. I. 551 = D. candidissima.

NEFFLEA, Spach, Hist. Vég. Phan. ix. 266 (1840) = **Celsia**, Linn. (Scrophul.).

NEFRAKIS, Rafin. Sylva Tellur. 69 (1838) = **Brya**, P. Br. (Legumin.).
Ebenus, Rafin. l. c. = B. Ebenus.

NEGRETIA, Ruiz & Pav. Prod. 98. t. 21 (1794) = **Mucuna**, Adans. (Legumin.).
elliptica, Ruiz & Pav. Syst. Veg. 176 = M. elliptica.
inflexa, Ruiz & Pav. l. c. = M. inflexa.
mitis, Blanco, Fl. Filip. ed. I. 588 = M. nivea.
mitis, Ruiz & Pav. Syst. Veg. 177 = M. mitis.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 444 = M. mollis.
Mutisiana, H. B. & K. l. c. 442 = M. Mutisiana.
nigricans, Poir. Encyc. Suppl. iv. 74 = M. nigricans.
platycarpa, Ruiz & Pav. Syst. Veg. 178 = M. platycarpa.
pruriens, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 63 = M. pruriens.
sericea, Willd. ex Steud. Nom. ed. II. ii. 164, 188 = M. urens.
urens, Blanco, Fl. Filip. ed. I. 586 = M. monosperma.
urens, Tussac, Fl. Antill. ii. 52. t. 13 = M. urens.

NEGRIA, F. Muell. Fragm. vii. 151 (1871). *GESNERACEAE*, Benth. & Hook. f. ii. 1011.
rhabdothamnoides, *F. Muell. l. c.* 152.—Austral.

NEGUNDIUM, Rafin. in Med. Repos. N. York, v. (1808) 352; et ex Desv. Journ. Bot. ii. (1809) 166 = **Negundo**, Moench (Sapind.).
americanum, DC. ex Loud. Hort. Brit. 398 = Negundo aceroides.
fraxineum, Steud. Nom. ed. II. ii. 188 = seq.
fraxinifolium, Rafin. in Med. Repos. N. York, v. (1808) 352; et ex Desv. Journ. Bot. ii. (1809) 170 = Negundo aceroides?
lobatum, Rafin. New Fl. Am. i. 48 = Negundo aceroides?
trifoliatum, Rafin. Atl. Journ. 176 = Negundo aceroides?

NEGUNDO, Moench, Meth. 334 (1794). *SAPINDACEAE*, Benth. & Hook. f. i. 409.
NEGUNDIUM, Rafin. in Med. Repos. N. York, v. (1808) 352; et ex Desv. Journ. Bot. ii. (1809) 166.
aceroides, *Moench, Meth.* 334.—Am. bor.
aceroides, Torr. in Pacif. Rail. Rep. iv. 74 = californicum.
californicum, *Torr. & Gray, Fl. N. Am.* i. 250.—Am. bor. occ.
cissifolium, Sieb. & Zucc. in Abh. Akad. Muench, iv. II. (1843) 159 = Acer cissifolium.
cochinchinense, DC. Prod. i. 596 = Acer pinnatum.
fraxinifolium, Nutt. Gen. Am. i. 253 = aceroides.
Fraxinus, Bourg. in Journ. Linn. Soc. iv. (1859) 9 = aceroides.
mandshuricum, *Budischtschew, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 171, nomen.—Manchuria.
mexicanum, *DC. Prod.* i. 596.—Mexic.
nikoënse, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 90 = Acer Maximowiczianum.
sessilifolium, Miq. l. c. 89 = Acer palmatum.
virginianum, Medic. Pfl. Anat. 439 = aceroides.

NEILLIA, D. Don, Prod. Fl. Nep. 228 (1825). *ROSACEAE*, Benth. & Hook. f. i. 612.
ADENILEMA, Blume, Bijdr. 1121 (1826).
EPICOSTORUS, Rafin. Atl. Journ. 144 (1832).
ICOTORUS, Rafin. in Ser. Bull. Bot. i. (1830) 216.
OPULASTER, Medic. Pfl. Anat. 109 (1799).
PHYSOCARPA, Rafin. New Fl. Am. iii. 73 (1836).
Adenilema, Miq. Fl. Ind. Bat. i. 1. 391 = thyrsiflora.
fallax, *Blume, Mél. Bot.* (1855).—*Cf. Flora*, xli. (1858) 254 = thyrsiflora.

NEILLIA :—
opulifolia, *Benth. & Hook. f. Gen.* i. 612.—Am. bor.
rubiflora, *D. Don, Prod. Fl. Nep.* 228.—Reg. Himal.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* i. 121.—Japon.
thyrsiflora, *D. Don, Prod. Fl. Nep.* 228.—Reg. Himal.; Java.
Torreyi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 136.—Am. bor. occ.
virgata, Wall. Cat. n. 7108 = thyrsiflora.

NEILREICHIA, Fenzl, in Denkschr. Akad. Wien, i. (1850) 258. t. 30 = **Schistocarpha**, Less. (Compos.).
eupatorioides, Fenzl, l. c.—Peruv.

NEILREICHIA, Kotule, in Spraw. Kom. Fis. Krakow. xvii. (1883) 136 = **Carex**, Dill. (Cyperac.).
umbrosa, Kotule, l. c. = C. praecox.

NEIOSPERMA, Rafin. Sylva Tellur. 162 (1838) = **Ochrosia**, Juss. (Apocyn.).
muricata, Rafin. l. c. = O. borbonica.
musculiformis, Rafin. l. c. = Banksia musculiformis.

NEIPPERGIA, C. Morr. in Ann. Soc. Bot. Gand, v. (1849) 375. t. 282 = **Acineta**, Lindl. (Orchid.).
chrysantha, C. Morr. l. c. = A. chrysantha.

NEISANDRA, Rafin. Sylva Tellur. 163 (1838) = **Hopea**, Roxb. (Dipteroc.).
indica, Rafin. l. c. = H. odorata.

NEJA, D. Don, in Sweet, Hort. Brit. ed. II. 299 (1830); et Brit. Flow. Gard. Ser. II. t. 78 (1831) = **Hysterionica**, Willd. (Compos.).
ciliaris, DC. Prod. v. 326 = H. pinifolia.
falcata, Nees, Del. Sem. Hort. Vratisl. (1839).—Cf. Linnaea, xiv. (1840) Litt. 168.—Bras.
filiformis, Nees, l. c. = H. pinifolia.
gracilis, D. Don. in Sweet, Hort. Brit. ed. II. 299; et Brit. Flow. Gard. Ser. II. t. 78 = H. linearifolia.
linearifolia, DC. Prod. v. 325 = H. linearifolia.
macrocephala, DC. l. c. (= *Asteropsis macrocephala*).—Bras.
montevidensis, Sch. Bip. in Seem. Bot. Voy. Herald, 302 = H. montevidensis.
nidorelloides, DC. Prod. v. 325 = H. nidorelloides.
Sprengelii, Sch. Bip. in Seem. Bot. Voy. Herald, 302 = H. montevidensis.
subvillosa, DC. Prod. v. 325 = H. montevidensis.
tenuifolia, DC. l. c. 326 = H. montevidensis.

NEKEMIAS, Rafin. Sylva Tellur. 87 (1838) = **Vitis**, Tourn. (Ampel.).
bipinnata, Rafin. l. c. = V. arborea.

NELEIXA, Rafin. New Fl. Am. iv. 55 (1836) = **Faramea**, Aubl. (Rubiac.).
prostrata, Rafin. l. c. = F. odoratissima.

NELENSIA, Poir. Illustr. Gen. iii. 568 (1823) = **Pedicularis**? Tourn. (Scrophul.).
bibracteata, Poir. l. c. (= *Enslenia bibracteata*, Rafin.).—Am. bor.

NELIPUS, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentib.).
bifida, Rafin. l. c. = U. bifida.
biloba, Rafin. l. c. = U. biloba.
limosa, Rafin. l. c. = U. limosa.
spiralis, Rafin. l. c. = U. spiralis.

NELITRIS, Gaertn. Fruct. i. 134. t. 27 (1788) = **Timonius**, Rumph. (Rubiac.).
Jambosella, Gaertn. l. c. = T. Jambosella.

NELITRIS, Spreng. Syst. ii. 488 (1825) = **Decaspermum**, Forst. (Myrtac.).
alba, Blume, Mus. Bot. Lugd. Bat. i. 74.—Ins. Amboina.
alternifolia, Miq. Fl. Ind. Bat. i. 1. 476.—Java.
Billardieri, Seem. Fl. Vit. 81.—N. Caled.
bracteata, Blume, Mus. Bot. Lugd. Bat. i. 74.—Ins. Molucc.
confinis, Blume, l. c.—Malaya.
Coriandri, Blume, l. c.—N. Guin.
Forsteri, Seem. Fl. Vit. 81 = D. fruticosum.
fruticosa, A. Gray, Bot. U. St. Expl. Exped. i. 547. t. 60.—Ins. Fiji.

NELITRIS :—
humilis, G. Don, Gen. Syst. ii. 829.—Austral.
ingens, F. Muell. ex Benth. Fl. Austral. iii. 294 = (Memecylonis sp. ?).—Austral.
laxiflora, Blume, Mus. Bot. Lugd. Bat. i. 74.—N. Guin.
leucocoma, Miq. Fl. Ind. Bat. i. 1. 474.—Sumatra.
lucida, Blume, Mus. Bot. Lugd. Bat. i. 74.—Java.
myrsinoides, Blume, l. c. 75.—Malaya.
pallescens, Miq. Fl. Ind. Bat. Suppl. 314 = Decaspermum paniculatum.
paniculata, Lindl. Coll. Bot. 16 = D. paniculatum.
parviflora, Blume, Mus. Bot. Lugd. Bat. i. 75.—Java.
polygama, Blume, l. c.—Malaya.
polygama, Spreng. Syst. ii. 488 = D. paniculatum.
polymorpha, Blume, Mus. Bot. Lugd. Bat. i. 75 = D. paniculatum.
psidioides, G. Don, Gen. Syst. ii. 829 = Rhodomyrtus psidioides.
pubescens, Miq. Fl. Ind. Bat. i. 1. 475.—Borneo.
pyrifolia, Miq. l. c.—Java.
rubra, Blume, Mus. Bot. Lugd. Bat. i. 73 = D. rubrum ?
rubra, Vidal, Sin. Pl. Leñ. t. 50. f. C = D. Blancoi.
trinervia, Spreng. Syst. ii. 488 = Eugenia dumetorum.
Urvillei, DC. Prod. iii. 231.—Ins. Sechell.
vitiensis, A. Gray, U. St. Expl. Exped. i. 548. t. 60.—Ins. Fiji.

NELLICA, Rafin. Sylva Tellur. 92 (1838) = **Phyllanthus**, Linn. (Euphorb.).
maderaspatana, Rafin. l. c. = P. maderaspatensis.

NELSONIA, R. Br. Prod. 480 (1810). *ACANTHACEAE*, Benth. & Hook. f. ii. 1073.
albicans, H. B. & K. Nov. Gen. et Sp. ii. 234 = campestris.
campestris, *R. Br. Prod.* 481.—Amphig. trop.
canescens, Spreng. Syst. i. 42 = campestris.
hirsuta, Roem. & Schult. Syst. i. 172 = Hemigraphis hirsuta.
lamiifolia, Spreng. Syst. i. 42 = campestris.
nummulariaefolia, Roem. & Schult. Syst. i. 173 = campestris.
origanoides, Roem. & Schult. l. c. = campestris.
Pohlii, Nees, in Mart. Fl. Bras. ix. 15 = campestris.
rotundifolia, R. Br. Prod. 481 = campestris.
senegalensis, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 119 = campestris.
Smithii, Oerst. l. c. = campestris.
tomentosa, A. Dietr. Sp. Pl. i. 419 = campestris.
vestita, Schult. Mant. i. 145 = campestris.
villosa, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 118 = campestris.

NELTUMA, Rafin. Sylva Tellur. 119 (1838) = **Prosopis**, Linn. (Legumin.).
arenosa, Rafin. l. c. = Mimosa arenosa.
juliflora, Rafin. l. c.—P. juliflora.

NELUMBIUM, Juss. Gen. 68 (1789). *NYMPHAEACEAE*, Benth. & Hook. f. i. 47.
CYAMUS, Sm. Exot. Bot. i. 59. t. 31, 32 (1804).
NELUMBO, [Tourn.] Adans. Fam. ii. 76 (1763).
TAMARA, Roxb. ex Steud. Nom. ed. II. ii. 661 (1841).
album, Bercht. & Presl, Rostl. i. Nelumbiac. 2 = speciosum.
asiaticum, Rich. in Ann. Mus. Par. xvii. (1811) 249 = speciosum.
caspicum, Fisch. ex DC. Syst. ii. 45 = speciosum.
caspium, Eichw. Casp. Cauc. 2 = praec.
codophyllum, Rafin. Fl. Ludov. 22 = luteum.
discolor, Steud. Nom. ed. II. ii. 188 = speciosum.
indicum, Poir. in Lam. Encyc. iv. 453 (Nelumbo) = speciosum.
indicum, Tussac, Fl. Antill. iii. 83 = luteum.
jamaicense, DC. Syst. ii. 41 = luteum.
javanicum, Poir. in Lam. Encyc. ii. 454 (Nelumbo) = speciosum.
luteum, *Willd. Sp. Pl.* ii. 1259.—Am. bor.; Ind. occ.
marginatum, Steud. Nom. ed. II. ii. 188 = speciosum.
nuciferum, Gaertn. Fruct. i. 73 (Nelumbo) = speciosum.
pentapetalum, Willd. Sp. Pl. ii. 1259 = luteum.
reniforme, Willd. l. c. 1260 = luteum.
Rheedii, Presl, Rel. Haenk. ii. 183 in obs. = speciosum ?

NELUMBIUM :—
speciosum, *Willd. Sp. Pl.* ii. 1258.—Afr. bor.; As. trop.
Tamara, Sweet, Hort. Brit. ed. I. 14 = speciosum.
transversum, Presl, Rel. Haenk. ii. 83 = speciosum.
turbinatum, Blanco, Fl. Filip. ed. I. 458 = speciosum.
venosum, Presl, Rel. Haenk. ii. 183, in obs. = speciosum ?

NELUMBO, [Tourn.] Adans. Fam. i. 73 (1763) = **Nelumbium**, Juss. (Nymph.).

NEMACAULIS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 168. *POLYGONACEAE*, Benth. & Hook. f. iii. 94.
denudata, Nutt. l. c. = Nuttallii.
foliosa, Nutt. l. c. 169 = Nuttallii.
Nuttallii, *Benth. in DC. Prod.* xiv. 23.—Calif.

NEMACLADUS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 254. *CAMPANULACEAE*, Benth. & Hook. f. ii. 554.
capillaris, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 196.—Calif.
longiflorus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 60.—Calif.
montanus, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 197.—Calif.
pinnatifidus, *Greene, l. c.*—Calif.
ramosissimus, *Nutt. in Trans. Am. Phil. Soc.* N. S. viii. (1843) 254.—Calif.
rigidus, *Curran, in Bull. Calif. Acad.* i. iii. (1885) 154.—Calif.
rubescens, *Greene, l. c.* iv. (1885) 197.—Calif.
tenuissimus, *Greene, l. c.* 198.—Calif.

NEMACONIA, Knowles & Westc. Flor. Cab. ii. (1838) 127 = **Ponera**, Lindl. (Orchid.).
graminifolia, Knowles & Westc. l. c. 129 = P. graminifolia.

NEMALLOSIS, Rafin. Fl. Tellur. iii. 34 (1836) = **Mollugo**, Linn. (Ficoid.).
erecta, Rafin. l. c.—Hab. ?
prostrata, Rafin. l. c. = M. oppositifolia.

NEMAMPSIS, Rafin. Fl. Tellur. iv. 16 (1836) = **Dracaena**, Vand. (Liliac.).
ternifolia, Rafin. l. c. = D. surculosa.

NEMANTHERA, Rafin. Fl. Tellur. iv. 80 (1836) = **Ipomoea**, Linn. (Convolv.).
bufalina, Rafin. l. c. = I. bufalina.

NEMASTACHYS, Steud. Syn. Pl. Gram. 357 (1855) = **Pollinia**, Trin. (Gramin.).
taitensis, Steud. l. c. = P. glabrata.

NEMASTYLIS, Nutt. in Trans. Am. Phil. Soc. v. (1837) 157. *IRIDEAE*, Benth. & Hook. f. iii. 696.
CHLAMYDOSTYLUS, Baker, in Journ. Bot. xiv. (1876) 185.
EUSTYLIS, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1847) 236.
acuta, Herb. in Bot. Mag. sub t. 3779 = geminifolia.
coelestina, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 157.—Mexic.
furcata, *Klatt, in Linnaea*, xxxi. (1861–62) 560.—Reg. Argent.
geminiflora, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 157.—Am. bor.
Medusa, *Benth. & Hook. f. Gen.* iii. 696 (Medicae).—Bras.
multiflora, *Benth. & Hook. f. l. c.*—Mexic.
nana, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 160.—Mexic.
punctata, *Hemsl. Biol. Centr. Am. Bot.* iii. 329.—Mexic.
purpurea, *Herb. in Bot. Mag.* sub t. 3779.—Am. bor.
spathacea, *Griseb. in Goett. Abh.* xix. (1874) 222.—Bolivia.
tenuis, Benth. & Hook. f. Gen. iii. 696 = coelestina.
triflora, *Herb. in Benth. Pl. Hartw.* 95.—Guatemala.

NEMASTYLUS, Baker, in Journ. Linn. Soc. xvi. (1877) 102 = praec.

NEMATANTHERA, Miq. in Linnaea, xviii. (1844) 606. t. 11 = **Piper**, Linn.
guianensis, Miq. l. c. 607 = P. Nematanthera.

NEMATANTHUS, Nees, in Linnaea, v. (1830) 661 = **Willdenowia**, Thunb. (Restiac.).
Ecklonii, Nees, l. c. 662 = W. striata.

NEMATANTHUS, Schrad. in Goett. Gel. Anz. i. (1821) 718; Mart. Nov. Gen. et Sp. iii. 46. t. 220 (1829). *GESNERACEAE*, Benth. & Hook. f. ii. 1010.
calcaratus, *Miq. in Linnaea*, xxii. (1849) 472.—Guiana.
calycinus, *Presl, Bot. Bemerk.* 145.—Bras.
chloronema, Hort. ex Tenore, in Parl. Giorn. Bot. Ital. ii. II. (1847) 202 = Hypocyrta perianthomega.
chloronema, Mart. Nov. Gen. et Sp. iii. 47. t. 219 = longipes.
corticola, *Schrad. in Goett. Gel. Anz.* i. (1821) 718.—Bras.
erianthus, *Benth. Pl. Hartw.* 231.—Ecuador.
erythrocalyx, *Hort. ex Lem. in Fl. des Serres*, ii. (Juill. 1846) sub t. 9; (Août, sub t. 5).—Hab.?
Guilleminianus, Hort. ex Paxt. Mag. Bot. x. (1843) 5 = longipes.
heterophyllus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 4. t. 103.—Peruv.
heterophyllus, *Presl, Bot. Bemerk.* 145.—Bras.
jonema, Mart. Nov. Gen. et Sp. iii. 48 = corticola.
longipes, *DC. Prod.* vii. 544.—Bras.
Morrelliana, Hort. ex Hook. Bot. Mag. sub t. 4460 = corticola.
pereskiaefolia, *Presl, Bot. Bemerk.* 143 (err. typ. 145).—Bras.
radicans, *Presl, l. c.*—Bras.
serrulatus, *Presl, l. c.* 144.—Bras.

NEMATOCERAS, Hook. f. Fl. N. Zel. i. 249. t. 57 (1855) = **Corysanthes**, R. Br. (Orchid.).
macrantha, Hook. f. l. c. 250 = C. macrantha.
oblonga, Hook. f. l. c. = C. oblonga.
rivularis, Hook. f. l. c. 251 = C. rivularis.
rotundifolia, Hook. f. l. c. = C. rotundifolia.
triloba, Hook. f. l. c. 250 = C. triloba.

NEMATOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 158. *RUTACEAE*, Benth. & Hook. f. i. 295.
SYMPHYOPETALUM, J. Drumm. in Hook. Kew Journ. vii. (1855) 54.
Euphemiae, *F. Muell. Fragm.* iii. 149. t. 25.—Austral.
phebalioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 158.—Austral.

NEMATOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 20 = **Templetonia**, R. Br. (Legumin.).
Hookeri, F. Muell. l. c. = T. Hookeri.

NEMATOPUS, A. Gray, in Hook. Kew Journ. iii. (1851) 150 = **Gnephosis**, Cass. (Compos.).
effusus, A. Gray, l. c. = G. arachnoides.
foliatus, Sond. in Linnaea, xxv. (1852) 486 = G. arachnoides.

NEMATOPYXIS, Miq. Fl. Ind. Bat. i. I. 630 (1855) = **Ludwigia**, Linn. (Onagrar.).
fruticulosa, Miq. l. c. = L. prostrata.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 95 = L. prostrata.
prostrata, Miq. Fl. Ind. Bat. i. I. 630 = L. prostrata.
pusilla, Miq. l. c. = L. prostrata.

NEMATOSPERMUM, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = **Lacistema**, Sw.
laevigatum, Rich. l. c. = L. myricoides.

NEMATOSTIGMA, A. Dietr. Sp. Pl. ii. 509 (1833) = **Libertia**, Spreng. (Irid.).
ixioides, A. Dietr. l. c. 510 = L. ixioides.
paniculatum, A. Dietr. l. c. = L. paniculata.
pulchellum, A. Dietr. l. c. = L. pulchella.

NEMATOSTIGMA, Planch. ex Benth. & Hook. f. Gen. iii. 356 (1880) = NEMOSTIGMA, Planch. = **Gironniera**, Gaudich. (Urtic.).

NEMATOSTYLIS, Hook. f. in Benth. & Hook. f. Gen. ii. 110 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
anthophylla, Baill. Hist. Pl. vii. 432 = loranthoides.
loranthoides, *Hook. f. in Benth. & Hook. f. Gen.* ii. 110.—Madag.

NEMATURIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 254 = **Enslenia**, Nutt. (Asclep.).
volubilis, Turcz. l. c. = E. volubilis.

NEMAUCHENES, Cass. in Bull. Soc. Philom. (1818) 77 = **Crepis**, Linn. (Compos.).
aculeata, Cass. in Dict. Sc. Nat. xxxiv. 362 = C. aspera.
ambigua, Cass. in Bull. Soc. Philom. (1818) 77 = C. aspera.
aspera, Steud. Nom. ed. II. ii. 189 = C. aspera.
inermis, Cass. in Dict. Sc. Nat. xxxiv. 363 = C. aspera.

NEMAULAX, Rafin. Fl. Tellur. iii. 52 (1836) = **Albuca**, Linn. (Liliac.).
fastigiata, Rafin. l. c. = A. fastigiata.

NEMEDRA, A. Juss. in Mém. Mus. Par. xix. (1830) 223 = **Aglaia**, Lour. (Meliac.).
elaeagnoidea, A. Juss. l. c. = A. elaeagnoidea.
Nimmonii, Dalz. in Dalz. & Gibs. Bomb. Fl. 37 = Amoora Lawii.

NEMEPIODON, Rafin. Fl. Tellur. iv. 22 (1836) = **Hymenocallis**, Salisb. (Amaryll.).
caribeum, Rafin. l. c. = H. caribaea.
mexicanum, Rafin. l. c. = H. lacera.
speciosum, Rafin. l. c. = H. speciosa.

NEMEPIS, Rafin. Fl. Tellur. iv. 91 (1836) = **Cuscuta**, Tourn. (Convolv.).
africana, Rafin. l. c. = C. ungulata.
americana, Rafin. l. c. = C. americana.
odorata, Rafin. l. c. = C. odorata.
prolifera, Rafin. l. c. (Quid?).—Hab.?

NEMESIA, Vent. Jard. Malm. 41. t. 41 (1803). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 931.
acuminata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 22.—Afr. austr.
affinis, Benth. l. c. 21 = floribunda.
anisocarpa, *E. Mey. ex Benth. l. c.* 19.—Afr. austr.
barbata, *Benth. l. c.*—Afr. austr.
bicornis, *Pers. Syn.* ii. 159.—Afr. austr.
calcarata, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 20.—Afr. austr.
chamaedrifolia, *Vent. Jard. Malm.* sub t. 42.—Afr. austr.
Cheiranthus, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 19.—Afr. austr.
cynanchifolia, *Benth. l. c.* 21.—Afr. austr.
dentata, *G. Don, Gen. Syst.* iv. 534.—Afr. austr.
diffusa, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 22.—Afr. austr.
divergens, *Benth. l. c.*—Afr. austr.
floribunda, *Lehm. Del. Sem. Hort. Hamb.* (1833); *ex Benth. in Bot. Reg.* (1838) t. 39.—Afr. austr.
foetens, *Vent. Jard. Malm.* 41. t. 41.—Afr. austr.
frutescens, G. Don, in Sweet, Hort. Brit. ed. III. 515 = Linaria fruticosa.
fruticans, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 22.—Afr. austr.
gracilis, *Benth. l. c.* 20.—Afr. austr.
hastata, *Benth. l. c.* 19.—Afr. austr.
ligulata, *E. Mey. ex Benth. l. c.* 20.—Afr. austr.
linearis, *Vent. Jard. Malm.* sub t. 41.—Afr. austr.
longicornis, *Pers. Syn.* ii. 159.—Afr. austr.
longicornis, Vent. ex Steud. Nom. ed. II. i. 501, in syn. = Diascia Thunbergiana.
lucida, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 19.—Afr. austr.
melissaefolia, *Benth. l. c.* 22.—Afr. austr.
monomotapensis, Benth. in DC. Prod. x. 264 = dentata.
natalitia, *Sond. in Linnaea*, xxiii. (1850) 82.—Afr. austr.
parviflora, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 20.—Afr. austr.
patens, *G. Don, Gen. Syst.* iv. 534.—Afr. austr.
pinnata, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 20.—Afr. austr.
pubescens, *Benth. l. c.* 22.—Afr. austr.
saccata, *E. Mey. ex Benth. l. c.* 19.—Afr. austr.
strumosa, *Benth. l. c.* 18.—Afr. austr.
Thunbergii, G. Don, Gen. Syst. iv. 534 = linearis.
versicolor, *E. Mey. ex Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 20.—Afr. austr.
viscosa, *E. Mey. ex Benth. l. c.* 21.—Afr. austr.

NEMEXIA, Rafin. Neogenyt. 3 (1825) = **Smilax**, Tourn.
cerulea, Rafin. l. c. = S. herbacea.
nigra, Rafin. l. c. = S. herbacea.

NEMIA, Berg. Desc. Pl. Cap. 160 (1767) = **Manulea**, Linn. (Scrophul.).
capensis, J. F. Gmel. Syst. 936 = M. Cheiranthus.
Cheiranthus, Berg. Desc. Pl. Cap. 160 = M. Cheiranthus.
rubra, Berg. l. c. 161 = M. rubra.

NEMITIS, Rafin. Fl. Tellur. iv. 33 (1836) = **Apteria**, Nutt. (Burman.).
setacea, Rafin. l. c. = A. setacea.

NEMOCHARIS, Beurl. in Bot. Notiser (1853), 52 = **Scirpus**, Tourn. (Cyperac.).
radicans, Burl. l. c. = S. radicans.
sylvatica, Beurl. l. c. = S. sylvaticus.

NEMOCHLOA, [Nees, in] Meyen, Reise, i. (1834) 108; et in Mart. Fl. Bras. ii. I. 148. t. 14, 15 (1842) = **Pleurostachys**, Brongn. (Cyperac.).
Beyrichii, Nees, in Mart. Fl. Bras. ii. I. 152 = P. Beyrichii.
elegans, Nees, l. c. 149 = P. elegans.
extenuata, Nees, l. c. 152 = P. extenuata.
foliosa, Nees, l. c. 152 = P. foliosa.
Gaudichaudi, Nees, l. c. = P. Gaudichaudii.
macrantha, Nees, l. c. 151 = P. macrantha.
Martiana, Nees, l. c. 150 = P. Martiana.
millegrana, Nees, l. c. 150 = P. Martiana.
Orbigniana, Nees, ex Steud. Nom. ed. II. ii. 189 = P. Orbigniana.
Sellovii, Nees, in Mart. Fl. Bras. ii. I. 152 = P. Sellowii.
sparsiflora, Nees, l. c. = P. sparsiflora.
stricta, Nees, l. c. = P. stricta.
tenuiflora, Nees, l. c. 149 = P. tenuiflora.
turbinata, [Nees, in] Meyen, Reise, i. 108 = P. tenuiflora.
Urvillei, Nees, in Mart. Fl. Bras. ii. I. 150 = P. Urvillei.

NEMOCTIS, Rafin. Fl. Tellur. iv. 106 (1836) = **Daphne**, Linn. (Thymel.).
angustifolia, Rafin. l. c. = D. oleoides.
buxifolia, Rafin. l. c. = D. oleoides.

NEMODAPHNE, Meissn. in DC. Prod. xv. I. 109 (1864) = **Ocotea**, Aubl. (Laurin.).
cuneata, Meissn. l. c.—Cuba.

NEMODON, Griff. Notul. iv. 286 (1854) = **Lepistemon**, Blume (Convolv.).

NEMOLEPIS, Vilm. Fl. Pl. Terre, ed. II. 396 (1866) = **Heliopsis**, Pers. (Compos.).
canescens, Vilm. l. c. = H. buphthalmoides.

NEMOPANTHES, Rafin. in Journ. Phys. lxxxviii. (1819) 96 = seq.

NEMOPANTHUS, Rafin. in Am. Monthly Mag. (1819) 357. *ILICINEAE*, Benth. & Hook. f. i. 357.
ILICIOIDES, Dum.-Cours. Bot. Cult. ed. I. iv. 127 (1804).
NUTTALLIA, DC. Rapp. Jard. Genèv. (1821) 44.
ambiguus, Wood, Class-Book (1861), 497 = Ilex ambigua.
Andersonii, Hort. ex Pasquale, Cat. Ort. Nap. (1866) adn. = Ilex dipyrena.
canadensis, DC. in Mém. Soc. Genèv. i. (1821) 44 = fascicularis.
fascicularis, *Rafin. in Journ. Phys.* lxxxviii. (1819) 96.—Am. bor.
lucida, C. Koch, Dendrol. ii. 229 = fascicularis.

NEMOPHILA, Nutt. ex Barton, Fl. N. Am. ii. 71. t. 61 (1882); et in Journ. Acad. Philad. ii. (1822) 179. *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 827.
PHOLISTOMA, Lilja, in Bot. Notiser (1839), 40.
atomaria, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 42 = Menziesii.
auriculaeflora, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 746 = Menziesii.
aurita, *Lindl. Bot. Reg.* t. 1601.—Calif.
breviflora, *A. Gray, in Proc. Am. Acad.* x. (1875) 315.—Am. bor. occ.

NEMOPHILA:—
crambeoides, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 586 = Menziesii.
discoidalis, Lem. in Fl. des Serres, ii. (Août, 1846) t. 7 [75] = Menziesii.
evanescens Darby, Bot. South. St. 444 = microcalyx.
heterophylla, Fisch. & Mey. Sert. Petrop. t. 6 = parviflora.
hirsuta, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 462 = phacelioides.
insignis, *Benth. in Trans. Hort. Soc.* N. S. i. (1835) 479; *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1836) 275.—Calif.
liniflora, Fisch. & Mey. Sert. Petrop. t. 5 = Menziesii.
maculata, *Benth. ex Lindl. in Journ. Hort. Soc.* iii. (1848) 320.—Calif.
Menziesii, *Hook. & Arn. Bot. Beech. Voy.* 152.—Am. bor. occ.
microcalyx, *Fisch. & Mey. Sert. Petrop.* t. 6.—Am. bor.
modesta, Kellogg, in Proc. Calif. Acad. vii. (1877) 93 = Menziesii.
nemorosa, Reuben-Heynes, ex Steud. Nom. ed. II. ii. 189 = phacelioides.
Nuttallii, Colla, Hort. Ripul. App. i. 135. t. 5 = phacelioides.
paniculata, Spreng. Syst. i. 569 = Hydrophyllum appendiculatum.
parviflora, *Dougl. ex Benth. in Trans. Linn. Soc.* xvii. (1836) 275.—Am. bor. occ.
pedunculata, Dougl. l. c. = parviflora.
phacelioides, *Barton, Fl. N. Am.* ii. 71. t. 61; *Nutt. in Journ. Acad. Philad.* ii. (1822) 179.—Am. bor. occ.
pilosa, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 463 = phacelioides.
racemosa, *Nutt. ex A. Gray, in Proc. Am. Acad.* x. (1875) 315.—Calif.
speciosa, Hartw. ex Lindl. in Journ. Hort. Soc. iii. (1848) 319 = maculata.

NEMOPHILLA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 462 = praec.

NEMOPOGON, Rafin. Fl. Tellur. ii. 27 (1836) = **Bulbine**, Linn. (Liliac.).
glaucum, Rafin. l. c. = B. semibarbata.

NEMORINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 172 = **Luzula**, DC. (Juncac.).
flavescens, Fourr. l. c. = L. Hostii.
Forsteri, Fourr. l. c. = L. Forsteri.
vernalis, Fourr. l. c. = L. vernalis.

NEMOSTIGMA, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 265 = **Gironniera**, Gaudich. (Urtic.).

NEMOSTIMA, Rafin. Fl. Tellur. iv. 82 (1836) = **Convolvulus**, Tourn.
canariensis, Rafin. l. c. = C. canariensis.
Cantabrica, Rafin. l. c. = C. Cantabrica.
Hystrix, Rafin. l. c. = C. Hystrix.
lanata, Rafin. l. c. = C. sericeus?
lineata, Rafin. l. c. = C. lineatus.
spicaefolia, Rafin. l. c. = C. lineatus.

NEMOSTYLIS, Herb. in Bot. Mag. sub t. 3779 (1840) = **Nemastylis**, Nutt. (Irid.).

NEMOSTYLIS, Stev. Verz. Taur. Halbinsel, 187, in obs. (1857) = **Phuopsis**, Benth. & Hook. f. (Rubiac.) (1873).
ciliaris, Stev. l. c. = P. stylosa.

NEMUARON, Baill. Adansonia, x. (1873) 351. *MONIMIACEAE*, Benth. & Hook. f. iii. 145.
Humboldtii, *Baill. l. c.* 354.—N. Caled.
Vieillardi, *Baill. l. c.* 353.—N. Caled.

NEMUM, Desv. in Ham. Prod. Pl. Ind. Occ. 13 (1825) = **Scirpus**, Tourn. (Cyperac.).
spadiceum, Desv. l. c. = S. spadiceus.

NEMURANTHES, Rafin. Fl. Tellur. ii. 61 (1836) = **Habenaria**, Willd. (Orchid.).
Habenaria, Rafin. l. c. 62 = H. macroceratitis.
longicauda, Rafin. l. c. 61 = H. longicauda.

NENAX, Gaertn. Fruct. i. 165. t. 32 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 140.
AMBRARIA, Cruse, Rub. Cap. 16. t. 1. ff. 3, 4 (1825).
acerosa, *Gaertn. Fruct.* i. 165. t. 32. f. 7.—Afr. austr.

NENGA, H. Wendl. & Drudè, in Linnaea, xxxix. (1875) 182. *PALMAE*, Benth. & Hook. f. iii. 888.
affinis, Becc. Malesia, i. (1877) 29 = Leptophoenix affinis.
celebica, Becc. in Ann. Jard. Buitenz. ii. (1885) 86 = *selebica*.
Geelvinkiana, Becc. Malesia, i. (1877) 28 = Adelonenga Geelvinkiana.
gracilis, Becc. l. c. 25 = Pinanga gracilis.
intermedia, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 81, 85.—Malacca.
latisecta, Scheff. l. c. i. (1876) 120; Becc. Malesia, i. (1877) 25 = Pinanga latisecta.
nagensis, Scheff. l. c.; Becc. l. c. = Areca nagensis.
pinangoides, Becc. l. c. 28 = Leptophoenix pinangoides.
pumila, H. Wendl. in Kerch. Palm. 251 = Wendlandiana.
Schefferiana, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 81, 84.—Sumatra.
selebica, Becc. Malesia, i. (1877) 30 = Gronophyllum selebicum.
variabilis, Becc. l. c. 26 = Adelonenga variabilis.
Wendlandiana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 153.—Java.

NENGELLA, Becc. Malesia, i. (1877) 32. t. 1. *PALMAE*, Benth. & Hook. f. iii. 886.
flabellata, *Becc. l. c.* 34.—N. Guin.
montana, *Becc. l. c.* 33.—Malaya.
paradoxa, Becc. l. c. 32, in obs. = Ophiria paradoxa.

NENNINGIA, Opiz, Seznam, 68 (1852) = **Campanula**, Tourn.
paniculata, Opiz, l. c. = C. sibirica.

NENUPHAR, Link, Enum. Hort. Berol. ii. 70 (1822) = **Nuphar**, Sm. (Nymph.).
advena, Link, l. c. = Nuphar advena.
luteum, Link, l. c. = Nuphar luteum.
minimum, Link, l. c. = Nuphar minimum.
pumila, Bluff & Fingh. Comp. Fl. Germ. i. 705 = Nuphar minimum.

NEOBARONIA, Baker, in Journ. Linn. Soc. xxi. (1884) 336 = **Phylloxylon**, Baill. Adansonia, ii. (1861) 54.
phyllanthoides, Baker, l. c. 337.—Madag.

NEOBOUTONIA, Muell. Arg. in Journ. Bot. ii. (1864) 336. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 317.
africana, *Muell. Arg. l. c.*—Afr. trop.
Mannii, *Benth. & Hook. f. Gen.* iii. 317.—Afr. trop.

NEOCEIS, Cass. in Bull. Soc. Philom. (1820) 90 = **Erechtites**, Rafin. (Compos.).
carduifolia, Cass. in Dict. Sc. Nat. xxxiv. 386 = E. praealta.
hieracifolia, Cass. l. c. 387 = E. praealta.
microcephala, Cass. l. c. 388 = E. arguta?
rigidula, Cass. l. c. 387 = E. praealta.

NEOCLIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 (Quid?).

NEODRYAS, Reichb. f. in Bot. Zeit. x. (1852) 834. *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
densiflora, *Reichb. f. in Gard. Chron.* (1875) i. 492.—Peruv.
Mandonii, *Reichb. f. Xenia Orch.* iii. 21.—Bolivia.
rhodoneura, *Reichb. f. in Bot. Zeit.* x. (1852) 834.—Bolivia.

NEOGAYA, Meissn. Gen. 144 (1838) = **Ligusticum**, Linn. (Umbell.).
mucronata, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 80.—Sibir.
simplex, Meissn. Gen. Comm. 104 = L. simplex.

NEOGYNA, Reichb. f. in Bot. Zeit. x. (1852) 931 = **Coelogyne**, Lindl. (Orchid.).
Gardneriana, Reichb. f. l. c. = C. Gardneriana.

NEOHALLIA, Hemsl. Biol. Centr. Am. Bot. ii. 519 (1882). *ACANTHACEAE*.
Borrerae, *Hemsl. l. c.*—Mexic.

NEOLACIS, Wedd. in DC. Prod. xvii. 59 (1873) = **Apinagia**, Tul. (Podost.).
capillacea, Wedd. l. c. 63.—Bras.
corymbosa, Wedd. l. c. 60 = Ligea corymbosa.
dissecta, Wedd. l. c. 62 = Ligea dissecta.
divaricata, Wedd. l. c. 60 = A. divaricata.
fucoides, Wedd. l. c. 61 = A. fucoides.
Gardneriana, Wedd. l. c. 61 = A. Gardneriana.
membranacea, Wedd. l. c. 64 = A. membranacea.
myriophylla, Wedd. l. c. 63.—Costa Rica.
nitelloides, Wedd. l. c.—Bras.
peruviana, Wedd. l. c. 64 = A. striata.
psyllophora, Wedd. l. c. 60 = A. psyllophora.
pusilla, Wedd. l. c. 63 = A. pusilla.
pygmaea, Wedd. l. c. 60 = A. pygmaea.
Richardiana, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96 = Ligea Richardiana.
Riedelii, Wedd. in DC. Prod. xvii. 61 = A. Riedelii.
ruppioides, Wedd. l. c. = A. ruppioides.
secundiflora, Wedd. l. c. 62 = Ligea secundiflora.
Spruceana, Wedd. l. c. 64.—Bras.

NEOLEXIS, Salisb. Gen. Pl. Fragm. 64 (1866) = **Smilacina**, Desf. (Liliac.).

NEOROEPERA, Muell. Arg. in DC. Prod. xv. II. 488 (1866). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 276.
Banksii, *Benth. Fl. Austral.* vi. 117.—Austral.
buxifolia, *Muell. Arg. in DC. Prod.* xv. II. 489.—Austral.

NEORTHOSIS, Rafin. Fl. Tellur. iv. 125 (1836) = **Ipomoea**, Linn. (Convolv.).
coccinea, Rafin. l. c. = I. coccinea.
tigrina, Rafin. l. c. = I. Pes-tigridis.

NEOSPARTON, Griseb. in Goett. Abh. xix. (1874) 245 (Pl. Lorentz. 197). *VERBENACEAE*, Benth. & Hook. f. ii. 1144.
Darwinii, *Benth. & Hook. f. Gen.* ii. 1144.—Bras.
ephedroides, *Griseb. in Goett. Abh.* xix. (1874) 245.—Reg. Argent.

NEOTIA, Scop. Ann. IV. Hist. Nat. 99 (1770) = **Neottia**, Linn. (Orchid.).

NEOTINEA, Reichb. f. Poll. Orch. Comm. 18, 29 (1852) = **Habenaria**, Willd. (Orchid.).
intacta, Reichb. f. l. c. = H. intacta.

NEOTTIA, Linn. Syst. ed. I (1735). *ORCHIDEAE*, Benth. & Hook. f. iii. 595.
NEOTTIDIUM, Schlecht. Fl. Berol. i. 454 (1823).
SYNOPLECTRIS, Rafin. Fl. Tellur. ii. 89 (1836).
abortiva, [Clairv]. Man. Herb. 264 = Limodorum abortivum.
abortiva, S. F. Gray, Nat. Arr. Brit. Pl. ii. 209 = N. Nidus-avis.
acaulis, Sm. Exot. Bot. ii. 91 = Spiranthes picta.
Adnaria, Rafin. Fl. Tellur. ii. 87 = Pelexia adnata.
adnata, Sw. Fl. Ind. Occ. iii. 1409 = Pelexia adnata.
aestivalis, Lam. Encyc. iv. 567; DC. Fl. Fr. iii. 258 = Spiranthes aestivalis.
aestivalis, Oakes, Cat. Pl. Verm. 28; et ex A. Gray, Fl. N. U. St. ed. V. 504 = Spiranthes latifolia.
alsinefolia, Reinw. ex Blume, Orch. Archip. Ind. 77 = Myrmechis gracilis.
amoena, Bieb. Fl. Taur. Cauc. iii. 606 = Spiranthes australis.
amphylla, Hook. Bot. Mag. t. 2797 = Spiranthes orchioides.
aurantiaca, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 2 = Spiranthes aurantiaca.
australis, R. Br. Prod. 319 = Spiranthes australis.
australis, Reichb. f. in Walp. Ann. iii. 595 = Listera australis.
autumnalis, Pers. Syn. ii. 510 = Spiranthes autumnalis.
Banksiana, Reichb. f. in Walp. Ann. iii. 595 = Listera Eschscholziana.
bicolor, Ker.-Gawl. in Bot. Reg. t. 794 = Spiranthes bicolor.
bifida, Blume, Bijdr. 408 = Goodyera bifida.

NEOTTIA:—
bracteosa, Steud. Nom. ed. II. ii. 189 = Spiranthes bracteosa.
calcarata, Sw. Fl. Ind. Occ. 1413 = Pelexia setacea.
camtschatea, Reichb. f. Ic. Fl. Germ. xiii., xiv. 146 = seq.
camtschatica, *Spreng. Syst.* iii. 707.—Kamtschat.
cernua, Sw. in Schrad. Neues Journ. i. (1805) 52; Willd. Sp. Pl. iv. 75 = Spiranthes cernua.
cinnabarina, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.). 3 = Spiranthes cinnabarina.
convallarioides, Rich. in Mém. Mus. Par. iv. (1818) 59 = Listera convallarioides.
cordata, Rich. l. c. = Listera cordata.
crispata, Blume, Bijdr. 406 = Spiranthes australis.
discolor, Steud. Nom. ed. II. ii. 189 = Haemaria discolor.
diuretica, Willd. Sp. Pl. iv. 73 = Spiranthes diuretica.
elata, Sw. in Schrad. Neues Journ. i. (1805) 51 = Spiranthes elata.
Eschscholziana, Steud. Nom. ed. II. ii. 189 = Listera Eschscholziana.
flava, Sw. Fl. Ind. Occ. iii. 1417 = Corymbis flava.
flexuosa, Sm. in Rees, Cycl. xxiv. n. 9 = Spiranthes australis.
gemmipara, Sm. Engl. Fl. iv. 36 = Spiranthes Romanzoffiana.
glandulosa, Sims, Bot. Mag. t. 842 = Ponthieva glandulosa.
gracilis, Bigel. Fl. Bost. ed. III. 345 = Spiranthes gracilis.
grandiflora, Hook. Bot. Mag. tt. 2730, 2956 = Spiranthes grandiflora.
kamtschatica, Lindl. Gen. et Sp. Orch. 458 = camtschatica.
lanceolata, Willd. Sp. Pl. iv. 73 = *Stenorhynchus lanceolatus*).—Guian.
latifolia, Rich. in Mém. Mus. Par. iv. (1818) 59 = Listera ovata.
Lindleyana, Decne. in Jacquem. Voy. Bot. 163. t. 163 = listeroides.
listeroides, *Lindl. in Royle, Illustr. Bot. Himal.* 368. —Reg. Himal.
macrantha, Sweet, Hort. Brit. ed. II. 485 = Spiranthes grandiflora.
macrophylla, D. Don, Prod. Fl. Nep. 27 = Herminium congestum.
macrostelis, Peterm. in Flora, xxvii. (1844) 369 = N. Nidus-avis.
maculata, *Linden, Illustr. Hortic.* xxix. (1812) 83.—N. Granat.
Michuacana, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 3 = Spiranthes Mechoacana.
micrantha, G. Don, in Sweet. Hort. Brit. ed. III. 654, sphalm. = *macrantha*.
micrantha, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.). 5 = Spiranthes Llaveana.
micrantha, *Lindl. Gen. et Sp. Orch.* 458.—Sibir.
minor, Jacq. Coll. iii. 172 = Spiranthes elata.
monophylla, D. Don, Prod. Fl. Nep. 27 = Herminium gramineum.
Nidus-avis, *Rich. in Mém. Mus. Par.* iv. (1818) 59.—Europ.; As. bor.
odorata, Nutt. in Journ. Acad. Philad. vii. (1834) 98 = Spiranthes odorata.
orchioides, Sw. Fl. Ind. Occ. iii. 1411. t. 28 = Spiranthes orchioides.
orobanchoidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 61, 70, 130 = N. Nidus-avis.
ovata, Bluff & Fingerh. Comp. Fl. Germ. ed. II. ii. 435 = Listera ovata.
pantanis, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 173 = Pachystoma pantanum.
papulosa, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 4.—Mexic.
parviflora, Sm. in Rees, Cycl. xxiv. n. 10 = Spiranthes australis.
Petola, Blume, Bijdr. 407. t. 2 = Macodes Petola.
picta, Sims, Bot. Mag. t. 1562; [R. Br. in] Ait. Hort. Kew. ed. II. v. 199 = Spiranthes picta.
plantaginea, D. Don, ex Spreng. Syst. iii. 708 = Spiranthes plantaginea.
plantaginea, Hook. Exot. Fl. t. 226 = Spiranthes orchioides.
polystachya, Sw. Fl. Ind. Occ. iii. 1415 (= *Chloidia vernalis*).—Jamaic.
procera, Ker.-Gawl. in Bot. Reg. t. 639 = Goodyera procera.

NEOTTIA:—
pubera, Steud. Nom. ed. II. ii. 189 = Ponthieva glandulosa.
pubescens, Willd. Sp. Pl. iv. 76 = Goodyera pubescens.
pudica, Sweet, Hort. Brit. ed. II. 485 = Spiranthes australis.
quadridentata, Willd. Sp. Pl. iv. 73 = Spiranthes quadridentata.
reniformis, Spreng. Syst. iii. 707 (= Habenariae sp.). —Ind. or.
repens, Pursh, Fl. Am. Sept. ii. 589 = Goodyera pubescens.
repens, Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 226 = Goodyera repens.
rubicunda, Blume. Bijdr. 408 = Goodyera rubicunda.
Salassia, Steud. Nom. ed. II. ii. 189 = Liparis purpurascens.
sinensis, Pers. Syn. ii. 510 = Spiranthes australis.
speciosa, Jacq. Ic.Pl.Rar. iii. t. 600 = Spiranthes colorata.
spiralis, Sw. in Schrad. Neues Journ. i. (1805) 51 = Spiranthes aestivalis, autumnalis.
squamosa, Dulac, Fl. Hautes-Pyr. 120 = N. Nidus-avis.
squamulosa, H. B. & K. Nov. Gen. et Sp. i. 332. t. 71 = Spiranthes orchioides.
strateumatica, R. Br. Prod. 319 = Zeuxine sulcata.
sulphurea, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.). 4 = Spiranthes sulphurea.
tortilis, Pursh, Fl. Am. Sept. ii. 589 = Spiranthes gracilis.
tortilis, Sw. in Schrad. Neues Journ. i. (1805) 51; Fl. Ind. Occ. iii. 1406 = Spiranthes tortilis.
vaginata, H. B. & K. Nov. Gen. et Sp. i. 331 (= *Stenorrhynchus vaginatus*).—Peruv.
viridiflora, Blume, Bijdr. 408 = Goodyera viridiflora.

NEOTTIDIUM, Schlecht. Fl. Berol. i. p. lxv. 454 (1823) = **Neottia**, Linn. (Orchid.).
Nidus-avis, Schlecht. l. c. 454 = Neottia Nidus-avis.

NEOU, Adans. ex Juss. Gen. 342 (1789) = **Parinarium**, Juss. (Rosac.).

NEOWEDIA, Schrad. in Neuwied, Reise Bras. ii. 343 (1821) = **Ruellia**, Linn. (Acanthac.).
affinis, Schrad. in Goett. Gel. Anz. i. (1821) 706 = R. speciosa.
Neesiana, Mart. ex Nees, in Mart. Fl. Bras. ix. 37 (= *Dipteracanthus Neesianus*.).—Bras.
speciosa, Mart. ex Nees, l. c. 30 = R. speciosa.
speciosa, Schrad. in Neuwied, Reise Bras. ii. 343 (= *Dipteracanthus Neowedia*).—Bras.

NEPA, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 286 = **Ulex**, Linn. (Legumin.).
aragonensis, Losc. ex Willk. & Lange, Prod. Fl. Hisp. iii. 427 = Genista hispanica.
Boivini, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 287 = U. Boivini.
Cossonii, Webb, l. c. = U. Cossonii.
Escayracii, Webb, l. c. = U. Escayracii.
lurida, Webb, l. c. 286 = U. Webbianus.
megalorites, Webb, l. c. 287 = U. Boivini.
Salzmanni, Webb, l. c. = U. Boivini.
Vaillantii, Webb, l. c. = U. Boivini.
Webbiana, Webb, l. c. 286 = U. Webbianus.

NEPENTHES, Linn. Syst. ed. I (1735); Gen. ed. I 273 (1737). *NEPENTHACEAE*, Benth. & Hook. f. iii. 116.
BANDURA, Burm. Thes. Zeyl. t. 17 (1737).
PHYLLAMORPHA, Lour. Fl. Cochinch. 606 (1790).
alata, *Blanco, Fl. Filip.* ed. I. 805.—Ins. Philipp.
albo-marginata, *Lobb, ex Lindl. in Gard. Chron.* (1849) 580.—Singapore.
ampullacea, Jack, in Wall. Cat. n. 2243 = ampullaria.
ampullaria, *Jack, Desc. Malay. Pl.* iii. [1823?] 23.—Malaya.
angustifolia, *Mast. in Gard. Chron.* (1881) II. 524.—Malaya.
atro-sanguinea ×, Hort.—Cf. Gard. Chron. (1882) I. 826. f. 125.
Bernaysii, *F. M. Bailey, in Proc. Linn. Soc. N. S. W.* v. (1881) 186.—Austral.
bicalcarata, *Hook. f. in DC. Prod.* xvii. 97.—Borneo.
Blancoi, *Blume, Mus. Bot. Lugd. Bat.* ii. 10.—Ins. Philipp.
Bongso, *Korth. Verh. Nat. Gesch. Bot.* 19. t. 14.—Sumatra.
Boschiana, *Korth. l. c.* 25. t. 2.—Borneo.

NEPENTHES:—
Burbidgei, Hook. f. ex Burb. in Gard. Chron. (1882) I. 56 = N. Phyllamphora.
celebica, *Hook. f. in DC. Prod.* xvii. 100.—Ins. Celebes.
Chelsoni ×, Hort. Veitch, ex Masters, in Gard. Chron. (1872) 542.
cincta, *Mast. l. c.* (1884) I. 576.—Borneo.
coccinea, *Hort.*—Cf. Gard. Chron. (1882) II. 169. f. 29. —Hab.?
Courtii ×, Hort. Veitch.—Cf. l. c. (1881) II. 844. f. 160.
cristata, *Brongn. in Ann. Sc. Nat.* Sér. I. i. (1824) 48. —Madag.
distillatoria, Auct. ex Steud. Nom. ed. II. ii. 190 = N. Phyllamphora.
distillatoria, Brion, in Belg. Hortic. v. (1855) 196 = cristata?
distillatoria, R. Grah. in Edinb. N. Phil. Journ. (Jul.–Sept. 1827) 371; et in Bot. Mag. t. 2798 = khasiana.
distillatoria, *Linn. Sp. Pl.* 955.—Zeylan.
Dominii ×, Hort. Veitch. ex Mast. in Gard. Chron. (1872) 542.
Dormanniana ×, Hort.—Cf. Gard. Chron. (1882) I. 525. f. 81.
Dyak, S. Moore, in Journ. Bot. xviii. (1880) 1 = bicalcarata.
echinostoma, *Hook. f. in DC. Prod.* xvii. 95.—Borneo.
Edwardsiana, *Low, ex Hook. f. in Trans. Linn. Soc.* xx. (1851) 420.—Borneo.
eustachys, *Miq. Pl. Ind. Bat.* i. I. 1074.—Sumatra.
fimbriata, Blume, Mus. Bot. Lugd. Bat. ii. 7 = N. Phyllamphora.
gracilis, *Korth. Verh. Nat. Gesch. Bot.* 22. t. 1.—Borneo.
gymnamphora, *Miq. Pl. Jungh.* 169.—Java.
gymnamphora, Reinw. ex Nees, in Ann. Sc. Nat. Sér. I. iii. (1824) 366 = melamphora.
Harryana ×, Burb. in Gard. Chron. (1882) I. 56.—Borneo.
hirsuta, *Hook. f. in DC. Prod.* xvii. 29.—Borneo.
Hookeri, Alphand, ex Hook. f. l. c. 96 = Rafflesiana.
Hookeriana, *Low, Sarawak*, 68; *ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 593, nomen.—Borneo.
hybrida ×, Hort. Veitch. ex Mast. in Gard. Chron. (1872) 542.
indica, Poir. in Lam. Encyc. iv. 458 = distillatoria.
intermedia ×, Hort. ex Veitch. in Gard. Chron. (1875) I. 551; (1882) I. 178. f. 29.
Kennedyana, *F. Muell. Fragm.* v. 154.—Austral.
Kennedyi, Benth. Fl. Austral. vi. 40 = praec.
khasiana, *Hook. f. in DC. Prod.* xvii. 102.—Reg. Himal.
Korthalsiana, Miq. Fl. Ind. Bat. i. I. 1071 = gracilis.
laevis, Korth. ex Hook. f. in DC. Prod. xvii. 104 = gracilis.
laevis, *Lindl. in Gard. Chron.* (1848) 655.—Malaya.
laevis, C. Morr. in Belg. Hortic. ii. (1852) 234 = Teysmanniana.
lanata, *Hort. ex Linden, Illustr. Hortic.* xxiii. (1876) 192 [t. 261].—Borneo.
Lawrenciana ×, Hort.—Cf. Gard. Chron. (1880) II. 40. f. 8.
Lindleyana, *Low, ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 593, nomen.—Borneo.
Loddigesii, *W. Baxt. l. c.*—Borneo.
Lowii, *Hook. f. in Trans. Linn. Soc.* xii. (1859) 420. —Borneo.
macrostachya, Blume, Mus. Bot. Lugd. Bat. ii. 7 = N. Phyllamphora.
madagascariensis, *Poir. in Lam. Encyc.* iv. 459.—Madag.
Mastersiana ×, Hort.—Cf. Gard. Chron. (1881) II. 749. f. 148.
maxima, *Reinw. in Ann. Sc. Nat.* Sér. I. iii. (1824) 369. t. 20.—Ins. Celebes.
melamphora, *Reinw. ex Blume, Cat. Gew. Buitenz.* 111.—Java.
Morganiana ×, Hort.—Cf. Gard. Chron. (1881) II. 381.
nigropurpurea, Hort. l. c. (1882) I. 429. f. 70.
Northiana, *Hook. f. l. c.* (1881) II. 717.—Borneo.
Outramiana ×, Hort.—Cf. l. c. (1879) II. 505. f. 10.
Pervillei, Blume, Mus. Bot. Lugd. Bat. ii. 10.—Ins. Seychell.
Phyllamphora, Hook. f. & Thoms. ex Hook. f. in DC. Prod. xvii. 102 = khasiana.
Phyllamphora, Reinw. ex Miq. Fl. Ind. Bat. i. I. 1073 = melamphora.

NEPENTHES:—
Phyllamphora, *Willd. Sp. Pl.* ii. 874.—Cochinch.; Ins. Molucc.
pumila, *Griff. Notul.* iv. 349.—Malaya.
Rafflesiana, *Jack, in Malay. Misc. App. Ined.* [1823?] 21; *et in Hook. Comp. Bot. Mag.* i. (1835) 271.—Ind. or.
Rajah, *Hook. f. in Trans. Linn. Soc.* xxii. (1859) 421.—Borneo.
Ratcliffiana ×, Hort.—Cf. Gard. Chron. (1882) I. 178. f. 28.
Reinwardtiana, *Miq. Pl. Jungh.* 168.—Malaya.
robusta ×, Hort.—Cf. Gard. Chron. (1880) II. 40. f. 11.
rubra, Hort. ex Rafarin, in Rév. Hortic. (1869) 270 = khasyana.
rubro-maculata ×, Hort. Veitch, ex Mast. in Gard. Chron. (1882) I. 143. f. 24.
sanguinea, *Lindl. in Gard. Chron.* (1849) 580.—Malaya.
Sedeni ×, Hort.—Cf. Flor. & Pomel. (1872) 54; Mast. in Gard. Chron. (1872) 542.
superba ×, Hort. ex Flor. Mag. N. S. (1881) t. 434.
tentaculata, *Hook. f. in DC. Prod.* xvii. 101.—Borneo.
Teysmanniana, *Miq. Fl. Ind. Bat.* i. 1. 1073.—Malaya.
tomentella, Miq. l. c. 1075 = albo-marginata.
trichocarpa, *Miq. l. c.* 1072.—Sumatra.
Veitchii, *Hook. f. in Trans. Linn. Soc.* xxii. (1859) 421.—Borneo.
ventricosa, *Blanco, Fl. Filip.* ed. I. 807.—Ins. Philipp.
Vieillardi, *Hook. f. in DC. Prod.* xvii. 104.—N. Caled.
villosa, Hook. Bot. Mag. t. 5080 = Veitchii.
villosa, *Hook. f. in Hook. Ic. Pl.* t. 888.—Borneo.
Wardii, E. P. Wright, in Trans. Roy. Irish Acad. xxiv. (1869) 576. tt. 29, 30 = Pervillei.
Williamsii ×, Hort. ex Gard. Chron. (1880) II. 40. f. 9.
Wrigleyana ×, Hort. Veitch, ex Mast. l. c. (1882) I. 143. f. 23.
zeylanica, Rafin. Fl. Tellur. iv. 101 = distillatoria.

NEPETA, Riv. ex Linn. Gen. ed. I. 170 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1199.
CATARIA, Adans. Fam. ii. 192 (1763).
CHAMAECLEMA, Moench, Meth. 393 (1794).
GLECHOMA, Linn. Syst. ed. I (1735).
LOPHANTHUS, Adans. Fam. ii. 194 (1763).
MARMORITES, Benth. in Hook. Bot. Misc. iii. (1833) 377.
OXYNEPETA, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxi. (1873) n. 1 (Lab. Pers.) 58.
SACCILABIUM, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 294.
SAUSSURIA, Moench, Meth. 388 (1794).
abyssinica, *Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* xi. 77.—Abyssin.
acerosa, Webb, Iter Hisp. 20 = Apuleii.
acinifolia, Spreng. Pugill. i. 41 = cyanea.
agrestis, *Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 417.—Corsica; Hispan.
alba, Steud. Nom. ed. I. 552 = nuda.
algeriensis, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 581.—Alger.
altissima, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 54 = Lophanthus nepetoides.
amboinica, Linn. f. Suppl. 273 = Anisomelis ovata.
americana, Vitm. Summa Pl. iii. 419 = N. Cataria?
amethystina, *Desf. ex Poir. Encyc. Suppl.* iii. 206.—Hispan.
amoena, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 45.—Persia.
andrica, Boiss. & Heldr. ex Benth. in DC. Prod. xii. 389 = cilicica.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3; Vahl, Symb. Bot. i. 41 = N. Nepetella.
annua, Pall. in Act. Acad. Petrop. ii. (1779) 263. t. 12 = botryoides.
Apuleii, *Ucria, in Roem. Arch.* i. 1. 69.—Reg. Mediterr. occ.
aragonensis, Lam. Encyc. i. 711 = N. Nepetella.
argentea, Hort. ex Benth. Lab. Gen. et Sp. 480 = grandiflora, Mussini.
argolica, Bory & Chaub. Exp. Mor. 36. n. 889. t. 16 = Sibthorpii.
aristata, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 656.—Armenia.
aristata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Hyptis pectinata.
arragonensis, Benth. Lab. Gen. et Sp. 478 = N. Nepetella.

NEPETA:—
atlantica, *Ball, in Journ. Bot.* xiii. (1875) 175.—Marocc.
Aucheri, *Boiss. in Tchihat. Asie Min. Bot.* ii. 155, nomen.—Armenia.
austriaca, *Host, Fl. Austr.* ii. 154.—Europ.
azurea, *R. Br. in Salt, Abyss. App.* 64, nomen; *Benth. Lab. Gen. et Sp.* 479.—Abyssin.
Balbisii, *Colla, Herb. Pedem.* iv. 442. t. 85 (Quid?).—Italia.
ballotaefolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 196.—Abyssin.
barbata, *Regel & Winkl. in Act. Hort. Petrop.* vi. (1879) 361.—Turkestan.
basilicoides, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 294.—Guian.
betonicaefolia, *C. A. Mey. Verz. Pfl. Cauc.* 92.—Reg. Caucas.
betonicoides, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 45.—Persia.
biloba, *Hochst. ex Benth. in DC. Prod.* xii. 375.—Abyssin.
bipinnata, Cav. Ic. i. 36. t. 49 = botryoides.
bipinnatifida, Ledeb. Fl. Ross. iii. 372 = botryoides.
Bodeana, *Bunge, Lab. Pers.* 53.—Persia.
Boissieri, Willk. in Bot. Zeit. xv. (1857) 219 = amethystina.
bombaiensis, *Dalz. in Dalz. & Gibs. Bomb. Fl.* 209.—Ind. or.
botryoides, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 287.—Reg. Himal.; Soongar.
bracteata, *Benth. in DC. Prod.* xii. 395.—Persia; Beluchist.
bracteata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 114 = Dysophylla crassicaulis.
brevifolia, *C. A. Mey. Verz. Pfl. Cauc.* 93.—Reg. Caucas.
cadmea, *Boiss. Diagn.* Ser. I. v. 21.—As. Min.
caesarea, *Boiss. Fl. Orient.* iv. 655.—Cappadoc.
Calamintha, *Tausch, in Flora*, xxiv. (1831) 219.—Hab.?
calaminthoides, Benth. in DC. Prod. xii. 382 = ruderalis.
Calverti, *Boiss. Diagn.* Ser. II. iv. 25.—Armen.
calycina, *Fenzl, in Flora*, xxvi. (1843) 400.—Mesopot.
campestris, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 97, nomen.—Rossia.
campestris, *Benth. Lab. Gen. et Sp.* 734.—Reg. Himal.
campestris, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 658 = nervosa.
camphorata, *Boiss. & Heldr. Diagn.* Ser. I. vii. 49.—Graecia.
canescens, J. F. Gmel. Syst. 900 = italica.
Cataria, *Linn. Sp. Pl.* 570.—Europ.; Oriens; Himal.
Cephalotes, *Boiss. Diagn.* Ser. I. v. 20.—Persia.
chenopodiifolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 46, 47.—Persia.
chionophila, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 643.—Persia.
ciliaris, *Benth. in Wall. Pl. As. Rar.* i. 64.—Reg. Himal.
cilicica, *Boiss. ex Benth. in DC. Prod.* xii. 368.—As. Min.; Syria.
circinnata, Willd. Enum. Hort. Berol. 602 = Stachys nepetaefolia.
citriodora, Dum. Fl. Belg. 47; Becker, Fl. Frankf. i. 218 = N. Cataria.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 663.—Reg. Himal.
clinopodioides, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 379 = ruderalis.
coerulea, [Soland. in] Ait. Hort. Kew. ed. I. ii. 285 = latifolia.
coerulescens, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 529.—China.
colorata, Willd. Enum. Hort. Berol. 601 = grandiflora.
concolor, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 386.—As. Min.
congesta, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 34 = curvidens.
connata, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 378.—Reg. Himal.
cordifolia, C. Koch, ex Boiss. Fl. Orient. iv. 658 = glabrescens.
crassifolia, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 175 = Mussini.
cretica, F. G. Dietr. Vollst. Lexik. Gaertn. vi. 348 = Micromeria marifolia.

NEPETA:—
crinita, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 46.—As. Min.
crispa, *Willd. Sp. Pl.* iii. 50.—Persia.
croatica, Ledeb. Fl. Ross. iii. 351 = Micromeria marifolia.
croatica, Spreng. Syst. ii. 727 = Micromeria rupestris.
cryptantha, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 669.—Syria; Persia.
curvidens, *Boiss. & Bal. Diagn.* Ser. II. iv. 26.—As. Min.
curviflora, *Boiss. l. c.* Ser. I. v. 22.—Syria.
cyanea, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 265.—Reg. Caucas.
cyanotricha, Tausch, in Flora, xiv. (1831) 218 = Mussini.
daënensis, *Boiss. Diagn.* Ser. I. vii. 53.—Persia.
dasyangula, *Lenné & C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 12.—Hab.?
delphinensis, Mutel, Fl. Fr. iii. 46 = N. Nepetella.
densiflora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 725.—Soongaria.
denudata, *Benth. in DC. Prod.* xii. 390.—Persia.
depauperata, *Benth. l. c.* 392.—Persia.
diffusa, Fisch. ex Spreng. Syst. ii. 729 = Mussini.
dirphya, Heldr. ex Boiss. Fl. Orient. iv. 652 = Sibthorpii.
discolor, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 378.—Reg. Himal.
distans, *Royle, l. c.* 379.—Reg. Himal.
disticha, Blume, Bijdr. 823 = Anisomeles ovata.
elata, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 378 = spicata.
elliptica, *Royle, l. c.* 378.—Reg. Himal.
erecta, *Benth. Lab. Gen. et Sp.* 482.—Reg. Himal.
eriostachys, *Benth. l. c.* 734.—Reg. Himal.
erodiifolia, Boiss. Diagn. Ser. I. v. 24 = Lallemantia Royleana.
Everandi, *S. Moore, in Journ. Bot.* xvi. (1878) 135.—China.
fissa, Benth. Lab. Gen. et Sp. 471 = supina.
fissa, C. A. Mey. Verz. Pfl. Cauc. 33 = teucriifolia.
floccosa, *Benth. Lab. Gen. et Sp.* 736.—Reg. Himal.
foliosa, *Moris, Stirp. Sard. Elench.* iii. 10.—*Cf. Linnaea*, v. (1830) *Litt.* 97.—Sardin.
frutescens, *Link, Enum. Hort. Berol.* ii. 103, nomen.—Ins. Cap. Virid.
glabra, Benth. Lab. Gen. et Sp. 483 = teucriifolia.
glabrescens, *Boiss. Fl. Orient.* iv. 658.—As. Min.
Glechoma, *Benth. Lab. Gen. et Sp.* 485.—Europ.; As. bor.
globifera, *Bunge, Del. Sem. Hort. Dorpat.* (1859).—*Cf. Linnaea*, xxx. (1859–60) 752.—Persia.
globiflora, Bunge, l. c.; et Lab. Pers. 58 = praec.
glomerata, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 46.—Syria; Cappadoc.
glomerulosa, *Boiss. Diagn.* Ser. I. v. 21.—Persia.
glutinosa, *Benth. Lab. Gen. et Sp.* 735.—Reg. Himal.
Govaniana, *Benth. l. c.* 482.—Reg. Himal.
Goviana, Steud. Nom. ed. II. ii. 190, sphalm. = praec.
gracilescens, *Boiss. Fl. Orient.* iv. 647.—Armenia.
graciliflora, *Benth. in Wall. Pl. As. Rar.* i. 65.—Reg. Himal.
graciliflora, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 648 = leptantha.
gracilis, Salisb. Prod. 78 = botryoides.
granatensis, *Boiss. Elench.* 76.—Hispan.
grandiflora, *Bieb. Fl. Taur. Cauc.* ii. 42.—Reg. Caucas.
grandiflora, Lapeyr. Hist. Abr. Pl. Pyr. 329 = latifolia.
grandiflora, Mutel, Fl. Dauph. 343 = N. Nepetella.
grandis, Hort. ex Benth. Lab. Gen. et Sp. 480 = grandiflora.
grata, *Benth. l. c.* 735.—Reg. Himal.
graveolens, Vill. Hist. Pl. Dauph. ii. 366 = N. Nepetella.
hederacea, Trev. Prosp. Fl. Eug. 26 = N. Glechoma.
heliotropifolia, *Lam. Encyc.* i. 711.—Persia; Mesopot.
hirsuta, Jan, ex Nym. Consp. 586 = tuberosa.
hirsuta, Linn. Sp. Pl. 571 = N. Scordotis.
hispanica, *Boiss. & Reut. Diagn.* Ser. II. iv. 26.—Hispan.
humilis, *Benth. in DC. Prod.* xii. 382.—Persia.
humilis, Salisb. Prod. 78 = N. Nepetella.
hymenodonta, *Boiss. Fl. Orient.* iv. 665.—Persia.
imbricata, Lag. ex Spreng. Syst. ii. 729 = italica.
Imobus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 115 = Elsholtzia incisa.

NEPETA:—
incana, Bieb. Fl. Taur. Cauc. ii. 40 = cyanea.
incana, Hort. ex Benth. Lab. Gen. et Sp. 480 bis = Mussini, grandiflora.
incana, Sieber, ex Reichb. Fl. Germ. Excurs. 328 = Calamintha cretica.
incana, [Soland. in] Ait. Hort. Kew. ed. I. ii. 285 = N. Nepetella.
incana, Thunb. Fl. Jap. 244 = Caryopteris Mastachanthus.
incana, Willd. Sp. Pl. iii. 52 = italica.
indica, Burm. f. Fl. Ind. 126 = Leucas biflora.
indica, Linn. Sp. Pl. 571 = Anisomeles ovata.
intermedia, Lej. ex Reichb. Fl. Germ. Excurs. 329 = Calamintha officinalis.
isaurica, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 378.—As. Min.
ispahanica, *Boiss. Diagn.* Ser. I. v. 23.—Persia.
italica, *Linn. Sp. Pl.* 571.—Europ. austr.; Reg. Caucas.
italica, Sibth. & Sm. Fl. Graec. vi. t. 548 = Sibthorpii.
japonica, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 468.—Japon.
japonica, Willd. Sp. Pl. iii. 52 = Caryopteris Mastachanthus.
juncea, *Benth. in DC. Prod.* xii. 379.—Afghanist.
kokamirica, *Regel, in Act. Hort. Petrop.* vi. (1879) 358.—Turkestan.
kokanica, *Regel, Pl. Nov. Fedsch.* 65.—Turkestan.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 50.—Persia.
laconicea, Orph. ex Nym. Consp. 586 = parnassica.
lagopsis, *Benth. in DC. Prod.* xii. 379.—Afghan.
lamiifolia, Adam, ex Hoffm. Hort. Mosq. (1808) n. 2169 = grandiflora.
lamiifolia, Benth. in DC. Prod. xii. 390 = brevifolia.
lamiifolia, *Willd. Enum. Hort. Berol.* 602.—Armenia.
lamiopsis, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 659.—Reg. Himal.
lanata, Jacq. Obs. Bot. iii. 21 = tuberosa.
lanceolata, Kral. ex Nym. Consp. 586 = agrestis.
lanceolata, Lam. Encyc. i. 710 = N. Nepetella.
lasiocephala, *Benth. in DC. Prod.* xii. 380.—Persia.
latifolia, *DC. Fl. Fr.* iii. 528.—Mont. Pyren.
lavandulacea, *Linn. f. Suppl.* 272.—Sibir.
lavandulifolia, J. F. Gmel. Syst. 1590, sphalm. = praec.
laxa, Benth. Lab. Gen. et Sp. 483 = teucriifolia.
laxiflora, *Benth. in DC. Prod.* xii. 393.—Persia.
leptantha, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 648.—As. Min.
leptoclada, *Trautv. in Act. Hort. Petrop.* ii. (1873) 480.—Armenia.
leucolaena, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 662.—Reg. Himal.
leucophylla, *Benth. Lab. Gen. et Sp.* 476.—Reg. Himal.
leucostegia, *Boiss. & Heldr. Diagn.* Ser. I. xii. 62.—As. Min.; Syria.
linearis, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 377.—Reg. Himal.
lobata, Rudolph, in Schrad. Journ. ii. (1799) 287 = lavandulacea.
longibracteata, *Benth. Lab. Gen. et Sp.* 737.—Reg. Himal.
longicaulis, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 429 = N. Nepetella.
longiflora, Fisch. ex Benth. Lab. Gen. et Sp. 484 = brevifolia.
longiflora, Kotschy, ex Benth. in DC. Prod. xii. 388 = cilicica.
longiflora, Sims, Bot. Mag. t. 923 = Mussini.
longiflora, *Vent. Jard. Cels*, 66. t. 66.—Persia.
lophantha, Fisch. ex Benth. Lab. Gen. et Sp. 464 = Lophanthus chinensis.
lusitanica, *Rouy, in La Natural.* (1882); (*Matér. Rév. Fl. Portug.* 32).—Lusitan.
lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 99.—Lycia.
macrantha, *Fisch. Hort. Gorenk.* ed. II. (1812) 22.—Sibir. altaic.
macrantha, Franch & Sav. Enum. Pl. Jap. ii. 464 = subsessilis.
macrantha, Ledeb. ex Hook. f. Fl. Brit. Ind. iv. 663 = erecta.
macrosiphon, *Boiss. Diagn.* Ser. I. vii. 51.—Persia.
macrostachya, Jan, ex Benth. Lab. Gen. et Sp. 736 = italica.
macrura, Ledeb. ex Spreng. Syst. ii. 729 = N. Cataria.

NEPETA:—
madagascariensis, Lam. Encyc. i. 712 = Plectranthus rotundifolius.
malabarica, Linn. Mant. ii. 566 = Anisomeles malabarica.
mallophora, *Webb & Heldr. ex Nym. Consp.* 585.—Hispan.
manchuriensis, *S. Moore, in Journ. Bot.* xviii. (1880) 5.—Manchuria.
maracandia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1847) 434.—Turkestan.
Mariae, *Regel, in Act. Hort. Petrop.* vi. (1879) 359.—Turkestan.
marifolia, *Boiss. & Huet, Diagn.* Ser. II. iv. 24.—Armenia.
marifolia, Cav. in Anal. Cienc. Nat. ii. (1800) 192 = Micromeria marifolia.
marrubioides, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 387.—Graecia; As. Min.
marrubioides, Willd. Enum. Hort. Berol. 603 = italica.
meda, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 46.—Persia.
melissaefolia, *Lam. Encyc.* i. 711.—Ins. Creta; Melos.
melissaefolia, Pers. Syn. ii. 115 = grandiflora.
menthoides, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 174.—Persia.
Meyeri, Benth. Lab. Gen. et Sp. 478 = micrantha.
micrantha, *Bunge, in Ledeb. Fl. Alt.* ii. 401.—Oriens; Sibir.
microphylla, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 44.—Persia.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = N. Cataria.
mollis, *Benth. Lab. Gen. et Sp.* 734.—Reg. Himal.
mollis, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 218. t. 14 = violacea?
mollis, Salisb. Prod. 78 = N. Cataria.
mollissima, *Tausch, in Flora*, xiv. (1831) 219 = cyanea.
montana, Turra, Fl. Ital. Prod. 66 = nuda.
multibracteata, *Desf. Fl. Atlant.* ii. 11. t. 123.—Alger.
multibracteata, Webb, Iter Hisp. 20 = granatensis.
multifida, Linn. Sp. Pl. 572 = lavandulacea.
multifida, Linn. f. Suppl. 273 = botryoides.
murcica, Guir. ex Willk. in Bot. Zeit. xv. (1857) 218 = amethystina.
Mussini, *Spreng. ex Henck. Adumbr. Pl. Hort. Hal.* 15.—Reg. Caucas.; Persia.
mutabilis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Hyptis spicata.
nepalensis, *Spreng. Syst.* ii. 730.—Nepal.
Nepetella, Asso, Syn. Pl. Arag. 73; Boiss. Voy. Bot. Espagne, 503 = amethystina.
Nepetella, *Linn. Syst.* ed. X. 1096.—Europ. austr.
Nepetellae, *Forsk. Fl. Aegypt. Arab.* 212.—Arab.
nervosa, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 378.—Reg. Himal.
nivalis, *Benth. Lab. Gen. et Sp.* 737.—Reg. Himal.
nova, Bieb. ex Steud. Nom. ed. II. ii. 191 = Micromeria Pulegium.
nuda, *Linn. Sp. Pl.* 570.—Europ. austr.; Oriens.
obtusata, E. Mey. ex Benth. in DC. Prod. xii. 385 = Mussini.
obtusicrena, Boiss. Fl. Orient. iv. 647 = Calverti.
Olgae, *Regel, Pl. Nov. Fedsch.* 64.—Turkestan.
orientalis, Mill. Gard. Dict. ed. VIII. n. 10 = italica.
Orlichiana, *Lenné & Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 12.—Reg. Himal.
orphanidea, Boiss. Diagn. Ser. II. iv. 22 = parnassica.
ouroumitanensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 230.—Turkest.
oxyodonta, *Boiss. Diagn.* Ser. I. vii. 51.—Persia.
pallida, C. Koch, in Linnaea, xxi. (1848) 675 = micrantha.
pallida, Salisb. Prod. 78 = Anisomeles malabaria.
paniculata, Crantz, Stirp. Austr. ed. II. fasc. iv. 270 = nuda.
paniculata, Mill. Gard. Dict. ed. VIII. n. 4 = N. Nepetella.
pannonica, Linn. Sp. Pl. 570 = nuda.
parnassica, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. iv. 22.—Graecia.
parviflora, Bieb. Fl. Taur. Cauc. ii. 41 = ucranica.
parviflora, Unio Itin. ex Benth. in DC. Prod. xii. 393 = heliotropifolia.
parvifolia, Hort. ex Benth. Lab. Gen. et Sp. 383 = Micromeria marifolia.
patavina, Fisch. ex Steud. Nom. ed. I. 552 = Hyptis polyantha?

NEPETA:—
Patella, All. ex Loud. Hort. Brit. 234, sphalm. = N. Nepetella?
pauciflora, Spreng. in Schrad. Journ. (1800) II. 198 = Hyptis polyantha?
pectinata, Linn. Syst. ed. X. 1096 = Hyptis pectinata.
Perretii, *Colla, Herb. Pedem.* iv. 441. t. 85. f. 1.—Hab.?
persica, *Boiss. Diagn.* Ser. I. xii. 66.—Persia.
persica, Poit. ex Benth. in DC. Prod. xii. 396 = Hyptis pectinata.
petraea, *Benth. l. c.* 394.—Persia.
pinetorum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 183.—Afghan.
pinnatifida, Fisch. Hort. Gorenk. ed. II. (1812) 22 = Dracocephalum botryoides.
pisidica, Boiss. & Heldr. Diagn. Ser. I. xii. 65 = cilicica.
podostachys, *Benth. in DC. Prod.* xii. 372.—Afghanist.
polystachya, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 378 = elliptica.
pontica, C. Koch, in Linnaea, xxi. (1848) 676 = nuda.
procumbens, Mill. Gard. Dict. ed. VIII. n. 11 = N. Scorditis.
prostrata, *Benth. in DC. Prod.* xii. 380.—Persia.
pubescens, *Benth. l. c.* 373.—Afghan.
pumila, Spreng. Nov. Prov. 29 = Micromeria rupestris.
pungens, *Benth. Lab. Gen. et Sp.* 487.—Persia.
pusilla, Benth. l. c. 488 = pungens.
pycnantha, Boiss. ex Benth. in DC. Prod. xii. 388 = cilicica.
racemosa, *Lam. Encyc.* i. 711.—Armen.; Persia.
raphanorhiza, *Benth. Lab. Gen. et Sp.* 734.—Reg. Himal.
rarifolia, C. Koch, in Linnaea, xxi. (1848) 676 = teucriifolia.
Reichenbachiana, Fisch. & Mey. Ind. Sem. Hort. Petrop. viii. 68 = racemosa.
reticulata, *Desf. Fl. Atlant.* ii. t. 124.—Alger.
robusta, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 212.—Afr. trop.
rosea, Salzm. ex Benth. Lab. Gen. et Sp. 471 = Apuleii.
rotundifolia, Benth. in DC. Prod. xii. 392 = floccosa.
ruderalis, Boiss. Fl. Orient. iv. 643 = N. Cataria.
ruderalis, *Buch.-Ham. ex Benth. in Wall. Pl. As. Rar.* i. 64.—Europ.; Oriens.
rugosa, *Benth. in DC. Prod.* xii. 384.—Arab.
Ruprechti, *Boiss. Fl. Orient.* iv. 661.—Reg. Caucas.
Sabinei, J. A. Schmidt, in Journ. Bot. vi. (1868) 228, 238 = discolor.
saccharata, *Bunge, Lab. Pers.* 56.—Persia.
salviaefolia, Pers. Syn. ii. 115 = Mussini.
salviaefolia, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 379.—Reg. Himal.
satureioides, *Boiss. Diagn.* Ser. I. v. 24.—Persia; Afghan.
scabra, Tausch, in Flora, xiv. (1831) 218 = tuberosa.
scabridifolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 44.—Persia.
schiraziana, *Boiss. Diagn.* Ser. I. vii. 52.—Persia.
Schtschurowskiana, *Regel, Pl. Nov. Fedsch.* 63, 66.—Turkestan.
scordifolia, Sieber, ex Mutel, Fl. Fr. iii. 45 = agrestis.
scordiifolia, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 53 (Quid?).—Sibir.
scordioides, Fisch. ex Schrank, l. c. = praec.
Scordotis, *Linn. Cent. Pl.* ii. 20; *Amoen. Acad.* iv. 317.—Ins. Creta.
Scorodonia, Boiss. & Heldr. Diagn. Ser. I. xii. 63 = isaurica.
Scorodotis, Hill, Veg. Syst. xviii. 36 = N. Scordotis.
secunda, Wall. Cat. n. 2122 = ruderalis.
septemcrenata, Benth. in DC. Prod. xii. 390, partim = persica.
septemcrenata, *Ehrenb. ex Benth. Lab. Gen. et Sp.* 484.—Aegypt.; Arab.
serpyllifolia, Bieb. Fl. Taur. Cauc. ii. 40 = Micromeria serpyllifolia.
sessilifolia, *Bunge, Lab. Pers.* 54.—Persia.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 360.—Turkestan.
sibirica, Aschers. Fl. Brandenb. i. 520 = macrantha.
sibirica, Bieb. Fl. Taur. Cauc. ii. 41 = ucranica.
sibirica, *Linn. Sp. Pl.* 572.—Sibir.
Sibthorpii, *Benth. Lab. Gen. et Sp.* 474.—Graecia.
sidaefolia, J. F. Gmel. Syst. 898 = Hyptis polyantha?

NEPETA:—
spathulifera, *Benth. in DC. Prod.* xii. 380.—Afghan.
speciosa, *Boiss. & Noé, Diagn.* Ser. II. iv. 34.—Persia.
spicata, *Benth. in Wall. Pl. As. Rar.* i. 64.—Reg. Himal.
Spruneri, *Boiss. Diagn.* Ser. II. iv. 23.—Graecia.
stachyoides, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 65, nomen.—Alger.
stenantha, *Kotschy & Boiss. ex Boiss. Fl. Orient.* iv. 661.—Armenia.
suaveolens, Roem. ex Steud. Nom. ed. I. 552 = Hyptis suaveolens?
subhastata, *Regel, Pl. Nov. Fedsch.* 64.—Turkestan.
subincisa, *Benth. in DC. Prod.* xii. 373.—Afghan.
subsessilis, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 469.—Japon.
sulphurea, C. Koch, in Linnaea, xxi. (1848) 677 = nuda.
supina, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 265.—Reg. Cauc. et Himal.
syriaca, *Hasselq. Iter Palaest.* 479.—Syria.
tenuicaulis, Boiss. Fl. Orient. iv. 661 = marifolia.
tenuifolia, *Benth. Lab. Gen. et Sp.* 468.—China.
tenuifolia, Steud. Nom. ed. II. ii. 191, sphalm. = seq.
teucriifolia, *Willd. Enum. Hort. Berol.* 602.—As. Min.; Persia.
teucrioides, Hort. ex Benth. in DC. Prod. xii. 385 = grandiflora, Mussini.
teucrioides, Lam. Encyc. i. 711 = italica.
teydea, *Webb & Berth. Phyt. Canar.* iii. 85. t. 165.—Ins. Canar.
thibetica, *Benth. Lab. Gen. et Sp.* 737.—Reg. Himal.
Thomsoni, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 658.—Reg. Himal.
tmolea, *Boiss. Diagn.* Ser. I. v. 22.—As. Min.
tolypantha, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 98.—Lycia.
tomentosa, Sieber, ex Benth. Lab. Gen. et Sp. 473 = N. Scorditis.
tomentosa, Vitm. Summa Pl. iii. 419 = N. Cataria?
Trautvetteri, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 175.—Reg. Caucas.
trichocalyx, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 656 = aristata.
trichodonta, Boiss. & Noé, Diagn. Ser. II. iv. 25 = heliotropifolia.
tuberosa, Desf. Fl. Atlant. ii. 10 = Apuleii.
tuberosa, *Linn. Sp. Pl.* 571.—Reg. Mediterr. occ.
tymphestea, Heldr. & Sart. ex Boiss. Fl. Orient. iv. 654 = Spruneri.
ucranica, Bieb. Fl. Taur. Cauc. ii. 41 = nuda.
ucranica, *Linn. Sp. Pl.* 570.—Europ. occ.; As. bor.
Vaillantii, Benth. Lab. Gen. et Sp. 471 = tuberosa.
versicolor, Trev. in Nov. Act. Nat. Cur. xiii. (1826) 183 = Craniotome versicolor.
violacea, Brot. Fl. Lusit. i. 173 = multibracteata.
violacea, Hort. ex Tausch, in Flora, xiv. (1831) 220 = Mussini, Calamintha grandiflora.
violacea, Lapeyr. Hist. Abr. Pl. Pyr. 329 = latifolia.
violacea, *Linn. Sp. Pl.* 570.—Hispan.
violacea, Vill. Hist. Pl. Dauph. ii. 367 = nuda.
violaefolia, *Boiss. & Noé, Diagn.* Ser. II. iv. 27.—Kurdistan.
virgata, C. Koch, in Linnaea, xxi. (1848) 676 = teucriifolia.
virginica, *Linn. Sp. Pl.* 571.—Am. bor.
virginica, Willd. Sp. Pl. iii. 56 = Pycnanthemum lanceolatum.
viscida, *Boiss. Diagn.* Ser. I. v. 20.—As. Min.
viscosa, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 294.—Guian.
vulgaris, Lam. Fl. Fr. ii. 398 = N. Cataria.
Wildenowiana, Hort. ex Benth. Lab. Gen. et Sp. 480 = Mussini.
yesoënsis, *Franch. & Sav. Enum. Pl. Jap.* ii. 464.—Japon.

NEPETTA, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 218 = praec.

NEPHELAPHYLLUM, Blume, Bijdr. 372. t. 22 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 514.
CYTHERIS, Lindl. Gen. et Sp. Orch. 128 (1831).
cordifolium, *Lindl. in Journ. Linn. Soc.* iii. (1859) 23.—Reg. Himal.
pulchrum, *Blume, Bijdr.* 373.—Malaya.
scapigerum, *Hook. f. Bot. Mag.* t. 5390.—Borneo.
tenuiflorum, *Blume, Bijdr.* 373.—Malaya.

NEPHELEPHYLLUM, Blume, Fl. Jav. praef. p. vii (1828) = praec.

NEPHELIUM, Linn. Mant. i. 18 (1767). *SAPINDACEAE*, Benth. & Hook. f. i. 405.
CORVINIA, Stadtm. ex Willem. in Usteri, Ann. Bot. xviii. (1796) 28.
DIMOCARPUS, Lour. Fl. Cochinch. 233 (1790), partim.
DIPHEROCARPUS, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 501.
EUPHORIA, Comm. ex Juss. Gen. 248 (1789), partim.
LAETJI, Osb. ex Steud. Nom. ed. I. 460 (1821).
LITCHI, Sonner. Voy. Ind. iii. 255 (1782).
OTONEPHELIUM, Radlk. Sapind. Holland. Ind. Nachtr. 71 (1879).
PSEUDONEPHELIUM, Radlk. l. c. (1879).
SCYTALIA, Gaertn. Fruct. i. 197. t. 42 (1788).
acuminatum, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 164.—Borneo.
altissimum, *Teijsm. & Binn. Nederl. Kruidk. Arch.* iii. (1855) 410.—Java.
Beccarianum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.
Beckleri, *Benth. Fl. Austral.* i. 467.—Austral.
benghalense, G. Don, Gen. Syst. i. 670 = N. Longana.
bifoliolatum, Thw. Enum. Pl. Zeyl. 57 = Sapindus bifoliolatus.
capense, Baill. Hist. Pl. v. 351 = Pappaea capensis.
chryseum, *Blume, Rumphia*, iii. 105.—Borneo.
compressum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 98.—Borneo.
connatum, *F. Muell. ex Benth. Fl. Austral.* i. 465.—Austral.
coriaceum, *Benth. l. c.* 466.—Austral.
costatum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 688.—Malacca.
cuspidatum, *Blume, Rumphia*, iii. 110.—Borneo.
daedaleum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.
Danura, G. Don, Gen. Syst. i. 671 = Sapindus Danura.
Dimocarpus, *Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind.* i. 688 = N. Lit-chi.
distyle, *F. Muell. Fragm.* ix. 99.—Austral.
divaricatum, *F. Muell. ex Benth. Fl. Austral.* i. 467.—Austral.
diversifolium, *F. Muell. Fragm.* x. 82.—Austral.
duriocarpum, T. Anders. ex Hook. f. Fl. Brit. Ind. i. 688 = N. Lit-chi.
echinatum, Noronha, in Verh. Batav. Gen. v. (1798) ed. I. Art. IV. 21 = lappaceum.
erectum, Thw. Enum. Pl. Zeyl. 57 = Sapindus erectus.
eriopetalum, *Miq. Fl. Ind. Bat. Suppl.* 509.—Malacca.
eximium, Thw. Enum. Pl. Zeyl. 57 = Pometia eximia.
ferrugineum, *F. Muell. Papuan Pl.* i. 21.—N. Guin.
foveolatum, *F. Muell. ex Benth. Fl. Austral.* i. 466.—Austral.
fumatum, *Blume, Rumphia*, iii. 111.—Borneo.
fuscatum, Thw. Enum. Pl. Zeyl. 58, 408 = Glenniea unijuga.
Gardneri, *Thw. l. c.* 58.—Zeylan.
glabrum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 21; *Walp. Rep.* i. 420.—Java.
Griffithianum, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 303 = mutabile.
hamulatum, *Radlk. Sapind. Holland. Ind. Nachtr.* 78.—Malaya.
hypoleucum, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 50 = N. Longana.
informe, Cambess. in Mém. Mus. Par. xviii. (1829) 30 = Euphoria informis.
juglandifolium, *Blume, Rumphia*, iii. 106.—Java.
lappaceum, *Linn. Mant.* ii. 566.—Malaya.
laurinum, *Blume, Rumphia*, iii. 109.—Java.
Leichhardti, *F. Muell. App. Rep. Intercol. Exhib.* (1867) 25.—Austral.
leiocarpum, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 25.—Austral.
Lichi, Steud. Nom. ed. II. ii. 191 = seq.
Lit-chi, *Cambess. in Mém. Mus. Par.* xviii. (1829) 30.—China.
Longana, *Cambess. l. c.*—Ind. or.; Burma.
macrophyllum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.

NEPHELIUM:—
malaiense, *Griff. Notul.* iv. 549.—Malacca.
Mangayi, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 688.—Malacca.
melanomiscum, *Radlk. Sapind. Holland. Ind. Nachtr.* 74.—Malaya.
microphyllum, *Benth. Fl. Austral.* i. 468.—Austral.
Mora, Gardn. ex Thw. Enum. Pl. Zeyl. 58 = N. Longana.
multijugum, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 164 = Capura multijuga (in Addend.).
multinerve, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.
mutabile, *Blume, Rumphia*, iii. 104.—Borneo.
Noronhianum, Cambess. in Mém. Mus. Par. xviii. (1829) 30 = Xerospermum Noronhianum.
ophiodes, *Radlk. Sapind. Holland. Ind. Nachtr.* 78.—Malaya.
oppositifolium, *G. Don, Gen. Syst.* i. 671.—Ins. Molucc.
parviflorum, *G. Don, l. c.*—Ins. Molucc.
pinnatum, Cambess. in Mém. Mus. Par. xviii. (1829) 30 = Schmidelia pinnata.
pupillum, *Wight, Illustr. Ind. Bot.* i. 141 = N. Longana.
reticulatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.
rimosum, *G. Don, Gen. Syst.* i. 671.—Ind. or.
rubescens, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 688.—Malaya.
Rubia, *Walp. Rep.* v. 365.—Ind. or.
rubrum, G. Don, Gen. Syst. i. 671 = Sapindus attenuatus.
semicinereum, *F. Muell. Fragm.* iv. 158.—Austral.
simplicifolium, Thw. Enum. Pl. Zeyl. 57 = Sapindus Thwaitesii.
stipulaceum, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 39.—Ind. or.
subdentatum, *F. Muell. ex Benth. Fl. Austral.* i. 465.—Austral.
sufferugineum, *Radlk. Sapind. Holland. Ind. Nachtr.* 77.—Malaya.
tomentosum, *F. Muell. in Trans. Vict. Inst.* ii. (1858) 64.—Austral.
variabile, *Wall. ex Voigt, Hort. Suburb. Calc.* 95.—Reg. Himal.
verticillatum, G. Don, Gen. Syst. i. 670 = Sapindus Danura.
xanthioides, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 97.—Borneo.
xerocarpum, Cambess. in Mém. Mus. Par. xviii. (1829) 30 = Xerospermum Noronhianum.

NEPHELOCHLOA, Boiss. Diagn. Ser. I. v. 72 (1844). *GRAMINEAE*, Benth. & Hook. f. iii. 1195.
altaica, Griseb. in Ledeb. Fl. Ross. iv. 367 = Poa soongarica.
breviglumis, Trautv. in Act. Hort. Petrop. vii. (1880) 523 = Sphenopus Gouani.
orientalis, *Boiss. Diagn.* Ser. I. v. 73.—As. Min.
persica, Griseb. in Ledeb. Fl. Ross. iv. 366 = Poa persica.
soongarica, Griseb. l. c. 367 = Poa soongarica.
tripolitana, Boiss. & Blanche, Diagn. Ser. II. iv. 133 = Poa persica.

NEPHRADENIA, Decne. in DC. Prod. viii. 604 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 771.
acerosa, *Decne. l. c.*—Bras.
asparagoides, *Fourn. in Mart. Fl. Bras.* vi. IV. 330.—Bras.
laurifolium, *Benth. & Hook. f. Gen.* ii. 756 (sphalm. linearifolium).—Bras.
linearis, *Benth. & Hook. f. l. c.* 756, 771.—Bras.
neriifolia, *Benth. & Hook. f. l. c.* 771.—Mexic.

NEPHRAEA, Hassk. Cat. Hort. Bog. Alt. 283 (1844) = NEPHREA, Noronha = **Pterocarpus**, Linn. (Legumin.).

NEPHRAËLES, Rafin. Fl. Tellur. iii. 70 (1836) = **Commelina**, Linn.
parviflora, Rafin. l. c. = C. pallida?

NEPHRANDRA, Willd. in Cothen. Disp. Veg. 8 (1790) = **Vitex**, Tourn. (Verben.).
dubia, Willd. l. c.; J. F. Gmel. Syst. 946 = V. umbrosa.

NEPHRANTHERA, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 145 = **Renanthera**, Lour. (Orchid.).
matutina, Hassk. l. c. = R. matutina.

NEPHREA, Noronha, in Verh. Batav. Gen. v. (1790) Art. IV. 3 = **Pterocarpus**, Linn. (Legumin.).

NEPHROCODIUM, Benth. & Hook. f. Gen. iii. 457 (1883), sphalm. = seq.

NEPHROCODUM, C. Muell. in Walp. Ann. vi. 40 (1861); Becc. Malesia, i. (1878) 242 = seq.

NEPHROCOELIUM, Turcz. in Bull. Soc. Nat. Mosc. (1853) I. 287 = **Burmannia**, Linn.
malaccensis, Turcz. l. c. = B. caelestis.

NEPHROIA, Lour. Fl. Cochinch. 565 (1790) = **Cocculus**, DC. (Menisp.).
sarmentosa, Lour. l. c. = C. Nephroia.

NEPHROICA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42 = **Cocculus**, DC. (Menisp.).
bantamensis, Miers, l. c. = C. bantamensis.
caudata, Miers, l. c. Ser. III. xix. (1867) 26 = C. diversifolia? *Miq.* (in Addend.).
cuneifolia, Miers, l. c. = C. cuneatus.
cynanchoides, Miers, l. c.; et in Hook. Kew Journ. iii. (1851) 259 = C. Thunbergii.
dianthera, Miers, l. c. Ser. II. vii. (1851) 42 = C. Thunbergii.
dilatata, Miers, l. c. Ser. III. xix. (1867) 26 = C. Thunbergii.
Ferrandiana, Miers, l. c. Ser. II. vii. (1851) 42 = C. Ferrandianus.
hastata, Miers, l. c. Ser. III. xix. (1867) 26 = C. Thunbergii.
hexagyna, Miers, l. c. = C. Thunbergii.
laurifolia, Miers, l. c. Ser. II. vii. (1851) 42 = C. laurifolius.
mollis, Miers, Contrib. Bot. S. Am. iii. 265 = C. mollis.
ovalifolius, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42 = C. Thunbergii.
pubinervis, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 259 = C. Thunbergii.
pycnantha, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 26 = C. Thunbergii.
sarmentosa, Miers, l. c. 25 = C. Thunbergii.
Thunbergii, Miers, l. c. 26 = C. Thunbergii.
triloba, Miers, l. c. Ser. II. vii. (1851) 42 = C. Thunbergii.

NEPHROMEDIA, Kostel. Ind. Hort. Prag. 92 (1844) = **Trigonella**, Linn. (Legumin.).
radiata, Kostel. l. c. = T. radiata.

NEPHROMISCHUS, Klotzsch, Ind. Sem. Hort. Berol. (1855) App. 1 = **Begonia**, Linn.
rutilans, Klotzsch, l. c. = B. rutilans.

NEPHROPHYLLUM, A. Rich. Tent. Fl. Abyss. ii. 77. t. 76 (1850). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 876.
HYGROCHARIS, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 77 (1851).
abyssinicum, *A. Rich. l. c.*—Abyssin.

NEPHROSIS, Rich. ex DC. Prod. ii. 420 (1825) = **Drepanocarpus**, G. F. W. Mey. (Legumin.).
aculeata, Rich. ex DC. l. c. = D. lunatus.

NEPHROSPERMA, Balf. f. in Baker, Fl. Maurit. 386 (1877). *PALMAE*. Benth. & Hook. f. iii. 907.
Van-Houtteana, *Balf. f. l. c.*—Ins. Seychelles.

NEPHROSTIGMA, Griff. Notul. iv. 717 (1854). *ANONACEAE*?—Cf. Benth. & Hook. f. i. 22.

NEPHTHYTIS, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 406. *AROIDEAE*, Benth. & Hook. f. iii. 982.
OLIGOGYNIUM, Engl. Bot. Jahrb. iv. (1883) 64.

NEPHTHYTIS:—
Afzelii, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 406.—Afr. trop.
constricta, *N. E. Br. in Gard. Chron.* (1881) I. 790.—Ins. Fernando Po.
liberica, *N. E. Br. l. c.*—Afr. trop.

NEPSERA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 28. *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
aquatica, *Naud. l. c.*— Am. trop.
glandulosa, *Naud. l. c.* 29.—Guian.
pendulifolia, Naud. l. c. = Aciotis pendulifolia.

NEPTUNIA, Lour. Fl. Cochinch. 653 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 592.
HEMIDESMAS, Rafin. Sylva Tellur. 119 (1838)
acinaciformis, *Miq. Fl. Ind. Bat.* i. I. 51 (sp. dub.).—Ins. Timor.
cinerea, F. Muell. Census, 43 = Dichrostachys cinerea.
gracilis, *Benth. in Hook. Journ. Bot.* iv. (1842) 355.—Austral.
hexapetala, *Micheli, in Mém. Soc. Phys. Hist. Nat. Genèv.* xxviii. (1883) n. VII. 50.—Paraguay.
javanica, *Miq. Fl. Ind. Bat.* i. I. 51.—Java.
lutea, *Benth. in Hook. Journ. Bot.* iv. (1842) 356.—Am. bor.
lutea, Torr. in Bot. Mex. Bound. 60, partim = pubescens.
monosperma, *F. Muell. ex Benth. Fl. Austral.* ii. 300.—Austral.
oleracea, *Lour. Fl. Cochinch.* 654.—Reg. trop.
plena, *Benth. in Hook. Journ. Bot.* iv. (1842) 355.—Am. trop.
plena, Lindl. Bot. Reg. (1846) t. 3, partim = oleracea.
polyphylla, Benth. in Hook. Journ. Bot. ii. (1840) 129 = plena.
prostrata, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 356.—Madag.
pubescens, *Benth. in Hook. Journ. Bot.* iv. (1842) 356.—Ind. occ.
spicata, F. Muell. Fragm. iii. 151 = Dichrostachys Muelleri.
stolonifera, Guill. & Perr. Fl. Seneg. Tent. i. 239 = oleracea.
surinamensis, Steud. in Flora, xxvi. (1843) 759 = plena.
tenuis, *Benth. in Hook. Journ. Bot.* iv. (1842) 355.—Texas.
triquetra, *Benth. l. c.*—Ind. or.

NERAUDIA, Gaudich. in Freyc. Voy. Bot. 500. t. 117 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 389.
glabra, Meyen, Reise, ii. 124 = melastomaefolia.
melastomaefolia, *Gaudich. in Freyc. Voy. Bot.* 500.—Ins. Sandvic.
melastomaefolia, Mus. Par. ex Blume, Mus. Bot. Lugd. Bat. ii. 56 = Pilea insignis.
ovata, Gaudich. in Freyc. Voy. Bot. 501 = melastomaefolia.
sericea, Gaudich. Bot. Voy. Bonite, t. 133 = melastomaefolia.

NERETIA, Moq. in DC. Prod. xiii. II. 58 (1849) = **Oreobliton**, Moq. & Dur. (Chenop.).
thesioides, Moq. l. c. 59 = O. thesioides.

NERIACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1096 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
Purdieanus, *Benth. l. c.; et in Hook. Ic. Pl.* t. 1200.—Jamaic.

NERIANDRA, A. DC. in DC. Prod. viii. 422 (1844) = **Skytanthus**, Meyen (Apocyn.).
angustifolia, A. DC. l. c. = S. acutus.
aurantiaca, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) 356.—Am. austr.
hancorniaefolia, A. DC. in DC. Prod. viii. 422 = S. hancorniaefolius.
havanensis, Muell. Arg. in Linnaea, xxx. (1859–60) 401 = Cameraria havanensis (in Addend.).
Martiana, Muell. Arg. in Mart. Fl. Bras. vi. I. 62 = S. Martianus.
suberecta, A. DC. in DC. Prod. viii. 422 = Urechites suberecta.

NERIJA, 'Roxb.' ex Endl. Gen. 1088 (1841) = NEERIJA, Roxb. = **Elaeodendron**, Linn. (Liliac.).

NERINE, Herb. in Bot. Mag. t. 2124 (1820). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 728.
GALATEA, Herb. in Bot. Mag. sub t. 2113 (1820).
LATICOMA, Rafin. Fl. Tellur. iv. 12 (1836).
LOXANTHES, Salisb. Gen. Pl. Fragm. 117 (1866).
aucta, M. Roem. Syn. Ensat. 107 = undulata.
aurea, Sweet, Hort. Brit. ed. I. 403 = Lycoris aurea.
cochinchinensis, M. Roem. Syn. Ensat. 104 = sarniensis.
corusca, Herb. in Bot. Mag. sub t. 2124 = sarniensis.
curvifolia, *Herb. l. c.*—Afr. austr.
excellens ×, T. Moore, in Florist (1882), t. 567.
filifolia, *Baker, in Bot. Mag.* t. 6547.—Afr. austr.
flexuosa, *Herb. in Bot. Mag.* sub t. 2124.—Afr. austr.
Fothergilli, M. Roem. Syn. Ensat. 104 = curvifolia.
humilis, *Herb. in Bot. Mag.* sub t. 2124.—Afr. austr.
hybrida, Herb. App. [Bot. Reg.] 51. t. 3 = versicolor.
Jacquini, M. Roem. Syn. Ensat. 105 = sarniensis.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 139 = Lycoris radiata.
lucida, *Herb. Amaryll.* 283. t. 26. f. 3.—Afr. austr.
marginata, *Herb. l. c.*—Afr. austr.
pudica, *Hook. f. Bot. Mag.* t. 5901.—Afr. austr.
pulchella, Herb. App. [Bot. Reg.] 19 = flexuosa.
radiata, Sweet, Hort. Brit. ed. I. 403 = Lycoris radiata.
rosea, Herb. in Bot. Mag. t. 2124 = sarniensis.
sarniensis, *Herb. l. c.* sub t. 2124.—Afr. austr.
undulata, *Herb. l. c.*—Afr. austr.
venusta, Herb. l. c. = sarniensis.
versicolor, *Herb. App.* [*Bot. Reg.*] 51.—Afr. austr.

NERION, [Tourn.] St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = **Nerium**, Linn. (Apocyn.).
oleandrum, St. Lag. l. c. = Nerium Oleander.

NERISSA, Rafin. Fl. Tellur. ii. 89 (1836) = **Ponthieva**, R. Br. (Orchid.).
glandulosa, Rafin. l. c. = P. glandulosa.

NERISSA, Salisb. Gen. Pl. Fragm. 131 (1866) = **Haemanthus**, Linn. (Amaryll.).
multiflorus, Salisb. l. c. = H. multiflorus.

NERIUM, Linn. Syst. ed. I (1735). *APOCYNACEAE*, Benth. & Hook. f. ii. 713.
OLEANDER, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 381.
antidysentericum, Lepech. Reise, i. 270 = Apocynum venetum.
antidysentericum, Linn. Sp. Pl. 209 = Wrightia zeylanica.
carneum, Hort. ex Dum.-Cours. Bot. Cult. ed. II. iii. 268 = N. Oleander.
caudatum, Lam. Encyc. iii. 458 = Strophanthus dichotomus.
caudatum, Roxb. Hort. Beng. [84]; Fl. Ind. ii. 9 = Strophanthus Wallichii.
chinense, Hunter, ex Roxb. ll. cc. = Strophanthus divergens.
coccineum, Roxb. ll. cc. 19; 2 = Wrightia coccinea.
Coraea, Buch.-Ham. ex A. DC. in DC. Prod. viii. 407, 421 = Wrightia tomentosa.
coronarium, Jacq. Coll. i. 138 = Tabernaemontana coronaria.
divaricatum, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iii. 654 = Wrightia zeylanica.
divaricatum, Linn. Sp. Pl. 209 = Tabernaemontana coronaria.
divaricatum, Lour. Fl. Cochinch. i. 142 (Quid?).—Cochinch.
divaricatum, Thunb. Fl. Jap. 110 = Rhynchospermum jasminoides.
flavescens, Spin, Jard. S. Sebast. (1812); ex Roem. & Schult. Syst. iv. 410 = odorum.
floridum, Salisb. Prod. 147 = N. Oleander.
grandiflorum, Desf. Tabl. ed. II. 92 = N. Oleander.
grandiflorum, Roxb. Hort. Beng. 19; Fl. Ind. ii. 10 = Cryptostegia grandiflora.
indicum, Mill. Gard. Dict. ed. VIII. n. 2 = odorum.
jaspideum, Span. in Linnaea, xv. (1841) 325 = Wrightia tinctoria.
Kotschii, Boiss. Diagn. Ser. I. vii. 21 = odorum.
latifolium, Mill. Gard. Dict. ed. VIII. n. 3 = odorum.
lauriforme, Lam. Fl. Fr. ii. 299 = N. Oleander.
luteum, Nois. ex Steud. Nom. ed. I. 553 = odorum.
macrocarpum, Span. in Linnaea, xv. (1841) 325 = Wrightia tinctoria.
mascatense, *A. DC. in DC. Prod.* viii. 421.—Arab.; Persia.

NERIUM:—
obesum, Forsk. Fl. Aegypt. Arab. 205 = Adenium obesum.
odoratum, Lam. Encyc. iii. 456 = seq.
odorum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 297.—Persia; Ind. or.; Japon.
Oleander, *Linn. Sp. Pl.* 209.—Reg. Mediterr.; Oriens.
pauciflorum, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 6 = Cryptolepis elegans.
piscidium, Roxb. Hort. Beng. 19; Fl. Ind. ii. 7 = Melodinus monogynus.
piscidium, Wall. Cat. n. 8177 = Cosmostigma racemosa.
reticulatum, Roxb. Hort. Beng. 19; Fl. Ind. ii. 8 = Cryptolepis Buchanani.
salicinum, *Forsk. Fl. Aegypt. Arab.* 205.—Arabia.
scandens, Lour. Fl. Cochinch. 116 = Strophanthus dichotomus.
scandens, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 148 (Alafiae sp. ?).—Afr. trop.
sibiricum, Medic. Beobacht. (1782) 15 = Apocynum cannabinum.
splendens, Hort. ex Paxt. Mag. Bot. iii. (1837) 73 = N. Oleander.
thyrsiflorum, *Paxt. l. c.*—Nepal.
tinctorium, Hort. ex Sweet, Hort. Brit. ed. I. 274 = Alstonia scholaris, Allamanda verticillata.
tinctorium, Roxb. Hort. Beng. 19; Fl. Ind. ii. 4 = Wrightia tinctoria.
tomentosum, Roxb. ll. cc. [84]; 6 = Wrightia tomentosa.
undulatum, Salisb. Prod. 148 = Celastrus undulatus.
verecundum, Salisb. l. c. 147 = odorum.
zeylanicum, Linn. Cent. Pl. ii. 12 = Wrightea zeylanica.

NEROPHILA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 119. *MELASTOMACEAE*, Benth. & Hook. f. i. 744.
gentianoides, *Naud. l. c.* 120.—Afr. trop.

NERTERA, Banks & Soland. ex Gaertn. Fruct. i. 124. t. 26 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 138.
CUNINA, Clos, in C. Gay, Fl. Chil. iii. 201. t. 34 (1849).
ERYTHRODANUM, Thou. Desc. Tristan d'Acugna, 41. tt. 10, 11 (1811).
GOMOZIA, Mutis, ex Linn. f. Suppl. 17 (1781).
LEPTOSTIGMA, Arn. in Hook. Journ. Bot. iii. (1841) 270.
NERTERIA, Sm. Ic. Ined. ii. 28. t. 28 (1790).
adsurgens, DC. Prod. ii. 452 = *assurgens*.
alsinoides, Cham. & Schlecht. in Linnaea, vi. (1831) 413 = depressa.
assurgens, Thou. Desc. Tristan d'Acugna, 42. t. 11 = depressa.
Cunninghamii, *Hook. f. Fl. N. Zel.* 112.—N. Zel.; Ins. Philipp.
depressa, *Banks & Soland. ex Gaertn. Fruct.* i. 124. t. 26.—Am. austr.; N. Zel.; Ins. Trist. Acun.
dichondraefolia, *Hook. f. Fl. N. Zel.* 112. t. 28 A.—N. Zel.
gracilis, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1864) 121 = dichondraefolia.
pusilla, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 331.—N. Zel.
repens, Ruiz & Pav. Fl. Peruv. i. 60 = depressa.
reptans, *F. Muell. ex Benth. Fl. Austral.* iii. 431.—Austral.
scapanioides, *Lange, Ind. Sem. Hort. Haun.* (1868) 22.—N. Zel.
setulosa, *Hook. f. Fl. N. Zel.* 112. t. 28 B.—N. Zel.
tetrasperma, H. B. & K. Nov. Gen. et Sp. iii. 379 = Mitchella ovata.

NERTERIA, Sm. Ic. Ined. ii. 28. t. 28 (1790) = praec.

NERVILIA, Comm. ex Gaudich. in Freyc. Voy. Bot. 422. t. 35 (1826) = **Pogonia**, Juss. (Orchid.).
Aragoana, Gaudich. l. c. = P. flabelliformis.
ovata, Gaudich. l. c. = P. ovata.

NESAEA, Comm. ex Juss. Gen. 332 (1789). *LYTHRARIEAE*, Benth. & Hook. f. i. 779.
CHRYSOLIGA, Willd. ex DC. Prod. iii. 89, in syn. (1828).

NESAEA:—
DECODON, J. F. Gmel. Syst. 677 (1791).
HEIMIA, Link & Otto, Ic. Pl. 63. t. 28 (1822).
TOLYPEUMA, E. Mey. in Drège, Zwei Pfl. Docum. 226 (1843), nomen.
TROTULA, Comm. ex DC. Prod. iii. 90 (1828).
anagalloides, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 327.—Afr. austr.
andongensis, *Welw. ex Hiern, in Oliver, Fl. Trop. Afr.* ii. 473.—Afr. trop.
arnhemica, *F. Muell. Fragm.* vii. 146.—Austral.
aspera, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 327.—Afr. trop.
brevipes, *Koehne, l. c.* 326.—Zeylan.
Candollei, *Guill. & Perr. Fl. Seneg. Tent.* i. 307.—Afr. trop.
capitellata, Presl, in Oken, Isis, xxi. (1828) 273 = triflora.
cordata, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 475.—Afr. trop.
crassicaulis, Koehne, in Engl. Bot. Jahrb. iii. (1882) 324 = polyantha.
crinipes, *Koehne, l. c.* 337.—Austral.
dodecandra, Koehne, l. c. 334 = Candollei.
erecta, *Guill. & Perr. Fl. Seneg. Tent.* i. 305.—Afr. trop.
floribunda, *Sond. in Harv. & Sond. Fl. Cap.* ii. 517.—Afr. austr.
heptamera, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 472.—Afr. trop.
humilis, Klotzsch, in Peters, Reise Mossamb. Bot. 68 = erecta.
icosandra, *Kotschy & Peyr. Pl. Tinn.* 10. t. 5a.—Afr. trop.
lanceolata, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 325.—Zeylan.
linariaefolia, *Steud. in Flora*, xxv. (1842) 474.—Am. austr.
linearis, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 475.—Afr. trop.
linifolia, *Welw. ex Hiern, l. c.* 472.—Afr. trop.
loandensis, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 325.—Afr. trop.
longipes, *A. Gray, Pl. Wright.* i. 68.—Texas.
lythroides, *Welw. ex Hiern, in Oliver, Fl. Trop. Afr.* ii. 471.—Afr. trop.
myrtifolia, *Desf. ex A. St. Hil. Fl. Bras. Mer.* iii. 138.—Bras.
passerinoides, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 338.—Afr. trop.
pedicellata, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 472.—Afr. trop.
polyandra, Steud. in Flora, xxv. (1842) 472 = Pleurophora polyandra.
polyantha, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 132.—Madag.
pusilla, Steud. in Flora, xxv. (1842) 473 = Pleurophora pusilla.
racemosa, Klotzsch, in Peters, Reise Mossamb. Bot. 68, in adnot. = erecta.
radicans, *Guill. & Perr. Fl. Seneg. Tent.* i. 306. t. 70.—Afr. trop.
recta, Steud. in Flora, xxv. (1842) 473 = Pleurophora pungens.
rigidula, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 333.—Afr. austr.
Robertsii, *F. Muell. Fragm.* vii. 145.—Austral.
sagittifolia, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 339.—Afr. austr.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 192.—Am. trop.
sarcophylla, *Koehne, in Engl. Bot. Jahrb.* iii. (1882) 328.—Afr. trop.
speciosa, H. B. & K. Nov. Gen. et Sp. vi. 191 = Diplusodon speciosus.
squarrosa, Steud. in Flora, xxv. (1842) 473 = Pleurophora polyandra.
syphilitica, *Steud. l. c.* 474.—Mexic.
triflora, *H. B. & K. Nov. Gen. et Sp.* vi. 191.—Am. austr.; Ins. Mascar.
verticillata, *H. B. & K. l. c.*—Am. bor. et austr.

NESCIDIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 192 = **Myonima**, Comm. (Rubiac.).
myrtifolia, A. Rich. l. c. 193 = M. myrtifolia.

NESIOTA, Hook. f. in Benth. & Hook. f. Gen. i. 380 (1862). *RHAMNEAE*, Benth. & Hook. f. l. c.
elliptica, *Hook. f. l. c.*; *et Ic. Pl.* t. 1052.—Ins. S. Helena.

NESLEA, Aschers. Fl. Brandenb. 63 (1864) = seq.

NESLIA, Desv. Journ. Bot. iii. (1814) 162. *CRUCIFERAE*, Benth. & Hook. f. i. 95.
VOGELIA, Medic. Pflanzeng. 32. t. 1. f. 6 (1792).
apiculata, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 68 = paniculata.
paniculata, *Desv. Journ. Bot.* iii. (1814) 162.—Europ.; Oriens; Ind. bor. occ.

NESODAPHNE, Hook. f. Fl. N. Zel. i. 217 (1853) = **Beilschmiedia**, Nees (Laurin.).
obtusifolia, Benth. Fl. Austral. v. 299.—Austral.
Tarairi, Hook. f. Fl. N. Zel. i. 217.—N. Zel.
Tawa, Hook. f. l. c.—N. Zel.

NESOGENES, A. DC. in DC. Prod. xi. 703 (1847). *VERBENACEAE*, Benth. & Hook. f. ii. 1141.
decumbens, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 19; *et in Baker, Fl. Maurit.* 252.—Ins. Rodriguez.
euphrasioides, *A. DC. in DC. Prod.* xi. 703.—Ins. Sandvic.

NESOPANAX, Seem. Journ. Bot. ii. (1864) 249 = **Plerandra**, A. Gray (Araliac.).
vitiensis, Seem. l. c. = P. vitiensis.

NESSEA, Steud. Nom. ed. I. 553 (1821) = **Nesaea**, Comm. (Lythrar.).

NESTEGIS, Rafin. Sylva Tellur. 10 (1838) = **Olea**, Tourn.
elliptica, Rafin. l. c. = O. americana ?

NESTLERA, E. Mey. ex Walp. Ann. iii. 856 (1852) = **Leucosidea**, Eckl. & Zeyh. (Rosac.).
acaenoides, E. Mey. l. c. = L. acaenoides.

NESTLERA, Spreng. Anleit. ii. II. 568 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 325.
COLUMELLEA, Jacq. Hort. Schoenb. iii. 28. t. 301 (1798).
POLYCHAETIA, Less. Syn. Comp. 371 (1832).
STEPHANOPAPPUS, Less. in Linnaea, vi. (1831) 234.
acerosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 296.—Afr. austr.
biennis, *Spreng. Syst.* iii. 589.—Afr. austr.
conferta, *DC. Prod.* vi. 284.—Afr. austr.
Dregeana, *Harv. in Harv. & Sond. Fl. Cap.* iii. 296.—Afr. austr.
Garnotii, Harv. l. c. = tenuifolia.
humilis, *Less. Syn. Comp.* 372; *DC. Prod.* vi. 283.—Afr. austr.
minuta, DC. l. c. = humilis.
muriculata, *DC. l. c.* 284.—Afr. austr.
oppositifolia, *DC. l. c.* 283.—Afr. austr.
prostrata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 296.—Afr. austr.
reflexa, DC. Prod. vi. 283 = biennis.
rigida, DC. l. c. 284 = humilis.
tenuifolia, *DC. l. c.*—Afr. austr.
tricephala, *Harv. in Harv. & Sond. Fl. Cap.* iii. 297.—Afr. austr.

NESTLERA, Willd. ex Steud. Nom. ed. II. ii. 192 (1841) = **Bouteloua**, Lag. (Gramin.).
festucaeformis, Willd. l. c. = B. bromoides.

NESTRONIA, Rafin. New Fl. Am. iii. 12 (1836) = **Buckleya**, Torr. (Santal.).
cuneata, Rafin. l. c. 14 = Ceanothus cuneatus.
umbellulata, Rafin. l. c. 13 (= *Darbya umbellulata*).—Am. bor.
undulata, Rafin. l. c. = B. distichophylla ?

NESTYLIX, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
amygdalina, Rafin. l. c. = S. triandra.
arborea, Rafin. l. c. = S. triandra.
falcata, Rafin. l. c. = S. nigra.
Hopeana, Rafin. l. c. = S. triandra.
Villarsiana, Rafin. l. c. = S. triandra.

NESYNSTYLIS, Rafin. Fl. Tellur. iv. 123 (1836) = **Strumaria**, Jacq. (Amaryll.).
filifolia, Rafin. l. c. = Nessea filifolia.
undulata, Rafin. l. c. = S. undulata.

NETOUXIA, G. Don, Gen. Syst. iv. 454 (1837) = **Nectouxia**, H. B. & K. (Solan.).

NETTLERA, Rafin. Sylva Tellur. 147 (1838) = **Cephaëlis**, Sw. (Rubiac.).
guianensis, Rafin. l. c.—Guiana.

NETTOA, Baill. in Adansonia, vi. (1866) 238. t. 7. *TILIACEAE*, Benth. & Hook. f. i. 986.
chrozophorifolia, *Baill. l. c.* 238, 242.—Austral.

NEUBECKIA, Alef. in Bot. Zeit. xxi. (1863) 290, 297 = **Iris**, Tourn.
caucasica, Alef. l. c. 297 = I. caucasica.
cristata, Alef. l. c. = I. cristata.
decora, Klatt, in Linnaea, xxxiv. (1865–66) 588 = I. nepalensis.
fulva, Alef. in Bot. Zeit. xxi. (1863) 297 = I. fulva.
humilis, Alef. l. c. = I. humilis.
reticulata, Alef. l. c. = I. reticulata.
scorpioides, Alef. l. c. = I. alata.
stylosa, Alef. l. c. = I. unguicularis.
sulcata, Klatt, in Linnaea, xxxiv. (1865–66) 590 = I. nepalensis.
tenuifolia, Alef. in Bot. Zeit. xxi. (1863) 297 = I. tenuifolia.
unguicularis, Alef. ex Nym. Consp. 702, Index 980 = I. unguicularis.

NEUBERIA, Eckl. Top. Verz. 37 (1827) = **Watsonia**, Mill. (Irid.).
humilis, Eckl. l. c. = W. humilis.
laccata, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 158 = W. humilis.
longifolia, Eckl. Top. Verz. 37 = Gladiolus Ecklonii.
marginata, Eckl. l. c. = W. marginata.
pyramidalis, Eckl. l. c. = W. Meriana.
rosea, Eckl. l. c. = W. rosea.

NEUBURGHIA, Walp. Ann. iii. 30 (1852) = seq.

NEUBURGIA, Blume, Mus. Bot. Lugd. Bat. i. 156 (1850). *APOCYNACEAE*, Benth. & Hook. f. ii. 700.
musculiformis, *Miq. Fl. Ind. Bat.* ii. 403.—Ins. Molucc.; N. Guin.
tuberculata, *Blume, Mus. Bot. Lugd. Bat.* i. 157.—Ins. Molucc.
tubiflora, *Blume, l. c.* 156.—N. Guin.

NEUDORFIA, Adans. Fam. ii. 219 (1763) = **Nolana**, Linn. (Solan.).
peruviana, Adans. l. c. = Nolana prostrata.

NEUHOFIA, Stokes, Bot. Mat. Med. i. 439 (1812) = **Baeckea**, Linn. (Myrtac.).
rosmarinifolia, Stokes, l. c. = B. cochinchinensis.

NEUMANNIA, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 369 = **Pitcairnia**, L'Hérit. (Bromel.).
Altensteinii, Griseb. in Goett. Nachr. (1864) 14 = P. Altensteinii.
atro-rubens, C. Koch, Allg. Gartenz. (1858) 269 = P. atro-rubens.
imbricata, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 369 = P. imbricata, *Baker*.
maidifolia, Koch, Ind. Sem. Hort. Berol. (1856) App. 2 = P. maidifolia.
nigra, E. Morr. Cat. Jard. Univ. Liège (1873), 11; Carr. in Rev. Hortic. (1881) 390.—Hab. ?
ochroleuca, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 2 = P. ochroleuca.
petiolata, Koch & Bouché, l. c. = P. petiolata.
sulphurea, C. Koch, l. c. = P. Wendlandi.

NEUMANNIA, A. Rich. Ess. Fl. Cub. 96, in adnot. (1845) = **Aphloia**, Benn. (Bixin.).
integrifolia, A. Rich. l. c. 97 = A. mauritiana.
theaeformis, A. Rich. l. c. = A. mauritiana.

NEUMAYERA, Reichb. Ic. Fl. Germ. xv. 30 (1841) = **Arenaria**, Rupp. (Caryoph.).
austriaca, Reichb. l. c. = A. austriaca.
filicaulis, Reichb. l. c. = A. austriaca.
Villarsii, Reichb. l. c. = A. Villarsii.

NEUMAYERA, Reichb. f. in Flora, lv. (1872) 278 = NIEMAYERA, F. Muell. = **Apostasia**, Blume (Orchid.).
stylidioides, Reichb. f. l. c. = A. stylidioides.

NEURACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 76 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1093.
aculeatus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 86.—Ins. Socotra.
africanus, *T. Anders. ex S. Moore, in Journ. Bot.* xviii. (1880) 37.—Afr. trop.
capitatus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 86.—Ins. Socotra.
decorus, *S. Moore, in Journ. Bot.* xviii. (1880) 307.—Afr. trop.
grandiflorus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 95.—Burma.
Lawii, Wight, Ic. iv. t. 1532 = sphaerostachyus.
Neesianus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 491.—Ind. or.
niveus, *S. Moore, in Journ. Bot.* xviii. (1880) 37.—Afr. trop.
parviflorus, *T. Anders. in Journ. Agr. Hort. Soc. Ind.* N. S. i. (1868) 276.—Ins. Maurit. ?
scaber, *S. Moore, in Journ. Bot.* xviii. (1880) 307.—Afr. trop.
sphaerostachys, *Dalz. in Hook. Kew Journ.* ii. (1850) 140.—Ind. or.
subuninervis, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 95 = Lepidagathis subuninervia.
tetragonostachys, Wall. Cat. n. 7168, partim = Justicia dasycarpa.
tetragonostachyus, *Nees, in Wall. Pl. As. Rar.* iii. 97.—Burma.
trinervius, *Wight, Ic.* iv. t. 1532.—Ind. or.

NEURACHNE, R. Br. Prod. 196 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1123.
alopecuroides, *R. Br. l. c.*—Austral.
meneritana, Boj. Hort. Maurit. 366 = Isachne miliacea.
meneritana, R. Br. ex Roem. & Schult. Syst. ii. 475 = Isachne australis.
Mitchelliana, *Nees, in Hook. Lond. Journ. Bot.* ii. (1843) 410.—Austral.
montana, Gaudich. in Freyc. Voy. Bot. t. 26 = Panicum nubigenum.
Munroi, *F. Muell. Fragm.* viii. 200.—Austral.
paradoxa, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 89 = Spinifex paradoxus.
phleoides, *R. Br. ex Nees, in Hook. Lond. Journ. Bot.* ii. (1843) 411.—Austral.
torrida, Gaudich. ex Kunth, Enum. Pl. i. 114 = Panicum torridum.

NEURACTIS, Cass. in Dict. Sc. Nat. xxxiv. 496 (1825) = **Chrysanthellum**, Rich. (Chrysanth.).
Leschenaultii, Cass. l. c.—Java.

NEURADA, Linn. Gen. ed. II. 185 (1742). *ROSACEAE*, Benth. & Hook. f. i. 625.
FIGARAEA, Viv. Pl. Aegypt. Dec. 11 (1830).
NEURAS, Adans. Fam. ii. 293 (1763).
TRIBULASTRUM, B. Juss. ex Linn. Gen. ed. II. 185 (1742).
procumbens, *Linn. Sp. Pl.* 441.—Afr. bor.; Oriens; Ind. or.

NEURAS, Adans. Fam. ii. 293 (1763) = praec.

NEURELMIS Rafin. New Fl. Am. iv. 72 (1836).—Cf. JALAMBICA, Rafin. (Compos.).
pumila, Rafin. l. c. (= *J. pumila*).—Am. bor.

NEUROCALYX, Hook. Ic. Pl. t. 174 (1837). *RUBIACEAE*, Benth. & Hook. f. ii. 54.
capitata, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 47.—Zeylan.
Championii, *Benth. ex Thw. Enum. Pl. Zeyl.* 139.—Zeylan.
Gardneri, *Thw. l. c.*—Zeylan.
Hookerianus, Wight, Ic. i. t. 52 = Wightii.
Wightii, *Arn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 22.—Ind. or.; Zeylan.
zeylanicus, *Hook. Ic. Pl.* t. 174.—Zeylan.

NEUROCARPAEA, R. Br. in Salt, Abyss. App. 64 (1814) = **Pentas**, Benth. (Rubiac.).
lanceolata, R. Br. l. c. = P. carnea.

NEUROCARPUM, Desv. Journ. Bot. i. (1813) 119; iii. (1814) 75 = **Clitoria**, Linn. (Legumin.).
angustifolium, Kunth, Mimos. 218. t. 60 = C. guianensis.
argenteum, Duch. & Walp. in Flora, xxxvi. (1853) 228 = C. glycinoides.
barbatum, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 412.—Bras.
bracteatum, Mart. ex Walp. Rep. i. 753 = C. stipularis.
cajanifolium, Presl, Symb. Bot. i. 17. t. 9 = C. cajanifolia.
densiflorum, Benth. in Ann. Wien. Mus. ii. (1838) 117 = C. densiflora.
ellipticum, Desv. in Journ. Bot. i. (1813) 119 = C. glycinoides.
erectum, Voigt, Hort. Suburb. Calc. 213 = C. erecta.
falcatum, DC. Prod. ii. 236 = C. glycinoides.
flagellare, Benth. in Hook. Journ. Bot. ii. (1840) 58 = C. flagellaris.
frigidulum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 116 = C. guianensis.
glycinoides, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 413 = C. glycinoides.
guianense, Desv. Journ. Bot. i. (1813) 119 (*gajanense*) = C. guianensis.
janense, Desv. l. c. iii. (1814) 75 = praec.
javitense, H. B. & K. Nov. Gen. et Sp. vi. 409 = C. javitensis.
laurifolium, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = C. cajanifolia.
longifolium, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 116 = C. guianensis.
macrophyllum, H. B. & K. Nov. Gen. et Sp. vi. 409 = C. macrophylla.
mexicanum, Steud. Nom. ed. II. ii. 193.—Mexic.
multiflorum, Hook. & Arn. Bot. Beech. Voy. 286.—Mexic.
resupinatum, Bunb. in Proc. Linn. Soc. i. (1841) 110.—Reg. Argent.
retusum, Hassk. Cat. Hort. Bog. Alt. 276 = C. cajanifolia.
rubiginosum, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = C. glycinoides.
rufescens, Benth. in Ann. Wien. Mus. ii. (1838) 116 = C. rufescens.
simplicifolium, Kunth, Mimos. 213. t. 59 = C. simplicifolia.
speciosum, Klotzsch, in Schomb. Fauna & Fl. Gui. 1202, nomen.—Guian.
villosum, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 413. = C. glycinoides.

NEUROCHLAENA, Less. Syn. Comp. 388 (1832) = **Neurolaena**, R. Br. (Compos.).

NEUROCTOLA, Rafin. ex Steud. Nom. ed. II. ii. 193 (1841) = NEVROCTOLA, Rafin. = **Uniola**, Linn. (Gramin.).

NEUROLA, Rafin. New Fl. Am. iv. 92 (1836) = **Sabbatia**, Adans. (Gentian.).
arkansica, Rafin. l. c.—Am. bor. occ.

NEUROLAENA, R. Br. in Trans. Linn. Soc. xii. (1817) 120. *COMPOSITAE*, Benth. & Hook. f. ii. 437.
CALEA, Sw. Prod. Veg. Ind. Occ. 113 (1788).
NEUROCHLAENA, Less. Syn. Comp. 388 (1832).
integrifolia, Cass. in Dict. Sc. Nat. xxxiv. 501 = lobata.
integrifolia, *Klatt, in Leopoldina*, xx. (1884) 94.—Ecuador.
lobata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 120.—Am. trop.
macrocephala, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 233.—Mexic.
semidentata, *Klatt, in Leopoldina*, xx. (1884) 94.—Mexic.
Suriani, Cass. in Journ. Phys. (Juillet, 1818) 29 = lobata.

NEUROLOMA, Andrz. ex DC. Prod. i. 156 (1824) = **Parrya**, R. Br. (Crucif.).
Adamsianum, Andrz. ex Ledeb. Fl. Ross. i. 131 = P. macrocarpa.
arabidiflorum, DC. Prod. i. 156 = P. Linnaeana.
articum, Spreng. Syst. ii. 888 = P. arctica.
exscapum, Steud. Nom. ed. II. ii. 193 = P. exscapa.
Haleniana, Andrz. ex Ledeb. Fl. Ross. i. 131 = P. macrocarpa.
nudicaule, DC. Prod. i. 156 = P. macrocarpa.
scapigerum, DC. l. c. = P. macrocarpa.
speciosum, Steud. Nom. ed. II. ii. 193 = P. integerrima.

NEUROLOMA, Endl. Gen. 99 (1836) in syn. = NEVROLOMA, Rafin. = **Briza**, Linn. (Gramin.).

NEUROPELTIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 43 (1824). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 878.
acuminata, *Benth. & Hook. f. Gen.* ii. 876, in nota, 878.—Afr. trop.
bracteata, Griff. Notul. iv. 285 = racemosa.
intermedia, Griff. l. c. = racemosa.
ovata, Wall. Cat. n. 1323 = racemosa.
racemosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 43; *Cat.* n. 1322.—Ind. or.; Malaya.

NEUROPHYLLUM, Torr. & Gray, Fl. N. Am. i. 612 (1840) = **Peucedanum**, Linn. (Umbell.).
longifolium, Torr. & Gray, l. c. = Peucedanum ternatum.

NEUROSCAPHA, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 137 = **Lonchocarpus**, H. B. & K. (Legumin.).
Guilleminiana, Tul. l. c. = L. Neuroscapha.
Martiana, Tul. l. c. 138 = L. Neuroscapha.
pubigera, Tul. l. c. = L. Neuroscapha.

NEUROSPERMUM, Bartl. Ord. 275 (1830) = NEVROSPERMA, Rafin. = **Momordica**, Linn. (Cucurb.).

NEUROTHECA, Salisb. ex Benth. & Hook. f. Gen. ii. 812,(1876). *GENTIANEAE*, Benth. & Hook. f. l. c.
OCTOPLEURA, Spruce, ex Prog. in Mart. Fl. Bras. vi. i. 212 (1865).
loeselioides, *Oliver* (*in Herb. Kew.*).—Afr. trop.; Bras.

NEUSTANTHUS, Benth. in Miq. Pl. Jungh. 234 (1851–55) = **Pueraria**, DC. (Legumin.).
chinensis, Benth. Fl. Hongk. 86 = P. Thunbergiana.
javanicus, Benth. in Miq. Pl. Jungh. 235 = P. phaseoloides.
peduncularis, Benth. l. c. = P. peduncularis.
phaseoloides, Benth. l. c. = P. phaseoloides.
sericans, Miq. Fl. Ind. Bat. i. 1. 218 = P. phaseoloides.
subspicatus, Benth. in Miq. Pl. Jungh. 234 = P. phaseoloides.

NEUWIEDIA, Blume, in Hoev. & De Vriese, Tijdschr. i. (1834) 140. *ORCHIDEAE*, Benth. & Hook. f. iii. 635.
Griffithii, *Reichb. f. Xenia Orch.* ii. 215.—Malaya.
veratrifolia, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 142.—Java.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 58.—Java.

NEVIUSA, Benth. & Hook. f. Gen. i. 613 (1865), sphalm. = seq.

NEVIUSIA, A. Gray, in Mem. Am. Acad. N. S. vi. (1858) 374. *ROSACEAE*, Benth. & Hook. f. i. 613.
alabamensis, *A. Gray, l. c.*—Alabama.

NEVOSMILA, Rafin. Sylva Tellur. 107 (1838) = **Crataeva**, Linn. (Cappar.).
arborea, Rafin. l. c. = C. gynandra.

NEVRILIS, Rafin. Sylva Tellur. 138 (1838) = **Millingtonia**, Linn. f. (Bignon.).
suberosa, Rafin. l. c. = M. hortensis.

NEVROCTOLA, Rafin. Neogenyt. 4 (1825) = **Uniola**, Linn. (Gramin.).
maritima, Rafin. l. c. = U. paniculata.
paniculata, Rafin. l. c. = U. paniculata.

NEVROLOMA, Rafin. in Journ. Phys. lxxxix. (1819) 106 = **Briza**, Linn. (Gramin.).
canadensis, Rafin. l. c. = B. canadensis.

NEVROLOMA, Spreng. Syst. ii. 888 (1825) = NEUROLOMA, Andrz. = **Parrya**, R. Br. (Crucif.).

NEVROSPERMA, Rafin. in Am. Monthly Mag. (1818) 40 = **Momordica**, Linn. (Cucurb.).
cuspidata, Rafin. l. c. = M. Balsamina.

NEWBERRYA, Torr. in Ann. Lyc. N. York, viii. (1867) 55, in adnot. *MONOTROPEAE*, Benth. & Hook. f. ii. 606.
HEMITOMES, A. Gray, in Pacif. Rail. Rep. vi. 80. t. 12 (1855).
congesta, *Torr. in Ann. Lyc. N. York*, viii. (1867) 55.—Am. bor. occ.
spicata, *A. Gray, in Proc. Am. Acad.* xv. (1880) 44.—Calif.

NEWBOULDIA, Seem. Journ. Bot. i. (1863) 225. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1045.
SPATHOTECOMA, Bur. Monog. Bign. 49 (1864).
laevis, *Seem. Journ. Bot.* i. (1863) 226; *et* viii. (1870) 210, 337.—Afr. trop.
pentandra, *Seem. l. c.* viii. 338.—Afr. trop.

NEWCASTLIA, F. Muell. in Hook. Kew Journ. ix. (1857) 22 (Newcastelia). *VERBENACEAE*, Benth. & Hook. f. ii. 1139.
bracteosa, *F. Muell. Fragm.* viii. 49.—Austral.
cephalantha, *F. Muell. l. c.* ix. 4.—Austral.
chrysotricha, *F. Muell. l. c.* 15.—Austral.
cladotricha, *F. Muell. in Hook. Kew Journ.* ix. (1857) 22.—Austral.
hexarrhena, *F. Muell. Fragm.* x. 16.—Austral.
spodiotricha, *F. Muell. l. c.* iii. 21. t. 21.—Austral.

NEXILIS, Rafin. New Fl. Am. iv. 9 (1836) = **Ammannia**, Houst. (Lythrar.).

NEZERA, Rafin. New Fl. Am. iv. 64 (1836) = **Linum**, Tourn.
albiflora, Rafin. l. c. 65 = L. striatum.
carnea, Rafin. l. c. = L. carneum?
cuspidata, Rafin. l. c. 66 = L. cuspidatum.
imbricata, Rafin. l. c. = L. imbricatum.

NHANDIROBA, Plum. ex Adans. Fam. ii. 139 (1763) = **Fevillea**, Linn. (Cucurb.).
scandens, Desc. Fl. Antill. viii. Index, 66 [t. 198] = F. cordifolia.

NIARA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Myrsine**, Linn.
montanensis, Dennst. l. c.—Ind. or.

NIBBISIA, Walp. Rep. i. 59 (1842), sphalm. = NIRBISIA, G. Don = **Aconitum**, Linn. (Ranunc.).

NIBO, Steud. Nom. ed. I. 553 (1821) = VIBO, Medic. = **Emex**, Neck. (Polygon.).

NIBORA, Rafin. Fl. Ludov. 36 (1817) = **Gratiola**, Rupp. (Scrophul.).
aquatica, Rafin. l. c. = G. Nibora.

NICANDRA, Adans. Fam. ii. 219 (1763). *SOLANACEAE*, Benth. & Hook. f. ii. 897.
CALYDERMOS, Ruiz & Pav. Fl. Per. et Chil. ii. 43 (1799).
anomala, Link & Otto, Ic. Pl. Sel. 77. t. 35 = Scopola lurida.
indica, Roem. & Schult. Syst. iv. 682 = Physalis minima.
minor, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 81 = physaloides.
physaloides, *Gaertn. Fruct.* ii. 237. t. 131. f. 2.—Peruv.

NICANDRA, Schreb. Gen. 283 (1789) = **Potalia**, Aubl. (Logan.).
amara, J. F. Gmel. Syst. 677; Willd. Sp. Pl. ii. 341 = P. amara.

NICHOLSONIA, Span. in Hook. Comp. Bot. Mag. i. (1835) 346 = NICOLSONIA, DC. = **Desmodium**, Desv. (Legumin.).

NICIPE, Rafin. Fl. Tellur. iii. 54 (1836) = **Ornithogalum**, Tourn. (Liliac.).
nivea, Rafin. l. c. = O. niveum.

NICODEMIA, Tenore, Cat. Ort. Napol. 88 (1845). *LOGANIACEAE*, Benth. & Hook. f. ii. 794.
diversifolia, *Tenore, l. c.*—Ins. Mascar.
rondeletiaeflora, *Benth. in Journ. Linn. Soc.* i. (1857) 108.—Ins. Comor.

NICOLAIA, Horan. Prod. Scitam. 32. t. 1 (1862) = **Amomum**, Linn. (Scitam.).
elatior, Horan. l. c. = Alpinia elatior.
hemisphaerica, Horan. l. c. (= *Elettaria hemisphaerica*).—Java.
imperialis, Horan. l. c. = A. magnificum.
pallida, Horan. l. c. (= *Elettaria pallida*).—Java.
solaris, Horan. l. c. (= *Elettaria solaris*).—Java.
speciosa, Horan. l. c. (= *Elettaria speciosa*).—Java.

NICOLETTIA, Benth. & Hook. f. Gen. ii. 409 (1873) = seq.

NICOLLETIA, A. Gray, in Rep. Frém. Exp. ii. 315 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 409.
Edwardsii, *A. Gray, Pl. Wright.* i. 119.—Mexic.
occidentalis, *A. Gray, in Rep. Frém. Exp.* ii. 316.—Calif.

NICOLSONIA, DC. Prod. ii. 325; Mém. Lég. vii. 311. t. 51 (1825) = **Desmodium**, Desv. (Legumin.).
barbata, DC. ll. cc. 325; 313 = D. barbatum.
caffra, E. Mey. Comm. Pl. Afr. Austr. 123 = D. Dregeanum.
cayennensis, DC. Mém. Lég. vii. 314. t. 51 = D. barbatum.
congesta, Wight, Ic. t. 1056 = Leptodesmia congesta.
major, Steud. in Flora, xxvi. (1843) 757 = D. barbatum.
oxalidifolia, Span. in Hook. Comp. Bot. Mag. i. (1835) 346, nomen.—Malaya.
radicans, Steud. in Flora, xxvi. (1843) 757 = D. barbatum.
reptans, Meissn. in Linnaea, xxi. (1848) 260 = D. triflorum.
setigera, E. Mey. Comm. Pl. Afr. Austr. 124 = D. hirtum.
styracifolia, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 418 = D. retroflexum.
triflora, Griseb. in Goett. Abh. vii. (1857) 202 = D. triflorum.
venustula, DC. Prod. ii. 325 = D. venustulum.
villosa, Cham. & Schlecht. in Linnaea, v. (1830) 584 = D. villosum.

NICOTIA, Opiz, Seznam, 68 (1852) = **Nicotiana**, Linn. (Solanac.).
rustica, Opiz, l. c. = Nicotiana rustica.

NICOTIANA, Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 906.
AMPHIPLEIS, Rafin. Fl. Tellur. iii. 75 (1836).
CAPNOREA, Rafin. l. c. 74 (1836).
CODYLIS, Rafin. in Am. Monthly Mag. (1819) 191.
EUCAPNIA, Rafin. Fl. Tellur. iii. 74 (1836).
LANGSDORFIA, Rafin. l. c. (1836).
LEHMANNIA, Spreng. Anleit. ii. II. 458 (1817).
NICOTIA, Opiz, Seznam, 68 (1852).
NICOTIDENDRON, Griseb. in Abh. Goett. xix. (1874) 216.
PERIETERIS, Rafin. Fl. Tellur. iii. 74 (1836).
POLYDICLIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 361.
SAIRANTHUS, G. Don, Gen. Syst. iv. 467 (1837).
SIPHAULAX, Rafin. Fl. Tellur. iii. 74 (1836).
TABACINA, Reichb. Handb. 201 (1837).
TABACUM, Gilib. Fl. Lituan. i. 39 (1781).
acuminata, *Hook. Bot. Mag.* t. 2919.—Am. austr.
acuta, *Griseb. in Goett. Abh.* xix. (1874) 215.—Reg. Argent.
acutiflora, *A. St. Hil. Pl. Rém. Bras.* 209, 223.—Bras.
acutifolia, Burb. in Gard. Chron. (1881) II. 183 = alata.
acutifolia, Walp. Rep. iii. 12, in syn. = acutiflora.
affinis, Hort.—Cf. Gard. Chron. (1881) II. 141, 271 = alata.
alata, *Link & Otto, Ic. Pl. Rar.* i. 63. t. 32.—Bras.
alba, Mill. Gard. Dict. ed. VIII. n. 5 = N. Tabacum.
alipes, Steud. Nom. ed. I. 553 = N. Tabacum.
alpina, Poepp. ex Dun. in DC. Prod. xiii. 1. 564 = monticola.
andicola, *H. B. & K. Nov. Gen. et Sp.* iii. 1.—Peruv.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3; Ehrh. Beitr. v. 169 = N. Tabacum.
angustifolia, Ruiz & Pav. Fl. Peruv. ii. 16. t. 130 = pusilla.
anisandra, *Vest, in Flora*, iv. (1821) 147.—Hab.?
anisoloba, *Sweet, ex Steud. Nom.* ed. II. ii. 194, nomen.—Chili.

NICOTIANA:—
asiatica, Schult. Obs. Bot. 43 = rustica.
attenuata, Steud. Nom. ed. I. 554 = N. Tabacum.
attenuata, *Torr. ex S. Wats. in Bot. King's Exped.* 276. t. 27.—Am. bor. occ.
auriculata, Agardh, ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 60 = maxima.
auriculata, *Bertol. in Moris, Stirp. Sard. Elench. App.* 7.—Sardinia.
Australasiae, R. Br. in Tuckey, Congo, App. 472 = suaveolens.
axillaris, Lam. Tabl. Encyc. ii. 7 = Petunia nyctaginiflora.
Berteriana, *Hort. ex G. Don, in Sweet, Hort. Brit.* ed. III. 504, nomen; *Dun. in DC. Prod.* xiii. 1. 568.—Chili?
Bigelovii, *S. Wats. in Bot. King's Exped.* 276. t. 27.—Am. bor. occ.
bonariensis, *Lehm. Nicot.* 27.—Reg. Argent.
brasilia, Hort. ex Steud. Nom. ed. II. ii. 194 = rustica.
brasiliensis, Link & Otto, ex Loud. Hort. Brit. 594; Mart. Fl. Bras. x. 168 = alata.
breviflora, Gill. ex Miers, in Hook. Lond. Journ. Bot. v. (1846) 175 = Bouchetia erecta.
breviflora, Poir. Encyc. Suppl. iv. 94 = undulata.
capensis, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 856 = N. Tabacum.
caudata, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 181 = N. Tabacum?
Cavanillesii, *Dun. in DC. Prod.* xiii. 1. 572.—Mexic.
cerinthoides, Hornem. Hort. Hafn. i. 214 = paniculata.
cerinthoides, Vitm. ex Viv. Elench. Pl. Dinegro, 26 = plumbaginifolia.
chinensis, *Fisch. ex Lehm. Nicot.* 18.—China.
cirrhoides, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 180.—Chili.
Clevelandi, *A. Gray, Syn. Fl. N. Am.* ii. 1. 242.—Calif.
commutata, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* vi. 61.—Hab.?
cordifolia, *Phil. in Bot. Zeit.* xiv. (1856) 646.—Ins. J. Fernand.
corymbosa, *Remy, in C. Gay, Fl. Chil.* v. 57.—Chili.
crispa, Cav. Desc. 105 = Cavanillesii.
crispa, Desf. Tabl. ed. I. 67; Pers. Syn. i. 217 = plumbaginifolia.
crispa, *Phil. Fl. Atac.* 41.—Chili.
crispula, Hort. ex Steud. Nom. ed. II. ii. 194 = N. Tabacum.
decurrens, Agardh, Några ord om Tobaks, 17 = alata.
decurrens, *Tausch, in Flora*, xix. (1836) 388.—Hab.?
dilatata, *Link, Enum. Hort. Berol.* i. 179.—Am. austr.
diversifolia, *Nees, ex Dun. in DC. Prod.* xiii. 1. 572, nomen.—Hab.?
Doniana, *Dun. l. c.* 570.—Cuba.
dubia, Jacquem. Voy. Bot. 114 = Physochlaina praealta.
exasperata, *Lehm. in Hamb. Gartenz.* xiv. (1858) 439.—Hab.?
fastigiata, Nees, in Lehm. Pl. Preiss. i. 343 = suaveolens.
florida, Salisb. Prod. 132 = N. Tabacum.
Forsteri, Roem. & Schult. Syst. iv. 323 = suaveolens.
fragrans, Bernh. ex Steud. Nom. ed. I. 554, nomen = suaveolens?
fragrans, *Hook. Bot. Mag.* t. 4865.—N. Caled.
frigida, *Phil. Fl. Atac.* 41.—Chili.
frutescens, Hort. Amstel. ex Lehm. Nicot. 23 = N. Tabacum.
fruticosa, Forst. f. Prod. 17 = suaveolens?
fruticosa, Linn. Syst. ed. X. 932 = N. Tabacum.
fruticosa, Lour. Fl. Cochinch. 111 = chinensis.
fruticosa, Moç. & Sesse, ex Dun. in DC. Prod. xiii. 1. 559 = pilosa.
gigantea, Hort. Dorpat. ex Lehm. Nicot. 19 = N. Tabacum.
glandulosa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 6 = trigonophylla.
glauca, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.–June 1828) 175; *et in Bot. Mag.* t. 2837.—Reg. Argent.
glutinosa, *Linn. Sp. Pl.* 181.—Am. austr.
graciliflora, *A. Br. Del. Sem. Hort. Frib.* (1849) 3, *adnot.*; *ex Walp. Ann.* iii. 150.—Mexic.
gracilipes, Steud. Nom. ed. I. 554 = N. Tabacum.
guatemalensis, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 856 = N. Tabacum.
havanensis, Hort. Matrit. ex Lehm. Nicot. 21 = N. Tabacum.

NICOTIANA:—
hispida, Domb. ex Dun. in DC. Prod. xiii. 1. 568 = pusilla.
humilis, *Bertol. Misc. Bot.* xiii. 14.—Am. bor.
humilis, *Link, Enum. Hort. Berol.* i. 178.—Aegypt.
humilis, Mill. Gard. Dict. ed. VIII. n. 10 = pusilla.
humilis, Steud. Nom. ed. I. 554 = rustica.
ipomopsiflora, *Dun. in DC. Prod.* xiii. 1. 559.—Mexic.
ipomopsiflora, A. Gray, in Proc. Am. Acad. v. (1862) 166 = trigonophylla.
lancifolia, Agardh, ex Steud. Nom. ed. II. ii. 194 = chinensis.
lancifolia, Willd. ex Lehm. Nicot. 26 = N. Tabacum.
Langsdorffii, *Schrank, Pl. Rar. Hort. Monac.* t. 72; *Weinm. ex Roem. & Schult. Syst.* iv. 323.—Bras.
latissima, Mill. Gard. Dict. ed. VIII. n. 1 = N. Tabacum.
Lehmanni, Agardh, ex Dun. in DC. Prod. xiii. 1. 559 = N. Tabacum.
Lingua, Steud. Nom. ed. I. 554 = N. Tabacum.
linoides, Gill. ex Miers, in Hook. Lond. Journ. Bot. v. (1846) 174 = Leptoglossis linifolia.
longiflora, *Cav. Desc.* 106.—Chili.
loxensis, *H. B. & K. Nov. Gen. et Sp.* iii. 3.—Am. austr.
lychnoides, *Remy, in C. Gay, Fl. Chil.* v. 57.—Chili.
lyrata, *H. B. & K. Nov. Gen. et Sp.* iii. 4.—Cuba.
Macgillivrayi, Seem. Fl. Vit. 179 = fragrans.
macrophylla, Spreng. Ind. Hort. Hal. (1807) 45 = N. Tabacum.
marylandica, Schuebl. in Hort. Tubing. ex Dun. in DC. Prod. xiii. 1. 572 = N. Tabacum.
maxima, *Hoffmgg. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* vi. 60.—Hab.?
megalosiphon, Heurck & Muell. Arg. in Heurck, Obs. Bot. 126 = suaveolens.
mendocina, *Phil. in Linnaea*, xxxiii. (1864–65) 199.—Chili.
mexicana, *Schlecht. Ind. Sem. Hort. Halens.* (1840).—*Cf. Linnaea*, xv. (1841) *Litt.* 95.—Mexic.
micrantha, *Haw. ex Steud. Nom.* ed. II. ii. 194.—Hab.?
Miersii, *Remy, in C. Gay, Fl. Chil.* v. 56.—Chili.
militaris, Linn. Sp. Pl. ed. II. 259, in syn. = glutinosa.
minima, Molina, Sagg. Chile, 153 = Nierembergia repens.
minima, *Phil. in Linnaea*, xxxiii. (1864–65) 198.—Chili.
monticola, *Dun. in DC. Prod.* xiii. 1. 564.—Chili.
multiflora, *Phil. in Linnaea*, xxxiii. (1864–65) 197.—Chili.
multiflora, Torr. in Pacif. Rail. Rep. v. 362 = trigonophylla.
multivalvis, Lindl. Bot. Reg. t. 1057 = quadrivalvis.
nana, Lindl. l. c. t. 833 = Hesperochiron californicus.
Neesii, Lehm. Pl. Preiss. i. 344 = suaveolens.
nepalensis, *Link & Otto, ex Loud. Hort. Brit.* 594; *Dun. in DC. Prod.* xiii. 1. 572.—Nepal.
noctiflora, *Hook. Bot. Mag.* t. 2785.—Chili.
nudicaulis, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 128.—N. Mexic.
nyctaginiflora, Lehm. Nicot. 47 = Petunia nyctaginiflora.
obtusifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 129.—Mexic.
odorata, Hort.—Cf. Gard. Chron. (1881) ii. 183 = suaveolens?
otophora, *Griseb. in Goett. Abh.* xxiv. (1879) 243.—Reg. Argent.
oulophylla, *Dun. in DC. Prod.* xiii. 1. 560.—Chili.
pallescens, Steud. Nom. ed. I. 554 = N. Tabacum.
pallida, Salisb. Prod. 132 = paniculata.
Palmeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 242.—Arizona.
pandurata, Dun. in DC Prod. xiii. 1. 569 = repanda.
paniculata, *Linn. Sp. Pl.* 180.—Am. austr.
parviflora, Lehm. Nicot. 48 = Petunia parviflora.
parviflora, *Phil. in Linnaea*, xxxiii. (1864–65) 198.—Chili.
pauciflora, *Remy, in C. Gay, Fl. Chil.* v. 52.—Chili.
Pavoni, *Dun. in DC. Prod.* xiii. 1. 561.—Peruv.
persica, Lindl. Bot. Reg. t. 1592 = alata.
petiolaris, *Schlecht. ex Dun. in DC. Prod.* xiii. 1. 558.—Hab.?
petiolata, *Agardh, Några ord om Tobaks*, 17.—Am. austr.
petuniaeflora, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 209.—Ins. Guadalup.

NICOTIANA:—
pilosa, *Dun. in DC. Prod.* xiii. 1. 559.—Mexic.
plantaginea, *DC. ex Dun. l. c.*—Mexic.
plumbaginifolia, *Viv. Elench. Pl. Hort. Dinegro*, 26. t. 5.—Am. bor. et trop.
plumbaginifolia, Willd. Enum. Hort. Berol. 230 = Dunalia ramiflora.
Pseudodecurrens, Steud. Nom. ed. II. ii. 194 = alata.
pulmonarioides, H. B. & K. Nov. Gen. et Sp. iii. 2 = undulata.
pumila, Steud. Nom. ed. I. 554 = rustica.
pusilla, Blanco, Fl. Filip. ed. I. 100 = rustica.
pusilla, *Linn. Syst.* ed. X. 933.—Am. trop.
pusilla, Moç. & Sesse, ex Dun. in DC. Prod. xiii. 1. 559 = plantaginea.
quadrivalvis, *Pursh, Fl. Am. Sept.* i. 141.—Am. bor. occ.
repanda, Sims, Bot. Mag. t. 2484 = Doniana.
repanda, *Willd. ex Lehm. Nicot.* 40. t. 3.—Mexic.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 767 = repanda.
rotundifolia, Lindl. Bot. Reg. (1838) Misc. 59 = suaveolens.
rugosa, Mill. Gard. Dict. ed. VIII. n. 7 = rustica?
ruralis, Vell. Fl. Flum. 76; ii. 72 = Langsdorffii.
rustica, *Linn. Sp. Pl.* 180.—Mexic.
sanguinea, *Link & Otto, ex Loud. Hort. Brit.* 594; *ex Dun. in DC. Prod.* xiii. 1. 572.—Bras.
scabra, Hort. Matrit. ex Lehm. Nicot. 34 = rustica.
scapigera, *Phil. Fl. Atac.* 41.—Chili.
Sellowii, *Link & Otto, ex Steud. Nom.* ed. II. ii. 194; *Maxim. in Ind. Sem. Hort. Petrop.* (1869) 21.—Reg. Argent.
serotina, Steud. Nom. ed. I. 554 = N. Tabacum.
sibirica, Hort. ex Lehm. Nicot. 34 = rustica.
sileniflora, Hort. ex Dun. in DC. Prod. xiii. 1. 567 = acuminata.
silenoides, Dum.-Cours. Bot. Cult. ed. I. Suppl. 148 = acuminata.
solanifolia, *Walp. Rep.* iii. 12.—Chili.
sordida, *Lehm. Del. Sem. Hort. Hamb.* (1843) 8.—Mexic.
suaveolens, *Lehm. Nicot.* 43.—Austral.
tabaca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = seq.
Tabacum, *Linn. Sp. Pl.* 180.—Am. austr.
tatarica, Hort. ex Lehm. Nicot. 34 = rustica.
tenella, Cav. Desc. 105 = pusilla.
texana, *Hort. Par. ex Maxim. in Ind. Sem. Hort. Petrop.* (1869) 21.—Texas.
tomentosa, *Ruiz & Pav. Fl. Per.* ii. 16.—Peruv.
trigonophylla, *Dun. in DC. Prod.* xiii. 1. 562.—Am. bor. occ.
tristis, *Sm. in Rees, Cycl.* xv. n. 6.—Reg. Argent.
turcica, Haberl. ex Steud. Nom. ed. II. ii. 194, in syn. = chinensis.
turcica, Hort. ex Dun. in DC. Prod. xiii. 1. 564 = rustica.
ulophylla, Phil. Fl. Atac. 41 = oulophylla.
undulata, *Ruiz & Pav. Fl. Per.* ii. 16.—Peruv.
undulata, Vent. Jard. Malm. 10 = suaveolens.
urens, Linn. Syst. ed. X. 932 = Wigandia crispa.
Verdon, Steud. Nom. ed. I. 554 = N. Tabacum.
vincaeflora, *Lag. ex Link, Enum. Hort. Berol.* i. 179.—Am. austr.
virginica, *Agardh, Några ord om Tobaks*, 18.—Am. bor.
viridiflora, Hort. Madrit. ex Lehm. Nicot. 36 = paniculata.
viscosa, *Lehm. l. c.* 28.—Reg. Argent.
viscoso-cordata, Hort. ex Dun. in DC. Prod. xiii. 1. 567 = acuminata.
wigandioides, *C. Koch & Fint. Wochenschr.* i. (1858) 65.—N. Granat.
ybarrensis, H. B. & K. Nov. Gen. et Sp. iii. 4 = N. Tabacum.

NICOTIDENDRON, Griseb. in Abh. Goett. xix. (1874) 216 = **Nicotiana**, Linn. (Solan.).
glauca, Griseb. l. c. = Nicotiana glauca.

NICTANTHES, All. in Misc. Taur. ii. 50 (1761) = **Nyctanthes**, Linn. (Oleac.).

NICTITELLA, Rafin. Sylva Tellur. 128 (1838) = **Cassia**, Tourn. (Legumin.).
amena, Rafin. l. c. = C. nictitans.
aspera, Rafin. l. c. = C. nictitans.
mimosoides, Rafin. l. c. = C. mimosoides.

NIDORELLA, Cass. in Dict. Sc. Nat. xxxvii. 469 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 282.
altissima, *Benth. & Hook. f. Gen.* ii. 282, in obs. sub Microglossa.—Madag.
amplexicaulis, *DC. Prod.* v. 324.—Afr. austr.
anomala, *Steetz, in Peters, Reise Mossamb. Bot.* 400.—Afr. trop.
auriculata, *DC. Prod.* v. 322.—Afr. austr.
choënsis, *Delile, in Rochet, Sec. Voy. Choa*, 341.—Afr. trop.
Chrysocoma, *DC. Prod.* v. 322 (Conyzae sp. ?).—Madag.
compressa, *Cass. ex Steud. Nom.* ed. II. ii. 194.—Ins. Maurit.
conyzoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 90.—Afr. austr.
depauperata, Harv. l. c. = anomala.
diversifolia, Spreng. ex Sch. Bip. in Walp. Rep. ii. 958 = Sprengelii.
exilis, *DC. Prod.* v. 324.—Afr. austr.
floribunda, *Lehm. Ind. Sem. Hort. Hamb.* (1851) 8; *ex Steetz, in Linnaea*, xxv. (1852) 310.—Afr. austr.
foetida, DC. Prod. v. 323 = foliosa.
foliosa, *Cass. in Dict. Sc. Nat.* xxxvii. 470.—Afr. austr.
gariepina, *DC. Prod.* v. 324 (Feliciae sp. ?).—Afr. austr.
hirta, *DC. l. c.* 322.—Afr. austr.
hispida, *Benth. & Hook. f. Gen.* ii. 282, in obs. sub Microglossa.—Madag.
hottentotica, *DC. Prod.* v. 323.—Afr. austr.
hyssopifolia, DC. l. c. = foliosa.
Kraussii, Sch. Bip. in Walp. Rep. ii. 958 = obovata.
linifolia, *DC. Prod.* v. 322.—Afr. austr.
longifolia, DC. l. c. 324 = amplexicaulis.
membranifolia, Steetz, in Peters, Reise Mossamb. Bot. 400 = Microcephala.
mespilifolia, DC. Prod. v. 321 = Microglossa mespiloides.
microcephala, *Steetz, in Peters, Reise Mossamb. Bot.* 406.—Afr. trop.
mucronata, *DC. Prod.* v. 323.—Madag.
natalensis, Sch. Bip. ex Walp. Rep. ii. 958 = Microglossa mespiloides.
nubigena, *C. Bolle, in Bonplandia*, vii. (1859) 294.—Ins. Cap. Virid.
obovata, DC. Prod. v. 322 = auriculata.
otophora, *Griseb. in Goett. Abh.* xxiv. (1879) 243.—Reg. Argent.
pinnatilobata, *DC. Prod.* v. 321.—Afr. austr.
polycephala, *DC. l. c.* 323.—Afr. austr.
punctulata, DC. l. c. = Psiadia arabica.
rapunculoides, DC. l. c. = resedifolia.
resedifolia, *DC. l. c.* 322.—Afr. austr.
scandens, Sch. Bip. in Walp. Rep. ii. 958 = Microglossa mespiloides.
senecionidea, DC. Prod. v. 322 = auriculata.
solidaginea, *DC. l. c.*—Afr. austr.
Sprengelii, *Sch. Bip. in Walp. Rep.* ii. 958.—Afr. austr.
Steetzii, *J. A. Schmidt, Beitr. Fl. Cap Verd Ins.* 185.—Ins. Cap. Virid.
triloba, DC. Prod. v. 321 = Erigeron trilobus.
undulata, Sond. in Harv. & Sond. Fl. Cap. iii. 90 = amplexicaulis.
varia, *J. A. Schmidt, Beitr. Fl. Cap Verd. Ins.* 184.—Ins. Cap. Virid.
vernonioides, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 384.—Abyssin.

NIDULARIUM, Lem. Jard. Fleur. iv. (1854) Misc. 60 = **Karatas**, Adans. (Bromel.).
acanthocrater, E. Morr. in Belg. Hortic. xxxiv. (1884) 140. t. 9 = K. acanthocrater.
albo-roseum, Griseb. in Goett. Nachr. (1864) 12.—Venezuela.
amazonicum, Linden & Morr. Cat. (1873) 11.—Bras.
ampullaceum, E. Morr. in Belg. Hortic. xxx. (1880) 242.—Bras.
Antoineanum, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 113.—Bras.
Binoti, E. Morr. in Belg. Hortic. xxx. (1880) 90 = K. Binoti.
bivittatum, Lem. Illustr. Hortic. ix. (1862) Misc. 58 = Cryptanthus bivittatus.
Carolinae, Lem. in Rev. Hortic. Sér. IV. vii. (1858) 398 = K. Carolinae.
chlorostictum, E. Morr. in Belg. Hortic. xxviii. (1878) 207.—Bras. ?
coeruleum, Lem. in Rev. Hortic. (1861) 153.—Hab. ?

NIDULARIUM :—
cruentum, Regel, Gartenfl. (1859) 267 = Karatas cruenta.
cyaneum, Linden & André, Illustr. Hortic. xx. (1873) 184.—Bras.; N. Granat.
denticulatum, Regel, Gartenfl. (1870) 268.—Am. trop.
discolor, Beer, Bromel. 74 = K. purpurea.
Ferdinandocoburgi, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 112.—Bras.
fulgens, Lem. Jard. Fleur. iv. (1854) t. 411. Misc. 60 = K. fulgens.
giganteum, Baker, in Journ. Bot. xviii. (1880) 50.—Am. trop.
Innocentii, Lem. Illustr. Hortic. ii. (1855) Misc. 13 = K. Innocentii.
Johannis, Carr. in Rev. Hortic. (1884) 432.—Bras.
Karatas, Lem. ex Griseb. Fl. Brit. W. Ind. 591 = K. Plumieri.
Laurentii, Regel, Ind. Sem. Hort. Petrop. (1866) 80; Gartenfl. (1867) t. 529 = K. Laurentii.
Lindeni, Regel, Gartenfl. (1868) 78; Ann. Sc. Nat. Sér. V. x. (1869) 379.—Am. trop.
marmoratum, Hort.—Cf. Gard. Chron. (1869) 561.—Bras. ?
Meyendorffii, Regel, Gartenfl. (1859) 264 = K. Carolinae.
neglectum, André, in Illustr. Hortic. xxiv. (1877) 178.—Bras.
pictum, Hort.—Cf. Gard. Chron. (1869) 561 = K. fulgens.
Pinelianum, Lem. Illustr. Hortic. (1860) Misc. 71.—Bras.
purpureum, Beer, Bromel. 75; Regel, Ind. Sem. Hort. Petrop. (1857) 28 = K. purpurea.
rutilans, È. Morr. in Belg. Hortic. (1885) 81.—Bras.
sarmentosum, Regel, Gartenfl. (1870) 268.—Am. trop.
Scheremetiewii, Regel, Ind. Sem. Hort. Petrop. (1857) 28 = K. Scheremetiewii.
spectabile, Hort. in Rev. Hortic. (1872) 120 = K. Meyendorffii.
spectabile, T. Moore, in Gard. Chron. (1873) 8 = K. spectabilis.
splendens, Hort. ex Beer, Bromel. 78 = Caraguata lingulata.
triste, Regel, Gartenfl. xv. (1866) 356.—Bras.

NIEBUHRIA, DC. Prod. i. 243 (1824). *CAPPARIDEAE*, Benth. & Hook. f. i. 107.
acutifolia, E. Mey. in Drège, Zwei Pfl. Docum. 144 = Boscia caffra.
aethiopica, Fenzl, in Sitzb. Acad. Wien, Math.-Nat. li. Abth. II. (1865) 141 = Maerua aethiopica.
angustifolia, *Harv. Gen. S. Afr. Pl.* ed. II. 12.—Afr. austr.
arenaria, DC. Prod. i. 244 = Maerua ovalifolia.
avicularis, DC. l. c. 243 = triphylla.
caffra, DC. l. c. = triphylla.
linearis, *DC. l. c.* 244.—Ind. or.
linifolia, Voigt, Hort. Suburb. Calc. 74 = linearis.
madagascariensis, *DC. Prod.* i. 244.—Madag.
nervosa, Hochst. in Flora, xxvii. (1844) 289 = Maerua nervosa.
oblongifolia, DC. Prod. i. 244 = Maerua oblongifolia.
oblongifolia, Royle, Illustr. Bot. Himal. 73 = Maerua ovalifolia.
oleoides, *DC. Prod.* i. 244.—Afr. austr.
pedunculosa, Hochst. in Flora, xxvii. (1844) 289 = Boscia caffra.
rosmarinoides, *Sond. in Linnaea*, xxiii. (1850) 7.—Afr. austr.
siamensis, *Kurz, For. Fl. Brit. Burma*, i. 59.—Burma.
triphylla, *Wendl. f. in Bartl. & Wendl. f. Beitr.* ii. 29.—Afr. austr.
undulata, Zeyh. ex Sond. in Linnaea, xxiii. (1850) 8 = Boscia caffra.
variabilis, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 68.—Burma.
venosa, *Bunbury, in Hook. Lond. Journ. Bot.* iii. (1844) 261.—Afr. austr.
Woodii, *Oliver, in Hook. Ic. Pl.* t. 1386.—Afr. austr.

NIEBUHRIA, Neck. Elem. i. 30 (1790) = **Wedelia**, Jacq. (Compos.).

NIEBUHRIA, Scop. Introd. 134 (1777) = **Baltimora**, Linn. (Compos.).

NIEDERLEINIA, Hieron. in Roca, Exped. Rio-negro, ii. 183 (1881). *FRANKENIACEAE.*
juniperoides, *Hieron. l. c.*—Patagon.

NIEMEYERA, F. Muell. Fragm. vi. 96 (1867) = **Apostasia**, Blume (Orchid.).
stylidioides, F. Muell. l. c. = A. stylidioides.

NIEMEYERA, F. Muell. Fragm. vii. 114 (1870) = **Chrysophyllum**, Linn. (Sapot.).
prunifera, F. Muell. l. c. = C. pruniferum.

NIEREMBERGIA, Ruiz & Pav. Prod. 23 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 908.
Blencocoës, Rafin. Fl. Tellur. iii. 75 (1836).
Stimenes, Rafin. l. c. 76 (1836).
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 9. t. 198.—Am. austr.
anomala, Dun. in DC. Prod. xiii. 1. 588 = Bouchetia erecta, Petunia parviflora.
anomala, Miers, in Hook. Lond. Journ. Bot. v. (1846) 175 = Bouchetia erecta.
aristata, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. iii. t. 255.—Reg. Argent.
Athinsiana ×, Sweet, l. c. t. 268.
browallioides, *Griseb. in Goett. Abh.* xix. (1874) 214.—Reg. Argent.
calycina, *Hook. Bot. Mag.* t. 3371.—Reg. Argent.
calycina, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 752 = gracilis.
coerulea, *Gill. ex Miers, in Hook. Lond. Journ. Bot.* v. (1846) 173.—Chili.
Coulteri, *Hemsl. Biol. Centr. Am. Bot.* ii. 437.—Mexic.
ericoides, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 168.—Bras.
filicaulis, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 752 = gracilis.
filicaulis, Lindl. Bot. Reg. t. 1649 = linariaefolia.
frutescens, *Dur. Ind. Sem. Hort. Burdig.* (1866) 25.—Chili.
gracilis, *Hook. Bot. Mag.* t. 3108.—Reg. Argent.
graveolens, A. St. Hil. Pl. Rem. Brés. 221. t. 21 = Petunia viscidula.
hippomanica, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 168.—Reg. Argent.
intermedia, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1833) 175 = Salpiglossis linearis.
linariaefolia, *R. Grah. l. c.* (Jul.-Sept. 1831) 378.—Chili.
linifolia, Miers, in Hook. Lond. Journ. Bot. v. (1846) 174 = Leptoglossis linifolia.
micrantha, Domb. ex Dun. in DC. Prod. xiii. 1. 588 = Bouchetia erecta.
Miersiana, Sendt. in Mart. Fl. Bras. x. 200 = Bouchetia erecta.
montana, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 395.—Reg. Argent.
nana, Miers, in Hook. Lond. Journ. Bot. v. (1846) 177 = Hesperochiron californicus.
petiolata, Remy, in C. Gay, Fl. Chil. v. 46 = repens.
petunioides, *Dun. in DC. Prod.* xiii. 1. 588.—Peruv.
phoenicea, G. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 193 = Petunia violacea.
pinifolia, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 168.—Bras.
procumbens, Benth. & Hook. f. Gen. ii. 908 = Bouchetia procumbens.
prunellaefolia, *Dun. in DC. Prod.* xiii. 1. 583.—Chili.
pubescens, Spreng. Syst. i. 615 = Petunia viscidula.
pulchella, *Gill. ex Miers, in Hook. Lond. Journ. Bot.* v. (1846) 173.—Reg. Argent.
punicea, Hort. ex Dun. in DC. Prod. xiii. 1. 573, 589 = Petunia violacea.
repens, *Ruiz & Pav. Fl. Per.* ii. 13. t. 123.—Chili.
rigida, *Miers, Trav. Chile*, ii. 532; *et in Hook. Lond. Journ. Bot.* v. (1846) 172.—Reg. Argent.
rivularis, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 167.—Reg. Argent.
scoparia, *Sendt. in Mart. Fl. Bras.* x. 178.—Bras.
spathulata, H. B. & K. Nov. Gen. et Sp. iii. 8 = repens.
staticaefolia, Sendt. in Mart. Fl. Bras. x. 179 = Bouchetia erecta.
stricta, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 171.—Reg. Argent.
subdentata, *Meyen, Reise*, i. 314.—Chili.
Veitchii, *Berkeley, ex Hook. Bot. Mag.* t. 5599.—Am. trop.
viscidula, H. B. & K. Nov. Gen. et Sp. vii. 205 = Petunia viscidula.
viscosa, *Torr. Bot. Mex. Bound.* 155.—N. Mexic.

NIETNERIA, Klotzsch & Rich. Schomb. in Rich. Schomb. Reise Brit. Gui. 1066 (1848), nomen. *LILIACEAE*, Benth. & Hook. f. iii. 825.
corymbosa, *Klotzsch & Rich. Schomb. l. c.*, nomen.—Guiana.

NIETOA, Schaffn. in La Naturaleza, iii. (1876) 343 = **Hanburia**, Seem. (Cucurb.).
mexicana, Schaffn. l. c. = H. mexicana.

NIGELLA, [Tourn.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 8.
EROBATOS, Reichb. Handb. 277 (1837).
GARIDELLA, Tourn. ex Linn. Gen. ed. I. 131 (1737).
MELANTHIUM, Medic. Phil. Bot. i. 96 (1789).
NIGELLASTRUM, Moench, Meth. 311 (1794).
agrestis, J. & C. Presl, Delic. Prag. 225 = arvensis.
amoena, Salisb. Prod. 374 = hispanica.
aristata, Sibth. & Sm. Fl. Graec. Prod. i. 373 = arvensis.
armena, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 274 = segetalis.
arvensis, *Linn. Sp. Pl.* 584.—Europ.; Reg. Mediterr.; Orient.
arvensis, Pall. ex Bieb. Fl. Taur. Cauc. ii. 16 = segetalis.
aspera, C. Koch, in Linnaea, xix. (1847) 48 = arvensis.
assyriaca, *Boiss. Fl. Orient.* i. 67.—Assyria.
bicolor, Boiss. & Heldr. Diagn. Ser. I. viii. 7 = segetalis.
Bourgaei, Jord. Pugill. Pl. Nov. (1852) 2 = damascena.
Catenae, Tenore, Fl. Neap. Syll. App. v. 14 = arvensis.
ciliaris, *DC. Syst.* i. 327.—Syria; Ins. Cyprus.
cilicica, Boiss. & Bal. Diagn. Ser. II. v. 11 = stellaris.
coarctata, *C. C. Gmel. Fl. Bad.* ii. 502, in adnot.—Europ.
coerulea, Lam. Fl. Fr. iii. 312 = damascena.
confusa, Salle, ex Nym. Consp. 16 = hispanica.
corniculata, *DC. Syst.* i. 326.—Orient?
cretensis, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) I. 282, in textu = arvensis.
cretica, Mill. Gard. Dict. ed. VIII. n. 4 = arvensis.
damascena, *Linn. Sp. Pl.* 584.—Reg. Mediterr.
deserti, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 359.—Arabia.
divaricata, Beaupré, ex DC. Syst. i. 329 = arvensis.
diversifolia, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 220.—Turkest.
doliata, Pall. ex Bieb. Fl. Taur. Cauc. ii. 16 = arvensis.
elata, *Boiss. Diagn.* Ser. I. i. 66.—As. Min.; Ins. Cyprus.
elegans, Salisb. Prod. 374 = damascena.
foeniculacea, DC. Syst. i. 328 = arvensis.
foeniculacea, Hohen. ex Boiss. Fl. Orient. i. 65 = segetalis.
fumariaefola, *Kotschy, in Unger, Ins. Cyp.* 319.—Ins. Cyprus.
gaditana, Soleir. ex Nym. Consp. 16 = hispanica.
gallica, Jord. Pugill. Pl. Nov. (1852) 3 = hispanica.
Garidella, *Spenn. Monog. Nigell.* 11.—Europ.
glauca, Schkuhr, ex Wender. in Linnaea, xiv. (1840) 587 = arvensis.
hispanica, Gren. & Godr. Fl. Fr. i. 44 = arvensis.
hispanica, *Linn. Sp. Pl.* 584.—Hispan.; Afr. bor.
indica, Roxb. ex Flem. in As. Res. xi. (1810) 173 = sativa.
integrifolia, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) I. 246.—As. centr.
intermedia, C. Koch, in Linnaea, xix. (1847) 48 = elata.
involucrata, C. Koch, l. c. = elata.
involucrata, Moench, Meth. 314 = damascena.
laevis, Delile, ex DC. Syst. i. 329 = arvensis.
latifolia, Mill. Gard. Dict. ed. VIII. n. 5 = arvensis.
multifida, Gater. Desc. Pl. Montaub. 100 = damascena.
Nigellastrum, Willk. in Willk. & Lange, Fl. Hisp. iii. 963 = N. Garidella.
Noëana, Boiss. Diagn. Ser. II. i. 19 = oxypetala.
orientalis, *Linn. Sp. Pl.* 584.—As. Min.; Syria; Reg. Caucas.
oxypetala, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 357.—Oriens.
persica, Boiss. l. c. 358 = oxypetala.
polygynia, Moench, Meth. Suppl. 111 = hispanica.
pygmaea, DC. Syst. i. 333 = coarctata.
sativa, *Linn. Sp. Pl.* 584.—Reg. Mediterr.
segetalis, *Bieb. Fl. Taur. Cauc.* ii. 16.—Tauria; Oriens.
stellaris, *Boiss. Diagn.* Ser. I. viii. 8.—Syria.
taurica, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 283 = damascena.

NIGELLA :—
tenuiflora, Gilib. Fl. Lituan. ii. 285 = arvensis.
truncata, Viv. Pl. Aegypt. Dec. 6 = sativa.
tuberculata, Griseb. Spicil. Fl. Rumel. i. 310 = arvensis.
unguicularis, *Spenn. Monog. Nigell.* 12.—Syria.
unguiculata, Stokes, Bot. Mat. Med. iii. 229 = arvensis.
verrucosa, *C. Koch, in Linnaea*, xv. (1841) 246.—Reg. Caucas.

NIGELLASTRUM, Moench, Meth. 311 (1794) = **Nigella**, Tourn. (Ranunc.).
ciliare, Bercht. & Presl, Rostl. i. Ranunc. 102 = Nigella ciliaris.
corniculatum, Bercht. & Presl, l. c. = Nigella corniculata.
flavum, Moench, Meth. 312 = Nigella orientalis.
orientale, Bercht. & Presl, Rostl. i. Ranunc. 102 = Nigella orientalis.

NIGRINA, Linn. Mant. i. 5 (1767) = **Melasma**, Berg (Scrophul.).
vesicaria, Burm. f. Fl. Cap. Prod. 5 = M. scabrum?
viscosa, Linn. Mant. i. 42 = M. scabrum.

NIGRINA, Thunb. Nov. Gen. 58 (1783); Fl. Jap. 5 (1784) = **Chloranthus**, Sw. (1787).
serrata, Thunb. in Nov. Act. Soc. Sc. Upsal. vii. (1815) 142 = C. serratus.
spicata, Thunb. Nov. Gen. 59 = C. inconspicuus.
spicifera, Lam. Illustr. i. 295. t. 71 = C. inconspicuus.

NIGRITELLA, Rich. in Mém. Mus. Par. iv. (1818) 48 = **Habenaria**, Willd. (Orchid.).
angustifolia, Rich. l. c. 56 = H. nigra.
brachystachya, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 224 = H. nigra.
fragrans, Saut. ex Reichb. Ic. Pl. Crit. viii. f. 1010; Fl. Germ. Excurs. 121 = H. nigra.
globosa, Reichb. Fl. Germ. Excurs. 121 = Orchis globosa.
Heufleri, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 225 = H. nigra.
hybrida, Schur, Enum. Pl. Transs. 647 = H. nigra.
megastachya, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 222 = H. nigra.
micrantha ×, Kern. l. c. 227.—Europ.
nigra, Reichb. f. Ic. Fl. Germ. xxiii., xxiv. 102 = H. nigra.
suaveolens, Koch, Syn. Fl. Germ. ed. I. 690 = H. nigra.

NIGROLEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Ebermaiera**? Nees (Acanthac.).

NIL, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 210 = **Ipomoea**, Linn. (Convolv.).

NIMA, Buch.-Ham. ex A. Juss. in Mém. Mus. Par. xii. (1825) 516 = **Picrasma**, Blume (Simarub.).
quassioides, Buch.-Ham. ex A. Juss. l. c. 517 = P. quassioides.

NIMBO, Dennst. Schluess. Hort. Malab. 30 (1818) = **Murraya**, Linn. (Rutac.).
melioides, Dennst. l. c. = M. Koenigii.

NIMMOIA, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 13 = **Amoora**, Roxb. (Meliac.).
Lawii, Wight, l. c. = A. Lawi.

NIMMOIA, Wight, in Madr. Journ. Sc. vii. (1837) 311. t. 20 = **Ammannia**, Linn. (Lythrar.).
floribunda, Wight, l. c. 312 = A. floribunda.

NIMMONIA, Wight, Illustr. i. 206, in obs. (1840) = praec.

NIMPHAEA, Neck. Delic. Gallo-Belg. i. 227 (1768) = **Nymphaea**, Linn.

NIMPHEA, Nocca, in Usteri, Ann. Bot. v. (1793) 49 = praec.

NINANGA, Rafin. Fl. Tellur. iii. 76 (1836) = **Gomphrena**, Linn. et **Froelichia**, Moench (Amarant.).
bicolor, Rafin. l. c. 77 = G. perennis.
interrupta, Rafin. l. c. = F. lanata.

NINTOOA, Sweet, Hort. Brit. ed. II. 258 (1830) = **Lonicera**, Linn. (Caprifol.).
canescens, Webb, Iter Hisp. 42 = L. biflora.
confusa, Sweet, Hort. Brit. ed. II. 258 = L. confusa.
japonica, Sweet, l. c. = L. japonica.
longiflora, Sweet, l. c. = L. macrantha.

NIOBE, Salisb. in Trans. Hort. Soc. i. (1812) 335 = **Funkia**, Spreng. (Liliac.).
cordifolia, Salisb. l. c. = F. subcordata.
japonica, Salisb. Gen. Pl. Fragm. 82 = F. ovata.

NIOBEA, Willd. ex Schult. f. Syst. vii. 762 (1830) = **Hypoxis**, Linn.
nemorosa, Willd. l. c. = H. decumbens.
pratensis, Willd. l. c. = H. decumbens.

NIOTA, Adans. Fam. ii. 172 (1763) = **Ceropegia**, Linn. (Asclep.).

NIOTA, Lam. Illustr. t. 299 (1796?); Encyc. iv. 490 (1797) = **Samadera**, Gaertn. (Simarub.).
Commersonii, Pers. Syn. i. 416 = S. indica.
globosa, Blanco, Fl. Filip. ed. II. 214 = Claoxylon indicum.
Lamarkiana, Blume, Bijdr. 251, 260 = S. indica.
lucida, Wall. Pl. As. Rar. ii. t. 168 = S. lucida.
pentapetala, Poir. in Lam. Encyc. iv. 490 = S. indica.
polyandra, Wight & Arn. Prod. 63 = Brownlowia lanceolata.
tetrapetala, Poir. in Lam. Encyc. iv. 490 = S. indica.

NIOTOUTT, Adans. Voy. 162 (1759) = **Commiphora**, Jacq. (Burser.).

NIPA, Benth. & Hook. f. Gen. i. 484 (1865), sphalm. = NEPA, Webb = **Ulex**, Linn. (Legumin.).

NIPA, Thunb. in Vet. Akad. Nya Handl. Stockh. iii. (1782) 231; Nov. Gen. 90 (1784). *PALMAE*, Benth. & Hook. f. iii. 920.
NYPA, Wurmb. in Verh. Batav. Gen. i. (1779) 349, nomen prius.
arborescens, Wurmb. ex H. Wendl. in Kerch. Palm. 252 = fruticans.
fruticans, *Thunb. in Vet. Akad. Nya Handl. Stockh.* iii. (1782) 231.—Ost. Ganget.; Malaya.
litoralis, Blanco, Fl. Filip. ed. I. 662 = fruticans.

NIPHAEA, Lindl. Bot. Reg. (1841) Misc. 80. *GESNERACEAE*, Benth. & Hook. f. ii. 997.
MENEGHINIA, Vis. in Atti 8va Riun. Scienz. Ital. 560 (1846).
albo-lineata, Hook. Bot. Mag. t. 4282 = Phinaea albo-lineata.
anoectochilifolia, Warsz. ex Hanst. in Linnaea, xxvii. (1854) 705 = Phinaea albo-lineata.
argyroneura, Planch. & Linden, ex Hanst. l. c. = Phinaea albo-lineata.
caripensis, *Klotzsch, ex Hanst. l. c.* 704.—Venezuela.
crenata, *Karst. Fl. Columb.* i. 161. t. 80.—N. Granat.
cupreo-virens, *Lem. Illustr. Hortic.* vii. (1860) *Misc.* 72.—Mexic.
discolor, Warsz. ex Hanst. in Linnaea, xxvii. (1854) 705 = Phinaea albo-lineata.
elata, Lem. Hort. Univ. iii. (1841) 301 = oblonga?
oblonga, *Lindl. Bot. Reg.* (1841) *Misc.* 80.—Guatemala.
parviflora, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1851) *App.* 14; *et in Linnaea*, xxv. (1852) 299.—Guatemala.
pulchella, *Griseb. Cat. Pl. Cub.* 198.—Cuba.
Roezli, *Regel, Gartenfl.* (1877) 67.—Am. trop.
rubida, *Lem. in Fl. des Serres*, iii. (1847) t. 251.—Hab.?
rubra, Baxt. in Loud. Hort. Brit. Suppl. iii. 594 = praec.
Warszewiczii, Hort. Berol. ex Hanst. in Linnaea, xxvii. (1854) 705 = parviflora.

NIPHOGETON, Schlecht. in Linnaea, xxviii. (1856) 481 = **Oreosciadium**, Wedd. (Umbellif.).
andicola, Schlecht. l. c. = O. dissectum.

NIPHUS, Rafin. ex Steud. Nom. ed. II. i. 133, in syn. (1840) = **Aristolochia**, Tourn.

NIRARATHAMNOS, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 513. *UMBELLIFERAE.*
asarifolius, *Balf. f. l. c.*—Ins. Socotra.

NIRBISIA, G. Don, Gen. Syst. i. 63 (1831) = **Aconitum**, Linn. (Ranunc.).
Bisma, G. Don, l. c. = A. palmatum.
Codua, G. Don, l. c. 64 = A. ferox.
Hamiltonii, G. Don, l. c. = A. ferox.

NIRURI, Adans. Fam. ii. 356 (1763) = **Phyllanthus**, Linn. (Euphorb.).

NIRURIS, Rafin. Sylva Tellur. 91 (1838) = **Phyllanthus**, Linn. (Euphorb.).
annua, Rafin. l. c. = P. Niruri.
indica, Rafin. l. c. = P. Niruri.

NIRWAMIA, Rafin. Sylva Tellur. 35 (1838). *EUPHORBIACEAE*?—Cf. Thunb. Fl. Jap. 367. n. 98.
pellucida, Rafin. l. c. (Quid?).—Japon.

NISA, Noronha, ex Thou. Nov. Gen. Madag. 24 (1806) = **Homalium**, Jacq. (Samyd.).
albiflora, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 71 = H. albiflorum.
involucrata, DC. Prod. ii. 55 = H. involucratum.
leucophlaea, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 72.—Madag.
nudiflora, DC. Prod. ii. 55.—Madag.
retusa, Blume, Mus. Bot. Lugd. Bat. ii. 28.—Madag.
sanguinea, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 70.—Madag.
scleroxylon, Tul. l. c. 69.—Madag.

NISOMENES, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
diffusa, Rafin. l. c. = E. diffusa.

NISORALIS, Rafin. Sylva Tellur. 74 (1838) = **Helicteres**, Pluk. (Stercul.).
jamaicensis, Rafin. l. c. = H. jamaicensis.

NISSOLIA, Jacq. Enum. Pl. Carib. 7 (1760). *LEGUMINOSAE*, Benth. & Hook. f. i. 513.
aculeata, DC. Prod. ii. 250 = Machaerium aculeatum.
aculeata, Spreng. Syst. iii. 190 = Berteroniana.
aculeata, Vell. Fl. Flum. 296; vii. t. 79 = Machaerium Vellozianum.
aculeata, Willd. ex Vog. in Linnaea, xi. (1837) 194 = Machaerium Humboldtianum.
acuminata, DC. Prod. ii. 258 = Machaerium acuminatum.
acuminata, *Vell. Fl. Flum.* vii. t. 76.—Bras.
arborea, Jacq. Enum. Pl. Carib. 27 = Machaerium arboreum.
Arrabidae, Steud. Nom. ed. II. ii. 196 = Machaerium Vellozianum.
Berteroniana, *Steud. l. c.*—N. Granat.
bicallosa, *Vog. in Linnaea*, xi. (1837) 178.—Bras.?
debilis, Vell. Fl. Flum. 297; vii. t. 81 = Machaerium pedicellatum.
declinata, Vell. l. c. 296; vii. t. 77 = Machaerium discolor.
diadelpha, DC. Prod. ii. 258 = Machaerium ferrugineum.
dubia, *Poir. Encyc. Suppl.* iv. 99.—Guian.
ferruginea, Willd. Sp. Pl. iii. 900 = Machaerium ferrugineum.
firma, Vell. Fl. Flum. 297; vii. t. 83 = Machaerium firmum.
fruticosa, Humb. & Bonpl. ex Vog. in Linnaea, xi. (1837) 178, partim = bicallosa.
fruticosa, *Jacq. Enum. Pl. Carib.* 27.—Am. austr.
fruticosa, Vell. Fl. Flum. 298; vii. t. 86 = Machaerium nigrum.
glabrata, Link, Enum. Hort. Berol. ii. 221 = Machaerium arboreum.
hirsuta, DC. Prod. ii. 257 = fruticosa.
hirta, Vell. Fl. Flum. 296; vii. t. 78 = Machaerium eriocarpum.
incorruptibilis, Vell. l. c. 297; vii. t. 82 = Machaerium incorruptibile.
lanceolata, Vell. l. c. 299; vii. t. 87 = Machaerium secundiflorum.
legalis, Vell. l. c. 298; vii. t. 84 = Machaerium legale.
leiophylla, DC. Prod. ii. 258 = Machaerium leiophyllum.

NISSOLIA:—
microptera, Poir. Encyc. Suppl. iv. 98 = Machaerium micropterum.
nyctitans, Vell. Fl. Flum. 295; vii. t. 75 = Machaerium nictitans.
platycalyx, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 344.—N. Mexic.
platycarpa, *Benth. in Mart. Fl. Bras.* xv. 1. 77, in nota.—Mexic.
polyphylla, Poir. Encyc. Suppl. iv. 98 = Machaerium polyphyllum.
polysperma, Bert. ex DC. Prod. ii. 257 = fruticosa.
punctata, *Lam. Illustr.* ii. 159. t. 600. f. 1.—Am. austr.
quinata, Aubl. Pl. Gui. ii. 743. t. 297 = Machaerium ferrugineum.
racemosa, DC. Prod. ii. 257 = fruticosa.
reticulata, Lam. Illustr. ii. 160. t. 600 = Machaerium reticulatum.
reticulata, Vell. Fl. Flum. 299; vii. t. 88 = Pterocarpus violaceus.
retusa, Willd. Enum. Hort. Berol. 742 (Machaerii sp.?).—Am. austr.
robiniaefolia, DC. Prod. ii. 258 = Machaerium robiniaefolium.
robusta, Vell. Fl. Flum. 298; vii. t. 85 = Centrolobium robustum.
scandens, *Koen. ex Spreng. Syst.* iii. 191.—Ind. or.
Schottii, *A. Gray, in Journ. Linn. Soc.* v. (1861) 26.—Mexic.
spinosa, Vell. Fl. Flum. 296; vii. t. 80 = Machaerium spinosum.
stipitata, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 99 = Machaerium stipitatum.
uncinata, Vell. Fl. Flum. 295; vii. t. 76 = Machaerium uncinatum.
Wislizeni, *A. Gray, in Journ. Linn. Soc.* v. (1861) 25.—Mexic.

NISSOLIA, [Tourn.] Linn. Syst. ed. I (1735) = **Lathyrus**, Tourn. (Leguminin.).
parviflora, Moench, Meth. Suppl. 49 = L. inconspicuus.
uniflora, Moench, Meth. 140 = L. Nissolia.

NISSOLIUS, Medic. in Vorles. Churpf. Phys. Gen. ii. (1787) 389 = **Nissolia**, Jacq. (Legumin.).

NITELIUM, Cass. in Dict. Sc. Nat. xxxv. 11 (1825) = **Dicoma**, Cass. (Compos.).
rubescens, Cass. l. c. 12 = D. diacanthoides.

NITRAPIA, Pall. Reise, i. 494 (1771), err. typ. = seq.

NITRARIA, Linn. Syst. ed. X. 1044 (1759). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 265.
Billardieri, DC. Prod. iii. 456 = Schoberi.
caspica, Willd. ex Pall. Fl. Ross. t. 50 = Schoberi.
Oliverii, Jaub. & Spach, Illustr. iii. 143 = Schoberi.
retusa, Aschers. in Verh. Bot. Ver. Brandenb. xviii. (1876) 94 = tridentata.
Schoberi, *Linn. Syst.* ed. X. 1044.—Rossia; As. bor.; Austral.
senegalensis, Poir. in Lam. Encyc. iv. 493 = tridentata.
sericea, Jaub. & Spach, Illustr. iii. 142 = tridentata.
sibirica, Poir. in Lam. Encyc. iv. 493 = Schoberi.
sphaerocarpa, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 74.—Mongol.
tridentata, *Desf. Fl. Atlant.* i. 372.—Syria; Afr. bor. et trop.

NITROPHILA, S. Wats. in Bot. King's Exped. 297 (1871). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 59.
occidentalis, *S. Wats. l. c.*—Am. bor. occ.

NIVARIA, Heist. Syst. 5, 19 (1748); Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 421 = **Leucojum**, Linn. (Amaryll.).
aestivalis, Moench, Meth. Suppl. 93 = L. aestivum.
hexanthera, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 422 = L. vernum.
monadelphia, Medic. l. c. = L. aestivum.
verna, Moench, Meth. 279 = L. vernum.

NIVENIA, R. Br. in Trans. Linn. Soc. x. (1810) 133. *PROTEACEAE*, Benth. & Hook. f. iii. 171.
PARANOMUS, Salisb. Parad. Lond. sub n. 66 (1805), nomen prius.
alopecuroides, Lam. ex St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 130 = N. Sceptrum.
candicans, R. Br. in Trans. Linn. Soc. x. (1810) 221 = Serruria candicans?
candicans, Roem. & Schult. Syst. iii. 389 = Serrura albicans.
capitata, *R. Br. in Trans. Linn. Soc.* x. (1810) 138.—Afr. austr.
concava, *R. Br. l. c.* 221; *Roem. & Schult. Syst.* iii. 388.—Afr. austr.
crithmifolia, *R. Br. l. c.* 136.—Afr. austr.
Dregei, *Buek, ex Meissn. in DC. Prod.* xiv. 300.—Afr. austr.
intermedia, Steud. Nom. ed. II. ii. 196 = media.
lagopus, *R. Br. in Trans. Linn. Soc.* x. (1810) 137.—Afr. austr.
marginata, *R. Br. l. c.* 134.—Afr. austr.
marginata, Drège, ex Meissn. in DC. Prod. xiv. 300 = spathulata.
media, *R. Br. in Trans. Linn. Soc.* x. (1810) 137.—Afr. austr.
mollissima, *R. Br. l. c.* 138.—Afr. austr.
parvifolia, *R. Br. l. c.* 135.—Afr. austr.
parvifolia, Buek, ex Meissn. in DC. Prod. xiv. 300 = spathulata.
Sceptrum, *R. Br. in Trans. Linn. Soc.* x. (1810) 134.—Afr. austr.
spathulata, *R. Br. l. c.* 135.—Afr. austr.
spathulata, Drège, ex Meissn. in DC. Prod. xiv. 300 = parvifolia.
spicata, *R. Br. in Trans. Linn. Soc.* x. (1810) 136.—Afr. austr.

NIVENIA, Vent. Dec. Gen. Nov. 3 (1808) = **Aristea**, Soland. (Irid.).
binata, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 382 = A. corymbosa (in Addend.).
corymbosa, Baker, in Journ. Linn. Soc. xvi. (1877) 109 = A. corymbosa.
filamentosa, Salisb. in Trans. Hort. Soc. i. (1812) 311 = A. corymbosa?
fruticosa, Baker, in Journ. Linn. Soc. xii. (1877) 109 = A. fruticosa.
stylosa, Salisb. in Trans. Hort. Soc. i. (1812) 311 = A. corymbosa.

NOAEA, Moq. in DC. Prod. xiii. II. 207 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 72.
NOEA, Boiss. & Bal. Diagn. Ser. II. iv. 76 (1859).
aretioides, Bunge, Anab. Rev. 35 = Anabasis aretioides.
Aucheri, Moq. in DC. Prod. xiii. II. 207 = Salsola canescens.
canescens, Moq. l. c. 208 = Salsola canescens.
daghestanica, *Turcz. ex Bunge, Anab. Rev.* 26.—Reg. Caucas.
Griffithii, *Bunge, l. c.* 22.—Afghan.
major, *Bunge, l. c.* 27.—Afghan.
malacophylla, Moq. in DC. Prod. xiii. II. 207 = Halimocnemis Kirilowii.
minuta, *Boiss. & Bal. Diagn.* Ser. II. iv. 76.—Cappadoc.
oppositiflora, Moq. in DC. Prod. xiii. II. 209 = Girgensohnia oppositiflora.
oppositifolia, Boiss. ex Bunge, Anab. Rev. 30 = Girgensohnia fruticulosa.
Regelii, *Bunge, in Act. Hort. Petrop.* vi. (1879) 440.—Turkest.
spinosissima, *Moq. in DC. Prod.* xiii. II. 209.—Reg. Mediterr.; Oriens.
tomentosa, Moq. l. c. 208 = Salsola canescens, aurantiaca.
Tournefortii, *Moq. l. c.*—As. Min.; Persia.

NOBULA, Adans. Fam. ii. 145 (1763) = **Phyllis**, Linn. (Rubiac.).

NOCCA, Cav. Ic. iii. 12. t. 224 (1794) = **Lagascea**, Cav. (Compos.).
helianthifolia, Cass. in Dict. Sc. Nat. xxv. 104 = L. suaveolens.
latifolia, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 31 = L. suaveolens.
macrophylla, Zucc. ex DC. Prod. v. 92 = L. suaveolens.
mollis, Jacq. Fragm. 58 = L. mollis.
rigida, Cav. Ic. iii. 12 = L. rubra.
rubra, Cass. in Dict. Sc. Nat. xxv. 104 = L. rubra.
suaveolens, Cass. l. c. 105 = L. suaveolens.

NOCCAEA, Moench, Meth. Suppl. 89 (1802); Reichb. Fl. Germ. Excurs. 663 (1832); et Ic. Fl. Germ. ii. 5. *CRUCIFERAE*, Benth. & Hook. f. i. 86.
CRUCIUNDULA, Rafin. Fl. Tellur. ii. 100 (1836).
alpina, Reichb. Fl. Germ. Excurs. 663 = Hutchinsia alpina.
Auerswaldii, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 780.—Hispan.
brevicaulis, Reichb. Fl. Germ. Excurs. 663 = Hutchinsia alpina.
brevistyla, Steud. Nom. ed. II. ii. 196 = Thlaspi rivale.
cepaeaefolia, Reichb. Fl. Germ. Excurs. 663 = Thlaspi cepaeaefolium.
hastulata, Steud. Nom. ed. II. ii. 196 = Thlaspi hastulatum.
petraea, Steud. l. c. 27, in syn., sphalm. = seq.
procumbens, Reichb. Fl. Germ. Excurs. 663 = Capsella procumbens.
pumila, Steud. Nom. ed. II. ii. 197 = Thlaspi pumilum.
rotundifolia, Moench, Meth. Suppl. 89 = Thlaspi rotundifolium.
stylosa, *Reichb. Fl. Germ. Excurs.* 663.—Europ. austr.
trinervia, Steud. Nom. ed. II. ii. 197 = Iberidella trinervia.

NOCCAEA, Willd. Sp. Pl. iii. 2393 (1803); Jacq. Fragm. 58. t. 85 (1805) = NOCCA, Cav. = **Lagascea**, Cav. (Compos.).

NOCHOTTA, S. G. Gmel. Reise, iii. 23. t. 3 (1774) = **Cicer**, Tourn. (Legumin.).
oleracea, S. G. Gmel. l. c. = C. arietinum.

NODOCARPAEA, A. Gray, in Proc. Am. Acad. xix. (1883) 79. *RUBIACEAE.*
radicans, *A. Gray, l. c.*—Cuba.

NOËA, Boiss. & Bal. Diagn. Ser. II. iv. 76 (1859) = **Noaea**, Moq. (Chenop.).

NOISETTIA, H. B. & K. Nov. Gen. et Sp. v. 382. 499 (1821). *VIOLARIEAE*, Benth. & Hook. f. t. i. 116.
IONIDIOPSIS, Presl, Bot. Bemerk. 13 (1844).
VIOLAEOIDES, Michx. ex DC. Prod. i. 280 (1824).
acuminata, DC. Prod. i. 290 = Ionidium concolor.
frangulaefolia, H. B. & K. Nov. Gen. et Sp. v. 384. t. 499 (= *Calyptrion frangulaefolium*).—Peruv.
galeopsifolia, A. St. Hil. Pl. Rem. Brés. 287 = longifolia.
longifolia, *H. B. & K. Nov. Gen. et Sp.* v. 384, in obs.—Guiana.
orchidiflora, Ging. in DC. Prod. i. 290 = longifolia.
orinocensis, H. B. & K. Nov. Gen. et Sp. v. 384 = Corynostylis Hybanthus.
pyrifolia, Mart. Nov. Gen. et Sp. i. 24 = Anchietea salutaris.
Roquefeuilliana, A. St. Hil. Pl. Rem. Brés. 289 = Anchietea salutaris.

NOLANA, Linn. Sp. Pl. ed. II. 202 (1762). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 879.
GUBLERIA, Gaudich. Bot. Voy. Bonite, t. 104 (1844–52).
NEUDORFIA, Adans. Fam. ii. 219 (1763).
SOREMA, Lindl. Bot. Reg. (1844) sub t. 46.
TEGANIUM, Schmid. Ic. 67. t. 18 (1747).
WALKERIA, Mill. ex Ehret, in Phil. Trans. liii. (1763) 130.
ZWINGERA, Neck. Elem. iii. 385 (1790).
acuminata, *Miers, ex Dun. in DC. Prod.* xiii. 1. 12.—Chili.
angustifolia, *Phil. in Linnaea*, xxix. (1857–58) 26.—Chili.
atriplicifolia, *Hort. ex D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 305.—Peruv.
baccata, *Dun. in DC. Prod.* xiii. 1. 14.—Chili.
carnosa, *Miers, ex Dun. l. c.*—Chili.
Chastenayana, *Gaudich. Bot. Voy. Bonite*, t. 33.—Peruv.?
coelestis, *Miers, ex Dun. in DC. Prod.* xiii. 1. 13.—Chili.
cordata, *Dun. l. c.*—Bolivia; Peruv.
coronata, *Ruiz & Pav. Fl. Per.* ii. 7. t. 112.—Peruv.
crassulifolia, Kunze, ex Walp. Rep. vi. 551 = Dolia tomentosa.
ericifolia, *Miers, ex Dun. l. c.* 14.—Chili.

NOLANA:—
fruticosa, Penny, Hort. Epsom. 34; ex Loud. Gard. Mag. v. (1829) 471 = tenella.
gallinacea, Pers. Syn. i. 168 = prostrata.
glandulosa, *Miers, ex Dun. in DC. Prod.* xiii. 1. 14.—Chili.
grandiflora, Lehm. ex G. Don, Gen. Syst. iv. 479 = atriplicifolia.
inflata, *Ruiz & Pav. Fl. Per.* ii. 7. t. 112.—Peruv.
lanceolata, *Miers, ex Dun. in DC. Prod.* xiii. 1. 12.—Chili.
linearis, *Miers, ex Dun. l. c.* 13.—Chili.
littoralis, *Miers, ex Dun. l. c.* 12.—Chili.
longifolia, *Miers, ex Dun. l. c.*—Chili.
microphylla, Miers, ex Dun. l. c. 15 = Phrodus microphyllus.
obtusa, *Miers, ex Dun. l. c.* 14.—Chili.
paradoxa, Bert. ex Dun. l. c. 17 = Dolia tomentosa.
paradoxa, Hook. Bot. Mag. t. 2604 = tenella.
paradoxa, *Lindl. in Bot. Reg.* t. 865.—Chili.
prostrata, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 11 = paradoxa.
prostrata, *Linn. Sp. Pl.* ed. II. 202.—Peruv.
revoluta, Mathews, ex Dun. in DC. Prod. xiii. 1. 17 = Dolia tomentosa.
revoluta, Ruiz & Pav. Fl. Per. ii. 8. t. 113. f. b = Dolia revoluta.
rostrata, *Miers, ex Dun. in DC. Prod.* xiii. 1. 14.—Chili.
rupicola, *Gaudich. Bot. Voy. Bonite*, t. 28.—Hab.?
sedifolia, Kunze, ex Walp. Rep. vi. 550 = Dolia vermiculata.
spathulata, *Ruiz & Pav. Fl. Per.* ii. 7. t. 112.—Peruv.
tenella, *Lindl. in Trans. Hort. Soc.* vii. (1830) 252.—Chili.
triquetra, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) 12.—Hab.?

NOLINA, Michx. Fl. Bor. Am. i. 207 (1803). *LILIACEAE*, Benth. & Hook. f. iii. 780.
BEAUCARNEA, Lem. Illustr. Hortic. viii. (1861) Misc. 59.
NOLINAEA, Baker, in Journ. Linn. Soc. xiii. (1873) 292.
NOLINEA, Pers. Syn. i. 399 (1805).
PINCENECTITIA, Hort. ex Lem. Illustr. Hortic. viii. (1861) Misc. 48, 59.
ROULINIA, Brongn. in Ann. Sc. Nat. Sér. II. xiv. (1840) 320.
Bigelovii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 247.—Arizona.
erumpens, *S. Wats. l. c.* 248.—Texas.
georgiana, *Michx. Fl. Bor. Am.* i. 208.—Am. bor.
Hartwegiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 371.—Mexic.
humilis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 248.—Mexic.
humilis, S. Wats. l. c. 248, partim = Watsoni.
javanica, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 120 = Chlorophytum laxum.
Lindheimeriana, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 247.—Mexic.
longifolia, *Hemsl. Biol. Centr. Am. Bot.* iii. 372.—Mexic.
microcarpa, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 247.—Arizona.
Palmeri, *S. Wats. l. c.* 248.—Calif.
Parryi, S. Wats. l. c. 247 = Bigelovii.
parviflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 372.—Mexic.
recurvata, *Hemsl. l. c.*—Mexic.
texana, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 248.—Texas.
Watsoni, *Hemsl. Biol. Centr. Am. Bot.* iii. 372.—Mexic.

NOLINAEA, Baker, in Journ. Linn. Soc. xiii. (1873) 292 = praec.

NOLINEA, Pers. Syn. i. 399 (1805) = praec.

NOLLETIA, Cass. in Dict. Sc. Nat. xxxvii. 479 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 285.
LEPTOTHAMNUS, DC. Prod. v. 367 (1836).
chrysocomoides, *Cass. in Dict. Sc. Nat.* xxxvii. 479.—Afr. bor. occ.
ciliaris, *Steetz, in Peters, Reise Mossamb. Bot.* 404.—Afr. austr.
ligulata, *Steetz, l. c.*—Afr. austr.
rarifolia, *Steetz, l. c.*—Afr. austr.

NOLTEA, Reichb. Consp. 145 (1828). *RHAMNEAE*, Benth. & Hook. f. i. 381.
HOLLIA, Heynh. Nom. i. 880 (1840); ii. 803 (1846).
VITMANNIA, Wight & Arn. Prod. 166 (1834).
WILLEMETIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 370. t. 16. f. 1.
africana, *Reichb. ex Harv. & Sond. Fl. Cap.* i. 478.—Afr. austr.

NOLTIA, Benth. & Hook. f. Gen. i. 381 (1862), sphalm. = praec.

NOLTIA, Eckl. ex Steud. Nom. ed. II. ii. 197 (1841) = **Selago**, Linn.

NOLTIA, Schum. & Thonn. Beskr. Guin. Pl. 189 (1827) = **Diospyros**, Linn. (Ebenac.).
tricolor, Schum. & Thonn. l. c. = D. tricolor.

NOMAPHILA, Blume, Bijdr. 804 (1826). *ACANTHACEAE*, Benth. & Hook. f. ii. 1075.
ciliata, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 21.—Afr. trop.
corymbosa, *Blume, Bijdr.* 804.—Java.
glandulosa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 198.—Afr. trop.
laevis, *Nees, in DC. Prod.* xi. 85.—Afr. trop.
Lyalliana, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 22.—Madag.
nodosa, Zipp. ex Span. in Linnaea, xv. (1841) 329 = petiolata.
Parishii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 455.—Burma; Java.
petiolata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 387.—Ins. Timor.
pinnatifida, Dalz. in Hook. Kew Journ. iii. (1851) 38 = Cardanthera pinnatifida.
pubescens, Kurz, in Journ. As. Soc. Beng. xl. (1871) ii. 74 = Parishii.
quadrangularis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 197.—Afr. trop.
stricta, *Nees, in DC. Prod.* xi. 84, excl. syn.—Ind. or.?; Malaya.

NOMISMIA, Wight & Arn. Prod. 236 (1834) = **Rhynchosia**, Lour. (Legumin.).
aurea, Wight & Arn. l. c. 237 = R. aurea.
capitata, Wight & Arn. l. c. = R. aurea.
nummularia, Wight & Arn. l. c. = R. nummularia.

NOMOCHLOA, Beauv. in Lestib. Essai Fam. Cyp. 37 (1819) = **Scirpus**, Linn. (Cyperac.).
compressa, Beauv. l. c. = S. compressus.

NOMOCHLOA, Nees, in Linnaea, ix. (1834) 299; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 110 = **Pleurostachys**, Brongn. (Cyperac.).
elegans, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 113 = P. elegans.
Orbigniana, Nees, in Linnaea, ix. (1834) 299 = P. Orbigniana.
stricta, Nees, in Hook. Journ. Bot. ii. (1840) 398 = P. stricta.
tenuiflora, Nees, in Linnaea, ix. (1834) 299 = P. tenuiflora.
turbinata, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 112 = P. tenuiflora.

NONATELIA, Aubl. Pl. Gui. i. 182. t. 70 (1775) = **Palicourea**, Aubl. (Rubiac.).
clusiaefolia, Reichb. ex DC. Prod. iv. 467, 532 = Chasalia clusiaefolia.
divaricata, Bartl. ex DC. Prod. iv. 466 = *panamensis*.
divaricata, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 207.—Guian.
filamentosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 137 = Hamiltonia suaveolens.
formosa, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 207.—Guian.
grandiflora, H. B. & K. Nov. Gen. et Sp. iii. 422.—Am. austr.
hispida, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 187 = Lasianthus Wallichii.
longiflora, Aubl. Pl. Gui. i. 185. t. 71 = P. longiflora.
lutea, Aubl. l. c. 190. t. 74.—Guian.
macrophylla, H. B. & K. Nov. Gen. et Sp. iii. 423.—Venezuela.
officinalis, Aubl. Pl. Gui. i. 182. t. 79 = Psychotria officinalis.

NONATELIA:—
officinalis, Spreng. ex DC. Prod. iv. 515 = Psychotria brachiata.
panamensis, DC. l. c. 466.—Panama.
paniculata, Aubl. Pl. Gui. i. 184 = Psychotria flexuosa.
pubescens, Spreng. Syst. i. 751 = Psychotria pubescens.
racemosa, Aubl. Pl. Gui. i. 186. t. 72 = Psychotria racemosa, *Willd.*
secundiflora, Spreng. Syst. i. 751 = Retiniphyllum secundiflorum.
violacea, Aubl. Pl. Gui. i. 188. t. 73 = Palicourea violacea.

NONEA, Medic. = seq.

NONNEA, (Nonea) Medic. Phil. Bot. i. 31 (1789); Moench, Meth. 421 (1794); Bernh. Syst. Verz. Erf. 98 (1800). *BORAGINEAE*, Benth. & Hook. f. ii. 856.
ECHIOIDES, Desf. Fl. Atlant. i. 163 (1800).
ELIZALDIA, Willk. Strand- u. Steppengeb. Iber. Halb. 128 (1852).
ONOCHILIS, Mart. in Denkschr. Akad. Muench. v. (1817) 177.
OSKAMPIA, Moench, Meth. 420 (1794).
aegyptiaca, Moench, ex Steud. Nom. ed. I. 556 = Anchusa flava.
alba, *DC. Fl. Fr. Suppl.* 420.—Europ. austr.; Oriens.
alpestris, *G. Don, Gen. Syst.* iv. 336.—Reg. Caucas.
amplexicaulis, Hochst. ex Boiss. Fl. Orient. iv. 170 = alba.
anchusoides, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 152.—Persia.
armena, Boiss. & Huet, ex Boiss. Fl. Orient. iv. 167 = pulla.
armena, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 574 = melanocarpa.
arvensis, DC. ex Steud. Nom. ed. I. 556 = Lycopsis arvensis.
atra, Griseb. Spicil. Fl. Rumel. ii. 94 = pulla.
auriculata, Boiss. Diagn. Ser. I. xi. 95 = Pulmonaria mollis.
Bourgaei, Coss. Notes Crit. 122 = micrantha.
capensis, G. Don, Gen. Syst. iv. 338 = Lycopsis capensis.
caspica, G. Don, l. c. 336 = picta.
ciliata, *DC. Fl. Fr.* iii. 626; *Roem. & Schult. Syst.* iv. 71, 754.—Oriens.
Colsmanniana, *G. Don, Gen. Syst.* iv. 337.—Hab.?
confertiflora, G. Don, l. c. 338 = Lycopsis confertiflora.
decumbens, *Moench, Meth.* 422.—Reg. Mediterr. occ.
decurrens, *G. Don, Gen. Syst.* iv. 337; *Boiss. Fl. Orient.* iv. 168.—Reg. Caucas.
diffusa, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 152 = picta.
dubia, Steud. Nom. ed. II. i. 85 = Alkanna lutea.
echioides, Roem. & Schult. Syst. iv. 71 = Macrotomia echioides.
Edgeworthii, A. DC. in DC. Prod. x. 30 = pulla.
elongata, *G. Don, Gen. Syst.* iv. 337.—Aegypt.
erecta, Bernh. Syst. Verz. Erf. 127 = pulla.
euphratica, Boiss. Diagn. Ser. I. xi. 97 = macrosperma.
flavescens, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 17; G. Don, Gen. Syst. iv. 337 = Alkanna lutea.
grandiflora, Steud. Nom. ed. II. ii. 197 = rosea.
intermedia, *Ledeb. Fl. Ross.* iii. 111.—Reg. Caucas.
lamprocarpa, Griseb. Spicil. Fl. Rumel. ii. 93 = obtusifolia.
longiflora, *Wettstein, in Denkschr. Acad. Wien*, l. (1885) 27.—Persia.
lutea, Bory & Chaub. Nouv. Fl. Pélop. 12; Auch. ex A. DC. in DC. Prod. x. 98 = Alkanna graeca.
lutea, DC. Fl. Fr. iii. 626; Suppl. 420 = Alkanna lutea.
macrosperma, *Boiss. & Heldr. Diagn.* Ser. I. xi. 97.—As. Min.
macrosperma, Griseb. Spicil. Fl. Rumel. ii. 93 = alba.
melanocarpa, *Boiss. Diagn.* Ser. I. xi. 96.—Syria; Persia.
micrantha, *Boiss. & Reut. Diagn. Pl. Hisp.* 21.—Hispan.
multicolor, *Kunze, in Flora*, xxix. (1846) 691.—Hispan.
nigricans, DC. Fl. Fr. iii. 626 = decumbens.
obtusifolia, *DC. l. c.*; *Roem. & Schult. Syst.* iv. 71.—Graec.; As. Min.
persica, *Boiss. Diagn.* Ser. I. vii. 32.—Persia.
phaneranthera, *Viv. Fl. Lib. Spec.* 9. t. 1. f. 3.—Afr. bor.
philistaea, *Boiss. Diagn.* Ser. I. xi. 96.—Palestina.
picta, *Sweet, Hort. Brit.* ed. I. 292.—Reg. Caucas.; Soongar.
pulla, *DC. Fl. Fr.* iii. 626.—Europ.; As. occ.; Ind. or.
pulmonarioides, *Boiss. & Bal. ex Boiss. Pl. Or. Nov. Dec.* ii. 4.—As. Min.
purpurea, Opiz, in Lotos, iii. (1853) 212 = pulla?
Reuteriana, *Boiss. Diagn.* Ser. I. xi. 95.—Hab.?
rosea, *Link, Enum. Hort. Berol.* i. 167.—Reg. Caucas.
rossica, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 572.—Rossia.
rubra, *Steud. Nom.* ed. II. ii. 198.—Italia.
Schultesii, G. Don, Gen. Syst. iv. 338 = phaneranthera.
setosa, Roem. & Schult. Syst. iv. 754 = Alkanna lutea.
Sibthorpiana, G. Don, Gen. Syst. iv. 336 = alba.
sordida, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 43 = picta.
stenantha, Opiz, in Lotos, iii. (1853) 213 (Quid?).—Europ.
stenophylla, Opiz, l. c. ii. (1852) 229 (Quid?).—Europ.
stenosolen, *Boiss. & Bal. Diagn.* Ser. II. vi. 123.—Cappadoc.
Szovitziana, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 574 = picta.
taurica, Ledeb. Fl. Ross. iii. 112 = pulla.
ventricosa, Griseb. Spicil. Fl. Rumel. ii. 93 = alba.
versicolor, Sweet, Hort. Brit. ed. I. 292 = rosea.
vesicaria, Reichb. Fl. Germ. Excurs. 338 = decumbens.
violacea, Auch. ex A. DC. in DC. Prod. x. 100 = Alkanna tinctoria.
violacea, DC. Fl. Fr. iii. 626 = decumbens.
Viviani, *A. DC. in DC. Prod.* x. 31.—Aegypt.

NONNIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = praec.

NOPALEA, Salm-Dyck, Cact. Hort. Dyck. ed. II. 63, 233 (1850). *CACTEAE*, Benth. & Hook. f. i. 850.
angustifrons, Lindb. in Act. Soc. Sc. Fenn. x. (1871) 123 = Opuntia leucacantha?
Auberi, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 64, 233.—Cuba.
coccinellifera, *Salm-Dyck, l. c.* 64.—Mexic.
dejecta, *Salm-Dyck, l. c.* 64, 233.—Cuba.

NORANTEA, Aubl. Pl. Gui. 554. t. 220 (1775). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 181.
ASCIUM, Schreb. Gen. 358 (1789).
SCHWARTZIA, Vell. Fl. Flum. v. t. 84 (1827).
adamantium, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 312. t. 62.—Bras.
anomala, *H. B. & K. Nov. Gen. et Sp.* vii. 218. t. 647 b.—Nicaragua.
aurantiaca, Spruce, ex Wittm. in Mart. Fl. Bras. xii. i. 240 = anomala.
bahiensis, Mart. ex Wittm. l. c. 245 = brasiliensis.
Berterii, *G. Don, Gen. Syst.* i. 624.—Ind. occ.
brasiliensis, *Choisy, in DC. Prod.* i. 566.—Bras.
cacabifera, *G. Don, Gen. Syst.* i. 625.—Peruv.
Caccabion, Delp. in Atti Soc. Ital. Sc. Nat. Milano, xii. (1869) 182 = cacabifera.
cordachida, *G. Don, Gen. Syst.* i. 625.—Peruv.
cuneifolia, *Delp. in Atti Soc. Ital. Sc. Nat. Milano*, xii. (1869) 182.—Bras.
Delpiniana, *Wittm. in Mart. Fl. Bras.* xii. i. 236.—Bras.
goyasensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 313.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 554. t. 220.—Am. trop.
indica, *Sweet, Hort. Brit.* ed. II. 79, nomen.—Maurit.
japurensis, *Mart. Nov. Gen. et Sp.* iii. 179. t. 295.—Bras.
Jussiaei, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 373.—Ind. occ.
macrocarpa, G. Don, Gen. Syst. i. 625 = Marcgravia rectiflora.
mixta, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 374.—N. Granat.
obovata, *G. Don, Gen. Syst.* i. 625.—Peruv.
oxystilis, *Baill. Adansonia*, x. (1872) 243.—N. Granat.
paraënsis, Mart. Nov. Gen. et Sp. iii. 180. t. 296 = guianensis.
pedunculatis, *Poepp. ex Wittm. in Mart. Fl. Bras.* xii. i. 238.—Peruv.

NORANTEA:—
rosea, Mart. ex Wittm. l. c. 243, in obs. = japurensis.
Selloi, G. Don, Gen. Syst. i. 624 = cuneifolia, brasiliensis.
sessiliflora, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 373 = anomala.
violacea, Poir. Encyc. Suppl. iv. 108 = guianensis.
Weddelliana, *Baill. Adansonia*, x. (1872) 242.—Bolivia.

NORATILEA, Walp. Ann. ii. 760 (1852), sphalm. in syn. = NONATELIA, Roxb. = **Lasianthus**, Jack (Rubiac.).

NORDMANNIA, Fisch. & Mey. ex C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 355 = **Daphnopsis**, Mart. & Zucc. (Thymel.).
tinifolia, Fisch. & Mey. l. c. 357 = D. tinifolia.

NORDMANNIA, Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 312 = **Trachystemon**, D. Don (Borag.).
cordifolia, Ledeb. ex Nordm. l. c. = T. orientale.

NORISCA, Dyer, in Hook. f. Fl. Brit. Ind. i. 253 (1874), sphalm. = NORYSCA, Spach = **Hypericum**, Tourn.

NORMANBYA, F. Muell. Fragm. xi. 57, in obs. (1878); Becc. in Ann. Jard. Buitenz. ii. (1885) 91 = **Ptychosperma**, Labill. (Palm.).

NORMANDIA, Hook. f. in Hook. Ic. Pl. t. 1121 (1872). *RUBIACEAE*, Benth. & Hook. f. ii. 139.
neo-caledonica, *Hook. f. in Hook. Ic. Pl.* t. 1121.—N. Caled.

NORMANIA, Lowe, Man. Fl. Mad. ii. 70 (1868) = **Solanum**, Tourn.
triphylla, Lowe, l. c. 87 = S. trisectum.

NORNA, Wahlenb. Fl. Suec. ed. II. 561 (1826) = **Calypso**, Salisb. (Orchid.).
borealis, Wahlenb. l. c. = C. borealis.

NORONHEA, Hook. Bot. Misc. ii. (1831) t. 88, sphalm. = seq.

NORONHIA, Stadm. in Thou. Gen. Nov. Madag. 8 (1806). *OLEACEAE*, Benth. & Hook. f. ii. 679.
Binia, Roem. & Schult. Syst. i. 72 = emarginata.
Broomeana, *Horne, ex Oliver, in Hook. Ic. Pl.* t. 1365.—Ins. Maurit.
cernua, G. Don, Gen. Syst. iv. 49 = Olea obtusifolia.
chartacea, Stadm. ex Hook. Bot. Misc. ii. (1831) 167 = emarginata.
emarginata, *Thou. Gen. Nov. Madag.* 8.—Madag.

NORRISIA, Gardn. in Hook. Kew Journ. i. (1849) 326. *LOGANIACEAE*, Benth. & Hook. f. ii. 796.
malaccensis, *Gardn. l. c.* 327.—Malacca; Borneo.

NORTA, Adans. Fam. ii. 417 (1763) = **Sisymbrium**, Tourn. (Crucif.).
juncea, Schur, Enum. Pl. Transs. 54 = S. strictissimum.
strictissima, Schur, l. c. = S. strictissimum.

NORTENIA, Thou. Gen. Nov. Madag. 9 (1806) = **Torenia**, Linn. (Scrophul.).
Thouarsii, Cham. & Schlecht. in Linnaea, iii. (1828) 18 = T. parviflora.

NORTHEA, Hook. f. in Hook. Ic. Pl. t. 1473 (1884). *SAPOTACEAE*.
seychellana, *Hook. f. l. c.*—Ins. Seychell.

NORYSCA, Spach, in Ann. Sc. Nat.. Sér. II. v. (1836) 363 = **Hypericum**, Tourn.
angustifolia, Blume, Mus. Bot. Lugd. Bat. ii. 22 = H. lanceolatum.
aurea, Blume, l. c. 23 = H. chinense.
calycina, Blume, l. c. 22 = H. calycinum.
cernua, Voigt, Hort. Suburb. Calc. 90 = H. cernuum.
chinensis, Spach, Hist. Vég. Phan. v. 427 = H. chinense.
cordifolia, Blume, Mus. Bot. Lugd. Bat. ii. 22 = H. cordifolium.

NORYSCA :—
coriacea, Blume, Mus. Bot. Lugd. Bat. ii. 24 = Hypericum coriaceum.
Hookeriana, Wight, Illustr. i. 113 = H. Hookeriana.
javanica, Blume, Mus. Bot. Lugd. Bat. ii. 24 = H. patulum.
lanceolata, Blume, l. c. 22 = H. lanceolatum.
Leschenaultii, Blume, l. c. 24 = H. Hookeriana.
madagascariensis, Blume, l. c. = H. madagascariensis.
myrtifolia, Spach, Hist. Vég. Phan. v. 428 = H. mysorense.
mysorensis, Wight, Ic. t. 56 = H. mysorense.
nervosa, Miq. Fl. Ind. Bat. i. II. 515 = H. nervosum.
oblongifolia, Blume, Mus. Bot. Lugd. Bat. ii. 22 = H. cernuum.
patula, Voigt, Hort. Suburb. Calc. 90 = H. patulum.
punctata, Blume, Mus. Bot. Lugd. Bat. ii. 23 = H. chinense.
rorida, Blume, l. c. 24.—Java.
salicifolia, Blume, l. c. 23 = H. chinense.
venosa, Blume, l. c. 22 = H. calycinum.

NOSTELIS, Rafin. Sylva Tellur. 76 (1838) = **Satureia** ?, Tourn. (Labiat.).
arborea, Rafin. l. c.—Hab. ?
minor, Rafin. l. c.—Hab. ?
viminea, Rafin. l. c. = Micromeria obovata.

NOTANTHERA, G. Don, Gen. Syst. iii. 428 (1834) = **Loranthus**, Linn.
acuminatus, G. Don, l. c. 429 = L. acuminatus.
anceps, G. Don, l. c. 430 = L. oblongo-ellipticus.
attenuatus, G. Don, l. c. = L. attenuatus.
buxifolius, G. Don, l. c. 429 = L. buxifolius.
caesius, G. Don, l. c. 428 = L. caesius.
conduplicatus, G. Don, l. c. 430 = L. conduplicatus.
ellipticus, G. Don, l. c. 429 = L. ellipticus.
grandiflorus, G. Don, l. c. 428 = L. grandiflorus, *Ruiz & Pav.*
heterophyllus, G. Don, l. c. 429 = L. heterophyllus.
lanceolatus, G. Don, l. c. 430 = L. lancifolius.
ligustrifolius, G. Don, l. c. 429 = L ligustrifolius.
longibracteatus, G. Don, l. c. 428 = L. longibracteatus.
orbicularis, G. Don, l. c. 429 = L. orbicularis.
ovatus, G. Don, l. c. 430 = L. ovatus.
oxyphyllus, G. Don, l. c. = L. oxyphyllus.
Poeppigii, G. Don, l. c. 428 = L. Poeppigii.
Pohlii, G. Don, l. c. = L. Pohlii.
radicans, G. Don, l. c. 429 = L. radicans.
salicifolius, G. Don, l. c. = L. salicifolius.
sarmentosus, G. Don, l. c. = L. sarmentosus.
Sellowianus, G. Don, l. c. = L. Sellowianus.
Sternbergianus, G. Don, l. c. 428 = L. Sternbergianus.
suaveolens, G. Don, l. c. 430 = L. suaveolens.
vegetus, G. Don, l. c. 429 = L. venetus.
verticillatus, G. Don, l. c. 428 = L. verticillatus, *Ruiz & Pav.*

NOTARISIA, Pestal. ex Cesati, in Bot. Zeit. xiv. (1856) 531 = **Ricotia**, Linn. (Crucif.).
securigera, Pestal. ex Cesati, l. c. = R. Pestalotiana.

NOTELAEA, Vent. Choix, 25. t. 25 (1803). *OLEACEAE*, Benth. & Hook. f. ii. 678.
HENSLOWIA, Lowe, ex DC. Prod. viii. 288 (1844).
PICCONIA, DC. l. c. (1844).
POSTUERA, Rafin. Sylva Tellur. 10 (1838).
RHYSOSPERMUM, Gaertn. f. Fruct. iii. 232. t. 224 (1805).
austro-caledonica, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 345.—N. Caled.
Badula, *Vieill. ex Panch. & Sebert, Not. Bois Nouv. Caléd.* 184.—N. Caled.
excelsa, *Webb & Bert. Phyt. Canar.* iii. 163.—Ins. Canar.
flavicans, *A. Dietr. Sp. Pl.* i. 246.—Hab. ?
lanceolata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 33.—Java.
laurifolia, *Kunth, Ind. Sem. Hort. Berol.* (1846) 12.—N. Zel.
ligustrina, Sieber, ex Benth. in DC. Prod. x. 191 = Duboisia myoporoides.
ligustrina, *Vent. Choix*, sub t. 25.—Austral.
linearis, *Benth. Fl. Austral.* iv. 300.—Austral.
longifolia, *Vent. Choix*, 25. t. 25.—Austral.
microcarpa, *R. Br. Prod.* 524.—Austral.
nervosa, *Steud. Nom.* ed. II. ii. 198.—Austral.

NOTELAEA :—
ovata, *R. Br. Prod.* 524.—Austral.
ovata, Endl. Iconog. t. 55 = longifolia.
Posua, D. Don, Prod. Fl. Nep. 107 (Linocierae sp. ?).—Nepal.
punctata, *R. Br. Prod.* 524.—Austral.
punctata, A. Cunn. ex DC. Prod. viii. 291 = longifolia.
reticulata, DC. l. c. = longifolia.
rigida, Desf. Hist. Arb. i. 120 = longifolia.
rostrata, *Teijsm. & Binn. Cat. Hort. Bogor.* 122.—Java.
venosa, F. Muell in Trans. Vict. Inst. (1855) 131 = longifolia.
zeylanica, Gardn. ex Thw. Enum. Pl. Zeyl. 188 = Olea polygama.
Zollingeriana, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 33 = Olea maritima.

NOTEROPHILA, Mart. Nov. Gen. et Sp. iii. 110. t. 254 (1829) = **Acisanthera**, P. Br. (Melastom.).
Beccabunga, Mart. in Flora, xxiv. (1841) II. Beibl. 69 = A. brevifolia.
brevifolia, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 280 = A. trivalvis.
inundata, Mart. Nov. Gen. et Sp. iii. 111. t. 254 = A. inundata.
inundata, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 279 = A. limnobios.
limnobios, Cham. in Linnaea, ix. (1834) 385 = A. fluitans.
limnobios, Mart. Nov. Gen. et Sp. iii. 112 = A. limnobios.
pusilla, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 281. t. 13. f. 1 = A. limnobios.
tetramera, Naud. l. c. 280 = A. tetramera.

NOTHAPHOEBE, Blume, Mus. Bot. Lugd. Bat. i. 328 (1851) = **Alseodaphne**, Nees (Laurin.).
canescens, Blume, l. c. 329.—Sumatra.
chartacea, Blume, l. c. 328.—Java.
cuneata, Blume, l. c.—Malaya.
falcata, Blume, l. c. 329.—Sumatra.
macrocarpa, Meissn. in DC. Prod. xv. I. 227 = *cuneata*.
macrophylla, Blume, Mus. Bot. Lugd. Bat. i. 329.—Malaya.
petiolaris, Meissn. in DC. Prod. xv. I. 59.—Assam.
spathulata, Meissn. l. c. 58.—Java.
superba, Blume, Mus. Bot. Lugd. Bat. i. 329.—Malaya.
umbelliflora, Blume, l. c. 328.—Malaya.

NOTHAPODYTES, Blume, Mus. Bot. Lugd. Bat. i. 248 (1850) = **Apodytes**, E. Mey. (Olacin.).
montana, Blume, l. c.—Java.

NOTHITES, Cass. in Dict. Sc. Nat. xxxv. 163 (1825) = **Stevia**, Cav. (Compos.).
angustifolia, Cass. l. c. 164 = S. satureiaefolia.
baccharidea, DC. Prod. v. 187 = Ophryosporus triangularis.
breviflora, Cass. in Dict. Sc. Nat. xxxv. 165.—Bras.
latifolia, Cass. l. c. 163 = *melissaefolia*.
melissaefolia, DC. Prod. v. 186.—Peruv.
ovatifolia, DC. l. c. 187 = Ophryosporus ovatifolius.
petiolata, Cass. in Dict. Sc. Nat. xxxv. 166.—Peruv.
satureiaefolia, DC. Prod. v. 186 = S. satureiaefolia.

NOTHOCESTRUM, A. Gray, in Proc. Am. Acad. vi. (1862) 48. *SOLANACEAE*, Benth. & Hook. f. ii. 894.
breviflorum, *A. Gray, l. c.* 49.—Ins. Sandvic.
latifolium, *A. Gray, l. c.* 48.—Ins. Sandvic.
longifolium, *A. Gray, l. c.*—Ins. Sandvic.
subcordatum, *H. Mann, l. c.* vii. (1867) 191.—Ins. Sandvic.

NOTHOCNESTIS, Miq. Fl. Ind. Bat. Suppl. 530 (1860) = **Kurrimia**, Wall. (Celastr.).
sumatrana, Miq. l. c. 531 = K. pulcherrima.

NOTHOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 137, in nota, t. 14 [1856] = **Pipturus**, Wedd. (Urtic.).
repanda, Blume, l. c. t. 14 = P. repandus.

NOTHODERRIS, Blume, ex Miq. Fl. Ind. Bat. i. I. 148 (1855) = **Derris**, Lour. (Legumin.).
elliptica, Blume, l. c. = D. elliptica.
glaucescens, Blume, l. c. 149 = D. elliptica.
purpurea, Blume, l. c. = D. elliptica.

NOTHOFAGUS, Blume, Mus. Bot. Lugd. Bat. i. 307 (1850) = **Fagus**, Tourn. (Cupulif.).
alpina, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 354 = F. alpina.
antarctica, Oerst. l. c. = F. antarctica.
betuloides, Blume, Mus. Bot. Lugd. Bat. i. 307 = F. betuloides.
cliffortioides, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 355 = F. cliffortioides.
Cunninghami, Oerst. l. c. = F. Cunninghamii.
Dombeyi, Blume, Mus. Bot. Lugd. Bat. i. 307 = F. Dombeyi.
dubia, Blume, l. c. = F. betuloides.
fusca, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 355 = F. fusca.
Gunnii, Oerst. l. c. 354 = F. Gunnii.
Menziesii, Oerst. l. c. 355 = F. Menziesii.
obliqua, Blume, Mus. Bot. Lugd. Bat. i. 307 = F. obliqua.
procera, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 354 = F. procera.
Solanderi, Oerst. l. c. 355 = F. Solandri.

NOTHOLIRION, Wall. ex Voigt, Hort. Suburb. Calc. 654 (1845) = **Lilium**, Tourn.
macrophyllum, Boiss. Fl. Orient. v. 190 = L. roseum.
roseum, Wall. ex Voigt, Hort. Suburb. Calc. 654 = L. roseum.

NOTHONIA, Endl. Ench. 245 (1841) = **Notonia**, DC. (Compos.).

NOTHOPANAX, Miq. Fl. Ind. Bat. i. I. 765 (1855) = **Panax**, Linn. (Araliac.).
anisum, Miq. l. c. 766 = P. anisum.
anomalum, Seem. Journ. Bot. iv. (1866) 296 = P. anomalum.
arboreum, Seem. l. c. = P. arboreum.
cephalobotrys, Seem. l. c. = P. cephalobotrys.
cochleatum, Miq. Fl. Ind. Bat. i. I. 766 = P. cochleatum.
Colensoi, Seem. in Journ. Bot. iv. (1866) 296 = P. Colensoi.
Cumingii, Seem. Fl. Vit. 114 = P. Cumingianum.
elegans, Seem. l. c. = P. elegans.
farinosum, Seem. Journ. Bot. iv. (1866) 295 = P. pinnatum.
fruticosum, Miq. Fl. Ind. Bat. i. I. 765 = P. fruticosum.
Gunnii, Seem. Journ. Bot. iv. (1866) 296 = P. Gunnii.
Macgillivrayi, Seem. Fl. Vit. 114 = P. Macgillivraei.
molle, Seem. Journ. Bot. iv. (1866) 295 = P. molle.
multijugum, Seem. Fl. Vit. 115, 114 = P. multijugum.
Murrayi, Seem. l. c. 114 = P. Murrayi.
obtusum, Miq. Fl. Ind. Bat. i. I. 766 = P. obtusum.
pinnatum, Miq. l. c. = Arthrophyllum pinnatum.
sambucifolium, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 77 = P. sambucifolium.
samoënse, Seem. Fl. Vit. 116 = P. samoënse.
Schimperi, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 78 = P. pinnatum.
simplex, Seem. Journ. Bot. iv. (1866) 296 = P. simplex.
Sinclairi, Seem. l. c. = P. Sinclairii.
tricochleatum, Miq. Fl. Ind. Bat. Suppl. 340 = Polyscias pinnata.
Zippelianum, Seem. Fl. Vit. 115 = P. Zippelianum.

NOTHOPEGIA, Blume, Mus. Bot. Lugd. Bat. i. 203 (1850). *ANACARDIACEAE*, Benth. & Hook. f. i. 425.
GLYCYCARPUS, Dalz. in Journ. As. Soc. Bomb. iii. (1849) 69.
aureo-fulva, *Bedd. ex Hook. f. Fl. Brit. Ind.* ii. 40.—Ind. or.
Colebrookiana, *Blume, Mus. Bot. Lugd. Bat.* i. 203.—Ind. or.
travancorica, *Bedd. ex Hook. f. Fl. Brit. Ind.* ii. 40.—Ind. or.

NOTHOPROTIUM, Miq. Fl. Ind. Bat. Suppl. 527 (1860). *BURSERACEAE*, Benth. & Hook. f. i. 326.
sumatranum, *Miq. l. c.*—Sumatra.

NOTHOSAERVA, Wight, Ic. vi. 1 (1853). *AMARANTACEAE*, Benth. & Hook. f. iii. 34.
PSEUDANTHUS, Wight, Ic. v. 3. t. 1776 bis B (1852).
brachiata, *Wight, Ic.* vi. 1.—Ind. or.; Burma; Afr. trop.

NOTHOSCORDUM, Kunth, Enum. Pl. iv. 457 (1848). *LILIACEAE*, Benth. & Hook. f. iii. 802.
CALOSCORDUM, Herb. in Bot. Reg. (1844) Misc. 66.
HESPEROCLES, Salisb. Gen. Pl. Fragm. 85 (1866).
OLIGOSMA, Salisb. l. c. (1866).
PSEUDOSCORDUM, Herb. Amaryll. 11 (1837), nomen prius.
andicolum, Kunth, Enum. Pl. iv. 463 = Milla andicola.
aureum, Hook. f. Bot. Mag. t. 5896 = Bloomeria aurea.
borbonicum, Kunth, Enum. Pl. iv. 462 = fragrans.
euosmum, Kunth, l. c. 460 = striatum.
flavescens, Kunth, l. c. 459 = striatum.
fragrans, *Kunth, l. c.* 461.—Am. bor.; Mexic.; Afr.; Maurit.
Gaudichaudianum, Kunth, l. c. 458 = Allium Sellovianum.
macrostemon, Kunth, l. c. 463 = Milla macrostemon.
neriniflorum, *Benth. & Hook. f. Gen.* iii. 802.—China.
ornithogaloides, Kunth, Enum. Pl. iv. 460 = striatum.
Philippianum, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1853) 17.—Chili.
Poiretii, *Kunth, Enum. Pl.* iv. 464.—Reg. Argent.
pulchellum, Kunth, l. c. 458 = striatum.
Sellowianum, Kunth, l. c. 460 = Allium Sellovianum.
siculum, Hort.—Cf. Gard. Chron. (1884) II. 174 = Nectaroscordum siculum.
striatellum, Kunth, Enum. Pl. iv. 458 = striatum.
striatum, *Kunth, l. c.* 459.—Am. bor.; Mexic.
strictum, C. Gay, Fl. Chil. vi. 114 = praec.
subbiflorum, Walp. Ann. iii. 636 = Allium subbiflorum.
Sulvia, Kunth, Enum. Pl. iv. 462 = fragrans.

NOTHOSMYRNIUM, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 58 (1867). *UMBELLIFERAE.*
japonicum, *Miq. l. c.*—Japon.

NOTHOSPARTIUM, Pritz. Ic. Ind. ii. 195 (1806) = **Notospartium**, Hook. f. (Legumin.).

NOTHOSPERMUM, Hort. ex Gard. Chron. (1883) II. 54 = **Notospartium**, Hook. f. (Legumin.).

NOTHRIA, Berg. Desc. Pl. Cap. 171 (1767) = **Frankenia**, Linn.
capensis, J. F. Gmel. Syst. 1012 = F. hirsuta.
hirsuta, Steud. Nom. ed. I. 556 = F. hirsuta.
repens, Berg. Desc. Pl. Cap. 171 = F. hirsuta.

NOTICASTRUM, DC. Prod. v. 279 (1836) = **Aster**, Tourn. (Compos.).
adscendens, DC. l. c. = A. marginatus.
album, Phil. in Anal. Univ. Chil. (1865) II. 317.—Chili.
antucense, Phil. in Linnaea, xxx. (1859–60) 192.—Chili.
erectum, Remy, in C. Gay, Fl. Chil. iv. 19.—Chili.
eriophorum, Remy, l. c. 20 = Leucopsis sericea.
Haplopappus, Remy, l. c. = Leucopsis sericea.
Philippi, Sch. Bip. in Bonplandia, iv. (1856) 54, nomen.—Peruv.
pradense, Phil. in Linnaea, xxxiii. (1864–65) 130 = A. pradensis.
sericeum, Less. ex Phil. l. c.—Chili.

NOTIOPHRYS, Lindl. in Journ. Linn. Soc. i. (1857) 189 = **Platylepis**, A. Rich. (Orchid.).
Commelynae, Lindl. l. c. = P. Commelynae.
glandulosa, Lindl. l. c. vi. (1862) 188 = P. glandulosa.
occulta, Lindl. l. c. i. (1857) 189 = P. goodyeroides.

NOTOBASIS, Cass. in Dict. Sc. Nat. xxv. 225 (1822); xxxv. 170 (1825) = **Cnicus**, Tourn. (Compos.).
syriaca, Cass. ll. cc. = C. syriacus.

NOTOBUXUS, Oliver, in Hook. Ic. Pl. xiv. 78. t. 1400 (1882). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 1222.
natalensis, *Oliver, l. c.*—Afr. austr.

NOTOCENTRUM, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 131 = **Meriania**, Sw. (Melastom.).
arboreum, Naud. l. c. = M. arborea.

NOTOCERAS, R. Br. in Ait. Hort. Kew. ed. II. iv. 117 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 71.
DICERATIUM, Lag. Elench. Madr. 20 (1815); Gen. et Sp. Nov. 20 (1816).

NOTOCERAS:—
bicorne, Amo, Fl. Iber. vi. 536 = canariense.
canariense, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 117.—Reg. Mediterr.; Oriens; Ind. bor. occ.
canescens, Walp. Rep. i. 124 = Diceratella canescens.
cardaminaefolium, DC. Syst. ii. 205 = Andrzeiowskia Cardamine.
floccosum, Walp. Rep. i. 124 = Diceratella floccosa.
hispanicum, DC. Syst. ii. 204 = canariense.
quadricorne, DC. l. c. = Tetracme quadricornis.
Sibthorpii, Spreng. Syst. ii. 894 = Mathiola bicornis.
sinuata, *Franch. in Revoil, Çomali*, 9.—Afr. trop.

NOTOCHAETE, Benth. in Wall. Pl. As. Rar. i. 63 (1830). *LABIATAE*, Benth. & Hook. f. ii. 1215.
hamosa, *Benth. in Wall. Pl. As. Rar.* i. 63.—Reg. Himal.

NOTOCLES, Salisb. Gen. Pl. Fragm. 54 (1866) = **Iphigenia**, Kunth (Liliac.).
indica, Salisb. l. c. = I. indica.

NOTONEMA, Rafin. Neogenyt. 4 (1825) = **Agrostis** Linn. (Gramin.).
arachnoides, Rafin. l. c. = A. arachnoides.

NOTONERIUM, Benth. in Benth. & Hook. f. Gen. ii. 698 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
Gossei, *Benth. in Hook. Ic. Pl.* t. 1180.—Austral.

NOTONIA, DC. in Guill. Arch. Bot. ii. (1833) 518. *COMPOSITAE*, Benth. & Hook. f. ii. 446.
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 444 (al. gen. ?).—Abyssin.
balsamica, *Dals. & Gibs. Bomb. Fl.* 133.—Ind. or.
coccinea, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 407.—Afr. trop.
corymbosa, DC. Prod. vi. 442 = grandiflora.
crassissima, *DC. l. c.*—Burma.
crassissima, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 337 = grandiflora.
grandiflora, *DC. in Guill. Arch. Bot.* ii. (1833) 518.—Ind. or.
Grantii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 407.—Afr. trop.
Hildebrandtii, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 197 = abyssinica.
kleinioides, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 151 = Senecio kleinioides.
Schweinfurthii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 407.—Afr. trop.
semperviva, Aschers. in Schweinf. Beitr. Fl. Aethiop. 152 = trachycarpa.
trachycarpa, *Klotzsch, in Sitzb. Acad. Wien, Math.-Nat.* li. *Abth.* II. (1865) 370.—Afr. trop.
Walkeri, *C. B. Clarke, Comp. Ind.* 176.—Ind. or.

NOTONIA, Wight & Arn. Prod. 207 (1834) = **Glycine**, Linn. (Legumin.).
Wightii, Wight & Arn. l. c. = G. javanica.

NOTOPHAENA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 267 = **Discaria**, Hook. (Rhamn.).
andina, Miers, l. c. 272.—Chili.
cognata, Miers, l. c. 270.—Chili.
discolor, Miers, l. c. = Colletia discolor.
foliosa, Miers, l. c. 269 = Colletia discolor.
magellanica, Miers, l. c. 271.—Reg. Magellan.
serratifolia, Miers, l. c. 268 = D. serratifolia.
tomentosa, Miers, l. c. 271 = Colletia tomentosa.
Toumatou, Miers, l. c. = D. Toumatou.

NOTOPHILUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325 = **Ranunculus**, Tourn.
muricatus, Fourr. l. c. = R. muricatus.
parviflorus, Fourr. l. c. = R. parviflorus.
vulgaris, Fourr. l. c. = R. sardous.

NOTOPORA, Hook. f. in Hook. Ic. Pl. t. 1159 (1873). *VACCINIACEAE*, Benth. & Hook. f. ii. 567.
Schomburgkii, *Hook. f. in Hook. Ic. Pl.* t. 1159.—Guian.

NOTOSCEPTRUM, Benth. in Benth. & Hook. f. Gen. iii. 775 (1883). *LILIACEAE*, Benth. & Hook. f. l. c.
alooides, *Benth. l. c.*—Afr. austr.
angolense, *Benth. l. c.*—Afr. trop.
benguellense, *Benth. l. c.*—Afr. trop.

NOTOSPARTIUM, Hook. f. in Hook. Kew Journ. ix. (1857) 176. t. 3. *LEGUMINOSAE*, Benth. & Hook. f. i. 502.
Carmicheliae, *Hook. f. in Hook. Kew Journ.* ix. (1857) 176. t. 3.—N. Zel.
Exsul, F. Muell. Fragm. vii. 126 = Carmichaelia Exsul.

NOTOTHIXOS, Oliver, in Journ. Linn. Soc. vii. (1864) 103. *LORANTHACEAE*, Benth. & Hook. f. iii. 214.
cornifolius, *Oliver, l. c.*—Austral.
floccosus, *Oliver, l. c.* 104.—Zeylan.
incanus, *Oliver, l. c.*—Austral.
subaureus, *Oliver, l. c.* 103.—Austral.

NOTOTHLASPI, Hook. f. in Benth. & Hook. f. Gen. i. 90 (1862). *CRUCIFERAE*, Benth. & Hook. f. l. c.
australe, *Hook. f. Handb. N. Zeal. Fl.* 15.—N. Zel.
notabile, *F. Buch. in Trans. N. Z. Inst.* xiv. (1882) 344.—N. Zel.
rosulatum, *Hook. f. Handb. N. Zeal. Fl.* 15.—N. Zel.

NOTOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 567 = **Malvastrum**, A. Gray (Malvac.).
cheilanthifolia, Turcz. l. c.—Bolivia.
discolor, Turcz. l. c.—Bolivia.
incana, Turcz. l. c. 568.—Bolivia.
incisa, Turcz. l. c.—Bolivia.

NOTYLIA, Lindl. Bot. Reg. sub t. 930 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 586.
TRIDACHNE, Liebm. ex Lindl. & Paxt. Flow. Gard. iii. (1852–53) 45.
albida, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 281.—Am. centr.
amazonica, *Linden, Illustr. Hortic.* xxix. (1882) 83, nomen.—Bras.
Arachnites, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 326.—Bolivia.
aromatica, *Barker, ex Lindl. Bot. Reg.* (1841) *Misc.* 40.—Guian.
Barkeri, *Lindl. l. c.* (1838) *Misc.* 90.—Mexic.
bicolor, *Lindl. in Benth. Pl. Hartw.* 93.—Guatemala.
bipartita, *Reichb. f. Xenia Orch.* i. 47.—Mexic.
cardioglossa, *Reichb. f. l. c.* 49.—Guian.
fragrans, *H. C. Focke, in Bot. Zeit.* xi. (1853) 342.—Guian.
hemitricha, *Rodrig. Orch. Nov.* ii. 223.—Bras.
Huegelii, *Fenzl, in Denkschr. Akad. Wien. Math.-Nat.* ii. (1850) II. 255.—Mexic.
incurva, *Lindl. Bot. Reg.* (1838) *Misc.* 90.—Ins. Trinit.
inversa, *Rodrig. Orch. Nov.* ii. 223.—Bras.
laxa, *Reichb. f. in Gard. Chron.* (1881) II. 620.—Bras.
laxiflora, *Westcott, in Phytologist*, i. (1841) 55.—Bras.
micrantha, *Lindl. Bot. Reg.* (1838) *Misc.* 90.—Guian.
multiflora, *Hook. Lond. Journ. Bot.* iii. (1844) 315. t. 10.—Mexic.
multiflora, *Lindl. Bot. Reg.* sub t. 930.—N. Granat.
nemorosa, *Rodrig. Orch. Nov.* ii. 224.—Bras.
odontonotos, *Reichb. f. & Warm. in Otia Bot. Hamb.* 87.—Bras.
orbicularis, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
pentachne, *Reichb. f. in Bonplandia*, ii. (1854) 90.—Venezuela.
pubescens, *Lindl. Bot. Reg.* (1842) *Misc.* 62.—Bras.
punctata, *Lindl. l. c.* sub t. 930.—Ins. Trinit.
replicata, *Linden, Illustr. Hortic.* xxix. (1882) 83, nomen.—Bras.
sagittifera, Hook. Lond. Journ. Bot. iii. (1844) 316 = multiflora, *Lindl.*
stenantha, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 299.—Bras.
tamaulipensis, *Reichb. f. l. c.* xvi. (1860) 281.—Mexic.
tenuis, *Lindl. Bot. Reg.* (1838) *Misc.* 90.—Guian.
Tridachne, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 45 = Huegelii.
trisepala, *Lindl. & Paxt. l. c.*—Costa Rica.
Wullschlaegeliana, *H. C. Focke, in Bot. Zeit.* xi. (1853) 343.—Guian.

NOVATILEA, Wight, in Calc. Journ. Nat. Hist. vi. (1846) 503, sphalm. = NONATELIA, Roxb. = **Lasianthus**, Jack (Rubiac.).

NOVELLA, Rafin. Sylva Tellur. 39 (1838) = **Cordia**, Linn. (Borag.).
nigra, Rafin. l. c. = C. subcordata.

NOWODWORSKYA, J. & C. Presl, Rel. Haenk. i. 351 (1830) = **Polypogon**, Desf. (Gramin.).
agrostoides, J. & C. Presl, l. c. = P. elongatus.
imberbis, Phil. in Anal. Univ. Chil. (1873) 562.—Chili.

NOYERA, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 135. t. 5. *URTICACEAE*, Benth. & Hook. f. iii. 373.
NAUCLEOPSIS, Miq. in Mart. Fl. Bras. iv. 1. 120. t. 35 (1853).
OGCODEIA, Bur. in DC. Prod. xvii. 282 (1873).
ONCODEIA, 'Bur.' emend. a Benth. & Hook. f. Gen. iii. 373 (1880).
rubra, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 136.—Guian.

NUBIGENA, Rafin. Sylva Tellur. 23 (1838) = **Cytisus**, Linn. (Legumin.).
tenerifa, Rafin. l. c. = C. fragrans.

NUDILUS, Rafin. Atl. Journ. 176 (1833) = **Forestiera**?, Poir. (Oleac.).
paradoxus, Rafin. l. c.—Texas.

NUFAR, Wallr. Beitr. Fl. Hercyn. 212 (1840) = **Nuphar**, Sibth. & Sm. (Nymph.).

NUMISAUREUM, Rafin. Fl. Tellur. iii. 32 (1836) = **Linum**, Tourn.
acuminatum, Rafin. l. c. = Reinwardtia tetragyna.
petiolatum, Rafin. l. c. = L. gallicum.
repens, Rafin. l. c., nomen = Reinwardtia trigyna?

NUMMULARIA, Riv. ex Rupp. Fl. Jen. ed. Hall. 18 (1745) = **Lysimachia**, Tourn. (Primul.).
centimorbia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 115 = L. Nummularia.
prostrata, Opiz, Seznam, 68 = L. Nummularia.
repens, Gilib. Fl. Lituan. i. 29 = L. Nummularia.
sylvatica, S. F. Gray, Nat. Arr. Brit. Pl. ii. 300 = L. nemorum.

NUMULARIA, Gilib. Fl. Lituan. i. 29 (1781) = praec.

NUNNEZHARIA, Ruiz & Pav. Prod. 147. t. 31 (1794) = **Chamaedorea**, Willd. (1805) (Palm.).
fragrans, Ruiz & Pav. Syst. Veg. 294 = C. fragrans.
geonomaeformis, Hook. f. Bot. Mag. t. 6088 = C. geonomaeformis.
geonomoides, Spruce, in Journ. Linn. Soc. xi. (1871) 122 = C. geonomoides.
tenella, Hook. f. Bot. Mag. t. 6584 = C. tenella.

NUNNEZIA, Willd. Sp. Pl. iv. 1154 (1805) = **Chamaedorea**, Willd. (Palm.).
fragrans, Willd. l. c. = C. fragrans.

NUPHAR, Sibth. & Sm. Fl. Graec. Prod. i. 361 [1808]. *NYMPHAEACEAE*, Benth. & Hook. f. i. 66.
NYMPHOZANTHUS, Rich. in Ann. Mus. Par. xvii. (1811) 230.
ROPALON, Rafin. New Fl. Am. ii. 17 (1836).
advena, *Ait. Hort. Kew.* ed. II. iii. 295.—Am. bor.
americanum, Provanch. Fl. Canad. 28 = advena.
intermedium, *Ledeb. Fl. Alt.* ii. 274.—Sibir. altaic.
japonicum, *DC. Syst.* ii. 62.—Japon.
Kalmianum, Ait. Hort. Kew. ed. II. iii. 295 = minimum.
latifolium, Steud. Nom. ed. II. ii. 199, sphalm. = seq.
longifolium, Sm. in Rees, Cycl. xxv. n. 5 = sagittaefolium.
luteum, *Sibth. & Sm. Fl. Graec. Prod.* i. 361.—Reg. bor. temp.
minimum, *Sm. Engl. Bot.* t. 2292.—Europ.; Am. bor.
minus, Dum. Fl. Belg. Prod. 131 = minimum?
polysepalum, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1866) 282.—Am. bor. occ.
pumilum, DC. Syst. ii. 61 = minimum.
rivulare, Dum. in Bull. Soc. Bot. Belg. iii. (1864) 5 = luteum.

NUPHAR:—
sagittaefolium, *Pursh, Fl. Am. Sept.* ii. 370.—Am. bor.
sericeum, Láng, in Syll. Ratisb. i. (1824) 180 = luteum.
spathuliferum, Reichb. Pl. Crit. ii. 10 = luteum.
Spennerianum, Gaud. Fl. Helv. iii. 439 = minimum.
systylum, Wallr. Beitr. Fl. Hercyn. 212 = luteum.
tenellum, Reichb. Pl. Crit. ii. 10 = luteum.
tomentosum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 58 = advena.
variegatum, Engelm. ex Clinton, 19th Rep. N. York St. Univ. 73; ex S. Wats. Bibl. Ind. N. Am. Bot. 37 = advena.

NUTTALLA, Rafin. in Am. Monthly Mag. (1818) 175 = **Bartonia**?, Muhl. (Gentian.).

NUTTALLIA, Barton, Fl. N. Am. ii. 74. t. 62 (1822) = **Callirhoë**, Nutt. (Malvac.).
ambigua, Croom, in Am. Journ. Sc. Ser. I. xxvi. (1834) 313 = C. Papaver.
cordata, Lindl. Bot. Reg. t. 1938 = C. digitata.
cordifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 98 = C. triangulata.
digitata, Barton, Fl. N. Am. ii. 74. t. 62 = C. digitata.
grandiflora, Paxt. Mag. Bot. v. (1838) 217 = C. Papaver.
involucrata, Nutt. ex Torr. in Ann. Lyc. N. York, ii. (1828) 172 = C. involucrata.
malvaeflora, Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iii. 41 = Sidalcea malvaeflora.
Munroana, Nutt. in Journ. Acad. Philad. vii. (1834) 16 = Sphaeralcea Munroana.
palmata, Torr. in Ann. Lyc. N. York, ii. (1828) 171, sphalm. pro *pedata*.
Papaver, R. Grah. in Bot. Mag. t. 3287 = C. Papaver.
pedata, Nutt. ex Hook. Exot. Fl. iii. t. 172 = C. digitata.
triangulata, Hook. Journ. Bot. i. (1834) 197 = C. triangulata.

NUTTALLIA, DC. Rapp. Jard. Genèv. (1821) 44 = **Nemopanthus**, Rafin. (Ilicin.).
canadensis, DC. l. c. = Nem. fascicularis.

NUTTALLIA, Spreng. Neue Entdeck. ii. 158 (1821) = **Trigonia**, Aubl. (Vochys.).
villosa, Spreng. l. c. = T. villosa.

NUTTALLIA, Torr. & Gray, ex Hook. Bot. Beech. Voy. 336. t. 82 (1840). *ROSACEAE*, Benth. & Hook. f. i. 611.
cerasiformis, *Torr. & Gray, l. c.* 337.—Am. bor. occ.
Davidiana, *Baill. Adansonia*, xi. (1875) 328.—Am. bor.

NUX, Tourn. ex Adans. Fam. ii. 497 (1763) = **Juglans**, Linn.

NUXIA, Comm. ex Lam. Illustr. i. 295. t. 71 (1791). *LOGANIACEAE*, Benth. & Hook. f. ii. 792.
LACHNOPYLIS, Hochst. in Flora, xxvi. (1843) 77.
capitata, *Baker, in Journ. Bot.* xx. (1882) 172.—Madag.
congesta, *R. Br. in Salt, Abyss. App.* 62, nomen; *et ex Fresen. in Flora*, xxi. (1838) 606.—Abyss.
corrugata, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 60 = Chilianthus corrugatus.
dentata, *R. Br. in Salt, Abyss. App.* 62, nomen; *et Benth. in DC. Prod.* x. 435.—Abyss.
dysophylla, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 60 = Chilianthus dysophyllus.
elata, Pers. Syn. i. 132 = Aegiphila elata.
emarginata, *Sond. in Linnaea*, xxiii. (1850) 83.—Afr. austr.
floribunda, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 59.—Afr. austr.
lobulata, Benth. l. c. 60 = Chilianthus lobulatus.
oppositifolia, *Benth. in DC. Prod.* x. 435.—Abyss.
pubescens, *Sond. in Linnaea*, xxiii. (1850) 84.—Afr. austr.
saligna, Benth. in Hook. Comp. Bot. Mag. ii. (1836) 59 = Chilianthus oleaceus.
tomentosa, *Sond. in Linnaea*, xxiii. (1850) 84.—Afr. austr.
verticillata, *Lam. Illustr.* i. 296.—Ins. Mascar.

NUYTSIA, R. Br. in Journ. Geogr. Soc. i. (1831) 17. *LORANTHACEAE*, Benth. & Hook. f. iii. 206.
floribunda, *R. Br. l. c.*—Austral.
ligustrina, A. Cunn. ex Lindl. Swan River App. 39 = Loranthus epigaeus.

NYALELIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Aglaia**, Lour. (Meliac.).
racemosa, Dennst. l. c. = A. edulis.

NYCTAGINIA, Choisy, in DC. Prod. xiii. II. 429 (1849). *NYCTAGINEAE*, Benth. & Hook. f. iii. 4.
capitata, *Choisy, l. c.*—Texas.
obtusa, *Choisy, l. c.*—Texas.
ovata, Choisy, in Mém. Soc. Phys. Genèv. xii. (1849) 167. t. 1 = capitata.
Torreyana, Choisy, in DC. Prod. xiii. II. 430 = Mirabilis multiflora.

NYCTAGO, Juss. Gen. 90 (1789) = **Mirabilis**, Linn. (Nyctag.).
angulata, DC. ex Choisy, in DC. Prod. xiii. II. 429 = Quamoclidion angulatum.
cordifolia, Moç. & Sesse, ex Choisy, l. c. 432 = Oxybaphus cordifolius.
dichotoma, Juss. ex Roem. & Schult. Syst. iv. 1 = M. dichotoma.
hortensis, Juss. ex Roem. & Schult. l. c. = M. Jalapa.
Jalapae, DC. Fl. Fr. iii. 426 = M. Jalapa.
longiflora, Salisb. Prod. 57 = M. longiflora.
parviflora, Salisb. l. c. = Oxybaphus viscosus.
versicolor, Salisb. l. c. = M. Jalapa.

NYCTANDRA Prior, in Proc. Linn. Soc. (1883–86) 5, sphalm. = **Nectandra**, Roland. (Laurin.).

NYCTANTHES, Linn. Gen. ed. I. 333 (1737). *OLEACEAE*, Benth. & Hook. f. ii. 675.
BRUSCHIA, Bertol. in Mem. Acad. Sc. Bolog. viii. (1857) 238.
PARIATICU, Adans. Fam. ii. 223 (1763).
PARILIUM, Gaertn. Fruct. i. 234. t. 51. f. 1 (1788).
acuminata, Burm. f. Fl. Ind. 5 = Tabernaemontana coronaria.
angustifolia, Linn. Sp. Pl. 6 = Jasminum angustifolium.
Arbor-tristis, *Linn. l. c.*—Ind. or.
dentata, *Blume, Mus. Bot. Lugd. Bat.* i. 282.—Sumatra.
elongata, Linn. f. Suppl. 82 = Jasminum elongatum.
glauca, Linn. f. l. c. = Jasminum glaucum.
Goa, Hort. ex Steud. Nom. ed. II. ii. 199 = Jasminum Sambac.
grandiflora, Lour. Fl. Cochinch. 21 = Jasminum arborescens.
hirsuta, Linn. Sp. Pl. 6 = Jasminum pubescens, Guettarda speciosa.
multiflora, Burm. f. Fl. Ind. 5. t. 3. f. 1 = Jasminum pubescens.
pubescens, Retz. Obs. v. 9 = Jasminum pubescens.
Sambac, Linn. Sp. Pl. 6 = Jasminum Sambac.
scandens, Retz. Obs. v. 9 = Jasminum scandens.
triflora, Burm. f. Fl. Ind. 4. t. 4 = Jasminum angustifolium.
tristis, Salisb. Prod. 11 = N. Arbor-tristis.
undulata, Linn. Sp. Pl. 6 = Jasminum Sambac.
viminea, Retz. Obs. v. 9 = Jasminum angustifolium.

NYCTANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

NYCTELEA, Scop. Introd. 183 (1777) = **Ellisia**, Linn. (Hydroph.).

NYCTERANTHUS, Neck. Elem. ii. 82 (1790) = **Mesembryanthemum**, Dill. (Ficoid.).

NYCTERINIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 239 (1834) = **Zaluzianskya**, F. W. Schmidt (Scrophul.).
africana, D. Don, l. c. sub t. 239 = Z. villosa.
capensis, Benth. in Hook. Comp. Bot. Mag. i. (1835) 370 = Z. capensis.
coriacea, Benth. l. c. 369 = Z. coriacea.
dentata, Benth. l. c. 370 = Z. dentata.
divaricata, Benth. l. c. 371 = Z. divaricata.
fragrans, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 239 = Z. lychnidea.

NYCTERINIA:—
longiflora, Benth. in Hook. Comp. Bot. Mag. i. (1835) 370 = Zaluzianskya longiflora.
lychnidea, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 239 = Z. lychnidea.
maritima, Benth. in Hook. Comp. Bot. Mag. i. (1835) 369 = Z. maritima.
natalensis, Bernh. in Flora, xxvii. (1844) 834 = Z. natalensis.
ovata, Benth. in Hook. Comp. Bot. Mag. i. (1835) 370 = Z. ovata.
peduncularis, Benth. l. c. 371 = Z. peduncularis.
pumila, Benth. l. c. 370 = Z. pumila.
pusilla, Benth. l. c. 371 = Z. pusilla.
selaginoides, Benth. l. c. 370 = Z. selaginoides.
spathacea, Benth. l. c. 369 = Z. spathacea.
tristis, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 239 = Erinus tristis.
villosa, Benth. in Hook. Comp. Bot. Mag. i. (1835) 371 = Z. Benthamiana.

NYCTERISITION, Ruiz & Pav. Prod. 30. t. 5 (1794) = **Chrysophyllum**, Linn. (Sapot.).
argenteum, H. B. & K. Nov. Gen. et Sp. iii. 238. t. 244 = C. granatense.
ferrugineum, Ruiz & Pav. Fl. Per. ii. 47. t. 187 = C. ferrugineum.
lanceolatum, Blume, Bijdr. 676 = C. javanicum.

NYCTERIUM, Vent. Jard. Malm. t. 85 (1803) = **Solanum**, Tourn.
amazonium, Sims, Bot. Mag. t. 1801 = S. amazonium.
cardaminifolium, Vent. Jard. Malm. sub t. 85 = S. cornutum.
cordifolium, Vent. l. c. t. 85 = S. Vespertilio.
cornutum, Link, Enum. Hort. Berol. i. 189 = S. cornutum.
flavum, Lindl. in Donn, Hort. Cantabr. ed. XI. 76 = S. heterandrum.
Fontanesianum, Sweet, Hort. Brit. ed. I. 301 = S. Fontanesianum.
heterandrum, Heynh. Nom. ii. 440 = S. heterandrum.
heterodoxum, Link, Enum. Hort. Berol. i. 189 = S. heterodoxum.
Lamberti, Lindl. in Donn, Hort. Cantabr. ed. XI. 76 = S. amazonium.
lobatum, Sweet, Hort. Brit. ed. I. 301 = S. heterandrum.
luteum, Torr. ex Steud. Nom. ed. II. ii. 199 = S. heterandrum.
pubescens, G. Don, in Sweet, Hort. Brit. ed. III. 501 = S. pubescens.
rostratum, Link, Enum. Hort. Berol. i. 189 = S. rostratum.
triphyllum, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimpr. 14 = S. trisectum.
Vespertilio, Link, Enum. Hort. Berol. i. 189 = S. Vespertilio.
violaceum, Engelm. ex Walp. Ann. iii. 167 = S. heterodoxum.

NYCTOCALOS, Teijsm. & Binn. in Journ. Bot. Néerl. i. (1861) 366. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1040.
brunfelsiaeflorus, *Teijsm. & Binn. in Journ. Bot. Néerl.* i. (1861) 367.—Java.
cuspidatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 249. —Ins. Molucc.; Celebes.
jucundum, Seem. in Journ. Bot. viii. (1870) 149 = Haussmannia jucunda.
macrosiphon, *Teijsm. & Binn. Cat. Hort. Bog.* 155, nomen.—Ins. Celebes.
Thomsoni, *Hook. f. Bot. Mag.* t. 5678.—Assam.

NYCTOPHYLA, Zipp. in Alg. Konst. en Letter-bode i. (1829) 298. 'Nomen delendum.'—Cf. Benth. & Hook. f. Gen. iii. 639.
alba, Zipp. l. c.—Ins. Timor.

NYCTOSMA, Rafin. Fl. Tellur. ii. 9 (1836) = **Epidendrum**, Linn. (Orchid.).
nocturna, Rafin. l. c. = E. nocturnum.

NYLANDTIA, Dum. Comm. Bot. 31 (1822) = **Mundtia**, Kunth (Polygal.).

NYMPHAEA, [Tourn.] Linn. Syst. ed. I (1735). *NYMPHAEACEAE*, Benth. & Hook. f. i. 46. CASTALIA, Salisb. Parad. Lond. i. t. 14 (1805).
abbreviata, Guill. & Perr. Fl. Seneg. Tent. i. 16 = stellata.
acutiloba, *DC. Prod.* i. 116.—China.
advena, [Soland. in] Ait. Hort. Kew. ed. I. ii. 226 = Nuphar advena.
aegyptiaca, Opiz, Natural. x. (1825) 216 = N. Lotus.
alba, *Linn. Sp. Pl.* 510.—Reg. bor. temp.
alba, Nutt. Gen. Am. ii. 13 = tuberosa.
alba, Patrin, ex Bunge, in Ledeb. Fl. Alt. ii. 272 = pauciradiata.
alba, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 93 = Basniniana.
alba, Walt. Fl. Carol. 155 = odorata.
albo-rosea, *Teijsm. ex Regel, Ind. Sem. Hort. Petrop.* (1866) 101.—Hab. ?
alboviridis, A. St. Hil. Voy. Distr. Diam. ii. 426 = amazonum.
amazonum, Hort. ex Casp. in Mart. Fl. Bras. iv. II. 161 = Rudgeana.
amazonum, *Mart. & Zucc. in Abh. Akad. Muench.* i. (1832) 363.—Bras.
ampla, *DC. Syst.* ii. 54.—Am. bor. ; Ind. occ.
ampla, Kotschy, ex Casp. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 244 = stellata.
arifolia, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 71 = Nuphar advena.
Bashiniana, Steud. Nom. ed. II. ii. 200 = seq.
Basniniana, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 93.—Sibir. baical.
bella, *Lehm. in Hamb. Gartenz.* ix. (1853) 212 (Quid ?). —Ind. or.
Bernieriana, Planch. in Rev. Hortic. Sér. IV. ii. (1853) 65 = stellata.
biradiata, Somm. in Flora, xvi. (1833) 625 = alba.
blanda, Hook. Bot. Mag. sub t. 4823 = amazonum.
blanda, *G. F. W. Mey. Prim. Fl. Esseq.* 201.—Am. austr.
blanda, Planch. in Ann. Sc. Nat. Sér. III. xix. (1853) 47 = Rudgeana.
Boucheana ×, Planch. in Fl. des Serres, x. (1854–55) t. 1033–1034.
cachemiriana, Cambess. in Jacquem. Voy. Bot. ii. t. 10 = alba.
caerulea, Andr. Bot. Rep. t. 197 = capensis.
caerulea, Savign. in Ann. Mus. Par. i. (1802) 366 = stellata.
Cahlara, Donn, Hort. Cantabr. ed. VII. 159 = stellata.
candida, J. & C. Presl, Delic. Prag. 224 = alba.
Candolliana, Lehm. in Hamb. Gartenz. ix. (1853) 203 = ampla.
capensis, *Thunb. Prod. Pl. Cap.* 92.—Afr. austr.
coerulea, Auct. vide *caerulea*.
Coteka, Roxb. ex Salisb. Parad. Lond. i. sub t. 14 = N. Lotus.
crenulata, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid ?).—Am. bor.
cyanea, Roxb. Hort. Beng. 41; Fl. Ind. ii. 577 = stellata.
dentata, Schum. & Thonn. Beskr. Guin. Pl. 249 = N. Lotus.
Devoniensis, Hook. Bot. Mag. t. 4665 = N. Lotus.
discolor, Lehm. in Hamb. Gartenz. ix. (1853) 214 = stellata.
Edgeworthii, Lehm. l. c. viii. (1852) 372 = stellata.
edulis, DC. Syst. ii. 52 = N. Lotus.
elegans, *Hook. Bot. Mag.* t. 4604.—Texas.
emirnensis, Planch. in Rev. Hortic. Sér. IV. ii. (1853) 65 = stellata.
erythrocarpa, Hentze, in Bot. Zeit. x. (1852) 747 = alba.
esculenta, Roxb. Hort. Beng. 41; Fl. Ind. ii. 578 = N. Lotus.
exumbonata, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xii. (1854) 219 = alba.
Fenzliana, Lehm. in Hamb. Gartenz. ix. (1853) 217 = blanda.
flava, *Leitner, ex Audubon, Birds of Am.* ed. fol. iv. t. 411.—Florida.
flavo-virens, *Lehm. Ind.Sem. Hort. Hamb.* (1852) 9; *et in Ann. Sc. Nat.* Sér. III. xix. (1853) 361.—Hab. ?
foetida, Gardn. ex Planch. in Fl. des Serres, Sér. II. i. (1856) 21 = amazonum.
fragrans, Gardn. ex Casp. in Mart. Fl. Bras. iv. II. 176 = Gardneriana.

NYMPHAEA:—
fragrantissima, *Welw. ex Casp. in Forn. Sc. Math. Lisboa*, n. xvi. (1873) reimpr. 11.—Afr. trop.
Gardneriana, *Planch. in Rev. Hortic.* Sér. IV. ii. (1853) 66; *et in Ann. Sc. Nat.* Sér. III. xix. (1853) 50.—Bras.
gigantea, *Hook. Bot. Mag.* t. 4647.—Austral.
glandulifera, Rodsch. ex G. F. W. Mey. Prim. Fl. Esseq. 201 = blanda.
Goudotiana, Planch. in Rev. Hortic. Sér. IV. ii. (1853) 66 = amazonum.
gracilis, *Zucc. in Abh. Akad. Muench.* i. (1832) 362.—Mexic.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 248 = stellata.
hastata, Michx. ex DC. Syst. ii. 62 = Nuphar sagittaefolium.
Heudelotii, Planch. in Rev. Hortic. Sér. IV. ii. (1853) 65 = stellata.
hirta, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Indie*, xxvii. (1864) 38.—Sumatra.
Hookeriana, Lehm. in Hamb. Gartenz. ix. (1853) 213 = stellata.
integrifolia, Salzm. ex Planch. in Fl. des Serres, xi. (1856) 21 = amazonum.
intermedia, Weiker, ex Reichb. Fl. Saxon. 10, in obs. = alba.
Jamesoniana, *Planch. in Rev. Hortic.* Sér. IV. ii. (1853) 66; *et in Ann. Sc. Nat.* Sér. III. xix. (1853) 51.—Ecuador.
Kalmiana, Sims, Bot. Mag. t. 1243 = Nuphar minimum.
Kosteletzkyi, Paillard, ex Lehm. Ind. Sem. Hort. Hamb. (1852) 10; et in Ann. Sc. Nat. Sér. III. xix. (1853) 362 = alba.
lasiophylla, *Mart. & Zucc. in Abh. Akad. Muench.* i. (1832) 364.—Bras.
Leiboldiana, Lehm. in Hamb. Gartenz. ix. (1853) 209 = ampla, Rudgeana.
lineata, A. St. Hil. Voy. Distr. Diam. ii. 425 = ampla.
longifolia, Michx. Fl. Bor. Am. i. 312 = Nuphar sagittaefolium.
Lotus, Aubl. Pl. Gui. i. 533 = ampla.
Lotus, *Linn. Sp. Pl.* 511.—As. et Afr. trop.
lutea, Linn. l. c. 510 = Nuphar luteum.
lutea, Thunb. Fl. Jap. 223 = Nuphar japonicum.
lutea, Walt. Fl. Carol. 154 = Nuphar advena.
maculata, Rafin. Med. Fl. ii. 45 = tuberosa ?
maculata, Schum. & Thonn. Beskr. Guin. Pl. 247 = stellata.
madagascariensis, DC. Syst. ii. 50 = stellata.
malabarica, Poir. in Lam. Encyc. iv. 457 = stellata.
Maximiliani, *Lehm. in Hamb. Gartenz.* ix. (1853) 215. —Bras.
membranacea, Wall. ex Casp. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 245 = stellata.
mexicana, A. Gray, Pl. Wright. i. 7 = elegans.
mexicana, *Zucc. in Abh. Akad. Muench.* i. (1832) 365. —Mexic.
micrantha, Guill. & Perr. Fl. Seneg. Tent. i. 16 = stellata.
microphylla, Pers. Syn. ii. 63 = Nuphar minimum.
Milletii, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 37 = alba.
minima, Reichb. Fl. Germ. Excurs. 14 = Nuphar minimum.
minor, DC. Syst. ii. 58 = odorata.
Naccheli, J. F. Gmel. Syst. 1591, sphalm. = *N. Nouchali*.
neglecta, Hausleutner, in Bot. Zeit. viii. (1850) 906 = alba.
Nelumbo, Linn. Sp. Pl. 511 = Nelumbium speciosum.
Nelumbo, Schoepf. Mat. Med. 88; Lunan, Hort. Jamaic. ii. 272 = Nelumbium luteum.
Nelumbo, Vell. Fl. Flum. 224; v. t. 95 = Victoria regia ?
nervosa, Lehm. in Hamb. Gartenz. ix. (1853) 211 = ampla.
nitida, *Sims, Bot. Mag.* t. 1359.—Sibir.
nocturna, March, ex Hook. Bot. Mag. sub t. 4823 = amazonum.
Nouchali, Burm. f. Fl. Ind. 120 = stellata ?
nubica, *Lehm. in Hamb. Gartenz.* ix. (1853) 211.—Afr. trop.
odorata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 227.—Am. bor.
odorata, Willd. Hort. Berol. t. 39 = nitida.
officinalis, Gater. Pl. Montaub. 99 = alba.

NYMPHAEA:—
Ortgiesiana, Planch. in Fl. des Serres, viii. (1852–53) 68 = N. Lotus.
oxypetala, *Planch. in Rev. Hortic.* Sér. IV. ii. (1853) 66; *et in Ann. Sc. Nat.* Sér. III. xix. (1853) 51.—Ecuador.
Parkeriana, *Lehm. Ind. Sem. Hort. Hamb.* (1853) 10; *et in Hamb. Gartenz.* ix. (1853) 529.—Guian.
parviflora, Hentze, in Bot. Zeit. vi. (1848) 700 = alba.
Passiflora, Lehm. in Hamb. Gartenz. ix. (1853) 217 = Gardneriana.
pauciradiata, *Bunge, in Ledeb. Fl. Alt.* ii. 272.—Sibir.
pentapetala, Walt. Fl. Carol. 155 = Nelumbium luteum.
permixta, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 37 = alba.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. 152 = stellata.
poecila, Lehm. Ind. Sem. Hort. Hamb. (1852) 11.—Cf. Hamb. Gartenz. viii. (1852) 371, 425 = stellata.
Pseudo-pygmaea, Lehm. in Hamb. Gartenz. ix. (1853) 208 = stellata.
pubescens, Willd. Sp. Pl. ii. 1154 = N. Lotus.
pulchella, DC. Syst. ii. 51 = ampla.
pulchella, Lehm. ex Casp. in Mart. Fl. Bras. iv. II. 177 = tenuinervia.
pumila, Hoffm. Deutschl. Fl. ed. II. i. I. 241 = Nuphar minimum.
punctata, Edgew. in Trans. Linn. Soc. xx. (1846) 29 = stellata.
punctata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 376.—As. centr.
pygmaea, [Dryand. in] Ait. Hort. Kew. ed. II. iii. 293 = tetragona.
radiata, Bercht. & Opiz, Natural. x. (1825) 216 = stellata.
Raja, Lehm. in Hamb. Gartenz. ix. (1853) 208 = oxypetala.
Regina, Schomb. ex Lindl. Monog. 3 = Victoria regia.
reniformis, DC. Syst. ii. 55 = tuberosa?
reniformis, Walt. Fl. Carol. 155 = odorata.
rhodantha, Lehm. in Hamb. Gartenz. ix. (1853) 214 = stellata.
rosea, Rafin. Med. Fl. ii. 45 = odorata?
rosea, *Sweet, Hort. Brit.* ed. I. 15.—Ind. or.
rotundifolia, Hentze, in Bot. Zeit. vi. (1848) 700 = alba.
rubra, Roxb. ex Salisb. Parad. Lond. i. sub t. 14 = N. Lotus.
Rudgeana, *G. F. W. Mey. Prim. Fl. Esseq.* 198.—Ind. occ.; Guiana.
rufescens, Guill. & Perr. Fl. Seneg. Tent. i. 15 = stellata.
rufescens, *Willd. ex Casp. in Jorn. Sc. Math. Lisboa*, n. 16 (1873) reimpr. 8, nomen.—Hab.?
sagittariaefolia, Lehm. in Hamb. Gartenz. ix. (1853) 216 = Jamesoniana.
sagittata, Edgew. in Trans. Linn. Soc. xx. (1846) 29 = N. Lotus.
sagittata, Pers. Syn. ii. 63 = Nuphar sagittaefolium.
sagittifolia, Walt. Fl. Carol. 155 = Nuphar sagittaefolium.
scutifolia, DC. Syst. ii. 50 = capensis.
semiaperta, Klinggraeff, Fl. Preuss. 20 = alba.
semisterilis, Lehm. in Hamb. Gartenz. ix. (1853) 215 = stellata.
sinuata, Salzm. ex Planch. in Ann. Sc. Nat. Sér. III. xix. (1853) 45 = ampla.
speciosa, Mart. & Zucc. in Abh. Akad. Muench. i. (1832) 361 = ampla.
spiralis, Rafin. Med. Fl. ii. 45 = tuberosa?
splendens, Hentze, in Bot. Zeit. vi. (1848) 603 = alba.
stellata, F. Muell. Fragm. ii. 142 = gigantea.
stellata, *Willd. Sp. Pl.* ii. 1153.—As. et Afr. trop.
stenaspidota, *Casp. in Mart. Fl. Bras.* iv. II. 175.—Bras.
suaveolens, Dum. in Bull. Soc. Bot. Belg. iii. (1864) 7 = alba.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 387.—Sumatra.
tenuinervia, *Casp. in Mart. Fl. Bras.* iv. II. 177.—Bras.
tetragona, *Georgi, Reise Russ. Reich.* i. 220.—Reg. Himal.; Sibir.; China.
thermalis, DC. Syst. ii. 54 = N. Lotus.
trisepala, *Gaudich. in Mém. Sav. Etr. Pétersb.* viii. (1841) 83.—Am. trop.

NYMPHAEA:—
tropaeolifolia, Lehm. in Hamb. Gartenz. ix. (1853) 209 = ampla, Rudgeana.
tuberosa, *Paine, Cat. Pl. Oneida*, 132.—Am. bor.
tussilagifolia, *Lehm. Ind. Sem. Hort. Hamb.* (1853) 10; *et in Hamb. Gartenz.* ix. (1853) 530.—Mexic.
umbilicalis, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 71 = Nuphar luteum.
undulata, *Lehm. in Hamb. Gartenz.* ix. (1853) 210.—Mexic.
urceolata, Hentze, in Bot. Zeit. vi. (1848) 699 = alba.
venusta, Hentze, l. c. = alba.
versicolor, Sims, Bot. Mag. t. 1189 = stellata.
Victoria, Schomb. ex Lindl. Monog. 3 = Victoria regia.
violacea, Lehm. in Hamb. Gartenz. ix. (1853) 218 = gigantea.
vivipara, *Lehm. Novit. Pl. Hort. Bot. Hamb.* (1852) 11.—*Cf. Hamb. Gartenz.* viii. (1852) 370.—Hab.?
Wenzelii, *Maack, ex Regel, in Mém. Acad. Pétersb.* Sér. VII. iv. (1861) n. IV. 17.—Reg. Amur.
zanzibariensis, *Casp. in Bot. Zeit.* xxxv. (1877) 203.—Afr. trop.

NYMPHAEOLA, Heist. Syst. 7 (1748) = **Hydrocharis**? Linn.

NYMPHANTHUS, Lour. Fl. Cochinch. 543 (1790) = **Phyllanthus**, Linn. (Euphorb.).
chinensis, Lour. l. c. 544 = P. villosus.
Niruri, Lour. l. c. 545 = P. Niruri.
pilosa, Lour. l. c. 544 = P. pilosus.
rubra, Lour. l. c. = P. ruber.
squamifolia, Lour. l. c. = P. squamifolius.

NYMPHEA, Rafin. in Med. Repos. N. York, v. (1808) 354 = **Nymphaea**, Tourn.

NYMPHOIDES, Tourn. ex Medic. Phil. Bot. i. 35 (1789) = **Limnanthemum**, S. P. Gmel. (Gentian.).
europaea, Fisch. ex Steud. Nom. ed. I. 557 = L. nymphoides.
orbiculata, Gilib. Fl. Lituan. i. 33 = L. nymphoides.

NYMPHOSANTHUS, Steud. Bot. ed. II. ii. 200 (1841) = seq.

NYMPHOZANTHUS, Rich. in Ann. Mus. Par. xvii. (1811) 230 = **Nuphar**, Sibth. & Sm. (Nymph.).
vulgaris, Rich. l. c. = Nuphar luteum.

NYPA, Wurmb. in Verh. Batav. Gen. i. (1779) 349 = **Nipa**, Thunb. (Palm.).
fruticans, Wurmb. l. c. = Nipa fruticans.

NYPHA, Buch.-Ham. in Mem. Wern. Soc. v. II. (1826) 318 = praec.

NYPHAR, Walp. Rep. i. 108 (1842), sphalm. = **Nuphar**, Sibth. & Sm. (Nymph.).

NYROPHYLLA, Neck. Elem. ii. 240 (1790). *LAURIN*, Gen. dub.

NYSSA, Gronov. ex Linn. Syst. ed. I (1735). *CORNACEAE*, Benth. & Hook. f. i. 952.
AGATHISANTHES, Blume, Bijdr. 645 (1825).
CERATOSTACHYS, Blume, l. c. 644 (1825).
DAPHNIPHYLLOPSIS, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 201.
TUPELO, Adans. Fam. ii. 80 (1763).
angulisans, Michx. Fl. Bor. Am. ii. 259 = uniflora.
angulosa, Poir. in Lam. Encyc. iv. 507 = uniflora.
aquatica, M. A. Curt. Bot. N. Carol. 62 = caroliniana.
aquatica, Linn. Sp. Pl. 1058 = multiflora, uniflora.
biflora, Michx. Fl. Bor. Am. ii. 259 = multiflora.
biflora, Walt. Fl. Carol. 253 = caroliniana?
canadensis, Poir. in Lam. Encyc. iv. 507 = multiflora?
candicans, Michx. Fl. Bor. Am. ii. 259 = capitata.
capitata, *Walt. Fl. Carol.* 253.—Am. bor.
caroliniana, *Poir. in Lam. Encyc.* iv. 507.—Am. bor.
ciliata, Rafin. Atl. Journ. 176 (Quid?).—Am. bor.
coccinea, Bartr. Trav. ed. II. (1794) 17 = capitata.
denticulata, Ait. Hort. Kew. ed. I. iii. 446 = uniflora.
glauca, Hort. Par. ex Roem. & Schult. Syst. v. 575 = multiflora.
grandidentata, Michx. f. Hist. Arb. For. Am. Sept. ii. 252 = uniflora.
integrifolia, Ait. Hort. Kew. ed. I. iii. 446 = multiflora.
montana, Gaertn. f. Fruct. iii. 201. t. 216 = capitata.

NYSSA:—
montana, Hort. ex Pursh, Fl. Am. Sept. i. 177 = multiflora.
multiflora, Ell. Sketch, ii. 684 = caroliniana?
multiflora, *Wangenh. Beitr. Forstw. Nordam. Holz.* 46. t. 16.—Am. bor.
Ogeche, Marsh. Arbust. 97 = capitata.
Oye-chee, Steud. Nom. ed. I. 558 = praec.
palustris, Salisb. Prod. 175 = uniflora.
sessiliflora, *Hook. f. & Thoms. in Benth & Hook. f. Gen.* i. 952.—Reg. Himal.
sylvatica, Marsh. Arbust. 97 = multiflora.
tomentosa, Michx. Fl. Bor. Am. ii. 259 = uniflora.
tomentosa, Poir. in Lam. Encyc. iv. 508 = capitata.
uniflora, *Wangenh. Beitr. Forstw. Nordam. Holz.* 83. t. 27.—Am. bor.
villosa, Bigel. Fl. Bost. ed. II. 380 = multiflora.
villosa, Michx. Fl. Bor. Am. ii. 258 = multiflora.

NYSSANTHES, R. Br. Prod. 418 (1810). *AMARANTACEAE*, Benth. & Hook. f. iii. 36.
diffusa, *R. Br. Prod.* 418.—Austral.
erecta, *R. Br. l. c.*—Austral.
media, R. Br. l. c. = diffusa.

O

OAKESIA, Tuckerm. in Hook. Lond. Journ. Bot. i. (1842) 445 = **Corema**, D. Don (Empetr.).
Conradii, Tuckerm. l. c. 446 = C. Conradii.

OAKESIA, S. Wats. in Proc. Am. Acad. xiv. (1879) 269. *LILIACEAE*.
puberula, *S. Wats. l. c.*—Am. bor.
sessilifolia, *S. Wats. l. c.*—Am. bor.

OBAEJACA, Cass. in Dict. Sc. Nat. xxxv. 270 (1825) = **Senecio**, Tourn. (Compos.).
sylvatica, Cass. l. c. 271 = S. sylvaticus.
viscosa, Cass. l. c. 270 = S. viscosus.

OBBEA, Hook. f. in Hook. Ic. Pl. t. 1070 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 102.
timonioides, *Hook. f. l. c.*—Ins. Sandvic.

OBECKIA, Griff. Notul. iv. 675 (1854), sphalm. = **Osbeckia**, Linn. (Melastom.).

OBELANTHERA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 148 = **Saurauja**, Willd. (Ternstr.).
melastomacea, Turcz. l. c. 149 = S. villosa.

OBELISCARIA, Cass. in Dict. Sc. Nat. xxxv. 272 (1825) = **Lepachys**, Rafin. (Compos.).
columnaris, DC. Prod. v. 559 = L. columnaris.
Drummondii, Maund, Botanist, v. t. 201 = Rudbeckia Drummondii.
peduncularis, Walp. Rep. ii. 979 = L. peduncularis.
pinnata, Cass. in Dict. Sc. Nat. xxxv. 273 = L. pinnatifida.
pulcherrima, DC. Prod. v. 559 = L. columnaris.
pulcherrima, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 744 = Rudbeckia Drummondii.
Tagetes, DC. Prod. v. 559 = L. Tagetes.

OBELISCOTHECA, Vaill. ex Adans. Fam. ii. 128 (1763) = **Rudbeckia**, Linn. (Compos.).

OBENTONIA, Vell. Fl. Flum. 18 (1825); i. t. 46 (1827) = **Cusparia**, Humb. (Rutac.).
castrata, Vell. ll. cc. = C. macrophylla.

OBERNA, Adans. Fam. ii. 255 (1763) = **Silene**, Linn. (Caryoph.).

OBERONIA, Lindl. Gen. et Sp. Orch. 15 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 494.
TITANIA, Endl. Prod. Fl. Norf. 31 (1838).
acaulis, Griff. Notul. iii. 275 = myriantha.
acaulis, Hook. Bot. Mag. t. 5056 = ensiformis.
anceps, *Lindl. Sert. Orchid.* sub t. 8 B.—Burma; Malaya.

OBERONIA:—
angustifolia, *Lindl. Fol. Orch. Ober.* 5.—Reg. Himal.
anthropophora, *Lindl. in Wall. Cat.* n. 1951.—Burma.
aporophylla, *Reichb. f. in Bonplandia*, iii. (1855) 223.—Ins. Philipp.
Arnottiana, Wight, Ic. t. 1628 = Wightiana.
bicornis, *Lindl. in Wall. Cat.* n. 1949.—Reg. Himal.
brachystachys, *Lindl. Sert. Orch.* sub t. 8 B.—Burma.
brevifolia, *Lindl. Gen. et Sp. Orch.* 16.—Ins. Societ.; Mascar.
Brunoniana, *Wight, Ic.* t. 1622.—Ind. or.
caulescens, *Lindl. Gen. et Sp. Orch.* 15.—Reg. Himal.
cimicina, Griff. Notul. iii. 275 = Griffithiana.
crassiuscula, F. Muell. ex Benth. Fl. Austral. vi. 290 = Phreatia limenophylax.
cylindrica, *Lindl. Bot. Reg.* (1840) *Misc.* 20.—Ins. Philipp.
demissa, *Lindl. Fol. Orch. Ober.* 4.—Reg. Himal.
denticulata, Wight, Ic. t. 1625 = iridifolia.
ensiformis, *Lindl. Fol. Orch. Ober.* 4.—Reg. Himal.; Burma.
forcipata, *Lindl. l. c.* 2.—Zeylan.
Gardneriana, *Thw. Enum. Pl. Zeyl.* 296.—Zeylan.
gladiata, A. Rich. Sert. Astrol. 6. t. 2 = Eria gladiata.
glandulifera, *Ridl. in Forbes, Nat. Wand. East Archip.* 518.—Ins. Timor.
glandulosa, *Lindl. Fol. Orch. Ober.* 6.—Ins. Pacif.
Griffithiana, *Lindl. Sert. Orch.* t. 8 B.—Burma.
Griffithii, Wight, Ic. t. 1629 = anceps.
imbricata, *Lindl. Gen. et Sp. Orch.* 17.—Java.
imbricata, Wight, Ic. t. 1629 = anceps.
iridifolia, *Lindl. in Wall. Cat.* n. 1948; *et in Gen. et Sp. Orch.* 15.—Ind. or.
Jenkinsiana, *Griff. ex Lindl. Fol. Orch. Ober.* 4.—Reg. Himal.
Lindlei, Brongn. in Duperr. Voy. Coq. Bot. t. 40. f. a = seq.
Lindleyana, A. Brongn. l. c. p. 200 = Phreatia Lindleyi.
Lindleyana, *Wight, Ic.* t. 1624.—Ind. or.
Lindleyi, Benth. & Hook. f. Gen. iii. 511, lapsu = *Lindleyana*, Brongn.
Lobbiana, *Lindl. Fol. Orch. Ober.* 7.—Borneo.
longibracteata, *Lindl. Gen. et Sp. Orch.* 15.—Zeylan.
lunata, *Lindl. l. c.* 17.—Java.
micrantha, A. Rich. Sert. Astrol. 7. t. 3 = Eria Richardiana.
microphylla, *Lindl. Gen. et Sp. Orch.* 17.—Java.
miniata, *Lindl. Bot. Reg.* (1843) *Misc.* 6.—Singapore; Ins. Philipp.
monstruosa, *Lindl. Gen. et Sp. Orch.* 17.—Java.
Myosurus, *Lindl. in Wall. Cat.* n. 1947.—Reg. Himal.; Burma.
myriantha, *Lindl. Fol. Orch. Ober.* 4.—Reg. Himal.
obcordata, *Lindl. l. c.* 7.—Reg. Himal.
pachystachya, *Lindl. l. c.* 6.—Ins. Pacif.
palmicola, F. Muell. Fragm. ii. 24 = O. Titania.
platycaulon, *Wight, Ic.* t. 1623.—Ind. or.
Pumilio, *Reichb. f. in Bonplandia*, v. (1857) 58.—Java.
pyrulifera, *Lindl. Fol. Orch. Ober.* 3.—Reg. Himal.
recurva, *Lindl. Bot. Reg.* (1839) *Misc.* 14.—Ind. or.
rufilabris, *Lindl. Sert. Orch.* t. 8 A.—Burma.
Scyllae, *Lindl. Fol. Orch. Ober.* 5.—Zeylan.
setifera, Lindl. l. c. 3 = recurva.
sikkimensis, Lindl. l. c. 4 = myriantha.
similis, *Lindl. Gen. et Sp. Orch.* 17.—Java.
spathulata, *Lindl. l. c.* 16.—Java.
spiralis, *Griff. Notul.* iii. 275.—Hab. ?
spiralis, *Lindl. Fol. Orch. Ober.* 7.—Java.
stachyoides, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 15 = Wightiana.
tahitensis, *Lindl. Fol. Orch. Ober.* 2.—Ins. Tahiti.
tenuis, *Lindl. l. c.* 3.—Zeylan.
Thisbe, *Reichb. f. in Bonplandia*, iii. (1855) 223.—Ins. Philipp.
Titania, *Lindl. Fol. Orch. Ober.* 8.—Java; Austral.
trilobata, Griff. Notul. iii. 273 = ensiformis.
truncata, *Lindl. Fol. Orch. Ober.* 3.—Zeylan.
verticillata, *Wight, Ic.* t. 1626.—Ind. or.
Wightiana, *Lindl. Bot. Reg.* (1839) *Misc.* 14.—Ind. or.; Zeylan.

OBESIA, Haw. Syn. Pl. Succ. 42 (1812) = **Piaranthus**, R. Br. (Asclep.).
decora, Haw. l. c. 43 = P. decorus.
geminata, Haw. l. c. 42 = P. geminatus.
punctata, Haw. l. c. 43 = P. punctatus.
serrulata, Sweet, Hort. Brit. ed. I. 278 = P. serrulatus.

OBETIA, Gaudich. Bot. Voy. Bonite, t. 82 ('1844'). *URTICACEAE*, Benth. & Hook. f. iii. 382.
ficifolia, *Gaudich. l. c.*—Madag.
ficifolia, Wedd. ex Baker, in Journ. Linn. Soc. xx. (1883) 263 = morifolia.
laciniata, *Baker, l. c.* 264.—Madag.
madagascariensis, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 178.—Madag.
morifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 263.—Madag.
pinnatifida, *Baker, l. c.* 264.—Madag.
Radula, *Baker, l. c.* 265.—Madag.

OBIONE, Gaertn. Fruct. ii. 198. t. 126 (1791) = **Atriplex**, Linn. (Chenop.).
acanthocarpa, Torr. Bot. Mex. Bound. 183 = A. acanthocarpa, Greggii.
arenaria, Moq. Chenop. Enum. 71 = A. arenaria.
argentea, Moq. l. c. 76 = A. argentea.
Barclayana, Benth. Bot. Voy. Sulph. 48 = A. Barclayana.
Belangerii, Moq. Chenop. Enum. 71 = A. repens.
Berlandieri, Moq. in DC. Prod. xiii. II. 114 = A. canescens.
Billardieri, Moq. Chenop. Enum. 72 = A. Billardieri.
bracteosa, Dur. & Hilg. in Journ. Acad. Philad. Ser. II. iii. (1855) 44 = A. bracteosa.
canescens, Moq. Chenop. Enum. 74 = A. canescens.
confertifolia, Torr. & Frém. Rep. 318 = A. confertifolia.
coriacea, Moq. Chenop. Enum. 71 = A. coriacea.
Coulteri, Moq. in DC. Prod. xiii. II. 113 = A. Coulteri.
crispa, Moq. Chenop. Enum. 73 = A. Halimus.
cristata, Moq. l. c. 73 = A. cristata.
elegans, Moq. in DC. Prod. xiii. II. 113 = A. elegans.
fera, Moq. l. c. 107 = A. lenticularis.
Gardneri, Moq. l. c. 114 = A. Gordoni.
glauca, Moq. l. c. 108 = A. glauca.
graeca, Moq. l. c. = A. graeca.
hymenelytra, Torr. in Pacif. Rail. Rep. iv. 129. t. 20 = A. hymenelytra.
imbricata, Moq. in DC. Prod. xiii. II. 108 = A. imbricata.
Konigii, Moq. l. c. 109 = A. repens.
Kunthiana, Moq. Chenop. Enum. 72 = A. muricata.
Lampa, Moq. in DC. Prod. xiii. II. 110 = A. Lampa.
lenticularis, Moq. Chenop. Enum. 70 = A. lenticularis.
lentiformis, Torr. in Sitgreaves, Rep. 169 = A. lentiformis.
leucophylla, Moq. in DC. Prod. xiii. II. 109 = A. leucophylla.
linifolia, Moq. Chenop. Enum. 74 = A. linifolia.
microcarpa, Benth. Bot. Voy. Sulph. 48 = A. microcarpa, *D. Dietr.*
montevidensis, Moq. in DC. Prod. xiii. II. 113 = A. montevidensis.
muricata, Gaertn. Fruct. ii. 198. t. 126. f. 5 = A. sibirica.
muricata, Moq. in DC. Prod. xiii. II. 109 = A. muricata.
nummularia, Moq. Chenop. Enum. 72 = A. repens.
occidentalis, Moq. in DC. Prod. xiii. II. 112 = A. canescens.
parvifolia, Moq. Chenop. Enum. 73 = A. parvifolia.
patagonica, Moq. in DC. Prod. xiii. II. 111 = A. patagonica.
pedunculata, Moq. Chenop. Enum. 75 = A. pedunculata.
phyllostegia, Torr. ex S. Wats. in Bot. King's Exp. 291 = A. phyllostegia.
polycarpa, Torr. in Emory, Notes Mil. Reconnois. 149, nomen = A. polycarpa.
polygama, Moq. in DC. Prod. xiii. II. 114 = A. polygama.
portulacoides, Moq. Chenop. Enum. 75 = A. portulacoides.
pusilla, Torr. ex S. Wats. in Bot. King's Exp. 291 = A. pusilla.
radiata, Torr. Bot. Mex. Bound. 183 = A. elegans.
rotundifolia, Moq. Chenop. Enum. 70 = A. rotundifolia.
salicifolia, Moq. l. c. 74 = A. polygama.
sibirica, Fisch. Cat. Hort. Gorenk. (1808) 25 = A. sibirica.
spinosa, Moq. in DC. Prod. xiii. II. 108 = A. spinosa.
Stocksii, Wight, Ic. t. 1789 = A. Stocksii.
stylosa, Moq. Chenop. Enum. 77 = A. stylosa.
Suckleyana, Torr. ex A. Gray, in Pacif. Rail. Rep. xii. II. (1860) 47. t. 4 = A. Suckleyana.

OBIONE:—
tetraptera, Benth. Bot. Voy. Sulph. 48 = Atriplex canescens.
thunbergiaefolia, Boiss. & Noé, Diagn. Ser. II. iv. 74 = A. thunbergiaefolia.
Torreyi, S. Wats. in Bot. King's Exp. 290 = A. Torreyi.
truncata, Torr. ex S. Wats. l. c. 291 = A. truncata.
turcomanica, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 451 = A. turcomanica.
undulata, Moq. in DC. Prod. xiii. II. 110 = A. undulata.
Verbasci, Auct. ex Trautv. in Act. Hort. Petrop. ix. (1884) 127 = seq. ?
verrucifera, Moq. Chenop. Enum. 76 = A. verrucifera.

OBLETIA, Rozier, in Journ. Phys. i. (1773) 367 = **Verbena**, Linn.

OBLIXILIS, Rafin. Fl. Tellur. iii. 49 (1836) = **Laportea**, Gaudich. (Urtic.).
canadensis, Rafin. l. c. = L. canadensis.
divaricatus, Rafin. l. c. = L. canadensis.

OBOEJACA, Steud. Nom. ed. II. ii. 201 (1841) = OBAEJACA, Cass. = **Senecio**, Tourn. (Compos.).

OBOLARIA, Linn. Sp. Pl. ed. I. 632 (1753). *GENTIANEAE*, Benth. & Hook. f. ii. 818.
OBULARIA, Linn. Hort. Cliff. 323 (1737).
SCHULTZIA, Rafin. in Med. Repos. N. York, v. (1808) 350.
caroliniana, Walt. Fl. Carol. 166 = Herpestis amplexicaulis.
virginica, *Linn. Sp. Pl.* 632.—Am. bor.

OBOSKON, Rafin. Fl. Tellur. iii. 93 (1836) = **Salvia**, Tourn. (Labiat.).
cretica, Rafin. l. c. = S. cretica.

OBSITILA, Rafin. Fl. Tellur. ii. 27 (1836) = **Anthericum**, Tourn. (Liliac.).

OBULARIA, Linn. Hort. Cliff. 323 (1737) = **Obolaria**, Linn. (Gentian.).

OCALIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 195 = **Croton**, Linn. (Euphorb.).
angustifolia, Klotzsch, l. c. = C. antisiphiliticus.
betulina, Klotzsch, l. c. = C. repens.
cordiaefolia, Klotzsch, l. c. = C. antisiphiliticus.
echiifolia, Klotzsch, l. c. = C. antisiphiliticus.
grandifolia, Klotzsch, l. c. = C. antisiphiliticus.
lantanaefolia, Didr. in Kjoeb. Vidensk. Meddel. (1857) 139 = C. urticaefolius.
Sellowiana, Klotzsch, in Wiegm. Archiv, vii. (1841) 195, nomen = C. antisiphiliticus.

OCAMPOA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 31 = **Cranichis**, Sw. (Orchid.).
mexicana, A. Rich. & Gal. l. c.—Mexic.

OCHAGAVIA, Phil. in Bot. Zeit. xiv. (1856) 647. *BROMELIACEAE*, Benth. & Hook. f. iii. 661.
elegans, *Phil. l. c.*—Ins. J. Fernand.

OCHANOSTACHYS, Mast. in Hook. f. Fl. Brit. Ind. i. 576 (1875). *OLACINEAE*.
amentacea, *Mast. l. c.* 577.—Malacca; Borneo.

OCHETOCARPUS, Meyen, Reise, i. 310 (1834) = **Grammatocarpus**, Presl (Loaseae).
striata, Meyen, l. c. = G. volubilis.

OCHETOPHILA, Poepp. ex Reiss. in Endl. Gen. 1099 (1840) = **Discaria**, Hook. (Rhamn.).
divergens, Poepp. l. c. = Sageretia divergens.
Hookeriana, Reiss. ex C. Gay, Fl. Chil. ii. 39 = Colletia Doniana.
parvifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 379.—Chili.
prostrata, Miers, l. c. = Colletia nana.
riparia, Poepp. ex Endl. Gen. 1099 = Sageretia riparia.
trinervis, Poepp. ex Endl. l. c. = Colletia Doniana.

OCHLANDRA, Thw. Enum. Pl. Zeyl. 376 (1864). *GRAMINEAE*, Benth. & Hook. f. iii. 1215.
BEESHA, Munro, in Trans. Linn. Soc. xxvi. (1868) 144.
stridula, *Thw. Enum. Pl. Zeyl.* 376.—Zeylan.

OCHNA, Linn. Gen. ed. I. 364 (1737); Schreb. Gen. 354 (1789). *OCHNACEAE*, Benth. & Hook. f. i. 317.
DIPORIDIUM, Bartl. & Wendl. f. Beitr. ii. 24 (1825).
Afzelii, *R. Br. ex Oliver, Fl. Trop. Afr.* i. 319.—Afr. trop.
andamanica, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 295 = Wallichii.
arborea, *Burch. ex DC. Prod.* i. 736.—Afr. austr.
ardisioides, *Webb, Fragm. Fl. Aethiop.* 59.—Afr. trop.
atropurpurea, *DC. in Ann. Mus. Par.* xvii. (1811) 412.—Afr. austr.
brevipes, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 652.—Burma.
capensis, Hort. Berol. ex Kunth, in Linnaea, xix. (1847) 394 = Elaeodendron verrucosum.
capensis, Page, Prod. Southhamp. Bot. Gard. 149 = arborea.
cardiosperma, Lam. Encyc. iv. 511 (*cordiosperma.*) = Gomphia cardiosperma.
chrysantha, Mundt, ex Steud. Nom. ed. II. ii. 201 = arborea.
ciliata, *Lam. Encyc.* iv. 511.—Madag.
collina, Edgew. in Trans. Linn. Soc. xx. (1846) 43 = pumila.
cordata, Thw. Enum. Pl. Zeyl. 70, 409 = squarrosa.
cordiosperma, vide *cardiosperma*.
crocea, Griff. Notul. iv. 463 = Gomphia sumatrana.
delagoënsis, *Walp. Rep.* i. 528.—Afr. trop.
dubia, Guill. & Perr. Fl. Seneg. Tent. i. 137 = multiflora.
fasciculares, Blanco, Fl. Filip. ed. II. 245 = Brackenridgea fascicularis.
fruticulosa, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 295 = pumila.
guianensis, Lam. Encyc. iv. 511 = Gomphia guyannensis.
Heyneana, Wight & Arn. Prod. 152 = Wightiana.
humilis, Wall. Cat. n. 3762 = pumila.
integrifolia, *Presl, in Oken, Isis*, xxi. (1828) 273.—Ins. Maurit.
itineraria, Vell. Fl. Flum. 224; v. t. 94 = Gomphia olivaeformis.
Jabotapita, Linn. Sp. Pl. 513 = Gomphia Jabotapita.
Jabotapitta, Vell. Fl. Flum. 223; v. t. 90 = Gomphia parviflora.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 317.—Afr. trop.
laevigata, Vahl, Symb. Bot. ii. 49 = Gomphia obtusifolia.
lanceolata, *Spreng. Syst.* ii. 597.—Ind. or.
leptoclada, *Oliver, Fl. Trop. Afr.* i. 318.—Afr. trop.
leucophloeos, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 129.—Abyss.
longifolia, Lam. Encyc. iv. 511 = Gomphia longifolia.
lucida, Lam. Encyc. iv. 510 = squarrosa.
macrocalyx, *Oliver, Fl. Trop. Afr.* i. 319.—Afr. trop.
madagascariensis, *DC. in Ann. Mus. Par.* xvii. (1811) 413.—Madag.
mauritiana, *Lam. Encyc.* iv. 512.—Ins. Maurit.
Mechowiana, *O. Hoffm. in Linnaea*, xliii. (1880–82) 123.—Afr. trop.
membranacea, *Oliver, Fl. Trop. Afr.* i. 316.—Afr. trop.
Moonii, Thw. Enum. Pl. Zeyl. 70 = Wightiana.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* i. 88. t. 16.—Afr. trop.
multiflora, *DC. in Ann. Mus. Par.* xvii. (1811) 412.—Afr. trop.
nana, Buch.-Ham. in Wall. Cat. n. 3761 = pumila.
natalitia, Walp. Rep. ii. 826 = atropurpurea.
nitida, Sw. Prod. Veg. Ind. Occ. 67 = Gomphia guyannensis.
nitida, Thunb. Prod. Pl. Cap. 67 = squarrosa.
nitida, Wall. Cat. n. 2804 = Wallichii.
obtusata, DC. in Ann. Mus. Par. xvii. (1811) 411 = squarrosa.
obtusifolia, Lam. Encyc. iv. 510 = Gomphia obtusifolia.
parviflora, Griff. Notul. iv. 464 = Wallichii.
parvifolia, Vahl, Symb. Bot. i. 33 = Wightiana.
polycarpa, *Baker, in Journ. Bot.* xx. (1882) 49.—Madag.
pulchra, *Hook. Ic. Pl.* t. 588.—Afr. austr.
pumila, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 224.—Ind. or.; Burma.
rufescens, Thw. Enum. Pl. Zeyl. 70 = squarrosa.
serratifolia, *Baker, in Journ. Linn. Soc.* xxi. (1884) 329.—Madag.

OCHNA:—
serrulata, Walp. Rep. v. 400 = atropurpurea.
squarrosa, Kurz, Andam. Rep. App. 74 = Wallichii.
squarrosa, *Linn. Sp. Pl.* ed. II. 732.—Ind. or.; Burma.
stipulacea, Colebr. in Wall. Cat. n. 2806 = Wallichii.
stipulata, Vell. Fl. Flum. 223; v. t. 91 = Gomphia stipulacea.
terminalis, *Vell. l. c.* 223; v. t. 92.—Bras.
vaccinioides, *Baker, in Journ. Linn. Soc.* xxi. (1884) 329.—Madag.
verticillata, Vell. Fl. Flum. 222; v. t. 89 = Gomphia verticillata.
vulgaris, *Vell. l. c.* 223; v. t. 93.—Bras.
Walkerii, Planch. in Hook. Lond. Journ. Bot. v. (1846) 653 = Wightiana.
Wallichii, *Planch. l. c.* 650.—Burma.
Wightiana, *Wall. Cat.* n. 2808.—Ind. or.
zeylanica, Lam. Encyc. iv. 512 = Gomphia angustifolia.

OCHRADENUS, Delile, Fl. Egypte, 236. t. 31. f. 1 (1813). *RESEDACEAE*, Benth. & Hook. f. i. 112.
HOMALODISCUS, Bunge, ex Boiss. Fl. Orient. i. 422 (1867).
Aucheri, *Boiss. Diagn.* Ser. II. i. 50.—Arab.
baccatus, *Delile, Fl. Egypte*, 236. t. 31.—Aegypt.; Oriens; Scind.
Boissieri, *Muell. Arg. in DC. Prod.* xvi. II. 588.—Persia.
minor, *Muell. Arg. l. c.*—Persia.
rostratus, Ehrenb. ex Muell. Arg. in Neue Denkschr. Schw. Ges. Zuerich, xvi. (1858) (Monog. Resed.) 95 = baccatus.

OCHRANTHE, Lindl. Bot. Reg. t. 1819 (1836) = **Turpinia**, Vent. (Sapind.).
arguta, Lindl. l. c. = T. arguta.

OCHROCARPOS, Thou. Gen. Nov. Madag. 15 (1806). *GUTTIFERAE*, Benth. & Hook. f. i. 175.
CALYSACCION, Wight, Illustr. i. 130 (1840).
africana, *Oliver, Fl. Trop. Afr.* i. 169.—Afr. trop.
Chapelieri, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 366.—Madag.
decipiens, *Baill. Adansonia*, xi. (1876) 370.—Madag.
evonymoides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 366.—Madag.
Goudotianus, *Planch. & Triana, l. c.* 365.—Madag.
Harmandii, *Pierre, Fl. For. Cochinch.* t. 93.—Cochinch.
longifolius, *Benth. & Hook. f. ex T. Anders. in Hook. f. Fl. Brit. Ind.* i. 270.—Ind. or.
macrophyllus, *O. Hoffm. Sert. Madag.* 7.—Madag.
madagascariensis, *DC. Prod.* i. 560.—Madag.
multiflorus, *O. Hoffm. Sert. Madag.* 7.—Madag.
nervosus, *Kurz, For. Fl. Brit. Burma*, i. 94.—Burma.
ovalifolius, *T. Anders. ex Hemsl. Bot. Voy. Challenger*, i. III. 122.—Ins. Timor Laut.
pentapetalus, *Blanco, ex Vill. in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 17.—Ins. Philipp.
siamensis, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 270.—Burma.

OCHROLASIA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 3 = **Hibbertia**, Andr. (Dillen.).
Drummondii, Turcz, l. c. 4 = H. Ochrolasia.

OCHROMA, Sw. Prod. Veg. Ind. Occ. 97 (1788). *MALVACEAE*, Benth. & Hook. f. i. 212.
lagopodum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = seq.
Lagopus, *Sw. Prod. Veg. Ind. Occ.* 98.—Am. trop.
tomentosa, *Willd. Enum. Hort. Berol.* 695.—Am. austr.

OCHRONELIS, Rafin. Atl. Journ. 153 (1832) = **Verbesina**, Linn. (Compos.).
sulfurea, Rafin. l. c. = V. nudicaulis.

OCHROSIA, Juss. Gen. 144 (1789). *APOCYNACEAE*, Benth. & Hook. f. ii. 700.
BLECKERIA, Hassk. Retzia, i. 38 (1855).
DIDEROTA, Comm. ex A. DC. in DC. Prod. viii. 356 (1844).
LACTARIA, Rumpf, ex Rafin. Sylva Tellur. 162 (1838); Miq. Fl. Ind. Bat. ii. 414 (1856).
NEIOSPERMA, Rafin. Sylva Tellur. 162 (1838).

OCHROSIA:—
Ackeringae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 138.—Ins. Banca.
borbonica, *J. F. Gmel. Syst.* 439.—Ins. Mascar.
calocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 139.—Austral.?
coccinea, *Miq. l. c.* 138.—Ins. Ceram.
elliptica, *Labill. Sert. Austro-Caled.* 25. t. 30.—N. Caled.
Kilneri, *F. Muell. Fragm.* vii. 129.—Austral.
maculata, Jacq. Coll. iv. 218 = borbonica.
mariannensis, *A. DC. in DC. Prod.* viii. 357.—Ins. Marian.
Moorei, *F. Muell. ex Benth. Fl. Austral.* iv. 310.—Austral.
Mulsanti, *Montr. in Mém. Acad. Lyon*, x. (1860) 235.—N. Caled.
parviflora, G. Don, Gen. Syst. iv. 99 = elliptica.
parviflora, Henslow, in Ann. Nat. Hist. Ser. I. i. (1838) 345 = elliptica.
platyspermos, A. DC. in DC. Prod. viii. 356 = borbonica.
salubris, Blume, Mus. Bot. Lugd. Bat. i. 158 = borbonica.
sandwicensis, *A. DC. in DC. Prod.* viii. 357.—Ins. Sandvic.
Thiollierei, *Montr. in Mém. Acad. Lyon*, x. (1860) 235.—N. Caled.
undulata, Boj. Hort. Maurit. 206 = borbonica.

OCHROSION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.

OCHROXYLUM, Schreb. Gen. 826 (1791) = **Zantho xylum**, Linn. (Rutac.).
punctatum, Mart. in Denkschr. Akad. Muench. (1816) 151 = Z. Ochroxylum.

OCHRUS, Tourn. ex Adans. Fam. ii. 330 (1763) = **Lathyrus**, Linn. (Legumin.).
hirsutus, Pritz. Ic. Ind. i. 751, sphalm. = *Orobus hirsutus*.
lathyroides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 357 = L. Ochrus.
pallida, Pers. Syn. ii. 305 = L. Ochrus.
uniflora, Moench, Meth. 163 = L. Ochrus.

OCHTHOCHARIS, Blume, in Flora, xiv. (1831) 523. *MELASTOMACEAE*, Benth. & Hook. f. i. 752.
bornensis, *Blume, Mus. Bot. Lugd. Bat.* i. 40.—Borneo.
buruensis, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 426 = bornensis.
javanica, *Blume, in Flora*, xiv. (1831) 523.—Burma; Malaya.
javanica, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 40 = bornensis.
paniculata, *Korth. in Verh. Nat. Gesch. Bot.* 247. t. 64.—Malaya.

OCHTHOCOSMUS, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 366. *LINEAE*, Benth. & Hook. f. i. 245.
africanus, Hook. f. in Hook. Ic. Pl. t. 773 = Phyllocosmus africanus.
Roraimae, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 366.—Guian.
sessiliflorus, Baill. Adansonia, x. (1873) 367 = Phyllocosmus sessiliflorus.

OCHTHODIUM, DC. Syst. ii. 423 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 97.
aegyptiacum, *DC. l. c.*—Cilicia; Syria; Aegypt.

OCHTOCHARIS, Walp. Ann. ii. 600 (1851–52) = **Ochthocharis**, Blume (Melastom.).

OCIMUM, Linn. Hort. Cliff. 315 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1171.
BECIUM, Lindl. Bot. Reg. (1842) Misc. 43.
OCYMUM, [Tourn.] Linn. Syst. ed. I (1735).
abyssinicum, Hort. Par. ex Lam. Encyc. i. 387 = filamentosum.
acrocephalum, Blume, Bijdr. 834 = Acrocephalus capitatus.
acutum, *Thunb. Fl. Jap.* 248.—Japon.
adscendens, Wight, ex Hook. f. Fl. Brit. Ind. iv. 614 = Orthosiphon tomentosus.

OCIMUM:—
adscendens, *Willd. Sp. Pl.* iii. 166.—Ind. or.
affine, *Hochst. ex Benth. in DC. Prod.* xii. 36.—Abyssin.
africanum, *Lour. Fl. Cochinch.* 370.—Afr. trop.
album, Linn. Mant. i. 85 = O. Basilicum.
album, Roxb. Hort. Beng. 44; Fl. Ind. iii. 15 = canum.
americanum, Auct. ex Benth. Lab. Gen. et Sp. 13 = micranthum.
americanum, Jacq. Hort. Vindob. iii. 45. t. 86 = O. Basilicum.
americanum, Linn. Cent. Pl. i. 15; Amoen. Acad. iv. 276 = canum.
angustifolium, *Benth. in DC. Prod.* xii. 37.—Afr. austr.
anisatum, Hort. ex Benth. Lab. Gen. et Sp. 4 = O. Basilicum.
anisodorum, F. Muell. Fragm. iv. 46 = sanctum.
anosurum, Fenzl, in Flora, xxvii. (1844) 312 = gratissimum.
arborescens, Boj. ex Benth. in DC. Prod. xii. 34 = gratissimum.
aristatum, *Blume, Bijdr.* 833.—Java.
asperum, Roth, Nov. Pl. Sp. 268 = Coleus barbatus.
atropurpureum, Tenore, ex Steud. Nom. ed. II. ii. 202, nomen.—Hab. ?
atrovirens, Bartl. Ind. Sem. Hort. Gotting. (1841).—Cf. Linnaea. xvi. (1842) Litt. 103 = carnosum.
Auricula, Forsk. ex Benth. in DC. Prod. xii. 43, 68 = Plectranthus madagascariensis.
Barrelieri, Roth, Nov. Pl. Sp. 278 = O. Basilicum.
Basilicum, *Linn. Sp. Pl.* 597.—As. occ. et trop.
bengalense, Poir. in Lam. Encyc. iv. 608 = Pogostemon plectranthoides.
Bojeri, *Benth. Lab. Gen. et Sp.* 8.—Madag.
brachiatum, *Blume, Bijdr.* 833.—Java.
bracteosum, *Benth. Lab. Gen. et Sp.* 14.—Afr. trop.
bullatum, Lam. Encyc. i. 384 = O. Basilicum.
Burchellianum, *Benth. Lab. Gen. et Sp.* 8.—Afr. austr.
campechianum, Chapm. Fl. S. U. St. 312 = micranthum.
campechianum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. centr.
canum, *Sims, Bot. Mag.* t. 2452.—As. et Afr. trop.
capitatum, Roth, Nov. Pl. Sp. 276 = Acrocephalus capitatus.
capitellatum, Linn. f. Suppl. 276 = Acrocephalus capitatus.
carnosum, *Link & Otto, ex Benth. Lab. Gen. et Sp.* 11.—Am. trop.
caryophyllatum, Roxb. Hort. Beng. 45; Fl. Ind. iii. 16 = O. Basilicum.
caryophyllatum, *Schweig. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 55.—Hab. ?
caryophyllinum, F. Muell. Fragm. iv. 46 = sanctum.
Cephalotes, Buch.-Ham. ex Wall. Cat. p. 86, sub n. 1563 = Acrocephalus capitatus.
Chandra, Buch.-Ham. ex Wall. l. c. = Acrocephalus capitatus.
chinense, Hort. ex Steud. Nom. ed. II. ii. 202 = Coleus scutellarioides.
ciliare, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 608 = O. Basilicum.
ciliatum, Hornem. Hort. Hafn. 565 = O. Basilicum.
cinereum, R. Br. in Salt, Abyss. App. 64 = Coleus barbatus.
cinnamomeum, *Boj. Hort. Maurit.* 253.—Afr. trop.
citriodorum, *Vis. Sem. Hort. Patav.* (1840).—*Cf. Linnaea*, xv. (1841) *Litt.* 102.—Nubia.
citrodorum, Blanco, Fl. Filip. ed. II. 591 = O. Basilicum.
citronatum, Buch.-Ham. ex Wall. Cat. n. 2720 = gratissimum.
cochleatum, Desf. Tabl. ed. I. 220 = O. Basilicum.
Coetsa, Spreng. Syst. iv. Cur. Post. 223 = Plectranthus Coetsa.
coloratum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 116 = Geniosporum strobiliferum.
coloratum, Hochst. in Flora, xxiv. (1841) 1. Intell. 23 = Orthosiphon Schimperi.
congestum, Spreng. ex Steud. Nom. ed. II. ii. 202 = Mosla ocimoides.
cordifolium, Boj. ex Loud. Hort. Brit. 240 = Bojeri.
cordifolium, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 116 = Plectranthus incanus.
cordifolium, Herb. Madr. ex Wall. Cat. n. 2724 C = Orthosiphon tomentosus.

OCIMUM:—
crispum, *Thunb. Fl. Jap.* 248.—Japon.
cristatum, Buch.-Ham. & Koen. ex Roxb. Fl. Ind. iii. 19 = adscendens.
densiflorum, Roth, Nov. Pl. Sp. 275 = Plectranthus rugosus.
dentatum, Moench, Meth. 413 = O. Basilicum ?
depauperatum, *Vatke, in Linnaea*, xliii. (1880–82) 84.—Afr. trop.
dichotomum, *Hochst. ex Benth. in DC. Prod.* xii. 39.—Abyss.
diffusum, Benth. in Wall. Pl. As. Rar. ii. 14 = Orthosiphon diffusus.
dimidiatum, Schum. & Thonn. Beskr. Guin. Pl. 267 = Moschosma polystachyum.
ebracteatum, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 72 = Sellowii.
fastigiatum, Roth, Nov. Pl. Sp. 277 = Salvia plebeia.
febrifugum, Lindl. in Bot. Reg. t. 753 = viride.
filamentosum, *Forsk. Fl. Aegypt. Arab.* 108.—Arab.; Afr. trop.
fimbriatum, Hort. ex Benth. Lab. Gen. et Sp. 5 = O. Basilicum.
flaccidum, A. Rich. Tent. Fl. Abyss. ii. 179 = Platystema flaccidum.
flexuosum, *Blanco, Fl. Filip.* ed. I. 481.—Ins. Philipp.
flexuosum, *Thunb. Pl. Bras. Dec.* ii. 23, nomen.—Bras.
fluminense, *Vell. Fl. Flum.* 243; vi. t. 11.—Bras.
foetidum, Banks, ex Benth. Lab. Gen. et Sp. 35 = Plectranthus foetidus.
formigense, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxxiii. (1884) 321.—Bras.
Forskolei, *Benth. Lab. Gen. et Sp.* 6; *et in DC. Prod.* xii. 33.—Arab.
frutescens, Burm. f. Fl. Ind. 129 = sanctum.
frutescens, Linn. Sp. Pl. 597 = Perilla ocimoides.
frutescens, Mill. Gard. Dict. ed. VIII. n. 6 = gratissimum.
fruticulosum, *Burch. Trav.* ii. 264.—Afr. austr.
glaucum, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 614 = Orthosiphon tomentosus.
gracile, *Benth. Lab. Gen. et Sp.* 12.—Afr. trop.
grandiflorum, Blume, Bijdr. 835 = Orthosiphon stamineus.
grandiflorum, Lam. Encyc, i. 387; L'Hérit. Stirp. Nov. 89. t. 43 = filamentosum.
gratissimum, Forsk. Fl. Aegypt. Arab. 110 = Plectranthus ovatus.
gratissimum, *Linn. Sp. Pl.* 1197.—Ind. or.
gratissimum, Seem. Fl. Vit. 191 = O. Basilicum.
graveolens, *A. Br. in Flora*, xxiv. (1841) 265.—Abyss.
guineense, Schum. & Thonn. Beskr. Guin. Pl. 264 = viride.
hadiense, Forsk. Fl. Aegypt. Arab. 109 = Plectranthus madagascariensis.
helianthemifolium, *Hochst. in Flora*, xxviii. (1845) 67.—Afr. austr.
heptodon, Beauv. Fl. Owar. ii. 59. t. 94 = viride.
herbaceum, *Roxb. Hort. Beng.* [95] nomen (Quid ?).—Nepal.
hians, Benth. in DC. Prod. xii. 36 = obovatum.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 83.—Afr. trop.
hirsutum, Benth. in Wall. Pl. As. Rar. ii. 14 = sanctum.
hirsutum, Wall. Cat. n. 2717, partim = Orthosiphon diffusus.
hirtum, Banks, ex Benth. Lab. Gen. et Sp. 38 = Plectranthus hirtus.
hispidulum, Schum. & Thonn. Beskr. Guin. Pl. 266 = canum.
hispidum, Lam. Encyc. i. 384 = O. Basilicum.
holosericeum, J. F. Gmel. Syst. 918 = gratissimum.
incanescens, Mart. Reise Bras. ii. 790 = canum.
indicum, Roth, Nov. Pl. Sp. 273 = adscendens.
inflexum, *Thunb. Fl. Jap.* 250.—Japon.
inodorum, Burm. f. Fl. Ind. 130 = sanctum.
integerrimum, Willd. Sp. Pl. iii. 162 = O. Basilicum.
Knyanum, *Vatke, in Linnaea*, xxxvii. (1871–73) 315.—Abyss.
lacerum, [Heyne], ex Wall. Cat. n. 2713 = O. Basilicum.
lamiifolium, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 23, nomen; *et ex Benth. in DC. Prod.* xii. 37.—Abyss.
lanceolatum, Schum. & Thonn. Beskr. Guin. Pl. 268 = O. Basilicum.
laxum, Vahl, ex Benth. Lab. Gen. et Sp. 5 = O. Basilicum.
longiflorum, Buch.-Ham. ex Wall. Cat. n. 2727 = Orthosiphon stamineus.

OCIMUM:—
longistylum, Hochst. ex Benth. in DC. Prod. xii. 34 = menthaefolium.
macrostachyum, Poir. Encyc. Suppl. i. 592 = Geniosporum prostratum.
madagascariense, Pers. Syn. ii. 135 = Plectranthus madagascariensis.
majus, Hort. ex Benth. Lab. Gen. et Sp. 4 = O. Basilicum.
maypurense, Roth, Nov. Pl. Sp. 271 (*mayporense*) = Plectranthus incanus.
medium, Mill. Gard. Dict. ed. VIII. n. 3 = O. Basilicum.
membranaceum, *Benth. Lab. Gen. et Sp.* 15.—Afr. trop.
menthaefolium, Benth. in DC. Prod. xii. 34, partim = O. Basilicum.
menthaefolium, *Hochst. ex Benth. l. c.*—Abyssin.; Arab.
menthoides, Heyne, ex Roth, Nov. Pl. Sp. 274 = viscosum.
menthoides, Linn. Sp. Pl. 598 = Geniosporum prostratum.
mexicanum, *Hort. ex Pasq. Cat. Ort. Bot. Nap.* 71, nomen.—Mexic.
micranthum, *Willd. Enum. Hort. Berol.* 630.—Am. bor. et austr.
minimum, Linn. Sp. Pl. 597 = O. Basilicum.
Miria, Buch.-Ham. in Wall. Cat. n. 2722 = Orthosiphon rubicundus.
molle, Ait. Hort. Kew. ed. I. ii. 322 = Plectranthus incanus.
mollissimum, Wall. ex Hook. f. Fl. Brit. Ind. iv. 615 = Orthosiphon rubicundus.
monachorum, Linn. Mant. i. 85 = sanctum.
monadelphum, R. Br. in Salt, Abyss. App. 64, nomen; Roth, Nov. Pl. Sp. 267 = Coleus spicatus.
monostachyum, Beauv. Fl. Owar. ii. 60 = Plectranthus Palisoti.
montanum, Hook. Bot. Mag. t. 2996 = micranthum.
moschatum, Salisb. Prod. 87 = Moschosma polystachyum.
Nelsonii, *Zipp. ex Span. in Linnaea*, xv. (1841) 333, nomen.—Ins. Timor.
nigrum, Thou. ex Benth. Lab. Gen. et Sp. 5 = O. Basilicum.
nudicaule, *Benth. l. c.* 14.—Bras.
obovatum, *E. Mey. Comm. Pl. Afr. Austr.* 226.—Afr. austr.
obtusifolium, *E. Mey. l. c.* 227.—Afr. austr.
odorum, Salisb. Prod. 87 = O. Basilicum.
ovatum, *Benth. Lab. Gen. et Sp.* 13.—Bras.
ovatum, Willd. ex Benth. in Linnaea, xi. (1837) 344 = Perilomia scutellarioides.
paniculatum, Boj. Hort. Maurit. 253 = gratissimum.
paniculatum, Pers. Syn. ii. 135 = Coleus Persoonii.
peltatum, Hort. ex Steud. Nom. ed. II. ii. 203 = O. Basilicum, micranthum.
peltatum, Schweig. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 55 = Coleus scutellarioides.
petiolare, Lam. Encyc. i. 385 = gratissimum.
Petitianum, *A. Rich. Tent. Fl. Abyss.* ii. 176.—Abyss.
pilosum, Willd. Enum. Hort. Berol. 629 = O. Basilicum.
Plectranthus, J. F. Gmel. Syst. 917 = Coleus galeatus.
polycladum, Link, Enum. Hort. Berol. ii. 119 = Mosla ocymoides.
polystachyon, Linn. Mant. ii. 567 = Moschosma polystachyum.
prostratum, Linn. l. c. 566 = Geniosporum prostratum.
pubescens, Boj. ex Benth. in DC. Prod. xii. 35 = adscendens.
pubescens, Mill. ex Benth. Lab. Gen. et Sp. 13 = micranthum.
punctatum, *Blume, ex Steud. Nom.* ed. II. ii. 203.—Java.
punctatum, Linn. f. Suppl. 275 = Plectranthus punctatus.
punctatum, Thunb. Fl. Jap. 249 = Mosla punctata.
punctulatum, J. F. Gmel. Syst. 917, sphalm. = praec.
racemosum, Thunb. Prod. Pl. Cap. 96 = Plectranthus Thunbergii.
rariflorum, Hochst. in Flora, xxviii. (1845) 67 = obtusifolium.
reflexum, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 126.—Afr. trop.
rigidum, *Benth. Lab. Gen. et Sp.* 14.—Afr. trop.
rigidum, Buch.-Ham. ex Wall. Cat. n. 2721 = Orthosiphon rubicundus.
robustum, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 608 = gratissimum.
rubicundum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 116 = Orthosiphon rubicundus.

OCIMUM:—
rugosum, *Thunb. Fl. Jap.* 249.—Japon.
saccharoides, Willd. ex Wall. Cat. n. 2728 = Coleus barbatus.
salinum, Molina, Sagg. Chile, 139 = O. Basilicum.
salvioides, Heyne, ex Roth, Nov. Pl. Sp. 272 = Plectranthus salvioides.
sanctum, *Linn. Mant.* i. 85.—Geront. trop.
scabrum, Thunb. in Trans. Linn. Soc. ii. (1794) 338 = Mosla punctata.
scabrum, Wight, ex Hook. f. Fl. Brit. Ind. iv. 608 = O. Basilicum.
scutellarioides, Linn. Sp. Pl. ed. II. 833 = Coleus scutellarioides.
scutellarioides, Willd. ex Benth. in Linnaea, xi. (1837) 344 = sanctum.
Selloi, *Benth. Lab. Gen. et Sp.* 6.—Am. austr.
sericeum, Medic. in Act. Acad. Theod. Palat. iv. Phys. (1780) 199. t. 7 = gratissimum.
serpyllifolium, *Forsk. Fl. Aegypt. Arab.* 110.—Arab.
stamineum, Sims, Bot. Mag. t. 2452 = canum.
striatum, Hochst. in Flora, xxviii. (1845) 66 = obovatum.
suave, Willd. Enum. Hort. Berol. 629 = gratissimum.
subserratum, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 609 = sanctum.
suffrutescens, Schum. & Thonn. in Dansk. Vidensk. Selsk. Afh. iv. (1829) 330 = Orthosiphon glabratus.
suffruticosum, Benth. Lab. Gen. et Sp. 709, sphalm. = praec.
sylvaticum, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 270 = Geniosporum Palisoti.
tenellum, *Benth. Lab. Gen. et Sp.* 12.—Madag.
tenuiflorum, Burm. f. Fl. Ind. 129 = Moschosma polystachyum.
tenuiflorum, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 610 = Geniosporum prostratum.
tenuiflorum, Linn. Sp. Pl. 597 = sanctum.
tereticaule, *Poir. Encyc. Suppl.* i. 595.—Afr. trop.
ternifolium, Spreng. Syst. iv. Cur. Post. 224 = Plectranthus ternifolius.
teucriifolium, Hochst. in Flora, xxviii. (1845) 66 = (Orthosiphonis sp.).—Afr. austr.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 265 = tereticaule.
Thonningii, Schum. & Thonn. l. c. 269 = Orthosiphon glabratus.
thymiflorum, Roth, Nov. Pl. Sp. 269 = Orthosiphon tomentosus.
thyrsiflorum, Linn. Mant. i. 84 = O. Basilicum.
tomentosum, Lam. Encyc. i. 387 = sanctum.
triste, Roth, Nov. Pl. Sp. 270 = Orthosiphon tomentosus.
tuberosum, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 18 = Orthosiphon rubicundus.
Tweedianum, *Benth. in DC. Prod.* xii. 38.—Uruguay.
urticaefolium, Hort. ex Benth. Lab. Gen. et Sp. 5 = O. Basilicum.
urticifolium, Roth, Catalect. fasc. ii. 52 = gratissimum.
Vaalae, *Forsk. Fl. Aegypt. Arab.* 111.—Arab.
verticillatum, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 609 = adscendens.
verticillatum, Linn. f. Suppl. 276 = Plectranthus Thunbergii.
villosum, Roxb. Hort. Beng. 44; Fl. Ind. iii. 13 = sanctum.
villosum, Weinm. in Syll. Ratisb. i. (1824) 121 = gratissimum.
virgatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 116 = Orthosiphon rubicundus.
virgatum, *Thunb. Fl. Jap.* 250.—Japon.
viride, *Willd. Enum. Hort. Berol.* 629.—Afr. trop.
viridiflorum, *Roth, Catalect.* fasc. ii. 54.—Hab.?
viscosum, *Roth, Nov. Pl. Sp.* 274.—Ind. or.
Zatarhendi, Forsk. Fl. Aegypt. Arab. 109 = Coleus spicatus.
zeylanicum, Medic. in Act. Acad. Theod. Palat. iv. Phys. (1780) 197. t. 6 = gratissimum.

OCKEA, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. i. 105 (1815) = **Adenandra**, Willd. (Rutac.).

OCKENIA, Steud. Nom. ed. II. i. 24 (1840) = praec.

OCKIA, Bartl. & Wendl. f. Diosm. 69 (1824) = praec.

OCNERON, Rafin. Fl. Tellur. iv. 120 (1836) = **Staehelina**, Linn. (Compos.).
Aquifolium, Rafin. l. c. (Quid?).—Hab.?
cordatum, Rafin. l. c. (Quid?).—Hab.?

OCOTEA, Aubl. Pl. Gui. ii. 780. t. 310 (1775).
LAURINEAE, Benth. & Hook. f. iii. 157.
AGRIODAPHNE, Steud. Nom. ed. II. i. 38 (1840).
ANIBA, Aubl. Pl. Gui. i. 327 (1775).
BALANOPSIS, Rafin. Sylva Tellur. 134 (1838).
CALYCODAPHNE, Boj. Hort. Maurit. 273 (1837).
CAMPHOROMOEA, Nees, Laurin. Expos. 16 (1833).
CANNELLA, Schott, ex Meissn. in DC. Prod. xv. i. 103, in syn. (1864).
CEDROTA, Schreb. Gen. i. 259 (1789).
CERAMOCARPIUM, Nees, ex Meissn. in DC. Prod. xv. i. 111 (1864).
CERAMOPHORA, Nees, l. c. (1864).
EVONYMODAPHNE, Nees, Syst. Laurin. 263 (1836).
GYMNOBALANUS, Nees & Mart. in Linnaea, viii. (1833) 38.
LEPTODAPHNE, Nees, Laurin. Expos. (1833) 16; Syst. Laurin. 358 (1836).
MESPILODAPHNE, Nees, in Linnaea, viii. (1833) 45.
NEMODAPHNE, Meissn. in DC. Prod. xv. i. 109 (1864).
OREODAPHNE, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 39.
PETALANTHERA, Nees, Laurin. Disp. Prog. 15 (1833); Syst. Laurin. 346 (1836).
STRYCHNODAPHNE, Nees, in Linnaea, viii. (1833) 39.
TELEIANDRA, Nees, l. c. (1836) 46.
acuminata, *Baker, in Journ. Linn. Soc.* xx. (1883) 242.—Madag.
aenea, Mart. ex Nees, Syst. Laurin. 394 = tristis.
amara, Mart. Reise Bras. iii. 1128 = Nectandra cymbarum.
amazonum, Mart. ex Nees, Syst. Laurin. 282 = Nectandra amazonum.
angustifolia, *Pohl, ex Meissn. in DC. Prod.* xv. i. 162.—Bras.
angustifolia, Schrad. in Goett. Gel. Anz. i. (1821) 711 = Nectandra angustifolia.
attenuata, Nees, ex Wall. Pl. As. Rar. ii. 71 = Phoebe attenuata.
bifida, Poepp. ex Nees, Syst. Laurin. 351 = Pleurothyrium bifidum.
Bofa, *H. B. & K. Nov. Gen. et Sp.* ii. 161.—N. Granat.
botryophylla, *Klotzsch & Karst. ex Nees, in Linnaea*, xxi. (1848) 524.—N. Granat.
bracteolata, *Mart. ex Nees, Syst. Laurin.* 347.—Bras.
bullata, *E. Mey. ex Meissn. in DC. Prod.* xv. i. 118.—Afr. austr.
caracasana, Klotzsch, ex Nees, in Linnaea, xxi. (1848) 516, partim = Nectandra discolor.
carolinensis, Kostel. Allg. Med. Pharm. Flora, ii. 482 = Persea carolinensis.
caryophyllacea, Kostel. l. c. = Dicypellium caryophyllatum.
chrysophylla, Poepp. ex Nees, Syst. Laurin. 351 = Pleurothyrium chrysophyllum.
citrosmoides, *Mart. ex Nees, l. c.* 384.—Bras.
commutata, *Nees, ex Meissn. in DC. Prod.* xv. i. 120, in syn.—Guian.
Cujumary, *Mart. Reise Bras.* iii. 1128.—Bras.
cuspidata, Mart. ex Nees, Syst. Laurin. 330 = Nectandra cuspidata.
cymbarum, H. B. & K. Nov. Gen. et Sp. ii. 166 = Nectandra cymbarum.
cymbarum, Poepp. ex Nees, Syst. Laurin. 120 = Phoebe cubensis.
cymbarum, Spreng. ex Nees, l. c. 213 = Cryptocarya moschata.
daphnoides, Mart. ex Nees, l. c. 402 = lanceolata.
dealbata, Poepp. ex Meissn. in DC. Prod. xv. i. 88 = Aydendron permolle.
declinata, Blume, Bijdr. 572 = Phoebe declinata.
desertorum, Mart. ex Nees, Syst. Laurin. 260 = Aydendron desertorum.
discolor, H. B. & K. Nov. Gen. et Sp. ii. 163 = Nectandra discolor.
effusa, *Hemsl. Biol. Centr. Am. Bot.* iii. 73.—Mexic.
elongata, Nees, ex Meissn. in DC. Prod. xv. i. 31 = Phoebe montana.
excelsa, Blume, Bijdr. 572 = Phoebe excelsa.
firmula, *Mart. ex Nees, Syst. Laurin.* 250.—Venezuela.

OCOTEA:—
floribunda, *Benth. & Hook. f. Gen.* iii. 158.—Bras.
foetens, *Benth. & Hook. f. l. c.*—Ins. Canar.
glaucescens, Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe pallida.
globosa, Schlecht. & Cham. in Linnaea, vi. (1831) 366 = Nectandra sanguinea.
glomerata, *Benth. & Hook. f. Gen.* iii. 158.—Guiana.
guianensis, *Aubl. Pl. Gui.* ii. 781.—Guiana.
helicterifolia, *Hemsl. Biol. Centr. Am. Bot.* iii. 73.—Mexic.
Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 453.—Madag.
incana, Schott, ex Meissn. in Mart. Fl. Bras. v. ii. 252 = Nectandra rigida.
indecora, *Schott, ex Meissn. in DC. Prod.* xv. i. 102.—Bras.
indica, Kostel. Allg. Med. Pharm. Flora, ii. 482 = Persea indica.
japurensis, Mart. ex Nees, Syst. Laurin. 335 = Nectandra japurensis.
Klotzschiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 73.—Mexic.
lanceolaria, Voigt, Hort. Suburb. Calc. 309 = Phoebe lanceolata.
lanceolata, Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe lanceolata.
lanceolata, *Nees, Syst. Laurin.* 474.—Bras.
latifolia, H. B. & K. Nov. Gen. et Sp. ii. 165 = Nectandra polita.
Leucoxylon, *Benth. & Hook. f. Gen.* iii. 158.—Ind. occ.
Lhotzkii, *Nees, Syst. Laurin.* 475.—Bras.
ligustrina, Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe declinata.
lineata, H. B. & K. Nov. Gen. et Sp. ii. 165 = Persea lineata.
longifolia, *H. B. & K. l. c.* 164.—N. Granat.
lucida, Mart. ex Nees, Syst. Laurin. 334 = Nectandra lucida.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 165 = Nectandra macrophylla.
Maximilianea, Mart. ex Nees, Syst. Laurin. 472 = puberula.
maynensis, Poepp. ex Meissn. in DC. Prod. xv. i. 32 = Phoebe peruviana.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* iii. 73.—Mexic.
microcarpa, *Poepp. ex Meissn. in DC. Prod.* xv. i. 112.—Bras.; Peruv.
minarum, Mart. ex Nees, Syst. Laurin. 303, ex parte = Nectandra psammophila.
mollis, H. B. & K. Nov. Gen. et Sp. ii. 164 = Nectandra villosa.
mucosa, Hemsl. Biol. Centr. Am. Bot. iii. 73 = Acrodiclidium salicifolium.
nitidula, Mart. ex Nees, Syst. Laurin. 313 = Nectandra nitidula.
obtusifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 165.—N. Granat.
obtusifolia, Mart. ex Nees, Syst. Laurin. 260 = Aydendron desertorum.
opifera, *Mart. Reise Bras.* iii. 1128.—Bras.
pallida, Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe pallida.
paniculata, Nees, l. c. = Phoebe paniculata.
Pichurim, H. B. & K. Nov. Gen. et Sp. ii. 166 = Aydendron Laurel.
Plumieri, Kostel. Allg. Med. Pharm. Flora, ii. 482 = Persea carolinensis.
polyantha, *Mart. ex Nees, Syst. Laurin.* 458.—Bras.
pretiosa, *Benth. & Hook. f. Gen.* iii. 158.—Bras.
psammophila, Mart. ex Nees, Syst. Laurin. 303 = Nectandra psammophila.
psychotrioides, Billb. ex Meissn. in DC. Prod. xv. i. 150 = Nectandra amazonum.
psychotrioides, *H. B. & K. Nov. Gen. et Sp.* ii. 162.—Mexic.
puberula, *Nees, Syst. Laurin.* 472.—Am. austr.
pubescens, Nees, in Wall. Pl. As. Rar. ii. 71 = Phoebe paniculata.
Puchury-major, Mart. Reise Bras. iii. 1127 = Nectandra Puchury-major.
Puchury-minor, Mart. l. c. = Nectandra Puchury-minor.
pulchella, *Mart. in Buch. Rep.* xxxv. (1830) 172.—Bras.
ramentacea, Mart. ex Nees, Syst. Laurin. 553 = Litsea glaucescens.

OCOTEA:—
rigida, H. B. & K. Nov. Gen. et Sp. ii. 163 = Nectandra rigida.
riparia, Mart. ex Nees, Syst. Laurin. 330 = Nectandra cuspidata.
rufo-tomentosa, *Mart. ex Nees, l. c.* 153.—Bras.
salicifolia, Hook. & Arn. Bot. Beech. Voy. 309 = Nectandra glabrescens.
salicifolia, H. B. & K. Nov. Gen. et Sp. ii. 166 = Nectandra sanguinea.
salicifolia, *Nees, Syst. Laurin.* 298.—Bras.
sanguinea, J. S. Presl, Rostl. ii. 60 = Nectandra sanguinea.
Schomburgkiana, *Benth. & Hook. f. Gen.* iii. 158.—Guiana.
sericea, H. B. & K. Nov. Gen. et Sp. ii. 162 = guianensis.
sericea, Nees, in Wall. Pl. As. Rar. ii. 71 = Machilus sericea.
Sieberi, *Hemsl. Biol. Centr. Am. Bot.* iii. 73.—Mexic.
squarrosa, *Mart. ex Nees, Syst. Laurin.* 326.—Bras.
striata, Buek, Ind. ad DC. Prod. iv. 234 = Nectandra striata.
suaveolens, *Benth. & Hook. f. ex Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 437.—Reg. Argent.
subacris, *Mart. ex Nees, Syst. Laurin.* 410.—Bras.
subtriplinervia, *Hemsl. Biol. Centr. Am. Bot.* iii. 74.—Mexic.
tampicensis, *Hemsl. l. c.*—Mexic.
trichophlebia, *Baker, in Journ. Linn. Soc.* xx. (1883) 242.—Madag.
tristis, *Mart. ex Nees, Syst. Laurin.* 304.—Bras.
turbacensis, H. B. & K. Nov. Gen. et Sp. ii. 162 = Nectandra turbacensis.
turbacensis, Poepp. ex Nees, Syst. Laurin. 368 (= *Goeppertia longifolia*).—Peruv.
umbelliflora, Blume, Bijdr. 573 = Phoebe umbelliflora.
umbrosa, H. B. & K. Nov. Gen. et Sp. ii. 161 (Nectandrae sp.).—N. Granat.
umbrosa, *Mart. ex Nees, Syst. Laurin.* 388.—Bras.
variabilis, *Mart. l. c.* 429.—Bras.
velutina, *Mart. l. c.* 386.—N. Granat.
venulosa, *Benth. & Hook. f. Gen.* iii. 158.—Bras.
virgultosa, *Mart. ex Nees, Syst. Laurin.* 304.—Bras.
xanthocalyx, *Mart. l. c.* 312.—Bras.

OCTADENIA, R. Br. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 41 (1836) = **Alyssum**, Tourn. (Crucif.).
lybica, R. Br. ex Fisch. & Mey. l. c. = A. libycum.
maritima, Fisch. & Mey. l. c. = A. maritimum.

OCTADESMIA, Benth. in Journ. Linn Soc. xviii. (1881) 311. *ORCHIDEAE*, Benth. & Hook. f. iii. 525.
Dilomilis, Rafin. Fl. Tellur. iv. 43 (1836), nomen prius.
elata, *Benth. & Hook. f. Gen.* iii. 525.—Jamaica.
monophylla, *Benth. & Hook. f. l. c.* 526.—Ind. occ.
montana, Benth. & Hook. f. l. c. 525 = serratifolia.
serratifolia, *Benth. in Journ. Linn. Soc.* xviii. (1881) 311.—Ind. occ.

OCTANEMA, Rafin. Sylva Tellur. 112 (1838) = **Capparis**, Linn.
angustifolia, Rafin. l. c. = C. angustifolia.
crotonoides, Rafin. l. c. = C. crotonoides.
incana, Rafin. l. c. = C. incana.
scabrida, Rafin. l. c. = C. scabrida.

OCTARILLUM, Lour. Fl. Cochinch. 90 (1790) = **Elaeagnus**, Tourn.
fruticosum, Lour. l. c.—Cochinch.

OCTARRHENA, Thw. Enum. Pl. Zeyl. 305 (1861) = **Phreatia**, Lindl. (Orchid.).
parvula, Thw. l. c. 305.—Zeylan.

OCTAS, Jack, in Malay. Misc. ii. vii. (1822). (*INC. SED.*).
spicata, *Jack, l. c.*—Malaya.

OCTAVIA, DC. Prod. iv. 464 (1830) = **Lasianthus**, Jack (Rubiac.).
sessiliflora, DC. l. c.—Guiana.

OCTELISIA, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legumin.).
aurea, Rafin. l. c. 130 = C. Sophera.

OCTELLA, Rafin. Sylva Tellur. 96 (1838) = **Melastoma**, Linn. etc.
alpina, Rafin. l. c. 97 = Charianthus coccineus.
angustifolius, Rafin. l. c. 96 = Tetragygia angustifolia.
axillaris, Rafin. l. c. 97 (Quid ?).—Hab. ?
capillaris, Rafin. l. c. = Sagraea capillaris.
coccinea, Rafin. l. c. = Charianthus purpureus.
fascicularis, Rafin. l. c. = Henriettella fascicularis.
glabrata, Rafin. l. c. = Calycogonium glabratum.
hispida, Rafin. l. c. = Tetrazygia hispida.
lateriflora, Rafin. l. c. = Henriettella lateriflora.
microphyla, Rafin. l. c. = Ossaea microphylla, *Triana.*
repens, Rafin. l. c. = M. repens.
umbrosa, Rafin. l. c. = Sagraea umbrosa.
vaccinoides, Rafin. l. c. = Miconia vaccinioides.
verticillata, Rafin. l. c. = Ossaea verticillata.
virgata, Rafin. l. c. = Mecranium virgatum.

OCTHOCHARIS, G. Don, Gen. Syst. ii. 731 (1832) = **Ochthocharis**, Blume (Melastom.).

OCTIMA, Rafin. Alsog. Am. 42 (1838) = **Populus**, Linn. (Salicin.).
canescens, Rafin. l. c. = P. alba.

OCTOCERAS, Bunge, in Arbeit. Nat. Ver. Riga, i. (1847) 172. *CRUCIFERAE*, Benth. & Hook. f. i. 97.
Lehmannianum, *Bunge, l. c.*—Persia; Afghan.

OCTOCLINIS, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 21 = **Callitris**, Vent. (Conif.).
Backhousi, W. Hill, in Cat. Nat. & Industr. Prod. Queensl. (1862) 16 = C. calcarata.
juniperoides, Carr. Conif. ed. II. 790.—N. Caled.
Macleyana, Muell. in Trans. Phil. Inst. Vict. ii. (1858) 22 = C. Macleyana.

OCTODON, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 74 (1827). *RUBIACEAE*, Benth. & Hook. f. ii. 145.
filifolium, *Schum. & Thonn. l. c.*—Afr. trop.
setosum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 242.—Afr. trop.

OCTOGONIA, Klotzsch, in Linnaea, xii. (1838) 233 = **Simocheilus**, Klotzsch (Ericac.).
glabella, Klotzsch, l. c. = S. glabellus.
hirta, Klotzsch, l. c. = S. hirtus.

OCTOLEPIS, Oliver, in Journ. Linn. Soc. viii. (1865) 161. t. 12. *THYMELAEACEAE*, Benth. & Hook. f. iii. 201.
Casearia, *Oliver, l. c.*—Afr. trop.

OCTOLOBUS, Welw. in Trans. Linn. Soc. xxvii. (1869) 17. *STERCULIACEAE*, Benth. & Hook. f. i. 982.
spectabilis, *Welw. l. c.* 18. t. 6.—Afr. trop.

OCTOMELIS, Miq. Fl. Ind. Bat. Suppl. 336 (1860). *DATISCEAE*, Benth. & Hook. f. i. 845.
sumatranus, *Miq. l. c.*—Sumatra.

OCTOMERIA, R. Br. in Ait. Hort. Kew. ed. II. v. 211 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 493.
Aspegrenia, Poepp. & Endl. Nov. Gen. et Sp. ii. 12. t. 116 (1837).
Enothrea, Rafin. Fl. Tellur. iv. 43 (1836).
aetheoantha, *Rodrig. Orch. Nov.* ii. 106.—Bras.
albina, *Rodrig. l. c.* 107.—Bras.
albopurpurea, *Rodrig. l. c.*—Bras.
aloëfolia, *Rodrig. l. c.* 113.—Bras.
alpina, *Rodrig. l. c.* 102.—Bras.
Baueri, *Lindl. Gen. et Sp. Orch.* 10.—Ind. occ.
caespitosa, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 596, nomen.—Ins. Trinit.
candida, *Lodd. Cat. l. c.*, nomen.—Bras.
chamaeleptotes, *Reichb. f. in Linnaea*, xxii. (1849) 817.—Bras.
cochlearis, *Reichb. f. in Gard. Chron.* (1881) i. 266.—Bras.
concolor, *Rodrig. Orch. Nov.* ii. 100.—Bras.
convallarioides, Wall. ex Lindl. Gen. et Sp. Orch. 70 = Eria convallarioides.
crassifolia, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Bras.
densiflora, *Rodrig. Orch. Nov.* ii. 97.—Bras.
diaphana, *Lindl. Bot. Reg.* (1839) *Misc.* 91.—Bras.
ementosa, *Rodrig. Orch. Nov.* ii. 102.—Bras.
exchlorophyllata, *Rodrig. l. c.* 103.—Bras.
fasciculata, *Rodrig. l. c.* i. 32.—Bras.
flava, Wall. ex Steud. Nom. ed. II. ii. 204 = Eria flava.
geraënsis, *Rodrig. Orch. Nov.* ii. 295.—Bras.
Glazioveana, *Regel, in Act. Hort. Petrop.* viii. (1883) 277.—Bras.
gracilis, *Lodd. ex Lindl. Bot. Reg.* (1838) *Misc.* 36.—Bras.
gracilis, *Rodrig. Orch. Nov.* ii. 108.—Bras.
graminifolia, *R. Br. in Ait. Hort. Kew.* ed. II. v. 211.—Ind. occ.
grandiflora, *Lindl. Bot. Reg.* (1842) *Misc.* 64.—Bras.
helvola, *Rodrig. Orch. Nov.* ii. 110.—Bras.
juncifolia, *Rodrig. l. c.*—Bras.
lancaefolia, *Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 596, nomen.—Bras.
leptophylla, *Rodrig. Orch. Nov.* ii. 112.—Bras.
lichenicola, *Rodrig. l. c.*—Bras.
linearifolia, *Rodrig. l. c.* 106.—Bras.
lithophila, *Rodrig. l. c.* 295.—Bras.
lobulosa, *Reichb. f. in Hamb. Gartenz.* xiv. (1858) 215.—Hab. ?
Loddigesii, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Bras. ?
micrantha, *Rodrig. Orch. Nov.* i. 33.—Bras.
montana, *Rodrig. l. c.* ii. 108.—Bras.
multiflora, *Rodrig. l. c.* 98.—Bras.
ochroleuca, *Rodrig. l. c.* i. 31.—Bras.
oxychela, *Rodrig. l. c.* ii. 99.—Bras.
paucifolia, Breda, Kuhl & Hasselt, Orch. [t. 11] = Eria paucifolia.
petulans, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 59.—Hab. ?
pinicola, *Rodrig. Orch. Nov.* ii. 101.—Bras.
praestans, *Rodrig. l. c.* 112.—Bras.
pubescens, Spreng. Syst. iv. Cur. Post. 310 = Eria flava.
pusilla, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Bras.
racemosa, Breda, Kuhl & Hasselt, Orch. [t. 12] = Eria micrantha.
rigida, *Rodrig. Orch. Nov.* ii. 104.—Bras.
robusta, *Rodrig. l. c.* 97.—Bras.
Rodeiensis, *Rodrig. l. c.* 105.—Bras.
rosea, Spreng. Syst. iv. Cur. Post. 310 = Eria rosea.
rubrifolia, *Rodrig. Orch. Nov.* i. 31.—Bras.
sarcophylla, *Rodrig. l. c.* ii. 104.—Bras.
Saundersiana, *Reichb. f. in Gard. Chron.* (1880) i. 264.—Bras.
scirpoidea, *Reichb. f. in Bot. Zeit.* x. (1852) 856.—Peruv.
secunda, Wall. ex Lindl. Gen. et Sp. Orch. 63 = Eria stricta.
semiteres, *Regel, in Ann. Sc. Nat.* Sér. IV. vi. (1856) 373.—Bras.
serratifolia, Hook. Bot. Mag. t. 2823 = Octadesmia serratifolia.
spatulata, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 424.—Hab. ?
stellaris, *Rodrig. Orch. Nov.* ii. 99.—Bras.
stellata, Spreng. Syst. iv. Cur. Post. 310 = Eria stellata.
surinamensis, *H. C. Focke, in Tijdschr. voor Wetensch.* ii. (1849) 200.—Guian.
teretifolia, *Batem. ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 651.—Bras.
teretifolia, Rodrig. Orch. Nov. i. 32 = leptophylla.
tricolor, *Reichb. f. in Gard. Chron.* (1872) 1035.—Bras.
tridentata, *Lindl. Bot. Reg.* (1839) *Misc.* 35.—Guiana.
truncicola, *Rodrig. Orch. Nov.* ii. 101.—Bras.
vaginata, Breda, Kuhl & Hasselt, Orch. [t. 5] = Eria vaginata.
Warmingii, *Reichb. f. Otia Bot. Hamb.* 94.—Bras.

OCTOMERIA, D. Don, Prod. Fl. Nep. 31 (1825) = **Eria**, Lindl. (Orchid.).
spicata, D. Don. l. c. = E. convallaroides.

OCTOMERIA, Pfeiff. Nom. ii. 474 (1874) sphalm. = **Otomeria**, Benth. (Rubiac.).

OCTOMERIS, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 52; et xvii. (1852) 378 = **Heterotrichum**, DC. et **Miconia**, Ruiz & Pav. (Melastom.).
Bonplandii, Naud. l. c. iv. (1845) 54 = H. Octonum.
bracteosa, Naud. l. c. xvii. (1852) 379 = Anaectocalyx bracteosa.
macrodon, Naud. l. c. iv. (1845) 53 = H. macrodon.
paupercula, Naud. l. c. xvii. (1852) 281 = M. paupercula.
polyandra, Naud. l. c. 382 = M. Benthamiana.
rostrata, Naud. l. c. iv. (1845) 53 = M. tuberculata.
Schlimii, Naud. l. c. xvii. (1852) 379 = H. Schlimii.
tuberculata, *Naud. l. c.* iv. (1845) 53 = M. tuberculata.

OCTONUM, Rafin. Sylva Tellur. 95 (1838) = **Heterotrichum**, DC. (Melastom.).
Humboldti, Rafin. l. c. = H. Octonum.

OCTOPERA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 156 = **Erica**, Tourn.
Bergiana, D. Don, l. c. = E. Bergiana.

OCTOPLEURA, Griseb. Fl. Brit. W. Ind. 260 (1860). *MELASTOMACEAE*, Benth. & Hook. f. i. 769.
ciliata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 146.—Ins. Trinit.
diversifolia, *Triana, l. c.* 145.—N. Granat.
Epibaterium, Ind. Kew. supra, i. 563 in syn., sphalm. = Ossaea Epibaterium.
macrophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 145.—Am. centr.
micrantha, *Griseb. Fl. Brit. W. Ind.* 260.—Jamaic.
quinquenervia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 145.—Am. centr.
robusta, *Triana, l. c.* 146.—N. Granat.; Peruv.
rubescens, *Triana, l. c.*—N. Granat.

OCTOPLEURA, Spruce, ex Prog. in Mart. Fl. Bras. vi. 1. 212. t. 58. f. 1 (1865) = **Neurotheca**, Salisb. (Gentian.).
loeselioides, Spruce, ex Prog. l. c. = N. loeselioides.

OCTOPLIS, Rafin. Fl. Tellur. iv. 105 (1836) = **Cansjera**, Juss. (Olacin.).
polistachya, Rafin. l. c. = C. Rheedii.

OCTOTROPIS, Bedd. Fl. Sylv. t. 327 (1873); et Forest. Man. Bot. 134/12. *RUBIACEAE*, Benth. & Hook. f. ii. 1229.
travancorica, *Bedd. ll. cc.* 327; 134/13.—Ind. or.

OCYMUM, Tourn. ex Linn. Gen. ed. I. 173 (1737) = **Ocimum**, Tourn. (Labiat.).

ODACMIS, Rafin. New Fl. Am. iv. 39 (1836) (Gen. sedis incert.).
fascicularis, Rafin. l. c. 40 (Quid?).—Am. bor.

ODICARDIS, Rafin. Fl. Tellur. iv. 55 (1836) = **Veronica**, Tourn. (Scroph.).
Crista-galli, Rafin. l. c. = V. Crista-galli.

ODINA, Netto, in Ann. Sc. Nat. Sér. V. v. (1866) 85 = **Marupa**, Miers (Simarub.).
Francoana, Netto, l. c. = M. Francoana.

ODINA, Roxb. Hort. Beng. 29 (1814); Fl. Ind. ii. 293 (1832). *ANACARDIACEAE*, Benth. & Hook. f. i. 423.
CALESIAM, Adans. Fam. ii. 446 (1763).
HABERLIA, Dennst. Schluess. Hort. Malab. 30 (1818).
KOKKIA, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 206 (1850).
LANNEA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 153. t. 42 (1832).
WIRTGENIA, Jungh. in Flora, xxvii. (1844) 624.
acida, *Walp. Rep.* v. 412.—Afr. trop.
asplenifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 508.—Ins. Socotra.
Barteri, *Oliver, Fl. Trop. Afr.* i. 446.—Afr. trop.
discolor, *Sond. in Linnaea*, xxiii. (1850) 25.—Afr. austr.
edulis, *Sond. in Harv. & Sond. Fl. Cap.* i. 503.—Afr. austr.
fraxinifolia, Fenzl, ex Oliver, Fl. Trop. Afr. i. 446 = fruticosa?

ODINA:—
fruticosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 141.—Abyss.
gummifera, *Blume, Mus. Bot. Lugd. Bat.* i. 206.—Java.
humilis, *Oliver, Fl. Trop. Afr.* i. 447.—Afr. trop.
obovata, *Hook. f. ex Oliver, l. c.*—Afr. trop.
Oghigee, Hook. f. in Hook. Niger Fl. 286 = acida.
ornifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 507.—Ins. Socotra.
pinnata, Rottl. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 209, in obs. = O. Wodier?
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 140.—Abyss.
Schweinfurthii, *Engl. in DC. Monog. Phan.* iv. 273.—Afr. trop.
speciosa, *Blume, Mus. Bot. Lugd. Bat.* i. 206.—N. Guin.
triphylla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 140.—Abyss.
velutina, *Endl. ex Walp. Rep.* i. 550.—Afr. trop.
Wodier, *Roxb. Hort. Beng.* 29; *Fl. Ind.* ii. 293.—Ind. or.; Burma.

ODISCA, Rafin. Sylva Tellur. 80 (1838) = **Colea**, Boj. (Bignoniac.).
rosea, Rafin. l. c. = C. mauritiana.

ODODECA, Rafin. Neogenyt. 3 (1825) = **Homalium**, Jacq. (Samyd.).

ODOGLOSSA, Rafin. New Fl. Am. ii. 15 (1836) = **Coreopsis**?, Linn. (Compos.).
heterophylla, Rafin. l. c. 16 (Quid?).—Am. bor.

ODOLLAM, Adans. Fam. ii. 171 (1763) = **Cerbera**, Linn. (Apocyn.).

ODOLLAMIA, Rafin. Sylva Tellur. 162 (1838) = **Cerbera**, Linn. (Apocyn.).
malabarica, Rafin. l. c. = C. Odollam?
Manghas, Rafin. l. c. = C. Odollam.
moluca, Rafin. l. c. = C. Odollam?

ODONECTIS, Rafin. in Med. Repos. N. York, v. (1808) 357 = **Pogonia**, Juss. (Orchid.).
verticillata, Rafin. l. c. = P. verticillata.

ODONIA, Bertol. Lucub. 35 (1822) = **Galactia**, P. Br. (Legumin.).
speciosa, Spreng. Syst. iv. Cur. Post. 279 (= *Collaea speciosa*).—Peruv.
tomentosa, Bertol. Lucub. 35 = G. tomentosa.
trinervia, Spreng. Syst. iv. Cur. Post. 279 = Atylosia Candollei.

ODONTADENIA, Benth. in Hook. Journ. Bot. iii. (1841) 242. *APOCYNACEAE*, Benth. & Hook. f. ii. 723.
ANISOLOBUS, A. DC. in DC. Prod. viii. 395 (1844).
CYLICADENIA, Lem. Illustr. Hortic. ii. (1855) Misc. 9.
angustifolia, A. DC. in DC. Prod. viii. 360 = cordata.
cordata, *A. DC. l. c.*—Bras.; Guian.
coriacea, Muell. Arg. in Linnaea, xxx. (1859–60) 450 = Angadenia coriacea.
formosa, Miers, Apocyn. S. Am. 127 = grandiflora.
geminata, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 119.—Bras.; Guian.
grandiflora, Miers, Apocyn. S. Am. 127 = speciosa.
grandiflora, *Miq. Stirp. Surin. Sel.* 166.—Guian.
Harrisii, Miers, Apocyn. S. Am. 128 = speciosa.
Hoffmannseggiana, Klotzsch, ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 117 = speciosa.
hypoglauca, *Muell. Arg. l. c.* 118.—Bras.
lucida, Muell. Arg. l. c. 120, in obs. = Rhabdadenia lucida.
macrocalyx, Miers, Apocyn. S. Am. 128 = Tabernaemontana macrocalyx.
nitida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 118 = cordata.
Poeppigii, Muell. Arg. l. c. 119.—Bras.
speciosa, *Benth. in Hook. Journ. Bot.* iii. (1841) 242.—Bras.; Guian.
sylvestris, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 117.—Bras.; Guian.

ODONTANDRA, Willd. ex Roem. & Schult. Syst. v. p. xli. 511 (1819) = **Trichilia**, P. Br. (Meliac.).
acuminata, Humb. & Bonpl. ex Roem. & Schult. l. c. 511 = T. acuminata.
acuminata, Karst. Fl. Columb. ii. 49. t. 125 = T. Karstenii.
appendiculata, Triana & Planch. in Ann. Sc. Nat. Sér. V. xv. (1872) 375 = T. appendiculata.
elegans, Triana & Planch. l. c. 376, in obs. = T. elegans.
Karstenii, Triana & Planch. l. c. 373 = T. Karstenii.
parviflora, Triana & Planch. l. c. 375 = T. parviflora.
Pseudostipulare, Triana & Planch. l. c. in obs. = T. Pseudostipularis.
quadrijuga, Triana & Planch. l. c. 374 = T. quadrijuga.
tuberculata, Triana & Planch. l. c. = T. tuberculata.

ODONTANDRIA, G. Don, Gen. Syst. i. 686 (1831), sphalm. = praec.

ODONTANTHERA, Wight, in Lindl. Veg. Kingd. 626 (1847), nomen. *ASCLEPIADEAE*.—Cf. Benth. & Hook. f. ii. 739.

ODONTARRHENA, C. A. Mey. in Ledeb. Fl. Alt. iii. 58 (1831) = **Alyssum**, Tourn. (Crucif.).
alpestris, Ledeb. Fl. Ross. i. 142 = A. alpestre.
argentea, Ledeb. l. c. 141 = A. argenteum.
Bourgaei, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 4 = A. alpestre.
castellana, Jord. & Fourr. l. c. 2 = A. alpestre.
clypeolata, Jord. & Fourr. l. c. 7 = A. argenteum.
depressa, Jord. & Fourr. l. c. 6 = A. argenteum.
diffusa, Jord. & Fourr. l. c. 5 = A. argenteum.
Fischeriana, C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 245 = A. Fischerianum.
granatensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 3 = A. alpestre.
hirta, Schrenk, Enum. Pl. Nov. 95.—Sibir. altaic.
lycia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 3 = A. alpestre.
Marschalliana, C. A. Mey. Verz. Pfl. Cauc. 182 = A. Marschallianum.
microphylla, C. A. Mey. in Ledeb. Fl. Alt. iii. 63 = A. alpestre.
muralis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 5 = A. argenteum.
muralis, Schur, Enum. Pl. Transs. 61 = A. argenteum.
murcica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 2 = A. alpestre.
obovata, C. A. Mey. in Ledeb. Fl. Alt. iii. 61 = A. sibiricum.
obtusifolia, C. A. Mey. Verz. Pfl. Cauc. 181 = A. alpestre.
pyrenaica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 2 = A. alpestre.
rubricaulis, Jord. & Fourr. l. c. 6 = A. argenteum.
saxatilis, Jord. & Fourr. l. c. 1 = A. alpestre.
surculosa, Schott & Kotschy, ex Boiss. Fl. Orient. i. 268 = A. alpestre.
tmolea, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 7 = A. argenteum.
tortuosa, C. A. Mey. in Ledeb. Fl. Alt. iii. 60 = A. tortuosum.

ODONTEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 391 = **Bupleurum**, Tourn. (Umbell.).
aristata, Fourr. l. c. = B. aristatum.

ODONTEILEMA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 587 = **Acalypha**, Linn. (Euphorb.).
Clausseni, Turcz. l. c. 588 = A. Clausseni.

ODONTITES, [Riv.] Hall. Enum. Stirp. Helv. ii. 627 (1742) = **Bartsia**, Linn. (Scrophul.).
albidula, Mabille, Rech. Pl. Corse, fasc. ii. 44 = *corsica*.
alpestris, Jour. & Fourr. Brev. Pl. Nov. fasc. ii. 87 = B. purpurea.
aspera, Boiss. Voy. Bot. Espagne, ii. 473 = B. aspera.
Aucheri, Boiss. Diagn. Ser. I. iv. 74.—Graecia; Oriens.
Biancae, Guss. ex Nym. Consp. 550.—Sicil.
Bocconi, Kotschy, in Unger & Kotschy, Cypern, 292 = *cypria*.
Bocconi, Walp. Rep. iii. 400.—Sicil.

ODONTITES :—
breviflora, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) 1. 105.—As. temp.
caucasica, C. Koch, in Linnaea, xvii. (1843) 289 = Euphrasia officinalis.
chrysantha, Bor. Fl. Centr. Fr. ed. III. ii. 491 = *Jaubertiana*.
ciliata, Pomel, Nouv. Mat. Fl. Atl. 302.—Alger.
Citardae, Todaro, ex Caruel, in Parl. Fl. Ital. vi. 459 = *rigidifolia*.
corsica, G. Don, Gen. Syst. iv. 611.—Corsica.
cretica, Boiss. Fl. Orient. iv. 477.—Ins. Creta.
cypria, Boiss. l. c.—Ins. Cyprus.
discolor, Pomel, Nouv. Mat. Fl. Atl. 301.—Alger.
divergens, Jord. in F. Schult. Arch. Fl. Fr. et Allem. 191 = Bartsia Odontitis.
Dukerleyi, Gren. & Paill. in Billotia, i. (1864) 82.—Alger.
Fradini, Pomel, Nouv. Mat. Fl. Atl. 303.—Alger.
glutinosa, Benth. in DC. Prod. x. 549 = Euphrasia glutinosa.
granatensis, Boiss. Elench. 71 = B. purpurea.
hispanica, Boiss. & Reut. Pugill. 91.—Hispan.
Holliana, Benth. in DC. Prod. x. 550 = B. scordifolia.
ixodes, Boiss. Diagn. Ser. I. vii. 55 = *glutinosa*.
Jaubertiana, D. Dietr. Syn. Pl. iii. 544.—Gallia.
Kochii, F. Schultz, in Flora, xxxvi. (1853) 557 = B. Odontites.
lanceolata, Reichb. Fl. Germ. Excurs. 862 = B. lutea.
linifolia, G. Don, Gen. Syst. iv. 611.—Europ.
Linkii, Heldr. & Sart. in Boiss. Diagn. Ser. II. iii. 177.—Graecia.
litoralis, Fries, Summa Veg. Scand. i. 19 = B. verna.
longiflora, Webb, Iter Hisp. 24 = Euphrasia longiflora.
lutea, [Clairv.] Man. Herb. 207 = B. lutea.
lutea, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4.—Rossia.
purpurea, G. Don, Gen. Syst. iv. 611 = B. purpurea.
Reboudii, Pomel, Nouv. Mat. Fl. Atl. 303.—Alger.
Recordoni, Burnat & Barbey, Voy. Bot. Baléar. 42. t. 1.—Hispan.
rigidifolia, Benth. in DC. Prod. x. 550 = Euphrasia rigidifolia.
rotundata, Ball, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 30 = B. verna.
rubra, Gilib. Fl. Lituan. i. 126 = B. Odontites.
rubra, Pers. Syn. ii. 150 = B. verna.
rubra, Willk. in Flora, xxxv. (1852) 264 = *virgata*.
santolinifolia, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 13 = B. santolinaefolia.
serotina, Dum. Fl. Belg. 32; Reichb. Fl. Germ. Excurs. 359 = B. Odontites.
serotina, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 617 = *virgata*.
sicula, Todaro, ex Nym. Consp. 550 = *Jaubertiana*.
simplex, Krok, ex Nym. l. c. 551 = B. verna.
tenuifolia, G. Don, Gen. Syst. iv. 611 = Euphrasia tenuifolia.
Triboutii, Gren. & Paill. in Billotia, i. (1864) 81.—Alger.
verna, Dum. Fl. Belg. 32; Reichb. Fl. Germ. Excurs. 359 = B. verna.
violacea, Pomel, Nouv. Mat. Fl. Atl. 301.—Alger.
virgata, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 617.—Hispan.
viscosa, [Clairv.] Man. Herb. 207; Reichb. Fl. Germ. Excurs. 360 = B. purpurea.
vulgaris, Moench, Meth. 439 = B. Odontites.
vulgaris, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4 = B. verna.

ODONTITES, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 258 = **Bupleurum**, Tourn. (Umbellif.).
glauca, Spreng. in Schult. Syst. vi. 383 = B. glaucum.
gracilis, Bieb. ex Hoffm. Gen. Umb. ed. II. 209 = B. Marschallianum.
luteola, Hoffm. l. c. = B. semicompositum.
luteola, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 258 = B. aristatum.
procumbens, Spreng. l. c. = B. tenuissimum.
semicomposita, Spreng. l. c. = B. semicompositum.
tenuissima, Spreng. l. c. = B. tenuissimum.

ODONTITIS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 88 = praec.

ODONTOCARPA, Neck. Elem. i. 123 (1790) = **Valerianella**, Moench.

ODONTOCARPHA, DC. Prod. v. 71 (1836) = **Gutierrezia**, Lag. (Compos.).
Gayana, Remy, in C. Gay, Fl. Chil. iv. 36 (= *Brachyris Gayana*).—Chili.
Poeppigii, DC. Prod. v. 72 = G. paniculata.

ODONTOCARYA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 38. *MENISPERMACEAE*, Benth. & Hook. f. i. 960.
CHONDODENDRON, Ruiz & Pav. Prod. 132 (1794).
acuparata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 100, nomen; *et Contrib.* iii. 61.—Bras.
convolvulacea, *Miers, ll. cc.* 100; 62.—Peruv.
filipendula, Miers, Contrib. iii. 65 = Cocculus filipendula.
hederaefolia, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 100; et Contrib. iii. 64 = tamoides.
Hernandia, Miers, l. c. Ser. II. vii. (1851) 38 = Cissampelos Hernandia.
macrostachya, Miers, l. c. Ser. III. xiv. (1864) 100 = O. Vitis.
sagittata, *Miers, l. c.* 100, nomen; *et Contrib.* iii. 64.—Guiana.
scabra, *Miers, ll. cc.* 100; 65.—Bras.
tamoides, *Miers, ll. cc.* 100; 63.—Bras.
Vitis, *Miers, l. c.* Ser. II. vii. (1851) 38.—Bras.

ODONTOCHILUS, Blume, Orch. Archip. Ind. 79. tt. 29, 36 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 600.
CYSTOPUS, Blume, Orch. Archip. Ind. 82 (1858).
flavescens, *Blume, Orch. Archip. Ind.* 80.—Java.
flavus, Benth. & Hook. f. Gen. iii. 598, sphalm. pro *luteo* = lanceolatus.
grandiflorus, *Benth. & Hook. f. l. c.*—Reg. Himal.
Jaubertii, *Blume, Orch. Archip. Ind.* 81.—Ins. Sandvic.
lanceolatus, *Blume, l. c.* 80.—Reg. Himal.
longiflorus, *Benth. & Hook. f. Gen.* iii. 598.—Ins. Vit.

ODONTOCYCLUS, Turcz. in Bull. Soc. Nat. Mosc. (1840) 65. *CRUCIFERAE*, Benth. & Hook. f. i. 75.
kurilensis, *Turcz. l. c.*—Ins. Kurile.

ODONTOGLOSSUM, H. B. & K. Nov. Gen. et Sp. i. 350. t. 85 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 561.
CUITLAUZINA, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 32 (1825).
LICHTERVELDIA, Lem. Illustr. Hortic. ii. (1855) t. 59.
MESOSPINIDIUM, Reichb. f. in Bot. Zeit. x. (1852) 929.
acuminatissimum ×, Reichb. f. in Gard. Chron. (1882) i. 256.—Hab.?
acuminatum, Hort. ex Lindl. Fol. Orch. Odontog. 9 = apterum.
Alexandrae, Batem. in Gard. Chron. (1864) 1083; et in Proc. Roy. Hort. Soc. iv. (1864) 186 = crispum.
Amphiglottii, *Reichb. f. in Linnaea*, xli. (1877) 2.—N. Granat.
anceps, Klotzsch, in Otto & Dietr. Allg. Gartenz. xix. (1851) 250 = Miltonia anceps.
anceps, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 45.—Mexic.
Andersonianum ×, Reichb. f. in Gard. Chron. (1868) 599.—N. Granat.
angustatum, *Lindl. Bot. Reg.* sub t. 1992.—N. Granat.
angustatum, Lindl. Orch. Linden. 17 = ramosissimum.
anthoxanthum, *Reichb. f. in Gard. Chron.* (1869) 388.—Peruv.
apterum, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*). 35.—Mexic.
armatum, *Reichb. f. in Linnaea*, xli. (1877) 32.—Ecuador.
Arminii, *Reichb. f. in Bonplandia*, iii. (1855) 66.—Venezuela.
Aspasia, Reichb. f. in Walp. Ann. vi. 851 = Aspasia epidendroides.
aspersum ×, Reichb. f. in Gard. Chron. (1879) i. 266.
astranthum, *Linden & Reichb. f. l. c.* (1867) 404.—Reg. Andin.
atropurpureum, *Reichb. f. ex Beer, Prakt. Orch.* 274.—Hab.?

ODONTOGLOSSUM :—
auropurpureum, *Reichb. f. in Linnaea*, xxii. (1849) 848.—Peruv.
axinopterum, *Reichb. f. l. c.* xli. (1877) 68.—Ecuador.
baphicanthum ×, Reichb. f. in Gard. Chron. (1876) ii. 260.—N. Granat.
Barkeri, Reichb. f. in Walp. Ann. vi. 854 = Gomeza Barkeri.
beloglossum, *Reichb. f. Xenia Orch.* ii. 149. t. 158.—Mexic.
Bergamii, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 596, nomen.—Mexic.
Biberianum, Reichb. f. in Walp. Ann. vi. 851 = Aspasia Biberiana.
bicolor, Lindl. Bot. Reg. (1845) Misc. 59 = Oncidium aureum.
bictoniense, *Lindl. Bot. Reg.* (1840) t. 66.—Mexic.
blandum, *Reichb. f. in Gard. Chron.* (1870) 1342.—Hab.?
Bluntii, *Reichb. f. in Bot. Zeit.* xxii. (1864) 415.—N. Granat.
brachypterum ×, Reichb. f. in Linnaea, xli. (1877) 26.—Peruv.
Brassia ×, Reichb. f. in Gard. Chron. (1885) i. 501.
brevifolium, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
caerulescens, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 27.—Mexic.
calodryas, *Reichb. f. in Linnaea*, xli. (1877) 3.—N. Granat.
candidum, *Linden & André, in Illustr. Hortic.* xxii. (1875) 58.—N. Granat.
cariniferum, *Reichb. f. in Bot. Zeit.* x. (1852) 638.—Am. centr.
Cervantesii, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 34.—Mexic.
chaetostroma ×, Reichb. f. in Gard. Chron. (1883) i. 592.
cheetostroma ×, Reichb. f. l. c. 562 = praec.
Chestertoni, *Hort.—Cf. Gard. Chron.* (1876) i. 374.—Hab.?
chiriquense, *Reichb. f. in Bot. Zeit.* x. (1852) 692.—Panama.
cimiciferum, Reichb. f. in Linnaea, xxii. (1849) 849 = Oncidium cimiciferum.
cinnamomeum, Hort. ex H. Williams, in Orchid. Alb. iii. (1884) sub t. 143 = Schillerianum.
cirrhosum, *Lindl. Gen. et Sp. Orch.* 211.—Peruv.
citrosmum, *Lindl. Bot. Reg.* (1842) *Misc.* 60.—Mexic.
claviceps, *Reichb. f. in Gard. Chron.* (1876) ii. 516.—Ecuador.
Clowesii, Lindl. Bot. Reg. (1839) Misc. 93 = Miltonia Clowesii.
Cochlioda, Reichb. f. in Walp. Ann. vi. 853 = Cochlioda densiflora.
coerulescens, Auct. vide caerulescens.
compactum, *Reichb. f. in Gard. Chron.* (1875) i. 492.—Peruv.
confertum, *Reichb. f. l. c.* (1879) i. 298.—Ecuador.
connivens, *Scheidw. in Journ. Hort. Prat.* iii. (1846) 148.—Mexic.
constrictum, *Lindl. Bot. Reg.* (1843) *Misc.* 17.—Venezuela.
Coradinei ×, Reichb. f. in Gard. Chron. (1872) 1068.—N. Granat.
cordatum, *Lindl. Bot. Reg.* (1838) *Misc.* 50.—Mexic.
coronarium, *Lindl. Fol. Orch. Odontog.* 21.—N. Granat.
costatum, *Lindl. l. c.* 11.—Venezuela.
crinitum, *Reichb. f. Xenia Orch.* ii. 207.—N. Granat.
crispatulum, Reichb. f. in Walp. Ann. vi. 853 = Gomesa crispa.
crispum, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 256.—N. Granat.
cristatellum, *Reichb. f. in Gard. Chron.* (1878) ii. 716.—Hab.?
cristatum, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
crocatum, *Linden & Reichb. f. l. c.* (1867) 404.—Hab.?
crocidipterum, *Reichb. f. l. c.* (1871) 1129.—N. Granat.
cruentum, *Reichb. f. Xenia Orch.* ii. 174.—N. Granat.
cuspidatum, *Reichb. f. l. c.* 174.—N. Granat.
Dawsonianum, *Reichb. f. in Gard. Chron.* (1865) 1226.—Mexic.
deltoglossum ×, Reichb. f. l. c. (1881) i. 202.
Denisonae ×, Hort. ex Flor. Mag. N. S. (1872) t. 26.—N. Granat.

ODONTOGLOSSUM:—
densiflorum, *Lindl. Orch. Linden.* 17.—Mexic.
dipterum, *Lindl. l. c.* 16.—N. Granat.
distans, *Reichb. f. in Linnaea*, xxii. (1849) 848.—Venezuela.
divaricatum, Lindl. Orch. Linden. 17 = Oncidium refractum.
Dormannianum, *Reichb. f. in Gard. Chron.* (1884) i. 11.—N. Granat.
Edwardi, *Reichb. f. l. c.* (1878) ii. 74.—Ecuador.
Egertoni, *Lindl. Bot. Reg.* (1845) *Misc.* 50.—Guatemala.
Ehrenbergii, Hort. Angl. ex Reichb. f. in Gard. Chron. (1879) i. 138 = Dawsonianum.
Ehrenbergii, *Link, Klotzsch & Otto, Ic. Pl. Hort. Berol.* 38. t. 16.—Mexic.
Ehrenbergii, Planch. in Fl. des Serres, viii. (1852–53) t. 846 = apterum.
elegans ×, Reichb. f. in Gard. Chron. (1879) i. 462.—Ecuador.
epidendroides, *H. B. & K. Nov. Gen. et Sp.* i. 351.—Peruv.
epidendroides, Lindl. Fol. Orch. Odontog. 2 = Lindleyanum.
erosum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 27; Reichb. f. & Warsz. in Bonplandia, ii. (1854) 99 = stellatum.
eugenes ×, Hort.—Cf. Gard. Chron. (1883) ii. 120.
excellens ×, Reichb. f. l. c. (1881) ii. 426.—N. Granat.
facetum ×, Reichb. f. l. c. i. 563.
ferrugineum ×, Reichb. f. l. c. (1883) i. 814.
festatum, Reichb. f. in Bonplandia, ii. (1854) 100; iv. (1856) 212 = Oncidium aureum.
foliosum, Reichb. f. in Walp. Ann. vi. 854 = Gomesa foliosa.
forcipatum, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 279.—N. Granat.
fractum, *Reichb. f. in Linnaea*, xli. (1877) 26.—N. Granat.
Galeottianum, *A. Rich. in Ann. Sc. Nat.* Sér. III. iii. (1845) 27.—Mexic.
Ghiesbreghtianum, *A. Rich. l. c.*—Mexic.
gloriosum, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.
gracile, *Lindl. in Benth. Pl Hartw.* 151.—Peruv.
grande, *Lindl. Bot. Reg.* (1840) *Misc.* 47.—Guatemala.
grande-pallidum, Klotzsch, ex Reichb. f. in Gard. Chron. (1865) 1082 = Schlieperianum.
Hallii, *Lindl. Bot. Reg.* sub t. 1992.—Ecuador.
hastatum, Batem. Orch. Mex. & Guat. t. 20 = Oncidium hastatum.
hastilabium, *Lindl. Orch. Linden.* 16; *et in Bot. Mag.* t. 4272.—N. Granat.
hebraicum ×, Reichb. f. in Gard. Chron. (1879) i. 462.
hemichrysum, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 100 = Oncidium aureum.
Hinnus ×, Reichb. f. Xenia Orch. ii. 153.—Ecuador.
histrionicum ×, Reichb. f. in Gard. Chron. (1882) i. 178.
Hookeri, Lem. Illustr. Hortic. iii. (1856) Misc. 41 = cordatum.
Horsmanni, *Reichb. f. in Gard. Chron.* (1880) i. 41.—N. Granat.
Humeanum ×, Reichb. f. l. c. (1876) i. 170.
Insleayi, *Lindl. Fol. Orch. Odontog.* 4.—Mexic.
ioplocon, *Reichb. f. in Gard. Chron.* (1884) i. 445.—Hab.?
ixioides, *Lindl. Bot. Reg.* (1845) *Misc.* 55.—N. Granat.
jamaicense, *Griseb. Fl. Brit. W. Ind.* 633.—Ind. occ.
Jenningsianum ×, Reichb. f. in Gard. Chron. (1878) i. 366.
Karwinskii, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Mexic.
Kegeljani, *E. Morr. in Belg. Hortic.* xxvii. (1877) 212. t. 10.—Hab.?
Krameri, *Reichb. f. in Gard. Chron.* (1868) 98.—Costa Rica.
lacerum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.
laeve, *Lindl. Bot. Reg.* (1844) t. 39.—Guatemala.
laxiflorum, Reichb. f. in Walp. Ann. vi. 853 = Gomesa chrysostoma.
Leeanum ×, Reichb. f. in Gard. Chron. (1882) i. 525.
Lehmanni, *Reichb. f. Otia Bot. Hamb.* 4.—Ecuador.
lepidum, *Reichb. f. in Gard. Chron.* (1883) ii. 526.—N. Granat.
leucomelas, *Reichb. f. in Bot. Zeit.* xxii. (1864) 415.—Mexic.
leucopterum, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 279.—N. Granat.
ligulare ×, Reichb. f. in Gard. Chron. (1882) i. 558.
limbatum, *Reichb. f. l. c.* (1870) 417.—Hab.?
Lindenii, *Lindl. Fol. Orch. Odontog.* 16.—N. Granat.
Lindleyanum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 99.—N. Granat.
Lindleyi, Galeotti, ex Lindl. Fol. Orch. Oncid. 33 = Oncidium maculatum.
Londesboroughianum, *Reichb. f. in Gard. Chron.* (1876) ii. 772.—Mexic.
longifolium, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
longipes, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 100.—Peruv.
Lueddemanni, Regel, Gartenfl. (1859) 353. t. 275 = cordatum.
lunatum, Reichb. f. in Walp. Ann. vi. 849 = Aspasia lunata.
luteo-purpureum, *Lindl. Orch. Linden.* 16.—N. Granat.
lyratum, Reichb. f. in Walp. Ann. vi. 852 = Aspasia lyrata.
lyroglossum ×, Reichb. f. in Flora, lxv. (1882) 534.
macrospilum ×, Reichb. f. in Gard. Chron. (1885) ii. 70.
macrum, *Lindl. Fol. Orch. Odontog.* 12.—Am. centr.
maculatum, Hook. Bot. Mag. t. 4878 = cordatum.
maculatum, *La Llave, ex Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 35.—Mexic.
madrense, *Reichb. f. in Gard. Chron.* (1874) ii. 804.—Mexic.
marginellum, *Reichb. f. l. c.* (1882) ii. 680.—Hab.?
Marriottianum ×, Reichb. f. l. c. (1881) i. 168.
maxillare, Hook. f. Bot. Mag. t. 6144 = madrense.
maxillare, *Lindl. Bot. Reg.* (1847) sub t. 62.—Mexic.
Measuresianum, *O'Brien, in Gard. Chron.* (1885) ii. 619.—Hab.?
megalophium, *Lindl. Orch. Linden.* 15.—Peruv.
melanthes, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 100.—Peruv.
membranaceum, Lindl. Sert. Orch. sub t. 25 = Cervantesii.
mentigerum, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 100 = Oncidium mentigerum.
mirandum, *Reichb. f. in Gard. Chron.* (1882) i. 143.—N. Granat.
multistellare, *Reichb. f. in Linnaea*, xli. (1877) 25.—Peruv.
Mulus ×, Reichb. f. Xenia Orch. ii. 151.—Hab.?
myanthum, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
myrianthum, *Reichb. f. Xenia Orch.* i. 189; *et in Bonplandia*, iv. (1856) 212.—Peruv.
mystacinum, *Lindl. Fol. Orch. Odontog.* 6.—Peruv.
Murrellianum ×, Reichb. f. in Gard. Chron. (1875) i. 653.—N. Granat.
naevium, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 87.—N. Granat.
nebulosum, *Lindl. Sert. Orch.* sub t. 25.—Mexic.
nevadense, *Reichb. f. Illustr. Hortic.* xvii. (1870) 243. t. 45.—N. Granat.
nobile, *Reichb. f. in Linnaea*, xxii. (1849) 850.—N. Granat.
odoratum, *Lindl. Orch. Linden.* 16.—N. Granat.
Oerstedii, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Costa Rica.
oliganthum, *Reichb. f. l. c.* iv. (1856) 321.—Guatemala.
orientale, *Reichb. f. in Gard. Chron.* (1879) i. 366.—Ecuador.
pardinum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.; N. Granat.
parviflorum, *Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 596, nomen.—Mexic.
pendulum, Batem. Monog. Odont. t. 6 = citrosmum.
Pescatorei, Linden, ex Lindl. & Paxt. Flow. Gard. iii. (1852–53) 83. t. 90 = nobile.
Phalaenopsis, Linden & Reichb. f. in Bonplandia, ii. (1854) 278 = Miltonia Phalaenopsis.
phyllochilum, C. Morr. in Ann. Soc. Bot. Gand, v. (1849) 295. t. 271 = Oncidium hastatum.
planifolium, Reichb. f. in Walp. Ann. vi. 853 = Gomesa planiflora.
platyodon, *Reichb. f. in Gard. Chron.* (1871) 1163.—N. Granat.
platyotum, Hort. ex Reichb. f. l. c. = praec.
Pollettianum ×, O'Brien, l. c. (1885) ii. 680.
polyxanthum, Hort.—Cf. Gard. Chron. (1881) ii. 460 = Kegeljani.
praenitens, *Reichb. f. in Gard. Chron.* (1875) i. 524.—N. Granat.
praestans, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 99.—N. Granat.
prasinum, *Linden & Reichb. f. in Gard. Chron.* (1870) 987.—Ecuador.
principissa, Reichb. f. in Walp. Ann. vi. 852 = Aspasia principissa.
prionopetalon ×, O'Brien, in Gard. Chron. (1885) ii. 620.
psittacinum, Reichb. f. in Linnaea, xli. (1877) 102 = Aspasia psittacina.
pulchellum, *Batem. ex Lindl. Bot. Reg.* (1841) t. 48.—Guatemala.
purum, *Reichb. f. in Gard. Chron.* (1872) 1328.—N. Granat.
pygmaeum, *Lindl. in Benth. Pl. Hartw.* 82.—Guatemala.
radiatum, *Reichb. f. in Gard. Chron.* (1865) 746.—N. Granat.
ramosissimum, *Lindl. Fol. Orch. Odontog.* 16.—Venezuela.
ramulosum, *Lindl. l. c.* 13.—N. Granat.
recurvum, Reichb. f. in Walp. Ann. vi. 853 = Gomesa recurva.
Reichenheimii, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 214.—Mexic.
retusum, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
revolutum, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 256.—N. Granat.
rigidum, *Lindl. in Benth. Pl. Hartw.* 152.—Peruv.
ringens, *Reichb. f. in Saund. Refug. Bot.* t. 99; *Xenia Orch.* ii. 150.—Peruv.
robustum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 100.—N. Granat.
Roezlii, Reichb. f. Xenia Orch. ii. 191 = Miltonia Roezlii.
Rolliae, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 596, nomen.—Guatemala.
roseum, Lindl. in Benth. Pl. Hartw. 151 = Cochlioda rosea (in Addend.).
Rossii, *Lindl. Sert. Orch.* sub t. 25; *Knowles & Westc. Fl. Cab.* ii. (1838) 186.—Mexic.
rubescens, *Lindl. in Journ. Hort. Soc.* v. (1850) 35.—Nicaragua.
Rueckerianum ×, Reichb. f. in Gard. Chron. (1873) 105.—Hab.?
Sanderianum, *Reichb. f. l. c.* (1881) ii. 524.—Am. trop.
Sceptrum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 99.—N. Granat.
Schillerianum, *Reichb. f. l. c.* 12.—Venezuela.
Schlieperianum, *Reichb. f. in Gard. Chron.* (1865) 1082.—Costa Rica.
Schlimii, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.
Schroederianum ×, Reichb. f. in Gard. Chron. (1882) i. 700.
Scottii, Hort.—Cf. l. c. (1883) i. 222.
Shuttleworthii ×, Hort.—Cf. l. c. (1884) i. 490.
spathaceum, *Lindl. Fol. Orch. Odontog.* 18.—N. Granat.
spectatissimum, *Lindl. l. c.* 19.—N. Granat.
spilotanthum, *Linden & Reichb. f. Xenia Orch.* ii. 175.—N. Granat.
stellatum, *Lindl. Bot. Reg.* (1841) *Misc.* 9.—Am. centr.
stellimicans ×, Reichb. f. in Gard. Chron. (1884) ii. 680.
stenochilum, *Linden & Reichb. f. l. c.* (1872) 969.—Hab.?
subuligerum, *Reichb. f. in Linnaea*, xli. (1877) 27.—Peruv.
tentaculatum ×, Reichb. f. in Gard. Chron. (1883) i. 814.
tetraplasium, *Reichb. f. l. c.* (1875) i. 558.—Peruv.
tigrinum, Lindl. Fol. Orch. Odontog. 5 = Oncidium tigrinum.
tripudians, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 100.—Peruv.
triumphans, *Reichb. f. l. c.* 99.—N. Granat.
ulopterum, *Linden & Reichb. f. in Gard. Chron.* (1872) 731.—N. Granat. *vel* Ecuador.
umbrosum, Reichb. f. in Linnaea, xxii. (1849) 848 = Oncidium linguiforme.
Uroskinneri, *Lindl. in Gard. Chron.* (1859) 708.—Guatemala.
variegatum, Reichb. f. in Walp. Ann. vi. 851 = Aspasia variegata.

ODONTOGLOSSUM:—
velleum, *Reichb. f. in Gard. Chron.* (1874) I. 406.—Ecuador.
vexativum, Reichb. f. l. c. (1876) II. 808.—Mexic.
vexillarium, Reichb. f. l. c. 901 = Miltonia vexillaria.
Victor, *Reichb. f. l. c.* (1883) I. 532.—Hab. ?
viminale, *Reichb. f. l. c.* (1885) I. 108.—N. Granat.
Vuylstekeanum ×, Reichb. f. l. c. (1884) II. 7.
Wageneri, *Reichb. f. in Bonplandia*, ii. (1854) 12.—Venezuela.
Wallisii, *Linden & Reichb. f. in Gard. Chron.* (1870) 104.—N. Granat.
Warneri, Lindl. Bot. Reg. (1845) Misc. 54 = Oncidium Warneri.
Warnerianum, *Reichb. f. in Bot. Zeit.* xxii. (1864) 297.—Mexic.
Warscewiczianum, Hemsl. Biol. Centr. Am. Bot. iii. 277, sphalm. = seq.
Warszewiczii, Reichb. f. in Bot. Zeit. x. (1852) 692 = Miltonia Warszewiczii.
Weirii, *Reichb. f. in Gard. Chron.* (1875) I. 461.—N. Granat.
Weltoni, Hort.—Cf. Gard. Chron. (1870) 417 = Miltonia Warszewiczii.
Wilckeanum ×, Reichb. f. in Gard. Chron. (1880) I. 298.
Williamsianum ×, Reichb. f. l. c. (1881) II. 134.
zebrinum, Reichb. f. in Linnaea, xxii. (1849) 849 = Oncidium zebrinum.

ODONTOLOMA, H. B. & K. Nov. Gen. et Sp. iv. 43. t. 319 (1820) = **Oliganthes**, Cass. (Compos.).
acuminata, H. B. & K. l. c. 44.—Venezuela.

ODONTOLOPHUS, Cass. in Dict. Sc. Nat. l. 252 (1827) = **Centaurea**, Linn. (Compos.).
cyanoides, Cass. l. c. 253 = C. trinervia.

ODONTONEMA, Nees, in Linnaea, xvi. (1842) 300 = **Thyrsacanthus**, Nees (Acanth.).
lucidum, Nees, l. c. = T. bracteolatus.

ODONTOPTERA, Cass. in Dict. Sc. Nat. xxxv. 396 (1825) = **Arctotis**, Linn. (Compos.).
sulphurea, Cass. l. c. = A. aspera ?

ODONTOSIPHON, M. Roem. Syn. Hesper. 85, 106 (1846) = **Trichilia**, Linn. (Meliac.).
elegans, M. Roem. l. c. 106 = T. elegans.
Richardianum, M. Roem. l. c. = T. Richardiana.

ODONTOSPERMUM, Neck. Elem. i. 20 (1790). *COMPOSITAE*, Benth. & Hook. f. ii. 340.
ASTERISCUS, Moench, Meth. 592 (1794).
NAUPLIUS, Cass. in Dict. Sc. Nat. xxiii. 566 (1822); xxxiv. 272 (1825).
SAULCYA, Michon, Voy. Reliq. Or. ii. 383 (1854).
aquaticum, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 232, 233.—Reg. Mediterr.
Daltoni, *Webb, in Hook. Niger Fl.* 140.—Ins. Cap. Virid.
graveolens, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 232.—Afr. trop.; Ins. Canar.
imbricatum, *Sch. Bip. l. c.*—Marocc.
intermedium, *Sch. Bip. l. c.* 232, 236.—Ins. Canar.
lanuginosum, *Sch. Bip. l. c.* 232.—Syria.
maritimum, *Sch. Bip. l. c.* 233.—Reg. Mediterr.; Ins. Canar.
odorum, *Sch. Bip. l. c.* 238.—Marocc.
Schultzii, *C. Bolle, in Bonplandia*, vii. (1859) 295.—Ins. Canar.
sericeum, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 234.—Ins. Canar.
Smithii, *Webb, in Hook. Niger Fl.* 139.—Ins. Cap. Virid.
stenophyllum, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 237.—Ins. Canar.
Vogelii, *Webb, in Hook. Niger Fl.* 140.—Ins. Cap. Virid.

ODONTOSTEMMA, Benth. in Wall. Cat. n. 645 (1828); et ex G. Don, Gen. Syst. i. 449 (1831) = **Arenaria**, Rupp. (Caryoph.).
glandulosum, Benth. ll. cc. = A. Benthamii.

ODONTOSTEMUM, Baker, in Journ. Linn. Soc. xi. (1871) 436 = **Odontostomum**, Torr. (Haemod.).

ODONTOSTIGMA, A. Rich. Fl. Cub. Fanerog. ii. 86 (1853) = **Stemmaderia**, Benth. (Apocyn.).
Galeottianum, A. Rich. l. c. 87 = S. Galeottianum.

ODONTOSTIGMA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 573 = **Gymnostachyum**, Nees (Acanth.).
subacaule, Zoll. & Mor. l. c. = G. Cumingianum.

ODONTOSTOMUM, Torr. in Pacif. Rail. Rep. iv. 150 (1856). *HAEMODORACEAE*, Benth. & Hook. f. iii. 680.
Hartwegii, *Torr. in Pacif. Rail. Rep.* iv. 150.—Calif.

ODONTOSTYLES, Breda, Kuhl & Hasselt, Ge . et Sp. Orch. t. [4] (1827) = **Bulbophyllum**, Thou. (Orchid.).
minor, Breda, l. c. [sub t. 4.] = *multiflora*.
multiflora, Breda, l. c. [t. 4].—Java.
triflora, Breda, l. c.—Java.

ODONTOTRICHUM, Zucc. in Abh. Akad. Muench. i. (1832) 311 = **Senecio**, Tourn. (Compos.).
cirsiifolium, Zucc. l. c. = S. cervariaefolius.

ODOPTERA, Rafin. Cat. 15 (1824) = **Corydalis**, Vent. (Papav.).
aurea, Rafin. l. c. = C. aurea.

ODOSTELMA, Rafin. Fl. Tellur. iv. 104 (1836) = **Tacsonia**, Juss. (Passifl.).
adulterina, Rafin. l. c. = T. adulterina.
pedunculata, Rafin. l. c. = T. peduncularis.

ODOSTEMON, Rafin. in Am. Monthly Mag. (1817) 192; et in Journ. Phys. lxxxix. (1819) 259 = **Berberis**, Linn.

ODOSTEMUM, Steud. Nom. ed. II. ii. 204 (1841) = praec.

ODOTALON, Rafin. Sylva Tellur. 66 (1838) = **Argithamnia**, Sw. (Euphorb.).
lanceolata, Rafin. l. c. 67 = A. tricuspidata.
tricuspidata, Rafin. l. c. 66 = A. tricuspidata.

OECEOCLADES, Lindl. Bot. Reg. sub t. 1522 (1832); et Gen. et Sp. Orch. 235 (1833) = **Saccolabium**, Blume (Orchid.).
falcata, Lindl. Gen. et Sp. Orch. 237 = Angraecum falcatum.
flexuosa, Lindl. in Wall. Cat. n. 7333 (Cleisostomatis sp.).—Reg. Himal.; Burma.
funalis, Lindl. Gen. et Sp. Orch. 237 = Dendrophylax funale.
gracilis, Lindl. l. c. = Angraecum gracile.
javanica, Teijsm. & Binn. in Tijdschr. Nederl. Indie, xxiv. (1862) 326.—Java.
Lindleyana, Regel, Ind. Sem. Hort. Petrop. (1865) 43 = seq.
Lindleyi, Regel, Gartenfl. (1866) 70 = Angraecum falcatum.
maculata, Lindl. Gen. et Sp. Orch. 237 = Eulophia monophylla.
paniculata, Lindl. l. c. 236 = Saccolabium buccosum.
parviflora, Lindl. l. c. = Angraecum parviflorum.
pusilla, Lindl. in Wall. Cat. n. 7332 = Saccolabium Pumilio.
Retzii, Lindl. Gen. et Sp. Orch. 237 (Sarcochili sp.).—Ind. or.
tenera, Lindl. l. c. 236 (Cleisostomatis sp.).—Ind. or.

OECHMEA, Jaume St. Hil. Expos. i. 103 (1805) = **Aechmea**, Ruiz & Pav. (Bromel.).

OECOECLADES, Franch. & Sav. Enum. Pl. Jap. ii. 28 (1879), sphalm. = **Oeceoclades**, Lindl. (Orchid.).

OEDEMATOPUS, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 249 = **Havetiopsis**, Planch. (Guttif.).
dodecandrus, Planch. & Triana, l. c. 251.—Bras.
obovatus, Planch. & Triana, l. c. 250.—Bras.
octandrus, Planch. & Triana, l. c.—Peruv.

OEDERA, Crantz, Duab. Dracon. Arb. 30 (1768) = **Dracaena**, Linn. (Liliac.).
dragonalis, Crantz, l. c. = D. Draco.

OEDERA, Linn. Mant. ii. 159 (1767). *COMPOSITAE*, Benth. & Hook. f. ii. 418.
OEDERIA, DC. Prod. vi. 1 (1837).
aliena, Linn. f. Suppl. 390 = Heterolepis decipiens.
alienata, Thunb. Prod. Pl. Cap. 169 = Hirpicium echinulatum.
apiculata, E. Mey. ex DC. Prod. vi. 289 = Relhania quinquenervis.
hirta, Thunb. Prod. Pl. Cap. 169.—Afr. austr.
imbricata, Lam. Encyc. ii. 345 = latifolia.
intermedia, DC. Prod. vi. 2 = latifolia.
laevis, *DC. l. c.*—Afr. austr.
lanceolata, Cass. in Dict. Sc. Nat. xxxv. 402 = prolifera.
latifolia, *Less. Syn. Comp.* 247.—Afr. austr.
obtusifolia, Cass. in Dict. Sc. Nat. xxxv. 402 = latifolia.
prolifera, *Linn. Mant.* ii. 159.—Afr. austr.
speciosa, E. Mey. ex DC. Prod. vi. 289 = Relhania speciosa.
trinervia, [*Spreng.*] *Bot. Gart. Halle*, 63.—Hab. ?

OEDERIA, DC. Prod. vi. 1 (1837) = praec.

OEDIPACHNE, Link, Hort. Berol. i. 51 (1827) = **Eriochloa**, H. B. & K. (Gramin.).
punctata, Link, l. c. = E. polystachya.

OEDMANNIA, Thunb. Nov. Gen. 136 (1800) = **Rafnia**, Thunb. (Legumin.).
lancea, Thunb. l. c. 137 = R. lancea.
lanceolata, Steud. Nom. ed. II. ii. 205 = R. lancea.

OEGINETIA, Wight, Ic. iii. textu n. 895 [1843] = **Aeginetia**, Linn. (Orobanch.).

OENANTHE, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 905.
CYSSOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxii (1849) II. 25.
DASYLOMA, DC. Prod. iv. 140 (1830).
PHELLANDRIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 77 (1737).
ambigua, Nutt. Gen. Am. i. 189 = Peucedanum ternatum.
amoena, *Lehm. ex Sweet, Hort. Brit.* ed. II. 247, nomen.—Hab. ?
anatolica, C. Koch, in Linnaea, xix. (1847) 38 = angulosa.
angulosa, *Griseb. Spicil. Fl. Rumel.* i. 352.—Macedon.; As. Min.; Syria.
anomala, *Dur. & Coss. Notes Crit.* 133.—Alger.
apiifolia, Brot. Fl. Lusit. i. 420 = crocata.
approximata, Mérat, Nouv. Fl. Par. 115 = Lachenalii.
approximata, Tenore, Fl. Nap. Prod. App. v. 10 = Jordani.
aquatica, Poir. in Lam. Encyc. iv. 530 = O. Phellandrium.
australiana, F. Muell. Fragm. v. 182 = stolonifera.
australis, Wulf. ex Steud. Nom. ed. II. ii. 205; et ej. Fl. Nor. Phan. 360 = silaifolia.
banatica, *Heuff. in Flora*, xxxvii. (1854) 291.—Ins. Banat.
benghalensis, *Benth. & Hook. f. Gen.* i. 906.—Ind. or.
biloba, Dulac, Fl. Hautes-Pyr. 353 = peucedanifolia.
californica, *S. Wats. in Proc. Am. Acad.* xi. (1876) 139.—Calif.
callosa, *Salzm. ex DC. Prod.* iv. 137.—Marocc.
calycina, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 473 = globulosa.
capensis, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza capensis.
capensis, Houtt. Handleid. viii. 140 = Lichtensteinia pyrethrifolia.
carolinensis, Pers. Syn. i. 318 = Peucedanum teretifolium.
carviformis, Schur, Enum. Pl. Transs. 255 = banatica.
chaerophyllaea, Gouan, Hort. Monsp. 144 = Seseli tortuosum ?
chaerophylloides, Pourr. in Mém. Acad. Toul. iii. (1788) 323 = pimpinelloides.
corticata, Edgew. in Trans. Linn. Soc. xx. (1846) 53 = stolonifera.
crocata, Kit. in Rochel, Pl. Banat. Rar. 27, nomen = banatica *vel* media.
crocata, *Linn. Sp. Pl.* 254.—Europ.
dacica, Kow. ex Schur, Enum. Pl. Transs. 255 = banatica.

OENANTHE:—
daucifolia, Link, ex Schult. Syst. vi. 423 = virgata.
densa, *J. Woods, Tour. Fl.* 147.—Europ. austr.
diffusa, Lag. Gen. et Sp. Nov. 13 = globulosa.
digitata, Nutt. ex DC. Prod. iv. 141 = Cynosciadium digitatum.
diversifolia, Dulac, Fl. Hautes-Pyr. 352 = pimpinelloides.
Dregeana, E. Mey. in Drège, Zwei Pfl. Docum. 205 = filiformis.
dubia, Roth, ex Schult. Syst. vi. 422 = peucedanifolia.
elata, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza elata.
exaltata, Thunb. Prod. Pl. Cap. 50 = Rhyticarpus difformis.
ferulacea, *Kotschy & Boiss. in Fl. Orient.* ii. 958.—Kurdistan.
ferulacea, Thunb. Prod. Pl. Cap. 50 = Peucedanum ferulaceum.
filicaulis, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza filicaulis.
filiformis, *Poir. in Lam. Encyc.* iv. 529.—Afr. austr.
filiformis, Walt. Fl. Carol. 113 = Peucedanum teretifolium.
filipendula, *Dum. Fl. Belg. Prod.* 80.—Europ.
filipenduloides, Thuill. Fl. Par. ed. II. 146 = peucedanifolia.
fistulifolia, Stokes, Bot. Mat. Med. ii. 111 = fistulosa.
fistulosa, Asso, Syn. Stirp. Arag. 36 = peucedanifolia.
fistulosa, *Linn. Sp. Pl.* 254.—Europ.; Reg. Casp.
fluviatilis, Colem. in Ann. & Mag. Nat. Hist. Ser. I. xiii. (1844) 188 = O. Phellandrium.
gigantea, *Zumag. Fl. Pedem.* i. 421.—Ital.
glauca, DC. ex Schult. Syst. vi. 428; et ex DC. Prod. iv. 137 = silaifolia.
globosa, Georgi, Geogr. Beschr. Russ. Reich. III. iv. 852 = pimpinelloides.
globosa, Hill, Veg. Syst. vi. 44 = praec.?
globularis, Link, Diss. Bot. Suerin (1795) 55 = globulosa.
globulosa, Bové, ex Dur. & Coss. Notes Crit. 135 = peucedanifolia.
globulosa, *Linn. Sp. Pl.* 255.—Europ. austr.; Afr. bor.
globulosa, Pall. Tabl. Taur. 49 = pimpinelloides.
graminifolia, *Gaud. Fl. Helv.* ii. 359, in obs.—Europ.
gymnorrhiza, Brign. Fasc. Pl. Forojul. 21 = Lachenalii.
Heldreichii, *Boiss. Fl. Orient.* ii. 957.—As. Min.
hirsuta, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza hirsuta.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 697.—Reg. Himal.
Huanaca, Spreng. Umbell. Prod. 37 = Huanaca Cavanillesii.
humilis, *Rafin. Préc. Découv.* 41.—Sicil.
incrassans, Bory & Chaub. Nouv. Fl. Pélop. t. 9 = incrassata.
incrassans, Heldr. ex Nym. Consp. 298 = media.
incrassata, *Bory & Chaub. Nouv. Fl. Pélop.* 19.—Graecia.
inebrians, Thunb. Prod. Pl. Cap. 49 (= *Glia gummifera*).—Afr. austr.
interrupta, Thunb. l. c. 50 = Lichtensteinia pyrethrifolia.
involucrata, Cav. ex Schult. Syst. vi. 421 = pimpinelloides.
javanica, DC. Prod. iv. 138 = stolonifera.
Jordani, *Tenore, Fl. Nap.* iii. 316.—Ital.
Karsthia, Hacq. Pl. Alp. Carniol. 17. t. 3 = silaifolia.
Kunzei, Willk. in Flora, xxxiv. (1851) 725 = globulosa.
Lachenalii, *C. C. Gmel. Fl. Bad.* i. 678.—Europ.; Reg. Casp.
laciniata, Zoll. Syst. Verz. Ind. Archip. 139 = stolonifera.
lanceolata, Poir. Encyc. Suppl. iv. 135 = fistulosa.
linearis, *Wall. Cat.* n. 586; *et ex DC. Prod.* iv. 138.—Reg. Himal.
Lisaea, Moris, in Mem. Acc. Torino, xxxviii. (1835) p. xxvii = pimpinelloides.
macrocarpa, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza macrocarpa.
macrosciadia, Willk. in Linnaea, xxv. (1852) 28 = crocata.
marginata, *Vis. Fl. Dalm.* iii. 38.—Hispan.
media, Bor. ex Nym. Consp. 298 = peucedanifolia.
media, *Griseb. Spicil. Fl. Rumel.* i. 352.—Europ. austr. or.; Syria.

OENANTHE:—
megapolitana, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 297 = Lachenalii.
meifolia, Schloss. & Vukot. Bilinar, 215 = fistulosa.
meoides, *Panč. Elem. Fl. Bulg.* 32.—Bulgaria.
michelfeldensis, Lachen. ex Mert. & Koch, Deutschl. Fl. ii. 416 = Lachenalii.
microsperma, Guss. ex Bertol. Fl. Ital. iii. 241 = silaifolia.
millefolia, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 177.—Thracia.
montana, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza montana.
Mutellina, DC. Prod. iv. 139 = Ligusticum Mutellina.
nodiflora, Schousb. Vextr. Marokko, 135 (= *Sclerosciadium nodiflorum*).—Marocc.
obscura, Spreng. Syst. i. 890 = Lichtensteinia pyrethrifolia.
patens, Moench, Meth. 91 = peucedanifolia.
peucedanifolia, Heuff. ex Boiss. Fl. Orient. ii. 958 = media.
peucedanifolia, *Pollich, Hist. Pl. Palat.* i. 289. f. 3.—Europ.
peucedanifolia, Schleich. ex DC. Prod. iv. 137 = Lachenalii.
peucedanoides, Roth, Beitr. i. 19 = peucedanifolia.
peucedanoides, Wredow, Fl. Meklenb. i. 516 = Lachenalii.
Phellandrium, *Lam. Fl. Fr.* iii. 432.—Europ.; As. bor.
Phellandrium, Nutt. Gen. Am. i. 189 = sarmentosa.
pimpinelloides, *Linn. Sp. Pl.* 255.—Europ.; As. Min.; Afr. bor.
pimpinelloides, Sm. Engl. Bot. t. 347 = Lachenalii.
Pollichii, C. C. Gmel. Fl. Bad. i. 679 = peucedanifolia.
prolifera, Hort. ex Steud. Nom. ed. I. 561 = virgata.
prolifera, Host, ex Reichb. Fl. Germ. Excurs. 463 = silaifolia.
prolifera, *Linn. Sp. Pl.* 254.—Europ. or.; Syria.
pteridifolia, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 30.—Ins. Madera.
purpurea, Poir. in Lam. Encyc. iv. 530 = Ligusticum Mutellina.
quitensis, Spreng. Syst. i. 888 = Ottoa oenanthoides.
Rarania, J. F. Gmel. Syst. 485, in syn. = silaifolia.
rhenana, DC. Fl. Fr. Suppl. 506 = Lachenalii.
rigida, Crantz, Cl. Umb. Emend. 85 = Peucedanum ternatum.
rigido-striata, Steud. Nom. ed. II. ii. 205 = Seseli tortuosum.
sarmentosa, Boland. Cat. 14; ex S. Wats. Bibliog. Ind. Bot. N. Am. 426 = californica.
sarmentosa, *Presl, ex DC. Prod.* iv. 138.—Am. bor. occ.
seseloides, Presl, Bot. Bemerk. 74 = Peucedanum ferulaceum.
silaifolia, Ball, in Ann. & Mag. Nat. Hist. Ser. I. xiv. (1844) 6 = peucedanifolia.
silaifolia, *Bieb. Fl. Taur. Cauc.* iii. 232.—Europ. austr.; Reg. Caucas.
silaifolia, Heuff. ex Rochel, Reise Bot. 66 = banatica.
Smithii, H. C. Wats. in Phytol. ii. (1845) 14 = peucedanifolia.
spuria, D. Dietr. Syn. Pl. ii. 953 = Annesorhiza spuria.
stenoloba, Schur, Enum. Pl. Transs. 255 = media.
stolonifera, *Wall. Cat.* n. 585; *DC. Prod.* iv. 138.—Ind. or.; Java; China.
stylosa, Pers. Syn. i. 318 = virgata.
Tabernaemontana, C. C. Gmel. Fl. Bad. i. 676 = fistulosa.
tenuifolia, *Boiss. & Orph. Diagn.* Ser. II. vi. 79.—Macedon.
tenuifolia, Froel. ex DC. Prod. iv. 137 = silaifolia.
tenuifolia, Thunb. Prod. Pl. Cap. 49 (= *Glia gummifera*).—Afr. austr.
teretifolia, Muhl. Cat. 31 = Peucedanum teretifolium.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 697.—Reg. Himal.
thracica, *Griseb. Spicil. Fl. Rumel.* i. 355.—Thracia.
virgata, Bory & Chaub. Nouv. Fl. Pélop. 19 = Carum multiflorum.
virgata, Griseb. Spicil. Fl. Rumel. i. 355 = thracica.
virgata, Koch, ex DC. Prod. iv. 137 = silaifolia.
virgata, *Poir. in Lam. Encyc.* iv. 529.—Afr. bor.

OENOCARPUS, Mart. Hist. Nat. Palm. ii. 21 (1823). *PALMAE*, Benth. & Hook. f. iii. 897.
altissimus, *Klotzsch, ex H. Wendl. Ind. Palm.* 30.—Venezuela.
Bacaba, *Mart. Hist. Nat. Palm.* ii. 24. t. 26. ff. 1, 2.—Guiana.
Bataua, *Mart. l. c.* 23.—Bras.
Batawa, Wallace, Palm-trees Amaz. 31. t. 10 = praec.
bolivianus, *Hort. Kew. ex H. Wendl. Ind. Palm.* 30.—Am. austr.
Calaber, Hort. ex H. Wendl. l. c. = caracasanus.
caracasanus, *H. Wendl. l. c.*—Venezuela.
Chiragua, *Hort. ex H. Wendl. l. c.*—Venezuela?
circumtextus, *Mart. Hist. Nat. Palm.* ii. 26. t. 26. ff. 3, 4.—Bras.
Cubarro, *Hort. ex H. Wendl. Ind. Palm.* 30.—Hab.?
dealbatus, Hort. ex H. Wendl. in Kerch. Palm. 252 = Mauritia armata.
distichus, *Mart. Hist. Nat. Palm.* ii. 22. tt. 22, 23.—Am. austr.
frigidus, Spreng. Syst. ii. 140 = Oreodoxa frigida.
glaucus, Lodd. ex H. Wendl. Ind. Palm. 30 = caracasanus.
gracilis, *Hort. ex H. Wendl. in Kerch. Palm.* 252, nomen.—Hab.?
Mapora, *Karst. in Linnaea*, xxviii. (1856) 274.—Venezuela.
minor, *Mart. Hist. Nat. Palm.* ii. 25. t. 27.—Bras.
multicaulis, *Spruce, in Journ. Linn. Soc.* xi. (1871) 142.—Bras.
pulchellus, *Linden, Illustr. Hortic.* xxviii. (1881) 32.—N. Granat.
regius, Spreng. Syst. ii. 140 = Oreodoxa regia.
Sancona, Spreng. l. c. = Oreodoxa Sancona.
Tarampabo, *Mart. Palmet. Orbign.* 12. t. 8. f. 3.—Bolivia.
utilis, Klotzsch, in Linnaea, xx. (1847) 447 = Euterpe acuminata.

OENONE, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96 = **Ligea**, Tul. (Podost.).
alcicornis, Wedd. in DC. Prod. xvii. 58 = L. alcicornis.
flexuosa, Wedd. l. c. = L. flexuosa.
longifolia, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 96 = L. longifolia.

OENOPLEA, Hedw. f. Gen. i. 151 (1806) = **Berchemia**, Neck. (Rhamn.).
lineata, Roem. & Schult. Syst. v. 333 = B. racemosa.
volubilis, Roem. & Schult. l. c. 332 = B. volubilis.

OENOPLIA, Roem. & Schult. Syst. v. p. xxvii; 332 (1819) = praec.

OENOSCIADIUM, Pomel, Nouv. Mat. Fl. Atl. 141 (1874). *UMBELLIFERAE*.
anomalum, *Pomel, l. c.*—Afr. bor.

OENOTHERA, Linn. Syst. ed. I (1735). *ONAGRARIAE*, Benth. & Hook. f. i. 789.
AENOTHERA, Lam. Encyc. iv. 550 (1797).
AGASSIZIA, Spach, Hist. Vég. Phan. iv. 347 (1835).
ANOGRA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 339.
BAUMANNIA, Spach, Hist. Vég. Phan. iv. 351 (1835).
BLENNODERMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 406.
BOISDUVALIA, Spach, Hist. Vég. Phan. iv. 383 (1835).
CALYLOPHUS, Spach, l. c. 349 (1835).
CAMISSONIA, Link, in Jahrb. Gew. (1818) 186.
CRATERICARPIUM, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 397.
DICTYOPETALUM, 'Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 45 (1835)' [§] ex Baill. Dict. ii. 410.
GODETIA, Spach, Hist. Vég. Phan. iv. 386 (1835).
HARTMANNIA, Spach, l. c. 370 (1835).
HOLOSTIGMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 332.
KNEIFFIA, Spach, Hist. Vég. Phan. iv. 373 (1835).
LAVAUXIA, Spach, l. c. 366 (1835).
MEGAPTERIUM, Spach, l. c. 363 (1835).
MERIOLIX, Rafin. in Am. Monthly Mag. (1819) 192.
ONAGRA, Spach, Hist. Vég. Phan. iv. 357 (1835).
PACHYLOPHUS, Spach, l. c. 365 (1835).

OENOTHERA:—
SPHAEROSTIGMA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 49 (1835).
XYLOPLEURUM, Spach, Hist. Vég. Phan. iv. 378 (1835).
acaulis, *Cav. Ic.* iv. 68. t. 399.—Chili.
acaulis, Lindl. in Bot. Reg. t. 763 = cuspidata.
affinis, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 269 = mollissima?
affinis, Penny, ex Loud. Gard. Mag. v. (1829) 470 = amoena.
Agassizia, *Steud. Nom.* ed. II. ii. 206.—Chili.
alata, Nutt. Gen. Am. i. 248 = missouriensis.
alata, Rafin. Fl. Ludov. 95 = Jussieua decurrens.
alba, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 = Ludwigia leucorrhiza?
albescens, *S. Wats. in Proc. Am. Acad.* viii. (1873) 597.—Calif.
albicans, *Lam. Encyc.* iv. 552.—Peruv.
albicaulis, *Fras. Cat.* (1813); *ex Nutt. Gen. Am.* i. 245.—Am. bor.
albicaulis, Pursh, Fl. Am. Sept. ii. 733 = pinnatifida, *Nutt.*
alyssoides, *Hook. & Arn. Bot. Beech. Voy.* 340.—Am. bor. occ.
ambigua, Spreng. Syst. ii. 229 = fruticosa.
ambigua, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 293. —Am. bor. occ.
amoena, *Lehm. Ind. Sem. Hort. Hamb.* (1821) 8; *Pugill. Pl.* i. 22.—Am. bor.
andina, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 512.—Am. bor. occ.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = biennis.
anisoloba, *Sweet, Brit. Flow. Gard.* Ser. II. t. 105.—Ins. Chiloe.
anomala, Curt. Bot. Mag. t. 388 = Gaura mutabilis.
arborea, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 32.—Calif.
arcuata, Kellogg, l. c. i. (1855) 58 = amoena.
Arnottii, Torr. & Gray, Fl. N. Am. i. 503 = decumbens.
ascendens, *Willd. ex Spreng. Syst.* ii. 230.—Am. austr.
asperifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 509 = hirta.
aurantia, Willd. ex Spreng. Syst. ii. 229 = multicaulis.
australis, *Salisb. Prod.* 278.—Reg. Magellan.
Berlandieri, Steud. Nom. ed. II. ii. 206 = serrulata.
Berlandieri, Walp. Rep. ii. 85 = Delessertiana.
Berteriana, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 343.—Chili.
biennis, *Linn. Sp. Pl.* 346.—Am. bor.
biennis, Walt. Fl. Carol. 129 = sinuata.
bifrons, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 386 = heterophylla.
bifrons, Lindl. in Bot. Reg. t. 1405 = amoena.
biloba, Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 87.—Calif.
bistorta, A. Gray, in Proc. Bost. Soc. Nat. Hist. vii. (1859–61) 146 = hirta.
bistorta, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 508.—Am. bor. occ.
Boothii, *Dougl. in Hook. Fl. Bor. Am.* i. 213.—Am. bor. occ.
Bottae, Steud. Nom. ed. II. ii. 206 = bistorta.
Bottae, *Torr. & Gray, Fl. N. Am.* i. 505.—Calif.
brachycarpa, *A. Gray, Pl. Wright.* i. 70.—N. Mexic.
brachysepala, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 345.—Chili.
bracteata, *Phil. in Anal. Univ. Chil.* (1862) II. 394.—Chili.
brasiliensis, Gardn. ex Micheli, in Mart. Fl. Bras. xiii. II. 155 = Jussieua nervosa.
Braunii ×, Doell, Fl. Bad. 1077, adnot.—Europ.
breviflora, *Torr. & Gray, Fl. N. Am.* i. 506.—Am. bor. occ.
brevipes, *A. Gray, ex Torr. in Pacif. Rail. Rep.* iv. 87. —Calif.
caesia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 514 = Gayophytum racemosum.
caespitosa, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 311 = subulata.
caespitosa, *Nutt. in Fras. Cat.* (1813); *ex Nutt. Gen. Am.* i. 246.—Am. bor. occ.
californica, D. Dietr. Syn. Pl. ii. 1288 = Bottae.
californica, *S. Wats. in Bot. Calif.* i. 223.—Calif.
campylocalyx, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) 17.—Hab.?

OENOTHERA:—
campylocarpa, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 291 = praec.
canadensis, Goldie, in Edinb. Phil. Journ. vi. (Apr. 1822) 325 = fruticosa.
canescens, *Torr. in Frém. Rep.* 315.—Am. bor. occ.
capensis, Hort. ex Steud. Nom. ed. II. ii. 206 = tetraptera.
capillifolia, Scheele, in Linnaea, xxi. (1848) 576 = serrulata.
cardiophylla, *Torr. in Pacif. Rail. Rep.* v. 360.—Am. bor. occ.
catharinensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 270.—Bras.
cedrosensis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 187.—Calif.
centaureifolia, *Steud. Nom.* ed. II. ii. 206.—Reg. Argent.
chamaenerioides, *A. Gray, Pl. Wright.* ii. 58.—N. Mexic.
Chamissonis, Link, Enum. Hort. Berol. i. 378 = dentata.
cheiranthifolia, A. Gray, in Ives, Rep. Colorado River, Bot. 12 = bistorta.
cheiranthifolia, *Hornem. ex Spreng. Syst.* ii. 228.—Chili.
cheiranthifolia, Lindl. Bot. Reg. t. 1040 = O. Agassizia.
chilensis, *D. Dietr. Syn. Pl.* ii. 1285.—Chili.
chrysantha, Michx. Fl. Bor. Am. i. 225 = pumila.
cinnabarina, *Hort. ex E. Morr. Belg. Hortic.* xiv. (1864) 75.—Hab.?
clavaeformis, Torr. & Frém. Rep. 314 = scapoidea.
clavaeformis, Torr. Bot. Mex. Bound. 66, partim = brevipes.
cochinchinensis, *Lour. ex Steud. Nom.* ed. I. 561, nomen.—China.
concinna, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 183.—Chili.
contorta, Boland. Cat. 12; ex S. Wats. in Proc. Am. Acad. viii. (1873) 606 = cheiranthifolia.
contorta, Dougl. ex Lehm. in Hook. Fl. Bor. Am. i. 214 = strigulosa.
coquimbensis, *C. Gay, Fl. Chil.* ii. 331.—Chili.
cordata, *Steud. Nom.* ed. II. ii. 206.—Chili; Ins. Juan Fernand.
coronopifolia, A. Gray, Pl. Wright. ii. 56 = pinnatifida.
coronopifolia, *Torr. & Gray, Fl. N. Am.* i. 495.—Am. bor. occ.
corymbosa, *Lam. Encyc.* iv. 554.—Mexic.
corymbosa, Sims, Bot. Mag. t. 1974 = Simsiana.
crassifolia, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 188.—Calif.
crassipes, Hort. Berol. ex Steud. Nom. ed. II. ii. 206, in syn. = elata.
crispa, *Schult, Obs. Bot.* 73.—Hab.?
cruciata, Nutt. ex G. Don, Gen. Syst. ii. 686 = biennis.
cruciformis, Kellogg, in Proc. Calif. Acad. ii. (1863) 227. f. 71 = scapoidea.
cuneifolia, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
cuprea, *Schlecht. in Linnaea*, xii. (1838) 269.—Mexic.
cuspidata, *Steud. Nom.* ed. II. ii. 206.—Am. austr.
decumbens, *Dougl. ex Hook. Bot. Mag.* t. 2889.—Calif.
Delessertiana, *Steud. Nom.* ed. II. ii. 206.—Mexic.
deltoides, Torr. & Frém. Rep. 313 = trichocalyx.
densiflora, *Lindl. Bot. Reg.* t. 1593.—Calif.
dentata, *Cav. Ic.* iv. 67. t. 398.—Am. occ., bor. et austr.
dentata, Link, ex Steud. Nom. ed. II. ii. 206 = heterophylla, *Steud.*
dentata, Ser. in DC. Prod. iii. 46, partim = hirta.
dentata, S. Wats. Bot. King's Exped. 111 = strigulosa.
diffusa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 513 = Gayophytum diffusum.
dissecta, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 357.—Mexic.
distorta, Veitch, ex Vilm. Fl. Pl. Terre, ed. III. 370 = bistorta.
divaricata, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 125.—Calif.
Drummondii, *Hook. Bot. Mag.* t. 3361.—Am. bor. occ.
Drummondii, Hook. Lond. Journ. Bot. vi. (1847) 221 = missouriensis.
Drummondii, Schnizl. Iconog. t. 265. f. 1 = speciosa.
Drummondii, Walp. Rep. ii. 85 = Spachiana.
dubia, Hort. ex Steud. Nom. ed. II. ii. 206 = tetraptera.
elata, *H. B. & K. Nov. Gen. et Sp.* vi. 90.—Mexic.

OENOTHERA:—
eminens, *Tausch, in Flora*, xxii. (1839) 559.—Hab.?
epilobiifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 92.—N. Granat.
epilobioides, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 511.—Am. bor.
erosa, *Lehm. in Linnaea*, iii. (1828) *Litt.* 8.—Afr. austr.
eximia, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 45 = caespitosa.
Fendleri, A. Gray, l. c. = Hartwegii.
fissifolia, *Steud. Nom.* ed. II. ii. 206.—Mexic.
florida, Salisb. Prod. 278 = fruticosa.
Fraseri, Pursh, Fl. Am. Sept. ii. 734 = glauca.
Fremontii, *S. Wats. in Proc. Am. Acad.* viii. (1873) 587.—Am. bor. occ.
fruticosa, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 44 = serrulata.
fruticosa, *Linn. Sp. Pl.* 346.—Am. bor.
gauraeflora, *Torr. & Gray, Fl. N. Am.* i. 510.—Am. bor. occ.
gauroides, Hornem. Hort. Hafn. i. 362 = biennis.
Gayana, *Steud. Nom.* ed. II. ii. 206.—Chili.
glabella, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 505.—Am. bor.
glabra, Mill. Gard. Dict. ed. VIII. n. 3 = biennis.
glauca, *Michx. Fl. Bor. Am.* i. 224.—Am. bor.
Glazioviana, *Micheli, in Mart. Fl. Bras.* xiii. II. 178. —Bras.
globularis, *Hort. Chels. ex G. Don, in Loud. Hort. Brit.* 154, nomen.—Hab.?
Godetia, Steud. Nom. ed. II. ii. 206 = Bottae.
graciliflora, *Hook. & Arn. Bot. Beech. Voy.* 341.—Calif.
graciliflora, Torr. in Pacif. Rail. Rep. iv. 87 = bistorta.
gracilis, Schrad. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 44 = pumila.
grandidentata, *Phil. in Linnaea*, xxxiii. (1864–65) 68.—Chili.
grandiflora, Ruiz & Pav. Fl. Per. iii. 78 = acaulis.
grandiflora, [Soland. in] Ait. Hort. Kew. ed. I. ii. 2; Lam. Encyc. iv. 554 = biennis.
grandiflora, S. Wats. in Proc. Am. Acad. viii. (1873) 596 = Whitneyi.
graveolens, Gilib. Fl. Lituan. ii. 186 = biennis.
Greggii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 46.—Mexic.
guadalupensis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 115, 137.—Calif.
guttata, *Molina, Sagg. Chile*, ed. II. 134.—Chili.
guttulata, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 222 = canescens.
Hartwegii, *Benth. Pl. Hartw.* 5.—Mexic.
Havardi, *S. Wats. in Proc. Am. Acad.* xx. (1885) 366. —Texas.
heterantha, *Nutt. in Journ. Acad. Philad.* vii. (1834) 22.—Am. bor. occ.
heterochroma, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 373.—Calif.
heterophylla, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 341 = hirta.
heterophylla, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 348.—Am. bor occ.
heterophylla, *Steud. Nom.* ed. II. ii. 207.—Chili.
Hilgardi, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 41.—Am. bor.
hirsuta, D. Dietr. Syn. Pl. ii. 1289 = biennis.
hirsuta, *Steud. Nom.* ed. II. ii. 206.—Mexic.
hirta, *Link, Enum. Hort. Berol.* i. 378.—Am. bor. occ.; Chili; Peruv.
hirta, Linn. Syst. ed. X. 998 = Jussieua peruviana.
hispidula, *S. Wats. in Proc. Am. Acad.* viii. (1873) 599. —Am. bor. occ.
holosericea, *Tausch, in Flora*, xxii. (1839) 558.—Hab.?
Hookeri, Torr. & Gray, Fl. N. Am. i. 493 = biennis.
humifusa, Lindl. Bot. Reg. t. 1829 = subulata.
humifusa, *Nutt. Gen. Am.* i. 245.—Am. bor.
humilis, Donn, Cat. Hort. Cantab. ed. I. 41; Willd. Sp. Pl. ii. 311 = purpurea.
hybrida, Michx. Fl. Bor. Am. i. 225 = fruticosa.
hyssopifolia, *Molina, Sagg. Chile*, ed. II. 134.—Chili.
incana, Nutt. Gen. Am. i. 247 = fruticosa.
indecora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 268.—Bras.
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 493.—Am. bor. occ.

OENOTHERA :—
Johnsoni, *Parry, in Am. Natural.* ix. (1875) 270.—Am. bor. occ.
laciniata, Hill, Hort. Kew. 172/4 = sinuata.
Lamarkiana, Ser. in DC. Prod. iii. 47 = biennis.
lampasana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 454 = Greggii ?
lasiocarpa, *Griseb. in Goett. Abh.* xix. (1874) 143.—Reg. Argent.
latiflora, [*Moç. & Sesse*] *ex Ser. in DC. Prod.* iii. 50.—Mexic.
lavandulaefolia, Torr. & Gray, Fl. N. Am. i. 501 = Hartwegii.
leona, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 455 = heterophylla, *Spach.*
lepida, D. Dietr. Syn. Pl. ii. 1287; Hook. & Arn. Bot. Beech. Voy. 342 = decumbens.
leptophylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 495 = albicaulis.
leucocarpa, Comien, ex Lehm. in Hook. Fl. Bor. Am. i. 210 = serrulata.
Lindleyana, Steud. Nom. ed. II. ii. 207 = seq.
Lindleyii, Dougl. in Bot. Mag. t. 2832 = amoena.
linearis, Michx. Fl. Bor. Am. i. 225 = fruticosa.
linifolia, *Nutt. in Journ. Acad. Philad.* ii. (1821) 120.—Am. bor.
lithospermoides, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 341 = Boothii.
littoralis, *Schlecht. in Linnaea*, xii. (1838) 268.—Mexic.
longiflora, *Linn. Mant.* ii. 227.—Reg. Argent.
longiflora, Scheele, in Roemer, Texas, 430; ex S. Wats. in Proc. Am. Acad. viii. (1873) 614 = sinuata.
macrantha, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 342 = amoena.
macrantha, *Sellow, ex Salm-Dyck, Hort. Dyck.* 182, nomen.—Hab. ?
macrocarpa, Pursh, Fl. Am. Sept. ii. 734 = missouriensis.
macrosceles, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 43.—Am. bor. occ.
macrosiphon, *Lehm. in Hamb. Gartenz.* xiv. (1858) 439.—Hab. ?
maculata, *Hort. Angl. ex Steud. Nom.* ed. II. ii. 207, nomen.—Am. bor.
malacophylla, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 344.—Chili.
marginata, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 342 = caespitosa.
maritima, Nutt. ex Hook. & Arn. l. c. 341 = graciliflora.
media, Link, Enum. Hort. Berol. i. 377 = fruticosa.
mendocinensis, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 310.—Chili.
mexicana, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 347 = sinuata.
micans, *Tausch, in Flora*, xxii. (1839) 559.—Hab. ?
micrantha, Hornem. ex Spreng. Syst. ii. 228 = hirta.
micrantha, Nutt. ex Torr. & Gray, Fl. N. Am. i. 514 = Gayophytum racemosum.
micrantha, *Presl, Rel. Haenk.* ii. 31.—Chili.
micrantha, Walp. Rep. ii. 84 = fissifolia.
minima, Pursh, Fl. Am. Sept. i. 262 = sinuata.
minutiflora, *D. Dietr. Syn. Pl.* ii. 1285.—Chili.
missouriensis, *Sims, Bot. Mag.* t. 1592, et sub t. 1674 (missourensis).—Am. bor. occ.
mollissima, Bert. ex Steud. Nom. ed. II. ii. 207 = cordata.
mollissima, Hook. & Arn. Bot. Beech. Voy. 23 = odorata.
mollissima, *Linn. Sp. Pl.* 346.—Reg. Argent.
mollissima, Walt. Fl. Carol. 129 = fruticosa.
montana, Nutt. in Torr. & Gray, Fl. N. Am. i. 500 = caespitosa.
multicaulis, *Ruiz & Pav. Fl. Per.* iii. 80. t. 317.—Peruv.
multijuga, *S. Wats. in Am. Natural.* vii. (1873) 300.—Am. bor.
muricata, Linn. Syst. ed. XII. 263 = biennis.
mutabilis, Steud. Nom. ed. I. 561 = tetraptera.
nana, *Griseb. in Goett. Abh.* xix. (1874) 143.—Reg. Argent.
nervosa, *Hornem. ex Sweet, Hort. Brit.* ed. II. 199, nomen.—Hab. ?
nevadensis, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 224. f. 70.—Calif.
nocturna, *Jacq. Coll.* iii. 205; *Ic. Pl. Rar.* iii. 3. t. 455.—Afr. austr.
nocturna, Willd. ex Spreng. Syst. ii. 228 = mollissima.

OENOTHERA :—
Nuttallii, Sweet, Hort. Brit. ed. II. 199 = albicaulis.
Nuttallii, Torr. & Frém. Rep. 89 = breviflora.
Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 506.—Am. bor. occ.
Oakesiana, *Robbins, ex S. Wats. Bibliog. Ind. N. Am. Bot.* 383.—Am. bor.
obtusifolia, D. Dietr. Syn. Pl. ii. 1289 = speciosa.
octovalvis, Jacq. Enum. Pl. Carib. 19 = Jussieua suffruticosa.
odorata, Hook. & Arn. Bot. Beech. Voy. 343 = biennis.
odorata, *Jacq. Ic. Pl. Rar.* iii. 3. t. 456; *Coll. Suppl.* 107.—Chili.
odoratissima, *Tausch, in Flora*, xxii. (1839) 557.—Hab. ?
ovata, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 507.—Calif.
pallida, Lindl. Bot. Reg. t. 1142 = albicaulis.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 251.—Am. bor. occ.
paludosa, Featherm. Rep. Louis. Univ. (1870) 71 = Jussieua decurrens.
Parryi, *S. Wats. in Am. Natural.* ix. (1875) 19; 270.—Am. bor. occ.
parviflora, Linn. Syst. ed. X. 998 = biennis.
parvula, Nutt. ex Torr. & Gray, Fl. N. Am. i. 511 = strigulosa.
perampla, *R. Grah. in Edinb. N. Phil. Journ.* (Oct. 1832) 362.—Peruv.
perennis, Linn. Syst. ed. X. 998 = pumila.
pilosella, Rafin. Ann. Nat. 15 = fruticosa.
pinnatifida, Hort. ex Steud. Nom. ed. II. ii. 207 = fissifolia.
pinnatifida, *H. B. & K. Nov. Gen. et Sp.* vi. 91.—Mexic.
pinnatifida, *Nutt. Gen. Am.* i. 245.—Am. bor.
pinnatifida, Torr. in Ann. Lyc. N. York, ii. (1828) 201 = albicaulis.
primiveris, *A. Gray, Pl. Wright.* ii. 58.—N. Mexic.
propinqua, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 343.—Chili.
prostrata, Ruiz & Pav. Fl. Per. iii. 79. t. 315 = sinuata.
psychrophila, *Ball, in Journ. Linn. Soc.* xxii. (1885) 38.—Peruv.
pterosperma, *S. Wats. in Bot. King's Exped.* 112. t. 14.—Am. bor. occ.
pubescens, Nees, in Wied-Neuwied, Reise N. Am. ii. 435 = biennis.
pubescens, *Willd. ex Spreng. Syst.* ii. 229.—Am. austr.
pumila, *Linn. Sp. Pl.* ed. II. 493.—Am. bor.
purpurea, Benth. Pl. Hartw. 310 = albescens.
purpurea, *Curt. Bot. Mag.* t. 352.—Am. bor.
purpurea, Durand, in Journ. Acad. Philad. Ser. II. iii. (1855) 88 = decumbens, quadrivulnera.
purpurea, A. Gray, in Proc. Am. Acad. viii. (1873) 384, partim = tenella.
purpurea, Hort. ex Lam. Encyc. iv. 554 = rosea.
Purshiana, Steud. Nom. ed. II. ii. 207 = seq.
Purshii, G. Don, Gen. Syst. ii. 688 = pinnatifida, *Nutt.*
pusilla, Michx. Fl. Bor. Am. i. 225 = pumila.
pygmaea, Dougl. ex Lehm. in Hook. Fl. Bor. Am. i. 213 = Boothii.
quadricolumnata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 21 = Ludwigia parviflora.
quadrivulnera, *Dougl. ex Lindl. Bot. Reg.* t. 1119.—Am. bor.
racemosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 514 = Gayophytum racemosum.
ramosissima, Nutt. l. c. 513 = Gayophytum ramosissimum.
ramulosa, *Steud. Nom.* ed. II. ii. 207.—Chili.
refracta, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 373.—Calif.
repanda, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 198. t. 8 = sinuata.
rhizocarpa, Spreng. Syst. ii. 230 = triloba.
rhombipetala, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 216 = heterophylla, *Spach.*
rhombipetala, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 493.—Am. bor.
rigida, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 311 = micrantha.
riparia, Lehm. in Hook. Fl. Bor. Am. i. 212 = pumila.
riparia, Nutt. Gen. Am. i. 247 = fruticosa.
Roemeriana, Scheele, in Linnaea, xxii. (1849) 154 = triloba.

OENOTHERA :—
Romanzowii, *Ledeb. ex Hornem. Hort. Hafn. Suppl.* 133.—Am. bor.
rosea, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 3.—Am. bor. occ.; Peruv.
roseo-alba, Bernh. Ind. Sem. Hort. Erford. (1824); ex Reichb. Ic. Bot. Exot. i. 34. t. 47 = amoena.
rubicunda, Durand, in Journ. Acad. Philad. Ser. II. iii. (1855) 87 = hispidula.
rubicunda, Hook. & Arn. Bot. Beech. Voy. 342; D. Dietr. Syn. Pl. ii. 1287 = amoena.
rubicunda, Torr. & Gray, Fl. N. Am. i. 502 = Bottae.
rubicunda, Torr. & Gray, in Pacif. Rail. Rep. ii. 121 = epilobioides.
rubra, Cav. Ic. iii. 68 = rosea.
salicifolia, Desf. Tabl. ed. II. 271 = elata.
salicina, Nutt. ex Torr. & Gray, Fl. N. Am. i. 505 = densiflora.
Sandiana, *Hassk. in Flora*, xxxix. (1856) 516.—Peruv.
scapigera, Pursh, Fl. Am. Sept. i. 263 = caespitosa.
scapoidea, *Torr. & Gray, Fl. N. Am.* i. 506.—Am. bor. occ.
scoparia, Ruiz & Pav. ex D. Don, in Edinb. N. Phil. Journ. (July–Sept. 1831) 277 = Balbisia verticillata.
Sellowii, *Link & Otto, ex Steud. Nom.* ed. II. ii. 207, nomen.—Am. austr.
serotina, *Lehm. Sem. Hort. Hamb.* (1825) 17.—Hab. ?
serotina, Sweet, Brit. Flow. Gard. Ser. I. ii. t. 184 = fruticosa.
serrulata, *Nutt. Gen. Am.* i. 246.—Am. bor.
siliquosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 512 = strigulosa.
Simsiana, *Ser. in DC. Prod.* iii. 47.—Mexic.
sinuata, *Linn. Mant.* ii. 228.—Am. bor.
Spachiana, Steud. Nom. ed. II. ii. 207 = serrulata.
Spachiana, *Torr. & Gray, Fl. N. Am.* i. 498.—Am. bor. occ.
Spachii, D. Dietr. Syn. Pl. ii. 1289 = speciosa.
speciosa, *Nutt. in Journ. Acad. Philad.* ii. (1821) 119.—Am. bor.
spectabilis, *Bernh. ex Steud. Nom.* ed. I. 562, nomen.—Mexic.
spectabilis, Hort. ex Spach, Hist. Vég. Phan. iv. 358 = Simsiana.
spinulosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 502 = serrulata.
spiralis, Hook. Fl. Bor. Am. i. 213 = cheiranthifolia.
stenophylla, Steud. Nom. ed. II. ii. 208 = chilensis.
striata, *Ledeb. in Link, Enum. Hort. Berol.* i. 377.—Hab. ?
stricta, Steud. Nom. ed. II. ii. 208, sphalm. = praec.
strigosa, Willd. ex Spreng. Syst. ii. 228 = tenuifolia.
strigulosa, Benth. Pl. Hartw. 310 = dentata.
strigulosa, *Torr. & Gray, Fl. N. Am.* i. 512; *D. Dietr. Syn. Pl.* ii. 1284.—Am. bor. occ.
suaveolens, Desf. Tabl. ed. I. 169; et Pers. Syn. i. 408 = biennis.
subincana, *Hort. ex Steud. Nom.* ed. II. ii. 208, nomen.—Hab. ?
subulata, *Ruiz & Pav. Fl. Per.* iii. 82. t. 316.—Chili.
tanacetifolia, Torr. & Gray, in Pacif. Rail. Rep. ii. 121 = Nuttallii.
taraxacifolia, Hort. ex Sweet, Brit. Flow. Gard. Ser. I. t. 294 = acaulis.
tarquensis, *H. B. & K. Nov. Gen. et Sp.* vi. 91.—Ecuador.
tasmanica, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 475.—Tasmania.
tenella, Bert. ex Steud. Nom. ed. II. ii. 208 = ramulosa.
tenella, *Cav. Ic.* iv. 66. t. 396. f. 2.—Chili.
tenella, A. Gray, in Proc. Am. Acad. viii. (1873) 384 = epilobioides.
tenella, A. Gray, in Proc. Bost. Soc. Nat. Hist. Ser. vii. (1859–61) 164 = Bottae.
tenuifolia, Bert. ex D. Dietr. Syn. Pl. ii. 1285 = chilensis.
tenuifolia, *Cav. Ic.* iv. 67. t. 397.—Chili.
tetragona, Roth, Catalect. fasc. ii. 39 = fruticosa.
tetraptera, *Cav. Ic.* iii. 40. t. 279.—Mexic.
Torreyi, *S. Wats. in Proc. Am. Acad.* viii. (1873) 600.—Calif.
trichocalyx, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 494.—Am. bor. occ.
triloba, Hook. Lond. Journ. Bot. vi. (1847) 223 = heterantha.

OENOTHERA:—
triloba, *Nutt. in Journ. Acad. Philad.* ii. (1821) 118.—Am. bor. occ.
tubicula, *A. Gray, Pl. Wright.* i. 71.—N. Mexic.
tubifera, [*Moç. & Sesse*] *ex Ser. in DC. Prod.* iii. 50.—Mexic.
uncinata, Scheele, in Linnaea, xxi. (1848) 578 = Spachiana.
undulata, Ait. Hort. Kew. ed. II. ii. 342 = odorata.
uniflora, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 311 = subulata.
uniflora, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen (Quid ?).—Am. bor.
variifolia, Steud. Nom. ed. II. ii. 208 = heterophylla, *Spach.*
versicolor, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 291 = campylocalyx.
versicolor, *Lehm. Ind. Sem. Hort. Hamb.* (1855) 7.—*Cf. Linnaea*, xxviii. (1856) 359.—Hab.?
villosa, *Thunb. Prod. Pl. Cap.* 75.—Afr. austr.
viminea, *Dougl. ex Hook. Bot. Mag.* t. 2873.—Calif.
viminea, Torr. in Pacif. Rail. Rep. iv. 86, partim = quadrivulnera.
viminea, S. Wats. Bot. King's Exped. 108 = epilobioides.
vinosa, Torr. & Gray, Fl. N. Am. i. 503; D. Dietr. Syn. Pl. ii. 1287 = amoena.
vinosa, Torr. Bot. Mex. Bound. 66 = epilobioides.
virgata, *Ruiz & Pav. Fl. Per.* iii. 79.—Peruv.
viridescens, Hook. Fl. Bor. Am. i. 214 = cheiranthifolia.
viscosa, Rafin. Fl. Ludov. 96 = sinuata ?
Webbiana, Steud. Nom. ed. II. ii. 208 = speciosa.
Whitneyi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 340.—Calif.
Williamsoni, *S. Wats. l. c.* viii. (1873) 597.—Calif.
Wrightii, *A. Gray, Pl. Wright*, ii. 57.—Texas.
Wrightii, Torr. Bot. Mex. Bound, 65 = Fremontii.

OEOLLANTHUS, G. Don, Gen. Syst. iv. 685 (1837) = **Aeolanthus**, Mart. (Labiat.).

OEONIA, Lindl. Orch. Scel. 14 (1826) = **Aeonia**, Lindl. (Orchid.).

OERSTEDELLA, Reichb. f. in Bot. Zeit. x. (1852) 932 = **Epidendrum**, Linn. (Orchid.).
centradenia, Reichb. f. l. c. = E. centropetalum.
centropetala, Reichb. f. l. c. = E. centropetalum.

OESCHINOMENE, Poir. in Lam. Encyc. iv. 447 (1797) = **Aeschynomene**, Linn. (Legumin.).

OESCULUS, Neck. Elem. ii. 232 (1790) = **Aesculus**, Linn. (Sapind.).

OETHIONEMA, Knowles & Westc. Fl. Cab. i. (1837) 42 = **Aethionema**, R. Br. (Crucif.).

OFAISTON, Rafin. Fl. Tellur. iii. 46 (1836).
CHENOPODIACEAE, Benth. & Hook. f. iii. 71.
OPSIESTON, Bunge, in Mém. Sav. Etr. Petersb. vii. (1847) 474 (Rel. Lehm. 298).
monandrum, Moq. in DC. Prod. xiii. II. 203 = pauciflorum.
pauciflorum, Moq. l. c. = paucifolium.
paucifolium, *Rafin. Fl. Tellur.* iii. 46.—As. bor.

OFTIA, Adans. Fam. ii. 199 (1763). *MYOPORINEAE*, Benth. & Hook. f. ii. 1125.
BATINDUM, Rafin. Sylva Tellur. 81 (1838).
SPIELMANNIA, Medik. in Act. Acad. Theod. Palat. iii. (1775) 196. t. 15.
africana, *Bocq. ex Baill. Adansonia*, ii. (1861–62) 11.—Afr. austr.
revoluta, *Bocq. ex Baill. l. c.* 12.—Afr. austr.

OGCEROSTYLUS, Cass. in Dict. Sc. Nat. xxiii. 563 (1822); xlix. 221 (1827) = **Angianthus**, Wendl. (Compos.).

OGCODEIA, Bur. in DC. Prod. xvii. 282 (1873) = **Noyera**, Tréc. (Urtic.).

OGIERA, Cass. in Bull. Soc. Philom. (1818) 32 = **Eleutheranthera**, Poit. (Compos.).
Eleuthranthera, Steud. Nom. ed. II. i. 549 = E. ovata.

OGIERA:—
leiocarpa, Cass. in Dict. Sc. Nat. xliii. 371 = Eleutheranthera ovata.
ruderalis, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 513 = E. ovata.
triplinervia, Bert. ex DC. Prod. v. 501, 546 = Podanthus Mitiqui.
triplinervia, Cass. in Dict. Sc. Nat. xxxv. 445 = E. ovata.
triplinervia, Spreng. Syst. iii. 674 = Podanthus ovatifolius.

OGINETIA, Wight, Ic. iii. t. 895 [1843], sphalm. = **Aeginetia**, Linn. (Orobanch.).

OGLIFA, Cass. in Bull. Soc. Philom. (1819) 143 = **Filago**, Linn. (Compos.).
arvensis, Cass. in Dict. Sc. Nat. xxxv. 448 = F. arvensis.
minima, Reichb. f. Ic. Fl. Germ. xvi. 27 = F. minima.
neglecta, Reichb. f. l. c. = F. neglecta.
Soyerii, Godr. Fl. Lorr. ed. I. ii. 34 = F. neglecta.

O-HIGGENSIA, Steud. Nom. ed. II. ii. 208 (1841) = seq.

OHIGGINSIA, Ruiz & Pav. Fl. Per. i. 55 (1798) = **Hoffmannia**, Sw. (Rubiac.).
aggregata, Ruiz & Pav. l. c. (= *Euosmia aggregata*).—Peruv.
obovata, Ruiz & Pav. l. c. 56.—Peruv.
verticillata, Ruiz & Pav. l. c. 55.—Peruv.

OHLENDORFFIA, Lehm. Del. Sem. Hort. Hamb. (1835).—Cf. Linnaea, xi. (1837) Litt. 91 = **Aptosimum**, Burch. (Scroph.).
caracasana, Hort. ex Pasq. Cat. Ort. Bot. Nap. 71.—Hab.?
procumbens, Schrad. Del. Sem. Hort. Hamb. (1835).—Cf. Linnaea, xi. (1837) Litt. 91 = A. depressum.
rosulata, Nees, ex Benth. in DC. Prod. x. 345 = A. indivisum.

OIANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 775 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
Beddomei, *Hook. f. Fl. Brit. Ind.* iv. 49.—Ind. or.
disciflorus, *Hook. f. l. c.*—Ind. or.
urceolatus, *Benth. in Hook. Ic. Pl.* t. 1191.—Ind. or.

OILEUS, Haw. Monog. Narciss. 4 (1831) = **Narcissus**, Tourn. (Amaryll.).
abscissus, albus, hexangularis, minor, pumilus, Haw. l. c. omnes = N. Pseudo-Narcissus.

OIOSPERMUM, Less. in Linnaea, iv. (1829) 339. *COMPOSITAE*, Benth. & Hook. f. ii. 225.
involucratum, *Less. l. c.*—Bras.
Metzianum, Sch. Bip. ex Hook. f. in Fl. Brit. Ind. iii. 228 = Centratherum Metzianum.
molle, Hook. f. l. c. 227, in syn. = Centratherum molle.
nigritianum, Benth. in Hook. Niger Fl. 425 = Gutenbergia nigritiana.
Wightianum, DC. Prod. v. 11 = Carpesium cernuum.

OISODIX, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
decipiens, Rafin. l. c. = S. fragilis.
fulva, Rafin. l. c. 16 (Quid ?).—Am. bor.
glauca, Rafin. l. c. 13 = S. glauca.
holosericea, Rafin. l. c. = S. Smithiana.
longifolia, Rafin. l. c. = S. longifolia.
oblongifolia, Rafin. l. c. 16 (Quid ?).—Am. bor.
reticulata, Rafin. l. c. 13 = S. reticulata.
serpyllifolia, Rafin. l. c. = S. retusa.

OKEA, Steud. Nom. ed. I. 562 (1821) = **Adenandra**, Willd. (Rutac.).

OKENIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 307 (1819) = **Adenandra**, Willd. (Rutac.).

OKENIA, Schlecht. & Cham. in Linnaea, v. (1830) 92. *NYCTAGINEAE*, Benth. & Hook. f. iii. 6.
hypogaea, *Schlecht. & Cham. l. c.*—Mexic.

OLAX, Linn. Diss. Dass. 3 (1747); Amoen. Acad. i. 387 (1749). *OLACINEAE*, Benth. & Hook. f. i. 347.
FISSILIA, Comm. ex Juss. Gen. 260 (1789).

OLAX:—
LOPADOCALYX, Klotzsch, in Lehm. Pl. Preiss. i. 178 (1844).
PSEUDALEIOIDES, Thou. Gen. Nov. Madag. 15 (1806).
PSEUDALIEA, Thou. l. c. (1806).
ROXBURGHIA, Koen. ex Roxb. Fl. ed. Carey, i. 168 (1820).
SPERMAXYRUM, Labill. Nov. Holl. Pl. ii. 84. t. 233 (1806).
acuminata, *Wall. Cat.* n. 6781.—Reg. Himal.
aphylla, *R. Br. Prod.* 358.—Austral.
Bador, Buch.-Ham. in Wall. Cat. n. 6778 = scandens.
Baticulin, Blanco, Fl. Filip. ed. II. 589 = Litsea obtusata.
Benthamiana, *Miq. in Lehm. Pl. Preiss.* i. 228.—Austral.
Bernieriana, *Baill. Adansonia*, iii. (1862–63) 121.—Madag. occ.
Boiviniana, *Baill. l. c.*—Madag.
Breonii, *Baill. l. c.*—Ins. Borbon.
dissitiflora, *Oliver, Fl. Trop. Afr.* i. 350.—Afr. trop.
emirnensis, *Baker, in Journ. Linn. Soc.* xxi. (1884 331.—Madag.
Gambecola, *Baill. Adansonia*, iii. (1862–63) 121.—Afr. trop.
Gardneriana, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 375 = Liriosma Gardneriana.
Heyneana, Wall. Cat. n. 6780 = acuminata; Gomphandra axillaris.
humilis, Royle, ex Decne. in DC. Prod. viii. 525 = Cynanchum Arnottianum ?
hypoleuca, *Baill. Adansonia*, viii. (1868) 345.—N. Caled.
imbricata, *Roxb. Hort. Beng.* 5; *Fl. Ind.* i. 164.—Burma; Malaya.
longifolia, Wall. Cat. n. 6782 = acuminata, Gomphandra axillaris.
loranthiformis, *Griff. Notul.* iv. 691.—Ind. or.
lucida, Heyne, in Wall. Cat. n. 6777 = Wightiana.
macrophylla, *Benth. in Proc. Linn. Soc.* i. (1840) 89.—Guian.
Mannii, *Oliver, Fl. Trop. Afr.* i. 349.—Afr. trop.
merguensis, *Planch. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 576.—Burma; Malacca.
multiflora, *Rich. ex Baill. Adansonia*, iii. (1862–63) 121.—Ins. Philipp.
nana, *Wall. Cat.* n. 6733.—Reg. Himal.
obtusa, Blume, Bijdr. 131 = imbricata, scandens.
pauciflora, Benth. in Proc. Linn. Soc. i. (1840) 89 = Liriosma pauciflora.
Pervilleana, *Baill. Adansonia*, iii. (1862–63) 120.—Madag.
phoenicarpa, *Roxb. Hort. Beng.* 5, nomen.—Ind. or.
Phyllanthi, *R. Br. Prod.* 358.—Austral.
Pseudoleia, *Willd. ex Steud. Nom.* ed. I. 562.—Madag.
Pseudoleioides, *Willd. l. c.*—Madag.
psittacorum, Vahl, Enum. ii. 33 = scandens, Wightiana.
quercina, *Baill. Adansonia*, iii. (1862–62) 120.—Ins. Madag.
retusa, *F. Muell. ex Benth. Fl. Austral.* i. 392.—Austral.
scandens, *Roxb. Pl. Corom.* ii. 2. t. 102.—Ind. or.; Malaya.
Schomburgkii, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1177, nomen.—Guian.
sphaerocarpa, *Griff. Notul.* iv. 689.—Ind. or.
stricta, *R. Br. Prod.* 358.—Austral.
subscorpioides, *Oliver, Fl. Trop. Afr.* i. 350.—Afr. trop.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 342 = Cansjera Rheedii.
Thouarsiana, *Baill. Adansonia*, iii. (1862–63) 121.—Ins. Maurit.?
triplinervia, *Oliver, Fl. Trop. Afr.* i. 349.—Afr. trop.
uliginosa, Klotzsch, in Lehm. Pl. Preiss. ii. 230 = Benthamiana.
Velloziana, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 375 = Liriosma Velloziana.
viridis, *Oliver, Fl. Trop. Afr.* i. 349.—Afr. trop.
Wightiana, *Wall. Cat.* n. 6779.—Ind. or.; Zeylan.
zeylanica, *Linn. Sp. Pl.* 34.—Zeylan.
zeylanica, Wall. Cat. n. 6777 B. = Wightiana.

OLBIA, Medic. Malv. 41 (1787) = **Lavatera**, Linn. (Malvac.).
canescens, Moench, Meth. Suppl. 200 = L. maritima.
deflexa, Moench, l. c. = L. punctata.
flava, Presl, Fl. Sic. 180 = L. flava.

OLBIA :—
hastata, Medic. Malv. 42 = Lavatera Olbia.
heterophylla, Moench, Meth. Suppl. 201 = L. thuringiaca.
hispida, Presl, Fl. Sic. 179 = L. Olbia.
micans, Moench, Meth. 613 = L. micans.
obtusa, Medic. Malv. 42 (Quid ?).—Hab. ?
thuringica, Medic. l. c. = L. thuringiaca.
triloba, Medic. l. c. = L. triloba.

OLDENBURGIA, Less. in Linnaea, v. (1830) 252.
COMPOSITAE, Benth. & Hook. f. ii. 494.
SCYTALA, E. Mey. ex DC. Prod. vii. 12 (1838).
Arbuscula, *DC. Prod.* vii. 12.—Afr. austr.
grandis, Baill. Hist. Pl. viii. 97 = O. Arbuscula.
Papionum, *DC. Prod.* vii. 12.—Afr. austr.
paradoxa, *Less. in Linnaea*, v. (1830) 252.—Afr. austr.

OLDENLANDIA, P. Br. Hist. Jamaic. 208 (1756) = **Jussieua**, Linn. (Onagr.).

OLDENLANDIA, Linn. Gen. ed. I. 362 (1737).
RUBIACEAE, Benth. & Hook. f. ii. 58.
AGATHISANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 294 (1861).
EDRASTENIA, Rafin. New Fl. Am. iv. 101 (1836).
GERONTOGEA, Cham. & Schlecht. in Linnaea, iv. (1829) 154.
GONOTHECA, Blume, ex DC. Prod. iv. 429 (1830).
KARAMYSCHEWIA, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 266.
KOHAUTIA, Cham. & Schlecht. in Linnaea, iv. (1829) 156.
PELTOSPERMUM, Benth. in Hook. Niger Fl. 400 (1849).
PENTODON, Hochst. in Flora, xxvii. (1844) 552.
LISTERIA, Neck. Elem. i. 206 (1790).
STELMOTIS, Rafin. New Fl. Am. iv. 101 (1836).
THEYODIS, A. Rich. Tent. Fl. Abyss. i. 364 (1847).
abyssinica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 57.—Afr. trop.
acerosa, A. Gray, Pl. Wright. ii. 67 = Houstonia acerosa.
affinis, DC. Prod. iv. 428 = dichotoma.
alata, *Herb. Mus. Par. ex Miq. Fl. Ind. Bat.* ii. 193 ; *et Hook. f. Fl. Brit. Ind.* iii. 70.—Ind. or.
alata, Koen. ex Roxb. Hort. Beng. 11 ; Fl. Ind. i. 421 paniculata.
alsinifolia, G. Don, Gen. Syst. iii. 529 = corymbosa.
angustifolia, Benth. Fl. Hongk. 151 = Hedyotis tenelliflora.
angustifolia, A. Gray, Pl. Wright. ii. 68 = Houstonia angustifolia.
arenarioides, Willd. ex Schult. Mant. iii. 144 = Hedyotis arenarioides.
arguta, Hook. f. Fl. Brit. Ind. iii. 67 = linoides.
Arnottiana, Miq. Fl. Ind. Bat. ii. 194 = Anotis Leschenaultiana.
aspera, *DC. Prod.* iv. 428.—Ind. or. ; Afr. trop.
asperula, DC. l. c. 425 = Heynei.
Auricularia, F. Muell. Census, 74 = Hedyotis Auricularia.
biflora, Lam. Encyc. iv. 533 = corymbosa.
biflora, *Linn. Sp. Pl.* 119.—Ind. or.
biflora, Roth, Nov. Pl. Sp. 97 = Vahlia oldenlandioides.
Blumeana, DC. Prod. iv. 423, in syn. = paniculata.
Bojeri, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 53.—Afr. trop.
Boscii, *Chapm. Fl. S. U. St.* 181.—Am. bor.
brachiata, *Hook. f. Fl. Brit. Ind.* iii. 66.—Ind. or.
brachypoda, DC. Prod. iv. 424 = diffusa.
brachypoda, Zoll. ex Miq. Fl. Ind. Bat. ii. 192 = paniculata.
Burmanniana, G. Don, Gen. Syst. iii. 529 = corymbosa.
caerulea, A. Gray, Man. Bot. N. U. St. ed. II. 174 = Houstonia caerulea.
caespitosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 61.—Afr. trop.
caffra, *Eckl. & Zeyh. Enum.* 360.—Afr. trop. et austr.
callitrichoides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 506.—Cuba.
capensis, *Linn. f. Suppl.* 127.—Syria ; Afr. trop. et austr.
capillaris, DC. Prod. iv. 426 = corymbosa ?
capillipes, *Griseb. Cat. Pl. Cub.* 130.—Cuba.
ciliolata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 152.—Borneo.

OLDENLANDIA :—
coccinea, *Hook. f. Fl. Brit. Ind.* iii. 69.—Reg. Himal.
coerulescens, F. Muell. Census, 74 = Hedyotis caerulescens.
congesta, *Balf. f. in Baker, Fl. Maurit.* 138.—Ins. Mascar.
conostyla, DC. Prod. iv. 428 = Kadua Menziesiana.
cordata, Vell. Fl. Flum. 53 ; i. t. 143 = Coccocypselum umbellatum.
corymbosa, Ait. Hort. Kew. ed. I. i. 163 = lactea.
corymbosa, Herb. Madr. ex Wight. & Arn. Prod. 408 = Hedyotis pruinosa.
corymbosa, *Linn. Sp. Pl.* 119.—Reg. trop.
crassifolia, *DC. Prod.* iv. 427.—Ins. Philipp.
Crouchiana, F. Muell. Census, 74 = Hedyotis Crouchiana.
crystallina, *Roxb. Hort. Beng.* 11 ; *Fl. Ind.* i. 422.—Ind. or. ; Malaya.
debilis, *Forst. f. Prod.* 10.—Ins. Tonga.
decumbens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 54.—Afr. trop.
decumbens, Spreng. Pugill. ii. 36 = Vahlia viscosa.
Deppeana, *DC. Prod.* iv. 428.—Mexic.
depressa, Willd. Sp. Pl. i. 675 = corymbosa ?
dichotoma, *Hook. f. Fl. Brit. Ind.* iii. 67.—Ind. or.
dichotoma, Roxb. ex Wight & Arn. Prod. 416 = Heynii.
dichotoma, Spreng. Pugill. ii. 36 = Vahlia oldenlandioides.
dichotoma, Willd. ex Cham. & Schlecht. in Linnaea, iv. (1829) 155 = biflora.
diffusa, *Roxb. Hort. Beng.* 11 ; *Fl. Ind*, i. 423.—Ind. or. ; Malaya.
digyna, Retz. Obs. iv. 22 = Vahlia viscosa.
effusa, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 84. t. 48.—Afr. trop.
elatinoides, F. Muell. Census, 74 = Hedyotis elatinoides.
erythraeoides, *Miq. Fl. Ind. Bat.* ii. 353.—Sumatra.
flosculosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 60.—Afr. trop.
foetida, *Forst. f. Prod.* 10.—Ins. Tonga.
galioides, *Benth. & Hook. f. Gen.* ii. 58.—Austral.
globosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 54.—Afr. trop.
glomerata, *Michx. Fl. Bor. Am.* i. 83.—Am. bor. ; Ind. occ.
gracilis, *Hook. f. Fl. Brit. Ind.* iii. 68.—Ind. or.
graminifolia, DC. Prod. iv. 425 = stricta.
grandiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 57.—Afr. trop.
Greenei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 77.—N. Mexic.
Halaei, Wood, Class-Book (1861), 404 = seq. ?
Halei, *Chapm. Fl. S. U. St.* 181.—Am bor.
hedyotoides, *Boiss. Fl. Orient.* iii. 11.—Reg. Casp. ; Afr. trop.
herbacea, DC. Prod. iv. 425 = corymbosa.
herbacea, Roxb. Hort. Beng. 11 ; Fl. Ind. i. 424 = Heynei.
heterophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 109.—Japon.
Heynei, Oliver, in Trans. Linn. Soc. xxix. (1873) 84 = seq.
Heynii, *G. Don, Gen. Syst.* iii. 531.—Ind. or. ; Malaya ; Afr. trop.
hirsuta, Linn. f. Suppl. 127 = Hedyotis hirsuta.
hirsuta, *Vell. Fl. Flum.* 54 ; i. t. 144.—Bras.
hispida, *DC. Prod.* iv. 428 (Quid ?).—Ind. or.
hispida, Lam. Encyc. iv. 536 = Hedyotis hispida.
Hornei, *Baker, Fl. Maurit.* 139.—Ins. Seychelles.
Horneriana, *Miq. Fl. Ind. Bat.* ii. 190.—Sumatra.
humifusa, A. Gray, Pl. Wright. ii. 68 = Houstonia Wrightii.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 109 = Hedyotis hirsuta.
Korthalsiana, *Miq. l. c.* iv. 228.—Sumatra.
lactea, *DC. Prod.* iv. 427 (sp. dub.).—Ind. or.
lancifolia, *DC. l. c.* 425 ; *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 61.—Afr. trop.
lasiocarpa, *Hiern, l. c.* 55.—Afr. trop.
latifolia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 409.—Madag.
latifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. I. (1844) 235.—Mexic.
laxiflora, Benth. in Hook. Niger Fl. 404 = macrophylla.
Lechleriana, *Benth. & Hook. f. Gen.* ii. 58.—Peruv.

OLDENLANDIA :—
linarifolia, Hook. f. Fl. Brit. Ind. iii. 67, sphalm. = seq.
linearifolia, Herb. Madr. ex Wight & Arn. Prod. 416 = dichotoma.
linearifolia, *Thunb. Fl. Javan.* 9.—Java.
linearis, DC. Prod. iv. 425 = Heynii.
linifolia, DC. l. c. = stricta, Heynii.
linoides, *Griff. Notul.* iv. 265.—Burma.
longiflora, Lam. Encyc. iv. 534 = Rondeletia pilosa.
longifolia, DC. Prod. iv. 426 = corymbosa.
macrophylla, *DC. l. c.* 427.—Afr. trop.
madagascariensis, *Desf. Tabl.* ed. I. 112, nomen.—Madag.
maritima, G. Don, Gen. Syst. iii. 531 = stricta ?
maritima, Roth, Nov. Pl. Sp. 97 = Peplidium humifusum.
microtheca, *DC. Prod.* iv. 428 ; *Cham. & Schlecht. in Linnaea*, vi. (1831) 414.—Mexic.
mitrasacmoides, F. Muell. in Trans. Phil. Soc. N. S. W. xiv. (1881) 90 ; Census, 74 = Hedyotis mitrasacmoides.
monanthes, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 60.—Afr. trop.
multiflora, DC. Prod. iv. 427 = paniculata.
mysorensis, G. Don, Gen. Syst iii. 531 = Anotis monosperma.
noctiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 57.—Afr. trop.
nudicaulis, *Roth, Nov. Pl. Sp.* 96.—Ind. or. ; Malaya.
nudiflora, *Thunb. Fl. Javan.* 9.—Java.
obtusiloba, *Hiern, in Oliver, Fl. Trop. Afr.* iii.—Afr. trop.
ovata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 97.—Mexic.
ovatifolia, DC. Prod. iv. 427 = nudicaulis.
oxyphylla, G. Don, Gen. Syst. iii. 531 = Anotis oxyphylla.
paniculata, *Linn. Sp. Pl.* 1667.—As. trop. ; Ins. Pacif.
Parishii, *Hook. f. Fl. Brit. Ind.* iii. 70.—Burma.
parviflora, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 84.—Afr. trop.
patens, *Chapm. Fl. S. U. St. Suppl.* 625.—Am. bor.
pauciflora, Roxb. ex Wight & Arn. Prod. 415 = diffusa.
peltospermum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 53.—Afr. trop.
pentandra, DC. Prod. iv. 427 = macrophylla.
pentandra, Retz. Obs. iv. 22 = Vahlia oldenlandioides.
polyclada, F. Muell. Census, 74 = Hedyotis polyclada.
pterita, Miq. Fl. Ind. Bat. ii. 193 = alata.
pterospora, F. Muell. Census, 74 = Hedyotis pterospora.
puberula, G. Don, Gen. Syst. iii. 530 = umbellata.
pumila, DC. Prod. iv. 425 = crystallina.
purpurea, A. Gray, Man. Bot. N. U. St. ed. II. 173 = Houstonia purpurea.
pusilla, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 216 = brachiata.
radicans, Roxb. ex Wight & Arn. Prod. 413 = paniculata.
ramosa, Roxb. Hort. Beng. 11 ; Fl. Ind. i. 424 = corymbosa.
ramosissima, Hohen. ex Walp. Rep. ii. 502 = hedyotoides.
ramosissima, Hook. f. Fl. Brit. Ind. iii. 69, sphalm. in syn. = paniculata.
ramosissima, *Spreng. Mant.* i. 35.—Hab. ?
recurva, *Miq. Fl. Ind. Bat.* ii. 195.—Borneo.
repens, Burm. f. Fl. Ind. 38 = trinervia.
repens, Linn. Mant. i. 40 = Dentella repens.
retrorsa, *Boiss. Fl. Orient.* iii. 12.—Sind ; Arab.
rigida, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 55.—Afr. trop.
riparia, DC. Prod. iv. 424 = capensis ?
rotundifolia, G. Don, Gen. Syst. iii. 531 = nudicaulis.
rotundifolia, A. Gray, Man. Bot. N. U. St. ed. II. 174 = Houstonia rotundifolia.
rubioides, Miq. Fl. Ind. Bat. ii. 353 = Hedyotis capitellata.
rubra, A. Gray, Pl. Wright. ii. 68 = Houstonia rubra.
rupestris, Lam. Encyc. iv. 535 = Rachicallis rupestris.
sabulosa, DC. Prod. iv. 424 = capensis ?
Salzmanni, *Benth. & Hook. f. Gen.* ii. 58.—Bras.
scabra, Wight, in Wight & Arn. Prod. 418 = Hedyotis caerulea.
scabrida, DC. Prod. iv. 425 = corymbosa.

OLDENLANDIA :—
Schimperi, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 21.—Arab.; Afr. trop.
scleranthoides, *Benth. & Hook. f. Gen.* ii 58.—Austral.
senegalensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 56.—Ind. or.?; Afr. trop.
serpyllifolia, A. Gray, Man. Bot. N. U. St. ed. II. 174 = Houstonia serpyllifolia.
sessiliflora, Sm. in Rees, Cycl. xxv. n. 2 = Vahlia viscosa.
Shuteri, *Hook. f. Fl. Brit. Ind.* iii. 69.—Ind. or.
Sieberi, *Baker, Fl. Maurit.* 138.—Maurit.
spergulacea, DC. Prod. iv. 428 = nudicaulis.
spermacocoides, F. Muell. Census, 74 = Hedyotis spermacocoides.
Stocksii, *Hook. f. Fl. Brit. Ind.* iii. 67.—Ind. or.
stricta, *Linn. Mant.* ii. 200.—Ind. or.
stricta, Willd. ex Schult. Mant. iii. 144 = Willdenowiana.
strigulosa, *DC. Prod.* iv. 427.—Ins. Marian.
strumosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 58.—Afr. trop.
subulata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 153.—Borneo.
subviscosa, Wright, ex A. Gray, Pl. Wright. ii. 68 = Houstonia subviscosa.
succulenta, *Wright, ex Griseb. Cat. Pl. Cub.* 285.—Cuba.
tenuifolia, Burm. f. Fl. Ind. 37 = corymbosa?
tenuifolia, *Forst. f. Prod.* 11.—Ins. Tanna.
tenuissima, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 61.—Afr. trop.
Teysmanniana, Miq. Fl. Ind. Bat. ii. 354 = Anotis Wightiana.
tillaeacea, F. Muell. Census, 74 = Hedyotis tillaeacea.
trachymenioides, F. Muell. l. c. = Hedyotis trachymenoides.
triflora, *Roxb. Hort. Beng.* 11, nomen.—Ind. or.
triflora, *Thunb. Fl. Javan.* 9.—Java.
trinervia, *Retz. Obs.* iv. 23.—As. et Afr. trop.
tuberosa, Lam. Encyc. iv. 535 = Lucya tuberosa.
tubulosa, G. Don, Gen. Syst. iii. 531 = paniculata, Anotis quadrilocularis, lancifolia.
umbellata, Hort. ex Steud. Nom. ed. I. 562 = lactea.
umbellata, *Linn. Sp. Pl.* 119.—Ind. or.; Abyss.
uniflora, Linn. l. c. = glomerata.
uniflora, *Ruiz & Pav. Fl. Per.* i. 57.—Chili.
verticillata, Bacl. ex DC. Prod. iv. 572 = Mitracarpum scabrum.
verticillata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 152.—Java.
verticillata, Linn. Mant. i. 40 = Hedyotis Crataeogonum.
virgata, *DC. Prod.* iv. 425.—Afr. trop.
wauensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 64.—Afr. trop.
Wightii, *Hook. f. Fl. Brit. Ind.* iii. 66.—Ind. or.
Willdenowiana, *Schult. Mant.* iii. 144.—Madag.
Zanguebariae, Lour. Fl. Cochinch. 78 = Hedyotis Zanguebariae.

OLDFIELDIA, Benth. & Hook. f. in Hook. Kew Journ. ii. (1850) 184. t. 6. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 281 (cf. i. 999).
africana, *Benth. & Hook. f. in Hook. Kew Journ.* ii. (1850) 185.—Afr. trop.

OLEA, [Tourn.] Linn. Syst. ed. I (1735). *OLEACEAE*, Benth. & Hook. f. ii. 679.
ENAIMON, Rafin. Sylva Tellur. 9 (1838).
GYMNELAEA, Spach, Hist. Vég. Phan. viii. (1839) 258.
NESTEGIS, Rafin. Sylva Tellur. 10 (1838).
PACHYDERMA, Blume, Bijdr. 682 (1825).
PAUSIA, Rafin. Sylva Tellur. 9 (1838).
PICRICARYA, Dennst. Schluess. Hort. Malab. 30 (1818).
POGENDA, Rafin. Sylva Tellur. 10 (1838).
STEREODERMA, Blume, Fl. Jav. Praef. 9 (1828).
TETRAPILUS, Lour. Fl. Cochinch. 611 (1790).
acuminata, *Wall. Cat.* n. 2809.—Ind. or.
africana, Mill. Gard. Dict. ed. VIII. n. 4 = verrucosa.
alba, Lam. ex Steud. Nom. ed. I. 562 = europaea.
americana, Linn. Mant. i. 24 = Osmanthus americana.
amygdalina, Gouan, Herbor. 264 = europaea.
angulosa, Gouan, l. c. = europaea.
angustifolia, Rafin. Sylva Tellur. 9 = europaea.
angustifolia, Salisb. Prod. 13 = Phillyrea angustifolia.
apetala, Andr. Bot. Rep. t. 316 = Notelaea longifolia.
apetala, *Vahl, Symb. Bot.* iii. 3.—N. Zel.; Ins. Norfolk.
Aquifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 166 = Osmanthus Aquifolium.

OLEA :—
arborea, *Lodd. Cat. ex Loud. Hort. Brit.* 5, nomen.—Hab.?
argentata, Clem. ex Steud. Nom. ed. II. ii. 209 = europaea.
arolensis, Clem. ex Steud. l. c. = europaea.
asiatica, Desf. Cat. Hort. Par. ed. III. 88 = cuspidata.
atrorubens, Gouan, Herbor. 264 = europaea.
attenuata, Wall. Cat. n. 2839 = Linociera macrophylla.
bifera, Rafin. Sylva Tellur. 8 (Quid?).—Hab.?
brevifolia, Rafin. l. c. 9 (Quid?).—Hab.?
Buchanani, 'Herb.' Lamb. ex D. Don, Prod. Fl. Nep. 107 (= *Notelaea Posua*).—Nepal.
buxifolia, Mill. Gard. Dict. ed. VIII. n. 5 = capensis.
buxifolia, Steud. Nom. ed. I. 562 = europaea.
cajetana, Petagn. Inst. Bot. ii. 19 = europaea.
capensis, Buchoz, ex Roem. & Schult. Syst. v. 417 = Pterocelastrus tricuspidatus.
capensis, *Linn. Sp. Pl.* 8.—Afr. austr.
cassinifolia, Salisb. Prod. 13 = capensis.
cayana, Rafin. Sylva Tellur. 9 (Quid?).—Guian.?
cernua, Vahl, Symb. Bot. iii. 3 = obtusifolia.
chrysophylla, *Lam. Tabl. Encyc.* i. 29; *Encyc.* iv. 544.—Ins. Mascar.; Afr. trop.
clavata, G. Don, Gen. Syst. iv. 48 = Linociera intermedia.
communis, Steud. Nom. ed. I. 562 = europaea.
compacta, Wall. Cat. n. 2819 = Ligustrum compactum.
concolor, *E. Mey. Comm. Pl. Afr. Austr.* 176.—Afr. austr.
consanguinea, Hance, ex Walp. Ann. iii. 18 = Ligustrum Stauntoni.
craniomorpha, Gouan, Herbor. 264 = europaea.
Cunninghamii, *Hook. f. Fl. N. Zel.* i. 175.—N. Zel.
cuspidata, *Wall. Cat.* n. 2817.—Reg. Himal.
dentata, *Wall. l. c.* n. 2840.—Burma.
depauperata, Steud. Nom. ed. II. ii. 209 = maritima.
dioica, *Roxb. Hort. Beng.* 3; *Fl. Ind.* i. 106.—Ind. or.
divaricata, Desf. Tabl. ed. II. 63 = Phillyrea divaricata.
emarginata, Lam. Tabl. Encyc. i. 29. t. 8. f. 2 = Noronhia emarginata.
Endlicheri, *F. Muell. in Journ. Bot.* xxiii. (1885) 354.—Ins. Norfolk.
europaea, *Linn. Sp. Pl.* 8.—Reg. Mediterr.; Oriens.
europaea, Thunb. Prod. Pl. Cap. 2 = verrucosa.
exasperata, Jacq. Hort. Schoenb. iii. 1. t. 251.—Afr. austr.
exasperata, Willd. Enum. Hort. Berol. Suppl. 1 = lancea.
excelsa, Ait. Hort. Kew. ed. I. i. 14 = Notelaea excelsa.
ferruginea, Hort. ex Steud. Nom. ed. II. ii. 209 = chrysophylla.
ferruginea, Royle, Illustr. Bot. Himal. 267 = cuspidata.
ferruginea, Wall. ex Aitch. in Journ. Linn. Soc. viii. (1865) 67 = europaea.
floribunda, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 225.—Ins. Salomon.
foveolata, *E. Mey. Comm. Pl. Afr. Austr.* 176.—Afr. austr.
fragrans, Thunb. Fl. Jap. 18. t. 2 = Osmanthus fragrans.
gallica, Mill. Gard. Dict. ed. VIII. n. 1 = europaea.
Gamblei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 613.—Reg. Himal.
Gardneri, Thw. Enum. Pl. Zeyl. 188 = polygama.
glabella, Herb. Banks, ex Lowe, in Trans. Camb. Phil. Soc. vi. III. (1838), reimpr. 15 = exasperata.
glandulifera, *Desf. Cat. Hort. Par.* ed. III. 391.—Reg. Himal.
glandulosa, DC. Prod. viii. 285 = glandulifera.
grandiflora, Wall. Cat. n. 2820 = Ligustrum nepalense.
Heyneana, Wall. l. c. n. 2823 = dioica.
hispalensis, Clem. ex Steud. ed. II. ii. 209 = europaea.
hispanïca, Mill. Gard. Dict. ed. VIII. n. 2 = europaea.
humilis, *Eckl. in S. Afr. Quart. Journ.* i. (1830) 370.—Afr. austr.
humilis, Salisb. Prod. 13 = Ligustrum vulgare.
ilicifolia, Hassk. Cat. Hort. Bog. Alt. 118 = Osmanthus Aquifolium.
intermedia, Tausch, in Flora, xix. (1836) 393 = laurifolia.
japonica, Hassk. Cat. Hort. Bog. Alt. 118 = Osmanthus Aquifolium?
laeta, Salisb. Prod. 13 = americana.
laevis, *Stokes, Bot. Comment.* i. 30 (Quid?).—Ind. or.
lancea, *Lam. Tabl. Encyc.* i. 29; *Encyc.* iv. 544.—Ins. Maurit.

OLEA :—
lanceolata, *Hook. f. Fl. N. Zel.* i. 176.—N. Zel.
lancifolia, Moench, Meth. 478 = europaea.
latifolia, Salisb. Prod. 13 = Phillyrea latifolia.
laurifolia, *Lam. Tabl. Encyc.* i. 29.—Afr. austr.
ligustrina, F. Muell. Fragm. v. 20 = Ligustrum australianum.
Lindleyi, Wall. Cat. n. 6305 = Ligustrum Massalongianum.
linocieroides, Wight, Ic. t. 1241 = Linociera Wightii.
longifolia, Steud. Nom. ed. I. 563 = europaea.
Lorentii, *Hochst. in Lorent, Wanderung.* 332.—As. Min.
macrophylla, *Baker, Fl. Maurit.* 219.—Ins. Maurit.
maderiensis, Hort. Cels, ex Steud. Nom. ed. I. 563 = Notelaea excelsa.
malabarica, Kostel. Allg. Med.-Pharm. Flora, iii. 1000 = dioica?
marginata, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 330 = Osmanthus marginata.
maritima, *Wall. Cat.* n. 2813.—Malaya.
microcarpa, Lour. ex DC. Prod. viii. 288, 293 = Phillyrea Lowei.
microcarpa, *Vahl, Enum.* i. 43.—Cochinch.
montana, *Hook. f. Fl. N. Zel.* i. 176. t. 46.—N. Zel.
myrtifolia, *Wall. ex Voigt, Hort. Suburb. Calc.* 547.—Ind. or.
nigra, *Loisel. Herb. Amat.* t. 256.—Hab.?
obliqua, Steud. Nom. ed. I. 563 = europaea.
oblonga, Gouan, Herbor. 264 = europaea.
obovata, *Baker, Fl. Maurit.* 219.—Ins. Maurit.
obtusifolia, *Lam. Tabl. Encyc.* i. 28.—Madag.
odorata, Rozier, ex Roem. & Schult. Syst. i. 69 = europaea.
officinarum, Crantz, Inst. ii. 379 = europaea.
Oleaster, Hoffmgg. & Link, Fl. Port. i. 387 = europaea.
ovalis, *Miq. in Journ. Bot. Néerl.* i. (1861) 111.—China.
pallida, Salisb. Prod. 13 = europaea.
paniculata, *R. Br. Prod.* 528.—Austral.
paniculata, Roxb. Hort. Beng. [80]; Fl. Ind. i. 105 = Linociera intermedia.
pauciflora, Wall. Cat. n. 2812 = Linociera pauciflora.
polygama, *Wight, Ic.* t. 1239.—Ind. or.
polymorpha, Risso, ex Schult. Mant. i. 75 = europaea.
Posua, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 107 (= *Notelaea Posua*).—Reg. Himal.
praecox, Gouan, Herbor. 264 = europaea.
procera, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1830) 178, nomen (Quid?).—N. Guin.
racemosa, Gouan, Herbor. 264 = europaea.
regia, Rozier, ex Roem. & Schult. Syst. i. 69 = europaea.
rigida, Hort. Cels, ex Steud. Nom. ed. I. 563 = microcarpa.
robusta, Sweet, Hort. Brit. ed. I. 489; Wall. Cat. n. 2822 = Ligustrum robustum.
Roxburghiana, Schult. Mant. i. 77 = seq.
Roxburghii, Spreng. Neue Entdeck. iii. 78 = Linociera intermedia.
Roxburghii, Wall. Cat. n. 2816 E = Ligustrum neilgherrense.
salicifolia, Wall. l. c. n. 2821 = dentata.
sandwicensis, A. Gray, in Proc. Am. Acad. v. (1861–62) 331 = Osmanthus sandwicensis.
sativa, Hoffmgg. & Link, Fl. Port. i. 388 = europaea.
similis, Burch. Trav. i. 177; ii. 264 = verrucosa.
sphaerica, Gouan, Herbor. 264 = europaea.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 3 = europaea.
terniflora, Kurz, For. Fl. Brit. Burma, ii. 157 = Linociera terniflora.
Thozetii, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 183.—N. Caled; Austral.
undulata, Jacq. Hort. Schoenb. i. 1. t. 2 = laurifolia.
variegata, Gouan, Herbor. 264 = europaea.
verrucosa, *Link, Enum. Hort. Berol.* i. 33.—Afr. austr.
verrucosa, Rafin. Sylva Tellur. 8 = europaea.
viridula, Gouan, Herbor. 264 = europaea.
vitiensis, *Seem. Fl. Vit.* 155.—Ins. Fiji.
Walpersiana, Hance, ex Walp. Ann. iii. 17 = Ligustrum sinense.
Wightiana, Wall. Cat. n. 2815 = dioica.

OLEANDER, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 381 = **Nerium**, Linn. (Apocyn.).
indica, Medic. l. c. = N. odorum.
vulgaris, Medic. l. c. = N. Oleander.

OLEARIA, Moench, Meth. Suppl. 254 (1802). *COMPOSITAE*, Benth. & Hook. f. ii. 276.
EURYBIA, Cass. in Dict. Sc. Nat. xvi. 46 (1820).
HAXTONIA, Caley, ex D. Don, in Edinb. N. Phil. Journ. (Jul.-Sept. 1831) 272.
ORESTION, Rafin., Fl. Tellur. ii. 48 (1836).
SHAWIA, Forst. Char. Gen. 95. t. 48 (1776), nomen prius.
SPONGOTRICHUM, Nees, Gen. et Sp. Ast. 176 (1833)
STEETZIA, Sond. in Linnaea, xxv. (1852) 450.
STEIRACTIS, DC. Prod. v. 345 (1836).
adenolasia, *F. Muell. Fragm.* v. 67.—Austral.
adenophora, *F. Muell. l. c.* 78.—Austral.
affinis, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 109 = stellulata.
albida, *Hook. f. Handb. N. Zeal. Fl.* 128.—N. Zel.
Allomii, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 92; iii. (1871) 179.—N. Zel.
alpicola, *F. Muell. Fragm.* v. 70.—Austral.
angulata, *T. Kirk, in Trans. N. Z. Inst.* xiii. (1881) 384.—N. Zel.
angustata, Armst. l. c. 337 = oleifolia.
angustifolia, *Hook. f. Fl. N. Zel.* i. 115.—N. Zel.
argophylla, *F. Muell. Fragm.* v. 68.—Austral.
arguta, *Benth. Fl. Austral.* iii. 487.—Austral.
asterotricha, *F. Muell. Fragm.* v. 79.—Austral.
avicenniaefolia, *Hook. f. Handb. N. Zeal. Fl.* 127.—N. Zel.
axillaris, *F. Muell. Fragm.* v. 64.—Austral.
Ballii, F. Muell. l. c. viii. 143 = Aster Balli.
calcarea, *F. Muell. ex Benth. Fl. Austral.* iii. 481.—Austral.
capillaris, *J. Buch. in Trans. N. Z. Inst.* iii. (1871) 212.—N. Zel.
Cassiniae, *F. Muell. Fragm.* v. 68.—Austral.
chrysophylla, *Benth. Fl. Austral.* iii. 468.—Austral.
ciliata, *F. Muell. Fragm.* v. 79.—Austral.
Colensoi, *Hook. f. Fl. N. Zel.* i. 115.—N. Zel.
colorata, *Colenso, in Trans. N. Z. Inst.* xii. (1880) 362.—N. Zel.
conocephala, *F. Muell. Fragm.* v. 79.—Austral.
Cunninghamii, *Hook. f. Handb. N. Zeal. Fl.* 126.—N. Zel.
cyanodiscalis, F. Muell. Fragm. v. 82 = ramosissima.
cydoniaefolia, *Benth. Fl. Austral.* iii. 470.—Austral.
decurrens, *Benth. l. c.* 481.—Austral.
dentata, Hook. f. Handb. N. Zeal. Fl. 126 = macrodonta.
dentata, *Moench, Meth. Suppl.* 254.—Austral.
elliptica, *DC. Prod.* v. 271.—Austral.
excorticata, *J. Buch. in Trans. N. Z. Inst.* vi. (1874) 241.—N. Zel.
exiguifolia, *F. Muell. Fragm.* v. 67.—Austral.
exilifolia, *F. Muell. l. c.* 69.—Austral.
Ferresii, *F. Muell. l. c.* 75.—Austral.
floribunda, *Benth. Fl. Austral.* iii. 477.—Austral.
Forsteri, *Hook. f. Handb. N. Zeal. Fl.* 127.—N. Zel.
furfuracea, *Hook. f. l. c.* 125.—N. Zel.
glandulosa, *Benth. Fl. Austral.* iii. 483.—Austral.
glutescens, F. Muell. Fragm. v. 77 = glutinosa.
glutinosa, *Benth. Fl. Austral.* iii. 482.—Austral.
grandiflora, *Hook. Ic. Pl.* t. 862.—Austral.
gravis, *F. Muell. Fragm.* v. 82.—Austral.
Gunniana, Hook. f. ex Hook. Bot. Mag. t. 4638 = stellulata.
Haastii, *Hook. f. Handb. N. Zeal. Fl.* 126.—N. Zel.
Hectori, *Hook. f. l. c.* 128.—N. Zel.
heleophila, *F. Muell. Fragm.* v. 66.—Austral.
homolepis, *F. Muell. l. c.* 65.—Austral.
Hookeri, *Benth. Fl. Austral.* iii. 483.—Austral.
hygrophila, *Benth. l. c.* 474.—Austral.
ilicifolia, *Hook. f. Handb. N. Zeal. Fl.* 126.—N. Zel.
illita, F. Muell. Fragm. v. 76 = elliptica.
imbricata, *Benth. Fl. Austral.* iii. 483.—Austral.
insignis, *Hook. f. Fl. N. Zel.* ii. 331.—N. Zel.
iodochroa, *F. Muell. Fragm.* v. 81.—Austral.
lacunosa, *Hook. f. Handb. N. Zeal. Fl.* 732.—N. Zel.
ledifolia, *Benth. Fl. Austral.* iii. 472.—Austral.
lepidophylla, *Benth. l. c.* 477.—Austral.
Lyallii, *Hook. f. Fl. N. Zel.* i. 116.—N. Zel.
macrodonta, *Baker, in Gard. Chron.* (1884) i. 604.—N. Zel.
magniflora, *F. Muell. Fragm.* v. 80.—Austral.
marginata, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 321.—N. Zel.
megalophylla, *F. Muell. Fragm.* v. 70.—Austral.
Mooneyi, F. Muell. l. c. viii. 144 = Aster Mooneyi.

OLEARIA:—
moschata, *Hook. f. Handb. N. Zeal. Fl.* 127.—N. Zel.
Muelleri, *Benth. Fl. Austral.* iii. 481.—Austral.
multibracteolata, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 242.—N. Zel.
muricata, *Benth. Fl. Austral.* iii. 484.—Austral.
myrsinoides, *F. Muell. Fragm.* v. 69.—Austral.
Nernstii, *F. Muell. l. c.* 81.—Austral.
nitida, *Hook. f. Handb. N. Zeal. Fl.* 125.—N. Zel.
nummularifolia, *Hook. f. l. c.* 127.—N. Zel.
obcordata, *Benth. Fl. Austral.* iii. 471.—Tasman.
oleifolia, *T. Kirk, in Trans. N. Z. Inst.* xi. (1879) 463.—N. Zel.
oliganthema, *F. Muell. ex Benth. Fl. Austral.* iii. 469.—Austral.
operina, *Hook. f. Fl. N. Zel.* i. 114.—N. Zel.
oraria, F. Muell. Fragm. v. 78 = glutinosa.
pannosa, *Hook. Ic. Pl.* sub t. 862.—Austral.
passerinoides, *Benth. Fl. Austral.* iii. 482.—Austral.
paucidentata, *F. Muell. Fragm.* v. 66.—Austral.
persoonioides, *Benth. Fl. Austral.* iii. 471.—Tasman.
phlogopappa, DC. Prod. v. 272 = stellulata.
picridifolia, *Benth. Fl. Austral.* iii. 487.—Austral.
pimeleoides, *Benth. l. c.* 479.—Austral.
pinifolia, *Benth. l. c.* 471.—Tasman.
populifolia, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 243.—N. Zel.
quercifolia, DC. Prod. v. 272 = stellulata.
ramosissima, *Benth. Fl. Austral.* iii. 479.—Austral.
ramulosa, *Benth. l. c.* 476.—Austral.
revoluta, *F. Muell. ex Benth. l. c.* 476.—Austral.
rosmarinifolia, *Benth. l. c.* 468.—Austral.
rotundifolia, DC. Prod. v. 271 = dentata.
rudis, *F. Muell. Fragm.* v. 75.—Austral.
semidentata, *Decne. ex Hook. Fl. N. Zel.* i. 115.—N. Zel.
Siemssenii, F. Muell. Fragm. v. 71 = viscidula.
Solandri, *Hook. f. Handb. N. Zeal. Fl.* 128.—N. Zel.
Steetzii, F. Muell. Fragm. v. 66 = strigosa.
stellulata, *DC. Prod.* v. 272.—Austral.
stricta, *Benth. Fl. Austral.* iii. 485.—Austral.
strigosa, *Benth. l. c.*—Austral.
Stuartii, *F. Muell. Fragm.* v. 76.—Austral.
subspicata, *F. Muell. l. c.* 78; *Benth. Fl. Austral.* iii. 478.—Austral.
tenuifolia, *Benth. l. c.* 486.—Austral.
teretifolia, *F. Muell. Fragm.* v. 77.—Austral.
tomentosa, DC. Prod. v. 252 = dentata.
Traillii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 372.—N. Zel.
Traversii, *F. Muell. ex J. Buch. l. c.* vii. (1875) 337.—N. Zel.
tubuliflora, *Benth. Fl. Austral.* iii. 475.—Austral.
Turczaninovii, F. Muell. Fragm. v. 67 = imbricata.
vernicosa, F. Muell. l. c. = passerinoides.
virgata, *Hook. f. Handb. N. Zeal. Fl.* 128.—N. Zel.
viscidula, *Benth. Fl. Austral.* iii. 474.—Austral.
viscosa, *Benth. l. c.* 468.—Austral.
xerophila, *F. Muell. Fragm.* v. 76.—Austral.

OLEASTER, Heist. Syst. 11, 16 (1748) = **Hippophaë**, Linn. (Elaeagn.).

OLEOBACHIA, Hort. ex Mast. in Gard. Chron. (1880) ii. 232 = **Sterculia**, Linn.
macrophylla, Hort.—Cf. l. c. (1883) i. 510 = S. rupestris.
palustris, Hort. ex Mast. l. c. (1880) ii. 232 = S. rupestris.

OLEOXYLON, Wall. Cat. p. 157. n. 953 (1831) = **Dipterocarpus**, Gaertn.
balsamiferum, Wall. l. c. = D. alatus.

OLFA, Adans. Fam. ii. 458 (1763) = **Isopyrum**, Linn. (Ranunc.).

OLGASIS, Rafin. Fl. Tellur. ii. 51 (1836) = **Oncidium**, Sw. (Orchid.).
triquetra, Rafin. l. c. = O. triquetrum.

OLIGACOCE, Willd. ex DC. Prod. iv. 632 (1830) = **Valeriana**, Tourn.
carnosa, Willd. ex Steud. Nom. ed. II. ii. 210 = Valeriana carnosa.
hyalinorrhiza, Willd. ex DC. Prod. iv. 638 = Valeriana hyalinorrhiza.
polystachya, Willd. ex DC. l. c. 639 = Valerianopsis polystachya.

OLIGACTIS, Cass. in Dict. Sc. Nat. xxxvi. 16 (1825) = **Liabum**, Adans. (Compos.).
apodocephala, Cass. l. c. 17 = L. sessiliflorum.
nubigena, Cass. l. c. = L. nubigenum.
sessiliflora, DC. Prod. v. 96 = L. sessiliflorum.
sessilifolia, Steud. Nom. ed. II. ii. 210 = praec.
volubilis, Cass. in Dict. Sc. Nat. xxxvi. 17 = L. volubile.

OLIGACTIS, Rafin. Fl. Tellur. ii. 44 (1836) = **Sericocarpus**, Nees (Compos.).
conyzoides, Rafin. l. c. = S. conyzoides.
linifolia, Rafin. l. c. = Aster acris?
tortifolia, Rafin. l. c. = S. tortifolius.

OLIGAERION, Cass. in Dict. Sc. Nat. ii. Suppl. 75 (1816) = **Ursinia**, Gaertn. (Compos.).

OLIGANDRA, Less. in Linnaea, ix. (1834) 199 = **Chenopodium**, Linn.
atriplicoides, Less. l. c. = C. polyspermum.

OLIGANDRA, Less. Syn. Comp. 123 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 302.
HYMENOPHOLIS, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 88.
Chrysocoma, *Wedd. Chlor. And.* i. 158.—N. Granat.
lucilioides, Less. Syn. Comp. 124 = lycopodioides.
lycopodioides, *Less. l. c.*—Bras.
pachymorpha, *Wedd. Chlor. And.* i. 230.—N. Granat.

OLIGANTHEMUM, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 56 = **Pterigeron**, A. Gray (Compos.).
filifolium, F. Muell. l. c. = P. filifolium.

OLIGANTHERA, Endl. Gen. Suppl. i. 1377 (1841) = **Chenopodium**, Linn.

OLIGANTHES, Cass. in Bull. Soc. Philom. (1817) 10. *COMPOSITAE*, Benth. & Hook. f. ii. 233.
ADENOCYCLUS, Less. in Linnaea, iv. (1829) 337.
DIALESTA, H. B. & K. Nov. Gen. et Sp. iv. 45. t. 320 (1820).
ODONTOLOMA, H. B. & K. l. c. 43. t. 319 (1820).
POLLALESTA, H. B. & K. l. c. 46. t. 321 (1820).
acuminata, *Sch. Bip. in Linnaea*, xx. (1847) 502.—Venezuela.
condensata, *Sch. Bip. l. c.* 501.—Ins. Trinit.
discolor, *Sch. Bip. l. c.* 502.—N. Granat.
Karstenii, *Sch. Bip. l. c.* xxx. (1859–60) 166.—N. Granat.
Karwinskii, *Sch. Bip. l. c.* xx. (1847) 505.—Mexic.
macrophylla, *Sch. Bip. l. c.* 503.—N. Granat.
oxylepis, *Benth. in Benth. & Hook. f. Gen.* ii. 233.—Mexic.
rufescens, Sch. Bip. in Pollichia, xx–xxi. (1863) 338 = Eupatorium domingense.
Schomburgkii, *Sch. Bip. in Linnaea*, xx. (1847) 504.—Guian.
Spruceana, *Benth. in Benth. & Hook. f. Gen.* ii. 233.—Bras.
triflora, *Cass. in Bull. Soc. Philom.* (1818) 58; *et in Dict. Sc. Nat.* xxxvi. 19.—Madag.

OLIGARRHENA, R. Br. Prod. 549 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 615.
micrantha, *R. Br. l. c.*—Austral.

OLIGLORON, Rafin. Sylva Tellur. 109 (1838) = **Capparis**, Tourn.
zeylanica, Rafin. l. c. = C. acuminata.

OLIGOCARPHA, Cass. in Bull. Soc. Philom. (1817) 151 = **Brachylaena**, R. Br. (Compos.).
nereifolia, Cass. in Dict. Sc. Nat. xxxvi. 22 = B. nereifolia.

OLIGOCARPUS, Less. Syn. Comp. 90 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 455.
XENISMIA, DC. Prod. v. 509 (1836).
calendulaceus, *Less. Syn. Comp.* 90.—Afr. austr.
tripteroides, *DC. Prod.* vi. 455.—Afr. austr.

OLIGOCHAETA, C. Koch, in Linnaea, xvii. (1843) 42 = **Volutarella**, Cass. (Compos.).
divaricata, C. Koch, l. c. 43 = V. divaricata.

OLIGODORA, DC. Prod. vi. 282 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 342.
dentata, *DC. l. c.* 283.—Afr. austr.

OLIGODORELLA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 187 = **Marasmodes**, DC. (Compos.).
teretifolia, Turcz. l. c. 188 = M. polycephalus ?

OLIGOGLOSSA, DC. Prod. vi. 76 (1837) = **Phymaspermum**, Less. (Compos.).
acicularis, DC. l. c. = P. aciculare.

OLIGOGYNE, DC. Prod. v. 629 (1836) = **Blainvillea**, Cass. (Compos.).
bahiensis, DC. l. c. = B. bahiensis.
Burchelli, Hook. Ic. Pl. t. 101 = B. rhomboidea.
megapotamica, DC. Prod. v. 629 = B. biaristata.
synedrelloides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 316 = B. biaristata.
tampicana, DC. Prod. v. 629 = B. tampicana.

OLIGOGYNIUM, Engl. Bot. Jahrb. iv. (1883) 64 = **Nephthytis**, Schott (Aroid.).
Poissoni, Engl. l. c. 65.—Afr. trop.

OLIGOLEPIS, Cass. ex DC. Prod. v. 370, in syn. (1836) = **Sphaeranthus**, Vaill. (Compos.).
amaranthoides, Wight, Ic. t. 1149 = S. amaranthoides.
angustifolia, Steetz, in Peters, Reise Mossamb. Bot. 417 = S. Steetzii.
indicus, Cass. ex Steud. Nom. ed. II. ii. 210 = S. indicus.
Sprunnera, Steetz, in Peters, Reise Mossamb. Bot. 418 = S. nubicus.

OLIGOMERIS, Cambess. in Jacquem. Voy. Bot. 23. t. 25 (1838). *RESEDACEAE*, Benth. & Hook. f. i. 112.
DIPETALIA, Rafin. Fl. Tellur. iii. 73 (1836), nomen prius.
ELLIMIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 125 (1838).
HOLOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 51.
RESEDELLA, Webb & Berth. Phyt. Canar. i. 107. t. 9 ('1835–60').
Burchellii, *Harv. in Harv. & Sond. Fl. Cap.* i. 65.—Afr. austr.
capensis, *Harv. l. c.* 64.—Afr. austr.
dipetala, Muell. Arg. in DC. Prod. xvi. II. 585 = capensis.
dipetala, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 330 = glaucescens.
dispersa, Muell. Arg. in Neue Denkschr. Schw. Ges. Zuerich, xvi. (1858); Monog. Resed. 214 = glaucescens.
Dregeana, *Muell. Arg. l. c.* 216.—Afr. austr.
Ellimia, Webb, Fragm. Fl. Aethiop. 27 = glaucescens.
glaucescens, *Cambess. in Jacquem. Voy. Bot.* 24. t. 25.—Afr. bor.; As. or.; Calif.; N. Mexic.
Resedella, Webb, Fragm. Fl. Aethiop. 26 = glaucescens.
spathulata, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 65.—Afr. austr.
subulata, Webb, Fragm. Fl. Aethiop. 26; Boiss. Fl. Orient. i. 435 = glaucescens.

OLIGOPHOLIS, Wight, Ic. iv. 5. t. 1422 (1850) = **Christisonia**, Gardn. (Orobanch.).
tubulosa, Wight, l. c. = C. tubulosa.

OLIGORON, Rafin. New Fl. Am. iv. 60 (1836) = **Asclepias**, Linn.
longifolium, Rafin. l. c. = A. angustifolia.
tenuifolium, Rafin. l. c. = praec. ?

OLIGOSCIAS, Seem. Journ. Bot. iii. (1865) 179 = **Panax**, Linn. (Araliac.).
madagascariensis, Seem. l. c. = P. Maralia ?

OLIGOSMA, Salisb. Gen. Pl. Fragm. 85 (1866) = **Nothoscordum**, Kunth (Liliac.).
bivalve, Salisb. l. c. 86 = N. striatum.

OLIGOSMILAX, Seem. Journ. Bot. vi. (1868) 258. t. 83 = **Heterosmilax**, Kunth (Liliac.).
Gaudichaudiana, Seem. l. c. = H. Gaudichaudiana.

OLIGOSPORUS, Cass. in Bull. Soc. Philom. (1817) 33 = **Artemisia**, Linn. (Compos.).
adfinis, Less. in Linnaea, ix. (1834) 164, 190 = A. campestris.
alpinus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 75 = A. campestris.

OLIGOSPORUS:—
argyreus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 75 = Artemisia campestris.
brachyphyllus, Jord. & Fourr. l. c. 78 = A. campestris.
brevicaulis, Jord. & Fourr. l. c. 76 = A. campestris.
campestris, Cass. in Bull. Soc. Philom. (1817) 33 = A. campestris.
collinus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 80 = A. campestris.
condimentarius, Cass. in Dict. Sc. Nat. xxxvi. 26 = A. Dracunculus.
delphinensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 77 = A. campestris.
emarginatus, Cass. in Ann. Sc. Nat. Sér. I. v. (1825) 104 = Abrotanella emarginata.
eoythrocladus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 78 = A. campestris.
floribundus, Jord. & Fourr. l. c. 81 = A. campestris.
fuscatus, Jord. & Fourr. l. c. 79 = A. campestris.
glutinosus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 89 = A. glutinosa.
griseus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 80 = A. campestris.
implexus, Jord. & Fourr. l. c. 81 = A. campestris.
laxatus, Jord. & Fourr. l. c. 76 = A. campestris.
littoreus, Jord. & Fourr. l. c. 82 = A. campestris.
Marschallianus, Less. in Linnaea, ix. (1834) 154, 191 = A. campestris.
mexicanus, Less. Syn. Comp. 264 = A. mexicana.
monospermus, Decne. ex DC. Prod. vi. 95 = A. Delileana.
monspeliensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 82 = A. glutinosa.
monticolus, Jord. & Fourr. l. c. 77 = A. campestris.
orophilus, Jord. & Fourr. l. c. 76 = A. campestris.
parvulus, Jord. & Fourr. l. c. 74 = A. campestris.
pubescens, Jord. & Fourr. l. c. 81 = A. campestris.
pycnocephalus, Less. in Linnaea, vi. (1831) 524 = A. pycnocephala.
pyramidatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 83 = A. campestris.
scoparius, Less. in Linnaea, ix. (1834) 191 = A. scoparia.
stenocladus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 79 = A. campestris.
suberectus, Jord. & Fourr. l. c. 77 = A. campestris.
subsericeus, Jord. & Fourr. l. c. 74 = A. campestris.
tenuifolius, Jord. & Fourr. l. c. 79 = A. campestris.
virescens, Jord. & Fourr. l. c. 80 = A. campestris.
xylopodus, Jord. & Fourr. l. c. 82 = A. campestris.

OLIGOSTEMON, Benth. in Benth. & Hook. f. Gen. i. 570 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
pictus, *Benth. in Trans. Linn. Soc.* xxv. (1865) 305. t. 39.—Afr. trop.

OLIGOSTEMON, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 447 = **Meliosma**, Blume (Sabiac.).
Schlimii, Turcz. l. c.—N. Granat.

OLIGOTHRIX, DC. Prod. vi. 304 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 452.
gracilis, *DC. Prod.* vi. 304; *et in Deless. Ic. Sel.* iv. t. 57.—Afr. austr.

OLINIA, Thunb. in Roem. Arch. ii. i. 4 (1799). *LYTHRARIEAE*, Benth. & Hook. f. i. 785.
CREMASTOSTEMON, Jacq. Fragm. 68 (1809).
acuminata, Klotzsch, in Otto & Dietr. Allg. Gartenz. iv. (1836) 27 = cymosa.
capensis, Klotzsch, l. c. = cymosa.
cymosa, *Thunb. in Roem. Arch.* ii. i. 5.—Afr. austr.
micrantha, *Decne. Caract. Olinées*, 14.—Afr. austr.
Rochetiana, *A. Juss. in Comp. Rend. Par.* xxii. (1846) 812.—Abyss.

OLISBAEA, Benth. & Hook. f. Gen. i. 772 (1867) = seq.

OLISBEA, DC. Prod. iii. 31 (1828) = **Mouriria**, Juss. (Melastom.).
rhizophoraefolia, DC. l. c. = M. rhizophoraefolia.

OLISIA, Spach, Hist. Vég. Phan. ix. 166 (1840) = **Stachys**, Tourn. (Labiat.).
annua, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 136 = S. annua.
delphinensis, Fourr. l. c. = S. annua.

OLIVAEA, Sch. Bip. ex Benth. in Hook. Ic. Pl. t. 1103 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 397.
tricuspis, *Sch. Bip. l. c.*—Mexic.

OLIVERIA, Vent. Jard. Cels, t. 21 (1800). *UMBELLIFERAE*, Benth. & Hook. f. i. 890.
CALLISTROMA, Fenzl, in Flora, xxvi. (1843) 457; et in Endl. Gen. Syst. Suppl. iii. 82 (1843).
Aucheri, Jaub. & Spach, Illustr. v. 34 = decumbens.
Bruguieri, Jaub. & Spach, l. c. = decumbens.
decumbens, *Vent. Jard. Cels*, t. 21.—Persia.
orientalis, DC. Prod. iv. 234 = decumbens.

OLIVERIANA, Reichb. f. in Linnaea, xli. (1877) 111 = **Trichopilia**, Lindl. (Orchid.).
egregia, Reichb. f. l. c.—N. Granat.

OLMEDIA, Ruiz & Pav. Prod. 129. t. 28 (1794). *URTICACEAE*, Benth. & Hook. f. iii. 371.
PSEUDOLMEDIA, Karst. Fl. Columb. ii. 21. t. 111 (1862?).
affinis, Steud. ex Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 134 = Helicostylis Poeppigiana.
angustifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 30. t. 143.—Bras.
aspera, Poepp. & Endl. l. c. 31 = Poeppigiana.
aspera, *Ruiz & Pav. Syst. Veg.* 257.—Peruv.
calophylla, Poepp. & Endl. Nov. Gen. et Sp. ii. 32. t. 146 = Perebea calophylla.
cubensis, *Klotzsch, in Linnaea*, xx. (1847) 523.—Cuba.
ferruginea, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 31.—Bras.
grandifolia, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 128 = guianensis.
guianensis, *Tréc. l. c.* 129.—Guiana.
laevis, *Ruiz & Pav. Syst. Veg.* 258.—Peruv.
laurina, *Baill. Adansonia*, xi. (1875) 305.—N. Granat.
Maquira, Steud. Nom. ed. II. ii. 210 (= *Maquira guianensis*).—Guiana.
mollis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 31. t. 144.—Bras.
Poeppigiana, *Klotzsch, in Linnaea*, xx. (1847) 525.—Peruv.
Poeppigiana, Mart. in Flora, xxiv. (1841) II. Beibl. 93 = Helicostylis Poeppigiana.
rigida, *Klotzsch & Karst. in Linnaea*, xx. (1847) 524.—N. Granat.
tomentosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 32. t. 145 = Helicostylis Poeppigiana.
tovarensis, *Klotzsch & Karst. in Linnaea*, xx. (1847) 526.—N. Granat.

OLMEDIELLA, Baill. in Bull. Soc. Linn. Par. i. (1880) 253. *URTICACEAE*, Benth. & Hook. f. iii. 1224.
Cesatiana, *Baill. l. c.*—Bras. ?
ilicifolia, *Baill. l. c.*—Bras. ?

OLMEDIOPSIS, Karst. Fl. Columb. ii. 17. t. 109 (1862?) = **Pseudolmedia**, Tréc. (Urtic.).
lanceolata, Karst. l. c.—N. Granat.
obliqua, Karst. l. c.—N. Granat.

OLNEYA, A. Gray, in Mem. Am. Acad. N. S. v. (1855) 328. *LEGUMINOSAE*, Benth. & Hook. f. i. 500.
TESOTA, C. Muell. in Walp. Ann. iv. 479 (1857).
Tesota, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 328.—Mexic.

OLOFUTON, Rafin. Sylva Tellur. 108 (1838) = **Capparis**, Tourn.
racemosum, Rafin. l. c. = C. cantoniensis.

OLOPETALUM, Klotzsch, in Linnaea, x. (1835) 428 = **Monsonia**, Linn. (Geran.).
ovatum, Klotzsch, l. c. 429 = M. ovata.

OLOSTYLA, DC. Prod. iv. 440 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 66.
HOLOSTYLA, Endl. Gen. 546 (1838).
corymbosa, *DC. Prod.* iv. 440.—N. Caled.

OLSYNIUM, Rafin. New Fl. Am. i. 72 (1836) = **Sisyrinchium**, Linn. (Irid.).
grandiflorum, Rafin. l. c. = S. grandiflorum.
luteum, Rafin. Fl. Tellur. iv. 30 = S. californicum.

OLUNTOS, Rafin. Sylva Tellur. 58 (1838) = **Ficus**, Tourn. (Urtic.).
levigata, Rafin. l. c. = F. laevigata.
trigona, Rafin. l. c. = F. trigona.

OLYMPIA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 359; et Hist. Vég. Phan. v. 406 (1836) = **Hypericum**, Linn.
glauca, Spach, ll. cc. 359; 407 = H. olympicum.

OLYMPUSA, Klotzsch, in Schomb. Fauna et Fl. Brit. Gui. 1147 (1848), nomen. *ASCLEPIADEAE.*
tomentosa, Klotzsch, l. c. nomen.—Guian.

OLYNIA, Steud. Nom. ed. II. ii. 210 (1841) = **Olinia**, Thunb. (Sapot.).

OLYNTHIA, Lindl. Coll. Bot. 19, in obs. (1825) = **Eugenia**, Mich. (Myrtac.).
disticha, Lindl. l. c. = E. disticha.

OLYRA, Linn. Syst. ed. X. 1261 (1759). *GRAMINEAE*, Benth. & Hook. f. iii. 1110.
LITHACHNE, Beauv. Agrost. 135. t. 24 (1812).
MAPIRA, Adans. Fam. ii. 39 (1763).
RADDIA, Bertol. in Opusc. Sc. Bologn. iii. (1819) 410.
STREPHIUM, Schrad. in Nees, Agrost. Bras. 298 (1829).
arundinacea, H. B. & K. Nov. Gen. et Sp. i. 197 = latifolia.
axillaris, Lam. Encyc. iv. 547 = pauciflora.
brasiliensis, *Desv. Opusc.* 106.—Bras.
brasiliensis, Spreng. Syst. iv. Cur. Post. 29 = floribunda.
brevifolia, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 402.—Afr. trop.
capillata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) II. 114.—Bras.
caudata, *Trin. in Linnaea*, x. (1836) 292.—Peruv.
ciliatifolia, *Raddi, Agrost. Bras.* 19.—Bras.
corcovadensis, Wawra, Bot. Ergeb. Maxim. Bras. 180. t. 95 = glaberrima.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* i. 198.—Am. austr.
cuneatifolia, *Desv. Opusc.* 106.—Bras.
dimidiata, *Hochst. ex Steud. Syn. Pl. Gram.* 36.—Guian.
Durvillei, Doell, in Mart. Fl. Bras. ii. II. 323, in syn. = micrantha.
ecaudata, *Doell, l. c.* 326.—Guian.
fasciculata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) II. 113.—Bras.
filiformis, *Trin. l. c.* 115.—Bras.
flaccida, *Doell, in Mart. Fl. Bras.* ii. II. 326.—Bras.
floribunda, *Raddi, Agrost. Bras.* 20.—Bras.
floribunda, Trin. Gram. Panic. 250 = capillata.
glaberrima, *Raddi, Agrost. Bras.* 19.—Bras.
glauca, *Schomb. Faun. et Fl. Brit. Gui.* 1057, nomen. —Guian.
guineensis, *Steud. Syn. Pl. Gram.* 37.—Afr. trop.
hirsuta, Trin. Gram. Panic. 250 = micrantha.
humilis, *Nees, Agrost. Bras.* 304.—Bras.
latifolia, *Linn. Syst.* ed. X. 1261.—Am. trop.; Ind. occ.
longifolia, Hochst. ex Steud. Syn. Pl. Gram. 36 = obliquifolia.
longifolia, *H. B. & K. Nov. Gen. et Sp.* i. 198.—Bras.; Guian.
malaccensis, Wall. Cat. n. 3540 B = Scleria caricina.
media, *Desv. Opusc.* 106.—Bras.
micrantha, *H. B. & K. Nov. Gen. et Sp.* 199.—Am. austr.
nana, *Doell, in Mart. Fl. Bras.* ii. II. 329.—Am. trop.
obliqua, *Desv. Opusc.* 106.—Bras.
obliquifolia, *Steud. Syn. Pl. Gram.* 36.—Guian.
orientalis, *Lour. Fl. Cochinch.* 552.—Cochinch.
ovata, *Ham. Prod. Pl. Ind. Occ.* 7.—Guian.
paniculata, Sw. Obs. Bot. 347 = latifolia.
pauciflora, *Sw. Prod. Veg. Ind. Occ.* 21.—Ind. occ.
Pineti, *Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 532.—Cuba.
polypodioides, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) II. 117.—Bras.
pubescens, Raddi, Agrost. Bras. 18 = latifolia.
ramosissima, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) II. 116.—Bras.
sarmentosa, *Doell, in Mart. Fl. Bras.* ii. II. 319.—Bras.; Peruv.

OLYRA :—
scabra, *Nees, Agrost. Bras.* 306.—Bras.
scrobiculata, Schrad. ex Nees, l. c. 303 = micrantha.
semiovata, *Trin. Gram. Panic.* 249.—Bras.
strephioides, *Griseb. Cat. Pl. Cub.* 229.—Cuba.
strictiflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 510.—Mexic.
surinamensis, Hochst. ex Steud. Syn. Pl. Gram. 36 = latifolia.
sympodica, *Doell, in Mart. Fl. Bras.* ii. II. 322.—Guian.
Urvillei, Steud. Syn. Pl. Gram. 36 = micrantha.
ventricosa, Nees, Agrost. Bras. 303 = micrantha.

OLYTHIA, Steud. Nom. ed. II. ii. 210 (1841) = OLYNTHIA, Lindl. = **Eugenia**, Mich. (Myrtac.).

OMALANTHUS, A. Juss. Euph. Tent. 50. t. 16 (1824) = **Homalanthus**, A. Juss. (Euphorb.).

OMALANTHUS, Less. Syn. Comp. 260 (1832) = **Tanacetum**, Tourn. (Compos.).
camphoratus, Less. l. c. = T. camphoratum.

OMALIUM, Auct. ex Steud. Nom. ed. II. i. 774, in syn. (1840) = **Homalium**, Jacq. (Samyd.).

OMALOCLADOS, Hook. f. in Hook. Ic. Pl. t. 1128 (1873) = HOMALOCLADOS, Hook. f. = **Faramea**, Aubl. (Rubiac.).

OMALOCLINE, Cass. in Dict. Sc. Nat. xlviii. 431 (1827) = **Crepis**, Linn. (Compos.).
montana, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 321 = C. montana.
prunellaefolia, Cass. in Dict. Sc. Nat. xlviii. 432 = C. pygmaea.
pygmaea, Reichb. f. Ic. Fl. Germ. xix. 51 = C. pygmaea.
Schimperi, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 321 = C. Schimperi.
succisaefolia, Monn. Ess. Hierac. 78 = C. succisaefolia.
terglovensis, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 321 = C. hyoseridifolia.

OMALOTES, DC. Prod. vi. 83 (1837) = **Tanacetum**, Tourn. (Compos.)
camphorata, DC. l. c. 84 = T. camphoratum.

OMALOTHECA, Cass. in Dict. Sc. Nat. lvi. 218 (1828) = **Gnaphalium**, Linn. (Compos.).
Einseleana, Schultz frat. Arch. Fl. 311 = G. sylvaticum.
Hoppeana, Schultz frat. l. c. = G. Hoppeanum.
norvegica, Schultz frat. l. c. = G. norvegicum.
supina, Cass. in Dict. Sc. Nat. lvi. 218 = G. supinum.
sylvatica, Schultz frat. Arch. Fl. 311 = G. sylvaticum.

OMBROPHYTUM, Poepp. in Leipz. Literaturz. (1833) 1874 [ex Endl. Gen. 73]. *BALANOPHOREAE*, Benth. & Hook. f. iii. 239.
peruvianum, *Poepp. & Endl. Nov. Gen.* ii. 40.—Peruv.
zamioides, *Wedd. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 184. t. 10.—Peruv.

OMENTARIA, Salisb. Gen. Pl. Fragm. 87 (1866) = **Tulbaghia**, Linn. (Liliac.).
cepacea, Salisb. l. c. 88 = T. cepacea.

OMMATODIUM, Lindl. Gen. et Sp. Orch. 365 (1838) = **Pterygodium**, Sw. (Orchid.).
Volucris, Lindl. l. c. = P. Volucris.

OMOEA, Blume, Bijdr. 359 (1825) = **Saccolabium**, Blume (Orchid.).
micrantha, Blume, l. c. (= Saccolabii sp. ?).—Malaya.

OMOLOCARPUS, Neck. Elem. ii. 40 (1790) = **Nyctanthes**? Linn. (Oleac.).

OMONOIA, Rafin. Fl. Tellur. ii. 92 (1836) = **Eschscholzia**, Cham. (Papav.).
californica, Rafin. l. c. = E. californica.

OMOSCLERIA, Nees, in Mart. Fl. Bras. ii. I. 180 (1842) = **Scleria**, Berg. (Cyperac.).
Flagellum, Nees, l. c. = S. Flagellum.
tenacissima, Nees, l. c. 181 = S. tenacissima.

OMPHACARPUS, Korth. Verh. Nat. Gesch. Bot. 192. t. 42 (1839–42) = **Grewia**, Linn. (Liliac.).
africanus, Hook. f. in Hook. Niger Fl. 237 = G. africana.
hirsutus, Korth. Verh. Nat. Gesch. Bot. 193 = G. omphacarpa.
opacus, Korth. l. c. = G. opaca.

OMPHACOMERIA, A. DC. in DC. Prod. xiv. 680 (1857). *SANTALACEAE*, Benth. & Hook. f. iii. 227.
acerba, *A. DC. l. c.* 681.—Austral.
psilotoides, *A. DC. l. c.*—Austral.

OMPHALANDRIA, P. Br. Hist. Jamaic. 335 (1756) = seq.

OMPHALEA, Linn. Syst. ed. X. 1264 (1759). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 332.
DUCHOLA, Adans. Fam. ii. 357 (1763).
HEBECOCCA, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 146.
HECATEA, Thou. Hist. Vég. Isles Fr. 27. t. 5 (1804).
OMPHALANDRIA, P. Br. Hist. Jamaic. 335 (1756).
RONNOWIA, Buchoz, Pl. Nouv. Découv. 6. t. 4 (1779).
alternifolia, *Baill. Etud. Gén. Euph.* 529.—Afr. trop.
axillaris, Sw. Prod. Veg. Ind. Occ. 95 = Phyllanthus axillaris.
biglandulosa, *Baill. Etud. Gén. Euph.* 529.—Afr. or.
brasiliensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 86. —Bras.
cardiophylla, *Hemsl. in Pharm. Journ. & Trans.* (1882) 301.—Am. centr.
cauliflora, Sw. Prod. Veg. Ind. Occ. 95 = Phyllanthus cauliflorus.
commutata, *Muell. Arg. in Linnaea*, xxxii. (1863) 86. —Ind. occ.
cordata, Sw. Prod. Veg. Ind. Occ. 95 = diandra.
diandra, *Linn. Syst.* ed. X. 1264.—Ind. occ.
diandra, Vell. Fl. Flum. x. t. 12 = Excoecaria diandra.
eglandulosa, *Vell. l. c.*—Bras.
Epistylium, Poir. Encyc. Suppl. iv. 140 = Phyllanthus cauliflorus.
frondosa, *Juss. ex Baill. Etud. Gén. Euph.* 529.—Ind. occ.
glandulata, Vell. Fl. Flum. x. t. 14 = Sapium aucuparium.
hypoleuca, *Griseb. in Goett. Nachr.* (1865) 177.—Cuba.
lactescens, *Vell. Fl. Flum.* x. t. 11.—Bras.
laevigata, Desf. Cat. Hort. Par. ed. III. 411 = triandra.
nucifera, Sw. Prod. Veg. Ind. Occ. 95 = triandra.
oleifera, *Hemsl. in Pharm. Journ. & Trans.* (1882) 301.—Am. centr.
triandra, *Linn. Syst.* ed. X. 1264.—Am. trop.
triandra, Tussac, Fl. Antill. iv. 18. t. 9 = commutata.
trichotoma, *Muell. Arg. in Linnaea*, xxxii. (1863) 86. —Cuba.
verticillata, Vell. Fl. Flum. x. t. 15 = Senefeldera multiflora.

OMPHALISSA, Salisb. Gen. Pl. Fragm. 134 (1866) = **Hippeastrum**, Herb. (Amaryll.).
aulica, Salisb. l. c. 135 = H. aulicum.
calyptrata, Salisb. l. c. = H. calyptratum.

OMPHALIUM. Roth, Enum. Pl. Germ. i. 590 (1827) = **Omphalodes**, Tourn. (Borag.).
scorpioides, Roth, l. c. 591 = O. scorpioides.
vernum, Roth, l. c. 590 = O. verna.

OMPHALOBIUM, Gaertn. Fruct. i. 217. t. 46 (1788) = **Connarus**, Linn.
africanum, DC. Prod. ii. 85 = C. africanus.
Beyrichii, Vog. ex Baker, in Mart. Fl. Bras. xiv. II. 189 = C. Beyrichii.
brasiliense, A. Rich. ex Baker, l. c. 185 = C. marginatus.
carneum, Zipp. ex Span. in Linnaea, xv. (1841) 189 = C. Spanoghei.
discolor, Sond. in Linnaea, xxiii. (1850) 24 = Cnestis natalensis.
fasciculatum, DC. Prod. ii. 86 = C. erianthus.
Gaudichaudii, DC. l. c. 85 = C. monocarpus.
indicum, Gaertn. Fruct. i. 217. t. 46 = C. monocarpus.
Lambertii, DC. Prod. ii. 85 = C. guianensis.

OMPHALOBIUM:—
lucidum, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1188, nomen.—Guian.
magicum, Mart. ex Baker, in Mart. Fl. Bras. xiv. II. 189 = Connarus Patrisii.
micranthum, Klotzch, in Schomb. Faun. & Fl. Brit. Gui. 1189, nomen.—Guian.
nervosum, G. Don, Gen. Syst. ii. 90 = Agelaea villosa.
obliquum, Presl, Epim. Bot. 207 = C. obliquus.
opacum, Klotzch, in Schomb. Faun. & Fl. Brit. Gui. 1188, nomen.—Guian.
ovatifolium, Mart. Herb. Fl. Bras. n. 1266 = C. Blanchetii.
Patrisii, DC. Prod. ii. 86 = C. Patrisii.
pentagonum, DC. l. c. = Agelaea Lamarckii.
Perrottetii, DC. l. c. = C. Perrottetii.
pictum, Blanco, Fl. Filip. ed. II. 271 = C. monocarpus.
pinnatum, DC. Prod. ii. 86; Wall. Cat. n. 8532 = C. monocarpus.
rubrum, Poepp. & Endl. Nov. Gen. et Sp. iii. 76. t. 288 = C. ruber.
scandens, Boj. Hort. Maurit. 84.—Afr. trop.
Smeathmanni, DC. Prod. ii. 86 = C. Smeathmanni.
Thonningii, DC. l. c. = C. floribundus.
turgidum, Wall. Cat. n. 8540; ex Planch. in Linnaea, xxiii. (1850) 436 = C. turgidus.
villosum, DC. Prod. ii. 86 = Agelaea villosa.

OMPHALOBIUM, Jacq. ex DC. Prod. ii. 508 (1825); et Eclog. Pl. ii. 6. t. 126 (1844) = **Schotia**, Jacq. (Legumin.).
Schotia, Jacq. ll. cc. = S. latifolia.

OMPHALOCARPUM, Beauv. Fl. Owar. i. 6. t. 5, 6 (1805). *SAPOTACEAE*, Benth. & Hook. f. ii. 653 (cf. i. 185).
elatum, *Miers, in Trans. Linn. Soc.* Ser. II. i. (1875) 16.—Afr. trop.
procerum, *Beauv. Fl. Owar.* i. 6. t. 5, 6.—Afr. trop.

OMPHALOCARYON, Klotzsch, in Linnaea, xii. (1838) 243 = **Scyphogyne**, Brongn. (Ericac.).
capitatum, Klotzsch, l. c. 244 = S. capitata.
eglandulosum, Klotzsch, l. c. = S. eglandulosa.
muscosum, Klotzsch, l. c. 243 = S. inconspicua.

OMPHALOCOCCA, Willd. ex Schult. Mant. iii. 10 (1827) = **Aegiphila**, Jacq. (Verbenac.).
cornifolia, Willd. l. c. 132 = A. cornifolia.

OMPHALODES, Tourn. ex Moench, Meth. 419 (1794). *BORAGINEAE*, Benth. & Hook. f. ii. 847. OMPHALIUM, Roth, Enum. Pl. Germ. i. 590 (1827). PICOTIA, Roem. & Schult. Syst. iv. 10 (1819).
aliena, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 377.—Mexic.
amplexicaulis, *Lehm. in Ges. Naturf. Fr. Berl. Mag.* viii. (1814) 98.—Hispan.
blepharolepis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 504.—China.
brassicaefolia, Sweet, Hort. Brit. ed. I. 293 = amplexicaulis.
cappadocica, DC. Prod. x. 161 = cornifolia.
cardiophylla, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 377.—Mexic.
cariensis, Boiss. Diagn. Ser. I. iv. 41 = Paracaryum myosotoides.
controversa, *DC. Prod.* x. 168.—Sibir.
cornifolia, *Lehm. in Ges. Naturf. Fr. Berl. Mag.* viii. (1814) 97. t. 5.—As. Min.
cristata, Schrank, in Denkschr. Muench. Akad. iii. (1812) 221 = Paracaryum cristatum.
diffusa, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 504.—China.
Fortisii, *G. Don, Gen. Syst.* iv. 352.—Europ.
glochidiata, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 413 = Paracaryum micranthum.
hirsuta, DC. Prod. x. 159 = Paracaryum hirsutum.
Howardi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 263.—Am. bor. occ.
Icumae, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 453.—Japon.
intermedia, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 255 = Paracaryum micranthum.
intermedia, Schimp. ex DC. Prod. x. 159 = papillosa.
japonica, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 452.—Japon.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 452.—Japon.

OMPHALODES:—
linifolia, *Moench, Meth.* 419.—Europ. austr.
linophyllum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
littoralis, *Lehm. in Ges. Naturf. Fr. Berl. Mag.* viii. (1814) 98.—Gallia.
longiflora, DC. Prod. x. 158 = Lindelofia spectabilis.
Luciliae, *Boiss. Diagn.* Ser. I. iv. 41.—Graec.; As. Min.
lusitanica, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1864) 28 = nitida.
lusitanica, Schrank, in Denkschr. Muench. Akad. iii. (1812) 221 = amplexicaulis.
micrantha, DC. Prod. x. 159 = Paracaryum micranthum.
myosotoides, Fresen. in Mus. Senckenb. i. (1834) 170 = Paracaryum rugulosum.
myosotoides, Schimp. ex DC. Prod. x. 160 = rugulosa.
myosotoides, Schrank, in Denkschr. Muench. Akad. iii. (1812) 222 = Paracaryum myosotoides.
nana, A. Gray, in Proc. Am. Acad. xx. (1885) 263 = Eritrichium nanum.
nervosa, Edgew. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 158 = Cynoglossum nervosum.
nitida, *Hoffmgg. & Link, Fl. Port.* i. 194. t. 25.—Lusitan.
papillosa, *DC. Prod.* x. 159.—As. Min.
Pavoniana, *Boiss. Diagn.* Ser. I. xi. 128.—Hispan.
persica, Boiss. l. c. viii. 30 = Paracaryum rugulosum.
physodes, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1847) 413 = Paracaryum micranthum.
pontica, C. Koch, in Linnaea, xxii. (1849) 646 = Paracaryum ponticum.
repens, Schrank, in Denkschr. Muench. Akad. iii. (1812) 221 = verna.
rugulosa, *DC. Prod.* x. 160.—As. Min.
rupestris, *Rupr. ex Boiss. Fl. Orient.* iv. 267.—Reg. Caucas.
scorpioides, *Schrank, in Denkschr. Muench. Akad.* iii. (1812) 222.—Europ.
sempervirens, D. Don, Prod. Fl. Nep. 101 = Anchusa sempervirens.
sericea, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 453.—Japon.
stricta, C. Koch, in Linnaea, xvii. (1843) 302 = Paracaryum hirsutum.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 155.—Reg. Himal.
trichocarpa, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 500.—China.
verna, *Moench, Meth.* 420.—Europ.
Wittmanniana, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 607 = cornifolia.

OMPHALOPHTHALMUM, Karst. Fl. Columb. ii. 119. t. 163 (1865?). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 767.
rubra, *Karst. l. c.*—N. Granat.

OMPHALOPUS, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 277. *MELASTOMACEAE*, Benth. & Hook. f. i. 759.
fallax, *Naud. l. c.*—Malaya.
leprosus, *Naud. l. c.*—Java.
reticulatus, Naud. l. c. = Dissochaeta inappendiculata.

OMPHALOSPORA, Bartl. Ord. 171 (1830) = **Veronica**, Tourn. (Scrophul.).
praecox, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 129 = V. praecox.
triphylla, Fourr. l. c. = N. triphyllos.

OMPHALOSTIGMA, Reichb. Nom. 132 (1841) = **Leianthus**, Griseb. (Gentian.).

OMPHALOTHALMA, Pritz. Ic. Ind. ii. 198 (1866), sphalm. = **Omphalophthalmum**, Karst. (Asclep.).

OMPHALOTHECA, Hassk. in Bull. Congr. Bot. Amst. 1865 (1866) 103, 105 = **Commelina**, Linn.

OMPHALOTRIX, Maxim. Prim. Fl. Amur. 208. t. 10 (1859). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 977.
longipes, *Maxim. l. c.* 209.—Manchuria.

ONAGRA, [Tourn.] Adans. Fam. ii. 85 (1763) = **Oenothera**, Linn. (Onagrar.).
biennis, Scop. Fl. Carn. ed. II. i. 269 = Oenothera biennis.

ONAGRA:—
chrysantha, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 355 = Oenothera biennis.
erosa, Spach, Hist. Vég. Phan. iv. 361 = Oenothera erosa.
europaea, Spach, l. c. 359 = Oenothera biennis.
Kunthiana, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 353 = Oenothera elata.
Lehmanniana, Spach, l. c. = Oenothera erosa.
Linkiana, Spach, l. c. = Oenothera fruticosa.
media, Spach, Hist. Vég. Phan. iv. 360 = Oenothera fruticosa.
mollissima, Moench, Meth. 675 = Oenothera mollissima.
muricata, Moench, l. c. = Oenothera biennis.
nocturna, Moench, Meth. Suppl. 287 = Oenothera nocturna.
parviflora, Moench, l. c. = Oenothera biennis.
pellucida, Moench, Meth. 675 = Oenothera longiflora.
purpurea, Moench, Meth. Suppl. 288 = Oenothera purpurea.
salicifolia, Spach, Hist. Vég. Phan. iv. 361 = Oenothera elata.
sinuata, Moench, Meth. 676 = Oenothera sinuata.
spectabilis, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 352 = Oenothera Simsiana.
undulata, Moench, Meth. Suppl. 287 = Oenothera odorata.
vulgaris, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 353 = Oenothera biennis.

ONCERUM, Dulac, Fl. Hautes-Pyr. 255 (1867) = **Silene**, Linn. (Caryoph.).
acaule, Dulac, l. c. 257 = S. acaulis.
alpinum, Dulac, l. c. = S. alpestris.
ciliatum, Dulac, l. c. 256 = S. ciliata.
cuspidatum, Dulac, l. c. 257 = S. Saxifraga.
gallicum, Dulac, l. c. 256 = S. gallica.
glareosum, Dulac, l. c. 257 = S. Cucubalus.
globosum, Dulac, l. c. 258 = S. Cucubalus.
nutans, Dulac, l. c. 256 = S. nutans.
quadrifidum, Dulac, l. c. 257 = S. quadrifida.
rubellum, Dulac, l. c. = S. rubella.
rupestre, Dulac, l. c. = S. rupestris.

ONCIDIUM, Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 239. *ORCHIDEAE*, Benth. & Hook. f. iii. 562.
COPPENSIA, Dum. in Mém. Ac. Brux. ix. (1835) (Sur Maelenia) 10, in nota.
CUITLANZINA, Lindl. Orch. Scel. 15 (1826).
CYRTOCHILUM, H. B. & K. Nov. Gen. et Sp. i. 349 (1815).
LOPHIARIS, Rafin. Fl. Tellur. iv. 40 (1836).
OLGASIS, Rafin. Fl. Tellur. iv. 51 (1836).
PALUMBINA, Reichb. f. in Walp. Ann. vi. 699 (1861).
PSYCHOPSIS, Rafin. Fl. Tellur. iv. 40 (1836).
TOLUMNIA, Rafin. l. c. ii. 101 (1836).
XARITONIA, Rafin. l. c. iv. 9 (1836).
XEILYATHUM, Rafin. l. c. ii. 62 (1836).
abortivum, *Reichb. f. in Linnaea*, xxii. (1849) 847.—Venezuela.
abruptum, *Linden & Reichb. f. in Illustr. Hortic.* xxix. (1882) 98, nomen.—N. Granat.
acinaceum, *Lindl. Sert. Orch.* sub t. 48.—Peruv.
acrobotryum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 289 = Harrisoniana.
advena, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 422.—Venezuela.
aemulum, *Reichb. f. in Bonplandia*, ii. (1854) 102.—N. Granat.
alboviolaceum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 27 = incurvum.
alcicorne, *Reichb. f. in Gard. Chron.* (1872) 969.—N. Granat.
Allemanii, *Rodrig. Orch. Nov.* ii. 185.—Bras.
altissimum, Lindl. Bot. Reg. t. 1651 = Baueri.
altissimum, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 240.—Ind. occ.
amabile, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 301.—Bras.
amictum, *Lindl. Bot. Reg.* (1847) t. 66.—Bras.
amoenum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
ampliatum, *Lindl. Gen. & Sp. Orch.* 202.—Am. centr.
anceps, Reichb. f. in Walp. Ann. vi. 758 = Miltonia anceps.
andigenum, *Linden & Reichb. f. in Gard. Chron.* (1869) 416.—Ecuador?

ONCIDIUM:—
annulare, *Reichb. f. in Gard. Chron.* (1875) I. 396.—N. Granat.
anomalum, Reichb. f. in Linnaea, xxii. (1849) 845 = panchrysum.
ansiferum, *Reichb. f. in Bot. Zeit.* x. (1852) 696.—N. Granat.
anthérotes, Reichb. f. in Gard. Chron. (1879) II. 782 = Brassia antherotes.
anthocrene, *Reichb. f. in Linnaea*, xli. (1877) 102.—N. Granat.
apterum, Beer, Prakt. Orch. 281 = Odontoglossum apterum?
armillare, *Lindl. Fol. Orch. Oncid.* 51.—Peruv.
ascendens, *Lindl. Bot. Reg.* (1842) sub t. 4.—Mexic.
atropurpureum, Beer, Prakt. Orch. 281 = Odontoglossum atropurpureum.
aurarium, *Reichb. f. in Gard. Chron.* (1884) II. 394.—Bolivia.
aureum, *Lindl. Sert. Orch.* sub t. 25.—Mexic.
auriculatum, *Schiller, ex Lindl. Fol. Orch. Oncid.* 58, nomen.—Hab.?
auriferum, *Reichb. f. in Linnaea*, xxii. (1849) 847.—N. Granat.
aurosum, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 108 = excavatum.
Baldeviamae, *Reichb. f. in Gard. Chron.* (1873) 915.—N. Granat.
Barbaceniae, *Lindl. Fol. Orch. Oncid.* 32.—Bras.
barbatum, *Lindl. Coll. Bot.* t. 27.—Am. trop.
barbatum, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 30 f. 150 = trichodes.
Barkeri, Lindl. in Bot. Reg. (1841) Misc. 81 = tigrinum.
Batemannianum, *Parment. ex Knowles & Westc. Flor. Cab.* iii. (1840) 183. t. 137.—Bras.
Baueri, *Lindl. Gen. et Sp. Orch.* 200.—Ind. occ.
Berenyce, *Reichb. f. in Bot. Zeit.* xx. (1862) 215.—Ind. occ.
bicallosum, *Lindl. Pl. Hartw.* 94.—Guatemala.
bicolor, Lindl. Bot. Reg. (1843) 66 = Martianum.
bicornutum, Hook. Bot. Mag. t. 3109 = O. Pubes.
bictoniense, Hort. ex Lindl. Fol. Orch. Oncid. 59, nomen (Odontoglossi sp.).—Hab.?
bidens, Reichb. f. in Walp. Ann. vi. 769 = Brassia bidens.
biflorum, *Rodrig. Orch. Nov.* ii. 187.—Bras.
bifolium, *Sims, Bot. Mag.* t. 1491.—Reg. Argent.
bifrons, *Lindl. in Gard. Chron.* (1857) 84.—Mexic.
Blancheti, *Reichb. f. in Linnaea*, xxii. (1849) 845.—Bras.
Bluntii, Reichb. f. in Gard. Chron. (1879) II. 489 = *Miltonia Bluntii*.
Boothianum, *Reichb. f. in Bonplandia*, ii. (1854) 14.—Venezuela.
Boydii, Hort. ex Lindl. Fol. Orch. Oncid. 41 = luridum.
brachiatum, Reichb. f. in Walp. Ann. vi. 768 = Brassia brachiata.
brachyandrum, *Lindl. Sert. Orch.* sub t. 25.—Mexic.
brachyphyllum, Lindl. Bot. Reg. (1842) sub t. 4 = O. Cebolleta.
brachypterum, *Linden & Reichb. f. in Illustr. Hortic.* xxix. (1882) 99, nomen.—N. Granat.
bracteatum, *Warsz. ex Reichb. f. in Bot. Zeit.* x. (1852) 695.—N. Granat.
Brassia, Reichb. f. in Walp. Ann. vi. 765 = Brassia maculata.
brevifolium, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
Brienianum, *Reichb. f. in Gard. Chron.* (1881) I. 40.—Paraguay.
Brunleesianum, *Reichb. f. Otia Bot. Hamb.* 87.—Bras.
brunnipetalum, *Rodrig. Orch. Nov.* ii. 190.—Bras.
bryolophotum, *Reichb. f. in Gard. Chron.* (1871) 738.—Am. centr.
caesium, Reichb. f. in Regel, Gartenfl. (1854) 75. t. 80 = Geertianum.
calanthum, *Reichb. f. in Gard. Chron.* (1870) 39.—Ecuador *vel* Peruv.
caldense, *Reichb. f. in Linnaea*, xxii. (1849) 846.—Bras.
calobulbon, *Linden & Reichb. f. in Illustr. Hortic.* xxix. (1882) 99, nomen.—Hab.?
caloglossum, *Reichb. f. in Gard. Chron.* (1885) II. 166.—Am. trop.
caminiophorum, *Reichb. f. in Bot. Zeit.* x. (1852) 857.—Venezuela.

ONCIDIUM:—
Candelabrum, Linden, Cat. n. 6 (1851) 8 = Odontoglossum coronarium.
candidum, *Lindl. Bot. Reg.* (1843) *Misc.* 56.—Guatemala.
candidum, Reichb. f. in Walp. Ann. vi. 763 = Miltonia candida.
Carderi, *Reichb. f. in Gard. Chron.* (1875) I. 748.—N. Granat.
cardiochilum, *Lindl. Fol. Orch. Oncid.* 27.—N. Granat.
cardiostigma, *Reichb. f. in Bonplandia*, ii. (1854) 91.—Mexic.
carinatum, Knowles & Westc. Flor. Cab. ii. (1838) 31 = Leochilus carinatus.
cariniferum, Beer, Prakt. Orch. 283 = Odontoglossum cariniferum.
carthaginense, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 240.—Am. trop.
castaneum, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 99, nomen.—N. Granat.
caudatum, Reichb. f. in Walp. Ann. vi. 766 = Brassia caudata.
Cavendishianum, *Batem. Orch. Mex. et Guatem.* t. 3.—Guatemala.
Cebolleta, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 240.—Bras.
centifrancum, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 13.—Hab.?
Cepula, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 833 = O. Cebolleta.
cerebriferum, *Reichb. f. in Bot. Zeit.* x. (1852) 696.—Panama.
Cervantesii, Beer, Prakt. Orch. 283 = Odontoglossum Cervantesii.
Cheiri, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 99, nomen.—N. Granat.
cheirophorum, *Reichb. f. in Bot. Zeit.* x. (1852) 695.—N. Granat.
chiriquense, Beer, Prakt. Orch. 283 = Odontoglossum chiriquense.
chrysomorphum, *Lindl. Fol. Orch. Oncid.* 54.—Venezuela.
chrysopterum, Reichb. f. ex Lindl. l. c. 48 = micropogon.
chrysopyramis, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 108.—N. Granat.
chrysormis, Reichb. f. in Gard. Chron. (1880) II. 620 = ochthodes.
chrysothyrsus, *Reichb. f. ex Warn. Sel. Orch.* Ser. II. t. 5.—Bras.
ciliatulum, Hoffmgg. ex Lindl. Fol. Orch. Oncid. 16 = barbatum.
ciliatum, Lindl. Gen. et Sp. Orch. 200 = barbatum.
ciliolatum, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 834 = barbatum.
cimiciferum, *Reichb. f. ex Lindl. Fol. Orch. Oncid.* 9.—N. Granat.
cimiferum, Reichb. f. ex Linden, in Illustr. Hortic. xxix. (1882) 99 = praec.
cinnabarinum, Reichb. f. in Walp. Ann. vi. 770 = Brassia cinnabarina.
cirrhosum, Beer, Prakt. Orch. 284 = Odontoglossum cirrhosum.
citrinum, *Lindl. Bot. Reg.* t. 1758.—Ins. Trinit.
citrosmum, Beer, Prakt. Orch. 284 = Odontoglossum citrosmum.
Clowesii, Reichb. f. in Walp. Ann. vi. 760 = Miltonia Clowesii.
cocciferum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 101.—Peruv.
cochleatum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.
coerulescens, Beer, Prakt. Orch. 285 = Odontoglossum coerulescens.
Columbae, Hort. fide Reichb. ex Lindl. Fol. Orch. Oncid. 36 = stramineum.
Columbiae, Hort. fide Reichb. l. c. = praec.
concolor, *Hook. Bot. Mag.* t. 3752.—Bras.
confragosum, Lindl. Bot. Reg. (1838) Misc. 51 = viperinum.
confusum, *Reichb. f. Xenia Orch.* i. 234.—Panama.
constrictum, Beer, Prakt. Orch. 285 = Odontoglossum constrictum.
convolvulaceum, Lindl. & Paxt. Flow. Gard. i. (1850–51) 102 = globuliferum.
cordatum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.
cornigerum, *Lindl. Bot. Reg.* t. 1542.—Bras.
coronarium, Beer, Prakt. Orch. 285 = Odontoglossum coronarium.

ONCIDIUM:—
corynephorum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.
costatum, Beer, Prakt. Orch. 285 = Odontoglossum costatum.
cosymbephorum, C. Morr. in Ann. Soc. Bot. Gand, v. (1849) 333. t. 275 = luridum.
crispum, *Lodd. Bot. Cab.* t. 1854.—Bras.
Crista-galli, *Reichb. f. in Bot. Zeit.* x. (1852) 697.—Peruv.; Mexic.
cristatum, Beer, Prakt. Orch. 283 = Odontoglossum cristatum.
crocodiliceps, *Reichb. f. in Gard. Chron.* (1885) II. 360.—Mexic.
Croesus, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 314.—Am. trop.
cruciatum, *Reichb. f. in Gard. Chron.* (1878) I. 138.—Bras.
cruciferum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 109.—N. Granat.
cryptocopis, *Reichb. f. in Gard. Chron.* (1870) 827.—N. Granat.
cucullatum, *Lindl. Sert. Orch.* sub t. 21.—Peruv.; N. Granat.
cuencanum, *Reichb. f. in Linnaea*, xli. (1877) 32.—Ecuador.
cultratum, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 384.—Peruv.
cuneatum, Lindl. Coll. Bot. sub t. 27 = luridum.
cuneatum, *Scheidw. in Otto & Dietr. Allg. Gartenz.* x. (1842) 309.—Bras.
curtum, *Lindl. Bot. Reg.* (1847) t. 68.—Bras.
cycnicolle, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 99, nomen.—N. Granat.
dactylopterum, *Reichb. f. in Gard. Chron.* (1875) I. 684.—N. Granat.
dasytyle, *Reichb. f. in Gard. Chron.* (1873) 253, 432.—Bras.
Davisii, *Reichb. f. in Linnaea*, xli. (1877) 24.—Peruv.
decipiens, Lindl. Fol. Orch. Oncid. 22 = O. Crista-galli.
deltoideum, *Lindl. Bot. Reg.* t. 2006.—Peruv.
delumbe, *Lindl. Fol. Orch. Oncid.* 48.—Guatemala.
densiflorum, Beer, Prakt. Orch. 285 = Odontoglossum densiflorum.
densiflorum, [Schiller,] ex Lindl. Fol. Orch. Oncid. 58, nomen (Quid?).—Hab.?
dentatum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 234 = micropogon.
Diadema, Linden, Cat. n. 20 (1866) = serratum?
diaphanum, Reichb. f. in Linnaea, xli. (1877) 67 (Leochili sp.).—Mexic.
diceratum, *Lindl. Fol. Orch. Oncid.* 6.—Peruv.
dichromaticum, *Reichb. f. in Bonplandia*, iii. (1855) 215.—N. Granat.
digitatum, Lindl. in Benth. Pl. Hartw. 93 = leucochilum.
dimorphum, *Regel, Ind. Sem. Hort. Petrop.* (1869) 22; *et Gartenfl.* xix. (1870) 2. t. 637. f. 4–6.—Bras.
diodon, *Reichb. f. in Gard. Chron.* (1880) II. 69.—Hab.?
dipterum, Beer, Prakt. Orch. 285 = Odontoglossum dipterum.
disciferum, *Lindl. Fol. Orch. Oncid.* 24.—Bolivia.
distans, Beer, Prakt. Orch. 285 = Odontoglossum distans.
divaricatum, *Lindl. Bot. Reg.* t. 1050.—Bras.
Donianum, *Batem. ex W. Baxt. in Lond. Hort. Brit. Suppl.* iii. 598.—Guatemala.
dubium, *André, Illustr. Hortic.* xx. (1873) 184.—Hab.?
echinatum, H. B. & K. Nov. Gen. et Sp. i. 345. t. 79 = Erycina echinata.
echinophorum, *Rodrig. Orch. Nov.* ii. 195.—Bras.
Egertonii, Beer, Prakt. Orch. 285 = Odontoglossum Egertoni.
Ehrenbergii, Beer, l. c. = Odontoglossum Ehrenbergii.
elegans, Beer, l. c. = Tetramicra rigida.
elegantissimum, *Reichb. f. in Gard. Chron.* (1877) I. 13.—Bras.
elephantotis, *Reichb. f. in Linnaea*, xli. (1877) 25.—Ecuador.
emarginatum, *G. F. W. Mey. Prim. Pl. Esseq.* 259 (sp. dub.).—Guiana.
endocharis, *Reichb. f. in Gard. Chron.* (1884) I. 206.—Hab.?
Engelii, *Reichb. f. in Linnaea*, xli. (1877) 22.—Venezuela.
ensatum, *Lindl. Bot. Reg.* (1842) *Misc.* 17.—Guatemala.

ONCIDIUM:—
epidendroides, Beer, Prakt. Orch. 285 = Odontoglossum epidendroides.
erosum, Beer, l. c. = Odontoglossum stellatum.
euodes, Reich. f. in Gard. Chron. (1880) I. 680 = Brassia euodes.
Eurycline, *Reichb. f. l. c.* (1883) II. 812.—Hab.?
euxanthinum, *Reichb. f. l. c.* (1869) 1158.—Bras.
examinans, *Lindl. Fol. Orch. Oncid.* 9.—Venezuela.
exasperatum, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 99.—N. Granat.
excavatum, *Lindl. Sert. Orch.* sub t. 25.—Peruv.
falcipetalum, *Lindl. Orch. Linden.* 14.—Venezuela.
fasciferum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 108.—Peruv.
festatum, *Reichb. ex Warsz. l. c.* 100.—Peruv.
filipes, Lindl. Sert. Orch. sub t. 48 = graminifolium.
fimbriatum, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 834 = barbatum.
fimbriatum, *Lindl. Gen. et Sp. Orch.* 199.—Bras.
flabelliferum, *Pinel, in Paxt. Mag. Bot.* xvi. (1849) 65.—Bras.
flabellulatum, *Reichb. f. & Linden, in Illustr. Hortic.* xxix. (1882) 99, nomen.—Bras.
flavescens, Reichb. f. in Walp. Ann. vi. 757 = Miltonia flavescens.
flexuosum, Lindl. Sert. Orch. sub t. 48 = cimiciferum.
flexuosum, *Sims, Bot. Mag.* t. 2203.—Bras.
Forbesii, *Hook. l. c.* t. 3705.—Bras.
Forkelii, *Scheidw. in Otto & Dietr. Allg. Gartenz.* x. (1842) 309 (sp. dub.).—Mexic.
formosissimum, *Reichb. f. in Bot. Zeit.* x. (1852) 694.—Peruv.
foveatum, *Lindl. Fol. Orch. Oncid.* 52.—Peruv.
Funcki, *Hort. Belg. ex Lubbers, Cat. Pl. Rar. San Donato* (1880), 18.—Hab.?
funereum, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 37.—Mexic.
funereum, Lindl. Gen. et Sp. Orch. 201 = reflexum.
fuscans, *Reichb. f. Otia Bot. Hamb.* 86.—Bras.
fuscatum, Reichb. f. in Walp. Ann. vi. 763 = Miltonia Warszewiczii.
Gaertneri, Beer, ex Lindl. Fol. Orch. Oncid. 58, nomen (Quid?).—Hab.?
galeatum, Beer, l. c. (Quid?).—Hab.?
galeatum, *Scheidw. in Otto & Dietr. Allg. Gartenz.* vii. (1839) 406.—Bras.
Galeottianum, Beer, Prakt. Orch. 286 = Odontoglossum Galeottianum.
gallopavinum, C. Morr. in Ann. Soc. Bot. Gand, i. (1845) 13. t. 1. = Batemannianum.
Gardneri, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 662.—Bras.
Gardnerianum, Hort.—Cf. Gard. Chron. (1880) I. 759; (1881) II. 86 = praec.
Gargantua, *Reichb. f. in Linnaea*, xli. (1877) 24.—Peruv.
Gautieri, Regel, Ind. Sem. Hort. Petrop. (1868) 80 = unicorne.
Geertianum, *C. Morr. in Ann. Soc. Bot. Gand*, iv. (1848) 55. t. 179.—Am. centr.
geraënse, *Rodrig. Orch. Nov.* ii. 193.—Bras.
Ghiesbreghtianum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 27.—Mexic.
Gireoudianum, Reichb. f. in Walp. Ann. vi. 768 = Brassia Gireoudiana.
globuliferum, *H. B. & K. Nov. Gen. et Sp.* i. 347.—N. Granat.
glossomystax, *Reichb. f. in Bot. Zeit.* x. (1852) 696.—Mexic.
glumaceum, Reichb. f. in Walp. Ann. vi. 769 = Brassia glumacea.
gracile, *Lindl. Bot. Reg.* sub t. 1920.—Bras.
graminifolium, *Lindl. Sert. Orch.* sub t. 48.—Peruv.
grandiflorum, *Reichb. f. in Gard. Chron.* (1881) I. 782.—N. Granat.
Guibertianum, A. Rich. Fl. Cub. Fanerog. ii. 244 = Lemonianum.
Gundlachii, *Wright, ex Griseb. Cat. Pl. Cub.* 266.—Cuba.
guttatum, Reichb. f. in Walp. Ann. vi. 782 = luridum.
guttulatum, *Reichb. f. ex Lindl. Fol. Orch. Oncid.* 50.—Hab.?
gyriferum, Reichb. f. in Gard. Chron. (1878) I. 558, in obs. = seq.
gyrobulbon, *Reichb. f. l. c.* (1869) 838.—Hab.?
haematochilum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 21. t. 6.—N. Granat.

ONCIDIUM:—
haematochrysum, *Reichb. f. in Linnaea*, xxii. (1849) 844.—Bras.
haematoxanthum, *Reichb. f. ex Lindl. Fol. Orch. Oncid.* 25.—Hab.?
Hallii, Beer, Prakt. Orch. 287 = Odontoglossum Hallii.
halteratum, *Lindl. Orch. Linden.* 14.—N. Granat.
Harrisonianum, *Lindl. Bot. Reg.* t. 1569.—Bras.
Hartwegii, *Lindl. in Benth. Pl. Hartw.* 151.—Peruv.; Ecuador.
hastatum, *Lindl. in Paxt. Flow. Gard.* i. (1850–51) 9. n. 2.—Am. centr.; N. Granat.
hastiferum, Hort.—Cf. Gard. Chron. (1875) I. 492 = macranthum.
hastiferum, *Reichb. f. in Bonplandia*, ii. (1854) 102.—Peruv.
hastilabium, Beer, Prakt. Orch. 287 = Odontoglossum hastilabium.
hebraicum, *Reichb. f. in Gard. Chron.* (1875) I. 780.—N. Granat.
Helenae, Reichb. f. in Walp. Ann. vi. 771 = Brassia Warszewiczii.
hemichrysum, Reichb. f. l. c. 733 = aureum.
Henchmannii, Lodd. Orch. 21; et ex Lindl. Fol. Orch. Oncid. 40 = carthaginense.
Henckenii, *Schomb. ex Lindl. l. c.* 11.—Ins. S. Doming.
herbaceum, Reichb. f. in Walp. Ann. vi. 772 = Leochilus herbaceus.
heteranthum, *Poepp. & Endl. Nov. Gen. et Sp.* i. 34. t. 60.—Peruv.
hians, *Lindl. Bot. Reg.* (1838) *Misc.* 65.—Peruv.
hieroglyphicum, *Reichb. f. & Warsz. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 274.—Peruv.
Hirundo, *Reichb. f. in Bot. Zeit.* xv. (1857) 158.—Hab.?
holochrysum, Reichb. f. in Hamb. Gartenz. xviii. (1862) 33 = onustum.
Hrubyanum, *Reichb. f. in Gard. Chron.* (1883) I. 562.—Hab.?
Huebschii, *Reichb. f. l. c.* (1885) II. 650.—Ecuador.
Huntianum, Hook. Bot. Mag. t. 3806 = carthaginense.
Huntleyanum, Beer, ex Lindl. Fol. Orch. Oncid. 58 (Quid?).—Hab.?
hyalinobulbon, *La Llave & Lex. Nov. Veg. Desc. Fasc.* ii. (*Orch. Opusc.*) 38 (Leochili sp.).—Mexic.
hydrophilum, *Rodrig. Orch. Nov.* i. 92.—Bras.
hyphaematicum, *Reichb. f. in Gard. Chron.* (1869) 814.—N. Granat.?
imbricatum, Reichb. f. in Walp. Ann. vi. 770 = Brassia intricata.
Imperatoris-Maximiliani, *Reichb. f. in Bot. Ergeb. Maxim. Bras.* 154.—Bras.
incurvum, *Barker, ex Lindl. in Bot. Reg.* (1840) *Misc.* 75; (1845) t. 64.—Mexic.
inferlobum, Hort.—Cf. Gard. Chron. (1872) 904 = superbiens.
insculptum, *Reichb. f. l. c.* (1872) 1035.—Am. trop.
Insleayi, Barker, ex Lindl. Bot. Reg. (1840) Misc. 19 = Odontoglossum Insleayi.
intermedium, Knowles & Westc. Flor. Cab. ii. (1838) 53. t. 60 = luridum.
ionodon, *Reichb. f. in Linnaea*, xli. (1877) 23.—Peruv.
ionopterum, *Reichb. f. l. c.* 20.—Peruv.
ionosmum, Hort. ex Lindl. in Gard. Chron. (1853) 726 = tigrinum.
iricolor, *Reichb. f. in Bonplandia*, ii. (1854) 91.—Hab.?
iridifolium, *H. B. & K. Nov. Gen. et Sp.* i. 344.—Am. trop.
iridifolium, Lindl. Bot. Reg. t. 1911 = O. Cristagalli.
isopterum, *Lindl. l. c.* sub t. 1920.—Bras.
ixioides, Beer, Prakt. Orch. 287 = Odontoglossum ixioides.
Jamesoni, Reichb. f. in Gard. Chron. (1877) II. 680 = seq.
Jamiesoni, *Lindl. in Lindl. & Paxt. Flow. Gard.* i. (1850–51) 104.—Peruv.
janeirense, Reichb. f. in Bonplandia, ii. (1854) 90 = longipes.
Jonesianum, *Reichb. f. in Gard. Chron.* (1883) II. 781.—Paraguay.
Josstianum, Reichb. f. in Walp. Ann. vi. 930 = Brassia Josstiana.
juncifolium, Lindl. Coll. Bot. sub t. 27 = O. Cebolleta.

ONCIDIUM:—
Jurgensenianum, Lem. Jard. Fleur. iv. (1854) t. 368, in nota = hastatum.
Kappleri, *Reichb. f. ex Lindl. Fol. Orch. Oncid.* 45.—Guiana.
Karwinskii, Lindl. Sert. Orch. sub t. 20 = Odontoglossum Karwinskii.
Keilianum, Reichb. f. in Walp. Ann. vi. 770 = Brassia Keiliana.
Kienastianum, *Reichb. f. in Gard. Chron.* (1878) I. 558.—Peruv.
Klotzschianum, Reichb. f. in Bot. Zeit. x. (1852) 695 = nebulosum.
Kramerianum, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 9.—Am. austr.
labiatum, Reichb. f. in Walp. Ann. vi. 741 = Leochilus cochlearis.
lacerum, Lindl. Bot. Reg. (1844) Misc. 30 = stipitatum.
laeve, Beer, Prakt. Orch. 288 = Odontoglossum laeve.
lamelligerum, *Reichb. f. in Gard. Chron.* (1876) II. 808.—Ecuador.
Lanceanum, *Lindl. in Trans. Hort. Soc.* N. S. ii. (1842) 100. t. 7.—Guiana.
lancifolium, *Lindl. ex Benth. Pl. Hartw.* 151.—Ecuador?
Lansbergii, *Reichb. f. in Gard. Chron.* (1876) I. 460.—Venezuela.
Lawrenceanum, Reichb. f. in Walp. Ann. vi. 766 = Brassia Lawrenceana.
Leiboldi, *Reichb. f. l. c.* 718.—Cuba.
Lemonianum, *Lindl. Bot. Reg.* t. 1789.—Cuba.
lentiginosum, *Reichb. f. in Bonplandia*, ii. (1854) 13.—Venezuela.
leopardinum, *Lindl. Fol. Orch. Oncid.* 28.—Peruv.
lepidum, *Linden & Reichb. f. in Gard. Chron.* (1870) 1053.—Ecuador.
leucochilum, *Batem. in Bot. Reg.* sub t. 1920; *et Orch. Mex. et Guatem.* t. 1.—Guatemala.
leucostomum, Hoffmgg. ex Lindl. Fol. Orch. Oncid. 36 = hians.
leucotis, *Reichb. f. in Gard. Chron.* (1879) II. 424.—N. Granat.
Lietzei, *Regel, in Act. Hort. Petrop.* vii. (1880) 387.—Bras.
Limminghii, *C. Morr. ex Lindl. Fol. Orch. Oncid.* 56.—Venezuela.
Lindeni, Hort. ex Reichb. f. fide Lindl. l. c. 36 = stramineum.
Lindenii, Lodd. ex Lindl. Sert. Orch. sub t. 48. n. 101 = luridum.
Lindleyanum, *Hort. Linden.—Cf. Gard. Chron.* (1869) 587.—Hab.?
lineoligerum, *Reichb. f. ex Warsz. in Bonplandia*, ii. (1854) 109.—Peruv.
linguiforme, *Lindl. Orch. Linden.* 14.—Venezuela.
litum, *Reichb. f. in Gard. Chron.* (1883) ii. 328.—Bras.
longicuspe, Reichb. f. Otia Bot. Hamb. 23 = Brassia longicuspis.
longifolium, Lindl. Bot. Reg. (1841) Misc. 22 = O. Cebolleta.
longipes, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 46.—Bras.?
loxense, *Lindl. l. c.* ii. (1851–52) 128.—Ecuador.
ludens, *Reichb. f. in Gard. Chron.* (1885) I. 756.—Hab.?
lunatum, *Lindl. Bot. Reg.* t. 1929.—Guiana.
luridum, Knowles & Westc. Flor. Cab. iii. (1840) 21. t. 97, partim = carthaginense.
luridum, *Lindl. in Bot. Reg.* t. 727.—Am. trop.
luteo-purpureum, Beer, Prakt. Orch. 288 = Odontoglossum luteo-purpureum.
macrantherum, Hook. Bot. Mag. t. 3845 = Leochilus oncidioides.
macranthum, *Lindl. Gen. et Sp. Orch.* 205.—Am. centr.
macronyx, *Reichb. f. Otia Bot. Hamb.* 95.—Bras.
macropetalum, Klotzsch, in Otto & Dietr. Allg. Gartenz. (1855) 274 = micropogon.
macropetalum, *Lindl. Sert. Orch.* sub t. 48.—Bras.
macropterum, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 26 = suave.
macropus, *Linden & Reichb. f. in Gard. Chron.* (1868) 125.—Ecuador.
macrum, Beer, Prakt. Orch. 289 = Odontoglossum macrum.
maculatum, Beer, l. c. = Odontoglossum maculatum.
maculatum, *Lindl. Sert. Orch.* sub t. 48.—Mexic.

ONCIDIUM:—
maculosum, *Lindl. Bot. Reg.* sub t. 1920.—Bras.
Magdalenae, *Reichb. f. in Bonplandia*, iii. (1855) 66.—Venezuela.
maizaefolium, *Lindl. Orch. Linden.* 14.—Venezuela.
mandibulare, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 279.—N. Granat.
Mandonii, *Reichb. f. Xenia Orch.* iii. 21.—Bolivia.
Marshallianum, *Reichb. f. in Gard. Chron.* (1866) 682.—Am. trop.
Martianum, *Lindl. Bot. Reg.* sub t. 1920.—Bras.
Massangei, *E. Morr. in Belg. Hortic.* xxvii. (1877) 124. t. 617.—Am. centr.
Mathieuanum, *Reichb. f. & Warsz. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 321.—Peruv.
maxillare, Beer, Prakt. Orch. 289 = Odontoglossum maxillare.
maxilligerum, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 43.—Bras.
megalophium, Beer, Prakt. Orch. 289 = Odontoglossum megalophium.
Meirax, *Reichb. f. in Bonplandia*, ii. (1854) 12.—Venezuela.
melanops, *Reichb. f. in Gard. Chron.* (1880) ii. 620.—Ecuador.
meliosmum, *Reichb. f. l. c.* (1882) i. 796.—Hab.?
membranaceum, Beer, Prakt. Orch. 289 = Odontoglossum Cervantesii.
mentigerum, *Lindl. Fol. Orch. Oncid.* 34.—Peruv.
metallicum, *Reichb. f. in Gard. Chron.* (1876) i. 394.—N. Granat.
Methonica, *Reichb. f. in Linnaea*, xli. (1877) 21.—Bolivia.
micranthum, Skinner, ex Lindl. Fol. Orch. Oncid. 31 = Hartwegii.
microchilum, *Batem. ex Lindl. Bot. Reg.* (1840) *Misc.* 82.—Guatemala.
microglossum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 233 = barbatum.
micropogon, *Reichb. f. in Bonplandia*, ii. (1854) 90.—Hab.?
microstigma, *Reichb. f. l. c.*—Hab.?
microxiphium, *Reichb. f. in Linnaea*, xli. (1877) 21.—Bolivia.
Millianum ×, Reichb. f. in Gard. Chron. (1878) ii. 364.—N. Granat.
minax, *Reichb. f. in Linnaea*, xli. (1877) 22.—Peruv.
mirabile, Reichb. f. in Bot. Zeit. x. (1852) 697 = Lockhartia mirabilis.
miserrimum, *Reichb. f. in Bonplandia*, iii. (1855) 66.—Venezuela.
nitidum, *Rodrig. Orch. Nov.* ii. 193.—Bras.
monachicum, *Reichb. f. in Gard. Chron.* (1883) i. 368.—N. Granat.
monoceras, Hook. Bot. Mag. t. 3890 = unicorne.
montanum, *Rodrig. Orch. Nov.* i. 93.—Bras.
murinum, *Reichb. f. in Linnaea*, xli. (1877) 20.—Ecuador.
mystacinum, Hort. ex Lindl. Fol. Orch. Oncid. 59, nomen (Odontoglossi sp.).—Hab.?
naevium, Beer, Prakt. Orch. 289 = Odontoglossum naevium.
nanum, *Lindl. Bot. Reg.* (1842) *Misc.* 37.—Guatemala.
nebulosum, *Lindl. l. c.* (1841) *Misc.* 81.—Guatemala.
nigratum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 78.—Guiana.
nodosum, E. Morr. Belg. Hortic. xxiv. (1874) 258. t. 12; et Cogn. & March. Pl. Ornem. ii. (1874) t. 55 = Kramerianum.
Normani, *Hort. ex Pritz. Ic. Ind.* i. 758.—*Cf. Glenny, Journ. Hort.* ii. (1840) 149, cum tab.—Hab.?
nubigenum, Lindl. Gen. et Sp. Orch. 197; et in Hook. Journ. Bot. i. (1834) 13 = cucullatum.
nudum, *Batem. in Bot. Reg.* sub t. 1994.—Venezuela.
oblongatum, *Lindl. l. c.* (1844) *Misc.* 4.—Mexic.
obovatum, *Presl, Rel. Haenk.* i. 99.—Mexic.
obryzatum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 108.—Peruv.
obsoletum, Galeotti, ex Lindl. Fol. Orch. Oncid. 41 = carthaginense.
ocanense, Reichb. f. in Walp. Ann. vi. 769 = Brassia ocanensis.
ochmatochilum, *Reichb. f. in Bot. Zeit.* x. (1852) 698.—N. Granat.
ochthodes, *Reichb. f. in Gard. Chron.* (1871) 1483.—Ecuador.
odontochilum, *Rodrig. Orch. Nov.* ii. 189.—Bras.

ONCIDIUM:—
odoratum, Beer, Prakt. Orch. 290 = Odontoglossum odoratum.
Oerstedii, Reichb. f. in Bonplandia, ii. (1854) 91 = carthaginense.
olivaceum, *H. B. & K. Nov. Gen. et Sp.* i. 347.—N. Granat.
onustum, *Lindl. Gen. & Sp. Orch.* 203; *et in Hook. Journ. Bot.* i. (1834) 7.—Panama.
orgyale, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 102.—N. Granat.
ornithocephalum, *Lindl. Fol. Orch. Oncid.* 30.—N. Granat.
ornithopodum, *Reichb. f. in Gard. Chron.* (1879) ii. 200.—Hab.?
ornithorynchum, *H. B. & K. Nov. Gen. et Sp.* i. 345. t. 80.—Mexic.
oxyacanthosmum, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 100, nomen.—N. Granat.
pachyphyllum, *Hook. Bot. Mag.* t. 3807.—Mexic.
pallidum, Lindl. Bot. Reg. (1840) Misc. 50 = Harrisonianum.
panchrysum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 267.—N. Granat.
panduriferum, H. B. & K. Nov. Gen. et Sp. i. 346. t. 82. = carthaginense.
pantherinum, Hoffmgg. ex Lindl. Fol. Orch. Oncid. 35 = Harrisonianum.
Papilio, *Lindl. Bot. Reg.* t. 910.—Ins. Trinit.
papilioniforme, *Regel, in Act. Hort. Petrop.* vi. (1879) 292.—N. Granat.
pardalis, *Reichb. f. in Bonplandia*, ii. (1854) 13.—Venezuela.
pardinum, Beer, Prakt. Orch. 291 = Odontoglossum pardinum.
pardothyrsus, *Reichb. f. & Warsz. in Bot. Zeit.* xv. (1857) 158.—Peruv.
Pastasae, *Reichb. f. in Linnaea*, xli. (1877) 21.—Ecuador.
pauciflorum, Lindl. Coll. Bot. sub t. 27 = quadripetalum.
Pavonii, *Reichb. f. in Lindl. Fol. Orch. Oncid.* 5.—Venezuela.
pectorale, *Lindl. Sert. Orch.* t. 39.—Bras.
pelicanum, Hort. Monac. ex Lindl. Bot. Reg. (1840) Misc. 89; Mart. ex Lindl. Bot. Reg. (1847) t. 70 = reflexum.
peliogramma, *Linden & Reichb. f. in Gard. Chron.* (1871) 1451.—Am. centr.
pentadactylon, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 383.—Peruv.
pentaspilum, Hoffmgg. ex Lindl. Fol. Orch. Oncid. 35 = Harrisonianum.
pentecostale, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 56.—Bras.
pergamineum, *Lindl. in Benth. Pl. Hartw.* 93.—Guatemala.
peruvianum, Reichb. f. in Walp. Ann. vi. 767 = Brassia peruviana.
Pescatorii, Beer, Prakt. Orch. 291 = Odontoglossum nobile.
Phalaenopsis, *Linden & Reichb. f. in Gard. Chron.* (1869) 416.—Ecuador.
Phalaenopsis, Linden, ex Reichb. f. l. c. in syn., nomen. (Quid?).—Hab.?
phantasmaticum, *Lem. Illustr. Hortic.* iv. (1857) *Misc.* 77.—N. Granat.
phyllochilum, Beer, Prakt. Orch. 292 = hastatum.
phylloglossum, *Reichb. f. in Gard. Chron.* (1881) i. 169.—N. Granat.
phymatochilum, *Lindl. l. c.* (1848) 139.—Mexic.
pictum, *H. B. & K. Nov. Gen. et Sp.* i. 346. t. 81.—Peruv.
picturatum, *Reichb. f. in Bonplandia*, ii. (1854) 13.—Venezuela.
Pinellianum, Lindl. in Journ. Hort. Soc. iii. (1848) Proc. 15. tab. B 17 = Batemannianum.
pirarense, *Reichb. f. in Linnaea*, xxii. (1849) 846.—Guian.
plagianthum, *Reichb. f. in Gard. Chron.* (1873) 915.—N. Granat.
planilabre, *Lindl. in Journ. Hort. Soc.* vi. (1851) 59.—Bras.
platyglossum, *Reichb. f. ex Linden, in Illustr. Hortic.* xxix. (1882) 100, nomen.—N. Granat.
plicigerum, *Reichb. f. in Gard. Chron.* (1873) 1141.—Ecuador.
polyadenium, *Lindl. Fol. Orch. Oncid.* 27.—Peruv.

ONCIDIUM:—
polychromum, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xii. (1844) 219.—Bras.
polycladium, *Reichb. f. ex Lindl. Fol. Orch. Oncid.* 47.—N. Granat.
porrigens, *Reichb. f. in Gard. Chron.* (1868) 125.—N. Granat.
praestans ×, Reichb. f. l. c. (1880) ii. 296.—Hab.?
praetextum, *Reichb. f. l. c.* (1873) 1206.—Bras.
psittacinum, Linden, ex Lindl. Fol. Orch. Oncid. 34 = maculatum.
puberum, Spreng. Syst. iv. Cur. Post. 308 = O. Pubes.
Pubes, *Lindl. Bot. Reg.* t. 1007.—Bras.
pulchellum, *Hook. Bot. Mag.* t. 2773.—Guian.; Jamaic.
pulvinatum, *Lindl. Bot. Reg.* (1838) *Misc.* 61.—Bras.
Pumilio, *Reichb. f. in Bot. Zeit.* x. (1852) 697.—Bras.
pumilum, *Lindl. Bot. Reg.* t. 920.—Bras.
pusillum, Reichb. f. in Walp. Ann. vi. 714 = iridifolium.
pygmaeum, Beer, Prakt. Orch. 292 = Odontoglossum pygmaeum.
pyramidale, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 384.—Peruv.
pyxidophorum, *Reichb. f. in Gard. Chron.* (1879) ii. 136.—Hab.?
quadricorne, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 249 = hians.
quadripetalum, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 240.—Mexic.
racemosum, Reichb. f. in Walp. Ann. vi. 771 = Solenidium racemosum.
ramiferum, Hort. ex Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 289 = Harrisonianum.
ramosissimum, Beer, Prakt. Orch. 290 = Odontoglossum ramosissimum.
ramosum, Lindl. Bot. Reg. sub t. 1920 = Batemannianum.
ramulosum, Beer, Prakt. Orch. 290 = Odontoglossum ramulosum.
raniferum, *Lindl. Bot. Reg.* sub t. 1920; (1838) t. 48.—Bras.
rariflorum, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
reflexum, *Lindl. Bot. Reg.* sub t. 1920.—Mexic.
refractum, *Reichb. f. in Bonplandia*, ii. (1854) 12.—N. Granat.
Regnellii, Reichb. f. in Walp. Ann. vi. 760 = Miltonia Regnelli.
Reichenbachii, *Lindl. Fol. Orch. Oncid.* 49.—N. Granat.
Retemeyerianum, *Reichb. f. in Bot. Zeit.* xiv. (1856) 513.—Mexic.
retusum, *Lindl. Bot. Reg.* sub t. 1920.—Peruv.
revolutum, Beer, Prakt. Orch. 290 = Odontoglossum revolutum.
Rhinoceros, *Reichb. f. in Bot. Zeit.* xiv. (1856) 514.—Hab.?
Rigbyanum, Hort. ex Paxt. Mag. Bot. xvi. (1849) 257, cum ic. = sarcodes.
rigidum, Beer, Prakt. Orch. 290 = Odontoglossum rigidum.
Rodgersii, Hort. ex Batem. in Gard. Chron. (1868) 1317; et in Journ. Hortic. (1869) 522, cum fig. = varicosum.
roseum, Beer, Prakt. Orch. 290 = Cochlioda rosea.
roseum, Lodd. Cat. ex Lindl. Fol. Orch. Oncid. 40 = carthaginense.
Rossii, Beer, Prakt. Orch. 290 = Odontoglossum apterum.
rostrans, *Reichb. f. in Gard. Chron.* (1875) i. 748.—N. Granat.
rotundatum, *Reichb. f. l. c.* (1873) 978.—Hab.?
rupestre, *Lindl. in Benth. Pl. Hartw.* 151.—Am. centr.
Russellianum, Lindl. Bot. Reg. t. 1830 = Miltonia Russelliana.
rusticum, *Linden & Reichb. f. in Gard. Chron.* (1870) 1053.—Ecuador.
saltabundum, *Reichb. f. in Flora*, lxv. (1882) 534.—N. Granat.
Saltator, *Lem. in Fl. des Serres*, iii. (1847) sub t. 237.—Bras.
salvum, *Reichb. f. in Flora*, xlviii. (1865) 278.—Ind. occ.
sanguineum, Lindl. Sert. Orch. t. 27 = carthaginense.
sarcodes, *Lindl. in Journ. Hort. Soc.* iv. (1849) 266.—Bras.

ONCIDIUM:—
Scansor, Reichb. f. in Linnaea, xxii. (1849) 844 = globuliferum.
Schillerianum, *Reichb. f. in Bonplandia*, ii. (1854) 91. —Am. trop.?
Schlimii, *Linden, ex Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 168.—Venezuela.
Sciurus, *Scheidw. in Otto & Dietr. Allg. Gartenz.* vii. (1839) 406.—Bras.
scriptum, Reichb. f. in Walp. Ann. vi. 772 = Leochilus scriptus.
sculptum, *Reichb. f. ex Linden, Cat.* n. 90 (1873) 11, nomen.—Ecuador.
Semele, *Linden, ex Reichb. f. in Gard. Chron.* (1870) 39.—Ecuador.
serpens, *Lindl. Gen. et Sp. Orch.* 204.—Peruv.
serratum, *Lindl. Sert. Orch.* sub t. 48. n. 17.—Peruv.; Ecuador.
sessile, Lindl. & Paxt. Flow. Gard. i. (1850–51) 101. t. 21 = lancifolium.
signatum, Reichb. f. in Gard. Chron. (1881) II. 6 = Brassia signata.
Skinneri, Lindl. l. c. (1855) 629 = rupestre.
spathaceum, Reichb. f. in Walp. Ann. vi. 770 = Brassia spathacea.
speciosum, Reichb. f. l. c. 761 = Miltonia cuneata.
spectabile, *Beer, Prakt. Orch.* 293; *Reichb. f. l. c.* 759. —Bras.
sphacelatum, *Lindl. Sert. Orch.* sub t. 48.—Am. centr.
sphegiferum, *Lindl. Bot. Reg.* (1843) *Misc.* 17.—Bras.
spilopterum, *Lindl. Bot. Reg.* (1844) *Misc.* 75.—Bras.
spilotanthum, Reichb. f. ex Linden, Illustr. Hortic. xxix. (1882) 100 (Quid?).—N. Granat.
splendidum, *A. Rich. ex Duch. in Journ. Soc. Hortic. Par.* viii. (1862) 50.—Guatemala.
Sprucei, *Lindl. Fol. Orch. Oncid.* 14.—Bras.
stellatum, Beer, Prakt. Orch. 293 = Miltonia flavescens.
stelligerum, *Reichb. f. in Gard. Chron.* (1873) 1398.—Mexic.
stenochilum, *Linden & Reichb. f. in Illustr. Hortic.* xxix. (1882) 100, nomen.—N. Granat.
stenopetalum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 273 = Batemannianum.
stenotis, *Reichb. f. in Linnaea*, xli. (1877) 67.—Costa Rica.
stipitatum, *Lindl. in Benth. Bot. Voy. Sulph.* 172.—Panama.
stramineum, *Batem. ex Lindl. Bot. Reg.* (1838) *Misc.* 39.—Mexic.
suave, *Lindl. l. c.* (1843) *Misc.* 16.—Mexic.
suaveolens, Reichb. f. in Walp. Ann. vi. 765 = Brassia Lanceana.
subciliatum, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 834 = barbatum.
superbiens, *Reichb. f. in Linnaea*, xxii. (1849) 843.—N. Granat.
superfluum, *Reichb. f. in Walp. Ann.* vi. 721.—Bras.
suscephalum, *Rodrig. Orch. Nov.* ii. 188.—Bras.
Suttoni, *Batem. in Lindl. Bot. Reg.* (1842) *Misc.* 4.—Am. centr.
sylvestre, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 332.—Cuba.
Tayleuri, Hort. ex Lindl. Fol. Orch. Oncid. 46 = suave.
tectum, *Reichb. f. in Gard. Chron.* (1875) I. 780.—N. Granat.
tenense, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 107.—N. Granat.
tenue, *Lindl. in Journ. Hort. Soc.* iii. (1848) 76.—Am. centr.
tenue, Lindl. l. c. vii. (1852) 271, partim = delumbe.
teretifolium, *Hort. ex Flor. & Pomol.* (1882) 90.—Hab.?
tetracopis, *Reichb. f. in Gard. Chron.* (1873) 915.—N. Granat.
tetrapetalum, Willd. Sp. Pl. iv. 112 = quadripetalum.
tetrotis, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 109.—N. Granat.
thyrsiflorum, *Rodrig. Orch. Nov.* ii. 194.—Bras.
tigratum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 109.—Peruv.
tigrinum, Hort. ex Lindl. Fol. Orch. Oncid. 33 = maculatum.

ONCIDIUM:—
tigrinum, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 36.—Mexic.
tipuloides, *Reichb. f. in Bot. Zeit.* x. (1852) 856.—Peruv.
trichodes, *Lindl. Fol. Orch. Oncid.* 15.—Bras.
tricolor, Hook. Bot. Mag. t. 4130 = quadripetalum.
tricolorum, *Beer, ex Lindl. Fol. Orch. Oncid.* 58, nomen.—Hab.?
tricuspidatum, *Reichb. f. Beitr. Orch. Centr. Am.* 72. —Costa Rica.
trifurcatum, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 384.—Peruv.
trilingue, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 42. —Peruv.
tripterygium, *Reichb. f. in Bot. Zeit.* x. (1852) 694.—Peruv.
triquetrum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 216.—Jamaic.
Trulla, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 101.—N. Granat.
trulliferum, *Lindl. Bot. Reg.* (1839) t. 57.—Bras.
tucumanense, *Reichb. f. in Linnaea*, xli. (1877) 23.—Reg. Argent.
uliginosum, *Rodrig. Orch. Nov.* i. 92.—Bras.
ulopterum, *Linden & Reichb. f. Illustr. Hortic.* xxix. (1882) 100, nomen.—Ecuador.
umbonatum, *Reichb. f. in Linnaea*, xli. (1877) 24.—Ecuador.
umbrosum, Reichb. f. in Bonplandia, ii. (1854) 12 = linguiforme.
undulatum, *Lindl. Sert. Orch.* sub t. 48.—N. Granat.
undulatum, Salisb. in Trans. Hort. Soc. i. (1812) 295 = carthaginense.
unguiculatum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xvii. (1849) 9 = concolor.
unguiculatum, Lindl. in Journ. Hort. Soc. i. (1846) 303 = tigrinum.
unicorne, *Lindl. Bot. Reg.* (1839) *Misc.* 55.—Bras.
unicornutum, *Knowles & Westc. Flor. Cab.* ii. (1838) 143.—Bras.
uniflorum, *Booth, ex Lindl. Bot. Reg.* (1843) t. 43.—Bras.
urophyllum, *Lodd. Cat.* ed. II. n. 1381; *ex Lindl. Sert. Orch.* sub t. 48. n. 90.—Bras.
usneoides, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 333.—Cuba.
ustulatum, *Reichb. f. in Gard. Chron.* (1883) I. 340. —Am. trop.
varicosum, *Lindl. Bot. Reg.* sub t. 1920.—Bras.
variegatum, Lindl. Orch. Linden. 13 = velutinum.
variegatum, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 240.—Ind. occ.
velutinum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 166. t. 33.—Cuba.
Ventilabrum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 101.—Peruv.
venustum, *C. Morr. in Hortic. Belg.* iii. (1836) 28. t. 49.—Bras.
vernixium, *Linden & Reichb. f. in Gard. Chron.* (1870) 1053.—Ecuador.
verrucosum, Reichb. f. in Walp. Ann. vi. 768 = Brassia verrucosa.
villosum, Reichb. f. l. c. 767 = Brassia villosa.
viperinum, *Lindl. Gen. et Sp. Orch.* 197.—Uruguay.
virgulatum, *Reichb. f. in Gard. Chron.* (1876) II. 452. —Ecuador.
volubile, *Linden, ex Lindl. Fol. Orch. Oncid.* 58, nomen. —Hab.?
Volvox, *Reichb. f. in Bonplandia*, ii. (1854) 13.—Venezuela.
Wageneri, Reichb. f. in Walp. Ann. vi. 767 = Brassia Wageneri.
Warmingii, *Reichb. f. in Otia Bot. Hamb.* 86.—Bras.
Warszewiczii, *Reichb. f. in Bot. Zeit.* x. (1852) 693.—N. Granat.
Warneri, *Lindl. Fol. Orch. Oncid.* 36.—Mexic.
Weddelli, *Lindl. l. c.* 39.—Bolivia.
Weltoni, Hort. Bull, ex Regel, Gartenfl. (1876) 339 = Miltonia Warszewiczii.
Wendlandianum, Reichb. f. in Bonplandia, ii. (1854) 91 = suave.
Wentworthianum, *Batem. ex Lindl. Bot. Reg.* (1840) *Misc.* 82.—Guatemala.
Widgreni, *Lindl. Fol. Orch. Oncid.* 17.—Bras.
Wrayae, Hook. Bot. Mag. t. 3854 = graminifolium.
Wydleri, *Reichb. f. in Berl. Deutsch. Bot. Ges.* iii. (1885) 276.—Porto Rico.

ONCIDIUM:—
xanthocentron, *Reichb. f. in Gard. Chron.* (1880) I. 104.—Reg. Andin.
xanthochlorum, Klotzsch, Ind. Sem. Hort. Berol. (1852) 12 = oblongatum.
xanthodon, *Reichb. f. in Gard. Chron.* (1868) 1338.—Am. austr.
xanthornis, *Reichb. f. ex Linden, Cat.* n. 21 (1867) 16. —N. Granat.
zebrinum, *Reichb. f. in Bonplandia*, ii. (1854) 12.—Venezuela.

ONCINEMA, Arn. in Edinb. N. Phil. Journ. xvii. (1834) 261 = **Glossostephanus**, E. Mey. (Asclep.).
Roxburghii, Arn. l. c. = G. linearis?

ONCINOCALYX, F. Muell. in Wing's South Sc. Record, iii. (March 1883). *VERBENACEAE.*
Betchei, *F. Muell. l. c.*—Austral.

ONCINOTIS, Benth. in Hook. Niger Fl. 451 (1849). *APOCYNACEAE*, Benth. & Hook. f. ii. 718.
hirta, *Oliver, in Hook. Ic. Pl.* t. 1230.—Afr. trop.
nitida, *Benth. in Hook. Niger Fl.* 451.—Afr. trop.
tomentella, *Radlk. in Bremen, Abh.* viii. (1883) 405. —Madag.

ONCINUS, Lour. Fl. Cochinch. 123 (1790) = **Melodinus**, Forst. (Apocyn.).
cochinchinensis, Lour. l. c. = M. monogynus?

ONCOBA, Forsk. Fl. Aegypt. Arab. 103 (1775). *BIXINEAE*, Benth. & Hook. f. i. 125, 971.
CHLANIS, Klotzsch, in Peters, Reise Mossamb. Bot. 144 (1861).
HEPTACA, Lour. Fl. Cochinch. 657 (1790).
LINDACKERIA, Presl, Rel. Haenk. ii. 89. t. 65 (1831).
LUNDIA, Schum. & Thonn. Beskr. Guin. Pl. 231 (1828).
MAYNA, Aubl. Pl. Gui. 921. t. 352 (1775).
VENTENATIA, Beauv. Fl. Owar. i. 29. t. 17 (1805).
XYLOTHECA, Hochst. in Flora, xxvi. (1843) 69.
africana, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 296.—Afr. trop.
aristata, *Oliver, Fl. Trop. Afr.* i. 118.—Afr. trop.
brachyanthera, *Oliver, l. c.* 116.—Afr. trop.
capreaefolia, *Baker, in Journ. Linn. Soc.* xxi. (1884) 320.—Madag.
dentata, *Oliver, Fl. Trop. Afr.* i. 119.—Afr. trop.
echinata, *Oliver, l. c.* 118.—Afr. trop.
glauca, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 296.—Afr. trop.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 116.—Afr. trop.
Kraussiana, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 296.—Afr. austr.
lasiocalyx, *Oliver, in Hook. Ic. Pl.* t. 1485.—Afr. trop.
latifolia, *Benth. ex Eichl. in Mart. Fl. Bras.* xiii. I. 440. —Bras.
lophocarpa, *Oliver, Fl. Trop. Afr.* i. 118.—Afr. trop.
Mannii, *Oliver, l. c.* 117.—Afr. trop.
maynensis, *Poepp. & Endl. ex Eichl. in Mart. Fl. Bras.* xiii. I. 441.—Bras.
monacantha, Steud. Nom. ed. II. ii. 212 = spinosa.
ovalis, *Oliver, Fl. Trop. Afr.* i. 118.—Afr. trop.
ovata, *Benth. ex Eichl. in Mart. Fl. Bras.* xiii. I. 441. —Bras.
pauciflora, *Benth. ex Eichl. l. c.* 442.—Bras.
Petersiana, *Oliver, Fl. Trop. Afr.* i. 116.—Afr. trop.
spinosa, *Forsk. Fl. Aegypt. Arab.* 103; *A. Rich. in Guill. & Perr. Fl. Seneg. Tent.* 32. t. 10.—Arab.
stipulata, *Oliver, in Trans. Linn. Soc.* xxix. (1872) 31. —Afr. trop.
tettensis, *Hook. f. ex Harv. & Sond. Fl. Cap.* ii. 584. —Afr. trop. et austr.
Welwitschii, *Oliver, Fl. Trop. Afr.* i. 117.—Afr. trop.

ONCOCALAMUS, Mann & H. Wendl. in Kerch. Palm. 252 (1878). *PALMAE*, Benth. & Hook. f. iii. 936.
Mannii, *H. Wendl. in Kerch. Palm.* 252.—Afr. trop.

ONCOCARPUS, A. Gray, Bot. U. St. Expl. Exped. i. 364. t. 43 (1854). *ANACARDIACEAE*, Benth. & Hook. f. i. 424.
atra, Seem. Miss. Viti, 435 = vitiensis.
Teysmannianus, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 70, nomen.—N. Guin.
vitiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 365.—Ins. Fiji.

ONCOCYCLUS, Siemss. in Bot. Zeit. iv. (1846) 706 = **Iris**, Tourn.
acutilobus, Siemss. l. c. 709 = I. acutiloba.
Helena, C. Koch, in Linnaea, xxi. (1848) 639 = I. acutiloba.
ibericus, Siemss. in Bot. Zeit. iv. (1846) 708 = I. iberica.
paradoxus, Siemss. l. c. 706 = I. paradoxa.
Sarii, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 373 = I. Sarii.
susianus, C. Koch, in Linnaea, xxi. (1848) 639 = I. susiana.

ONCODEIA, Bur. in DC. Prod. xvii. 282 (1873) (Ogcodeia) = **Noyera**, Trécul. (Urtic.).

ONCODIA, Lindl. Fol. Orch. (1853) = **Brachtia**, Reichb. f. (Orchid.).
glumacea, Lindl. l. c. = B. glumacea.

ONCOMA, Spreng. Syst. iv. Cur. Post. 11, 18 (1827) = **Oxera**, Labill. (Verbenac.).
pulchellum, Spreng. l. c. 18 = Oxera pulchella.

ONCORHIZA, Pers. Syn. i. 374 (1805) = **Oncus**, Lour. (Dioscor.).
esculentus, Pers. l. c. = Oncus esculentus.

ONCORHYNCHUS, Lehm. Ind. Sem. Hort. Hamb. (1832).—Cf. Linnaea, viii. (1833) Litt. 5 = **Orthocarpus**, Nutt. (Scrophul.).
pinnatifidus, Lehm. l. c. = Orth. australis.
tenellus, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 35.—Chili.

ONCOSPERMA, Blume, Rumphia, ii. 96. t. 82, 103 (1836). *PALMAE*, Benth. & Hook. f. iii. 895.
KEPPLERIA, Meissn. Gen. 355 (1842).
cambodianum, *Hance, in Journ. Bot.* xiv. (1876) 261.—Cambodia.
fasciculatum, *Thw. Enum. Pl. Zeyl.* 328.—Zeylan.
filamentosum, *Blume, Rumphia*, ii. 97.—Malaya.
horridum, *Scheff. in Tijdschr. Nederl. Indie*, xxxii. (1871) 191.—Sumatra.
Van-Houtteanum, H. Wendl. in Fl. des Serres, xvii. (1867–68) t. 1798 = Nephrosperma Van-Houtteana.

ONCOSPORUM, Putterl. in Endl. Nov. Stirp. Dec. 9 (1839); et Syn. Pittosp. 21 = **Marianthus**, Huegel (Pittosp.).
bicolor, Putterl. ll. cc. = M. pictus.
cuneatum, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 561 = M. cuneatus.
Drummondianum, Putterl. in Lehm. Pl. Preiss. i. 194 = M. Drummondianus.
granulatum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 366 = M. granulatus.
microphyllum, Turcz. l. c. 365 = M. microphyllus.
villosum, Turcz. l. c. = M. villosus.

ONCOSTEMA, Rafin. Fl. Tellur. ii. 13 (1836) = **Scilla**, Linn. (Liliac.).
villosa, Rafin. l. c. = S. villosa.

ONCOSTEMON, Spach, Hist. Vég. Phan. ix. 374 (1840) = seq.

ONCOSTEMUM, A. Juss. in Mém. Mus. Par. xix. (1830) 133. t. 19. *MYRSINEAE*, Benth. & Hook. f. ii. 648.
arthriticum, *Baker, in Journ. Linn. Soc.* xx. (1883) 202.—Madag.
Bojeranum, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 82.—Madag.
Capelieranum, *A. Juss. in Mém. Mus. Par.* xix. (1830) 137. t. 19.—Madag.
Commersonianum, *A. Juss. l. c.* 136. t. 19.—Madag.
divaricatum, *A. DC. in DC. Prod.* viii. 89.—Madag.
Goudotianum, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 82.—Madag.
myrtifolium, *A. DC. in DC. Prod.* viii. 90.—Madag.
nemorosum, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 82.—Madag.?
neriifolium, *Baker, in Journ. Linn. Soc.* xxi. (1885) 421.—Madag.
pauciflorum, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 82.—Madag.

ONCOSTEMUM:—
pedicellatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 202.—Madag.
phyllanthoides, *Baker, l. c.* 203.—Madag.
platycladum, *Baker, l. c.* xxi. (1885) 421.—Madag.
venulosum, *Baker, l. c.* 422.—Madag.

ONCOSTYLIS, Nees, in Mart. Fl. Bras. ii. i. 80 (1842) = **Fimbristylis**, Vahl (Cyperac.).
arenaria, Nees, l. c. 87 = F. capillaris.
bivalvis, Nees, l. c. 205.—Bras.
ciliata, Nees, l. c. 84 = F. capillaris.
circinata, Nees, l. c. 82.—Bras.
conifera, Nees, l. c. 81.—Bras.
dubia, Nees, l. c. 84 = F. capillaris.
fimbriata, Nees, l. c. 88.—Bras.
hirtella, Nees, l. c. 85.—Bras.
hispida, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 240 = F. vestita.
junciformis, Nees, in Mart. Fl. Bras. ii. i. 85.—Bras.
Kunthiana, Nees, l. c. 87 (= *Bulbostylis consanguinea*).—Bras.
paradoxa, Nees, l. c. 81 (= *Bulbostylis paradoxa*).—Bras.
pauciflora, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 241.—Mexic.
tenella, Nees, in Mart. Fl. Bras. ii. i. 82 = F. tenella.
tenuifolia, Nees, l. c. 83 = F. capillaris.
truncata, Nees, l. c. = F. capillaris.
vestita, Nees, l. c. 88, in nota = F. vestita.

ONCUFIS, Rafin. Sylva Tellur. 114 (1838) = **Cleome**, Linn. (Capparid.).

ONCUS, Lour. Fl. Cochinch. 194 (1790). *DIOSORACEAE*, Benth. & Hook. f. iii. 745.
ONCORHIZA, Pers. Syn. i. 374 (1805).
esculentus, *Lour. Fl. Cochinch.* 194.—Cochinch.

ONDETIA, Benth. in Hook. Ic. Pl. t. 1112 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
linearis, *Benth. in Hook. Ic. Pl.* t. 1112.—Afr. austr.

ONEFERA, Rafin. Fl. Tellur. iii. 30 (1836) = **Chironia**, Linn. (Gentian.).
coccinea, Rafin. l. c. = C. liniodes.

ONISTIS, Rafin. New Fl. Am. iv. 63 (1836) = **Asclepias**? Linn.
longifolia, Rafin. l. c.—Am. bor.

ONITES, Rafin. Fl. Tellur. iii. 86 (1836) = **Origanum**, Tourn. (Labiat.).
tomentosa, Rafin. l. c. = Orig. Onites.

ONIX, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378 = **Astragalus**, Tourn. (Legumin.).
galegiformis, Medic. l. c. = A. galegiformis.

ONIXOTIS, Rafin. Fl. Tellur. ii. 32 (1836) = **Dipidax**, Laws. (Liliac.).
ciliata, Rafin. l. c. = D. ciliata.
secunda, Rafin. l. c. = D. ciliata.
viridis, Rafin. l. c. = Ornithoglossum glaucum.

ONKERIPUS, Rafin. Fl. Tellur. iv. 42 (1836) = **Xylobium**, Lindl. (Orchid.).
pallidus, Rafin. l. c.—Ins. S. Vinc.

ONOBROMA, Gaertn. Fruct. ii. 380. t. 160 (1791) = **Carduncellus**, Adans. (Compos.).
arborescens, Spreng. Syst. iii. 391 = Carthamus arborescens.
arenosum, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.—Alger.
armenium, Spreng. Syst. iii. 392 = Carthamus flavescens.
atractyloides, Pomel, Nouv. Mat. Fl. Atl. 24.—Alger.
caeruleum, Gaertn. Fruct. ii. 380 = C. caeruleus.
calvum, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.—Alger.
Carduncellus, Steud. Nom. ed. II. ii. 213 = C. monspeliensium.
carlinoides, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.—Alger.
carthamoides, Pomel, l. c.—Alger.
Choulettianum, Pomel, l. c.—Alger.
coeruleum, Gaertn. Fruct. ii. t. 160 = C. caeruleus.

ONOBROMA:—
corymbosum, Spreng. Syst. iii. 393 = Cardopatium corymbosum.
cynaroides, Spreng. l. c. 392 = Cousinia cynaroides.
dentatum, Spreng. l. c. = Carthamus dentatus.
depauperatum, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.—Alger.
eriocephalum, Pomel, Nouv. Mat. Fl. Atl. 24.—Alger.
flavescens, Spreng. Syst. iii. 392 = Carthamus flavescens.
glaucum, Spreng. l. c. 393 = Carthamus glaucus.
helenioides, Spreng. l. c. 392 = Carthamus helenioides.
hispanicum, Pomel, in Bull. Soc. Bot. Fr. xvii. (1870) 235.—Alger.
lanatum, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 328 = Carthamus lanatus.
leucocaulon, Spreng. Syst. iii. 392 = Carthamus leucocaulos.
lucidum, Aucher, ex DC. Prod. vi. 553 = Cousinia lucida.
macrocephalum, C. A. Mey. Verz. Pfl. Cauc. 67 = Cousinia macrocephala.
mitissimum, Spreng. Syst. iii. 392 = Carduncellus mitissimus.
monspeliense, Spreng. l. c. = C. monspeliensium.
multifidum, Spreng. l. c. 393 = C. caeruleus.
pectinatum, Spreng. l. c. 392 = C. pectinatus.
persicum, DC. Prod. vi. 613 = Carthamus flavescens.
pinnatum, Spreng. Syst. iii. 393 = C. pinnatus.
salicifolium, Link, Enum. Hort. Berol. ii. 304 = Carlina salicifolia.
stricta, Pomel, Nouv. Mat. Fl. Atl. 278.—Alger.
tingitana, Sweet, Hort. Brit. ed. I. 216 = Carthamus caeruleus.

ONOBRUCHUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 372 = seq.

ONOBRYCHIS, Linn. Syst. ed. I (1735); Adans. Fam. ii. 327 (1763). *LEGUMINOSAE*, Benth. & Hook. f. i. 511.
ERIOCARPAEA, Bertol. in Nov. Comm. Acad. Bonon. vi. (1843) 234; (Misc. Bot. ii. 20).
SARTORIA, Boiss. Diagn. Ser. I. ix. 109 (1849).
aequidentata, Urv. in Mém. Soc. Linn. Par. i. (1822) 346 = cretica.
affinis, *Verlot, Cat. Hort. Grenoble* (1852), 11.—Hab.?
alba, Bor. Fl. Centr. Fr. ed. III. 168, in adnot. = viciaefolia.
alba, *Desv. Journ. Bot.* iii. (1814) 83.—Europ. austr.
alectorocephale, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = O. Caput-galli.
alectorolopha, St. Lag. l. c. 70 = C. Crista-galli.
Allionii, *Jord. Hort. Gratian.* (1851).—Cf. Linnaea, xxv. (1852) 304.—Gallia.
arenaria, *DC. Prod.* ii. 345.—Hungar.; Sibir.
arenaria, Koch, Syn. Fl. Germ. ed. II. 211 = gracilis.
argaea, *Boiss. & Bal. Diagn.* Ser. II. vi. 61.—Cappadoc.
argentea, *Boiss. Voy. Bot. Espagne*, ii. 188.—Hispan.
argyrea, *Boiss. Diagn.* Ser. I. ix. 107.—Cilicia.
armena, *Boiss. l. c.* Ser. II. ii. 36.—Armenia.
arnacantha, *Bunge, ex Boiss. Fl. Orient.* ii. 538.—Persia.
atropatana, *Boiss. l. c.* 541.—Armenia.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 94.—Persia.
aurantiaca, *Boiss. l. c.* Ser. I. ix. 105.—Syria.
aurea, *Boiss. ex Tchihatch. Asie Min. Bot.* i. 104, nomen.—As. Min.
Balansae, *Boiss. Diagn.* Ser. II. vi. 64.—As. Min.
Belangeri, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 76 = melanotricha.
Bergeriana, *DC. Prod.* ii. 346.—Ital.
Buhseana, *Bunge, ex Boiss. Fl. Orient.* ii. 541.—Persia.
Bungei, *Boiss. l. c.* 535.—Persia.
Buxbaumiana, Desv. Journ. Bot. iii. (1814) 81 = Pallasii.
Buxbaumiana, Desv. ex Eichw. Nov. Casp. Cauc. 23 = vaginalis.
Buxbaumii, Bieb. Fl. Taur. Cauc. ii. 179 = radiata.
cadmea, *Boiss. Diagn.* Ser. I. ii. 96.—As. Min.
candidum, Steud. Nom. ed. II. i. 730, in syn., sphalm. = Hedysarum candidum.
cappadocica, *Boiss. Fl. Orient.* ii. 550.—As. Min.
Caput-galli, *Lam. Fl. Fr.* ii. 651.—Reg. Mediterr. et Cauc.
Caput-galli, Schur, Enum. Pl. Transs. 165 = carpathica.
carnea, Boiss. et Heldr. Diagn. Ser. II. vi. 62 = ebenoides.

ONOBRYCHIS:—
carnea, Schleich. Cat. Pl. Helvet. ed. IV. 24 = arenaria.
carpathica, *DC. Prod.* ii. 346.—Mont. Carpath.
carpathica, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 784 = viciaefolia.
chorassanica, *Bunge, ex Boiss. Fl. Orient.* ii. 550.—Persia.
circinnata, *Desv. Journ. Bot.* iii. (1814) 81.—Reg. Caucas.; As. Min.
collina, Jord. Cat. Hort. Grenoble (1851).—Cf. Linnaea, xxv. (1852) 304 = viciaefolia.
comosa, Spreng. Syst. iii. 205 = Uraria crinita.
conferta, Desv. Journ. Bot. iii. (1814) 83 = viciaefolia.
conferta, Stev. ex Boiss. Fl. Orient. ii. 536 = gracilis.
conferta, Tenore, Fl. Nap. v. 128 = supina.
cornuta, *Desv. Journ. Bot.* iii. (1814) 81.—As. Min.; Persia.
cretica, *Desv. l. c.* 83. t. 23.—Europ. austr.; Persia.
crinita, *Desv. l. c.* 82.—Oriens.
crinita, Spreng. Syst. iii. 205 = Uraria crinita.
Crista-galli, Gaertn. Fruct. ii. 318. t. 148 = Gaertneriana.
Crista-galli, *Lam. Fl. Fr.* ii. 652.—Reg. Mediterr.
cristata, DC. Prod. ii. 346 = cretica.
cristata, *Pomel, Nouv. Mat. Fl. Atl.* 201.—Alger.
cuneifolia, DC. Prod. ii. 348 = Indigofera echinata.
dasycephala, *Baker, in Journ. Linn. Soc.* xviii. (1880) 48.—Afghan.
dealbata, *Stocks, in Hook. Kew Journ.* iv. (1852) 146.—Beluchist.
declivium, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1868) II. 149.—Rossia.
decumbens, Wallr. Beitr. Fl. Hercyn. 246; Jord. Cat. Hort. Grenoble (1851).—Cf. Linnaea, xxv. (1852) 305 = viciaefolia.
depauperata, *Boiss. Fl. Orient.* ii. 543.—Persia.
depressa, *Presl, Fl. Sicul.* p. xxii.—Sicilia.
diffusa, Cambess. in Jacquem. Voy. Bot. 41 = Taverniera nummularia.
ebenoides, *Boiss. & Sprun. Diagn.* Ser. I. ii. 97.—Graecia.
echinata, D. Dietr. Syn. Pl. iv. 1160 = alba.
echinata, *G. Don, Gen. Syst.* ii. 305.—Calabria.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = O. Caput-galli.
Echinus, DC. Prod. ii. 346 = O. Caput-galli?
elata, *Boiss. & Bal. Diagn.* Ser. II. vi. 62.—Graecia; As. Min.
elegans, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 265.—Turkestan.
elymaitica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 538.—Persia.
eriophora, *Desv. Journ. Bot.* iii. (1814) 82.—Reg. Mediterr. occ.
eriophora, A. Rich. Tent. Fl. Abyss. i. 206 = Richardi.
eubrychidea, *Boiss. Fl. Orient.* ii. 546.—Afghan.
expansa, Boiss. & Heldr. Diagn. Ser. I. ix. 105 = Tournefortii.
Fontanesii, G. Don, Gen. Syst. ii. 304 = Hedysarum humile.
foveolata, DC. Prod. ii. 346 = cretica.
Gaertneriana, *Boiss. Diagn.* Ser. I. ix. 108.—Syria.
Gaillardoti, *Boiss. Fl. Orient.* ii. 548.—Mesopot.
galegifolia, *Boiss. Diagn.* Ser. I. ii. 91.—Syria.
Gaudiniana, Jord. Cat. Gr. Jard. Grenoble, (1850) 10.—Cf. Linnaea, xxv. (1852) 305 = gracilis.
geminiflora, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 64.—Bras.
glabra, Desv. Journ. Bot. iii. (1814) 82 = viciaefolia.
gracilis, *Bess. Enum. Pl. Volh.* 74.—Graec.; As. Min.
Haussknechtii, *Boiss. Fl. Orient.* ii. 542.—Assyr.
Heldreichii, Bunge, ex Boiss. l. c. 530 = laconica.
heliocarpa, *Boiss. Diagn.* Ser. I. ii. 93.—Persia.
hemicycla, *Blanche, ex Boiss. Fl. Orient.* ii. 553.—Syria.
heterophylla, *C. A. Mey. Verz. Pfl. Cauc.* 146.—Reg. Caucas.; Persia.
heterophylla, Schrank, ex Steud. Nom. ed. II. ii. 213 = Desmodium triflorum.
Hohenackeriana, C. A. Mey. in Bull. Sc. Acad. Pétersb. ii. (1837) 36 = Michauxii.
horrida, *Desv. Journ. Bot.* iii. (1814) 83.—Afr. bor.
Huetiana, *Boiss. Diagn.* Ser. II. ii. 35.—Armenia.
Huetiana, Boiss. & Huet, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 75 = atropatana.
hypargyrea, *Boiss. Diagn.* Ser. I. ii. 91.—As. Min.
hypoleuca, De Roem. ex Willk. & Lange, Prod. Fl. Hisp. iii. 267 = stenorrhiza.

ONOBRYCHIS:—
incana, Gueldenst. Reisen, ii. 19 = viciaefolia.
incurvata, *DC. Prod.* ii. 346.—Hab.?
inermis, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 165 = viciaefolia.
kachetica, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 75.—Persia.
Kotschyana, *Fenzl, Pugill. Pl. Nov. Syr.* 3.—Syria.
laconica, *Orph. ex Boiss. Fl. Orient.* ii. 530.—Graecia.
lanata, *Boiss. Diagn.* Ser. I. ii. 92.—Persia.
lasiostachya, *Boiss. l. c.* 96.—As. Min.
laxiflora, *Baker, in Journ. Linn. Soc.* xix. (1882) 159.—Afghan.
lineata, Desv. Journ. Bot. iii. (1814) 80 = Flemingia lineata.
longespicata, Boiss. & Bal. ex Boiss. Fl. Orient. ii. 549 = hypargyrea.
longipes, *Bunge, ex Boiss. l. c.* 546.—Persia.
lunata, *Boiss. Diagn.* Ser. I. ii. 95.—Assyria.
major, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 533 = Balansae.
mareotica, Stev. ex DC. Prod. ii. 345 = viciaefolia.
matritensis, Boiss. & Reut. Pl. Nov. Hisp. Diagn. 11 = argentea.
megalophylla, Griseb. Spicil. Fl. Rumel. i. 68 = Tournefortii.
megataphros, *Boiss. Diagn.* Ser. I. ii. 97.—Syria.
melanotricha, *Boiss. l. c.* vi. 43.—Persia.
Michauxii, *DC. Prod.* ii. 347.—Armen.; Persia.
micrantha, *Schrenk, Enum. Pl. Nov.* 85.—Soongaria.
microptera, *Baker, in Journ. Linn. Soc.* xviii. (1880) 48.—Afghan.
miniata, Stev. ex Boiss. Fl. Orient. ii. 536 = gracilis.
montana, DC. Fl. Fr. iv. 611 = viciaefolia.
neglecta, *Godet, in Dub. Voy. Cauc.* iv. 541, nomen.—Reg. Caucas.
nitida, *Boiss. Diagn.* Ser. I. ii. 92.—Cappadoc.
nummularia, *Stocks, in Hook. Kew Journ.* iv. (1852) 146.—Beluchist.
Olivieri, *Boiss. Fl. Orient.* ii. 547.—Mesopot.
orientalis, Jaume St. Hil. Expos. ii. 239 = cornuta.
ornata, Boiss. & Buhse, in Mém. Soc. Nat. Mosc. xii. (1860) 75 = atropatana.
ornata, *Desv. Journ. Bot.* iii. (1814) 81.—Galatia.
ornata, Ledeb. Fl. Ross. i. 711 = argyrea.
oxyodonta, *Boiss. Diagn.* Ser. I. ii. 98.—Cappadoc.
oxyptera, *Boiss. l. c.* vi. 44.—Persia.
oxytropoides, *Bunge, ex Boiss. Fl. Orient.* ii. 530.—Armenia.
Pallasii, *Bieb. Cent. Pl. Ross.* t. 35.—Tauria.
pallens, *Lång, Enum.* 2; *ex Neilr. Aufzaehl. Ung. Gefaesspfl.* 349.—Europ.
pallescens, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 183 = viciaefolia.
pallida, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* ii. 530.—Cilicia.
pannosa, Boiss. Diagn. Ser. I. ix. 106 = Hedysarum pannosum.
parviflora, Boiss. & Reut. ex Nym. Consp. 199 = supina.
paucidentata, *Pomel, Nouv. Mat. Fl. Atl.* 200.—Alger.
peduncularis, *DC. Prod.* ii. 346.—Afr. bor.
peduncularis, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 264 = saxatilis.
petraea, Bess. Enum. Pl. Volh. 29 = gracilis.
petraea, *Desv. Journ. Bot.* iii. (1814) 82.—Reg. Caucas.
petraea, [Kit.] Host, Fl. Austr. ii. 351 = carpathica.
picta, Desv. ex DC. Prod. ii. 347 = Michauxii.
pilosa, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 250 = Tournefortii.
pisidica, *Boiss. Diagn.* Ser. I. ix. 107.—Pisidia.
procumbens, Bunge, ex Boiss. Fl. Orient. ii. 535 = Bungei.
procumbens, Fisch. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 64 = viciaefolia.
psoraleaefolia, *Boiss. Diagn.* Ser. I. ii. 93.—Persia.
ptolemaica, Boiss. & Gaill. ex Boiss. Fl. Orient. ii. 548 = Gaillardoti.
ptolemaica, *DC. Prod.* ii. 347.—Aegypt.; Arab.
pulchella, Heldr. in Boiss. Diagn. Ser. II. vi. 64 = laconica.
pulchella, *Schrenk, Enum. Pl. Nov.* 87.—Soongaria.
pumila, *Desv. Journ. Bot.* iii. (1814) 84.—Hispan.
radiata, Bieb. Cent. Pl. Ross. t. 55 = vaginalis.

ONOBRYCHIS:—
radiata, *Fisch. Cat. Hort. Gorenk.* (1812) 73.—Reg. Caucas.
radiata, Steud. Nom. ed. II. ii. 213 = circinnata, Michauxii.
reticulosa, Opiz, Natural. ix. (1825) 134 (Quid?).—Europ.
Richardi, *Baker, in Oliver, Fl. Trop. Afr.* ii. 141.—Afr. trop.
rotundifolia, Desv. Journ. Bot. iii. (1814) 84 = Indigofera echinata.
sativa, Lam. Fl. Fr. ii. 652 = viciaefolia.
saxatilis, *Lam. l. c.* 653.—Europ. occ.
scapiformis, C. Koch, in Linnaea, xix. (1847) 67 = cretica.
scrobiculata, *Boiss. Diagn.* Ser. I. ii. 95.—Persia.
sericea, Spreng. Syst. iii. 205 = Ebenus pinnata.
sibirica, *Turcz. ex Bess. in Flora*, xvii. (1834) I. *Beibl.* 10.—Sibir.
simplicifolia, *R. Br. in Salt, Abyss. App.* 65.—Abyss.
sororia, Desv. Journ. Bot. iii. (1814) 80 = Eleiotis sororia.
spicata, Moench, Meth. 122 = viciaefolia.
spinosissima, *Baker, in Journ. Linn. Soc.* xviii. (1880) 49.—Afghan.
Spruneriana, Hampe, in Flora, xxv. (1842) 61 = ebenoides.
squarrosa, Viv. Fl. Aegypt. Arab. Diagn. 14 = Gaertneriana.
stenorhiza, *DC. Prod.* ii. 346.—Hispan.
Stewartii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 141.—Ind. bor. occ.
subacaulis, *Boiss. Fl. Orient.* ii. 544.—Persia.
sulphurea, *Boiss. & Bal. Diagn.* Ser. II. vi. 63.—Cappadoc.
supina, *DC. Fl. Fr. Suppl.* 612.—Europ.
supina, Gaud. Fl. Helv. iv. 570 = arenaria.
Szovitsii, *Boiss. Fl. Orient.* ii. 542.—Persia.
tanaitica, Fisch. ex Steud. Nom. ed. II. ii. 214 = viciaefolia.
tanaitica, Spreng. Neue Entdeck. ii. 162 = petraea, viciaefolia.
tavernieraefolia, *Stocks, ex Boiss. Fl. Orient.* ii. 545.—Afghan.
tenuifolia, Moench, Meth. Suppl. 41 = saxatilis.
tenuis, Boiss. & Bal. Diagn. Ser. II. vi. 65 = cadmea.
Tommasinii, Borb. ex Nym. Consp. 589 = alba.
Tommasinii, Jord. Cat. Gr. Jard. Grenoble (1851).—Cf. Linnaea, xxv. (1852) 303 = gracilis.
Tournefortii, *Desv. Journ. Bot.* iii, (1814) 81.—Macedon.; As. Min.
tragacanthoides, DC. Prod. ii. 347 = cornuta.
trichocalycina, *Boiss. Diagn.* Ser. II. v. 90.—Armenia.
trilophocarpa, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 223, nomen = O. Crista-galli.
vaginalis, *C. A. Mey. Verz. Pfl. Cauc.* 145.—Reg. Caucas.
velutina, Boiss. Fl. Orient. ii. 549 = hypargyrea.
venosa, *Desv. Journ. Bot.* iii. (1814) 81.—Reg. Tunit.; Ins. Cyprus.
viciaefolia, *Scop. Fl. Carn.* ed. II. ii. 76.—Europ.; As. or. et bor.
vicioides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 372 = viciaefolia.
virgata, *Presl, Delic. Prag.* 43.—Sicilia.
Visianii, Borb. ex Nym. Consp. 198 = alba.
vulgaris, Gueldenst. Reisen, ii. 19, 33; Gaud. Fl. Helv. iv. 569 = viciaefolia.
Wiedemanni, *Boiss. Fl. Orient.* ii. 549.—Oriens.

ONOCHILIS, Mart. in Denkschr. Akad. Muench. v. (1817) 177 = **Nonnea**, Medic. (Borag.).
pulla, Mart. l. c. = N. pulla.

ONOCTONIA, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 276 = **Poteranthera**, Bong. (Melastom.).
calcarata, Naud. l. c. 277 = P. calcarata.
crassipes, Naud. l. c. = P. crassipes.
pauciflora, Naud. l. c. = P. pauciflora.

ONODONTEA, G. Don, Gen. Syst. i. 148 (1831), sphalm. = ANODONTEA, Sweet = **Alyssum**, Tourn. (Crucif.).

ONOEA, Franch. & Sav. Enum. Pl. Jap. ii. 178 (1879) = **Diarrhena**, Rafin. (Gramin.).
japonica, Franch. & Sav. l. c. = D. japonica.

ONONIS, Linn. Syst. ed. ed. I. (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 485.
ANONIS, Tourn. ex Adans. Fam. ii. 323 (1763).
BONAGA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 345; Phil. Bot. 206 (1789).
BUGRANOPSIS, Pomel, Nouv. Mat. Fl. Atl. 171 (1874).
NATRIX, Moench, Meth. 158 (1794).
PASSAEA, Adans. Fam. ii. 509 (1763).
adenotricha, *Boiss. Diagn.* Ser. I. ii. 14.—As. Min.; Syria.
adglutinans, Presl, Bot. Bemerk. 50 = O. Natrix.
aethiopica, Burm. f. Fl. Cap. Prod. 21 = Melolobium cernuum.
aggregata, Asso, Syn. Stirp. Arag. 97. t. 6. f. 1 = striata.
alba, *Poir. Voy. Barb.* ii. 210.—Reg. Mediterr.
alopecuroides, *Coss. Notes Crit.* 33.—Ind. or.
alopecuroides, *Linn. Sp. Pl.* 717.—Reg. Mediterr.
alopecuroides, Pall. Tabl. Taur. 55 = hircina.
altissima, Colm. Genist. & Anthyll. Portug. 69 = arvensis.
altissima, Lam. Encyc. i. 506 = hircina.
ambigua, *Lange, in Kjoeb. Vidensk. Meddel.* (1877) 238.—Hispan.
angustifolia, G. Don, Gen. Syst. ii. 160 = seq.
angustissima, *Lam. Encyc.* i. 508.—Reg. Mediterr. occ.
Anil, Mill. Gard. Dict. ed. VIII. n. 14 (Indigoferae sp.).—Ind. occ.
anomala, *Pomel, Nouv. Mat. Fl. Atl.* 165.—Alger.
antennata, *Pomel, l. c.* 315.—Alger.
anthylloides, Ser. in DC. Prod. ii. 168 = Lotononis umbellata.
antiquorum, *Linn. Sp. Pl.* ed. II. 1006.—Europ. austr.; Oriens.
antiquorum, Pollini, ex Reichb. Fl. Germ. Excurs. 517 = spinosa.
antiquorum, Willk. Sert. 41; ex Willk. & Lange, Prod. Fl. Hisp. iii. 393 = arvensis.
aphylla, Lam. Encyc. i. 509 = variegata.
arachnoidea, Lapeyr. Hist. Abr. Pl. Pyren. 409 = O. Natrix.
aragonensis, *Asso, Syn. Stirp. Arrag.* 96. t. 6. f. 2.—Hispan.
arborescens, Clem. ex Willk. & Lange, Prod. Fl. Hisp. iii. 416 = speciosa.
arborescens, *Desf. Fl. Atlant.* ii. 149. t. 193.—Alger.
Arbuscula, Desv. Journ. Bot. iii. (1814) 76 = tridentata.
arenaria, DC. Cat. Hort. Monsp. 128 = O. Natrix.
argentea, Linn. f. Suppl. 324 = Rhynchosia microscias.
aristata, Steud. Nom. ed. II. ii. 214, sphalm. = *longearistata*.
arragonensis, Coss. ex Boiss. & Reut. Pugill. Pl. Nov. 31 = Reuteri.
arthropodia, Brot. Phyt. Lusit. i. 141. t. 58 = pubescens.
arvensis, *Linn. Syst.* ed. XII. 478.—Europ.; Sibir.
arvensis, Retz. Obs. ii. 21 = hircina?
aspalathoides, G. Don, Gen. Syst. ii. 164 = seq.
asphaltoides, Ser. in DC. Prod. ii. 167 = Lotononis involucrata.
atlantica, *Ball, in Journ. Bot.* xi. (1873) 304.—Marocc.
Aucheri, Jaub. & Spach, Illustr. i. 85 = Lotus Garcini.
avellana, *Pomel, Nouv. Mat. Fl. Atl.* 168.—Alger.
baetica, Roxas Clem. Ensay. Vid, 291 = hirta.
balearica, Pourr. ex Nym. Consp. 161 = pubescens.
barbata, Cav. Ic. ii. 42 = minutissima.
Barrelieri, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 324 = tridentata.
Bernhardii, Vis. in Orto Bot. Padov. (1842) 101 = spinosa.
bicolor, Moris, Stirp. Sard. Elench. iii. 7 = biflora.
biflora, *Desf. Fl. Atlant.* ii. 143.—Reg. Mediterr.
Bonanniana, Presl, Fl. Sicul. p. xix = oligophylla.
Bourgaei, *Boiss. & Reut. Pugill. Pl. Nov.* 31.—Hispan.; Lusit.
brachycarpa, *Ser. in DC. Prod.* ii. 160.—Hispan.
brachystachya, *Vis. Fl. Dalm.* iii. 274.—Dalmat.
breviflora, Ser. in DC. Prod. ii. 160 = viscosa.
Broteriana, *Ser. l. c.* 162.—Hispan.
Broussonetii, DC. Prod. ii. 160 = pendula.
caduca, Vill. Hist. Pl. Dauph. iii. 428 = arvensis.
calycina, Lam. Encyc. i. 506 = pubescens.
calycina, *Viv. Fl. Lyb. Spec.* 41. t. 18. f. 3.—Afr. bor.
campestris, Koch & Ziz, Cat. Pl. Palat. 22 = spinosa, arvensis.
campestris, *Sieber, ex Webb & Berth. Phyt. Canar.* ii. 23.—Palaestina.

ONONIS:—
canescens, Steud. Nom. ed. II. ii. 214 = tridentata.
capensis, Linn. Pl. Rar. Afr. 16; Amoen. Acad. vi. 95 (Lotononis sp.).—Afr. austr.
capillaris, Thunb. Prod. Pl. Cap. 129 = Aspalathus capillaris.
capitata, Cav. Ic. ii. 43. t. 159 = Columnae.
cenisia, *Linn. Mant.* ii. 267.—Europ. austr.
cephalantha, *Pomel, Nouv. Mat. Fl. Atl.* 168.—Alger.
Cephalotes, *Boiss. Elench.* 33.—Hispan.
cernua, Linn. Pl. Rar. Afr. 16; Amoen. Acad. vi. 95 = Melolobium cernuum.
cerrifolia, Reichb. ex DC. Prod. ii. 317 = Stylosanthes procumbens.
cheiranthoides, *Lapeyr. ex Steud. Nom.* ed. II. ii. 214 (Quid?).—Pyren.
Cherleri, Desf. Fl. Atlant. ii. 148; Sibth. & Sm. Fl. Graec. vii. 72. t. 677 = reclinata.
Cherleri, Forsk. Fl. Aegypt. Arab. 131 = vaginalis.
Cherleri, *Linn. Sp. Pl.* ed. II. 1007.—Reg. Mediterr.
Cherleri, Linn. l. c., partim = Columnae.
chorassanica, *Bunge, in Boiss. Fl. Orient.* ii. 62.—Persia.
cicerifolia, *DC. Mém. Lég.* vi. 222; *Prod.* ii. 165.—Hab.?
cintrana, *Brot. Phyt. Lusit.* 138. t. 57.—Reg. Mediterr. occ.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 315.—Alger.
Columnae, *All. Fl. Pedem.* i. 318. t. 20. f. 3.—Reg. Mediterr.; Oriens.
compressa, *Lag. & Rodr. in Anal. Sc. Nat.* iv. (1801) 262.—Hab.?
confusa, Bor. Fl. Centr. Fr. ed. III. ii. 144, in obs. = spinosa?
conjugata, *Herb. Lamb. ex G. Don, Gen. Syst.* iii. 469.—Mexic.
coriifolia, Reichb. ex Guill. & Perr. Fl. Seneg. Tent. 204 = Stylosanthes erecta.
Cossoniana, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 33.—Reg. Mediterr. occ.
crassifolia, Duf. ex DC. Prod. ii. 161 = tridentata.
crinita, *Pomel, Nouv. Mat. Fl. Atl.* 170.—Alger.
crispa, *Linn. Sp. Pl.* ed. II. 1010.—Hispan.
crispa, Sibth. & Sm. Fl. Graec. vii. t. 680 = O. Natrix.
cristata, Mill. Gard. Dict. ed. VIII. n. 9 = cenisia.
crotalarioides, *Coss. Notes Crit.* 155.—Hispan.
cuspidata, *Desf. Fl. Atlant.* ii. 144.—Algeria.
decumbens, Mill. Gard. Dict. ed. VIII. n. 15 = *Milleri*.
decumbens, Sieber, ex DC. Prod. ii. 217 = Psoralea decumbens.
decumbens, Thunb. Prod. Pl. Cap. 129 = Lotononis porrecta.
Dehnhardtii, Tenore, Ind. Sem. Hort. Neap. (1825) 11. n. 18; Syll. 348 = serrata.
dentata, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 34 = reclinata.
Desfontainii, *Duf. ex DC. Prod.* ii. 162.—Hispan.; Ital.
diacantha, Sieber, ex Reichb. Pl. Crit. i. 9. t. 7 = antiquorum.
diffusa, Coss. Not. Crit. 35 = Cossoniana.
diffusa, Tenore, Prod. Fl. Nap. p. xli = serrata.
dumosa, Lapeyr. Hist. Abr. Pl. Pyr. 410 = aragonensis.
elegans, Clem. ex Willk. & Lange, Prod. Fl. Hisp. iii. 416 = speciosa.
ellipticifolia, *Willk. l. c.* 397.—Hispan.
elongata, Thunb. Prod. Pl. Cap. 129 = Lotononis prostrata.
emarginata, Boj. ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 576 = Crotalaria spinosa.
ervoides, Sieber, Reise Kreta, ii. 325 = Cicer ervoides.
euphrasiaefolia, *Desf. Fl. Atlant.* ii. 141. t. 184.—Alger.
excisa, Thunb. Fl. Cap. ed. Schult. 586 = Crotalaria humilis.
excisa, Thunb. Prod. Pl. Cap. 129 = geminata, Lotononis excisa.
falcata, Viv. Fl. Lyb. Spec. 41. t. 18 = angustissima.
fasciculata, Thunb. Prod. Pl. Cap. 130 = Aspalathus undulata.
filicaulis, *Salzm. ex Boiss. Voy. Bot. Espagne*, 153. t. 46.—Reg Mediterr. occ.
filiformis, Linn. Mant. ii. 266 = Indigofera sarmentosa.
flava, *Forsk. Fl. Aegypt. Arab.* 130.—Arab.
flexipes, *Webb & Berth. Phyt. Canar.* ii. 27.—Ins. Canar.
foetens, All. Fl. Pedem. i. 317. t. 41. f. 1 = hircina.

ONONIS:—
foetida, Schousb. ex Balb. in DC. Prod. ii. 161 = porrigens.
foliosa, *Willk. & Costa, in Linnaea*, xxx. (1859-60) 96.—Hispan.
fruticosa, *Linn. Sp. Pl.* 718.—Europ.
geminata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 28.—Afr. austr.
geminiflora, Lag. Gen. et Sp. Nov. 22 = biflora.
gibraltarica, Boiss. Elench. 32 = O. Natrix.
gibraltarica, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 410 = Cossoniana.
glabra, Thunb. Prod. Pl. Cap. 130 = Lotononis trichopoda.
glandulifera, Weinm. Cat. Hort. Dorpat. (1810) 107 = rotundifolia.
glaucescens, *Pomel, Nouv. Mat. Fl. Atl.* 167.—Alger.
glutinosa, Mart. in Denkschr. Akad. Muench. vi. (1816-17) 154 = Crotalaria paniculata.
glutinosa, Presl, Delic. Prag. 35 = viscosa.
grandiflora, *Munby, in Bull. Soc. Bot. Fr.* xi. (1864) 45.—Alger.
Hackelii, Lange, in Kjoeb. Vidensk. Meddel. (1877) 239 = biflora.
hebecarpa, *Webb & Berth. Phyt. Canar.* ii. 25.—Ins. Canar.
heterophylla, Scheele, in Linnaea, xxv. (1852) 266.—Dalmat.
heterophylla, Thunb. Prod. Pl. Cap. 129 = Lotononis erostrata.
hircina, *Jacq. Hort. Vindob.* i. 40.—Europ.; As. bor.
hirsuta, Thunb. Prod. Pl. Cap. 129 = Euchlora serpens.
hirta, *Desf. Hort. Par*; *ex Poir. Encyc. Suppl.* i. 741.—Hispan.; Cilicia; Syria.
hispanica, Linn. f. Suppl. 324 = O. Natrix.
hispida, *Desf. Fl. Atlant.* ii. 146. t. 189.—Alger.
hybrida, Schrank, Baier. Fl. ii. 251 = pubescens?
inaequifolia, Ser. in DC. Prod. ii. 165 = O. Natrix.
inaperta, Moench, Meth. Suppl. 54 = Columnae.
incisa, *Coss. & Dur. in Munby, Cat. Pl. Alger.* 8, nomen.—Alger.
inclusa, *Pourr. ex Willk. & Lange, Prod. Fl. Hisp.* iii. 401.—Corsic.
inermis, Huds. Fl. Angl. ed. II. 313 = hircina.
inermis, *Pall. Reise Suedl. Statth. Russl.* ii. 87, 88.—Tauria.
insignis, Coss. ex Nym. Consp. 161.—Hispan.
intermedia, C. A. Mey. ex Beck. in Bull. Soc. Nat. Mosc. (1858) i. 28 = arvensis.
involucrata, Berg. Desc. Pl. Cap. 213; Linn. f. Suppl. 324 = Lotononis involucrata.
juncea, *Asso, Syn. Stirp. Arrag.* 96. t. 5. f. 2.—Hispan.
Kotschyana, *Fenzl, Pugill. Pl. Nov. Syn.* 5.—Syria.
Kotschyana, Kotschy, ex Boiss. Fl. Orient. ii. 58 = adenotricha.
laevis, *Pall. Reise Suedl. Statth. Russ. Reich.* i. 350.—Rossia.
Lagopus, Thunb. Prod. Pl. Cap. 130 = Aspalathus lotoides.
latifolia, Asso, Syn. Stirp. Arag. Mant. t. 11. f. 1 = rotundifolia.
laxiflora, *Desf. Fl. Atlant.* ii. 146. t. 190.—Afr. bor.; Ins. Canar.
laxiflora, Lowe, ex Ball, in Journ. Linn. Soc. xvi. (1878) 408 = pendula.
laxiflora, Viv. Fl. Cors. Sp. 13 = reclinata.
legitima, Delarb. Fl. Auv. ed. II. 446 = spinosa, arvensis.
leiosperma, *Boiss. Fl. Orient.* ii. 57.—Syria; Persia.
leucotricha, *Coss. Notes Crit.* 34.—Hispan.
lingulata, *Munby, Cat. Pl. Alger.* 8.—Alger.
litoralis, *Moris, ex Spreng. Syst.* iv. *Cur. Post.* 346.—Sardin.
littoralis, Vahl, ex Willk. & Lange, Prod. Fl. Hisp. iii. 398 = serrata.
longearistata, Presl, Fl. Sic. p. xix = viscosa.
longifolia, Brouss. ex Willd. Enum. Hort. Berol. 750 = angustissima.
lusitanica, Spreng. Syst. iii. 182 = Broteriana.
macracantha, Bernh. Sel. Sem. Hort. Erf. (1835).—Cf. Linnaea, xi. (1837) Litt. 86 = spinosa.
macracantha, Clarke, Trav. ii. 354 = antiquorum.
macrophylla, *Spreng. Fl. Hal. Mant.* 47.—Lusit.
maritima, Cav. ex Nym. Consp. 160 = O. Natrix.
maritima, Dum. in Bull. Soc. Bot. Belg. i. (1862) 113, in nota = arvensis.
Masquillierii, *Bertol. in Nov. Comm. Acad. Sc. Bonon.* iv. (1840) 19. t. 5.—Italia.

ONONIS:—
massesylia, *Pomel, Nouv. Mat. Fl. Atl.* 316.—Alger.
mauritanica, Linn. Mant. ii. 266 = Indigofera coriacea.
mauritanica, *Pomel, Nouv. Mat. Fl. Atl.* 164.—Alger.
mauritanica, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 411 = alopecuroides.
Maweana, *Ball, in Journ. Bot.* xi. (1873) 304.—Marocc.
megalostachys, *Munby, in Bull. Soc. Bot. Fr.* xi. (1864) 45.—Alger.
micrantha, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 25.—Ins. Madera.
micrantha, Thunb. Prod. Pl. Cap. 130 = Lotononis micrantha.
microphylla, Linn. f. Suppl. 324; Thunb. Prod. Pl. Cap. 129 = Melolobium microphyllum.
microphylla, *Pomel, Nouv. Mat. Fl. Atl.* 164.—Alger.
microphylla, Presl, Bot. Bemerk. 50 = O. Natrix.
microsperma, Lowe, Man. Fl. Mad. i. 598 = micrantha.
Milleri, Ser. in DC. Prod. ii. 168 (Indigoferae sp).—Mexic.
Miniana, Plan. Fl. Galleg. 166 = arvensis.
minuta, Pall. Tabl. Taur. 55; et Phys. Gem. Taur. 115 = Columnae?
minutissima, Jacq. Fl. Austr. iii. 23. t. 240 = Columnae.
minutissima, *Linn. Sp. Pl.* 717.—Reg. Mediterr. occ.
mitis, Mill. Gard. Dict. ed. VIII. n. 2; C. C. Gmel. Fl. Bad. ii. 162 = arvensis.
mitis, Pall. Reise, i. 236 = hircina.
mitissima, *Linn. Sp. Pl.* 717.—Reg. Mediterr.
mollis, Lag. Gen. et Sp. Nov. 22 = Desfontainii.
mollis, Savi, in Mem. Soc. Ital. Mod. ix. (1802) 351. t. 8 = reclinata.
monophylla, Desf. Fl. Atlant. ii. 145 = alopecuroides.
montana, *Coss. Notes Crit.* 103.—Hispan.
Morissonii, Gouan, Illustr. 47 = pubescens.
natricoides, Coss. & Dur. ex Munby, Cat. Pl. Alger. 8 = lingulata.
Natrix, *Linn. Sp. Pl.* 717.—Reg. Mediterr.
occidentalis, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 394 = arvensis.
oligophylla, *Tenore, Prod. Fl. Nap. Suppl.* ii. 69.—Ital.
ornithopodioides, *Linn. Sp. Pl.* 718.—Reg. Mediterr.
ornithopodioides, Sieber, ex Tausch, in Flora, xiv. (1831) 210 = Medicago minima.
pallida, *Willd. ex Hornem. Hort. Hafn.* ii. 676.—Hab.?
paniculata, *Cav. in Anal. Cienc. Nat.* iv. (1801) 69.—Afr. bor.
paradoxa, Welw. ex Ball, in Journ. Linn. Soc. xvi. (1878) 411 = alopecuroides.
parviflora, *Berg. Desc. Pl. Cap.* 214.—Afr. austr.
parviflora, Lam. Encyc. i. 510 = Columnae.
parvula, Steud. ex Baker, in Oliver, Fl. Trop. Afr. ii. 48 = reclinata.
peduncularis, *Lindl. Bot. Reg.* t. 1447.—Ins. Canar.
pendula, *Desf. Fl. Atlant.* ii. 147. t. 191.—Reg. Mediterr. occ.
pendula, Sieber, ex Ser. in DC. Prod. ii. 162 = Sieberi.
persica, Burm. f. Fl. Ind. 157. t. 49 = sicula.
phyllocephala, *Boiss. Fl. Orient.* ii. 63.—Syria.
Picardi, *Boiss. Elench.* 33.—Hispan.
Picardi, Coss. ex Boiss. & Reut. Pugill. Pl. Nov. (1852) 31 = Bourgaei.
picta, Desf. Fl. Atlant. ii. 144. t. 187 = O. Natrix.
pilosa, Bartl. in Bartl. & Wendl. f. Beitr. ii. 77 = reclinata.
pinguis, Linn. Sp. Pl. 1006 = O. Natrix.
pinnata, *Brot. Fl. Lusit.* ii. 99.—Lusitan.
polymorpha, Guss. Pl. Rar. Sic. 291 = Sieberi.
polyphylla, *Ball, in Journ. Bot.* xi. (1873) 304.—Marocc.
porrigens, *Salzm. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 406.—Marocc.
praecox, Bianca, Pl. Nov. 14; ex Guss. Fl. Sic. Syn, ii. 851 = biflora.
procumbens, Link, Handb. ii. 512, sphalm. = *procurrens*, Wallr.
procurrens, Benth. in Royle, Illustr. Bot. Himal. 197 = hircina.
procurrens, Wallr. Sched. Crit. 381 = arvensis.
prostrata, Burm. f. Fl. Cap. Prod. 21 = Lotononis prostrata.
psammophila, Dur. ex Munby, Cat. Pl. Alger. 8 = lingulata.

ONONIS:—
Pseudo-hircina, Schur, Enum. Pl. Transs. 150 = hircina.
Pseudo-repens, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 170 = praec.?
pubescens, *Linn. Mant.* ii. 267.—Reg. Mediterr.
pulchella, Boiss. Fl. Orient. ii. 58 = adenotricha.
pungens, *Pomel, Nouv. Mat. Fl. Atl.* 166.—Alger.
purpurea, Gilib. Fl. Lituan. ii. 80 = arvensis.
purpurea, Mill. ex Steud. Nom. ed. II. ii. 215 = fruticosa.
pusilla, Linn. Syst. ed. X. 1159 = Columnae?
pygmaea, C. Koch, in Linnaea, xix. (1847) 62 = viscosa.
pyramidale, *Cav. in Anal. Cienc. Nat.* iv. (1801) 71.—Afr. bor.
pyrenaica, *Willk. & Costa, in Linnaea*, xxx. (1859–60) 97.—Mont. Pyren.
quinata, Forsk. Fl. Aegypt. Arab. 130 = Dorycnium latifolium.
quinata, Thunb. Prod. Pl. Cap. 130 = Lotononis quinata, *Harv.*
racemosa, Brot. Fl. Lusit. ii. 96 = O. Natrix.
racemosa, *Thunb. Prod. Pl. Cap.* 129.—Afr. austr.
ramosissima, Desf. Fl. Atlant. ii. 142. t. 186 = O. Natrix.
reclinata, Brot. Fl. Lusit. ii. 97 = pendula.
reclinata, DC. Fl. Fr. iv. 511, in syn. = striata.
reclinata, *Linn. Sp. Pl.* ed. II. 1011.—Reg. Mediterr.
reflexa, Roem. Hort. Turic.; ex Steud. Nom. ed. II. ii. 215 (Quid?).—Lusitan.
repens, Linn. Sp. Pl. 717 = arvensis.
Reuteri, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 30.—Hispan.
rhinanthoides, *Lapeyr. Hist. Abr. Pl. Pyren.* 407.—Mont. Pyren.
rigida, Kunze, in Flora, xxix. (1846) 634 = fruticosa.
rosaefolia, *DC. Mém. Lég.* vi. 221; Prod. ii. 165.—Hispan.
rosea, *Dur. in Duch. Rev. Bot.* ii. (1846) 437.—Afr. bor.
rostrata, Ucria, in Roem. Arch. i. 1. 69 = Trigonella prostrata?
rotundifolia, *Linn. Sp. Pl.* 719.—Europ. austr.
rotundifolia, Linn. l. c. ed. II. 1050, excl. syn. = tribracteata.
Salzmanniana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 34 = alopecuroides.
saxatilis, Lam. Encyc. i. 509 = minutissima.
saxicola, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 32.—Hispan.
scabra, *Lapeyr. Hist. Abr. Pl. Pyren.* 407.—Mont. Pyren.
scariosa, Moench, Meth. 158 = mitissima.
Schouwii, Ser. in DC. Prod. ii. 162 = pendula.
secunda, Thunb. Prod. Pl. Cap. 130 = Lotononis peduncularis.
senescens, Lapeyr. Hist. Abr. Pl. Pyren. 405 = spinosa.
sericea, Thunb. Prod. Pl. Cap. 129 = Argyrolobium lanceolatum.
serotina, *Pomel, Nouv. Mat. Fl. Atl.* 166.—Alger.
serrata, *Forsk. Fl. Aegypt. Arab.* 131.—Reg. Mediterr.
serrata, Gren. & Godr. Fl. Fr. i. 375 = diffusa.
sicula, *Guss. Adnot. Cat. Pl. Boccad.* (1821) 10.—Reg. Mediterr.; Persia.
Sieberi, *Bess. ex DC. Prod.* ii. 162.—Europ. austr.
speciosa, *Lag. Nov. Gen. et Sp.* 22.—Hispan.
spicata, *Munby, Fl. Algér.* 76 = rosea.
spicata, Thunb. Prod. Pl. Cap. 129 = Aspalathus heterophylla.
spinosa, Bieb. Fl. Taur. Cauc. ii. 149 = hircina.
spinosa, Cav. Desc. 533 = antiquorum.
spinosa, Hasselq. Iter, ed. Gall. i. 138; ii. 187; ex DC. Prod. ii. 352 = Alhagi maurorum.
spinosa, *Linn. Sp. Pl.* 716.—Europ.
spinosaeformis, *Simk. in Oestr. Bot. Zeitschr.* xxvii. (1877) 158.—Europ. austr.
spinulosa, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 57 = atlantica.
stipularis, Duf. ex Nym. Consp. 161 = sicula.
stipulata, Thunb. Prod. Pl. Cap. 129 = Melolobium stipulatum.
striata, *Gouan, Illustr.* 47.—Europ. austr.
stricta, *Pomel, Nouv. Mat. Fl. Atl.* 169.—Alger.
strigosa, Burm. f. Fl. Cap. Prod. 21 = Psoralea hirta.
strigosa, Thunb. Prod. Pl. Cap. 130 = Lotononis umbellata.
subaristata, DC. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 652 = cenisia.

ONONIS:—
subcordata, *Cav. in Anal. Cienc. Nat.* iv. (1801) 70 (sp. dub.).—Hispan.
subocculta, Vill. Prosp. 41 = Columnae.
subspicata, *Lag. ex Colm. Genist. et Anthyll. Portug.* 73.—Hispan.
syriaca, Jaub. & Spach, Illustr. ii. 72. t. 155 = adenotricha.
tingitana, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 410 = Cossoniana.
Tournefortii, *Coss. Notes Crit.* 34.—Reg. Mediterr. occ.
tribracteata, *DC. Fl. Fr. Suppl.* 553.—Carinthia.
tribracteolata, Link, Handb. ii. 157, sphalm. = praec.
tridentata, *Linn. Sp. Pl.* 718.—Hispan.
trifoliata, Linn. Syst. ed. XII. 479 = Trifolium bracteatum.
Tuna, *Pomel, Nouv. Mat. Fl. Atl.* 317.—Alger.
ulicina, *Webb & Berth. Phyt. Canar.* ii. 23.—Ins. Canar.
umbellata, Linn. Mant. ii. 266 = Lotononis peduncularis.
umbrosa, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 169 = spinosa.
vaginalis, *Vahl, Symb. Bot.* i. 53.—Afr. bor.; Syria.
variegata, *Linn. Sp. Pl.* 717.—Reg. Mediterr.
variegata, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 411 = alopecuroides.
vestita, Viv. Fl. Lyb. Spec. 40. t. 18 = vaginalis.
villosa, Linn. Syst. ed. X. 1160 = viscosa.
villosa, Sieber, ex Presl, Bot. Bemerk. 49 = Rhynchosia microscias.
villosa, Thunb. Prod. Pl. Cap. 129 = Lotononis azurea.
villosissima, *Desf. Fl. Atlant.* ii. 147. t. 192.—Alger.
violacea, Peterm. in Rabenh. Bot. Centralbl. (1846) 45 = spinosa.
virgata, Burm. f. Fl. Cap. Prod. 21 = Psoralea decumbens.
virgata, *Kunze, in Flora*, xxix. (1846) 760.—Hispan.
viscidula, *Pomel, Nouv. Mat. Fl. Atl.* 318.—Alger.
visciflora, *Ser. in DC. Prod.* ii. 168.—Afr. austr.
viscosa, Coss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 406 = porrigens.
viscosa, *Linn. Sp. Pl.* 718.—Europ. austr.
viscosa, Sieber, ex Presl, Bot. Bemerk. 50 = pubescens.
viscosa, Welden, ex Reichb. Fl. Germ. Excurs. 517 = O. Natrix.
vulgaris, Gueldenst. Reisen, ii. 25 = hircina.

ONOPIX, Rafin. Fl. Ludov. 59 (1817). Genus dubium *COMPOSITARUM*.
ONOPYXOS, Spreng. Syst. iii. 386 (1826).
intybaceus, Rafin. Fl. Ludov. 60.—Louisiana.
sericeus, Rafin. l. c.—Louisiana.

ONOPORDON, Linn. Sp. Pl. ed. I. 827 (1753). *COMPOSITAE*, Benth. & Hook. f. ii. 469.
ACANOS, Adans. Fam. ii. 116 (1763).
acanthifolium, C. Koch, in Linnaea, xxiv. (1851) 690 = Carduus lanuginosus.
acanthiifolia, Gilib. Fl. Lituan. i. 190 = O. Acanthium.
Acanthium, *Linn. Sp. Pl.* 827.—Europ.; Afr. bor.; Oriens.
acaule, Willd. Sp. Pl. iii. 1688, in syn. = seq.
acaulon, *Linn. Sp. Pl.* ed. II. 1159.—Hab.?
alexandrinum, Boiss. Diagn. Ser. I. x. 93 = Sibthorpianum.
algeriense, *Pomel, Nouv. Mat. Fl. Atl.* 21.—Alger.
ambiguum, *Fresen. in Mus. Senckenb.* i. (1833) 85.—Arab.; Syria.
anatolicum, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 562 = Sibthorpianum.
anisacanthum, Boiss. Diagn. Ser. I. x. 93 = heteracanthum.
arabicum, Bové, ex DC. Prod. vi. 618 = illyricum.
arabicum, *Linn. Sp. Pl.* 827.—Europ. austr.
araneosum, Boiss. ex Boiss. Fl. Orient. iii. 562 = Sibthorpianum.
arenarium, *Pomel, Nouv. Mat. Fl. Atl.* 20.—Alger.
argolicum, Boiss. Diagn. Ser. I. x. 91 = tauricum.
berardioides, *Sch. Bip. in Linnaea*, xix. (1847) 329.—Persia; Beluchist.
bracteatum, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. x. 91.—Pisidia.
Carduelium, *C. Bolle, in Bonplandia*, vii. (1859) 297.—Ins. Canar.
carduiforme, Boiss. Diagn. Ser. I. x. 92 = Sibthorpianum.
Cardunculus, Boiss. Fl. Orient. iii. 561 = illyricum.
carthamoides, Buch.-Ham. ex Wall. Cat. n. 3922 = Vernonia Roxburghii.

ONOPORDON :—
caulescens, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 361.—Ins. Melos.
corymbosum, *Boiss. Fl. Orient.* iii. 560.—Lycia; Cilicia.
corymbosum, Willk. in Linnaea, xxx. (1859–60) 108 = tauricum.
cynarocephalum, *Boiss. & Blanche, Diagn.* Ser. II. iii. 48.—Syria.
cynaroides, Stev. ex Link, Enum. Hort. Berol. ii. 303 = Coussinia cynaroides.
Delortii, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. i. (1872–73) 385 = illyricum.
deltoides, Ait. Hort. Kew. ed. I. iii. 146 = Serratula atriplicifolia.
Diacantha, Labill. ex Steud. Nom. ed. II. ii. 216 = Cnicus Diacantha.
elatum, Sibth. & Sm. Fl. Graec. ix. 23 = tauricum.
elongatum, Lam. Fl. Fr. ii. 6 = illyricum.
excelsum, Salisb. Prod. 185 = arabicum.
floccosum, Boiss. Diagn. Ser. I. x. 92 = illyricum.
glomeratum, Costa, Introd. Fl. Catal. 135 = illyricum.
graecum, Bory & Chaub. ex Boiss. Fl. Orient. iii. 562 = Sibthorpianum.
graecum, Gouan, Illustr. 64. t. 25 = illyricum.
graecum, *Linn. f. Suppl.* 349.—Graecia; Cyprus.
gratum, Bess. ex Steud. Nom. ed. II. ii. 216 (Quid?). —Hab.?
heteracanthum, *C. A. Mey. Vers. Pfl. Cauc.* 68.—Reg. Caucas.; Persia.
horridum, Viv. Fl. Lib. Spec. 68 = illyricum.
Ilex, *Janka, in Termész. Füzetek*, ii. (1878) 29, nomen.—*Cf. Oestr. Bot. Zeitschr.* xxviii. (1878) 176. —Graecia.
illiricum, Vill. Hist. Pl. Dauph. iii. 26 = seq.
illyricum, *Linn. Sp. Pl.* 827.—Europ. austr.; As. Min.
intermedium, *Schult. Obs. Bot.* 198.—Hab.?
lanatum, Herb. Madr. ex DC. Prod. vii. 36 = Dicoma tomentosa.
lanatum, Moench, Meth. Suppl. 225 = arabicum.
lanigerum, Boiss. Fl. Orient. iii. 564 = leptolepis.
leptolepis, *DC. Prod.* vi. 619.—Persia.
libanoticum, Boiss. & Blanch. Diagn. Ser. II. iii. 48 = illyricum.
lyratum, Froel. ex. Sch. Bip. in Linnaea, xix. (1847) 329 = Jurinea depressa.
macrocanthum, *Schousb. Vextr. Marokko*, 198. t. 5.—Reg. Mediterr. occ.
macrocanthum, Schult. Obs. Bot. 198 = O. Acanthium.
macrocanthum, Sibth. & Sm. Fl. Graec. ix. 23 = Sibthorpianum.
macrocanthum, Urv. in Mém. Soc. Linn. Par. i. (1822) 361 = illyricum.
maximum, *Schult. Obs. Bot.* 199.—Hab.?
meridionale, *Link, Handb.* i. 685.—Arab.
myriacanthum, Boiss. Diagn. Ser. II. vi. 114 = illyricum.
nervosum, *Boiss. Voy. Bot. Espagne*, ii. 357. t. 108.—Hispan.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 51.—Turkestan.
orientale, Mill. Gard. Dict. ed. VIII. n. 4 = illyricum.
parnassicum, *Boiss. & Heldr. Diagn.* Ser. II. iv. 114. —Graecia.
polyacanthum, Eichw. Pl. Nov. Casp. Cauc. 3 = Cousinia onopordioides.
polycephalum, *Boiss. Diagn.* Ser. II. iii. 47.—As. Min.
pyrenaicum, DC. Fl. Fr. Suppl. 457 = acaulon.
ramosum, Dulac, Fl. Hautes-Pyr. 525 = O. Acanthium.
rhodense, Boiss. Fl. Orient. iii. 560 = illyricum.
rotundifolium, All. Fl. Pedem. i. 144 = Berardia subacaulis.
Schultesii, Bretting. ex Koch, Syn. Fl. Germ. ed. I. 404 = O. Acanthium.
Sibthorpianum, *Boiss. & Heldr. in Boiss. Fl. Orient.* iii. 561.—Graecia; Syria; Afr. bor.
stenostegium, *Boiss. Fl. Orient.* iii. 563.—Mesopot.
tauricum, *Willd. Sp. Pl.* iii. 1687.—Europ. austr.
teres, Buch.-Ham. ex Wall. Cat. n. 2926 = Vernonia teres.
tomentosum, Moench, Meth. 551 = illyricum.
tomentosum, Salisb. Prod. 185 = O. Acanthium.
uniflorum, Cav. Ic. i. 60. t. 88 = acaulon.
virens, DC. Fl. Fr. Suppl. 456 = tauricum.
viride, Desf. Tabl. ed. II. 92 = O. Acanthium.
viscosum, Hornem. ex Spreng. Syst. iii. 386 = tauricum.
Weissianum, Aschers. ex E. Weiss, in Verh. Zool. Bot. Ges. Wien, xix. (1869) 43 = Sibthorpianum.

ONOPORDUM, Vaill. ex Linn. Gen. ed. I. 243 (1737) = praec.

ONOPYXOS, Spreng. Syst. iii. 386 (1826) = ONOPIX, Rafin. (Compos.).

ONOPYXUS, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 319 = **Carduus**, Tourn. (Compos.).
hamulosus, Bubani, l. c. = C. hamulosus.

ONOSERIS, DC. in Ann. Mus. Par. xix. (1812) 65. t. 12. *COMPOSITAE*, Benth. & Hook. f. ii. 486.
Caloseris, Benth. Pl. Hartw. 88 (1841).
Centroclinium, D. Don, in Trans. Linn. Soc. xvi. (1830) 254.
Chaetachlaena, D. Don, l. c. 256 (1830).
Cladoseris, Spach, Hist. Vég. Phan. x. 35 (1841).
Cursonia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 422.
Hilairia, DC. Prod. vii. 33 (1838).
Hipposeris, Cass. in Dict. Sc. Nat. xxxiii. 464 (1824).
Isotypus, H. B. & K. Nov. Gen. et Sp. iv. 11. t. 307 (1820).
Rhodoseris, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 84.
Schaetzellia, Klotzsch, in Otto & Dietr. Allg. Gartenz. xvii. (1849) 81.
Seris, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139.
acerifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 8.—Am. austr.
adpressa, *Less. Syn. Comp.* 120.—Peruv.
annua, *Less. in Linnaea*, v. (1830) 341.—Am. austr.
brevifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 246.—Bras.
Castelnaeana, *Wedd. Chlor. And.* i. 10.—Peruv.
corymbosa, *Benth. & Hook. f. Gen.* ii. 487.—Bras.
Cumingii, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 103 = odorata?
denticulata, Willd. ex DC. Prod. vii. 22 = Lycoseris denticulata.
Drakeana, *André, in Rev. Hortic.* (1882) 181.—N. Granat.
eriocephala, Benth. Pl. Hartw. 211 = Hieracium erianthum.
hastata, *Wedd. Chlor. And.* i. 9.—Bolivia.
heterophylla, Spreng. Syst. iii. 503 = Trichocline heterophylla.
hieracioides, Bert. ex DC. Prod. vii. 29 = Chaetanthera Berteriana.
hieracioides, *H. B. & K. Nov. Gen. et Sp.* iv. 7. t. 304.—Am. austr.
hyssopifolia, *H. B. & K. l. c.* t. 306.—Ecuador.
integrifolia, *Less. in Linnaea*, v. (1830) 343.—Peruv.
Isotypus, *Benth. & Hook. f. Gen.* ii. 487.—Mexic.
lanuginosa, Wall. Cat. n. 2929 = Gerbera lanuginosa.
linifolia, Bert. in Merc. Chil. (1829).—Cf. Am. Journ. Sc. Ser. I. xxiii. (1833) 258 = Chaetanthera linifolia.
macrophylla, Wall. ex Steud. Nom. ed. II. ii. 216 = Gerbera nepalensis.
mexicana, Willd. Sp. Pl. iii. 1703 = Lycoseris mexicana.
montevidensis, Spreng. Syst. iii. 502 = Leucopsis diffusa.
nepalensis, Steud. Nom. ed. II. ii. 216 = Gerbera nepalensis.
odorata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 103.—Peruv.
ovalifolia, Wall. Cat. n. 2930 = Gerbera piloselloides.
paniculata, [Moç. & Sesse,] ex DC. Prod. vii. 33 = O. Isotypus.
purpurata, *Willd. Sp. Pl.* iii. 1702.—N. Granat.
purpurea, DC. Prod. vii. 34 = praec.
reflexa, *Less. Syn. Comp.* 119.—Peruv.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 7.—Am. austr.
speciosa, *H. B. & K. l. c.* 7. t. 305.—Ecuador.
stricta, Spreng. Syst. iii. 503 = Trixis stricta.
turbacensis, Spreng. l. c. 502 = Lycoseris mexicana.

ONOSMA, Linn. Sp. Pl. ed. II. 196 (1762). *BORAGINEAE*, Benth. & Hook. f. ii. 864.
Colsmannia, Lehm. in Ges. Naturf. Fr. Berl. Mag. viii. (1817–18). 92, t. 4.
Maharanga, DC. Prod. x. 71 (1846).
Podonosma, Boiss. Diagn. Ser. I. xi. 113 (1849).
Sava, Adans. Fam. ii. p. (21) (1763).

ONOSMA :—
albo-roseum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 38.—As. Min.; Syria.
aleppicum, *Boiss. Diagn.* Ser. I. xi. 107.—Syria; Mesopot.
angustifolium, Lehm. Asperif. 361 = stellulatum.
arenarium, Waldst. & Kit. Pl. Rar. Hung. iii. 308. t. 279 = echioides.
armenum, *DC. Prod.* x. 59.—As. Min.
asperifolium, Lehm. Asperif. ii. 372 = Arnebia cornuta.
astroites, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 585, in nota = giganteum.
Aucherianum, DC. Prod. x. 60 = stellulatum.
atrocyaneum, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 214.—As. centr.
aurantiacum, Janka, ex Boiss. Fl. Orient. iv. 201 = stellulatum.
auriculatum, *DC. Prod.* x. 61.—Mesopot.
bicolor, *Wall. Cat.* n. 939.—Reg. Himal.
bifrons, Stev. ex Ledeb. Fl. Ross. iii. 128 = polyphyllum.
bilabiatum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 153.—Persia.
Bodeanum, *Boiss. Fl. Orient.* iv. 187.—Persia.
Bourgaei, *Boiss. Pl. Or. Nov. Dec.* ii. 2.—Armenia.
brachyanthum, Boiss. Diagn. Ser. I. vii. 34 = rostellatum.
brachysolen, Boiss. Diagn. l. c. xi. 104 = flavum.
bracteatum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 13. —Reg. Himal.
brevifolium, DC. ex Boiss. Fl. Orient. iv. 202 = stellulatum.
bulbotrichum, *DC. Prod.* x. 64.—Persia.
caerulescens, *Boiss. Diagn.* Ser. I. xi. 110.—Syria.
calycinum, Stev. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 81 = setosum.
canescens, Presl, Fl. Sic. p. xxxiv = stellulatum.
caspica, Willd. Sp. Pl. i. 775 = Nonnea picta.
caspium, Gruner, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 441 = setosum.
cassium, *Boiss. Diagn.* Ser. I. xi. 102.—Syria.
cespitosum, *Kotschy, in Unger & Kotschy, Cypern*, 282.—Ins. Cyprus.
chlorotrichum, *Boiss. & Noé, Diagn.* Ser. II. iii. 137. —Kurdistan.
chorassanicum, Bunge, Helioc. 12, nomen = bulbotrichum.
ciliatum, DC. Prod. x. 61 = molle.
cinereum, Schreb. in Nov. Act. Nat. Cur. iii. (1767) 474 = stellulatum.
circinnatum, Bourg. ex Boiss. Pl. Or. Nov. Dec. ii. 3 = Bourgaei.
coerulea, Willd. Sp. Pl. i. 775 = Moltkia coerulea.
coerulescens, vide caerulescens.
concinnum, DC. Prod. x. 64 = microspermum.
congestum, DC. l. c. 61 = albo-roseum.
cyaneum, Less. ex Ledeb. Fl. Ross. iii. 123 = staminеum.
dalmaticum, Scheele, in Flora, xxvi. (1843) 560 = stellulatum.
dasytrichum, *Boiss. Diagn.* Ser. I. vii. 33.—Persia.
decipiens, Schott & Kotschy, ex Boiss. Pl. Or. Nov. Dec. ii. 8 = nanum.
dichroanthum, Boiss. Diagn. Ser. I. xi. 107 = setosum.
diffusum, Boiss. & Huet, l. c. Ser. II. iii. 137 = rupestre.
divaricatum, Lehm. Asperif. 372 = Arnebia cornuta.
echinatum, Auch. ex DC. Prod. x. 58 = aleppicum.
echinatum, *Desf. Fl. Atlant.* i. 161. t. 43.—Afr. bor.
echioides, Kotschy, ex Boiss. Diagn. Ser. I. xi. 108 = aleppicum.
echioides, *Linn. Sp. Pl.* ed. II. 196.—Europ.; As. temp.
echioides, Petagn. Inst. Bot. ii. 320 = stellulatum.
echioides, Sibth. & Sm. Fl. Graec. ii. t. 172 = frutescens.
echioides, Sieber, ex Nym. Consp. 516 = graecum.
elegans, C. Koch, in Linnaea, xvii. (1843) 306 = sericeum.
elivendicum, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 28.—Persia.
Emodi, Wall. Cat. n. 940 = Wallichianum.
Emodi, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 11.—Reg. Himal.
erectum, Sibth. & Sm. Fl. Graec. Prod. i. 121 = stellulatum.
fallax, Borb. ex Nym. Consp. 516 = setosum.
flavidum, Boiss. Diagn. Ser. I. xi. 100 = flavum.
flavum, *Vatke, ex Boiss. Fl. Orient.* iv. 186.—As. Min.; Persia.

ONOSMA:—
frutescens, *Lam. Illustr.* i. 407.—Graecia; Syria.
fruticosum, *Labill. Pl. Syr. Dec.* iii. 10. t. 6.—Ins. Cyprus.
giganteum, *Lam. Illustr.* i. 407.—Cilicia; Syria.
Gmelini, Ledeb. Fl. Alt. i. 184 = echioides.
gracile, *Trautv. in Act. Hort. Petrop.* ii. (1873) 478.—Armenia.
graecum, *Boiss. Diagn.* Ser. I. xi. 106.—Graecia.
halophilum, *Boiss. & Heldr. l. c.* 105.—Lycaonia.
hebebulbum, *DC. Prod.* x. 60.—Mesopot.
helveticum, Boiss. Diagn. Ser. I. xi. 111 = stellulatum.
heterophyllum, Griseb. Spicil. Fl. Rumel. ii. 80 = stellulatum.
hispidum, Wall. Cat. n. 938; Stev. ex Ledeb. Fl. Ross. iii. 125 = echioides.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 178.—Reg. Himal.
ibericum, Fisch. ex Ledeb. Fl. Ross. iii. 127 = setosum.
incanescens, DC. Prod. x. 58 = rostellatum.
intermedium, Schur, Enum. Pl. Transs. 469 = setosum.
isauricum, *Boiss. & Heldr. Diagn.* Ser. I. xi. 101.—As. Min.
kilouyense, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* ii. 1.—Persia.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 34.—Persia.
laconicum, Boiss. Pl. Or. Nov. Dec. ii. 8 = stellulatum.
lanceolatum, *Boiss. & Haussk. ex Boiss. l. c.* 1.—Kurdistan.
latifolium, *Boiss. & Haussk. ex Boiss. l. c.* 3.—Kurdistan.
liparioides, *DC. Prod.* x. 65.—Cappadoc.
longilobum, *Bunge, Helioc.* 12.—Oriens.
macrocephala, D. Don, Prod. Fl. Nep. 99 = bracteatum.
megalospermum, Boiss. ex Boiss. Pl. Or. Nov. Dec. i. 8 = Megacaryon armenum.
micranthos, Pall. Reise, ii. 734 = Heliotropium micranthum.
microcarpum, DC. Prod. x. 62 = microspermum.
microspermum, *Stev. in Bull. Soc. Nat. Mosc.* (1838) 305.—As. Min.; Persia.
mite, *Boiss. & Heldr. Diagn.* Ser. I. xi. 111.—Lycia; Cyprus.
molle, *DC. Prod.* x. 61.—Cappadoc.
montanum, Sibth. & Sm. Fl. Graec. Prod. i. 121 = stellulatum.
mutabile, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* ii. 3.—Cataonia.
nanum, *DC. Prod.* x. 65.—Cilicia.
Olivieri, *Boiss. Pl. Or. Nov. Dec.* ii. 2.—Persia.
oreodoxum, *Boiss. & Heldr. Diagn.* Ser. I. xi. 108.—Pisidia.
orientale, Habl. Desc. Phys. Taur. 116 = stellulatum.
orientale, Lehm. Asperif. ii. 376 = frutescens.
orientale, *Linn. Sp. Pl.* ed. II. 196.—Aegypt.
orientale, Pall. Reise, t. 50 = Arnebia cornuta.
Orphanideum, Boiss. Pl. Or. Nov. Dec. ii. 8 = stellulatum.
ovalifolium, *Kotschy & Boiss. ex Boiss. Fl. Orient.* iv. 196.—Cilicia.
pachypodum, *Boiss. Diagn.* Ser. I. xi. 104.—Persia.
pallidum, Boiss. Diagn. Ser. I. xi. 112 = stellulatum.
pallidum, Salisb. Prod. 114 = echioides.
paradoxum, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 180 = stellulatum.
pilosum, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 246.—Syria.
polyanthum, *DC. Prod.* x. 63.—Cappadoc.
polyphyllum, *Ledeb. Ic. Fl. Ross.* i. 7. t. 24.—Tauria.
procerum, *Boiss. Fl. Orient.* iv. 193.—Armenia.
Pseudoarenarium, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 76 = setosum.
pulla, Schrank, ex DC. Prod. x. 67 = Nonnea pulla.
pustulatum, Schur, Enum. Pl. Transs. 469 = stellulatum.
pyrenaicum, Timb. ex Nym. Consp. 516 = echioides.
Rascheyanum, *Boiss. Diagn.* Ser. I. xi. 110.—Syria; Persia.
rigidum, Ledeb. Fl. Alt. i. 180 = stellulatum.
rostellatum, *Lehm. Asperif.* 374.—Mesopot.; Persia.
Roussaei, *DC. Prod.* x. 59.—Syria.
rupestre, *Bieb. Fl. Taur. Cauc.* i. 132.—As. Min.; Reg. Caucas.
saxatile, *Lehm. Asperif.* 371.—Sibir.
scaberrimum, Boiss. Pl. Or. Nov. Dec. ii. 8 = stellulatum.
scorpioides, Boiss. & Huet, Diagn. Ser. II. iii. 137 = stellulatum.

ONOSMA:—
sericeum, Boiss. Diagn. Ser. I. xi. 101 = isauricum.
sericeum, *Willd. Sp. Pl.* i. 774.—As. Min.; Persia.
setigerum, Clarke, Trav. ii. 261 = rostellatum.
setosum, Herb. Unio It. (1828); ex Steud. Nom. ed. II. ii. 217 = echioides.
setosum, *Ledeb. Fl. Alt.* i. 181.—Reg. Caucas.
sibiricum, Lam. Illustr. i. 407 = simplicissimum.
simplex, Gaertn. Fruct. i. 325 = simplicissimum.
simplicissimum, *Linn. Sp. Pl.* ed. II. 196.—Reg. Caucas.; Soongar.
Spruneri, *Boiss. Diagn.* Ser. I. xi. 109.—Graecia.
spathulatum, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 29.—Persia.
stamineum, *Ledeb. in Eichw. Pl. Nov. Casp. Cauc.* 10. t. 3.—Reg. Casp.
Stapfii, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 29.—Persia.
stellulatum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 189. t. 173.—Europ.; Oriens.
stenosiphon, *Boiss. Diagn.* Ser. I. vii. 35.—Persia.
strigosissimum, *Boiss. l. c.* xi. 102.—Pamphylia.
strigosum, H. B. & K. Nov. Gen. et Sp. iii. 93 = Onosmodium strigosum.
strigosum, Stev. ex Boiss. Fl. Orient. iii. 181 = setosum.
suffruticosum, Schrank, ex Steud. Nom. ed. II. ii. 217 = simplicissimum.
synantherum, *Fisch. ex Sweet, Hort. Brit.* ed. II. 375 (Quid?).—Sibir. altaic.
syriacum, *Labill. Pl. Syr. Dec.* iii. 8. t. 5.—Syria; Persia.
Szovitzii, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 596.—Armenia.
tauricum, Pall. in Nov. Act. Petrop. (1792) 306 = stellulatum.
tauricum, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. ii. (1799) 122 = echioides.
taygeteum, *Boiss. & Heldr. Diagn.* Ser. I. xi. 101.—Graecia.
tenuiflorum, Willd. Sp. Pl. i. 775 = rupestre.
tenuifolium, *Tausch, in Syll. Ratisb.* ii. (1828) 246.—Carniola.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 178.—Reg. Himal.
tinctorium, *Bieb. Fl. Taur. Cauc.* i. 131.—Tauria.
tinctorium, Bory & Chaub. Fl. Pélop. 13 = graecum.
Tournefortii, Griseb. Spicil. Fl. Rumel. ii. 80 = frutescens.
trachyanthum, Boiss. Pl. Or. Nov. Dec. ii. 8 = Kotschyi.
trachytricha, Boiss. Diagn. Ser. I. xi. 103 = flavum.
transsilvanicum, Schur, ex Nym. Consp. 516 = setosum.
trapezuntinum, Huet, ex Boiss. Fl. Orient. iv. 201 = stellulatum.
tricerosperma, Lag. Gen. et Sp. Nov. 10 = echioides.
trinervium, Lehm. Asperif. 378. t. 9 = Onosmodium strigosum.
tripartitum, Hochst. in Lorent, Wanderungen, 334 = sericeum.
Troodi, *Kotschy, in Unger & Kotschy, Cypern*, 283.—Ins. Cyprus.
tuberculatum, *Kit. ex Rochel, Pl. Banat. Rar.* 26, nomen; *et in Linnaea*, xxxii. (1863) 436 (Quid?).—Hungar.
vaudense, Gremli, Excurs. Schw. ed. II. 291 = echioides.
velutinum, *Boiss. Diagn.* Ser. I. xi. 105.—Bithynia.
verrucosum, F. G. Dietr. Vollst. Lexik. Gaertn. vi. 465 = echinatum.
versicolor, Schott & Kotschy, ex Boiss. Pl. Or. Nov. Dec. ii. 8 = armenum.
vestitum, Wall. Cat. n. 937 = Emodi.
Visianii, Clem. in Atti 3a Riun. Sc. Ital. Firenze (1852), 519 = setosum.
Wallichianum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 179.—Reg. Himal.
xanthotrichum, Boiss. Diagn. Ser. I. xi. 109 = aleppicum.

ONOSMIDIUM, Walp. Ann. iii. 134 (1852–53), err. typ. = seq.

ONOSMODIUM, Michx. Fl. Bor. Am. i. 132. t. 15 (1803). *BORAGINEAE*, Benth. & Hook. f. ii. 859.
OSMODIUM, Rafin. in Med. Repos. N. York, v. (1808) 352.
PURSHIA, Spreng. Anleit. ii. i. 450 (1817).

ONOSMODIUM:—
bejariense, *A. DC. in DC. Prod.* x. 70.—Texas.
carolinianum, A. DC. l. c. = molle.
carolinianum, Torr. Bot. Mex. Bound. 139 = bejariense.
eriocaulon, *A. DC. in DC. Prod.* x. 70.—Mexic.
hispidum, *Michx. Fl. Am. Bor.* i. 133.—Am. bor.
molle, *Michx. l. c.* t. 15.—Am. bor.
scabrum, *Roem. & Schult. Syst.* iv. 57 (Quid?).—Virginia.
strigosum, *G. Don, Gen. Syst.* iv. 317.—Mexic.
Thurberi, *A. Gray, Syn. Fl. N. Am.* ii. i. 205.—N. Mexic.
virginianum, A. DC. in DC. Prod. x. 70 = hispidum.

ONOSURIS, Rafin. Fl. Ludov. 95 (1817) = **Oenothera**, Linn. (Onagrar.).
acuminata, Rafin. l. c. 96 (Quid?).—Louisiana.
Chamissonis, DC. Prod. iii. 64 = Oe. dentata.

ONOTROPHE, Cass. in Dict. Sc. Nat. xxxvi. 145 (1825) = **Cnicus**, Tourn. (Compos.).
acaulis, Cass. l. c. 146 = C. acaulis.
oleracea, Cass. l. c. = C. oleraceus.
palustris, Cass. l. c. = C. palustris.

ONURIS, Phil. in Anal. Univ. Chil. (1872) 675. *CRUCIFERAE*.
graminifolia, *Phil. l. c.*—Chili.

ONYCHACANTHUS, Nees, in DC. Prod. xi. 217 (1847) = **Bravaisia**, DC. (Acanth.).
arboreus, Karst. Fl. Columb. ii. 111. t. 159.—Venezuela.
Berlandierianus, Nees, in DC. Prod. xi. 217 = B. floribunda.
Cumingii, Nees, l. c. = B. floribunda.
speciosus, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 131 = B. floribunda.

ONYCHIUM, Blume, Bijdr. 323. t. 10 (1825) = **Dendrobium**, Sw. (Orchid.).
affine, Decne. in Nouv. Ann. Mus. Par. iii. (1836) 365 = D. affine.
connatum, Blume, Bijdr. 328 = D. connatum.
crumenatum, Blume, l. c. 326 = D. crumenatum.
fimbriatum, Blume, l. c. 325 = D. Blumii.
flavescens, Blume, l. c. = D. flavescens.
gracile, Blume, l. c. 327 = D. gracile.
japonicum, Blume, l. c. 328 = D. moniliforme.
lamellatum, Blume, l. c. 326 = D. lamellatum.
mutabile, Blume, l. c. 324 = D. mutabile.
nudum, Blume, l. c. 325 = D. nudum.
rigidum, Blume, l. c. 324 = D. firmum.
subulatum, Blume, l. c. 328 = D. subulatum.
tenellum, Blume, l. c. 327 = D. tenellum.
tetraedre, Blume, l. c. = D. tetraedre.
tricuspe, Blume, l. c. 326 = D. tricuspe.

ONYCHOSEPALUM, Steud. Syn. Pl. Cyp. 249 (1855). *RESTIACEAE*, Benth. & Hook. f. iii. 1034.
laxiflorum, *Steud. l. c.*—Austral.

ONYX, Medic. Phil. Bot. i. 207 (1789) = **Astragalus**, Linn. (Legumin.).

OOCARPON, Micheli, in Flora, lvii. (1874) 303. *ONAGRARIAE*.
jussiaeoides, *Micheli, l. c.*—Am. trop.

OOCLINIUM, DC. Prod. v. 133 (1836) = **Eupatorium**, Tourn. (Compos.).
capillare, DC. l. c. 134 = E. capillare.
clavatum, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 108.—Guian.
depressum, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 437 = E. decumbens.
grandiflorum, DC. Prod. v. 134 = E. kleinioides.
paucidentatum, DC. l. c. = E. kleinioides.
pedunculare, DC. l. c. = E. kleinioides.
quinquepartitum, Gardn. ex Baker, in Mart. Fl. Bras. vi. ii. 341 = E. capillare.
rupestre, Gardn. ex Baker, l. c. 343 = E. kleinioides.
rigidum, DC. Prod. v. 134 = E. rigidum.
Sideritis, DC. in Deless. Ic. Sel. iv. t. 14 = E. kleinioides.
villosum, DC. Prod. v. 134 = E. urticaefolium.

OOSTERDYKIA, Burm. Rar. Afr. Pl. 259 (1738) = **Cunonia**, Linn. (Saxifr.).
capensis, Crantz, Inst. ii. 452 = C. capensis.

OPA, Lour. Fl. Cochinch. 308 (1790) = **Eugenia**, Mich. (Myrtac.).
integerrima, Seem. Journ. Bot. i. (1863) 281 = Raphiolepis japonica.
japonica, Seem. l. c. = Raphiolepis japonica.
Mertensii, Seem. l. c. = Raphiolepis indica.
Metrosideros, Lour. Fl. Cochinch. 309 = Raphiolepis indica.
odorata, Lour. l. c. (= *Syzygium odoratum*).—Cochinch.
spiralis, Seem. Journ. Bot. i. (1863) 281 = Raphiolepis indica.

OPALATOA, Aubl. Pl. Gui. t. 147 (APALATOA, in textu, p. 382) (1775) = **Crudia**, Schreb. (Legumin.).

OPANEA, Rafin. Sylva Tellur. 106 (1838) = **Eugenia**, Mich. (Myrtac.).
Billardiana, Rafin. l. c. = Myrcia Billardiana.
trinervia, Rafin. l. c. = E. dumetorum.

OPELIA, Pers. Syn. i. 241 (1805) = **Opilia**, Roxb. (Olacin.).

OPERCULARIA, Gaertn. Fruct. i. 112. t. 24 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 141.
CRYPTOSPERMUM, Young, in Trans. Linn. Soc. iii. (1797) 30.
apiciflora, *Labill. Nov. Holl. Pl.* i. 35. t. 48.—Austral.
aspera, *Gaertn. Fruct.* i. 112. t. 24.—Austral.
aspera, Juss. in Ann. Mus. Par. iv. (1804) 427. t. 70. f. 1 = hispida.
Banksiana, Schult. Mant. iii. 261 = Pomax umbellata.
brachyphylla, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 110 = varia.
coprosmoides, F. Muell. l. c. 108 = hispida.
diphylla, *Gaertn. Fruct.* i. 113.—Austral.
echinocephala, *Benth. Fl. Austral.* iii. 436.—Austral.
ecliptoides, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 110 = varia.
hirsuta, *F. Muell. ex Benth. Fl. Austral.* iii. 434.—Austral.
hirtella, DC. Prod. iv. 616 = hispida?
hispida, *Spreng. Syst.* i. 385.—Austral.
hispidula, *Endl. in Enum. Pl. Hueg.* 58.—Austral.
hispidula, Miq. in Nederl. Kruidk. Arch. iv. (1859) 108 = hispida.
hyssopifolia, Juss. in Ann. Mus. Par. iv. (1804) 428. t. 71 = aspera.
hyssopifolia, Miq. in Nederl. Kruidk. Arch. iv. (1859) 109 = varia.
Lamarckiana, Schult Mant. iii. 261 = Pomax umbellata.
liberiflora, F. Muell. Fragm. iv. 92 = Eleuthranthes opercularina.
ligustrifolia, Juss. in Ann. Mus. Par. iv. (1804) 428. t. 71 = aspera.
micromerioides, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 108 = ovata.
multicaulis, Bartl. in Lehm. Pl. Preiss. i. 369 = vaginata.
ocimifolia, Juss. in Ann. Mus. Par. iv. (1804) 428. t. 71 = aspera.
ovata, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 465.—Austral.
paleata, Young, in Trans. Linn. Soc. iii. (1797) 30. t. 5 = aspera.
pauciflora, Endl. in Enum. Pl. Hueg. 57 = hispidula.
purpurea, Bartl. in Lehm. Pl. Preiss. i. 369 = hispidula.
rubioides, *Juss. in Ann. Mus. Par.* iv. (1804) 428.—Austral.
scabrida, *Schlecht. in Linnaea*, xx. (1847) 604.—Austral.
sessiliflora, Juss. in Ann. Mus. Par. iv. (1804) 427. t. 70 = diphylla.
spermacocea, *Labill. Nov. Holl. Pl.* i. 45. t. 47.—Austral.
Stuartiana, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 110 = varia.
turpis, F. Muell. l. c. 109 = varia.
umbellata, Gaertn. Fruct. i. 112. t. 24 = Pomax umbellata.
vaginata, *Labill. Nov. Holl. Pl.* i. 34. t. 46.—Austral.
varia, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 466.—Austral.
volubilis, *R. Br. ex Benth. Fl. Austral.* iii. 435.—Austral.

OPERCULINA, Silva Manso, Enum. Subst. Bras. 16 (1836) = **Ipomoea**, Linn. (Convolv.).
altissima, Meissn. in Mart. Fl. Bras. vii. 213 = I. altissima.
Convolvulus, Silva Manso, Enum. Subst. Bras. 12, 49 = I. operculata.
flammea, Meissn. in Mart. Fl. Bras. vii. 212 = I. flammea.
pterodes, Meissn. l. c. 213 = I. pterodes.
pteropis, Meissn. l. c. 214.—Am. trop.
surinamensis, Meissn. l. c. 214 = I. surinamensis.
tuberosa, Meissn. l. c. 212 = I. tuberosa.
Turpethum, Silva Manso, Enum. Subst. Bras. 16, 49 = I. Turpethum.

OPETIOLA, Gaertn. Fruct. i. 14. t. 2 (1788) = **Mariscus**, Gaertn. (Cyperac.).
myosuroides, Gaertn. l. c. = M. umbellatus

OPHELIA, D. Don, ex G. Don, Gen. Syst. iv. 178 (1837); D. Don, in Trans. Linn. Soc. xvii. (1837) 522, 524 = **Swertia**, Linn. (Gentian.).
affinis, Wight & Arn. in Wight, Illustr. ii. 175. t. 157, bis = S. affinis.
alata, Griseb. Gen. et Sp. Gent. 321 = S. alata.
alba, Wight & Arn. in Wight, Illustr. t. 157 bis = S. decussata.
angustifolia, D. Don, ex G. Don, Gen. Syst. iv. 178 = S. angustifolia.
Baxteri, Griseb. Gen. et Sp. Gent. 318 (Quid?).—Austral.
bimaculata, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 159 = S. bimaculata.
chinensis, Bunge, ex Griseb. in DC. Prod. ix. 126 = S. diluta.
Chirata, Griseb. Gen. et Sp. Gent. 320 = S. Chirata.
Chirayta, D. Don, ex G. Don, Gen. Syst. iv. 178 = S. cordata.
ciliata, D. Don, ex G. Don, l. c. = S. purpurascens.
coerulescens, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 569.—Java.
cordata, D. Don, ex G. Don, Gen. Syst. iv. 178 = S. cordata.
corymbosa, Griseb. Gen. et Sp. Gent. 317 = S. corymbosa.
Dalhousiana, Griseb. l. c. 313 = S. purpurascens.
densifolia, Griseb. in DC. Prod. ix. 125 = S. decussata.
diluta, Ledeb. Fl. Ross. iii. 73 = S. diluta.
elegans; Wight, Ic. iv. t. 1331 = S. affinis.
fimbriata, Hochst. in Flora, xxvii. (1844) 28 = S. Schimperi.
florida, G. Don, Gen. Syst. iv. 178 = S. pulchella.
gracilis, Bedd. in Trans. Linn. Soc. xxv. (1865) 221.—Ind. or.
Grisebachiana, Wight, Ic. t. 1330 = S. corymbosa.
japonica, Griseb. Gen. et Sp. Gent. 321 = S. rotata.
javanica, Hassk. Cat. Hort. Bog. Alt. 127 = S. javanica.
Lawii, Wight & Arn. in Wight, Illustr. ii. 175 = S. corymbosa.
lucida, G. Don, Gen. Syst. iv. 179, sphalm. = seq.
lurida, D. Don, in Trans. Linn. Soc. xvii. (1837) 527 = S. lurida.
macrosperma, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 448 = S. macrosperma.
minor, Griseb. in DC. Prod. ix. 126 = Pleurogyne minor.
multiflora, Dalz. in Hook. Kew Journ. ii. (1850) 135 = S. decussata.
nervosa, Griseb. Gen. et Sp. Gent. 317 = S. nervosa.
nuda, Klotzsch, in Bot. Ergeb. Waldem. Reise, 91 = S. cordata.
oxyphylla, Miq. Fl. Ind. Bat. ii. 562.—Java.
paniculata, D. Don, in Trans. Linn. Soc. xvii. (1837) 525 = S. paniculata.
papillosa, Franch. & Sav. Enum. Pl. Jap. ii. 450.—Japon.
pauciflora, Dalz. in Hook. Kew Journ. iii. (1851) 211 = S. corymbosa.
porrigens, G. Don, Gen. Syst. iv. 178 = S. pulchella.
pratensis, Edgew. in Trans. Linn. Soc. xx. (1846) 85 = S. pulchella.
pulchella, D. Don, ex G. Don, Gen. Syst. iv. 525 = S. pulchella.
purpurascens, D. Don, in Trans. Linn. Soc. xvii. (1837) 526 = S. purpurascens.

OPHELIA :—
racemosa, Griseb. Gen. et Sp. Gent. 319 = Swertia racemosa.
teres, D. Don, ex G. Don, Gen. Syst. iv. 178.—Reg. Himal.
tetragona, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 122 = S. tetragona.
trichotoma, Wight & Arn. ex Wight, Illustr. sub t. 157 b = S. trichotoma.
trilobata, Pritz. Ic. Ind. 762 = praec.
umbellata, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 446 = S. corymbosa.
vacillans, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 229 = S. vacillans.
Wallichii, G. Don, Gen. Syst. iv. 178 = S. paniculata.
Wilfordii, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 102.—Manchur.
yesoënsis, Franch. & Sav. Enum. Pl. Jap. ii. 451.—Japon.
zeylanica, Griseb. Gen. et Sp. Gent. 316 = S. zeylanica.

OPHELUS, Lour. Fl. Cochinch. 412 (1790) = **Adansonia**, Linn. (Malvac.).
sitularius, Lour. l. c. = A. Situla.

OPHIANTHE, Hanst. in Linnaea, xxvi. (1853) 205 = **Pentarhaphia**, Lindl. (Gesner.).
libanensis, Hanst. l. c. = P. libanensis.

OPHIOCARYON, Schomb. in Endl. Gen. Suppl. i. 1425 (1841). *SABIACEAE*, Benth. & Hook. f. i. 415.
paradoxum, *Schomb. in Hook. Lond. Journ. Bot.* iv. (1845) 377.—Guian.

OPHIOCAULON, Hook. f. in Benth. & Hook. f. Gen. i. 813 (1867). *PASSIFLOREAE*, Benth. & Hook. f. l. c.
cissampeloides, *Hook. f. l. c.* 814.—Afr. trop.
cynanchifolius, *Hook. f. l. c.*—Afr. trop.
gummifer, *Harv. Gen. S. Afr. Pl.* ed. II. 121.—Afr. trop.

OPHIOCAULON, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legumin.).
serpens, Rafin. l. c. = C. serpens.

OPHIOLYZA, Salisb. Gen. Pl. Fragm. 142 (1866) = **Gladiolus**, Tourn. (Irid.).
alata, Salisb. l. c. = G. alatus.
namaquensis, Salisb. l. c. = G. alatus.
orchidiflora, Salisb. l. c. = G. orchidiflorus.
viperata, Salisb. l. c. = G. orchidiflorus.

OPHIOMERIS, Miers, in Proc. Linn. Soc. i. (1847) 328; et in Trans. Linn. Soc. xx. (1851) 373. t. 15 = **Thismia**, Griff. (Burman.).
iguassuensis, Miers, in Proc. Linn. Soc. i. (1847) 329.—Bras.
macahensis, Miers, l. c. 329.—Bras.

OPHIONE, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 101. *AROIDEAE*, Benth. & Hook. f. iii. 997.
Purdieana, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 101.—N. Granat.

OPHIOPOGON, Ker-Gawl. in Bot. Mag. t. 1063 (1807). *HAEMODORACEAE*, Benth. & Hook. f. iii. 678.
CHLOOPSIS, Blume, Enum. Pl. Jav. 14 (1827).
FLUEGGIA, Rich. in Schrad. Neues Journ. ii. (1807) 8. t. 1.
SLATERIA, Desv. Journ. Bot. i. (1808) 243.
gracilis, Kunth, Enum. Pl. v. 298 = Liriope spicata.
indicus, *Royle, Illustr. Bot. Himal.* i. 382, nomen.—Ind. or.
indicus, Wight, Ic. t. 2050 = intermedius.
intermedius, *D. Don, Prod. Fl. Nep.* 48.—Ind. or.
Jaburan, *Lodd. Bot. Cab.* t. 1876.—Japon.
japonicus, *Ker-Gawl. in Bot. Mag.* t. 1063.—Japon.
japonicus, Wall. Cat. n. 5139 A = longifolius.
longifolius, *Decne. in Fl. des Serres*, xvii. (1867–68) 182.—Reg. Himal.
Malcolmsonii, *Royle, Illustr. Bot. Himal.* i. 382.—Burma.
minor, Royle, l. c. = intermedia?
mollis, Royle, l. c. = Theropogon pallidus?

OPHIOPOGON:—
Muscari, Decne. in Fl. des Serres, xvii. (1867-68) 181 = Liriope spicata.
pallidus, Wall. Cat. n. 2138 = Theropogon pallidus.
prolifera, *Lindl. in Journ. Hort. Soc.* i. (1846) 76.—Ins. Penang.
spicatus, Hook. Bot. Mag. t. 5348 = Liriope spicata.
spicatus, Ker-Gawl. in Bot. Reg. t. 593.—Cf. Bot. Mag. sub t. 1063 = Liriope spicata.
Thunbergii, Schult. f. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 151 = Reineckia carnea.
umbraticola, Hance, in Journ. Bot. vi. (1868) 115 = japonicus.

OPHIOPRASON, Salisb. Gen. Fl. Fragm. 72 (1866) = **Asphodelus**, Tourn. (Liliac.).
fistulosum, Salisb. l. c. = A. fistulosus.

OPHIORHIZA, Linn. Fl. Zeyl. 190 (1747) = seq.

OPHIORRHIZA, Linn. Sp. Pl. ed. I. 150 (1753). *RUBIACEAE*, Benth. & Hook. f. ii. 63.
MUNGOS, Adans. Fam. ii. 225 (1763).
acuminata, *DC. Prod.* iv. 416.—Ins. Philipp.
acuminatissima, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 132.—Sumatra.
affinis, *Zoll. & Mor. Syst. Verz. Zoll.* 62.—Java.
albida, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 126.—Sumatra.
angustifolia, Thw. Enum. Pl. Zeyl. 140 = O. Mungos.
argentea, R. Br. in Wall. Cat. n. 6229 = Harrisiana.
australiana, *Benth. Fl. Austral.* iii. 407.—Austral.
Blumeana, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 130.—Java.
bracteata, *Korth. l. c.* 131.—Java.
bracteolata, Wall. Cat. n. 6228 = fasciculata.
Brunonis, *Wight & Arn. Prod.* 404.—Ind. or.
calcarata, *Hook. f. Fl. Brit. Ind.* iii. 84.—Reg. Himal.
canescens, *Blume, Bijdr.* 977.—Java.
cantonensis, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 222.—China.
corymbosa, Buch.-Ham. ex Wall. Cat. n. 6232 = discolor.
Croomii, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 128 = Mitreola sessilifolia.
cryptantha, *Zoll. & Mor. Syst. Verz. Pfl.* 62.—Java.
decumbens, Gardn. ex Thw. Enum. Pl. Zeyl. 419 = Harrisiana.
discolor, *R. Br. in Wall. Cat.* n. 6232 A.—Ins. Penang.
discolor, R. Br. l. c. n. 6232 B = rosea.
elongata, Reinw. ex Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 128 = Richardiana.
eriantha, *Wight, Ic.* t. 1067.—Ind. or.
erubescens, *Wall. Cat.* n. 6233.—Burma.
Eyrei, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 170 = japonica.
falcata, *Bedd. in Madr. Journ. Sc.* Ser. II. xxii. (1861) 72.—Ind. or.
fasciculata, *D. Don, Prod. Fl. Nep.* 136.—Reg. Himal.
filistipula, *Miq. Fl. Ind. Bat.* ii. 351.—Java.
fruticulosa, *Nadeaud, Enum. Pl. Tahiti*, 53.—Ins. Tahiti.
fucosa, *Hance, in Journ. Bot.* xv. (1877) 334.—Cambodia.
geminata, Wall. Cat. n. 6237 = Polyura geminata.
glechomaefolia, *Thw. Enum. Pl. Zeyl.* 140.—Zeylan.
gracilis, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 311.—Burma.
grandiflora, *Wight, Ic.* t. 1069.—Ind. or.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 82.—Reg. Himal.
Harrisiana, *Heyne, ex Wall. Cat.* n. 6236.—Ind. or.
Harrisoniana, Steud. Nom. ed. II. ii. 218, sphalm. = praec.
Hasseltii, *Blume, ex Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 128.—Malaya.
heterophylla, *Jack, in Malay. Misc.* ii. (1822) n. VII. p. 85.—Sumatra.
hirsutula, *Wight, ex Hook. f. Fl. Brit. Ind.* iii. 81.—Ind. or.
hispida, *Hook. f. l. c.* 83.—Reg. Himal.
hispidula, Wall. Cat. n. 6234 = trichocarpa.
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 170 = longiflora.
imbricata, Gardn. ex Hook. f. Fl. Brit. Ind. iii. 81 = pectinata.
inconspicua, *Miq. Fl. Ind. Bat.* ii. 174.—Java.
Jackiana, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 127.—Sumatra.

OPHIORRHIZA:—
japonica, *Blume, Bijdr.* 978.—Japon.
Junghuhniana, *Miq. Fl. Ind. Bat.* ii. 352.—Java.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 232.—Java.
lanceolata, Ell. Sketch, i. 238 = Mitreola petiolata.
lanceolata, Forsk. Fl. Aegypt. Arab. 42 = Mussaenda luteola.
latifolia, *Miq. Fl. Ind. Bat.* ii. 172.—Java.
laxa, *A. Gray, in Proc. Am. Acad.* iv. (1858-59) 312.—Ins. Pacif.
leptanthā, *A. Gray, l. c.*—Ins. Pacif.
littorea, *Miq. Fl. Ind. Bat.* ii. 350.—Java.
longiflora, *Blume, Bijdr.* 977.—Java.
lurida, *Hook. f. Fl. Brit. Ind.* iii. 82.—Reg. Himal.
marginata, *Blume, Bijdr.* 976.—Java.
Mitreola, Linn. Sp. Pl. 150 = Mitreola petiolata.
Mitreola, Michx. Fl. Bor. Am. i. 148 = Mitreola sessilifolia.
Mungos, Bartl. ex DC. Prod. iv. 415 = oblongifolia.
Mungos, *Linn. Sp. Pl.* 150.—Ind. or.; Malaya.
Mungos, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 544 = pectinata.
nana, Edgew. in Trans. Linn. Soc. xx. (1846) 60 = Clarkella nana.
neglecta, *Blume, ex DC. Prod.* iv. 416.—Java.
Nelsoni, *Seem. Fl. Vit.* 126.—Ins. Tahiti.
nemorosa, Thw. Enum. Pl. Zeyl. 139 = O. Mungos.
nutans, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 84.—Reg. Himal.
oblongifolia, *DC. Prod.* iv. 415.—Ins. Philipp.
ochroleuca, *Hook. f. Fl. Brit. Ind.* iii. 78.—Reg. Himal.
oligantha, *Miq. Fl. Ind. Bat.* ii. 350.—Sumatra.
oppositifolia, *Hook. f. Fl. Brit. Ind.* iii. 80.—Reg. Himal.
ovalifolia, Muhl. Cat. 20 = Mitreola sessilifolia.
pallida, *Thw. Enum. Pl. Zeyl.* 140.—Zeylan.
parviflora, *Reinw. ex Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 127.—Ins. Celebes.
pauciflora, *Hook. f. Fl. Brit. Ind.* iii. 84.—Reg. Himal.
pectinata, *Arn. in Nov. Act. Nat. Cur.* xviii. I. (1836) 338.—Ind. or.
peploides, *A. Gray, in Proc. Am. Acad.* iv. (1858-59) 311.—Ins. Pacif.
perpusilla, *Blume, ex DC. Prod.* iv. 417.—Java.
prostrata, D. Don, Prod. Fl. Nep. 136 = Harrisiana.
pumila, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 169.—China.
quadrifida, *Blume, Bijdr.* 977.—Java.
radicans, *Gardn. ex Thw. Enum. Pl. Zel.* 139.—Zeylon.
reticulata, *Korth. in Nederl. Kruid. Arch.* ii. II. (1851) 130.—Sumatra.
Richardiana, *Gaudich. Freyc. Voy. Bot.* 473. t. 97.—Ins. Molucc.
rosea, *Hook. f. Fl. Brit. Ind.* iii. 78.—Reg Himal.
Roxburghiana, *Wight, Ic.* t. 1068.—Ind. or.
rubella, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 126.—Borneo.
rugosa, Blume, Bijdr. 976 = bracteata.
rugosa, Hook.f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 82 = Thomsoni.
rugosa, *Soland. ex Seem. Fl. Vit.* 127.—Ins. Tahiti.
rugosa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 546; Cat. n. 6235 = Harrisiana.
sanguinea, *Blume, Bijdr.* 978.—Java.
scorpioidea, *Nadeaud, Enum. Pl. Tahiti*, 53.—Ins. Tahiti.
Solandri, *Seem. Fl. Vit.* 127.—Ins. Tahiti.
subbracteata, *Korth. in Nederl Kruid. Arch.* ii. II. (1851) 132.—Sumatra.
subcapitata, *Wall. Cat.* n. 6238.—Reg. Himal.
subfalcifolia, *Miq. Fl. Ind. Bat.* ii. 167.—Ins. Celebes.
subumbellata, Bartl. ex DC. Prod. iv. 417 = acuminata.
subumbellata, *Forst. f. Prod.* 12.—Ins. Tahiti.
succirubra, *King, ex Hook. f. Fl. Brit. Ind.* iii. 82.—Reg. Himal.
tahitensis, *Seem. Fl. Vit.* 127.—Ins. Tahiti.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 351.—Java.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iii. 82.—Reg. Himal.
tomentosa, *Jack, in Roxb. Fl. Ind. ed. Carey*, ii. 546.—Ins. Penang; Sumatra.
torrentium, *Nadeaud, Enum. Pl. Tahiti*, 54.—Ins. Tahiti.

OPHIORRHIZA:—
Treutleri, *Hook. f. Fl. Brit. Ind.* iii. 83.—Reg. Himal.
triandra, *Blanco, Fl. Filip.* ed. II. 65.—Ins. Philipp.
trichocarpa, *Blume, Bijdr.* 977.—Ind. or.; Malaya.
villosa, Kurz, in Journ. As. Soc. xlvi. (1877) II. 130 = trichocarpa.
villosa, Reinw. ex Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 132 = longiflora.
villosa, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 702.—Ind. or.
villosa, Wall. Cat. n. 6230 B = fasciculata.
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 79.—Reg. Himal.; Burma.

OPHIORRHIZIPHYLLON, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 76. *ACANTHACEAE*, Benth. & Hook. f. ii. 1074.
macrobotryum, *Kurz, l. c.*—Burma.

OPHIOSCORODON, Wallr. Sched. Crit. 129 (1822) = **Allium**, Tourn. (Liliac.).
magicum, Wallr. l. c., in obs. = A. roseum.
tricoccon, Wallr. l. c. = A. tricoccum.
ursinum, Wallr. l. c. 130 = A. ursinum.

OPHIOSPERMUM, Reichb. Consp. 82 (1828) = OPHISPERMUM, Lour. = **Aquilaria**, Lam. (Thymel.).

OPHIOSTACHYS, [Delile, in] Red. Lil. t. 464 (1816) = **Chamaelirium**, Willd. (Liliac.).
virginica, [Delile,] l. c. = C. luteum.

OPHIOXYLON, Linn. Fl. Zeyl. 239 (1747) = **Rauwolfia**, Plum. (Apocyn.).
album, Gaertn. Fruct. ii. 129 = R. serpentina.
arboreum, Koen. ex Wight & Arn. Prod. 434 = Gaertnera Koenigii.
belgaumense, Wight, Ic. iv. II. 2.—Ind. or.
ceylanicum, Auct. = *zeylanicum*.
chinense, Hance, in Journ. Bot. iii. (1865) 380 (= *Dissolaena verticillata*).—China.
densiflorum, Thw. Enum. Pl. Zeyl. 191 = R. densiflora.
macrocarpum, Wight, Ic. iv. II. 1 = R. densiflora.
majus, Hassk. in Flora, xxviii. (1845) [263 bis] 295.—Java.
micranthum, Wight, ex Hook. f. Fl. Brit. Ind. iii. 633 = R. micrantha.
neilgheerense, Wight, Ic. iv. t. 1292 = R. densiflora.
obversum, Miq. Fl. Ind. Bat. ii. 405 = R. serpentina.
Ochrosia, Pers. Syn. i. 266 = Ochrosia borbonica.
salutiferum, Salisb. Prod. 146 = R. serpentina.
serpentinum, Linn. Sp. Pl. 1043 = R. serpentina.
trifoliatum, Gaertn. Fruct. ii. 123 = R. serpentina.
zeylanicum, Wight, Ic. iv. t. 1291 = R. densiflora.

OPHIRA, Linn. Mant. ii. 150 (1767) = **Grubbia**, Berg. (Santal.).
stricta, Lam. Encyc. iv. 565. [t. 293] = G. stricta.
stricta, Linn. Mant. ii. 229 = G. rosmarinifolia.

OPHIRIA, Becc. in Ann. Jard. Buitenz. ii. (1885) 128 = **Pinanga**, Blume (Palm.).
paradoxa, Becc. l. c. = P. paradoxa.

OPHISMENUS, Poir. Encyc. Suppl. iv. 271 (1816) sphalm. = **Oplismenus**, Beauv. (Gramin.).

OPHISPERMUM, Lour. Fl. Cochinch. 280 (1790) = **Aquilaria**, Lam. (Thymel.).
sinense, Lour. l. c. 281 = A. Ophispermum.

OPHIURINELLA, Desv. Opusc. 75. t. 5. f. 4 (1831) = **Psilurus**, Trin. (Gramin.).
micrantha, Desv. l. c.—Ins. Borbon.

OPHIURUS, Gaertn. f. Fruct. iii. 3. t. 181 (1805); R. Br. Prod. 206 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1130.
DIPERIUM, Desv. Opusc. 76 (1831).
MNESITHEA, Kunth, Rév. Gram. i. 154 (1830).
THYRIDOSTACHYUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 379 (1836).
aethiopicus, Rupr. ex Steud. Syn. Pl. Gram. 360 = papillosus.
appendiculatus, Steud. l. c. 359 = Rottboellia exaltata.
auriculatus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. Math. (1833) 246 = Ratzeburgia pulcherrima.
compressus, *Presl, Cyp. et Gram. Sic.* 51.—Sicil.

OPHIURUS:—
corymbosus, *Gaertn. f. Fruct.* iii. 4. t. 181. f. 4.—Ind. or.; Austral.
cylindricus, Beauv. Agrost. 116 = Lepturus cylindricus.
erectus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 88; Link, Hort. Berol. ii. 172 = Lepturus filiformis.
filiformis, Roem. & Schult. Syst. ii. 797 = Lepturus filiformis.
gracilis, J. Gay, in Schult. Mant. iii. 663, in obs. = Lepturus filiformis.
incurvatus, Beauv. Agrost. 116 = Lepturus incurvatus.
laevis, Benth. in Journ. Linn. Soc. xix. (1881) 69 = perforatus.
loricatus, Nees, ex Steud. Syn. Pl. Gram. 360 = Rottboellia loricata.
monostachyus, *J. & C. Presl, Rel. Haenk.* i. 330.—China; Ins. Philipp.
muricatulus, Steud. Syn. Pl. Gram. 360 = Rottboellia glandulosa.
pannonicus, Beauv. Agrost. 116 = Lepturus pannonicus.
papillosus, *Hochst. in Flora*, xxvii. (1844) 248.—Afr. trop.
perforatus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. *Math.* (1833) 246.—Ind. or.; Malaya.
plicatus, Nees, ex Steud. Nom. ed. II. ii. 218 = Rottboellia loricata.
radicans, Steud. Syn. Pl. Gram. 430 = (Lepturi sp.?).—Ins. Nossibé.
subulatus, Link, Handb. i. 2 = Lepturus cylindricus.
undatus, Nees, in Hook. Kew Journ. ii. (1850) 100 = monostachyus.
undulatus, Benth. in Journ. Linn. Soc. xix. (1881) 69, sphalm. = praec.

OPHRIS, [Tourn.] Linn. Syst. ed. I (1735) = **Ophrys**, Linn. (Orchid.).

OPHRYOCOCCUS, Oerst. in Kjoeb. Vidensk. Meddel. (1852) 52. *RUBIACEAE*, Benth. & Hook. f. ii. 70.
gesnerioides, *Oerst. l. c.* 53.—Nicaragua.

OPHRYOSCLERIA, Nees, in Mart. Fl. Bras. ii. 1. 182. t. 25 (1842) = **Scleria**, Berg. (Cyperac.).
asperrima, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 261 = S. asperrima.
lucida, Nees, in Mart. Fl. Bras. ii. 1. 183 = S. mitis.
microcarpa, Nees, l. c. 184 = S. microcarpa.
mitis, Nees, l. c. 185 = S. mitis.
paludosa, Nees, l. c. = S. paludosa.
racemosa, Nees, l. c. 183 = S. racemosa.
riparia, Nees, l. c. 184 = S. riparia.
Schiedeana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 261 = S. Schiedeana, *Schlecht.* (p. 489, non 494).
sororia, Nees, in Mart. Fl. Bras. i. 11. 184 = S. sororia.

OPHRYOSPORUS, Meyen, Reise, i. 402 (1834). *COMPOSITAE*, Benth. & Hook. f. ii. 239.
OPHYROSPORUS, Baker, in Mart. Fl. Bras. vi. 11. 186 (1876).
Burchellii, *Baker, in Mart. Fl. Bras.* vi. 11. 187.—Bras.
clavulatus, *Griseb. in Goett. Abh.* xxiv. (1879) 173.—Reg. Argent.
Cumingii, *Benth. & Hook. f. Gen.* ii. 239.—Bolivia.
Freyreissii, *Baker, in Mart. Fl. Bras.* vi. 11. 188.—Bras.
laxiflorus, *Baker, l. c.* 189.—Bras.
macrodon, *Griseb. in Goett. Abh.* xxiv. (1879) 173.—Reg. Argent.
Mandonii, Benth. & Hook. f. Gen. ii. 239 = piquerioides.
ovatifolius, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 79.—Mexic.
pachychaeta, *Baker, in Mart. Fl. Bras.* vi. 11. 187.—Bras.
paradoxus, *Benth. & Hook. f. Gen.* ii. 239.—Chili.
piquerioides, *Benth. & Hook. f. l. c.*—Peruv.
Regnellii, *Baker, in Mart. Fl. Bras.* vi. 11. 188.—Bras.
triangularis, *Meyen, Reise*, i. 402.—Chili.

OPHRYS, Linn. Gen. ed. I. 272 (1737). *ORCHIDEAE*, Benth. & Hook. f. iii. 621.
ARACHNITES, F. W. Schmidt, Fl. Boëm. i. 74 (1793), partim.
MYODIUM, Salisb. in Trans. Hort. Soc. i. (1812) 289.
adrachnites, Mill. Gard. Dict. ed. VIII. n. 7 = O. Arachnites.
aestiva, Balb. Elench. 96 = Spiranthes aestivalis.

OPHRYS:—
aestivalis, Michx. Fl. Bor. Am. ii. 157 = Spiranthes tortilis.
aestivalis, Poir. in Lam. Encyc. iv. 567 = Spiranthes aestivalis.
aestrifera, Boiss. Fl. Orient. v. 79 = oestrifera.
alaris, Linn. f. Suppl. 404 = Pterygodium catholicum.
alata, Linn. l. c. = Pterygodium alatum.
albiflora, Sprun. ex Boiss. Fl. Orient. v. 79 = apifera.
aloidea, Poir. in Lam. Encyc. iv. 559 = guianensis.
alpina, Linn. Sp. Pl. 948 = Herminium alpinum.
andracnitis, Bory & Chaub. Nouv. Fl. Pélop. 62. t. 34 = O. Ferrum-equinum.
anthropomorpha, Willd. Sp. Pl. iv. 63 = Aceras anthropophora.
anthropophora, Linn. Sp. Pl. 948 = Aceras anthropophora.
anthropophora, Oed. Fl. Dan. t. 103 = Orchis ustulata.
apicula, *J. C. Schmidt, ex Reichb. f. Ic. Fl. Germ.* xiii. 79 (an hybr.?).—Helvet.
apiculata, Rich. in Mém. Mus. Par. iv. (1818) 55 = O. Arachnites, bombyliflora.
apifera, *Huds. Fl. Angl.* ed. I. 340.—Europ.; Afr. bor.
apiformis, Steud. Nom. ed. I. 567 = sphegifera.
aquisgranensis, *Kaltenb. in Verh. Rheinl.* vii. (1850) 519. t. 7.—Europ.
arachnis, *Vell. Fl. Flum.* ix. t. 60.—Bras.
Arachnites, *Lam. Fl. Fr.* iii. 515; *Murr. Syst.* ed. XIV. 813.—Europ.; As. Min.; Syria.
Arachnites, Link, in Schrad. Journ. ii. (1799) 325 = tenthredinifera.
Arachnites, Reichard, Fl. Moeno-Franc. ii. 89 = apifera.
arachnitiformis, Gren. & Philippi, in Mém. Soc. Emul. Doubs, Sér. III. iv. (1860) 391 = O. Arachnites.
arachnoides, *Vell. Fl. Flum.* ix. t. 59.—Bras.
Araneola, Reichb. Pl. Crit. ix. 22; Fl. Germ. Excurs. 140^{19} = aranifera.
aranifera, *Huds. Fl. Angl.* ed. II. ii. 392.—Europ.; As. Min.
argentea, *Vell. Fl. Flum.* ix. t. 61.—Bras.
arundinacea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 = Arundina densiflora.
asilifera, *Vayr. in Anal. Soc. Esp. Hist. Nat.* ix. (1880) 98.—Hispan.
atlantica, *Munby, in Bull. Soc. Bot. Fr.* iii. (1856) 108.—Algeria.
atrata, *Lindl. Bot. Reg.* t. 1807.—Europ. austr.; As. Min.
atrata, Linn. Mant. i. 121 = Ceratandra chloroleuca.
Atropos, Barla, Ic. Orch. t. 59 = O. Scolopax.
attica, Boiss. & Orph. Diagn. Ser. II. iv. 91 = O. Arachnites.
aurita, *Brot. Phyt. Lusit.* ii, 34. t. 91. f. II.—Lusit.
austriaca, Wiesb. ex Dichtl, in Deutsch. Bot. Monatsb. i. (1883) 148 = apifera.
autumnalis, Balb. Elencho, 96 = Spiranthes autumnalis.
barbata, Walt. Fl. Carol. 221 = Calopogon pulchellus.
Bertolonii, *Moretti, Dec.* vi. 9; *in Brugn. Giorn. Fis.* ii. (1823).—Italia.
Biancae, Macch. in Nuov. Giorn. Bot. xiii. (1881) 315 = O. Arachnites.
biceratia, Delile, ex Nym. Consp. 697 = atrata.
bicolor, Thunb. Prod. Pl. Cap. 2 = Corycium bicolorum.
bicornis, Sadl. ex Nendtv. Pl. Quinq. Eccl. (1836) 35.—Cf. in Linnaea, x. (1836) Litt. 205 = oestrifera.
bifolia, Lam. Fl. Fr. iii. 516; Gilib. Exercit. ii. 486 = Listera ovata.
bilunulata, *Risso, Fl. Nice*, 463.—Gallia.
bivalvata, Linn. f. Suppl. 403 = Disa melaleuca.
bombylifera, Willd. Sp. Pl. iv. 68 = seq.
bombyliflora, *Link, in Schrad. Journ.* ii. (1799) 325.—Reg. Mediterr.
bombyliflora, Spreng. Syst. iii. 702 = O. Arachnites.
brachyotus, Reichb. Fl. Germ. Excurs. 128 = O. Arachnites.
bracteata, Linn. f. Suppl. 403 = Satyrium bracteatum.
bremifera, Stev. ex Bieb. Fl. Taur. Cauc. i. 370, in obs.; et in Mém. Soc. Nat. Mosc. ii. (1809) 174 = oestrifera.
Broteroi, Welw. ex Nym. Consp. 698 = apifera.
bullata, Cyril. ex Steud. Nom. ed. II. ii. 219 = O. Scolopax.
caffra, Linn. Pl. Rar. Afr. 28; Amoen. Acad. vi. 111 = Pterygodium caffrum.
campestris, *Vell. Fl. Flum.* ix. t. 58.—Bras.
camtschatea, Linn. Sp. Pl. 948 = Neottia camtschatica.

OPHRYS:—
canaliculata, Viv. App. Fl. Cors. Prod. 7 = bombyliflora.
caroliniana, Raeusch. Nom. ed. III. 261 = Liparis liliifolia?
catholica, Linn. Pl. Rar. Afr. 27; Amoen. Acad. vi. 110 = Pterygodium catholicum.
cernua, Linn. Sp. Pl. ed. II. 946 = Spiranthes cernua.
chlorantha, Hegetschw. Fl. Schw. 876 = apifera.
ciliata, Bivona, Sic. Pl. Cent. i. 60 = O. Speculum.
ciliata, H. B. & K. Nov. Gen. et Sp. i. 334. t. 74 = Cranichis ciliata.
circumflexa, Linn. Pl. Afr. Rar. 27; Amoen. Acad. vi. 111 = Disperis secunda.
Corallorhiza, Linn. Sp. Pl. 945 = Corallorhiza innata.
Corallorhiza, Michx. Fl. Bor. Am. ii. 158 = Corallorhiza odontorhiza.
cordata, Linn. Sp. Pl. 946 = Listera cordata.
corniculata, Brot. Phyt. Lusit. ii. 38. t. 93 = O. Scolopax.
cornuta, Stev. ex Bieb. Fl. Taur. Cauc. i. 370, in obs.; et in Mém. Soc. Nat. Mosc. ii. (1809) 175 = oestrifera.
corsica, *Viv. Fl. Cors. Sp. Nov.* 16.—Corsica.
corsica, Soleir. ex Bertol. Fl. Ital. ix. 595 = lutea.
crabronifera, Sebast. & Mauri, Rom. Pl. Cent. 13. t. 2. f. 1 = O. Arachnites.
cristata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 (Quid?).—Java.
crucigera, Jacq. Ic. Pl. Rar. i. 18. t. 185 = aranifera.
cyanogramma, Welw. ex Nym. Consp. 697 = atrata.
densiflora, Desf. in Ann. Mus. Par. x. (1807) 228. t. 16 = Habenaria intacta.
devenensis ×, Reichb. f. Ic. Fl. Germ. xiii-xiv. 87.—Europ.
discors, Bianca, in Todaro, Orch. Sic. 84 = O. Arachnites.
disthoma, Bivona, Sic. Pl. Cent. i. 59 = bombyliflora.
distoma, Tenore, Syll. 131 = praec.
epeirophora ×, Peter, in Flora, lxvi. (1883) 10.—Europ.
episcopalis, Poir. Encyc. Suppl. iv. 170 = tenthredinifera.
exaltata, Tenore, Cat. Pl. Hort. Neap. (1819) App. ii. 83 = O. Arachnites.
Ferrum-equinum, *Desf. in Ann. Mus. Par.* x. (1807) 226. t. 15.—Graecia; Reg. Caucas.
fimbriata, Walt. Fl. Carol. 221 = Habenaria fissa?
flavicans, Vis. Fl. Dalm. i. 178 = tenthredinifera.
fucifera, Sm. Engl. Bot. t. 2649 = aranifera.
fuciflora, Curt. Fl. Lond. fasc. vi. t. 67 = aranifera.
fuciflora, Moench, Meth. Suppl. 311 = O. Arachnites.
funerea, Viv. Fl. Cors. Sp. Nov. 15 = fusca.
fusca, *Link, in Schrad. Journ.* ii. (1799) 324.—Reg. Mediterr.
fusca, Reichb. Ic. Pl. Germ. xiii. 73 = tenthredinifera.
fusca, Tenore, Fl. Nap. ii. 303. t. 92 = lutea.
galactostictos, Heldr. & Sart, ex Boiss. Fl. Orient. v. 79 = Spruneri.
glabra, Pers. Syn. ii. 510 = vespifera.
grandiflora, Tenore, Cat. Pl. Hort. Neap. (1819) App. ii. 83; Fl. Nap. ii. 309 = tenthredinifera.
grassensis, Jauvy, ex Steud. Nom. ed. I. 568 (Quid?).—Gallia.
guianensis, *Aubl. Pl. Gui.* ii. 816.—Guian.
Herminium, Gren. in Gren. & Godr. Fl. Fr. iii. 300 = Herminium Monorchis.
hiulca, Sebast. & Mauri, Cent. Rom. Pl. xiii. 43 = bombyliflora.
hiulca, Sprun. ex Reichb. f. Ic Fl. Germ. xiii. 93. t. 101. f. 2 = Spruneri.
hybrida ×, Pokorny, ex Reichb. f. l. c. 79.—Austria.
incubacea, Bianca, ex Tod. Orch. Sic. 75 = atrata.
insectifera, Gueldenst. Reisen, i. 422 = oestrifera.
insectifera, Linn. Sp. Pl. 948 = sp. plurimae.
integra, *Sacc. in Nuov. Giorn. Bot.* iii. (1871) 165.—Italia.
inversa, Thunb. Prod. Pl. Cap. 2 = Pterygodium inversum.
Insengae, Nym. Syll. Suppl. 61 = Bertolonii.
iricolor, Desf. in Ann. Mus. Par. x. (1807) 224. t. 13 = fusca.
kamtschatica, Georgi, Beschr. Russ. Reich. v. 1273 = Neottia camtschatica.
labrofossa, Brot. Phyt. Lusit. ii. 29. t. 88 = bombyliflora.
laetea, Willk. & Lange, Prod. Fl. Hisp. i. 174 = fusca.
lancea, *Thunb. ex Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 223.—Java.

OPHRYS:—
lilifolia, Huds. Fl. Angl. ed. I. 339; ed. II. 389 = Liparis Loeselii.
lilifolia, Linn. Sp. Pl. 946 = Liparis liliifolia.
limbata, Link, Handb. i. 247 = Serapias Lingua.
Loeselii, Linn. Sp. Pl. 947 = Liparis Loeselii.
lunulata, *Parl. in Giorn. Sc. Sicil.* lxii. (1838) 4.—Europ. austr.
lutea, Bivona, Cent. ii. 41. t. 5 = fusca.
lutea, *Cav. Ic.* ii. 46. t. 160.—Reg. Mediterr.
macrostachya, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 9 = Microstylis macrostachya.
mammosa, Desf. in Ann. Mus. Par. x. (1807) 222. t. 12 = aranifera.
monophyllos, Linn. Sp. Pl. 947 = Microstylis monophyllos.
monophyllos, Pav. ex Lindl. Gen. et Sp. Orch. 23 = Microstylis macrostachya.
Monorchis, Linn. Sp. Pl. 947 = Herminium Monorchis.
muscaria, Lam. Fl. Fr. iii. 515 = muscifera.
muscaria, Pall. Reise Suedl. Statth. Russ. Reich. ii. 172 = oestrifera ?
muscifera, *Huds. Fl. Angl.* ed. I. 340.—Europ.
myodes, Jacq. Misc. ii. 273 = muscifera.
myodes, Lapeyr. Hist. Abr. Pl. Pyren. ii. 551 = fusca.
neglecta, *Parl. Fl. Ital.* iii. 548.—Italia; Pyren.
nervosa, Thunb. Fl. Jap. 27 = Liparis nervosa.
Nidus-avis, Linn. Sp. Pl. 945 = Neottia Nidus-avis.
obscura, *Beck, in Oestr. Bot. Zeitschr.* xxix. (1879) 353.—Europ.
oestrifera, *Bieb. Fl. Taur. Cauc.* ii. 369; *Stev. in Mém. Soc. Nat. Mosc.* ii. (1809) 176. t. 11. f. 4, 5.—Europ. austr.; Oriens.
oestrifera, Reichb. Fl. Germ. Excurs. 128 = O. Arachnites.
ovata, Linn. Sp. Pl. 946 = Listera ovata.
oxyrhynchos, Todaro, in Imparziale Giorn. Sc. Lett. Art. (1840) 74; ex eod. Orch. Sic. 82 = O. Arachnites.
paleacea, H. B. & K. Nov. Gen. et Sp. i. 334 = Altensteinia paleacea.
pallida, *Rafin. Caratteri*, 87.—Sicilia.
paludosa, Linn. Sp. Pl. 947 = Malaxis paludosa.
paludosa, O. F. Muell. in Fl. Dan. t. 877 = Liparis Loeselii.
palustris, Huds. Fl. Angl. ed. I. 339 = praec.
patens, Linn. f. Suppl. 404 = Disa tenuifolia.
Pectus, *Mutel, in Ann. Sc. Nat.* Sér. II. iii. (1835) 242.—Algeria.
peruviana, Aubl. Pl. Gui. ii. 816 (Spiranthes sp. ?).—Peruv.
Philippi, Gren. in Mém. Soc. Emul. Doubs, Sér. III. iv. (1860) 401 = oestrifera.
picta, Bory & Chaub. Nouv. Fl. Pélop. 62. t. 33; ex Boiss. Fl. Orient. v. 80 = oestrifera.
picta, Link, in Schrad. Journ. ii. (1799) 325 = O. Scolopax.
picta, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 673 = bombyliflora.
prostrata, Steud. Nom. ed. II. ii. 249, sphalm. = *rostrata*.
Pseudospeculum, DC. Fl. Fr. Suppl. 332 = aranifera.
pubescens, *Presl, Rel. Haenk.* 91.—Peruv.
pubescens, Reinw. ex Lindl. in Journ. Linn. Soc. iii. (1859) 15 = Dendrobium truncatum.
pulchella, Salisb. Prod. 8 = Liparis Loeselii.
pulla, Tenore, Fl. Nap. Prod. p. liii; ii. 311. t. 97 = bombyliflora.
purpurea, Tausch, in Flora, xiv. (1831) 222 = apifera.
quinquelobata, Poir. in Lam. Encyc. iv. 568 = peruviana.
rosea, Desf. ex Boiss. Fl. Orient. v. 77 = tenthredinifera.
rostrata, Tenore, Ind. Sem. Hort. Neap. (1830) 15 = apifera.
Reinholdi, Sprun. ex Boiss. Fl. Orient. v. 79 = Spruneri.
Salassia, Comm. ex A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 47 = Liparis purpurascens.
Scolopax, Bory & Chaub. Nouv. Fl. Pélop. 62 t. 24. f. 4 = oestrifera.
Scolopax, *Cav. Ic.* ii. 46. t. 161.—Reg. Mediterr.
Scolopax, Tenore, Fl. Neap. Prod. 53; Fl. Nap. ii. 396 = Bertolonii.
Scolopax, Willd. Sp. Pl. iv. 69 = O. Speculum.
secundiflora, Steud. Nom. ed. II. i. 768; ii. 219 = Habenaria intacta.
sicula, Tineo, Pl. Rar. Sicil. 13 = lutea.
Speculum, Bertol. Amoen. Ital. 201 = Bertolonii.

OPHRYS:—
Speculum, *Link, in Schrad. Journ.* ii. (1799) 324.—Reg. Mediterr.
Speculum, Steud. Nom. ed. II. ii. 219 = lutea.
sphegifera, *Willd. Sp. Pl.* iv. 65.—Afr. bor.
sphegodes, Mill. Gard. Dict. ed. VIII. n. 8 = apifera ?
spiralis, Georgi, Reise, i. 232 = Spiranthes australis.
spiralis, Linn. Sp. Pl. 945 = Spiranthis sp. variae.
Spruneri, *Nym. Syll.* 360.—Graecia.
squamata, Forst. f. Prod. 59 = Dipodium squamatum.
tabanifera, Sieber, ex Nym. Consp. 697 = tenthredinifera.
tabanifera, Willd. Sp. Pl. iv. 68 = bombyliflora.
Tenoreana, Lindl. Bot. Reg. t. 1093 = tenthredinifera.
tenthredinifera, Tenore, Fl. Nap. i. 308. t. 93 = neglecta.
tenthredinifera, *Willd. Sp. Pl.* iv. 67.—Reg. Mediterr.
Todaroana ×, Macch. in Nuov. Giorn. Bot. xiii. (1881) 314.—Sardin.
Tommasinii, *Vis. Fl. Dalm.* iii. 354.—Dalmat.
tricolor, Desf. ex Nym. Consp. 699 = fusca.
trifolia, Walt. Fl. Carol. 221 = Liparis liliifolia.
trigona, Gilib. Exercit. ii. 488 = Liparis Loeselii.
triphylla, Thunb. Prod. Pl. Cap. 2 = Pterygodium Volucris.
Trollii, Hegetschw. Fl. Schw. 874. t. 8 = apifera.
truncata, Dulac, Fl. Hautes-Pyr. 128 = O. Arachnites.
umbilicata, Desf. in Ann. Mus. Par. x. (1807) 227. t. 15 = bombyliflora.
unifolia, Forst. f. Prod. 59 = Microtis porrifolia.
unilateralis, Poir. in Lam. Encyc. iv. 569 = Spiranthes diuretica.
vernixia, Brot. Fl. Lusit. i. 24 = O. Speculum.
vespifera, Brot. l. c. = lutea.
vespifera, *Willd. Sp. Pl.* iv. 65.—Barbar.
vetula, *Risso, Fl. Nice*, 464.—Gallia.
villosa, Desf. in Ann. Mus. Par. x. (1807) 225. t. 14 = tenthredinifera.
Volucris, Linn. f. Suppl. 403 = Pterygodium Volucris.
Volucris, Thunb. Prod. Pl. Cap. 2 = Corycium vestitum.

OPHTHALMACANTHUS, Nees, in DC. Prod. xi. 219 (1847) = **Ruellia**, Plum. (Acanth.).
gemmiparus, Nees, l. c. 220 = R. gemmipara.
parvus, Nees, l. c. = R. parva.
roseus, Nees, l. c. = R. rosea.
speciosus, Nees, l. c. = R. pulcherrima.

OPHTHALMOBLAPTON, Fr. Allem. Tract. Rio Jan. (1849) 4. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 333.
brasiliense, *Walp. Ann.* iii. 362.—Bras.
crassipes, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 532.—Bras.
macrophyllum, *Fr. Allem. Tract. Rio Jan.* 4.—Bras.
megalophyllum, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 707.—Bras.
pedunculare, *Muell. Arg. l. c.* 533.—Bras.

OPHYOSTACHYS, Steud. Nom. ed. II. ii. 220 (1841) = OPHIOSTACHYS, Delile = **Chamaelirium**, Willd. (Liliac.).

OPHYRA, Steud. Nom. ed. II. ii. 220 (1841) = OPHIRA, Linn. = **Grubbia**, Berg. (Santal.).

OPHYROSPORUS, Baker, in Mart. Fl. Bras. vi. II. 186 (1876), sphalm. = **Ophryosporus**, Meyen (Comp.).

OPICRINA, Rafin. New Fl. Am. iv. 85 (1836). Gen. dub. *COMPOSITARUM*.
angustifolia, Rafin. l. c. = *crepidinea*.
crepidinea, Rafin. l. c.—Am. bor.
latifolia, Rafin. l. c.—Am. bor.

OPILIA, Roxb. Pl. Corom. ii. 31. t. 158 (1799). *OLACINEAE*, Benth. & Hook. f. i. 350.
GROUTIA, Guill. & Perr. Fl. Seneg. Tent. 100. t. 22 (1832).
PENTITDIS, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 246 (1850).
acuminata, Wall. Cat. n. 7206 F = Lepionurus oblongifolius.
amentacea, *Roxb. Pl. Corom.* ii. 31. t. 158.—Ind. or.
amentacea, Wall. Cat. n. 2331 = Cansjera Rheedii.
celtidifolia, Endl. ex Walp. Rep. i. 377 = amentacea.
Cumingiana, *Baill. Adansonia*, iii. (1862–63) 124.—Ins. Philipp.
javanica, Miq. Fl. Ind. Bat. i. 1. 784 = amentacea.

OPILIA:—
manillana, *Baill. Adansonia*, iii. (1862–63) 124.—Ins. Philipp.
odorata, Spreng. Syst. i. 766 = Aglaia odorata.
odoratissima, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 43 = Aglaia Diepenhorstii.
Pentitdis, *Blume, Mus. Bot. Lugd. Bat.* i. 246.—N. Guin.
senegalensis, *Baill. Adansonia*, ii. (1861–62) 381. t. 10.—Afr. trop.
umbellulata, *Baill. l. c.* viii. (1868) 199.—Afr. trop.

OPISTHOCENTRA, Hook. f. in Benth. & Hook. f. Gen. i. 749 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
clidemioides, *Hook. f. l. c.*—Bras.

OPIZIA, J. & C. Presl, Rel. Haenk. i. 293. t. 41. f. 1 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1173.
stolonifera, *Presl, l. c.*—Mexic.

OPIZIA, Rafin. New Fl. Am. ii. 29 (1836) = **Capsella**, Moench (Crucif.).
bursoides, Rafin. l. c. = C. Bursa-pastoris.

OPLEXION, Rafin. Fl. Tellur. iv. 86 (1836) = **Lobostemon**, Lehm. (Borag.).
ferox, Rafin. l. c. = L. ferocissimus.

OPLISMENUS, Beauv. Fl. Owar. ii. 14. t. 58 (1807). *GRAMINEAE*, Benth. & Hook. f. iii. 1104.
HEKATEROSACHNE, Steud. Syn. Pl. Gram. 118 (1855).
OPHISMENUS, Poir. Encyc. Suppl. iv. 271 (1816).
ORTHOPOGON, R. Br. Prod. 194 (1810).
PANICULATUM, Ard. Animad. ii. 14 (1764).
abortivus, Auct. ex Roem. & Schult. Syst. ii. 808 = Andropogon squarrosus.
acuminatus, Nees, ex Steud. Syn. Pl. Gram. 45 = undulatifolius.
aemulans, Roem. & Schult. Syst. ii. 487 = undulatifolius.
aemulus, Kunth, Rév. Gram. i. 44 = undulatifolius.
affinis, J. & C. Presl, Rel. Haenk. i. 323 = Burmannii.
affinis, Schult. Mant. ii. 273 = Panicum Schultesii.
africanus, Beauv. Fl. Owar. ii. 15. t. 67 = compositus.
albus, Roem. & Schult. Syst. ii. 890 = Burmannii.
angustifolius, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 501 (Panici sp.).—Mexic.
brasiliensis, Raddi, Agrost. Bras. 40 = compositus.
brevisetus, Nees, ex Steud. Syn. Pl. Gram. 46 = Panicum brevisetum.
bromoides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 452.—Madag.
bromoides, Beauv. Agrost. 54 = Burmannii.
Burmannii, *Beauv. l. c.*—Amphig. trop.
capensis, *Hochst. in Flora*, xxix. (1846) 114.—Afr. austr.
chondrosioides, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 501 (Panici sp.).—Mexic.
colonus, H. B. & K. Nov. Gen. et Sp. i. 108 = Panicum colonum.
compositus, *Beauv. Agrost.* 54.—Cosmop. trop.
cristatus, J. & C. Presl, Rel. Haenk. i. 323 = Burmannii.
Crus-galli, Dum. Obs. Gram. Belg. 138 = Panicum Crus-galli.
Crus-pavonis, H. B. & K. Nov. Gen. et Sp. i. 108 = Panicum Crus-galli.
cubensis, Kunth, Rév. Gram. i. 45 = Panicum cubense.
cuspidatus, Kunth, l. c. = Panicum cuspidatum.
decompositus, Nees, in Endl. Prod. Fl. Norf. 19 = compositus.
depauperatus, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 501.—Mexic.
discolor, Kunth, Rév. Gram. i. 45 = Panicum discolor.
dubius, *Kunth, l. c.* 44.—Ind. or.
echinatus, Kunth, l. c. 45 = Panicum Crus-galli.
elatior, Beauv. Agrost. 54 = compositus.
erianthos, Kunth, Rév. Gram. i. 45 = Panicum Crus-galli ?
fasciculatus, Roem. & Schult. Syst. ii. 487 = Panicum fasciculiflorum.
festucoides, Kunth, Enum. Pl. i. 146 = Chaetium festucoides.
flaccidus, Roem. & Schult. Syst. ii. 487 = undulatifolius.
foliaceus, Beauv. Agrost. 54 = compositus.

OPLISMENUS:—
frumentaceus, Kunth, Rév. Gram. i. 45 = Panicum Crus-galli.
gracilis, *Schlecht. in Linnaea*, xxiv. (1851) 649.—Hab.?
helvolus, Beauv. Agrost. 54 = Setaria helvola.
hirsutus, Schult. Mant. ii. 271 = Panicum javanicum.
hirtellus, Roem. & Schult. Syst. ii. 481 = compositus.
hirtiflorus, J. & C. Presl, Rel. Haenk. i. 320 = rariflorus?
hirtus, Heyne, ex Roem. & Schult. Syst. ii. 484 = Panicum Rothii.
hispidulus, Kunth, Rév. Gram. i. 44 = Panicum Crus-galli.
holciformis, H. B. & K. Nov. Gen. et Sp. i. 107 = Panicum holciforme.
Hookeri, Parl. in Atti 8va Riun. Sc. Ital. (1846) 586 = Burmannii.
Humboldtianus, Nees, Agrost. Bras. 264 = Burmannii.
imbecillis, Roem. & Schult. Syst. ii. 487 = undulatifolius.
indicus, Roem. & Schult. l. c. 484 = compositus.
intermedius, Kunth, Rév. Gram. i. 45 = Setaria italica.
Jacquini, Sieber, ex Steud. Syn. Pl. Gram. 45 = compositus.
jamaicensis, Kunth, Enum. Pl. i. 147 = Panicum spectabile.
javanicus, *Klotzsch, ex Schlecht. in Linnaea*, xxxi. (1861–62) 309.—Java.
javanicus, Roem. & Schult. Syst. ii. 891 = Panicum javanicum.
lanceolatus, Kunth, Rév. Gram. i. 45 = compositus.
latifolius, Haenke, ex Steud. Nom. ed. II. ii. 220 = compositus.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 502.—Mexic.
limosus, J. & C. Presl, Rel. Haenk. i. 321 = Panicum Crus-galli.
loliaceus, Beauv. Agrost. 168 = compositus.
longisetus, Kunth, Rév. Gram. i. 45 = Panicum Crus-galli.
magellanicus, Roem. & Schult. Syst. ii. 485 = Centotheca lappacea.
margaritaceus, Kunth, Rév. Gram. i. 44 = Panicum margaritaceum.
Minarum, Nees, Agrost. Bras. 268 = Ichnanthus Minarum.
mollissimus, Hochst. ex Steud. Syn. Pl. Gram. 44 = Burmannii.
multisetus, A. Rich. Tent. Fl. Abyss. ii. 377 = Burmannii.
muricatus, Kunth, Rév. Gram. i. 44 = Panicum Crus-galli.
oahuaensis, Nees & Meyen, ex Steud. Nom. ed. II. ii. 220 = compositus.
parvifolius, Kunth, Rév. Gram. i. 45 = undulatifolius.
penicillatus, Kunth, l. c. = Ichnanthus bambusiflorus.
pictus, Kunth, Enum. Pl. i. 144 = Panicum Crus-galli.
polystachyus, *H. B. & K. Nov. Gen. et Sp.* i. 107.—Mexic.
pratensis, Schult. Mant. ii. 597 = compositus.
Preslei, Kunth, Enum. Pl. i. 141 = Burmannii.
prostratus, Edgew. in Journ. Linn. Soc. vi. (1862) 195 = Panicum prostratum.
Pseudocolonus, Kunth, Rév. Gram. i. 44 = Panicum colonum.
Pseudoundulatifolius, Kunth, l. c. = Panicum Pseudoundulatifolium.
rariflorus, *J. & C. Presl, Rel. Haenk.* i. 320.—Mexic.
repens, J. & C. Presl, l. c. 321 = Panicum prorepens.
sabulicolus, Kunth, Enum. Pl. i. 145 = Panicum Crus-galli.
scaber, Kunth, Rév. Gram. i. 44 = Panicum Crus-galli.
secundus, *J. & C. Presl, Rel. Haenk.* i, 322.—Peruv.
semialatus, Desv. Opusc. 81 = Panicum semialatum.
setarius, Roem. & Schult. Syst. ii. 481 = undulatifolius.
spectabilis, Kunth, Enum. Pl. i. 145 = Panicum spectabile.
stagninus, Kunth, Rév. Gram. i. 44 = Panicum Crus-galli.
strictus, Schult. Mant. ii. 272 = Panicum bengalense.
sylvaticus, Roem. & Schult. Syst. ii. 481 = compositus.
tenuis, J. & C. Presl, Rel. Haenk. i. 319 = Ichnanthus candicans.
Thiebauti, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 502.—Mexic.
tomentosus, Schult. Mant. ii. 272 = Panicum tomentosum.

OPLISMENUS:—
undulatifolius, *Beauv. Agrost.* 54.—Reg. trop. et subtrop.
velutinus, Schult. Mant. ii. 271 = Burmannii?
Walteri, Kunth, Rév. Gram. i. 45 = Panicum caroliniánum.
zelayensis, H. B. & K. Nov. Gen. et Sp. i. 108 = Panicum zelayense.

OPLONIA, Rafin. Fl. Tellur. iv. 64 (1836) = **Anthacanthus**, Nees (Acanth.).
spinosa, Rafin. l. c. 65 = A. spinosus.

OPLOPANAX, Miq. Ann. Mus. Bot. Lugd. Bat. i. 16 (1863) = **Fatsia**, Decne. (Araliac.).
horridum, Miq. l. c. = F. horrida.

OPLOTHECA, Nutt. Gen. Am. ii. 78 (1818) = **Froelichia**, Moench (Amarant.).
amarantoides, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 297 = Iresine amarantoides.
decumbens, Mart. l. c. = Celosia trigyna.
floridana, Nutt. Gen. Am. ii. 79 = F. floridana.
gracilis, Hook. Ic. Pl. sub t. 256 = F. gracilis.
interrupta, Nutt. Gen. Am. ii. 79 = F. lanata.
lactea, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 295 = Aerva sanguinolenta.
lanata, Mart. Nov. Gen. et Sp. ii. 48. t. 146 = F. lanata.
Poiretiana, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 297 = F. Humboldtiana.
sericea, Mart. Nov. Gen. et Sp. ii. 49 = F. sericea.
texana, A. Br. in Ann. Sc. Nat. Sér. III. xii. (1849) 355.—Texas.
tomentosa, Mart. Nov. Gen. et Sp. ii. 48 = F. tomentosa.

OPLUKION, Rafin. Sylva Tellur. 53 (1838) = **Lycium**, Linn. (Solan.).
afrum, Rafin. l. c. = L. afrum.
horridum, Rafin. l. c. = L. horridum, *Thunb.*

OPNITHOGALUM, Roem. Arch. i. III. (1798) 42, err. typ. = **Ornithogalum**, Tourn. (Liliac.).

OPOIDIA, Lindl. Bot. Reg. (1839) Misc. 66 = **Peucedanum**, Linn. (Umbell.).
galbanifera, Lindl. l. c.—Persia.

OPOPANAX, Koch, in Nov. Act. Nat. Cur. xii. I. (1824) 96. *UMBELLIFERAE*, Benth. & Hook. f. i. 922.
CRENOSCIADIUM, Boiss. & Heldr. Diagn. Ser. I. x. 30 (1849).
Chironium, Guss. Fl. Sic. Syn. i. 352 = hispidum.
Chironium, *Koch, in Nov. Act. Nat. Cur.* xii. I. (1824) 96.—Reg. Mediterr.
glabrum, Bernh. Sel. Sem. (1837); ex Steud. Nom. ed. II. ii. 220 = O. Chironium.
hispidum, *Griseb. Spicil. Fl. Rumel.* i. 378.—Graec.; As. Min.
orientale, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 330 = hispidum.
persicum, *Boiss. & Heldr. Diagn.* Ser. I. x. 36.—Persia.
siculum, Huet, ex Nym. Consp. 288 = hispidum.

OPORANTHUS, Herb. App. [Bot. Reg.] 38 (1821) = **Sternbergia** Waldst. & Kit. (Amaryll.).
citrinus, Herb. l. c. = S. citrina.
Clusianus, Herb. l. c. = S. colchiciflora.
colchiciflorus, Herb. l. c. = S. colchiciflora.
exiguus, Herb. l. c. = Tapeinanthus humilis.
Fischerianus, Herb. Amaryll. 412. t. 47. f. 2 = S. Fischeriana.
luteus, Herb. App. [Bot. Reg.] 38 = S. lutea.
siculus, Parl. Fl. Ital. iii. 97 = S. lutea.

OPORINEA, Baxt. in Loud. Hort. Brit. Suppl. iii. 599 (1850) = seq.

OPORINIA, D. Don, in Edinb. N. Phil. Journ. (Jan.-March, 1829) 309 = **Leontodon**, Linn. (Compos.).
autumnalis, Coss. ex Willk. & Lange, Prod. Fl. Hisp. ii. 215 = L. carpetana.
autumnalis, D. Don, in Edinb. N. Phil. Journ. (Jan.-March, 1829) 309 = L. autumnalis.
hispidula, DC. Prod. vii. 109 = L. hispidulus.

OPORINIA:—
keretina, Walp. Rep. vi. 345 = Leontodon pyrenaicus.
laciniata, Bertol. Misc. Bot. ii. 21.—Mesopot.
microcephala, Boiss. ex DC. Prod. vii. 109 = L. microcephalus.
norvegica, Griseb. in Wiegm. Archiv, x. (1844) I. 12 (Quid?).—Norvegia.
pratensis, Less. Syn. Comp. 132 = L. autumnalis.
pyrenaica, Sch. Bip. ex Willk. & Lange, Prod. Fl. Hisp. ii. 216 = L. pyrenaicus.

OPSAGO, Rafin. Sylva Tellur. 54 (1838) = **Withania**, Pauq. (Solan.).
cordata, Rafin. l. c. = W. frutescens.
suberosa, Rafin. l. c. = W. frutescens.

OPSANTHA, Delarb. Fl. Auv. ed. II. 80 (1800) = **Gentiana**, Linn.
Amarella, Delarb. l. c. = G. Amarella.

OPSANTHE, Renealm. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 120 = **Gentiana**, Linn. f.
flava, Fourr. l. c. = G. germanica.
germanica, Fourr. l. c. = G. germanica.
obtusifolia, Fourr. l. c. = G. germanica.

OPSIANTHES, Lilja, Fl. Sverig. Suppl. 25 (1840) = **Clarkia**, Pursh. (Onagr.).
gauroides, Lilja, l. c. = C. rhomboidea.

OPSIESTON, Bunge in Mém. Sav. Etr. Pétersb. vii. (1851) 474 (Rel. Lehm. 298) = **Ofaiston**, Rafin. (Chenop.).

OPSOPAEA, Neck. Elem. ii. 305 (1790) = **Helicteres**, Linn. (Stercul.).

OPULASTER, Medic. Ph. Anat. 109 (1799) = **Neillia**, D. Don (Rosac.).
bullatus, Medic. l. c. = N. opulifolia.

OPULUS, Tourn. ex Linn. Gen. ed. I. 86 (1737) = **Viburnum**, Linn. (Caprifol.).
edulis, J. S. Presl, Rostl. ii. 98 = V. Opulus.
glandulosa, Moench, Meth. 505 = V. Opulus.
lobatofolia, Gilib. Exercit. i. 4 = V. Opulus.
mollis, J. S. Presl, Rostl. ii. 97 = V. molle.
oxycoccos, J. S. Presl, l. c. = V. Opulus.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 489 = V. Opulus.
rosea, Schur, Enum. Pl. Transs. 941 (Index) = V. Opulus.
trilobifolia, Gilib. Fl. Lituan. i. 2; ex Ledeb. Fl. Ross. ii. 384 = V. Opulus.
vulgaris, Borckh. in Roem. Archiv, i. II. 20 = V. Opulus.

OPUNTIA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752). *CACTEAE*, Benth. & Hook. f. i. 851.
CONSOLEA, Lem. in Rev. Hortic. (1862) 174.
FICINDICA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70.
TEPHROCACTUS, Lem. Cact. 88 (1868).
acanthocarpa, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 51. t. 18. f. 1–3.—Am. bor. occ.
acracantha, Walp. Rep. ii. 354 = aoracantha.
albicans, *Salm-Dyck, Hort. Dyck.* 361.—Mexic.
Alfagayucca, Karw. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 68 = O. Amyclaea.
alpina, Gill. ex Pfeiff. Enum. Diagn. Cact. 146 = platyacantha.
Amyclaea, *Tenore, Fl. Neap. Prod. App.* v. 15.—Mexic.
andicola, *Hort. Angl. ex Pfeiff. Enum. Diagn. Cact.* 145.—Chili.
andicola, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 71 = glomerata.
angusta, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 30.—Mexic.
angustata, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 39. t. 7.—Calif.
aoracantha, *Lem. Cact. Aliq. Nov. Desc.* 34.—Reg. Argent.
arborea, *Steud. Nom.* ed. II. ii. 220.—Bras.
arborescens, *Engelm. in Wislis. Tour N. Mexic.* 90.—N. Mexic.
Arbuscula, *Engelm. in Proc. Am. Acad.* iii. (1857) 309.—Mexic.
arenaria, *Engelm. l. c.* 301.—Texas.

OPUNTIA:—
argentina, *Griseb. in Goett. Abh.* xxiv. (1879) 140.—Reg. Argent.
articulata, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 103 = Cereus articulatus.
atacamensis, *Phil. Fl. Atac.* 24.—Chili.
Auberi, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 282.—Cuba.
aurantiaca, *Lindl. Bot. Reg.* t. 1606.—Chili.
Bartrami, Rafin. Atl. Journ. 146 (Quid ?).—Am. bor.
basilaris, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 43.—Am. bor. occ.
bicolor, *Phil. in Linnaea*, xxxiii. (1864–65) 83.—Chili.
Bigelovii, *Engelm. in Pacif. Rail. Rep.* iv. 50.—Calif.
boliviana, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 387.—Bolivia.
Bonplandii, Hort. ex Pfeiff. Enum. Diagn. Cact. 161 = O. Tuna.
brachyarthra, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 47.—Am. bor. occ.
brasiliensis, *Haw. Suppl. Pl. Succ.* 79.—Bras.
bulbispina, Engelm. in Proc. Am. Acad. iii. (1857) 304 = tunicata.
caespitosa, Auct. = *cespitosa*.
calacantha, *Forbes, ex Baxt. in Loud. Hort. Brit. Suppl.* iii. 599, nomen.—Hab. ?
californica, Engelm. in Emory, Notes Mil. Reconnoiss. 158 = acanthocarpa.
calva, Lem. Cact. Gen. Sp. Nov. 73 = Turpinii.
camanchica, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 40.—Am. bor. occ.
candelabriformis, *Hort. Monac. ex Pfeiff. Enum. Diagn. Cact.* 159.—Mexic.
caracassana, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 238.—Venezuela.
carolina, *Forbes, Hort. Tour Germ.* 160, nomen.—Am. bor.
catocantha, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 166.—Ins. S. Thomas.
cespitosa, Rafin. in Ser. Bull. Bot. (1830) 216 = Rafinesquii.
chlorotica, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 38.—Calif.
clavarioides, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 173.—Mexic.
clavata, *Engelm. in Wisliz. Tour. N. Mexic.* 95.—Mexic.
clavata, *Phil. in Anal. Univ. Chil.* (1872) 722.—Chili.
coccinea, Hort. ex Pfeiff. Enum. Diagn. Cact. 161 = O. Tuna.
coccinellifera, Steud. Nom. ed. II. ii. 220 = Nopalea coccinellifera.
cochenillifera, DC. Pl. Grass. t. 137 = nigricans.
cochinelifera, Mill. Gard. Dict. ed. VIII. n. 6 = Nopalea coccinellifera.
coerulea, Gill. ex Pfeiff. Enum. Diagn. Cact. 155 = sericea.
corrugata, *Hort. Angl. ex Salm-Dyck, Hort. Dyck.* 360; *et ex D. Dietr. Syn. Pl.* iii. 111.—Am. austr.
corrugata, Hort. Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 43 = Parmentieri.
costigera, *Miq. in Nederl. Kruidk. Arch.* iv. (1859) 338.—Mexic.
crassa, *Haw. Suppl. Pl. Succ.* 81.—Mexic.
crinifera, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 157.—Bras.
cristata, Salm-Dyck, Cact. Hort. Dyck. ed. I. 49; et ed. II. 73 = imbricata.
cruciata, Hort. ex Pfeiff. Enum. Diagn. Cact. 167 = spinosissima.
curassavica, *Mill. Gard. Dict.* ed. VIII. n. 7.—Ind. occ.
cylindrica, *DC. Prod.* iii. 471.—Peruv.
cymochila, Engelm. & Bigel. in Pacif. Rail. Rep. iv. 42 = Rafinesquii.
Darwinii, *Hensl. in Mag. Zool. & Bot.* i. (1837) 466.—Patagonia.
Davisii, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 49.—Am. bor.
decipiens, DC. in Mém. Mus. Par. xvii. (1828) 118 = imbricata.
decumana, *Haw. Rev. Pl. Succ.* 71.—Am. austr.
decumbens, *Salm-Dyck, Hort. Dyck.* 361.—Mexic.
deflexa, Lem. Cact. Gen. Sp. Nov. 68 = gracilior.
dejecta, *Salm-Dyck, Hort. Dyck.* 361.—Cuba.

OPUNTIA:—
Demorenia, Forbes, Hort. Tour Germ. 160, nomen = seq.
Demoriana, *Hort. Monac. ex Foerst. Handb. Cact.* ed. I. 508, nomen.—Hab. ?
Deppei, *Wendl. ex Foerst. l. c.* 508, nomen.—Mexic. ?
diademata, *Lem. Cact. Aliq. Nov. Desc.* 36.—Reg. Argent.
dichotoma, *Hort. ex Forbes, Hort. Tour Germ.* 157.—Hab. ?
diffusa, Hort. ex Pfeiff. Enum. Diagn. Cact. 159 = dejecta.
Dillenii, *Haw. Suppl. Pl. Succ.* 79.—Am. austr.
dimorpha, *Foerst. in Hamb. Gartenz.* xvii. (1861) 167.—Peruv.
dolabriformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 167 = ferox.
Drummondii, *R. Grah. in Maund, Botanist*, v. t. 246; *et ex Pritz. Ic. Ind.* i. 767 (Quid ?).—Am. bor.
dulcis, Engelm. in Cact. Mex. Bound. 48 = Engelmanni.
eborina, Foerst. ex Haage, Verz. Cact. 29 = seq. ?
eburnea, Lem. Cact. Aliq. Nov. Desc. 35 = corrugata.
echinocarpa, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 49.—Am. bor. occ.
elata, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 434, nomen; *Salm-Dyck, Hort. Dyck.* 361.—Bras.
elatior, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. austr.
Ellemetiana, *Miq. in Nederl. Kruidk. Arch.* iv. (1859) 337.—Chili.
elongata, Haw. Suppl. Pl. Succ. 81 = decumana, fulvispina.
Emoryi, *Engelm. in Proc. Am. Acad.* iii. (1857) 303.—Am. bor. occ.
Engelmanni, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 235.—Am. bor. occ.
erinacea, Engelm. & Bigel. in Pacif. Rail. Rep. iv. 47 = rutila.
erythrocentron, Lem. ex Foerst. Handb. Cact. ed. I. 492 = fulvispina.
extensa, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 147.—Hab. ?
exuviata, DC. in Mém. Mus. Par. xvii. (1828) 118 = tunicata.
exuviata, Salm-Dyck, Cact. Hort. Dyck. ed. I. 48; ed. II. 73 = imbricata.
exuviato-stellata, Lem. ex Labour. Cact. 492 = arborescens.
ferox, *Haw. Suppl. Pl. Succ.* 82.—Am. austr.
festiva, Sencke, ex Haage, Verz. Cact. 29 (Quid ?).—Hab. ?
Ficus-indica, Mill. Gard. Dict. ed. VIII. n. 2 = O. Tuna ?
filipendula, *Engelm. in Proc. Am. Acad.* iii. (1857) 294.—Texas.
flavicans, Lem. Cact. Gen. Sp. Nov. 61 = robusta.
flavispina, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 508, nomen.—Hab. ?
flexibilis, Hort. ex Pfeiff. Enum. Diagn. Cact. 161 = O. Tuna.
floccosa, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 388.—Peruv.
floribunda, Lem. Cact. Gen. Sp. Nov. 68 = Salmiana.
foliosa, *Salm-Dyck, ex DC. Prod.* iii. 471.—Am. austr.
fragilis, *Haw. Suppl. Pl. Succ.* 82.—Am. bor. occ.
fragilis, Hort. ex Foerst. Handb. Cact. ed. I. 489 = missouriensis.
frutescens, Engelm. in Bost. Journ. Nat. Hist. v. (1845) 245 = leptocaulis.
fulgens, Engelm. in Bot. Calif. i. 250 = seq.
fulgida, *Engelm. in Proc. Am. Acad.* iii. (1857) 306.—Mexic.
fulvispina, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 164.—Am. austr.
furiosa, Wendl. Cat. Hort. Herrenh. (1835) = tunicata.
fusco-atra, *Engelm. in Proc. Am. Acad.* iii. (1857) 297.—Texas.
fusiformis, Engelm. & Bigel. in Pacif. Rail. Rep. iv. 43 = Rafinesquii.
galapageia, *Hensl. in Mag. Zool. & Bot.* i. (1837) 467.—Ins. Galapag.
Galeottii, *De Smet, ex Miq. in Nederl. Kruidk. Arch.* iv. (1859) 337.—Mexic.
Gilliesii, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 102 = Cereus ovatus.
glaberrima, Hort. Berol. ex Pfeiff. l. c. 153 = crassa.
glauca, *Hort. ex Pfeiff. l. c.* 153 = parvula.

OPUNTIA:—
glaucescens, Hort. ex Pfeiff. Enum. Diagn. Cact. 155 = grandis.
glaucescens, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 434; *Salm-Dyck, Hort. Dyck.* 362.—Mexic.
glaucophylla, *Wendl. Cat. Hort. Herrnh.* (1835); *ex Pfeiff. Diagn. Cact.* 162.—Mexic. ?
glomerata, *Haw. in Phil. Mag.* vii. (1830) 111.—Reg. Argent.
gracilior, *Lem. Cact. Gen. et Sp. Nov.* 68.—Hab. ?
gracilis, *Hort. Monac. ex Pfeiff. Enum. Diagn. Cact.* 172.—Mexic.
Grahami, *Engelm. in Proc. Am. Acad.* iii. (1857) 304.—Mexic.
grandiflora, *Engelm. l. c.* 295.—Texas.
grandis, *Hort. Angl. ex Pfeiff. Enum. Diagn. Cact.* 155.—Mexic.
grata, *Phil. in Linnaea*, xxx. (1859–60) 211.—Chili.
Hernandezii, *DC. in Mém. Mus. Par.* xvii. (1828) 69. t. 16.—Mexic.
Hieronymi, *Griseb. in Goett. Abh.* xxiv. (1879) 140.—Reg. Argent.
Hitchenii, *Forbes, Hort. Tour Germ.* 159, nomen.—Hab. ?
horizontalis, Gill. ex Pfeiff. Enum. Diagn. Cact. 145 = andicola.
horizontalis, Hort. ex Pfeiff. l. c. 159 = dejecta.
horrida, Salm-Dyck, ex DC. Prod. iii. 472 = O. Tuna.
humifusa, Rafin. Fl. Med. ii. 247 = vulgaris.
humilis, Haw. Syn. Pl. Succ. 189 = O. Tuna.
hystricina, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 44.—Mexic.
Hystrix, *Griseb. Cat. Pl. Cub.* 117.—Cuba.
Hystrix, Hort. ex Pfeiff. Enum. Diagn. Cact. 148 = foliosa.
imbricata, *DC. Prod.* iii. 471.—Mexic.
inermis, DC. l. c. 473 = stricta.
inermis, Moris & De Not. Fl. Caprar. 58 = vulgaris.
intermedia, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 206 = Rafinesquii.
intermedia, Salm-Dyck, Hort. Dyck. 364 = vulgaris.
involuta, Otto, ex Foerst. Handb. Cact. ed. I. 505 = floccosa.
irrorata, Mart. ex Pfeiff. Enum. Diagn. Cact. 154 = decumbens.
italica, Tenore, ex Pfeiff. l. c, 149 = vulgaris.
Jussieuii, *Haage, ex Foerst. Handb. Cact.* ed. I. 508, nomen.—Hab. ?
Karwinskiana, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 239.—Mexic.
Kleiniae, *DC. in Mém. Mus. Par.* xvii. (1828) 118.—Mexic.
lanceolata, *Haw. Syn. Pl. Succ.* 192.—Am. austr.
lanigera, Parm. ex Pfeiff. Enum. Diagn. Cact. 157 = fulvispina.
lanigera, Salm-Dyck, Cact. Hort. Dyck, ed. II. 65, 234 = crinifera.
lasiacantha, Hort. Vindob. ex Pfeiff. Enum. Diagn. Cact. 160 = megacantha.
leptocaulis, *DC. in Mém. Mus. Par.* xvii. (1828) 118.—Mexic.
leucacantha, Salm-Dyck, Hort. Dyck. 362 = megacantha.
leucantha, *Link & Otto, in Verh. Ver. Befoerd. Gartenb.* vi. (1830) 434, nomen; *Pfeiff. Enum. Diagn. Cact.* 167.—Mexic.
leucostata, Forbes, Hort. Tour Germ. 160, nomen = seq. ?
leucosticta, Wendl. Cat. Hort. Herrnh. (1835); ex Pfeiff. Enum. Diagn. Cact. 167 = megacantha.
leucotricha, *DC. in Mém. Mus. Par.* xvii. (1828) 119.—Mexic.
Lindheimeri, Engelm. in Bost. Journ. Nat. Hist. vi. (1850) 207 = Engelmanni.
longispina, *Haw. in Phil. Mag.* vii. (1830) 111.—Chili.
longispina, Hort. ex Pfeiff. Enum. Diagn. Cact. 156 = orbiculata.
macracantha, *Griseb. Cat. Pl. Cub.* 116.—Cuba.
macrocentra, *Engelm. in Proc. Am. Acad.* iii. (1857) 292.—Mexic.
macrorhiza, Engelm. in Bost. Journ. Nat. Hist. vi. (1850) 206 = Rafinesquii.
maculacantha, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 166.—Reg. Argent.
Maelenii, Hort. Belg. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 46 = sericea.
magnifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 22 = Nopalea coccinellifera.

OPUNTIA:—
mamillata, *Schott, ex Engelm. in Proc. Am. Acad.* iii. (1857) 308.—Mexic.
maritima, Rafin. Fl. Med. ii. 247 = vulgaris.
maxima, Mill. Gard. Dict. ed. VIII. n. 5 = decumana.
maxima, Salm-Dyck, ex DC. Prod. iii. 474 = O. Amyclaea.
media, Haw. Suppl. Pl. Succ. 82 = missourensis.
megacantha, *Salm-Dyck, Hort. Dyck.* 363.—Mexic.
mesacantha, Rafin. in Ser. Bull. Bot. (1830) 216 = Rafinesquii.
Metternichii, Picol. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 46 = orbiculata.
mexicana, Hort. ex Pfeiff. Enum. Diagn. Cact. 150, 160 = megacantha, Nopalea coccinellifera.
microcarpa, *Engelm. in Emory, Notes Mil. Reconnois.* 157.—Am. bor. occ.
microdasys, *Pfeiff. Enum. Diagn. Cact.* 154.—Mexic.
microthele, Hort. ex Pfeiff. l. c. 173 = clavarioides.
minor, C. Muell. in Walp. Ann. v. 50, sphalm. = tenuispina.
Miquelii, Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 49 = pulverulenta.
missouriensis, *DC. Prod.* iii. 472.—Am. bor.
mohavensis, Engelm. & Bigel. in Pacif. Rail. Rep. iv. t. 9. f. 6–8 = seq.
mojavensis, *Engelm. & Bigel. l. c.* iv. 40.—Colorado.
monacantha, *Haw. Suppl. Pl. Succ.* 81.—Am. austr.
moniliformis, Haw. ex Steud. Nom. ed. II. 221 = Cereus moniliformis.
montana, Sencke, in Haage, Verz. Cact. 29 (Quid ?).—Hab. ?
monticola, *Phil. in Linnaea*, xxxiii. (1864–65) 82.—Chili.
myriacantha, *Link & Otto, ex Steud. Nom.* ed. II. ii. 221, nomen.—Mexic.
nana, DC. Pl. Grass. sub t. 138 = vulgaris.
nigricans, *Haw. Syn. Pl. Succ.* 189.—Am. austr.
nigricans, Hort. Par. ex Foerst. Handb. Cact. ed. I. 496 = elatior.
Nopalilla, Karw. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 68 = Karwinskiana.
oblongata, *Wendl. Cat. Hort. Herrnh.* (1835); *ex Pfeiff. Enum. Diagn. Cact.* 161.—Mexic.
occidentalis, Engelm. & Bigel. in Pacif. Rep. iv. 38 = Engelmanni.
oligacantha, Hort. Vindob. ex Pfeiff. Enum. Diagn. Cact. 137 = spinulifera.
orbiculata, *Salm-Dyck, ex Pfeiff. l. c.* 156.—Chili.
Ottonis, G. Don, Gen. Syst. iii. 172 = Echinocactus Ottonis.
Ottonis, Salm-Dyck, Cact. Hort. Dyck. ed. I. 48, nomen. = Schomburgkii.
Ovallei, *C. Gay, Fl. Chil.* iii. 29.—Chili.
ovata, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 102 = Cereus ovatus.
ovata, *Pfeiff. l. c.* 144.—Chili.
ovoides, Lem. Cact. Gen. Sp. Nov. 73 = ovata.
papyracantha, *Phil. in Regel, Gartenfl.* (1872) 129.—Chili.
Parmentieri, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 276.—Paraguay.
Parryi, *Engelm. in Am. Journ. Sc.* Ser. II. xiv. (1852) 339.—Am. bor. occ.
parvispina, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 238.—Mexic ?
parvula, Salm-Dyck, l. c. 69 = crassa.
parvula, *Salm-Dyck, Hort. Dyck.* 364.—Chili.
patagonica, *Phil. in Linnaea*, xxxiii. (1864–65) 82.—Patagon.
pelaguensis, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 71 = platyacantha.
Pentlandii, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 387.—Bolivia.
Pes-corvi, *Leconte, ex Chapm. Fl. S. U. St.* 145.—Am. bor.
phaeacantha, *Engelm. in Mem. Am. Acad.* N. S. iv. (1849) 51.—N. Mexic.
phyllacantha, *Salm-Dyck, ex Foerst. Handb. Cact.* ed. I. 508, nomen.—Hab. ?
Phyllanthus, Mill. Gard. Dict. ed. VIII. n. 9 = Phyllocactus Phyllanthus.
Pintadera, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 47 = crinifera.
platyacantha, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 103 = diademata.
platyacantha, *Hort. Dyck. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 371.—Chili.

OPUNTIA:—
Poeppigii, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 174.—Chili.
polyacantha, Haw. Suppl. Pl. Succ. 82 = missouriensis.
polyantha, *Haw. Syn. Pl. Succ.* 190.—Am. austr.
polymorpha, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 103 = Cereus articulatus.
Pottsii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 236.—Mexic.
praecox, *Forbes, Hort. Tour Germ.* 160, nomen.—Hab. ?
Prate, Sabine, ex Pfeiff. Enum. Diagn. Cact. 155 = albicans.
procumbens, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 39. t. 7. f. 4–5.—N. Mexic.
prolifera, *Engelm. in Am. Journ. Sc.* Ser. II. xiv. (1852) 338.—Am. bor. occ.
prostrata, Monv. & Lem. ex Foerst. Handb. Cact. ed. I. 478 = intermedia.
protracta, *Lem. l. c.* 508, nomen.—Hab. ?
pruinosa, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 67 = albicans.
Pseudo-tuna, *Salm Dyck, Obs. Bot.* (1822) 7.—Am. austr.
puberula, *Hort. Vindob. ex Pfeiff. Enum. Diagn. Cact.* 156.—Mexic.
pubescens, *Wendl. Cat. Hort. Herrnh.* (1835); *ex Pfeiff. l. c.* 149.—Mexic.
pulchella, *Engelm. in Trans. Acad. St. Louis*, ii. (1863) 201.—Am. bor. occ.
pulverulenta, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 407.—Am. austr.
pulvinata, DC. in Mém. Mus. Par. xvii. (1828) 119 = microdasys.
pusilla, Haw. Syn. Pl. Succ. 195 = foliosa.
pusilla, *Salm-Dyck, Obs. Bot.* (1822) 10.—Am. austr.
Rafinesquii, *Engelm. in Pacif. Rail. Rep.* iv. 41. t. 11.—Am. bor. occ.
ramosissima, Engelm. in Am. Journ. Sc. Ser. II. xiv. (1852) 339 = tesselata.
ramulifera, *Salm-Dyck, Hort. Dyck.* 360.—Mexic.
repens, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 277.—Porto Rico.
repens, Karw. ex Salm-Dyck, Hort. Dyck. 361 = decumbens.
reptans, *Karw. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 599, nomen.—Mexic.
retrospinosa, Lem. Cact. Aliq. Nov. Desc. 35 = Parmentieri.
robusta, *Wendl. Cat. Hort. Herrnh.* (1835); *ex Pfeiff. Enum. Diagn. Cact.* 165.—Mexic.
rosea, *DC. Prod.* iii. 471.—Mexic.
Roxburghiana, Voigt, Hort. Suburb. Calc. 62 = Dillenii.
rubescens, *Salm-Dyck, ex DC. Prod.* iii. 474.—Bras.
rufescens, Salm-Dyck, ex Foerst. Handb. Cact. ed. I. 493 = fulvispina.
rufida, *Engelm. in Proc. Am. Acad.* iii. (1857) 298.—Mexic.
rutila, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 555.—Am. bor. occ.
Sabinii, Hort. ex Pfeiff. Enum. Diagn. Cact. 147 = fragilis.
salicornoides, Spreng. ex Pfeiff. l. c. 141 = Rhipsalis salicornoides.
Salmiana, *Parm. ex Pfeiff. l. c.* 172.—Bras.
Salmii, *Hort. ex Forbes, Hort. Tour Germ.* 158, nomen.—Hab. ?
Schomburgkii, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. I. 48, nomen.—Guian. ?
Schottii, *Engelm. in Proc. Am. Acad.* iii. (1857) 304.—Texas.
Segethi, *Phil. in Anal. Univ. Chil.* lv. (1879) 263; *ex Phil. in Regel, Gartenfl.* (1883) 260.—Hab. ?
senilis, Parm. ex Pfeiff. Enum. Diagn. Cact. 157 = crinifera.
senilis, *Roezl, in Belg. Hortic.* xxiv. (1874) 39.—Peruv.
sericea, *G. Don, ex Salm-Dyck, Hort. Dyck.* 363.—Chili.
serpentina, *Engelm. in Am. Journ. Sc.* Ser. II. xiv. (1852) 338.—Am. bor. occ.
setispina, *Engelm. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 239.—Mexic.
speciosa, Steud. Nom. ed. II. ii. 222 = Phyllocactus phyllanthoides.
sphaerica, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 167.—Peruv.

OPUNTIA:—
sphaerocarpa, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 47.—N. Mexic.
spinalba, Rafin. Atl. Journ. 147 (Quid ?).—Am. bor.
spinaurea, Karw. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 46 = O. Pseudo-tuna.
spiniflora, *Phil. in Linnaea*, xxx. (1859–60) 211.—Chili.
spinosissima, *Mill. Gard. Dict.* ed. VIII. n. 8.—Ind. occ.
spinulifera, *Salm-Dyck, Hort. Dyck.* 364.—Mexic.
spinulosa, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 46, nomen.—Hab. ?
spirocentra, *Engelm. & Bigel. ex Haage, Verz. Cact.* 30.—N. Mexic.
splendens, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 159 = missouriensis ?
Stanlyi, *Engelm. in Emory, Notes Mil. Reconnois.* 157.—N. Mexic.
Stapeliae, *DC. in Mém. Mus. Par.* xvii. (1828) 117.—Mexic.
stellata, Salm-Dyck, Cact. Hort. Dyck. ed. I. 48; ed. II. 73 = arborescens.
stenochila, Engelm. & Bigel. in Pacif. Rail. Rep. iv. 43 = Rafinesquii.
stenopetala, *Engelm. in Proc. Am. Acad.* iii. (1857) 289.—Mexic.
straminea, *Sencke, ex Haage, Verz. Cact.* 30.—Hab. ?
streptacantha, *Lem. Cact. Gen. Sp. Nov.* 62.—Mexic.
stricta, *Haw. Syn. Pl. Succ.* 191.—Am. austr.
Strigil, *Engelm. in Proc. Am. Acad.* iii. (1857) 290.—Texas.
subferox, Schott, ex Pfeiff. Enum. Diagn. Cact. 167 = leucacantha.
subinermis, *Link, ex Steud. Nom.* ed. II. ii. 222, nomen.—Am. austr.
subulata, *Engelm. in Gard. Chron.* (1883) i. 627.—Am. austr.
sulphurea, *G. Don, ex Loud. Hort. Brit.* 196; *Salm-Dyck, Hort. Dyck.* 360.—Chili.
syringacantha, Pfeiff. ex Foerst. Handb. Cact. ed. I. 470 = diademata.
tenuispina, *Engelm. in Proc. Am. Acad.* iii. (1857) 294.—Mexic.
tesselata, *Engelm. in Pacif. Rail. Rep.* iv. 52. t. 21.—Calif.
Thurberi, *Engelm. in Proc. Am. Acad.* iii. (1857) 308.—Mexic.
Tidballii, Bigel. in. Pacif. Rail. Rep. iv. 11 = chlorotica.
tomentosa, *Salm-Dyck, Obs. Bot.* (1822) 8.—Am. austr.
tortispina, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 41.—Am. bor. occ.
triacantha, Sweet, Hort. Brit. ed. I. 172 = curassavica.
tuberculata, *Haw. Suppl. Pl. Succ.* 80.—Am. austr.
tuberosa, Hort. Angl. ex Pfeiff. Enum. Diagn. Cact. 146 = platyacantha.
Tuna, DC. Pl. Grass. sub t. 138 = monacantha.
Tuna, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. austr.
tunicata, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 170.—Mexic.
Turpinii, *Lem. Cact. Aliq. Nov. Desc.* 36.—Reg. Argent.
Umbrella, Steud. Nom. ed. II. ii. 222 = curassavica ?
undulata, Hort. ex Pfeiff. Enum. Diagn. Cact. 170 = tunicata.
vaginata, Engelm. in Wisliz. Tour N. Mexico, 100 = leptocaulis.
vestita, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 388.—Peruv.
violacea, *Engelm. in Emory, Notes Mil. Reconnois.* 157.—Arizona.
virgata, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 173 = leptocaulis.
virgata, Hort. Vindob. ex Foerst. Handb. Cact. ed. I. 506 = gracilis.
vulgaris, *Mill. Gard. Dict.* ed. VIII. n. 1.—Am. bor.
vulgaris, Tenore, Syll. Fl. Neap. 239 = O. Tuna ?
Whipplei, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 50.—Calif.
Wrightii, Engelm. in Proc. Am. Acad. iii. (1857) 308 = Kleiniae.

ORABANCHE, Losc & Pardo, Ser. Inconfect. Pl. Arag. 79 (1863), sphalm. = **Orobanche**, Tourn.

ORANIA, Zipp. in Alg. Konst- en Letter-bode (1829), i. 297. *PALMAE*, Benth. & Hook. f. iii. 918.
ARAUSIACA, Blume, Rumphia, ii. praef. p. 8 (1836), tt. 119, 122.
MACROCLADUS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 489.

ORANIA:—
aruensis, *Becc. Malesia*, i. (1877) 76.—N. Guin.
macrocladus, *Mart. Hist. Nat. Palm.* iii. 186. t. 177.—Malacca; Java.
moluccana, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 163.—Ins. Molucc.
nicobarica, Kurz, in Journ. Bot. xiii. (1875) 331 = Bentinckia nicobarica.
philippinensis, *Scheff. ex Becc. in Ann. Jard. Buitenz.* ii. (1885) 156.—Ins. Philipp.
porphyrocarpa, Blume, ex Mart. Hist. Nat. Palm. iii. 190. t. 157 = Didymosperma porphyrocarpa.
regalis, *Blume, ex Zipp. in Alg. Konst- en Letterbode* (1829), i. 297.—Java.
regalis, Miq. Palm. Archip. Ind. 4 = moluccana.

ORAOMA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 411 = **Amoora**, Roxb. (Meliac.).
canarana, Turcz. l. c. = A. canarana.

ORBEA, Haw. Syn. Pl. Succ. 37 (1812) = **Stapelia**, Linn. (Asclep.).
anguinea, Haw. l. c. 41 = S. picta.
bisulca, Haw. l. c. 39 = S. variegata.
bufonia, Haw. l. c. 40 = S. variegata.
clypeata, Haw. Suppl. Pl. Succ. 13 = S. clypeata.
Curtisii, Haw. Syn. Pl. Succ. 40 = S. variegata.
conspurcata, Schult. Syst. vi. 834 = S. conspurcata.
decora, Steud. Nom. ed. II. ii. 222 = Piaranthus decorus.
inodora, Haw. Suppl. Pl. Succ. 12 = S. inodora.
lepida, Haw. l. c. 13 = S. lepida.
maculosa, Haw. Syn. Pl. Succ. 37 = S. maculosa.
marginata, Schult. Syst. vi. 834; et G. Don, Gen. Syst. iv. 120 = S. marginata.
marmorata, Schult. Syst. vi. 834 = S. marmorata.
mixta, Haw. Syn. Pl. Succ. 38 = S. mixta.
mutabilis, G. Don, in Loud. Hort. Brit. 96; Gen. Syst. iv. 121 = S. mutabilis.
normalis, Schult. Syst. vi. 834; G. Don, l. c. 120 = S. normalis.
orbicularis, Haw. Syn. Pl. Succ. 40 = S. orbicularis.
picta, Haw. l. c. 42 = S. picta.
planiflora, G. Don, Gen. Syst. iv. 120 = S. planiflora.
quinquenervia, Haw. Syn. Pl. Succ. 38 = S. planiflora.
retusa, Haw. l. c. 41 = S. retusa.
rugosa, G. Don, in Loud. Hort. Brit. 96 = S. rugosa.
trisulca, G. Don, ex Steud. Nom. ed. II. ii. 222 = S. trisulca.
variegata, Haw. Syn. Pl. Succ. 40 = S. variegata.
Wendlandiana, Schult. Syst. vi. 834; G. Don, Gen. Syst. iv. 120 = S. Wendlandiana.
Woodfordiana, Haw. Syn. Pl. Succ. 42 = S. Woodfordiana.

ORBECKIA, G. Don, Gen. Syst. ii. 730 (1832), err. typ. = **Osbeckia**, Linn. (Melast.).

ORBEXILUM, Rafin. Atl. Journ. 145 (1832) = **Psoralea**, Linn. (Legumin.).
latifolia, Rafin. l. c. = P. Onobrychis.

ORBICULARIA, Baill. Etud. Gén. Euph. 616 (1858) = **Phyllanthus**, Linn. (Euphorb.).
phyllanthoides, Baill. l. c. 617 = P. orbicularis.

ORBIGNIA, Bert. Herb. ex Steud. Nom. ed. II. ii. 222 (1841) = seq.

ORBIGNYA, Bert. in Merc. Chil. (1829) 737; ex Am. Journ. Sc. Sér. I. xxiii. (1833) 259 = **Llagunoa**, Ruiz & Pav. (Sapind.).
trifolia, Bert. ex Steud. Nom. ed. II. ii. 222 = L. glandulosa.
trifoliata, Bert. in Merc. Chil. (1829) 737; ex Am. Journ. Sc. Ser. I. xxiii. (1833) 259 = L. glandulosa.

ORBIGNYA, Mart. ex Endl. Gen. 257 (1837). *PALMAE*, Benth. & Hook. f. iii. 948.
Cuci, Kunth, ex H. Wendl. in Kerch. Palm. 252 = phalerata.
dubia, *Mart. Hist. Nat. Palm.* iii. 304. t. 169.—Bras.
Eichleri, *Drude, in Mart. Fl. Bras.* iii. II. 449.—Bras.
humilis, *Mart. Palmet. Orbign.* 129. t. 10. f. 2.—Bolivia.
Lydiae, *Drude, in Mart. Fl. Bras.* iii. II. 448.—Bras.

ORBIGNYA:—
phalerata, *Mart. Palmet. Orbign.* 126. t. 13. f. 2.—Bolivia.
Pixuna, Barb.-Rodr. Prot. App. (1879) 49 = Attalea spectabilis.
racemosa, *Drude, in Mart. Fl. Bras.* iii. II. 448.—Bras.

ORBINDA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 (Quid?).

ORCHIASTRUM, Lem. in Illustr. Hortic. ii. (1855) Misc. 96, 100 = **Lachenalia**, Jacq. (Liliac.).
Aitoni, Lem. l. c. 100 = L. orchioides.
glaucinum, Lem. l. c. = L. glaucina.
mutabile, Lem. l. c. = L. orchioides.
pallidum, Lem. l. c. = L. pallida.
pulchellum, Lem. l. c. = L. orchioides.
virentiflavum, Lem. l. c. = L. orchioides.

ORCHIDION, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 218 = **Arethusa**, Gronov. (Orchid.).

ORCHIDIUM, Sw. Summa Veg. Scand. 32 (1814); et in Svensk Bot. t. 518 (1819) = **Calypso**, Salisb. (Orchid.).
americanum, Steud. Nom. ed. II. ii. 222 = C. borealis.
arcticum, Sw. Summa Veg. Scand. 32 = C. borealis.
boreale, Sw. in Svensk. Bot. t. 518 = C. borealis.

ORCHIDOCARPUM, Michx. Fl. Bor. Am. i. 329 (1803) = **Asimina**, Adans. (Anonac.).
arietinum, Michx. l. c. = A. triloba.
grandiflorum, Michx. l. c. 330 = A. grandiflora.
parviflorum, Michx. l. c. 329 = A. parviflora.
pygmaeum, Michx. l. c. 330 = A. pygmaea.

ORCHIDOFUNCKIA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 24 = **Cryptarrhena**, R. Br. (Orchid.).
pallidiflora, A. Rich. & Gal. l. c. = C. pallidiflora.

ORCHIOPS, Salisb. Gen. Pl. Fragm. 21 (1866) = **Lachenalia**, Jacq. (Liliac.).
orchioides, Salisb. l. c. = L. orchioides.

ORCHIPEDA, Blume, Bijdr. 1026 (1826) = **Voacanga**, Thou. (Apocyn.).
foetida, Blume, Bijdr. 1027 = V. foetida.
gracilipes. Miq. Ann. Mus. Bot. Lugd. Bat. i. 316.—Ins. Molucc.
grandifolia, Miq. l. c. = V. grandifolia.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 553.—Sumatra.

ORCHIPEDIUM, Benth. in Journ. Linn. Soc. xviii. (1881) 344, err. typ. = seq.

ORCHIPEDUM, Breda, Kuhl & Hasselt, Gen. et Sp. Orch. [t. 10] (1827) = **Physurus**, Rich. (Orchid.).
plantaginifolium, Breda, l. c.—Java.

ORCHIS, [Tourn.] Linn. Syst. ed. I (1735). *ORCHIDEAE*, Benth. & Hook. f. iii. 620.
ABROCHIS, Neck. Elem. iii. 130 (1790).
ANACAMPTIS, Rich. in Mém. Mus. Par. iv. (1818) 41, 47.
BARLIA, Parl. Nuov. Gen. Piant. Monocot. 5 (1858); Fl. Ital. iii. 445 (1858).
COMPERIA, C. Koch, in Linnaea, xxii. (1849) 287.
DACTYLORHIZA, Neck. Elem. iii. 129 (1790).
HIMANTOGLOSSUM, Spreng. Syst. iii. 675 (1826).
LOROGLOSSUM, Rich. in Mém. Mus. Par. iv. (1818) 41, 47.
MESICERA, Rafin. Neogenyt. 4 (1825).
ORCHITES, Schur, Enum. Pl. Transs. 942 (1866).
STRATEUMA, Salisb. in Trans. Hort. Soc i. (1812) 290.
TRAUNSTEINERA, Reichb. Nom. 50 (1841); Fl. Sax. 87 (1844).
ZOOPHORA, Bernh. Syst. Verz. Erf. 308 (1800).
abortiva, Leyss. Fl. Hal. 226 = Orobanche purpurea.
abortiva, Linn. Sp. Pl. 943 = Limodorum abortivum.
acuminata, Desf. Fl. Atlant. ii. 318 = tridentata.
acuta, Banks, ex Presl, Fl. Am. Sept. ii. 588 = Habenaria dilatata.
aetnensis, Tineo, in Guss. Fl. Sic. Syn. ii. 876 = tridentata.
affinis, C. Koch, in Linnaea, xxii. (1849) 284 = latifolia.

ORCHIS:—
agastachys, Fisch. ex Lindl. Gen. et Sp. Orch. 287 = Habenaria dilatata.
alata, Fleury, Orch. Rennes (1819), 17 = laxiflora.
alba, Lam. Fl. Fr. iii. 502 = Habenaria bifolia.
albida, Frivald. ex Hampe, in Flora, xx. (1837) 230 (= *Gymnadenia Frivaldskyana*).—Rumel.
albida, Scop. Fl. Carn. ed. II. ii. 201 = Habenaria albida.
alpina, Scop. l. c. 196 = Herminium alpinum.
ambigua, Degl. ex Le Gall, Fl. Morb. 585 = laxiflora.
ambigua ×, Kern. in Verh. Zool. Bot. Ges. Wien, xv. (1865) 205.—Europ.
ambigua, Martr. Fl. Tarn, 705 (Quid?).—Gallia.
amoena, Crantz, Stirp. Austr. ed. II. fasc. vi. 490 = ustulata.
anatolica, *Boiss. Diagn.* Ser. I. v. 56.—As. Min.; Syria.
angusticruris ×, Franch. in Humnicki, Mon. Pl. Orl. 27, nomen.—Gallia.
angustifolia, Bieb. Fl. Taur. Cauc. ii. 368 = iberica.
angustifolia, Loisel. ex Reichb. Pl. Crit. ix. 17. 1140 = latifolia.
angustifolia, Wimm. & Grab. Fl. Siles. ii. 252 = incarnata.
anthropomorpha, Steud. Nom. ed. I. 567 = Ophrys anthropomorpha.
anthropophora, All. Fl. Pedem. ii. 148 = Aceras anthropophora.
aphylla, Forsk. Fl. Aegypt. Arab. 156 = Habenaria aphylla, *R. Br.*
aphylla, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 240 = Epipogum aphyllum.
apifera, Salisb. Prod. 7 = Ophrys apifera.
appendiculata, Stokes, Bot. Mat. Med. iv. 291 = pyramidalis.
Arachnites, Scop. Fl. Carn. ed. II. ii. 194 = Ophrys Arachnites.
aristata, Fisch. ex Lindl. Gen. et Sp. Orch. 262 = latifolia.
Aschersoniana ×, Haussk. in Mitth. Geogr. Ges. Jena, ii. (1885) 223.—Europ.
atlantica, Willd. Sp. Pl. iv. 42 = Habenaria intacta.
atropurpurea, Tausch, in Flora, xiv. (1831) 223 = Habenaria nigra.
austriaca ×, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 139 = *Dietrichiana*.
barbata, Linn. f. Suppl. 399 = Disa barbata.
batrachites, Schrank, Baier. Reise, 86 = Habenaria viridis.
Beyrichii ×, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 208.—Europ.
bicornis, Gilib. Exercit. ii. 473 = pyramidalis.
bicornis, Jacq. Hort. Schoenb. ii. 26 = Satyrium ochroleucum.
bicornis, Linn. Pl. Afr. Rar. 26; Amoen. Acad. vi. 109 = Satyrium bicorne.
bicornis, Linn. Sp. Pl. ed. II. 1330 = Satyrium coriifolium.
bidentata, Ell. Sketch, ii. 488 = Habenaria herbiola.
Biermanni, Ortm. in Bot. Zeit. viii. (1850) 738 = maculata.
biflora, Linn. Sp. Pl. ed. II. 1330 = Disa torta.
bifolia, Linn. l. c. 939 = Habenaria bifolia.
bipunctata, Rafin. Préc. Découv. 43 = quadripunctata.
Bivonae, Todaro, in Imparziale Giorn. Sc. Lett. Art. (1840) 34 = longicruris.
blephariglottis, Willd. Sp. Pl. iv. 9 = Habenaria ciliaris.
Bonanniana, Tod. Orch. Sic. 57 = maculata.
Bornemanni ×, Aschers. in Oestr. Bot. Zeitschr. xv. (1865) 70 = longicornu.
Boryi, Reichb. f. Ic. Fl. Germ. xiii–xiv. 19 = quadripunctata.
brachiata, Gilib. Exercit. ii. 477 = militaris.
brachystachys, Urv. in Mém. Soc. Linn. Par. i. (1822) 377 = pyramidalis.
bracteata, Tenore, Prod. Fl. Nap. p. lii = O. Pseudosambucina.
bracteata, Willd. Sp. Pl. iv. 34 = Habenaria viridis.
Brancifortii, Bivona, Stirp. Rar. Sicil. i. [11] t. 1. f. 1 = quadripunctata.
Braunii ×, Halac. in Oestr. Bot. Zeitschr. xxxi. (1881) 137.—Europ.
brevicornis, Viv. Ann. Bot. i. II. 184 = patens.
brevicornu, Viv. Fl. Ital. Fragm. 11. t. 12. f. 2 = praec.
brevilabris, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 30 = tridentata.

ORCHIS:—
Bruniana ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 118.—Europ.
Burmanniana, Linn. Pl. Afr. Rar. 26; Amoen. Acad. vi. 108 = Bartholina pectinata.
calcarata, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) = Satyrium coriifolium.
calcarata, *Walt. Fl. Carol.* 220.—Am. bor.
camtschatica, *Cham. in Linnaea*, iii. (1828) 27.—Kamtschatka.
canariensis, *Lindl. Gen. et Sp. Orch.* 263.—Ins. Canar.
candidissima, Krock. Fl. Siles. iii. 16. t. 2 = Habenaria bifolia?
caprina, Bieb. Fl. Taur. Cauc. iii. 602 = hircina.
carnea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 294.—Afr. austr.
caspia, Trautv. in Act. Hort. Petrop. ii. (1873) 484 = laxiflora.
cassidea, Bieb. Fl. Taur. Cauc. iii. 600 = coriophora.
Cataphracta, *Sieber, ex Steud. Nom.* ed. II. ii. 223 (Quid?).—Hab.?
caucasica, *Regel, Ind. Sem. Hort. Petrop. App.* (1868) 22.—Reg. Caucas.
cercopithaca, Poir. in Lam. Encyc. iv. 593 = O. Simia.
cernua, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) (Quid?).—Afr. austr.
Champagneuxii, Barn. in Ann. Sc. Nat. Sér. II. xx. (1843) 380 = O. Morio.
chlorantha, Willk. & Lange, Prod. Fl. Hisp. i. 171, in syn. = Habenaria bifolia?
Chusua, *D. Don, Prod. Fl. Nep.* 23.—Reg. Himal.
cicadaria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 21 (Quid?).—Java.
ciliaris, Linn. Sp. Pl. 939 = Habenaria ciliaris.
cimicina, Bréb. Fl. Norm. ed. II. 296; Des Moul. in Act. Linn. Soc. Bord. xx. (1855) 706 = olida.
cimicina, Crantz, Stirp. Austr. ed. II. fasc. vi. 498 = coriophora.
cinerea, Schrank, Baier. Fl. i. 241 = militaris.
circumflexa, Bolus, in Journ. Linn. Soc. xix. (1882) 346, in syn. = Disperis secunda.
clavellata, Michx. Fl. Bor. Am. ii. 155 = Habenaria tridentata.
collina, [Soland. in] Russell, Aleppo, ed. II. ii. 264; Banks, ex Lindl. Gen. et Sp. Orch. 262 = saccata.
Columnae, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 227 = ustulata.
commelinefolia, Roxb. Hort. Beng. 63; Fl. Ind. iii. 451 = Habenaria commelinifolia.
commutata, Tod. Orch. Sic. 24 = tridentata.
comosa, Schur, Enum. Pl. Transs. 641 = angustifolia.
comosa, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 233 = maculata.
comosa, Scop. Fl. Carn. ed. II. ii. 198 = latifolia.
Comperiana, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* i. (18) 259.—As. Min.; Tauria.
compressiflora, Stokes, Bot. Mat. Med. iv. 292 = mascula.
condensa, Schur, Enum. Pl. Transs. 642 = latifolia.
condensata, Desf. Fl. Atlant. ii. 316 = pyramidalis.
conica, Willd. Sp. Pl. iv. 14 = tridenta.
conopea, Gren. Fl. Ch. Jurass. 751 = *conopsea*, Linn.
conopsea, Asso, Syn. Stirp. Arag. 130 = odoratissima.
conopsea, Linn. Sp. Pl. 942 = Habenaria conopsea.
convallariaefolia, Fisch. ex Lindl. Gen. et Sp. Orch. 217 (= *Platanthera convallariaefolia*).—Kamtschat.
cordata, Willd. Sp. Pl. iv. 27 = Habenaria cordata.
cordigera, *Fries, Novit. Fl. Succ. Mant.* iii. 130.—Europ.
coreosmus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = coriophora.
coriophora, *Linn. Sp. Pl.* 940.—Europ.; Oriens.
cornopica, Mill. Gard. Dict. ed. VIII. n. 8, sphalm. = *conopsea*, Linn.
cornuta, Houtt. Handleid. xii. 456 = Satyrium coriifolium.
cornuta, Linn. Pl. Afr. Rar. 27; Amoen. Acad. vi. 109 = Disa cornuta.
cornuta, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 21 (Quid?).—Java.
corsica, Viv. Fl. Cors. Sp. Nov. 16 = tridentata.
crenulata, Gilib. Exercit. ii. 474 = O. Morio.
cristata, Michx. Fl. Bor. Am. ii. 156 = Habenaria cristata.
crocea, Rafin. Atl. Journ. 119 (Quid?).—Am. bor.
cruenta, O. F. Muell. in Fl. Dan. t. 876; Retz. Prod. Fl. Scand. ed. II. 205 = incarnata.
cruenta, Roch, Pl. Banat. Rar. 34. t. 1. f. 1 = cordigera.

ORCHIS:—
cubitalis, Linn. Sp. Pl. 940 = Habenaria cubitalis.
cucullata, Linn. l. c. 939 = Habenaria cucullata.
Cupani, Tod. Orch. Sic. 56 (sp. dub.).—Sicil.
cuprina, Schur, Enum. Pl. Transs. 645, sphalm. = caprina.
curvifolia, F. Nyland. Spicil. Pl. Fennic. ii. 12. n. 25 = maculata.
cylindrica, *C. Koch, in Linnaea*, xix. (1847) 12.—As. Min.
Cyrilli, Tenore, Prod. Fl. Nep. p. lii = provincialis.
decipiens, Bianca, Nov. Pl. Sp. 1 = papilionacea.
densiflora, Wahlenb. in Vet. Acad. Handl. Stockh. (1806) 68 = Habenaria conopsea, *Benth.*
dentata, Sw. l. c. xxi. (1800) 207 (= *Plantanthera dentata*).—China.
depsata, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) = Satyrium candidum?
Dietrichiana ×, Bogenhard, Taschenb. Fl. Jena, 351.—Europ.
dilatata, Pursh, Fl. Am. Sept. ii. 588 = Habenaria dilatata.
discolor, Pursh, l. c. 586 = Tipularia discolor.
divaricata, Bor. Fl. Centr. Fr. ed. II. 522; ed. III. ii. 646, in obs. = angustifolia.
divaricata, Rich. ex Loisel. Fl. Gall. ed. I. 606; ed. II. ii. 267 = latifolia.
dolichorhiza, Fisch. ex Lindl. Gen. & Sp. Orch. 287 = Habenaria hyperborea.
Draconis, Linn. f. Suppl. 400 = Disa Draconis.
dubia ×, Camus, in Bull. Soc. Bot. Fr. xxxii. (1885) 216.—Gallia.
dubia, Le Gall. Fl. Morb. 585 = laxiflora.
Duquesnii, Nym. Syll. 358 = pyramidalis.
Durandii, Boiss. & Reut. Pugill. 111 = latifolia.
ecalcarata, Costa & Vayr, in Anal. Soc. Esp. Hist. Nat. ix. (1880) 97 = Habenaria albida.
elata, Hartm. Handb. Skand. Fl. ed. X. 202 = latifolia.
elata, Poir. Voy. Barb. ii. 248 (= *Gymnadenia elata*).—Afr. bor.
elatior, Afzel. ex Reichb. f. Ic. Fl. Germ. xiii–xiv. 51 = incarnata.
elatior, *Fries, Summa Veg. Scand.* 61, 217.—Fennia.
elegans, Heuff. ex Rochel, Bot. Reise Banat. 67; et in Verh. Zool. Bot. Ges. Wien, viii. (1858) 202 = palustris.
elodes, Griseb. Bild. Torfs. (in Goett. Studien, 1845); Gesam. Abh. 67 = incarnata.
ennensis, Guss. ex Tineo, Pl. Rar. Sicil. 10 = tridentata.
ensifolia, Vill. Hist. Pl. Dauph. ii. 29 = laxiflora.
entomantha, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 8 = Habenaria entomantha.
epidendroides, Willem. in Usteri, Ann. Bot. xviii. (1796) 53 (Quid?).—Ins. Maurit.
erubescens, Zucc. ex Lindl. Gen. & Sp. Orch. 277 = Habenaria odoratissima.
expansa, Tenore, Ind. Sem. Hort. Neap. (1829) 17; et Syll. 455 = papilionacea.
extensa, Hartm. ex Pritz. Ic. Ind. 768 = latifolia.
falcata, Thunb. Fl. Jap. 26 = Angraecum falcatum.
fallax, De Not. ex Willk. & Lange, Prod. Fl. Hisp. i. 168 = patens.
fasciculata, Tineo, in Guss. Fl. Sic. Syn. ii. 875 = O. Pseudo-sambucina.
fastigiata, Spreng. Syst. iii. 687 = Cynorchis fastigiata.
ferruginea, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 238 = Habenaria viridis.
festiva, Heuff. ex Rochel, Bot. Reise Banat. 67 = palustris.
filicornis, Linn. f. Suppl. 400 = Disa patens.
filicornis, Schum. & Thonn. Beskr. Guin. Pl. 397 = Habenaria filicornis.
fimbriata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 297 = Habenaria fimbriata.
fimbriata, Pursh, Fl. Am. Sept. ii. 588 = Habenaria fissa?
fissa, Muhl. ex Willd. Sp. Pl. iv. 40 = Habenaria fissa.
fissa, Pursh, Fl. Am. Sept. ii. 589 = Habenaria peramoena.
fistulata, Stokes, Bot. Mat. Med. iv. 296 = seq.
fistulosa, Moench, Meth. 713 = latifolia.
flava, Linn. Sp. Pl. 942 = Habenaria herbiola.
flava, Nutt. Gen. Am. ii. 188 = Habenaria integra.
flavescens, C. Koch, in Linnaea, xxii. (1849) 281 = O. Pseudo-sambucina.
flexicaulis, Linn. f. Suppl. 398; J. F. Gmel. Syst. 51 = Disa torta.

ORCHIS:—
flexuosa, Linn. Pl. Afr. Rar. 26; Amoen. Acad. vi. 108 = Disa flexuosa.
foliacea, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) = Satyrium carneum.
foliosa, Masson, ex [Ker-Gawl.] in Bot. Reg. sub t. 357 = longibracteata.
foliosa, Schur, Enum. Pl. Transs. 642 = cordigera.
foliosa, *Soland. ex Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 13.—Ins. Madera.
foliosa, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 206 = Habenaria foliosa.
formosa, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1812) 106.—Reg. Caucas.
fragrans, Pollini, Elem. Bot. ii. 157 = coriophora.
fragrans, Tenore, Fl. Nap. Prod. p. lii = longibracteata.
fuciflora, Crantz, Stirp. Austr. ed. II. fasc. vi. 483 = Ophrys Arachnites.
fusca, Jacq. Fl. Austr. iv. 4 = purpurea.
fuscata, Pall. Reise, ii. 124 = purpurea.
fuscescens, Linn. Sp. Pl. 943; Pursh, Fl. Am. Septr. ii. 587 = Habenaria herbiola.
fuscescens, Steph. Fl. Mosq. n. 610; ex Ledeb. Fl. Ross. iv. 61 = purpurea?
galeata, Poir. in Lam. Encyc. iv. 593 = militaris.
geniculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 25 = Habenaria geniculata.
Gennarii, Reichb. f. Ic. Fl. Germ. xiii–xiv. 172 = papilionacea.
germanorum, Moritz, Fl. Schweiz (1844), 509 = palustris.
Gervasiana, Tod. Orch. Sic. 57 = maculata?
gigantea, Sm. Exot. Bot. ii. 79 = Habenaria Susannae.
glariosa, Rafin. Atl. Journ. 150 (Quid?).—Am. bor.
glaucophylla, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 101 = mascula?
globosa, Brot. Fl. Lusit. i. 19 = tridentata.
globosa, *Linn. Syst.* ed. X. 1242.—Europ.; As. Min.
gracilis, Fisch. ex Lindl. Gen. et Sp. Orch. 285 = Habenaria tipuloides.
graminea, Crantz, Stirp. Austr. ed. II. fasc. vi. 480 = Herminium alpinum.
grandiflora, Bigel. Fl. Bost. ed. III. 343 = Habenaria fimbriata.
grandiflora, Heyne, in Wall. Cat. sub n. 7032 = Habenaria rotundifolia.
Grisebachii, *Pant. in Verh. Ver. Natur- u. Heilk. Presb.* ii. (1871–72) 27.—Montenegr.
Gussonii, Tod. in Atti Acc. Sc. Palerm. N. S. i. (1845) 8 = tridentata.
Habenaria, Linn. Syst. ed. X. 1242 = Habenaria macroceratitis.
haematodes, Reichb. Fl. Germ. Excurs. 126 = incarnata.
Halleri, Crantz, Stirp. Austr. ed. II. fasc. vi. 488 = globosa.
Hanrici, Hénon, in Ann. Soc. Agr. ix. (1846) 722; ex Gren. & Godr. Fl. Fr. iii. 288 = seq.
Hanrii, Jord. Obs. Pl. Crit. i. (1846) 27 = tridentata.
Hatagirea, D. Don, Prod. Fl. Nep. 23 = latifolia.
Hausknechtii ×, M. Schulze, in Mitth. Geogr. Ges. Jena, ii. (1885) 225.—Europ.
Heinzeliana ×, Reichhardt, in Verh. Zool.-Bot. Ges. Wien, xxvi. (1876) 464.—Europ.
herbiola, Pursh, Fl. Am. Sept. ii. 743 = Habenaria herbiola.
Herminium, Gren. in Gren. & Godr. Fl. Fr. iii. 300 = Herminium Monorchis.
heroica, *Clarke, Trav.* ii. 117.—Troas.
Heuffeliana, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 71 = palustris.
hircina, *Crantz, Stirp. Austr.* ed. II. fasc. vi. 484.—Europ.; Afr. bor.
hirtella, Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 207; Fl. Ind. Occ. iii. 139 = Physurus hirtellus.
hispida, Thunb. Prod. Pl. Cap. 4 = Holothrix parvifolia.
hispidula, Linn. Suppl. 401 = Holothrix parvifolia.
Hookeri, Wood, Am. Bot. & Flor. 327 = seq.
Hookeriana, Oakes, Cat. Verm. Pl. 27 = Habenaria Hookeriana.
Hostii, Tratt. Arch. Gew. 107. t. 122 (1814) = quadripunctata.
humilis, Michx. Fl. Bor. Am. ii. 155 = spectabilis.
huronensis, Nutt. Gen. Am. ii. 189 = Habenaria hyperborea.
hybrida, Boenngh. ex Reichb. Fl. Germ. Excurs. 125 = purpurea.

ORCHIS :—
hyemalis, *Rafin. Fl. Ludov.* 168; *et ex Lindl. Gen. et Sp. Orch.* 274.—Sicilia.
hyperborea, Linn. Mant. i. 121 = Habenaria hyperborea.
iberica, *Bieb. ex Willd. Sp. Pl.* iv. 25.—Graecia; Oriens.
ichneumonea, *Afzel. ex Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 207, nomen; *et ex Pers. Syn.* ii. 506.—Afr. trop.
imbricata, Vest, in Syll. Ratisb. i. (1824) 80 = ustulata.
impudica, Crantz, Stirp. Austr. ed. II. fasc. vi. 497 (sp. dub.).—Europ.
incarnata, *Linn. Fl. Suec.* 812.—Europ.; Oriens.
incarnata, Willd. Sp. Pl. iv. 30 = sambucina.
incisa, Muhl. ex Willd. l. c. iv. 40; Nutt. Gen. Am. ii. 189 = Habenaria fissa.
insectifera, Crantz, Stirp. Austr. ed. II. fasc. vi. 481 = *Ophrys insectifera*.
intacta, Link, in Schrad. Journ. ii. (1799) 322 = Habenaria intacta.
integra, Nutt. Gen. Am. ii. 188 = Habenaria integra.
intermedia, Meigen & Weniger, Syst. Verz. Pfl. Rheins, 106 = O. Morio.
italica, Poir. in Lam. Encyc. iv. 600 = longicruris.
Jacquini, Godr. Fl. Lorr. ed. I. iii. 33 = purpurea.
japonica, Thunb. Fl. Jap. 26 = Habenaria japonica.
kamtschatica, Ledeb. Fl. Ross. iv. 63 = camtschatica.
Koenigii, Gunner, Fl. Norveg. ii. 103; Retz. Fl. Scand. ed. I. 168 = Habenaria hyperborea.
lacera, Michx. Fl. Am. Bor. ii. 156 = Habenaria lacera.
lactea, Poir. in Lam. Encyc. iv. 594 = tridentata.
laeta, Steinh. in Ann. Sc. Nat. Sér. II. ix. (1838) 209 = provincialis.
Lagotis, Reichb. f. Ic. Fl. Germ. xiii. t. 403 = latifolia.
lanceata, A. Dietr. Fl. Boruss. t. 5 (1833) = incarnata.
lanceolata, A. Dietr. Fl. March. 807 (Index) sphalm. = praec.
lancibracteata, C. Koch, in Linnaea, xxii. (1849) 282 = maculata.
lapponica, Laest. ex Reichb. f. Ic. Fl. Germ. xiii. 59 = maculata.
lata, *Walt. Fl. Carol.* 220.—Carolina.
latifolia, *Linn. Sp. Pl.* 941.—Europ.; Afr. bor.; Oriens; Reg. Himal.
latifolia, Scop. Fl. Carn. ed. II. ii. 196 = incarnata.
laurentina, Bolós, ex Vayr. in Anal. Soc. Esp. Hist. Nat. ix. (1880) 96. t. 1 = sambucina.
laxiflora, *Lam. Fl. Fr.* iii. 504.—Europ.; Oriens.
laxiflora, C. A. Mey. Verz. Pfl. Cauc. 38 = palustris.
leptophylla, C. Koch, in Linnaea, xxii. (1849) 282 = iberica.
leucophaea, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 161 = Habenaria leucophaea.
leucostachya, Griseb. Spicil. Fl. Rumel. ii. 359 = provincialis.
Lindackeri, *Sieber, ex Steud. Nom.* ed. II. ii. 224 (Quid?).—Creta.
Lingua, All. Fl. Pedem. ii. 148 = Serapias Pseudocordigera.
Lingua, Scop. Fl. Carn. ed. II. ii. 187 = Serapias Lingua.
longebracteata, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 233. t. 2 = maculata.
longibracteata, *Bivona, Pl. Sic. Cent.* i. 57.—Europ. austr.
longibracteata, Schur, Enum. Pl. Transs. 643 = maculata.
longicornis, Lam. Encyc. iv. 591; Desf. Fl. Atlant. ii. 317. t. 246 = longicornu.
longicornu, Heyne, ex Wall. Cat. sub n. 7027 = Habenaria longicornu.
longicornu, *Poir. Voy. Barb.* ii. 247.—Reg. Mediterr.
longicornu, Salzm. ex Lindl. Gen. et Sp. Orch. 329 = Habenaria pauciflora, *Reichb. f.*
longicruris, *Link, in Schrad. Journ.* ii. (1799) 323.—Reg. Mediterr.
Loreziana ×, Bruegg. Beitr. Kenntn. Chur, 56; et in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 118.—Europ.
lucana, Spreng. Pugill. ii. 79 = O. Pseudo-sambucina.
lusitanica, Steud. Nom. ed. II. ii. 224 = latifolia.
lutea, Dulac, Fl. Hautes-Pyr. 125 = sambucina.
macedonica, Griseb. Spicil. Fl. Rumel. ii. 361 = maculata.
macra, Koch, Syn. Fl. Germ. ed. II. 789 = spuria.
macra, Lindl. Synop. ed. II. 260 = O. Simia.
macra, Schur, Enum. Pl. Transs. 641 = incarnata.

ORCHIS :—
macronectar, Vell. Fl. Flum. ix. t. 45 (Habenariae sp.).—Bras.
macrostachys, Tineo, Pl. Rar. Sicil. 7 = maculata.
maculata, *Linn. Sp. Pl.* 942.—Europ.; As. Min.
majalis, Reichb. Pl. Crit. vi. II. 7 = latifolia.
Markusii, Tineo, Pl. Rar. Sicil. 9 = O. Pseudo-sambucina.
Martrinii, Timb. in Bull. Soc. Bot. Fr. iii. (1856) 92 = coriophora.
mascula, Alsch. Fl. Jadrens. 210 = provincialis.
mascula, Crantz, Stirp. Austr. ed. II. fasc. vi. 500 = palustris.
mascula, *Linn. Fl. Suec.* 810.—Europ.; Afr. bor.; Oriens.
masculo-latifolia, *Lange, in Kjoeb. Vidensk. Meddel.* (1860) 78.—Hispan.
matodes, Reichb. f. Ic. Fl. Germ. xiii. t. 398 = incarnata.
mauritiana, Poir. in Lam. Encyc. iv. 601 = Angraecum gladiifolium?
mauritiana, Sieber, ex Steud. Nom. ed. II. i. 711; ii. 224 = Cynorchis fastigiata.
maxima, C. Koch, in Linnaea, xix. (1847) 14 = purpurea.
mediterranea, Guss. Pl. Rar. 365 = palustris.
mediterranea, Tenore, Syll. 455 = laxiflora.
membranacea, Afzel. ex Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 207, nomen; et ex Pers. Syn. ii. 506 = Habenaria membranacea.
Michauxii, Wood, Am. Bot. & Flor. 328 = Habenaria Michauxii.
micranthema, Buch.-Ham. ex Wall. Cat. sub n. 7069 (= *Peristylus Hamiltonianus*).—Reg. Himal.
militaris, Hornem. in Fl. Dan. t. 1277 = purpurea.
militaris, *Linn. Sp. Pl.* 941.—Europ.; As. bor.
militaris, Poir. in Lam. Encyc. iv. 592 = longicruris.
militaris, Puccin. ex Parl. Fl. Ital. iii. 521 = incarnata.
militaris, Sm. Engl. Bot. t. 1873 = O. Simia.
mimusops, Thuill. Fl. Par. ed. II. 458 = militaris.
miniata, Crantz, Stirp. Austr. ed. II. fasc. vi. 486 = Habenaria nigra.
mixta, Retz. Fl. Scand. ed. I. 167 = incarnata, sambucina.
Monorchis, Crantz, Stirp. Austr. ed. II. fasc. vi. 478 = Herminium Monorchis.
monorrhiza, Sw. Prod. Veg. Ind. Occ. 118 = Habenaria brachyceras, maculosa.
montana, F. W. Schmidt, Fl. Boëm. i. 35. t. 56 = Habenaria nigra.
moravica, Jacq. Ic. Rar. i. t. 182 = purpurea.
Moria, Retz. Fl. Scand. ed. I. 166, sphalm. = seq.
Morio, *Linn. Sp. Pl.* 940.—Europ.; As. med. et bor.
Morio, Ucria, Hort. Reg. Panorm. 382 = laxiflora.
moriopapilionacea ×, Timb. in Mém. Acad. Toul. Sér. IV. iv. (1854) 277 = papilionacea.
Moritziana, Bruegg. Beitr. Kennt. Chur, 58; et in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 118 = Habenaria nigra.
Munbyana, *Boiss. & Reut. Pugill. Pl. Nov.* 112.—Alger.
muscaria, Scop. Fl. Carn. ed. II. ii. 193 = Ophrys muscifera.
muscifera, Salisb. Prod. 7 = Ophrys muscifera.
musciflora, Schrank, Baier. Fl. i. 242 = Ophrys muscifera.
mutabilis, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 244 = formosa.
myodes, Bernh. Syst. Verz. Erf. 315, in syn. = Ophrys muscifera.
mysorensis, Heyne, ex Wall. Cat. n. 7038 B = Habernaria digitata, marginata.
Natalis, Tineo, Pl. Rar. Sicil. 8 = O. Pseudo-sambucina.
natolica, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 30 = angustifolia.
nervata, Marchand, in Hall, Bijdr. Nat. Wetensch. ii. (1827) 436 = militaris.
Nicodemi, Cyrill. ex Tenore, Prod. Fl. Nap. p. liii = O. Morio.
Nicodemi, Sieber, ex Steud. Nom. ed. II. ii. 224 (= *Gymnadenia humilis*).—Ital.
nigra, Scop. Fl. Carn. ed. II. ii. 200 = Habenaria nigra.
nivea, Nutt. Gen. Am. ii. 188 = Habenaria nivea.
obcordata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 23 = Habenaria galeandra.

ORCHIS :—
obcordata, Willem. in Usteri, Ann. Bot. xviii. (1796) 52 = (Cynorchidis sp.?).—Ins. Maurit.
obsoleta, Willd. Sp. Pl. iv. 34 = Habenaria obsoleta.
obtusa, Schur, Enum. Pl. Transs. 942 (Index) = mascula?
obtusata, Banks, ex Pursh, Fl. Am. Sept. ii. 588 = Habenaria obtusata.
obtusiflora, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 71 = mascula.
obtusifolia, Schur, l. c. 72 = maculata.
ochroleuca, Schur, Enum. Pl. Transs. 641 = O. Pseudo-sambucina.
ochroleuca, Tenore, Add. Fl. Neap. Syll. App. v. 45 = Habenaria bifolia.
odoratissima, *Linn. Syst.* ed. X. 1243.—Europ.
oestrifera, Bieb. Fl. Taur. Cauc. iii. 605 = Ophrys oestrifera.
officinalis, Salisb. Prod. 6 = O. Morio.
olbiensis, Ardoino, ex Moggr. Contr. Fl. Ment. t. 18 = provincialis?
olbiensis, Reut. ex Ardoino, Fl. Alpes-Marit. 353 = mascula.
olida, *Bréb. Fl. Normand.* ed. I. 259.—Gallia.
olivetorum, Gren. ex Nym. Consp. 693 = mascula.
ophioglossoides, Walt. Fl. Carol. 220 = Microstylis ophioglossoides.
orbiculata, Pursh, Fl. Am. Sept. ii. 588 = Habenaria Hookeriana.
ornithis, Jacq. Fl. Austr. ii. 23 = Habenaria conopsea, *Benth.*
ovalis, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 224. t. 1 = mascula.
pallens, *Linn. Mant.* ii. 292.—Europ. austr.; As. Min.
pallens, Moritzi, Fl. Schw. 508; Puccin. Syn. Fl. Luc. 476 = sambucina.
pallens, Savi, Fl. Pis. ii. 300 = provincialis.
palmata, Gilib. Exercit. ii. 479 = latifolia?
paludosa, Pall. Reise, iii. 320 = Malaxis paludosa.
palustris, *Jacq. Coll.* i. 75.—Europ.; Afr. bor.; Oriens.
panormitana, Tineo, in Guss. Fl. Sic. Syn. ii. 875 = patens.
papilionacea, *Linn. Syst.* ed. X. 1242.—Reg. Mediterr.
parasitica, Willem. in Usteri, Ann. Bot. xviii. (1796) 53 (Quid?).—Ins. Maurit.
Parlatoris, Tineo, Pl. Rar. Sicil. 29 = tridentata.
Parreissii, Presl, Bot. Bemerk. 112 = mascula.
parviflora, Poir. in Lam. Encyc. iv. 599 = Habenaria albida.
parviflora, Steud. Nom. ed. II. ii. 224 = Satyrium parviflorum.
parviflora, Tenore, Prod. Fl. Neap. p. liii = tridentata.
parviflora, Willd. Sp. Pl. iv. 27 = ustulata.
parvifolia, *Chaub. in St. Amans, Fl. Agen.* 369. t. 7.—Gallia.
patens, *Desf. Fl. Atlant.* ii. 318. t. 248.—Reg. Mediterr.
pauciflora, Fisch. ex Lindl. Gen. et Sp. Orch. 280 = (= *Gymradenia pauciflora*).—Dalmat.
pauciflora, Tenore, Prod. Fl. Neap. p. lii = provincialis.
pectinata, Sm. Exot. Bot. ii. 77 = Habenaria pectinata.
pectinata, Thunb. Prod. Pl. Cap. 4 = Bartholina pectinata.
peloria, Fouc. ex Poir. Encyc. Suppl. iv. 179 = Habenaria conopsea, *Benth.*
peramoena, Wood, Am. Bot. & Flor. 327 = Habenaria peramoena.
picta, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 431 = O. Morio.
picta, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
picta, Reinw. ex Lindl. in Journ. Linn. Soc. i. (1857) 179 = Anaectochilus setaceus.
pinetorum, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* v. 68.—Cilicia.
piramidalis, Neck. Delic. Gallo-Belg. ii. 371 = pyramidalis.
plantaginea, Roxb. Pl. Corom. i. 32. t. 37 = Habenaria platyphylla.
plantaginea, Sw. in Vet. Handl. Stockh. xxi. (1800) 207 = Spiranthes orchioides, Physurus plantagineus.
platychila, C. Koch, in Linnaea, xix. (1847) 13 = laxiflora.

ORCHIS:—
platyphyllos, Willd. Sp. Pl. iv. 11 = Habenaria platyphylla.
Polliniana, Spreng. Pugill. ii. 78 = coriophora.
pratensis, Salzm. ex Lindl. Gen. et Sp. Orch. 328 = Habenaria pratensis.
procera, Afzel. ex Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 207, nomen; et ex Pers. Syn. ii. 506 = Habenaria procera.
provincialis, *Balb. Misc. Alt.* 20. t. 2.—Reg. Mediterr.
Pseudo-apifera, *Caldesi, in Nuov. Giorn. Bot.* xii. (1880) 258.—Ital.
Pseudo-conopea, Gren. Fl. Ch. Jurass. 751 = Habenaria conopsea, *Benth.*
Pseudo-pallens, C. Koch, in Linnaea, xix. (1847) 13 = pallens.
Pseudo-pallens, Tod. Orch. Sic. 58 = provincialis.
Pseudo-sambucina, *Tenore, Syn.* ed. I. 72; ed. II. 64; ex ej. Fl. Nap. ii. 284.—Europ. austr.; Oriens.
psychodes, Linn. Sp. Pl. 943 = Habenaria fissa?
psychodes, Muhl. Cat. ed. I. 80 = Habenaria lacera,
punctulata, *Stev. in Lindl. Gen. et Sp. Orch.* 273.—As. Min.
purpurascens, Spreng. Syst. iii. 687 = Cynorchis purpurascens.
purpurea, *Huds. Fl. Angl.* ed. I. 334.—Europ.; As. Min.
pyramidalis, Bieb. Fl. Taur. Cauc. ii. 263 = tridentata.
pyramidalis, *Linn. Sp. Pl.* 940.—Europ.; Afr. bor.
pyrenaica, Philippe, Fl. Pyrén. ii. 354 = Habenaria conopsea, *Benth.*
quadripunctata, *Cyrill. ex Tenore, Prod. Fl. Neap.* p. liii.—Europ. austr.
quinqueseta, Michx. Fl. Bor. Am. ii. 155 = Habenaria Michauxii.
Raddeana, *Regel, Ind. Sem. Hort. Petrop.* (1869) 22.—Reg. Caucas.
radiata, Pers. Syn. ii. 502 = Habenaria radiata.
rariflora, C. Koch, in Linnaea, xix. (1847) 13 = anatolica.
recurva, Nyland. ex Fries, Summa Veg. Scand. i. 61 = maculata.
Regeliana ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 118.—Europ.
Reichenbachii, Mutel, Fl. Fr. iii. 245, 403 = Habenaria nigra.
repens, Eyster. ex Poir. Encyc. vi. 581 = Goodyera repens.
repens, *Wood, Am. Bot. & Flor.* 328 = Habenaria repens.
resupinata, *Naumb. in Roem. Arch.* ii. (1799) 238.—Europ.
Ricasoliana, Parl. in Atti 5a Riun. Sc. Ital. (1843) 730, nomen; Fl. Ital. iii. 474 = tridentata.
Richardi, *Tratt. ex Steud. Nom.* ed. II. ii. 224.—Hab.?
Rivini, Gouan, Illustr. 74 = militaris.
rivularis, Heuff. ex Schur, Enum. Pl. Transs. 642 = cordigera.
Robertiana, Loisel. Fl. Gall. ed. I. 606 = longibracteata.
romana, Sebast. & Mauri, Rom. Pl. Fasc. i. 12. t. 3 = O. Pseudo-sambucina.
rotundifolia, Banks, ex Pursh, Fl. Am. Sept. ii. 588 = Habenaria rotundifolia.
Roxburghii, Pers. Syn. ii. 503 = Habenaria platyphylla.
rubra, Jacq. Ic. Pl. Rar. 183 = papilionacea.
saccata, Reichb. Fl. Germ. Excurs. 122 = sambucina.
saccata, *Tenore, Prod. Fl. Neap.* p. liii.—Reg. Mediterr.; Oriens.
saccata, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 375 (an praec.?).—Colchis.
saccifera, Brongn. in Exp. Sc. Mor. Bot. 259 = maculata.
sagittalis, Linn. f. Suppl. 399 = Disa sagittalis.
sagittata, Munby, Fl. Algér. 100 = Habenaria intacta.
salina, Fronius, in Verh. Siebenb. Ver. Naturf. viii. (1857) 102 = sambucina.
salina, *Turcz. ex Lindl. Gen. et Sp. Orch.* 259.—Sibir. baical.
sambucina, Brot. Fl. Lusit. i. 21; Bieb. Fl. Taur. Cauc. ii. 367 = O. Pseudo-sambucina.
sambucina, *Linn. Fl. Suec.* 312.—Europ.
sancta, *Linn. Fl. Palaest.* n. 512; *Amoen. Acad.* iv. 464.—As. Min.; Syria.
satyroides, *Linn. Amoen. Acad.* vi. 109.—Afr. trop.
Schleicheri, Sweet, Hort. Brit. ed. I. 382; Brit. Flow. Gard. Ser. I. ii. t. 199 = sambucina.
Scopoli, Timb. Diagn. (1850); ex Gren. & Godr. Fl. Fr. iii. 288 = tridentata.

ORCHIS:—
scutellata, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 161 = Habenaria herbiola.
secunda, Thunb. Prod. Pl. Cap. 4 = Holothrix secunda.
secundiflora, Bertol. Pl. Rar. Ital. Dec. ii. 42 = Habenaria intacta.
sepulchralis, Boiss. & Heldr. Diagn. Ser. I. xiii. 10 = punctulata.
sesquipedalis, Willd. Sp. Pl. iv. 30 = latifolia.
setacea, Gilib. Exercit. ii. 482 = Habenaria conopsea, *Benth.*
setacea, Jacq. Enum. Pl. Carib. 28 = Habenaria maculosa.
sicula, Tineo, Pl. Rar. Sicil. 8 = O. Pseudo-sambucina.
signifera, Vest, in Syll. Ratisb. i. (1824) 79 = militaris.
Simia, *Lam. Fl. Fr.* iii. 507.—Europ.; Oriens.
Simia, Vill. Hist. Pl. Dauph. ii. 33 = tridentata.
Smithii, Sweet, Hort. Brit. ed. I. 381 = O. Simia.
solida, Moench, Meth. 713 = maculata.
sparsiflora, Tenore, ex Boiss. Fl. Orient. v. 67 = saccata.
spathulata, Linn. f. Suppl. 398 = Disa spathulata.
spathulata, Reichb. f. ex Benth. in Journ. Linn. Soc. xviii. (1880) 355 = Habenaria spathulata.
speciosa, Heuff. ex Rochel, Bot. Reise Banat. 67 = palustris.
speciosa, Host, Fl. Austr. ii. 527 = mascula.
speciosa, Linn. f. Suppl. 401; Thunb. Prod. Pl. Cap. 4 = Habenaria Bonatea.
spectabilis, *Linn. Sp. Pl.* 943.—Am. bor.
sphaerica, Bieb. Fl. Taur. Cauc. ii. 362 = globosa.
Spitzelii, *Sauter, ex Koch, Syn. Fl. Germ.* ed. I. 686.—Austria.
spuria, *Reichb. f. in Bot. Zeit.* vii. (1849) 891.—Europ.
squamosa, Poir. in Lam. Encyc. iv. 601 = Cynorchis squamosa.
Stabiana, Tenore, Syll. 453 = mascula.
stercoraria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 (Quid?).—Java.
Steveni, *Reichb. f. in Bot. Zeit.* vii. (1849) 892.—Reg. Caucas.; Persia.
Steveniana, *Comper. ex Léveillé, in Démid. Voy. Russ. Mérid.* ii. 168.—Tauria.
strateumatica, Linn. Sp. Pl. 943 = Zeuxine sulcata.
strictiflora, Opiz, Natural. x. (1825) 217 = angustifolia.
strictifolia, Opiz, Seznam, 70 = incarnata.
stylomisantha, *Vell. Fl. Flum.* ix. t. 46.—Bras.
suaveolens, Salisb. Prod. 7 = Habenaria conopsea, *Benth.*
suaveolens, Vill. Hist. Pl. Dauph. ii. 38 = Habenaria nigra.
sulphurea, Sims, Bot. Mag. t. 2569 = pallens.
sulphurea, Spreng. Syst. iii. 688, in syn. = O. Pseudo-sambucina.
Susannae, Linn. Sp. Pl. 939 = Habenaria Susannae.
Susannae, Thunb. Fl. Jap. 25 = Habenaria radiata.
syriaca, Boiss. & Bal. ex Boiss. Fl. Orient. v. 60 = O. Morio.
Tabernaemontani, C. C. Gmel. Fl. Bad. iii. 542 = laxiflora.
taurica, Lindl. Gen. et Sp. Orch. 271 = tridentata.
tenella, Linn. f. Suppl. 400 = Disa tenella.
Tenoreana, Guss. Fl. Sic. Syn. ii. 533 = tridentata.
tenuifolia, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26). (Quid?).—Afr. austr.
tenuifolia, C. Koch, in Linnaea, xxii. (1849) 281 = O. Pseudo-sambucina.
tenuis, Rottl. ex Wall. Cat. sub n. 7053 = Habenaria plantaginea.
tephrosanthos, Desf. Fl. Atlant. ii. 319 = longicruris.
tephrosanthos, Vill. Prosp. 16; Hist. Pl. Dauph. ii. 32 = O. Simia.
tetragona, Heuff. in Flora, xvi. (1833) 363 = maculata.
Tharandina, Reichb. f. Ic. Fl. Germ. xiii. t. 403 = latifolia.
Timbali ×, Velen. in Sitzb. Boehm. Acad. (1882) 254.—Europ.
tipuloides, Linn. f. Suppl. 401 = Tipularia camtschatica.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 72 = maculata.
Traunsteineri, Sauter, ex Koch, Syn. Fl. Germ. ed. I. 688 = angustifolia.
trichocera, Brongn. in Exp. Sc. Mor. Bot. 260 = quadripunctata.
tridactylites, Webb & Berth. Phyt. Canar. iii. 307 = Habenaria tridactylites.

ORCHIS:—
tridentata, Muhl. ex Willd. Sp. Pl. iv. 41 = Habenaria tridentata.
tridentata, *Scop. Fl. Carn.* ed. II. ii. 190.—Europ.
tripetaloides, Linn. f. Suppl. 398 = Disa venosa, *Lindl.*
triphylla, C. Koch, in Linnaea, xxii. (1849) 283 = latifolia.
triphylla, Spreng. Syst. iii. 68 (= *Gymnadenia triphylla*).—Ins. Maurit.
triplicata, Willem. in Usteri, Ann. Bot. xviii. (1796) 52 = Calanthe veratrifolia.
Uechtritziana ×, Hausskn. in Mitth. Geogr. Ges. Jena, ii. (1885) 225.—Europ.
umbrosa, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 504 = latifolia.
undulata, Poir. in Lam. Encyc. iv. 588 = Disa flexuosa.
undulatifolia, Bivona, Sic. Pl. Cent. ii. 44 = longicruris.
uniflora, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 25 = Habenaria triflora?
uniflora, Roxb. Hort. Beng. 63; Fl. Ind. iii. 452 = Diplomeris pulchella.
Urvilleana, Steud. Nom. ed. II. ii. 225 = sancta.
Urvillei, Steud. l. c. 224, in syn. = praec.
ustulata, *Linn. Sp. Pl.* 941.—Europ.
vallesiaca ×, Spiess, in Oestr. Bot. Zeitschr. xxvii. (1877) 352.—Helvet.
varia, J. F. Gmel. Syst. 53 (Quid?).—Guian.
variegata, All. Fl. Pedem. ii. 147 = tridentata.
variegata, Schrank, Baier. Fl. i. 241 = Habenaria nigra?
ventricosa, Steud. Nom. ed. II. ii. 225 = Serapias ventricosa.
vernalis, Salisb. Prod. 6 = mascula.
vestita, *Cav. in Anal. Cienc. Nat.* vi. (1803) 142.—Hab.?
virens, Scop. Fl. Carn. ed. II. ii. 199 = Habenaria viridis.
virescens, Willd. Sp. Pl. iv. 37 = Habenaria herbiola.
virescens, Zollik. ex Gaud. Fl. Helv. v. 497 = Habenaria chlorantha.
viridiflora, Rottl. ex Pers. Syn. ii. 503. (= seq.?).—Ind. or.
viridiflora, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 206 = Habenaria viridiflora.
viridis, Crantz, Stirp. Austr. ed. II. fasc. vi. 491 = Habenaria viridis.
Welwitschii, *Reichb. f. Ic. Fl. Germ.* xxiii-xxiv. 183.—Lusitan.
zoophora, Thuill. Fl. Par. ed. II. 459 = O. Simia.

ORCHITES, Schur, Enum. Pl. Transs. 942 (1866) = **Orchis**, Tourn.
globosa, Schur, l. c. = O. globosa.

ORCYA, Vell. Fl. Flum. 344 (1825) = **Acanthospermum**, Schrank (Compos.).
adhaerescens, Vell. l. c.; et viii. t. 83 = A. Brasilum.

OREACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1104 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
Manni, *Benth. l. c.*; *et in Hook. Ic. Pl.* t. 1211.—Afr. trop.

OREAMUNOA, Oerst. in Kjoeb. Vidensk. Meddel. (1870) 166. t. 2.—(Cf. OREOMUNNEA) = **Engelhardtia**, Leschen. (Jugland.).
pterocarpa, Oerst. l. c. 167 = E. Oreamunnea.

OREANTHES, Benth. Pl. Hartw. 140 (1844). *VACCINIACEAE*, Benth. & Hook. f. ii. 569.
buxifolius, *Benth. Pl. Hartw.* 140.—Peruv.

OREANTHUS, Rafin. in Ser. Bull. Bot. i. (1830) 216 = **Heuchera**, Linn. (Saxifr.).
bracteatus, Rafin. l. c. = H. bracteata.

OREAS, Cham. & Schlecht. in Linnaea, i. (1826) 29. t. 1 = **Braya**, Sternb. & Hoppe (Crucif.).
involucrata, Cham. & Schlecht. l. c. = B. involucrata.

OREINOTINUS, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 281 = **Viburnum**, Linn. (Caprifol.).
acutifolius, Oerst. l. c. 293. t. 6. f. 25 = V. acutifolium.
ayavacensis, Oerst. l. c. 290 = V. ayavacense.
costa-ricanus, Oerst. l. c. = V. costa-ricanum.
discolor, Oerst. l. c. 287 = V. discolor.
divaricatus, Oerst. l. c. = V. divaricatum.
ferrugineus, Oerst. l. c. 286 (err. typ. 386).—N. Granat.
furcatus, Oerst. l. c. 284.—N. Granat.
fuscus, Oerst. l. c. 289 = V. fuscum.
glabratus, Oerst. l. c. 290 = V. glabratum.
Halli, Oerst. l. c. 292.—Ecuador.

OREINOTINUS:—
Hartwegi, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 289 = Viburnum Hartwegi.
Jamesonii, Oerst. l. c. 288.—Ecuador.
lasiophyllus, Oerst. l. c. 287 = V. lasiophyllum.
laurifolius, Oerst. l. c. 291.—Peruv.
Mathewsii, Oerst. l. c.—Peruv.
membranaceus, Oerst. l. c. 284 = V. membranaceum.
microcarpus, Oerst. l. c. 282 = V. microcarpum.
microphyllus, Oerst. l. c. 293 = V. microphyllum.
obovatus, Oerst. l. c. 286 (err. typ. 386) = V. obovatum, *Ruiz & Pav.*
pichinchensis, Oerst. l. c. 287 = V. pichinchense.
reticulatus, Oerst. l. c. 286 (err. typ. 386) = V. reticulatum.
rhombifolius, Oerst. l. c. 283 = V. rhombifolium.
stellato-tomentosus, Oerst. l. c. = V. stellato-tomentosum.
stellatus, Oerst. l. c. 292 = V. stellatum.
stenocalyx, Oerst. l. c. 285 = V. stenocalyx.
sulcatus, Oerst. l. c. 287 = V. sulcatum.
tiliaefolius, Oerst. l. c. 282. t. 6 = V. tiliaefolium.
tinoides, Oerst. l. c. 291 = V. tinoides.
triphyllus, Oerst. l. c. 290 = V. triphyllum.
undulatus, Oerst. l. c. 288.—N. Granat.
villosus, Oerst. l. c. = V. villosum.
Wendlandi, Oerst. l. c. 283 = V. Wendlandi.

ORELIA, Aubl. Pl. Gui. i. 270. t. 106 (1775) = **Allamanda**, Linn. (Apocyn.).
grandiflora, Aubl. l. c. 271 = A. cathartica.

OREOBLITON, Dur. & Moq. in Duch. Rev. Bot. ii. (1847) 428. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 52.
NERETIA, Moq. in DC. Prod. xiii. II. 58 (1849).
chenopodioides, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 367.—Alger.
thesioides, *Dur. & Moq. in Duch. Rev. Bot.* ii. (1847) 428.—Alger.

OREOBOLUS, R. Br. Prod. 235 (1810). *CYPERACEAE*, Benth. & Hook. f. iii. 1057.
clandestinus, Phil. in Linnaea, xxix. (1857-58) 81 = obtusangulus.
distichus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 109 = Pumilio.
furcatus, *H. Mann, in Proc. Am. Acad.* vii. (1867) 210.—Ins. Sandvic.
obtusangulus, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 99; *et Atlas*, 63. t. 2. f. 1.—Reg. Magell.
pectinatus, *Hook. f. Fl. Antarct.* 87. t. 49.—N. Zel.; Ins. Antarct.
Pumilio, *R. Br. Prod.* 236.—Austral.; N. Zel.
strictus, *Berggr. in Minneskr. Fisiog. Sällsk. Lund* (1877), n. 8. p. 25.—N. Zel.

OREOBULUS, Boeck. in Linnaea, xxxviii. (1874) 230, sphalm. = praec.

OREOCALLIS, R. Br. in Trans. Linn. Soc. x. (1810) 196 = **Embothrium**, Forst. (Proteac.).
grandiflora, R. Br. l. c. 197 = E. grandiflorum.
Ruizii, Klotzsch, in Linnaea, xx. (1847) 474.—Peruv.
Wickhami, F. Muell. Fragm. viii. 164 = E. Wickhami.

OREOCHARIS, Benth. in Benth. & Hook. f. Gen. ii. 1021 (1876). *GESNERACEAE*, Benth. & Hook. f. l. c.
Auricula, *C. B. Clarke, in DC. Monog. Phan.* v. 64.—China.
Benthami, *C. B. Clarke, l. c.* 63.—China.
filipes, *Hance, in Journ. Bot.* xxi. (1883) 166.—China.
lanuginosus, *Benth. & Hook. f. Gen.* ii. 1021.—China.
Maximowiczii, C. B. Clarke, in DC. Monog. Phan. v. 63 = lanuginosus.
notha, *C. B. Clarke, l. c.* 64.—Ins. Philipp.
primuloides, *Benth. & Hook. f. Gen.* ii. 1021.—Japon.

OREOCHARIS, Lindl. Veg. Kingd. 656 (1847) = **Mertensia**, Roth (Borag.).

OREOCHLOA, Link, Hort. Berol. i. 44 (1827). *GRAMINEAE*, Benth. & Hook. f. iii. 1188.
disticha, *Link, l. c.*—Europ.
pedemontana, *Reut. in Boiss. & Reut. Pugill. Pl. Nov.* 126.—Hispan.

OREOCNIDA, Miq. Pl. Jungh. i. 39 (1851) = **Villebrunea**, Gaudich. (Urtic.).
acuminata, Kurz, For. Fl. Brit. Burma, ii. 427 = V integrifolia.
frutescens, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 131 = V. frutescens.
integrifolia, Miq. l. c. iv. 306 = V. integrifolia.
major, Miq. Pl. Jungh. 41 = V. scabra.
murina, Miq. Fl. Ind. Bat. i. II. 271 = V. murina.
pilosula, Miq. Pl. Jungh. 41 = V. glomerata.
rhodopleura, Miq. Fl. Ind. Bat. i. II. 271 = V. rhodopleura.
rubescens, Blume, ex Miq. in Zoll. Syst. Verz. Ind. Archip. 101 = V. scabra.
rubescens, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 306 = V. scabra.
rufescens, Miq. Fl. Ind. Bat. i. II. 271 = V. rufescens.
semicrenata, Miq. l. c. 270 = V. semicrenata.
sylvatica, Miq. Pl. Jungh. 40 = V. integrifolia.
trinervis, Miq. Fl. Ind. Bat. i. II. 270 = V. trinervis.

OREOCOME, Edgew. in Proc. Linn. Soc. i. (1845) 252; et in Trans. Linn. Soc. xx. (1846) 54 = **Selinum**, Linn. (Umbell.).
Candolliana, Edgew. in Trans. Linn. Soc. xx. (1846) 55 = S. tenuifolium.
Candollii, Edgew. in Proc. Linn. Soc. i. (1845) 253 = S. tenuifolium.
cicutaria, Edgew. l. c. = S. tenuifolium.
elata, Edgew. l. c. = S. tenuifolium.
filicifolia, Edgew. l. c. = S. tenuifolium.

OREOCOSMUS, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 37 = **Tibouchina**, Aubl. (Melastom.).
Galeottianus, Naud. l. c. 38 (= *Pleroma Galeottianum, scabriusculum*).—Mexic.
Ghiesbreghtii, Naud. l. c. 39 (= *Pleroma scabriusculum*).—Mexic.
monticola, Naud. l. c. 38 (= *Pleroma Naudinianum*).—Mexic.
Naudinianus, Naud. l. c. 39 (= *Pleroma Naudinianum*).—Mexic.
tortuosus, Naud. l. c. 38 (= *Pleroma mexicanum*).—Mexic.

OREODAPHNE, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 39 = **Ocotea**, Aubl. (Laurin.).
aciphylla, Nees & Mart. ex Nees, l. c. 43.—Bras.
acutangula, Miq. ex Meissn. in DC. Prod. xv. I. 121.—Bras.
acutifolia, Nees, in Linnaea, viii. (1833) 42.—Bras.; Reg. Argent.
Adenotrachelium, Nees, Syst. Laurin. 431.—Bras.
alba, A. Rich. Fl. Cub. Fanerog. ii. 189 = Phoebe elongata.
alnifolia, Meissn. in DC. Prod. xv. I. 132.—N. Granat.
amazonica, Meissn. l. c. 123.—Bras.
amoena, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 44.—Bras.; Reg. Argent.
angustifolia, Miq. ex Meissn. in DC. Prod. xv. I. 158 = Nectandra puberula.
angustifolia, Nees, Syst. Laurin. 418 = Ocotea angustifolia.
antioquiana, Meissn. in DC. Prod. xv. I. 116 = Persea costata.
bahiensis, Meissn. l. c. 130.—Bras.
Benthamiana, Nees, in Linnaea, xxi. (1848) 521 = Phoebe Hartwegii.
Beyrichii, Nees, Syst. Laurin. 435.—Bras.
Blanchetii, Meissn. in DC. Prod. xv. I. 113.—Bras.
Bofo, Nees, Syst. Laurin. 441 = Ocotea Bofo.
Boissieriana, Meissn. in DC. Prod. xv. I. 123.—Bras.
Bowiei, Walp. Ann. i. 576 = Cryptocarya australis.
brachybotrya, Meissn. in DC. Prod. xv. I. 127.—Bras.
bracteolata, Meissn. l. c. 124 = Ocotea bracteolata.
bracteosa, Meissn. l. c. 114.—Bras.
bullata, Nees, Syst. Laurin. 449 = Ocotea bullata.
californica, Nees, l. c. 463 = Umbellularia californica.
canescens, Meissn. in DC. Prod. xv. I. 122 = Nectandra psammophila.
caracasana, Nees, in Linnaea, xxi. (1848) 522.—N. Granat.
caudata, Miq. Stirp. Surin. Sel. 202 (= *Camphoromoea surinamensis*?).—Guian.
caudata, Nees, in Linnaea, xxi. (1848) 519.—Guian.
cayennensis, Meissn. in DC. Prod. xv. I. 133.—Guian.
celastroides, Meissn. l. c. 126.—N. Granat.

OREODAPHNE:—
cernua, Griseb. Fl. Brit. W. Ind. i. 284.—Ins. Dominic.
cernua, Nees, Syst. Laurin. 424 = Ocotea Sieberi.
citrosmoides, Nees, l. c. 384 = Ocotea citrosmoides.
colombiana, Meissn. in DC. Prod. xv. I. 137.—N. Granat.
commutata, Nees, Syst. Laurin. 428 = Ocotea commutata.
confertiflora, Meissn. in DC. Prod. xv. I. 131.—Bras.
confusa, Meissn. l. c. 126.—Bras.
coriacea, Griseb. ex Meissn. l. c. 159 = Nectandra membranacea.
coriacea, Nees, Syst. Laurin. 454 = Nectandra Willdenoviana, patens.
costulata, Nees, in Linnaea, xxi. (1848) 520 = *Neesiana*, Nectandra cuspidata.
crassifolia, Nees, l. c. 520 = Misanteca crassiramea.
crassiramea, Meissn. in DC. Prod. xv. I. 117.—Bras.
cuneata, Nees, Syst. Laurin. 385.—Peruv.
cuneifolia, Nees, in Linnaea, xxi. (1848) 518 = *obovata*.
cuprea, Meissn. in DC. Prod. xv. I. 114.—Peruv.
cupularis, Nees, Syst. Laurin. 438 (= *Mespilodaphne cupularis*).—Ins. Mascar.
cymosa, Nees, l. c. 437 (= *Mespilodaphne cupularis*).—Ins. Mascar.
daphnifolia, Meissn. in DC. Prod. xv. I. 127.—Bras.
declinata, Meissn. l. c. 114.—Bras.
diospyrifolia, Meissn. l. c. 126.—Bras.
dispersa, Mart. in Flora, xxiv. (1841) II. Beibl. 7 = Nectandra polita.
dispersa, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 43 = *confusa, japurensis*.
dispersa, Nees, ex Meissn. in DC. Prod. xv. I. 129 = *maynensis*.
divaricata, Nees, Syst. Laurin. 393.—Guian.?
domingensis, Nees, l. c. 439 = Ocotea botryophylla.
dominicana, Meissn. in DC. Prod. xv. I. 139.—Ins. Dominic.
effusa, Meissn. l. c. 120 = Ocotea effusa.
estrellensis, Meissn. l. c. 126.—Bras.
Evonymodaphne, Meissn. l. c. 139.—Peruv.
exaltata, Nees, Syst. Laurin. 406 = Nectandra sanguinea.
fallax, Miq. Stirp. Surin. Sel. 203.—Guian.
fasciculata, Nees, in Linnaea, xxi. (1848) 521 = Ocotea firmula.
Fensliana, Meissn. in DC. Prod. xv. I. 117 = Ocotea velutina.
floribunda, Benth. in Benth. & Hook. f. Gen. Pl. iii. 158 = Ocotea Cujumary.
floribunda, Nees, Syst. Laurin. 411.—Bras.
florulenta, Meissn. in DC. Prod. xv. I. 125.—Bras.
foetens, Nees, Syst. Laurin. 449 = Ocotea foetens.
fragrans, Meissn. in DC. Prod. xv. I. 127.—Jamaic.
frondosa, Meissn. in Mart. Fl. Bras. v. II. 309.—Bras.
gaboonensis, Meissn. in DC. Prod. xv. I. 130 = Beilschmiedia gaboonensis (in Addend.).
glaberrima, Meissn. l. c. 119 = Ocotea lanceolata.
glauca, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 42 (= *Mespilodaphne glauca*).—Bras.
glaucina, Meissn. in DC. Prod. xv. I. 134.—Bras.
glomerata, Nees, in Linnaea, xxi. (1848) 515.—Guian.
gracilis, Meissn. in DC. Prod. xv. I. 128.—Am. trop.
grandifolia, Nees, in Linnaea, xxi. (1848) 517 = Ocotea microcarpa.
guianensis, Nees, l. c. 516 = Ocotea guianensis.
helicterifolia, Meissn. in DC. Prod. xv. I. 123 = Ocotea helicterifolia.
Hookeriana, Meissn. l. c. 131.—Bras.
Hostmanniana, Miq. Stirp. Surin. Sel. 202 = *Martiniana*.
Humboldti, Meissn. in DC. Prod. xv. I. 129 = Ocotea umbrosa, *H. B. & K.*
indecora, Nees, Syst. Laurin. 407 = Ocotea indecora.
Jacobinae, Meissn. in DC. Prod. xv. I. 122.—Bras.
Jacquini, Meissn. l. c. 114.—Ins. Martinic.
japurensis, Meissn. l. c. 124.—Bras.
kavaiensis, H. Mann, in Mem. Bost. Soc. Nat. Hist. i. IV. (1869) 536 = Cryptocarya kawaiensis (in Addend.).
Klotzschiana, Nees, in Linnaea, xxi. (1848) 523 = Ocotea Klotzschiana.
Kunthiana, Meissn. in DC. Prod. xv. I. 119.—Bras.
lanata, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 43 (= *Mespilodaphne lanata*).—Bras.
lancifolia, Nees, Syst. Laurin. 410 = Ocotea subacris.
Langsdorffii, Meissn. in DC. Prod. xv. I. 128.—Bras.
leptobotra, Nees, in Linnaea, xxi (1848) 520.—Peruv.

OREODAPHNE:—
Leucoxylon, Nees, Syst. Laurin. 413 = Ocotea Leucoxylon.
Lindeniana, A. Rich. Fl. Cub. Fanerog. ii. 189 = Ocotea botryophylla.
Lobbii, Meissn. in DC. Prod. xv. 1. 136.—Bras.
longifolia, Nees, Syst. Laurin. 391 = Ocotea longifolia.
lucida, Meissn. in DC. Prod. xv. 1. 127.—Bras.
macrophylla, Meissn. l. c. 131.—Bras.
macrothyrsus, Meissn. l. c. 115 = Ocotea Cujumary.
Mannii, Meissn. l. c. 130 = Beilschmiedia Mannii (in Addend.).
maranhana, Meissn. l. c. 138.—Bras.
marginata, Nees, Syst. Laurin. 460 (= *Mespilodaphne cupulans*).—Ins. Mascar.
marowynensis, Miq. Stirp. Surin. Sel. 201 (= *Camphoromoea surinamensis*).—Guian.
Martiana, Nees, in Linnaea, viii. (1833) 41 = Ocotea lanceolata.
Martiniana, Nees, Syst. Laurin. 415.—Guian.
Maximilianea, Nees & Mart. ex Nees, l. c. 451 (= *Mespilodaphne prolifera*).—Bras.
maynensis, Meissn. in DC. Prod. xv. 1. 129.—Bras.
maynensis, Poepp. ex Meissn. l. c. 32 = Phoebe peruviana.
megaphylla, Meissn. l. c. 131.—Bras.
mexicana, Meissn. l. c. 118 = Ocotea mexicana.
microbotrys, Meissn. l. c. 125.—Bras.
minarum, Steud. Nom. ed. II. ii. 226 = Ocotea umbrosa.
minutiflora, Meissn. in DC. Prod. xv. 1. 130 = Beilschmiedia minutiflora (in Addend.).
Moritziana, Nees, in Linnaea, xxi. (1848) 515.—Bras.
mucosa, Nees, Syst. Laurin. 401 = Acrodiclidium salicifolium.
myriantha, Meissn. in DC. Prod. xv. 1. 120.—Bras.
Neesiana, Meissn. l. c. 114.—Guian.
nitidula, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 41.—Bras.
notata, Nees & Mart. l. c. 42 (= *Mespilodaphne notata*).—Bras.
nutans, Nees, Syst. Laurin. 421 (= *Mespilodaphne nutans*).—Bras.
oblongo-obovata, Nees, in Linnaea, xxi. (1848) 518 = *cuneata*.
obovata, Meissn. in DC. Prod. xvi. 1. 119.—Peruv.
obovata, A. Rich. Fl. Cub. Fanerog. ii. 189 (= *Nemodaphne cuneata*).—Cuba.
obtusata, Nees, Syst. Laurin. 447 (= *Mespilodaphne cupularis*).—Ins. Maurit.
odorata, Meissn. in DC. Prod. xv. 1. 116.—Bras.
opifera, Nees, Syst. Laurin. 390 = Ocotea opifera.
pallida, Meissn. in DC. Prod. xv. 1. 115.—Bras.
paraënsis, Meissn. l. c. 126.—Bras.
parviflora, Griseb. in Goett. Abh. vi. (1857) 219; et ex Meissn. l. c. 174 = Aydendron argenteum.
parviflora, Nees, Syst. Laurin. 415.—Jamaic.
pauciflora, Nees, l. c. 409.—Bras.
perseoides, Nees, ex Meissn. in DC. Prod. xv. 1. 141 = Ocotea puberula.
Petalanthera, Meissn. l. c. 125.—Bras.
phillyreoides, Nees, Syst. Laurin. 400 (= *Mespilodaphne phillyraeoides*).—Bras.
Poeppigiana, Nees, l. c. 404.—Peruv.
polyantha, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 44 = Ocotea polyantha.
pomaderroides, Meissn. in DC. Prod. xv. 1. 128.—Bras.
porosa, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 44.—Peruv.
psychotrioides, Benth. Pl. Hartw. 50 = Ocotea Klotzschiana.
psychotrioides, Nees, Syst. Laurin. 452 = Ocotea psychotrioides.
pubescens, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 45.—Bras.
pulchella, Nees & Mart. ex Nees, l. c. 40 = Ocotea pulchella.
rariflora, Meissn. in DC. Prod. xv. 1. 123.—Bras.
regalis, Regel, Gartenfl. (1856) 366.—Hab.?
Regeliana, Meissn. in DC. Prod. xv. 1. 132 = Ocotea lanceolata.
retroflexa, Nees, Syst. Laurin. 412 = Ocotea botryophylla.
Riedelii, Meissn. in DC. Prod. xv. 1. 130.—Bras.
rigens, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 40 = Ocotea tristis.

OREODAPHNE:—
rigida, Meissn. in DC. Prod. xv. 1. 115.—Bras.
rufo-tomentosa, Meissn. l. c. 45 = Ocotea rufo-tomentosa.
Schomburgkiana, Nees, in Linnaea, xxi. (1848) 519.—Guian.
Schottii, Meissn. in DC. Prod. xv. 1. 133.—Bras.
Sellowii, Meissn. l. c.—Bras.
semecarpifolia, Meissn. l. c. 120.—Bras.
semicompleta, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 43.—Bras.
sericea, Nees, l. c. xxi. (1848) 516 = Ocotea guianensis.
Sieberi, Meissn. in DC. Prod. xv. 1. 137 = Ocotea Sieberi.
spectabilis, Meissn. l. c. 119.—Bras.
Spixiana, Nees, Syst. Laurin. 382 = Ocotea rufo-tomentosa.
splendens, Meissn. in DC. Prod. xv. 1. 129.—Guian.
strumosa, Griseb. Fl. Brit. W. Ind. 284 = Nectandra membranacea.
suaveolens, Meissn. in DC. Prod. xv. 1. 136.—Paraguay.
subalpina, Meissn. l. c. 137.—Bras.
sublanuginosa, Nees, in Linnaea, xxi. (1848) 515.—Peruv.
subtriplinervia, Meissn. in DC. Prod. xv. 1. 125 = Ocotea subtriplinervia.
sylvatica, Meissn. in Mart. Fl. Bras. v. 11. 228.—Bras.
tampicensis, Meissn. in DC. Prod. xv. 1. 136 = Ocotea tampicensis.
tarapotana, Meissn. l. c. 129.—Peruv.
teleiandra, Meissn. l. c. 138.—Bras.
terminalis, Nees, ex Meissn. in DC. Prod. xv. 1. 139 = *O. Evonymodaphne*.
terminalis, Nees, Syst. Laurin. 426 = *leptobotra*.
tetragona, Meissn. in Kjoeb. Vidensk. Meddel. (1870) 137.—Bras.
thymelaeodes, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 42 = Ocotea lanceolata.
tinctoria, 'Arruda,' ex Rosenth. Syn. Pl. Diaph. 235.—Bras.
tovarensis, Klotzsch & Karst. ex Nees, in Linnaea, xxi. (1848) 511.—Venezuela.
tristis, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 40 = Ocotea tristis.
Tweediei, Meissn. in DC. Prod. xv. 1. 136.—Bras.
umbrosa, Nees, Syst. Laurin. 388 = Ocotea umbrosa.
umbrosa, Nees, l. c. 456 (nec p. 388) (Nectandrae sp.).—N. Granat.
undulata, Meissn. in DC. Prod. xv. 1. 115.—Bras.
urophylla, Meissn. l. c. 139.—Guian.
vaccinioides, Meissn. l. c. 127.—Bras.
variabilis, Nees, Syst. Laurin. 429 = Ocotea variabilis.
Velloziana, Meissn. in DC. Prod. xv. 1. 132.—Bras.
velutina, Nees, Syst. Laurin. 386 = Ocotea velutina.
venulosa, Meissn. in DC. Prod. xv. 1. 126.—Bras.
verruculosa, Meissn. l. c. 127.—Bras.
vesiculosa, Nees & Mart. ex Nees, in Linnaea, viii. (1833) 44.—Bras.
Warmingii, Meissn. in Kjoeb. Vidensk. Meddel. (1870) 136 = Ocotea puberula.
Willdenoviana, Nees, Syst. Laurin. 412 = Ocotea botryophylla.

OREODOXA, Willd. in Mém. Acad. Berl. 1804 (1807) 34, partim. *PALMAE*, Benth. & Hook. f. iii. 899.
acuminata, Willd. in Mém. Acad. Berl. 1804 (1807), 35 = Euterpe acuminata.
frigida, *H. B. & K. Nov. Gen. et Sp.* i. 304.—Ecuador.
Ghiesbreghtii, Hort.—Cf. Gard. Chron. (1870) 561 = Gaussia Ghiesbrechti?
Manaele, *Mart. Hist. Nat. Palm.* iii. 310.—Ins. S. Doming.
oleracea, *Mart. l. c.* 166. t. 156. f. 1, 2.—Ind. occ.
praemorsa, Willd. in Mém. Acad. Berl. 1804 (1807) 36 = Catoblastus praemorsus.
regia, *H. B. & K. Nov. Gen. et Sp.* i. 305.—Cuba; Panama.
Sancona, *H. B. & K. l. c.* 304.—N. Granat.
Sarcona, Steud. Nom. ed. II. ii. 226 = praec.
ventricosa, Hort. ex H. Wendl. in Kerch. Palm. 252 = Gaussia Ghiesbreghti.
ventricosa, *H. Wendl. ex Linden, Illustr. Hortic.* xxviii. (1881), 32, nomen.—Mexic.

OREOMUNNEA, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 33 = **Engelhardtia**, Leschen. (Jugland.).
pterocarpa, Oerst. l. c. 34 = E. Oreomunnea.

OREOMYRRHIS, Endl. Gen. Pl. 787 (1839). *UMBELLIFERAE*, Benth. & Hook. f. i. 897.
CALDASIA, Lag. Amen. Acad. Madr. ii. 98 (1821).
andicola, *Endl. Gen. Pl.* 787.—Reg. Andin.; Austral.; N. Zel.
andina, Hemsl. Biol. Centr. Am. Bot. i. 567 = praec.
argentea, Hook. f. in Hook. Ic. Pl. t. 300 = andicola.
brachycarpa, Hook. f. l. c. sub t. 300 = andicola.
ciliata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 471 [bis] = andicola.
Colensoi, Hook. f. Fl. N. Zel. i. 92 = andicola.
daucoides, *Urban, in Linnaea*, xliii. (1880–82) 303.—Peruv.
eriopoda, Hook. f. Fl. Tasm. i. 162 = andicola.
gracilipes, *Hemsl. Diagn. Pl. Nov.* i. 16.—Mexic.
Haastii, Hook. f. Handb. N. Zeal. Fl. 91 = andicola.
lasiopetala, *A. Gray, Bot. U. St. Expl. Exped.* i. 712.—Peruv.
planipetala, *Hemsl. Diagn. Pl. Nov.* i. 16.—Mexic.
pulvinifica, *F. Muell. Fragm.* viii. 185.—Austral.
ramosa, Hook. f. Handb. N. Zeal. Fl. 91 = andicola.
sessiliflora, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 47 [bis] = andicola.

OREOPANAX, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 107. *ARALIACEAE*, Benth. & Hook. f. i. 946.
MONOPANAX, Regel, Gartenfl. xviii. (1869) 35. t. 606.
acerifolium, *Seem. Journ. Bot.* iii. (1865) 275.—Am. austr.
Aleurites, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 108, nomen = Pavonii.
amplum, *Decne. & Planch. l. c.*, nomen.—Venezuela.
Andreanum, *March. in Bull. Soc. Bot. Belg.* xix. (1880) 90.—Ecuador.
angulare, *Seem. in Journ. Bot.* iii. (1865) 276.—Am. austr.
argentatum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
avicenniaefolium, *Decne. & Planch. l. c.*—Ecuador.
boliviense, *Seem. in Journ. Bot.* iii. (1865) 272.—Bolivia.
Bonplandianum, Planch. & Linden, ex Seem. l. c. 274 = Humboldtianum.
brachybotryum, Planch. & Linden, ex Seem. l. c. = seq.
brachystachyum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
Brownii, Witte, in Fl. Jard. Pays-bas, iv. (1861) 90; et ex Regel, Gartenfl. (1863) 146. t. 399 = Dendropanax japonicum.
brunneum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
bulbosum, Seem. in Journ. Bot. iii. (1865) 275 = seq.
bullosum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
capitatum, *Decne. & Planch. l. c.*—Am. austr.
catalpaefolium, Decne. & Planch. l. c. = capitatum.
Cecropia, *Decne. & Planch. l. c.*, nomen.—N. Granat.
cheirophyllum, *Seem. in Journ. Bot.* iii. (1865) 275.—Am. austr.
chrysoleucum, *Planch. & Linden, ex Seem. l. c.* 273.—Venezuela.
confusum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 85.—Ecuador.
cordatum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
coriaceum, *Decne. & Planch. l. c.* (1854), 321, nomen.—N. Granat.
costaricense, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 89.—Costa Rica.
crassinervium, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
cumanense, *Decne. & Planch. l. c.*—Venezuela.
dactylifolium, *Hort.—Cf. Gard. Chron.* (1864) 418.—Hab.?
Destructor, Seem. in Journ. Bot. vii. (1869) 351 = capitatum.
discolor, *Decne. & Planch. Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
divulsum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 90.—Peruv.
Dombeyannm, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Peruv.

OREOPANAX:—
Echinops, *Decne. & Planch. in Rev. Hortic.* Sér. iv. iii. (1854) 108.—Mexic.
ecuadoriense, *Seem. Journ. Bot.* iii. (1865) 272.—Ecuador.
elegans, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Hab.?
Epremesnilianum, *André, l. c.* (1882) 557.—Hab.?
farinosum, *Decne. & Planch. l. c.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
flabellatum, *Decne. & Planch. l. c.*, nomen.—Ecuador.
flaccidum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 84.—Mexic.
floribundum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
fraternum, *Decne. & Planch. l. c.*, nomen.—Venezuela.
fulvum, *March. in Mart. Fl. Bras.* xi. 1. 254.—Bras.
geminatum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 94.—Nicaragua.
guatemalense, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Guatemala.
Horquetero, *Decne. & Planch. l. c.*, nomen.—Venezuela.
Humboldtiana, *Decne. & Planch. l. c.*, nomen.—Venezuela.
hypargyreum, *Decne. & Planch. l. c.*, nomen (hyphargyrum).—N. Granat.
hypoleucum, Decne. & Planch. l. c., nomen = praec.
ilicifolium, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 82.—Bolivia.
incisum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Hab.?
jatrophaefolium, *Decne. & Planch. l. c.*—Mexic.
laciniosum, *Decne. & Planch. l. c.*, nomen.—N. Granat.
lancifolium, *Planch. & Linden, ex Seem. in Journ. Bot.* iii. (1865) 273.—Venezuela.
lanigerum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
Lechleri, *Seem. in Journ. Bot.* iii. (1865) 271.—Peruv.
Liebmannii, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 87.—Mexic.
Lindenii, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
Macleani, *Seem. in Journ. Bot.* iii. (1865) 272.—Peruv.
macrocephalum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen; *et ex Wedd. Chlor. And.* ii. 207.—Bolivia.
macrophyllum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Hab.?
Mathewsii, Seem. in Journ. Bot. iii. (1865) 272 = Pavonii.
meridense, *Planch. & Linden, ex Seem. l. c.* 275, nomen.—Venezuela.
Mutisianum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
myriocarpum, *Decne. & Planch. l. c.*, nomen.—N. Granat.
obtusilobum, *Decne. & Planch. l. c.*—Ecuador.
ocanense, *Decne. & Planch. l. c.*, nomen.—N. Granat.
Oerstedianum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 85.—Costa Rica.
oxyodon, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Mexic.
pachycephalum, *Planch. & Linden, ex Seem. in Journ. Bot.* iii. (1865) 275.—N. Granat.
Pavonii, *Seem. l. c.* 272.—Peruv.
peltatum, *Linden, Cat.* n. 14 (1859) 8; *et ex Regel, Gartenfl.* (1862) 170.—Mexic.
Pentlandiana, *Decne. & Planch. ex Wedd. Chlor. And.* ii. 207.—Bolivia.
permixtum, *March. in Mart. Fl. Bras.* xi. 1. 254.—Bras.
platanifolium, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—Reg. Andin.
platyphyllum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 88.—Mexic.
Pseudo-Jatropha, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
Pseudo-Platanus, *Decne. & Planch. l. c.*, nomen.—Hab.?
reticulatum, *Decne. & Planch. l. c.*, nomen; *et ex Seem. in Journ. Bot.* iii. (1865) 275.—Am. austr.
Ruizii, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—Mexic.
Salvinii, *Hemsl. Diagn. Pl. Nov.* 16.—Guatemala.
sclerophyllum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108.—N. Granat.
Seemannianum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 80.—Ecuador.
septemnervium, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 108 = capitatum.

OREOPANAX:—
Serra, *André, in Rev. Hortic.* (1884) 321.—Ecuador.
sessiliflorum, *Decne. & Planch. l. c.*—Peruv.
Sprucei, *Seem. in Journ. Bot.* iii. (1865) 272.—Ecuador.
Thibautii, *Hook. f. Bot. Mag.* t. 6340.—Mexic.
Trianae, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 108, nomen.—N. Granat.
turbacense, *Decne. & Planch. l. c.* 108.—N. Granat.
xalapense, *Decne. & Planch. l. c.*—Mexic.

OREOPHEA, Auct. ex Steud. Nom. ed. II. ii. 226 (1841) = **Orophea**, Blume (Anonac.).

OREOPHILA, D. Don, in Trans. Linn. Soc. xvi. (1830) 178; et in Phil. Mag. xi. (1832) 388 = **Hypochoeris**, Linn. (Compos.).
apargioides, D. Don, in Phil. Mag. xi. (1832) 388 = H. apargioides.
Meyeniana, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 292 = H. Meyeniana.
odorata, Poepp. & Endl. Nov. Gen. et Sp. iii. 57. t. 263 = H. odorata.
picroides, D. Don, in Phil. Mag. xi. (1832) 388 = H. apargioides.
sessiliflora, D. Don, in Trans. Linn. Soc. xvi. (1830) 178 = H. sessiliflora.
sibirica, C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 95 = H. grandiflora.
taraxacifolia, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 291.—Peruv.
taraxacoides, D. Don, in Phil. Mag. xi. (1832) 388 = H. Scorzonerae.
taraxacoides, Walp. Rep. vi. 336, in syn. = *taraxacifolia*.
tenuifolia, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 31 = H. tenuifolia.

OREOPHILA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 258 (1840) = **Pachystima**, Rafin. (Celastr.).
myrtifolia, Nutt. ex Torr. & Gray, l. c. 259 = P. Myrsinites.
myrtifolia, Scheele, Roem. Texas, 432; ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 158 = Ilex Cassine.

OREOPOLUS, Schlecht. in Linnaea, xxviii. (1856) 493. *RUBIACEAE*, Benth. & Hook. f. ii. 97.
citrinus, *Schlecht. l. c.*—Reg. Andin.

OREORCHIS, Lindl. in Journ. Linn. Soc. iii. (1859) 26. *ORCHIDEAE*, Benth. & Hook. f. iii. 498.
foliosa, *Lindl. l. c.* 27.—Reg. Himal.
gracilis, *Franch. & Sav. Enum. Pl. Jap.* ii. 27.—Japon.
lancifolia, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 410.—Japon.
micrantha, *Lindl. in Journ. Linn. Soc.* iii. (1859) 27.—Reg. Himal.
patens, *Lindl. l. c.*—Japon.

OREOSCIADIUM, Wedd. Chlor. And. ii. 203 (1857). *UMBELLIFERAE*, Benth. & Hook. f. i. 889.
NIPHOGETON, Schlecht. in Linnaea, xxviii. (1856) 481, nomen prius.
acaule, *A. Gray, in Proc. Am. Acad.* vii. (1868) 343.—Am. bor. occ.
dissectum, *Wedd. Chlor. And.* ii. 204.—N. Granat.
Lingula, *Wedd. l. c.* 205.—N. Granat.
montanum, *Wedd. l. c.* 204.—N. Granat.

OREOSELINUM, [Tourn.] Adans. Fam. ii. 100 (1763); Bieb. Fl. Taur. Cauc. iii. 209 (1819) = **Peucedanum**, Tourn. (Umbellif.).
album, Hoffm. Gen. Umb. ed. I. 155 = Silaus longifolius.
austriacum, Hoffm. l. c. = P. austriacum.
capense, Eckl. & Zeyh. Enum. 350 = P. capense.
caucasicum, Bieb. Fl. Taur. Cauc. iii. 209 = P. caucasicum.
Chabraei, Bieb. l. c. = P. carvifolium.
dahuricum, Bess. ex Steud. Nom. ed. II. ii. 226 = P. terebinthaceum.
Elaphoboscum, Delarb. Fl. Auv. ed. II. i. 429 = P. Cervaria.
elegans, Hoffm. Gen. Umb. ed. I. 155 = P. austriacum.
glaucum, Presl, Bot. Bemerk. 75 (= *Bubon montanus, hypoleucus*).—Afr. austr.
humile, Bess. ex Steud. Nom. ed. II. ii. 226 = P. salinum.

OREOSELINUM:—
latifolium, Bieb. Fl. Taur. Cauc. iii. 211 = Peucedanum latifolium.
legitimum, Bieb. l. c. 210 = P. Oreoselinum.
montanum, Hoffm. Gen. Umb. ed. I. 155 = P. austriacum.
nigricans, Hoffm. l. c. = P. austriacum.
nigrum, Delarb. Fl. Auv. ed. II. i. 428 = P. Oreoselinum.
peucedanoides, Hoffm. Gen. Umb. ed. I. 155 = P. involucratum.
podolicum, Bess. Enum. Pl. Volh. 12 = P. Chabraei.
Pseudo-carvifolium, Hoffm. Gen. Umb. ed. I. 155 = Selinum Carvifolia.
pyrenaeum, Hoffm. l. c. = Selinum pyrenaeum.
rablense, Hoffm. l. c. = P. austriacum.
uliginosum, Eckl. & Zeyh. Enum. 350 = P. uliginosum.

OREOSERIS, DC. Prod. vii. 17 (1838) = **Gerbera**, Gronov. (Compos.).
lanuginosa, DC. l. c. = G. lanuginosa.
nivea, DC. l. c. 18 = G. nivea.
ovalifolia, Wall. Cat. n. 2930 = G. piloselloides.
pusilla, DC. Prod. vii. 17 = G. lanuginosa.

OREOSOLEN, Hook. f. Fl. Brit. Ind. iv. 318 (1884). *SCROPHULARINEAE.*
Wattii, *Hook. f. l. c.* 319.—Reg. Himal.

OREOSPHACUS, Phil. in Leyb. Escurs. a las Pampas, 45 (1873). *LABIATAE*, Benth. & Hook. f. ii. 1184.
parvifolia, *Phil. l. c.*—Chili.

OREOSPLENIUM, Zahlb. ex Endl. Gen. 814, 1446 (1841) = **Zahlbrucknera**, Reichb. (Saxif.).

OREOSTYLIDIUM Berggr. in Minneskr. Fisiog. Sällsk. Lund (1877), n. VIII. 1. *STYLIDEAE.*
subulatum, *Berggr. l. c.* 2.—N. Zel.

OREOSYCE, Hook. f. in Oliver, Fl. Trop. Afr. ii. 548 (1871). *CUCURBITACEAE*
africana, *Hook. f. l. c.*—Afr. trop.

OREOTRYS, Rafin. Atl. Journ. 145 (1832); Fl. Tellur. ii. 74 (1836) = **Heuchera**, Linn. (Saxifr.).
bracteata, Rafin. Atl. Journ. 145 = H. bracteata.

OREOXIS, Rafin. in Ser. Bull. Bot. i. (1830) 217 = **Cymopterus**, Rafin. (Umbell.).
humilis, Rafin. l. c. = C. alpinus.

ORESCIA, Reinw. in Syll. Ratisb. ii. (1828) 15. Gen. ignot.; an *PRIMULACEAE*?
montana, Reinw. l. c. (Quid?).—Java.

ORESIGONIA, Schlecht. ex Less. Syn. Comp. 393 (1832) = **Culcitium**, H. B. & K. (Compos.).

ORESIGONIA, Willd. ex Less. Syn. Comp. 393 (1832); DC. Prod. vi. 323 (1837) = **Werneria**, H. B. & K. (Compos.).
brevifolia, Willd. ex Wedd. Chlor. And. i. 82 = W. humilis.
latifolia, Willd. ex Less. Syn. Comp. 393 = W. disticha.
pycnophylla, Bonpl. ex Wedd. Chlor. And. i. 83 = W. rigida.

ORESITROPHE, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 106 (Enum. Pl. Chin. Bor. 31). *SAXIFRAGEAE*, Benth. & Hook. f. i. 639.
rupifraga, *Bunge, l. c.*—China.

ORESTION, Kunze, ex Berg, in Linnaea, xxvii. (1854) 394 = **Myrtus**, Tourn.
vaccinioides, Kunze, ex Berg, l. c. = M. bullata.

ORESTION, Rafin. Fl. Tellur. ii. 48 (1836) = **Olearia**, Moench (Compos.).
dentata, Rafin. l. c. = O. dentata.

OREXIS, Salisb. Gen. Pl. Fragm. 117 (1866) = **Lycoris**, Herb. (Amaryll.).
radiata, Salisb. l. c. 118 = L. radiata.

ORFILEA, Baill. Etud. Gén. Euph. 452 (1858) = **Alchornea**, Sw. (Euphorb.).
coriacea, Baill. l. c. 453 = A. coriacea.

ORIASTRUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 50. t. 257 (1842) = **Chaetanthera**, Ruiz & Pav. (Compos.).
chilense, Wedd. Chlor. And. i. 30.—Chili.
cochlearifolium, A. Gray, in Proc. Am. Acad. v. (1861) 144.—Peruv.
gnaphalioides, Wedd. Chlor. And. i. 30.—Chili.
lycopodioides, Wedd. l. c.—Chili.
pulvinatum, Phil. in Linnaea, xxxiii. (1864–65) 112.—Chili.
pusillum, Poepp. & Endl. Nov. Gen. et Sp. iii. 50.—Chili.

ORIBA, Adans. Fam. ii. 459 (1763) = **Anemone**, Tourn. (Ranunc.).
sylvestris, Opiz, Seznam, 70 = A. sylvestris.

ORIBASIA, [Moç. & Sesse,] ex DC. Prod. vi. 323 (1837) = **Werneria**, H. B. & K. (Compos.).
acaulis, [Moç. Sesse & Cerv.] ex DC. l. c. 324 = W. Mocciniana.

ORIBASIA, Schreb. Gen. 123 (1789) = **Palicourea**, Aubl. (Rubiac.).
longiflora, J. F. Gmel. Syst. 367 = Pal. longiflora.
lutea, J. F. Gmel. l. c. (= *Nonatelia lutea*).—Guian.
officinalis, J. F. Gmel. l. c. = Psychotria officinalis.
paniculata, J. F. Gmel. l. c. = Psychotria flexuosa.
racemosa, J. F. Gmel. l. c. = Psychotria racemosa.
violacea, J. F. Gmel. l. c. = Pal. violacea.

ORIGANON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Origanum**, Tourn. (Labiat.).
majoranum, St. Lag. l. c. = Origanum Majorana.
onitum, St. Lag. l. c. = Origanum Onites.

ORIGANUM, Tourn. ex Linn. Syst. ed. I (1735).
LABIATAE, Benth. & Hook. i. ii. 1185.
AMARACUS, Gleditsch, Syst. 189 (1764).
BELTOKON, Rafin. Fl. Tellur. iii. 86 (1836).
MAJORANA, Rupp. Fl. Jen. ed. Hall. 235 (1745).
ONITES, Rafin. Fl. Tellur. iii. 86 (1836).
OROGA, Rafin. l. c. (1836).
SCHIZOCALYX, Scheele, in Flora, xxvi. (1843) 575.
ZATARENDIA, Rafin. Fl. Tellur. iii. 86 (1836).
acinacifolium, Steud. Nom. ed. II. ii. 226 = majoricum.
acinifolium, Wallr. ex Benth. Lab. Gen. et Sp. 728 = majoricum.
aegyptiacum, Linn. Sp. Pl. 588 = O. Maru.
albiflorum, C. Koch, in Linnaea, xxi. (1848) 662 = vulgare.
album, Salisb. Prod. 85 = O. Onites.
anglicum, Hill, Veg. Syst. xvii. 35 = vulgare.
angustifolium, C. Koch, in Linnaea, xxi. (1848) 661 = vulgare.
balearicum, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1863) 5 = majoricum.
bengalense, Heyne, ex Wall. Cat. n. 1532 = Pogostemum Patchouly.
benghalense, Burm. f. Fl. Ind. 128 = Pogostemon plectranthoides.
calcaratum, Juss. Gen. 115 = Tournefortii.
capitatum, Willd. ex Benth. in Linnaea, xi. (1837) 339 = vulgare.
ciliatum, *Boiss. & Kotschy, Diagn.* Ser. II. iv. 10.—As. Min.
ciliatum, Willd. Sp. Pl. iii. 133 (Acanthacea).—Afr. trop.
cinereum, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 579.—Afr. bor.
clinopodioides, Walt. Fl. Carol. 165 = Pycnanthemum aristatum.
compactum, *Benth. Lab. Gen. et Sp.* 334.—Afr. bor.
confertum, Savi, Osserv. Origan. 12 = O. Majorana.
cordifolium, *Vog. in Linnaea*, xv. (1841) 76.—Syria; Ins. Cyprus.
creticum, Linn. Sp. Pl. 589 = vulgare.
creticum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 610 = compactum.
creticum, Sieber, ex Benth. Lab. Gen. & Sp. 336 = hirtum.
decipiens, Wallr. ex Benth. l. c. 728 = vulgare.
dictamnifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = seq.
Dictamnus, *Linn. Sp. Pl.* 589.—Ins. Creta.
dubium, *Boiss. Fl. Orient.* iv. 553.—As. Min.
Ehrenbergii, *Boiss. l. c.* 551.—Syria.

ORIGANUM:—
elegans, Steph. ex Steud. Nom. ed. II. ii. 227 = Dracocephalum pinnatum.
flexuosum, Walt. Fl. Carol. 165 = Pycnanthemum linifolium.
floribundum, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 286.—Alger.
floridum, Salisb. Prod. 85 = vulgare.
fortuitum, Savi, Osserv. Origan. 14 = paniculatum.
glandulosum, Desf. Fl. Atlant. ii. 27 = hirtum.
glandulosum, Salzm. ex Benth. Lab. Gen. et Sp. 335 = compactum.
gracile, C. Koch, in Linnaea, xxi. (1848) 661 = vulgare.
Haussknechtii, *Boiss. Fl. Orient.* iv. 550.—Cataonia.
heracleoticum, Benth. Lab. Gen. et Sp. 336 = hirtum.
heracleoticum, Koch, in Mert. & Koch, Deutschl. Fl. iv. 305 = O. Orega.
heracleoticum, *Linn. Sp. Pl.* 589.—Reg. Mediterr.
heracleoticum, Reichb. Fl. Germ. Excurs. 313 = vulgare.
hirtum, *Link, Enum. Hort. Berol.* ii. 114.—Europ.
hirtum, Noé, ex Nym. Consp. 592 = vulgare.
humile, Mill. Gard. Dict. ed. VIII. n. 4 = vulgare.
hybridum, Mill. l. c. n. 12 = sipyleum.
illyricum, Scheele, in Flora, xxvi. (1843) 574 = hirtum.
incanum, Walt. Fl. Carol. 165 = Pycnanthemum aristatum.
indicum, Roth, Nov. Pl. Sp. 265 = Pogostemum plectranthoides, P. Patchouly.
laevigatum, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 252.—As. Min.; Syria.
lanatum, Boj. ex Nees, in DC. Prod. xi. 721 = Nelsonia canescens.
latifolium, Mill. Gard. Dict. ed. VIII. n. 3 = vulgare.
latifolium, Scheele, in Flora, xxvi. (1843) 574 = hirtum.
laxiflorum, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 376 = vulgare.
leptocladum, *Boiss. Fl. Orient.* iv. 549.—Cilicia.
libanoticum, *Boiss. Diagn.* Ser. I. v. 14.—Syria.
Loureiri, *Kostel. Allg. Med.-Pharm. Fl.* iii. 768.—Cochinch.
lusitanicum, Rouy, in La Natural. ii. (1882) (Matér. Rév. Fl. Portug. 36, in syn.) = majoricum.
lusitanicum, *Rouy, l. c.* 37.—Ins. Balear.
macrostachyum, Hoffmgg. & Link, Fl. Port. i. 120 = vulgare.
Majorana, *Linn. Sp. Pl.* 590.—Europ.
majoranoides, Willd. Sp. Pl. iii. 137 = O. Majorana.
majoricum, *Cambess. in Mém. Mus. Par.* xiv. (1827) 296.—Ins. Balear.
Maru, *Linn. Sp. Pl.* 825.—Syria; Arabia.
Maru, Sibth. & Sm. Fl. Graec. vi. 59. t. 573 = microphyllum.
megastachyum, Link, Enum. Hort. Berol. ii. 114 = vulgare.
micranthum, Hort. ex Colla, Hort. Ripul. App. iv. 53 = vulgare?
micranthum, *Vog. in Linnaea*, xv. (1841) 77.—As. Min.
microphyllum, *Sieber, ex Benth. Lab. Gen. et Sp.* 338.—Ins. Creta.
neglectum, Vog. in Linnaea, xv. (1841) 81 = hirtum.
nervosum, Vog. l. c. 78 = O. Maru.
normale, D. Don, Prod. Fl. Nep. 113 = vulgare.
nutans, Willd. ex Benth. in Linnaea, xi. (1837) 339 = vulgare.
oblongatum, Link, Enum. Hort. Berol. ii. 114 = vulgare.
odorum, Salisb. Prod. 85 = O. Majorana.
officinale, Gueldenst. Reisen, ii. 287 = vulgare.
Onites, Lam. Encyc. iv. 608 = O. Majorana.
Onites, *Linn. Sp. Pl.* 590.—Europ. austr. or.; As. Min.; Syria.
Orega, *Vog. in Linnaea*, xv. (1841) 78.—Hab.?
orientale, Mill. Gard. Dict. ed. VIII. n. 5 = vulgare.
pallidum, Desf. Tabl. ed. I. 62, nomen; Cat. Hort. Par. ed. III. 395 = O. Onites.
paniculatum, *Koch, in Mert. & Koch, Deutschl. Fl.* iv. 306.—Europ.
parviflorum, Urv. in Mém. Soc. Linn. Par. i. (1822) 327 = hirtum.
pruinosum, C. Koch, in Linnaea, xxi. (1848) 663 = vulgare.
pulchellum, *Boiss. Diagn.* Ser. II. iv. 9.—Oriens.
pulchrum, *Boiss. & Heldr. l. c.* 11.—Graecia.
punctatum, Poir. Encyc. Suppl. iv. 186 = Pycnanthemum incanum.
purpurescens, Gilib. Fl. Lituan. i. 64 = vulgare.
rotundifolium, *Boiss. Diagn.* Ser. II. iv. 11.—Armenia.
salvifolium, *Roth, Catalect.* fasc. ii. 48.—Hab.?

ORIGANUM:—
sardoum, Nym. Consp. 592 = hirtum.
saxatile, Salisb. Prod. 85 = O. Dictamnus.
scabrum, *Boiss. & Heldr. Diagn.* Ser. I. vii. 48.—Graecia.
serpylliforme, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 59.—Hab.?
siculum, Nym. Consp. 592 = virens.
sipyleum, *Linn. Sp. Pl.* 589.—As. Min.
smyrnaeum, Linn. l. c. = O. Onites.
smyrnaeum, Sibth. & Sm. Fl. Graec. vi. 57. t. 571 = hirtum.
stoloniferum, Bess. ex Reichb. Fl. Germ. Excurs. 313 = vulgare.
suffruticosum, Hort. ex Steud. Nom. ed. I. 572 = O. Majorana.
syriacum, Linn. Sp. Pl. 590 = O. Maru?
thymiflorum, Reichb. Fl. Germ. Excurs. 313 = vulgare.
Tournefortii, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 311.—Archip. Graec.
Tragoriganum, *Zucc. ex Steud. Nom.* ed. II. ii. 227, nomen (Quid?).—Italia?
turbinatum, *Vog. in Linnaea*, xv. (1841) 77.—Hab.?
venosum, Willd. ex Benth. l. c. xi. (1837) 239 = vulgare.
vestitum, Clarke, Trav. ii. 451 = O. Maru.
virens, *Hoffmgg. & Link, Fl. Port.* i. 119.—Europ.
virens, C. A. Mey. Verz. Pfl. Cauc. 90 = vulgare.
virescens, Poir. Encyc. Suppl. iv. 186 = virens.
viride, Willd. ex Benth. in Linnaea, xi. (1837) 339 = vulgare.
viridulum, Martr. Fl. Tarn, 551 = vulgare.
vulgare, *Linn. Sp. Pl.* 590.—Europ.; As. et Afr. bor.
vulgare, O. F. Muell. in Fl. Dan. t. 638 = Mentha aquatica.
Wallichianum, Benth. in Wall. Cat. n. 1565 = O. Majorana.
watariense, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 627 = Anisochilus carnosus.
Watsoni, Schlagint. ex T. A. Schmidt, in Journ. Bot. vi. (1868) 227, 234 = vulgare.

ORIMARIA, Rafin. in Ser. Bull. Bot. i. (1830) 218.
UMBELLIF. gen. dub.
filiformis, Rafin. l. c. (Quid?).—Am. bor.

ORIOPHORUM, Gunn. Fl. Norveg. ii. 84 (1772), err. typ. = **Eriophorum**, Linn. (Cyper.).

ORITES, Banks & Soland. ex Hook. f. Fl. Antarct. 282 (1847) = **Donatia**, Forst. (Saxifrag.).
depressa, Banks & Soland. l. c. = D. fascicularis.

ORITES, R. Br. in Trans. Linn. Soc. x. (1810) 189.
PROTEACEAE, Benth. & Hook. f. iii. 180.
AMPHIDERRIS, Spach, Hist. Vég. Phan. x. 402 (1841).
ORITINA, R. Br. in Trans. Linn. Soc. x. (1810) 224.
PATAGUA, Poepp. ex Baill. Adansonia, x. (1871) 49.
TROPOCARPA, D. Don, ex Meissn. in DC. Prod. xiv. 423 (1857).
acicularis, *Roem. & Schult. Syst.* iii. 427; *R. Br. Prot. Nov.* 32.—Austral.
diversifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 190.—Tasman.
excelsa, *R. Br. Prot. Nov.* 32.—Austral.
lancifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 108.—Austral.
Milligani, *Meissn. in Hook. Kew Journ.* iv. (1852) 209, nomen; *et in DC. Prod.* xiv. 424.—Tasman.
myrtoides, *Benth. & Hook. f. Gen.* iii. 180.—Chili.
revoluta, *R. Br. in Trans. Linn. Soc.* x. (1810) 190.—Tasman.

ORITHALIA, Blume, Fl. Jav. Praef. 6 (1828) = **Agalmyla**, Blume (Gesner.).

ORITHYIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 336 (1836) = **Tulipa**, Tourn. (Liliac.).
biflora, Kunth, Enum. Pl. iv. 227 = T. biflora.
dasystemon, Regel, in Act. Hort. Petrop. v. (1877) 261 = T. dasystemon.
edulis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 158 = T. edulis.
heterophylla, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) i. 440 = T. heterophylla.
nutans, Trautv. Pl. Imag. 15. t. 10 = T. uniflora.

ORITHYIA :—
oxypetala, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. sub t. 336 = T. uniflora.
oxypetala, A. Gray, in Perry, Jap. Exped. ii. 322 = Tulipa edulis.
uniflora, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 336 = T. uniflora.

ORITINA, R. Br. in Trans. Linn. Soc. x. (1810) 224 = **Orites**, R. Br. (Proteac.).
acicularis, R. Br. l. c. = Orites acicularis.

ORIUM, Desv. Journ. Bot. iii. (1814) 161. t. 25. f. 10 = **Clypeola**, Linn. (Crucif.).
lanuginosum, Desv. l. c. 162 = C. eriocarpa.

ORIXA, Thunb. Nov. Gen. 56 (1783) = **Celastrus**, Linn.
japonica, Thunb. l. c. 57 = C. Orixa.
ternata, Blanco, Fl. Filip. ed. I. 62 = Evodia robusta.

ORIZA, Franch. & Sav. Enum. Pl. Jap. ii. 155 (1879), sphalm. = **Oryza**, Tourn. (Gramin.).

ORLAYA, Hoffm. Gen. Umb. ed. I. 58 (1814) = **Daucus**, Tourn.
anisopoda, Boiss. Diagn. Ser. I. x. 46 = D. littoralis.
Bubania, Philippe, Fl. Pyren. i. 407 = D. pumilus.
cretica, Nym. Consp. 278 = D. pumilus.
grandiflora, Hoffm. Gen. Umb. ed. I. 58 = D. grandiflorus, *Scop.*
intermedia, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 49 = D. muricatus.
maritima, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 79 = D. pumilus.
platycarpos, Koch, l. c. 79 = D. muricatus.

ORLEANISIA, Rodrig. Orch. Nov. i. 62 (1877). *ORCHIDEAE.*
amazonica, *Rodrig. l. c.* 64.—Bras.

ORLOWIA, Gueldenst. ex Georgi, Beschr. Russ. Reich. III. v. 1089 (1800) = **Phlomis**, Linn. (Labiat.).
rossica, Gueldenst. l. c. = P. tuberosa.

ORMENIS, Cass. in Dict. Sc. Nat. xxix. 180 (1823) = **Anthemis**, Mich. (Compos.).
aurea, Coss. & Dur. Expl. Alg. t. 61 = A. aurea.
aurea, Lowe, Man. Fl. Mader. i. 467 = A. nobilis.
bicolor, Cass. in Dict. Sc. Nat. xxxvi. 355 = A. mixta.
coronopifolia, Less. Syn. Comp. 249 = A. mixta.
eriolepis, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61, nomen.—Marocc.
fuscata, Sch. Bip. ex Nym. Consp. 363 = A. fuscata.
heterophylla, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61, nomen = A. heterophylla.
mixta, Dum. Fl. Belg. 69; DC. Prod. vi. 18 = A. mixta.
nobilis, J. Gay, ex Coss. & Germ. Fl. Par. ed. I. 397 = A. nobilis.

ORMIASTIS, Rafin. Fl. Tellur. iii. 93 (1836) = **Salvia**, Tourn. (Labiat.).
pinnata, Rafin. l. c. = S. pinnata.

ORMILIS, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
Horminum, Rafin. l. c. = S. Horminum.
viridis, Rafin. l. c. = S. viridis.

ORMISCUS, Eckl. & Zeyh. Enum. 9 (1835) = **Heliophila**, Burm. f. (Crucif.).
amplexicaulis, Eckl. & Zeyh. l. c. = H. amplexicaulis.
pendulus, Spach, Hist. Vég. Phan. vi. 453 = H. pendula.
pinnatus, Eckl. & Zeyh. Enum. 10 = H. pendula.
pusillus, Eckl. & Zeyh. l. c. 9 = H. pusilla.
tenuisiliqua, Eckl. & Zeyh. l. c. = H. pusilla.

ORMOCARPUM, Beauv. Fl. Owar. i. 95. t. 58 (1805). *LEGUMINOSAE*, Benth. & Hook. f. i. 515.
ACROTAPHROS, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 207. t. 38 (1847).
DIPHACA, Lour. Fl. Cochinch. 453 (1790).
HORMOCARPUS, Spreng. Gen. ii. 594 (1831).
RATHKEA, Schum. & Thonn. Beskr. Guin. Pl. 365 (1828).
RUSSELLIA, Koen. ex Roxb. Fl. Ind. iii. 364 (1832).

ORMOCARPUM :—
aristatum, Desv. in Mém. Soc. Linn. Par. iv. (1826) 308 = Pictetia aristata.
australe, Desv. l. c. 307 = Desmodium umbellatum.
bibracteatum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 143. —Afr. trop.
caeruleum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 511.—Ins. Socotra.
cassioides, Desv. Journ. Bot. i. (1813) 122. t. 5. f. 16 = sennoides.
coccinea, *G. Don, Gen. Syst.* ii. 279.—Mexic.
coronilloides, G. Don, l. c. = sennoides.
discolor, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 223.—Afr. trop.
elegans, *G. Don, Gen. Syst.* ii. 279.—Mexic.
glabrum, *Teijsm. & Binn. in Tijdschr. Nederl. Indie*, xxvii. (1864) 56.—Ins. Celebes.
intermedium, *Teijsm. & Binn. l. c.* 55.—Java.
Kirkii, *S. Moore, in Journ. Bot.* xv. (1877) 290.—Afr. trop.
mimosoides, *S. Moore, l. c.* 291.—Afr. trop.
oblongum, Desv. in Mém. Soc. Linn. Par. iv. (1825) 307 = Desmodium umbellatum.
ochroleucum, *Zoll. & Mor. Syst. Verz. Zoll.* 6.—Java.
pluriflorum, Vatke, in Oestr. Bot. Zeitschr. xxviii. (1878) 215 = Kirkii.
rugosum, *Desv. Journ. Bot.* i. (1813) 122, nomen.—Hab.?
sennoides, *DC. Prod.* ii. 315.—As. et Afr. trop.
singapurianum, *Wall. Cat.* n. 5659.—Singapore.
striatum, Desv. in Mém. Soc. Linn. Par. iv. (1826) 307, sphalm. = sulcatum.
suberosum, *Teijsm. & Binn. in Tijdschr. Nederl. Indie*, xxvii. (1864) 55.—Java.
sulcatum, *Beauv. Fl. Owar.* i. 96.—Ins. S. Doming.
Thurberi, *Benth. & Hook. Gen.* i. 515.—Mexic.
umbellatum, Desv. in Mém. Soc. Linn. Par. iv. (1826) 307 = Desmodium umbellatum.
verrucosum, *Beauv. Fl. Owar.* i. 96. t. 58.—Afr. trop.

ORMOSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 95. *UMBELLIFERAE*, Benth. & Hook. f. i. 925.
Aucheri, *Boiss. l. c.*—As. Min.

ORMOSIA, Jacks. in Trans. Linn. Soc. x. (1811) 360. t. 25. *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
LAVIA, Hook. & Arn. Bot. Beech. Voy. 183. t. 38 (1833).
MACROTROPIS, DC. Prod. ii. 98 (1825).
TOULICHIBA, Adans. Fam. ii. 326 (1763).
acuminata, R. Grah. in Wall. Cat. n. 5973 = Sophora acuminata.
acuta, Vog. in Linnaea, xi. (1837) 405 = dasycarpa.
angolensis, Baker, in Oliver, Fl. Trop. Afr. ii. 255.—Afr. trop.
calavensis, *Azaola, in Blanco, Fl. Filip.* ed. II. 230.—Ins. Philipp.
coarctata, *Jacks. in Trans. Linn. Soc.* x. (1811) 363. t. 27.—Guian.
coccinea, *Jacks. l. c.* 360. t. 25.—Am. trop.
dasycarpa, *Jacks. l. c.* 362. t. 26.—Ind. occ.
discolor, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. 1. 318.—Bras.
emarginata, *Benth. in Hook. Kew Journ.* iv. (1852) 77. —Ins. Hongkong.
excelsa, *Benth. in Mart. Fl. Bras.* xv. 1. 318.—Bras.
fastigiata, *Tul. in Arch. Mus. Par.* iv. (1844) 108.—Am. trop.
floribunda, Wall. Cat. n. 5337 = robusta.
glauca, *Wall. Pl. As. Rar.* ii. 23. t. 125.—Reg. Himal.
laxiflora, *Benth. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 255.—Afr. trop.
macrodisca, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 253. —Malacca.
macrophylla, *Benth. in Ann. Wien. Mus.* ii. (1838) 88. —Bras.
microsperma, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 253. —Malacca.
minor, Vog. in Linnaea, xi. (1837) 405 = dasycarpa.
nitida, *Vog. l. c.* 405.—Bras.
nobilis, *Tul. in Arch. Mus. Par.* iv. (1844) 106.—Bras.
pachycarpa, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 76.—Ins. Hongkong.

ORMOSIA :—
pacimonensis, Spruce, ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 119 = Andira Amazonum.
panamensis, *Benth. in Seem. Bot. Voy. Herald*, 111.—Panama.
parvifolia, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 253.—Malacca.
robusta, *Baker, l. c.* 252.—Reg. Himal.; Burma.
semicastrata, *Hance, in Journ. Bot.* xx. (1882) 78.—China.
subsimplex, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. 1. 316.—Bras.
travancorica, *Bedd. Fl. Sylv.* i. t. 45.—Ind. or.
venosa, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 254.—Malacca.

ORMOSOLENIA, Tausch, in Flora, xvii. (1834) 348 = **Peucedanum**, Tourn. (Umbell.).
cretica, Tausch, l. c. = P. creticum.
pisidica, Boiss. & Heldr. Diagn. Ser. I. x. 34 = P. pisidicum.

ORMOSTEMA, Rafin. Fl. Tellur. iv. 38 (1836) = **Dendrobium**, Sw. (Orchid.).
albiflora, Rafin. l. c. = D. moniliforme.
purpurea, Rafin. l. c. = D. Linawianum.

ORMYCARPUS, Neck. Elem. iii. 82 (1790) = **Raphanus**, Tourn. (Crucif.).

ORNANTHES, Rafin. New Fl. Am. iii. 93 (1836) = **Fraxinus**, Tourn. (Oleac.).
americana, Rafin. l. c. = F. americana.
florida, Rafin. Alsog. Am. 40 = F. Ornus.
lutea, Rafin. l. c. = F. Ornus.
mannifera, Rafin. New Fl. Am. iii. 93 = F. rotundifolia.
parviflora, Rafin. Alsog. Am. 40 = F. Ornus?
rotundifolia, Rafin. New Fl. Am. iii. 93 = F. rotundifolia.

ORNITHARIUM, Lindl. & Paxt. Flow. Gard. i. (1850–51) 188. f. 117 = **Sarcochilus**, R. Br. (Orchid.).
striatulum, Lindl. & Paxt. l. c.—Ind. or.

ORNITHIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 293. *ORCHIDEAE*, Benth. & Hook. f. iii. 556.
SIAGONANTHUS, Poepp. & Endl. Nov. Gen. et Sp. i. 40. t. 69 (1835).
acaule, *Hoffmgg. Verz. Orch.—Cf. Bot. Zeit.* i. (1843) 834.—Bras.
aggregatum, *Reichb. f. in Bonplandia*, ii. (1854) 18.—N. Granat.
album, Hook. Bot. Mag. t. 3306 = Maxillaria Camaridii.
anceps, *Reichb. f. Beitr. Orch. Centr. Am.* 75.—Costa Rica.
aureum, *Poepp. & Endl. Nov. Gen. et Sp.* i. 57.—Peruv.
bicolor, Reichb. f. in Walp. Ann. vi. 489, sphalm. = Odontoglossum bicolor.
bracteatum, Wall. ex Lindl. Fol. Orch. Sunipia = Sunipia scariosa.
ceriferum, *Rodrig. Orch. Nov.* ii. 209.—Bras.
chloroleucum, *Rodrig. l. c.* 208.—Bras.
coccineum, *Salisb. in Trans. Hort. Soc.* i. (1812) 293. —Ind. occ.
confertum, *Griseb. Fl. Brit. W. Ind.* 626.—Ind. occ.
Cordyline, *Reichb. f. in Linnaea*, xli. (1877) 34.—Ecuador.
croceorubens, *Reichb. f. l. c.* 35.—Guian.? Ind. occ.?
densum, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Mexic.
distichum, *Lindl. in Benth. Pl. Hartw.* 153.—Peruv.
divaricatum, *Rodrig. Orch. Nov.* ii. 209.—Bras.
flavoviridium, *Rodrig. l. c.*—Bras.
fulgens, *Reichb. f. Beitr. Orch. Centr. Am.* 76.—Costa Rica.
giganteum, *Lindl. in Benth. Pl. Hartw.* 153.—Peruv.
histrionicum, *Reichb. f. in Bonplandia*, iv. (1856) 324. —Mexic.
inflexum, Reichb. f. in Walp. Ann. vi. 928 = Camaridium inflexum.
Jamesoni, *Reichb. f. in Linnaea*, xli. (1877) 35.—Ecuador.
Jenischianum, *Reichb. f. in Bonplandia*, ii. (1854) 18. —Venezuela.

ORNITHIDIUM:—
miniatum, *Lindl. Bot. Reg.* (1845) *Misc.* 62.—N. Granat.
multicaule, *Reichb. f. in Bonplandia*, ii. (1854) 18.—Peruv.
niveum, *Lindl. Orch. Linden.* 22.—Venezuela.
nubigenum, *Reichb. f. in Walp. Ann.* vi. 488.—N. Granat.
paleatum, *Reichb. f. in Linnaea*, xli. (1877) 36.—Am. centr.
parviflorum, *Reichb. f. in Bonplandia*, ii. (1854) 19.—Venezuela.
quitense, *Reichb. f. in Linnaea*, xli. (1877) 34.—Ecuador.
ruberrimum, *Reichb. f. in Walp. Ann.* vi. 489.—Venezuela.
sanguinolentum, *Lindl. Orch. Linden.* 22.—N. Granat.
semiscabrum, *Lindl. in Benth. Pl. Hartw.* 153.—Ecuador.
serrulatum, *Lindl. l. c.*—Ecuador.
Sophronitis, *Reichb. f. in Bonplandia*, ii. (1854) 18.—Venezuela.
squamatum, *Rodrig. Orch. Nov.* ii. 209.—Bras.
strumatum, *Endres & Reichb. f. in Gard. Chron.* (1874) II. 772.—Costa Rica.
Tafallae, *Reichb. f. in Bonplandia*, ii. (1854) 18.—Peruv.
vestitum, Reichb. f. in Walp. Ann. vi. 491 = parviflorum.

ORNITHOBOEA, Parish, ex C. B. Clarke, in DC. Monog. Phan. v. 147 (1883). *CYRTANDREAE.*
Parishii, *C. B. Clarke, l. c.* 148.—Burma.

ORNITHOCEPHALOCHLOA, Kurz, in Journ. Bot. xiii. (1875) 332. t. 171 = **Thuarea**, Pers. (Gramin.).
arenicola, Kurz, l. c. = T. sarmentosa.

ORNITHOCEPHALUS, Hook. Exot. Fl. ii. t. 127 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 568.
apiculatus, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 383.—Peruv.
Avicula, *Reichb. f. in Walp. Ann.* vi. 495.—Guian.
bicornis, *Lindl. in Benth. Bot. Voy. Sulph.* 172.—Panama.
Bonplandi, *Reichb. f. in Walp. Ann.* vi. 494.—Hab.?
chloroleucus, *Reichb. f. in Bonplandia*, iii. (1855) 226.—Bras.
ciliatus, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 383.—Guian.
Cruegeri, *Reichb. f. ex Griseb. Fl. Brit. W. Ind.* 635.—Ins. Trinit.
cujetifolia, *Rodrig. Orch. Nov.* i. 133.—Bras.
dolabratus, *Reichb. f. in Linnaea*, xli. (1877) 106.—Ecuador.
Elephas, *Reichb. f. in Walp. Ann.* vi. 493.—Guatemala.
falcatus, *Focke, in Tijdschr. Wetensch.* i. (1848) 211.—Bras.
gladiatus, *Hook. Exot. Fl.* ii. t. 127.—Ins. Trinit.
grandiflorus, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 383.—Bras.
Ibis, Reichb. f. in Walp. Ann. vi. 493 = gladiatus.
inflexus, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 384.—Mexic.
iridifolius, *Reichb. f. in Walp. Ann.* vi. 494.—Mexic.
Kruegeri, Reichb. f. l. c. 495 = Cruegeri.
mexicanus, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 24 = inflexus.
microphyllus, *Rodrig. Orch. Nov.* i. 134.—Bras.
myrticola, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 383.—Bras.
navicularis, Rodrig. Orch. Nov. i. 135 = Zygostates lunata.
Oberonia, *Reichb. f. in Gard. Chron.* (1869) 988.—Ins. Trinit.
planifolius, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 315.—Venezuela.
polyodon, *Reichb. f. in Linnaea*, xli. (1877) 33.—Ecuador.
Pseudobulbiferus, *Rodrig. Orch. Nov.* i. 134.—Bras.
pygmaeus, *Reichb. f. & Warm. in Otia Bot. Hamb.* 88.—Bras.
Salvinii, *Reichb. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 291.—Guatemala.
trichorhizus, *Batem. ex G. Don, in Sweet, Hort. Brit.* ed. III. 642.—Ins. Trinit.
tripollinicus, *Rodrig. Orch. Nov.* i. 136.—Bras.

ORNITHOCHILUS, Wall. ex Lindl. Gen. et Sp. Orch. 242 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 581.
eublepharon, Hance, in Journ. Bot. xxii. (1884) 364 = fuscus.
fuscus, *Wall. ex Lindl. Gen. et Sp. Orch.* 242.—Reg. Himal.; China.
Hillii, *Benth. in Journ. Linn. Soc.* xviii. (1881) 334.—Austral.
striatulus, Hort. Calc. ex Lindl. & Paxt. Flow. Gard. i. (1850–51) 188 (= Sarcochili sp.).—Hab.?

ORNITHOGALON, Rafin. Fl. Tellur. ii. 22 (1836) = seq.

ORNITHOGALUM, [Tourn.] Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 815.
ARDERNIA, ASPASIA, BERYLLIS, BRIZOPHILA, CATHISSA, EUSTACHYS, MYANTHE, OSMYNE, PHAEOCLES, TAENIOLA, UROPHYLLON, Salisb. Gen. Pl. Fragm. 33–35 (1866).
ALBUCEA, Reichb. Fl. Germ. Excurs. 109 (1830).
CARUELIA, Parl. Nouv. Gen. e Spec. 21 (1854).
CELSIA, Heist. Syst. 5, 20 (1748).
COILONOX, Rafin. Fl. Tellur. ii. 28 (1836).
ELIOKARMOS, Rafin. l. c. 24; iii. 58 (1836).
ETHESIA, Rafin. l. c. iii. 60 (1836).
FENELONIA, Rafin. Atl. Journ. 145 (1832).
HONORIUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 177 (1821).
IFUON, Rafin. Fl. Tellur. iv. 12 (1836).
LOMARESIS, Rafin. l. c. iii. 59 (1836).
LONCODILUS, Rafin. l. c. ii. 24 (1836).
LONCOMAELOS, Rafin. l. c. (1836).
LONCOXIS, Rafin. l. c. iii. 58 (1836).
MELOMPHIS, Rafin. l. c. ii. 21 (1836).
MYOGALUM, Link, Handb. i. 163 (1829).
NICIPE, Rafin. Fl. Tellur. iii. 54 (1836).
OZIROË, Rafin. l. c. 53 (1836).
PARTHENOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 158.
RAXAMARIS, Rafin. Fl. Tellur. iii. 54 (1836).
SIMIRA, Rafin. l. c. iv. 28 (1836).
TOMOXIS, Rafin. l. c. iii. 54 (1836).
TRIMELOPTER, Rafin. l. c. ii. 24 (1836).
TRITRIELA, Rafin. l. c. iii. 59 (1836).
abyssinicum, Hort. ex Schult. f. Syst. vii. 500 = Albuca abyssinica.
aciphyllum, *Baker, in Journ. Bot.* xii. (1874) 365.—Afr. austr.
acuminatum, Baker, in Saund. Refug. Bot. t. 177 = Eckloni, *Schlecht.*
acuminatum, Schur, Enum. Pl. Transs. 664 = orthophyllum.
acutipetalum, Herb. ex Schult. f. Syst. vii. 540 = Gagea minima.
Adalgisae, *Groves, in Nuov. Giorn. Bot.* ix. (1877) 71.—Italia.
aemulum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 161 = oligophyllum.
aequipetalum, Bert. in Merc. Chil. xvi. 788; ex Am. Journ. Sc. Ser. I. xxiii. (1833) 259 = biflorum.
affine, Bor. Fl. Centr. Fr. ed. III. ii. 625 = umbellatum.
affine, *Schult. f. Syst.* vii. 1700.—Afr. austr.
afrum, *Zuccagni, in Roem. Collect.* 137.—Aegypt.
albovirens, *Baker, in Gard. Chron.* (1878) II. 364.—Afr. austr.
albucoides, Hort. Par. ex Poir. Encyc. Suppl. iv. 191 = Albuca abyssinica.
albucoides, Thunb. Prod. Pl. Cap. 62 = suaveolens.
album, Kit. ex Schult. f. Syst. vii. 536, nomen (Quid?).—Europ.
algeriense, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 52 = orthophyllum.
alliaceum, *Lodd. Bot. Cab.* t. 1818.—Chili.
alpestre, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 53 = monticolum.
alpigenum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 79.—Lycia.
altaicum, Laxm. in Nov. Comm. Acad. Petrop. xviii. (1774) 530. t. 7. f. 1 = Lloydia alpina.
altissimum, Linn. f. Suppl. 199 = Urginea altissima.
andinum, Poepp. Fragm. Syn. Pl. Chil. 9 = Allium Poeppigi.
angustifolium, *Bor. Fl. Centr. Fr.* ed. III. ii. 625.—Gallia.
anomalum, *Baker, in Saund. Refug. Bot.* t. 178.—Afr. austr.

ORNITHOGALUM:—
anthericoides, Link, ex Steud. Nom. ed. I. 573 = Urginea anthericoides.
arabicum, Brot. Fl. Lusit. i. 531 = narbonense.
arabicum, [DC. in] Red. Lil. t. 63 = thyrsoides.
arabicum, *Linn. Sp. Pl.* 307.—Reg. Mediterr.
arcuatum, *Stev. in Mém. Soc. Nat. Mosc.* vii. (1829) 271.—Reg. Caucas.
aridum, Poepp. Fragm. Syn. Pl. Chil. 9 = arabicum.
armeniacum, *Baker, in Gard. Chron.* (1879) I. 748.—As. Min.
arvense, Pers. in Usteri, Ann. Bot. xi. (1794) 8 = Gagea stellaris.
asphodeloides, Schur, Enum. Pl. Transs. 664 = pyrenaicum.
atticum, Boiss. & Orph. ex Boiss. Fl. Orient. v. 217 = montanum.
Aucheri, Boiss. Diagn. Ser. I. v. 65 = oligophyllum.
Aucheri, Boiss. Fl. Orient. v. 222 (non Diagn.) = Balansae.
aurantiacum, *Baker, in Gard. Chron.* (1878) II. 748.—Afr. austr.
aureum, Curt. Bot. Mag. t. 190 = thyrsoides.
autumnale, Lam. Fl. Fr. iii. 274 = Scilla autumnalis.
baeticum, Boiss. Elench. 84 = umbellatum.
Balansae, *Boiss. Fl. Orient.* v. 222.—As. Min.
barbatum, *Jacq. Hort. Schoenb.* i. 47. t. 91.—Afr. austr.
belgicum, Lej. Rev. Fl. Spa, 67 = Gagea bracteolaris, pygmaea.
benguellense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 248.—Afr. trop.
Bergii, *Schlecht. in Linnaea*, i. (1826) 254.—Afr. austr.
Bertolonii, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 58 = exscapum.
bicolor, Haw. Misc. Nat. 177 = thyrsoides.
biflorum, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. iv. t. 246.—Chili; Peruv.
biflorum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 58 = exscapum.
bifolium, C. Koch, in Linnaea, xix. (1847) 10 = oligophyllum.
bifolium, Lodd, Bot. Cab. t. 1802 = biflorum.
bifolium, Neck. in Act. Acad. Theod. Palat. ii. (1770) 461 = Scilla bifolia.
bivalve, Linn. Sp. Pl. 306 = Nothoscordum striatum.
boeticum, vide *baeticum*.
bohemicum, Gaud. Fl. Helv. ii. 507 = Gagea saxatilis.
bohemicum, Lag. ex Willk. & Lange, Prod. Fl. Hisp. i. 218 = Gagea bracteolaris.
bohemicum, Loisel. Fl. Gall. ed. II. i. 243 = Gagea Liotardi.
bohemicum, Tenore, Fl. Nap. i. 172 = Gagea chrysantha.
bohemicum, Zauschn. in Abh. Privatges. Prag, ii. (1776) 120. t. 4 = Gagea pygmaea.
Bolusianum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 279.—Afr. austr.
bonariense, Pers. Syn. i. 363 = Nothoscordum Poiretii.
Boraeanum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 55 = umbellatum.
Boucheanum, Aschers. in Verh. Bot. Ver. Brand. viii. (1866) 165 = nutans.
Bourgaeanum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 52 = orthophyllum.
brachystachys, *Hort. Gorenk. ex Schult. f. Syst.* vii. 536, nomen.—Dahuria.
brachystachys, C. Koch, in Linnaea, xxii. (1849) 248 = narbonense.
bracteatum, Thunb. Prod. Pl. Cap. 62 = longebracteatum.
bracteatum, Torr. in Ann. Lyc. N. York, ii. (1828) 251 = Lloydia alpina.
brevipedicellatum, *Boiss. ex Baker, in Journ. Linn. Soc.* xiii. (1873) 263.—Lycia; Ins. Cyprus.
brevistylum, Wolfner, in Oestr. Bot. Wochenbl. vii. (1857) 230 = narbonense.
bulbiferum, Pall. Reise, ii. 736 = Gagea bulbifera.
bulbigerum, Herb. ex Schult. f. Syst. vii. 552 = Gagea bulbifera.
bulbosum, Herb. ex Schult. f. l. c. = praec.
Bungei, *Boiss. Fl. Orient.* v. 213.—Persia.
busambarense, Tineo, in Guss. Fl. Sic. Syn. ii. 813 = Gagea foliosa?
byzanthinum, Strangways & Tenore, ex Parl. Giorn. Bot. Ital. ii. (1847) VII. 106 = montanum.
byzantinum, Hort. ex Baker, in Journ. Linn. Soc. xiii. (1873) 264 = cuspidatum.

ORNITHOGALUM:—
caeruleum, Steud. Nom. ed. II. ii. 228 = *ceruleum*.
calcaratum, *Baker, in Gard. Chron.* (1874) I. 723.—Afr. austr.
callosum, Kit. in Schult. Oestr. Fl. ed. II. i. 557 = Gagea minima.
canadense, Linn. Sp. Pl. 308 = Albuca major, minor.
canaliculatum, Baker, in Saund. Refug. Bot. t. 177, ad calcem = Eckloni, *Schlecht.*
canaliculatum, Lag. Gen. et Sp. Nov. 14 = graminifolium.
capense, Linn. Sp. Pl. 308 = Eriospermum latifolium.
capitatum, *Hook. f. Bot. Mag.* t. 5388.—Afr. austr.
carolinianum, Hort. ex Schult. f. Syst. vii. II. 1128 = Nothoscordum striatum.
carpaticum, Kit. in Linnaea, xxxii. (1863) 336 = villosum.
caucasicum, *Boiss. ex Tchihat. Asie Min. Bot.* ii. 547, nomen.—Armenia.
caudatum, *Jacq. Coll.* ii. 315.—Afr. austr.
cepaefolium, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 248.—Afr. trop.
ceruleum, Rafin. Caratteri, 85 = Scilla Cupani.
cespititium, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 56 = umbellatum.
chilense, *Phil. in Linnaea*, xxix. (1857–58) 73.—Chili.
chinense, Schult. f. Syst. vii. 555, in obs. = Scilla chinensis.
chloranthum, *Baker, in Gard. Chron.* (1875) II. 323.—Afr. austr.
chloranthum, Bieb. Fl. Taur. Cauc. ii. 264 = Gagea chlorantha.
chloranthum, Sauter, ex Koch, Taschenb. 508 = nutans.
chloroleucum, Lindl. Bot. Reg. t. 1853 = biflorum.
chrysanthum, Jan, Elench. 5; Guss. Suppl. Fl. Sic. Prod. 100 = Gagea chrysantha.
ciliare, Fisch. ex R. Grah. in Edinb. N. Phil. Jou n. (Apr.–Jun. 1831) 191 = fimbriatum.
ciliatum, [Eckl.] ex Schult. f. Syst. vii. II. 1697 = Bergii.
ciliatum, Linn. f. Suppl. 199 = Urginea ciliata.
circinnatum, Linn. f. l. c. = Gagea reticularis.
Clusii, Tausch, in Flora, xi. (1828) 431 = Gagea pusilla.
coarctatum, *Jacq. Ic. Pl. Rar.* ii. 20. t. 435.—Afr. austr.
coccineum, Pritz. Ic. Ind. 773, sphalm. = Ornithidium coccineum.
coeruleum, Parl. Fl. Ital. ii. 467 = *ceruleum*.
collinum, *Guss. Ind. Sem. Hort. Bocc.* (1825); *Fl. Sic. Prod.* i. 412.—Europ. austr.
collinum, Koch, Syn. Fl. Germ. ed. II. 821 = tenuifolium.
collinum, Phil. ex Kunth, Enum. Pl. iv. 361 = comosum.
comosum, *Linn. Cent. Pl.* ii. 15; *Amoen. Acad.* iv. 312.—Europ. or.
comosum, Sadler, Fl. Pest. ed. I. i. 262 = collinum.
compositum, Knaf, ex Opiz, Seznam, 45, in syn. = Gagea bracteolaris.
comptum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 274.—Afr. austr.
concinnum, Salisb. Prod. 240 = unifolium.
conicum, Jacq. Coll. iii. 232 = lacteum.
conicum, Willd. ex Kunth, Enum. Pl. iv. 353 = coarctatum.
consanguineum, Kunth, l. c. 368 = tuberosum.
contortum, Tenore, Syll. 595 = comosum.
Cooperi, *Baker, in Journ. Linn. Soc.* xiii. (1873) 284.—Afr. austr.
corniculatum, *Hort. Belved. ex Schult. f. Syst.* vii. II. 1702, nomen.—Hab.?
cornosum, Vis. in Flora, xii. (1829) I. Erg. 11, sphalm. = comosum?
corsicum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 133 = biflorum.
corymbosum, Ruiz & Pav. Fl. Per. iii. 68. t. 300 = arabicum.
crenulatum, *Linn. f. Suppl.* 198.—Afr. austr.
croceum, Ell. Sketch, i. 397 = Anthericum croceum.
cuspidatum, *Bertol. Fl. Ital.* iv. 99, in obs.—Mesopot.
cuspidatum, Griseb. Spicil. Fl. Rumel. ii. 392 = montanum.
Cydni, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* iv. (1854) 153.—Cilicia.
decipiens, Goiron. 'Spec. Morph. Veg.' 21 (Quid?).—Hab.?

ORNITHOGALUM:—
declinatum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 57 = divergens.
deltoideum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 281.—Afr. austr.
densum, Boiss. & Blanche, in Boiss. Diagn. Sér. II. iv. 107 = narbonense.
dichotomum, Labill. Nov. Holl. Pl. i. 83. t. 109 = Thysanotus dichotomus.
divaricatum, Lindl. Bot. Reg. (1841) Misc. 53 = Chlorogalum pomeridianum.
divergens, *Bor. Fl. Centr. Fr.* 507; *et* ed. III. ii. 625.—Europ. austr.
divergens, *Sang. in Atti Acc. Pont. Lincei*, Sér. I. xviii. (1865) t. 5 (an praec.?).—Italia.
Dregeanum, Kunth, Enum. Pl. iv. 351 = graminifolium.
dubium, Houtt. Handleid. xii. 309 = thyrsoides.
Eckloni, *Schlecht. in Linnaea*, xxv. (1852) 177.—Afr. austr. et trop.
Ecklonii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 62.—Afr. austr.
edule, Siebold, in Verh. Batav. Gen. xii. (1830) 16 = Tulipa edulis.
elatum, *Andr. Bot. Rep.* t. 528.—Aegypt.
ensifolium, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 173.—Afr. trop.
erubescens, Bess. Enum. Pl. Volh. 45 = Gagea Liotardi.
etruscum, Parl. Fl. Ital. ii. 443 = orthophyllum.
exscapum, Gasp. ex Tenore, Syll. 173 = refractum.
exscapum, F. Schult. in Flora. x. (1827) 664 = Gagea bracteolaris.
exscapum, *Tenore, Prod. Fl. Nap.* p. xxii.—Europ. austr.
exuviatum, Kunth, Enum. Pl. iv. 369 = Urginea exuviata.
falcatum, Heyne, ex Wall. Cat. n. 5057 = Chlorophytum laxum.
fasciculatum, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 126.—Gallia.
fibrosum, Desf. Fl. Atlant. i. 294 = Gagea fibrosa.
filifolium, Kunth, Enum. Pl. iv. 369 = Urginea filifolia.
filiforme, *Ledeb. Fl. Alt.* ii. 30.—Sibir.
fimbriatum, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 420.—Graec.; As. Min.
fistulosum, Ram. ex DC. Fl. Fr. iii. 215 = Gagea Liotardi.
fistulosum, Roth, Enum. i. II. 42 = Gagea saxatilis.
flavescens, Jacq. Coll. iii. 229 = thyrsoides.
flavescens, Lam. Fl. Fr. iii. 277 = pyrenaicum.
flavissimum, Jacq. Ic. Pl. Rar. ii. 20. t. 436 = thyrsoides.
flavovirens, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. austr.
flavum, *Forsk. Fl. Aegypt. Arab.* 209.—Arab.
foliosum, J. & C. Presl, Delic. Prag. 149 = Gagea chrysantha.
fragiferum, Vill. Hist. Pl. Dauph. ii. 270 = Gagea Liotardi.
fragrans, Kunth, Enum. Pl. iv. 366 = Urginea fragrans.
fuscatum, *Jacq. Ic. Pl. Rar.* ii. 19. t. 429.—Afr. austr.
fuscescens, *Boiss. & Gaill. Diagn.* Sér. II. iv. 107.—Syria.
gadense, Wight, ex Wall. Cat. sub n. 5062 = Urginea indica.
garganicum, Tenore, Ind. Sem. Hort. Neap. (1827) 3; et in Mem. Acc. Sc. Borb. Nap. iii. II. (1832) 110. t. 6 = comosum.
geminiflorum, Herb. ex Lindl. Bot. Reg. (1838) Misc. 56 = biflorum.
giganteum, Jacq. Hort. Schoenb. i. 45. t. 87 = Urginea altissima.
glaucophyllum, *Baker, in Gard. Chron.* (1875) II. 36.—As. Min.
glaucum, Dethard. ex Schult. f. Syst. vii. II. 1702 = Gagea bracteolaris.
gracile, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. austr.
gracile, Hagen, Chlor. Boruss. 122 = Gagea minima.
gracile, Hort.—Cf. Gard. Chron. (1883) II. 506; (1885) I. 480 = lacteum.
graciliflorum, C. Koch, in Linnaea, xxii. (1849) 248 = montanum.
gramineum, Lam. Fl. Fr. iii. 278 = Anthericum Liliago.
gramineum, Poepp. ex Kunth, Enum. Pl. iv. 325 = biflorum.
gramineum, Sims, Pot. Mag. t. 2419 = Nothoscordum striatum.

ORNITHOGALUM:—
graminifolium, *Thunb. Prod. Pl. Cap.* 61.—Afr. austr.
Granatelli, Parl. in Diar. l'Occhio (1839), n. xi. p. 85 = Gagea Granatelli.
Grimaldiae, Nocca, Pl. Sel. Tic. 7. t. 4 = thyrsoides.
griseum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 281.—Afr. austr.
Gussonii, Tenore, Fl. Nap. iii. 371 = tenuifolium.
haworthioides, *Baker, in Journ. Bot.* xvi. (1878) 322.—Afr. austr.
hexapterum, Fenzl, ex Boiss. Fl. Orient. v. 221 = oligophyllum.
Heynii, Roth, in Roem. Archiv, i. III. 42 = Gagea spathacea.
hirsutum, Linn. Sp. Pl. 306 = Hypoxis erecta.
hispidum, *Hornem. Hort. Hafn.* 331.—Afr. austr.
horologicum, Stokes, Bot. Mat. Med. ii. 242 = umbellatum.
hortense, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 54 = umbellatum.
Huetii, Boiss. Diagn. Ser. II. iv. 108 = montanum.
humifusum, *Baker, in Gard. Chron.* (1874) I. 500.—Afr. austr.
hyalinum, Granier, ex Brouss. Elench. Hort. Monsp. (1805) 41, nomen = nutans.
indicum, Willd. ex Schult. f. Syst. vii. 535 = Chlorophytum attenuatum.
intermedium, Schleich. ex Steud. Nom. ed. II. ii. 228 = Gagea Liotardi.
intermedium, Schultz, ex Schult. f. Syst. vii. 537 = Gagea bracteolaris.
ixioides, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 257 = Brodiaea ixioides.
japonicum, Thunb. Fl. Jap. 137 = Scilla japonica.
Joannoni, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 54 = umbellatum.
junceum, Poir. Encyc. Suppl. iv. 197, in syn. = graminifolium.
juncifolium, *Jacq. Hort. Schoenb.* i. 46. t. 90.—Afr. austr.
juncifolium, Ker-Gawl. in Bot. Mag. t. 972 = graminifolium.
Kirkii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 279.—Afr. trop.
Kochii, Parl. Fl. Ital. ii. 440 = tenuifolium.
lacteum, *Jacq. Ic. Pl. Rar.* ii. 20. t. 434; *Coll. Suppl.* 76.—Afr. austr.
lacteum, Vill. Hist. Pl. Dauph. ii. 272 = narbonense.
lanceolatum, Kunth, Enum. Pl. iv. 360, partim = cuspidatum.
lanceolatum, *Labill. Ic. Pl. Syr. Dec.* v. 11.—Syria.
latifolium, Linn. Herb. ex Baker, in Journ. Linn. Soc. xiii. (1873) 275 = pyrenaicum.
latifolium, *Linn. Sp. Pl.* 307.—Arab.; Aegypt.
latifolium, Ucria, Hort. Reg. Panorm. 155 = arabicum.
libanoticum, *Boiss. Diagn.* Ser. II. iv. 106.—Syria.
liliforme, Lam. Fl. Fr. iii. 278 = Paradisia Liliastrum.
Liotardi, Sternb. in Denkschr. Bot. Ges. Regensb. ii. (1818) 56. t. 3 = Gagea Liotardi.
longebracteatum, *Jacq. Hort. Vindob.* iii. 18. t. 29.—Afr. austr.
Luschani, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 79.—Mesopot.
luteum, Linn. Sp. Pl. 306 = Gagea stellaris, fasciculatis, bracteolaris.
luteum, Tenore, Fl. Nap. iii. 370, excl. syn. = Gagea pusilla.
macranthum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 280.—Afr. austr.
maculatum, *Jacq. Coll.* ii. 368. t. 18. f. 3.—Afr. austr.
maculatum, Thunb. Prod. Pl. Cap. 62 = Thunbergianum.
majus, Gilib. Exercit. ii. 467 = *luteum*, Linn.
marginatum, Lindl. Bot. Reg. (1845) t. 21 = cuspidatum.
marginatum, Pall. Tabl. Taur. 49 = Gagea pusilla.
maritimum, Lam. Fl. Fr. iii. 276 = Urginea maritima.
Massoni, J. F. Gmel. Syst. 551 = caudatum.
Melleri, *Baker, in Journ. Linn. Soc.* xiii. (1873) 280.—Afr. trop.
merizostylum, *Steud. Nom.* ed. II. ii. 229.—Chili.
millegranum, Janka, in Termész. Füzetek, viii. (1884) 29 = umbellatum.
miniatum, Jacq. Coll. iii. 233 = thyrsoides.
minimum, Heldr. ex Nym. Consp. 728 = exscapum.
minimum, Linn. Sp. Pl. 306 = Gagea minima.
minimum, O. F. Muell. in Fl. Dan. t. 612 = Gagea spathacea.
minimum, Willd. Sp. Pl. ii. 114 = Gagea stellaris.

ORNITHOGALUM:—
minus, Linn. Mant. ii. 364 = umbellatum.
minutum, Pall. Reise, ii. 14 = Gagea minima.
monile, *Pall. Reise in Suedl. Statth. Russ. Reich.* ii. 80, 89.—Tauria.
monspeliense, Gueldenst. Reisen, ii. 25 = narbonense.
montanum, *Tenore, Prod. Fl. Nap.* p. xxii.—Europ. austr.; Oriens.
monticolum, *Jord. & Fourr. Brev. Pl. Nov. fasc.* i. 54.—Gallia.
multifolium, *Baker, in Journ. Linn. Soc.* xiii. (1873) 271.—Afr. austr.
Mundianum, Kunth, Enum. Pl. iv. 351 = pilosum.
mutabile, De Not. in Mem. Acc. Sc. Torin. II. ix. (1848) 441 = exscapum.
nanum, Brot. Fl. Lusit. i. 529 = unifolium.
nanum, *Sibth. & Sm. Fl. Graec. Prod.* i. 230; *Fl. Graec.* iv. 28. t. 333.—Graecia; As. Min.
nanum, Spreng. Pugill. ii. 57 = exscapum.
nanum, Tenore, Ind. Sem. Hort. Neap. (1830); ex ej. Syll. 173 = umbellatum.
nanum, Thunb. Prod. Pl. Cap. 62 = Eucomis nana.
narbonense, Bertol. Amoen. Ital. 143 = pyrenaicum.
narbonense, Falk, Beitr. ii. 161 = Eremurus spectabilis.
narbonense, *Linn. Cent. Pl.* ii. 15; *Amoen. Acad.* iv. 312.—Reg. Mediterr.; Oriens.
narbonense, Reichard, Fl. Moeno-Fr. ii. 180 = umbellatum.
narbonense, Stev. in Mém. Soc. Nat. Mosc. vii. (1829) 272 = arcuatum.
nebrodense, Tod. Rar. Pl. Sic. Decad. i.; in Atti Acc. Sc. Palermo, N. S. i. (1845) 7 = Gagea saxatilis.
nervosum, Burch. Trav. i. 537 = Scilla rigidifolia.
neurostegium, *Boiss. & Blanche, ex Boiss. Fl. Orient.* v. 222.—Syria.
nivale, *Boiss. Diagn.* Ser. I. v. 65.—Graecia.
niveum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 440.—Afr. austr.
Nocceanum, *Balb. in Brugnat. Giorn. Fis.* ix. (1816) 446.—Hab. ?
notatum, Schult. f. Syst. vii. 528 = maculatum.
nudiflorum, Steud. Nom. ed. II. i. 652, in syn., sphalm. = seq.
nudiscapum, F. Schultz, in Flora, x. (1827) 664 = Gagea bracteolaris.
nutans, *Linn. Sp. Pl.* 308.—Europ.; As. Min.
Nuttallianum, Kunth, Enum. Pl. iv. 372 = Schoenolirion croceum.
nyssanum, *Petrović, Fl. Agr. Nyss.* 799.—Serbia.
octandrum, Fingerh. in Linnaea, iv. (1829) 383. t. 4 = Gagea stellaris.
odoratum, Jacq. Ic. Pl. Rar. ii. 20. t. 432; Coll. Suppl. 78 = suaveolens.
oligophyllum, *Clarke, Trav.* iv. 555.—Graecia; As. Min.; Reg. Caucas.
orthophyllum, *Tenore, Syll. Fl. Neap.* 594.—Italia.
ovatum, *Thunb. Prod. Pl. Cap.* 62.—Afr. austr.
oxypetalum, Ledeb. Fl. Alt. ii. 27 = Tulipa uniflora.
pallidum, Salisb. Prod. 239 = pyrenaicum.
paludosum, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. austr.
palustre, *Eckl. Top. Verz.* 6.—Afr. austr.
pannonicum, Chaix, ex Vill. Prosp. 18 = Gagea minima.
paradoxum, Jacq. Coll. Suppl. 81 = Eriospermum paradoxum.
parviflorum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 56 = angustifolium.
parviflorum, Licht. ex Kunth, Enum. Pl. iv. 571 = Bulbinella robusta.
Pater-familias, *Godr. in Mém. Soc. Emul. Doubs*, Sér. II. v. (1854) 15.—Gallia.
patulum, Rafin. Préc. Découv. 44 = montanum.
pauciflorum, Rafin. l. c. = exscapum.
pauciflorum, Turcz. in Flora, xvii. (1834) Beibl. i. 25 = Lloydia triflora.
pedicellare, Boiss. & Kotschy, in Unger & Kotschy, Cypern, 195.—Ins. Cyprus.
pedunculare, Presl, Delic. Prag. 150 = Gagea pygmaea.
Persoonii, Hoppe, in Bot. Zeit. Regensb. vi. (1807) 138 = Gagea fascicularis.
Peyrei, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 125.—Gallia.
pilosum, *Linn. f. Suppl.* 199.—Afr. austr.
platyphyllum, Boiss. Diagn. Ser. I. v. 64 = montanum.
podolicum, Bess. ex Schult. f. Syst. vii. 545 = Gagea pusilla ?

ORNITHOGALUM:—
polymorphum, F. Schultz, in Arch. Fl. Fr. et Allem. 18 = bracteolaris.
polyphyllum, Herb. Heyne, ex Wall. Cat. n. 5062 F (Urgineae sp.).—Ind. or.
polyphyllum, Jacq. Ic. Pl. Rar. ii. 19. t. 430; Coll. Suppl. 79 = tuberosum.
polyphyllum, Link, ex Kunth, Enum. Pl. iv. 361 = montanum.
praetextum, Stev. in Hort. Berol. (1842); ex Kunth, l. c. 363 = umbellatum.
prasandrum, Griseb. Spicil. Fl. Rumel. ii. 390 = nutans.
prasinum, *Ker-Gawl. in Bot. Reg.* t. 158.—Afr. austr.
pratense, Pers. in Usteri, Ann. Bot. xi. (1794) 8 = Gagea bracteolaris.
pratense, Sadler, Fl. Pest. i. 260 = Gagea pusilla.
procerum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 14.—Persia.
proliferum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 57 = divergens.
proliferum, Pall. Tabl. Taur. 49 = Gagea scillaris.
propinquum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 53 = monticolum.
Pseudo-tenuifolium, *Goiron*, '*Spec. Morph. Veg.*' 22 (Quid ?).—Hab. ?
pterocarpum, Kotschy & Boiss. ex Baker, in Journ. Linn. Soc. xiii. (1873) 263 = oligophyllum.
pubescens, *Baker, l. c.* 282.—Afr. austr.
pulchellum, Salisb. Prod. 239 = Nothoscordum striatum.
punctatum, Thunb. Prod. Pl. Cap. 62 = Eucomis punctata.
pusillum, Noé, ex Baker, in Journ. Linn. Soc. xiii. (1873) 269 = nanum.
pusillum, F. W. Schmidt, Fl. Boëm. Cent. iv. 41 = Gagea pusilla.
pygmaeum, Willd. Enum. Hort. Berol. 367 = Gagea pygmaea.
pyramidale, Linn. Sp. Pl. 307 = narbonense.
pyramidale, Tenore, Viag. Garg. 366. t. 1 = comosum.
pyrenaicum, Desf. Fl. Atlant. i. 295 = narbonense.
pyrenaicum, *Linn. Sp. Pl.* 306.—Europ. austr.; Oriens.
ramosum, Lam. Fl. Fr. iii. 279 = Anthericum ramosum.
refractum, Guss. Suppl. Fl. Sic. Prod. 101 = divergens.
refractum, Moris, Stirp. Sard. i. 46 = exscapum.
refractum, Noé, ex Nym. Consp. 728 = nanum.
refractum, *Willd. Enum. Hort. Berol. Suppl.* 18.—Hungar.; Reg. Caucas.
reticulatum, Pall. Reise, iii. 727 = Gagea reticularis.
revolutum, *Jacq. Hort. Schoenb.* i. 46. t. 89.—Afr. austr.
rigidulum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 53 = monticolum.
rigidum, Sprun. ex Boiss. Diagn. Ser. I. vii. 109 = Gagea reticularis.
roccense, Link, in Schrad. Journ. ii. (1799) 320 = unifolium.
Roegnerianum, C. Koch, in Linnaea, xxii. (1849) 246 = fimbriatum.
rubrum, Ruiz & Pav. ex D. Don, in Edinb. N. Phil. Journ. (Oct. 1832) 235 = Xiphidium rubrum.
Rudolphi, Jacq. Eclog. i. 31, 151. t. 20 = graminifolium.
rupestre, *Linn. f. Suppl.* 199.—Afr. austr.
rupestre, Rudolphi, in Schrad. Journ. ii. (1799) 281 = graminifolium.
rupestre, Schlecht. ex Kunth, Enum. Pl. iv. 569 = Bulbinella triquetra.
rusticum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 55 = umbellatum.
ruthenicum, *Bouché, ex Kunth, Enum. Pl.* iv. 363.—Europ. austr.
ruthenicum, Griseb. Spicil. Fl. Rumel. ii. 393 = oligophyllum.
saxatile, Koch, ex Schult. f. Syst. vii. 549 = Gagea saxatilis.
saxatile, Vis. in Flora, xii. (1829) i. Erg. 10 = collinum.
Schreberi, Reichb. l. c. v. (1822) 587 = Gagea bracteolaris.
scilloides, *Jacq. Hort. Schoenb.* i. 46. t. 88.—Afr. austr.
secundum, Jacq. Ic. Pl. Rar. ii. 20. t. 433; Coll. Suppl. 79 = Urginea secunda.

ORNITHOGALUM:—
serotinum, Reichb. in Moessl. Handb. ed. II. i. 554 = Lloydia alpina.
sessile, *Pacho, Voy. Marm.* t. 96 (Quid ?).—Graecia.
sessiliflorum, *Desf. Fl. Atlant.* i. 295.—Afr. bor. occ.
setifolium, Kunth, Enum. Pl. iv. 351 = niveum.
siculum, *Ucria, in Roem. Arch.* i. 1. 68.—Sicilia.
simplex, Becker, Fl. Frankf. i. 141 = Gagea bracteolaris.
sinense, Lour. Fl. Cochinch. 206 = Scilla japonica.
Soleirolii, Mutel, Fl. Fr. iii. 298, et Index, 93 = Gagea foliosa.
sororium, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* iv. (1854) 105.—Cilicia.
spathaceum, Hayne, in Usteri, Ann. Bot. xxi. (1797) 11 = Gagea spathacea.
spathaceum, Poir. Encyc. iv. 618 = Nothoscordum Poiretii.
spathaceum, Sibth. & Sm. Fl. Graec. iv. 26. t. 331 = Gagea foliosa.
speciosum, Rafin. Caratt. 85 = Scilla peruviana.
speciosum, Salisb. Prod. 240 = arabicum.
sphaerocarpum, A. Kern. in Oestr. Bot. Zeitschr. xxviii. (1878) 15 = pyrenaicum.
spicatum, Plan. Ens. Fl. Gallega, 381 = unifolium.
spirale, Wimm. in Linnaea, xvi. (1842) Litt. 274 = Eriospermum spirale.
squamosum, Lam. Fl. Fr. iii. 274 = Scilla Liliohyacinthus.
Squilla, Ker.-Gawl. in Bot. Mag. t. 918 = Urginea maritima.
stachyodes, [Soland. in] Ait. Hort. Kew. ed. I. i. 441 = narbonense.
stachyoides, Host, Fl. Austr. i. 432 = pyrenaicum.
stellare, Salisb. Prod. 239 = umbellatum.
stenopetalum, Fries, Novit. Fl. Suec. ed. II. 87 = Gagea bracteolaris.
stenophyllum, Schur, Enum. Pl. Transs. 942 (Index), sphalm. = praec.
Sternbergii, Hoppe, in Bot. Zeit. Regensb. v. (1806) 339, cum tab. = Gagea minima.
striatum, Willd. Sp. Pl. ii. 112 = Lloydia alpina.
suaveolens, *Jacq. Coll.* ii. 316.—Afr. austr.
subalpinum, J. Gay, ex Baker, in Journ. Linn. Soc. xiii. (1873) 266 = monticolum.
subulatum, *Baker, in Gard. Chron.* (1874) i. 723.—Afr. austr.
sulfureum, Schult. f. Syst. vii. 518 = pyrenaicum.
sulphureum, Auct. = praec.
sylvaticum, Pers. in Usteri, Ann. Bot. xi. (1794) 7 = Gagea fascicularis.
sylvaticum, Schleich. ex Kunth, Enum. Pl. iv. 236 = Gagea bracteolaris.
Szovitsii, Láng, in Flora, x. (1827) i. Beil. 64 = Gagea pygmaea.
tardans, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 56 = umbellatum.
tenellum, *Jacq. Coll.* ii. 316.—Afr. austr.
tenuifolium, [DC. in] Red. Lil. t. 312 = graminifolium.
tenuifolium, Gren. & Godr. Fl. Fr. iii. 191, partim = monticolum.
tenuifolium, *Guss. Fl. Sicul. Prod.* i. 413.—Europ. austr.; As. Min.; Aegypt.
tenuifolium, Schur, ex Nym. Consp. 727 = umbellatum.
texanum, Scheele, in Linnaea, xxiii. (1850) 146 = Schoenolirion croceum.
Thirkeanum, C. Koch, l. c. xix. (1847) 17 = nutans.
Thunbergianum, *Baker, in Journ. Linn. Soc.* xiii. (1873) 269.—Afr. austr.
Thunbergii, Kunth, Enum. Pl. iv. 369 = Urginea nematodes.
thyrsoides, *Jacq. Hort. Vind.* iii. 17. t. 28.—Afr. austr.
transbaicalense, Pall. in Nov. Act. Acad. Petrop. x. (1795) 309 = Gagea transbaicalensis.
transversale, Pall. l. c. = Gagea spathacea.
triandrum, Labill. Nov. Holl. Pl. i. 84. t. 110 = Thysanotus triandrus.
trichophyllum, Boiss. & Heldr. in Boiss. Diagn. Ser. II. iv. 108 = tenuifolium.
triflorum, Ledeb. in Mém. Acad. Pétersb. v. (1812) 529 = Lloydia triflora.
trigonophyllum, Auct. ex Schult. f. Syst. vii. 544 = Gagea pusilla.
trigynum, [Delile, in] Red. Lil. t. 417 = narbonense.
tropicale, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. trop.

ORNITHOGALUM:—
tuberosum, *Mill. Gard. Dict.* ed. VIII. n. 10.—Afr. austr.
tunicatum, J. & C. Presl, Delic. Prag. 216 = Gagea pusilla.
umbellatum, *Linn. Sp. Pl.* 307.—Europ.; As. Min.; Afr. bor.
undulatum, Thunb. Prod. Pl. Cap. 62 = Eucomis undulata.
uniflorum, Linn. Mant. i. 62 = Tulipa uniflora.
uniflorum, Zauschn. in Abh. Privatges. Prag, ii. (1776) 120 = Gagea pusilla.
unifolium, *Ker-Gawl. in Bot. Mag.* t. 935.—Reg. Mediterr. occ.
unifolium, *Retz. Obs.* ii. 17.—Afr. austr.
villosum, Bieb. Fl. Taur. Cauc. i. 274 = Gagea stellaris.
villosum, *Kit. in Linnaea*, xxxii. (1863) 386.—Hungar.
virens, *Lindl. in Bot. Reg.* t. 814.—Afr. austr.
virgineum, *Soland. ex Baker, in Journ. Linn. Soc.* xiii. (1873) 271.—Afr. austr.
Visianicum, *Tommas. ex Vis. in Mém. Ist. Venet.* xx. (1876) 176.—Dalmat.
vittatum, *Kunth, Enum. Pl.* iv. 367.—Afr. austr.
Wiedemanni, *Boiss. Fl. Orient.* v. 221.—Anatolia.
Zauschneri, Pohl, Fl. Bohem. ii. 14 = Gagea pygmaea.
Zeyheri, *Baker, in Journ. Linn. Soc.* xiii. (1873) 281.—Afr. austr.
zeylanicum, Hort. Belved. ex Schult. f. Syst. vii. 1702, nomen (Quid?).—Zeylan.

ORNITHOGLOSSUM, Salisb. Parad. Lond. t. 54 (1806). *LILIACEAE*, Benth. & Hook. f. iii. 824.
CYMATION, Spreng. Syst. ii. 142 (1825).
LICHTENSTEINIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 19. t. 1.
glaucum, *Salisb. Parad. Lond.* t. 54.—Afr. austr.
Lichtensteinii, Schlecht. in Linnaea, i. (1826) 91 = glaucum.
undulatum, Spreng. Syst. iv. Cur. Post. 143 = glaucum.
viride, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 327 = glaucum.

ORNITHOPHORA, Rodrig. Orch. Nov. ii. 225. *ORCHIDEAE*.
quadricolor, *Rodrig. l. c.* 226.—Bras.

ORNITHOPODIUM, [Tourn.] Linn. Syst. ed. I (1735) = **Ornithopus**, Linn. (Legumin.).
compressum, All. Fl. Pedem. i. 336 = Ornithopus compressus.
perpusillum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 369 = Ornithopus perpusillus.
scorpioides, Scop. Fl. Carn. ed. II. ii. 73 = Coronilla scorpioides.
triphyllum, Moench, Meth. 121 = Coronilla scorpioides.

ORNITHOPUS, Linn. Gen. ed. I. 226 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 509.
ANTOPETITIA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 261.
ARTROLOBIUM, Desv. Journ. Bot. iii. (1814) 121. t. 4.
ASTROLOBIUM, DC. Prod. ii. 311 (1825).
ORNITHOPODIUM, [Tourn.] Linn. Syst. ed. I (1735).
compressus, *Linn. Sp. Pl.* 744.—Reg. Mediterr.; Oriens.
compressus, Sieber, ex Steud. Nom. ed. II. ii. 230 = Coronilla scorpioides.
coriandrinus, *Hochst. & Steud. in Flora*, xxiv. (1841) i. *Intell.* 32.—Abyssin.
durus, *Cav. Ic.* i. 31. t. 41.—Lusitan.
durus, DC. Fl. Fr. iv. 603 = ebracteatus.
ebracteatus, *Brot. Fl. Lusit.* ii. 159. t. 68.—Europ. occ.; Reg. Mediterr.
exstipulatus, Thore, Chlor. Land. 311 = ebracteatus.
heterophyllus, Brot. Fl. Lusit. ii. 160 = durus.
intermedius, Roth, Tent. Fl. Germ. i. 319 = perpusillus.
intermedius, Salzm. ex Baker, in Journ. Linn. Soc. xvi. (1878) 426 = isthmocarpus.
isthmocarpus, *Coss. Notes Crit.* ii. 36.—Reg. Mediterr. occ.
laevigatus, Sm. in Rees, Cycl. xxv. n. 6 = ebracteatus.
lotoides, Viv. Ann. Bot. i. II. 177; et Fragm. Fl. Ital. i. 7. t. 9 = Coronilla repanda.
nodosus, Mill. Gard. Dict. ed. VIII. n. 2 = perpusillus.

ORNITHOPUS:—
nudiflorus, Lag. in Varied. Cienc. ii. IV. (1805) 40 = ebracteatus.
perpusillus, *Linn. Sp. Pl.* 743.—Europ.
pusillus, Lepech. Reise Versch. Prov. Russ. Reich. iii. 193 = perpusillus?
pygmaeus, Viv. Fragm. Fl. Ital. i. 13. t. 14. f. 2 = ebracteatus.
repandus, Poir. Voy. Barb. ii. 215; et in Lam. Encyc. iv. 580 = Coronilla repanda.
roseus, Duf. in Ann. Sc. Nat. Sér. I. v. (1825) 82 = perpusillus.
ruber, Lour. Fl. Cochinch. ii. 452 = Desmodium rubrum.
sativus, *Brot. Fl. Lusit.* ii. 160.—Hispan.
sativus, Gren. & Godr. Fl. Fr. i. 499 = perpusillus.
scorpioides, Linn. Sp. Pl. 744 = Coronilla scorpioides.
subumbellatus, Gilib. Fl. Lituan. ii. 108 = perpusillus.
tetraphyllus, Linn. Syst. ed. X. 1168 = Zornia myriadenia.
trifoliatus, Lam. Fl. Fr. ii. 659 = Coronilla scorpioides.

ORNITHORYNCHIUM, Roehl. Deutschl. Fl. ed. II. 356 (1812-13) = **Euclidium**, R. Br. (Crucif.).
syriacum, Roehl. l. c. = E. syriacum.

ORNITHOSPERMA, Rafin. Fl. Ludov. 149 (1817) = **Ipomoea**, Linn. (Convolv.).
autumnalis, Rafin. Fl. Tellur. iv. 72 = I. hederacea.
serotina, Rafin. Fl. Ludov. 149 = I. serotina.

ORNITHOXANTHUM, Link, Handb. i. 161 (1829) = **Gagea**, Salisb. (Liliac.).
arvense, Link, l. c. = G. arvensis.
bohemicum, Link, l. c. = G. bohemica.
fistulosum, Link, l. c. = G. Liotardi.
gracile, Link, l. c. 162 = G. minima.
Liotardi, Link, l. c. = G. Liotardi.
luteum, Link, l. c. 161, 162 = G. lutea.
minimum, Link, l. c. 162 = G. minima.
Persoonii, Link, l. c. = G. lutea.
pratense, Link, l. c. 161 = G. pratensis.
pusillum, Link, l. c. 162 = G. pusilla.
spathaceum, Link, l. c. = G. spathacea.
stenopetalum, Link, l. c. 161 = G. stenopetala.
Sternbergii, Link, l. c. 162 = G. minima?
sylvaticum, Link, l. c. = G. lutea.
villosum, Link, l. c. 161 = G. arvensis.
Zauschneri, Link, l. c. = G. bohemica.

ORNITHROPHUS, Boj. ex Engl. in Linnaea, xxxvi. (1869-70) 636 = **Weinmannia**, Linn. (Saxifr.).

ORNITOPUS, Krock. Fl. Siles. Suppl. ii. 214 (1823) = **Ornithopus**, Linn. (Legumin.).

ORNITROPE, Pers. Syn. i. 412 (1805) = seq.

ORNITROPHE, Comm. ex Juss. Gen. 247 (1789) = **Schmidelia**, Linn. (Sapind.).
Allophyllus, Pers. Syn. i. 412 = Allophylus zeylanicus.
aporetica, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 264 = Allophylus Cobbe.
asiatica, Hort. ex Steud. Nom. ed. II. ii. 230 = Allophylus Cobbe.
borbonica, J. F. Gmel. Syst. 613 = Allophylus Cobbe?
Cobbe, Willd. Sp. Pl. ii. 322 = Allophylus Cobbe.
Cominia, Willd. l. c. 323 = S. Cominia.
glabra, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 267 = Allophylus Cobbe.
glabrata, Poir. in Dict. Sc. Nat. xxxvi. 450 = S. glabrata.
integrifolia, Willd. Sp. Pl. ii. 322 = Allophylus Cobbe.
macrophylla, Poir. Encyc. viii. 263 = S. macrophylla.
magica, Schum. & Thonn. Beskr. Guin. Pl. 186 = S. magica.
malabarica, Herb. Madr. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 674 = Allophylus Cobbe.
mauritiana, Boj. ex Baker, Fl. Maurit. 57 = Allophylus Cobbe.
Merularia, Comm. ex DC. Prod. i. 610 = Pittosporum Senacia.
mollis, Poir. in Dict. Sc. Nat. xxxvi. 451 = S. mollis.
occidentalis, Willd. Sp. Pl. ii. 323 = S. occidentalis.
panigera, Labill. Sert. Austro-Caled. 52. t. 52 = S. Cominia.
pinnata, Poir. Encyc. viii. 366 = Pometia pinnata.
repanda, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 269.—Ins. Molucc.

ORNITROPHE:—
rigida, Willd. Sp. Pl. ii. 324 = Schmidelia rigida.
Schmidelia, Pers. Syn. i. 412 = Allophylus Cobbe.
serrata, Roxb. Pl. Corom. i. 44. t. 61 = Allophylus Cobbe.
spicata, Poir. Encyc. viii. 265 = Allophylus Cobbe.
thyrsoides, Schum. & Thonn. Beskr. Guin. Pl. 185 = S. thyrsoides.
triandra, Blanco, Fl. Filip. ed. I. 291 = Schleichera trijuga.
tristachyos, Schum. & Thonn. Beskr. Guin. Pl. 188 = Allophylus zeylanica.
villosa, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 265 = Allophylus Cobbe.

ORNUS, Neck. Elem. ii. 375 (1790) = **Fraxinus**, Tourn. (Oleac.).
americana, Bosc. in Mém. Instit. Fr. ix. 1808 (1811), 216; Pursh, Fl. Am. Sept. i. 9 = F. Ornus.
cappadocica, A. Dietr. Sp. Pl. i. 249 = F. Ornus.
dipetala, Nutt. Sylva iii. 66. t. 101 = F. dipetala.
europaea, Pers. Syn. i. 8 = F. Ornus.
floribunda, Sweet, Hort. Brit. ed. I. 271 = F. floribunda.
mammifera, Steud. Nom. ed. I. 575; ed. II. ii. 230, sphalm. loco *mannifera* = F. rotundifolia.
Moorcroftiana, G. Don, Gen. Syst. iv. 57 = F. excelsior.
rotundifolia, Moench, Meth. 55 = F. rotundifolia.
rotundifolia, Pers. Syn. ii. 605 = F. Ornus.
striata, Sweet, Hort. Brit. ed. I. 271 = F. strigata.
strigata, A. Dietr. Sp. Pl. i. 249 = F. angustifolia.
urophylla, G. Don, Gen. Syst. iv. 57 = F. floribunda.
xanthoxyloides, G. Don, l. c. = F. xanthoxyloides.

OROBA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 400 = OROBUS, Tourn. = **Lathyrus**, Linn. (Legumin.).

OROBANCHE, [Tourn.] Linn. Syst. ed. I (1735). *OROBANCHACEAE*, Benth. & Hook. f. ii. 984.
BOULARDIA, F. Schultz, Arch. Fl. Fr. et Allem. 103 (1847).
CATODIACRUM, Dulac, Fl. Hautes-Pyr. 369 (1867).
CERATOCALYX, Coss. in Ann. Sc. Nat. Sér. III. ix. (1848) 145. t. 10.
CHOROBANCHE, Benth. & Hook. f. Gen. ii. 985 (1876), lapsu.
CHOROBANE, Presl, Bot. Bemerk. 91 (1844); in Abh. Boehm. Ges. V. iii. (1845) 521.
KOPSIA, Dum. Comm. Bot. 16 (1822).
PHELIPANCHE, Pomel, Nouv. Mat. Fl. Atl. 102 (1874).
THALESIA, Rafin. Neogenyt. 3. (1825).
abortiva, Reut. in DC. Prod. xi. 37 = arenaria.
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 137 = minor, *Sutt.*
acaulis, Roxb. Hort. Beng. 45; Fl. Ind. iii. 28 = Aeginetia pedunculata.
adenostemon, Reichb. Fl. Germ. Excurs. 354 = caryophyllacea.
Aeginetia, Linn. Sp. Pl. ed. II. 883 = Aeginetia indica.
aegyptiaca, *Pers. Syn.* ii. 181.—Aegypt.
alba, Bieb. Fl. Taur. Cauc. ii. 82 = crenata.
alba, Reichb. Pl. Crit. vii. 40. f. 912 = reticulata.
alba, *Steph. ex Willd. Sp. Pl.* iii. 450.—Europ. or.; As. bor.
alba, Wierzb. in Flora, xxviii. (1845) 324 = alsatica.
albiflora, Gren. & Godr. Fl. Fr. ii. 628 = ramosa.
albolanata, Steud. Nom. ed. II. ii. 230 = caerulescens.
Alectra, D. Dietr. Syn. Pl. iii. 624 = Harveya capensis.
Alexandri, *Tineo, in Guss. Fl. Sic. Syn. Add.* ii. II. 845.—Sicil.
alostensis, Demoor. ex T. Durand & Donck. in Bull. Soc. Bot. Belg. xii. (1873) 397 = Carotae.
alpestris, F. W. Schultz, in Flora, xxiii. (1840) 128 = Salviae.
alpigena, *C. Koch, in Linnaea*, xxii. (1849) 668.—As. Min.
alsatica, *Kirschl. Prod. Fl. Alsat.* 109; *F. Schultz, Arch. Fl. Fr. et Allem.* 69.—Europ.
altissima, Noé, ex Nym. Consp. 558 = crenata.
ambigua, Moris, Diagn. Stirp. Sard. (1857) 2 = Picridis.
ambigua, *Pomel, Nouv. Mat. Fl. Atl.* 109.—Alger.
americana, Linn. Mant. i. 88 = Conopholis americana.
amethystea, Maly, ex Reut. in DC. Prod. xi. 718 = major.
amethystea, *Thuill. Fl. Par.* ed. II. i. 317.—Europ.; As. Min.; Afr. bor.
amethystea, Thuill. l. c. partim = gracilis.

OROBANCHE :—
amethystina, Reichb. Ic. Fl. Germ. vii. 42 = amethystea.
ammophila, C. A. Mey. in Ledeb. Fl. Alt. ii. 454 = caerulescens.
amoena, *C. A. Mey. l. c.* 457.—Sibir.
anatolica, *Boiss. & Reut. ex Reut. in DC. Prod.* xi. 17.—As. Min.; Reg. Cauc.; Persia.
angustisepala, *F. Schultz, in Flora*, xxvi. (1843) 130.—Graecia.
annulata, Griseb. Reise, ii. 188 = gracilis.
antirrhina, Reut. in DC. Prod. xi. 21 = reticulata.
apiculata, Griseb. Spicil. Fl. Rumel. ii. 58 = alsatica.
apiculata, Wallr. Sched. Crit. 310 = minor.
arachnoidea, F. Schultz, in Linnaea, xix. (1847) 26 = pubescens.
*araliocto*na, Meissn. in Bot. Zeit. xxiv. (1866) 17 = Hederae.
arcuata, *F. Schultz, in Flora*, xxvi. (1843) 1 9.—Graecia.
arenaria, Auct. ex Wallr. Orob. Gen. 36 = caryophyllacea.
arenaria, Bieb. Fl. Taur. Cauc. ii. 83 = cumana.
arenaria, *Borckh. in Roem. Neues Mag.* i. (1794) 6; *et ex Gaertn. Mey. & Scherb. Fl. Wett.* ii. 405.—Europ.
Artemisiae, Gren. in Gren. & Godr. Fl. Fr. ii. 638 = loricata.
Artemisiae-campestris, Gaud. Fl. Helvet. iv. 179 = loricata.
Artemisae-vulgaris, F. Schultz, in Ann. Gew. Regensb. v. (1830) 5 = arenaria.
artemisi-epiphyta, St. Lag. Cat. Fl. Rhône (1883), 611 = loricata.
arvensis, Dum. Fl. Belg. Prod. 33 = minor.
asiatica, Weinm. in Bull. Soc. Nat. Mosc. xxiii. (1850) 1. 552 = borealis.
atropurpurea, F. Schultz, in Flora, xxiii. (1840) 128 = seq.
atrorubens, F. Schultz, l. c. = Teucrii.
attica, Reut. in DC. Prod. xi. 28 = amethystea.
australiana, *F. Muell. Census*, 98, nomen.—Austral.
australis, Moris, ex Bertol. Fl. Ital. vi. 440 = canescens.
Avellanae, Pfund, in Flora, xxvi. (1843) 183 = alsatica.
banatica, Nym. Consp. 559 = alsatica.
barbata, Brot. Fl. Lusit. i. 183 = foetida, major.
barbata, *Poir. in Lam. Encyc.* iv. 621.—Reg. Mediterr. occ.
barbata, Reichb. Pl. Crit. vii. 31 (excl. syn.) t. 208 = minor.
barbata, Sm. Engl. Bot. t. 2859 = Hederae.
Bartlingii, Griseb. Spicil. Fl. Rumel. ii. 57 = alsatica.
Benthami, Timb. in Mém. Acad. Toul. Sér. VII. vi. (1874) 643 = O. Rapum-genistae.
Berberidis, Facchini, ex Reut. in DC. Prod. xi. 23 = lucorum.
Berthelotii, *Webb & Berth. Phyt. Canar.* iii. 155.—Ins. Canar.
bicolor, C. A. Mey. in Ledeb. Fl. Alt. ii. 458 = cernua.
bicolor, Salzm. ex Moris, Fl. Sard. iii. 249 = sanguinea.
biflora, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21. (Quid?).—Java.
biflora, Nutt. Gen. Am. ii. 59 = Aphyllon uniflorum.
bipontina, F. Schultz, Beitr. 7; et in Flora, xii. (1829) 464 = caryophyllacea.
bohemica, *Čelak. in Sitzb. Boehm. Ges. Wiss.* (1874) 61.—Europ.
Boissieri, *Reichb. f. Ic. Fl. Germ.* xx. 118. t. 1838.—Algeria.
borealis, Reut. in DC. Prod. ix. 32 = caerulescens.
borealis, *Turcz. in Bull. Soc. Nat. Mosc.* (1841) 722.—Reg. Himal.; As. centr.
Borkhausenii, Andrz. ex Bess. in Flora, xv. (1832) II. Beibl. 19 = arenaria.
Bovei, Reut. in DC. Prod. xi. 24 = minor.
brachysepala, F. Schultz, in Arch. Fl. Fr. et Allem. 69, 89 = alsatica.
bracteata, *Viv. in Ann. Stor. Nat.* iv. (1830) 235.—Corsica.
bracteata, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. VII. 65 = reticulata.
bracteosa, Nym. Consp. 551 = O. Rapum-genistae.
breviflora, *F. Schultz, in Flora*, xxvi. (1843) 130.—Graecia.
Buekiana, Koch, Syn. Fl. Germ. ed. II. 619 = alsatica.
Buekii, A. Dietr. Fl. Boruss. iii. t. 145 = lutea.
Buhsei, Reut. in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 169 = O. Epithymum.

OROBANCHE :—
caerulea, Bory, & Chaub. ex Reut. in DC. Prod. xi. 7, 37 = lavandulacea.
caerulea, Fisch. ex Wallr. Orob. Gen. 57 = caerulescens.
caerulea, Moris, Elench. fasc. i. 35 = stricta.
caerulea, Plan. Ens. Fl. Gallega, 325 = amethystea.
caerulea, Vill. Hist. Pl. Dauph. ii. 406 = purpurea.
caerulescens, C. Koch, in Linnaea, xvii. (1843) 291 = connata.
caerulescens, *Steph. ex Willd. Sp. Pl.* iii. 349.—Europ.; As. bor.
caesia, *Guss. Fl. Sic. Syn.* ii. 138.—Europ.
Calendulae, *Pomel, Nouv. Mat. Fl. Atl.* 110.—Alger.
californica, Cham. & Schlecht. in Linnaea, iii. (1828) 134 = Aphyllon californicum.
Calotropidis, Edgew. in Hook. Journ. Bot. ii. (1840) 285 (*Calatropidis*) = Cistanche tubulosa.
camptolepis, *Boiss. & Reut. ex Boiss. Fl. Orient.* iv. 515.—Cilicia.
canescens, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 124 = caerulescens.
canescens, *Presl, Delic. Prag.* 72.—Sicilia.
canescens, F. Schultz, in Flora, xxvi. (1843) 129 = crenata.
Cannabios, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = seq.
Cannabis, Vauch. in Duby, Bot. Gall. i. 351 = ramosa.
capensis, Burm. f. Fl. Cap. Prod. 17 = seq.?
capensis, Thunb. Prod. Pl. Cap. 97 = Harveya capensis.
Cardui, Saut. in Flora, xxxiv. (1851) 49 = reticulata.
carlinoides, Miégev. in Bull. Soc. Bot. Fr. xii. (1865) 347 = reticulata.
Carotae, Des Moul. in Ann. Sc. Nat. Sér. II. iii. (1835) 78 = Picridis.
Carotae, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxvii. (1877) 394 = loricata.
caryophyllacea, C. A. Mey. Verz. Pfl. Cauc. 103 = loricata.
caryophyllacea, Quorund. ex Wallr. Orob. Gen. 46 = rubens.
caryophyllacea, F. Schultz, Beitr. Kenntn. Deutsch. Orob. 8 = gracilis.
caryophyllacea, *Sm. in Trans. Linn. Soc.* iv. (1798) 169.—Europ.; Oriens.
caryophyllacea, Strohm. ex Wallr. Orob. Gen. 39 = gracilis.
castellana, Reut. in DC. Prod. xi. 29 = amethystea.
caudata, *De Not. Prospett. Fl. Ligust.* 53 (sp. dub.).—Italia.
Centaureae-scabiosae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = major.
centaurina, Bertol. Fl. Ital. vi. 430 = Picridis.
cernua, *Loefl. It. Hisp.* 152.—Reg. Mediterr.; Oriens; Ind. or.; Austral.
cernua, Pall. Ind. Taur. ex Wallr. Orob. Gen. 61, partim = bicolor.
Cervariae, Kirschl. in Flora, xviii. (1835) 303; Suard, in Godr. Fl. Lorr. ed. I. ii. 180 = alsatica.
cervarihaerens, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
Chaboissaei, Lacroix, in Bull. Soc. Bot. Fr. vi. (1859) 565 = Hederae.
Chironii, *Lojac. Contrib. Fl. Sicil.* 12.—Sicil.
Chrysanthemi, F. Schultz, ex Reut. in DC. Prod. xi. 27 = pubescens.
cistanchoides, *Beck, in Denkschr. Acad. Wien*, l. (1885) 26.—Persia.
Cirsii, Fries, Mant. iii. 58 = reticulata.
Cirsii-oleracei, Casp. in Schrift. Phys. Ges. Koenigsb. ii. (1861) 46 = reticulata.
citrina, A. Dietr. Fl. Boruss. vii. t. 441 = caryophyllacea.
citrina, Nym. Consp. 558 = gracilis.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 326.—Reg. Himal.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 107.—Alger.
coccinea, Bieb. Casp. 58; Fl. Taur. Cauc. ii. 84 = Phelipaea coccinea.
coelestis, *Boiss. & Reut. ex Reut. in DC. Prod.* xi. 5, 37.—As. Min.
coerulea, Auct. vide *caerulea*.
coerulescens, Auct. vide *caerulescens*.
colorata, C. Koch, in Linnaea, xvii. (1843) 291 = alba.
Columbariae, Gren. & Godr. Fl. Fr. ii. 634 = minor.
commutata, Zumag. Fl. Pedem. i. 202 = ramosa?
comosa, Hook. Fl. Bor. Am. ii. 92 = Aphyllon comosum.
comosa, Schur, Enum. Pl. Transs. 505 = major.
comosa, Sprun. ex Reut. in DC. Prod. xi. 36 = alba.

OROBANCHE :—
comosa, Wallr. Sched. Crit. 314 = arenaria.
compacta, *Viv. Fl. Lib. Spec.* 34. t. 15. f. 2.—Afr. bor.
concolor, Duby, Bot. Gall. 350 = minor.
condensata, Ball, in Journ. Linn. Soc. xvi. (1878) 606 = tetuanensis.
condensata, Griseb. Spicil. Fl. Rumel. ii. 56 = minor.
condensata, Koch, ex Nym. Consp. 558 = gracilis.
condensata, Lojac. Crit. Orob. 35 = Sicil.
condensata, Moris, Stirp. Sard. ii. 8 = variegata.
confusa, F. Schultz, Fl. Pfalz (1846), 339, in syn.; in Flora, xxx. (1847), 66. t. 2. f. 2 = major.
congesta, Reichb. f. Ic. Fl. Germ. xx. 117. t. 1835 = crenata.
connata, *C. Koch, in Linnaea*, xxii. (1849) 669 (sp. dub.).—Armenia.
crassistyla, F. Schultz, in Flora, xxvi. (1843) 131 = alba.
crenata, *Forsk. Fl. Aegypt. Arab.* 113.—Arabia.
crinita, Benth. Cat. Pl. Pyr. 109 = O. Rapum-genistae.
crinita, Reichb. Pl. Crit. vii. f. 922 = densiflora.
crinita, Viv. Fl. Cors. Sp. Nov. 11 = sanguinea.
Crithmi, Bertol. Fl. Ital. vi. 424 = seq.
Crithmi-maritimi, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = minor.
cruenta, Bertol. Rar. It. Pl. Dec. iii. 56 = gracilis.
cruenta, Boiss. Voy. Bot. Espagne, ii. 475 = reticulata.
cumana, Guss. Suppl. Fl. Sic. Prod. 206 = cernua.
cumana, *Wallr. Orob. Gen.* 58.—Ross. austr.; Reg. Caucas.
cuprea, *Boiss. & Bal. ex Boiss. Fl. Orient.* iv. 510.—Cilicia.
curvata, *Pomel, Nouv. Mat. Fl. Atl.* 111.—Alger.
curviflora, Viv. Pl. Aegypt. Dec. 22 = cernua.
Cypria, *Reut. in Unger & Kotschy, Cypern*, 294.—Ins. Cyprus.
Cytisi, Duby, Bot. Gall. 349 = O. Rapum-genistae.
Cytisi-laburni, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = variegata.
Cytisi-scoparii, Vauch. ex Duby, Bot. Gall. 349 = O. Rapum-genistae.
decora, Moris, ex Reut. in DC. Prod. xi. 26 = denudata.
Delilei, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 201 = aegyptiaca.
densiflora, Moris, Fl. Sard. iii. 249 = minor.
densiflora, *Salzm. ex Reut. in DC. Prod.* xi. 19.—Reg. Mediterr. occ.
denudata, *Moris, App. ad Stirp. Sard. Elench.* 1.—Ins. Sardoa.
Deucalion, Reichb. f. Ic. Fl. Germ. xx. 92. t. 1527 = reticulata.
diaphana, F. Schultz, in Flora, xxvi. (1843) 130 = alba.
Dorycnii-suffruticosi, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = gracilis.
Echinopis, Panč. in Oestr. Bot. Zeitschr. xviii. (1868) 80 = major.
elatior, Auct. ex Wallr. Orob. Gen. 34 = minor.
elatior, DC. Fl. Fr. iii. 490 = amethystea.
elatior, A. Dietr. Fl. Boruss. t. 148 = alsatica.
elatior, Fluegge, ex Wallr. Orob. Gen. 42 = reticulata.
elatior, Hornem. in Fl. Dan. t. 1338, excl. syn. = O. Rapum-genistae.
elatior, Koch & Ziz, Cat. Pl. Pal. 12; & Reichb. Pl. Crit. f. 901, 902 = lutea.
elatior, Schleich. ex Reichb. Fl. Germ. Excurs. 355 = loricata.
elatior, Sutton, in Trans. Linn. Soc. iv. (1798) 178. t. 17 = major.
electra, D. Dietr. Syn. Pl. iii. 624 = alsatica.
epithymoides, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 28 = alba.
epithymophyta, St. Lag. in Soc. Bot. Lyon, vii. (1880) 131 = seq.
Epithymum, DC. Fl. Fr. iii. 490 = alba.
erubescens, Saut. in Flora, xxv. (1852) 577 = caryophyllacea.
Eryngii, Duby, Bot. Gall. 350 = amethystea.
Eryngii-campestris, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = praec.
Esulae, *Panč. Add. Fl. Princ. Serb.* 194 = Serbia.
euglossa, *Reichb. f. Ic. Fl. Germ.* xx. 117. t. 217.—Byzant.
fasciculata, Nutt. Gen. Am. ii. 59 = Aphyllon fasciculatum.
ferruginea, C. Koch, in Linnaea, xvii. (1843) 291 = alba.
flava, *Mart. ex F. Schultz, Beitr. Kenntn. Deutsch. Orob.* 9.—Europ.

OROBANCHE:—
flavidiflora, Guss. Fl. Sic. Syn. ii. 11. 748 = canescens.
foetida, Bivona, Stirp. Rar. Sic. Manip. i. n. = variegata.
foetida, Duby, Bot. Gall. i. 347 = O. Rapum-genistae.
foetida, Klotzsch, in Reise Princ. Waldem. Bot. 111 = Hansii.
foetida, Lapeyr. Hist. Abr. Pl. Pyr. i. 358 = gracilis, O. Rapum-genistae.
foetida, *Poir. Voy. Barb.* ii. 195; *Desf. Fl. Atlant.* ii. 59. t. 144.—Reg. Mediterr. occ.
foetida, Schimp. ex Reut. in DC. Prod. xi. 18 = variegata.
foetida, Viv. Fl. Lib. Spec. 34 = crenata.
forojuliensis, Nym. Consp. 560 = major.
Fraasii, F. Schultz, in Flora, xxvi. (1843) 127 = lavandulacea.
fragrans, Griseb. Spicil. Fl. Rumel. ii. 58 = pubescens.
fragrans, Koch, in Mert. & Koch, Deutschl. Fl. iv. 463 = major.
fragrantissima, Bertol. Fl. Ital. vi. 486 = lutea.
Freynii, Nym. Consp. 561 = Picridis.
Froelichii, Reichb. f. Ic. Fl. Germ. xx. 106 = flava.
fuliginosa, *Reut. ex Jord. Obs. Pl. Crit.* iii. (1846) 225. —Gallia; Graecia.
fusca, Lam. Encyc. iv. 621 = foetida.
galeopsiflora, Zumag. Fl. Pedem. i. 197 (Quid?) = Italia.
Galii, Duby, Bot. Gall. 349 = caryophyllacea.
Galii, Klotzsch, Reise Princ. Waldem. Bot. 111 = Solmsii.
Galii-molluginis, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = *Galii*, Duby.
galionepiphyta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
gallica, Gren. in Acad. Sc. Besanç. Séances Publ. (1838) 144 = cernua.
gamosepala, *Reut. in DC. Prod.* xi. 15.—Reg. Caucas.
Genistae, Stokes, Bot. Mat. Med. iii. 407 = major.
Genistae-cinereae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = praec.
Genistae-germanicae, F. Schultz, l. c. = praec.
Genistae-sagittalis, F. Schultz, l. c. = praec.
Genistae-tinctoriae, F. Schultz, l. c. = praec.
gilva, A. Dietr. Fl. Boruss. vii. t. 439 = caryophyllacea.
glaberrima, Guss. ex Reut. in DC. Prod. xi. 719 = Hederae.
glabra, Gaud. ex Reichb. f. Ic. Fl. Germ. xx. 103 = minor.
glabra, Hook. Fl. Bor. Am. ii. 91 = Boschniakia glabra.
glabrata *C. A. Mey. in Eichw. Pl. Casp.* 16. t. 17.—Reg. Caucas.; Persia.
glaucantha, Trautv. in Act. Hort. Petrop. i. (1871) 26 = cumana.
gracilis, Bess. ex Ledeb. Fl. Ross. iii. 318 = reticulata.
gracilis, *Sm. in Trans. Linn. Soc.* iv. (1798) 172.—Europ.; Reg. Caucas.
gracilis, Welw. ex Nym. Consp. 562 = minor.
grandiflora, Bory & Chaub. Nouv. Fl. Pélop. 40. t. 23 = crenata.
grandiflora, Presl, Fl. Sicul. i. 34 = variegata.
grandisepala, *F. Schultz, in Flora*, xxvi. (1843) 128.—Graecia.
grandiuscula, Moris, Diagn. Stirp. Sard. (1857) 2; et Fl. Sard. iii. 246 = gracilis.
Grenieri, Nym. Consp. 562 = minor.
Grenieri, F. Schultz, in Flora, xxviii. (1845) 739 = cernua.
Grisebachii, Reut. in DC. Prod. xi. 28 = minor?
Gussonii, Nym. Consp. 558 = cruenta.
Haenseleri, *Reut. in DC. Prod.* xi. 22.—Hispan.
Hansii, *Kern. in Ber. Naturw. Ver. Innsbruck*, i. (1870) 111.—Reg. Himal.
Hederae, *Duby, Bot. Gall.* 350.—Europ.
Hederae-helicis, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = praec.
Helicis, Rota, Prospett. Piant. Pav. 287 = Hederae.
Hellebori, *Miégev. in Bull. Soc. Bot. Fr.* xii. (1865) 347. —Hispan.
hians, Reut. ex Boiss. Fl. Orient. iv. 508 = Teucrii.
hians, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 266, nomen; xxx. (1857) 354 = alba.
Hieracii-pilosellae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = Picridis.

OROBANCHE:—
hirta, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 21 (Quid?).—Java.
hispanica, Boiss. Voy. Bot. Espagne, ii. 476 = cernua.
Hookeriana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 605.—Marocc.
hyalina, Sprun. ex Reut. in DC. Prod. xi. 24 = minor.
hygrophila, Bruegg. in Sendtner, Veget. Suedbay. 836 = reticulata.
hymenocalyx, Reut. in DC. Prod. xi. 20 (sp. dub.).—Reg. Caucas.
Hyperici, Unger, Einfl. Bodens Verth. Gew. (1836) 321 = reticulata?
Hypopitys, Hill, Hort. Kew. 228 = Monotropa Hypopitys.
ianthina, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 227.—As. centr.
Illicis, Haensel. in Bot. Zeit. iv. (1846) 313, sphalm. = *Ulicis*.
incurva, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = caryophyllacea.
indica, Buch.-Ham. ex Roxb. Hort. Beng. 45; Fl. Ind. iii. 27 = aegyptiaca.
indica, Spreng. Syst. ii. 817 = Striga orobanchoides.
indica, Wall. Cat. n. 3966 = cernua.
insignis, Clarke, Trav. iii. 42 (Cistanches sp.).—Aegypt.
interrupta, *Pers. Syn.* ii. 181.—Afr. austr.
ionantha, *Kern. in Oestr. Bot. Zeitschr.* xxiv. (1874) 236.—Europ.
kashmirica, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 324.—Reg. Himal.
Klugei, Schmitz & Regel, Fl. Bonn. 190 = minor.
Knappii, Pant. in Verh. Ver. Natur- u. Heilk. Presb. ii. (1871–72) 73 = amethystea.
Kochii, F. Schultz, in Flora, xxx. (1847) 66 = major.
Kotschyi, *Reut. in DC. Prod.* xi. 33.—Persia.
Krausei, A. Dietr. Pl. Boruss. vii. t. 437 = caryophyllacea.
Krylowi, G. Beck, in Oestr. Bot. Zeitschr. xxxi. (1881) 309 = major.
kurdica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 505.—Persia.
labellata, Dum. Fl. Belg. 33 = gracilis.
lactea, Eckl. & Zeyh. ex Presl, Bot. Bemerk. 91 = Harveya lactea.
laevis, Geners. ex Wallr. Orob. Gen. 36 = caryophyllacea.
laevis, Lam. Fl. Fr. ii. 327 = caerulea.
laevis, Linn. Sp. Pl. 632 = purpurea, arenaria.
Lamyana, De Cassac, ex Nym. Consp. 562 = amethystea?
Laserpitii, Reut. ex Jord. Obs. Pl. Crit. iii. (1846) 223 = seq.
Laserpitii-sileris, *Reut. in DC. Prod.* xi. 25.—Helvetia.
latepubescens, Guss. Fl. Sic. Syn. ii. 749 = caryophyllacea.
latisquama, Reut. ex Boiss. Fl. Orient. iv. 516 = camptolepis.
laurina, Bonap. ex Bertol. Fl. Ital. vi. 424 = minor.
laurina, Reichb. f. Ic. Fl. Germ. xx. 104. t. 181 = Hederae.
lavandulacea, Hohen. ex Reut. in DC. Prod. xi. 8 = aegyptiaca.
lavandulacea, *Reichb. Pl. Crit.* vii. 48. f. 935.—Reg. Mediterr.
laxa, Pomel, Nouv. Mat. Fl. Atl. 109 (sp. dub.).—Alger.
laxiflora, Reichb. Pl. Crit. vii. 39 = caryophyllacea.
lazica, Reut. ex Boiss. Fl. Orient. iv. 508 = Teucrii.
Leonuri, Rota, Prosp. Piant. Pav. 270, 287 = minor.
leptantha, Pomel, Nouv. Mat. Fl. Atl. 110 (sp. dub.). —Alger.
leucantha, Griseb. Spicil. Fl. Rumel. ii. 57 = reticulata.
leucantha, Reichb. Ic. Fl. Germ. xx. 97. t. 167 = alsatica.
leucopogon, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 504.—Cataonia.
Levieri, Lojac. Contrib. Fl. Sic. 14 = gracilis.
Libanotidis, Rupr. Beitr. Pfl. Russl. iv. 70 = alsatica.
Ligustri, Suard, ex Godr. Fl. Lorr. ed. I. ii. 178 = caryophyllacea.
lilacina, F. Schultz, in Flora, xxvi. (1843) 129 = Picridis.
litorea, Guss. Fl. Sic. Prod. ii. 184 = minor.
livida, Sendtn. ex Reichb. f. Ic. Fl. Germ. xx. 104 = minor.

OROBANCHE:—
Lobelii, Noul. Fl. Sous-Pyr. 481 = gracilis.
longiflora, Pers. Syn. ii. 181 = aegyptiaca.
loricata, *Reichb. Pl. Crit.* vii. 41.—Europ.; Reg. Caucas.
Loti-corniculati, *F. Schultz, in Ann. Gew. Regensb.* v. (1830) 500 (sp. dub.).—Europ.
Loti-cytisoidis, F. Schultz, l. c. = sanguinea.
lucorum, *A. Br. ex F. Schultz, l. c.* 504.—Bavaria.
ludoviciana, Nutt. Gen. Am. ii. 58 = Aphyllon ludovicianum.
lutea, *Baumg. Enum. Stirp. Transs.* ii. 215.—Europ.
lutea, Eckl. & Zeyh. ex Steud. Nom. ed. II. ii. 231 = Harveya lutea.
lutea, Lojac. in Natural. Sicil. i. (1882) 256 = variegata.
lutea, Nym. Syll. 134 = Cistanche lutea.
lycica, *F. Schultz, in Flora*, xxvi. (1843) 128.—Graecia.
macrantha, A. Dietr. Fl. Boruss. vii. t. 438 = caryophyllacea.
macrolepis, *Coss. in Ann. Sc. Nat.* Sér. III. ix. (1848) 146.—Hispan.
macrolepis, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 356 = pycnostachyum.
macrosepala, F. Schultz, in Arch. Fl. Fr. et Allem. 70 = minor.
major, Auct. ex Wallr. Orob. Gen. 59 = comosa.
major, Braun, ex Wallr. l. c. 46 = lutea.
major, DC. Fl. Fr. iii. 488 = Rapum-genistae.
major, Duby, Bot. Gall. ii. 349 = gracilis.
major, Gorter. Fl. Ingr. 100 = alsatica.
major, Langsd. ex Wallr. Orob. Gen. 44 = foetida.
major, *Linn. Sp. Pl.* 632.—Europ.
major, Loefl. It. Hisp. 151 = Hederae.
major, Pollich, Hist. Pl. Palat. ii. 200 = caryophyllacea.
major, Pourr. ex Costa, Cat. Fl. Catal. 191 = sanguinea.
major, Wolf. ex Wallr. Orob. Gen. 48 = alba.
marginata, Reut. ex Boiss. Fl. Orient. iv. 508 = Teucrii.
Mathenii, F. Schultz, in Bot. Zeit. ii. (1844) 857 = reticulata.
Mathesii, Nym. Consp. 559 = praec.
media, Desf. Fl. Atlant. ii. 59 = cernua.
Medicaginis, Duby, Bot. Gall. 349 = lutea.
Medicaginis, Reichb. Pl. Crit. vii. 41. f. 919 = Hederae.
Medicaginis-sativae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = praec.
micrantha, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 235 = Salviae.
micrantha, Wallr. Sched. Crit. 315 = ramosa.
Millefolii, Steud. Nom. ed. II. ii. 231 = caerulea.
minor, Facch. Fl. Suedtir. 76 = Hederae.
minor, *Sutt. in Trans. Linn. Soc.* iv. (1798) 179; *Sm. Engl. Bot.* t. 422.—Europ.; Afr. bor.
minor, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
minor, Vahl, in Fl. Dan. t. 1219 = loricata.
Morisiana, Nym. Syll. Suppl. 24 = Muteli.
multiflora, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 179 = Aphyllon multiflorum.
Muteli, A. Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 48, 49 = ramosa.
Muteli, *F. Schultz, in Mutel, Fl. Fr.* ii. 353. t. 43.—Europ.; Oriens.
mutila, Dum. Fl. Belg. 33 = cernua?
nana, *Noé, ex Reut. in DC. Prod.* xi. 9.—Reg. Mediterr.
nebrodensis, Tineo, in Guss. Fl. Sic. Syn. ii. 845 = sanguinea.
neottioides, Saut. in Flora, xxxv. (1852) 578 = Salviae.
Nicotianae, Wight, Illustr. ii. 179. t. 158 = cernua.
Nočana, Nym. Consp. 558 = gracilis.
nuda, Koch, ex Wallr. Orob. Gen. 48 = alba.
nudiflora, Wallr. Sched. Crit. i. 310 = minor.
olbiensis, Nym. Syll. 133 = nana.
ombrochares, Hance, in Journ. Linn. Soc. xiii. (1873) 84 = coerulescens.
Ozanonis, F. Schultz, Arch. Fl. Fr. 358 (1866) = Salviae.
palaestina, Boiss. Fl. Orient. v. 506, partim = Grisebachii.
palaestina, Reut. in DC. Prod. xi. 718 = minor.
palatina, F. Schultz, Arch. Fl. Fr. 161 = O. Rapum-genistae.
pallens, F. Schultz, in Flora, xiv. (1840) 128 = Picridis.
pallida, F. Schultz, l. c. = praec.

OROBANCHE:—
pallidiflora, Wimm. & Grab. Fl. Siles, ii. 233 = reticulata.
palustris, Bruegg. in Flora, xxxviii. (1855) 34 = reticulata.
papyracea, DC. ex F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = caryophyllacea.
paradoxa, Koch, ex Wallr. Orob. Gen. 48 = alba.
Pareysi, G. Beck, in Oestr. Bot. Zeitschr. xxxi. (1881) 310 = purpurea.
parviflora, E. Mey. ex Steud. Nom. ed. II. i. 723, in syn.; ii. 231 = Harveya parviflora.
parviloba, *F. Schultz, in Flora*, xxvi. (1843) 128 (Quid?).—Graecia.
pedunculata, Roxb. Hort. Beng. 45; Fl. Ind. iii. 29 = Aeginetia pedunculata.
pedunculata, Viv. Pl. Aegypt. Dec. 22 = aegyptiaca.
Pelargonii, *Caldesi, in Nuov. Giorn. Bot.* xii. (1880) 175.—Italia.
Peisonis, G. Beck, in Halác. & Braun, Nachtr. Fl. Nieder-Oestr. 121 = caesia.
Petasites, Borb. in Oestr. Bot. Zeitschr. xxxv. (1885) 76, nomen = flava.
Phelipaea, Wallr. ex Reut. in DC. Prod. xi. 38 = Cistanche violacea (in Addend.).
Phelypaea, Willd. Sp. Pl. iii. 352 = Phelypaea foliata.
picridiphyta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = seq.
Picridis, *F. Schultz, in Ann. Gew. Regensb.* v. (1830) 504.—Europ.
Picridis-hieracioidis, F. Schultz, l. c. 500 = praec.
pinetorum, S. Wats. in Bot. Calif. i. 585, sphalm. = seq.
pinorum, Geyer, ex Hook. Kew Journ. iii. (1851) 297 = Aphyllon pinetorum.
platylepis, Reut. ex Boiss. Fl. Orient. iv. 516 = camptolepis.
platystigma, Reichb. Fl. Germ. Excurs. 353 = gracilis.
platystigma, Reichb. Pl. Crit. vii. 31 = reticulata.
pogonanthera, Reut. in DC. Prod. xi. 33 = cernua.
polymorpha, Schrank, ex Steud. Nom. ed. II. ii. 231 = spp. europaeae omnes.
pratensis, Eckl. & Zeyh. ex Presl, Bot. Bemerk. 91 = Harveya pratensis.
procera, Koch, in Mert. & Koch, Deutschl. Fl. iv. 438 = reticulata.
pruinosa, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 87 = crenata.
psila, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 327.—Reg. Himal.
psilandra, C. Koch, in Linnaea, xxii. (1849) 668 = reticulata.
pubescens, Gilib. Fl. Lituan. i. 129 = major.
pubescens, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 332.—Europ.; Oriens.
pumila, Graaf, ex Reut. in DC. Prod. xi. 29; Koch & Noé, in Reichb. f. Ic. Fl. Germ. xx. 104 = minor, *Sutt.*
punctata, F. Schultz, in Flora, xxvi. (1843) 131 = alba.
purpurascens, J. F. Gmel. Syst. 954 = arenaria.
purpurea, *Jacq. Enum. Stirp. Vindob.* 108, 252.—Europ.
purpurea, Linn. f. Suppl. 228; Thunb. Prod. Pl. Cap. 97 = Harveya purpurea.
purpurea, Rafin. Caratt. 80. n. 196 = variegata.
pycnostachya, *Hance, in Journ. Linn. Soc.* xiii. (1873) 84.—China.
pyrrha, Reichb. f. Ic. Fl. Germ. xx. 105 = minor.
quadrifida, C. Koch, in Linnaea, xxii. (1849) 665 = caryophyllacea.
quadrivalvis, Regel, in Act. Hort. Petrop. vi. (1879) 354 = caerulescens.
racemosa, Schleich. Cat. Pl. Helv. (1821) 25 = minor, Hederae.
racemosa, Steph. ex C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 157 = Boschniakia glabra.
ramosa, Delile, ex Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 201 = aegyptiaca.
ramosa, Duby, Bot. Gall. 351 = lavandulacea, Muteli.
ramosa, *Linn. Sp. Pl.* 633.—Europ.; Kashmir.
ramosa, Thunb. Prod. Pl. Cap. 97 = interrupta.
Rapum, Reichb. Pl. Crit. vii. 43 = seq.
Rapum-genistae, *Thuill. Fl. Par.* ed. II. 317.—Europ.
Reichardiae, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxvii. (1878) 392 = Picridis.

OROBANCHE:—
reticulata, *Wallr. Orob. Gen.* 42.—Reg. Mediterr. occ.
Reuteri, F. W. Schultz, ex Reut. in DC. Prod. xi. 24 = minor.
rhizopus, Wallr. ex Ledeb. Fl. Ross. iii. 314 (= *Phelypaea salsa*).—Rossia.
rigens, *Loisel. Fl. Gall.* ed. I. 384.—Corsica.
Ritro, Gren. & Godr. Fl. Fr. ii. 635 = major.
robusta, A. Dietr. Fl. Boruss. iii. t. 151 = arenaria.
rossica, Cham. & Schlecht. in Linnaea, iii. (1828) 132 = Boschniakia glabra.
rubens, Wallr. Sched. Crit. 307 = lutea.
Rubi, Duby, Bot. Gall. 350 = lucorum.
Rubi-fruticosi, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = praec.
rubiginosa, A. Dietr. Fl. Boruss. iii. t. 252 = alba.
rubra, Rafin. Chlor. Aetn. 10 = variegata?
rubra, Sm. Engl. Bot. t. 1786; Hook. Fl. Scot. 19 = alba.
rufescens, Nym. Syll. 133 = Muteli.
sabulicola, Lojac. in Natural. Sicil. ii. (1883) 61 = fuliginosa.
Salisii, Req. ex Coss. Notes Crit. 9 = minor.
Salviae, *F. Schultz, in Ann. Gew. Regensb.* v. (1830) 505.—Europ.
salvihaerens, St. Lag. Cat. Pl. Rhône, 611 = praec.
sambucina, Janka, in Termész. Füzetek, viii. (1884) 29; et in Oestr. Bot. Zeitschr. xxxiv. (1884) 336 = reticulata.
sanguinea, *Presl, Delic. Prag.* 71.—Sicilia.
Santolinae, *Losc. & Pard. Ser. Inconf.* 79.—Hispan.
sarothamnophyta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = O. Rapum-genistae.
Satureiae-montanae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = alba.
Satyrus, De Not. Prospett. Fl. Ligust. 35 = variegata.
Sauteri, F. Schultz, in Flora, xxxiv. (1851) 49 = reticulata.
Scabiosae, Koch, in Mert. & Koch, Deutschl. Fl. iv. 440 = reticulata.
Scabiosae-columbariae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = minor, *Sutt.*
scabioshaerens, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = *Scabiosae.*
scarlatina, E. Mey. ex Steud. Nom. ed. II. i. 723, in syn.; ii. 231 = Harveya scarlatina.
Schultzii, *Mutel, Fl. Fr.* ii. 352.—Reg. Mediterr.; As. Min.
Scolymi, *Pomel, Nouv. Mat. Fl. Atl.* 108 (sp. dub.).—Alger.
segetum, Sprun. in Flora, xxvi. (1843) 129 = crenata.
serotina, Kirschl. in Pollichia, iii. (1845) 14 = reticulata.
Serpylli, Vauch. ex Des Moul. in Ann. Sc. Nat. Sér. II. iii. (1835) 76 = alba.
sicula, Lojac. in Natural. Sicil. i. (1882) 255 = variegata.
sicula, *Spenn. ex Walp. Rep.* iii. 463.—Sicilia.
Solmsii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 325.—Reg. Himal.
sordida, C. A. Mey. in Ledeb. Fl. Alt. ii. 455 = lutea.
sparsiflora, Wallr. Sched. Crit. i. 309 = alba.
Spartii, Guss. Fl. Sic. Prod. 182 = variegata.
Spartii-juncei, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = gracilis.
speciosa, DC. Fl. Fr. Suppl. 393 = crenata.
speciosa, A. Dietr. Fl. Boruss. t. 149 = reticulata.
speciosa, Moris, Stirp. Sard. Elench. fasc. i. 35 = condensata.
speciosa, Salis, in Flora, xvii. (1834) II. Beibl. 20 = minor.
spectabilis, Benth. ex E. Mey. in Drège, Zwei Pfl. Docum. 205, nomen = Harveya spectabilis.
spectabilis, Reut. in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 169 = Kotschyi.
Spruneri, Lojac. in Natural. Sicil. i. (1882) 215 = Todari.
Spruneri, F. Schultz, in Flora, xxvi. (1843) 130 = gracilis.
squalida, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 354 = alba.
squamaria, Fisch. ex Ledeb. Fl. Ross. iii. 314 (= *Phelypaea salsa*).—Rossia.
squamata, G. Don, Gen. Syst. iv. 633 = seq.
squamosa, Thunb. Prod. Pl. Cap. 97 = Harveya squamosa.
stans, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 (Quid?).—Java.

OROBANCHE:—
stenantha, Lojac. in Natural. Sicil. ii. (1883) 62 = Hederae.
stigmatodes, Wimm. Fl. Schles. (1841) 280 = major.
Stocksii, Boiss. Fl. Orient. iv. 505 = cistanchoides.
striata, Reut. in DC. Prod. xi. 28 = minor?
stricta, Moris, ex Bertol. Fl. Ital. vi. 450 = Schultzii.
strigosa, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 604 = variegata.
strobiligena, Reichb. Pl. Crit. vii. 38 = caryophyllacea.
strumosa, Rogow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 83, nomen = alsatica.
subacaulis, Benth. ex Walp. Ann. iii. 208 = Christisonia subacaulis.
subverticillata, *F. Schultz, in Flora*, xxvi. (1843) 128 (Quid?).—Graecia.
superba, De Not. Prospett. Fl. Ligust. 35 = variegata.
superficialis, Griseb. ex Reut. in DC. Prod. xi. 28 = pubescens.
sylvatica, F. Schultz, in Flora, xxiii. (1840) 128 = Salviae.
tetuanensis, *Ball, in Journ. Linn. Soc.* xvi. (1878) 606.—Marocc.
Teucrii, *Holand. Fl. Moselle*, ed. I. ii. 322; *F. Schultz, in Flora*, xviii. (1835) 200.—Europ.; As. Min.
teucrionepiphyta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
thapsoides, Lojac. in Natural. Sicil. ii. (1883) 60 = pubescens.
Thymi-serpylli, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = alba.
Thymi-vulgaris, F. Schultz, l. c. = alba.
thyrsoidea, Moris, ex Bertol. Fl. Ital. vi. 428 = rigens.
tinctoria, Forsk. Fl. Aegypt. Arab. 112 = Cistanche lutea.
tinctoria, Willd. Sp. Pl. iii. 353 = Cistanche lutea.
Todari, *Lojac. in Natural. Sicil.* i. (1882) 215.—Sicilia.
Tommasinii, Reichb. f. Ic. Fl. Germ. xx. 92 = Picridis.
torquata, Reichb. Pl. Crit. ix. 12. f. 1128 = caryophyllacea.
transsilvanica, F. Porc. Fl. Phan. Naseud. (1881) 73 = alba.
Trifolii-pratensis, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = minor.
Trotzkii, *Wagn. ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 64 (Quid?).—Rossia.
tubata, E. Mey. ex Benth. in DC. Prod. x. 505 = Harveya tubata.
tuberosa, Hook. Fl. Bor. Am. ii. 92 = Boschniakia Hookeri.
tubiflora, A. Dietr. Fl. Boruss. iii. t. 147 = caryophyllacea.
tulbaghensis, Eckl. & Zeyh. ex Presl, Bot. Bemerk. 91 = Harveya tulbaghensis.
tussilaginis, Mutel, Fl. Fr. ii. 349 = flava.
uitenhagensis, Eckl. ex Hook. Ic. Pl. sub t. 351 = Harveya purpurea.
Ulicis, Des Moul. in Ann. Sc. Nat. Sér. II. iii. (1835) 71 = gracilis?
Ulicis, Haensel. ex Willk. in Bot. Zeit. iv. (1846) 313; Reut. in DC. Prod. xi. 720 = gracilis.
Ulicis-europaeae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = gracilis.
unicolor, Bor. Fl. Centr. Fr. ed. III. ii. 501 = minor.
uniflora, Linn. Sp. Pl. 633 = Aphyllon uniflorum.
vagabunda, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = pubescens.
varia, E. Mey. ex Steud. Nom. ed. II. i. 723, in syn.; ii. 231 = Harveya varia.
variegata, *Wallr. Orob. Gen.* 40.—Europ. med.; Marocc.
Vaucherii, Noulet, Fl. Sous-Pyr. 487 = Hederae.
versicolor, F. Schultz, in Flora, xxvi. (1843) 129 = pubescens.
versicolor, Welw. Fl. Lusit. Excurs. n. 398 = gracilis.
Viciae-fabae, F. Schultz, in Ann. Gew. Regensb. v. (1830) 500 = crenata.
villiflora, Koch, ex Reut. in DC. Prod. xi. 27 = pubescens.
villosa, F. Schultz, ex Guss. Enum. Pl. Inar. 218 = pubescens.
villosiflora, F. Schultz, in Flora, xxviii. (1845) 737, 740 = pubescens.
violacea, Wallr. Orob. Gen. 70 = Cistanche violacea (in Addend.).
virginiana, Linn. Sp. Pl. 633 = Epifagus americanus.
virginica, Hill, Veg. Syst. xviii. 71 = praec.

OROBANCHE:—
viridis, *Hazlin. in Verh. Zool. Bot. Ges. Wien*, ii. (1852) *Abh.* 3, nomen (Quid ?).—Europ.
Vitalbae, Bertol. Fl. Ital. vi. 441 = pubescens.
vulgaris, Gaud. Fl. Helv. iv. 176 = gracilis.
vulgaris, Poir. in Lam. Encyc. iv. 621 = caryophyllacea.
Weinmanni, Steud. Nom. ed. II. ii. 231 = reticulata.
Welwitschii, Nym. Consp. 562 = gracilis.
Wiedemanni, *Boiss. Fl. Orient.* iv. 510.—Anatolia.
Wierzbickii, F. Schultz, in Flora, xxviii. (1845) 324 = gracilis.
Yuccae, Savi f. ex Bertol. Fl. Ital. vi. 423 = minor.

OROBANCHE, Vell. Fl. Flum. 254 (1825); vi. tt. 60–79 (1827) = **Gesneria**, Linn. etc.
brasiliensis, Vell. l. c. t. 63 (Besleriae sp. ?).—Bras.
carnosa, Vell. l. c. 255; vi. t. 61 (Besleriae sp. ?).—Bras.
cernua, Vell. l. c. 258; vi. t. 71 = Sinningia speciosa.
crenata, Vell. l. c. 254; vi. t. 60.—Bras.
ferox, DC. Prod. vii. 541 = *pedunculata*.
fissa, Vell. Fl. Flum. 257; vi. t. 69 (Hypocyrtae sp.).—Bras.
fluminensis, Vell. l. c. 255; vi. t. 62 (Besleriae sp. ?).—Bras.
fornix, Vell. l. c. 259; vi. t. 75 = G. allagophylla.
hirta, Vell. l. c. 261; vi. t. 79 = G. allagophylla.
melancholica, Vell. l. c. 260; vi. t. 77 (Hypocyrtae sp. ?).—Bras.
monanthera, Vell. l. c. 256; vi. t. 66 (Hypocyrtae sp. ?).—Bras.
pedunculata, Vell. l. c. 258; vi. t. 73 (Hypocyrtae sp. ?).—Bras.
perianthomega, Vell. l. c. 257; vi. t. 68 = Alloplectus dichrous.
petraea, Vell. l. c.; vi. t. 60.—Bras.
racemosa, Vell. l. c. 258; vi. t. 72 = Rytidophyllum prasinatum.
serpens, Vell. l. c. 257; vi. t. 67 (Hypocyrtae sp. ?).—Bras.
spicata, Vell. l. c. 260; vi. t. 78 = G. allagophylla.
tuberosa, Vell. l. c. 257; vi. t. 70 (Gloxiniae sp. ?).—Bras.
tubulosa, Vell. l. c. 259; vi. t. 76 = G. Douglasii.
umbellata, Vell. l. c. 256; vi. t. 65 = G. confertiflora.
ventricosa, Vell. l. c. 259; vi. t. 74 (Besleriae sp. ?).—Bras.
verticillata, Vell. l. c. 255; vi. t. 64 = G. maculata.

OROBANCHIA, Vand. Fl. Lusit. 41. ff. 18, 19 (1788) = **Alloplectus**, Mart. (Gesner.).
lanceolata, Poir. Encyc. Suppl. iv. 202.—Bras.
radicans, Poir. l. c. 203.—Bras.

OROBELLA, Presl, in Weitenw. Beitr. ii. (1837) 27. t. 4 = **Vicia**, Tourn. (Legumin.).
vicioides, Presl, l. c. 28 = V. dumetorum.

OROBIUM, Reichb. Consp. 185 (1828) = **Braya**, Sternb. & Hoppe (Crucif.).

OROBIUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 164 (1842) = **Lagenocarpus**, Nees (Cyperac.).
adamantinum, Schrad. ex Nees, l. c. 165 = L. adamantinus.

OROBOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = OROBUS, Tourn.
piscidius, St. Lag. l. c. = *Orobus Piscidia*.

OROBUS, [Tourn.] Linn. Syst. ed. I (1735) = **Lathyrus**, Tourn. (Legumin.).
acutangulus, Wender. ex Steud. Nom. ed. II. ii. 232 = Vicia amoena ?
aestivalis, Schur, in Oestr. Bot. Zeitschr. x. (1860) 325 = *venetus*.
alatus, A. Br. Ind. Sem. Hort. Berol. (1853) 13 = Lathyrus alatus.
alatus, Maxim. Prim. Fl. Amur. 83.—Manchuria.
albus, Linn. f. Suppl. 327 = Lathyrus pannonicus.
alpestris, Waldst. & Kit. Pl. Rar. Hung. ii. 133. t. 126 = Lathyrus canescens ?
americanus, Mill. Gard. Dict. ed. VIII. n. 9 = Abrus precatorius.
angulatus, Philippe, Fl. Pyren. i. 266 = Lathyrus angulatus.
angustifolius, Linn. Sp. Pl. 729.—Sibir.
angustifolius, Vill. Fl. Delphin. 75 = Lathyrus canescens.
anomalus, C. Koch, in Linnaea, xxiv. (1851) 96 = Vicia truncatula.

OROBUS :—
argenteus, Mill. Gard. Dict. ed. VII. n. 10.—Mexic.
aristatus, Lapeyr. Hist. Abr. Pyr. Suppl. 106 = Vicia Orobus.
armenus, Boiss. & Huet, Diagn. Ser. II. ii. 43.—Armenia.
articulatus, A. Br. Ind. Sem. Hort. Berol. (1853) 13 = Lathyrus Clymenum.
asphodeloides, Gouan, Illustr. 48 = Lathyrus pannonicus.
atropurpureus, Desf. Fl. Atlant. ii. 157. t. 196 = Vicia sicula.
atropurpureus, Lapeyr. Hist. Abr. Pl. Pyr. 413 = Lathyrus canescens.
aurantius, Stev. ex Bieb. Fl. Taur. Cauc. iii. 462 = Vicia aurantia.
aureus, Stev. ex Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iii. 42 = Lathyrus montanus ?
austriacus, Crantz, Stirp. Austr. ed. II. fasc. v. 374 = Lathyrus pannonicus.
bimarginatus, Stokes, Bot. Mat. Med. iv. 36 = Lathyrus sativus.
bulbosus, Stokes, l. c. 38 = Lathyrus montanus.
californicus, Alef. in Bonplandia, ix. (1861) 146 = Lathyrus venosus ?
canescens, Linn. f. Suppl. 327 = Lathyrus canescens.
capensis, Spreng. Syst. iii. 260 = Lessertia pulchra.
caucasicus, Spreng. l. c. 261 = Vicia truncatula.
ciliatus, Alef. in Flora, xlv. (1862) 275 = Lathyrus ciliatus.
Clusii, Spreng. Syst. iii. 259 = Vicia oroboides.
Clymenum, A. Br. Ind. Sem. Hort. Berol. (1853) 13 = Lathyrus Clymenum.
coccineus, Mill. Gard. Dict. ed. VIII. n. 12; Lodd. Bot. Cab. t. 883 = Lathyrus sphaericus.
croceus, Desf. in Ann. Mus. Par. xii. (1808) 59. t. 9 = Vicia aurantia.
cyaneus, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 51 = Lathyrus cyaneus.
cyaneus, Wahlenb. in Berggr. Resor, ii. Bihang, 41 = *sessilifolius*.
Denhardtii, Tenore, ex Steud. Nom. ed. II. ii. 232 (Quid ?).—Italia.
diffusus, Nutt. in Fras. Cat. (1813) = Vicia americana.
digitatus, Bieb. Fl. Taur. Cauc. ii. 153 = *sessilifolius*.
dispar, Nutt. Gen. Am. ii. 95 = Astragalus multiflorus.
dissitifolius, Alef. in Bonplandia, ix. (1861) 145 = Vicia americana.
divaricatus, Lapeyr. in Mém. Mus. Par. ii. (1815) 302. t. 11 = Lathyrus montanus.
domingensis, Spreng. Syst. iii. 261.—Ins. S. Doming.
emodi, Wall. Cat. n. 5948 = Lathyrus luteus.
enervius, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 796 = *venosus*.
ensifolius, Lapeyr. in Mém. Mus. Par. ii. (1815) 303. t. 12 = Lathyrus canescens.
erectus, Poir. in Lam. Encyc. iv. 627 = Vicia truncatula.
Ewaldi, Meinsh. in Bull. Soc. Nat. Mosc. xli. (1868) 1. 354.—Ingria.
exaltatus, Tenore & Guss. Fl. Neap. Syll. App. v. 25.—Italia.
Faba, Brot. Fl. Lusit. ii. 147 = Vicia Faba.
filiformis, Lam. Fl. Fr. ii. 568 = Lathyrus canescens.
Fischeri, Sweet, Brit. Flow. Gard. Ser. I. iii. t. 289 = Vicia sicula.
flaccidus, Kit. ex Reichb. Fl. Germ. Excurs. 536 = *vernosus*.
flaccidus, Radius, ex DC. Prod. ii. 377 = Lathyrus vernus.
formosus, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 50 = Pisum formosum.
Friedrichsthalii, Griseb. Spicil. Fl. Rumel. ii. 498 = alpestris.
fruticosus, Pers. ex DC. Prod. ii. 264 = Coursetia tomentosa.
glaberrimus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 20 = Lathyrus montanus ?
glabratus, Nym. Consp. 204 = Lathyrus hirsutus.
Gmelini, Fisch. ex DC. Prod. ii. 378.—Sibir.
Gmelini, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. vii. 66 = Lathyrus pannonicus.
gracilis, Alef. in Bonplandia, ix. (1861) 141.—Chili.
gracilis, Gaud. Fl. Helv. iv. 500 = Lathyrus montanus.
graminifolius, Becker, Fl. Frankf. i. 408 = Lathyrus canescens.
grandiflorus, Boiss. Fl. Orient. ii. 622.—Syria.
hirsutus, Linn. Sp. Pl. 728 = Lathyrus hirsutus.
hispidulus, Alef. in Bonplandia, ix. (1861) 141 = Lathyrus inconspicuus.
Hookeri, Alef. l. c. 144 = Lathyrus magellanicus.
humilis, Ser. in DC. Prod. ii. 371 = Lathyrus altaicus.

OROBUS :—
inconspicuus, A. Br. Ind. Sem. Hort. Berol. (1853) 13 = Lathyrus inconspicuus.
incurvus, A. Br. l. c. = Lathyrus incurvus.
intermedius, C. A. Mey. in Ledeb. Fl. Alt. iii. 360 = Lathyrus heterophyllus ?
japonicus, Alef. in Bonplandia, ix. (1861) 143 = Vicia pallida.
Jordani, Griseb. Spicil. Fl. Rumel. i. 74 = Lathyrus canescens ?
Jordani, Tenore, Prod. Fl. Nap. App. v. 21.—Italia.
Kolenatii, C. Koch, in Linnaea, xxiv. (1851) 96 = Lathyrus montanus.
lacteus, Bieb. Fl. Taur. Cauc. ii. 152 = Lathyrus pannonicus.
laevigatus, Waldst. & Kit. Pl. Rar. Hung. iii. 270 = Lathyrus montanus ?
lathyroides, Habl. Taur. 125 = Lathyrus hirsutus.
lathyroides, Linn. Sp. Pl. 728 = Vicia unijuga.
lathyroides, Sibth. & Sm. Fl. Graec. Prod. ii. 63 = Vicia oroboides.
laxiflorus, Desf. in Ann. Mus. Par. xii. (1808) 57. t. 8 = Lathyrus hirsutus.
Lens, Stokes, Bot. Mat. Med. iv. 37 = Ervum Lens.
linifolius, Reich. ex Gaertn. Mey. & Scherb. Fl. Wett. iii. 25 = Lathyrus canescens.
linifolius, Reichb. ex Nym. Consp. 205 = Lathyrus montanus.
linifolius, Wender. in Linnaea, x. (1830) Litt. 55 = praec.?
littoralis, A. Gray, in Pacif. Rail. Rep. xii. ii. 58 = Lathyrus littoralis, *Endl.*
longifolius, Kit. in Linnaea, xxxii. (1863) 632 = *pauciflorus*, Kit.
longifolius, Nutt. Gen. Am. ii. 95 = Astragalus pictus.
luteus, Linn. Sp. Pl. 728 = Lathyrus montanus.
macrorrhizus, Dulac, Fl. Hautes-Pyr. 291 = Lathyrus montanus.
magellanicus, Alef. in Bonplandia, ix. (1861) 144 = Lathyrus magellanicus.
maritimus, Reichb. Fl. Germ. Excurs. 538 = Lathyrus maritimus.
montanus, Hoppe & Hornsch. in Flora, xxi. (1838) 38, 485 = Lathyrus pannonicus.
montanus, Scop. Fl. Carn. ed. II. ii. 60 = Lathyrus montanus.
Muehlenbergi, Alef. in Bonplandia, ix. (1861) 146 = Lathyrus venosus.
multiflorus, Sieber, in Flora, iv. (1821) 97 = *venetus*.
mutabilis, Alef. in Bonplandia, ix. (1861) 144.—Sibir.
myrtiflorus, Alef. in Bot. Zeit. xviii. (1860) 162 = Lathyrus palustris.
nervosus, Boiss. Diagn. Ser. I. vi. 45 = Lathyrus nervosus.
niger, Linn. Sp. Pl. 729 = Lathyrus niger.
Nissolia, Doell, Rhein. Fl. 788 = Lathyrus Nissolia.
ochroleucus, Waldst. & Kit. Pl. Rar. Hung. ii. 123. t. 118 = Lathyrus ochroleucus.
Ochrus, A. Br. Ind. Sem. Hort. Berol. (1853) 13 = Lathyrus Ochrus.
orientalis, Boiss. Diagn. Ser. I. ii. 106 = Lathyrus montanus.
pallescens, Bieb. Fl. Taur. Cauc. ii. 153 = Lathyrus canescens.
palustris, Reichb. Fl. Germ. Excurs. 537 = Lathyrus palustris.
pannonicus, Jacq. Enum. Stirp. Vindob. 128 = Lathyrus pannonicus.
pannonicus, Pall. Tabl. Taur. 55 = Lathyrus canescens.
paradensis, Kit. & Kanitz, in Verh. Zool.-Bot. Ges. Wien, xiii. (1863) 554.—Hungar.
pastoënsis, Spreng. Syst. iv. Cur. Post. 283 = Coursetia dubia.
pauciflorus, Hort. ex Link, Enum. Hort. Berol. ii. 236 = *venosus*.
pauciflorus, Kit. in Linnaea, xxxii. (1863) 632.—Hungar.
pellucidus, Spreng. Syst. iii. 260 = Lessertia falciformis.
persicus, Boiss. Diagn. Ser. I. ii. 107 = Vicia iranica.
Philippi, Alef. in Bonplandia, ix. (1861) 144.—Chili.
Piscidia, Spreng. Pugill. i. 47 (Quid ?).—N. Caled.
pisiformis, Wender. Ind. Sem. Hort. Marb. (1837); ex Steud. Nom. ed. II. ii. 233; A. Br. Ind. Sem. Berol. (1853) 13 = Lathyrus pisiformis.
Plukenetii, Lapeyr. in Mém. Mus. Par. ii. (1815) 299 = Lathyrus montanus.
polymorphus, Alef. in Bonplandia, ix. (1861) 146 = Lathyrus polymorphus.
praecox, Kit. in Linnaea, xxxii. (1863) 632 = seq. ?
praecox, Lilja, in Bot. Notiser (1839), 32 = Lathyrus vernus ?

OROBUS:—
pratensis, Stokes, Bot. Mat. Med. iv. 39; Doell, Rhein. Fl. 787 = Lathyrus pratensis.
procumbens, Mill. Gard. Dict. ed. VIII. n. 11.—Mexic.
prostratus, Host, Fl. Austr. ii. 322 = Lathyrus montanus.
Pseudo-Clusii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 193.—Europ.
pubescens, Alef. in Bonplandia, ix. (1861) 143 = Lathyrus pubescens.
purpureo-caeruleus, Gilib. Fl. Lituan. ii. 100 = Lathyrus vernus.
pyrenaeus, Philippe, Fl. Pyren. i. 263 = seq.
pyrenaicus, Linn. Sp. Pl. 729 = Lathyrus montanus.
pyrenaicus, Scop. Fl. Carn. ed. II. ii. 59 = *venetus*.
quadrangulus, Spreng. Syst. iii. 258 = Lathyrus canescens.
Rafinesquei, Presl, Delic. Prag. 41 = Lathyrus canescens.
ramuliflorus, Maxim. Prim. Fl. Amur. 83.—Manchuria.
rigidus, Láng, ex Rochel, Pl. Banat. Rar. 54 = Lathyrus vernus.
roseus, Ledeb. Fl. Ross. i. 687 = Lathyrus roseus, *Stev.*
rupensis, Lerchenf. ex Schur, Enum. Pl. Transs. 167 = Vicia oroboides.
ruscifolius, Willd. ex Poir. Encyc. Suppl. iv. 203 = *venetus*.
saxatilis, Vent. Jard. Cels, t. 94 = Lathyrus ciliatus.
Semenovii, Regel & Herd. in Bull. Soc. Nat. Mosc. xxxix. (1866) II. 35.—Turkestan.
sericeus, Boiss. & Bal. ex Boiss. Fl. Orient. ii. 621.—As. Min.
serotinus, Presl, Delic. Prag. 41 = *venetus*.
sessiliflorus, Sm. ex DC. Prod. ii. 377.—Hab.?
sessilifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 64.—Graec.; As. Min.
setifolius, Philippe, Fl. Pyrén. i. 266; Alef. in Bonplandia, ix. (1861) 141 = Lathyrus setifolius.
setiformis, Schleich. ex Gaud. Fl. Helv. iv. 500 = Lathyrus canescens.
siculus, Rafin. Caratt. 72. n. 182 = Lathyrus canescens.
silvestris, Auct. = *sylvestris*.
sparsiflorus, Cocc. & Cug. in Mem. Acad. Sc. Bolog. Ser. III. vii. (1876) 559.—Europ.
sphaericus, Philippe, Fl. Pyrén. i. 266; Dulac, Fl. Hautes-Pyr. 293 = Lathyrus sphaericus.
stenophyllus, Steud. Nom. ed. II. ii. 233.—Sibir.
stipulaceus, Hook. Bot. Mag. t. 2937 = *venosus*.
styriacus, Gremli, ex Dalla Torre, Atl. Alpenfl. 89.—Europ.
subalpinus, Herbich, Stirp. Rar. Bucov. 49 = Lathyrus montanus.
subvillosus, Ledeb. Fl. Alt. iii. 359 = Vicia subvillosa.
sylvaticus, Bast. Fl. Maine-et-Loire, Suppl. 7 = Vicia cassubica.
sylvaticus, Baumg. Enum. Stirp. Transs. ii. 329 = Lathyrus vernus.
sylvaticus, Linn. Cent. Pl. i. 23; Amoen. Acad. iv. 284 = Vicia Orobus.
sylvestris, Wender. ex Steud. Nom. ed. II. ii. 233 = Lathyrus pisiformis.
tenuifolius, Baumg. Enum. Stirp. Transs. ii. 328 = Lathyrus pannonicus.
tenuifolius, Roth, Beitr. i. 78; Tent. Fl. Germ. i. 305; ii. 170 = Lathyrus montanus.
tomentosus, Desf. Tabl. ed. I. 195 = Coursetia tomentosa.
Tournefortii, Lapeyr. in Mém. Mus. Par. ii. (1815) 298. t. 10 = Lathyrus montanus?
trachycarpus, Boiss. Diagn. Ser. I. vi. 45 = Lathyrus trachycarpus.
transylvanicus, Spreng. Syst. iii. 260 = Lathyrus montanus.
tricuspidata, Janka, in Termész. Füzetek, ix. (1885) 145 = Vicia tricuspidata?
tristis, Láng, ex Reichb. Fl. Germ. Excurs. 538 = Lathyrus niger.
tuberosus, Linn. Sp. Pl. 728 = Lathyrus montanus.
uninervius, Turcz. ex Boiss. in Flora, xvii. (1834) I. Beibl. 11 = Lathyrus nervosus.
variabilis, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 617.—Cilicia.
variegatus, Lapeyr. Hist. Abr. Pl. Pyr. 414 = Lathyrus montanus.
variegatus, Tenore, Fl. Nap. Prod. Suppl. i. 62 = *venetus*.

OROBUS:—
varius, Hill, Veg. Syst. xiii. 63; Soland. ex Sims, Bot. Mag. t. 675 = Lathyrus pannonicus.
varius, Schleich. ex Reichb. Fl. Germ. Excurs. 529 = Vicia pallens.
venetus, Mill. Gard. Dict. ed. VIII. n. 8.—Europ. austr.
venosus, Willd. ex Link, Enum. Hort. Berol. ii. 236 = Lathyrus venosus.
vernus, Linn. Sp. Pl. 728 = Lathyrus vernus.
vernus, Sibth. & Sm. Fl. Graec. vii. t. 691 = *venetus*.
versicolor, Bieb. ex Steud. Nom. ed. II. ii. 233 = *venetus*.
versicolor, J. F. Gmel. Syst. 1108 = Lathyrus pannonicus.
vestitus, Alef. in Bonplandia, ix. (1861) 145 = Lathyrus vestitus.
viciaeformis, Lag. Gen. et Sp. Nov. 22 = Vicia calcarata.
viciaeformis, Vill. ex Willk. & Lange, Prod. Fl. Hisp. iii. 322 = Lathyrus canescens.
vicioides, Ser. in DC. Prod. ii. 377 = Vicia oroboides.
vicioides, Vill. Prosp. 43 = Lathyrus canescens.
villosus, Torr. in A. Gray, Bot. U. St. Expl. Exped. ii. 267 = Lathyrus Torreyi.

OROGA, Rafin. Fl. Tellur. iii. 86 (1836) = **Origanum**, Tourn. (Labiat.).
heracleotica, Rafin. l. c. = Origanum vulgare.

OROGENIA, S. Wats. in Bot. King's Exped. 120. t. 15. f. 1–3 (1871). *UMBELLIFERAE*.
linearifolia, *S. Wats. l. c.*—Am. bor. occ.

OROLLANTHUS, E. Mey. Comm. Pl. Afr. Austr. ii. 230 (1837), sphalm. = **Aeolanthus**, Mart. (Labiat.).

ORONTIUM, Linn. Diss. Chen. 26 (1751); Amoen. Acad. iii. 17 (1756). *AROIDEAE*, Benth. & Hook. f. iii. 994.
ARONIA, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 206.
aquaticum, *Linn. Sp. Pl.* 324.—Am. bor.
cochinchinense, Lour. Fl. Cochinch. 208 = Acorus Calamus.
japonicum, Thunb. Fl. Jap. 144 = Rohdea japonica.
liliifolium, Salisb. Prod. 263 = Rohdea japonicum.

ORONTIUM, Pers. Syn. ii. 158 (1807) = **Antirrhinum** Linn. (Scrophul.).
arabicum, Ehrenb. ex Boiss. Fl. Orient. iv. 386 = Schweinfurthia pterosperma.
arvense, Pers. Syn. ii. 158 = A. Orontium.
Asarina, Pers. l. c. 159 = A. Asarina.
calycinum, Pers. l. c. 158 = A. Orontium.
majus, Pers. l. c. = A. majus.
molle, Pers. l. c. = A. molle.
papilionaceum, Pers. l. c. = Schweinfurthia sphaerocarpa.
pinnatum, Pers. l. c. 159 = Nemesia pinnata.
sempervirens, Pers. l. c. 158 = A. sempervirens.
siculum, Pers. l. c. = A. siculum.
supinum, Willem. ex Steud. Nom. ed. I. 433, in syn. = Linaria repens.

OROPETIUM, Trin. Fund. Agrost. 98. t. 3 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1206.
thomaeum, *Trin. l. c.*—Ind. or.

OROPHEA, Blume, Bijdr. 18 (1825). *ANONACEAE*, Benth. & Hook. f. i. 29.
acuminata, *A. DC. Mém. Anonac.* 39.—Burma.
anceps, *Pierre, Fl. For. Cochinch.* t. 46.—Cambodia.
aurantiaca, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 25.—N. Guinea.
Beccarii, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 18.—N. Guin.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 26.—Borneo.
Brandisii, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 91.—Burma.
celebica, *Miq. Fl. Ind. Bat.* i. II. 30.—Ins. Celebes.
chrysocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 24.—Borneo.
coriacea, Thw. Enum. Pl. Zeyl. 8 = Bocagea coriacea.
corymbosa, *Miq. Fl. Ind. Bat.* i. II. 29.—Java.
costata, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 19.—N. Guin.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 170.—Ins. Philipp.

OROPHEA:—
Desmos, *Pierre, Fl. For. Cochinch.* t. 43.—Cambodia.
Diepenhorstii, *Scheff. in Flora*, lii. (1869) 302; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 10.—Java.
dodecandra, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 25.—Borneo.
enneandra, *Blume, Bijdr.* 18.—Malaya.
enterocarpa, *Maing. ex Hook. f. Fl. Brit.* i. 92.—Malacca.
enterocarpoidea, *Vidal, Phan. Cuming. Philipp.* 171.—Ins. Philipp.
erythrocarpa, *Bedd. in Madr. Journ. Sc.* Ser. II. xxii. (1861) 71.—Ind. or.
Harmandiana, *Pierre, Fl. For. Cochinch.* t. 44.—Cambodia.
hexandra, *Blume, Bijdr.* 18.—Java.
Heyneana, Hook. f. & Thoms. Fl. Ind. i. 110 = Mitrephora Heyneana.
katschallica, *Kurz, in Journ. Bot.* xiii. (1875) 323.—Ins. Nicobar.
latifolia, *Miq. Fl. Ind. Bat.* i. II. 30.—Java.
macrocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 26.—Ins. Celebes.
obliqua, Hook. f. & Thoms. Fl. Ind. i. 112 = Bocagea obliqua.
ovata, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 3.—N. Guin.
pamateonis, *Miq. Fl. Ind. Bat.* i. II. 24.—Borneo.
polycarpa, *A. DC. Mém. Anonac.* 39.—Burma.
polycephala, *Pierre, Fl. For. Cochinch.* t. 46.—Cambodia.
reticulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 23.—Java.
rugosa, *Miq. l. c.*—Java.
sumatrana, *Miq. l. c.* 24.—Sumatra.
Thomsoni, *Bedd. in Madr. Journ.* Ser. II. xxii. (1861) 72.—Ind. or.
Thorelii, *Pierre, Fl. For. Cochinch.* t. 44.—Cambodia.
trachycarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 23.—Sumatra.
trigyna, *Miq. l. c.* 25.—Borneo.
undulata, *Pierre, Fl. For. Cochinch.* t. 45.—Cambodia.
uniflora, *Hook. f. & Thoms. Fl. Ind.* i. 110.—Ind. or.
zeylanica, *Hook. f. & Thoms. l. c.* 111.—Zeylan.

OROPHOMA, Drude, in Mart. Fl. Bras. iii. II. 294. t. 66 (1882) = **Mauritia**, Linn. f. (Palm.).
Caranà, Spruce, ex Drude, l. c. 295 = M. Caranà.
subinermis, Drude, l. c. 296 = M. subinermis.

OROPOGON, Neck. Elem. iii. 211 (1790) = **Andropogon**, Linn. (Gramin.).

OROSTACHYS, [Fisch.] Hort. Gorenk. ed. I (1808), 99 = **Cotyledon**, Tourn. (Crassul.).
chlorantha, [Fisch.] l. c. = C. spinosa.
malachophylla, [Fisch.] l. c. = C. malachophylla.
serrata, Sweet, Hort. Brit. ed. II. 225 = C. malachophylla.
spinosa, Sweet, l. c. = C. spinosa.
thyrsiflora, [Fisch.] Cat. Hort. Gorenk. ed. I (1808), 99 = C. leucantha, Crassula thyrsiflora.

OROSTACHYS, Steud. Nom. ed. II. ii. 233 (1841) = ORTHOSTACHYS, Ehrh. = **Elymus**, Linn. (Gramin.).

OROTHAMNUS, Pappe, ex Hook. Bot. Mag. t. 4357 (1848) = **Mimetes**, Salisb. (Proteac.).
Zeyheri, Pappe, ex Hook. l. c. = M. Zeyheri.

OROXYLON, Vent. Dec. Gen. Nov. (1808) 8. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1040.
CALOSANTHES, Blume, Bijdr. 760 (1826).
HIPPOXYLON, Rafin. Sylva Tellur. 78 (1838).
indicum, *Vent. Dec. Gen. Nov.* 8.—As. trop.

ORPHANIDESIA, Boiss. & Bal. ex Boiss. Pl. Or. Nov. Dec. i. 3 (1875). *ERICACEAE*, Benth. & Hook. f. ii. 586.
gaultherioides, *Boiss. & Bal. l. c.* 4.—As. Min.

ORPHIUM, E. Mey. Comm. Pl. Afr. Austr. 181 (1837). *GENTIANEAE*, Benth. & Hook. f. ii. 805.
VALERANDA, Neck. Elem. ii. 33 (1790).
arenarium, Presl, Bot. Bemerk. 104 = Chironia arenaria.
frutescens, *E. Mey. Comm. Pl. Afr. Austr.* 181.—Afr. austr.

ORSIDICE, Reichb. f. in Bonplandia, ii. (1854) 93 = **Sarcochilus**, R. Br. (Orchid.).
amplexicaulis, Reichb. f. l. c. = S. lilacinus.
lilacina, Reichb. f. l. c. = S. lilacinus.

ORSINA, Bertol. in Ann. Stor. Nat. Bologn. I. ii. (1829) 362 = **Inula**, Linn. (Compos.).
camphorata, Bertol. l. c. = I. viscosa.

ORSINIA, Bertol. ex DC. Prod. v. 104 (1836) = **Clibadium**, Linn. (Compos.).
Eupatoria, DC. l. c. = C. rotundifolium.

ORSOPEA, Rafin. Sylva Tellur. 73 (1838) = **Sterculia**, Linn.
fetida, Rafin. l. c. = S. Helicteres.

ORTACHNE, Nees, in Seem. Bot. Voy. Herald, 225 (1852–57) = **Aristida**, Linn. (Gramin.).
pilosa, Nees, l. c. = A. jorullensis.
retorta, Nees, ex Steud. Syn. Pl. Gram. 121.—Patagon.
scabra, Fourn. in Bull. Soc. Bot. Fr. xxvii. (1880) 295 = A. scabra.
tenuis, Fourn. l. c. = A. tenuis.

ORTEGA, Linn. Sp. Pl. ed. I. 560 (1753). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
Mosina, Adans. Fam. ii. 272 (1763).
Terogia, Rafin. Fl. Tellur. iii. 56 (1836).
dichotoma, *Linn. Mant.* ii. 174.—Italia.
hispanica, *Linn. Sp. Pl.* 560.—Hispan.; Afr. bor.

ORTEGIA, Loefl. Iter Hisp. 112 (1758) = praec.

ORTEGIOIDES, Soland. ex DC. Prod. iii. 76 (1828) = **Ammannia**, Houst. (Lythrar.).
decussata, Soland. l. c. = A. pentandra.

ORTGIESIA, Regel, Ind. Sem. Hort. Petrop. (1866) 80; et Gartenfl. (1867) 193. t. 547. *BROMELIACEAE*, [Benth. & Hook. f. iii. 662].
tillandsioides, *Regel, ll. cc.* 81; 193.—Bras.; Uraguay.

ORTHAEA, Klotzsch, in Linnaea, xxiv. (1851) 23. *VACCINIACEAE*, Benth. & Hook. f. ii. 568.
panurensis, Spruce, ex Benth. & Hook. f. Gen. ii. 568 = Satyria panurensis.
secundiflora, *Klotzsch, in Linnaea*, xxiv. (1851) 24.—Peruv.

ORTHANTHE, Lem. Illustr. Hortic. iii. (1856) sub t. 81 = **Gloxinia**, L'Hérit. (Gesner.).
Fyfiana, Lem. l. c. = G. *Fyfiana*.

ORTHANTHERA, Wight, Contrib. 48 (1834). *ASCLEPIADEAE*, [Benth. & Hook. f. ii. 778.]
viminea, *Wight, l. c.*—Ind. bor. occ.

ORTHOCARPUS, Nutt. Gen. Am. ii. 56 (1818). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 974.
Oncorhynchus, Lehm. Ind. Sem. Hort. Hamb. (1832).—Cf. Linnaea, viii. (1833) Litt. 5.
Physocheilus, Nutt. ex Benth. in DC. Prod. x. 536, in syn. (1846).
Triphysaria, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 52 (1835).
attenuatus, *A. Gray, in Pacif. Rail. Rep.* iv. 121.—Am. bor. occ.
australis, *Benth. in DC. Prod.* x. 537.—Peruv.
Beldingi, *Greene, in Bull. Calif. Acad.* i. III. (1885) 123.—Calif.
Bidwelliae, *A. Gray, in Proc. Am. Acad.* xv. (1880) 51.—Calif.
bracteosus, *Benth. Scroph. Ind.* 13.—Am. bor. occ.
campestris, *Benth. Pl. Hartw.* 329.—Calif.
castillejoides, *Benth. Scroph. Ind.* 13.—Am. bor. occ.
densiflorus, *Benth. l. c.*—Calif.
erianthus, *Benth. l. c.* 12.—Calif.
faucibarbatus, A. Gray, in Pacif. Rail. Rep. iv. 121 = erianthus.
floribundus, *Benth. Scroph. Ind.* 12.—Calif.
gracilis, *Benth. l. c.*—Calif.
hispidus, *Benth. l. c.* 13—Am. bor. occ.
imbricatus, *Torr. ex S. Wats. in Bot. King's Exped.* 458.—Am. bor. occ.
lacerus, *Benth. Pl. Hartw.* 329.—Calif.
lasiorhynchus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 82.—Calif.

ORTHOCARPUS:—
linearifolius, Benth. ex A. Gray, in Proc. Am. Acad. xix. (1884) 95, sphalm. = tenuifolius.
linearilobus, *Benth. Pl. Hartw.* 330.—Calif.
lithospermoides, *Benth. Scroph. Ind.* 13.—Am. bor. occ.
luteus, *Nutt. Gen. Am.* ii. 57.—Am. bor. occ.
mexicanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 463.—Mexic.
pachystachyus, *A. Gray, Syn. Fl. N. Am.* ii. 1. 300.—Calif.
pallescens, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 339.—Am. bor. occ.
pallescens, A. Gray, in Bot. King's Exped. 231 = pilosus.
Parishii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 229.—Calif.
Parryi, A. Gray, in Am. Natural. viii. (1874) 214 = pallescens.
pilosus, *S. Wats. in Bot. King's Exped.* 231, 459.—Calif.
purpurascens, *Benth. Scroph. Ind.* 13.—Calif.; Arizon.
purpureo-albus, *S. Wats. in Bot. King's Exped.* 458.—Am. bor. occ.
pusillus, *Benth. Scroph. Ind.* 12.—Calif.
strictus, Benth. l. c. = luteus.
tenuifolius, *Benth. l. c.* 12, in nota.—Am. bor. occ.
tenuifolius, A. Gray, in Bot. Calif. i. 577 = imbricatus.
Tolmiei, *Hook. & Arn. Bot. Beech. Voy.* 379.—Am. bor. occ.

ORTHOCENTRON, Cass. in Dict. Sc. Nat. xxxvi. 480 (1823) = **Cnicus**, Linn. (Compos.).
glomeratum, Cass. l. c. 481 = C. polyanthemus.

ORTHOCERAS, R. Br. Prod. 316 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 608.
Solandri, Lindl. Gen. et Sp. Orch. 512 = strictum.
strictum, *R. Br. Prod.* 317.—Austral.; N. Zel.

ORTHOCHILUS, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 284. t. 82 (1851) = **Eulophia**, R. Br. (Orchid.).
abyssinicus, Hochst. l. c. = E. abyssinica.
Mechowii, Reichb. f. in Flora, lxv. (1882) 532.—Afr. trop.
Renschianus, Reichb. f. l. c.—Afr. trop.
Welwitschii, Reichb. f. l. c. xlviii. (1865) 186.—Afr. trop.

ORTHOCLADA, Beauv. Agrost. 69. t. 14. f. 9 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1190.
laxa, *Beauv. l. c.* 70.—Guian.; Bras.
laxa, Kunth, Rév. Gram. t. 71 = rariflora.
rariflora, *Beauv. Agrost.* 70.—Am. trop.

ORTHODANUM, E. Mey. Comm. Pl. Afr. Austr. 131 (1835) = **Rhynchosia**, Lour. (Legumin.).
argenteum, E. Mey. l. c. 132 = R. Orthodanum.
glabratum, Meissn. in Hook Lond. Journ. Bot. ii. (1843) 91 = R. Orthodanum.
latifolium, E. Mey. Comm. Pl. Afr. Austr. i. 131 = R. Orthodanum.
Muehlenbeckii, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 92 = R. Orthodanum.
sordidum, E. Mey. Comm. Pl. Afr. Austr. 131 = R. Orthodanum.

ORTHODON, Benth. & Oliver, in Journ. Linn. Soc. ix. (1867) 167 = **Mosla**, Buch.-Ham. (Labiat.).
japonicum, Benth. & Oliver, l. c. = M. japonica.

ORTHOGYNIUM, Baill. in Bull. Soc. Linn. Par. i. (1885) 459. *MENISPERMACEAE*.
gomphioides, *Baill. l. c.*—Madag.

ORTHOLOMA, Hanst. in Linnaea, xxvi. (1853–55) 209 = **Columnea**, Linn. (Gesner.).
acuminatum, Hanst. l. c.; Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 126 = C. acuminata.
ochroleucum, Klotzsch, ex Oerst. l. c. 125 = C. ochroleuca.
pendulum, Klotzsch, ex Oerst. l. c. 126 = C. pendula.
vestitum, Klotzsch, ex Oerst. l. c. 124 = C. Warszewicziana.
Warscewiczianum, Klotzsch, ex Oerst. l. c. 125 = C. Warszewicziana.
Warczewiczii, Hanst. in Linnaea, xxvi. (1853) 216. t. 2. f. 5–6 = C. Warszewicziana.

ORTHOLOTUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 363 = **Dorycnium**, Linn. (Legumin.).
rectus, Fourr. l. c. = D. rectum.

ORTHOPETALUM, Beer, Bromel. 70 (1857) = **Pitcairnia**, L'Hérit. (Bromel.).
inerme, Beer, l. c. 72 = P. inermis.
lanuginosum, Beer, l. c. 70 = P. lanuginosa.
pulverulentum, Beer, l. c. 71 = P. pulverulenta.

ORTHOPHYTUM, Beer, in Flora, xxvii. (1854) 347 = **Pitcairnia**, L'Hérit. (Bromel.).

ORTHOPOGON, R. Br. Prod. 194 (1810) = **Oplismenus**, Beauv. (Gramin.).
abortivus, Spreng. Syst. i. 306 = Chamaeraphis spinescens.
aemulus, R. Br. Prod. 194 = Oplism. undulatifolius.
africanus, Sweet, Hort. Brit. ed. I. 448 = Oplism. compositus.
agrostoides, Trev. ex Steud. Nom. ed. II. ii. 234 = Arundinella nepalensis.
bromoides, Loud. Hort. Brit. 25 = Oplism. Burmannii.
Burmanni, R. Br. Prod. 194 = Oplism. Burmannii.
compositus, R. Br. l. c. = Oplism. compositus.
Crus-galli, Spreng. Syst. i. 307 = Panicum Crus-galli.
cubensis, Spreng. l. c. 306 = Panicum cubense.
dichotomus, Llanos, Fragm. 38 = Panicum Crus-galli.
echinatus, Spreng. Syst. i. 307 = Panicum Crus-galli.
flaccidus, R. Br. Prod. 194 = Oplism. undulatifolius.
gonyrrhizus, Miq. Fl. Ind. Bat. iii. 443.—Java.
hirsutus, Spreng. ex Steud. Nom. ed. II. ii. 234 = Panicum spectabile.
hirtellus, R. Br. Prod. 194 = Oplism. compositus.
hirtellus, Llanos, Fragm. 37 = Panicum lanceolatum.
hirtellus, Nutt. Gen. Am. i. 55 = Panicum Nuttallianum.
hispidus, Spreng. Syst. i. 307 = Panicum Crus-galli.
holciformis, Spreng. l. c. = Panicum holciforme.
imbecillis, R. Br. Prod. 194 = Oplism. undulatifolius.
loliaceus, Spreng. Syst. i. 306 = Oplism. compositus.
longeracemosus, Miq. Fl. Ind. Bat. iii. 443 = Oplism. compositus.
parvifolius, Nutt. Gen. Am. ii., in Erratis = Panicum Nuttallianum.
pratensis, Spreng. Syst. i. 306 = Oplism. compositus.
remotus, Trin. Fund. Agrost. 181 = Oplism. compositus.
Retzii, Spreng. Syst. i. 307 = Panicum Crus-galli.
setarius, Spreng. l. c. 306 = Oplism. undulatifolius.
squarrosus, Spreng. l. c. 307 = Panicum Crus-galli.
stagninus, Spreng. l. c. = Panicum Crus-galli.
subverticillatus, Llanos, Fragm. 38 = Panicum colonum.
sylvaticus, Miq. Fl. Ind. Bat. iii. 443 = Oplism. compositus.
undulatifolius, Spreng. Syst. i. 306 = Oplism. undulatifolius.
velutinus, Spreng. l. c. = Oplism. Burmannii ?

ORTHORAPHIUM, Nees, in Proc. Linn. Soc. i. (1841) 94 = **Stipa**, Linn. (Gramin.).
Roylei, Nees, l. c.—Ind. or.

ORTHOSANTHUS, Steud. Nom. ed. II. ii. 234 (1840) = **Orthrosanthus**, Sweet (Irid.).

ORTHOSELIS, Spach, Hist. Vég. Phan. vi. 453 (1838) = **Heliophila**, Linn. (Crucif.).
pilosa, Spach, l. c. 454 = H. pilosa.

ORTHOSIA, Decne. in DC. Prod. viii. 526 (1844) = **Vincetoxicum**, Moench (Asclep.).
acuminata, Griseb. Cat. Pl. Cub. 175.—Cuba.
arenosa, Decne. in DC. Prod. viii. 527.—Bras.
congesta, Decne. l. c. 527.—Bras.
Decaisnei, Fourn. in Mart. Fl. Bras. vi. IV. 222.—Bras.
Eichleri, Fourn. l. c.—Bras.
Kunthii, Decne. in DC. Prod. viii. 527 = V. Kunthii.
multiflora, Fourn. in Mart. Fl. Bras. vi. IV. 222.—Bras.
oblongata, Griseb. Cat. Pl. Cub. 176.—Cuba.

ORTHOSIA :—
paniculata, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 954, nomen.—Guian.
serpyllifolia, Decne. in DC. Prod. viii. 527.—Ecuador.
suberosa, Decne. l. c. = Metastelma suberosum.
umbrosa, Decne. l. c.—Bras.
urceolata, Fourn. in Mart. Fl. Bras. vi. iv. 223.—Bras.
virgata, Fourn. l. c.—Bras.

ORTHOSIPHON, Benth. in Bot. Reg. sub t. 1300 (1830); et Lab. Gen. et Sp. 25 (1832). *LABIATAE*, Benth & Hook. f. ii. 1174.
adscendens, [Benth.] ex Sweet, Hort. Brit. ed. II. 413 = Ocimum adscendens.
ambiguus, *Bolus, in Journ. Linn. Soc.* xviii. (1881) 394.—Afr. austr.
aristatus, *Miq. Fl. Ind. Bat.* ii. 943.—Java.
asperus, [Benth.] ex Sweet, Hort. Brit. ed. II. 413 = Coleus barbatus.
australis, *Vatke, in Linnaea*, xl. (1876) 179.—Afr. trop.
bracteatus, Wight, Ic. t. 1428 = comosus.
brevicaulis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 433.—Madag.
breviflorus, *Vatke, in Linnaea*, xliii. (1880–82) 88.—Afr. bor.
capitatus, Benth. Lab. Gen. et. Sp. 29 = Catopheria capitata.
cleistocalyx, *Vatke, in Linnaea*, xxxvii. (1871–73) 317.—Abyssin.
coloratus, *Vatke, l. c.* xliii. (1880–82) 86.—Afr. trop.
comosus, *Wight, ex Benth. in DC. Prod.* xii. 52.—Ind. or.
diffusus, *Benth. l. c.* 50.—Ind. or.
Ehrenbergii, *Vatke, in Linnaea*, xxxvii. (1871–73) 316.—Abyssin.
emirnensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 433.—Madag.
ferrugineus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotr.
glabratus, Benth. in Wall. Pl. As. Rar. ii. 14; et in Cat. n. 2724 = tomentosus.
Hildebrandtii, *Vatke, in Bremen, Abh.* ix. (1885) 134.—Madag.
hispidus, Benth. in DC. Prod. xii. 50; et in Wall. Cat. n. 2718 = diffusus.
incurvus, *Benth. in Wall. Pl. As. Rar.* ii. 15.—Reg. Himal.
inodorus, Koen. ex Hook. f. Fl. Brit. iv. 613 = pallidus.
linearis, *Benth. in Hook. Ic. Pl.* t. 1274.—Afr. trop.
pallidus, *Benth. Lab. Gen. et Sp.* 708; *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 370.—Ind. or.; Arab.
parvifolius, *Vatke, in Linnaea*, xliii. (1880–82) 87.—Afr. trop.
petiolaris, *Miq. Fl. Ind. Bat.* ii. 943.—Java.
physocalycinus, *A. Rich. Tent. Fl. Abyss.* ii. 180.—Abyssin.
reflexus, Vatke, in Linnaea, xliii. (1880–82) 85 = Ehrenbergii.
robustus, *Hook. f. Fl. Brit. Ind.* iv. 614.—Reg. Himal.
rubicundus, *Benth. in Wall. Pl. As. Rar.* ii. 14.—Ind. or.; Burma.
scapiger, *Benth. l. c.* 15.—Reg. Himal.
Schimperi, *Benth. in DC. Prod.* xii. 51.—Abyssin.
secundiflorus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 433.—Madag.
somalensis, *Vatke, in Linnaea*, xliii. (1880–82) 87.—Afr. trop.
spicatus, *Benth. in DC. Prod.* xii. 53.—N. Granat.
stamineus, *Benth. in Wall. Pl. As. Rar.* ii. 15.—Ind. or.; Malaya; Austral.
tenuiflorus, *Benth. in DC. Prod.* xii. 50.—Arab.
tomentosus, *Benth. in Wall. Pl. As. Rar.* ii. 14.—Ind. or.
tristis, Benth. in Bot. Reg. sub t. 1300 = tomentosus.
verticillatus, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 613 = pallidus.
virgatus, Benth. in Wall. Pl. As. Rar. ii. 14 = rubicundus.
viscosus, Benth. in Wall. Cat. n. 2723 = tomentosus.
viscosus, *Benth. in Wall. Pl. As. Rar.* ii. 14.—Ind. or.

ORTHOSPERMUM, Opiz, Seznam, 70 (1852); Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 63 = **Chenopodium**, Tourn.
acuminatum, Schur, Enum. Pl. Transs. 571 = C. rubrum.

ORTHOSPERMUM :—
Bonus-Henricus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 63 = Chenopodium Bonus-Henricus.
crassifolium, Schur. l. c. = C. rubrum.
glaucum, Opiz, Seznam, 71 = C. glaucum.
rubrum, Opiz, l. c. 70 = C. rubrum.

ORTHOSPORUM, T. Nees, Gen. Fl. Germ. Monochl. n. 58 (1835) = **Chenopodium**, Tourn.
ambrosioides, Kostel. Allg. Med.-Pharm. Fl. iv. 1433 = C. ambrosioides.
anthelminticum, Kostel. l. c. = C. ambrosioides.
Bonus-Henricus, T. Nees, Gen. Fl. Germ. Monochl. n. 58 = C. Bonus-Henricus.
glaucum, Peterm. Fl. Bienitz, 94 = C. glaucum.
multifidum, Kostel. Allg. Med.-Pharm. Fl. iv. 1434 = Roubieva multifida.
rubrum, T. Nees, Gen. Fl. Germ. Monochl. n. 58 = C. rubrum.
suffruticosum, Kostel. Allg. Med.-Pharm. Fl. iv. 1433 = C. ambrosioides.

ORTHOSTACHYS, Ehrh. Beitr. iv. 146 (1789) = **Elymus**, Linn. (Gramin.).

ORTHOSTACHYS, Spach, Hist. Vég. Phan. ix. 32 (1840) = OROSTACHYS, Fisch. = **Cotyledon**, Tourn. (Crassul.).

ORTHOSTEMMA, Wall. ex Voigt, Hort. Suburb. Calc. 384 (1845) = **Pentas**, Benth. (Rubiac.).
comorensis, Boiv. ex Baill. Adansonia, xii. (1879) 234 = Cremaspora comorensis.
roseum, Wall. ex Voigt, Hort. Suburb. Calc. 384 = P. carnea?

ORTHOSTEMON, Berg, in Linnaea, xxvii. (1854) 440 = **Feijoa**, Berg (Myrtac.).
obovatus, Berg, in Mart. Fl. Bras. xiv. i. 468 = F. obovata.
Sellowianus, Berg, l. c. 467 = F. Sellowiana.

ORTHOSTEMON, R. Br. Prod. 451 (1810) = **Canscora**, Lam. (Gentian.).
erectum, R. Br. l. c. = C. diffusa.
Huegelii, Griseb. in DC. Prod. ix. 63 = C. diffusa.

ORTHOTACTUS, Nees, in Mart. Fl. Bras. ix. 131. t. 21 (1847) = **Dianthera**, Gronov. (Acanth.).
aequilabris, Nees, l. c. 134.—Bras.
Arnottianus, Nees, l. c. 132.—Bras.
ciliatus, Nees, l. c. 135 = Beloperone ciliata?
Felisbertianus, Nees, l. c. 132 = Adhatoda Felisbertiana.
glandulosus, Nees, l. c.—Bras.
montanus, Nees, l. c. 134.—Bras.
nemorosus, Nees, in Benth. Bot. Voy. Sulph. 147, in obs.—Hab.?
oblongus, Nees, l. c. 147 = D. oblonga.
Pohlianus, Nees, in Mart. Fl. Bras. ix. 134.—Bras.
roseus, Nees, l. c. 133.—Bras.
strobilaceus, Nees, l. c.—Bras.
venosus, Nees, l. c.—Bras.

ORTHOTHECIUM, Schott & Endl. Melet. 31 (1832) = **Helicteres**, Pluk. (Stercul.).
hirsutum, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 135 = H. hirsuta.
javanense, Hassk. l. c. xii. (1845) 105 = H. javanensis.
Lhotskyanum, Schott & Endl. Melet. 31.—Am. austr.
viscidum, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 135 = H. viscida.

ORTHOTHELIUM, Walp. Ann. i. 105 (1848), sphalm. = praec.

ORTHOTROPIS, Benth. in Lindl. Swan River App. 16 (1839) = **Chorizema**, Labill. (Legumin.).
pungens, Benth. l. c. = C. Henchmanni.

ORTHROSANTHES, Rafin. Fl. Tellur. iv. 30 (1836) = **Orthrosanthus**, Sweet (Irid.).
ixioides, Rafin. l. c. = Orthrosanthus multiflorus.

ORTHROSANTHUS, Sweet, Fl. Austral. t. 11 (1827). *IRIDEAE*, Benth. & Hook. f. iii. 697.
EVELTRIA, Rafin. Fl. Tellur. iv. 30 (1836).
caeruleus, *Mangles, ex Herb. in Journ. Hort. Soc.* ii. (1847) 92 (Quid?).—Austral.
chilensis, Klotzsch, ex Baker, in Journ. Linn. Soc. xvi. (1877) 121 = Solenomelus chilensis.
chimboracensis, *Baker, in Gard. Chron.* (1876) ii. 67.—Am. trop.
gracilis, *Herb. in Journ. Hort. Soc.* ii. (1847) 92, nomen.—Austral.
gramineus, *Benth. Fl. Austral.* vi. 411.—Austral.
laxus, *Benth. l. c.*—Austral.
Moritzianus, Klotzsch, ex Baker, in Journ. Linn. Soc. xvi. (1877) 113 = chimboracensis.
Muelleri, *Benth. Fl. Austral.* vi. 411.—Austral.
multiflorus, *Sweet, Fl. Austral.* t. 11.—Austral.
polystachyus, *Benth. Fl. Austral.* vi. 411.—Austral.
spicatus, *Baker, in Journ. Linn. Soc.* xvi. (1877) 113.—Bras.

ORTIGA, Neck. Elem. ii. 400 (1790) = **Loasa**, Adans.

ORTMANNIA, Opiz, in Flora, xvii. (1834) 592 = **Geodorum**, Jacks. (Orchid.).
cernua, Opiz, l. c. = G. dilatatum.

ORTOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvii. (1869) 136 = **Stachys**, Tourn. (Labiat.).
recta, Fourr. l. c. = S. recta.

ORUCARIA, Juss. ex DC. Prod. ii. 420 (1825) = **Drepanocarpus**, G. F. W. Mey. (Legumin.).
lunata, Juss. l. c. = D. lunatus.

ORVALA, Linn. Gen. ed. I. 174 (1737) = **Lamium**, Tourn. (Labiat.).
garganica, Linn. Sp. Pl. 578 = L. Orvala.
lamioides, DC. Fl. Fr. iii. 539 = L. Orvala.

ORXERA, Rafin. Fl. Tellur. iv. 37 (1836) = **Aërides**, Lour. (Orchid.).
cornuta, Rafin. l. c. = A. odoratum.

ORYCHOPHRAGMUS, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 81; (Enum. Pl. Chin. Bor. 7) = **Moricandia**, DC. (Crucif.).
sonchifolius, Bunge, ll. cc. = M. sonchifolia.

ORYCTANTHUS. Eichl. in Mart. Fl. Bras. v. ii. 87. t. 29, 30 (1868) = **Loranthus**, Linn.
amplexicaulis, Eichl. l. c. 88 = L. amplexicaulis.
botryostachyus, Eichl. l. c. 89.—Am. trop.
chloranthus, Eichl. l. c. = L. chloranthus.
glaberrimus, Eichl. l. c. = L. glaberrimus.
occidentalis, Eichl. l. c. = L. occidentalis.
ruficaulis, Eichl. l. c. 90 = L. ruficaulis.
scabridus, Eichl. l. c. 91.—Am. trop.
spicatus, Eichl. l. c. 89 = L. spicatus.

ORYCTES, S. Wats. Bot. King's Exped. 274. t. 28 (1871). *SOLANACEAE*, Benth. & Hook. f. ii. 893.
nevadensis, *S. Wats. l. c.*—Am. bor. occ.

ORYGIA, Forsk. Fl. Aegypt. Arab. 103 (1775). *FICOIDEAE*, Benth. & Hook. f. i. 856.
AXONOTECHIUM, Fenzl, in Ann. Wien. Mus. i. (1836) 354; ii. (1838) 299.
decumbens, *Forsk. Fl. Aegypt. Arab.* 103.—As. occ.; Ind. or.; Afr. trop.
mucronata, Klotzsch, in Peters, Reise Mossamb. Bot. 140 = decumbens.
portulacaefolia, Forsk. Fl. Aegypt. Arab. 103 = Talinum cuneifolium.

ORYMARIA, Meissn. Gen. 103 (1838) = **Orimaria**, Rafin.

ORYSA, Desv. Journ. Bot. i. (1813) 76 = **Oriza**, Linn. (Gramin.).

ORYTHIA, Endl. Gen. Suppl. i. 1408 (1841) = **Agalmyla**, Blume (Gesner.).

ORYZA, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1116.
PADIA, Zoll. & Mor. Syst. Verz. Pfl. Zoll. 103 (1846).
caudata, Doell, in Mart. Fl. Bras. ii. ii. 8 = sativa?
clandestina, A. Br. ex Aschers. Fl. Brandenb. 799 = Leersia oryzoides.

ORYZA :—
coarctata, *Roxb. Hort. Beng.* 87 ; *Fl. Ind.* ii. 206.—Reg. Himal.
communissima, Lour. Fl. Cochinch. 215 = sativa.
denudata, Steud. Nom. ed. I. 577 = sativa.
elongata, Steud. l. c. = sativa ?
fatua, Koen. ex Trin. in Mém. Acad. Pétersb. Sér. VI. ii. II. Sc. Nat. (1839) 177 = sativa.
filiformis, Buch.-Ham. ex Steud. Syn. Pl. Gram. 3 = sativa ?
glaberrima, *Steud. l. c.*—Afr. trop.
glumaepatula, *Steud. l. c.*—Guiana.
glutinosa, Lour. Fl. Cochinch. 215 = sativa.
granulata, Nees, in Wall. Cat. n. 8634; Steud. Syn. Pl. Gram. 3 = sativa ?
hexandra, Doell, in Mart. Fl. Bras. ii. II. 10 = Leersia hexandra.
minuta, *J. & C. Presl, Rel. Haenk.* i. 208.—Ins. Philipp.
monandra, Doell, in Mart. Fl. Bras. ii. II. 9 = Leersia monandra.
montana, Buch.-Ham. in Wall. Cat. n. 8633 = sativa ?
montana, Lour. Fl. Cochinch. 215 = sativa.
mutica, Steud. Nom. ed. I. 577 = sativa.
nepalensis, G. Don, ex Loud. Hort. Brit. 139 = sativa.
officinalis, Wall. Cat. n. 8635 ; et Steud. Syn. Pl. Gram. 3 = sativa.
palustris, Salisb. Prod. 25 = sativa.
parviflora, Beauv. Agrost. 27 = sativa.
perennis, Moench, Meth. 197 = sativa.
platyphylla, Schult. f. Syst. vii. 1364 = latifolia.
praecox, Lour. Fl. Cochinch. 215 = sativa.
prehensilis, *Steud. Syn. Pl. Gram.* 3.—Afr. austr.
pubescens, Steud. Nom. ed. I. 577 = sativa ?
pumila, Hort. ex Steud. l. c. ed. II. ii. 234 = sativa.
punctata, *Kotschy, ex Steud. Syn. Pl. Gram.* 3.—Afr. trop.
repens, Buch.-Ham. ex Steud. l. c. = sativa.
rubra, Hort. Pillnitz. ex Walp. Ann. i. 924 = Panicum colonum.
rubribarbis, Steud. Nom. ed. I. 577 = sativa.
rufipogon, Griff. Notul. iii. 5 ; Ic. Pl. Asiat. 145. f. 2 = sativa ?
sativa, Hochst. ex Steud. Syn. Pl. Gram. 3 = glumaepatula, punctata.
sativa, *Linn. Sp. Pl.* 333.—As. trop.
segetalis, Russ. ex Steud. Syn. Pl. Gram. 3 = sativa.
sorghoidea, Steud. Nom. ed. I. 577 = sativa.
sorghoides, Steud. Syn. Pl. Gram. 3 = praec.
subulata, Nees, Agrost. Bras. 518 = sativa ?
triandra, Heyne, ex Steud. Syn. Pl. Gram. 3 = sativa ?
triticoides, Griff. Notul. iii. 8 = coarctata.

ORYZOPSIS, Michx. Fl. Bor. Am. i. 51. t. 9 (1803).
GRAMINEAE, Benth. & Hook. f. iii. 1142.
CARYOCHLOA, Spreng. Syst. iv. Cur. Post. 22, 30 (1827).
DILEPYRUM, Rafin. in Med. Repos. N. York, v. (1808) 352.
ERIOCOMA, Nutt. Gen. Am. i. 40 (1818).
FENDLERIA, Steud. Syn. Pl. Gram. 419 (1855).
NASSELLA, E. Desv. in C. Gay, Fl. Chil. vi. 263. t. 75. f. 1 (1853).
PIPTATHERUM, Beauv. Agrost. 17. t. 5. f. 10. 11 (1812).
PIPTOCHAETIUM, J. & C. Presl, Rel. Haenk. i. 222. t. 37 (1830).
URACHNE, Trin. Fund. Agrost. 109 (1820).
asperifolia, [Rich. in] Michx. Fl. Bor. Am. i. 51.—Am. bor.
canadensis, *Torr. ex A. Gray, Man. Bot. N. U. St.* ed. I. 583.—Am. bor.
coerulescens, *Hack. in Denkschr. Acad. Wien*, l. (1885) 75.—As. Min.
cuspidata, *Benth. ex Vasey, Gram. U. St.* 23.—Am. bor.
exigua, *Thurb. ex Torr. Bot. U. St. Expl. Exped.* ii. 481.—Am. bor. occ.
fimbriata, *Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
Fournieriana, *Hemsl. l. c.*—Mexic.
holciformis, *Hack. in Denkschr. Acad. Wien*, l. (1885) 8.—Persia.

ORYZOPSIS :—
leucosperma, Walp. Ann. iii. 728 = asperifolia.
melanocarpa, *Muhl. Desc. Gram.* 79.—Am. bor.
micrantha, *Thurb. ex Porter & Coult. Syn. Fl. Colorado*, 145.—Am. bor. occ.
mutica, Link, Enum. Hort. Berol. i. 41, in obs. = asperifolia.
paradoxa, *Nutt. in Journ. Acad. Philad.* iii. (1823) 128.—Europ. ; Reg. Caucas.
parviflora, Nutt. l. c. 126 = canadensis.
pubiflora, *Hack. in Denkschr. Acad. Wien*, l. (1885) 8.—Persia.
setacea, *Rich. in Dict. Sc. Class.* xii. 445.—Reg. Argent.
Webberi, *Benth. ex Vasey, Gram. U. St.* 23.—Am. bor.

OSBECKIA, Linn. Sp. Pl. ed. I. 345 (1753).
MELASTOMACEAE, Benth. & Hook. f. i. 744.
AMBLYANTHERA, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849).
ASTEROSTOMA, Blume, l. c. (1849).
BECKIA, Rafin. Sylva Tellur. 103 (1838).
CERAMICALYX, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849).
HEDUSA, Rafin. Sylva Tellur. 101 (1838).
KADALI, Adans. Fam. ii. 234 (1763).
alpestris, DC. Prod. iii. 140 = Pterolepis alpestris.
alveolata, Bedd. Ic. Pl. Ind. Or. i. 37 = reticulata.
angustifolia, D. Don, Prod. Fl. Nep. 221 = chinensis.
antennina, Sm. in Rees, Cycl. xxv. n. 3 = Dissotis antennina.
Antherotoma, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 56 = Antherotoma Naudini.
aspera, Benth. in Wall. Cat. n. 4073 = glauca.
aspera, *Blume, in Flora*, xiv. (1831) 474.—Ind. or.
aspericaulis, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 55.—Burma.
Aubletiana, Spreng. Syst. ii. 311 = Rhynchanthera grandiflora.
australiana, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 59.—Austral.
Bonplandiana, Spreng. Syst. ii. 313 = Monochaetum multiflorum.
brachystemon, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 57 = cupularis.
bracteolaris, DC. Prod. iii. 141 = Marcetia bracteolaris.
brasila, Schrank, ex DC. Prod. iii. 154 = Leandra sericea.
brasiliana, Schrank, ex Steud. Nom. ed. II. ii. 234 = Leandra scabra.
buxifolia, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 309.—Zeylan.
buxifolia, Thw. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 519 = Walkeri.
campestris, Buch.-Ham. in Wall. Cat. n. 4063 = rostrata.
canescens, R. Grah. in Edinb. N. Phil. Journ. xxviii. (1840) 399 = Dissotis incana.
capitata, *Benth. in Wall. Cat.* n. 4072 ; *et ex Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850). 68.—Reg. Himal.
capitata, D. Don, ex Wight & Arn. Prod. 323 = Wightiana.
cernua, Spreng. Syst. ii. 312 = Brachyotum cernuum.
chinensis, Gaertn. Fruct. ii. 202 = zeylanica.
chinensis, [Ker-Gawl.] Bot. Reg. t. 542 = octandra.
chinensis, *Linn. Sp. Pl.* 345.—As. et Austral. trop.
Chulesis, D. Don, Prod. Fl. Nep. 221 = nepalensis.
ciliaris, Ser. ex DC. Prod. iii. 132 (= *Pleroma ciliaris*).—N. Granat.
confertiflora, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 59 = cupularis.
coronata, D. Don, ex Wight & Arn. Prod. 322 = truncata.
crinita, *Benth. in Wall. Cat.* n. 4066.—Reg. Himal. ; Burma.
crinita, Naud. in Sc. Nat. Sér. III. xiv. (1850) 72 = stellata.
cupularis, *D. Don, ex Wight & Arn. Prod.* 323.—Ind. or.
debilis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 65 = zeylanica.
debilis, Sond. in Linnaea, xxiii. (1850) 47 = Dissotis debilis.
decandra, DC. Prod. iii. 143 = Dissotis decandra.
decora, Wall. Cat. n. 4070 = chinensis.
decumbens, DC. Prod. iii. 143 = Dissotis decumbens.

OSBECKIA :—
divaricata, Spreng. Syst. ii. 312 = Pterogastra divaricata.
dolichophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 64.—Ins. Celebes.
elliptica, Naud. l. c. 62 = rubicunda.
erythrocephala, Naud. l. c. 58 = cupularis.
excelsa, Spreng. Syst. ii. 312 = Centronia excelsa.
eximia, Sond. in Linnaea, xxiii. (1850) 48 = Dissotis eximia.
Gardneriana, Wight, Ic. t. 997 = Leschenaultiana.
glabrata, Benth. in Wall. Cat. n. 4071 = chinensis.
glauca, *Benth. l. c.* n. 4073.—Ind. or.
glauca, Wall. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 68 = capitata.
glomerata, DC. Prod. iii. 141 = Pterolepis glomerata.
glutinosa, Spreng. Syst. ii. 313 = Bucquetia glutinosa.
gracilis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 45 ; et in Trans. Linn. Soc. xxv. (1865) 216.—Ind. or.
grandiflora, Sm. in Rees, Cycl. xxv. n. 6 = Dissotis grandiflora.
heteranthera, Spreng. Syst. ii. 312 = Melastoma repens.
hirta, Gaertn. Fruct. ii. t. 126 ; Sm. in Rees, Cycl. xxv. sub n. 2 = chinensis ?
hispidissima, *Wight, Ic.* t. 1612.—Ind. or.
inappendiculata, D. Don, ex Wight & Arn. Prod. 323 = octandra.
incana, E. Mey. ex Hochst. in Flora, xxvii. (1844) 424 = Dissotis incana.
japonica, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 70 = chinensis.
Jucunda, DC. Prod. iii. 139 = Miconia Jucunda.
Kleinii, Wight & Arn. Prod. 323 = aspera.
Koti-gueda, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 64 = Wightiana.
lanceolata, Spreng. Syst. ii. 312 = Tibouchina longifolia.
Langsdorffiana, Spreng. l. c. = Tibouchina Langsdorffiana.
Leschenaultiana, Dalz. & Gibs. Bomb. Fl. 92 = truncata.
Leschenaultiana, *DC. Prod.* iii. 142.—Ind. or.
Leschenaultiana, Thw. Enum. Pl. Zeyl. 104, partim = parvifolia.
Leschenaultiana, Wight, Ic. t. 996 = cupularis.
linearis, Blume, in Flora, xiv. (1831) 473 = chinensis.
longicollis, Benth. in Wall. Cat. n. 4065 = rostrata.
manillana, DC. Prod. iii. 142 = Otanthera manillana.
marginulata, *Wall. Cat.* n. 4064.—Ind. or.
maritima, A. St. Hil. Voy. Dist. Diam. ii. 417 = Pterolepis maritima.
microphylla, DC. Prod. iii. 140 = Chaetolepis microphylla.
minor, Triana, in Trans. Linn. Soc. xxviii. (1871) 55 = aspera.
Moonii, *Thw. Enum. Pl. Zeyl.* 105.—Zeylan.
multiflora, Blanco, Fl. Filip. ed. I. 293 = chinensis.
multiflora, Sm. in Rees, Cycl. xxv. n. 7 = Dissotis multiflora.
muralis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 56 = truncata.
myrtifolia, Blume, Mus. Bot. Lugd. Bat. i. 51 = chinensis.
nepalensis, *Hook. Exot. Fl.* t. 31.—Reg. Himal. ; Burma.
nutans, *Wall. Cat.* n. 4068.—Reg. Himal.
octandra, *DC. Prod.* iii. 142.—Ind. or.
oleaefolia, DC. l. c. 139 = Microlepis oleaefolia.
ornata, Sw. Fl. Ind. Occ. ii. 647 (= *Pleroma strigosum*).—Ind. occ.
ovata, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 586 = Sarcopyramis nepalensis.
parnassifolia, DC. Prod. iii. 140 = Pterolepis parnassifolia.
parvifolia, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 308.—Zeylan.
perangusta, *F. Muell. Fragm.* v. 181.—Austral.
Perrottetii, *DC. ex Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 71.—Ins. Philipp.
peruviana, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 315 = Centronia laurifolia.
phaeotricha, Hochst. in Flora, xxvii. (1844) 424 = Dissotis phaeotricha.
pityrophylla, DC. Prod. iii. 139 = Pyramia pityrophylla.
polycephala, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 67 = octandra.
Princeps, DC. Prod. iii. 140 = Dissotis Princeps.

OSBECKIA :—
pulchella, Benth. in Wall. Cat. n. 4059 = rostrata.
pumila, DC. Prod. iii. 141 = Pterolepis pumila.
pusilla, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 586 = chinensis.
quaterna, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 221 = rostrata.
repanda, DC. Prod. iii. 141 = Pterolepis repanda.
repens, DC. l. c. 142 = Melastoma repens.
reticulata, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 45, nomen; *et in Trans. Linn. Soc.* xxv. (1865) 216. —Ind. or.
Rheedii, *Gardn. ex Thw. Enum. Pl. Zeyl.* 104.—Zeylan.
rostrata, *D. Don, Prod. Fl. Nep.* 221.—Reg. Himal.
rotundifolia, Sm. in Rees, Cycl. xxv. n. 204 = Dissotis rotundifolia.
Royeni, Miq. Comm. Phyt. 69 = Melastoma polyanthum.
rubicunda, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 309.—Zeylan.
rubicunda, Thw. Enum. Pl. Zeyl. 105 = cupularis.
senegambiensis, Guill. & Perr. Fl. Seneg. Tent. i. 310 = Dissotis senegambiensis.
septemnervia, Buch.-Ham. ex Wall. Cat. n. 4062 B = stellata.
serialis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 67 = zeylanica.
sessilifolia, Stokes, Bot. Mat. Med. ii. 348 = chinensis.
Simsii, DC. Prod. iii. 140 = octandra.
sinensis, J. F. Gmel. Syst. 605 = chinensis.
sipaneoides, DC. Prod. iii. 141 = Pterolepis sipaneoides.
speciosa, D. Don, Prod. Fl. Nep. 222 = nepalensis.
speciosa, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 73 = stellata.
stellata, Buch.-Ham. ex D. Don, in Bot. Reg. t. 674 = crinita.
stellata, *Wall. Cat.* n. 4062.—Reg. Himal.
striphnocalyx, DC. Prod. iii. 140 = Pterolepis striphnocalyx.
ternifolia, D. Don, Prod. Fl. Nep. 221 = rostrata.
tetrandra, Roxb. Hort. Beng. 88; Fl. Ind. ii. 224 = Anplectrum glaucum.
tortuosa, Spreng. Syst. ii. 312 (= *Pleroma mexicanum*). —Mexic.
truncata, Arn. in Hook. Comp. Bot. Mag. ii. (1836) 308 = cupularis.
truncata, *D. Don, ex Wight & Arn. Prod.* 322.—Ind. or.
tubulosa, Sm. in Rees, Cycl. xxv. n. 5 = Dissotis tubulosa.
umlaasiana, Hochst. in Flora, xxvii. (1844) 424 = Dissotis incana.
urceolaris, DC. Prod. iii. 139 = Tibouchina urceolaris.
virgata, D. Don, ex Wight & Arn. Prod. 323 = octandra.
virusana, Baill. Hist. Pl. vii. 34 = Tristemma virusanum.
Walkeri, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 309.—Ind. or.
Wightiana, *Benth. in Wall. Cat.* n. 4060.—Ind. or.
Wightiana, Benth. l. c. n. 4074 = cupularis, octandra.
wynaadensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 521.—Ind. or.
zanzibariensis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 55 = Dissotis prostrata.
zeylanica, DC. ex Naud. l. c. 71 = chinensis.
zeylanica, [Ker-Gawl.] in Bot. Reg. t. 565 = parvifolia.
zeylanica, *Linn. f. Suppl.* 215.—Zeylan.
zeylanica, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 56 = truncata.

OSBECKIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 118 = **Dissotis**, Benth. (Melastom.).
Heudelotii, Naud. l. c. 119 = D. grandiflora.

OSBORNIA, F. Muell. Fragm. iii. 30 (1862). *MYRTACEAE*, Benth. & Hook. f. i. 711.
octodonta, *F. Muell. l. c.* 31.—Austral.; Ins. Philipp.

OSCARIA, Lilja, in Lindbl. Bot. Notiser (1839), 89 = **Primula**, Linn.
chinensis, Lilja, l. c. = P. sinensis.
cortusoides, Lilja, Flora oefver Sverig. odl. Vext. 19 = P. cortusoides.

OSCHATZIA, Walp. Ann. i. 340 (1849) = **Azorella**, Lam. (Umbell.).
Saxifraga, Walp. l. c. = A. Saxifraga.

OSERYA, Tul. & Wedd in Ann. Sc. Nat. Sér. III. xi. (1849) 105. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 114.
biceps, *Tul. & Wedd. l. c.* 106.—Bras.
Coulteriana, *Tul. l. c.*—Mexic.
flabellifera, *Tul. & Wedd. l. c.*—Bras.
sphaerocarpa, *Tul. l. c.*—Guian.

OSKAMPIA, Moench, Meth. 420 (1794) = **Nonnea**, Moench (Borag.).
trichotoma, Moench, l. c. = Alkanna lutea.

OSKAMPIA, Rafin. Sylva Tellur. 123 (1838) = **Tournefortia**, Linn. (Borag.).
hirsuta, Rafin. l. c. = T. hirsutissima ?
scandens, Rafin. l. c. = T. scandens.

OSMADENIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 391 = **Hemizonia**, DC. (Compos.).
tenella, Nutt. l. c. 392 = H. tenella.

OSMANTHUS, Lour. Fl. Cochinch. 28 (1790). *OLEACEAE*, Benth. & Hook. f. ii. 677.
CARTREMA, Rafin. Sylva Tellur. 184 (1838).
acuminata, Wall. Cat. n. 2809 = fragrans.
americana, *Benth. & Hook. f. Gen.* ii. 677.—Am. bor.
aquifolia, C. Koch in Belg. Hortic. xii. (1862) 112; et Dendrol. ii. 1. 278 = seq.
Aquifolium, *Siebold, ex Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 166.—Japon.
fragrans, *Lour. Fl. Cochinch.* 29.—Reg. Himal.; China; Japon.
heterophylla, Hort. ex C. Koch, Dendrol. ii. 1. 278 = O. Aquifolium.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 86 = O. Aquifolium ?
marginata, *Benth. & Hook. f. Gen.* ii. 677.—Ins. Hongkong.
sandwicensis, *Benth. & Hook. f. l. c.*—Ins. Sandvic.
suavis, *King, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 607.—Reg. Himal.

OSMELIA, Thw. Enum. Pl. Zeyl. 20 (1858). *SAMYDACEAE*, Benth. & Hook. f. i. 797.
STACHYCRATER, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 464.
conferta, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 89.—Ins. Philipp.
Gardneri, *Thw. Enum. Pl. Zeyl.* 20.—Zeylan.
philippinensis, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 89.—Ins. Philipp.
zeylanica, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 595 = Gardneri.

OSMIA, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 251 = **Eupatorium**, Tourn. (Compos.).
alternans, Sch. Bip. l. c. 252 = E. laevigatum.
arrayana, Sch. Bip. l. c. 253 = E. arrayanum.
ascendens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 296 = E. ascendens.
asclepiadea, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = E. bupleurifolium.
asperrima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 293 = E. asperrimum.
atriplicifolia, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 252 = E. atriplicifolium.
bartsiaefolia, Sch. Bip. l. c. 253 = E. bartsiaefolium.
Bertholdii, Sch. Bip. l. c. = E. Bertholdii.
caleoides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 282 = E. squalidum.
cinereo-viridis, Sch. Bip. ex Baker, l. c. 294 = E. cinereoviride.
Clematitis, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 252 = E. conyzoides.
conyzoides, Sch. Bip. l. c. = E. conyzoides.
crenata, Sch. Bip. l. c. 253 = E. squalidum.
divergens, Sch. Bip. l. c. 252 = E. conyzoides.
extensa, Sch. Bip. l. c. 253 = E. extensum.
ferruginea, Sch. Bip. l. c. = E. ferrugineum.
floribunda, Sch. Bip. l. c. 252 = E. conyzoides.
foliata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 278 = E. conyzoides.
furcata, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 252 = E. furcatum.
glaberrima, Sch. Bip. l. c. = E. glaberrimum.
graciliflora, Sch. Bip. l. c. = E. graciliflorum.
gracillima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 290 = E. ivaefolium.

OSMIA :—
hexantha, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = Eupatorium bartsiaefolium.
hypericifolia, Sch. Bip. l. c. = E. hypericifolium.
integrifolia, Sch. Bip. l. c. 252 = E. integrifolium.
ivaefolia, Sch. Bip. l. c. 250, 252 = E. ivaefolium.
laevigata, Sch. Bip. l. c. 252 = E. laevigatum.
leptocephala, Sch. Bip. l. c. 253 = E. leptocephalum.
liatridea, Sch. Bip. l. c. 252 = E. liatrideum.
macrantha, Sch. Bip. l. c. = E. macranthum.
macrodon, Sch. Bip. l. c. = E. laevigatum.
Martiusii, Sch. Bip. l. c. = E. squalidum.
Maximiliani, Sch. Bip. l. c. = E. conyzoides.
mucronata, Sch. Bip. l. c. 253 = E. conyzoides.
multiflosculosa, Sch. Bip. l. c. 252 = E. multiflosculosum.
myriocephala, Sch. Bip. l. c. 253 = E. myriocephalum.
obscura, Sch. Bip. l. c. 252 = E. ivaefolium.
odorata, Sch. Bip. l. c. = E. odoratum, conyzoides.
Osseana, Sch. Bip. l. c. = E. Osseanum.
oxylepis, Sch. Bip. l. c. = E. oxylepis.
palmaris, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 294 = E. cinereo-viride.
pedalis, Sch. Bip. ex Baker, l. c. 295 = E. pedale.
perforata, Sch. Bip. ex Baker, l. c. 289 = E. perforatum.
phlogifolia, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = E. phlogifolium.
picta, Sch. Bip. l. c. = E. pictum.
polyantha, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 285 = E. subserratum ?
propinqua, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 252 = E. laevigatum.
pulchella, Sch. Bip. l. c. 253 = E. pulchellum.
punctata, Sch. Bip. l. c. 252 = E. punctatum.
punctulata, Sch. Bip. l. c. 253 = E. punctulatum.
ramosissima, Sch. Bip. l. c. = E. squalidum.
repanda, Sch. Bip. l. c. 252 = E. repandum.
rhinanthacea, Sch. Bip. l. c. 253 = E. rhinanthaceum.
scabra, Sch. Bip. l. c. = E. scabrum.
sericea, Sch. Bip. l. c. 252 = E. sericeum.
serratuloides, Sch. Bip. l. c. = E. serratuloides.
serrulata, Sch. Bip. l. c. 253 = E. serrulatum.
smilacina, Sch. Bip. l. c. 252 = E. smilacinum.
squalida, Sch. Bip. l. c. = E. squalidum.
stenocephala, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 299 = E. punctulatum.
stenophylla, Sch. Bip. ex Baker, l. c. 290 = E. ivaefolium.
subserrata, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = E. subserratum.
subtruncata, Sch. Bip. l. c. = E. subtruncatum.
subvelutina, Sch. Bip. l. c. 252 = E. squalidum.
tecta, Sch. Bip. l. c. 253 = E. tectum.
tomentosa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 282 = E. squalidum.
tozziaefolia, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = E. tozziaefolium.
trichophora, Sch. Bip. l. c. = E. bartsiaefolium.
trifoliata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 282 = E. squalidum.
verbenacea, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 253 = E. verbenaceum.

OSMITES, Linn. Pl. Rar. Afr. 24 (1760); Amoen. Acad. vi. 105 (1764). *COMPOSITAE*, Benth. & Hook. f. ii. 341.
BELLIDIASTRUM, [Vaill.] Less. Syn. Comp. 383 (1832).
BELLIDIOPSIS, Spach. Hist. Vég. Phan. x. 25 (1841).
SPANOTRICHUM, E. Mey. ex DC. Prod. vi. 291 (1837).
angustifolia, *DC. Prod.* vi. 291.—Afr. austr.
anthemoides, *DC. l. c.*—Af. austr.
asteriscoides, Linn. Pl. Rar. Afr. 24; Amoen. Acad. vi. 105 = Osmitopsis asteriscoides.
Bellidiastrum, DC. Prod. vi. 291 (excl. syn.) = hirsuta.
Bellidiastrum, *Linn. Pl. Rar. Afr.* 24; *Amoen. Acad.* vi. 105; *Thunb. Fl. Cap. ed. Schult.* 701.—Afr. austr.
calycina, Linn. f. Suppl. 380 = Peyrousea calycina.
camphorata, Thunb. Prod. Pl. Cap. 163 = Osmitopsis asteriscoides.
camphorina, Gaertn. Fruct. ii. 442. t. 74. f. 3 = dentata.
camphorina, *Linn. Pl. Rar. Afr.* 24; *Amoen. Acad.* vi. 105.—Afr. austr.
camphorina, Thunb. Fl. Cap. ed. Schult. 700 = Osmitopsis asteriscoides.

OSMITES:—
dentata, *Thunb. Prod. Fl. Cap.* 163.—Afr. austr.
hirsuta, *Less. Syn. Comp.* 385.—Afr. austr.
lancea, Thunb. ex Harv. & Sond. Fl. Cap. iii. 304 = O. Bellidiastrum.
parviflora, *DC. Prod.* vi. 291.—Afr. austr.
pinnatifida, *DC. l. c.*—Afr. austr.

OSMITIPHYLLUM, Sch. Bip. in Flora, xxvii. (1844) 675 = **Peyrousea**, DC. (Compos.).
Kraussii, Sch. Bip. l. c. = P. calycina.

OSMITOPSIS, Cass. in Bull. Soc. Philom. (1817) 154; et in Dict. Sc. Nat. xxix. 180 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 341.
asteriscoides, *Cass. in Dict. Sc. Nat.* xxxvii. 6.—Afr. austr.
camphorina, Less. Syn. Comp. 387 = asteriscoides.

OSMODIUM, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Onosmodium**, Michx. (Borag.).
nigrum, Rafin. Atl. Journ. 148 = Onosmodium virginianum?

OSMORHIZA, Rafin. in Journ. Phys. lxxxix. (1819) 257. *UMBELLIFERAE*, Benth. & Hook. f. i. 897.
SCHUDIA, Molina, ex C. Gay, Fl. Chil. iii. 143 (1847).
SPERMATURA, Reichb. Consp. 141 (1828).
URASPERMUM, Nutt. Gen. Am. i. 192 (1818).
amurensis, *F. Schmidt, in Maxim. Prim. Fl. Amur.* 129.—As. or.
Berterii, *DC. Prod.* iv. 232.—Chili.
brachypoda, *Torr. ex Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 89.—Am. bor. occ.
brevistylis, Bong. Vég. Sitcha, 142 = nuda.
brevistylis, *DC. Prod.* iv. 232.—Reg. Himal.; As. bor.; Am. bor.
chilensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. 355.—Chili.
Claytoni, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 690 = brevistylis.
cordata, Rafin. Med. Bot. ii. 249 = longistylis.
divaricata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 639 = nuda.
dulcis, Rafin. Med. Bot. ii. 249 = brevistylis.
edulis, Rafin. l. c. = Carum Gairdneri.
glabrata, *Phil. in Linnaea*, xxviii. (1856) 653.—Chili.
japonica, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 203.—Japon.
laxa, Royle, Illustr. Bot. Himal. 233 = brevistylis.
longistylis, *DC. Prod.* iv. 232.—Am. bor.
mexicana, *Griseb. in Goett. Abh.* xxiv. (1879) 147.—Reg. Andin.
nuda, *Torr. in Pacif. Rail. Rep.* iv. 93.—Am. bor. occ.
occidentalis, Torr. Bot. Mex. Bound. 71 = Myrrhis occidentalis.
villosa, Rafin. Med. Bot. ii. 249 = longistylis.

OSMOSCLERIA, Lindl. Veg. Kingd. 119 (1847) = OMOSCLERIA, Nees = **Scleria**, Berg. (Cyperac.).

OSMOTHAMNUS, DC. Prod. vii. 715 (1839) = **Rhododendron**, Linn. (Ericac.).
fragrans, DC. l. c. = R. Anthopogon.
pallidus, DC. l. c.—R. Anthopogon.

OSMOXYLON, Miq. Ann. Mus. Lugd. Bat. i. 5 (1863). *ARALIACEAE*, Benth. & Hook. f. i. 944.
amboinense, *Miq. l. c.*—Ins. Amboin.
barbatum, *Becc. Malesia*, i. (1878) 197.—Malaya.
carpophagarum, *Becc. l. c.* 196.—Malaya.
Geelvinkianum, *Becc. l. c.*—Malaya.
helleborinum, *Becc. l. c.* 198.—Borneo.
Insidiator, *Becc. l. c.* 195.—N. Guin.
insigne, *Becc. l. c.*—Ins. Molucc.
moluccanum, *Becc. l. c.*—Ins. Amboin.
novo-guineense, *Becc. l. c.* 197.—N. Guin.
Zippelianum, *Becc. l. c.* 195.—Ins. Amboin.

OSMYNE, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Tourn. (Liliac.).
odorata, Salisb. l. c. = Ornith. suaveolens.

OSSAEA, DC. Prod. iii. 168 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 770.
DICLEMIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 102.
GONEMA, Rafin. Sylva Tellur. 96 (1838).
SAGRAEA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 92.

OSSAEA:—
acuminata, *DC. Prod.* iii. 169.—Ins. S. Doming.
amygdalina, DC. l. c. = Mecranium amygdalinum.
amygdaloides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bras.
angustifolia, *Triana, l. c.*—Bras.
asperifolia, *Triana, l. c.*—Ind. occ.
birimosa, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 335 = Mecranium birimosum.
brachystachya, Naud. l. c. 337 = Mecranium integrifolium.
brachystachya, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bras.
bracteata, *Triana, l. c.*—N. Granat.
capillaris, *Wright, in Sauv. Fl. Cub.* 46.—Cuba.
cinnamomifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bras.
confertiflora, *Triana, l. c.*—Bras.
coriacea, *Triana, l. c.*—Bras.
cuspidata, Poepp. ex Triana, l. c. 137 = Sagraea sessiliflora.
echinata, Benth. & Hook. f. Gen. i. 762 = Calycogonium echinatum.
Epibaterium, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bras.
euphorbioides, *Triana, l. c.*—Bras.
fascicularis, Griseb. Fl. Brit. W. Ind. 246 = Henriettella fascicularis.
flavescens, DC. Prod. iii. 169 = Henriettella flavescens.
glomerata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 146.—Ind. occ.
Grisebachii, *Triana, l. c.*—Cuba.
heteronervis, *Triana, l. c.*—Peruv.
hirsuta, *Triana, l. c.*—Jamaic.
hirtella, *Triana, l. c.*—Jamaic.
hirtella, Wright, in Sauv. Fl. Cub. 46 = Wrightii?
hypoglauca, Benth. & Hook. f. Gen. i. 762 = Calycogonium hypoglaucum.
integrifolia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 337 = Mecranium integrifolium.
involucrata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Jamaic.
lanata, *Triana, l. c.* 146; *Wright, in Sauv. Fl. Cub.* 45.—Cuba.
lateriflora, DC. Prod. iii. 169 = Henriettella lateriflora.
leptopus, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Peruv.
Lima, *Triana, l. c.*—Jamaic.
marginata, *Triana, l. c.*—Bras.
micrantha, Macfad. Fl. Jamaic. ii. 49, 100 = Octopleura micrantha.
microphylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 146.—Jamaic.
microphylla, *Wright, in Sauv. Fl. Cub.* 45.—Cuba.
multiflora, DC. Prod. iii. 169 = Mecranium multiflorum.
muricata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Cuba.
neurotricha, *Wright, in Sauv. Fl. Cub.* 46.—Cuba.
penninervis, *Wright, l. c.*—Cuba.
petiolaris, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bolivia.
pratensis, *Macfad. Fl. Jamaic.* ii. 48.—Jamaic.
punctata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Henriettella punctata.
purpurascens, DC. Prod. iii. 169 = Mecranium purpurascens.
purpurascens, Macfad. Fl. Jamaic. ii. 49 = Mecranium virgatum.
retropila, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 147.—Bras.
rufescens, *Triana, l. c.* 146; *Wright, in Sauv. Fl. Cub.* 45.—Cuba.
rufibarbis, *Triana, l. c.* 147.—N. Granat.
salicifolia, Crueg. in Linnaea, xx. (1847) 105 = Sagraea salicifolia.
scabrosa, *DC. Prod.* iii. 169.—Jamaic.
scalpta, *DC. l. c.* 168.—Ins. S. Doming.
sparsiflora, DC. l. c. 169 = Henriettella lateriflora.
trichopetala, *Wright, in Sauv. Fl. Cub.* 45.—Cuba.
trinitensis, Crueg. in Linnaea, xx. (1847) 105 = Sagraea trinitensis.
verrucosa, Benth. & Hook. f. Gen. i. 762 = Calycogonium verrucosum.
verticillata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 146.—Ind. occ.
Wrightii, *Triana, l. c.*—Cuba.

OSTACHYRIUM, Steud. Nom. ed. II. ii. 236 (1841) = OTACHYRIUM, Nees = **Panicum**, Tourn. (Gramin.).

OSTEIZA, Steud. Nom. ed. II. ii. 236 (1841) = OTEIZA, La Llave = **Calea**, Linn. (Compos.).

OSTEOCARPUM, F. Muell. Second Gen. Rep. 15 (1854) = **Threlkeldia**, R. Br. (Chenop.).
salsuginosum, F. Muell. l. c. = T. salsuginosa.

OSTEOCARPUS, Phil. in Regel, Gartenfl. (1884) 39, 356. *SOLANACEAE*.
rostratus, *Phil. l. c.* 356.—Chili.

OSTEOMELES, Lindl. in Trans. Linn. Soc. xiii. (1821) 98. t. 8. *ROSACEAE*, Benth. & Hook. f. i. 628.
ELEUTHEROCARPUM, Schlecht. in Lechl. Berb. Am. Austr. 59 (1857).
HESPEROMELES, Lindl. Bot. Reg. sub t. 1956 (1837).
anthyllidifolia, *Lindl. in Trans. Linn. Soc.* xiii. (1821) 99. t. 1.—Ins. Pacif.; China.
anthyllifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 131 = praec.
cordata, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 181 = latifolia.
escalloniaefolia, *Decne. l. c.* 184.—Peruv.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* vi. 211.—Ecuador.
Gayana, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 183.—Peruv.
glabrata, *H. B. & K. Nov. Gen. et Sp.* vi. 210. t. 553.—N. Granat.
Goudotiana, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 182.—Peruv.
latifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 212. t. 554.—Peruv.
obtusifolia, Kunth, ex Steud. Nom. ed. II. i. 433; ii. 236 = glabrata.
Pentlandiana, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 183.—Peruv.
pernettyoides, *Decne. l. c.* 184.—N. Granat.; Bolivia; Peruv.
Persoonii, H. B. & K. Nov. Gen. et Sp. vi. 211, in obs. = glabrata.
pyracantha, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 183 = Photinia Fortuneana.
rufescens, *Decne. l. c.* 181.—N. Granat.
subrotunda, C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 250 = anthyllidifolia.

OSTEOSPERMUM, Linn. Gen. ed. I. 264 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 455.
ERIOCLINE, Cass. in Dict. Sc. Nat. xv. 191 (1819).
GIBBARIA, Cass. in Bull. Soc. Philom. (1817) 139; et in Dict. Sc. Nat. xviii. 526 (1820).
LEPISIPHON, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 180.
MONILIFERA, Adans. Fam. ii. 127 (1763).
XEROTHAMNUS, DC. Prod. v. 311 (1836).
aciphyllum, DC. Prod. vi. 463 = triquetrum.
amplexicaule, Steud. Nom. ed. II. ii. 236 = corymbosum.
arctotioides, Linn. f. Suppl. 385 = Venidium arctotioides.
arctotioides, Thunb. Prod. Pl. Cap. 167 = Venidium hispidulum.
aritotoides, J. F. Gmel. Syst. 1595, sphalm. = praec.
asperum, *Less. Syn. Comp.* 89.—Afr. austr.
Bidens, *Thunb. Prod. Pl. Cap.* 166.—Afr. austr.
bipinnatum, Thunb. l. c. 167 = Garuleum bipinnatum.
Burchellii, *DC. Prod.* vi. 461.—Afr. austr.
caeruleum, Jacq. Coll. i. 78 = Garuleum viscosum.
calendulaceum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 440.—Afr. austr.
calendulaceum, Harv. l. c. 626 (Index), sphalm. = Venidium hirsutum.
calendulaceum, Linn. f. Suppl. 386 = Oligocarpus calendulaceus.
caulescens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 442.—Afr. austr.
ciliatum, *Berg. Desc. Pl. Cap.* 332.—Afr. austr.
coeruleum, Auct. vide *caeruleum*.
connatum, DC. Prod. vi. 466 = Tripteris amplexicaulis.
coriaceum, *DC. l. c.* 461.—Afr. austr.
corymbosum, *Linn. Mant.* ii. 290.—Afr. austr.
cuspidatum, DC. Prod. vi. 463 = Tripteris tomentosa.

OSTEOSPERMUM :—
dentatum, Burm. f. Fl. Cap. Prod. 29 (err. typ. 25) = Tripteris arborescens.
dichotomum, E. Mey. ex DC. Prod. vi. 462 = corymbosum.
ephedroides, DC. l. c. 463 = O. Bidens.
falcatum, E. Mey. ex l. c. 466 = Gymnostephium ciliare.
fallax, Spreng. ex DC. l. c. 467 = Gamolepis munita.
floridum, E. Mey. ex DC. l. c 73 = Dimorphotheca chrysanthemifolia.
foveolatum, DC. l. c. 462 = imbricatum.
glabratum, Less. Syn. Comp. 89 = Tripteris macroptera.
glandulosum, *DC. Prod.* vi. 463.—Afr. austr.
glandulosum, Spreng. f. Tent. Suppl. Syst. 26 = corymbosum.
grandidentatum, *DC. Prod.* vi. 465.—Afr. austr.
grandiflorum, *DC. l. c.* 464.—Afr. austr.
helichrysoides, *DC. l. c.* 466.—Afr. austr.
herbaceum, *Linn. f. Suppl.* 385.—Afr. austr.
heterophyllum, DC. Prod. vi. 467 = Gamolepis brachypoda.
hirsutum, *Thunb. Prod. Pl. Cap.* 166.—Afr. austr.
hispidum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 438.—Afr. austr.
ilicifolium, *Linn. Syst.* ed. X. 1234.—Afr. austr.
imbricatum, *Linn. Mant.* ii. 290.—Afr. austr.
incanum, *Burm. f. Fl. Cap. Prod.* 29 (err. typ. 25) (Quid?).—Afr. austr.
incanum, Thunb. Prod. Pl. Cap. 166 = Tripteris incana.
junceum, Berg. Desc. Pl. Cap. 334 = Phymaspermum junceum.
junceum, Thunb. ex Harv. & Sond. Fl. Cap. iii. 437 = coriaceum.
lanatum, Spreng. ex Sch. Bip. in Flora, xxvii. (1844) 768 = corymbosum.
lanceolatum, *DC. Prod.* vi. 463.—Afr. austr.
lavandulaceum, *DC. l. c.* 466.—Afr. austr.
laxum, *DC. l. c.* 465.—Afr. austr.
leiocarpum, *DC. l. c.* 467.—Afr. austr.
macrocarpum, DC. l. c. 461 = coriaceum.
marginatum, *Spreng. f. Tent. Suppl. Syst.* 27.—Afr. austr.
microphyllum, *DC. Prod.* vi. 463.—Afr. austr.
moniliferum, *Linn. Sp. Pl.* 923.—Afr. austr.
muricatum, *E. Mey. in DC. Prod.* vi. 464.—Afr. austr.
natalense, Sch. Bip. ex Harv. & Sond. Fl. Cap. iii. 440 = grandidentatum.
nervatum, *DC. Prod.* vi. 462.—Afr. austr.
niveum, Linn. f. Suppl. 386 = Cryptostemma niveum.
pachypteris, DC. Prod. vi. 460 = Tripteris pachypteris.
paniculatum, Less. ex Harv. & Sond. Fl. Cap. iii. 437 = coriaceum.
perfoliatum, Linn. f. Suppl. 386 = Venidium perfoliatum.
perfoliatum, Thunb. ex Harv. & Sond. Fl. Cap. iii. 460 = Venidium discolor.
picridioides, *DC. Prod.* vi. 465.—Afr. austr.
piliferum, Thunb. Fl. Cap. ed. Schult. 715 = *pisiferum*, Linn.
pinifolium, *Regel, Cat. Pl. Hort. Aksakov.* 99.—Afr. austr.
pinnatifidum, L'Hérit. Stirp. Nov. 11 = Garuleum viscosum.
pinnatipartitum, *DC. Prod.* vi. 465.—Afr. austr.
pisiferum, Burm. ex DC. Prod. vi. 461 = coriaceum.
pisiferum, Linn. Syst. ed. X. 1234 = moniliferum.
plebeium, *DC. Prod.* vi. 464.—Afr. austr.
polygaloides, *Linn. Sp. Pl.* 924.—Afr. austr.
polygaloides, Thunb. Prod. Pl. Cap. 166 = imbricatum.
pterospermum, *E. Mey. ex DC. Prod.* vi. 460.—Afr. austr.
pungens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 437.—Afr. austr.
retirugum, *DC. Prod.* vi. 462.—Afr. austr.
rigidum, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 275.—Afr. austr.
rotundatum, DC. Prod. vi. 461 = moniliferum.
rubens, Burm. f. Fl. Cap. Prod. 29 (err. typ. 25) = ciliatum.
scabridum, *DC. Prod.* vi. 466.—Afr. austr.
scabrum, *Thunb. Prod. Pl. Cap.* 166.—Afr. austr.
scariosum, DC. Prod. vi. 464 = Tripteris flexuosa.
sonchifolium, *DC. l. c.* 465.—Afr. austr.
spinescens, DC. l. c. 459 = moniliferum.
spinescens, Thunb. ex DC. l. c 137 = Pentzia spinescens.
spinescens, Thunb. Fl. Cap. ed. Schult. 713 = Tripteris spinescens.

OSTEOSPERMUM :—
spinescens, *Thunb. Prod. Pl. Cap.* 166.—Afr. austr.
spinescens, Willd. Sp. Pl. iii. 2365 = spinosum.
spinosum, Jacq. Hort. Schoenb. iii. 66 = spinescens.
spinosum, *Linn. Sp. Pl.* 923.—Afr. austr.
squarrosum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 440.—Afr. austr.
subauritum, *DC. Prod.* vi. 464.—Afr. austr.
subcanescens, *DC. l. c.* 464.—Afr. austr.
subulatum, *DC. l. c.* 463.—Afr. austr.
tenuifolium, Harv. & Sond. Fl. Cap. iii. 607, sphalm. = *teretifolium*.
tenuilobum, DC. Prod. vi. 467 = Euryops punctatus.
teretifolium, Thunb. Prod. Pl. Cap. 166 = Pteronia baccharoides, Gamolepis trifurcata?
thymelaeoides, *DC. Prod.* vi. 462.—Afr. austr.
tricuspe, *E. Mey. ex DC. l. c.* 467 = tridens.
tridens, *DC. l. c.*—Afr. austr.
trigonospermum, *DC. l. c.* 460.—Afr. austr.
trigonum, Spreng. Syst. iii. 628 = Pteronia baccharoides.
tripteroideum, DC. Prod. vi. 463 = Tripteris spinescens.
triquetrum, DC. l. c. 465 = ciliatum.
triquetrum, *Linn. f. Suppl.* 385.—Afr. austr.
Uvedalia, *Linn. Sp. Pl.* 923.—Afr. austr.
Zeyheri, Spreng. ex DC. Prod. vi. 466 = herbaceum.

OSTERDAMIA, Neck. Elem. iii. 218 (1790) = **Zoysia**, Willd. (Gramin.).

OSTERDICKIA, Burm. Rar. Afr. Pl. 259. t. 96 (1737) = **Cunonia**, Linn. (Saxifr.).

OSTERDYCKIA, Reichb. Consp. 159 (1828) = praec.

OSTERICUM, Hoffm. Gen. Umb. ed. I. 262, in nota (1814); ed. II. 162 (1816) = **Angelica**, Riv. (Umbell.).
palustre, Bess. Enum. Pl. Volh. 94 = A. palustris.
pratense, Hoffm. Gen. Umb. ed. II. 164 = A. palustris.
verticillare, Reichb. Fl. Germ. Excurs. 455 = Peucedanum verticillare.

OSTINIA, [Clairv.] Man. Herb. 162 (1811) (Rosac.).
Cotoneaster, [Clairv.] l. c. = Cotoneaster integerrima.
lanata, [Clairv.] l. c.—Europ.
Mespylus, [Clairv.] l. c. = Pyrus germanica.
Oxyacantha, [Clairv.] l. c. = Crataegus Oxyacantha.

OSTODES, Blume, Bijdr. 619 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 299.
DESMOSTEMON, Thw. Enum. Pl. Zeyl. 278 (1861).
FAHRENHEITIA, Reichb. f. & Zoll. in Linnaea, xxviii. (1856) 600.
corniculata, Baill. Etud. Gén. Euph. 391 = paniculata.
Helferi, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 215.—Burma.
macrophylla, *Benth. & Hook. f. Gen.* iii. 299.—Malacca.
minor, Muell. Arg. in Linnaea, xxxiv. (1865–66) 214 = zeylanica.
paniculata, *Blume, Bijdr.* 620.—Reg. Himal.; Burma; Java.
zeylanica, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 214.—Ind. or.

OSTROWSKIA, Regel, in Act. Hort. Petrop. viii. (1884) 686. *CAMPANULACEAE*.
magnifica, *Regel, l. c.*—As. centr.

OSTRUTHIUM, Link, Handb. i. 360 (1829) = **Peucedanum**, Tourn. (Umbellif.).
officinale, Link, l. c. = P. Ostruthium.

OSTRYA, Mich. ex Linn. Syst. ed. I (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 406.
ZUGILUS, Rafin. Fl. Ludov. 159 (1817).
americana, Michx. f. Hist. Arb. Am. iii. 54 = virginica.
carpinifolia, *Scop. Fl. Carn.* ed. II. ii. 244.—Europ. austr.; As. Min.
italica, Spach, Hist. Vég. Phan. xi. 216 = carpinifolia.
mandshurica, *Budisch. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 166, nomen.—As. or.
virginiana, C. Koch, Dendrol. ii. II. 6 = seq.
virginica, *Willd. Sp. Pl.* iv. 469.—Am. bor.
vulgaris, Wats. Dendr. t. 143 = virginica.
vulgaris, Willd. Sp. Pl. iv. 469 = carpinifolia.

OSTRYOCARPUS, Hook. f. in Hook. Niger Fl. 316 (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 548.
riparius, *Hook. f. in Hook. Niger Fl.* 316.—Afr. trop.
Welwitschii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 240.—Afr. trop.

OSTRYODIUM, Desv. Journ. Bot. iii. (1814) 119. t. 4 = **Flemingia**, Roxb. (Legumin.).

OSTRYOPSIS, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 155. *CUPULIFERAE*, Benth. & Hook. f. iii. 405.
Davidiana, *Decne. l. c.*—As. or.

OSWALDA, Cass. in Dict. Sc. Nat. lix. 322 (1829) = **Clibadium**, Linn. (Compos.).
baillierioides, Cass. l. c. = C. surinamense.

OSWALDIA, Less. Syn. Comp. 215 (1832) = praec.

OSYRICERA, Blume, Bijdr. 307. t. 58 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 505.
crassifolia, *Blume, l. c.*—Java.

OSYRIDICARPOS, A. DC. in DC. Prod. xiv. 635. (1857). *SANTALACEAE*, Benth. & Hook. f. iii. 222.
natalensis, *A. DC. l. c.*—Afr. austr.
Schimperianus, *A. DC. l. c.*—Abyssin.

OSYRIS, Linn. Syst. ed. I (1735). *SANTALACEAE*, Benth. & Hook. f. iii. 227.
abyssinica, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 22, nomen.—Abyssin.
alba, *Linn. Sp. Pl.* 1022.—Reg. Mediterr.
arborea, *Wall. Cat.* n. 4035.—Ind. or.
compressa, A. DC. in DC. Prod. xiv. 634 = Colpoon compressum.
japonica, Thunb. Fl. Jap. 31 = Helwingia rusciflora.
lanceolata, *Hochst. & Steud. ex A. DC. in DC. Prod.* xiv. 633.—Hispan.
nepalensis, Griff. Notul. iv. 376 = arborea.
peltata, Roxb. Hort. Beng. 71; Fl. Ind. iii. 755 = Macaranga Roxburghii.
pendula, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 93.—Ins. Socotra.
quadrifida, Salzm. ex A. DC. in DC. Prod. xiv. 633 = lanceolata.
quadripartita, Salzm. ex Decne. in Ann. Sc. Nat. Sér. II. vi. (1836) 65 = lanceolata.
rhamnoides, Scop. Fl. Carn. ed. II. ii. 261 = Hippophaë rhamnoides.
rotundata, Griff. Notul. iv. 742; Ic. Pl. Asiat. t. 627. f. 11 = Henslowia buxifolia.
Schoberi, Pall. ex A. DC. in DC. Prod. xiv. 634 = Nitraria Schoberi.
spinescens, *Mart. & Eichl. in Mart. Fl. Bras.* xiii. I. 236.—Bras.
Wightiana, Wall. Cat. n. 4036 = arborea.

OTACANTHUS, Lindl. in Fl. des Serres, xv. (1862?) 53. t. 1526. *ACANTHACEAE*, Benth. & Hook. f. ii. 1076.
caeruleus, *Lindl. l. c.*—Bras.

OTACHYRIUM, Nees, Agrost. Bras. 273 (1829) = **Panicum**, Tourn. (Gramin.).
junceum, Nees, l. c. 272 = P. Pterygodium.

OTAMPLIS, Rafin. New Fl. Am. iv. 8 (1836) = **Menispermum**?, Tourn.
vitifolia, Rafin. l. c. = M. canadense?

OTANDRA, Salisb. in Trans. Hort. Soc. i. (1812) 298 = **Geodorum**, Jacks. (Orchid.).
cernua, Salisb. l. c. = G. dilatatum.

OTANEMA, Rafin. New Fl. Am. iv. 61 (1826) = **Asclepias**, Tourn.
lanceolata, Rafin. l. c. = A. paupercula?
latifolia, Rafin. l. c. = A. latifolia.
ovata, Rafin. l. c. = Vincetoxicum officinale?

OTANTHERA, Blume, in Flora, xiv. (1831) 488. *MELASTOMACEAE*, Benth. & Hook. f. i. 746.
LACHNOPODIUM, Blume, in Flora, xiv. (1831) 477, in nota; et Mus. Bot. Lugd. Bat. i. 56 (1849).
MALABATHRIS, Rafin. Sylva Tellur. 97 (1838).

OTANTHERA :—
bracteata, *Korth. Verh. Nat. Gesch. Bot.* 235. t. 51.—Sumatra.
celebica, *Blume, Mus. Bot. Lugd. Bat.* i. 56.—Ins. Celebes.
crinita, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 354.—Ins. Philipp.
cyanoides, Triana, in Trans. Linn. Soc. xxviii. (1871) 55 = moluccana.
Fordii, *Hance, in Journ. Bot.* xix. (1881) 46.—Ins. Hongkong.
gracilis, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 353.—Ins. Amboin.
manillana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 56.—Ins. Philipp.
moluccana, *Blume, in Flora*, xiv. (1831) 489.—Ins. Molucc.
nicobarensis, *Teijsm. & Binn. in Tijdschr. Nederl. Indie*, xxv. (1863) 427.—Ins. Nicobar.
rubro-limbata, Triana, in Trans. Linn. Soc. xxviii. (1871) 56 = bracteata ?

OTANTHUS, Hoffmgg. & Link, Fl. Port. ii. 364 (1809) = **Diotis**, Desf. (Compos.).
maritimus, Hoffmgg. & Link, l. c. 365 = D. candidissima.

OTARIA, Kunth, in H. B. & K. Nov. Gen. et Sp. iii. 192 (1818) = **Asclepias**, Linn.
auriculata, G. Don, Gen. Syst. iv. 144 = A. auriculata.
fragrans, Benth. Pl. Hartw. 89 = A. fragrans.
laxiflora, Benth. l. c. 67 = A. laxiflora.

OTEIZA, La Llave, Reg. Trim. Mex. (1832) 41 ; ex DC. Prod. vii. 262 = **Calea**, Linn. (Compos.).
acuminata, La Llave, l. c. = C. elegans.

OTHAKE, Rafin. New Fl. Am. iv. 73 (1836) = **Gaillardia**, Fouger. (Compos.).
longifolium, Rafin. l. c. 74.—Am. bor.
tenuifolium, Rafin. l. c. = Polypteris callosa.

OTHANTHERA, G. Don, Gen. Syst. iv. 107 (1837), sphalm. = **Orthanthera**, Wight (Asclep.).

OTHERA, Thunb. Nov. Gen. 56 (1783) = **Ilex**, Linn. (Ilicin.).
japonica, Thunb. l. c. = I. Othera.
Orixa, Lam. Illustr. i. 310 = Celastrus Orixa.

OTHLIS, Schott, in Spreng. Syst. iv. App. 407 (1827) = **Doliocarpus**, Roland. (Dillen.).
castaneaefolia, Schott, l. c. = D. grandiflorus.

OTHOCALLIS, Salisb. Gen. Pl. Fragm. 28 (1866) = **Scilla**, Linn. (Liliac.).
amoena, Salisb. l. c. = S. amoena.
sibirica, Salisb. l. c. = S. sibirica.

OTHONNA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 453.
ARISTOTELA, Adans. Fam. ii. 125 (1763).
CALTHOIDES, B. Juss. ex DC. Prod. vi. 473 (1837).
CERADIA, Lindl. Bot. Reg. (1845) Misc. 12.
DORIA, Less. in Linnaea, vi. (1831) 94.
abrotanifolia, Linn. Sp. Pl. 926 = Euryops abrotanifolius.
acutiloba, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
adscendens, DC. Prod. vi. 480 = tuberosa.
alata, Sch. Bip. in Flora, xxvii. (1844) 769 = Othonnopsis alata.
amplexicaulis, Sims, Bot. Mag. t. 1312 = amplexifolia.
amplexicaulis, Thunb. Prod. Pl. Cap. 167 = denticulata.
amplexifolia, *DC. Prod.* vi. 480.—Afr. austr.
angustifolia, DC. l. c. 477 = Othonnopsis angustifolia.
arborescens, *Linn. Sp. Pl.* ed. II. 1310.—Afr. austr.
Arbuscula, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
Athanasiae, Jacq. Hort. Schoenb. ii. 62. t. 242 = Euryops Athanasiae.
Athanasiae, Linn. f. Suppl. 386 = Euryops abrotanifolius.
auriculaefolia, *Licht. ex Less. in Linnaea*, vi. (1831) 93.—Afr. austr.
Bainesii, Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 423 = coriifolia.

OTHONNA :—
Bergeri, Spreng. Syst. iii. 626 = coronopifolia.
brachypoda, DC. Prod. vi. 480 = heterophylla.
bulbosa, Linn. Sp. Pl. 924 = sp. plurim.
bupleuroides, DC. Prod. vi. 475 = parviflora.
cacalioides, *Linn. f. Suppl.* 388.—Afr. austr.
cakilefolia, *DC. Prod.* vi. 482.—Afr. austr.
Calthoides, Mill. Gard. Dict. ed. VIII. n. 2 = Othonnopsis cheirifolia.
campanulata, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
capillaris, Linn. f. Suppl. 388 = Gymnodiscus capillaris.
carnosa, *Less. Syn. Comp.* 88.—Afr. austr.
cheirifolia, Linn. Sp. Pl. 926 = Othonnopsis cheirifolia.
chromachaeta, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
ciliata, Linn. f. Suppl. 386 = Othonnopsis ciliata.
Cineraria, *Linn. Sp. Pl.* 925.—Afr. austr.
cluytiaefolia, Sch. Bip. in Flora, xxvii. (1844) 769 = Othonnopsis cluytiaefolia.
cneorifolia, Sch. Bip. l. c. = Othonnopsis cneorifolia.
coriifolia, *Sond. in Harv. & Sond. Fl. Cap.* iii. 333.—Afr. austr.
coronopifolia, *Linn. Sp. Pl.* 926.—Afr. austr.
coronopifolia, Thunb. Prod. Pl. Cap. 167 = arborescens.
crassifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 336.—Afr. austr.
crassifolia, Linn. Mant. i. 118 = Othonnopsis cheirifolia.
crassifolia, E. Mey. ex DC. Prod. vi. 476 = denticulata.
crispa, *Roxb. Hort. Beng.* 62, nomen.—Afr. austr.
cuneata, *DC. Prod.* vi. 481.—Afr. austr.
cyanoglossa, DC. l. c. = auriculaefolia.
cylindrica, *DC. l. c.* 477.—Afr. austr.
cymbalarifolia, Linn. Pl. Rar. Afr. 24 = Senecio cymbalarifolius.
dentata, *Linn. Sp. Pl.* 926.—Afr. austr.
denticulata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 276.—Afr. austr.
digitata, *Linn. f. Suppl.* 386.—Afr. austr.
diversifolia, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
Dregeana, DC. Prod. vi. 475 = denticulata.
Eckloneana, DC. l. c. 476.—Afr. austr.
elliptica, DC. l. c. 480 = tuberosa.
ericoides, Linn. f. Suppl. 388 ; Thunb. Prod. Pl. Cap. 167 = Gamolepis ericoides.
eriocarpa, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
filicaulis, Eckl. ex Harv. & Sond. Fl. Cap. iii. 336 = crassifolia.
filicaulis, *Jacq. Hort. Schoenb.* ii. 62. t. 241.—Afr. austr.
filifolia, Burm. f. ex DC. Prod. vi. 278 = Leyssera gnaphaloides.
flabellifolia, Lodd. Bot. Cab. t. 728 = Euryopus virgineus.
frutescens, Drège, ex Harv. & Sond. Fl. Cap. iii. 331 = leptodactyla.
frutescens, Eckl. ex Harv. & Sond. Fl. Cap. iii. 323 = Othonnopsis Kraussii.
frutescens, *Linn. Mant.* ii. 288.—Afr. austr.
geifolia, *Linn. Sp. Pl.* 924.—Afr. austr.
glauca, Spreng. ex Sch. Bip. in Flora, xxvii. (1844) 696 = Senecio othonnaeflorus.
grandidentata, DC. Prod. vi. 474 = pluridentata.
Gymnodiscus, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
helenitis, Linn. Sp. Pl. 925 = Senecio brachychaetus.
heterophylla, *Linn. f. Suppl.* 387.—Afr. austr.
imbricata, Thunb. Prod. Pl. Cap. 167 = Euryops lateriflorus.
incisa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 341.—Afr. austr.
integrifolia, *Linn. Sp. Pl.* 925.—Afr. austr.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 769 = Othonnopsis Kraussii.
lactucaefolia, *DC. Prod.* vi. 482.—Afr. austr.
lasiocarpa, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
lateriflora, Linn. f. Suppl. 387 ; Thunb. Prod. Pl. Cap. 167 = Euryops lateriflorus.
Laureola, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
laxa, *Savign. in Lam. Encyc.* iv. 668.—Afr. austr.
leptodactyla, *Harv. in Harv. & Sond. Fl. Cap.* iii. 331.—Afr. austr.
leptophylla, *DC. Prod.* vi. 477.—Afr. austr.

OTHONNA :—
lineariifolia, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
Lingua, Herb. Thunb. ex Harv. & Sond. Fl. Cap. iii. 342 = tuberosa.
Lingua, *Linn. f. Suppl.* 387 ; *Thunb. Prod. Pl. Cap.* 167.—Afr. austr.
linifolia, Burm. f. Fl. Cap. Prod. 29 (err. typ. 25) = Euryops tenuissimus.
linifolia, *Linn. f. Suppl.* 388.—Afr. austr.
longifolia, DC. Prod. vi. 476 = parviflora.
lyrata, *DC. l. c.* 481.—Afr. austr.
macrophylla, *DC. l. c.* 480.—Afr. austr.
macrosperma, *DC. l. c.* 478.—Afr. austr.
maritima, *Linn. Sp. Pl.* 925.—Afr. austr.
membranifolia, *DC. Prod.* vi. 476.—Afr. austr.
minima, *DC. l. c.* 479.—Afr. austr.
mucronata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 331.—Afr. austr.
multicaulis, *Harv. l. c.* 333.—Afr. austr.
multifida, Thunb. Prod. Pl. Cap. 168 = Euryops multifidus.
munita, Linn. f. Suppl. 388 = Gamolepis munita.
nana, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
natalensis, *Sch. Bip. l. c.* 771.—Afr. austr.
nodulosa, *Sch. Bip. l. c.* 769.—Afr. austr.
nudata, *DC. Prod.* vi. 475 (sp. dub.).—Afr. austr.
nudicaulis, *Savign. in Lam. Encyc.* iv. 668.—Afr. austr.
oblongifolia, DC. Prod. vi. 480 = tuberosa.
obtusiloba, *Harv. in Harv. & Sond. Fl. Cap.* iii. 337.—Afr. austr.
osteospermoides, *DC. Prod.* vi. 474.—Afr. austr.
pallens, DC. l. c. 476 = Othonnopsis pallens.
palustris, Linn. Sp. Pl. 924 = Senecio palustris.
parviflora, Berg. Desc. Pl. Cap. 335 = denticulata.
parviflora, *Linn. Mant.* i. 89.—Afr. austr.
Pavonia, *E. Mey. ex DC. Prod.* vi. 473.—Afr. austr.
pectinata, Linn. Sp. Pl. 926 = Euryops pectinatus.
pectinata, E. Mey. ex DC. Prod. vi. 484 = Ruckeria othonnoides.
perfoliata, *Jacq. Hort. Schoenb.* ii. 61. t. 240.—Afr. austr.
petiolaris, *DC. Prod.* vi. 480.—Afr. austr.
picridioides, DC. l. c. 481 = auriculaefolia.
pinnata, *Linn. f. Suppl.* 387.—Afr. austr.
pinnatifida, Thunb. Prod. Pl. Cap. 168 = Cineraria othonnoides.
pinnatilobata, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
pluridentata, *DC. Prod.* vi. 474.—Afr. austr.
primulina, *DC. l. c.* 479.—Afr. austr.
procera, Salisb. Prod. 211 = arborescens.
pteronioides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 332.—Afr. austr.
purpurascens, *Harv. l. c.* 331.—Afr. austr.
quercifolia, *DC. Prod.* vi. 478.—Afr. austr.
quinquedentata, DC. l. c. 474 = parviflora.
quinquedentata, *Thunb. Prod. Pl. Cap.* 168.—Afr. austr.
quinqueradiata, *DC. Prod.* vi. 478.—Afr. austr.
racemosa, E. Mey. ex DC. l. c. 446 = Euryops oligoglossus.
ramosa, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
ramulosa, *DC. Prod.* vi. 477.—Afr. austr.
reticulata, *DC. l. c.* 481.—Afr. austr.
retrofracta, *Jacq. Hort. Schoenb.* iii. 66. t. 376.—Afr. austr.
retrorsa, *DC. Prod.* vi. 479.—Afr. austr.
rhamnoides, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
rigens, *Linn. Pl. Afr. Rar.* 24.—Afr. austr.
rigida, Burm. ex DC. Prod. vi. 510 = Gazania Pavonia.
rigida, DC. l. c. 476 = denticulata.
rosea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 341.—Afr. austr.
rotundifolia, Harv. l. c. 344, sphalm. = seq.
rotundiloba, *DC. Prod.* vi. 481.—Afr. austr.
rufibarbis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 341.—Afr. austr.
scapigera, *Harv. Thes. Cap.* i. 10.—Afr. austr.
sedifolia, *DC. Prod.* vi. 479.—Afr. austr.
semicalva, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
senecionea, DC. Prod. vi. 475 = denticulata.
sibirica, Linn. Sp. Pl. 924 = Senecio Ligularia.
sonchifolia, *DC. Prod.* vi. 482.—Afr. austr.
sonchifolia, Linn. Sp. Pl. 924 = Cineraria sonchifolia.

OTHONNA :—
speciosa, Spreng. ex Steud. Nom. ed. II. ii. 237 = Euryops tenuissimus.
spinescens, *DC. Prod.* vi. 476.—Afr. austr.
spinescens, D. Dietr. Syn. Pl. iv. 1662 = rhamnoides.
subumbellata, DC. Prod. vi. 474 = quinquedentata.
sulcata, Thunb. Prod. Pl. Cap. 167 = Euryops sulcatus.
Tagetes, Linn. Syst. ed. X. 1234 = Gamolepis annua.
taraxacoides, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
tenuissima, Haw. Rev. Pl. Succ. 315 = cylindrica.
tenuissima, Linn. Mant. i. 98 = Euryops tenuissimus.
tephrosioides, *Sond. in Harv. & Sond. Fl. Cap.* iii. 344.—Afr. austr.
tomentosa, Salisb. Prod. 211 = Euryops pectinatus.
tortuosa, *Sch. Bip. in Flora*, xxvii. (1844) 769.—Afr. austr.
trifida, Linn. f. Suppl. 387 ; Thunb. Prod. Pl. Cap. 168 = Euryops trifidus.
trifida, Thunb. ex Harv. & Sond. Fl. Cap. iii. 414 = Euryops oligoglossus.
trifurcata, Less. in Linnaea, vi. (1831) 94 = Gamolepis trifurcata.
trifurcata, Linn. f. Suppl. 387 ; Thunb. Prod. Pl. Cap. 168 = Euryops trifurcatus.
trinervia, *DC. Prod.* vi. 477.—Afr. austr.
triplinervis, *DC. l. c.* 478.—Afr. austr.
tuberosa, DC. l. c. 479 (excl. syn.) = heterophylla.
tuberosa, *Thunb. Prod. Pl. Cap.* 168.—Afr. austr.
umbelliformis, *DC. Prod.* vi. 474.—Afr. austr.
undulata, *E. Mey. ex Steud. Nom.* ed. II. ii. 238.—Afr. austr.
uniflora, *Savign. in Lam. Encyc.* iv. 670.—Afr. austr.
vestita, *DC. Prod.* vi. 482.—Afr. austr.
viminea, *E. Mey. ex DC. l. c.* 477.—Afr. austr.
virginea, Linn. f. Suppl. 389 ; Thunb. Prod. Pl. Cap. 168 = Euryops virgineus.
Zeyheri, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 339.—Afr. austr.

OTHONNOPSIS, Jaub. & Spach. Illustr. iv. 90. t. 357 (1852). *COMPOSITAE*, Benth. & Hook. f. ii. 451.
HERTIA, Neck. Elem. i. 8 (1790); Less. in Linnaea, vi. (1831) 94.
alata, *Benth. & Hook. f. Gen.* ii. 451.—Afr. austr.
angustifolia, *Jaub. & Spach, Illustr.* iv. 90. t. 357.—Persia.
cheirifolia, *Benth. & Hook. f. Gen.* ii. 451.—Algeria.
ciliata, *Benth. & Hook. f. l. c.*—Afr. austr.
cluytiaefolia, *Benth. & Hook. f. l. c.*—Afr. austr.
cneorifolia, *Benth. & Hook. l. c.*—Afr. austr.
intermedia, *Boiss. Diagn.* Ser. II. vi. 106.—Persia; Beluchist.
Kraussii, *Benth. & Hook. f. Gen.* ii. 451.—Afr. austr.
pallens, *Benth. & Hook. f. l. c.*—Afr. austr.

OTHOSTEMMA, Pritz. Ic. Ind. i. 782 (1855) = OTOSTEMMA, Blume = **Hoya**, R. Br. (Asclep.).

OTHRYS, Noronha, ex Thou. Nov. Gen. Madag. 13 (1806) = **Crataeva**, Linn. (Cappar.).

OTIDIA, Lindl. ex Sweet, Geran. t. 98 (1820) = **Pelargonium**, L'Hérit. (Geran.).
alternans, Sweet, l. c. t. 286 = P. alternans.
Burmanniana, Eckl. & Zeyh. Enum. 69 = P. ferulaceum.
carnosa, Lindl. ex Sweet, Geran. t. 98 = P. carnosum.
ceratophylla, Hoffmgg. Verz. Pfl. 84 = P. ceratophyllum.
corallina, Eckl. & Zeyh. Enum. 69 = P. corallinum.
crithmifolia, Sweet, Hort. Brit. ed. I. 75; Geran. t. 354 = P. crithmifolium.
dasycaulon, Eckl. & Zeyh. Enum. 68 = P. dasycaule.
ferulacea, Eckl. & Zeyh. l. c. 69 = P. ferulaceum.
laxa, Sweet, Geran. t. 196 = P. carnosum.
microphylla, Eckl. & Zeyh. Enum. 69 = P. microphyllum.

OTILLIS, Gaertn. Fruct. i. 275. t. 57 (1788) = **Leea**, Royen (Ampel.).
zeylanica, Gaertn. l. c. = L. sambucina.

OTIOPHORA, Zucc. in Abh. Akad. Muench. i. (1832) 315. *RUBIACEAE*, Benth. & Hook. f. ii. 137.
cupheoides, *N. E. Br. in Hook. Ic. Pl.* t. 1453.—Afr. austr.
pauciflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 170.—Madag.
scabra, *Zucc. in Abh. Akad. Muench.* i. (1832) 316.—Madag.

OTITES, Adans. Fam. ii. 255 (1763) = **Silene**, Linn. (Caryoph.).
nemoralis, Opiz, Seznam, 71 = S. italica.
nutans, Opiz, l. c. = S. nutans.
parviflorus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 346 = S. Otites.
viridiflora, Opiz, in Lotos, iv. (1854) 42 = S. viridiflora.

OTOCARPUS, Dur. in Duch. Rev. Bot. ii. (1847) 435 = **Rapistrum**, Desv. (Crucif.).
virgatus, Dur. l. c. 436.—Alger.

OTOCHILUS, Lindl. Gen. et Sp. Orch. 35 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 519.
BROUGHTONIA, Wall. ex Lindl. Gen. et Sp. Orch. 35 (1830).
TETRAPELTIS, Wall. ex Lindl. Bot. Reg. sub t. 1522 (1832).
alba, *Lindl. in Wall. Cat.* n. 1967.—Reg. Himal.
fusca, *Lindl. l. c.*—Reg. Himal.
lancifolia, Griff. Notul. iii. 278 ; Ic. Pl. Asiat. t. 289 = porrecta.
latifolia, Griff. ll. cc. 279 ; t. 288 = fusca.
porrecta, *Lindl. in Wall. Cat.* n. 1968.—Reg. Himal.; Burma.

OTOCHLAMYS, DC. Prod. vi. 77 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 428.
Eckloniana, *DC. Prod.* vi. 77.—Afr. austr.
majuscula, Harv. ex Harv. & Sond. Fl. Cap. iii. 177 = Eckloniana.

OTOGLYPHIS, Pomel, Nouv. Mat. Fl. Atl. 56 (1874) = **Cotula**, Tourn. (Compos.).
pubescens, Pomel, l. c. 57 = C. pubescens.

OTOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 572 = **Capura**, Blanco (Sapind.).
nigrescens, Turcz. l. c. 573 = C. pinnata.

OTOMERIA, Benth. in Hook. Niger Fl. 405 (1849) *RUBIACEAE*, Benth. & Hook. f. ii. 55.
dilatata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 50.—Afr. trop.
guineensis, *Benth. in Hook. Niger Fl.* 405.—Afr. trop.
lanceolata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 50.—Afr. trop.
madiensis, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 83. t. 47.—Afr. trop.
oculata, *S. Moore, in Journ. Bot.* xviii. (1880) 4.—Afr. trop.

OTONEPHELIUM, Radlk. Sapind. Holland-Ind. Nachtr. 71 (1879) = **Nephelium**, Linn. (Sapind.).
stipulaceum, Radlk. l. c. = N. stipulaceum.

OTONYCHIUM, Blume, Rumphia, iii. 180 (1850) = **Harpullia**, Roxb. (Sapind.).
imbricatum, Blume, l. c. = H. cupanioides.
retusum, Miq. Fl. Ind. Bat. i. II. 572.—Java.

OTOPAPPUS, Benth. in Benth. & Hook. f. Gen. ii. 380 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
curviflorus, *Hemsl. Biol. Centr. Am. Bot.* ii. 191.—Mexic.
epaleaceus, *Hemsl. l. c.*—Mexic.
robustus, *Hemsl. l. c.*—Mexic.
verbesinoides, *Benth. in Hook. Ic. Pl.* t. 1153.—Nicaragua.

OTOPETALUM, Miq. Fl. Ind. Bat. ii. 400 (1856); et in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 191. *APOCYNACEAE*, Benth. & Hook. f. ii. 692.
micranthum, *Miq. ll. cc.*—Java.

OTOPHORA, Blume, Rumphia, iii. 142 (1850). *SAPINDACEAE*, [Benth. & Hook. f. i. 405.]
alata, *Blume, l. c.* 145.—Borneo.
amoena, Blume, l. c. 142 = Melicocca amoena.
Blancoi, Blume, l. c. = Sapindus baccatus.
confinis, *Blume, l. c.* 143.—Borneo.
cordigera, *Radlk. Sapind. Holland-Ind. Nachtr.* 85.—Borneo.
erythrocalyx, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 680.—Malacca.

OTOPHORA :—
fruticosa, Blume, Rumphia, iii. 142 = Sapindus fruticosus.
imbricata, *Blume, l. c.* 144.—Borneo.
javanica, *Miq. Fl. Ind. Bat.* i. II. 561.—Java.
nigrescens, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 52 = Capura pinnata.
paradoxa, *Blume, Rumphia*, iii. 146.—Ins. Philipp.
paucijuga, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 680.—Malacca.
pubescens, *Blume, Rumphia*, iii. 145.—Borneo.
ramiflora, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879), 102.—Borneo.
spectabilis, *Blume, Rumphia*, iii. 142.—Java.
Zollingeriana, *Teijsm. & Binn. in Tijdschr. Nederl. Indie*, xxv. (1863) 423.—Java.

OTOPHYLLA, Benth. in DC. Prod. x. 512 (1846) = **Gerardia**, Linn. (Scrophul.).
Drummondii, Benth. l. c. = G. densiflora.
Michauxii, Benth. l. c. = G. auriculata.

OTOPTERA, DC. Mém. Lég. 249. t. 42 (1825) = **Vigna**, Savi (Legumin.).
Burchelli, DC. l. c. = V. Burchellii.

OTOSEMA, Benth. in Miq. Pl. Jungh. 248 (1851) 55 = **Milletia**, Wight & Arn. (Legumin.).
caudata, Benth. in Miq. Pl. Jungh. 249 = M. caudata.
extensa, Benth. l. c. = M. extensa.
fruticosa, Benth. l. c. = M. fruticosa.
macrophylla, Benth. l. c. = M. auriculata.

OTOSMA, Rafin. New Fl. Am. ii. 90 (1836); et Fl. Tellur. iv. 8 (1836) = **Richardia**, Kunth (Aroid.).
ethiopica, Rafin. ll. cc. = R. africana.

OTOSPERMUM, Willk. in Bot. Zeit. xxii. (1864) 251 = **Matricaria**, Tourn. (Compos.).
glabrum, Willk. l. c. = M. glabra.

OTOSTEGIA, Benth. Lab. Gen. et Sp. 601 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1213.
DICTILIS, Rafin. Fl. Tellur. iii. 89 (1836).
arabica, *Jaub. & Spach, Illustr.* iv. 121. t. 380.—Oriens.
Aucheri, *Boiss. Diagn.* Ser. I. v. 40.—Persia; Beluchist.
Benthamiana, *Jaub. & Spach, Illustr.* iv. 119. t. 379.—Oriens.
integrifolia, *Benth. Lab. Gen. et Sp.* 602.—Abyssin.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 59.—Persia.
limbata, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 680.—Ind. bor. occ.
microphylla, *Boiss. Diagn.* Ser. I. v. 39.—Persia.
moluccoides, Jaub. & Spach, Illustr. iv. 117. t. 378 = scariosa.
persica, Boiss. Fl. Orient. iv. 777 = microphylla.
repanda, *Benth. Lab. Gen. et Sp.* 602.—Abyssin.
scariosa, *Benth. l. c.*—Arab.; Abyssin.
Schimperi, *Boiss. Fl. Orient.* iv. 776.—Arab.
Steudneri, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 682.—Afr. bor.
tomentosa, *A. Rich. Tent. Fl. Abyss.* ii. 198.—Abyssin.

OTOSTEMMA, Blume, Rumphia, iv. 30 (1848) = **Hoya**, R. Br. (Asclep.).
lacunosum, Blume, l. c. = H. lacunosa.

OTOTROPIS, Nees, Del. Sem. Hort. Vratisl. (1838); ex Linnaea, xiii. (1839) Litt. 120 = **Desmodium**, Desv. (Legumin.).
microphyllus, Benth. in Ann. Wien. Mus. ii. (1838) 142 = Indigofera gracilis, *Spreng.*
recurvatus, Nees, Del. Sem. Hort. Vratisl. (1838); ex Linnaea, xiii. (1839) Litt. 121, in obs. = D. laxiflorum.
rufescens, Nees, l. c. 121 = D. rufescens.
sambuensis, Nees, l. c. 120 = D. floribundum.

OTTELIA, Pers. Syn. i. 400 (1805). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 453.
DAMASONIUM, Schreb. Gen. 242 (1789).
HYMENOTHECA, Salisb. in Trans. Hort. Soc. i. (1812) 268.
alismoides, *Pers. Syn.* i. 400.—As. et Austral. trop.
brasiliensis, *Walp. Ann.* iii. 510.—Bras.
cygnorum, Walp. l. c. = ovalifolia.

OTTELIA:—
ensiformis, *Blanco, Fl. Filip.* ed. I. 460.—Ins. Philipp.
Hoffmeisteri, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 41.—Reg. Himal.
indica, Planch. ex Dalz. & Gibs. Bomb. Fl. 278 = ovalifolia.
japonica, *Miq. in Verh. Akad. Wetensch.* II. ii. (1868) 89.—Japon.
javanica, *Miq. Fl. Ind. Bat.* iii. 240.—Java.
lactucaefolia, Walp. Ann. iii. 510 = alismoides.
lanceolata, Wight, ex Wall. Cat. n. 7244 = alismoides.
lancifolia, *A. Rich. Tent. Fl. Abyss.* ii. 280.—Abyssin.
ovalifolia, *Rich. in Mém. Inst. Par.* (1811) 27, 78. t. 7.—Austral.
scabra, *Baker, in Trans. Linn. Soc.* xxix. (1875) 151.—Afr. trop.
tenera, *Benth. Fl. Austral.* vi. 257.—Austral.
ulvaefolia, *Wdlp. Ann.* iii. 510.—Madag.

OTTILIS, Endl. Gen. 797 (1839), sphalm. = OTILLIS, Gaertn. = **Leea**, Royen (Ampel.).

OTTOA, H. B. & K. Nov. Gen. et Sp. v. 20. t. 423 (1821). *UMBELLIFERAE*, Benth. & Hook. f. i. 899.
oenanthoides, *H. B. & K. l. c.*—Mexic.; N. Granat.

OTTONIA, Spreng. Neue Entdeck. i. 255 (1820) = **Piper**, Linn.
Anisum, Mart. ex Miq. in Mart. Fl. Bras. iv. i. 66 = P. corcovadense.
Anisum, Spreng. Neue Entedck. i. 255 = P. Jaborandi.
Blanchetii, Moric. Pl. Nouv. Am. 86 = P. corcovadense.
carpinifolia, Presl, Epim. Bot. 229 = P. carpinifolium.
Carpunga, Miq. Syst. Pip. 547 = P. Carpunga.
consanguinea, Presl, Epim. Bot. 230 = P. punctatissimum.
corcovadensis, Miq. in Linnaea, xx. (1847) 175 = P. corcovadense.
diversifolia, Kunth, l. c. xiii. (1839) 578 = P. Selloi.
eucalyptifolia, Kunth, l. c. 582 = P. eucalyptiphyllum.
glaucescens, Miq. in Seem. Bot. Voy. Herald, 200 = P. darienense.
Hookeriana, Miq. in Hook. Journ. Bot. iv. (1842) 40 = P. Gardneri.
Jaborandi, Kunth, in Linnaea, xiii. (1839) 579 = P. Jaborandi.
Klotzschiana, Kunth, l. c. 589 = P. Klotzschianum.
laeta, Kunth, l. c. 587 = P. laetum.
leptostachya, Kunth, l. c. 586 = P. Boucheanum.
Lessertiana, Miq. Syst. Pip. 546 = P. Lessertianum.
macrophylla, Kunth, in Linnaea, xiii. (1839) 583 = P. Heyneanum.
macrostachya, Mart. ex Miq. in Linnaea, xx. (1847) 176 = P. sebastianopolitense.
Martiana, Miq. l. c. 178 = P. Miquelianum.
padifolia, Kunth, l. c. xiii. (1839) 580 = P. padifolium.
Pohliana, Miq. l. c. xx. (1847) 179 = P. Pohlianum.
propinqua, Kunth, l. c. xiii. (1839) 583 = P. corcovadense.
propinqua, Miq. in Mart. Fl. Bras. iv. i. 66, 222 = P. Baptisianum.
pteropoda, Moric. Pl. Nouv. Am. 88 = P. pteropodon.
punctata, Griseb. Cat. Pl. Cub. 65 = P. cubense.
sphaerocarpa, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 176 = P. sphaerostachyum.
sphaerostachya, C. DC. in DC. Prod. xvi. i. 326 = P. sphaerostachyum.
stipulacea, Presl, Epim. Bot. 231 = P. stipulaceum.
unguiculata, Heynh. Nom. ii. 464 = P. unguiculatum.
Vahlii, Kunth, in Linnaea, xiii. (1839) 585 = P. ovatum.
Warakabacoura, Miq. l. c. xx. (1847) 180 = P. Warakaboura.

OUDEMANSIA, Miq. Pl. Jungh. 295 (1854) = **Helicteres**, Pluk. (Stercul.).
hirsuta, Miq. l. c. 297 = H. spicata?
integerrima, Miq. l. c. 296 = H. angustifolia.
javanensis, C. Muell. in Walp. Ann. iv. 320 = seq.
javensis, Hassk. Retzia, i. 134 = H. angustifolia.
Korthalsii, Miq. Fl. Ind. Bat. i. ii. 170 = H. integerrima.
lanata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 252.—Siam.
microcarpa, Miq. Fl. Ind. Bat. i. ii. 171 = H. microcarpa.
obversa, Miq. Pl. Jungh. 297 = H. sundaica.
spicata, Teijsm. & Binn. Cat. Hort. Bog. 196 = H. spicata.
virgata, Hassk. Retzia, i. 134 = H. angustifolia.
viscida, Miq. Pl. Jungh. 297 = H. viscida.

OUDNEYA, R. Br. in Denh. & Clapp. Trav. App. 220 (1826) = **Moricandia**, DC. (Crucif.).
africana, R. Br. l. c. = M. suffruticosa.

OUGEINIA, Benth. in Miq. Pl. Jungh. 216 (1851–55). *LEGUMINOSAE*, Benth. & Hook. f. i. 518.
dalbergioides, *Benth. in Miq. Pl. Jungh.* 216—Ind. or.

OURAGOGA, Linn. Hort. Cliff. 486 (1737) = **Uragoga**, Linn. (Rubiac.).

OURATEA, Aubl. Pl. Gui. i. 397. t. 152 (1775) = **Gomphia**, Schreb. (1789) (Ochnac.).
acuminata, Engl. in Mart. Fl. Bras. xii. ii. 318 = G. acuminata.
alternifolia, Engl. l. c. 339 = G. alternifolia.
aquatica, Engl. l. c. 343 = G. aquatica.
Blanchetiana, Engl. l. c. 320 = G. Blanchetiana.
Candollei, Engl. l. c. 314, in adnot. = G. guyannensis.
caracasana, Engl. l. c. 315 = G. caracasana.
cardiosperma, Engl. l. c. 348 = G. cardiosperma.
cassinefolia, Engl. l. c. 319 = G. cassinefolia.
castaneaefolia, Engl. l. c. 309 = G. castaneaefolia.
caudata, Engl. l. c. 313.—Am. trop.
cinnamomea, Wawra, in Oestr. Bot. Zeitschr. xxix. (1879) 215.—Bras.
coccinea, Engl. in Mart. Fl. Bras. xii. ii. 310 = G. coccinea.
conduplicata, Engl. l. c. 343 = G. conduplicata.
confertiflora, Engl. l. c. 328 = G. confertiflora.
crassifolia, Engl. l. c. 311 = G. crassifolia.
crassinervia, Engl. l. c. 337.—Am. trop.
cuspidata, Engl. l. c. 345 = G. cuspidata.
decorans, Baill. Hist. Pl. iv. 357 = G. olivaeformis.
Duparquetiana, Baill. l. c. 359 = G. Duparquetiana.
ferruginea, Engl. in Mart. Fl. Bras. xii. ii. 335.—Am. trop.
Fieldingiana, Engl. l. c. 309 = G. Fieldingiana.
floribunda, Engl. l. c. 331 = G. floribunda.
gigantophylla, Engl. l. c. 338 = G. gigantophylla.
glaucescens, Engl. l. c. 328 = G. glaucescens.
globosa, Engl. l. c. 323.—Am. trop.
grandiflora, Engl. l. c. 336 = G. grandiflora.
grandifolia, Engl. l. c. 346 = G. grandifolia.
guatemalensis, Engl. l. c. 345.—Am. trop.
guianensis, Aubl. Pl. Gui. i. 397 = G. guyannensis.
hexasperma, Baill. Hist. Pl. iv. 366 = G. hexasperma.
humilis, Engl. in Mart. Fl. Bras. xii. ii. 330 = G. humilis.
ilicifolia, Baill. Hist. Pl. iv. 366 = G. ilicifolia.
inundata, Engl. in Mart. Fl. Bras. xii. ii. 335 = G. inundata.
Jabotapita, Engl. l. c. 350 = G. Jabotapita.
Juergensii, Engl. l. c. 351 = G. Jurgensenii.
lanceolata, Engl. l. c. 327 = G. lanceolata.
laurifolia, Engl. l. c. 350 = G. laurifolia.
longifolia, Engl. l. c. 316 = G. longifolia.
lucens, Engl. l. c. 350 = G. lucens.
lucidula, Engl. l. c. 349 = G. lucidula.
Magdalenae, Engl. l. c. 315 = G. Magdalenae.
membranacea, Engl. l. c. 342 = G. membranacea.
mexicana, Engl. l. c. 312 = G. mexicana.
microdonta, Engl. l. c. = G. microdonta.
Miersii, Engl. l. c. 341 = G. Miersii.
multiflora, Engl. l. c. = G. multiflora.
nana, Engl. l. c. 326 = G. nana.
nervosa, Engl. l. c. 332 = G. nervosa.
nitida, Engl. l. c. 310, 314, in adnot. = G. guyannensis.
odora, Poepp. ex Engl. l. c. 334 = G. odora.
oleaefolia, Engl. l. c. 325 = G. oleaefolia.
olivaeformis, Engl. l. c. 344 = G. olivaeformis.
opaca, Engl. l. c. 349.—Am. trop.
ovalis, Engl. l. c. 326 = G. ovalis.
parviflora, Baill. Hist. Pl. iv. 366 = G. parviflora.
parvifolia, Engl. in Mart. Fl. Bras. xii. ii. 332 = G. parvifolia.
patens, Engl. l. c. 317.—Am. trop.
pendula, Engl. l. c. 339 = G. pendula.
pisiformis, Engl. l. c. 343.—Am. trop.
polita, Engl. l. c. 349 = G. polita.
polyantha, Engl. l. c. 317 = G. polyantha.
polygyna, Engl. l. c. 342.—Am. trop.
pubescens, Engl. l. c. 327 = G. pubescens.
pulchella, Engl. l. c. 340 = G. pulchella.
pycnostachys, Engl. l. c. 315 = G. pycnostachys.
pyrifolia, Engl. l. c. 320 = G. pyrifolia.
revoluta, Engl. l. c. 346 = G. revoluta.
Riedeliana, Engl. l. c. 322.—Am. trop.

OURATEA:—
rigida, Engl. in Mart. Fl. Bras. xii. ii. 307.—Am. trop.
Roraimae, Engl. l. c. 308.—Am. trop.
rotundifolia, Engl. l. c. 307 = Gomphia rotundifolia.
rufidula, Engl. l. c. 320 = G. Blanchetiana.
rupununiensis, Engl. l. c. 318 = G. rupununiensis.
salicifolia, Engl. l. c. 324 = G. salicifolia.
Schomburgkii, Engl. l. c. 333 = G. Schomburgkii.
Sellowii, Engl. l. c. 347 = G. Sellowii.
semiserrata, Engl. l. c. 323 = G. semiserrata.
spectabilis, Engl. l. c. 330 = G. spectabilis.
Spruceana, Engl. l. c. 337.—Am. trop.
squamosa, Engl. l. c. 318 = G. squamosa.
stipulacea, Engl. l. c. 347 = G. stipulacea.
suaveolens, Engl. l. c. 349 = G. suaveolens.
subscandens, Engl. l. c. 322 = G. subscandens.
superba, Engl. l. c. 334.—Am. trop.
tenuifolia, Engl. l. c. 324.—Am. trop.
Theophrasta, Baill. Hist. Pl. iv. 359 = G. gigantophylla.
thyrsoidea, Engl. in Mart. Fl. Bras. xii. ii. 313.—Am. trop.
vaccinioides, Engl. l. c. 329 = G. vaccinioides.
Vasivae, Engl. l. c. 336 = G. Vasivae.
verruculosa, Engl. l. c. 335.—Am. trop.
verticillata, Engl. l. c. 349 = G. verticillata.

OURET, Adans. Fam. ii. 268 (1763) = **Aerua**, Forsk. (Amarant.).

OURISIA, Comm. ex Juss. Gen. 100 (1789). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 962.
DICHROMA, Cav. in Anal. Cienc. Nat. iii. (1801) 232; et Ic. vi. 59. t. 582 (1801).
alpina, *Poepp. & Endl. Nov. Gen. et Sp.* i. 3. t. 6.—Chili.
antartica, Hook. f. Fl. Antarct. 335. t. 18 = breviflora.
biflora, *Wedd. Chlor. And.* ii. 117.—Bolivia.
breviflora, *Benth. in DC. Prod.* x. 493.—Reg. Magell.
caespitosa, *Hook. f. Fl. N. Zel.* i. 197.—N. Zel.
californica, Benth. Pl. Hartw. 327 = Hesperochiron californicus.
chamaedrifolia, *Benth. in DC. Prod.* x. 493.—N. Granat.; Peruv.
coccinea, *Pers. Syn.* ii. 169.—Ins. Chiloe.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 218.—N. Zel.
Diazi, *Phil. in Linnaea*, xxxiii. (1864–65) 211.—Chili.
elegans, *Phil. l. c.* xxx. (1859–60) 197.—Chili.
fragrans, *Phil. l. c.* xxxiii. (1864–65) 211.—Chili.
glabra, Molina, Sagg. Chile, ed. II. 291 = magellanica.
glandulosa, *Hook. f. Handb. N. Zeal. Fl.* 219.—N. Zel.
integrifolia, *R. Br. Prod.* 439.—Austral.
macrocarpa, *Hook. f. Fl. N. Zel.* i. 197.—N. Zel.
macrophylla, *Hook. Ic. Pl.* tt. 545, 546.—N. Zel.
magellanica, *Gaertn. f. Fruct.* iii. 44. t. 185.—Reg. Magellan.
magellanica, Poepp. & Endl. Nov. Gen. et Sp. i. 2. t. 4 = Poeppigii.
melospernoides, *Phil. in Anal. Univ. Chil.* (1873) 527.—Chili.
microphylla, *Poepp. & Endl. Nov. Gen. et Sp.* i. 3. t. 7.—Chili.
muscosa, *Benth. in DC. Prod.* x. 493.—N. Granat.
nana, *Benth. l. c.*—Ecuador.
pallens, *Poepp. & Endl. Nov. Gen. et Sp.* i. 3. t. 5.—Chili.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864–65) 210.—Chili.
pinnata, Wall. ex Benth. Scroph. Ind. 47 = Sibthorpia pinnata.
Poeppigii, *Benth. in DC. Prod.* x. 492.—Chili.
polyantha, *Poepp. & Endl. Nov. Gen. et Sp.* i. 4.—Chili.
pulchella, *Wedd. Chlor. And.* ii. 116.—Bolivia.
pygmaea, *Phil. in Linnaea*, xxix. (1857–58) 27.—Chili.
racemosa, *Clos, in C. Gay, Fl. Chil.* v. 130.—Chili.
rancoana, *Phil. in Linnaea*, xxxiii. (1864–65) 210.—Chili.
robusta, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 246.—N. Zel.
rosea, *Phil. in Linnaea*, xxix. (1857–58) 26.—Chili.
rupicola, *Wedd. Chlor. And.* ii. 115.—Bolivia.

OURISIA:—
serpyllifolia, *Benth. in DC. Prod.* x. 493.—Chili.
sessilifolia, *Hook. f. Handb. N. Zeal. Fl.* 218.—N. Zel.
uniflora, *Phil. in Linnaea*, xxix. (1857-58) 27.—Chili.
villosa, Molina, Sagg. Chile, ed. II. 291 = coccinea.

OUROUPARIA, Aubl. Pl. Gui. i. 177. t. 68 (1775) = **Uncaria**, Schreb. (1789) (Rubiac.).
acida, Baill. Hist. Pl. vii. 375 = U. acida.
africana, Baill. in Bull. Soc. Linn. Par. i. (1879) 228 = U. africana.
Gambir, Baill. Hist. Pl. vii. 350, 375 = U. Gambier.
guianensis, Aubl. Pl. Gui. i. 177 = U. tomentosa.
madagascariensis, Baill. in Bull. Soc. Linn. Par. i. (1879) 228 = U. africana?
Perrottetii, Baill. l. c. 227 (= *Sabicea Perrottetii*).—Afr. trop.
polycephala, Baill. l. c. 229 = Nauclea Cinchonae.

OUSTROPIS, G. Don, Gen. Syst. ii. 214 (1832) = **Indigofera**, Linn. (Legumin.).
microphylla, G. Don, l. c. 215 = I. gracilis, *Spreng.*

OUTARDA, Dum. Anal. Fam. 32 (1829), sphalm. = **Coutarea**, Aubl. (Rubiac.).

OUTEA, Aubl. Pl. Gui. 28. t. 9 (1775) = **Macrolobium**, Schreb. (1789) (Legumin.).
acaciaefolia, Benth. in Hook. Journ. Bot. ii. (1840) 94 = M. acaciaefolium.
bijuga, DC. Prod. ii. 511 = Afzelia bijuga.
gracilis, Spruce, ex Benth. in Mart. Fl. Bras. xv. II. 223 = M. gracile.
guianensis, Aubl. Pl. Gui. i. 28. t. 9 = M. Utea.
humilis, Spruce, ex Benth. in Mart. Fl. Bras. xv. II. 223 = M. gracile.
multijuga, DC. Prod. ii. 510 = M. multijugum.
racemigera, Tul. in Arch. Mus. Par. iv. (1844) 174 = M. pendulum.

OUTREYA, Jaub. & Spach, Illustr. i. 131. t. 68 (1842) = **Jurinea**, Cass. (Compos.).
carduiformis, Jaub. & Spach, l. c. = J. carduiformis.

OUVIRANDRA, Thou. Gen. Nov. Madag. 2 (1806) = **Aponogeton**, Thunb. (Naiad.).
Bernieriana, Decne. in Deless. Ic. Sel. i. 62. t. 100 = A. Bernerianum.
fenestralis, Poir. Encyc. Suppl. iv. 237 = A. fenestrale.
Heudelotii, Kunth, Enum. Pl. iii. 593 (sp. dub.).—Afr. austr.
Hildebrandtii, Hort. Berol. ex Kurtz, in Verh. Bot. Ver. Brandenb. xix. (1877) p. viii; et ex Eichl. in Monatsschr. Befoerd. Gartenb. xxii. (1879) 6. t. 1 = A. Bernerianum.
undulata, Edgew. in Hook. Lond. Journ. Bot. iii. (1844) 405 = A. crispum.

OVERSTRATIA, Deschamps, ex R. Br. in Benn. Pl. Jav. Rar. 171 (1840) = **Saurauja**, Willd. (Ternstr.).

OVIDIA, Meissn. in DC. Prod. xiv. 524 (1857). *THYMELAEACEAE*, Benth. & Hook. f. iii. 190.
andina, *Meissn. l. c.*—Chili.
Humboldti, *Meissn. l. c.* 525.—N. Granat.
parviflora, *Meissn. l. c.*—Chili.
Pillo-Pillo, Hohen. ex Meissn. l. c. = parviflora.
Pillo-Pillo, *Meissn. l. c.* 524.—Chili.

OVIDIA, Rafin. Fl. Tellur. iii. 68 (1836) = **Commelina**, Plum.
gracilis, Rafin. l. c. = C. fasciculata?

OVIEDA, Linn. Gen. ed. I. 59 (1737) = **Clerodendron**, Linn. (Verben.).
inermis, Burm. f. Fl. Ind. 136. t. 43. f. 1 = C. Siphonanthus.
mitis, Linn. Sp. Pl. ed. II. 888 = C. Siphonanthus.
ovalifolia, A. Juss. in Ann. Mus. Par. vii. (1806) 76 = C. floribundum.
Pillo-Pillo, Meissn. in Lechl. Berb. Am. Austr. 53, sphalm. = Ovidia Pillo-Pillo.
spinosa, Linn, Sp. Pl. 637 = C. spinosum.
verticillatum, Roxb. ex D. Don, Prod. Fl. Nep. 102 = C. Siphonanthus.

OVIEDA, Spreng. Anleit. ii. 1. 258 (1817) = **Lapeyrousia**, Pourr. (Irid.).
aculeata, Klatt, in Linnaea, xxxii. (1863) 777 = L. compressa.
anceps, Spreng. Syst. i. 147 = L. compressa.
bracteata, Spreng. l. c. = L. fissifolia.
corymbosa, Spreng. l. c. = L. corymbosa.
erythrantha, Klotzsch, ex Klatt, in Peters, Reise Mossamb. Bot. 516 = L. erythrantha.
Fabricii, Spreng. Syst. i. 147 = L. Fabricii.
falcata, Spreng. l. c. = L. falcata.
fasciculata, Spreng. l. c. = L. fasciculata.
fissifolia, Spreng. l. c. = L. fissifolia.
fistulosa, Spreng. ex Klatt, in Linnaea, xxxii. (1863) 781 = L. fistulosa.
micrantha, E. Mey. ex Klatt, l. c. = L. micrantha.
purpureo-lutea, Klatt, l. c. 780 = L. purpureo-lutea.
silenoides, Spreng. Syst. i. 147 = L. silenoides.

OVILLA, Adans. Fam. ii. 134 (1763) = **Jasione**, Linn. (Campan.).

OVOSTIMA, Rafin. New Fl. Am. ii. 70 (1836) = **Gerardia**? Linn. (Scrophul.).
petiolata, Rafin. l. c. (Quid?).—Am. bor.

OWENIA, Hilsenb. ex Meissn. in DC. Prod. xiv. 89 (1856) = **Oxygonum**, Burch. (Polygon.).

OWENIA, F. Muell. in Hook. Kew Journ. ix. (1857) 303. *MELIACEAE*, Benth. & Hook. f. i. 337.
acidula, *F. Muell. l. c.* 304.—Austral.
cepiodora, *F. Muell. Fragm.* xi. 82.—Austral.
cerasifera, *F. Muell. in Hook. Kew Journ.* ix. (1857) 305.—Austral.
reticulata, *F. Muell. l. c.*—Austral.
venosa, *F. Muell. l. c.* 304.—Austral.
vernicosa, *F. Muell. Fragm.* iii. 15, in obs.—Austral.
xerocarpa, F. Muell. l. c. 13 = reticulata.

OXALIS, Linn. Gen. ed. I. 134 (1737). *GERANIACEAE*, Benth. & Hook. f. i. 276.
OXYS, Tourn. ex Adans. Fam. ii. 388 (1763).
SASSIA, Molina, Sagg. Chile, 145 (1782).
abyssinica, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 432 = O. Pes-caprae.
abyssinica, *Walp. Ann.* ii. 241.—Afr. trop.
Acetosella, Blanco, Fl. Filip. ed. I. 388 = corniculata.
Acetosella, Hook. Fl. Bor. Am. i. 118, partim = oregana.
Acetosella, *Linn. Sp. Pl.* 433,—Reg. bor. temp.
acuminata, Schlecht. in Linnaea, v. (1830) 224 = Lindenii.
acuminata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 429.—Mexic.
acutifolia, *Prog. in Mart. Fl. Bras.* xii. II. 514.—Bras.
adenodes, *Sond. in Harv. & Sond. Fl. Cap.* i. 334.—Afr. austr.
adenophylla, *Gill. in Hook. Bot. Misc.* iii. (1833) 165.—Chili.
adoënsis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 273.—Abyssin.
adscendens, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823-24) 164 = medicaginea.
adspersa, *Eckl. & Zeyh. Enum.* 94.—Afr. austr.
aemula, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 496 = densifolia.
aeschynomenifolia, *O. Hoffm. in Bremen, Abh.* vii. (1882) 244.—Madag.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 328.—Afr. austr.
aganophila, *Sond. l. c.* 339.—Afr. austr.
aganophylla, Eckl. & Zeyh. Enum. 86 = ebracteata.
alata, *Zucc. in Denkschr. Akad. Muench.* ix. (1823-24) 155.—Bras.
alba, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 398.—Am. austr.
alba, Steud. Nom. ed. I. 578; ed. II. ii. 238 = O. Acetosella.
albicans, *H. B. & K. Nov. Gen. et Sp.* v. 244.—Mexic.
albida, Sond. in Harv. & Sond. Fl. Cap. i. 325 = leucotricha.
albizzioides, *O. Hoffm. in Bremen, Abh.* vii. (1882) 242.—Madag.

OXALIS:—
algoënsis, *Eckl. & Zeyh. Enum.* 90.—Afr. austr.
aliena, Spreng. Neue Entdeck. iii. 58 = mandioccana.
alsinoides, *Walp. in Nov. Act. Nat. Cur.* xix. Suppl. i. (1843) 319.—Chili.
alta, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 504 = elatior.
amara, *A. St. Hil. Fl. Bras. Mer.* i. 119.—Bras.
amazonica, *Prog. in Mart. Fl. Bras.* xii. II. 503.—Bras.
ambigua, Jacq. Oxal. 80. t. 43 = mutabilis.
ambigua, A. Rich. Bot. Voy. Astrol. i. 296 = corniculata.
ambigua, Salisb. in Trans. Linn. Soc. ii. (1794) 242 = corniculata.
americana, Bigel. ex DC. Prod. i. 700 = O. Acetosella.
amoena, *Jacq. Hort. Schoenb.* ii. 42.—Afr. austr.
amoena, Salisb. Prod. 321 = purpurea.
andicola, *Gill. in Hook. Bot. Misc.* iii. (1833) 161.—Chili.
anemonoides, Eckl. & Zeyh. Enum. 85 = heterophylla.
angusta, *Sond. in Harv. & Sond. Fl. Cap.* i. 329.—Afr. austr.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 249.—Mexic.
anthelmintica, A. Rich. Tent. Fl. Abyss. i. 124 = O. Pes-caprae.
antucensis, *Phil. in Anal. Univ. Chil.* (1872) 685.—Chili.
Apodoscias, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 430.—Java.
approximata, *Sond. in Harv. & Sond. Fl. Cap.* i. 326.—Afr. austr.
aptera, *Zucc. ex Prog. in Mart. Fl. Bras.* xii. II. 514.—Bras.
Aracatcha, Hort. ex Zucc. in Abh. Akad. Muench. i. (1829-30) 245 = crassicaulis.
arachnoides, *Prog. in Mart. Fl. Bras.* xii. II. 494.—Bras.
arborea, Hort.—Cf. Gard. Chron. (1881) II. 183 = stricta.
arborescens, *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 430.—Hab.?
arborescens, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 133.—Guian.
Arbuscula, *Barn. in C. Gay, Chil.* i. 443. t. 14.—Chili?
arcuata, Jacq. Oxal. 67. t. 31 = linearis.
arenaria, *Bert. in Merc. Chil.* n. xvi. 739, 740; *ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 48.—Chili.
aretioides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 436 = linearis.
argentea, Eckl. & Zeyh. Enum. 85 = bifurca.
Arracacha, G. Don. Gen. Syst. i. 756 = crassicaulis.
arthrophylla, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 433 = laburnifolia.
articulata, *Savign. in Lam. Encyc.* iv. 686.—Reg. Argent.
asinina, *Jacq. Oxal.* 59. t. 24.—Afr. austr.
aureo-flava, Steud. in Flora, xxxix. (1856) 443; et ex Phil. in Linnaea, xxviii. (1856) 617 = parvifolia.
autumnalis, *Poepp. ex Prog. in Mart. Fl. Bras.* xii. II. 477.—Peruv.; Mexic.
autumnalis, A. St. Hil. Fl. Bras. Mer. i. 128 = eriorhiza.
bahiensis, *Prog. in Mart. Fl. Bras.* xii. II. 501.—Bras.
balsamifera, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 335.—Afr. austr.
Barrelieri, *Linn. Sp. Pl.* ed. II. 624; *Jacq. Oxal.* 24. t. 3.—Am. austr.
Barrelieri, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823-24) 174 = sepium.
Berlandieri, *Torr. in Bot. Mex. Bound.* 41.—Am. bor.
Berteroana, *Barn. in C. Gay, Fl. Chil.* i. 437. t. 14.—Chili.
bicolor, Savign. in Lam. Encyc. iv. 687 = megalorrhiza.
bifida, *Thunb. Prod. Pl. Cap.* 82.—Afr. austr.
bifida, Vell. Fl. Flum. 196; iv. t. 183 = rhombeo-ovata.
bifida, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823-24) 162 = pubescens.
biflora, Burm. f. Fl. Cap. Prod. 13 = O. Pes-caprae.
bifolia, *Eckl. & Zeyh. Enum.* 92.—Afr. austr.
bifolia, Page, Prod. Southamp. Bot. Gard. 177 (Quid?)—Afr. austr.

OXALIS:—
bifrons, *Prog. in Mart. Fl. Bras.* xii. II. 495.—Bras.
bifurca, Eckl. & Zeyh. Enum. 85 = heterophylla.
bifurca, *Lodd. Bot. Cab.* t. 1056.—Afr. austr.
biglandulosa, *Steud. in Flora*, xxxix. (1856) 444.—Chili.
bipartita, *A. St. Hil. Fl. Bras. Mer.* i. 125.—Bras.
bipunctata, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1827) 176; & in Hook. Bot. Mag. t. 2781 = corymbosa.
bisfracta, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 595.—Bolivia.
bisulca, E. Mey. in Drège, Zwei Pfl. Docum. 44, nomen = stellata.
Blanchetiana, Moric. ex Prog. in Mart. Fl. Bras. xii. II. 518 = Eichleria Blanchetiana.
Blumei, *Zucc, in Abh. Akad. Muench.* i. (1829–30) 276.—Java.
borjensis, *H. B. & K. Nov. Gen. et Sp.* v. 249.—Am. austr.
Bowieana, Lodd. Bot. Cab. t. 1782 = purpurata.
Bowiei, Lindl. Bot. Reg. t. 1585 = praec.
brasiliensis, *Lodd. Bot. Cab.* t. 1962.—Bras.
Brehmiana, Eckl. & Zeyh. Enum. 86 = bifurca.
brevicaulis, *Sond. in Harv. & Sond. Fl. Cap.* i. 344.—Afr. austr.
brevicaulis, *Steud. Nom.* ed. II. ii. 239.—Chili.
breviscapa, Eckl. & Zeyh. Enum. 91 = variabilis.
breviscapa, *Jacq. Oxal.* 95. t. 58.—Afr. austr.
Bridgesii, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 49.—Chili.
bryoides, *Phil. in Linnaea*, xxviii. (1856) 678.—Chili.
bupleurifolia, *A. St. Hil. Fl. Bras. Mer.* i. 117.—Bras.
Burkei, *Sond. in Harv. & Sond. Fl. Cap.* i. 320.—Afr. austr.
Burmanni, *Jacq. Oxal.* 41.—Afr. austr.
Bustillosii, *Phil. in Linnaea*, xxxviii. (1856) 614.—Chili.
caesia, *Phil. Fl. Atac.* 13.—Chili.
caespitosa, *A. St. Hil. Fl. Bras. Mer.* i. 122.—Bras.
cajanifolia, A. St. Hil. l. c. 110 = Barrelieri.
calcaria, Eckl. & Zeyh. Enum. 93 = punctata.
caledonica, *Sond. in Harv. & Sond. Fl. Cap.* i. 341.—Afr. austr.
calliantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 310.—Japon.
calligera, Sond. in Harv. & Sond. Fl. Cap. i. 333 = foveolata.
calophylla, *Prog. in Mart. Fl. Bras.* xii. II. 517.—Bras.
calthaefolia, Pohl, ex Prog. l. c. 512 = hirsutissima.
calva, *Prog. in Mart. Fl. Bras.* xii. II. 486.—Bras.
campestris, A. St. Hil. Fl. Bras. Mer. i. 116 = densifolia.
campestris, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 172.—Bras.
cana, *Sond. in Harv. & Sond. Fl. Cap.* i. 345.—Afr. austr.
candida, Eckl. & Zeyh. Enum. 94 = Smithiana.
Candolleana, Steud. Nom. ed. II. ii. 239 = Biophytum sensitivum.
canescens, Jacq. Oxal. 45. t. 11 = tubiflora.
capillacea, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 345.—Afr. austr.
caprina, Eckl. & Zeyh. Enum. 84 = livida.
caprina, E. Mey. ex Harv. & Sond. Fl. Cap. i. 349 = cernua.
caprina, Thunb. Diss. Oxal. 13; Prod. Pl. Cap. 82 = O. Pes-caprae.
carnosa, *Molina, Sagg. Chile*, ed. II. 288; *Lindl. Bot. Reg.* t. 1063.—Chili.
casta, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 183.—Bras.
cataractae, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 315 = magellanica.
cearensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 272.—Bras.
ceratilis, E. Mey. in Drège, Zwei. Pfl. Docum. 72, nomen = corniculata.
cernua, *Thunb. Diss. Oxal.* 14.—Afr. austr.
cespitosa, Rafin. New Fl. Am. ii. 27 = corniculata?
checoënsis, *Meyen, Reise*, i. 406.—Chili.
chilensis, Pers. Syn. i. 518 = rosea.
chinensis, Haw. ex G. Don, in Loud. Hort. Brit. Suppl. i. 595, nomen = stricta?
chrysantha, *Prog. in Mart. Fl. Bras.* xii. II. 491.—Bras.

OXALIS:—
ciliariflora, Eckl. & Zeyh. Enum. 93 = obtusa.
ciliaris, Eckl. & Zeyh. l. c. 88 = glabra.
ciliaris, *Jacq. Oxal.* 66. t. 30.—Afr. austr.
ciliaris, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 326, 328 = approximata.
ciliata, Spreng. ex Prog. in Mart. Fl. Bras. xii. II. 512 = hirsutissima.
ciliata, Spreng. Syst. ii. 423 = alata.
ciliifera, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
cineracea, *A. St. Hil. Fl. Bras. Mer.* i. 123.—Bras.
cinerea, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 156 = Haenkeana.
clandestina, *Phil. in Linnaea*, xxviii. (1856) 616.—Chili.
clandestina, *Prog. in Mart. Fl. Bras.* xii. II. 479.—Bras.
clavifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 330.—Afr. austr.
clavigera, Sond. l. c. 539 = praec.
cognata, Steud. in Lehm. Pl. Preiss. i. 160 = corniculata.
collina, *Eckl. & Zeyh. Enum.* 90.—Afr. austr.
Commersonii, *Pers. Syn.* i. 519.—Reg. Argent.
commutata, *Sond. in Harv. & Sond. Fl. Cap.* i. 333.—Afr. austr.
comosa, *E. Mey. ex Harv. & Sond. l. c.* 340.—Afr. austr.
comosa, *Prog. in Mart. Fl. Bras.* xii. II. 506.—Bras.
compacta, *Gill. in Hook. Bot. Misc.* iii. (1833) 162.—Chili.
compressa, *Linn. f. Suppl.* 243; *Thunb. Diss. Oxal.* 11.—Afr. austr.
concinna, Salisb. Prod. 322 = cernua.
conferta, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 516 = bupleurifolia.
confertissima, *A. St. Hil. Fl. Bras. Mer.* i. 122.—Bras.
conorrhiza, *Jacq. Oxal.* 26.—Paraguay.
Consolii, *Tod. Nuovi Gen. e Spec.* 31.—Afr. austr.?
controversa, *Tod. l. c.* 33.—Afr. austr.?
convexula, *Jacq. Oxal.* 92. t. 55.—Afr. austr.
Coppolerii, *Tod. Nuovi Gen. e Spec.* 32.—Afr. austr.?
coquimbana, *Phil. in Linnaea*, xxxiii. (1864–65) 34.—Chili.
cordata, *A. St. Hil. Pl. Us. Brés.* t. 45.—Bras.
Cornelli, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 246.—Ins. Galap.
corniculata, *Linn. Sp. Pl.* 435.—Amphig. temp. et trop.
corymbosa, *DC. Prod.* i. 696.—Ins. Mascar.
crassicaulis, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 163.—Peruv.; Mexic.
crassifolia, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 317 = corniculata.
crassifolia, Eckl. & Zeyh. Enum. 91 = truncatula.
crassipes, *Urb. in Eichl. Jahrb. Berl.* iii. (1884) 242.—Afr. trop.
cratensis, *Hook. Ic. Pl.* t. 361.—Bras.
crenata, *Jacq. Oxal.* 27.—Peruv.
crispa, Jacq. l. c. 58. t. 23 = fabiaefolia.
crispula, *Sond. in Harv. & Sond. Fl. Cap.* i. 325.—Afr. austr.
cruentata, Eckl. & Zeyh. Enum. 91 = truncatula.
cruentata, *Jacq. f. Eclog. Am.* i. 66. t. 45.—Afr. austr.
Cumingiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 426.—Ins. Philipp.
Cummingii, Herb. in Bot. Reg. t. 1545 = pubescens?
cumulata, *Pohl, ex Prog. in Mart. Fl. Bras.* xii. II. 506.—Bras.
cuneata, Herb. Un. It. ex Harv. & Sond. Fl. Cap. i. 337 = obtusa.
cuneata, *Jacq. Oxal.* 76. t. 40.—Afr. austr.
cuneata, Thunb. Fl. Cap. ed. Schult. 540 = glabra.
cuneifolia, Jacq. Oxal. 77. t. 41 = cuneata.
cuneifolia, E. Mey. ex Harv. & Sond. Fl. Cap. i. 329 = angusta.
cuprea, Lodd. Bot. Cab. t. 824 = obtusa.
cylindrica, E. Mey. in Drège, Zwei Pfl. Docum. 99 nomen = tenella.
cytisoides, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 178 = Barrelieri.
daphnaeformis, *Mikan, ex Zucc. in Abh. Akad. Muench.* i. (1829–30) 269.—Bras.
Darvalliana, *Knowles & Westc. Flor. Cab.* iii. (1840) 5.—Hab.?

OXALIS:—
Darwinii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 214.—Patagon.
debilis, *H. B. & K. Nov. Gen. et Sp.* v. 236. t. 466.—N. Granat.
decaphylla, *H. B. & K. l. c.* 238. t. 468.—Mexic.
decipiens, *Prog. in Mart. Fl. Bras.* xii. II. 507.—Bras.
deflexa, *Poepp. ex Prog. l. c.* 479.—Chili.
delicata, *Pohl, ex Prog. l. c.* 498.—Bras.
delta, Vell. Fl. Flum. 195; iv. t. 178 = papilionacea.
dendroides, *H. B. & K. Nov. Gen. et Sp.* v. 250.—N. Granat.
densiflora, Sond. in Harv. & Sond. Fl. Cap. i. 339 = seq.
densifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 431.—Afr. austr.
densifolia, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 169.—Bras.
dentata, Eckl. & Zeyh. Enum. 84 = O. Pes-caprae.
dentata, *Jacq. Oxal.* 38. t. 7.—Afr. austr.
depauperata, *Prog. in Mart. Fl. Bras.* xii. II. 505.—Bras.
Deppei, *Lodd. Bot. Cab.* t. 1500.—Am. austr.
Deppei, Schlecht. ex Hemsl. Biol. Centr. Am. Bot. i. 165 = tetraphylla.
Deppii, Sweet, Hort. Brit. ed. II. 102, nomen = Deppei?
depressa, Eckl. & Zeyh. Enum. 91 = collina, convexula.
dichondraefolia, *A. Gray, Pl. Wright.* i. 72.—Texas; Mexic.
dichotomiflora, *Steud. in Flora*, xxxix. (1856) 442.—Chili.
diffusa, Bor. Fl. Centr. Fr. ed. III. ii. 136, in obs. = corniculata.
diffusa, *Pohl, ex Prog. in Mart. Fl. Bras.* xii. II. 500.—Bras.
digitata, Hort. ex Poir. Encyc. Suppl. iv. 254 = pentaphylla.
Dillenii, Jacq. Oxal. 28 = corniculata.
discolor, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 257.—Mexic.
dissecta, E. Mey. in Drège, Zwei Pfl. Docum. 54, nomen = bifurca.
distans, *A. St. Hil. Fl. Bras. Mer.* i. 115.—Bras.
disticha, *Jacq. Oxal.* 52. t. 18.—Afr. austr.
divaricata, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 169.—Bras.
divergens, *Benth. Pl. Hartw.* 9.—Mexic.
divergens, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
divergens, Eckl. & Zeyh. Enum. 92 = tenella.
dormiens, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 182.—Bras.
Dregei, *Sond. in Harv. & Sond. Fl. Cap.* i. 318.—Afr. austr.
droseroides, *E. Mey. ex Harv. & Sond. l. c.* 330.—Afr. austr.
Drummondii, *A. Gray, Pl. Wright.* ii. 25.—Am. bor.
dumetorum, *Barn. in C. Gay, Fl. Chil.* i. 448.—Chili.
ebracteata, *Savign. in Lam. Encyc.* iv. 682.—Afr. austr.
Echegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 19.—Reg. Argent.
Eckloniana, *Presl, Bot. Bemerk.* 29.—Afr. austr.
Eckloniana, Steud. Nom. ed. II. ii. 239 = tenella.
Ehrenbergii, *Schlecht. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 313.—Mexic.
elatior, *Prog. in Mart. Fl. Bras.* xii. II. 504.—Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* v. 234. t. 466.—Peruv.
elegantula, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 434 = imbricata?
elongata, Eckl. & Zeyh. Enum. 88 = lineatis.
elongata, Jacq. Oxal. 73. t. 37 = versicolor.
enneaphilla, *Cav. Ic.* v. 7. t. 411.—Ins. Falkland.
erecta, Savign. in Lam. Encyc. iv. 685 = O. Pes-caprae.
eremobia, *Phil. in Anal. Univ. Chil.* (1865) II. 332.—Chili.
ericoides, Hort. Angl. ex Steud. Nom. ed. II. ii. 239 (Quid?).—Afr. austr.
eriocarpa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 28.—Reg. Magellan.
erioclados, *Prog. in Mart. Fl. Bras.* xii. II. 494.—Bras.
eriolepis, *Wedd. Chlor. And.* ii. 290.—Am. austr.
eriorhiza, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 142.—Reg. Argent.

OXALIS:—
erubescens, E. Mey. in Drège, Zwei Pfl. Docum. 103, nomen = commutata.
erythrorhiza, *Gill. in Hook. Bot. Misc.* iii. (1833) 162. —Chili.
euphorbioides, *A. St. Hil. Fl. Bras. Mer.* i. 113.—Bras.
europaea, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 309, 311 = corniculata.
exaltata, Jacq. Oxal. 86. t. 49 = mutabilis.
excisa, *Prog. in Mart. Fl. Bras.* xii. II. 496.—Bras.
exilis, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
fabaefolia, *Jacq. Oxal.* 62. t. 27.—Afr. austr.
falcata, Eckl. & Zeyh. Enum. 95 = pentaphylla.
falcata, Sond. in Harv. & Sond. Fl. Cap. i. 322 = linearis.
Falconiana, *Steud. Nom.* ed. II. ii. 239.—Chili.
fallax, Eckl. & Zeyh. Enum. 91 = purpurea.
fallax, Jacq. Oxal. 107. t. 67 = luteola.
fasciculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 272.—Mexic.
favosa, Turcz. l. c. xxxi. (1858) I. 435 = punctata.
ferruginata, Jacq. Hort. Schoenb. iii. 13 = mutabilis.
filicaulis, Jacq. l. c. ii. 42. t. 205 = bifida.
filifolia, Jacq. l. c. iii. 12. t. 273 = polyphylla.
filiformis, *H. B. & K. Nov. Gen. et Sp.* v. 245. t. 464. —N. Granat.
filiformis, Steud. Nom. ed. II. ii. 239, sphalm. = Indigofera sarmentosa.
flabellifolia, *Jacq. Oxal.* 117. t. 74.—Afr. austr.
flaccida, Jacq. l. c. 88. t. 51 = mutabilis.
flava, *Linn. Sp. Pl.* 433.—Afr. austr.
flavo-punctata, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 510 = Zuccarinii.
floribunda, Lehm. ex Lindl. Bot. Reg. t. 1123 = rosea.
floribunda, *Lehm. Ind. Sem. Hort. Hamb.* (1826) 17; *Link & Otto, Ic. Pl. Rar.* i. 19. t. 10.—Bras.
floribunda, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 510 = physocalyx.
florida, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 328.—Afr. austr.
florida, Salisb. Prod. 322 = corniculata.
fontana, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 87 = corniculata.
foveolata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 435.—Afr. austr.
fragrans, *Tod. Nuovi Gen. e Spec.* 34.—Hab.?
frutescens, Linn. Sp. Pl. 435 = Plumieri, *Jacq.*
frutescens, *Ruiz & Pav. ex G. Don, Gen. Syst.* i. 754. —Peruv.
fruticosa, Raddi, in Mem. Soc. Ital. Mod. xviii. Fis. (1820) 401 = rusciformis.
fruticosa, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 171 = borjensis.
Fruticula, *Phil. Fl. Atac.* 13.—Chili.
fulgida, Lindl. Bot. Reg. t. 1073 = hirta.
fulva, A. St. Hil. Pl. Us. Brés. t. 44 = hirsutissima.
furcata, Ell. Sketch, i. 527 = corniculata.
fuscata, *Jacq. Oxal.* 82. t. 45.—Afr. austr.
Galeotti, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 433.—Mexic.
Gardneriana, *Prog. in Mart. Fl. Bras.* xii. II. 497.—Bras.
Gaudichaudii, *Barn. in C. Gay, Fl. Chil.* i. 425.—Chili.
Gayana, *Phil. in Linnaea*, xxviii. (1856) 615.—Chili.
gemella, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 146 = debilis.
geminata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 163.—Chili.
geniculata, *Knowles & Westc. Flor. Cab.* i. (1837) 43.—Bras.
gigantea, *Barn. in C. Gay, Fl. Chil.* i. 433.—Chili.
glabella, E. Mey. in Drège, Zwei Pfl. Docum. 96 = punctata.
glabra, Meissn. ex Harv. & Sond. Fl. Cap. i. 321 = versicolor.
glabra, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
glandulitega, Salisb. Parad. Lond. t. 66 = linearis.
glandulitheca, Steud. Nom. ed. I. 579 = praec.
glandulosa, *Jacq. Oxal.* 83. t. 46.—Afr. austr.
glauca, *H. B. & K. Nov. Gen. et Sp.* v. 248. t. 471.—Bras.
glauca, Rafin. ex DC. Prod. i. 692 = corniculata.
glaucovirens, Sond. in Harv. & Sond. Fl. Cap. i. 336 = stenoptera.
Glazioviana, *Prog. in Mart. Fl. Bras.* xii. II. 494.—Bras.

OXALIS:—
glomerata, Hook. & Arn. Bot. Beech. Voy. 13 = tortuosa.
glutinosa, *Phil. in Linnaea*, xxviii. (1856) 614, 615.—Chili.
goniorhiza, *Eckl. & Zeyh. Enum.* 89.—Afr. austr.
goyazensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) I. 272.—Bras.
gracilenta, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 68 = Biophytum Reinwardtii.
gracilicaulis, Eckl. & Zeyh. Enum. 91 = Smithiana.
gracilis, Eckl. & Zeyh. l. c. 88 = polyphylla.
gracilis, *Jacq. Oxal.* 69. t. 33.—Afr. austr.
Grahamiana, *Benth. Pl. Hartw.* 9.—Mexic.
grammopetala, *Sond. in Harv. & Sond. Fl. Cap.* i. 335.—Afr. austr.
grandiflora, Jacq. Oxal. 91. t. 54 = variabilis.
grandifolia, *DC. Prod.* i. 696.—Mexic.
Griffithii, *Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind.* i. 436.—Reg. Himal.
grisea, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 28.—Bras.
gymnoclada, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 432.—Afr. austr.
gyrorrhiza, *Bert. in Merc. Chil.* n. xvi. 739; *ex Am. Journ. Sc.* Ser. I. xxiii. (1833) 261.—Chili.
Haenkeana, Spreng. Syst. iv. Cur. Post. 186 = melilotoides.
Haenkeana, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 233.—Peruv.
hapalconidea, Barn. in C. Gay, Fl. Chil. i. 450 = geminata.
hedysarifolia, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 504 = elatior.
hedysarifolia, Raddi, in Mem. Soc. Ital. Mod. xviii. Fis. (1820) 401 = Neaei.
hedysaroides, *H. B. & K. Nov. Gen. et Sp.* v. 247.—Peruv.; Bras.
Hernandesii, [Moç. & Sesse,] ex DC. Prod. i. 695 = decaphylla.
herpestica, *Schlecht. in Linnaea*, xxvii. (1854) 525.—Guatemala.
heterophylla, *DC. Prod.* i. 694.—Afr. austr.
heterophylla, Eckl. & Zeyh. Enum. 94 = stenodactyla.
heterophylla, E. Mey. in Drège, Zwei Pfl. Docum. 145 = bifurca.
Hieronymi, *Griseb. in Goett. Abh.* xxiv. (1879) 72.—Reg. Argent.
hilliifolia, Walp. Rep. i. 483, sphalm. = trilliifolium.
hirsuta, *Sond. in Harv. & Sond. Fl. Cap.* i. 327.—Afr. austr.
hirsuta, Steud. Nom. ed. I. 579 = hirta.
hirsutissima, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 179.—Bras.
hirta, *Linn. Sp. Pl.* 434.—Afr. austr.
hirtella, Jacq. Oxal. 48. t. 14 = hirta.
hirtella, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 160 = parvifolia.
hispida, Zucc. l. c. 170 = distans.
hispidula, *Zucc. l. c.* 143.—Reg. Argent.
holosericea, E. Mey. in Drège, Zwei Pfl. Docum. 84, nomen = truncatula.
holosericea, *Phil. in Linnaea*, xxviii. (1856) 678.—Chili.
holosericea, *Sond. in Harv. & Sond. Fl. Cap.* i. 338. —Afr. austr.
Hookeriana, *Pohl, ex Prog. in Mart. Fl. Bras.* xii. II. 478.—Chili.
humilis, *Thunb. Prod. Pl. Cap. App.* 190.—Afr. austr.
humistrata, Willd. ex Steud. Nom. ed. II. ii. 240 = corniculata.
imbricata, *Eckl. & Zeyh. Enum.* 93.—Afr. austr.
impatiens, *Vell. Fl. Flum.* 186; iv. t. 181.—Bras.
incana, Eckl. & Zeyh. Enum. 85 = bifurca.
incana, *Phil. in Linnaea*, xxviii. (1856) 677.—Chili.
incarnata, *Linn. Sp. Pl.* 433.—Afr. austr.
inops, Eckl. & Zeyh. Enum. 92 = convexula.
inscripta, E. Mey. in Drège, Zwei Pfl. Docum. 103, nomen = variabilis.
insipida, *A. St. Hil. Fl. Bras. Mer.* i. 109.—Bras.
intermedia, *A. Rich. Ess. Fl. Cub.* 315.—Cuba.
intermedia, Steud. Nom. ed. II. ii. 240 = Biophytum intermedium.
isosepala, *Tod. Nuovi Gen. e Spec.* 36.—Am. austr.?
Jacquiniana, *H. B. & K. Nov. Gen. et Sp.* v. 235. t. 466.—Mexic.
jamaicensis, *Macfad. Fl. Jamaic.* i. 182.—Jamaic.

OXALIS:—
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 308.—Japon.
javanica, *Blume, Bijdr.* 243.—Java.
laburnifolia, *Jacq. Oxal.* 63. t. 28.—Afr. austr.
lacicola, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
laciniata, *Cav. Ic.* v. 7. t. 412.—Am. austr.
lactea, Hook. Comp. Bot. Mag. i. (1835) 276 = magellanica.
lacunosa, Eckl. & Zeyh. Enum. 93 = obtusa.
laevigata, *Willd. Enum. Hort. Berol. Suppl.* 26.—Afr. austr.
lanata, *Linn. f. Suppl.* 244.—Afr. austr.
lancaefolia, Tod. Ind. Sem. Hort. Panorm. (1858) 24 = Coppolerii.
lanceaefolia, Jacq. Oxal. 61. t. 26 = asinina.
lasiandra, R. Grah. in Hook. Bot. Mag. t. 3896 = floribunda?
lasiandra, *Zucc. in Otto & Dietr. Allg. Gartenz.* ii. (1834) 245.—Mexic.
lasiopetala, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 149.—Reg. Argent.
lasiophylla, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 30.—Bras.
lateriflora, *Jacq. Hort. Schoenb.* ii. 41. t. 204.—Afr. austr.
latifolia, *H. B. & K. Nov. Gen. et Sp.* v. 237. t. 467.—Mexic.
latipes, Mart. ex Steud. Nom. ed. II. ii. 240 (Quid?).—Bras.
Laureola, *Prog. in Mart. Fl. Bras.* xii. II. 509.—Bras.
laxa, *Hook. & Arn. Bot. Beech. Voy.* 13.—Chili.
laxa, E. Mey. ex Harv. & Sond. Fl. Cap. i. 340 = ebracteata.
laxula, Eckl. & Zeyh. Enum. 90 = purpurea.
laxula, Jacq. Oxal. 94. t. 57 = variabilis.
laxula, Thunb. ex Harv. & Sond. Fl. Cap. i. 335 = luteola.
Lechleri, Steud. in Flora, xxxix. (1856) 443, in obs. = dumetorum.
lepida, Jacq. Oxal. 55. t. 21 = monophylla.
leporina, Jacq. l. c. 60. t. 25 = asinina.
leporina, Tod. Ind. Sem. Hort. Panorm. (1858) 24 = Consolii.
leptocalyx, *Sond. in Harv. & Sond. Fl. Cap.* i. 320.—Afr. austr.
leptophylla, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 172.—Bras.
leptopodes, *G. Don, Gen. Syst.* i. 755.—Peruv.
lespedezioides, *G. Don, l. c.* 754.—Peruv.
leucotricha, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 436.—Hab.?
libyca, Viv. Fl. Lib. Spec. 24. t. 13. f. 1 = cernua.
lignosa, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 179 = psoralioides.
ligulata, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 324.—Afr. austr.
lilacina, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 258.—Mexic.
limbata, E. Mey. in Drège, Zwei Pfl. Docum. 101, nomen = tenuifolia.
limosa, *Prog. in Mart. Fl. Bras.* xii. II. 487.—Bras.
Lindenii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 429.—Mexic.
lindsaeaefolia, *Hook. Lond. Journ. Bot.* iii. (1844) 141. —N. Granat.
linearis, Eckl. & Zeyh. Enum. 88 = glabra.
linearis, *Jacq. Oxal.* 68. t. 32.—Afr. austr.
linearis, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 168 = Zuccariniana.
lineata, *Gill. in Hook. Bot. Misc.* iii. (1833) 162.—Chili.
liniflora, Eckl. & Zeyh. Enum. 85 = lateriflora.
liniflora, *Prog. in Mart. Fl. Bras.* xii. II. 489.—Bras.
littoralis, Eckl. & Zeyh. Enum. 90 = minuta.
littoralis, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 501 = pentantha.
livescens, E. Mey. ex Harv. & Sond. Fl. Cap. i. 333 = commutata.
livida, Herb. ex Harv. & Sond. l. c. 338 = lanata.
livida, *Jacq. Oxal.* 39. t. 8.—Afr. austr.
lobata, *Sims, Bot. Mag.* t. 2386.—Chili.
Loddigesiana, Eckl. & Zeyh. Enum. 87 = tenuifolia.
longiflora, Linn. Sp. Pl. 433 = violacea.
longiscapa, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 164 = peduncularis.
longisepala, *Tod. Nuovi Gen. e Spec.* 77.—Afr. austr.

OXALIS:—
lotoides, *H. B. & K. Nov. Gen. et Sp.* v. 241.—N. Granat.
loxensis, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 145 = elegans.
Lujula, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 = javanica.
lunulata, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 200.—Mexic.
lupinifolia, *Jacq. Oxal.* 115. t. 72.—Afr. austr.
lupulina, H. B. & K. Nov. Gen. et Sp. v. 243 = corniculata.
lutea, Steud. Nom. ed. I. 579 = corniculata.
luteola, *Jacq. Oxal.* 103. t. 65.—Afr. austr.
luteola, E. Mey. ex Harv. & Sond. Fl. Cap. i. 338 = lanata.
lybica, Willk. & Lange, Prod. Fl. Hisp. iii. 521 = cernua.
Lyoni, Pursh, Fl. Am. Sept. i. 322 = corniculata.
macrogonya, Jacq. Oxal. 110. t. 70 = luteola.
macrogonya, Sieber, ex Presl, Bot. Bemerk. 29 = punctata.
macromischos, Spreng. Syst. iv. Cur. Post. 185 = hirta.
macrophylla, Dougl. ex Hook. Fl. Bor. Am. i. 118 = trilliifolia.
macrophylla, Hornem. Hort. Hafn. i. 428 = O. Pescaprae.
macrophylla, H. B. & K. Nov. Gen. et Sp. v. 237 = grandifolia.
macropoda, *Baker, in Journ. Linn. Soc.* xxi. (1884) 328.—Madag.
macropus, *Phil. in Linnaea*, xxviii. (1856) 617.—Chili.
macrorrhiza, *Gill. in Hook. Bot. Misc.* iii. (1833) 162.—Chili.
macrostylis, Jacq. Oxal. 43. t. 9 = tubiflora.
maculata, *Desf. Tabl.* ed. II. 168 (Quid?).—Afr. austr.
magellanica, *Forst. in Comm. Gotting.* ix. (1789) 33.—Reg. Magellan.
Majoranae, *Tod. Ind. Sem. Hort. Panorm.* (1858) 27; *Ann. Sc. Nat.* Sér. IV. xi. (1859) 381.—Hab.?
mallobolba, *Cav. Ic.* iv. 64. t. 393. f. 2.—Am. austr.
mandioccana, *Raddi, in Mem. Soc. Ital. Mod.* xviii. *Fis.* (1820) 400.—Bras.
marginata, Jacq. Oxal. 108. t. 68 = pulchella.
marginata, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 160 = albicans.
maritima, *Barn. in C. Gay, Fl. Chil.* i. 424.—Chili.
Martiana, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 144 = corymbosa.
mauritiana, *Lodd. Bot. Cab.* t. 1780.—Ins. Maurit.
medicaginea, *H. B. & K. Nov. Gen. et Sp.* v. 242.—N. Granat.
megalorrhiza, Bert. ex Steud. in Flora, xxxix. (1856) 441 = reticulata.
megalorrhiza, *Jacq. Oxal.* 33.—Peruv.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 184.—Bras.
Meissneri, *Sond. in Harv. & Sond. Fl. Cap.* i. 345.—Afr. austr.
melanosticta, *Sond. l. c.* 333.—Afr. austr.
melilotoides, A. St. Hil. Fl. Bras. Mer. i. 112 = campestris.
melilotoides, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 165.—Peruv.
Metziana, Miq. ex Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind. i. 438 = Biophytum Reinwardtii.
Meyeri, *Sond. in Harv. & Sond. Fl. Cap.* i. 328.—Afr. austr.
Meyeriana, Presl, Bot. Bemerk. 29 = goniorhiza?
micrantha, Bert. ex Colla, in Mem. Acc. Torin. xxxvii. (1834) 50 = laxa.
micrantha, Boj. ex Prog. in Mart. Fl. Bras. xii. II. 493 = corniculata.
microcarpa, *Benth. Pl. Hartw.* 115.—Am. austr.
microphylla, A. Cunn. ex Hook. f. Handb. N. Zeal. Fl. 38 = corniculata.
microphylla, H. B. & K. Nov. Gen. et Sp. v. 245 = parvifolia.
microphylla, Mart. ex Prog. in Mart. Fl. Bras. xii. II. 497 = densifolia.
microphylla, *Phil. in Linnaea*, xxviii. (1856) 678.—Chili.
microphylla, Poir. Encyc. Suppl. iv. 248 = corniculata.
microstachys, *Prog. ex Warm. in Kjoeb. Vidensk. Meddel.* (1879–80) 23.—Bras.
mimosoides, *A. St. Hil. Fl. Bras. Mer.* i. 107.—Bras.
miniata, Jacq. Oxal. 71. t. 35 = gracilis.
miniata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 320 = glabra.
minima, Eckl. & Zeyh. Enum. 92 = commutata.

OXALIS:—
minima, *Sond. in Harv. & Sond. Fl. Cap.* i. 336.—Afr. austr.
minima, Steud. Nom. ed. I. 579 = corniculata.
minor, Eckl. & Zeyh. Enum. 88 = glabra.
minuta, *Thunb. Diss. Oxal.* 8.—Afr. austr.
minutifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 431 = gymnoclada.
minutula, *Cesati, in Atti Soc. Acc. Sc. Nap.* v. (171) n. VII. p. 5.—Chili.
Mirbeli, *Dehnh. Mem. Sp. Oxal.* 2; *ex Walp. Rep.* i. 492.—Hab.?
modesta, *Phil. in Linnaea*, xxviii. (1856) 617.—Chili.
mollis, *H. B. & K. Nov. Gen. & Sp.* v. 241.—Reg. Andin.
monadelpha, Roxb. ex Wight & Arn. Prod. 142 = corniculata.
monophylla, *Linn. Mant.* ii. 241.—Afr. austr.
montana, Rafin. in Am. Monthly Mag. (1818) 266; Ann. Nat. 12 = O. Acetosella.
montevidensis, *Prog. in Mart. Fl. Bras.* xii. II. 491.—Am. trop.
Mooniana, Steud. Nom. ed. II. ii. 241 = Biophytum polyphyllum.
multibulbosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 595.—Ins. Maurit.
multicaulis, *Eckl. & Zeyh. Enum.* 87.—Afr. austr.
multiflora, Jacq. Oxal. 49. t. 15 = hirta.
multifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 434 = imbricata.
Mundtii, *Sond. in Harv. & Sond. Fl. Cap.* i. 323.—Afr. austr.
muscoides, *Phil. in Anal. Univ. Chil.* (1870) II. 166.—Chili.
mutabilis, *Sond. in Harv. & Sond. Fl. Cap.* i. 327.—Afr. austr.
myriophylla, *O. Hoffm. in Bremen, Abh.* vii. (1882) 243.—Madag.
myriophylla, *A. St. Hil. Fl. Mer.* i. 121.—Bras.
namaquana, *Sond. in Harv. & Sond. Fl. Cap.* i. 325.—Afr. austr.
nana, *A. St. Hil. ex Spreng. Syst.* iv. *Cur. Post.* 184.—Bras.
natans, Eckl. & Zeyh. Enum. 92 = obtusa.
natans, *Linn. f. Suppl.* 243.—Afr. austr.
Navieri, Jord. in F. Schultz, Arch. Fr. Fl. et Allem. 310 = stricta.
Neaei, *DC. Prod.* i. 691.—Am. trop.
nematodes, Spreng. Syst. ii. 429 = filiformis.
nemoralis, Salisb. Prod. 321 = O. Acetosella.
nemorosa, Eckl. & Zeyh. Enum. 94 = O. Smithiana.
nervosa, *Vell. Fl. Flum.* 196; iv. t. 180.—Bras.
Neuwiedii, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 267.—Bras.
nidulans, Eckl. & Zeyh. Enum. 92 = purpurea.
nidulans, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 436 = minima?
nigrescens, *A. St. Hil. Fl. Bras. Mer.* i. 113.—Bras.
nigricans, *Pohl, ex Warm. in Kjoeb. Vidensk. Meddel.* (1879–80) 21.—Bras.
nigricans, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 497 = Zuccariniana.
Noronhae, *Oliver, in Hook. Ic. Pl.* xiv. 21. t. 1226.—Ins. Fern. Noronha.
nubigena, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* I. (1843) 320.—Bolivia.
nuda, *Arn. l. c.* xviii. I. (1836) 326.—Zeylan.
nudiflora, [*Moç. & Sesse,*] *ex DC. Prod.* i. 695.—Mexic.
nutans, Hill. Veg. Syst. ix. 4 (Quid?).—Afr. austr.
obliquifolia, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 123.—Abyss.
obtriangulata, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 62.—Japon.
obtusa, *Jacq. Oxal.* 106. t. 79. f. 1.—Afr. austr.
oregana, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 211.—Am. bor.
ornata, *Phil. Fl. Atac.* 13.—Chili.
ornata, *Poepp. ex Prog. in Mart. Fl. Bras.* xii. II. 476.—Bras.
ornithopus, *Phil. Fl. Atac.* 13.—Chili.
Ortgiesi, *Regel, Gartenfl.* (1875) 1; *et in Act. Hort. Petrop.* iii. (1875) 117.—Peruv.
Ottonis, Klotzsch, in Otto & Dietr. Allg. Gartenz. vii. (1839) 313 = lobata.
Ottonis, *C. Morr. Belg. Hortic.* vi. (1856) 291. t. 73.—Cuba.
ovata, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 154.—Bras.
oxyptera, *Prog. in Mart. Fl. Bras.* xii. II. 489.—Bras.

OXALIS:—
pachyphylla, *Phil. in Linnaea*, xxviii. (1856) 610.—Chili.
pachyrhiza, *Wedd. Chlor. And.* ii. 291.—Peruv.
pallens, Eckl. & Zeyh. Enum. 89 = goniorhiza, versicolor.
paludosa, *A. St. Hil. Fl. Bras. Mer.* i. 121.—Bras.
palustris, *A. St. Hil. l. c.* 127.—Bras.
papilionacea, *Hoffmgg. ex Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 148.—Bras.
paposana, *Phil. Fl. Atac.* 13.—Chili.
paradoxa, Mart. ex Prog. in Mart. Fl. Bras. xii. II. 514 = mandioccana.
pardalis, *Sond. in Harv. & Sond. Fl. Cap.* i. 345.—Afr. austr.
parviflora, Lej. Fl. Spa, ii. 307 = O. Acetosella.
parvifolia, *DC. Prod.* i. 693.—Ecuador.
parvifolia, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 496 = densifolia.
parvula, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 354.—Mexic.
patula, Eckl. & Zeyh. Enum. 87 = tenuifolia.
paucifolia, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 505 = depauperata.
Pavonii, *G. Don, Gen. Syst.* i. 754.—Peruv.
Pearcei, *Phil. in Anal. Univ. Chil.* (1872) 684.—Chili.
pectinata, Jacq. Oxal. 118. t. 75 = flava.
pedunculoris, *H. B. & K. Nov. Gen. et Sp.* v. 239.—Ecuador.
penicillata, *Phil. in Anal. Univ. Chil.* (1872) 685.—Chili.
pentantha, *Jacq. Oxal.* 21. t. 1.—Am. austr.
pentaphylla, E. Mey. ex Harv. & Sond. Fl. Cap. i. 346 = variifolia.
pentaphylla, *Sims, Bot. Mag.* t. 1549.—Afr. austr.
pentaphylloides, Sond. in Harv. & Sond. Fl. Cap. i. 346 = variifolia.
Perdicaria, Bert. in Merc. Chil. n. xvi. 739; ex Am. Journ. Sc. Ser. I. xxiii. (1833) 260 = lobata.
perennans, Haw. Misc. 181 = corniculata.
Pes-caprae, *Linn. Sp. Pl.* 434.—Afr. austr.
Pes-caprae, Savign. in Lam. Encyc. iv. 685 = cernua.
Petersianum, C. Muell. in Walp. Ann. vii. 502 = Biophytum sensitivum.
Petersii, Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind. i. 437 = Biophytum Apodiscias.
petiolaris, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 350 = purpurata.
phellandrioides, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 54 = lateriflora.
physocalyx, *Zucc. ex Prog. in Mart. Fl. Bras.* xii. II. 510.—Bras.
pichinchensis, *Benth. Pl. Hartw.* 166.—Ecuador.
Pickeringii, *A. Gray, Bot. U. St. Exped.* i. 323.—Peruv.
pilosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 212 = corniculata.
pilosa, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 498 = pilulifera.
pilosissima, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 427.—Bras.
pilosiuscula, H. B. & K. Nov. Gen. et Sp. v. 243 = corniculata.
pilulifera, *Prog. in Mart. Fl. Bras.* xii. II. 498.—Bras.
Piottae, Colla, Hort. Ripul. 98. t. 1 = punctata?
plagiantha, E. Mey. in Drège, Zwei Pfl. Docum. 95, nomen = obtusa.
platensis, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 30.—Bras.
platycaulis, *Steud. Nom.* ed. II. ii. 241.—Chili.
platylepis, *Wedd. Chlor. And.* ii. 240.—Bolivia.
platypila, *Gill. in Hook. Bot. Misc.* iii. (1833) 163.—Chili.
Plumieri, *Jacq. Oxal.* 23.—Am. austr.
Plumieri, *Lindl. ex Prog. in Mart. Fl. Bras.* xii. II. 480.—Ins. Martinic.
Plumierii, Sieber, ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 174 = sepium.
Poeppigii, *Prog. l. c.* 511.—Bras.
Pohliana, *Zucc. ex Prog. l. c.* 512.—Bras.
polyantha, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 319 = geminata.
polymorpha, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 174 = Neuwiedii, rhombeo-ovata, roselata.
polyphylla, *Jacq. Oxal.* 75. t. 39.—Afr. austr.
polyscapa, Eckl. & Zeyh. Enum. 84 = stellata.
polytricha, *Sond. in Harv. & Sond. Fl. Cap.* i. 324.—Afr. austr.

OXALIS:—
praecox, *Lehm. Ind. Sem. Hort. Hamb.* (1826) 17.—*Cf. Linnaea*, iii. (1828) *Litt.* 11.—Hab. ?
praecox, *Prog. in Mart. Fl. Bras.* xii. II. 487.—Bras.
praetexta, *Prog. l. c.* 513.—Bras.
pratensis, Eckl. & Zeyh. Enum. 89 = minuta.
Preissiana, Steud. in Lehm. Pl. Preiss. i. 160 = corniculata.
primulaefolia, *Raddi, in Mem. Soc. Ital. Mod.* xviii. *Fis.* (1820) 400.—Bras.
procumbens, Steud. ex A. Rich. Tent. Fl. Abyss. i. 123 = corniculata.
prolifera, Arn. in Nov. Act. Nat. Cur. xviii. I. (1836) 327 = Biophytum proliferum.
prolongata, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 505 = sepium.
propinqua, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
prorepens, *Phil. in Linnaea*, xxviii. (1856) 618.—Chili.
prostrata, Haw. Misc. 183 = corniculata.
psilopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 434.—Afr. austr.
psilotricha, *Turcz. l. c.* 427.—Mexic.
psoraleoides, Mart. in Flora, xx. (1837) II. Beibl. 124, nomen (an seq. ?).—Bras.
psoralioides, *H. B. & K. Nov. Gen. et Sp.* v. 246. t. 470.—Bras.
puberula, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 43 = Neaei.
puberula, *Phil. in Anal. Univ. Chil.* (1865) II. 332.—Chili.
pubescens, Bert. ex Steud. Nom. ed. II. ii. 241 = brevicaulis, *Steud.*, laxa, platycaulis, O. Vinaquillo.
pubescens, *H. B. & K. Nov. Gen. et Sp.* v. 240.—Peruv.
pubescens, Stokes, Bot. Mat. Med. ii. 558 = corniculata.
pulchella, Eckl. & Zeyh. Enum. 91, partim = obtusa.
pulchella, *Jacq. Oxal.* 109. t. 69.—Afr. austr.
pulchella, Salisb. Prod. 321 = violacea.
pumila, Nutt. ex Torr. & Gray, Fl. N. Am. i. 212 = corniculata.
pumila, Urv. in Mém. Soc. Linn. Par. iv. (1826) 616 = enneaphilla.
punctata, *Linn. f. Suppl.* 243.—Afr. austr.
purpurata, *Jacq. Hort. Schoenb.* iii. 57. t. 356.—Afr. austr.
purpurea, *Linn. Sp. Pl.* 433; *Thunb. Diss. Oxal.* 12; *Jacq. Oxal.* 93. t. 56.—Afr. austr.
pusilla, Jacq. Oxal. 79 = glabra.
pusilla, E. Mey. ex Harv. & Sond. Fl. Cap. i. 319 = goniorhiza.
pusilla, Salisb. Prod. 322 = corniculata.
pycnophylla, *Wedd. Chlor. And.* ii. 291.—Bolivia.
pygmaea, *A. Gray, Bot. U. St. Expl. Exped.* 322.—Peruv.
pygmaea, E. Mey. in Drège, Zwei Pfl. Docum. 105 = punctata.
pygmaea, *Phil. in Linnaea*, xxviii. (1856) 614.—Chili.
quinata, *Savign. in Lam. Encyc.* iv. 688.—Afr. austr.
quinquepunctata, *Desf. Tabl.* ed. II. 168 (Quid ?).—Afr. austr.
racemosa, Savign. in Lam. Encyc. iv. 684 = rosea.
radiata, *Pohl, ex Prog. in Mart. Fl. Bras.* xii. II. 506.—Bras.
radicosa, A. Rich. Tent. Fl. Abyss. i. 123 = corniculata.
ramigera, Sond. in Harv. & Sond. Fl. Cap. i. 327 = gymnoclada.
ramosa, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 504 = elatior.
recisa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 21 = repens.
reclinata, Jacq. Oxal. 70. t. 34 = gracilis.
rectangularis, E. Mey. in Drège, Zwei Pfl. Docum. 147, nomen = Smithiana.
recticaulis, *Sond. in Harv. & Sond. Fl. Cap.* i. 330.—Afr. austr.
recurva, Ell. Sketch, i. 526 = corniculata.
refracta, *A. St. Hil. Fl. Bras. Mer.* i. 119.—Bras.
Regnellii, *Miq. in Linnaea*, xxii. (1849) 545.—Bras.
Reinwardtii, Zucc. in Abh. Akad. Muench. i. (1829–30) 274 = Biophytum Reinwardtii.
repens, Thunb. Diss. Oxal. 16; Prod. Pl. Cap. 82 = corniculata.
reptans, *Soland. ex Forst. f. Prod.* 90.—Ins. Norfolk.

OXALIS:—
reptatrix, Jacq. Oxal. 54 = variabilis.
reptatrix, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 344 = brevicaulis.
reticulata, *Steud. in Flora*, xxxix. (1856) 441.—Chili.
revoluta, E. Mey. in Drège, Zwei Pfl. Docum. 101, nomen = polyphylla.
rhombeo-ovata, *A. St. Hil. Fl. Bras. Mer.* i. 108.—Bras.
rhombifolia, *Jacq. Oxal.* 22. t. 2.—Venezuela.
Richardiana, Steud. Nom. ed. II. ii. 241 = corniculata.
rigidula, Jacq. Oxal. 96 = variabilis.
rosacea, Jacq. Oxal. 51. t. 17 = hirta.
rosacea, Tod. Ind. Sem. Hort. Panorm. (1858) 24 = controversa,
rosea, *Jacq. Oxal.* 25.—Chili.
rosea, Sims, Bot. Mag. t. 2415 = Simsii.
roselata, *A. St. Hil. Fl. Bras. Mer.* i. 109.—Bras.
rostrata, Jacq. Oxal. 57. t. 22 = monophylla.
rostrata, E. Mey. in Drège, Zwei Pfl. Docum. 67, nomen = Dregei.
rubella, *Jacq. Oxal.* 50. t. 16 = hirta.
rubens, Haw. Misc. 182 = corniculata.
rubra, *A. St. Hil. Fl. Bras. Mer.* i. 124.—Bras.
rubrocincta, *Lindl. Bot. Reg.* (1842) t. 64.—Guatemala.
rubro-flava, Jacq. Oxal. 87. t. 50 = mutabilis.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 429.—Peruv.
rufescens, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 162 = mollis.
rufo-pilosa, *Schult. ex Prog. in Mart. Fl. Bras.* xii. II. 477.—Reg. Andin.
rugulosa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 436 = natans.
rupestris, Rafin. New Fl. Am. ii. 26 (Quid ?).—Am. bor.
rupestris, *A. St. Hil. Fl. Bras. Mer.* i. 126.—Bras.
rusciformis, *Mikan, Delect. Fl. et Faun. Bras.* fasc. iii. [sine num.].—Bras.
Rutenbergii, *O. Hoffm. in Bremen, Abh.* vii. (1882) 242.—Madag.
saliciformis, *Mikan, ex Zucc. in Abh. Akad. Muench.* i. (1829–30) 270.—Bras.
sanguinea, Jacq. Oxal. 64. t. 29 = laburnifolia.
sanguinolaria, Rafin. Fl. Ludov. 89 (Quid ?).—Am. bor.
sarmentosa, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 244.—Bras.
saxatilis, *A. St. Hil. Fl. Bras. Mer.* i. 114.—Bras.
scandens, *H. B. & K. Nov. Gen. et Sp.* v. 242.—N. Granat.
Schiedeana, *Zucc. ex Steud. Nom.* ed. II. ii. 241 (Quid ?).—Mexic.
Schomburgkiana, *Prog. in Mart. Fl. Bras.* xii. II. 500.—Bras.
Schraderiana, *H. B. & K. Nov. Gen. et Sp.* v. 236. t. 466.—N. Granat.
secunda, Jacq. Oxal. 46. t. 12 = tubiflora.
segetalis, Poepp. ex Prog. in Mart. Fl. Bras. xii. II. 511 = Poeppigii.
Selloi, Spreng. Syst. v. 496 (Index) = Zuccariniana.
Sellowiana, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 198.—Bras.
semiglandulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 330.—Afr. austr.
semiglauca, Eckl. & Zeyh. Enum. 88 = goniorhiza.
semiloba, *Sond. in Harv. & Sond. Fl. Cap.* i. 350.—Afr. austr.
sensitiva, Herb. ex Edgew. & Hook. f. in Fl. Brit. Ind. i. 439 = Biophytum intermedium.
sensitiva, Linn. Sp. Pl. 434 = Biophytum sensitivum.
sensitiva, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 183. t. 5 B = Biophytum Reinwardtii.
sepium, *A. St. Hil. Fl. Bras. Mer.* i. 111.—Panama.
sericea, *Linn. f. Suppl.* 243.—Afr. austr.
serpens, *A. St. Hil. Fl. Bras. Mer.* i. 120.—Bras.
sessilifolia, Linn. Mant. ii. 241 = hirta.
sessilis, Buch.-Ham. in Wall. Cat. n. 4344 = Biophytum Apodiscias.
setosa, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 334.—Afr. austr.
sexenata, Savign. in Lam. Encyc. iv. 687 = Commersonii.
Simsii, *Sweet, Hort. Brit.* ed. II. 102.—Chili.
simulans, *Baker, in Journ. Linn. Soc.* xx. (1883) 112.—Madag.
Smithiana, *Eckl. & Zeyh. Enum.* 94.—Afr. austr.
Smithii, Sond. in Harv. & Sond. Fl. Cap. i. 342 = praec.

OXALIS:—
soldanellaefolia, Pohl, ex Prog. in Mart. Fl. Bras. xii. II. 513 = Pohliana.
somnians, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 181.—Bras.
speciosa, Eckl. & Zeyh. Enum. 90 = purpurea.
speciosa, Jacq. Oxal. 97. t. 60 = variabilis.
spectabilis, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 653 = brasiliensis.
sphacelata, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 170 = glauca.
spiralis, *G. Don, Gen. Syst.* i. 755.—Peruv.
spodiophylla, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* v. (1843) 318.—Chili.
Spruceana, *Prog. in Mart. Fl. Bras.* xii. II. 481.—Peruv.
squamata, *Zucc. in Denkschr. Akad. Muench.* ix. (1823–24) 157.—Chili.
squamoso-radicosa, *Steud. in Flora*, xxxix. (1856) 443.—Reg. Magell.
squarrosa, *Barn. in C. Gay, Fl. Chil.* i. 445.—Chili.
staphyloides, Mart. ex Steud. Nom. ed. II. ii. 242 = seq. ?
staphyleoides, *Prog. in Mart. Fl. Bras.* xii. II. 509.—Bras.
stellata, *Eckl. & Zeyh. Enum.* 84.—Afr. austr.
stenodactyla, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 437 = variifolia.
stenophylla, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 54 = monophylla.
stenoptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 433.—Afr. austr.
Sternbergii, Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 156 = Commersonii.
stictophylla, *Sond. in Harv. & Sond. Fl. Cap.* i. 335.—Afr. austr.
stricta, Linn. Sp. Pl. 435 = corniculata.
strictula, *Steud. in Flora*, xxxix. (1856) 443.—Chili.
strigulosa, *Prog. in Mart. Fl. Bras.* xii. II. 484.—Bras.
strumosa, Jacq. Oxal. 102. t. 64 = mutabilis.
strumosa, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 335 = luteola.
stylosa, E. Mey. in Drège, Zwei Pfl. Docum. 95, nomen = obtusa.
subacaulis, *Gill. in Hook. Bot. Misc.* iii. (1833) 163.—Chili.
subcarnosa, *Klotzsch, ex Walp. Rep.* ii. 822.—Chili.
succulenta, *Barn. in C. Gay, Fl. Chil.* i. 442.—Chili.
suffruticosa, *Salzm. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 427.—Bolivia.
suggillata, Jacq. Oxal. 98. t. 61 = variabilis.
sulphurea, Eckl. & Zeyh. Enum. 91 = Eckloniana.
sulphurea, *Jacq. Oxal.* 100. t. 63.—Afr. austr.
sulphurea, E. Mey. in Drège, Zwei Pfl. Docum. 207 = luteola.
sylvestris, Steud. Nom. ed. I. 579; ed. II. ii. 242 = versicolor.
tenella, Eckl. & Zeyh. Enum. 92 = commutata.
tenella, *Jacq. Oxal.* 53. t. 19.—Afr. austr.
tenella, Lodd. Bot. Cab. t. 1096 = minuta.
tenera, *Phil. in Anal. Univ. Chil.* (1870) II. 166.—Chili.
tenera, Spreng. Syst. ii. 424; Lindl. Bot. Reg. t. 1046 = lobata.
tenera, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823–24) 145 = Jacquiniana.
tenuicaulis, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 316 = corniculata.
tenuifolia, Eckl. & Zeyh. Enum. 87 = multicaulis.
tenuifolia, *Jacq. Oxal.* 74. t. 38.—Afr. austr.
tenuifolia, E. Mey. in Drège, Zwei Pfl. Docum. 105, nomen = versicolor.
tephrodes, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 427.—Mexic.
teretifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 322.—Afr. austr.
tetraphylla, *Cav. Ic.* iii. 19.—Mexic.
tetraphylla, Link & Otto, Ic. Pl. Rar. i. 21. t. 11 = Deppei.
thermarum, Eckl. & Zeyh. Enum. 93 = obtusa.
Thunbergiana, Eckl. & Zeyh. l. c. 87 = tenuifolia.
Thunbergiana, *Montr. in Mém. Acad. Lyon*, x. (1860) 192.—N. Caled.
tinctoria, *Poepp. ex Prog. in Mart. Fl. Bras.* xii. II. 479.—Chili.
tomentella, *Pohl, ex Prog. l. c.* 511.—Bras.

OXALIS:—
tomentosa, *Linn. f. Suppl.* 244.—Afr. austr.
tortuosa, *Lindl. Bot. Reg.* t. 1249.—Chili.
triangularis, *A. St. Hil. Fl. Bras. Mer.* i. 128.—Bras.
trichocalyx, *Phil. Fl. Atac.* 18.—Chili.
trichocalyx, *Steud. Nom.* ed. II. ii. 242.—Chili.
tricolor, Eckl. & Zeyh. Enum. 94 = affinis.
tricolor, Jacq. Oxal. 84 = mutabilis.
trilliifolia, *Hook. Fl. Bor. Am.* i. 118.—Am. bor.
tristis, Eckl. & Zeyh. Enum. 93 = Smithiana.
tropaeoloides, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 584 = corniculata.
truncata, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1843-44) 148 = papilionacea.
truncatula, *Jacq. Oxal.* 99. t. 62.—Afr. austr.
tuberosa, *Molina, Sagg. Chile*, 132, 352; *Savign. in Lam. Encyc.* iv. 684.—Chili.
tubiflora, Eckl. & Zeyh. Enum. 86, ex parte = hirta.
tubiflora, *Jacq. Oxal.* 44. t. 10.—Afr. austr.
tubistipula, Steud. ex Phil. f. Cat. Pl. Vasc. Chil. 40 = rosea.
uitenhagensis, Sond. in Harv. & Sond. Fl. Cap. i. 337 = psilopoda.
umbraticola, *A. St. Hil. Fl. Bras. Mer.* i. 111.—Bras.
umbrosa, Willd. ex Zucc. in Denkschr. Akad. Muench. ix. (1823-24) 147 = grandifolia.
undulata, Jacq. Oxal. 81. t. 44 = mutabilis.
urbica, A. St. Hil. Fl. Bras. Mer. i. 126 = corymbosa.
Urvillei, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 315 = corniculata.
valdiviana, *Hort. Veitch, ex Vilm. Fl. Pl. Terre*, ed. III. 807 (an seq. ?).—Chili.
valdiviensis, *Barn. in C. Gay, Fl. Chil.* i. 446.—Chili.
variabilis, *Jacq. Oxal.* 89.—Afr. austr.
variifolia, *Steud. Nom.* ed. II. ii. 242.—Afr. austr.
venosa, Savign. in Lam. Encyc. iv. 681 = glabra.
venusta, Lowe, Man. Fl. Mad. i. 595 = purpurea.
verecunda, Salisb. Prod. 322 = incarnata.
versicolor, Jacq. Ic. Pl. Rar. t. 478 = polyphylla.
versicolor, *Linn. Sp. Pl.* 434.—Afr. austr.
versicolor, Tod. Ind. Sem. Hort. Panorm. (1858) 24 = fragrans.
verticillata, [*Moç. & Sesse,*] *ex DC. Prod.* i. 691.—Mexic.
verticillata, Vell. Fl. Flum. 195; iv. t. 179 = primulaefolia.
vespertilionis, A. Gray. in Mem. Am. Acad. iv. (1849) 27 = violacea.
vespertilionis, Torr. & Gray, Fl. N. Am. i. 679 = Drummondii.
vespertilionis, *Zucc. in Otto & Dietr. Allg. Gartenz.* ii. (1834) 245.—Mexic.
villosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 112.—Madag.
villosa, Bieb. Fl. Taur. Cauc. i. 355 = corniculata.
villosa, *Prog. in Mart. Fl. Bras.* xii. II. 495.—Bras.
villosissima, Pohl, ex Prog. l. c. 512 = goyazensis.
Vinaquillo, *Steud. Nom.* ed. II. ii. 242; *et in Flora*, xxxix. (1856) 442.—Chili.
violacea, *Linn. Sp. Pl.* 434.—Am. bor.
violacea, Savign. in Lam. Encyc. iv. 686 = bipunctata.
virginea, *Jacq. Hort. Schoenb.* iii. 13. t. 275.—Afr. austr.
virginea, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 328 = florida.
virginica, Steud. Nom. ed. I. 580 = virginea.
virgosa, Molina, Sagg. Chile, 132 = gigantea.
viscosa, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 344.—Afr. austr.
viva, Willd. ex Steud. Nom. ed. II. ii. 242 = dendroides.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 630 = O. Acetosella.
Wallichiana, Gill. in Hook. Bot. Misc. iii. (1833) 162 = andicola.
Wrightii, *A. Gray, Pl. Wright.* i. 27.—N. Mexic.
xiphophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 112.—Madag.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 347.—Afr. austr.
zonata, *Liebm. Ind. Sem. Hort. Haun.* (1853) 20; *et in Ann. Sc. Nat.* Sér. IV. i. (1854) 329.—Chili?
Zuccariniana, *Steud. Nom.* ed. II. ii. 242.—Bras.
Zuccarinii, *Prog. in Mart. Fl. Bras.* xii. II. 510.—Bras.

OXALISTYLIS, Baill. Etud. Gén. Euph. 628 (1858) = **Phyllanthus**, Linn. (Euphorb.).
Kunthiana, Baill. l. c. 629 = P. salicifolius.

OXALLIS, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 21 = **Oxalis**, Linn. (Geran.).

OXANDRA, A. Rich. Ess. Fl. Cub. 45. t. (1842) = **Bocagea**, A. St. Hil. (Anonac.).
aromatica, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 36 = B. aromatica.
laurifolia, A. Rich. Ess. Fl. Cub. 49.—Cuba.
leucodermis, Warm. in Kjoeb. Vidensk. Meddel. (1878) 147 = B. leucodermis.
Reinhardtiana, Warm. l. c.—Bras.
virgata, A. Rich. Ess. Fl. Cub. 47.—Ind. occ.

OXANTHERA, Montr. in Mém. Acad. Lyon, x. (1860) 186 = **Citrus**, Tourn. (Rutac.).
fragrans, Montr. l. c.—N. Caled.

OXERA, Labill. Sert. Austro-Caled. 23. t. 28. (1824).
VERBENACEAE, Benth. & Hook. f. ii. 1155.
ONCOMA, Spreng Syst. iv. Cur. Post. 11. (1827).
MAOUTIA, Montr. in Mém. Acad. Lyon, x. (1860) 241.
baladica, *Vieill. in Bull. Soc. Linn. Normand.* vii. (1863) 91.—N. Caled.
cordifolia, *Vieill. l. c.* 95.—N. Caled.
glandulosa, *Vieill. l. c.* 93.—N. Caled.
longifolia, *Vieill. l. c.* 92.—N. Caled.
Morieri, *Vieill. l. c.* 93.—N. Caled.
oblongifolia, *Vieill. l. c.* 94.—N. Caled.
ovata, *Vieill. l. c.* 95.—N. Caled.
pulchella, *Labill. Sert. Austro-Caled.* i. 23. t. 28.—N. Caled.
robusta, *Vieill. in Bull. Soc. Linn. Normand.* vii. (1863) 92.—N. Caled.
subverticillata, *Vieill. l. c.* 91.—N. Caled.

OXEROSTYLUS, Steud. Nom. ed. II. ii. 242. (1841) sphalm. = OGCEROSTYLUS, Cass. = **Angianthus**, Wendl. (Compos.).

OXIA, Reichb. Nom. 69 (1841), sphalm. = AXIA, Lour. = **Boerhaavia**, Linn. (Nyctag.).

OXICEROS, Lour. Fl. Cochinch. 150 (1790), sphalm. = OXYCEROS, Lour. = **Randia**, Linn. (Rubiac.).

OXICOCCUS, Neck. Elem. i. 217 (1790) = **Oxycoccus**, Tourn. (Vaccin.).

OXIPHOERIA, Hort. Angl. ex Dum.-Cours. Bot. Cult. ed. II. iv. 205 (1811) = **Humea**, Sm. (Compos.).
foetida, Hort. Angl. l. c. = H. elegans.

OXIS, Medic. Malv. 113 (1787) = OXYS, Tourn. = **Oxalis**, Linn. (Geran.).

OXISMA, Rafin. Sylva Tellur. 94 (1838) = **Loreya**, DC. (Melastom.).
arborescens, Rafin. l. c. = L. arborescens.
flavescens, Rafin. l. c. = Henriettella flavescens.

OXLEYA, A. Cunn. in Hook. Bot. Misc. i. (1830) 246. t. 54 = **Flindersia**, R. Br. (Meliac.).
xanthoxyla, A. Cunn. l. c. = F. Oxleyana.

OXODIUM, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
callosum, Rafin. l. c. = P. callosum.

OXODON, Steud. Nom. ed. II. ii. 75, in syn. (1841), sphalm. = OXYDON, Less. = **Chaptalia**, Vent. (Compos.).

OXYACANTHA, Medic. Phil. Bot. i. 150 (1789) = **Crataegus**, Tourn. (Rosac.).
apiifolia, M. Roem. Syn. Rosifl. 108 = C. Oxyacantha.
Azorella, M. Roem. l. c. 107 = C. Oxyacantha.
elegans, M. Roem. l. c. 108 = C. Oxyacantha.
granatensis, M. Roem. l. c. 106 = C. granatensis.
kyrtostyla, M. Roem. l. c. = C. Oxyacantha.
laciniata, M. Roem. l. c. 109 = C. laciniata.
macrocarpa, M. Roem. l. c. 107 = C. Oxyacantha.
media, M. Roem. l. c. 108 = C. Oxyacantha.
monogyna, M. Roem. l. c. 107 = C. Oxyacantha.
obtusata, M. Roem. l. c. = C. Oxyacantha.
Pallasii, M. Roem. l. c. 106 = C. pentagyna.
sicula, M. Roem. l. c. 109 = C. laciniata.
vulgaris, M. Roem. l. c. = C. Oxyacantha.

OXYADENIA, Spreng. Syst. i. 351, in syn. (1825) = OXYDENIA, Nutt. = **Leptochloa**, Beauv. (Gramin.).

OXYANDRA, Reichb. Handb. 303 (1837) = **Sloanea**, Linn. (Tiliac.).

OXYANTHE, Steud. Syn. Pl. Gram. 197 (1855) = **Phragmites**, Trin. (Gramin.).
japonica, Steud. l. c. = P. communis.

OXYANTHERA, Brongn. in Duperr. Voy. Coq. Bot. 197. t. 37 (1829) = **Thelasis**, Blume (Orchid.).
micrantha, Brongn. l. c.—Ins. Molucc.

OXYANTHUS, DC. in Ann. Mus. Par. ix. (1807) 218. *RUBIACEAE*, Benth. & Hook. f. ii. 91.
MEGACARPHA, Hochst. in Flora, xxvii. (1844) 551.
breviflorus, *Benth. in Hook. Niger Fl.* 388.—Afr. trop.
cymosus, Reichb. ex DC. Prod. iv. 372 = Mussaenda Landia.
formosus, *Hook. f. in Hook. Ic. Pl.* t. 785.—Afr. trop.
Gerrardi, *Sond. in Harv. & Sond. Fl. Cap.* iii. 3.—Afr. austr.
gracilis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 109.—Afr. trop.
hirsutus, DC. Prod. iv. 376 = tubiflorus.
latifolius, *Sond. in Harv. & Sond. Fl. Cap.* iii. 4.—Afr. austr.
longiflorus, Hort. ex Lem. in Fl. des Serres, ii. (Sept. 1846) 148 = versicolor.
macrophyllus, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 110.—Afr. trop.
multiflorus, Hassk. Cat. Hort. Bog. Alt. 115 = Randia longiflora.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 50.—Afr. trop.
platystylis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 109.—Afr. trop.
querimbensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 292.—Afr. trop.
rubiflorus, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 108.—Afr. trop.
Smithii, *Hiern, l. c.* 107.—Afr. trop.
speciosus, Ait. Hort. Kew. ed. II. i. 371 = tubiflorus.
speciosus, *DC. in Ann. Mus. Par.* ix. (1807) 218.—Afr. trop.
sulcatus, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 108.—Afr. trop.
Thonningii, *Benth. in Hook. Niger Fl.* 388.—Afr. trop.
tubiflorus, *DC. Prod.* iv. 376.—Afr. trop.
tubiflorus, Hort. ex Lem. in Fl. des Serres, ii. (1846) sub t. 148 = versicolor.
unilocularis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 110.—Afr. trop.
versicolor, *Lindl. Bot. Reg.* (1840) *Misc.* 69.—Cuba.
villosus, *G. Don, Gen. Syst.* iii. 494.—Afr. trop.

OXYBAPHUS, L'Hérit. ex Willd. Sp. Pl. i. 185 (1797); Vahl, Enum. ii. 39 (1806). *NYCTAGINEAE*, Benth. & Hook. f. iii. 4.
ALLIONIA, Loefl. It. 181 (1758); Linn. Syst. ed. X. 890 (1759).
CALYMENIA, Pers. Syn. i. 36 (1805).
CALYXHYMENIA, Orteg. Hort. Matr. Dec. 5. t. 1 (1798).
VITMANIA, Turra, ex Cav. Ic. iii. Add. [53] (1794).
aggregatus, *Vahl, Enum.* ii. 41.—Mexic.
albidus, *Sweet, Hort. Brit.* ed. II. 429.—Am. bor.
angustifolius, *Sweet, l. c.* ed. I. 334.—Am. bor. occ.
bracteosus, *Griseb. in Goett. Abh.* xix. (1874) 86.—Reg. Argent.
californicus, *Benth. & Hook. f. Gen.* iii. 4.—Calif.
campestris, *Griseb. in Goett. Abh.* xix. (1874) 87.—Reg. Argent.
Cervantesii, Lag. ex Choisy, in DC. Prod. xiii. II. 432 = nyctagineus.
chilensis, Bert. ex Sweet, Hort. Brit. ed. III. 768 = ovatus.
chilensis, *Steud. Nom.* ed. II. ii. 213.—Chili.
coccineus, Torr. in Bot. Mex. Bound. 169 = Mirabilis coccinea.
cordifolius, *Kunze, ex Choisy, in DC. Prod.* xiii. II. 432.—Chili.
cucullatus, *Choisy, l. c.* 434.—Hab.?
decumbens, Sweet, Hort. Brit. ed. I. 334 = angustifolius.
elegans, *Choisy, in DC. Prod.* xiii. II. 431.—Chili.

OXYBAPHUS:—
expansus, *Vahl, Enum.* ii. 41.—Peruv.
floribundus, *Choisy, in DC. Prod.* xiii. II. 433.—Am. bor.
Froebellii, Behr, in Proc. Calif. Acad. i. (1855) 69 = Mirabilis Froebelii.
glaber, *S. Wats. in Am. Natural.* vii. (1873) 301.—Am. bor.
glabrifolius, *Vahl, Enum.* ii. 40.—Am. bor. occ.; Mexic.
himalaicus, *Edgew. in Trans. Linn. Soc.* xx. (1846) 87.—Reg. Himal.
hirsutus, *Sweet, Hort. Brit.* ed. I. 334; *Hook. Fl. Bor. Am.* ii. 124.—Am. bor. occ.
laevis, *Benth. Bot. Voy. Sulph.* 44 (an californicus?).—Calif.
linearifolius, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 375.—Am. bor.
micranthus, *Choisy, in DC. Prod.* xiii. II. 432.—Chili.
multiflorus, Torr. in Ann. Lyc. N. York, ii. (1828) 237 = Mirabilis multiflora.
nyctagineus, *Sweet, Hort. Brit.* ed. I. 334.—Am. bor.
ovatus, *Vahl, Enum.* ii. 40.—Peruv.
pauciflorus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 7 = nyctagineus.
pilosus, Sweet, Hort. Brit. ed. I. 334 = glabrifolius?
prostratus, Poepp. ex Choisy, in DC. Prod. xiii. II. 432 = cordifolius.
prostratus, *Vahl, Enum.* ii. 40.—Peruv.
Toscae, *Lor. in Roca, Exped. Rio-Negro*, ii. 195.—Patag.
violaceus, *Choisy, in DC. Prod.* xiii. II. 432; *Schlecht. in Linnaea*, xxvi. (1853–55) 644.—N. Granat.
viscosus, *L'Hérit. ex Choisy, in DC. Prod.* xiii. II. 430.—Peruv.
Wrightii, *Hemsl. Biol. Centr. Am. Bot.* iii. 3.—Texas.

OXYBASIS, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 738 = **Chenopodium**, Tourn.
minutiflora, Kar. & Kir. l. c. 739 = C. rubrum.

OXYCARPUS, Lour. Fl. Cochinch. 647 (1790) = **Garcinia**, Linn. (Guttif.).
celebica, Poir. Encyc. Suppl. iv. 258 = G. cornea, indica.
cochinchinensis, Lour. Fl. Cochinch. 648 = G. cochinchinensis.
gangetica, Buch.-Ham. in Mem. Wern. Soc. v. (1824) 344 = G. Cowa.
indica, Buch.-Ham. ex Wall. Cat. sub n. 4852 = G. cornea.
indica, Poir. Encyc. Suppl. iv. 257 = G. indica.
Sopsopia, Buch.-Ham. in Mem. Wern. Soc. v. (1824) 345 (Quid?).—Ind. or.

OXYCARYUM, Nees, in Mart. Fl. Bras. ii. 1. 90 (1842) = **Scirpus**, Linn. (Cyper.).
Schomburgkianum, Nees, l. c. = S. cubensis.

OXYCEDRUS, Hort. ex Carr. Conif. ed. II. 13 (1867) = **Juniperus**, Tourn. (Conif.).
echinoformis, Hort. ex Carr. l. c. = J. Oxycedrus.
Withmanniana, Hort. ex Carr. l. c. = J. Oxycedrus.

OXYCEROS, Lour. Fl. Cochinch. 150 (1790) = **Randia**, Houst. (Rubiac.).
chinensis, Lour. l. c. 151 = R. sinensis.
horridus, Lour. l. c. = R. horrida.
sinensis, Roem. & Schult. Syst. v. 248, in syn. = R. sinensis.

OXYCHLOË, Phil. Fl. Atac. 52. t. 6 (1860) = **Distichia**, Nees & Meyen (Juncac.).
andina, Phil. l. c. = D. macrocarpa.

OXYCLADIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 20 = **Mirbelia**, Sm. (Legumin.).
semiseptatum, F. Muell. l. c. = M. oxyclada.

OXYCLADUS, Miers, in Proc. Linn. Soc. ii. (1851) 156; et in Trans. Linn. Soc. xxi. (1853) 146. t. 18 = **Monttea**, C. Gay (Scrophul.).
aphyllus, Miers, ll. cc.—Am. austr.

OXYCOCA, Rafin. Med. Fl. ii. 48 (1830) = **Oxycoccus**, Tourn. (Vaccin.).
berberidacea, Rafin. l. c. 49 = seq.?
macrocarpa, Rafin. l. c. 48 = Oxycoccus macrocarpus.
vulgaris, Rafin. l. c. 49 = Oxycoccus palustris.

OXYCOCCOS, Hedw. f. Gen. 275 (1806) = seq.

OXYCOCCUS, Tourn. ex Adans. Fam. ii. 164 (1763). *VACCINIACEAE*, Benth. & Hook. f. ii. 575.
SCHOLLERA, Roth, Tent. Fl. Germ. i. 170 (1788).
erectus, Pursh, Fl. Am. Sept. i. 264 = Vaccinium erythrocarpum.
erythrocarpus, Pers. Syn. i. 419 = Vaccinium erythrocarpum.
europaeus, Pers. l. c. = palustris.
hispidulus, Pers. l. c. = Chiogenes serpyllifolia.
macrocarpus, *Pers. l. c.*—Reg. temp.
microcarpus, Turcz. ex Rupr. in Beitr. Pfl. Russ. Reich. iv. 56 = palustris.
palustris, *Pers. Syn.* i. 419.—Reg. bor.
quadripetala, Gilib. Fl. Lituan. i. 5 = palustris.
vulgaris, Bong. Veg. Ins. Sitcha, 152 = palustris.

OXYDENDRON, D. Dietr. Syn. Pl. ii. 1370, 1389 (1840) = seq.

OXYDENDRUM, DC. Prod. vii. 601 (1839). *ERICACEAE*, Benth. & Hook. f. ii. 585, 1239.
arboreum, *DC. l. c.*—Am. bor.

OXYDENIA, Nutt. Gen. Am. i. 76 (1818) = **Leptochloa**, Beauv. (Gramin.).
attenuata, Nutt. l. c. = L. mucronata.
filiformis, Nutt. l. c. = L. filiformis.
virgata, Nutt. l. c. = L. virgata?

OXYDIUM, Benn. Pl. Jav. Rar. 156 (1840) = **Desmodium**, Desv. (Legumin.).
oxalidifolium, Walp. Rep. i. 747 = D. Rottleri.

OXYDON, Less. in Linnaea, v. (1830) 357 = **Chaptalia**, Vent. (Compos.).
bicolor, Less. l. c. = C. runcinata.

OXYGONUM, Burch. Trav. i. 548 (1822). *POLYGONACEAE*, Benth. & Hook. f. iii. 96.
CERATOGONON, Meissn. in Wall. Pl. As. Rar. iii. 63 (1832).
OWENIA, Hilsenb. ex Meissn. in DC. Prod. xiv. 39 (1856).
Acetosella, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 60.—Afr. trop.
alatum, *Burch. Trav.* i. 548.—Afr. austr.
calcaratum, *Burch. ex Meissn. in DC. Prod.* xiv. 38.—Afr. austr.
canescens, *Sond. in Linnaea*, xxiii. (1850) 100.—Afr. austr.
dentatum, Meissn. l. c. xiv. (1840) 488, in obs. = calcaratum.
Dregeanum, *Meissn. l. c.* 487.—Afr. austr.
Dregei, Meissn. in DC. Prod. xiv. 38 = praec.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 100.—Afr. austr.

OXYGRAPHIS, Bunge, Verz. Suppl. Fl. Alt. 46 (1836). *RANUNCULACEAE*, Benth. & Hook. f. i. 6.
glacialis, *Bunge, Verz. Suppl. Fl. Alt.* 47.—Sibir. altaic.
polypetala, *Hook. f. & Thom. Fl. Ind.* i. 27.—Reg. Himal.
Shaftoana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 149.—Afghan.

OXYLAENA, Benth. in Hook. Ic. Pl. t. 1109 (1872) = **Anaglypha**, DC. (Compos.).
acicularis, Benth. l. c. = A. acicularis.

OXYLAPATHON, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = **Rumex**, Linn. (Polygon.).

OXYLEPIS, Benth. Pl. Hartw. 87 (1841) = **Helenium**, Linn. (Compos.).
lanata, Benth. l. c. = H. integrifolium.

OXYLOBIUM, Andr. Bot. Rep. t. 492 (1809). *LEGUMINOSAE*, Benth. & Hook. f. i. 467.
CALLISTACHYS, Vent. Jard. Malm. t. 115 (1803).
PODOLOBIUM, R. Br. in Ait. Hort. Kew. ed. II. iii. 9 (1811).
aciculiferum, *Benth. Fl. Austral.* ii. 25.—Austral.
acutum, *Benth. l. c.* 24.—Austral.
alpestre, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 38.—Austral.

OXYLOBIUM:—
angustifolium, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 70; et in Fl. Austral. ii. 17 = ellipticum.
arborescens, R. Br. in Ait. Hort. Kew. ed. II. iii. 9 = ellipticum.
argenteum, Kunze, in Linnaea, xx. (1847) 61 = ellipticum.
atropurpureum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 1. 250.—Austral.
Batillum, Hook. Ic. Pl. t. 612 = Gastrolobium polystachyum.
Callistachys, *Benth. Fl. Austral.* ii. 16.—Austral.
capitatum, *Benth. in Enum. Pl. Hueg.* 69.—Austral.
carinatum, *Benth. Fl. Austral.* ii. 18.—Austral.
cordifolium, *Andr. Bot. Rep.* t. 492.—Austral.
cuneatum, *Benth. in Lindl. Swan River App.* 41.—Austral.
dilatatum, Benth. l. c. 43 = cuneatum.
dilatatum, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 79, 80 = Pultenaea ternata.
Drummondii, Meissn. in Lehm. Pl. Preiss. i. 30 = cuneatum.
ellipticum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 10.—Austral.
ferrugineum, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 163, nomen (Quid?).—Afr. austr.
genistoides, Meissn. in Bot. Zeit. xiii. (1855) 12 = Chorizema humile.
hamulosum, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 379.—Austral.
heterophyllum, *Benth. Fl. Austral.* ii. 25.—Austral.
Leichhardtii, F. Muell. Fragm. iv. 20 = Isotropis filicaulis.
lineare, *Benth. Fl. Austral.* ii. 17.—Austral.
microphyllum, *Benth. l. c.* 19.—Austral.
nervosum, Meissn. in Bot. Zeit. xiii. (1855) 12 = capitatum.
obovatum, Benth. in Lindl. Swan River App. 40 = cuneatum.
obtusifolium, *Sweet, Fl. Austral.* t. 5.—Austral.
ovalifolium, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 63. t. 85 = pyramidale.
ovalifolium, Meissn. in Lehm. Pl. Preiss. i. 28 = retusum.
parviflorum, *Benth. in Lindl. Swan River App.* 40.—Austral.
procumbens, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 37.—Austral.
Pulteneae, *DC. Mém. Lég.* v. 170; *Prod.* ii. 104.—Austral.
Pulteneae, Lodd. Bot. Cab. t. 1947 = ellipticum.
reticulatum, *Meissn. in Lehm. Pl. Preiss.* i. 29.—Austral.
retusum, *R. Br. in Bot. Reg.* t. 913.—Austral.
scandens, *Benth. in Ann. Wien. Mus.* ii. (1838) 70.—Austral.
sericeum, Benth. l. c. = Brachysema undulatum.
spathulatum, *Benth. Fl. Austral.* ii. 18.—Austral.
spectabile, *Endl. Nov. Stirp. Dec.* 2.—Austral.
spinosum, DC. Prod. ii. 104 = Pultenaea ternata.
staurophyllum, *Benth. Fl. Austral.* ii. 26.—Austral.
tricuspidatum, *Meissn. in Lehm. Pl. Preiss.* i. 30.—Austral.
trilobatum, *Benth. Fl. Austral.* ii. 25.—Austral.
virgatum, *Hort. Kew. ex Benth. l. c.* 22.—Austral.

OXYLOBUS, [Moç.] ex DC. Prod. v. 115 (1836) = **Ageratum**, Linn. (Compos.).
trinervius, [Moç.] ex DC. Prod. v. 115 (= *Phania trinervia*).—Mexic.

OXYMERIS, DC. Prod. iii. 190 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 761.
CHRYSOPHORA, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94, 95, 147.
CLIDEMIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 87.
TSCHUDYA, DC. Prod. iii. 168 (1828).
acutiflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 93.—Bras.
agrestis, *Triana, l. c.* 95.—Guian.
alpestris, *Triana, l. c.* 92.—Bras.
angustifolia, *Triana, l. c.*—Bras.
anisophylla, *Triana, l. c.* 96.—Guian.
aptera, *Triana, l. c.* 91.—Bras.
aristigera, *Triana, l. c.* 94.—Peruv.
asperifolia, *Triana, l. c.* 96.—Ind. occ.

OXYMERIS:—
asperiuscula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 95.—Guiana.
aurea, *Triana, l. c.* 91.—Bras.
australis, *Triana, l. c.* 90.—Bras.
barbinervis, *Cham. ex Triana, l. c.* 93.—Bras.
Brackenridgei, *Triana, l. c.* 92.—Bras.
calvescens, *Triana, l. c.* 91.—Bras.
calyptrata, Cham. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 179 = Miconia Chamissois.
carassana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 91.—Bras.
chaetocalyx, *Triana, l. c.* 92.—Bras.
chaetodon, *Triana, l. c.* 94.—Bras.
ciliata, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 38.—Bras.
cinerea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 92.—Bras.
cinnamomea, *Triana, l. c.* 94.—Panama.
congestiflora, *Triana, l. c.* 92.—Bras.
cordifolia, *Triana, l. c.* 90.—Bras.
cordigera, *Triana, l. c.* 94.—Bras.
cornioides, *Triana, l. c.* 93.—Mexic.
crenata, *Triana, l. c.* 91.—Peruv.; Bras.
cuneata, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 38.—Bras.
cuspidata, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 109 = Miconia cuspidata.
cuspidata, *Triana, l. c.* 93.—Bras.
debilis, *Triana, l. c.* 94.—Bras.
deflexa, *Triana, l. c.*—Bras.
dendroides, *Triana, l. c.*—Bras.
dichotoma, *Triana, l. c.* 95.—Peruv.
dispar, *Triana, l. c.* 91.—Bras.
dolichantha, *Triana, l. c.* 90.—Bras.
dolichondris, *Cham. ex Triana, l. c.* 93, nomen.—Bras.
dolichostachya, *Triana, l. c.* 94.—Bras.
dubia, *Triana, l. c.* 92.—Bras.
erostrata, *Triana, l. c.*—Bras.
fallax, *Triana, l. c.* 91.—Bras.
floribunda, *Triana, l. c.*—Bras.
foveolata, *Triana, l. c.* 90.—Bras.
furfuracea, Poepp. ex Triana, l. c. 107 = Miconia Poeppigii.
glandulifera, *Triana, l. c.* 95.—Bras.
heterobasis, *Triana, l. c.*—Mexic.
hirta, *Triana, l. c.* 90.—Bras.
inaequalifolia, *Triana, l. c.* 96.—Bras.
inopogon, *Triana, l. c.* 91.—Bras.
intermedia, *Triana, l. c.* 95.—Bras.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxxii. (1882) 37.—Bras.
laevigata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 94.—Bras.
Lindeniana, *Triana, l. c.* 90.—Venezuela.
longibarbis, *Triana, l. c.*—Bras.
macrophylla, Benth. Bot. Voy. Sulph. 95 = Octopleura macrophylla.
macropora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 94.—Bras.
megalophylla, *Wawra, in Oestr. Bot. Zeitschr.* xxxii. (1882) 37.—Bras.
melanodesma, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 92.—Mexic.
Miconiastrum, *Triana, l. c.* 91.—Bras.
micropetala, *Triana, l. c.* 96.—Bras.
multiplinervis, *Triana, l. c.* 94.—Mexic.
nervosa, *Triana, l. c.* 92.—Peruv.
Nianga, *Triana, l. c.* 90.—Bras.
oocarpa, *Triana, l. c.* 96.—Bras.
ossaeoides, *Triana, l. c.* 93.—Venezuela.
paraguayensis, *Triana, l. c.* 96.—Paraguay.
penduliflora, *Triana, l. c.* 93.—Bras.
pennipilis, *Triana, l. c.* 92.—Bras.
Pickeringii, *Triana, l. c.* 93.—Bras.
polystachya, *Triana, l. c.* 92.—Bras.
pubescens, *Triana, l. c.* 95.—Bras.
pulchra, *Cham. ex Triana, l. c.* 93.—Bras.
pulverulenta, *Triana, l. c.* 96.—Guian.
purpurascens, *Triana, l. c.* 91.—Bras.
quinquedentata, *DC. Prod.* iii. 190.—Bras.
quinquenodis, *DC. l. c.*—Bras.
Regnellii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 91.—Bras.
reversa, *Triana, l. c.* 95.—Bras.
rhodopogon, *Triana, l. c.* 90.—Bras.
ribesiaeflora, *Triana, l. c.* 92.—Bras.
Riedeliana, *Triana, l. c.* 92.—Bras.

OXYMERIS:—
rufescens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 95.—Bras.
secunda, *Triana, l. c.* 95.—Costa Rica; Peruv.
secundiflora, *Triana, l. c.* 95.—Bras.
sessilifolia, *Triana, l. c.* 93.—N. Granat.
simplicicaulis, *Triana, l. c.* 92.—Bras.
strigilliflora, *Triana, l. c.*—Bras.
subseriata, *Triana, l. c.*—Mexic.
sylvestris, *Triana, l. c.*—Bras.
tetraquetra, *Triana, l. c.* 91.—Bras.
urophylla, *Triana, l. c.* 94.—Bras.
velutina, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 347.—Bras.
Widgrenii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 94.—Bras.
xanthocoma, *Triana, l. c.* 91.—Bras.
xantholasia, *Triana, l. c.* 90.—Bras.
xanthopogon, *Triana, l. c.* 91.—Bras.

OXYMITUS, Presl, (Bot. Bemerk. 92) (1844); in Abh. Boehm. Ges. V. iii. (1845) 522 = **Argylia**, D. Don (Bignon.).
argylioides, Presl, ll. cc.—Chili.

OXYMYRRHINE, Schau. in Linnaea, xvii. (1843) 240 = **Baeckea**, Linn. (Myrtac.).
gracilis, Schau. l. c. = B. polyandra.

OXYMITRA, Hook. f. & Thoms. Fl. Ind. i. 145 (1855). *ANONACEAE*, Benth. & Hook. f. i. 26.
affinis, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 70.—Malacca.
bassiaefolia, *Teijsm. & Binn. in Tijdschr. Nederl.* Indie, xxv. (1863) 419.—Malaya.
biglandulosa, *Scheff. l. c.* xxxi. (1870) 341.—Malaya.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 30.—Borneo.
caesia, *Miq. l. c.* 31.—Sumatra.
cuneiformis, *Zoll. in Linnaea*, xxix. (1857-58) 324; *Miq. Fl. Ind. Bat.* i. II. 50.—Java.
diadena, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 31.—Borneo.
excisa, *Miq. l. c.* 32.—Sumatra.
filipes, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 71.—Malacca.
fornicata, *Hook. f. & Thoms. Fl. Ind.* i. 146.—Reg. Himal.; Burma.
Gabriaciana, *Baill. Adansonia*, x. (1871) 106.—Cochinch.
Gardneri, Baill. Hist. Pl. i. 236 = Goniothalamus Gardneri.
glauca, *Hook. f. & Thoms. Fl. Ind.* i. 146.—Ins. Philipp.
gracilipes, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 471.—Afr. trop.
Grayana, Baill. Hist. Pl. i. 237 = Richella monosperma.
hamata, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 471.—Afr. trop.
hirta, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 30.—Ins. Celebes.
Korthalsiana, *Miq. l. c.* 29.—Borneo.
latifolia, *Hook. f. & Thoms. Fl. Ind.* i. 145.—Malaya.
Maclellandii, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 70.—Burma.
macrophylla, A. DC. Mém. Anon. 42 = Goniothalamus sesquipedalis.
Motleyana, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 155.—Borneo.
myristicifolia, *Oliver, Fl. Trop. Afr.* i. 33.—Afr. trop.
obtusata, *Baill. Adansonia*, viii. (1867-68) 178.—N. Caled.
oxyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 29.—Borneo.
patens, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 472.—Afr. trop.
platypetala, *Benth. l. c.*—Afr. trop.
stenopetala, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 71.—Burma.
unonaefolia, *Hook. f. & Thoms. Fl. Ind.* i. 146.—Burma.

OXYNEPETA, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxi. (1873) I. (Lab. Pers.) 58 = **Nepeta**, Linn. (Labiat.).
heliotropifolia, Bunge, l. c. 59 = N. heliotropifolia.
involucrata, Bunge, l. c. = N. cryptantha.
parviflora, Bunge, l. c. = N. ucranica.

OXYNIX, Noronha, in Verh. Batav. Gen. v. (1790) ed. I Art. IV. 3 = **Averrhoa**, Linn. (Geraniac.).

OXYODON, DC. Prod. vii. 43 (1838) lapsu = OXYDON, Less. = **Chaptalia**, Vent. (Compos.).

OXYPAPPUS, Benth. Bot. Voy. Sulph. 118. t. 42 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
scabra, *Benth. Bot. Voy. Sulph.* 118.—Mexic.

OXYPETALUM, R. Br. in Mem. Wern. Soc. i. (1809) 41. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 752.
GOTHOFREDA, Vent. Choix, t. 60 (1803).
PACHYGLOSSUM, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 345, in obs.
SCHIZOSTEMMA, Decne. l. c. (1838) 344.
TWEEDIA, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 291.
aequaliflorum, *Fourn. in Mart. Fl. Bras.* vi. IV. 283.—Bras.
albiflorum, *Decne. in DC. Prod.* viii. 587.—Peruv.
ampliflorum, *Fourn. in Mart. Fl. Bras.* vi. IV. 276.—Bras.
appendiculatum, *Mart. Nov. Gen. et Sp.* i. 48. t. 30.—Bras.
arachnoideum, *Fourn. in Mart. Fl. Bras.* vi. IV. 272.—Bras.
Arnottianum, *Buek, ex Fourn. l. c.* 278.—Bras.
Banksii, *Schult. Syst.* vi. 91.—Bras.
Berterianum, *Spreng. Syst.* i. 854.—N. Granat.
Berterii, Spreng. ex Decne. in DC. Prod. viii. 588 = Metastelma Berterianum.
brevipes, *Phil. in Linnaea*, xxxiii. (1864-65) 177.—Chili.
Brunonis, *Decne. in DC. Prod.* viii. 587.—Reg. Argent.
campestre, *Decne. l. c.* 585.—Bras.
capitatum, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 288 = Arnottianum.
capitatum, *Mart. Nov. Gen. et Sp.* i. 50.—Bras.
coalitum, *Fourn. in Mart. Fl. Bras.* vi. IV. 282.—Bras.
coccineum, *Griseb. in Goett. Abh.* xix. (1874) 206.—Reg. Argent.
coeruleum, *Decne. in DC. Prod.* viii. 585.—Reg. Argent.
confertiflorum, *Decne. l. c.* 588.—Chili.
coriaceum, *Decne. l. c.* 585.—Bras.
corymbiferum, *Fourn. in Mart. Fl. Bras.* vi. IV. 267.—Bras.
crispum, *Wight, ex Hook. Journ. Bot.* i. (1834) 288.—Bras.
deltoideum, *Fourn. in Mart. Fl. Bras.* vi. IV. 269.—Bras.
densiflorum, *Decne. in DC. Prod.* viii. 584.—Bras.
dentatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 263.—Bras.
Dombeyanum, *Decne. in DC. Prod.* viii. 583.—Bras.
Echegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 54.—Reg. Argent.
erectum, *Mart. Nov. Gen. et Sp.* i. 50.—Bras.
erianthum, *Decne. in DC. Prod.* viii. 584.—Bras.
erostre, *Fourn. in Mart. Fl. Bras.* vi. IV. 265.—Bras.
floribundum, *Decne. in DC. Prod.* viii. 586.—Bras.
foliosum, *Mart. Nov. Gen. et Sp.* i. 50.—Bras.
glomeratum, *Fourn. in Mart. Fl. Bras.* vi. IV. 281.—Bras.
Gothofreda, *Schult. Syst.* vi. 92.—N. Granat.
grandiflorum, *Fourn. in Mart. Fl. Bras.* vi. IV. 272.—Bras.
Guilleminianum, *Decne. in DC. Prod.* viii. 582.—Bras.
Hilariana, *Fourn. l. c.* 283.—Bras.
Hookeri, *Decne. in DC. Prod.* viii. 587.—Chili.
incanum, *Fourn. in Mart. Fl. Bras.* vi. IV. 276.—Bras.
integrifolium, *Fourn. l. c.* 279.—Bras.
Jacobinae, *Decne. in DC. Prod.* viii. 584.—Bras.
lagoënse, *Fourn. in Mart. Fl. Bras.* vi. IV. 264.—Bras.
lanatum, *Decne. ex Fourn. l. c.* 266.—Bras.
ligulatum, *Fourn. l. c.* 258.—Bras.
Lindenianum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 318.—N. Granat.
lineare, *Decne. in DC. Prod.* viii. 587.—Bras.
Luschnathii, *Fourn. in Mart. Fl. Bras.* vi. IV. 261.—Bras.
lutescens, *Fourn. l. c.* 273.—Bras.
macrolepis, *Decne. in DC. Prod.* viii. 585.—Bras.
maritimum, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 288.—Bras.

OXYPETALUM:—
Martii, *Fourn. in Mart. Fl. Bras.* vi. IV. 280.—Bras.
megapotamicum, *Spreng. Syst.* iv. *Cur. Post.* 111.—Bras.
microphyllum, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 289.—Reg. Argent.
microstemma, *Fourn. in Mart. Fl. Bras.* vi. IV. 282.—Bras.
Minarum, *Fourn. l. c.* 258.—Bras.
molle, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 289.—Peruv.
monanthum, *Decne. in DC. Prod.* viii. 584.—Bras.
montanum, *Mart. Nov. Gen. et Sp.* iii. 49.—Bras.
mucronatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 271.—Bras.
multicaule, *Fourn. l. c.* 284.—Bras.
muticum, *Fourn. l. c.* 277.—Bras.
nigrescens, *Fourn. l. c.* 264.—Bras.
niveum, *Griseb. in Goett. Abh.* xix. (1874) 206.—Reg. Argent.
pachyglossum, *Decne. in DC. Prod.* viii. 585.—Bras.
pachygynum, *Decne. l. c.* 583.—Bras.
pallidum, *Fourn. in Mart. Fl. Bras.* vi. IV. 259.—Bras.
paludosum, *Decne. in DC. Prod.* viii. 585.—Bras.
pannosum, *Decne. l. c.* 583.—Bras.
pardense, *Fourn. in Mart. Fl. Bras.* vi. IV. 277.—Bras.
parviflorum, *Decne. in DC. Prod.* viii. 587.—Bras.
parvifolium, *Fourn. in Mart. Fl. Bras.* vi. IV. 272.—Bras.
patulum, *Fourn. l. c.* 278.—Bras.
pauperculum, *Fourn. l. c.* 265.—Bras.
pedicellatum, *Decne. in DC. Prod.* viii. 582.—Bras.
pilosum, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 539.—Bras.
pratense, *Griseb. in Goett. Abh.* xxiv. (1879) 231.—Reg. Argent.
proboscideum, *Fourn. in Mart. Fl. Bras.* vi. IV. 281.—Bras.
propinquum, *Decne. in DC. Prod.* viii. 582.—Bras.
pulchrum, *Decne. l. c.* 584.—Bras.
riparium, *H. B. & K. Nov. Gen. et Sp.* iii. 197.—N. Granat.
saxatile, *Decne. in DC. Prod.* viii. 588.—Chili.
Schottii, *Fourn. in Mart. Fl. Bras.* vi. IV. 273.—Bras.
siliculae, *Fourn. l. c.* 66.—Bras.
solanoides, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 289.—Reg. Argent.
squamulatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 270.—Bras.
stigmatosum, *Fourn. l. c.* 280.—Bras.
strictum, *Mart. Nov. Gen. et Sp.* i. 50.—Bras.
suaveolens, *Fourn. in Mart. Fl. Bras.* vi. IV. 264.—Bras.
subhirtellum, *Miq. in Linnaea*, xxii. (1849) 568.—Bras.
tenuiflorum, *Griseb. in Goett. Abh.* xxiv. (1879) 230.—Reg. Argent.
ternifolium, *Fourn. in Mart. Fl. Bras.* vi. IV. 279.—Bras.
tomentosum, *Wight, ex Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 288.—Bras.
umbellatum, *Gardn. in Hook. Lond. Journ.* i. (1842) 179.—Bras.
variegatum, *Griseb. in Goett. Abh.* xxiv. (1879) 230.—Reg. Argent.
villosum, *Fourn. in Mart. Fl. Bras.* vi. IV. 262.—Bras.
Wightianum, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 288.—Bras.

OXYPHAERIA, Steud. Nom. ed. I. 138 (1821), sphalm. = seq.

OXYPHERIA, DC. Prod. vi. 157 (1837) = OXIPHOERIA, Hort. Angl. = **Humea**, Sm. (Compos.).

OXYPHYLLUM, Phil. Fl. Atac. 28. t. 4 (1860). *COMPOSITAE*, Benth. & Hook. f. ii. 499.
ulicinum, *Phil. l. c.*—Chili.

OXYPOGON, Rafin. in Journ. Phys. lxxxix. (1819) 98 = **Lathyrus**, Linn. (Legumin.).
elegans, Rafin. l. c. = L. venosus.
elegans, Rafin. ex DC. Prod. ii. 355 = Vicia dumetorum.

OXYPOLIS, Rafin. Neogenyt. 2 (1825) = **Peucedanum**, Tourn. (Umbell.).
caroliniana, Rafin. in Ser. Bull. Bot. i. (1830) 214 = P. teretifolium.
denticulata, Rafin. Neogenyt. 2 = P. ternatum.
rigida, Rafin. l. c. = P. ternatum.
teretifolia, Rafin. l. c. = P. teretifolium.
tricuspidata, Rafin. l. c. = P. ternatum.
triquinata, Rafin. l. c. = Archangelica hirsuta.

OXYRAMPHIS, Wall. Cat. n. 5348 (1831) = **Lespedeza**, Michx. (Legumin.).
macrostyla, Lindl. Bot. Reg. (1846) t. 28 = L. eriocarpa.
macrostyla, Wall. Cat. n. 5348 = L. sericea.
sericea, R. Grah. in Wall. l. c. n. 5349 = L. sericea.
stenocarpa, Klotzsch, in Bot. Ergeb. Waldem. Reise, 159 = L. sericea.
virgata, R. Grah. in Wall. Cat. n. 5350 = L. eriocarpa.

OXYRHAMPHIS, Reichb. Nom. 147 (1841) = praec.

OXYRIA, Hill, Veg. Syst. x. 24. t. 24 (1765). *POLYGONACEAE*, Benth. & Hook. f. iii. 100.
DONIA, R. Br. in Ross, Voy. ed. I. p. xli (1819).
OXYLAPATHON, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159.
acida, R. Br. ex Hook. & Arn. Brit. Fl. ed. VI. 360 = digyna.
digyna, *Hill, Hort. Kew.* 158.—Reg. bor. alp. et arct.
elatior, R. Br. in Wall. Cat. n. 1726 = digyna.
reniformis, Hook. Fl. Scot. 111 = digyna.
rotundifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 277 = digyna.

OXYS, Tourn. ex Adans. Fam. ii. 388 (1763) = **Oxalis**, Linn. (Geran.).
Acetosella, Scop. Fl. Carn. ed. II. i. 326 = Oxalis Acetosella.
alba, Lam. Fl. Fr. iii. 60; Gilib. Fl. Lituan. i. 62 = Oxalis Acetosella.
corniculata, Scop. Fl. Carn. ed. II. i. 326 = Oxalis corniculata.
incarnata, Medic. Malv. 113; Moench, Meth. 47 = Oxalis incarnata.
lutea, Lam. Fl. Fr. iii. 60 = Oxalis corniculata.
lutea, Moench, Meth. 48 = Oxalis corniculata.
stricta, All. Fl. Pedem. ii. 89; Medic. Malv. 113 = Oxalis corniculata.
vulgaris, Rupr. Fl. Caucas. 264 = Oxalis Acetosella.

OXYSEPALUM, Wight, Ic. v. 17. t. 1736 (1852) (Oxysepala) = **Bulbophyllum**, Thou. (Orchid.).
ovalifolium, Wight, l. c. = B. clandestinum.

OXYSPERMUM, Eckl. & Zeyh. Enum. 364 (1837) = **Galopina**, Thunb. (Rubiac.).
asperum, Eckl. & Zeyh. l. c. 365 = G. Oxyspermum.

OXYSPORA, DC. Prod. iii. 123 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 751.
ALLOZYGIA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 309. t. 15.
HOMOCENTRIA, Naud. l. c. (1851) 308.
HYLOCHARIS, Miq. Fl. Ind. Bat. Suppl. 319 (1860).
cernua, *Hook. f. & Thoms. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 73.—Reg. Himal.
macrophylla, *Triana, l. c.*—Sumatra.
paniculata, *DC. Prod.* iii. 123.—Reg. Himal.
pauciflora, *Benth. Fl. Hongk.* 116.—Ins. Hongkong.
vagans, Hook. Bot. Mag. t. 4553 = paniculata.
vagans, *Wall. Pl. Rar. As.* i. 78.—Reg. Himal.

OXYSTELMA, R. Br. in Mem. Wern. Soc. i. (1809) 40. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 749.
Alpini, Decne. in DC. Prod. viii. 543 = esculentum.
bifidum, Llanos, Fragm. 64 = Tylophora bifida.
bornouense, *R. Br. in Denh. & Clapp. Trav. App.* 239.—Afr. trop.
carnosum, R. Br. Prod. 462 = Vincetoxicum carnosum.
caudatum, Buch.-Ham. ex Wall. Cat. n. 8213 D = Pentatropis spiralis.
esculentum, *R. Br. in Mem. Wern. Soc.* i. (1809) 40.—Austral.
esculentum, Wall. ex Decne. in DC. Prod. viii. 543 = Wallichii.

OXYSTELMA:—
Hooperianum, *Blume, Bijdr.* 1053.—Java.
pulchellum, D. Dietr. Syn. Pl. ii. 902 = Raphistemma pulchellum.
senegalense, Decne. in DC. Prod. viii. 543 = bornouense.
Wallichii, Wight, Contrib. 54 = esculentum.
Zippelianum, Blume, Bijdr. 1053 = esculentum.

OXYSTEMON, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 314 = **Clusia**, Linn. (Guttif.).
nervosus, Planch. & Triana, l. c. xiv. (1860) 226.—N. Granat.

OXYSTOPHYLLUM, Blume, Bijdr. 335. t. 38 (1825) = **Dendrobium**, Sw. (Orchid.).
atropurpureum, Blume, Rumphia, iv. 41. tt. 193. f. 4; 198 f. C = D. atropurpureum.
carnosum, Blume, Bijdr. 336 = D. atropurpureum.
excavatum, Blume, l. c. = D. excavatum.
macrostoma, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 122 = D. Serra.
rigidum, Blume, Bijdr. 336 = D. rigens.

OXYSTYLIS, Torr. & Frém. in Rep. Frém. App. 312 (1845). *CAPPARIDEAE*, Benth. & Hook. f. i. 107.
lutea, *Torr. & Frém. l. c.* 313.—Am. bor. occ.

OXYTANDRUM, Neck. Elem. ii. 255 (1790) = **Apeiba**, Aubl. (Tiliac.).

OXYTENANTHERA, Munro, in Trans. Linn. Soc. xxvi. (1868) 126. *GRAMINEAE*, Benth. & Hook. f. iii. 1211.
abyssinica, *Munro, l. c.* 127.—Afr. trop.
albo-ciliata, *Munro, l. c.* 129.—Ind. or.
monostigma, *Bedd. Forest. Man. Bot.* 233.—Ind. or.
nigrociliata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 128.—Ind. or.
Stocksii, *Munro, l. c.* 130.—Ind. or.
Thwaitesii, *Munro, l. c.* 129.—Zeylan.

OXYTENIA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 172. *COMPOSITAE*, Benth. & Hook. f. ii. 353.
acerosa, *Nutt. l. c.*—Calif.

OXYTHECA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 169. *POLYGONACEAE*, Benth. & Hook. f. iii. 92.
BRISEGNOA, Remy, in C. Gay, Fl. Chil. v. 291 (1849).
TETRARHAPHIS, Miers, in Trans. Linn. Soc. xxi. (1853) 141. t. 17.
apiculata, S. Wats. in Bot. Calif. ii. 82, in syn., sphalm. = *spiculata*.
caryophylloides, *Parry, in Proc. Davenp. Acad. Nat. Sc.* i. (1876) 175.—Calif.
dendroidea, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 169.—Am. bor. occ.
foliosa, *Nutt. l. c.*—Am. bor.
glandulosa, *Nutt. l. c.* 170.—Calif.
inermis, *S. Wats. in Proc. Am. Acad.* xii. (1877) 273.—Calif.
Parishii, *Parry, in Proc. Davenp. Acad. Nat. Sc.* i. (1876) 176.—Calif.
perfoliata, *A. Gray, ex S. Wats. Bot. King's Exped.* 311. t. 34.—Am. bor. occ.
Reddingiana, *Jones, in Bull. Torrey Bot. Club*, ix. (1882) 32.—Am. bor.
spiculata, Miers, in Proc. Linn. Soc. ii. (1851) 155; et Trans. Linn. Soc. xxi. (1855) 144. t. 17 = dendroidea.
trilobata, *A. Gray, in Proc. Am. Acad.* xii. (1877) 83.—Calif.
Watsoni, *Torr. & Gray, ex S. Wats. Bot. King's Exp.* 311. t. 34. f. 5–7.—Am. bor. occ.

OXYTHECE, Miq. in Mart. Fl. Bras. vii. 105 (1863) = **Lucuma**, Juss. (Sapot.).
leptocarpa, Miq. l. c. 106 = L. leptocarpa.
Pseudo-sideroxylon, Miq. l. c. 106.—Bras.

OXYTRIA, Rafin. Fl. Tellur. ii. 26 (1836) = **Schoenolirion**, Torr. (Liliac.).
crocea, Rafin. l. c. = S. croceum.

OXYTROPIS, DC. Astrag. 24, 66. tt. 2–6, 8 (1802). *LEGUMINOSAE*, Benth. & Hook. f. i. 507.
aciphylla, *Ledeb. Fl. Alt.* iii. 279.—Sibir. altaic.
adproximata, Less. in Linnaea, ix. (1834) 154 = campestris.
aequipetala, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 11.—Turkestan.
ajanensis, *Bunge, l. c.* 95.—Sibir. or.
albana, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 54.—Reg. Caucas.
albiflora, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 96.—Sibir.
algida, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 9.—As. bor.
alpicola, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 86 = Tschujae.
alpicola, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 743 = uralensis.
alpina, Bunge, Ind. Sem. Hort. Dorpat. (1840) 8 = uralensis.
altaica, Pers. Syn. ii. 333 = brevirostra.
ambigua, *DC. Astrag.* 70.—Sibir.
ammophila, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 66.—Sibir.
amoena, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 327 = lapponica.
ampullacea, Turcz. l. c. 744 = mixotriche.
ampullata, Less. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 133 = polyphylla.
ampullata, *Pers. Syn.* ii. 333.—Sibir.
ampullata, Turcz. ex Bunge, Verz. Alt. Pfl. 89 = mixotriche.
annularis, DC. Astrag. 93 = Astragalus annularis.
approximata, Less. in Linnaea, ix. (1834) 175 = campestris.
arctica, R. Br. in Parry, 1st Voy. App. 278 = uralensis.
arctica, Trautv. Fl. Taimyr. 49 = sordida.
arctobia, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1871) I. 114.—Am. bor.
argaea, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 505 = albana.
argentata, Ledeb. Fl. Ross. i. 592 = recognita.
argentata, *Pers. Syn.* ii. 331.—Desert. Kirgis.
argentata, Pursh, Fl. Am. Sept. ii. 473 = nana?
argentata, Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 77 = ajanensis.
argyraea, DC. Prod. ii. 276 = argentata.
argyrophylla, Ledeb. Fl. Ross. Ill. t. 54 = argentata.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 41.—Persia.
baicalensis, *Pall. ex Bess. in Flora*, xvii. (1834) I. *Beibl.* 10.—Sibir. baical.
baicalia, *Pers. Syn.* ii. 333.—Sibir. baical.
bicolor, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 91.—China.
borealis, DC. Prod. ii. 275 = campestris.
borealis, Trautv. ex C. A. Mey. Fl. Ochot. 26 = Trautvetteri.
brachybotrys, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 53.—Sibir.
bracteosa, Bunge, l. c. 143, in textu = oxyphylla?
brevicaulis, *Ledeb. Fl. Alt.* iii. 284.—Sibir. altaic.
brevifolia, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 752 = dubia.
brevirostra, *DC. Astrag.* 82. t. 6.—Sibir.
cabulica, *Boiss. Fl. Orient.* ii. 507.—Afghan.
cachemiriana, *Cambess. in Jacquem. Voy. Bot.* 38. t. 44.—Reg. Himal.
caerulea, *DC. Astrag.* 68.—Sibir.
caespitosa, *Pers. Syn.* ii. 333.—Dahuria.
campestris, *DC. Astrag.* 74.—Reg. bor. temp. et arct.
campestris, Hook. in Lond. Journ. Bot. vi. (1847) 212 = Lambertii.
cana, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 3.—As. bor.
candicans, *DC. Astrag.* 72.—Sibir.
Capusii, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 263.—Turkestan.
carinthiaca, Fischer-Oost. in Flora, xxxvii. (1854) 99 = lapponica.
carpatica, Uechtr. in Oestr. Bot. Zeitschr. xiv. (1864) 218 = montana.
caucasica, *Regel, Ind. Sem. Hort. Petrop.* (1865) 40.—Reg. Caucas.
caudata, *DC. Astrag.* 73.—Sibir. altaic.
chiliophylla, Royle, Illustr. Bot. Himal. 198 = microphylla.
chinensis, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 35 = manshurica.

OXYTROPIS:—
chionobia, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 148.—Turkestan.
chionophylla, *Schrenk, Enum. Pl. Nov.* 77.—Sibir.
chrysocarpa, *Boiss. Diagn.* Ser. I. vi. 34.—Persia.
chrysotricha, *Franch. Pl. Davidian.* i. (1884) 90.—China.
ciliata, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 186.—Mongolia.
cinerascens, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 26.—Reg. Himal.
citricans, Thomas, ex Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 63 = *hybrida*.
coerulea, Auct. vide caerulea.
coerulea, Turcz. Fl. Baic.-Dahur. i. 304 = filiformis.
collina, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 741 = grandiflora.
confusa, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 227, in textu.—Mont. Altai.
curviflora, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 10 = caerulea.
cuspidata, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 70.—Soongaria.
cyanea, *Bieb. Fl. Taur. Cauc.* iii. 502.—Reg. Caucas.
cyanea, Gaud. Fl. Helvet. iv. 536 = Gaudini.
cymbicarpos, *Pers. Syn.* ii. 333.—Lusitan.
dasypoda, *Rupr. ex Boiss. Fl. Orient.* ii. 506.—Reg. Caucas.
Davidi, *Franch. Pl. Davidian.* i. (1884) 89.—China.
dealbata, Pers. Syn. ii. 334 = Astragalus albicaulis.
deflexa, *DC. Astrag.* 96.—Reg. bor. temp. et arct.
densa, *Benth. ex Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 24.—Reg. Himal.
diantha, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 470.—Mongolia.
dichoptera, DC. Astrag. 97 = triphylla.
dichroantha, *Schrenk, Enum. Pl. Nov.* 78.—Sibir.
didymophysa, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 61.—Turkestan.
diffusa, Ledeb. Fl. Alt. iii. 281 = glabra.
dioritica, *Boiss. Diagn.* Ser. II. v. 84.—As. Min.
dissitiflora, *Ledeb. Fl. Ross.* i. 597.—Sibir. or.
Drakeana, *Franch. Pl. Davidian.* i. (1884) 89.—China.
dubia, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 752.—Sibir. baical.
elongata, Turcz. in Bull. Soc. Nat. Mosc. (1840) 67 = candicans.
eriocarpa, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 122.—Mont. Altai.
falcata, *Bunge, l. c.* 156.—China.
Fetisowi, *Bunge, in Act. Hort. Petrop.* vii. (1880) 367.—Turkestan.
filiformis, *DC. Astrag.* 80. t. 4.—Dahuria.
Fischeri, DC. Prod. ii. 281 = vaginata.
floribunda, Benth. in Royle, Illustr. Bot. Himal. 198 = mollis.
floribunda, *DC. Astrag.* 94.—Turkestan.
foetida, *DC. l. c.* 75.—Europ.
foliolosa, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 101 = macrosema.
foliolosa, Hook. Fl. Bor. Am. i. 146 = deflexa.
follicularis, Pers. ex DC. Prod. ii. 301 = Astragalus follicularis.
frigida, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xiv. (1841) 402.—Soongaria.
fruticulosa, *Bunge, l. c.* xxxix. (1866) II. 7.—As. bor.
Gaudini, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 226.—Helvet.
Gebleri, *Fisch. ex Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 89.—Mont. Altai.
Gebleriana, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 307 = mugodsharica.
gelida, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 29, in textu = albana.
generosa ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 66.—Europ.
glabra, *DC. Astrag.* 95. t. 8.—Reg. Himal.; Sibir.
glacialis, Benth. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 18 = lapponica.
glacialis, Strach. & Winterb. ex Bunge, l. c. 26 = cinerascens.
glandulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 754.—Sibir.
glauca, *Sweet, Hort. Brit.* ed. I. 123.—Tauria.
globifera, Franch. in Ann. Sc. Nat. Sér. VI. xv. (1883) 262 = seq.
globiflora, *Bunge, in Rupr. Sert. Tiansch.* 43.—Turkestan.

OXYTROPIS:—
Gmelini, Fisch. Hort. Gorenk. ed. II (1812) 71, nomen = longirostra.
gracillima, *Bunge, in Linnaea*, xvii. (1843) 5.—Mongol.
grandiflora, *DC. Astrag.* 71.—Sibir.
Griffithii, *Bunge, ex Boiss. Fl. Orient.* ii. 502.—Afghan.
gymnogyne, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 25.—Turkestan.
Halleri, Bal. ex Boiss. Fl. Orient. ii. 500 = lazica.
Halleri, Bunge, Verz. Alt. Suppl. 2 = uralensis.
Hallii, *Bunge, in Mém. Acad. Sc. Pétersb.* Sér. VII. xxii. (1874) I. 162.—Am. bor.
hedysaroides, Pers. Syn. ii. 334 = Hedysarum obscurum.
heratensis, *Bunge, ex Boiss. Fl. Orient.* ii. 507.—Persia.
heterophylla, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 470.—Mongolia.
heteropoda, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 4.—As. bor.
heterotricha, *Turcz. l. c.* xv. (1842) 752.—Sibir. baical.
hirsuta, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 55.—Soongaria.
hirta, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 91.—China.
Hookeriana, Nutt. ex Torr. & Gray, Fl. N. Am. i. 340 = Lambertii.
humifusa, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 535 = albana.
hybrida ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 63.—Europ.
hypsophila, *Bunge, ex Boiss. Fl. Orient.* ii. 501.—Persia.
Hystrix, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 254.—Soongaria.
inaria, *DC. Prod.* ii. 279.—Sibir. altaic.
inaria, Ledeb. Fl. Alt. iii. 273 = pumila.
integripetala, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 69.—Turkestan.
intermedia, *Bunge, Del. Sem. Hort. Dorpat.* (1839) 8.—*Cf. in Linnaea*, xiv. (1840) *Litt.* 120.—Sibir. altaic.
Jacquinii, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 226 = montana.
kansuensis, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 38.—China.
Kasbecki, *Bunge, ex Boiss. Fl. Orient.* ii. 506.—Reg. Caucas.
kashemiriana, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 43 = cachemiriana.
Kotschyana, *Boiss. & Hohen. Diagn.* Ser. I. ix. 36.—Persia.
Lagopus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 17.—Am. bor.
Lambertii, *Pursh, Fl. Am. Sept.* ii. 740.—Am. bor.
lanata, *DC. Astrag.* 89.—Sibir.
lanceolata, Benth. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 72, in textu = mollis.
lancifolia, Vis. ex Bunge, l. c. 107 = campestris?
lapponica, *Gaud. Fl. Helvet.* iv. 543.—Europ.; As. bor.
lasiopoda, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 151.—Mongolia.
lazica, *Boiss. Fl. Orient.* ii. 499.—As. Min.
Lehmanni, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 225.—Turkestan.
leptophylla, *DC. Astrag.* 77.—Sibir.
leptophysa, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 61.—Turkestan.
leucantha, *Pers. Syn.* ii. 331.—Sibir.
leucocyanea, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 22.—Turkestan.
leucopodia, C. A. Mey. in Ledeb. Fl. Alt. iii. 279 = squammulosa.
leucotricha, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 745.—Sibir.
longibracteata, *Kar. & Kir. l. c.* (1841) 403.—Sibir.
longicuspis, Ledeb. Ind. Sem. Dorpat.; ex ej. Fl. Alt. iii. 283 = vaginata.
longipes, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 227 = revoluta.
longipes, *Fisch. ex Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 118.—Sibir.
longirostra, *DC. Astrag.* 80. t. 5.—Sibir. baical.
lunata, Pers. Syn. ii. 333 = Astragalus lunatus.
lycotriche, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 134 = aciphylla.

OXYTROPIS:—
macrobotrys, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 53.—Mont. Altai.
macrocarpa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 326.—Turkestan.
macroptera, Fisch. ex DC. Prod. ii. 283 = Astragalus macropterus.
macrosema, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 101.—Sibir. altaic.
malacophylla, *Bunge, l. c.*—Mont. Altai.
manshurica, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 12, in obs.—Manchuria.
Maydelliana, *Trautv. in Act. Hort. Petrop.* vi. (1879) 16.—As. bor.
Meinshauseni, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 254.—Reg. Himal; Sibir.
melaleuca, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 15.—As. bor.
melanocalyx. *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 8.—China.
melanotricha, *Bunge, l. c.* 26.—Turkestan.
merkensis, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 11.—As. bor.
Mertensiana, *Turcz, l. c.* (1840) 68.—Fret. Behring.
Meyeri, *Bunge, ex Boiss. Fl. Orient.* ii. 504.—Reg. Caucas.
micans, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 138, in textu = splendens.
micrantha, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 470.—Mongolia.
microphylla, *DC. Astrag.* 83.—Sibir.
microphylla, Hook. f. & Thoms. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 155 = tibetica.
microrhyncha, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 137.—Reg. Himal.
microrhyncha, Strach. & Winterb. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 8 = lapponica.
microsphaera, *Bunge, l. c.* 64.—Turkestan.
Middendorffii, *Trautv. in Middend. Sibir. Reise*, 49. t. 7.—Sibir.
mixotriche, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 589; *Verz. Suppl. Fl. Alt.* 89.—Dahuria.
Moellendorffii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 469.—China.
mollis, *Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.
mollissima, C. A. Mey. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xv. (1869) I. 87 = Astragalus saganlugensis.
montana, Benth. Cat. Pl. Pyr. 110 = pyrenaica.
montana, *DC. Astrag.* 66.—Europ.
montana, Guss. Pl. Rar. 393 = lapponica.
montana, Spreng. Fl. Hal. 207 = Astragalus hypoglottis.
montana, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 54 = cyanea.
monticola, *A. Gray, in Proc. Am. Acad.* xx. (1885) 6.—Am. bor.
mugodsharica, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 68.—Soongar.
multiceps, A. Gray, ex Bunge, l. c. 162, partim = Hallii.
multiceps, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 341.—Am. bor.
muricata, *DC. Astrag.* 86.—Sibir.
myriophylla, *DC. l. c.* 87.—Sibir.
nana, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 410 = multiceps.
nana, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 340.—Am. bor.
neglecta, J. Gay, ex Tenore, Syll. Fl. Nap. 368 = lapponica.
nigrescens, *Fisch. ex DC. Prod.* ii. 278.—Sibir.; Am. bor. et arct.
nitens, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 746.—Sibir. baical.
nivea, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 85.—Mont. Altai.
nutans, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 17.—Sibir.
ochotensis, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 95.—Sibir.
ochrantha, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 188.—Mongol.
ochrocephala, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 57.—China.
ochroleuca, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 6.—As. bor.

OXYTROPIS:—
oligantha, Bunge, in Bull. Soc. Nat. Mosc. xxxix. (1866) II. 18 = chionobia.
oligantha, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 586; *Verz. Suppl. Fl. Alt.* 85.—Sibir. altaic.
oreophila, *A. Gray, in Proc. Am. Acad.* xx. (1885) 3.—Am. bor.
Owerini, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1875) I. 105.—Reg. Caucas.
oxyphylla, *DC. Astrag.* 84.—As. bor.
oxyphylla, Richards. in Frankl. Journ. App. ed. II. 28 = splendens.
pagobia, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 27.—Turkestan.
Pallasii, *Pers. Syn.* ii. 334.—Ross. austr.; Reg. Caucas.
pallens, Turcz. ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 100, in textu = candicans.
Parryi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 4.—Mexic.
Parvopassuae, Parl. Viag. Cat. Monte Bianco, 21 = Gaudini.
pauciflora, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 227 = O. Tschujae.
pellita, *Bunge, in Mém. Acad. Sc. Pétersb.* Sér. VII. xxii. (1874) I. 147.—Soongar.
persica, *Boiss. Diagn.* Ser. I. ii. 40.—Persia.
petraea, Steud. Nom. ed. II. i. 731, in syn., sphalm. = Onobrychis petraea.
physocarpa, *Ledeb. Fl. Alt.* iii. 272.—Sibir. altaic.
physodes, DC. Astrag. 77 = ampullata, caespitosa.
pilosa, *DC. l. c.* 91.—Europ.; As. bor.
platonychia, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 44.—Kokan.
plattensis, Nutt. in Torr. & Gray, Fl. N. Am. i. 340 = Lambertii.
platysema, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 254.—Sibir.
podocarpa, *A. Gray, in Proc. Am. Acad.* vi. (1864) 234.—Am. bor. occ.
podoloba, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 327.—Soongar.
polaris, Seem. in Narr. Voy. Herald, ii. 24 = Astragalus polaris.
polyphylla, *Ledeb. Fl. Alt.* iii. 277.—Sibir. altaic.
proboscidea, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 17.—Tibet.
prostrata, *DC. Astrag.* 85.—Sibir.
psammocharis, *Hance, in Journ. Linn. Soc.* xiii. (1873) 78.—China.
pumila, *Fisch. ex DC. Prod.* ii. 279.—Sibir. altaic.
pumila, Ledeb. Fl. Alt. iii. 275 = stenophylla.
Pumilio, *Ledeb. Fl. Ross.* i. 589. = Ins. Curil.
pusilla, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 27.—Tibet.
pyrenaica, *Gren. & Godr. Fl. Fr.* i. 449.—Europ. austr.
racemosa, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 187.—Mongol.
recognita, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 79.—Soongaria.
revoluta, *Ledeb. Fl. Ross.* i. 588.—Kamtschatka.
rhaetica ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 55.—Europ.
rhynchophysa, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 196.—Soongar.
rosea, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 69.—Turkestan.
rupifraga, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 8.—Turkestan.
samurensis, *Bunge, ex Boiss. Fl. Orient.* ii. 504.—Reg. Caucas.
savellanica, *Bunge, ex Boiss. l. c.* 503.—Persia.
Schmidtii, *Meinsh. in F. Schmidt, Mammuth.* 97.—Sibir. arct.
Schrenkii, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) I. 486.—As. bor.
selengensis, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 143.—Mongolia.
Semenowii, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 13.—As. bor.
sericea, Nutt. in Torr. & Gray, Fl. N. Am. i. 339 = Lambertii.
sericea, Schur, Enum. Pl. Transs. 162 = uralensis.
setosa, *DC. Astrag.* 71.—Mont. Alta.
Sewerzowii, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 36.—Turkestan.

OXYTROPIS:—
songarica, *DC. Astrag.* 73.—Sibir.
sordida, Gaud. ex Nym. Consp. 196 = campestris.
sordida, *Pers. Syn.* ii. 332.—Europ.
splendens, *Dougl. in Hook. Fl. Bor. Am.* i. 147.—Am. bor. occ.
squammulosa, *DC. Astrag.* 79. t. 3.—Dahuria.
ssojutica, Less. ex Walp. Rep. ii. 862 = uralensis.
stenophylla, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 587.—Sibir. alt.
Stracheyana, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 62, in adnot.—Reg. Himal.
strobilacea, *Bunge, l. c.* 103.—China.
subfalcata, *Hance, in Journ. Linn. Soc.* xiii. (1873) 78.—China.
submutica, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 50, in adnot.—Kurdistan.
subverticillaris, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 274.—Sibir. altaic.
sulfurea, Bunge, in Mém. Acad. Sc. Pétersb. Sér. VII. xxii. (1874) I. 80 = seq.
sulphurea, *Fisch. ex Steud. Nom.* ed. II. ii. 244; *Ledeb. Fl. Alt.* iii. 285, partim.—Sibir. altaic.
sulphurea, Ledeb. l. c. partim = recognita.
sylvatica, *DC. Astrag.* 82.—Sibir.
sylvicola, Bunge, in Mém. Acad. Pétersb. Sér. VII. xxii. (1874) I. 138 = sylvatica.
Szovitsii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 57.—Persia.
tachtensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 263.—Turkestan.
tatarica, *Cambess. ex Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 16.—Reg. Himal.; Turkest.
tatarica, Hook. f. & Thoms. ex Bunge, l. c. 17 = proboscidea.
taurica, Pers. ex DC. Prod. ii. 284 = Astragalus tauricus.
tenella, Fisch. ex DC. l. c. 281 = floribunda.
teres, DC. Astrag. 92 = vaginata.
Thomasii, Gaud. Fl. Helv. iv. 543 = lapponica.
Thomsoni, *Benth. ex Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 72.—Ind. or.
tianschanica, *Bunge, in Rupr. Sert. Tiansch.* 43.—As. centr.
tibetica, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 155.—Tibet.
Tilingii, *Bunge, l. c.* 94.—Sibir.
tragacanthoides, *Fisch. ex DC. Prod.* ii. 280.—Mont. Alta.
Trautvetteri, *Meinsh. in F. Schmidt, Mammuth.* 97.—Sibir.
trichocalycina, *Bunge, ex Boiss. Fl. Orient.* ii. 502.—Turkestan.
trichophysa, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 158.—Turkestan.
triflora, *Hoppe, ex Sturm, Deutschl. Fl.* Heft 49.—Tyrol.
triphylla, *DC. Astrag.* 77.—Sibir. or.
Tschujae, *Bunge, in Mém. Acad. Pétersb.* Sér. VII. xxii. (1874) I. 86.—Turkestan.
tuberculata, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 10 = muricatus.
uncata, Pers. Syn. ii. 332 = Astragalus uncatus.
uralensis, Bieb. Fl. Taur. Cauc. iii. 503 = Owerini.
uralensis, *DC. Astrag.* 68.—Europ.; As. bor.
uralensis, Ledeb. Fl. Alt. iii. 289 = ambigua.
uralensis, C. A. Mey. Verz. Pl. Cauc. 140 = Meyeri.
uratensis, Franch. Pl. Davidian. i. (1884) 91, sphalm. = uralensis.
vaginata, *Fisch. ex DC. Prod.* ii. 281.—Sibir. altaic.
velutina, Schur, Enum. Pl. Transs. 162 = uralensis.
venosa, Steud. Nom. ed. II. i. 732, in syn., sphalm. = Onobrychis venosa.
versicolor, Fisch. ex DC. Prod. ii. 281 = floribunda.
verticillaris, *DC. Astrag.* 88.—Sibir.
verticillaris, Ledeb. Fl. Ross. i. 579 = oxyphylla.
viscida, Nutt. ex Torr. & Gray, Fl. N. Am. i. 341 = campestris.
viscosa, Pers. Syn. ii. 332 = foetida.
viscosa, Turcz. ex Steud. Nom. ed. II. ii. 244 (Quid?).—Sibir. baical.

OXYURA, DC. Prod. v. 693 (1836) = **Layia**, Hook. & Arn. (Compos.).
chrysanthemoides, DC. Prod. v. 693 = L. chrysanthemoides.

OXYURA :—
chrysanthemoides, Lindl. Bot. Reg. t. 1850 = Layia calliglossa.
Douglasii, E. Mey. Ind. Sem. Hort. Regiom. (1845) = L. Douglasii.

OYEDAEA, DC. Prod. v. 576 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 374.
SERPAEA, Gardn. in Hook. Lond. Journ. vii. (1848) 296.
acuminata, *Benth. & Hook. f. Gen.* ii. 374; *et ex Hemsl. Biol. Centr. Am. Bot.* ii. 176.—Mexic.
ampeloides, *Hemsl. l. c.*—Mexic.
angustifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 293.—Bras.
bahiensis, *Baker, in Mart. Fl. Bras.* vi. III. 206.—Bras.
Bonplandiana, *Benth. & Hook. f. Gen.* ii. 374.—Bras.
buphthalmoides, *DC. Prod.* v. 577.—Peruv.
Cuerviana, *Triana, in Ann. Sc. Nat.* Sér. IV. ix. (1858) 39.—N. Granat.
helianthoides, *Triana, l. c.* 38.—N. Granat.
Humboldtiana, *Benth. & Hook. f. Gen.* ii. 374.—Bras.
lippioides, *Baker, in Mart. Fl. Bras.* vi. III. 208.—Bras.
macrophylla, *Benth. & Hook. f. Gen.* ii. 374; *et ex Hemsl. Biol. Centr. Am. Bot.* ii. 176.—Mexic.
ovalifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 183.—Mexic.
ovata, *Benth. & Hook. f. Gen.* ii. 374.—Bras.
rotundifolia, *Baker, in Mart. Fl. Bras.* vi. III. 208.—Bras.
Seemanni, A. Gray, in Proc. Am. Acad. xix. (1883) 10 = Viguiera Seemannii.
verbesinoides, *DC. in Deless. Ic. Sel.* iv. t. 34.—Venezuela.
vestita, *Baker, in Mart. Fl. Bras.* vi. III. 207.—Bras.

OZANTHES, Rafin. Sylva Tellur. 133 (1838) = **Lindera**, Thunb. (Laurin.).
Benzoin, Rafin. l. c. = L. Benzoin.

OZARTHRIS, Rafin. Fl. Tellur. iv. 92 (1836) = **Cassytha**, Linn. (Laurin.).
capense, Rafin. l. c. = C. capensis.
opuntioides, Rafin. l. c.—Hab. ?
paradoxa, Rafin. l. c.—Hab. ?
triquetra, Rafin. l. c. = Hab. ?

OZIROË, Rafin. Fl. Tellur. iii. 53 (1836) = **Ornithogalum**, Tourn. (Liliac.).
leuchlora, Rafin. l. c. 54 = O. biflorum.

OZODIA, Wight & Arn. Prod. 375 (1834) = **Foeniculum**, Tourn. (Umbell.).
foeniculacea, Wight & Arn. l. c. = F. vulgare.

OZODYCUS, Rafin. Atl. Journ. 145 (1832) = **Cucurbita**, Tourn.
perennis, Rafin. l. c. = C. foetidissima.

OZOMELIS, Rafin. Fl. Tellur. ii. 73 (1836) = **Mitella**, Tourn. (Saxifr.).
varians, Rafin. l. c. = M. trifida.

OZOPHYLLUM, Schreb. Gen. 452 (1791) = **Ticorea**, Aubl. (Rutac.).
foetidum, Raeusch. Nom. ed. III. 191 = T. foetida.
trifoliatum, Willd. Sp. Pl. iii. 585 = T. foetida.

OZOROA, Delile, in Ann. Sc. Nat. Sér. III. xx. (1843) 91. t. 1 = **Anaphrenium**, E. Mey. (Anacard.).
insignis, Delile, l. c. = A. abyssinicum.

OZOTHAMNUS, R. Br. in Trans. Linn. Soc. xii. (1816) 125 = **Helichrysum**, Vaill. (Compos.).
adnatus, DC. Prod. vi. 166 = H. adnatum.
antennaria, Hook. f. Fl. Tasman. i. 203 = H. antennarium.
Backhousii, Hook. f. l. c. 204 = H. Backhousii.
Becklerii, F. Muell. Fragm. i. 183 = H. Becklerii.
bracteolatus, Hook. f. Fl. Tasman. i. 203 = H. bracteolatum.
cinereus, DC. Prod. vi. 165 = H. cinereum.
coralloides, Hook. f. Fl. N. Zel. ii. 332.—N. Zel.
decurrens, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 59 = H. decurrens.

OZOTHAMNUS :—
depressus, Hook. f. Fl. N. Zel. 134.—N. Zel.
diosmaefolius, DC. Prod. vi. 166 = Helichrysum diosmaefolium.
diotophyllus, F. Muell. Fragm. v. 150 = H. diotophyllum.
ericaefolius, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 119 = seq.
ericifolius, A. Cunn. ex DC. Prod. vi. 166 = H. diosmaefolium.
ferrugineus, DC. l. c. 165 = Helipterum citrinum.
glomeratus, Hook. f. Fl. N. Zel. 133.—N. Zel.
Gunnii, Hook. f. Fl. Tasman. i. 205 = H. Gunnii.
Hookeri, Sond. in Linnaea, xxv. (1852) 509 = H. baccharoides.
lanceolatus, J. Buch. in Trans. N. Z. Inst. ii. (1870) 88.—N. Zel.
ledifolius, Hook. f. Fl. Tasman. i. 204 = H. ledifolium.
lepidophyllus, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 120 = H. baccharoides.
lepidophyllus, Steetz, in Lehm. Pl. Preiss. i. 468 = H. lepidophyllum.
lycopodioides, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 119 = H. lycopodioides.
microphyllus, Hook. f. Fl. N. Zel. 134. t. 35 A.—N. Zel.
obcordatus, DC. Prod. vi. 165 = H. obcordatum.
pholidotus, F. Muell. Fragm. ii. 131 = H. pholidotum.
pinifolius, DC. Prod. vi. 166; R. Br. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 129.—N. Zel.
purpurascens, DC. l. c. 165 = H. diosmaefolium.
reticulatus, DC. l. c. 164 = H. reticulatum.
retusus, Sond. & F. Muell. in Linnaea, xxv. (1852) 510 = H. adnatum.
rosmarinifolius, DC. Prod. vi. 165 = H. diosmaefolium.
rufescens, DC. l. c.—Austral.
scaber, F. Muell. in Linnaea, xxv. (1852) 407 = H. adnatum.
scutellifolius, Hook. f. Fl. Tasman. i. 202. t. 56 = H. scutellifolium.
selaginioides, Sond. & F. Muell. in Linnaea, xxv. (1852) 510 = H. selaginioides.
Selago, Hook. f. Fl. N. Zel. i. 332.—N. Zel.
tephrodes, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 79 = H. obtusifolium.
thyrsoideus, DC. Prod. vi. 165 = H. diosmaefolium.
turbinatus, DC. l. c. 164 = H. cinereum.
Vauvilliersii, Hombron & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 38 = Cassinia Vauvilliersii.

OZOTIS, Rafin. Alsog. Am. 71 (1838) = **Aesculus**, Linn. (Sapind.).
trifoliata, Rafin. l. c. = Ae. glabra.

OZOXETA, Rafin. Sylva Tellur. 75 (1838) = **Helicteres**, Pluk. (Stercul.).
ovata, Rafin. l. c. = H. pentandra.

P

PACHEA, Pourr. ex Steud. Nom. ed. II. i. 449, in syn. (1840) = **Crypsis**, Ait. (Gramin.).

PACHIDENDRON, Haw. Rev. Pl. Succ. 35 (1821) = **Aloë**, Tourn. (Liliac.).
africanum, Haw. l. c. 36 = A. africana.
angustifolium, Haw. l. c. 38 = Gasteria disticha.
ferox, Haw. l. c. = A. ferox.
Principis, Haw. l. c. 37 = A. Salm-Dyckiana.
Pseudoafricanum, Haw. ex Steud. Nom. ed. II. ii. 245 = A. africana.
Pseudoferox, Haw. Rev. Pl. Succ. 38 = A. ferox.
supralaeve, Haw. l. c. 40 = A. supralaevis.

PACHILA, Rafin. in Loud. Gard. Mag. viii. (1832) 246 = **Erucaria**, Gaertn. (Crucif.).

PACHILOMA, Rafin. New Fl. Am. iv. 33 (1836) = **Polytaenia**, DC. (Umbell.).
Nuttali, Rafin. l. c. = P. Nuttallii.
verticillata, Rafin. l. c. = P. Nuttallii.

PACHIPHILLUM, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 42 = **Pachyphyllum**, H. B. & K. (Orchid.).

PACHIRA, Aubl. Pl. Gui. ii. 725. tt. 291, 292 (1775). *MALVACEAE*, Benth. & Hook. f. i. 210.
CAROLINEA, Linn. f. Suppl. 51 (1781).
PACHYRA, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 209.
SOPHIA, Linn. Pl. Surin. n. 86 (1775); Amoen. Acad. viii. 259.
affinis, *Decne. in Fl. des Serres*, xxiii. (1880) 52.—Bras.
alba, *Walp. Rep.* i. 329.—N. Granat.
angusta, Duchass. ex Decne. in Fl. des Serres, xxiii. (1880) 52 = sessilis.
aquatica, *Aubl. Pl. Gui.* ii. 726. tt. 291, 292.—Am. trop.
arenaria, *A. St. Hil. Fl. Bras. Mer.* i. 261.—Bras.
aurea, *Decne. in Fl. des Serres*, xxiii. (1880) 47.—Mexic.
Barrigon, *Seem. Bot. Voy. Herald*, 83.—Panama.
bracteolata, *Decne. in Fl. des Serres*, xxiii. (1880) 51.—Venezuela.
campestris, *Decne. l. c.* 52.—Bras.
Commersonii, *Planch. ex Decne. l. c.* 48.—Bras.
cyathophora, *Casar. Nov. Stirp. Bras. Dec.* 21.—Bras.
decaphylla, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 209.—Bras.
emarginata, *A. Rich. Ess. Fl. Cub.* 176.—Cuba.
fastuosa, *Decne. in Fl. des Serres*, xxiii. (1880) 48.—Mexic.
Fendleri, Seem. Bot. Voy. Herald, 83 = Bombax Fendleri.
glabra, *Pasq. in Rendic. Acc. Nap.* vii. (1868) 18.—Hab. ?
grandiflora, *Tussac, Fl. Antill.* iv. 12. tt. 3, 4.—Ind. occ.
humilis, Spruce, ex Benth. in Journ. Linn. Soc. vi. (1862) 108 = Bombax humile.
insignis, *Savign. in Lam. Encyc.* iv. 690.—Mexic.
Loddigesii, *Decne. in Fl. des Serres*, xxiii. (1880) 51.—Ind. occ.
longiflora, *Decne. l. c.* 52.—Bras.
longifolia, *Hook. Bot. Mag.* t. 4549.—Mexic.
macrantha, *A. St. Hil. Fl. Bras. Mer.* i. 261.—Bras.
macrantha, *Spruce, ex Decne. in Fl. des Serres*, xxiii. (1880) 47.—Bras.
macrocarpa, *Walp. Rep.* i. 329.—Mexic.
marginata, *A. St. Hil. Fl. Bras. Mer.* i. 260.—Bras.
minor, *Hemsl. Biol. Centr. Am. Bot.* i. 124.—Mexic.
nitida, H. B. & K. Nov. Gen. et Sp. v. 302 = aquatica.
oleagina, *Decne. in Fl. des Serres*, xxiii. (1880) 49.—Hab. ?
pulchra, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 320.—N. Granat.
rufescens, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 210 = Bombax carolinum.
sessilis, *Benth. Bot. Voy. Sulph.* 70.—Panama.
speciosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 319.—N. Granat.
Spruceana, *Decne. in Fl. des Serres*, xxiii. (1880) 46.—Bras.
stenopetala, *Casar. Nov. Stirp. Bras. Dec.* 21.—Bras.
tomentosa, *Endl. ex Walp. Rep.* i. 329.—Bras.

PACHISTIMA, Rafin. in Am. Monthly Mag. (1818) 176. *CELASTRINEAE*, Benth. & Hook. f. i. 361.
OREOPHILA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 258 (1840).
PACHYSTIGMA, Meissn. Gen. Comm. 348 (1843), sphalm.
Canbyi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 623.—Am. bor.
Myrsinites, *Rafin. in Am. Monthly Mag.* (1818) 176.—Am. bor.

PACHITES, Lindl. Gen. et Sp. Orch. 301 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 629.
appressa, *Lindl. l. c.*—Afr. austr.

PACHYANTHUS, A. Rich. Ess. Fl. Cub. i. 559 (1845). *MELASTOMACEAE*, Benth. & Hook. f. i. 762.
CHALYBEA, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 99.
SARCOMERIS, Naud. l. c. xv. (1851) 340.

PACHYANTHUS:—
angustifolius, *Griseb. Cat. Pl. Cub.* 101.—Cuba.
corymbosus, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 97.—N. Granat.
cubensis, *A. Rich. Ess. Fl. Cub.* i. 560.—Cuba.
oleifolius, *Griseb. Cat. Pl. Cub.* 100.—Cuba.
Poiretii, *Griseb. l. c.* 101.—Cuba.
Wrightii, Griseb. l. c. = Miconia Wrightii.

PACHYCALYX, Klotzsch, in Linnaea, xii. (1838) 230 = **Simocheilus**, Klotzsch (Ericac.).
glaber, Klotzsch, l. c. 231 = S. glaber.
hispidus, Klotzsch, l. c. = S. hispidus.
inaequalis, Klotzsch, l. c. = S. glaber.
pubescens, Klotzsch, l. c. 230 = S. Ecklonianus.

PACHYCARPUS, E. Mey. Comm. Pl. Afr. Austr. 209 (1837) = **Gomphocarpus**, R. Br. (Asclep.).
albens, E. Mey. l. c. 214 = G. albens.
appendiculatus, E. Mey. l. c. 210 = G. appendiculatus.
asperifolius, Meissn. in Hook. Lond. Journ. Bot. ii. (1848) 544 (err. typ. 444) = G. asperifolius (cf. Addend.).
concolor, E. Mey. Comm. Pl. Afr. Austr. 210 = G. concolor.
corniculatus, Hochst. in Flora, xxvii. (1844) 101 = Asclepias micrantha.
coronarius, E. Mey. Comm. Pl. Afr. Austr. 209 = G. coronarius.
crispus, E. Mey. l. c. 214 = G. crispus.
dealbatus, E. Mey. l. c. 211 = G. dealbatus.
gomphocarpoides, E. Mey. l. c. 213.—Afr. austr.
grandiflorus, E. Mey. l. c. 209 = G. grandiflorus.
humilis, E. Mey. l. c. 212 = G. humilis.
ligulatus, E. Mey. l. c. 211 = G. dealbatus.
marginatus, E. Mey. l. c. 213 = G. marginatus.
orbicularis, E. Mey. l. c. 212 = Xysmalobium orbiculare.
reflectens, E. Mey. l. c. 210 = G. reflectens.
reflexus, Steud. Nom. ed. II. ii. 245 = praec.
rigidus, E. Mey. Enum. Pl. Afr. Austr. 211 = G. rigidus.
vexillaris, E. Mey. l. c. 212 = G. dealbatus.
vexillatus, Steud. Nom. ed. II. ii. 245 = praec.
viridiflorus, E. Mey. Comm. Pl. Afr. Austr. 214 = G. viridiflorus.

PACHYCENTRIA, Blume, in Flora, xiv. (1831) 519. *MELASTOMACEAE*, Benth. & Hook. f. i. 760.
constricta, *Blume, l. c.* 520.—Java.
cordata, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 24 = tuberculata.
elliptica, *Blume, l. c.* 23.—Borneo.
glauca, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 89.—Borneo.
Junghuhniana, *Miq. Fl. Ind. Bat.* i. 1. 552.—Java.
laxiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 23.—Sumatra.
rigida, *Blume, l. c.* 23.—Sumatra.
tuberculata, *Korth. Verh. Nat. Gesch. Bot.* 246. t. 63.—Borneo.
varingaefolia, *Blume, in Flora*, xiv. (1831) 520.—Java.
varingaefolia, Korth. in Verh. Nat. Gesch. Bot. t. 63. f. 2 = laxiflora.
varingaefolia, Mor. Syst. Verz. Zoll. 11 = Zollingeriana.
Walkeri, Thw. Enum. Pl. Zeyl. 107 = Kendrickia Walkeri.
Zollingeriana, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 301.—Java.

PACHYCENTRON, Pomel, Nouv. Mat. Fl. Atl. 32 (1874) = **Centaurea**, Linn. (Compos.).

PACHYCHAETA, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 186 (1876) = **Ophryosporus**, Meyen (Compos.).
eupatorioides, Sch. Bip. ex Baker, l. c. 187 = O. Pachychaeta.

PACHYCHILUS, Blume, Fl. Jav. Praef. 7 (1828) = **Pachystoma**, Blume (Orchid.).
chinensis, Blume, Mus. Bot. Lugd. Bat. ii. 173 = Pachystoma chinensis.
montanus, Blume, l. c. = Pachystoma senile.

PACHYCHILUS:—
pantanus, Blume, Mus. Bot. Lugd. Bat. ii. 173 = Pachystoma pantanum.
pubescens, Blume, l. c. = Pachystoma pubescens.
senilis, Blume, l. c. = Pachystoma senile.
Smithianus, Blume, l. c. = Pachystoma senile.
speciosum, Blume, Orch. Archip. Ind. 29. t. 22 = Pachystoma senile.

PACHYCLADON, Hook. f. Handb. N. Zeal. Fl. 724 (1864). *CRUCIFERAE*, Benth. & Hook. f. i. 967.
glabra, *J. Buch. in Trans. N. Z. Inst.* xiv. (1882) 344.—N. Zel.
Novae-Zelandiae, *Hook. f. Handb. N. Zeal. Fl.* i. 724.—N. Zel.

PACHYCORNIA, Hook. f. in Benth. & Hook. f. Gen. iii. 65 (1880). *CHENOPODIACEAE*, Benth. & Hook. f. l. c.
robusta, *Hook. f. l. c.*—Austral.

PACHYDENDRON, Dum. Anal. Fam. 60 (1829) = PACHIDENDRON, Haw. = **Aloë**, Linn. (Liliac.).

PACHYDERIS, Cass. in Dict. Sc. Nat. lvi. 170 (1828) = **Pteronia**, Linn. (Compos.).
oblongifolia, Steud. Nom. ed. II. ii. 245, sphalm. = seq.
obtusifolia, Cass. in Dict. Sc. Nat. lvi. 170 = Pteronia leptalepis.

PACHYDERMA, Blume, Bijdr. 682 (1825) = **Olea**, Tourn.
javanicum, Blume, l. c. = O. maritima.

PACHYGLOSSUM, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 345, in obs. = **Oxypetalum**. R. Br. (Asclep.).

PACHYGONE, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43. *MENISPERMACEAE*, Benth. & Hook. f. i. 38.
HYPERBAENA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44.
KOON, Gaertn. Fruct. ii. 486. t. 180 (1791), nomen prius.
adversa, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 321; Contrib. iii. 332 = ovata.
brachystachys, Miers, l. c. Ser. II. vii. (1851) 43; Ser. III. xix. (1867) 321; Contrib. iii. 334 = ovata.
columbica, *Eichl. in Flora*, xlvii. (1864) 394.—N. Granat.
concinna, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 321; Contrib. iii. 331 = ovata.
cubensis, *Griseb. Cat. Pl. Cub.* 4.—Cuba.
dasycarpa, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 62.—Burma.
domingensis, *Eichl. in Flora*, xlvii. (1864) 394.—Ins. S. Doming.
graciliflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xi. (1881) 141.—Guian.
hebephylla, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 321; Contrib. iii. 333 = ovata.
Hullsii, *F. Muell. Fragm.* ix. 81.—Austral.
leptostachys, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43; Ser. III. xix. (1867) 321; Contrib. iii. 335 = ovata.
oblongifolia, *Eichl. in Flora*, xlvii. (1864) 394.—Bras.
odorifera, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 321; Contrib. iii. 333 = ovata.
ovata, *Miers, ex Hook. f. & Thoms. Fl. Ind.* i. 203.—Ind. or.; Malaya.
Plukenetii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 43; Ser. III. xix. (1867) 321; *Contrib.* iii. 330.—Ind. or.
pubescens, *Benth. Fl. Austral.* i. 58.—Austral.
Wightiana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43 = ovata.

PACHYLAENA, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 106. *COMPOSITAE*, Benth. & Hook. f. ii. 495.
CHIONOPTERA, DC. Prod. vii. 14 (1838).
atriplicifolia, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 106.—Chili.
elegans, *Phil. in Linnaea*, xxxiii. (1864–65) 118.—Chili.

PACHYLEPIS, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1831) 189 = **Callitris**, Vent. (Conif.).
Commersonia, Brongn. l. c. (= *Widdringtonia Commersonii*).—Madag.
cupressoides, Brongn. l. c. = C. cupressoides.
juniperoides, Brongn. l. c. = C. arborea.

PACHYLEPIS, Less. Syn. Comp. 139 (1832) = **Rodigia**, Spreng. (Compos.).
Kalmii, Less. l. c.—Am. bor.

PACHYLOBUS, G. Don, Gen. Syst. ii. 89 (1832) = **Canarium**, Linn. (Burser.).
edulis, G. Don, l. c. = C. edule.

PACHYLOMA, DC. Prod. iii. 122 (1828) = **Comolia**, DC. (Melastom.).
coriaceum, DC. l. c. 123.—Bras.
huberioides, Triana, in Trans. Linn. Soc. xxviii. (1871) 64 = Urodesmium huberioides.

PACHYLOMA, Spach, Hist. Vég. Phan. vii. 194 (1839) = **Ranunculus**, Tourn.
arvense, Spach, l. c. 195 = R. arvensis.

PACHYLOPHUS, Spach, Hist. Vég. Phan. iv. 365 (1835) = **Oenothera**, Linn. (Onagrar.).
Nuttalliana, Spach, in Ann. Mus. Par. Sér. III. iv. (1835) 356. t. 30 = seq.
Nuttallii, Spach, Hist. Vég. Phan. iv. 365 = Oe. caespitosa.

PACHYMERIA, Benth. Pl. Hartw. 130 (1844) = **Meriania**, Sw. (Melastom.).
rigida, Benth. l. c. = M. rigida.

PACHYMITRA, Nees, in Mart. Fl. Bras. ii. 1. 115 (1842) = **Rynchospora**, Vahl (Cyperac.).
candida, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 606 = R. candida.
conferta, Nees, ex Boeck. l. c. 615 = R. velutina.
rufa, Nees, ex Boeck. l. c. 616 = R. rufa.
velutina, Nees, in Mart. Fl. Bras. ii. 1. 115 = R. velutina.

PACHYNE, Salisb. in Trans. Hort. Soc. i. (1812) 299 = **Phaius**, Lour. (Orchid.),
spectabilis, Salisb. l. c. = Ph. grandifolius.

PACHYNEMA, R. Br. ex DC. Syst. i. 411 (1818). *DILLENIACEAE*, Benth. & Hook. f. i. 15.
HUTTIA, Drumm. & Harv. in Hook. Kew Journ. vii. (1855) 51.
complanatum, *R. Br. ex DC. Syst.* i. 412.—Austral.
conspicuum, *Benth. Fl. Austral.* i. 47.—Austral.
dilatatum, *Benth. l. c.* 48.—Austral.
junceum, *Benth. l. c.* i. 47.—Austral.
sphenandrum, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* v. (1882) 79.—Austral.

PACHYNEURUM, Bunge, Del. Sem. Hort. Dorp. (1839) 8; et in Linnaea, xiv. (1840) Litt. 121 = **Parrya**, (Crucif.).
grandiflorum, Bunge, ll. cc. = Parrya microcarpa.

PACHYNOCARPUS, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 159. t. 22. *DIPTEROCARPEAE*, Benth. & Hook. f. i. 192.
umbonatus, *Hook. f. l. c.*—Borneo.

PACHYPHRAGMA, Reichb. Nom. 179 (1841) = **Thlaspi**, Dill. (Crucif.).

PACHYPHYLLUM, H. B. & K. Nov. Gen. et Sp. i. 338. t. 77. (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 571.
crystallinum, *Lindl. Orch. Linden.* 18.—Venezuela.
Cuencae, *Reichb. f. in Linnaea*, xli. (1877) 31.—Ecuador.
distichum, *H. B. & K. Nov. Gen. et Sp.* i. 339. t. 77.—Peruv.
distichum, Lindl. ex Reichb. f. in Bonplandia, iii. (1855) 219 = Hartwegii.
echinocarpon, *Spreng. Syst.* iii. 731.—Am. austr.
falcifolium, *Reichb. f. in Linnaea*, xli. (1877) 32.—Peruv.
Hartwegii, *Reichb. f. in Bonplandia*, iii. (1855) 219.—Ecuador.

PACHYPHYLLUM :—
parvifolium, *Lindl. in Hook. Ic. Pl.* ii. (1837) t. 117.—Peruv.
Pastii, *Reichb. f. in Bonplandia*, iii. (1855) 239.—Peruv.
pectinatum, *Reichb. f. Xenia Orch.* iii. 22.—Chili.
procumbens, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 601, nomen.—Mexic.
Pseudo-dichaea, *Reichb. f. Xenia Orch.* iii. 22.—Bolivia.
scandens, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 28.—Mexic.
Serra, *Reichb. f. in Bonplandia*, iii. (1855) 219.—Peruv.
squarrosum, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 107.—N. Granat.
uniflorum, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc*) 42.—Mexic.

PACHYPHYTUM, Link, Klotzsch & Otto, in Otto & Dietr. Allg. Gartenz. ix. (1841) 9 = **Cotyledon**, Linn. (Crassul.).
bracteosum, Link, Klotzsch & Otto, l. c. 10 = C. Pachyphytum.
Lingua, Hort. ex Baker, in Saund. Refug. Bot. sub t. 58 = C. linguaefolia.
roseum, Hort. ex Baker, l. c. sub t. 60 = C. adunca.

PACHYPLEURUM, Ledeb. Fl. Alt. i. 296 (1829) = **Ligusticum**, Linn. (Umbell.).
albo-marginatum, Rupr. Sert. Tiansch. 49 = L. simplex.
alpinum, Ledeb. Fl. Alt. i. 297.—Sibir. altaic.
pyrenaicum, Mutel, Fl. Fr. ii. 31, 403 = Seseli nanum.
simplex, Reichb. Fl. Germ. Excurs. 471 = L. simplex.

PACHYPODIUM, Lindl. Bot. Reg. t. 1321 (1830). *APOCYNACEAE*, Benth. & Hook. f. ii. 722.
BELONITES, E. Mey. Comm. Pl. Afr. Austr. 187 (1837).
bispinosum, *A. DC. in DC. Prod.* viii. 424.—Afr. austr.
glabrum, G. Don, Gen. Syst. iv. 77 = bispinosum.
Lealii, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 45.—Afr. trop.
namaquanum, *Welw. l. c.*—Afr. austr.
obesum, A. DC. in DC. Prod. viii. 412, in syn. = Adenium obesum.
rosulatum, *Baker, in Journ. Bot.* xx. (1882) 219.—Madag.
Rutenbergianum, *Vatke, in Bremen, Abh.* ix. (1885) 125.—Madag.
succulentum, Steud. Nom. ed. II. i. 196; ii. 245 = tomentosum.
tomentosum, *G. Don, Gen. Syst.* iv. 78.—Afr. austr.
tuberosum, Lindl. Bot. Reg. t. 1321 = tomentosum ?

PACHYPODIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 96 (1838) = **Thelypodium**, Endl. (Crucif.).
brachycarpum, Torr. Bot. Am. Expl. Exped. ii. t. 1 = T. brachycarpum.
integrifolium, Nutt. ex Torr. & Gray, Fl. N. Am. i. 96 = T. integrifolium.
laciniatum, Nutt. l. c. = T. laciniatum.
linearifolium, A. Gray, in Proc. Am. Acad. vi. (1866) 187 = T. linearifolium.
sagittatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 97 = T. sagittatum.

PACHYPODIUM, Webb & Berth. Phyt. Canar. i. 74 (1836-40) = **Sisymbrium**, Tourn. (Crucif.).
bursifolium, Endl. Gen. ii. 874 = S. bursifolium.
Columnae, Webb, Iter Hisp. 75 = S. Columnae.
erysimoides, Webb & Berth. Phyt. Canar. i. 75 = S. erysimoides.
pannonicum, Endl. Gen. ii. 874 = S. pannonicum.

PACHYPTERA, DC. in Meissn. Gen. 299 ; Comm. 207 (1840) ; et Prod. ix. 175 (1845) = **Adenocalymna**, Mart. (Bignon.).
dasyantha, DC. Prod. ix. 176.—Bras.
foveolata, DC. l. c. 175.—Guian.
Perrottetii, DC. l. c. 176.—Guian.
puberula, DC. l. c. 175.—Bras.
striata, DC. l. c. 176.—Bras.
umbelliformis, DC. l. c. 175.—Bras.

PACHYPTERIS, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 158 = **Pachypterygium**, Bunge, (Crucif.).
multicaule, Kar. & Kir. l. c. 159 = Pachypterygium multicaule.

PACHYPTERYGIUM, Bunge, in Del. Sem. Hort. Dorpat. (1843) 8 ; et in Linnaea, xviii. (1844) 155. *CRUCIFERAE*, Benth. & Hook. f. i. 94.
PACHYPTERIS, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) i. 159.
brevipes, *Bunge, Del. Sem. Hort. Dorpat.* (1843) 8 ; *et in Linnaea*, xviii. (1844) 155.—Persia ; Turkestan.
densiflorum, *Bunge, ex Boiss. Fl. Orient.* i. 373.—Persia.
heterotrichum, *Bunge, l. c.* 374.—Persia.
lamprocarpum, *Bunge, Del. Sem. Hort. Dorpat.* (1843) 8 ; *et in Linnaea*, xviii. (1844) 155.—Persia ; Turkestan.
multicaule, *Bunge, in Linnaea*, xviii. (1844) 155.—Turkestan.
stelligerum, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 227.—Turkestan.

PACHYRA, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 209 = **Pachira**, Aubl. (Malvac.).

PACHYRAPHEA, Presl, Bot. Bemerk. 126 (1844) ; in Abh. Boehm. Ges. V. iii. (1845) 556 = **Aspalathus**, Linn. (Legumin.).
propinqua, Presl, Bot. Bemerk. 126 = A. propinqua.
triquetra, Presl, l. c. = A. triquetra.
Zeyheriana, Presl, l. c. = A. triquetra.

PACHYRHIZUS, Rich. ex DC. Mém. Lég. 379 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 540.
CACARA, Rumph. ex Thou. in Dict. Sc. Nat. v. 35 (1805).
ROBYNSIA, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 193.
TAENIOCARPUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 420.
angulatus, *Rich. ex DC. Prod.* ii. 402.—As. trop.
articulatus, Duchass. ex Walp. in Flora, xxxvi. (1853) 226 = angulatus.
bulbosus, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 246 = angulatus.
Jicamas, Blanco, Fl. Filip. ed. I. 579 = angulatus.
mollis, Hassk. in Flora, xxv. (1842) II. Beibl. 74 ; Cat. Hort. Bog. Alt. 280 = Pueraria phaseoloides.
montanus, DC. Prod. ii. 402 = Pueraria phaseoloides.
orbicularis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 208.—Afr. trop.
palmatilobus, *Benth. & Hook. f. Gen.* i. 540.—Mexic.
teres, Blanco, Fl. Filip. ed. I. 580 = Pueraria phaseoloides.
Thunbergianus, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 237 = Pueraria Thunbergiana.
trilobus, DC. Prod. ii. 402 = angulatus, Pueraria Thunbergiana.
tuberosus, *Spreng. Syst.* iv. *Cur. Post.* 281.—Ind. occ.

PACHYRHYNCHUS, DC. Prod. vi. 255 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 313.
xeranthemoides, *DC. l. c.*—Afr. austr.

PACHYRRHIZOS, Spreng. Syst. iv. Cur. Post. 281 (1827) = **Pachyrhiza**, Rich. (Legumin.).

PACHYSA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = **Erica**, Tourn.
ardens, D. Don, l. c. = E. ardens.
aristata, G. Don, Gen. Syst. iii. 806 = E. Vernix.
baccans, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = E. baccans.
formosa, G. Don, Gen. Syst. iii. 806 = E. formosa.
glomiflora, G. Don, l. c. = E. reflexa.
Lambertiana, G. Don, l. c. = E. physodes.
physodes, G. Don, l. c. = E. physodes.
Vernix, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = E. Vernix.
vesicularis, G. Don, Gen. Syst. iii. 806 = E. vesicularis.

PACHYSANDRA, Michx. Fl. Bor. Am. ii. 177 (1803). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 267.
coriacea, Hook. Exot. Fl. ii. t. 148 = Sarcococca pruniformis.
erecta, Rafin. ex Baill. Monog. Buxin. 56 = procumbens.
procumbens, *Michx. Fl. Am. Bor.* ii. 178. t. 45.—Am. bor.
terminalis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 142.—Japon.

PACHYSANTHUS, Presl, Bot. Bemerk. 87 (1844) ; in Abh. Boehm. Ges. V. iii. (1845) 517 = **Rudgea**, Salisb. (Rubiac.).
macrophyllus, Presl, ll. cc. = R. Pachysanthus.

PACHYSTACHYS, Nees, in Mart. Fl. Bras. ix. 99 (1847) = **Jacobinia**, Moric. (Acanth.).
asperula, Nees, in DC. Prod. xi. 320 = J. coccinea.
coccinea, Nees, l. c. 319 = J. coccinea.
latior, Nees, l. c. 320 = J. coccinea.
lutea, Nees, l. c.—Peruv.
nutans, Nees, l. c. = J. nutans.
Riedeliana, Nees, in Mart. Fl. Bras. ix. 99.—Bras.

PACHYSTEMON, Blume, Bijdr. 626 (1825) = **Macaranga**, Thou. (Euphorb.).
bancanus, Miq. Fl. Ind. Bat. Suppl. 462 = M. bancana.
depressus, Muell. Arg. in Flora, xlvii. (1864) 465 = M. depressa.
populifolius, Miq. Fl. Ind. Bat. Suppl. 462 = M. populifolia.
trilobus, Blume, Bijdr. 626 = M. triloba.

PACHYSTIGMA, Hochst. in Flora, xxv. (1842) 234 = **Fadogia**, Schweinf. (Rubiac.).
lasianthum, Sond. in Linnaea, xxiii. (1850) 55.—Afr. austr.
latifolium, Sond. l. c. 56.—Afr. austr.
venosa, Hochst. in Flora, xxv. (1842) 235.—Afr. austr.
Zeyheri, Sond. in Linnaea, xxiii. (1850) 56 = F. Zeyheri (cf. Addend.).

PACHYSTIGMA, Hook. Ic. Pl. t. 698 (1844) = **Peltostigma**, Walp. (Rutac.).
pteleoides, Hook. l. c. = Peltost. pteleoides.

PACHYSTIGMA, Meissn. Gen. Comm. 348 (1843) sphalm. = **Pachistima**, Rafin. (Celastr.).

PACHYSTIMA, 'Rafin.' ex Endl. Gen. Suppl. i. 1424 (1841) = **Pachistima**, Rafin. (Celastr.).

PACHYSTOMA, Blume, Bijdr. 376 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 511.
APATURIA, Lindl. Gen. et Sp. Orch. 130 (1831).
IPSEA, Lindl. l. c. 124 (1831).
PACHYCHILUS, Blume, Fl. Jav. Praef. 7 (1828).
angustifolium, Reichb. f. in Bonplandia, v. (1857) 54, in nota = Tainia angustifolia.
chinensis, *Reichb. f. l. c.* iii. (1855) 251.—Ins. Hongkong.
Edgworthii, Reichb. f. in Walp. Ann. iii. 464 = senile.
Fortuni, Reichb. f. l. c. = senile.
Josephi, Reichb. f. l. c. = senile.
Lindleyanum, Reichb. f. in Bonplandia. iii. (1855) 251 = senile.
malabaricum, Reichb. f. in Walp. Ann. vi. 462 = senile.
montanum, Reichb. f. in Bonplandia, iii. (1855) 251 = senile.
pantanum, *Miq. Fl. Ind. Bat.* iii. 675.—Borneo.
parvifolium, Reichb. f. in Walp. Ann. vi. 463 = senile.
pubescens, *Blume, Bijdr.* 376 ; *Tabell.* t. 29.—Java.
senile, *Reichb. f. in Bonplandia*, iii. (1858) 251.—As. trop. ; Malaya.
Smithianum, Reichb. f. l. c. = senile.
speciosum, Reichb. f. l. c. 250 = senile.
Thompsoni, Hort. in Gard. Chron. (1879) II. 505, 627. ff. 102, 103 = seq.
Thompsonianum, *Reichb. f. l. c.* 582.—Afr. trop.
Wightii, Reichb. f. in Walp. Ann. vi. 464 = senile.

PACHYSTROMA, [Klotzsch] Muell. Arg. in Linnaea, xxxiv. (1865-66) 177. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 307.
ACANTHOLOMA, Gaudich. ex Baill. Adansonia, vi. (1866) 231. t. 1.
ilicifolium, *Muell. Arg. in Linnaea* xxxiv. (1865-66) 178.—Bras.

PACHYSTYLUM, Eckl. & Zeyh. Enum. 13 (1835) = **Heliophila**, Burm. f. (Crucif.).
glabrum, Eckl. & Zeyh. l. c. = H. glabra.

PACHYSURUS, Steetz, in Lehm. Pl. Preiss. i. 441 (1845) = **Calocephalus**, R. Br. (Compos.).
aervoides, F. Muell. Fragm. iii. 154 = C. aeruoides.
angianthoides, Steetz, in Lehm. Pl. Preiss. i. 442 = C. angianthoides.

PACHYSURUS :—
Francisii, F. Muell. Fragm. iii. 155 = Calocephalus Francisii.
multiflorus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 192 = C. multiflorus.
platycephalus, F. Muell. Fragm. iii. 154 = C. platycephalus.

PACHYTHELIA, Steetz, in Peters, Reise Mossamb. Bot. 453 (1863) = **Epaltes**, Cass. (Compos.).
brasiliensis, Steetz, l. c. 455 = E. brasiliensis.
mexicana, Steetz, l. c. 454 = E. mexicana.

PACHYTHRIX, Hook. f. Fl. Antarct. i. 32, in syn. (1844) = **Pleurophyllum**, Hook. f. (Compos.).

PACHYTROPHE, Bur. in DC. Prod. xvii. 234 (1873). *URTICACEAE*, Benth. & Hook. f. iii. 363.
Dimepate, *Bur. l. c.*—Madag.
obovata, *Bur. l. c.* 235.—Madag.

PACOUREA, Hook. f. Fl. Brit. Ind. iii. 624 (1882), sphalm. = seq.

PACOURIA, Aubl. Pl. Gui. i. 268. t. 105 (1775) = **Landolphia**, Beauv. (Apocyn.).
Gudara, Buch.-Ham. ex Wall. Cat. n. 4465 = Willughbeia edulis.
guianensis, Aubl. Pl. Gui. i. 269. t. 105.—Guian.
Roxburghii, Kostel. Allg. Med.-Pharm. Fl. iii. 1070 = Willughbeia edulis.

PACOURINA, Aubl. Pl. Gui. ii. 800. t. 316 (1775).
COMPOSITAE, Benth. & Hook. f. ii. 224.
HAYNEA, Willd. Sp. Pl. iii. 1787 (1803).
MEISTERIA, Scop. Introd. 124 (1777).
PACOURINOPSIS, Cass. in Bull. Soc. Philom. (1817) 151 ; et in Dict. Sc. Nat. xxxvii. 212 (1825).
cirsiifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 30.—Ecuador.
edulis, *Aubl. Pl. Gui.* ii. 800.—Guian.

PACOURINOPSIS, Cass. in Bull. Soc. Philom. (1817) 151 ; et in Dict. Sc. Nat. xxxvii. 212 (1825) = **Pacourina**, Aubl. (Compos.).
dentata, Cass. in Bull. Soc. Philom. (1817) 151 = Pacourina cirsiifolia.
integrifolia, Cass. l. c. = Pacourina edulis.

PADBRUGGEA, Miq. Fl. Ind. Bat. i. i. 150 (1855) = **Millettia**, Wight & Arn. (Legumin.).
dasyphylla, Miq. l. c. = Java.

PADBRUGGIA, Baker, in Hook. f. Fl. Brit. Ind. ii. 110 (1876), sphalm. = praec.

PADIA, Zoll. & Mor. Syst. Verz. Zoll. 103 (1845-46) = **Oryza**, Linn. (Gramin.).
Meyeriana, Zoll. & Mor. l. c. = O. sativa.

PADOSTEMON, Griff. Notal. iv. 376 (1854) err. typ. = **Podostemon**, Michx.

PADOTA, Adans. Fam. ii. 505 (1763) = **Marrubium**, Tourn. (Labiat.).

PADUS, Linn. Syst. ed. I (1735) ; Gen. ed. I. 142 (1737) = **Prunus**, Tourn. (Rosac.).
avium, Mill. Gard. Dict. ed. VIII. n. 1 = Prunus Padus.
canadensis, M. Roem. Syn. Rosifl. 88 = Prunus americana ?
Capollin, M. Roem. l. c. 87 = Prunus Capollin.
Capulinos, Hamel. in Rev. Hort. (1884) 111 = praec.
Carolina, Borck. in Roem. Arch. i. II. 38 = seq.
caroliniana, Mill. Gard. Dict. ed. VIII. n. 6 = Prunus caroliniana.
cartilaginea, M. Roem. Syn. Rosifl. 86 = Prunus serotina.
cornuta, Carr. in Rev. Hortic. (1869) 275. f. 64 = Prunus Padus.
demissa, M. Roem. Syn. Rosifl. 87 = Prunus demissa.
densiflora, M. Roem. l. c. 84 = Prunus virginiana.
Duerinchii, M. Roem. l. c. = Prunus virginiana.
eglandulosa, Moench, Meth. 672 = Prunus lusitanica.
elliptica, M. Roem. Syn. Rosifl. 88 = Prunus elliptica.
fimbriata, M. Roem. l. c. 84 = Prunus virginiana.
germanica, Borck. in Roem. Arch. i. II. 38 = Prunus Padus.
glaucifolia, M. Roem. Syn. Rosifl. 88 = Prunus nepaulensis.
hirsuta, M. Roem. l. c. 87 = Prunus virginiana.
Laurocerasus, Mill. Gard. Dict. ed. VIII. n. 4 = Prunus Laurocerasus.
lusitanica, Mill. l. c. n. 5 = Prunus lusitanica.
Mahaleb, Borck. in Roem Arch. i. II. 38 = Prunus Mahaleb.
micrantha, M. Roem. Syn. Rosifl. 84 = Prunus micrantha.
nana, Borck. in Roem. Arch. i. II. 38 = Prunus Padus.
oblonga, Moench, Meth. 672 = Prunus virginiana.
obovata, M. Roem. Syn. Rosifl. 86 = Prunus virginiana.
oxypyrena, M. Roem. l. c. 84 = Prunus Padus.
petraea, M. Roem. l. c. 87 = Prunus Padus.
racemosus, Gilib. Fl. Lituan. ii. 231 = Prunus Padus.
retusa, M. Roem. Syn. Rosifl. 85 = Prunus serotina ?
rubra, Mill. Gard. Dict. ed. VIII. n. 2 = Prunus virginiana.
serotina, Borck. in Roem. Arch. i. II. 38 = Prunus serotina.
virginalis, M. Roem. Syn. Rosifl. 86 = Prunus virginiana.
virginiana, Mill. Gard. Dict. ed. VIII. n. 3 = Prunus serotina.
virginiana, M. Roem. Syn. Rosifl. 86 = Prunus virginiana.
virginica, Borck. in Roem. Arch. i. II. 38 = Prunus virginiana.
vulgaris, Borck. Forstbot. ii. 1426 = Prunus Padus.

PAEDERIA, Linn. Mant. i. 7, 52 (1767). *RUBIACEAE*, Benth. & Hook. f. ii. 133.
HONDBESSEN, Adans. Fam. ii. 158 (1763).
REUSSIA, Dennst. Schluess. Hort. Malab. 33 (1818).
aestivalis, Buch.-Ham. ex Wall. Cat. n. 6247 = foetida.
amboinensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 254. —Ins. Amboin.
argentifolia, Boj. ex A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 195 = Lecontea argentea.
autumnalis, Buch.-Ham. ex Wall. Cat. n. 6247 = foetida.
barbulata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 255 = tomentosa.
brasiliana, *DC. Prod.* iv. 472.—Bras.
calycina, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 74.—Burma.
chinensis, Hance, in Journ. Bot. xvi. (1878) 228 = tomentosa.
corymbosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 = foetida.
densiflora, *Miq. Fl. Ind. Bat.* ii. 259.—Malaya.
erecta, *Roxb. Hort. Beng.* 15 ; *Fl. Ind.* i. 685 (Pyschotriae sp. ?).—Ind. or.
erecta, Spreng. Neue Entdeck. iii. 34 = brasiliana.
foetida, *Linn. Mant.* i. 52.—Ind. or. ; Malaya.
foetida, Thunb. Fl. Jap. 106 = tomentosa.
foetida, Wall. Cat. n. 6247 F = calycina.
fragrans, Lam. Encyc. ii. 260 = Landolphia owariensis.
Gardneri, *Hook. f. in Benth. & Hook. f. Gen.* ii. 184. —Bras.
lanuginosa, Wall. Cat. n. 7293 = Wallichii.
lanuginosa, *Wall. Pl. As. Rar.* ii. 52. t. 165.—Burma.
linearis, *Hook. f. Fl. Brit. Ind.* iii. 197.—Burma.
Lingua, Boj. ex A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 195 = Lecontea Bojeriana.
macrocarpa, Wall. Cat. n. 7292 = lanuginosa.
magnifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 = tomentosa.
malayana, F. Villar, in Blanco, Fl. Filip. ed. III. Nov. App. 112 = Psychotria Tacpo.
ovata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 255 = foetida.
owariensis, Spreng. Syst. i. 669 = Landolphia owariensis.
pilifera, *Hook. f. Fl. Brit. Ind.* iii. 196.—Burma.
recurva, Roxb. Hort. Beng. 15 ; Fl. Ind. i. 684 = Ixora recurva.
Roxburghii, Spreng. Syst. iv. Cur. Post. 70 = erecta.
secundiflora, Poir. Encyc. Suppl. ii. 450 = Manettia secundiflora.
sessiliflora, Poir. l. c. 449 = foetida.

PAEDERIA :—
Tacpo, Blanco, Fl. Filip. ed. I. 160 = Psychotria Tacpo.
ternata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 520 = Saprosma ternata.
tetrandra, Wall. Cat. n. 6249 = Gynochthodes sublanceolata.
Thouarsiana, *Baill. Adansonia*, xii. (1879) 233.—Madag.
tomentosa, *Blume, Bijdr.* 968.—As. trop.
Valli-kara, Juss. in Mém. Mus. Par. vi. (1820) 381 = tomentosa.
verticillata, *Blume, Bijdr.* 968.—Malaya.
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 196.—Burma.

PAEDEROTA, Linn. Gen. ed. I. 4 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 964.
BONAROTA, Adans. Fam. ii. 209 (1763).
Ageria, *Linn. Mant.* i. 171.—Europ.
Amherstiana, Wall. Cat. n. 410 = Wulfenia Amherstiana.
angustifolia, *Turcz. ex Bess. in Flora*, xvii. (1834) i. Beibl. 21.—Sibir. baical.
axillaris, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 144.—Japon.
Bonae-spei, Linn. Sp. Pl. ed. II. 20 = Diascia Bergiana.
Bonarota, Jacq. Hort. Vindob. ii. t. 121 = P. Ageria.
Bonarota, *Linn. Sp. Pl.* ed. II. 20.—Europ.
borealis, Lam. ex Choisy, in DC. Prod. xii. 25 = Lagotis glauca.
bracteata, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 145 = villosula ?
caerulea, Scop. Ann. II. Hist. Nat. 41 = P. Bonarota.
chamaedrifolia, Steud. Nom. ed. II. ii. 246, sphalm. = *cochlearifolia*.
chamaedrifolia, Suffrén, Princ. Bot. 108 = P. Bonarota.
cochlearifolia, Koen. ex Rottl. in Wall. Cat. n. 3920 = Peplidium humifusum.
densifolia, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 107 = Veronica densifolia.
humilis, Willd. ex Link, Jahrb. i. III. (1820) 45 = Veronica densiflora.
lutea, Linn. f. Suppl, 84 = P. Ageria.
minima, Koen. ex Retz. Obs. v. 10 = Microcarpaea muscosa.
minima, Wight, ex Wall. Cat. n. 2488 ; Retz. fide Herb. Sm. ex Benth. in DC. Prod. x. 426 = Glossostigma spathulatum.
nudicaulis, Lam. Illustr. i. 48 ; Savign. in Lam. Encyc. iv. 693 = Wulfenia carinthiaca.
obliqua, A. Dietr. Sp. Pl. i. 563 = Rhinchoglossum obliquum.
pontica, *Rupr. ex Boiss. Fl. Orient.* iv. 434.—Oriens.
racemosa, Houtt. Handleid. vii. 110 = Hemimeris montana.
sibirica, Walp. Rep. iii. 369 = Veronica virginica.
tubiflora, Walp. l. c. 370 = Veronica tubiflora.
urticaefolia, Brign. Fasc. Pl. Forojul. 8 = P. Ageria.
variegata, Raeusch. Nom. ed. III. 5 (Quid ?).—Maurit.
villosula, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 118.—Japon.
virginica, Torr. Fl. N. York, ii. 44 = Veronica virginica.
Wulfenia, Willd. Sp. Pl. i. 76, in syn. = Wulfenia carinthiaca.
Zannichellii, Brign. Fasc. Pl. Forojul. 9 = P. Ageria.

PAENNAEA, Meerb. Pl. Rar. Viv. (1789) Desc. ad t. 51 = **Penaea**, Linn.

PAEONIA, [Tourn.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 157 (1737). *RANUNCULACEAE*, Benth. & Hook. f. i. 10.
MOUTAN, Reichb. in Moessl, Handb. ed. II. i. p. lx. (1827).
albiflora, *Pall. Fl. Ross.* i. II. 92.—Sibir.
albiflos, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 369 = microcarpa.
anomala, *Linn. Mant.* ii. 247.—Sibir.
arborea, Donn, Cat. Hort. Cantab. ed. III. 102 = P. Moutan.
arietina, Anders. in Trans. Linn. Soc. xii. (1817) 275 = peregrina.
banatica, Rochel, Pl. Banat. Rar. 48. t. 11 = officinalis.
Baxteri, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 370 = peregrina.
Biebersteiniana, *Rupr. Fl. Cauc.* 47, 288.—Reg. Caucas.

PAEONIA:—
Broteri, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 4 = corallina.
Brownii, *Dougl. in Hook. Fl. Bor. Am.* i. 27.—Am. bor. occ.
byzantina, Hort. ex DC. Prod. i. 66 = decora.
californica, Nutt. ex Torr. & Gray, Fl. N. Am. i. 41 = Brownii.
Cambessedesii, Willk. Illustr. Fl. Hisp. t. 65 a = corallina.
chinensis, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 892 = albiflora.
commutata, Wender. in Flora, xii. (1829) i. Erg. 29 = officinalis.
corallina, *Retz. Obs.* iii. 34.—Europ.; As. Min.
coriacea, *Boiss. Elench.* 7.—Hispan.
corsica, Sieber, ex Tausch, in Flora, xi. (1828) 88 = corallina.
cretica, Sabine, in Bot. Reg. t. 819 = peregrina.
daurica, Anders. in Bot. Mag. t. 1441 = corallina.
decora, *Anders. in Trans. Linn. Soc.* xii. (1817) 273.—Thrac.; As. Min.
edulis, Salisb. Parad. Lond. t. 78 = albiflora.
elegans, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 370 = officinalis.
Emodi, Wall. Cat. n. 4727 = anomala.
festa, Hort. ex Steud. Nom. ed. II. ii. 247 = seq.?
festiva, Tausch, in Flora, xi. (1828) 84 = officinalis.
fimbriata, *Hort. Angl. ex Sabine, in Trans. Hort. Soc.* ii. (1817) 276.—Oriens.
flavescens, Presl, Delic. Prag. 5 = corallina.
foemina, Mill. Gard. Dict. ed. VIII. n. 2 = corallina.
foemina, Vilm. Fl. Pl. Terre, ed. III. 886 = officinalis.
foliosa, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 370 = peregrina.
fragrans, Vilm. Fl. Pl. Terre, ed. III. 892 = albiflora.
frutescens, Link, Enum. Hort. Berol. ii. 77, sphalm. = seq.
fruticosa, Dum.-Cours. Bot. Cult. ed. II. iv. 462 = P. Moutan.
fulgens, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 648 = officinalis.
fulgida, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 370 = officinalis.
Hartwissiana, Hort. ex Trautv. in Act. Hort. Petrop. viii. (1883) 61, nomen = Wittmanniana.
hirsuta, Mill. Gard. Dict. ed. VIII. n. 4 = peregrina.
Humei, Vilm. Fl. Pl. Terre, ed. III. 892 = albiflora.
humilis, Retz. Obs. iii. 35 = peregrina.
humilis, Willd. Sp. Pl. ii. 1222; Enum. Hort. Berol. 572 = paradoxa.
hybrida, Pall. Fl. Ross. i. II. 94 = tenuifolia.
integra, Murr. in Comm. Gotting. vii. (1786) 92 = corallina.
integrifolia, Link, Enum. Hort. Berol. ii. 77, sphalm. = praec.
intermedia, C. A. Mey. in Ledeb. Fl. Alt. ii. 277 = anomala.
laciniata, Pall. Reise, ii. 244 = tenuifolia.
laciniata, Siev. in Pall. N. Nord. Beitr. vii. 346 = anomala.
lactea, Pall. Reise, iii. 321 = albiflora.
lactiflora, Pall. l. c. 286 = albiflora.
lanceolata, Salm-Dyck, Hort. Dyck. (1834) 367 = corallina.
lobata, Boiss. Voy. Bot. Espagne, ii. 15 = corallina.
lobata, Desf. Tabl. ed. I. 126 = officinale.
lobata, Pall. Reise, ii. 553 = anomala.
lobata, Reichb. Ic. Fl. Germ. iv. f. 4741 = peregrina.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 6 = corallina?
lusitanica, Tausch, in Flora, xi. (1828) 86 = officinalis.
Makoya, Marn. in Loud. Hort. Brit. Suppl. iii. 601 = albiflora.
mascula, Desf. Tabl. ed. I. 126 = corallina.
mascula, Mill. Gard. Dict. ed. VIII. n. 1 = officinalis.
microcarpa, Boiss. & Reut. Pugill. Pl. Nov. (1852) 3 = officinalis.
microcarpa, Salm-Dyck, Hort. Dyck. (1834) 368 = corallina.
mollis, *Anders. in Trans. Linn. Soc.* xii. (1817) 282 (sp. dub.).—Sibir.
Moutan, *Sims, Bot. Mag.* t. 1154.—China.
multifida, Gueldenst. Reisen, ii. 19 = tenuifolia.
multifida, Salm-Dyck, Hort. Dyck. (1834) 369 = peregrina.
nemoralis, Salisb. Prod. 375 = officinalis.
obovata, *Maxim. Prim. Fl. Amur.* 29.—Manchuria.

PAEONIA:—
officinalis, Falk, Beitr. ii. 198; Hook. f. & Thoms. Fl. Ind. i. 60 = anomala.
officinalis, Gouan, Fl. Monsp. 266 = peregrina.
officinalis, Hayne, Arzneigew. v. t. 26 = paradoxa.
officinalis, *Linn. Sp. Pl.* 530.—Europ.
officinalis, Pall. Fl. Ross. ii. 93; Brot. Fl. Lusit. ii. 299 = corallina.
officinalis, Thunb. Fl. Jap. 230 = P. Moutan.
oreogeton, S. Moore, in Journ. Linn. Soc. xvii. (1879) 376 = Wittmanniana.
pallens, Sims, ex Boiss. Fl. Orient. i. 1000 (Index), sphalm. = *pubens*.
papaveracea, Andr. Bot. Rep. t. 463 = P. Moutan.
paradoxa, *Anders. in Trans. Linn. Soc.* xii. (1817) 280.—Europ. austr.
paradoxa, Tenore, Syll. Fl. Neap. 261 = officinalis.
peregrina, *Mill. Gard. Dict.* ed. VIII. n. 3.—Oriens.
peregrina, Tenore, Viagg. Abruzzi, 71 = paradoxa.
porrigens, Reichb. Fl. Germ. Excurs. 752 = officinalis.
promiscua, Reichb. l. c. 751; Ic. Fl. Germ. iv. 28. t. 126 = corallina.
promiscua, Tausch, in Flora, xi. (1828) 83 = officinalis.
pubens, Sims, Bot. Mag. t. 2264 = officinalis.
pullens, Boiss. Fl. Orient. i. 97, sphalm. = praec.
quinquecapsularis, Pall. Reise, iii. 316 = anomala.
Reevesiana, Loud. Hort. Brit. Suppl. iii. 601 = albiflora.
romanica, Brandza, Prod. Fl. Romane, 38 = decora.
rosea, Host, Fl. Austr. ii. 64 = officinalis.
rotundifolia, Hort. ex Steud. Nom. ed. II. ii. 247 = corallina.
rubra, Hort. ex Steud. l. c. = officinalis.
Russi, Amo, Fl. Iber. 746 = coriacea.
Russi, Bivona, Man. Fl. Sic. iv. 12 = corallina.
sessiliflora, Sims, Bot. Mag. t. 2648 = officinalis.
sibirica, Pall. Fl. Ross. Ic. t. 85 = anomala.
sinensis, Hort. ex Steud. Nom. ed. II. ii. 247 = albiflora.
splendens, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 648 = officinalis.
splendens, Sabine, ex Salm-Dyck, Hort. Dyck. (1834) 370 = peregrina.
subternata, *Salm-Dyck, l. c.* 367.—Hab.?
suffruticosa, Andr. Bot. Rep. v. t. 373 = P. Moutan.
tatarica, Mill. Gard. Dict. ed. VIII. n. 5 = peregrina.
tenuifolia, *Linn. Syst.* ed. X. 1079.—Europ. or.; As. bor.
triternata, *Pall. Tabl. Taur.* 52.—Rossia.
versicolor, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 653, nomen = officinalis?
villosa, Desf. Tabl. ed. I. 126 = peregrina.
villosa, Sweet, Hort. Brit. ed. I. 10 = officinalis.
Wittmanniana, *Hartwiss, ex Lindl. Bot. Reg.* (1846) t. 9; *Stev. in Bull. Soc. Nat. Mosc.* xxi. (1848) II. 275.—Reg. Caucas.

PAEPALANTHUS, Mart. in Nov. Act. Nat. Cur. xvii. I. (1835) 13. *ERIOÇAULEAE*, Benth. & Hook. f. iii. 1022.
CLADOCAULON, Gardn. in Hook. Ic. Pl. t. 528 (1843).
DUPATYA, Vell. Fl. Flum. 35 (1825).
LASIOLEPIS, Boeck. in Flora, lxi. (1873) 90.
LIMNOXERANTHEMUM, Salzm. ex Steud. Syn. Pl. Cyp. 281 (1855).
STEPHANOPHYLLUM, Guill. in Deless. Ic. Sel. iii. 61. t. 98, in obs. (1837).
aciphyllus, *Koern. in Mart. Fl. Bras.* iii. I. 438.—Bras.
affinis, *Kunth, Enum. Pl.* iii. 514.—Bras.
albidus, *Gardn. in Hook. Ic. Pl.* t. 525.—Bras.
alpinus, *Koern. in Mart. Fl. Bras.* iii. I. 409.—Am. trop.
alsinoides, *Wright, in Sauv. Fl. Cub.* 163.—Cuba.
amoenus, *Koern. in Mart. Fl. Bras.* iii. I. 316.—Bras.
andicola, *Koern. l. c.* 408.—Am. trop.
androsaceus, *Griseb. Cat. Pl. Cub.* 225.—Cuba.
angustifolius, *Koern. in Mart. Fl. Bras.* iii. I. 424.—Bras.
anomalus, *Koern. l. c.* 458.—Bras.
anthemidiflorus, *Kunth, Enum. Pl.* iii. 535.—Bras.
appressus, *Koern. in Mart. Fl. Bras.* iii. I. 468.—Bras.
aquatilis, Mart. ex Koern. l. c. 495 = Eriocaulon aquatile.
arenarius, *Walp. Ann.* i. 890.—Bras.

PAEPALANTHUS:—
argenteus, *Koern. in Mart. Fl. Bras.* iii. I. 343.—Bras.
argyrolinon, *Koern. l. c.* 374.—Bras.
Armeria, *Mart. ex Koern. l. c.* 377.—Bras.
atrovirens, *Koern. l. c.* 445.—Bras.
bahiensis, *Kunth, Enum. Pl.* iii. 517.—Bras.
bifidus, *Kunth, l. c.* 512.—Bras.
bisulcatus, *Koern. in Mart. Fl. Bras.* iii. I. 436.—Bras.
Blanchetianus, *Koern. l. c.* 422.—Bras.
blepharocnemis, *Mart. ex Koern. l. c.* 376.—Bras.
blepharophorus, *Kunth, Enum. Pl.* iii. 499.—Bras.
Bongardi, *Kunth, l. c.* 519.—Bras.
brachypus, *Kunth, l. c.* 516.—Bras.
brasiliensis, *Walp. Ann.* i. 890.—Bras.
breviscapus, Koern. in Mart. Fl. Bras. iii. I. 370 = zosterifolius.
brizoides, Kunth, Enum. Pl. iii. 534 = gracilis.
bryoides, *Kunth, l. c.* 520.—Bras.
caespititius, *Mart. ex Koern. in Mart. Fl. Bras.* iii. I. 365.—Bras.
caespitosus, *Koern. l. c.* 431.—Bras.
calvus, *Koern. l. c.* 391.—Bras.
canescens, *Koern. l. c.* 381.—Bras.
capillaceus, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1063, nomen.—Am. trop.
capillaris, *Koern. in Mart. Fl. Bras.* iii. I. 367.—Bras.
Capito, *Koern. l. c.* 392.—Bras.
caricifolius, *Gardn. in Hook. Ic. Pl.* t. 524.—Bras.
caryophyllus, Mart. ex Koern. in Mart. Fl. Bras. iii. I. 337 = plumosus.
caulescens, *Kunth, Enum. Pl.* iii. 537.—Bras.
centauroides, *Koern. in Mart. Fl. Bras.* iii. I. 434.—Bras.
chrysanthus, Koern. l. c. 454 = morulus.
ciliatus, *Kunth, Enum. Pl.* iii. 517.—Bras.
circinnatus, *Koern. in Mart. Fl. Bras.* iii. I. 429.—Bras.
Claussenianus, *Koern. l. c.* 323.—Bras.
campactus, *Gardn. in Hook. Ic. Pl.* t. 527.—Bras.
conduplicatus, *Koern. in Mart. Fl. Bras.* iii. I. 414.—Bras.
confusus, *Koern. l. c.* 433.—Bras.
congestus, *Kunth, Enum. Pl.* iii. 505.—Bras.
consanguineus, *Koern. in Mart. Fl. Bras.* iii. I. 411.—Bras.
corymbosus, *Kunth, Enum. Pl.* iii. 509.—Bras.
crassicaulis, *Koern. in Mart. Fl. Bras.* iii. I. 408.—Bras.
crassifolius, *Koern. l. c.* 420.—Bras.
curvifolius, Kunth, Enum. Pl. iii. 522.—Bras.
cuyabensis, Kunth, l. c. 520 = Philodice cuyabensis.
decussus, *Koern. in Mart. Fl. Bras.* iii. I. 318.—Bras.
dendroides, *Kunth, Enum. Pl.* iii. 507.—Am. trop.
densiflorus, *Koern. in Mart. Fl. Bras.* iii. I. 450.—Bras.
densus, *Koern. l. c.* 464.—Bras.
denudatus, *Koern. l. c.* 328.—Bras.
dianthoides, *Mart. in Koern. l. c.* 339.—Bras.
dichotomus, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1064, nomen.—Guian.
dictyophyllus, Mart. ex Koern. in Mart. Fl. Bras. iii. I. 486 = Eriocaulon dictyophyllum.
distichophyllus, *Mart. in Nov. Act. Nat. Cur.* xvii. I. (1835) 23.—Bras.
divaricatus, *Kunth, Enum. Pl.* iii. 515.—Bras.
dubius, *Koern. in Mart. Fl. Bras.* iii. I. 342.—Bras.
Dupatya, *Mart. ex Koern. l. c.* 410.—Bras.
eburneus, *Koern. l. c.* 430.—Bras.
elatus, *Koern. l. c.* 343.—Bras.
elegans, *Kunth, Enum. Pl.* iii. 527.—Bras.
elongatus, Koern. in Mart. Fl. Bras. iii. I. 312 = helichrysoides.
ensifolius, *Kunth, Enum. Pl.* iii. 501.—Am. trop.
Erigeron, *Mart. ex Koern. in Mart. Fl. Bras.* iii. I. 390.—Bras.
eriocephalus, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1064 = flavescens.
eriophyllus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. I. 463.—Bras.
exiguus, *Koern. l. c.* 314.—Bras.
falcatus, *Koern. l. c.* 387.—Bras.
falcifolius, *Koern. l. c.* 327.—Bras.
fasciculatus, Koern. l. c. 357 = congestus.
fasciculatus, Kunth, Enum. Pl. iii. 506 = Schraderi.

PAEPALANTHUS:—
fastigiatus, *Koern. in Mart. Fl. Bras.* iii. 1. 386.—Bras.
fertilis, *Koern. l. c.* 448.—Bras.
Fischerianus, Koern. l. c. 455 = nardifolius.
flaccidus, *Kunth, Enum. Pl.* iii. 511.—Bras.
flagellaris, *Kunth, l. c.* 526.—Bras.
flavescens, *Koern. in Mart. Fl. Bras.* iii. 1. 423.—Bras.
flaviceps, *Koern. l. c.* 412.—Bras.
flavidulus, *Kunth, Enum. Pl.* iii. 532.—Am. bor.
fluitans, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 415.—Bras.
foliosus, *Koern. l. c.* 333.—Bras.
Freyreissii, Koern. l. c. 370 = paludosus.
Funckeanus, *Koern. l. c.* 404.—Am. trop.
fuscoater, *Koern. l. c.* 382.—Bras.
Gardnerianus, *Walp. Ann.* i. 889.—Bras.
geniculatus, *Kunth, Enum. Pl.* iii. 521.—Bras.
glanduliferus, Mart. ex Koern. in Mart. Fl. Bras. iii. 1. 464 = eriophyllus.
glareosus, *Kunth, Enum. Pl.* iii. 521.—Bras.
glaucescens, *Koern. in Mart. Fl. Bras.* iii. 1. 391.—Bras.
goyazensis, *Koern. l. c.* 453.—Bras.
gracilis, *Koern. l. c.* 460.—Bras.
gramineus, *Kunth, Enum. Pl.* iii. 523.—Bras.
granatensis, *Koern. in Mart. Fl. Bras.* iii. 1. 403.—Am. trop.
guyanensis, *Klotzsch, ex Schomb. Faun. & Fl. Brit. Gui.* 1064, nomen.—Guian.
helichrysoides, *Kunth, Enum. Pl.* iii. 512.—Bras.
helminthorrhizus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 443.—Bras.
helodes, Mart. ex Koern. l. c. 357 = myocephalus.
heteropeplus, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 238.—Guian.
heterophyllus, *Koern. in Mart. Fl. Bras.* iii. 1. 442.—Bras.
Hilairei, *Koern. l. c.* 332.—Bras.
hirsutus, *Kunth, Enum. Pl.* iii. 530.—Bras.
hispidus, Klotzsch, ex Schomb. Faun. & Fl. Brit. Gui. 1116 = simplex.
homomallus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 340.—Bras.
Humboldtii, *Kunth, Enum. Pl.* iii. 535.—Am. trop.
imbricatus, *Koern. in Mart. Fl. Bras.* iii. 1. 430.—Bras.
incanus, *Koern. l. c.* 380.—Bras.
intermedius, *Koern. l. c.* 371.—Bras.
inundatus, *Koern. l. c.* 468.—Bras.
iridifolius, Kunth, Enum. Pl. iii. 502 = monticola.
ithyphyllus, *Walp. Ann.* i. 890.—Bras.
juniperinus, Kunth, Enum. Pl. iii. 512 = flaccidus.
Kegelianus, *Koern. in Mart. Fl. Bras.* iii. 1. 438.—Bras.
Klotzschianus, *Koern. l. c.* 389.—Bras.
lagopodioides, *Griseb. Cat. Pl. Cub.* 225.—Cuba.
Lamarckii, *Kunth, Enum. Pl.* iii. 506.—Bras.
lanato-albus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 342.—Bras.
lanceolatus, *Koern. l. c.* 396.—Bras.
Langsdorffii, *Koern. l. c.* 338.—Bras.
lanuginosus, *Koern. l. c.* 428.—Bras.
laricifolius, *Gardn. in Hook. Ic. Pl.* t. 524.—Bras.
latifolius, *Koern. in Mart. Fl. Bras.* iii. 1. 344.—Bras.
laxifolius, *Koern. l. c.* 395.—Bras.
laxifolius, Mart. ex Koern. l. c. 494 = Eriocaulon laxifolium.
Leprieuri, *Koern. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 239.—Guian.
leucoblepharus, *Koern. in Mart. Fl. Bras.* iii. 1. 388.—Bras.
leucophaeus, Mart. ex Koern. l. c. 441 = laricifolius.
Liebmannianus, *Koern. l. c.* 444.—Bras.
lingulatus, *Kunth, Enum. Pl.* iii. 522.—Bras.
longifolius, *Koern. in Mart. Fl. Bras.* iii. 1. 333.—Bras.
Lundii, *Koern. l. c.* 385.—Bras.
luxurians, *Koern. l. c.* 418.—Bras.
macaheensis, *Koern. in Kjoeb. Vidensk. Meddel.* (1871) 311.—Bras.
macrocephalus, *Koern. in Mart. Fl. Bras.* iii. 1. 379.—Bras.
macrorrhizus, *Kunth, Enum. Pl.* iii. 524.—Bras.
Martianus, *Koern. in Mart. Fl. Bras.* iii. 1. 325.—Bras.
Maximiliani, Kunth, Enum. Pl. iii. 513 = ramosus.
melaleucus, *Kunth, l. c.* 510.—Bras.

PAEPALANTHUS:—
meridensis, *Klotzsch, ex Koern. in Mart. Fl. Bras.* iii. 1. 407.—Am. trop.
microphyllus, *Kunth, Enum. Pl.* iii. 519.—Bras.
Minarum, *Koern. in Mart. Fl. Bras.* iii. 1. 378.—Bras.
minutulus, *Mart. ex Koern. l. c.* 359.—Bras.
mollis, Kunth, Enum. Pl. iii. 507 = plumosus.
monticola, *Mart. in Nov. Act. Nat. Cur.* xvii. 1. (1835) 10.—Bras.
morulus, *Kunth, Enum. Pl.* iii. 533.—Bras.
mucronatus, *Kunth, l. c.* 523.—Bras.
muscosus, *Koern. in Mart. Fl. Bras.* iii. 1. 348.—Am. trop.
myocephalus, *Mart. ex Koern. l. c.* 356.—Bras.
nardifolius, *Kunth, Enum. Pl.* iii. 532.—Bras.
neglectus, *Koern. in Mart. Fl. Bras.* iii. 1. 368.—Bras.
nigritella, Mart. ex Koern. l. c. 337 = plumosus.
nigro-niveus, *Kunth, Enum. Pl.* iii. 502.—Bras.
nitens, *Kunth, l. c.* iii. 531.—Bras.
nitidus, *Kunth, l. c.* 528.—Bras.
niveus, *Kunth, l. c.* 527.—Bras.
nubigena, *Kunth, l. c.* 538.—Bras.
oblongus, *Koern. in Mart. Fl. Bras.* iii. 1. 446.—Bras.
obtusifolius, *Koern. l. c.* 353.—Bras.
ochrocephalus, *Koern. l. c.* 329.—Bras.
Oerstedianus, *Koern. l. c.* 374.—Bras.
oligocephalus, *Koern. l. c.* 402.—Am. trop.
Ottonis, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1115, nomen = Lamarckii.
ovatus, *Koern. in Mart. Fl. Bras.* iii. 1. 367.—Bras.
oxycnemis, Mart. ex Koern. l. c. 461 = gracilis.
oxyphyllus, *Koern. l. c.* 313.—Bras.
pachyphyllus, *Koern. l. c.* 378.—Bras.
paludosus, *Kunth, Enum. Pl.* iii. 502.—Bras.
pauciflorus, *Koern. in Mart. Fl. Bras.* iii. 1. 398.—Bras.
perpusillus, *Kunth, Enum. Pl.* iii. 503.—Bras.
petraeus, *Koern. in Mart. Fl. Bras.* iii. 1. 403.—Am. trop.
philodicoides, *Koern. l. c.* 469.—Bras.
pilifer, *Kunth, Enum. Pl.* iii. 524.—Bras.
pilosus, *Kunth, l. c.* 518.—Am. trop.
pilulifer, *Koern. in Mart. Fl. Bras.* iii. 1. 425.—Bras.
planifolius, Koern. l. c. 413 = monticola.
plantagineus, Koern. l. c. 369 = zosterifolius.
plantaginoides, *Koern. l. c.* 470.—Am. trop.
Plantago, *Mart. ex Koern. l. c.* 426.—Bras.
plumosus, *Koern. l. c.* 337.—Bras.
polyanthus, *Kunth, Enum. Pl.* iii. 516.—Bras.
polygonus, *Koern. in Mart. Fl. Bras.* iii. 1. 393.—Bras.
polytrichoides, *Kunth, Enum. Pl.* iii. 504.—Bras.
procerus, Klotzsch, ex Schomb. Faun. & Fl. Brit. Gui. 1115, nomen = caulescens.
prolifer, *Koern. in Mart. Fl. Bras.* iii. 1. 417.—Bras.
propinquus, *Koern. l. c.* 418.—Bras.
prostratus, *Koern. l. c.* 350.—Bras.
puberulus, *Koern. l. c.* 410.—Bras.
pubescens, *Koern. l. c.* 384.—Bras.
pulcher, *Koern. l. c.* 452.—Bras.
pullus, *Koern. l. c.* 366.—Bras.
pungens, *Griseb. Cat. Pl. Cub.* 224.—Cuba.
pusillus, *Koern. in Mart. Fl. Bras.* iii. 1. 459.—Bras.
radicans, Mart. ex Koern. l. c. 370 = zosterifolius.
ramosus, *Kunth, Enum. Pl.* iii. 514.—Bras.
reclinatus, *Koern. in Mart. Fl. Bras.* iii. 1. 447.—Bras.
regalis, *Mart. ex Koern. l. c.* 393.—Bras.
Regelianus, *Koern. l. c.* 386.—Bras.
repens, *Koern. l. c.* 371.—Ind. occ.
retusus, *Wright, in Sauv. Fl. Cub.* 164.—Cuba.
Riedelianus, Koern. in Mart. Fl. Bras. iii. 1. 383 = Vauthierianus.
rigidulus, Mart. in Nov. Act. Mat. Cur. xvii. 1. (1835) 15. t. 2. f. 1.—Bras.
rigidus, *Kunth, Enum. Pl.* iii. 515.—Bras.
rosulatus, Mart. ex Koern. in Mart. Fl. Bras. iii. 1. 487 = Eriocaulon rosulatum.
rufulus, *Kunth, Enum. Pl.* iii. 530.—Bras.
rupestris, *Gardn. in Hook. Ic. Pl.* t. 525.—Bras.
Ruprechtianus, *Koern. in Mart. Fl. Bras.* iii. 1. 437.—Bras.

PAEPALANTHUS:—
saxatilis, *Koern. in Mart. Fl. Bras.* iii. 1. 365.—Bras.
saxicola, *Koern. l. c.* 455.—Bras.
Schlechtendalii, *Koern. l. c.* 425.—Bras.
Schlimii, *Koern. l. c.* 405.—Am. trop.
Schnechianus, *Koern. l. c.* 369.—Bras.
Schomburgkii, *Klotzsch, ex Schomb. Faun. & Fl. Brit. Gui.* 1064, nomen.—Guian.
Schraderi, *Koern. in Mart. Fl. Bras.* iii. 1. 360.—Bras.
scirpeus, *Mart. ex Koern. l. c.* 364.—Bras.
sedoides, *Koern. l. c.* 352.—Bras.
selaginoides, *Benth. Pl. Hartw.* 260.—N. Granat.
selaginoides, *Koern. in Mart. Fl. Bras.* iii. 1. 362.—Am. trop.
Sellowianus, *Koern. l. c.* 326.—Bras.
seslerioides, *Griseb. Cat. Pl. Cub.* 224.—Cuba.
sessiliflorus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 361.—Bras.
simplex, *Miq. Stirp. Surin. Sel.* 222.—Guian.
spadiceus, *Koern. in Mart. Fl. Bras.* iii. 1. 452.—Bras.
spathulatus, *Koern. l. c.* 389.—Bras.
speciosus, *Gardn. in Hook. Ic. Pl.* t. 512.—Bras.
speciosus, *Koern. in Mart. Fl. Bras.* iii. 1. 315.—Bras.
spiralis, *Koern. l. c.* 420.—Bras.
Spixianus, Mart. in Nov. Act. Nat. Cur. xvii. (1835) 1. 14. t. 1 = Eriocaulon quinquangulare?
splendens, Mart. ex Koern. in Mart. Fl. Bras. iii. 1. 466 = caulescens.
stellaris, Kunth, Enum. Pl. iii. 525 = plumosus.
Steudelianus, *Koern. in Mart. Fl. Bras.* iii. 1. 450.—Bras.
strictus, *Koern. l. c.* 319.—Bras.
subtilis, *Miq. Stirp. Surin. Sel.* 221.—Bras.
subulatus, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1116, nomen = subtilis.
succisus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 384.—Bras.
supinus, *Koern. l. c.* 352.—Bras.
surinamensis, Miq. in Linnaea, xix. (1847) 126 = caulescens.
tenuis, *Kunth. Enum. Pl.* iii. 534.—Am. trop.
tortilis, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 354.—Bras.
triangularis, *Koern. l. c.* 470.—Bras.
trichopetalus, *Koern. l. c.* 399.—Bras.
trichophyllus, *Koern. l. c.* 318.—Bras.
Trinianus, *Walp. Ann.* i. 890.—Bras.
truxillensis, *Koern. in Mart. Fl. Bras.* iii. 1. 406.—Am. trop.
tuberosus, *Kunth, Enum. Pl.* iii. 508.—Bras.
umbellatus, *Kunth, l. c.* 537.—Am. trop.
uncinatus, *Gardn. in Hook. Ic. Pl.* t. 523.—Bras.
vaginatus, *Koern. in Mart. Fl. Bras.* iii. 1. 313.—Bras.
Vauthierianus, *Kunth, Enum. Pl.* iii. 500.—Bras.
vellozioides, *Koern. in Mart. Fl. Bras.* iii. 1. 401.—Bras.
vernonioides, *Kunth, Enum. Pl.* iii. 528.—Bras.
verticillatus, *Kunth, l. c.* 536.—Bras.
villosulus, *Mart. ex Koern. in Mart. Fl. Bras.* iii. 1. 400.—Bras.
viridis, *Koern. l. c.* 355.—Bras.
viviparus, *Mart. ex Koern. l. c.* 417.—Bras.
Wahlbergii, *Koern. l. c.* 459.—Afr. austr.
Weddellianus, *Koern. l. c.* 317.—Bras.
Widgrenianus, *Koern. l. c.* 454.—Bras.
xeranthemoides, *Koern. l. c.* 432.—Bras.
zosterifolius, *Kunth, Enum. Pl.* iii. 504.—Bras.

PAGAEA, Griseb. in DC. Prod. ix. 70 (1845). *GENTIANEAE*, Benth. & Hook. f. ii. 814.
BRACHYCODON, Prog. in Mart. Fl. Bras. vi. 1. 229. t. 42 (1865).
breviflora, *Benth. & Hook. f. Gen.* ii. 814.—Guian.
Poeppigii, *Griseb. in DC. Prod.* ix. 71.—Am. trop.
recurva, *Benth. & Hook. f. Gen.* ii. 814.—Bras.
subcordata, *Benth. & Hook. f. l. c.*—Bras.

PAGAMEA, Aubl. Pl. Gui. i. 112. t. 44 (1775). *LOGANIACEAE*, Benth. & Hook. f. ii. 798.
capitata, *Benth. in Journ. Linn. Soc.* i. (1857) 85, 109.—Guian.
coriacea, *Benth. l. c.* 110.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 113.—Guian.
hirsuta, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1857) 111.—Bras.
macrophylla, *Spruce, l. c.* 110.—Bras.
plicata, *Spruce, l. c.* 109.—Bras.
sessiliflora, *Spruce, l. c.* 110.—Bras.
thyrsiflora, *Spruce, l. c.*—Bras.

PAGERIA, Juss. in Dict. Sc. Nat. iii. 53 (1816) = **Lapageria**, Ruiz & Pav. (Liliac.).

PAGESIA, Rafin. Fl. Ludov. 48 (1817) = **Gerardia**? Linn. (Scrophul.).
leucantha, Rafin. l. c. 49.—Am. bor.

PAGETIA, F. Muell. Fragm. v. 178 (1866). *RUTACEAE*, Benth. & Hook. f. i. 991.
medicinalis, *F. Muell. l. c.*—Austral.

PAHUDIA, Miq. Fl. Ind. Bat. i. 1. 86 (1855). *LEGUMINOSAE*, Benth. & Hook. f. i. 580.
Hasskarliana, *Miq. l. c.* Add. 1080.—Java.
javanica, *Miq. l. c.* 87.—Java.
xylocarpa, *Kurz, For. Fl. Brit. Burma*, i. 413.—Burma.

PAIVA, Vell. Fl. Flum. 104 (1825); iii. t. 16 (1827) = **Sabicea**, Aubl. (Rubiac.).
verticillata, Vell. ll. cc.—Bras.

PAIVAEA, Berg, in Mart. Fl. Bras. xiv. 1. 614 (1857). *MYRTACEAE*, Benth. & Hook. f. i. 712.
Langsdorffi, *Berg, in Mart. l. c.*—Bras.

PAIVAEUSA, Welw. in Trans. Linn. Soc. xxvii. (1869) 20. *BURSERACEAE*, Benth. & Hook. f. i. 993.
dactylophylla, *Welw. l. c.* 21.—Afr. trop.

PAJANELIA, DC. in Bibl. Univ. Genèv. xvii. (1838) 130; (Rev. Bign. 14). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1047.
multijuga, *DC. Prod.* ix. 227.—Ind. or.; Burma.
Rheedii, Wight, Ic. t. 1343 = multijuga.

PALA, Juss. in Ann. Mus. Par. xv. (1810) 346 = **Alstonia**, R. Br. (Apocyn.).

PALAFOXIA, Lag. Gen. et Sp. Nov. 26 (1816) *COMPOSITAE*, Benth. & Hook. f. ii. 405.
Lomaxeta, Rafin. New Fl. Am. iv. 72 (1836).
Paleolaria, Cass. in Bull. Soc. Philom. (1816) 198.
callosa, Torr. & Gray, Fl. N. Am. ii. 369 = Polypteris collina.
fastigiata, Less. Syn. Comp. 156 = Polypteris integrifolia.
Feayi, *A. Gray, in Proc. Am. Acad.* xii. (1877) 59.—Am. bor.
Hookeriana, Torr. & Gray, Fl. N. Am. ii. 368 = Polypteris Hookeriana.
integrifolia, Torr. & Gray, l. c. 369 = Polypteris integrifolia.
latifolia, *DC. Prod.* v. 125 (al. gen.?).—Mexic.
leucophylla, A. Gray, in Proc. Am. Acad. viii. (1870) 291 = linearis.
Lindenii, *A. Gray, Pl. Wright.* i. 120.—Mexic.
linearis, *Lag. Gen. et Sp. Nov.* 26.—Am. bor. occ.; Mexic.
texana, DC. Prod. v. 125 = Polypteris texana.
texana, Hook. Ic. Pl. t. 148 = Polypteris Hookeriana.

PALAMOSTIGMA, Benth. & Hook. f. Gen. iii. 294 (1880), sphalm. = seq.

PALANOSTIGMA, Mart. ex Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 49 = **Croton**, Linn. (Euphorb.).
abutiloides, Baill. Etud. Gén. Euph. 359 = C. abutiloides.
crotonoides, Mart. ex Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 48 = C. Palanostigma.
hibiscifolium, Baill. Etud. Gén. Euph. 359 = C. gossipiifolius.
Martiana, Baill. l. c. = C. Palanostigma.
xalapense, Baill. l. c. = C. xalapensis.

PALAQUIUM, Blanco, Fl. Filip. ed. I. 403 (1837); ed. II. 282 = **Dichopsis**, Thw. (Sapotac.).
acuminatum, Burck, in Ann. Jard. Buitenz. v. (1885) 40.—Sumatra.
amboinense, Burck, l. c. 37.—Ins. Amboin.
argentatum, Pierre, ex Burck, l. c. 30 = D. argentatum.
bancanum, Burck, l. c. 43.—Ins. Banca.
Beauvisagei, Burck, l. c. 38.—Sumatra?
borneënse, Burck, l. c. 26.—Borneo.

PALAQUIUM:—
borneënse, Pierre, in Bull. Soc. Linn. Par. i. (1885) 499.—Borneo.
calophyllum, Pierre, ex Burck, in Ann. Jard. Buitenz. v. (1885) 28 = Bassia caloneura?
celebicum, Burck, l. c. 32.—Ins. Celebes.
cinereum, Burck, l. c. 42.—Sumatra.
cuneatum, Vidal, Sinops. t. 62. f. A = Bassia cuneata.
cupreum, Burck, in Ann. Jard. Buitenz. v. (1885) 42.—Ins. Amboin.
formosum, Pierre, in Bull. Soc. Linn. Par. i. (1885) 498.—Malacca.
gloegoerense, Burck, in Ann. Jard. Buitenz. v. (1885) 40.—Sumatra.
Gutta, Burck, l. c. 24 = D. Gutta.
javense, Burck, l. c. 36. – Java.
lanceolatum, Blanco, Fl. Filip. ed. I. 403 (an seq.?).—Ins. Philipp.
lanceolatum, Burck, in Ann. Jard. Buitenz. v. (1855) 43 = D. lanceolata.
latifolium, Blanco, Fl. Filip. ed. I. 404 = D. latifolia.
latifolium, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 125 = D. luzoniensis.
linggense, Burck, in Ann. Jard. Buitenz. v. (1885) 42.—Ins. Lingga.
Lobbianum, Burck, l. c 29.—Ins. Ternate.
macrocarpum, Burck, l. c. 32.—Sumatra.
malaccense, Pierre, in Bull. Soc. Linn. Par. i. (1885) 498.—Malacca.
membranaceum, Burck, in Ann. Jard. Buitenz. v. (1885) 42.—Sumatra.
Minahassae, Burck, l. c.—Ins. Celebes.
Montgommerianum, Burck, l. c. 34.—Sumatra.
Njatoh, Burck, l. c. 41.—Java.
oblongifolium, Burck, l. c. 25.—Malacca.
obscurum, Burck, l. c. 40.—Sumatra.
obtusifolium, Burck, l. c. 33.—Ins. Celebes.
oleiferum, Blanco, Fl. Filip. 405 = D. oleifera.
Oxleyanum, Burck, in Ann. Jard. Buitenz. v. (1885) 33.—Ins. Banca.
Oxleyanum, Pierre, in Bull. Soc. Linn. Par. i. (1885) 498.—Singapore.
parviflorum, Burck, in Ann. Jard. Buitenz. v. (1885) 37.—Sumatra.
parvifolium, Burck, l. c. 36.—Ins. Banca.
Pierrei, Burck, l. c. 31.—Borneo.
Pisang, Burck, l. c. 41.—Sumatra.
Princeps, Pierre, in Bull. Soc. Linn. Par. i. (1885) 499.—Borneo.
quercifolium, Burck, in Ann. Jard. Buitenz. v. (1885) 41.—Borneo.
rostratum, Burck, l. c. 39.—Ins. Banca.
Selendit, Burck, l. c. 41.—Sumatra.
sumatranum, Burck, l. c. 34.—Sumatra.
Teysmannianum, Burck, l. c. 38.—Sumatra.
Treubii, Burck, l. c. 27.—Ins. Banca.
Verstegii, Burck, l. c. 35.—Ins. Banca.
Vrieseanum, Burck, l. c. 28.—Sumatra.
xanthochymum, Pierre, ex Burck, l. c. 30.—Borneo.

PALAUA, Cav. Diss. 40. t. 11. f. 4, 5 (1785). *MALVACEAE*, [Benth. & Hook. f. i. 200].
declinata, Moench, Meth. 609 = malvaefolia.
dissecta, *Benth. in Journ. Linn. Soc.* vi. (1862) 101.—Peruv.
flexuosa, *Mast. in Gard. Chron.* (1866) 435.—Peruv.
malvaefolia, *Cav. Diss.* 40.—Burma.
moschata, *Cav. l. c.* 41.—Peruv.
prostrata, G. Don, in Loud. Hort. Brit. 290 = moschata.
rhombifolia, *R. Grah. in Edinb. N. Phil. Journ.* (July–Sept., 1830) 369.—Peruv.

PALAUA, Ruiz & Pav. Prod. 100. t. 22 (1794) = **Saurauja**, Willd. (Ternstr.).
biserrata, Ruiz & Pav. Syst. Veg. 181 = S. biserrata.
glabrata, Ruiz & Pav. l. c. 182 = S. serrata.
lanceolata, Ruiz & Pav. l. c. 181 = S. Ruiziana.
scabra, Kunth, Syn. Pl. Aequin. 213 = S. excelsa.
tomentosa, Kunth, l. c. = S. tomentosa.

PALAVA, Juss. Gen. 271 (1789) = **Palaua**, Cav. (Malvac.).

PALAVA, Pers. Syn. ii. 91 (1807) = PALAUA, Ruiz & Pav. = **Saurauja**, Willd. (Ternstr.).

PALAVIA, Schreb. Gen. ii. 464 (1791) = **Palaua**, Cav. (Malvac.).

PALEISTA, Rafin. New Fl. Am. ii. 43 (1836) = **Eclipta**, Linn. (Compos.).
brachypoda, Rafin. l. c. = E. erecta.
procumbens, Rafin. l. c. = E. erecta.

PALENGA, Thw. in Hook. Kew Journ. viii. (1856) 270. t. 7 = **Putranjiva**, Wall. (Euphorb.).
zeylanica, Thw. l. c. 271 = Putranjiva Roxburghii.

PALEOLARIA, Cass. in Bull. Soc. Philom. (1816) 198 = **Palafoxia**, Lag. (Compos.).
carnea, Cass. l. c. = Palafoxia linearis.
fastigiata, Less. Syn. Comp. 156 = Polypteris integrifolia.

PALETUVIERA, Thou. ex DC. Prod. iii. 32 (1828) = **Bruguiera**, Lam. (Rhizoph.).

PALEYA, Cass. in Dict. Sc. Nat. xxxix. 393 (1826) = **Crepis**, Vaill. (Compos.).
albida, Cass. l. c. = C. albida.
billotioides, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 320.—Abyss.
oligocephala, Sch. Bip. l. c.—Ind. or.

PALIAVANA, Vell. ex Vand. Fl. Lusit. 40. t. 3. f. 17 (1788). *GESNERACEAE*, Benth. & Hook. f. ii. 1002.
Codonophora, Lindl. Bot. Reg. sub t. 1110 (1827).
Prasanthea, Decne. in Rev. Hortic. (1848) 467.
prasinata, *Benth. & Hook. f. Gen.* ii. 1003.—Bras.
sericifolia, *Benth. & Hook. f. l. c.*—Bras.

PALICOUREA, Aubl. Pl. Gui. i. 172. t. 66 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 125.
Colladonia, Spreng. Syst. i. 516 (1825).
Nonatelia, Aubl. Pl. Gui. i. 182. t. 70 (1775).
Oribasia, Schreb. Gen. 123 (1789).
Rhodostoma, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 286.
Stephanium, Schreb. Gen. 124 (1789).
alpina, *DC. Prod.* iv. 528.—Ind. occ.; Am. austr.
amethystina, *DC. l. c.* 527.—Peruv.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 367.—Venezuela.
apicata, *H. B. & K. l. c.* t. 285.—Venezuela.
aurantiaca, Miq. Stirp. Sel. Surin. 181 = crocea.
aurata, Mart. Reise Bras. i. 544 = Psychotria rigida.
barbinervia, *DC. Prod.* iv. 530.—Am. austr.
Blanchetiana, Schlecht. in Linnaea, xxviii. (1856) 531 = Psychotria Blanchetiana.
brasiliensis, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 70 = Psychotria Muelleriana.
breviflora, Schlecht. in Linnaea, xxviii. (1856) 530 = Psychotria breviflora.
cajennensis, Ham. Prod. Pl. Ind. Occ. 29 = pedunculosa.
calophylla, *DC. Prod.* iv. 529.—Guian.
calycina, *Benth. Pl. Hartw.* 133.—Peruv.
cayennensis, DC. Prod. iv. 526, in syn. = praec.
chionantha, DC. l. c. 526 = Luschnathii.
coccinea, *DC. l. c.* 529 (= crocea?).—Ins. S. Doming.
coerulea, *Roem. & Schult. Syst.* v. 194.—Peruv.
costa-ricensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 38.—Am. centr.
costata, *H. B. & K. Nov. Gen. et Sp.* iii. 366.—Venezuela.
crocea, *Roem. & Schult. Syst.* v. 193.—Ind. occ.; Am. austr.
crocea, Schlecht. in Linnaea, xxviii. (1856) 525 = Psychotria subcrocea.
croceoides, *Ham. Prod. Pl. Ind. Occ.* 29 (= crocea?).—Ind. occ.
cujabensis, Schlecht. in Linnaea, xxviii. (1856) 525 = Psychotria subcrocea.
densiflora, Mart. in Flora, xxi. (1838) ii. Beibl. 85 = Rudgea viburnoides.
densiflora, Wawra, in Oestr. Bot. Zeitschr. xiii. (1863) 220 = Psychotria longepedunculata.
didymocarpa, *Griseb. Cat. Pl. Cub.* 138.—Cuba.
diuretica, Mart. Reise Bras. i. 544 = Psychotria rigida.
divaricata, Schlecht. in Linnaea, xxviii. (1856) 529 = Psychotria Schlechtendaliana.
domingensis, *DC. Prod.* iv. 529.—Ins. S. Doming.

PALICOUREA:—
elliptica, *H. B. & K. Nov. Gen. et Sp.* iii. 368.—Venezuela.
eriantha, *DC. Prod.* iv. 528.—Ins. S. Doming.
fastigiata, Cham. & Schlecht. in Linnaea, iv. (1829) 16 = Psychotria Sellowiana.
fastigiata, *H. B. & K. Nov. Gen. et Sp.* iii. 368.—Venezuela.
flavescens, *H. B. & K. l. c.* 366.—Peruv.
Galeottiana, *Martens, in Bull. Acad. Brux.* xi. 1. (1844) 136.—Mexic.
gardenioides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 52.—Mexic.
guianensis, *Aubl. Pl. Gui.* i. 173. t. 66.—Ind. occ; Am. austr.
Haenkeana, *DC. Prod.* iv. 530.—Peruv.
hebeantha, DC. l. c. 525 = Psychotria Marcgravii.
hyacinthiflora, *Roem. & Schult. Syst.* v. 194.—Peruv.
intermedia, *Oerst. ex Polak. in Linnaea*, xli. (1877) 571.—Costa Rica.
laevigata, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 133.—Am. austr.
lanceolata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 39.—Costa Rica.
lasiorhachis, *Benth. ex Oerst. l. c.* 39.—Costa Rica.
laxa, *Roem. & Schult. Syst.* v. 195.—Peruv.
lineata, *Benth. Pl. Hartw.* 192.—Ecuador.
longepedunculata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 109 = Psychotria longepedunculata.
longibracteata, *DC. Prod.* iv. 527.—Panama.
longiflora, *DC. l. c.* 528.—Guian.
longifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 369.—N. Granat.
longifolia, A. St. Hil. Pl. Rem. Brés. 232. t. 22 = Psychotria verticillata.
macrobotrys, *Roem. & Schult. Syst.* v. 184.—Peruv.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* iii. 369.—N. Granat.
Marcgravii, A. St. Hil. Pl. Rem. Brés. 231. t. 22 = Psychotria Marcgravii.
metallica, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 207 = Psychotria Blanchetiana.
mexicana, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 37 = alpina.
mitis, *DC. Prod.* iv. 527.—Peruv.
nicotianaefolia, Cham. & Schlecht. in Linnaea, iv. (1829) 18 = Psychotria tabacifolia.
nigrescens, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 136.—Am. centr.
noxia, Mart. Reise Bras. i. 544 = Psychotria Marcgravii.
obovata, *DC. Prod.* iv. 527.—Peruv.
officinalis, *Mart. Reise Bras.* i. 544.—Bras.
parviflora, *Benth. Bot. Voy. Sulph.* 107.—Peruv.
Pavetta, *DC. Prod.* iv. 525.—Ind. occ.
pedunculosa, *DC. l. c.* 526.—Guian.
pedunculosa, Miq. in Linnaea, xix. (1847) 444 = Psychotria longepedunculata.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* iii. 370.—Mexic.
popayanensis, *Benth. Pl. Hartw.* 193.—N. Granat.
pulchra, *Griseb. Fl. Brit. W. Ind.* 345.—Jamaic.
punicea, *DC. Prod.* iv. 526.—Peruv.
quadrifolia, *Rudge, Pl. Gui.* 27. t. 42.—Guian.
rigida, *H. B. & K. Nov. Gen. et Sp.* iii. 370.—Am. austr.
riparia, Benth. in Hook. Journ. Bot. iii. (1841) 224 = crocea.
Sellowiana, DC. Prod. iv. 525 = Psychotria Sellowiana.
sonans, *Mart. Reise Bras.* i. 544.—Bras.
speciosa, *H. B. & K. Nov. Gen. et Sp.* iii. 368.—N. Granat.
stipularis, *Benth. Pl. Hartw.* 133.—Ecuador.
strepens, Mart. Reise Bras. i. 544 = rigida.
subrubra, *Polak. in Linnaea*, xli. (1877) 571.—Costa Rica.
sulphurea, *DC. Prod.* iv. 528.—Peruv.
tabernaefolia, *DC. l. c.* 525.—Ins. S. Doming.
tetraphylla, Cham. & Schlecht. in Linnaea, iv. (1829) 17 = Psychotria tetraphylla.
thyrsiflora, *DC. Prod.* iv. 528.—Peruv.
tinctoria, *Roem. & Schult. Syst.* v. 194.—Peruv.
tinifolia, *DC. Prod.* iv. 529.—Am. austr.
triphylla, DC. l. c. 526 = Psychotria triphylla.
umbellata, *DC. l. c.*—Ind. occ.; Guian.
vaginata, *Benth. Pl. Hartw.* 193.—N. Granat.

PALICOUREA:—
velutina, Schlecht. in Linnaea, xxviii. (1856) 528 = Psychotria Marcgravii.
verticillata, DC. Prod. iv. 526 = Psychotria verticillata.
violacea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 175.—Guian.
viridis, Roem. & Schult. Syst. v. 195 = Psychotria viridis.
xanthina, *DC. Prod.* iv. 525.—Guian.

PALICUREA, Roem. & Schult. Syst. v. 193 (1819) = praec.

PALICURIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 86 = praec.

PALIMBIA, Bess. Enum. Pl. Volh. 55, 94 (1822) = **Peucedanum**, Tourn. (Umbellif.).
alpigenum, C. Koch, ex Boiss. Fl. Orient. ii. 102 = Peuced. alpigenum.
carvifolia, Koch, ex DC. Prod. iv. 176 = Peuced. carvifolium.
Chabraei, Bertol. Fl. Ital. iii. 354, partim = Peuced. Schottii.
Chabraei, DC. Prod. iv. 176 = Peuced. carvifolium.
chrysantha, C. Koch, ex Boiss. Fl. Orient. ii. 1021 = Peuced. palimbioides.
decussata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 30 = Peuced. carvifolium.
ramosissima, DC. Prod. iv. 176 = Peuced. ramosissimum.
ramosissima, Thw. Enum. Pl. Zeyl. 131 = Peuced. ceylanicum.
salsa, Bess. Enum. Pl. Volh. 55, 94 = Peuced. redivivum.

PALINETES, Salisb. Gen. Pl. Fragm. 116 (1866) = **Ammocharis**, Herb. (Amaryll.).
coranica, Salisb. l. c. = A. falcata.
falcata, Salisb. l. c. = A. falcata.

PALIRIS, Dum. Fl. Belg. 134 (1827) = **Liparis**, Rich. (Orchid.).
Loeselii, Dum. l. c. = L. Loeselii.

PALISOTA, Reichb. Consp. 59 (1828). *COMMELINACEAE*, Benth. & Hook. f. iii. 847.
DUCHEKIA, Kostel. Allg. Med.-Pharm. Fl. i. 213 (1831).
ambigua, *C. B. Clarke, in DC. Monog. Phan.* iii. 131.—Afr. trop.
Barteri, *Hook. f. in Bot. Mag.* t. 5318.—Afr. trop.
bicolor, *Mast. in Gard. Chron.* (1878) 527.—Afr. trop.
bracteosa, *C. B. Clarke, in DC. Monog. Phan.* iii. 133.—Afr. trop.
Mannii, *C. B. Clarke, l. c.* 132.—Afr. trop.
prionostachys, *C. B. Clarke, l. c.* 134.—Afr. trop.
Schweinfurthii, *C. B. Clarke, l. c.* 132.—Afr. trop.
thyrsiflora, *Benth. in Hook. Niger Fl.* 544.—Afr. trop.
thyrsoidea, Baker, in Journ. Linn. Soc. xiv. (1875) 518 = thyrsiflora.

PALISSYA, Baill. Etud. Gén. Euph. 502 (1858) = **Alchornea**, Sw. (Euphorb.).
castaneaefolia, Baill. l. c. 503 = A. madagascariensis.

PALIUROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Paliurus**, Tourn. (Rhamn.).
Aubletius, St. Lag. l. c. = Paliurus ramosissima.

PALIURUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752). *RHAMNEAE*, Benth. & Hook. f. i. 375.
AUBLETIA, Lour. Fl. Cochinch. 283 (1790).
aculeatus, *Lam. Illustr.* ii. 347 n. 611.—Europ. austr.; As. occ.
asiaticus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 25 (Zizyphi sp. ?).—Java.
Aubletia, Roem. & Schult. Syst. v. 343 = ramosissimus.
australis, Gaertn. Fruct. i. 23. t. 43 = aculeatus.
clypeiformis, Medic. Beobacht. (1782) 327 = aculeatus.
Dao, Blanco, Fl. Fllip. ed. I. 174 = Dracontomelon mangiferum.
dubius, Blanco, l. c. 175 = Harrisonia paucijuga.
edulis, Blanco, l. c. 173 = Dracontomelon mangiferum.
inermis, Hort. Par. ex DC. Prod. ii. 31 = Ceanothus reclinatus.
Lamujo, Blanco, Fl. Filip. ed. II. 122 = Dracontomelon mangiferum.
lucidus, *Carr. in Rev. Hortic.* (1866) 380.—China.

PALIURUS:—
perforatus, Blanco, Fl. Filip. ed. I. 174 = Harrisonia paucijuga.
Petasus, Dum.-Curs. Bot. Cult. ed. II. vi. 266 = aculeatus.
ramosissimus, *Poir. Encyc. Suppl.* iv. 262.—China.
reticulatus, Vahl, Eclog. Am. iii. 6 = Zizyphus reticulata.
solitarius, Stokes, Bot. Mat. Med. ii. 169 = aculeatus.
Spina-Christi, Mill. Gard. Dict. ed. VIII. = aculeatus.
texanus, Scheele, in Linnaea, xxi. (1848) 594 = Zizyphus obtusifolia.
trinervatus, Moench, Meth. 74 = aculeatus.
virgatus, D. Don, in Bot. Mag. t. 2535 = Zizyphus incurva.
vulgaris, Steud. Nom. ed. II. ii. 249 = aculeatus.

PALLADIA, Lam. Illustr. t. 285 (1792) = **Blackwellia**, Gaertn. (genus dubium).
antarctica, Lam. Encyc. iv. 698; et Illustr. ii. 426 = B. antarctica.

PALLADIA, Moench, Meth. 429 (1794) = **Lysimachia**, Linn. (Primul.).
atropurpurea, Moench, l. c. = L. atropurpurea.

PALLASIA, Houtt. Handleid. iv. 382 (1775) = **Calodendrum**, Thunb. (Rutac.).
capensis, [Christm. in] Houtt. Pfl. Syst. iii. 318 = C. capense.

PALLASIA, Klotzsch, in Monatsb. Akad. Berl. (1853) 498 (reimpr. 5). *RUBIACEAE*, Benth. & Hook. f. ii. 48.
Stanleyana, *Klotzsch, l. c.*—Guian.

PALLASIA, L'Hérit. Diss. (1784); ex Ait. Hort. Kew. ed. I. iii. 498 (1789) = **Encelia**, Adans. (Compos.).
dentata, Humb. & Bonpl. Pl. Aequin. ii. 101. t. 111 = Verbesina dentata.
grandiflora, Willd. Sp. Pl. iii. 2261 = E. halimifolia.
halimifolia, L'Hérit. ex Ait. Hort. Kew. ed. I. iii. 498 = E. canescens.
procumbens, Spreng. Syst. iii. 610 = E. parvifolia.
serratifolia, Sm. in Rees, Cycl. xxvi. n. 3 = Verbesina encelioides.

PALLASIA, Linn. f. Suppl. 87 (1781) = **Calligonum**, Linn. (Polygon.).
caspica, Linn. f. Suppl. 252 = C. Pallasia.
comosa, Raeusch. Nom. ed. III. 141 = C. comosum.
polygonoides, Pall. Reise, iii. 530 = C. Pallasia.
Pterococcus, Pall. Fl. Ross. i. II. 70 = C. Pallasia.

PALLASIA, Scop. Introd. 72 (1777) = **Crypsis**, Ait. (Gramin.).

PALLASTEMA, Salisb. Gen. Pl. Fragm. 36 (1866) = **Albuca**, Linn. (Liliac.).
abyssinica, Salisb. l. c. = A. abyssinica.

PALLAVIA, Vell. Fl. Flum. 151 (1825); iv. t. 12 (1827) **Pisonia**, Linn. (Nyctag.).
aculeata, Vell. ll. cc. = Pisonia aculeata.

PALLAVICINIA, De Not. in Flora, xxx. (1847) 567 = **Cyphomandra**, Sendt. (Solan.).
fragrans, De Not. l. c. 568 = C. fragrans.

PALLENIS, Cass. in Dict. Sc. Nat. xxiii. 566 (1822); xxxvii. 275 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 340.
ASTERISCUS, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 229 (1835-60).
ATHALMUS, Neck. Elem. i. 20 (1790).
aurea, Nym. Consp. 391 = spinosa.
aurea, *Pomel, Nouv. Mat. Fl. Atl.* 38.—Alger.
canescens, *Viv. Mem. Alc. Plag.* (1838) 11.—Aegypt.
cuspidata, *Pomel, Nouv. Mat. Fl. Atl.* 38.—Alger.
spinosa, *Cass. in Dict. Sc. Nat.* xxxvii. 275.—Reg. Mediterr.; Oriens.

PALMA, Plum. ex Mill. Gard. Dict. ed. VI (1752) (Farrago Palmarum, etc.).
altissima, Mill. l. c. n. 4 = Elaeis occidentalis.
americana, Mill. l. c. n. 10 (Quid ?).—Ind. occ.
argentea, Jacq. Fragm. 38. t. 43 = Thrinax argentea.
Avoira, Aubl. Pl. Guin. Suppl. 95 (Quid ?).—Guian.
Bache, Aubl. l. c. 103 (Quid ?).—Guian.

PALMA:—
Cocos, Mill. Gard. Dict. ed. VIII. n. 2 = Cocos nucifera.
Comon, Aubl. Pl. Gui. Suppl. 102.—Guian.
dactylifera, Mill. Gard. Dict. ed. VIII. n. 1 = Phoenix dactylifera.
Draco, Mill. l. c. n. 11 = Dracaena Draco.
gracilis, Mill. l. c. n. 5 = Bactris minor.
Maripa, Aubl. Pl. Gui. ii. 974.—Guian.
Mocaia, Aubl. l. c. 976.—Guian.
Mocayba Correa, in Ann. Mus. Par. x. (1807) t. 7 = praec.
oleosa, Mill. Gard. Dict. ed. VIII. n. 6 = Elaeis guineensis?
Paripou, Aubl. Pl. Gui. ii. 974 (Quid?).—Guian.
Patavoua, Aubl. Pl. Gui. Suppl. 102 (Quid?).—Guian.
Pinao, Aubl. Pl. Gui. ii. 974 (Quid?).—Guian.
polypodiifolia, Mill. Gard. Dict. ed. VIII. n. 8 = Cycas circinalis.
prunifera, Mill. l. c. n. 7 = Chamaerops humilis.
pumila, Mill. l. c. n. 9 = Zamia furfuracea.
Sancona, Kunth, Enum. Pl. iii. 182 = Oreodoxa Sancona.
spinosa, Mill. Gard. Dict. ed. VIII. n. 3 = Bactris minor; Elaeis guineensis.
thebaica, Jacq. Fragm. 83 = Hyphaene thebaica.
Zagueneti, Aubl. Pl. Gui. ii. 976 (Quid?).—Guian.

PALMERELLA, A. Gray, in Proc. Am. Acad. xi. (1876) 80. *CAMPANULACEAE*, Benth. & Hook. f. ii. 1238.
debilis, *A. Gray*, *l. c.*—Calif.

PALMERIA, F. Muell. Fragm. iv. 151 (1864); v. 2 (1865). *MONIMIACEAE*, Benth. & Hook. f. iii. 140.
arfakiana, *Becc. Malesia*, i. (1877) 186.—N. Guin.
racemosa, *A. DC. in DC. Prod.* xvi. II. 657.—Austral.
scandens, *F. Muell. Fragm.* iv. 152.—Austral.

PALMIA, Endl. Gen. 653 (1839) = **Hewittia**, Wight & Arn. (Convolv.).
bicolor, Endl. ex Dalz. & Gibs. Bomb. Fl. 163 = H. bicolor.

PALMONARIA, Boiss. Pl. Or. Nov. Dec. ii. 8 (1875), err. typ. = **Pulmonaria**, Tourn. (Borag.).

PALMORCHIS, Rodrig. Orch. Nov. i. 169 (1877). *ORCHIDEAE*.
pubescentis, *Rodrig. l. c.* 170.—Bras.
sobralioides, *Rodrig. l. c.*—Bras.

PALMSTRUCKIA, Retz. f. Obs. Bot. Pugill. 15 (1810) = **Chaenostoma**, Benth. (Scrophul.).
foetida, Retz. f. l. c. = C. foetidum.

PALMSTRUCKIA, Sond. in Harv. & Sond. Fl. Cap. i. 35 (1859). *CRUCIFERAE*, Benth. & Hook. f. i. 96.
capensis, *Sond. l. c.*—Afr. austr.

PALOUE, Aubl. Pl. Gui. 365. t. 141 (1775) = seq.

PALOVEA, Aubl. Pl. Gui. 365. t. 141 (1775) (Paloue); Juss. Gen. 350 (1789). *LEGUMINOSAE*, Benth. & Hook. f. i. 578.
GINANNIA, Scop. Introd. 300 (1777).
guianensis, *Aubl. Pl. Gui.* i. 365. t. 141.—Guian.

PALTORIA, Ruiz & Pav. Prod. 13 (1794) = **Ilex**, Tourn.
ovalis, Ruiz & Pav. Fl. Per. i. 54 = I. Paltoria.

PALUDANA, Giseke, Prael. Linn. 199, 207 (1792) = **Amomum**, Linn. (Scitam.).
crispa, Giseke, l. c. 207 = A. echinatum.

PALUDANA, Salisb. Gen. Pl. Fragm. 53 (1866) 53, in obs. = seq.

PALUDARIA, Salisb. Gen. Pl. Fragm. 53 (1866) = **Colchicum**, Tourn. (Liliac.).

PALUMBINA, Reichb. f. in Walp. Ann. vi. 699 (1861) = **Oncidium**, Sw. (Orchid.).
candida, Reichb. f. l. c. = O. candidum.

PALURA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 145 (1825) = **Symplocos**, Linn. (Styrac.).
odorata, Buch.-Ham. ex D. Don, l. c. = S. crataegoides.
pendula, Miers, in Journ. Linn. Soc. xvii. (1879) 297 = S. pendula.
sinica, Miers, l. c. = S. sinica.

PAMEA, Aubl. Pl. Gui. ii. 946. t. 359 (1775) = **Terminalia**, Linn. (Combret.).
guianensis, Aubl. l. c. = T. Pamea.

PAMPHALEA, DC. in Ann. Mus. Par. xix. (1812) 68. *COMPOSITAE*, Benth. & Hook. f. ii. 503.
CERATOLEPIS, Cass. in Bull. Soc. Philom. (1819) 111.
PANPHALEA, Lag. Amen. Nat. i. 34 (1811).
bupleurifolia, *Less. in Linnaea*, v. (1830) 8.—Bras.
cardaminefolia, *Less. l. c.* 9.—Bras.
Commersonii, *Cass. in Bull. Soc. Philom.* (1819) 111.—Reg. Argent.
heterophylla, *Less. in Linnaea*, v. (1830) 8.—Bras.
maxima, *Less. l. c.* 9.—Bras.
Tweedii, *Gardn. & Field. Sert. Pl.* t. 21.—Uruguay.

PAMPHILIA, Mart. Herb. Fl. Bras. n. 902 (1837). *STYRACEAE*, Benth. & Hook. f. ii. 670.
aurea, *Mart. l. c.*; *et ex A. DC. in DC. Prod.* viii. 271.—Bras.
styracifolia, *A. DC. l. c.*—Bras.

PANACEA, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) App. 221 = **Panax**, Linn. (Araliac.).

PANAETIA, Cass. in Ann. Sc. Nat. Ser. I. xvii. (1829) 417; et in Dict. Sc. Nat. lx. (1830) 593 = **Podolepis**, Labill. (Compos.).
athrixioides, Sond. & F. Muell. in Linnaea, xxv. (1852) 506 = Athrixia tenella.
fulva, Lindl. in Bot. Reg. (1838) Misc. 47.—Austral.
Lessonii, Cass. in Ann. Sc. Nat. Ser. I. xvii. (1829) 417 = Podolepis Lessoni.
Muelleri, Sond. in Linnaea, xxv. (1852) 505 = Podolepis Lessoni.

PANARGYRUM, D. Don, in Phil. Mag. xi. (1832) 390 = seq.

PANARGYRUS, Lag. Amen. Nat. i. 33 (1811) = **Nassauvia**, Juss. (Compos.).
abbreviatus, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 43 = N. abbreviata.
Darwinii, Hook. & Arn. l. c.—Reg. Magellan.
densifolius, Phil. in Linnaea, xxx. (1859–60) 192 = *glomeratum*.
Diazi, Phil. l. c. xxxiii. (1864–65) 117.—Chili.
glomeratus, Gill. ex D. Don, in Phil. Mag. xi. (1832) 390.—Chili.
heteraphyllus, Phil. in Linnaea, xxxiii. (1864–65) 118.—Patagonia.
Lagascae, DC. Prod. vii. 54 = N. Lagascae.
latifolius, Phil. in Linnaea, xxxiii. (1864–65) 119.—Chili.
oligocephalus, DC. Prod. vii. 54.—Chili.
pectinatus, Phil. in Anal. Univ. Chil. (1861) I. 63.—Chili.
spinosus, D. Don, in Phil. Mag. xi. (1832) 390.—Chili.
uniflorus, D. Don, l. c. = *oligocephalus*.

PANAX, Linn. Syst. ed. I (1735). *ARALIACEAE*, Benth. & Hook. f. i. 938.
CHEIRODENDRON, Nutt. ex Seem. in Journ. Bot. v. (1867) 236.
GINSEN, Adans. Fam. ii. 102 (1763).
MARALIA, Thou. Gen. Nov. Madag. 13 (1806).
NOTHOPANAX, Seem. in Journ. Bot. iv. (1866) 293.
OLIGOSCIAS, Seem. l. c. iii. (1865) 179.
PANACEA, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) App. 221.
RAUKANA, Seem. in Journ. Bot. iv. (1866) 352.
SCIADOPANAX, Seem. l. c. iii. (1865) 73. t. 27.
aculeatum, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 448 = Acanthopanax aculeatum.
americanum, Rafin. New Fl. Am. iv. 58 = Aralia quinquefolia.
amplifolium, *Baker, in Journ. Linn. Soc.* xxi. (1884) 351.—Madag.

PANAX:—
angustifolium, F. Muell. in Trans. Phil. Inst. Vict. i. (1855) 42 = sambucifolium.
Anisum, *DC. Prod.* iv. 254.—Ins. Molucc.
anomalum, *Hook. Lond. Journ. Bot.* ii. (1843) 422. t. 12.—N. Zel.
arboreum, *Forst. f. Prod.* 75.—N. Zel.
armatum, Wall. Cat. n. 4933 = Aralia armata.
attenuatum, *Sw. Prod. Veg. Ind. Occ.* 54.—Ind. occ.
australasium, Pers. Syn. i. 298 = arboreum.
australe, *Spreng. ex Schult. Syst.* vi. 215 (sp. dub.).—Austral.
bandanense, Zipp. ex Span. in Linnaea, xv. (1841) 208 = Arthrophyllum pinnatum.
Bastardianum, *Decne. Bot. Voy. Vénus*, 24.—Ins. Tahiti.
bijugum, Wall. Cat. n. 4937 = Pentapanax Leschenaultii.
bipinnatifidum, Seem. in Journ. Bot. vi. (1868) 54 = Aralia bipinnatifida.
Boivini, *Seem. l. c.* iii. (1865) 74.—Madag.
calvum, Cham. in Linnaea, viii. (1833) 332 = Didymopanax calvum.
cephalobotrys, *F. Muell. Fragm.* ii. 83.—Austral.
chrysophyllum, Vahl, Eclog. Am. i 33 = Didymopanax Morototoni.
cissiflorum, *Baker, in Journ. Linn. Soc.* xx. (1883) 154.—Madag.
cissodendron, *C. Moore & F. Muell. Fragm.* vii. 96; viii. 280.—Austral.
cochleatum, *DC. Prod.* iv. 253.—Ins. Malay.
Colensoi, *Hook. f. Fl. N. Zel.* 94. t. 21.—N. Zel.
conchifolium, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 77.—Ind. or.
confertifolium, *Baker, in Journ. Linn. Soc.* xxi. (1884) 351.—Madag.
coriaceum, *Regel, Gartenfl.* (1859) 45.—N. Zel.
crassifolium, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 105 = Pseudopanax crassifolium.
crenatum, Dietr. ex Steud. Nom. ed. II. ii. 249 (Quid?).—Hab.?
crenatum, *Planch. & Sebert, Not. Bois Nouv. Caléd.* 202.—N. Caled.
Cumingianum, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 310.—Ins. Philipp.
curcifolium, Griff. Itin. Notes, 145 = Brassaiopsis Hainla.
decaphyllum, Sagot, ex Seem. in Journ. Bot. iii. (1865) 266 = Sciadophyllum decaphyllum.
decompositum, Wall. Cat. n. 4935 = Aralia cachemirica.
dendroides, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 42 = sambucifolium.
digitatum, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 75.—Ind. or.
discolorum, *T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 178.—N. Zel.
divaricatum, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 200 = Acanthopanax divaricatum.
dumosum, Hort. Bull.—Cf. Gard. Chron. (1883) I. 404 = fruticosum.
Edgerleyi, *Hook. f. Fl. N. Zel.* 94.—N. Zel.
elegans, *C. Moore & F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 68.—Austral.
fallax, *Miq. in Journ. Bot. Néerl.* i. (1861) 118.—China.
ferox, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 493, in obs. = Pseudopanax crassifolium.
ferrugineum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 28.—Afr. trop.
Finlaysonianum, *Wall. Cat.* n. 4936.—Ind. or.
fissum, Bull, Cat. (1884) 14 = fruticosum.
foliolosum, Wall. Cat. n. 4928 = Aralia foliolosa.
Forsteri, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 105 = Polyscias pinnata.
fragrans, Roxb. Hort. Beng. 21; Fl. Ind. ii. 76 = Heteropanax fragrans.
fruticosum, *Linn. Sp. Pl.* ed. II. 1513.—As. trop.; Malay.; Polynes.
fulvum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 28.—Afr. trop.
Gaudichaudii, *DC. Prod.* iv. 253.—Ins. Sandvic.
Gerrardi, *Harv. Gen. S. Afr. Pl.* ed. II. 147.—Afr. austr.
Ginseng, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 340 = Aralia quinquefolia.
glabratum, *H. B. & K. Nov. Gen. et Sp.* v. 10.—Am. austr.
Guilfoylei, Cogn. & March. Pl. Ornament. ii. (1874) t. 58 = Aralia Guilfoylei.

PANAX:—
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 466.—Tasman.
Hainla, DC. Prod. iv. 253 = Brassaiopsis Hainla.
Heyneanum, Wall. Cat. n. 4927 = Polyscias pinnata.
horridum, Sm. in Rees, Cycl. xxvi. n. 10 = Fatsia horrida.
innovans, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 198 = Acanthopanax innovans.
Jackianum, Wall. Cat. n. 4931 = Arthrophyllum diversifolium.
japonicum, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 340 = Aralia quinquefolia.
laciniatum, Hort.—Cf. Gard. Chron. (1876) I. 735; (1880) I. 759 = fruticosum.
lanceolatum, Rafin. New Fl. Am. iv. 57 = Aralia trifolia.
Leschenaultii, DC. Prod. iv. 254 = Pentapanax Leschenaultii.
Lessonii, DC. Prod. iv. 253 = Pseudopanax Lessonii.
lineare, *Hook. f. Fl. N. Zel.* 93.—N. Zel.
longepetiolatum, Pohl, ex DC. Prod. iv. 253 = Didymopanax longepetiolatum.
longissimum, *Hook. f. Handb. N. Zeal. Fl.* 102.—N. Zel.
Loureirianum, DC. Prod. iv. 252 = Acanthopanax aculeatum.
Macgillivraei, *Benth. Fl. Austral.* iii. 382.—Austral.
Macgillivrayi, Seem. Revis. Heder. 101, sphalm. = praec.
macrocarpum, Schlecht. in Linnaea, i. (1826) 404 = Didymopanax macrocarpum.
macrosciadeum, F. Muell. Fragm. ii. 108 = Mackinlaya macrosciadea.
Manguette, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 66.—N. Caled.
Maralia, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 105.—Madag.
margaritiferum, *Vis. Ind. Sem. Hort. Patav.* (1855). —*Cf. Linnaea*, xxviii. (1856) 365.—Hab.?
micranthum, *Wall. Cat.* n. 4938.—Ind. or.
microphyllum, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 328; xvii. (1885) 240.—N. Zel.
molle, *Benth. Fl. Austral.* iii. 382.—Austral.
Morototoni, Aubl. Pl. Gui. ii. 949 = Didymopanax Morototoni.
multibracteatum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 351.—Madag.
multijugum, *Benth. & Hook. f. Gen.* i. 938.—Ins. Fiji.
Murrayi, *F. Muell. Fragm.* ii. 106.—Austral.
obtusum, *Blume, Bijdr.* 880.—Java.
ornifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 155.—Madag.
ovatum, *Hook. & Arn. Bot. Beech. Voy.* 84.—Ins. Sandvic.
palmatum, Roxb. Hort. Beng. 21; Fl. Ind. ii. 74 = Brassaiopsis palmata.
papyriferum, *F. Muell. Fragm.* iv. 122.—Austral.
parviflorum, *Mart. & Zucc. in Abh. Akad. Muench.* i. (1832) 319.—Bras.
pentadactylon, *Decne. ex Regel, Cat. Pl. Hort. Aksakov.* 100.—N. Zel.
pentamerum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 352.—Madag.
pinnatum, Baill. Hist. Pl. vii. 164 (err. typ. 464) 197 = Polyscias pinnata.
pinnatum, Lam. Encyc. ii. 715 = Arthrophyllum pinnatum.
pinnatum, *A. Rich. Tent. Fl. Abyss.* i. 335.—Abyss.
platyphyllum, *Hook. & Arn. Bot. Beech. Voy.* 84.—Ins. Sandvic.
plumatum, Hort.—Cf. Gard. Chron. (1878) I. 439 = fruticosum.
polycarpum, Wall. Cat. n. 4930 = Arthrophyllum diversifolium.
Pseudo-ginseng, Wall. Pl. As. Rar. ii. 30 = Aralia Pseudo-ginseng.
pusillum, Sims, Bot. Mag. t. 1334 = Aralia trifolia.
quinquefolium, Linn. Sp. Pl. 1058 = Aralia quinquefolia.
repens, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 64. —Japon.
ricinifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 199 = Acanthopanax ricinifolium.
rotundifolium, Hort. ex Link, Enum. Hort. Berol. ii. 448, nomen (Quid?).—Hab.?
sambucifolium, *Sieber, ex DC. Prod.* iv. 255.—Austral.

PANAX:—
samoënse, *A. Gray, Bot. U. St. Expl. Exped.* i. 717.—Ins. Samoa.
scutellarioides, Reinw. in Blume, Bijdr. 880 = cochleatum.
secundum, Schult. Syst. vi. 213 = Arthrophyllum pinnatum.
sericeum, Pohl, in DC. Prod. iv. 254 = Didymopanax vinosum.
serratum, Wall. Cat. n. 4915 = Macropanax oreophilum.
sessiliflorum, Panch. ex Sebert, Not. Bois Nouv. Caléd. 202; et ex Carr. in Rev. Hortic. (1874) 248 = Murrayi.
sessiliflorum, Rupr. & Maxim. in Bull. Phys.-Math. Pétersb. xv. (1856) 133; et Maxim. Prim. Fl. Amur. 131 = Acanthopanax sessiliflorum.
sessilifolium, C. Koch, Dendrol. i. 675, sphalm. = praec.
simplex, *Forst. f. Prod.* 75.—N. Zel.
simplicifolium, F. G. Dietr. Vollst. Lexik. Gaertn. vi. 633; ex DC. Prod. iv. 253 = simplex.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 103.—N. Zel.
speciosum, Willd. Sp. Pl. iv. 1126 = Didymopanax Morototoni.
spinosum, Linn. f. Suppl. 441 = Acanthopanax spinosum.
spinosum, Poir. Encyc. Suppl. ii. 778 = Didymopanax Morototoni.
splendens, *H. B. & K. Nov. Gen. et Sp.* v. 11.—Am. austr.
tahitense, *Nadeaud, Enum. Pl. Tahiti*, 63.—Ins. Tahiti.
tomentosum, Wall. ex DC. Prod. iv. 254 = Heptapleurum impressum.
tridactylon, Decne. et Planch. ex Seem. in Journ. Bot. iii. (1865) 179 = Pseudopanax crassifolium.
trifolium, Linn. Sp. Pl. 1059 = Aralia trifolia.
tripinnatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 153.—Madag.
tripinnatum, Wall. Cat. n. 4934 = Aralia cachemirica.
undulatum, H. B. & K. Nov. Gen. et Sp. v. 11. t. 417. f. 2 = Didymopanax Morototoni.
valdiviense, *Baill. Hist. Pl.* vii. 164 (err. typ. 464).—Chili.
Victoriae, *André, in Illustr. Hortic.* xxxi. (1884) t. 521. —Ins. Pacif.
Victoriae, Hort. Bull.—Cf. Gard. Chron. (1883) I. 404 = fruticosum.
vinosum, Cham. & Schlecht. in Linnaea, i. 402 = Didymopanax vinosum.
Weinmanniae, *Baill. Adansonia*, xii. (1878) 150.—N. Caled.
zanthoxyloides, *Baker, in Journ. Linn. Soc.* xx. (1883) 154.—Madag.
Zippelianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 15. —N. Guinea.

PANAXUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = **Panax**, Linn. (Araliac.).

PANCALUM, Ehrh. Beitr. iv. 147 (1789) = **Hypericum**, Linn.

PANCHERIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. ix. (1862) 74; et in Ann. Sc. Nat. Sér. V. i. (1864) 347. *SAXIFRAGEAE*, Benth. & Hook. f. i. 649.
alaternoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 75.—N. Caled.
elegans, *Brongn. & Gris, l. c.*—N. Caled.
ferruginea, *Brongn. & Gris, l. c.* 76.—N. Caled.
obovata, *Brongn. & Gris, l. c.* 75.—N. Caled.
pirifolia, *Brongn. & Gris, l. c.*—N. Caled.
pyrifolia, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. i. (1864) 376 = praec.
ternata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 76.—N. Caled.
Vieillardi, *Brongn. & Gris, l. c.* 75.—N. Caled.

PANCHEZIA, Montr. in Mém. Acad. Lyon, x. (1860) 223 = **Ixora**, Linn. (Rubiac.).
collina, Montr. l. c.—N. Caled.

PANCIATICA, Picciv. Hort. Panciat. 9 (1783) = **Cadia**, Forsk. (Legumin.).
purpurea, Picciv. l. c. = C. purpurea.

PANČIČIA, Vis. Sem. Hort. Patav. (1857) 9.—Cf. Linnaea, xxix. (1857-58) 732 = **Pimpinella**, Riv. (Umbell.).
serbica, Vis. ll. cc.—Europ.

PANCOVIA, Heist. ex Adans. Fam. ii. 294 (1763) = **Potentilla**, Linn. (Rosac.).
angustifolia, Rafin. Fl. Tellur. ii. 56 = Pot. palustris.
digitata, Rafin. l. c. 55 = Pot. palustris.
palustris, Rafin. l. c. = Pot. palustris.

PANCOVIA, Willd. Sp. Pl. ii. 285 (1799) = **Erioglossum**, Blume (Sapind.).
africana, Baill. Adansonia, ix. (1870) 229 = E. cauliflorum.
bijuga, Willd. Sp. Pl. ii. 285 = E. cauliflorum.
rubiginosa, Baill. ex Kurz, For. Fl. Brit. Burma, i. 296 = E. edule.
tomentosa, Kurz, l. c. = Sapindus tomentosus.
turbinata, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 270.—Afr. trop.

PANCRATIO-CRINUM, Herb. ex Steud. Nom. ed. II. ii. 250 (1841) = **Crinum**, Linn. (Amaryll.).
disticho-capense, Herb. ex Steud. l. c. = C. longifolium.

PANCRATIUM, Dill. ex Linn. Syst. ed. I (1735).
AMARYLLIDEAE, Benth. & Hook. f. iii. 733.
ALMYRA, Salisb. in Trans. Hort. Soc. i. (1812) 336.
BOLLAEA, Parl. in Bull. Soc. Bot. Fr. v. (1858) 509.
HALMYRA, Herb. Amaryll. 202 (1837); Parl. Nov. Gen. Monoc. 28 (1854).
TIARANTHUS, Herb. l. c. (1837).
TROXISTEMON, Rafin. Fl. Tellur. iv. 23 (1836).
ZOUCHIA, Rafin. l. c. 22 (1836).
abchasicum, *Regel, Cat. Pl. Hort. Aksakov.* 101.—Reg. Caucas.
acutifolium, Sweet, Hort. Brit. ed. I. 406 = Hymenocallis littoralis.
aegyptiacum, *M. Roem. Syn. Ensat.* 180.—Aegypt.
Amancaes, [Ker-Gawl.] in Bot. Mag. t. 1224 = Hymenocallis Amancaes.
amboinense, Linn. Sp. Pl. 291 = Eurycles sylvestris.
americanum, Mill. Gard. Dict. ed. VIII. n. 7 = Hymenocallis littoralis.
amoenum, Andr. Bot. Rep. t. 556 = Hymenocallis ovata.
amoenum. Salisb. in Trans. Linn. Soc. ii. (1794) 71. t. 10 = Hymenocallis caribaea.
angustifolium, M. Roem. Syn. Ensat. 178 = maritimum.
angustum, Ker-Gawl. in Bot. Reg. t. 221 = Hymenocallis caribaea.
aurantiacum, H. B. & K. Nov. Gen. et Sp. i. 280 = Stenomesson aurantiacum.
australasicum, Ker-Gawl. in Bot. Reg. t. 715 = Eurycles sylvestris.
australe, Spreng. Syst. ii. 47 = Eurycles sylvestris.
biflorum, Roxb. Hort. Beng. [23]; Fl. Ind. ii. 125 = verecundum.
Boschianum, Walp. Ann. i. 835 = Hymenocallis undulata.
Bovei, Steud. Nom. ed. II. ii. 250 = parviflorum.
calathiforme, [Delar. in] Red. Lil. t. 352 = Hymenocallis calathina.
calathinum, [Ker-Gawl. in] Bot. Reg. t. 215 = Hymenocallis calathina.
cambayense, Herb. Amaryll. 202, 207. t. 42. f. 1 = longiflorum.
canariense, [*Ker-Gawl. in*] *Bot. Reg.* t. 174.—Ins. Canar.
caribaeum, Curt. Bot. Mag. t. 826 = Hymenocallis speciosa.
caribaeum, Linn. Sp. Pl. 291 = Hymenocallis caribaea.
carolinianum, Linn. Sp. Pl. 291 = maritimum.
carribaeum, Mill. Gard. Dict. ed. VIII. n. 4 = Hymenocallis lacera.
Chapmanni, Harv. Gen. S. Afr. Pl. ed. II. 384 = trianthum.
Chlidanthus, Steud. Nom. ed. II. ii. 250 = Stenomesson luteum, Chlidanthus fragrans.
coccineum, Ruiz & Pav. Fl. Peruv. iii. 54 = Stenomesson coccineum.
collinum, *Coss. & Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 228.—Afr. bor.

PANCRATIUM:—
coronarium, Le Conte, in Ann. Lyc. N. York, iii. (1828–36) 145. t. 4. f. 7–9 = Hymenocallis crassifolia.
crassifolium, *Baker, in Saund. Refug. Bot.* t. 331.—Florida.
crassifolium, Schult. f. Syst. vii. 921 = Hymenocallis crassifolia.
croceum, Savign. in Lam. Encyc. iv. 725 (err. typ. 735); [DC. in] Red. Lil. iv. t. 187 = Stenomesson croceum.
Cunninghami, Steud. Nom. ed. II. ii. 251 = Eurycles Cunninghamii.
curvidentatum, Hort. ex Steud. l. c. 250 = Stenomesson flavum.
declinatum, Jacq. Select. Am. 99 = Hymenocallis caribaea.
disciforme, [DC. in] Red. Lil. t. 155 = Hymenocallis lacera.
distichum, Sims, Bot. Mag. t. 1879 = Hymenocallis littoralis.
Dryandri, Ker-Gawl. in Quart. Journ. Sc. & Arts, iii. (1817) 326 = Hymenocallis littoralis.
excisum, Linn. f. ex Kunth, Enum. Pl. v. 669 = Hymenocallis caribaea.
expansum, Sims, Bot. Mag. t. 1941 = Hymenocallis expansa.
flavum, Ruiz & Pav. Fl. Per. iii. 54 = Stenomesson croceum.
fluitans, Fraser, ex Schult. f. Syst. vii. 930, in obs. = nutans.
foetidum, *Pomel, Nouv. Mat. Fl. Atl.* 253.—Alger.
formosum, Hort. ex M. Roem. Syn. Ensat. 167 = = Hymenocallis speciosa.
fragrans, Salisb. in Trans. Linn. Soc. ii. (1794) 72 = Hymenocallis ovata.
fragrans, Willd. Sp. Pl. ii. 42 = Hymenocallis caribaea.
glaucum, Zucc. in Abh. Akad. Muench. ii. (1837) 317 = Hymenocallis glauca.
guadelupense, *Hort. ex M. Roem. Syn. Ensat.* 182 (Hymenocallidis sp. ?).—Ins. Guadelup.
guianense, [Ker-Gawl. in] Bot. Reg. t. 265 = Hymenocallis tubiflora.
Herbertianum, Spreng. Syst. iv. Cur. Post. 132 = Stenomesson flavum.
humile, Cav. Ic. iii. t. 4 = Tapeinanthus humilis.
illyricum, Blanco, Fl. Filip. ed. I. 251 = Hymenocallis littoralis.
illyricum, Forsk. Fl. Aegypt. Arab. p. lxv = aegyptiacum.
illyricum, *Linn. Sp. Pl.* 291.—Europ. austr.
incarnatum, H. B. & K. Nov. Gen. et Sp. i. 280 = Stenomesson incarnatum.
Knightii, Steud. Nom. ed. II. ii. 251 = nutans ?
latifolium, Mill. Gard. Dict. ed. VIII. n. 8 = Hymenocallis speciosa.
latifolium, Ruiz & Pav. Fl. Per. iii. 54 = Urceolina latifolia.
limense, Ker-Gawl. in Quart. Journ. Sc. & Arts, iii. (1817) 332 = Stenomesson coccineum.
liriosme, Rafin. Fl. Ludov. 19 (Quid ?).—Am. bor.
littorale, Jacq. Select. Am. 99 = Hymenocallis littoralis.
longiflorum, *Buch.-Ham. ex Roxb. Hort. Beng.* [23]; *Fl. Ind.* ii. 125.—Ins. Molucc.
luteum, Pav. ex Steud. Nom. ed. II. ii. 251 = Stenomesson luteum.
luteum, Poir. ex Spreng. Syst. ii. 48 = Chlidanthus fragrans.
Macleanum, Steud. Nom. ed. II. ii. 251 = Ismene Macleana.
Macquaria, Lewin, ex Steud. Nom. ed. II. ii. 251 = Calostemma purpureum.
malabathricum, Herb. Amaryll. 202, 206 = triflorum.
maritimum, *Linn. Sp. Pl.* 291.—Reg. Mediterr.; Am. bor.
maximum, *Forsk. Fl. Aegypt. Arab.* 72.—Arabia.
mexicanum, Lindl. Bot. Reg. t. 940 = Hymenocallis littoralis.
mexicanum, Linn. Sp. Pl. 290 = Hymenocallis lacera.
narcissiflorum, Jacq. Fragm. 68 = Hymenocallis calathina.
nervifolium, Salisb. Parad. Lond. t. 84 = Eurycles sylvestris.
nutans, *Ker-Gawl. in Quart. Journ. Sc. & Arts*, iii. (1817) 324.—Am. bor. et austr.

PANCRATIUM:—
obtusatum, *Griseb. in Linnaea*, xxi. (1848) 277.—Guiana.
occidentale, Le Conte, in Ann. Lyc. N. York, iii. (1828–36) 146 = Hymenocallis crassifolia.
ornatum, *Bouché, in Linnaea*, xiii. (1839) 79.—Hab. ?
ovalifolium, Steud. Nom. ed. II. ii. 251 = Hymenocallis ovalifolia.
ovatifolium, Stokes, Bot. Mat. Med. ii. 215 = Eurycles sylvestris.
ovatum, Mill. Gard. Dict. ed. VI. [in 4to] (1771) n. 9 = Hymenocallis ovata.
parviflorum, *Decne. in Ann. Sc. Nat.* Sér. II. iv. (1835) 346.—Syria.
parviflorum, Desf. Tabl. ed. II. 272 = Vagaria parviflora.
parvum, *Dals. in Hook. Kew Journ.* ii. (1840) 144.—Ind. or.
patens, [Delile, in] Red. Lil. vii. sub t. 414 = Hymenocallis caribaea.
patens, Lindl. in Trans. Hort. Soc. vi. (1820) 87 = Hymenocallis caymanensis.
pedale, Schult. f. Syst. vii. 916 = Hymenocallis pedalis.
petiolatum, Willd. in Schult. f. l. c. 912, in obs. = Hymenocallis tubiflora.
plicatum, Livinst. ex Steud. Nom. ed. II. ii. 251 (Quid ?).—Ind. or.
quitense, Willd. ex Schult. f. Syst. vii. 908 = Stenomesson incarnatum.
quitoënse, Schult. f. l. c. 923 = Hymenocallis quitoënsis.
recurvatum, Ruiz & Pav. Fl. Per. iii. 54 = Stenomesson recurvatum.
recurvum, Stokes, Bot. Mat. Med. ii. 214 = Hymenocallis caribaea.
ringens, Ruiz & Pav. Fl. Per. iii. 53 = Elisena ringens.
rotatum, Ker-Gawl. in Bot. Mag. t. 827 = Hymenocallis lacera.
rubrum, Pav. ex Steud. Nom. ed. II. ii. 251 = Stenomesson rubrum.
Sickenbergeri, *Aschers. & Schweinf. ex Boiss. Fl. Orient.* v. 153.—Egypt.; Arabia.
speciosum, Linn. f. ex Salisb. in Trans. Linn. Soc. ii. (1794) 73; Salisb. Prod. 227 = Hymenocallis speciosa.
Staplesi, Steud. Nom. ed. II. ii. 251 = Hymenocallis littoralis.
stellare, Salisb. in Trans. Linn. Soc. ii. (1794) 74 = illyricum.
Tagliabue, *Colla, Hort. Ripul. App.* iv. 60.—Hab. ?
Teneriffae, Willd. in Schult. f. Syst. vii. 925, adnot. = canariense.
tenuiflorum, Herb. ex Steud. Nom. ed. II. ii. 251 = Hymenocallis tenuiflora.
tenuifolium, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 312 = trianthum.
tiaraeflorum, Salisb. Parad. Lond. t. 86 = zeylanicum.
tortifolium, Boiss. Diagn. Ser. I. xiii. 18 = tortuosum.
tortuosum, *Herb. in Ann. Nat. Hist.* Ser. I. iv. (1840) 28.—Arab.
trianthum, *Herb. l. c.*—Afr. trop.; Arab.
trichromum, *Cerv. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 20.—Mexic.
triflorum, *Roxb. Hort. Beng.* [23]; *Fl. Ind.* ii. 126.—Ind. or.
triphyllum, Willd. ex M. Roem. Syn. Ensat. 169 = Hymenocallis undulata.
tristylum, *De Rijk, in Rev. Hortic.* (1880) 446.—Java.
tubiflorum, Schult. f. Syst. vii. 923 = Hymenocallis tubiflora.
tubulosum, *Willd. ex Kunth, Enum. Pl.* v. 667.—Am. austr.
undulatum, H. B. & K. Nov. Gen. et Sp. i. 280 = Hymenocallis undulata.
uniflorum, Stokes, Bot. Mat. Med. ii. 213; Donn, Hort. Cantabr. ed. IV. 68 = zeylanicum.
variegatum, *Ruiz & Pav. Fl. Per.* iii. 55.—Peruv.
verecundum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 412.—Ind. or.
viridiflorum, Ruiz & Pav. Fl. Per. iii. 55 = Stenomesson viridiflorum.
zeylanicum, *Linn. Sp. Pl.* 290.—Ind. or.

PANCTENIS, Rafin. New Fl. Am. ii. 60 (1836) = **Gerardia**, Linn. (Scrophul.).
pectinata, Rafin. l. c. 61 = G. Pedicularia.
pedicularis, Rafin. l. c. = G. Pedicularia.
pumila, Rafin. l. c. (Quid ?).—Am. bor.

PANDACA, Noronha, ex Thou. Gen. Nov. Madag. 10 (1806) = **Tabernaemontana**, Linn. (Apocyn.).

PANDANOPHYLLUM, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 118 = **Mapania**, Aubl. (Cyperac.).
humile, Hassk. l. c. 119 = M. humilis.
hypolytroides, F. Muell. Fragm. ix. 16 (Thoracostachyi sp.).—Austral.
immersum, Thw. Enum. Pl. Zeyl. 433 = M. immersa.
longifolium, Boeck. in Flora, lviii. (1875) 112.—Malaya; Ins. Samoa.
macrocephalum, Boeck. l. c.—N. Guin.
Miquelianum, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) ii. 81.—Malaya.
palustre, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 119 = M. palustris.
squamatum, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) ii. 80.—Java.
Wendlandi, Hort.—Cf. Gard. Chron. (1884) i. 711 = M. humilis.
zeylanicum, Thw. Enum. Pl. Zeyl. 345 = M. zeylanica.
Zippelianum, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 224 = M. humilis.

PANDANUS, Rumph. ex Linn. f. Suppl. 64 (1781).
PANDANEAE, Benth. & Hook. f. iii. 949.
Athrodactylis, Forst. Char. Gen. 149 (1776).
Barrotia, Gaudich. Bot. Voy. Bonite, t. 13 (1844–52).
Bryantia, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 287. t. 15.
Doornia, De Vriese, in Hook. Kew Journ. vi. (1854) 266; et in Tuinb. Fl. (1855) 174.
Dorystigma, Gaudich. Bot. Voy. Bonite, t. 13 (1844–52).
Eydouxia, Gaudich. l. c. t. 18 (1844–52).
Fisquetia, Gaudich. l. c. t. 4, 5 (1844–52).
Foullioya, Gaudich. l. c. t. 26 (1844–52).
Hasskarlia, Walp. Ann. i. 753 (1849).
Heterostigma, Gaudich. Bot. Voy. Bonite, t. 25 (1844–52).
Hombronia, Gaudich. l. c. t. 22 (1844–52).
Jeanneretia, Gaudich. l. c. t. 25 (1844–52).
Keura, Forsk. Fl. Aegypt. Arab. 95, 122, 172 (1775).
Marquartia, Hassk. in Flora, xxv. (1842) ii. Beibl. 14.
Russinia, Gaudich. Bot. Voy. Bonite, t. 21 (1844–52).
Rykia, De Vriese, in Verh. Kon. Akad. Wetensch. ii. (1854) 203.
Sussea, Gaudich. Bot. Voy. Bonite, tt. 24, 25, 38 (1844–52).
Tuckeya, Gaudich. l. c. t. 26 (1844–52).
Vinsonia, Gaudich. l. c. tt. 17, 23, 31 (1844–52).
acuminatus, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 40, nomen.—Madag.
acuminatus, *Hort. ex H. Wendl. Ind. Palm.* (1854) 45.—Hab. ?
affinis, *Kurz, in Journ. Bot.* v. (1867) 101.—Zeylan.
albus, *Hort. ex Steud. Nom.* ed. II. ii. 251.—Ind. or.
altissimus, Panch. ex Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 277 = P. Minda ?
altissimus, *Solms, in Linnaea*, xlii. (1878–79) 43.—N. Caled.
amaryllidifolius, Hort. ex H. Wendl. Ind. Palm. (1854) 46 = laevis ?
amaryllidifolius, Voigt, in Syll. Ratisb. ii. (1828) 52 = seq. ?
amaryllifolius, *Roxb. Hort. Beng.* 71; *Fl. Ind.* iii. 743.—Ins. Amboin.
amarylloides, Parment. ex Desf. Cat. Hort. Par. ed. III. 9 = praec. ?
Amherstiae, *Hort. ex H. Wendl. Ind. Palm.* (1854) 45.—Hab. ?
andamanensis, Hort. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 41, nomen = seq. ?
andamanensium, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) ii. 148.—Ins. Andam.
angustifolius, *Hort. ex Steud. Nom.* ed. II. ii. 251.—Hab. ?

PANDANUS:—
aquaticus, *F. Muell. in Hook. Kew Journ.* viii. (1856) 329; *Fragm.* v. 40.—Austral.
aragoënsis, *Solms, in Linnaea*, xlii. (1878–79) 45; *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 41.—N. Caled.
atrocarpus, *Griff. Notul.* iii. 160.—Ind. or.
australis, *Prestoe, Cat. Hort. Trinit.* 77.—Austral.?
Bagea, Miq. Fl. Ind. Bat. iii. 159 = dubius?
Balansae, Solms, in Linnaea, xlii. (1878–79) 45; *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 42.—N. Caled.
Barkleyi, *Balf. f. in Baker, Fl. Maurit.* 397.—Ins. Mascar.
Beccarii, *Solms, in Ann. Jard. Buitenz.* iii. (1883) 97.—N. Guin.
Bidur, *Jungh. ex Miq. Pl. Jungh.* i. 166 = latissimus.
Blancoi, Kunth, Enum. Pl. iii. 583 = odoratissimus?
Boivini, *Solms, in Linnaea*, xlii. (1878–79) 26.—Madag.
Boryi, Gaudich. Bot. Voy. Bonite, t. 22. f. 15 = odoratissimus?
Boucheanus, *C. Koch, in C. Koch & Fint. Wochenschr.* (1858) 131.—Madag.
bromeliaefolius, *Lodd. ex Desf. Cat. Hort. Par.* ed. III. 9.—Ins. Borbon.
butyrophorus, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 150.—Hab.?
calathiphorus, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 42.—Ins. Salomon.
Candelabrum, *Beauv. Fl. Owar.* i. 37. tt. 21, 22.—Afr. trop.
Candelabrum, Hook. Bot. Mag. t. 5014 = utilis?
Candelabrum, Kurz, in Journ. Bot. v. (1867) 127 = P. Kaida.
caricosus, *Spreng. Syst.* iii. 897.—Ins. Molucc.
cauliflorus, *Carm. ex Balf. f. in Journ. Linn. Soc.* xvii. (1878) 44, nomen.—Ins. Maurit.
ceramicus, Kunth, Enum. Pl. iii. 98 = conoideus?
ceratophorus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 448.—Madag.
ceylanicus, *Solms, in Linnaea*, xlii. (1878–79) 16.—Zeylan.
Chamissonis, Gaudich. Bot. Voy. Bonite, t. 22. f. 9 = odoratissimus?
concidens, Pritz. Ic. Ind. i. 793, sphalm. = conoideus.
concretus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 448.—Madag.
conglomeratus, *Balf. f. in Baker, Fl. Maurit.* 403.—Ins. Mascar.
conoideus, De Vriese, in Hook. Kew Journ. vi. (1854) 264 = montanus?
conoideus, *Lam. Encyc.* i. 372.—Malaya.
conoideus, Thou. in Bull. Soc. Philom. (1808) 5 = prostratus?
crassipes, Wall. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 46 = furcatus?
cylindricus, *Hort. ex Balf. f. l. c.* 44, nomen.—Hab.?
decorus, *Hort. ex C. Koch, Wochenschr.* xiii. (1870) 166.—N. Caled.
decumbens, *Solms, in Linnaea*, xlii. (1878–79) 47; *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 44.—N. Caled.
deflexus, Hort. ex H. Wendl. Ind. Palm. (1854) 45 = Doornianus?
demissus, Soland. ex Kunth, Enum. Pl. iii. 104 = Freycinetia demissa.
distichus, Hort.—Cf. Illustr. Hortic. xix. (1872) 55 = utilis?
Doornianus, *De Vriese, ex H. Wendl. Ind. Palm.* (1854) 45.—Ins. Mascar.
Douglasii, Gaudich. Bot. Voy. Bonite, t. 22. f. 16 = odoratissimus?
drupaceus, *Thou. in Bull. Soc. Philom.* (1808) 4.—Ins. Mascar.
dubius, *Spreng. Syst.* iii. 897.—Ins. Amboin.
edulis, De Vriese, in Hook. Kew Journ. vi. (1854) 264 = dubius?
edulis, *Thou. in Bull. Soc. Philom.* (1808) 4.—Madag.
elegans, Thou. l. c. = sylvestris?
elegantissimus, Hort. ex Proc. Roy. Hort. Soc. iii. (1863) 108 = utilis?
ensifolius, *Thou. in Bull. Soc. Philom.* (1808) 4.—Ins. Mascar.
erigens, Thou. l. c. 5 = montanus.
exaltatus, *Blanco, Fl. Filip.* ed. I. 778.—Ins. Philipp.
Eydouxia, *Balf. f. in Baker, Fl. Maurit.* 401.—Ins. Mascar.

PANDANUS:—
fascicularis, Lam. Encyc. i. 372 = odoratissimus?
flabelliformis, Carr. in Rev. Hortic. (1866) 271 = utilis?
foetidus, *Roxb. Hort. Beng.* 71; *Fl. Ind.* iii. 742.—Ind. or.
Forsteri, *C. Moore & F. Muell. in F. Muell. Fragm.* viii. 220.—Ins. Dom. Howe.
fragrans, *Gaudich. Bot. Voy. Bonite*, t. 22. f. 10.—Ins. Marian.
freycinetioides, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 151.—Hab.?
furcatus, *Roxb. Hort. Beng.* 71; *Fl. Ind.* iii. 744.—Ind. or.
furcatus, Thw. Enum. Pl. Zeyl. 327 = ceylanicus.
glaucescens, *Hort.—Cf. Proc. Roy. Hort. Soc.* v. (1865) 143.—Hab.?
glaucus, Hort. ex H. Wendl. Ind. Palm. (1854) 47 = sessilis.
globuliferus, Thou. in Bull. Soc. Philom. (1808) 5 = sphaeroideus?
gracilis, *Blanco, Fl. Filip.* ed. I. 778.—Ins. Philipp.
gramineus, Hort. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 48 = Freycinetia graminea.
graminifolius, Hort. ex Miq. Anal. Bot. Ind. ii. 17 = pygmaeus.
graminifolius, *Kurz, in Journ. Bot.* v. (1867) 104.—Burma.
helicopus, *Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 54. t. 2.—Ins. Banca.
heterocarpus, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 22; *Baker, Fl. Maurit.* 399.—Ins. Mascar.
Heudelotianus, *Balf. f. l. c.* xvii. (1878) 49.—Afr. trop.
Hoffa, *Chapel. ex Boj. Hort. Maurit.* 303.—Madag.
Hornei, *Balf. f. in Baker, Fl. Maurit.* 397.—Ins. Seychell.
horridus, Reinw. ex Blume, Cat. Gew. Buitenz. 111 = furcatus.
Houlletii, *Carr. in Rev. Hortic.* (1868) 210. f. 23.—Singapore.
humilis, Jacq. Fragm. 21. t. 14. f. 2 = sylvestris?
humilis, Lour. Fl. Cochinch. 603 = polycephalus.
humilis, *Rumph. ex Miq. Fl. Ind. Bat.* iii. 160.—Ind. or.; China.
Iceryi, *Horne, ex Balf. f. in Baker, Fl. Maurit.* 400.—Ins. Seychell.
inclinans, Soland. ex Kunth, Enum. Pl. iii. 102 = Freycinetia Banksii.
inermis, *Blanco, Fl. Filip.* ed. II. 537.—Ins. Philipp.
inermis, Roxb. Hort. Beng. 71; Fl. Ind. iii. 744 = laevis.
integrifolius, *Lour. ex Steud. Nom.* ed. II. ii. 251 (Quid?).—China; Ind. or.
javanicus, Hort. ex H. Wendl. Ind. Palm. (1854) 45 = P. Candelabrum?
Joskei, *Horne, ex Balf. f. in Journ. Linn. Soc.* xx. (1883) 416.—Ins. Fiji.
Kaida, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 148.—Ind. or.
Korthalsii, *Solms, in Linnaea*, xlii. (1878–79) 12.—Borneo.
Kurzianus, Solms, l. c. 4 = polycephalus.
labyrinthicus, *Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 53.—Sumatra.
laevis, *Lour. Fl. Cochinch.* 604.—Cochinch.; Java.
lagenaeformis, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 50.—Hab.?
Lais, *Hort. ex Balf. f. l. c.* nomen.—Ins. Molucc.
Lais, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 218 = furcatus.
latifolius, Hassk. in Flora, xxv. (1842) II. Beibl. 13 = amaryllifolius.
latifolius, *Hort. Angl. ex Link, Enum. Hort. Berol.* ii. 412, nomen (Quid?).—Hab.?
latifolius, Hort. Lodd. ex C. Koch, Wochenschr. (1858) 131 = Boucheanus.
latifolius, Loud. ex Steud. Nom. ed. II. ii. 251 = polycephalus.
latifolius, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 134.—Ins. Philipp.
latissimus, *Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 52.—Malaya.
leonensis, *Hort. ex H. Wendl. Ind. Palm.* (1854) 46.—Afr. trop.
Leram, *Jones, ex Voigt, Hort. Suburb. Calc.* 683.—Ins. Nicobar.
Leram, Kurz, in Journ. Bot. v. (1867) 185 = andamanensium.

PANDANUS:—
leucanthus, Hassk. in Flora, xxv. (1842) II. Beibl. 14 = odoratissimus?
leucanthus, Hort. ex H. Wendl. Ind. Palm. (1854) 46 = odoratissimus?
Linnaei, Gaudich. Bot. Voy. Bonite, t. 22. ff. 1–8 = odoratissimus?
Linnei, *Hort. ex Balf. f. in Journ. Linn. Soc.* xvii. (1878) 51, nomen.—Hab.?
littoralis, Jungh. Topog. Nat. Reise Java, 61 = odoratissima?
littoralis, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 150.—Malaya.
lividus, *Hort. ex H. Wendl. Ind. Palm.* (1854) 46.—Hab.?
longifolius, Hort. Berol. ex H. Wendl. l. c. = laevis?
longifolius, Hort. Lodd. ex Steud. Nom. ed. II. ii. 251 = Doornianus?
Loureiri, Gaudich. Bot. Voy. Bonite, t. 22. f. 13 = odoratissimus?
lucidus, Wall. ex Kurz, in Journ. As. Soc. Beng. xxxviiii. (1869) II. 149 = sylvestris?
macrocarpus, *Hort. ex Balf. f. in Journ. Linn. Soc.* xvii. (1878) 52, nomen.—Hab.?
macrocarpus, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 51.—N. Caled.
madagascariensis, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 52.—Madag.
malatensis, *Blanco, Fl. Filip.* ed. II. 536.—Ins. Philipp.
marginatus, *Roxb. Hort. Beng.* 71.—Ins. Maurit.
maritimus, Thou. in Bull. Soc. Philom. (1808) 4 = utilis?
mauritianus, Hort. Berol. ex H. Wendl. Ind. Palm. (1854) 47 = sylvestris?
mauritianus, Hort. Kew. ex Lem. Illustr. Hortic. vii. (1860) t. 265 = utilis?
Menziesii, Gaudich. Bot. Voy. Bonite, t. 22. f. 14 = odoratissimus?
microcarpus, *Balf. f. in Baker, Fl. Maurit.* 396.—Ins. Mascar.
microcephalus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 447.—Madag.
microstigma, *Balf. f. l. c.* xvii. (1878) 53.—Madag.
militaris, *Balf. f. l. c.*—Singapore.
Millore, Roxb. Hort. Beng. 71, nomen = P. Leram?
Minda, Panch. ex Brongn. in Compt.-rend. Par. lxxx. (1875) 1196; et in Ann. Sc. Nat. Sér. VI. i. (1875) 288 = oblongus?
Minda, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 51.—N. Caled.
minor, *Buch.-Ham. ex Wall. Cat.* n. 8592.—Ind. or.
monodon, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 53.—Cochinch.
montanus, *Bory, Voy.* i. 313.—Ins. Borbon.
montanus, Miq. Fl. Ind. Bat. iii. 161 = atrocarpus.
monticola, *F. Muell. Fragm.* v. 40; vii. 63; viii. 220.—Austral.
Moorei, *F. Muell. Census*, 140.—Austral.
moschatus, Hort. ex H. Wendl. Ind. Palm. (1854) 45 = amaryllifolius?
moschatus, Miq. Fl. Ind. Bat. iii. 165 = laevis?
Motleyanus, *Solms, in Linnaea*, xlii. (1878–79) 21.—Borneo.
multispicatus, *Balf. f. in Baker, Fl. Maurit.* 403.—Ins. Seychell.
muricatus, *Thou. in Bull. Soc. Philom.* (1808) 6.—Madag.
nitidus, Kurz, in Journ. Bot. v. (1867) 103 = stenophyllus?
nudus, Thou. in Bull. Soc. Philom. (1808) 4 = utilis?
obeliscus, *Thou. l. c.* 6.—Madag.
oblongus, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 54.—N. Caled.
odoratissimus, Blume, Cat. Gew. Buitenz. 111 = amaryllifolius?
odoratissimus, *Hort. Herrenh. ex H. Wendl. Ind. Palm.* (1854) 46.—Hab?
odoratissimus, Jacq. Fragm. 21. t. 13, 14. f. 1 = utilis?
odoratissimus, *Linn. f. Suppl.* 424.—Ind. or.; Arab.
odoratissimus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = laevis.
odoratus, Salisb. Prod. 3 = odoratissimus?
odoratus, *Thunb. Fl. Jav.* 19.—Java.
Olango, *Blanco, ex Vidal & Garcia, Cat. Sem. Manil.* 14.—Ins. Philipp.

PANDANUS:—
oligocephalus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 448.—Madag.
ornatus, *Hort. Bull.—Cf. Journ. Roy. Hort. Soc.* N. S. i. (1866) *Misc.* i.—Ins. Rodriguez.
ornatus, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 147.—Malacca.
ovatus, *Kurz, l. c.*—Malacca.
palustris, *Thou. in Bull. Soc. Philom.* ii. (1808) 6.—Ins. Mascar.
Pancheri, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 57.—N. Caled.
papuanus, *Solms, in Ann. Jard. Buitenz.* iii. (1883) 93.—N. Guin.
parasiticus, Noronha, in Verh. Batav. Gen. v. (1790) ed I. Art. IV. 25 (Freycinetiae sp.).—Java.
pedunculatus, *R. Br. Prod.* 341.—Austral.
Pervilleanus, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) II. 149 = Boucheanus ?
Pervilleanus, *Solms, in Linnaea*, xlii. (1878-79) 35.—Madag.
polycephalus, *Lam. Encyc.* i. 372.—Malaya.
polyryzos, *Noronha, ex Thou. Prod. Phyt.* 2, *in Mélang.* (1811), nomen.—Madag.
Porteanus, *Lescuyer, ex Hérincq, L'Hortic. Français*, (1866) 16.—Ins. Philipp.
Princeps, *Hort. Bull.—Cf. Gard. Chron.* (1878) I. 440.—Austral.
proliferus, *Hort. ex H. Wendl. Ind. Palm.* (1854) 47.—Hab. ?
prostratus, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 59.—Ins. Maurit.
purpurascens, *Thou. in Bull. Soc. Philom.* (1808) 3.—Ins. Mascar.
pygmaeus, Hook. Bot. Mag. t. 4736 = seq. ?
pygmaeus, *Thou. in Bull. Soc. Philom.* (1808) 6.—Ins. Mascar.
pyramidalis, *Barkly, ex Balf. f. in Baker, Fl. Maurit.* 399.—Ins. Mascar.
pyramidatus, Carm. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 53 = montanus.
racemosus, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 150.—Hab. ?
radicans, *Blanco, Fl. Filip.* ed. I. 780.—Ins. Philipp.
reflexus, *Lodd. ex Desf. Cat. Hort. Par.* ed. III. 9; *C. Koch & Fint. Wochenschr.* i. (1858) 132.—Ins. Mascar.
repens, Miq. Fl. Ind. Bat. iii. 165 = P. Samak.
reticulatus, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 52.—N. Caled.
Rheedei, Gaudich. Bot. Voy. Bonite, t. 22. f. 12 = odoratissimus ?
Rumphii, Gaudich. l. c. f. 11 = odoratissimus ?
Sabotan, *Blanco, Fl. Filip.* ed. I. 779.—Ins. Philipp.
Samak, *Hassk. in Flora* (1842), *Beibl.* II. 13; *Cat. Hort. Bog.* 61.—Java.
sativus, Thou. in Bull. Soc. Philom. (1808) 3 = utilis ?
sechellarum, *Balf. f. in Baker, Fl. Maurit.* 402.—Ins. Seychell.
sessilis, *Boj. Hort. Maurit.* 302.—Afr. trop.
sessilis, *Hort. ex H. Wendl. Ind. Palm.* (1854) 47.—Austral.
sphaerocephalus, *Panch. ex Brongn. in Ann. Sc. Nat.* Sér. VI. i. (1875) 284, partim.—N. Caled.
sphaerocephalus, Panch. ex Brongn. l. c., partim = Pancheri.
sphaeroideus, *Thou. in Bull. Soc. Philom.* (1808) 5.—Ins. Mascar.
spinifructus, Dennst. Schluess. Hort. Malab. 27 = furcatus ?
spiralis, Blanco, Fl. Filip. ed. I. 777 = odoratissimus ?
spiralis, R. Br. Prod. 341 = odoratissimus ?
spiralis, Hort. ex H. Wendl. Ind. Palm. (1854) 47 = utilis ?
spurius, Miq. Anal. Bot. Ind. ii. 15 = utilis ?
spurius, Miq. Fl. Ind. Bat. iii. 157 = odoratissimus ?
spurius, Steud. Nom. ed. II. ii. 251 = P. Candelabrum.
stenocarpus, *Solms, in Ann. Jard. Buitenz.* iii. (1883) 91.—N. Guin.
stenophyllus, *Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 53.—Java.
striatus, Carm. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 60 = pyramidalis ?
strigilis, Carm. ex Balf. f. l. c. 45 = drupaceus ?
subumbellatus, *Becc. ex Solms, in Ann. Jard. Buitenz.* iii. (1883) 96.—Malaya.
Sussea, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 62.—Hab. ?

PANDANUS:—
sylvestris, *Bory, Voy.* ii. 260.—Ins. Mascar.
sylvestris, Kunth, Enum. Pl. iii. 98 = conoideus ?
tectorius, Soland. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 63, 56 = odoratissimus.
tenuifolius, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 22.—Ins. Rodriguez.
tenuifolius, *Linden, ex Dufresne, in Illustr. Hortic.* xx. (1873) 70.—Hab. ?
tetrodon, *Balf. f. in Journ. Linn. Soc.* xvii. (1878) 63.—Hab. ?
turbinatus, *Lodd. ex Steud. Nom.* ed. II. ii. 251.—Ind. or.
unguifer, Hook. f. Bot. Mag. t. 6347 = minor.
unipapillatus, *Dennst. Schluess. Hort. Malab.* 27.—Ind. or.
urophyllus, Hance, in Journ. Bot. xiii. (1875) 68 = furcatus ?
utilis, *Bory, Voy.* ii. 3.—Madag.
utilis, Hort. ex H. Wendl. Ind. Palm. (1854) 47 = odoratissimus ?
Vacqua, Carm. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 64 = utilis ?
Vandermeeschii, *Balf. f. in Baker, Fl. Maurit.* 398.—Ins. Mascar.
variegatus, Miq. Anal. Bot. Ind. ii. 28 = P. Samak ?
Veitchi, *Hort.—Cf. Gard. Chron.* (1868) 349; *Hort. ex Dallière, Pl. Ornament.* i. 28 (1873).—Polynes.
viscidus, *Panch. ex Brongn. in Ann. Sc. Nat.* Sér. VI. i. (1875) 287.—N. Caled.
vittariifolius, *Boj. Hort. Maurit.* 302.—Madag.
Yvanii, *Solms, in Linnaea*, xlii. (1878-79) 20.—Malacca.

PANDERIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 46 (1835). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 57.
PTEROCHLAMYS, Fisch. ex Endl. Gen. 294 (1837).
divaricata, C. Koch, in Linnaea, xxii. (1849) 186 = pilosa.
monticola, *Boiss. ex Bunge, Pfl. Geogr. Betracht. Fam. Chenop.* 9.—Turkestan.
pilosa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 46.—As. Min.; Oriens.

PANDIAKA, Benth. & Hook. f. Gen. iii. 35 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
cylindrica, *Benth. & Hook. f. l. c.* 36.—Afr. trop.
Heudelotii, *Benth. & Hook. f. l. c.*—Afr. trop.
involucrata, *Benth. & Hook. f. l. c.*—Afr. trop.

PANDORA, Noronha, ex Thou. Vég. Isl. Afr. 48 (1806) = **Rhodolaena**, Thou. (Chlaenac.).

PANDOREA, Spach, Hist. Vég. Phan. ix. 136 (1840) = **Tecoma**, Juss. (Bignon.).
australis, Spach, l. c. = T. australis.
austro-caledonica, Seem. in Gard. Chron. (1870) 1085 = T. austro-caledonica.

PANEGUIA, Rafin. Fl. Tellur. iv. 34 (1836) = **Sisyrinchium**, Linn. (Irid.).
palmifolia, Rafin. l. c. = S. palmifolium.
striata, Rafin. l. c. = S. striatum.

PANEMATA, Rafin. Fl. Tellur. iv. 64 (1836) = **Diciliptera**, Juss. (Acanthac.).
zeylanica, Rafin. l. c. = D. zeylanica.

PANETOS, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 227; vi. 81 = **Hedyotis**, Linn. (Rubiac.).

PANGIUM, Reinw. in Syll. Ratisb. ii. (1828) 12. *BIXINEAE*, Benth. & Hook. f. i. 129.
edule, *Reinw. l. c.*—Java.

PANICASTRELLA, Moench, Meth. 205 (1794) = **Echinaria**, Desf. (Gramin.).
capitata, Moench, l. c. 206 = Sesleria capitata.
muricata, Moench, l. c. = E. capitata.

PANICULARIA, Heist. Syst. 12 (1748) (Quid ?).

PANICULUM, Ard. Animad. Spec. Alt. 14 (1764), err. typ. = **Oplismenus**, Beauv. (Gramin.).
undulatifolium, Ard. l. c. = O. compositus ?

PANICUM, Linn. Syst. ed. I (1735); Gen. ed. I. 17 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1100.
ACICARPA, Raddi, Agrost. Bras. 31 (1823).
BLUFFIA, Nees, Del. Sem. Hort. Hamb. (1835).—Cf. Linnaea, x. (1836) Litt. 114; et Fl. Afr. Austr. 61 (1841).
BRACHIARIA, Griseb. in Ledeb. Fl. Ross. iv. 469 (1853).
COLEATAENIA, Griseb. in Goett. Abh. xxiv. (1879) 308.
CORIDOCHLOA, Nees, in Edinb. N. Phil. Journ. xv. (1833) 381.
DIGITARIA, Heist. ex Adans. Fam. ii. 38, 550 (1763); Scop. Fl. Carn. ed. II. i. 52 (1772).
DILEUCADEN, Rafin. ex Steud. Nom. ed. II. i. 508 (1840).
DIMORPHOSTACHYS, Fourn. in Compt. Rend. Acad. Par. lxxx. (1875) 441.
ECHINOCHLOA, Beauv. Agrost. 53. t. 11 (1812).
ERIACHNE, Phil. Sert. Mendoc. Alt. 49 (1871).
ERIOLYTRUM, Desv. ex Kunth, Rév. Gram. i. 219 (1829).
HOLOSETUM, Steud. in Flora, xxxiii. (1850) I. 228, nomen; Syn. Pl. Gram. 118 (1855).
HYMENACHNE, Beauv. Agrost. 48. t. 10. f. 8 (1812).
MESOSETUM, Steud. in Flora, xxxiii. (1850) I. 228, nomen; Syn. Pl. Gram. 118 (1855).
MILIUM, [Tourn.] Adans. Fam. ii. 34 (1763).
OTACHYRIUM, Nees, Agrost. Bras. 273 (1829).
PARACTAENUM, Beauv. Agrost. 47. t. 10 (1812).
PSEUDORAPHIS, Griff. Notul. iii. 29 (1851).
RHYNCHELYTHRUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
SANGUINELLA, Gleichen, Mikrosk. Untersuch. 4 [bis] t. 8 (1764).
STEINCHISMA, Rafin. in Ser. Bull. Bot. i. (1830) 220.
STREPTOSTACHYS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190.
SYNTHERISMA, Walt. Fl. Carol. 76 (1788).
TEMA, Adans. Fam. ii. 496 (1763).
THALASIUM, Spreng. Syst. iv. Cur. Post. 30 (1827).
THRASYA, H. B. & K. Nov. Gen. et Sp. i. 120. t. 39 (1815).
TRICHACHNE, Nees, Agrost. Bras. 85 (1829).
TRICHOLAENA, Schrad. in Schult. Mant. ii. 163 (1824).
TYLOTHRASYA, Doell, in Mart. Fl. Bras. ii. II. 295. t. 37 (1877).
UROCHLOA, Beauv. Agrost. 52. t. 11. f. 1 (1812).
VALOTA, Adans. Fam. ii. 495 (1763).
abludens, *Roem. & Schult. Syst.* ii. 457.—Ind. or.
abortivum, R. Br. Prod. 193 = Chamaeraphis spinescens.
abyssinicum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 360.—Abyss.
acariferum, Trin. Sp. Gram. Ic. i. t. 87 = Thysanolaena Agrostis.
accrescens, Trin. l. c. t. 88 = trigonum ?
aciculare, *Desv. in Poir. Encyc. Suppl.* iv. 274.—Ind. or.
acroanthum, *Steud. Syn. Pl. Gram.* 87.—Ind. or.; China; Japon.
acromelaenum, *Hochst. in Flora*, xxxviii. (1855) 198.—Abyss.
acrotrichum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 226.—Afr. trop.
acuminatissimum, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 194 = latissimum.
acuminatissimum, Steud. Syn. Pl. Gram. 45 = Oplismenus acuminatus.
acuminatum, Muhl. Desc. Gram. 125 = Muehlenbergii.
acuminatum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 234 = P. Myurus.
acuminatum, Sw. Prod. Veg. Ind. Occ. 23 = dichotomum.
acutatum, *Steud. Syn. Pl. Gram.* 86.—Chili.
acutiflorum, *Poir. Encyc. Suppl.* iv. 283.—Hab. ?
acutiflorum, Steud. Syn. Pl. Gram. 93 = Ichnanthus candicans.
acutifolium, Willd. ex Spreng. Syst. i. 320 = capillare, hirtum.
acutiglumum, Steud. Syn. Pl. Gram. 66 = amplexicaule.
acutum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Reimaria acuta.
adhaerens, Forsk. Fl. Aegypt. Arab. 20 = Setaria verticillata.

PANICUM:—
adoënse, Steud. Nom. ed. II. ii. 252 = Pennisetum adoënse.
adpressum, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 193 = sanguinale.
adscendens, Cham. & Schlecht. in Linnaea, vi. (1831) 32 = sanguinale.
adscendens, Hoffmgg. ex Schult. Mant. ii. 592 = parvifolium.
adscendens, *H. B. & K. Nov. Gen. et Sp.* i. 97.—Am. trop.
adspersum, Trin. Gram. Panic. 146 = grossarium.
adstans, Steud. Syn. Pl. Gram. 94 = Isachne miliacea.
adtenuatum, *Moench, Meth. Suppl.* 67.—Hab.?
adustum, *Nees, Agrost. Bras.* 101.—Bras.
aegyptiacum, Hort. ex Roem. & Schult. Syst. ii. 493 = Setaria italica.
aegyptiacum. Retz. Obs. iii. 8 = sanguinale.
aegyptium, J. F. Gmel. Syst. 1596, sphalm. = praec.
aemulum, Steud. Nom. ed. II. ii. 252 = Oplismenus setarius.
aequatum, *Nees, ex Steud. Syn. Pl. Gram.* 98.—Ind. or.
aequinerve, *Nees, Fl. Afr. Austr.* 40.—Afr. austr.
affine, Hook. & Arn. Bot. Beech. Voy. 100 = nubigenum.
affine, Nees, Agrost. Bras. 113 = paspaloides.
affine, Poir. Encyc. Suppl. iv. 273 = Oplismenus prostratus.
africanum, Poir. l. c. 275 = Oplismenus compositus?
Afzelii, *Sw. Adnot. Bot.* 5.—Afr. trop.
agglutinans, Kunth, Enum. Pl. i. 120 = lanatum.
agrioides, Trin. ex Doell, in Mart. Fl. Bras. ii. II. 271 = Isachne miliacea.
agrostideum, Salzm. ex Steud. Nom. ed. II. ii. 252 = Ichnanthus pallens.
agrostidiforme, Lam. Illustr. i. 172 = laxum.
Agrostis, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 213 = laxum.
agrostoides, Salzm. ex Steud. Syn. Pl. Gram. 93 = Ichnanthus candicans.
agrostoides, *Spreng. Pugill.* ii. 4.—Am. bor.
airoides, R. Br. Prod. 190 = repens.
airoides, Fluegge, ex Nees, Agrost. Bras. ii. 175, in adnot. = madagascariense.
alabamense, Trin. ex Steud. Syn. Pl. Gram. 64 = floridanum.
albens, Trin. ex Steud. Nom. ed. II. ii. 252; Syn. Pl. Gram. 96 = Isachne albens.
albidulum, *Steud. l. c.* 69.—Nubia.
album, Poir. Encyc. Suppl. iv. 274 = Oplismenus Burmanni.
alectorocnemum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = P. Crus-galli.
alectromerum, Dulac, Fl. Hautes-Pyr. 76 = P. Crus-galli.
almadense, Nees, ex Trin. Gram. Panic. 186 = Ichnanthus almadensis.
alopecuroides, Koen. ex Trin. l. c. 162 = Setaria glauca.
alopecuroides, Linn. Sp. Pl. 55 = Pennisetum compressum, Linnaei.
alopecuroides, Spreng. Syst. i. 309 = Neurachne alopecuroides.
alopecuroides, Walt. Fl. Carol. 72 = Setaria verticillata.
alopecuroideum, Schreb. ex Steud. Syn. Pl. Gram. 50 = Setaria flava.
Alopecuros, Fisch. ex Spreng. Nov. Prov. 38 = Setaria macrostachya.
Alopecuros, Lam. Illustr. i. 169 = Pennisetum setosum.
alsinoides, Griseb. Fl. Brit. W. Ind. 550 = Ichnanthus alsinoides.
alternatum, *Brouss. Elench. Hort. Monsp.* (1805) 42, nomen.—Hab.?
altissimum, Brouss. l. c. = maximum.
altissimum, *G. F. W. Mey. Prim. Fl. Esseq.* 63.—Guian.
amabile, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 324 = decompositum.
amarum, *Ell. Sketch*, i. 121.—Am. bor.
ambiguum, Haussk. in Oestr. Bot. Zeitschr. xxv. (1875) 345 = P. Pseudoholcus.
ambiguum, Lapeyr. Hist. Abr. Pl. Pyr. 31 = glabrum.
ambiguum, *Sang. in Atti Acc. Pont. Lincei*, Ser. I. xviii. (1865) 219. t. 6.—Italia.
ambiguum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 243.—As. trop.; Maurit.; Ins. Pacif.
ambitiosum, *Fourn. ex Hemsl. Biol. Centr. Am.* iii. 485.—Mexic.
americanum, Linn. Sp. Pl. 56 = Pennisetum typhoideum.

PANICUM:—
ammophilum, F. Muell. in Trans. Vict. Inst. i. (1855) 46 = divaricatissimum.
ammophilum, *Steud. Syn. Pl. Gram.* 92.—Afr. austr.
ammophilum, Trin. ex Nees, Agrost. Bras. 164 = Setaria brachiata.
amnigenum, *Trin. ex Steud. Nom.* ed. II. ii. 252.—Hab.?
amphibium, Steud. Syn. Pl. Gram. 61 = molle.
amphibolum, Steud. l. c. 51 = Setaria composita.
amphistemon, *Wright, in Sauv. Fl. Cub.* 198.—Cuba.
amplectens, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 20.—Am. bor.
amplexicaule, Poir. ex Kunth, Enum. Pl. i. 113 = ovalifolium.
amplexicaule, *Rudge, Pl. Gui.* i. 21.—Amphig. trop.
amplexifolium, *Hochst. in Flora*, xxxviii. (1855) 194.—Abyss.
amplifolium, Steud. Syn. Pl. Gram. 53 = Setaria composita.
amplissimum, Steud. l. c. 54 = plicatum.
anceps, *Michx. Fl. Bor. Am.* i. 48.—Am. bor.
ancylocarpum, *Nees, ex Steud. Nom.* ed. II. ii. 252.—Hab.?
angustifolium, Chapm. Fl. S. U. St. 374 = brachyanthum.
angustifolium, Ell. Sketch, i. 129 = dichotomum.
angustifolium, Le Conte, ex Torr. Cat. Pl. N. York, 91 = debile.
angustissimum, *Hochst. ex Steud. Syn. Pl. Gram.* 66.—Am. trop.
angustum, Trin. Sp. Gram. Ic. iii t. 334 = indicum.
angustum, Wall. Cat. n. 1642 = myosuroides.
anisostachium, *Boj. Hort. Maurit.* 364.—Ins. Maurit.
annulatum, *A. Rich. Tent. Fl. Abyss.* ii. 370.—Abyss.
anomalum, Walt. Fl. Carol. 72 = dichotomum.
ansatum, *Trin. Sp. Gram. Ic.* iii. t. 279.—Bras.
antidotale, *Retz. Obs.* iv. 17.—Geront. calid.
antillarum, Poir. Encyc. Suppl. iv. 275 = Setaria setosa.
antipodum, Spreng. Syst. i. 314 = Isachne australis.
Aparine, Steud. Syn. Pl. Gram. 52 = Setaria verticillata.
apiculatum, Salzm. ex Steud. l. c. 65 = pilosum.
appressum, Doell, in Mart. Fl. Bras. ii. II. 184 = paspaloides.
appressum, Forsk. Fl. Aegypt. Arab. 20 = fluitans?
appressum, Kunth, Enum. Pl. i. 84 = *adpressum*.
aquariorum, Buch.-Ham. ex Wall. Cat. n. 8727 = ramosum.
aquaticum, Bosc, ex Spreng. Syst. ii. 319 = gibbum.
aquaticum, Muhl. Desc. Gram. 126 = hydrophilum.
aquaticum, *Poir. Encyc. Suppl.* iv. 281.—Portorico.
aquaticum, *A. Rich. Tent. Fl. Abyss.* ii. 373.—Abyss.
arabicum, Nees, ex Steud. Nom. ed. II. ii. 252 = colonum.
araneosum, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 180.—Ind. or.
arborescens, Linn. Sp. Pl. ed. I. 59 = sparsicomum, Arundinaria glaucescens.
arborescens, Sieber, ex Trin. Gram. Panic. 208 = latifolium.
arbusculum, Sieber, ex Griseb. Fl. Brit. W. Ind. 553 = Isachne rigens.
arcuatum, R. Br. Prod. 189 = indicum.
arenarium, Bieb. Fl. Taur. Cauc. i. 52 = glabrum.
arenarium, Brot. Fl. Lusit. i. 82 = repens.
arenarium, *Cham. & Schlecht. in Linnaea*, vi. (1831) 35.—Mexic.
argenteum, *R. Br. Prod.* 190.—Austral.
argyrograptum, Nees, Fl. Afr. Austr. 27 = sanguinale.
argyrostachyum, *Steud. Syn. Pl. Gram.* 40.—Java.
argyrotrichum, *Anderss. in Peters, Reise Mossamb. Bot.* 548.—Afr. trop.
aristatum, Cav. ex Willk. & Lange, Prod. Fl. Hisp. i. 57 = Polypogon monspeliensis.
aristatum, Macfad. in Hook. Bot. Misc. ii. (1831) 115 = spectabile.
aristatum, *Rasp. in Ann. Sc. Nat.* Sér. I. v. (1825) 299, absque desc.—Hab.?
aristatum, Retz. Obs. iv. 17 = Oplismenus compositus.
Aristella, *Doell, in Mart. Fl. Bras.* ii. II. 221.—Am. trop.
aristidoides, Willd. ex Spreng. Syst. i. 302 = Pennisetum compressum.
aristulatum, *Steud. Syn. Pl. Gram.* 42.—Afr. trop.
Arnacites, *Trin. Sp. Gram. Ic.* iii. t. 317.—Bras.

PANICUM:—
Arnottianum, Nees, in Steud. Syn. Pl. Gram. 59 = nodosum.
Arnottianum, Nees, in Steud. l. c. 98 = nubigenum.
arundinaceum, Sw. Prod. Veg. Ind. Occ. 24 = Isachne disperma.
Arundinariae, *Trin. ex Steud. Nom.* ed. II. ii. 253; *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 485.—Mexic.
arvense, Kunth, Rév. Gram. ii. t. 109 = ramosum.
ascendens, Hemsl. Biol. Centr. Am. Bot. iii. 485 = adscendens.
ascendens, Willd. ex Spreng. Syst. i. 321 = parvifolium.
asiaticum, Hort. ex Schult. Mant. ii. 598, in obs. = Setaria italica.
asperatum, Kunth, Rév. Gram. i. 39 = plicatum.
asperrimum, Fisch. Cat. Hort. Gorenk. ed. II. 3; ex Jacq. Eclog. Gram. 46. t. 31; Lag. ex Link, Enum. Hort. Berol. i. 77 = miliaceum.
asperum, Koen. in Naturf. xxiii. (1788) 209 = Chamaeraphis spinescens.
asperum, Lam. Fl. Fr. iii. 577 = Setaria verticillata.
asperum, Link, Enum. Hort. Berol. i. 75 = Pennisetum asperum.
astracanicum, Hort. ex Steud. Nom. ed. II. ii. 253 = Koeleria brachystachya.
atrichum, *Steud. l. c.*—Abyss.
atro-purpureum, *Desv. Opusc.* 97.—Hab.?
atrosanguineum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 375.—Abyss.
atroviolaceum, *A. Rich. l. c.* 368.—Abyss.
atrovirens, Trin. in Spreng. Neue Entdeck. ii. 88 = Isachne australis.
attenuatum, Hort. ex Trin. Gram. Panic. 166 = Setaria italica.
attenuatum, Willd. Enum. Hort. Berol. 1033 = miliare.
aturense, Balb. ex Kunth, Enum. Pl. i. 119 = oryzoides.
aturense, *H. B. & K. Nov. Gen. et Sp.* i. 103. t. 33.—Venezuel.
aurantiacum, Buch.-Ham. ex Wall. Cat. n. 8714 = paludosum.
Aurelianum, *Hale, ex Wood, Class Book* (1861), 787.—Am. bor.
aureum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 196.—Bras.
auricomum, *Nees, ex Trin. Gram. Panic.* 204.—Bras.
auriculatum, Willd. ex Spreng. Syst. i. 322 = Ichnanthus panicoides.
auritum, Hassk. Pl. Jav. Rar. 22 = amplexicaule.
auritum, *Presl, ex Nees, Agrost. Bras.* 176.—As. trop.
australe, Kasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Isachne australis.
australe, Spreng. Syst. i. 309 = parviflorum.
austro-caledonicum, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 326.—N. Caled.
autumnale, *Bosc, ex Spreng. Syst.* i. 320.—Am. bor.
autumnale, F. Muell. Fragm. viii. 196 = papposum.
avenaceum, *H. B. & K. Nov. Gen. et Sp.* i. 99.—Am. trop.
axillare, Nees, Agrost. Bras. 141 = Ichnanthus pallens.
axipilium, *Steud. Syn. Pl. Gram.* 79.—Guian.
Baileyi, *Benth. Fl. Austral.* vii. 471.—Austral.
Balanites, Trin. in Linnaea, x. (1835-36) 297 = streptostachys.
Balbisianum, Schult. Mant. ii. 254 = oryzoides.
Balfourii, *Baker, Fl. Maurit.* 438.—Ins. Rodrig.
bambusaefolium, *Desv. Opusc.* 83.—Bras.
bambusiflorum, Trin. Gram. Panic. 188 = Ichnanthus bambusiflorus.
bambusoides, *Ham. Prod. Pl. Ind. Occ.* 10.—Portorico.
Bansnium, Buch.-Ham. ex Wall. Cat. n. 8725 A = ramosum.
barbatulum, Steud. Nom. ed. II. ii. 253, 255, in syn., sphalm. = *barbulatum*.
barbatum, *Kunth, Rév. Gram.* i. 33.—Malaya; China.
barbatum, Lam. Illustr. i. 171 = prostratum
barbatum, Roxb. Hort. Beng. 7; Fl. Ind. i. 282 = Pennisetum barbatum.
barbifultum, Hochst. ex Schlecht. in Linnaea, xxxi. (1861-62) 307 = Oplismenus undulatifolius.
barbigerum, *Bertol. in Mem. Acc. Sc. Bologn.* iii. (1851) 250.—Afr. trop.?

PANICUM:—
barbinode, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 256 = molle.
barbulatum, Michx. Fl. Bor. Am. i. 49 = dichotomum.
basisetum, *Steud. Syn. Pl. Gram.* 52.—Afr. trop.
batavicum, Steud. l. c. 96 = Isachne australis.
beckmanniaeforme, Mikan, ex Trin. in Spreng. Neue Entdeck. ii. 83 = paspaloides.
Beecheyi, *Hook. & Arn. Bot. Beech. Voy.* 100.—Ins. Sandvic.
bellum, Steud. Nom. ed. II. ii. 253 = Isachne pulchella.
bengalense, *Spreng. Syst.* i. 311.—Ind. or.
Benjamini, Steud. Syn. Pl. Gram. 96 = Isachne australis.
Benthami, *Steud. l. c.* 43.—Afr. trop.
Berchtholdiae, Doell, in Mart. Fl. Bras. ii. II. 150, in obs. = Chaetium bromoides.
bermudianum, Steud. Nom. ed. II. ii. 253 = gibbum.
Berteronianum, *Steud. Syn. Pl. Gram.* 50.—Ins. S. Doming.
Bertolonianum, Schult. Mant. ii. 240 = Ichnanthus candicans.
Beyreichii, Kunth, Rév. Gram. i. 281. t. 27 = rugulosum.
bicolor, *R. Br. Prod.* 191.—Austral.
bicolor, Moench, Meth. 206 = Setaria viridis.
bicorne, Kunth, Rév. Gram. i. 33 = sanguinale.
bicorne, Sieber, ex Steud. Syn. Pl. Gram. 41 = didactylum.
bicrululum, *Salzm. ex Steud. Nom.* ed. II. ii. 273 (Quid ?).—Bras.
bidentatum, Steud. Syn. Pl. Gram. 45 = Oplismenus compositus.
bifasciculatum, Trin. Gram. Panic. 76 = Schraderi.
bifidum, *Bertol. in Mem. Acad. Sc. Bolog.* ii. (1850) 598.—Am. bor.
biflorum, Lam. Illustr. i. 174 = Isachne miliacea.
biforme, Kunth, Rév. Gram. i. 33 = sanguinale.
Billbergianum, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 113 = aturense.
bipustulatum, *Schlecht. in Linnaea*, xxvi. (1853-55) 135.—N. Granat.
bistipulatum, Steud. Syn. Pl. Gram. 417, sphalm. = praec.
bisulcatum, *Thunb. in Nov. Act. Soc. Sc. Upsal.* vii. (1815) 141 (sp. dub.).—Japon.
blephariphyllum, *Trimen, in Journ. Bot.* xxiii. (1885) 272.—Zeylan.
blepharophorum, J. & C. Presl, Rel. Haenk. i. 312 = aturense.
Bobartii, Lam. Encyc. iv. 748 = dichotomum ?
Bonplandianum, Steud. Syn. Pl. Gram. 48 = Oplismenus polystachyus.
Boscianum, Spreng. Syst. i. 317 = striatum.
Boscii, Poir. Encyc. Suppl. iv. 282 = dichotomum ?
Bourgaei, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 486.—Mexic.
brachiariaeforme, *Steud. Syn. Pl. Gram.* 54.—Afr. trop.
brachiatum, *Bosc, ex Spreng. Syst.* i. 321.—Ins. Bermud.
brachiatum, Poir. Encyc. Suppl. iv. 282 = Setaria brachiata.
brachyanthum, *Steud. Syn. Pl. Gram.* 67.—Am. bor.
brachyclados, Reichb. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 251 = stoloniferum.
brachyglume, Hochst. ex Steud. Syn. Pl. Gram. 95 = Coelachne pulchella.
brachylachnum, *Steud. l. c.* 62.—Afr. trop.
brachyphyllum, *Steud. l. c.* 42.—Mexic.
brachystachyum, *Trin. Gram. Panic.* 138.—Bras.
brachytrichum, *Steud. in Lechl. Berb. Am. Austr.* 52, nomen.—Chili.
brasiliense, Spreng. Syst. i. 321 = parvifolium.
Braunii, *Steud. Syn. Pl. Gram.* 93.—Abyss.
brevifolium, Balb. ex Nees, Fl. Afr. Austr. 40 = miliare.
brevifolium, Boj. Hort. Maurit. 365 = parvifolium.
brevifolium, Jahn, ex Schrank, in Syll. Ratisb. i. (1824) 193 = Jahnii.
brevifolium, *Kunth, Rév. Gram.* i. 32.—Mexic.
brevifolium, *Linn. Sp. Pl.* 59.—Am. trop.
brevifolium, Roxb. Fl. Ind. i. 306 = ovalifolium.
brevifolium, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 290 = microspermum.
brevisetum, Doell, Fl. Grossh. Baden, 233 = Setaria italica.

PANICUM:—
brevisetum, *Steud. Syn. Pl. Gram.* 46.—Venezuela.
brizaeforme, J. & C. Presl, Rel. Haenk. i. 302 = paspaloides.
brizanthum, *Hochst. in Flora*, xxiv. (1841) I. Intell. 19, nomen ; *et ex A. Rich. Tent. Fl. Abyss.* ii. 363. —Abyss.
brizoides, Jacq. Eclog. Gram. 2. t. 2; Trin. Gram. Panic. 128 = flavidum.
brizoides, Lam. Illustr. i. 170 = paspaloides.
brizoides, Linn. Mant. i. 184; Sw. Obs. Bot. 36 = colonum.
brizoides, Retz. Obs. iii. 8 = fluitans.
brizoides, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 274 = Isachne ventricosa.
bromoides, Lam. Illustr. i. 170 = Oplismenus Burmanni.
Brownei, Roem. & Schult. Syst. ii. 462 = lanatum.
Brownianum, Wight & Arn. ex Steud. Syn. Pl. Gram. 34, 98 = subeglume.
Brunonianum, Griff. Notul. iii. 29; Ic. Pl. As. iii. t. 145 (Chamaeraphidis sp.).—Ind. or.
Buchingeri, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 486.—Mexic.
bulbosum, *H. B. & K. Nov. Gen. et Sp.* i. 99.—Mexic.
Buncei, *F. Muell. ex Benth. Fl. Austral.* vii. 487.—Austral.
bunophilum, Steud. Syn. Pl. Gram. 76 = Ichnanthus calvescens.
Burmanni, Balb. Misc. Bot. i. 8; Schleich. ex Steud. Nom. ed. II. ii. 253 = Oplismenus undulatifolius.
Burmanni, Bieb. Fl. Taur. Cauc. i. 51 = P. Crus-galli.
Burmanni, Heyne, ex Wall. Cat. n. 8734 A = Isachne Kunthiana.
Burmanni, Retz. Obs. iii. 10 = Oplismenus Burmannii.
caesioglaucum, *Nees, ex Steud. Syn. Pl. Gram.* 75.—Java.
caesium, Hook. & Arn. Bot. Beech. Voy. 235; Nees, ex Steud. Syn. Pl. Gram. 47 = colonum.
caespititium, *Lam. Illustr.* i. 173; *Encyc.* iv. 743.—Am. austr.
caespitosum, Spreng. ex Steud. Nom. ed. II. ii. 253 = grossarium.
caespitosum, Sw. Fl. Ind. Occ. i. 146 = prostratum.
caffrorum, Retz. Obs. ii. 7 = Sorghum vulgare.
cajennense, *Lam. Illustr.* i. 173.—Guian.
calaccanzense, *Steud. Syn. Pl. Gram.* 65.—Ins. Philipp.
californicum, *Benth. Bot. Voy. Sulph.* 55.—Calif.
callosum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 374 = miliaceum.
Calumet, Brouss. Elench. Hort. Monsp. (1805) 42, nomen (Quid ?).—Hab. ?
calvescens, Nees, ex Trin. Gram. Panic. 193 = Ichnanthus calvescens.
campestre, *Nees, ex Trin. l. c.* 197.—Bras.
camporum, *Kunth, Enum. Pl.* i. 88.—Bras.
campylogrammum, *Buese, in Miq. Pl. Jungh.* 375.—Sumatra.
canaliculatum, *Nees, ex Steud. Syn. Pl. Gram.* 55.—Ind. or.
Canarae, Steud. Syn. Pl. Gram. 58 = Paspalum compactum.
candicans, Nees, Agrost. Bras. 133 = Ichnanthus candicans.
candidum, Nees, ex Trin. Gram. Panic. 176 = Ichnanthus pallens.
canescens, Roth, Nov. Sp. Pl. 54 = ramosum ?
capense, *Licht. ex Roem. & Schult. Syst.* ii. 457.—Afr. austr.
capillaceum, Lam. Illustr. i. 173 = brevifolium.
capillare, Hook. & Arn. Bot. Beech. Voy. 100, nomen = nephelophilum.
capillare, *Linn. Sp. Pl.* 58.—Am. bor. et austr.
capillare, F. Muell. Fragm. viii. 191 = effusum.
capillare, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 271 = commelinaefolium.
capillare, Vahl, ex Hornem. Hort. Hafn. i. 84 = commelinaefolium ?
capillipes, *Benth. Fl. Austral.* vii. 484.—Austral.
Careyanum, Nees, Agrost. Bras. 118 = paspaloides
caricoides, *Nees, ex Trin. Gram. Panic.* 149.—Bras.
carinatum, *J. & C. Presl, Rel. Haenk.* i. 309.—Ins. Philipp.
carinatum, Torr. ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 410 = Curtisii.

PANICUM:—
carnosum, Salzm. ex Steud. Syn. Pl. Gram. 60 = paspaloides.
carolinianum, *Spreng. Syst.* i. 310.—Am. bor.
carthaginense, *Sw. Fl. Ind. Occ.* i. 148.—Am. trop.
cartilagineum, Muhl. Desc. Gram. 128 = dichotomum.
cartilagineum, Nees, Agrost. Bras. 112 = subulatum.
caucasicum, Trin. Sp. Gram. Ic. iii. t. 262 = P. Isachne.
Cauda-ratti, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 59.—Afr. trop.
caudatum, Lam. Illustr. i. 171 = Setaria setosa.
caudatum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 233 = vilfoides.
caudatum, Thunb. Prod. Pl. Cap. 19 = interruptum.
caudatum, Willd. ex Steud. Nom. ed. II. ii. 251 = amplexicaule.
caudigerum, *Trin. ex Steud. Nom.* ed. II. ii. 254.—Hab. ?
cayennense, *Lam. Illustr.* i. 173.—Am. trop.
cenchroides, Ell. Sketch, i. 111 = Cenchrus Elliotii.
cenchroides, Lam. Encyc. iv. 737 = Trachys mucronata.
cenchroides, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Pennisetum setosum.
cernuum, *Willd. ex Spreng. Syst.* i. 318. = Ecuador.
certificandum, Steud. Syn. Pl. Gram. 44 = Oplismenus compositus.
chaetium, *Steud. Nom.* ed. II. ii. 254.—Bras.
chaetophorum, Roem. & Schult. Syst. ii. 884 = setigerum.
chamaelonche, Trin. Gram. Panic. 242 = dichotomum.
Chamaeraphis, Nees, ex A. Br. Ind. Sem. Hort. Berol. (1855) App. 20 = rhachitrichum.
Chamaeraphis, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 217 = Chamaeraphis hordeacea.
Chapmani, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 61.—Am. bor.
chartaginense, Sw. Prod. Veg. Ind. Occ. 22 = carthaginense.
Chauvinii, *Steud. Syn. Pl. Gram.* 68.—Mexic.
chinense, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 225.—China.
chloroleucum, *Griseb. in Goett. Abh.* xix. (1874) 262. —Reg. Argent.
chlorostachyum, *Doell, in Mart. Fl. Bras.* ii. II. 173. —Am. trop.
chloroticum, Nees, Agrost. Bras. 164 = Setaria brachiata.
chnoodes, *Trin. Gram. Panic.* 211.—Bras.
chondrachne, Steud. Syn. Pl. Gram. 51 = Setaria setosa.
Chondrachyrum, Trin. ex Steud. Nom. ed. II. ii. 254 (Melicae sp. ?).—Bras.
chrysanthum, Steud. Nom. ed. II. ii. 254 = Setaria glauca.
chrysites, Steud. Syn. Pl. Gram. 38 = Paspalum chrysites.
chrysoblephare, Steud. l. c. = Paspalum exasperatum.
chrysochaetum, *Steud. l. c.* 50.—Ins. Borbon.
chrysodactylon, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 197 = Paspalum chrysodactylon.
chrysostachyum, Trin. l. c. 196 = Paspalum chrysostachyum.
ciliare, Retz. Obs. iv. 16 = sanguinale.
ciliatiflorum, Wood, Class Book (1861), 786 = Anthaenantia villosa.
ciliatifolium, Kunth, Rév. Gram. i. 36 = dichotomum.
ciliatum, Ell. Sketch, i. 126 = dichotomum.
ciliatum, Thunb. Mus. Nat. Acad. Upsal. x. 148, nomen (Quid ?).—Hab. ?
cimicinum, *Retz. Obs.* iii. 9.—Ind. or.
cinerascens, *Doell, in Mart. Fl. Bras.* ii. II. 189.—Am. trop.
clandestinum, *Linn. Sp. Pl.* 58.—Am. bor.
clandestinum, *Wall. Cat.* n. 8680.—Burma.
coarctatum, Stev. ex Trin. Gram. Panic. 137 = P. Crus-galli ?
coccospermum, Steud. Syn. Pl. Gram. 62 = villosum.
coenicolum, *F. Muell. in Trans. Vict. Inst.* i. (1855) 45.—Austral.
coenosum, *Doell, in Mart. Fl. Bras.* ii. II. 191.—Am. trop.
coeruleum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Peruv.
cognatissimum, *Steud. Syn. Pl. Gram.* 69.—Afr. trop.
cognatum, Roem. & Schult. Syst. ii. 235 = autumnale.
collare, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 60.—Afr. trop.

PANICUM:—
collinum, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 323.—N. Caled.
colonum, *Linn. Syst.* ed. X. 870.—Reg. calid.
coloratum, Cav. Ic. ii. 6. t. 110 = repens.
coloratum, Kunth, Enum. Pl. i. 104 = Kunthii.
coloratum, *Linn. Mant.* i. 30.—Afr. bor.
coloratum, F. Muell. Fragm. viii. 192 = melananthum, *F. Muell.*
coloratum, Thou. ex Spreng. Syst. i. 317 = madagascariense.
comatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 376.—Abyss.
commelinaefolium, Kunth, Enum. Pl. i. 114 = microspermum.
commelinaefolium, *Rudge, Pl. Guian.* 21. t. 28.—Guian.
commixtum, *Steud. Syn. Pl. Gram.* 59.—Austral.
commutatum, Nees, in Linnaea, vii. (1832) 274 = sanguinale.
commutatum, Roem. & Schult. Syst. ii. 242 = dichotomum.
comosum, *Steud. Syn. Pl. Gram.* 53.—Ins. Philipp.
comosum, *Steud. l. c.* 417.—Japon.
compactum, Kit. in Schult. Oesterr. Fl. ed. II. i. 212 = Setaria italica.
compactum, *Sw. Adnot. Bot.* 14; *et Griseb. Fl. Brit. W. Ind.* 552.—Ind. occ.; Mexic.
composito-proximum, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1804) 224 = Oplismenus compositus.
compositum, Linn. Sp. Pl. 57 = Oplismenus compositus.
compositum, Nees, Agrost. Bras. 244 = Setaria composita.
compositum, Rottl. ex Steud. Syn. Pl. Gram. 44 = peninsularum.
compressum, Balb. ex Steud. Nom. ed. II. ii. 254 = Setaria glauca.
compressum, *Bivona, Stirp. Sic. Manip.* iv. 6.—Sicilia.
compressum, Forsk. Fl. Aegypt. Arab. 18 = Eleusine indica.
conchatum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 487.—Mexic.
concinnum, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 179.—Ind. or.
concinnum, Nees, in Hook. Kew Journ. ii. (1850) 97 = sarmentosum.
condensatum, *Bertol. in Bologn. Opusc. Sc.* iii. (1819) 408.—Bras.
confertum, *Desv. ex Poir. Encyc. Suppl.* iv. 279.—Ind. occ.
confine, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 373 = trypheron.
confusum, Trin. ex Steud. Nom. ed. II. ii. 254 = maximum, *Jacq.*
conglomératum, *Boj. Hort. Maurit.* 364.—Ins. Maurit.
conglomeratum, *Linn. Mant.* ii. 324; *Willd. Sp. Pl.* i. 341 (sp. dub.).—Ind. or.
conjugatum, Roxb. Hort. Beng. 82; Fl. Ind. i. 288 = Paspalum conjugatum.
connivens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 206.—Bras.
consanguineum, Kunth, Rév. Gram. i. 36 = dichotomum.
contractum, *Trin. ex Steud. Nom.* ed. II. ii. 254.—Ind. occ.
contractum, Wight & Arn. ex Steud. l. c. = indicum.
controversum, *Steud. Syn. Pl. Gram.* 60.—Abyss.
convallium, F. Muell. in Trans. Vict. Inst. i. (1855) 46 = effusum.
convolutum, Beauv. ex Spreng. Syst. i. 319 = repens.
cordatum, *Buese, in Miq. Pl. Jungh.* 376.—Java.
cordatum, *Doell, in Mart. Fl. Bras.* ii. III. 239.—Bras.
cordifolium, *Desv. Opusc.* 90.—Am. bor.
cordifolium, Steud. Syn. Pl. Gram. 85 = sciurotis.
cordovense, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 487.—Mexic.
coriophorum, Auct. = *coryophorum.*
corrugatum, Ell. Sketch, i. 123 = Setaria corrugata.
corvipes, Stokes, Bot. Mat. Med. i. 141 = P. Crus-galli.
corymbiferum, *Nees, ex Steud. Syn. Pl. Gram.* 76.—Afr. trop.
corymbosum, Roxb. Hort. Beng. 7; Fl. Ind. i. 292 = sanguinale.
coryophorum, Kunth, Rév. Gram. ii. t. 107 = curvatum.
costatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 312 = flavescens.

PANICUM:—
courtallense, Nees & Wight, ex Steud. Syn. Pl. Gram. 83 = montanum.
crinitum, Willd. ex Steud. Nom. ed. II. ii. 255 = Setaria purpurea.
Crinum-ursi, Bory, ex Steud. l. c. = Zoysia tenuifolia.
crispum, Llanos, Fragm. 42 = prostratum.
Crista-galli, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 357 = P. Crus-galli?
cristatum, *Anderss. in Peters, Reise Mossamb. Bot.* 548.—Afr. trop.
cristatum, Steud. Syn. Pl. Gram. 46 = Oplismenus Burmannii.
cruciatum, Nees, ex Steud. l. c. 39 = sanguinale.
cruciforme, Steud. Nom. ed. II. ii. 255, sphalm. = *eruciforme.*
Crus-ardeae, *Willd. ex Nees, Agrost. Bras.* 253.—Am. austr.
Crus-corvi, Linn. Syst. ed. X. 870 = P. Crus-galli.
Crus-galli, *Linn. Sp. Pl.* 56.—Reg. calid.
Crus-pavonis, Nees, Agrost. Bras. 259 = P. Crus-galli.
Crus-pici, Willd. ex Doell, in Mart. Fl. Bras. ii. II. 143, in obs. = P. Crus-galli.
ctenanthum, *F. Muell. Fragm.* viii. 153.—Austral.
ctenodes, Trin. Sp. Gram. Ic. ii. t. 171 = stoloniferum.
cubense, *Steud. Nom.* ed. II. ii. 255.—Cuba.
cujabense, Steud. l. c. = cuyabense.
culicinum, F. Muell. Fragm. viii. 189 = obtusum.
cultratum, Trin. Gram. Panic. 126 = monostachyum.
Cumingianum, *Steud. Syn. Pl. Gram.* 58.—Ins. Philipp.
Curtisii, Chapm. Fl. S. U. St. 573 = hydrophilum?
Curtisii, *Steud. Syn. Pl. Gram.* 66.—Am. bor.
curvatum, *Linn. Syst.* ed. XII. 730.—Ind. or.; Afr. austr.
curvatum, Roxb. Fl. Ind. i. 286 = myosuroides.
curviflorum, *Hornem. Hort. Hafn. Suppl.* 116.—Hab.?
cuspidatum, *Roxb. Hort. Beng.* 7; *Fl. Ind.* i. 298.—Ind. or.
cuspidatum, Willd. ex Steud. Nom. ed. II. ii. 255 = Isachne australis.
cuspidiglume, Steud. Syn. Pl. Gram. 96 = Isachne Kunthiana.
cuyabense, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 206.—Bras.
cyanescens, *Nees, Agrost. Bras.* 220.—Bras.
cylindricum, Link, Enum. Hort. Berol. i. 76 = Setaria setosa.
cylindricum, Thunb. Mus. Nat. Acad. Upsal. x. 148 (Quid?).—Java.
Cynodon, *Reichardt, in Sitzb. Akad. Wien*, lxxvi. (1878) 724.—Ins. Sandvic.
cynosuroides, *Linn. Syst.* ed. X. 870.—Hab.?
cynosuroides, Scop. Fl. Carn. ed. II. i. 50 = Setaria viridis.
cynotis, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 325 = Ichnanthus inconstans.
Dactylon, Linn. Sp. Pl. 58 = Cynodon Dactylon.
Daltoni, Parl. ex Webb, in Hook. Niger Fl. 185 = colonum.
dasyanthum, Licht. ex Nees, Fl. Afr. Austr. 19 = Tricholaena capensis.
dasytrichum, Spreng. Syst. i. 317 = cayennense.
dasyurum, Nees, Agrost. Bras. 241 = Pennisetum Sieberi.
dasyurum, Willd. ex Nees, Fl. Afr. Austr. 56 = Setaria imberbis.
debile, Desf. Fl. Atlant. i. 59 = sanguinale.
debile, *Poir. Encyc. Suppl.* iv. 283.—Am. bor.
debile, Torr. ex Steud. Nom. ed. II. ii. 255 = dichotomum.
decipiens, Kunth, Enum. Pl. i. 92 = despiciens.
decipiens, *Nees, Agrost. Bras.* 193.—Bras.
decolorans, *H. B. & K. Nov. Gen. et Sp.* i. 100.—Mexic.
decompositum, *R. Br. Prod.* 191.—Ind. or.; Austral.
decumbens, *Roem. & Schult. Syst.* ii. 429.—Am. austr.
deflexum, Guss. in Tenore, Fl. Nap. v. 320 = Ehrharta panicea.
deflexum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 63.—Afr. trop.
delicatulum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 381.—Nubia.
demissum, *Trin. Sp. Gram. Ic.* iii. t. 319.—Bras.

PANICUM:—
densepilosum, Steud. Syn. Pl. Gram. 72.—Japon.
densiflorum, *Willd. ex Spreng. Syst.* i. 320.—Am. austr.
densiglume, Steud. Syn. Pl. Gram. 41, sphalm. = uniglume.
densispica, Poir. Encyc. Suppl. iv. 273 = Pennisetum setosum.
densum, Muhl. Desc. Gram. 122 = dichotomum.
denudatum, Kunth. Rév. Gram. i. 32 = violascens?
depauperatum, *Muhl. Desc. Gram.* 112.—Am. bor.
desertorum, *A. Rich. Tent. Fl. Abyss.* ii. 365.—Abyss.
despiciens, *Link, Hort. Berol.* i. 207.—Hab.?
Despreauxii, *Steud. Syn. Pl. Gram.* 58.—Afr. trop.
deustum, Brickell & Enslin, ex Muhl. Desc. Gram. 119 = dichotomum.
deustum, *Thunb. Prod. Pl. Cap.* 19.—Afr. austr.
diagonale, *Nees, Fl. Afr. Austr.* 23.—Afr. austr.
diamesum, *Steud. Syn. Pl. Gram.* 42.—Afr. trop.
diandrum, Kunth, Rév. Gram. ii. t. 110 = laxum.
dichotomiflorum, Michx. Fl. Bor. Am. i. 48 = autumnale.
dichotomum, Forsk. Fl. Aegypt. Arab. 19 = Pennisetum dichotomum.
dichotomum, *Linn. Sp. Pl.* 58.—Am. bor.
dictyoneurum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 329. t. 8.—Nubia.
didactylum, Kunth, Rév. Gram. i. 33; Enum. Pl. i. 84 = sanguinale.
didymostachyum, *Steud. Syn. Pl. Gram.* 97.—Chili.
difforme, *Roth, Nov. Pl. Sp.* 52.—Ind. or.
diffusulum, *Salzm. ex Steud. Syn. Pl. Gram.* 89.—Am. trop.
diffusum, Pursh, Fl. Am. Sept. i. 68 = capillare.
diffusum, *Sw. Prod. Veg. Ind. Occ.* 23.—Ind. occ.; Afr. austr.
Digitaria, Latirr. Ami des Champs (1825), 339 = Paspalum distichum.
digitarioides, Carpenter, ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 410 = Curtisii.
digitarioides, Rasp. ex Steud. Nom. ed. II. ii. 255 = Paspalum Michauxianum.
digitatum, Gilib. Exercit. 517 = P. Crus-galli.
dilatatum, *Steud. Syn. Pl. Gram.* 39.—Ind. or.
dimidiatum, Burm. f. Fl. Ind. 25. t. 8. f. 3 = Trachys mucronata.
dimidiatum, Linn. Sp. Pl. 57 = Stenotaphrum complanatum.
dimidiatum, Walt. Fl. Carol. 72 = carolinianum.
dioecum, *Spreng. Syst.* i. 322.—Bras.
disciferum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 488.—Mexic.
discolor, Muhl. Desc. Gram. 114 = dichotomum.
discolor, *Spreng. Fl. Hal. Mant.* 31.—Am. bor.
discolor, *Trin. ex Nees, Agrost. Bras.* 144.—Bras.
discrepans, *Doell, in Mart. Fl. Bras.* ii. II. 252.—Am. trop.
disjunctum, *Steud. Syn. Pl. Gram.* 417.—Madag.
dispar, *Steud. l. c.* 96.—Nepal.
dispermum, Lam. Illustr. i. 173 = Isachne disperma.
dispersum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1885) 282 = rugulosum.
dissectum, Linn. Sp. Pl. 57 = Paspalum scrobiculatum.
dissitiflorum, *Steud. Nom.* ed. II. ii. 255.—Mexic.
distachyum, *Linn. Mant.* i. 138.—As. et Austral. trop.
distans, Salzm. in Steud. Syn. Pl. Gram. 61 = plantagineum, *Link.*
distans, Trin. Sp. Gram. Ic. ii. t. 172 = flavidum.
distans, Willd. ex Spreng. Syst. i. 305 = pilosum.
distantiflorum, *A. Rich. Fl. Cub. Fanerog.* ii. 304.—Cuba.
distichophyllum, Nees, Agrost. Bras. 234 = subulatum.
distichophyllum, *Trin. Gram. Panic.* 147.—Afr. trop.
distichum, Gaudich. ex Steud. Nom. ed. II. ii. 255 = glabrum.
distichum, Lam. Encyc. iv. 731 (err. typ. 741) = pilosum.
divaricatissimum, *R. Br. Prod.* 192.—Austral.
divaricatum, Kunth, ex Griseb. Fl. Brit. W. Ind. 552 = martinicense.
divaricatum, *Linn. Syst.* ed. X. 871.—Am. austr.
divaricatum, Michx. Fl. Bor. Am. i. 50 = debile.
divaricatum, Presl, ex Griseb. Fl. Brit. W. Ind. 551 = latifolium.

PANICUM:—
divergens, Afzel. ex Sw. Adnot. Bot. 5 = Afzelii.
divergens, *H. B. & K. Nov. Gen. et Sp.* i. 102.—Ecuador.
divergens, Muhl. Desc. Gram. 120 = autumnale.
diversinerve, *Nees, Fl. Afr. Austr.* 23.—Afr. austr.
divisum, Forsk. ex J. F. Gmel. Syst. 156 = Pennisetum setosum.
domingense, *Zuccagni, in Roem. Collect.* 123.—Ins. S. Doming.
donacifolium, Raddi, Agrost. Bras. 44 = Ichnanthus panicoides.
Dregeanum, *Nees, Fl. Afr. Austr.* 42.—Afr. austr.
Drummondii, *Nees, ex Steud. Syn. Pl. Gram.* 63.—Am. bor.
dubium, Lam. Encyc. iv. 743 = Isachne miliacea.
dubium, Sieber, ex Steud. Nom. ed. II. ii. 256 = P. Crus-galli.
dubium, Steud. l. c. = Oplismenus dubius.
Duchaissingii, Steud. Syn. Pl. Gram. 93 = lanatum.
dumetorum, *Rich. ex Steud. l. c.* 49.—Ind. occ.
Dumus, *Desv. Opusc.* 88.—Am. calid.
durum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 533.—Ind. occ.
eburneum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 268.—Nepal.
echinatum, Willd. Enum. Hort. Berol. 1032 = P. Crus-galli.
echinatum, Willd. ex Doell, in Mart. Fl. Bras. ii. II. 193 = uncinatum.
Echinochloa, Steud. Nom. ed. II. i. 316, in syn., sphalm. = seq.
Echinolaena, Nees, Agrost. Bras. 128 = Echinolaena hirta.
Ecklonii, *Nees, Fl. Afr. Austr.* 43.—Afr. austr.
effusum, *R. Br. Prod.* 191.—Austral.
elatius, Kunth, Rév. Gram. i. 38 = altissimum.
elatius, Linn. f. Suppl. 107 = Oplismenus compositus.
elatum, Willd. ex Steud. Nom. ed. II. ii. 256 = decolorans.
elegans, *Wight & Arn. ex Steud. l. c.*—Zeylan.
elephantipes, *Nees, Agrost. Bras.* 165.—Bras.
Elliotianum, Schult. Mant. ii. 256 = gibbum.
Elliotii, Spreng. ex Steud. Nom. ed. II. ii. 256 = dichotomum.
Elliottii, Trin. ex Nees, Agrost. Bras. 170 = proliferum.
elongatum, Poir. Encyc. Suppl. iv. 278 = P. Crus-ardeae.
elongatum, Pursh, Fl. Am. Sept. i. 69 = virgatum.
elongatum, Salisb. Prod. 18 = Setaria italica.
Elythroblepharum, Steud. in Zoll. Syst. Verz. Pfl. Ind. Archip. 54; Miq. Fl. Ind. Bat. iii. 439 = barbatum.
emergens, *Doell, in Mart. Fl. Bras.* ii. II. 269.—Am. trop.
emergens, *Hochst. in Flora*, xxxviii. (1855) 196.—Abyss.
eminens, Steud. Syn. Pl. Gram. 43 = barbatum.
enneaneurum, *Griseb. in Goett. Abh.* xix. (1874) 261.—Reg. Argent.
ensifolium, Baldw. in Ell. Sketch, i. 126 = dichotomum?
Enslini, Trin. Gram. Panic. 230 = dichotomum.
equinum, Salzm. ex Steud. Syn. Pl. Gram. 67 = molle.
Equisetum, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 206 = maximum, *Jacq.*
equitans, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 365.—Abyssin.
erianthum, Poir. Encyc. Suppl. iv. 284 = P. Crus-galli.
erichrysoides, Pritz. Ic. Ind. i. 795, sphalm. = seq
eriochrysoides, *Nees, ex Trin. Gram. Panic.* 160.—Bras.
eriogonum, Schrad. Pl. Hort. Gotting. fasc. 1; ex Schult. Mant. ii. 253 = sanguinale.
eriophorum, Link, ex Nees, Agrost. Bras. 148 = Ichnanthus Hoffmannseggii.
eriophorum, Schult. Mant. ii. 591 = dichotomum.
erubescens, Willd. Enum. Hort. Berol. 1031 = Pennisetum erubescens.
eruciforme, Sibth. & Sm. Fl. Graec. i. 44. t. 59 = P. Isachne.
erythrospermum, Vahl, ex Hornem. Hort. Hafn. i. 82 = Setaria italica.
esculentum, *A. Br. Ind. Sem. Hort. Berol.* (1861) *App.* 3.—Japon (cult.).
Esenbeckii, Steud. Syn. Pl. Gram. 90 = Olyra nana.
euchroum, Steud. l. c. 98 = montanum.
exaratum, *Trin. Gram. Panic.* 160.—Bras.
exasperatum, *Nees, ex Steud. Syn. Pl. Gram.* 62.—Afr. trop.
excelsum, Nees, Agrost. Bras. 180 = rivulare.

PANICUM:—
excelsum, Spreng. ex Steud. Nom. ed. II. ii. 256 = Paspalum barbatum.
excurrens, Trin. Sp. Gram. Ic. i. t. 89 = plicatum.
exiguiflorum, *Griseb. Cat. Pl. Cub.* 234.—Cuba.
exile, Steud. Nom. ed. II. ii. 256 = Ichnanthus candicans.
expansum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* i. 488.—Mexic.
expansum, *Trin. ex Steud. Nom.* ed. II. ii. 256 (Quid?).—Mexic.
extensum, *Desv. Opusc.* 84.—Aegypt.
extensum, Nees, ex Steud. Syn. Pl. Gram. 39 = dilatatum.
extensum, *Steud. l. c.* 72.—Ins. Philipp.
falcatum, *J. F. Gmel. Syst.* 163.—Guian.
falciferum, *Trin. Gram. Panic.* 127.—Afr. trop.
fallax, Kunth, Rév. Gram. i. 35 = decipiens.
fallax, Spreng. Nachr. I. Bot. Gart. Halle, 29 = sanguinale.
falsum, *Steud. Syn. Pl. Gram.* 67.—Cuba.
fasciculare, Schrad. ex Kunth, Enum. Pl. i. 80 = Schraderi.
fasciculatum, Griseb. Fl. Brit. W. Ind. 549 = maximum, *Jacq.*
fasciculatum, Hort. ex Trin. Gram. Panic. 76 = Schraderi.
fasciculatum, Lam. Encyc. iv. 731 (err. typ. 741), = fasciculiflorum.
fasciculatum, *Sw. Prod. Veg. Ind. Occ.* 22.—Am. calid.
fasciculiflorum, *Steud. Nom.* ed. II. ii. 256.—Jamaic.
fastigiatum, Poir. Encyc. Suppl. iv. 277 = fasciculatum.
fatmense, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 256; et Syn. Pl. Gram. 100 (Eriochloae sp.).—Ind. or.; Arab.
fatuum, *Fig. & De Not. in Mém. Acc. Torin.* Ser. II. xiv. (1854) 346.—Nubia.
fenestratum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 361.—Abyssin.
ferrugineum, Kunth, Rév. Gram. i. 39 = phaeothrix.
ferrugineum, Trin. Gram. Panic. 159 = eriochryseoides.
festucoides, *Poir. Encyc. Suppl.* iv. 283.—Ind. or.
Figarei, *Tod. Ind. Sem. Hort. Panorm.* (1866) 33.—Aegypt.
filamentosum, Pers. Syn. i. 83 = brevifolium.
filiforme, Garcke, Fl. Deutschl. ed. II. 360 = glabrum.
filiforme, Jacq. Obs. iii. 18. t. 70 = sanguinale.
filiforme, *Linn. Sp. Pl.* 57.—Am. bor.
filiforme, Poir. Voy. Barb. ii. 93 = debile.
fimbriatum, Kunth, Rév. Gram. i. 33; J. & C. Presl, Rel. Haenk. i. 298 = sanguinale.
fimbriatum, Willd. ex Spreng. Syst. i. 316 = divergens.
firmandum, Steud. Syn. Pl. Gram. 418 = microcarpon?
firmifolium, Trin. ex Kunth, Enum. Pl. i. 110 = cyanescens.
firmum, Kunth, Rév. Gram. i. 37 = Oryzopsis canadensis.
fistulosum, *Hochst. ex Steud. Syn. Pl. Gram.* 71.—Am. trop.
flabellatum, *Fourn. in Bull. Soc. Bot. Fr.* xxvii. (1880) 293.—Nicaragua.
flabellatum, Steud. Syn. Pl. Gram. 53 = Setaria sulcata.
flaccidum, Steud. Nom. ed. II. ii. 256 = Oplismenus setarius.
flavescens, Moench, Meth. 206 = Setaria glauca.
flavescens, Sieber, ex Presl, Bot. Bemerk. 120 = Eriochloa polystachya.
flavescens, *Sw. Prod. Veg. Ind. Occ.* 23.—Ind. occ.
flavidum, *Retz. Obs.* iv. 15.—As. trop.; Austral.
flavum, Nees, ex Trin. Gram. Panic. 162 = Setaria flava.
flexirameum, Steud. Nom. ed. II. ii. 254, in syn., sphalm. = *flexuosum*, Willd.
flexuosum, Retz. Obs. iii. 9 = Isachne ventricosa.
flexuosum, Willd. ex Schult. Mant. ii. 591 = capense.
floribundum, Rich. ex Lam. Encyc. iv. 742 = cayennense.
floribundum, *Willd. ex Spreng. Syst.* i. 306.—Ind. or.
floridanum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 248.—Am. bor.
fluitans, Brickell, ex Muhl. Desc. Gram. 126 = hydrophilum.
fluitans, *Retz. Obs.* iii. 8.—Geront. trop.
fluitans, Retz. l. c. v. 18, in obs. = paspaloides.
fluviicola, *Steud. Syn. Pl. Gram.* 89.—Afr. trop.
foliaceum, Steud. Nom. ed. II. ii. 256, sphalm. = *loliaceum*.
foliolosum, *Wall. Cat.* n. 8680.—Burma.
foliosum, *R. Br. Prod.* 191.—Malaya; Austral.

PANICUM:—
Forbesianum, Nees, in Steud. Syn. Pl. Gram. 98 (Setariae sp.).—Ind or.
fragile, Kunth, Enum. Pl. i. 36 = autumnale.
Francavillanum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 489.—Mexic.
Francoi, Steud. Syn. Pl. Gram. 44 = Oplismenus Burmannii.
fraudulentum, *Steud. l. c.* 419.—Ins. Borbon.
frondescens, G. F. W. Mey. Prim. Fl. Esseq. 56 = stoloniferum.
frondescens, Trin. Sp. Gram. Ic. ii. t. 218 = Ichnanthus panicoides.
frumentaceum, Roxb. Hort. Beng. 7; Fl. Ind. i. 304 = colonum.
frumentaceum, Salisb. Prod. 18 = Sorghum vulgare.
fruticosum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 207 = latifolium.
fugax, *J. F. Gmel. Syst.* i. 162.—Ind. or.?
Funckianum, *Steud. Syn. Pl. Gram.* 77.—N. Granat.
fuscatum, Presl, ex Steud. Nom. ed. II. ii. 256 = fasciculatum.
fuscescens, Willd. ex Steud. l. c. = Setaria imberbis.
fusciflorum, Steud. Syn. Pl. Gram. 93 = Anthaenantia lanata.
fuscinode, Steud. l. c. 418.—Ins. Comor.
fusco-rubens, Lam. Illustr. i. 171 = fasciculatum.
fusco-viride, *Steud. Syn. Pl. Gram.* 80.—Hab.?
fuscum, Doell, in Mart. Fl. Bras. ii. II. 130 = violascens.
fuscum, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. (1842) 239 = carthaginense.
fuscum, Sieber, ex Griseb. Fl. Brit. W. Ind. 552 = martinicense.
fuscum, *Sw. Prod. Veg. Ind. Occ.* 23.—Jamaic.
Galli, *Thunb. Prod. Pl. Cap.* 18.—Afr. austr.
Gardneri, *Thw. Enum. Pl. Zeyl.* 359.—Zeylan.
Gaudichaudii, *Kunth, Rév. Gram.* ii. t. 106.—Ins. Marian.
Gavanianum, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 137 = lanatum.
Gayanum, *Kunth, l. c.* i. 239. t. 31.—Afr. trop.
geminatum, Forsk. Fl. Aegypt. Arab. 18 = paspaloides.
geminatum, Hochst. in Flora, xxiv. (1841) I. Intell. 19 = Hochstetterianum.
geniculatum, Forsk. ex Spreng. Syst. i. 309, in syn. = fluitans.
geniculatum, Lam. Encyc. iv. 727 (err. typ. 737) = Setaria geniculata.
geniculatum, Muhl. Desc. Gram. 123; et ex Spreng. Syst. i. 319, in syn. = proliferum.
geniculatum, Thunb. Fl. Cap. i. 388 = Pennisetum Thunbergi.
georgicum, Spreng. Syst. i. 308 = Eriochloa punctata.
germanicum, Mill. Gard. Dict. ed. VIII. n. 1 = Setaria italica.
Ghiesbreghtii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 489.—Mexic.
gibbosum, *R. Br. Prod.* 193.—Austral.
gibbum, *Ell. Sketch*, i. 116.—Am. bor.
gibbum, Steud. Syn. Pl. Gram. 87 = trigonum.
giganteum, Scheele, in Linnaea, xxii. (1849) 340 = virgatum.
Gilesii, *Benth. Fl. Austral.* vii. 477.—Austral.
glaberrimum, Steud. Syn. Pl. Gram. 94 = gymnocarpum?
glabrescens, *Steud. l. c.* 71.—Afr. trop.
glabripes, *Doell, in Mart. Fl. Bras.* ii. II. 216.—Am. trop.
glabrum, Ell. ex Steud. Nom. ed. II. ii. 257 = gibbum.
glabrum, *Gaud. Agrost.* i. 22.—Reg. bor. temp. et trop.
gladiatum, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 170 = Oplismenus dubius.
glandulosum, Nees, ex Trin. Gram. Panic. 174 = uncinatum.
glareae, F. Muell. in Linnaea, xxv. (1852) 445 = lanatum.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* i. 104.—Am. austr.
glaucescens, Lam. Encyc. iv. 749 = Arundinaria glaucescens.
glaucescens, Nees, Agrost. Bras. 100 = sanguinale.
glaucescens, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 157 = Setaria imberbis.
glaucescens, Willd. ex Doell, l. c. 275 = Isachne disperma.
glaucum, Linn. Sp. Pl. 56 = Setaria glauca.
glaucum, Rupr. ex Galeotti, in Bull. Acad. Brux ix. (1842) 241 = Setaria geniculata.

PANICUM:—
glaucum, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 157 = Setaria imberbis.
globulare, Steud. Nom. ed. II. ii. 51 = Setaria globularis.
globuliferum, *Steud. Syn. Pl. Gram.* 51.—Am. trop.
glomeratum, Moench, Meth. 207 = Setaria italica.
glomeratum, Trin. Gram. Panic. 170 = conglomeratum.
glumaepatulum, *Steud. Syn. Pl. Gram.* 41.—Ins. Philipp.
glumare, *Trin. Gram. Panic.* 143.—N. Zel.
glutinosum, Lam. Illustr. i. 174 = latifolium.
glutinosum, Presl, ex Griseb. Fl. Brit. W. Ind. 552 = martinicense.
glutinosum, *Sw. Prod. Veg. Ind. Occ.* 24.—Am. austr.
Gmelini, [Honck,] Verz. Aller Gew. Teutschl. 164 (Quid ?).—Europ.
gonatodes, Steud. Syn. Pl. Gram. 95 = Isachne australis.
gongylodes, Jacq. Eclog. Gram. 30. t. 21 = maximum, *Jacq.*
gonyrrhizum, Steud. Syn. Pl. Gram. 44 (= *Orthopogon gonyrrhizus*).—Java.
gossypinum, Hook. & Arn. Bot. Beech. Voy. 100 = montanum.
gossypinum, *A. Rich. Tent. Fl. Abyss.* ii. 365.—Afr. bor. et austr.
Gouini, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 489.—Mexic.
gracile, *R. Br. Prod.* 190.—Austral.
gracile, Nees, ex Spreng. Syst. iv. Cur. Post. 33 = distachyum.
gracile, Nym. Syll. Suppl. 70 = sanguinale.
gracile, Willd. ex Spreng. Syst. i. 320 = micranthum.
gracilentum, Poir. Encyc. Suppl. iv. 276 = madagascariense.
gracilescens, Desv. ex Poir. l. c. 279 = microcarpon ?
grandiflorum, Trin. ex Nees, Agrost. Bras. 143 = oryzoides.
grandifolium, *Doell, in Mart. Fl. Bras.* ii. II. 195.—Am. trop.
granulare, Boj. Hort. Maurit. 364 = fluitans.
granulare, Lam. Illustr. i. 170 = flavidum.
granuliferum, H. B. & K. Nov. Gen. et Sp. i. 19 = parvifolium.
grossarium, Griff. Notul. iii. 36; Ic. Pl. Asiat. t. 139. f. 113 = colonum.
grossarium, Koen. in Naturf. xxiii. (1788) 205 = prostratum *vel* trypheron.
grossarium, *Linn. Syst.* ed. X. 871.—Am. austr.
grossarium, Roxb. Hort. Beng. 82; Fl. Ind. i. 297 = paspaloides ?
grossularium, Thunb. Fl. Jap. 48 = bisulcatum.
grossum, Salisb. Prod. 18 = P. Crus-galli.
grumosum, *Nees, Agrost. Bras.* 182.—Am. austr.
guadaloupense, Steud. Syn. Pl. Gram. 61 = molle.
guadalupense, *Spreng. ex Steud. Nom.* ed. II. ii. 256.—Ins. Guadalup.
guaraniticum, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1813) 129.—Reg. Argent.
guayaquilense, *Steud. Syn. Pl. Gram.* 85.—Ecuador.
guineense, Desv. ex Poir. Encyc. Suppl. iv. 279 = ovalifolium.
gymnocarpum, *Ell. Sketch*, i. 117.—Am. bor.
haematodes, Presl, Fl. Sic. i. 43 = colonum.
Haenkeanum, *J. & C. Presl, Rel. Haenk.* i. 304.—Mexic.
Hallii, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 61.—Texas.
hamatum, *Desv. Opusc.* 83.—Bras.
Hamiltonii, *Kunth, Rév. Gram.* i. 33.—Ind. occ.
Hasskarlii, Steud. ex Zoll. Syst. Verz. Ind. Archip. 54 = amplexicaule.
havaïense, Reichardt, in Sitzb. Akad. Wien, lxxvi. I. (1878) 723 = nephelophilum.
hebotes, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 301.—Bras.
Helopus, Trin. in Spreng. Neue Entdeck. ii. 84 = javanicum.
helvolum, Linn. f. Suppl. 107 = Setaria glauca.
hemignostum, Steud. Syn. Pl. Gram. 77 = Ichnanthus pallens.
hemitomum, Schult. Mant. ii. 227 = carolinianum.
hermaphroditum, *Steud. Syn. Pl. Gram.* i. 67.—As. trop.; Austral.
heteranthum, Link, Hort. Berol. i. 212 = uncinatum.

PANICUM:—
heteranthum, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 174 = barbatum.
heterophyllum, Bosc, ex Nees, Agrost. Bras. 227 = dichotomum.
heterophyllum, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 160, nomen = dichotomum ?
heterophyllum, Spreng. Neue Entdeck. iii. 13 = pilosum.
heterophyllum, Sw. Adnot. Bot. 6 = Kalmii.
Heynii, *Roth, Nov. Sp. Pl.* 49.—Ind. or.
hians, Ell. Sketch, i. 118 = debile.
hians, Spruce, ex Griseb. Fl. Brit. W. Ind. 548 = stenodes.
hirsutissimum, *Steud. Syn. Pl. Gram.* 72.—Afr. austr.
hirsutum, Heyne, ex Wall. Cat. n. 8734 B = prostratum.
hirsutum, Koen. ex Roxb. Hort. Beng. 7; Fl. Ind. i. 300 = javanicum.
hirsutum, Lam. Encyc. iv. 741 = rugulosum.
hirsutum, *Sw. Fl. Ind. Occ.* i. 173.—Ind occ.
hirsutum, Vahl, ex Griseb. Fl. Brit. W. Ind. 548 = diffusum.
hirsutum, Willd. ex Spreng. Syst. i. 317 = cayennense.
hirtellum, All. Fl. Pedem. ii. 240 = Oplismenus undulatifolius.
hirtellum, Burm. f. Fl. Ind. 24. t. 13; Thunb. Fl. Jap. 46 = Oplismenus Burmannii.
hirtellum, Linn. Syst. ed. X. 870 = Oplismenus loliaceus.
hirtellum, Walt. Fl. Carol. 72 = P. Crus-Galli.
hirticalycinum, Steud. Nom. ed. II. ii. 257 = seq.
hirticalycum, Bosc, ex Spreng. Syst. i. 315 = P. Crus-galli.
hirticaulum, J. & C. Presl, Rel. Haenk. i. 308 = cayennense.
hirticulmum, Doell, in Mart. Fl. Bras. ii. II. 219 = cayennense.
hirtum, *Lam. Encyc.* iv. 741.—Am. trop.
hirtum, Nees, Agrost. Bras. 212 = brevifolium.
hirtum, Roth, Nov. Sp. Pl. 46 = Rothii.
hirtum, Willd. ex Kunth, Enum. Pl. i. 112 = Isachne miliacea.
hispidulum, Lam. Illustr. i. 171; Retz. Obs. v. 18 = P. Crus-galli.
hispidum, Forst. f. Prod. 7; Muhl. Desc. Gram. 107 = P. Crus-galli.
Hochstetteri, *Steud. Syn. Pl. Gram.* 90.—Abyss.
Hochstetterianum, *A. Rich. Tent. Fl. Abyss.* ii. 369.—Abyss.
Hoffmannseggii, Roem. & Schult. Syst. ii. 450 = Ichnanthus Hoffmannseggii.
holciforme, *Steud. Nom.* ed. II. ii. 257.—Mexic.
holcoides, Hort. ex Trin. Gram. Panic. 163 = P. Pseudoholcus.
holcoides, Jacq. Eclog. Gram. t. 22 = Setaria glauca, Pennisetum barbatum.
holcoides, Roxb. Hort. Beng. 7; Fl. Ind. i. 285 = Pennisetum holcoides.
holochrysum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 195 = Paspalum radiatum.
holosericeum, *R. Br. Prod.* 190.—Austral.
holosericeum, Nees, Fl. Afr. Austr. 30 = gossypinum.
homonymum, Steud. Syn. Pl. Gram. 48 = rhachitrichum.
Hookerianum, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 326.—N. Caled.
hordeaceum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Chamaeraphis hordeacea.
hordeiforme, Thunb. Fl. Jap. 46 = Pennisetum compressum.
hordeoides, *Lam. Illustr.* i. 170.—Afr. trop.
horizontale, G. F. W. Mey. Prim. Fl. Esseq. 54 = sanguinale.
horridum, Salzm. ex Steud. Syn. Pl. Gram. 47 = P. Crus-galli.
horticolum, *Steud l. c.* 95.—Hab. ?
Hostii, Bieb. Fl. Taur. Cauc. iii. 57 = P. Crus-galli.
humedorum, *Buch.-Ham. in Wall. Cat.* n. 8721.—Ind. or.
humifusum, Kunth, Rév. Gram. i. 33 = glabrum.
humile, Aitch. Cat. Punjab Pl. 159 = hydaspicum.
humile, *Nees, ex Steud. Syn. Pl. Gram.* 84.—Ind. or.
humile, Thunb. ex Trin. Gram. Panic. 164 = Setaria viridis.
humile, Trin. l. c. 167 = Setaria verticillata.
hybridum, *Trin. ex Steud. Nom.* ed. II. ii. 257.—Ins. Guadalup.
hydaspicum, *Edgew. in Journ. Linn. Soc.* vi. (1862) 207.—Ind. or.

PANICUM:—
hydrophilum, Salzm. ex Steud. Nom. ed. II. ii. 71 = Setaria brachiata.
hydrophilum, *Schult. Mant.* ii. 237.—Am. bor.
hydrophilum, Trin. ex Nees, Agrost. Bras. 208 = gibbum.
hygrocharis, F. Muell. Fragm. viii. 195 = obseptum.
hygrocharis, Steud. Syn. Pl. Gram. 72 = repens.
hygrophilum, Salzm. ex Steud. l. c. 71 = Setaria brachiata.
hymeniochilum, *Nees, Fl. Afr. Austr.* 46.—Afr. austr.
Hystrix, *Steud. Syn. Pl. Gram.* 95.—Afr. trop.
ichnanthoides, *Fourn. ex Hemsl. Biol. Centr. Am.* iii. 491.—Mexic.
Ichnanthus, Nees, Agrost. Bras. 149 = Ichnanthus panicoides.
ichnodes, *Griseb. Fl. Brit. W. Ind.* 551.—Ind. occ.
ignoratum, Kunth, Rév. Gram. i. 55. t. 20 = Anthaenantia villosa.
illinoniense, *Desv. Opusc.* 91.—Am. bor.
imbecille, Trin. Sp. Gram. Ic. ii. t. 191 = Oplismenus undulatifolius.
imbelle, Spreng. Syst. i. 321 = tenellum.
imberbe, Poir. Encyc. Suppl. iv. 272 = Setaria viridis.
imberbe, *Schlecht. in Linnaea*, xxxi. (1861–62) 431.—Am. bor.; Bras.
immersum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 197 = Paspalum aureum.
imperfectum, Roxb. ex Kunth, Enum. Pl. i. 61 = Paspalum compactum.
impressum, Nees, Agrost. Bras. 247 = Setaria impressa.
inaequale, *Fourn. ex Hemsl. Biol. Centr. Am.* iii. 490.—Mexic.
inaequale, *F. Muell. Fragm.* viii. 189.—Austral.
incanum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 60.—Afr. trop.
incertum, Bosc, ex Steud. Nom. ed. II. ii. 258 = colonum.
incomptum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 344.—Nubia.
incomptum, F. Muell. Fragm. x. 76 = Mitchellii ?
incomtum, Trin. Gram. Panic. 200 = sarmentosum.
inconstans, Trin. ex Nees, Agrost. Bras. 132 = Ichnanthus inconstans.
incurvum, Linn. Herb. ex Munro, in Journ. Linn. Soc. vi. (1862) 39 = indicum.
indicum, *Linn. Mant.* ii. 184; *Retz. Obs.* iii. 9.—Geront. trop.
indicum, Mill. Gard. Dict. ed. VIII. n. 3 = Setaria italica.
indutum, *Steud. Syn. Pl. Gram.* 68.—Afr. trop.
infestum, *Anderss. in Peters, Reise Mossamb. Bot.* 546.—Afr. trop.
infidum, Steud. Syn. Pl. Gram. 63 = ambiguum, *Trin.* ?
insculptum, *Steud. l. c.* 49.—Afr. trop.
insigne, *Steud. Nom.* ed. II. ii. 258.—Abyss.
insulare, G. F. W. Mey. Prim. Fl. Esseq. 60 = lanatum.
insularum, Steud. Syn. Pl. Gram. 61 = prostratum.
insulicola, Steud. l. c. 78 = aureum.
intermedium, Roth, Nov. Pl. Sp. 47 = Setaria glauca.
intermedium, Salzm. ex Steud. Syn. Pl. Gram. 51 = Setaria macrostachya.
intermedium, Vahl, ex Hornem. Hort. Hafn. i. 82 = Setaria italica.
interruptum, Steud. Syn. Pl. Gram. 66, partim = indicum.
interruptum, *Willd. Sp. Pl.* i. 341.—Ind. or.; Afr. austr.
inundatum, Kunth, Rév. Gram. i. 34 = interruptum.
involucratum, Roxb. Hort. Beng. 7; Fl. Ind. i. 284 = Pennisetum typhoideum.
involutum, Torr. Fl. U. St. 144 = depauperatum.
Isachne, *Roth, Nov. Pl. Sp.* 54.—Reg. Mediterr.; Ind. or.
isachnoides, *Munro, in Journ. Bot.* vii. (1869) 178, nomen.—Ins. Sandvic.
ischaemoides, Herb. Heyne, ex Wall. Cat. n. 8679 = Chamaeraphis spinescens.
ischaemoides, Koen. ex Trin. Gram. Panic. 97 = Paspalum furcatum.
ischaemoides, Retz. Obs. iv. 17 = repens.
Ischaemum, Schreb. ex Schweigg. Spec. Fl. Erlang. (1804) 16 = glabrum.
ischnocaulon, Steud. Syn. Pl. Gram. 45 = Oplismenus gracilis.
isocalycinum, *G. F. W. Mey. Prim. Fl. Esseq.* 59.—Guian.

PANICUM:—
italicum, Linn. Sp. Pl. 56 = Setaria italica.
italicum, Ucria, Hort. Reg. Panorm. 54 = Setaria verticillata.
Itieri, Steud. Syn. Pl. Gram. 51 = Setaria Itieri.
Jacobinae, *Steud. l. c.* 55.—Am. austr.
Jahnii, *Steud. l. c.* 54.—Ind. or.
jamaicense, Steud. Nom. ed. II. ii. 257 = spectabile.
japonicum, Kunth, Rév. Gram. i. 205, in syn., sphalm. = javanicum.
japonicum, Steud. in Flora, xxix. (1846) 18 = Oplismenus Burmannii.
Jardini, *Steud. Syn. Pl. Gram.* 95.—Afr. trop.
javanicum, *Poir. Encyc. Suppl.* iv. 274.—Geront. trop.
javanicum, Thw. Enum. Pl. Zeyl. 358 = ambiguum, *Trin.*
javanum, Nees, ex Buese, in Miq. Pl. Jungh. 376 = Isachne australis.
jejunum, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 71 = debile.
Johannae, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 39 = indicum.
jubatum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 331.—Nubia.
jubiflorum, Trin. Gram. Panic. 180 = gracile.
jumentorum, Pers. Syn. i. 83 = maximum, *Jacq.*
jumentorum, A. Rich. Tent. Fl. Abyss. ii. 373 = trypheron.
junceum, *Nees, Agrost. Bras.* 159.—Bras.
junceum, Trin. Gram. Panic. 220 = depauperatum.
junciforme, Steud. Syn. Pl. Gram. 82 = caricoides.
Junghuhnianum, *Nees, ex Steud. l. c.* 63.—Java.
juniperinum, Nees, Agrost. Bras. 236 = lycopodioides.
Kalmii, *Sw. Adnot. Bot.* 6.—Am. bor.
Kappleri, Steud. Syn. Pl. Gram. 90 = micranthum.
Kegelii, *Steud. l. c.* 65.—Guatemala.
Kleinianum, Nees, ex Anderss. in Walp. Ann. vi. 945 = plicatum.
Koenigii, Herb. Wight, ex Nees, Fl. Afr. Austr. 30 = P. Isachne.
Koenigii, Spreng. Syst. i. 311 = javanicum.
Kohautianum, *Presl, Bot. Bemerk.* 119.—Ins. Martinic.
Kotschyanum, Hochst. in Flora, xxv. (1842) 1. Beibl. 133 = ramosum.
Kraussii, *Steud. Syn. Pl. Gram.* 45.—Afr. austr.
Kunthianum, *Wight & Arn. ex Steud. l. c.* 96.—Java.
Kunthii, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 490 = arenarium.
Kunthii, *Steud. Nom.* ed. II. ii. 258.—Ind. or.
lachnanthum, *A. Gray, in Pacif. Rail. Rep.* vii. 21.—Am. bor. occ.
lachnanthum, *Hochst. in Flora*, xxxviii. (1855) 195.—Abyss.
lachneum, Nees, Agrost. Bras. 248 = Setaria setosa.
lachnophyllum, *Benth. Fl. Austral.* vii. 486.—Austral.
lacunarium, F. Muell. in Trans. Vict. Inst. i. (1855) 47 = P. Crus-galli.
laetum, *Kunth, Rév. Gram.* ii. t. 113.—Afr. trop.
laeve, Lam. Illustr. i. 172 = maximum, *Jacq.*
laevicaule, Steud. Syn. Pl. Gram. 82, sphalm. = *laevinode.*
laevigatum, Ell. Sketch, i. 112 = Setaria glauca.
laevigatum, Lam. Fl. Fr. iii. 578 = Setaria viridis.
laevigatum, Muhl. Desc. Gram. 100 = Setaria laevigata.
laevinode, Lindl. in Mitch. Three Exped. i. 238 = decompositum.
Lagopus, Willd. ex Spreng. Syst. i. 265 = Saccharum cayennense.
Lagotis, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1835) 326 = Ichnanthus petiolatus.
Lamarckii, *Kunth, Rév. Gram.* i. 41.—Hab.?
lanatum, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 269.—Am. trop.
lanceaefolium, *Desv. Opusc.* 83.—Bras.
lanceaefolium, *Desc. l. c.* 94 (sp. diversa).—Bras.
lancearum, Trin. Agrost. Bras 246 = dichotomum.
lanceolatum, Retz. Obs. v. 17 = Oplismenus compositus.
lancifolium, Salzm. ex Doell, in Mart. Fl. Bras. ii. 11. 250 = sciurotis.
Langei, *Fourn. in Hemsl. Biol. Centr. Am. Bot.* iii. 490.—Mexic.
laniflorum, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 410 = lanatum.
lanuginosum, Ell. Sketch, i. 123 = dichotomum.

PANICUM:—
lanuginosum, J. & C. Presl, Rel. Haenk. i. 306 = molliconum.
lappaceum, Willd. ex Spreng. Syst. i. 306 = Oplismenus Burmannii.
lasianthum, Trin. Sp. Gram. Ic. iii. t. 245 = rugulosum.
laterale, Nees, ex Doell, in Mart. Fl. Bras ii. 11. 329 = Olyra nana.
laterale, *J. & C. Presl, Rel. Haenk.* i. 305.—Am. trop.
laticomum, *Nees, Fl. Afr. Austr.* 43.—Afr. austr.
latifolium, H. B. & K. Nov. Gen. et Sp. i. 100 = oryzoides.
latifolium, *Linn. Sp. Pl.* 58.—Am. bor.; Ind. occ.
latiglume, *Doell, in Mart. Fl. Bras.* ii. 11. 257.—Am. trop.
latissimum, *Mikan, ex Trin. in Spreng. Neue Entdeck.* ii. 87.—Bras.
latum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 61.—Afr. trop.
laxiflorum, Lam. Encyc. iv. 748 = dichotomum.
laxum, *Sw. Prod. Veg. Ind. Occ.* 23.—Am. trop.
Leandri, Trin. Sp. Gram. Ic. iii. t. 335 = plantagineum.
leersioides, *Hochst. in Flora*, xxxviii. (1855) 196.—Abyss.
leiocarpon, Spreng. Neue Entdeck. i. 243 = Ichnanthus leiocarpus.
leiogonum, *Delile, Fl. Aegypt. Illustr.* 51.—Aegypt.
leiogonum, Sieber, ex Link, Hort. Berol. i. 206 = numidianum.
leiophyllum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 490.—Mexic.
leiophyllum, *Nees, Agrost. Bras.* 249.—Bras.
lene, Steud. Syn. Pl. Gram. 54 = plicatum.
lentigerum, *J. & C. Presl, Rel. Haenk.* i. 299.—Hab.?
lepidotum, Steud. in Flora, xxix. (1846) 19 = Isachne australis.
Leprieurii, Steud. Syn. Pl. Gram. 65 = stoloniferum.
Leptachne, *Doell, in Mart. Fl. Bras.* ii. 11. 195.—Am. trop.
Leptachyrium, *Doell, l. c.* 150.—Am. trop.
leptanthum, *Steud. Syn. Pl. Gram.* 79.—Ins. Philipp.
leptocaulon, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. 11. (1835) 275.—Afr. trop.
Leptochloa, *Nees, ex Steud. Syn. Pl. Gram.* 62.—Ind. or.
leptomerum, *J. & C. Presl, Rel. Haenk.* i. 311.—Hab.?
leptostachyum, *J. & C. Presl, l. c.*—Mexic.
leucanthum, *A. Rich. Tent. Fl. Abyss.* ii. 372; *Steud. Syn. Pl. Gram.* 92.—Abyss.
leucitis, Trin. Gram. Panic. 85 = mollissimum.
leucoblepharis, Trin. Clav. Agrost. 234 = dichotomum.
leucophaeum, H. B. & K. Nov. Gen. et Sp. i. 97 = lanatum.
Liebmannianum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 491.—Mexic.
ligulare, Nees, ex Trin. Gram. Panic. 206 = scaberrimum.
limnaeum, Steud. Syn. Pl. Gram. 72 = molle.
limosum, Presl, ex Nees, Agrost. Bras. 257 = P. Crus-galli.
Lindenbergianum, *Nees, Fl. Afr. Austr.* 47.—Afr. austr.
Lindeni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 491.—Mexic.
Lindleyanum, *Nees, ex Steud. Syn. Pl. Gram.* 91.—Afr. trop.
lineare, Burm. f. Fl. Ind. 25. t. 10. f. 2 = Paspalum brevifolium.
lineare, Krock. Fl. Siles. i. 95 = glabrum.
lineare, *Linn. Sp. Pl.* ed. II. 85.—Am. bor.
lineatum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 61.—Afr. trop.
lineatum, *Trin. Sp. Gram. Ic.* ii. t. 233.—Afr. trop.
Linkianum, Kunth, Rév. Gram. i. 33 = sanguinale.
Linkii, Steud. Nom. ed. II. ii. 259 = Setaria longifolia.
linoides, *Steud. Syn. Pl. Gram.* 418.—Ins. Comor.
litigosum, Steud. l. c. 89 = Isachne miliacea.
liton, *Schult. Mant.* ii. 250 = dichotomum.
littorale, *C. Mohr, ex Vasey, in Coult. Bot. Gaz.* iii. (1878) 106.—Am. bor.
lobatum, *Brouss. ex Schult. Mant.* ii. 258 (Quid?).—Hab.?
loliaceum, Bertol. in Bologn. Opusc. Sc. iii. (1819) 408 = Ichnanthus candicans.

PANICUM:—
loliaceum, Lam. Illustr. i. 170 = Oplismenus loliaceus.
loliiforme, *Hochst. ex Steud. Syn. Pl. Gram.* 56.—Am. trop.
Lolium, Nees, Agrost. Bras. 102 = rottboellioides.
longebrachiatum, *Steud. Syn. Pl. Gram.* 60.—Guian.
longeracemosum, Steud. l. c. 45 = Oplismenus compositus.
longiflorum, J. F. Gmel. Syst. 158 = Paspalum brevifolium.
longiflorum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1835) 317 = Ichnanthus longiflorus.
longifolium, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 63.—Afr. trop.
longifolium, *Torr. Fl. U. St.* 149.—Am. bor.
longisetum, Poir. Encyc. Suppl. iv. 275 = Pennisetum borbonicum.
longisetum, Torr. in Am. Journ. Sc. Ser. I. iv. (1822) 58 = P. Crus-galli.
longispiculum, Doell, in Mart. Fl. Bras. ii. 11. 261 = Ichnanthus longiflorus.
loreum, *Trin. Gram. Panic.* 211.—Bras.
lutescens, Weig. Obs. 20 = Setaria glauca.
luteum, Gueldenst. Reisen, ii. 58, nomen = Setaria glauca.
luxurians, Willd. ex Spreng. Syst. i. 305 = Setaria luxurians.
luzonense, *J. & C. Presl, Rel. Haenk.* i. 308.—Ins. Philipp.
lycopodioides, *Bory, ex Nees, Agrost. Bras.* 236.—Ins. Borbon.
macilentum, *J. & C. Presl, Rel. Haenk.* i. 310.—Ins. Philipp.
macractinium, *Benth. Fl. Austral.* vii. 468.—Austral.
macranthum, *Trin. Gram. Panic.* 209.—Bras.
macrocarpon, *Torr. Fl. U. St.* i. 143.—Am. bor.
macrocarpum, Le Conte, ex Torr. Cat. Pl. N. York, 91 = scoparium.
macrochaetum, Link, Enum. Hort. Berol. i. 76 = Setaria italica.
macrochaetum, Voigt, Hort. Suburb. Calc. 702 = Setaria macrochaeta.
macrophyllum, Raddi, Agrost. Bras. 46 = streptostachys.
macrospermum, Willd. ex Spreng. Syst. i. 317 = olyroides.
macrostachyum, Doell, in Mart. Fl. Bras. ii. 11. 166 = Setaria composita.
macrostachyum, Nees, Agrost. Bras. 245 = Setaria macrostachya.
macrotrichum, *Steud. Syn. Pl. Gram.* 92.—Abyssin.
macrourum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1835) 227 = Setaria macrostachya.
macrum, Kunth, Rév. Gram. i. 40 = dichotomum.
maculatum, Aubl. Pl. Gui. i. 51 = latifolium.
maculatum, Reichb. ex Schlecht. in Linnaea, vi. (1831) 33 = divaricatum.
madagascariense, *Spreng. Syst.* i. 317.—Ins. Mascar.
magellanicum, *Lam. Encyc.* iv. 732 (err. typ. 742).—Am. austr.
majusculum, *F. Muell. ex Benth. Fl. Austral.* vii. 482.—Austral.
malaccense, Trin. Gram. Panic. 204 = Isachne pulchella.
malacotrichum, *Steud. Syn. Pl. Gram.* 418.—Ins. Nossibé.
mandshuricum, Maxim. Prim. Fl. Amur. 328 = Arundinella anomala.
mangaloricum, *Steud. Syn. Pl. Gram.* 78.—Ind. or.
margaritaceum, *Link, Hort. Berol.* i. 205.—Hab.?
marginatum, *R. Br. Prod.* 190.—Austral.
marginatum, Vahl, ex Wall. Cat. n. 8732 B = prostratum.
Mariae, *Steud. Syn. Pl. Gram.* 419.—Madag.
maritimum, Hort. ex Lam. Encyc. iv. 727 (err. typ. 737) = Setaria italica.
Martianum, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1835) 325 = Ichnanthus petiolatus.
martinicense, *Griseb. Fl. Brit. W. Ind.* 552.—Ind. occ.
mauritianum, Willd. ex Spreng. Syst. i. 305 = plicatum.
Maximilianum, *Schrad. ex Schult. Mant.* ii. 255.—Bras.
maximum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 373 = trypheron.
maximum, *Jacq. Ic. Pl. Rar.* i. t. 13; *Coll.* i. 76.—Am. trop.; Geront. trop. (inquil.).
maximum, *Nees, Fl. Afr. Austr.* 36.—Afr. austr.
mayarense, *Wright, in Sauv. Fl. Cub.* 197.—Cuba.

PANICUM:—
maynense, *Trin. in Linnaea*, x. (1836) 298.—Peruv.
megacarpum, Steud. in Lechl. Berb. Am. Austr. 56 = latifolium.
megalanthum, *Steud. Syn. Pl. Gram.* 93.—Ind. or.
megaphyllum, *Steud. l. c.* 53.—Afr. trop.
megapotamicum, *Spreng. Syst.* iv. *Cur. Post.* 34.—Bras.
megapotamicum, Spreng. Syst. iv. Cur. Post. 34 = debile.
megastachyum, *Nees, ex Trin. Gram. Panic.* 205.—Bras.
megastachyum, J. & C. Presl, Rel. Haenk. i. 305 = Preslii.
megiston, Schult. Mant. ii. 248 = maximum, *Jacq.*
melananthum, *F. Muell. in Trans. Vict. Inst.* i. (1855) 47.—Austral.
melananthum, *Trin. ex Steud. Nom.* ed. II. ii. 259 (Quid?).—Reg. Argent.
Mel-frugum, Hort. ex Schult. Mant. ii. 599, in obs. = Setaria italica.
melicarium, Michx. Fl. Bor. Am. i. 50 = debile.
melicoides, *Poir. Encyc. Suppl.* iv. 284.—Bras.
Melinis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 291.—Bras.
Meneri, Koen. ex Nees, Fl. Afr. Austr. 40 = miliare.
Meneritana, Spreng. Syst. i. 321 = Isachne australis.
Mertensii, *Roth, Nov. Pl. Sp.* 48.—Guian.
mesocomum, *Nees, Fl. Afr. Austr.* 34.—Afr. austr.
Metzii, Hochst. ex Steud. Syn. Pl. Gram. 95 = Isachne Kunthiana.
Meyerianum, *Nees, Fl. Afr. Austr.* 32.—Afr. austr.
Meyerianum, Trin. ex Steud. Nom. ed. II. ii. 259 (Quid?).—Hab.?
Michauxii, Roem. & Schult. Syst. ii. 427 = Eriochloa punctata.
micranthum, *H. B. & K. Nov. Gen. et Sp.* i. 105.—Venezuela.
microbachne, *J. & C. Presl, Rel. Haenk.* i. 298.—Am. austr.
microcarpon, *Muhl. Desc. Gram.* 111.—Am. bor.
microgrostum, Steud. Syn. Pl. Gram. 78 = sarmentosum.
microspermum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 492.—Mexic.
microstachyum, Lam. Illustr. i. 170 = indicum.
miliaceoides, *Roxb. Hort. Beng.* 8, nomen.—Sumatra.
miliaceum, Blanco, Fl. Filip. ed. I. 39 = Setaria italica.
miliaceum, *Linn. Sp. Pl.* 58.—Reg. calid.
miliaceum, Trin. ex Kunth, Enum. Pl. i. 113 = Isachne pulchella.
miliaceum, Walt. Fl. Carol. 72 = proliferum.
miliare, *Lam. Illustr.* i. 173.—As. trop. (cult.).
miliiforme, *J. & C. Presl, Rel. Haenk.* i. 300.—Ins. Philipp.
milioides, Nees, ex Trin. Gram. Panic. 225 = debile.
Milium, Pers. Syn. i. 83 = miliaceum.
Millegrana, Poir. Encyc. iv. 278 = rugulosum.
Minarum, Steud. Nom. ed. II. ii. 259 = Ichnanthus Minarum.
minutiflorum, *Doell, in Mart. Fl. Bras.* ii. II. 253.—Am. trop.
minutiflorum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 350.—Nubia.
minutiflorum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 362 = Paspalum ischnocaulon.
minutiflorum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Melinis minutiflora.
minutiflorum, *A. Rich. Fl. Cub. Fanerog.* ii. 305.—Cuba.
minutulum, *Desv. Opusc.* 87.—Carolina.
minutulum, Gaudich. in Freyc. Voy. Bot. 410 = Isachne miliacea.
minutum, R. Br. Prod. 191 = brevifolium, *Linn.*
mirabile, A. Br. Del. Sem. Hort. Carlsruh. (1846).—Cf. Walp. Ann. i. 924 = colonum.
Mitchelli, *Benth. Fl. Austral.* vii. 489.—Austral.
mite, *Steud. Syn. Pl. Gram.* 68.—Nubia.
molinioides, *Trin. Gram. Panic.* 238.—Bras.
molle, Michx. Fl. Bor. Am. i. 47 = Eriochloa punctata.
molle, *Sw. Prod. Veg. Ind. Occ.* 22.—Am. trop.; As. trop. (inquil.).
molliconum, *Kunth, Enum. Pl.* i. 108.—Peruv.
mollifolium, *Link, Hort. Berol.* ii. 215.—Afr. austr.
mollissimum, *Kunth, Rév. Gram.* i. 33.—Mexic.
Monachne, *Trin. in Spreng. Neue Entdeck.* ii. 86.—Ins. Borbon.
monachnoides, *Desv. Opusc.* 86.—Bras.

PANICUM:—
monobotrys, *Trin. ex Steud. Syn. Pl. Gram.* 55.—Bras.
monodactylum, *Nees, Fl. Afr. Austr.* 21.—Afr. austr.
monostachyum, *H. B. & K. Nov. Gen. et Sp.* i. 96.—Venezuela.
monostachyum, Salzm. ex Steud. Nom. ed. II. ii. 260 = monobotrys.
montanum, Gaudich. in Freyc. Voy. Bot. 411 = nubigenum.
montanum, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 313.—As. trop.
montevidense, Spreng. ex Doell, in Mart. Fl. Bras. ii. II. 260 = racemosum.
monticolum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 226.—Afr. trop.
movaiense, Steud. Syn. Pl. Gram. 78 = concinnum.
mucronatum, Heyne, ex Wall. Cat. n. 8717 = trypheron.
mucronatum, *Roth, Nov. Pl. Sp.* 45 (sp. dub.).—Ind. or.
Muehlenbergianum, *Schult. Mant.* ii. 230.—Am. bor.
Muehlenbergii, *Spreng. Syst.* i. 314.—Am. bor.
multibrachiatum, Hochst. ex Steud. Syn. Pl. Gram. 74 = *Perrottetii*.
multiculmum, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 133.—Ins. Galap.
multiflorum, Ell. Sketch, i. 122 = microcarpum.
multiflorum, Poir. Encyc. Suppl. iv. 282 = agrostoides.
multinerve, *Desv. ex Poir. Encyc. Suppl.* iv. 279.—Ind. occ.
multinode, Lam. Encyc. iv. 747 = trigonum.
multinode, J. & C. Presl, Rel. Haenk. i. 303 = nodosum.
multisetum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 377 = Oplismenus Burmannii.
munitum, *Trin. ex Steud. Nom.* ed. II. ii. 260.—Mexic.
Munroi, F. Muell. Fragm. v. 204 = Neurachne Munroi.
muricatum, Michx. Fl. Bor. Am. i. 47 = P. Crus-galli.
muricatum, Retz. Obs. iv. 18 = Isachne australis.
musaefolium, *Klotzsch, ex Schlecht. in Linnaea*, xxxi. (1861–62) 417.—N. Granat.
muscarium, *Trin. Sp. Gram. Ic.* ii. t. 235.—Afr. trop.
musciparum, Linn. Herb. ex Munro, in Journ. Linn. Soc. vi. (1862) 39 = miliare.
Musei, *Steud. Syn. Pl. Gram.* 58.—Guian.
muticum, *Forsk. Fl. Aegypt. Arab.* 20.—Aegypt.; Arab.
muticum, Link, Hort. Berol. i. 206 = numidianum.
Myosotis, Steud. Syn. Pl. Gram. 96 = Isachne Myosotis.
myosuroides, *R. Br. Prod.* 189.—As. trop.; Austral.
Myosurus, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = indicum.
myrianthum, *Buese, in Miq. Pl. Jungh.* 374.—Java.
Myuros, Kunth, Enum. Pl. i. 86 = seq.
Myurum, G. F. W. Mey. Prim. Fl. Esseq. 50 = amplexicaule.
Myurum, Wight, ex Steud. Syn. Pl. Gram. 55 = canaliculatum.
Myurus, Lam. Illustr. i. 172 = *P. Myosurus*.
nanum, Herb. Royle, ex Steud. Syn. Pl. Gram. 62 = villosum.
natalense, *Hochst. in Flora*, xxix. (1846) 113.—Afr. austr.
natans, Koen. ex Trin. Gram. Panic. 237 = paludosum, decompositum.
naviculare, Nees, Agrost. Bras. 123 = Ichnanthus Martianus.
Neesianum, Wight & Arn. ex Steud. Syn. Pl. Gram. 74 = trypheron.
Neesii, Kunth, Rév. Gram. i. 33 = sanguinale.
neglectum, *Roem. & Schult. Syst.* ii. 451.—Afr. trop.
nemorale, Schrad. ex Schult. Mant. ii. 255 = Ichnanthus leiocarpus.
nemorosum, Reichb. ex Steud. Nom. ed. II. ii. 260 = stoloniferum.
nemorosum, Sw. Prod. Veg. Ind. Occ. 22 = Ichnanthus nemorosus.
nemorosum, Willd. ex Mart. Fl. Bras. ii. II. 193 = uncinatum.
nepalense, Spreng. Syst. i. 321 = plicatum.
nephelophilum, *Gaudich. in Freyc. Voy. Bot.* 411.—Ins. Sandvic.
nervosum, *Lam. Encyc.* iv. 747.—Guian.
nervosum, Muhl. Desc. Gram. 116 = dichotomum.
nervosum, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 194 = sanguinale.

PANICUM:—
nervosum, Roxb. Hort. Beng. 8; Fl. Ind. i. 311 = plicatum.
neuranthum, *Griseb. Cat. Pl. Cub.* 232.—Cuba; Mexic.
neurodes, Schult. Mant. ii. 228 = plicatum.
neurophyllum, Spruce, ex Doell, in Mart. Fl. Bras. ii. II. 254 = P. Pterigodium.
nigrescens, Salzm. ex Doell, l. c. 213 = laxum.
nigricans, Willd. ex Spreng. Syst. i. 310 = fasciculatum.
nigrirostre, *Nees, Fl. Afr. Austr.* 55.—Afr. austr.
nigropedatum, *Munro, ex Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 29.—Afr. austr.
nilagiricum, Steud. Syn. Pl. Gram. 62 = villosum.
nitidum, Lam. Illustr. i. 172; Encyc. iv. 748 = dichotomum.
nodibarbatum, Hochst. ex Steud. Syn. Pl. Gram. 95 = Isachne australis.
nodiflorum, Lam. Encyc. iv. 744 = dichotomum.
nodiflorum, *Pursh, Fl. Am. Sept.* i. 68.—Am. bor.
nodosum, *Kunth, Enum. Pl.* i. 97.—As. trop.
nodosum, Willd. ex Steud. Nom. ed. II. ii. 260 = bulbosum.
norfolkianum, *Endl. Prod. Fl. Norf.* 18.—Ins. Norfolc.
nossibense, *Steud. Syn. Pl. Gram.* 419.—Ins. Nossibé.
notatum, Retz. Obs. iv. 18 = montanum?
Novae-Hollandiae, *Desv. Opusc.* 83.—Austral.
nubicum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 352.—Nubia.
nubicum, *Steud. Syn. Pl. Gram.* 88 (sp. diversa?).—Nubia.
nubigenum, *Kunth, Enum. Pl.* i. 98.—Ins. Sandvic.
nudiglume, Hochst. in Flora, xxvii. (1844) 253 = ramosum.
nudum, Walt. Fl. Carol. 73 = autumnale?
numaeënse, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 325.—N. Caled.
numidianum, *Lam. Illustr.* i. 172; *Encyc.* iv. 749.—Afr. bor.; Syria.
numidianum, Presl, Cyp. et Gram. Sic. 19 = colonum.
numidianum, Sieber, ex Schult. Mant. ii. 267 = paspaloides.
nutans, Desv. Opusc. 93 = Ichnanthus pallens.
Nuttallianum, Steud. Nom. ed. II. ii. 260 = Oplismenus undulatifolius.
oahuaense, Steud. l. c. 260 = Oplismenus compositus.
oaxacense, *Steud. Syn. Pl. Gram.* 73.—Mexic.
obliquum, Roth, Nov. Pl. Sp. 51 = radicans.
obliquum, Zoll. ex Steud. Syn. Pl. Gram. 96 = Isachne Kunthiana.
oblongatum, *Griseb. in Goett. Abh.* xix. (1874) 261.—Reg. Argent.
oblongiflorum, *Desv. Opusc.* 89.—Carolina.
obovatum, *Doell, in Mart. Fl. Bras.* ii. II. 256.—Am. trop.
obseptum, *Trin. ex Nees, Agrost. Bras.* 113.—Austral.; Bras.
obtectum, *J. & C. Presl, Rel. Haenk.* i. 301.—Mexic.
obtusiflorum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 367.—Abyss.
obtusiflorum, *A. Rich. Fl. Cub. Fanerog.* ii. 305.—Cuba.
obtusifolium, *Delile, Fl. Egypt.* 150. t. 5.—Alger.; Aegypt.
obtusiforme, D. Dietr. Syn. Pl. i. 256 = obtusifolium.
obtusum, *H. B. & K. Nov. Gen. et Sp.* i. 98.—Mexic.
ocreatum, Willd. ex Spreng. Syst. i. 305 = Setaria ocreata.
oliganthum, *Schlecht. in Linnaea*, xxvi. (1853–55) 137.—Venezuela.
oligosanthes, Schult. Mant. ii. 256 = dichotomum.
oligostachyum, *Steud. Syn. Pl. Gram.* 97.—Chili.
oligotrichum, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 333.—Nubia.
Olyrachne, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 112 = aturense.
olyraefolium, Raddi, Agrost. Bras. 43 = stoloniferum.
olyroides, *H. B. & K. Nov. Gen. et Sp.* i. 102.—Mexic.
Onurus, Willd. in Nees, Agrost. Bras. 251 = Setaria Onurus.
orientale, Willd. Enum. Hort. Berol. 1031 = Pennisetum orientale.
orinocense, Willd. ex Spreng. Syst. i. 316 = latifolium.
ornatum, *Ham. Prod. Pl. Ind. Occ.* 11.—Portorico.
ornithopus, Trin. in Spreng. Neue Entdeck. ii. 83 = sanguinale.

PANICUM:—
orthum, Voigt, Hort. Suburb. Calc. 701 = Kunthii.
oryzinum, J. F. Gmel. Syst. i. 157 = P. Crus-galli.
oryzoides, Ard. Aminad. Spec. Alt. 16. t. 5 = P. Crus-galli.
oryzoides, Nees, Agrost. Bras. 154 = grandiflorum.
oryzoides, *Sw. Prod. Veg. Ind. Occ.* 23.—Reg. trop.
ovale, *R. Br. in Salt, Abyss. App.* 62, nomen.—Abyss.
ovale, Ell. Sketch, i. 123 = microcarpon?
ovaliferum, Steud. Nom. ed. II. ii. 260, sphalm. = ovuliferum.
ovalifolium, *Poir. Encyc. Suppl.* iv. 279.—As. et Afr. trop.
ovatum, Steud. Nom. ed. II. ii. 260, sphalm. = ovale.
ovuliferum, *Trin. Gram. Panic.* 191.—Am. trop.
oxyanthum, *Steud. Syn. Pl. Gram.* 41.—Ins. S. Doming.
oxyphyllum, Hochst. ex Steud. l. c. 65 = pilipes.
ozogonum, *Steud. l. c.* 68.—Afr. trop.
pabulare, Aitch. & Hemsl. in Journ. Linn. Soc. xix. (1882) 190 = sanguinale.
pachystachys, *Franch. & Sav. Enum. Pl. Jap.* ii. 594. —Japon.
pallens, Cham. & Schlecht. in Linnaea, vi. (1831) 35 = Schlechtendalii.
pallens, Sw. Prod. Veg. Ind. Occ. 23 = Ichnanthus pallens.
pallide-fuscum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 58.—Afr. trop.
pallidifolium, *Desv. Opusc.* 85.—Am. calid.
palmaefolium, Koen. in Naturf. xxiii. (1788) 208 = plicatum.
palmifolium, *Poir. Encyc. Suppl.* iv. 282.—Ind. occ.
paludicolum, Nees, Agrost. Bras. 179 = Ichnanthus panicoides.
paludicolum, Steud. ex Griseb. Fl. Brit. W. Ind. 547 = fuscum.
paludosum, Hochst. ex Steud. Syn. Pl. Gram. 72 = repens.
paludosum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 307.—As. trop.
palustre, Trin. Gram. Panic. 181 = Ichnanthus panicoides.
pamplemoussense, Steud. Syn. Pl. Gram. 71 = maximum, *Jacq.*
pangerangense, *Steud. l. c.* 95.—Java.
paniculiferum, Steud. l. c. 54 = Setaria paniculifera.
papillosum, Fenzl, ex Steud. l. c. 100 = javanicum.
pappophorum, *Nees, Agrost. Bras.* 104.—Bras.
papposum, *R. Br. Prod.* 192.—Austral.
paractaenoides, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 219.—Ind. occ.
Paractaenum, Kunth, Rév. Gram. i. 41; Enum. Pl. i. 134 = gracile.
paradoxum, R. Br. Prod. 193 = Chamaeraphis paradoxa.
paradoxum, Roth, ex Steud. Syn. Pl. Gram. 75 = abludens.
paraguayense, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 189 = numidianum.
parciflorum, Steud. Syn. Pl. Gram. 45 = Oplismenus rariflorus.
Parlatorei, *Steud. l. c.* 40.—Ins. Canar.
parviflorum, *R. Br. Prod.* 192.—Austral.
parviflorum, Lam. ex Spreng. Syst. i. 321 = tenellum.
parvifolium, *Lam. Illustr.* i. 173; *Encyc.* iv. 752.—Reg. trop.
parvulum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 205 = Paspalum brevifolium.
parvum, *Buese, in Miq. Pl. Jungh.* 373.—Java.
paspaliforme, *J. & C. Presl, Rel. Haenk.* i. 296. —Peruv.
paspaloides, *Pers. Syn.* i. 81.—Reg. trop.
patagonicum, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 377.—Reg. Argent.
patens, Boj. Hort. Maurit 365 = auritum.
patens, Burm. f. Fl. Ind. t. 10. f. 2 = trigonum.
patens, Hochst. ex Steud. Syn. Pl. Gram. 90 = Hochstetteri.
patens, *Linn. Sp. Pl.* 58.—Ind. or.
patentissimum, *Desv. ex Poir. Encyc. Suppl.* iv. 283.—Am. trop.
patentissimum, Roem. & Schult. Syst. ii. 448 = debile.
pauciflorum, Bory, ex Nees, Agrost. Bras. 236 = parvifolium.
pauciflorum, Bosc, ex Spreng. Syst. i. 313 = dichotomum.
pauciflorum, *R. Br. Prod.* 191.—Austral.

PANICUM:—
pauciflorum, Ell. Sketch, i. 120 = scoparium.
pauciflorum, Willd. ex Spreng. Syst. i. 312 = depauperatum, pygmaeum.
paucisetum, Steud. Syn. Pl. Gram. 52 = plicatum.
pauperulum, *Steud. l. c.* 58.—Afr. trop.
pedunculare, Willd. ex Steud. Nom, ed. II. ii. 260; et Syn. Pl. Gram. 77 = cayennense.
pedunculatum, Torr. Fl. U. St. 141 = clandestinum, *Linn.*
pellitum, Trin. Gram. Panic. 198 = nubigenum.
penicillatum, Nees, Agrost. Bras. 145 = Ichnanthus bambusiflorus.
penicillatum, Trin. Gram. Panic. 196 = discolor.
penicillatum, Willd. ex Nees, Agrost. Bras. 242 = Setaria glauca.
peninsularum, *Steud. Syn. Pl. Gram.* 44.—Ind. or.
pennatum, Hochst. in Flora, xxxviii. (1855) 197 (Paspali sp.).—Afr. bor.; As. occ.
Pennisetum, Roth, Nov. Pl. Sp. 55 = pilosum.
pensylvanicum, Spreng. Nachr. I. Bot. Gart. Halle, 29 = anceps.
perdensum, *Steud. Syn. Pl. Gram.* 65.—Reg. Argent.
peregrinum, *Steud. l. c.* 63.—Hab.?
perfoliatum, Nees, Agrost. Bras. 142 = streptostachys.
perforatum, Nees, l. c. 126 = trachystachyum.
perpusillum, *Nees, ex Steud. Syn. Pl. Gram.* 96.—Zeylan.
Perrottetii, Kunth, Rév. Gram. ii. 395. t. 111 (Paspali sp.).—Ind. or.; Afr. trop.
persicum, *Fensch. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 358, nomen.—Reg. Transcauc.
petiolatum, Nees, Agrost. Bras. 140 = Ichnanthus petiolatus.
Petiverii, Kotschy, ex Griseb. Fl. Brit. W. Ind. 545 = colonum.
Petiverii, Thw. Enum. Pl. Zeyl. 359 = semiverticillatum.
Petiverii, Trin. Gram. Panic. 144 = ramosum, remotum.
petrosum, *Trin. Sp. Gram. Ic.* iii. t. 280.—Bras.
phaeocarpum, *Nees, Fl. Afr. Austr.* 22.—Afr. austr.
phaeothrix, *Trin. Sp. Gram. Ic.* iii. t. 91.—Bras.
phaetrix, Steud. Nom. ed. II. ii. 261, sphalm. = praec.
phalaroides, Roem. & Schult. Syst. ii. 452 = indicum.
philadelphicum, Bernh. ex Trin. Gram. Panic. 216 = capillare.
phleiforme, *J. & C. Presl, Rel. Haenk.* i. 302.—Mexic.
phleoides, R. Br. Prod. 189 = indicum.
Phragmites, Nees, Agrost. Bras. 190 = discolor.
phyllanthum, Steud. Syn. Pl. Gram. 47 = spectabile.
phyllomacrum, *Steud. l. c.* 53.—Afr. trop.
pictigluma, Steud. l. c. 73 = numidianum.
pictum, Koen. in Naturf. xxiii. (1788) 204 = P. Crus-galli.
piliferum, *Koen. l. c.* 235; *ex J. F. Gmel. Syst.* 157.—Hab.?
piligerum, *F. Muell. ex Benth. Fl. Austral.* vii. 477.—Austral.
pilipes, Nees, ex Thw. Enum. Pl. Zeyl. 359 = hermaphroditum, trigonum.
pilipes, *Nees & Arn. ex Buese, in Miq. Pl. Jungh.* 376 (an seq.?).—Java.
pilisparsum, G. F. W. Mey. Prim. Fl. Esseq. 57 = pilosum.
pilosissimum, Roth, Nov. Pl. Sp. 48 = cayennense.
pilosum, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. (1842) 239 = Paspalum fastigiatum.
pilosum, *Sw. Prod. Veg. Ind. Occ.* 22.—Am. austr.
pilulíferum, *Nees, ex Steud. Syn. Pl. Gram.* 94.—Ind. or.
piptopilum, *Steud. l. c.* 417.—Ins. Nossibé.
Planopteris, Trin. ex Doell, in Mart. Fl. Bras. ii. II. 222 = Maximiliani.
Planotis, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 322 = Ichnanthus Planotis.
plantagineum, *Link, Hort. Berol.* i. 206.—Am. trop.
plantagineum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 64.—Afr. trop.
platycarphum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 198.—Ins. Bonin.
platyphyllum, *Munro, ex Wright, in Sauv. Fl. Cub.* 197.—Cuba.
pleiophyllum, *Link, Hort. Berol.* i. 208.—Hab.?
plicatile, *Hochst. in Flora*, xxxviii. (1855) 198.—Afr. trop.
plicatum, *Lam. Illustr.* i. 171; *Encyc.* iv. 736; *Schrank, Pl. Rar. Hort. Monac.* t. 19.—Reg. trop.

PANICUM:—
plicatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 311 = asperatum.
plumosum, Presl, Fl. Sic. p. xliii = Teneriffae.
poaeforme, Boj. Hort. Maurit. 365 = Stenotaphrum complanatum.
poaeforme, Poir. Encyc. Suppl. iv. 284 = melicoides.
poaeforme, *Willd. ex Spreng. Syst.* i. 318.—Ins. Mascar.
poaemorphum, *J. & C. Presl, Rel. Haenk.* i. 310.—Peruv.
Poiretianum, *Schult. Mant.* ii. 229.—Bras.
polyanthes, Schult. l. c. 207 = dichotomum.
polycomum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 306.—Guian.
polygamum, Forsk. Fl. Aegypt. Arab. 27 (Penniseti sp.).—Arab.
polygamum, Sw. Prod. Veg. Ind. Occ. 24 = maximum, *Jacq.*
polygonatum, *Kunth, Rév. Gram.* ii. t. 209 = laxum.
polygonatum, Schrad. in Schult. Mant. ii. 256 = pilosum.
polygonatum, Sieber, ex Griseb. Fl. Brit. W. Ind. 548 = gibbum.
polygonoides, Lam. Encyc. iv. 742 = Isachne miliacea.
polygonoides, *C. Muell. in Bot. Zeit.* xix. (1861) 323. —Am. bor.
polyneuron, Steud. Syn. Pl. Gram. 91 = dichotomum.
polyphyllum, *R. Br. Prod.* 190.—Austral.
polyrrhizum, J. & C. Presl, Rel. Haenk. i. 296 = Paspalum distichum.
polystachion, Linn. Syst. ed. X. 870 = Andropogon caricosum.
polystachion, Ucria, Hort. Reg. Panorm. 54 = repens.
polystachyum, Burm. ex Kunth, Enum. Pl. i. 478 = Pogonatherum saccharoides.
polystachyum, *J. & C. Presl, Rel. Haenk.* i. 312.—Peruv.
polystachyum, Rich. ex Kunth, Enum. Pl. i. 93 = pyramidale.
polystachyum, Schult. Mant. ii. 146, in obs. = holcoides.
polystachyum, Steud. Nom. ed. I. 589; ed. II. ii. 261 = Oplismenus polystachyus.
polythyrsum, Nees, ex Steud. Syn. Pl. Gram. 93 = Ichnanthus candicans.
porphyrium, Trin. ex Nees, Agrost. Bras. 198 = capillare.
porphyrrhizos, Steud. Syn. Pl. Gram. 72 = trypheron.
porranthum, *Steud. l. c.* 42.—Afr. trop.
portoricense, *Ham. Prod. Pl. Ind. Occ.* 11.—Portorico.
potamicum, Steud. Syn. Pl. Gram. 71 = seq.
potamium, Trin. Gram. Panic. 239 = gibbum.
praealtum, *Afzel. ex Sw. Adnot. Bot.* 5.—Afr. trop.
praegnans, *Steud. Syn. Pl. Gram.* 74.—Bras.
praelongum, *Steud. l. c.* 73.—Afr. trop.
pratense, Steud. Nom. ed. II. ii. 261 = Oplismenus compositus.
praticola, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 203 = maximum, *Jacq.*
Preslii, *Kunth, Enum. Pl.* i. 121.—Peruv.
Prionitis, *Nees, Agrost. Bras.* 162.—Bras.
probandum, Steud. Syn. Pl. Gram. 76 = rugulosum.
proboscideum, Trin. Gram. Panic. 184 = olyroides.
procumbens, Nees, Agrost. Bras. 109 = prostratum.
procurrens, *Nees, ex Trin. Gram. Panic.* 183.—Bras.
proliferum, *Lam. Encyc.* iv. 747.—Am. bor.
proliferum, F. Muell. Fragm. viii. 191 = decompositum.
prolisetum, *Steud. Syn. Pl. Gram.* 52.—Afr. trop.
prolutum, *F. Muell. in Trans. Vict. Inst.* i. (1855) 46. —Austral.
propinquum, R. Br. Prod. 193 = Paspalum brevifolium.
prorepens, *Steud. Syn. Pl. Gram.* 46.—Mexic,
prostratum, *Lam. Illustr.* i. 171.—Amphig. trop.
proximum, *Steud. Syn. Pl. Gram.* 64.—Guian.
pruinosum, Bernh. ex Trin. Gram. Panic. 191 = antidotale.
pruriens, Trin. l. c. 77 = sanguinale.
Pseudagrostis, Trin. l. c. 197 = nubigenum.
Pseudoagrostis, *Steud. Syn. Pl. Gram.* 71.—Afr. trop.
Pseudocolonum, Roth, Nov. Pl. Sp. 47 = colonum.
Pseudo-duroa, Nees, Fl. Afr. Austr. 21 = Paspalum brevifolium.
Pseudoholcus, *Steud. Nom.* ed. II. ii. 262; Syn. Pl. Gram. 50.—Am. austr.
Pseudo-neurachne, F. Muell. Fragm. viii. 199 = Spinifex paradoxus.

PANICUM:—
Pseudopaspalus, *Nees, Del. Sem. Hort. Vrat.* (1850). —Cf. Linnaea, xxiv. (1851) 236.—Hab.?
Pseudoryzoides, Steud. Syn. Pl. Gram. 75 = oryzoides.
Pseudosetaria, *Steud. l. c.* 39.—Ind. or.
Pseudoundulatifolium, *Roem. & Schult. Syst.* ii. 437. —Guian.
psilanthum, Steud. Syn. Pl. Gram. 66 = laxum.
psilocaulon, Steud. l. c. 50 = Setaria viridis.
psilopodium, *Trin. Gram. Panic.* 217.—Ind. or.
Pterigodium, *Trin. l. c.* 227.—Bras.
puberulum, Kunth, Rév. Gram. i. 32 = Paspalum ischnocaulon.
puberulum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 277 = rugulosum.
pubescens, R. Br. Prod. 190 = foliosum.
pubescens, Hort. ex Kunth, Enum. Pl. i. 59 = Paspalum elegans.
pubescens, Lam. Encyc. iv. 748 = dichotomum.
pubescens, Michx. Fl. Bor. Am. i. 49 = dichotomum.
pubescens, Presl, ex Steud. Nom. ed. II. ii. 262 = glutinosum.
pubescens, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1805) 301 = parvifolium.
pubiflorum, Trin. Gram. Panic. 150 = gracile.
pubigerum, Roem. & Schult. Syst. ii. 460 = foliosum.
pubinode, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 363 = P. Isachne.
pulchellum, *Raddi, Agrost. Bras.* 42.—Am. trop.
pulchellum, Spreng. Syst. i. 322 = Isachne pulchella.
pulchrum, Willd. ex Spreng. Syst. i. 272; et ex Trin. Gram. Panic. 88 = Paspalum pulchrum.
pulipes, C. Muell. in Walp. Ann. vi. 953, sphalm. = pilipes.
pumilum, Bosc, ex Nees, Agrost. Bras. 228 = dichotomum.
pumilum, Link, Enum. Hort. Berol. i. 76 = Setaria italica.
pumilum, Poir. Encyc. Suppl. iv. 273 = Setaria glauca.
pumilum, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen (Quid?).—Am. bor.
punctatum, *Burm. f. Fl. Ind.* 26.—Ind. or.
punctulatum, Arn. ex Steud. Syn. Pl. Gram. 62 = molle.
pungens, *Poir. Encyc. Suppl.* iv. 273.—Am. bor.
pungens, *Trin. Gram. Panic.* 210.—Bras.
purpurascens, Opiz, in Flora, v. (1822) 266; Raddi, ex Nees, Agrost. Bras. 240 = Setaria viridis.
purpureum, Ruiz & Pav. Fl. Per. i. 48 = Setaria purpurea.
pusillum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 227.—Afr. trop.
pycnanthum, Steud. Syn. Pl. Gram. 70 = rivulare.
pygmaeum, Boj. Hort. Maurit. 365 = foliosum.
pygmaeum, *R. Br. Prod.* 191.—Austral.
pygmaeum, Spreng. Syst. i. 312 = parvifolium.
pyramidale, *Lam. Illustr.* i. 171; *Encyc.* iv. 735 (err. typ. 745).—Afr. trop.
quadrifarium, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et in A. Rich. Tent. Fl. Abyss.* ii. 367. —Abyss.
quale-Linnaei, Krock. Fl. Siles. i. 88 = Setaria viridis.
quinquangulare, Buch.-Ham. ex Wall. Cat. n. 8725 B = ramosum.
racemiferum, *Steud. Syn. Pl. Gram.* 64.—Guian.
racemiferum, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 171.—Bras.
racemosum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = reptans.
racemosum, *Spreng. Syst.* i. 313.—Bras.
Raddianum, Steud. Syn. Pl. Gram. 45 = Oplismenus compositus.
radiatum, R. Br. Prod. 192 = divaricatissimum.
radicans, *Retz. Obs.* iv. 18.—Ind. or.
radicosum, J. & C. Presl, Rel. Haenk. i. 297 = sanguinale.
Rafinesquianum, *Schult. Mant.* ii. 257.—Am. bor.
ramosissimum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 312 = parvifolium.
ramosum, *Linn. Mant.* i. 29.—Ind. or.
ramulare, Trin. Gram. Panic. 244 = parviflorum.
ramuliflorum, Hochst. ex Steud. Syn. Pl. Gram. 65 = laxum.
ramulosum, Michx. Fl. Bor. Am. i. 50 = dichotomum.
rariflorum, Lam. Encyc. iv. 746 = Orthoclada rariflora.

PANICUM:—
rariflorum, *Rupr. ex Galeotti, in Bull. Acad. Brux.* ix. II. (1842) 240.—Mexic.
rariflorum, Steud. Syn. Pl. Gram. 51 = Setaria rariflora.
raripilum, *Kunth, Rév. Gram.* ii. t. 114.—Afr. trop.
rarisetum, *Steud. Syn. Pl. Gram.* 52.—Ins. Mascar.
rarum, *R. Br. Prod.* 189.—Austral.
recalvum, *Kunth, Rév. Gram.* i. 39.—Bras.
reclinatum, Vill. Hist. Pl. Dauph. ii. 64 = Setaria viridis.
rectum, Roem. & Schult. Syst. ii. 457 = depauperatum.
redevivum, *Trin. ex Steud. Nom.* ed. II. ii. 262.—Mexic.
reflexopilum, *Steud. Syn. Pl. Gram.* 84.—Mexic.
regulare, *Nees, Fl. Afr. Austr.* 41.—Afr. trop.
relaxatum, Nees, ex Steud. Nom. ed. II. ii. 262 = remotum.
reimarioides, *Anderss. in Peters, Reise Mossamb. Bot.* 547.—Afr. trop.
remotum, *Retz. Obs.* iv. 17.—Ind. or.
Renggeri, Steud. Syn. Pl. Gram. 89 = aturense.
repandum, *Nees, Agrost. Bras.* 98.—Bras.
repens, Burm. f. Fl. Ind. 2. f. 1 = prostratum.
repens, *Linn. Sp. Pl.* ed. II. 87.—Reg. calid.
repens, Nees, Agrost. Bras. 171 = parvifolium.
repens, Thw. Enum. Pl. Zeyl. 360 = paludosum.
reptans, Kunth, Rév. Gram. i. 219. t. 21 = racemosum.
reptans, Linn. Syst. ed. X. 870 = repens?
rescissum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 218.—Afr. trop.
respiciens, *Hochst. ex Steud. Syn. Pl. Gram.* 52.—Abyss.; Java.
restitutum, *Steud. l. c.* 53.—Ins. S. Doming.
reticulatum, *Griseb. in Goett. Abh.* vii. (1857) 264.—Ind occ.?
reticulatum, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 271.—Zeylan.
reticulatum, *Torr. in Marcy, Bot. Exp. Red Riv. Louis.* (1852), 299.—Am. bor.
retroflexum, *Steud. Syn. Pl. Gram.* 74.—Guian.
retrofractum, *Delile, ex Desv. Opusc.* 96.—Am. bor.
reversipilum, *Steud. Syn. Pl. Gram.* 76.—Chili.
reversum. *F. Muell. Fragm.* viii. 152.—Austral.
rhabdinum, Steud. Syn. Pl. Gram. 96 = Isachne virgata.
rhachitrichum, *Hochst. in Flora*, xxvii. (1844) 254.—As. et Afr. trop.
rhigiophyllum, Steud. Syn. Pl. Gram. 76 = cayennense.
rhignon, Steud. l. c. 95 = Isachne rigida.
rhizophorum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 495.—Mexic.
Riedelii, Trin. Sp. Gram. Ic. iii. t. 323 = Ichnanthus Riedelii.
rigens, Salzm. ex Steud. Syn. Pl. Gram. 76 = cayennense.
rigens, Sw. Prod. Veg. Ind. Occ. 23 = Isachne rigens.
rigidifolium, *Kunth, Rév. Gram.* i. 37.—Ins. S. Doming.
rigidifolium, Trin. Sp. Gram. Ic. ii. t. 214 = dioecum.
rigidulum, Bosc, ex Nees, Agrost. Bras. 163 = agrostoides, anceps.
rigidum, *Balf. in Proc. Roy. Soc. Edinb.* xii. (1884) 97.—Ins. Socotra.
rigiophyllum, Doell, in Mart. Fl. Bras. ii. II. 220 = cayennense.
rivulare, *Trin. Gram. Panic.* 213.—Bras.
Rohrii, Nees, ex Steud. Syn. Pl. Gram. 76 = streptostachys.
roseum, *Steud. l. c.* 92.—Afr. austr.
roseum, Willd. ex Spreng. Syst. i. 251 = Milium roseum.
rostellatum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 288 = rugulosum.
rostratum, Muhl. ex Willd. Enum. Hort. Berol. 1032 = anceps.
Rothii, *Spreng. Syst.* i. 310.—Ind. or.
rottboelloides, *H. B. & K. Nov. Gen. et Sp.* i. 96.—Venezuela.
Rottleri, Kunth, Rév. Gram. i. 33 = sanguinale.
Rottleri, Nees, Fl. Afr. Austr. 53 = Setaria glauca.
Rottleri, *Sw. Adnot. Bot.* 7.—Ind. or.
Roxburghianum, Schult. Mant. ii. 245 = uliginosum.
Roxburghii, Spreng. Syst. i. 320 = trypheron.
Royleanum, Nees, ex Steud. Syn. Pl. Gram. 47 = Paspalum ischnocaulon.
rubigerum, Steud. Nom. ed. II. i. 567, in syn., sphalm. = nubigenum.

PANICUM:—
rubiginosum, *Steud. Syn. Pl. Gram.* 50.—Ins. Philipp.
rude, Lam. ex Steud. Nom. ed. I. 589, 772 = Setaria verticillata.
rude, *Nees, Agrost. Bras.* 158.—Bras.
Rudgei, Roem. & Schult. Syst. ii. 444 = cayennense.
rudimentosum, *Steud. Syn. Pl. Gram.* 51.—Afr. trop.
rufum, Kunth, Rév. Gram. i. 35 = Anthaenantia rufa.
Rugelii, *Griseb. Cat. Pl. Cub.* 233.—Cuba.
rugosum, Bosc, ex Nees, Agrost. Bras. 226 = verrucosum.
rugosum, Hemsl. Biol. Centr. Am. Bot. iii. 506 = Setaria semirugosa.
rugulosum, *Trin. Gram. Panic.* 195.—Bras.
rupestre, *Trin. l. c.* 109.—Bras.
Ruprechtii, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 495 = rariflorum.
ruscifolium, *H. B. & K. Nov. Gen. et Sp.* i. 101.—Mexic.
sabulicolum, Nees, Agrost. Bras. 258 = P. Crus-galli.
sabulorum, *Lam. Encyc.* iv. 744.—Reg. Argent.
sacchariferum, Salisb. Prod. 18 = Sorghum vulgare.
sacchariflorum, *Steud. Syn. Pl. Gram.* 97.—Bras.
saccharoides, Kunth, Rév. Gram. i. 237. t. 30 = Paspalum saccharoides.
saccharoides, *A. Rich. Fl. Cub. Fanerog.* ii. 306.—Cuba.
saccharoides, Trin. Gram. Panic. 245 = Teneriffae.
saccharoides, Willd. ex Hemsl. Biol. Centr. Am. Bot. iii. 490 = lanatum.
sagittaefolium, *Hochst. ex Steud. Syn. Pl. Gram.* 54. —Abyss.
Salzmanni, Trin. ex Steud. l. c. 95 = Isachne ventricosa.
Sanctae-Marthae, *Steud. l. c.* 45.—N. Granat.
sanguinale, Burm. f. Fl. Ind. t. 10. f. 1 = javanicum.
sanguinale, *Linn. Sp. Pl.* 57.—Cosmop.
sanguinale, Pollich, Hist. Pl. Palat. i. 57 = glabrum.
sanguineum, Gueldenst. Reisen, ii. 278 = sanguinale.
sarmentosum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Thuarea sarmentosa.
sarmentosum, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 308.—As. trop.
saxatile, *Salzm. ex Steud. Nom.* ed. II. ii. 263.—Bras.
saxatile, Steud. Syn. Pl. Gram. 97 = Isachne albens.
scaberrimum, *Lag. Gen. et Sp. Nov.* 2.—Mexic.
scabridum, *Doell, in Mart. Fl. Bras.* ii. II. 201.—Am. trop.
scabrifolium, *Nees, Agrost. Bras.* 246.—Bras.
scabriusculum, *Ell. Sketch*, i. 121.—Am. bor.
scabrum, Lam. Illustr. i. 171 = P. Crus-galli.
scandens, Trin. Gram. Panic. 166 = Setaria scandens.
scariosum, *Trin. ex Steud. Nom.* ed. II. ii. 263.—Mexic.
Scheelii, Steud. Syn. Pl. Gram. 51 = Setaria polystachya.
Schiedeanum, *Trin. ex Steud. Nom.* ed. II. ii. 263.—Mexic.
Schiedei, Spreng. ex Steud. l. c. = Zeugites mexicanus.
Schimperianum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 371.—Abyss.
Schlechtendalii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 496.—Mexic.
Schraderi, *Kunth, Rév. Gram.* i. 32.—Hab.?
Schultesii, *Steud. Nom.* ed. II. ii. 263.—N. Granat.
sciaphilum, *Rupr. ex Galeotti, in Bull. Acad. Brux.* ix. II. (1842) 240.—Mexic.
scindens, *Nees, ex Steud. Syn. Pl. Gram.* 47.—Am. bor.
sciurotis, *Trin. Gram. Panic.* 228.—Bras.
Sclerochloa, *Trin. Sp. Gram. Ic.* iii. t. 283.—Bras.
scoparium, *Lam. Encyc.* iv. 744.—Am. bor.
scoparium, Michx. Fl. Bor. Am. 49 = latifolium.
scoparium, Rudge, Pl. Guian. Rar. 21. t. 29 = cayennense.
scopuliferum, Trin. Sp. Gram. Ic. ii. t. 165 = serratum.
secundum, Steud. Nom. ed. II. ii. 263 = Oplismenus secundus.
secundum, Trin. Sp. Gram. Ic. iii. t. 324 = latissimum.
Sellowii, Nees, Agrost. Bras. 153 = rugulosum.
semialatum, *R. Br. Prod.* 192.—As. trop.; Afr. austr.; Austral.
semirugosum, Nees, Agrost. Bras. 243 = Setaria semirugosa.
semitonsum, *F. Muell. ex Benth. Fl. Austral.* vii. 483.—Austral.
semiundulatum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19. nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 49.—Abyss.

PANICUM:—
semiverticillatum, *Rottl. in Ainslie, Mat. Med. Hindost.* ed. I. 219.—Ind. or.
senegalense, *Desv. Opusc.* 86.—Afr. trop.
senescens, *Trin. ex Steud. Nom.* ed. II. ii. 263.—Hab. ?
sericatum, Steud. Syn. Pl. Gram. 58 = lanatum.
sericeum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. 88.—Ind. or.
serotinum, Hort. ex Schult. Mant. iii. 598, in obs. = Setaria italica.
serotinum, Trin. Gram. Panic. 166 = Paspalum serotinum.
serpens, Kunth, Rév. Gram. i. 38 = parvifolium.
serraefolium, *Hochst. in Flora*, xxxviii. (1855) 196.—Abyss.
serrarium, *Fig. & De Not. in Mem. Acc. Torin.* Ser. II. xiv. (1854) 347.—Nubia.
serratulatum, Buch.-Ham. ex Wall. Cat. n. 8699 = amplexicaule.
serratum, *R. Br. Prod.* 189; *Spreng. Syst.* i. 309.—Afr. austr.
serrulatum, Hochst. ex Steud. Syn. Pl. Gram. 78 = mangaloricum.
serrulatum, Roxb. Hort. Beng. 8; Fl. Ind. i. 307 = amplexicaule.
serrulatum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 62.—Afr. trop.
setaceum. Hort. ex. Trin. Gram. Panic. 166 = Setaria italica.
setaceum, Muhl. Desc. Gram. 99 = dichotomum ?
setarioides, *Steud. Syn. Pl. Gram.* 74.—Guian.
setarium, Lam. Illustr. i. 170 = Oplismenus undulatifolius.
setifolium, Nees, Agrost. Bras. 233 = subulatum.
setifolium, Trin. ex Steud. Nom. ed. II. ii. 263 = rupestre.
setigerum, *Beauv. Fl. Owar.* i. t. 49.—Afr. trop.
setigerum, Retz. Obs. iv. 16 = prostratum ?
setosum, Hort. ex Trin. Gram. Panic. 166 = Setaria italica.
setosum, Sw. Prod. Veg. Ind. Occ. 22 = Setaria setosa.
setosum, Trin. ex Doell, in Mart. Fl. Bras. ii. II. 169 = Setaria impressa.
sibiricum, Hort. ex Roem. & Schult. Syst. ii. 493, in syn. = Setaria italica.
siccaneum. *Trin. in Linnaea*, x. (1836) 298.—Bras.
Sieberi, Link, Hort. Berol. i. 207 = prostratum.
simplex, Rottl. ex Trin. Gram. Panic. 216 = miliare.
simplex, Willd. ex Spreng. Syst. i. 318 = *sumatrense*.
simplicissimum, *Koen. in Naturf.* xxiii. (1788) 209 (Quid ?).—Ind. or.
simpliciusculum, Wight & Arn. ex Steud. Syn. Pl. Gram. 96 = Coelachne pulchella.
singulare, Steud. Syn. Pl. Gram. 60 = marginatum.
sordidum, *Thw. Enum. Pl. Zeyl.* 443.—Zeylan.
Sloanei, Griseb. Fl. Brit. W. Ind. 551 = latifolium.
Sorghi, *Delile, Fl. Aeg. Illustr.* 51, nomen.—Aegypt.
sorghoideum, *Ham. Prod. Pl. Ind. Occ.* 10.—Portorico; Mexic.
Sorghum, Steud. Syn. Pl. Gram. 58 = P. Sorghi.
spanianthum, *Steud. l. c.* 69.—Hab. ?
sparsicomum, *Nees, ex Steud. l. c.* 83.—Zeylan.
sparsiflorum, *Doell, in Mart. Fl. Bras.* ii. II. 243.—Am. trop.
sparsum, Nees, ex Steud. Syn. Pl. Gram. 63 = peregrinum ?
sparsum, Rottl. ex Steud. l. c. 70 = antidotale.
sparsum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 64.—Afr. trop.
spathellosum, *Doell, in Mart. Fl. Bras.* ii. II. 241.—Bras.
speciosum, Nees, ex Trin. Gram. Panic. 169 = Poiretianum.
speciosum, *Walt. Fl. Carol.* 78.—Am. bor.
spectabile, *Nees, ex Trin. Gram. Panic.* 138.—Am. et Afr. trop.
sphacelatum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 58.—Afr. trop.
sphacelatum, *Steud. Syn. Pl. Gram.* 92 (sp. diversa).—Afr. trop.
sphaerocarpon, Ell. Sketch, i. 125 = microcarpum.
sphaerocarpum, *Salzm. ex Steud. Syn. Pl. Gram.* 51.—Bras.
spicatum, Roxb. Hort. Beng. 7; Fl. Ind. i. 283 = Pennisetum typhoideum.
spiciforme, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 359 = interruptum ?

PANICUM:—
spinescens, R. Br. Prod. 193 = Chamaeraphis spinescens.
spithameum, Willd. ex Steud. Nom. ed. II. ii. 263 = fasciculatum.
spithamineum, Willd. ex Steud. Syn. Pl. Gram. 80 = carthagenense.
Sprengelianum, Schult. Mant. ii. 234 = pilosum.
Sprengelii, Kunth, Rév. Gram. i. 39 = Muehlenbergii.
spretum, Schult. Mant. ii. 248 = dichotomum.
squamatum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 496.—Mexic.
squarrosum, Lam. Encyc. iv. 743 = Chamaeraphis spinescens.
squarrosum, Retz. Obs. iv. 15 = Trachys mucronata.
stagninum, Retz. l. c. 17; Host, Gram. Austr. iii. t. 2. 51 = P. Crus-galli.
stenanthum, *Steud. Syn. Pl. Gram.* 96.—Madag.
stenocladium, Trin. Gram. Panic. 200 = subulatum.
stenodes, *Griseb. Fl. Brit. W. Ind.* 547.—Jamaic.
stenostachyum, *Benth. Fl. Austral.* vii. 470.—Austral.
stenostachyum, Thw. Enum. Pl. Zeyl. 436 = canaliculatum.
stenotaphrodes, *Nees, ex Steud. Syn. Pl. Gram.* 41.—'Ins. Choin.'
stigmosum, *Trin. Gram. Panic.* 194.—Bras.
stipatum, *J. & C. Presl, Rel. Haenk.* i. 297.—Mexic.
stoloniferum, *Poir. Encyc. Suppl.* iv. 274.—Am. austr.
streptostachys, *Spreng. Syst.* i. 316.—Am. austr.
striatulum, Schult. Mant. ii. 252 = Isachne australis.
striatum, R. Br. Prod. 192 = parviflorum.
striatum, Chapm. Fl. S. U. St. Suppl. 666 = fasciculatum.
striatum, *Lam. Illustr.* i. 172.—Am. bor.
striatum, Muhl. ex Hemsl. Biol. Centr. Am. Bot. iii. 492, in syn. = neuranthum.
striatum, Spreng. Syst. i. 318 = miliaceum.
strictissimum, *Afzel. ex Sw. Adnot. Bot.* 4.—Afr. trop.
strictum, Bosc, ex Trin. Gram. Panic. 241 = dichotomum.
strictum, R. Br. Prod. 190 = marginatum.
strictum, Pursh, Fl. Am. Sept. i. 69 = depauperatum.
strictum, Roxb. Hort. Beng. 7; Fl. Ind. i. 303 = bengalense.
strigosum, Muhl. in Ell. Sketch, i. 126 = capillare.
striolatum, Schult. Mant. iii. 589 = striatum.
strumosum, *J. & C. Presl, Rel. Haenk.* i. 303.—Calif.
subalbidum, Kunth, Rév. Gram. ii. 397. t. 112 = antidotale.
subangustum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 59.—Afr. trop.
subaristulatum, Steud. Syn. Pl. Gram. 79 = Ichnanthus candicans.
subcordatum, Roth, Nov. Pl. Sp. 46 = prostratum ?
subeglume, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 292.—Ind. or.
subfalcatum, *Doell, in Mart. Fl. Bras.* ii. II. 181.—Am. trop.
subpellucidum, *Steud. Syn. Pl. Gram.* 77.—Bras.
subpetiolatum, Steud. l. c. 70 = Ichnanthus petiolatus.
subquadriparum, Trin. Gram. Panic. 145 = distachyum.
subsphaerocarpum, Salzm. ex Schlecht. in Linnaea, xxxi. (1861–62) 483 = Setaria macrostachya.
subspicatum, *Desv. Opusc.* 84.—Am.
subtile, Nees, ex Steud. Nom. ed. II. ii. 264 = distachyum.
subulatum, *Spreng. Syst.* i. 319.—Bras.
subuniflorum, *Bosc, ex Spreng. l. c.* 313.—Carolina.
suffrutescens, *Steud. Syn. Pl. Gram.* 92.—Arab.
sulcatum, Aubl. Pl. Gui. i. 50 = Setaria sulcata.
sulcatum, Bertol. in Opusc. Scient. iv. (1823) 230 = P. Crus-ardeae.
sulphureum, *Nees & Mey. ex Steud. Nom.* ed. II. ii. 264.—China.
sumatrense, Roth, Nov. Pl. Sp. 50 (Paspali sp. ?).—Ind. or.
surinamense, Hochst. ex Steud. Syn. Pl. Gram. 42 = Paspalum scoparium.
sylvaticum, Lam. Encyc. iv. 733 (err. typ. 743) = Oplismenus compositus.
Syzigachne, *Steud. in Flora*, xxix. (1846) 19.—Japon.
tabulatum, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 234.—Austral.
taitense, *Steud. Syn. Pl. Gram.* 418.—Ins. Taiti.
taurinum, *Steud. l. c.* 74.—Tauria.
tectum, Willd. ex Spreng. Syst. i. 313 = dichotomum.
Teff, Desv. Opusc. 97 = Poa abyssinica.

PANICUM:—
tejucense, Nees, ex Trin. Gram. Panic. 166 = Setaria viridis.
telmatodes, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 324.—N. Caled.
tenacissimum, Nees, Agrost. Bras. 238 = Setaria scandens.
tenax, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Setaria setosa.
tenellum, Griff. Notul. iv. 21 = humile.
tenellum, Hort. ex Willd. Enum. Hort. Berol. 89 = Paspalum elegans.
tenellum, *Lam. Illustr.* i. 173; *Encyc.* iv. 753.—Afr. trop.
tenellum, Roxb. Hort. Beng. 7; Fl. Ind. i. 306 = trypheron.
Teneriffae, *R. Br. Prod.* 189.—Reg. Mediterr.; Oriens.
Teneriffae, Parl. in Hook. Niger Fl. 183 = megalanthum.
tenerrimum, *Kunth, Rév. Gram.* i. 39.—Bras.
tenerum, Beyreich, ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 341 = anceps.
tenue, Muhl. Desc. Gram. 118 = dichotomum.
tenue, *Roxb. Fl. Ind. ed. Carey*, i. 313.—Ind. or.
tenuiculmum, G. F. W. Mey. Prim. Fl. Esseq. 58 = laxum.
tenuiflorum, R. Br. Prod. 193 = Paspalum brevifolium.
tenuiflorum, *Schrank, in Syll. Ratisb.* i. (1824) 192.—Bras.
tenuifolium, Hook. & Arn. Bot. Beech. Voy. 100 = nephelophilum.
tenuissimum, *Benth. Fl. Austral.* vii. 470.—Austral.
tenuissimum, Mart. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 26 = Sporobolus minutiflorus.
tenuissimum, Vest, ex Steud. Nom. ed. II. ii. 264 = Polypogon monspeliensis.
ternatum, Hochst. in Flora, xxiv. (1841) I. Intell. 19, nomen; et ex Steud. Syn. Pl. Gram. 40 = glabrum.
tetraspermum, Steud. Nom. ed. II. ii. 257, in syn., sphalm. = *trachyspermum*.
tetrastichum, Forsk. Fl. Aegypt. Arab. 19 = colonum.
texanum, *Vasey, Agric. Grasses U. St.* 36. t. 7.—Texas.
thermale, Boland. in Proc. Calif. Acad. ii. (1863) 181 = dichotomum.
Thouarsianum, Nees, ex Steud. Syn. Pl. Gram. 62 = flavescens.
Thrasya, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 228; *Steud. Syn. Pl. Gram.* 55.—Am. austr.
thrasyoides, *Trin. Gram. Panic.* 126.—Bras.
Thuarii, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = Pennisetum compressum.
timorense, *Kunth, Enum. Pl.* i. 83.—Ins. Timor.
tjicoyaënse, *Steud. Syn. Pl. Gram.* 70.—Java.
tomentosum, *Roxb. Hort. Beng.* 7; *Fl. Ind.* i. 304.—Ind. or.
tonsum, *Steud. Syn. Pl. Gram.* 92.—Afr. austr.
Torreyanum, Wight & Arn. ex Steud. Nom. ed. II. ii. 264 = subeglume.
Torreyi, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 497.—Mexic.
torridum, *Gaudich. in Freyc. Voy. Bot.* 411.—Ins. Sandvic.
trachypus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 258.—Nepal.
trachyrhachis, *Benth. Fl. Austral.* vii. 490.—Austral.
trachyspermum, Nees, Agrost. Bras. 212 = Isachne miliacea.
trachystachyum, *Nees, l. c.* 125.—Bras.
tremulum, Spreng. Neue Entedeck. ii. 103, 202 = brachyanthum.
tricarinatum, Steud. Nom. ed. II. ii. 264 = ovalifolium.
trichanthum, Nees, Agrost. Bras. 210 = microspermum.
trichanthum, A. Rich. Tent. Fl. Abyss. ii. 375 = Hochstetteri.
trichidiachus, *Doell, in Mart. Fl. Bras.* ii. II. 339.—Am. trop.
trichocladum, Reichb. ex Kunth, Enum. Pl. i. 89 = stoloniferum.
trichocondylum, Steud. Syn. Pl. Gram. 74 = maximum, *Jacq.*
trichogonum, Willd. Steud. Nom. ed. II. ii. 264 = pilosum.
trichoides, Cham. & Schlecht. in Linnaea, vi. (1831) 33 = microspermum.

PANICUM:—
trichoides, Sw. Prod. Veg. Ind. Occ. 24 = brevifolium.
tricholaenoides, Steud. Syn. Pl. Gram. 68 = junceum, *Linn.*
trichophorum, Schrad. ex Schult. Mant. ii. 247 = pilosum.
trichopiptum, Steud. Syn. Pl. Gram. 85 = Isachne miliacea.
trichopodon, *A. Rich. Tent. Fl. Abyss.* ii. 369.—Abyss.
trichopus, Hochst. in Flora, xxvii. (1844) 254 = javanicum.
trichostachyum, Hort. ex Steud. Nom. ed. II. ii. 264 = sanguinale.
triflorum, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 179.—Ind. or.
trigonum, *Retz. Obs.* iii. 9.—Ind. or.
trigonum, Wight, ex Steud. Syn. Pl. Gram. 83 = sparsicomum.
trinapalium, Wall. Cat. sub n. 8719, 8722, 8739 = montanum.
trinerve, *Trin. Sp. Gram. Ic.* ii. t. 226.—Bras.
Trinii, *Kunth, Enum. Pl.* i. 88.—Bras.
Trinii, Mor. Syst. Verz. Zoll. 102 = uncinatum.
triquetrum, Willd. ex Doell, in Mart. Fl. Bras. ii. II. 161 = Setaria setosa.
tristachyoides, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 71.—Afr. trop.
triticeum, Willd. ex Spreng. Syst. ii. 609 = Pariana angustifolia.
triticoides, Poir. Encyc. Suppl. iv. 274 = Pennisetum setosum.
truncatum, *Nees, Agrost. Bras.* 215.—Bras.
truncatum, Trin. Gram. Panic. 130 = paspaloides.
trypheron, *Schult. Mant.* ii. 244.—Ind. or.
tuberculatum, J. & C. Presl, Rel. Haenk. i. 307 = maximum, *Jacq.*
tuberculiflorum, Steud. Syn. Pl. Gram. 59 = Eriochloa villosa.
tubulatum, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 234. —Austral.
tumescens, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 316 = aturense.
tumidum, *Koen. in Naturf.* xxiii. (1788) 205; *J. F. Gmel. Syst.* 157.—Hab.?
turgidum, *Forsk. Fl. Aegypt. Arab.* 18.—Afr. bor.; Oriens.
turgidum, Hochst. ex Steud. Syn. Pl. Gram. 88 = nubicum, *Steud.*
turritum, Thunb. Mus. Nat. Acad. Upsal. x. 148, nomen = interruptum.
uliginosum, Roth, Nov. Pl. Sp. 50 = interruptum.
uliginosum, Roxb. Hort. Beng. 8; Fl. Ind. i. 308 = repens.
umbellatum, Trin. Gram. Panic. 238 = parvifolium.
umbraculum, Bosc, ex Spreng. Syst. i. 314 = verrucosum.
umbraticola, *Kunth, Rév. Gram.* i. 33.—Bras.
umbrosum, Le Conte, ex Torr. Cat. Pl. N. York, 91 = verrucosum.
umbrosum, Retz. Obs. iv. 16 = prostratum.
umbrosum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 210 = stoloniferum.
uncinatum, *Raddi, Agrost. Bras.* 41.—Reg. trop.
uncinulatum, *R. Br. Prod.* 191.—Austral.
unciphyllum, Trin. Gram. Panic. 242 = dichotomum.
undatum, Steud. Nom. ed. II. ii. 264 = Oplismenus compositus.
undulatifolium, Ard. Animad. Spec. Alt. 14 = Oplismenus undulatifolius.
unguiculatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 275.—Afr. austr.
unguinosum, Herb. Banks, ex Roxb. Fl. Ind. i. 294 = Oplismenus compositus.
uniglume, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 370.—Abyss.
unilaterale, *Desv. Opusc.* 86.—Am. austr.
unisetum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 217, in adnot. = Setaria uniseta.
Urochloa, *Desv. Opusc.* 82.—Ins. Maurit.
Urochloa, *Steud. Nom.* ed. II. ii. 264.—Ins. Philipp.
Urvilleanum, *Kunth, Rév. Gram.* i. 35; ii. 405. t. 115. —Chili.
utriculatum, Steud. Nom. ed. II. ii. 264 = Setaria vaginata.
vacillans, Steud. Syn. Pl. Gram. 75 = sarmentosum.
vaginaeflorum, Steud. l. c. 80 = decumbens.
vaginatum, Gren. & Godr. Fl. Fr. iii. 462 = Paspalum distichum.

PANICUM:—
vaginatum, Nees, Agrost. Bras. 156 = macranthum.
vagum, Scop. Fl. Carn. ed. II. i. 51 = Polypogon monspeliensis.
Valenzuelanum, *A. Rich. Fl. Cub. Fanerog.* ii. 304.—Cuba.
variegatum, Hort.—Cf. Gard. Chron. (1867) 458 = Oplismenus Burmannii.
velutinosum, *Nees, ex Trin. Gram. Panic.* 144.—Am. trop.
velutinum, *Bosc, ex Spreng. Syst.* i. 315 = dichotomum.
velutinum, Kunth, Enum. Pl. i. 92 = velutinosum.
velutinum, G. F. W. Mey. Prim. Fl. Esseq. 51 = Oplismenus Burmannii?
Ventenatii, Steud. Nom. ed. II. ii. 265 = Setaria glauca.
ventricosum, Lam. Illustr. i. 173; Encyc. iv. 745 = Isachne ventricosa.
verrucosum, *Muhl. Desc. Gram.* 113.—Am. bor.
versicolor, Doell, in Mart. Fl. Bras. ii. II. 254 = truncatum.
verticillatum, Linn. Sp. Pl. ed. II. 82 = Setaria verticillata.
verticillatum, Rottl. ex Spreng. Syst. i. 304 = Setaria glauca.
verticillatum, Ucria, Hort. Reg. Panorm. 54 = Agrostis stolonifera.
vestitum, *Kunth, Rév. Gram.* i. 39.—Bras.
vestitum, Nees, ex Steud. Syn. Pl. Gram. 62 = villosum.
viaticum, Griff. Ic. Pl. Asiat. t. 145. f. 2 = semialatum.
viaticum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 155 = prostratum.
vicarium, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 498.—Mexic.
vilfiforme, *Wood, Class Book* (1861), 785.—Am. bor.
vilfoides, *Trin. Gram. Panic.* 171 (vilvoides).—Am. trop.
villiferum, Nees, in Linnaea, xix. (1847) 687 = adscendens.
villosum, R. Br. Prod. 192 = lanatum.
villosum, Ell. Sketch, i. 124 = dichotomum.
villosum, Hoffmgg. ex Steud. Nom. ed. II. ii. 265 = Ichnanthus Hoffmannseggii.
villosum, *Lam. Illustr.* i. 173.—As. trop.
villosum, Presl, Cyp. & Gram. Sic. 18 = Teneriffae.
vilvoides, Trin. vide vilfoides.
vimineum, *Schrad. in Schult. Mant.* ii. 255.—Bras.
violaceum, Klein, ex Thiele, in Linnaea, ix. (1834) 307 = Isachne australis.
violaceum, Lam. ex Spreng. Syst. i. 303 = Pennisetum typhoideum.
violaceum, Llanos, Fragm. 42 = nodosum.
violaceum, Rottl. ex Sw. Adnot. Bot. 7 = Rottleri.
violascens, *Kunth, Rév. Gram.* i. 33.—Am. trop.
virescens, *Poir. Encyc. Suppl.* iv. 278.—Hab.?
virescens, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 157 = Setaria viridis.
virgatum, *Linn. Sp. Pl.* 59.—Am. bor.
virgatum, F. Muell. Fragm. viii. 192 = trachyrhachis.
virgatum, Muhl. Desc. Gram. 120 = coloratum.
virgatum, Roxb. ex Steud. Nom. ed. II. ii. 265 = psilopodium.
virgatum, Willd. ex Spreng. Syst. i. 318 = avenaceum.
viride, Desf. Fl. Atlant. i. 58 = Setaria verticillata.
viride, Linn. Syst. ed. X. 870 = Setaria viridis.
viride, Pollich, Hist. Pl. Palat. i. 55 = Setaria glauca.
viridescens, *Steud. Syn. Pl. Gram.* 51.—China.
viridiflorum, Nees, Agrost. Bras. 135 = aturense.
Virletii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 498.—Mexic.
viscidum, *Ell. Sketch*, i. 123.—Am. bor.
viviparum, *Nees, ex Steud. Syn. Pl. Gram.* 97 (an seq.?). —Afr. trop.
viviparum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 62.—Afr. trop.
vulgare, Wallr. in Linnaea, xiv. (1840) 542; et Beitr. Fl. Hercyn. 172 = Setaria italica.
vulpinum, Willd. Enum. Hort. Berol. 1031 = Pennisetum cenchroides.
vulpisetum, *Lam. Encyc.* iv. 735 (err. typ. 745).—Ins. S. Doming.
Walkeri, Steud. Syn. Pl. Gram. 97 = Isachne Walkeri.
Wallichianum, Nees, Fl. Afr. Austr. 49 = plicatum, *Perrottetii.*
Walteri, Ell. Sketch, i. 115 = Curtisii.
Walteri, Muhl. Desc. Gram. 108 = carolinianum.
Walteri, Poir. Encyc. Suppl. iv. 282 = latifolium.

PANICUM:—
Walteri, Pursh, Fl. Am. Sept. i. 66 = P. Crus-galli.
Wightianum, Arn. & Nees, ex Nees, Fl. Afr. Austr. 30 = seq.
Wightii, Nees, l. c. 29 = P. Isachne.
Willdenowii, Steud. Nom. ed. II. ii. 265 = Setaria luxurians.
Williamsii, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 250.—China.
xalapense, *H. B. & K. Nov. Gen. et Sp.* i. 103.—Mexic.
xanthophysum, *A. Gray, Man. Bot. N. U. St.* ed. I. 613.—Am. bor.
xanthorrhizum, *Steud. Syn. Pl. Gram.* 59.—Java?
Zenkowskii, *Rupr. Ind. Sem. Hort. Petrop.* (1851). —*Cf. Linnaea*, xxvii. (1854) 477.—Hab.?
Zeyheri, *Nees, Fl. Afr. Austr.* 25.—Afr. austr.
zeylayense, *Steud. Nom.* ed. II. ii. 265.—Mexic.
zizanioides, H. B. & K. Nov. Gen. et Sp. i. 100 = oryzoides.
Zollingeri, Steud. Syn. Pl. Gram. 96 = Isachne albens.
zonale, Guss. Fl. Sic. Prod. i. 82 = colonum.

PANIOPSIS, Rafin. Fl. Tellur. ii. 49 (1836) = **Inula**, Linn. (Compos.).
fetidum, Rafin. l. c. = Nidorella foliosa.
glutinosum, Rafin. l. c. = Grindelia glutinosa.
graveolens, Rafin. l. c. = I. graveolens.
viscosum, Rafin. l. c. = I. viscosa.

PANIOS, Adans. Fam. ii. 124 (1763) = **Erigeron**, Linn. (Compos.).

PANISEA, Lindl. Fol. Orch. Panis. i. (1854). *ORCHIDEAE*, Benth. & Hook. f. iii. 507.
ANDROGYNE, Griff. Notul. iii. 279 (1851).
apiculata, *Lindl. Fol. Orch. Panis.* 2.—Burma.
bilamellata, *Reichb. f. in Walp. Ann.* vi. 240.—Ins. Philipp.
parviflora, *Lindl. Fol. Orch. Panis.* 1.—Reg. Himal.
reflexa, Lindl. l. c. = parviflora.
uniflora, Lindl. l. c. 2 = Coelogyne uniflora.

PANISIA, Rafin. Sylva Tellur. 128 (1838) = **Cassia**, Tourn. (Legumin.).
biflora, Rafin. l. c. 129 = C. biflora.

PANKE, Molina, Sagg. Chile, 143 (1782) = **Gunnera**, Linn. (Halorag.).
acaulis, Molina, l. c. 145 = G. chilensis.
caulescens, J. F. Gmel. Syst. 652 = G. chilensis?
sonchifolia, Willd. Sp. Pl. ii. 487 = Francoa sonchifolia.
tinctoria, Molina, Sagg. Chile, 143 = G. chilensis.

PANKEA, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 189, 192 = **Gunnera**, Linn. (Halorag.).
bracteata, Oerst. l. c. 192 = G. bracteata.
chilensis, Oerst. l. c. = G. chilensis.
insignis, Oerst. l. c. = G. insignis.

PANOPE, Rafin. Fl. Tellur. ii. 103 (1836) = **Lippia**, Linn. (Verben.).
stechadifolia, Rafin. l. c. = L. stoechadifolia.

PANOPIA, Noronha, ex Thou. Gen. Nov. Madag. 26 (1806) = **Macananga**, Thou. (Euphorb.).

PANOPSIS, Salisb. in Knight, Prot. 104 (1809). *PROTEACEAE*, Benth. & Hook. f. iii. 177.
ANDRIAPETALUM, Pohl, Pl. Bras. Ic. i. 113 (1827).
hameliaefolia, *Knight, Prot.* 104.—Guian.
hamelifolia, Spreng. Syst. i. 483 = praec.

PANOXIS, Rafin. Med. Fl. ii. 109 (1830) = **Veronica**, Tourn. (Scrophul.).
cataractae, Rafin. l. c. = V. cataractae.
macrocarpa, Rafin. l. c. = V. macrocarpa.
salicifolia, Rafin. l. c. = V. salicifolia.

PANPHALEA, Lag. Amen. Nat. i. 34 (1811) = **Pamphalea**, DC. (Compos.).

PANSTENUM, Rafin. Fl. Tellur. ii. 20 (1836) = **Allium**, Tourn. (Liliac.).
monspessulanum, Rafin. l. c. 21 = A. nigrum?

PANSTREPIS, Rafin. Fl. Tellur. iv. 41 (1836) = **Coryanthes**, Hook. (Orchid.).
paradoxa, Rafin. l. c. = C. macrantha.

PANTASACHME, Endl. Gen. 590 (1838) sphalm. = **Pentasachme**, Wall. (Asclep.).

PANTATHERA, Phil. in Bot. Zeit. xiv. (1856) 649.
GRAMINEAE, Benth. & Hook. f. iii. 1200.
avenacea, Hemsl. Bot. Voy. Challeng. t. 61 = fernandeziana.
fernandeziana, *Phil. in Bot. Zeit.* xiv. (1856) 649.—Ins. J. Fernand.

PANTERPA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 195 = **Bignonia**, Linn.
dichotŏma, Miers, l. c. 196 = B. dichotoma.
extensa, Miers, l. c., nomen = B. extensa.
Leucopogon, Miers, l. c. = B. Leucopogon.
mollissima, Miers, l. c. = B. mollissima.
palpebrosa, Miers, l. c., nomen.—Bras.
Prieuri, Miers, l. c. = B. Prieurei.
prolixa, Miers, l. c., nomen.—Bras.
puberula, Miers, l. c. = Cuspidaria puberula.
trajecta, Miers, l. c., nomen.—Bras.
varians, Miers, l. c. = B. varians.

PANTHOCARPA, Rafin. Neogenyt. 2 (1825) = **Mimosa**? Linn. (Legumin.).
chionacantha, Rafin. l. c. = *M. chionacantha*.

PANTOCSEKIA, Griseb. ex Pantoč. in Oestr. Bot. Zeitzschr. xxiii. (1873) 267 = **Convolvulus**, Tourn.
illyrica, Griseb. l. c.—Europ. austr.

PANUREA, Spruce, ex Benth. & Hook. f. Gen. i. 553 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
longifolia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxv. (1865) 301.—Am. trop.

PANZA, Salisb. Gen. Pl. Fragm. 99 (1866) = **Narcissus**, Tourn. (Amaryll.).
bicolor, Salisb. l. c. = N. Macleayi.

PANZERA, Willd. Sp. Pl. ii. 540 (1799) = **Eperua**, Aubl. (Legumin.).
falcata, Willd. l. c. = E. falcata.

PANZERIA, J. F. Gmel. Syst. 247 (1791) = **Lycium**, Linn. (Solan.).
caroliniana, J. F. Gmel. l. c. = L. carolinianum.

PANZERIA, Moench, Meth. 402 (1794) = **Leonurus**, Linn. (Labiat.).
canescens, Bunge, Del. Sem. Hort. Dorpat. (1839) 8 = L. lanatus.
lanata, Bunge, l. c. (1836) 5 = L. lanatus.
multifida, Moench, Meth. Suppl. 137 = L. sibiricus.
sibirica, Hort. ex Steud. Nom. ed. II. ii. 265 = L. sibiricus.
tatarica, Hort. ex Steud. l. c = L. tataricus.
tomentosa, Moench, Meth. 402 = L. lanatus.
tripartita, Moench, l. c. = L. tataricus.
viridis, Bunge, Del. Sem. Hort. Dorpat. (1839) 8.—Hab.?

PAPAVER, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 150 (1737). *PAPAVERACEAE*, Benth. & Hook. f. i. 51.
CALOMECON, Spach, Hist. Vég. Phan. vii. 7. (1839).
CERASTITES, S. F. Gray, Nat. Arr. Brit. Pl. ii. 703 (1821).
CLOSTERANDRA, Boiv. ex Bélang. Voy. Indes Or., tab. absque num. (1839).
aculeatum, *Thunb. Prod. Pl. Cap.* 92.—Afr. austr.; Austral.
agrivagum, Jord. Diagn. i. 96 = P. Rhoeas.
album, Mill. Gard. Dict. ed. VIII. n. 9 = somniferum.
album-&-nigrum, Crantz, Stirp. Austr. ed. I. fasc. ii. 129; ed. II. 138 = somniferum.
alpinum, Linn. Sp. Pl. 507 = nudicaule.
alpinum, Salisb. Prod. 377 = Meconopsis cambrica.
ambiguum, Jord. & Four. Brev. Pl. Nov. fasc. i. 3 = dubium.
amoenum, Lindl. Bot. Reg. (1839) Misc. 56 = somniferum.
amplexicaule, Stokes, Bot. Mat. Med. iii. 181 = somniferum.
angustulum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 2 = dubium.

PAPAVER:—
apulum, *Tenore, Fl. Neap. App.* v. 16.—Europ. austr.; Oriens.
arenarium, *Bieb. Fl. Taur. Cauc.* iii. 364.—Reg. Caucas. et Casp.
Argemone, *Linn. Sp. Pl.* 506.—Europ.; Afr. bor.
argemonium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
argemonoides, Cesati, in Bibl. Ital. xci. (1838) 346 = apulum.
armeniacum, Lam. Encyc. i. 247 = caucasicum.
arvaticum, Jord. Diagn. i. 95 = P. Rhoeas.
arvense, Borkh. in Rhein. Mag. i. (1793) 439 = P. Argemone.
arvense, Salisb. Prod. 376 = P. Rhoeas.
atropurpureum, Gilib. Fl. Lituan. ii. 214 = P. Rhoeas.
aurantiacum, Loisel. in Desv. Journ. Bot. ii. (1809) 340 = nudicaule.
Belangeri, *Boiss. Fl. Orient.* i. 117.—Persia.
bipinnatum, *C. A. Mey. Verz. Pfl. Cauc.* 175.—Reg Caucas.
bracteatum, *Lindl. Coll. Bot.* t. 23.—Reg. Caucas.; Persia.
breviflorum, Gilib. Fl. Lituan. ii. 214 = dubium.
Burseri, Crantz, Stirp. Austr. ed. I. fasc. ii. 129; ed. II. 138 = nudicaule.
cambricum, Linn. Sp. Pl. 508 = Meconopsis cambrica.
caucasicum, *Bieb. Fl. Taur. Cauc.* ii. 5.—Reg. Caucas.
caucasicum, J. Henning, in Mém. Soc. Nat. Mosc. vi. (1823) 73 = arenarium.
caudatifolium, *Timb. in Bull. Soc. Hist. Nat. Toul.* iv. (1870) 163.—Gallia.
cereale, Jord. Diagn. i. 97 = P. Rhoeas.
chelidoniaefolium, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 11.—Persia.
clavatum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 112.—Syria.
clavatum, Gilib. Fl. Lituan. ii. 215 = P. Argemone.
clavigerum, Lam. Fl. Fr. iii. 175 = P. Argemone.
collinum, Bogenh. ex Bisch. Del. Sem. Hort. Heidelb. (1849) 4 = dubium.
commutatum, Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iv. 41 = P. Rhoeas.
confine, Jord. Diagn. i. 89 = dubium.
corniculatum, Pall. Reise, iii. 591 = Glaucium corniculatum.
cornigerum, Stocks, in Hook. Kew Journ. iv. (1852) 142 = pavoninum.
corsicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 4 = dubium.
corsicum, *Thouin, ex Schrank, in Syll. Ratisb.* i. (1824) 219.—Corsica.
croceum, Ledeb. Fl. Alt. ii. 271 = nudicaule.
cruciatum, Jord. Diagn. i. 97 = P. Rhoeas.
davuricum, *Fisch. ex Schrank, in Syll. Ratisb.* i. (1829) 219.—Dahuria.
Decaisnei, *Hochst. & Steud. ex Elkan, Tent. Monog. Papav.* 26.—Arab.; Persia; Afghan.
depressum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 4 = dubium.
dodecandrum, *Medic. Beobacht.* (1783) 307.—Hab.?
Dodonei, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1870) 161 = P. Rhoeas.
dubium, *Linn. Sp. Pl.* 1196.—Europ.
dubium, Tenore, Fl. Nap. iv. 306 = pinnatifidum.
erosulum, Jord. Diagn. i. 88 = laevigatum.
errabundum, Jord. l. c. 93 = dubium.
erraticum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 705; Jord. l. c. 96 = P. Rhoeas.
erroneum, Jord. l. c. 91 = dubium.
flavum, Moench, Meth. 247 = Meconopsis cambrica.
floribundum, Desf. in Ann. Mus. Par. vi. (1805) 376 = caucasicum.
frenatum, Steud. Nom. ed. II. ii. 266, sphalm. = seq.
frentanum, *Tenore, Relaz. Viag. Abruzz.* 124.—Italia.
Fuchsii, Timb. in Bull. Soc. Hist. Nat. Toulouse, vi. (1870) 162 (*Fuschii*) = P. Rhoeas.
fugax, Poir. Encyc. v. 118 = caucasicum.
gariepinum, Burch. ex DC. Syst. ii. 79; et in Burch. Trav. i. 318 = aculeatum.
glabellum, Stev. ex DC. l. c. 78 = laevigatum.
glabrum, Gilib. Fl. Lituan. 213; Exercit. Bot. i. 337 = somniferum.
glabrum, Royle, Illustr. Bot. Himal. 67 = dubium.
glaucioides, *Roux, in Bull. Soc. Bot. et Hort. Prov.* i. (1879) 83.—Gallia.
glaucum, *Boiss. & Hausskn. ex Boiss. Fl. Orient.* i. 116.—Syria.

PAPAVER:—
gracile, *Auch. ex Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 372.—As. Min.
grandiflorum, Moench, Meth. 247 = orientale.
Heldreichii, *Boiss. Fl. Orient.* i. 108.—As. Min.
hispidum, Desf. ex Elkan, Tent. Monog. Papav. 22 = caucasicum.
hispidum, Lam. Fl. Fr. iii. 174 = hybridum.
Hookeri, *Baker, ex Hook. f. Bot. Mag.* t. 6729.—Hab.?
horridum, DC. Syst. ii. 79 = aculeatum.
hortense, Hussen, Chard. Nanc. Fl. 39 = somniferum.
hybridum, *Linn. Sp. Pl.* 506.—Europ.; Syria; Afr. bor.
hybridum, Spenn. Fl. Friburg. iii. 979 = P. Argemone.
hyoscyamifolium, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 110.—Syria.
improperum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 3 = dubium.
indehiscens, Dum. Fl. Belg. Prod. 130 = somniferum.
inornatum, Schott & Kotrchy, ex Schott, in Oestr. Bot. Wochenbl. iv. (1854) 233 = polytrichum.
insignitum, Jord. Diagn. i. 94 = P. Rhoeas.
integrifolium, *Viquier, Hist. Nat. Pavots*, 38.—Hispan.
intermedium, Becker, Fl. Frankf. i. 386 = P. Rhoeas.
intermedium, DC. 7me Not. Pl. Rar. Jard. Genèv. 37 = bracteatum.
laevigatum. *Bieb. Fl. Taur. Cauc.* iii. 364.—Graecia; Oriens.
Lamottei, Bor. Fl. Centr. Fr. ed. III. ii. 30 = dubium.
lateritium, *C. Koch, Ind. Sem. Hort. Berol.* (1854) 14.—Armenia.
Lecoqii, Lamotte, Not. Pap. Dub. in Ann. Auvergne, (1851) 429 = dubium.
leiocarpum, Turcz. in Bull. Soc. Nat. Mosc. xi. (1838) 1. 86 = nudicaule.
libanoticum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 373.—Syria.
luteo-rubrum, Jord. Diagn. i. 92 = dubium.
luteum, Lam. Fl. Fr. iii. 173 = Meconopsis cambrica.
macrostomum, *Boiss. & Huet, Diagn.* Ser. II. v. 15.—As. Min.; Persia.
maculosum, Schur, Enum. Pl. Transs. 36 = dubium.
maritimum, With. Arr. Brit. Pl. ed. III. 486 = P. Argemone.
micranthum, Bor. Fl. Centr. Fr. ed. III. ii. 29 = P. Argemone.
microcarpum, DC. Syst. ii. 71 = nudicaule.
miniatum, Reichb. Pl. Crit. viii. 19 = nudicaule.
mixtum, Jord. Diagn. i. 91, in obs; Ic. t. 66 = dubium.
modestum, Jord. Pugill. Pl. Nov. 4 = laevigatum.
monanthum, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 393.—Reg. Caucas.
multiflorum, Burm. f. Fl. Cap. Prod. 15 = aculeatum?
nothum, Stev. ex Nym. Consp. 24 = laevigatum.
nudicaule, *Linn. Sp. Pl.* 507.—Reg. alp. et arct.
nudum, Burm. f. Fl. Cap. Prod. 15 = aculeatum?
obtusifolium, *Desf. Fl. Atlant.* i. 407.—Alger.
officinale, C. C. Gmel. Fl. Bad. ii. 479 = somniferum.
oligophyllum, Bélang. Voy. Indes Or., tab. absque num. = tenuifolium.
olympicum, Sibth. Fl. Graec. v. 75 = pilosum.
opiiferum, Forsk. Fl. Aegypt. Arab. p. cxiii. n. 336 = somniferum.
oreophilum, *Rupr. Fl. Cauc.* 51.—Reg. Caucas.
orientale, Bieb. Fl. Taur. Cauc. ii. 5; iii. 365 = bracteatum.
orientale, *Linn. Sp. Pl.* 508.—As. Min.; Persia.
paniculatum, D. Don, Prod. Fl. Nep. 197 = Meconopsis napalensis.
parviflorum, Lam. Fl. Fr. iii. 173 = dubium.
pavoninum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 10 = apulum.
pavoninum, *Mey. in Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. 82.—Afghan.; Turkestan.
persicum, Hort. Dijon, ex Elkan, Tent. Monog. Papav. 21 = caucasicum.
persicum, *Lindl. Bot. Reg.* t. 1570.—Persia.
pilosum, *Sibth. & Sm. Prod.* i. 360; *Fl. Graec.* v. 75. t. 492.—Bithynia.
pinnatifidum, *Moris, Fl. Sard.* i. 74.—Ital.; Sardoa.
polycephalum, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 594 = somniferum.
polychaetum, *Schott & Kotschy, ex Boiss. Fl. Orient.* i. 111.—As. Min.
polytrichum, *Boiss. & Kotschy, Diagn.* Sér. II. v. 14.—As. Min.; Syria.

PAPAVER:—
pulcherrimum, Fisch. ex Steud. Nom. ed. II. ii. 266 = bracteatum.
pyrenaicum, Willd. Enum. Hort. Berol. 563 = nudicaule.
radicatum, Rottb. ex DC. Prod. i. 118 = nudicaule.
ramosissima, Hort. Prag. ex Elkan, Tent. Monog. Papav. 21 = caucasicum.
raripilum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 3 = dubium.
Rhoeas, *Linn. Sp. Pl.* 507.—Europ.; Oriens; Afr. bor.
Rhoeas, Pall. Reise, iii. 546 = arenarium.
Roubiaei, Viquier, Hist. Nat. Pavots, 39 = P. Rhoeas.
rubro-aurantiacum, Fisch. ex Steud. Nom. ed. II. ii. 266 = nudicaule.
rupifragum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 6. Hispan.; Marocc.
rusticum, Jord. Diagn. i. 99 = P. Rhoeas.
segetale, Jord. l. c. 98 = P. Rhoeas.
segetale, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 979 = P. Rhoeas, dubium.
setigerum, DC. Fl. Fr. Suppl. 585 = somniferum.
siculum, Guss. Fl. Sic. Syn. ii. 6 = hybridum.
simplicifolium, D. Don, Prod. Fl. Nep. 197 = Meconopsis simplicifolia.
somniferum, *Linn. Sp. Pl.* 508.—Graecia; Oriens.
spectabile, Link, ex Elkan, Tent. Monog. Papav. 22 = caucasicum.
spectabile, Salisb. Prod. 377 = orientale.
spicatum, *Boiss. & Bal. Diagn.* Ser. II. v. 14.—As. Min.
strictum, *Boiss. & Bal. l. c.* vi. 8.—Phrygia.
strigosum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 66 = P. Rhoeas.
stylatum, *Boiss. & Bal. Diagn.* Ser. II. v. 13.—Cilicia.
suaveolens, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 72 = nudicaule.
sylvestre, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 23 (Quid?).—Gallia.
syriacum, Boiss. & Blanche, Diagn. Ser. II. vi. 8 = P. Rhoeas.
tauricolum, *Boiss. Fl. Orient.* i. 109.—As. Min.
tauricum, Hort. Kiel. ex Elkan, Tent. Monog. Papav. 21 = caucasicum.
tenue, *Ball, in Journ. Bot.* xi. (1873) 296.—Marocc.
tenuifolium, *Boiss. & Hohen. Diagn.* Ser. I. viii. 10.—Persia.
trilobum, Wallr. Ann. Bot. 149 = P. Rhoeas.
triniaefolium, *Boiss. Fl. Orient.* i. 110.—Cappadoc.
turbinatum, *DC. Syst.* ii. 84.—Assyria.
turbinatum, Fresen. in Mus. Senkenb. i. (1834) 173 = Decaisnei.
umbonatum, *Boiss. Diagn.* Ser. I. viii. 11.—Syria.
umbrosum, Hort.—Cf. Gard. Chron. (1876) II. 16 = P. Rhoeas.
uniflorum, Balb. ex Spenn. Fl. Friburg. iii. 980 = P. Rhoeas, dubium.
vagum, Jord. Diagn. i. 90 = dubium.
villosum, C. Koch, in Linnaea, xix. (1847) 49 = pilosum.
virgatum, Sm. in Rees, Cycl. xxvi. n. 9 = caucasicum.

PAPAYA, Tourn. ex Linn. Syst. ed. I (1735) = **Carica**, Linn. (Passifl.).
Carica, Gaertn. Fruct. ii. t. 122 = C. Papaya.
cauliflora, Poir. Encyc. Suppl. iv. 296 = C. cauliflora.
citriformis, A. DC. in DC. Prod. xv. 1. 414 = C. citriformis.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = C. Papaya.
cucumerina, Noronha, l. c. = C. Papaya.
edulis, Boj. Hort. Maurit. 277.—Hab.?
microcarpa, Poir. Encyc. Suppl. iv. 296 = C. microcarpa.
monoica, Poir. l. c. 297 = C. monoica.
nana, A. DC. in DC. Prod. xv. 1. 415 = C. nana.
Posoposa, DC. in Poir. Encyc. v. 3 = C. Posoposa.
quercifolia, Baill. Hist. Pl. iv. 285 (= *Vasconcellia quercifolia*.).—Am. trop.
sativa, Tuss. Fl. Antill. iii. 45 = C. Papaya.
spinosa, DC. Poir. Encyc. v. 3 = C. spinosa.
vulgaris, DC. l. c. v. 2 = C. Papaya.

PAPEDA, Hassk. in Flora, xxv. (1842) II. Beibl. 42; et Cat. Hort. Bog. Alt. 216 (1844) = **Citrus**, Linn. (Rutac.).
Rumphii, Hassk. ll. cc. = C. Hystrix.

PAPHIA, Seem. in Journ. Bot. ii. (1864) 77 = **Agapetes**, G. Don (Vaccin.).
vitiensis, Seem. l. c.—Ins. Fiji.

PAPHINIA, Lindl. Bot. Reg. (1843) Misc. 14 = **Lycaste**, Lindl. (Orchid.).
cristata, Lindl. l. c. = L. cristata.
grandiflora, Rodrig. Orch. Nov. i. 124.—Bras.
grandis, Reichb. f. ex Warn. Orch. Alb. iv. t. 145 = *grandiflora*.
nutans, Houllet, in Rev. Hort. (1878) 188 = *grandiflora*?
rugosa, Reichb. f. in Linnaea, xli. (1877) 110.—N. Granat.

PAPILLARIA, Dulac, Fl. Hautes-Pyr. 45 (1867) = **Scheuchzeria**, Linn. (Naiad.).
palustris, Dulac, l. c. = S. palustris.

PAPIRIA, Thunb. in Phys. Sällsk. Handl. Lund, i. (1776) 110 = **Gethyllis**, Plum. (Amaryll.).
ciliaris, Thunb. l. c. 111 = G. ciliaris.
lanceolata, Thunb. l. c. 112 = Apodolirion lanceolatum.
spiralis, Thunb. l. c. 111 = G. spiralis.
villosa, Thunb. l. c. = G. villosa.

PAPPEA, Eckl. & Zeyh. Enum. 53 (1835). *SAPINDACEAE*, Benth. & Hook. f. i. 407.
capensis, *Eckl. & Zeyh. l. c.*—Afr. austr.

PAPPEA, Sond. & Harv. Fl. Cap. ii. 562 (1862) = **Choritaenia**, Benth. (Umbell.).
capensis, Sond. & Harv. l. c.—Afr. austr.

PAPPERITZIA, Reichb. f. in Bot. Zeit. x. (1852) 670. *ORCHIDEAE*.
Leiboldi, *Reichb. f. l. c.*—Mexic.

PAPPOCHROMA, Rafin. Fl. Tellur. ii. 48 (1836) = **Erigeron**, Linn. (Compos.).
uniflora, Rafin. l. c. = E. Pappochroma.

PAPPOPHORUM, Schreb. Gen. ii. 787 (1791). *GRAMINEAE*, Benth. & Hook. f. iii. 1174.
ENNEAPOGON, Desv. ex Beauv. Agrost. 81 (1812); et Journ. Bot. iii. (1814) 70.
EURAPHIS, Trin. ex Lindl. Veg. Kingd. 115 (1847).
POLYRHAPHIS, Lindl. l. c. (1847).
abyssinicum, *Hochst. in Flora*, xxxviii. (1855) 202.—Abyss.
alopecuroideum, *Vahl, Symb. Bot.* iii. 10. t. 51.—Am. austr.
apertum, *Munro, ex Scribner, in Bull. Torrey Bot. Club*, ix. (1882) 148.—N. Mexic.
arabicum, Hochst. ex Steud. Syn. Pl. Gram. 199 = brachystachyum.
Aucheri, Jaub. & Spach, Illustr. iv. 32 = persicum.
avenaceum, *Lindl. in Mitch. Journ. Trop. Austral.* 320.—Austral.
bicolor, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 568.—Mexic.
boreale, *Griseb. in Ledeb. Fl. Ross.* iv. 404.—Sibir. baical.
brachystachyum, *Jaub. & Spach, Illustr.* iv. 34.—Afr. bor.; Arab.
bulbosum, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 254 = brachystachyum.
caerulescens, Gaudich. in Freyc. Voy. Bot. 409 = nigricans.
cenchroides, *Licht. in Roem. & Schult. Syst.* ii. 616.—Afr. austr.
commune, F. Muell. Fragm. viii. 200 = nigricans.
elegans, *Nees, ex Steud. Syn. Pl. Gram.* 199.—Ind. or.
elongatum, Spreng. Syst. iv. Cur. Post. 34 = alopecuroideum.
eutrianoides, Trin. ex Nees, Agrost. Bras. 414 = Bouteloua multiseta.
Figarianum, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 254 = brachystachyum.
flavescens, Lindl. in Mitch. Journ. Trop. Austral. 34 = nigricans.
glandulosum, Berg. ex Nees, Fl. Afr. Austr. 298 = Danthonia Buekeana.
glumosum, *Hochst. in Flora*, xxxviii. (1855) 202.—Afr. trop.
gracile, R. Br. Prod. 185 = nigricans.

PAPPOPHORUM:—
Jaminianum, Coss. & Dur. ex Boiss. Fl. Orient. v. 558 = brachystachyum.
laguroideum, Schrad. in Schult. Mant. ii. 342 = alopecuroideum.
macrostachyum, Schrad. l. c. = alopecuroideum.
megapotamicum, Spreng. Syst. iv. Cur. Post. 34 = alopecuroideum.
mexicanum, *Griseb. ex Hemsl. Biol. Centr. Am. Bot.* iii. 568.—Mexic.
molle, *Kunth, Enum. Pl.* i. 255.—Afr. austr.
mucronulatum, *Nees, in Mart. Fl. Bras.* ii. 412.—Bras.
nanum, Steud. ex J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 144; Syn. Pl. Gram. 200 = Vincentianum.
nigricans, *R. Br. Prod.* 185.—Austral.; N. Guin.
pallidum, R. Br. l. c. = nigricans.
persicum, *Steud. Syn. Pl. Gram.* 200.—Persia; Afghan.; Turkest.
phleoides, [*Lag.*] *Hort. Matr. ex Roem. & Schult. Syst.* ii. 616.—Arabia.
phleoides, Turcz. in Bull. Soc. Nat. Mosc. xxix. (1856) 1. 30 = boreale.
polystachyum, Kunth, Rév. Gram. ii. t. 131 = alopecuroideum.
Pumilio, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 92 = Boissiera bromoides.
pungens, Trin. l. c. = Triraphis pungens.
purpurascens, R. Br. Prod. 185 = nigricans.
saccharoides, *Griseb. in Goett. Abh.* xxiv. (1879) 301.—Paraguay.
scabrum, *Kunth, Enum. Pl.* i. 255.—Afr. austr.
Schimperianum, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 403.—Abyss.
senegalense, *Steud. Syn. Pl. Gram.* 199.—Afr. trop.
sinaicum, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. II. (1838) 54 = Boissiera bromoides.
squarrosum, [Soland. in] Russ. Aleppo, ed. II. ii. 244 = Boissiera bromoides.
turcomanicum, *Trautv. in Act. Hort. Petrop.* i. (1871) 27.—Turkestan.
vaginatum, *Phil. Sert. Mendoc. Alt.* 48.—Chili.
Vincentianum, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 144.—Ins. Cap. Virid.
virens, Lindl. in Mitch. Journ. Trop. Austral. 360 = nigricans.
Wrightii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 178.—Mexic.

PAPULARIA, Forsk. Fl. Aegypt. Arab. 69 (1775) = **Trianthema**, Linn. (Ficoid.).
crystallina, Forsk. l. c. = T. crystallina.

PAPYRIUS, Lam. Illustr. t. 762 (1798) = **Broussonetia**, Vent. (Urtic.).
japonica, Lam. Encyc. v. 3 = B. papyrifera.
polimorphus, Cav. Desc. 343 = B. papyrifera.

PAPYRUS, Willd. in Abh. Acad. Berl. (1812–13) 70 = **Cyperus**, Mich.
aequalis, Boj. Hort. Maurit. 381 = C. aequalis.
antiquorum, Willd. in Abh. Acad. Berl. (1812) 70 = C. Papyrus.
comosus, Willd. l. c. 74 = C. comosus.
corymbosus, Nees, in Wight, Contrib. 89 = C. tegetum.
dehiscens, Nees, in Wight, l. c. = C. tegetum.
domesticus, Poir. in Dict. Sc. Nat. xxxvii. 447 = C. Papyrus.
elatus, Nees, in Wight, Contrib. 88 = C. auricomus.
elegans, G. Don, ex Loud. Hort. Brit. 22 = C. elegans.
elegans, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 59 = C. comosus.
giganteus, Schrad. ex Nees, l. c. 58 = C. giganteus.
junciformis, Schrad. ex Nees, l. c. 57. t. 3 = C. spicigerus.
latifolius, Willd. in Abh. Acad. Berl. (1812) 74 = C. platyphyllus.
laxiflorus, Spreng. Syst. i. 232 = C. aequalis.
madagascariensis, Willd. in Abh. Acad. Berl. (1812) 72 = C. madagascariensis.
mossambicensis, Parl. in Peters, Reise Mossamb. Bot. 582 (Index), t. 59 = C. Papyrus.
odoratus, Willd. ex Nees, in Mart. Fl. Bras. ii. 1. 58 = C. giganteus.
Pangorei, Nees in Wight, Contrib. 88 = C. corymbosus.

PAPYRUS :—
Pohlii, Nees, in Mart. Fl. Bras. ii. 1. 59 (Marisci sp.). —Bras.
radiatus, Schrad. ex Nees, l. c. 57 = C. comosus.
siculus, Parl. in Hook. Kew Journ. iii. (1851) 189 = C. Papyrus.
spectabilis, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 58 = C. giganteus.
tuberiferus, Schrad. Ind. Sem. Hort. Gotting. (1835). —Cf. Steud. Nom. ed. II. ii. 266 = C. digitatus.
venustus, Nees, in Linnaea, x. (1836) 138 = C. exaltatus.

PAQUERINA, Cass. in Dict. Sc. Nat. xxxvii. 492 (1825) = **Brachycome**, Cass. (Compos.).
graminea, Cass. l. c. 464 = B. graminea.

PARABAENA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 39. *MENISPERMACEAE*, Benth. & Hook. f. i. 34.
ferruginea, Miers, l. c. = sagittata.
heterophylla, Miers, l. c. = sagittata.
oleracea, Miers, l. c. = sagittata.
sagittata, *Miers, l. c.*—Reg. Himal.; Burma.
tuberculata, *Becc. Malesia*, i. (1877) 137.—Malaya; N. Guin.

PARABESLERIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 126 (Gesn. Centr. Am. 52. t. 6) = **Besleria**, Linn. (Gesnerac.).
costaricensis, Oerst. l. c. 127 = B. costaricensis.
triflora, Oerst. l. c. = B. triflora.

PARABOTRYS, C. Muell. in Walp. Ann. vii. 59 (1868) sphalm. = PARARTABOTRYS, Miq. = **Xylopia**, Linn. (Anonac.).

PARACARYUM, Boiss. Diagn. Ser. I. xi. (1849) 128. *BORAGINEAE*, Benth. & Hook. f. ii. 849.
anchusoides, Benth. & Hook. f. Gen. ii. 850 = heliocarpum.
ancyritanum, *Boiss. Fl. Orient.* iv. 260.—Anatolia.
angustifolium, *Boiss. Diagn.* Ser. I. xi. 131.—Graecia; As. Min.
asperum, *Stocks, in Hook. Kew Journ.* iv. (1852) 175. —Afghan.
Aucheri, *Boiss. Diagn.* Ser I. xi. 131.—Caria.
azureum, Boiss. & Heldr. Diagn. Ser. I. xi. 131 = angustifolium.
caelestinum, *Benth. & Hook. f. Gen.* ii. 850.— Ind. or.
calathicarpum, *Boiss. Fl. Orient.* iv. 254.—Beluchist.
calycinum, *Boiss. & Bal. Diagn.* Ser. II. vi. 125.— Cappadoc.
cappadocicum, *Boiss. & Bal. l. c.* 124.—Cappadoc.
Capusii, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 218.—Turkest.
cariense, Boiss. Diagn. Ser. I. xi. 130 = myosotoides.
corymbiforme, *Boiss. l. c.* 131.—As. Min.
corymbiforme, Tchihat. Asie Min. Bot. ii. 115. t. 21 = angustifolium.
cristatum, *Boiss. Diagn.* Ser. I. xi. 131.—Armenia.
denticulatum, Boiss. l. c. Ser. II. iii. 140 = cristatum.
erigerifolium, Schott, ex Kotschy, in Boiss. l. c. = angustifolium.
erysimifolium, *Boiss. Fl. Orient.* iv. 259.—Armenia.
glastifolium, *Boiss. Diagn.* Ser. I. xi. 131.—Armenia.
glochidiatum, *Benth. & Hook. f. Gen.* ii. 850.—Reg. Himal.
heliocarpum, *Kern. in Ber. Naturw. Ver. Innsbruck*, i. (1870) 105.—Reg. Himal.
himalayense, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 162.—Reg. Himal.
hirsutum, *Boiss. Diagn.* Ser. I. xi. 130.—As. Min.
incanum, *Boiss. Fl. Orient.* iv. 262.—Reg. Caucas.
Lambertianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 161.—Ind. or.
lamprocarpum, *Boiss. Diagn.* Ser. I. xi. 131.—Syria.
laxiflorum, *Trautv. in Act. Hort. Petrop.* iii. 11. (1875) 274.—Armenia.
leptophyllum, *Boiss. Diagn.* Ser. I. xi. 131.—Armenia.
longiflorum, Boiss. l. c. 132 = Lindelofia spectabilis.
longipes, *Boiss. l. c.* Ser. II. vi. 126.—Cappadoc.
malabaricum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 160.—Ind. or.
micranthum, *Boiss. Diagn.* Ser. I. xi. 129.—Aegypt.; Persia; Turkest.
microcarpum, *Boiss. l. c.* II. iii. 139.—Reg. Himal.; Afghan.

PARACARYUM:—
modestum, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* ii. 5.—Persia.
myosotoides, *Boiss. Diagn.* Ser. I. xi. 130.—Graec.; As. Min.; Syria.
obtusifolium, *Boiss. & Huet, in Tchihat. Asie Min. Bot.* ii. 115, nomen.—As. Min.
persicum, Boiss. Diagn. Ser. I. xi. 129 = rugulosum.
ponticum, *Boiss. Pl. Or. Nov. Dec.* ii. 9.—As. Min.
Reuteri, *Boiss. & Haussk. ex Boiss. l. c.* 6.—Cataonia.
rubriflorum, Stocks, in Hook. Kew Journ. iv. (1852) 175 = rugulosum.
rugulosum, *Boiss. Diagn.* Ser. I. xi. 129.—Persia; Afghan.
salsum, *Boiss. Fl. Orient.* iv. 255.—Persia.
stenolophum, *Boiss. Diagn.* Ser. II. vi. 124.—Cappadoc.
strictum, Boiss. Fl. Orient. iv. 256 = hirsutum.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 161.—Reg. Himal.
tibeticum, *C. B. Clarke, l. c.* 162.—Reg. Himal.
undulatum, *Boiss. Diagn.* Ser. I. xi. 129.—Persia.

PARACELASTRUS, Miq. Fl. Ind. Bat. i. 11. (1859) 590 = **Microtropis**, Wall. (Celastr.).
bivalvis, Miq. l. c. = M. bivalvis.

PARACELSEA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 18 = **Exacum**, Linn. (Gentian.).
amoena, Zoll. & Mor. l. c.—Java.
fruticosa, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 172.—Ins. Sunda.

PARACELSIA, Hassk. in Flora, xxx. (1847) 664 = praec.

PARACELSIA, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 314, 316 (1857) = **Mollinedia**, Ruiz & Pav. (Monim.).
arguta, Mart. l. c. 314 = M. brasiliensis.
obovata, Mart. l. c. 316 = M. cinerea.

PARACEPHAËLIS, Baill. Adansonia, xii. (1879) 316. *RUBIACEAE*.
tiliacea, *Baill. l. c.*—Madag.

PARACROTON, Miq. Fl. Ind. Bat. i. 11. 382 (1859). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 299.
pendulus, *Miq. l. c.*—Java.

PARACTAENIUM, Benth. & Hook. f. Gen. iii. 1102 (1883), sphalm. = seq.

PARACTAENUM, Beauv. Agrost. 47. t. 10 (1812) = **Panicum**, Linn. (Gramin.).
Novae-Hollandiae, Beauv. l. c. = Panic. gracile.

PARADENOCLINE, Muell. Arg. in DC. Prod. xv. 11. 1141 (1866) = **Adenocline**, Turcz. (Euphorb.).
procumbens, Muell. Arg. l. c.—Afr. austr.

PARADIGMA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 30. t. 8 = **Cordia**, Linn. (Borag.).
Galeottiana, Miers, l. c. 31 = C. Galeottiana.

PARADISANTHUS, Reichb. f. in Bot. Zeit. x. (1852) 930. *ORCHIDEARUM* gen. dub.—Cf. Benth. & Hook. f. iii. 477.
bahiensis, *Reichb. f. l. c.* 93.—Bras.
Moseni, *Reichb. f. in Gard. Chron.* (1881) 1. 298.— Bras.
paranaënsis, *Rodrig. Orch. Nov.* ii. 215.—Bras.
paulensis, *Rodrig. l. c.*—Bras.

PARADISEA, Mazzuc. Viag. Alp. Giul. 27 (1814). *LILIACEAE*, Benth. & Hook. f. iii. 783.
ALLOBROGIA, Tratt. Fl. Kaiserth. ii. 91. n. 184 (1820).
CZACKIA, Andrz. Diss. (1818).
HYPEROGYNE, Salisb. Gen. Pl. Fragm. 81 (1866).
LILIASTRUM, [Tourn.] Link, Handb. i. 173 (1829).
PLEISOLIRION, Rafin. Fl. Tellur. ii. 28 (1836).
hemeroanthericoides, Mazzuc. Viag. Alp. Giul. 27 = P. Liliastrum.
Liliastrum, *Bertol. Fl. Ital.* iv. 133.—Europ.

PARADRYMONIA, Hanst. in Linnaea, xxvi. (1853–55) 207 = **Episcia**, Mart. (Gesnerac.).
glabra, Hanst. l. c. = E. glabra.

PARAGENIPA Baill. in Bull. Soc. Linn. Par. ii. (1879) 207. *RUBIACEAE*.
cervorum, *Baill. l. c.*—Afr. trop.

PARAGNATHIS, Spreng. Syst. iii. 695 (1826) = **Diplomeris**, D. Don (Orchid.).
pulchella, Spreng. l. c. = D. pulchella.

PARAGONIA, Bur. in Bull. Soc. Bot. Fr. xix. (1872) 17 = **Bignonia**, Linn.

PARALEA, Aubl. Pl. Gui. i. 576. t. 231 (1775) = **Diospyros**, Linn. (Ebenac.).
guianensis, Aubl. l. c. = D. Paralea.

PARALLOSA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 359 = **Vicia**, Linn. (Legumin.).
dianthos, Boiss. Fl. Orient. ii. 593, in syn. = V. smyrnaea.
monanthos, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 359 = V. smyrnaea.

PARAMERIA, Benth. in Benth. & Hook. f. Gen. ii. 715 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
glandulifera, *Benth. l. c.*—Malaya; Cambodia.
glandulifera, Vidal, Sin. Pl. Leñ. t. 47. f. D = philippinensis.
pedunculosa, *Benth. in Benth. & Hook. f. Gen.* ii. 715. —Malaya.
philippinensis, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* xiv. [1884] 518.—Ins. Philipp.
polyneura, *Hook. f. Fl. Brit. Ind.* iii. 660.—Malaya.
vulneraria, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* xiv. [1884] 519.—Ins. Philipp.

PARAMESUS, Presl, Symb. Bot. i. 45 (1830) = **Trifolium**, Tourn. (Legumin.).
laevigatus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. strictum.
strictus, Presl, Symb. Bot. i. 46 = T. strictum.

PARAMIGNYA, Wight, Illustr. i. 108. t. 42 (1838). *RUTACEAE*, Benth. & Hook. f. i. 305.
ARTHROMISCHUS, Thw. Enum. Pl. Zeyl. 47 (1858).
angulata, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 135 = longispina.
armata, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 43.—Zeylan.
Blumei, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 137.—Java.
citrifolia, *Hook. f. Fl. Brit. Ind.* i. 510.—Ind. or.
citrifolia, Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 42 = Griffithii.
grandiflora, *Oliver, l. c.*—Burma; Singapore.
Griffithii, *Hook. f. Fl. Brit. Ind.* i. 510.—Reg. Himal.; Burma; Malaya.
littoralis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.— Java.
longispina, *Hook. f. Fl. Brit. Ind.* i. 511.—Malacca.
micrantha, Kurz, in Andam. Rep. App. B 4.—Cf. Journ. As. Soc. Beng. xliv. (1875) II. 135 = citrifolia.
monophylla, *Wight, Illustr.* i. 109. t. 42.—Ind. or.

PARANEPHELIUM, Miq. Fl. Ind. Bat. Suppl. 509 (1860). *SAPINDACEAE*, Benth. & Hook. f. i. 400.
MILDEA, Miq. Ann. Mus. Bat. Lugd. Bat. iii. 88 (1867).
gibbosum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1866) 254.—Sumatra.
xestophyllum, *Miq. Fl. Ind. Bat. Suppl.* 509.— Sumatra.

PARANEPHELIUS, Poepp. & Endl. Nov. Gen. et Sp. iii. 42. t. 248 (1842) = **Liabum**, Adans. (Compos.).
bullatus, A. Gray, ex Wedd. Chlor. And. i. 214.— Peruv.
ovatifolius, A. Gray, ex Wedd. l. c. = L. ovatum.
ovatus, Wedd. l. c. = L. ovatum.
uniflorus, Poepp. & Endl. Nov. Gen. et Sp. iii. 42, t. 248 = L. uniflorum.

PARANOMUS, Salisb. Parad. Lond. sub t. 66 (1807) = **Nivenia**, R. Br. (1810) (Proteac.).
abrotanifolius, Knight, Prot. 68.—Afr. austr.
adiantifolius, Knight, l. c. 70 = N. spathulata.
argenteus, Knight, l. c. 68.—Afr. austr.

PARANOMUS:—
bracteolaris, Knight, Prot. 68.—Afr. austr.
crithmifolius, Knight, l. c. 69 = Nivenia spicata.
cumuliflorus, Knight, l. c. 68.—N. Lagopus.
flabellifer, Knight, l. c. 70.—Afr. austr.
longicaulis, Knight, l. c.—Afr. austr.
sceptriformis, Knight, l. c. 69 = N. Sceptrum.
sceptriformis, Salisb. ex Spreng. Syst. i. 469 = N. media.

PARAPANAX, Miq. Fl. Ind. Bat. Suppl. 338 (1860) = **Trevesia**, Vis. (Araliac.).
littorale, Miq. l. c. 339.—Sumatra.
sessile, Miq. l. c.—Sumatra.

PARAPETALIFERA, Wendl. Coll. Bot. i. 49 (1805) = **Barosma**, Willd. (Rutac.).
odorata, Wendl. l. c. 50. t. 15 = B. crenata.
serrata, Wendl. l. c. 92. t. 34 = B. serratifolia.

PARAPODIUM, E. Mey. Comm. Pl. Afr. Austr. 221 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 749.
costatum, E. Mey. l. c.—Afr. austr.

PARAPOTTSIA, Miq. Fl. Ind. Bat. ii. 1080 (1856) = **Pottsia**, Hook. & Arn. (Apocyn.).
Hookeriana, Miq. l. c. = P. cantoniensis.
laxiflora, Miq. l. c = P. cantoniensis.

PARAPYROLA, Miq. Ann. Mus. Lugd. Bat. iii. 191 (1867) = **Epigaea**, Linn. (Ericac.).
trichocarpa, Miq. l. c. = E. asiatica.

PARARTABOTRYS, Miq. Fl. Ind. Bat. Suppl. 154 (1860) = **Xylopia**, Linn. (Anonac.).
hexagynus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 48.—Sumatra.
sumatranus, Miq. Fl. Ind. Bat. Suppl. 154, 374 = X. malayana.

PARARTOCARPUS, Baill. Adansonia, xi. (1875) 294. *URTICACEAE*, Benth. & Hook. f. iii. 375.
Beccariana, *Baill. l. c.*—Borneo.

PARASHOREA, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65 = **Shorea**, Roxb. (Dipterocarp.).
longisperma, Kurz, l. c. 66 = S. longisperma.
lucida, Kurz, l. c. = S. lucida.
stellata, Kurz, l. c. = S. stellata.

PARASIA, Rafin. Fl. Tellur. iii. 78 (1836) = **Belmontia**, E. Mey. (Gentian.).
cordata, Rafin. l. c. = B. cordata.

PARASPALATHUS, Presl, Bot. Bemerk. 129 (1844); in Abh. Boehm. Ges. V. iii. (1845) 559 = **Aspalathus**, Linn. (Legumin.).
acicularis, Presl, ll. cc. 130; 560 = A. acicularis.
aemula, Presl, ll. cc. = A. aemula.
albanensis, Presl, ll. cc. = A. frankenioides.
albiflora, Presl, ll. cc. = A. thymifolia.
araneosa, Presl, ll. cc. = A. araneosa.
arenarioides, Presl, ll. cc. = A. thymifolia.
argentea, Presl, ll. cc. 129; 559 = A. argentea.
ascendens, Presl, ll. cc. = A. elongata.
callosa, Presl, ll. cc. 130; 560 = A. callosa.
cancellata, Presl, ll. cc. 130, 133; 560, 563 = A. araneosa.
capitata, Presl, ll. cc. 130; 560 = A. capitata.
carnosa, Presl, ll. cc. = A. carnosa.
Cephalotes, Presl, ll. cc. = A. Cephalotes.
cerrhantha, Presl, ll. cc. = A. spicata.
cinerea, Presl, ll. cc. 129; 559 = A. cytisoides.
crocea, Presl, ll. cc. 130; 560 = A. abietina.
cytisoides, Presl, ll. cc. 129; 559 = A. cytisoides.
elongata, Presl, ll. cc. 129; 559 = A. elongata.
ericifolia, Presl, ll. cc. 130; 560 = A. ericifolia.
ericoides, Presl, ll. cc. = A. ericifolia.
erythrodes, Presl, ll. cc. = A. erythrodes.
fagonioides, Presl, ll. cc. = A. callosa.
filicaulis, Presl, ll. cc. = A. filicaulis.
frankenioides, Presl, ll. cc. = A. frankenioides.
galeata, Presl, ll. cc. = A. galeata.
globosa, Presl, ll. cc. = A. globosa.
heterophylla, Presl, ll. cc. 129; 559 = A. lotoides.
hispida, Presl, ll. cc. 130; 560 = A. thymifolia.
humifusa, Presl, ll. cc. 129; 559 = A. procumbens.
incurva, Presl, ll. cc. 130; 560 = A. incurva.
intermedia, Presl, Bot. Bemerk. 130 (1854); in Abh. Boehm. Ges. V. iii. (1845) 560 = Aspalathus frankenioides.
Jacobaea, Presl, ll. cc. 129; 559 = A. Jacobaea.
leucocephala, Presl, ll. cc. 130; 560 = A. virgata.
lotoides, Presl, ll. cc. = A. lotoides.
melanoides, Presl, ll. cc. = A. nigra.
Meyeriana, Presl, ll. cc. = A. ciliaris.
micrantha, Presl, ll. cc. = A. thymifolia.
microcarpa, Presl, ll. cc. = A. microcarpa.
mollis, Presl, ll. cc. = A. mollis.
multiflora, Presl, ll. cc. = A. thymifolia.
nigra, Presl, ll. cc. = A. nigra.
nigrescens, Presl, ll. cc. = A. nigra.
pilosa, Presl, ll. cc. 129; 559 = A. pilosa.
Pluckenetiana, Presl, ll. cc. = A. rugosa.
procumbens, Presl, ll. cc. = A. lotoides.
psoraloides, Presl, ll. cc. = A. cytisoides.
purpurascens, Presl, ll. cc. = A. purpurea.
Scholliana, Presl, ll. cc. = A. Scholliana.
securifolia, Walp. Ann. i. 212 = A. securifolia.
sericea, Presl, (Bot. Bemerk. 130) in Abh. Boehm. Ges. V. iii. (1845) 560 = A. sericea.
Simsiana, Presl, ll. cc. = A. araneosa.
spatulata, Presl, ll. cc. 129; 559 = A. securifolia.
spicata, Presl, ll. cc. 130; 560 = A. spicata.
stellaris, Presl, ll. cc. 129; 559 = A. stellaris.
stenophylla, Presl, ll. cc. = A. stenophylla.
thymifolia, Presl, ll. cc. 130; 560 = A. thymifolia.
villosa, Presl, ll. cc. 129; 559 = A. villosa.
virgata, Presl, ll. cc. = A. virgata.

PARASPONIA, Miq. Pl. Jungh. 68 (1851). *URTICACEAE*, Benth. & Hook. f. iii. 355.
Andersonii, *Planch. in DC. Prod.* xvii. 195.—Ins. Fiji.
aspera, Blume, Mus. Bot. Lugd. Bat. ii. 66 = parviflora.
eurhyncha, *Miq. Fl. Ind. Bat. Suppl.* 411.—Ins. Banca.
parviflora, *Miq. Pl. Jungh.* 69.—Malaya.
rugosa, Blume, Mus. Bot. Lugd. Bat. ii. 66 = parviflora.
similis, Blume, l. c. = parviflora.

PARASTEMON, A. DC. in Ann. Sc. Nat. Sér. II. xviii. (1842) 208. *ROSACEAE*, Benth. & Hook. f. i. 607.
urophyllus, *A. DC. l. c.*—Burma; Malaya.

PARASTRANTHUS, G. Don, Gen. Syst. iii. 716 (1834) = **Lobelia**, Linn. (Campan.).
capitatus, A. DC. in DC. Prod. vii. 354 = L. lutea.
ericoides, A. DC. l. c. = L. lutea.
euphrasioides, A. DC. l. c. 355 = L. lutea.
flavus, A. DC. l. c. 785 = L. flava.
luteus, A. DC. l. c. 354 = L. lutea.
simplex, G. Don, Gen. Syst. iii. 716 = L. lutea.
stellarioides, Vatke, in Linnaea, xxxviii. (1874) 717 = L. stellarioides.
thermalis, A. DC. in DC. Prod. vii. 354 = L. leptocarpa.
unidentatus, G. Don, Gen. Syst. iii. 716 = L. tenella.
variifolius, G. Don, l. c. = L. variifolia.

PARASTREPHIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 449. *COMPOSITAE*, Benth. & Hook. f. ii. 286.
ericoides, *Nutt. l. c.* 450.—Peruv.

PARATHERIA, Griseb. Cat. Pl. Cub. 236 (1866). *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
prostrata, *Griseb. l. c.*—Ind. occ.

PARATHESIS, Hook. f. in Benth. & Hook. f. Gen ii. 645 (1876). *MYRSINEAE*, Benth. & Hook. f. l. c.
adenanthera, *Hook. f. l. c.*—Peruv.
corymbosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 291.—Mexic.
crenulata, *Hook. f. ex Hemsl. l. c.*—Mexic.
melanosticta, *Hemsl. l. c.*—Mexic.
trichogyne, *Hemsl. l. c.*—Nicaragua.

PARATROPHIS, Blume, Mus. Bot. Lugd. Bat. ii. 81. 1852 [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 364.
UROMORUS, Bur. in DC. Prod. xvii. 236 (1873).
heterophylla, *Blume, Mus. Bot. Lugd. Bat.* ii. 81.—N. Zel.
philippinensis, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 198.—Ins. Philipp.

PARATROPIA, DC. Prod. iv. 265 (1830) = **Heptapleurum**, Gaertn. (Araliac.).
acutissima, Miq. Ann. Mus. Bot. Lugd. Bat. i. 20 = H. acutissimum.
aesculifolia, C. Koch, in Wochenschr. ii. (1859) 365 = Brassaiopsis aculeata.
apiculata, Miq. Ann. Mus. Bot. Lugd. Bat. i. 219 = H. apiculatum.
aromatica, Miq. l. c. 760 (= *Agalma aromaticum*).—Java.
assamicum, C. Koch, in Wochenschr. ii. (1859) 365 = Sciadophyllum assamicum.
avenis, Miq. Ann. Mus. Bot. Lugd. Bat. i. 19 = H. avene.
brachybotrya, Miq. Fl. Ind. Bat. i. 1. 755 = H. scandens.
calophylla, Miq. l. c. 758 = H. heterophyllum.
cantoniensis, Hook. & Arn. Bot. Beech. Voy. 189 = H. octophyllum.
capitata, Wight & Arn. Prod. 378 = Brassaia capitata.
confine, Miq. in Bonplandia, iv. (1856) 138; et Fl. Ind. Bat. i. 1. 757 = H. confine.
Corona-sylvae, Miq. l. c. 755 = H. Coronae-sylvae.
crassa, Blanco, Fl. Filip. ed. II. 158 = H. venulosum.
Cumingiana, Presl, Epim. Bot. 250 = Panax Cumingianum.
digitata, Voigt, Hort. Suburb. Calc. 25 = H. venulosum.
divaricata, Miq. Fl. Ind. Bat. i. 1. 754 = H. confine.
elata, C. Koch, in Wochenschr. ii. (1859)365; Hook. f. in Journ. Linn. Soc. vii. (1864) 196 = H. elatum.
elliptica, Miq. Fl. Ind. Bat. i. 1. 755 = H. venulosum.
emarginata, Regel, Gartenfl. (1865) 305 = H. emarginatum.
eurhyncha, Miq. Ann. Mus. Bot. Lugd. Bat. i. 21 = H. eurhynchum.
farinifera, Hort. Linden, ex C. Koch, Wochenschr. ii. (1859) 365 = H. tomentosum.
farinosa, Teijsm. & Binn. Cat. Hort. Bogor. 167; et in Tijdsch. Nederl. Indie, xxix. (1867) 251.—Ins. Molucc.
fastigiata, Miq. Ann. Mus. Bot. Lugd. Bat. i. 24 = H. fastigiatum.
floribunda, C. Koch, Wochenschr. ii. (1859) 365 = Brassaiopsis speciosa.
gracilis, Miq. Ann. Mus. Bot. Lugd. Bat. i. 22 = H. gracile.
heterophylla, Presl, Epim. Bot. 250 = H. heterophyllum.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 761 (= *Agalma Horsfieldii*).—Java.
Junghuhniana, Miq. l. c. 758 = H. Junghuhniana.
latifolia, C. Koch, in Wochenschr. ii. (1859) 365 = Polyscias pinnata.
littoralis, Teijsm. & Binn. Cat. Hort. Bogor. 167 (= *Parapanax littorale*).—Sumatra.
longifolia, DC. Prod. iv. 266 = H. longifolium.
lucescens, Miq. Fl. Ind. Bat. i. 1. 754.—Java.
lucida, Miq. l. c. = H. rigidum.
lutescens, C. Koch, in Wochenschr. ii. (1859) 365 = *lucescens*.
macrantha, Miq. Fl. Ind. Bat. i. 1. 756 = H. venulosum.
macrostachya, Miq. l. c. 760.—N. Guin.
Mannii, Hook. f. in Journ. Linn. Soc. vi. (1862) 10 = H. Mannii.
micrantha, Miq. Fl. Ind. Bat. Suppl. 337 = H. micranthum.
multijuga, A. Gray, Bot. U. St. Expl. Exped. i. 722.—Ins. Fiji.
nodosa, DC. Prod. iv. 265 = Polyscias nodosa.
obovata, C. Koch, in Wochenschr. ii. (1859) 365 = H. stellatum.
obtusa, Blanco, Fl. Filip. ed. II. 159 = H. venulosum.
oxyphylla, Miq. Fl. Ind. Bat. Suppl. 338 = H. oxyphyllum.
parasitica, Miq. l. c. i. 1. 757 = H. parasiticum.
pergamacea, DC. Prod. iv. 266 = H. pergamaceum.
petiolosa, Miq. Ann. Mus. Bot. Lugd. Bat. i. 24 = H. petiolosum.
polita, Miq. l. c. 22 = H. politum.
polybotrya, Miq. Fl. Ind. Bat. i. 1. 755 = H. polybotryum.
polydactylis, Montr. in Mém. Acad. Lyon, x. (1860) 212.—N. Caled.
polyphylla, Miq. Fl. Ind. Bat. i. 1. 760 = H. polyphyllum.
pubigera, Brongn. ex C. Koch, in Wochenschr. ii. (1859) 365 = Sciadophyllum assamicum.
pulchra, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 106 = Sciadophyllum pulchrum.

PARATROPIA:—
rigida, DC. Prod. iv. 266 = Heptapleurum rigidum.
rostrata, Wight, Ic. t. 1013–14 = H. rostratum.
rotundifolia, Tenore, Cat. Orto Nap. (1845) 89 = H. venulosum ?
samoënsis, A. Gray, Bot. U. St. Expl. Exped. i. 722. —Ins. Samoa.
scandens, Miq. Fl. Ind. Bat. i. 1. 757 = H. scandens.
serrata, Miq. l. c. = H. serratum.
singalensis, Miq. Ann. Mus. Bot. Lugd. Bat. i. 23 = H. singalangense.
subcordata, C. Koch, Wochenschr. ii. (1859) 365 = Pentapanax subcordatum.
subulata, Miq. Ann. Mus. Bot. Lugd. Bat. i. 22 = H. subulatum.
Teijsmaniana, Hort. ex C. Koch, in Wochenschr. ii. (1859) 365 = H. tomentosum.
terebinthinacea, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 338 = H. stellatum.
tomentosa, Miq. Fl. Ind. Bat. i. 1. 753 = H. tomentosum.
trifoliata, C. Koch, in Wochenschr. ii. (1859) 365 = Pentapanax Leschenaultii.
umbraculifera, Wall. ex Voigt, Hort. Suburb. Calc. 25 = Polyscias nodosa.
undulata, C. Koch, Wochenschr. ii. (1859) 365 = Macropanax undulatum.
venulosa, Wight & Arn. Prod. 377 = H. venulosum.
verticillata, C. Koch, Wochenschr. ii. (1859) 365 = H. verticillatum.
Wallichiana, C. Koch, l. c. = H. tomentosum.
Wallichiana, Wight & Arn. Prod. 377 = H. Wallichianum.

PARAVINIA, Hassk. in Flora, xxxi. (1848) 580 = **Praravinia**, Korth. (Rubiac.).

PARDANTHUS, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 246 = **Belamcanda**, Adans. (Irid.).
bulbiferus, Klatt, in Linnaea, xxxi. (1861–62) 561 = Sparaxis bulbifera.
chinensis, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 247 = B. punctata.
dichotomus, Ledeb. Fl. Ross. iv. 106 = Iris dichotoma.
nepalensis, Sweet, Hort. Brit. ed. I. 395 = B. punctata.
sinensis, Van Houtte, in Fl. des Serres, xxvi. (1865–67) t. 1632 = B. punctata.

PARDINIA, Herb. in Lindl. Bot. Reg. (1844) Misc. 66, in obs. = **Hydrotaenia**, Lindl. et **Tigridia**, Juss. (Irid.).

PARDISIUM, Burm. f. Fl. Cap. Prod. 26 (1768) = **Gerbera**, Gronov. (Compos.).
capense, Burm. f. l. c.—Afr. austr.

PARDUYNA, Salisb. Gen. Pl. Fragm. 58 (1866) = **Schelhammera**, R. Br. (Liliac.).
multiflora, Salisb. l. c. = S. multiflora.

PARECHITES, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1859) 193 = **Trachelospermum**, Lem. (Apocyn.).
adnascens, Hance, in Journ. Bot. vi. (1868) 299 = T. jasminoides.
borneana, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 193.—Borneo.
Bowringii, Hance, in Journ. Bot. vi. (1868) 299.—China.
Thunbergii, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 403 = T. jasminoides.

PARENTUCELLIA, Viv. Fl. Lib. Spec. 31. t. 21. f. 3 (1824) = **Bartsia**, Linn. (Scrophul.).
floribunda, Viv. l. c. 32 = B. latifolia.

PARFONSIA, Scop. Introd. 210 (1777), err. typ. = PARSONSIA, P. Br. = **Cuphea**, P. Br. (Lythrar.).

PARIANA, Aubl. Pl. Gui. ii. 876. t. 337 (1775).
GRAMINEAE, Benth. & Hook. f. iii. 1112.
APHONINA, Neck. Elem. iii. 237 (1790).
EREMITIS, Doell, in Mart. Fl. Bras. ii. II. 338. t. 48 (1877).
angustifolia, *Spreng. Syst.* ii. 609.—Venezuela.
campestris, *Aubl. Pl. Gui.* ii. 877. t. 337.—Guian.; Bras.
glauca, *Nees, Agrost. Bras.* 294.—Bras.
gracilis, *Doell, in Mart. Fl. Bras.* ii. II. 337.—Am. trop.
imberbis, *Nees, Agrost. Bras.* 297.—Bras.

PARIANA:—
inaequalis, *Miq. in Linnaea*, xix. (1847) 127.—Guian.
intermedia, *Doell, in Mart. Fl. Bras.* ii. II. 337.—Am. trop.
lanceolata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 107.—Bras.
lunata, *Nees, Agrost. Bras.* 295.—Bras.
mollis, *Nees, l. c.* 296.—Bras.
parviflora, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) 105.—Bras.
radiciflora, *Sagot, ex Doell, in Mart. Fl. Bras.* ii. II. 336.—Guian.
scabra, *Nees, Agrost. Bras.* 294.—Bras.
sylvestris, *Nees, l. c.* 295.—Bras.
zinziberina, *Doell, in Mart. Fl. Bras.* ii. II. 337.—Am. trop.

PARIATICU, Adans. Fam. ii. 223 (1763) = **Nyctanthes**, Linn. (Oleac.).

PARIETARIA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 317 (1737). *URTICACEAE*, Benth. & Hook. f. iii. 392.
FREIREA, Gaudich. in Freyc. Voy. Bot. 502 (1826).
THAUMURIA, Gaudich. l. c. (1826).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 258 = Pouzolzia abyssinica.
affinis, Hornem. Hort. Hafn. (1850); ex Blume, Mus. Bot. Lugd. Bat. ii. 250 = cretica.
alsinefolia, *Delile, Fl. Egypt.* 281. t. 50. f. 2.—Aegypt.; Oriens.
angustifolia, *Boj. Hort. Maurit.* 295.—Hab. ?
appendiculata, Webb & Berth. Phyt. Canar. iii. 265 = debilis.
arborea, L'Hérit. in Journ. Phys. xxxiii. (1788) 55 = Gesnouinia arborea.
aspera, Leschen. ex Gaudich. in Freyc. Voy. Bot. 510 = Fatoua pilosa.
assurgens, Poir. Encyc. v. 15 = officinalis.
australis, Blume, Mus. Bot. Lugd. Bat. ii. 256 = debilis.
boehmerioides, *Mart. ex Miq. in Mart. Fl. Bras.* iv. I. 194.—Bras.
bracteata, Moench, Meth. Suppl. 115 = cretica.
bracteata, Wight, ex Wall. Cat. n. 4600 = Pouzolzia caudata.
canescens, Blume, Mus. Bot. Lugd. Bat. ii. 249 = officinalis.
canescens, Poir. Encyc. v. 17 = Pouzolzia laevigata.
capensis, Thunb. Prod. Pl. Cap. 31 = Boehmeria capensis.
carnosula, Blume, Mus. Bot. Lugd. Bat. ii. 255 = debilis.
circaeafolia, Pourr. ex Wedd. in DC. Prod. xvi. I. 235^{15} = mauritanica.
cochinchinensis, Lour. Fl. Cochinch. 654 = Pouzolzia indica.
cretica, *Linn. Sp. Pl.* 1052.—Graecia; Ins. Creta.
cretica, Loisel. Fl. Gall. 693 = lusitanica.
cretica, Moris, Stirp. Sard. Elench. fasc. i. 41 = Helxine Soleirolii.
cretica, Willd. Sp. Pl. iv. 956 = judaica.
cryptorum, C. Koch, in Linnaea, xxii. (1849) 606 = judaica.
cuneata, Eckl. & Zeyh. ex Krauss, in Flora, xxviii. (1845) 88 = Droguetia ambigua.
cuneata, E. Mey. ex Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 212 = Australina acuminata.
debilis, *Forst. f. Prod.* 73.—Reg. trop. et subtrop.
Decaisnei, Steud. Nom. ed. II. ii. 267 = Pouzolzia indica.
diffusa, Mert. & Koch, Deutschl. Fl. i. 827 = judaica.
diffusa, Schur, Enum. Pl. Transs. 602 = officinalis.
elliptica, C. Koch, in Linnaea, xxii. (1849) 606 = judaica.
erecta, Mert. & Koch, Deutschl. Fl. i. 825 = officinalis.
Feliciana, *Phil. in Bot. Zeit.* xxviii. (1870) 501.—Chili.
filamentosa, Webb & Berth. Phyt. Canar. iii. 263 = Gesnouinia filamentosa.
filiformis, Tenore, Syn. Fl. Neap. 77 = lusitanica.
floridana, Nutt. Gen. Am. ii. 208 = pensylvanica.
gracilis, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 16 = debilis.
guineensis, G. Don, ex Hook. Niger Fl. 518 = Pouzolzia guineensis.
Huetia, Comm. ex Wedd. in DC. Prod. xvi. I. 132 = Pilea verbascifolia.

PARIETARIA:—
humifusa, Blume, Mus. Bot. Lugd. Bat. ii. 252 = debilis.
indica, Linn. Mant. i. 128 = Pouzolzia indica.
indica, Portensch. ex Blume, Mus. Bot. Lugd. Bat. ii. 232 = Pouzolzia auriculata.
judaica, *Linn. Sp. Pl.* ed. II. 1492.—Graecia; Oriens; Tibet.
judaica, Pers. Syn. ii. 555 = lusitanica.
judaica, Vill. Hist. Pl. Dauph. ii. 346 = officinalis.
judaica, Welw. ex Blume, Mus. Bot. Lugd. Bat. ii. 252 = mauritanica.
lanceolata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 17 = Australina capensis.
levigata, Poir. Encyc. v. 17 = Pouzolzia laevigata.
linguaefolia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 313 = pensylvanica.
lusitanica, Janka, in Oester. Bot. Zeitschr. viii. (1858) 200 = officinalis.
lusitanica, *Linn. Sp. Pl.* 1052.—Reg. Mediterr.
lusitanica, Sieber, ex Blume, Mus. Bot. Lugd. Bat. ii. 250 = cretica.
lusitanica, Viv. Fl. Cors. Prod. App. 7 = Helxine Soleirolii.
maderensis, Reichb. in Flora, xiii. (1830) 131 = officinalis *vel* debilis.
mauritanica, *Dur. in Duch. Rev. Bot.* ii. (1847) 427.—Reg. Mediterr. occ.
micrantha, Ledeb. Ic. Pl. Fl. Ross. i. 7. t. 22 = debilis.
microphylla, Linn. Syst. ed. X. 1308 = Pilea microphylla.
multicaulis, Boiss. & Heldr. Diagn. Ser. I. xii. 106 = judaica.
muralis, Salisb. Prod. 69 = officinalis.
nitens, C. Koch, in Linnaea, xxii. (1849) 607 = judaica.
nitida, *Miq. in Mart. Fl. Bras.* iv. I. 204.—Bras.
ocimifolia, Poir. Encyc. v. 15 = mauritanica.
officinalis, *Linn. Sp. Pl.* 1052.—Europ. austr.; Oriens.
oppositifolia, F. Muell. Fragm. iv. 88 = Pouzolzia indica.
Orizabae, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 313 = pensylvanica.
parviflora, Friedrich, ex Blume, Mus. Bot. Lugd. Bat. ii. 249 = lusitanica.
pensylvanica, *Muhl. ex Willd. Sp. Pl.* iv. 955.—Am. bor.
pilosa, *Willd. l. c.* 954.—Afr. austr.
platyphyllos, Link, ex Steud. Nom. ed. II. ii. 267 (Quid ?).—Hab. ?
polygonoides, *Willd. Sp. Pl.* iv. 956.—Armenia.
populifolia, *Nym. in Linnaea*, xviii. (1844) 661.—Ins. Melita.
prostrata, Hort. Berol. ex Spreng. Nov. Prov. 30 = Pilea serpyllifolia.
punctata, Willd. Sp. Pl. iv. 953 = judaica.
ramiflora, Moench, Meth. 327 = judaica.
reclinata, *Moon, Cat. Pl. Ceyl.* 72.—Zeylan.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 (= Pouzolziae sp. ?).—Java.
repens, Soleir. ex Mutel, Fl. Fr. iii. 172 = Helxine Soleirolii.
rubicunda, *Pohl, ex Miq. in Mart. Fl. Bras.* iv. I. 194. —Bras.
serbica, Panč. ex Kanitz, in Just, Bot. Jahresb. iii. 1875 (1877), 701 = lusitanica.
serpyllifolia, Poir. Encyc. v. 16 = Pilea serpyllifolia.
Soleirolii, Spreng. Syst. iv. Cur. Post. 318 = Helxine Soleirolii.
Sonnerati, Poir. Encyc. v. 15 = Phenax vulgaris.
squalida, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 285 = debilis.
tibethana, Blume, Mus. Bot. Lugd. Bat. ii. 251 = judaica.
urticaefolia, Linn. f. Suppl. 434 = Pilea rupipendia.
urticaefolia, E. Mey. ex Wedd. in DC. Prod. xvi. I. 235^{58} = Droguetia ambigua.
velutina, Blume, Mus. Bot. Lugd. Bat. ii. 251 = judaica.
verbascifolia, Poir. Encyc. v. 16 = Pilea verbascifolia.
Welwitschii, Blume, Mus. Bot. Lugd. Bat. ii. 251 = mauritanica.
zeylanica, Linn. Sp. Pl. 1052 = Pouzolzia indica.

PARILIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Elaeodendron**, Jacq. (Tiliac.).
malabarica, Dennst. l. c. (Quid?).—Ind. or.

PARILIUM, Gaertn. Fruct. i. 234. t. 51. f. 1 (1788) = **Nyctanthes**, Linn. (Oleac.).
Arbor-tristis, Gaertn. l. c.; et ii. 265 = N. Arbor-tristis.

PARILLAX, Rafin. Neogenyt. 8 (1825) = **Smilax**, Tourn. (Liliac.).
laurifolia, Rafin. l. c. = S. laurifolia.
pumila, Rafin. l. c. = S. pumila.

PARILTARIA, Burm. f. Fl. Ind. 221 (1768), err. typ. = **Parietaria**, Tourn. (Urtic.).

PARINARI, Aubl. Pl. Gui. i. 514 (1775) = seq.

PARINARIUM, Aubl. Pl. Gui. i. 514 (Parinari) (1775); Juss. Gen. 342 (1789). *ROSACEAE*, Benth. & Hook. f. i. 607.
BALANTIUM, Desv. in Ham. Prod. Pl. Ind. Occ. 34 (1825).
DUGORTIA, Scop. Introd. 217 (1777).
ENTOSIPHON, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 44.
EXITELIA, Blume, Fl. Jav. Praef. p. vii (1828).
FEROLIA, Aubl. Pl. Gui. Suppl. 7 (1775).
GRYMANIA, Presl, Epim. Bot. 193 (1849).
LEPIDOCARPA, Korth. in Nederl. Kruidk. Arch. iii. (1855) 385.
LEPIDOCARYA, Korth. ex Miq. Fl. Ind. Bat. i. 1. 353 (1855).
MAMPATA, 'Adans.' ex Steud. Nom. ed. II. ii. 98 (1841).
MARANTHES, Blume, Bijdr. 89 (1825).
NEON, Adans. ex Juss. Gen. 342 (1789).
PETROCARYA, Schreb. Gen. 245 (1789).
THELIRA, Thou. Gen. Nov. Madag. 21 (1806).
anamense, *Hance, in Journ. Bot.* xv. (1877) 333.—China.
asperulum, *Miq. Fl. Ind. Bat. Suppl.* 307.—Sumatra.
brachystachyum, Benth. in Hook. Journ. Bot. ii. (1840) 213 = excelsum.
brasiliense, *Hook. f. in Mart. Fl. Bras.* xiv. II. 51.—Bras.
campestre, *Aubl. Pl. Gui.* i. 514.—Guian.
capense, *Harv. in Harv. & Sond. Fl. Cap.* ii. 597.—Afr. austr.
Chapelieri, *Baill. Adansonia*, ix. (1868–70) 148.—Madag.
chrysophyllum, *Oliver, Fl. Trop. Afr.* ii. 369.—Afr. trop.
congense, *Didr. in Kjoeb. Vidensk. Meddel.* (1854) 197.—Afr. trop.
cordatum, *Hook. f. in Mart. Fl. Bras.* xiv. II. 50.—Bras.
coriaceum, *Benth. in Hook. Journ. Bot.* ii. (1840) 213.—Guian.
corymbosum, *Miq. Fl. Ind. Bat.* i. 1. 356.—Java.
costatum, *Blume, Mélang. Bot.* (1855).—*Cf. Flora*, xli. (1858) 255; *et ex Miq. l. c.* 354.—Malaya.
curatellaefolium, *Planch. ex Benth. in Hook. Niger Fl.* 333.—Afr. trop.
dillenifolium, R. Br. in Wall. Cat. n. 7520 = Dipterocarpus cornutus.
excelsum, *Sabine, in Trans. Hort. Soc.* v. (1824) 451.—Afr. trop.
fragile, *Teijsm. & Binn. Cat. Hort. Bog.* 253.—Sumatra.
Gardneri, *Hook. f. in Mart. Fl. Bras.* xiv. II. 50.—Bras.
glaberrimum, *Hassk. in Flora*, xxvii. (1844) 583.—Java.
glabrum, *Oliver, Fl. Trop. Afr.* ii. 370.—Afr. trop.
Glaziovianum, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 72.—Bras.
Griffithianum, *Benth. in Hook. Niger Fl.* 334.—Burma; Malaya.
Helfeii, *Hook. f. Fl. Brit. Ind.* ii. 311.—Burma.
indicum, *Bedd. Ic. Pl. Ind. Or.* 22. t. 109; *Fl. Sylv.* i. t. 191.—Ind. or.
insularum, *A. Gray, Bot. U. St. Expl. Exped.* i. 488. t. 54.—Ins. Samoa.
Jackianum, *Benth. in Hook. Niger Fl.* 335.—Malacca.
lanceolatum, *Teijsm. & Binn. Cat. Hort. Bog.* 253.—Java.

PARINARIUM:—
laurinum, *A. Gray, Bot. U. St. Expl. Exped.* i. 490.—Ins. Fiji.
macrophyllum, *Sabine, in Trans. Hort. Soc.* v. (1824) 452.—Afr. trop.
macrophyllum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 256.—Malaya.
Maranthes, *Blume, Mélang. Bot.* (1855).—*Cf. Flora*, xli. (1858) 256; *et in Mus. Bot. Lugd. Bat.* ii. 99.—Java.
Margarata, *A. Gray, Bot. U. St. Expl. Exped.* i. 489.—Ins. Fiji.
Mobola, *Oliver, Fl. Trop. Afr.* ii. 368.—Afr. trop.
montanum, *Aubl. Pl. Gui.* i. 514.—Guian.
multiflorum, Miq. Fl. Ind. Bat. i. 1. 356 = Griffithianum.
nitidum, *Hook. f. Fl. Brit. Ind.* ii. 310.—Malaya.
nonda, *F. Muell. ex Benth. Fl. Austral.* ii. 426.—Austral.
oblongifolium, *Hook. f. Fl. Brit. Ind.* ii. 309.—Malacca.
obtusifolium, *Hook. f. in Mart. Fl. Bras.* xiv. II. 52.—Bras.
ovale, Blume, ex Miq. Fl. Ind. Bat. i. 1. 353 = costatum.
Pohlii, *Hook. f. in Mart. Fl. Bras.* xiv. II. 51.—Bras.
polyandrum, *Benth. in Hook. Niger Fl.* 333.—Afr. trop.
polyneurum, *Miq. Fl. Ind. Bat. Suppl.* 306.—Sumatra.
robustum, *Oliver, Fl. Trop. Afr.* ii. 370.—Afr. trop.
salicifolium, *Miq. Fl. Ind. Bat.* i. 1. 357.—Ins. Philipp.
scabrum, *Hassk. in Flora*, xxvii. (1844) 585.—Java.
senegalense, Perr. ex DC. Prod. ii. 527 = macrophyllum, *Sabine*.
Sprucei, *Hook. f. in Mart. Fl. Bras.* xiv. II. 53.—Bras.
subcordatum, *Oliver, Fl. Trop. Afr.* ii. 367.—Afr. trop.
sumatranum, Benth. in Hook. Niger Fl. 335 = costatum.
sumatranum, Kurz, For. Fl. Brit. Burma, i. 433 = Helferi.
travancoricum, *Bedd. Ic. Pl. Ind. Or.* i. 43—Ind. or.
Wallichianum, *R. Br. in Wall. Cat.* n. 7520.—Singapore.

PARIPON, Voigt, in Syll. Ratisb. ii. (1828) 51. *PALMAE*.
Palmiri, Voigt, l. c. (Quid?).—Hab.?

PARIS, [Rupp.] Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 833.
ALOPICARPUS, Neck. Elem. ii. 369 (1790).
CARTALINIA, Szov. ex Kunth, Enum. Pl. v. 119 (1850).
DAISWA, Rafin. Fl. Tellur. iv. 18 (1836).
DEMIDOVIA, Hoffm. Hort. Mosq. (1808) f. 2.
EUTHYRA, Salisb. Gen. Pl. Fragm. 61 (1866).
apetala, Hoffm. in Comm. Phys. Med. Mosc. i. (1808) 1. p. 5 = incompleta.
dahurica, *Fisch. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 105.—Dahuria.
Daisua, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 49 = polyphylla.
hexaphylla, *Cham. in Linnaea*, vi. (1831) 568.—Sibir.
incompleta, *Bieb. Fl. Taur. Cauc.* i. 306.—Reg. Caucas.; Armenia.
obovata, *Ledeb. Ic. Pl. Ross.* i. t. 16.—Sibir.
octophylla, Hoffm. Hort. Mosq. (1808) 27 = incompleta.
pentafolia, Renault, Fl. Dép. Orne, 26 = quadrifolia.
polyphylla, *Sm. in Rees, Cyclop.* xxvi. n. 2.—Reg. Himal.; China.
quadrifolia, *Linn. Sp. Pl.* 367.—Europ.; As. bor.
tetraphylla, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 412.—Japon.
trifolia, Renault, Fl. Dép. Orne, 26 = quadrifolia.
verticillata, *Bieb. Fl. Taur. Cauc.* iii. 287.—Reg. Caucas.; Sibir.

PARISHELLA, A. Gray, in Coult. Bot. Gaz. vii. (1882) 94. *CAMPANULACEAE*.
californica, *A. Gray, l. c.*—Am. bor. occ.

PARISHIA, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 169. t. 26. *ANACARDIACEAE*, Benth. & Hook. f. i. 424.
insignis, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 170. t. 26.—Burma.

PARISHIA:—
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 30.—Malacca.
pubescens, *Hook. f. l. c.*—Malacca.
paucijuga, *Engl. in DC. Monog. Phan.* iv. 309.—Malaya.

PARITA, Scop. Introd. 282 (1777) = seq.

PARITI, Adans. Fam. ii. 401 (1763) = seq.

PARITIUM, A. Juss. in A. St. Hil. Fl. Bras. Mer. i. 255 (1827) = **Hibiscus**, Linn. (Malvac.).
abutiloides, G. Don, Gen. Syst. i. 485 = H. abutiloides.
Azanzae, G. Don, l. c. = H. Azanzae.
Bealii, Wall. ex Voigt, Hort. Suburb. Calc. 120, nomen.—Japon.
circinnatum, G. Don, Gen. Syst. i. 485 = H. tiliaceus.
elatum, G. Don, l. c. = H. elatus.
gangeticum, G. Don, l. c. = Thespesia Lampas.
guineense, G. Don, l. c. = H. tiliaceus.
macrophyllum, G. Don, l. c. = H. macrophyllus.
pernambucense, G. Don, l. c. = H. pernambucensis.
purpurascens, Seem. in Bonplandia, ix. (1861) 254.—Ins. Fiji.
quinquelobum, Hook. f. in Hook. Niger Fl. 227 = H. quinquelobus.
simile, G. Don, Gen. Syst. i. 485 = H. elatus.
sterculiaefolium, Guill. & Perr. Fl. Seneg. Tent. i. 60 = H. quinquelobus.
tiliaceum, A. St. Hil. Fl. Bras. Mer. i. 256 = H. tiliaceus.
tricuspe, G. Don, Gen. Syst. i. 485 = H. tricuspis.
virgatum, Guill. & Perr. Fl. Seneg. Tent. i. 59 = H. quinquelobus.
Wrayae, Walp. Rep. i. 311 = H. Huegelii.

PARIVOA, Aubl. Pl. Gui. ii. 756. t. 303 (1775) = **Eperua**, Aubl. (Legumin.).
grandiflora, Aubl. l. c. ii. 757. t. 303 = E. grandiflora.
tomentosa, Aubl. l. c. 759 = Crudia Parivoa.

PARKIA, R. Br. in Denh. & Clapp. Trav. App. 234 (1826). *LEGUMINOSAE*, Benth. & Hook. f. i. 588.
PARYPHOSPHAERA, Karst. Fl. Columb. ii. 7. t. 104 (1862).
africana, *R. Br. in Denh. & Clapp. Trav. App.* 234.—Afr. trop.
auriculata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 363; *et in Mart. Fl. Bras.* xv. II. 267.—Bras.
biglandulosa, *Wight & Arn. Prod.* 279.—Malaya.
biglobosa, Benth. in Hook. Journ. Bot. iv. (1842) 328 = Roxburghii, africana.
Brunonis, R. Grah. in Wall. Cat. n. 5288 = Roxburghii.
discolor, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 363; *et in Mart. Fl. Bras.* xv. II. 267.—Bras.
filicina, *Benth. ex Walp. Rep.* v. 577.—Am. trop.
filicoidea, *Welw. in Oliver, Fl. Trop. Afr.* ii. 324.—Afr. trop.
fraterna, *Vatke, in Linnaea*, xliii. (1880–82) 339.—Madag.
grandis, Hassk. in Flora, xxv. (1842) II. Beibl. 55 = Roxburghii.
Hoffmanni, *Vatke, in Linnaea*, xliii. (1880–82) 338.—Madag.
insignis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 74.—Burma.
intermedia, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 149 = Roxburghii?
intermedia, *Oliver, Fl. Trop. Afr.* ii. 324.—Afr. trop.
leiophylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 73.—Burma.
macrocarpa, *Miq. Fl. Ind. Bat. Suppl.* 284.—Sumatra.
multijuga, *Benth. in Trans. Linn. Soc.* xxx. (1875) 362; *et in Mart. Fl. Bras.* xv. II. 265.—Am. trop.
nitida, *Miq. Stirp. Surin. Sel.* 7.—Guian.
oppositifolia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 363; *et in Mart. Fl. Bras.* xv. II. 267.—Bras.

PARKIA:—
Parrii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 359.—Ins. Fiji.
Paryphosphaera, *Benth. in Trans. Linn. Soc.* xxx. (1875) 363.—Am. trop.
pectinata, *Benth. l. c.* 362; *et in Mart. Fl. Bras.* xv. II. 266.—Am. trop.
pendula, *Benth. ex Walp. Rep.* v. 577.—Bras.
platycephala, *Benth. in Hook. Journ. Bot.* iv. (1842) 329.—Bras.
Roxburghii, *G. Don, Gen. Syst.* ii. 397.—As. trop.
singularis, *Miq. Fl. Ind. Bat.* i. 1. 1078.—Sumatra.
speciosa, Hassk. in Flora, xxv. (1842) II. Beibl. 55 = Roxburghii?
streptocarpa, *Hance, in Journ. Bot.* xiv. (1876) 259.—Siam.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 284.—Sumatra.
uniglobosa, G. Don, Gen. Syst. ii. 396 = africana.

PARKINSONIA, Plum. ex Linn. Gen. ed. I. 342 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 570.
aculeata, *Linn. Sp. Pl.* 375.—Am. trop.
africana, *Sond. in Linnaea*, xxiii. (1850) 38.—Afr. austr.
florida, *S. Wats. in Proc. Am. Acad.* xi. (1876) 135.—Am. bor. occ.
inermis, *Spreng. Syst.* ii. 345.—Reg. Argent.
microphylla, *Torr. in Pacif. Rail. Rep.* iv. 82.—Am. bor. occ.
orientalis, Spreng. Syst. iv. Cur. Post. 170 = Ormocarpum sennoides.
spinosa, H. B. & K. Nov. Gen. et Sp. vi. 335 = aculeata.
texana, *S. Wats. in Proc. Am. Acad.* xi. (1876) 136.—Am. bor. occ.
Torreyana, *S. Wats. l. c.* 135.—Am. bor. occ.

PARLATOREA, Rodrig. Orch. Nov. i. 141 (1877). *ORCHIDEAE.*
discolor, *Rodrig. l. c.*—Bras.

PARLATORIA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 72. *CRUCIFERAE*, Benth. & Hook. f. i. 102.
brachycarpa, Boiss. l. c. 73 = Sobolewskia clavata.
cakiloides, *Boiss. l. c.* 72.—Assyria.
clavata, Boiss. l. c. 73 = Sobolewskia clavata.
Griffithiana, *Boiss. Diagn.* Ser. II. i. 23.—Afghan.
rostrata, *Boiss. & Hohen. l. c.* Ser. I. viii. 22.—Persia.

PARMENTIERA, DC. in Bibl. Univ. Genèv. xvii. (1838) 135 (Rev. Bign. 19). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1051.
ZENKERIA, Reichb. Nom. 236 (1841).
aculeata, Seem. Bot. Voy. Herald, 183 = edulis.
alata, *Miers, in Trans. Linn. Soc.* xxvi. (1868) 166.—Mexic.
cereifera, *Seem. Bot. Voy. Herald*, 182. t. 32.—Panama.
edulis, *DC. Prod.* ix. 244.—Mexic.
foliolosa, Miers, in Trans. Linn. Soc. xxvi. (1870) 166 = edulis.
lanceolata, Miers, l. c. 167 = edulis.

PARNASSIA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 87 (1737). *SAXIFRAGEAE*, Benth. & Hook. f. i. 639.
affinis, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 81 = ovata.
alpina, Dalla Torre, Atl. Alpenfl. 71 = palustris.
americana, Muhl. Cat. 33, nomen = caroliniana.
asarifolia, *Vent. Jard. Malm.* 39. t. 39.—Am. bor.
cabulica, *Planch. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 403.—Afghan.
caroliniana, *Michx. Fl. Bor. Am.* i. 184.—Am. bor.
ciliata, Gilib. Fl. Lituan. ii. 139 = palustris.
europaea, Pers. Syn. i. 331 = palustris.
fimbriata, *Banks, in Kon. & Sims, Ann. Bot.* i. (1805) 391.—Am. bor. occ.
foliosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 79.—Reg. Himal.
glauca, Rafin. Autikon Bot. 42 = caroliniana.
grandiflora, Rafin. l. c. = caroliniana.
grandifolia, DC. Prod. i. 320 = caroliniana.
Kotzebuei, *Cham. ex Spreng. Syst.* i. 951.—Am. bor. occ.
Laxmanni, Pall. ex Schult. Syst. vi. 696 = ovata.
maisorensis, Henders. & Hume, Lahore, 320; et ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 403 = ovata.
mucronata, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 169 = palustris.
mysorensis, *Heyne, ex Wall. Cat.* n. 3754; *et Wight & Arn. Prod.* 35.—Ind. or.
nana, Griff. Notul. iv. 427; Ic. Pl. Asiat. t. 570. f. 2 = Wightiana.
nubicola, *Wall. Cat.* n. 1246.—Reg. Himal.
nudata, Rafin. Autikon Bot. 41 = asarifolia.
nummularia, Maxim. in Bull. Acad. Pétersb. xi. (1867) 430 = foliosa.
obtusiflora, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 23. t. 3.—Ross. arct.
oreophila, *Hance, in Journ. Bot.* xvi. (1878) 106.—China.
ornata, Wall. Cat. n. 1247 = Wightiana.
ovata, *Ledeb. in Mém. Acad. Pétersb.* v. (1815) 528.—Reg. Himal.; Sibir.
ovata, Muhl. Cat. ed. II. 32, nomen. = caroliniana.
palustris, *Linn. Sp. Pl.* 273.—Europ.; As. et Am. bor.
parviflora, *DC. Prod.* i. 320.—Am. bor.
polynectaria, Forsk. Fl. Aegypt. Arab. 207 = Swertia decumbens.
pusilla, *Wall. Cat.* n. 1245.—Reg. Himal.
Schmidii, Zenk. Pl. Ind. Dec. i. 3. t. 5 = Wightiana.
speciosa, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 653 = caroliniana.
subacaulis, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 164 = ovata.
tenella, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 80.—Reg. Himal.; Soongar.
tenuis, DC. Prod. i. 320, in syn. = palustris.
trinervis, Drude, in Linnaea, xxxix. (1875) 322 = ovata.
Turczaninowii, Ledeb. Fl. Ross. i. 263 = ovata.
vulgaris, Dum.-Cours. Bot. Cult. ed. II. iv. 546 = palustris.
Wightiana, *Wall. Cat.* n. 3755; *et ex Wight & Arn. Prod.* 35.—Ind. or.; Reg. Himal.

PAROCHETUS, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 240 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 485.
COSMIUSA, Alef. in Bot. Zeit. xxiv. (1866) 146.
communis, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 240.—Ind. or.; Malaya; Afr. trop.
maculata, Benn. Pl. Jav. Rar. i. 162 = communis.
major, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 241 = communis.
oxalidifolius, Royle, Illustr. Bot. Himal. 201. t. 35. f. 1 = communis.

PAROLINIA, Endl. Gen. Suppl. i. 1372 (1841) = **Callitris**, Vent. (Conif.).
juniperoides, Endl. ex Gord. Pinet. Suppl. 107 = C. arborea.

PAROLINIA, Webb, in Ann. Sc. Nat. Sér. II. xiii. (1840) 133. t. 3. *CRUCIFERAE*, Benth. & Hook. f. i. 71.
ornata, *Webb, l. c.*—Ins. Canar.

PARONYCHIA, [Tourn.] Linn. Syst. ed. I (1735); Adans. Fam. ii. 272 (1763); Juss. in Mém. Mus. Par. ii. (1815) 389. *ILLECEBRACEAE*, Benth. & Hook. f. iii. 15.
BUINALIS, Rafin. New Fl. Am. iv. 40 (1836).
PLOTTOZIA, Arn. in Lindl. Introd. Nat. Syst. ed. II. 441 (1836).
Achyrantha, Desf. Tabl. ed. II. 54 = Alternanthera Achyrantha.
albella, *Juss. in Mém. Mus. Par.* ii. (1815) 390.—Arabia.
alsinefolia, *Juss. l. c.*—Hispan.
americana, Fenzl, ex Walp. Rep. i. 262 = Siphonychia americana.
andina, *A. Gray, Bot. U. St. Expl. Exped.* i. 128.—Reg. Andin.
andina, *Phil. in Anal. Univ. Chil.* (1870) II. 172.—Chili.
appressa, *Phil. in Linnaea*, xxxiii. (1864–65) 79.—Chili.
arabica, *DC. Cat. Hort. Monsp.* (1813) 130; *Delile, Fl. Egypte*, 187. t. 18.—Syria; Arabia; Aegypt.
arborea, Walp. Rep. i. 261 = Acronychia arborea.

PARONYCHIA:—
Arbuscula, *C. Gay, Fl. Chil.* ii. 520.—Chili.
aretioides, *DC. Prod.* iii. 371.—Hispan.
argentea, *Lam. Fl. Fr.* iii. 230.—Reg. Mediterr.
argentea, Schur, Enum. Pl. Transs. 223 = capitata.
argyrocoma, *Nutt. Gen. Am.* i. 160.—Am. bor.
aristata, DC. in Poir. Encyc. v. 23 = Polycarpaea aristata.
arragonica, *Schult. in Roem. & Schult. Syst.* v. 520.—Hispan.
aurasiaca, Webb, ex Coss. in Bull. Soc. Bot. Fr. xi. (1864) 160 = argentea.
australis, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 337.—Chili.
Baldwinii, *Fenzl, in Walp. Rep.* i. 262.—Am. bor.
bengalensis, Roem. & Schult. Syst. v. 521 = Achyranthes bengalensis.
bogotensis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 147.—N. Granat.
bonariensis, *DC. Prod.* iii. 370.—Reg. Argent.
brasiliana, DC. in Poir. Encyc. v. 23 = bonariensis.
brevistipulata, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 99.—Hispan.
bryoides, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 302.—Abyss.
Bungei, *Boiss. Fl. Orient.* i. 744.—Persia.
camphorosmoides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 187.—Bras.
canadensis, Wood, Class-Book (1861), 262 = Anychia dichotoma.
canariensis, *Juss. in Mém. Mus. Par.* ii. (1815) 390.—Ins. Canar.
capensis, Eckl. ex Steud. Nom. ed. II. ii. 269 = Cyathula globulifera.
capitata, *Lam. Fl. Fr.* iii. 229.—Europ. austr.; Oriens.
capitata, Reichb. Fl. Germ. Excurs. 564 = P. Cephalotes.
Cephalotes, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 184.—Tauria.
chilensis, *DC. Prod.* iii. 370.—Chili.
chionaea, Boiss. Diagn. Ser. I. iii. 9 = capitata.
communis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 186.—Bras.
confertissima, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 44; *Contrib.* 96.—Reg. Argent.
coquimbensis, *C. Gay, Fl. Chil.* ii. 520.—Chili.
Cossoniana, *J. Gay, ex Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 486.—Alger.
curdica, Walp. Rep. v. 75 = kurdica.
cymosa, *DC. in Poir. Encyc.* v. 26.—Reg. Mediterr. occ.
depressa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 171 = Jamesii.
desertorum, *Boiss. Diagn.* Ser. I. iii. 11.—Aegypt.; Arab.; Persia.
dichotoma, DC. in Poir. Encyc. v. 25 = Herniaria dichotoma.
dichotoma, *Nutt. Gen. Am.* i. 159.—Am. bor.
dichotoma, Torr. in Ann. Lyc. N. York, ii. (1828) 206 = Jamesii.
divaricata, Juss. in Mém. Mus. Par. ii. (1815) 390 = Polycarpaea Teneriffae.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 170.—Am. bor. occ.
dubia, Hort. ex Moq. in DC. Prod. xiii. II. 361 = Alternanthera sessilis?
echinata, *Lam. Fl. Fr.* iii. 232.—Reg. Mediterr.
ficoidea, Desf. Tabl. ed. I. 44; ed. II. 54 = Alternanthera sessilis.
flavescens, Boiss. Diagn. Ser. I. x. 14 = sinaica.
frutescens, Desf. Tabl. ed. I. 41; ed. II. 54 = Alternanthera frutescens.
fruticosa, *Lam. Fl. Fr.* iii. 230.—Reg. Mediterr. occ.
glomerata, Moench, Meth. 315 = argentea.
Hartwegiana, *Rohrb. in Linnaea*, xxxvii. (1871–73) 204.—Ecuador.
herniarioides, *Nutt. Gen. Am.* i. 159.—Am. bor.
hispanica, DC. in Poir. Encyc. v. 24 = argentea.
hispida, Steud. ex Boiss. Fl. Orient. i. 744 = kurdica.
hungarica, Griseb. Spicil. Fl. Rumel. i. 215 = P. Cephalotes.
illecebroides, *Webb, in Hook. Niger Fl.* 106. t. 7.—Ins. Cap. Virid.
imbricata, *Boiss. & Haussk. in Boiss. Fl. Orient.* i. 744.—Syria.
imbricata, Reichb. Fl. Germ. Excurs. 564 = capitata.
italica, Roem. & Schult. Syst. v. 518 = argentea.

PARONYCHIA:—
Jamesii, *Torr. & Gray, Fl. N. Am.* i. 170.—Am. bor occ.
Kapela, *A. Kern. in Oestr. Bot. Zeitschr.* xix. (1869) 367.—Europ. austr.
Kochiana, Boiss. Diagn. Ser. I. x. 13 = capitata.
kurdica, *Boiss. l. c.* iii. 10.—Oriens.
lanata, Moench, Meth. Suppl. 112 = Aerva lanata.
lanuginosa, *Poir. Encyc. Suppl.* iv. 303; Tuss. Fl. Antilles, ii. 90. t. 21.—Ins. S. Doming.
laurifolia, Walp. Rep. i. 261 = Acronychia laurifolia.
Lindheimeri, *Engelm. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 152.—N. Mexic.
linearifolia, DC. in Poir. Encyc. v. 26 = Polycarpaea linearifolia.
longifolia, *Steud. Nom.* ed. II. ii. 269.—Chili.
longiseta, Webb & Berth. Phyt. Canar. i. 163, in textu = arabica.
macrosepala, *Boiss. Diagn.* Ser. I. iii. 11; *Ball, in Journ. Bot.* xiii. (1875) 204.—Reg. Mediterr.
Mandoniana, *Rohrb. in Linnaea*, xxxvii. (1871–73) 208.—Bolivia.
maritima, DC. ex G. Don, in Loud. Hort. Brit. 91 = capitata.
mexicana, *Hemsl. Diagn. Pl. Nov.* 36.—Mexic.
mutica, *Phil. in Linnaea*, xxxiii. (1864–65) 79.—Chili.
nitida, Gaertn. Fruct. ii. t. 128 = argentea.
nivea, DC. in Poir. Encyc. iv. 25 = capitata.
nivea, C. A. Mey. Verz. Pfl. Cauc. 154 = kurdica.
paniculata, *Benth. & Hook. f. Gen.* iii. 16.—Hispan.
polycnemoides, Schlecht. in Linnaea, xiii. (1839) 407 = Pentacaena polycnemoides.
polygonifolia, *DC. Fl. Fr.* iii. 403.—Europ. austr.
pubescens, DC. l. c. = Herniaria hirsuta.
pulvinata, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 58.—Am. bor. occ.
ramosissima, DC. Mém. Paronych. 12. t. 4 = Pentacaena polycnemoides.
rigida, Moench, Meth. 315 = capitata.
riparia, *Chapm. Fl. S. U. St. Suppl.* 607.—Am. bor.
Rosetta, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 188.—Bras.
Rugelii, Shuttl. ex Chapm. Fl. S. U. St. 47 = Siphonychia Rugelii.
sclerocarpa, Steud. Nom. ed. II. ii. 269 = seq.
sclerocephala, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 262 = Sclerocephalus arabicus.
sedifolia, *R. Br. in Salt, Abyss. App.* 64; *Roem. & Schult. Syst.* v. 523.—Abyss.
serpyllifolia, Auct. ex Nym. Consp. 255 = capitata.
serpyllifolia, *DC. Fl. Fr.* iii. 404; *et in Poir. Encyc.* v. 24.—Europ. austr.
sessiliflora, *Nutt. Gen. Am.* i. 160.—Am. bor.
sessilis, Desf. Tabl. ed. II. 45 = Alternanthera sessilis.
setacea, *Torr. & Gray, Fl. N. Am.* i. 170.—Am. bor. occ.
sinaica, *Fresen. in Mus. Senckenb.* i. (1833) 180.—Syria.
Smithii, Choisy, ex Link, in Buch, Beschr. Canar. Ins. 142 = Polycarpaea Smithii.
splendens, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 184 = kurdica.
striata, DC. in Poir. Encyc. v. 25 = Polycarpon tetraphyllum.
subulata, DC. l. c. = Polycarpaea corymbosa.
suffruticosa, DC. l. c. = fruticosa.
tenella, Desf. Cat. Hort. Par. ed. III. 407 = Alternanthera tenella?
tenuifolia, *Link, Enum. Hort. Berol.* i. 151; *DC. Prod.* iii. 372.—Hab.?
tetragona, Moench, Meth. 315 = Alternanthera sessilis.
trinervis, Dulac, Fl. Hautes-Pyr. 364 = polygonifolia.
urceolata, Shuttl. ex Chapm. Fl. S. U. St. 47 = Siphonychia americana.
verticillata, Lam. Fl. Fr. iii. 231; DC. in Poir. Encyc. v. 23 = Illecebrum verticillatum.
virginica, Spreng. Syst. i. 822 = dichotoma.

PAROPHIORRHIZA, C. B. Clarke, ex Hook. f. Fl. Brit. Ind. iii. 85 (1880) = **Mitreola**, Linn. (Logan.).
khasiana, C. B. Clarke, l. c. = M. pedicellata.

PAROPSIA, Noronha, ex Thou. Hist. Vég. Isles Afr. 59. t. 19 (1805). *PASSIFLOREAE*, Benth. & Hook. f. i. 812.
TRICHODIA, Griff. Notul. iv. 570 (1854).
edulis, *Thou. Hist. Vég. Isles Afr.* ii. 59. t. 19.—Madag.

PAROPSIA:—
grewioides, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* ii. 505.—Afr. trop.
guineensis, *Oliver, in Journ. Linn. Soc.* viii. (1865) 161.—Afr. trop.
malayana, *Planch. ex Mast. in Trans. Linn. Soc.* xxvii. (1871) 639.—Malacca.
obscura, *O. Hoffm. Sert. Madag.* 20.—Madag.
vareciformis, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 639.—Malacca.

PAROSELA, Cav. Desc. 185 (1802) = **Dalea**, Linn. (Legumin.).
Lagopus, Cav. l. c. 187 = D. Lagopus.
lutea, Cav. l. c. 186 = D. lutea.
mutabilis, Cav. l. c. = D. mutabilis.
tuberculata, Cav. Elench. Hort. Madr. (1805); ex DC. Prod. ii. 245 = D. thymoides.

PAROSELLA, Cav. ex DC. Prod. ii. 244 (1825) = praec.

PARQUI, Adans. Fam. ii. 219 (1763) = **Cestrum**, Linn. (Solanac.).

PARRIA, Steud. Nom. ex II. ii. 269 (1841) = **Parrya**, R. Br. (Crucif.).

PARROTIA, C. A. Mey. Verz. Pfl. Cauc. 46 (1831). *HAMAMELIDEAE*, Benth. & Hook. f. i. 666.
Jacquemontiana, *Decne. in Jacquem. Voy. Bot.* 73. t. 82.—Reg. Himal.
persica, *C. A. Mey. Verz. Pfl. Cauc.* 46.—Persia; Reg. Caucas.
siderodendron, C. A. Mey. ex Ledeb. Fl. Ross. ii. 377 = persica.

PARROTIA, Walp. Ann. i. 754 (1849?), sphalm. = BARROTIA, Gaudich. = **Pandanus**, Linn.

PARRYA, R. Br. in Parry, Voy. App. 268 (1824). *CRUCIFERAE*, Benth. & Hook. f. i. 67.
ERMANIA, Cham. in Linnaea, vi. (1831) 533.
NEUROLOMA, Andrz. ex DC. Prod. i. 156 (1824).
PACHYNEURUM, Bunge, in Linnaea, xiv. (1840) 121.
arctica, *R. Br. in Parry, Voy. App.* 270.—Reg. arct.
arctica, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 242 = macrocarpa.
arenicola, *Hook. f. in Trans. Linn. Soc.* xxiii. (1861) 285, 315.—Am. arct.
Ermani, *Ledeb. Fl. Ross.* i. 132.—Sibir.
eriocalyx, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 234.—Turkestan.
exscapa, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 28.—Tibet. occ.; Mont. Altai.
flabellata, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) I. 261.—Rossia.
fruticulosa, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 237.—Kokan.
integerrima, *G. Don, Gen. Syst.* i. 173.—Sibir.
lanuginosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 136.—Reg. Himal.
Linnaeana, *Ledeb. Fl. Ross.* i. 131.—Kamtschatk.
macrocarpa, *R. Br. in Parry, Voy. App.* 270.—Reg. arct.; As. centr.
microcarpa, *Ledeb. Fl. Ross.* i. 132.—Mont. Altai.
nudicaulis, Boiss. Fl. Orient. i. 159 = macrocarpa.
pinnatifida, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 147.—Mont. Altai.; Afghan.
platycarpa, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 136.—Reg. Himal.
pumila, *Kurz, in Flora*, lv. (1872) 285.—Tibet.
scapigera, *G. Don, Gen. Syst.* i. 173.—Sibir.
stenocarpa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 147.—Mont. Altai.
stenoloma, Schrenk, ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 69 = stenocarpa.

PARRYELLA, Torr. & Gray, ex A. Gray, in Proc. Am. Acad. vii. (1868) 397. *LEGUMINOSAE*.
filifolia, *Torr. & Gray, l. c.*—Mexic.

PARSONSIA, P. Br. Hist. Jamaic. 199. t. 21 (1756) = **Cuphea**, P. Br. (Lythrar.).
herbacea, Jaume St Hil. Expos. ii. 178 = C. Parsonsia.
prostrata, Bert. ex DC. Prod. iii. 86 = C. Parsonsia.

PARSONSIA, R. Br. in Mem. Wern. Soc. i. (1809) 64. *APOCYNACEAE*, Benth. & Hook. f. ii. 711.
GASTRANTHUS, F. Muell. Fragm. vi. 130 (1868).
HELICANDRA, Hook. & Arn. Bot. Beech. Voy. 198, in nota (1833).
HELIGME, Blume, Fl. Jav. Praef. p. vii (1828).
HELYGIA, Blume, Bijdr. 1043 (1826).
SPIROSTEMON, Griff. Notul. iv. 80 (1854).
acuminata, Wall. Cat. n. 1634 = spiralis.
albiflora, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 118.—N. Zel.
barbata, Blume, Bijdr. 1042 = Parameria glandulifera.
bracteata, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 287.—Reg. Argent.
capsularis, *R. Br. in Mem. Wern. Soc.* i. (1809) 65.—Austral.; N. Zel.
capsularis, Endl. ex Deless. Ic. Sel. v. t. 49 = albiflora.
capsularis, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 118 = rosea.
corymbosa, *R. Br. in Mém. Wern. Soc.* i. (1809) 65.—Ins. S. Doming.
Cumingiana, *A. DC. in DC. Prod.* viii. 402.—Ins. Philipp.
diaphanophlebia, F. Muell. Fragm. ii. 158 = Lyonsia diaphanophlebia.
eucalyptifolia, Benth. Fl. Austral. iv. 323 = seq.
eucalyptophylla, F. Muell. l. c. 159 = Lyonsia eucalyptifolia.
floribunda, R. Br. in Mem. Wern. Soc. i. (1809) 65 = Forsteronia floribunda, *G. F. W. Mey.*
Forsteri, G. Don, Gen. Syst. iv. 79 = rosea, albiflora.
glaucescens, F. Muell. Fragm. vi. 126 = lanceolata.
Helicandra, *Hook. & Arn. Bot. Beech. Voy.* 197.—China.
heterophylla, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 46 = albiflora.
induplicata, F. Muell. Fragm. vi. 129 = Lyonsia induplicata.
javanica, Blume, Bijdr. 1041 (= *Chavannesia javanica*).—Java.
lanceolata, *R. Br. Prod.* 466.—Austral.
Langiana, F. Muell. Fragm. vi. 128 = Lyonsia Langiana.
Leichhardtii, *F. Muell. l. c.*—Austral.
leptocarpa, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 287.—Bras.
lilacina, F. Muell. Fragm. vi. 127 = Lyonsia lilacina.
macrocarpa, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 331.—N. Zel.
mollis, R. Br. Prod. 466 = lanceolata.
myrtifolia, *Roem. & Schult. Syst.* iv. 403.—Madag.
oblonga, Wall. Cat. n. 1632 = spiralis.
ovata, Wall. l. c. n. 1630 = Pottsia cantonensis.
pauciflora, Wall. l. c. n. 1633 = spiralis.
Rheedii, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 130.—Ins. Philipp.
rosea, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 118.—N. Zel.
spicata, *R. Br. in Mem. Wern. Soc.* i. (1809) 65.—N. Granat.
spiralis, Vidal, Sin. Fl. Leñ. t. 66. f. E = Rheedii.
spiralis, *Wall. Cat.* n. 1631.—As. trop.
variabilis, Lindl. in Journ. Hort. Soc. v. (1850) 196 = rosea?
velutina, *R. Br. Prod.* 466.—Austral.
ventricosa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 71.—Austral.

PARTHENICE, A. Gray, Pl. Wright. ii. 85 (in Smithson. Contrib. v.) (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 352.
mollis, *A. Gray, l. c.*—N. Mexic.

PARTHENIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 265 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 351.
ARGYROCHAETA, Cav. Ic. iv. 54, t. 378 (1791).
BOLOPHYTA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 347.
HYSTEROPHORUS, Adans. Fam. ii. 128 (1763).
VILLANOVA, Orteg. Hort. Matr. Dec. 47. t. 6 (1789).
alpinum, *Torr. & Gray, Fl. N. Am.* ii. 285.—Am. bor. occ.
amplectens, Rafin. New Fl. Am. ii. 26 = integrifolium.
angustifolium, Rafin. l. c. 25 (Quid?).—Am. bor.
annuum, Crantz, Inst. i. 345 = Iva annua.
argentatum, *A. Gray, in Torr. Bot. Mex. Bound.* 86.—Texas.

PARTHENIUM:—
confertum, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 216.—Mexic.
Dillenianum, *Rafin. New Fl. Am.* ii. 26 (Quid ?).—Am. bor.
Elliotanum, *Rafin. l. c.* (Quid ?).—Am. bor.
frutescens, Crantz, Inst. i. 345 = Iva frutescens.
fruticosum, *Less. in Linnaea*, v. (1830) 152.—Am. austr.
hispidum, *Rafin. New. Fl. Am.* ii. 35 (Quid ?).—Am. bor.
Hysterophorus, *Linn. Sp. Pl.* 988.—Am. bor. et austr.
incanum, *H. B. & K. Nov. Gen. et Sp.* iv. 260. t. 391.—Am. bor. occ.
integrifolium, *Linn. Sp. Pl.* 988.—Am. bor.
lobatum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457 = P. Hysterophorus.
luteum, Spreng. Nov. Prov. 31 = Guizotia abyssinica.
lyratum, *A. Gray, Syn. Fl. N. Am.* i. II. 244.—Texas; Mexic.
Matricaria, Gueldenst. Reisen, ii. 25 = Chrysanthemum Parthenium.
pinnatifidum, Stokes, Bot. Mat. Med. iv. 278 = P. Hysterophorus.
pumilum, *Rafin. New Fl. Am.* ii. 25 (Quid ?).—Am. bor.
ramosissimum, DC. Prod. v. 532 = incanum.
sinuatum, *Rafin. New Fl. Am.* ii. 25 (Quid ?).—Am. bor.
tomentosum, *DC. Prod.* v. 532.—Mexic.

PARTHENOPSIS, Kellogg, in Proc. Calif. Acad. v. (1873) 100 = **Venegazia**, DC. (Compos.).
maritimus, Kellogg, l. c. 101 = V. carpesioides.

PARTHENOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 158 = **Ornithogalum**, Tourn. (Liliac.).
narbonensis, Fourr. l. c. = O. narbonnense.
pyrenaica, Fourr. l. c. = O. pyrenaicum.

PARTHENOXYLON, Blume, Mus. Bot. Lugd. Bat. i. 322 (1851) = **Cinnamomum**, Tourn. (Laurin.).
porrectum, Blume, l. c. 323 = C. Parthenoxylon.
pruinosum, Blume, l. c.—Java.
Pseudo-sassafras, Blume, l. c. = C. Parthenoxylon.

PARVATIA, Decne. in Arch. Mus. Par. i. (1837) 190. t. 12 A. *BERBERIDEAE*, Benth. & Hook. f. i. 42.
Brunoniana, *Decne. l. c.*; *et in Ann. Sc. Nat.* Sér. II. xii. (1839) 104.—Reg. Himal.

PARYPHANTHA, Schau. in Linnaea, xvii. (1843) 235 = **Thryptomene**, Endl. (Myrtac.).
camphorata, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 116 = T. saxicola.
cuspidata, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 321 = T. tenella.
Mitchelliana, Schau. in Linnaea, xvii. (1843) 235 = T. Mitchelliana.

PARYPHANTHE, Benth. Fl. Austral. iii. 62 (1866) = praec.

PARYPHOSPHAERA, Karst. Fl. Columb. iii. 7. t. 104 (1862) = **Parkia**, R. Br. (Legumin.).
arborea, Karst. l. c. = Parkia Paryphosphaera.

PASANIA, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81 = **Quercus**, Tourn. (Cupul.).
acuminata, Oerst. l. c. 83 = Q. acuminata.
acuminatissima, Oerst. l. c. 84 = Q. Junghuhniana.
Amherstiana, Oerst. l. c. = Q. Amherstiana.
cornea, Oerst. l. c. 83 = Q. cornea.
crassinervia, Oerst. l. c. 84 = Q. crassinervia.
cuspidata, Oerst. l. c. = Q. cuspidata.
dealbata, Oerst. l. c. = Q. dealbata.
densiflora, Oerst. l. c. 83 = Q. densiflora.
Eyrei, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. Eyrei.
fenestrata, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 84 = Q. fenestrata.
fissa, Oerst. l. c. = Q. fissa.
glabra, Oerst. l. c. 83 = Q. glabra.
glomerata, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. glomerata.
glutinosa, Oerst. l. c. = Q. glutinosa.

PASANIA:—
Harlandi, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 83 = Quercus Harlandi.
Irwinii, Oerst. l. c. = Q. Irwinii.
Korthalsii, Oerst. l. c. = Q. Korthalsii.
lanceaefolia, Oerst. l. c. 84 = Q. lanceaefolia.
lappacea, Oerst. l. c. = Q. lappacea.
lithocarpaea, Oerst. l. c.—Assam.
littoralis, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. litoralis.
mixta, Oerst. in Kjoeb. Vidensk. Meddel. 1866) 83 = Q. spicata.
Molucca, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 373 = Q. Molucca.
neurophylla, Oerst. l. c. 379 = Q. sundaica ?
oligoneura, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 84 = Q. oligoneura.
pallida, Oerst. l. c. 83 = Q. pallida.
Pinanga, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. Pinanga.
placentaria, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 83 = Q. spicata.
plumbea, Oerst. l. c. 84 = Q. plumbea.
polystachya, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. polystachya.
pruinosa, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 83 = Q. pruinosa.
Pseudo-molucca, Oerst. l. c. = Q. Pseudo-molucca.
rotundata, Oerst. l. c. = Q. rotundata.
sessilifolia, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Q. sessilifolia.
spicata, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 83 = Q. spicata.
sundaica, Oerst. l. c. = Q. sundaica.
thalassica, Oerst. l. c. = Q. thalassica.
urceolaris, Oerst. l. c. = Q. Lamponga ?
variabilis, Regel, Cat. Pl. Hort. Aksakov. 102 = Q. variabilis.

PASCALIA, Orteg. Hort. Matr. Dec. 39. t. 4 (1797). *COMPOSITAE*, Benth. & Hook. f. ii. 369.
LORENTZIA, Griseb. in Goett. Abh. xix. (1874) 182.
baccata, Spreng. Syst. iii. 602 = Wulffia stenoglossa.
glandulosa, *Fisch. ex Steud. Nom.* ed. I. 592.—Hab. ?
glauca, *Orteg. Hort. Matr. Dec.* 39.—Chili.

PASCHANTHUS, Burch. Trav. i. 543 (1822) = **Modecca**, Lam. (Passifl.).
repandus, Burch. l. c. = M. Paschanthus.

PASINA, Adans. Fam. ii. 189 (1763) = **Horminum**, Linn. (Labiat.).

PASITHEA, D. Don, in Edinb. N. Phil. Journ. xiii. (Oct. 1832) 236. *LILIACEAE*, Benth. & Hook. f. iii. 791.
coerulea, *D. Don, in Edinb. N. Phil. Journ.* xiii. (Oct. 1832) 237.—Chili.

PASPALANTHIUM, Desv. Opusc. 59 (1831) = **Paspalum**, Linn. (Gramin.).
stoloniferum, Desv. l. c. = Paspalum racemosum.

PASPALUM, Linn. Syst. ed. X. 855 (1759). *GRAMINEAE*, Benth. & Hook. f. iii. 1097.
ANACHYRIS, Nees, in Hook. Kew Journ. ii. (1850) 103.
ANASTROPHUS, Schlecht. in Bot. Zeit. viii. (1850) 681.
AXONOPUS, Beauv. Agrost. 12 (1812).
CABRERA, Lag. Gen. et Sp. Nov. 5 (1816).
CERESIA, Pers. Syn. i. 85 (1805).
CLEACHNE, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 285.
CYMATOCHLOA, Schlecht. in Bot. Zeit. xii. (1854) 821.
LAPPAGOPSIS, Steud. Syn. Pl. Gram. 112 (1855).
MAIZILLA, Schlecht. in Bot. Zeit. viii. (1850) 601.
MOENCHIA, Wender. ex Steud. Nom. ed. II. ii. 153 (1841).
PASPALANTHIUM, Desv. Opusc. 59 (1831).
SABSAB, Adans. Fam. ii. 31 (1763).
WIRTGENIA, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 40 (1877).
abbreviatum, *Trin. ex Steud. Nom.* ed. II. ii. 270.—Mexic.
abstrusum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 135.—Bras.
acuminatum, *Raddi, Agrost. Bras.* 25.—Bras.

PASPALUM:—
adelogaeum, *Steud. Syn. Pl. Gram.* 27.—Java.
adpressum, Pers. Syn. i. 85, in syn. = Panicum paspaloides.
aegyptiacum, Poir. Encyc. Suppl. iv. 314 = Panicum sanguinale.
affine, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 125.—Costa Rica.
affine, Steud. Syn. Pl. Gram. 24 = tenellum ?
africanum, *Poir. Encyc. Suppl.* iv. 314.—Afr. trop.
alternans, Steud. Syn. Pl. Gram. 26 = scrobiculatum.
alterniflorum, *A. Rich. Fl. Cub. Fanerog.* ii. 299.—Cuba.
altissimum, *Le Conte, in Journ. Phys.* xci. (1820) 285.—Am. bor.
amazonicum, Trin. in Linnaea, x. (1836) 294 = plicatulum.
ambiguum, DC. Fl. Fr. iii. 16 = Panicum glabrum.
ambiguum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 59 = pleostachyum.
ammodes, *Trin. Gram. Panic.* 120.—Bras.
ancylocarpum, *Nees, ex Steud. Syn. Pl. Gram.* 27.—Bras.
angustifolium, *Le Conte, in Journ. Phys.* xci. (1820) 285.—Am. bor.
angustifolium, Nees, ex Trin. Gram. Panic. 99 = Neesii.
annulatum, Fluegge, Gram. Monog. 133 = Eriochloa annulata.
antillense, Husnot, in Bull. Soc. Linn. Normand. Sér. II. v. (1870) 260 = undulatum.
apiculatum, *Doell, in Mart. Fl. Bras.* ii. II. 48.—Am. trop.
appendiculatum, *J. & C. Presl, Rel. Haenk.* i. 211.—Panama.
appressum, Lam. Illustr. i. 176 = Panicum paspaloides.
approximatum, *Doell, in Mart. Fl. Bras.* ii. II. 82.—Am. trop.
arenarium, *Schrad. ex Schult. Mant.* ii. 172.—Bras.
arenicolum, *C. Muell. in Bot. Zeit.* xix. (1861) 325.—Guian.
argyrocondylon, *Steud. Syn. Pl. Gram.* 22.—Guian.
aristatum, Moench, Meth. 178 = Beckmannia erucaeformis.
arundinaceum, *Poir. Encyc. Suppl.* iv. 310.—m. Abor.
aspidiotes, *Trin. Sp. Gram. Ic.* iii. t. 269.—Bras.
atrocarpum, *Steud. Syn. Pl. Gram.* 25.—Hab. ?
attenuatum, J. & C. Presl, Rel. Haenk. i. 212 = scoparium ?
aureum, *H. B. & K. Nov. Gen. et Sp.* i. 93. t. 27.—Am. austr.
aureum, Trin. Gram. Panic. 88 = pulchrum.
auriculatum, J. & C. Presl, Rel. Haenk. i. 217 = scrobiculatum.
axicilium, Steud. Syn. Pl. Gram. 20 = Ischaemum Urvilleanum.
barbatum, Nees, Agrost. Bras. 27 = barbigerum.
barbatum, Schult. Mant. ii. 558 = distichum.
barbatum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 53.—Afr. trop.
barbigerum, *Kunth. Rév. Gram.* i. 24.—Bras.
barbigreum, Steud. Nom. ed. II. ii. 260. in syn., sphalm. = praec.
bicorne, Lam. Illustr. i. 176 = Panicum sanguinale.
bicrurulum, Salzm. ex Steud. Syn. Pl. Gram. 21 = seq. ?
bicrurum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 55 = conjugatum.
bifarium, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 179 = scrobiculatum ?
biflorum, *Rasp. in Ann. Sc. Nat.* Sér. I. v. (1825) 301 (sp. dub.).—Hab. ?
biglume, *Steud. Syn. Pl. Gram.* 24.—Hab. ?
bistipulatum, Hochst. ex Steud. Syn. Pl. Gram. 29 = repens.
blepharophorum, Roem. & Schult. Syst. ii. 292 = ciliatum.
Blodgettii, *Chapm. Fl. S. U. St.* 571.—Ind. occ.
Boivini, Steud. Syn. Pl. Gram. 416 = heteropodium.
Bonplandianum, *Fluegge, Gram. Monog.* 71.—Ecuador.
borbonicum, *Steud. Syn. Pl. Gram.* 27.—Ins. Mascar.
Boryanum, *J. & C. Presl, Rel. Haenk.* i. 209.—Ins. Philipp.
Boscianum, Fluegge, Gram. Monog. 170 = undulatum.
brachiatum, Trin. ex Nees, Agrost. Bras. 62 = distichum.
brachystachyum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 133 = Eriochloa distachya.
bracteatum, Dufour, ex Kunth, Enum. Pl. i. 53 = distichum ?

PASPALUM :—
brasiliense, Spreng. Syst. i. 249 = dissitiflorum.
brevifolium, *Fluegge, Gram. Monog.* 150.—Geront. trop.
brunneum, Bosc, ex Poir. Encyc. Suppl. iv. 312 = undulatum.
Burchellii, *Munro, ex Doell, in Mart. Fl. Bras.* ii. II. 42.—Bras.
caespitosum, *Fluegge, Gram. Monog.* 161.—Ind. occ.
caespitosum, Hochst. ex Doell, in Mart. Fl. Bras. ii. II. 76, 79 = plicatulum, undulatum, dissectum.
campestre, *Schlecht. in Linnaea*, xxvi. (1853) 131.—Bras.
campestre, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 144 = pumilum.
canaliculatum, *Nees, Agrost. Bras.* 33.—Bras.
candidum, *Kunth, in Mém. Mus. Par.* ii. (1803) 68; *H. B. & K. Nov. Gen. et Sp.* i. 87.—Ecuador.
canescens, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 132.—Ins. Galapag.
canescens, Nees, ex Trin. Gram. Panic. 89 = chrysodactylon.
capillare, Boj. Hort. Maurit. 362 = heteropodium.
capillare, *Lam. Illustr.* i. 176.—Am. austr.
carinatum, *Fluegge, Gram. Monog.* 65.—N. Granat.
carolinianum, Poir. Encyc. Suppl. iv. 311 = Panicum sanguinale?
cartilagineum, J. & C. Presl, Rel. Haenk. i. 216 = scrobiculatum.
castaneum, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 348.—Bolivia.
caudicatum, *Wright, in Sauv. Fl. Cub.* 196.—Cuba.
chepica, Steud. Syn. Pl. Gram. 21 = distichum.
chinense, Nees, in Hook. & Arn. Bot. Beech. Voy. 231; et in Steud. Syn. Pl. Gram. 41 = brevifolium.
chrysites, *Doell, in Mart. Fl. Bras.* ii. II. 117.—Am. trop.
chrysoblephare, Doell, l. c. 119 = exasperatum.
chrysocomum, Trin. ex Nees, Agrost. Bras. 81 = chrysostachyum.
chrysodactylon, *Doell, in Mart. Fl. Bras.* ii. II. 118.—Am. trop.
chrysostachyum, *Schrad. in Schult. Mant.* ii. 176.—Ins. Philipp.
chrysotrichum, *J. & C. Presl, Rel. Haenk.* i. 211.—Ins. Philipp.
chrysotrichum, *Schrad. ex Schult. Mant.* ii. 176.—Bras.
ciliare, DC. Fl. Fr. Suppl. 250 = Panicum sanguinale.
ciliatifolium, Michx. Fl. Bor. Am. i. 44 = setaceum.
ciliatifolium, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 340 = conjugatum.
ciliatum, *H. B. & K. Nov. Gen. et Sp.* i. 87. t. 24.—Bras.
ciliatum, Lam. Illustr. i. 175 = conjugatum.
ciliatum, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 285 (Quid?).—Guian.
ciliatum, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. (1842) 238 = Humboldtianum.
clavuliferum, *Wright, in Sauv. Fl. Cub.* 195.—Cuba.
cochinchinense, *Willd. in Fluegge, Gram. Monog.* 198.—Cochinch.
cognatissimum, Steud. Syn. Pl. Gram. 18 = paniculatum.
cognatum, *Steud. l. c.* 28.—Java.
coloratum, Rich. ex Doell, in Mart. Fl. Bras. ii. II. 78 = scrobiculatum.
comans, *Trin. ex Doell, l. c.* 109.—Bras.
Commersonii, Lam. Illustr. i. 175 = scrobiculatum.
Commersonii, Zuccagni, in Roem. Collect. 122 = membranaceum.
commutatum, *Nees, Agrost. Bras.* 59.—Bras.
compactum, *Roth, Nov. Pl. Sp.* 36.—Ind. or.
complanatum, Nees, ex Trin. Gram. Panic. 96 = platycaule.
complicatum, D. Dietr. Syn. Pl. i. 242 = platycaule.
compressicaule, Raddi, Agrost. Bras. 29 = paniculatum.
compressum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301; Nees, ex Trin. Gram. Panic. 96 = platycaule.
concinnum, Steud. Syn. Pl. Gram. 19 = ischnocaulon.
confertum, Le Conte, in Journ. Phys. xci. (1820) 285 = undulatum.
confertum, Willd. ex Steud. Nom. ed. II. ii. 270 = candidum.
conjugatum, *Berg. in Act. Helv.* vii. (1772) 129. t. 8.—Reg. trop.

PASPALUM :—
consanguineum, Kunth, Enum. Pl. i. 46 = Panicum sanguinale.
conspersum, Schrad. ex Schult. Mant. ii. 174 = virgatum.
convexum, *Fluegge, Gram. Monog.* 175.—Mexic.
convexum, Willd. ex Doell, in Mart. Fl. Bras. ii. II. 79 = ancylocarpum.
corcovadense, Raddi, Agrost. Bras. 27 = laxum.
corcovadense, Trin. Sp. Gram. Ic. ii. t. 153 = densiflorum.
cordovense, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 477.—Mexic.
coromandelinum, Lam. Illustr. i. 175 = scrobiculatum.
corymbosum, Kunth, Rév. Gram. i. 25 = Panicum sanguinale.
coryphaeum, *Trin. Gram. Panic.* 114.—Bras.
costatum, Hochst. ex Steud. Syn. Pl. Gram. 58. = compactum.
cristatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 152.—Chili.
cromyorhizon, *Trin. ex Doell, in Mart. Fl. Bras.* ii. II. 74.—Am. trop.
ctenostachyum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 133 = grandiflora.
cubense, *Spreng. Neue Entdeck.* iii. 12.—Ind. occ.
cujabense, Trin. Sp. Gram. Ic. iii. t. 284 = stellatum.
Cumingii, Nees, ex Steud. Syn. Pl. Gram. 23 = dasypleurum.
Curtisianum, *Steud. l. c.* 26.—Am. bor.
curvistachyum, Raddi, Agrost. Bras. 26 = Panicum decumbens.
cuyabense, Auct. = *cujabense*.
cymbiforme, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 477.—Mexic.
cynosuroides, Brot. Fl. Lustit. i. 85 = Spartina stricta.
Dactylon, DC. Fl. Fr. iii. 16 = Cynodon Dactylon.
Dactylon, Lam. Illustr. i. 176 = Panicum glabrum.
dasyphyllum, Ell. Sketch, i. 105 = setaceum.
dasypleurum, *Kunze, ex Desv. in C. Gay, Fl. Chil.* i. 242.—Chili.
debile, Michx. Fl. Bor. Am. i. 44 = setaceum.
debile, Muhl. Desc. Gram. 91 = arenarium.
debile, Poir. Encyc. v. 34 = Panicum sanguinale.
decumbens, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 285 (Quid?).—Guian.
decumbens, Sagot, ex Doell, in Mart. Fl. Bras. ii. II. 77 = undulatum.
decumbens, Sw. Prod. Veg. Ind. Occ. 22 = Panicum decumbens.
Delochei, Steud. Syn. Pl. Gram. 23 = fasciculatum.
densiflorum, *Doell, in Mart. Fl. Bras.* ii. II. 52.—Am. trop.
densum, *Poir. Encyc.* v. 32.—Bras.
denticulatum, *Trin. Gram. Panic.* 111.—Am. austr.
depauperatum, *J. & C. Presl, Rel. Haenk.* i. 215.—Peruv.
depressum, Steud. Syn. Pl. Gram. 20 = distichum.
didactylum, Salzm. ex Steud. l. c. = distichum.
difforme, *Le Conte, in Journ. Phys.* xci. (1820) 284.—Am. bor.
diffusum, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 285 (Quid?).—Guian.
Digitaria, *C. Muell. in Bot. Zeit* xix. (1861) 324.—Am. bor.
Digitaria, Poir. Encyc. Suppl. iv. 316 = dichotomum.
digitarioides, *Desv. Opusc.* 56.—Bras.
digitatum, *Kunth, Rév. Gram.* i. 24.—Jamaic.
dilatatum, *Poir. Encyc.* v. 35.—Bras.
dilatatum, Trin. in Linnaea, x. (1835) 294 = dasypleurum.
dimidiatum, Linn. Syst. ed. X. 855 = scrobiculatum.
dissectum, *Linn. Sp. Pl.* ed. II. 81.—Am. austr.
dissectum, Sw. ex Roem. & Schult. Syst. ii. 308 = caespitosum.
dissectum, Thunb. Fl. Jap. 45 = scrobiculatum.
dissitiflorum, *Trin. Gram. Panic.* 92.—Bras.
distachyon, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 57 = pumilum.
distachyon, Willd. ex Doell, l. c. 73 = distichum.
distachyum, *Poit. ex Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 142.—Ins. S. Doming.
distans, Nees, ex Trin. Gram. Panic. 83 = Panicum sanguinale.
distichophyllum, *H. B. & K. Nov. Gen. et Sp.* i. 86.—Am. austr.
distichum, Houtt. Handleid. xiii. 167. t. 89. f. 4 = Eriochloa villosa.

PASPALUM :—
distichum, *Linn. Amoen. Acad.* v. 391; *Burm. f. Fl. Ind.* 23.—Reg. trop. et subtrop.
divergens, *Doell, in Mart. Fl. Bras.* ii. II. 71.—Am. trop.
dolichopus, Trin. ex Steud. Nom. ed. II. ii. 271 = conjugatum.
dolichostachyum, Trin. ex Nees, Agrost. Bras. 30 = pellitum.
Dombeyi, *Spreng. Nachr.* I. *Bot. Gart. Halle*, 31 (Quid?).—Hab.?
Drummondi, *C. Muell. in Bot. Zeit.* xix. (1861) 332.—Am. bor.
dubium, *DC. Cat. Hort. Monsp.* 130.—Hab.?
effusum, *Nees, in Hook. Kew Journ.* ii. (1850) 104.—Bras.
effusum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Milium effusum.
elatum, *Rich. ex Doell, in Mart. Fl. Bras.* ii. II. 78.—Am. trop.
elegans, Fluegge, Gram. Monog. 183 = tenellum.
elegans, Hort. ex Roem. & Schult. Syst. ii. 290 = membranaceum.
elegantulum, *J. & C. Presl, Rel. Haenk.* i. 211.—Peruv.
ellipticum, *Doell, in Mart. Fl. Bras.* ii. II. 71.—Am. trop.
elongatum, *Griseb. in Goett. Abh.* xix. (1874) 260.—Reg. Argent.
eminens, Nees, Agrost. Bras. 30 = barbigerum.
erianthum, *Nees, ex Trin. Gram. Panic.* 121.—Bras.
eriophorum, *Schult. Mant.* ii. 560 = dilatatum.
eriophorum, Willd. ex Nees, Agrost. Bras. 56 = conjugatum.
eriostachyum, D. Dietr. Syn. Pl. i. 254, in syn., sphalm. = *curvistachyum*.
erucaeforme, *Spreng. Nachr.* I. *Bot. Gart. Halle*, 31 (Quid?).—Hab.?
eruciferum, Trin. ex Steud. Nom. ed. ii. 271, 272 = quadrifarium.
eucanum, Steud. l. c. 271 = seq.
eucomum, *Nees, Agrost. Bras.* 37.—Am. trop.
exaltatum, *J. & C. Presl, Rel. Haenk.* i. 219.—Chili.
exasperatum, *Nees, Agrost. Bras.* 81.—Bras.
excavatum, Nees, ex Trin. Gram. Panic. 88 = aureum.
exile, *Kippist, in Proc. Linn. Soc.* i. (1842) 157.—Afr. trop.
expansum, *Doell, in Mart. Fl. Bras.* ii. II. 81.—Am. trop.
extenuatum, *Nees, Agrost. Bras.* 25.—Bras.
falcatum, *Nees, ex Steud. Syn. Pl. Gram.* 31.—Bras.
Falcula, *Doell, in Fl. Bras.* ii. II. 60.—Am. trop.
familiare, *Steud. Syn. Pl. Gram.* 24.—N. Granat.
fasciculatum, *Willd. ex Fluegge, Gram. Monog.* 69.—Am. trop.
fastigiatum, *Nees, ex Trin. Gram. Panic.* 100.—Bras.
fernandezianum, Colla, in Mem. Acc. Torin. xxxix. (1836) 27 = distichum.
ferrugineum, Trin. Sp. Gram. Ic. ii. t. 136 = quadrifarium.
filiculme, Nees, ex Thw. Enum. Pl. Zeyl. 358 = brevifolium.
filifolium, *Nees, ex Steud. Syn. Pl. Gram.* 22.—Bras.
filifolium, Kunth, Enum. Pl. i. 582 (Index), sphalm. = fissifolium.
filiforme, Fluegge, Gram. Monog. 139 = Panicum filiforme.
filiforme, Steud. Nom. ed. I. 586 = Panicum sanguinale.
filiforme, *Sw. Prod. Veg. Ind. Occ.* 22.—Ind. occ.
filostachyum, *Rich. ex Steud. Syn. Pl. Gram.* 20.—Ind. occ.
fimbriatum, *H. B. & K. Nov. Gen. et Sp.* i. 93.—Am. austr.
firmum, Trin. Gram. Panic. 105 = scrobiculatum.
fissifolium, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 104 = distichum.
fissifolium, *Raddi, Agrost. Bras.* 26.—Bras.
flaccidum, *Nees, Agrost. Bras.* 48.—Bras.
flavum, *J. & C. Presl, Rel. Haenk.* i. 220.—Hab.?
flexuosum, Klein, ex J. & C. Presl, l. c. 215 = scrobiculatum.
floribundum, *Desv. Opusc.* 58.—Ind. occ.
floridanum, *Michx. Fl. Bor. Am.* i. 44.—Am. bor.
fluitans, *Kunth, Rév. Gram.* i. 24.—Am. bor.
foliosum, *Kunth, l. c.* 25.—Cuba.
Forsterianum, *Fluegge, Gram. Monog.* 172.—N. Caled.
foveolatum, *Steud. Syn. Pl. Gram.* 22.—Guian.
fragile, *Steud. l. c.* 17.—Venezuela.
Frankii, *Steud. l. c.* 19.—Am. bor.

PASPALUM:—
frumentaceum, Rottl. ex Roem. & Schult. Syst. ii. 296 = scrobiculatum.
furcatum, Cham. & Schlecht. in Linnaea, vi. (1831) 32 = obtusifolium.
furcatum, Fluegge, Gram. Monog. 114 = distichum.
furcatum, Spreng. ex Steud. Nom. ed. II. ii. 271 = Panicum hybridum.
fuscatum, Spreng. ex Steud. Nom. ed. II. ii. 257, in syn. = praec.
fuscescens, *J. & C. Presl, Rel. Haenk.* i. 213.—Peruv.
fuscum, *J. & C. Presl, l. c.* 214.—Am. austr.
Gardnerianum, *Nees, in Hook. Kew Journ.* ii. (1850) 103.—Bras.
Gayanum, *E. Desv. in C. Gay, Fl. Chil.* vi. 240.—Chili.
geminiflorum, Steud. Syn. Pl. Gram. 25 = undulatum.
geniculatum, Rafin. Fl. Ludov. 15 (Quid ?).—Am. bor.
geniculatum, Steud. Syn. Pl. Gram. 18 = pusillum.
germanicum, Baumg. Enum. Stirp. Transs. iii. 277 = Setaria italica.
glabrum, *Poir. Encyc.* v. 30.—Ind. occ.
globosum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Airopsis globosa.
gnaphalioideum, *C. Muell. in Bot. Zeit.* xix. (1861) 332.—Guian.
gracile, Le Conte, in Journ. Phys. xci. (1820) 284 = tenue.
gracile, Poir. Encyc. Suppl. iv. 313 = Poiretii.
gracile, Rudge, Pl. Guian. Ic. et Desc. 20. t. 26 = repens.
gracile, *Schlecht. in Linnaea*, xxvi. (1853–55) 134.—N. Granat.
granulare, *Trin. in Spreng. Neue Entdeck.* ii. 47 (sp. dub.).—Ind. or.
Griffithianum, *C. Muell. in Bot. Zeit.* xix. (1861) 325.—Ind. or.
guadaloupense, *Steud. Syn. Pl. Gram.* 18.—Ind. occ.
guineense, Steud. l. c. = paniculatum.
guttatum, *Trin. Gram. Panic.* 91.—Bras.
haemisphericum, Poir. Encyc. v. 31 = paniculatum.
Haenkeanum, Hort. Vrat. ex Nees, Cat. Sem. Hort. Vrat. (1850).—Cf. Linnaea, xxiv. (1851) 236 = Panicum Pseudopaspalus.
Haenkeanum, *J. & C. Presl, Rel. Haenk.* i. 210.—Peruv.
Hartwegianum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 479.—Mexic.
hemicryptum, *Wright, in Sauv. Fl. Cub.* 196.—Cuba.
heterophyllum, *Desv. ex Poir. Encyc. Suppl.* iv. 315.—Ins. S. Doming.
heteropodium, *Steud. Syn. Pl. Gram.* 19.—Ins. Maurit.
heterotrichum, *Trin. Sp. Gram. Ic.* iii. t. 285.—Bras.
hirsutum, Poir. Encyc. v. 28 = conjugatum.
hirsutum, Retz. Obs. ii. 7 = scrobiculatum.
hirtigluma, *Steud. Syn. Pl. Gram.* 18.—Guian.
hirtum, *H. B. & K. Nov. Gen. et Sp.* i. 89.—N. Granat.
holosericeum, Blume, ex Griseb. in Ledeb. Fl. Ross. iv. 467 = Eriochloa succincta.
horticola, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 60 = P. Falcula.
Houttuynii, H. C. Hall, ex De Vriese, Pl. Ind. Bat. Reinw. 113 = scrobiculatum.
Humboldtianum, *Fluegge, Gram. Monog.* 67.—Am. trop.
humifusum, Heyne, in Wall. Cat. n. 8755 = Panicum glabrum.
humifusum, *Poir. Encyc. Suppl.* iv. 316 = undulatum.
humile, Steud. Syn. Pl. Gram. 25 = dissectum.
hyalinum, *Nees, ex Trin. Gram. Panic.* 103.—Bras.
immersum, Nees, Agrost. Bras. 82 = aureum.
inaequale, Link, Enum. Hort. Berol. i. 103 = Panicum sanguinale.
inaequivalve, *Raddi, Agrost. Bras.* 28.—Bras.
incertum, *Roem. & Schult. Syst.* ii. 308.—Hab. ?
incomptum, Nees, ex Steud. Nom. ed. II. ii. 271 (an seq. ?).—Bras.
incomtum, Kunth, ex Doell, in Mart. Fl. Bras. ii. II. 68 = obscurum.
infirmum, Roem. & Schult. Syst. ii. 307 = setaceum ?
inflatum, *A. Rich. Fl. Cub. Fanerog.* ii. 298.—Cuba.
interruptum, Wood, Class Book (1861), 783 = Panicum floridanum.
iridifolium, *Poepp. Reise*, ii. 324.—Chili.
ischnocaulon, *Trin. Sp. Gram. Ic.* ii. t. 126.—As. trop.
Jardini, *Steud. Syn. Pl. Gram.* 18.—Afr. trop.

PASPALUM:—
jubatum, *Griseb. in Goett. Nachr.* (1868) 84.—Reg. Himal.
Kappleri, Hochst. ex Doell, in Mart. Fl. Bras. ii. II. 96 = carinatum.
Karwinskii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 479.—Mexic.
Kegelii, *C. Muell. in Bot. Zeit.* xix. (1861) 324.—Guian.
Kleinianum, J. & C. Presl, Rel. Haenk. i. 209 = distichum.
koleopodum, *Steud. Syn. Pl. Gram.* 18.—Ins. Guadeloupe.
Kora, Forst. ex Spreng. Syst. i. 244 = distichum.
Kora, Willd. Sp. Pl. i. 332; Hort. Berol. t. 74 = scrobiculatum.
lachneum, *Nees, ex Steud. Syn. Pl. Gram.* 26.—Bras.
lachneum, Trin. ex Doell, in Mart. Fl. Bras. ii. II. 77 = undulatum.
laeve, *Michx. Fl. Bor. Am.* i. 44.—Am. bor.
laevigatum, Bosc, ex Poir. Encyc. Suppl. iv. 313 = floridanum.
Lagascae, Roem. & Schult. Syst. ii. 317 = quadrifarium.
lanatum, H. B. & K. Nov. Gen. et Sp. i. 94. t. 29 = Anthaenantia lanata.
lanatum, Spreng. Syst. iv. Cur. Post. 30 = dilatatum.
lanceolatum, Mikan, ex Trin. in Spreng. Neue Entdeck. ii. 48 = laxum.
lanceaefolium, *Desv. Opusc.* 58.—Hispan.
lanciflorum, *Trin. Sp. Gram. Ic.* iii. t. 286.—Bras.
lanuginosum, *Nees, Agrost. Bras.* 63.—Bras.
lanuginosum, Willd. ex Steud. Nom. ed. II. ii. 271 = lentiferum.
lasiogonum, *Link, Hort. Berol.* i. 48.—Hab. ?
laticulmum, Spreng. Syst. i. 245 = platycaule.
latifolium, *Le Conte, in Journ. Phys.* xci. (1820) 284.—Bras.
latifolium, Spreng. Syst. i. 248 = virgatum.
laxiflorum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 148.—Bras.
laxum, *Lam. Illustr.* i. 176.—Am. austr.
laxum, Rich. ex Steud. Syn. Pl. Gram. 17 = Richardi.
Leconteanum, *Schult. Mant.* ii. 168.—Am. bor.
Lenormandi, *Husnot, in Bull. Soc. Linn. Normand.* Sér. II. v. (1870) 259.—Ind. occ.
lenticulare, *H. B. & K. Nov. Gen. et Sp.* i. 92.—Mexic.
lentiferum, *Lam. Illustr.* i. 175.—Am. bor.
lentiferum, Munro, ex Benth. Pl. Hartw. 346 = Hartwegianum.
lentiginosum, *J. & C. Presl, Rel. Haenk.* i. 218.—Mexic.
lepton, Schult. Mant. ii. 173 = tenue.
leptostachyum, DC. Cat. Hort. Monsp. (1813) 130 = incertum.
leptostachyum, *Fluegge, Gram. Monog.* 123.—Venezuela.
leptostachyum, H. B. & K. Nov. Gen. et Sp. i. 90 = scoparium.
leptostachyum, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 108 = barbigerum.
leucocheilum, *Wright, in Sauv. Fl. Cub.* 194.—Cuba.
leucotrichum, Steud. Syn. Pl. Gram. 16 = falcatum.
Liebmannii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 479.—Mexic.
ligulare, *Nees, Agrost. Bras.* 60.—Bras.
Lindenianum, *A. Rich. Fl. Cub. Fanerog.* ii. 299.—Cuba.
lineare, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 479.—Mexic.
lineare, Sw. ex Steud. Nom. ed. II. ii. 272 = caespitosum.
lineare, Trin. Gram. Panic. 99 = Neesii.
litorale, Spreng. Syst. i. 244 = seq.
littorale, R. Br. Prod. 188 = distichum.
lividum, *Trin. ex Schlecht. in Linnaea*, xxvi. (1853–55) 383.—Mexic.
loliaceum, Willd. ex Steud. Nom. ed. II. ii. 272 = Microchloa setacea.
longepedunculatum, Le Conte, in Journ. Phys. xci. (1820) 284 = arenarium.
longiflorum, Raddi, Agrost. Bras. 25 = platycaule.
longiflorum, Retz. Obs. iv. 15; ex Roxb. Fl. Ind. i. 279 = distichum.
longiflorum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 307 = Ichnanthes longiflorus.
longifolium, Roxb. Hort. Beng. 7; Fl. Ind. i. 280 = scrobiculatum.

PASPALUM:—
longifolium, *Steud. Syn. Pl. Gram.* 21.—Jamaic.
longispica, *Doell, in Mart. Fl. Bras.* ii. II. 105.—Am. trop.
longissimum, Hochst. ex Steud. Syn. Pl. Gram. 19 = conjugatum.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* i. 92.—N. Granat.
macropodium, Steud. Syn. Pl. Gram. 19 = platycaule.
macrospermum, Fluegge, Gram. Monog. 172 = floridanum.
maculosum, Trin. Gram. Panic. 98; Sp. Gram. Ic. iii. t. 113 = distichum.
malacophyllum, *Trin. l. c.* t. 271.—Bras.
mandiocanum, *Trin. Gram. Panic.* 113.—Panama; Bras.
marginatum, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 348.—Bolivia.
marginatum, Spreng. ex Steud. Nom. ed. II. ii. 272 = undulatum.
marginatum, *Trin. Gram. Panic.* 90.—Bras.
maritimum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 148.—Bras.
mauritanicum, Nees, ex Steud. Syn. Pl. Gram. 26 = scrobiculatum.
melanospermum, *Desv. ex Poir. Encyc. Suppl.* iv. 315.—Guian.
membranaceum, *Lam. Illustr.* i. 177.—Peruv.
membranaceum, Walt. Fl. Carol. 75 = Walterianum.
metabolon, Steud. Syn. Pl. Gram. 19 = scrobiculatum.
Metzii, Steud. l. c. 21 = scrobiculatum.
Meyerianum, Spreng. Syst. i. 248 = quadrifarium, Eriochloa Kunthii.
Michauxianum, Kunth, Rév. Gram. i. 25 = distichum.
micranthum, *Desv. Opusc.* 56.—Ins. Mascar.
micranthum, *Doell, in Mart. Fl. Bras.* ii. II. 51.—Am. trop.
microstachyum, *J. & C. Presl, Rel. Haenk.* i. 215.—Hab. ?
miliare, Spreng. Syst. i. 247 = milioideum.
miliaria, *C. Muell. in Bot. Zeit.* xix. (1861) 332.—Ind. or.
milioideum, *Desv. ex Poir. Encyc. Suppl.* iv. 315.—Ind. occ.
millegranum, Schrad. ex Schult. Mant. ii. 175 = densum.
minimum, *Meyen, Reise*, i. 484, nomen.—Peruv.
minus, *Fourn. ex Hemsl. Biol. Centr. Bot.* iii. 480.—Mexic.
minutiflorum, *Desv. Opusc.* 57.—Bras.
minutiflorum, Steud. Syn. Pl. Gram. 17 = brevifolium.
minutum, *Trin. in Linnaea*, x. (1836) 293.—Peruv.
molle, *Poir. Encyc.* v. 34.—Ins. S. Thomas.
mollicomum, Kunth, Enum. Pl. i. 47 = molle.
mollipilum, Steud. Syn. Pl. Gram. 20 = scrobiculatum.
mononeurum, *Steud. l. c.* 24.—Mexic.
monostachyum, Hort. ex Walp. Ann. iii. 714 = Panicum Pseudopaspalus.
monostachyum, *Vasey, ex Chapm. Fl. S. U. St. Suppl.* 665.—Florida.
monostachyum, Willd. ex Steud. Nom. ed. II. ii. 272 = Panicum monostachyum.
montevidense, Spreng. Syst. i. 246 = undulatum.
mucronatum, Muhl. Cat. ex Doell, in Mart. Fl. Bras. ii. II. 98 = repens.
mucronatum, *Muhl. Desc. Gram.* 96.—Am. bor.
multicaule, *Poir. Encyc. Suppl.* iv. 309.—Bras.
multiflorum, *Desv. Opusc.* 58.—Bras.
multiflorum, *Doell, in Mart. Fl. Bras.* ii. II. 90.—Am. trop.
multispica, Steud. Syn. Pl. Gram. 18 = paniculatum.
nanum, *Wright, ex Griseb. Cat. Pl. Cub.* 230.—Cuba.
natans, Le Conte, in Journ. Phys. xci. (1820) 285 = mucronatum.
Neesii, *Kunth, Rév. Gram.* i. 25.—Bras.
nematodes, Schult. Mant. ii. 71 = brevifolium.
nepalense, Spreng. ex Steud. Nom. ed. II. ii. 272 = Panicum plicatum.
notatum, Fluegge, Gram. Monog. 106 = distichum.
nutans, Lam. Illustr. i. 175 = conjugatum.
oajacense, Hemsl. Biol. Centr. Am. Bot. iii. 499 = seq.
oaxacense, Steud. Syn. Pl. Gram. 21 = Panicum oaxacense.
obscurum, *Nees, ex Steud. l. c.* 30.—Am. trop.
obtusatum, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 103 = distichum.
obtusifolium, *Raddi, Agrost. Bras.* 23.—Mexic.
obtusifolium, Trin. Sp. Gram. i. t. 115 = distichum.

PASPALUM:—
oligostachyum, Salzm. ex Steud. Syn. Pl. Gram. 23 = undulatum.
orbiculare, Forst. f. Prod. 7 = scrobiculatum.
orbiculatum, Poir. Encyc. v. 32 = pusillum.
orizaeforme, Steud. Syn. Pl. Gram. 27 = scoparium.
orthos, Kunth, Enum. Pl. i. 57, in syn. = tenue.
ovale, *Nees, ex Steud. Syn. Pl. Gram.* 22.—Bras.
ovatum, Nees, ex Trin. Gram. Panic. 113 = dilatatum.
oxyanthum, Steud. Syn. Pl. Gram. 27 = Panicum sanguinale.
pachyrrhizum, *Steud. l. c.* 23.—Chili.
pallidum, *H. B. & K. Nov. Gen. et Sp.* i. 88.—Ecuador.
paniculatum, *Linn. Syst.* ed. X. 855.—Am. austr.
paniculatum, Poir. Encyc. v. 32, partim = quadrifarium.
paniculatum, Walt. Fl. Carol. 75 = repens.
papillosum, Spreng. Nov. Prov. 47 = multicaule.
paradisiacum, *Steud. Syn. Pl. Gram.* 23.—Chili.
parviflorum, Desv. Opusc. 58 = heterophyllum.
parviflorum, *Rhodé, ex Fluegge, Gram. Monog.* 98.—Am. austr.
patulum, Hornem. Hort. Hafn. i. 78 = Panicum parviflorum.
pauperculum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 481.—Mexic.
pectinatum, *Nees, Agrost. Bras.* 34.—Bras.
pedicellare, *Trin. ex Steud. Nom.* ed. II. ii. 272.—Ind. or.
pedunculare, *J. & C. Presl, Rel. Haenk.* i. 217.—Hab. ?
pedunculatum, Desv. in Poir. Encyc. Suppl. iv. 315 = Panicum decumbens.
pellitum, *Nees, ex Trin. Gram. Panic.* 89.—Bras.
pellitum, Willd. ex Steud. Nom. ed. II. ii. 272 = pallidum.
penicillatum, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 171.—Ins. Galap.
pilosum, Lam. Illustr. i. 175 = Panicum monostachyum.
pilosum, Roth, Catalect. fasc. iii. 12 = parviflorum.
pilosum, Spreng. ex Steud. Nom. ed. II. ii. 272 = Panicum thrasioides.
planifolium, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 481.—Mexic.
plantagineum, Nees, Agrost. Bras. 69 = densiflorum, laxum.
platense, Spreng. Syst. i. 247 = dilatatum.
platycaule, *Poir. Encyc.* v. 34 (platicaule).—Afr. et Am. trop.
platycaule, Willd. ex Steud. Nom. ed. II. ii. 272 = distichum.
platyculmum, Thou. ex Nees, Agrost. Bras. 24 = platycaule.
platyphyllum, *Griseb. Cat. Pl. Cub.* 230.—Cuba.
platyphyllum, Schult. Mant. ii. 557 = virgatum.
pleostachyum, *Doell, in Mart. Fl. Bras.* ii. II. 58.—Am. trop.
plicatulum, Michx. Fl. Bor. Am. i. 45 = undulatum.
plicatum, Pers. Syn. i. 86 = praec.
pluriracemosum, *Steud. Syn. Pl. Gram.* 27.—Ins. Phillipp.
Poeppigianum, *Trin. ex Steud. Nom.* ed. II. ii. 272.—Chili.
Poiretii, *Roem. & Schult. Syst.* ii. 878.—Ins. S. Doming.
polydactylon, Steud. Syn. Pl. Gram. 19 = scoparium.
polyphyllum, Nees, ex Trin. Gram. Panic. 114 = distichophyllum.
polystachyum, R. Br. Prod. 188 = scrobiculatum.
polystachyum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Eriochloa polystachya.
polystachyum, Salzm. ex Steud. Syn. Pl. Gram. 18 = paniculatum.
praecox, *Walt. Fl. Carol.* 75.—Am. bor.
Preslii, *Kunth, Enum. Pl.* i. 47.—Peruv.
pressum, Nees, ex Steud. Syn. Pl. Gram. 23 = scoparium.
protensum, Trin. Gram. Panic. 108 = conjugatum.
pruinosum, Trin. Sp. Gram. Ic. iii. t. 272 = coryphaeum.
Pseudo-duroa, Nees, ex Baker, Fl. Maurit. 431 = longiflorum.
puberulum, Roem. & Schult. Syst. ii. 316 = scrobiculatum.
pubescens, R. Br. Prod. 188 = scrobiculatum.
pubescens, Lag. Gen. et Sp. Nov. 2. n. 21 = quadrifarium.
pubescens, Muhl. ex Willd. Enum. Hort. Berol. 89 = conjugatum.

PASPALUM:—
pubescens, J. & C. Presl, Rel. Haenk. i. 214 = Preslii.
pubescens, Sm. in Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 37 = granulare.
pubiflorum, *Rupr. ex Galeotti, in Bull. Acad. Brux.* ix. (1842) 237.—Mexic.
pubifolium, *J. & C. Presl, Rel. Haenk.* i. 219.—Peruv.
pulchellum, *H. B. & K. Nov. Gen. et Sp.* i. 90.—Am. austr.
pulchellum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Coelachne pulchella.
pulchrum, *Nees, Agrost. Bras.* 79.—Bras.
pumilum, *Nees, l. c.* 52.—Bras.
punctatum, Fluegge, Gram. Monog. 127 = Eriochloa polystachya.
punctulatum, *Bertol. Misc. Bot.* xi. 13.—Am. bor.?
purpurascens, Ell. Sketch, i. 108 = undulatum.
purpureum, Ruiz & Pav. Fl. Per. i. 47 = racemosum.
pusillum, *Vent. ex Fluegge, Gram. Monog.* 100.—Am. austr.
pyramidale, Nees, Agrost. Bras. 77 = repens.
quadrifarium, *Lam. Illustr.* i. 176.—Reg. Argent.
quadrifarium, Nees, Agrost. Bras. 71 = denticulatum.
quadrifarium, Trin. Sp. Gram. Ic. ii. t. 128 = falcatum.
racemosum, *Lam. Illustr.* i. 176.—Bras.
racemosum, Nutt. in Trans. Am. Phil. Soc. v. (1837) 145 = Panicum floridanum.
racemulosum, Chapm. Fl. S. U. St. 571 = Panicum floridanum.
radiatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 195, in syn.—Bras.
ramosissimum, Nees, Agrost. Bras. 80 = pulchrum, chrysites.
rectum, *Nees, in Hook. Kew Journ.* ii. (1850) 104.—Bras.
reduncum, *Nees, ex Steud. Syn. Pl. Gram.* 31.—Bras.
reimarioides, Brongn. Bot. Voy. Coquille, t. 20 = Panicum sanguinale.
reimarioides, *Chapm. Fl. S. U. St. Suppl.* 665.—Florida.
remotum, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 349.—Bolivia.
Remyi, *Steud. Syn. Pl. Gram.* 28.—Bolivia.
Renggeri, Steud. l. c. 17 = conjugatum.
repens, *Berg. in Act. Helvet.* vii. (1772) 129. t. 7.—Am. calid.
repens, Frank, ex Steud. Syn. Pl. Gram. 19 = Frankii.
reptans, Poir. ex Doell, in Mart. Fl. Bras. ii. II. 75 = distichum.
rhizomatosum, *Steud. Syn. Pl. Gram.* 17.—Ins. Guadalup.
Richardi, *Steud. l. c.*—Ind. occ.
riparium, *Nees, Agrost. Bras.* 56.—Bras.
robustum, Link, ex Steud. Nom. ed. II. ii. 273 = tenellum.
Rodschiedianum, G. F. W. Mey. ex Steud. l. c. (Quid ?).—Guian.
rottboellioides, *Wright, in Sauv. Fl. Cub.* 195.—Cuba.
Royleanum, Nees, ex Thw. Enum. Pl. Zeyl. 358 = ischnocaulon.
rudimentosum, *Steud. Syn. Pl. Gram.* 24.—Mexic.
rufum, *Nees, ex Steud. l. c.* 26.—Bras.
rupestre, *Trin. in Linnaea*, x. (1836) 293.—Cuba.
saccharoides, *Nees, ex Trin. Spec. Gram. Ic.* i. t. 107.—Panama.
Salzmanni, *Doell, in Mart. Fl. Bras.* ii. II. 49.—Bras.
sanguinale, Lam. Illustr. i. 176 = Panicum sanguinale.
sanguinale, Schult. Mant. ii. 291 = Panicum javanicum.
sanguinolentum, *Trin. Gram. Panic.* 116.—Bras.
savannarum, *Schlecht. in Linnaea*, xxvi. (1853-55) 132.—N. Granat.
saxatile, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 76 = undulatum.
scalare, *Trin. Sp. Gram. Ic.* iii. t. 274.—Bras.
Schaffneri, *Griseb. ex Hemsl. Biol. Centr. Am. Bot.* iii. 482.—Mexic.
sciaphilum, Steud. Syn. Pl. Gram. 18 = molle.
scoparium, *Fluegge, Gram. Monog.* 124.—Am. austr.
scorbiculatum, Steud. Nom. ed. I. 594, sphalm. = seq.
scrobiculatum, *Linn. Mant.* i. 29.—Geront. trop.
scutatum, *Nees, ex Trin. Gram. Panic.* 105.—Bras.
Selloi, Spreng. ex Nees, Agrost. Bras. 43 = dilatatum.
semirugosum, *Nees, ex Steud. Nom.* ed. II. ii. 273.—Bras.
senescens, *Doell, in Mart. Fl. Bras.* ii. II. 119.—Am. trop.

PASPALUM:—
sericeum, *Scheele, in Linnaea*, xxii. (1849) 341.—Texas.
serotinum, *Fluegge, Gram. Monog.* 145.—Carolina.
serpens, *Presl, ex Trin. Gram. Panic.* 102; *Nees, Agrost. Bras.* 50.—Bras.
serpentinum, Hochst. ex Steud. Syn. Pl. Gram. 22 = distichum.
sesquiglume, Benth. in Journ. Linn. Soc. xix. (1881) 36 = subsesquiglume.
setaceum, *Michx. Fl. Bor. Am.* i. 43.—Am. bor.
setaceum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Microchloa setacea.
setifolium, Doell, in Mart. Fl. Bras. ii. II. 61 = Kegelii.
setifolium, Nees, ex Steud. Nom. ed. II. ii. 273 (Quid ?).—Hab. ?
siccum, *Nees, Agrost. Bras.* 28.—Bras.
Sieberianum, *Steud. Syn. Pl. Gram.* 17.—Ind. occ.
singulare, *Link, Hort. Berol.* i. 48.—Bras.
sinuosum, *Desv. Opusc.* 57.—Ind. occ.
spathaceum, *Desv. ex Poir. Encyc. Suppl.* iv. 314.—Ins. S. Thomas.
squamatum, *Steud. Syn. Pl. Gram.* 21.—Afr. trop.
squamulatum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 482.—Mexic.
stellatum, *Fluegge, Gram. Monog.* 62.—Am. austr.
stellatum, Trin. Gram. Panic. 100 = carinatum.
stoloniferum, Bosc, in Trans. Linn. Soc. ii. (1794) 83 = racemosum.
stoloniferum, *E. Desv. in C. Gay, Fl. Chil.* vi. 241.—Chili.
strictum, Brot. Phyt. Lusit. 13. t. 5 = Spartina stricta.
strictum, Pers. Syn. i. 86 = paniculatum.
strictum, Spreng. ex Steud. Nom. ed. II. ii. 273 = mandiocanum.
subarticulatum, Beauv. ex Steud. l. c. ed. I. 594 (Quid ?).—Hab. ?
subsesquiglume, *Doell, in Mart. Fl. Bras.* ii. II. 43.—Am. trop.
succinctum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 131 = Eriochloa succincta.
suffultum, Mikan, ex Trin. in Spreng. Neue Entdeck. ii. 46 = scoparium.
sumatrense, Roth, Nov. Pl. Sp. 35 = scrobiculatum.
Sumichrastii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 492.—Mexic.
superbum, Spreng. Syst. i. 248 = Chloris distichophylla.
supinum, Bosc, ex Poir. Encyc. v. 29 = conjugatum.
supinum, Hornem. Hort. Hafn. i. 77 = dubium.
supinum, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. (1842) 237 = paniculatum.
supinum, Sieber, ex Baker, Fl. Maurit. 431 = heteropodium.
surinamense, *Hochst. ex Benth. in Journ. Linn. Soc.* xix. (1881) 35.—Guian.
Swartzianum, Fluegge, Gram. Monog. 96 = caespitosum.
taphrophyllum, *Steud. Syn. Pl. Gram.* 19.—Ins. Martinic.
tectum, *Steud. l. c.* 29.—Florida.
tenax, *Trin. Gram. Panic.* 122.—N. Zel.
tenellum, *Willd. Enum. Hort. Berol.* 89.—Mexic.
tenerum, Nees, Agrost. Bras. 32 = dissitiflorum.
tenue, Gaertn. Fruct. ii. 2. t. 80 = conjugatum.
tenue, *Kunth, Rév. Gram.* i. 26.—Am. bor.
tenue, Willd. ex Steud. Nom. ed. II. ii. 273 = platycaule.
Thouarsianum, *Fluegge, Gram. Monog.* 148.—Madag.
Thunbergii, Kunth, ex Steud. Syn. Pl. Gram. 28 = scrobiculatum.
tomentosum, *Poir. Encyc. Suppl.* iv. 308.—Hab. ?
trachycaleon, *Steud. Syn. Pl. Gram.* 28.—Am. trop.
tridentatum, Trin. Gram. Panic. 119 (= *Helopus tridentatus*).—Am. austr.
triglume, *Steud. Syn. Pl. Gram.* 27.—Mexic.
tristachyum, Lam. Illustr. i. 176 = platycaule.
tristachyum, Le Conte, in Journ. Phys. xci. (1820) 285 = distichum.
tristachyum, Willd. ex Steud. Nom. ed. II. ii. 273 = pulchellum.
tropicum, *Doell, in Mart. Fl. Bras.* ii. II. 83.—Am. trop.
umbellatum, Lam. Illustr. i. 177 = Panicum Dactylon.
umbrosum, Salzm. ex Doell, in Mart. Fl. Bras. ii. II. 50 = molle.

PASPALUM:—
umbrosum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 153.—Bras.
undulatum, *Poir. Encyc.* v. 29.—Am. bor. et austr.
undulatum, Spreng. Fl. Hal. Mant. 30 = Forsterianum.
undulosum, Le Conte, in Journ. Phys. xci. (1820) 284 = Leconteanum.
uniflorum, *Steud. Nom.* ed. II. ii. 273.—Bras.
uniseriatum, *Steud. in Lechl. Berb. Am. Austr.* 55.—Peruv.
Urvillei, *Steud. Syn. Pl. Gram.* 24.—Hab. ?
vaginatum, Ell. Sketch, i. 409 = Walterianum.
vaginatum, Sw. Prod. Veg. Ind. Occ. 21 = distichum.
vaginiflorum, Steud. Syn. Pl. Gram. 19 = Panicum decumbens.
variegatum, Link, Hort. Berol. i. 50 = quadrifarium.
velutinum, *Kunth, Rév. Gram.* i. 27.—Mexic.
venustum, Forst. f. ex Roem. & Schult. Syst. ii. 297 = scrobiculatum.
vestitum, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 45 = parviflorum.
villifolium, Steud. Syn. Pl. Gram. 20 = ancylocarpum.
villosum, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 311 = Panicum flavidum.
villosum, Thunb. Fl. Jap. 45 = Eriochloa villosa.
villosum, Willd. ex Steud. Nom. ed. II. ii. 273 = Panicum filiforme.
virgatum, Cham. & Schlecht. in Linnaea, vi. (1831) 32 = lentiginosum.
virgatum, Le Conte, in Journ. Phys. xci. (1820) 285 = undulatum.
virgatum, *Linn. Syst.* ed. X. 855.—Am. austr.
Virletii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 483.—Mexic.
vulnerans, Salzm. ex Steud. Nom. ed. II. ii. 273 = densum.
Wagnerianum, *Schlecht. in Linnaea*, xxvi. (1853–55) 133.—N. Granat.
Walteri, Chapm. Fl. S. U. St. 570, sphalm. = seq.
Walterianum, *Schult. Mant.* ii. 166.—Am. bor.
Zollingeri, Steud. Syn. Pl. Gram. 28 = scrobiculatum.

PASPALUS, Fluegge, Gram. Monog. 53 (1810) = **Paspalum**, Linn. (Gramin.).

PASSAEA, Adans. Fam. ii. 509 (1763) = **Ononis**, Linn. (Legumin.).

PASSAEA, Baill. Etud. Gén. Euph. 507. t. 18 (1858) = **Bernardia**, Houst. (Euphorb.).
spartioides, Baill. l. c. 508 = B. spartioides.

PASSALIA, Soland. ex R. Br. in Tuckey, Congo, 440 (1818) = **Alsodeia**, Thou. (Violar.).

PASSAVERIA, Mart. & Eichl. ex Miq. in Mart. Fl. Bras. vii. 85. t. 38; et t. 47. f. 1, 3 (1863) = **Ecclinusa**, Mart. (Sapot.).
lanceolata, Mart. & Eichl. l. c. 86.—Bras.
lancifolia, Mart. & Eichl. l. c.—Bras.
obovata, Mart. & Eichl. l. c. 87.—Bras.

PASSERINA, Linn. Gen. ed. I. 341 (1737). *THYMELAEACEAE*, Benth. & Hook. f. iii. 194.
BALENDASIA, STEIROCTIS, TRIMEIANDRA, Rafin. Fl. Tellur. iv. 105 (1836).
Ammodendron, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 444 = Stellera Lessertii.
annua, Auch. ex Meissn. in DC. Prod. xiv. 552 = Thymelaea Aucheri.
annua, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 320 = Thymelaea arvensis.
anthylloides, Linn. f. Suppl. 226 = Lasiosiphon anthylloides.
arvensis, Ball, in Journ. Linn. Soc. xvi. (1878) 653 = Thymelaea arvensis.
axillaris, Thunb. Mus. Nat. Acad. Upsal. xiii. 106, nomen = Pimelea virgata.
bruniades, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 579 = Lachnaea penicillata.
brunioides, Eckl. & Zeyh. l. c. 563, 579 = praec.
calocephala, Meissn. in Linnaea, xiv. (1840) 393 = Arthrosolen calocephalus.
calycina, DC. Fl. Fr. iii. 360 = Thymelaea calycina.
campanulata, E. Mey. ex Presl, Bot. Bemerk. 108 = Cryptadenia breviflora.
canescens, Schousb. Vext. Marok. 190 = Thymelaea canescens.

PASSERINA:—
cantabrica, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 299 = Thymelaea coridifolia.
capitata, Linn. Pl. Rar. Afr. 11; Amoen. Acad. vi. 88 = Lachnaea capitata.
cephalophora, Thunb. Prod. Pl. Cap. 75 = Lasiosiphon eriocephalus.
Chamaedaphne, Bunge, Enum. Pl. Chin. Bor. 58 = Wikstroemia canescens.
Chamaejasme, Fisch. ex Meissn. in DC. Prod. xiv. 549 = Stellera Chamaejasme.
ciliata, Linn. Sp. Pl. 559 = Struthiola lucens ?
ciliata, Thunb. Fl. Cap. ex Schult. 377 = Cryptadenia ciliata.
concinna, Walp. Ann. i. 583 = Thymelaea arvensis.
conglomerata, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 322 = ericoides.
coridifolia, Wikstr. l. c. 334 = Thymelaea coridifolia.
corymbosa, Eckl. ex Meissn. in DC. Prod. xiv. 562 = filiformis.
costata, Griff. Notul. iv. 367 = Diarthron vesiculosum.
cupressina, *Wendl. ex Bartl. in Linnaea*, xiv. (1840) 404.—Afr. austr.
cupressoides, Steud. Nom. ed. II. ii. 273 = praec.
diarthronoides, Griff. Notul. iv. 365 = Thymelaea arvensis.
dichotoma, Steud. Nom. ed. II. ii. 273 = Stellera Chamaejasme.
dioica, Ramond, in Bull. Soc. Philom. ii. (1800) 139 = Thymelaea dioica.
dodecandra, Linn. Cent. Pl. i. 10 = Struthiola erecta.
elliptica, Boiss. Voy. Bot. Espagne, ii. 556. t. 158 = Thymelaea elliptica.
empetrifolia, Lapeyr. Hist. Abr. Pl. Pyr. 212 = Thymelaea dioica.
ericoides, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 563 = rigida.
ericoides, *Linn. Mant.* i. 236.—Afr. austr.
eriocephala, Thunb. Prod. Fl. Cap. 75 = Lachnaea globulifera.
filiformis, *Linn. Sp. Pl.* 559.—Afr. austr.
filiformis, Mill. Gard. Dict. ed. VIII. n. 1 = Struthiola erecta.
Ganpi, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 201 = Wikstroemia canescens.
globosa, Lam. Illustr. ii. 431 = Lachnaea alpina.
glomerata, Thunb. Prod. Pl. Cap. 75 = ericoides.
Gnidia, Linn. f. Suppl. 226 = Pimelea Gnidia.
grandiflora, Curt. Bot. Mag. t. 292 = Cryptadenia breviflora.
grandiflora, Linn. f. Suppl. 226 = Cryptadenia grandiflora.
gymnostachya, Meissn. in Linnaea, xiv. (1840) 397 = Arthrosolen gymnostachys.
hirsuta, Asso, Stirp. Arag. 49 = Thymelaea tinctoria.
hirsuta, *Linn. Sp. Pl.* 559.—Reg. Mediterr.
incana, Pourr. ex Willk. & Lange, Prod. Pl. Hisp. i. 300 = Thymelaea velutina.
inconspicua, Meissn. in Linnaea, xiv. (1840) 397 = Arthrosolen inconspicuus.
incurva, *Wendl. ex Bartl. l. c.* 404.—Afr. austr.
involucrata, Spreng. ex Meissn. in DC. Prod. xiv. 595 = Lasiosiphon linifolius.
involucrata, Thunb. Mus. Nat. Acad. Upsal. xiii. 106 = Pimelia linifolia.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 200 = Wikstroemia japonica.
javanica, Thunb. Fl. Javan. ii. 19 (Quid ?).—Java.
juniperifolia, Lapeyr. Hist. Abr. Pl. Pyr. 213 = Thymelaea tinctoria, calycina.
kalifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 301 = Thymelaea canescens.
laevigata, Linn. Cent. Pl. ii. 15 = Gnidia oppositifolia.
lateriflora, Hort. ex Wikstr. in Vet. Acad. Handl. Stockh. (1818) 347 = Arthrosolen spicatus.
laxa, Linn. f. Suppl. 226 = Arthrosolen laxus.
Lessertii, Wikstr. in Vet. Akad. Handl. Stockh. (1818) 341 = Stellera Lessertii.
linariaefolia, Pourr. ex Wikstr. l. c. 333 = Thymelaea thesioides.
linearifolia, Wikstr. l. c. 343 = Gnidia linoides.
linearis, Wendl. ex Bartl. in Linnaea, xiv. (1840) 404 (Quid ?).—Afr. austr.
linoides, Thunb. Prod. Pl. Cap. 75 = Gnidia linoides.
longiflora, Thunb. l. c. 189 = Pimelea longifolia.
longifolia, Thunb. Mus. Nat. Acad. Upsal. xiii. 106, nomen = Pimelia longifolia.
Matnak, J. F. Gmel. Syst. 1597, sphalm. = seq.
Metkan, J. F. Gmel. l. c. 634, sphalm. = seq.

PASSERINA:—
Metnan, Forsk. Fl. Aegypt. Arab. 81 = Thymelaea hirsuta.
microphylla, Coss. & Dur. in Bull. Soc. Bot. Fr. iii. (1856) 744 = Thymelaea microphylla.
nervosa, Thunb. Prod. Pl. Cap. 75 = Lachnaea nervosa.
nervosa, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 328 = Lachnaea striata.
nitida, Desf. Fl. Atlant. i. 331. t. 94 = Thymelaea nitida.
nivalis, Ram. in Bull. Soc. Philom. ii. (1800) 131. t. 9 = Thymelaea tinctoria.
orientalis, Willd. Sp. Pl. ii. 431 = Thymelaea orientalis.
paleacea, *Wikstr. in Vet. Acad. Handl. Stockh.* (1818) 323.—Afr. austr.
pectinata, Hort. ex Lodd. Cat. (1816) 18 = filiformis.
pendula, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 563 = rigida.
pentandra, Thunb. Prod. Pl. Cap. 76 = Lonchostoma obtusiflorum.
persica, Boiss. Diagn. Ser. I. vii. 85 = Stellera Lessertii.
pilosa, Linn. f. Suppl. 226; Forst. f. Prod. 28 = Pimelea virgata.
planifolia, Burm. f. Fl. Cap. Prod. 12 = Lachnaea alpina.
polycephala, E. Mey. ex Meissn. in Linnaea, xiv. (1840) 390 = Arthrosolen polycephalus.
polygalaefolia, Lapeyr. Hist. Abr. Pl. Pyr. 214 = hirsuta.
prostrata, Linn. f. Suppl. 227 = Pimelea prostrata.
pubescens, Guss. Fl. Sic. Syn. i. 451 = Thymelaea arvensis.
pubescens, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 332 = Thymelaea virgata.
purpurea, Wikstr. l. c. 323 = Lasiosiphon eriocephalus.
racemosa, Wikstr. l. c. 320 = Stellera Chamaejasme.
rigida, *Wikstr. l. c.* 326.—Afr. austr.
salsolaefolia, Herb. Berol. ex Meissn. in DC. Prod. xiv. 664 (Bruniacea ? *vel* Phylicae sp.).—Hab. ?
salsolaefolia, Poir. Encyc. v. 41 = paleacea.
sericea, Linn. Cent. Pl. ii. 15 = Gnidia sericea.
setosa, Thunb. Prod. Pl. Cap. 75 = Gnidia setosa.
spicata, Linn. f. Suppl. 226 = Arthrosolen spicatus.
stachyoides, Meissn. in DC. Prod. xiv. 550, in syn. = Stellera stachyoides.
Stellera, Ram. ex DC. Fl. Fr. iii. 361 = Thymelaea arvensis.
Stelleri, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 321 = Stellera Chamaejasme.
striata, Lam. Illustr. ii. 430 = Lachnaea striata.
stricta, Thunb. Prod. Pl. Cap. 75 = Gnidia stricta, Wikstroemiana ?
subspicata, Meissn. in Linnaea, xiv. (1840) 395 = Wikstroemia subspicata.
Tartonraira, Schrad. Neues Journ. iv. I. (1810) 89 = Thymelaea Tartonraira.
tenuiflora, Willd. Enum. Hort. Berol. 426 = Arthrosolen laxus.
tetragona, Steud. Nom. ed. II. ii. 274 = Struthiola erecta.
thesioides, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 333 = Thymelaea thesioides.
Thomasii, Duby, Bot. Gall. i. 406 = Thymelaea Tartonraira.
Thunbergii, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 343 = Gnidia sericea.
Thymelaea, DC. Fl. Fr. Suppl. 366 = Thymelaea Sanamunda.
tinctoria, Pourr. in Mém. Acad. Toul. iii. (1788) 323 = Thymelaea tinctoria.
tingitana, Salzm. ex Meissn. in DC. Prod. xiv. 554 = Thymelaea villosa.
tomentosa, Wikstr. in Vet. Akad. Handl. Stockh. (1818) 332 = Thymelaea Tartonraira.
uniflora, Drège, ex Meissn. in DC. Prod. xiv. 574 = Cryptadenia ciliata.
uniflora, Linn. Sp. Pl. 560 = Cryptadenia uniflora.
velutina, Boiss. Elench. 81 = Thymelaea elliptica.
velutina, [Pourr.] Cambess. in Mém. Mus. Par. xiv. (1827) 304 = Thymelaea velutina.
vesiculosa, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1839) 170 = Diarthron vesiculosum.
villosa, Thunb. in Mus. Nat. Acad. Upsal. xiii. 106, nomen = Pimelea arenaria.
villosa, Wikstr. in Vet. Acad. Handl. Stockh. (1818) 332 = Thymelaea villosa.
virescens, Coss. & Dur. ex Meissn. in DC. Prod. xiv. 554 = Thymelaea virescens.
virgata, Desf. Fl. Atlant. i. 331. t. 95 = Thymelaea virgata.

PASSIFLORA, Linn. Syst. ed. I (1735). *PASSIFLOREAE*, Benth. & Hook. f. i. 810.
ANTHACTINIA, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 139.
ASTEPHANANTHES, Bory, l. c. (1819) 138.
ASTROPHEA, Reichb. Consp. 132 (1828).
BALDWINIA, Rafin. in Am. Monthly Mag. (1818) 267.
BLEPHISTELMA, Rafin. Fl. Tellur. iv. 103 (1836).
CERATOSEPALUM, Oerst. Réch. Fl. Am. Centr. 18. t. 17 (1863).
CIECA, Medic. Malv. 97 (1787).
DECALOBA, M. Roem. Syn. Pepon. 131, 152 (1846).
DISEMMA, Labill. Sert. Austro-Caled. 78. t. 79 (1824).
DISTEMMA, Lem. Hort. Van Houtte, fasc. ii. t. 11 (1847 ?).
DISTEPHANA, Juss. in Ann. Mus. Par. vi. (1805) 396.
DISTEPHANIA, Steud. Nom. ed. II. i. 521 (1840).
DISTEPHIA, Salisb. ex DC. Prod. iii. 335 (1828).
DYSOSMIA, M. Roem. Syn. Pepon. 149 (1846).
ERNDELIA, Neck. Elem. ii. 235 (1790).
GRANADILLA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 166 (1745).
MACROPHORA, Rafin. Fl. Tellur. iv. 103 (1836).
MEIOPERIS, Rafin. l. c. (1836).
MONACTINEIRMA, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 138.
MURUCUJA, Tourn. ex Medic. Malv. 97 (1787).
PENTARIA, M. Roem. Syn. Pepon. 131, 187 (1846).
PEREMIS, Rafin. Fl. Tellur. iv. 104 (1836).
PERICODIA, Rafin. l. c. (1836).
POLYANTHEA, DC. in Mém. Soc. Phys. Genèv. i. II. (1822) 435.
PSILANTHUS, Juss. in Ann. Mus. Par. vi. (1805) 396.
SYNACTILA, Rafin. Fl. Tellur. iv. 104 (1836).
TETRAPATHAEA, Reichb. Consp. 132 (1828).
TRIPSILINA, Rafin. Fl. Tellur. iv. 103 (1836).
XEROGONA, Rafin. l. c. (1836).
acerifolia, Cham. & Schlecht. in Linnaea, v. (1830) 89 = adenopoda.
acerifolia, Hort.—Cf. Darwin, Climb. Plants, ed. I. 154 = sicyoides.
actinia, Hook. Bot. Mag. t. 4009.—Bras.
aculeata, *Hort. ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 144.—Hab. ?
acuminata, DC. Prod. iii. 328 = laurifolia.
adenophylla, Mast. in Mart. Fl. Bras. xiii. i. 568 = alba.
adenopoda, [*Moç. & Sesse, ex*] *DC. Prod.* iii. 330. —Mexic.
adiantifolia, *Ker-Gawl. in Bot. Reg.* t. 233.—Ins. Norfolk.
Adiantum, Willd. Enum. Hort. Berol. 698 = glabra.
adulterina, Linn. f. Suppl. 408 = Tacsonia adulterina.
affinis, *Engelm. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 233.—Texas.
alata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 306.—Peruv.
alba, *Link & Otto, Ic. Pl. Rar. Hort. Berol.* t. 33.—Bras.
albicans, *Hort. ex Loud. Hort. Brit.* 596; *et ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 144.—Mexic.
albida, Ker-Gawl. in Bot. Reg. t. 677 = mucronata.
alnifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 136.—N. Granat.
alnifolia, Mast. in Mart. Fl. Bras. xiii. i. 549, partim = bogotensis.
amabilis, *Hook. Bot. Mag.* t. 4406.—Bras.
amabilis, Hort. ex Lem. in Fl. des Serres, iii. (1847) 209 = populifolia.
amabilis, Hort. ex Mart. in Journ. Hort. Soc. N. S. iv. (1877) 144 = racemosa.
amethystina, *Mikan, Delic. Fl. et Faun. Bras.* 20.—Bras.
Andersonii, *DC. Prod.* iii. 326.—Ind. occ.
Andreana, *Mast. in Journ. Linn. Soc.* xx. (1883) 37.—N. Granat.
anfracta, *Mast. & André, l. c.* 38.—N. Granat.
angulata, Sweet, Hort. Brit. ed. I. 354, nomen; Hort. ex Mart. in Journ. Hort. Soc. N. S. iv. (1877) 137 = alata.
angustifolia, Sw. Prod. Veg. Ind. Occ. 97 = suberosa.
antioquiensis, Karst. in Linnaea, xxx. (1859-60) 162 = Tacsonia Van-Volxemii.
appendiculata, G. F. W. Mey. Prim. Fl. Esseq. 223 = auriculata.

PASSIFLORA:—
arborea, *Spreng. Syst.* iii. 42.—N. Granat.; Ecuador.
aristulata, *Mast. in Mart. Fl. Bras.* xiii. i. 570.—Peruv.
arvensis, *Vell. Fl. Flum.* ix. t. 71.—Bras.
asperata, *Regel, Cat. Pl. Hort. Aksakov.* 104.—Hab. ?
atomaria, Planch. ex Mast. in Mart. Fl. Bras. xiii. i. 570 = alba.
atropurpurea, *Hort.—Cf. Gard. Chron.* xix. (1883) 499.—Hab. ?
aurantia, Andr. Bot. Rep. t. 295 = glabra.
aurantia, *Forst. f. Prod.* 326.—N. Caled.
auriculata, *H. B. & K. Nov. Gen. et Sp.* ii. 131.—Venezuela.
bahiensis, *Klotzsch, in Linnaea*, xiv. (1840) 293.—Bras.
Banksii, *Benth. Fl. Austral.* iii. 312.—Austral.
Baraquiniana, Lem. Illustr. Hortic. vii. (1860) t. 276 = foetida.
Barclayi, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 634.—Ins. Fiji.
Baueriana, *Mast. l. c.*—Ins. Norfolk.
bauhiniaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 132 (bauhinifolia).—Ecuador.
Belottii ×, Hort. ex Moore & Ayres, Gard. Mag. Bot. i. (1850) 9, cum ic.
Berteriana, *Balb. ex DC. Prod.* iii. 325.—Ins. S. Doming.
bicornis, *Mill. Gard. Dict.* ed. VIII. n. 13.—N. Granat.
bicuspidata, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 635; *et Mart. Fl. Bras.* xiii. i. 558.—N. Granat.
biflora, Domb. ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xiii. (1873) 156 = P. Niorbo.
biflora, Lam. Encyc. iii. 36 = lunata.
biformis, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 128 = coriacea.
bifurca, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 162.—N. Granat.
biglandulosa, Caley, ex DC. Prod. iii. 333 = Herbertiana.
bilobata, *A. Juss. in Ann. Mus. Par.* vi. (1805) 107. t. 37. f. 2.—Ins. S. Doming.
bilobata, Vell. Fl. Flum. ix. t. 78 = capsularis.
bogotensis, Benth. Pl Hartw. 184 = alnifolia.
brachystephana, *F. Muell. ex Benth. Fl. Austral.* iii. 312.—Austral.
bracteosa, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 137 (Tacsoniae sp. ?).—N. Granat.
brasiliana, Desf. Cat. Hort. Par. ed. III. 411 = alata.
brasiliensis, Steud. Nom. ed. II. ii. 274, sphalm. = praec.
bryonifolia, Spreng. Syst. iii. 42 = seq.
bryonioides, *H. B. & K. Nov. Gen. et Sp.* ii. 140.—Mexic.
Buchanani, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 144 = vitifolia.
Buonapartea ×, Baxt. in Loud. Hort. Brit. Suppl. iii. 603; Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 145.
caerulea, *Linn. Sp. Pl.* 959.—Bras.
caerulea, Lour. ex DC. Prod. iii. 330 = chinensis.
calcarata, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 638.—Madag.
caloneura, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 95, 94 = Aristolochia trilobata.
campanulata, *Mast. in Mart. Fl. Bras.* xiii. i. 568, 615. —Bras.
candida, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 629. —Bras.
capsularis, Lam. Encyc. iii. 36 = rubra.
capsularis, *Linn. Sp. Pl.* 957.—Am. trop.
caracasana, *Jacq. ex Willd. Enum. Hort. Berol.* 697. —Venezuela.
cardinalis, *Hort. ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 145.—Hab. ?
Cavanillesii, DC. Prod. iii. 323 = cuprea.
celtifolia, Salisb. Prod. 154 = serratifolia.
cephaleima, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 152. t. 22. f. 2 = perfoliata.
ceratosepala, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 630.—Am. trop.
chelidonea, *Mast. in Gard. Chron.* (1879) II. 40.—Ecuador.
chilensis, Miers, Trav. Chil. ii. 522 = Tacsonia pinnatistipulata.

PASSIFLORA:—
chinensis, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 145 = caerulea.
chinensis, *Sweet, Hort. Brit.* ed. I. 355; ed. II. 217.—China.
ciliaris, 'Curt.' ex Steud. Nom. ed. I. 595 = *ciliata*.
ciliata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 310 = foetida.
cincinnata, *Mast. in Gard. Chron.* (1868) 966.—Bras.
cinerea, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 57. t. 177. —Guian.
cinnabarina, *Lindl. in Gard. Chron.* (1855) 724.—Austral.
cirrhiflora, *A. Juss. in Ann. Mus. Par.* vi. (1805) 115. t. 41. f. 1.—Guian.
citrifolia, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 629.—Guian.
citrifolia, Salisb. Prod. 154 = alata.
clathrata, *Mast. in Mart. Fl. Bras.* xiii. i. 547, 580.—Bras.
clypeata, Sm. in Rees, Cycl. xxvi. n. 20 = coriacea.
coccinea, *Aubl. Pl. Gui.* ii. 828. t. 324.—Guian.
coccinea, Blanco, Fl. Filip. ed. I. 650 = Modecca heterophylla.
coccinea, Soland. ex Benth. Fl. Austral. iii. 312 = Banksii.
cochinchinensis, Spreng. Syst. iv. Cur. Post. 346 (Malpigiac. ?).—Cochinch.
coerulea, Auct. vide caerulea.
colorata, *Mast. in Mart. Fl. Bras.* xiii. i. 562, 601.—Bras.
colubrina, Poepp. & Endl. Nov. Gen. et Sp. ii. 58 = triloba.
Colvillei ×, Sweet, Brit. Flow. Gard. Ser. II. i. t. 126.
Contrayerva, Sm. in Rees, Cycl. xxvi. n. 23 = bilobata.
coriacea, *Juss. in Ann. Mus. Par.* vi. (1805) 109. t. 39. f. 23.—Am. austr.
coriacea, *A. Rich. Ess. Fl. Cub.* 599.—Cuba.
cornuta, *Mast. in Mart. Fl. Bras.* xiii. i. 566.—Bras.
costata, *Mast. l. c.* 545, 573.—Bras.
cuellensis, Goudot, ex Planch. & Triana, in Ann. Sc. Nat. Sér. V. xvii. (1873) 154 = pilosa.
cuneata, *Willd. Enum. Hort. Berol.* 696.—Venezuela.
cuneifolia, Cav. Diss. x. t. 292 = edulis ?
cuprea, *Linn. Sp. Pl.* 955.—Ind. occ.
cyanea, *Mast. in Mart. Fl. Bras.* xiii. i. 571.—Venezuela.
cyathophora, Desv. in Ham. Prod. Pl. Ind. Occ. 48 = auriculata.
decipiens, *Mast. in Journ. Bot.* xxi. (1883) 34.—Guian.
dentata, Vell. Fl. Flum. ix. t. 94 = Raddiana ?
diaden, Vell. l. c. 90 = edulis.
Dictamo, [*Moç. & Sesse, ex*] *DC. Prod.* iii. 324.—Mexic.
difformis, H. B. & K. Nov. Gen. et Sp. ii. 136 = coriacea, *Juss.*
digitata, Linn. Sp. Pl. ed. II. 1355 = serrata.
discolor, Link & Otto, Ic. Pl. Sel. 13. t. 5 = Maximiliana.
divaricata, Griseb. in Bonplandia, vi. (1858) 7 = pulchella.
divaricata, Linn. Mant. ii. 491 = perfoliata.
edulis, *Sims, Bot. Mag.* t. 1989.—Bras.
Eichleriana, *Mast. in Mart. Fl. Bras.* xiii. i. 568.—Bras.
elata, Fisch. Hort. Gorenk. ed. II. (1812) 32, sphalm. = alata ?
elegans, *Mast. in Mart. Fl. Bras.* xiii. i. 621.—Bras.
elegans, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 133.—N. Granat.
elliptica, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 173.—Bras.
elongata, Poir. Encyc. Suppl. ii. 839 = oblongata.
emarginata, *Humb. & Bonpl. Pl. Aequin.* i. t. 23.—Peruv.
eminula, *Mast. in Journ. Linn. Soc.* xx. (1883) 32.—Am. trop.
erubescens, Macfad. Fl. Jamaic. ii. 154 = rubra.
erubescens, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 161.—N. Granat.
erythrophylla, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 633.—Am. trop.
exoperculata, *Mast. in Mart. Fl. Bras.* xiii. i. 556.—Bolivia.
exsudans, Zucc. in Abh. Akad. Muench. ii. (1831-36) 342 = bryonioides.

PASSIFLORA:—
ferruginea, *Mast. in Mart. Fl. Bras.* xiii. 1. 556.—Am. trop.
filamentosa, *Cav. Diss.* x. t. 294.—Am. austr.
filipes, *Benth. Pl. Hartw.* 118.—Mexic.; Ecuador.
flexipes, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 142 = Tacsonia flexipes.
flexuosa, Gardn. in Hook. Journ. Bot. i. (1834) 174 = suberosa.
floribunda, Lem. in Fl. des Serres, iv. (1848) 335 b = sexflora.
foetida, *Linn. Sp. Pl.* 959.—Bras.
foetida, Vell. Fl. Flum. ix. t. 86 = Vellozii.
fragrans, Regel, Cat. Hort. Aksakov. (1860) 104; et ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 145 = Middletoniana.
fulgens, Wallis, ex E. Morr. in Belg. Hortic. xv. (1865) 103; xvi. (1866) 193 = coccinea.
furcata, *Mast. in Mart. Fl. Bras.* xiii. 1. 550.—Venezuela.
fuscinata, *Mast. l. c.* 551.—Mexic.
Garckei, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 639.—Bras.
Gardneri, *Mast. in Mart. Fl. Bras.* xiii. 1. 566, 614.—Bras.
Gaudichaudiana, Planch. ex Planch. & Triana, in Ann. Sc. Nat. Sér. V. xvii. (1873) 142 = Tacsonia peduncularis.
Gaudichaudii, Planch. l. c. = praec.
geminiflora, DC. Prod. iii. 323 = P. Vespertilio.
geminiflora, D. Don, Prod. Fl. Nep. 63 = napalensis.
glaberrima, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 132 = Tacsonia glaberrima.
glabra, Mill. Gard. Dict. ed. VIII. n. 4 = minima.
glabra, *Wendl. Coll.* i. 55. t. 17.—Ins. Norfolk.
glabrata, H. B. & K. Nov. Gen. et Sp. ii. 135 = lunata.
glandulosa, *Cav. Diss.* x. 453. t. 281.—Bras.
glauca, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 308.—Mexic.
glauca, Humb. & Bonpl. Pl. Aequin. i. 76. t. 22 = arborea.
globosa, Vell. Fl. Flum. ix. t. 85 = suberosa.
gossypifolia, Desv. in Ham. Prod. Pl. Ind. Occ. 48 = foetida.
gossypifolia, *Lindl. Bot. Reg.* t. 1634.—Ind. occ.; Mexic.; Peruv.
Goudotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 132.—N. Granat.
gracilis, *Jacq. Eclog. Am.* t. 168; *Link, Enum. Hort. Berol.* ii. 182.—Bras.
grandiflora, Salisb. Prod. 154 = quadrangularis.
gritensis, *Karst. in Linnaea*, xxx. (1859–60) 163.—Venezuela.
guazumaefolia, *Jacq. Fragm.* 13; *Juss. in Ann. Mus. Par.* vi. (1805) 112. t. 39.—N. Granat.
guianensis, Miq. in Linnaea, xviii. (1844) 751 = quadriglandulosa.
haematostigma, *Mast. in Mart. Fl. Bras.* xiii. 1. 545.—Am. trop.
Hahnii, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 638.—Am. trop.
hainanensis, Hance, in Journ. Bot. xvi. (1878) 227 = ligulifolia.
hamosa, Goudot, ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 169 = acerifolia.
Hartwiesiana, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 145 = caerulea.
hastata, Bertol. Fl. Guatim. 27 = foetida.
hastata, Ruiz & Pav. ex Mast. in Mart. Fl. Bras. xiii. 1. 569 = reflexiflora.
hederacea, Cav. Diss. x. 448 = suberosa.
hederaefolia, Lam. Encyc. iii. 38 = minima, suberosa.
helleborifolia, *Wallis, ex Mast. in Mart. Fl. Bras.* xiii. 1. 622.—Am. trop.
Helleri, *Peyr. in Linnaea*, xxx. (1859–60) 54.—Mexic.
hemicycla, G. F. W. Mey. Prim. Fl. Esseq. 225 = P. Vespertilio.
Herbertiana, [*Ker.-Gawl. in*] *Bot. Reg.* t. 737.—Austral.
Hermanni, *DC. Prod.* iii. 332.—Ins. Curaçao.
heterophylla, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 309 = suberosa.
heterophylla, *Lam. Encyc.* iii. 41.—Ins. S. Doming.
hexangularis, Raeusch. Nom. ed. III. 85 (Quid?).—Ind. occ.

PASSIFLORA:—
Heyneana, Wall. Cat. n. 1233 = Erythropalum populifolium?
hibiscifolia, Lam. Encyc. iii. 39 = foetida.
hircina, Sweet, Hort. Brit. ed. I. 354; Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 146 = foetida.
hirsuta, Linn. Sp. Pl. 958 = suberosa.
hirsuta, Lodd. Bot. Cab. t. 138 = foetida.
hispida, *DC. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 172.—Reg. trop.
hispidula, Knowles & Westc. Flor. Cab. iii. (1839) 126 = suberosa.
holosericea, *Linn. Sp. Pl.* 958.—Mexic.
holosericea, Ruiz & Pav. ex Mast. in Journ. Linn. Soc. xx. (1883) 44 = alba.
Hondala, Steud. Nom. ed. II. ii. 275 = Modecca palmata.
Horsfieldii, *Blume, Rumphia*, 170. t. 52.—Java.
Hulletti, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 137 = macrocarpa.
hyacinthiflora, *Planch. & Linden, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. V. xvii. (1873) 179.—N. Granat.
hybrida, Nees, 'Sammlung, t. 36'; ex Pritz. Ic. Ind. ii. 206 (Quid?).—Hab.?
ianthina, *Mast. in Journ. Bot.* xxi. (1883) 36.—Bolivia.
ichthyura, *Mast. in Mart. Fl. Bras.* xiii. 1. 549.—Bras.
inamoena, *A. Gray, Pl. Wright.* ii. 59.—Mexic.
incana, Seem. ex Mast. in Journ. Linn. Soc. xx. (1883) 40 = Seemanni.
incarnata, Ker.-Gawl. in Bot. Reg. t. 332 = edulis.
incarnata, *Linn. Sp. Pl.* 959.—Am. trop.
indecora, *H. B. & K. Nov. Gen. et Sp.* ii. 134.—N. Granat.
Innesii ×, Mast. in Gard. Chron. (1870) 891.
insignis, Hook. Bot. Mag. t. 6069 = Tacsonia insignis.
Itondala, Steud. Nom. ed. I. 595 = *P. Hondala*.
Jileki, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 110; *et in Bot. Ergeb. Maxim. Bras.* 59.—Am. trop.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* ii. 133.—Mexic.
Kalbreyeri, *Mast. in Journ. Bot.* xxi. (1883) 36.—N. Granat.
Karsteniana, A. Dietr. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 42 = bryonoides.
Karwinskii, *Mast. in Mart. Fl. Bras.* xiii. 1. 555.—Am. trop.
Kegeliana, Garcke, in Linnaea, xxii. (1849) 60 = suberosa.
Kerii, Spreng. Syst. iii. 39 = edulis.
kermesina, Link & Otto, in Verh. Ver. Befoerd. Gartenb. ii. (1826) 403. t. 15 = Raddiana.
Kohautiana, Presl, Bot. Bemerk. 72 = suberosa.
Lamberti, *Hort. ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 146 (Quid?).—Hab.?
lanata, Poir. Encyc. Suppl. ii. 843 = Tacsonia lanata.
lancearia, *Mast. in Journ. Bot.* xxiii. (1885) 114.—Costa Rica.
lanceolata, G. Don, Gen. Syst. iii. 54 = seq.
lancifolia, *Desv. in Ham. Prod. Pl. Ind. Occ.* 48.—Ind. occ.
latifolia, DC. Prod. iii. 328 = alata.
laurifolia, *Linn. Sp. Pl.* 956.—Am. trop.
Lawsoniana ×, Mast. in Gard. Chron. (1868) 1288.—Bras.
Lehmanni, *Mast. in Journ. Bot.* xxiii. (1885) 115.—N. Granat.
Lemichesiana ×, Mast. in Journ. Hort. Soc. N. S. iv. (1877) 146.
lepidota, *Mast. in Mart. Fl. Bras.* xiii. 1. 547, 581.—Am. trop.
Leschenaultii, *DC. Prod.* iii. 326.—Ind. or.
Liebmanni, *Mast. in Mart. Fl. Bras.* xiii. 1. 547.—Am. trop.
ligularis, *A. Juss. in Ann. Mus. Par.* vi. (1805) 113. t. 40.—Am. trop.
ligulifolia, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 632.—Ins. Hongkong.
limbata, Tenore, Ind. Sem. Hort. Neap. (1839) 12. n. 10; et in Ann. Sc. Nat. Sér. II. xiii. (1840) 380 = suberosa?
Lindeniana, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 182.—Venezuela.

PASSIFLORA:—
linearifolia, Hook. f. in Trans. Linn. Soc. xx. (1347) 222 = suberosa.
littoralis, H. B. & K. Nov. Gen. et Sp. ii. 133 = suberosa.
Lobbii, *Mast. in Mart. Fl. Bras.* xiii. 1. 553.—N. Granat.
Lockharti, G. Don, ex M. Roem. Syn. Pepon. 183 = quadriglandulosa.
longiflora, Lam. Encyc. iii. 39 = Tacsonia mixta.
longifolia, Lam. l. c. 40 = suberosa.
longipes, *A. Juss. in Ann. Mus. Par.* vi. (1805) 111. t. 33. f. 1.—N. Granat.
lorifera, *Mast. & André, in Journ. Linn. Soc.* xx. (1883) 42.—Ecuador.
Loudoni ×, Loud. Encyc. Pl. 1234; Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 146.
Loureirii, G. Don, Gen. Syst. iii. 54 = caerulea?
Lowei, Heer, in Regel, Gartenfl. i. (1852) 69. t. 9 = ligularis.
lunata, *Juss. ex DC. Prod.* iii. 331.—Peruv.
lunata, Poepp. & Endl. Nov. Gen. et Sp. ii. 58 = Poeppigii.
lunata, Vell. Fl. Flum. ix. t. 80 = capsularis.
lunata, *Willd. Sp. Pl.* iii. 612 (Quid?).—Am. trop.
lutea, *Linn. Sp. Pl.* 958.—Ind. occ.
lutea, Ruiz & Pav. ex Mast. in Journ. Linn. Soc. xx. (1883) 44 = alba.
lyraefolia, Tussac, Fl. Antill. i. 70. t. 4 = oblongata.
macrocarpa, *Mast. in Gard. Chron.* (1869) 1012.—Bras.
macrophylla, *Spruce, ex Mast. in Journ. Linn. Soc.* xx. (1883) 31.—Ecuador.
maculata, Donn, Hort. Cantabr. ed. XIII. 436 = peltata.
maculata, *Scanag. ex Colla, Hort. Ripul.* 101.—Ind. occ.
Magdalenae, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 166.—N. Granat.
malacophylla, *Mast. in Mart. Fl. Bras.* xiii. 1. 563.—Am. trop.
maliformis, *Linn. Sp. Pl.* 956.—Ind. occ.
maliformis, Vell. Fl. Flum. ix. t. 73 = alata.
manicata, Pers. Syn. ii. 221 = Tacsonia manicata.
Manora, *Scott, in Journ. Linn. Soc.* viii. (1865) 202.—Hab.?
Mansoi, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 629.—Bras.
marginata, *Mast. in Mart. Fl. Bras.* xiii. 1. 562, 603.—Am. trop.
Marigouja, Perr. ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 172 = hispida.
marmorea, Linden, Cat. n. 8 (1853), 3; Hort. ex Gard. Chron. (1871) 234 = Ophiocaulon cissampeloides.
mascarensis, Presl, Bot. Bemerk. 72 = alata.
mauritiana, Thou. in Ann. Mus. Par. vi. (1805) 457. t. 65 = alata.
Maximiliana, *Bory, in Ann. Gén. Sc. Phys.* ii. (1819) 129. t. 24.—Bras.
Mayana, Veitch, ex Voigt, Hort. Suburb. Calc. 81; Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 146 = caerulea.
mediterranea, *Vell. Fl. Flum.* ix. t. 72.—Bras.
Medusae, Lem. in Fl. des Serres, iii. (1848) 373 b = jorullensis.
membranacea, *Benth. Pl. Hartw.* 88.—Guatemal.
menispermacea, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 168.—N. Granat.
menispermifolia, H. B. & K. Nov. Gen. et Sp. ii. 137 = pilosa.
meridensis, *Karst. in Linnaea*, xxx. (1859–60) 165.—N. Granat.
mexicana, *A. Juss. in Ann. Mus. Par.* vi. (1805) 108. t. 38. f. 2.—Mexic.
microcarpa, *Mast. in Mart. Fl. Bras.* xiii. 1. 556, 593.—Am. trop.
micropetala, *Mast. l. c.* 549.—Am. trop.
Middletoniana, *Paxt. Mag. Bot.* ix. (1842) t. 51 (an hybr.?).—Hab.?
Miersii, *Mast. in Mart. Fl. Bras.* xiii. 1. 561, 599.—Bras.
minima, Blanco, Fl. Filip. ed. I. 647 = serrulata?
minima, Linn. Sp. Pl. 950 = suberosa.
misera, *H. B. & K. Nov. Gen. et Sp.* ii. 136.—Am. austr.
mixta, Linn. f. Suppl. 408 = Tacsonia mixta.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 137.—N. Granat.

PASSIFLORA:—
moluccana, *Reinw. in Blume, Bijdr.* 938.—Ins. Molucc.
Monnieri, Buchoz, Herb. Coll. Am. (1788) t. 35 = coccinea.
Mooreana, *Hook. Bot. Mag.* t. 3773.—Bras.
morifolia, *Mast. in Mart. Fl. Bras.* xiii. 1. 555.—Bras.
Moritziana, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 175.—N. Granat.
mucronata, *Lam. Encyc.* iii. 33.—Bras.
Muelleriana, *Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 133.—Austral.
multiflora, *Linn. Sp. Pl.* 956.—Ins. S. Doming.
multiformis, *Jacq. Fragm.* 50. t. 67. f. 1.—Venezuela.
Munroii ×, Mast. in Gard. Chron. (1868) 1288.
Murucuia, Mill. Gard. Dict. ed. VIII. n. 14 = perfoliata.
Murucuja, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 131 = lunata, *Juss.*?
Murucuja, *Linn. Sp. Pl.* 957.—Ind. occ.
myriadenia, *Van Houtte, Hortus*, t. 5; *ex Pritz. Ic. Ind.* i. 806 (Quid?).—Hab.?
napalensis, *Wall. Tent. Fl. Nep.* 20. t. 11.—Reg. Himal.
naviculata, *Griseb. in Goett. Abh.* xix. (1874) 149.—Am. austr.
Neillii, *Regel, in Flora*, xxxii. (1849) 184.—Hab.?
nepalensis, Walp. Rep. ii. 221 = napalensis.
Neumannii ×, Paxt. Mag. Bot. xv. (1849) 270, cum t.
nigelliflora, Hook. Bot. Mag. t. 3635 = foetida.
nigra, Jacq. Obs. Bot. ii. 27. t. 46. f. 3 = suberosa.
Niorbo, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 156.—Peruv.
nipaulensis, G. Don, Gen. Syst. iii. 47 = napalensis.
nitida, *H. B. & K. Nov. Gen. et Sp.* ii. 130.—Venezuela.
normalis, Linn. Syst. ed. X. 1248 = perfoliata.
nymphaeoides, Karst. in Linnaea, xxx. (1859–60) 165 = nitida.
oblonga, DC. Prod. iii. 327 = seq.
oblongata, *Sw. Prod. Veg. Ind. Occ.* 97.—Jamaic.
obscura, Lindl. in Trans. Hort. Soc. vii. (1830) 148 = foetida.
obtusa, Vell. Fl. Flum. ix. t. 81 = organensis?
obtusiloba, *Mast. in Mart. Fl. Bras.* xiii. 1. 554.—Peruv.
ocanensis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 183.—N. Granat.
octandra, Thomps. ex DC. Prod. iii. 337 = Deidamia Thompsoniana.
odora, Link & Otto, Ic. Pl. Rar. 93. t. 47 = sicyoides.
Oerstedii, *Mast. in Mart. Fl. Bras.* xiii. 1. 562.—Costa Rica.
olivaeformis, Mill. Gard. Dict. ed. VIII. n. 6; Vell. Fl. Flum. ix. t. 83 = suberosa.
onychina, Lindl. Bot. Reg. (1838) Misc. 1. t. 21 = amethystina.
orbiculata, Cav. Diss. x. 456 = P. Murucuja.
orbifolia, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 150.—N. Granat.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 104.—Bras.
ornata, H. B. & K. Nov. Gen. et Sp. ii. 129 = maliformis.
ovalis, *Vell. Fl. Flum.* ix. t. 75.—Bras.
ovata, *Martin, ex DC. Prod.* iii. 322.—Guian.
oviformis, M. Roem. Syn. Pepon. 167 = alata.
Pala, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 162.—N. Granat.
pallens, *Poepp. ex Mast. in Trans. Linn. Soc.* xxvii. 1871) 638.—Am. trop.
pallida, Linn. Sp. Pl. 955 = suberosa.
pallida, Lour. Fl. Cochinch. 644 = *cochinchinensis*.
pallida, Vell. Fl. Flum. ix. t. 70 = mucronata.
pallidiflora, Bertol. Syll. Pl. Hort. Bonon. (1827) 6; ex Moris, in Mem. Acc. Torin. xxxvi. (1833) 178. t. 6 = edulis.
palmata, Lodd. Bot. Cab. 97 = filamentosa.
palmatisecta, *Mast. in Mart. Fl. Bras.* xiii. 1. 564.—Reg. Argent.
pamplonensis, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 137 (Tacsoniae sp.).—N. Granat.

PASSIFLORA:—
pannosa, *Sm. in Rees, Cycl.* xxvi. n. 28.—Ind. occ.
parviflora, Blanco, Fl. Filip. ed. I. 649 = Modecca cardiophylla.
parviflora, Sw. Prod. Veg. Ind. Occ. 97 = suberosa.
pavonis, *Mast. in Journ. Bot.* xxi. (1883) 35.—Mexic.
pectinata, *Griseb. Fl. Brit. W. Ind.* 294.—Ind. occ.
pedata, *Linn. Sp. Pl.* 960.—Ins. S. Doming.
pedunculatis, Cav. Ic. v. 15 = Tacsonia pedunculatis.
pedunculata, Hort. ex Mast. in Journ. Hort. Soc. iv. (1877) 137 = alata.
peltata, Cav. Diss. x. 447. t. 274 = suberosa.
penangiana, Wall. Cat. n. 1233 = Modecca nicobarica.
penduliflora, *Bert. ex DC. Prod.* iii. 326.—Jamaic.
pennipes, M. Roem. Syn. Pepon. 194 = Tacsonia pinnatistipula.
pentagona, *Mast. in Mart. Fl. Bras.* xiii. 1. 545, 575.—Am. trop.
perfoliata, *Linn. Sp. Pl.* 956.—Ind. occ.
pertusa, Vell. Fl. Flum. ix. t. 79 = organensis?
phoenicea, Lindl. Bot. Reg. t. 1603 = alata.
picta, Hort. ex Steud. Nom. ed. I. 596 = tuberosa.
picturata, *Ker-Gawl. in Bot. Reg.* t. 673.—Bras.
piligera, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 173 = capsularis.
pilosa, *Ruiz & Pav. ex DC. Prod.* iii. 330.—Mexic.
Pinardia, Buchoz, Herb. Pl. Med. Chin. t. 30 (Quid?).—China.
pinnatifolia, Molina, Sagg. Chile, Ed. II. 292 = Tacsonia pinnatistipula.
pinnatistipula, Cav. Ic. v. 16 = Tacsonia pinnatistipula.
Pisonis, Kostel. Allg. Med.-Pharm. Fl. iv. 1389 = cincinnata?
platystyla, *Mart. in Journ. Bot.* xxi. (1883) 35.—Bras.
Poeppigii, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 630.—Am. trop.
Pohlii, *Mast. in Mart. Fl. Bras.* xiii. 1. 549.—Bras.
Polignacia, Buchoz, Herb. Pl. Méd. Chin. t. 38 (Quid?).—China.
polyaden, Vell. Fl. Flum. ix. t. 92 = foetida.
pomifera, M. Roem. Syn. Pepon. 179 = edulis.
populifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 150.—N. Granat.
porophylla, *Vell. Fl. Flum.* ix. t. 82.—Bras.
Princeps, Lodd. Bot. Cab. t. 84 = racemosa.
Pseudo-suberosa, Fisch. Ind. Sem. Hort. Petrop. ix. 82 = suberosa.
pubera, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 185.—N. Granat.
puberula, Hook. f. in Trans. Linn. Soc. xx. (1847) 222 = suberosa.
pubescens, H. B. & K. Nov. Gen. et Sp. ii. 132 = capsularis.
pulchella, *H. B. & K. l. c.* ii. 134.—Venezuela.
punctata, *Linn. Sp. Pl.* 957.—Peruv.
punctata, Lodd. Bot. Cab. t. 101 = tuberosa.
punicea, Ruiz & Pav. ex DC. Prod. iii. 329 = vitifolia.
pyriformis, DC. l. c. 331 = alata?
quadrangularis, *Linn. Syst.* ed. X. 1248.—Am. trop.
quadrangularis, Triana & Planch in Ann. Sc. Nat. Sér. V. xvii. (1873) 147 = macrocarpa.
quadriglandulosa, *Rodschied, Obs.* 77.—Am. trop.
racemosa, Brot. in Trans. Linn. Soc. xii. (1817) 71.—Bras.
Raddiana, *DC. Prod.* iii. 329.—Bras.
recurva, *Mast. in Mart. Fl. Bras.* xiii. 1. 565, 608.—Bras.
reflexiflora, *Cav. Ic.* v. 15. t. 425.—Ecuador.
regalis, Macfad. Fl. Jamaic. ii. 151 = lancifolia.
resticulata, *Mast. & André, in Journ. Linn. Soc.* xx. (1883) 42.—Am. trop.
reticulata, *Sauv. Fl. Cub.* 56.—Cuba.
retusa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 325 = Maximiliana.
rhamnifolia, *Mast. in Mart. Fl. Bras.* xiii. 1. 545, 575.—Am. trop.
rigidula, Jacq. Eclog. Am. ii. 5. t. 124 = incarnata?
riparia, *Mart. ex Mast. in Mart. Fl. Bras.* xiii. 1. 599.—Am. trop.
Rohrii, DC. Prod. iii. 326 = auriculata.
rotundifolia, Jacq. Obs. t. 46. f. 1 = pulchella.
rotundifolia, *Linn. Sp. Pl.* 957.—Ind. occ.
rotundifolia, Sw. Obs. Bot. 337 = Swartzii.
rubra, Lam. Encyc. iii. 35 = capsularis.

PASSIFLORA:—
rubra, *Linn. Sp. Pl.* 956.—Ind. occ.
rubricaulis, Jacq. Eclog. Am. ii. t. 169 = edulis.
rugosa, Planch. & Triana, in Ann. Sc. Nat. Sér. V. xvii. (1873) 136 = Tacsonia rugosa.
sanguinea, Colla, Hort. Ripul. App. iii. 45. t. 6 = *hybrida*?
sanguinea, Sm. in Rees, Cycl. xxvi. n. 45 = vitifolia.
sanguinolenta, *Mast. & Linden, in Gard. Chron.* (1868) 1162.—N. Granat.?; Peruv.?
saponaria, Blanco, Fl. Filip. ed. I. 650 = Modecca trilobata.
Schlimiana, *Regel, Cat. Pl. Hort. Aksakov.* 105; *et Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 130.—N. Granat.
Seemanni, *Griseb. in Bonplandia*, vi. (1858) 7.—Am. centr.
Selloi, Dehnh. Revista Napol. i. III. 180; ex Walp. Rep. ii. 220 = serrata.
semiciliosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 145.—N. Granat.
semiorbicularis, *Hort. ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 146, nomen.—Bras.?
sengaporeana, Steud. Nom. ed. II. ii. 276 = Modecca singaporeana.
septenata, DC. Prod. iii. 323 = cirrhiflora.
serrata, *Linn. Syst.* ed. X. 1248.—Ins. Martinic.
serratifolia, *Linn. Sp. Pl.* 955.—Am. austr.
serratistipula, [Moç. & Sesse, ex] DC. Prod. iii. 328 = ligularis.
serrato-digitata, Linn. Sp. Pl. 960 = serrata.
serrulata, *Jacq. Obs.* ii. t. 46. f. 2.—N. Granat.
servitensis, Karst. in Linnaea, xxx. (1859–60) 163 = vitifolia.
setacea, *DC. Prod.* iii. 329.—Bras.
sexflora, *A. Juss. in Ann. Mus. Par.* vi. (1805) t. 37. f. 1.—Ins. S. Doming.
sexocellata, Schlecht. in Linnaea, xxvii. (1854) 521 = mexicana.
sicyoides, *Cham. & Schlecht. in Linnaea*, v. (1830) 88.—Mexic.
sidaefolia, M. Roem. Syn. Pepon. 173 = tetraden.
silvestris, *Vell. Fl. Flum.* ix. t. 74.—Bras.
singaporeana, Wall. Cat. n. 1232 = Modecca singaporeana.
spathulata, *Mast. in Mart. Fl. Bras.* xiii. 1. 552.—Am. trop.
speciosa, *Gardn. in Field. Sert. Pl.* t. 17.—Bras.
sphaerocarpa, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 184.—N. Granat.
spicata, *Mast. in Mart. Fl. Bras.* xiii. 1. 545, 576.—Am. trop.
spinosa, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 630.—Bras.
Sprucei, *Mast. in Mart. Fl. Bras.* xiii. 1. 568.—Ecuador.
stenophylla, Hoffmgg. Verz. Pfl. 85 = caerulea.
stipulata, *Aubl. Pl. Gui.* ii. 830. t. 325.—Guian.
stipulata, Benth. ex Griseb. Cat. Pl. Cub. 113; et ex Mast. in Mart. Fl. Bras. xiii. 1. 567 = pallens.
stipulata, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 152 = alba, glauca.
suberosa, *Linn. Sp. Pl.* 958.—Am. trop.
subpeltata, Orteg. Hort. Matr. Dec. 78 = subulata?
subrotunda, *Mast. in Mart. Fl. Bras.* xiii. 1. 561, 601.—Bras.
subulata, *Mast. l. c.* 567.—Peruv.
sulcata, Jacq. Select. Am. 232 = quadrangularis.
Sulivani, Booth, ex Baxt. in Loud. Hort. Brit. Suppl. ii. 654 = amethystina.
sumatrana, *Blume, Rumphia*, i. 170, in obs.—Sumatra.
surinamensis, Miq. in Linnaea, xviii. (1844) 363 = P. Vespertilio.
Sururuca, Vell. Fl. Flum. ix. t. 88 = setacea.
Swartzii, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 634 (Swarzii); *et in Mart. Fl. Bras.* xiii. 1. 556.—Bras.
sylvestris, Auct. = silvestris.
Tacso, Cav. Diss. x. 450 = Tacsonia mixta.
tacsonioides, *Griseb. Fl. Brit. W. Ind.* 295.—Jamaic.
tenuiloba, *Engelm. in A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 192.—Texas.
tetradena, Vand. ex DC. Prod. iii. 331 = alata.
tetradena, *Vell. Fl. Flum.* ix. t. 91.—Bras.
tetragona, M. Roem. Syn. Pepon. 165 = alata.
tetrandra, *Banks & Soland. ex DC. Prod.* iii. 323.—N. Zel.

PASSIFLORA :—
theobromaefolia, DC. Prod. iii. 331 = guazumaefolia.
tiliaefolia, *Cav. Diss.* x. 455. t. 285.—Peruv.
tiliaefolia, Linn. Sp. Pl. 956 = ligularis *vel* triloba.
timorensis, M. Roem. Syn. Pepon. 191 = moluccana.
timoriana, Span. in Linnaea, xv. (1841) 207 = moluccana.
tinifolia, Juss. in Ann. Mus. Par. vi. (1805) 113. t. 41. f. 2 = laurifolia.
tomentosa, Cav. Diss. x. 450. tt. 275, 276 = Tacsonia mixta.
tomentosa, Lam. Encyc. iii. 40 = Tacsonia tomentosa.
torta, *Mast. in Mart. Fl. Bras.* xiii. 1. 548.—Bras.
tricuspis, *Mast. l. c.* 550.—Bras.
tridactylites, Hook. f. in Trans. Linn. Soc. xx. (1847) 222 = suberosa.
trifasciata, *Lem. Illustr. Hortic.* xv. (1868) t. 544.—Hab. ?
triflora, *Macfad. Fl. Jamaic.* ii. 149.—Jamaic.
trifoliata, Cav. Ic. v. 16 = Tacsonia trifoliata.
trigona, Ruiz & Pav. ex DC. Prod. iii. 334 = Tacsonia trifoliata.
triloba, *Ruiz & Pav. ex DC. l. c.* 330.—Peruv.
trinervia, *Poir. Encyc. Suppl.* ii. 843; *Spreng. Syst.* iii. 40; *Mast. in Trans. Linn. Soc.* xxvii. (1871) 635.—N. Granat.
tripartita, *Breit. Hort. Breit.* 327, nomen (Quid ?).—Am. austr.
triquetra, *Lam. ex Mart. in Trans. Linn. Soc.* xxvii. (1871) 633 = rubra.
trisecta, *Mast. in Mart. Fl. Bras.* xiii. 1. 564.—Bolivia.
trisetosa, [*Moç. & Sesse ex,*] *DC. Prod.* iii. 324.—Mexic.
truncata, *Regel, Gartenfl.* viii. (1859) 356.—Bras.
truxillensis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 135.—Venezuela.
tuberosa, *Jacq. Hort. Schoenb.* iv. 49. t. 496.—Am. trop.
tubiflora, H. B. & K. Nov. Gen. et Sp. ii. 139 = viridiflora.
tucumanensis, *Hook. Bot. Mag.* t. 3636.—Chili.
ulmifolia, *Hort. ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 146.—Hab. ?
Van-Volxemi, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 141 = Tacsonia Van-Volxemii.
Varanzof, Carr. in Rev. Hortic. (1872) 320 = racemosa ?
variegata, Mill. Gard. Dict. ed. VIII. n. 8 = foetida.
variolata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 58. t. 179.—Bras.
velata, *Mast. in Mart. Fl. Bras.* xiii. 1. 560.—Venezuela.
Vellozii, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 103.—Bras.
velutina, DC. Prod. iii. 327 = coccinea.
verecunda, Salisb. Prod. 155 = rubra.
verrucifera, Lindl. Bot. Reg. (1840) Misc. 49 = edulis.
verrucosula, Steud. Nom. ed. II. ii. 276, sphalm. = seq.
verruculosa, *Weinm. in Syll. Ratisb.* i. (1824) 228.—Austral.
Vespertilio, Ker-Gawl. in Bot. Reg. t. 597 = Maximiliana.
Vespertilio, Lawrence, Passif. t. 8; ex DC. Prod. iii. 326 = lunata.
Vespertilio, *Linn. Sp. Pl.* 957.—Am. trop.
vetutina, G. Don, Gen. Syst. iii. 51, sphalm. = coccinea.
villosa, Macfad. Fl. Jamaic. ii. 151 = suberosa.
villosa, *Vell. Fl. Flum.* ix. t. 87.—Bras.
violacea, *Vell. l. c.* t. 84.—Bras.
viridiflora, *Cav. Ic.* v. 15. t. 424.—Mexic.
viridiflora, Hort. ex Tenore, Ind. Sem. Hort. Neap. (1839) 12. n. 10, partim = limbata.
vitiensis, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 634.—Ins. Fiji.
vitifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 138.—Am. austr.
Walkeri, Wight, Illustr. Ind. Bot. ii. 39. t. 108 = suberosa.
Warei, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 297 = suberosa.
Warmingii, *Mast. in Mart. Fl. Bras.* xiii. 1. 554, 91.—Am. trop.

PASSIFLORA :—
Weberiana, *André, in Rev. Hortic.* (1885) 113.—Reg. Argent.
Zucca, Blanco, Fl. Filip. ed. I. 648 = Modecca palmata.

PASSOURA, Aubl. Pl. Gui. Suppl. 21. t. 380 (1775) = **Alsodeia**, Thou. (Violar.).
guianensis, Aubl. l. c. = A. guianensis.

PASSOVIA, Karst. ex Klotzsch, in Bot. Zeit. iv. (1846) 107 = **Loranthus**, Linn.
odorata, Karst. l. c. x. (1852) 305 = L. Theobromae.
suaveolens, Karst. l. c. iv. (1846) 107 (an praec. ?).—N. Granat.

PASSOWIA, Karst. in Bot. Zeit. x. (1852) 305 = praec.

PATAGONICA, Dill. ex Adans. Fam. ii. 179 (1763) = **Patagonula**, Linn. (Borag.).

PATAGONIUM, E. Mey. Comm. Pl. Afr. Austr. 203. t. 37 (1835) = **Aeschynomene**, Linn. (Legumin.).
racemosum, E. Mey. l. c. = A. micrantha.

PATAGONIUM, Schrank, in Denkschr. Akad. Muench. (1808) 93 = **Adesmia**, DC. (1825) (Legumin.).
hedysaroides, Schrank, l. c. = A. muricata.

PATAGONULA, Linn. Gen. ed. I. 45 (1737).
BORAGINEAE, Benth. & Hook. f. ii. 839.
PATAGONICA, Dill. ex Adans. Fam. ii. 179 (1763).
americana, *Linn. Sp. Pl.* 149.—Patagonia.
australis, Salisb. Prod. 111 = americana.
bahiensis, *Moric. Pl. Nouv. Am.* 144. t. 86.—Bras.
glabra, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 29 = americana.
Tweediana, *Miers, l. c.*—Bras.

PATAGUA, Poepp. ex Baill. Adansonia, x. (1871) 49 = **Orites**, R. Br. (Proteac.).
chilensis, Poepp. ex Baill. l. c. = R. myrtoidea.

PATAMOGETON, Honck. Syn. 110 (1793), err. typ. = **Potamogeton**, Tourn. (Naiad.).

PASTINACA, Linn. Gen. ed. I. 81 (1737) = **Peucedanum**, Tourn. (Umbell.).
altissima, Lam. Fl. Fr. iii. 465 = Opopanax Chironium.
ambigua, Torr. Fl. U. St. 315 = Peuced. ternatum.
Anethum, Spreng. in Schult. Syst. vi. 587 = Peuced. graveolens.
angulosa, Dulac, Fl. Hautes-Pyr. 343 = Peuced. sativum.
armena, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 324.—Reg. Caucas.
arvensis, Steud. Nom. ed. I. 596 = Peuced. sativum.
atropurpurea, Steud. ex A. Rich. Tent. Fl. Abyss. i. 329 = Peuced. atropurpureum.
Candolleana, Wight & Arn. Prod. 372 = Heracleum rigens.
capensis, Sond. in Harv. & Sond. Fl. Cap. ii. 562.—Afr. austr.
dasyantha, C. Koch, in Linnaea, xvi. (1842) 359 = Malabaila pimpinellifolia.
dasycarpa, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 598.—Turkestan.
denticulata, D. Dietr. Syn. Pl. ii. 971 = Peuced. ternatum.
dissecta, Vent. Hort. Cels, 78 = Malabaila Sekakul.
divaricata, Desf. Tabl. ed. II. 139; Cat. Hort. Par. ed. III. 200, 405 = Peuced. divaricatum.
esculenta, Salisb. Prod. 167 = Peuced. sativum.
ferulacea, Walp. Rep. iii. 412, in syn. = Peuced. foeniculaceum.
Fleischmanni, Hladnik, in D. Dietr. Syn. Pl. ii. 970.—Carinth.
Fleischmanni, Reichb. ex Steud. Nom. ed. II. ii. 276 = Peuced. caruifolium.
foeniculacea, Spreng. in Schult. Syst. vi. 587 = Peuced. foeniculaceum.
glandulosa, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 1062.—Cataonia.
glauca, Dalz. in Hook. Kew Journ. iv. (1852) 293 = Peuced. Dana.
glauca, Scop. Fl. Carn. ed. II. i. 203 = farrago spp. Peucedani et Trinia vulgaris.
grandis, Dalz. & Gibs. Bomb. Fl. 107 = Peuced. grande.

PASTINACA :—
graveolens, Bernh. Syst. Verz. Erf. 171 = Peucedanum graveolens.
graveolens, Bieb. Fl. Taur. Cauc. i. 237 = Malabaila graveolens.
graveolens, Salzm. ex Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 91 = Peuced. divaricatum.
hirsuta, Panč. Fl. Princ. Sorbiae (1874), 359 = Peuced. sativum.
hispida, Fenzl, in Flora, xxvi. (1843) 462 = Opopanax hispidum.
Hookeriana, Wight, Ic. t. 1010 = Heracleum Hookerianum.
incana, Willd. ex DC. Prod. iv. 196 = Zosimia orientalis.
intermedia, Fisch. & Mey. ex Boiss. Fl. Orient. ii. 1061.—Reg. Caucas.
involucrata, Cesati, ex Nym. Consp. 288 = Malabaila involucrata.
involucrata, C. Koch, in Linnaea, xvi. (1842) 359 (nom. delend.—Cf. Boiss. Fl. Orient. ii. 1062).
involucrata, Stokes, Bot. Mat. Med. ii. 132 = Peuced. Opopanax.
Kochii, Duby, Bot. Gall. i. 220 = Peuced. divaricatum.
latifolia, DC. Prod. iv. 189.—Corsica.
latifolia, Ledeb. Fl. Ross. ii. 318 = *teretiuscula*, Boiss.
ligusticifolia, Wight & Arn. Prod. 372 = Heracleum rigens.
lucida, Auct. ex Nym. Consp. 286 = *latifolia*.
lucida, Linn. Mant. i. 58.—Ins. Balear.
lutea, Gilib. Fl. Lituan. ii. 39 = Peuced. sativum.
Mazurevskii, Kalen. in Bull. Soc. Nat. Mosc. xviii. (1845) 1. 232.—Reg. Caucas.
nudicaulis, Spreng. in Schult. Syst. vi. 587 = Peuced. nudicaule.
obtusifolia, DC. Prod. iv. 190 = Malabaila obtusifolia.
Olgae, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 37.—Turkestan.
opaca, Bernh. ex Hornem. Hort. Hafn. 961 = Peuced. sativum.
Opopanax, Linn. Sp. Pl. 262 = Opopanax hispidum.
panacifolia, Fisch. ex Sweet, Hort. Brit. ed. II. 251.—Tauria.
parvifolia, Kral. ex Nym. Consp. 289 = Peuced. divaricatum.
Petteri, Vis. ex Reichb. Fl. Germ. Excurs. 866 = Peuced. coriaceum.
pimpinellifolia, Bieb. Fl. Taur. Cauc. i. 237 = Malabaila pimpinellifolia.
pimpinellifolia, Bory & Chaub. Nouv. Fl. Pélop. 19 = Malabaila aurea.
pimpinellifolia, Griseb. Spicil. Fl. Rumel. i. 376 = Malabaila involucrata.
pratensis, H. Mart. Fl. Mosq. 55; ex Ledeb. Fl. Ross. ii. 318 = Peuced. sativum.
propinqua, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 286, in obs. = Peuced. sativum.
rectistyla, Cesati, in Linnaea, xi. (1837) 321 = Malabaila aurea.
rectistylis, Boiss. Fl. Orient. ii. 1053 = praec.
rigens, Wight, Ic. t. 1009 = Heracleum rigens.
rigida, Spreng. in Schult. Syst. vi. 586 = Peuced. ternatum.
sativa, Linn. Sp. Pl. 262 = Peuced. sativum.
sativa, Thomas, ex DC. Prod. iv. 189 = Peuced. divaricatum.
Secacul, [Soland. in] Russ. Aleppo, ed. II. ii. 249.—Syria; Aegypt.
Sekakul, DC. Prod. iv. 189, partim. = Malabaila pumila.
selinoides, Vis. in Flora, xii. (1829) 1. Erg. 10 = Peuced. caruifolium.
Sprengeliana, Wight, Ic. t. 1008 = Heracleum Sprengelianum.
stenocarpa, DC. Prod. iv. 189.—Sibir.
sulcata, C. Koch, ex Boiss. Fl. Orient. ii. 1055 = Malabaila sulcata.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 1 = Peuced. sativum.
taraxacifolia, Sweet, Hort. Brit. ed. II. 251 = Peuced. sativum.
tereticaulis, Bor. ex Nym. Consp. 289 = Peuced. sativum.
teretiuscula, Boiss. Fl. Orient. ii. 1060.—As. Min.; Syria.
teretiuscula, Jord. ex Nym. Consp. 289 = Peuced. sativum.

PASTINACA:—
tolucensis, Spreng. Syst. i. 913 = Peucedanum tolucense.
tricuspidata, D. Dietr. Syn. Pl. ii. 971 = Peuced. ternatum.
triquinata, Spreng. Sp. Umbellif. 68 = Archangelica hirsuta.
umbrosa, Stev. ex DC. Prod. iv. 189 = Peuced. sativum.
urens, Req. ex Gren. & Godr. Fl. Fr. i. 694 = Peuced. sativum.
velutina, Koch, ex DC. Prod. iv. 189 = Peuced. divaricatum.
zozimoides, Fenzl, in Tchihat. Asie Min. Bot. i. 436. —Cilicia.

PASTINACHA, Hill, Herb. Brit. i. 132 (1769) = praec.

PASTORAEA, Tod. in Nuov. Gen. e Sp. Piante, i. 17 (1858) = seq.

PASTOREA, Tod. ex Bertol. Fl. Ital. x. 520 (1854) = **Cochlearia**, Tourn. (Crucif.).
praecox, Tod. l. c. (= *Ionopsidium albiflorum*).—Sicil.
Savianum, Ball, ex Nym. Consp. 66 = Bivonaea Saviana.

PATABEA, Aubl. Pl. Gui. i. 110. t. 43 (1775) = **Psychotria**, Linn. (Rubiac.).
alba, H. B. & K. Nov. Gen. et Sp. iii. 375 = Psych. maypurensis.
capitellata, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 209 = Pysch. barbiflora.
coccinea, Aubl. Pl. Gui. i. 111.—Guian.
coriacea, Cham. in Linnaea, ix. (1835) 234 = Psych. xanthophylla.
lanceolata, Presl, Bot. Bemerk. 80.—Bras.
tenerior, Cham. in Linnaea, ix. (1835) 236 = Psych. tenerior.
tenuiflora, DC. Prod. iv. 538.—Guian.

PATAGNANA, Steud. Nom. ed. II. ii. 277 (1841), sphalm. = PETAGNANA, J. F. Gmel. = **Smithia**, Ait. (Legumin.).

PATERSONIA, R. Br. Prod. 303 (1810). *IRIDEAE*, Benth. & Hook. f. iii. 699.
GENOSIRIS, Labill. Nov. Holl. Pl. i. 13. t. 9 (1804), nomen prius.
babianoides, *Benth. Fl. Austral.* vi. 408.—Austral.
bicolor, F. Muell. ex Benth. l. c. 403 = longiscapa.
compar, Endl. in Lehm. Pl. Preiss. ii. 30 = occidentalis.
Diesingii, Endl. l. c. = occidentalis.
Drummondii, *F. Muell. ex Benth. Fl. Austral.* vi. 407. —Austral.
flaccida, Endl. in Lehm. Pl. Preiss. ii. 30 = occidentalis.
glabrata, R. Br. in Bot. Reg. t. 51 = sericea.
glabrata, *R. Br. Prod.* 304.—Austral.
glauca, *R. Br. l. c.*—Austral.
graminea, *Benth. Fl. Austral.* vi. 408.—Austral.
inaequalis, *Benth. l. c.*—Austral.
juncea, *Lindl. Swan River App.* 58.—Austral.
lanata, *R. Br. Prod.* 303.—Austral.
limbata, *Endl. in Lehm. Pl. Preiss.* ii. 29.—Austral.
longifolia, *R. Br. Prod.* 303.—Austral.
longiscapa, *Sweet, Fl. Austral.* t. 39.—Austral.
macrantha, *Benth. Fl. Austral.* vi. 407.—Austral.
Maxwellii, *F. Muell. ex Benth. l. c.* 405.—Austral.
media, R. Br. Prod. 304 = glabrata.
montana, Endl. in Lehm. Pl. Preiss. ii. 31 = occidentalis.
nana, Endl. l. c. 30 = occidentalis.
occidentalis, *R. Br. Prod.* 304.—Austral.
pannosa, Endl. in Lehm. Pl. Preiss. ii. 29 = lanata.
pygmaea, *Lindl. Swan River App.* 58.—Austral.
Roei, Endl. in Lehm. Pl. Preiss. ii. 31 = juncea?
rudis, *Endl. l. c.* 29.—Austral.
sapphirina, Lindl. Bot. Reg. (1839) t. 60 = occidentalis.
sericea, *R. Br. in Bot. Mag.* t. 1041.—Austral.
sylvestris, Endl. in Lehm. Pl. Preiss. ii. 30 = occidentalis.
tenuispatha, Endl. l. c. 31 = occidentalis.
turfosa, Endl. l. c. = occidentalis.
umbrosa, *Endl. l. c.*—Austral.
xanthina, *F. Muell. Fragm.* i. 214.—Austral.

PATERSONIA, *vel* PATHERSONIA, Poir. Encyc. Suppl. iv. 323 (1816) = PATTERSONIA, J. F. Gmel. = **Ruellia**, Linn. (Acanth.).

PATIENTIA, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Linn. (Polygon.).

PATIMA, Aubl. Pl. Gui. i. 196. t. 77 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 73.
Forsythii, *DC. Prod.* iv. 444 (an seq.?).—Guian.
guianensis, *Aubl. Pl. Gui.* i. 197. t. 77.—Guian.
laxiflora, Benth. in Hook. Journ. Bot. iii. (1841) 220 = Kutchubaea insignis.

PATONIA, Wight, Illustr. i. 18 (1838) = **Xylopia**, Linn. (Anonac.).
parvifolia, Wight, l. c. 19 = X. parvifolia.
Walkerii, Wight, l. c. = Diospyros Gardneri.

PATRINIA, Juss. in Ann. Mus. Par. x. (1807) 311. *VALERIANEAE*, Benth. & Hook. f. ii. 153.
FEDIA, Adans. Fam. ii. 152 (1763).
MOUFFETTA, Neck. Elem. i. 124 (1790).
ceratophylla, Hook. Fl. Bor. Am. i. 290 = Valeriana edulis.
coronata, Fisch. Hort. Gorenk. ed. II. (1812) 43 = sibirica.
gibbosa, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 67.—Japon.
graveolens, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 224 = heterophylla.
heterophylla, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 109.—China.
hispida, *Bunge, Pl. Mongholico-Chin.* 25. t. 3 = scabiosaefolia.
intermedia, *Roem. & Schult. Syst.* iii. 90.—Sibir.
japonica, Miq. in Arch. Néerl. v. (1870) 96 = Valeriana japonica.
Jatamansi, D. Don, Prod. Fl. Nep. 159 = Nardostachys Jatamansi.
longifolia, Macnab, in Edinb. N. Phil. Journ. xix. (1835) 59 = Valeriana edulis.
monandra, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 210.—Reg. Himal.
nudiuscula, Fisch. Hort. Gorenk. ed. II. (1812) 43 = intermedia.
ovata, Bunge, Pl. Mongholico-Chin. 23. t. 2 = villosa.
palmata, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 66.—Japon.
palmatifida, Miq. in Verh. Kon. Akad. Wetensch. II. iv. (1870) 22, in syn., sphalm. = praec.
parviflora, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 195 = scabiosaefolia.
rupestris, Bunge, in Ledeb. Fl. Alt. i. 130 = intermedia.
rupestris, Steud. Nom. ed. II. ii. 277, sphalm. = seq.
ruthenica, Juss. in Ann. Mus. Par. x. (1807) 311 = sibirica.
scabiosaefolia, [*Fisch. ex*] *Link, Enum. Hort. Berol.* i. 131.—Dahuria.
scabra, *Bunge, Pl. Mongholico-Chin.* 20. t. 1.—China.
serratulaefolia, Fisch. ex DC. Prod. iv. 624 = scabiosaefolia.
sibirica, *Juss. in Ann. Mus. Par.* x. (1807) 311.—Sibir.
triloba, Miq. in Verh. Kon. Akad. Wetensch. II. iv. (1870) 22 = palmata.
villosa, *Juss. in Ann. Mus. Par.* x. (1807) 311.—Japon.

PATRINIA, Rafin. in Journ. Phys. lxxxix. (1819) 97 = **Sophora**, Linn. (Legumin.).
sericea, Rafin. l. c. = S. sericea.

PATRISIA, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = **Ryania**, Vahl (Bixin.).
affinis, H. B. & K. Nov. Gen. et Sp. v. 357 = R. dentata.
bicolor, DC. Prod. i. 256. = Guian.
dentata, H. B. & K. Nov. Gen. et Sp. v. 357 = R. dentata.
parviflora, DC. Prod. i. 256 = R. parviflora.
pyrifera, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = R. speciosa.

PATRISIA, Rohr, ex Steud. Nom. ed. II. i. 342 (1840) = **Chailletia**, DC.

PATROCLES, Salisb. Gen. Pl. Fragm. 100 (1866) = **Narcissus**, Tourn. (Amaryll.).
orientalis, Salisb. l. c. = N. Tazetta.

PATSJOTTI, Adans. Fam. ii. 84 (1763) = **Strumpfia**, Jacq. (Rubiac.).

PATTARA, Adans. Fam. ii. 447 (1763) = **Embelia**, Burm. f. (1768) (Myrsin.).
acuta, Rafin. Sylva Tellur. 11 = E. robusta.
Basal, Rafin. l. c. = E. robusta.

PATTERSONIA, J. F. Gmel. Syst. 925 (1791) = **Ruellia**, Linn. (Acanth.).
carolinensis, J. F. Gmel. l. c. = R. strepens?

PATTONIA, Wight, Ic. t. 1750 (1852) = **Grammatophyllum**, Blume (Orchid.).
macrantha, Wight, l. c. = G. speciosum.

PATYA, Neck. Elem. i. 296 (1790) = **Verbena**, Linn.

PAULETIA, Cav. Ic. v. 5. t. 409 (1799) = **Bauhinia**, Linn. (Legumin.).
aculeata, Cav. l. c. 6. t. 410 = B. Pauletia.
bicolor, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 117. t. 6 = B. bicolor.
cheilantha, Bong. l. c. 120 = B. cheilantha.
cuyabensis, Bong. l. c. 125 = B. cuyabensis.
dodecandra, Bong. l. c. 129 = B. dodecandra.
ferruginea, Bong. l. c. 119 = B. Bongardi.
fusconervis, Bong. l. c. 116 = B. fusconervis.
glandulosa, H. B. & K. Nov. Gen. et Sp. vi. 314 = B. glandulosa.
glaucescens, H. B. & K. l. c. 316 = B. glaucescens.
grandifolia, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 130 = B. grandifolia.
hirsuta, Bong. l. c. 134 = B. hirsuta.
holophylla, Bong. l. c. 121 = B. holophylla.
inermis, Cav. Ic. v. 6 = B. inermis.
longiflora, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 122 = B. forficata.
longifolia, Bong. l. c. 128 = B. longifolia.
marginata, Bong. l. c. 127 = B. marginata.
mollis, Bong. l. c. 133 = B. mollis.
multinervia, H. B. & K. Nov. Gen. et Sp. vi. 316 = B. multinervia.
ovata, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 132 = B. ovata.
pentandra, Bong. l. c. 126 = B. pentandra.
picta, H. B. & K. Nov. Gen. et Sp. vi. 316 = B. picta.
rufa, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 123 = B. rufa, *Steud.*

PAULINIA, Gled. in Mém. Acad. Berl. (1751) 125 = seq.

PAULLINIA, Linn. Gen. ed. I. 116 (1737). *SAPINDACEAE*, Benth. & Hook. f. i. 394.
ENOUREA, Aubl. Pl. Gui. i. 587 (1775).
SEMARILLARIA, Ruiz & Pav. Prod. iv. 54. t. 9 (1794).
acutangula, *Pers. Syn.* i. 443.—Peruv.
affinis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 377. t. 78 A.—Bras.
africana, G. Don, Gen. Syst. i. 661 = Serjania curassavica.
alata, *G. Don, l. c.* 660.—Peruv.
angustifolia, Poir. Encyc. v. 101 = Serjania angustifolia, *Willd.*
anisoptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 397.—Guian.
asiatica, Linn. Sp. Pl. 365 = Toddalia aculeata.
aurea, Hort. Calc. ex Radlk. Monog. Serjan. 59 = Koelreuteria paniculata.
australis, A. St. Hil. Pl. Rém. Brés. 236 = Serjania perulacea.
baetadensis, A. Gray, Bot. U. St. Expl. Exped. i. 249 = Serjania clematidifolia.
barbadensis, *Jacq. Enum. Pl. Carib.* 36; *Obs. Bot.* t. 61. f. 9.—Ind. occ.
belangeroides, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 338 = Serjania Laruotteana.
Berteriana, Balb. ex DC. Prod. i. 602 = Urvillea Berteriana.
bipinnata, *Poir. Encyc.* v. 99.—Bras.
biternata, D. Dietr. Syn. Pl. ii. 1314, sphalm. = Serjania diversifolia.
brachystachya, *Griseb. in Goett. Abh.* xix. (1874) 109. —Reg. Argent.
brasiliensis, *Lodd. ex Radlk. Monog. Serjan.* 67.—Bras.
Cambessedesii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 352, in obs.—Bras.
capitata, *Benth. ex Triana & Planch. l. c.* 352.—N. Granat.

PAULLINIA:—
capreolata, Radlk. Monog. Serjan. 34, 70 = connarifolia.
caracasana, Jacq. Hort. Schoenb. i. 52 = Serjania caracasana.
caribaea, Jacq. Obs. Bot. iii. 11 = Serjania mexicana.
carpopoda, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 376. t. 78 B.—Bras.
cartagenensis, Balb. Cat. Hort. Taur. (1807) 43 = Serjania polyphylla.
cartagenensis, Dum.-Cours. Bot. Cult. ed. II. iv. 549 = Serjania subdentata.
cartagenensis, Jacq. Obs. Bot. iii. 11 = Serjania curassavica.
cartagenensis, Vis. Ort. Bot. Padov. (1842) 104 = Serjania triquetra.
caudata, *Vell. Fl. Flum.* 160; iv. t. 31.—Bras.
cauliflora, *Jacq. Ic. Pl. Rar.* iii. 3. t. 458.—Am. austr.
clavigera, *Schlecht. in Linnaea*, x. (1835–36) 239. —Mexic.
connaracea, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 354.—N. Granat.
connarifolia, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 349.—Guian.
coriacea, *Casar. Nov. Stirp. Bras. Dec.* iii. 27.—Bras.
costata, *Schlecht. & Cham. in Linnaea*, v. (1830) 216. —Mexic.
Cupana, *H. B. & K. Nov. Gen. et Sp.* v. 117.—Venezuela.
cupaniaefolia, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 349.—Venezuela.
curassavica, *Jacq. Obs. Bot.* iii. 12. t. 61; *Sw. Obs.* 151. —Ind. or.
curassavica, Linn. Sp. Pl. 366, partim = Serjania curassavica.
curassavica, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Serjania polyphylla.
Cururu, Linn. Sp. Pl. 365 = Serjania nodosa.
densiflora, *Sm. in Rees, Cycl.* xxvi. n. 3.—N. Granat.
dentata, Vell. Fl. Flum. 159; iv. t. 28 = Serjania dentata.
discolor, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 337.—Bras.
divaricata, Sw. Prod. Veg. Ind. Occ. 64 = Serjania mexicana, *Willd.*
diversifolia, Jacq. Enum. Pl. Carib. 36 = Serjania diversifolia.
diversifolia, Miq. in Hoev. & De Vriese, Tijdschr. x. (1843) 87 = Serjania curassavica.
elegans, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 370. —Bras.
elegans, Griseb. in Goett. Abh. xvii. (1879) 79 = Serjania caracasana.
enneaphylla, *G. Don, Gen. Syst.* i. 662.—Peruv.
erianthra, *Benth. ex Radlk. Monog. Serjan.* 75.—Peruv.
eriocarpa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 354.—N. Granat.
erythrocarpa, *Casar. Nov. Stirp. Bras. Dec.* iii. 27.—Bras.
faginea, *Radlk. Monog. Serjan.* 76.—N. Granat.
falcata, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 530.—Bras.
ferruginea, *Casar. Nov. Stirp. Bras. Dec.* iii. 28.—Bras.
fibulata, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 349.—Guian.
fluminensis, Vell. Fl. Flum. 161; iv. t. 37 = thalictrifolia.
fraxinifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 363.—N. Granat.
fusca, Griseb. in Bonplandia, vi. (1858) 3 = velutina.
fuscescens, H. B. & K. Nov. Gen. et Sp. v. 120 = velutina.
gigantea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 37.—Chili.
glabra, Bertol. Fl. Guatim. 13 = Serjania caracasana.
granatensis, *Radlk. Monog. Serjan.* 76.—N. Granat.
grandiflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 372.—Bras.
grandifolia, *Benth. ex Radlk. Monog. Serjan.* 75.—Bras.
Guarumima, Vell. Fl. Flum. 160; iv. t. 35 = Serjania cuspidata.
guatemalensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 268.—Guatemala.
hispida, *Jacq. Hort. Schoenb.* iii. 268.—Am. austr.
hispida, Rossi, Cat. Pl. Hort. Reg. Modoet. (1826) 49 = Serjania triquetra.

PAULLINIA:—
Hostmanni, Steud. in Flora, xxvii. (1844) 725 = Serjania curassavica.
ingaefolia, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 349.—Ind. occ.; Guian.
interrupta, *Benth. in Hook. Kew Journ.* iii. (1851) 195.—Bras.
jamaicensis, Macfad. Fl. Jamaic. i. 158 = curassavica.
japonica, Thunb. Fl. Jap. 170 = Vitis serjaniaefolia.
lachnocarpa, *Benth. ex Radlk. Monog. Serjan.* 75.—Ecuador.
laciniata, *Hort. Cels, ex C. C. Gmel. Hort. Bad.* (1811) 197, nomen.—Am. austr.
lactescens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 37. —Peruv.
latifolia, *Benth. ex Radlk. Monog. Serjan.* 75.—Bras.
leiocarpa, *Griseb. Fl. Brit. W. Ind.* 124.—Ind. occ.
lucida, Poir. Encyc. v. 102 = Serjania polyphylla.
lucida, Salisb. Prod. 280 = curassavica.
lupulina, Poir. Encyc. v. 101 = Serjania diversifolia.
macrophylla, Cambess. in Mém. Mus. Par. xviii. (1829) 23 = Schmidelia macrophylla.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 114.—N. Granat.
macropoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = longipes.
macrostachya, Hemsl. Biol. Centr. Am. Bot. i. 210, sphalm. = Serjania polystachya, *Radlk.*
marginata, *Casar. Nov. Stirp. Bras. Dec.* iii. 28.—Bras.
maritima, *Vell. Fl. Flum.* 161; iv. t. 36.—Bras.
meliaefolia, *Juss. in Ann. Mus. Par.* iv. (1804) 347.—Bras.
mexicana, Jacq. Obs. Bot. iii. 11. t. 61. f. 5 = Serjania angustifolia.
mexicana, Linn. Sp. Pl. 366 = Serjania mexicana, *Willd.*
Meyeniana, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 312 = Serjania cuspidata.
micrantha, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 373.—Bras.
micropterygia, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 89.—Guian.
mollis, H. B. & K. Nov. Gen. et Sp. v. 120 = Serjania amplifolia.
monogyna, *Radlk. Monog. Serjan.* 74.—Hab.?
multiflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 379.—Bras.
neglecta, *Radlk. Monog. Serjan.* 71.—Hab.?
nitida, Cambess. ex Radlk. l. c. 72 = tricornis.
nitida, *H. B. & K. Nov. Gen. et Sp.* v. 115.—Venezuela.
nitida, Steud. in Flora, xxvi. (1843) 756 = Serjania curassavica.
nodosa, Jacq. Enum. Pl. Carib. 35; Obs. Bot. iii. 11 = Serjania nodosa.
obliqua, Ruiz & Pav. ex Radlk. Monog. Serjan. 54 = Thouinia obliqua.
obovata, *Pers. Syn.* i. 443.—Peruv.
obtusa, *Vell. Fl. Flum.* 161; iv. t. 38.—Bras.
oceanica, *Hort.—Cf. Gard. Chron.* (1875) i. 589.—Ins. Pacif.
pachycarpa, *Benth. in Hook. Kew Journ.* iii. (1851) 196.—Bras.
paullinioides, Radlk. Monog. Serjan. 75 = Castanella paullinioides.
pinnata, Linn. Sp. Pl. 366 = Serjania curassavica.
Plumierii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 360.—N. Granat.
podocarpa, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1180, nomen.—Guian.
polyphylla, Aubl. ex Steud. Nom. ed. I. 597 = Serjania polyphylla.
polyphylla, Linn. Sp. Pl. 366 = Serjania diversifolia.
polyphylla, *Schum. in Act. Soc. Hist. Nat. Hafn.* iii. ii. (1792) 462.—Ind. occ.
polyphylla, Sm. in Rees, Cyclop. xxvi. n. 8 = Serjania trachygona.
polystachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 268.—Mexic.
protracta, Steud. in Flora, xxvii. (1844) 725 = Serjania paucidentata.
pterocarpa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 356.—N. Granat.
pterophylla, *Triana & Planch. l. c.* 354.—N. Granat.
pteropoda, [*Moç. & Sesse, ex*] *DC. Prod.* i. 605.—Mexic.
pubescens, *DC. l. c.* 606.—Am. austr.

PAULLINIA:—
racemosa, Poir. Encyc. v. 101 = Serjania racemosa.
racemosa, *Vell. Fl. Flum.* 159; iv. t. 29.—Bras.
racemosa, *Wawra, in Oestr. Bot. Zeitschr.* xxix. (1879) 215.—Bras.
rhizantha, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 36.—Peruv.
riparia, H. B. & K. Nov. Gen. et Sp. v. 116 = Serjania nodosa.
riparia, Radlk. Monog. Serjan. 75 = Castanella riparia.
Rossii, Lodd. ex Radlk. l. c. 67 (Quid?).—Hab.?
rubicaulis, Pav. ex Radlk. l. c. 254 = Serjania rubicaulis.
rubiginosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 371.—Bras.
rufescens, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 349.—Guian.
rugosa, *Benth. ex Radlk. Monog. Serjan.* 75.—Bras.
scabra, *Benth. l. c.*—Bras.
seminuda, *Radlk. l. c.* 47.—Bras.
senegalensis, Juss. in Ann. Mus. Par. iv. (1804) 348 = Serjania curassavica.
Seriana, Gaertn. Fruct. i. 381 = barbadensis.
Seriana, Linn. Sp. Pl. 365 = Serjania sinuata.
Seriana, Vell. Fl. Flum. 159; iv. t. 27 = coriacea.
sericea, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 374. t. 77 A.—Bras.
serjaniaefolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 356.—N. Granat.
sinuata, Lam. Illustr. ii. 446 = Serjania sinuata.
sorbilis, Mart. Reise Bras. i. 311, nomen = P. Cupana.
spectabilis, Poir. Encyc. v. 101 = Serjania mexicana, *Willd.*
sphaerocarpa, *Rich. ex Juss. in Ann. Mus. Par.* iv. (1804) 348.—Ind. occ.; Guian.
spicata, *Benth. in Hook. Kew Journ.* iii. (1851) 193. —Bras.
stenopetala, *Sagot, in Ann. Sc. Nat.* Sér. VI. xii. (1881) 194.—Guian.
stipularis, *Benth. ex Radlk. Monog. Serjan.* 75.—Bras.
subalata, *A. Gray, in S. Wats. Bibliog. Ind. N. Am. Bot.* i. 180.—Am. bor.
subcordata, *Benth. ex Radlk. Monog. Serjan.* 75.—Bras.
subrotunda, *Pers. Syn.* i. 443.—Peruv.
Tamuya, *Vell. Fl. Flum.* 160; iv. t. 34.—Bras.
tenera, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 37. t. 243. f. a, b.—Peruv.
tetragona, *Aubl. Pl. Gui.* i. 355.—Guian.
thalictrifolia, DC. Prod. i. 605, sub n. 12 = Serjania filicifolia.
thalictrifolia, *Juss. in Ann. Mus. Par.* iv. (1804) 347. —Am. trop.
Timbo, Vell. Fl. Flum. 160; iv. t. 32 = multiflora.
tomentosa, *Jacq. Enum. Pl. Carib.* 37; *Mill. Gard. Dict.* ed. VIII. n. 6.—Ind. occ.
tricornis, *Radlk. Monog. Serjan.* 44.—Guina.
trigonia, *Vell. Fl. Flum.* 159; iv. t. 30.—Bras.
triptera, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 356.—N. Granat.
triquetra, Hort. Madr. ex Dum.-Cours. Bot. Cult. ed. II. iv. 549 = Serjania triquetra.
triternata, H. B. & K. Nov. Gen. et Sp. v. 120 = Serjania diversifolia.
triternata, Jacq. Select. Am. 110 = Serjania polyphylla.
triternata, Linn. Mant. ii. 236 = Serjania triternata.
triternata, Steud. in Flora, xxvi. (1843) 756 = Serjania paucidentata.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* v. 114.—N. Granat.
uvata, Schum. & Thonn. Beskr. Guin. Pl. 195 = Serjania curassavica.
velutina, *DC. Prod.* i. 605.—N. Granat.
velutina, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 397 = Serjania polystachya.
Vespertilio, *Sw. Prod. Veg. Ind. Occ.* 64.—Ind. occ.
vetulina, G. Don, Gen. Syst. i. 662 = velutina.
weinmanniaefolia, A. Gray, in Bot. U. St. Expl. Exped. i. 247 = discolor.
weinmanniaefolia, *Mart. in Flora*, xx. (1837) II. *Beibl.* 91.—Bras.

PAULO-WILHELMIA, Hochst. in Flora, xxvii. (1844) 17 = **Mollugo**, Linn. (Ficoid.).
triphylla, Hochst. l. c. = M. Pauli-Wilhelmi.

PAULO-WILHELMIA, Hochst. in Flora, xxvii. (1844) Beibl. 4. *ACANTHACEAE*, Benth. & Hook. f. ii. 1079.
EPISANTHERA, Hochst. ex Nees, in DC. Prod. xi. 208, in syn. (1847).
polysperma, *Benth. in Hook. Niger Fl.* 479.—Afr. trop.
speciosa, *Hochst. in Flora*, xxvii. (1844) *Beibl.* 5.—Abyssin.

PAULOWNIA, Sieb. & Zucc. Fl. Jap. i. 25. t. 10 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 939.
imperialis, *Sieb. & Zucc. l. c.*—Japon.
tomentosa, Steud. Nom. ed. II. ii. 278 = imperialis.

PAURIDIA, Harv. Gen. S. Afr. Pl. 341 (1838). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 716.
hypoxidioides, *Harv. l. c.* 342.—Afr. austr.

PAURIDIANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 69 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
canthiifolia, *Hook. f. l. c.* 70.—Afr. trop.

PAUSANDRA, Radlk. in Flora, liii. (1870) 92. t. 2. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 298.
Martini, *Baill. Adansonia*, xi. (1873) 92.—Guian.
megalophylla, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 504.—Bras.
Morisiana, *Radlk. in Flora*, liii. (1870) 92. t. 2.—Bras.
Trianae, *Baill. Adansonia*, xi. (1873) 92.—N. Granat.

PAUSIA, Rafin. Fl. Tellur. iv. 105 (1836) = **Thymelaea**, Tourn.
orientalis, Rafin. l. c. = T. orientalis.

PAUSIA, Rafin. Sylva Tellur. 9 (1838) = **Osmanthus**, Lour. (Oleac.).
americana, Rafin. l. c. = O. americana.
odorata, Rafin. l. c. = O. fragrans.

PAUTSAUVIA, Juss. in Mém. Mus. Par. iii. (1817) 443 = **Marlea**, Roxb. (Cornac.).

PAVETTA, Linn. Diss. Dass. 4 (1747); Amoen. Acad. i. 388 (1749). *RUBIACEAE*, Benth. & Hook. f. ii. 114.
ACMOSTIMA, Rafin. Sylva Tellur. 26 (1838).
BACONIA, DC. in Ann. Mus. Par. ix. (1807) 219.
CHARPENTIERA, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 346.
CRINITA, Houtt. Handleid. vii. 361. t. 40. f. 1 (1777).
VERULAMIA, DC. ex Poir. Encyc. viii. 543 (1808).
abyssinica, *Fresen. in Mus. Senckenb.* ii. (1837) 166; *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 352.—Abyssin.
Ackeringae, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 31 = Ixora cuneifolia.
acuminata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 263.—Malaya.
acutifolia, *Reinw. ex Korth. l. c.* 262.—Java.
adelensis, *Delile, in Rochet, Sec. Voy. Choa*, 343.—Afr. trop.
alba, Vahl, Symb. Bot. iii. 11 = indica.
amboinica, Blume, Bijdr. 949 = Ixora amboinica.
amplexicaulis, *Pers. Syn.* i. 131.—Ind. or.
angustifolia, *Roem. & Schult. Syst.* iii. 175.—Ind. or.
angustifolia, Thw. Enum. Pl. Zeyl. 156 = hispidula.
anthophylla, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 181.—Madag.
arborea, Blanco, Fl. Filip. ed. I. 61 = Ixora stricta.
arborescens, *Hassk. Retzia*, i. 22.—Java.
arenosa, *Lour. Fl. Cochinch.* 73.—China.
aristata, Wall. Cat. n. 6169 = Webera stellulata?
assimilis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 20.—Afr. austr.
Baconia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 176.—Afr. trop.
Bandhuca, Miq. Fl. Ind. Bat. ii. 266 = Ixora coccinea.
barbata, Sm. in Rees, Cyclop. xxvi. n. 6 (= *Polyosus barbatus*).—'Ins. Hominoa.'
bidentata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 170.—Afr. trop.

PAVETTA:—
borbonica, *Hort.—Cf. Gard. Chron.* (1878) II. 535.—Ins. Borbon.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 200.—Borneo.
Bowkeri, *Harv. Thes. Cap.* ii. 21.—Afr. austr.
brachycalyx, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 169.—Afr. trop.
breviflora, *DC. Prod.* iv. 491.—Ind. or.
Brunonis, Miq. Fl. Ind. Bat. ii. 27 = Ixora Brunonis.
Brunonis, *Wall. Cat.* n. 6172.—Ind. or.
Brunonis, Wight, Ic. t. 1065 = indica.
caffra, *Linn. f. Suppl.* 121; *Thunb. Prod. Pl. Cap.* 29.—Afr. austr.
calycina, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 265.—Sumatra.
canarica, Bedd. Forest. Man. Bot. 134/8 = Webera canarica?
canescens, R. Br. in Wall. Cat. n. 6181 = indica.
canescens, *DC. Prod.* iv. 492.—Afr. trop.
cerberaefolia, Miq. Fl. Ind. Bat. ii. 279 = Webera odorata.
cerniflora, Zipp. ex Span. in Linnaea, xv. (1841) 318; et ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 195 = indica.
chinensis, Roem. & Schult. Syst. iii. 526 = Ixora chinensis.
ciliolata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 261.—Borneo.
cinerea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 181.—Madag.
coccinea, Blume, Bijdr. 950 = Ixora coccinea.
coffeacea, Wall. Cat. n. 7290 = Webera attenuata?
compactiflora, *Hook. f. Fl. Brit. Ind.* iii. 153.—Burma.
congesta, R. Br. in Salt, Abyss. App. 62 = abyssinica.
congesta, Miq. Fl. Ind. Bat. ii. 269 = Ixora congesta.
Cooperi, *Harv. & Sond. Fl. Cap.* iii. 19.—Afr. austr.
coriacea, Boj. Hort. Maurit. 172 = Rutidea coriacea.
Cornelia, Reichb. ex DC. Prod. iv. 473 (= *Canthium Cornelia*).—Afr. trop.
corymbosa, Houtt. ex Roem. & Schult. Syst. iii. 175, in syn. = caffra.
crebrifolia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 172.—Afr. trop.
diversifolia, Boj. Hort. Maurit. 172 = Rutidea coriacea.
dolichosepala, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 174.—Afr. trop.
doreensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 31.—N. Guin.
dubia, *Endl. in Ann. Wien. Mus.* i. (1836) 176.—N. Hebrid.; Ins. Amic.
edentula, *Sond. in Harv. & Sond. Fl. Cap.* iii. 20.—Afr. austr.
elliptica, Hochst. in Flora, xxvii. (1844) 98 = Feretia apodanthera.
erubescens, Miq. Fl. Ind. Bat. ii. 270 = Ixora nigricans.
Finlaysoniana, Wall. Cat. n. 6177 = indica.
floribunda, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 193.—Borneo.
foetidissima, Cyril. Pl. Rar. Neap. Fasc. i. 7 = Putoria calabrica.
fulgens, Miq. Fl. Ind. Bat. ii. 204 = Ixora fulgens.
gardeniaefolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 351.—Abyss.
genipaefolia, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 78 = P. Baconia.
Gerrardii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 20.—Afr. austr.
glauca, Boj. Hort. Maurit. 172 = cinerea.
glaucescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 171.—Afr. trop.
glaucina, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 245.—Ins. Banca.
Gleniei, *Thw. ex Hook. f. Fl. Brit. Ind.* iii. 152.—Zeylan.
graciliflora, Wall. Cat. n. 6178 = indica.
gracilipes, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 179.—Afr. trop.
gracilis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 290.—Afr. trop.
gracilis, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 262.—Borneo.
gracilis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 181.—Madag.
grandifolia, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 264.—Sumatra.

PAVETTA:—
hispida, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 175.—Afr. trop.
hispidula, *Wight & Arn. Prod.* 431.—Ind. or.
Hookeriana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 176.—Afr. trop.
humilis, *Hook. f. Fl. Brit. Ind.* iii. 151.—Malacca.
hypsophila, Miq. Fl. Ind. Bat. ii. 273 = longituba.
incana, Klotzsch, in Peters, Reise Mossamb. Bot. 289 = gracilis, *Klotzsch.*
incarnata, Blume, Bijdr. 950 = Ixora coccinea.
indica, Burm. f. Fl. Ind. 35. t. 13. f. 3 = angustifolia.
indica, *Linn. Sp. Pl.* 110.—As. et Austral. trop.
involucrata, *Thw. Enum. Pl. Zeyl.* 156.—Zeylan.
jambosaefolia, *Teijsm. & Binn. Cat. Hort. Bog.* 112.—Ins. Celebes.
javanica, Blume, Bijdr. 949 = Ixora javanica.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 197.—Sumatra.
Kroneana, *Miq. in Journ. Bot. Néerl.* i. (1861) 107.—China.
Kurziana, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 246.—Ins. Banca.
laevis, *Benth. in Hook. Niger Fl.* 415.—Afr. trop.
lanceolata, *Eckl. in South Afr. Quart. Journ.* (1830) 374.—Afr. austr.
lateriflora, G. Don, Gen. Syst. iii. 575 = Memecylon Donianum.
leptantha, *Zipp. ex Span. in Linnaea*, xv. (1841) 318, nomen.—Ins. Timor.
leucoxylon, *Miq. Fl. Ind. Bat.* ii. 275.—Java.
littorea, *Miq. l. c.* 272.—Ins. Flores.
Lobbii, Teijsm. & Binn. Cat. Hort. Bog. 112 = Ixora fulgens.
longiflora, Blume, Bijdr. 952 = macrantha.
longiflora, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 351 = gardeniaefolia.
longiflora, *Vahl, Symb. Bot.* iv. 12.—Arabia.
longifolia, Miq. Fl. Ind. Bat. ii. 275 = Webera longifolia.
longipes, *DC. Prod.* iv. 490.—Ins. Timor.
longituba, *Miq. Fl. Ind. Bat.* ii. 278.—Java.
lucens, R. Br. in Wall. Cat. n. 6168 = Webera lucens.
macrantha, *Steud. Nom.* ed. II. ii. 278.—Java.
macrocoma, Miq. Fl. Ind. Bat. ii. 274 = Ixora grandifolia.
macrophylla, Blume, Bijdr. 953 = Ixora grandifolia.
macroptera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 197.—Sumatra.
macrosepala, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 172.—Afr. trop.
macrothyrsa, *Teijsm. & Binn. in Tijdsch. Nederl. Ind.* xxix. (1867) 246.—Java.
manillensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 351.—Ins. Philipp.
Mannii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 169.—Afr. trop.
membranacea, Blanco, Fl. Filip. ed. I. 59 = angustifolia.
mollis, *Afzel. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 174.—Afr. trop.
mollis, Wall. Cat. n. 6179 = indica.
montana, *Reinw. ex Blume, Bijdr.* 952.—Java.
monticola, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 170.—Afr. trop.
natalensis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 19.—Afr. austr.
naucleaeflora, G. Don, Gen. Syst. iii. 575 = seq.
naucleiflora, *R. Br. in Wall. Cat. n.* 6171 (naucliflora).—Ind. or.; Burma; Malaya.
neurocarpa, *Benth. in Hook. Niger Fl.* 414.—Afr. trop.
nigricans, *Miq. Fl. Ind. Bat.* ii. 272.—Java.
nilagirica, Bedd. Forest. Man. Bot. 134/8 = Webera nilagirica.
oblonga, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 263.—Borneo.
oblonga, Wall. Cat. n. 6170 (Weberae sp.).—Siam.
obovata, *E. Mey. in Drège, Zwei Pfl. Docum.* 208, nomen.—Natal.
obtusa, Pers. Syn. i. 131 = indica.
odorata, *Blume, Bijdr.* 952.—Java.
odorata, Zoll. ex Miq. Fl. Ind. Bat. ii. 204 (= *Stylocoryna polycarpa*).—Java.
Oliveriana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 174.—Afr. trop.
opaca, Miq. Fl. Ind. ii. 270 = Ixora opaca.
opulina, *DC. Prod.* iv. 492.—N. Caled.

PAVETTA:—
owariensis, *Beauv. Fl. Owar.* i. 87. t. 52.—Afr. trop.
paludosa, *Blume, Bijdr.* 954.—Java.
paludosa, Miq. Fl. Ind. Bat. ii. 271 = Ixora paludosa.
parasitica, *Lour. Fl. Cochinch.* 92.—Cochinch.
parviflora, *Afzel. Rem. Guin.* viii. 47.—Afr. trop.
parviflora, Smeathm. ex DC. Prod. iv. 492 = Smeathmanni.
parvifolia, *Vidal, Phan. Cuming. Philipp.* 183.—Ins. Philipp.
pauciflora, *Blume, ex DC. Prod.* iv. 487; *Miq. Fl. Ind. Bat.* ii. 268.—Java.
pendula, Miq. l. c. 269 = Ixora pendula.
pentandra, Sw. Fl. Ind. Occ. i. 233 = Palicourea Pavetta.
petiolaris, Wall. Cat. n. 6186 = indica.
polita, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 196.—Borneo.
polyantha, *R. Br. in Salt, Abyss. App.* 62.—Abyss.
polyantha, Wall. Cat. n. 6176 = indica.
Pseudo-amboinica, *Korth. in Kruidk. Nederl. Arch.* ii. II. (1851) 261.—Borneo.
puberula, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 171.—Afr. trop.
pulcherrima, *Teijsm. & Binn. Cat. Hort. Bog.* 113.—Ins. Banca; Sumatra.
reflexa, R. Br. in Salt, Abyss. App. 62 = gardeniaefolia.
reticulata, *Blume, Bijdr.* 953.—Java.
revoluta, *Hochst. in Flora*, xxv. (1842) 237.—Afr. austr.
rigida, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 178.—Afr. trop.
Rothiana, DC. Prod. iv. 491 = indica.
rubra, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = Ixora coccinea.
salicifolia, Blume, Bijdr. 951 = Ixora fulgens.
sambucina, Blanco, Fl. Filip. ed. II. 41 = angustifolia.
sambucina, *DC. Prod.* iv. 492.—Ins. Amicor.
secundiflora, *Zipp. ex Span. in Linnaea*, xv. (1841) 318, nomen.—Ins. Timor.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 107 = indica.
siphonantha, Dalz. in Hook. Kew Journ. ii. (1850) 133 = hispidula.
Smeathmanni, *DC. Prod.* iv. 492.—Afr. trop.
stenophylla, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 262.—Borneo.
stricta, Blume, Bijdr. 950 = Ixora stricta.
subcana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 172.—Afr. trop.
subcapitata, *Hook. f. Fl. Brit. Ind.* iii. 150.—Reg. Himal.
subcuneata, *Miq. Fl. Ind. Bat. Suppl.* 546.—Sumatra.
subglabra, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 78.—Afr. trop.
subulata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 403.—Java.
subvelutina, *Miq. Fl. Ind. Bat.* ii. 276.—Java.
sumatrensis, *Roth, Nov. Pl. Sp.* 88.—Sumatra.
sylvatica, *Blume, Bijdr.* 953.—Java.
syringoides, *Webb, in Hook. Kew Journ.* ii. (1850) 371.—Ins. Cap. Virid.
Teijsmanni, Hassk. in Nederl. Kruidk. Arch. iv. (1859) 4 = Ixora grandifolia?
tenuiflora, *Miq. Fl. Ind. Bat.* ii. 268.—Ins. Molucc.
tenuifolia, *Benth. in Hook. Niger Fl.* 415.—Afr. trop.
ternata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 246.—Malaya.
ternifolia, *Hook. f. ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 86. t. 51.—Afr. trop.
thyrsiflora, Thunb. ex DC. Prod. iv. 492 = caffra.
timorensis, *Miq. Fl. Ind. Bat.* ii. 278.—Ins. Timor.
tomentosa, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 181 = canescens.
tomentosa, Roxb. ex Sm. in Rees, Cyclop. xxvi. n. 2 = indica.
triflora, *DC. Prod.* iv. 492.—Ins. Tahiti.
undulata, *Lehm. in Hamb. Gartenz.* xiv. (1858) 437; *et in Linnaea*, xxix. (1857–58) 726.—Hab.?
velutina, Wall. Cat. n. 6174 = indica.
villosa, Heyne, in Roth, Nov. Pl. Sp. 88 = indica.
villosa, *Vahl, Symb. Bot.* iii. 12.—Arabia.
Wallichiana, Steud. Nom. ed. II. ii. 279 = indica.
weberaefolia, R. Br. in Wall. Cat. n. 6182 = Webera odorata *et* Wallichii.
Wightiana, Wall. l. c. n. 6167 = Webera corymbosa.

PAVETTA:—
Wightii, *Hook. f. Fl. Brit. Ind.* iii. 152.—Ind. or.
Wyckii, Hassk. in Retzia, i. 21 = Ixora jucunda.
Wyckii, *Miq. Fl. Ind. Bat.* ii. 356.—Java.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* iii. 21.—Afr. austr.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 201.—N. Guin.

PAVIA, Boerh. ex Mill. Gard. Dict. ed. VI (1752) = **Aesculus**, Linn. (Sapind.).
alba, Poir. Encyc. v. 95 = Ae. parviflora.
americana, Jaume St. Hil. Expos. ii. 16 = Ae. Pavia.
arguta, Rafin. Alsog. Am. 74 = Ae. arguta.
atropurpurea, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 58 = Ae. Pavia.
bicolor, Rafin. Alsog. Am. 74 = Ae. flava.
californica, Hartw. in Journ. Hort. Soc. ii. (1847) 123 = Ae. californica.
carnea, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 53 = Ae. glabra.
discolor, Sweet, Hort. Brit. ed. I. 69 = Ae. flava.
edulis, Poit. Arb. Fr. t. 88; ex DC. Prod. i. 598 = Ae. parviflora.
flava, Moench, Meth. 66; DC. l. c. = Ae. flava.
fulva, Rafin. Alsog. Am. 74 = Ae. octandra (in Addend.).
glabra, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 54 = Ae. glabra.
glauca, Rafin. Alsog. Am. 71.—Am. bor.
humilis, G. Don, in Loud. Hort. Brit. 143 = Ae. Pavia?
hybrida, DC. Prod. i. 598 = Ae. flava.
indica, Wall. ex Jacquem. Voy. Bot. 31 = Ae. indica.
intermedia, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 60 = Ae. Pavia.
khassyana, Voigt, Hort. Suburb. Calc. 97 = Ae. punduana.
Lindleyana, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 59 = Ae. Pavia.
livida, Spach, l. c. 56 = Ae. Pavia?
longiflora, Rafin. Alsog. Am. 72.—Am. bor.
lucida, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 60 = Ae. Pavia.
lutea, Poir. Encyc. v. 95 = Ae. flava.
macrocarpa, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 765, nomen = Ae. Pavia.
macrostachya, DC. Prod. i. 598 = seq.
macrostachys, Loisel. Herb. Amat. iii. t. 212 = Ae. parviflora.
Michauxii, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 61 = Ae. Pavia.
mollis, Rafin. Alsog. Am. 71.—Am. bor.
mutabilis, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 57 = Ae. Pavia?
nana, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 603, nomen = Ae. Pavia.
neglecta, G. Don. ex Loud. Hort. Brit i. 143; et Gen. Syst. i. 653 = Ae. flava.
octandria, Mill. Gard. Dict. ed. VIII. n. 1 = Ae. Pavia.
ohioënsis, Michx. f. N. Am. Sylv. ii. 101. t. 29 = Ae. glabra.
pallida, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 54 = Ae. glabra.
parviflora, Rafin. Alsog. Am. 73 = Ae. parviflora.
punctata, Rafin. l. c. 72.—Am. bor.
punduana, Wall. ex Voigt, Hort. Suburb. Calc. 97 = Ae. punduana.
reticulata, Rafin. Alsog. Am. 73 = Ae. flava.
rubra, Moench, Meth. 65 = Ae. Pavia.
versicolor, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 57 = Ae. Pavia?
Watsoniana, Spach, l. c. 53 = Ae. glabra?
Whitleyi, C. Koch, Dendrol. i. 511 = Ae. Pavia.
Willdenowiana, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 59 = Ae. Pavia.

PAVIANA, Rafin. Fl. Ludov. 87 (1817) = **Aesculus**, Linn. (Sapind.).
coccinea, Rafin. l. c. = Ae. Pavia.
flava, Rafin. l. c. = Ae. flava.

PAVINDA, Thunb. ex Bartl. Ord. 374 (1830) = **Audouinia**, Brongn. (Bruniac.).
capensis, Retz. ex Steud. Nom. ed. II. ii. 279 = A. capitata.

PAVONIA, Cav. Diss. ii. App. II. (1786); et iii. 132. t. 45 (1787). *MALVACEAE*, Benth. & Hook. f. i. 205.
Asterochlaena, Garcke, in Bot. Zeit. viii. (1850) 666.
Brehmia, Schrank, in Syll. Ratisb. i. (1824) 85.
Diplopenta, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 10.
Greevesia, F. Muell. in Hook. Kew Journ. viii. (1856) 8.
Lebretonia, Schrank, Pl. Rar. Hort. Monac. t. 90 (1819).
Lopimia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 96.
Malache, B. Vog. in Trew, Pl. Select. 50. t. 90 (1772).
Pentameris, E. Mey. in Drège, Zwei Pfl. Docum. 210 (1843).
Thorntonia, Reichb. Consp. 202 (1828).
Typhalea, Neck. Elem. ii. 412 (1790).
acerifolia, Link & Otto, Ic. Pl. Sel. i. 5. t. 1 = Hibiscus collinus.
achanioides, *Griseb. Cat. Pl. Cub.* 27.—Cuba.
acuminata, Spreng. Syst. iii. 98 = Kosteletzkya acuminata.
affinis, A. St. Hil. Fl. Bras. Mer. i. 229 = sagittata.
alba, Seem. Bot. Voy. Herald, 81 = paniculata.
alnifolia, A. St. Hil. Fl. Bras. Mer. i. 241 = Goethea alnifolia.
angustifolia, *Benth. in Hook. Kew Journ. Bot.* iv. (1842) 119.—Guian.
arabica, *Hochst. ex Steud. Nom.* ed. II. ii. 279.—Sind.; Arab.; Abyss.
arachnoides, *Presl, Rel. Haenk.* ii. 129.—Mexic.
arenaria, *Roth, Catalect.* fasc. ii. 80.—Ind. or.
aristata, Cav. Diss. iii. 133 = spinifex.
Armitiana, F. Muell. Fragm. x. 78 = Urena Armitiana.
assimilis, C. Muell. in Walp. Ann. vii. 399, sphalm. = Pavetta assimilis.
begoniaefolia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 333 = rosea.
Berterii, Spreng. Syst. iii. 99 = Kosteletzkya pentasperma.
betonicifolia, *Presl, Rel. Haenk.* ii. 128.—Peruv.
Blanchetiana, *Miq. in Linnaea*, xxii. (1849) 801.—Bras.
Bojeri, *Baker, in Journ. Bot.* xx. (1882) 45.—Madag.
brachypoda, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 563 = P. Typhalaea.
brachysepala, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 44 = rosea.
bracteosa, Benth. in Hook. Journ. Bot. iv. (1842) 118 = sessiliflora.
brasiliensis, Spreng. Syst. iii. 100 = Hibiscus brasiliensis.
calyculosa, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 40.—Bras.
cancellata, *Cav. Diss.* iii. 135.—Venezuela.
caracasana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 188 = paniculata.
cardiosepala, Turcz. l. c. = varians.
castanaefolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 44.—Ind. occ.; Bras.
Cavanillesii, Spreng. Syst. v. 513 = Kosteletzkya hastata, *Garcke*.
ceratocarpa, *Dalz. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 331.—Sind.
cernua, *Walp. Rep.* ii. 790.—Ins. Timor.
clathrata, *Mast. in Oliver, Fl. Trop. Afr.* i. 193.—Afr. trop.
coccinea, *Cav. Diss.* iii. 140. t. 47. f. 1.—Ins. S. Doming.
Columella, *Cav. l. c.* 138. t. 48. f. 3.—Ins. Borbon.
communis, A. St. Hil. Fl. Bras. Mer. i. 224 = spinifex.
commutata, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 212.—Bras.
conferta, *A. St. Hil. Fl. Bras. Mer.* i. 238.—Bras.
Cooperi, C. Muell. in Walp. Ann. vii. 399, sphalm. = Pavetta Cooperi.
cordifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 27.—Cuba.
corymbosa, Willd. Sp. Pl. iii. 836 = paniculata.
crenata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 53 = Kraussiana.
crenata, Hochst. in Flora, xxvii. (1844) 100; et ex Mast. in Oliver, Fl. Trop. Afr. i. 179 = Sida triloba.

PAVONIA:—
cuneifolia, Cav. Diss. iii. 139 = praemorsa.
cymbalaria, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 42.—Bras.
dasypetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 189.—Venezuela.
deltoidea, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 59.—Bras.
dictyocarpa, Hochst. ex Webb, Fragm. Fl. Aethiop. 46 = Hibiscus dictyocarpus.
distinguenda, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 42 = sagittata.
diuretica, *A. St. Hil. Pl. Us. Brés.* t. 53.—Bras.
diversifolia, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 134; xii. (1845) 79.—Java?
Dregei, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 218.—Afr. austr.
Drummondii, Torr. & Gray, Fl. N. Am. i. 682 = Malvaviscus Drummondii.
edentula, C. Muell. in Walp. Ann. vii. 399, sphalm. = Pavetta edentula.
elegans, *Garcke, in Eichl. Jahrb. Berl.* ii. (1883) 332.—Afr. trop.
flava, Spring, ex Mart. in Flora, xx. (1837) II. Beibl. 96 = sepium.
flavispina, Miq. in Linnaea, xxii. (1849) 551 = sepium.
foliosa, Mart. ex Garcke, in Eichl. Jahrb. Berl. i. (1881) 217 = microphylla.
Gardneriana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 187 = rosea.
geminiflora, Moric. Pl. Nouv. Am. 120. t. 73 = paniculata.
Gerrardii, C. Muell. in Walp. Ann. vii. 399, sphalm. = Pavetta Gerrardi.
glandulosa, *Franch. in Revoil, Çomali*, 18.—Afr. trop.
glandulosa, *Presl, Reliq. Haenk.* ii. 129.—Mexic.
glechomifolia, *Garcke, in Schweinf. Beitr. Fl. Aethiop.* 54.—As. et Afr. trop.
glechomoides, *A. St. Hil. Fl. Bras. Mer.* i. 227. t. 45.—Bras.
glomerata, Casar. Nov. Stirp. Bras. Dec. 38 = P. Typhalaea.
Goudotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 160.—N. Granat.
grandiflora, *A. St. Hil. Fl. Bras. Mer.* i. 232.—Bras.
grandifolia, *Walp. Rep.* i. 300.—Bras.
grewioides, Hochst. ex Boiss. Fl. Orient. i. 837 = propinqua.
grisea, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 41 = speciosa.
hastata, *Cav. Diss.* iii. 138. t. 47. f. 2.—Bras.
hastata, Spreng. Syst. iii. 99 = Kosteletzkya hastata, *Garcke*.
heterophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 188.—Mexic.
hirsuta, *Guill. & Perr. Fl. Seneg. Tent.* i. 51.—Afr. trop.
hirta, Klotzsch, in Linnaea, xiv. (1840) 301 = cancellata.
hirta, *Schweigg. Enum. Pl. Hort. Regiom.* 52.—Hab.?
hirta, Spreng. Syst. iii. 99 = glechomoides.
hirtiflora, Benth. Pl. Hartw. 7 = glandulosa, *Presl*.
hispida, Spreng. Syst. iii. 99 = Kosteletzkya hispidula.
hispida, *Wright, in Sauv. Fl. Cub.* 12.—Cuba.
Hostmanni, *Miq. in Linnaea*, xxii. (1849) 469.—Guian.
humifusa, *A. St. Hil. Fl. Bras. Mer.* i. 235.—Bras.
humilis, Jacq. Fragm. 78 = odorata.
insignis, Fenzl, ex Webb, Fragm. Fl. Aethiop. 42 = hirsuta.
intermedia, *A. St. Hil. Fl. Bras. Mer.* i. 223.—Bras.
intermixta, *A. Rich. Ess. Fl. Cub.* 128.—Cuba.
involucrata, Klotzsch, in Linnaea, xiv. (1840) 301 = sessiliflora.
Jonesii, Feay, ex Wood, Class-Book (1862), 269 = hastata.
Kotschyi, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 43; *et in Webb, Fragm. Fl. Aethiop* 43.—Abyss.
Kraussiana, *Hochst. in Flora*, xxvii. (1844) 293.—Afr. trop. et austr.
lanceolata, *Schlecht. in Linnaea*, xi. (1837) 356.—Mexic.
lappacea, Casar. Nov. Stirp. Bras. Dec. 37 = rosea.
lasiopetala, *Scheele, in Linnaea*, xxi. (1848) 470.—Texas.
latifolia, *Spreng Syst.* iii. 98.—Bras.

PAVONIA:—
laxiflora, A. St. Hil. Fl. Bras. Mer. i. 226 = paniculata.
Le-Contei, Torr. & Gray, in Mem. Am. Acad. N. S. iv. (1849) 16 = hastata.
leptocarpa, Cav. Diss. vi. 351 = P. Typhalaea.
leucantha, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 211.—Bras.
linearis, *A. Rich. Ess. Fl. Cub.* 127.—Cuba.
longifolia, *A. St. Hil. Fl. Bras. Mer.* i. 239.—Bras.
Luschnathiana, Klotzsch, in Linnaea, xiv. (1840) 301, nomen = microphylla.
macrophylla, E. Mey. ex Harv. & Sond. Fl. Cap. i. 169 = Kraussiana.
macrotis, *Baker, in Journ. Linn. Soc.* xx. (1883) 98.—Madag.
Makoyana, E. Morr. in Belg. Hortic. xxviii. (1878) 59. t. 3 = Goethea Mackoyana.
malacophylla, Hemsl. Biol. Centr. Am. Bot. i. 117, in syn. = velutina.
malvaviscoides, *A. St. Hil. Fl. Bras. Mer.* i. 237.—Bras.
megalophylla, Wright, ex Griseb. Cat. Pl. Cub. 27 = velutina.
mexicana, H. B. & K. Nov. Gen. et Sp. v. 284 = Kosteletzkya mexicana.
Meyeri, Mast. in Oliver, Fl. Trop. Afr. i. 191 = P. Columella.
microphylla, *Casar. Nov. Stirp. Bras. Dec.* 38.—Am. trop.
microphylla, E. Mey. ex Harv. & Sond. Fl. Cap. i. 169 = Dregei.
modesta, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 59.—Bras.
mollis, H. B. & K. Nov. Gen. et Sp. v. 282 = Mutisii.
mollis, E. Mey. ex Harv. & Sond. Fl. Cap. i. 169 = P. Columella.
monatherica, *Casar. Nov. Stirp. Bras. Dec.* 37.—Bras.
montana, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 222, nomen.—Bras.
multiflora, *A. St. Hil. Fl. Bras. Mer.* i. 239. t. 47.—Bras.
muricata, A. St. Hil. l. c. 225. t. 44 = Schrankii.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* v. 283.—N. Granat.
natalensis, C. Muell. in Walp. Ann. vii. 400, sphalm. = Pavetta natalensis.
nemoralis, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 43 = rosea.
nudicaulis, *A. St. Hil. Fl. Bras. Mer.* i. 231.—Bras.
obovata, C. Muell. in Walp. Ann. vii. 400, sphalm. = Pavetta obovata.
odorata, *Willd. Sp. Pl.* iii. 837.—As. et Afr. trop.
ovata, Spreng. Syst. iii. 99 = Hibiscus aethiopicus.
paniculata, *Cav. Diss.* iii. 135. t. 46. f. 2.—Am. austr.
papilionacea, *Cav. l. c.* 140. t. 49. f. 2.—Ins. Societ.
paraibica, *Wawra, in Oestr. Bot. Zeitschr.* xxxii. (1882) 39.—Bras.
parviflora, Desf. Tabl. ed. I. 221; et ex Poir. Encyc. Suppl. iv. 335 = Kosteletzkya hispidula.
pellita, *H. B. & K. Nov. Gen. et Sp.* v. 282.—Venezuela.
pentacarpos, Poir. Encyc. Suppl. iv. 335 = Kosteletzkya pentacarpa.
pilosa, Willd. Enum. Hort. Berol. Suppl. 50 = Kosteletzkya hispidula.
plantanifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 99.—Madag.
platanifolia, Willd. in Ges. Naturf. Fr. Berl. Mag. iv. (1810) 220 = Hibiscus collinus.
plumosa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 190 = velutina.
Poeppigii, Schlecht. in Linnaea, xi. (1837) 357, in textu = Hibiscus Poeppigii.
polymorpha, A. St. Hil. Fl. Bras. Mer. i. 232 = speciosa.
praemorsa, *Cav. Diss.* iii. 139.—Afr. austr.
procumbens, Casar. Nov. Stirp. Bras. Dec. 39 = cancellata.
procumbens, Walp. Rep. i. 301 = glechomifolia.
propinqua, *Garcke, in Schweinf. Beitr. Fl. Aethiop.* 55.—Sind.; Beluchist.
prostrata, *Moric. Pl. Nouv. Am.* 119. t. 72.—Bras.
Pseudo-typhalaea, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 160.—N. Granat.
racemiflora, *Hook. & Arn. Bot. Beech. Voy.* 277.—Mexic.
racemosa, *Sw. Fl. Ind. Occ.* ii. 1215.—Am. trop.

PAVONIA:—
Regnelliana, Miq. in Linnaea, xxii. (1849) 551 = sepium?
repanda, Spreng. Syst. iii. 98 = Urena repanda.
reticulata, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 212.—Bras.
Romborna, Steud. Nom. ed. II. ii. 279, sphalm = seq.
Romborua, Buch.-Ham. in Wall. Cat. n. 2690 = odorata.
Rosa-campestris, *A. St. Hil. Fl. Bras. Mer.* i. 236. t. 46.—Bras.
rosea, *Schlecht. in Linnaea*, xi. (1837) 355.—Am. trop.
rosea, Wall. Cat. n. 1887; Moris, in Mem. Acc. Torin. xxxvi. (1833) 194 = odorata.
rubiformis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 189.—Ins. Philipp.
sagittata, *A. St. Hil. Fl. Bras. Mer.* i. 229.—Bras.
scabra, Benth. ex Garcke, in Eichl. Jahrb. Berl. i. (1881) 221 = speciosa.
scabra, Presl, Rel. Haenk. ii. 129 = paniculata.
Schiedeana, Steud. Nom. ed. II. ii. 279 = rosea.
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 52.—Abyssin.
Schrankii, *Spreng. Syst.* iii. 98.—Bras.
semiserrata, *Steud. Nom.* ed. II. ii. 279.—Bras.
semperflorens, *Garcke, in Eichl. Jahrb. Berl.* i. (1881) 222.—Bras.
sepium, *A. St. Hil. Fl. Bras. Mer.* i. 225.—Bras.
serrata, *Franch. in Revoil, Çomali*, 18.—Afr. trop.
sessiliflora, *H. B. & K. Nov. Gen. et Sp.* v. 281.—N. Granat.
sidaefolia, *H. B. & K. l. c.* 283.—Venezuela.
sidoides, Hornem. Hort. Hafn. ii. 658 = odorata.
somalensis, *Franch. in Revoil, Çomali*, 17.—Afr. trop.
speciosa, *H. B. & K. Nov. Gen. et Sp.* v. 281.—Venezuela.
spicata, Cav. Diss. iii. 136 = racemosa.
spinifex, *Cav. l. c.* 133.—Am. austr.
spinigera, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
spiralis, Cav. Ic. v. 20 = Malvaviscus ciliatus.
stellata, Spreng. Syst. iii. 97 = P. Typhalaea.
subhastata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 163.—N. Granat.
subpandurata, *Wright, ex Griseb. Cat. Pl. Cub.* 26.—Cuba.
subrotunda, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 42.—Bras.
sulphurea, Steud. Nom. ed. II. ii. 279 = cancellata.
surinamensis, Miq. in Hoev. & De Vriese, Tijdschr. x. (1843) 77 = P. Typhalaea.
tomentosa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 52 = Schimperiana.
tricalycaris, *A. St. Hil. Fl. Bras. Mer.* i. 240.—Bras.
triloba, Guill. & Perr. Fl. Seneg. Tent. i. 50 = zeylanica.
triloba, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 43; *et ex Webb, Fragm. Fl. Aethiop.* 43.—Afr. trop.
Typhalaea, *Cav. Diss.* iii. 134.—Ind. occ.
Typhalaea, A. St. Hil. Fl. Bras. Mer. i. 223 = rosea.
typhalaeoides, H. B. & K. Nov. Gen. et Sp. v. 279 = P. Typhalaea.
urens, *Cav. Diss.* iii. 137. t. 49. f. 1; v. 283.—Ins. Mascar.
urticaefolia, *Presl, Rel. Haenk.* ii. 128.—Mexic.
varians, *Moric. Pl. Nouv. Am.* 122. t. 74.—Bras.
velutina, *A. St. Hil. Fl. Bras. Mer.* i. 233.—Bras.
veneta, Spreng. Syst. iii. 98 = Kosteletzkya pentacarpa.
virginica, Spreng. l. c. = Kosteletzkya virginica.
viscida, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1170 = paniculata.
viscidula, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 42 = sagittata.
viscosa, *A. St. Hil. Fl. Bras. Mer.* i. 236.—Bras.
Wioti, E. Morr. in Belg. Hortic. xxv. (1875) 113. t. 7 = multiflora.
Wrightii, A. Gray, Gen. Pl. Am. Ill. ii. 76. t. 130 = lasiopetala.
Zeyheri, C. Muell. in Walp. Ann. vii. 400 = Pavetta Zeyheri.
zeylanica, *Cav. Diss.* iii. 134.—Ind. or.; Afr. trop.

PAVONIA, Domb. ex Lam. Tabl. Encyc. (Illustr.) i. 421 (1800) = **Cordia**, Linn. (Borag.).
lutea, Domb. l. c. = C. lutea.

PAVONIA, Ruiz & Pav. Prod. 127. t. 28 (1794) = **Laurelia**, Juss. (Monim.).
sempervirens, Ruiz & Pav. Syst. Veg. 253 = L. aromatica.

PAXTONIA, Lindl. Bot. Reg. (1838) Misc. 61 = **Spathoglottis**, Blume (Orchid.).
rosea, Lindl. l. c.—Ins. Philipp.

PAYANELIA, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 384 (1884) = **Pajanelia**, DC. (Bignon.).

PAYENA, A. DC. in DC. Prod. viii. 196 (1844). *SAPOTACEAE*, Benth. & Hook. f. ii. 659.
AZAOLA, Blanco, Fl. Filip. ed. I. 402 (1837), nomen prius.
CACOSMANTHUS, Miq. Fl. Ind. Bat. ii. 1040 (1857).
KAKOSMANTHUS, Hassk. Retzia, 97 (1855); Walp. Ann. v. 474 (1858).
KERATEPHORUS, Hassk. in Flora xxxviii. (1855), 578; Retzia, 100 (1855).
acuminata, Pierre, in Bull. Soc. Linn. Par. i. (1885) 528 = Mimusops acuminata.
Balem, Pierre, l. c. 530 = Bassia Balem.
bankensis, *Burck, in Ann. Jard. Buitenz.* v. (1885) 54.—Ins. Banca.
Bawun, *Scheff. l. c.* i. (1876) 33.—N. Guin.
Beccarii, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 525.—Borneo.
Benjamina, Pierre, l. c. 524 = Isonandra Benjamina.
Betis, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 125.—In. Philipp.
Boelageana, *Burck, in Ann. Jard. Buitenz.* v. (1885). 54.—Java?
Croixiana, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 524.—Malacca.
dasyphylla, Pierre, l. c. 527 = Isonandra dasyphylla.
glutinosa, *Pierre, l. c.* 529.—Borneo.
Griffithii, Kurz, For. Fl. Brit. Burma, ii. 121 = Dichopsis hexandra.
Griffithii, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 525.—Malacca.
Hillii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 368.—Madag.
Junghuhniana, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 530.—Java.
lamponga, Burck, in Ann. Jard. Buitenz. v. (1885) 48, 59 = Isonandra lamponga.
latifolia, *Burck, l. c.* 58.—Ins. Sunda.
Leerii, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 69; *Burck, l. c.* 48, 56.—Burma.
longipetiolata, *Kurz, l. c.*—Ind. or.
Lowiana, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 525.—Malaya.
lucida, *A. DC. in DC. Prod.* viii. 197.—Penang; Singapore.
lucida, Herb. Traj. ex Pierre, in Bull. Soc. Linn. Par. i. (1885) 526 = Griffithii.
lucida, Pierre, l. c. 525 = Lowiana.
macrophylla, *Burck, in Ann. Jard. Buitenz.* v. (1885) 51.—Java.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 547.—Malacca.
malaccensis, *C. B. Clarke, l. c.*—Malacca.
microphylla, Burck, in Ann. Jard. Buitenz. v. (1885) 99, 60 = Isonandra microphylla.
nigro-punctata, *Burck, l. c.* 53.—Java.
obscura, *Burck, l. c.* 60.—Sumatra.
paralleloneura, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 70.—Burma.
polyandra, *Benth. & Hook. f. Gen.* ii. 659.—Ind. or.
puberula, Pierre, ex Burck, in Ann. Jard. Buitenz. v. (1885) 50 = Isonandra puberula.
rubro-pedicellata, *Burck, l. c.* 55.—Java?
sericea, De Vriese, Pl. Reinw. 62; Miq. Fl. Ind. Bat. ii. 1039 = Bassia sericea.
stipularis, *Burck, in Ann. Jard. Buitenz.* v. (1885) 48.—Ins. Celebes.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 582.—Sumatra.
sumatrana, Pierre, in Bull. Soc. Linn. Par. i. (1885) 527 = Teysmanniana.
Suringariana, Burck, in Ann. Jard. Buitenz. v. (1885) 49 = Bassia sericea.
Teysmanniana, *Pierre, in Bull. Soc. Linn. Par.* i. (1885) 527.—Sumatra.

PAYERIA, Baill. Adansonia, i. (1860–61) 50. t. 3 = **Quivisia**, Comm. (Meliac.).
chrysogyne, Muell. Arg. in Linnaea, xxxiv. (1865–66) 65.—Ins. Borbon.
excelsa, Baill. Adansonia, i. (1860–61) 51. t. 3.—Madag.

PAYPAYROLA, Aubl. Pl. Gui. i. 249. t. 99 (1775). *VIOLARIEAE*, Benth. & Hook. f. i. 118.
LIGNONIA, Scop. Introd. 292 (1777).
PAYROLA, Juss. Gen. 472 (1789).
PERICLISTIA, Benth. in Hook. Journ. Bot. iv. (1842) 108.
WIBELIA, Pers. Syn. i. 210 (1805).
Blanchetiana, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 371.—Bras.
brasiliensis, *Steud. Nom.* ed. II. ii. 280.—Bras.
confertiflora, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 373.—Guian.
grandiflora, *Tul. l. c.* 371.—Guian.
guianensis, *Aubl. Pl. Gui.* i. 249. t. 99.—Guian.
longifolia, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 372.—Guian.
Sprengelii, Tul. l. c. 371 = brasiliensis.
ventricosa, *Tul. l. c.* xi. (1849) 153.—Peruv.

PAYROLA, Juss. Gen. 472 (1789) = **Paypayrola**, Aubl. (Violar.).
guianensis, Lam. Illustr. ii. t. 72 = Paypayrola guianensis.

PEARCEA, Regel, Gartenfl. xvi. (1867) 388 = **Isoloma**, Decne. (Gesner.).
hypocyrtiflora, Regel. l. c. = I. hypocyrtiflorum.

PECCANA, Rafin. Sylva Tellur. 114 (1838) = **Euphorbia**, Linn.
glauca, Rafin. l. c. 115 = E. nudiflora.

PECHEA, 'Arrab.' ex Steud. Nom. ed. II. ii. 280 (1841) sphalm. = PECKIA, Vell. = **Cybianthus**, Mart. (Myrsin.).

PECHEA, Pour. Chlor. Narb. n. 103; ex Kunth, Enum. Pl. i. 22, in syn. (1833) = **Heleochloa**, Host (Gramin.).
subcylindrica, Pourr. l. c. = H. schoenoides.

PECHEYA, Scop. Introd. 143 (1777) = **Coussarea**, Aubl. (Rubiac.).

PECKEYA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87 = praec.

PECKIA, Vell. Fl. Flum. 51; Ic. i. t. 134, 135 (1825) = **Cybianthus**, Mart. (Myrsin.).
megaphylla, Vell. l. c. 52; i. t. 135.—Bras.
verticillata, Vell. l. c. 51; i. t. 134 = C. cuneifolius.

PECKOLTIA, Fourn. in Mart. Fl. Bras. vi. VI. 310 (1885). *ASCLEPIADEAE*.
pedalis, *Fourn. l. c.*—Bras.

PECTANISIA, Rafin. Fl. Tellur. iii. 72 (1836) = **Reseda**, Linn.
Phyteuma, Rafin. l. c. 73 = R. Phyteuma.

PECTEILIS, Rafin. Fl. Tellur. ii. 37 (1836) = **Habenaria**, Willd. (Orchid.).
gigantea, Rafin. l. c. 38 = H. Susannae.
radiata, Rafin. l. c. = H. radiata.
Susanna, Rafin. l. c. = H. Susannae.

PECTEN, Lam. Fl. Fr. iii. 437 (1778) = **Scandix**, Tourn. (Umbell.).
Veneris, Lam. l. c. = S. Pecten-Veneris.

PECTIANTIA, Rafin. Fl. Tellur. ii. 72 (1836) = **Mitella**, Tourn. (Saxifr.).
mitelloides, Rafin. l. c. = M. pentandra.

PECTIDIUM, Less. in Linnaea, vi. (1831) 706 = **Pectis**, Linn. (Compos.).
punctatum, Less. l. c. = Pectis punctata.

PECTIDOPSIS, DC. Prod. v. 98 (1836) = **Pectis**, Linn. (Compos.).
angustifolia, DC. l. c. = Pectis angustifolia.

PECTINARIA, Bernh. Syst. Verz. Erf. 113 (1800) = **Scandix**, Tourn. (Umbell.).
vulgaris, Bernh. l. c. 168 = S. Pecten-Veneris.

PECTINARIA, Haw. Syn. Pl. Succ. Suppl. 14 (1819). *ASCLEPIADEAE*, [Benth. & Hook. f. ii. 782].
articulata, *Haw. l. c.*—Afr. austr.
mammillaris, Sweet, Hort. Brit. ed. II. 357 = Bouce-rosia mammillaris.

PECTINASTRUM, Cass. in Dict. Sc. Nat. xliv. 38 (1826) = **Centaurea**, Linn. (Compos.).
napifolium, Cass. l. c. = C. napifolia.

PECTINEA, Gaertn. Fruct. ii. 136. t. 111. f. 3 (1791) = **Erythrospermum**, Lam. (Bixin.).
zeylanica, Gaertn. l. c. = E. phytolaccoides?

PECTIS, Linn. Syst. ed. X. 1221 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 412.
CHEILODISCUS, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 36.
CHTHONIA, Cass. in Dict. Sc. Nat. ix. 173 (1817).
CRYPTOPETALON, Cass. in Bull. Soc. Philom. (1817) 12; et Dict. Sc. Nat. xii. 123 (1818).
CRYPTOPETALUM, DC. Prod. v. 101, in syn. (1836).
HELIOREOS, Rafin. Atl. Journ. 145 (1832).
LORENTEA, Lag. Gen. et Sp. Nov. 28 (1816).
PECTIDIUM, Less. in Linnaea, vi. (1831) 706.
PECTIDOPSIS, DC. Prod. v. 98 (1836).
SEALA, Adans. Fam. ii. 131 (1763).
STAMMARIUM, Willd. ex DC. Prod. v. 102 (1836).
acicularis, D. Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 320 = Hymenatherum tenuifolium.
affinis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Bras.
anceps, Less. in Linnaea, vi. (1831) 715 = ciliaris.
angustifolia, *Torr. in Ann. Lyc. N. York*, ii. (1828) 214.—Am. bor. occ.
apodocephala, Baker, in Mart. Fl. Bras. vi. III. 288 = brevipedunculata.
arenaria, *Benth. Bot. Voy. Sulph.* 110.—Mexic.
auricularis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
Berlandieri, *DC. Prod.* v. 100.—Mexic.
bibracteata, *Klatt, in Leopoldina*, xx. (1884) 92.—Mexic.
bipinnata, Steud. Nom. ed. II. ii. 280 = Schkuhria abrotanoides.
Bonplandiana, *H. B. & K. Nov. Gen. et Sp.* iv. 293.—Mexic.
brevipedunculata, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. III. 287.—Bras.
canescens, *H. B. & K. Nov. Gen. et Sp.* iv. 263. t. 393.—Mexic.
canescens, Less. in Linnaea, v. (1830) 135 = auricularis.
capillaris, *DC. Prod.* v. 99.—Mexic.
capillipes, *Hemsl. Biol. Centr. Am. Bot.* ii. 225.—Nicaragua.
carthusianorum, *Less. in Linnaea*, vi. (1831) 712.—Ins. S. Doming.
ciliaris, *Linn. Sp. Pl.* ed. X. 1221.—Ind. occ.
ciliaris, Sw. Obs. Bot. 307 = Swartziana.
ciliata, Steud. Nom. ed. II. i. in syn. = ciliaris.
congesta, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309; *Baker, in Mart. Fl. Bras.* vi. III. 287.—Bras.
costata, *Ser. & Merc. in DC. Prod.* v. 100.—Mexic.
Coulteri, *Harv. & Gray, in Mem. Am. Acad.* N. S. iv. (1849) 62.—Arizona.
cryptocephala, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Peruv.
cubensis, *Griseb. Cat. Pl. Cub.* 156.—Cuba.
decumbens, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309; *Baker, in Mart. Fl. Bras.* vi. III. 286.—Bras.
dichotoma, *Klatt, in Leopoldina*, xx. (1884) 92.—Mexic.
diffusa, *Hook. & Arn. Bot. Beech. Voy.* 296.—Mexic.
discoidea, Hornem. Suppl. Hort. Hafn. 100 = Leyssera capillifolia.
elongata, *H. B. & K. Nov. Gen. et Sp.* iv. 262. t. 392.—N. Granat.
fasciculata, Poir. Encyc. v. 120 = prostrata.
fasciculiflora, *DC. Prod.* v. 100.—Mexic.

PECTIS:—
fastigiata, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 62 = angustifolia.
febrifuga, *Van Hall, in Fl. Jard. Pays-Bas*, iv. (1861) 33.—Ins. Curaçoa.
filipes, Harv. & Gray, in Mem. Am. Acad. N. S. iv. (1849) 62 = taliscana.
floribunda, *A. Rich. Fl. Cub. Fanerog.* ii. 36.—Cuba.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 287.—Bras.
gibbosa, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 84.—Mexic.
gracilis, *Baker, in Mart. Fl. Bras.* vi. III. 290.—Bras.
Granierii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
Haenkeana, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
humifusa, *Sw. Prod. Veg. Ind. Occ.* 114.—Ind. occ.
imberbis, *A. Gray, Pl. Wright.* ii. 70.—Arizona; Mexic.
jaliscana, Hemsl. Biol. Centr. Am. Bot. ii. 225 = taliscana.
latisquama, Sch. Bip. ex A. Gray, in Proc. Am. Acad. v. (1862) 181 = canescens.
Liebmannii, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 226.—Mexic.
linearis, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 84.—Mexic.
linearis, Willd. ex Less. in Linnaea, vi. (1831) 712 = Bonplandiana.
linifolia, Linn. Amoen. Acad. v. 407 = punctata.
linifolia, *Linn. Syst.* ed. X. 1221.—Ind. occ.
longipes, *A. Gray, Pl. Wright.* ii. 69.—Arizona.
maritima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 289 = oligocephala.
meifolia, Wall. Cat. n. 3198 = Glossocardia linearifolia.
minuta, Schreb. ex Steud. Nom. ed. I. 103, 509 = Bellium minutum.
minuta, Willd. ex Less. in Linnaea, vi. (1831) 713 = pygmaea.
multifida, Orteg. Hort. Matr. Dec. 45 = Schkuhria abrotanoides.
multiflosculosa, Sch. Bip. in Herb. ex Benth. & Hook. f. Gen. ii. 412 = arenaria.
multiflosculosa, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Costa Rica.
multiseta, *Benth. Bot. Voy. Sulph.* 20.—Calif.
odorata, *Griseb. in Goett. Abh.* xxiv. (1879) 200.—Reg. Argent.
oligocephala, Sch. Bip. in Herb. ex Benth. & Hook. f. Gen. ii. 412 = rigida.
oligocephala, *Sch. Bip. in Seem. Bot. Voy Herald*, 309; *Baker, in Mart. Fl. Bras.* vi. III. 289.—Bras.
papposa, *Harv. & Gray, in Mem. Am. Acad.* N. S. iv. (1849) 62.—Am. bor. occ.
patula, Linn. f. ex DC. Prod. v. 99 = Swartziana.
pinnata, Lam. in Journ. Hist. Nat. Par. ii. (1792) 150. t. 31 = Schkuhria abrotanoides.
Plumieri, *Griseb. Fl. Brit. W. Ind.* 378.—Ind. occ.
polycephala, Sch. Bip. in Seem. Bot. Voy. Herald, 309 = elongata.
pratensis, *Wright, in Sauv. Fl. Cub.* 81.—Cuba.
prostrata, *Cav. Ic.* iv. 12. t. 324.—Am. bor. occ.; Mexic.
prostrata, Sieber, ex Less. in Linnaea, vi. (1831) 717 = humifusa.
punctata, *Jacq. Enum. Pl. Carib.* 28.—Calif.; Ind. occ.; Ins. Galap.
punctata, Sw. ex DC. Prod. v. 99 = Swartziana.
pygmaea, *H. B. & K. Nov. Gen. et Sp.* iv. 262.—N. Granat.
ramosissima, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Bras.
rigida, *Baker, in Mart. Fl. Bras.* vi. III. 288.—Bras.
rubiacea, *Baker, l. c.* 287.—Bras.
Rusbyi, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. II. 361.—Arizona.
satureioides, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
Seemannii, Sch. Bip. l. c. = Oxypappus scaber.
serpyllifolia, Less. in Linnaea, vi. (1831) 715 = humifusa.
sessiliflora, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Hab. ?
Sieberi, Less. in Linnaea, vi. (1831) 717 = humifusa.
stricta, Willd. ex Less. l. c. 710 = elongata.

PECTIS:—
subsquarrosa, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
Swartziana, *Less. in Linnaea*, vi. (1831) 711.—Ind. occ.; Panama.
taliscana, Benth. Pl. Hartw. 121 = diffusa.
taliscana, *Hook. & Arn. Bot. Beech. Voy.* 296.—Mexic.
tenella, *DC. Prod.* v. 99.—Mexic.
tenella, Rothrock, in Wheeler, Rep. vi. 171 = papposa.
tenuifolia, *Sch. Bip. in Seem. Bot. Voy. Herald*, 309.—Mexic.
uniaristata, *DC. Prod.* v. 99.—Mexic.
violacea, *Griseb. ex Lorentz, Veget. Entre-Rios*, 78, nomen; *et in Goett. Abh.* xxiv. (1879) 200.—Reg. Argent.

PECTOCARYA, DC. ex Meissn. Gen. 279 (1840).
BORAGINEAE, Benth. & Hook. f. ii. 847.
GRUVELIA, A. DC. in DC. Prod. x. 119 (1846).
KTENOSPERMUM, Lehm. Del. Sem. Hort. Hamb. (1837) 17.
chilensis, DC. Prod. x. 120 = linearis.
lateriflora, *DC. l. c.*—Peruv.
linearis, *DC. l. c.*—Calif.; Chili.
penicillata, *A. DC. l. c.*—Am. bor. occ.
pusilla, *A. Gray, in Proc. Am. Acad.* xii. (1877) 81.—Calif.; Chili.
setosa, *A. Gray, l. c.*—Calif.

PECTOPHYLLUM, Reichb. Consp. 145 (1828), sphalm. = seq.

PECTOPHYTUM, H. B. & K. Nov. Gen. et Sp. v. 28. t. 425 (1821) = **Azorella**, Lam. (Umbell.).
pedunculare, H. B. & K. l. c. 29 = A. pedunculata.

PEDALIUM, Adans. Fam. ii. 277 (1763) = **Atraphaxis**, Linn. (Polygon.).

PEDALIUM, Royen, ex Linn. Syst. ed. X. 1123 (1759).
PEDALINEAE, Benth. & Hook. f. ii. 1056.
CACATALI, Adans. Fam. ii. 43 (1763).
Caillaudii, Hochst. ex DC. Prod. ix. 256, in adnot. = Rogeria adenophylla.
filiforme, *Pav. ex DC. l. c.* 256, 257.—Mexic.
longiflorum, Decne. in Ann. Sc. Nat. Sér. V. iii. (1865) 331 = Rogeria longiflora.
microcarpum, Decne. l. c. 332 = Rogeria microcarpa.
Murex, *Linn. Syst.* ed. X. 1123.—Ind. or.; Afr. trop.
muricatum, Salisb. Prod. 104 = praec.
Rogeria, Decne. in Ann. Sc. Nat. Sér. V. iii. (1865) 331 = Rogeria adenophylla.

PEDASTIS, Rafin. Sylva Tellur. 87 (1838) = **Vitis**, Linn. (Ampel.).
indica, Rafin. l. c. = V. pedata.

PEDDIEA, Harv. in Hook. Journ. Bot. ii. (1840) 265. t. 10. *THYMELAEACEAE*, Benth. & Hook. f. iii. 199.
CYATHODISCUS, Hochst. in Flora, xxv. (1842) 240.
PSILOSOLENA, Presl, Bot. Bemerk. 102 (1844).
africana, *Harv. in Hook. Journ. Bot.* ii. (1840) 266.—Afr. austr.
africana, Plant, ex Meissn. in DC. Prod. xiv. 528 = Dregei.
Dregei, *Meissn. l. c.*—Afr. austr.
Harveyi, Meissn. l. c. = africana.
involucrata, *Baker, in Journ. Linn. Soc.* xx. (1883) 244.—Madag.
parviflora, *Hook. f. l. c.* vi. (1862) 20.—Afr. trop.

PEDERIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = **Paederia**, Linn. (Rubiac.).

PEDERLEA, Rafin. Sylva Tellur. 54 (1838) = **Acnistus**, Schott (Solan.).
aggregata, Rafin. l. c. = A. aggregatus.
arborescens, Rafin. l. c. = A. aggregatus.
cestroides, Rafin. l. c. = Lycium cestroides.

PEDEROTA, Scop. Ann. II. Hist. Nat. (1769) 41 = **Paederota**, Linn. (Scrophul.).

PEDICELLARIA, Schrank, in Roem. & Usteri, Mag. Bot. iii. St. VIII. (1790) 10 = **Gynandropsis**, DC. (1824) (Capparid.).
pentaphylla, Schrank, l. c. 11 = G. pentaphylla.

PEDICELLIA, Lour. Fl. Cochinch. 655 (1790) = **Ratonia** ?, DC. (Sapind.).
oppositifolia, Lour. l. c.—Cochinch.

PEDICULARIS, [Tourn.] Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 978.
PROSOPIA, Reichb. Consp. 121 (1828).
SCEPANIUM, Ehrh. Beitr. iv. 148 (1789).
SCEPTRUM-CAROLINUM, Linn. Fl. Lapp. 197 (1737).
abrotanifolia, *Bieb. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 22.—Sibir.
abrotanifolia, Henders. & Hume, Yarkand, 330; Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 309 = cheiranthifolia.
abrotanifolia, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 718 = violascens.
acaulis, Salzer, Reiseb. 323 = Oederi.
acaulis, *Scop. Fl. Carn.* ed. II. i. 439.—Europ. austr.
acaulis, *Vell. Fl. Flum.* 270; vi. t. 107.—Bras.
achilleaefolia, Host, ex Boiss. Fl. Orient. iv. 492 = comosa.
achilleaefolia, *Steph. ex Willd. Sp. Pl.* iii. 219.—Sibir.
achilleaefolia, Turcz. ex Bunge, in Ledeb. Fl. Ross. iii. 293 = fissa.
acmodonta, Boiss. Diagn. Ser. I. iv. 83 = comosa.
adscendens, *Gaud. Fl. Helvet.* iv. 145.—Europ.
adscendens, Hoppe & Sternb. in Denkschr. Bot. Ges. Regensb. i. II. (1818) 122 = tuberosa.
adunca, *Bieb. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 29.—Sibir.; Kamtschat.
aequinoctialis, H. B. & K. Nov. Gen. et Sp. ii. 332 = canadensis.
alaschanica, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 59.—Reg. Himal.; Mongolia.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 353.—As. centr.
Allionii, Reichb. f. Ic. Fl. Germ. xxx. 77. t. 1760 = rosea.
alopecuroides, Adams, ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 49 = Langsdorffi.
altaica, Steph. ex Bunge, in Ledeb. Fl. Ross. iii. 291 = uliginosa.
altaica, *Steph. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 48. t. 14 a.—Sibir. altaic.
altissima, Pall. Reise, iii. 297 = uncinata.
amoena, *Adams, ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 25.—Sibir.; Kamtschat.
angustifolia, *Benth. Pl. Hartw.* 22.—Mexic.
anthemifolia, Fisch. in Ledeb. Fl. Ross. iii. 303 = amoena.
araratica, Bunge, in Mém. Acad. Pétersb. Sér. VI. vii. (1858) 594 = crassirostris.
arctica, Adams, ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 51 = hirsuta.
arctica, Bieb. ex Stev. l. c. 25 = amoena.
arctica, *R. Br. in Parry, Voy. App.* 280; *et in Frankl. Journ. App.* 742 = Langsdorffi.
armata, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 56.—China.
armena, Boiss. & Huet, Diagn. Ser. II. iii. 175 = caucasica.
armena, Bunge, in Mém. Acad. Pétersb. Sér. VI. vii. (1858) 595 = crassirostris.
Artselaeri, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 84.—Mongolia.
ascendens, Schleich. Cat. Pl. Helv. ed. III. 21 = tuberosa.
asparagoides, Lapeyr. Hist. Abr. Pl. Pyren. 349 = comosa.
asplenifolia, Baumg. Enum. Stirp. Transs. ii. 202 = Oederi.
asplenifolia, Croom, in Am. Journ. Sc. Ser. I. xxvi. (1834) 315 = canadensis.
asplenifolia, *Floerk. ex Willd. Sp. Pl.* iii. 208.—Austria.
asplenifolia, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 173 = Oederi ?
asplenifolia, Wall. Cat. n. 416; Hook. f. Fl. Brit. Ind. iv. 314 (excl. syn.) = Wallichii.
atropurpurea, *Nordm. in Bull. Phys.-Math. Acad. Pétersb.* ii. (1837) 313.—As. Min.
atrorubens, Salzer, Reiseb. 323 = Oederi.
atrorubens, *Schleich. Cat. Pl. Helv.* (1815) 21.—Helvet.
attenuata. Benth. in DC. Prod. x. 574 = densiflora.
attollens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 384.—Calif.
auriculata, Sm. in Rees, Cycl. xxvi. [n. 6] = lanceolata.
avana, Wall. Cat. n. 1007 = Cardanthera avana.

PEDICULARIS:—
Barrelieri, Reichb. Fl. Germ. Excurs. 362 = adscendens.
bella, *Hook. f. Fl. Brit. Ind.* iv. 313.—Reg. Himal.
Besseriana, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 74, nomen (Quid ?).—Rossia.
bicornuta, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 109.—Reg. Himal.
bicuspidata, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 325 = comosa.
Bonjeani, Colla, Herb. Pedem. iv. 363 = cenisia.
borealis, Stev. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 337 = palustris.
Borgeaui, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 519.—Armenia.
brachyodonta, *Schloss. & Vukot. Fl. Croat.* 652.—Macedon.
brachystachys, *Bunge, in Ledeb. Fl. Alt.* ii. 429.—Mont. Altai.
bracteosa, *Benth. in Hook. Fl. Bor. Am.* ii. 110.—Am. bor. occ.
bracteosa, Pall. ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 252 = fissa.
breviflora, *Regel, in Act. Hort. Petrop.* vi. (1879) 352.—As. centr.
brevifolia, *D. Don, Prod. Fl. Nep.* 94.—Reg. Himal.
Brunoniana, Wall. Cat. n. 422 = gracilis.
bulbosa, Pall. Reise, ii. 87 = comosa.
Bungii, *Tchihat. Asie Min. Bot.* ii. 45.—As. Min.
cabulica, Benth. in DC. Prod. x. 575 = dolichorrhiza.
cadmea, *Boiss. Diagn.* Ser. I. iv. 82.—As. Min.
caespitosa, Sieber, ex Reichb. Fl. Germ. Excurs. 364 = rostrata.
caespitosa, Webb, Iter Hisp. 24 = verticillata.
callicoma, Herb. Gorenk. ex Stev. in Mem. Soc. Nat. Mosc. vi. (1823) 55 = Wilhelmsiana.
callosa, Adams, ex Stev. l. c. 25 = verticillata.
campestris, *Griseb. & Schenk, in Wiegm. Archiv*, xviii. (1852) I. 324.—Transsylv.
campylosiphon, C. Koch, in Linnaea, xxii. (1849) 682 = condensata.
canadensis, *Linn. Mant.* i. 86.—Am. bor.; Mexic.
capitata, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 100.—As. et Am. arct.
carnosa, *Wall. Pl. As. Rar.* ii. 44. t. 154.—Reg. Himal.
caucasica, *Bieb. Fl. Taur. Cauc.* ii. 72.—Reg. Caucas.; Persia.
cenisia, *Gaud. Fl. Helvet.* iv. 132.—Helvet.
centranthera, *A. Gray, in Torr. Bot. Mex. Bound.* 120.—Am. bor. occ.
Chamissonis, *Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 20. t. 4. f. 1.—Sibir.; Ins. Aleut.
cheilanthifolia, *Schrenk, Enum. Pl. Nov.* ii. 19; *et in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 79.—Reg. Himal.; Soongar.; China.
chinensis, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 57.—China.
chorgonica, *Regel, in Act. Hort. Petrop.* vi. (1879) 350.—As. centr.
chorgossica, Regel & Winkler, ex Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 514 = praec.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 310.—Reg. Himal.
comosa, Baumg. Enum. Stirp. Transs. ii. 202 = campestris.
comosa, Griseb. Spicil. Fl. Rumel. ii. 17 = brachyodonta.
comosa, *Linn. Sp. Pl.* 609.—Europ.; Oriens; As. bor.
comosa, Mattus. Enum. Stirp. Siles. 160 = sudetica.
comosa, Scop. Fl. Carn. ed. II. i. 440 = foliosa.
comosa, *Vell. Fl. Flum.* 270; vi. t. 106.—Bras.
compacta, *Steph. ex Willd. Sp. Pl.* iii. 219.—Soongar.; Sibir.
condensata, *Bieb. Fl. Taur. Cauc.* ii. 72.—As. Min.; Reg. Caucas.
conica, Pall. ex Bunge, in Ledeb. Fl. Ross. iii. 295 = physocalyx.
conica, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 46 = comosa.
contorta, *Benth. in Hook. Fl. Bor. Am.* ii. 108.—Am. bor. occ.
coronensis, Schur, in Oestr. Bot. Zeitschr. x. (1860) 183 = campestris.
corymbosa, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 42 = euphrasioides.
cranolopha, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 55.—China.
crassirostris, *Bunge, in Bull. Sc. Acad. Pétersb.* viii. (1840) 248.—Reg. Caucas.; Armenia.

PEDICULARIS:—
crenulata, *Benth. in DC. Prod.* x. 568.—Am. bor. occ.
cristata, Vitm. Summa Pl. iii. 481 = gyroflexa.
curvipes, *Hook. f. Fl. Brit. Ind.* iv. 316.—Reg. Himal.
curvituba, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 60.—China.
dasystachys, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 195 = sudetica.
densiflora, *Benth. in Hook. Fl. Bor. Am.* ii. 110.—Calif.
denudata, *Hook. f. Fl. Brit. Ind.* iv. 309.—Reg. Himal.
discolor, Adams, ex Ledeb. Fl. Ross. iii. 286 = sudetica.
distans, Willd. ex Walp. Rep. iii. 412 = verticillata.
dolichorrhiza, Reichb. Ic. Fl. Germ. xx. 71 = campestris.
dolichorrhiza, *Schrenk, Enum. Pl. Nov.* ii. 23; *et in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 80.—Reg. Himal.; As. centr.
Dollingeriana, Nordm. ex Benth. in DC. Prod. x. 573 = condensata.
elata, Pursh, Fl. Am. Sept. ii. 425 = bracteosa ?
elata, *Willd. Sp. Pl.* iii. 210.—Sibir.
elegans, Ledeb. ex Sweet, Hort. Brit. ed. II. 395 (Quid ?).—Sibir.
elegans, *Tenore, Fl. Nap.* v. 37.—Ital.
elephantoides, *Benth. Scroph. Ind.* 23.—Reg. Himal.
Elephas, Boiss. Diagn. Ser. I. iv. 81 = siphonantha.
Elephas, Boiss. Fl. Orient. iv. 489, partim = rhinanthoides.
elongata, *Kern. in Zeitschr. Ferd. Tirol.* III. xv. (1870) 260.—Europ.
Elwesii, *Hook. f. Fl. Brit. Ind.* iv. 312.—Reg. Himal.
erecta, Gilib. Exercit. i. 135 = palustris.
eriophora, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 333.—Reg. Himal.
eriostachys, Ledeb. ex Spreng. Syst. ii. 280 = sudetica.
erubescens, Kern. ex Caruel, in Parl. Fl. Ital. vi. 445 = Jacquini.
euphrasioides, Cham. in Linnaea, vi. (1831) 590 = lapponica.
euphrasioides, *Stev. ex Willd. Sp. Pl.* iii. 204.—Sibir.; Am. arct.
exaltata, *Bess. in Flora*, xv. (1832) II. *Beibl.* 19.—Ross. med.
exaltata, Fisch. ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 252 = condensata.
exaltata, Ledeb. ex Sweet, Hort. Brit. ed. II. 395 (Quid ?).—Sibir.
excelsa, *Hook. f. Fl. Brit. Ind.* iv. 311.—Reg. Himal.
eximia, Watt, in Journ. Linn. Soc. xviii. (1881) 381 = bicornuta.
fasciculata, Bell. ex Willd. Sp. Pl. iii. 218 = gyroflexa.
fasciculata, Poll. ex Benth. in DC. Prod. x. 570 = Friderici-Augusti.
Fetisowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 349.—Am. centr.
filicifolia, Adams, ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 252 = Oederi.
fissa, Hook. f. Fl. Brit. Ind. iv. 316 = dolichorrhiza.
fissa, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 339.—Sibir. baical.
fistulosa, Turcz. l. c. 350 = P. Sceptrum-Carolinum.
flagellaris, *Benth. in DC. Prod.* x. 581.—Reg. Himal.
flammea, *Linn. Sp. Pl.* 609.—Europ. arct.; Am. bor. occ.
flammea, Oed. in Fl. Dan. t. 30 = Oederi.
flava, Ledeb. Fl. Alt. ii. 433 = physocalyx.
flava, *Pall. Reise*, iii. 736.—Sibir.
flava, Sibth. & Sm. Fl. Graec. Prod. i. 429 = comosa.
flexuosa, *Hook. f. Fl. Brit. Ind.* iv. 308.—Reg. Himal.
folio-ceterac, Gilib. Fl. Lituan. i. 133 = P. Sceptrum-Carolinum.
foliosa, Georgi, Beschr. Russ. Reich. v. 1104 = exaltata.
foliosa, Gunn. Fl. Norveg. ii. 146 = Oederi.
foliosa, *Linn. Mant.* i. 86.—Europ.
foliosa, Schur, Enum. Pl. Transs. 509 = condensata.
Friderici-Augusti, *Tommasini, in Linnaea*, xiii. (1839) 74. t. 2.—Austria.
frondosa, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 46 = comosa.
Furbishiae, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 375.—Am. bor.
furfuracea, *Wall. Cat.* n. 412.—Reg. Himal.
Ganderi, Huter, in Oestr. Bot. Zeitschr. xxiii. (1873) 122 = Jacquini.
Gaudini, Arv. in Bull. Soc. Dauph. vii. (1880) 273, in obs. = adscendens.

PEDICULARIS:—
geminata, *Portenschl. ex J. N. Gebhard, Verz.* (1821) 190.—Alp. austr.
geminiflora, Reichb. Fl. Germ. Excurs. 361, sphalm. = praec.
gladiata, Michx. Fl. Bor. Am. ii. 18 = canadensis.
globifera, *Hook. f. Fl. Brit. Ind.* iv. 308.—Reg. Himal.
gloriosa, *Bisset & S. Moore, in Journ. Bot.* xv. (1877) 295.—Japon.
gracilis, *Wall. Cat.* n. 413.—Reg. Himal.
graeca, *Bunge, in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 376.—Graecia.
grandiflora, Cham. ex Ledeb. Fl. Ross. iii. 302 = capitata.
grandiflora, *Fisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 68.—Dahuria.
groenlandica, *Retz. Prod. Fl. Scand.* ed. II. 145.—Am. bor. occ. et arct.
gyroflexa, DC. ex Nym. Consp. 553 = pyrenaica.
gyroflexa, Lapeyr. Hist. Abr. Pl. Pyr. 349 = rostrata.
gyroflexa, Reichb. Fl. Germ. Excurs. 362 = cenisia.
gyroflexa, *Vill. Hist. Pl. Dauph.* ii. 426. t. 9.—Europ. med.
Hacquetii, Graf, in Flora, xvii. (1834) 42 = condensata ?
Hacquetii, C. Koch, in Linnaea, xvii. (1843) 42 = condensata.
hamata, Bieb. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 20. t. 4. f. 1 = Chamissonis.
Hausmanni, Huter, in Oestr. Bot. Zeitschr. xxiii. (1873) 126 = rosea.
heterodonta, *Panč. ex Janka, in Termész. Füzetek*, iv. (1881) 318.—Europ.
heterophylla, Patrin, ex Stev. in Mém. Soc. Nat. Mosc. 42 = euphrasioides.
himilayea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 107 = siphonantha.
hirsuta, All. Fl. Pedem. i. 63 = rosea.
hirsuta, Haenke, in Jirasek, Riesengeb. 86 = sudetica.
hirsuta, *Linn. Sp. Pl.* 609.—Reg. arct.
hirsuta, Linn. l. c. ed. II. 847, partim = Oederi.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 108 = megalantha.
Hookeriana, Wall. Cat. n. 421 = siphonantha.
Howellii, *A. Gray, in Proc. Am. Acad.* xx. (1885) 307.—Calif.
Huguenini, Reichb. f. Ic. Fl. Germ. xxx. 73. t. 1777 = gyroflexa.
Huteri ×, A. Kern. in Zeitschr. Ferd. Tirol. III. xv. (1870) 263.—Europ.
imbricata, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 46 = comosa.
incarnata, Baumg. Enum. Stirp. Transs. ii. 203 = Jacquini.
incarnata, *Jacq. Fl. Austr.* ii. 24. t. 140.—Europ.
incarnata, Krock. Fl. Siles. ii. 378 = sudetica.
incarnata, *Linn. Sp. Pl.* 609.—Sibir.
incarnata, Pall. Reise, ii. 404 = laeta.
incarnata, Retz. Prod. Fl. Scand. ed. I. 117; et Obs. iv. 27. t. 1 = groenlandica.
incurva, *Benth. in DC. Prod.* x. 566.—N. Granat.
insubrica, Rota, Prosp. Fl. Bergamo, 102 = palustris.
integrifolia, *Hook. f. Fl. Brit. Ind.* iv. 308.—Reg. Himal.
interrupta, *Steph. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 23.—Sibir.
Jacquini, *Koch, in Roehl. Deutschl. Fl.* ed. III. iv. 363.—Austria.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 121.—Japon.
jucunda, Schott & Kotschy, ex Boiss. Fl. Orient. iv. 492 = comosa.
Kanei, Dur. in Journ. Acad. Philad. Ser. II. iii. (1856) 195 = Langsdorffi.
kansuensis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 516.—China.
Kaufmanni, *Pinzger, Krit. Vergl. Gouv. Moskau Wildw. Pfl.* 17. t. 1.—Rossia.
Keiskei, *Franch. & Sav. Enum. Pl. Jap.* ii. 459.—Japon.
Kerneri, Dalla Torre, Anleit. Alpenpfl. (1882) 176 = rostrata.
Kerneri, *Huter, in Oestr. Bot. Zeitschr.* xxiii. (1873) 122, 123.—Europ.
Korolkowi, Regel, in Act. Hort. Petrop. vi. (1879) 349 = violascens.
labellata, Benth. in DC. Prod. x. 565 = siphonantha, rhinanthoides.
labellata, Boiss. Fl. Orient. iv. 489 = siphonantha.

PEDICULARIS:—
labellata, Decne. in Jacquem. Voy. Bot. 118. t. 123 = rhinanthoides.
labradorica, [Panzer, in] Houtt. Pflanzensyst. viii. 39 = euphrasioides.
lachnoglossa, *Hook. f. Fl. Brit. Ind.* iv. 311.—Reg. Himal.; China.
laciniosa, Pall. ex Benth. in DC. Prod. x. 570 = fissa.
laciniosa, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 46 = comosa.
laeta, Stev. ex Steud. Nom. ed. II. ii. 281 = sudetica.
laevicaulis, Láng, ex Walp. Rep. iii. 420 = palustris.
lanata, Pall. ex Pursh, Fl. Am. Sept. ii. 426 = verticillata.
lanata, Willd. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 49 = Langsdorffi.
lanceolata, *Michx. Fl. Am. Bor.* ii. 18.—Am. bor.
Langsdorffi, *Fisch. ex Stev. in Mém. Soc. Hist. Nat. Mosc.* vi. (1823) 49.—Reg. arct.
laniflora, Stev. ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 252 = Wilhelmsiana.
lanigera, Wall. Cat. n. 419 = brevifolia.
lapponica, *Linn. Sp. Pl.* 609.—Reg. arct.
lasiophrys, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 68.—Mongolia.
lasiostachys, *Bunge, in Ledeb. Fl. Alt.* ii. 434.—Sibir. altaic.
lasiostachys, Less. in Linnaea, ix. (1834) 157 = sudetica.
lepidota, *Weinm. in Bull. Soc. Nat. Mosc.* xxiii. (1850) i. 551.—Rossia.
leptorhiza, *Rupr. Sert. Tiansch.* 62.—Turkestan.
Letourneuxii, Personn. in Bull. Soc. Bot. Fr. viii. (1861) 461 = rostrata.
leucodon, *Griseb. Spicil. Fl. Rumel.* ii. 17.—Macedon.
leucodon, Reichb. f. Ic. Fl. Germ. xxx. 70. t. 133 = comosa.
limnogena, *Kern. in Oestr. Bot. Zeitschr.* xiii. (1863) 362.—Europ.
longiflora, *Rudolph, in Mém. Acad. Pétersb.* iv. (1811) 345. t. 3.—Sibir.; Reg. Himal.
Ludwigi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) i. 107.—As. bor.
lusitanica, *Hoffmgg. & Link, Fl. Port.* i. 306. t. 61.—Lusitan.
macrantha, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 108.—Reg. Himal.
macrantha, Spreng. Syst. iv. Cur. Post. 233 = grandiflora.
macrocarpa, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 57 = tristis.
macrodon, Steud. Nom. ed. II. ii. 281 = seq.
macrodontis, *Richards. in Frankl. 1st Journ. App.* ed. II. 24.—Sinus Hudson.
macrostachya, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = P. Sceptrum-Carolinum.
Malyi, *Janka, in Termész. Füzetek*, iv. (1880) 318.—Europ.
mandshurica, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 79.—Manchur.
Mariae, *Regel, in Act. Hort. Petrop.* vi. (1879) 351.—As. centr.
megalantha, *D. Don, Prod. Fl. Nep.* 94.—Reg. Himal.
melampyroides, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
Menziesii, *Benth. in DC. Prod.* x. 563.—Am. bor. occ.
mexicana, *Zucc. ex Bunge, in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 384, nomen.—Mexic.
microcalyx, *Hook. f. Fl. Brit. Ind.* iv. 315.—Reg. Himal.
mixta, Gren. ex Gren. & Godr. Fl. Fr. ii. 617 = pyrenaica.
mollis, *Wall. Cat.* n. 415.—Reg. Himal.
montenegrina, Janka, ex Nym. Consp. 554 = Frideric-Augusti.
moschata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 516.—Mongolia.
mucida, C. Koch, ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 251 = Frederici-Augusti.
Murithiana, *Arv. in Bull. Soc. Dauph.* vii. (1880) 275.—Europ.
muscicola, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 54.—China.
myriophylla, Pall. Reise, iii. 737 = abrotanifolia.
myrtophylla, Steud. Nom. ed. II. ii. 281, sphalm. = praec.
nasuta, *Bieb. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 43. t. 15. f. 1.—Ins. Curile.

PEDICULARIS:—
nasuta, Bong. Vég. Ins. Sitcha, 159 = pedicellata.
nasuta, Turcz. ex Bess. in Flora, xvii. (1834) i. Beibl. 21 = sudetica.
Nelsonii, *R. Br. in Frankl. 1st Journ. App.* 25.—Am. arct.
Nordmanniana, *Bunge, in Ledeb. Fl. Ross.* iii. 277.—Reg. Caucas.; As. Min.
nudicaulis, C. Koch, in Linnaea, xvii. (1843) 289 = caucasica.
numidica, *Pomel, Nouv. Mat. Fl. Atl.* 300.—Alger.
obliqua, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 248 = lapponica.
obsoleta, Crantz, Stirp. Austr. ed. II. fasc. iv. 316 = recutita.
occulta, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 180 = leucodon.
ochroleuca, Schloss. in Reichb. Ic. Fl. Germ. xx. 71. t. 1755 = comosa.
Oederi, *Vahl, in Hornem. Dansk. Oek. Plantel.* ed. II. 580.—Reg. bor. frigid.
Olgae, Regel, in Act. Hort. Petrop. vi. (1879) 349 = pycnantha.
oliosa, Vill. Fl. Delphin. 64 (err. typ. 65), sphalm. = foliosa.
olympica, *Boiss. Diagn.* Ser. I. iv. 83.—Bithyn.
Orizabae, Benth. Pl. Hartw. 47 = mexicana.
Orizabae, *Cham. & Schlecht. in Linnaea*, v. (1830) 103.—Mexic.
ornithorhyncha, *Benth. in Hook. Fl. Bor. Am.* ii. 118.—Am. bor. occ.
orthantha, *Griseb. Spicil. Fl. Rumel.* ii. 15.—Macedon.
pallescens, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 102 = Vulpii.
pallida, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 424 = lanceolata.
pallida, *Nutt. Gen. Am.* ii. 50.—Am. bor.
palustris, *Linn. Sp. Pl.* 607.—Reg. bor. temp. et arct.
paniculata, Pall. Reise, iii. 34 = euphrasioides.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1860) 250.—Am. bor. occ.
parviflora, Sm. in Rees, Cycl. xxvi. [n. 4] = palustris.
pectinata, *Wall. Cat.* n. 420.—Reg. Himal.
pedicellata, *Bunge, in Bull. Sc. Acad. Pétersb.* viii. (1841) 250.—Am. arct.
pennina, Gaud. Fl. Helvet. iv. 139 = atrorubens.
Perrottetii, *Benth. in DC. Prod.* x. 565.—Ind. or.
persica, Willd. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 26 = caucasica.
petiolaris, *Tenore, Fl. Nap.* v. *Syll. App.* iv. 11.—Italia.
philippica, *Gay, ex Benth. in DC. Prod.* x. 575.—Mexic.
physocalyx, *Bunge, in Bull. Sc. Acad. Pétersb.* viii. (1841) 252.—Sibir.
physodes, [Fisch.] Herb. Gorenk. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 36 = compacta.
pilostachya, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 64.—China.
platyrhyncha, *Schrenk, Enum. Pl. Schrenk.* ii. 21; *et in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 79.—As. centr.
polygaloides, *Hook. f. Fl. Brit. Ind.* iv. 317.—Reg. Himal.
pontica, *Boiss. Fl. Orient.* iv. 485.—As. Min.
porrecta, *Wall. Cat.* n. 423.—Reg. Himal.
Portenschlagii, Saut. ex Reichb. Fl. Germ. Excurs. 361 = geminata.
pratensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 56 = campestris.
proboscidea, Gaud. Fl. Helv. iv. 133 = cenisia.
proboscidea, *Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 33.—Sibir.
procera, Adams, ex Stev. l. c. 55 = fissa.
procera, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 251.—Am. bor. occ.
procumbens, Gilib. Exercit. i 135 = sylvatica.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 55.—Reg. Himal.; China.
Pseudocomosa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 68 = campestris.
psyllifolia, *Poir. Encyc. Suppl.* iv. 338 (gen. dub.).—Oriens.
pulchella, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 21.—Sibir. baical.
pumila, Mielichh. ex Saut. in Oestr. Bot. Wochenbl. vi. (1856) 107 = geminata.
punctata, Decne. in Jacquem. Voy. Bot. 117 (err. typ. 11) t. 122 = siphonantha.

PEDICULARIS:—
punicea, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 53 = elata.
purpurascens, Cham. ex Spreng. Syst. ii. 781 = Langsdorffi.
purpurea, Walp. Rep. iv. 617 (Index) = myriophylla.
pycnantha, *Boiss. Diagn.* Ser. I. xii. 45.—Turkestan; Persia; Reg. Himal.
pyramidata, Pall. ex Stev. in Mém Soc. Nat. Mosc. vi. (1823) 46 = comosa.
pyramidata, *Royle, ex Benth. Scroph. Ind.* 52.—Reg. Himal.
pyrenaica, *J. Gay, in Ann. Sc. Nat.* Sér. I. xxvi. (1832) 210.—Mont. Pyren.
racemosa, *Dougl. ex Hook. Fl. Bor. Am.* ii. 108.—Am. bor. occ.
ramosa, Wormsk. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 42 = euphrasioides.
ramosa, *Wormsk. in Fl. Dan.* t. 1758.—Groenl.
recutita, Baumg. Enum. Stirp. Transs. ii. 201, 202 = verticillata.
recutita, *Linn. Sp. Pl.* 608.—Europ.
recutita, Pursh, Fl. Am. Sept. ii. 425 = bracteosa?
refracta, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 517.—China.
resupinata, *Linn. Sp. Pl.* 608.—Sibir.
resupinata, Pursh, Fl. Am. Sept. ii. 425 = lanceolata?
rhaetica, Kern. Sched. Fl. Exsicc. ii. 116 = rostrata.
rhinanthoides, Hook. f. Fl. Brit. Ind. iv. 314, partim = elephantoides.
rhinanthoides, *Schrenk, Enum. Pl. Nov.* i. 22.—Sibir.; Reg. Himal.
Roborowskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 512.—China.
robusta, *Hook. f. Fl. Brit. Ind.* iv. 306.—Reg. Himal.
Romanzovii, Cham. ex Spreng. Syst. ii. 778 = Chamissonis.
rosea, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 16 = orthantha.
rosea, Tenore, ex Benth. in DC. Prod. x. 570 = Frideric-Augusti.
rosea, *Wulf. in Jacq. Misc.* ii. 57.—Alp. europ.
rostrata, Georgi, Beschr. Russ. Reich. v. 1102; C. A. Mey. Verz. Pfl. Cauc. 108 = Nordmanniana.
rostrata, Guss. ex Tenore, Fl. Nap. iv. Syll. 87 = elegans.
rostrata, Jacq. Fl. Austr. t. 205 = Jacquini.
rostrata, *Linn. Sp. Pl.* 607.—Europ.
rostrato-capitata, Crantz, Stirp. Austr. ed. II. fasc. iv. 320, 323 = rostrata.
rostrato-spicata, Crantz, l. c. fasc. iv. 317, 323 = incarnata.
Roylei, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 517.—Reg. Himal.; China.
rubens, Bunge, in Ledeb. Fl. Alt. ii. 435 = uliginosa.
rubens, Steph. ex Willd. Sp. Pl. iii. 219 = fissa.
rubra, Gueldenst. Reisen, i. 427 = crassirostris?
rudis, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 67.—Mongolia.
rupestris, Boiss. & Orph. Diagn. Ser. II. iii. 175 = graeca.
rupestris, Turcz. ex Bunge, Verz. Alt. Pfl. 60 = abrotanifolia.
sajanensis, *Steph. ex Bunge, in Bull. Sc. Acad. Pétersb.* viii. (1841) 252.—Sibir. baical.
salina, Turcz. ex Bunge, l. c. = fissa.
sarawschanica, *Regel, Pl. Nov. Fedsch.* 61.—Turkestan.
Sceptrum, Schrank, Baier. Fl. ii. 162 = seq.
Sceptrum-Carolinum, *Linn. Sp. Pl.* 608.—Europ.
Sceptrum-Carolinum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 122 = gloriosa.
Sceptrum-Marianum, *Vell. Fl. Flum.* 269; vi. t. 104 (Quid?).—Bras.
Schlosseri, Tommas. ex Nym. Consp. 554 = comosa.
Scolopax, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 513.—China.
scopulorum, *A. Gray, Syn. Fl. N. Am.* ii. 1. 308.—Am. bor. occ.
Seliwanowii, *Czern. Consp. Pl. Charcow*, 47; *ex Trautv. in Act. Hort. Petrop.* ix. (1884) 78, nomen.—Rossia.
Semenowi, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) i. 108 = pycnantha.
semibarbata, *A. Gray, in Proc. Am. Acad.* vii. (1868) 385.—Calif.
serotina, Adams, ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 251 = sudetica.

PEDICULARIS:—
serotina, Muhl. Cat. 58; ed. II. 60; et ex Benth. in DC. Prod. x. 582 = lanceolata.
sessilis, Vell. Fl. Flum. 270; vi. t. 105 = Hygrophila costata.
Sibthorpii, Boiss. Diagn. Ser. I. iv. 83 = comosa.
sima, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 514.—China.
siphonantha, *D. Don, Prod. Fl. Nep.* 65.—Reg. Himal.; Persia.
siphonantha, Hook. f. Fl. Brit. Ind. iv. 313, partim = rhinanthoides.
soongarica, *Schrenk, Enum. Pl. Nov.* ii. 25; *et in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 79.—Soongar.
speciosa, Ledeb. ex Sweet, Hort. Brit. ed. II. 305 (Quid ?).—Sibir.
Spica-foliosa, Jacq. Fl. Austr. ii. 24 = foliosa.
spicata, *Pall. Reise*, iii. 738.—Dahuria.
Stelleriana, Pall. ex Stev. in Mém. Soc. Hist. Nat. Mosc. vi. (1823) 54 = capitata.
Steveni, Adams, ex Stev. l. c. 44 = sudetica.
Steveni, Bunge, in Ledeb. Fl. Alt. ii. 427 = verticillata.
striata, *Pall. Reise*, iii. 737.—Dahur.
stricta, Wall. Cat. n. 414 = gracilis.
subnuda, Benth. in DC. Prod. x. 577 = pedicellata.
subrostrata, *C. A. Mey. Verz. Pfl. Cauc.* 108.—Reg. Caucas.
sudetica, *Willd. Sp. Pl.* iii. 209.—Europ.; Reg. arct.
sulphurea, Pall. Reise, iii. 223 = flava.
sumana, Pall. ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 252 = condensata.
sumana, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 34; Spreng. Pugill. ii. 70 = tuberosa.
surrecta, Benth. in Hook. Fl. Bor. Am. ii. 107 = groenlandica.
sylvatica, *Linn. Sp. Pl.* 607.—Europ.
tanacetifolia, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 102 = sudetica.
Tatarinowi, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 60.—China.
tenuifolia, Adams, ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 251 = palustris.
tenuirostris, Aitch. in Journ. Linn. Soc. xviii. (1880) 84; xix. (1882) 181 = pyramidata.
tenuirostris, *Benth. Scroph. Ind.* 52.—Reg. Himal.
ternata, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 64.—Mongolia.
Tommasinii, Kern. ex Marches. in Boll. Soc. Adriat. Sc. Nat. Trieste, ii. (1880) reimpr. 22, nomen = Friderici-Augusti ?
teucriifolia, *Bieb. ex Stev. in Mém. Soc. Nat. Mosc.* vi. (1823) 31.—Ins. Curila.
tianschanica, *Rupr. Sert. Tiansch.* 63.—Turkestan.
transsilvanica, Schur, in Oestr. Bot. Zeitschr. xi. (1861) 361 = exaltata.
trichoglossa, *Hook. f. Fl. Brit. Ind.* iv. 310.—Reg. Himal.
tripinnata, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 34.—Mexic.
tristis, *Linn. Sp. Pl.* 608.—Sibir.
tuberosa, All. Fl. Pedem. i. 64 = elegans.
tuberosa, *Linn. Sp. Pl.* 610.—Europ.
tuberosa, Vill. Fl. Delphin. 65 = adscendens.
tubiflora, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 58 = longiflora.
tubiformis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 106 = longiflora.
uliginosa, *Bunge, Del. Sem. Hort. Dorpat.* (1839) 8; *et in Bull. Sc. Acad. Pétersb.* viii. (1841) 251.—Sibir. altaic.
uncinata, Poir. Encyc. v. 133 (excl. syn.) = tuberosa.
uncinata, *Steph. in Willd. Sp. Pl.* iii. 213.—Sibir.
venosa, Pall. ex Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 20 = striata.
venusta, Schang. ex Bunge, in Bull. Sc. Acad. Pétersb. viii. (1841) 251 = fissa.
versicolor, Wahlenb. Veg. Helvet. 118 = Oederi.
verticillata, Benth. Scroph. Ind. 53 = Roylei.
verticillata, Bunge, in Ledeb. Fl. Alt. ii. 427 = amoena.
verticillata, *Linn. Sp. Pl.* 608.—Reg. bor. et arct.
verticillata, Pursh, Fl. Am. Sept. ii. 426 = capitata.
verticillata, Tchihat. Asie Min. Bot. ii. 46 = pontica.
villobracteata, C. Koch, in Linnaea, xxii. (1849) 682 = atropurpurea.
villosa, *Ledeb. ex Spreng. Syst.* ii. 780.—Sibir.
violascens, *Schrenk, Enum. Pl. Nov.* ii. 22; *et in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 79.—As. centr.

PEDICULARIS:—
virescens, Wahlenb. Fl. Suec. i. 389 = Oederi.
virginica, Poir. Encyc. v. 126 = lanceolata.
Vulpii ×, Solms, in Oestr. Bot. Zeitschr. xv. (1865) 174.—Europ.
Wallichiana, Steud. Nom. ed. II. ii. 282 = seq.
Wallichii, *Bunge, in Bull. Sc. Acad. Pétersb.* viii. (1843) 251, nomen.—Reg. Himal.
Wilhelmsiana, *Fisch. ex Bieb. Fl. Taur. Cauc.* iii. 412.—Reg. Caucas.
Wlassoviana, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 27. t. 9. f. 1 = palustris.
yezoënsis, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 69.—Japon.
zeylanica, *Benth. Scroph. Ind.* 54.—Ind. or.; Zeylan.

PEDILANTHUS, Neck. Elem. ii. 354 (1790). *EUPHORBIACEAE*, Benth. & Hook. iii. 257.
CREPIDARIA, Haw. Syn. Pl. Succ. 136 (1812).
DIADENARIA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 254; et in Abh. Akad. Berl. (1860) 107.
HEXADENIA, Klotzsch & Garcke, ll. cc. 253 (1859); 107 (1860).
VENTENATIA, Tratt. Gen. Pl. Disp. 86 (1802).
anacampseroides, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253 = padifolius.
angustifolius, *Poit. in Ann. Mus. Par.* xix. (1812) 393.—Hab. ?
aphyllus, *Boiss. in Klotzsch & Garcke, in Monatsb. Akad. Berl.* (1859) 253, nomen; *et in Abh. Akad. Berl.* (1860) 106.—Mexic.
articulatus, *Boiss. in DC. Prod.* xv. II. 6.—Mexic.
bracteatus, *Boiss. l. c.*—Mexic.
calcaratus, *Schlecht. in Linnaea*, xix. (1847) 255.—Mexic.
canaliculatus, *Sweet, Hort. Brit.* ed. I. 355.—Am. austr.
carinatus, *Spreng. Syst.* iii. 802.—Ind. occ.
cordatus, Spreng. l. c.—angustifolius.
cordellatus, *Steud. Nom.* ed. II. ii. 282.—Am. austr.
cymbiferus, *Schlecht. in Linnaea*, xix. (1847) 253.—Mexic.
Fendleri, *Boiss. in DC. Prod.* xv. II. 5.—Venezuela.
Finkii, *Boiss. l. c.* 1261.—Mexic.
Ghiesbreghtianus, *Baill. Adansonia*, i. (1860–61) 340.—Mexic.
Houlletii, *Baill. l. c.* 341.—Hab. ?
involucratus, *Boiss. in DC. Prod.* xv. II. 6.—Mexic.
linearifolius, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 161.—Ind. occ.
macrocarpus, *Benth. Bot. Voy. Sulph.* 49. t. 23.—Calif.
myrsifolius, Rafin. Fl. Tellur. iv. 117 = seq. ?
myrtifolius, Link, Enum. Hort. Berol. ii. 18 = tithymaloides.
Oerstedii, *Klotzsch & Garcke, in Monatsb. Akad. Berl.* (1859) 253, nomen.—Nicaragua.
padifolius, *Poit. in Ann. Mus. Par.* xix. (1812) 393.—Am. austr.
parasiticus, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253, nomen = ramosissimus.
Pavonis, *Boiss. in DC. Prod.* xv. II. 6.—Mexic.
ramosissimus, *Boiss. l. c.*—Mexic.
retusus, *Benth. in Hook. Kew Journ.* vi. (1854) 321.—Bras.
subcarinatus, *Steud. Nom.* ed. II. ii. 282.—Am. austr.
tithymaloides, *Poit. in Ann. Mus. Par.* xix. (1812) 390. t. 19.—Am. austr.

PEDILEA, Lindl. Orch. Scel. 27 (1826) = **Microstylis**, Nutt. (Orchid.).

PEDILONIA, Presl, Pedil. Nov. Pl. Gen. (1829); et in Flora, xii. (1829) 568 = **Wachendorfia**, Linn. (Haemod.).
violacea, Presl, l. c. = W. hirsuta.

PEDILONUM, Blume, Bijdr. 320. t. 36 (1825) = **Dendrobium**, Sw. (Orchid.).
biflorum, Blume, l. c. 322 = D. gemellum.
erosum, Blume, l. c. 323 = D. erosum.
Hasseltii, Blume, l. c. 321 = D. Hasseltii, *Lindl.*
Kuhlii, Blume, l. c. 321 = D. Kuhlii.
secundum, Blume, l. c. 322 = D. secundum.
undulatum, Blume, l. c. = D. hymenophyllum.

PEDINA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 144 = **Astragalus**, Tourn. (Legumin.).
aspera, Stev. l. c. = A. asper.

PEDOCHELUS, Wight, Ic. v. t. 1748. f. 2 (1852), sphalm. = **Podochilus**, Blume (Orchid.).

PEDROSIA, Lowe, in Hook. Kew Journ. viii. (1856) 292 = **Lotus**, Tourn. (Legumin.).
arenaria, Lowe, in Journ. Linn. Soc. v. (1861) 38 = L. arenarius.
argentea, Lowe, in Hook. Kew Journ. viii. (1856) 293 = L. argenteus.
Bertholetii, Webb, ex Masf. in Anal. Soc. Esp. Hist. Nat. x. (1881) 160 = L. Bertholetii.
florida, Lowe, in Hook. Kew Journ. viii. (1856) 294 = L. floridus.
glauca, Lowe, l. c. = L. glaucus.
leptophylla, Lowe, Man. Fl. Mad. i. 177.—Ins. Mader.
Loweana, Lowe, l. c. 181 = L. Loweanus.
macrantha, Lowe, in Hook. Kew Journ. viii. (1856) 293 = L. macranthus.
neglecta, Lowe, Man. Fl. Mader. i. 172 = L. neglectus.
Paivae, Lowe, Fl. Salv. Tent. 10.—Ins. Salvad.
porto-sanctana, Lowe, in Hook. Kew Journ. viii. (1856) 293 = L. Loweanus.
sessilifolia, Lowe, Man. Fl. Mad. i. 177, in obs. = L. sessilifolius.
tenella, Lowe, l. c. 177.—Ins. Mader.

PEGAMEA, Vitm. Summa Pl. i. 360 (1789) = **Pagamea**, Aubl. (Logan.).

PEGANON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Peganum**, Linn. (Rutac.).
harmalum, St. Lag. l. c. = Peganum Harmala.

PEGANUM, Linn. Syst. ed. I (1735); Gen. ed. I. 157 (1737). *RUTACEAE*, Benth. & Hook. f. i. 287.
HARMALA, Tourn. ex Adans. Fam. ii. 343 (1763).
MALACOCARPUS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 78 (1843).
crithmifolium, Georgi, ex DC. Prod. i. 712 = Adonidis sp. ?
crithmifolium, Retz. Obs. iii. 34 = P. Harmala.
dauricum, Pall. Reise, i. 418 = P. Harmala.
davurica, Linn. Sp. Pl. 445 = Ruta dahurica.
Harmala, *Linn. l. c.* 444.—Reg. Mediterr.; As. med.
mexicanum, *A. Gray, Pl. Wright.* i. 30; ii. 306.—Mexic.
Nigellastrum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 87.—China.
retusum, Forsk. Fl. Aegypt. Arab. 211 = Nitraria tridentata.

PEGESIA, 'Rafin.' ex Steud. Nom. ed. II. ii. 282 (1841) = PAGESIA, Rafin. = **Gerardia**, Linn. (Scrophul.).

PEGIA, Coleb. in Trans. Linn. Soc. xv. (1827) 364 = **Tapiria**, Juss. (Anacard.).
Colebrookiana, Wight, Ic. t. 236 = Nothopegia Colebrookiana.
nitida, Colebr. in Trans. Linn. Soc. xv. (1827) 364 = Tapiria hirsuta.

PEGOLETTIA, Cass. in Dict. Sc. Nat. xxxviii. 230 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 334.
acuminata, DC. Prod. v. 481 = Iphiona acuminata.
baccharidifolia, Less. Syn. Comp. 200 = Iphiona baccharidifolia.
javanica, *Mig. Fl. Ind. Bat.* ii. 63.—Java.
lanceolata, Harv. in Harv. & Sond. Fl. Cap. iii. 123 = Bojeria lanceolata (in Addend.).
mucronata, *Benth. in Hook. Niger Fl.* 432.—Afr. trop.
oxyodonta, *DC. Prod.* v. 481.—Afr. austr.
polygalaefolia, Less. Syn. Comp. 200 = Iphiona polygalaefolia.
senegalensis, *Cass. in Dict. Sc. Nat.* xxxviii. 230.—Afr. trop.
tenella, *DC. Prod.* v. 482.—Afr. austr.

PEGOLLETIA, Less. Syn. Comp. 200 (1832) = praec.

PEIRANISIA, Rafin. Sylva Tellur. 127 (1838) = **Cassia**, Tourn. (Legumin.).
aversifolia, Rafin. l. c. = C. biflora.

PEIRESCIA, Zucc. in Abh. Akad. Muench. ii. (1837) 695 = **Pereskia**, Plum. (Cact.).

PEIRESKIA, Steud. Nom. ed. II. ii. 282 (1841), sphalm. = praec.

PEIXOTOA, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 60. t. 172 (1833). *MALPIGHIACEAE*, Benth. & Hook. f. i. 257.
cordistipula, *A. Juss. in Arch. Mus. Par.* iii. (1843) 431.—Bras.
discolor, *Griseb. in Linnaea*, xiii. (1839) 216.—Bras.
glabra, Griseb. l. c. 212 = hispidula.
glabra, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 60. t. 172.—Bras.
hirta, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 430.—Bras.
hispidula, *A. Juss. l. c.* 429.—Bras.
Jussieuana, *Mart. ex A. Juss. l. c.* 432.—Bras.
lateritia, *A. Juss. in Walp. Rep.* v. 251.—Bras.
leptoclada, *A. Juss. in Arch. Mus. Par.* iii. (1843) 431.—Bras.
macrophylla, *Griseb. in Linnaea*, xiii. (1839) 214.—Bras.
microphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 584.—Bras.
paludosa, Turcz. l. c. xxxi. (1858) i. 293 = Jussieuana.
parviflora, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 62.—Bras.
reticulata, *Griseb. in Linnaea*, xiii. (1839) 213.—Bras.
tomentosa, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 61.—Bras.

PEKEA, Aubl. Pl. Gui. i. 594 (1775) = **Caryocar**, Linn. (Ternstr.).
butyrosa, Aubl. l. c. = C. butyrosum.
Couroupita, Juss. ex DC. Prod. iii. 294 = Couroupita guianensis.
ternata, Poir. Encyc. v. 148 = C. glabrum.
tuberculosa, Aubl. Pl. Gui. i. 597 = C. tomentosum.
villosa, Poir. Encyc. v. 148 = C. villosum.

PEKIA, Steud. Nom. ed. II. ii. 282 (1841), sphalm. = praec.

PELAGODENDRON, Seem. Fl. Vit. 124 (1866). *RUBIACEAE*, Benth. & Hook. f. ii. 92.
vitiense, *Seem. l. c.*—Ins. Fiji.

PELAPHIA, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I ii. (1839) 206 = **Coprosma**, Forst. (Rubiac.).
acerosa, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 109 = C. acerosa.
grandifolia, Banks & Soland. ex Hook. f. l. c. 104 = C. grandifolia.
laeta, Banks & Soland. ex Hook. f. l. c. = C. grandifolia.
laurifolia, Banks & Soland. ex Hook. f. l. c. = C. lucida.
parvifolia, Banks & Soland. ex Hook. f. l. c. 109 = C. propinqua.
retusa, Banks & Soland. ex Hook. f. l. c. 104 = C. Baueri.

PELARGONION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Pelargonium**, L'Hérit. (Geran.).
althaeoideum, St. Lag. l. c. = Pelargonium althaeoides.
ladanomum, St. Lag. l. c. = Pelargonium Ladanoma.
radulum, St. Lag. l. c. = Pelargonium Radula.

PELARGONIUM, L'Hérit. Geran. tt. 7–35, 43, 44 (1787). *GERANIACEAE*, Benth. & Hook. f. i. 273.
Campylia, Lindl. ex Sweet, Geran. sub t. 43 (1820).
Ciconium, Sweet, in Colvill, Cat.; et Geran. t. 13 (1820).
Corthumia, Reichb. Nom. 201 (1841).
Cortusina, Eckl. & Zeyh. Enum. 77 (1835).
Dibrachya, Eckl. & Zeyh. l. c. 74 (1835).
Dimacria, Lindl. in Sweet, Geran. t. 46 (1820).
Eumorpha, Eckl. & Zeyh. Enum. 77 (1835).
Grenvillea, Sweet, Geran. sub t. 262 (1825).
Hoarea, Sweet, l. c. t. 18 (1820).
Isopetalum, Sweet, l. c. t. 126 (1820).
Jenkinsonia, Sweet, l. c. t. 79 (1820).
Ligularia, Eckl. & Zeyh. Enum. 69 (1835).
Myrrhidium, Eckl. & Zeyh. l. c. 71 (1835).
Otidia, Lindl. ex Sweet, Geran. t. 98 (1820).
Peristera, Eckl. & Zeyh. Enum. 72 (1835).
Phymatanthus, Sweet, Geran. t. 43 (1820).
Polyactium, Eckl. & Zeyh. Enum. 65 (1835).

PELARGONIUM:—
Polyschisma, Turcz. in Bull. Soc. Nat. Mosc. (1859) i. 269.
Seymouria, Sweet, Geran. t. 206 (1830).
abbreviatum ×, Tratt. Pelarg. ii. t. 62.
abrotanifolium, *Jacq. Hort. Schoenb.* ii. t. 36.—Afr. austr.
abutiloides ×, Sweet, Geran. Suppl. t. 3.
abutiloides ×, Tratt. Pelarg. i. t. 30.
abyssinicum, *R. Br. in Salt, Abyss. App.* 65.—Abyss.
acanthifolium, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 63.—Afr. austr. ?
acaule, *DC. Prod.* i. 680.—Afr. austr.
accedens ×, Colv. ex Sweet, Hort. Brit. ed. I. 79.
acerifolium, *L'Hérit. Geran.* t. 21.—Afr. austr.
acerinum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 63 = palmatum.
aceroides ×, Sweet, Geran. t. 242.
acerosum, Hort. ex Hoffmgg. Verz. Pfl. 94, in syn. = palmatum.
acetabulosum ×, Sweet, Geran. Suppl. t. 44.
acetosellum, *Hort. ex Steud. Nom.* ed. II. ii. 291.—Hab. ?
acetosum, *Soland. in Ait. Hort. Kew.* ed I. ii. 430.—Afr. austr.
aciculare, *Hoffmgg. Verz. Pfl.* 86.—Afr. austr.
acidum ×, Sweet, Geran. t. 261.
aconitophyllum, *Steud. Nom.* ed. II. ii. 283.—Afr. austr.
Actinogramma ×, Tratt. Pelarg. i. t. 46.
acugnaticum, Thou. Fl. Tristan d'Acugn. 44. t. 13 = australe.
aculeatum, *Hort. ex Steud. Nom.* ed. II. ii. 283.—Afr. austr.
acuminatum, *DC. Prod.* i. 680.—Afr. austr.
acutangulum ×, Tratt. Pelarg. i. t. 35.
acutidentatum ×, Sweet, Geran. Suppl. t. 40.
acutifolium ×, Sweet, Hort. Brit. ed. I. 81.
acutilobum ×, Sweet, Geran. t. 184.
Adelinae ×, Tratt. Pelarg. vi. t. 255.
adscendens ×, Sweet, Hort. Brit. ed. I. 78, nomen.
adulterinum, *L'Hérit. Geran.* t. 34.—Afr. austr.
adulterinum, Wendl. ex Hoffmgg. Verz. Pfl. 92 = limbatum.
adventitium ×, Sweet, Geran. Suppl. t. 80.
aemulum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 63 = australe.
aemulum ×, Sweet, Geran. t. 160.
aequaliflorum, *Colv. ex Steud. Nom.* ed. II. ii. 283.—Hab. ?
affine, *Colv. ex Sweet, Hort. Brit.* ed. I. 79; *et Geran.* t. 277.—Hab. ?
affluens ×, Sweet, l. c. t. 194.
agathosmum ×, Tratt. Pelarg. iii. t. 119.
agonizans ×, Tratt. l. c. t. 98.
alatum, Steud. Nom. ed. II. ii. 291 = elatum.
albidum, *Colv. ex Steud. l. c.* 283.—Hab. ?
albiflorum, *Spin, Cat.* (1818) 30.—Hab. ?
albinotatum ×, Sweet, Geran. t. 359.
alchemillaefolium, Salisb. Prod. 312 = seq.
alchemilloides, *L'Hérit. ex* [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 419.—Afr. austr.
alienatum ×, Tratt. Pelarg. ii. t. 60.
Allenii ×, Sweet, Geran. t. 229.
alneum, Hoffmgg. Verz. Pfl. 86 = cordatum.
alnifolium, *Willd. Enum. Hort. Berol.* 710.—Hab. ?
Aloysianum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Afr. austr.
alpinum, *Eckl. & Zeyh. Enum.* 80.—Afr. austr.
Alstromeriae ×, Tratt. Pelarg. t. 127.
alternans, *Wendl. Hort. Herrenh.* i. 14.—Afr. austr.
althaeaefolium, Salisb. Prod. 313 = seq.
althaeoides, *L'Hérit. Geran.* t. 10.—Afr. austr.
altum ×, Sweet, Geran. Suppl. t. 26.
alutaceum, *Hoffmgg. Verz. Pfl.* 86.—Afr. austr.
amabile, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 64.—Afr. austr.
amabile ×, Tratt. Pelarg. i. t. 2.
amaryllidis ×, Tratt. l. c. t. 42.
amatymbicum, *Steud. Nom.* ed. II. ii. 283.—Afr. austr.
amauron ×, Tratt. Pelarg. t. 111.
ambiguum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Afr. austr. ?
ambrosiacum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 64.—Afr. austr. ?
amicum, *Hoffmgg. ex F. G. Dietr. l. c.*—Afr. austr. ?
amoenum, Hort. ex F. G. Dietr. l. c. = zonale.

PELARGONIUM:—
amoenum ×, Sweet, Geran. t. 121.
ampliatum ×, Tratt. Pelarg. t. 167.
amplissimum, *Wendl. Coll.* ii. 67. t. 65; *Willd. Enum. Hort. Berol. Suppl.* 48.—Afr. austr.
anacampton ×, Sweet, Geran. Suppl. t. 64.
Anaïs ×, Chauv. ex Poit. in Rev. Hortic. Sér. II. i. (1841) 121.
anceps, DC. Prod. i. 660 = australe.
anceps, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 420.—Afr. austr.
Andrewsii, *G. Don, Gen. Syst.* i. 729.—Afr. austr.
anemonifolium, Harv. in Harv. & Sond Fl. Cap. i. 288 = urbanum.
anemonifolium, Jacq. Ic. Pl. Rar. iii. 535 = myrrhifolium.
anethifolium, *Steud. Nom.* ed. II. ii. 283.—Afr. austr.
angulatum, Link, Handb. ii. 334 = seq.
angulosum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 426 = acerifolium.
angustifolium, *DC. Prod.* i. 680.—Afr. austr.
angustifolium, L'Hérit. ex Harv. & Sond. Fl. Cap. i. 293 = coronopifolium.
angustissimum, E. Mey, in Drège, Zwei Pfl. Docum. 208 = coronopifolium.
angustum, *Hort. ex Steud. Nom.* ed. II. ii. 283.—Afr. austr.
anisodonton ×, Sweet, Geran. t. 399.
Annesleyanum ×, Sweet, Geran. Suppl. t. 56.
Annleyanum, Pritz. Ic. Ind. i. 812, sphalm. = praec.
anodaefolium, *Hoffmgg. Verz. Pfl.* 86.—Afr. austr. ?
anomalanthum ×, Tratt. Pelarg. t. 182.
anomalum ×, Sweet, Hort. Brit. ed. I. 78; Geran. t. 315.
anomalum ×, Tratt. Pelarg. t. 38.
anosma, Hoffmgg. Verz. Pfl. 86 = citriodorum.
anosmon ×, Tratt. Pelarg. t. 175.
anthriscifolium ×, Sweet, Geran. t. 233.
antidysentericum, *Kostel. Allg. Med.-Pharm. Flora*, v. 1896.—Afr. austr.
Antonianum ×, Tratt. Pelarg. t. 109.
aphanoides, *DC. Prod.* i. 680.—Afr. austr.
aphanoides, Harv. in Harv. & Sond. Fl. Cap. i. 296 = alchemilloides.
apiifolium, *Jacq. f. Eclog.* i. t. 27.—Afr. austr.
apioides, *Steud. Nom.* ed. II. ii. 291.—Afr. austr.
appendiculatum, *Willd. Sp. Pl.* iii. 651.—Afr. austr.
aquilinum ×, Tratt. Pelarg. t. 66.
ardens, *Lodd. Bot. Cab.* t. 139.—Afr. austr.
ardescens ×, Sweet, Geran. t. 231.
Arembergianum, Klier, in Hamb. Gartenz. i. (1845–46) 78. t. 2 (Quid ?).—Hab. ?
arenarium, *DC. Prod.* i. 680.—Afr. austr.
arenarium, Harv. in Harv. & Sond. Fl. Cap. i. 272 = lobatum.
arenicola, *Steud. Nom.* ed. II. ii. 283.—Afr. austr.
argutum ×, Sweet, Hort. Brit. ed. I. 471; Geran. t. 344.
aristatum, *G. Don, Gen. Syst.* i. 728.—Afr. austr.
armatum, *Colv. in Sweet, Geran.* t. 214.—Afr. austr.
aromaticum, *Hoffmgg. Verz. Pfl.* 87.—Afr. austr.
artemisiaefolium, *DC. Prod.* i. 661.—Afr. austr.
artemisiaefolium, E. Mey. in Drège, Zwei Pfl. Docum. 208 = abrotanifolium.
articulatum, Harv. in Harv. & Sond. Fl. Cap. i. 296 = alchemilloides.
articulatum, *Willd. Sp. Pl.* iii. 655.—Afr. austr.
asarifolium, *G. Don, Gen. Syst.* i. 731.—Afr. austr.
asperifolium, *Sweet, Geran.* t. 169.—Afr. austr.
asperum, Ehrh. ex Spreng. Syst. iii. 62 = graveolens.
asperum, Willd. Sp. Pl. iii. 678 = quercifolium.
astragalifolium, *Jacq. Ic. Pl. Rar.* iii. 8.—Afr. austr.
astrantiaefolium, *Hoffmgg. Verz. Pfl.* 87.—Afr. austr.
athamanthoides, *L'Hérit. Geran.* 47; *ex DC. Prod.* i. 660.—Afr. austr.
Atkinsianum, *Sweet, Geran. Suppl.* t. 99.—Hab. ?
atrofulgens, *Kern. Hort.* t. 804.—Hab. ?
atrofuscum, *Sweet, Geran.* t. 82.—Afr. austr.
atromaculatum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Afr. austr. ?
atropurpureum ×, Sweet, Geran. t. 152.
atrorubens ×, Colv. ex Sweet, Hort. Brit. ed. I. 97.
atrosanguineum, *DC. Prod.* i. 653.—Afr. austr.
atrosanguineum, F. G. Dietr. in N. Bot. Garten-Journ. i. 163; ex Vollst. Lexik. Gaertn. Nachtr. vi. 61 = sanguineum.
atrovirens ×, Sweet, Geran. Suppl. t. 74.
atrum, *L'Hérit. Geran.* t. 44.—Afr. austr.
attenuatum, *Harv. in Harv. & Sond. Fl. Cap.* i. 266.—Afr. austr.

PELARGONIUM:—

augustissimum ×, Tratt. Pelarg. t. 193.
augustum, *Link, Enum. Hort. Berol.* ii. 192.—Afr. austr.
aulicum ×, Sweet, Hort. Brit. ed. I. 77.
aurantiacum ×, Sweet, Geran. t. 198.
aurantiadorum, *Hoffmgg. Verz. Pfl.* 87.—Afr. austr.?
auriculaeflorum ×, Tratt. Pelarg. t. 232.
auriculatum ×, Forbes, Hort. Woburn. 149.
auriculatum, Willd. Sp. Pl. iii. 643 = longifolium.
auritum, *Willd. Sp. Pl.* iii. 644.—Afr. austr.
auroreum ×, Tratt. Pelarg. t. 8.
australe, *Jacq. Eclog. Pl.* t. 100; *Sweet, Geran.* t. 68.—Reg. austr.
autumnale, Hort. ex DC. Prod. i. 676 = floridum.
Avronianum ×, Sweet, Geran. t. 394.
Azaleae ×, Tratt. Pelarg. t. 87.
Bacchantum, De Kleist, ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 65 = glutinosum?
Baileyanum, *Sweet, Geran.* t. 87.—Afr. austr.
Bakerianum ×, Sweet, l. c. t. 240.
Balbisianum, *Spin, Cat. Hort. Sebast.* (1818) 30.—Afr. austr.
Baldinianum ×, Tratt. Pelarg. t. 219.
balsameum, Jacq. Ic. Pl. Rar. iii. t. 543; Coll. Suppl. 136 = scabrum.
barbatum, *Jacq. l. c.* t. 513; *Coll.* 197.—Afr. austr.
Barclayanum ×, Sweet, Hort. Brit. ed. I. 84; Geran. t. 384.
Barmanum, *Voigt, ex Hoffmgg. Verz. Pfl.* 87.—Afr. austr.?
Barnardianum, *Sweet, Geran.* t. 127.—Afr. austr.
Barringtonii, *Willd. Enum. Hort. Berol.* 706.—Afr. austr.
basilicum ×, Sweet, Hort. Brit. ed. I. 471; Geran. t. 360.
Bathoniae, Hoare, in Sweet, Geran. sub t. 74 = *Thynneae.*
Baumannianum ×, Tratt. Pelarg. t. 154.
Beadoniae ×, Sweet, Geran. t. 191.
Beaufortianum, *Pers. Syn.* ii. 231.—Afr. austr.
Belladonna ×, Sweet, Geran. t. 270.
Bellardi, *Spin, Cat. Hort. St. Sebast.* (1818) 30.—Afr. austr.
Bellermanni, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Hab.?
bellulum, *Sweet, Geran.* t. 60.—Afr. austr.
bellum, Hort. ex Tratt. Pelarg. sub t. 22 = *chamaerhodon.*
Bentinckianum, DC. Prod. i. 664 = fulgidum?
Bentinkianum, Hort. ex Willd. Enum. Hort. Berol. 704 = praec.?
Berterianum, *Spin, Cat. Hort. Sebast. Suppl.* (1823) 12; *et ex Colla, Hort. Ripul. App.* iii. 8. t. 4.—Afr. austr.
Besseri, Link, Enum. Hort. Berol. ii. 195 (Quid?).—Afr. austr.?
betonicum, Jacq. Coll. Suppl. 127 = myrrhifolium.
betulaefolium, Schrank, ex Hoffmgg. Verz. Pfl. 87 = cordatum.
betulinum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 429.—Afr. austr.
Beudoniae, Steud. Nom. ed. II. ii. 292, sphalm. = *Beadoniae.*
bicolor, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 425.—Afr. austr.
biflorum ×, Sweet, Geran. t. 287.
bifolium, *Willd. Sp. Pl.* iii. 645.—Afr. austr.
bijugum, *Steud. Nom.* ed. II. ii. 284.—Afr. austr.
bipartitum, *DC. Prod.* i. 653.—Afr. austr.
bipinnatifidum, *Steud. Nom.* ed. II. ii. 284.—Afr. austr.
bipinnatifidum ×, Sweet, Geran. Suppl. t. 62.
Birolii, *Spin, ex Colla, Hort. Ripul. App.* ii. 353.—Afr. austr.?
Bishopae ×, Sweet, Geran. t. 272.
Bivonae, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Afr. austr.?
bivulnerum ×, Tratt. Pelarg. t. 178.
Blandfordianum, *Sweet, Geran.* t. 101.—Afr. austr.
blandum, Hornem. Hort. Hafn. Suppl. 76 = Hornemanni.
blandum, *Sweet, Geran.* t. 4.—Afr. austr.
blattarium, Jacq. Hort. Schoenb. ii. 3. t. 131 = ovale.
Bluecherianum, *Hort. ex Hoffmgg. Verz. Pfl.* 87.—Hab.?
Bluntianum ×, Sweet, Geran. Suppl. t. 79.
Boleyniae ×, Forbes, Hort. Woburn. 150.

PELARGONIUM:—

Boscaweniae ×, Colv. ex Sweet, Hort. Brit. ed. I. 79.
Bowkeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 592.—Afr. austr.
Boyleae, Sweet, Geran. t. 50 = formosissimum.
bracteosum, *DC. Prod.* i. 664.—Afr. austr.
brassicatum ×, Sweet, Hort. Brit. ed. I. 78.
Brayanum ×, Tratt. Pelarg. t. 141.
Breesianum, *Sweet, Geran.* t. 64.—Afr. austr.
brevipes ×, Sweet, Hort. Brit. ed. I. 84, nomen.
brevirostre, E. Mey. in Drège, Zwei Pfl. Docum. 208 = australe.
Brightianum ×, Sweet, Geran. t. 227.
Broadleyae ×, Colv. ex Sweet, Hort. Brit. ed. I. 81; G. Don, Gen. Syst. i. 740.—Hab.?
Broughtoniae ×, Sweet, Geran. t. 181.
Brownii, *Sweet, l. c.* t. 146.—Afr. austr.
bryoniaefolium ×, Sweet, l. c. t. 320.
bubonifolium, *Pers. Syn.* ii. 227.—Afr. austr.
Buffonianum, *Spin, ex Colla, Hort. Ripul. App.* ii. 353.—Hab.?
bullatum, Jacq. Ic. Pl. Rar. iii. 530; Coll. v. 124 = myrrhifolium.
Burdini, *Colla, Hort. Ripul. App.* iii. 6. t. 3.—Afr. austr.
Burmanni, Hoffmgg. Verz. Pfl. 87 = Barmanum.
Burmannianum, Steud. Nom. ed. II. ii. 284 = ferulaceum.
Burnelianum, Steud. l. c., sphalm. 292 = seq.
Burnettianum ×, Sweet, Geran. t. 369.
Byronianum ×, Forbes, Hort. Woburn. 150.
caespitosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 419.—Afr. austr.
caffrum, *Steud. Nom.* ed. II. ii. 284.—Afr. austr.
calamistratum ×, Sweet, Geran. Suppl. t. 36.
calliston ×, Sweet, l. c. t. 9.
callodon ×, Tratt. Pelarg. t. 76.
callosum, E. Mey. in Drège, Zwei Pfl. Docum. 208 = anethifolium.
calocephalon ×, Sweet, Geran. t. 201.
calographicum ×, Tratt. Pelarg. t. 25.
calycinum ×, Sweet, Geran. t. 81.
Camaldulense ×, Dehnh. Cat. Pl. Hort. Camald. ed. II. 21.
campestre, *Steud. Nom.* ed. II. ii. 284.—Afr. austr.
camphoratum, *Hoffmgg. Verz. Pfl.* 87.—Afr. austr.
campyliaeflorum ×, Sweet, Geran. t. 251.
campylosepalum ×, Sweet, l. c. 379.
canaliculatum, Zuccagni, in Roem. Collect. 146 = myrrhifolium.
canariense, *Willd. Hort. Berol.* t. 17; *Enum. Hort. Berol.* 702 (Quid?).—Ins. Canar.
candicans, *Spreng. Syst.* iii. 57.—Afr. austr.
candidum, *Hoare, ex Sweet, Geran.* t. 128.—Afr. austr.?
candidum, Link, Enum. Hort. Berol. ii. 194 = Blandfordianum.
canescens, Sweet, Geran. sub t. 93 = incisum.
cantabrigiense, Forbes, Hort. Woburn. 150.
canum, Pers. Syn. ii. 229 = ovale.
capense, Link, Enum. Hort. Berol. ii. 195 (Quid?).—Afr. austr.
capillare, Harv. in Harv. & Sond. Fl. Cap. i. 293 = violarium.
capillare, *Willd. Sp. Pl.* iii. 660.—Afr. austr.
capitatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 425.—Afr. austr.
capnoides, L'Hérit. ex DC. Prod. i. 660 = minimum.
carbasinum ×, Sweet, Geran. Suppl. t. 38.
carbonatum ×, Tratt. Pelarg. t. 100.
cardiifolium, *Sweet, Geran.* t. 15.—Afr. austr.
cardiophyllum, Harv. in Harv. & Sond. Fl. Cap. i. 284 = setulosum.
carinatum, *Sweet, Geran.* t. 21.—Afr. austr.
carinatum, *Wendl. Coll.* ii. 79. t. 71.—Afr. austr.
carnescens ×, Sweet, Hort. Brit. ed. I. 79.
carneum, *Jacq. Ic. Pl. Rar.* iii. 8. t. 512; *Coll.* iv. 189.—Afr. austr.
carnosum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 421.—Afr. austr.
carolinaeum ×, Tratt. Pelarg. t. 218.
carotaefolium, Hoffmgg. Verz. Pl. 87 = flavum.
cartilagineum, Sweet, Hort. Brit. ed. I. 471; Geran. t. 382.
cartilagineum ×, Tratt. Pelarg. t. 7.
caryophyllaceum ×, Sweet, Hort. Brit. ed. I. 77.
castaliaefolium ×, Tratt. Pelarg. t. 68.
castellianum, *Spin, Cat. Hort. Sebast.* (1818) 30.—Afr. austr.
castum ×, Tratt. Pelarg. t. 15.

PELARGONIUM:—

cataractae, Steud. Nom. ed. II. ii. 284 = patulum.
caucalifolium, *Jacq. Ic. Pl. Rar.* iii. 529; *Coll.* v. 145.—Afr. austr.
cavrianum ×, Tratt. Pelarg. t. 243.
centauroides, *L'Hérit. Geran.* t. 12.—Afr. austr.
cephalanthum ×, Tratt, Pelarg. t. 159.
cephalicum, Hoffmgg. Verz. Pfl. 87 = scabrum.
Cephalotes, *Hoffmgg. l. c.* 88.—Afr. austr.
ceratophyllum, *L'Hérit. Geran. Ined.* 12; *ex DC. Prod.* i. 652.—Afr. austr.
cerinum, DC. Prod. i. 659 = inquinans.
cernuum ×, Tratt. Pelarg. t. 209.
chaerophyllum ×, Sweet, Geran. t. 257.
chamaedrifolium, *Harv. in Harv. & Sond. Fl. Cap.* i. 290.—Afr. austr.
chamaedrifolium, *Jacq. Ic. Pl. Rar.* iii. 523; *Coll. Suppl.* 130.—Afr. austr.
chamaerhodon ×, Tratt. Pelarg. t. 22.
Chandleri ×, Colv. ex Sweet, Hort. Brit. ed. I. 79; Forbes, Hort. Woburn. 151.
Charlwoodii ×, Sweet, Geran. t. 380.
chelidoniifolium ×, Sweet, Hort. Brit. ed. I. 471; Geran. t. 341.
Chelidonium, *DC. Prod.* i. 650.—Afr. austr.
chenopodifolium ×, Sweet, Geran. t. 328.
chrysanthemifolium ×, Sweet, l. c. t. 124.—Afr. austr.
Chrysosplenium ×, Tratt. Pelarg. t. 262.
ciliatum, Jacq. Coll. iv. 195 = longifolium.
ciliatum, Willd. Sp. Pl. iii. 643 = parnassioides.
cinctum, *Baker, in Saund. Refug. Bot.* t. 151.—Afr. austr.
circumflexum ×, Sweet, Hort. Brit. ed. I. 78, nomen.
circumscriptum ×, Tratt. Pelarg. t. 102.
citriodorum, *Hort. ex Breit. Hort. Breit.* 331; *Schrank, in Syll. Ratisb.* ii. (1828) 67.—Afr. austr.
citrosum, Voigt, ex Breit. l. c. = citriodorum.
clandestinum, L'Hérit. ex DC. Prod. i. 160 = australe.
clarum, Hoffmgg. Verz. Pfl. 88 = zonale.
clarum ×, Sweet, Geran. t. 366.
clathratum ×, Sweet, l. c. Suppl. t. 85.
clausum ×, Colv. ex Sweet, Hort. Brit. ed. I. 77.
clavatum, *L'Hérit. ex DC. Prod.* i. 660.—Afr. austr.
Clementinae ×, Tratt. Pelarg. t. 196.
Clintoniae ×, Sweet, Geran. t. 392.
clypeatum, Steud. Nom. ed. II. ii. 284 = peltatum.
coarctatum ×, Sweet, Geran. t. 70.—Afr. austr.
Coburgiae ×, Tratt. Pelarg. t. 148.
Coburgianum ×, Tratt. l. c. t. 147.
coccineum, Ehrh. Beitr. vii. 162 = hybridum.
cochleatum, Willd. Enum. Hort. Berol. Suppl. 48 = acerifolium.
Codringtonii ×, Forbes, Hort. Woburn. 150.
coeleste ×, Tratt. Pelarg. t. 17.
coilophyllum, Sweet, Hort. Brit. ed. II. 98.
Collae, *Spin, Cat. Hort. Sebast.* (1818) 31.—Afr. austr.
Colleyanum ×, Sweet, Geran. Suppl. t. 83.
columbinum, Jacq. Hort. Schoenb. ii. 4. t. 133 = australe.
columbinum, E. Mey. ex Harv. in Harv. & Sond. Fl. Cap. i. 291 = fumarioides.
coluteaefolium, *Steud. Nom.* ed. II. ii. 284.—Afr. austr.
Colvillii ×, Sweet, Geran. t. 86.—Afr. austr.
commixtum ×, Sweet, l. c. Suppl. t. 71.
compactum ×, Sweet, Hort. Brit. ed. I. 84; et l. c. t. 70.
compactum ×, Tratt. Pelarg. t. 116.
compar ×, Colv. ex Sweet, Hort. Brit. ed. I. 81.
compositum, L'Hérit. ex DC. Prod. i. 667 = laevigatum.
compson ×, Tratt. Pelarg. t. 234.
Comptoniae ×, Sweet, Geran. t. 122.—Afr. austr.
comptum ×, Sweet, l. c. t. 255.
concavifolium, *Pers. Syn.* ii. 226.—Afr. austr.
concavifolium, Wendl. Coll. t. 64; Hort. ex DC. Prod. i. 672 = acerifolium.
concavum, Hoffmgg. Verz. Pfl. 88 = Dennisianum.
concavum ×, Sweet, Geran. t. 237.
conchaefolium ×, Tratt. Pelarg. t. 187.
conchiifolium ×, Sweet, Hort. Brit. ed. I. 82.
conchyllatum ×, Sweet, Geran. Suppl. t. 95.
concinnum, Dehnh. Cat. Pl. Hort. Camald. ed. II. 21 (Quid?).—Afr. austr.?
concinnum ×, Sweet, Geran. t. 108.—Afr. austr.

PELARGONIUM:—
conclausum ×, Sweet, Geran. t. 305.
concolor ×, Sweet, l. c. t. 140.—Afr. austr.
concretum ×, Sweet, l. c. Suppl. t. 24.
condensatum, Pers. Syn. ii. 227.—Afr. austr.
conduplicatum, *Willd. Enum. Hort. Berol.* 705.—Afr. austr.
conduplicatum-crispum, Hort. ex Hoffmgg. Verz. Pfl. 97 = vestitum.
confertifolium ×, Sweet, Hort. Brit. ed. I. 77.
conforme ×, Tratt. Pelarg. t. 54.
confusum, *DC. Prod.* i. 661.—Afr. austr.
congestum, *Sweet, Geran.* t. 325; *G. Don, Gen. Syst.* i. 727.—Hab.?
coniophyllum, Steud. Nom. ed. II. ii. 285 = flavum.
Conquestor ×, Stroud, ex Forbes, Hort. Woburn. 150.
consanguineum, *Willd. Enum. Hort. Berol.* 710.—Afr. austr.
consimile ×, Sweet, Hort. Brit. ed. I. 78.
conspicuum, *G. Don, Gen. Syst.* i. 731.—Afr. austr.
contiguum ×, Sweet, Geran. Suppl. t. 73.
convolutum, *Hoffmgg. Verz. Pfl.* 88.—Hab.?
corallinum, *Steud. Nom.* ed. II. ii. 285.—Afr. austr.
cordatum, *L'Hérit. Geran.* t. 22.—Afr. austr.
cordatum-varium, Van Eeden, ex Hoffmgg. Verz. Pfl. 97 = P. Volucris.
cordifolium, Curt. Bot. Mag. t. 165 = cordatum.
cordifolium, Hort. ex Willd. Enum. Hort. Berol. 705 = conduplicatum.
cordiforme ×, Sweet, Geran. Suppl. t. 49.
coriaceum ×, Tratt. Pelarg. t. 26.
coriandrifolium, Jacq. Ic. Pl. Rar. iii. 528; Coll. v. 142 = myrrhifolium.
Cornvallii ×, Forbes, Hort. Woburn. 150.
coronillaefolium, Fisch. Hort. Gorenk. ed. II. (1812) 54, nomen = rapaceum.
coronillaefolium, *Pers. Syn.* ii. 227.—Afr. austr.
coronopifolium, *Jacq. Ic. Pl. Rar.* iii. t. 526; *Coll.* v. 132.—Afr. austr.
cortusaefolium, Jacq. Ic. Pl. Rar. iii. t. 539 = saniculaefolium.
cortusaefolium, *L'Hérit. Geran.* t. 25.—Afr. austr.
cortusaefolium, E. Mey. in Drège, Zwei Pfl. Docum. 208 = ionidiflorum.
coruscans, *Hoare, ex Sweet, Geran.* t. 173.—Afr. austr.
corydaliflorum, DC. Prod. i. 652 = rapaceum.
corydalifolium, Harv. in Harv. & Sond. Fl. Cap. i. 269, sphalm. = praec.
corymbosum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 422 = tomentosum.
Cosmianum ×, Sweet, Geran. t. 189.
cosmon ×, Tratt. Pelarg. t. 239.
Cotyledon, G. Don, Gen. Syst. i. 732 = seq.
cotyledonifolium, Salisb. Prod. 315 = seq.
Cotyledonis, *L'Hérit. Geran.* t. 27.—Ins. S. Helen.
Couttsiae ×, Sweet, Geran. t. 269.
crassicaule, *L'Hérit. Geran.* t. 26; Curt. Bot. Mag. t. 477.—Afr. austr.
crassicaule, E. Mey. in Drège, Zwei Pfl. Docum. 209 = carnosum.
crassifolium, Hoffmgg. Verz. Pfl. 88 = crispum.
crassipes, *Harv. in Harv. & Sond. Fl. Cap.* i. 281.—Afr. austr.
crataegifolium, *Eckl. & Zeyh. Enum.* 82.—Afr. austr.
crataegifolium ×, Sweet, Geran. Suppl. t. 19.
crateriifolium, Hoffmgg. Verz. Pfl. 88 = calycinum.
crenaeflorum ×, Colv. ex Sweet, Hort. Brit. ed. I. 82.
crenatum, Hoffmgg. Verz. Pfl. 88 = zonale.
crenatum, *Link, Enum. Hort. Berol.* ii. 189.—Afr. austr.
crenulatum ×, Sweet, Geran. t. 162.
crinitum, *Harv. in Harv. & Sond. Fl. Cap.* i. 263.—Afr. austr.
crinitum, Nees, in Lehm. Pl. Preiss. i. 163 = australe.
crispilobum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 65.—Afr. austr.
crispum, *L'Hérit. Geran.* t. 32.—Afr. austr.
crispum-minus, Hort. ex Hoffmgg. Verz. Pfl. 91 = Hospes.
crithmifolium, *Sm. Ic. Pict.* i. t. 13.—Afr. austr.
crucifolium, *Hoffmgg. Verz. Pfl.* 88.—Afr. austr.?
cruentum, *Sweet, Geran.* t. 170.—Afr. austr.
Cuae, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Afr. austr.?
cucullatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 426.—Afr. austr.

PELARGONIUM:—
cucumerinum ×, Tratt. Pelarg. t. 63.
cucurbitaceum ×, Tratt. l. c. t. 123.
Cummingiae, *Sweet, Geran. Suppl.* t. 39.—Hab.?
cuneatum ×, Hoffmgg. Verz. Pfl. 88 = *mixtum*.
cuneatum, Spin, Cat. Hort. Sebast. (1818) 31 = praec.?
cuneiflorum ×, Sweet, Geran. t. 330.
curinbeticum, *Hort. Cels, ex Hoffmgg. Verz. Pfl.* 88.—Afr. austr.
Curtisianum, Balb. & Spin, ex DC. Prod. i. 667; Spreng. Syst. iii. 52 = pulchellum.
Curtisii, Hort. ex Hoffmgg. Verz. Pfl. 89 = adulterinum.
Curtisii, Jacq. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 51 = pulchellum.
curvatum, Steud. Nom. ed. II. i. 677, in syn. = aristatum, &c.
cuspidatum, Willd. Enum. Hort. Berol. Suppl. 47 = glaucum.
cycloides, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 65.—Hab.?
cynosbatifolium, Willd. Hort. Berol. t. 78 = malvaefolium.
Czerniniae ×, Tratt. Pelarg. t. 253.
Czerninianum ×, Tratt. l. c. t. 117.
daedaleum ×, Sweet, Geran. Suppl. t. 100.
damnoides, Hort. ex Hoffmgg. Verz. Pfl. 96 = thamnoides.
Darnleyanum ×, Sweet, Geran. Suppl. t. 63.
dasycaule, *Haw. Rev. Pl. Succ.* 309.—Afr. austr.
dasyphyllum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = crithmifolium.
daucifolium, Salisb. Prod. 312 = flavum.
daucifolium, Steud. Nom. ed. II. ii. 285 = triste.
daucoides, Jacq. Coll. Suppl. 143 = flavum.
Daveyanum ×, Sweet, Geran. t. 32.—Afr. austr.
Deburghae ×, Sweet, l. c. Suppl. t. 21.
decipiens, Steud. Nom. ed. II. ii. 285 = discipes?
decipiens ×, Tratt. Pelarg. t. 27.
decoloratum ×, Tratt. l. c. t. 120.
decorum ×, Colv. ex Sweet, Hort. Brit. ed. I. 81; et Geran. Suppl. t. 15.
degener, *Hoffmgg. Verz. Pfl.* 89.—Afr. austr.
delectabile, *Hoffmgg. l. c.*—Afr. austr.
deletum ×, Tratt. Pelarg. t. 59.
delicatum ×, Hoare, ex Sweet, Geran. t. 267.
delphinifolium, *Willd. Enum. Hort. Berol.* 708.—Afr. austr.
deltoideum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 65 (Quid?).—Hab.?
deltoideum, L'Hérit. ex DC. Prod. i. 650 = P. Ficaria.
dendriticum ×, Tratt. Pelarg. t. 90.
Dennisianum, *Sweet, Geran.* t. 20.—Afr. austr.
densiflorum, Steud. Nom. ed. II. ii. 285 = zonale.
dentatum ×, Sweet, Hort. Brit. ed. I. 81.—Hab.?
denticulatum, *Jacq. Hort. Schoenb.* ii. t. 135.—Afr. austr.
dependens ×, Sweet, Geran. t. 195.
depressum, Jacq. Coll. iv. 193 = longifolium.
depressum, Licht. ex Spreng. Syst. iii. 60 = lignosum.
dianthiflorum ×, Sweet, Hort. Brit. ed. I. 82.
diaphanum ×, Tratt. Pelarg. t. 101.
dichondraefolium, *DC. Prod.* i. 656.—Afr. austr.
dichotomum, *Otto & Dietr. Allg. Gartenz.* iii. (1835) 306.—Hab.?
didymum ×, Tratt. Pelarg. t. 52.
difforme ×, Sweet, Geran. t. 105.
dilatatum ×, Sweet, l. c. t. 314.
dilatilobum ×, Sweet, l. c. t. 378.
dilectum ×, Sweet, Hort. Brit. ed. I. 78, nomen.
dilutum ×, Sweet, l. c. 84; et Geran. t. 293.
dimacriaeflorum ×, Sweet, Geran. t. 220.
dimidiatum, Steud. Nom. ed. II. ii. 285 = alchemilloides.
diminutum ×, Tratt. Pelarg. t. 130.
dioicum, R. Br. in Ait. Hort. Kew. ed. II. iv. 162 = atrum.
dipetalum, *L'Hérit. Geran.* t. 43.—Afr. austr.
discipes, *Haw. Rev. Pl. Succ.* 305.—Afr. austr.
dissectum, Hort. ex Link, Enum. Hort. Berol. ii. 190 = Willdenowii.
dissectum, *Steud. Nom.* ed. II. ii. 285.—Afr. austr.
dissectum ×, Sweet, Geran. t. 247.
dissimile ×, Sweet, l. c. Suppl. t. 60.
distans, *L'Hérit. Geran. Ined.* 35; *ex DC. Prod.* i. 660.—Afr. austr.

PELARGONIUM:—
distantifolium, Hort. ex Steud. Nom. ed. II. ii. 285 = amplissimum.
distentifolium, Hort. ex Link, Enum. Hort. Berol. ii. 189 = amplissimum.
distichum, Dehnh. Cat. Pl. Hort. Camald. ed. II. 23 = strigosum, *Dehnh.*
distichum, *Hoffmgg. Verz. Pfl.* 89.—Afr. austr.
divaricatum, *DC. Prod.* i. 682.—Afr. austr.
divaricatum, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 52 = egregium.
divergens ×, Sweet, Geran. Suppl. t. 39.
diversiflorum ×, Tratt. Pelarg. t. 53.
diversifolium, Wendl. Bot. Beobacht. 52 = glaucum.
diversilobum ×, Sweet, Geran. t. 361.
Dobraeanum ×, Sweet, l. c. t. 253.
Dodonaei, *Hoffmgg. Verz. Pfl.* 89.—Afr. austr.
dondiaefolium, *Link, Enum. Hort. Berol.* ii. 187.—Afr. austr.
Drakeae ×, Sweet, Geran. Suppl. t. 96.
Dregeanum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 422 = cordatum.
Drummondii, *Turcz. l. c.* 421.—Austral.
dubium ×, Tratt. Pelarg. t. 11.
dumosum ×, Sweet, Geran. t. 19.
durum, *Hoffmgg. Verz. Pfl.* 89.—Afr. austr.
dysgenon ×, Tratt. Pelarg. t. 214.
echinatum, *Curt. Bot. Mag.* t. 309.—Afr. austr.
Ecklonii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 419.—Afr. austr.
egregium, *Hort. Belved.* 71; *ex Link, Enum. Hort. Berol.* ii. 195.—Afr. austr.?
elatum, *DC. Prod.* i. 657; *Colla, Hort. Ripul.* 103. t. 5.—Afr. austr.
elatum, G. Don, Gen. Syst. i. 731 = violarium.
elatum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = caespitosum.
electum ×, Sweet, Geran. t. 238.
elegans, Hort. ex Hoffmgg. Verz. Pfl. 86 = anodaefolium.
elegans, *Willd. Sp. Pl.* iii. 655.—Afr. austr.
elegantissimum, Hort. Burdin, ex Colla, Hort. Ripul. App. ii. 354 = Burdini.
elegantulum, *Steud. Nom.* ed. II. ii. 292.—Afr. austr.?
Elisae ×, Tratt. Pelarg. t. 205.
ellipticum, *DC. Prod.* i. 681.—Afr. austr.
elongatum, Steud. Nom. ed. II. ii. 285 = tabulare.
emancipatum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 66.—Afr. austr.
emarginatum, Hort. ex F. G. Dietr. l. c. (Quid?).—Hab.?
emarginatum, *Moench, Meth. Suppl.* 296.—Hab.?
emarginatum, *Wendl. ex Breit. Hort. Breit.* 334.—Afr. austr.
Endlicherianum, *Fenzl, Pugill. Pl. Nov. Syr.* 6; *et in Belg. Hortic.* vi. (1856) 133.—Oriens.
enigmaticum ×, Tratt. Pelarg. t. 86.
ensatum, *DC. Prod.* i. 680.—Afr. austr.
epiphanon ×, Tratt. Pelarg. t. 251.
eratinum ×, Sweet, Geran. Suppl. t. 30.
erectum ×, Sweet, Geran. t. 187.
eriocaulon ×, Sweet, l. c. t. 357.
eriophorum ×, Sweet, Hort. Brit. ed. I. 79.
eriophyllum ×, Sweet, Geran. t. 141.
eriosepalum ×, Sweet, l. c. t. 375.
eriostemum, Jacq. Hort. Schoenb. ii. 4. t. 132 = ovale.
erodioides, Hook. Journ. Bot. i. (1834) 252 = australe.
erosum ×, Tratt. Pelarg. t. 264.
errans, Hort. ex Link, Enum. Hort. Berol. ii. 193 = P. Exsul.
erubescens, *Spin, Cat. Hort. Sebast.* (1809) 27.—Hab.?
Eryngii ×, Tratt. Pelarg. t. 84.
erythraeum, *DC. Prod.* i. 681.—Afr. austr.
erythraeum, Hoffmgg. Verz. Pfl. 89 = *saepeflorens*.
erythrophyllum, *Steud. Nom.* ed. II. ii. 285.—Afr. austr.
Essexiae ×, Colv. ex Sweet. Hort. Brit. ed. i. 82, nomen.
eugenianum ×, Tratt. Pelarg. t. 223.
eupatoriaefolium, *Steud. Nom.* ed. II. ii. 286.—Afr. austr.
euphraseum, *Hornem. Hort. Hafn.* ii. 634.—Afr. austr.
euryphyllum ×, Tratt. Pelarg. t. 129.
exaratum ×, Tratt. l. c. 211.
exetelon ×, Tratt. l. c. 238.
exile ×, Sweet, Hort. Brit. ed. I. 78.
eximium, Hoffmgg. Verz. Pfl. 90 = delectabile.
eximium ×, Sweet. Geran. t. 26.

PELARGONIUM:—
exochum ×, Tratt. Pelarg. t. 131.
exornatum ×, Sweet, Hort. Brit. ed. I. 80.
expansum ×, Sweet, l. c. 82; Steud. Nom. ed. II. ii. 286.—Afr. austr.
exquisitum, *Hoffmgg. Verz. Pfl.* 90.—Afr. austr.
exquisitum ×, Sweet, Geran. Suppl. t. 55.
exstipulatum, *L'Hérit. Geran.* t. 35.—Afr. austr.
Exsul, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 66.—Afr. austr.
Fairliae ×, Sweet, Geran. t. 178.
familiare, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 66.—Afr. austr.?
fasciculatum ×, Colv. ex Sweet, Hort. Brit. ed. I. 79.
fastuosum ×, Sweet, Geran. Suppl. t. 93.
fastuosum ×, Tratt. Pelarg. t. 4.
Fauxiae ×, Forbes, Hort. Woburn. 152.
Felderi, Hort. Wendl. ex Hoffmgg. Verz. Pfl. 89, in syn. = seq.
Felteri, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 52 = egregium.
fenestratum ×, Tratt. Pelarg. t. 94.
Ferdinandeum ×, Tratt. l. c. t. 49.
ferulaceum, *Willd. Sp. Pl.* iii. 687.—Afr. austr.
festivum, Hoffmgg. Verz. Pfl. 90 = cordatum.
Ficaria, E. Mey. ex Harv. & Sond. Fl. Cap. i. 262 = Meyeri.
Ficaria, *Willd. Sp. Pl.* iii. 643.—Afr. austr.
ficifolium, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 66.—Afr. austr.
ficifolium ×, Tratt. Pelarg. t. 206.
filipendulifolium, Sweet, Geran. t. 85 = triste.
fimbriatum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = amatymbicum.
Fischerianum, Dehnh. Cat. Pl. Hort. Camald. ed. II. 21 (Quid?).—Afr. austr.
fissifolium, *Pers. Syn.* ii. 227.—Afr. austr.
fissum, *Baker, in Saund. Refug. Bot.* t. 149.—Afr. austr.
flabellatum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 21.—Hab.?
flabellifolium, *Harv. in Harv. & Sond. Fl. Cap.* i. 276.—Afr. austr.
flabellifolium ×, Sweet, Geran. Suppl. t. 48.
flaccidum ×, Sweet, Hort. Brit. ed. I. 471; Geran. t. 337.
flagrans, Sweet, Geran. t. 16.
flammeum ×, Colv. ex Sweet, Hort. Brit. ed. I. 79; Forbes, Hort. Woburn. 151.
flavescens, Steud. Nom. ed. II. ii. 286 = tetragonum.
flavum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 418.—Afr. austr.
flexuosum, *DC. Prod.* i. 680.—Afr. austr.
flexuosum ×, Sweet, Geran. t. 180.
floccosum ×, Sweet, l. c. 129.
floribundum, Breit. Hort. Breit. 334 = undulatum.
floribundum, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 163.—Afr. austr.
floridum, *Sweet, Geran.* t. 41.—Afr. austr.?
Flos-jambeae, Steud. Nom. ed. II. ii. 293 = *Foljambeae*.
foliolosum, DC. Prod. i. 653 = astragalifolium.
foliosum ×, Sweet, Hort. Brit. ed. I. 471; Geran. t. 340.
Foljambeae ×, Sweet, Geran. t. 312.
Footianum ×, Forbes, Hort. Woburn. 151.
formosissimum, *Pers. Syn.* ii. 231.—Afr. austr.
formosum, Balb. ex Spreng. Syst. iii. 61 = scabrum.
formosum, Desf. Arb. i. 459 = formosissimum.
formosum, Sweet, Geran. t. 120 = Willdenowii.
formosum-grandiflorum, Cels, ex Hoffmgg. Verz. Pfl. 92 = magniflorum.
Forsterianum ×, Forbes, Hort. Woburn. 152.
Fothergillii, Sweet, ex Kern. Hort. t. 469.—Hab.?
fragile, Hort. ex Hoffmgg. Verz. Pfl. 92 = laeve.
fragile, Willd. Sp. Pl. iii. 686 = tripartitum.
fragrans, E. Mey. in Drège, Zwei Pfl. Docum. 209 = althaeoides.
fragrans, *Willd. Hort. Berol.* i. t. 77.—Hab.?
fragrantissimum, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 54 = praec.?
fragrantissimum, *Link, Enum. Hort. Berol.* ii. 192.—Afr. austr.
Francisceum ×, Tratt. Pelarg. t. 198 = seq.?
Francisii ×, Sweet, Hort. Brit. ed. I. Add. 471.
fraternum, *Schrank, Pl. Rar. Hort. Monac.* t. 16.—Afr. austr.
Friderici-Guilelmi, *Klier, ex Otto & Dietr. Allg. Gartenz.* v. (1837) 202.—Hab.?
fruticosum, DC. Prod. i. 661 = myrrhifolium.
fruticosum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = divaricatum.

PELARGONIUM:—
fruticosum, *Willd. Sp. Pl.* iii. 689.—Afr. austr.
Fuerstenbergianum ×, Tratt. Pelarg. t. 222.
Fuerstianum ×, Tratt. l. c. t. 228.
fulgens, Hoffmgg. Verz. Pfl. 90 = formosissimum.
fulgens, *Steud. Nom.* ed. II. ii. 293.—Hab.?
fulgidum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 422.—Afr. austr.
fulminans, *Hoffmgg. Verz. Pfl.* 90.—Afr. austr.?
fumarioides, *L'Hérit. ex Harv. & Sond. Fl. Cap.* i. 290.—Afr. austr.
fuscatum, Jacq. Ic. Pl. Rar. iii. 450 = saniculaefolium.
fuscatum, Sweet, Geran. t. 210 = seq.
fusciflorum, Sweet, Geran. errat. sub t. 215.
Galatheae ×, Tratt. Pelarg. t. 213.
gangrenosum ×, Tratt. l. c. t. 112.
Gardeniae, Hort. ex Steud. Nom. ed. II. ii. 286 = seq.
Gardneriae ×, Colv. ex Sweet, Hort. Brit. ed. I. 80.
geifolium, E. Mey. in Drège, Zwei Pfl. Docum. 209 = heracleifolium.
Geislernianum ×, Tratt. Pelarg. t. 121.
geniculatum, Dehnh. Cat. Pl. Hort. Camald. ed. II. 22 (Quid?).—Afr. austr.?
geranioides, Steud. Nom. ed. II. ii. 286 = alchemilloides.
germanum, Hort. ex Tratt. Pelarg. sub t. 42 = Amaryllidis.
Ghilini, *Spin, Cat. Hort. Sebast.* (1818) 31.—Hab.?
gibbosum, *L'Hérit. ex* [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 422.—Afr. austr.
giganteum ×, Tratt. Pelarg. t. 215.
Giondani, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22 (Quid?).—Hab.?
glabellum, Steud. Nom. ed. II. ii 293 = seq.
glabrellum, *Link, Enum. Hort. Berol.* ii. 190.—Hab.?
glabrescens ×, Sweet, Geran. Suppl. t. 45.
glabrifolium, *Steud. Nom.* ed. II. ii. 293.—Afr. austr.?
glabrum, *Hoffmgg. Verz. Pfl.* 90.—Afr. austr.?
glaciale ×, Tratt. Pelarg. t. 58.
gladioloides ×, Tratt. l. c. t. 29.
glaucifolium, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22 (an seq.?).—Hab.?
glauciifolium ×, Sweet, Geran. t. 179.
glaucoides, Wendl. Coll. 63. t. 63 = glaucum.
glaucophyllum, Steud. Nom. ed. II. ii. 286, sphalm. = praec.
glaucum, *L'Hérit. Geran.* t. 29.—Afr. austr.
glechomoides, *A. Rich. Tent. Fl. Abyss.* i. 118.—Afr. trop.
glomeratum, Jacq. f. Eclog. i. 146. t. 98 = australe.
glorianum ×, Sweet, Geran. Suppl. t. 82.
gloriosum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 67.—Afr. austr.
glutinosum, *L'Hérit. Geran.* t. 20.—Afr. austr.
glutinosum, Van Eeden, ex Hoffmgg. Verz. Pfl. 89 = Dodonaei.
Gordoniae ×, Forbes, Hort. Woburn. 152.
gossypiifolium ×, Sweet, Hort. Brit. ed. II. 98.
Goweri, *Sweet, l. c.* ed. I. Add. 471.—Afr. austr.?
gracile, *Steud. Nom.* ed. II. ii. 286.—Afr. austr.
gracilescens ×, Sweet, Hort. Brit. ed. I. 78.
Graftoniae ×, Forbes, Hort. Woburn. 152.
grandidentatum ×, Sweet, Geran. t. 217.
grandiflorum, *Willd. Sp. Pl.* iii. 674.—Afr. austr.
graphicum ×, Sweet, Geran. Suppl. t. 12.
gratum, *Willd. Enum. Hort. Berol.* 710.—Hab.?
graveolens, *L'Hérit. Geran.* t. 17.—Afr. austr.
Grenvilleae, *Harv. in Harv. & Sond. Fl. Cap.* i. 265.—Afr. austr.
Grevillianum, Hort. ex Steud. Nom. ed. II. ii. 286 = conspicuum.
Grimaldiae, Spin, Cat. Hort. Sebast. (1818) 31 (Quid?).—Hab.?
grossulariaefolium, Salisb. Prod. 313 = seq.
grossularioides, [Soland. in] Ait. Hort. Kew. ed. I. ii. 42 = australe.
gruinale ×, Tratt. Pelarg. t. 165.
Gurneyanum ×, Sweet, Geran. t. 393.
Gussonii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Hab.?
gymnophyllum ×, Tratt. Pelarg. t. 134.
haemanthon ×, Sweet, Hort. Brit. ed. I. 78.
haemastictum ×, Sweet, Geran. Suppl. t. 18.
hamatum, Jacq. Hort. Schoenb. ii. 7 = echinatum.
Hammersleiae ×, Sweet, Geran. t. 225.
hardenbergense, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Hab.?

PELARGONIUM:—
Harewoodiae ×, Sweet, Geran. t. 389.
hederaefolium, Salisb. Prod. 315 = lateripes.
Hellerii, *Hort. ex Hoffmgg. Verz. Pfl.* 90.—Afr. austr.
Henricae ×, Tratt. Pelarg. t. 146.
hepaticaefolium, Hoffmgg. Verz. Pfl. 91 = lateripes.
hepaticaefolium, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 31; *Nov. Prov.* 31.—Afr. austr.
heracleifolium, *Lodd. Bot. Cab.* t. 437.—Afr. austr.
heraldicum ×, Tratt. Pelarg. t. 56.
Heritieri, *Sweet, Hort. Brit.* ed. I. 83; ed. II. 100.—Afr. austr.
hermanniaefolium, Jacq. Ic. Pl. Rar. iii. t. 545; Coll. Suppl. 137 = crispum.
Herminae ×, Tratt. Pelarg. t. 242.
Heselrigii, Sweet, Hort. Brit. ed. I. 84 = seq.
Hesilrigeae ×, Sweet, Geran. t. 319.
hesperideum ×, Tratt. Pelarg. t. 18.
heterogamum, *L'Hérit. Geran.* t. 18.—Afr. austr.
heterolobum, *DC. Prod.* i. 680.—Afr. austr.
heteromallum ×, Sweet, Hort. Brit. ed. I. 79.
heterophyllum, *Jacq. Ic. Pl. Rar.* iii. 516; *Coll.* iv. 197.—Afr. austr.
heterophyllum, L'Hérit. ex DC. Prod. i. 652 = atrum.
heterotrichum ×, Sweet, Hort. Brit. ed. I. 84; et Geran. t. 303.
hibiscifolium ×, Sweet, Hort. Brit. ed. I. 79.
hibiscinum ×, Tratt. Pelarg. t. 99.
hibridaefolium, Steud. Nom. ed. II. ii. 286 = heracleifolium.
hilare, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 67.—Afr. austr.?
Hillianum ×, Sweet, Geran. Suppl. t. 51.
Hippocrepis, L'Hérit. ex DC. Prod. i. 668 = saniculaefolium.
hirsutum, [Soland. in] Ait. Hort. Kew. ed. I. ii. 417 = atrum.
hirtum, Jacq. Ic. Pl. Rar. iii. 536; Coll. v. 144 = tenuifolium.
hirtum, Willd. Sp. Pl. iii. 644 = longifolium.
hispidum, *Willd. l. c.* 677.—Afr. austr.
Histrio, Hoffmgg. Verz. Pfl. 91 = elatum.
hoaraeflorum ×, Sweet, Geran. t. 133.
Hoareanum ×, Sweet, l. c. t. 80.
Hoffmannseggii, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 32.—Afr. austr.
Hollandianum ×, Forbes, Hort. Woburn. 152.
holosericeum, Hoffmgg. Verz. Pfl. 91 = ovale.
homonymum, *Steud. Nom.* ed. II. ii. 293.—Afr. austr.?
homophyllum ×, Tratt. Pelarg. t. 106.
honestum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 67.—Afr. austr.
Hornemanni, *DC. Prod.* i. 681.—Afr. austr.?
Hospes, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 67.—Afr. austr.
Humei ×, Sweet, Geran. t. 324.
humifusum, DC. Prod. i. 654 = australe.
humifusum, Willd. Enum. Hort. Berol. Suppl. 47 = parvulum.
humile, Hort. ex Hoffmgg. Verz. Pfl. 91 = scandens.
huraefolium, Sweet, Geran. sub t. 281 = luridum.
Husseyanum ×, Sweet, l. c. t. 92.
hybridaefolium, Harv. in Harv. & Sond. Fl. Cap. i. 272 = heracleifolium.
hybridum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 424.—Afr. austr.
hypoleucum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 421.—Afr. austr.
hypotrachys, *Hoffmgg. Verz. Pfl.* 91.—Afr. austr.
Hystrix, *Harv. in Harv. & Sond. Fl. Cap.* i. 280.—Afr. austr.
iatrophaefolium, Steud. Nom. ed. II. ii. 287 = jatrophaefolium.
ichorosum ×, Tratt. Pelarg. t. 118.
iconicum ×, Sweet, Geran. Suppl. t. 88.
Idion ×, Tratt. Pelarg. t. 115.
ignescens, *Lodd. Bot. Cab.* ii. t. 109.—Afr. austr.?
Illecebrum ×, Tratt, Pelarg. t. 186.
illustre, *Steud. Nom.* ed. II. ii. 287.—Afr. austr.
imbricatum ×, Sweet, Geran. t. 65.
imbutum, *Colv. ex Sweet, Hort. Brit.* ed. I. 81; *Hort. ex Steud. Nom.* ed. II. ii. 293.—Afr. austr.
Immelinianum ×, Tratt. Pelarg. t. 155.
imperatoriae ×, Tratt. l. c. t. 110.
imperiale, Hoffmgg. Verz. Pfl. 91 = cordatum.
imperiale ×, Sweet, Geran. t. 365.
implicatum ×, Sweet, Geran. Suppl. t. 86.
impurum ×, Tratt. Pelarg. t. 157.

PELARGONIUM:—
incanescens, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.?
incanescens ×, Sweet, Geran. t. 208.
incanum ×, Willd. Enum. Hort. Berol. Suppl. 47.
incarnatum, *Moench, Meth. Suppl* 295.—Afr. austr.?
incarnatum ×, Sweet, Geran. t. 308.
incisum, *Willd. Sp. Pl.* iii. 686.—Afr. austr.
incomparabile ×, Sweet, Hort. Brit. ed. I. 79.
incrassatum, *Sims, Bot. Mag.* t. 761.—Afr. austr.
incurvum ×, Sweet, Geran. t. 249.
Infans ×, Tratt. Pelarg. t. 73.
infundibuliforme ×, Tratt. l. c. t. 103.
inodorum, Willd. Enum. Hort. Berol. 702 = australe.
inquinans, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 424.—Afr. austr.
inscriptum ×, Sweet, Geran. t. 193.
insculptum ×, Sweet, Geran. Suppl. t. 65.
insigne, *Hoffmgg. Verz. Pfl.* 91.—Afr. austr.
insignitum ×, Sweet, Geran. t. 300.
instratum ×, Sweet, l. c. Suppl. t. 78.
integrifolium ×, Tratt. Pelarg. t. 72.
interlitum ×, Sweet, Hort. Brit. ed. I. 80.
intermedium ×, Sweet, l. c. 78; et Tratt. Pelarg. t. 170.
intermixtum ×, Sweet, Hort. Brit. ed. I. 82.
intertextum ×, Sweet, Geran. t. 185.
involucratum ×, Sweet, l. c. t. 33.
iocastum, Steud. Nom. ed. II. ii. 287 = grossularioides.
ionidiflorum, *Steud. l. c.*—Afr. austr.
Irbyana ×, Colv. ex Sweet, Hort. Brit. ed. I. 81; Hort. ex Steud. l. c.
irregulare ×, Sweet, Hort. Brit. ed. I. 79.
ixiaeflorum ×, Tratt. Pelarg. t. 67.
Jacquinianum ×, Tratt. l. c. t. 125.
Jannackii, *Hoffmgg. Verz. Pfl.* 91.—Afr. austr.
jatrophaefolium, *DC. Prod.* i. 679.—Afr. austr.
Jenkinsoni ×, Sweet, Geran. t. 154.
Jenseni, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.?
Jeschekianum ×, Tratt. Pelarg. t. 156.
Joanneum ×, Tratt. l. c. t. 200.
jocundum ×, Hoare, ex Sweet, Hort. Brit. ed. I. 81.
jonquillinum ×, Sweet, Geran. t. 241.
Josephinae ×, Lee, ex Sweet, Hort. Brit. ed. I. 80.
Josephinum ×, Tratt. Pelarg. t. 145.
Julianae ×, Tratt. l. c. t. 257.
Kalmianum, Van Eeden, ex Hoffmgg. Verz. Pfl. 96 = spurium.
Keglevicsianum ×, Tratt. Pelarg. t. 256.
Kenrickae ×, Sweet, Geran. Suppl. t. 58.
Kernhoferianum ×, Tratt. Pelarg. t. 39.
Kingii ×, Sweet, Geran. t. 248.
Klierianum ×, Tratt. Pelarg. t. 132.
Knappianum ×, Tratt. l. c. t. 151.
Knipeae ×, Sweet, Geran. t. 372.
Kolowrathiae ×, Tratt. Pelarg. t. 220.
Kolowrathianum ×, Tratt. l. c. 137.
Krebnerianum ×, Tratt. l. c. t. 150.
Kunthii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.
labyrinthicum ×, Sweet, Hort. Brit. ed. I. 80; G. Don, Gen. Syst. i. 726.—Afr. austr.?
lacerum, Jacq. Ic. Pl. Rar. iii. t. 532; *Coll.* v. 122 = myrrhifolium.
laciniatum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = barbatum.
laciniatum, Pers. Syn. ii. 228 = longifolium.
laciniatum, *Steud. Nom.* ed. II. ii. 293.—Afr. austr.
Laconiae ×, Sweet, Geran. t. 374.
ladanosma, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 68.—Afr. austr.
laetum ×, Tratt. Pelarg. t. 37.
laeve, *Otto, ex Hoffmgg. Verz. Pfl.* 92.—Afr. austr.
laevigatum, *Willd. Sp. Pl.* iii. 685.—Afr. austr.
Lamberti ×, Sweet, Geran. t. 104.
lanatum, DC. Prod. i. 681 = cordatum.
lanceaefolium, G. Don, Gen. Syst. i. 729 = longifolium.
lanceolatum, Hort. ex Kern. Hort. t. 632 = glaucum.
lanceolatum, *Steud. Nom.* ed. II. ii. 287.—Hab.?
lancifolium, *Steud. l. c.*—Afr. austr.
lanosum ×, Sweet, Geran. Suppl. t. 84.
lanuginosum, *Hoffmgg. Verz. Pfl.* 92.—Afr. austr.
lappaceum, *Hoffmgg. l. c.*—Afr. austr.
lasiocaulon ×, Sweet, Geran. t. 364.
lasiophyllum ×, Sweet, Hort. Brit. ed. I. 83; et Geran. t. 296.
lateripedatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = lateripes.

PELARGONIUM:—
lateripes, *L'Hérit. Geran.* t. 24.—Afr. austr.
lateritium, Willd. Enum. Hort. Berol. Suppl. 47 = malvaefolium.
Lathyrus ×, Tratt. Pelarg. t. 259.
latidentatum ×, Sweet, Geran. Suppl. t. 28.
latifolium ×, Sweet, Geran. t. 335.
latilobum ×, Sweet, l. c. t. 236.
lautum ×, Sweet, l. c. Suppl. t. 91.
lavateraefolium, *Steud. Nom.* ed. II. ii. 287.—Afr. austr.
Lawranceanum ×, Sweet, Geran. Suppl. t. 23.
laxatum, Harv. in Harv. & Sond. Fl. Cap. i. 274 = triste.
laxiflorum ×, Sweet, Geran. t. 216.
laxulum ×, Sweet, l. c. Suppl. t. 75.
laxum, G. Don, Gen. Syst. i. 730 = crithmifolium.
laxum, Steud. Nom. ed. II. ii. 287 = carnosum.
laxum ×, Tratt. Pelarg. t. 260.
Lechianum, *Spin, Cat. Hort. Sebast.* (1818) 32.—Afr. austr.?
Leeanum, *Loud. Hort. Brit.* 271.—Afr. austr.
Leghkeckae ×, Sweet, Geran. t. 377.
leion ×, Tratt. Pelarg. t. 247.
Leontinae ×, Tratt. l. c. t. 221.
Leopoldinum, Hoffmgg. Verz. Pfl. 92 = *Thynneae*.
lepidum ×, Sweet, Geran. t. 156.
leptopetalum, *Steud. Nom.* ed. II. ii. 287.—Afr. austr.
lessertiaefolium, Steud. l. c. = astragalifolium.
leucanthum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 48 = zonale.
leucophyllum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 421.—Afr. austr.
leve, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 38 = ranunculifolium.
lignosum, *Spreng. Syst.* iii. 60.—Afr. austr.
lingulatum ×, Sweet, Hort. Brit. ed. I. 84; Geran. t. 301.
lilacinum ×, Tratt. Pelarg. t. 164.
Lima ×, Tratt. Pelarg. t. 135.
limbatum, *Hoffmgg. Verz. Pfl.* 92.—Afr. austr.
limbatum, E. Mey, in Drège, Zwei Pfl. Docum. 209 = barbatum.
limoneum ×, Sweet, Hort. Brit. ed. I. 81; Geran. t. 278.
lineare, *Pers. Syn.* ii. 228.—Afr. austr.
lineatum ×, Sweet, Geran. t. 16.
lineolatum ×, Tratt. Pelarg. t. 169.
litigiosum ×, Tratt. l. c. t. 85.
Littleanum ×, Sweet, Geran. Suppl. t. 72.
littorale, Huegel, in Enum. Pl. Hueg. 44 = australe.
lividum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 68 = nothum.
lividum ×, Sweet, Hort. Brit. ed. I. 80.
lobatum, Hort. ex Hoffmgg. Verz. Pfl. 92 = lappaceum.
lobatum, *Willd. Sp. Pl.* iii. 650.—Afr. austr.
Locusta ×, Tratt. Pelarg. t. 248.
longicaule, *Jacq. Ic. Pl. Rar.* iii. t. 533; Coll. v. 125.—Afr. austr.
longiflorum, Jacq. l. c. t. 521 = seq.
longifolium, *Jacq. l. c.* t. 518; *Coll.* iv. 194.—Afr. austr.
longipedicellatum ×, Tratt. Pelarg. t. 163.
longipes, *Hoffmgg. Verz. Pfl.* 92.—Afr. austr.
longipetalum ×, Tratt. Pelarg. t. 9.
Loudonianum ×, Sweet, Geran. Suppl. t. 17.
Lousadianum, *Sweet, Geran.* t. 44.
lucidum, *Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 68.—Afr. austr.
lucidum ×, Sweet, Geran. t. 373.
luculentum ×, Sweet, Hort. Brit. ed. I. 78.
Ludovicae ×, Tratt. Pelarg. t. 195.
Ludovicianum ×, Tratt. l. c. t. 69.
luridum, *Sweet, Geran.* t. 281.—Afr. austr.
luteum, *G. Don, Gen. Syst.* i. 726.—Afr. austr.
luxurians ×, Tratt. Pelarg. t. 173.
macranthum, Hoffmgg. Verz. Pfl. 92 = *magniflorum*.
macranthum ×, Sweet, Geran. t. 83.
macrolobum, *Hoffmgg. Verz. Pfl.* 92.—Afr. austr.
macrostephanon ×, Tratt. Pelarg. t. 143.
magnificiens, Van Eeden, in Hoffmgg. Verz. Pfl. 92 = magnificum.
magnificum, *Cels, ex Hoffmgg. l. c.*—Afr. austr.?
magniflorum, *Hoffmgg. l. c.* 92.—Hab.?
magnifolium ×, Sweet, Geran. Suppl. t. 10.
magnistipulatum ×, Sweet, Hort. Brit. ed. II. 93.
mahernium, L'Hérit. ex DC. Prod. i. 656 = capillare.
majale ×, Tratt. Pelarg. t. 41.
majestum, Hort. ex Hoffmgg. Verz. Pfl. 97 = *villosum*.

PELARGONIUM:—
malachraefolium ×, Sweet, l. c. Suppl. t. 2.
malacophyllon ×, Tratt. Pelarg. t. 216.
malacophyllum ×, Sweet, Geran. t. 397.
malvaceum ×, Sweet, Hort. Brit. ed. I. 82.
malvaefolium, *Jacq. f. Eclog.* i. 145. t. 97.—Afr. austr.
mamillaricaule, Hort. ex Steud. Nom. ed. II. ii. 287 = crassicaule.
mammulosum, *Wendl. Coll.* ii. 77. t. 70.—Afr. austr.
marginatum, *Link, Enum. Hort. Berol.* ii. 189.—Afr. austr.
Mariae ×, Tratt. Pelarg. t. 204.
Markianum ×, Tratt. l. c. t. 10.
marmoratum, Steud. Nom. ed. II. ii. 287 = patulum.
Mathildinae ×, Tratt. Pelarg. t. 254.
Mattocksianum ×, Sweet, Geran. t. 234.
Maurii, Dehnh. Cat. Pl. Hort. Camald. ed. II. 22. (Quid?).—Hab.?
maximum, *Hort. ex Hoffmgg. Verz. Pfl.* 93.—Afr. austr.?
medusulum ×, Tratt. Pelarg. t. 57.
megalanthum ×, Sweet, Geran. Suppl. t. 4.
megaleion ×, Sweet, l. c. Suppl. t. 5.
megalostictum ×, Sweet, l. c. t. 8.
megiston ×, Tratt. Pelarg. t. 210.
meifolium, Meissn. ex Krauss, in Flora, xxvii. (1844) 299 = artemisiaefolium.
melalophon ×, Tratt. Pelarg. t. 13.
melananthum, Jacq. Ic. Pl. Rar. iii. t. 514 = atrum.
melancholicum ×, Sweet, Geran. Suppl. t. 53.
melanostictum, *Sweet, l. c.* t. 6.—Hab.?
melissimum ×, Sweet, Geran. t. 5.
metallicum ×, Tratt. Pelarg. t. 263.
metempsychodes ×, Tratt. l. c. t. 3.
Metternichianum ×, Tratt. l. c. t. 77.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* i. 262.—Afr. austr.
micans ×, Tratt. Pelarg. t. 184.
micranthum, *G. Don, Gen. Syst.* i. 733.—Afr. austr.
micranthum, Eckl. & Zeyh. Enum. 79 = tomentosum.
micranthum, Presl, Bot. Bemerk. 27 = australe.
micropetalum, E. Mey, in Drège, Zwei Pfl. Docum. 209 = australe.
microphyllum, *Steud. Nom.* ed. II. ii. 287.—Afr. austr.
militare, Hoffmgg. Verz. Pfl. 93.—Afr. austr.
millefoliatum, Sweet, Geran. t. 230 = triste.
Milleri ×, Sweet, Hort. Brit. ed. I. 84.
minimum, *Willd. Sp. Pl.* iii. 664.—Afr. austr.
mirabile ×, Tratt. Pelarg. t. 136.
mirabile ×, Young, ex Sweet, Geran. Suppl. t. 57.
mitellaefolium, *Sweet, Hort. Brit.* ed. II. 97.—Afr. austr.
mixtum ×, Sweet, Geran. t. 71.
modestum ×, Sweet, l. c. t. 204.
moestum, Hoffmgg. Verz. Pfl. 93 = triste.
molle, *Hoffmgg. l. c.*—Afr. austr.
mollifolium ×, Sweet, Geran. Suppl. t. 77.
moniliforme, *E. Mey, in Drège, Zwei Pfl. Docum.* 209; *et ex Harv. in Harv. & Sond. Fl. Cap.* i. 264.—Afr. austr.
monsoniaefolium, Fisch. ex Steud. Nom. ed. I. 601 = incisum.
monstrosum, Pers. Syn. ii. 231 = seq.
monstrum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 425.—Afr. austr.
Moreanum ×, Sweet, Hort. Brit. ed. I. 84.
morifolium ×, Tratt. Pelarg. t. 107.
moschatum, Andr. ex Sweet, Hort. Brit. ed. I. 80 = bellulum.
Mostynae ×, Sweet, Geran. t. 10.
mucronatum ×, Sweet, l. c. t. 275.
multibracteatum, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 29; *et ex A. Rich. Tent. Fl. Abyss.* i. 118.—Abyss.
multicaule, *Jacq. Ic. Pl. Rar.* iii. 534; *Coll.* v. 126.—Afr. austr.
multidentatum ×, Sweet, Hort. Brit. ed. I. 82.
multifidum, *Harv. in Harv. & Sond. Fl. Cap.* i. 282.—Afr. austr.
multifidum, *Hort. ex Hoffmgg. Verz. Pfl.* 93.—Afr. austr.
multiflorum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 48.—Afr. austr.
multinerve ×, Sweet, Geran. t. 17.
multiradiatum, Harv. in Harv. & Sond. Fl. Cap. i. 274, in syn. = triste.

PELARGONIUM:—
multiradiatum, E. Mey. ex Harv. & Sond. Fl. Cap. i. 273 = radulaefolium.
multiradiatum, *Wendl. Coll.* ii. 56. t. 59; *Link, Enum. Hort. Berol.* ii. 191.—Afr. austr.
mundulum ×, Sweet, Hort. Brit. ed. I. 84.
munitum, Burch. Trav. i. 225 = crithmifolium.
Murrayanum, *Colla, Hort. Ripul. App.* iii. 5.—Afr. austr?
mutabile, Dehnh. Cat. Pl. Hort. Camal. 26; ed. II. 22. (Quid?).—Hab. ?
mutabile, E. Mey. in Drège, Zwei Pfl. Docum. 209 = australe.
mutabile ×, Sweet, Geran. t. 213.
Muthianum ×, Tratt. Pelarg. t. 226.
myrrhifolium, *L'Hérit. ex* [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 421.—Afr. austr.
Myrtillum ×, Haw. ex Sweet, Hort. Brit. ed. I. 81.
Nairnii ×, Sweet, Geran. t. 376.
nanum, L'Hérit. ex DC. Prod. i. 660 = parvulum.
nanum ×, Sweet, Geran. t. 102.
neapolitanum, Van Eeden, ex Steud. Nom. ed. II. ii. 288 = adulterinum.
nervifolium, *Jacq. Ic. Pl. Rar.* iii. t. 517; *Coll.* iv. 198.—Afr. austr.
nervosum ×, Sweet, Geran. t. 47.
Newshamianum ×, Sweet, l. c. t. 144.
nigrescens, *Sweet, ex Steud. Nom.* ed. II. ii. 288 (Quid?).—Afr. austr.?
nitidum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.?
nissonianum, Lecchi, ex Colla, Hort. Ripul. App. iii. 8, 10 = Berterianum.
nitens ×, Tratt. Pelarg. t. 33.
nitidum ×, Sweet, Hort. Brit. ed. I. 77.
Niveni, *Harv. in Harv. & Sond. Fl. Cap.* i. 271.—Afr. austr.
niveum, *Loud. Hort. Brit.* 277.—Afr. austr.
niveum ×, Tratt. Pelarg. t. 23.
nobile, *Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 68.—Afr. austr.
nodosum, *Sweet, Geran. Suppl.* t. 68.
Nonfordium ×, Forbes, Hort. Woburn. 1 3.
notabile ×, Sweet, Hort. Brit. ed I. 81.
notatum ×, Sweet, Geran. t. 208.
nothum, *Willd. Enum. Hort. Berol.* 710.—Afr. austr.?
novum, Hort. ex Steud. Nom. ed. II. ii. 294 = zonale.
nubilum ×, Sweet, Geran. Suppl. t. 20.
nummulariaefolium, Harv. in Harv. & Sond. Fl. Cap. i. 289 = australe.
nummulifolium, Salisb. Parad. Lond. t. 23 = australe?
nutans, DC. Prod. i. 651 = rapaceum.
nutans ×, Sweet, Geran. Suppl. t. 66.
obcordatum ×, Sweet, Hort. Brit. ed. I. 83.
oblatum ×, Sweet, Geran. t. 35.
obliquum ×, Tratt. Pelarg. t. 240.
oblongatum, *E. Mey. in Drège, Zwei Pfl. Docum.* 209.—Afr. austr.
obovatum ×, Sweet, Geran. t. 367.
Obrienianum ×, Sweet, l. c. Suppl. t. 11.
obscurum ×, Sweet, Geran. t. 89.
obsoletum ×, Tratt. Pelarg. t. 172.
obtusidentatum ×, Sweet, Geran. Suppl. t. 92.
obtusifolium ×, Sweet, Geran. t. 25.—Afr. austr.
obtusilobum, *Willd. Enum. Hort. Berol.* 11.—Afr. austr.
obtusum, *Hoffmgg. Verz. Pfl.* 93.—Afr. austr.?
ochroleucum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 263.—Afr. austr.
odontophyllum ×, Tratt. Pelarg. t. 55.
odoratissimum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 419.—Afr. trop.
odoratum, *Hoffmgg. Verz. Pfl.* 93.—Afr. austr.?
odoriferum, Sweet, Hort. Brit. ed. I. 81; G. Don, Gen. Syst. i. 739 = odoratissimum.
odorum, Salisb. Prod. 312 = odoratissimum.
oenanthifolium ×, Sweet, Geran. Suppl. t. 13.
Oenotherae, *Jacq. Ic. Pl. Rar.* iii. 9. t. 525.—Afr. trop.
oleatum, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 52 = egregium.
olidum, *Hoffmgg. Verz. Pfl.* 93.—Afr. austr.
optabile ×, Sweet, Geran. t. 62.
optimum, *Hoffmgg. Verz. Pfl.* 93.—Afr. austr.
opulifolium ×, Sweet, Geran. t. 53.
opulinum, Hoffmgg. Verz. Pfl. 94 = praec.
orbiculatum ×, Sweet, Hort. Brit. ed. I. 78.
orchideum ×, Tratt. Pelarg. t. 75.
ornatum ×, Sweet, Geran. t. 39.

PELARGONIUM:—
ornithopifolium, *Steud. Nom.* ed. II. ii. 288.—Afr. austr.
orobifolium, Steud. l. c. (Quid?).—Afr. austr.?
orphanum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 68 = *mixtum*.
Otites, E. Mey. in Drège, Zwei Pfl. Docum. 209 = longifolium.
ovale, *L'Hérit. Geran.* t. 28.—Afr. austr.
ovalifolium, *DC. Prod.* i. 649.—Afr. austr.
ovalifolium, Harv. in Harv. & Sond. Fl. Cap. i. 262 = Meyeri.
ovifolium, Salisb. Prod. 316 = ovale.
oxalidifolium, *Pers. Syn.* ii. 227.—Afr. austr.
oxaloides, *Willd. Sp. Pl.* iii. 642.—Afr. austr.
oxoniense, Van Eeden, ex Hoffmgg. Verz. Pfl. 89 = malvaefolium.
oxyaloides, Ait. ex Steud. Nom. ed. II. ii. 288 = malvaefolium.
oxyphyllum, DC. Prod. i. 667 = laevigatum.
Palkii ×, Sweet, Geran. t. 224.
pallens, Sweet, l. c. t. 148 = stipulaceum.
pallidulum ×, Sweet, l. c. Suppl. t. 97.
pallidum, *Willd. Enum. Hort. Berol.* 710.—Hab.?
palmatum, *Hort. Par. ex Breit. Hort. Breit.* 337.—Afr. austr.
palmatum, Van Eeden, ex Hoffmgg. Verz. Pfl. 91 = illustre.
panduraeforme, Eckl. & Zeyh. Enum. 82 = quercifolium.
paniculatum, *Hornem. Hort. Hafn. Suppl.* 76.—Afr. austr.
paniculatum, Jacq. Hort. Schoenb. ii. 6 = crithmifolium.
pannifolium ×, Sweet Geran. t. 9.
papaveriflorum, Steud. Nom. ed. II. ii. 288 = triste.
papilionaceum, *Ait. Hort. Kew.* ed. I. ii. 423.—Afr. austr.
papyraceum ×, Sweet, Geran. Suppl. t. 22.
paradoxum ×, Sweet, Hort. Brit. ed. I. 77.
paradoxum ×, Tratt. Pelarg. t. 21.
parameson ×, Tratt. l. c. t. 233.
Parmentieri ×, Colv. ex Sweet, Hort. Brit. ed. I. 82.
parnassioides, *L'Hérit. Geran. Ined.* 1; *ex DC. Prod.* i. 650.—Afr. austr.
particeps ×, Sweet, Geran. t. 49.
parviflorum, *Wendl. Coll.* ii. 69. t. 66.—Afr. austr.
parvulum, *DC. Prod.* i. 660.—Afr. austr.
parvulum, Harv. in Harv. & Sond. Fl. Cap. i. 291 = fumarioides.
patens ×, Sweet, Geran. t. 125.
patentissimum, *Wendl. Coll.* ii. 75. t. 69.—Afr. austr.
paternum ×, Tratt. Pelarg. t. 65.
patulum, *Jacq. Ic. Pl. Rar.* iii. t. 541; *Coll.* iv. 187.—Afr. austr.
patulum, Spin, ex DC. Prod. i. 667 = albiflorum.
paucidentatum ×, Sweet, Geran. t. 186.
pavoninum, Hoffmgg. Verz. Pfl. 94 = anodaefolium.
pavoninum ×, Sweet, Geran. t. 40.
pectinifolium, Sweet, l. c. t. 66 = crispum.
pedicellatum, Sweet, l. c. t. 250 = pulverulentum.
pedunculare, Dehnh. Cat. Pl. Hort. Camald. ed. II. 22 = seq.?
pedunculatum ×, Sweet, Hort. Brit. ed. I. 470; et Geran. t. 346.
Peelii ×, Forbes, Hort. Woburn. 154.
pellitum ×, Tratt. Pelarg. t. 97.
peltatum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 427.—Afr. austr.
penasticum. Steud. Nom. ed. II. ii. 294, sphalm. = *pentasticum*.
pendulum, *G. Don, Gen. Syst.* i. 732.—Afr. austr.
penicillatum, Willd. Hort. Berol. t. 37 = betulinum.
pennaeforme, *Harv. in Harv. & Sond. Fl. Cap.* i. 267, in textu.—Afr. austr.
penniforme, *Pers. Syn.* ii. 227.—Afr. austr.
pentastictum ×, Sweet, Geran. Suppl. t. 25.
peramplum ×, Sweet, l. c. t. 61.
Percyanum ×, Forbes, Hort. Woburn. 154.
perigraphon ×, Tratt. Pelarg. t. 237.
Perryanum ×, Colv. ex Sweet, Hort. Brit. ed. I. 81.
petroselinifolium, *G. Don, Gen. Syst.* i. 728.—Afr. austr.
peucedanifolium, Steud. Nom. ed. II. ii. 288 = anethifolium.
Peytoniae ×, Sweet, Geran. Suppl. t. 46.
phaedimon ×, Tratt. Pelarg. t. 252.
Phellandrium, E. Mey. in Drège, Zwei Pfl. Docum. 209 = senecioides.
phellodes, *Hoffmgg. Verz. Pfl.* 94.—Afr. austr.
phoeniceum ×, Sweet, Geran. t. 207.

PELARGONIUM:—
phymatanthoides, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 172.—Afr. austr.
pictum, Pers. Syn. ii. 227 = pulchellum.
picturatum ×, Sweet, Geran. Suppl. t. 34.
Pignatellii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.
piliferum, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 69.—Afr. austr.
piligerum, Hoffmgg. Verz. Pfl. 94 = praec.
pilipes, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 69.—Afr. austr.
pilosellaeforme, Steud. Nom. ed. II. i. 771, in syn. = seq.
pilosellaefolium, *Steud. l. c.* ii. 288.—Afr. austr.
pilosum, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 69 = piliferum.
pilosum, *Pers. Syn.* ii. 227.—Afr. austr.
pingue, *DC. Prod.* i. 680.—Afr. austr.
pinguifolium, Sweet, Geran. t. 52 = lateripes.
pinnatum, *L'Hérit. Geran.* t. 8.—Afr. austr.
pinnatum, Willd. Sp. Pl. iii. 647 = astragalifolium.
piperatum, *Spin, Cat. Hort. Sebast.* (1818) 32.—Afr. austr.?
piperitum, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 69 = praec.?
placidum, Hoffmgg. Verz. Pfl. 94 = *amoenum*.
planiflorum, Colv. ex Sweet, Hort. Brit. ed. I. 79 = seq.
planifolium ×, Sweet, Geran. t. 219.
platanifolium, *Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 69.—Afr. austr.
platanifolium ×, Sweet, Geran. t. 326.
platypetalum, *Sweet, l. c.* t. 116.—Afr. austr.
platyphyllum, Hoffmgg. Verz. Pfl. 94 = lappaceum.
plectophyllum ×, Sweet, Geran. Suppl. t. 41.
plicatum, *DC. Prod.* i. 680.—Afr. austr.
plicatum, Hoffmgg. Verz. Pfl. 94 = acerifolium.
poculifolium ×, Sweet, Geran. Suppl. t. 31.
podophylloides, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.
politum ×, Sweet, Geran. Suppl. t. 87.
polyanthon ×, Sweet, Hort. Brit. ed. I. 84.
polyanthum, Krauss, in Flora, xxvii. (1844) 298 = ardens.
polycephalum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = ferulaceum.
polygrammon ×, Tratt. Pelarg. t. 235.
polymorphum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = aconitiphyllum.
polyodon ×, Tratt. Pelarg. t. 185.
polyphyllum ×, Tratt. l. c. t. 180.
polytrichum ×, Sweet, Geran. t. 274.
populifolium, Eckl. & Zeyh. Enum. 81 = ribifolium.
porphyreon ×, Sweet, Geran. Suppl. t. 89.
Porphyria, *Hoffmgg. Verz. Pfl.* 94.—Afr. austr.?
Potteri ×, Sweet, Geran. t. 147.
praeclarum ×, Sweet, Geran. Suppl. t. 67.
Prestonii ×, Forbes, Hort. Woburn. 154.
primulaeforme, Steud. Nom. ed. II. ii. 289 = pulverulentum.
primulinum, Sweet, Geran. sub t. 14, in textu = crassicaule.
Princeanum ×, Sweet, Geran. t. 386.
principissae, *Sweet, l. c.* t. 139.
procumbens, DC. Prod. i. 654 = australe.
procumbens, E. Mey. in Drège, Zwei Pfl. Docum. 209 = parvulum.
procumbens, *Pers. Syn.* ii. 229.—Afr. austr.?
proliferum, Steud. Nom. ed. II. ii. 289 = barbatum.
prophanon ×, Tratt. Pelarg. t. 208.
propinquum ×, Colv. ex Sweet, Hort. Brit. ed. I. 81.
propugnatum, Hoffmgg. Verz. Pfl. 94 = fulminans.
pruinosum, *Hoffmgg. l. c.*—Afr. austr.
Pseudociconium ×, Tratt. Pelarg. t. 188.
psilophyllum ×, Sweet, Geran. t. 356.
pubescens, *Andr. ex Link, Enum. Hort. Berol.* ii. 195.—Afr. austr.
pubescens, Hort. ex Tratt. Pelarg. sub t. 13 = *melalophum*.
pubescens, *Spin, Cat. Hort. Sebast.* (1818) 32.—Afr. austr.?
pudicum ×, Tratt. Pelarg. t. 43.
pulchellum, *Sims, Bot. Mag.* t. 524.—Afr. austr.
pulchellum, Willd. Enum. Hort. Berol. Suppl. 47 = Willdenowii.
pulcherrimum ×, Sweet, Geran. t. 134.
pulchrum, Hoffmgg. Verz. Pfl. 95 = formosissimum.

PELARGONIUM:—
pulchrum × , Sweet, Geran. t. 107.
pullaceum × , Sweet, Geran. Suppl. t. 76.
pulverulentum, *Colv. ex Sweet, Geran.* t. 218.—Afr. austr.
pumilum, Willd. Enum. Hort. Berol. 704 = scandens.
punctatum, *Willd. Sp. Pl.* iii. 645.—Afr. austr.
puniceum × , Colv. ex Sweet, Hort. Brit. ed. I. 77.
puniceum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 48.—Afr. austr.
purpurascens, Pers. Syn. ii. 227 = longifolium.
purpurascens, Van Eeden, ex Hoffmgg. Verz. Pfl. 96 = spectabile.
purpureum, *Hoffmgg. l. c.* 95.—Afr. austr.
purpureum, Hort. ex Steud. Nom. ed. II. ii. 289 = longifolium.
purum × , Sweet, Geran t. 334.
pusillum × , Colv. ex Sweet, Hort. Brit. ed. I. 77.
pustulosum, Sweet, Geran. t. 11 = crispum.
pygmaeum × , Colv. ex Sweet, Hort. Brit. ed. I. 77.
pyrethrifolium × , Sweet, Geran. t. 153; Hoffmgg. Verz. Pfl. 95.
quadriflorum × , Sweet, l. c. t. 321.
Queenii × , Forbes, Hort. Woburn. 154.
querceolatum, *Hort. ex Hoffmgg. Verz. Pfl.* 95.—Afr. austr. ?
quercifolium, Baum. ex Hoffmgg. Verz. Pfl. 97, in syn. = ustulatum.
quercifolium, Hort. Cels, ex Hoffmgg. l. c. 94, in syn. = phellodes.
quercifolium, Hort. ex Hoffmgg. l. c. 95, in syn. = quercinum.
quercifolium, *L'Hérit. Geran.* t. 14.—Afr. austr.
quercinum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.
quinatum, *Sims, Bot. Mag.* t. 547.—Afr. austr.
quinquelobatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 118.—Abyss.
quinquelobum, *Colla, Hort. Ripul.* 105. t. 33.—Afr. austr.
quinquevulnerum, *Willd. Enum. Hort. Berol.* 703.—Afr. austr.
radiatum, *Pers. Syn.* ii. 226.—Afr. austr.
radiatum, Steud. Nom. ed. II. ii. 289 = seq.
radicatum, *Vent. Jard. Malm.* n. 65.—Afr. austr.
Radula, *L'Hérit. Geran.* t. 16.—Afr. austr.
radulaefolium, *Steud. Nom.* ed. II. ii. 289.—Afr. austr.
raduloides, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 70 = P. Radula.
Rainerianum × , Tratt. Pelarg. t. 201.
ramigerum × , Sweet, Hort. Brit. ed. I. 471; et Geran. t. 352.
ramosissimum, *Willd. Sp. Pl.* iii. 688.—Afr. austr.
ramulosum × , Sweet, Geran. t. 177.
ranunculifolium, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 38.—Afr. austr.
ranunculifolium, Steud. Nom. ed. II. ii. 289 = alchemilloides.
ranunculophyllum, *Baker, in Saund. Refug. Bot.* iii. t. 150.—Afr. austr.
ranunculophyllum, Harv. in Harv. & Sond. Fl. Cap. i. 296 = alchemilloides.
rapaceum, *Jacq. Ic. Pl. Rar.* iii. t. 510.—Afr. austr.
rapaefolium × , Sweet, Hort. Brit. ed. I. 78.
raphanosmon × , Tratt. Pelarg. t. 212.
rarum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.
Razoumovianum, *Spreng. Fl. Hal. Mant.* 46.—Afr. austr. ?
recurvatum, *Sweet, Geran.* t. 223.—Afr. austr.
recurvifolium × , Sweet, Hort. Brit. ed. I. 471; et Geran. t. 343.
reflexum, E. Mey. ex Harv. & Sond. Fl. Cap. i. 264 = ochroleucum.
reflexum, *Pers. Syn.* ii. 227.—Afr. austr.
reformatum × , Tratt. Pelarg. t. 138.
regale × , Hoffmgg. Verz. Pfl. 95 = *villosum*.
regium × , Sweet, Geran. t. 368.
reniforme, *Curt. Bot. Mag.* t. 493.—Afr. austr.
reniforme, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 46 = Willdenowii.
reniforme, *Spreng. Fl. Hal. Mant.* 46.—Afr. austr.
repandum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.
reticulatum, *DC. Prod.* i. 649; *G. Don, Gen. Syst.* i. 726.—Afr. austr.
retinervium, *DC. Prod.* i. 659.—Afr. austr.
resurgens × , Tratt. Pelarg. t. 158.
retirugum × , Tratt. l. c. t. 82,
retusum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.

PELARGONIUM:—
retusum, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 58 = spurium.
retusum, Wendl. ex Hoffmgg. Verz. Pfl. 95, in syn. = reniforme.
revolutiflorum × , Tratt. Pelarg. t. 166.
revolutum, *Pers. Syn.* ii. 226.—Afr. austr.
Rheinwardtianum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.
Rhodiola × , Tratt. Pelarg. t. 161.
rhodolentum, Sweet, Hort. Brit. ed. I. 84; et Geran. t. 291.
rhodopetalon × , Sweet, Geran. Suppl. t. 14.
rhoeadifolium, *Steud. Nom.* ed. II. ii. 289.—Afr. austr.
ribifolium, Baumg. ex Hoffmgg. Verz. Pfl. 87, in syn. = aurantiodorum.
ribifolium, DC. Prod. i. 658, in syn. = zonale.
ribifolium, *Jacq. Ic. Pl. Rar.* iii. t. 538; *Coll.* v. 141.—Afr. austr.
ribiodorum, Steud. Nom. ed. II. ii. 295 = aurantiodorum.
Richianum × , Sweet, Geran. t. 370.
rigescens × , Sweet, l. c. t. 112.
rigidum, Wendl. ex Hoffmgg. Verz. Pfl. 96, in syn. = sidaefolium.
rigidum, Willd. Sp. Pl. iii. 681 = crispum.
ringens × , Sweet, Geran. t. 256.
Robertianum, *Hoffmgg. Verz. Pfl.* 95.—Afr. austr.
Robinsoni × , Sweet, Geran. t. 150.
robustum × , Colv. ex Sweet, Hort. Brit. ed. I. 79.
Rochlederianum × , Tratt. Pelarg. t. 152.
Rodneyanum, *Lindl. in Mitch. Three Exped.* ii. 144.—Austral.
Rollisoni × , Sweet, Geran. t. 371.
Rosenthalianum × , Tratt. Pelarg. t. 153.
Rosetta, Hort. ex DC. Prod. i. 675 = bellulum.
roseum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 161.—Afr. austr.
roseum, Ehrh. Beitr. vii. 161 = hybridum.
roseum, Willd. Sp. Pl. iii. 679 = P. Radula.
rosodorum, Hoffmgg. Verz. Pfl. 95 = P. Radula.
Rosula, Hoffmgg. l. c. = zonale.
rotundilobum × , Sweet, Geran. t. 252.
Rowanium × , Hoffmgg. Verz. Pfl. 95 = *Daveyanum*.
rubefactum × , Tratt. Pelarg. t. 231.
rubens, *Wendl. Coll.* ii. 45. t. 54.—Afr. austr.
rubescens × , Jenkins. ex Sweet, Geran. t. 30.
rubicundum × , Sweet, Hort. Brit. ed. I. 78.
rubidum, *Sweet, l. c.* 81.—Afr. austr.
rubrocinctum, Link, Enum. Hort. Berol. ii. 191 = cordatum.
rubromaculatum, *Andr. Bot. Rep.* t. 168.—Afr. austr.
rubromaculatum, Poir. Encyc. Suppl. ii. 757 = pulchellum.
Rudolphianum × , Tratt. Pelarg. t. 202.
rugosum, *Colv. ex Sweet, Hort. Brit.* ed. I. 79; *G. Don, Gen. Syst.* i. 740.—Afr. austr.
rumicifolium, *Sweet, Geran.* t. 318.—Afr. austr.
Russelianum × , Sweet, l. c. t. 385.
rutaceum × , Sweet, Hort. Brit. ed. I. 84; Geran. t. 279.
rutaefolium, *Baker, in Saund. Refug. Bot.* t. 27.—Afr. austr.
rutilum × , Colv. ex Sweet, Hort. Brit. ed. I. 82.
Sabini, Haw. ex Sweet, l. c. 81.
saepeflorens × , Sweet, Geran. t. 58.
Salamonce, Hort. ex Dehnh. Cat. Pl. Hort. Camald. ed. II. 22 = hardenbergense.
salebrosum × , Sweet, l. c. t. 309.
sanguineum, *Wendl. Coll.* ii. 43. t. 53; *Sweet, Geran.* t. 76.—Afr. austr.
saniculaefolium, *Willd. Sp. Pl.* iii. 673.—Afr. austr.
saturatum × , Sweet, Geran. Suppl. t. 37.
Saundersii × , Sweet, l. c. t. 205.
Savii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.
Savonitianum × , Tratt. Pelarg. t. 246.
scabridum, *Colv. ex Sweet, Hort. Brit.* ed. I. 78.—Afr. austr.
scabrum, Hort. ex Hoffmgg. Verz. Pfl. 86, in syn. = alutaceum.
scabrum, *L'Hérit. Geran.* t. 31.—Afr. austr.
scandens, *Ehrh. Beitr.* vii. 161; *Willd. Sp. Pl.* iii. 666.—Afr. austr.
scapifolium, L'Hérit. ex DC. Prod. i. 649 = reticulatum.
scaposum, DC. Prod. i. 656 = ovale.

PELARGONIUM:—
Scarboroviae × , Sweet, Geran. t. 117.
scariosum, *Wendl. ex Hoffmgg. Verz. Pfl.* 96.—Afr. austr.
sceleratum × , Colv. ex Sweet, Hort. Brit. ed. I. 82.
schizopetalum, *Sweet, Geran.* t. 232.—Afr. austr.
schizophyllum × , Sweet, l. c. t. 289.
Schottii, *Hook. f. Bot. Mag.* t. 5777.—Hab. ?
Schraderi, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 22.—Afr. austr.
Schreibersianum × , Tratt. Pelarg. t. 126.
Schulzianum × , Tratt. l. c. t. 229.
Schwarzenbergianum × , Tratt. l. c. t. 1.
scintillans × , Sweet, Geran. t. 28.
scintillare, Steud. Nom. ed. II. ii. 289 = fulgidum ?
scitulum × , Sweet, Geran. t. 390.
Sckellii, Hort. ex Steud. Nom. ed. II. ii. 295 = Skelii.
Scottii × , Sweet, Geran. t. 264.
scutatum, DC. Prod. i. 666 = peltatum.
scutatum, *Sweet, Geran.* t. 95.—Afr. austr.
selectum × , Sweet, Geran. t. 190.
selenifolium × , DC. Prod. i. 652.
Selinum, Pers. Syn. ii. 228 = rapaceum.
semicirculare × , Tratt. Pelarg. t. 81.
semitrilobum, Jacq. Hort. Schoenb. ii. t. 130 = adulterinum.
senecioides, *L'Hérit. Geran.* t. 11.—Afr. austr.
sericeum, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 282.—Afr. austr.
serotinum, Colv. ex DC. Prod. i. 673 = Watsoni.
Serpesianum × , Tratt. Pelarg. t. 244.
Serra, *Hoffmgg. Verz. Pfl.* 96.—Hab. ?
serratifolium × , Sweet, Geran. t. 221.
serratum, *DC. Prod.* i. 681.—Afr. austr.
setiferum, *DC. l. c.* i. 680.—Afr. austr.
setigerum, *Steud. Nom.* ed. II. ii. 290.—Afr. austr.
setosum, *DC. Prod.* i. 652.—Afr. austr.
setosum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = barbatum.
setulosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 422.—Afr. austr.
severum × , Tratt. Pelarg. t. 174.
Seymouriae × , Sweet, Geran. t. 37.
sibthorpiaefolium, *Harv. in Harv. & Sond. Fl. Cap.* i. 301.—Afr. austr.
siccifolium, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 70 = zonale.
sidaefolium, *Willd. Enum. Hort. Berol.* 706.—Afr. austr.
sidoides, *DC. Prod.* i. 680.—Afr. austr.
signatum, Sweet, Geran. t. 265.
simplex × , Tratt. Pelarg. t. 74.
sinapifolium × , Sweet, Hort. Brit. ed. I. 78.
sisonifolium, *Baker, in Saund. Refug. Bot.* t. 28.—Afr. austr.
sisymbriifolium, *Steud. Nom.* ed. II. ii. 295.—Afr. austr.
Skelii, *Hoffmgg. Verz. Pfl.* 96.—Afr. austr.
Smithianum, *Steud. Nom.* ed. II. ii. 295.—Afr. austr. ?
Smithii × , Sweet, Geran. t. 110.
smyrniifolium × , Tratt. Pelarg. t. 64.
solubile × , Sweet, Geran. t. 24.
somalense, *Franch. in Revoil, Çomali*, 19.—Afr. trop.
Sophiae × , Tratt. Pelarg. t. 199.
sororium × , Willd. Enum. Hort. Berol. Suppl. 48.
Southcotianum × , Sweet, Hort. Brit. ed. I. 84; et Geran. t. 348.
spathulatum, *Pers. Syn.* ii. 226.—Afr. austr.
spatulatum, G. Don, Gen. Syst. i. 729 = praec.
speciosum, Van Eeden, ex Hoffmgg. Verz. Pfl. 95, in syn. = rarum.
speciosum, *Willd. Enum. Hort. Berol.* 706.—Afr. austr.
spectabile, Hort. ex Link, l. c. ii. 192 = sidaefolium.
spectabile, *Sweet, Geran.* t. 136.—Afr. austr.
Speculum × , Sweet, l. c. t. 52.
Speculum × , Tratt. Pelarg. t. 192.
sphaerocephalum × , Sweet, Geran. t. 317.
sphondilifolium, Steud. Nom. ed. II. ii. 290 = lobatum.
sphondylifolium × , Sweet, Geran. t. 246.
sphondyliifolium, Salisb. Prod. 312 = lobatum.
Spinii, Colla, Hort. Ripul. 105. t. 7 = quercifolium.
spinosum, *Willd. Sp. Pl.* iii. 681.—Afr. austr.
spirans, *Hoffmgg. Verz. Pfl.* 96.—Afr. austr.
splendens, Willd. Hort. Berol. ii. 76 = ignescens.
splendidum × , Sweet, Hort. Brit. ed. I. 78.

PELARGONIUM:—
splendidum ×, Tratt. Pelarg. t. 216.
spondylifolium, Harv. in Harv. & Sond. Fl. Cap. i. 272 sphalm. = lobatum.
spurium, *Willd. Enum. Hort. Berol.* 709.—Afr. austr.?
Stamfordianeae ×, Forbes, Hort. Woburn. 155.
staphysagrioides, *Sweet, Geran. Suppl.* t. 98.—Afr. austr.
Stapletoni ×, Sweet, l. c. t. 212.
staticaefolium, *Steud. Nom.* ed. II. ii. 290.—Afr. austr.
stenanthum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 419 = australe.
stenopetalum, Ehrh. Beitr. vii. 161 = zonale.
Stephanianum ×, Tratt. Pelarg. t. 241.
Stewartii ×, Sweet, Geran. t. 353.
Stiftii ×, Tratt. Pelarg. t. 224.
stipulaceum, *Willd. Sp. Pl.* iii. 655.—Afr. austr.
striatum ×, Sweet, Geran. t. 1.
strictum, Dietr. ex Steud. Nom. ed. II. ii. 290 = crispum.
strigosum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.?
strigosum, *Steud. Nom.* ed. II. ii. 290.—Afr. austr.
suaveolens, *Desf. Arb.* ii. 465; *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 70.—Afr. austr.
subuniflorum ×, Tratt. Pelarg. t. 160.
succulentum ×, Sweet, Geran. Suppl. t. 69.
suffultum ×, Tratt. Pelarg. t. 207.
suffusum ×, Sweet, Hort. Brit. ed. I. 471; et Geran. l. c. t. 47.
sulphureum, *Steud. Nom.* ed. II. ii. 295.—Afr. austr.
superbum, F. G. Dietr. Aesth. Pfl. 44; ex ej. Vollst. Lexik. Gaertn. Nachtr. vi. 43 = Willdenowii.
Sweetianum, *DC. Prod.* i. 653.—Afr. austr.
Sweetianum ×, Sm. ex Sweet, Hort. Brit. ed. II. 588 = praec.?
Synnotii, *G. Don, Gen. Syst.* i. 731.—Afr. austr.
tabulare, *L'Hérit. Geran.* t. 9.—Afr. austr.
tanacetifolium ×, Sweet, l. c. 336.
Tankervillei, Hort. ex Hoffmgg. Verz. Pfl. 96 = glaucum.
tectum, *DC. Prod.* i. 680.—Afr. austr.
tenellum, *Loud. Hort. Brit.* 272; *G. Don, Gen. Syst.* i. 729.—Afr. austr.
Tenorii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.
tenuiflorum, *Schweigg. Enum. Pl. Regiom.* 53.—Afr. austr.?
tenuifolium, *L'Hérit. Geran.* t. 12.—Afr. austr.
tenuilobum, Steud. Nom. ed. II. ii. 290 = patulum.
terebintaceum, Harv. & Sond. Fl. Cap. i. 306, in syn. = graveolens.
ternatifidum, *Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 70.—Afr. austr.
ternatum, *Jacq. Ic. Pl. Rar.* iii. 544; *Coll. Suppl.* 135.—Afr. austr.
terpnon ×, Tratt. Pelarg. t. 171.
testaceum, E. Mey. in Drège, Zwei Pfl. Docum. 209 = pulverulentum.
tetragonum, *L'Hérit. Geran.* t. 23.—Afr. austr.
thamnoides, *Hoffmgg. Verz. Pfl.* 96.—Afr. austr.?
theianthum, Steud. Nom. ed. II. ii. 290 = ochroleucum.
Theresiae ×, Tratt. Pelarg. t. 203.
Thibautii, *Regel, Gartenfl.* (1866) 356.—Hab.?
Thynneae ×, Sweet, Geran. t. 74.
Tibbitsianum ×, Sweet, Geran. t. 158.
tinctum, *Sweet, Geran. Suppl.* t. 29.
tinctum ×, Tratt. Pelarg. t. 142.
Tinei, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.
tomentellum, *DC. Prod.* i. 680.—Afr. austr.
tomentosum, Eckl. & Zeyh. Enum. 613 = papilionaceum.
tomentosum, *Jacq. Ic. Pl. Rar.* 10. t. 537; *Coll. Suppl.* 140.—Afr. austr.
tomentosum, L'Hérit. ex DC. Prod. i. 656 = ovale.
Tormanni, F. G. Dietr. Beschr. Gaert. ed. III. 253 = Barringtonii.
torrefactum ×, Sweet, Geran. t. 243.
tortuosum ×, Sweet, Geran. Suppl. t. 43.
tottum, *DC. Prod.* i. 681.—Afr. austr.
trachyphyllum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
tragacanthoides, Burch. Trav. ii. 98 = ramosissimum.
translucens ×, Sweet, Geran. t. 400.
transmarinum, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 32 = Hoffmanseggii.
Trattinickiae ×, Tratt. Pelarg. t. 245.
triangulare, *Steud. Nom.* ed. II. ii. 290.—Afr. austr.

PELARGONIUM:—
trichostemum, Jacq. Ic. Pl. Rar. iii. t. 524 = ovale.
tricolor, Curt. Bot. Mag. t. 240 = violareum.
tricuspidatum, Hort. ex Steud. Nom. ed. I. 602 = acerifolium.
tricuspidatum, *L'Hérit. Geran.* t. 30.—Afr. austr.
trifidum, Jacq. Hort. Schoenb. ii. 5 = tripartitum.
trifidum, *Willd. Sp. Pl.* iii. 646.—Afr. austr.
trifoliatum, *Harv. in Harv. & Sond. Fl. Cap.* i. 271.—Afr. austr.
trifoliatum, Steud. Nom. ed. II. ii. 290 = *astragalifolium*.
trifoliatum, Sweet, Geran. t. 294 = laevigatum.
trifurcatum ×, Colv. ex Sweet, Hort. Brit. ed. I. 81.
trilingue, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 58.—Afr. austr.?
trilobatum, Eckl. & Zeyh. Enum. 81 = ribifolium.
trilobatum, Schrad. Hort. Gotting. i. 8. t. 2 = semitrilobum.
trilobatum, Van Eeden, ex Hoffmgg. Verz. Pfl. 92, in syn. = lobatum.
trilobum, *DC. Prod.* i. 680.—Afr. austr.
trilobum, Van Eeden, ex Hoffmgg. Verz. Pfl. 95, in syn. = Rheinwardtianum.
trinum ×, Tratt. Pelarg. t. 61.
tripartitum, Hort. ex Hoffmgg. Verz. Pfl. 92 = laeve.
tripartitum, *Willd. Sp. Pl.* iii. 683.—Afr. austr.
triphyllum, *Jacq. Ic. Pl. Rar.* iii. 515; *Coll.* iv. 199.—Afr. austr.
trisectum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
triste, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 418.—Afr. austr.
triternatum ×, Tratt. Pelarg. t. 45.
triumphans ×, Colv. ex Sweet, Hort. Brit. ed. I. 82.
tuberculatum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.
tuberosum, Hort. ex Steud. Nom. ed. II. ii. 290 = triste.
tussilaginifolium ×, Tratt. Pelarg. t. 14.
tyrianthinum ×, Sweet, Geran. t. 183.
uitenhagense, Steud. Nom. ed. II. ii. 290 = schizopetalum.
umbellatum ×, Breit. Hort. Breit. 340.
umbellatum, *Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 48.—Afr. austr.?
undatum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
undulaeflorum ×, Sweet, Geran. t. 329.
undulatipetalum, Pasq. Cat. Orto Bot. Nap. 76 = seq.
undulatum, Harv. in Harv. & Sond. Fl. Cap. i. 262, in syn. = longifolium.
undulatum, [*Soland. in*] *Ait. Hort. Kew.* ed. II. iv. 160.—Afr. austr.
undulifolium, Hoffmgg. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 70 = undulatum.
unguiculatum ×, Tratt. Pelarg. t. 93.
unicolorum, Willd. Enum. Hort. Berol. Suppl. 48 = Willdenowii.
uniflorum, *Breit. Hort. Breit.* 340; *Link, Enum. Hort. Berol.* ii. 193.—Afr. austr.
urbanum, *Steud. Nom. Bot.* ed. II. ii. 290.—Afr. austr.
urbanum ×, Sweet, Geran. Suppl. t. 50.
urenaefolium, *Steud. Nom.* ed. II. ii. 290.—Afr. austr.
ursinum ×, Sweet, Geran. Suppl. t. 94.
ustulatum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
Valleti ×, Colv. ex Sweet, Hort. Brit. ed. I. 79.
Vandesiae ×, Sweet, Geran. t. 7.
varians, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.?
variegatum, Hort. ex Steud. Nom. ed. II. ii. 291 = marginatum.
variegatum, *Willd. Sp. Pl.* iii. 674.—Afr. austr.
variifolium, *Steud. Nom.* ed. II. ii. 291.—Afr. austr.
variifolium ×, Sweet, Geran. t. 280.
Vaudricourtii, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.
Veitchianum ×, Sweet, Geran. Suppl. t. 81.
velutinum, Hoffmgg. Verz. Pfl. 97 = ovale.
velutinum, *L'Hérit. Geran. Ined.* iv. 82; *ex DC. Prod.* i. 650.—Afr. austr.
verbasciflorum, DC. l. c. 657 = ovale.
verbascifolium, Steud. Nom. ed. II. i. 680, in syn., sphalm. = praec.
veniferum ×, Sweet, Geran. t. 322.
veniflorum ×, Sweet, l. c. t. 258.
venosum ×, Steud. Nom. ed. II. ii. 295.
venulosum ×, Sweet, Hort. Brit ed. I. 77.
venustum, Hoffmgg. Verz. Pfl. 97 = imbricatum.

PELARGONIUM:—
venustum ×, Sweet, Geran. t. 167.
verbenaefolium ×, Sweet, Geran. t. 149 = piperatum.
verecundum ×, Sweet, l. c. t. 316.
veronicaefolium, *Steud. Nom.* ed. II. ii. 291.—Afr. austr.
versicolor ×, Sweet, Geran. t. 78.
vespertinum ×, Sweet, l. c. t. 239.
vestifluum ×, Sweet, l. c. Suppl. t. 27.
vestitum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
viciaefolium, L'Hérit. Geran. Ined. n. 15; ex DC. Prod. i. 653 = pinnatum.
Victorianum ×, Sweet, Geran. Suppl. t. 7.
villiferum, *DC. Prod.* i. 680.—Afr. austr.
villosum, Spin, Cat. Hort. Sebast. (1818) 83 = tomentosum.
villosum ×, Sweet, Geran. t. 100.
violaeflorum, *DC. Prod.* i. 652.—Afr. austr.
violareum, *Jacq. Ic. Pl. Rar.* t. 527.—Afr. austr.
virgatum ×, Tratt. Pelarg. t. 217.
virgineum, Pers. Syn. ii. 226 = longifolium.
viridifolium ×, Sweet, Geran. t. 331.
viscidum, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* vi. 45.—Afr. austr.
viscidum, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
viscosissimum, *Sweet, Geran.* t. 118.—Afr. austr.
viscosum, Harv. & Sond. Fl. Cap. i. 306, in syn. = glutinosum.
vitifolium, *L'Hérit. Geran.* t. 19.—Afr. austr.
volatiflorum ×, Sweet, Hort. Brit. ed. I. 84; et Geran. t. 284.
volubile, *Sweet, ex Drapiez, Herb. Amat. Fl.* i. (1828) t. 43.—Afr. austr.?
Volucris, *Hoffmgg. Verz. Pfl.* 97.—Afr. austr.?
vulpinum ×, Tratt. Pelarg. t. 34.
Wachendorfiae ×, Tratt. l. c. t. 183.
Watsoni, *Link, Enum. Hort. Berol.* ii. 192.—Afr. austr.?
Wedelii, *Hort. ex Hoffmgg. Verz. Pfl.* 97.—Afr. austr.
Wellsianum ×, Sweet, Geran. t. 175.
Willdenowii, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 46 = Barringtonii.
Willdenowii, *Link, Enum. Hort. Berol.* ii. 190.—Afr. austr.
Yarburghii ×, Forbes, Hort. Woburn. 155.
Yeatmanianum ×, Sweet, Geran. Suppl. t. 59.
Wuerthianum ×, Tratt. Pelarg. t. 225.
Youngii ×, Sweet, Geran. t. 131.
Zeyheri, *Harv. in Harv. & Sond. Fl. Cap.* i. 276.—Afr. austr.
Zeyherianum, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Afr. austr.
Zichyanum ×, Tratt. Pelarg. t. 149.
zingiberinum ×, Sweet, Geran. Suppl. t. 42.
zonale, *L'Hérit.* [*ex Soland.*] *in Ait. Hort. Kew.* ed. I. ii. 424.—Afr. austr.

PELEA, A. Gray. in Bot. U. St. Expl. Exped. i. 339 (1854) = **Melicope**, Forst. (Rutac.).
anisata, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 314.—Ins. Sandvic.
auriculaefolia, A. Gray, Bot. U. St. Expl. Exped. i. 343. t. 36.—Ins. Sandvic.
calophylla, Baill. Adansonia, x. (1871–72) 322.—N. Caled.
clusiaefolia, A. Gray, Bot. U. St. Expl. Exped. i. 340. t. 35.—Ins. Sandvic.
Deplanchei, Baill. Adansonia, x. (1871–72) 322.—N. Caled.
hawaiensis, Wawra, in Flora, lvi. (1873) 110.—Ins. Sandvic.
Kaalae, Wawra, l. c. = M. elliptica.
kavaiensis, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 313.—Ins. Sandvic.
lucida, A. Gray, Bot. U. St. Expl. Exped. i. 348. t. 34.—Ins. Sandvic.
madagascarica, Baill. Adansonia, x. (1871–72) 323.—Madag.
myrtoidea, Baill. l. c.—N. Caled.
oblongifolia, A. Gray, Bot. U. St. Expl. Exped. i. 343.—Ins. Sandvic.
rotundifolia, A. Gray, l. c. 344. t. 37.—Ins. Sandvic.
sandwicensis, A. Gray, l. c. 345. t. 37.—Ins. Sandvic.
sapotaefolia, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 312.—Ins. Sandvic.
volcanica, A. Gray, Bot. U. St. Expl. Exped. i. 346. t. 38.—Ins. Sandvic.
Waialealae, Wawra, in Flora, lvi. (1873) 108.—Ins. Sandvic.

PELECINUS, Tourn. ex Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378 = **Biserrula**, Linn. (Legumin.).
Biserrula, Moench, Meth. 169 = B. Pelecinus.
vulgaris, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378 = B. Pelecinus.

PELECYNTHIS, E. Mey. Comm. Pl. Afr. Austr. 13 (1835) = **Rafnia**, Thunb. Legumin.).
axillaris, E. Mey. l. c. 14 = R. axillaris.
corymbosa, E. Mey. l. c. = R. cuneifolia.
dichotoma, E. Mey. l. c.—Afr. austr.
diffusa, E. Mey. l. c. 15 = R. diffusa.
gibba, E. Mey. l. c. 14 = R. dichotoma.
opposita, E. Mey. l. c. = R. opposita.
retroflexa, E. Mey. l. c. 15 = R. retroflexa.
racemosa, Walp. Rep. i. 579 = R. racemosa.
rhomboidea, E. Mey. Comm. Pl. Afr. Austr. 14 = R. cuneifolia.

PELECYPHORA, Ehrenb. in Bot. Zeit. i. (1843) 737. *CACTEAE*, Benth. & Hook. f. i. 848.
aselliformis, *Ehrenb. l. c.*—Mexic.

PELEXIA, [Poit.] Rich. in Mém. Mus. Par. iv. (1818) 59. *ORCHIDEAE*, Benth. & Hook. f. iii. 597.
ADNULA, Rafin. Fl. Tellur. ii. 87 (1836).
COLLEA, Lindl. Bot. Reg. sub t. 760 (1823).
ELTROPLECTRIS, Rafin. Fl. Tellur. ii. 51 (1836).
acianthiformis, *Reichb. f. & Warm. in Otia Bot. Hamb.* 83.—Bras.
adnata, *Spreng. Syst.* iii. 704.—Ind. occ.
bursaria, Lindl. in Ann. Nat. Hist. Ser. I. v. (1840) 116 = Physurus plantaginensis.
corymbosa, *Lindl. Gen. et Sp. Orch.* 482.—Peruv.
cranichoides, *Griseb. Cat. Pl. Cub.* 269.—Cuba.
domingensis, *Lindl. Gen. et Sp. Orch.* 482.—Ins. S. Doming.
falcata, Spreng. Syst. iii. 704 = Cephalanthera falcata.
foliosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 17.—Bras.
Hoffmanni, Reichb. f. Beitr. Orch. Centr. Am. 102 = Pseudocentrum Hoffmanni.
japonica, Spreng. Syst. iii. 704 = Cephalanthera pallens.
laxa, *Lindl. Gen. et Sp. Orch.* 482.—Peruv.
pauciflora, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 17. t. 124.—Bras.
repens, *Poepp. & Endl. l. c.* ii. 17. t. 124. f. a–c.—Peruv.
roseoalba, *Reichb. f. in Bonplandia*, ii. (1854) 11.—N. Granat.
setacea, *Lindl. Gen. et Sp. Orch.* 482.—Bras.
spiranthoides, Lindl. Bot. Reg. t. 985 = adnata.
stenorrhynchoides, *Griseb. Cat. Pl. Cub.* 269.—Cuba.
triloba, *Lindl. Gen. et Sp. Orch.* 483.—Bras.

PELIANTHUS, E. Mey. ex Moq. in DC. Prod. xiii. ii. 246 (1849) = **Hermbstaedtia**, Reichb. (Amarant.).
celosioides, E. Mey. ex Moq. l. c. = H. caffra.

PELIOSANTHES, Andr. Bot. Rep. tt. 605, 634 (1808). *HAEMODORACEAE*, Benth. & Hook. f. iii. 678.
BULBISPERMA, Reinw. ex Blume, Cat. Hort. Buitenz. 59 [1823].
BULBOSPERMUM, Blume, Enum. Pl. Jav. 15 (1828).
TETA, Roxb. Hort. Beng. 24 (1814); Fl. Ind. ii. 165 (1832).
altissima, *Donn, Hort. Cantab.* ed. VI. 89.—Ind. or.
campanulata, Wall. ex Baker, in Journ. Linn. Soc. xvii. (1879) 504 = violacea.
courtallensis, *Wight, Ic.* t. 2051.—Ind. or.
Griffithii, *Baker, in Journ. Linn. Soc.* xvii. (1879) 506.—Reg. Himal.
humilis, *Andr. Bot. Rep.* t. 634.—Penang; Malacca.
humilis, *Baker, in Journ. Linn. Soc.* xvii. (1879) 505.—Reg. Himal.
javanica, *D. Dietr. Syn. Pl.* ii. 1123.—Java.
longifolia, Steud. ex Baker, in Journ. Linn. Soc. xvii. (1879) 503 = neilgherriensis.
macrophylla, *Wall. ex Baker, l. c.* 505.—Reg. Himal.
macrostegia, *Hance, in Journ. Bot.* xxiii. (1885) 328.—China.
neilgherriensis, *Wight, Ic.* t. 2052.—Ind. or.
petiolaris, *Wall. ex Voigt, Hort. Suburb. Calc.* 650, nomen.—Hab. ?
Teta, *Andr. Bot. Rep.* t. 605.—Ind. or.; Burma; Penang.
Teta, Wall. Cat. n. 5083 = violacea, courtallensis.
violacea, *Wall. Cat.* n. 5084.—Reg. Himal.; Burma.

PELIOSTOMUM, E. Mey. ex Benth. in Bot. Reg. sub t. 1882 (1836). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 927.
leucorrhizum, *E. Mey. ex Benth. l. c.*—Afr. austr.
origanoides, *E. Mey. ex Benth. l. c.*—Afr. austr.
scoparium, *E. Mey. ex Benth. l. c.*—Afr. austr.
virgatum, *E. Mey. ex Benth. l. c.*—Afr. austr.
viscosum, *E. Mey. ex Benth. l. c.*—Afr. austr.

PELLA, Gaertn. Fruct. i. 143. t. 28. f. 8 (1788) = **Salvadora**, Garcin.
Ribesioides, Gaertn. l. c. = S. persica.

PELLACALYX, Korth. in Hoev. & De Vriese, Tijdschr. iii. (1836) 20. t. 3. *RHIZOPHOREAE*, Benth. & Hook. f. i. 680.
axillaris, *Korth. l. c.*—Malaya.

PELLEA, André, in Illustr. Hortic. xxvii. (1880) 177 = **Pellionia**, Gaudich. (Urtic.).

PELLETERIA, Poir. in Dict. Sc. Nat. xxxviii. 324 (1825) = seq.

PELLETIERA, A. St. Hil. in Mém. Mus. Par. ix. (1822) 365 = **Asterolinon**, Hoffmgg. & Link (Primul.).
verna, A. St. Hil. l. c. = A. serpyllifolium.

PELLICIERA, Planch. & Triana, in Benth. & Hook. f. Gen. i. 186 (1862). *TERNSTROEMIACEAE*, Benth. & Hook. f. l. c.
Rhizophoreae, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 381.—N. Granat.

PELLINIA, Molina, Sagg. Chile, ed. II. 160 (1810) = **Eucryphia**, Cav. (Rosac.).
cordifolia, Molina, l. c. = E. cordifolia.

PELLIONIA, Gaudich. in Freyc. Voy. Bot. 494. t. 119 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 385.
POLYCHROA, Lour. Fl. Cochinch. 559 (1790).
ambigua, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 285.—Reg. Himal.
australis, *Wedd. in DC. Prod.* xvi. i. 169.—Ind. occ.
brevifolia, *Benth. Fl. Hongk.* 330.—China.
Daveauana, *N. E. Br. in Gard. Chron.* (1880) ii. 262.—Burma; Malaya; Cochinch.
divaricata, *Gaudich. in Freyc. Voy. Bot.* 494.—Ins. Marian.
elatostemoides, *Gaudich. l. c.* t. 119.—Ins. Molucc.
filicoides, *Seem. Fl. Vit.* 239.—Ins. Fiji.
Griffithiana, *Wedd. in DC. Prod.* xvi. i. 165.—Reg. Himal.
Grijsii, *Hance, in Journ. Bot.* vi. (1868) 49.—China.
Helferiana, *Wedd. in DC. Prod.* xvi. i. 170.—Burma.
heteroloba, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 283.—Reg. Himal.
Heyneana, *Wedd. l. c.* 287.—Ind. or.
javanica, *Wedd. l. c.* 288.—Malaya.
procridifolia, *Kurz, in Journ. Bot.* xiii. (1875) 330.—Ins. Nicobar.
pulchra, *N. E. Br. in Gard. Chron.* (1882) ii. 712.—Cochinch.
radicans, *Wedd. in DC. Prod.* xvi. i. 167.—Japon.
scabra, *Benth. Fl. Hongk.* 330.—China.
stenocarpa, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 284.—Reg. Himal.
umbellata, *Wedd. in DC. Prod.* xvi. i. 167.—Japon.
vitiensis, *A. Gray, ex Wedd. l. c.*—Ins. Fiji.

PELONASTES, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 474 = **Myriophyllum**, Linn. (Halorag.).
integrifolia, Hook. f. l. c. 475 = M. integrifolium.
tillaeacea, F. Muell. in Hook. Kew Journ. viii. (1856) 65 = M. integrifolium.
tuberculata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 474 = M. Drummondii.

PELTA, Dulac, Fl. Hautes-Pyr. 43 (1867) = **Zannichellia**, Mich. (Naiad.).
palustris, Dulac, l. c. = Z. palustris.

PELTACTILA, Rafin, New Fl. Am. iv. 28 (1836) = **Daucus**, Tourn. (Umbell.).
aurea, grandiflora, hispida, parviflora, Rafin. l. c. = D. Carota *vel* D. pusillus.

PELTANDRA, Rafin. in Journ. Phys. lxxxix. (1819) 102. *AROIDEAE*, Benth. & Hook. f. iii. 975.
LECONITA, W. Cooper, ex Torr. Compend. 358 [bis] (1826).
RENSSELAERIA, Beck, Bot. U. St. 382 (1833).
alba, *Rafin. New Fl. Am.* i. 88.—Am. bor.
angustifolia, Rafin. l. c. 89 = virginica.
canadensis, Rafin. l. c. 87 = virginica.
hastata, Rafin. l. c. 88 = virginica.
heterophyla, Rafin. l. c. = virginica.
latifolia, Rafin. l. c. 87 = virginica.
sagittaefolia, Rafin. in Journ. Phys. lxxxix. (1819) 102 = Xanthosoma sagittaefolium.
undulata, *Rafin. l. c.*—Am. bor.
undulata, Schott, Melet. i. 19 = virginica.
virginica, *Rafin. in Journ. Phys.* lxxxix. (1819) 102.—Am. bor.
virginica, Schott, Syn. Aroid. 50 = undulata.
Walteri, Rafin. New Fl. Am. i. 88 = virginica.

PELTANDRA, Wight, Ic. v. ii. 24 (1852) = **Phyllanthus**, Linn. (Euphorb.).
flexuosa, Thw. Enum. Pl. Zeyl. 281 = Phyll. Thwaitesianus.
longipes, Wight, Ic. t. 1891 = Phyll. longipes.
parvifolia, Wight, l. c. t. 1892 = Phyll. Thwaitesianus.

PELTANTHERA, Benth. in Benth. & Hook. f. Gen. ii. 797 (1876). *LOGANIACEAE*, Benth. & ok. f. l. c.
floribunda, *Benth. l. c.*—Peruv.

PELTANTHERA, Roth, Nov. Sp. Pl. 132 (1821) = **Vallaris**, Burm. f. (Apocyn.).
solanacea, Roth, l. c. = V. Heynei.

PELTARIA, Burm. ex DC. Prod. ii. 420 (1825) = **Wiborgia**, Thunb. (Legumin.).
capensis, Burm. ex DC. l. c. = W. fusca.

PELTARIA, Jacq. Enum. Hort. Vindob. 260 (1762); Linn. Sp. Pl. ed. II. 910 (1763). *CRUCIFERAE*, Benth. & Hook. f. i. 93.
BOHADSCHIA, Crantz, Stirp. Austr. ed I. fasc. i. 2 (1762); ed. II. 5 (1769).
aleppica, Hort. ex Steud. Nom. ed. I. 603 = Isatis aleppica.
alliacea, *Jacq. Enum. Hort. Vindob.* 260.—Europ. or.
angustifolia, *DC. Syst.* ii. 329.—As. Min.; Persia.
aspera, Grauer, Pl. Min. Cogn. Dec. 6 (Quid ?).—Oriens.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 175.—Armenia.
capensis, Linn. f. Suppl. 296 = Heliophila diffusa.
caramaniensis, Sibth. ex DC. Syst. ii. 285 = Ricotia tenuifolia.
deflexa, DC. l. c. 337 = Texiera glastifolia.
Garcini, Burm. f. Fl. Ind. 139 = Fortuynia Garcini.
glastifolia, DC. Syst. ii. 330 = Texiera glastifolia.
obcordata, Steud. Nom. ed. I. 603 = Alyssum obcordatum.

PELTICALYX, Griff. Notul. ii. 706 (1854). Genus dub. *ANONACEARUM*.—Cf. Benth. & Hook. f. i. 22.
argentea, Griff. l. c.—Ind. or.

PELTIDIUM, Zollikof. in Nat. Anz. Schw. Ges. iv. (1820) 13 = **Chondrilla**, Tourn. (Compos.).
apargioides, Zollikof. l. c. = C. Peltidium.

PELTIMELA, Rafin. Atl. Journ. 199 (1833) = **Glossostigma**, Arn. (1836) (Scrophul.).
cuneata, Rafin. l. c. = G. spathulatum.

PELTISPERMUM, Moq. Clenop. Enum. 102 (1840) = **Anthochlamys**, Fenzl (Chenop.).
Aucheri, Moq. in DC. Prod. xiii. ii. 143 = A. polygaloides.

PELTOBRYON, Klotzsch, ex Miq. Syst. Pip. 369 (1843) = **Piper**, Linn.
attenuatum, Klotzsch, l. c. 371 = Pip. Peltobryon.
callosum, Miq. l. c. 369 = Pip. callosum.
cyrtopodium, Miq. in Mart. Fl. Bras. iv. i. 219 = Pip. cyrtopodon.
dichotomum, Klotzsch, ex Miq. Syst. Pip. 372 = Pip. dichotomum.

PELTOBRYON :—
exserens, Miq. in Mart. Fl. Bras. iv. 1. 31 = Piper exserens.
Guilleminianum, Miq. l. c. 30 = Pip. crassinervium.
Hookeri, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 442 = Pip. munchanum.
longifolium, Miq. Syst. Pip. 372 = Pip. longifolium.
Martianum, Miq. in Linnaea, xx. (1847) 133 = Pip. egense.
Mathewsi, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 441 = Pip. tenuistylum.
phytolaccaefolium, Presl, Epim. Bot. 224 = Pip. phytolaccaefolium.
Poeppigii, Klotzsch, ex Miq. Syst. Pip. 370 = Pip. Poeppigii.
pubescens, Miq. in Linnaea, xx. (1847) 134 = Pip. brasiliense.

PELTODON, Pohl, Pl. Bras. Ic. i. 66. tt. 54–56 (1827). *LABIATAE*, Benth. & Hook. f. ii. 1177.
longipes, *A. St. Hil. ex Benth. Lab. Gen. & Sp.* 63.—Bras.
pusillus, *Pohl, Pl. Bras. Ic.* i. 67. t. 54.—Bras.
radicans, *Pohl, l. c.* 68. t. 55.—Bras.
tomentosus, *Pohl, l. c.* 69. t. 56.—Bras.

PELTOGYNE, Vog. in Linnaea, xi. (1837) 410. *LEGUMINOSAE*, Benth. & Hook. f. i. 582.
confertiflora, *Benth. in Mart. Fl. Bras.* xv. II. 232.—Bras.
densiflora, *Spruce, ex Benth. l. c.*—Bras.
discolor, *Vog. in Linnaea*, xi. (1837) 410.—Bras.
latifolia, *Benth. in Mart. Fl. Bras.* xv. II. 232.—Bras.
paniculata, *Benth. in Hook. Journ. Bot.* ii. (1840) 96.—Am. austr.
parvifolia, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 233.—Bras.
pauciflora, *Benth. l. c.* 234.—Bras.
porphyrocardia, *Griseb. ex Benth. l. c.* 234, in obs.—Am. trop.
pubescens, *Benth. in Hook. Journ. Bot.* ii. (1840) 96.—Guian.
venosa, *Benth. in Mart. Fl. Bras.* xv. II. 233, in obs.—Guian.

PELTOMESA, Rafin. Sylva Tellur. 126 (1838) = **Loranthus**, Linn.
acuminata, Rafin. l. c. = L. acuminatus.

PELTOPHORA, Benth. & Hook. f. Gen. iii. 1129 (1883), sphalm. = PELTOPHORUS, Desv.

PELTOPHORUM, Walp. Rep. i. 811 (1842). *LEGUMINOSAE*, Benth. & Hook. f. i. 565.
adnatum, *Griseb. Fl. Brit. W. Ind.* 206.—Ins. Bahamas.
africanum, *Sond. in Linnaea*, xxiii. (1850) 35.—Afr. trop.
dasyrachis, *Kurz, ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 257.—Malacca.
ferrugineum, *Benth. Fl. Austral.* ii. 279.—Austral.
inerme, Naves, ex Villar, in Blanco, Fl. Filip. ed. III. Nov. App. 69 = ferrugineum.
Linnaei, *Benth. ex Griseb. Fl. Brit. W. Ind.* 206.—Jamaic.
Vogelianum, *Walp. Rep.* i. 811.—Bras.

PELTOPHORUS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = **Rottboellia**, Linn. (Gramin.).
elegans, Beauv. Agrost. 172 = R. Myurus.
Myurus, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = R. Myurus.

PELTOPHYLLUM, Gardn. in Proc. Linn. Soc. i. (1843) 176 ; et in Trans. Linn. Soc. xix. (1845) 157. t. 15 = **Triuris**, Miers.
luteum, Gardn. ll. cc. = T. lutea.

PELTOPSIS, Rafin. in Journ. Phys. lxxxix. (1819) 102 = **Potamogeton**, Tourn. (Naiad.).
perfoliata, Rafin. l. c. 103 = Pot. perfoliatus.

PELTOSPERMUM, Benth. in Hook. Niger Fl. 400 (1849) = **Oldenlandia**, Linn. (Rubiac.).
paniculatum, Benth. l. c. = O. Peltospermum.

PELTOSPERMUM, DC. in Bibl. Univ. Genèv. xvii. (1838) 133 (Rev. Bign. 17) = **Aspidosperma**, Mart. & Zucc. (Bignon.).
latisiliquum, DC. ex A. DC. in Prod. viii. 676 = A. latisiliqua.
orbiculatum, DC. l. c. 191 = Anemopaegma orbiculatum.
Patrisii, DC. l. c. viii. 676 = A. latisiliqua.

PELTOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 223. Gen. *INCERT. SED.* (Cf. Benth. & Hook. f. Gen. i. 217).
parviflora, *Turcz. l. c.* 224.—Bras.

PELTOSTIGMA, Walp. Rep. v. 387 (1846). *RUTACEAE*, Benth. & Hook. f. i. 300.
PACHYSTIGMA, Hook. Ic. Pl. t. 698 (1844).
ptelioides, *Walp. Rep.* v. 387.—Jamaic.

PEMPHIS, Forst. Char. Gen. 67. t. 34 (1776). *LYTHRARIEAE*, Benth. & Hook. f. i. 780.
MACLELLANDIA, Wight, Ic. t. 1996 (1853).
MILLANIA, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 128 ('1852') [1856].
acidula, *Forst. Char. Gen.* 68. t. 34.—As. trop. ; Ins. Pacif.
angustifolia, Roxb. Hort. Beng. [91] ; Fl. Ind. ii. 465 = acidula.
hexandra, Mart. ex Koehne, in Mart. Fl. Bras. xiii. II. 324 = Diplusodon hexander.
setosa, Blanco, Fl. Filip. ed. I. 410 = acidula.
stachydifolia, Mart. ex Koehne, in Mart. Fl. Bras. xiii. II. 332 = Diplusodon villosissimus.

PENAEA, Linn. Hort. Cliff. 37 (1737). *PENAEACEAE*, Benth. & Hook. f. iii. 202.
STYLAPTERUS, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 22, 23. t. 1.
acuta, Thunb. in Hoffmgg. Phytog. Blaetter, i. (1803) 16 = Sarcocolla acuta.
acutifolia, *A. Juss. in Ann. Sc. Nat.* Sér. III. vi. (1846) 22.—Afr. austr.
affinis, *Endl. Gen. Suppl.* iv. II. 73.—Afr. austr.
barbata, *Endl. l. c.*—Afr. austr.
Cneorum, Meerb. Ic. Pl. Sel. 51 ; Lam. Illustr. i. 317 = myrtoides.
Cneorum, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 556 (err. typ. 456) = myrtifolia.
Dregei, *Endl. Gen. Suppl.* iv. II. 73.—Afr. austr.
ericoides, *Endl. l. c.*—Afr. austr.
formosa, *Thunb. Prod. Pl. Cap.* 187.—Afr. austr.
fruticulosa, *Linn. f. Suppl.* 121.—Afr. austr.
fucata, Herb. Boiv. ex A. DC. in DC. Prod. xiv. 484 = mucronata.
fucata, Herb. Lamb. ex A. DC. l. c. 490 (= *Glischrocolla Lessertiana*).—Afr. austr. ?
fucata, Herb. Vit. ex A. DC. l. c. 487 = Sarcocolla imbricata.
fucata, *Linn. Mant.* i. 199.—Afr. austr.
fucata, Zeyh. ex A. DC. in DC. Prod. xiv. 489 = Sarcocolla squamosa.
furcata, Murr. Syst. ed. XIII. 130, sphalm. = fucata.
fuscata, Auct. ex Steud. Nom. ed. I. 603 = fucata.
fuscata, Poir. Encyc. vi. 540 = Sarcocolla imbricata.
imbricata, Eckl. ex A. DC. in DC. Prod. xiv. 484 = mucronata.
imbricata, R. Grah. in Edinb. N. Phil. Journ. (Apr.–Jun. 1827) 177 ; et in Bot. Mag. t. 2809 = Sarcocolla imbricata.
intermedia, *Endl. Gen. Suppl.* iv. II. 73.—Afr. austr.
lateriflora, *Linn. f. Suppl.* 121.—Afr. austr.
Linnaei, A. Juss. ex A. DC. in DC. Prod. xiv. 486 = Sarcocolla fucata.
longiflora, Meerb. Ic. Pl. Sel. t. 51 = Sarcocolla longiflora.
marginata, *Linn. Mant.* i. 199.—Afr. austr.
mucronata, *Linn. Sp. Pl.* 111.—Afr. austr.
myrtifolia, *Endl. Gen. Suppl.* iv. II. 73.—Afr. austr.
myrtilloides, Herb. Berol. ex A. DC. in DC. Prod. xiv. 485 = acutifolia.
myrtilloides, Thunb. Prod. Pl. Cap. 30 = seq.
myrtoides, *Linn. f. Suppl.* 122.—Afr. austr.
nitida, *Lour. Fl. Cochinch.* 72.—Cochinch.
ovata, *Eckl. & Zeyh. ex A. Juss. in Ann. Sc. Nat.* Sér. III. vi. (1846) 22.—Afr. austr.

PENAEA :—
punicea, *Hort. ex Steud. Nom.* ed. II. ii. 297.—Afr. austr.
Sarcocolla, Lam. ex A. DC. in DC. Prod. xiv. 486 = squamosa.
Sarcocolla, Linn. Sp. Pl. 111 ; Berg. Desc. Fl. Cap. 35 = fucata.
scandens, *Lour. Fl. Cochinch.* 73.—Cochinch.
squamosa, Auct. ex A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 25 = Sarcocolla imbricata.
squamosa, *Linn. Sp. Pl.* 112.—Afr. austr.
tetragona, Berg. Desc. Pl. Cap. 36 = fucata.
tomentosa, *Thunb. Prod. Pl. Cap.* 30.—Afr. austr.

PENANTHES, Vell. Fl. Flum. 350 (1825), sphalm. = **Prenanthes**, Vaill. (Compos.).

PENDULINA, Willk. in Linnaea, xxv. (1852) 2 = **Diplotaxis**, DC. (Crucif.).
crassifolia, Willk. l. c. = D. pendula.
Fontanesii, Willk. l. c. 4.—Afr. bor.
hispida, Willk. l. c. = D. pendula.
intricata, Willk. l. c. = D. intricata.
Lagascana, Willk. l. c. 2 = D. Lagascana.
Webbiana, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 868.—Hispan.

PENIANTHUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 124 ; xx. (1867) 171. Gen. dub. *MENISPERMACEARUM.*
longifolius, *Miers, ll. cc.* 124 ; 172.—Afr. trop.

PENICILLANTHEMUM, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 94 = **Hugonia**, Linn. (Lineae).
latifolium, Vieill. l. c. 95.—N. Caled.
neo-caledonicum, Vieill. l. c. 96.—N. Caled.
racemosum, Vieill. l. c. 95.—N. Caled.

PENICILLARIA, Willd. Enum. Hort. Berol. 1036 (1809) = **Pennisetum**, Rich. (Gramin.).
alopecuroides, Sweet, Hort. Brit. ed. I. 440 ; A. Br. Ind. Sem. Hort. Berol. (1855) App. 25 = Pennis. typhoideum.
arabica, A. Br. l. c. 23.—Arab.
chinensis, Nees, ex A. Br. l. c. 26 = Pennis. chinense.
ciliata, Willd. Enum. Hort. Berol. 1037 = Pennis. typhoideum.
compacta, A. Br. Ind. Sem. Hort. Berol. (1855) App. 23.—Nubia.
cordofana, A. Br. l. c. 23.—Afr. trop.
cylindrica, Roem. & Schult. Syst. ii. 498 = Pennis. typhoideum.
deflexa, Anderss. ex A. Br. Ind. Sem. Hort. Berol. (1855) App. 25.—Afr. trop.
Doche, A. Br. l. c. 23, in obs.—Hab. ?
fallax, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 371.—Nubia.
Gymnothrix, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 26.—Afr. trop.
involucrata, Schult. Mant. ii. 283 ; Schlecht. in Linnaea, xxv. (1852) 565 = Pennis. typhoideum.
leucostachys, Klotzsch, ex A. Br. Ind. Sem. Hort. Berol. (1855) App. 25.—Afr. trop.
macrostachya, Klotzsch, l. c. = Pennis. megastachyum.
mossambicensis, Klotzsch, l. c.—Afr. trop.
nigritarum, Schlecht. in Linnaea, xxv. (1852) 561.—Afr. trop.
nubica, A. Br. Ind. Sem. Hort. Berol. (1855) App. 23.—Afr. trop.
Perrottetii, Klotzsch, ex A. Br. l. c. 25.—Afr. trop.
Plukeneti, Link, Hort. Berol. i. 221 = Pennis. typhoideum.
Raddiana, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 373, in obs. = Pennis. typhoideum.
Roxburghii, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 26 = Pennis. typhoideum.
Sieberiana, Schlecht. in Linnaea, xxv. (1852) 565.—Aegypt.
socia, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 24.—Afr. trop. ?
solitaria, Stokes, Bot. Mat. Med. i. 129 = Pennis. typhoideum.
speciosa, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 23.—Afr. trop.
spicata, Willd. Enum. Hort. Berol. 1037 ; Wall. ex Anderss. in Walp. Ann. vi. 966 = Pennis. typhoideum.

PENICILLARIA:—
stenostachya, Klotzsch, ex A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 25.—Afr. trop.
typhoidea, Fig. & Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 371 = Pennisetum typhoideum.
vulpina, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1855) App. 24.—Afr. trop.
Willdenowii, Klotzsch, ex. A. Br. & Bouché, l. c. 25.—Afr. trop.

PENNANTIA, Forst. Char. Gen. 133. t. 67 (1776).
OLACINEAE, Benth. & Hook. f. i. 351.
corymbosa, Endl. Prod. Fl. Norf. 80 = Endlicheri.
corymbosa, *Forst. Char. Gen.* 134.—N. Zel.
Cunninghamii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. ix. (1852) 491.—Austral.
Endlicheri, *Reiss. in Linnaea*, xvi. (1842) 341. t. 13.—Ins. Norfolk.
odorata, Raoul, in Ann. Soc. Sc. Nat. Sér. III. ii. (1844) 123 = corymbosa.

PENNISETUM, Rich. in Pers. Syn. i. 72 (1805).
GRAMINEAE, Benth. & Hook. f. iii. 1105.
AMPHOCHAETA, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 136.
BECKEROPSIS, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xiv. (1854) 365.
CATATHEROPHORA, Steud. in Flora, xii. (1829) 465.
ERIOCHAETA, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xiv. (1854) 374.
GYMNOTRIX, Beauv. Agrost. 59. t. 13. f. 6. (1812).
MACROCHAETA, Steud. in Zoll. Syst. Verz. Ind. Archip. 60 (1854).
PENICILLARIA, Willd. Enum. Hort. Berol. 1036 (1809).
PENTASTACHYA, Hochst. ex Steud. Nom. ed. II. ii. 299 (1841).
SERICURA, Hassk. in Flora, xxv. (1842) II. Beibl. 2.
adoënse, *Steud. Nom.* ed. II. ii. 252.—Abyssin.
alopecuroides, Ham. Prod. Pl. Ind. Occ. 11 = Hamiltonii.
alopecuroides, Spreng. Syst. i. 303 = typhoideum.
alopecuroides, *Steud. in Flora*, xii. (1829) 472.—Afr. austr.
Alopecuros, Jacq. Eclog. Gram. t. 45 = Setaria macrostachya.
Alopecuros, *Steud. Syn. Pl. Gram.* 102.—Ind. or.
amethystinum, *Beauv. Agrost.* 59, nomen.—Hab. ?
amoenum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 386.—Abyssin.
angustifolium, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 599 = longistylum.
antillarum, *Desv. Opusc.* 76.—Ind. occ.
araneosum, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 180 = lanuginosum.
arnhemicum, *F. Muell. Fragm.* viii. 109.—Austral.
articulare, *Trin. in Spreng. Neue Entdeck.* ii. 77.—Ins. Marques.
asperifolium, Hochst. & Steud. n. 100; ex Steud. Nom. ed. II. ii. 297 = Ruppellii.
asperifolium, Hochst. & Steud. n. 308; ex Steud. l. c. = elatum.
asperifolium, *Kunth, Rév. Gram.* i. 49.—Syria; Afr. bor.
asperum, Link, Hort. Berol. i. 214 = compressum.
asperum, *Schult. Mant.* ii. 149.—Afr. austr.
aureum, *Link, Hort. Berol.* i. 1215.—Ind. or.
aureum, *A. Rich. Tent. Fl. Abyss.* ii. 378.—Abyss.
bambusiforme, *Hemsl. Biol. Centr. Am. Bot.* iii. 507.—Mexic.
barbatum, *Schult. Mant.* ii. 147.—Ind. or.
Benthami, *Steud. Syn. Pl. Gram.* 105.—Afr. trop.
borbonicum, Kunth, Rév. Gram. i. 259 = setosum.
breve, *Nees, Agrost. Bras.* 281.—Bras.
breviflorum, *Steud. Syn. Pl. Gram.* 107.—Afr. trop.
calyculatum, Spreng. Syst. i. 303 = Cenchrus calyculatus.
carneum, *Hornem. ex Steud. Nom.* ed. II. ii. 297.—Hab. ?
cenchroides, *Rich. in Pers. Syn.* i. 72.—Amphig. trop. et subtrop.
cereale, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 185 = typhoideum.
chinense, *Steud. Syn. Pl. Gram.* 108.—China.
ciliare, Link, Hort. Berol. i. 213; Handb. i. 85 = cenchroides.

PENNISETUM:—
ciliatum, Parl. ex Webb, in Hook. Niger Fl. 184 = cenchroides.
cognatum, *Steud. Syn. Pl. Gram.* 107.—Afr. trop.
complanatum, *Hemsl. Biol. Centr. Am. Bot.* iii. 507.—Guatemala.
compressum, *R. Br. Prod.* 195.—Austral.
corrugatum, Nutt. Gen. Am. i. 55 = Setaria corrugata.
crinitum, Spreng. Syst. i. 302 = Setaria purpurea.
Crus-galli, Baumg. Enum. Stirp. Transs. iii. 277 = Panicum Crus-galli.
cynosuroides, Wall. Cat. n. 8563 = P. Alopecuros.
dasystachyum, *Desv. Opusc.* 76.—Guian.
dichotomum, *Delile, Fl. Egypte*, 159. t. 8. f. 1.—Aegypt.; Arab.; Ind. or.
Dilloni, *Steud. Syn. Pl. Gram.* 107, in. obs.—Abyss.
dioicum, A. Rich. Tent. Fl. Abyss. ii. 380 = adoënse.
distachyum, Rupr. in Bull. Acad. Brux. ix. II. (1842) 242 = tristachyum.
distylum, Guss. Ind. Sem. Hort. Bocc. (1826) 8; Pl. Sic. Syn. i. 115 = cenchroides.
domingense, Spreng. Syst. i. 302 = Setaria Ventenatii.
elatum, Hochst. ex Steud. Nom. ed. II. ii. 297 = dichotomum.
elegans, *Nees, ex Steud. Syn. Pl. Gram.* 105.—Afr. trop.
elegans, Rupr. ex Steud. l. c. = mollissimum.
erubescens, *Link, Hort. Berol.* i. 215.—Ind. occ.
fasciculatum, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 181 = orientale.
flaccidum, Griseb. in Goett. Nachr. (1868) 86 = incomptum.
flavescens, *J. & C. Presl, Rel. Haenk.* i. 316.—Mexic.
flavisetum, *Steud. ex Jard. in Mém. Soc. Sc. Nat. Cherb.* v. (1857) 299.—Ins. Pacif.
geniculatum, Jacq. Eclog. Gram. t. 27 = Setaria geniculata.
germanicum, Baumg. Enum. Stirp. Transs. iii. 277 = Setaria italica.
giganteum, *Regel, in Act. Hort. Petrop.* viii. (1884) 643.—Hab. ?
giganteum, *A. Rich. Tent. Fl. Abyss.* ii. 382.—Abyss.
giganteum, Tenore, ex Steud. Nom. ed. II. ii. 297 = typhoideum.
glabrum, *Steud. Syn. Pl. Gram.* 104.—Abyss.
glaucifolium, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19; *et ex A. Rich. Tent. Fl. Abyss.* ii. 382.—Abyss.
glaucum, R. Br. Prod. 195 = Setaria glauca.
gracile, *Benth. in Hook. Niger Fl.* 564.—Afr. trop.
gracilescens, *Hochst. in Flora*, xxxviii. (1855) 199.—Abyss.
Hamiltonii, *Steud. Nom.* ed. II. ii. 297.—Ind. occ.
helvolum, R. Br. Prod. i. 195 = Setaria helvola.
hirsutum, *Nees, Agrost. Bras.* 284.—Bras.
Hohenackeri, Hochst. ex Steud. Syn. Pl. Gram. 103 = P. Alopecuros.
holcoides, *Schult. Mant.* ii. 148.—Ind. or.
hordeiforme, Spreng. Syst. i. 302 = compressum.
Humboldtianum, *Hemsl. Biol. Centr. Am. Bot.* iii. 508.—Mexic.
humile, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 383.—Abyss.
identicum, *Steud. ex Jard. in Mém. Soc. Sc. Nat. Cherb.* v. (1857) 299.—Ins. Pacif.
imberbe, Edgew. in Journ. As. Soc. Beng. xxi. (1852) 181 = parviflorum.
implicatum, *Steud. Syn. Pl. Gram.* 107.—Afr. trop.
incomptum, *Nees, ex Steud. l. c.* 105.—Reg. Himal.
intertextum, *Schlecht. in Bot. Zeit.* ix. (1851) 878.—Ins. Cap. Virid.
italicum, R. Br. Prod. 195 = Setaria italica.
japonicum, Trin. in Spreng. Neue Entdeck. ii. 76 = compressum.
Karwinskyi, *Schrad. in Linnaea*, xii. (1838) 431.—Mexic.
lanatum, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 65.—Reg. Himal.
lanuginosum, *Hochst. in Flora*, xxv. (1842) *Beibl.* i. 133, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 385.—Abyss.; Ind. or.
latifolium, *Spreng. Syst.* i. 302.—Reg. Argent.
Lechleri, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Linnaei, Kunth, Rév. Gram. i. 49 = typhoideum.

PENNISETUM:—
longifolium, Fenzl, ex Steud. Syn. Pl. Gram. 109 = Cenchrus longifolius.
longistylum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 388.—Abyss.
macrochaetum, Jacq. Eclog. Gram. t. 25 = Setaria italica.
macrostachyon, Fresen. in Mus. Senckenb. ii. (1837) 135 = Ruppellii.
macrostachyum, Benth. in Hook. Niger Fl. 563 = Benthami.
macrostachyum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 177.—Ins. Molucc.
macrourum, *Trin. l. c.* 178.—Afr. austr.
megastachyum, *Steud. Syn. Pl. Gram.* 108.—Hab. ?
mexicanum, *Hemsl. Biol. Centr. Am. Bot.* iii. 508.—Mexic.
mollissimum, *Hochst. in Flora*, xxvii. (1844) 253.—Aegypt.
multiflorum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 508.—Mexic.
myosuroides, Spreng. Syst. i. 303 = Cenchrus myosuroides.
Myurus, *Parl. in Atti 8va Riun. Sc. Ital.* 586; *et in Hook. Niger Fl.* 183.—Ins. Cap. Virid.
nepalense, *Spreng. Syst.* i. 303.—Reg. Himal.
nervosum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 177.—Bras.
nicaraguense, *Fourn. in Bull. Soc. Bot. Fr.* xxvii. (1880) 293.—Nicaragua.
nigricans, *Trin. ex Steud. Nom.* ed. II. ii. 297.—Ins. Philip.
numidicum, *Paris, in Bull. Soc. Bot. Fr.* xviii. (1871) 263.—Afr. trop.
orientale, Link, Hort. Berol. i. 214 = Triticum scaberrimum.
orientale, *Rich. in Pers. Syn.* i. 72.—Alger.; Oriens; Ind. or.
ovale, *Rupr. ex Steud. Syn. Pl. Gram.* 104.—Afr. trop.
pallidum, *Nees, Agrost. Bras.* 285.—Bras.
parviflorum, *Trin. Sp. Gram. Ic.* iii. t. 288.—Afr. trop.
pauperum, *Steud. Syn. Pl. Gram.* 102.—Ins. Galapag.
pedicellatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 184.—Ins. Cap. Virid.
pentastachyum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 387.—Abyss.
persicum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 232 = orientale.
peruvianum, *Trin. in Linnaea*, x. (1836) 295.—Peruv.
petraeum, Steud. Syn. Pl. Gram. 106 = cenchroides.
phalariforme, *Steud. Syn. Pl. Gram.* 106.—Ind. or.
phalaroides, Schult. Mant. ii. 147 = dichotomum.
plumosum, *Hochst. & Steud. Syn. Pl. Gram.* 201, in syn.—Arab.
polystachyum, Schult. Mant. ii. 146 = holcoides.
Preslii, *Trin. ex Steud. Nom.* ed. II. ii. 298.—Peruv.
Prieurii, *Kunth, Rév. Gram.* ii. 411. t. 119.—Afr. trop.; Ind. or.
pungens, Nutt. Gen. Am. i. 54 = Cenchrus Elliotii.
purpurascens, H. B. & K. Nov. Gen. et Sp. i. 113 = setosum.
purpureum, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 44.—Afr. trop.
Quartinianum, *A. Rich. Tent. Fl. Abyss.* ii. 384.—Abyss.
ramosissimum, *Steud. Syn. Pl. Gram.* 105.—Afr. trop.
ramosum, *C. Sm. in Tuckey, Congo*, 251, nomen.—Ins. Cap. Virid.
respiciens, A. Rich. Tent. Fl. Abyss. ii. 379 = Setaria verticillata.
refractum, F. Muell. Fragm. viii. 109 = Plagiosetum refractum.
Richardii, Kunth, Rév. Gram. i. 49 = setosum.
riparioides, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 19, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 383.—Abyss.
riparium, *Hochst. ex A. Rich. l. c.* 381.—Abyss.
rufescens, Spreng. Syst. i. 302 = cenchroides.
Ruppellii, *Steud. Nom.* ed. II. ii. 298.—Abyss.
sagittifolium, *A. Rich. Tent. Fl. Abyss.* ii. 379.—Abyss.
scandens, Jacq. f. Cat. Hort. Vindob. (1801); ex Roem. & Schult. Syst. ii. 279 = scandens.

PENNISETUM:—
Schimperi, *Steud. Nom.* ed. II. ii. 298.—Abyss.
sciureum, *Trin. ex Steud. l. c.*—Hab. ?
setosum, Raddi, Agrost. Bras. 50 = breve.
setosum, *Rich. in Pers. Syn.* i. 72.—Reg. trop.
Sieberi, *Kunth, Rév. Gram.* i. 50.—Ins. Martinic.
sinaicum, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 11 = orientale.
spectabile, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 248 = Prieurii.
spicatum, Roem. & Schult. Syst. ii. 499 = typhoideum.
squamulatum, *Fresen. in Mus. Senckenb.* ii. (1837) 137.—Abyss.
Swartzii, F. Muell. Fragm. viii. 110 = Setaria macrostachya.
Teneriffae, *Steud. Syn. Pl. Gram.* 105.—Ins. Canar.
tenue, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 246 = orientale.
tenuispiculatum, *Steud. Syn. Pl. Gram.* 107.—Afr. trop.
Thunbergii, *Kunth, Rév. Gram.* i. 50.—Afr. austr.
tiberiadis, Boiss. Diagn. Ser. I. xiii. 43 = asperifolium.
triflorum, Nees, ex Steud. Syn. Pl. Gram. 107 = orientale
tristachyon, *Spreng. Syst.* i. 302.—Ecuador.
tristachyum, Steud. Syn. Pl. Gram. 103 = praec.
triticoides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 453.—Madag.
triticoides, Roem. & Schult. Syst. ii. 877 = setosum.
typhoideum, *Rich. in Pers. Syn.* i. 72.—Reg. trop.
uniflorum, H. B. & K. Nov. Gen. et Sp. i. 114. t. 34 = setosum.
unisetum, *Benth. in Journ. Linn. Soc.* xix. (1881) 49, 47.—Afr. austr.
Vahlii, *Kunth, Rév. Gram.* i. 49.—Arabia.
variabile, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 248 = orientale.
versicolor, *Schrad. Ind. Hort. Gotting.* (1827).—*Cf. Linnaea*, xii. (1838) 432.—Hab. ?
verticillatum, R. Br. Prod. 195 = Setaria verticillata.
villosum, *R. Br. in Salt, Abyss. App.* 62, nomen; *et in Fresen. Mus. Senckenb.* ii. (1837) 134.—Abyss.
villosum, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 599 = longistylum.
violaceum, Rich. in Pers. Syn. i. 72 = typhoideum.
viride, R. Br. Prod. 195 = Setaria viridis.

PENSTEMON, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 214; D. Don, in Edinb. N. Phil. Journ. xix. (1835) 113 = **Pentstemon**, Mitch. (Scrophul.).

PENTABOTHRA, Hook. f. Fl. Brit. Ind. iv. 18 (1883). *ASCLEPIADEAE.*
nana, *Hook. f. l. c.* 19.—Reg. Himal.

PENTABRACHION, Muell. Arg. in Flora, xlvii. (1864) 532 = **Microdesmis**, Planch. (Euphorb.).
reticulatum, Muell. Arg. l. c. 533.—Afr. trop.

PENTABRACHIUM, Muell. Arg. in DC. Prod. xv. II. 223 (1866) = praec.

PENTACAELIUM, Franch. & Sav. Enum. Pl. Jap. i. 368 (1875) = PENTACOELIUM, Sieb. & Zucc. = **Myoporum**, Banks & Soland.

PENTACAENA, Bartl. in Presl, Rel. Haenk. ii. 5. t. 49 (1830). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 14.
ACANTHONYCHIA, Rohrb. in Mart. Fl. Bras. xiv. II. 249. t. 56 (1872).
CARDIONEMA, DC. Prod. iii. 372 (1828).
camphorosmoides, *Walp. Rep.* i. 261.—Bras.
congesta, *Benth. Pl. Hartw.* 186.—Ecuador.
polycnemoides, *Bartl. in Presl, Rel. Haenk.* ii. 5. t. 49. f. 1.—Mexic.
ramosissima, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 338 = polycnemoides.
rosetta, *Walp. Rep.* i. 261.—Bras.
tenuior, Steud. Nom. ed. II. ii. 298 = polycnemoides.

PENTACALIA, Cass. in Dict. Sc. Nat. xlviii. 461 (1827) = **Senecio**, Tourn. (Compos.).
arborea, Cass. l. c. (= *Psacalium arboreum*).—Am. austr.
peltata, Cass. ex Steud. Nom. ed. II. ii. 298 = S. peltiferus.

PENTACARYA, DC. ex Meissn. Gen. 279; Comm. 187 (1840); DC. Prod. ix. 559 (1845) = **Heliotropium**, Tourn. (Borag.).
heliotropoides, DC. Prod. ix. 559 = H. anomalum.

PENTACE, Hassk. Hort. Bog. Desc. i. 110 (1858). *TILIACEAE*, Benth. & Hook. f. i. 231.
burmanica, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 47.—Burma.
polyantha, *Hassk. Hort. Bog. Desc.* i. 110.—Java.
triptera, *Mast. in Hook. f. Fl. Brit. Ind.* i. 382.—Malacca.

PENTACERAS, Hook. f. in Benth. & Hook. f. Gen. i. 298 (1862). *RUTACEAE*, Benth. & Hook. f. l. c.
australis, *Hook. f. l. c.*—Austral.

PENTACERAS, Roem. & Schult. Syst. v. p. xlviii; 570 (1818) = seq.

PENTACEROS, G. F. W. Mey. Prim. Fl. Esseq. 136 (1818) = **Buettneria**, Linn.
aculeatus, G. F. W. Mey. l. c. = B. brevipes.

PENTACHAETA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 336. *COMPOSITAE*, Benth. & Hook. f. ii. 251.
alsinoides, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 109.—Calif.
aphantochaeta, Greene, in Coult. Bot. Gaz. viii. (1883) 255 = exilis.
aurea, A. Gray, in Pacif. Rail. Rep. iv. 99 = bellidiflora.
aurea, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 336.—Calif.
bellidiflora, *Greene, in Bull. Calif. Acad.* i. III. (1885) 86.—Calif.
exilis, *A. Gray, in Proc. Am. Acad.* viii. (1873) 633.—Calif.
gracilis, *Benth. in Hook. Ic. Pl.* t. 1101.—Mexic.
Lyoni, *A. Gray, Syn. Fl. N. Am.* i. II. 445.—Calif.
paleacea, Greene, in Bull. Calif. Acad. i. IV. (1885) 189 = aurea.

PENTACHONDRA, R. Br. Prod. 549 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 612.
ericaefolia, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 271.—Tasman.
involucrata, *R. Br. Prod.* 549.—Tasman.
javanica, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 576.—Java.
mucronata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 270 = Leucopogon Fraseri.
pumila, *R. Br. Prod.* 549.—Austral.; N. Zel.
vaccinioides, Sond. in Linnaea, xxvi. (1853–55) 252 = pumila.
verticillata, *Hook. f. Fl. Tasman.* i. 256. t. 77 B.—Tasman.

PENTACLATHRA, Endl. Gen. Suppl. ii. 108 (1842) = **Polyclathra**, Bertol. ? (Cucurb.).

PENTACLETHRA, Benth. in Hook. Journ. Bot. ii. (1840) 127. *LEGUMINOSAE*, Benth. & Hook. f. i. 588.
brevifila, Benth. in Hook. Journ. Bot. ii. (1840) 128 = filamentosa.
filamentosa, *Benth. l. c.* 127.—Guian.; Ind. occ.
Griffoniana, *Baill. Adansonia*, vi. (1866) 206.—Afr. trop.
macrophylla, *Benth. in Hook. Journ. Bot.* iv. (1842) 330.—Afr. trop.

PENTACME, A. DC. in DC. Prod. xvi. II. 626 (1868) = **Shorea**, Roxb. (Dipteroc.).
siamensis, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 66 = S. siamensis.
suavis, A. DC. in DC. Prod. xvi. II. 626 = S. siamensis.

PENTACOCCA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 600 = **Phyllocosmus**, Klotzsch (Lineae).
leonenis, Turcz. l. c. 601 = Phyllocosmus africanus.

PENTACOELIUM, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 151 = **Myoporum**, Banks.
bontioides, Sieb. & Zucc. l. c. = M. bontioides.

PENTACRASPEDON, Steud. Syn. Pl. Gram. 151 (1855) = **Amphipogon**, R. Br. (Gramin.).
amphipogonoides, Steud. l. c.—Austral.

PENTACROPHYS, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 261 = **Acleisanthes**, A. Gray (Nyctag.).
Wrightii, A. Gray, l. c. = A. Wrightii.

PENTACRYPTA, Lehm. Sem. Hort. Hamb. (1828) 16.—Cf. Flora, xii. (1829) Erg. 52 = **Arracacia**, Bancr. (Umbellif.).
atropurpurea, Lehm. l. c. 17 = A. atropurpurea.

PENTADACTYLON, Gaertn. f. Fruct. iii. 219. t. 220 (1805) = **Persoonia**, Sm. (Proteac.).
angustifolium, Gaertn. f. l. c. = Persoonia linearis.

PENTADENIA, Hanst. in Linnaea, xxvi. (1853) 211 = **Columnea**, Plum. (Gesnerac.).
aurantiaca, Hanst. l. c. = C. aurantiaca.
nervosa, Hanst. l. c. 216. t. 1. f. 65; Klotzsch, ex Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 131 = C. nervosa.
strigosa, Hanst. l. c. 211 = C. strigosa.

PENTADESMA, Sabine, in Trans. Hort. Soc. v. (1824) 457. *GUTTIFERAE*, Benth. & Hook. f. i. 174.
butyracea, *Sabine, l. c.*—Afr. trop.

PENTADESMOS, Spruce, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 295 = **Moronobea**, Aubl. (Guttif.).
riparius, Spruce, l. c. 296 = M. riparia.

PENTADYNAMIS, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 76 (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 408.
incana, *R. Br. l. c.*—Austral.

PENTAGLOSSUM, Forsk. Fl. Aegypt. Arab. 11 (1775) = **Lythrum**, Linn.
linifolium, Forsk. l. c. = L. Hyssopifolia.

PENTAGLOTTIS, Tausch, in Flora, xii. (1829) 643 = **Anchusa**, Linn. (Borag.).
sempervirens, Tausch, l. c. = A. sempervirens.

PENTAGLOTTIS, Wall. Cat. n. 1156 (1831) = **Melhania**, Forsk. (Stercul.).
suberifolia, Wall. l. c. 1169, partim = Pterospermum Heyneanum.
tomentosa, Wall. l. c. 1156 = M. Hamiltoniana.

PENTAGONASTER, Klotzsch, in Otto & Dietr Allg. Gartenz. iv. (1836) 112 = **Kunzea**, Reichb. (Myrtac.).
Baxteri, Klotzsch, l. c. 115 = K. Baxteri.

PENTAGONIA, Benth. Bot. Voy. Sulph. 105. t. 39 (1844)). *RUBIACEAE*, Benth. & Hook. f. ii. 78.
MEGAPHYLLUM, Spruce, ex Baill. Hist. Pl. vii. 456, in nota (1880).
SEEMANNIA, Hook. Lond. Journ. Bot. vii. (1848) 567.
macrophylla, *Benth. Bot. Voy. Sulph.* 105. t. 39.—Panama.
macrophylla, H. Wendl. ex Hook. Bot. Mag. t. 5230 = Wendlandi.
pinnatifida, *Seem. in Hook. Lond. Journ. Bot.* vii. (1848) 566.—Panama.
Tinagita, Hook. Bot. Mag. t. 5230 = seq.
Tinajita, *Seem. in Hook. Kew Journ.* i. (1849) 187.—N. Granat.
Wendlandi, *Hook. Bot. Mag.* t. 5230.—Mexic.

PENTAGONIA, Vent. ex Steud. Nom. ed. II. ii. 298 (1841) = **Specularia**, Heist. (Campan.).
Speculum, Vent. ex Steud. l. c. = S. Speculum.

PENTAGONIUM, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 364 = **Philibertia**, H. B. & K. (Asclep.).
flavum, Schau. l. c. = Phil. flava.

PENTAGONOCARPUS, Mich. ex Parl. Fl. Ital. v. 105 (1872) = **Kosteletzkya**, Presl (Malvac.).
cordatus, Parl. Fl. Ital. v. 106 = K. cordata.
hastatus, Parl. l. c. = K. hastata.
hispidus, Parl. l. c. = K. hispida.
palmatus, Parl. l. c. = K. palmata.
paniculatus, Parl. l. c. = K. paniculata.
pentacarpos, Reichb. ex Nym. Consp. 127 = K. pentacarpa.
sagittatus, Parl. Fl. Ital. v. 106 = K. sagittata.
virginicus, Parl. l. c. = K. virginica.
Zannichellii, Parl. l. c. = K. pentacarpa.

PENTAKE, Rafin. Fl. Tellur. iv. 90 (1836) = **Cuscuta**, Linn. (Convolv.).
chinense, Rafin. l. c. = C. chinensis.

PENTALEPIS, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 496; in Edinb. N. Phil. Journ. N.S. xvii. (Jan.–Apr. 1863) 230 = **Chrysogonum**, Linn. (Compos.).
ecliptoides, F. Muell. ll. cc. = C. ecliptoides.
trichodesmoides, F. Muell. ll. cc. = C. trichodesmoides.

PENTALINON, Voigt, Hort. Suburb. Calc. 523 (1845) = **Prestonia**, R. Br. (Apocyn.).
suberectum, Voigt, l. c. (= *Haemadictyon suberectum*). —Ind. occ.

PENTALOBA, Lour. Fl. Cochinch. 154 (1790) = **Alsodeia**, Thou. (Violar.).
benghalensis, Wall. Cat. n. 4896 = A. bengalensis.
ceylanica, Arn. in Mag. Zool. Bot. ii. (1838) 543 = A. zeylanica.
corylifolia, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 341 = A. corylifolia.
fasciculata, Turcz. l. c. = A. fasciculata.
lanceolata, Wall. Cat. n. 4023 = A. lanceolata.
macrophylla, Wall. l. c. 4024 = A. Wallichiana.
Roxburghii, Wall. l. c. 7189 = A. Roxburghii.
semigyrata, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 342 = A. semigyrata.
sessilis, Lour. Fl. Cochinch. 154 = A. sessilis.
zeylanica, Thw. Enum. Pl. Zeyl. 21 = A. zeylanica.

PENTALONCHA, Hook. f. in Benth. & Hook. f. Gen. ii. 73 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
humilis, *Hook. f. l. c.*—Afr. trop.

PENTALOPHUS, A. DC. in DC. Prod. x. 86 (1846) = **Lithospermum**, Tourn. (Borag.).
longiflorus, DC. l. c. = L. angustifolium.
mandanensis, DC. l. c. 87 = L. angustifolium.

PENTAMEREA, Klotzsch, ex Baill. Adansonia, ii. 39 (1861) = seq.

PENTAMERIA, Klotzsch. ex Baill. Etud. Gén. Euph. 584 (1858) = **Bridelia**, Willd. (Euphorb.).
melanthesoides, Klotzsch, ex Baill. l. c. = B. cathartica.

PENTAMERIS, Beauv. Agrost. 92. t. 18. f. 8 (1812) = **Danthonia**, DC. (Gramin.).
airoides, Steud. Nom. ed. II. ii. 298 = D. airoides.
angulata, Steud. l. c. = D. angulata.
angustifolia, Steud. l. c. = D. angustifolia.
barbata, Steud. l. c. = D. barbata.
Buekeana, Steud. l. c. = D. Buekeana.
cirrhosula, Steud. l. c. = D. cirrhosula.
collinita, Steud. l. c. = D. collinita.
crispa, Steud. l. c. = D. crispa.
curvifolia, Nees, in Linnaea, vii. (1832) 313 = D. curvifolia.
cyathophora, Steud. Nom. ed. II. ii. 298 = D. cyathophora.
densiflora, Steud. l. c. = D. densiflora.
denudata, Steud. l. c. = D. densiflora.
distichophylla, Nees, in Linnaea, vii. (1832) 314 = D. distichophylla.
elegans, Steud. Nom. ed. II. ii. 298 = D. elegans.
eriostoma, Steud. l. c. = D. eriostoma.
filiformis, Steud. l. c. = D. filiformis.
glandulosa, Nees, in Linnaea, vii. (1832) 310 = D. glandulosa.
heptameris, Steud. Nom. ed. II. ii. 298 = D. heptamera.
hirsuta, Steud. l. c. = D. hirsuta.
involucrata, Nees, in Linnaea, vii. (1832) 313 = D. involucrata.
Lima, Steud. Nom. ed. II. ii. 299 = D. Lima.
longiglumis, Steud. l. c. = D. longiglumis.
macrantha, Nees, in Linnaea, vii. (1832) 312 = D. macrantha.
micrantha, Steud. Nom. ed. II. ii. 299 = D. micrantha.
mutica, Steud. l. c. = D. mutica.
nutans, Steud. l. c. = D. nutans.
pallescens, Nees in Linnaea, vii. (1832) 312 = D. pallescens.
papillosa, Nees, l. c. 311 = D. papillosa.
patula, Steud. Nom. ed. II. ii. 299 = D. patula.
porosa, Steud. l. c. = D. porosa.

PENTAMERIS:—
procumbens, Steud. Nom. ed. II. ii. 298, in syn = Danthonia Buekeana.
propinqua, Steud. l. c. 299 = D. propinqua.
rupestris, Steud. l. c. = D. rupestris.
scabra, Steud. l. c. = D. scabra.
speciosa, Steud. l. c. = D. speciosa.
stricta, Nees, in Linnaea, vii. (1832) 313 = D. stricta.
Thouarii, Steud. Nom. ed. I. 604 = *Thuarii*.
Thouarsii, Hochst. ex Krauss, in Flora, xxix. (1846) 120 = D. trichotoma.
Thouarsii, Kunth, Rév. Gram. i. 309. t. 66 = seq.
Thuarii, Beauv. Agrost. 93. t. 18 = D. tortuosa.
tortuosa, Nees, in Linnaea, vii. (1832) 311 = D. tortuosa.
trichotoma, Steud. Nom. ed. II. ii. 299 = D. trichotoma.
villosa, Nees, in Linnaea, vii. (1832) 310 = D. Thunbergii.
viscidula, Steud. Nom. ed. II. ii. 299 = D. viscidula.

PENTAMERIS, E. Mey. in Drège, Zwei Pfl. Docum. 210 (1843) = **Pavonia**, Cav. (Malvac.).
macrophylla, E. Mey. l. c. = Pav. Kraussiana.
microphylla, E. Mey. l. c. = Pav. microphylla.

PENTAMORPHA, Scheidw. in Bull. Acad. Brux. ix. 1. (1842) 20 = **Erythrochiton**, Nees & Mart. (Rutac.).
graveolens, Scheidw. l. c. 21 = E. brasiliensis.

PENTANEMA, Cass. in Bull. Soc. Philom. (1818) 74 = **Vicoa**, Cass. (Compos.).
Aucheri, Boiss. Diagn. Ser. I. vi. 75 = Inula pulicariaeformis.
divaricatum, Cass. in Bull. Soc. Philom. (1818) 75 (= seq. ?).—Oriens.
divaricatum, Jaub. & Spach, Illustr. t. 350 = Inula divaricata.
multicaule, Boiss. Diagn. Ser. I. vi. 75 = Inula pulicariaeformis.
pulicariaeforme, Jaub. & Spach, Illustr. iv. 88. t. 351 = Inula pulicariaeformis.
radiatum, Boiss. Diagn. Ser. II. iii. 15 = V. vestita.

PENTANISIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 21. *RUBIACEAE*, Benth. & Hook. f. ii. 104.
DIOTOCARPUS, Hochst. in Flora, xxvi. (1843) 70.
angustifolia, Hochst. in Flora, xxvii. (1844) 555 = variabilis.
cymosa, Klotzsch, in Peters, Reise Mossamb. Bot. 287, 288 = Pentas carnea.
glaucescens, Harv. in Hook. Lond. Journ. Bot. i. (1842) 22 = variabilis.
nervosa, Klotzsch, in Peters, Reise Mossamb. Bot. 287, 288 = Pentas carnea.
ouranogyne, *S. Moore, in Journ. Bot.* xviii. (1880) 4.—Afr. trop.
prunelloides, Walp. Rep. ii. 941 = variabilis.
Schweinfurthiana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 131.—Afr. trop.
suffruticosa, Klotzsch, in Peters, Reise Mossamb. Bot. 287, 288 = Pentas carnea.
variabilis, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 22.—Afr. austr.
zanzibarica, Klotzsch, in Peters, Reise Mossamb. Bot. 286 = Pentas purpurea.

PENTANOME, [Moç. & Sessé, ex] DC. Prod. i. 725, in syn. (1824) = **Zanthoxylum**, Linn. (Rutac.).
simplicifolia, [Moç. & Sessé,] l. c. = Z. Pentanome.

PENTANTHUS, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 32 = **Senecio**, Tourn. (Compos.).
jungioides, Hook. & Arn. l. c. 33.—Peruv.

PENTANTHUS, Less. Syn. Comp. 397 (1832) = **Nassauvia**, Comm. (Compos.).
aculeatus, Less. l. c. 398 = N. aculeata.

PENTANTHUS, Rafin. Fl. Tellur. iv. 70 (1836) = **Jacquemontia**, Choisy (Convolv.).
violaceus, Rafin. l. c. = J. violacea.

PENTANURA, Blume, Mus. Bot. Lugd. Bat. i. 125. t. 21 (1849). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 739.
khasiana, *Kurz, For. Fl. Brit. Burma*, ii. 196.—Reg. Himal.; Burma.
sumatrana, *Blume, Mus. Bot. Lugd. Bat.* i. 125.—Sumatra.

PENTAPANAX, Seem. Journ. Bot. ii. (1864) 294. *ARALIACEAE*, Benth. & Hook. f. i. 936.
angelicifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 144. —Am. austr.
Leschenaultii, *Seem. in Journ. Bot.* ii. (1864) 296.—Ind. or.; Burma.
micranthum, Hook. f. Fl. Brit. Ind. ii. 779 (Index), sphalm. = Leschenaultii.
parasiticum, *Seem. in Journ. Bot.* ii. (1864) 296.—Reg. Himal.
racemosum, *Seem. l. c.* 295.—Reg. Himal.
subcordatum, *Seem. l. c.*—Reg. Himal.
umbellatum, Seem. l. c. = Leschenaultii.

PENTAPELTIS, Bunge, in Lehm. Pl. Preiss. i. 292 (1845) = **Xanthosia**, Rudge (Umbell.).
peltigera, Bunge, l. c. = X. peltigera.

PENTAPERA, Klotzsch, in Linnaea, xii. (1838) 497. *ERICACEAE*, Benth. & Hook. f. ii. 589.
sicula, *Klotzsch, l. c.*—Sicil.

PENTAPETES, Linn. Diss. Dass. 23 (1747); Amoen. Acad. i. 407 (1749). *STERCULIACEAE*, Benth. & Hook. f. i. 222.
ERIORHAPHE, Miq. Pl. Jungh. 289 (1854).
MORANDA, Scop. Introd. 289 (1777).
acerifolia, Linn. Sp. Pl. 698 = Pterospermum acerifolium.
acutangula, Poir. Encyc. v. 155 = Dombeya acutangula.
alba, Hill, Veg. Syst. xv. 18 (Quid?).—Ind. or.
angulosa, Poir. Encyc. v. 155 = Dombeya acutangula.
angustifolia, Blume, Bijdr. 87 = phoenicea.
bracteosa, Steud. Nom. ed. II. ii. 299 = Melhania Denhami.
cebuana, Blanco, Fl. Filip. ed. I. 529 = phoenicea.
coccinea, Blanco, l. c. = phoenicea.
decanthera, Poir. Encyc. v. 157 = Trochetia decanthera.
Erythroxylon, Forst. f. in Comm. Gotting. ix. (1789) 61 = Melhania Erythroxylon.
Erythroxylon, Sims, Bot. Mag. t. 1000 = Melhania Melanoxylon.
ferruginea, Poir. Encyc. v. 159 = Dombeya ferruginea.
Leprieurii, Steud. Nom. ed. II. ii. 299 = Melhania abyssinica.
Melanoxylon, Soland. ex Sims, Bot. Mag. sub t. 1000 = Melhania Melanoxylon.
ovata, Poir. Encyc. v. 158 = Dombeya ovata.
ovata, Willd. Enum. Hort. Berol. 719 = Melhania abyssinica.
palmata, Poir. Encyc. v. 154 = Dombeya acutangulata.
parviflora, Fisch. ex Steud. Nom. ed. I. 604 = Melhania abyssinica?
phoenicea, *Linn. Sp. Pl.* 698.—As. trop.
populnea, Poir. Encyc. v. 154 = Dombeya populnea.
punctata, Poir. l. c. 157 = Dombeya punctata.
suberifolia, Linn. Sp. Pl. 698 = Pterospermum suberifolium.
tiliaefolia, Poir. Encyc. v. 156 = Dombeya tiliaefolia.
tomentosa, Poir. l. c. = Dombeya tomentosa.
tomentosa, 'Wall. Cat.' ex Mast. in Hook. f. Fl. Brit. Ind. i. 373 (cf. Index 728), sphalm. = Pentaglottis tomentosa.
umbellata, Poir. Encyc. v. 158 = Dombeya umbellata.
velutina, Vahl, Symb. Bot. i. 49 = Melhania velutina.

PENTAPHILTRUM, Reichb. Nom. 121 (1841) = **Physalis**, Linn. (Solan.).

PENTAPHORUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 296 = **Gochnatia**, H. B. & K. (Compos.).
fascicularis, D. Don, in Phil. Mag. (Apr. 1832) 392 = G. fascicularis.
foliolosus, D. Don, in Trans. Linn. Soc. xvi. (1830) 297 = G. rigida?
glutinosus, D. Don, in Phil. Mag. (Apr. 1832) 392 = G. glutinosa, *D. Don.*
pyrifolius, D. Don, l. c. = G. pyrifolia.
rosmarinifolius, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 109 = Nardophyllum revolutum.

PENTAPHRAGMA, Wall. Cat. n. 1313 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 558.
begoniaefolium, *Wall. Cat.* n. 1313; *G. Don, Gen. Syst.* iii. 781.—Burma; Penang.

PENTAPHRAGMA:—
grandiflorum, *Kurz, in Flora*, lv. (1872) 136.—Ins. Molucc.
macrophyllum, *Oliver, in Journ. Linn. Soc.* xv. (1875) 29.—N. Guin.

PENTAPHRAGMA, Zucc. ex Reichb. Consp. 131 (1828) = **Araujia**, Brot. (Asclep.).

PENTAPHYLAX, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 244. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 183.
euryoides, *Gardn. & Champ. l. c.*—China.

PENTAPHYLLON, Pers. Syn. ii. 352 (1807) = **Trifolium**, Tourn. (Legumin.).
Ammani, Ledeb. Ind. Sem. Hort. Dorpat. (1823) 5 = T. Lupinaster.
Lupinaster, Pers. Syn. ii. 352 = T. Lupinaster.
macrocephalum, Spreng. Syst. iii. 286 = T. megacephalum.

PENTAPHYLLUM, Gaertn. Fruct. i. 349. t. 73 (1788) = **Potentilla**, Linn. (Rosac.).
norwegicum, Gaertn. l. c. = Pot. norvegica.

PENTAPLE, Reichb. Nom. 205; et Ic. Fl. Germ. v. 37. t. 227 (1841) = **Cerastium**, Linn. (Caryoph.).
mantica, Reichb. Ic. Fl. Germ. v. 37. t. 227 = C. manticum.

PENTAPOGON, R. Br. Prod. 173 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1154.
Billardieri, Beauv. Agrost. 34 = Deyeuxia quadriseta.
Billardieri, *R. Br. Prod.* 173.—Austral.
Drummondii, Steud. Syn. Pl. Gram. 193 = Deyeuxia cylindrica.

PENTAPTELION, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 193 = **Leucopogon**, R. Br. (Epacrid.).
involucratum, Turcz. l. c. 194.—Austral.

PENTAPTERA, Roxb. Hort. Beng. 34 (1814); Fl. Ind. ii. 437 (1832) = **Terminalia**, Linn. (Combret.).
alata, Banks, ex G. Don, Gen. Syst. ii. 661 = T. paniculata.
angustifolia, Roxb. Hort. Beng. 34; Fl. Ind. ii. 437 = T. Arjuna.
Arjuna, Roxb. ll. cc. 34; 438 = T. Arjuna.
bialata, Roxb. ll. cc. 34; 441 = T. bialata.
coriacea, Roxb. ll. cc. 34; 438 = T. tomentosa.
crenulata, Roxb. ll. cc. = T. tomentosa.
cuneata, DC. Prod. iii. 15 = T. tomentosa.
glabra, Roxb. Hort. Beng. 34; Fl. Ind. ii. 440 = T. Arjuna.
gracilis, Presl, Epim. Bot. 214.—Burma.
Huliva, Buch.-Ham. ex Wall. Cat. n. 3980 = T. paniculata.
macrocarpa, Wall. l. c. n. 3982 = T. tomentosa.
Maradu, G. Don, Gen. Syst. ii. 660 = T. tomentosa.
mollis, Presl, Epim. Bot. 214 = T. mollis, *Vidal*.
obovata, DC. Prod. iii. 14 = T. Arjuna.
paniculata, Roxb. Hort. Beng. 34; Fl. Ind. ii. 442 = T. paniculata.
pyrifolia, Presl, Epim. Bot. 215 = T. pyrifolia.
pyrifolia, Wall. Cat. n. 3985 = Combretum pyrifolium.
Roxburghii, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 84 = Combretum decandrum.
Saja, Buch.-Ham. ex Wall. Cat. n. 3983 = T. myriocarpa.
tomentosa, Roxb. Hort. Beng. 34; Fl. Ind. ii. 440 = T. tomentosa.
triopteris, Heyne, ex Steud. Nom. ed. II. ii. 299 = T. paniculata.
triphylla, Presl, Bot. Bemerk. 142.—Ind. or.
undulata, Buch.-Ham. ex Wall. Cat. n. 3984 = Combretum flagrocarpum.

PENTAPTERYGIUM, Klotzsch, in Linnaea, xxiv. (1851) 47. *VACCINIACEAE*, Benth. & Hook. f. ii. 572.
flavum, *Hook. Bot. Mag.* t. 4910.—Reg. Himal.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 450.—Reg. Himal.
Listeri, *King, ex C. B. Clarke, l. c.*—Reg. Himal.
rugosum, *Hook. Bot. Mag.* t. 5198.—Reg. Himal.
serpens, *Klotzsch, in Linnaea*, xxiv. (1851) 47.—Reg. Himal.

PENTAPYXIS, Hook. f. in Benth. & Hook. f. Gen. ii. 6 (1873). *CAPRIFOLIACEAE*, Benth. & Hook. f. l. c.
stipulata, *Hook. f. Fl. Brit. Ind.* iii. 17.—Reg. Himal.

PENTARHAPHIA, Lindl. Bot. Reg. sub t. 1110 (1827) (Pentaraphia). *GESNERACEAE*, Benth. & Hook. f. ii. 1005.
CHORISANTHERA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 109.
CODONOPHORA, Lindl. Bot. Reg. sub t. 1110 (1827).
CODONORAPHIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1859) 82.
CONRADIA, Mart. Nov. Gen. et Sp. iii. 38 (1829).
DITTANTHERA, G. Don, ex Steud. Nom. ed. II. i. 521 (1840).
DUCHARTREA, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 109. t. 8.
GESNERIA, Linn. Gen. ed. I. 179 (1737).
HENRINCQUIA, Benth. & Hook. f. Gen. ii. 1006 (1876), sphalm.
HERINCQUIA, Decne. ex Jacques & Hérincq, Man. Pl. ii. 562 [1850].
MEGAPLEILIS, Rafin. Fl. Tellur. ii. 57 (1836).
OPHIANTHE, Hanst. in Linnaea, xxvi. (1853-55) 205.
STYROSINIA, Rafin. Fl. Tellur. ii. 95 (1836).
TULISMA, Rafin. l. c. 98 (1836).
VAUPELLIA, Griseb. Fl. Brit. W. Ind. 460 (1861).
albiflora, *Decne. in Ann. Sc. Nat.* Sér. III. vi. (1846) 101.—Porto Rico.
calycina, *Hanst. in Linnaea*, xxxiv. (1865–66) 307.—Jamaic.
calycosa, *Decne. in Ann. Sc. Nat.* Sér. III. vi. (1846) 99.—Jamaic.
catalpaeflora, *Decne. l. c.* 100.—Ind. occ.
corymbosa, *Hanst. in Linnaea*, xxxiv. (1865–66) 307.—Jamaic.
craniolaria, *Decne. in Ann. Sc. Nat.* Sér. III. vi. (1846) 99.—Ins. S. Doming.
cubensis, *Decne. l. c.* 108.—Cuba.
cumanensis, *Hanst. in Linnaea*, xxxiv. (1865–66) 300.—Venezuela.
cuneifolia, *Hanst. l. c.* 294.—Mexic.
duchartraeoides, *Wright, in Sauv. Fl. Cub.* 92.—Cuba.
exserta, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 104 = longiflora.
ferruginea, *Wright, in Sauv. Fl. Cub.* 93.—Cuba.
floribunda, *Carr. in Rév. Hortic.* (1878) 30, cum tab.; *et in Gard. Chron.* (1878) II. 593.—Cuba.
florida, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 106. t. 7 = longiflora.
glandulosa, *Griseb. Cat. Pl. Cub.* 199.—Cuba.
Herminieri, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 106 = longiflora.
hispida, Hanst. in Linnaea, xxxiv. (1865–66) 293 = Sloanei.
humilis, *Hanst. l. c.* 294.—Ind. occ.
incurva, *Griseb. Cat. Pl. Cub.* 199.—Cuba.
Lessertiana, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 101 = catalpaeflora.
libanensis, *Hanst. in Linnaea*, xxxiv. (1865–66) 293.—Ind. occ.
Lindleyana, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 103 = longiflora.
longiflora, *Lindl. Bot. Reg.* sub t. 1110.—Ind. occ.
montana, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 105 = longiflora.
neglecta, *Hanst. in Linnaea*, xxxiv. (1865–66) 305.—Jamaic.
parviflora, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 102 = catalpaeflora.
pedunculosa, *Decne. l. c.* 108.—Mexic.
pumila, *Hanst. in Linnaea*, xxxiv. (1865–66) 305.—Jamaic.
salicifolia, *Griseb. Cat. Pl. Cub.* 199.—Cuba.
scabra, *Hanst. in Linnaea*, xxxiv. (1865–66) 308.—Jamaic.
Sloanei, *Hanst. l. c.* 295.—Jamaic.
Swartzii, Decne. in Ann. Sc. Nat. Sér. III. vi. (1846) 103 = longiflora.
tenera, *Hanst. in Linnaea*, xxxiv. (1865–66) 304.—Costa Rica.
triflora, *Griseb. Cat. Pl. Cub.* 199.—Cuba.
ventricosa, Hanst. in Linnaea, xxxiv. (1865–66) 229 = longiflora.

PENTARHAPHIA:—
verrucosa, *Decne. in Ann. Sc. Nat.* Sér. III. vi. (1846) 107.—Cuba.
viridiflora, *Hanst. in Linnaea*, xxxiv. (1865–66) 306.—Cuba.

PENTARIA, M. Roem. Syn. Pepon. 131, 187 (1846) = **Passiflora**, Linn.
orbiculata, M. Roem. l. c. 187 = P. Murucuja.

PENTARRHAPHIS, H. B. & K. Nov. Gen. et Sp. i. 177. t. 60 (1815). Gen. dub. *GRAMINEARUM*.—Cf. Benth. & Hook. f. Gen. iii. 1096.
STROMBODURUS, Willd. ex Steud. Nom. ed. II. ii. 647 (1841).
scabra, *H. B. & K. Nov. Gen. et Sp.* i. 178. t. 60.—Mexic.

PENTARRHINUM, E. Mey. Comm. Pl. Afr. Austr. 199 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 758.
abyssinicum, *Decne. in DC. Prod.* viii. 553.—Abyss.
insipidum, *E. Mey. Comm. Pl. Afr. Austr.* 200.—Afr. austr.

PENTAS, Benth. in Bot. Mag. t. 4086 (1844). *RUBIACEAE*, Benth. & Hook. f. ii. 54.
NEUROCARPAEA, R. Br. in Salt, Abyss. App. 64 (1814).
ORTHOSTEMMA, Wall. ex Voigt, Hort. Suburb. Calc. 384 (1845).
VIGNALDIA, A. Rich. Tent. Fl. Abyss. i. 357 (1847).
arvensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 47.—Afr. trop.
carnea, *Benth. in Bot. Mag.* t. 4086.—Afr. trop.
elatior, Benth. in Hook. Niger Fl. 402, in obs.; et Walp. Rep. vi. 57 = Virecta procumbens?
Klotzschii, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 231 = carnea.
micrantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 408.—Madag.
mussaendoides, *Baker, l. c.* xx. (1883) 165.—Madag.
occidentalis, *Benth. & Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 46.—Afr. trop.
parviflora, *Benth. in Bot. Mag.* sub t. 4086.—Afr. trop.
parvifolia, *Hiern, in Journ. Linn. Soc.* xvi. (1877) 262.—Afr. trop.
pentandra, Benth. in Hook. Niger Fl. 402, in obs. = Oldenlandia macrophylla.
purpurea, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 83.—Afr. trop.
Quartiniana, Oliver, l. c. 82. t. 46 = carnea.
Schimperiana, *Vatke, in Linnaea*, xl. (1876) 192.—Afr. trop.
Thonningii, Walp. Rep. vi. 58 = Oldenlandia macrophylla.
zanzibarica, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 232 = purpurea.

PENTASACHME, Wall. ex Wight, Contrib. 60 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 769.
caudatum, *Wall. l. c.*—Reg. Himal.; Burma.
Championi, *Benth. in Hook. Kew Journ.* v. (1853) 54.—China.
glaucescens, *Decne. in DC. Prod.* viii. 627.—China.
Stauntonii, *Decne. l. c.*—China.
Wallichii, *Wight, Contrib.* 60.—Reg. Himal.

PENTASACME, G. Don, Gen. Syst. iv. 159 (1837) = praec.

PENTASCYPHUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 495. *SAPINDACEAE*.
thyrsiflorus, *Radlk. l. c.* 539.—Guian.

PENTASPADON, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 168. t. 24. *ANACARDIACEAE*, Benth. & Hook. f. i. 419.
Motleyi, *Hook. f. l. c.*—Borneo.
velutinus, Hook. f. Fl. Brit. Ind. ii. 28 = Microstemon velutina.

PENTASTACHYA, Hochst. ex Steud. Nom. ed. II. ii 299 (1841) = **Pennisetum**, Rich. (Gramin.).
abyssinica, Hochst. l. c. = Pennis. pentastachyum.

PENTASTEMON, Batsch, Tab. Affin. 193 (1802) = **Pentstemon**, Mitch. (Scrophul.).

PENTASTEMUM, Steud. Nom. ed. II. ii. 299 (1841) = praec.

PENTASTICHA, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 330 = **Fuirena**, Rottb. (Cyperac.).
madagascariensis, Turcz. l. c. = F. stricta.

PENTATAPHRUS, Schlecht. in Linnaea, xx. (1847) 618 = **Astroloma**, R. Br. (Epacrid.).
Behrii, Schlecht. l. c. = A. conostephioides.

PENTATAXIS, D. Don, in Mem. Wern. Soc. v. (1826) 550 = **Helichrysum**, Vaill. (Compos.).
micrantha, D. Don, l. c. = H. Zeyheri.

PENTATROPIS, R. Br. in Salt, Abyss. App. 64 (1814). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 764.
EUTROPIS, Falc. in Proc. Linn. Soc. i. (1839) 15.
RHYNCHARRHENA, F. Muell. Fragm. i. 128 (1859).
atropurpurea, *Benth. Fl. Austral.* iv. 329.—Austral.
cynanchoides, *R. Br. in Salt, Abyss. App.* 64.—Abyss.; Persia; Ind. bor. occ.
Kempeana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 172.—Austral.
linearis, *Decne. in DC. Prod.* viii. 536.—Austral.
madagascariensis, *Decne. l. c.*—Madag.
microphylla, Wall. Cat n. 8213 D = cynanchoides.
microphylla, *Wight & Arn. Contrib.* 52.—Ind. or.
quinquepartita, *Benth. Fl. Austral.* iv. 329.—Austral.
senegalensis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 328 = cynanchoides?
spiralis, Decne. l. c. 327 = cynanchoides.

PENTECA, Rafin. Sylva Tellur. 62 (1838) = **Croton**, Linn. (Euphorb.).
tomentosa, Rafin. l. c. = C. dioicus.

PENTELESIA, Rafin. Sylva Tellur. 146 (1838) = **Bignonia**, Linn.
discolor, Rafin. l. c. = B. carichanensis.

PENTENA, Rafin. Fl. Tellur. iv. 96 (1836) = **Scabiosa**, Tourn. (Dipsac.).
ochroleuca, Rafin. l. c. = S. ochroleuca.
uniseta, Rafin. l. c. = S. gramuntia.

PENTHEA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Disa**, Berg. (Orchid.).
atricapilla, Harv. ex Lindl. in Hook. Lond. Journ. Bot. i. (1842) 17 = D. atricapilla.
elegans, Sond. in Linnaea, xx. (1847) 220, nomen = D. elegans.
filicornis, Lindl. Gen. et Sp. Orch. 361 = D. patens.
melaleuca, Lindl. l. c. = D. melaleuca.
minor, Sond. in Linnaea, xix. (1847) 104 = D. minor.
obtusa, Lindl. Gen. et Sp. Orch. 361 = D. Richardiana.
patens, Lindl. l. c. 362 = D. tenuifolia.
Pumilio, Lindl. in Journ. Linn. Soc. vi. (1862) 138 = Brachycorythis Pumilio.
reflexa, Lindl. Gen. et Sp. Orch. 361 = D. patens.
triloba, Sond. in Linnaea, xix. (1847) 104 = D. oligantha.
virginea, Harv. Gen. S. Afr. Pl. 325 = D. Richardiana.

PENTHEA, Spach, Hist. Vég. Phan. x. 9 (1841) = **Barnadesia**, Mutis (Compos.).
laxa, Steud. Nom. ed. II. ii. 300 = B. rosea.

PENTHORUM, Gronov. ex Linn. Coroll. Gen. 8 (1737). *CRASSULACEAE*, Benth. & Hook. f. i. 661.
chinense, Pursh, Fl. Am. Sept. i. 323 = sedoides.
circinale, Salisb. Prod. 308 = sedoides.
humile, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 65.—Reg. Amur.
intermedium, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. VII. 152.—China.
sedoides, *Linn. Sp. Pl.* 432.—Am. bor.

PENTHYSA, Rafin. Fl. Tellur. iv. 86 (1836) = **Lobostemon**, Lehm. (Borag.).
glauca, Rafin. l. c. = L. glaucophyllus.
levigata, Rafin. l. c. = L. laevigatus.
strigosa, Rafin. l. c. = L. strigosus.

PENTIPHRAGMA, Hook. Bot. Misc. i. (1830) t. 57, err. typ. = **Pentaphragma**, Wall. (Compos.).

PENTITDIS, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 246 (1850) = **Opilia**, Roxb. (Olacin.).
scandens, Zipp. ex Blume, l. c. = O. Pentitdis.

PENTLANDIA, Herb. in Bot. Reg. (1839) t. 68 = **Urceolina**, Reichb. (Amaryll.).
miniata, Herb. l. c. = U. miniata.
sulivanica, M. Roem. Syn. Ensat. 55 = U. miniata.

PENTOCNIDE, Rafin. Fl. Tellur. iii. 48 (1836) = **Pouzolzia**, Gaudich. (Urtic.).
glomerata, Rafin. l. c. = P. indica.

PENTODON, Ehrenb. ex Boiss. Fl. Orient. iv. 925 (1879) = **Kochia**, Roth (Chenopod.).
barbatus, Ehrenb. l. c. = K. monticola.

PENTODON, Hochst. in Flora, xxvii. (1844) 552 = **Oldenlandia**, Linn. (Rubiac.).
abyssinicus, Hochst. l. c. = O. macrophylla.
decumbens, Hochst. l. c. = O. macrophylla.
Halei, A. Gray, Syn. Fl. N. Am. i. II. 28 = O. Halei.
pentandrus, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 231 = O. macrophylla.

PENTOPETIA, Decne. in DC. Prod. viii. 500 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 741.
androsaemifolia, *Decne. l. c.*—Madag.
Cotoneaster, *Decne. l. c.*—Madag.
gracilis, *Decne. l. c.*—Madag.

PENTOSTEMON, Rafin. Atl. Journ. 176 (1833) = **Pentstemon**, Mitch. (Scrophul.).

PENTRIAS, Benth. & Hook. f. Gen. iii. 28 (1880), sphalm. = seq.

PENTRIUS, Rafin. Fl. Tellur. iii. 42 (1836) = **Amaranthus**, Linn.
oleraceus, Rafin. l. c. = A. oleraceus.

PENTROPIS, Rafin. Fl. Tellur. ii. 80 (1836) = **Campanula**, Tourn.
saxatilis, Rafin. l. c. = C. saxatilis.

PENTSTEMON, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 214 (Penstemon). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 940.
DASANTHERA, Rafin. in Am. Monthly Mag. (1818) 267.
ELMIGERA, Reichb. Consp. 123 (1828).
LEIOSTEMON, Rafin. Neogenyt. 2 (1825).
LEPIDOSTEMON, Lem. Illustr. Hortic. ix. (1862) t. 315.
LEPTEIRIS, Rafin. New Fl. Am. ii. 73 (1836).
PENTASTEMON, Batsch, Tab. Affin. 193 (1802).
acuminatus, *Dougl. in Bot. Reg.* t. 1285.—Am. bor. occ.
albidus, *Nutt. Gen. Am.* ii. 53.—Am. bor occ.
albidus, Torr. in Ann. Lyc. N. York, ii. (1828) 229, partim = Jamesii.
alpinus, Torr. l. c. i. (1824) 35 = glaber.
ambiguus, *Torr. l. c.* ii. (1828) 228.—Am. bor. occ.
amoenus, Kunze, Hort. Lips. Sem. (1840).—Cf. Linnaea, xvi. (1842) Litt. 107 = venustus.
amplexicaulis, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 461 = acuminatus.
amplexicaulis, Moench, Meth. 442 = pubescens.
angustifolius, Fraser, ex Pursh, Fl. Am. Sept. ii. 738 = caeruleus.
angustifolius, Poir. in Dict. Sc. Nat. xxxviii. 385 = campanulatus.
antirrhinoides, *Benth. in DC. Prod.* x. 594.—Calif.
argutus, Paxt. Mag. Bot. vi. (1839) 271 = diffusus?
atropurpureus, G. Don, in Loud. Hort. Brit. 243 = campanulatus.
attenuatus, *Dougl. in Bot. Reg.* t. 1295.—Am. bor. occ.
azureus, *Benth. Pl. Hartw.* 327.—Am. bor. occ.
baccharifolius, *Hook. Bot. Mag.* t. 4627.—Texas.
barbatus, *Roth, Catalect.* fasc. iii. 49; *Nutt. Gen. Am.* ii. 53.—Am. bor. occ.
Bradburii, Pursh, Fl. Am. Sept. ii. 738 = grandiflorus.
breviflorus, *Lindl. Bot. Reg.* t. 1946.—Calif.
brevilabris, A. Gray, in Proc. Am. Acad. xvii. (1881–82) 229 = cerrosensis.
Bridgesii, *A. Gray, l. c.* vii. (1868) 379.—Am. bor. occ.
caeruleus, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 651 = Hartwegii.
caeruleus, Nutt. vide coeruleus.
caesius, *A. Gray, in Proc. Am. Acad.* xix. (1883) 92.—Calif.

PENTSTEMON:—
caespitosus, *Nutt. ex A. Gray, Proc. Am. Acad.* vi. (1862) 66.—Am. bor. occ.
campanulatus, *Willd. Sp. Pl.* iii. 228.—Mexic.
canoso-barbatus, Kellogg, in Proc. Calif. Acad. ii. (1863) 15 = breviflorus.
carinatus, Kellogg, l. c. i. (1855) 63 = breviflorus.
centranthifolius, *Benth. Scroph. Ind. Introd.* 7, adnot.—Calif.
centranthifolius, S. Wats. Bot. King's Exped. 219 = Eatoni.
cerrosensis, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 19.—Calif.
Clevelandi, *A. Gray, l. c.* xi. (1876) 94.—Calif.; Mexic.
Cobaea, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 182.—Am. bor. occ.
coccineus, Engelm. in Wisliz. Tour N. Mexic. 107 (Sketch, 23) = barbatus.
coccineus, Hoffmgg.—Cf. Linnaea, xvi. (1842) Litt. 281 = Hartwegii.
coeruleus, *Nutt. Gen. Am.* ii. 52.—Am. bor. occ.
coeruleus, Torr. in Ann. Lyc. N. York, ii. (1828) 229 = barbatus.
comarrhenus, A. Gray, in Proc. Am. Acad. xii. (1877) 81 = strictus.
confertus, *Dougl. in Bot. Reg.* t. 1260.—Am. bor. occ.
connatus, Deppe, in Otto & Dietr. Allg. Gartenz. xii. (1844) 276 = perfoliatus.
cordatus, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 17 = perfoliatus.
cordatus, Walp. Rep. iii. 249 = cordifolius.
cordifolius, *Benth. Scroph. Ind. Introd.* 7.—Calif.
corymbosus, *Benth. in DC. Prod.* x. 593.—Calif.
crassifolius, Lindl. Bot. Reg. (1838) t. 16 = Menziesii.
crassifolius, Shuttl. ex Benth. in DC. Prod. x. 327 = pubescens.
cristatus, *Fras. Cat.* (1813); *ex Nutt. Gen. Am.* ii. 52.—Am. bor. occ.
Cusickii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 106.—Am. bor. occ.
cyananthus, Hook. Bot. Mag. t. 4464; et in Moore & Ayres, Gard. Mag. Bot. iii. (1851) 249 = glaber.
dasyphyllus, *A. Gray, in Torr. Bot. Mex. Bound.* 112.—N. Mexic.
deustus, *Dougl. in Bot. Reg.* t. 1318.—Am. bor. occ.
diffusus, *Dougl. l. c.* t. 1132.—Am. bor. occ.
Digitalis, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 181 = laevigatus.
dissectus, *Ell. Sketch*, ii. 129.—Am. bor.
Douglasii, Hook. Fl. Bor. Am. ii. 98 = Menziesii.
Eatoni, *A. Gray, in Proc. Am. Acad.* viii. (1872) 395.—Am. bor. occ.
elegans, Poir. in Dict. Sc. Nat. xxxviii. 384 = campanulatus.
erianthera, Fras. Cat. (1813); ex Nutt. Gen. Am. ii. 52 = glaber.
erianthera, Pursh, Fl. Am. Sept. ii. 737 = cristatus.
Fendleri, Torr. & Gray, in Pacif. Rail. Rep. ii. 168 = acuminatus.
filifolius, Nutt. ex Benth. in DC. Prod. x. 322 = laricifolius.
Fremonti, A. Gray, in Bot. Calif. i. 622 = heterodoxus.
Fremonti, *Torr. & Gray, in Proc. Am. Acad.* vi. (1862) 60.—Am. bor. occ.
frutescens, *Lamb. in Trans. Linn. Soc.* x. (1811) 259. t. 6.—Unalashk.; Kamtsch.; Japon.
Gairdneri, *Hook. Fl. Bor. Am.* ii. 99.—Am. bor. occ.
gentianoides, Lindl. Bot. Reg. (1838) t. 3 = Hartwegii.
gentianoides, *Poir. in Dict. Sc. Nat.* xxxviii. 385.—Mexic.
giganteus, *Morr. in Ann. Soc. Bot. Gand*, ii. (1846) 143. t. 63.—Am. bor.
glaber, *Pursh, Fl. Am. Sept.* ii. 738.—Am. bor. occ.
glaberrimus, *Bevis, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 654 (Quid?).—Am. bor.?
glandulosus, *Dougl. ex Lindl. Bot. Reg.* t. 1262.—Am. bor. occ.
glaucifolius, A. Gray, in Pacif. Rail. Rep. vi. 82 = azureus.
glaucophyllus, Scheele, in Linnaea, xxi. (1848) 763 = laevigatus.
glaucus, *R. Grah. in Edinb. N. Phil. Journ.* (July–Oct. 1829) 348.—Am. bor. occ.
Gordoni, Hook. Bot. Mag. t. 4319 = glaber.
Gordonianus, A. Gray, in Pacif Rail. Rep. xii. II. Bot. 45 = praec.
gracilentus, *A. Gray, l. c.* vi. 82.—Am. bor. occ.

PENTSTEMON:—
gracilis, *Nutt. Gen. Am.* ii. 52.—Am. bor. occ.
grandiflorus, *Fras. Cat.* (1813); *ex Nutt. l. c.* 53.—Am. bor.
Hallii, *A. Gray, in Proc. Am. Acad.* vi. (1862) 70.—Am. bor. occ.
Harbourii, *A. Gray, l. c.* 71.—Am. bor. occ.
Hartwegii, *Benth. Pl. Hartw.* 48.—Mexic.
Havardi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 306.—Texas.
heterander, Torr. & Gray, in Pacif. Rail. Rep. ii. 123 = deustus.
heterodoxus, *A. Gray, Syn. Fl. N. Am.* ii. 1. 269.—Calif.
heterophyllus, *Lindl. Bot. Reg.* t. 1899.—Am. bor. occ
heterophyllus, S. Wats. Bot. King's Exp. 222 = azureus.
hirsutus, Willd. Sp. Pl. iii. 227 = pubescens.
Humboldtii, G. Don, Gen. Syst. iv. 640 = imberbis.
humilis, *Nutt. ex A. Gray, in Proc. Am. Acad.* vi. (1862) 69.—Am. bor. occ.
ilicifolius, Nutt. ex A. DC. in DC. Prod. x. 593 = deustus.
imberbis, *Trautv. in Bull. Sc. Acad. Pétersb.* v. (1839) 345.—Mexic.
inermis, Poir. in Dict. Sc. Nat. xxxviii. 385 = imberbis.
Jaffrayanus, Hook. Bot. Mag. t. 5045 = azureus.
Jamesii, *Benth. in DC. Prod.* x. 325.—Am. bor. occ.
Jeffreyanus, A. Murr. Circ. Oreg. Comm. 2 = azureus.
Kingii, *S. Wats. Bot. King's Exp.* 222.—Am. bor. occ.
Kleei, Greene, in Bull. Torrey Bot. Club, x. (1883) 127 = Rattani.
Kunthii, G. Don, in Loud. Hort. Brit. 243 = campanulatus.
labrosus, *Hort.—Cf. Gard. Chron.* (1883) ii. 536; *Hook. f. Bot. Mag.* t. 6738.—Calif.
laetus, *A. Gray, in Proc. Am. Acad.* vi. (1866) 76.—Calif.
laevigatus, *Ait. Hort. Kew.* ed. I. ii. 361.—Am. bor.
lanceolatus, *Benth. Pl. Hartw.* 22.—Mexic.
laricifolius, *Hook. & Arn. Bot. Beech. Voy.* 376.—Am. bor. occ.
latifolius, Hoffmgg. Verz. Pfl. 98 = pubescens.
Lemmoni, *A. Gray, in Bot. Calif.* i. 557.—Calif.
Lewisii, Benth. in DC. Prod. x. 321 = Menziesii.
linarioides, *A. Gray, in Torr. Bot. Mex. Bound.* 112.—N. Mexic.
Lobbii, Hort. ex Lem. Illustr. Hortic. (1862) t. 315 = antirrhinoides.
longifolius, Scheele, in Linnaea, xxi. (1848) 764 = pubescens.
Macewani ×, Paxt. Mag. Bot. xiv. (1848) 174.
Mackayanus, Knowles & Westc. Flor. Cab. ii. (1838) 117 = pubescens.
Menziesii, *Hook. Fl. Bor. Am.* ii. 98.—Am. bor. occ.
mexicanus, Hort. ex A. DC. in DC. Prod. x. 331, in nota = Tetranema mexicanum.
micranthus, Nutt. in Journ. Acad. Philad. vii. (1834) 45 = confertus.
micranthus, Torr. in Frém. Rep. 89 = strictus.
microphyllus, *A. Gray, in Pacif. Rail. Rep.* iv. 119.—Arizona.
miniatus, *Lindl. Bot. Reg.* (1847) t. 14.—Mexic.
multiflorus, Chapm. ex Benth. in DC. Prod. x. 327 = pubescens.
Murrayanus, *Hook. Bot. Mag.* t. 3472.—Texas.
nemorosus, Trautv. in Bull. Sc. Acad. Pétersb. v. (1839) 345 = Chelone nemorosa.
Newberryi, A. Gray, in Pacif. Rail. Rep. vi. 82 = Menziesii.
nitidus, Dougl. ex Benth. in DC. Prod. x. 323 = acuminatus.
nudiflorus, *A. Gray, in Proc. Am. Acad.* xx. (1885) 206.—Arizona.
Nuttallii, Beck, in Am. Journ. Sc. Ser. I. xiv. (1828) 120 = laevigatus?
oregonensis, *A. Murr. Circ. Oreg. Comm.* 2.—Oregon.
ovatus, *Dougl. in Bot. Mag.* t. 2903.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 379.—Arizona.
Parishii, *A. Gray, l. c.* xvii. (1881–82) 228.—Calif.
Parryi, *A. Gray, Syn. Fl. N. Am.* ii. 1. 264.—Am. bor. occ.
pauciflorus, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 461 = Phlox pilosa.
pauciflorus, Greene, in Coult. Bot. Gaz. vi. (1881) 218 = lanceolatus.

PENTSTEMON:—
pedunculatus, Nutt. ex A. DC. in DC. Prod. x. 593 = Menziesii.
perfoliatus, *A. Brongn. in Hortic. Univ.* (Janv. 1844) cum ic.; *ex Benth. in DC. Prod.* x. 326.—Mexic.
pinifolius, *Greene, in Coult. Bot. Gaz.* vi. (1881) 218.—Arizona.
Pononi, Hort.—Cf. Gard. Chron. (1872) 733 = confertus *vel* acuminatus.
procerus, Dougl. ex R. Grah. in Edinb. N. Phil. Journ. (July-Oct. 1829) 348 = confertus.
pruinosus, *Dougl. in Bot. Reg.* t. 1280.—Am. bor. occ.
pubescens, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 360.—Am. bor.
pulchellus, Lindl. Bot. Reg. t. 1138 = campanulatus.
pumilus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 46.—Am. bor. occ.
puniceus, *A. Gray, in Torr. Bot. Mex. Bound.* 113.—Arizona.
puniceus, Lilja, in Linnaea, xvii. (1843) 111 = Hartwegii.
Rattani, *A. Gray, in Proc. Am. Acad.* xv. (1880) 50.—Calif.
Richardsonii, *Dougl. in Lindl. Bot. Reg.* t. 1121.—Am. bor. occ.
rivularis, Dougl. ex Trautv. in Bull. Sc. Acad. Pétersb. v. (1839) 346 = Richardsonii.
Roezli, *Regel, Gartenfl.* (1872) 239.—Am. bor. occ.
roseus, G. Don, in Loud. Hort. Brit. 243 = campanulatus.
rostriflorus, Kellogg, in Proc. Calif. Acad. ii. (1863) 15 = Bridgesii.
Rothrockii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 260.—Calif.
rubescens, *A. Gray, in Proc. Am. Acad.* xix. (1883) 92.—Am. bor.
Scouleri, Dougl. in Bot. Reg. t. 1277 = Menziesii.
secundiflorus, *Benth. in DC. Prod.* x. 325.—Am. bor. occ.
serrulatus, Menzies, in Hook. Fl. Bor. Am. ii. 95 = diffusus.
speciosus, Dougl. in Bot. Reg. t. 1270 = glaber.
spectabilis, *Thurb. ex Torr. & Gray, in Pacif. Rail. Rep.* iv. 119.—Calif.
spicatus, Hort.—Cf. Gard. Chron. (1872) 733 = confertus *vel* acuminatus.
staticifolius, Lindl. Bot. Reg. t. 1770 = glandulosus.
stenophyllus, *A. Gray, in Torr. Bot. Mex. Bound.* 112.—Mexic.
strictus, *Benth. in DC. Prod.* x. 324.—Am. bor. occ.
suffruticosus, Dougl. ex Benth. l. c. 330 = venustus.
tenellus, Kellogg, in Proc. Calif. Acad. i. (1855) 56 = azureus.
tenuifolius, *Benth. Pl. Hartw.* 22.—Mexic.
teretiflorus, Fras. Cat. (1813); ex Nutt. Gen. Am. ii. 53 = albidus.
ternatus, *Torr. ex A. Gray, in Torr. Bot. Mex. Bound.* 115.—Calif.
Themisteri, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 657 = diffusus.
Thurberi, Torr. in Pacif. Rail. Rep. vii. 15 = ambiguus.
Tolmiei, Hook. Fl. Bor. Am. ii. 98 = confertus.
Torreyi, Benth. in DC. Prod. x. 324 = barbatus.
Torreyi, *Hort. ex Regel, Ind. Sem. Hort. Petrop.* (1860) 40.—Texas.
triphyllus, *Dougl. in Bot. Reg.* t. 1245.—Am. bor. occ.
tubiflorus, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 181.—Am. bor. occ.
venustus, *Dougl. in Bot. Reg.* t. 1309.—Am. bor. occ.
verticillatus, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 18 = perfoliatus.
virgatus, *A. Gray, in Torr. Bot. Mex. Bound.* 112.—N. Mexic.
viscidulus, Nees, in Neuwied, Reise Nord. Am. ii. 444 = albidus.
Wardii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 82.—Am. bor. occ.
Watsoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 267.—Am. bor. occ.
Whippleanus, *A. Gray, in Proc. Am. Acad.* vi. (1862) 73.—N. Mexic.
Wrightii, *Hook. Bot. Mag.* t. 4601.—Am. bor. occ.

PENTSTEMONACANTHUS, Nees, in Mart. Fl. Bras. ix. 159 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1082.
modestus, *Nees, l. c.*—Bras.

PENTSTEIRA et PENTSTERIA, Griff. Notul. iv. 11 (1854) = **Torenia**, Linn. (Scrophul.).
dichotoma, Griff. l. c. 119 = T. parviflora.
grandiflora, Griff. l. c. 118 = T. vagans.
paniculata, Griff. l. c. 119 = Vandellia crustacea.
racemosa, Griff. l. c. 120 = T. flava.

PENTSTEMUM, Steud. Nom. ed. II. ii. 299 (1841) = **Pentstemon**, Mitch. (Scrophul.).

PENTULOPS, Rafin. Fl. Tellur. iv. 42 (1836) = **Xylobium**, Lindl. (Orchid.).
discolor, Rafin. l. c. (= *Maxillaria decolor*).—Ind. occ.

PENTZIA, Thunb. Prod. Pl. Cap. 145 (1800). *COMPOSITAE*, Benth. & Hook. f. ii. 433.
annua, *DC. Prod.* vi. 138.—Afr. austr.
Burchellii, Fenzl, ex Harv. & Sond. Fl. Cap. iii. 172 = Matricaria Burchellii.
cinerascens, *DC. Prod.* vi. 138.—Afr. austr.
Cooperi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 173.—Afr. austr.
cotuloides, *DC. Prod.* vi. 138.—Afr. austr.
crenata, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
dichotoma, *DC. Prod.* vi. 138.—Afr. austr.
elegans, *DC. l. c.* 136.—Afr. austr.
flabelliformis, Willd. Sp. Pl. iii. 1808 = crenata.
globosa, *Less. Syn. Comp.* 266.—Afr. austr.
microphylla, *DC. Prod.* vi. 137.—Afr. austr.
nana, *Burch. Trav.* i. 400; *et ex DC. Prod.* vi. 137.—Afr. austr.
pinnatifida, *Oliver, in Hook. Ic. Pl.* t. 1340.—Afr. austr.
punctata, Harv. in Harv. & Sond. Fl. Cap. iii. 172 = Matricaria Burchellii.
quinquefida, *Less. Syn. Comp.* 266.—Afr. austr.
sphaerocephala, *DC. Prod.* vi. 138.—Afr. austr.
spinescens, *Less. Syn. Comp.* 266.—Afr. austr.
tortuosa, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 174.—Afr. austr.
trifurcata, *Sch. Bip. in Flora*, xxvii. (1844) 675.—Afr. austr.
virgata, *Less. Syn. Comp.* 266.—Afr. austr.

PEPERIDIA, Kostel. Allg. Med. Pharm. Fl. ii. 455 (1831) = **Piper**, Linn.
Afzeliana, Kostel. l. c. 456 = Piper subpeltatum.
peltata, Kostel. l. c. 455 = Piper peltatum.
sidaefolia, Kostel. l. c. = Piper umbellatum.
subpeltata, Kostel. l. c. 456 = Piper subpeltatum.
umbellata, Kostel. l. c. 455 = Piper umbellatum.

PEPERIDIA, Reichb. Consp. 212[a] (1828) = **Chloranthus**, Sw.

PEPERIDIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Renealmia**, Linn. (Scitam.).
tubulatum, Lindl. l. c. = R. exaltata.

PEPEROMIA, Ruiz & Pav. Prod. 8 (1794). *PIPERACEAE*, Benth. & Hook. f. iii. 132.
Acrocarpidium, Miq. Syst. Pip. 51 (1843).
Erasmia, Miq. in Inst. Versl. & Med. 1842 (1843) 81.
Micropiper, Miq. in Bull. Sc. Phys. Mat. Néerl. (1839) 447.
Phylloberyon, Miq. Syst. Pip. 50 (1843).
Tildenia, Miq. in Inst. Versl. et Med. 1842 (1843) 82.
Troxirum, Rafin. Sylva Tellur. 85 (1838).
abyssinica, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 419.—Abyss.
acuminata, A. Dietr. Sp. Pl. i. 148 = nemorosa.
acuminata, Griseb. Cat. Pl. Cub. 63 = guadaloupensis.
acuminata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 416 = Rupertiana.
acuminata, *Ruiz & Pav. Fl. Per.* i. 32. t. 51. f. a.—Peruv.
acuminatissima, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 416.—Bras.
acutifolia, *C. DC. in Journ. Bot.* iv. (1866) 142.—Peruv.
adscendens, *C. DC. l. c.* 140.—Venezuela.
agapatensis, *C. DC. in DC. Prod.* xvi. 1. 455.—Mexic.; Peruv.

PEPEROMIA :—
aguacatensis, *C. DC. in Linnaea*, xxxvii. (1871–73) 376.—Costa Rica.
alata, *Ruiz & Pav. Fl. Per.* i. 31. t. 48. f. b.—Am. trop.
alismaefolia, *Presl, Epim. Bot.* 222.—Peruv.
alpina, *A. Dietr. Sp. Pl.* i. 157.—Jamaic.
alpina, Griseb. Cat. Pl. Cub. 64 = P. Pseudopereskiaefolia.
alpina, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 19 = quadrifolia.
alsinoides, *A. Dietr. Sp. Pl.* i. 168.—Am. bor.
amantlanensis, *C. DC. in Linnaea*, xxxvii. (1871–73) 369.—Mexic.
amplexicaulis, *A. Dietr. Sp. Pl.* i. 144.—Jamaic.
amplexifolia, *A. Dietr. l. c.* 145.—Jamaic.
andinacea, *C. DC. in Journ. Bot.* iv. (1866) 134.—Ecuador.
angularis, *C. DC. in DC. Prod.* xvi. I. 415.—Ecuador.
angulata, *H. B. & K. Nov. Gen. et Sp.* i. 66.—Venezuela.
angustata, *H. B. & K. l. c.* 68.—Venezuela.
angustifolia, C. DC. in DC. Prod. xvi. I. 438, in syn. = Roxburghiana.
antillarum, *C. DC. l. c.* 447.—Cuba.
aphylla, *C. DC. l. c.* 465.—N. Granat.
arabica, *Decne. ex Miq. Syst. Pip.* 121.—Afr. austr.
arboricola, *C. DC. in Linnaea*, xxxvii. (1871–73) 374.—Mexic.
argyrea, Hort.—Cf. Gard. Chron. (1865) 934 = seq.?
argyreia, *E. Morr. in Belg. Hortic.* xvii. (1867) 2. t. 2.—Am. austr.
arifolia, *Miq. Syst. Pip.* 72.—Bras.
asarifolia, *Schlecht. in Linnaea*, v. (1830) 75.—Mexic.
atropunctata, *Miq. in Arch. Néerl.* vi. (1871) 171; *et in Verh. Kon. Acad. Wetensch.* II. v. (1871) 233.—Bras.
augescens, *Miq. ll. cc.*—Bras.
australis, Phil. in Linnaea, xxx. (1859–60) 200 = portulacaefolia.
Balansana, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 313.—Paraguay.
bangrooana, *C. DC. in Journ. Bot.* iv. (1866) 134.—Afr. trop.
barbarana, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 314.—Paraguay.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1883) 239.—Madag.
basellaefolia, *H. B. & K. Nov. Gen. et Sp.* i. 66.—Peruv.
Baueriana, *Miq. Syst. Pip.* 120.—Ins. Norfolk.
Belangeri, *C. DC. in DC. Prod.* xvi. I. 411.—Ins. Martinic.
Berlandieri, Miq. Syst. Pip. 158 = reflexa.
Bernieriana, *Miq. l. c.* 124.—Madag.
Bernouillii, *C. DC. in Linnaea*, xxxvii. (1871–73) 367.—Guatemala.
Berteroana, *Miq. Syst. Pip.* 114.—Ins. J. Fernand.
bilineata, *Miq. l. c.* 106.—Java.
biuncialis, *H. B. & K. Nov. Gen. et Sp.* i. 65.—N. Granat.
blanda, *H. B. & K. l. c.* 67.—Venezuela.
blattarum, A. Dietr. Sp. Pl. i. 150 = Piper blattarum.
Boivinii, *C. DC. in Journ. Bot.* iv. (1866) 146.—Ins. Comor.
boliviensis, *C. DC. in DC. Prod.* xvi. I. 453.—Bolivia.
borbonensis, *Miq. Syst. Pip.* 109.—Ins. Borbon.
Botterii, *C. DC. in Journ. Bot.* iv. (1866) 146.—Mexic.
Bourgeaui, *C. DC. in Linnaea*, xxxvii. (1871–73) 370.—Mexic.
brachyphylla, Sweet, Hort. Brit. ed. I. 380 = trinervis.
brachystachya, *A. Dietr. Sp. Pl.* i. 181.—Peruv.
bracteiflora, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 317.—Ins. Martinic.
brasiliensis, *Miq. in Mart. Fl. Bras.* iv. I. 10.—Bras.
brevipes, *C. DC. in DC. Prod.* xvi. I. 397.—Am. trop.
caffra, E. Mey. ex Miq. Syst. Pip. 121 = arabica.
Caldasiana, C. DC. in Linnaea, xxxvii. (1871–73) 365 = Martiana.
caledonica, *C. DC. in Journ. Bot.* iv. (1866) 135.—N. Caled.
campinasana, *C. DC. in Linnaea*, xxxvii. (1871–73) 381.—Bras.
camptotricha, *Miq. Syst. Pip.* 144.—Mexic.
candida, *Miq. l. c.* 105.—Java.
Candolleana, Miq. l. c. 146 (err. typ. 246) = portulacoides.

PEPEROMIA :—
capensis, Loud. Hort. Brit. 13 = Piper capense.
cardiophylla, C. DC. in Journ. Bot. iv. (1866) 141.—N. Granat.
carlosiana, *C. DC. l. c.* 140.—Venezuela.
carthaginensis, *C. DC. in Linnaea*, xxxvii. (1871–73) 377.—Costa Rica.
casapiana, *C. DC. in Journ. Bot.* iv. (1866) 146.—Peruv.
Casarettii, *C. DC. l. c.* 141.—Bras.
Casimiri, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 111.—Ins. Martinic.
catharina, *Miq. Syst. Pip.* 127.—Bras.
caulibarbis, *Miq. l. c.* 98.—Bras.
ceylanica, Miq. in Hook. Lond. Journ. Bot. v. (1846) 550 = Wightiana.
chachapoyasensis, *C. DC. in DC. Prod.* xvi. I. 449.—Peruv.
chilensis, *C. DC. in Journ. Bot.* iv. (1866) 144.—Chili.
choroniana, *C. DC. l. c.* 141.—Venezuela.
chrysotricha, *Miq. Syst. Pip.* 163.—Peruv.
ciliaris, *C. DC. in Journ. Bot.* iv. (1866) 144.—N. Granat.
ciliata, *H. B. & K. Nov. Gen. et Sp.* i. 68.—Mexic.
ciliata, Mor. ex C. DC. in DC. Prod. xvi. I. 411 = lanceolato-peltata.
ciliolata, *Miq. in Hook. Lond. Journ. Bot.* vi. (1847) 459.—Mexic.
ciliosa, *C. DC. in Journ. Bot.* iv. (1866) 144.—N. Granat.
circinnata, *Link, Bot. Jahrb.* i. III. (1820) 64.—Bras.
circularis, *Hensch. in Nov. Act. Soc. Sc. Upsal.* Ser. III. viii. n. VIII. (1873) 24.—Bras.
claytonioides, *Kunth, Ind. Sem. Hort. Berol.* (1847) 11.—Guatemala.
clusiaefolia, Hook. Bot. Mag. t. 2943 = obtusifolia.
colorata, *H. B. & K. Nov. Gen. et Sp.* i. 64. t. 11.—N. Granat.
columbiana, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 429.—N. Granat.
Commersonii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 479.—Madag.
concava, *Ruiz & Pav. Fl. Per.* i. 31. t. 46. f. C.—Peruv.
concinna, *A. Dietr. Sp. Pl.* i. 164.—Jamaic.
congesta, *H. B. & K. Nov. Gen. et Sp.* i. 68.—Peruv.
conjugata, *H. B. & K. l. c.* 63.—Peruv.
consoquitlana, *C. DC. in Linnaea*, xxxvii. (1871–73) 379.—Mexic.
convexa, *Miq. Syst. Pip.* 107.—Java.
Cookiana, *C. DC. in DC. Prod.* xvi. I. 450.—Ins. Sandvic.
corcovadensis, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 187.—Bras.
cordifolia, *A. Dietr. Sp. Pl.* i. 154.—Jamaic.
cordifolia, Opiz, in Presl, Rel. Haenk. i. 164 = dependens.
cordovana, *C. DC. in Linnaea*, xxxvii. (1871–73) 374.—Mexic.
cordulata, *C. DC. in Journ. Bot.* iv. (1866) 137.—Panama.
coriacea, *Loud. Hort. Brit.* 13.—Ind. or.
Cotyledon, *Benth. Pl. Hartw.* 145.—N. Granat.
Coulteri, *C. DC. in DC. Prod.* xvi. I. 424.—Mexic.
courtallensis, *Miq. in Hook. Lond. Journ. Bot.* v. (1846) 549.—Ind. or.
crystallina, *Ruiz & Pav. Fl. Per.* i. 30.—Peruv.
cubana, *C. DC. in DC. Prod.* xvi. I. 413.—Cuba.
cubensis, *C. DC. in Journ. Bot.* iv. (1866) 142.—Cuba.
cuneata, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 429.—Ins. S. Vincent.
cuneifolia, A. Dietr. Sp. Pl. i. 146 = obtusifolia.
cyclophylla, *C. DC. in DC. Prod.* xvi. I. 444.—Peruv.
cyclophylla, *Miq. in Mart. Fl. Bras.* iv. I. 219.—Am. trop.
dasystachya, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 416.—Peruv.
Decaisnei, *Miq. Syst. Pip.* 133.—Ins. Borbon.
decurrens, *C. DC. in Journ. Bot.* iv. (1866) 143.—Venezuela.
defoliata, *C. DC. l. c.* 138.—N. Granat.
delicatula, *Hench. in Nov. Act. Soc. Sc. Upsal.* Sér. III. viii. n. VIII. (1873) 22.—Hab.?
demissa, *C. DC. in DC. Prod.* xvi. I. 423.—Venezuela.
dendrophila, Griseb. Cat. Pl. Cub. 64 = guadaloupensis, alata.
dendrophila, *Schlecht. in Linnaea*, v. (1830) 74.—Mexic.

PEPEROMIA :—
dependens, *Ruiz & Pav. Fl. Per.* i. 33. t. 48. f. a.—Peruv.
Deppeana, *Schlecht. & Cham. in Linnaea*, v. (1830) 75.—Mexic.
diaphana, *Miq. in Arch. Néerl.* vi. (1871) 170; *et in Verh. Kon. Acad. Wetensch.* II. v. (1871) 232.—Bras.
dichotoma, *Regel, in Bull. Soc. Nat. Mosc.* xxxi. (1858) II. 545.—Bras.
diffusa, *C. DC. in Journ. Bot.* iv. (1866) 133.—Venezuela.
dindygulensis, *Miq. Syst. Pip.* 122.—Ind. or.
discolor, *C. DC. in DC. Prod.* xvi. I. 463.—Jamaic.
discolor, *Loud. Hort. Brit.* 13.—Am. austr.
dissimilis, H. B. & K. Nov. Gen. et Sp. i. 67 = blanda.
distachya, *A. Dietr. Sp. Pl.* i. 156.—Am. austr.
distachya, Griseb. Cat. Pl. Cub. 64 = producta, cubensis.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* i. 64.—Peruv.
Doelli, *Phil. Fl. Atac.* 49.—Chili.
dolabriformis, *H. B. & K. Nov. Gen. et Sp.* i. 60. t. 4.—Peruv.
dominicana, *C. DC. in Journ. Bot.* iv. (1866) 149.—Ins. S. Doming.
donaguiana, *C. DC. in Linnaea*, xxxvii. (1871–73) 382.—Mexic.
drusophila, *C. DC. l. c.* 372.—Mexic.
eburnea, *Linden, Cat.* n. 87 (1871) 5.—N. Granat.
edulis, *Miq. in Linnaea*, xviii. (1844) 711.—Mexic.
elegans, *C. DC. in DC. Prod.* xvi. I. 430.—Peruv.
elliptica, *A. Dietr. Sp. Pl.* i. 167.—Ins. Borbon.
elliptica, Link, Bot. Jahrb. i. III. (1820) 64 = mauritiana.
ellipticifolia, *C. DC. in DC. Prod.* xvi. I. 463.—Venezuela.
elliptico-rhombea, *C. DC. l. c.* 407.—Peruv.
elongata, *H. B. & K. Nov. Gen. et Sp.* i. 62.—Venezuela.
elongata, Miq. in Linnaea, xx. (1847) 127 = japurensis.
emarginata, *Ruiz & Pav. Fl. Per.* i. 33. t. 49. f. a.—Peruv.
emarginella, *C. DC. in DC. Prod.* xvi. I. 437.—Ins. Guadaloup.
emarginulata, *C. DC. l. c.* 433.—Peruv.
Endlicheri, *Miq. Syst. Pip.* 102.—Ins. Norfolk.
epidendron, *C. DC. in Linnaea*, xxxvii. (1871–73) 379.—Mexic.
Erasmia, *C. DC. in DC. Prod.* xvi. I. 397.—Mexic.
Ernstiana, *C. DC. l. c.* 422.—Venezuela.
estrellensis, *C. DC. l. c.* 421.—Bras.
exigua, Griseb. Cat. Pl. Cub. 63 = minima.
exigua, *Miq. Syst. Pip.* 77.—Java.
exilis, *Griseb. Fl. Brit. W. Ind.* 164.—Ind. occ.
exilis, *Miq. in Arch. Néerl.* vii. (1871) 169; *et in Verh. Kon. Akad. Wetensch.* II. v. (1871) 231.—Bras.
Fendleriana, *C. DC. in DC. Prod.* xv. i. 460.—Venezuela.
Fenzlei, *Regel, in Act. Hort. Petrop.* vi. (1879) 293.—Hab.?
fernandeziana, *Miq. Syst. Pip.* 139.—Ins. J. Fernand.
fernandopoana, C. DC. in DC. Prod. xvi. I. 397 = seq.
fernandopoiana, *C. DC. in Journ. Bot.* iv. (1866) 134.—Afr. trop.
filiformis, A. Dietr. Sp. Pl. i. 177 = obtusifolia.
filiformis, *Ruiz & Pav. Fl. Per.* i. 31. t. 47. f. b.—Peruv.
fimbriata, *Miq. Syst. Pip.* 178.—Bras.
flagelliformis, *Hook. f. ex Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 423; *et in Trans. Linn. Soc.* xx. (1846) 181.—Ins. Galapag.
flavescens, *C. DC. in DC. Prod.* xvi. I. 436.—Peruv.
flexicaulis, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 240; *et Bot. Ergeb. Waldem. Reise*, 134. t. 19. f. A.—Bras.
foliiflora, *Ruiz & Pav. Fl. Per.* i. 30. t. 45. f. c.—Peruv.
foliosa, *H. B. & K. Nov. Gen. et Sp.* i. 63.—N. Granat.
fontinalis, *A. Dietr. Sp. Pl.* i. 176.—Bras.
fragrans, *C. DC. in Journ. Bot.* iv. (1866) 140.—Venezuela.
Fraseri, *C. DC. l. c.* 134.—Ecuador.
freireaefolia, *A. Rich. Tent. Fl. Abyss.* ii. 274.—Abyssin.

PEPEROMIA:—
fugax, *C. DC. in Linnaea*, xxxvii. (1871–73) 370.—Mexic.
furcata, *Opiz, in Presl, Rel. Haenk.* i. 164.—Peruv.
galapagensis, *Hook. f. ex Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 426.—Ins. Galap.
Galeottiana, *Hook. Ic. Pl.* t. 327.—Mexic.
Galeottii, *Miq. Syst. Pip.* 190.—Mexic.
galioides, *H. B. & K. Nov. Gen. et Sp.* i. 71. t. 17.—N. Granat.
Gardneriana, *Miq. Syst. Pip.* 73.—Bras.
Gaudichaudii, *Miq. l. c.* 137.—Ins. Sandvic.
gemella, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 83, 85.—Java.
glabella, *A. Dietr. Sp. Pl.* i. 156.—Jamaic.
glabella, Griseb. Fl. Brit. W. Ind. 165 = nigro-punctata.
glabra, *C. DC. in Journ. Bot.* iv. (1866) 143.—Venezuela.
Glaziovi, *C. DC. in Linnaea*, xxxvii. (1871–73) 380.—Bras.
globulifera, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 93; *Contrib.* 103.—Paraguay.
Goudotii, *Miq. Syst. Pip.* 133.—Ins. Borbon.
grandifolia, A. Dietr. Sp. Pl. i. 166 = Piper subpeltatum.
Grisebachii, *C. DC. in Linnaea*, xxxvii. (1871–73) 382.—Ind. occ.
guadaloupensis, *C. DC. in Journ. Bot.* iv. (1866) 139.—Mexic.
guayaquilensis, *C. DC. in DC. Prod.* xvi. 1. 400.—Ecuador.
Guildingiana, A. Dietr. Sp. Pl. i. 162 = scandens.
Haenkeana, *Opiz, in Presl, Rel. Haenk.* i. 162.—Peruv.
Hahnii, *C. DC. in Linnaea*, xxxvii. (1871–73) 368.—Ins. Martinic.
Hamiltoniana, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 417.—Jamaic.
Hartwegiana, *Miq. l. c.* 425.—N. Granat.
Haworthiana, *A. Dietr. Sp. Pl.* i. 171.—Ind. or. ?
Haworthii, Sweet, Hort. Brit. ed. II. 482 = praec.
Haycockii, *C. DC. in DC. Prod.* xvi. 1. 404.—Zeylan.
hederacea, *Miq. in Mart. Fl. Bras.* iv. 1. 20.—Bras.
hemionitidifolia, *Ham. Prod. Pl. Ind. Occ.* 2.—Ins. S. Doming.
Herminieri, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 316.—Ins. Guadeloup.
hernandiaefolia, *A. Dietr. Sp. Pl.* i. 157; *Miq. Syst. Pip.* 72.—Ind. occ.
hernandiaefolia, Griseb. Fl. Brit. W. Ind. 166; Cat. Pl. Cub. 64 = Ponthieui.
Hesperomannii, *Wawra, in Flora*, lviii. (1875) 227.—Ins. Sandvic.
heterophylla, Griseb. ex C. DC. in DC. Prod. xvi. 1. 456 = agapatensis.
heterophylla, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 415.—Peruv.
heterostachya, *A. Dietr. Sp. Pl.* i. 717 (Index).—Peruv.
Heyneana, *Miq. Syst. Pip.* 123.—Ind. or.
Heyneana, Wight, Ic. t. 1922 = dindigulensis.
Hilariana, *Miq. Syst. Pip.* 89.—Bras.
hircina, *C. DC. in DC. Prod.* xvi. 1. 415.—Ins. Borbon.
hirsuta, *Miq. Syst. Pip.* 143.—Bras.
hirta, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 20.—Ins. Rodriguez.
hirta, *C. DC. in DC. Prod.* xvi. 1. 412.—Cuba.
hirtella, Griseb. Cat. Pl. Cub. 64 = hirta, *C. DC.*
hirtella, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 414.—Ins. Dominic.
hispidula, *A. Dietr. Sp. Pl.* i. 165.—Ind. occ.
Hoffmannii, *C. DC. in Journ. Bot.* iv. (1866) 133.—Costa Rica.
huatoscoana, *C. DC. in Linnaea*, xxxvii. (1871–73) 383.—Mexic.
humilis, *A. Dietr. Sp. Pl.* i. 168.—Ins. S. Crucis.
hydrocotyloides, *Miq. in Linnaea*, xx. (1847) 118. t. 1. f. 1.—Bras.
hymenophylla, Miq. Syst. Pip. 78 = exigua.
hypoleuca, *Miq. l. c.* 136.—Ins. Sandvic.
inaequalifolia, *Ruiz & Pav. Fl. Per.* i. 30. t. 46. f. a.—Peruv.
incana, *A. Dietr. Sp. Pl.* i. 166.—Bras.
increscens, Miq. in Linnaea, xx. (1847) 124 = Langsdorffii.
insularum, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 422.—Ins. Sandvic.
ionophylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 175.—Ind. occ.; Venezuela.

PEPEROMIA:—
Jamesoni, *Regel, in Bull. Soc. Nat. Mosc.* xxxi. (1858) II. 544.—Ecuador.
Jamesoniana, *C. DC. in Journ. Bot.* iv. (1866) 137; *in DC. Prod.* xvi. 1. 412.—Ecuador.
japurensis, *C. DC. in DC. Prod.* xvi. 1. 407.—Bras.
jarisiana, *C. DC. in Linnaea*, xxxvii. (1871–73) 382.—Costa Rica.
javanica, *Miq. Pl. Jungh.* 15.—Java.
Junghuhniana, *Miq. l. c.*—Java.
Klotzschiana, *Miq. Syst. Pip.* 145.—Peruv.
Knoblecheriana, Schott, in Bot. Zeit. ix. (1851) 225 = Verhuellia Knoblecheriana.
kravangensis, *Miq. Syst. Pip.* 108.—Java.
Kuhliana, *Miq. l. c.* 104.—Java.
Kunthiana, *C. DC. in DC. Prod.* xvi. 1. 464.—Ecuador.
laevifolia, *Miq. Syst. Pip.* 107.—Java.
laevis, *C. DC. in Journ. Bot.* iv. (1866) 139.—Venezuela.
lanata, A. Dietr. Sp. Pl. i. 145 = Piper caninum.
lanceolata, *C. DC. in Journ. Bot.* iv. (1866) 145.—Ecuador.
lanceolato-peltata, *C. DC. l. c.* 136.—Costa Rica.
lancifolia, *Hook. Ic. Pl.* t. 332.—Mexic.
Langsdorffii, *Miq. Syst. Pip.* 116.—Bras.
larecajana, *C. DC. in DC. Prod.* xvi. 1. 406.—Bolivia.
latifolia, *Miq. Syst. Pip.* 128.—Ins. Sandvic.
laxiflora, *H. B. & K. Nov. Gen. et Sp.* i. 63. t. 10.—N. Granat.
lenticularis, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 93; *Contrib.* 102.—Paraguay.
leptophylla, *Miq. in Linnaea*, xviii. (1844) 712.—Mexic.
leptostachya, *Chapm. Fl. S. U. St. Suppl.* 645.—Am. bor.
leptostachya, *Hook. & Arn. Bot. Beech. Voy.* 96.—Ins. Sandvic.
leuconeura, Hort. Berol. ex C. DC. in DC. Prod. xvi. 1. 417 = trinervis.
Liebmannii, *C. DC. in Linnaea*, xxxvii. (1871–73) 383.—Mexic.
lignescens, *C. DC. in Journ. Bot.* iv. (1866) 137.—Costa Rica.
linatifolia, Rich. ex C. DC. in DC. Prod. xvi. 1. 410 = caulibarbis.
Lindeniana, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 59.—Mexic.
Lindleyana, Miq. Syst. Pip. 147 (err. typ. 247) = Perrottetiana.
linearis, *C. DC. in Journ. Bot.* iv. (1866) 145.—Venezuela.
lineatipila, *A. Rich. Fl. Cub. Fanerog.* ii. 229.—Cuba.
longepedunculata, *Opiz, in Presl, Rel. Haenk.* i. 164.—Peruv.
longiflora, *Link, ex A. Dietr. Sp. Pl.* i. 174.—Hab. ?
longifolia, *C. DC. in DC. Prod.* xvi. 1. 405.—Guian.
loxensis, *H. B. & K. Nov. Gen. et Sp.* i. 70.—Ecuador.
lunaria, *Ham. Prod. Pl. Ind. Occ.* 2.—Ins. S. Doming.
lunaris, A. Dietr. Sp. Pl. i. 167 = praec.
Lundii, *C. DC. in Linnaea*, xxxvii. (1871–73) 367.—Bras.
Lyalli, *C. DC. in Journ. Bot.* iv. (1866) 140.—Madag.
Macraeana, *C. DC. l. c.* 145.—Ins. Sandvic.
macropoda, *Miq. in Linnaea*, xx. (1847) 128.—Bras.
macrorhiza, *H. B. & K. Nov. Gen. et Sp.* i. 72.—Peruv.
macrostachya, *A. Dietr. Sp. Pl.* i. 149.—Guian.
macrothyrsa, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 419.—Peruv.
maculosa, *Hook. Exot. Fl.* t. 92.—Ins. S. Doming.
madagascariensis, *C. DC. in DC. Prod.* xvi. 1. 448.—Madag.
magnoliaefolia, A. Dietr. Sp. Pl. i. 153 = tithymalioides.
major, C. DC. in DC. Prod. xvi. 1. 432 = hederacea.
mandioccana, *Miq. in Linnaea*, xx. (1847) 125.—Bras.
Mandonii, *C. DC. in DC. Prod.* xvi. 1. 395.—Bolivia.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 217.—Afr. trop.
margaritifera, *Bert. ex Hook. Ic. Pl.* t. 91.—Ins. J. Fernand.
marginata, *Miq. Syst. Pip.* 129.—Ins. Borbon.
mariannensis, *C. DC. in DC. Prod.* xvi. 1. 442.—Ins. Marian.
maribiana, *C. DC. l. c.* 428.—Bras.

PEPEROMIA:—
marmorata, *Hook. f. Bot. Mag.* t. 5568.—Bras.
Martiana, Miq. in Arch. Néerl. iv. (1871) 173 = caldesiana.
Martiana, *Miq. Syst. Pip.* 189.—Bras.
mascharena, *C. DC. in Journ. Bot.* iv. (1866) 135.—Madag.
Mathewsiana, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 423.—Peruv.
Mathewsii, *C. DC. in Journ. Bot.* iv. (1866) 144.—Peruv.
matlalucaënsis, *C. DC. in Linnaea*, xxxviii. (1871–73) 375.—Mexic.
mauiensis, *Wawra, in Flora*, lviii. (1875) 225.—Ins. Sandvic.
mauritiana, *A. Dietr. Sp. Pl.* i. 171.—Ins. Maurit.
maypurensis, *H. B. & K. Nov. Gen. et Sp.* i. 66.—Venezuela.
melanostigma, *Miq. Syst. Pip.* 90.—Costa Rica.
membranacea, *Hook. & Arn. Bot. Beech. Voy.* 96.—Ins. Sandvic.
membranacea, Miq. Syst. Pip. 120 = mariannensis.
Menkeana, Miq. in Linnaea, xx. (1847) 124 = galioides.
metapalcoënsis, C. DC. in DC. Prod. xvi. 1. 416 = rotundifolia.
mexicana, *Miq. Syst. Pip.* 75.—Mexic.
microphylla, *H. B. & K. Nov. Gen. et Sp.* i. 69.—N. Granat.
minensis, *Hensch. in Nov. Act. Soc. Sc. Upsal.* Ser. III. viii. n. VIII. (1873) 29.—Bras.
minima, *C. DC. in DC. Prod.* xvi. 1. 399.—Cuba.
Miqueliana, *C. DC. in Journ. Bot.* iv. (1866) 138.—Ecuador.
Miqueliana, Griseb. Cat. Pl. Cub. 63 = metapalcoënsis.
miradoresiana, *C. DC. in DC. Prod.* xvi. 1. 395.—Mexic.
Moerenhouti, *C. DC. l. c.* 458.—Ins. Tahiti.
mollis, *H. B. & K. Nov. Gen. et Sp.* i. 65.—N. Granat.
Monini, *C. DC. in DC. Prod.* xvi. 1. 444.—Ins. Borbon.
monostachya, *Ruiz & Pav. Fl. Per.* i. 33.—Peruv.
monsterifolia, Griseb. Cat. Pl. Cub. 64 = septuplinervia.
montana, *C. DC. in DC. Prod.* xvi. 1. 426.—Peruv.
monticola, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 217.—Afr. trop.
monticola, *Miq. Syst. Pip.* 71.—Mexic.
Moritzii, *C. DC. in DC. Prod.* xvi. 1. 420.—Venezuela.
moulmeiniana, *C. DC. in Journ. Bot.* iv. (1866) 140.—Burma.
Muelleri, *C. DC. in Linnaea*, xxxvii. (1871–73) 366.—Mexic.
multiplex, Lepr. ex C. DC. in DC. Prod. xvi. 1. 406 = longifolia.
muscophila, *C. DC. in Journ. Bot.* iv. (1866) 133.—Mexic.
muscosa, *Link, Bot. Jahrb.* i. III. 64 (1820).—Bras.
myosuroides, *A. Dietr. Sp. Pl.* i. 157.—Guian.
Myosurus, Willd. ex A. Dietr. l. c. 160 = scandens.
myriocarpa, Miq. in Linnaea, xx. (1847) 128 = maribiana.
myriocarpa, *Miq. Syst. Pip.* 185.—Bras.
myrtifolia, *A. Dietr. Sp. Pl.* i. 147; *Miq. Syst. Pip.* 92.—Ins. S. Crucis.
Myrtillus, *Miq. l. c.* 154.—Venezuela.
nana, *C. DC. in Journ. Bot.* iv. (1866) 135.—Ins. Comoro.
naraugoana, *C. DC. in Linnaea*, xxxvii. (1871–73) 378.—Costa Rica.
nematostachya, *Link, Bot. Jahrb.* i. III. (1820) 63.—Bras.
nemorosa, *C. DC. in DC. Prod.* xvi. 1. 415.—Jamaic.
nemostachya, A. Dietr. Sp. Pl. i. 156 = nematostachya.
nervoso-venosa, *Miq. Syst. Pip.* 104.—Java.
nigro-punctata, *Miq. l. c.* 188.—Ins. Martinic.
nivalis, *Miq. l. c.* 112.—Peruv.
nuda, *C. DC. in Linnaea*, xxxvii. (1871–73) 368.—Bras.
nummularifolia, *H. B. & K. Nov. Gen. et Sp.* i. 66.—Am. trop.
nummularioides, *Griseb. in Goett. Abh.* vi. (1854) 132.—Chili.
oahuensis, *C. DC. in DC. Prod.* xvi. 1. 422.—Ins. Sandvic.
obcordata, *Presl, Epim. Bot.* 223.—Peruv.
obliqua, *Ruiz & Pav. Fl. Per.* i. 32. t. 51. f. c.—Peruv.
obtusifolia, *A. Dietr. Sp. Pl.* i. 154.—Am. trop.
obtusifolia, A. Dietr. l. c., partim = magnoliaefolia.

PEPEROMIA:—
obtusifolia, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 429 = emarginulata.
obversa, *A. Dietr. Sp. Pl.* i. 173 ; *Griseb. Fl. Brit. W. Ind.* 168.—Ind. occ.
obversa, C. DC. in DC. Prod. xvi. 1. 415 = Grisebachii.
Oerstedi, *C. DC. in Linnaea*, xxxvii. (1871–73) 375.—Costa Rica.
oleracea, Poepp. ex Miq. in Mart. Fl. Bras. iv. 1. 10 = pellucida.
olivacea, *C. DC. in Journ. Bot.* iv. (1866) 146.—Costa Rica.
Opiziana, A. Dietr. Sp. Pl. i. 162 = dependens.
orbicularis, *C. DC. in DC. Prod.* xvi. 1. 465.—Bras.
orbiculata, *A. Dietr. Sp. Pl.* i. 180.—Am. bor.
oreophila, *Hensch. in Nov. Act. Soc. Sc. Upsal.* Ser. III. viii. n. 8 (1873) 28.—Bras.
Ottoniana, *Kunth, ex Miq. Syst. Pip.* 100.—Mexic.
ovalifolia, *Hook. Exot. Fl.* t. 165.—Ins. S. Vincent.
ovato-peltata, *C. DC. in Journ. Bot.* iv. (1866) 132.—Costa Rica.
oxycarpa, *C. DC. in Linnaea*, xxxvii. (1871–73) 377.—Mexic.
pachyphylla, *Miq. Syst. Pip.* 137.—Ins. Sandvic.
pallens, *H. B. & K. Nov. Gen. et Sp.* i. 69.—N. Granat.
pallescens, *Miq. in Hook. Lond. Journ. Bot.* vi. (1847) 460 [bis].—Guatemala.
pallida, *A. Dietr. Sp. Pl.* i. 153.—Ins. Societ.
pallida, Seem. Fl. Vit. 259 = vitiana.
paniculata, *Regel, in Bull. Soc. Nat. Mosc.* xxxi. (1858) II. 542.—Bras.
papantlacensis, *C. DC. in Linnaea*, xxxvii. (1871–73) 366.—Mexic.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 94 ; *Contrib.* 104.—Paraguay.
parasitica, *C. DC. in DC. Prod.* xvi. 1. 421.—Ecuador.
Parkeriana, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 427.—Guian.
parnassiaefolia, *Miq. in Linnaea*, xx. (1847) 119.—Bras.
parvifolia, *C. DC. in Journ. Bot.* iv. (1866) 133.—Peruv. ; Boliv.
patula, *C. DC. l. c.* 139.—Venezuela.
Pavoniana, *C. DC. in DC. Prod.* xvi. 1. 460.—Peruv.
pedunculata, *C. DC. in Journ. Bot.* iv. (1866) 146.—Ins. Borbon.
peepuloides, A. Dietr. Sp. Pl. i. 145 = Piper peepuloides.
pellucida, *H. B. & K. Nov. Gen. et Sp.* i. 64.—Am. et Afr. trop.
peltata, A. Dietr. Sp. Pl. i. 142 = Piper peltatum.
peltoidea, *H. B. & K. Nov. Gen. et Sp.* i. 61. t. 7.—Venezuela.
pendula, *Willd. ex Schult. Mant.* i. 372 (Index).—Am. merid.
penninervia, *Baker, Fl. Maurit.* 298.—Ins. Mascar.
peploides, *H. B. & K. Nov. Gen. et Sp.* i. 69.—Ecuador.
pereskiaefolia, *H. B. & K. l. c.* 68.—Venezuela.
perforata, *Opiz, in Presl, Rel. Haenk.* i. 162.—Mexic.
pernambucensis, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 420.—Bras.
Perrottetiana, *Miq. Syst. Pip.* 146.—Ins. Maurit.
Perrottetiana, Sieber, ex C. DC. in DC. Prod. xvi. 1. 441 = marginata.
petiolaris, *C. DC. in Journ. Bot.* iv. (1866) 138.—Costa Rica.
petiolata, *Hook. f. in Trans. Linn. Soc.* xx. (1846) 181.—Ins. Galap.
petrophila, *C. DC. in Linnaea*, xxxvii. (1871–73) 369.—Mexic.
phyllantha, Opiz, in Presl, Rel. Haenk. i. 162 = foliiflora.
Pichinchae, *C. DC. in Journ. Bot.* iv. (1866) 144.—Ecuador.
pilosa, *Ruiz & Pav. Fl. Per.* i. 32. t. 50. f. a.—Peruv.
pilosula, *C. DC. in Linnaea*, xxxvii. (1871–73) 378.—Bras.
piperea, *C. DC. in Journ. Bot.* iv. (1866) 143.—Guian.
plantaginifolia, Opiz, in Presl, Rel. Haenk. i. 164 = Piper chilense.
plicata, *Opiz, l. c.* 163.—Mexic.
Poeppigii, *Miq. Syst. Pip.* 84.—Peruv.
polybotrya, *H. B. & K. Nov. Gen. et Sp.* i. 60.—Peruv.

PEPEROMIA:—
polystachya, *Hook. Exot. Fl.* t. 23.—Ind. occ.
Ponthieui, *Miq. Syst. Pip.* 186.—Ind. occ.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 147.—Panama.
portulacaefolia, *H. B. & K. Nov. Gen. et Sp.* i. 70.—Peruv.
portulacoides, *A. Dietr. Sp. Pl.* i. 172.—Ins. Maurit.
procumbens, *C. DC. in Journ. Bot.* iv. (1866) 141.—Peruv.
producta, *Griseb. Fl. Brit. W. Ind.* 166.—Ind. occ.
prostrata, *Hort. Williams, ex Gard. Chron.* (1879) i. 716 ; *et in Illustr. Hortic.* xxvi. (1879) 113.—N. Granat. ?
pruinosa, Kunth, Syn. Pl. Aequin. i. 124 = Piper peltatum.
Pseudoamplexicaulis, *C. DC. in DC. Prod.* xvi. 1. 432.—Mexic.
Pseudodependens, *C. DC. in Journ. Bot.* iv. (1866) 137.—Venezuela.
Pseudodindygulensis, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 307.—Paraguay.
Pseudofurcata, *C. DC. in DC. Prod.* xvi. 1. 399.—Peruv.
Pseudopeltoidea, *C. DC. l. c.* 434.—Venezuela.
Pseudopereskiaefolia, *C. DC. l. c.* 448.—Cuba.
Pseudorhombea, *C. DC. l. c.* 440.—Zeylan.
Pseudorhynchophoron, *C. DC. l. c.* 425.—Peruv.
Pseudovariegata, *C. DC. l. c.* 428.—N. Granat.
pterocaulis, Miq. Syst. Pip. 86 = alata.
puberula, *Baker, in Saund. Refug. Bot.* t. 302.—Guatemala.
pubescens, *Ruiz & Pav. Fl. Per.* i. 31.—Peruv.
pubifolia, *Hort. Veitch, in Proc. Hort. Soc.* v. (1865) 143.—Hab. ?
pulchella, *A. Dietr. Sp. Pl.* i. 177.—Jamaic.
pulicaris, Opiz, in Presl, Reliq. Haenk. i. 162. t. 30. f. 1 = scandens.
pumila, *Opiz, l. c.* 163. t. 30. f. 3.—Peruv.
punctulata, *Regel, in Bull. Soc. Nat. Mosc.* xxxi. (1858) II. 543.—Bras.
purpurascens, *C. DC. in DC. Prod.* xvi. 1. 417.—Ins. Sandvic.
purpurascens, Mor. ex C. DC. l. c. 420 = trinervula.
purpurea, *Ruiz & Pav. Fl. Per.* i. 32. t. 49. f. b.—Peruv.
purpurella, *C. DC. in DC. Prod.* xvi. 1. 414.—Venezuela.
pyrifolia, *Humb. & Bonpl. in Pl. Aequin.* i. 61.—Mexic.
quadrangularis, *A. Dietr. Sp. Pl.* i. 169.—Ins. Trinit.
quadrangularis, Griseb. Fl. Brit. W. Ind. 168 = muscosa.
quadrifolia, *H. B. & K. Nov. Gen. et Sp.* i. 69.—Am. trop.
quadrifolia, Miq. in Linnaea, xx. (1847) 571 = trineura.
quadrifolia, Mus. Par. ex Miq. Syst. Pip. 169 = reflexa.
quaternata, *Miq. in Seem. Bot. Voy. Herald*, 197.—Panama.
quintuplinervia, C. DC. in DC. Prod. xvi. 1. 470 (Index) = Pichinchae ?
quitensis, Benth. Pl. Hartw. 253 = blanda.
quitensis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 421 = ciliata.
radicans, *C. DC. in Mém. Soc. Phys. Genèv.* xxvii. II. (1881) 308.—Paraguay.
ramulosa, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 158.—Ins. Galap.
recurvata, *Miq. Syst. Pip.* 141.—Java.
reflexa, *A. Dietr. Sp. Pl.* i. 180.—Ind. occ. ; Afr. austr.
reflexa, H. B. & K. Nov. Gen. et Sp. i. 70 = fontinalis.
Regelii, *C. DC. in DC. Prod.* xvi. 1. 455.—Bras.
Regnelliana, *C. DC. in Linnaea*, xxxvii. (1871–73) 381.—Bras.
Regnelliana, *Miq. in Arch. Néerl.* vi. (1871) 173 ; *et in Verh. Kon. Akad. Wetensch.* II. v. (1871) 235.—Bras.
Reinwardtiana, *Miq. Syst. Pip.* 141.—Borneo.
reniformis, A. Dietr. Sp. Pl. i. 162 = Verhuellia elegans.
reniformis, Hook. Exot. Fl. t. 164 = scandens.
repens, H. B. & K. Nov. Gen. et Sp. i. 65 = scandens.
reptans, *C. DC. in Journ. Bot.* iv. (1866) 141.—N. Granat.
resedaeflora, *Linden & André, Illustr. Hortic.* xvii. (1870) t. 26.—N. Granat.

PEPEROMIA:—
reticulata, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 21 ; *et in Baker, Fl. Maurit.* 297.—Ins. Mascar.
retusa, *A. Dietr. Sp. Pl.* i. 155.—Afr. austr.
rhexiaefolia, *Mor. ex C. DC. in DC. Prod.* xvi. 1. 460.—Venezuela.
rhombea, *Ruiz & Pav. Fl. Per.* i. 31. t. 46. f. c.—Peruv.
rhomboidea, *Hook. & Arn. Bot. Beech. Voy.* 70.—Ins. Societ.
Riedeliana, *Regel, Ind. Sem. Hort. Petrop.* (1858) 35 ; *et in Gartenfl.* (1859) 229. t. 265.—Bras.
rigida, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 94 ; *Contrib.* 104.—Paraguay.
rodriguesiana, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 21.—Ins. Mascar.
Rodriguezi, Balf. f. in Phil. Trans. clxviii. (1879) 370 = praec.
Rollingo, *Link, Bot. Jahrb.* i. III. (1820) 65.—Hab. ?
rostrata, *A. Dietr. Sp. Pl.* i. 153.—Ins. Molucc.
rotundata, *H. B. & K. Nov. Gen. et Sp.* i. 67. t. 12.—N. Granat.
rotundifolia, Griseb. Cat. Pl. Cub. 63 = circinata.
rotundifolia, H. B. & K. Nov. Gen. et Sp. i. 65 = nummulariifolia.
rotundifolia, *Schlecht. & Cham. in Linnaea*, vi. (1831) 353.—Mexic.
Roxburghiana, *A. Dietr. Sp. Pl.* i. 144.—Malaya.
rubella, *Hook. Exot. Fl.* t. 58.—Am. austr.
rubella, Hook. & Arn. ex C. DC. in DC. Prod. xvi. 1. 380 = Piper purpurascens.
rubioides, *H. B. & K. Nov. Gen. et Sp.* i. 70.—Mexic.
rubricaulis, A. Dietr. Sp. Pl. i. 182 = pereskiaefolia.
rubrinodis, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 12.—Guatemala.
rufescens, *C. DC. in DC. Prod.* xvi. 1. 416.—Peruv.
Rupertiana, *C. DC. l. c.* 413.—Ins. Dominic.
rupestris, *H. B. & K. Nov. Gen. et Sp.* i. 62.—Mexic.
rupestris, Miq. in Linnaea, xx. (1847) 127 = nematostachya.
rupestris, Miq. Syst. Pip. 183, partim = distachya.
ruscifolia, *H. B. & K. Nov. Gen. et Sp.* i. 63.—Venezuela.
saligna, *H. B. & K. l. c.* 62.—N. Granat.
san-carlosiana, *C. DC. in Journ. Bot.* iv. (1866) 138.—Venezuela.
Sandersii, *C. DC. in DC. Prod.* xvi. 1. 400.—Bras.
sandwicensis, *Miq. Syst. Pip.* 126.—Ins. Sandvic.
san-joseana, *C. DC. in Linnaea*, xxxvii. (1871–73) 372.—Costa Rica.
sarmentosa, A. Dietr. Sp. Pl. i. 161 = Piper sarmentosum.
saxatilis, *A. Dietr. l. c.* 178.—Reg. Himal.
scandens, *Ruiz & Pav. Fl. Per.* i. 32. t. 51. f. b.—Peruv.
Schiedeana, Schlecht. in Linnaea, vii. (1832) 138 = tenerrima.
Schneepeana, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 95 ; *Contrib.* 104.—Paraguay.
Schomburgkii, *C. DC. in DC. Prod.* xvi. 1. 395.—Guiana.
scutata, A. Dietr. Sp. Pl. i. 143 = Piper peltatum.
scutellaefolia, *Ruiz & Pav. Fl. Per.* i. 29. t. 44. f. b.—Peruv.
secunda, *Ruiz & Pav. l. c.* 30. t. 45. f. a.—Peruv.
Seemanniana, *Miq. in Seem. Bot. Voy. Herald*, 198. t. 37.—Panama.
Sellowiana, Miq. Syst. Pip. 161 = Deppeana.
semiruensis, *C. DC. in DC. Prod.* xvi. 1. 467.—Malaya.
septemnervis, *Ruiz & Pav. Fl. Per.* i. 31. t. 47. f. c.—Peruv.
septuplinervia, *C. DC. in Journ. Bot.* iv. (1866) 142.—Cuba.
serpens, *C. DC. l. c.* 136.—N. Granat.
serpens, Griseb. Cat. Pl. Cub. 63 = subrotundifolia.
serpens, Loud. Hort. Brit. 13 = Verhuellia serpens.
serpentarioides, *Miq. in Linnaea*, xx. (1847) 120.—Bras.
serpyllifolia, *A. Dietr. Sp. Pl.* i. 174.—Ins. Maurit.
sessilifolia, *Humb. & Bonpl. ex Schult. Mant.* i. 372 (Index).—Am. austr.
sidaefolia, A. Dietr. Sp. Pl. i. 141 = Piper subpeltatum.
simplex, *Ham. Prod. Pl. Ind. Occ.* 2.—Jamaic.
spathulata, *Ham. l. c.*—Jamaic.
speciosa, Kunth, Syn. Pl. Aequin. i. 124 = Piper peltatum.
spectabilis, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 417.—Peruv.

PEPEROMIA:—
Spruceana, *Benth. ex C. DC. in DC. Prod.* xvi. 1. 402.—Bras.
Sprucei, *C. DC. in Journ. Bot.* iv. (1866) 133.—Peruv.
stelechophila, *C. DC. in DC. Prod.* xvi. 1. 433.—Ecuador.
stellata, *A. Dietr. Sp. Pl.* i. 175.—Jamaic.
stenocarpa, *Regel, Ind. Sem. Hort. Petrop.* (1858) 36; *et Gartenfl.* (1859) 291. t. 271.—Bras.
stolonifera, *H. B. & K. Nov. Gen. et Sp.* i. 71.—N. Granat.
striata, *Ruiz & Pav. Fl. Per.* i. 32. t. 52. f. b.—Peruv.
suaveolens, *Ham. Prod. Pl. Ind. Occ.* 2.—Cuba.
subpeltata, *C. DC. in Journ. Bot.* iv. (1866) 136.—Ecuador.
subpeltata, A. Dietr. Sp. Pl. i. 144 = Piper subpeltatum.
subrotunda, A. Dietr. l. c. 154 = magnoliaefolia.
subrotundifolia, *C. DC. in Journ. Bot.* iv. (1866) 136.—Cuba.
subsessilifolia, *C. DC. in Linnaea*, xxxvii. (1871–73) 375.—Bras.
subvillosa, *C. DC. Heurck & Muell. Arg. in Heurck, Obs. Bot.* 113.—Ins. Martinic.
succulenta, *C. DC. in Journ. Bot.* iv. (1866) 142.—Venezuela.
sumidoriana, *C. DC. in DC. Prod.* xvi. 1. 435.—Bras.
surinamensis, *C. DC. l. c.* 408.—Guian.
Swartziana, Miq. Syst. Pip. 155; Griseb. Fl. Brit. W. Ind. 166 = obtusifolia.
sylvatica, *C. DC. in DC. Prod.* xvi. 1. 435.—Peruv.
sylvestris, *C. DC. in Journ. Bot.* iv. (1866) 142.—Ecuador.
talinifolia, *H. B. & K. Nov. Gen. et Sp.* i. 62. t. 8.—N. Granat.
talinifolia, Link, Bot. Jahrb. i. 111. (1820) 64 = amplexifolia.
tanalensis, *Baker, in Journ. Bot.* xx. (1882) 244.—Madag.
tarapotana, *C. DC. in Journ. Bot.* iv. (1866) 137.—Peruv.
tenella, *A. Dietr. Sp. Pl.* i. 153.—Jamaic.
tenera, Miq. in Mart. Fl. Bras. iv. 1. 19 = hispidula.
tenerrima, *Schlecht. & Cham. in Linnaea*, vi. (1831) 353.—Mexic.
tenuiflora, *Opiz, in Presl, Rel. Haenk.* i. 163.—Mexic.
tenuifolia, *C. DC. in Linnaea*, xxxvii. (1871–73) 371.—Costa Rica.
tetragona, *Ruiz & Pav. Fl. Per.* i. 31. t. 47. f. a.—Peruv.
tetraphylla, Hook. & Arn. Bot. Beech. Voy. 96 = reflexa.
thrichocarpa, C. DC. in DC. Prod. xvi. 1. 455, sphalm. = trichocarpa.
Thwaitesii, C. DC. l. c. 448 = dindygulensis.
tithymaloides, *A. Dietr. Sp. Pl.* i. 152.—Am. trop.
tlapacoyoënsis, *C. DC. in Linnaea*, xxxvii. (1871–73) 376.—Mexic.
tomentosa, *A. Dietr. Sp. Pl.* i. 172.—Java.
tovariana, *C. DC. in DC. Prod.* xvi. 1. 404.—Venezuel.
transparens, *Miq. in Linnaea*, xx. (1847) 121.—Bras.
Trianae, *C. DC. in Journ. Bot.* iv. (1866) 135.—N. Granat.
tricarinata, *A. Dietr. Sp. Pl.* i. 181.—Ind. occ.
trichocarpa, *Miq. Syst. Pip.* 176.—Bras.
trichophylla, *Baker, in Journ. Linn. Soc.* xxi. (1885) 436.—Madag.
trifolia, *A. Dietr. Sp. Pl.* 173.—Guian.
trinervia, Miq. ex C. DC. in DC. Prod. xvi. 1. 470 (Index) = trineura.
trinervis, *Ruiz & Pav. Fl. Per.* i. 32. t. 50. f. b.—Peruv.
trinervula, *C. DC. in DC. Prod.* xvi. 1. 420.—Venezuel.
trineura, *Miq. Syst. Pip.* 175.—Bras.
tristachya, *H. B. & K. Nov. Gen. et Sp.* i. 61. t. 6.—N. Granat.
tristachya, Opiz, in Presl, Rel. Haenk. i. 163 = heterostachya.
tuberosa, *Opiz, l. c.* 164.—Mexic.
turialvensis, *C. DC. in Linnaea*, xxxvii. (1871–73) 380.—Costa Rica.
umbellata, Kunth, Syn. Pl. Aequin. i. 124 = Piper umbellatum.
umbellata, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 427.—Peruv.

PEPEROMIA:—
umbilicata, *Ruiz & Pav. Fl. Per.* i. 30. t. 45. f. b.—Mexic.; Peruv.
undeninervia, *C. DC. in DC. Prod.* xvi. 1. 441.—Peruv.
unguiculata, Steud. Nom. ed. II. ii. 341, in syn., sphalm. = umbilicata.
urocarpa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 42.—Bras.
Urvilleana, A. Rich. Bot. Voy. Astrolab. 356 = Endlicheri.
Vahlii, *A. Dietr. Sp. Pl.* i. 169.—Hab.?
valantoides, Miq. Syst. Pip. 174 = reflexa.
Vanheurcki, *C. DC. ex C. DC. Huerck & Muell. Arg. in Heurck, Obs. Bot.* 116.—Ins. Martinic.
variegata, Ruiz & Pav. Fl. Per. i. 33. t. 52. f. a = maculosa.
variifolia, *C. DC. in DC. Prod.* xvi. 1. 462.—Peruv.
Velloziana, Miq. in Linnaea, xx. (1847) 122 = alata.
Velloziana, *Miq. Syst. Pip.* 88.—Bras.
velutina, *Linden & André, Illustr. Hortic.* xix. (1872) t. 89.—Ecuador.
venezueliana, *C. DC. in Journ. Bot.* iv. (1866) 139.—Venezuela.
Ventenatii, *Miq. Syst. Pip.* 125.—Java.
Verhuellia, *C. DC. in DC. Prod.* xvi. 1. 398.—Bras.
Verschaffeltii, *Lem. Illustr. Hortic.* xvi. (1869) t. 598.—Bras.
verticillata, A. Dietr. Sp. Pl. i. 179 = pulchella.
Victoriana, *C. DC. in DC. Prod.* xvi. 1. 449.—Venezuela.
villosa, *C. DC. in Journ. Bot.* iv. (1866) 135.—Ecuador.
vincentiana, *Miq. in Hook. Lond. Journ.* iv. (1845) 415.—Ins. S. Vincent.
vitiana, *C. DC. in DC. Prod.* xvi. 1. 458.—Ins. Fiji.
Vogelii, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 413.—Afr. trop.
Wallichii, *Miq. Syst. Pip.* 82.—Ind. or.
Warmingii, *C. DC. in Linnaea*, xxxvii. (1871–73) 371.—Bras.
Weddelii, *C. DC. in Journ. Bot.* iv. (1866) 135.—Bras.
Wightiana, *Miq. in Hook. Lond. Journ. Bot.* v. (1846) 548.—Ind. or.
Wightiana, Wight, Ic. t. 1924 = dindygulensis?
Wrightiana, *C. DC. in DC. Prod.* xvi. 1. 431.—Cuba.
zeylanica, Walp. Ann. i. 598 = Wightiana.

PEPINIA, Brongn. ex André, Illustr, Hortic. xvii. (1870) 32. t. 5 = **Pitcairnia**, L'Hérit. (Bromel.).
aphelandraeflora, André, l. c. = Pitc. aphelandraeflora.
punicea, André, l. c. = Pitc. punicea.

PEPLIDIUM, Delile, Fl. Egypte, 148. t. 4 (1813). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 957.
capense, Spreng. Syst. i. 43 = Glossostigma spathulatum.
diandrum, Voigt, Hort. Suburb. Calc. 503 = Glossostigma spathulatum.
humifusum, *Delile, Fl. Egypte*, 148.—Aegypt.
Muelleri, *Benth. Fl. Austral.* iv. 500.—Austral.

PEPLIS, Linn. Syst. ed. I (1735). *LYTHRARIEAE*, Benth. & Hook. f. i. 776.
CHABRAEA, Adans. Fam. ii. 234 (1763).
MIDDENDORFIA, Trautv. in Mém. Sav. Etr. Acad. Pétersb. iv. (1842) 489.
PORTULA, Dill. ex Moench, Meth. 661 (1794).
alternifolia, *Bieb. Fl. Taur. Cauc. Suppl.* 277.—Rossia; Sibir.
alternifolia, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 63 = Tetradiclis salsa.
americana, Pursh, Fl. Am. Sept. i. 238 = Elatine americana.
australis, J. Gay, ex Schult. f. Syst. vii. 54 = erecta.
biflora, Salzm. ex DC. Prod. iii. 77 = erecta.
Boraei, *Ford. Obs. Pl. Crit.* iii. (1846) 81.—Gallia.
Boraei, Lange, in Kjoeb. Vidensk. Meddel. (1865) 140 = erecta.
borysthenica, *Bess. Enum. Pl. Volh.* 81.—Europ.
diandra, Nutt. ex DC. Prod. iii. 77 = Didiplis linearis.
erecta, *Req. ex Benth. Cat. Pl. Pyrén.* 111.—Reg. Mediterr.
Fradini, *Pomel, Nouv. Mat. Fl. Atl.* 156.—Alger.
glandulifera, Dulac, Fl. Hautes-Pyr. 327 = P. Portula.
hispidula, Dur. in Duch. Rev. Bot. ii. (1846–47) 431 = erecta.

PEPLIS:—
indica, Willd. Sp. Pl. ii. 244 = Ammannia peploides.
longedentata, Boiss. & Reut. ex Nym. Consp. 252 = P. Portula.
nummulariaefolia, Jord. Obs. Pl. Crit. iii. (1846) 95 = erecta.
occidentalis, Spreng. Syst. ii. 135 = Rotala ramosior.
Portula, *Linn. Sp. Pl.* 332.—Europ.; Reg. Caucas.
Portula, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1818) 14, nomen.—Cf. S. Wats. Bibliog. Ind. N. Am. Bot. 376 = Ludwigia palustris?
Portulaca, Crantz, Inst. i. 139 = P. Portula.
portulacaefolia, Gilib. Fl. Lituan. ii. 147 = P. Portula.
serpyllifolia, Rupr. Fl. Ingr. i. 387 = P. Portula.
tetrandra, Linn. Diss. Pugill. Jam. App.; Amoen. Acad. v. 413 = Lucya tuberosa.
Timeroyi, Jord. Obs. Pl. Crit. iii. (1846) 83 = erecta.
tithymaloides, Bertol. Fl. Ital. iv. 233 = erecta.
wolgensis, Fisch. ex DC. Prod. iii. 77 = alternifolia.

PEPLONIA, Decne. in DC. Prod. viii. 545 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 763.
amazonica, *Benth. in Hook. Ic. Pl.* t. 1234.—Bras.
Hilariana, *Fourn. in Mart. Fl. Bras.* vi. iv. 237.—Bras.
nitida, *Decne. in DC. Prod.* viii. 546.—Bras.

PEPO, [Tourn.] Linn. Syst. ed. I (1735) = **Cucurbita**, Tourn.
Citrullus, Sageret, in Ann. Sc. Nat. Sér. I. viii. (1826) 311 = C. Pepo.
eximius, Sageret, l. c. = C. moschata.
lagenarius, Moench, Meth. 658 = Lagenaria vulgaris.
macrocarpus, Rich. ex Spach, Hist. Vég. Phan. vi. 202 = C. moschata.
malabaricus, Sageret, in Ann. Sc. Nat. Sér. I. viii. (1826) 312 (Quid?).—Ind. or.
maximus, Peterm. Fl. Lips. Excurs. 562 = C. maxima.
Melopepo, Moench, Meth. 653 = Cucumis Melo.
moschatus, Sageret, in Ann. Sc. Nat. Sér. I. viii. (1826) 312 = C. moschata.
Potiron, Sageret, l. c. 311 = C. Pepo.
verrucosus, Moench, Meth. 653 = Cucumis Melo.
vulgaris, Moench, l. c. = C. Pepo.

PEPONIA, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 29. t. 3. *CUCURBITACEAE*, Benth. & Hook. f. i. 823.
Boivinii, *Cogn. in DC. Monog. Phan.* iii. 408.—Madag.
Bojeri, *Cogn. l. c.* 410.—Afr. trop.
caledonica, *Cogn. l. c.*—Afr. austr.
Cienkowskii, *Hook. f. in Oliver, Fl. Trop. Afr.* i. 527.—Afr. trop.
lagenarioides, *Hook. f. l. c.*—Afr. trop.
Mackennii, *Naud. in Ann. Sc. Nat.* Sér. V. v. (1866) 29.—Afr. austr.
Vogelii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 526.—Afr. trop.

PEPONOPSIS, Naud. in Ann. Sc. Nat. Sér. IV. xii. (1859) 88. *CUCURBITACEAE*, Benth. & Hook. f. i. 829.
adhaerens, *Naud. l. c.* 89.—Am.

PERA, Mutis, in Vet. Akad. Handl. Stockh. v. (1784) 299. t. 8. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 340.
CLISTANTHUS, Muell. Arg. in DC. Prod. xv. II. 1025 (1866).
CLISTRANTHUS, Poit. ex Baill. Etud. Gén. Euph. 434 (1858).
PERIDIUM, Schott, in Spreng. Syst. iv. Cur. Post. 410 (1827).
PERULA, Schreb. Gen. 703 (1791).
SCHISMATOPERA, Klotzsch, in Wiegm. Arch. i. (1841) 178.
SPIXIA, Leandr. Sacram. in Denkschr. Akad. Muench. vii. (1821) 231. t. 13.
anisotricha, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 426.—Bras.
arborea, Baill. Adansonia, v. (1864–65) 224 = glabrata.
arborea, *Mutis, in Vet. Akad. Handl. Stockh.* v. (1784) 299.—N. Granat.
Bailloniana, *Muell. Arg. in DC. Prod.* xv. II. 1030.—Bras.
bicolor, *Muell. Arg. l. c.* 1028.—Guian.

PERA:—
bumeliaefolia, Baill. Adansonia, v. (1864–65) 224 = Bailloniana.
bumelifolia, *Griseb. in Goett. Nachtr.* (1865) 180.—Cuba.
cinerea, Baill. Adansonia, v. (1864–65) 223 = tomentosa.
citriodora, *Baill. l. c.* 222.—Bras.
coccinea, *Muell. Arg. in DC. Prod.* xv. II. 1028.—Bras.
decipiens, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 201.—Guian.
distichophylla, *Baill. Etud. Gén. Euph.* 434.—Bras.
echinocarpa, *Baill. Adansonia*, v. (1864–65) 223.—Bras.
ferruginea, *Muell. Arg. in DC. Prod.* xv. II. 1031.—Bras.
furfuracea, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 426.—Bras.
glabrata, *Poepp. ex Baill. Etud. Gén. Euph.* 434.—Bras.
heterodoxa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 423.—Bras.
Klotzschiana, Baill. Adansonia, v. (1864-65) 225 = glabrata.
Leandri, Baill. Etud. Gén. Euph. 434 = obovata.
obovata, *Baill. Adansonia*, v. (1864–65) 225.—Bras.
obtusifolia, Muell. Arg. in DC. Prod. xv. II. 1030 = obovata.
oppositifolia, *Griseb. in Goett. Nachr.* (1865) 181; *Muell. Arg. in DC. Prod.* xv. II. 1029.—Cuba.
parvifolia, *Muell. Arg. l. c.* 1031.—Bras.
Schomburgkiana, *Muell. Arg. l. c.* 1027.—Guian.
tomentosa, *Muell. Arg. l. c.* 1028.—Bras.

PERACARPA, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 26. *CAMPANULACEAE*, Benth. & Hook. f. ii. 558.
carnosa, *Hook. f. & Thoms. l. c.*—Reg. Himal.

PERALTEA, H. B. & K. Nov. Gen. et Sp. vi. 469. t. 589 (1823) = **Brongniartia**, H. B. & K. (Legumin.).
lupinoides, Benth. Pl. Hartw. 10 = B. Benthamiana.
lupinoides, H. B. & K. Nov. Gen. et Sp. vi. 471. t. 589 = B. thermoides.
oxyphylla, DC. Mém. Lég. xiii. 463 = B. oxyphylla.

PERAMA, Aubl. Pl. Gui. i. 54. t. 18 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 147.
BUCHIA, H. B. & K. Nov. Gen. et Sp. ii. 269. t. 132 (1817).
MATTUSCHKAEA, Schreb. Gen. 788 (1791).
dichotoma, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 29.—Bras.
ericoides, Poepp. & Endl. l. c. = hirsuta.
galioides, *Poir. in Dict. Sc. Nat.* xxxviii. 414.—Venezuela.
hirsuta, *Aubl. Pl. Gui.* i. 54. t. 18.—Guian.
hispida, Poir. in Dict. Sc. Nat. xxxviii. 414 = hirsuta.
humilis, *Benth. in Hook. Journ. Bot.* iii. (1841) 239.—Guian.
setulosa, Miq. in Linnaea, xviii. (1844) 617 = hirsuta.
stricta, Benth. in Hook. Journ. Bot. ii. (1841) 239 = galioides.

PERAMIBUS, Rafin. Ann. Nat. 14 (1820) = **Coreopsis**, Linn. (Compos.).
acutus, Rafin. l. c. = Actinomeris squarrosa.
hirtus, Rafin. l. c. = Rudbeckia triloba.
scaber, Rafin. l. c. = C. scabra.

PERAMIUM, Salisb. in Trans. Hort. Soc. i. (1812) 301 = **Goodyera**, R. Br. (Orchid.).
repens, Salisb. l. c. = G. repens.

PERAPHORA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 20 = **Lophophyllum**, Griff. (Menisp.).
robusta, Miers, l. c. 21 = L. bicristatum.

PERAPHYLLUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 474 (1840). *ROSACEAE*, [Benth. & Hook. f. i. 628].
ramosissimum, *Nutt. ex Torr. & Gray, l. c.*—Am. bor. occ.

PERCEPIER, Dill. ex Moench, Meth. 690 (1794) = **Alchemilla**, Tourn. (Rosac.).
arvensis, Moench. l. c. = A. arvensis.

PERDICIUM, Linn. Pl. Afr. Rar. 22 (1762) = **Gerbera**, Gronov. et **Trixis**, P. Br. (Compos.).
Anandria, R. Br. in Ait. Hort. Kew. ed. II. v. 84 = Gerbera Anandria.
bocconiaefolium, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 393 = Jungia floribunda, *Less.*
brasiliense, R. Grah. in Edinb. N. Phil. Journ. (Jan.-Mar. 1827) 387 = Trixis divaricata.
brasiliense, Linn. Mant. i. 115 = Trixis Lessingii.
brasiliense, Mart. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 189 = Trixis glutinosa.
cacalioides, H. B. & K. Nov. Gen. et Sp. iv. 154 = Trixis cacalioides.
calycinum, Mart. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 188 = Trixis calycina.
capense, Linn. Pl. Afr. Rar. 22 = *Taraxaci.*
chilense, Willd. Sp. Pl. iii. 2118 = Chaetanthera chilensis.
cordatum, Cerv. in La Llave et Lex. Nov. Veg. Desc. fasc. i. 27; Moç. [& Sesse,] ex DC. Prod. vii. 66 = Perezia hebeclada.
corymbosum, Sesse & Moç. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 188 = Trixis corymbosa.
decurrens, Moç. & Sesse, ex DC. Prod. vii. 68 = Trixis decurrens.
divaricatum, H. B. & K. Nov. Gen. et Sp. iv. 155 = Trixis divaricata.
exscapum, Steud. Nom. ed. I. 604 = Chaptalia exscapa.
flexuosum, H. B. & K. Nov. Gen. et Sp. iv. 155 = Trixis divaricata.
glaciale, Poepp. ex DC. Prod. vii. 60 = Leuceria pulchella.
havanense, H. B. & K. Nov. Gen. et Sp. iv. 155 = Trixis frutescens.
lactucoides, Vahl, in Skrivt. Nat. Hist. Selsk. i. II. (1791) 11 = Perezia lactucoides.
laevigatum, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 322 = Perezia lactucoides.
laevigatum, Berg. in Vet. Acad. Handl. Stockh. (1772) 238 = Trixis frutescens.
leiocarpum, DC. Prod. vii. 39.—Afr. austr.
Leontodon, Crantz, Inst. i. 230.—Hab.?
lithospermifolium, Poepp. ex DC. Prod. vii. 60 (= *Chabraea lithospermifolia*).—Chili.
longifolium, Sesse & Moç. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 192 = Trixis longifolia.
lyratum, R. Br. ex Steud. Nom. ed. II. ii. 303 = Chaptalia lyrata.
magellanicum, Linn. f. Suppl. 376 = Perezia magellanica.
mexicanum, Alaman, ex DC. Prod. vii. 67 = Perezia Alamani.
mexicanum, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xvi. (1830) 201 = Perezia Humboldtii.
nervosum, Thunb. Prod. Pl. Cap. 160 = Landtia nervosa.
nutans, Poepp. ex DC. Prod. vii. 63 = Perezia nutans.
oblongum, Ruiz & Pav. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 240 = Chaptalia oblonga.
ovale, Ruiz & Pav. ex D. Don, l. c. = Chaptalia ovalis.
paniculatum, Poepp. ex DC. Prod. vii. 57 = Leuceria paniculata.
pedicularifolium, Poepp. ex Less. Syn. Comp. 410 = Perezia pedicularidifolia.
piloselloides, Juss. ex Steud. Nom. ed. II. ii. 303 = Chaptalia exscapa.
piloselloides, Vahl, in Skrivt. Nat. Hist. Selsk. ii. II. (1793) 38 = Chaptalia piloselloides.
populifolium, Sm. in Rees, Cycl. xxvi. n. 4 = Gerbera Anandria.
prenanthoides, Poepp. ex Less. Syn. Comp. 409 = Perezia prenanthoides.
Printzii, Cass. in Dict. Sc. Nat. xxxviii. 426 = *Taraxaci.*
purpureum, Vahl, in Skrivt. Nat. Hist. Selsk. i. II. (1791) 9 = Leuceria purpurea.
radiale, Linn. Sp. Pl. ed. II. 1248 = Trixis frutescens.
recurvatum, Poepp. ex Less. Syn. Comp. 412 = Perezia linearis.
recurvatum, Vahl, in Skrivt. Nat. Hist. Selsk. i. II. (1791) 13 = Perezia recurvata.
roseum, Poepp. ex Less. Syn. Comp. 405 = Leuceria runcinata.
sarmentosum, Steud. Nom. ed. I. 604 = Chevreulia stolonifera.
semifloscularе, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 166 = Gerbera nepalensis.

PERDICIUM:—
semifloscularе, Linn. Amoen. Acad. vi. 103 = *Taraxaci.*
semifloscularе, Wall. Fl. Carol. 204 = Chaptalia tomentosa.
senecioides, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 149 = Senecio perdicioides.
senecioides, Willd. ex Less. in Linnaea, v. (1830) 13 = Perezia Humboldtii.
serrulatum, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xvi. (1830) 200 = Perezia reticulata.
sinuatum, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 322 = Perezia magellanica.
squarrosum, Vahl, in Skrivt. Nat. Hist. Selsk. i. II. (1791) 11 = Perezia squarrosa.
suaveolens, Urv. in Mém. Soc. Linn. Par. iv. (1826) 611 = Leuceria gossypina.
Taraxaci, Vahl, in Skrivt. Nat. Hist. Selsk. i. II. (1791) 9.—Afr. austr.
tomentosum, Blanco, Fl. Filip. ed. I. 630 = Grangea maderaspatana.
tomentosum, Thunb. Fl. Jap. 319 = Gerbera Anandria.
triflorum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 169 = Ainsliaea pteropoda.
viscosum, Poepp. ex Less. Syn. Comp. 408 = Perezia viscosa.

PEREBEA, Aubl. Pl. Gui. ii. 952. t. 361. (1775). *URTICACEAE*, Benth. & Hook. f. iii. 373.
MIKANIA, Neck. Elem. ii. 217 (1790).
calophylla, *Benth. & Hook. f. Gen.* iii. 371, in obs.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 953. t. 361.—Guian.; Ind. occ.
integrifolia, *Karst. Fl. Columb.* ii. 23.—N. Granat.
laurifolia, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 133.—Guian.
xanthochyma, *Karst. Fl. Columb.* ii. 23.—N. Granat.

PEREILEMA, J. & C. Presl, Rel. Haenk. i. 233. t. 37. f. a (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1144.
brasilianum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. *Sc. Nat.* (1845) 382.—Bras.
ciliatum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 543.—Mexic.
crinitum, *J. & C. Presl, Reliq. Haenk.* i. 233. t. 37. f. a.—Mexic.
crinitum, Trin. Sp. Gram. Ic. iii. t. 358 = brasilianum.
setarioides, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 544.—Mexic.

PEREIRA, Hook. f. & Thoms. Fl. Brit. Ind. i. 99 (1872) sphalm. = seq.

PEREIRIA, Lindl. Fl. Med. 370 (1838) = **Coscinium**, Colebr. (Menisp.).
medica, Lindl. l. c. 370 = C. fenestratum.

PEREMIS, Rafin. Fl. Tellur. iv. 104 (1836) = **Passiflora**, Linn.
orbiculata, Rafin. l. c. = Passif. Murucuja.

PEREPUSA, Steud. Nom. ed. II. i. 303 (1841), sphalm. = **Prepusa**, Mart. (Gentian.).

PERESCIA, Lem. Cact. Hort. Monv. 14 (1838) = seq.

PERESKIA, Plum. ex Linn. Syst. ed. I (1735); Mill. Dict. ed. VI (1752). *CACTEAE*, Benth. & Hook. f. i. 851.
acardia, Parm. ex Pfeiff. Enum. Diagn. Cact. 176 = aculeata.
aculeata, *Mill. Gard. Dict.* ed. VIII; *Haw. Syn. Pl. Succ.* 198.—Ind. occ.
affinis, *Meinsh. in C. Koch & Fint. Wochenschr.* ii. (1859) 118.—Hab.?
Bleo, *DC. Prod.* iii. 475.—Mexic.; N. Granat.
brasiliensis, Hort. Hamb. ex Pfeiff. Enum. Diagn. Cact. 176 = aculeata.
calandriniaefolia, *Hort. Berol. ex Salm Dyck, Cact. Hort. Dyck.* ed. II. 252.—Mexic.
crassicaulis, Zucc. ex Pfeiff. Enum. Diagn. Cact. 176 = spathulata.
cruenta, Hort. ex Pfeiff. l. c. = P. Bleo.
glomerata, *Pfeiff. in Meyen, Reise*, i. 452.—Peruv.

PERESKIA:—
grandiflora, Hort. ex Pfeiff. Enum. Diagn. Cact. 177 = seq.
grandifolia, *Haw. Rev. Pl Succ.* 85.—Bras.
grandispina, *Hort. ex Forbes, Hort. Tour Germ.* 159, nomen.—Bras. ?
Haageana, *Meinsh. in C. Koch & Fint. Wochenschr.* ii. (1859) 118.—Hab. ?
horrida, *DC. Prod.* iii. 475.—Bras.
lanceolata, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 252.—Hab. ?
lanceolata, Hort. Hamb. ex Foerst. Handb. Cact. ed. I. 513 = aculeata.
longispina, Haw. Syn. Pl. Succ. 178 = aculeata.
lychnidiflora, *DC. in Mém. Mus. Par.* xvii. (1828) 75. t. 18.—Mexic.
ochnacarpa, *Miq. in Bull. Sc. Nat. Phys. Néerl.* (1838) 48.—Hab. ?
opuntiaeflora, *DC. in Mém. Mus. Par.* xvii. (1828) 76. t. 19.—Mexic.
Pititache, Karw. ex Labour. Cact. 503 = calandriniaefolia.
Pititache, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 176.—Mexic.
plantaginea, *Hort. Gotting. ex Pfeiff. Enum. Diagn. Cact.* 179.—Hab. ?
Poeppigii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 76, 252.—Am. austr.
portulacaefolia, *Haw. Syn. Pl. Succ.* 199.—Ind. occ.
rosea, *Hort. ex A. Dietr. in Otto & Dietr. Allg. Gartz.* xix. (1851) 153.—Mexic. ?
rotundifolia, *DC. in Mém. Mus. Par.* xvii. (1828) 77. t. 20.—Mexic.
Sacharosa, Griseb. in Goett. Abh. xxiv. (1879) 141 = aculeata.
spathulata, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 176.—Mexic.
subulata, *Muehlenpf. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 347.—Chili.
undulata, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 11.—Bras.
zinniaeflora, *DC. in Mém. Mus. Par.* xvii. (1828) 75. t. 17.—Mexic.

PERESKIA, Vell. Fl. Flum. 34 (1825); i. t. 81 (1827) = **Hippocratea**, Linn. (Celastr.).
paniculata, Vell. ll. cc. = H. ovata ?

PEREUPHORA. Hoffmgg. Verz. Pfl. Nachtr. ii. 173 (1826) = **Serratula**, Linn. (Compos.).
coronata, Hoffmgg, l. c. = S. coronata.

PEREZIA, Lag. Amen. Nat. i. 31 (1811). *COMPOSITAE*, Benth. & Hook. f. ii. 500.
ACOURTIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 203.
CLARIONEA, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 65.
DROZIA, Cass. in Dict. Sc. Nat. xxxiv. 217 (1825).
DUMERILIA, Less. in Linnaea, v. (1830) 13.
HETERANTHUS, Bonpl. ex Cass. in Dict. Sc. Nat. xxi. 110 (1821).
HOMANTHIS, H. B. & K. Nov. Gen. et Sp. iv. 12 (1820).
HOMOIANTHUS, Bonpl. ex DC. in Ann. Mus. Par. xix. (1812) 65.
ISANTHUS, DC. Prod. vii. 63, in syn. (1838).
POGONURA, DC. in Lindl. Introd. Nat. Syst. ed. II. 263 (1836).
SCOLYMANTHUS, Willd. ex DC. Prod. vii. 63 (1838).
acanthoides, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 33 = multiflora.
adnata, *A. Gray, Pl. Wright.* i. 127.—Mexic.
Alamani, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
arizonica, A. Gray, in Bot. Calif. i. 422 = Wrightii.
Beckii, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 34 = recurvata.
bidentata, *Meyen, Reise*, i. 470.—Peruv.
carduncelloides, *Griseb. in Goett. Abh.* xix. (1874) 198.—Reg. Argent.
carpholepis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 60.—Mexic.
carthamoides, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 34 = diversifolia.
ciliaris, *Hook. & Arn. l. c.*—Chili.
cirsiifolia, *Wedd. Chlor. And.* i. 41.—Am. austr.
coerulescens, *Wedd. l. c.* 39.—Peruv.
Coulteri, A. Gray, in Proc. Am. Acad. xv. (1880) 40 = Wrightii.
cubataënsis, *Less. in Linnaea*, v. (1830) 16.—Bras.
dicephala, *Less. l. c.* 21.—Hab. ?
diversifolia, *Meyen, Reise*, i. 311, 347.—Chili.
Doniana, Less. Syn. Comp. 412 = recurvata.
Dugesii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 60.—Mexic.
fruticosa, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 26.—Mexic.
Gayana, DC. in Deless. Ic. Sel. iv. t. 94 = diversifolia.
glandulosa, *Meyen, Reise*, i. 470.—Peruv.
hebeclada, *A. Gray, Pl. Wright.* i. 127.—Mexic.
Humboldtii, *A. Gray, l. c.* 128.—Mexic.
inermis, *Wedd. Chlor. And.* i. 39.—Chili.
integrifolia, *Wedd. l. c.* 40.—Peruv.
Kingii, *Baker, in Mart. Fl. Bras.* vi. III. 380.—Reg. Argent.
lactucoides, *Less. in Linnaea*, v. (1830) 22.—Reg. Magellan.
laevis, *Less. l. c.* 18.—Bras.
Lagascae, Cass. in Dict. Sc. Nat. xxxviii. 455 = magellanica.
lanigera, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 42.—Patagon.
linearis, *Less. Syn. Comp.* 412.—Chili.
lyrata, *Wedd. Chlor. And.* i. 42.—Chili.
magellanica, *Lag. Amen. Nat.* i. 31.—Reg. Magell.
microcephala, *A. Gray Pl. Wright.* i. 127.—Calif.
moschata, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 26.—Mexic.
multicapitata, *Wedd. Chlor. And.* i. 44.—Chili.
multiflora, *Less. in Linnaea*, v. (1830) 19.—Bras.
nana, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 111.—Am. bor. occ.
nivalis, *Wedd. Chlor. And.* i. 39.—Peruv.
nudicaulis, *A. Gray, Pl. Wright.* i. 127.—Guatemala.
nutans, *Less. Syn. Comp.* 409.—Chili.
ochroleuca, Less. in Linnaea, v. (1836) 22 = Trixis ochroleuca.
oxylepis, *Sch. Bip. ex A. Gray, in Proc. Am. Acad.* xv. (1880) 41.—Mexic.
Parryi, *A. Gray, l. c.* 40.—Mexic.
patens, *A. Gray, Pl. Wright.* i. 127.—Mexic.
pedicularidifolia, *Less. Syn. Comp.* 410.—Chili.
perfoliata, *Remy, in C. Gay, Fl. Chil.* iii. 415.—Chili.
pilifera, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 34.—Chili.
pinnatifida, *Lag. ex D. Don, in Trans. Linn. Soc.* xvi. (1830) 210, in syn.—Peruv.
platyphylla, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 111.—N. Mexic.
Poeppigii, *Less. Syn. Comp.* 411.—Chili.
prenanthoides, *Less. l. c.* 409.—Chili.
pristiphylla, *Wedd. Chlor. And.* i. 37.—Chili.
pungens, *Less. in Linnaea*, v. (1830) 20.—Peruv.
purpurata, *Wedd. Chlor. And.* i. 43.—Bolivia.
pygmea, *Wedd. l. c.* 40.—Am. austr.
recurvata, *Less. in Linnaea*, v. (1830) 21.—Chili.
reflexa, *Meyen, Reise*, i. 311.—Chili.
reticulata, *A. Gray, Pl. Wright.* i. 128.—Mexic.
rigida, *A. Gray, l. c.* 127.—Mexic.
runcinata, *Lag. ex D. Don, in Trans. Linn. Soc.* xvi. (1830) 207, in syn.—Texas; Mexic.
Seemannii, *A. Gray, Pl. Wright.* i. 127.—Mexic.
sonchifolia, *Baker, in Mart. Fl. Bras.* vi. III. 380.—Bras.
spathulata, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 33 = viscosa.
squarrosa, *Less. in Linnaea*, v. (1830) 15.—Bras.
Thurberi, *A. Gray, in Mem. Acad. Am.* N. S. v. (1855) 324.—Arizona.
thyrsoidea, *A. Gray, in Torr. Bot. Mex. Bound.* 104.—Mexic.
turbinata, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 25.—Mexic.
violacea, *Wedd. Chlor. And.* i. 42.—Bolivia.
virens, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 34 = Poeppigii.
viscosa, *Less. Syn. Comp.* 408.—Chili.
Wislizeni, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 111.—Mexic.
Wrightii, *A. Gray, Pl. Wright.* i. 127.—Am. bor. occ.

PERFOLIATA, Dod. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 390 = **Bupleurum**, Tourn. (Umbell.).
protracta, Fourr. l. c. = B. protractum.
rotundifolia, Fourr. l. c. = B. rotundifolium.

PERFONON, Rafin. Sylva Tellur. 29 (1838) = **Rhamnus**, Tourn.
ferrugineum, Rafin. l. c. = Colubrina ferruginosa.
laurifolium, Rafin. l. c. = R. californicus.

PERGULARIA, Linn. Mant. i. 8 (1767). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 773.
accedens, *Blume, Bijdr.* 1056.—Java.
albiflora, *Zipp. ex Span. in Linnaea*, xv. (1841) 323, nomen.—Ins. Timor.
angustifolia, D. Dietr. Syn. Pl. ii. 894 = Lygisma angustifolia.
angustifolia, *Fisch. Cat. Hort. Gorenk.* ed. II. (1812) 29 (Quid ?).—Hab. ?
bifida, *Zipp. in Ann. Sc. Nat.* Sér. II. ix. (1838) 276.—Ins. Timor.
Brunoniana, D. Dietr. Syn. Pl. ii. 894 = Marsdenia Brunoniana.
calesiana, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 37 = Marsdenia Calesiana.
campanulata, Buch.-Ham. ex Wall. Pl. As. Rar. ii. 50 = Raphistemma pulchellum.
cinerascens, Spreng. Syst. i. 844 = Marsdenia cinerascens.
clausa, Spreng. l. c. = Marsdenia clausa.
coromandeliana, Decne. in DC. Prod. viii. 619 = odoratissima ?
crocea, Zipp. ex Span. in Linnaea, xv. (1841) 323 = Marsdenia tenacissima.
divaricata, Lour. Fl. Cochinch. 169 = Strophanthus divergens.
edulis, *Thunb. Prod. Pl. Cap.* 38.—Afr. austr.
erecta, Spreng. Syst. i. 844 = Marsdenia erecta.
exilis, Spreng. l. c. = Tylophora exilis.
flavescens, D. Dietr. Syn. Pl. ii. 894 = Marsdenia flavescens.
glabra, Blanco, Fl. Filip. ed. II. 141 = procumbens.
glabra, Linn. Mant. i. 53 = Vallaris Pergulana.
glandulosa, Blanco, Fl. Filip. ed. I. 201 = minor.
Hamiltonii, D. Dietr. Syn. Pl. ii. 894 = Marsdenia Hamiltonii.
japonica, Thunb. Fl. Jap. 111 = Metaplexis Stauntoni.
jasminiflora, Burch. ex Decne. in DC. Prod. viii. 630 = Barrowia jasminiflora.
limbata, Wall. ex Wight, Contrib. 42 = odoratissima ?
lucida, *Fisch. Cat. Hort. Gorenk.* ed. II. (1812) 29 (Quid ?).—Hab. ?
minor, *Andr. Bot. Rep.* t. 184.—As. trop.
montana, Decne. in DC. Prod. viii. 619 = odoratissima ?
odora, Salisb. Prod. 151 = Daemia cordata.
odoratissima, Roxb. Hort. Beng. 20; Fl. Ind. ii. 46; Wight & Arn. Contrib. 43 = minor.
odoratissima, *Sm. Ic. Pict.* t. 16; *Andr. Bot. Rep.* t. 185.—China; Ind. or. ?
pallida, Wight & Arn. Contrib. 42 = odoratissima ?
parviflora, Blume, Bijdr. 1056 = Marsdenia tinctoria.
procumbens, *Blanco, Fl. Filip.* ed. I. 201.—Malaya.
puberula, Miq. Fl. Ind. Bat. ii. 495 = procumbens.
purpurea, *Vahl, Symb. Bot.* iii. 44.—Ind. or.; China.
rostrata, Spreng. Syst. i. 844 = Marsdenia rostrata.
Roylei, D. Dietr. Syn. Pl. i. 894 = Marsdenia Roylei.
sanguinolenta, Lindl. Bot. Reg. t. 2532 (Cryptolepidis sp.).—Afr. trop.
sinensis, Lour. Fl. Cochinch. 169 = Vallaris sinensis.
suaveolens, Spreng. Syst. i. 843 = Marsdenia suaveolens.
syringaefolia, Zipp. ex Decne. in DC. Prod. viii. 624 = Marsdenia syringaefolia.
tenacissima, D. Dietr. Syn. Pl. ii. 895 = Marsdenia tenacissima.
tinctoria, Spreng. Syst. i. 844 = Marsdenia tinctoria.
tomentosa, Linn. Mant. i. 53 = Daemia cordata.
tomentosa, Span. in Linnaea, xv. (1841) 323 = Marsdenia tenacissima.
velutina, Spreng. Syst. i. 844 = Marsdenia velutina.
villosa, *Blume, Bijdr.* 1057.—Java.
viridiflora, Spreng. Syst. i. 844 = Marsdenia viridiflora.
viridis, Buch.-Ham. ex Wight, Contrib. 42 = odoratissima ?

PERIANDRA, Cambess. in Jacquem. Voy. Bot. 27. t. 29 (1844) = **Thylacospermum**, Fenzl (Caryoph.).
caespitosa, Cambess. l. c. = T. rupifragum.

PERIANDRA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 120. *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
acutifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 121.—Bras.
angulata, Benth. l. c. = dulcis.

PERIANDRA:—
angustifolia, Benth. in Ann. Wien. Mus. ii. (1838) 121 = dulcis.
Berteriana, *Benth. l. c.* 122.—Ins. S. Doming.
coccinea, *Benth. l. c.*—Bras.
densiflora, *Benth. l. c.*—Bras.
dulcis, *Mart. ex Benth. l. c.* 121.—Bras.
heterophylla, *Benth. l. c.*—Bras.
mucronata, Mart. ex Benth. l. c. = dulcis.
racemosa, Benth. l. c. = dulcis.

PERIANTHOPODUS, Silva Manso, Enum. Subst. Bras. 28 (1836) = **Cayaponia**, Silva Manso (Cucurb.).
amazonicus, Cogn. in Mart. Fl. Bras. vi. IV. 96 = amazonica.
Bonplandii, Cogn. in Bull. Soc. Bot. Belg. xvii. (1878) 296 = C. Bonplandii.
Carijo, Silva Manso, Enum. Subst. Bras. 29 = C. Espelina.
Espelina, Silva Manso, l. c. 28 = C. Espelina.
Tomba, Silva Manso, l. c. 28 = C. Espelina.
Weddellii, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 203 = C. Weddellii.

PERIBAEA, Lindl. Veg. Kingd. 206 (1847) = PERIBOEA, Kunth = **Hyacinthus**, Tourn. (Liliac.).

PERIBALLANTHUS, Franch. & Sav. Enum. Pl. Jap. ii. 524 (1876) = **Polygonatum**, Adans. (Liliac.).
involucratus, Franch. & Sav. l. c. = Polyg. multiflorum.

PERIBALLIA, Trin. Fund. Agrost. 133 (1820) = **Deschampsia**, Beauv. (Gramin.).
hispanica, Trin. l. c. = D. involucrata.
involucrata, Janka, in Terméz. Füzetek (1877), 97 = D. involucrata.
poaeoides, Trin. ex Steud. Nom. ed. II. i. 492; ii. 304 = D. involucrata.

PERIBLEMA, DC. in Ann. Sc. Nat. Sér. II. xi. (1839) 296. *ACANTHACEAE*, Benth. & Hook. f. ii. 1091.
BOUTONIA, DC. in Bibl. Univ. Genèv. xvii. (1838) 143 (Rev. Bign. 18).
cuspidata, *DC. Prod.* ix. 243.—Madag.

PERIBOEA, Kunth, Enum. Pl. iv. 292 (1843) = **Hyacinthus**, Tourn. (Liliac.).
brevifolia, Kunth, l. c. 294 = H. brevifolius.
corymbosa, Kunth, l. c. 293 = H. corymbosus.
Gawleri, Kunth, l. c. = H. brevifolius.

PERICALIA, Cass. in Dict. Sc. Nat. xlviii. 459 (1827) = **Senecio**, Tourn. (Compos.).
cordifolia, Cass. l. c. = S. cardiophyllus.

PERICALLIS, Webb & Berth. Phyt. Canar. ii. 324 (1836–50) = **Senecio**, Tourn. (Compos.).
cruenta, Webb & Berth. l. c. 336; C. Bolle, in Bonplandia, viii. (1860) 133 = S. cruentus.
echinata, Webb & Berth. l. c. 335 = S. echinatus.
lanata, Webb & Berth. l. c. 324 = S. Heritieri.
multiflora, Webb & Berth. l. c. 326 = S. multiflorus.
papyracea, Webb & Berth. l. c. 339 = S. papyraceus.
populifolia, Webb & Berth. l. c. 328 = S. appendiculatus.
Tussilaginis, Webb & Berth. 331 = S. Tussilaginis.
Webbii, C. Bolle, in Bonplandia, viii. (1860) 133.—Ins. Canar.

PERICALYMMA, Endl. Gen. 1230 (1840) = **Leptospermum**, Forst. (Myrtac.).
crassipes, Schau. in Lehm. Pl. Preiss. i. 120 = L. crassipes.
ellipticum, Schau. l. c. = L. ellipticum.
floridum, Schau. l. c. 121 = L. floridum.
roseum, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 334.—Austral.
teretifolium, Turcz. l. c.—Austral.

PERICALYMNA, Meissn. Gen. Comm. 354 (1843) = praec.

PERICAMPYLUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40. *MENISPERMACEAE*, Benth. & Hook. f. i. 37.

PERICAMPYLUS:—
LEGNEPHORA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 88.
PSELIUM, Lour. Fl. Cochinch. 622 (1790).
aduncus, Miers, l. c. xiv. (1864) 371 = incanus.
assamicus, Miers, l. c. 371 = incanus.
incanus, *Miers, l. c.* Ser. II. vii. (1851) 40.—Ind. or.; Malaya.
lanuginosus, Miq. Fl. Ind. Bat. i. II. 83 = Cocculus lanuginosus.
membranaceus, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 371 = incanus.

PERICAULON, Rafin. New Fl. Am. ii. 50 (1836) = **Baptisia**, Vent. (Legumin.).
cordatum, Rafin. l. c. 51 = B. perfoliata.
microphylum, Rafin. l. c. = B. microphylla.
perfoliatum, Rafin. l. c. = B. perfoliata.

PERICHASMA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 21 = **Stephania**, Lour. (Menisp.).
laetificata, Miers, l. c. 22.—Afr. trop.

PERICLA, Rafin. Sylva Tellur. 112 (1838) = **Cleome**, Linn. (Capparid.).
imbricata, Rafin. l. c. = C. integrifolia.

PERICLISTIA, Benth. in Hook. Journ. Bot. iv. (1842) 108 = **Paypayrola**, Aubl. (Violar.).
latifolia, Benth. l. c. 109 = Payp. guianensis.
longifolia, Benth. l. c.—Guian.
longifolia, Miq. in Mart. Fl. Bras. xiii. I. 378, in syn. = Payp. guianensis.

PERICLYMENUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 250 (1745); Mill. Gard. Dict. ed. VI (1752) = **Lonicera**, Linn. (Caprifol.).
americanum, Mill. Gard. Dict. ed. VIII. n. 7 = L. grata.
germanicum, Mill. l. c. n. 4 = L. Periclymenum.
italicum, Mill. l. c. n. 5 = L. Caprifolium.
javanicum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = L. javanica.
perfoliatum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 487 = L. Caprifolium.
racemosum, Mill. Gard. Dict. ed. VIII. n. 2 = Chiococca racemosa.
sempervirens, Mill. l. c. n. 1 = L. sempervirens.
serotinum, Ait. ex Steud. Nom. ed. II. ii. 304 = L. Periclymenum.
verticillatum, Mill. Gard. Dict. ed. VIII. n. 3 = L. verticillata.
vulgare, Mill. l. c. n. 6 = L. Periclymenum.

PERICODIA, Rafin. Fl. Tellur. iv. 104 (1836) = **Passiflora**, Linn.
perfoliata, Rafin. l. c. = Passifl. perfoliata.

PERICOME, A. Gray, Pl. Wright. ii. 81 (in Smithson. Contrib. v.) (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 406.
GALINSOGEOPSIS, Sch. Bip. in Seem. Bot. Voy. Herald, 306 (1857 ?).
caudata, *A. Gray, Pl. Wright.* ii. 82.—N. Mexic.
spilanthoides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 215.—Mexic.

PERICOPSIS, Thw. Enum. Pl. Zeyl. 413 (1864). *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
Mooniana, *Thw. l. c.* 414.—Zeylan.

PERICTENIA, Miers, Apocyn. S. Am. 182 (1878). *APOCYNACEAE.*
stipellaris, *Miers, l. c.* 183.—Am. austr.

PERICYCLA, Blume, Rumphia, ii. 47. t. 94 (1843) = **Licuala**, Thunb. (Palm.).
penduliflora, Blume, l. c. = L. penduliflora, *Zipp.*

PERIDERAEA, Webb, Iter Hisp. 37 (1838) = **Anthemis**, Mich. (Compos.).
aurea, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 90 = Cotula aurea.
fuscata, Webb, Iter Hisp. 37 = A. fuscata.

PERIDERIDIA, Reichb. Handb. 219 (1837) = **Eulophus**, Nutt. (Umbell.).
americana, Reichb. l. c. = E. americanus.

PERIDISCUS, Benth. in Benth. & Hook. f. Gen. i. 127 (1862). *BIXINEAE*, Benth. & Hook. f. l. c.
lucidus, *Benth. l. c.*—Bras.

PERIDIUM, Schott, in Spreng. Syst. iv. Cur. Post. 410 (1827) = **Pera**, Mutis (Euphorb.).
bicolor, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 44 = Pera bicolor.
coccineum, Benth. in Hook. Kew Journ. vi. (1854) 323 = Pera coccinea.
ferrugineum, Schott, in Spreng. Syst. iv. Cur. Post. 410 = Pera ferruginea.
glabratum, Schott, l. c. = Pera glabrata.
oblongifolium, Benth. in Hook. Kew Journ. ii. (1850) 243 = Pera ferruginea.
obovatum, Klotzsch, in Wiegm. Arch. vii. (1841) 180 = Pera obovata.
obtusifolium, Schott, in Spreng. Syst. iv. Cur. Post. 410 = Pera glabrata.
ovale, Klotzsch, in Wiegm. Arch. vii. (1841) 180 = Pera ferruginea.
parvifolium, Klotzsch, l. c. = Pera parvifolia.
Schomburgkianum, Benth. in Hook. Kew Journ. vi. (1854) 323, in obs. = Pera Schomburgkiana.
Schomburgkii, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1089 = praec.

PERIEILEMA, Benth. & Hook. f. Gen. iii. 1144 (1883), sphalm. = **Pereilema**, Presl (Gramin.).

PERIÈTERIS, Rafin. Fl. Tellur. iii. 74 (1836) = **Nicotiana**, Linn. (Solan.).
tristis, Rafin. l. c. = N. tristis.
viridiflora, Rafin. l. c. = N. Langsdorffii.

PERIGARIA, Span. in Linnaea, xv. (1841) 204 = PIRIGARA, Aubl. = **Gustavia**, Linn. (Myrtac.).

PERIGLOSSUM, Decne. in DC. Prod. viii. 520 (1844) = **Cordylogyne**, E. Mey. (Asclep.).
angustifolium, DC. l. c.—Afr. austr.
M'Kenii, Harv. Thes. Cap. ii. 7. t. 111.—Afr. austr.
macrum, Decne. in DC. Prod. viii. 520 (= *Lagarinthus macer*).—Afr. austr.

PERIHEMIA, Rafin. Fl. Tellur. iv. 20 (1836) = **Haemanthus**, Tourn. (Amaryll.).
coarctata, Rafin. l. c. = H. coccineus.

PERILLA, Linn. Gen. ed. VI. 578 (1764). *LABIATAE*, Benth. & Hook. f. ii. 1182.
DENTIDIA, Lour. Fl. Cochinch. 369 (1790).
arguta, *Benth. in DC. Prod.* xii. 164.—China.
elata, D. Don, Prod. Fl. Nep. 115 = Elsholtzia blanda.
fruticosa, D. Don, l. c. = Elsholtzia polystachya.
lanceolata, *Benth. in DC. Prod.* xii. 164.—China.
leptostachya, D. Don, Prod. Fl. Nep. 115 = Elsholtzia incisa.
macrostachya, Benth. in Wall. Cat. n. 1559 = ocymoides.
marathrosma, Hort. ex Spreng. Nov. Prov. 24 = Lophanthus anisatus.
nankinensis, Decne. in Rev. Hortic. Sér. IV. i. (1852) 61 = arguta.
nankinensis, Hort. ex Wender.—Cf. Linnaea, xxix. (1857–58) 726 = Coleus ingratus.
ocymoides, *Linn. Gen.* ed. VI. 578.—Ind. or.
polystachya, D. Don, Prod. Fl. Nep. 115 = Elsholtzia cristata.
urticaefolia, Salisb. Prod. 80 = ocymoides.

PERILLULA, Maxim. in Bull. Acad. Pétersb. xx. (1875) 463. *LABIATAE*, Benth. & Hook. f. ii. 1182.
reptans, *Maxim. l. c.*—Japon.

PERILOBA, Rafin. Fl. Tellur. iv. 87 (1836) = **Nothoscordum**, Kunth (1843) (Liliac.).
paradoxa, Rafin. l. c. = N. fragrans.

PERILOMIA, H. B. K. Nov. Gen. et Sp. ii. 326, t. 159 (1817). *LABIATAE*, Benth. & Hook. f. ii. 1202.
THERESA, Clos, in C. Gay, Fl. Chil. iv. 496, t. 54 (1849).
cordifolia, Cham. & Schlecht. in Linnaea, vi. (1831) 374 = Scutellaria splendens.

PERILOMIA:—
dubia, *Spreng. Syst.* ii. 704.—Bras.
fruticosa, Cham. & Schlecht. in Linnaea, v. (1830) 102 = Scutellaria Mociniana.
gardoquioides, *Benth. & Hook. f. Gen.* ii. 1203.—Peruv.
ocymoides, Benth. in Bot. Reg. t. 1394 = scutellarioides?
ocymoides, *H. B. & K. Nov. Gen. et Sp.* ii. 328.—Ecuador.
scutellarioides, *H. B. & K. l. c.* 327. t. 157.—Peruv.
tomentosa, *Benth. Lab. Gen. et Sp.* 446.—Mexic.

PERIMA, Rafin. Sylva Tellur. 118 (1838) = **Entada**, Adans. (Legumin.).
odorata, Rafin. l. c. = E. scandens.

PERIMENIUM, Steud. Nom. ed. II. ii. 305 (1841) = **Perymenium**, Schrad. (Compos.).

PERINKA, Rafin. Sylva Tellur. 60 (1838) = **Elaeocarpus**, Burm. (Tiliac.).
grandiflora, Rafin. l. c. = E. grandiflorus, *Sm.*
reticulata, Rafin. l. c. = E. cyaneus.

PERIPEA, Steud. Nom. ed. II. ii. 305 (1841) = PIRIPEA, Aubl. = **Buchnera**, Linn. (Scrophul.).

PERIPHANES, Salisb. Gen. Pl. Fragm. 118 (1866) = **Hessea**, Herb. (Amaryll.).
crispa, Salisb. l. c. = H. crispa.
stellaris, Salisb. l. c. = H. stellaris.

PERIPHAS, Rafin. Fl. Tellur. iv. 85 (1836) = **Convolvulus**, Tourn.
pannifolius, Rafin. l. c. = C. canariensis.

PERIPHRAGMOS, Ruiz & Pav. Prod. 26. t. 24 (1794) = **Cantua**, Juss. (Polemon.).
dependens, Ruiz & Pav. Fl. Per. ii. 18 = C. buxifolia.
flexuosa, Ruiz & Pav. l. c. 17 = C. pyrifolia.
foetidus, Ruiz & Pav. l. c. = Vestia lycioides.
uniflorus, Ruiz & Pav. l. c. 18 = C. ovata.

PERIPLEXIS, Wall. Cat. sub n. 8022 (1832) = **Hemicyclia**, Wight & Arn. (Euphorb.).

PERIPLOCA, Tourn. ex Linn. Gen. ed. I. 65 (1737), partim. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 746.
APLOCA, Neck. Elem. i. 255 (1790).
CAMPELEPIS, Falc. in Trans. Linn. Soc. xix. (1842) 101.
africana, Linn. Sp. Pl. 211 (= *Cynoctonum pilosum*).—Afr. austr.
Afzelii, G. Don, Gen. Syst. iv. 163 = nigrescens?
albicans, *Poir. Encyc.* v. 191 (Quid?).—Ind. or.
alboflavescens, Dennst. Schluess. Hort. Malab. 35 (Parsonsiae sp.?).—Ind. or.
angustifolia, Labill. Ic. Pl. Syr. Dec. ii. 13. t. 7 = laevigata.
aphylla, *Decne. in Jacquem. Voy. Bot.* 109. t. 116.—Oriens; Ind. bor. occ.
arborea, Dennst. Schluess. Hort. Malab. 35 (Quid?).—Ind. or.
audicea, Raeusch. Nom. ed. III. 72 = laevigata.
bifida, D. Dietr. Syn. Pl. ii. 883 = Phyllanthera bifida.
calophylla, *Falc. in Proc. Linn. Soc.* i. (1841) 115.—Reg. Himal.
calumpitensis, Llanos, Fragm. 62 = Streptocaulon Baumii.
capensis, Roxb. ex Arn. in Edinb. N. Phil. Journ. xvii. (1834) 261 = Glossostephanus linearis?
capsularis, Forst. f. Prod. 20 = Parsonsia rosea.
chinensis, Decne. in DC. Prod. viii. 499 = *cochinchinensis*.
chinensis, Spreng. Syst. i. 836 = Vallaris sinensis.
ciliata, Leschen. ex Kostel. Allg. Med.-Pharm. Fl. iii. 1082 (Quid?).—Java?
cochinchinensis, Lour. Fl. Cochinch. 167 (Pergulariae sp.?).—Cochinch.
cordata, Poir. Encyc. v. 191 = Hemidesmus indicus.
divaricata, Spreng. Syst. i. 836 = Strophanthus divergens.
dubia, Burm. f. Fl. Ind. 70 = Cryptolepis Buchanani.
emetica, Retz. Obs. ii. 14 = Secamone emetica.
esculenta, Linn. f. Suppl. 168 = Oxystelma esculentum.
fasciculata, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = laevigata.
Forsteri, Decne. in DC. Prod. viii. 499 = Parsonsia rosea.

PERIPLOCA:—
fruticosa, Mill. Gard. Dict. ed. VIII. n. 3 (Quid?).—Mexic.
gracilis, *Boiss. Fl. Orient.* iv. 50.—Cilicia.
graeca, *Linn. Sp. Pl.* 211.—Europ. austr.; Oriens.
hastata, Decne. in DC. Prod. viii. 499 = Microloma sagittatum.
heterophylla, A. Cunn. ex Decne. l. c. = Parsonsia albiflora.
Humboldtiana, Schult. Syst. vi. 129 = Astephanus cubensis.
Hydaspidis, *Falc. in Proc. Linn. Soc.* i. (1841) 115.—Ind. bor. occ.
indica, Linn. Sp. Pl. 211 = Hemidesmus indicus.
khasiana, Benth. ex Hook. f. Fl. Brit. Ind. iv. 4 = Pentanura khasiana.
laeta, Salisb. Prod. 148 = graeca.
laevigata, *Ait. Hort. Kew.* ed. I. i. 301.—Reg. Mediterr.; Ins. Canar.
laevigata, Bal. ex Boiss. Fl. Orient. iv. 50 = gracilis.
linearifolia, *Dillon & A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 263.—Abyssin.
linearis, Hochst. in Flora, xxiv. (1841) I. Intell. 25 = linearifolia.
maculata, Moench, Meth. 46 = graeca.
malabarica, Burm. ex Decne. in DC. Prod. viii. 499 = Hemidesmus indicus.
mauritiana, Poir. Encyc. v. 188 = Tylophora laevigata.
montana, Rottl. ex Wight, Contrib. 45 = Gymnema montanum.
mucronata, *Willd. ex Schult. Syst.* vi. 129.—Am. austr.
multiflora, *Burm. f. Fl. Cap. Prod.* 7 (Quid?).—Afr. austr.
nigrescens, *Afzel. Stirp. Guin. Med. Sp. Nov.* i. 2.—Afr. trop.
oblongata, *Humb. & Bonpl. ex Schult. Syst.* vi. 129.—Am. austr.
oleaefolia, Salisb. Prod. 148 = laevigata.
ovata, Poir. ex Decne. in DC. Prod. viii. 508 = Astephanus ovatus.
pallida, Salisb. Prod. 148 (= *Cynoctonum pilosum*).—Afr. austr.
Palvalli, Dennst. Schluess. Hort. Malab. 35 (Quid?).—Ind. or.
parviflora, *Poir. Encyc.* v. 192.—Ind. or.
parviflora, Willd. ex Schult. Syst. vi. 129 = Astephanus cubensis.
Petersiana, *Vatke, in Oestr. Bot. Zeitschr.* xxvi. (1876) 147.—Afr. trop.
pubescens, *Willd. ex Schult. Syst.* vi. 129.—Am. austr.
punicaefolia, Cav. Ic. iii. 9. t. 217 = laevigata.
reticulata, Roth, Nov. Pl. Sp. 175 = Cynanchum Callialata?
rigida, Viv. Lib. Spec. 14. t. 6 = laevigata.
Secamone, Lam. ex Decne. in DC. Prod. viii. 498 = laevigata.
Secamone, Linn. Mant. ii. 216 = Oxystelma Alpini.
Secamone, Thunb. Prod. Pl. Cap. 47 = Secamone Thunbergii.
sepium, *Bunge, Enum. Pl. Chin. Bor.* 43.—China.
sinensis, Steud. Nom. ed. II. i. 552, in syn. = Vallaris sinensis.
sylvestris, Retz. Obs. ii. 15 = Gymnema sylvestre.
tenuifolia, Linn. Sp. Pl. 212 = Microloma lineare.
tenuifolia, Willd. ex Schult. Syst. vi. 129 = Gymnema subvolubile.
tunicata, Hort. Madr. ex Wight, Contrib. 56 = Cynanchum Callialata.
tunicata, Retz. Obs. ii. 15 = Cynanchum pauciflorum.
venosa, Hochst. ex Decne. in DC. Prod. viii. 493 = Tacazzea venosa.
virgata, Poir. Encyc. v. 192 (= *Orthosia virgata*).—Bras.
viridiflora, Kostel. Allg. Med.-Pharm. Fl. iii. 1082 = Cryptolepis Buchanani.
vomitoria, Leschen. ex Kostel. l. c. (Quid?).—Java?

PERIPTERA, DC. Prod. i. 459 (1824) (Malvac.).
megapotamica, G. Don, Gen. Syst. i. 490 = Abutilon megapotamicum.
punicea, DC. Prod. i. 459 = Anoda punicea.

PERIPTERYGIUM, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 142; et Cat. Hort. Bog. Alt. 234 (1844) = **Cardiopteris**, Wall. (Olacin.).
quinquelobum, Hassk. ll. cc. 142; 235 = C. lobata.

PERISTEIRA, Hook. f. Fl. Brit. Ind. iv. 277–9 (1884), sphalm. = PENTSTEIRA, Griff. = **Torenia**, Linn. (Scrophul.).

PERISTERA, Eckl. & Zeyh. Enum. 72 (1835) = **Pelargonium**, L'Hérit. (Geran.).
althaeoides, Eckl. & Zeyh. l. c. = Pelarg. althaeoides.
anceps, Eckl. & Zeyh. l. c. = Pelarg. anceps.
brevipes, Presl, Bot. Bemerk. 27 = Pelarg. australe.
columbina, Eckl. & Zeyh. Enum. 73 = Pelarg. australe.
grossularioides, Eckl. & Zeyh. l. c. 72 = Pelarg. australe.
humifusa, Eckl. & Zeyh. l. c. 73 = Pelarg. parvulum.
iocasta, Eckl. & Zeyh. l. c. = Pelarg. australe.
nummulariaefolia, Eckl. & Zeyh. l. c. 72 = Pelarg. australe.
parvula, Eckl. & Zeyh. l. c. 73 = Pelarg. parvulum.
procumbens, Eckl. & Zeyh. l. c. = Pelarg. australe.

PERISTERIA, Hook. Bot. Mag. t. 3116 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 550.
ECKARTIA, Reichb. Nom. 53 (1841).
Barkeri, Batem. Orch. Mex. & Guat. t. 8 = Acineta Barkeri.
cerina, *Lindl. Bot. Reg.* t. 1953.—Am. austr.
elata, *Hook. Bot. Mag.* t. 3116.—Panama.
Ephippium, *Reichb. f. in Gard. Chron.* (1883) II. 198.—Am. trop.
fuscata, Hort. ex Lindl. in Gard. Chron. (1856) 388 = Lycomormium squalidum.
graveolens, Hort. l. c. (1882) II. 151.—Hab.?
guttata, *Knowles & Westc. Flor. Cab.* ii. (1838) 99. t. 70.—Am. austr.
Humboldtii, Lindl. Bot. Reg. (1843) t. 18 = Acineta Humboldtii.
lentiginosa, Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit Suppl. iii. 604 = pendula?
longiscapa, Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 25 = Lacaena bicolor.
maculata, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1474 = pendula.
pendula, *Hook. Bot. Mag.* t. 3479.—Guian.
stapelioides, Loud. Hort. Brit. Suppl. iii. 604 = Zygopetalum stapelioides?

PERISTIMA, Rafin. Sylva Tellur. 89 (1838) = **Heliotropium**, Tourn. (Borag.).
bicolor, Rafin. l. c. = H. strigosum.

PERISTROPHE, Nees, in Wall. Pl. As. Rar. iii. 112 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1121.
RAMUSIA, Nees, in DC. Prod. xi. 309 (1847).
STREPSIPHUS, Rafin. Fl. Tellur. ii. 91 (1836).
acuminata, *Nees, in Wall. Pl. As. Rar.* iii. 113.—Burma; Malaya.
albiflora, *Hassk. Cat. Hort. Bog. Alt.* 152.—Java.
angustifolia, *Nees, in DC. Prod.* xi. 494.—Java.
bicalyculata, *Nees, in Wall. Pl. As. Rar.* iii. 113.—As. et Afr. trop.
Blumeana, Nees, in DC. Prod. xi. 495 = acuminata.
caulopsila, *Presl, Bot. Bemerk.* 95; *Nees, l. c.* 498.—Afr. trop.
cernua, Nees, in Linnaea, xv. (1841) 374 = caulopsila.
chinensis, *Nees, in DC. Prod.* xi. 494.—China.
ciliata, Royle, ex Nees, l. c. 499, 500 = Justicia pubigera.
commutata, *Nees, l. c.* 497.—Ins. Molucc.
contorta, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 157.—Ins. Philipp.
Cumingiana, Nees, in DC. Prod. xi. 498 = Hypoëstes Cumingiana.
Dalaora, Nees, l. c. 499 = tinctoria?
dichotoma, Hassk. Cat. Hort. Bog. Alt. 152 = Dianthera dichotoma.
fera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 556.—Reg. Himal.
fragilis, Nees, in Wall. Pl. As. Rar. iii. 113 = acuminata.
gracilis, Nees, l. c. = acuminata.
grandiflora, Parish, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 556 = lanceolaria.
jalappaefolia, *Nees, in DC. Prod.* xi. 494.—Java.
Kotschyana, Nees, l. c. 497 = bicalyculata.
Krebsii, *Presl, Bot. Bemerk.* 94.—Afr. austr.
lanceolaria, *Nees, in Wall. Pl. As. Rar.* iii. 114.—Ind. or.; Burma.
maculata, Hochst. ex Nees, in DC. Prod. xi. 485 = Dicliptera maculata.
montana, *Nees, in Wall. Pl. As. Rar.* iii. 113.—Ind. or.

PERISTROPHE:—
natalensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 48.—Afr. austr.
oblonga, Nees, in Linnaea, xv. (1841) 375 = Rhinacanthus oblongus.
pallida, *Nees, in DC. Prod.* xi. 494.—Java.
pubigera, Nees, in Wall. Pl. As. Rar. iii. 114 = Justicia pubigera.
pulchella, *Voigt, Hort. Suburb. Calc.* 492.—Hab.?
rivinoides, Wall. Cat. n. 2488 = bicalyculata.
salicifolia, *Hassk. Cat. Hort. Bog. Alt.* 152.—Java.
Schimperiana, Hochst. ex Nees, in DC. Prod. xi. 497 = bicalyculata.
speciosa, *Nees, in Wall. Pl. As. Rar.* iii. 113.—Reg. Himal.
tinctoria, *Nees, l. c.*—Ind. or.; Malaya.
tinctoria, Span. ex Nees, in DC. Prod. xi. 493 = fera.
undulata, *Nees, l. c.* 496.—Ind. or.

PERISTYLIS, Benth. & Hook. f. Gen. iii. 625 (1883), sphalm. = seq.

PERISTYLUS, Blume, Bijdr. 404. t. 30 (1825) = **Habenaria**, Willd. (Orchid.).
albidus, Lindl. Syn. Brit. Fl. ed. II. 261 = H. albida.
arachnoideus, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 267.—Abyss.
aristatus, Lindl. Gen. et Sp. Orch. 300 = H. aristata.
atlanticus, Lindl. l. c. = H. intacta.
brachyphyllus, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 70. t. 2.—Ind. or.
bracteatus, Lindl. Gen. et Sp. Orch. 298 = H. viridis.
brevilobus, Thw. Enum. Pl. Zeyl. 311 = H. breviloba.
chloranthus, Lindl. in Hook. Kew Journ. vii. (1855) 37 = H. lacertifera.
chorisianus, Lindl. Gen. et Sp. Orch. 297 = H. chorisiana.
citrinus, Lindl. l. c. 298 = H. citrina.
Commersonianus, Lindl. l. c. 297.—Ins. Borbon.
constrictus, Lindl. l. c. 300.—Reg. Himal.; Burma.
cordatus, Lindl. l. c. 298 = Orchis cordata.
densiflorus, Lindl. l. c. = H. intacta.
elatus, Dalz. in Hook. Kew Journ. iii. (1851) 344 = H. Wightii.
elatus, Wall. ex Voigt, Hort. Suburb. Calc. 631, nomen. —Hab.?
exilis, Wight, Ic. v. t. 1698 = H. aristatus.
fallax, Lindl. Gen. et Sp. Orch. 298 = Herminium fallax.
filiformis, Kraentz. in Bremen, Abh. vii. (1882) 258 = H. filiformis.
flexuosus, S. Moore, in Baker, Fl. Maurit. 336.—Ins. Mascar.
goodyeroides, Lindl. Gen. et Sp. Orch. 299 = H. goodyeroides.
gracilis, Blume, Bijdr. 404 = H. tipulifera?
gramineus, S. Moore, in Baker, Fl. Maurit. 336 = H. graminea.
grandis, Blume, Bijdr. 405. t. 30 = H. goodyeroides.
Hamiltonianus, Lindl. Gen. et Sp. Orch. 299.—Reg. Himal.
Horsfieldii, Blume, Mus. Bot. Lugd. Bat. ii. 189.—Java.
islandicus, Lindl. Gen. et Sp. Orch. 297 = H. viridis.
lancifolius, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 69.—Ind. or.
latifolius, Lindl. Gen. et Sp. Orch. 297 = H. chlorantha.
Lawii, Wight, Ic. v. t. 1698.—Ind. or.
Lefebvreanus, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 266. t. 16 = Herminium Lefebreanum.
maculatus, Reichb. f. Ic. Pl. Germ. xiii–xiv. 2, in syn. sphalm. = H. intacta.
montanus, Schur, Enum. Pl. Transs. 645 = H. viridis.
Novo-Ebudarum, F. Muell. Phyt. N. Hebrid. 22 = H. Novo-Ebudarum.
obsoletus, Lindl. Gen. et Sp. Orch. 297 = H. obsoleta.
Parishii, Reichb. f. in Trans. Linn. Soc. xxx. (1874) 139.—Ind. or.
Petitianus, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 266.—Abyss.
plantagineus, Lindl. Gen. et Sp. Orch. 300.—Ind. or.
purpureus, S. Moore, in Baker, Fl. Maurit. 335 = H. purpurea.
purpureus, Schur, Enum. Pl. Transs. 646 = H. cucullata.
Quartinianus, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 265.—Abyss.
Richardianus, Wight, Ic. t. 1697.—Ind. or.
robustior, Wight, l. c. v. t. 1699.—Ind. or.

PERISTYLUS:—
sacculatus, Balf. f. & S. Moore, in Journ. Bot. xiv. (1876) 293.—Ins. Borbon.
Sampsoni, Hance, in Journ. Bot. vi. (1868) 371.—China.
satyrioides, Reichb. f. in Bot. Zeit. vii. (1849) 868 (= *Coeloglossum satyrioides*).—Tauria.
secundiflorus, Boiv. ex S. Moore, in Journ. Bot. xiv. (1876) 293, nomen.—Ins. Borbon.
spiralis, S. Moore, in Baker, Fl. Maurit. 335 = H. spiralis, *A. Rich.*
spiralis, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 69. t. 2 B = H. spiralis, *Trimen.*
squamatus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 289.—Abyss.
tridentatus, Hook. f. in Journ. Linn. Soc. vii. (1864) 221.—Afr. trop.
unifolius, Hochst. ex Reichb. f. in Bonplandia, iii. (1855) 213 (= *Deroemera unifolia*).—Abyss.
virescens, Lindl. Gen. et Sp. Orch. 298 = H. herbiola.
viridis, Lindl. Syn. Brit. Fl. ed. II. 261 = H. viridis.

PERITASSA, Miers, in Trans. Linn. Soc. xxviii. (1872) 402 = **Salacia**, Linn. (Celastr.).
adamantina, Miers, l. c. 404.—Bras.
compta, Miers, l. c. 405 = S. laevigata.
dulcis, Miers, l. c. 403 = S. dulcis.
fluminensis, Miers, l. c. 407 = S. fluminensis.
rorida, Miers, l. c. 406 = S. laevigata.
rubicunda, Miers, l. c. 405 = S. laevigata.
smaragdina, Miers, l. c. 404 = S. laevigata.

PERITOMA, DC. Prod. i. 237 (1824) = **Cleome**, Linn. (Cappar.).
aureum, Nutt. in Journ. Acad. Philad. vii. (1834) 15 = C. lutea.
integrifolium, Nutt. l. c. 14 = C. integrifolia.
luteum, Rafin. Sylva Tellur. 112 = C. lutea.
serrulatum, DC. Prod. i. 237 = C. integrifolia.

PERITOMIA, G. Don, Gen. Syst. iv. 668 (1837), sphalm. = **Perilomia**, H. B. & K. (Labiat.).

PERITRIS, Rafin. New Fl. Am. ii. 23 (1836) = **Arnica**? Linn. (Compos.).
ovatifolia, Rafin. l. c. (Quid?).—Am. bor.

PERITTIUM, Vog. in Linnaea, xi. (1837) 408 = **Melanoxylon**, Schott (Legumin.).
ferrugineum, Vog. l. c.—M. Brauna.

PERITYLE, Benth. Bot. Voy. Sulph. 23. t. 15 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
Acmella, Harv. & Gray, in Mem. Am. Acad. N. S. iv. (1849) 77 = microglossa.
aglossa, A. Gray, Pl. Wright. ii. 107 = Laphamia aglossa.
californica, *Benth. Bot. Voy. Sulph.* 23. t. 15.—Calif.
californica, A. Gray, in Proc. Am. Acad. v. (1862) 159 = microglossa.
coronopifolia, *A. Gray, Pl. Wright.* ii. 82.—N. Mexic.
dissecta, *A. Gray, in Proc. Am. Acad.* ix. (1874) 195. —Texas; Mexic.
Emoryi, Torr. in Emory, Notes Mil. Reconnois. 142 = californica.
Fitchii, *Torr. ex A. Gray, in Pacif. Rail. Rep.* iv. 100. —Calif.
incana, *A. Gray, in Proc. Am. Acad.* xi. (1876) 78. —Calif.
leptoglossa, *Harv. & Gray, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 77.—Arizona.
microglossa, *Benth. Bot. Voy. Sulph.* 119.—Calif.
nuda, Torr. ex A. Gray, in Pacif. Rail. Rep. iv. 100 = californica.
Parryi, A. Gray, Pl. Wright. ii. 106 = Laphamia Parryi.
plumigera, *Harv. & Gray, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 77.—Arizona.

PERIZOMA, Miers, ex Lindl. Veg. Kingd. 622 (1847) = **Salpichroa**, Miers (Solan.).

PERLEBIA, DC. Mém. Omb. 67 (1829) = **Prangos**, Lindl. (Umbell.).
triquetra, DC. l. c. = P. triquetra.

PERLEBIA, Mart. Reise Bras. i. 555 (1828) = **Bauhinia**, Linn. (Legumin.).
bauhinoides, Mart. l. c. = B. microphylla.

PERNETTYA, Gaudich. in Ann. Sc. Nat. Sér. I. v. (1825) 102 (Pernettia); et in Freyc. Voy. Bot. 454. t. 67 (1826). *ERICACEAE*, Benth. & Hook. f. ii. 582.
angustata, Benth. Pl. Hartw. 212 = Pentlandii.
angustifolia, *Lindl. Bot. Reg.* (1840) t. 63.—Chili.
brasiliensis, Mart. Herb. Fl. Bras. n. 987; ex Meissn. in Mart. Fl. Bras. vii. 150 = myrtilloides.
breviflora, *Phil. in Linnaea*, xxix. (1857–58) 12.—Chili.
Bridgesii, Phil. in Bot. Zeit. xiv. (1856) 646 = rigida.
buxifolia, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 538.—Mexic.
buxifolia, *Phil. in Anal. Univ. Chil.* (1873) 508.—Chili.
Cavanillesiana, G. Don, Gen. Syst. iii. 837 = Gaultheria microphylla.
ciliaris, *D. Don, ex G. Don, l. c.*—Peruv.
congesta, *Klotzsch, ex Oerst. Amér. Centr.* 9.—Am. centr.
coriacea, *Klotzsch, in Linnaea*, xxiv. (1851) 85.—Bras.
crassifolia, *Phil. l. c.* xxix. (1857–58) 10.—Chili.
elegans, *Phil. l. c.* xxxiii. (1864–65) 173.—Chili.
elliptica, *DC. Prod.* vii. 587.—Peruv.?
empetrifolia, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 102.—Reg. Magellan.
floribunda, *Hort.—Cf. Gard. Chron.* (1882) ii. 649.—Chili?
furens, *Klotzsch, in Linnaea*, xxiv. (1851) 83.—Chili.
Gayana, *Decne. Bot. Voy. Astrol. et Zélée*, ii. *Dicot. Phan.* t. 23.—Chili.
Halliana, *Klotzsch, in Linnaea*, xxiv. (1851) 84.—Ecuador.
humilis, Planch. ex Wedd. Chlor. And. ii. 170 = Pentlandii.
ilicifolia, *Miq. in Lechl. Berb. Am. Austr.* 54, nomen. —Chili.
leucocarpa, *DC. Prod.* vii. 586.—Chili.
linifolia, *Phil. in Linnaea*, xxxiii. (1864–65) 172.—Chili.
litoralis, *Phil. in Anal. Univ. Chil.* (1865) ii. 349.—Chili.
melanocarpa, *Phil. l. c.* (1884) *reimpr.* 5.—Chili.
microphylla, Gaudich. ex G. Don, Gen. Syst. iii. 836 = Gaultheria microphylla.
minima, *Phil. in Linnaea*, xxix. (1857–58) 11.—Chili.
mucronata, *Gaudich. ex G. Don, Gen. Syst.* iii. 836. —Reg. Magellan.
Myrsinites, G. Don, l. c. = Gaultheria microphylla.
myrsinoides, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 325.—N. Granat.
myrtilloides, Griseb. ex Wedd. Chlor. And. ii. 170 = Pentlandii.
myrtilloides, *Meissn. in Mart. Fl. Bras.* vii. 149.—Bras.
myrtilloides, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 325.—Chili.
nitida, Planch. ex Wedd. Chlor. And. ii. 170 = Pentlandii.
nubigena, *Phil. in Anal. Univ. Chil.* i. (1861) 64.—Chili.
oblongifolia, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zélée, ii. Dicot. Phan. t. 22 bis = mucronata.
Oerstediana, *Klotzsch, ex Oerst. Amér. Centr.* 9.—Am. centr.
ovalifolia, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zelée, ii. Dicot. Phan. t. 22 bis = mucronata.
parvifolia, *Benth. Pl. Hartw.* 212.—N. Granat.
Pentlandii, *DC. Prod.* vii. 587.—Reg. Andin.
phillyreaefolia, *DC. l. c.*—Am. austr.
pilosa, *G. Don, Gen. Syst.* iii. 837.—Mexic.
Poeppigii, *Klotzsch, in Linnaea*, xxiv. (1851) 83.—Chili.
pumila, Hook. Ic. Pl. t. 9 = empetrifolia.
punctata, *Klotzsch, in Linnaea*, xxiv. (1851) 83.—Chili.
purpurea, D. Don, ex G. Don, Gen. Syst. iii. 837 = Pentlandii.
quadrifida, *Phil. in Linnaea*, xxix. (1857–58) 11.—Chili.
racemulosa, DC. Prod. vii. 588 = rigida.
repens, Zoll. & Mor. Syst. Verz. Zoll. 42 = Gaultheria nummularioides.
reticulata, *Phil. in Anal. Univ. Chil.* (1873) 507.—Chili.
rigida, *DC. Prod.* vii. 587.—Ins. J. Fernand.
robusta, *Wedd. Chlor. And.* ii. 171.—N. Granat.

PERNETTYA:—
rupicola, *Phil. in Linnaea*, xxix. (1857–58) 10.—Chili.
serpyllifolia, DC. Prod. vii. 587 = Gaultheria microphylla.
setigera, *Klotzsch, in Linnaea*, xxiv. (1851) 85.—Costa Rica.
tasmanica, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 268.—Tasmania.
tenuifolia, *Phil. in Linnaea*, xxix. (1857–58) 10.—Chili.

PERNETYA, Scop. Introd. 150 (1777) = **Canarina**, Linn. (Campan.).

PEROA, Pers. Syn. i. 174 (1805) = PEROJOA, Cav. = **Leucopogon**, R. Br. (Epacrid.).
microphylla, Pers. l. c. = L. microphyllus.

PEROBACHNE, J. & C. Presl, Rel. Haenk. i. 348. t. 48 (1830) = **Anthistiria**, Linn. f. (Gramin.).
secunda, J. & C. Presl. l. c. = A. gigantea.

PEROJOA, Cav. Ic. iv. 29. t. 349 (1797) = **Leucopogon**, R. Br. (Epacrid.).
microphylla, Cav. l. c. = L. microphyllus.

PERONEMA, Jack, in Malay. Misc. ii. (1822) n. VII. 46. *VERBENACEAE*, Benth. & Hook. f. ii. 1158.
canescens, *Jack, l. c.*—Sumatra.
heterophyllum, Miq. Fl. Ind. Bat. Suppl. 570 = canescens.

PERONIA, R. Br in Wall. Cat. n. 7543 (1832) = **Sarcosperma**, Hook. f. (Sapot.).
spicigera, R. Br. l. c.—Ind. or.

PERONIA, [Delar. in] Red. Lil. vi. t. 342 (1812) = **Thalia**, Linn.
stricta, [Delar.] l. c. = T. dealbata.

PEROSTEMA, Raeusch. Nom. ed. III. 224 (1797) = **Nectandra**, Roland. (Laurin.).
bijuga, Raeusch. l. c. = Ocotea guianensis.
sanguinea, Raeusch. l. c. = N. sanguinea.

PEROTIS, Ait. Hort. Kew. ed. I. i. 85 (1789); iii. 506. *GRAMINEAE*, Benth. & Hook. f. iii. 1123.
XYSTIDIUM, Trin. Fund. Agrost. 102 (1820).
Burmanni, Steud. Nom. ed. I. 605 = latifolia.
cubana, *Wright, in Sauv. Fl. Cub.* 202.—Cuba.
glabrata, Steud. Syn. Pl. Gram. 186 = latifolia.
hordeiformis, *Nees, ex Steud. Nom.* ed. II. ii. 306; *et ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 267.—Zeylan.
latifolia, *Ait. Hort. Kew.* ed. I. i. 85.—Ind. or.; Japon.
latifolia, Eckl. ex Steud. Nom. ed. II. ii. 306 = Pennisetum macrourum.
longiflora, *Nees, ex Hook. & Arn. Bot. Beech. Voy.* 247.—China.
patula, Nees, ex Steud. Nom. ed. II. ii. 306 = latifolia.
polystachya, Willd. Sp. Pl. i. 324 = Pogonatherum saccharoideum.
rara, *R. Br. Prod.* 172.—Austral.
scabra, *Willd. ex Trin. Gram. Unifl.* 172.—Afr. trop.
spicaeformis, Beauv. ex Steud. Nom. ed. I. 605 = latifolia.

PEROTRICHE, Cass. in Bull. Soc. Philom. (1818) 75. *COMPOSITAE*, Benth. & Hook. f. ii. 322.
GYMNACHAENA, Reichb. ex DC. Prod. vi. 264 (1837).
microphylla, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 181.—Afr. austr.
tortilis, *Cass. in Bull. Soc. Philom.* (1818) 75.—Afr. austr.

PEROVSKIA, Karel. in Bull. Soc. Nat. Mosc. (1841) 15. t. 1. *LABIATAE*, Benth. & Hook. f. ii. 1193.
abrotanoides, *Karel. l. c.*—Turkest.; Reg. Himal.
artemisioides, *Boiss. Diagn.* Ser. II. iv. 15.—Beluchist.
atriplicifolia, *Benth. in DC. Prod.* xii. 261.—Reg. Himal.; Afghan.
scrophulariaefolia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 433.—Turkestan.

PEROWSKIA, Benth. in DC. Prod. xii. 260 (1848) = praec.

PERPENSUM, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) (1768) = **Gunnera**, Linn. (Halorag.).
blitispermum, Burm. f. l. c. = G. perpensa.

PERRALDERIA, Coss. in Bull. Soc. Bot. Fr. vi. (1859) 394; et in Ann. Sc. Nat. Sér. IV. xviii. (1862) 209. t. 12 = **Grantia**, Boiss. (Compos.).
coronopifolia, Coss. Il. cc. 395; 210 = G. coronopifolia.

PERREYMONDIA, Barn. in Ann. Sc. Nat. Sér. III. iii. (1845) 168 = **Schizopetalon**, Sims (Crucif.).
Brongniartii, Barn. l. c.—Chili.
dentata, Barn. l. c.—Chili.
multifida, Barn. l. c.—Chili.
rupestris, Barn. l. c.—Chili.
viridis, Phil. in Linnaea, xxviii. (1856) 668.—Chili.

PERROTTETIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 95 = **Desmodium**, Desv. (Legumin.).
barbata, DC. l. c. 96 = D. barbatum.
cayennensis, DC. l. c.—Guian.
venustula, DC. l. c. = D. venustulum.

PERROTTETIA, H. B. & K. Nov. Gen. et Sp. vii. 73. t. 622 (1825). *CELASTRINEAE*, Benth. & Hook. f. i. 367.
THEAPHYLLUM, Nutt. ex Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 605.
lanceolata, *Karst. Fl. Columb.* ii. 47. t. 124.—N. Granat.
ovata, *Hemsl. Diagn. Pl. Nov.* 6.—Mexic.
quinduensis, *H. B. & K. Nov. Gen. et Sp.* vii. 75. t. 622.—N. Granat.
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 291.—Ins. Sandvic.

PERSEA, Plum. ex Linn. Gen. ed. I. 94 (1737); Gaertn. f. Fruct. iii. 222. t. 221 (1805). *LAURINEAE*, Benth. & Hook. f. iii. 156.
MENESTRATA, Vell. Fl. Flum. 199 (1825).
TAMALA, Rafin. Sylva Tellur. 136 (1838).
acuminata, Nees & Mart. ex Nees, Syst. Laurin. 170 = Nectandra berchemiaefolia.
alba, *Nees, in Linnaea*, viii. (1833) 51.—Bras.
alpigena, *Spreng. Syst.* ii. 268.—Jamaic.
americana, Mill. Gard. Dict. ed. VIII = gratissima.
amplexicaulis, Cham. & Schlecht. in Linnaea, v. (1830) 90.—Mexic.
angustifolia, Bartl. ex Meissn. in DC. Prod. xv. I. 162 = Nectandra angustifolia.
angustifolia, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 261; *Contrib.* 67.—Reg. Argent.
angustifolia, Schott, in Spreng. Syst. iv. Cur. Post. 405 = Ocotea angustifolia.
argentea, Spreng. Syst. ii. 269 = Ocotea guianensis.
aurata, *Miq. in Linnaea*, xxii. (1849) 805.—Bras.
Ayui-y, *Parodi, in Anal. Soc. Cient. Argent.* iv. (1877) 131; *Contrib.* 68.—Reg. Argent.
azorica, *Seub. Fl. Azor.* 29. t. 6.—Ins. Azor.
Benthamiana, *Meissn. in DC. Prod.* xv. I. 44.—Bras.
Bofo, Spreng. Syst. ii. 269 = Ocotea Bofo.
Borbonia, Spreng. l. c. 268 = carolinensis.
brevipes, *Meissn. in DC. Prod.* xv. I. 46.—Ecuador.
caesia, *Meissn. l. c.* 44.—Bras.
Camfora, Spreng. Syst. ii. 268 = Cinnamomum Camphora.
canariensis, Spreng. l. c. = Apollonias canariensis.
canescens, Mart. ex Nees, Syst. Laurin. 280 = Nectandra canescens.
carolinensis, *Nees, l. c.* 150.—Am. bor.
caryophyllata, Mart. Reise Bras. iii. 915 = Dicypellium caryophyllatum.
Cassia, Spreng. Syst. ii. 267 = Cinnamomum Cassia.
Catesbyana, Chapm. Fl. S. U. St. 393 = Nectandra Willdenoviana.
Cathia, Spreng. Syst. iv. Cur. Post. 156 = Phoebe paniculata.
caustica, Spreng. Syst. ii. 269 = Mauria simplicifolia.
causticans, Pasq. Cat. Orto Bot. Nap. 77 (= *Tetranthera causticans*).—Hab.?
ceanothifolia, Mart. ex Nees, Syst. Laurin. 236 (= *Mespilodaphne ceanothifolia*).—Bras.
cinnamomifolia, Hemsl. Biol. Centr. Am. Bot. iii. 71 = Phoebe mexicana.
cinnamomifolia, H. B. & K. Nov. Gen. et Sp. ii. 160 = Phoebe cinnamomifolia.

PERSEA:—
Cinnamomum, Spreng. Syst. ii. 267 = Cinnamomum zeylanicum.
cordata, *Meissn. in DC. Prod.* xv. I. 54.—Bras.
costata, *Meissn. l. c.* 50.—Peruv.
crassifolia, Spreng. Syst. ii. 269 = Ocotea puberula.
Cubeba, Spreng. l. c. = Litsea Cubeba.
cubensis, *Meissn. in DC. Prod.* xv. I. 51.—Peruv.
cuneata, *Meissn. l. c.* 46.—N. Granat.
cupularis, Spreng. Syst. ii. 269 (= *Mespilodaphne cupularis*).—Ins. Mascar.
curvifolia, *Spreng. l. c.* 268.—Cochinch.
densiflora, *Meissn. in DC. Prod.* xv. I. 52.—Bras.
drymifolia, Cham. & Schlecht. in Linnaea, vi. (1831) 365 = gratissima.
dulcis, Spreng. Syst. ii. 268 = Cinnamomum Burmanni.
edulis, Rafin. Sylva Tellur. 134 = gratissima?
effusa, Benth. & Hook. f. Gen. iii. 158, in nota; et ex Hemsl. Biol. Centr. Am. Bot. iii. 71 = Phoebe effusa.
erythropus, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 49.—Bras.
exaltata, Spreng. Syst. ii. 268 = Nectandra sanguinea.
ferruginea, Benth. & Hook. f. Gen. iii. 159 (= Ocoteae sp.).—Peruv.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* ii. 159.—Peruv.
floribunda, Schott, in Spreng. Syst. iv. Cur. Post. 405 = Ocotea floribunda.
foetens, Spreng. Syst. ii. 268 = Ocotea foetens.
fuliginosa, *Nees, Syst. Laurin.* 156.—Bras.
globosa, Spreng. Syst. ii. 269 = Nectandra sanguinea.
grandis, Bartl. ex Meissn. in DC. Prod. xv. I. 146 = Nectandra rigida.
grandis, Nees, in Wall. Pl. As. Rar. iii. 32 = Phoebe opaca.
granulosa, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 265; *Contrib.* 71.—Reg. Argent.
gratissima, *Gaertn. f. Fruct.* iii. 222.—Am. trop.
Hartwegii, Hemsl. Biol. Centr. Am. Bot. iii. 72 = Phoebe Hartwegii.
hirta, *Nees, Syst. Laurin.* 673.—Peruv.
hypericifolia, *Nees, l. c.* 165.—Hab.?
hypoglauca, Herb. Monac. ex Meissn. in DC. Prod. xv. I. 49 = pyrifolia, scrobiculata.
hypoglauca, Mart. ex Nees, Syst. Laurin. 132 = venosa.
incana, Schott, in Spreng. Syst. iv. Cur. Post. 405 = Nectandra villosa.
incrassata, Nees, Syst. Laurin. 127 = Dehaasia microcarpa.
indecora, Schott, in Spreng. Syst. iv. Cur. Post. 405 = Ocotea indecora.
indica, *Spreng. Syst.* ii. 268.—Ins. Canar.
laevigata, *H. B. & K. Nov. Gen. et Sp.* ii. 157.—N. Granat.
lancifolia, Schott, in Spreng. Syst. iv. Cur. Post. 405 = Ocotea subacris.
lancifolia, Willd. ex Meissn. in DC. Prod. xv. I. 57 = Nectandra turbacensis.
latifolia, Spreng. Syst. ii. 270 = Nectandra polita.
Laurel-moroti, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 263; *Contrib.* 69.—Reg. Argent.
laxa, Mart. ex Nees, Syst. Laurin. 468, 469 (= *Oreodaphne venulosa*).—Bras.
leucantha, Mart. l. c. 310 = Nectandra leucantha.
Leucoxylon, Spreng. Syst. ii. 268 = Ocotea Leucoxylon.
Ligé, Nees, ex Meissn. in DC. Prod. xv. I. 48 = P. Lingue.
lineata, *Spreng. Syst.* ii. 270.—Venezuela.
Lingue, *Nees, Syst. Laurin.* 157.—Chili.
longifolia, Spreng. Syst. ii. 270 = Ocotea longifolia.
longifolia, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 344 = Actinodaphne Rumphii.
longipeda, *Bertol. Misc. Bot.* xiii. 14.—Am. bor.
longipes, *Meissn. in DC. Prod.* xv. I. 55.—Mexic.
macrocarpa, Rafin. in Med. Repos. N. York, v. (1808) 352 = gratissima.
macrophylla, Blume, Bijdr. 568 = Phoebe attenuata.
macrophylla, Spreng. Syst. ii. 270 = Nectandra macrophylla.
macropoda, H. B. & K. Nov. Gen. et Sp. ii. 160 = Ocotea velutina.
marginata, Bartl. ex Meissn. in DC. Prod. xv. I. 142 = Ocotea puberula.
membranacea, Spreng. Syst. ii. 270 = Nectandra saligna.

PERSEA :—
mexicana, Hemsl. Biol. Centr. Am. Bot. iii. 72 = Phoebe mexicana.
Meyeniana, *Nees, in Linnaea*, viii. (1833) 5.—Chili.
microneura, *Meissn. in DC. Prod.* xv. 1. 53.—Bras.
mollis, Spreng. Syst. ii. 270 = Nectandra villosa.
montana, Spreng. l. c. ii. 268 = Phoebe montana.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* ii. 158.—N. Granat.
Nam-mu, *Oliver, in Hook. Ic. Pl.* t. 1316.—China.
nitida, Spreng. Syst. iv. Cur. Post. 156 = Cinnamomum iners.
nutans, Nees, Syst. Laurin. 141 = Phoebe cubensis.
obovata, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 51.—Bras.
Orizabae, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 358.—Mexic.
pachypoda, *Nees, in Linnaea*, xxi. (1848) 490.—Mexic.
paniculigera, Mart. ex Nees, Syst. Laurin. 313 = Nectandra nitidula.
parviflora, Spreng. Syst. ii. 270 (= *Oreodaphne parviflora*).—Jamaic.
patens, Spreng. Syst. ii. 270 = Nectandra patens.
pedunculares, Nees, Syst. Laurin. 125 (= *Haasia pedunculares*).—Ind. or.
pedunculosa, *Meissn. in DC. Prod.* xv. 1. 47.—Bras.
peruviana, *Nees, Syst. Laurin.* 155.—Peruv.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* ii. 159.—Am. austr.
Pichurim, Spreng. Syst. ii. 269 = Aydendron Laurel.
praecox, Poepp. ex Nees, Syst. Laurin. 129 = gratissima.
psammophila, Mart. ex Meissn. in DC. Prod. xv. 1. 161 = Nectandra psammophila.
Pseudoindica, Link, Handb. i. 389 = indica.
Pseudo-sassafras, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, i. (1844) 615 = Cinnamomum Parthenoxylon.
psychotrioides, Spreng. Syst. ii. 269 = Ocotea psychotrioides.
puberula, Schott, in Spreng. l. c. iv. Cur. Post. 405 = Nectandra puberula.
punctata, *Meissn. in DC. Prod.* xv. 1. 50.—Bras.
pyrifolia, *Nees, Syst. Laurin.* 145.—Bras.
pyrifolia, Spreng. Syst. iv. Cur. Post. 156 = Cinnamomum pyrifolium.
retroflexa, Spreng. l. c. ii. 269 = Ocotea botryophylla.
Richardiana, Cham. & Schlecht. in Linnaea, vi. (1831) 366 = Ocotea puberula.
Riedelii, *Meissn. in DC. Prod.* xv. 1. 54.—Bras.
rigida, Meissn. l. c. 46 = rufotomentosa.
rigida, *Nees, Syst. Laurin.* 154.—Bras.
rimosa, Zoll. ex Meissn. in DC. Prod. xv. 1. 42 = Machilus rimosa.
rufotomentosa, *Nees & Mart. ex Nees, Syst. Laurin.* 153.—Bras.
rugosa, Meissn. in DC. Prod. xv. 1. 47 = hirta, pedunculosa.
salicifolia, Hemsl. Biol. Centr. Am. Bot. iii. 72 = Phoebe salicifolia.
salicifolia, Spreng. Syst. ii. 270 = Nectandra sanguinea.
sanguinea, Spreng. l. c. 268 = Nectandra sanguinea.
sarcocalyx, Buek, Ind. DC. Prod. iv. 253 = Nectandra nitidula.
Sassafras, Spreng. Syst. ii. 270 = Sassafras officinalis.
Schiedeana, Nees, Syst. Laurin. 130 = gratissima.
scrobiculata, *Meissn. in DC. Prod.* xv. 1. 50.—Bras.
Sellowiana, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 50.—Bras.
sericea, *H. B. & K. Nov. Gen. et Sp.* ii. 159.—Peruv.
sericea, Poepp. ex Meissn. in DC. Prod. xv. 1. 46 = peruviana.
splendens, *Meissn. l. c.* 44.—Bras.
stenophylla, *Meissn. l. c.* 55.—Bras.
subcordata, *Nees, in Linnaea*, xxi. (1848) 492.—Peruv.
subsessilis, *Meissn. in DC. Prod.* xv. 1. 54.—Peruv.
subtriplinervia, Mart. ex Nees, Syst. Laurin. 466 (= *Camphoromoea subtriplinervia*).—Bras.
sulcata, *Meissn. in DC. Prod.* xv. 1. 54.—Peruv.
sumatrana, *Teijsm. & Binn. Cat. Hort. Bog.* 93.—Sumatra.
surinamensis, Spreng. Syst. ii. 269 = Ocotea pulchella.

PERSEA :—
sylvestris, *A. Rich. Fl. Cub. Fanerog.* ii. 186.—Cuba.
tabacifolia, Meissn. in DC. Prod. xv. 1. 51 = Ocotea umbrosa.
Tamala, Spreng. Syst. ii. 268 = Cinnamomum Tamala.
Teneriffae, F. Muell. Sel. Pl. Indust. Cult. 159 = indica.
tenuiflora, Mart. ex Nees, Syst. Laurin. 359 (= *Leptodaphne tenuiflora*).—Bras.
tigricida, Spreng. Syst. ii. 270 = Nectandra discolor.
tomentosa, Spreng. l. c. iv. Cur. Post. 156 = Phoebe paniculata.
tubigera, Mart. ex Nees, Syst. Laurin. 480 = Nectandra psammophila.
umbrosa, Spreng. Syst. ii. 269 (= *Ocotea umbrosa*, H. B. & K.).—N. Granat.
velutina, Benth. & Hook. f. Gen. iii. 159 (Ocoteae sp.).—N. Granat.
venosa, Benth. ex Meissn. in DC. Prod. xv. 1. 44 = Benthamiana.
venosa, *Nees & Mart. ex Nees, in Linnaea*, viii. (1833) 50.—Bras.
veraguasensis, *Seem. Bot. Voy. Herald*, 193.—Panama.
veraguensis, Meissn. in DC. Prod. xv. 1. 51 = praec.
viburnoides, *Meissn. l. c.* 52.—Bras.
Yacupeti, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 264; *Contrib.* 70.—Reg. Argent.

PERSICA, [Tourn.] Mill. Gard. Dict. ed. VI. (1752) = **Prunus**, Linn. (Rosac.).
Amygdalus, Mill. l. c. ed. VIII. n. 3 = Pr. Persica.
communis, Duham. ex Steud. Nom. ed. II. i. 81, in syn. = Pr. Persica.
Davidiana, Carr. in Rev. Hortic. (1872) 74 = Pr. Persica.
domestica, Risso, Hist. Nat. ii. 104 = Pr. Persica.
ispahamensis, Thouin, in Ann. Mus. Par. viii. (1806) 433 = Pr. Persica.
laevis, DC. Fl. Fr. iv. 487 = Pr. Persica.
nana, Mill. Gard. Dict. ed. VIII. n. 2 (= *Amygdalus nana*).—Europ.; Reg. Cauc.; Sibir.
necturina, Steud. Nom. ed. II. i. 81, in syn. = Pr. Persica.
nucicarpa, Steud. l. c. = Pr. Persica.
Nucipersica, Borckh. Hess. Hoelz. 205; ex Roem. Arch. i. II. (18) 37 = Pr. Persica.
Pavia, Delarb. Fl. Auv. ed. II. 328 = Pr. Persica.
pumila, Steud. Nom. ed. II. ii. 307 = Pr. sinensis.
violacea, Risso, Hist. Nat. ii. 119 = Pr. Persica.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = Pr. Persica.

PERSICANA, Tourn. ex Scop. Introd. 344 (1777), sphalm. = PERSICARIA, Tourn.

PERSICARIA, Neck. Elem. ii 210 (1790) = **Atraphaxis**, Linn. (Polygon.).

PERSICARIA, [Tourn.] Linn. Syst. ed. I (1735) = **Polygonum**, Tourn.
acris, Gilib. Exercit. ii. 431 = Polyg. Hydropiper.
amphibia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 268 = Polyg. amphibium.
Brittingeri, Opiz, in Lotos, iv. (1854) 42 = Polyg. lapathifolium.
dubia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 146 = Polyg. mite.
Fischeri, Opiz, in Lotos, iv. (1854) 43 (Quid?).—Europ.
Hydropiper, Opiz, Seznam, 72 = Polyg. Hydropiper.
incana, S. F. Gray, Nat. Arr. Brit. Pl. ii. 269 = Polyg. Persicaria.
interrupta, S. F. Gray, l. c. 732 = Polyg. Persicaria.
lapathifolia, S. F. Gray, l. c. 270 = Polyg. lapathifolium.
laxiflora, Opiz, Seznam, 72 = Polyg. mite.
Lorinseri, Opiz, l. c. 73 = Polyg. Lorinseri.
maculata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 270 = Polyg. Persicaria.
maculosa, S. F. Gray, l. c. 269 = Polyg. Persicaria.
minor, Opiz, Seznam, 72 = Polyg. minus.
mitis, Gilib. Exercit. ii. 431 = Polyg. Persicaria.
nodosa, Opiz, Seznam, 72 = Polyg. lapathifolium.
orientalis, Vilm. Fl. Pl. Terre, ed. II. 660 = Polyg. orientale.

PERSICARIA :—
polystachya, Opiz, in Lotos, ii. (1852) 229 (Quid?).—Europ.
pusilla, S. F. Gray, Nat. Arr. Brit. Pl. ii. 269 = Polyg. Persicaria.
rivularis, Opiz, Seznam, 72 = Polyg. Persicaria.
salicifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 270 = Polyg. Persicaria.
serrulata, Webb & Moq. in Webb & Berth. Phyt. Canar. iii. 219 = Polyg. serrulatum.
virginiana, Gaertn. Fruct. ii. 180 = Polyg. virginianum.
vulgaris, Webb & Moq. in Webb & Berth. Phyt. Canar. iii. 219 = Polyg. Persicaria.

PERSIMON, Rafin. Sylva Tellur. 164 (1838) = **Diospyros**, Linn. (Ebenac.).
virginiana, Rafin. l. c. = D. virginiana.

PERSONARIA, Lam. Illustr. t. 716 (1798?) = **Gorteria**, Linn. (Compos.).
personata, Lam. ex Poir. Illustr. iii. 291 (1823) = G. personata.

PERSONULA, Rafin. Fl. Tellur. iv. 110 (1836) = **Utricularia**, Linn. (Lentib.).
grandiflora, Rafin. l. c. = U. juncea.

PERSOONIA, Michx. Fl. Bor. Am. ii. 104. t. 43 (1805) = **Marshallia**, Schreb. (Compos.).
angustifolia, Michx. l. c. 106 = M. angustifolia.
lanceolata, Michx. l. c. 105 = M. lanceolata.
latifolia, Michx. l. c. = M. latifolia.

PERSOONIA, Sm. in Trans. Linn. Soc. iv. (1798) 215.
PROTEACEAE, Benth. & Hook. f. iii. 176.
LINKIA, Cav. Ic. iv. 61. t. 389 (1797).
PENTADACTYLON, Gaertn f. Fruct. iii. 219. t. 220 (1805)
abietina, A. Cunn. ex Meissn. in DC. Prod. xiv. 336 = curvifolia.
acerosa, *Sieber, ex Schult. Mant.* iii. 269.—Austral.
acicularis, *F. Muell. Fragm.* vi. 220.—Austral.
angulata, *R. Br. Prot. Nov.* 14.—Austral.
angustiflora, *Benth. Fl. Austral.* v. 386.—Austral.
angustifolia, Knight, Prot. 99 = linearis.
apiculata, Meissn. in Hook. Kew Journ. vii. (1855) 73 = nutans.
arborea, *F. Muell. Fragm.* v. 37.—Austral.
arida, Sieber, ex Schult. Mant. iii. 270; et Spreng. Syst. iv. Cur. Post. 45 = hirsuta.
articulata, *R. Br. in Trans. Linn. Soc.* x. (1810) 164.—Austral.
aspera, A. Cunn. ex Benth. Fl. Austral. v. 394 = hirsuta.
attenuata, R. Br. Prot. Nov. 16 = media *vel* lanceolata.
brachystylis, *F. Muell. Fragm.* vi. 221.—Austral.
brevifolia, Loud. Hort. Brit. Suppl. iii. 605 (Quid?).—Austral.
Caleyi, *R. Br. Prot. Nov.* 13.—Austral.
Caleyi, F. Muell. Pl. Vict. ii. t. 69 = P. Chamaepeuce.
Chamaepeuce, *Lhotsky, ex Meissn. in DC. Prod.* xiv. 336.—Austral.
Chamaepitys, *A. Cunn. in Field, N. S. Wales*, 329.—Austral.
comata, *Meissn. in Hook. Kew Journ.* vii. (1855) 71.—Austral.
confertifolia, *Benth. Fl. Austral.* v. 396.—Austral.
cornifolia, *A. Cunn. ex R. Br. Prot. Nov.* 16.—Austral.
Cunninghamii, *R. Br. l. c.* 15.—Austral.
curvifolia, *R. Br. l. c.* 13.—Austral.
daphnoides, A. Cunn. ex R. Br. l. c. 15 = prostrata.
diadena, *F. Muell. Fragm.* x. 46.—Austral.
dillwynioides, *Meissn. in DC. Prod.* xiv. 333.—Austral.
Drummondii, Lindl. Swan River App. 35 = longifolia.
elliptica, *R. Br. in Trans. Linn. Soc.* x. (1810) 164.—Austral.
falcata, *R. Br. l. c.* 162.—Austral.
fastigiata, *R. Br. Prot. Nov.* 13.—Austral.
fastigiata, A. Cunn. ex Meissn. in DC. Prod. xiv. 336 = curvifolia.
ferruginea, *Sm. Exot. Bot.* ii. 47. t. 83.—Austral.
filifolia, F. G. Dietr. Vollst. Lexik. Gaertn. vii. 100 = linearis.
flexifolia, R. Br. in Trans. Linn. Soc. x. (1810) 162 = scabra.

PERSOONIA:—
flexifolia, Lodd. Bot. Cab. t. 922 = nutans.
Fraseri, R. Br. Prot. Nov. 14 = saccata.
Fraseri, Meissn. in Lehm. Pl. Preiss. i. 532 = angustiflora.
fusca, Hort. ex Steud. Nom. ed. II. ii. 307 (Quid?).—Austral.
glauca, *Sm. ex Spreng. Syst.* i. 473.—Austral.
glaucescens, Sieber, ex Schult. Mant. iii. 271 = lanceolata.
gnidioides, Sieber, l. c. 269; et Spreng. Syst. iv. Cur. Post. 45 = P. Chamaepitys.
graminea, *R. Br. in Trans. Linn. Soc.* x. (1810) 164.—Austral.
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 283.—Tasmania.
hakeaeformis, *Meissn. in DC. Prod.* xiv. 330.—Austral.
heterophylla, Loud. Hort. Brit. Suppl. iii. 605 (Quid?).—Austral.
hirsuta, A. Cunn. ex Meissn. in DC. Prod. xiv. 339 = mollis.
hirsuta, Knight, Prot. 99 (= seq.?).—Austral.
hirsuta, *Pers. Syn.* i. 118.—Austral.
juniperina, *Labill. Nov. Holl. Pl.* i. 33. t. 45.—Austral.
lanceolata, *Andr. Bot. Rep.* t. 74.—Austral.
lanceolata, A. Cunn. ex Meissn. in DC. Prod. xiv. 343 = salicina.
latifolia, Andr. Bot. Rep. t. 280 = lanceolata.
latifolia, Knight, Prot. 100 = salicina.
Laureola, Lindl. Swan River App. 35 = elliptica.
laurina, Pers. Syn. i. 118 = ferruginea.
ledifolia, *A. Cunn. ex Meissn. in DC. Prod.* xiv. 339.—Austral.
ledifolia, Nois. ex Schult. Mant. iii. 273 (Quid?).—Hab.?
ligustrina, Knight, Prot. 100 = lanceolata.
linariifolia, A. Cunn. ex Benth. Fl. Austral. v. 402 = virgata.
linearis, *Andr. Bot. Rep.* t. 77.—Austral.
linearis, Sieber, ex Schult. Mant. iii. 271 = nutans.
longifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 164.—Austral.
lucida, *R. Br. l. c.* 161.—Austral.
macrostachya, Lindl. Swan River App. 35 = saccata.
marginata, *A. Cunn. ex R. Br. Prot. Nov.* 16.—Austral.
media, *R. Br. l. c.*—Austral.
microcarpa, *R. Br. in Trans. Linn. Soc.* x. (1810) 160.—Austral.
microphylla, R. Br. Prot. Nov. 15 = oxycoccoides.
mimosoides, A. Cunn. ex Benth. Fl. Austral. v. 386 = falcata.
Mitchellii, *Meissn. in Hook. Kew Journ.* vii. (1855) 73.—Austral.
mollis, *R. Br. in Trans. Linn. Soc.* x. (1810) 161.—Austral.
mollis, A. Cunn. ex Meissn. in DC. Prod. xiv. 337 = hirsuta.
Myrsinites, A. Cunn. l. c. 339 = mollis.
myrtilloides, *Sieber, ex Schult. Mant.* iii. 272.—Austral.
nutans, *R. Br. in Trans. Linn. Soc.* x. (1810) 162.—Austral.
obcordata, A. Cunn. ex Benth. Fl. Austral. v. 393 = marginata.
oblongata, *A. Cunn. ex R. Br. Prot. Nov.* 15.—Austral.
oleifolia, A. Cunn. ex Endl. Gen. Suppl. iv. II. 82 = myrtilloides.
oxycoccoides, *Sieber, ex Schult. Mant.* iii. 270; *et Spreng. Syst.* iv. *Cur. Post.* 45.—Austral.
pallida, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1828) 177 = acerosa.
pentadactylon, Steud. Nom. ed. II. ii. 308 = linearis.
pinifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 160.—Austral.
pinifolia, Sieber, ex Schult. Mant. iii. 271 = linearis.
planifolia, A. Cunn. ex Benth. Fl. Austral. v. 400 = oblongata.
prostrata, *R. Br. in Trans. Linn. Soc.* x. (1810) 163.—Austral.
pruinosa, A. Cunn. ex Steud. Nom. ed. II. ii. 308 = linearis.
pubes, F. Muell. ex Meissn. in DC. Prod. xiv. 336 = juniperina.
quinquenervis, *Hook. Ic. Pl.* t. 425.—Austral.
revoluta, *Sieber, ex Schult. Mant.* iii. 272.—Austral.
rigida, *R. Br. Prot. Nov.* 14.—Austral.

PERSOONIA:—
rudis, *Meissn. in Hook. Kew Journ.* vii. (1855) 73.—Austral.
rufiflora, *Meissn. l. c.* 72.—Austral.
saccata, *R. Br. Prot. Nov.* 12.—Austral.
salicina, *Pers. Syn.* i. 118.—Austral.
Saundersiana, *Kippist, ex Meissn. in Hook. Kew Journ.* vii. (1855) 72.—Austral.
scabra, *R. Br. in Trans. Linn. Soc.* x. (1810) 162.—Austral.
scabrella, *Meissn. in Hook. Kew Journ.* vii. (1855) 72.—Austral.
scoparia, Meissn. in DC. Prod. xiv. 329 = teretifolia.
sericea, *A. Cunn. ex R. Br. Prot. Nov.* 13.—Austral.
spathulata, R. Br. in Trans. Linn. Soc. x. (1810) 162 = scabra.
spathulata, Sieber, ex Schult. Mant. iii. 271 = rigida.
Strangeana, Kippist, ex Meissn. in DC. Prod. xiv. 334 = tenuifolia.
striata, *R. Br. Prot. Nov.* 13.—Austral.
striolata, Meissn. in Hook. Kew Journ. vii. (1855) 72 = striata.
suffruticosa, F. Muell. 1st Gen. Rep. 17 = P. Chamaepeuce.
sulcata, *Meissn. in DC. Prod.* xiv. 333.—Austral.
surrecta, F. Muell. ex Meissn. in DC. Prod. xiv. 336 = juniperina.
tenuifolia, *R. Br. Prot. Nov.* 12.—Austral.
tenuifolia, Meissn. in DC. Prod. xiv. 334 = virgata.
teretifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 160.—Austral.
thymifolia, A. Cunn. ex R. Br. Prot. Nov. 15 = oxycoccoides.
tinifolia, A. Cunn. ex Benth. Fl. Austral. v. 392 = cornifolia.
Tora, A. Cunn. in Ann. Hist. Nat. Ser. I. i. (1838) 378 = P. Toru.
tortifolia, *Meissn. in Hook. Kew Journ.* iv. (1852) 185.—Austral.
Toru, *A. Cunn. in Bot. Mag.* sub t. 3513.—N. Zel.
trinervis, *Meissn. in DC. Prod.* xiv. 332.—Austral.
velutina, A. Cunn. ex Meissn. in Lehm. Pl. Preiss. i. 534 = sericea.
virgata, *R. Br. in Trans. Linn. Soc.* x. (1810) 161.—Austral.

PERSOONIA, Willd. Sp. Pl. ii. 331 (1799) = **Carapa**, Aubl. (Meliac.).
guareoides, Willd. l. c. = C. guianensis.

PERTYA, Sch. Bip. in Bonplandia, x. (1862) 109. t. 10. *COMPOSITAE*, Benth. & Hook. f. ii. 493.
Aitchisoni, *C. B. Clarke, in Journ. Linn. Soc.* xviii. (1880) 72.—Afghan.
glabrescens, Sch. Bip. in Bonplandia, x. (1862) 109 = scandens.
ovata, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 217.—Japon.
paradoxa, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 153, nomen = Lactuca paradoxa.
scandens, *Sch. Bip. in Bonplandia*, x. (1862) 109.—Japon.

PERULA, Rafin. Sylva Tellur. 58 (1838) = **Ficus**, Tourn. (Urticac.).
benghalensis, Rafin. l. c. 59 = F. benghalensis.
retusa, Rafin. l. c. = F. retusa.
rubiginosa, Rafin. l. c. = F. rubiginosa.

PERULA, Schreb. Gen. ii. 703 (1791) = **Pera**, Mutis (Compos.).
arborea, Gaudich. ex Baill. Etud. Gén. Euph. 434 = Pera ferruginea.
arborea, Willd. Sp. Pl. iv. 836 = Pera arborea.

PERULARIA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Habenaria**, Willd. (Orchid.).
fuscescens, Lindl. l. c. = H. herbiola.

PERVILLAEA, Decne. in DC. Prod. viii. 613 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 774.
tomentosa, *Decne. l. c.*—Ins. Maurit.

PERVINCA, Tourn. ex Adans. Fam. ii. 172 (1763) = **Vinca**, Linn. (Apocyn.).
major, Scop. Fl. Carn. ed. II. i. 170 = V. major.
minor, Scop. l. c. = V. minor.
procumbens, Gilib. Exercit. i. 63 = V. minor.
rosea, Moench, Meth. 463 = V. rosea.

PERYMENIUM, Schrad. Ind. Sem. Hort. Gotting. (1830).—Cf. Linnaea, vi. (1831) Litt. 73. *COMPOSITAE*, Benth. & Hook. f. ii. 377.
Barclayanum, *DC. Prod.* v. 609.—Mexic.
Berlandierii, *DC. l. c.* 608.—Mexic.
buphthalmoides, *DC. l. c.* 609.—Mexic.
Cervantesii, *DC. l. c.*—Mexic.
discolor, *Schrad. Ind. Sem. Hort. Gotting.* (1830).—*Cf. Linnaea*, vi. (1831) *Litt.* 73.—Mexic.
gracile, *Hemsl. Biol. Centr. Am. Bot.* ii. 181.—Mexic.
grande, *Hemsl. l. c.*—Guatemala.
gymnolomoides, *DC. Prod.* v. 609.—Mexic.
Mendezii, *DC. l. c.* 608.—Mexic.
parvifolium, *A. Gray, in Proc. Am. Acad.* xv. (1880) 36.—Mexic.
tenellum, *A. Gray, l. c.*—Mexic.
Tuerckheimii, *Klatt, in Leopoldina*, xx. (1884) 95.—Mexic.
verbesinoides, *DC. Prod.* v. 608.—Mexic.

PERYTIS, Rafin. Alsog. Am. 29 (1838) = **Quercus**, Tourn. (Cupul.).
annulata, Rafin. l. c. = Q. annulata.
concentrica, Rafin. l. c. = Q. concentrica.
glauca, Rafin. l. c. = Q. glauca.
lamellosa, Rafin. l. c. = Q. lamellosa.

PESCATOREA, Reichb. f. in Gard. Chron. (1869) 710 = seq.

PESCATORIA, Reichb. f. in Bot. Zeit. x. (1852) 667 = **Zygopetalum**, Hook. (Orchid.).
Backhousiana, Reichb. f. in Gard. Chron. (1877) II. 456 = Z. Backhousianum.
bella, Reichb. f. l. c. (1878) I. 492 = Z. bellum.
cerina, Reichb. f. in Bot. Zeit. x. (1852) 667; et Xenia Orch. i. 184. t. 65 = Z. cerinum.
coronaria, Reichb. f. in Linnaea, xli. (1877) 108 = Z. coronarium.
Dayana, Reichb. f. in Gard. Chron. (1872) 1618 = Z. Dayanum.
Dormaniana, Reichb. f. l. c. (1881) I. 330.—N. Granat.
euglossa, Reichb. f. l. c. (1876) II. 808 = Z. Roezlii.
fimbrata, Regel, in Act. Hort. Petrop. vi. (1879) 293; et in Gartenfl. (1880) 129. t. 1008.—N. Granat.?
Gairiana, Reichb. f. in Gard. Chron. (1879) I. 684 = Z. Gairianum.
Klabochorum, Reichb. f. l. c. = Z. Klabochorum.
lamellosa, Reichb. f. l. c. (1875) II. 225 = Z. lamellosum.
Lehmanni, Reichb. f. l. c. (1879) II. 424 = Z. Lehmanni.
Roezlii, Reichb. f. l. c. (1874) I. 755 = Z. Roezlii.
Rueckeriana, Reichb. f. l. c. (1885) II. 424.—Hab.?
Russeliana, Reichb. f. l. c. (1878) II. 524 = Z. Russelianum.
triumphans, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 97 = Z. triumphans.
Vervaeti, Hort. ex T. Moore, Fl. & Pom. (1883) 10.—Hab.?
Wallisii, Linden & Reichb. f. in Gard. Chron. (1869) 710 = Z. Wallisii.

PESCHIERA, A. DC. in DC. Prod. viii. 360 (1844) = **Tabernaemontana**, Plum. (Apocyn.).
acuminata, Miers, Apocyn. S. Am. 43 = T. acuminata.
affinis, Miers, l. c. 40 = T. affinis.
albidiflora, Miers, l. c. 39.—Bras.
albiflora, Miq. Stirp. Surin. Sel. 165.—Guian.
australis, Miers, Apocyn. S. Am. 46 = T. australis.
blanda, Miers, l. c. 44.—Peruv.
brevifolia, Miers, l. c. 45 = T. breviflora.
catharinensis, Miers, l. c. 41 = T. catharinensis.
concinna, Miers, l. c. 44.—Peruv.
cuspidata, Miers, l. c. 37.—N. Granat.
diversifolia, Miq. Stirp. Surin. Sel. 164 = T. heterophylla, *Vahl.*
echinata, A. DC. in DC. Prod. viii. 360 = T. echinata.
fallax, Miers, Apocyn. S. Am. 40 = T. fallax.
florida, Miers, l. c. 41 = T. laeta.
fuchsiaefolia, Miers, l. c. 34 = T. fuchsiaefolia.
Gaudichaudii, Miers, l. c. 40 = T. Gaudichaudii.
gracillima, Miers, l. c. 41 = T. gracilis.
granulosa, Miers, l. c. 37.—Bras.
heterophylla, Miers, l. c. 38 = T. heterophylla, *Vahl.*
Hilariana, Miers, l. c. 41 = T. Hilariana.

PESCHIERA:—
Hystrix, A. DC. in DC. Prod. viii. 360 = Tabernaemontana Hystrix.
laeta, Miers, Apocyn. S. Am. 35 = T. laeta.
latiflora, Benth. ex Miers, l. c. 50 = T. flavicans.
lingulata, Miers, l. c. 42.—Peruv.
Linkii, Miers, l. c. 47 = T. Linkii.
litoralis, Miers, l. c. 45 = T. litoralis.
lorifera, Miers, l. c. 47.—Guian.
Lundii, Miers, l. c. 36 = T. Lundii.
multiflora, Spruce, ex Miers, l. c. 45 = T. Benthamiana.
muricata, A. DC. in DC. Prod. viii. 361 = T. muricata.
ochracea, Miers, Apocyn. S. Am. 42 = T. muricata.
praeclara, Miers, l. c. 47.—Venezuela.
psychotriaefolia, Miers, l. c. 42 = T. psychotrifolia.
puberiflora, Miers, l. c. 43.—Peruv.
Salzmanni, Miers, l. c. 40 = T. Salzmanni.
Solandri, Miers, l. c. 46 = T. cymosa?
solanifolia, Miers, l. c. 46 = T. solanifolia.
Spixiana, Miers, l. c. 36 = T. Spixiana.
stenoloba, Miers, l. c. 38 = T. stenoloba.
surinamensis, Miq. in Linnaea, xviii. (1844) 742.—Guian.
tenuiflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 70. t. 280 = T. tenuiflora, *Muell. Arg.*
umbrosa, Miers, Apocyn. S. Am. 44 = T. umbrosa.

PESOMERIA, Lindl. Bot. Reg. (1838) Misc. 5 = **Phaius**, Lour. (Orchid.).
tetragona, Lindl. l. c. = Ph. villosus.

PESSULARIA, Salisb. Gen. Pl. Fragm. 70 (1866) = **Anthericum**, Linn. (Liliac.).
ramosa, Salisb. l. c. = A. ramosum.

PESTALOZZIA, Zoll. & Mor. Syst. Verz. Zoll. 31 (1846) = **Gynostemma**, Blume (Cucurb.).
laxa, Thw. Enum. Pl. Zeyl. 124 = G. Wightianum.
pedata, Zoll. & Mor. Syst. Verz. Zoll. 31 = G. pedatum.

PETAGNANA, J. F. Gmel. Syst. 1078 (1791) = **Smithia**, Ait. (Legumin.).
sensitiva, J. F. Gmel. l. c. 1119 = S. sensitiva.

PETAGNIA, Guss. Fl. Sic. Prod. i. 311 (1826). *UMBELLIFERAE*, Benth. & Hook. f. i. 881.
HETEROSCIADIUM, DC. Prod. iv. 83, in syn. (1830).
saniculaefolia, *Guss. Fl. Sic. Prod.* i. 311.—Sicil.

PETALACTE, D. Don, in Mem. Wern. Soc. v. (1826) 522, ex syn. cit. *COMPOSITAE*, Benth. & Hook. f. ii. 314.
PETALOLEPIS, Less. Syn. Comp. 357 (1832).
bicolor, D. Don, in Mem. Wern. Soc. v. (1826) 553 = coronata.
canescens. DC. Prod. vi. 267 = Anaxeton canescens (in Addend.).
coronata, *D. Don, in Mem. Wern. Soc.* v. (1826) 553.—Afr. austr.

PETALANDRA, Hassk. Hort. Bog. Desc. 104 (1858) = **Hopea**, Roxb. (Dipteroc.).
micrantha, Hassk. l. c. 105 = H. micrantha.

PETALANDRA, F. Muell. ex Boiss. in DC. Prod. xv. II. 27, in syn. (1862) = **Euphorbia**, Linn.
euphorbioides, F. Muell. ex Boiss. l. c. = E. micradenia.

PETALANISIA, Rafin. Fl. Tellur. iii. 80 (1836) = **Hypericum**, Linn.

PETALANTHERA, Nees, Laurin. Disp. 15 (1833); Syst. Laurin. 346 (1836) = **Ocotea**, Aubl. (Laurin.).
bracteolata, Nees, Syst. Laurin. 347 = O. bracteolata.

PETALANTHERA, Nutt. in Journ. Acad. Philad. vii. I. (1834) 107 = **Cevallia**, Lag. (Loasac.).
hispida, Nutt. l. c. = C. sinuata.

PETALANTHERA, Rafin. Fl. Tellur. ii. 100 (1836) = **Justicia**, Houst. (Acanth.).
punctata, Rafin. l. c. = J. ventricosa.

PETALIDIUM, Nees, in Wall. Pl. As. Rar. iii. 75 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1081.
PSEUDOBARLERIA, T. Anders. in Journ. Linn. Soc. vii. (1863) 26.
barlerioides, *Nees, in Wall. Pl. As. Rar.* iii. 82.—Ind. or.
coccineum, *S. Moore, in Journ. Bot.* xviii. (1880) 225.—Afr. trop.
glandulosum, *S. Moore, l. c.* 226.—Afr. trop.
halimoides, *S. Moore, l. c.* 228.—Afr. trop.
Lepidagathis, *S. Moore, l. c.* 227.—Afr. trop.
linifolium, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 25.—Afr. austr.
loranthifolium, *S. Moore, in Journ. Bot.* xviii. (1880) 227.—Afr. trop.
patulum, Dalz. & Gibs. Bomb. Fl. 185 = Ruellia patula.
physaloides, *S. Moore, in Journ. Bot.* xviii. (1880) 225.—Afr. trop.
rupestre, *S. Moore, l. c.* 226.—Afr. trop.
Welwitschii, *S. Moore, l. c.* 227.—Afr. trop.

PETALINIA, Becc. Malesia, i. (1883) 257. *OLACINEAE*.
bancana, *Becc. l. c.* 258.—Ins. Banca.

PETALODISCUS, Baill. Etud. Gén. Euph. 571. t. 22 (1858) = **Savia**, Willd. (Euphorb.).
Laureola, Baill. l. c. 572 = S. Laureola.
platyrachis, Baill. l. c. = S. platyrachis.

PETALOGYNE, F. Muell. in Hook. Kew Journ. viii. (1856) 324 = **Petalostylis**, R. Br. (Legumin.).
cassioides, F. Muell. l. c. = Petalostylis labicheoides.
labicheoides, F. Muell. l. c. 325 = Petalostylis labicheoides.

PETALOLEPIS, Cass. in Bull. Soc. Philom. (1817) 134 = **Helichrysum**, Vaill. (Compos.).
ferruginea, Cass. in Dict. Sc. Nat. xxix. 195 = Helipterum citrinum.
rosmarinifolia, Cass. l. c. = Helichrysum diosmifolium.

PETALOLEPIS, Less. Syn. Comp. 357 (1832) = **Petalacte**, D. Don (Compos.).
bicolor, Less. l. c. 358 = Petalacte coronata.
coronata, Less. l. c. 357 = Petalacte coronata.
discolor, Less. l. c. 358 = Petalacte coronata.

PETALOMA, Rafin. ex Boiss. in DC. Prod. xv. II. 63 (1862) = **Euphorbia**, Linn.

PETALOMA, Roxb. Hort. Beng. [90] (1814); Fl. Ind. ii. 372 (1832) = **Lumnitzera**, Willd. (Combret.).
alba, Blanco, Fl. Filip. ed. I. 344 = L. racemosa.
albiflora, Zipp. ex Span. in Linnaea, xv. (1841) 203 = L. racemosa.
alternifolia, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 372 = L. racemosa.
coccinea, Blanco, Fl. Filip. ed. I. 345 = L. coccinea.

PETALOMA, Sw. Prod. Veg. Ind. Occ. 73 (1788); Fl. Ind. Occ. ii. 831 (1800) = **Mouriria**, Juss. (Melastom.).
cauliflora, Mart. ex DC. Prod. iii. 7 = M. cauliflora.
domingensis, Tussac, Fl. Antill. iii. 119. t. 37 = M. domingensis.
grandiflora, D. Dietr. Syn. Pl. ii. 1472 = M. grandiflora.
mexicana, D. Dietr. l. c. = M. mexicana.
Mouriri, Sw. Prod. Veg. Ind. Occ. 73 = M. guianensis.
myrtilloides, Sw. l. c. = M. myrtilloides.

PETALONYX, A. Gray, in Mem. Am. Acad. v. (1855) 319. *LOASEAE*, Benth. & Hook. f. i. 803.
crenatus, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 358.—Mexic.
linearis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 188.—Calif.
nitidus, *S. Wats. in Am. Natural.* vii. (1873) 300.—Utah.
Parryi, A. Gray, in Proc. Am. Acad. x. (1874) 72 = nitidus.
Thurberi, *A. Gray, in Mem. Acad. Am.* N. S. v. (1855) 319.—Mexic.

PETALOPOGON, Reiss. in Endl. Nov. Stirp. Dec. 82 (1839) = **Phylica**, Linn. (Rhamn.).
cuspidatus, Reiss. l. c. 83 = Phylica cuspidata.

PETALOSTEIRA, Rafin. Fl. Tellur. ii. 74 (1836) = **Tiarella**, Linn. (Saxifr.).
laciniata, Rafin. l. c. = T. trifoliata.
unifolia, Rafin. l. c. = T. unifoliata.

PETALOSTELMA, Fourn. in Mart. Fl. Bras. vi. IV. 328 (1885). *ASCLEPIADEAE*.
Martianum, *Fourn. l. c.*—Bras.

PETALOSTEMMA, R. Br. in Salt, Abyss. App. 44 (1814), nomen (Quid?).
Chenopodii, R. Br. l. c., nomen.—Abyssin.

PETALOSTEMON, Michx. Fl. Bor. Am. ii. 48. t. 37 (1803) (Petalostemum). *LEGUMINOSAE*, Benth. & Hook. f. i. 493.
CYLIPOGON, Rafin. in Journ. Phys. lxxxix. (1819) 97.
DALEA, Cram. Disp. Syst. 154 (1803).
GATESIA, Bertol. Misc. Bot. vii. 30. t. 1 (1848).
GAVESIA, Walp. Ann. ii. 950 (1852).
KUHNISTERA, Lam. Encyc. iii. 370 (1789).
alopecuroides, Pers. Syn. ii. 268 = Dalea alopecuroides.
amethystinus, *Schott, in Wien. Zeitschr.* iv. (1830) 1216; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 55.—Bras.
bicolor, Bertol. Misc. Bot. ix. 9 = gracilis.
candidus, *Michx. Fl. Bor. Am.* ii. 49. t. 37.—Am. bor.
capitatus, DC. Prod. ii. 244 = Dalea aurea.
carneus, *Michx. Fl. Bor. Am.* ii. 49.—Am. bor.
corymbosus, *Michx. l. c.* 50.—Am. bor.
decumbens, *Nutt. in Journ. Acad. Philad.* vii. I. (1834) 93.—Am. bor. occ.
emarginatus, *Torr & Gray, Fl. N. Am.* i. 311.—Am. bor. occ.
exilis, *A. Gray, Pl. Wright.* ii. 41.—N. Mexic.
Feayi, *Chapm. Fl. S. U. St. Suppl.* 615.—Florida.
flavescens, *S. Wats. in Am. Natural.* vii. (1873) 299.—Am. bor. occ.
foliosus, *A. Gray in Proc. Am. Acad.* vii. (1868) 336.—Am. bor.
gracilis, *Nutt. in Journ. Acad. Philad.* vii. I. (1834) 92.—Am. bor. occ.
griseus, *Torr. & Gray, Fl. N. Am.* i. 310.—Am. bor. occ.
laxiflorus, Steud. Nom. ed. II. ii. 308 = Dalea laxiflora.
macrostachyus, *Torr. in Ann. Lyc. N. York*, ii. (1828) 176.—Am. bor. occ.
macrostachyus, Torr. & Gray, Fl. N. Am. i. 309, partim = ornatus.
multiflorus, *Nutt. in Journ. Acad. Philad.* vii. I. (1834) 92.—Am. bor. occ.
obovatus, *Torr. & Gray, Fl. N. Am.* i. 310.—Am. bor. occ.
ornatus, *Dougl. ex Hook. Fl. Bor. Am.* i. 138.—Am. bor. occ.
ovatus, S. Wats. in Am. Natural. vii. (1873) 300, sphalm., in obs. = praec.
phleoides, *Torr. & Gray, Fl. N. Am.* i. 310.—Am. bor. occ.
reclinatus, Steud. Nom. ed. II. ii. 408, in syn. = Dalea reclinata.
roseus, *Nutt. in Am. Journ. Sc.* Ser. I. v. (1822) 298.—Florida.
Searlsiae, *A. Gray, in Proc. Am. Acad.* vii. (1872) 380.—Am. bor. occ.
sessilis, [Moç. & Sesse] ex DC. Prod. ii. 246 = Dalea Thouini.
tenuifolius, *A. Gray, in Proc. Am. Acad.* xi. (1876) 73.—N. Mexic.
villosus, *Nutt. Gen. Am.* ii. 85.—Am. bor.
violaceus, *Michx. Fl. Bor. Am.* ii. 50. t. 37. f. 2.—Am. bor. occ.
virgatus, Nees & Schwein. in Neuwied, Reise Nord-Am. ii. 432 = candidum.
virgatus, Scheele, in Linnaea, xxi. (1848) 461 = violaceus.

PETALOSTIGMA, F. Muell. in Hook. Kew Journ. ix. (1857) 16. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 277.
HYLOCOCCUS, R. Br. ex Benth. Fl. Austral. vi. 92 (1873).
australianum, Baill. Adansonia, vii. (1866–67) 356. t. 2 = quadriloculare.
quadriloculare, *F. Muell. in Hook. Kew Journ.* ix. (1857) 17.—Austral.
triloculare, Muell. Arg. in Flora, xlvii. (1864) 471 = quadriloculare.

PETALOSTIMA, Rafin. Fl. Tellur. ii. 79 (1836) = **Wahlenbergia**, Schrad. (Campan.).
capensis, Rafin. l. c. = W. capensis.

PETALOSTYLES, Benth. Fl. Austral. ii. 291 (1864) = seq.

PETALOSTYLIS, R. Br. in Sturt, Exped. Centr. Austral. App. ii. 79 (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 573.
PETALOGYNE, F. Muell. in Hook. Kew Journ. viii. (1856) 324.
labicheoides, *R. Br. in Sturt, Exped. Centr. Austr.* 80. —Austral.

PETALOSTYLIS, Lindl. Veg. Kingd. 614 (1847) = PETASOSTYLIS, Griseb. = **Leianthus**, Griseb. (Gentian.).

PETALOTOMA, DC. Prod. iii. 294 (1828) = **Carrallia**, Roxb. (Rhizoph.).
brachiata, DC. l. c. 295 = C. integerrima.

PETALOXIS, Rafin. Fl. Tellur. ii. 83 (1836) = **Dichorisandra**, Mikan (Commel.).
purpurea, Rafin. l. c. = D. oxypetala.

PETAMENES, Salisb. in Trans. Hort. Soc. i. (1812) 324 = **Antholyza**, Linn. (Irid.).
quadrangularis, Salisb. l. c. = A. quadrangularis.

PETASIOIDES, Vitm. Summa Veg. i. 349 (1789) PETESIOIDES, Jacq. = **Wallenia**, Sw. (Myrsin.).

PETASITES, [Tourn.] Linn. Syst. ed. I (1735); Gaertn. Fruct. ii. 406. t. 166. (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 439.
albus, *Gaertn. l. c.*—Europ.; As. bor.
albus, Wirzin, Casan, n. 265; ex Ledeb. Fl. Ross. ii. 470 = spurius.
alpestris ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 104.—Europ.
alpinus, Baumg. Enum. Stirp. Transs. iii. 92 = Homogyne alpina.
androgynus, Schur, Enum. Pl. Transs. 302 = officinalis.
consimilis, Jord. ex Nym. Consp. 397 = officinalis.
discolor, Baumg. Enum. Stirp. Transs. iii. 92 = Homogyne discolor.
elatior, Hill, Herb. Brit. i. 29 = officinalis.
Farfara, Baill. Hist. Pl. viii. 58, 309 = Tussilago Farfara.
fragrans, *Presl, Fl. Sicul.* i. 28.—Reg. Mediterr.
frigidus, *Fries, Syll.* 20; *et Summa Veg. Scand.* 182.—Reg. bor. et austr.
giganteus, Fuss, Fl. Transs. 310 = officinalis.
giganteus, *F. Schmidt, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 443.—Ins. Sachalin.
glabrescens, Hoppe, ex DC. Prod. v. 207 = albus.
hermaphroditus, Schur, Enum. Pl. Transs. 302 = officinalis.
hybridus, Gaertn. Mey. & Scherb. Fl. Wett. iii. 184 = officinalis.
intermedius, Hoppe, ex Nym. Consp. 397 = officinalis.
japonicus, *F. Schmidt, Reise Amurl.* 145.—Ins. Sachalin.
Kablikianus, Tausch, ex Lotos (1851), 120; et ex Reichb. Ic. Fl. Germ. xv. 4. t. 900 = officinalis.
laevigatus, *Reichb. Fl. Germ. Excurs.* 279.—Bohem.; Sibir.
Lorenzianus ×, Bruegg, in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 104.—Europ.
macrophyllus, Schur, Enum. Pl. Transs. 302 = officinalis.
niveus, *Baumg. Enum. Stirp. Transs.* iii. 94; *Cass. in Dict. Sc. Nat.* xxxix. 202.—Europ.
niveus, Trotzky, Ind. Sem. Hort. Casan. (1842); ex Ledeb. Fl. Ross. ii. 470 = spurius.
ochroleucus, Boiss. & Huet, Diagn. Ser. II. iii. 5 = officinalis.
officinalis, *Moench, Meth.* 568.—Europ.; As. bor.
ovatus, Hill, Herb. Brit. i. 28 = officinalis.
palmatus, *A. Gray, in Bot. Calif.* i. 407.—As. occ.; Am. bor.
paradoxus, Baumg. Enum. Stirp. Transs. iii. 94 = niveus.
pratensis, Jord. Pugill. Pl. 103 = officinalis.

PETASITES:—
ramosus, Baumg. Enum. Stirp. Transs. iii. 94 = albus.
Reuterianus, Jord. ex Nym. Consp. 397 = officinalis.
riparius, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 321 = officinalis.
sagittatus, *A. Gray, in Bot. Calif.* i. 407.—Am. bor.
sebethius, Tenore, Cat. Ort. Bot. Nap. (1845) 90 = officinalis.
spurius, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 168 = japonicus.
spurius, *Reichb. f. Ic. Fl. Germ.* xvi. 3. t. 896.—Europ.
stramineus, *Nym. Syll.* 20.—Europ.
tomentosus, DC. Prod. v. 207 = spurius.
tricholobus, *Franch. Pl. Davidian.* i. (1884) 172.—Mongol.
vulgaris, Desf. Fl. Atlant. ii. 270 = officinalis.

PETASOSTYLIS, Griseb. in DC. Prod. ix. 71 (1845) = **Leianthus**, Griseb. (Gentian.).
nigrescens, Griseb. l. c. = L. nigrescens.
saponarioides, Griseb. l. c. = L. saponarioides.

PETASTOMA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 194 = **Bignonia**, Tourn.
Clausseni, Miers, l. c. 195 = Arrabidaea Clausseni.
corymbiferum, Miers, l. c. 193 = B. corymbifera.
crenatum, Miers, l. c. 193.—Bras.
divaricatum, Miers, l. c. 195, nomen.—Bras.
Donianum, Miers, l. c. 195 = B. Doniana.
formosum, Bur. in Rev. Hortic. (1868) 154.—Bras.
glabrum, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 195 = B. samydoides.
jucundum, Miers, l. c., nomen.—Guian.
lachnaeum, Miers, l. c., nomen.—Bras.
laevigatum, Miers, l. c., nomen.—Bras.
leptopodum, Miers, l. c., nomen.—Bras.
mansuetum, Miers, l. c., nomen.—Bras.
obliquum, Miers, l. c. = B. obliqua.
ophthalmicum, Miers, l. c. = B. ophthalmica.
patelliferum, Miers, l. c. = B. patellifera.
phaseoloides, Miers, l. c. = B. phaseoloides.
physaloides, Miers, l. c. = B. physaloides.
platyphyllum, Miers, l. c. = Arrabidaea platyphylla.
pubescens, Miers, l. c. = B. pubescens.
puniceum, Miers, l. c. = B. punicea.
samydoides, Miers, l. c. = B. samydoides.
Sceptrum, Miers, l. c. = B. Sceptrum.
simplicifolium, Miers, l. c. 194.—Bras.
triplinerve, Miers, l. c. 195 = B. triplinervia.

PETASULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Trevesia**, Vis. (Araliac.).

PETERIA, A. Gray, Pl. Wright. i. 50; (in Smithson. Contrib. iii.) (1852). *LEGUMINOSAE*, Benth. & Hook. f. i. 495.
scoparia, *A. Gray, Pl. Wright.* i. 50.—N. Mexic.
Thomsonae, *S. Wats. in Am. Natural.* vii. (1873) 300.—Am. bor.

PETERMANNIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 124; et in Abh. Akad. Berl. (1855) 74. t. 6 C = **Begonia**, Linn.
Cumingiana, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 124 = B. Cumingiana.
fasciculata, Klotzsch, l. c. = B. fasciculata.
geniculata, Klotzsch, l. c. = B. fasciculata.
racemosa, Klotzsch, l. c. = B. racemosa.

PETERMANNIA, F. Muell. Fragm. ii. 92 (1860). *DIOSCOREACEAE*, Benth. & Hook. f. iii. 746.
cirrosa, *F. Muell. l. c.* 93.—Austral.

PETERMANNIA, Reichb. Nom. 236 (1841) = **Cycloloma**, Moq. (Chenop.).

PETERSIA, Klotzsch, in Peters, Reise Mossamb. Bot. 168. t. 30 (1861) = **Capparis**, Linn.
rosea, Klotzsch, l. c. = C. rosea.

PETERSIA, Welw. in Benth. & Hook. f. Gen. i. 721 (1867). *MYRTACEAE*, Benth. & Hook. f. l. c.
africana, *Welw. ex Benth. & Hook. f. l. c.*—Afr. trop.

PETESIA, P. Br. Hist. Jamaic. 143. tt. 2, 3 (1756) = **Rondeletia**, Linn. (Rubiac.).
carnea, Forst. f. Prod. 10 = Ixora carnea.
carnosa, Hook. & Arn. Bot. Beech. Voy. 64 = Bikkia grandiflora.
coriacea, Hook. & Arn. l. c. 85 = Gouldia sandwicensis.
grandis, Bartl. ex DC. Prod. iv. 395 = Sommera arborescens.
hispida, Bartl. l. c. (an Wendlandiae sp.?).—Ins. Philipp.
Hita, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 134 = Hedyotis scandens.
Lygistum, Linn. Syst. ed. X. 894 = Manettia Lygistum.
minor, Griseb. in Bonplandia, vi. (1858) 8.—Mexic.
nitida, Bartl. ex DC. Prod. iv. 395.—Ins. Philipp. et Marian.
simplicissima, Lour. Fl. Cochinch. 77.—Cochinch.
spicata, Sw. Fl. Ind. Occ. iii. 1945 = Gonzalea Petesia.
stipularis, Linn. Syst. ed. X. 894 = R. tomentosa.
terminalis, Hook. & Arn. Bot. Beech. Voy. 85 = Gouldia sandwicensis.
ternifolia, Bartl. ex DC. Prod. iv. 395.—Ins. Philipp.
tomentosa, Jacq. Select. Am. 18.—N. Granat.
trifida, Lour. Fl. Cochinch. 77.—Cochinch.
villosa, Sm. in Rees, Cycl. xxvii. n. 4 = R. umbellulata.

PETESIOIDES, Jacq. Select. Am. 17 (1763) = **Wallenia**, Sw. (1788) (Myrsin.).
laurifolium, Jacq. l. c. = W. laurifolia.

PETILIUM, Linn. Syst. ed. I (1735); Hort. Cliff. 119 (1737) = **Fritillaria**, Linn. (Liliac.).
imperiale, Jaume St. Hil. Expos. i. 120 = F. imperialis.

PETITIA, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1829) 219 = **Xatardia**, Meissn. (Umbellif.).
scabra, J. Gay, l. c. = X. scabra.

PETITIA, Jacq. Enum. Pl. Carib. 1 (1760). *VERBENACEAE*, Benth. & Hook. f. ii. 1151.
SCLEROON, Benth. in Lindl. Bot. Reg. (1843) Misc. 65.
domingensis, *Jacq. Enum. Pl. Carib.* 12.—Ins. S. Doming.
domingensis, Walp. Rep. iv. 71 = Poeppigii.
oleina, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 539.—Mexic.
Poeppigii, *Schau. in DC. Prod.* xi. 639.—Cuba.
quinduensis, H. B. & K. Nov. Gen. et Sp. ii. 248 = Aegiphila Humboldtii.
tenuifolia, Willd. ex Schult. Mant. iii. 50 = Aegiphila Humboldtii.

PETITIA, Neck. Elem. ii. 407 (1790) = **Hibiscus**, Linn. (Malvac.).

PETIVERIA, Plum. ex Linn. Gen. ed. I. 341 (1737). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 82.
MAPA, Vell. Fl. Flum. 59 (1825).
alliacea, *Linn. Sp. Pl.* 342.—Ind. occ.
foetida, Salisb. Prod. 214 = alliacea.
hexaglochin, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 35.—Bras.
ochroleuca, Moq. in DC. Prod. xiii. II. 9 = alliacea.
octandra, Linn. Sp. Pl. ed. II. 486 = alliacea.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 160; *Contrib.* 53.—Paraguay.
tetrandra, *Gomez, Obs. Med. Bot. Pl. Bras.* (1803) 13; *et in Mem. Ac. Lisboa*, iii. (1812) *Mem. Corresp.* 17.—Bras.

PETRACANTHUS, Nees, in DC. Prod. xi. 97 (1847) = **Gymnostachyum**, Nees (Acanth.).
Junghuhnii, Nees, l. c. 98 = G. Cumingianum.

PETRAEA, B. Juss. ex Juss. Gen. 108 (1789) = **Petrea**, Houst. (Verben.).

PETRAEOVITEX, Oliver, in Hook. Ic. Pl. xv. 15. t. 1420 (1883). *VERBENACEAE*.
Riedelii, *Oliver, l. c.* 16. t. 1420.—Malaya.

PETRAMNIA, Rafin. Fl. Tellur. iv. 68 (1836) = **Craniolaria**, Linn. (Pedalin.).

PETRANTHA, DC, Prod. v. 503 (1836), err. typ. = TETRANTHA, Poit. = **Riencourtia**, Cass. (Compos.).

PETRANTHE, Salisb. Gen. Pl. Fragm. 27 (1866) = **Scilla**, Linn. (Liliac.).
verna, Salisb. l. c. = S. verna.

PETREA, Houst. ex Linn. Gen. ed. I. 347 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1149.
arborea, *H. B. & K. Nov. Gen. et Sp.* ii. 282.—N. Granat.
aspera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 211.—Bras.
Blanchetiana, *Schau. in DC. Prod.* xi. 617.—Bras.
bracteata, *Steud. in Flora*, xxvi. (1843) 764.—Guian.
Candolleana, *Schau. in DC. Prod.* xi. 617.—Hab.?
dentata, Spreng. Syst. ii. 761 = Patagonula americana.
denticulata, Schrad. in Goett. Gel. Anz. i. (1821) 712 = Patagonula americana.
erecta, Lodd. Bot. Cab. t. 1606 = arborea.
guianensis, *Cham. in Linnaea*, vii. (1832) 367.—Guian.
insignis, *Schau. in DC. Prod.* xi. 620.—Bras.
Kohautiana, Presl, Bot. Bemerk. 99 = subserrata, *Cham.*
macrostachya, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 448 = guianensis.
Martiana, *Schau. in DC. Prod.* xi. 620.—Bras.
mexicana, Willd. ex Cham. in Linnaea, vii. (1832) 367 = arborea.
multiflora, Sm. in Rees, Cycl. xxvii. n. 2 (Quid?).—'Ins. Honim.'
oblonga, Spreng. Syst. ii. 761 (Banisteriae sp.).—Bras.
obtusifolia, *Benth. Pl. Hartw.* 246.—N. Granat.
ovata, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 328 = arborea.
pubescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 211.—Bras.
racemosa, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 72 = oblonga.
retusa, Presl, Bot. Bemerk. 99 = subserrata.
rugosa, *H. B. & K. Nov. Gen. et Sp.* ii. 282.—Venezuela.
Schomburgkiana, *Schau. in DC. Prod.* xi. 619.—Guian.
serrata, Presl, Bot. Bemerk. 99 = subserrata.
Stapeliae, *Paxt. Mag. Bot.* iv. (1838) 199 (Stapelsiae).—Am. austr.
subserrata, *Bárcena, Notic. Cient. Estad. Hidalg.* 31.—Mexic.
subserrata, *Cham. in Linnaea*, vii. (1832) 368.—Bras.
vincentina, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 212.—Ins. S. Vinc.
volubilis, Gaertn. Fruct. ii. 471 = Patagonula americana.
volubilis, *Linn. Sp. Pl.* 626; *Jacq. Select. Am.* 180. t. 114.—Am. trop.
volubilis, Vell. Fl. Flum. 254; vi. t. 59 = subserrata, *Cham.*
volubilis, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 381 = arborea.

PETROBIUM, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1838) t. 4 = **Lithobium**. Bong. (Melastom.).
cordatum, Bong. l. c. = L. cordatum.

PETROBIUM, R. Br. in Trans. Linn. Soc. xii. (1816) 113. *COMPOSITAE*, Benth. & Hook. f. ii. 355.
DRIMYPHYLLUM, Burch. ex DC. Prod. v. 501 (1836).
LAXMANNIA, Forst. Char. Gen. 94. t. 47 (1776).
arboreum, *R. Br. in Trans. Linn. Soc.* xii. (1816) 113.—Ins. S. Helen.
Forsteri, Cass. in Dict. Sc. Nat. xxxix. 307 = arboreum.

PETROCALLIS, R. Br. in Ait. Hort. Kew. ed. II. iv. 93 (1812) = **Draba**, Linn. (Crucif.).
araratica, Griseb. ex Tchihat. Asie Min. Bot. i. 317, nomen = D. araratica?
fenestrata, Boiss. & Hohen. Diagn. Ser. I. viii. 27.—Persia.
pyrenaica, R. Br. in Ait. Hort. Kew. ed. II. iv. 93 = D. pyrenaica.

PETROCARVI, Tausch, in Flora, xvii. (1834) 355 = **Athamantha**, Linn. (Umbell.).
cretense, Tausch, l. c. = A. cretensis.

PETROCARYA, Schreb. Gen. 245 (1789) = **Parinarium**, Aubl. (Rosac.).
brasiliensis, Schott, in Spreng. Syst. iv. App. 405 = Parin. montanum.
campestris, Willd. Sp. Pl. ii. 287 = Parin. campestre.

PETROCARYA :—
dillenifolia, Steud. Nom. ed. II. ii. 309 = Dipterocarpus cornutus.
excelsa, Jack, in Malay. Misc. ii. (1822) n. VII. p. 66 = Parinarium Jackianum.
excelsa, Steud. Nom. ed. II. ii. 309 = Parin. excelsum.
glaberrima, Miers, in Journ. Linn. Soc. xvii. (1879) 336 = Parin. glaberrimum.
Griffithiana, Miers, l. c. = Parin. Griffithianum.
macrophylla, Steud. Nom. ed. II. ii. 309 = Parin. macrophyllum, *Sabine*.
Maranthes, Miers, in Journ. Linn. Soc. xvii. (1879) 336 = Parin. Maranthes.
montana, Willd. Sp. Pl. ii. 287 = Parin. montanum.
scabra, Miers, in Journ. Linn. Soc. xvii. (1879) 336 = Parin. scabrum.
senegalensis, Steud. Nom. ed. II. ii. 309 = Parin. senegalense.
sumatrana, Jack, in Malay. Misc. ii. (1822) n. VII. p. 67.—Sumatra.
sumatrana, Miers, in Journ. Linn. Soc. xvii. (1879) 336 = Parin. costatum.

PETROCODON, Hance, in Journ. Bot. xxi. (1883) 167. *GESNERACEAE*.
dealbatus, *Hance, l. c.*—China.

PETROCOMA, Rupr. Fl. Cauc. 200 (1869) = **Silene**, Linn. (Caryoph.).
Hoefftiana, Rupr. l. c. = S. Hoefftiana.

PETROCOPTIS, A. Br. in Flora, xxvi. (1843) 370 = **Lychnis**, Tourn. (Caryoph.).
glaucifolia, Boiss. Diagn. Ser. II. i. 80.—Hispan.
Lagascae, Willk. in Flora, xxxiv. (1851) 601 = L. Lagascae.
pyrenaica, A. Br. l. c. xxvi. (1843) 370 = L. pyrenaica.

PETRODORA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
fruticulosa, Fourr. l. c. = V. fruticulosa.
saxatilis, Fourr. l. c. = V. saxatilis.

PETROGETON, Eckl. & Zeyh. Enum. 291 (1837) = **Crassula**, Dill.
alpinum, Eckl. & Zeyh. l. c. = C. alpina.
dentatum, Eckl. & Zeyh. l. c. = C. dentata.
nemorosum, Eckl. & Zeyh. l. c. 292 = C. nemorosa.
nivale, Eckl. & Zeyh. l. c. = C. nivalis.
patens, Eckl. & Zeyh. l. c. 291 = C. patens.
typicum, Eckl. & Zeyh. l. c. = C. dentata.
Umbella, Eckl. & Zeyh. l. c. = C. Umbella.

PETROMARULA, Vent. ex Hedw. f. Gen. 139 (1806) = **Phyteuma**, Linn. (Campan.).
pinnata, A. DC. Monog. Camp. 139 = Phyt. pinnatum.

PETRONIA, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 52, sphalm. = **Pteronia**, Linn. (Compos.).

PETRONIA, Rodrig. Orch. Nov. i. 106 (1877). *ORCHIDEAE*.
regia, *Rodrig. l. c.* 107.—Bras.

PETROPHILA, R. Br. in Trans. Linn. Soc. x. (1810) 67. *PROTEACEAE*, Benth. & Hook. f. iii. 172.
ATYLUS, Salisb. Parad. Lond. sub t. 67 (1807), partim.
acicularis, *R. Br. in Trans. Linn. Soc.* x. (1810) 69.—Austral.
adiantifolia, Sm. ex Meissn. in DC. Prod. xiv. 274 = diversifolia.
anceps, R. Br. Prot. Nov. 5 = linearis.
axillaris, Meissn. in Hook. Kew Journ. vii. (1855) 68 = P. Serruriae.
biloba, *R. Br. Prot. Nov.* 7.—Austral.
biternata, *Meissn. in Hook. Kew Journ.* vii. (1855) 69.—Austral.
brevifolia, Lindl. Swan River App. 35 = media.
canescens, A. Cunn. ex R. Br. Prot. Nov. 6 = sessilis.
carduacea, *Meissn. in DC. Prod.* xiv. 274.—Austral.
chrysantha, *Meissn. in Hook. Kew Journ.* vii. (1855) 68.—Austral.
circinnata, *Kippist, ex Meissn. l. c.* 67.—Austral.
colorata, *Meissn. in Lehm. Pl. Preiss.* ii. 246.—Austral.

PETROPHILA :—
conifera, *Meissn. in Hook. Kew Journ.* vii. (1855) 67.—Austral.
crassifolia, R. Br. Prot. Nov. 5 = teretifolia.
crispata, *R. Br. l. c.* 6.—Austral.
Cunninghami, Meissn. in Lehm. Pl. Preiss. i. 499 = squamata.
divaricata, *R. Br. Prot. Nov.* 7.—Austral..
diversifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 70.—Austral.
Drummondii, *Meissn. in Lehm. Pl. Preiss.* i. 496.—Austral.
dubia, R. Br. Prot. Nov. 7 = Isopogon roseus.
ericifolia, *R. Br. l. c.* 5.—Austral.
falcata, A. Cunn. ex Meissn. in DC. Prod. xiv. 267 = linearis.
fastigiata, *R. Br. in Trans. Linn. Soc.* x. (1810) 76.—Austral.
filifolia, R. Br. l. c. 69 = acicularis.
fucifolia, Knight, Prot. 92 = pulchella.
glanduligera, Lindl. Swan River App. 35 = P. Serruriae.
gracilis, A. Cunn. ex Endl. Gen. Suppl. iv. II. 76 = squamata.
heterophylla, *Lindl. Swan River App.* 35.—Austral.
inconspicua, *Meissn. in Hook. Kew Journ.* vii. (1855) 68.—Austral.
intricata, Lindl. Swan River App. 35 = divaricata.
juncifolia, Lindl. l. c. = media.
linearis, *R. Br. Prot. Nov.* 6.—Austral.
longifolia, *R. Br. l. c.* 5.—Austral.
longifolia, A. Cunn. ex Endl. Gen. Suppl. iv. II. 75 = teretifolia.
macrostachya, *R. Br. Prot. Nov.* 7.—Austral.
media, *R. Br. l. c.* 5.—Austral.
megalostegia, *F. Muell. Fragm.* x. 61.—Austral.
multisecta, *F. Muell. l. c.* vi. 242.—Austral.
pedunculata, *R. Br. in Trans. Linn. Soc.* x. (1810) 70.—Austral.
phylicoides, R. Br. Prot. Nov. 6 = ericifolia.
plumosa, *Meissn. in Hook. Kew Journ.* vii. (1855) 69.—Austral.
propinqua, *R. Br. Prot. Nov.* 7.—Austral.
pulchella, *R. Br. in Trans. Linn. Soc.* x. (1810) 69.—Austral.
rhoifolia, Sm. ex Meissn. in DC. Prod. xiv. 274 = diversifolia.
rigida, *R. Br. in Trans. Linn. Soc.* x. (1810) 69.—Austral.
Roei, *Endl. Gen. Suppl.* iv. II. 75, nomen.—Austral.
scabriuscula, Meissn. in Lehm. Pl. Preiss. i. 495 = ericifolia.
semifurcata, *F. Muell. ex Benth. Fl. Austral.* v. 335.—Austral.
seminuda, *Lindl. Swan River App.* 34.—Austral.
Serruriae, *R. Br. Prot. Nov.* 6.—Austral.
sessilis, *Sieber, ex Schult. Mant.* iii. 262.—Austral.
Shuttleworthiana, *Meissn. in Lehm. Pl. Preiss.* ii. 246.—Austral.
squamata, *R. Br. in Trans. Linn. Soc.* x. (1810) 70.—Austral.
striata, *R. Br. Prot. Nov.* 6.—Austral.
teretifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 68.—Austral.
trifida, *R. Br. in Trans. Linn. Soc.* x. (1810) 70.—Austral.
trifida, Lodd. Bot. Cab. t. 1883 = Isopogon trilobus.
triternata, Kippist, ex Meissn. in Hook. Kew Journ. vii. (1855) 67 = Drummondii.

PETROPHILE, Knight, Prot. 92 (1809) = **Petrophila**, R. Br. (Proteac.).

PETROPHYES, Webb & Berth. Phyt. Canar. i. 201. t. 36 (1836–40) = **Monanthes**, Haw. Crassul.).
agriostaphis, Webb & Berth. l. c. 203.—Ins. Canar.
brachycaulon, Webb & Berth. l. c. = M. brachycaulon.
icterica, Webb, ex Bolle, in Bonplandia, vii. (1859) 244.—Ins. Canar.
microbotrys, Bolle & Webb, l. c. 245.—Ins. Canar.
minima, Bolle, l. c.—Ins. Canar.
moralis, Bourg. ex Bolle, l. c. = *tilophila*.
muralis, Webb, ex Bolle, l. c. 244.—Ins. Canar.
polyphyllum, Webb & Berth. Phyt. Canar. i. 201 = M. polyphylla.
purpurascens, Bolle & Webb, in Bonplandia, vii. (1859) 245.—Ins. Canar.
tilophila, Bolle, l. c.—Ins. Canar.

PETRORHAGIA, Link, Handb. ii. 235 (1831) = **Tunica**, Hall. (Caryophyll.).
glomerata, Link, l. c. = Gypsophila glomerata.
multicaulis, Link, l. c. = T. Saxifraga.
rigida, Link, l. c. = T. Saxifraga.
Saxifraga, Link, l. c. = T. Saxifraga.

PETROSAVIA, Becc. in Nuov. Giorn. Bot. Ital. iii. (1871) 7. t. 1. *LILIACEAE*, Benth. & Hook. f. iii. 828.
stellaris, *Becc. l. c.* 8.—Borneo.

PETROSCIADIUM, Edgew. in Proc. Linn. Soc. i. (1845) 252; et in Trans. Linn. Soc. xx. (1846) 51 = **Pimpinella**, Linn. (Umbell.).
caespitosum, Edgew. ll. cc. = Pimpinella caespitosa.

PETROSELINUM, Hoffm. Gen. Umb. i. 78 (1814); DC. Prod. iv. 102 = **Carum**, Linn. (Umbell.).
ammoides, Reichb. f. Ic. Fl. Germ. xxi. 11 = C. ammoides.
australe, Hook. f. Fl. N. Zel. ii. 366 (Index) = Apium prostratum.
crispum, Nym. Consp. 309 = C. Petroselinum.
dissectum, Benth. Pl. Hartw. 188.—N. Granat.
filiforme, A. Rich. Ess. Fl. Nouv. Zél. 278 = Apium prostratum.
fractophyllum, Lag. ex Sweet, Hort. Brit. ed. II. 245, nomen.—Hab. ?
hortense, Hoffm. Gen. Umb. 163 = C. Petroselinum.
humile, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 531 = Apium humile.
intermedium, Reichb. f. Ic. Fl. Germ. xxi. 11 = C. inundatum.
peregrinum, Lag. Amen. Nat. ii. 100, 103 = C. peregrinum.
prostratum, DC. Prod. iv. 102 = Apium prostratum.
protractum, Walp. Rep. ii. 391, sphalm. = praec.
romanum, Sweet, Hort. Brit. ed. II. 245 = C. peregrinum; C. Petroselinum.
sativum, Hoffm. Gen. Umb. 177 = C. Petroselinum.
sativum, Hook. & Gill. in Hook. Bot. Misc. i. (1830) 335; iii. (1833) 354 = Seseli Gilliesii.
segetum, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 128 = C. segetum.
selinoides, DC. Prod. iv. 102 = C. peregrinum.
Seubertianum, H. C. Wats. in Hook. Lond. Journ. Bot. vi. (1847) 387.—Ins. Azor.
ternatum, Mutel, Fl. Fr. ii. 402 = Ligusticum scoticum.
Thermoeri, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. VII. 60.—Rossia.
Thorei, Coss. ex Reichb. Ic. Fl. Germ. xxi. 11 = C. inundatum.
trifoliatum, H. C. Wats. in Hook. Lond. Journ. Bot. iii. (1844) 502.—Ins. Azor.

PETROSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = **Silene**, Linn. (Caryoph.).
saxifragum, Fourr. l. c. = S. Saxifraga.

PETROSIMONIA, Bunge, Anab. Rev. 52 (1862). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 73.
brachiata, *Bunge, l. c.* 59.—Reg. Casp.
crassifolia, *Bunge, l. c.* 55.—Reg. Casp.
glauca, Bunge, l. c. 58 = crassifolia.
monandra, *Bunge, l. c.* 53.—Reg. Casp.
sibirica, *Bunge, l. c.* 60.—Rossia; Soongaria.
squarrosa, *Bunge, l. c.* 57.—Soongaria.
Volvox, *Bunge, l. c.* 52.—Reg. Casp.

PETROSTYLIS, Pritz. Ic. Ind. i. 828 (1855), sphalm. = **Pterostylis**, R. Br. (Orchid.).

PETROTHECA, Steud. Nom. ed. II. ii. 310 (1841), sphalm. = **Tetratheca**, Sm. (Tremand.).
ericaefolia, Steud. l. c. = T. ericifolia.
glandulosa, Steud. l. c. = T. pilosa.

PETRUSIA, Baill. in Bull. Soc. Linn. Par. i. (1881) 274. *OLACINEAE*?
madagascariensis, *Baill. l. c.*—Madag.

PETTERA, Reichb. Ic. Fl. Germ. v. 33. t. 220 (1841); Nom. 205 (1841) = **Arenaria**, Rupp. (Caryoph.).
graminifolia, Reichb. l. c. 33 = A. graminifolia.

PETTERIA, Presl, Bot. Bemerk. 139 (1844); in Abh. Boehm. Ges. V. iii. (1845) 569. *LEGUMINOSAE*, Benth. & Hook. f. i. 482.
ramentacea, *Presl, l. c.*—Dalmat.

PETTOSPERMUM, Roxb. Hort. Beng. 18 (1814), nomen, sphalm. = **Pittosporum**, Banks.

PETUNGA, DC. Prod. iv. 398 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 93.
HIGGINSIA, Blume, Bijdr. 988 (1826).
PHYLANTHERA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.
SPICILLARIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 252.
coniocarpa, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 173.—Borneo.
dasyoneura, *Korth. l. c.*—Sumatra.
glomerata, *DC. Prod.* iv. 399.—Java.
gracilis, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 173.—Borneo.
lanceolata, *Korth. l. c.* 171.—Borneo.
longifolia, *DC. Prod.* iv. 399.—Penang; Java.
microcarpa, DC. l. c. = Roxburghii.
Roxburghii, *DC. l. c.*—Ind. or.; Malaya.
rufidula, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 131.—Sumatra.
salicina, *Miq. l. c.* 132.—Borneo.
variabilis, Hassk. in Flora, xxv. (1842) II. Beibl. 24; xxviii. (1845) 232 = Roxburghii.
venulosa, *Hook. f. Fl. Brit. Ind.* iii. 121.—Malaya.

PETUNIA, Juss. in Ann. Mus. Par. ii. (1803) 215. t. 47. *SOLANACEAE*, Benth. & Hook. f. ii. 907.
CALIBRACHOA, La Llave & Lex. Nov. Veg. Desc. fasc. ii. 3 (1825).
LEPTOPHRAGMA, Benth. ex Dun. in DC. Prod. xiii. I. 578 (1852).
STIMOMPHIS, Rafin. Fl. Tellur. iii. 76 (1836).
STIMORYNE, Rafin. l. c. (1836).
WADDINGTONIA, Phil. Fl. Atac. 41. t. 5 (1857).
acuminata, R. Grah. in Edinb. N. Phil. Journ. (Jul.–Sept. 1828) 378 = Nicotiana acuminata.
Atkinsiana, D. Don, ex Loud. Hort. Brit. Suppl. ii. 655 = violacea.
caesia, *Sendtn. in Mart. Fl. Bras.* x. 173.—Bras.
calycina, *Sendtn. l. c.*—Bras.
cirrhoides, Miers, Trav. ii. 531 = Nicotiana cirrhoides.
Cumingiana, *Remy, in C. Gay, Fl. Chil.* v. 50.—Chili.
dichotoma, *Sendtn. in Mart. Fl. Bras.* x. 173.—Bras.
elegans, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 186.—Bras.
helianthemoides, *Sendtn. in Mart. Fl. Bras.* x. 174.—Bras.
heterophylla, Sendtn. l. c. 176 = Fabiana heterophylla.
humifusa, *Dun. in DC. Prod.* xiii. I. 574.—Chili.
hybrida ×, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 615.
intermedia, G. Don, Gen. Syst. iv. 468 = Salpiglossis linearis.
ledifolia, *Sendtn. in Mart. Fl. Bras.* x. 176.—Bras.
linearis, Herb. ex Paxt. Mag. Bot. ii. (1836) 219 = Salpiglossis linearis.
linoides, *Sendtn. in Mart. Fl. Bras.* x. 174.—Bras.
Meleagris ×?, Planch. in Fl. des Serres, vi. (1850–51) 39. t. 550.
mendozinensis, Gill. ex Miers, in Hook. Lond. Journ. Bot. v. (1846) 174 = Leptoglossis linifolia.
nyctaginiflora, *Juss. in Ann. Mus. Par.* ii. (1803) 216. t. 47. f. 2.—Reg. Argent.
ovalifolia, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 189.—Bras.
parviflora, *Juss. in Ann. Mus. Par.* ii. (1803) 216. t. 47.—Reg. Argent.
phoenicea, D. Don, ex Loud. Hort. Brit. Suppl. ii. 655 = violacea.
propinqua, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 185.—Reg. Argent.
punctata ×, Paxt. Mag. Bot. xi. (1842) 7.
Sellowiana, Sendtn. in Mart. Fl. Bras. x. 176 = Fabiana Sellowiana.
serpyllifolia, *Sendtn. l. c.* 175.—Bras.
thymifolia, *Sendtn. l. c.*—Bras.
villadiana, *Bárcena, ex Hemsl. Biol. Centr. Am. Bot.* ii. 436.—Mexic.
violacea, *Lindl. Bot. Reg.* t. 1626.—Reg. Argent.
viscidula, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 189.—Mexic. (cult.).
viscosa, Colla, in Mem. Acc. Torin. xxxviii. (1835) 135. t. 45 = Bouchetia erecta.
viscosa, Miers, Trav. ii. 531 = Nicotiana acuminata.

PEUCE, Rich. in Ann. Mus. Par. xvi. (1810) 298 (Conif.)
Abies, Rich. l. c. = Picea excelsa.

PEUCE:—
balsamea, Rich. in Ann. Mus. Par. xvi. (1810) 298 = Abies balsamea.
Cedrus, Rich. l. c. = Cedrus Libani.
Larix, Rich. l. c. = Larix europaea.
rubra, Rich. l. c. = Picea nigra.

PEUCEDANON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Peucedanum**, Tourn. (Umbell.).
cervarium, St. Lag. l. c. = Peucedanum Cervaria.

PEUCEDANUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 74 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 918.
ALVARDIA, Fenzl, in Flora, xxvii. (1844) 312.
ANALYRIUM, E. Mey. in Drège, Zwei Pfl. Docum. 163 (1843).
ANETHUM, Tourn. ex Linn. Gen. ed. I. 82 (1737).
ARCHEMORA, DC. Prod. iv. 188 (1830).
BUBON, Linn. Sp. Pl. ed. I. 233 (1753); Koch, in Nov. Act. Nat. Cur. xii. I. (1824) 95.
CAROSELINUM, Griseb. Spicil. Fl. Rumel. i. 374 (1843).
CERVARIA, Gaertn. Fruct. i. 90. t. 21 (1788).
COGSWELLIA, Schult. Syst. vi. p. xlviii (1820).
CYNORHIZA, Eckl. & Zeyh. Enum. 351 (1837).
DIPLOTAENIA, Boiss. in Ann. Sc. Nat. Nat. Sér. III. i. (1844) 308.
DREGEA, Eckl. & Zeyh. Enum. 350 (1837).
ELAEOCHYTRIS, Fenzl, in Russegg. Reise, i. 957. (1843); (Ill. Pl. Syr. 71. t. 19).
EULOPHUS, Nutt. in Journ. Acad. Philad. vii. (1834) 27.
EURYPTERA, Torr. & Gray, Fl. N. Am. i. 629 (1840).
GALIMBIA, Endl. Ench. 379 (1841), sphalm.
IFDREGEA, Steud. Nom. ed. II. i. 801 (1840).
IMPERATORIA, Linn. Syst. ed. I (1735).
LOMATIUM, Rafin. in Journ. Phys. lxxxix. (1819) 101.
MACROSELINUM, Schur. in Verh. Siebenb. Ver. Naturw. iv. (1853) 30.
NEUROPHYLLUM, Torr. & Gray, Fl. N. Am. i. 612 (1840).
OPOIDIA, Lindl. Bot. Reg. (1839) Misc. 66.
OREOSELINUM, [Tourn.] Adans Fam. ii. 100 (1763); Bieb. Fl. Taur. Cauc. iii. 209 (1819).
ORMOSOLENIA, Tausch, in Flora, xvii. (1834) 348.
OSTRUTHIUM, Link. Handb. i. 360 (1829).
OXYPOLIS, Rafin. Neogenyt. 2 (1825).
PALIMBIA, Bess. Enum. Pl. Volh. 55 (1822).
PASTINACA, Linn. Gen. ed. I. 81 (1737).
PLEUROTAENIA, Hohen. ex Benth. & Hook. f. Gen. i. 920 (1865).
POLYCYRTUS, Schlecht. in Linnaea, xvii. (1843) 126.
PORPHYROSCIAS, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 62 (1867).
PTEROSELINUM, Reichb. Fl. Germ. Excurs. 453 (1832).
SATARIA, Rafin. New Fl. Am. iv. 20 (1836).
SCHLOSSERIA, Vukot. in Oestr. Bot. Wochenbl. vii. (1857) 350.
SCIOTHAMNUS, Endl. Gen. 780 (1839).
STEGANOTAENIA, Hochst. in Flora, xxvii. (1844) Beibl. 4.
TAENIOPETALUM, Vis. Fl. Dalm. iii. 49 (1849).
THYSSELINUM, Moench, Meth. 84 (1794); Hoffm. Umb. Gen. ed. I. 153 (1814).
TIEDMANNIA, DC. Prod. iv. 187 (1830).
TOMMASINIA, Bertol. Fl. Ital. iii. 414 (1837).
XANTHOSELINUM, Schur, Enum. Pl. Transs. 264 (1866).
abbreviatum, *E. Mey. ex Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 535.—Afr. austr.
abrotanifolium, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 184 = ambiguum.
abyssinicum, *Vatke, in Linnaea*, xl. (1876) 187.—Abyss.
album, Hort. ex Hoffm. Gen. Umb. ed. I. 155 = Silaus alpestris.
album, Hort. Par. ex Poir. Encyc. v. 228 = alsaticum.
album, Spreng. Gesch. Bot. ii. 200 = salinum.
alliacea, Baill. Hist. Pl. vii 185 = Ferula alliacea.
alpestre, *Linn. Sp. Pl.* 246.—Bessarab.
alpestre, Spreng. Sp. Umbell. 56 = Silaus alpestris.
alpestre, Stev. ex DC. Prod. iv. 116 = Silaus carvifolius.
alpigenum, *Boiss. Fl. Orient.* ii. 1020.—As. Min.

PEUCEDANUM:—
alsaticum, *Linn. Sp. Pl.* ed. II. 354.—Europ. austr.; As. bor.
alsaticum, Poir. Encyc. v. 227 = officinale.
altissimum, Desf. Tabl. ed. I. 119 = officinale.
altum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 20.—Afr. trop.
ambiguum, Hook. Lond. Journ. Bot. vi. (1847) 235 = Geyeri.
ambiguum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 626.—Am. bor. occ.
Ammoniacum, Baill. Hist. Pl. vii. 186 = Dorema Ammoniacum.
anagymnodes, Pall. Reise in Suedl. Statth. Russ. Reich. i. 134 = paucifolium.
anamallayense, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 710.—Ind. or.
anatolicum, Boiss. Fl. Orient. ii. 985 = Ferula anatolica.
angelicaefolium, Turcz. in Bull. Soc. Nat. Mosc. (1838) i. 93; xvii. (1844) 738 = Angelica sylvestris.
angustifolium, *Reichb. f. Ic. Fl. Germ.* xxi. 62.—Italia.
araliaceum, *Benth. & Hook. f. ex Vatke, in Linnaea*, xl. (1876) 188.—Afr. trop.
arenarium, Hort. ex Spreng. Umb. Prod. 13, 32 = Silaus tenuifolius.
arenarium, *Waldst. & Kit. Pl. Rar. Hung.* i. 19. t. 20.—Hungar.
Asa-foetida, Baill. Hist. Pl. vii. 98, 185 = Ferula Assa-foetida.
atlanticum, *Pomel, Nouv. Mat. Fl. Atl.* 144.—Algeria.
atropurpureum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 21.—Afr. trop.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 315 = Dorema Aucheri.
augymnodes, Cusson, ex Ledeb. Fl. Ross. ii. 272 = paucifolium.
aureum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 341.—Ins. Canar.
aureum, Spreng. Syst. i. 911 = Ferula soongarica.
austriacum, *Koch, in Nov. Act. Nat. Cur.* xii. i. (1824). 94.—Europ.
baicalense, *Koch, l. c.*—Hungar.; Sibir.
Bellardi, *Lag. ex Sweet, Hort. Brit.* ed. II. 250 (Quid?).—Pedemont.
Bellardi, Nym. Consp. 285, in syn. = austriacum.
Besserianum, DC. Prod. iv. 177 = ruthenium.
bicolor, *S. Wats. in Bot. King's Exp.* 129.—Am. bor. occ.
Bourgaei, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 42 = P. Oreoselinum.
caledonicum, *Eckl. & Zeyh. Enum.* 349.—Afr. austr.
californicum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 628 = caruifolium.
campestre, *Janka, in Linnaea*, xxx. (1859–60) 570.—Hungar.
canescens, Ledeb. Fl. Alt. i. 307 = Ferula canescens.
capense, D. Dietr. Syn. Pl. ii. 967 (= *Bubon capensis*).—Afr. austr.
capense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 554.—Afr. austr.
capillaceum, DC. Prod. iv. 178 = Sieberianum.
capillaceum, *Thunb. Prod. Pl. Cap.* 50.—Afr. austr.
caruifolium, *Torr. & Gray, Fl. N. Am.* i. 628.—Am. bor. occ.
carvifolium, Bieb. ex Ledeb. Fl. Ross. ii. 285 = Silaus carvifolius.
carvifolium, *Vill. Prosp.* 25; *Hist. Pl. Dauph.* ii. 638.—Europ.
caspicum, Link, Enum. Hort. Berol. i. 272 = Ferula caspica.
caucasicum, *C. Koch, in Linnaea*, xvi. (1842) 358.—Reg. Caucas.
Cephalotes, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 314.—Rossia.
Cervaria, *Cusson, ex Lapeyr. Hist. Abr. Pl. Pyr.* 149.—Europ. med.; As. bor.
cervariaefolium, *C. A. Mey. Verz. Pfl. Cauc.* 126.—Persia.
ceylanicum, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 14.—Zeylan.
Chabraei, Reichb. ex Moesl. Handb. ed. II. 448 = carvifolium.
chaerophylloides, Less. ex Ledeb. Fl. Ross. ii. 290 = Selinum Benthami.
chrysanthum, *Boiss. & Bal. Diagn.* Ser. II. vi. 86.—As. Min.
cnidioides, *Boiss. & Heldr. l. c.* ii. 90.—Graecia.

PEUCEDANUM:—
collinum, *D. Dietr. Syn. Pl.* ii. 969.—Afr. austr.
connatum, *E. Mey. in Drège, Zwei Pfl. Docum.* 210, nomen; *et ex Sond. in Harv. & Sond. Fl. Cap.* ii. 557.—Afr. austr.
cordatum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 514.—Ins. Socotr.
coriaceum, *Reichb. Fl. Germ. Excurs.* 866.—Europ. austr.
creticum, DC. Prod. iv. 182 = Ferula nodosa.
creticum, *Spreng. Neue Entdeck.* ii. 148.—Ins. Creta.
Crouanorum, Bor. ex Nym. Consp. 286 = palustre.
cuneifolium, Vill. ex Steud. Nom. ed. I. 607 = Selinum Carvifolia.
cupulare, *Boiss. Diagn.* Ser. I. vi. 63.—Persia.
Cynorhiza, *Sond. in Harv. & Sond. Fl. Cap.* ii. 558.—Afr. austr.
Dana, *Buch.-Ham. in Wall. Cat.* n. 7216.—Ind. or.
dasycarpum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 254.—Turkestan.
dasycarpum, *Torr. & Gray, Fl. N. Am.* i. 628.—Calif.
daucifolium, Nutt. ex Torr. & Gray, l. c. 627 = foeniculaceum.
davuricum, Turcz. ex Ledeb. Fl. Ross. ii. 314 = terebinthaceum.
decussatum, Dulac, Fl. Hautes-Pyr. 343 = carvifolium.
depauperatum, *Boiss. & Bal. Diagn.* Ser. II. v. 98.—As. Min.; Syria.
dissectum, DC. Prod. iv. 178 = Selinum dissectum.
dissectum, Ledeb. Fl. Alt. i. 306 = Ferula dissecta.
distans, Griseb. Spicil. Fl. Rumel. i. 374 = Johrenia graeca.
divaricatum, *Benth. & Hook. f. ex Arcang. Comp. Fl. Ital.* 294.—Europ.
diversifolium, Benth. & Hook. f. Gen. i. 919 = Heracleum Wallichii.
Dregeanum, D. Dietr. Syn. Pl. ii. 969 = Ecklonianum?
dubium, Ledeb. Fl. Alt. i. 310 = Ferula gracilis.
Ecklonianum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 555.—Afr. austr.
Ecklonis, Walp. Rep. ii. 410 = capillaceum.
elatum, Ledeb. Fl. Alt. i. 304 = Ferula soongarica.
elongatum, *E. Mey. ex Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 535.—Afr. austr.
euryptera, *A. Gray, in Proc. Am. Acad.* vii. (1868) 348.—Calif.
falcaria, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) ii. 192.—As. centr.
farinosum, *Geyer, in Hook. Lond. Journ. Bot.* vi. (1847) 235.—Am. bor. occ.
Ferula, Baill. Hist. Pl. vii. 186 = Ferula communis.
ferulaceum, Eckl. & Zeyh. Enum. 349 = Sieberianum.
ferulaceum, *Thunb. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 556.—Afr. austr.
ferulaeoides, Steud. Nom. ed. II. ii. 311 = Ferula resinosa.
filiforme, Gilib. Fl. Lituan. ii. 15 = officinalis.
flavum, Salisb. Prod. 163 = Silaus flavescens.
Fleischmanni, *Benth. & Hook. f. Gen.* i. 920.—Carinth.
flexuosum, Kit. ex Rochel, Pl. Rar. Banat. 26 = Seseli annuum.
flexuosum, Moench, Meth. 82 = alsaticum.
foeniculaceum, Cooper, in Pacif. Rail. Rep. xii. 63 = millefolium, *S. Wats.*?
foeniculaceum, Newberry, in Pacif. Rail. Rep. vi. 75 = macrocarpum?
foeniculaceum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 627.—Am. bor. occ.
foeniculaceum, Torr. Bot. Mex. Bound. 70 = dasycarpum, *Torr. & Gray*.
fraxinifolium, *Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 79.—Afr. trop.
frutescens, E. Mey. in Drège, Zwei Pfl. Docum. 210 = capense.
galbanifluum, Baill. Hist. Pl. vii. 185 = Ferula galbaniflua.
Galbanum, *Benth. & Hook. f. Gen.* i. 920.—Afr. austr.
gallicum, *Latour. Chlor. Lugd.* 7; *Pers. Syn.* i. 310.—Europ.
geniculatum, Forst. f. Prod. 22 = Angelica geniculata.
Geyeri, *S. Wats. Bibliog. Ind. N. Am. Bot.* 428.—Am. bor. occ.
glaucifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = P. Cervaria.

PEUCEDANUM:—
glaucum, *DC. Prod.* iv. 179.—Reg. Himal.
glaucum, Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 712 = Heracleum cachemiricum.
glaucum, Link, Enum. Hort. Berol. i. 273 = Ferula glauca.
gracile, Ledeb. Fl. Alt. i. 308 = Ferula gracilis.
graminifolium, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 314.—Bithynia.
grande, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 710.—Ind. or.
Grantii, *Kingston, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 79.—Afr. trop.
graveolens, *Benth. & Hook. f. Gen.* i. 919.—Europ. et As. trop.
graveolens, *S. Wats. in Bot. King's Exped.* 128.—Am. bor. occ.
Hallii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 141.—Am. bor. occ.
heterophyllum, Vis. Cat. Sem. Hort. Patav. (1836) 3, 4 = carvifolium.
hispanicum, *Endl. ex Walp. Rep.* ii. 411.—Hispan.
Hooshe, Baill. Hist. Pl. vii. 186 = Ferula Assa-foetida?
Howellii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 369.—Am. bor.
humile, Turcz. in Bull. Soc. Nat. Mosc. (1838) i. 93; xvii. (1844) 745 = salinum.
Hystrix, *Bunge, Verz. Suppl. Fl. Alt.* 23; *et in Mém. Sav. Etr. Pétersb.* ii. (1835) 538.—Sibir.
iletense, *Spreng. Gesch. Bot.* ii. 200; *Sp. Umbell.* 57.—Sibir.
imperatorioides, DC. Prod. iv. 183, in syn. = angustifolium.
involucratum, *Koch, in Nov. Acad. Nat. Cur.* xii. i. (1824) 94.—Europ.
isetense, Spreng. ex Schult. Syst. vi. 569, sphalm. = iletense.
italicum, Mill. Gard. Dict. ed. VIII. n. 2 = officinale.
japonicum, Thunb. Fl. Jap. 117 = Ligusticum acutifolium.
junceum, Humb. & Bonpl. ex Spreng. in Schult. Syst. vi. 572 = Eulophus peucedanoides.
karatawiense, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 598.—Turkestan.
Kotschyi, Boiss. Diagn. Ser. I. vi. 62 = Dorema Aucheri.
laciniatum, Heynh. Nom. ii. 504 = Ferula dissecta.
laevigatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 627 = ambiguum.
lanceolatum, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 536 = capense.
lancifolium, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 39.—Hispan.
lasiocarpum, Boiss. Diagn. Ser. II. ii. 88 = Zosima lasiocarpa.
lateriflorum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 557.—Afr. austr.
latifolium, *DC. Prod.* iv. 181.—Europ. med.; Reg. Caucas.
latifolium, Nutt. ex Torr. & Gray, Fl. N. Am. i. 625 = Nuttallii.
lativittatum, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 14.—Soongaria.
Latourrettii, Cusson. ex Steud. Nom. ed. I. 607, in syn. = gallicum.
Ledebouri, G. Don, Gen. Syst. iii. 330 = officinale.
Ledebourii, Steud. Nom. ed. II. ii. 311 = Ferula dissecta.
leiocarpum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 626.—Am. bor. occ.
leptocarpum, Nutt. ex Torr. & Gray, l. c. 626 = triternatum, *Nutt.*
ligusticoides, Less. ex Ledeb. Fl. Ross. ii. 290 = Selinum Benthami.
longifolium, *Waldst. & Kit. Pl. Rar. Hung.* iii. 279.—Banat.
ludovicianum, Rafin. ex Torr. & Gray, Fl. N. Am. i. 629 = verticillare?
lusitanicum, Hoffmgg. ex Spreng. in Schult. Syst. vi. 571 = Ferula sulcata.
macedonicum, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 203.—Macedon.
macrocarpum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 627.—Am. bor. occ.
macrocarpum, Parry, in Am. Natural. ix. (1875) 272 = Parryi.
macrocoleum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 310 = Ferula macrocolea.

PEUCEDANUM:—
magalismontanum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 558.—Afr. austr.
marginatum, Benth. Pl. Hartw. 312 = caruifolium.
Matthioli, Spreng. in Schult. Syst. vi. 569 = Silaus tenuifolius.
meifolium, *Boiss. Diagn.* Ser. I. x. 34.—Syria.
membranaceum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 315.—Persia.
mexicanum, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 361.—N. Mexic.
Meyeri, Boiss. Fl. Orient. ii. 1018 = paucifolium.
microcoleum, Boiss. Diagn. Ser. I. vi. 62 = Ferula microcolea.
microphyllum, Sm. in Rees, Cycl. xxvii. n. 11 = salinum.
millefolium, *Sond. in Harv. & Sond. Fl. Cap.* ii. 559.—Afr. austr.
millefolium, *S. Wats. in Bot. King's Exped.* 129.—Am. bor. occ.
minus, All. Fl. Pedem. ii. 6 = carvifolium.
minus, Huds. Fl. Angl. ed. I. 101 = Trinia vulgaris.
minus, Poir. Encyc. v. 228 = Silaus longifolius.
montanum, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 94 = austriacum.
Morisoni, Bess. ex Schult. Syst. vi. 567 = ruthenicum.
Munbyi, *Boiss. Diagn.* Ser. II. ii. 89.—Afr. bor. occ.
neapolitanum, Baill. Hist. Pl. vii. 98 = Ferula glauca?
nebrodense, Nym. Syll. 153 = carvifolium.
Neumayeri, *Reichb. f. Ic. Fl. Germ.* xxi. 63.—Europ. or.
nevadense, *S. Wats. in Proc. Am. Acad.* xi. (1876) 143.—Calif.
Newberryi, S. Wats. in Am. Natural. vii. (1873) 301 = Ferula Newberryi.
nodosum, Linn. Sp. Pl. 246 = Ferula nodosa.
nudicaule, Boland. Cat. ex S. Wats. Bibliog. Ind. N. Am. Bot. 429 = macrocarpum.
nudicaule, A. Gray, in Proc. Am. Acad. viii. (1872) 385 = Hallii.
nudicaule, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 627.—Am. bor. occ.
nudicaule, S. Wats. in Bot. King's Exped. 130 = nevadense.
nudiceps, Cusson, ex Ledeb. Fl. Ross. ii. 272 = paucifolium.
Nuttallii, Walp. Rep. ii. 411 = triternatum, *Nutt.*
Nuttallii, *S. Wats. in Bot. King's Exped.* 128.—Calif.
obtusifolium, Sibth. & Sm. Fl. Graec. Prod. i. 189; Fl. Graec. iii. 70 = Malabaila obtusifolia.
officinale, Bess. Prim. Fl. Galic. i. 208 = Ferula sylvatica.
officinale, Gorter, Fl. Ingr. 42 = Selinum Benthami?
officinale, *Linn. Sp. Pl.* 245.—Europ. med.; As. bor.
officinale, Pall. Tabl. Taur. 48, nomen = ruthenicum.
officinale, Thuill. Fl. Par. ed. I. 140 = gallicum.
officinale, Ucria, Hort. Reg. Panorm. 128 = Seseli Bocconi.
oopodum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 100 = Ferula oopoda.
opacum, Franch. Fl. Loir-et-Cher, 248 = sativum.
orbiculare, Dulac, Fl. Hautes-Pyr. 344 = P. Oreoselinum.
Oreopansil, *Griseb. in Goett. Abh.* xxiv. (1879) 147.—Reg. Argent.
Oreoselinum, *Moench, Meth.* 82.—Europ. med. et austr.
orientale, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 309 = Ferula orientalis.
orientale, G. Don, Gen. Syst iii. 332 = Selinum dissectum.
Ostruthium, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 95.—Europ.
ovinum, Boiss. Diagn. Ser. I. vi. 61 = Ferula ovina.
Palimbia, Baill. Hist. Pl. vii. 100 = carvifolium?
palimboides, *Boiss. Fl. Orient.* ii. 1021.—As. Min.
Pallasii, Cusson, ex Ledeb. Fl. Ross. ii. 272 = paucifolium.
palustre, *Moench, Meth.* 82.—Europ.
palustre, Sieber, ex Spreng. Sp. Umbell. 73 = alsaticum.
paniculatum, Ledeb. Fl. Alt. i. 310 = Ferula resinosa.
paniculatum, *Loisel. Fl. Gall.* 722.—Corsica.
parisiense, DC. Fl. Fr. iv. 336 = gallicum.
parisiense, Koch, Syn. Fl. Germ. ed. I. 303 = coriaceum.
Parryi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 143.—Am. bor. occ.

PEUCEDANUM:—
parvifolium, *Torr. & Gray, Fl. N. Am.* i. 628.—Am. bor. occ.
Pastinaca, Baill. Hist. Pl. vii. 96, 188 = sativum.
pastinacaefolium, *Boiss. & Hohen. Diagn.* Ser. I. x. 35.—Persia.
paucifolium, *Ledeb. Fl. Ross.* ii. 312.—Reg. Caucas.; Persia.
persicum, Baill. Hist. Pl. vii. 186 = Ferula persica.
petiolare, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 312.—Persia.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 327.—Abyss.
petraeum, Noé, ex Koch, Syn. Fl. Germ. ed. I. 304 = Schottii.
Petroselinum, Desf. Tabl. ed. II. 272; Cat. Hort. Par. ed. III. 405 = Carum Petroselinum.
Petteri, Vis. in Cat. Sem. Hort. Patav. (1836) 3, 4. n. 8 = coriaceum.
pisidicum, *Boiss. Fl. Orient.* ii. 1021.—Pisidia.
platycarpum, *E. Mey. in Drège, Zwei Pfl. Docum.* 210, nomen; *et ex Harv. & Sond. Fl. Cap.* ii. 557.—Afr. austr.
platytaenium, Boiss. & Bal. Diagn. Ser. II. vi. 87 = ruthenicum.
podolicum, Eichw. Skizze, 155 = carvifolium.
polymorphum, Lag. ex Sweet, Hort. Brit. ed. II. 250 (Quid?).—Rossia.
polyphyllum, Ledeb. Fl. Alt. i. 314 = baicalense.
polyscias, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 318.—Persia.
pratense, Lam. Fl. Fr. iii. 469 = Silaus flavescens.
pschawicum, *Boiss. Fl. Orient.* ii. 1020.—Reg. Caucas.
Pseudoreoselinum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 36.—Turkestan.
puberulum, Turcz. in Bull. Soc. Nat. Mosc. (1838) 1. 93 = vaginatum.
pungens, *E. Mey. in Drège, Zwei Pfl. Docum.* 210, nomen; *et ex Sond. in Harv. & Sond. Fl. Cap.* ii. 557.—Afr. austr.
pyramidatum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 366 = Ferula paniculata.
pyrenaicum, Costa, ex Steud. Nom. ed. I. 209, in syn. = alsaticum.
rablense, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 94 = austriacum.
ramosissimum, *D. Dietr. Syn. Pl.* ii. 967.—Reg. Himal.
rapiferum, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) 1. 326.—As. centr.
redivivum, *Pall. in Nov. Act. Petrop.* ii. (1779) 252. t. 8.—Rossia.
refractum, Vill. ex Spreng. Sp. Umbell. 52 = gallicum.
Renardi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 596.—Turkestan.
rhombifolium, Stokes, Bot. Mat. Med. ii. 82 = P. Galbanum.
rigidius, Baill. Hist. Pl. vii. 187 = ternatum.
rigidum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 106.—China.
rigidum, Eckl. & Zeyh. Enum. 350 = capillaceum.
rigidum, E. Mey in Drège, Zwei. Pfl. Docum. 210; et ex Sond. in Harv. & Sond. Fl. Cap. ii. 554 = capense.
Rochelianum, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 27 = ruthenicum.
rubricaule, Baill. Hist. Pl. vii. 186 = Ferula rubricaulis.
rupestre, Boiss. & Bal. Diagn. Ser. II. vi. 85 = Ferula orientalis.
rupestre, *Hoffmgg. ex Spreng. in Schult. Syst.* vi. 572.—Lusitan.
ruthenicum, *Bieb. Fl. Taur. Cauc.* i. 215.—Rossia; Reg. Caucas.
ruthenicum, Kováts, ex Janka, in Linnaea, xxx. (1859–60) 570 = campestre.
salinum, *Pall. ex Spreng. Syst.* i. 910.—Sibir. alt.
salsum, Steud. Nom. ed. II. ii. 311 = redivivum.
sativum, *Benth. & Hook. f. Gen.* i. 920.—Europ.; Am. bor.
saxatile, Noé, ex Bertol. Fl. Ital. iii. 355 = carvifolium.
Schwereckii, Eichw. Skizze, 155 = palustre.
Schlechtendalii, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 311 = Ferula Schlechtendalii.
Schottii, *Bess. ex DC. Prod.* iv. 178.—Podolia.
sclerophyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1015.—Persia.
scoparium, *Boiss. Diagn.* Ser. II. ii. 90.—Persia.

PEUCEDANUM:—
selinoides, DC. Prod. iv. 181 (= *Pachypleurum alpinum*).—Sibir. altaic.
serotinum, Pers. Syn. i. 310 = Silaus tenuifolius.
seseloides, Turcz. in Bull. Soc. Nat. Mosc. (1838) 1. 93; xvii. (1844) 743 = vaginatum.
Shair, Baill. Hist. Pl. vii. 186 = Ferula Schair.
sibiricum, DC. Prod. iv. 178, partim = Ferula tatarica.
sibiricum, Waldst. & Kit. Pl. Rar. Hung. i. 60 = Ferula Sadleriana.
sibiricum, Willd. Sp. Pl. i. 1406 = Ferula soongarica.
Sieberianum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 556.—Afr. austr.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 63.—Japon.
sikkimense, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 710.—Reg. Himal.
silaifolium, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 20.—Afr. trop.
Silaus, Bieb. Fl. Taur. Cauc. i. 215 = Silaus alpestris.
Silaus, Linn. Sp. Pl. 246 = Silaus flavescens.
simplex, *Nutt. ex S. Wats. in Bot. King's Exped.* 129.—Am. bor. occ.
skardicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 711.—Reg. Himal.
soongaricum, G. Don, Gen. Syst. iii. 332 = Ferula paniculata.
Sowa, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 116 = graveolens, *Benth. & Hook. f.*
Spreitzenhoferi, *Dingl. in Flora*, lxvi. (1883) 210.—Oriens.
Sprengelii, *DC. Prod.* iv. 182.—Hab.?
stadense, Eckl. & Zeyh. Enum. 349 = ferulaceum.
stenocarpum, *Boiss. & Reut. ex Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 311.—Hispan.
striatum, Sond. in Harv. & Sond. Fl. Cap. ii. 555 = collinum.
Suksdorfii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 369.—Am. bor.
sulcatum, *Nym. Syll.* 152.—Ital.
sulcatum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 559.—Afr. austr.
Sumbul, Baill. Hist. Pl. vii. 186 = Ferula Sumbul.
sylvaticum, Baill. l. c. 187 = Ferula sylvatica.
sylvestre, DC. Prod. iv. 179 = palustre.
Szowitzianum, Baill. Hist. Pl. vii. 186 = Ferula Szowitziana.
tauricum, Bieb. Fl. Taur. Cauc. i. 215 = ruthenicum.
tauricum, Desf. Cat. Hort. Par. ed. III. 203, 361 = Silaus carvifolius.
tenuifolium, Eckl. & Zeyh. Enum. 349 = ferulaceum.
tenuifolium, Hort. Par. ex Poir. Encyc. v. 228 = Silaus tenuifolius.
tenuifolium, *Thunb. Prod. Pl. Cap.* 50.—Afr. austr.
tenuisectum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 37.—Turkest.
tenuissimum, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 235 = ambiguum.
terebinthaceum, *Fisch. ex Reichb. Fl. Germ. Excurs.* 454.—Europ.
teretifolium, *Wood, Am. Bot. & Flor.* 136.—Am. bor.
ternarium, Pall. ex Ledeb. Fl. Ross. ii. 271 = paucifolium.
ternatum, *Nutt. Gen. Am.* i. 182.—Am. bor.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 711.—Reg. Himal.
tolucense, *Hemsl. Biol. Centr. Am. Bot.* i. 570.—Mexic.
tomentosum, Benth. Pl. Hartw. 312 = dasycarpum, *Torr. & Gray.*
transiliense, *Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 78.—As. centr.
transsilvanicum, Schur, Enum. Pl. Transs. 265 = austriacum.
trilobatum, Gilib. Fl. Lituan. ii. 14 = Silaus flavescens.
triternatum, *Eckl. & Zeyh. Enum.* 349.—Afr. austr.
triternatum, A. Gray, in Proc. Am. Acad. viii. (1873) 385 = leiocarpum.
triternatum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 626.—Am. bor. occ.
triternatum, Torr. in Sitgreave, Rep. 160 = simplex.
triternatum, Torr. & Gray, in Pacif. Rail. Rep. ii. 121 = Carum Gairdneri.
uliginosum, *D. Dietr. Syn. Pl.* ii. 967.—Afr. austr.
utriculatum, *Nutt. ex Torr. & Gray, Fl. N. Am.* 628.—Am. bor. occ.

PEUCEDANUM:—
vaginale, Pall. ex Ledeb. Fl. Ross. ii. 300 = Ferula tatarica.
vaginatum, *Ledeb. Fl. Alt.* i. 312.—Sibir. alt.
varium, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 15.—Soongaria.
venetum, *Koch, Syn. Fl. Germ.* ed. I. 305.—Europ. austr.
verticillare, *Spreng. Syst.* iv. *Cur. Post.* 121.—Europ. austr.
verticillatum, Rafin. Fl. Ludov. 81 = Discopleura Nuttallii.
villosum, *Nutt. ex S. Wats. in Bot. King's Exp.* 131.—Calif.
virgatum, Cham. ex Schlecht. in Linnaea, i. (1826) 392 = capense.
vittijugum, *Boiss. Fl. Orient.* ii. 1018.—Graecia.
Vocontiorum, Spreng. Sp. Umbell. 51 = Heracleum minimum.
Wallichianum, DC. Prod. iv. 181 = Selinum tenuifolium.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* ii. 559.—Afr. austr.
Zeyheri, Steud. Nom. ed. II. ii. 312 = capillaceum.
zeylanicum, Thw. Enum. Pl. Zeyl. 131 = ceylanicum.

PEUCEPHYLLUM, A. Gray, in Bot. Mex. Bound. 74 (1858). *COMPOSITAE*, Benth. & Hook. f. ii. 438.
Schottii, *A. Gray, l. c.*—Am. bor. occ.

PEUDANUM, Dingl. in Flora, lxvi. (1883) 210, sphalm. = **Peucedanum**, Tourn. (Umbell.).

PEUMUS, Molina, Sagg. Chile, 185 (1782), partim; Pers. Syn. ii. 629 (1807). *MONIMIACEAE*, Benth. & Hook. f. iii. 141, 150.
Boldea, Juss. in Ann. Mus. Par. xiv. (1809) 134.
Boldu, Feuill. ex Adans. Fam. ii. 446 (1763).
Boldus, Schult. f. Syst. vii. p. x. 56 (1829).
Ruizia, Ruiz & Pav. Prod. 135. t. 29. (1794).
alba, Molina, Sagg. Chile, 185 = Cryptocarya Peumus.
Boldus, *Molina. l. c.* 185.—Chili.
fragrans, Pers. Syn. ii. 629 = P. Boldus.
mammosa, Molina, Sagg. Chile, 185 = Cryptocarya Peumus.
rubra, Molina, l. c. = Cryptocarya Peumus.

PEUROUSEA, Stend. Nom. ed. II. i. 312 (1841), sphalm. = **Peyrousea**, DC. (Compos.).

PEUTALIS, Rafin. New Fl. Am. iv. 47 (1836); Fl. Tellur. iii. 14 (1836) = **Polygonum**, Tourn.
divergens, Rafin. l. c. 48 (Quid?).—Am. bor.
floridana, Rafin. l. c. 47 (Quid?).—Am. bor.
glomerata, Rafin. l. c. 48 (Quid?).—Am. bor.
heterophyla, Rafin. l. c. (Quid?).—Am. bor.
Hydropiper, Rafin. Tellur. iii. 14 = Polyg. Hydropiper.
incana, Rafin. l. c. = Polyg. lapathifolium.
ligularis, Rafin. New Fl. Am. iv. 47 (Quid?).—Am. bor.
linearis, Rafin. Fl. Tellur. iii. 16 (Quid?).—Hab.?
longifolia, Rafin. New Fl. Am. iv. 48.—Am. bor.
minus, Rafin. Fl. Tellur. iii. 14 = Polyg. minus.
nodosa, Rafin. l. c. = Polyg. nodosum.
palustris, Rafin. New Fl. Am. iv. 50 (Quid?).—Am. bor.
Persicaria, Rafin. Fl. Tellur. iii. 14 = Polyg. Persicaria.
polystachya, Rafin. New Fl. Am. iv. 50 = Polyg. polystachyum.
punctata, Rafin. l. c. 48 = Polyg. punctatum, *Rafin.*
scabra, Rafin. l. c. = Polyg. lapathifolium?

PEVRAEA, Comm. ex Juss. Gen. 320 (1789) = POIVREA, Comm. = **Combretum**, Linn.

PEYRITSCHIA, Fourn. ex Benth. & Hook. f. Gen. iii. 1158 (1883) = **Deschampsia**, Beauv. (Gramin.).
koelerioides, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 555 = D. koelerioides.

PEYROUSEA, DC. Prod. vi. 76 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 432.
Lapeirousia, Thunb. Prod. Pl. Cap. 163 (1800).
Lapeyrousia, Spreng. Anleit. ii. II. 552 (1818).
Osmitiphyllum, Sch. Bip. in Flora, xxvii. (1844) 675.
calycina, *DC. Prod.* vi. 77.—Afr. austr.
oxylepis, DC. l. c. = calycina.

PEYROUSIA, Poir. in Dict. Sc. Nat. xxxix. 363 (1826) = **Lapeyrousia**, Pourr. (Irid.).
aculeata, Sweet, Brit. Flow. Gard. Ser. II. t. 39 = L. compressa.
anceps, Poir. in Dict. Sc. Nat. xxxix. 363 = L. compressa.
bracteata, Sweet, Hort. Brit. ed. II. 499 = L. fissifolia.
corymbosa, Poir. in Dict. Sc. Nat. xxxix. 365 = L. corymbosa.
Fabricii, Sweet, Hort. Brit. ed. II. 499 = L. Fabricii.
falcata, Sweet, l. c. = L. falcata.
fasciculata, Sweet, l. c. = L. fasciculata.
fissifolia, Poir. in Dict. Sc. Nat. xxxix. 364 = L. fissifolia.
juncea, Poir. l. c. = L. juncea.
marmorata, Poir. l. c. = L. juncea.
silenoides, Poir. l. c. 365 = L. silenoides.

PEYRUSA, Rich. ex Dun. in DC. Prod. vii. 560, in syn. (1839) = **Hornemannia**, Vahl (Vaccin.).
royenoides, Rich. ex Dun. l. c. = H. martinicensis.

PEYSSONELIA, Boiv. ex Webb & Berth. Phyt. Canar. ii. 35, in obs. ('1836–50') = **Cytisus**, Linn. (Legumin.).

PFAFFIA, Mart. Nov. Gen. et Sp. ii. 20. t. 122-124 (1826). *AMARANTACEAE*, Benth. & Hook. f. iii. 37.
Serturnera, Mart. Nov. Gen. et Sp. ii. 36. t. 136. (1826).
glabrata, *Mart. Nov. Gen. et Sp.* ii. 21. t. 122.—Bras.
glauca, *Spreng. Syst.* iv. *Cur. Post.* 107.—Bras.
gnaphaloides, *Mart. Nov. Gen. et Sp.* ii. 24.—Bras.
hirtula, *Mart. l. c.*—Bras.
iresinoides, Spreng. Syst. iv. Cur. Post. 103, 106 = Gomphrena iresinoides.
jubata, *Mart. Nov. Gen. et Sp.* ii. 24.—Bras.
lanata, Gibert, Enum. Pl. Montev. 73 = tomentosa.
luzulaeflora, D. Dietr. Syn. Pl. i. 868 = Gomphrena luzulaeflora.
puberula, Spreng. Syst. iv. Cur. Post. 106 = Telanthera puberula.
rufa, Spreng. l. c. = Telanthera rufa.
sericea, *Mart. Nov. Gen. et Sp.* ii. 22. t. 123.—Bras.
tomentosa, *Mart. l. c.* 24.—Bras.
velutina, *Mart. l. c.* 23. t. 124.—Bras.
villosa, Spreng. Syst. iv. Cur. Post. 106 = Telanthera Martii.

PFEIFFERA, Salm-Dyck, Cact. Hort. Dyck. ed. I. 40 (1845) = **Rhipsalis**, Gaertn. (Cact.).
cereiformis, Salm-Dyck, l. c. 41 = R. cereiformis.

PFEIFFERIA, Buching. in Ann. Sc. Nat. Sér. III. v. (1846) 88 = **Cuscuta**, Tourn. (Convolv.).
suaveolens, Buching. l. c. = C. racemosa.

PFUNDIA, Opiz, Seznam, 73 (1852) = **Ranunculus**, Tourn.
arvensis, Opiz, l. c.; et in Lotos, iv. (1854) 43 = R. arvensis.

PHACA, Linn. Coroll. Gen. 13 (1737) = **Astragalus**, Linn. (Legumin.).
abbreviata, Ledeb. Fl. Alt. iii. 268 = A. alpinus.
aboriginorum, Hook. Fl. Bor. Am. i. 143 = A. aboriginorum.
alpina, Linn. Sp. Pl. 755 = A. alpinus.
alpina, O. F. Muell. in Fl. Dan. t. 856 = A. frigidus.
amoena, Phil. in Linnaea, xxviii. (1856) 630.—Chili.
andicola, Gill. in Hook. Bot. Misc. iii. (1833) 105 = *carinata*.
andina, Nutt. ex Torr. & Gray, Fl. N. Am. i. 345 = A. alpinus.
annua, Geyer, in Hook. Lond. Journ. Bot. vi. (1847) 213 = A. Geyeri.
arenaria, Pall. Reise, iii. 748 = A. chorinensis.
argophylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 342 = A. triphyllus.
aristata, [Clairv.] Man. Herb. 133 = A. aristatus.
Arnottiana, Gill. in Hook. Bot. Misc. iii. (1833) 184.—Chili.
aspera, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = A. asper.
astragalina, DC. Astrag. 52. n. 9 = A. alpinus.
astragalina, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 51.—Mexic.

PHACA:—
atrata, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 760 = Astragalus australis.
australis, Ledeb. Fl. Alt. iii. 270 = A. vaginatus.
australis, Linn. Mant. i. 103; ii. 448 = A. lusitanicus.
austriaca, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = A. austriacus.
baetica, Linn. Sp. Pl. 755 = A. lusitanicus.
Bayeri, Janka, in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 431 = A. Haarbachii.
Berteriana, Moris, in Mem. Acc. Torin. xxxvii. (1834) 105 = *canescens*.
bisulcata, Hook. Fl. Bor. Am. i. 145 = A. bisulcatus.
brachycarpa, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 185 = A. zacharensis.
brachyptera, Phil. in Linnaea, xxxiii. (1864–65) 44.—Chili.
brachytropis, Stev. in Mém. Soc. Nat. Mosc. iv. (1812) 93 = A. brachytropis.
bracteosa, Kar. & Kir. l. c. xv. (1842) 323 = A. aksuensis.
Bustillosi, Phil. in Linnaea, xxviii. (1856) 680.—Chili.
cachemerica, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 126 = A. himalayanus.
caespitosa, Nutt. Gen. Am. ii. 98 = A. triphyllus.
campestris, Wahlenb. Veg. Helv. 130 = Oxytropis campestris.
candidissima, Benth. Bot. Voy. Sulph. 13 = A. candidissimus.
Candolleana, H. B. & K. Nov. Gen. et Sp. vi. 495. t. 586 = A. triflora.
canescens, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 185.—Chili.
canescens, Nutt. in Torr. & Gray, Fl. N. Am. i. 344 = A. leucopsis.
carinata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 185.—Chili.
Chamissonis, Vog. in Linnaea, x. (1836) 592.—Chili.
chilensis, *Nees, l. c.* x. (1836) *Litt.* 72.—Chili.
chrysantha, Moris, in Mem. Acc. Torin. xxxvii. (1834) 104.—Chili.
clandestina, Phil. Fl. Atac. 14.—Chili.
collina, Hook. Fl. Bor. Am. i. 141 = A. collinus.
concinna, Phil. in Linnaea, xxxiii. (1864–65) 44.—Chili.
coquimbensis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 184.—Chili.
cretacea, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 452 = A. missouriensis.
Crotalariae, Benth. Pl. Hartw. 307 = A. oocarpus.
Cruckshanksii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 184.—Chili.
curvicaulis, Clos, in C. Gay, Fl. Chil. ii. 102.—Chili.
debilis, Nutt. in Torr. & Gray, Fl. N. Am. i. 345 = A. debilis.
densifolia, Sm. in Rees, Cycl. xxvii. n. 9 = A. Menziesii.
densifolia, Torr. in Pacif. Rail. Rep. iv. 80 = A. oocarpus.
depauperata, Phil. Fl. Atac. 14.—Chili.
digitata, Torr. in Frém. Rep. (1845) 89 = A. tridactylicus.
Douglasii, Torr. & Gray, Fl. N. Am. i. 346 = A. Douglasii.
Edmonstonei, Hook. f. in Trans. Linn. Soc. xx. (1847) 227.—Ins. Galap.
elata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 185.—Chili.
elegans, Hook. Fl. Bor. Am. i. 144 = A. oroboides.
elongata, Hook. l. c. 140 = A. flexuosus.
elongata, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 51.—Mexic.
elongata, Phil. in Linnaea, xxxiii. (1864–65) 43.—Chili.
exaltata, Fisch. ex Sweet, Hort. Brit. ed. II. 145 = A. secundus?
fastidia, Kellogg, in Hesperian (June 1860), cum fig.; ex Curran, in Bull. Calif. Acad. i. III. (1885) 136 = A. Coulteri?
Fendleri, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 36 = A. Fendleri.
flava, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 186.—Chili.
flexuosa, Hook. Fl. Bor. Am. i. 140 = A. flexuosus.
floridana, Willd. Sp. Pl. iii. 1252 = Sesbania platycarpa.
frigida, Linn. Syst. ed. X. 1173 = A. frigidus.
Gerardi, Vill. Prosp. 43; Hist. Pl. Dauph. iii. 474 = A. australis.
glabra, Clar. in Bull. Soc. Philom. iii. (1811) 99 = A. australis.

PHACA :—
glabriuscula, Hook. Fl. Bor. Am. i. 144 = Astragalus glabriusculus.
gracilenta, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 35 = A. gracilentus.
grata, Clos, in C. Gay, Fl. Chil. ii. 92.—Chili.
halicacaba, Willd. Sp. Pl. iii. 1254 = A. Vulnerariae.
Halleri, Vill. Prosp. 42; Hist. Pl. Dauph. iii. 473 = A. australis.
helvetica, Vitm. Summa Pl. iv. 297 = A. australis.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 159 = A. adesmiaefolius.
Hookeriana, Torr. & Gray, Fl. N. Am. i. 693 = A. Hookerianus.
incana, Vahl, Symb. Bot. i. 57 = A. anthylloides.
inflata, Gill. in Hook. Bot. Misc. iii. (1833) 183.—Chili.
inflata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 343 = A. Menziesii.
lanata, Pall. Reise, iii. 746 = Oxytropis lanata.
Landbeckii, Phil. in Linnaea, xxxiii. (1864–65) 42.—Chili.
lapponica, Wahlenb. Veg. Helvet. 131, in obs. = A. arcticus.
Laxmanni, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = A. adsurgens?
leucophylla, Hook. & Arn. Bot. Beech. Voy. 333 = A. leucophyllus.
leucopsis, Torr. & Gray, Fl. N. Am. i. 694 = A. leucopsis.
longicaulis, Benth. ex Bunge, in Mém. Acad. Sc. Pétersb. Sér. VII. xv. (1869) n. 1. 28 = A. himalayanus.
longifolia, Nutt. in Torr. & Gray, Fl. N. Am. i. 346 = A. pictus, *A. Gray*.
lotiflora, Torr. & Gray, l. c. 349 = A. lotiflorus.
lutea, Ledeb. ex Sweet, Hort. Brit. ed. II. 145 (Quid?).—Sibir.
macrocarpa, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 36 = A. lonchocarpus.
macrocarpa, Phil. in Linnaea, xxviii. (1856) 619.—Chili.
macrodon, Hook. & Arn. Bot. Beech. Voy. 333 = A. lonchocarpus.
macrophysa, Phil. in Linnaea, xxviii. (1856) 619.—Chili.
macrostachys, Turcz. in Bull. Soc. Nat. Mosc. (1840) 66 = A. mongholicus.
magellensis, Huet, ex Bunge, in Mém. Acad. Sc. Pétersb. Sér. VII. xv. (1869) n. 1. 24 = A. australis.
megacarpa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 343 = A. megacarpus.
membranacea, Fisch. ex Link, Enum. Hort. Berol. ii. 252 = A. membranaceus.
microcarpa, Strach. & Winterb. ex Bunge, in Mém. Acad. Sc. Pétersb. Sér. VII. xv. (1869) n. 1. 22 = A. melanostachys.
microphylla, Pall. Reise, iii. 744 = A. polyphyllus.
minima, All. Fl. Pedem. i. 338 = A. alpinus.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 496 = A. Humboldtii.
mollissima, Nutt. ex Torr. & Gray, Fl. N. Am. i. 350 = A. Purshii.
montana, Crantz, Stirp. Austr. ed. II. fasc. v. 422 = Oxytropis montanus.
montana, Wahlenb. Fl. Lapp. 189 = Oxytropis montana, lapponica.
muricata, Pall. Reise, iii. 746 = Oxytropis muricata.
myriophylla, Pall. l. c. 745 = Oxytropis myriophylla.
neglecta, Torr. & Gray, Fl. N. Am. i. 344 = A. Cooperi.
nigrescens, Hook. Fl. Bor. Am. i. 143 = A. multiflorus.
nubigena, E. Mey. ex Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 16.—Chili.
Nuttallii, Torr. & Gray, Fl. N. Am. i. 343, 693 = A. Menziesii.
ochreata, Crantz, Stirp. Austr. ed. II. fasc. v. 419 = A. frigidus.
ochroleuca, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 186.—Chili.
oligantha, Phil. in Linnaea, xxxiii. (1864–65) 42.—Chili.
oreophila, Phil. l. c. xxviii. (1856) 681.—Chili.
oroboides, DC. Prod. ii. 274 = A. oroboides.
ovigera, Boiss. Diagn. Ser. I. vi. 35 = A. ovigerus.
oxyphylla, Pall. Reise, iii. 743 = Oxytropis oxyphylla.
parviflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 348 = A. oroboides.

PHACA :—
parviflora, Turcz. in Bull. Soc. Nat. Mosc. (1840) 66 = Astragalus secundus.
parvifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 348 = A. gracilis.
pauciflora, Nutt. l. c. = A. leptaleus.
pauciflora, Pers. Syn. ii. 331 = A. triflorus.
pectinata, Hook. Fl. Bor. Am. i. 141 = A. pectinatus, *Dougl.*
picta, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 36 = A. pictus.
pilosa, [Clairv.] Man. Herb. 133 = Oxytropis pilosa.
Pisissi, Phil. in Anal. Univ. Chil. (1862) 46.—Chili.
podocarpa, Hook. Fl. Bor. Am. i. 142 = A. sclerocarpus.
prostrata, Pall. Reise, iii. 744; Linn. f. Suppl. 336 = Oxytropis prostrata.
pulchella, Clos, in C. Gay, Fl. Chil. ii. 92.—Chili.
pygmaea, Nutt. ex Torr. & Gray, Fl. N. Am. i. 349 = A. Chamaeleuce.
quindecimjuga, Phil. in Linnaea, xxxiii. (1864–65) 41.—Chili.
Reedi, Phil. in Anal. Univ. Chil. (1870) ii. 168.—Chili.
reptans, Benth. ex Bunge, in Mém. Acad. Sc. Pétersb. Sér. VII. xv. (1869) n. 1. 28 = A. tenuicaulis.
Richii, A. Gray, Bot. U. St. Expl. Exped. i. 769 = A. Richii.
Robbinsii, Oakes, in Hovey, Mag. Hortic. vii. (1841) 181 = A. Robbinsii.
robusta, Phil. in Anal. Univ. Chil. (1872) 690.—Chili.
salsula, Pall. Reise, iii. 747 = Sphaerophysa salsula.
sericea, Nutt. ex Torr. & Gray, Fl. N. Am. i. 343 = A. sericoleucus.
sibirica, Linn. Sp. Pl. 755 = Oxytropis lanata.
simplicifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 350 = A. simplicifolius.
sordida, Fisch. ex Link, Enum. Hort. Berol. ii. 252 = Oxytropis campestris?
sordida, Wahlenb. Fl. Lapp. 190 = Oxytropis sordida.
striata, Clos, in C. Gay, Fl. Chil. ii. 92.—Chili.
subverticillaris, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 62 = A. subverticillaris (in Addend.).
sulcata, Linn. Syst. ed. X. 1173 = A. sulcatus.
sylvatica, Pall. Reise, iii. 744 = Oxytropis sylvatica.
tomentosa, Georgi, Bemerk. Reise Russ. Reich. i. 226 = Oxytropis lanata.
Tragacantha, All. Fl. Pedem. i. 338. = A. aristatus.
triangularis, Zea, in Hornem. Hort. Hafn. Suppl. 63 = A. triangularis.
trichopoda, Nutt. ex Torr. & Gray, Fl. N. Am. i. 343 = A. trichopodus.
tricolor, Clos, in C. Gay, Fl. Chil. ii. 105.—Chili.
triflora, DC. Astrag. 62. t. 1 = A. triflorus.
trifoliata, Linn. Mant. ii. 270.—China.
trigonocarpa, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 761 = A. trigonocarpus.
triphylla, Eaton & Wright, N. Am. Bot. ed. VIII. 351 = A. triphyllus.
umbellata, Nym. Consp. 196 = A. frigidus.
uralensis, Wahlenb. Fl. Carp. 223 = Oxytropis uralensis.
uspallatensis, Phil. in Anal. Univ. Chil. (1870) ii. 168.—Chili.
vaga, Clos, in C. Gay, Fl. Chil. ii. 99.—Chili.
velutina, Turcz. ex Bess. in Flora, xvii. (1834) 1. Beibl. 10, nomen.—As. centr.?
vesicaria, Linn. Mant. i. 103 = A. halicacabus.
vestita, Benth. Bot. Voy. Sulph. 13 = A. vestitus, *S. Wats.*
villosa, Nutt. Gen. Am. ii. 97 = A. villosus.
villosa, S. Wats. Bibliog. Ind. N. Am. Bot. 197 = A. mollissimus.
viscosa, [Clairv.] Man. Herb. 133 = Oxytropis foetida.
Vogelii, Webb, in Hook. Niger Fl. 123. t. 8 = A. prolixus.
zerdana, Boiss. Diagn. Ser. I. vi. 34 = A. zerdanus.

PHACELIA, Juss. Gen. 129 (1789). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 827.
Aldea, Ruiz & Pav. Prod. 19 (1794).
Cosmanthus, Nolte, ex A. DC. in DC. Prod. ix. 296 (1845).
Endiplus, Rafin. in Am. Monthly Mag. (1811) 356; (1819) 191.
Eutoca, R. Br. Frankl. Voy. App. 764. t. 27 (1823).

PHACELIA :—
Heteryta, Rafin. in Med. Repos. N. York, v. (1808) 352.
Microgenetes, A. DC. in DC. Prod. ix. 291 (1845).
Whitlavia, Harv. in Hook. Lond. Journ. Bot. v. (1846) 311.
affinis, *A. Gray, Syn. Fl. N. Am.* ii. 1. 417.—Calif.
Aldea, Loud. Hort. Brit. 59 = circinata.
arizonica, *A. Gray, Syn. Fl. N. Am.* ii. 1. 394.—Arizona.
artemisioides, *Griseb. in Goett. Abh.* xix. (1874) 230.—Reg. Argent.
bicolor, *Torr. ex S. Wats. in Bot. King's Exped.* 255.—Am. bor. occ.
bipinnatifida, Frank, ex A. DC. in DC. Prod. ix. 300 = Hydrophyllum macrophyllum.
bipinnatifida, *Michx. Fl. Bor. Am.* i. 134. t. 16.—Am. bor.
Bolanderi, *A. Gray, in Proc. Am. Acad.* x. (1875) 322.—Calif.
brachyantha, *Benth. in Trans. Linn. Soc.* xvii. (1837) 279.—Chili.
brachyloba, *A. Gray, in Proc. Am. Acad.* x. (1875) 324.—Calif.
Brannani, Kellogg, in Proc. Calif. Acad. vii. 1. (1877) 90 = Fremontii.
brevistylis, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 172 = bipinnatifida.
Breweri, *A. Gray, in Proc. Am. Acad.* x. (1875) 317.—Calif.
caerulea, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 122.—Am. bor. occ.
californica, Cham. in Linnaea, iv. (1829) 495 = circinata.
campanularia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 164.—Calif.
canescens, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 159 = circinata.
Cephalotes, *A. Gray, in Proc. Am. Acad.* x. (1875) 325.—Utah.
ciliata, *Benth. in Trans. Linn. Soc.* xvii. (1837) 280.—Am. bor. occ.
circinata, *Jacq. f. Eclog. Am.* i. 135. t. 91.—Am. bor. occ.; Chili.
circinatiformis, *A. Gray, in Proc. Am. Acad.* x. (1875) 325.—Calif.
clinopodioides, *Bert. in Merc. Chil.* (1829); *ex Am. Journ. Sc.* Ser. I. xxiii. (1833) 263.—Chili.
coerulea Auct. = caerulea.
conferta, G. Don, Gen. Syst. iv. 397 = congesta.
congesta, *Hook. Bot. Mag.* t. 3452.—Texas.
Cooperae, A. Gray, in Proc. Am. Acad. xv. (1880) 49 = gymnoclada?
crassifolia, Parry, in Am. Natural. ix. (1875) 16 = pulchella.
crassifolia, *Torr. ex S. Wats. in Bot. King's Exped.* 255.—Am. bor. occ.
crenulata, *Torr. ex S. Wats. l. c.* 251.—Am. bor. occ.
Cumingii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 169, in obs.—Chili.
curvipes, Parry, in Am. Natural. ix. (1875) 16 = P. Cephalotes.
curvipes, *Torr. ex S. Wats. in Bot. King's Exped.* 252.—Am. bor. occ.
Davidsonii, *A. Gray, in Proc. Am. Acad.* x. (1875) 324.—Calif.
demissa, *A. Gray, l. c.* 326.—N. Mexic.
distans, *Benth. Bot. Voy. Sulph.* 36.—Calif.
divaricata, *A. Gray, in Proc. Am. Acad.* x. (1875) 325.—Calif.
Douglasii, *Torr. in Bot. Mex. Bound.* 143.—Calif.
Endiplus, *Steud. Nom.* ed. II. ii. 313 (sp. dub.).—Am. bor.
fimbriata, *Michx. Fl. Bor. Am.* i. 134. t. 16.—Am. bor.
fimbriata, Pursh, Fl. Am. Sept. i. 140 = Purshii.
floribunda, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 200.—Calif.
Franklinii. *A. Gray, Man. Bot. N. U. St.* ed. II. 329.—Am. bor.
Fremontii, *Torr. in Ives, Rep. Colorado River Bot.* 21.—Am. bor. occ.
furcata, *Dougl. ex Hook. Fl. Bor. Am.* ii. 75.—Calif.
glabra, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 192.—Am. bor. occ.
glandulosa, Kellogg, in Proc. Calif. Acad. vii. (1877) 92 = Parryi?

PHACELIA:—
glandulosa, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 160.—Am. bor. occ.
grandiflora, *A. Gray, in Proc. Am. Acad.* x. (1875) 321.—Calif.
grisea, *A. Gray, l. c.* xii. (1877) 80.—Calif.
gymnoclada, *Torr. ex S. Wats. in Bot. King's Exped.* 255.—Calif.
hastata, Dougl. ex Lehm. Pugill. ii. 20 = circinata.
heterophylla, Pursh, Fl. Am. Sept. i. 140 = circinata.
hirsuta, Nutt. in Trans. Am. Phil. Soc. v. (1837) 191 = parviflora.
hispida, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 463 = patuliflora.
hispida, *A. Gray, Syn. Fl. N. Am.* ii. 1. 161.—Calif.
humilis, *Torr ex Gray, in Pacif. Rail. Rep.* ii. 122. t. 7.—Am. bor. occ.
hydrophylloides, *Torr. ex A. Gray, in Proc. Am. Acad.* vii. (1868) 400.—Calif.
infundibuliformis, *Torr. Bot. Mex. Bound.* 144.—N. Mexic.
integrifolia, *Torr. in Ann. Lyc. N. York*, ii. (1828) 222. t. 3.—Am. bor. occ.
invenusta, *A. Gray, in Proc. Am. Acad.* xx. (1885) 303.—Arizona.
Ivesiana, *Torr. in Ives, Rep. Colorado River Bot.* 21.—Am. bor. occ.
ixodes, *Kellogg, in Bull. Calif. Acad.* i. (1884) 6.—Calif.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 417.—Arizona.
leucophylla, Torr. in Frém. Rep. (1845) 93 = circinata.
loasaefolia, *Torr. Bot. Mex. Bound.* 144.—Calif.
longipes, *Torr. ex A. Gray, in Proc. Am. Acad.* x. (1875) 322.—Calif.
Lyoni, *A. Gray, l. c.* xx. (1885) 303.—Calif.
maculata, *Wood, Am. Bot. & Flor.* 255 (Quid?).—Am. bor.
malvaeflora, G. Don, Gen. Syst. iv. 396 = seq.
malvaefolia, *Cham. & Schlecht. in Linnaea*, iv. (1829) 494.—Am. bor. occ.
Menziesii, *Torr. ex S. Wats. in Bot. King's Exped.* 252.—Am. bor. occ.
micrantha, *Torr. Bot. Mex. Bound.* 144.—Am. bor. occ.
mohavensis, *A. Gray, Syn. Fl. N. Am.* ii. 1. 164.—Calif.
namatoides, *A. Gray, in Proc. Am. Acad.* x. (1875) 317.—Calif.
nana, *Wedd. Chlor. And.* ii. 86.—Bolivia.
neo-mexicana, *Thurb. ex Torr. Bot. Mex. Bound.* 143.—Am. bor. occ.
Orcuttiana, *A. Gray, in Proc. Am. Acad.* xix. (1883) 88.—Calif.
pachyphylla, *A. Gray, l. c.*—Calif.
Palmeri, *Torr. ex S. Wats. in Bot. King's Exped.* 251.—Calif.
Parishii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 88.—Calif.
Parryi, *Torr. Bot. Mex. Bound.* 143.—Calif.
parviflora, *Pursh, Fl. Am. Sept.* i. 140.—Am. bor.
patuliflora, *A. Gray, in Proc. Am. Acad.* x. (1875) 321.—Texas.
pedicellata, *A. Gray, Syn. Fl. N. Am.* ii. 1. 160.—Calif.
peruviana, Spreng. Syst. i. 584 = circinata.
phyllomanica, *A. Gray, in Proc. Am. Acad.* xi. (1876) 87.—Calif.
pimpinelloides, *A. Gray, l. c.* x. (1875) 321.—Mexic.
pinnatifida, *Griseb. ex Wedd. Chlor. And.* ii. 86.—Peruv.
platycarpa, *Spreng. Syst.* i. 584.—Mexic.
platyloba, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 223.—Calif.
Popei, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 172.—Am. bor. occ.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 223.—Calif.
procera, *A. Gray, l. c.* x. (1875) 323.—Am. bor. occ.
pubescens, Poir. Encyc. v. 239 = bipinnatifida.
pulchella, *A. Gray, in Proc. Am. Acad.* x. (1875) 326.—Utah.
Purshii, *Buckl in Am. Journ. Sc.* Ser. I. xlv. (1843) 171.—Am. bor.
pusilla, Buckl. l. c. 172 = parviflora.
pusilla, *Torr. ex S. Wats. in Bot. King's Exped.* 253.—Am. bor. occ.

PHACELIA:—
ramosissima, *Dougl. ex Lehm. Pugill.* ii. 21.—Am. bor. occ.
Rattani, *A. Gray, in Proc. Am. Acad.* xx. (1885) 302.—Am. bor. occ.
rotundifolia, *Torr. ex S. Wats. in Bot. King's Exped.* 253.—Am. bor. occ.
rudis, Dougl. ex A. DC. in DC. Prod. ix. 298 = circinata.
saxicola, *A. Gray, in Proc. Am. Acad.* xx. (1885) 304.—Arizona.
secunda, J. F. Gmel. Syst. 330 = circinata?
sericea, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 254.—Am. bor. occ.
simplex, Pers. Syn. i. 168 = bipinnatifida.
sinuata, *Phil. in Linnaea*, xxxiii. (1864–65) 185.—Chili.
strictiflora, *A. Gray, in Proc. Am. Acad.* x. (1875) 321.—Am. bor. occ.
suffrutescens, Parry, in Proc. Davenp. Acad. iv. (1884) 38 = ramosissima.
tanacetifolia, *Benth. in Trans. Linn. Soc.* xvii. (1837) 280.—Calif.
tanacetifolia, A. DC. in DC. Prod. ix. 299 = congesta.
tanacetifolia, Thurb. ex Torr. Bot. Mex. Bound. 143 = ramosissima.
tenuifolia, Harv. ex Torr. l. c. = tanacetifolia.
tripinnata, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 79 = tanacetifolia.
vinifolia, Paxt. Mag. Bot. iii. (1837) 121 = circinata?
viscida, *Torr. Bot. Mex. Bound.* 143.—Calif.
viscosa, *Phil. Fl. Atac.* 37.—Chili.
vitifolia, Steud. Nom. ed. II. ii. 313 = *vinifolia*.
Whitlavia, *A. Gray, in Proc. Am. Acad.* x. (1875) 321.—Calif.

PHACELLANTHUS, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 141 (Fam. Nat. Fl. Jap. ii. 17). *OROBANCHACEAE*, Benth. & Hook. f. ii. 983.
tubiflorus, *Sieb. & Zucc. l. c.*—Japon.

PHACELLANTHUS, Steud. in Zoll. Syst. Verz. Ind. Archip. 61 (1854) = **Gahnia**, Forst. (Cyper.).
multiflorus, Steud. l. c. = G. javanica.

PHACELLARIA, Benth. in Benth. & Hook. f. Gen. iii. 229 (1880). *SANTALACEAE*, Benth. & Hook. f. l. c.
compressa, *Benth. l. c.*—Burma.
rigidula, *Benth. l. c.*—Burma.

PHACELLARIA, Willd. ex Steud. Nom. ed. II. i. 353, in syn. (1840); ii. 313 (1841) = **Chloris**, Sw. (Gramin.).
panicea, Willd. ex Steud. ll. cc. = C. submutica.

PHACELLOTHRIX, F. Muell. Fragm. xi. 49 (1878). *COMPOSITAE*.
cladochaeta, *F. Muell. l. c.*—Austral.

PHACELURA, Benth. in Journ. Linn. Soc. xix. (1881) 68, sphalm. = seq.

PHACELURUS, Griseb. Spicil. Fl. Rumel. ii. 423 (1844) = **Rottboellia**, Linn. f. (Gramin.).
digitatus, Griseb. l. c. 424 = R. digitata.

PHACOCAPNOS, Bernh. in Linnaea, xii. (1838) 664 = **Corydalis**, DC. (Papav.).
Cracca, Bernh. l. c. = C. Cracca.
Dregeanus, Bernh. l. c. = C. Cracca.
pruinosus, Bernh. l. c. = C. pruinosa.

PHACOSPERMA, Haw. in Phil. Mag. i. (1827) 124 = **Calandrinia**, H. B. & K. (Portul.).
peruviana, Haw. l. c. = C. Phacosperma.

PHADROSANTHUS, Neck. Elem. iii. 153 (1790) = **Epidendrum**? Linn. (Orchid.).

PHAEANTHUS, Hook. f. & Thoms. Fl. Ind. i. 146 (1855). *ANONACEAE*, Benth. & Hook. f. i. 27.
crassipetalus, *Becc. in Nuov. Giorn. Bot.* iii. (1871) 191.—Borneo.
Cumingii, *Miq. Fl. Ind. Bat.* i. II. 51.—Ins. Philipp.
dioicus, Kurz, in Flora, liii. (1870) 274 = Miliusa Roxburghiana.

PHAEANTHUS:—
glabrescens, Baill. Hist. Pl. i. 246 = Piptostigma glabrescens.
heteropetalus, Baill. l. c. 245 = Heteropetalum brasiliense.
lanceolatus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 40.—Malaya.
malabaricus, *Bedd. Ic. Pl. Ind. Or.* i. 16. t. 76.—Ind. or.
malabaricus, Naves, in Blanco, Fl. Filip. ed. III. t. 193 = Polyalthia suberosa.
nutans, *Hook. f. & Thoms. Fl. Ind.* i. 147.—Malaya.
nutans, Scheff. in Ann. Jard. Buitenz. i. (1876) 5 = crassipetala.
pilosus, Baill. Hist. Pl. i. 246 = Piptostigma pilosum.
splendens, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 40.—Borneo.
sumatrana, *Miq. l. c.*—Sumatra.

PHAECASIUM, Cass. in Dict. Sc. Nat. xxxix. 387 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 515.
CYMBOSERIS, Boiss. Diagn. Ser. I. xi. 50 (1849).
IDIANTHES, Desv. Fl. Anj. 199 (1827).
SCLEROPHYLLUM, Gaud. Fl. Helv. v. 47 (1829).
lampsanoides, *Cass. in Dict. Sc. Nat.* xxxix. 387.—Europ.
pulchrum, Reichb. f. Ic. Fl. Germ. xix. 39 = lampsanoides.

PHAEDRA, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1850) = **Bernardia**, P. Br. (Euphorb.).

PHAEDRANASSA, Herb. in Bot. Reg. (1845) t. 17. Misc. 16. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 732.
Carmioli, *Baker, in Saund. Refug. Bot.* t. 46.—Costa Rica.
chloracra, *Herb. in Bot. Reg.* (1845) t. 17. *Misc.* 16.—Peruv.
eucrosioides, *Benth. & Hook. f. Gen.* iii. 733.—Ecuador.
Lehmanni, *Regel, Gartenfl.* (1883) 354.—N. Granat.
multiflora, Kunth, Enum. Pl. v. 502 = chloracra.
obtusa, Herb. in Bot. Reg. (1845) sub t. 17 = chloracra.
rubro-viridis, Baker, in Gard. Chron. (1875) II. 7 = Eustephia coccinea.
schizantha, *Baker, l. c.* (1880) II. 556.—Peruv.
viridiflora, *Regel, Gartenfl.* (1853) 357; *Baker, in Gard. Chron.* (1877) II. 134.—Peruv.

PHAEDRANTHUS, Miers, in Proc. Hort. Soc. iii. (1863) 182 = **Bignonia**, Tourn.
buccinatorum, Miers, l. c. = B. buccinatoria.
cinerascens, Miers, l. c. 183 = B. buccinatoria.
exsertus, Miers, l. c. = B. buccinatoria.
extatus, Miers, l. c. = B. buccinatoria?
Lindleyanus, Miers, l. c. 182 = B. buccinatoria.

PHAENICANTHUS, Thw. Enum. Pl. Zeyl. 242, in syn. (1861) = **Premna**, Linn. (Verben.).
zeylanicus, Thw. l. c. = Prem. purpurascens.

PHAENIX, Hill, Hort. Kew. 449 (1768) = **Phoenix**, Linn. (Palm.).

PHAENIXOPUS, Cass. in Dict. Sc. Nat. xxxix. 391 (1826) = **Lactuca**, Tourn. (Compos.).
acanthifolius, Cass. l. c. xlviii. 426 = L. acanthifolia.
chondrillaeflorus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 102 = L. chondrillaeflora.
decurrens, Cass. in Dict. Sc. Nat. xxxix. 391 = L. viminea.
intricatus, Pomel, Nouv. Mat. Fl. Atl. 264 = L. intricata.
intybaceus, Less. Syn. Comp. 137 = L. intybacea.
Kotschyi, Boiss. Fl. Orient. iii. 818 = L. brevirostris.
muralis, Koch, Syn. Fl. Germ. ed. I. 430 = L. muralis.
orientalis, Boiss. Voy. Bot. Espagne, ii. 390 = L. spinosa.
ramosissimus, Cass. in Dict. Sc. Nat. xxxix. 392; Reichb. Fl. Germ. Excurs. 272 = L. viminea.
spinosus, Cass. l. c. xlviii. 426 = L. spinosa.
triqueter, DC. Prod. vii. 176 = L. triquetra.
vimineus, Reichb. Fl. Germ. Excurs. 272 = L. viminea.

PHAEOCARPUS, Mart. Nov. Gen. et Sp. i. 61. t. 37, 38 (1824) = **Magonia**, A. St. Hil. (Sapind.).
agrestis, Mart. l. c. 63, in obs.—Bras.
campestris, Mart. l. c. 62 = M. pubescens.
glabratus, Spreng. Syst. iv. Cur. Post. 150 = M. glabrata.

PHAENOCODON, Salisb. Gen. Pl. Fragm. 58 (1866) = **Lapageria**, Ruiz & Pav. (Liliac.).

PHAENOCOMA, D. Don, in Mem. Wern. Soc. v. (1826) 554. *COMPOSITAE*, Benth. & Hook. f. ii. 313.
prolifera, *D. Don, l. c.*—Afr. austr.

PHAENOMERIA, Steud. Nom. ed. II. ii. 313 (1841) = PHAEOMERIA, Lindl. = **Amomum**, Linn. (Scitam.).

PHAENOPODA, Cass. in Dict. Sc. Nat. xlii. 84 (1826) = **Podotheca**, Cass. (Compos.).
angustifolia, Cass. l. c. 85 = Podotheca angustifolia.

PHAENOPUS, DC. Prod. vii. 176 (1838) = PHAENIXOPUS, Cass. = **Lactuca**, Tourn. (Compos.).

PHAENOPYRUM, M. Roem. Syn. Rosifl. 152 (1847) = **Crataegus**, Tourn. (Rosac.).
acerifolium, M. Roem. l. c. 157 = C. cordata.
arborescens, M. Roem. l. c. 153 = C. arborescens.
carolinianum, M. Roem. l. c. 152 = C. flava.
coccineum, M. Roem. l. c. 156 = C. coccinea.
corallinum, M. Roem. l. c. 154 = C. cordata.
cordatum, M. Roem. l. c. 157 = C. cordata.
ellipticum, M. Roem. l. c. 155 = C. flava.
hirsutum, M. Roem. l. c. 154 = C. mexicana.
melanocarpum, M. Roem. l. c. 155 = C. pentagyna.
mexicanum, M. Roem. l. c. 154 = C. mexicana.
nigrum, M. Roem. l. c. 157 = C. nigra.
odoratissimum, M. Roem. l. c. 158 = C. orientalis.
parvifolium, M. Roem. l. c. 152 = C. parvifolia.
pentagynum, M. Roem. l. c. 155 = C. pentagyna.
populifolium, M. Roem. l. c. 153 = C. coccinea.
pruinosum, M. Roem. l. c. 154 = C. glandulosa.
Schraderianum, M. Roem. l. c. 158 = C. sanguinea.
spathulatum, M. Roem. l. c. 155 = C. spathulata.
subvillosum, M. Roem. l. c. 154 = C. subvillosa.
tanacetifolium, M. Roem. l. c. 158 = C. tanacetifolia.
Tournefortii, M. Roem. l. c. = C. orientalis.
Trewianum, M. Roem. l. c. 154 = C. Crus-galli.
uniflorum, M. Roem. l. c. 153 = C. tomentosa.
virginicum, M. Roem. l. c. 155 = C. flava.
viride, M. Roem. l. c. 156 = C. Crus-galli.
Wendlandii, M. Roem. l. c. 154 = C. coccinea.

PHAENOPYRUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 166 (1842) = **Lagenocarpus**, Nees (Cyperac.).
paniculatum, Schrad. ex Nees, l. c. = L. guianensis.

PHAENOSPERMA, Munro, ex Benth. in Journ. Linn. Soc. xix. (1881) 59. *GRAMINEAE*, Benth. & Hook. f. iii. 1119.
globosa, *Munro, ex Benth. l. c.*—China.

PHAENOSTOMA, Steud. Nom. ed. II. i. 234, in syn. (1840), sphalm. = **Chaenostoma**, Benth. (Scrophul.).

PHAEOCLES, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Linn. (Liliac.).
maculata, Salisb. l. c. = O. maculatum.

PHAEOCORDYLIS, Griff. in Proc. Linn. Soc. i. (1844) 219; et in Trans. Linn. Soc. xx. (1846) 100. t. 8 = **Rhopalocnemis**, Jungh. (Balanoph.).
areolatus, Griff. in Proc. Linn. Soc. i. (1844) 220 = R. phalloides.

PHAEOMERIA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Amomum**, Linn. (Scitam.).
magnifica, Lindl. l. c. = A. magnificum.

PHAEOPAPPUS, Boiss. Diagn. Ser. I. vi. 122 (1845) = **Centaurea**, Linn. (Compos.).
amblyolepis, Boiss. Fl. Orient. iii. 601 = C. amblyolepis.
araneosus, Boiss. l. c. 598 = C. araneosa.

PHAEOPAPPUS :—
aristatus, Jaub. & Spach, Illustr. iii. 18. t. 213 = Centaurea aristata.
armenus, Jaub. & Spach, l. c. 13. t. 210 = C. armena.
cappadocicus, Boiss. Fl. Orient. iii. 594 = C. cappadocica.
carthamoides, Boiss. Diagn. Ser. I. vi. 125 = C. carthamoides.
cataonicus, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 596 = C. cataonica?
cheirolophus, Boiss. l. c. 595 = C. macrolopha.
chrysocollus, Nym. Consp. 418 = C. trinervia.
cichoriifolius, Jaub. & Spach, Illustr. iii. 14.—As. Min.
declinatus, Boiss. Fl. Orient. iii. 595.—Lycia.
decurrens, Boiss. l. c. 600 = C. decurrens.
dichrous, Boiss. l. c. 601 = C. dichroa.
drabifolius, Boiss. l. c. 594 = C. drabifolia.
floccosus, Boiss. l. c. 595 = C. macrolopha.
gymnocladus, Jaub. & Spach, Illustr. iii. 15. t. 211 = C. gymnocladus.
Haussknechtii, Boiss. Fl. Orient. iii. 599.—Mesopot.
helenioides, Boiss. l. c. 633 = C. helenioides.
inuloides, Nym. Consp. 418 = C. trinervia.
Kotschyi, Boiss. Fl. Orient. iii. 595 = C. macrolopha.
leuzeoides, Boiss. Diagn. Ser. I. vi. 124 = C. leuzeoides.
libanoticus, Boiss. Fl. Orient. iii. 594.—Syria.
macracanthus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 129 = C. leuzeoides.
macrocephalus, Boiss. Diagn. Ser. I. vi. 124 = C. macrocephala.
microcephalus, Jaub. & Spach, Illustr. iii. 16. t. 212 = C. microcephala.
microlonchoides, Boiss. Fl. Orient. iii. 602 = C. microlonchoides.
microlophus, Boiss. in Tchihat. Asie Min. Bot. ii. 308 = C. armena.
obtusifolius, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 599.—Mesopot.
ossicus, C. Koch, in Linnaea, xxiv. (1851) 436 = C. ossica.
rupestris, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 598.—Cataonia.
Ruprechtii, Boiss. l. c. 601.—Reg. Caucas.
salignus, Boiss. l. c. 596.—As. Min.
saxatilis, Boiss. l. c. 602 = C. saxatilis.
scoparius, Boiss. l. c. = C. scoparia.
spectabilis, Boiss. Diagn. Ser. I. vi. 124.—Armenia.
Steveni, C. Koch, in Linnaea, xxiv. (1851) 437 = C. Steveni.
Szowitsii, Boiss. Diagn. Ser. I. vi. 123 = C. leuzeoides.
trinervius, Boiss. Fl. Orient. iii. 600 = C. trinervia.

PHAEOPSIS, Nutt. ex Benth. in DC. Prod. xii. 555 (1848) = **Stenogyne**, Benth. (Labiat.).
montana, Nutt. ex Benth. l. c. 556 = S. Nelsonii.

PHAEOPTILUM, Radlk. in Bremen, Abh. viii. (1883) 435. *NYCTAGINEAE*.
spinosum, *Radlk. l. c.* 436.—Afr. austr.

PHAEOSPHAERIONA, Hassk. in Flora, xlix. (1866) 212 = **Athyrocarpus**, Schlecht. (Commel.).
efoveolatum, C. B. Clarke, in DC. Monog. Phan. iii. 136.—Venezuela.
leiocarpum, Hassk. in Flora, xlix. (1866) 212 = A. leiocarpus.
Mathewsii, C. B. Clarke, in DC. Monog. Phan. iii. 138.—Peruv.
persicariaefolium, C. B. Clarke, l. c. 137 = A. persicariaefolius.

PHAEOSTEMMA, Fourn. in Mart. Fl. Bras. vi. iv. 311 (1885). *ASCLEPIADEAE*.
Glaziovi, *Fourn. l. c.* 312.—Bras.
Riedelii, *Fourn. l. c.* 311.—Bras.

PHAEOSTOMA, Spach, Hist. Vég. Phan. iv. 392 (1835) = **Clarkia**, Pursh (Onagr.).
Douglasii, Spach, l. c. 393 = C. elegans.
elegans, Lilja, Fl. Sverig. Suppl. 25 = C. elegans.

PHAETHUSA, Gaertn. Fruct. ii. 425. t. 169. f. 3 (1791) = **Verbesina**, Linn. (Compos.).
americana, Gaertn. l. c. = V. occidentalis.
australis, Spreng. Syst. iii. 591 = Monactis flaverioides.
borealis, Spreng. l. c. = V. occidentalis.

PHAETUSA, Schreb. Gen. ii. 571 (1791) = praec.

PHAGNALON, Cass. in Bull. Soc. Philom. (1819) 174. *COMPOSITAE*, Benth. & Hook. f. ii. 304.
GNAPHALON, Lowe, Man. Fl. Mad. i. 438 (1868).
abyssinicum, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 26, nomen.—Abyss.
acuminatum, Aitch. in Journ. Linn. Soc. xix. (1882) 169 = niveum.
acuminatum, *Boiss. Fl. Orient.* iii. 222.—Beluchist.
affine, Boiss. Diagn. Ser. I. vi. 34 = persicum.
arabicum, Boiss. l. c. = Iphiona arabica.
atlanticum, *Ball, in Journ. Bot.* xi. (1873) 364.—Marocc.
Bennetti, Lowe, ex DC. Prod. v. 397 = rupestre.
bicolor, *Ball, in Journ. Bot.* xi. (1873) 364.—Marocc.
Borianianum, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 286, nomen.—Afr. trop.
Borianum, Oliver, Fl. Trop. Afr. iii. 338 = praec.
calycinum, DC. Prod. v. 397 = rupestre.
denticulatum, Decne. ex C. B. Clarke, Comp. Ind. 113 = niveum.
geminiflorum, Nym. Consp. 385 = rupestre.
graecum, *Boiss. & Heldr. Diagn.* Ser. I. xi. 6.—Graec.; As. Min.
hypoleucum, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 396 = abyssinicum.
intermedium, DC. ex Boiss. & Heldr. Diagn. Ser. I. xi. 7, in obs. = graecum.
Kotschyi, *Sch. Bip. ex Boiss. Fl. Orient.* iii. 221.—Syria; Assyria.
Lagascae, Cass. in Bull. Soc. Philom. (1819) 174 = saxatile.
lepidotum, *Pomel, Nouv. Mat. Fl. Atl.* 34.—Alger.
Lowei, DC. ex Lowe, Man. Fl. Mad. i. 443 = rupestre.
lucidum, *Webb, in Hook. Niger Fl.* 136.—Ins. Cap. Virid.
melanoleucum, *Webb, l. c.* 135. t. 9.—Ins. Cap. Virid.
nitidum, *Fresen. in Mus. Senckenb.* i. (1833) 81. t. 4. f. 2.—Arab.; Persia; Afghan.
niveum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 68.—Reg. Himal.
persicum, *Boiss. Diagn.* Ser. I. vi. 74.—Persia.
pumilum, *DC. Prod.* v. 397.—Ins. Creta.
purpurascens, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 212. t. 82.—Ins. Canar.
Quartinianum, *A. Rich. Tent. Fl. Abyss.* i. 396.—Abyss.
rupestre, *DC. Prod.* v. 396.—Reg. Mediterr.; Oriens.
saxatile, *Cass. in Bull. Soc. Philom.* (1819) 174.—Reg. Mediterr.
scariosum, *DC. ex Lowe, Man. Fl. Mad.* i. 443.—Ins. Mader.
Schweinfurthii, *Sch. Bip. ex Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 685.—Abyss.
scoparium, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 153, nomen.—Afr. trop.
sordidum, Reichb. Fl. Germ. Excurs. 224 DC. Prod. v. 396 = tricephalum.
spathulatum, Cass. in Dict. Sc. Nat. xxxix. 401 = rupestre.
subdentatum, Cass. l. c. 400 = saxatile.
telonense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 61 = tricephalum.
Tenorii, Presl, Fl. Sic. i. 29 = rupestre.
tricephalum, *Cass. in Dict. Sc. Nat.* xxxix. 401.—Reg. Mediterr.
umbelliforme, *DC. Prod.* v. 396.—Ins. Canar.
viride, *Uecht. in Oestr. Bot. Zeitschr.* xxiv. (1874) 134.—Hispan.
viridifolium, *Decne. ex Boiss. Fl. Orient.* iii. 220.—Arabia.

PHAIANTHES, Rafin. Fl. Tellur. iv. 30 (1836) = **Moraea**, Mill. (Irid.).
lurida, Rafin. l. c. 31 = M. lurida.

PHAIOPHLEPS, Rafin. Fl. Tellur. iv. 29 (1836) = **Solenomelus**, Miers (Irid.).
odoratissima, Rafin. l. c. = S. biflorus.

PHAIOSPERMA, Rafin. New Fl. Am. iv. 32 (1836) = **Polytaenia**, DC. (Umbell.).
pulverulenta, Rafin. l. c. = Polyt. Nuttallii.
trifida, Rafin. l. c.—Polyt. Nuttallii.

PHAIUS, Lour. Fl. Cochinch. 529 (1790). *ORCHIDEAE*, Benth. & Hook. f. iii. 512.
CYANORCHIS, Thou. ex Steud. Nom. ed. II. i. 457 (1840).
HECABE, Rafin. Fl. Tellur. iv. 44 (1836).
LIMATODIS, Blume, Bijdr. 375 (1825).
PACHYNE, Salisb. in Trans. Hort. Soc. i. (1812) 299.
PESOMERIA, Lindl. Bot. Reg. (1838) Misc. 5.
TANKERVILLIA, Link, Handb. i. 251 (1829).
THUNIA, Reichb. f. in Bot. Zeit. x. (1852) 764.
albus, *Lindl. in Wall. Cat.* n. 3749.—Reg. Himal.
amboinensis, *Blume, Mus. Bot. Lugd. Bat.* ii. 180.—Ins. Amboin.
Augustinianus, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiv. (1856) 9 = cupreus.
australis, F. Muell. Fragm. i. 42 = grandifolius.
Bensoniae, Benth. in Journ. Linn. Soc. xviii. (1881) 305; Hemsl. in Gard. Chron. (1882) II. 565 = albus.
Bernaysii, *F. Muell. ex Bernays, in Gard. Chron.* (1873) 361, 1244.—Austral.
bicolor, Lindl. Gen. et Sp. Orch. 128 = Wallichii.
Blumei, Lindl. l. c. 127 = Wallichii.
bracteosus, *Reichb. f. in Bonplandia*, v. (1857) 42.—Java.
callosus, *Lindl. Gen. et Sp. Orch.* 128.—Malaya.
Carroni, F. Muell. Pl. Burdek. Exped. 19 = Wallichii.
crispus, *Blume, Mus. Bot. Lugd. Bat.* ii. 180.—Java.
cupreus, *Reichb. f. in Bonplandia*, iii. (1855) 226.—Java
Dodgsonii, *Dean, in Flor. Mag.* N. S. (1878) t. 329.—Ind. or.
fimbriatus, *Hort.—Cf. Gard. Chron.* (1876) I. 735.—Hab.?
flavus, *Lindl. Gen. et Sp. Orch.* 128.—Java.
flexuosus, *Blume, Mus. Bot. Lugd. Bat.* ii. 179.—Java.
formosus, Hort. ex Hemsl. in Gard. Chron. (1882) II. 565 (Quid?).—Hab.?
giganteus, Hort. ex Hemsl. l. c. = grandifolius.
Graeffei, *Reichb. f. in Seem. Fl. Vit.* 299.—Ins. Samoa.
grandiflorus, Reichb. f. in Walp. Ann. vi. 459 = seq.
grandifolius, Lindl. in Wall. Cat. n. 3747 = Wallichii.
grandifolius, *Lour. Fl. Cochinch.* 529.—China; Austral.
gratus, *Blume, Mus. Bot. Lugd. Bat.* ii. 181.—Ins. Amboin.
Hookerianus, *Reichb. f. in Bonplandia*, iv. (1856) 328.—Am. austr.?
Humblotii, *Reichb. f. in Gard. Chron.* (1880) II. 812.—Madag.
indigoferus, *Hassk. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 140; *Reichb. f. Xenia Orch.* i. 202.—Java.
inquilinus ×, Reichb. f. in Gard. Chron. (1867) 544.
irroratus ×, Reichb. f. l. c. 264.
Kuhlii, *Reichhb. f. Xenia Orch.* ii. 81. t. 122.—Java.
leucophaeus, F. Muell. Fragm. iv. 163 = grandifolius.
luridus, Hort.—Cf. Gard. Chron. (1882) II. 566 (Quid?).—Hab.?
luridus, *Thw. Enum. Pl. Zeyl.* 300.—Zeylan.
maculatus, *Lindl. in Wall. Cat.* n. 3748; *et in Bot. Mag.* t. 3960.—Reg. Himal.; China.
Mannii, Reichb. f. Otia Bot. Hamb. 118 = Wallichii.
Marshalliae, Hort.—Cf. Gard. Chron. (1871) 807; Veitch, Cat. (1880) 68; ex Hemsl. in Gard. Chron. (1882) II. 566 = albus.
minor, *Blume, Mus. Bot. Lugd. Bat.* ii. 181.—Japon.
mishmensis, *Reichb. f. in Bonplandia*, v. (1857) 43.—Reg. Himal.
niveus, Hort. ex Hemsl. in Gard. Chron. (1882) II. 566 = albus.
pauciflorus, *Blume, Mus. Bot. Lugd. Bat.* ii. 181.—Java.
platychilus, *Reichb. f. in Bonplandia*, v. (1857) 42.—Java.
pulchellus, *Kraentz. in Bremen, Abh.* vii. (1882) 254.—Madag.
Robertsii, *F. Muell. in Wing's South. Sc. Record*, iii (18) 263.—N. Caled.
rosellus, *Lem. in Fl. des Serres*, iii. (1847) t. 301.—Guian.
Rumphii, Blume, Mus. Bot. Lugd. Bat. ii. 179 = callosus.
stuppeus, *Blume, Orch. Archip. Ind.* 14.—Ins. Borbon.
Tankervillii, Blume, Mus. Bot. Lugd. Bat. ii. 177 = Wallichii.

PHAIUS:—
tenuis, *Reichb. f. in Bonplandia*, v. (1857) 42.—Java.
tetragonus, *Reichb. f. l. c.* iii. (1855) 221.—Ins. Mascar.
tuberculatus, Blume, Orch. Archip. Ind. t. 11 = seq.
tuberculosus, *Blume, Mus. Bot. Lugd. Bat.* ii. 181.—Madag.
veratrifolius, *Lindl. in Wall. Cat.* n. 3746.—Reg. Himal.
vestitus, Reichb. f. in Gard. Chron. (1867) 264, in obs. = Calanthe vestita.
villosus, *Blume, Mus. Bot. Lugd. Bat.* ii. 182.—Ins. Maurit.
Wallichii, *Lindl. in Wall. Pl. As. Rar.* ii. 46. t. 158.—Ind. or.; Burma; Malaya.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 42.—Java.

PHAJUS, Hassk. Cat. Hort. Bog. Alt. 42 (1844) = praec.

PHAKELLANTHUS, Steud. Syn. Pl. Cyp. 153 (1855) = PHACELLANTHUS, Sieb. & Zucc. = **Gahnia**, Forst. (Cyperac.).

PHALACRAEA, DC. Prod. v. 105 (1836) = **Piqueria**, Cav. (Compos.).
coelestina, Regel, Gartenfl. (1854) 388. t. 107 = Ageratum conyzoides.
latifolia, DC. in Deless. Ic. Sel. iv. t. 8 = Ageratum conyzoides.
Lindenii, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 240 = Alomia ageratoides.
Wendlandii, Sch. Bip. ex Klatt, in Leopoldina, xx. (1884) 74 = Piq. latifolia.

PHALACROCARPUM, Willk. in Bot. Zeit. xxii. (1864) 252 = **Chrysanthemum**, Tourn. (Compos.).
oppositifolium, Willk. l. c. = C. oppositifolium.

PHALACRODERIS, DC. Prod. vii. 97 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 511.
coa, *DC. l. c.*—Ins. Cos.

PHALACRODISCUS, Less. Syn. Comp. 253 (1832) = **Chrysanthemum**, Tourn. (Compos.).
arvensis, Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop. iii. 44 = C. tingitanum?
cebennensis, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 245 = C. monspeliense.
ceratophylloides, Less. Syn. Comp. 254 = C. monspeliense.
corsicus, Less. l. c. = C. coronopifolium.
Froelichii, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 245 (Quid?).—Europ.?
graminifolius, Less. Syn. Comp. 253 = C. graminifolium.
lanceolatus, Less. l. c. = C. lanceolatum.
latifolius, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 245 = C. lacustre.
maximus, Less. Syn. Comp. 253 = C. maximum.
monspeliensis, Sch. Bip. Ueb. Tanacet. 44 = C. monspeliense.
montanus, Less. Syn. Comp. 254 = C. montanum.
pallens, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 245, nomen = C. pallens.
pyrethroides, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 354 = Anthemis chia.

PHALACROGLOSSUM, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 (1835–60) = **Chrysanthemum**, Tourn. (Compos.).

PHALACROLOMA, Cass. in Dict. Sc. Nat. xxxix. 404 (1826) = **Erigeron**, Linn. (Compos.).
acutifolium, Cass. l. c. 405 = E. annuus.
annuum, Dum. Fl. Belg. 67 = E. annuus.
Beyrichii, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 63 = E. strigosus.
obtusifolium, Cass. in Dict. Sc. Nat. xxxix. 404 = E. strigosus.

PHALACROMESUS, Cass. in Dict. Sc. Nat. liii. 235 (1828) = **Tessaria**, Ruiz & Pav. (Compos.).
riparia, Cass. l. c. = T. integrifolia.

PHALACROS, Wenzig, in Linnaea, xxxviii. (1874) 164 = **Crataegus**, Tourn. (Rosac.).
cordatus, Wenzig, l. c. = C. cordata.

PHALACROSERIS, A. Gray, in Proc. Am. Acad. vii. (1868) 364. *COMPOSITAE*, Benth. & Hook. f. ii. 507.
Bolanderi, *A. Gray, l. c.*—Calif.

PHALAENOPSIS, Blume, Bijdr. 294 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 573.
POLYCHILOS, Breda, Kuhl & Hass. Orch., sine num. (1827).
POLYSTYLUS, Hasselt, ex Hassk. Retzia, 3 (1855).
STAURITIS, Reichb. f. in Hamb. Gartenz. xviii. (1862) 34.
STAUROGLOTIS, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 432.
SYNADENA, Rafin. Fl. Tellur. iv. 9 (1836).
amabilis, *Blume, Bijdr.* 294. t. 44.—Malaya.
amabilis, Lindl. Bot. Reg. (1838) t. 34 = P. Aphrodite.
ambigua, Reichb. f. in Hamb. Gartenz. xviii. (1862) 35 = P. Aphrodite.
amethystina, *Reichb. f. in Gard. Chron.* (1865) 602.—Malaya.
antennifera, Reichb. f. l. c. (1879) I. 398 = P. Esmeralda.
Aphrodite, *Reichb. f. in Hamb. Gartenz.* xviii. (1862) 35.—Ins. Philipp.
bella, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1862) 321 = P. Hebe.
Boxallii, *Reichb. f. in Gard. Chron.* (1883) I. 274.—Ins. Philipp.
Brymeriana, Reichb. f. l. c. (1876) I. 366 = *intermedia*.
casta ×, Reichb. f. l. c. (1875) I. 590.—Ins. Philipp.
Corningiana, *Reichb. f. l. c.* (1879) I. 620.—Hab.?
Cornu-cervi, *Blume & Reichb. f. in Hamb. Gartenz.* xvi. (1860) 116.—As. trop.
Cumingiana, Reichb. f. in Gard. Chron. (1881) I. 562, sphalm. = Corningiana.
delicata ×, Reichb. f. l. c. (1882) I. 700.
deliciosa, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Java.
De-Vriesiana, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 116.—Java.
equestris, Reichb. f. in Linnaea, xxii. (1849) 864 = rosea.
Esmeralda, *Reichb. f. in Gard. Chron.* (1874) II. 582.—Cochinch.
fasciata, *Reichb. f. l. c.* (1882) II. 134.—Ins. Philipp.
fuscata, *Reichb. f. l. c.* (1874) II. 6.—Malaya.
grandiflora, Lindl. l. c. (1848) 39 = amabilis.
Hebe, *Reichb. f. in Hamb. Gartenz.* xviii. (1862) 35.—Java.
intermedia ×, Lindl. in Lindl. & Paxt. Flow. Gard. iii. (1852–53) 162, 163. f. 310.—Ins. Philipp.
leucorrhoda ×, Reichb. f. in Gard. Chron. (1875) I. 301, 366.—Ins. Philipp.
Lowii, *Reichb. f. in Bot. Zeit.* xx. (1862) 214.—Burma.
Lueddemanniana, *Reichb. f. l. c.* xxiii. (1865) 146.—Ins. Philipp.
Lueddemannii, Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 243 = praec.
maculata, *Reichb. f. in Gard. Chron.* (1881) II. 134.—Borneo.
Mannii, *Reichb. f. l. c.* (1871) 902.—Assam.
Mariae, *Burbidge, in Warn. Orch. Album*, ii. t. 80.—Ins. Philipp.
pallens, *Reichb. f. in Walp. Ann.* vi. 932.—Ins. Philipp.
pantherina, *Reichb. f. in Bot. Zeit.* xxii. (1864) 298.—Borneo.
Parishii, *Reichb. f. l. c.* xxiii. (1865) 146.—Burma.
Porteri, Hort.—Cf. Gard. Chron. (1876) I. 369 = *intermedia*.
Reichenbachiana, *Reichb. f. & Sander, in Gard. Chron.* (1882) II. 586.—Ins. Philipp.
rosea, *Lindl. in Gard. Chron.* (1848) 671.—Ins. Philipp.
Ruckeri, Hort.—Cf. l. c. (1881) I. 562 = Sarcochilus unguiculatus.
Sanderiana, *Reichb. f. l. c.* (1883) I. 656.—Ins. Philipp.
Schilleriana, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 115.—Ins. Philipp.
speciosa, *Reichb. f. in Gard. Chron.* (1881) I. 562.—Ins. Andaman.
Stobartiana, *Reichb. f. l. c.* (1877) II. 392.—Hab.?
Stuartiana, *Reichb. f. l. c.* (1881) I. 748.—Ins. Philipp.
sumatrana, *Korth. & Reichb. f. in Hamb. Gartenz.* xvi. (1860) 115.—Sumatra.

PHALAENOPSIS:—
tetraspsis, *Reichb. f. Xenia Orch.* ii. 146.—Ins. Andaman.
Valentini, *Reichb. f. in Gard. Chron.* (1883) II. 262.—Malaya.
Veitchiana, *Reichb. f. l. c.* (1872) 935.—Ins. Philipp.
violacea, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 320.—Sumatra.
Wightii, Reichb. f. in Bot. Zeit. xx. (1862) 214 = Doritis Wightii.
zebrina, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1862) 319 = sumatrana.

PHALANGION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = PHALANGIUM, Tourn.

PHALANGIUM, Burm. f. Prod. Cap. 3 (1768) = **Melasphaerula**, Ker-Gawl. (Irideae).
obsoletum, Burm. f. l. c. (Quid ?).—Afr. austr.
ramosum, Burm. f. l. c. = M. graminea.
spicatum, Burm. f. l. c. (Quid ?).—Afr. austr.

PHALANGIUM, [Tourn.] Adans. Fam. ii. 49 (1763); Juss. Gen. 52 (1789) = **Anthericum**, Linn. (Liliac.).
abyssinicum, Kunth, Enum. Pl. iv. 598 = Chlorophytum tetraphyllum.
acuminatum, Dulac, Fl. Hautes-Pyr. 114 = A. Liliago.
Adenanthera, Poir. Encyc. v. 252 = Dianella nemorosa.
alatum, Buch.-Ham. ex Wall. Cat. n. 5056 = Chlorophytum arundinaceum.
albucoides, Poir. Encyc. v. 249 = Ornithogalum suaveolens.
algeriense, Boiss. & Reut. Pugill. Pl. Nov. 115 = A. Liliago.
angustifolium, Schweinf. Beitr. Fl. Aethiop. 294 = A. angustifolium.
attenuatum, Wight, Ic. t. 2037 = Chlorophytum attenuatum.
baeticum, Boiss. Elench. 84 = A. Liliago.
bicolor, [DC. in] Red. Lil. t. 215 = Simethis bicolor.
bipedunculatum, Jacq. ex Poir. Encyc. v. 244 = A. triflorum.
caeruleum, Pers. Syn. i. 368 = Pasithea coerulea.
canaliculatum, Poir. Encyc. v. 249 = A. canaliculatum.
capillare, Poir. l. c. 247 = Bulbinella triquetra.
ciliatum, H. B. & K. Nov. Gen. et Sp. i. 276. t. 676 = A. Humboldtii.
coarctatum, Pers. Syn. i. 368 = Excremis ramosa.
coeruleum, Poir. Encyc. Suppl. iv. 381 = Pasithea coerulea.
comosum, Poir. Encyc. v. 252 = Chlorophytum Sternbergianum.
contortum, Pers. Syn. i. 368 = A. contortum.
croceum, Michx. Fl. Bor. Am. i. 196 = Schoenolirion croceum.
croceum, Nutt. Gen. Am. i. 220 = A. Nuttallianum.
eccremorrhizum, Poir. Encyc. Suppl. iv. 380 = A. eccremorrhizum.
elatum, Poir. Encyc. v. 248; [DC. in] Red. Lil. t. 191 = Chlorophytum elatum.
elongatum, Pers. Syn. i. 367 = A. Jacquinianum.
esculentum, Nutt. Gen. Am. i. 219, partim = Camassia esculenta.
exuviatum, Poir. Encyc. v. 243 = Urginea exuviata.
falcatum, Wall. Cat. n. 5057 = Chlorophytum laxum.
fasciculatum, Baker, in Journ. Linn. Soc. xv. (1876) 331 = Chlorophytum elatum.
fastigiatum, Poir. Encyc. v. 246 = Chlorophytum elatum.
filifolium, Poir. l. c. 542 = Urginea nematodes.
filiforme, Poir. l. c. 247 = Bulbinella triquetra.
flavescens, Kunth, Enum. Pl. iv. 596 = A. flavescens.
flexifolium, Poir. Encyc. v. 243 = A. Jacquinianum.
floribundum, Pers. Syn. i. 368 = Bulbinella latifolia.
fragrans, Poir. Encyc. v. 247 = A. glaucum.
glaucum, Poir. l. c. Suppl. iv. 384 = A. glaucum.
graecum, Poir. Encyc. v. 250 = Lloydia alpina.
graminifolium, Pers. Syn. i. 368 = A. undulatum.
hirsutum, Pers. l. c. = A. hirsutum.
holosericeum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 203 = Simethis bicolor.
humile, Schweinf. Beitr. Fl. Aethiop. 294 = A. humile.
indicum, Kunth, Enum. Pl. iv. 598 = Chlorophytum attenuatum.
japonicum, Poir. Encyc. v. 248 = Anthericum japonicum.
latifolium, H. B. & K. Nov. Gen. i. 276 = A. glaucum

PHALANGIUM:—
laxum, F. Muell. Fragm. vii. 71 = Chlorophytum laxum.
lilaceum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = A. Liliago.
liliaginoides, Schlecht. in Linnaea, xxiv. (1851) 187 = A. Liliago ?
Liliago, Schreb. Spicil. 36 = A. Liliago.
Liliastrum, Brot. Fl. Lusit. i. 534 = Paradisia Liliastrum.
lineare, Hort.—Cf. Rev. Hortic. (1879) 320.—Hab. ?
longifolium, R. Grah. in Edinb. N. Phil. Journ. (Oct.-Dec. 1830) 171.—Peruv.
longifolium, Poir. Encyc. v. 243 = A. longifolium.
luteum, Nutt. Gen. Am. i. 220 = Schoenolirion croceum ?
marginatum, Poir. Encyc. v. 250 = Urginea physodes.
milleflorum, Steud. Nom. ed. II. i. 141; ii. 314 = Arthropodium paniculatum.
nepalense, Lindl. in Trans. Hort. Soc. vi. (1826) 277 = Chlorophytum nepalense.
niveum, Poir. Encyc. v. 248 = Chlorophytum tuberosum.
oligospermum, Wight, Ic. t. 2038 = Chlorophytum Orchidastrum.
ornithogaloides, Schweinf. Beitr. Fl. Aethiop. 294 = Chlorophytum tuberosum.
ossifragum, Muhl. ex Steud. Nom. ed. II. ii. 184, 314 = Narthecium ossifragum.
paniculatum, [Delar. in] Red. Lil. t. 360; Poir. Encyc. Suppl. iv. 383 = Arthropodium paniculatum.
parviflorum, Wight, Ic. t. 2039 = Chlorophytum laxum.
pauciflorum, Poir. Encyc. v. 251 = Urginea pusilla.
pedunculatum, Baker, in Journ. Linn. Soc. xv. (1876) 315 (*bipedunculatum*) = A. triflorum.
pendulum, [Delar. in] Red. Lil. t. 360; Poir. Encyc. Suppl. iv. 380 = Arthropodium paniculatum.
pilosum, Poir. Encyc. v. 244 = Ornithogalum hispidum.
planifolium, Pers. Syn. i. 367 = Simethis bicolor.
pomeridianum, Sweet, Hort. Brit. ed. I. 416 = Chlorogalum pomeridianum.
Quamash, Pursh, Fl. Am. Sept. i. 226 = Camassia esculenta.
ramosissimum, Presl, Rel. Haenk. ii. 127 = Echeandia terniflora.
ramosum, Houtt. Handleid. xii. 114; Poir. Encyc. v. 250 = A. ramosum.
reflexum, Poir. Encyc. v. 249 = Echeandia terniflora.
Renarnii, Booth, ex Schlecht. in Linnaea, xxiv. (1851) 187 = A. Liliago ?
revolutum, Pers. Syn. i. 367 = Scilla revoluta.
revolutum, Poir. Encyc. v. 248; Pers. l. c. 368 = A. revolutum.
scabrellum, Benth. Pl. Hartw. 293 = A. scabrellum.
scilloides, Poir. Encyc. v. 251 = A. scilloides.
serotinum, Poir. l. c. 241 = Lloydia alpina.
spicatum, Houtt. Handleid. xii. 115. t. 80. f. 2 = Micranthus alopecuroideus.
spicatum, Poir. Encyc. v. 247 = Scilla revoluta.
spirale, Poir. l. c. 249 = Eriospermum spirale.
squameum, Poir. l. c. 246 = A. hispidum.
sulphureum, Poir. Encyc. Suppl. iv. 381 = Ornithogalum pyrenaicum.
triflorum, Pers. Syn. i. 368 = A. triflorum.
tuberosum, Dalz. & Gibs. Bomb. Fl. 251 = Chlorophytum malabaricum.
tuberosum, Kunth, Enum. Pl. iv. 598 = Chlorophytum tuberosum.
undulatum, Poir. Encyc. v. 242 = A. undulatum.
variegatum, Hort. ex Baker, in Gard. Chron. (1882) II. 108 = Chlorophytum elatum.
vespertinum, Poir. l. c. 249 = A. falcatum.
villosum, Poir. Encyc. Suppl. iv. 381 = Gagea pygmaea.
virgatum, Poir. Encyc. v. 246 = Nolina georgiana.
viviparum, Reinw. ex Kunth, Enum. Pl. iv. 607 = Chlorophytum Sternbergianum.

PHALARIDIUM, Nees, in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) (Pl. Meyen.) 161 = **Dissanthelium**, Trin. (Gramin.).
peruvianum, Nees & Meyen, l. c. = D. supinum.

PHALARIS, Linn. Syst. ed. I (1735); Gen. ed. I. 14 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1138.
BALDINGERA, Gaertn. Mey. & Schreb. Fl. Wett. i. 43 (1799).

PHALARIS:—
DIGRAPHIS, Trin. Fund. Agrost. 127 (1820).
ENDALLEX, Rafin. in Ser. Bull. Bot. i. (1830) 220.
TYPHOIDES, Moench, Meth. 201 (1794).
alpina, Haenke, in Jacq. Coll. ii. 91; Turra, Fl. Ital. Prod. 62 = Phleum Michelii.
ambigua, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 326 = minor.
americana, Ell. Sketch, i. 101 = intermedia.
americana, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 531 = arundinacea.
amethystina, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) 56.—Calif.
Ammophila, Link, Enum. Hort. Berol. i. 66 = Ammophila arundinacea.
angusta, *Hort. Gryph. ex Schlecht. in Linnaea*, xxiv. (1851) 187.—Hab. ?
angusta, Nees, Agrost. Bras. ii. 391 = intermedia.
appendiculata, Schult. Mant. ii. 216 = paradoxa.
aquatica, Ait. Hort. Kew. ed. I. i. 56 = minor.
aquatica, Delile, ex Boiss. Fl. Orient. v. 471 = canariensis.
aquatica, Desf. Fl. Atlant. i. 56 = tuberosa.
aquatica, Guss. Fl. Sic. Prod. i. 67 = truncata.
aquatica, *Linn. Cent. Pl. Rar.* i. 4; *Amoen. Acad.* iv. 264.—Europ.
aquatica, Sebast. & Mauri, Fl. Rom. Prod. i. 31 = coerulescens.
arenaria, Huds. Fl. Angl. ed. I. 21 = Phleum arenarium.
ariculata, Dicks. ex Roem. & Usteri, Mag. Bot. iv. 10tes St. (1790) 190, sphalm. = *utriculata* ?
aristata, Gouan, ex Beauv. Agrost. 172 = Polypogon monspeliensis.
aristata, Schousb. ex Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 413 = Alopecurus pratensis.
arundinacea, *Linn. Sp. Pl.* 55.—Reg. bor. temp.
arundinacea, Michx. Fl. Bor. Am. i. 43 = intermedia.
aspera, Retz. Obs. iv. 14 = Phleum asperum.
avicularis, Salisb. Prod. 17 = canariensis.
Barrelieri, Tenore, Prod. Fl. Nap. p. viii = Koeleria villosa.
Bellardi, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 415 = Phleum tenue.
Berteroniana, *Steud. Syn. Pl. Gram.* 11.—Chili.
brachystachys, *Link, in Schrad. Neues Journ.* i. III. (1806) 134.—Reg. Mediterr.; Oriens.
brevis, Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1839) 50 = amethystina.
bulbosa, Cav. Ic. i. 46. t. 64 = coerulescens.
bulbosa, Desf. Fl. Atlant. i. 35 = minor.
bulbosa, Linn. Cent. Pl. Rar. i. 4; Amoen. Acad. iv. 264 = Phleum tenue.
bulbosa, Tenore, Fl. Nap. iii. 60 = tuberosa.
caesia, *Nees, Fl. Afr. Austr.* 6.—Afr. austr.
caespitosa, *Thou. Fl. Tristan d'Acugn.* 37 (cespitosa).—Ins. Trist. Acunha.
californica, Hook. & Arn. Bot. Beech. Voy. 161 = intermedia.
canariensis, Brot. Fl. Lusit. i. 9 = brachystachys.
canariensis, *Linn. Sp. Pl.* 54.—Europ.; Afr. bor.
capensis, Thunb. Prod. Pl. Cap. 19 = minor.
caroliniana, Walt. Fl. Carol. 74 = intermedia.
chilensis, J. & C. Presl, Rel. Haenk. i. 245 = intermedia.
ciliaris, [Spreng.] Bot. Gart. Halle, 69 = seq. ?
ciliata, Lag. Gen. et Sp. Nov. 3 = Phleum Michelii.
ciliata, Pourr. in Mém. Acad. Toul. iii. (1788) 323 = Anthoxanthum odoratum, Koeleria villosa.
coerulescens, *Desf. Fl. Atlant.* i. 56.—Reg. Mediterr.
colchaguensis, *Phil. in Linnaea*, xxxiii. (1864–65) 276.—Chili.
colorata, Beauv. Agrost. 172 = arundinacea.
commutata, Roem. & Schult. Syst. ii. 403 = coerulescens.
cristata, Forsk. Fl. Aegypt. Arab. 17 = Polypogon monspeliensis.
crypsoides, Urv. in Mém. Soc. Linn. Par. i. (1822) 263 = Maillea crypsoides.
cuspidata, Hort. ex Steud. Nom. ed. I. 609 = Phleum Michelii.
cylindrica, DC. Fl. Fr. iii. 11 = Phleum tenue.
decumbens, Moench, Meth. 208 = minor.
dentata, *Linn. f. Suppl.* 106.—Afr. austr.
dentata, Sieber, ex Kunth. Enum. Pl. i. 33 = paradoxa.
disticha, Forsk. Fl. Aegypt. Arab. 17 = Sporobolus pungens.

PHALARIS:—
erucaeformis, Linn. Sp. Pl. 55 = Beckmannia erucaeformis.
explicata, Link, in Schrad. Journ. ii. (1799) 312 = Heleochloa alopecuroides.
geniculata, Sibth. & Sm. Fl. Graec. Prod. i. 38 = Heleochloa alopecuroides.
glomerata, Gueldenst. Reisen, ii. 214 = Dactylis glomerata.
gracilis, Parl. Pl. Nov. 36 = minor.
hispida, Thunb. Fl. Jap. 44 = Arthraxon ciliaris.
humilis, *Desv. Opusc.* 63.—Oriens.
intermedia, *Bosc, ex Poir. Encyc. Suppl.* i. 300.—Am. bor.
japonica, *Steud. Syn. Pl. Gram.* 11.—Japon.
laxa, Spreng. ex Steud. Nom. ed. II. ii. 315 = intermedia?
ludoviciana, Torr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1839) 56 = americana.
macrostachya, D. Dietr. Syn. Pl. i. 237, sphalm. = *microstachya*.
maritima, Nutt. Gen. Am. i. 48 = Ammophila arundinacea.
Michelii, Savi, in Nuov. Giorn. Pisa, iv. (1806) 230 = Phleum Michelii.
microstachya, DC. Cat. Hort. Monsp. 131 = intermedia.
minor, *Retz. Obs.* iii. 8.—Graecia; Oriens; Afr. austr.
mollis, Thou. Fl. Tristan d'Acugn. 37 = Polypogon intermedius.
mucronata, Desf. Tabl. ed. II. 12 = Phleum Michelii.
muricata, Forsk. Fl. Aegypt. Arab. 202 = Tragus racemosus.
nepalensis, Trin. Sp. Gram. i. t. 80 = minor.
nitida, Presl, Cyp. et Gram. Sic. 26 = brachystachys.
nodosa, Murr. Syst. ed. XIII. 88 = tuberosa.
obvallata, Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1839) 51 = paradoxa.
occidentalis, Nutt. ex Trin. l. c. 56 = intermedia.
oryzoides, Linn. Mant. ii. 183 = Andropogon squarrosus.
oryzoides, Linn. Sp. Pl. 55 = Leersia oryzoides.
oryzoides, Thunb. Fl. Jap. 44 = Beckmannia erucaeformis.
ovata, Moench, Meth. 208 = canariensis.
paniculata, Ait. Hort. Kew. ed. I. i. 87 = Phleum asperum.
paradoxa, *Linn. Sp. Pl.* ed. II. 1665.—Reg. Mediterr.; Oriens.
pectinata, *Roth, Nov. Pl. Sp.* 34.—Ind. or.
phleoides, Linn. Sp. Pl. 55 = Phleum Boehmeri.
phleoides, Savi, Fl. Pis. i. 58 = Phleum Michelii.
phleoides, Steud. Nom. ed. II. ii. 315 = Phleum arenarium.
picta, Hort. ex Steud. l. c. ed. I. 609 = arundinacea.
praemorsa, Lam. Fl. Fr. iii. 566 = paradoxa.
Pseudo-paradoxa, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 327 = paradoxa.
pubescens, Lam. Encyc. i. 92 = Koeleria villosa.
quadrivalvis, Lag. Gen. et Sp. Nov. 3 = brachystachys.
Robinsoniana, Nees, ex Steud. Syn. Pl. Gram. 11 = intermedia.
rubens, Ehrenb. ex Trin. in Mém. Acad. Petrop. Sér. VI. v. II. (1839) 50 = paradoxa.
sativa, Pers. Syn. i. 78 = Phleum tenue.
segetalis, *Steud. in Lechl. Berb. Am. Austr.* 52, nomen.—Chili.
semineutra, Roem. & Schult. Syst. ii. 406 = Poa semineutra.
semiverticillata, Forsk. Fl. Aegypt. Arab. 17 = Agrostis alba.
setacea, Forsk. l. c. = Pennisetum dichotomum.
Sibthorpii, Griseb. Spicil. Fl. Rumel. ii. 468, in obs. = paradoxa.
subulata, Savi, Fl. Pis. i. 57 = Phleum tenue.
tenuifolia, Hort. ex Steud. Nom. ed. I. 609 = Phleum asperum.
tenuis, Host, Gram. Austr. ii. 27 = Phleum tenue.
trigyna, Host, l. c. iv. 12. t. 20 = Phleum Michelii.
trivialis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. v. II. (1839) 55.—Am. bor.
truncata, *Guss. Fl. Sic. Prod. Suppl.* 18; *Fl. Sic. Syn.* i. 118.—Reg. Mediterr.
tuberosa, Link, in Schrad. Journ. ii. (1799) 312 = coerulescens.
tuberosa, *Linn. Mant.* ii. 557.—Reg. Mediterr.
utricularis, Salisb. Prod. 17 = seq.
utriculata, Linn. Syst. ed. X. 869 = Alopecurus utriculatus.

PHALARIS:—
utriculosa, Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 36 = paradoxa.
vaginiflora, Forsk. Fl. Aegypt. Arab. 18 = Heleochloa schoenoides.
variegata, Spreng. Neue Entdeck. ii. 101 = coerulescens.
velutina, Forsk. Fl. Aegypt. Arab. 17 = Panicum sanguinale.
villosa, Michx. Fl. Bor. Am. i. 43 = Anthaenantia villosa.
villosula, De Not. ex Parl. Fl. Ital. i. 73 = coerulescens.
zizanioides, Linn. Mant. ii. 183 = Andropogon squarrosus.

PHALERIA, Jack, in Malay. Misc. ii. (1822) n. VII. p. 59. *THYMELAEACEAE*, Benth. & Hook. f. iii. 199.
DRIMYSPERMUM, Reinw. in Syll. Ratisb. ii. (1828) 15.
PLUTONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.
Blumei, *Benth. Fl. Austral.* vi. 38.—Malaya; Austral.
capitata, *Jack, in Malay. Misc.* ii. (1822) n. VII. p. 59.—Sumatra.
cauliflora, *Bedd. Forest. Man. Bot.* 180; *Trimen, Cat. Pl. Ceyl.* 77.—Zeylan.
Clerodendron, *F. Muell. ex Benth. Fl. Austral.* vi. 38.—Austral.
coccinea, *F. Muell. Papuan Pl.* ii. 9.—N. Guin.
Cumingii, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 183.—Ins. Philipp.
disperma, *Baill. Adansonia*, xi. (1873-76) 318.—Ins. Tonga.
dubiosa, *Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie*, i. (1844) 616.—Malaya.
laurifolia, *Hook. f. Bot. Mag.* t. 5787.—Ins. Timor.
Neumanni, *F. Muell. ex Benth. Fl. Austral.* vi. 38.—Austral.
octandra, *Baill. Adansonia*, xi. (1873-76) 321.—Ind. or.
Perrottetiana, *Villar, in Blanco, Fl. Filip.* ed. III. *Nov. App.* 183.—Ins. Philipp.

PHALEROCARPUS, G. Don, Gen. Syst. iii. 841 (1834) = **Chiogenes**, Salisb. (Vaccin.).
hispidulus, Heynh. Nom. i. 612 = C. serpyllifolia.
serpyllifolia, G. Don, Gen. Syst. iii. 841 = C. serpyllifolia.

PHALLARIA, Schum. & Thonn. Beskr. Guin. Pl. 112 (1827) = **Plectronia**, Linn. (Rubiac.).
horizontalis, Schum. in Schum. & Thonn. l. c. (= *Canthium horizontale*).—Afr. trop.
lucida, Hochst. in Flora, xxv. (1842) 238 (= *Canthium obovatum*).—Afr. austr.
Schimperi, Hochst. l. c. xxiv. (1841) 1. Intell. 27 = P. Schimperi.
spinosa, Schum. & Thonn. Beskr. Guin. Pl. 113 (= *Canthium Thonningii*).—Afr. trop.

PHALLEROCARPUS, G. Don, Gen. Syst. iii. 788 (1834) = **Chiogenes**, Salisb. (Vaccin.)

PHALOCALLIS, Herb. in Bot. Mag. t. 3710 (1839) = **Cypella**, Herb. (Irid.).
plumbea, Herb. l. c. = C. plumbea.

PHALOË, Dum. Fl. Belg. 110 (1827) = **Sagina**, Linn. (Caryoph.).
nodosa, Dum. l. c. = S. nodosa.
saginoides, Dum. l. c. = S. spergella.
subulata, Dum. l. c. = S. subulata.

PHALOLEPIS, Cass. in Dict. Sc. Nat. l. 248 (1828) = **Centaurea**, Linn. (Compos.).
alba, Cass. l. c. 249 = C. alba.
nitens, Cass. l. c. = C. nitens.
splendens, Cass. l. c. = C. splendens.

PHALONA, Dum. Obs. Gram. Belg. 114 (1823) = **Cynosurus**, Linn. (Gramin.).
Castagnei, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 186 = C. echinatus.
echinata, Dum. Obs. Gram. Belg. 114 = C. echinatus.

PHANERA, Lour. Fl. Cochinch. 37 (1790) = **Bauhinia**, Linn. (Legumin.).
acuminatissima, Miq. Fl. Ind. Bat. Suppl. 287.—Sumatra.
albo-lutea, Miq. Fl. Ind. Bat. i. 1. 1079; et Suppl. 285.—Sumatra.
bicornuta, Miq. Fl. Ind. Bat. Suppl. 286.—Sumatra.
bidentata, Benth. in Miq. Pl. Jungh. 263 = B. bidentata.
Blancoi, Benth. in Miq. l. c. 264 = B. Blancoi.
bracteata, Benth. in Miq. l. c. 264 = B. bracteata.
Burkeana, Benth. in Miq. l. c. 264 = B. esculenta.
catalpaefolia, Miq. Fl. Ind. Bat. i. 1. 1079.—Sumatra.
Championii, Benth. in Hook. Kew Journ. iv. (1852) 78 = B. Championi.
coccinea, Lour. Fl. Cochinch. 37 = B. Cumingiana.
complicata, Miq. Fl. Ind. Bat. i. 1. 70 = B. diphylla.
cordifolia, Miq. l. c. 166 = B. emarginata.
corymbosa, Benth. in Hook. Kew Journ. iv. (1852) 77 = B. glauca.
corymbosa, Benth. in Miq. Pl. Jungh. 264 = B. corymbosa.
Cumingiana, Benth. l. c. 263 = B. Cumingiana.
Cunninghami, Benth. l. c. 264 = B. Cunninghamii.
dasycarpa, Miq. Fl. Ind. Bat. i. 1. 1078.—Sumatra.
debilis, Miq. l. c. 69 = B. debilis.
diphylla, Benth. in Miq. Pl. Jungh. 264 = B. diphylla.
diptera, Miq. Fl. Ind. Bat. i. 1. 70 = B. emarginata.
elongata, Benth. in Miq. Pl. Jungh. 262 = B. elongata.
excelsa, Blume, ex Miq. Fl. Ind. Bat. i. 1. 62 = B. ferruginea.
ferruginea, Benth. in Miq. Pl. Jungh. 262 = B. ferruginea.
Finlaysoniana, Benth. in Miq. l. c. = B. Finlaysoniana.
fulva, Benth. l. c. 262, 263 = B. fulva.
glabrifolia, Benth. l. c. 263 = B. glabrifolia.
glauca, Benth. l. c. 265 = B. glauca.
Griffithiana, Benth. l. c. 263 = B. ferruginea.
integrifolia, Benth. l. c. = B. integrifolia.
Junghuhniana, Benth. l. c.—Java.
Kockiana, Benth. l. c. = B. Kockiana.
Lingua, Miq. Fl. Ind. Bat. i. 1. 67 = B. Lingua.
lucida, Benth. in Miq. Pl. Jungh. 262 = B. emarginata.
macrostachya, Benth. l. c. = B. macrostachya, *Wall.*
maculata, A. Rich. ex Teijsm. & Binn. Cat. Hort. Bog. 268 = B. maculata?
nervosa, Benth. in Miq. Pl. Jungh. 262 = B. nervosa.
ochroleuca, Miq. Fl. Ind. Bat. i. 1. 69 = B. glauca.
parvifolia, Teijsm. & Binn. Cat. Hort. Bog. 268 = B. parvifolia.
phoenicea, Benth. in Miq. Pl. Jungh. 262 = B. phoenicea.
piperifolia, Benth. l. c. = B. anguina?
polyanthra, Miq. Fl. Ind. Bat. i. 1. 1079.—Sumatra.
purpurea, Benth. in Miq. Pl. Jungh. 262 = B. purpurea.
pyrrhaneura, Benth. l. c. 262 = B. pyrrhaneura.
retusa, Benth. l. c. 263 = B. retusa.
rosea, Rich. ex Teijsm. & Binn. Cat. Hort. Bog. 268.—Madag.
rufa, Benth. in Miq. Pl. Jungh. 263 = B. rufa, *R. Grah.*
scandens, Teijsm. & Binn. Cat. Hort. Bog. 268 = B. macrostachya, racemosa.
semibifida, Benth. in Miq. Pl. Jungh. 263 = B. semibifida.
speciosa, Blume, ex Miq. Fl. Ind. Bat. i. 1. 61 = B. elongata.
stipularis, Benth. in Miq. Pl. Jungh. 263 = B. stipularis.
sumatrana, Miq. Fl. Bat. i. 1. 1078.—Sumatra.
Vahlii, Benth. in Miq. Pl. Jungh. 263 = B. Vahlii.
variegata, Benth. l. c. 262 = B. variegata.
velutina, Benth. l. c. = B. velutina.

PHANERANDRA, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 20 = **Leucopogon**, R. Br. (Epacrid.).
esquamata, Stschegl. l. c. = L. esquamatus.

PHANERANTHERA, DC. ex Meissn. Gen. Comm. 189 (1840) = **Nonnea**, Medic. (Borag.).

PHANIA, DC. Prod. v. 114 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 238.
arbutifolia, DC. l. c. 115 = Ageratum arbutifolium.
dissecta, Hook. & Arn. Bot. Beech. Voy. 433 = Eupatorium dissectum.
domingensis, *Griseb. Cat. Pl. Cub.* 145.—Cuba.
matricarioides, *Griseb. l. c.*—Cuba.

PHANIA:—
multicaulis, *DC. in Deless. Ic. Sel.* iv. t. 11.—Am. austr.
trinervia, DC. Prod. v. 115 (Agerati sp.?).—Mexic.
urenifolia, Hook. & Arn. Bot. Beech. Voy. 297 = Fleischmannia urenaefolia.

PHANTIS, Linn. Fl. Zeyl. 191 (1747) = **Atalantia**? Correa (Rutac.).

PHARBITIS, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 438 (Convolv. Or. 56) = **Ipomoea**, Linn. (Convolv.).
acuminata, Choisy, in DC. Prod. ix. 348 = I. acuminata.
aristolochifolia, Rafin. Fl Tellur. iv. 72.—Hab.?
asteropila, Tenore, ex Pasquale, Ind. Sem. Hort. Neap. Adn. (1866).—Hab.?
barbata, G. Don, Gen. Syst. iv. 263 = I. hederacea.
barbigera, G. Don, l. c. 262 = Lettsomia barbigera.
bogotensis, Choisy, in DC. Prod. ix. 341 = I. bogotensis.
bracteata, Choisy, l. c. 344 = I. bracteata.
caerulescens, Sweet, Hort. Brit. ed. III. 482 = I. hederacea.
calycosa, A. Rich. Fl. Cub. Fanerog. ii. 128.—Cuba.
cathartica, Choisy, in DC. Prod. ix. 342 = I. cathartica.
cuspidata, G. Don, Gen. Syst. iv. 263 = I. bicolor.
dealbata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 272 = I. dealbata.
Dillenii, G. Don, Gen. Syst. iv. 263 = I. bicolor.
diversifolia, Lindl. Bot. Reg. t. 1988 = I. hederacea.
eriocalyx, Mart. in DC. Prod. ix. 342 = I. eriocalyx.
Forskoeli, G. Don, Gen. Syst. iv. 263 = I. scabra.
fragrans, Boj. Hort. Maurit. 227 = I. fragrans.
githaginea, Hochst. ex Choisy, in DC. Prod. ix. 341 = I. purpurea.
grandiflora, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 139 = I. grandiflora?
hederacea, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 440 = I. hederacea.
heterophylla, Choisy, in DC. Prod. ix. 344 = I. heterophylla.
heterosepala, Benth. Bot. Voy. Sulph. 134.—N. Granat.
hispida, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 440 = I. purpurea.
insularis, Choisy, l. c. 441 = I. congesta.
jamaicensis, Gib. Enum. Pl. Montev. 28 = I. jamaicensis.
laciniata, Dalz. in Hook. Kew Journ. iii. (1851) 178 = I. laciniata, *C. B. Clarke*.
Learii, Lindl. Bot. Reg. (1841) t. 56 = I. Leari.
lilacina, Schlecht. ex Kunze, in Linnaea, xx. (1847) 31 = I. Schechtendalii.
limbata, Lindl. in Journ. Hort. Soc. v. (1850) 33 = I. hederacea.
Lindeni, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 272.—Mexic.
livescens, Schlecht. ex Kunze, in Linnaea, xx. (1847) 32 = I. livescens.
longepedunculata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 271 = I. longepedunculata.
medians, Choisy, in DC. Prod. ix. 343 = I. villosa, *Ruiz & Pav.*
mollis, Choisy, l. c. 342 = I. mollis.
mutabilis, Boj. Hort. Maurit. 227 = I. bicolor.
Nil, Benth. ex Meissn. in Mart. Fl. Bras. vii. 227 = I. bicolor.
Nil, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 441 = I. hederacea.
Nil, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 180 = I. acuminata.
ostrina, Lindl. Bot. Reg. (1842) t. 51.—Cuba.
polymorpha, Siebold & De Vriese, Fl. Jard. Pays-Bas, i. (1858) 187. tt. 3-5.—Japon.
Preauxii, Webb & Berth. Phyt. Canar. iii. 26.—Ins. Canar.
pubescens, Choisy, in DC. Prod. ix. 344 = I. pubescens, *Lam.*
punctata, G. Don, Gen. Syst. iv. 263 = I. hederacea.
purpurea, Voigt, Hort. Suburb. Calc. 354 = I. purpurea.
Purshii, G. Don, Gen. Syst. iv. 163 = I. hederacea.
quinquefolia, Rafin. Fl. Tellur. iv. 81 = I. quinquefolia.
rosea, Choisy, in DC. Prod. ix. 342 = I. jamaicensis.
rubro-caerulea, Planch. in Fl. des Serres, ix. (1854) t. 966 = I. rubro-caerulea.
scabra, G. Don, Gen. Syst. iv. 263 = I. hederacea.
scabrida, G. Don, l. c. = I. hederacea.

PHARBITIS:—
serotina, Choisy, in DC. Prod. ix. 341 = Ipomoea serotina.
speciosa, Choisy, l. c. 343 = I. bicolor.
tomentosa, Choisy, l. c. 342 = I. jamaicensis.
triloba, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 93 = I. hederacea.
tyrianthina, Hook. Bot. Mag. t. 4024 = I. tyrianthina.
varia, G. Don, Gen. Syst. iv. 263 = I. pubescens, *Hornem.*
variifolia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 390 = I. hederacea.
villosa, G. Don, Gen. Syst. iv. 263 = I. villosa.
violacea, Boj. Hort. Maurit. 227 = I. violacea.

PHARETRELLA, Salisb. Gen. Pl. Fragm. 47 (1866) = **Cyanella**, Linn. (Haemod.).
alba, Salisb. l. c. = C. alba.

PHARIA, Steud. Nom. II. ii. 243 (1841), in syn., err. typ. = **Phania**, DC. (Compos.).

PHARIUM, Herb. in Bot. Reg. t. 1546 (1832) = **Bessera**, Schult. f. (Liliac.).
elegans, Steud. Nom. ed. II. ii. 316 = B. elegans.
fistulosum, Herb. in Bot. Reg. t. 1546 = B. elegans.
Herberti, Steud. Nom. ed. II. ii. 316 = B. elegans.

PHARMACOSYCEA, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 64. t. 2 = **Ficus**, Tourn. (Urtic.).
adhatodaefolia, Miq. l. c. 70 = F. adhatodaefolia.
angustifolia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 333 = F. Segoviae.
anthelmintica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 66 = F. anthelmintica.
dendroctona, Miq. l. c. 67 = F. dendrocida.
glaucescens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 332 = F. glaucescens.
grandaeva, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 70 = F. oblongata.
guyanensis, Miq. l. c. 67 = F. Parkeri.
Hernandezii, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 332 = F. Hernandezii.
jamaicensis, Liebm. l. c. 331, in nota = F. jamaicensis.
laurifolia, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 71 = F. daphniphylla.
mexicana, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 415 = F. mexicana.
obtusiuscula, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 69 = F. obtusiuscula.
Parkeriana, Miq. l. c. 71 = F. Parkeri.
perforata, Miq. l. c. 68 = F. perforata.
peruviana, Miq. l. c. 72.—Peruv.
Pseudoradula, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 414 = F. Pseudoradula.
Radula, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 64. t. 2 B.—Bras.
Radula, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 414 = F. Radula.
rigida, Miq. in Seem. Bot. Voy. Herald, 195 = F. coybana.
vermifuga, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 70 = F. vermifuga.

PHARNACEUM, Linn. Coroll. Gen. 5 (1737).
FICOIDEAE, Benth. & Hook. f. i. 857.
GINGINSIA, DC. Prod. iii. 362 (1828).
HYPERTELIS, E. Mey. ex Fenzl, in Ann. Wien. Mus. ii. (1839) 261.
acidum, *Hook. f. in Hook. Ic. Pl.* t. 1035.—Ins. S. Helena.
albens, *Linn. f. Suppl.* 186; *Thunb. Prod. Pl. Cap.* 53.—Afr. austr.
album, Walp. Rep. ii. 244, in syn. = albens.
arenarium, Spreng. Syst. i. 949 = Mollugo verticillata.
bellidifolium, Poir. Encyc. v. 262 = Mollugo nudicaulis.
Berterianum, Spreng. Syst. i. 949 = Mollugo verticillata.
brevicaule, Bartl. in Linnaea, vii. (1832) 625 = dichotomum.
Cerviana, Linn. Sp. Pl. 272 = Mollugo Cerviana.
Cerviana, Mart. ex Rohrb. in Mart. Fl. Bras. xiv. II. 241 = Mollugo verticillata.
confertum, Eckl. & Zeyh. Enum. 285 = incanum.
cordifolium, Linn. Pl. Afr. Rar. 9; Amoen. Acad. vi. 85 = Polycarpon tetraphyllum.
croceum, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* ii. (1839) 255.—Afr. austr.

PHARNACEUM:—
denticulatum, D. Dietr. Syn. Pl. ii. 1014 = Mollugo oppositifolia.
depressum, Linn. Mant. ii. 564 = Polycarpon depressum.
detonsum, *Fenzl, in Ann. Wien. Mus.* ii. (1839) 253.—Afr. austr.
dichotomum, *Linn. f. Suppl.* 186.—Afr. austr.
diffusum, Bartl. in Linnaea, vii. (1832) 625 = Adenogramma diffusa.
distichum, Linn. Mant. ii. 221 = Mollugo racemosa.
distichum, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 30.—Afr. austr.
fluviale, Eckl. & Zeyh. Enum. 286 = dichotomum.
galioides, Licht. ex Schult. Syst. vi. 692 = Adenogramma diffusa.
galioides, Willd. ex Schult. l. c. = Mollugo verticillata.
glabrum, Steud. Nom. ed. I. 609 = Mollugo Cerviana.
glomeratum, Linn. f. Suppl. 185; Thunb. Prod. Pl. Cap. 53 = Adenogramma galioides.
gracile, *Fenzl, in Ann. Wien. Mus.* ii. (1839) 256.—Afr. austr.
hirtum, Spreng. Syst. i. 949 = Mollugo hirta.
Hoffmannseggianum, Schult. Syst. vi. 692 = Mollugo verticillata.
holosteoides, Turra, Fl. Ital. Prod. 65 = Polycarpon tetraphyllum.
incanum, *Linn. Sp. Pl.* 272.—Afr. austr.
lanatum, *Bartl. in Linnaea*, vii. (1832) 625.—Afr. austr.
Lichtensteinianum, Schult. Syst. vi. 692 = Adenogramma diffusa.
lineare, Andr. Bot. Rep. t. 326 = reflexum.
lineare, Bert. ex Rohrb. in Mart. Fl. Bras. xiv. II. 241 = Mollugo verticillata.
lineare, *Linn. f. Suppl.* 185.—Afr. austr.
Linkii, D. Dietr. Syn. Pl. ii. 1014 = Mollugo pentaphylla.
marginatum, Thunb. Prod. Pl. Cap. 54 = Psammotropha parvifolia.
maritimum, Walt. Fl. Carol. 117 = Sesuvium pentandrum.
microphyllum, *Linn. f. Suppl.* 185.—Afr. austr.
Mollugo, Linn. Mant. ii. 561 = Mollugo oppositifolia.
mucronatum, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 29 = Psammotropha androsacea.
multiflorum, *G. Don, Gen. Syst.* i. 422.—China.
nudicaule, *D. Dietr. Syn. Pl.* ii. 1014 (Quid?).—Ind. or.
occultum, Forsk. Fl. Aegypt. Arab. 58 = Gisekia pharnaceoides.
oppositifolium, Spreng. Syst. i. 949 = Mollugo oppositifolia.
parviflorum, Roth, Nov. Pl. Sp. 186 = Mollugo oppositifolia.
patens, Eckl. & Zeyh. Enum. 285 = detonsum.
pentagonum, Roxb. Hort. Beng. 22; Fl. Ind. ii. 103 = Mollugo hirta.
pentaphyllum, Spreng. Syst. i. 949 = Mollugo pentaphylla.
pruinosum, *Haw. Succ. Pl. Suppl.* 15.—Afr. austr.
quadrangulare, Linn. f. Suppl. 185 = Psammotropha quadrangularis.
quinquefidum, Walp. Rep. ii. 246 = Coelanthum parviflorum.
radiatum, *D. Dietr. Syn. Pl.* ii. 1014.—Chili.
reflexum, *Eckl. & Zeyh. Enum.* 285.—Afr. austr.
rigidum, Bartl. in Linnaea, vii. (1832) 626 = Adenogramma rigida.
salsoloides, Burch. Trav. i. 286 = verrucosum.
scleranthoides, *Sond. in Harv. & Sond. Fl. Cap.* i. 143.—Afr. austr.
semiquinquefidum, Hook. Ic. Pl. t. 83 = Coelanthum parviflorum.
serpyllifolium, *Linn. f. Suppl.* 186.—Afr. austr.
spathulatum, Sw. Fl. Ind. Occ. i. 568 = Mollugo nudicaulis.
sperguloides, Poir. Encyc. v. 260 = Mollugo sperguloides.
strictum, Spreng. Syst. i. 949 = Mollugo stricta.
subtile, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* ii. (1839) 259.—Afr. austr.
suffruticosum, *Baker, in Journ. Linn. Soc.* xx. (1883) 151.—Madag.
suffruticosum, Pall. Reise, iii. 716 = Securinega ramiflora.
teretifolium, *Thunb. Prod. Pl. Cap.* 53.—Afr. austr.
teretifolium, Willd. ex Walp. Rep. ii. 250 = Adenogramma galioides.

PHARNACEUM:—
tomentosum, *Burm. f. Fl. Cap. Prod.* 8 (Quid?).—Afr. austr.
trigonum, *Eckl. & Zeyh. Enum.* 286.—Afr. austr.
triphyllum, Dennst. Schluess. Hort. Malab. 36 = Mollugo triphylla.
triphyllum, Spreng. Syst. i. 949 = Mollugo stricta.
umbellatum, Forsk. Fl. Aegypt. Arab. 58 = Mollugo Cerviana.
verrucosum, *Eckl. & Zeyh. Enum.* 286.—Afr. trop.
verticillatum, Spreng. Syst. i. 949 = Mollugo verticillata.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 141.—Afr. austr.

PHARSEOPHORA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 185 = **Macfadyena**, A. DC. (Bignon.).
allamandaeflora, Miers, l. c. 186 = Bignonia allamandaeflora.
bracteosa, Miers. l. c. = M. bracteosa.
coriacea, Miers, l. c. = Bignonia Selloi?
fallax, Miers, l. c. = Bignonia fallax.
involucrata, Miers, l. c.—Bras.

PHARUS, P. Br. Hist. Jamaic. 344 (1756). *GRAMINEAE*, Benth. & Hook. f. iii. 1110.
angustifolius, *Doell, in Mart. Fl. Bras.* ii. II. 23.—Am. trop.
aristatus, Retz. Obs. v. 23 = Hygroryza aristata.
Banksii, Spreng. Syst. ii. 114 = Leptaspis Banksii.
brasiliensis, *Raddi, Agrost. Bras.* 21.—Bras.
ciliatus, Retz. Obs. v. 23 = Zizania ciliata.
glaber, *H. B. & K. Nov. Gen. et Sp.* i. 196.—Am. trop.
glochidiatus. *J. & C. Presl, Rel. Haenk.* i. 345.—Peruv.
lancifolius, Ham. Prod. Pl. Ind. Occ. 8 = glaber.
lappulaceus, Aubl. Pl. Gui. ii. 859 = latifolius.
lappulaceus, Lam. ex Tuss. Fl. Antill. ii. t. 8 = glaber.
latifolius, *Linn. Syst.* ed. X. 1269.—Am. trop.
latifolius, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 305, 459. t. 9. f. 4 (Quid?).—Hab.?
latifolius, Trin. ex Griseb. Fl. Brit. Ind. 536 = glaber.
micranthus, *Schrad. ex Nees, Agrost. Bras.* ii. 302.—Bras.
natans, Russ. ex Steud. Syn. Pl. Gram. 5 = Hygroryza aristata.
ovalifolius, Ham. Prod. Pl. Ind. Occ. 8 = latifolius.
pubescens, Spreng. Neue Entdeck. i. 241 = brasiliensis.
scaber, H. B. & K. Nov. Gen. et Sp. i. 196 = latifolius.
urceolatus, *Roxb. Hort. Beng.* [104]; *Fl. Ind.* iii. 611.—Ind. or.
virescens, *Doell, in Mart. Fl. Bras.* ii. II. 21.—Am. trop.
vittatus, Lem. in Fl. des Serres, iii. (1847) t. 265, Misc. 50 = latifolius.

PHASELLUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 352 = **Phaseolus**, Tourn. (Legumin.).
farinosus, Medic. Phil. Bot. i. 205 = Phaseolus trinervius.
lanatus, Moench, Meth. 140 = Phaseolus trinervius.
lathyroides, Moench, l. c. = Phaseolus semierectus.
roseus, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 353 = Phaseolus trinervius.
scaber, Moench, Meth. 140 = Phaseolus scaber.
violaceus, Moench, l. c. 141 = Phaseolus violaceus.

PHASEOLUS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 216 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 538.
CANDELIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 352; Phil. Bot. i. 206 (1789).
CARACALLA, Tod. Ind. Sem. Panorm. (1861) 32; et in Illustr. Hortic. ix. (1862) Misc. 13.
LIPUSA, Alef. Landw. Fl. 26 (1866).
PHASELLUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 352.
STROPHOSTYLES, Ell. Sketch, ii. 229 (1822).
abyssinicus, Hort. ex DC. Prod. ii. 396 = violaceus.
abyssinicus, *Savi, in Mem. Acc. Torin.* xxxviii. (1835) 176.—Abyss.
acherensis, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 351.—Hab.?
aconitifolius, *Jacq. Obs. Bot.* iii. 2. t. 52.—Ind. or.; Arab.
aconitifolius, Roxb. Hort. Beng. 55; Fl. Ind. iii. 299 = trilobus.
acutifolius, *A. Gray, Pl. Wright.* i. 43.—N. Mexic.
adenanthus, *G. F. W. Mey. Prim. Fl. Esseq.* 239.—Ind. or. (ubique cult.).
alatus, Linn. Sp. Pl. 725 = Vigna vexillata?

PHASEOLUS:—
alatus, Roxb. Hort. Beng. 54; Fl. Ind. iii. 288 = adenanthus.
alatus, *Schrank, in Syll. Ratisb.* ii. (1828) 79 (Quid?).—Ind. or.
amarus, Roxb. ex Wight & Arn. Prod. 244 = adenanthus.
amazonicus, Benth. in Mart. Fl. Bras. xv. I. 181 = lunatus.
amoenus, Soland. ex Forst. f. Prod. 91 = adenanthus.
amplus, *Benth. Bot. Voy. Sulph.* 85.—Am. austr.
anguinus, *Bunge, Enum. Pl. Chin. Bor.* 20.—China.
angulosus, Orteg. Hort. Matr. Dec. 24 = diversifolius.
angulosus, Schuebl. & Martens, ex Martens, Gartenbohne, 47 = vulgaris.
angustatus, *Wall. Cat.* n. 5597.—Burma.
angustifolius, R. Grah. in Wall. Cat. n. 5596 = Vigna capensis.
angustifolius, Hassk. Cat. Hort. Bog. Alt. 279 = hispidulus.
angustifolius, *Roxb. Hort. Beng.* 55.—Ind. or.
angustissimus, *A. Gray, Pl. Wright.* ii. 33.—N. Mexic.
anisotrichus, *Schlecht. in Linnaea*, xii. (1838) 326.—Mexic.
appendiculatus, *Benth. in Ann. Wien. Mus.* ii. (1838) 137.—Bras.
arcuatus, *Zipp. ex Span. in Linnaea*, xv. (1841) 196, nomen.—Ins. Timor.
Arrabidae, *Steud. Nom.* ed. II. ii. 316.—Bras.
Asellus, Heyne, in Wall. Cat. n. 5588 G = trilobus.
Asellus, *Molina, Sagg. Chile*, 130.—Chili.
asparagioides, Schur, Enum. Pl. Transs. 177 = vulgaris.
asper, Benth. in Ann. Wien. Mus. ii. (1838) 139 = coriaceus.
asper, Benth. in Linnaea, xxii. (1849) 519 = linearis.
asperrimus, Schrank, in Syll. Ratisb. ii. (1828) 81 = Vigna capensis.
atropurpureus, [*Moç. & Sesse*], *ex DC. Prod.* ii. 395.—Mexic.
Atsuki, Siebold, in Verh. Batav. Gen. xii. (1830) 57 = P. Mungo.
aureus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 297 = P. Mungo.
Balansae, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 29.—Parag.
barbulatus, Benth. in Ann. Wien. Mus. ii. (1838) 138 = adenanthus.
bicolor, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 389 = multiflorus.
bicolor, *Vell. Fl. Flum.* 311; vii. t. 123.—Bras.
bilobatus, *Engelm. in Wisliz. Tour N. Mexic.* 109.—N. Mexic.
bipunctatus, Jacq. Hort. Vindob. i. 44 = lunatus.
bracteatus, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 27.—Bras.
bracteatus, *Spreng. Syst.* iii. 253 (an praec.?).—Bras.
bracteolatus, DC. Prod. ii. 391, sphalm. = bracteatus, *Nees & Mart.*
brevipes, Benth. in Ann. Wien. Mus. ii. (1838) 139 = adenanthus.
Bulai, Blanco, Fl. Filip. ed. I. 572 = Crotalaria verrucosa.
Bundoo, Siebold, in Verh. Batav. Gen. xii. (1830) 57 = P. Mungo.
caeduorum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 138 = adenanthus.
caffer, Haberle, ex Steud. Nom. ed. II. ii. 316 = Vigna capensis.
calcaratus, *Roxb. Hort. Beng.* 54; *Fl. Ind.* iii. 289.—Ind. or.; Malaya.
campestris, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Bras.
candidus, *Vell. Fl. Flum.* 311; vii. t. 125.—Bras.
canescens, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 196 = atropurpureus.
capensis, Burm. f. Fl. Cap. Prod. 21 = Rhynchosia glandulosa.
capensis, Hort. ex Schur, Enum. Pl. Transs. 177 = vulgaris.
capensis, Thunb. Prod. Pl. Cap. 130 = Vigna capensis.
Caracalla, *Linn. Sp. Pl.* 725.—Reg. trop.
caracallensis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
caribaeus, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Rhynchosia minima.
chrysanthos, *Savi, in Nuov. Giorn. Sc.* iii. (1822) 318; *et in Mem. Acc. Torin.* xxxviii. (1835) 177.—Hab.?
cinnabarinus, Salisb. Prod. 335 = semierectus.

PHASEOLUS:—
circinalis, Buch.-Ham. in Wall. Cat. n. 5594 = Dunbaria circinalis.
cirrosus, H. B. & K. Nov. Gen. et Sp. v. i. 448 = adenanthus.
citrinus, *Savi, in Mem. Acc. Torin.* xxxviii. (1835) 176.—Hab.?
clitorioides, *Mart. ex Benth. in Ann. Mus. Wien*, ii. (1838) 137.—Bras.
clitorius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = Clitoria Ternatea.
coccineus, Lam. Encyc. iii. 70 = multiflorus.
coccineus, *Linn. Sp. Pl.* 724; *Mill. Gard. Dict.* ed. VIII. n. 6.—Hab.?
coccineus, *Moç. & Sesse, ex G. Don, Gen. Syst.* ii. 350.—Mexic.
cochleatus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 262.—Ind. occ.
cochleatus, *Vell. Fl. Flum.* 312; vii. t. 126.—Bras.
communis, Pritz. Ic. Ind. 832 = vulgaris?
compressus, DC. Prod. ii. 392 = vulgaris.
compressus, Zoll. & Mor. Syst. Verz. Zoll. 3 = lunatus.
conspicuus, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1203, nomen.—Guian.
coriaceus, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 419.—Bras.
cornutus, Blume, ex Miq. Fl. Ind. Bat. i. I. 198 = trilobus.
cornutus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 (Lablab sp.?).—Java.
crassifolius, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 138.—Bras.
crotalarioides, Mart. ex Benth. l. c. 140 = semierectus.
cruentus, Hort. ex Schur, Enum. Pl. Transs. 177 = vulgaris.
Cumingii, Benth. in Ann. Wien. Mus. ii. (1838) 139 (*Cummingii*) = adenanthus.
cytisoides, Zoll. & Mor. Syst. Verz. Zoll. 3 = C. Caracalla.
decipiens, Salzm. ex Benth. in Mart. Fl. Bras. xv. I. 191 = bracteatus, *Nees & Mart.*
decurrens, R. Grah. in Wall. Cat. n. 5612 = Pueraria phaseoloides.
depauperatus, *R. Grah. in Wall. Cat.* n. 5614.—Burma.
derasus, *Schrank, Pl. Rar. Hort. Monac.* t. 89.—Bras.
difformis, Wall. Cat. n. 5599 = Vigna pilosa.
dimidiatus, Haberle, ex Schuebl. & Mart. Fl. Wuert. 472 = vulgaris.
discolor, *Span. ex Miq. Fl. Ind. Bat* i. I. 196.—Java.
diversifolius, *Pers. Syn.* ii. 296.—Am. bor.
dolichoides, Roxb. Hort. Beng. 54; Fl. Ind. iii. 290 = Vigna dolichoides.
dolichoides, Vell. Fl. Flum. 312; vii. t. 127 = Arrabidae.
domingensis, *Zuccagni, in Roem. Collect.* 150.—Ins. S. Doming.
dumosus, Macfad. Fl. Jamaic. i. 279 = vulgaris?
dysophyllus, *Benth. Pl. Hartw.* 287.—Mexic.
ellipticus, Schur, Enum. Pl. Trans.. 177 = Dolichos sphaerospermus.
erectus, *Fisch. Hort. Gorenk.* ed. II. (1812) 70, nomen.—Hab.?
erythroloma, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Bras.
esculentus, Salisb. Prod. 335 = vulgaris.
falcatus, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 304.—Mexic.
farinosus, Linn. Sp. Pl. 724 = trinervius.
ferrugineus, *Salzwed. ex Medic. Beobacht.* (1783) 213.—Hab.?
ficifolius, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 62.—Ind. or.
filiformis, *Benth. Bot. Voy. Sulph.* 13.—Calif.
firmulus, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 138.—Bras.
foecundus, Macfad. Fl. Jamaic. i. 281 = lunatus.
formosus, *H. B. & K. Nov. Gen. et Sp.* vi. 448.—Mexic.
frutescens, Eaton & Wright, Bot. N. Am. ed. VIII. 354 = Wistaria frutescens.
fuscus, *Wall. Pl. As. Rar.* i. 6. t. 6.—Burma.
fusiformis, R. Grah. in Wall. Cat. n. 5607 = Vigna capensis.
gibbosifolius, *Orteg. Hort. Matr. Dec.* 25.—Cuba.
gibbosus, *Blume, ex Miq. Fl. Ind. Bat.* i. I. 201.—Java.
glaber, Roxb. Hort, Beng. 55; Fl. Ind. iii. 291 = P. Mungo.

PHASEOLUS:—
glaber, *Schlecht. in Linnaea*, xii. (1838) 327.—Mexic.
glabrescens, Steud. Nom. ed. II. ii. 317 = P. Mungo.
glycinaeformis, *Weinm. in Flora*, iv. (1821) i. 29.—China.
gonospermus, Savi, in Mem. Phas. iii. (1826?) 21. f. 19 = vulgaris.
gracilis, *Poepp. ex Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Cuba.
Grahamianus, *Wight & Arn. Prod.* 244.—Ind. or.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 379.—Madag.
grandiflorus, *Steud. Nom.* ed. II. ii. 317.—Bras.
grandis, Buch.-Ham. in Wall. Cat. n. 5602 = velutinus.
grandis, *Dalz. & Gibs. Bomb. Fl.* 72.—Ind. or.
haematocarpus, Savi, Mem. Phas. iii. (1826?) 20. f. 17 = vulgaris.
hastaefolius, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 141 = semierectus.
helvolus, *Linn. Sp. Pl.* 724.—Am. bor.
helvulus, Crantz, Inst. ii. 21, sphalm. = praec.
Hernandesii, *Savi, in Nuov. Giorn. Sc.* iii. (1822) 310.—Mexic.
heterophyllus, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 753.—Mexic.
hircinus, Steud. Nom. ed. II. ii. 317, sphalm. = trinervius.
hirsutus, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 140 = lasiocarpus.
hirtus, Retz. Obs. iii. 38 = P. Mungo.
hirtus, Wall. Cat. n. 5593 = calcaratus.
hispanicus, *Gromow, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 256, nomen.—Rossia.
hispidulus, *Hassk. Cat. Hort. Bog. Alt.* 279.—Malaya.
humifusus, *Savi, in Mem. Acc. Torin.* xxxviii. (1835) 177.—Hab.?
humilis, Hassk. Cat. Hort. Bog. Alt. 278 = chrysantha.
hysterinus, *Dur. Ind. Sem. Hort. Burdig* (1866) 27.—China.
ilocanus, Blanco, Fl. Filip. ed. I. 572 = lunatus.
inamoenus, Linn. Sp. Pl. 724 = lunatus.
incanus, *Zoll. & Mor. Syst. Verz. Zoll.* 4.—Java.
ionocarpus, Fingerh. in Linnaea, x. (1836) 8 = vulgaris.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 194.—Afr. trop.
Lambertianus, *D. Dietr. Syn. Pl.* iv. 1196 (Quid?).—Mexic.
lanatus, Benth. in Ann. Wien. Mus. ii. (1838) 137, sphalm. = lunatus.
lanceolatus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 262.—Porto Rico.
lasiocarpus, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 140.—Am. trop.
lathyroides, Linn. Sp. Pl. ed. II. 1018 = semierectus.
latifolius, *Benth. in Ann. Wien. Mus.* ii. (1838) 139.—Bras.
latifolius, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Rhynchosia latifolia.
latisiliquus, Macfad. Fl. Jamaic. i. 282 = lunatus.
leiospermus, Torr. & Gray, Fl. N. Am. i. 280 = pauciflorus, *Benth.*
leptophyllus, *G. Don, Gen. Syst.* ii. 350.—Mexic.
leptospermus, *Lag. Gen. et Sp. Nov.* 22.—Mexic.
leptostachyus, *Benth. in Ann. Wien. Mus.* ii. (1838) 136.—Mexic.
lilacinus, Zuccagni, in Roem. Collect. 148 = vulgaris.
limensis, Macfad. Fl. Jamaic. i. 279 = lunatus?
linearis, *H. B. & K. Nov. Gen. et Sp.* vi. 445.—Bras.
lobatus, *Hook. Bot. Mag.* t. 4076.—Reg. Argent.
lobatus, Roxb. ex Wight & Arn. Prod. 247 = aconitifolius.
longepedunculatus, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Bras.
longifolius, *Benth. l. c.* 139.—Bras.
lucens, Wall. Cat. n. 5601 = Vigna lucens.
lunatus, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 122 = tenuiflorus.
lunatus, Haberle, Sel. Sem. Hort. Pesth. (1824); ex Martens, Gartenbohne, 47 = vulgaris.
lunatus, *Linn. Sp. Pl.* 724.—Reg. trop. (cult.).
lupinoides, *Fingerh. in Linnaea*, x. (1836) 24 (cult.).—Hab.?
luteus, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 200.—Java.
Macfadyeni, Steud. Nom. ed. II. ii. 317 = adenanthus.
macrocarpus, *Benth. in Ann. Wien. Mus.* ii. (1838) 140.—Mexic.

PHASEOLUS:—
macrocarpus, Moench, Meth. 155 = lunatus.
macrocarpus, *Poir. Encyc. Suppl.* iii. 6.—Hab.?
macropoides, *A. Gray. Pl. Wright.* ii. 33.—N. Mexic.
macropus, *Benth. in Ann. Wien. Mus.* ii. (1838) 140.—Mexic.
macrostachyus, Ell. in Journ. Acad. Philad. i. II. (1818) 324 = perennis.
maculatus, Mart. Hort. Reg. Monac. (1829) 185 (Quid?).—Hab.?
maculatus, Scheele, in Linnaea, xxi. (1848) 465 = retusus, *Benth.*
marinus, Burm. Ind. Alt. [18] = semierectus.
maritimus, Benth. in Ann. Wien. Mus. ii. (1838) 140 = semierectus.
maritimus, Salzm. ex Benth. in Mart. Fl. Bras. xv. 1. 183 = membranaceus.
Martii, *Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Bras.
Max, Linn. Sp. Pl. 725 = P. Mungo.
maximus, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 288 = lunatus?
melanospermus, Fingerh. in Linnaea, x. (1836) 8 = vulgaris.
membranaceus, *Benth. in Ann. Wien. Mus.* ii. (1838) 137.—Bras.
menispermoideus, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Rynchosia menispermoidea.
mesoleucus, *Schrank, ex Link, Enum. Hort. Berol.* ii. 232.—Bras.
mexicanus, Mart. Hort. Reg. Monac. (1829) 185 = vulgaris?
micranthus, *Hook. & Arn. Bot. Beech. Voy.* 287.—Mexic.
micranthus, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 196.—Mexic.
microcarpus, *Mart. Ausw. Merkw. Pfl.* 18. t. 12.—Mexic.
microspermus, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. i. 197 = Vigna membranacea.
microspermus, *Orteg. Hort. Matr. Dec.* 130.—Cuba.
minimus, *Roxb. Hort. Beng.* 54; *Fl. Ind.* iii. 290.—China.
Minoomvo, Roxb. ex Wight & Arn. Prod. 246 = P. Mungo.
modestus, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 138 = clitorioides.
mollis, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 228.—Ins. Galap.
monoicus, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Amphicarpaea monoica.
monophyllus, *Benth. in Ann. Wien. Mus.* ii. (1838) 140.—Bras.
multiflorus, *Willd. Sp. Pl.* iii. 1030.—Mexic.
Mungo, *Linn. Mant.* i. 101.—Reg. trop.
Mungo, Wall. Cat. n. 5589 C = calcaratus.
nanus, Linn. Cent. Pl. i. 23; Amoen. Acad. iv. 284 = vulgaris.
nigerrimus, Juss. ex Zuccagni, in Roem. Collect. 149 = vulgaris.
nigricans, Haberle, ex Martens, Gartenbohne, 47 = vulgaris.
obliquaefolius, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 137.—Bras.
oblongifolius, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 27.—Parag.
oblongus, Savi, Mem. Phas. iii. 17. t. 10. f. 14 = vulgaris.
obovatus, R. Grah. in Wall. Cat. n. 5609 = Vigna retusa.
obvallatus, *Schlecht. in Linnaea*, xii. (1838) 328.—Mexic.
opisotrichus, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 193 = trinervius.
ovalispermus, *Fingerh. in Linnaea*, x. (1836) 12, 21.—Hab.?
ovatus, *Benth. in Ann. Wien. Mus.* ii. (1838) 139.—Bras.
Pallar, *Molina, Sagg. Chil.* 130.—Chili.
palmatus, Forsk. Fl. Aegypt. Arab. 214 = aconitifolius.
panduratus, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 141.—Bras.
paniculatus, Michx. Fl. Bor. Am. ii. 60 = perennis.
parabolicus, Nutt. ex Barton, Comp. Fl. Philad. ii. 81 = helvolus.
parviflorus, Schlecht. in Linnaea, xii. (1838) 325 = heterophyllus.
parviflorus, *Stokes, Bot. Mat. Med.* iv. 18.—Jamaic.
parvulus, *Greene, in Coult. Bot. Gaz.* vi. (1881) 217.—N. Mexic.

PHASEOLUS:—
pascuorum, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 137 = peduncularis.
pauciflorus, *Benth. l. c.* 140.—Am. bor. occ.
pauciflorus, *Dalz. in Hook. Kew Journ.* iii. (1851) 209.—Ind. or.
pauciflorus, Sesse & Moç. ex G. Don, Gen. Syst. ii. 356 = Lambertianus.
pedicellatus, *Benth. in Ann. Wien. Mus.* ii. (1838) 137.—Mexic.
peduncularis, Barton, Comp. Fl. Philad. ii. 81 = helvolus.
peduncularis, *H. B. & K. Nov. Gen. et Sp.* vi. 447.—N. Granat.
pedunculatus, Spreng. Syst. iv. Cur. Post. 282 = praec.
perennis, *Walt. Fl. Carol.* 182.—Am. bor.
pictus, Cav. ex Steud. Nom. ed. II. ii. 317 = vulgaris.
pilosus, *H. B. & K. Nov. Gen. et Sp.* vi. 453.—Am. austr.
pisiformis, Fingerh. in Linnaea, x. (1836) 24 = lupinoides.
pisiformis, Schrad. ex Steud. Nom. ed. II. ii. 317 = P. Mungo.
pius, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 138.—Bras.
Pitcheri, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Amphicarpaea Pitcheri.
platyspermus, Haberle, ex Steud. Nom. ed. II. ii. 317 = lunatus.
polymorphus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 346.—Am. bor.
portoricensis, *Bert. ex Spreng. Syst.* iii. 253.—Portorico.
praecox, *Fingerh. in Linnaea*, x. (1836) 17.—Hab.?
prostratus, *Benth. in Mart. Fl. Bras.* xv. 1. 192.—Bras.
pruriens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = Mucuna urens.
psoraleoides, Wight & Arn. Prod. 244 = semierectus.
puberulus, H. B. & K. Nov. Gen. et Sp. vi. 451 = lunatus.
pubescens, Blume, Cat. Gew. Buitenz. 93 = calcaratus.
pulniensis, Wight, Ic. t. 202 = Vigna capensis.
quadrifarius, Baker, in Oliver, Fl. Trop. Afr. ii. 199 = seq.
quadriflorus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 216 = Vigna vexillata.
radiatus, Linn. Sp. Pl. 725 = P. Mungo.
radicans, Benth. in Ann. Wien. Mus. ii. (1838) 138 = adenanthus.
reniformis, Eaton & Wright, N. Am. Bot. ed. VIII. 353 = Rhynchosia tomentosa, *Hook. & Arn.*
repens, R. Grah. in Wall. Cat. n. 5608 = Vigna repens.
retusus, *Benth. Pl. Hartw.* 11.—Am. bor. occ.; Mexic.
retusus, *Moench, Meth.* 155.—Ind. or.
Ricciardianus, *Tenore, Ind. Sem. Hort. Neap.* (1833) 4; *et ex Savi, in Mem. Acc. Torin.* xxxviii. (1835) 173. t. 3.—Hab.?
romanus, Savi, Mem. Phas. iii. 17. t. 10. f. 20 = vulgaris.
rostratus, Wall. Pl. As. Rar. i. 50. t. 63 = adenanthus.
rotundifolius, *A. Gray, Pl. Wright.* ii. 34.—N. Mexic.
Roxburghii, Wight & Arn. Prod. 245 = P. Mungo.
rufus, *Jacq. Hort. Vindob.* i. 13.—Hab.?
rufus, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 29.—Parag.
saccharatus, Macfad. Fl. Jamaic. i. 282 = lunatus.
saccharatus, *Moench, Meth.* 155.—Hab.?
saccharatus, *Stokes, Bot. Mat. Med.* iv. 16.—Jamaic.
saponaceus, Savi, Mem. Phas. iii. 19. t. 10. f. 15 = vulgaris.
scaber, *Steud. Nom.* ed. I. 610.—Hab.?
scaberulus, *Miq. Fl. Ind. Bat.* i. 1. 197.—Java.
scabrellus, *Benth. ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 346.—Mexic.
Schiedeanus, Schlecht. in Linnaea, xii. (1838) 323 = atropurpureus.
Schottii, Benth. in Ann. Wien. Mus. ii. (1838) 139 = longifolius.
sclaraeoides, *Jacq. Enum. Pl. Carib.* 27.—Cuba.
semierectus, *Linn. Mant.* i. 100.—Reg. trop.
senegalensis, Guill. & Perr. Fl. Seneg. Tent. i. 217 = adenanthus.
sepiarius, Dalz. in Hook. Kew Journ. ii. (1850) 33 = Vigna capensis.
setulosus, Dalz. l. c. = P. Mungo.
sinensis, Hort. ex Schur, Enum. Pl. Transs. 177 = vulgaris.

PHASEOLUS:—
sinuatus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 279.—Am. bor.
sordidus, Salisb. Prod. 335 = P. Max.
speciosus, Boj. ex Benth. in Ann. Wien. Mus. ii. (1838) 139 = adenanthus.
speciosus, *H. B. & K. Nov. Gen. et Sp.* vi. 452.—Am. trop.
sphaericus, Savi, Mem. Phas. iii. 20 = vulgaris.
sphaerocarpus, Schur, Enum. Pl. Transs. 945 (Index) = seq.
sphaerospermus, Linn. Sp. Pl. ed. II. 1018 = Dolichos sphaerospermus.
Spixianus, Benth. in Linnaea, xxii. (1849) 518 = membranaceus.
Spixianus, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 137 = pedunculares.
stenophyllus, *Klotzsch, in Schomb. Faun. et Fl. Brit. Gui.* 1204, nomen.—Guian.
stipularis, *Lam. Encyc.* iii. 74.—Peruv.
strictus, A. Br. & Bouché, in Ind. Sem. Hort. Berol. (1852) App. 3 = semierectus.
subglobosus, Fingerh. in Linnaea, x. (1836) 20 = vulgaris.
sublobatus, *Buch.-Ham. in Wall. Cat.* n. 5598 C.—Burma.
sublobatus, Roxb. Hort. Beng. 54; Fl. Ind. iii. 288 = trilobus.
sublobatus, Wall. Cat. n. 5598 A, B = calcaratus.
sublobatus, Wall. l. c. 5598 D = tenuicaulis.
subtortus, Benth. in Ann. Wien. Mus. ii. (1838) 138 = adenanthus.
subtrilobus, Link, Enum. Hort. Berol. ii. 232 = gibbosifolius?
subvolubilis, Buch.-Ham. in Wall. Cat. n. 5605 = P. Mungo.
superbus, *A. DC. in Mém. Soc. Phys. Genèv.* vii. (1836) 310.—Hab. ?
surinamensis, Miq. in Ann. & Mag. Nat. Hist. Ser. I. xi. (1843) 14 = adenanthus.
sylvestris, *H. B. & K. Nov. Gen. et Sp.* vi. 450.—Mexic.
tenuicaulis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 201.—Burma.
tenuiflorus, *Beurl. in Vet. Akad. Handl. Stockh.* 1854, (1856) 121.—Panama.
tetraspermus, Heyne, ex Wall. Cat. sub n. 5536 = Dolichos Lablab.
tomentosus, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1854) 250.—Ins. Galap.
torosus, Roxb. Hort. Beng. 55; Fl. Ind. iii. 298 = calcaratus?
triangularis, Steud. Nom. ed. II. ii. 318 = seq.
triangulus, *Fingerh. in Linnaea*, x. (1836) 12.—Hab. ?
trichocarpus, *Wright, in Sauv. Fl. Cub.* 30.—Cuba.
trilobatus, Baill. in Bull. Soc. Linn. Par. i. (1883) 379 = trilobus.
trilobus, *Ait. Hort. Kew.* ed. I. iii. 30.—As. et Afr. trop.
trilobus, Michx. Fl. Bor. Am. ii. 60 = diversifolius.
trilobus, Wall. Cat. n. 5588 L–N = aconitifolius.
trinervius, *Heyne, ex Wall. Cat.* n. 5603.—Ind. or.; Malaya.
truncatus, Blume, ex Miq. Fl. Ind. Bat. i. 1. 200, = trinervius.
truxillensis, H. B. & K. Nov. Gen. et Sp. vi. 451 = adenanthus.
tuberosus, Eaton & Wright, N. Am. Bot. ed. VIII. 354 = Apios tuberosa.
tuberosus, *Lour. Fl. Cochinch.* 434.—Cochinch.
tuberosus, *Sesse & Moç. ex G. Don, Gen. Syst.* ii. 355.—Mexic.
tumidus, Savi, Mem. Phas. iii. 19. f. 16 = vulgaris.
tunkidus, Steud. Nom. ed. II. ii. 318, sphalm. = praec.
tunkinensis, Lour. Fl. Cochinch. 435 = lunatus.
velutinus, *R. Grah. in Wall. Cat.* n. 5615.—Reg. Himal.; Burma.
vestitus, *Hook. Bot. Misc.* ii. (1831) 216.—Peruv.
vexillatus, Blanco, Fl. Filip. ed. I. 574 = vulgaris.
vexillatus, Linn. Sp. Pl. 724 = Vigna capensis.
vexillatus, Walt. Fl. Carol. 182 = helvolus.
violaceus, *Moench, Meth.* 141; *Steud. Nom.* ed. I. 610.—Afr.
violaceus, Vell. Fl. Flum. 311; vii. t. 124 = grandiflorus.
viridissimus, Tenore, ex Miq. Fl. Ind. Bat. i. 1. 197 = P. Mungo.
virosus, Boj. Hort. Maurit. 106 = Canavalia ensiformis.
vulgaris, *Linn. Sp. Pl.* 723.—Ubique cult.
vulgaris, Wall. Cat. n. 5595 = lunatus.

PHASEOLUS:—
Wightianus, R. Grah. in Wall. Cat. n. 5591 = P. Mungo.
Wightii, Wight & Arn. Prod. 245 = praec.
Wrightii, *A. Gray, Pl. Wright.* i. 43.—N. Mexic.
Xuaresii, Zuccagni, in Roem. Collect. 149 = lunatus.
Zebra, Fingerh. in Linnaea, x. (1836) 9 = vulgaris?

PHASIOLOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.

PHASIOLUS, Moench, Meth. 141 (1794) = praec.

PHAULOTHAMNUS, A. Gray, in Proc. Am. Acad. xx. (1885) 293. *PHYTOLACCACEAE.*
spinescens, *A. Gray, l. c.* 294.—Mexic.

PHAYLOPSIS, Willd. Sp. Pl. iii. 342 (1800). *ACANTHACEAE*, Benth. & Hook. f. ii. 1081.
AETHEILEMA, R. Br. Prod. 478 (1810).
MICRANTHUS, Wendl. Bot. Beobacht. 38, 39 (1798), nomen prius.
angolana, *S. Moore, in Journ. Bot.* xviii. (1880) 229.—Afr. trop.
Barteri, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 27.—Afr. trop.; Ind. occ.
glutinosa, G. Don, ex Loud. Hort. Brit. 244 = parviflora.
imbricata, Sweet, Hort. Brit. ed. I. 327 = longifolia.
longifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1863) 26.—Ins. Madag.
longifolia, *Sims, Bot. Mag.* t. 2433.—Afr. trop.; Arab.; Ind. or.
obliqua, *T. Anders. ex S. Moore, in Journ. Bot.* xviii. (1880) 229.—Afr. trop.
parviflora, *Willd. Sp. Pl.* iii. 342.—Madag.

PHEBALIUM, Vent. Jard. Malm. 102 (1804). *RUTACEAE*, Benth. & Hook. f. i. 292.
amblycarpum, *Benth. Fl. Austral.* i. 345.—Austral.
anceps, DC. Prod. i. 719 = argenteum.
anceps, Sieber, ex A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 136, in nota = squamulosum.
argenteum, *Sm. in Rees, Cycl.* xxvii. n. 3.—Austral.
asteriscophorum, F. Muell. in Trans. Vict. Inst. i. (1855) 31 = Asterolasia Muelleri.
aureum, A. Cunn. in Field, N. S. Wales, 331 = squamulosum.
Baxteri, *Benth. Fl. Austral.* i. 345.—Austral.
Billardierii, *A. Juss. in Mém. Soc. Hist. Nat. Par.* ii. (1825) 134.—Austral.
bilobum, Bartl. in Lehm. Pl. Preiss. i. 172 = rude.
bilobum, *Lindl. in Mitch. Three Exped.* ii. 178.—Austral.
brachyphyllum, *Benth. Fl. Austral.* i. 341.—Austral.
buxifolium, A. Cunn. ex Benth. l. c. 351 = Asterolasia buxifolia.
correaefolium, A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 130 = Asterolasia correifolia.
Daviesii, Hook. f. Fl. Tasman. ii. 358 = glandulosum.
dentatum, *Sm. in Rees, Cycl.* xxvii. n. 2.—Austral.
diosmeum, *A. Juss. in Mém. Soc. Hist. Nat. Par.* ii. (1825) 135. t. 11. f. 3.—Austral.
Drummondii, *Benth. Fl. Austral.* i. 343.—Austral.
elaeagnifolium, A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 132 = squamulosum.
elaeagnoides, Sieber, ex A. Juss. l. c. 136, in nota = Billardierii.
elatius, *Benth. Fl. Austral.* i. 340.—Austral.
elatum, A. Cunn. in Field, N. S. Wales, 331 = Billardierii.
filifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 159.—Austral.
glandulosum, *Hook. in Mitch. Journ. Trop. Austral.* 199.—Austral.
grandiflorum, Hook. Ic. Pl. t. 724 = Asterolasia grandiflora.
hexapetalum, A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 131. t. 11. f. 1 = Asterolasia mollis.
lachnaeoides, *A. Cunn. in Field, N. S. Wales*, 332.—Austral.
lachnoides, Sweet, Hort. Brit. ed. I. 471, nomen = praec.?
lamprophyllum, *Benth. Fl. Austral.* i. 340.—Austral.
linearis, Hort. Mackay, ex Sweet, Hort. Brit. ed. I. 471, nomen (Quid?).—Austral.
microphyllum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 159.—Austral.
montanum, *Hook. Journ. Bot.* i. (1834) 255.—Tasman.
nudum, *Hook. Ic. Pl.* t. 568.—N. Zel.

PHEBALIUM:—
obcordatum, *A. Cunn. ex Benth. Fl. Austral.* i. 342.—Austral.
Oldfieldii, *F. Muell. l. c.* 340.—Austral.
ovatifolium, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 99.—Austral.
ovatum, Sieber, ex A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 136, in nota = Asterolasia correifolia.
ozothamnoides, *F. Muell. in Trans. Vict. Inst.* i. (1855) 31.—Austral.
phylicifolium, *F. Muell. l. c.* 32.—Austral.
phylicioides, Sieber, ex A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 136, in nota = diosmeum.
podocarpoides, F. Muell. in Trans. Vict. Inst. i. (1855) 31 = squamulosum.
pungens, *Benth. Fl. Austral.* i. 338.—Austral.
Ralstoni, *Benth. l. c.* 339.—Austral.
retusum, Hook. Journ. Bot. i. (1834) 254 = Billiardierii.
rotundifolium, *Benth. Fl. Austral.* i. 341.—Austral.
rude, *Bartl. in Lehm. Pl. Preiss.* i. 172.—Austral.
salicifolium, A. Juss. in Mém. Soc. Hist. Nat. Par. ii. (1825) 134 = dentatum.
sediflorum, F. Muell. in Trans. Vict. Inst. i. (1855) 30 = glandulosum.
squamuligerum, Hook. Ic. Pl. t. 727 = Asterolasia squamuligera.
squamulosum, *Vent. Jard. Malm.* 102. t. 102.—Austral.
truncatum, Hook. f. Fl. Tasm. i. 64 = bilobum.
tuberculosum, *Benth. Fl. Austral.* i. 343.—Austral.

PHEBOANTHA, Reichb. Fl. Germ. Excurs. Index 91 (1833), sphalm. = PHLEBOANTHA, Tausch = **Ajuga**, Linn. (Labiat.).

PHEBOLITIS, 'Gaertn.' ex DC. Prod. viii. 201 (1844) = **Phlebolithis**, Gaertn. (Gen. dub. Sapot.).

PHEDIMUS, Rafin. Cat. 15 (1824) = **Sedum**?, Tourn. (Crassul.).
sagittatus, Rafin. l. c. (Quid?).—Am. bor.

PHEGOPYRUM, Peterm. Fl. Bienitz, 92 (1841) = **Fagopyrum**, Tourn. (Polygon.).
esculentum, Peterm. l. c. = F. esculentum.
tataricum, Peterm. l. c. = F. tataricum.

PHEGOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = **Fagus**, Tourn. (Cupul.).

PHELANDRIUM, Neck. Delic. Gallo-Belg. i. 149 (1768) = PHELLANDRIUM, Tourn. = **Oenanthe**, Tourn. (Umbell.).

PHELIMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Michelia**, Linn. (Magnol.).

PHELIPACA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 131, sphalm. = **Phelypaea**, Tourn. (Orobanch.).

PHELIPAEA, Tourn. ex Desf. in Ann. Mus. Par. x. (1807) 298. t. 21 = **Phelypaea**, Tourn. (Orobanch.).

PHELIPANCHE, Pomel, Nouv. Mat. Fl. Atl. 102 (1874) = **Orobanche**, Tourn.
aegyptiaca, Pomel, l. c. 107 = O. aegyptiaca.
arenaria, Pomel, l. c. 103 = O. arenaria.
atlantica, Pomel, l. c. = O. arenaria?
cernua, Pomel, l. c. 105 = O. nana.
coerulea, Pomel, l. c. 103 = O. purpurea.
floribunda, Pomel, l. c. 106 = O. nana.
lavandulacea, Pomel, l. c. = O. lavandulacea.
Mutelii, Pomel, l. c. = O. Muteli.
pulchra, Pomel, l. c. 105 = O. nana.
ramosa, Pomel, l. c. 103 = O. ramosa.
Schultzii, Pomel, l. c. 107 = O. Schultzii.
tenuiflora, Pomel, l. c. 104 = O. Muteli?

PHELIPEA, Pers. Syn. ii. 181 (1807) = **Phelypaea**, Tourn. (Orobanch.).

PHELLANDRIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 77 (1737) = **Oenanthe**, Tourn. (Umbell.).
aquaticum, Linn. Sp. Pl. 255 = Oe. Phellandrium.
ciliatum, Herb. ex DC. Prod. iv. 81 = Spananthe paniculata.

PHELLANDRIUM:—
conioides, Nolte, ex Nym. Consp. 297 = Oenanthe Phellandrium.
divaricatum, Gilib. Fl. Lituan. ii. 33 = Oe. Phellandrium.
globulosum, Bertol. ex Nym. Consp. 299 = Oe. peucedanifolia.
montanum, S. G. Gmel. Reise durch Russl. iv. 48.—Turkestan.
Mutellina, Linn. Sp. Pl. 255 = Ligusticum Mutellina.
stoloniferum, Roxb. Hort. Beng. 21; Fl. Ind. ii. 93 = Oe. stolonifera.

PHELLANDRYUM, Gilib. Fl. Lituan. ii. 33 (1791) = praec.

PHELLINE, Labill. Sert. Austro-Caled. 35. t. 38 (1824). *RUTACEAE*, Benth. & Hook. f. i. 302.
comosa, *Labill. l. c.*—N. Caled.
lucida, *Vieill. ex Baill. Adansonia*, x. (1871–72) 332.—N. Caled.

PHELLOCARPUS, Benth. in Ann. Wien. Mus. ii. (1838) 106 = **Pterocarpus**, Linn. (Legumin.).
acutus, Benth. l. c. = Pteroc. ancylocalyx.
amazonum, Mart. ex Benth. l. c. = Pteroc. Rohrii.
floridus, Benth. l. c. = Pteroc. Rohrii.
laxiflorus, Benth. l. c. = Pteroc. violaceus.

PHELLODENDRON, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 353. *RUTACEAE*, Benth. & Hook. f. i. 301.
amurense, *Rupr. l. c.*—Reg. Amur.
japonicum, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 212.—Japon.

PHELLODERMA, Miers, in Trans. Linn. Soc. xxvii. (1870) 100. t. 27 = **Priva**, Adans. (Verben.).
cuneato-ovata, Miers, l. c. 101. t. 27 = Priva laevis.

PHELLOLOPHIUM, Baker, in Journ. Linn. Soc. xxi. (1884) 349. *UMBELLIFERAE*.
madagascariense, *Baker, l. c.*—Madag.

PHELLOPTERUS, Benth. in Benth. & Hook. f. Gen. i. 905 (1867). *UMBELLIFERAE*, Benth. & Hook. f. l. c.
GLEHNIA, F. Schmidt, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 61 (1867).
littoralis, *Benth. in Benth. & Hook. f. Gen.* i. 905.—Ins. Sachalin.

PHELLOPTERUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 623 (1840) = **Cymopterus**, Rafin. (Umbell.).
glaucus, Nutt. ex Torr. & Gray, l. c. 624 = C. glaucus.
montanus, Nutt. ex Torr. & Gray, l. c. = C. montanus.

PHELYPAEA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 80 (1737). *OROBANCHACEAE*, Benth. & Hook. f. ii. 982.
ALATRAEA, Neck. Elem. i. 374 (1790).
ANOPLANTHUS, Endl. Gen. 727 (1839), pro parte.
ANOPLON, Reichb. Consp. 212[b] (1828).
KOPSIA, Dum. Comm. Bot. 16 (1822).
PHELIPAEA, Tourn. ex Desf. in Ann. Mus. Par. x. (1807) 298. t. 21.
abbreviata, Steud. Nom. ed. II. ii. 318 = Aeginetia pedunculata.
aegyptiaca, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 327 = Orobanche ramosa.
aegyptiaca, Walp. Rep. iii. 463 = Orobanche aegyptiaca.
albiflora, Gren. & Godr. Fl. Fr. ii. 628 = Orobanche ramosa.
aleppensis, Reut. ex Boiss. Fl. Orient. iv. 499 = Orobanche aegyptiaca.
ambigua, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 429 (al. gen.).—Turkestan.
araneosa, Reut. ex Boiss. Fl. Orient. iv, 496 = Orobanche caesia.
arenaria, Walp. Rep. iii. 459 = Orobanche arenaria.
armena, C. Koch, in Linnaea, xvii. (1843) 291 = salsa.
Biebersteinii, Fisch. ex Walp. Rep. iii. 481 = coccinea.
biflora, Spreng. Syst. ii. 818 = Aphyllon uniflorum.
bohemica, Čelak. in Sitzb. Boehm. Ges. (1874) 61 = Orobanche bohemica.
Brunneri, Webb, in Hook. Niger Fl. 167 (al. gen.).—Ins. Cap. Virid.
caesia, Griseb. Spicil. Fl. Rumel. ii. 59 = Orobanche Muteli.
caesia, Guss. Fl. Sic. Syn. 4 = Orobanche Schultzii.

PHELYPAEA:—
caesia, Reut. in DC. Prod. xi. 6 = Orobanche caesia.
californica, G. Don, Gen. Syst. iv. 632 = Aphyllon californicum.
Calotropidis, Walp. Rep. iii. 462 = Cistanche tubulosa.
capensis, G. Don, Gen. Syst. iv. 633 = Harveya squamosa.
carnosa, C. A. Mey. ex Walp. Rep. iii. 463 = Orobanche sicula.
cedreti, Reut. ex Boiss. Fl. Orient. iv. 495 = Orobanche coerulea.
cernua, Pomel, Nouv. Mat. Fl. Atl. 105 = Orobanche nana.
coccinea, *Poir. Encyc.* v. 268.—As. Min.; Reg. Caucas.
coelestis, Reut. in DC. Prod. xi. 5 = Orobanche coelestis.
coerulea, C. A. Mey. Verz. Pfl. Cauc. 104 = Orobanche coerulea.
comosa, G. Don, Gen. Syst. iv. 632 = Orobanche arenaria.
comosa, Torr. & Gray, in Pacif. Rail. Rep. xii. II. Bot. 45 = Aphyllon comosum.
compacta, G. Don, Gen. Syst. iv. 632 (Quid?).—Afr. bor.
concolor, Moris, Diagn. Stirp. 3; et ej. Fl. Sard. iii. 243 = Orobanche Muteli.
cytinoides, Reut. in DC. Prod. xi. 14 = Christisonia neilgherrica.
Delilei, Walp. Rep. iii. 459 = Orobanche aegyptiaca.
elongata, Lojac. in Natural. Sicil. ii. (1883) 133 = Orobanche aegyptiaca.
emarginata, Reut. in DC. Prod. xi. 9 = Orobanche Muteli.
erianthera, Engelm. ex A. Gray, in Proc. Am. Acad. vii. (1868) 372 = Aphyllon multiflorum.
erianthera, S. Wats. Bot. King's Exp. 225 = Aphyllon californicum.
fasciculata, Spreng. Syst. ii. 818 = Aphyllon fasciculatum.
fissa, C. A. Mey. Verz. Pfl. Cauc. 104 (al. gen.).—Reg. Caucas.
flava, C. A. Mey. l. c. (al. gen.).—Reg. Casp.
floribunda, Pomel, Nouv. Mat. Fl. Atl. 106 = Orobanche nana.
foliata, *Lamb. in Trans. Linn. Soc.* x. (1811) 260. t. 7.—As. Min.; Reg. Caucas.
Fraasii, Walp. Rep. iii. 460 = Orobanche lavandulacea.
gigantea, Karel. in Bull. Soc. Nat. Mosc. (1839) 165 = *flava*.
gossypina, *Baker, ex Tristram, Surv. West. Palest.* 377, nomen.—Palaestin.
gratiosa, Webb & Berth. Phyt. Canar. iii. 153 (Quid?).—Ins. Canar.
Gussoneana, Lojac. in Natural. Sicil. i. (1882) 197 = Orobanche Schultzii.
Heldreichii, Reut. in DC. Prod. xi. 8 = Orobanche Heldreichii.
Hesperugo, Webb, in Hook. Niger Fl. 167 (al. gen.).—Afr. trop.
hirtiflora, Reut. in DC. Prod. xi. 10 = Orobanche aegyptiaca.
Hohenackeri, Reut. l. c. = Orobanche Muteli.
hyans, Reut. in Tchihat. Asie Min. Bot. ii. 57, nomen (Quid?).—As. Min.
incana, Payne, Expl. Palest. 116 = salsa.
indica, G. Don, Gen. Syst. iv. 632 = Orobanche aegyptiaca.
indica, Spreng. ex Steud. Nom. ed. II. ii. 318 = Aeginetia indica.
lanuginosa, C. A. Mey. in Ledeb. Fl. Alt. ii. 460 = Orobanche caesia.
lavandulacea, F. Schultz, Arch. Fl. Fr. et All. 99; Reut. in DC. Prod. xi. 7 = Orobanche Schultzii, lavandulacea.
longiflora, C. A. Mey. Verz. Pfl. Cauc. 104 = Orobanche aegyptiaca.
ludoviciana, G. Don, Gen. Syst. iv. 632 = Aphyllon ludovicianum.
lusitanica, Coss. Notes Crit. 43 = Cistanche lutea.
lutea, Brunn. in Flora, xxiii. (1840) II. Beibl. 4 = *Brunneri*.
lutea, Desf. Fl. Atlant. ii. 60. t. 146 = Cistanche lutea.
lutea, Parry, in Am. Natural. viii. (1874) 214 = Aphyllon fasciculatum.
macrantha, C. Koch, in Linnaea, xvii. (1843) 290 = Orobanche ramosa *vel* Muteli.

PHELYPAEA:—
mauritanica, *Coss. & Dur. in Bull. Soc. Bot. Fr.* (1857) 409.—Afr. bor.
Millefolii, Nym. Consp. 556 = Orobanche coerulea.
Muteli, Reut. in DC. Prod. xi. 8 = Orobanche Muteli.
nana, Reichb. f. Ic. Fl. Germ. xx. 88 = Orobanche nana.
obtusiloba, Reut. ex Boiss. Fl. Orient. iv. 496 = Orobanche arenaria.
olbiensis, Coss. Notes Crit. 8 = Orobanche nana.
oxyloba, Reut. in DC. Prod. xi. 9 (Orobanches sp.).—As. Min.
pallens, Bunge, ex Ledeb. Fl. Ross. iii. 312 (Orobanches sp.).—Sibir.
panormitana, Lojac. in Natural. Sicil. i. (1882) 174. t. 6 = Orobanche Muteli.
pedunculata, Walp. Rep. iii. 459 = Orobanche aegyptiaca.
pinetorum, A. Gray, in Proc. Am. Acad. vii. (1868) 371 = Aphyllon pinetorum.
pulchella, C. A. Mey. in Eichw. Pl. Casp. 17. t. 18 (Orobanches sp.).—Reg. Caucas.
pulchra, Pomel, Nouv. Mat. Fl. Atl. 105 = Orobanche nana.
purpurea, Aschers. Fl. Brandenb. 499 = Orobanche purpurea.
pyramidalis, Reut. in DC. Prod. xi. 7 = Orobanche Schultzii.
ramosa, C. A. Mey. Verz. Pfl. Cauc. 104 = Orobanche ramosa.
ramosissima, Genn. Sp. Fl. Sard. 10; ex Barbey, Fl. Sard. Comp. 233 = Orobanche Muteli.
Reuteri, Boiss. & Huet, ex Tchihat. Asie Min. Bot. ii. 57 (Quid?).—As. Min.
Reuteri, Moris, Fl. Sard. iii. 343 = Orobanche Muteli.
Reuteriana, Reichb. f. Ic. Fl. Germ. xx. 117. t. 1839 = Orobanche caesia.
robusta, Walp. Rep. iii. 460 = Orobanche arenaria.
Rosmarini, Welw. ex Nym. Consp. 557 = Orobanche Muteli.
rufescens, Griseb. Spicil. Fl. Rumel. ii. 59 = Orobanche Muteli.
rufescens, Lojac. Contrib. Fl. Sic. 4 = Orobanche Schultzii.
salsa, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 461.—As. bor. et occ.
Schultzii, F. Schultz, ex Reut. in DC. Prod. xi. 7 = Orobanche lavandulacea.
senegalensis, *Reut. in DC. Prod.* xi. 13.—Afr. trop.
serotina, Haussk. ex Boiss. Fl. Orient. iv. 499 = Orobanche aegyptiaca.
sicula, Walp. Rep. iii. 463 = Orobanche sicula.
stricta, [Moris,] Reut. in DC. Prod. xi. 7 = Orobanche Schultzii.
subacaulis, Benth. Scroph. Ind. 55 = Christisonia subacaulis.
syspirensis, C. Koch, in Linnaea, xxii. (1849) 662 = Orobanche coelestis.
tenuiflora, Pomel, Nouv. Mat. Fl. Atl. 104 = Orobanche Muteli.
tinctoria, Walp. Rep. iii. 462 = Cistanche lutea.
Tournefortii, Desf. in Ann. Mus. Par. x. (1807) 298. t. 21 = coccinea.
trichocalyx, Webb & Berth. Phyt. Canar. iii. 154 = Orobanche lavandulacea.
tricholoba, Reut. in DC. Prod. xi. 10 = Orobanche lavandulacea.
trivalvis, Trautv. in Act. Hort. Petrop. v. (1877) 467 (al. gen.).—Turkestan.
tuberosa, A. Gray, in Proc. Am. Acad. vii. (1868) 371 = Aphyllon tuberosum.
tubulosa, Schenk, Sp. Pl. Aegypt. 23 = Cistanche tubulosa.
violacea, Desf. Fl. Altant. ii. 60. t. 115 = Cistanche violacea (in Addend.).

PHELYPEA, Thunb. Nov. Gen. Pl. v. 91 (1784) = **Cytinus**, Linn. (Balanoph.).
sanguinea, Thunb. l. c. = C. dioicus.

PHEMERANTHUS, Rafin. in Med. Repos. N. York, v. (1808) 350 = **Talinum**, Adans. (Portul.).
teretifolius, Rafin. l. c. = T. teretifolium.

PHENAKOSPERMUM, Endl. Prod. Fl. Norf. 35, in obs. (1833) = **Ravenala**, Adans. (Scitam.).
guyanensis, Endl. l. c. = R. guyanensis.

PHENAX, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 191. *URTICACEAE*, Benth. & Hook. f. iii. 392.
angustifolius, *Wedd. l. c.* 193.—Costa Rica.
aquatica, Wedd. in DC. Prod. xvi. 1. 235², lapsu = Pouzolzia pentandra.
asper, *Wedd. l. c.* 235³⁹.—Cuba.
ballotaefolius, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 192.—N. Granat.
concinna, Wedd. in DC. Prod. xvi. 1. 235², lapsu = Pouzolzia Wightii.
Galeottianus, Blume, Mus. Bot. Lugd. Bat. ii. 227 = mexicanus.
Gaudichaudianus, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 193 = mexicanus.
haitensis, Wedd. l. c. 192 = laevigatus.
hirtus, Wedd. in DC. Prod. xvi. 1. 235³⁸ = urticaefolius.
hirtus, Wedd. l. c. 235², lapsu = Pouzolzia Bennettiana.
integrifolius, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 193.—Peruv.
laevigatus, *Wedd. l. c.* 191.—Am. trop.
laxiflorus, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 499.—Peruv.
mexicanus, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 193.—Mexic.
nilghirensis, Wedd. in DC. Prod. xvi. 1. 235², lapsu = Pouzolzia Wightii.
pentandrus, Blume, Mus. Bot. Lugd. Bat. ii. 169 = vulgaris.
pentandrus, Wedd. in DC. Prod. xvi. 1. 235², lapsu = Pouzolzia pentandra.
Pentlandianus, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 192 = urticaefolius.
petiolaris, Wedd. l. c. = urticaefolius.
rugosa, Wedd. in DC. Prod. xvi. 1. 235³⁸ = ballotaefolius.
Sonneratii, Wedd. l. c. 235³⁷ = vulgaris.
Stocksii, Wight, ex Wedd. l. c. 492, lapsu = Pouzolzia pentandra.
ternata, Benn. ex Wedd. l. c., lapsu = Pouzolzia Wightii.
tomentosus, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 192 = uliginosus.
tomentosa, Wight, ex Wedd. in DC. Prod. xvi. 1. 492, lapsu = Pouzolzia Bennettiana.
triandra, Wedd. l. c. 235², lapsu = Pouzolzia parvifolia.
uliginosus, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 192.—Venezuela.
urticaefolius, *Wedd. l. c.*—Jamaic.
urticifolius, Griseb. Fl. Brit. W. Ind. 159 = laevigatus.
urticifolius, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 174 = asper.
vulgaris, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 192.—Am. trop.

PHENIANTHUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83 = **Lonicera**, Linn. (Caprifol.).
sempervirens, Rafin. l. c. = L. sempervirens.

PHENOPUS, Hook. f. Fl. Brit. Ind. iii. 404 (1881), sphalm. = PHAENOPUS DC. = **Lactuca**, Tourn. (Compos.)

PHENOTRICHIS, Steud. Nom. ed. II. ii. 319 (1841), sphalm. = **Pherotrichis**, Decne. (Asclep.).

PHEROSPHAERA, Archer, in Hook. Kew Journ. ii. (1850) 52. *CONIFERAE*, Benth. & Hook. f. iii. 433.
Fitzgeraldi, *F. Muell. in Hook. Ic. Pl.* t. 1383.—Austral.
Hookeriana, *Archer, in Hook. Kew Journ.* ii. (1850) 52, partim.—Tasman.
Hookeriana, Archer, l. c., partim = Microcachys tetragona.

PHEROTRICHIS, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 322. *ASCLEPIADEAE*, [Benth. & Hook. f. ii. 767.]
villosa, *Meissn. Gen. Comm.* 176.—Mexic.

PHIALANTHUS, Griseb. Fl. Brit. W. Ind. 335 (1861). *RUBIACEAE*, Benth. & Hook. f. ii. 106.
myrtilloides, *Griseb. l. c.*—Ind. occ.
resinifluus, *Griseb. Cat. Pl. Cub.* 140.—Cuba.
rigidus, *Griseb. l. c.*—Cuba.
spicatus, Wright, in Sauv. Fl. Cub. 71 = Ceratopyxis verbenacea.
stillans, *Griseb. Cat. Pl. Cub.* 149.—Cuba.

PHILACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1102 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
Griffithii, *Benth. l. c.*—Reg. Himal.

PHIALIS, Spreng. Gen. ii. 631 (1831) = **Eriophyllum**, Lag. (Compos.).
caespitosa, Spreng. l. c. = E. caespitosum.

PHIALODISCUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 497, 655. *SAPINDACEAE*.
unijugatus, *Radlk. l. c.* 655.—Afr. trop.
zambesiacus, *Radlk. l. c.*—Afr. trop.

PHILACTIS, Schrad. Ind. Sem. Hort. Gotting. (1831). —Cf. Linnaea, viii. (1833) Litt. 24. *COMPOSITAE*, Benth. & Hook. f. ii. 356.
longipes, *A. Gray, in Proc. Am. Acad.* xv. (1880) 35. —Mexic.
zinnioides, *Schrad. Ind. Sem. Hort. Gotting.* (1831).—*Cf. Linnaea*, viii. (1833) *Litt.* 24.—Mexic.

PHILADELPHUS, [Riv.] Linn. Syst. ed. I (1735). *SAXIFRAGEAE*, Benth. & Hook. f. i. 642.
SYRINGA, Tourn. ex Adans. Fam. ii. 244 (1763).
acuminatus, Lange, Ind. Sem. Hort. Haun. (1875) 33; ex Bot. Tidsskr. Ser. III. ii. (1877–79) 131 = coronarius.
affinis, *Schlecht. in Linnaea*, xiii. (1839) 419.—Mexic.
aromaticus, Ait. Hort. Kew. ed. I. ii. 156 = Leptospermum ericoides.
aromaticus, Hort. Par. ex Steud. Nom. ed. I. 611 = coronarius.
asperifolius, *Koern. in Regel, Gartenfl.* (1867) 73.—Mexic.
californicus, Benth. Pl. Hartw. 309 = Lewisii.
californicus, Carr. in Rev. Hortic. (1866) 339 = Gordonianus.
chinensis, Hort. ex C. Koch, Dendrol. i. 339 = coronarius.
columbianus, Hort. l. c. 344 = Gordonianus.
cordatus, Hort. l. c. = Gordonianus.
cordifolius, Lange, in Bot. Tidsskr. Ser. III. ii. (1877–79) 132 = Gordonianus ?
coronarius, *Linn. Sp. Pl.* 473.—Europ. austr.
corymbosus, Wall. Cat. n. 3652 = Deutzia staminea.
dubius, Hort. ex C. Koch, Dendrol. i. 343 = grandiflorus.
floribundus, Schrad. ex DC. Prod. iii. 205 = grandiflorus.
floribundus, Usteri, ex Roem. Mag. iii. 7tes St. (1790) 177. t. 2 = Leptospermum scoparium.
globosus, Hort. ex C. Koch, Dendrol. i. 344 = Gordonianus.
Godokokerii, *Hort. Booth, ex C. Koch, l. c.* 346.—Am. bor. ?
Gordonianus, *Lindl. Bot. Reg.* (1838) *Misc.* 21.—Calif.
gracilis, Hort. ex DC. Prod. iii. 206 = hirsutus.
Grahami, Hort. ex C. Koch, Dendrol. i. 344 = Gordonianus.
grandiflorus, Lodd. ex Schrad. in Linnaea, xii. (1838) 393 = coronarius.
grandiflorus, *Willd. Enum. Hort. Berol.* 511; [*Ker-Gawl.*] *Bot. Reg.* t. 570.—Am. bor.
hirsutus, *Nutt. Gen. Am.* i. 301.—Am. bor.
humilis, Hort. ex DC. Prod. iii. 206 = grandiflorus.
imbricatus, Soland. ex Gaertn. Fruct. i. 174 = Baeckea crenulata.
inodorus, Hort. ex Poir. Encyc. Suppl. v. 135 = grandiflorus.
inodorus, *Linn. Sp. Pl.* 470.—Am. bor.
inodorus, Stuv. ex Schrad. in Linnaea, xii. (1838) 399 = hirsutus.
insignis, *Carr. in Rev. Hortic.* (1870) 40.—Hab. ?
intermedius, *Salm-Dyck, Hort. Dyck.* (1834) 204.—Hab. ?
Keteleerii, *Carr. in Rev. Hortic.* (1866) 44.—Hab. ?
laniger, Ait. Hort. Kew. ed. I. ii. 150 = Leptospermum pubescens.
latifolius, Schrad. ex DC. Prod. iii. 206 = grandiflorus.
laxus, Schrad. l. c. = grandiflorus.
Ledebourii, Hort. ex. C. Koch, Dendrol. i. 339 = coronarius.
Lewisii, *Pursh, Fl. Am. Sept.* i. 329.—Am. bor.
mexicanus, *Schlecht. in Linnaea*, xiii. (1839) 418.—Mexic.

PHILADELPHUS:—
microphyllus, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 54.—N. Mexic.
myrtoides, *Bertol. Fl. Guatim.* 21.—Guatemala.
nanus, Mill. Gard. Dict. ed. VIII. n. 2 = coronarius.
nepalensis, Wall. ex Loud. Hort. Brit. 196 = coronarius.
nivalis, Hort. ex C. Koch, Dendrol. i. 343 = pubescens.
niveus, Hort. l. c. = praec.
oreganus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 595 = Gordonianus.
parviflorus, *Carr. in Rev. Hortic.* (1870–71) 460.—China.
pekinensis, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 365 = coronarius.
pendulifolius, *Carr. in Rev. Hortic.* (1875) 360.—Hab. ?
primulaeflorus, Hort.—Cf. Gard. Chron. (1871) 5 = coronarius.
pubescens, Bosc, ex Torr. & Gray, Fl. N. Am. i. 595 = hirsutus.
pubescens, Loisel. in Herb. Amat. iv. 268 = grandiflorus.
Roylei, Hort. ex C. Koch, Dendrol. i. 339 = coronarius.
rubricaulis, *Carr. in Rev. Hortic.* (1870–71) 460.—China.
salicifolius, Hort. ex C. Koch, Dendrol. i. 337 = coronarius.
Salzmanni, Hort. l. c. 338 = coronarius.
satsumanus, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 99 = seq.
Satsumi, Siebold, ex Lindl. & Paxt. Flow. Gard. ii. (1851–52) 102 = coronarius.
Schrenkii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 365 = coronarius.
scoparius, Ait. Hort. Kew. ed. I. ii. 150 = Leptospermum scoparium.
serpyllifolius, *A. Gray, Pl. Wright.* i. 77.—N. Mexic.
speciosus, Hort. ex C. Koch, Dendrol. i. 339 = coronarius.
speciosus, Schrad. ex DC. Prod. iii. 206 = grandiflorus.
stamineus, Wall. Pl. As. Rar. ii. 82 = Deutzia staminea.
stenopetala, *Carr. in Rev. Hortic.* (1870) 340.—Hab. ?
tenuifolius, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 133 = coronarius.
tomentosus, D. Don, Gen. Syst. ii. 807; Wall. Cat. n. 3653 = coronarius.
tomentosus, James, ex Maxim. Hydrang. As. 29 = Deutzia staminea.
trichopetalus, *Koern. in Regel, Gartenfl.* (1867) 73.—Costa Rica.
triflorus, Wall. Cat. n. 3653 = coronarius.
trinervius, Schrad. in Linnaea, xii. (1838) 399 = hirsutus.
undulatus ×, Hort. ex Petz. & Kirch. Arbor. Muscav. 205.
uniflorus, Hort. l. c. 206 = coronarius ?
verrucosus, Schrad. ex DC. Prod. iii. 205 = coronarius.
villosus, Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 606 = hirsutus.
Zeyheri, Schrad. ex DC. Prod. iii. 205 = coronarius.

PHILAGERIA ×, Mast. in Gard. Chron. (1872) 358 (Lapageria × Philesia). *LILIACEAE*.
Veitchii, Mast. l. c.

PHILAGINOPSIS, Walp. Rep. ii. 972 (1843), sphalm. = FILAGINOPSIS, Torr. & Gray = **Diaperia**, Nutt. (Compos.).

PHILAGONIA, Blume, Cat. Gew. Buitenz. 20 (1823); Bijdr. 250 (1825) = **Evodia**, Forst. (Rutac.).
fraxinifolia, Hook. Ic. Pl. t. 710 = E. fraxinifolia.
procera, Blume, ex Nees, in Flora, viii. (1825) 125 = E. fraxinifolia.
sambucina, Blume, Cat. Gew. Buitenz. 20; Bijdr. 250 = E. fraxinifolia.

PHILAMMOS, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = **Astragalus**, Tourn. (Legumin.).
Arbuscula, Stev. l. c. = A. Arbuscula.
arcuatus, Stev. l. c. xxix. (1856) II. 146 = A. arcuatus.
arenarius, Stev. l. c. iv. (1832) 268 = A. arenaria.
ceratoides, Stev. l. c. = A. ceratoides.
compressus, Stev. l. c. xxix. (1856) II. 146 = A. compressus.
corniculatus, Stev. l. c. iv. (1832) 268 = A. corniculatus

PHILAMMOS:—
davuricus, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = Astragalus davuricus.
erioceras, Stev. l. c. xxix. (1856) II. 146 = A. erioceras.
eriolobus, Stev. l. c. = A. depauperatus.
Helmii, Stev. l. c. = A. Helmii.
horizontalis, Stev. l. c. = A. Arbuscula.
macroceras, Stev. l. c. = A. macrolobus.
macrolobus, Stev. l. c. iv. (1832) 268 = A. macrolobus.
pycnolobus, Stev. l. c. xxix. (1856) II. 146 = A. pycnolobus.
stenoceras, Stev. l. c. = A. stenoceras.
Stevenianus, Stev. l. c. iv. (1832) 268 = A. Stevenianus.
subulatus, Stev. l. c. = A. subulatus.

PHILASTREA, Pierre, in Bull. Soc. Linn. Par. i. (1885) 475. *STERCULIACEAE.*
pauciflora, *Pierre, l. c.*—Cambodia.

PHILENOPTERA, Fenzl, in Flora, xxvii. (1844) 312 = **Lonchocarpus**, H. B. & K. (Legumin.).
Heuglini, Pritz. Ic. Ind. ii. 212, cum cit. falsa = L. laxiflorus.
Kotschyana, Fenzl, in Flora, xxvii. (1844) 312 = L. laxiflorus.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 232 = L. laxiflorus.

PHILEOZERA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 459 = **Actinella**, Nutt. (Compos.).
multiflora, Buckl. l. c. = A. odorata.

PHILESIA, Comm. ex Juss. Gen. 41 (1789). *LILIACEAE*, Benth. & Hook. f. iii. 767.
buxifolia, Lam. ex Poir. Encyc. v. 269 = magellanica.
magellanica, *J. F. Gmel. Syst.* 1012.—Reg. Magellan.
rosea, D. Dietr. Syn. Pl. ii. 1122 = Lapageria rosea.

PHILETAERIA, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 283 = **Fouquieria**, H. B. & K. (Tamariscin.).
horrida, Liebm. l. c. = F. formosa.

PHILIBERTIA, H. B. & K. Nov. Gen. et Sp. iii. 195, t. 230 (1818). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 750.
CERAMANTHUS, Kunze, in Linnaea, xx. (1847) 26.
PENTAGONIA, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 364.
ZOSIMA, Phil. Sert. Mendoc. Alt. 29 (1871).
canescens, Decne. in Orb. Voy. Bot. i. t. 9 = solanoides.
crassifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 318.—Mexic.
crispa, *Hemsl. l. c.*—Mexic.
cumanensis, *Hemsl. l. c.*—Mexic.
cynanchoides, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Texas; Mexic.
elegans, *Hemsl. Biol. Centr. Am. Bot.* ii. 318.—Mexic.
flava, *Meyen, Reise*, i. 466.—Peruv.
Gilliesii, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 290 = solanoides.
gracilis, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. iv. t. 403.—Reg. Argent.
grandiflora, Hook. Bot. Mag. t. 3618 = gracilis.
incana, Decne. in DC. Prod. viii. 542 = solanoides.
Lindeniana, *Hemsl. Biol. Centr. Am. Bot.* ii. 318.—Mexic.
linearis, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Calif.; Arizon.
lurida, *Hemsl. Biol. Centr. Am. Bot.* ii. 319.—Mexic.
odorata, *Hemsl. l. c.*—Guatemala.
pannosa, *Hemsl. l. c.* 320.—Mexic.
Pavoni, *Hemsl. l. c.* 319.—Mexic.
rotata, *Griseb. in Goett. Abh.* xxiv. (1879) 233.—Reg. Argent.
rotundifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 320.—Mexic.
solanoides, Decne. in DC. Prod. viii. 542 = flava?
solanoides, *H. B. & K. Nov. Gen. et Sp.* iii. 196. t. 230.—Am. austr.
stellaris, *Griseb. in Goett. Abh.* xxiv. (1879) 284.—Reg. Argent.
Torreyi, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Texas.
undulata, *A. Gray, l. c.* 65.—N. Mexic.
viminalis, *A. Gray, l. c.* 64.—Ind. occ.; Guian.

PHILIPPIA, Klotzsch, in Linnaea, ix. (1834) 354. *ERICACEAE*, Benth. & Hook. f. ii. 591.
ELEUTHEROSTEMON, Klotzsch, in Linnaea, xii. (1838) 219.
abietina, *Klotzsch, l. c.* ix. (1834) 357:—Afr. austr.; Ins. Mascar.
arborescens, *Klotzsch, l. c.* 358.—Ins. Maurit.
aristata, *Benth. in DC. Prod.* vii. 696.—Madag.
brachyphylla, *Benth. l. c.* 697.—Ins. Maurit.
Chamissonis, *Klotzsch, in Linnaea*, ix. (1834) 356.—Afr. austr.
ciliata, *Benth. in DC. Prod.* vii. 695.—Madag.
densa, *Benth. l. c.*—Madag.
floribunda, *Benth. l. c.* 696.—Madag.
galioides, *Benth. l. c.*—Ins. Borbon.; Madag.
Goudotiana, *Klotzsch, in Linnaea*, ix. (1834) 355.—Madag.
Leeana, *Klotzsch, l. c.* xii. (1838) 213.—Afr. austr.
macrocalyx, *Baker, in Journ. Linn. Soc.* xx. (1883) 195.—Madag.
montana, *Klotzsch, in Linnaea*, ix. (1834) 354.—Ins. Maurit.; Borbon.
oophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 195.—Madag.
Parkeri, *Baker, in Journ. Bot.* xx. (1882) 171.—Madag.
parviflora, *Benth. in DC. Prod.* vii. 695.—Madag.
ramosissima, *Benth. l. c.* 696.—Ins. Borbon.
tenuifolia, *Benth. l. c.*—Madag.
tenuissima, *Klotzsch, in Linnaea*, ix. (1834) 358.—Madag.
viscosa, *Benth. in DC. Prod.* vii. 696.—Ins. Borbon.

PHILIPPODENDRON, Endl. Gen. 1004 (1840) = seq.

PHILIPPODENDRUM, Poit. in Ann. Sc. Nat. Sér. II. viii. (1837) 183. t. 3 = **Plagianthus**, Forst. (Malvac.).
regium, Poit. l. c. 184. t. 3 = Plag. betulinus.

PHILLYREA, Linn. Gen. ed. I. 2 (1737). *OLEACEAE*, Benth. & Hook. f. ii. 677.
PHYLLIREA, Tourn. ex Adans. Fam. ii. 223 (1763).
angustifolia, *Linn. Sp. Pl.* 7.—Reg. Mediterr.
brachiata, Stokes, Bot. Comm. i. 26 = angustifolia.
bracteolata, Herb. Lamb. ex D. Don, Prod. Fl. Nep. 107 = Ligustrum nepalense.
buxifolia, Link, Jahrb. i. I. (1820) 157 = media.
capensis, Burm. f. Fl. Cap. Prod. 1 = Polygala bracteolata, myrtifolia.
conferta, Stokes, Bot. Comm. i. 23 = angustifolia, latifolia.
coriacea, Link, Jahrb. i. I. (1820) 155 = latifolia.
decora, *Boiss. & Bal. in Vilm. Fl. Pl. Terre, Suppl.* (1867).—Lazistan.
divaricata, *Vis. Ort. Bot. Pad.* (1842) 144.—Hab.?
grandiflora, Wall. ex DC. Prod. viii. 289 = Ligustrum nepalense.
ilicifolia, Willd. Enum. Hort. Berol. 13 = latifolia.
indica, Lour. Fl. Cochinch. 19 = Olea microcarpa.
laevis, Willd. Enum. Hort. Berol. 12 = latifolia.
lanceolata, Steud. Nom. ed. I. 611 = angustifolia.
latifolia, *Linn. Sp. Pl.* 8.—Reg. Mediterr.
latifolia, Sibth. & Sm. Fl. Graec. i. 2. t. 2 = media.
laurifolia, Hort.—Cf. Gard. Chron. (1883) II. 494 = decora.
levis, Tenore, Fl. Nap. iii. 7 = *laevis*.
ligustrifolia, Mill. Gard. Dict. ed. VIII. n. 4 = media.
longifolia, Link, Jahrb. i. I. (1820) 154 = latifolia.
Lorentei, Walp. Rep. vi. 461 = Olea Lorentii.
Lowei, *DC. Prod.* viii. 293.—Ins. Mader.
major, Zumag. Fl. Pedem. i. 73 = latifolia.
media, *Linn. Syst.* ed. X. 887.—Reg. Mediterr.
media, Tenore, Fl. Nap. iii. 6 = angustifolia.
Medwedewi, Sredinski, ex Trautv. in Act. Hort. Petrop. ix. (1884) 13 = decora.
microphylla, J. Grah. Cat. Pl. Bomb. 108 = Linociera malabarica.
microphylla, Link, ex Steud. Nom. ed. I. 611 = media?
minor, Zumag. Fl. Pedem. i. 73 = angustifolia.
obliqua, Tenore, Fl. Nap. 6, in syn. = angustifolia.
obliqua, Willd. Enum. Hort. Berol. 12 = latifolia.
obtusata, Link, Jahrb. i. I. (1820) 156 = latifolia.
oleaefolia, Mill. Gard. Dict. ed. VIII. n. 5; Willd. Enum. Hort. Berol. 13 = media.

PHILLYREA:—
paniculata, Roxb. Hort. Beng. 3; Fl. Ind. i. 100 = Ligustrum lucidum.
pedunculata, Bory & Chaub. Nouv. Fl. Pélop. 7 = media.
pendula, Willd. Enum. Hort. Berol. 12 = media.
racemosa, Link, Jahrb. i. I. (1820) 160 = media.
ramiflora, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 611 = Linociera macrophylla.
robusta, Blume, Bijdr. 611 = Ligustrum obtusiusculum.
robusta, Roxb. Hort. Beng. 3; Fl. Ind. i. 101 = Ligustrum robustum.
rosmarinifolia, Mill. Gard. Dict. ed. VIII. n. 7 = angustifolia.
spinosa, Mill. l. c. n. 3 = latifolia.
stricta, Bertol. Fl. Ital. i. 43 = latifolia.
terminalis, Heyne, ex Wall. Cat. n. 2816 B = Ligustrum lucidum.
terminalis, Roxb. ex Wall. Cat. n. 2816 A = Ligustrum Roxburghii.
undulata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 314 = Ligustrum undulatum.
variabilis, Timb. ex Nym. Consp. 494 = spp. plurimae.
Vilmoriniana, Boiss. & Bal. ex Boiss. Fl. Orient. iv. 37 = decora.
virgata, Willd. Enum. Hort. Berol. 12 = media.
vulgaris, Caruel, Prod. Fl. Tosc. 444 = spp. plurimae.

PHILOCRENA, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1832) 80. t. 6 = **Tristicha**, Thou. (Podost.).
pusilla, Bong. l. c. = T. hypnoides.

PHILODENDRON, Schott, in Wien. Zeitschr. iii. (1829) 780; ex Endl. in Linnaea, vi. (1831) Litt. 53; Melet. 19 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 978.
BAURSEA, Hoffmgg. Verz. Pfl. 42 (1824).
CALOSTIGMA, Schott, Melet. 19 (1832).
ELOPIUM, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 34.
MECONOSTIGMA, Schott, Melet. 20 (1832).
SPHINCTEROSTIGMA, Schott, l. c. (1832).
TELIPODUS, Rafin. Fl. Tellur. iii. 66 (1836).
aceriferum, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 71 = squamiferum.
Acrocardium, Schott, l. c. viii. (1858) 179 = P. Oxycardium.
aculeatum, *Kunth, Enum. Pl.* iii. 48.—Guian.
acutatum, *Schott, Syn. Aroid.* 94.—Venezuela.
adamantinum, *Mart. ex Schott, l. c.* 114.—Bras.
adhatodaefolium, *Schott, l. c.* 81.—Venezuela.
advena, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 289.—Mexic.
aemulum, *Schott, in Journ. Bot.* ii. (1864) 4.—Bras.
affine, *Hemsl. Diagn. Pl. Nov.* 37.—Guatemala.
alatum, Klotzsch & Karst. ex Engl. in DC. Monog. Phan. ii. 362 = Karstenianum.
alatum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 87.—Peruv.
albo-vaginatum, C. Koch & Sello, Ind. Sem. Hort. Berol. (1853) App. 3 = lacerum.
alternans, *Schott, Syn. Aroid.* 75.—Bras.
amazonicum, Hort. ex Engl. in Mart. Fl. Bras. iii. II. 164 = laciniosum.
ambiguum, Schott, Prod. Aroid. 227 = P. Ochrostemon.
amphibium, *Kunth, Enum. Pl.* iii. 52.—Bras.
anatomicum, Hort. ex Kunth, Ind. Sem. Hort. Berol. (1847) 11 = Monstera deliciosa.
angustatum, *Schott, Syn. Aroid.* 103.—Am. trop.
anisotomum, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = Fenzlii.
arborescens, Kunth, Enum. Pl. iii. 48 = Montrichardia aculeata.
arboreum, Kunth, l. c. Montrichardia aculeata.
asperatum, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 4.—Bras.
Augustinum, *C. Koch, l. c.* (1853) *App.* 4.—Am. trop.
aurantiifolium, Schott, in Oestr. Bot. Zeitschr. viii (1858) 178 = guttiferum.
auritum, *Lindl. in Journ. Hort. Soc.* viii. (1853) 60.—Guatemala.
bipennifolium, Schott, in Oestr. Bot. Wochenbl. v. (1855) 289 = panduraeforme.
bipinnatifidum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Bras.

PHILODENDRON:—
Blanchetianum, *Schott, in Bonplandia*, vii. (1859) 164.—Bras.
brasiliense, *Engl. in Mart. Fl. Bras.* iii. II. 168.—Bras.
brevilaminatum, *Schott, in Journ. Bot.* ii. (1864) 4.—Bras.
brevispathum, *Schott, in Bonplandia*, vii. (1859) 29.—Panama.
callaefolium, Hort. ex Engl. in Mart. Fl. Bras. iii. II. 162 = P. Imbe.
calophyllum, *Brongn. ex Linden & André, Illustr. Hortic.* xviii. (1871) 172. t. 76.—Bras.
cannaefolium, *Mart. in Flora*, xiv. (1831) 451.—Bras.
cannaefolium, Sweet, Hort. Brit. ed. III. 632 = spathiphyllum.
Carderi, Hort.—Cf. Gard. Chron. (1878) II. 56; (1880) I. 759 = verrucosum.
cardiophyllum, C. Koch & Sello, Ind. Sem. Hort. Berol. (1853) App. 4 = eximium.
Clementis, *Griseb. Cat. Pl. Cub.* 220.—Cuba.
consanguineum, *Schott, Syn. Aroid.* 88.—Ins. S. Doming.
corcovadense, *Kunth, Enum. Pl.* iii. 49.—Bras.
cordatum, *Kunth, l. c.* 52.—Bras.
cordifolium, Moritz, ex Schott, Prod. Aroid. 246 = fraternum.
crassinervium, *Lindl. Bot. Reg.* t. 1958.—Bras.
crinipes, Hort. Amstel. ex C. Koch, in Ind. Sem. Hort. Berol. (1855) App. 4 = squamiferum.
crinitum, Hort. ex Engl. in DC. Monog. Phan. ii. 417 = squamiferum.
cruentum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 86.—Peruv.
cuneatum, *Engl. in Bot. Jahrb.* vi. (1885) 281.—N. Granat.
curvilobum, *Schott, Syn. Aroid.* 102.—Am. trop.
cuspidatum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) *App.* 7.—Mexic.; Panama.
cuspidifolium, Mart. ex Schott, Prod. Aroid. 243 = ecordatum.
daemonum, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 17.—Mexic.
Dagilla, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = tripartitum.
daguense, Linden & André, in Illustr. Hortic. (1871) 192. t. 79 = verrucosum.
deflexum, *Poepp. ex Schott, Syn. Aroid.* 101.—Peruv.
deltoideum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 87. t. 298.—Peruv.
deviatum, *Schott, in Bonplandia*, vii. (1859) 29.—Venezuela.
dilaceratum, Hort. ex Endl. in DC. Monog. Phan. ii. 265 = Monstera dilacerata.
dispar, Schott, Syn. Aroid. 79 = Karstenianum.
disparile, *Schott, in Bonplandia*, x. (1862) 86.—Bras.
dolosum, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 71.—Bras.
ecordatum, *Schott, Syn. Aroid.* 83.—Bras.
elaphoglossoides, Schott, l. c. 80 = heterophyllum.
elongatum, *Engl. in Mart. Fl. Bras.* iii. II. 161.—Bras.
erubescens, *C. Koch & Augustin, Ind. Sem. Hort. Berol.* (1854) *App.* 6.—N. Granat.
erythropus, Mart. ex Schott, Syn. Aroid. 76 (= *Adelonema erythropus*).—Bras.
eximium, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 378.—Bras.
fenestratum, *Linden, Cat.* n. 14 (1859) 7.—Mexic.
Fenzlii, *Engl. in Mart. Fl. Bras.* iii. II. 144.—Mexic.
fibrillosum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 86.—Peruv.
Fontanesii, Kunth, Enum. Pl. iii. 48 = Xanthosoma hastifolium.
Fontanesii, Kunth, ex Engl. in DC. Monog. Phan. ii. 385 = Simsii.
fragantissimum, *Kunth, Enum. Pl.* iii. 49.—Guian.
fraternum, *Schott, in Bonplandia*, vii. (1859) 29.—Venezuela.
Ghiesbrechtii, Linden, ex Engl. in DC. Monog. Phan. ii. 407 = sagittifolium.
giganteum, *Schott, Syn. Aroid.* 89.—Am. trop.
Glaziovii, *Hook. f. Bot. Mag.* t. 6813.—Bras.
gloriosum, *André, in Illustr. Hortic.* xxiii. (1876) 194. t. 262.—N. Granat.
gracile, *Schott, Prod. Aroid.* 244.—Costa Rica.
grandifolium, *Schott, in Wien. Zeitschr.* iii. 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Venezuela.

PHILODENDRON:—
guttiferum, *Kunth, Enum. Pl.* iii. 51.—Costa Rica.
hastaefolium, *Regel, in Gartenfl.* (1856) 131. t. 159.—Bras.
hastaefolium, Regel, Ind. Sem. Hort. Petrop. (1855) 21 = seq.
hastatum, *C. Koch & Sello, Ind. Sem. Hort. Berol.* (1854) *App.* 7.—Bras.
hastatum, Schott, Syn. Aroid. 101 = elongatum.
hederaceum, Kunth, Enum. Pl. iii. 49, partim = Hoffmanni.
hederaceum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Am. austr.
heterophyllum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 86. t. 297.—Peruv.
Hoffmanni, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 178.—Costa Rica.
Holtonianum, Mast. in Gard. Chron. (1876) II. 367 = tripartitum.
Holtonianum, *Schott, in Bonplandia*, vii. (1859) 29.—N. Granat.
Hookeri, Schott, Syn. Aroid. 93 = grandifolium.
Imbe, Hort. ex Engl. in DC. Monog. Phan. ii. 407, 408 = sagittifolium, sanguineum.
Imbe, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Am. austr.
imperiale, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 71 = asperatum.
impolitum, Schott, Prod. Aroid. 291 = radiatum.
inaequilaterum, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 16.—Mexic.
inciso-crenatum, Kunth, Enum. Pl. iii. 49 = lacerum.
inconcinnum, *Schott, Syn. Aroid.* 81.—Venezuela.
inops, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 99 = hastatum.
insigne, *Schott, Syn. Aroid.* 73.—Guian.
Isertianum, Schott, Prod. Aroid. 242 = P. Oxycardium.
Jacquinii, Schott, Syn. Aroid. 90 = Hoffmanni?
Karstenianum, *Schott, l. c.* 78.—Venezuela.
Krebsii, *Schott, in Bonplandia*, vii. (1859) 164.—Ins. S. Thomas.
lacerum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Ind. occ.
lacinatum, Engl. in Mart. Fl Bras. iii. II. 164 = laciniosum.
laciniosum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Bras.
lanceolatum, Schott, Syn. Aroid. i. 75 = crassinervium.
latifolium, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 3.—Venezuela.
latilobum, *Schott, Syn. Aroid.* 104.—Peruv.
latipes, C. Koch & Augustin, Ind. Sem. Hort. Berol. (1854) App. 6 = fragrantissimum.
Lechlerianum, *Schott, Prod. Aroid.* 250.—Peruv.
Lehmanni, *Engl. in Bot. Jahrb.* vi. (1885) 282.—N. Granat.
ligulatum, *Schott, Prod. Aroid.* 224.—Costa Rica.
Lindeni, Hort. ex Engl. in DC. Monog. Phan. ii. 388 = verrucosum.
Lindeni, *Wallis, in Belg. Hortic.* xvi. (1866) 202.—Ecuador.
Lindenianum, Wallis, l. c. xvi. (1867) 105 = praec.?
Lindenii, *Schott, Syn. Aroid.* 89.—N. Granat.
linguaeforme, C. Koch, ex Schott, Prod. Aroid. 269 = Simsii.
linguifolium, *Schott, Syn. Aroid.* 74.—Bras.
lingulatum, *Schott, l. c.* 77.—Ind. occ.
Linnaei, *Kunth, Enum. Pl.* iii. 47.—Bras.
longepetiolatum, *Engl. in Bot. Jahrb.* i. (1881) 483.—Guian.
longilaminatum, *Schott, in Bonplandia*, x. (1862) 5.—Bras.
Lundii, Warm. in Kjoeb. Vidensk. Meddel. (1867) 128 = bipinnatifidum.
luridum, Kunth, Enum. Pl. iii. 50 = Staurostigma concinnum.
macroglossum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 197.—Venezuela.
Mamei, *André, in Rev. Hortic.* (1883) 104. f. 21.—Ecuador.
Martini, Schott, Prod. Aroid. 225 = guttiferum.
Mathewsii, *Schott, in Bonplandia*, vii. (1859) 29.—Peruv.
megalophyllum, Schott, Prod. Aroid. 279 = deflexum.
melanochrysum, *Linden & André, Illustr. Hortic.* xx. (1873) 198. t. 149.—N. Granat.
Melinonoi, *Brongn. ex Regel, Gartenfl.* xxiii. (1874) 67. t. 789.—Am. trop.

PHILODENDRON:—
membranaceum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 87.—Peruv.
mexicanum, *Engl. in Mart. Fl. Bras.* iii. II. 143.—Mexic.
micans, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 7 = cuspidatum.
micranthum, *Poepp. ex Schott, in Oestr. Bot. Wochenbl.* v. (1855) 17.—Peruv.
microphyllum, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 7 = cuspidatum.
Minarum, *Engl. in Mart. Fl. Bras.* iii. II. 166.—Bras.
modestum, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 289.—Venezuela.
montanum, *Engl. in Bot. Jahrb.* vi. (1885) 282.—N. Granat.
muricatum, *Schott, in Oestr. Bot Wochenbl.* iv. (1854) 418.—Bras.
nervosum, *Kunth, Enum. Pl.* iii. 51 = Karstenianum?
nervosum, Schott, ex Engl. in DC. Monog. Phan. ii. 359 = P. Ochrostemon.
nigrum, Kunth, Enum. Pl. iii. 52 = Xanthosoma violaceum?
niveo-chermesinum, Linden & André, ex André, Illustr. Hortic. xviii. (1871) 172. sub t. 76 = calophyllum.
oblongum, *Kunth, Enum. Pl.* iii. 51.—Bras.
obtusilobum, *Miq. Delect. Sem. Hort. Amstelod.* (1853); *ex Schott, Syn. Aroid.* 95.—Venezuela?
Ochrostemon, *Schott, Prod. Aroid.* 229.—Bras.
ornatum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 378.—Bras.
Oxycardium, *Schott, Syn. Aroid.* 82.—Bras.
oxyprorum, *Schott, l. c.* 82.—Venezuela.
panduraeforme, *Kunth, Ennm. Pl.* iii. 51.—Bras.
Pearcei, *Hort.—Cf. Gard. Chron.* (1869) 587.—Hab.?
pedatum, Kunth, Enum. Pl. iii. 49 = laciniatum.
peltatum, Schott, Prod. Aroid. 253 = Homalomena peltata.
peregrinum, Kunth, Enum. Pl. iii. 51 = Alocasia macrorrhiza.
pertusum, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 11 = Monstera deliciosa.
pinnatifidum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Venezuela.
pinnatilobum, *Engl. in Mart. Fl. Bras.* iii. II. 162.—Bras.
placidum, Schott, in Bonplandia, vii. (1859) 164 = guttiferum.
planinervium, C. Koch, ex Schott, Syn. Aroid. 74 = crassinervium.
Poeppigii, *Schott, l. c.* 84.—Bras.
Poiteauanum, Schott, Prod. Aroid. 242 = ecordatum.
polytomum, *Schott, in Bonplandia*, vii. (1859) 164.—Mexic.
populneum, *C. Koch, ex Schott, Prod. Aroid.* 258.—Hab.?
Prieureanum, Brongn. ex Engl. in Mart. Fl. Bras. iii. II. 151 = calophyllum.
Prieurianum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 378.—Bras.
propinquum, *Schott, Syn. Aroid.* 78.—Bras.
pteropus, *Mart. ex Engl. in DC. Monog. Phan.* ii. 364.—Bras.
pterotum, *C. Koch & Augustin, Ind. Sem. Hort. Berol.* (1854) *App.* 6.—Venezuela.
punctatum, Kunth, Enum. Pl. iii. 48 = Alocasia indica.
quercifolium, Hort. ex Engl. in Mart. Fl. Bras. iii. II. 164 = laciniosum.
quinquenervium, Miq. Delect. Sem. Hort. Amstelod. (1853); ex Schott, Syn. Aroid. 95 = acutatum.
radiatum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 378.—Mexic.
recurvifolium, *Schott, in Bonplandia*, x. (1862) 5.—Bras.
Reichenbachianum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 197.—Venezuela.
Riedelianum, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 98.—Bras.
robustum, *Schott, l. c.* v. (1865) 33.—Am. trop.
Roezlii, *Bull, Cat.* (1872) 7.—N. Granat.
rotundatum, *Engl. in Kjoeb. Vidensk. Meddel.* (1880) 360.—Bras.
rubens, *Schott, Syn. Aroid.* 84.—Venezuela.
rubro-punctatum, *Hook. f. Bot. Mag.* t. 5948.—Bras.
Rudgeanum, *Schott, Syn. Aroid.* 78.—Guian.

PHILODENDRON:—
Ruizii, *Schott, in Oestr. Bot. Wochenbl.* iv. (1854) 418.—Peruv.
sagittifolium, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 17.—Mexic.
sanguineum, *Regel, Ind. Sem. Hort. Petrop.* (1868) 82; *et in Gartenfl.* (1869) 196. t. 621.—Mexic.
Saueranum, C. Koch, Ind. Sem. Hort. Berol. (1855) App. 3 = eximium.
scaberulum, *Wright, in Sauv. Fl. Cub.* 155.—Cuba.
scandens, C. Koch & Sello, Ind. Sem. Hort. Berol. (1853) 14 = cuspidatum.
Schottianum, *Wendl. ex Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 72.—Costa Rica.
Seguine, *Schott, in Bonplandia*, vii. (1859) 164.—Mexic.
Selloum, *C. Koch, in Bot. Zeit.* x. (1852) 276.—Bras.
Sellowianum, C. Koch, Ind. Sem. Hort. Berol. (1854) 8 = latifolium.
Sellowianum, Kunth, Enum. Pl. iii. 50 = P. Imbe.
serpens, *Hook. f. Bot. Mag.* t. 6375.—N. Granat.
serpens, Hort. ex Engl. in DC. Monog. Phan. ii. 421 = Warszewiczii.
Simsii, Hort. ex C. Koch, Ind. Sem. Hort. Berol. (1854) 8, 9 = hastatum, P. Imbe.
Simsii, *Sweet, Hort. Brit.* ed. III. 632; *Kunth, Enum. Pl.* iii. 48.—Guian.
Sodiroi, *Hort.—Cf. Gard. Chron.* (1883) i. 510.—N. Granat.
Sonderianum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 237.—Bras.
speciosum, *Schott, Melet.* 20.—Bras.
spectabile, Linden, ex Gard. Chron. (1869) 587 (Quid?).—N. Granat.
sphalerum, *Schott, Prod. Aroid.* 235.—Guian.
Splitgerberi, *Schott, l. c.* 251.—Guian.
squamiferum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 87.—Bras.
striatipes, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 11 = Caladium striatipes.
subincisum, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 99 = lacerum?
subovatum, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 289.—Mexic.
surinamense, *Engl. in DC. Monog. Phan.* ii. 361.—Guian.
Swartzianum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 237.—Ins. S. Christoph.
tanyphyllum, Schott, Prod. Aroid. 272 = sagittifolium.
tenue, *C. Koch & Augustin, Ind. Sem. Hort. Berol.* (1854) *App.* 7.—Venezuela.
thaliaefolium, *Schott, Syn. Aroid.* 81.—Venezuel.
trilobatum, Hort. ex Regel, Cat. Pl. Hort. Aksakov. 109 = seq.?
tripartitum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Venezuela.
Tweedieanum, *Schott, in Bonplandia*, vii. (1859) 29.—Bras.
undulatum, *Engl. in DC. Monog. Phan.* ii. 428.—Paraguay.
variifolium, *Schott, Syn. Aroid.* 100.—Peruv.
verrucosum, *Mathieu, ex Schott, l. c.* 85.—N. Granat.
Warszewiczii, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) *App.* 4.—Guatemala.
Weddellianum, Engl. Bot. Jahrb. i. (1881) 485 = laciniosum.
Wendlandii, *Schott, Prod. Aroid.* 221.—Am. centr.
Williamsii, *Hook. f. Bot. Mag.* t. 5899.—Bras.
Wrightii, *Griseb. Cat. Pl. Cub.* 219.—Cuba.
Wullschlaegelii, Schott, Syn. Aroid. 94 = acutatum.

PHILODICE, Mart. in Nov. Act. Nat. Cur. xvii. i. (1835) 16. t. 3. *ERIOCAULEAE*, Benth. & Hook. f. iii. 1023.
cuyabensis, *Koern. in Mart. Fl. Bras.* iii. i. 305.—Bras.
Hoffmannseggii, *Mart. in Nov. Act. Nat. Cur.* xvii. i. (1835) 17. t. 3.—Bras.

PHILOGLOSSA, DC. Prod. v. 567 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 347.
peruviana, *DC. l. c.*—Peruv.

PHILOGYNE, Salisb. in Trans. Hort. Soc. i. (1812) 355 = **Narcissus**, Tourn. (Amaryll.).
Broteri, M. Roem. Syn. Ensat. 212 = N. odorus.
calathina, Salisb. in Trans. Hort. Soc. i. (1812) 356 = N. odorus.
Campernelli, Haw. Monog. Narciss. 66 = N. odorus.
conspicua, Salisb. in Trans. Hort. Soc. i. (1812) 355 = N. odorus.

PHILOGYNE:—
Curtisii, Haw. Monog. Narciss. 6 = Narcissus odorus.
heminalis, Haw. Suppl. Pl. Succ. 136 = N. odorus.
heminalis, Salisb. in Trans. Hort. Soc. i. (1812) 356 = N. heminalis.
Infundibulum, M. Roem. Syn. Ensat. 213 = N. odorus.
interjecta, Haw. Suppl. Pl. Succ. 135 = N. odorus.
isometra, M. Roem. Syn. Ensat. 212 = N. odorus.
laeta, M. Roem. l. c. 211 = N. odorus.
lobata, M. Roem. l. c. 210 = N. odorus.
minor, Haw. Suppl. Pl. Succ. 137 = N. juncifolius.
odora, Haw. l. c. 134 = N. odorus.
rugulosa, Haw. l. c. 133 = N. odorus.
rupicola, M. Roem. Syn. Ensat. 211 = N. juncifolius.
triloba, Haw. Suppl. Pl. Succ. 135 = N. odorus.

PHILOMEDA, Noronha, ex Thou. Gen. Nov. Madag. 17 (1806) = **Gomphia**, Schreb. (Ochnac.).

PHILONOMIA, DC. ex Steud. Nom. ed. II. ii. 88, 320 (1841) = **Macromeria**, D. Don (Borag.).

PHILONOTION, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 421. *AROIDEAE*, Benth. & Hook. f. iii. 981.
Spruceanum, *Schott, l. c.*—Bras.

PHILOSTEMON, Rafin. Fl. Ludov. 107 (1817) = **Rhus**, Linn. (Anacard.).
radicans, Rafin. l. c. = R. Toxicodendron.

PHILOSTEMUM, Steud. Nom. ed. II. ii. 320 (1841) = praec.

PHILOSTIZUS, Cass. in Dict. Sc. Nat. xxxix. 498 (1826) = **Centaurea**, Linn. (Compos.).
Fontanesianus, Cass. l. c. = C. ferox.

PHILOTHECA, Rudge, in Trans. Linn. Soc. xi. (1815) 298. t. 21. *RUTACEAE*, Benth. & Hook. f. i. 293.
australis, *Rudge, l. c.*—Austral.
calida, *F. Muell. Fragm.* vii. 21.—Austral.
ciliata, Hook. in Mitch. Journ. Trop. Austral. 347 = australis.
ericoides, *F. Muell. Fragm.* vii. 21.—Austral.
Gaudichaudii, *G. Don, Gen. Syst.* i. 792 (sp. dub.).—Austral.
Hassellii, *F. Muell. in Wing's South. Sc. Record*, iii. (1883) 3.—Austral.
longifolia, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 16 = Reichenbachiana.
Reichenbachiana, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 253.—Austral.
Steudelii, *Buching. ex Steud. Nom.* ed. II. ii. 320 (Quid?).—Austral.

PHILOTRIA, Rafin. in Am. Monthly Mag. (1818) 175 = **Elodea**, Michx. (Hydroch.).

PHILOXERUS, R. Br. Prod. 416 (1810). *AMARANTACEAE*, Benth. & Hook. f. iii. 40.
BLUTAPARON, Rafin. New Fl. Am. iv. 45 (1836).
CARAXERON, Vaill. ex Rafin. Fl. Tellur. iii. 38 (1836).
aggregatus, *H. B. & K. Nov. Gen. et Sp.* ii. 203.—Am. austr.
brasiliana, Sm. in Rees, Cycl. xxvii. n. 4 = Mogiphanes straminea.
brasiliensis, R. Br. Prod. 416 = Telanthera dentata.
canescens, Poir. Encyc. Suppl. iv. 393 = Gomphrena canescens.
carnosus, *Benth. & Hook. f. Gen.* iii. 40.—Reg. Argent.
conicus, R. Br. Prod. 416 = Gomphrena conica.
crassifolius, *H. B. & K. Nov. Gen. et Sp.* ii. 203.—Am. austr.
diffusus, R. Br. Prod. 416.—Austral.
ferrugineus, Griff. Notul. iv. 341 = Psilotrichum ferrugineum.
flaccidus, Poir. Encyc. Suppl. iv. 392 = Gomphrena flaccida.
heliotropifolium, Griseb. in Goett. Abh. xix. (1874) 82 = Gomphrena Martiana.
humilis, Poir. Encyc. Suppl. iv. 392 = Gomphrena humilis.
lanatus, Poir. l. c. = Gomphrena Brownii.
maracaybiensis, *Klotzsch, ex Benth. & Hook. f. Gen.* iii. 40.—N. Granat.
parviflorus, *Benth. & Hook. f. l. c.*—Austral.

PHILOXERUS:—
portulacoides, A. St. Hil. Voy. Bras. ii. 436 = Iresine portulacoides.
surinamensis, Miq. ex Moq. in DC. Prod. xiii. ii. 339 = Iresine surinamensis.
tenellus, A. Cunn. ex Moq. l. c. = Gomphrena tenella.
vermicularis, Mart. ex Moq. l. c. 341 = Iresine portulacoides.
vermiculatus, R. Br. Prod. 416.—Mexic.
Wrightii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 40.—China.

PHILYEA, Linn. Gen. ed. I. 58 (1737), sphalm. = **Phylica**, Linn. (Rhamn.).

PHILYDRELLA, Caruel, in Nuov. Giorn. Bot. Ital. x. (1878) 91 = **Pritzelia**, F. Muell. (Philyd.).
pygmaea, Caruel, in DC. Monog. Phan. iii. 4 = Pritz. pygmaea.

PHILYDRUM, Banks, ex Gaertn. Fruct. i. 62. t. 16. (1788). *PHILYDRACEAE*, Benth. & Hook. f. iii. 840.
GARCIANA, Lour. Fl. Cochinch. 14 (1790).
glaberrimum, Hook. f. Bot. Mag. t. 6056 = Helmholtzia glaberrima.
Helmholtzii, F. Muell. Fragm. v. 203 = Helmholtzia acorifolia.
lanuginosum, *Banks, ex Gaertn. Fruct.* i. 62. t. 16. f. 10.—As. et Austral. trop.
pygmaeum, R. Br. Prod. 265 = Pritzelia pygmaea.

PHILYRA, Klotzch, in Wiegm. Archiv, i. (1841) 199 = **Argithamnia**, Sw. (Euphorb.).
brasiliensis, Klotzsch, l. c. = A. brasiliensis.

PHINAEA, Benth. in Benth. & Hook. f. Gen. ii. 997 (1876). *GESNERACEAE*, Benth. & Hook. f. l. c.
albo-lineata, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* ii. 472.—Am. centr.

PHIPPSIA, R. Br. in Parry's 1st Voy. App. 285 (err. typ. 185) (1824). *GRAMINEAE*, Benth. & Hook. f. iii. 1147.
algida, *R. Br. l. c.*—Reg. arct.
monandra, Hook. & Arn. Bot. Beech. Voy. 132, in syn. = algida.

PHISALIS, Nocca, in Usteri, Ann. Bot. v. (1793) 48 = **Physalis**, Linn. (Solan.).

PHITOPIS, Hook. f. in Hook. Ic. Pl. t. 1093 (1871). *RUBIACEAE*, Benth. & Hook. f. ii. 84.
multiflora, *Hook. f. in Hook. Ic. Pl.* t. 1093.—Peruv.

PHLEBANTHIA, Reichb. Nom. 205 (1843). *CARYOPHYLLEARUM* gen. dub.

PHLEBOANTHE, Tausch, in Flora, xi. (1828) 322 = **Ajuga**, Linn. (Labiat.).
Laxmanni, Tausch, l. c. 323 = A. Laxmanni.
salicifolia, Schur, Enum. Pl. Transs. 546 = A. salicifolia.

PHLEBOCALYMNA, Griff. ex Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 109. *OLACINEAE*, Benth. & Hook. f. i. 353.
Calleryana, *Baill. Adansonia*, ix. (1868–70) 147.—Ins. Philipp.
Griffithiana, *Benth. & Hook. f. Gen.* i. 353.—Burma.
Lobbiana, *Benth. & Hook. f. l. c.*—Burma.
lobospora, *F. Muell. Fragm.* ix. 151.—Austral.

PHLEBOCARYA, R. Br. Prod. 301 (1810). *HAEMODORACEAE*, Benth. & Hook. f. iii. 675.
ciliata, *R. Br. l. c.*—Austral.
filifolia, *F. Muell. Fragm.* viii. 23.—Austral.
laevis, Lindl. Swan River App. 43 = ciliata.
pilosissima, *F. Muell. Fragm.* viii. 23.—Austral.

PHLEBOCHITON, Wall. in Trans. Med. & Phys. Soc. Calc. vii. (1835) 230 = **Tapiria**, Juss. (Anacard.).
extensum, Wall. l. c. 231 = T. hirsuta.

PHLEBOLITHIS, Gaertn. Fruct. i. 201 (1788). *SAPOTACEARUM* gen. dub. Benth. & Hook. f. iii. 653.
indica, *Gaertn. l. c.*—Ind. or.

PHLEBOPHYLLUM, Nees, in Wall. Pl. As. Rar. iii. 75, 102 (1832) = **Strobilanthes**, Blume (Acanth.).
angustifolium, Benth. ex C. B. Clarke, in Hook. f. Fl. Ind. iv. 434 = S. Kunthianus.
apricum, Benth. in Hook. Kew Journ. v. (1853) 131 = S. apricus.
Kunthianum, Nees, in Wall. Pl. As. Rar. iii. 83 = S. Kunthianus.

PHLEBOSPORIUM, Hassk. in Flora, xxx. (1847) 508 = seq.

PHLEBOSPORUM, Jungh. Reise, 346 (1845) = **Lespedeza**, Michx. (Legumin.).
cytisoides, Jungh. l. c. = L. cytisoides.

PHLEBOTAENIA, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 156. *POLYGALEAE*, Benth. & Hook. f. i. 138.
cuneata, *Griseb. l. c.* 157.—Cuba.

PHLEDINIUM, Spach, Hist. Vég. Phan. vii. 351 (1839) = **Delphinium**, Linn. (Ranunc.).
virgatum, Spach, l. c. 353 = D. halteratum.

PHLEOBANTHE, Ledeb. Fl. Ross. iii. 449 (1846–51), sphalm. = PHLEBOANTHE, Tausch = **Ajuga**, Linn. (Labiat.).

PHLEUM, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1146.
ACHNODON, Link, Hort. Berol. i. 65 (1827).
ACHNODONTON, Beauv. Agrost. 24. t. 7. f. 5 (1812).
CHILOCHLOA, Beauv. l. c. 37. t. 7. f. 2 (1812).
PLANTINIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317.
STELEPHUROS, Adans. Fam. ii. 31 (1763).
aculeatum, Lam. Fl. Fr. iii. 563 = Crypsis aculeata.
Aegilops, Scop. Fl. Carn. ed. II. i. 55 = Triticum ovatum.
africanum, *Lour. Fl. Cochinch.* 61.—Afr. or.
alatum, Host, Fl. Austr. i. 84 = Phalaris coerulescens.
alopecuroides, Pill. & Mitter. It. Posegan. 147. t. 16 = Heleochloa alopecuroides.
alpinum, *Linn. Sp. Pl.* 59.—Reg. bor. et arct.; Am. austr.
ambiguum, Griseb. Spicil. Fl. Rumel. ii. 463 = Boehmeri.
ambiguum, Tenore, Fl. Nap. iii. 64 = Michelii.
annuum, Bieb. Fl. Taur. Cauc. i. 46 = asperum.
arcticum, Giesecke, ex Lange, Consp. Fl. Groenl. i. 155 = alpinum.
arenarium. *Linn. Sp. Pl.* 60.—Europ.
arvense, Pourr. in Mém. Acad. Toul. iii. (1788) 324 = pratense.
asperum, *Jacq. Coll.* i. 110.—Europ.; Oriens.
autumnale, Scop. Fl. Carn. ed. II. i. 56 = Sesleria argentea.
Bellardi, J. F. Gmel. Syst. 166 = geniculatum.
Bellardii, Willd. Enum. Hort. Berol. 85 = tenue.
Bertolonii, DC. Cat. Hort. Monsp. 132 = pratense.
Boehmeri, *Wibel, Prim. Fl. Werth.* 125.—Europ.; As. bor.
brevifolium, Dulac, Fl. Hautes-Pyr. 75 = Boehmeri.
bulbosum, Gouan, Fl. Monsp. 117 = pratense.
capitatum, Scop. Fl. Carn. ed. II. i. 56 = Alopecurus Gerardi.
ciliatum, Gilib. Exercit. ii. 518 = pratense.
ciliatum, Pourr. in Mém. Acad. Toul. iii. (1788) 324 = Boehmeri.
cochinchinense, Lour. Fl. Cochinch. 48 = Paspalum cochinchinense.
collinum, Schur, Enum. Pl. Transs. 945 (Index) = Boehmeri.
commutatum, Gaud. in Alpina, iii. (1808) 4 = alpinum.
crinitum, Roxb. Hort. Beng. 8; Fl. Ind. i. 313 = Polypogon maritimus.
crinitum, Schreb. Beschr. Graes. i. 151 = Polypogon monspeliensis.
cristatum, Scop. Fl. Carn. ed. II. i. 57 = Cynosurus cristatus.
cuspidatum, Willd. Enum. Hort. Berol. Suppl. 5 = Michelii.
Dactylon, Georgi, Beschr. Russ. Reich. iv. 681 = Cynodon Dactylon.

PHLEUM:—
Deckeri, Roem. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 124 = pratense.
dentatum, Pers. Syn. i. 79 = Phalaris dentata.
echinatum, *Host, Gram. Austr.* iii. 8. t. 11.—Europ. austr.
erucoides, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 450 = Beckmannia erucaeformis.
exaratum, Griseb. Spicil. Fl. Rumel. ii. 462 = graecum.
exaratum, *Hochst. ex Boiss. Fl. Orient.* v. 480.—As. Min.; Syria.
fallax, Janka, Koezlem. xii. (1876) 168; ex Just, Jahresb. iv. III. 1061 = pratense.
felinum, Sibth. & Sm. Prod. Fl. Graec. i. 32 = echinatum.
flavum, Scop. Fl. Carn. ed. II. i. 55 = Alopecurus agrestis.
geniculatum, *Bell. ex Vitm. Summa Pl.* i. 176 (Quid?).—Italia.
Gerardi, All. Fl. Pedem. ii. 232 = Alopecurus Gerardi.
Gerardi, *Georgi, Beschr. Russ. Reich.* iv. 684 (Quid?).—Tauria.
Gerardi, [Panz.] in Sturm, Deutschl. Fl. Heft 29 = alpinum.
gibbum, *Boiss. Diagn.* Ser. I. v. 69.—As. Min.
glabratum, Koch, Syn. Fl. Germ. ed. II. 1119 (Index), sphalm. = seq.
glabrum, Bernh. Syst. Verz. Erf. 38 = Boehmeri.
glomeruliflorum, *Steud. Syn. Pl. Gram.* 150.—Afr. trop.
graecum, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 42.—Graecia.
graecum, Willd. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 122 = echinatum.
Haenkeanum, J. & C. Presl, Rel. Haenk. i. 245 = alpinum.
hirsutum, Suter, Fl. Helv. i. 34 = Michelii.
Hostii, Jacq. Cat. Hort. Vindob.; ex Roem. & Schult. Syst. ii. 409 = Michelii.
intermedium, Jord. in F. Schultz, Arch. Fl. Fr et Allem. 325 = pratense.
japonicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 158.—Japon.
Kotschyi, Boiss. Fl. Orient. v. 483 = Boehmeri.
laeve, Bieb. Fl. Taur. Cauc. i. 46 = Boehmeri.
Lima, Hort. ex C. C. Gmel. Fl. Bad. i. 121 = asperum.
macrourum, Willd. ex Steud. Nom. ed. II. i. 365; ii. 321 = Epicampes macroura.
Mesopotamiae, Hort. ex Roem. & Schult. Syst. ii. 382 = tenue.
Michelii, *All. Fl. Pedem.* ii. 233.—Europ.; Reg. Caucas.
microstachyum, Ruiz, ex Nym. in Bot. Notiser (1851), 67 = pratense.
monspeliense, Koel. Desc. Gram. 57 = Polypogon monspeliensis.
montanum, C. Koch, in Linnaea, xxi. (1848) 383 = Boehmeri.
nigricans, Willd. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 123 = alpinum.
nodosum, Linn. Syst. ed. X. 871 = pratense.
odoratum, Hort. ex Roem. & Schult. Syst. ii. 410 = asperum.
palare, Tardent, Ess. Hist. Nat. Bessar. 34 = Boehmeri?
paniculatum, Huds. Fl. Angl. ed. I. 23 = asperum.
parnassicum, Boiss. & Heldr. ex Nym. Consp. 792 = pratense.
phalarideum, Vill. Hist. Pl. Dauph. ii. 60 = Michelii.
Phalaris, Pers. Syn. i. 79 = Boehmeri.
phalaroides, Koel. Desc. Gram. 52 = Boehmeri.
praecox, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 325 = pratense.
pratense, *Linn. Sp. Pl.* 59.—Reg. bor. temp.
ramosum, Gilib. Exercit. ii. 519 = arenarium.
reclinatum, Gorter, Fl. Belg. 18 = pratense.
schoenoides, Jacq. Fl. Austr. v. 29 = Crypsis aculeata.
schoenoides, Linn. Sp. Pl. 60 = Heleochloa schoenoides.
serotinum, Jord. Pugill. Pl. Nov. 141 = pratense.
serrulatum, Boiss. Diagn. Ser. II. iv. 125 = Boehmeri.
sibiricum, Hort. ex C. C. Gmel. Fl. Bad. i. 121 = asperum.
stoloniferum, Host, Fl. Austr. i. 80 = pratense.
strictum, Willd. ex Steud. Nom. ed. II. i. 365; ii. 321 = Epicampis macroura.
subulatum, Spreng. ex Steud. l. c. ii. 12, 321 = Lasiochloa hispida.
supinum, Lam. Fl. Fr. iii. 563 = Heleochloa schoenoides.

PHLEUM:—
tenue, *Schrad. Fl. Germ.* i. 191.—Europ. austr.; Oriens.
trigynum, Schrad. ex Mert. & Koch, Deutschl. Fl. i. 496 = Michelii.
tuberosum, Panz. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 124 = pratense.
vaginatum, Pall. Reise Suedl. Statth. Russ. Reich. ii. 192 = Alopecurus vaginatus.
veluchense, Boiss. Fl. Orient. v. 483 = Boehmeri.
velutinum, Steud. Nom. ed. I. 276, 612 = Panicum sanguinale.
ventricosum, Moench, Meth. 177 = asperum.
villosum, Opiz, Natural. x. (1825) 221 = pratense.
viride, All. Fl. Pedem. ii. 232 = asperum.

PHLOGA, Noronha, ex Thou. Prod. Phyt. 2; in Mélang. (1811) nomen. *PALMAE*, Benth. & Hook. f. iii. 909.
polystachia, *Noronha, l. c.*—Madag.

PHLOGACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 76, 99 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1100.
JANASIA, Rafin. Fl. Tellur. iv. 63 (1836).
LOXANTHUS, Nees, in Wall. Pl. As. Rar. iii. 76, 322 (1832).
albiflorus, Bedd. Ic. Pl. Ind. Or. i. 40 = Diotacanthus albiflorus.
asperulus, *Nees, in Wall. Pl. As. Rar.* iii. 99.—Reg. Himal.
asperulus, Wall. Cat. n. 7171 = Wallichii.
curviflorus, *Nees, in Wall. Pl. As. Rar.* iii. 99.—Reg. Himal.
cymosus, Kurz, For. Fl. Brit. Burma, ii. 246 = Cystacanthus cymosus.
elongatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 508.—Burma.
grandis, Bedd. Ic. Pl. Ind. Or. i. 39 = Diotacanthus grandis.
guttatus, *Nees, in Wall. Pl. As. Rar.* iii. 99.—Reg. Himal.
insignis, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 79 = Cystacanthus insignis.
Jenkinsii, *C. B. Clarke, in Hook. Fl. Brit. Ind.* iv. 513.—Assam; Burma.
latifolius, Wight, Ic. t. 1537 = Gymnostachyum latifolium.
parviflorus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 506.—Reg. Himal.
pubinervius, *T. Anders. l. c.* 508.—Reg. Himal.
pulcherrimus, T. Anders. l. c. 507 = Cystacanthus pulcherrimus.
thyrsiflorus, *Nees, in Wall. Pl. As. Rar.* iii. 99.—Reg. Himal.; Burma.
tubiflorus, *Nees, l. c.*—Reg. Himal.
vitellinus, T. Anders. in Journ. Linn. Soc. ix. (1867) 507 = asperulus.
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 511.—Reg. Himal.

PHLOIODICARPUS, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 14 = **Johrenia**, DC. (Umbell.).
dahuricus, Turcz. in Bull. Soc. Nat. Mosc. (1838) I. 93 = *nudiusculus*.
microcarpus, Ledeb. Fl. Ross. ii. 331.—Sibir. altaic.
nudiusculus, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 14.—Dahuria.
villosus, Turcz. l. c.; et in Bull. Soc. Nat. Mosc. (1838) I. 93; xvii. (1844) 727.—Sibir. baical.

PHLOMIDOPSIS, Link, Handb. i. 479 (1829) = **Phlomis**, Linn. (Labiat.).
tuberosa, Link, l. c. 480 = Phlomis tuberosa.

PHLOMIS, Linn. Syst. ed. I (1735); Gen. ed. I. 177 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1214.
ANEMITIS, BELOAKON, TRAMBIS, Rafin. Fl. Tellur. iii. 87 (1836).
ANTHOCOMA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 569.
BLEPHILOMA, Rafin. Fl. Tellur. iii. 95 (1836).
HERSILIA, Rafin. l. c. 88 (1836).
ORLOWIA, Gueldenst. ex Georgi, Beschr. Russ. Reich. v. 1089 (1800).
PHLOMIDOPSIS, Link, Handb. i. 479 (1829).
PHLOMITIS, Reichb. ex T. Nees, Gen. Fl. Germ. Gamop. ii. Trib. VII. n. 43 (1843).
PHLOMOIDES, Moench, Meth. 403 (1794).

PHLOMIS:—
africana, Beauv. Fl. Owar. ii. 82. t. 111 = Leonotis pallida.
agraria, *Bunge, in Ledeb. Fl. Alt.* ii. 411.—Sibir. altaic.
alba, Blanco, Fl. Filip. ed. I. 474 = Anisomeles ovata.
alba, Forsk. Fl. Aegypt. Arab. 107 = Ballota Forskâhlei.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 373.—Turkestan.
alpina, *Pall. in Act. Acad. Petrop.* ii. (1779) 265. t. 13.—Sibir altaic.
angustifolia, Boiss. & Bal. ex Boiss. Fl. Orient. iv. 783 = linearis.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = aurea.
anisodonta, *Boiss. Diagn.* Ser. I. v. 37.—Persia.
armeniaca, Benth. in DC. Prod. xii. 538 = orientalis.
armeniaca, *Willd. Sp. Pl.* iii. 119.—As. Min.
aspera, Willd. Enum. Hort. Berol. ii. 621 = Leucas aspera.
Aucheri, *Boiss. Diagn.* Ser. I. v. 36.—Persia.
aurea, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 251.—Arab.
Barrelieri, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 430 = crinita.
bicolor, Benth. Lab. Gen. et Sp. 629 = floccosa.
bicolor, Willd. ex Benth. in Linnaea, xi. (1837) 338 = discolor.
biflora, Roxb. Hort. Beng. 44; Fl. Ind. iii. 12 = Leucas procumbens.
biflora, Vahl, Symb. Bot. iii. 77 = Leucas biflora.
biloba, Desf. Fl. Atlant. ii. 25. t. 127 = crinita.
Bourgaei, *Boiss. Fl. Orient.* iv. 787.—Pamphylia.
Bovei, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 585.—Afr. bor.
brachystegia, *Bunge, Lab. Pers.* 77.—Turkestan.
bracteata, Herb. Madr. ex Wall. Cat. n. 2043 F = Leucas capitata.
bracteosa, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 383.—Reg. Himal.
breviflora, *Benth. in Wall. Pl. As. Rar.* i. 62.—Reg. Himal.
brevilabris, *Ehrenb. ex Boiss. Fl. Orient.* iv. 782.—Syria.
Bruguieri, *Desf. in Mém. Mus. Par.* xi. (1824) 9. t. 5.—Mesopot.; Persia.
cachemeriana, Benth. in DC. Prod. xii. 542 = spectabilis.
calycina, Roxb. Fl. Ind. iii. 11 = Roylea elegans.
cancellata, *Bunge, Lab. Pers.* 76.—Persia.
capensis, *Thunb. Prod. Fl. Cap.* 95.—Afr. austr.
capitata, *Boiss. Diagn.* Ser. II. iv. 96.—As. Min.
caribaea, Jacq. Coll. i. 154 = Leucas martinicensis.
cashmeriana, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 382.—Reg. Himal.
Cephalotes, Koen. ex Roxb. Hort. Beng. 46; Fl. Ind. iii. 10; et Roth, Nov. Sp. Pl. 262 = Leucas capitata.
chinensis, Blume, Bijdr. 829 = Leucas javanica.
chinensis, Retz. Obs. ii. 19 = Leucas chinensis.
chinensis, Wight, ex Wall. Cat. n. 2524 C = Leucas marrubioides.
chorassanica, *Bunge, Lab. Pers.* 76.—Persia.
chrysophylla, *Boiss. Diagn.* Ser. I. xii. 89.—Syria.
ciliata, Heyne, ex Wall. Cat. n. 2046 = Leucas ciliata.
clandestina, Bory & Chaub. Exp. Sc. Mor. 170 = Sideritis theezans.
collina, Salisb. Prod. 84 = fruticosa.
condensata, *Mart. Hort. Erlang.* (1814) 123, nomen.—Hab. ?
cordata, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 782.—Syria.
cordata, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 382 = bracteosa.
cretica, Presl, Delic. Prag. 84 = ferruginea.
crinita, *Cav. Ic.* iii. 25. t. 247.—Reg. Mediterr. occ.
cymifera, Boiss. Diagn. Ser. I. vii. 60 = persica.
decemdentata, Willd. Sp. Pl. iii. 124 = Leucas flaccida.
dentosa, *Franch. Pl. Davidian.* i. (1884) 243.—China.
diffusa, Herb. Madr. ex Wall. Cat. n. 2528 = Leucas dimidiata.
diffusa, Rottl. ex Hook. f. Fl. Brit. Ind. iv. 689 = Leucas diffusa.
dimidiata, Roth, Nov. Pl. Sp. 264 = Leucas dimidiata.
discolor, *Ledeb. Fl. Ross.* iii. 435.—Sibir.
elliptica, *Benth. Lab. Gen. et Sp.* 626.—Persia.
eriostoma, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 686 = Leucas eriostoma.

PHLOMIS:—
esculenta, Roxb. Hort. Beng. 44; Fl. Ind. iii. 10 = Leucas aspera.
ferruginea, Hort. ex Benth. in DC. Prod. xii. 540 = fruticosa.
ferruginea, Mill. Gard. Dict. ed. VIII. n. 12 = purpurea.
ferruginea, *Tenore, Prod. Fl. Neap.* 35; *Fl. Nap.* ii. 36. t. 57.—Ital.; Graecia.
flaccida, Steud. Nom. ed. I. 613 = Leucas flaccida.
flavescens, Mill. Gard. Dict. ed. VIII. n. 10 = aurea.
floccosa, *D. Don, in Bot. Reg.* t. 1300.—Afr. bor. or.
fruticosa, *Linn. Sp. Pl.* 584.—Europ. austr.
fruticosa, Sieber, ex Presl, Delic. Prag. 84 = ferruginea.
ghilanensis, *C. Koch, in Linnaea*, xxi. (1848) 698.—Persia.
ghilanica, Boiss. Fl. Orient. iv. 784 = praec.
glabrata, Vahl, Symb. Bot. i. 42 = Leucas glabrata.
glandulosa, Schenk, Pl. Sp. Aegypt. 20 = viscosa.
gracilis, Salisb. Prod. 84 = Leucas zeylanica.
Haussknechti, Bunge, Lab. Pers. 78 = capitata.
Herba-venti, *Linn. Sp. Pl.* 586.—Reg. Mediterr.
hirsuta, Wight, ex Wall. Cat. n. 2528 D = Leucas dimidiata.
hirta, Herb. Madr. ex Wall. l. c. 2053 B = Leucas helianthemifolia.
hirta, Heyne, ex Roth, Nov. Pl. Sp. 264 = Leucas hirta.
iberica, Vis. ex Boiss. Fl. Orient. iv. 793 = Eremostachys laciniata.
imbricata, Boiss. Fl. Orient. iv. 785 = lunarifolia.
indica, Herb. Madr. ex Wall. Cat. n. 2528 = Leucas dimidiata.
indica, Linn. Sp. Pl. 586 = Leucas indica, *R. Br.*
italica, *Linn. Syst.* ed. X. 1102; *Sm. Spic.* i. 6.—Ital.; Lusit.
laciniata, Linn. Sp. Pl. 585 = Eremostachys laciniata.
laciniata, Willd. ex Ledeb. Fl. Ross. iii. 440 = Eremostachys gymnocalyx.
lamiiflora, *Regel, in Act. Hort. Petrop.* vi. (1879) 374.—Turkestan.
lamiifolia, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 383 = bracteosa.
lanata, *Willd. Enum. Hort. Berol. Suppl.* 41.—Ital.; Creta.
lanceolata, Boiss. Diagn. Ser. I. v. 36 = armeniaca.
lanigera, Siev. in Pall. N. Nord. Beitr. vii. 295 = Eremostachys gymnocalyx.
latifolia, Mill. Gard. Dict. ed. VIII. n. 3 = fruticosa.
latifolia, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 383 = bracteosa.
Leonitis, Linn. Mant. i. 13 = Leonotis Leonitis.
Leonurus, Linn. Sp. Pl. 587 = Leonotis Leonurus.
linearis, *Boiss. & Bal. Diagn.* Ser. II. iv. 46.—As. Min.
linifolia, Roth, Nov. Pl. Sp. 260 = Leucas lavandulaefolia.
longifolia, Boiss. & Blanche, Diagn. Ser. II. iv. 47 = viscosa.
lunarifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 414.—As. Min.; Ins. Rhodos.
lunarifolia, Sibth. & Sm. l. c., partim = samia.
Lychnitis, *Linn. Sp. Pl.* 585.—Europ.
lycia, *D. Don, in Fellows, Discov. Lycia*, 293.—Lycia.
macrophylla, *Wall. Pl. As. Rar.* i. 62.—Reg. Himal.
major, Clos, in Bull. Soc. Bot. Fr. xv. (1868) 7 = *polymorpha*.
marrubioides, *Regel, in Act. Hort. Petrop.* vi. (1879) 375.—Turkestan.
martinicensis, Sw. Prod. Veg. Ind. Occ. 88 = Leucas martinicensis.
mauritanica, Munby, Fl. Algér. 60. t. 3 = crinita.
mesopotamica, *Boiss. Diagn.* Ser. II. iv. 47.—Mesopot.
micrantha, Burch. Trav. i. 340 = Stachys Burchellii.
microphylla, Sieber, Reise Kreta, i. 190; ii. 319. t. 8 = lanata.
mollis, Schum. & Thonn. Beskr. Guin. Pl. 263 = Leucas martinicensis.
moluccana, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 11 = Leucas flaccida.
moluccoides, Vahl, Symb. Bot. i. 42. t. 14 = Otostegia scariosa.
mongolica, *Turcz. in Bull. Soc. Nat. Mosc.* (1851) II. 406.—Mongolia.
montana, Koen. ex Roxb. Hort. Beng. [95]; Fl. Ind. iii. 11 = Leucas chinensis.
montana, Roth, Nov. Pl. Sp. 263 = Leucas montana.
montana, Rottl. ex Wall. Cat. n. 2524 = Leucas marrubioides.

PHLOMIS:—
nepetaefolia, Linn. Sp. Pl. 586 = Leonotis nepetaefolia.
nepetifolia, Linn. Syst. ed. XII. 398 = praec.
Nissolii, *Linn. Sp. Pl.* 585.—As. Min.; Syria.
nutans, Roth, Nov. Pl. Sp. 263 = Leucas nutans.
obliqua, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 690 = Leucas aspera.
oblongata, *Schrenk, Enum. Pl. Nov.* i. 29.—As. Min.
octodentata, Stokes, Bot. Mat. Med. iii. 355 = Leucas zeylanica.
ocymifolia, *Burm. f. Fl. Cap. Prod.* 16.—Afr. austr.
Olgae, *Regel, Pl. Nov. Fedsch.* 68.—Turkestan.
Olivieri, Benth. Lab. Gen. et Sp. 624 = orientalis.
oppositiflora, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 784.—Cataonia.
oreophila, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 426.—Sibir.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 9.—As. Min.
pallida, Schum. & Thonn. Beskr. Guin. Pl. 262 = Leonotis pallida.
parviflora, Benth. in Wall. Cat. n. 2066 = breviflora.
parvifolia, Burch. Trav. i. 225 = Stachys integrifolia.
parvifolia, Presl, Delic. Prag. 86 = lanata.
persica, *Boiss. Diagn.* Ser. I. v. 37.—Persia.
pilosa, Herb. Madr. ex Wall. Cat. n. 2526 = Leucas nepetaefolia.
pilosa, Roxb. Hort. Beng. 44; Fl. Ind. iii. 12 = Leucas pilosa.
Pluckenetii, Roth, Nov. Pl. Sp. 261 = Leucas aspera.
polymorpha, Clos, in Bull. Soc. Bot. Fr. xv. (1868) 7 = viscosa, ferruginea.
Portae, Kern. ex Nym. Consp. 581 = fruticosa.
pratensis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 426.—Sibir.
pungens, Willd. Sp. Pl. iii. 121 = P. Herba-venti.
purpurea, *Linn. Sp. Pl.* 585.—Hispan.; Lusit.
purpurea, Linn. l. c. ed. II. 818 = italica.
repens, Willd. ex Benth. in Linnaea, xi. (1837) 338 = Leucas decumbens.
repetifolia, Roxb. Fl. Ind. iii. 8, err. typ. = *nepetaefolia*.
rigida, *Labill. Ic. Pl. Syr.* iii. 15 t. 10.—As. Min.; Syria.
Rodontia, Buch.-Ham. ex Wall. Cat. n. 2525 = Leucas montana.
rotata, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 694.—Reg. Himal.
rotundifolia, Mill. Gard. Dict. ed. VIII. n. 13 = italica.
rugosa, *Benth. in Wall. Pl. As. Rar.* i. 63; *Cat.* n. 2067.—Reg. Himal.; Malaya.
Russeliana, Lag. ex Benth. Lab. Gen. et Sp. 629 = viscosa.
sagittata, *Regel, in Act. Hort. Petrop.* vi. (1879) 373.—Turkestan.
salicifolia, *Regel, l. c.* 371.—Turkestan.
salviaefolia, Jacq. Hort. Schoenb. iii. 58 = purpurea.
salvifolia, Stokes, Bot. Mat. Med. iii. 354 = fruticosa.
samia, *Linn. Sp. Pl.* 585.—Graec.; As. Min.
scariosa, Presl, Delic. Prag. 85 = fruticosa.
setigera, *Falc. ex Benth. in DC. Prod.* xii. 543.—Reg. Himal.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 372.—Turkestan.
sibirica, Medic. Beobacht. (1783) 124 = Leonurus sibiricus.
simplex, Royle, ex Benth. in Hook. Bot. Misc. iii. (1833) 382 = bracteosa.
sinensis, J. F. Gmel. Syst. 910 = *chinensis*.
speciosa, Salisb. Prod. 84 = Leonotis Leonurus.
spectabilis, *Falc. ex Benth. in DC. Prod.* xii. 543.—Reg. Himal.
Spica-venti, C. Koch, in Linnaea, xxi. (1848) 699, sphalm. = P. Herba-venti.
spinosa, *Vand. Fasc. Pl.* 15.—Bras.
Stewartii, *Hook. f. Fl. Brit. Ind.* iv. 692.—Ind. bor. occ.
stricta, Heyne, ex Hook. f. l. c. 688 = Leucas stricta.
superba, C. Koch, in Linnaea, xxi. (1848) 699 = samia.
syriaca, Boiss. Diagn. Ser. I. xii. 89 = Nissolii.
tarsensis, Boiss. Fl. Orient. iv. 781 = Nissolii.
taurica, *Hartwiss, ex Bunge, Lab. Pers.* 77, in obs.—Persia.
thapsoides, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 440.—Turkestan.
tomentosa, Herb. Madr. ex Wall. Cat. n. 2524 = Leucas marrubioides.
tomentosa, *Regel, Pl. Nov. Fedesch.* 69.—Turkestan.
tuberosa, *Linn. Sp. Pl.* 586.—Europ. austr.; As. or. et bor.

PHLOMIS:—
turkestana, *Bunge, Lab. Pers.* 78, nomen.—Turkestan.
umbrosa, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 76.—China.
urticifolia, Vahl, Symb. Bot. iii. 76 = Leucas urticaefolia.
ventosa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70, 131 = P. Herba-venti.
virens, DC. Cat. Hort. Monsp. 132 = viscosa.
viscosa, *Poir. Encyc.* v. 271.—Syria.
zeylanica, Blanco, Fl. Filip. ed. I. 475 = Leucas aspera.
zeylanica, Heyne, ex Wall. Cat. n. 2522 = Pluckenetii.
zeylanica, Linn. Sp. Pl. 586 = Leucas zeylanica.
zeylanica, Roxb. Hort. Beng. 44; Fl. Ind. iii. 9 = Leucas lavandulaefolia.

PHLOMITIS, Reichb. ex T. Nees, Gen. Fl. Germ. Gamop. ii. Trib. VII. n. 43 (1843) = **Phlomis**, Linn. (Labiat.).
pungens, Reichb. ex T. Nees, l. c. = Phlomis Herba-venti.
tuberosa, Reichb. ex T. Nees, l. c. = Phlomis tuberosa.

PHLOMOIDES, Moench, Meth. 403 (1794) = **Phlomis**, Linn. (Labiat.).
tuberosa, Moench, l. c. = Phlomis tuberosa.

PHLOMOSTACHYS, C. Koch, Ind. Sem. Hort. Berol. (1856) App. 3; Beer, Bromel. 45 (1857) = **Pitcairnia**, L'Hérit. (Bromel.).
Altensteinii, Beer, Bromel. 45 = Pitc. Altensteinii.
atrorubens, C. Koch, Ind. Sem. Hort. Berol. (1856) App. 3 = Pitc. atrorubens.
densiflora, Beer, Bromel. 46 = Pitc. densiflora.
Funkiana, Beer, l. c. 47 = Pitc. Funkiana.
gigantea, C. Koch, Ind. Sem. Hort. Berol. (1856) App. 3 = Pitc. Altensteinii.
imbricata, Beer, Bromel. 47 = Pitc. imbricata.
sulphurea, Beer, l. c. 46 = Pitc. Wendlandi.

PHLOX, Linn. Gen. ed. I. 52 (1737). *POLEMONIACEAE*, Benth. & Hook. f. ii. 821.
FONNA, Adans. Fam. ii. 214 (1763).
acuminata, Pursh, Fl. Am. Sept. ii. 730 = paniculata.
acutifolia, Sweet, Hort. Brit. ed. II. 368 = praec.?
adsurgens, *Torr. ex A. Gray, in Proc. Am. Acad.* viii. (1870) 256.—Am. bor. occ.
alba, Moench, Meth. Suppl. 173 = maculata.
altissima, Moench, l. c. 454 = glaberrima.
americana, Hort. ex Sweet, Brit. Flow. Gard. Ser. I. sub t. 248 = paniculata.
amoena, *Sims, Bot. Mag.* t. 1308.—Am. bor.
amplexicaulis, Rafin. Fl. Ludov. 36 (Quid?).—Am. bor.
aristata, Lodd. Bot. Cab. t. 1731 = subulata.
aristata, Michx. Fl. Bor. Am. i. 144 = pilosa.
atrocaulis ×, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 656.
bifida, *Beck, in Am. Journ. Sc.* Ser. I. xi. (1826) 170.—Am. bor.
biflora, *Ruiz & Pav. Fl. Per.* ii. 17.—Chili.
bimaculata, Hort. ex Sweet, Brit. Flow. Gard. Ser. I. sub t. 224 = odorata.
brevifolia, *Baum. ex Hoffmgg. Verz. Pfl.* 98 (Quid?).—Am. bor.?
Bridgesii ×, Hort. ex Marnock, Floric. Mag. (1842) t. 1.
Broughtoni ×, Hort. l. c. t. 3.
Brownii ×, Salm-Dyck, Hort. Dyck. 205.
bryoides, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 153.—Am. bor. occ.
caespitosa, *Nutt. l. c.* vii. (1834) 41.—Am. bor. occ.
Caldriana, Court. ex Steud. Nom. ed. II. ii. 322 (Quid?).—Hab.?
canadensis, Sweet, Brit. Flow. Gard. Ser. I. t. 221 = divaricata.
candida, Pers. Syn. i. 186 = maculata.
canescens, *Salm-Dyck, Hort. Dyck.* 205.—Hab.?
canescens, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 122.—Am. bor. occ.
carnea, Sims, Bot. Mag. t. 2155 = glaberrima.
carolina, Linn. Sp. Pl. ed. II. 216 = ovata.
carolina, Sims, Bot. Mag. t. 1344 = glaberrima.
carolina, Sweet, Brit. Flow. Gard. Ser. I. t. 190 = floridana.
carolina, Walt. Fl. Carol. 96 = maculata.
caroliniana, Hill, Veg. Syst. viii. 32 = ovata?

PHLOX:—
clarkioides, *Poit. in Rev. Hortic.* Sér. II. i. (1841-43) 69.—Hab.?
Coldryana ×, Paxt. Mag. Bot. vii. (1840) 197.
cordata, Ell. Sketch, ii. 40 = paniculata.
corymbosa, Hort. ex Sweet, Hort. Brit. ed. II. 368; et Brit. Flow. Gard. Ser. II. t. 114 = paniculata.
crassifolia, Lodd. Bot. Cab. t. 1596 = reptans.
cruenta, Curt. ex Steud. Nom. ed. II. 322 (an hybr.?).
cuspidata, Scheele, in Linnaea, xxiii. (1850) 139 = pilosa.
decussata, Lyon, ex Pursh, Fl. Am. Sept. ii. 730 = paniculata.
diffusa, Benth. Pl. Hartw. 325 = Douglasii.
disticha, Sabine, ex Sweet, Hort. Brit. ed. III. 478 (Quid?).—Am. bor.
divaricata, Durand, in Journ. Acad. Philad. N. S. iii. (1855) 97 = speciosa.
divaricata, *Linn. Sp. Pl.* 152.—Am. bor.
divergens, Wender. ex Steud. Nom. ed. II. ii. 322 (Quid?).—Hab.?
Douglassii, *Hook. Fl. Bor. Am.* ii. 73.—Am. bor. occ.
Drummondii, *Hook. Bot. Mag.* t. 3441.—Texas.
elata, *Penny, ex G. Don, in Loud. Hort. Brit. Suppl.* i. 596, nomen.—Am. bor.
excelsa, Penny, ex G. Don, in Sweet, Hort. Brit. ed. II. 368 (Hybr.?).—Am. bor.
floridana, *Benth. in DC. Prod.* ix. 304.—Am. bor.
frondosa, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 683 = subulata.
fruticosa, Hort. ex Steud. Nom. ed. II. ii. 322 = maculata?
glaberrima, *Linn. Sp. Pl.* 152.—Am. bor.
glomerata, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 196, nomen.—Am. bor.
glutinosa, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 177 = divaricata.
Hentzii, Nutt. in Journ. Acad. Philad. vii. (1834) 110 = subulata.
Heynholdii, Hort.—Cf. Gard. Chron. (1871) 1041 = Drummondii?
Hoodii, *Richards. in Frankl. Journ. App.* ed. II. 6. t. 28.—Am. bor. occ.
Hookeri, Dougl. ex Hook. Fl. Bor. Am. ii. 73 = Gilia pungens.
humilis, Dougl. ex Hook. l. c. 72 = longifolia.
hybrida ×, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen.—Am. bor.
Ingramiana ×, Hort. ex Loud. Hort. Brit. Suppl. ii. 656.
intermedia, *Lodd. ex Sweet, Hort. Brit.* ed. I. 285, nomen.—Am. bor.
involucrata, Nutt. ex Loud. Hort. Brit. Suppl. i. 596 (Quid?).—Am. bor.
involucrata, Wood, Class-book (1861), 568 = amoena.
laeta ×, Penny, ex G. Don, in Sweet, Hort. Brit. ed. III. 478, nomen.—Am. bor.
latifolia, Michx. Fl. Bor. Am. i. 143 = ovata.
Leopoldiana ×, Paxt. Mag. Bot. xiv. (1848) t. 266.
linearifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 133.—Am. bor. occ.
linearis, Cav. Ic. vi. 17 = Collomia coccinea.
Listoniana, *Sweet, Hort. Brit.* ed. II. 368; *Forbes, Hort. Woburn.* 37.—Am bor.
longiflora, Sweet, Brit. Flow. Gard. Ser. II. t. 31 = maculata.
longifolia, *Nutt. in Journ. Acad. Philad.* vii. (1834) 41.—Am. bor. occ.
macrantha, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 5 = Roemeriana.
macrophylla ×, Court. ex Steud. Nom. ed. II. ii. 323.
maculata, *Linn. Sp. Pl.* 152.—Am. bor.
melampyrifolia, Salisb. Prod. 123 = glaberrima.
missourica, Salm-Dyck, Hort. Dyck. 205 (Quid?).—Am. bor.
montana, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen.—Am. bor.
muscoides, *Nutt. in Journ. Acad. Philad.* vii. (1834) 42. t. 6. f. 2.—Am. bor. occ.
nana, *Nutt. l. c.* N. S. i. (1847) 153.—Am. bor. occ.
nitida, Pursh, Fl. Am. Sept. ii. 730 = glaberrima.
nivalis, Lodd. Bot. Cab. t. 780 = subulata.
Nuttalliana, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 767 = floridana.
Nuttallii, Court. ex Steud. Nom. ed. II. ii. 323 = praec.
obovata, Muhl. ex Roem. & Schult. Syst. iv. 362 = reptans.

PHLOX:—
occidentalis, Durand, ex Torr. in Pacif. Rail. Rep. iv. 125 = speciosa.
odorata, *Sweet, Brit. Flow. Gard.* Ser. I. t. 224.—Am. bor.
Oldryana, Walp. Rep. vi. 526, sphalm. = *Coldryana*.
omniflora ×, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 656.
ovata, *Linn. Sp. Pl.* 152.—Am. bor.
paniculata, *Linn. l. c.* 151.—Am. bor.
penduliflora, Sweet, Hort. Brit. ed. II. 368; Brit. Flow. Gard. Ser. II. t. 46 = maculata.
philadelphica, Hort. ex Steud. Nom. ed. II. ii. 323 (Quid?).—Am. bor.
pilosa, *Linn. Sp. Pl.* 152.—Am. bor.
pilosa, Walt. Fl. Carol. 96 = amoena.
pinnata, Cav. Ic. vi. 17 = Collomia Cavanillesiana.
procumbens, A. Gray, Man. Bot. N. U. St. ed. V. 372 = amoena.
procumbens ×, Lehm. Sem. Hort. Hamb. (1828) 17.
prostrata, Ait. ex Sweet, Hort. Brit. ed. I. 285 = reptans.
pulchella ×, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 656.
pyramidalis, Sm. Exot. Bot. ii. 55. t. 87 = maculata.
reflexa, Hort. ex Sweet, Brit. Flow. Gard. Ser. I. t. 232 = maculata.
reptans, *Michx. Fl. Bor. Am.* i. 145.—Am. bor.
revoluta, Aikin, ex A. Gray, Syn. Fl. N. Am. ii. 1. 130 = glaberrima.
Richardsoni, *Hook. Fl. Bor. Am.* ii. 73. t. 160.—Am. arct.
rigida, Benth. in DC. Prod. ix. 306 = caespitosa.
Roemeriana, *Scheele, in Linnaea*, xxi. (1848) 752.—Texas.
rosea, *Forbes, Hort. Woburn.* 38 (Quid?).—Am. bor.
Sabini, Dougl. in Hook. Fl. Bor. Am. ii. 72 = speciosa.
scabra, Sweet, Brit. Flow. Gard. Ser. I. t. 248 = paniculata.
setacea, Linn. Sp. Pl. 153 = subulata.
Shepherdii ×, Penny, ex Sweet, Hort. Brit. ed. II. 368.
sibirica, *Linn. Sp. Pl.* 153.—As. or.; Am. bor. arct.
Sickmanni, Lehm. Ind. Sem. Hort. Hamb. (1826) 17 = paniculata.
Siebmanni, Benth. in DC. Prod. ix. 303, sphalm. = praec.
speciosa, Lindl. in Bot. Reg. t. 1351 = linearifolia.
speciosa, *Pursh, Fl. Am. Sept.* i. 149.—Am. bor. occ.
Stellaria, *A. Gray, in Proc. Am. Acad.* viii. (1870) 252.—Am. bor.
stolonifera, Sims, Bot. Mag. t. 563 = reptans.
suaveolens, Ait. Hort. Kew. ed. I. i. 205 = maculata.
subulata, *Linn. Sp. Pl.* 152.—Am. bor.
suffruticosa, *Vent. Jard. Malm.* ii. 107.—Am. bor.
suffruticosa, Willd. Enum. Hort. Berol. 200 = glaberrima.
tardiflora, Penny, ex Loud. Hort. Brit. Suppl. i. 596 = maculata.
Thomsoni, *Court. ex Steud. Nom.* ed. II. ii. 323 (Quid?).—Am. bor.
tigrina, *Wender. ex Steud. l. c.* (Quid?).—Hab.?
triflora, Michx. Fl. Bor. Am. i. 143 = glaberrima?
triflora, Sweet, Brit. Flow. Gard. Ser. I. t. 29 = ovata.
triovulata, Thurb. ex Torr. in Bot. Mex. Bound. 145 = nana.
undulata, Ait. Hort. Kew. ed. I. i. 265; Moench, Meth. 454 = paniculata.
unidentata, Bert. ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 126. t. 39 = Collomia coccinea.
uniflora, Rafin. Fl. Ludov. 47 (Quid?).—Am. bor.
verna ×, Forbes, Hort. Woburn. 38.
vernalis, Salisb. Prod. 123 = divaricata.
Vernoniana ×, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 656.
virginica, *Lodd. ex Sweet, Hort. Brit.* ed. I. 285, nomen.—Am. bor.
Walteri, Chapm. Fl. S. U. St. 339 = amoena.
Wheeleriana ×, Sweet, Hort. Brit. ed. II. 368.

PHLOXUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = praec.

PHLYARODOXA, S. Moore, in Journ. Bot. xiii. (1875) 229 = **Ligustrum**, Tourn. (Oleac.).
leucantha, S. Moore, l. c. = L. molliculum.

PHOBEROS, Lour. Fl. Cochinch. 317 (1790) = **Scolopia**, Schreb. (Bixin.).
acuminatus, Gardn. ex Thw. Enum. Pl. Zeyl. 17, 400 = S. crenata.
Arnottiana, Thw. l. c. 16 = S. crenata.
Brownii, F. Muell. Fragm. iii. 11 = S. Brownii.
chinensis, Lour. Fl. Cochinch. 318 = S. crenata.
chinensis, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 334.—China.
cochinchinensis, Lour. Fl. Cochinch. 318 = S. cochinchinensis.
crenatus, Wight & Arn. Prod. 29 = S. crenata.
Dasyanthera, Benn. Pl. Jav. Rar. 192 = S. Dasyanthera.
Eckloni, Presl, Bot. Bemerk. 70; Harv. & Sond. Fl. Cap. i. 67.—Afr. austr.
Gaertneri, Thw. Enum. Pl. Zeyl. 17 = S. Gaertneri.
Hookerianus, Wight, ex Thw. l. c. = S. crenata.
lanceolatus, Wight & Arn. Prod. 30 = S. crenata.
macrophyllus, Wight & Arn. l. c. = S. Rhinanthera.
maritima, Miq. Fl. Ind. Bat. Suppl. 388.—Sumatra.
Mundii, Presl, Bot. Bemerk. 70 = seq.
Mundtii, Harv. & Sond. Fl. Cap. i. 67.—Afr. austr.
Rhinanthera, Benn. Pl. Jav. Rar. 187 = S. Rhinanthera.
Roxburghii, Benn. l. c. 192 = S. Roxburghii.
saevus, Hance, in Walp. Ann. iii. 825 = S. crenata.
Wightianus, Wight & Arn. Prod. 30 = S. crenata.
Zeyheri, Presl, Bot. Bemerk. 70; Harv. & Sond. Fl. Cap. i. 67.—Afr. austr.

PHOCEA, Seem. in Journ. Bot. viii. (1870) 68. *CELASTRINEAE*.
Andersonii, *Seem. l. c.*—N. Caled.

PHOEBE, Nees, Syst. Laurin. 98 (1836). *LAURINEAE* [Benth. & Hook. f. iii. 157].
angustifolia, *Meissn. in DC. Prod.* xv. I. 34.—Reg. Himal.
antillana, *Meissn. l. c.* 31.—Ind. occ.
attenuata, *Nees, Syst. Laurin.* 104.—Reg. Himal.
attenuata, Miq. ex Meissn. in DC. Prod. xv. I. 36 = opaca.
barbusana, *Webb & Berth. Phyt. Canar.* iii. 223.—Ins. Canar.
canescens, *Miq. Fl. Ind. Bat.* i. I. 912.—Sumatra.
chartacea, *Miq. l. c.*—Java.
cinnamomifolia, *Nees, in Linnaea*, xxi. (1848) 488.—Am. trop.
cinnamomifolia, Rich. ex Griseb. Cat. Pl. Cub. 111 = triplinervis.
cubensis, *Nees, Syst. Laurin.* 120.—Cuba.
cuneata, *Blume, Mus. Bot. Lugd. Bat.* i. 326.—Java.
cuspidata, *Blume, l. c.* 328.—Java.
declinata, *Nees, Syst. Laurin.* 114.—Malaya.
densiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Java.
effusa, *Meissn. in DC. Prod.* xv. I. 33.—Mexic.
elliptica, Blume, Mus. Bot. Lugd. Bat. i. 326 = macrophylla.
elongata, *Nees, Syst. Laurin.* 116.—Am. trop.
elongata, Nees, l. c., partim = montana.
excelsa, *Nees, l. c.* 102.—Java.
falcata, *Miq. Fl. Ind. Bat.* i. I. 912.—Sumatra.
glaucescens, Nees, Syst. Laurin. 100; Meissn. in DC. Prod. xv. I. 36 = pallida.
granatensis, Meissn. in DC. l. c. 32 = cinnamomifolia.
grandifolia, *Miq. Fl. Ind. Bat.* i. I. 913.—Malaya.
Hartwegii, *Meissn. in DC. Prod.* xv. I. 30.—Mexic.
holosericea, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Sumatra.
hypoleuca, A. Rich. Fl. Cub. Fanerog. ii. 185 = Persea cubensis.
impressa, *Meissn. in DC. Prod.* xv. I. 33.—Guian.
incerta, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Java.
insularis, *Meissn. in DC. Prod.* xv. I. 33.—Ins. Cocos.
javanica, *Meissn. l. c.* 37.—Java.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 359.—Sumatra.
lanceolata, *Nees, Syst. Laurin.* 109.—Ind. or.; Burma.
latifolia, Champ. in Hook. Kew Journ. v. (1853) 197 = Cinnamomum Parthenoxylon.
leiophylla, *Miq. Fl. Ind. Bat.* i. I. 911.—Ins. Celebes; Borneo.
ligustrina, Nees, Syst. Laurin. 113, partim = declinata.
Lingue, Steud. Nom. ed. II. ii. 16, in syn., sphalm. = Persea Lingue.
lucida, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Malaya.
macrophylla, *Blume, l. c.* 326.—Java.
malaccensis, Meissn. in DC. Prod. xv. I. 37 = opaca.
maynensis, Nees, Syst. Laurin. 118 = peruviana.

PHOEBE:—
membranacea, Macfad. ex Griseb. Fl. Brit. W. Ind. 283 = Ocotea Leucoxylon.
membranacea, Nees, Syst. Laurin. 118 = cubensis.
mexicana, *Meissn. in DC. Prod.* xv. I. 31.—Mexic.
montana, *Griseb. in Mem. Acad.* N. S. viii. (1861) 187.—Ind. occ.
multiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 326.—Java.
nitida, *Blume, l. c.* 328.—Sumatra.
obtusa, *Blume, ex Meissn. in DC. Prod.* xv. I. 35.—Java.
opaca, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Malaya.
opaca, Zoll. Syst. Verz. Ind. Arch. ii. 113 = javanica.
pallida, *Nees, Syst. Laurin.* 112.—Ind. or.
paniculata, *Nees, l. c.* 105.—Ind. or.
parviflora, *Blume, Mus. Bot. Lugd. Bat.* i. 327.—Java.
parviflora, Nees, Syst. Laurin. 120, in nota = triplinervis.
patens, *Nees, l. c.* 117.—Jamaic.
pedunculosa, *Meissn. in DC. Prod.* xv. I. 47.—Bras.
peruviana, *Meissn. l. c.* 32.—Peruv.
Poeppigii, *Meissn. l. c.* 33.—Peruv.
Poiretiana, Nees, Syst. Laurin. 119 = montana.
pubescens, Nees, l. c. 107 = paniculata.
rigida, *Miq. Fl. Ind. Bat.* i. I. 910.—Sumatra.
Ruiziana, *Nees, in Linnaea*, xxi. (1848) 489.—Peruv.
salicifolia, *Nees, l. c.* 488.—Mexic.
Sellowii, *Meissn. in DC. Prod.* xv. I. 30.—Bras.
sericea, Nees, Syst. Laurin. 99 = Machilus sericea.
spathulata, *Miq. Fl. Ind. Bat.* i. I. 911.—Java.
stereophylla, *Meissn. in DC. Prod.* xv. I. 29.—Bras.
sumatrana, *Miq. Pl. Jungh.* 175.—Guian.
superba, *Miq. Fl. Ind. Bat.* i. I. 912.—Malaya.
tomentosa, *Meissn. in DC. Prod.* xv. I. 30.—Bras.
triplinervis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 187.—Cuba.
umbelliflora, *Nees, Syst. Laurin.* 115.—Malaya.
Valenzuelana, A. Rich. Fl. Cub. Fanerog. ii. 185 = elongata.
villosa, Wight, Ic. t. 1822 (Machili sp.).—Reg. Himal.
Wightii, Meissn. in DC. Prod. xv. I. 38 = paniculata.

PHOENICANTHEMUM, Blume, in Schult. f. Syst. vii. 1729 (1830) = **Loranthus**, Linn.
Bennettianum, Miq. Fl. Ind. Bat. i. I. 826 = L. coccineus.
coccineum, Blume, in Schult. f. Syst. vii. 1729 = L. coccineus.
cylindricum, Miq. Fl. Ind. Bat. i. I. 825 = L. cylindricus.
longissimum, Miq. Fl. Ind. Bat. Suppl. 345 = I. setigerus ?
Lyndenianum, Miq. Fl. Ind. Bat. i. I. 824 = L. Lyndenianus.
Macklottianum, Miq. l. c. = L. Macklottianus.
pentapetalum, Blume, in Schult. f. Syst. vii. 1729 = L. pentapetalus.
setigerum, Miq. Fl. Ind. Bat. i. I. 826 = L. setigerus.
trianthum, Miq. l. c. = L. trianthus.
triste, Miq. l. c. = L. tristis.
Wallichianum, Blume, in Schult. f. Syst. vii. 1729 = L. Wallichianus.
xestophyllum, Miq. Fl. Ind. Bat. i. I. 823 = L. macrostachys, *Korth.*

PHOENICAULIS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 89 (1838) = **Cheiranthus**, Linn. (Crucif.).
cheiranthoides, Nutt. l. c. = C. Menziesii.
Menziesii, Steud. Nom. ed. II. ii. 323 = C. Menziesii.

PHOENICOCISSUS, Mart. ex Meissn. Gen. Comm. 209 (1840) = **Lundia**, DC. (Bignon.).

PHOENICOPHORIUM, H. Wendl. in Illustr. Hortic. xii. (1865) Misc. 5. t. 433 = **Stevensonia**, Dunc. (Palm.).
Sechellarum, H. Wendl. l. c. = S. grandifolia.
viridifolium, Hort. ex H. Wendl. in Kerch. Palm. 252 = Roscheria melanochaetes.

PHOENICOPUS, Spach, Hist. Vég. Phan. x. 6 (1841) = PHAENOPUS, DC. = **Lactuca**, Tourn. (Compos.).
intricatus, Pomel, Nouv. Mat. Fl. Atl. 264 = L. intricata.

PHOENICOSPERMUM, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 68. t. 3 (1865-66). *TILIACEAE*, Benth. & Hook. f. i. 987.
javanicum, *Miq. l. c.*—Java.

PHOENIX, Linn. Syst. ed. I (1735). *PALMAE*, Benth. & Hook. f. iii. 921.
DACHEB, Adans. Fam. ii. 25 (1763).
ELATE, Linn. Mus. Cliff. 12 (1737).
FULCHIRONIA, Lesch. ex Desf. Cat. Hort. Par. ed. III. 29 (1829).
PHAENIX, Hill, Hort. Kew. 449 (1768).
PHONIPHORA, Neck. Elem. iii. 302 (1790).
ZELONOPS, Rafin. Fl. Tellur. ii. 102 (1836).
acaulis, *Buch.-Ham. ex Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 783.—Ind. or.; Burma.
aequinoctialis, Hort.—Cf. Rev. Hortic. (1879) 340 (Quid ?).—Hab. ?
Ammonis, *Ehrenb. ex Steud. Nom.* ed. II. ii. 323.—Hab. ?
canariensis, *Hort. ex Chabaud, in La Provence Agric.* n. 19. (Oct. 1882) 293, ff. 66-68.—Ins. Canar.
cycadifolia, Hort. Athen. ex Regel, Gartenfl. xxviii. (1879) 131. t. 974 = dactylifera.
dactylifera, *Linn. Sp. Pl.* 1188.—Afr. bor.; Arab.
equinoxialis, Boj. Hort. Maurit. 306 = reclinata.
excelsa, Steud. Nom. ed. II. ii. 323, sphalm. = seq.
excelsior, Cav. Ic. ii. 13. t. 125 = dactylifera.
farinifera, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 87, partim = humilis.
farinifera, *Roxb. Pl. Corom.* i. 55. t. 74.—Ind. or.
ferruginea, Hort. ex H. Wendl. Ind. Palm. 31 (Quid ?).—Hab. ?
Hanceana, Hort. ex Wendl. in Kerch. Palm. 252 = humilis.
humilis, Cav. Ic. ii. 12. t. 115 = Chamaerops humilis.
humilis, *Royle, Illustr. Bot. Himal.* 394.—Ind. or.; Burma; China.
hybrida, Hort. Bog. ex Teijsm. & Binn. Cat. Hort. Bog. 72 (Quid ?).—Hab ?
Jubae, Webb, ex Christ, in Engl. Bot. Jahrb. vi. (1885) 469 = canariensis.
leonensis, Lodd. ex Kunth, Enum. Pl. iii. 257 = reclinata.
Loureiri, Kunth, l. c. = humilis.
Ouseleyana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 347 = humilis.
paludosa, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 789.—Ind. or.; Siam.
pedunculata, Griff. Palms Brit. Ind. 139 = humilis.
pumila, Hort.—Cf. Regel, Gartenfl. xx. (1871) 153 (Quid ?).—Hab. ?
pusilla, *Gaertn. Fruct.* i. 24. t. 9.—Ind. or.
pusilla, Lour. Fl. Cochinch. 614 = humilis.
pygmaea, Hort. ex Link, Enum. Hort. Berol. ii. 421 (an seq. ?).—Hab. ?
pygmaea, Raeusch. Nom. ed. III. 375 = farinifera.
reclinata, *Jacq. Fragm.* i. 27. t. 24.—Afr. trop. et austr.
rupicola, *T. Anders. in Journ. Linn. Soc.* xi. (1871) 13.—Reg. Himal.
senegalensis, Van Houtte, ex Salomon, in Regel, Gartenfl. (1882) 305 = reclinata.
siamensis, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. 5. p. 14 = paludosa.
spadicea, Wight, ex Mart. Hist. Nat. Palm. iii. 274, in obs. = farinifera.
spinosa, Schum. & Thonn. Beskr. Guin. Pl. 487 = reclinata.
sylvestris, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 787.—Ind. or.
sylvestris, Thw. Enum. Pl. Zeyl. 329 = pusilla.
tenuis, Versch. Cat. (1869) 13, cum ic. = canariensis.
Vigieri, Hort. ex Naud. in Rev. Hortic. (1885) 541, in nota = canariensis.
zeylanica, Trimen, in Journ. Bot. xxiii. (1885) 267 = pusilla.

PHOENIXOPUS, Reichb. Consp. 98 (1828) = **Lactuca**, Tourn. (Compos.).

PHOENOCOMA, G. Don, in Loud. Hort. Brit. 342 (1830) = **Phaenocoma**, D. Don (Compos.).

PHOENOPUS, Nym. Consp. 435 (1879) = PHAENOPUS, DC. = **Lactuca**, Tourn. (Compos.).

PHOLACILIA, Griseb. Fl. Brit. W. Ind. 129 (1859) = **Trichilia**, P. Br. (Meliac.).
diversifolia, Griseb. l. c. 130 = T. simplicifolia.
pteleaefolia, Griseb. l. c., in obs. = T. pteleaefolia.
trinitensis, Griseb. l. c. = T. trinitensis.

PHOLIDANDRA, Neck. Elem. i. 348 (1790) = **Raputia**, Aubl. (Rutac.).

PHOLIDIA, R. Br. Prod. 517 (1810). *MYOPORINEAE*, Benth. & Hook. f. ii. 1124.
DUTTONIA, F. Muell. in Hook. Kew Journ. viii. (1856) 73. t. 1.
PHOLIDIOPSIS, F. Muell. in Linnaea, xxv. (1852) 429.
PSEUDOPHOLIDIA, A. DC. in DC. Prod. xi. 704 (1847).
SENTIS, F. Muell. Fragm. iv. 47 (1863).
adenotricha, *F. Muell. ex Benth. Fl. Austral.* v. 14.—Austral.
Behriana, *F. Muell. in Linnaea*, xxv. (1852) 430.—Austral.
brevifolia, *Benth. Fl. Austral.* v. 12.—Austral.
crassifolia, *F. Muell. in Linnaea*, xxv. (1852) 430.—Austral.
Dalyana, *F. Muell. ex Benth. Fl. Austral.* v. 10.—Austral.
Delisserii, *F. Muell. l. c.* 11.—Austral.
densifolia, *F. Muell. l. c.* 13.—Austral.
divaricata, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 47.—Austral.
gibbifolia, *F. Muell. ex Benth. Fl. Austral.* v. 13.—Austral.
imbricata, *Benth. l. c.*—Austral.
microtheca, *F. Muell. ex Benth. l. c.* 14.—Austral.
polyclada, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 47 = Eremophila polyclada.
resinosa, *Endl. in Ann. Mus. Wien*, ii. (1838) 203.—Austral.
santalina, *F. Muell. ex Benth. Fl. Austral.* v. 15.—Austral.
scoparia, *R. Br. Prod.* 517.—Austral.
Woollsiana, *F. Muell. ex Benth. Fl. Austral.* v. 12.—Austral.

PHOLIDIOPSIS, F. Muell. in Linnaea, xxv. (1852) 429 = **Pholidia**, R. Br. (Myopor.).
santalina, F. Muell. l. c. = Pholidia santalina.

PHOLIDOCARPUS, Blume, in Schult. f. Syst. vii. 1308 (1830). *PALMAE*? Benth. & Hook. f. iii. 883.
Ihur, *Blume, Rumphia*, iii. 90.—Malaya.
Ihur, Miq. Fl. Ind. Bat. iii. 47 = Majadum.
Majadum, *Becc. Malesia*, i. (1877) 80.—Malaya.
tunicatus, *H. Wendl. in Kerch. Palm.* 235.—Cochinch.

PHOLIDOPHYLLUM, Vis. Sem. Hort. Patav. (1847) 4; et in Otto & Dietr. Allg. Gartenz. xvi. (1851) 30 = **Cryptanthus**, Otto & Dietr. (Bromel.).
zonatum, Vis. ll. cc. = C. zonatus.

PHOLIDOSTACHYS, H. Wendl. ex Benth. & Hook. f. Gen. iii. 915, in nota (1883). *PALMAE*, Benth. & Hook. f. l. c.
pulchra, *H. Wendl. l. c.*—Costa Rica.

PHOLIDOTA, Lindl. in Hook. Exot. Fl. ii. t. 138 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 520.
ACANTHOGLOSSUM, Blume, Bijdr. 381 (1825).
CRINONIA, Blume, l. c. 338 (1825).
PTILOCNEMA, D. Don, Prod. Fl. Nep. 33 (1825).
aculeata, Bot. Reg. ex Steud. Nom. ed. II. ii. 324 (Quid?).—Ind. or.
advena, Reichb. f. Otia Hamb. i. 47, in syn. = Coelogyne advena.
articulata, *Lindl. in Wall. Cat.* n. 1992.—Reg. Himal.
calceata, *Reichb. f. in Bonplandia*, iv. (1856) 329.—Reg. Himal.
camelostalix, *Reichb. f. Xenia Orch.* i. 217. t. 89.—Java.
carnea, *Lindl. Gen. et Sp. Orch.* 37.—Java.
chinensis, *Lindl. in Journ. Hort. Soc.* ii. (1847) 308.—China.
clypeata, *Lindl. l. c.* v. (1850) 37.—Borneo.
conchoidea, *Lindl. Bot. Reg.* (1840) *Misc.* 84.—Ins. Philipp.
crotalina, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 218.—Java.
gibbosa, *Lindl. ex De Vriese, Illustr. Orch.* [13]. t. 10.—Malaya?
globosa, *Lindl. Gen. et Sp. Orch.* 37.—Java.
imbricata, *Lindl. in Hook. Exot. Fl.* ii. t. 138.—Ind. or.
khasiyana, Reichb. f. in Bonplandia, iv. (1856) 329 = articulata.
loricata, *Reichb. f. l. c.* v. (1857) 43.—Java.
membranacea, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1858) 488.—Java.
nervosa, *Reichb. f. in Bonplandia*, v. (1857) 43.—Java.
pallida, Lindl. Bot. Reg. sub t. 1777 = imbricata.
Pholas, *Reichb. f. in Walp. Ann.* vi. 237.—Ins. Hongkong.
recurva, *Lindl. Gen. et Sp. Orch.* 37.—Reg. Himal.
rhombophora, *Reichb. f. in Linnaea*, xli. (1877) 117, in nota.—Ins. Philipp.
rubra, *Lindl. Gen. et Sp. Orch.* 37.—Reg. Himal.
suaveolens, Lindl. in Gard. Chron. (1856) 372 (Coelogynes sp.).—Ind. or.
triotos, Reichb. f. in Walp. Ann. vi. 238 = Coelogyne triotos.
tristis, *Lindl. ex Baxt. in Loud. Hort. Brit. Suppl.* iii. 607, nomen.—Ind. or.
undulata, Wall. ex Lindl. Bot. Reg. sub t. 1213 = rubra.
ventricosa, *Reichb. f. in Bonplandia*, v. (1857) 43.—Java.

PHOLISMA, Nutt. ex Hook. Ic. Pl. t. 626 (1844). *LENNOACEAE*, Benth. & Hook. f. ii. 622.
arenarium, *Nutt. l. c.*—Am. bor.
depressum, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 198.—Calif.

PHOLISTOMA, Lilja, in Bot. Notiser (1839), 40 = **Nemophila**, Nutt. (Hydroph.).
aurita, Lilja, l. c. = N. aurita.

PHOLIURUS, Trin. Fund. Agrost. 131 (1820) = **Lepturus**, R. Br. (Gramin.).
pannonicus, Trin. l. c. = L. pannonicus.

PHOLIURUS, Trin. in Spreng. Neue Entdeck. ii. 67 (1821) = **Rottboellia**, Linn. f. (Gramin.).
graecus, Trin. l. c. = R. digitata.

PHOLOMPHIS, Rafin. Sylva Tellur. 97 (1838) = **Miconia**, Ruiz & Pav. (Melastom.).
fragilis, Rafin. l. c. = M. Acinodendron.

PHONIPHORA, Neck. Elem. iii. 302 (1790) = **Phoenix**, Linn. (Palm.).

PHONUS, Hill, Veg. Syst. iv. 5 (1762) = **Carthamus**, Tourn. (Compos.).
lanatus, Hill, Hort. Kew. 56 = C. lanatus.

PHORADENDRON, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185. *LORANTHACEAE*, Benth. & Hook. f. iii. 214.
ALLOBIUM, Miers, in Ann. & Mag. Nat. Hist. Ser. II. viii. (1851) 178.
CASTREA, A. St. Hil. Morphol. 451 (1840).
SPICIVISCUM, Engelm. in A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 58.
acinacifolium, *Mart. ex Eichl. in Mart. Fl. Bras.* v. II. 117.—Am. trop.
affine, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = rubrum.
amplexicaule, *Eichl. l. c.* 110.—Am. trop.
angustifolium, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
annulatum, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 176.—Costa Rica.
bathyoryctum, *Eichl. in Mart. Fl. Bras.* v. II. 123.—Am. trop.
Berterianum, Griseb. Fl. Brit. W. Ind. 313 = Grisebachianum.
Berterianum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Cuba; Ins. S. Doming.
Bolleanum, *Eichl. in Mart. Fl. Bras.* v. II. 134 m.—Mexic.
brachystachyum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
brevifolium, *Oliver, in Kjoeb. Vidensk. Meddel.* (1846) 176.—Am. trop.
buxifolium, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Dendrophthora buxifolia.
californicum, *Nutt. l. c.*—Calif.
caripense, *Eichl. in Mart. Fl. Bras.* v. II. 111.—Am. trop.
cayennense, *Eichl. l. c.* 129, in obs.—Guian.; Venezuela.
cearense, *Eichl. in Mart. Fl. Bras.* v. II. 118.—Am. trop.
chrysocladon, *A. Gray, Bot. U. St. Expl. Exped.* i. 743.—Bras.
chrysostachyum, *Eichl. in Mart. Fl. Bras.* v. II. 107, in nota.—Peruv.
clavatum, *Eichl. l. c.*—Am. trop.
constrictum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 192 = Dendrophthora constricta.
cordifolium, *Eichl. in Mart. Fl. Bras.* v. II. 134, nomen.—Am. centr.
coriaceum, *Mart. ex Eichl. l. c.* 121.—Am. trop.
corynarthron, *Eichl. l. c.* 115, in obs.—Am. trop.
craspedophyllum, *Eichl. l. c.* 124.—Am. trop.
crassifolium, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Bras.
crassuloides, *Eichl, in Mart. Fl. Bras.* v. II. 107, in nota.—Am. trop.
dimidiatum, *Eichl. l. c.* 134 m.—Guian.; Venezuela.
dipterum, *Eichl. l. c.* 109.—Am. trop.
ellipticum, *Eichl. l. c.* 119.—Am. trop.
emarginatum, *Mart. ex Eichl. l. c.* 118.—Am. trop.
ensifolium, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
falcatum, *Eichl. in Mart. Fl. Bras.* v. II. 107, in nota.—Am. trop.
falcifrons, *Eichl. l. c.* 134 m.—Uruguay.
Fendlerianum, *Eichl. l. c.* 129, in obs.—Am. trop.
flavens, Griseb. Fl. Brit. W. Ind. 313 = chrysocladon.
flavescens, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Texas.
flavum, Eichl. in Mart. Fl. Bras. v. II. 106 = chrysocladon.
glandulosum, *Eichl. l. c.* 134 m.—Venezuela.
globuliflorum, *Eichl. l. c.*—N. Granat.
Grisebachianum, *Eichl. l. c.*—Ind. occ.
habrostachyum, *Eichl. l. c.* 111.—Am. trop.
hexastichon, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 191.—Am. trop.
holoxanthum, *Eichl. in Mart. Fl. Bras.* v. II. 116.—Am. trop.
interruptum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
juniperinum, *Engelm. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 59.—N. Mexic.; Ind. occ.
lanceolato-ellipticum, Eichl. in Mart. Fl. Bras. v. II. 114 = interruptum.
lanceolatum, *Engelm. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 59.—Texas.
latifolium, Griseb. l. c. viii. (1861) 191 = Schottii.
laurifolium, *Eichl. in Mart. Fl. Bras.* v. II. 107, in nota.—Am. trop.
leptostachya, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Dendrophthora leptostachya.
Liga, *Eichl. l. c.* 134 m.—Chili.
ligustrinum, *Eichl. l. c.*—Am. centr.
linearifolium, *Eichl. l. c.* 115.—Bras.
linifolium, *Eichl. l. c.* 134 m.—Am. trop.
longifolium, *Zucc. ex Eichl. l. c.* 107, in nota.—Am. trop.
macrarthrum, *Eichl. l. c.* 124.—Am. trop.
macrostachyum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Dendrophthora leptostachya.
Mandonii, *Eichl. in Mart. Fl. Bras.* v. II. 124, in obs.—Am. trop.
martinicense, Griseb. Fl. Brit. W. Ind. 314 = Berterianum.
martinicense, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 191 = hexastichon?
mesembryanthemifolium, *Griseb. in Lechl. Berb. Am. Austr.* 59.—N. Granat.
multifoveolatum, *Eichl. in Mart. Fl. Bras.* v. II. 110.—Am. trop.
myrtilloides, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 191 = Dendrophthora myrtilloides.
nervosum, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 175.—Mexic.
nitidum, *Eichl. in Mart. Fl. Bras.* v. II. 113.—Am. trop.
obliquum, *Eichl. l. c.* 134 m.—Am. centr.
oblongifolium, *Eichl. l. c.*—Ind. occ.
obtusissimum, *Eichl. l. c.*—Venezuela.
orbiculatum, *Engelm. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 59.—Texas.
Ottonis, *Eichl. in Mart. Fl. Bras.* v. II. 119, in obs.—Am. trop.
pachyarthron, *Eichl. l. c.* 122, in obs.—Am. trop.
pauciflorum, *Torr. in Pacif. Rail. Rep.* iv. 134.—Mexic.
pellucidulum, *Eichl. in Mart. Fl. Bras.* v. II. 112.—Am. trop.
pennivenium, *Eichl. l. c.* 128.—Am. trop.

PHORADENDRON :—
pepericarpum, *A. Gray, Bot. U. St. Expl. Exped.* i. 742.—Bras.
Perrottetii, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Bras.
peruvianum, *Eichl. in Mart. Fl. Bras.* v. II. 123, in obs.—Am. trop.
piperoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Schottii.
platycaulon, *Eichl. l. c.* 108.—Bras.
polygynum, *Eichl. l. c.* 124, in obs.—Am. trop.
pteroneuron, *Eichl. l. c.* 127.—Bras.
Reichenbachianum, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 175.—Am. trop.
robustissimum, *Eichl. in Mart. Fl. Bras.* v. II. 122.—Am. trop.
rubrum, Griseb. ex Eichl. l. c. 115, in obs. = corynarthron.
rubrum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
salicifolium, *Eichl. in Mart. Fl. Bras.* v. II. 110, in obs.—Am. trop.
sauruoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Schottii.
Schottii, *Nutt. l. c.*; *A. Gray, Bot. U. St. Expl. Exped.* i. 742.—Bras.
Selloi, *Eichl. in Mart. Fl. Bras.* v. II. 116.—Am. trop.
serpyllifolium, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 192 = Dendrophthora cupressoides.
sessilifolium, *Griseb. l. c.* 191.—Ind. occ.
squamigerum, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 176.—Costa Rica.
strongyloclados, *Eichl. in Mart. Fl. Bras.* v. II. 109.—Am. trop.
tereticaule, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 185 = Schottii.
testifolium, Wright, ex Griseb. Cat. Pl. Cub. 120 = Dendrophthora constricta.
tetrastachyum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 191.—Cuba.
tomentosum, *Engelm. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 59.—N. Mexic.
tomentosum, *Oliver, in Kjoeb. Vidensk. Meddel.* (1864) 176.—Mexic.
torulosum, *Eichl. in Mart. Fl. Bras.* v. II. 134 m.—Am. trop.
Trianae, *Eichl. l. c.* 117, obs. ii.—Am. trop.
trinervium, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Ind. occ.
tunaeforme, *Eichl. in Mart. Fl. Bras.* v. II. 108.—Bras.
ulophyllum, *Eichl. l. c.* 123.—Am. trop.
undulatum, *Eichl. l. c.* 122.—Bras.
velutinum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 185.—Am. trop.
illosum, Nutt. l. c.; Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 212 = flavescens.
Warmingii, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 209.—Bras.

PHORMIUM, Forst. Char. Gen. 47. t. 24 (1776). *LILIACEAE*, Benth. & Hook. f. iii. 773.
CHLAMYDIA, Banks, ex Gaertn. Fruct. i. 71. t. 18 (1788).
aloides, Linn. f. Suppl. 205 = Lachenalia tricolor.
bulbiferum, Cyrill. Pl. Rar. Neap. Fasc. i. 35 = Lachenalia pendula.
Colensoi, Hook. f. in Raoul, Choix, 41 = Cookianum.
Cookianum, *Le Jolis, in Hook. Lond. Journ. Bot.* vii. (1848) 536.—N. Zel.
Forsterianum, Hook. l. c. iii. (1844) 8 = Cookianum.
hirtum, Thunb. Nov. Gen. 96 = Lachenalia hirta.
hyacinthoides, Linn. f. Suppl. 204 = Lachenalia orchioides.
orchioides, Thunb. Nov. Gen. 95 = Lachenalia orchioides.
ramosum, Billb. in Linneska Samfund. i. 1832 (1833) 22 = tenax.
tenax, *Forst. Char. Gen.* 48.—N. Zel.
viride, Thunb. Nov. Gen. 127, in syn. = Dipcadi filamentosum.

PHORNOTHAMNUS, Baker, in Journ. Linn. Soc. xxi. (1884) 342. *MELASTOMACEAE.*
thymoides, *Baker, l. c.*—Madag.

PHOSANTHUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 82 = **Isertia**, Schreb. (Rubiac.).
coccinea, Rafin. l. c. = I. coccinea.

PHOTINIA, Lindl. in Trans. Linn. Soc. xiii. (1821) 103. t. 10. *ROSACEAE*, Benth. & Hook. f. i. 627. HETEROMELES, M. Roem. Syn. Rosifl. 105 (1847).
arbutifolia, *Lindl. in Trans. Linn. Soc.* xiii. (1821) 103. t. 10.—Calif.
arguta, Wall. Cat. n. 672 = Pourthiaea arguta.
bengalensis, Wall. ex G. Don, Gen. Syst. ii. 602 = Eriobotrya bengalensis.
Benthamiana, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 213.—China.
Blumei, Decne. in Nouv. Arch. Mus. Par. Sér. I. xi. (1874) 142 = Notoniana.
crenato-serrata, *Hance, in Journ. Bot.* xviii. (1880) 261.—China.
dasythyrsa, *Miq. Fl. Ind. Bat.* i. 1. 387.—Sumatra.
dubia, Lindl. in Trans. Linn. Soc. xiii. (1821) 104. t. 10 = Eriobotrya bengalensis.
eugeniaefolia, Lindl. Bot. Reg. t. 1956 = Notoniana.
foliolosa, Nutt. ex M. Roem. Syn. Rosifl. 105 = arbutifolia.
Fortuneana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 179.—China.
glabra, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 140.—Japon.
Griffithii, *Decne. l. c.* 142.—Reg. Himal.
integrifolia, *Lindl. in Trans. Linn. Soc.* xiii. (1821) 103. t. 10.—Reg. Himal.
laevis, *DC. Prod.* ii. 631.—Japon.
Lindleyana, *Wight & Arn. Prod.* 302.—Ind. or.
longifolia, Decne. in Nouv. Arch. Mus.. Par. Ser. I. x. (1874) 142 = Eriobotrya longifolia.
Maximowiczii, *Decne. l. c.* 143.—As. or.
melanostigma, *Hance, in Journ. Bot.* xx. (1882) 5.—China.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* i. 380.—Mexic.
micrantha, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 143 = Notoniana.
mollis, *Hook. f. Fl. Brit. Ind.* ii. 381.—Reg. Himal.
Notoniana, Wall. Cat. n. 671 = Lindleyana.
Notoniana, *Wight & Arn. Prod.* 302.—Ind. or.; Burma; Malaya.
nudiflora, Nutt. ex M. Roem. Syn. Rosifl. 105 = arbutifolia.
polyneura, King, ex Hook. f. Fl. Brit. Ind. ii. 377 = Pyrus rhamnoides.
prunifolia, *Lindl. Bot. Reg.* t. 1956.—China.
pustulata, *Lindl. Bot. Reg.* sub t. 1956.—China.
pustulata, S. Moore, in Journ. Bot. xvi. (1878) 138 = serrulata.
salicifolia, Presl, Epim. Bot. 204 = arbutifolia.
serrulata, *Lindl. in Trans. Linn. Soc.* xiii. (1821) 103. t. 10.—Japon; China.
serrulata, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 121 = glabra.
Sieboldi, G. Don, Gen. Syst. ii. 602 = Raphiolepis japonica.
subsessilis, King, ex Hook. f. Fl. Brit. Ind. ii. 371 = Eriobotrya Hookeriana.
tenuis, *Carr. in Rev. Hortic.* (1869) 370 (Quid?).—Japon.
villosa, *DC. Prod.* ii. 631.—Japon.

PHOXANTHUS, Benth. in Trans. Linn. Soc. xxii. (1857) 127. t. 23, 24. *SABIACEAE*, Benth. & Hook. f. i. 414.
heterophyllus, *Benth. l. c.*—Bras.

PHRAGMITES, Adans. Fam. ii. 34 (1763) = **Saccharum**, Linn. (Gramin.).

PHRAGMITES, Trin. Fund. Agrost. 134 (1820), partim. *GRAMINEAE*, Benth. & Hook. f. iii. 1179. ARUNDO, Beauv. Agrost. 60. t. 13. f. 2 (1812). CZERNYA, Presl, Cyp. et Gram. Sicul. 22 (1826). OXYANTHE, Steud. Syn. Pl. Gram. 197 (1855). TRICHOON, Roth, in Roem. Arch. i. III. (1798) 37.
altissimus, Mabille, Rech. Pl. Corse, fasc. ii. 39 = communis.
arundinacea, Allam. ex Munro, in Journ. Linn. Soc. vi. (1862) 49 = communis.
australis, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
Berlandieri, Fourn. in Bull. Soc. Bot. Fr. xxiv. (1877) 178 = communis.
breviglumis, *Pomel, Nouv. Mat. Fl. Atl.* 386.—Alger.
capensis, Nees, Fl. Afr. Austr. 356 = communis.

PHRAGMITES :—
capensis, Trin. ex Steud. Nom. ed. II. ii. 34 = P. xenochloa.
caudatus, *Nees, in Meyen, Reise*, i. 407, nomen.—Chili.
chilensis, Steud. Nom. ed. II. ii. 324 = communis.
chrysanthus, Mabille, Rech. Pl. Corse, fasc. ii. 37 = communis.
communis, *Trin. Fund. Agrost.* 134.—Amphigaea.
congesta, Lowe, in Trans. Camb. Phil. Soc. vi. III. (1838) 529 = communis.
disticha, *Buese, in Miq. Pl. Jungh.* 344.—Sumatra.
egmontianus, *Trin. ex Steud. Nom.* ed. II. ii. 324 (Quid?).—Ins. Falkland.
Emodi, *Arn. & Nees, ex Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 174.—Reg. Himal.
explanatus, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
filiformis, *Steud. Syn. Pl. Gram.* 196; *et in Zoll. Syst. Verz. Ind. Archip.* 55.—Java.
fissicha, *Steud. l. c.* 195.—Chili.
flavescens, Hegetschw. Fl. Schw. 68; Peterm. in Flora, xxvii. (1844) 227 = communis.
flexuosa, *Trin. ex Steud. Nom.* ed. II. ii. 324 (sp. dub.).—Ins. Molucc.
gigantea, J. Gay, ex F. Schultz, Arch. Fl. Fr. et Allem. 200 = communis.
graeca, Steud. Nom. ed. II. ii. 324 = communis.
hispanica, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 152 = communis.
humilis, De Not. in Cat. Hort. Genuen. (1846) 27.—Cf. Ann. Sc. Nat. Sér. III. ix. (1848) 325 = communis.
isiaca, Reichb. Fl. Germ. Excurs. 140[b]; Kunth, Rév. Gram. i. 80 = communis.
japonica, Steud. Syn. Pl. Gram. 196 = communis.
Karka, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
laxiflora, Steud. Syn. Pl. Gram. 196 = communis?
longivalvis, *Steud. l. c.*—Japon.
macer, Munro, in Journ. Bot. xv. (1877) 350 = communis?
maritimus, Mabille, Rech. Pl. Corse, fasc. ii. 42 = communis.
martinicensis, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
mauritanica, Kunth, Rév. Gram. i. 80 = communis.
Naya, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
nepalensis, Nees, ex Steud. Syn. Pl. Gram. 196 = communis.
nigricans, Mabille, Rech. Pl. Corse, fasc. ii. 44 = communis.
nudus, Nees, Fl. Afr. Austr. 356 = communis.
occidentalis, Trin. ex Steud. Nom. ed. II. ii. 324 = communis.
pumila, Willk. Iber Halb. 157 = communis.
Roxburghii, Steud. Nom. ed. II. ii. 324 = communis.
ruscinonensis, Mabille, Rech. Pl. Corse, fasc. ii. 41 = communis.
splendens, Mabille, l. c. 42 = communis.
vulgaris, Trin. Fund. Agrost. 134 = communis.
Willkommianus, Mabille, Rech. Pl. Corse, fasc. ii. 58, in obs. (*Wilkomianus*) = communis.
xenochloa, *Trin. ex Steud. Nom.* ed. II. ii. 324; *Syn. Pl. Gram.* 196 (Quid?).—Afr. austr.
Zollingeri, *Steud. Syn. Pl. Gram.* 196 (Quid?).—Java.

PHREATIA, Lindl. Gen. et Sp. Orch. 63 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 510. OCTARRHENA, Thw. Enum. Pl Zeyl. 305 (1861). PLEXAURE, Endl. Prod. Fl. Norf. 30 (1833).
carinata, Miq. Fl. Ind. Bat. iii. 655 = Thelasis carinata.
cauligera, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Ovalu.
contracta, Miq. Fl. Ind. Bat. iii. 655 = Thelasis contracta.
coriacea, Lindl. Gen. et Sp. Orch. 64 = Eria coriacea.
densiflora, *Lindl. l. c.*—Java.
elegans, *Lindl. l. c.* 63.—Reg. Himal.; Zeylan.
laxiflora, *Lindl. l. c.* 64.—Java.
limenophylax, *Benth. Fl. Austral.* vi. 290.—Austral.
Lindleyi, *Benth. & Hook. f. Gen.* iii. 511.—Hab.?
Matthewsii, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Societ.
microtidis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 62.—Java.
minutiflora, *Lindl. l. c.*—Malaya.
Myosurus, *Lindl. l. c.* 61.—Malaya.

PHREATIA :—
oreophylax, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Fiji.
paleata, *Reichb. f. in Linnaea*, xli. (1877) 87.—N. Caled.
prorepens, *Reichb. f. Otia Bot. Hamb.* 54.—Ins. Philipp.
pusilla, *Lindl. Gen. et Sp. Orch.* 64.—Java.
retusa, Lindl. l. c. = Eria retusa.
secunda, Lindl. l. c. = Eria secunda.
tahitensis, *Lindl. in Journ. Linn. Soc.* iii. (1859) 62.—Ins. Tahiti.
uniflora, Wight, Ic. t. 1734 = Eria pusilla, *Lindl.*

PHRENANTHES, [Weber, ex] Wigg. Prim. Fl. Holsat. 56 (1780) = **Prenanthes**, Linn. (Compos.).

PHRISSOCARPUS, Miers, Apocyn. S. Am. 71 (1878). *APOCYNACEAE.*
rigidus, *Miers, l. c.* 72.—Am. austr.

PHRODUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 33. *SOLANACEAE*, Benth. & Hook. f. ii. 896.
RHOPALOSTIGMA, Phil. Fl. Atac. 42. t. 6 (1860).
Bridgesii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 34.—Chili.
microphyllus, *Miers, l. c.* 33.—Chili.
nodosus, *Miers, l. c.* 35.—Chili.

PHRYGANOCYDIA, Mart. ex DC. Prod. ix. 198, in obs. (1845) = **Macfadyena**, A. DC. etc. (Bignon.).
antisyphilitica, Mart. l. c. 199 = Cybistax antisyphilitica.
Candolleana, Mart. l. c. 207 = Spathodea Candolleana.
dipleuropus, Mart. l. c. 204 = Spathodea Coito.
japurensis, Mart. l. c. 214 = M. japurensis.
longa, Mart. l. c. 181 = Lundia longa.
Pisoniana, Mart. l. c. 209 = Spathodea Pisoniana.
xanthophylla, Mart. l. c. 214 = Tabebuia xanthophylla.

PHRYGANTHUS, Baker, in Journ. Linn. Soc. xvii. (1879) 495 = PHYGANTHUS, Poepp. & Endl. = **Tecophilaea**, Bert. (Haemod.).

PHRYGIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 441 (1821) = **Centaurea**, Linn. (Compos.).
major, S. F. Gray, l. c. = C. Scabiosa.
nigra, S. F. Gray, l. c. 442 = C. nigra.
pratensis, S. F. Gray, l. c. = C. nigrescens.

PHRYGILANTHUS, Eichl. in Mart. Fl. Bras. v. II. 45 (1868) = **Loranthus**, Linn.
acutifolius, Eichl. l. c. 49 = L. acutifolius.
aphyllus, Eichl. l. c. 47 = L. aphyllus.
Berteroi, Eichl. l. c. = L. Berteroi.
Bidwillii, Eichl. l. c. 48 = L. Bidwillii.
celastroides, Eichl. l. c. = L. celastroides.
corymbosus, Eichl. l. c. 46 = L. corymbosus.
cuneifolius, Eichl. l. c. 49 = L. cuneifolius.
Destructor, Eichl. l. c. 48 = L. Destructor.
ellipticus, Eichl. l. c. = L. ellipticus.
eugenioides, Eichl. l. c. 50 = L. eugenioides.
flagellaris, Eichl. l. c. 51 = L. flagellaris.
grandiflorus, Eichl. l. c. 46 = L. grandiflorus.
heterophyllus, Eichl. l. c. 47 = L. heterophyllus.
lanceolatus, Eichl. l. c. 48 = L. lancifolius.
ligustrinus, Eichl. l. c. = L. ligustrifolius.
mutabilis, Eichl. l. c. 47 = L. mutabilis.
myrtifolius, Eichl. l. c. 48 = L. myrtifolius.
punctatus, Eichl. l. c. = L. punctatus.
puracensis, Eichl. l. c. = L. puracensis.
secundus, Eichl. l. c. 46 = L. secundus.
suaveolens, Eichl. l. c. 48 = L. suaveolens.
Tagua, Eichl. l. c. = L. Tagua.
tetrandrus, Eichl. l. c. 47 = L. tetrandrus.
verticillatus, Eichl. l. c. = L. verticillatus.

PHRYMA, Forsk. Fl. Aegypt. Arab. p. cxv (1775), fide Vahl = **Priva**, Adans. (Verben.).

PHRYMA, Linn. Diss. Chen. 33 (1751); Amoen. Acad. iii. 19 (1787). *VERBENACEAE*, Benth. & Hook. f. ii. 1137.
DENISAEA, Neck. Elem. i. 306 (1790).
LEPTOSTACHIA, Mitch. in Act. Phys. Med. Soc. Nat. Cur. viii. (1748) App. 212 (nomen prius).

PHRYMA :—
carolinensis, Walt. Fl. Carol. 166 = leptostachya.
dehiscens, Linn. f. Suppl. 277 = Bouchea cuneifolia.
leptostachya, *Linn. Sp. Pl.* 601.—As. bor.; Reg. Himal.; Am. bor.
media, Rafin. New Fl. Am. ii. 37 = leptostachya.
parvifolia, Rafin. l. c. 38 = leptostachya.
pubescens, Rafin. l. c. = leptostachya.

PHRYNIUM, Loefl. It. Hisp. 178 (1758); Willd. Sp. Pl. i. 17 (1797). *SCITAMINEAE*, Benth. & Hook. f. iii. 652.
ILYTHURIA, Rafin. Fl. Tellur. iv. 51 (1836).
PHYLLODES, Lour. Fl. Cochinch. 13 (1790).
Achira, Poepp. & Endl. Nov. Gen. et Sp. ii. 21 = Calathea capitata.
albo-vaginatum, *C. Koch & Linden, in C. Koch & Fint. Wochenschr.* viii. (1865) 369.—Bras.
Allouya, Rosc. Mon. Pl. Scit. t. 38 = Calathea Allouia.
altissimum, Poepp. & Endl. Nov. Gen. et Sp. ii. 20 = Calathea altissima.
angustifolium, *Lodd. Cat. ex Sweet, Hort. Brit.* ed. III. 658.—Ind. or.
bicolor, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea zebrina.
blandum, *C. Koch, l. c.* = Calathea blanda.
brachystachyum, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 108.—Afr. trop.
canniforme, Koern. in Regel, Gartenfl. (1858) 85 = Clinogyne dichotoma.
capitatum, *Willd. Sp. Pl.* i. 17.—Ind. or.; Malaya.
Casupo, Rosc. Mon. Pl. Scit. t. 43 = Calathea discolor.
chrysantha, Brongn. ex Horan. Prod. Scitam. 12 = Calathea chrysantha.
chrysoleucum, Poepp. & Endl. Nov. Gen. et Sp. ii. 19 = Calathea chrysoleuca.
clavatum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea grandifolia.
coloratum, Hook. Bot. Mag. t. 3010 = Calathea colorata.
comosum, Rosc. Mon. Pl. Scit. t. 35 = Calathea comosa.
compositum, Link, ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 59 = Maranta Riedeliana.
compressum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Ctenanthe Luschnathiana.
cylindricum, Rosc. Mon. Pl. Scit. t. 40 = Calathea grandifolia.
Daniellii, Benn. in Pharm. Journ. xiv. (1855) 161 = Thaumatococcus Daniellii.
densiflorum, *Blume, Enum. Pl. Jav.* 38.—Java.
densiflorum, Mor. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 101 = capitatum.
densiflorum, Mor. l. c. 102 = parviflorum.
densum, *C. Koch & Linden, in Belg. Hortic.* xv. (1865) 100.—Bras.
dicephalum, Poepp. & Endl. Nov. Gen. et Sp. ii. 20 = Calathea dicephala.
dichotomum, Roxb. in As. Res. xi. (1810) 324 = Clinogyne dichotoma.
discolor, Hort. ex C. Koch & Mathieu, in Ind. Sem. Hort. Berol. (1855) App. 12 = Calathea varians.
elegans, C. Koch, Wochenschr. vi. (1863) 359 = Calathea vittata.
ellipticum, Rosc. Mon. Pl. Scit. t. 44 = Ischnosiphon spicatus.
eximum, C. Koch & Bouché, in Ind. Sem. Hort. Berol. = Calathea propinqua.
exscapum, Poepp. & Endl. Nov. Gen. et Sp. ii. 18. t. 125 (1855) App. 11 = Calathea eximia.
fasciculatum, *Horan. Prod. Scit.* 11.—Ins. Philipp.
filipes, Benth. in Hook. Niger Fl. 532 = Clinogyne filipes.
flavescens, Sweet, Hort. Brit. ed. II. 494 = Calathea flavescens.
flexuosum, *Benth. in Hook. Niger Fl.* 531.—Afr. trop.
floribundum, Lem. Jard. Fleur. ii. (1852) Misc. 96. t. 189 = Calathea violacea.
giganteum, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 58.—N. Guin.
giganteum, *Teijsm. & Binn. Cat. Hort. Bog.* 60.—Ins. Amboin.
grandiflorum, Rosc. Mon. Pl. Scit. t. 33 = Calathea flavescens.
grandifolium, Sweet, Hort. Brit. ed. II. 494 = Calathea grandifolia.

PHRYNIUM :—
hirsutum, Hort. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 65 = Maranta setosa.
Houtteanum, *C. Koch, Wochenschr.* vi. (1863) 259.—Java.
imbricatum, *Roxb. Hort. Beng.* [1]; *Fl. Ind.* i. 6.—Ind. or.
jagoranum, *C. Koch, in Wochenschr.* vi. (1863) 358.—Malacc.
laterale, Poepp. & Endl. Nov. Gen. et Sp. i. 21 = Calathea lateralis.
latifolium, *Blume, Enum. Pl. Jav.* 37.—Java.
leptostachyum, Hort. Petrop. ex C. Koch, in Berl. Allg. Gartenz. (1857) 147 = Maranta leptostachya.
littorale, Ledeb. ex Sweet, Hort. Brit. ed. III. 658 = Calathea littoralis.
longibracteatum, Sweet, l. c. ed. II. 494 = Calathea longibracteata.
longifolium, C. Koch, in Berl. Allg. Gartenz. (1857) 146 = Calathea grandifolia.
Luschnathiana, Hort. ex C. Koch, l. c. 147 = Ctenanthe Luschnathiana.
luteum, Sweet, Hort. Brit. ed. II. 494 = Calathea lutea.
macilentum, Sweet, l. c. ed. III. 658 = Calathea macilenta.
macrostachyum, *Wall. Cat.* n. 6612 A, B.—Burma.
marantinum, Koern. ex Otto & Dietr. Allg. Gartenz. xxiii. (1855) 193 = Calathea Cassupito.
maximum, *Blume, Enum. Pl. Jav.* 37.—Java.
metallicum, C. Koch, in Allg. Gartenz. xxv. (1857) 147, nomen; et in Wochenschr. vi. (1863) 358 = Calathea metallica.
micans, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxii. (1854) 249 = Calathea micans.
microcephalum, Poepp. & Endl. Nov. Gen. et Sp. ii. 20. t. 128 = Calathea microcephala.
mischanthum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Maranta mischanta.
monophyllum, C. Koch, l. c. = Calathea monophylla.
monostachyum, Steud. Nom. ed. II. ii. 325 = macrostachyum.
Myrosma, Rosc. Mon. Pl. Scit. t. 39 = Myrosma cannaefolia.
nicobaricum, *Didr. in Kjoeb. Vidensk. Meddel.* (1854) 183.—Ins. Nicob.
nobile, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea nobilis.
obliquum, *Lodd. Cat. ex Sweet, Hort. Brit.* ed. III. 658.—Ind. or.
oblongum, Mart. in Flora, xxiv. (1841) II. Beibl. 59 = Calathea oblonga.
obscurum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 329.—Sumatra.
orbiculatum, Sweet, Hort. Brit. ed. III. 658 = Calathea orbiculata.
ornatum, C. Koch, in Allg. Gartenz. xxv. (1857) 147 = Calathea ornata.
ovatum, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) I. 27 = Calathea ovata.
pachystachyum, Poepp. & Endl. Nov. Gen. et Sp. ii. 19. t. 127 = Calathea pachystachya.
Parkeri, Rosc. Mon. Pl. Scit. t. 42 = Ischnosiphon Parkeri.
parviflorum, *Roxb. Hort. Beng.* [1]; *Fl. Ind.* i. 7.—Ind. or.
proliferum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Maranta prolifera.
propinquum, Poepp. & Endl. Nov. Gen. et Sp. ii. 18. t. 125 = Calathea propinqua.
pubigerum, *Blume, Enum. Pl. Jav.* 37.—Java.
pubinerve, *Blume, l. c.* 38.—Java.
pulchellum, Linden, ex C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 149 = Calathea zebrina.
pumilum, Hort. ex C. Koch, l. c. = Calathea vittata.
pumilum, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 918 = Calathea micans.
pumilum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 149 = Calathea pumila.
pusillum, *C. Koch, in Wochenschr.* i. (1863) 359.—Am. trop.
ramosissimum, *Benth. in Hook. Niger Fl.* 532.—Afr. trop.
repens, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 103.—Java.
Riedelianum, Hort. Hafn. ex Horan. Prod. Scitam. 12; Didr. in Kjoeb. Vidensk. Meddel. (1854) 185 = Calathea Koernickiana.
Riedelii, F. Didr. Ind. Sem. Hort. Hafn. (1853) = praec.

PHRYNIUM:—
Rossii, Sweet, Hort. Brit. ed. III. 658 = Calathea Rossii.
rotundifolium, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Ischnosiphon rotundifolius.
sanguineum, Hook. Bot. Mag. t. 4646 = Stromantha sanguinea.
setosum, Rosc. Mon. Pl. Scit. t. 41 = Myrosma setosum.
sinicum, *Miq. in Journ. Bot. Néerl.* i. (1861) 94.—China.
spicatum, *Griff. Notul.* iii. 408 (an seq.?).—Malacca.
spicatum, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 5.—Ind. or.; Burma.
strobiliferum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea strobilifera.
strobulaceum, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 111, sphalm. = praec.
sumatranum, *Miq. Fl. Ind. Bat. Suppl.* 616.—Java.
trifasciatum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147, 162 = Calathea trifasciata.
trifasciatum, *Poit. in Rev. Hortic.* Sér. II. iv. (1845–46) 409.—Mexic.
tuberosum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea tuberosa.
unilaterale, Baker, in Saund. Refug. Bot. t. 312 = Myrosma madagascariensis.
Van-den-Heckei, *Lem. Illustr. Hortic.* x. (1863) t. 380 = Calathea Vandenheckei.
varians, C. Koch & Mathieu, in Ind. Sem. Hort. Berol. (1855) App. 12 = Calathea varians.
variegatum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea variegata.
velutinum, Poepp. & Endl. Nov. Gen. et Sp. ii. 19. t. 126 = Calathea velutina.
villosulum, *Miq. Fl. Ind. Bat. Suppl.* 616.—Sumatra.
villosum, Sweet, Hort. Brit. ed. III. 658 = Calathea villosa.
violaceum, Rosc. Mon. Pl. Scit. t. 37 = Calathea violacea.
virgatum, Roxb. in As. Res. xi. (1810) 324 = Clinogyne virgata.
vittatum, Hort. ex C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 147 = Calathea vittata.
Warszewiczii, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 89 = Calathea Warscewiczii.
zebrinum, Rosc. Mon. Pl. Scit. t. 36 = Calathea zebrina.
zeylanicum, *Benth. & Hook. f. Gen.* iii. 653.—Zeylan.

PHTHEIROSPERMUM, Bunge, in Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 35 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 976.
EMMENOSPERMUM, C. B. Clarke, ex Hook. f. Fl. Brit. Ind. iv. 249, 304 (1883).
chinense, *Bunge, in Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 35.—China.
glandulosum, *Benth. & Hook. f. Gen.* ii. 976.—Reg. Himal.
Parishii, *Hook. f. Fl. Brit. Ind.* iv. 304.—Burma.

PHTHEIROTHECA, Maxim. ex Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 223 = **Caulophyllum**, Michx. (Berberid.).
cyanosperma, Maxim. ex Regel, l. c. = C. robustum.

PHTHIRUSA, Mart. in Flora, xiii. (1830) i. 110 = **Loranthus**, Linn.
adenostemon, Eichl. in Mart. Fl. Bras. v. II. 58.—Bras.
alternifolia, Eichl. l. c. 66.—Bras.
caucana, Eichl. l. c. 60, in nota.—N. Granat.
clandestina, Mart. in Flora, xiii. (1830) i. 110 = L. clandestinus.
cubensis, Eichl. in Mart. Fl. Bras. v. II. 55, in nota = L. cubensis.
domingensis, Eichl. l. c. = L. domingensis.
emarginata, Eichl. l. c. = L. emarginatus.
erythrocarpus, Eichl. l. c. 58 = L. erythrocarpus.
guadalupensis, Eichl. l. c. 55 = L. guadalupensis.
guyanensis, Eichl. l. c. 64.—Guian.
inconspicua, Eichl. l. c. 67, in nota = L. inconspicuus.
janeirensis, Eichl. l. c. 65.—Bras.
laxiflora, Eichl. l. c. 55, in nota = L. laxiflorus.
lepidobotrys, Eichl. l. c. 334 = L. lepidobotrys.
lucarquensis, G. Don, Gen. Syst. iii. 421 = L. lucarquensis.
Magdalenae, Eichl. in Mart. Fl. Bras. v. II. 55 = L. Magdalenae.
micrantha, Eichl. l. c. 65.—Bras.
microphylla, Blume, in Schult. f. Syst. vii. 1730 = L. microphyllus.

PHTHIRUSA:—
Myrsinites, Eichl. in Mart. Fl. Bras. v. II. 66.—Bras.
nitens, Eichl. l. c. 59 = Loranthus nitens.
orinocensis, Eichl. l. c. 60 = L. orinocensis.
ovata, Eichl. l. c. 60 = L. ovatus.
parvifolia, Eichl. l. c. 55, in nota = L. parvifolius.
pauciflora, Eichl. l. c. = L. pauciflorus, *Sw.*
phaeoclados, Eichl. l. c. 65.—Bras.
polystachya, Eichl. l. c. 57.—Bras.
portoricensis, Eichl. l. c. 55, in nota = L. portoricensis.
psilobotrys, Eichl. l. c. = L. psilobotrys.
pycnostachya, Eichl. l. c. 62.—Guian.
pyrifolia, Eichl. l. c. 63 = L. pyrifolius.
rufa, Eichl. l. c. 61 = L. rufus.
salicifolia, G. Don, Gen. Syst. iii. 421 = L. salicifolius.
santaremensis, Eichl. in Mart. Fl. Bras. v. II. 64.—Bras.
squamulosa, Eichl. l. c. 62.—Guian.
stenophylla, Eichl. l. c. 60.—Am. trop.
theloneura, Eichl. l. c. 59.—Bras.
Theobromae, Eichl. l. c. 56 = L. Theobromae.
uniflora, Eichl. l. c. 55, in nota = L. uniflorus.
virgata, Eichl. l. c. 55 = L. virgatus.

PHTIRIUM, Rafin. Princ. Somiol. 29 (1814) = **Delphinium**, Tourn. (Ranunc.).

PHU, Rupp. Fl. Jen. ed. Hall. 214 (1745) = **Valeriana**, Linn.

PHUCAGROSTIS, Willd. Sp. Pl. iv. 649 (1805) = **Cymodocea**, Kon. (Naiad.).
ciliata, Ehrenb. & Hempr. Symb. Phys. Bot. t. 6; ex Boiss. Fl. Orient. v. 23 = C. ciliata.
major, Cavolini, Phucagr. 13. t. 1 = C. aequorea.
minor, Cavolini, l. c. 14. t. 2 = Zostera nana.
rotundata, Ehrenb. & Hempr. Symb. Phys. Bot. t. 11; ex Boiss. Fl. Orient. v. 21 = C. rotundata.
tridentata, Ehrenb. & Hempr. ex Boiss. l. c. 24 = Halodule australis.

PHUOPSIS, Benth. & Hook. f. Gen. ii. 151 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
NEMOSTYLIS, Verz. Stev. Taur. Halbinsel, 187, in obs. (1857), nomen prius.
stylosa, *Benth. & Hook. f. Gen.* ii. 151.—Reg. Caucas.

PHUSICARPOS, Poir. Encyc. Suppl. iv. 399 (1816) = **Hovea**, R. Br. (Legumin.).
linearis, Poir. l. c. 400 = H. linearis.
longifolia, Poir. l. c. = H. longifolia.

PHYCELLA, Lindl. Bot. Reg. sub t. 928 (1825) = **Hippeastrum**, Herb. (Amaryll.).
angustifolia, Phil. in Anal. Univ. Chil. (1873) 543 = H. bicolor.
attenuata, Herb. Amaryll. 153 = H. bicolor.
bicolor, Herb. l. c. 154 = H. bicolor.
biflora, Lindl. Bot. Reg. (1838) Misc. 43 = H. bicolor.
brevituba, Herb. Amaryll. 154 = H. bicolor.
chloracra, Herb. l. c. 155 = Phaedranassa chloracra.
corusca, Lindl. in Trans. Hort. Soc. vii. (1830) 74 = H. bicolor.
cyrtanthoides, Lindl. in Bot. Reg. sub t. 928 = H. bicolor.
glauca, Lodd. Bot. Cab. t. 1746 = H. bicolor?
graciliflora, Herb. Amaryll. 152. t. 25. f. 4 = H. bicolor.
Herbertiana, Lindl. Bot. Reg. t. 1341 = H. Herbertianum.
ignea, Lindl. l. c. sub t. 928 = H. bicolor.
Macleanica, Baker, in Saund. Refug. Bot. t. 332 = H. advena.
Macraeana, Pritz. Ic. Ind. i. 838 = H. bicolor.
magnifica, Herb. Amaryll. 152 = H. bicolor.
obtusa, Lindl. Bot. Reg. (1844) Misc. 91 = Phaedranassa chloracra.
obtusifolia, Pritz. Ic. Ind. i. 838 = H. bicolor.

PHYGANTHUS, Poepp. & Endl. Nov. Gen. et Sp. ii. 71. t. 200 (1838) = **Tecophilaea**, Bert. (Haemod.).
vernus, Poepp. & Endl. l. c. = T. violaeflora.

PHYGELIUS, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. ii. (1836) 53. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 936.
capensis, *E. Mey. ex Benth. l. c.*—Afr. austr.

PHYLA, Lour. Fl. Cochinch. 66 (1790) = **Lippia**, Linn. (Verben.).
chinensis, Lour. l. c. = L. nodiflora.

PHYLACANTHUS, Benth. Scroph. Ind. 1 (1835) = **Angelonia**, H. B. & K. (Scrophul.).

PHYLACIUM, Benn. Pl. Jav. Rar. 159. t. 33 (1838?). *LEGUMINOSAE*, Benth. & Hook. f. i. 523.
HELEIOTIS, Hassk. Cat. Hort. Bog. Alt. 272, in syn. (1844).
bracteosum, *Benn. Pl. Jav. Rar.* 159.—Java.
scandens, *Hassk. in Hoev. & Vriese, Tijdschr.* x. i. (1843) 147.—Java.

PHYLANTHERA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Petunga**, DC. (Rubiac.).

PHYLAX, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Polygala**, Tourn.

PHYLICA, Linn. Sp. Pl. ed. I. 195 (1753). *RHAMNEAE*, Benth. & Hook. f. i. 380.
ALATERNOIDES, Adans. Fam. ii. 304 (1763).
CALOPHYLICA, Presl, Bot. Bemerk. 39 (1844).
PETALOPOGON, Reiss. in Endl. Nov. Stirp. Dec. 82 (1839).
PHILYCA, Linn. Gen. ed. I. 58 (1737).
SOULANGIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 377. t. 10. f. 3.
TRICHOCEPHALUS, Brongn. l. c. 374. t. 17. f. 1 (1827).
TYLANTHUS, Reiss. ex Endl. Gen. 1101 (1840).
WALPERSIA, Reiss. l. c. 1100 (1840).
abietina, *Eckl. & Zeyh. Enum.* 132.—Afr. austr.
abietina, E. Mey. in Drège, Zwei Pfl. Docum. 210 = Spatalla sericea.
acerosa, Sieber, ex Presl, Bot. Bemerk. 39 = parviflora.
acerosa, Willd. Enum. Hort. Berol. 252 = ericoides.
acmaephylla, *Eckl. & Zeyh. Enum.* 133.—Afr. austr.
acuminata, *Nois. ex Steud. Nom.* ed. I. 614, nomen.—Afr. austr.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 489.—Afr. austr.
albida, *Presl, Bot. Bemerk.* 38.—Afr. austr.
alpina, *Eckl. & Zeyh. Enum.* 133.—Afr. austr.
ambigua, *Sond. in Harv. & Sond. Fl. Cap.* i. 486.—Afr. austr.
angustifolia, Hort. ex Steud. Nom. ed. I. 614 = paniculata.
apiculata, *Sond. in Harv. & Sond. Fl. Cap.* i. 497.—Afr. austr.
arborea, Thou. Fl. Tristan d'Acugn. 45 = nitida.
arborescens, Link, ex Steud. Nom. ed. II. ii. 325 = nitida.
arborescens, Steud. l. c. = paniculata.
asperulata, Wender. ex Steud. l. c., nomen (Quid?).—Hab.?
atrata, Bernh. ex Krauss, in Flora, xxvii. (1844) 347 = callosa.
atrata, *Licht. ex Roem. & Schult. Syst.* v. 490.—Afr. austr.
australis, *Link, Enum. Hort. Berol.* i. 230.—Austral.?
axillaris, *Lam. Illustr.* ii. 78.—Afr. austr.
bicolor, *Linn. Mant.* ii. 208; *Murr. Syst.* ed. XIII. 196.—Afr. austr.
brachycephala, *Sond. in Harv. & Sond. Fl. Cap.* i. 500.—Afr. austr.
brevifolia, *Eckl. & Zeyh. Enum.* 133.—Afr. austr.
brunioides, Lam. Illustr. ii. 78 = eriophora.
buxifolia, *Linn. Sp. Pl.* 195.—Afr. austr.
callosa, *Linn. f. Suppl.* 153.—Afr. austr.
capitata, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 490 = rigida.
capitata, *Thunb. Prod. Pl. Cap.* 45.—Afr. austr.
cephalantha, *Sond. in Harv. & Sond. Fl. Cap.* i. 501.—Afr. austr.
Commelini, Spreng. in Ges. Naturf. Fr. Berl. Mag. viii. (1814) 104. t. 8 = plumosa.
comosa, *Steud. Nom.* ed. II. ii. 325.—Afr. austr.
cordata, Herb. ex Sond. in Harv. & Sond. Fl. Cap. i. 489 = reflexa.
cordata, Linn. Sp. Pl. ed. II. 283 = buxifolia.
cordifolia, Lam. ex Steud. Nom. ed. I. 614 = seq.?
cordifolia, Salisb. Prod. 140 = buxifolia.
corifolia, Salisb. l. c. = eriophora.

PHYLICA:—
cotinifolia, Portenschl. ex Reiss. in Endl. Nov. Stirp. Dec. 79 = Nesiota elliptica.
cryptandroides, *Sond. in Harv. & Sond. Fl. Cap.* i. 484.—Afr. austr.
cuspidata, *Eckl. & Zeyh. Enum.* 134.—Afr. austr.
cylindrica, Eckl. & Zeyh. l. c. 134 = excelsa.
cylindrica, *Wendl. Coll. Pl.* i. 29. t. 7.—Afr. austr.
debilis, *Eckl. & Zeyh. Enum.* 132.—Afr. austr.
dioica, *Linn. Syst.* ed. X. 938.—Afr. austr.
diosmoides, *Sond. in Harv. & Sond. Fl. Cap.* i. 496.—Afr. austr.
disticha, *Eckl. & Zeyh. Enum.* 133.—Afr. austr.
divaricata, *Vent. Jard. Malm.* sub t. 57.—Hab. ?
elliptica, Roxb. in Beats. Tracts St. Helena, App. 316 = Nesiota elliptica.
elongata, Salisb. Prod. 140 = spicata.
elongata, Willd. ex Roem. & Schult. Syst. v. 491 = Staavia globosa.
empetroides, Eckl. & Zeyh. Enum. 133 = disticha, rubra.
ericaefolia, Salisb. Prod. 140 = seq.
ericoides, *Linn. Sp. Pl.* 195.—Afr. austr.
ericoides, E. Mey. in Drège, Zwei Pfl. Docum. 210 = villosa.
eriophora, *Berg. Desc. Pl. Cap.* 52.—Afr. austr.
eriophoros, Thunb. Prod. Pl. Cap. 44 = praec.
excelsa, *Wendl. Coll. Pl.* iii. 3. t. 74.—Afr. austr.
fulva, *Eckl. & Zeyh. Enum.* 134.—Afr. austr.
glabrata, Thunb. Diss. Phyl. 3 = ericoides.
globosa, Thunb. l. c. 8 = Staavia globosa.
gnidioides, *Eckl. & Zeyh. Enum.* 135.—Afr. austr.
gracilis, *D. Dietr. Syn. Pl.* i. 875.—Afr. austr.
hirsuta, Thunb. Diss. Phyl. 6 = axillaris.
horizontalis, Vent. Jard. Malm. sub t. 57 = plumosa.
humilis, *Sond. in Harv. & Sond. Fl. Cap.* i. 495.—Afr. austr.
imberbis, *Berg. Desc. Pl. Cap.* 51.—Afr. austr.
imberbis, Drège, ex Presl, Bot. Bemerk. 37 = excelsa.
imberbis, Eckl. & Zeyh. Enum. 133 = ericoides.
imberbis, Thunb. ex Sond. in Harv. & Sond. Fl. Cap. i. 493 = cuspidata.
imbricata, Thunb. Prod. Pl. Cap. 45 = Brunia racemosa.
juniperifolia, Eckl. & Zeyh. Enum. 135 = gnidioides.
laevis, Steud. Nom. ed. II. ii. 326 = atrata.
lanceolata, Eckl. & Zeyh. Enum. 132 = lasiocarpa.
lanceolata, Thunb. Prod. Pl. Cap. 44 = eriophora.
lancifolia, *Steud. Nom.* ed. II. ii. 326.—Afr. austr.
lasiocarpa, *Sond. in Harv. & Sond. Fl. Cap.* i. 483.—Afr. austr.
ledifolia, *Desf. Tabl.* ed. I. 203, nomen.—Afr. austr.
linearis, *Desf. l. c.* ed. II. 232, nomen.—Afr. austr.
litoralis, D. Dietr. Syn. Pl. i. 875 = atrata.
longirostrata, Hort. ex Steud. Nom. ed. I. 614 = paniculata.
lutescens, *D. Dietr. Syn. Pl.* i. 876.—Afr. austr.
mauritiana, Boj. Hort. Maurit. 70, nomen; et ex Baker, Fl. Maurit. 53 = nitida.
Meyeri, *Sond. in Harv. & Sond. Fl. Cap.* i. 488.—Afr. austr.
microcephala, Eckl. & Zeyh. Enum. 133 = disticha.
microcephala, Roem. & Schult. Syst. v. 491 = ericoides.
microphylla, *D. Dietr. Syn. Pl.* i. 876.—Afr. austr.
montana, *Sond. in Harv. & Sond. Fl. Cap.* i. 492.—Afr. austr.
mucronata, *E. Mey. in Drège, Zwei Pfl. Docum.* 211, nomen.—Afr. austr.
myrtifolia, Poir. Encyc. v. 293 = paniculata.
nigrita, *Sond. in Harv. & Sond. Fl. Cap.* i. 498.—Afr. austr.
nitida, *Lam. Illustr.* ii. 77.—Afr. austr.
nitida, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 483 = lasiocarpa.
nuda, Burm. f. Fl. Cap. Prod. 6 = Staavia radiata ?
oblongifolia, Dum.-Cours. Bot. Cult. ed. I. 613 = ledifolia.
oblongifolia, Hort. ex Steud. Nom. ed. I. 614 = paniculata.
oleaefolia, *Vent. Jard. Malm.* sub t. 57.—Afr. austr.
oleoides, DC. Prod. ii. 36 = oleaefolia.
orientalis, *Willd. Enum. Hort. Berol. Suppl.* 12, nomen; *Loisel. Herb. Amat.* t. 233.—Hab. ?
ovata, *Hort. ex Steud. Nom.* ed. II. ii. 326, nomen.—Afr. austr.
paniculata, *Willd. Sp. Pl.* i. 1112.—Afr. austr.
papillosa, Wendl. Coll. Pl. iii. 5. t. 75 = excelsa.
parviflora, *Berg. Desc. Pl. Cap.* 48.—Afr. austr.

PHYLICA:—
parviflora, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 483 = lasiocarpa.
parviflora, E. Mey. in Drège, Zwei Pfl. Docum. 211 = rosmarinifolia.
pedicellata, DC. Prod. ii. 36 = villosa.
pinea, Thunb. Diss. Phyl. 6 = villosa, axillaris.
pinifolia, Linn. f. Suppl. 153 = Brunia pinifolia.
plumosa, *Linn. Sp. Pl.* 195.—Afr. austr.
plumosa, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 486 = ambigua.
propinqua, *Sond. in Harv. & Sond. Fl. Cap.* i. 496.—Afr. austr.
pubescens, Ait. Hort. Kew. ed. I. i. 268 = capitata.
pubescens, Lam. Illustr. ii. 78 = cylindrica.
pubescens, Willd. Enum. Hort. Berol. 252 = plumosa.
pumila, Wendl. ex Willd. Enum. Hort. Berol. 252 = plumosa.
purpurea, *Sond. in Harv. & Sond. Fl. Cap.* i. 485.—Afr. austr.
racemosa, Linn. Mant. ii. 209 = Brunia racemosa.
radiata, Linn. Cent. Pl. i. 8 = Staavia radiata.
ramosissima, *DC. Prod.* ii. 34.—Ins. S. Helena.
reclinata, *Bernh. ex Hochst. in Flora*, xxvii. (1844) 347.—Afr. austr.
reclinata, *Wendl. Coll. Pl.* ii. 49. t. 56.—Afr. austr.
recurvifolia, *Eckl. & Zeyh. Enum.* 135.—Afr. austr.
reflexa, *Lam. Illustr.* ii. 78.—Afr. austr.
retrorsa, *E. Mey. in Drège, Zwei Pfl. Docum.* 211, nomen; *et ex Sond. in Harv. & Sond. Fl. Cap.* i. 492.—Afr. austr.
rhizophora, D. Dietr. Syn. Pl. i. 875 = stipularis.
rigida, *Eckl. & Zeyh. Enum.* 133.—Afr. austr.
rigidifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 484.—Afr. austr.
rosmarinifolia, Lam. Illustr. ii. 78 = eriophora.
rosmarinifolia, E. Mey. in Drège, Zwei Pfl. Docum. 211 = cryptandroides.
rosmarinifolia, Roxb. in Beats. Tracts St. Helena, App. 316 = ramosissima.
rosmarinifolia, *Thunb. Diss. Phyl.* 8.—Afr. austr.
rubra, *Willd. ex Roem. & Schult. Syst.* v. 491.—Afr. austr.
secunda, Thunb. Prod. Pl. Cap. 44 = eriophora.
selaginoides, *Sond. in Harv. & Sond. Fl. Cap.* i. 499.—Afr. austr.
sessiliflora, Hort. ex Steud. Nom. ed. I. 615 = paniculata.
spicata, *Linn. f. Suppl.* 153.—Afr. austr.
spicata, Lodd. Bot. Cab. 323 = oleaefolia.
squamosa, Willd. ex Roem. & Schult. Syst. v. 491 = Raspalia phylicoides.
squarrosa, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 489 = Meyeri.
squarrosa, Vent. Jard. Malm. sub t. 57 = plumosa.
stipularis, *Linn. Mant.* ii. 208.—Afr. austr.
strigosa, Berg. Desc. Pl. Cap. 50 = cuspidata, bicolor.
strigulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 498.—Afr. austr.
tenuis, Spreng. ex Sond. l. c. 501 = eriophora.
Thunbergiana, *E. Mey. in Drège, Zwei Pfl. Docum.* 211, nomen; *et ex Sond. l. c.* 493.—Afr. austr.
thymifolia, *Vent. Jard. Malm.* 57. t. 57.—Ins. Pacif.
thyrsophora, Steud. Nom. ed. II. ii. 326 = Brunia pinifolia.
tortuosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 211, nomen; *et ex Sond. in Harv. & Sond. Fl. Cap.* i. 489.—Afr. austr.
trachyphylla, *D. Dietr. Syn. Pl.* i. 875.—Afr. austr.
trichotoma, Thunb. Prod. Pl. Cap. 187 = Staavia globosa.
velutina, *Sond. in Harv. & Sond. Fl. Cap.* i. 487.—Afr. austr.
verticillata, *D. Dietr. Syn. Pl.* i. 875.—Afr. austr.
villosa, *Thunb. Prod. Pl. Cap.* 44.—Afr. austr.
virgata, *D. Dietr. Syn. Pl.* i. 875.—Afr. austr.
Willdenowiana, *Eckl. & Zeyh. Enum.* 135.—Afr. austr.

PHYLIDRUM, Willd. Sp. Pl. i. 17 (1797) = **Philydrum**, Banks.

PHYLLACANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 78 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Grisebachianus, *Hook. f. l. c.*; *et Ic. Pl.* t. 1095.—Cuba.

PHYLLACHNE, Forst. Char. Gen. 115. t. 58 (1776). *STYLIDIEAE*, Benth. & Hook. f. ii. 535.
HELOPHYLLUM, Hook. f. Handb. N. Zeal. Fl. 167 (1864).
bellidifolia, *F. Muell. Fragm.* viii. 39.—Austral.
clavigera, *F. Muell. l. c.* 40.—N. Zel.
Colensoi, *Berggr. in Minneskr. Fisiog. Sällsk. Lund* (1877), n. 8. p. 11.—N. Zel.
Haastii, *Berggr. in Journ. Bot.* xviii. (1880) 104.—N. Zel.
sedifolia, *F. Muell. Fragm.* viii. 40.—N. Zel.
uliginosa, *Forst. Char. Gen.* 116.—Reg. Magell.

PHYLLACTINIA, Benth. in Benth. & Hook. f. Gen. ii. 488 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
Grantii, *Benth. ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 102.—Afr. trop.

PHYLLACTIS, Pers. Syn. i. 39 (1805). *VALERIANEAE*, Benth. & Hook. f. ii. 153.
AMBLYORHINUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 168.
PORTERIA, Hook. Ic. Pl. t. 864 (1852).
aretioides, *Wedd. Chlor. And.* ii. 30.—Ecuador.
bracteata, *Wedd. l. c.*—N. Granat.
cordifolia, *Wedd. l. c.* 32.—Venezuela.
corymbulosa, *Wedd. l. c.* 34.—Bolivia.
crassipes, *Wedd. l. c.* 29.—Bolivia.
densa, *Wedd. l. c.* 31.—Bolivia.
dinorrhiza, *Griseb. in Goett. Abh.* xix. (1874) 161.—Reg. Argent.
ferax, *Griseb. l. c.* xxiv. (1879) 159.—Reg. Argent.
Gilliesii, *Wedd. Chlor. And.* ii. 33.—Chili.
hispida, *Wedd. l. c.* 29.—Bolivia.
inconspicua, *Wedd. l. c.* 31.—Bolivia.
macrorrhiza, *Wedd. l. c.* 33.—Chili.
Mandoniana, *Wedd. l. c.* 34.—Bolivia.
mexicana, *Benth. & Hook. f. Gen.* ii. 154.—Mexic.
Mutisiana, *Wedd. Chlor. And.* ii. 32.—N. Granat.
obovata, Nutt. Gen. i. 21 = Valeriana edulis.
pinnatifida, *Wedd. Chlor. And.* ii. 33.—N. Granat.
polybotrya, *Griseb. in Goett. Abh.* xix. (1874) 162.—Reg. Argent.
polystachya, Benth. & Hook. f. Gen. ii. 154 = Valerianopsis polystachya.
pratensis, *Benth. & Hook. f. l. c.*—Mexic.
rigida, *Pers. Syn.* i. 39.—Am. austr.
sedifolia, *Wedd. Chlor. And.* ii. 31, in textu.—Reg. Magell.
spathulata, *Pers. Syn.* i. 39.—Peruv.
tenuifolia, *Pers. l. c.*—Peruv.

PHYLLACTIS, Steud. Nom. ed. II. i. 326 (1841) = **Philactis**, Schrad. (Compos.).

PHYLLAGATHIS, Blume, in Flora, xiv. (1831) 507. *MELASTOMACEAE*, Benth. & Hook. f. i. 754.
gymnantha, *Korth. Verh. Nat. Gesch. Bot.* 252.—Borneo.
rotundifolia, *Blume, in Flora*, xiv. (1831) 507.—Sumatra.

PHYLLAMPHORA, Lour. Fl. Cochinch. 606 (1790) = **Nepenthes**, Linn.
mirabilis, Lour. l. c. = N. Phyllamphora.

PHYLLANTHERA, Blume, Bijdr. 1048 (1826). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 739.
bifida, *Blume, l. c.*—Java.

PHYLLANTHERUM, Rafin. in Journ. Phys. xci. (1820) 72 = **Trillium**, Linn. (Liliac.).

PHYLLANTHIDEA, Didr. in Kjoeb. Vidensk. Meddel. (1857) 150 = **Andrachne**, Linn. (Euphorb.).
microphylla, Didr. l. c. 151 = A. microphylla.

PHYLLANTHOS, St.-Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Phyllanthus**, Linn. (Euphorb.).
peltandrum, St. Lag. l. c. = Phyllanthus Peltandra.

PHYLLANTHUS, Linn. Gen. ed. I. 282 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 272.
ANISONEMA, A. Juss. Euph. Tent. 19. t. 4 (1824).
ASTERANDRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 200.
CALOCOCCUS, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvi. (1864) 48.

PHYLLANTHUS:—
Cathetus, Lour. Fl. Cochinch. 607 (1790).
Ceramanthus, Hassk. Cat. Hort. Bogor. Alt. 240 (1844).
Cicca, Linn. Mant. i. 124 (1767).
Conami, Aubl. Pl. Gui. ii. 926. t. 354 (1775).
Diasperus, Linn. Syst. ed. I (1735).
Dichelactina, Hance, in Walp. Ann. iii. 375 (1852–53).
Emblica, Gaertn. Fruct. ii. 122. t. 108 (1791).
Episteira, Rafin. Sylva Tellur. 20 (1838).
Epistylium, Sw. Fl. Ind. Occ. ii. 1095. t. 22 (1800).
Eriococcus, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 143.
Frankia, Bert. ex Steud. Nom. ed. II. i. 646 (1840).
Geminaria, Rafin. Cat. 14 (1824).
Genesiphylla, L'Hérit. Sert. Angl. 29 (1788).
Hemicicca, Baill. Etud. Gén. Euph. 645 (1858).
Hexadena, Rafin. Sylva Tellur. 92 (1838).
Kirganelia, Juss. Gen. 387 (1789).
Lobocarpus, Wight & Arn. Prod. 7 (1834).
Lomanthes, Rafin. Sylva Tellur. 92 (1838).
Macraea, Wight, Ic. v. ii. 27 (1852).
Margaritaria, Linn. f. Suppl. 66 (1781), partim.
Meborea, Aubl. Pl. Gui. ii. 825. t. 323 (1775).
Menarda, Comm. ex A. Juss. Euph. Tent. 23. t. 6 (1824).
Moeroris, Rafin. Sylva Tellur. 91 (1838).
Nellica, Rafin. l. c. 92 (1838).
Niruris, Rafin. l. c. 91 (1838).
Nymphanthus, Lour. Fl. Cochinch. 543 (1790).
Orbicularia, Baill. Etud. Gén. Euph. 616 (1858).
Oxalistylis, Baill. l. c. 628 (1858).
Peltandra, Wight, Ic. v. ii. 24 (1852).
Prosorus, Dalz. in Hook. Kew Journ. iv. (1852) 345.
Reidia, Wight, Ic. v. ii. 27 t. 1903, 1904 (1852).
Rhopium, Schreb. Gen. 608 (1791).
Scepasma, Blume, Bijdr. 582 (1825).
Staurothyrax, Griff. Notul. iv. 476 (1854).
Synexemia, Rafin. Neogenyt. 2 (1825).
Synostemon, F. Muell. Fragm. i. 32 (1858).
Tephranthus, Neck. Elem. ii. 235 (1790).
Tricarium, Lour. Fl. Cochinch. 557 (1790).
Tricaryum, Spreng. Syst. iii. 848 (1826).
Urinaria, Medic. Malv. 80 (1787).
Williamia, Baill. Etud. Gén. Euph. 559. t. 27 (1858).
Wurtzia, Baill. Adansonia, i. (1860–61) 186. t. 7.
Xylophylla, Linn. Mant. ii. 147 (1771); Schreb. Gen. 200 (1789).
Zygospermum, Thw. ex Baill. Etud. Gén. Euph. 620. t. 27 (1858).
abnormis, *Baill. Adansonia*, i. (1860–61) 42.—Texas.
acidissimus, *Muell. Arg. in Linnaea*, xxxii. (1863) 50.—Ins. Philipp.
acidissimus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 22 = Sauropus albicans.
acuminatus, *Vahl, Symb. Bot.* ii. 95.—Am. trop.
acutifolius, *Poir. ex Spreng. Syst.* iii. 21.—Bras.
acutissimus, *Miq. Fl. Ind. Bat.* i. ii. 369.—Java.
acutus, *Wall. Cat.* n. 7931; *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 75.—Penang?
Adami, *Muell. Arg. in DC. Prod.* xv. ii. 327.—Austral.
adenodiscus, *Muell. Arg. in Linnaea*, xxxii. (1863) 23.—Mexic.
adenophyllus, *Muell. Arg. l. c.* 24.—Bras.
adianthoides, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 51.—Guian.
aeneus, *Baill. Adansonia*, ii. (1861–62) 231.—N. Caled.
affinis, *Muell. Arg. in Linnaea*, xxxii. (1863) 48.—Zeylan.
alaternoides, Reichb. ex Baill. Adansonia, i. (1860–61) 83 = reticulatus.
alatus, Blume, Bijdr. 594 = P. Urinaria.
albicans, Wall. Cat. n. 7937 = Fluggea Leucopyrus.
albiflorus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 70.—Austral.
albus, *Muell. Arg. in Flora*, xlviii. (1865) 387.—Ins. Philipp.
albus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 22 (Quid?).—Malaya.
almadensis, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 38.—Bras.
amarus, Schum. & Thonn. Beskr. Guin. Pl. 421 = P. Niruri.

PHYLLANTHUS:—
amentuliger, *Muell. Arg. in Flora*, xlviii. (1865) 390.—Ins. Fiji.
amoenus, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 66.—Bras.
anabaptizatus, *Muell. Arg. in DC. Prod.* xv. ii. 421.—Zeylan.
anceps, Vahl, Symb. Bot. ii. 95 = simplex.
anceps, Wall. Cat. n. 7896 C = Agyneia bacciformis.
anceps, Wall. l. c. n. 7906 F = maderaspatensis.
andamanicus, Kurz, Rep. Veg. Andam. 47 = Helferi.
Andersonii, Muell. Arg. in Flora, lv. (1872) 3 = assamicus.
Anderssoni, *Muell. Arg. in DC. Prod.* xv. ii. 395.—Venezuela.
andrachnoides, Willd. Sp. Pl. iv. 575 = maderaspatensis.
angolensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 329.—Afr. trop.
angulatus, Schum. & Thonn. Beskr. Guin. Pl. 415 = Fluggea microcarpa.
angustifolius, *Sw. Fl. Ind. Occ.* ii. 1111.—Ind. occ.
angustissimus, *Muell. Arg. in Linnaea*, xxxii. (1863) 55.—Bras.
anisolobus, *Muell. Arg. in DC. Prod.* xv. ii. 382.—Peruv.
anomalus, Muell. Arg. in Linnaea, xxxii. (1863) 52 = indicus.
antillanus, Muell. Arg. l. c. 51 = nobilis, virens.
aoraiensis, *Nadeaud, Enum. Pl. Tahiti*, 73.—Ins. Tahiti.
aquaticus, *Wright, in Sauv. Fl. Cub.* 124.—Cuba.
arabicus, Hochst. ex Steud. Nom. ed. II. ii. 326 = maderaspatensis.
arborescens, *Muell. Arg. in Flora*, xlviii. (1865) 370.—Java.
arboreus, Muell. Arg. l. c. 385 = Glochidion arboreum.
arboreus, Sesse & Moç. ex Baill. Adansonia, i. (1860–61) 35 = Moçinianus.
Arbuscula, J. F. Gmel. Syst. 204 = speciosus.
arenicola, *Casar. Nov. Stirp. Bras. Dec.* 88.—Bras.
argentatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 22 = distichus.
aridus, *Benth. Fl. Austral.* vi. 110.—Austral.
Armstrongii, *Benth. l. c.* 112.—Austral.
Arnottianus, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion Arnottianum.
arvensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 332.—Afr. trop.
asperus, *Muell. Arg. in Flora*, xlviii. (1865) 377.—Ind. or.
assamicus, *Muell. Arg. l. c.* 378.—Ind. or.
atropurpureus, *Boj. Hort. Maurit.* 280, nomen.—Ins. Comoro.
atroviridis, Muell. Arg. in Linnaea, xxxii. (1863) 8 = Bojerianus.
attenuatus, *Miq. in Linnaea*, xxi. (1848) 479.—Guiana.
attenuatus, Willd. ex Steud. Nom. ed. II. ii. 326 = acuminatus.
Augustini, *Baill. in Adansonia*, v. (1864–65) 354.—Bras.
australis, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 284.—Tasman.
averrhoaefolius, Hort. ex Muell. Arg. in DC. Prod. xv. ii. 329 = grandifolius.
averrhoaefolius, Hort. ex Steud. Nom. ed. II. ii. 326 = acuminatus.
avicularis, *Muell. Arg. in Linnaea*, xxxii. (1863) 32.—Bras.
axillaris, *Muell. Arg. in DC. Prod.* xv. ii. 412.—Jamaic.
baccatus, Gaertn. Fruct. ii. 123, in syn. = *bacciformis*, Linn.
baccatus, *F. Muell. ex Benth. Fl. Austral.* vi. 102.—Austral.
bacciformis, *Buch.-Ham. ex Wall. Cat.* n. 7909.—Reg. Himal.
bacciformis, Herb. Madr. ex Wall. Cat. n. 7896 B = rotundifolius.
bacciformis, Linn. Mant. ii. 294 = Agyneia bacciformis.
baeobotryoides, *Wall. Cat.* n. 7942.—Reg. Himal.; Burma.
bahiensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 20.—Bras.
Baillonianus, *Muell. Arg. l. c.* 47.—Zeylan.

PHYLLANTHUS:—
baladensis, *Baill. Adansonia*, ii. (1861–62) 233.—N. Caled.
Banksii, A. Cunn. ex Benth. Fl. Austral. vi. 117 = Neoroepera Banksii.
Beckleri, Muell. Arg. in Linnaea, xxxiv. (1865–66) 74 = simplex.
benguelensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 331.—Afr. trop.
Benthamianus, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion macrophyllum, *Benth.*
Benthamianus, Muell. Arg. in Linnaea, xxxii. (1863) 29 = graveolens.
Bernerianus, *Baill. Adansonia*, ii. (1861–62) 52, nomen; *Muell. Arg. in DC. Prod.* xv. ii. 361.—Madag.
Berteroanus, *Muell. Arg. in Linnaea*, xxxii. (1863) 44.—Ins. S. Doming.
bicolor, Muell. Arg. in Flora, xlviii. (1865) 389 = Glochidion acuminatum.
bicolor, *Vis. in Att. Ist. Venez. Sc.* Ser. III. iv. (1858–59) 139.—Hab.?
Billardieri, Muell. Arg. in Flora, xlviii. (1865) 375 = Glochidion Billardieri.
birmanicus, *Muell. Arg. in Linnaea*, xxxii. (1863) 47.—Burma.
Blanchetianus, *Muell. Arg. l. c.* 38.—Bras.
Blancoanus, *Muell. Arg. l. c.* 49.—Ins. Philipp.
Blumei, Steud. Nom. ed. II. ii. 326 = Breynia rubra.
Bojerianus, Baill. Adansonia, ii. (1861–62) 53 = fusco-luridus.
Bojerianus, *Muell. Arg. in DC. Prod.* xv. ii. 343.—Madag.
bornensis, *Muell. Arg. in Flora*, xlviii. (1865) 377.—Borneo.
bossiaeoides, A. Cunn. ex Benth. Fl. Austral. vi. 98 = Adami.
botryanthus, *Muell. Arg. in DC. Prod.* xv. ii. 323.—Venezuela.
Bourgeoisii, *Baill. Adansonia*, ii. (1861–62) 235.—N. Caled.
brachycladus, Muell. Arg. in Linnaea, xxxii. (1863) 35 = rupestris.
brachylobus, Muell. Arg. in Flora, xlviii. (1865) 373 = Glochidion brachylobum.
brachypodus, F. Muell. ex Benth. Fl. Austral. vi. 103 = maderaspatensis.
brasiliensis, Muell. Arg. in DC. Prod. xv. ii. 383 = P. Conami.
brasiliensis, Poir. Encyc. v. 296 = graveolens.
bupleuroides, *Baill. Adansonia*, ii. (1861–62) 233.—N. Caled.
Burchelli, *Muell. Arg. in Linnaea*, xxxii. (1863) 7.—Afr. austr.
buxifolius, *Reinw. in Blume, Cat. Gew. Buitenz.* 106, nomen; *Muell. Arg. l. c.* 50.—Malaya.
cacuminum, *Muell. Arg. in Flora*, xlviii. (1865) 371.—Java.
caledonicus, Muell. Arg. l. c. 376 = Glochidion caledonicum.
callitrichoides, Griseb. in Goett. Nachr. (1865) 166 = carnosulus.
calycinus, *Labill. Nov. Holl. Pl.* ii. 75. t. 225.—Austral.
calycinus, Wall. Cat. n. 7939 B (Breyniae sp.).—Ind. or.
canaranus, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion zeylanicum.
cantoniensis, Hornem. Enum. Pl. Hort. Hafn. (1807) 29 = P. Urinaria.
cantoniensis, *Schweigg. Enum. Pl. Hort. Regiom.* (1812) 54.—China.
cantoniensis, Zipp. ex Span. in Linnaea, xv. (1841) 347 = Zippelianus.
capensis, Spreng. ex Baill. Adansonia, iii. (1862–63) 163 = Andrachne ovalis.
capensis, Spreng. ex Sond. in Linnaea, xxiii. (1850) 135 = heterophyllus.
capillaris, *Schum. & Thonn. Beskr. Guin. Pl.* 417.—Afr. trop.
carnosulus, *Muell. Arg. in Linnaea*, xxxii. (1863) 30.—Cuba.
carolinensis, *Walt. Fl. Carol.* 228.—Am. bor.
carolinianus, Blanco, Fl. Filip ed. I. 691 = P. Niruri.
Carpentariae, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 72.—Austral.
Casticum, *Soyer-Willem. in Usteri, Ann. Bot.* xviii. (1796) 55.—Ins. Mascar.

PHYLLANTHUS:—
cataractarum, *Muell. Arg. in DC. Prod.* xv. II. 320.—N. Caled.
caudatus, *Muell. Arg. l. c.* 321.—N. Caled.
cauliflorus, Griseb. Fl. Brit. W. Ind. 33 = cladanthus.
cauliflorus, *Muell. Arg. in Linnaea*, xxxii. (1863) 46.—Jamaic.
celastroides, *Muell. Arg. in Flora*, xlviii. (1865) 390.—Java.
ceramica, Pers. Syn. ii. 591 = Exocarpus ceramicus.
cernuus, Poir. Encyc. v. 298 = Breynia cernua.
cernuus, Wall. Cat. n. 7911 (Breyniae sp.).—Ind. or.
Chamaecerasus, *Baill. Adansonia*, ii. (1861–62) 235.—N. Caled.
Chamissonis, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) 420 = reticulatus.
Chantrieri, *André, in Rev. Hortic.* (1883) 537. f. 106, 107.—Cochinch.
Chapelieri, Baill. Adansonia, ii. (1861–62) 52, nomen = madagascariensis.
Cheramela, Roxb. Hort. Beng. 104 = distichus.
Cheremila, Muell. Arg. in DC. Prod. xv. II. 414, in syn. = praec.
chlorophaeus, *Baill. Adansonia*, i. (1860–61) 27.—Mexico.
choretroides, *Muell. Arg. in Linnaea*, xxxii. (1863) 52.—Bras.
chrysanthus, *Baill. Adansonia*, ii. (1861–62) 238.—N. Caled.
Cicca, Griseb. in Goett. Nachr. (1865) 166 = P. Pseudo-cicca.
Cicca, Muell. Arg. in Linnaea, xxxii. (1863) 50 = distichus.
ciccoides, *Muell. l. c.* 13.—Ins. Pacif.
ciliaris, *Baill. Adansonia*, xi. (1876) 373.—N. Caled.
cinerascens, Hook. & Arn. Bot. Beech. Voy. 211 = fasciculatus.
cinerascens, Muell. Arg. in Flora, xlviii. (1865) 390 = Glochidion cinerascens.
cinerascens, Wall. Cat. n. 7915 = Breynia rhamnoides.
cinereus, *Muell. Arg. in Linnaea*, xxxii. (1863) 48.—Zeylan.
cladanthus, *Muell. Arg. l. c.* 46.—Jamaic.
cladotrichus, *Muell. Arg. l. c.* 25.—Bras.
Clausseni, *Muell. Arg. l. c.* 40.—Bras.
coccineus, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion coccineum.
coccineus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 (Quid?).—Java.
cochin-chinensis, *Muell. Arg. in DC. Prod.* xv. II. 417.—China.
cochin-chinensis, Spreng. Syst. iii. 21 = fasciculatus.
cognatus, Spreng. l. c. 23 = linearis.
columnaris, *Muell. Arg. in Linnaea*, xxxii. (1863) 15.—Burma.
colutaeoides, *Baill. Adansonia*, ii. (1861–62) 53, nomen; *et Muell. Arg. in DC. Prod.* xv. II. 335.—Madag.
Commersonii, Baill. Adansonia, ii. (1861–62) 53, nomen = phillyreaefolius.
compressicaulis, Muell. Arg. in Flora, xlviii. (1865) 376 = Glochidion compressicaule.
compressus, *H. B. & K. Nov. Gen. et Sp.* ii. 109.—Mexic.
Conami, *Sw. Prod. Veg. Ind. Occ.* 28.—Am. trop.
concolor, Muell. Arg. in Flora, xlviii. (1865) 374 = Glochidion concolor.
congestus, Benth. ex Muell. Arg. in Linnaea, xxxii. (1863) 25 = Securinega congesta.
consanguineus, *Muell. Arg. in DC. Prod.* xv. II. 378.—Madag.
conterminus, Muell. Arg. in Linnaea, xxxii. (1863) 32 = simplex.
corcovadensis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 30.—Bras.
cordatus, Muell. Arg. in Flora, xlviii. (1865) 376 = Glochidion cordatum.
cordifolius, Wall. Cat. n. 7913 = Andrachne cordifolia.
coriaceus, Muell. Arg. in Flora, xlviii. (1865) 386 = Glochidion coriaceum.
coriaceus, *Wall. Cat.* n. 7946.—Singapore.
cornifolius, H. B. & K. Nov. Gen. et Sp. ii. 115 = grandifolius.
cornutus, *Baill. Adansonia*, ii. (1861–62) 236.—N. Caled.
crassifolius, *Muell. Arg. in Flora*, xlvii. (1864) 513.—Austral.

PHYLLANTHUS:—
crotalaroides, *Zipp. ex Span. in Linnaea*, xv. (1841) 347, nomen.—Ins. Timor.
cryptophilus, *Muell. Arg. in Linnaea*, xxxii. (1863) 8.—Madag.
cumanensis, Willd. ex Steud. Nom. ed. II. ii. 327 = acuminatus.
Cumingii, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion Cumingii.
cuneatus, Willd. Enum. Hort. Berol. Suppl. 65 = maderaspatensis.
cuspidatus, *Muell. Arg. in Flora*, xlviii. (1865) 377.—Ins. Samoa.
cyanospermus, *Muell. Arg. in Linnaea*, xxxii. (1863) 51.—Zeylan.
cyclanthera, *Baill. Adansonia*, i. (1860–61) 31.—Cuba.
cygnorum, Endl. in Enum. Pl. Hueg. 19 = calycinus.
cyrtophylloides, *Muell. Arg. in DC. Prod.* xv. II. 1270.—Java.
cyrtophyllus, Muell. Arg. in Flora, xlviii. (1865) 390 = Glochidion crytophyllum.
cyrtostylus, Muell. Arg. l. c. 387 = Glochidion cyrtostylum.
dalbergioides, Wall. Cat. n. 7934 = reticulatus.
Dallachyanus, *Benth. Fl. Austral.* vi. 104.—Austral.
Daltonis, Muell. Arg. in Flora, xlviii. (1865) 388 = Glochidion Daltoni.
dasyanthus, Muell. Arg. l. c. 390 = Glochidion dasyanthum.
debilis, *Herb. Ham. ex Wall. Cat.* sub n. 7892.—Ind. or.; Afr. trop.
debilis, Herb. Wight, ex Wall. Cat. n. 7899 B = simplex.
debilis, Willd. Sp. Pl. iv. 582 = P. Niruri.
decipiens, *Muell. Arg. in DC. Prod.* xv. II. 347.—Madag.
deflexus, Klotzsch, in Peters, Reise Mossamb. Bot. 104 = Thwaitesianus.
Deplanchei, *Muell. Arg. in Linnaea*, xxxii. (1863) 13.—N. Caled.
depressus, Buch.-Ham. ex Dillw. Rev. Ref. Hort. Malab. 51 = pedunculatus.
depressus, Wall. Cat. n. 7898 = simplex.
dictyospermus, *Muell. Arg. in DC. Prod.* xv. II. 394.—Bras.
diffusus, *Klotzsch, in Seem. Bot. Voy. Herald*, 105.—Am. trop.
dilatatus, Klotzsch, in Peters, Reise Mossamb. Bot. 106 = pentandrus.
dioicus, Schum. & Thonn. Beskr. Guin. Pl. 416 = Fluggea microcarpa.
discoides, *Muell. Arg. in Linnaea*, xxxii. (1863) 51.—Afr. trop.
discolor, *Spreng. Syst.* iii. 21.—Cuba.
distichus, Hook. & Arn. Bot. Beech. Voy. 95 = sandwicensis.
distichus, *Muell. Arg. in DC. Prod.* xv. II. 413.—Ind. or.; As. trop.; Madag.
ditassoides, *Muell. Arg. in Flora*, xlvii. (1864) 487.—Austral.
diversifolius, *Miq. Fl. Ind. Bat. Suppl.* 448.—Sumatra.
dracunculoides, Baill. Adansonia, ii. (1861–62) 239 = Glochidion salicifolium.
Dregeanus, Scheele, in Linnaea, xxv. (1852) 585 = Andrachne ovalis.
dumetosus, *Poir. Encyc.* v. 303.—Madag.
echinatus, Wall. Cat. n. 7893 B = P. Urinaria.
echinospermus, *Wright, in Sauv. Fl. Cub.* 122.—Cuba.
elachophyllus, *F. Muell. ex Benth. Fl. Austral.* vi. 101.—Austral.
elegans, *Wall. Cat.* n. 7926.—Burma.
ellipticus, Baill. Adansonia, v. (1865) 354 = Augustini.
ellipticus, *Buckl. in Proc. Acad. Sc. Philad.* 1862 (1863) 7.—Texas.
ellipticus, *Desf. ex Lam. Encyc. Suppl.* iv. 404.—Hab.?
ellipticus, H. B. & K. Nov. Gen. et Sp. ii. 114 = Securinega elliptica.
elongatus, Steud. Nom. ed. I. 615 = angustifolius.
Emblica, *Linn. Sp. Pl.* 982.—As. trop.
emblicoides, Muell. Arg. in Linnaea, xxxii. (1863) 15 = polyphyllus.
emirnensis, Muell. Arg. l. c. 14 = Bojerianus.

PHYLLANTHUS:—
empetrifolius, Baill. Adansonia, v. (1864–65) 352 = ramillosus.
epiphyllanthus, *Linn. Sp. Pl.* ed. II. 1392.—Cuba.
epiphyllanthus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 (Quid?).—Java.
epistylium, Griseb. Fl. Brit. Ind. 33 = axillaris.
ericoides, *Torr. Bot. Mex. Bound.* 193.—Mexic.
eriocarpus, Muell. Arg. in Flora, xlviii. (1865) 387 = Glochidion eriocarpum, *Champ.*
erythrinus, *Muell. Arg. in DC. Prod.* xv. II. 332.—Cuba.
erythroxyloides, *Muell. Arg. l. c.* 418.—Madag.
essequiboënsis, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1187, nomen.—Guian.
fagifolius, Muell. Arg. in Flora, xlviii. (1865) 373 = Glochidion sphaerogynum.
Fagueti, Boiv. ex Baill. Adansonia, ii. (1861–62) 237 = Glochidion Fagueti.
falcatus, Sw. Fl. Ind. Occ. ii. 1115 = epiphyllanthus.
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 73.—Bras.
fasciculatus, *Muell. Arg. in DC. Prod.* xv. II. 350.—Cochinch.
fasciculatus, Poir. Encyc. v. 304 = P. Casticum.
fastigiatus, *Mart. ex Muell. Arg. in Linnaea*, xxxii. (1863) 45.—Bras.
Ferdinandi, *Muell. Arg. in Flora*, xlviii. (1865) 379.—Austral.
filicaulis, *Benth. Fl. Austral.* vi. 111.—Austral.
filiformis, Pav. ex Baill. Adansonia, (1860–61) 29 = lathyroides.
filipes, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 94.—Ins. Socotra.
fimbriatus, *Muell. Arg. in Linnaea*, xxxii. (1863) 47.—Ind. or.
Finlaysonianus, *Wall. Cat.* n. 7907.—Siam?
flaccidus, Thw. Enum. Pl. Zeyl. 283 = Rheedii.
flagellaris, *Benth. Fl. Austral.* vi. 106.—Austral.
flagelliformis, *Muell. Arg. in Linnaea*, xxxii. (1863) 54.—Bras.
flavidus, Muell. Arg. in Flora, xlviii. (1865) 390 = Glochidion flavidum.
flexuosus, Muell. Arg. in DC. Prod. xv. II. 324 = japonicus.
floribundus, H. B. & K. Nov. Gen. et Sp. ii. 116 = salviaefolius.
floribundus, *Muell. Arg. in Linnaea*, xxxii. (1863) 14.—Afr. trop.
flueggeiformis, Muell. Arg. in DC. Prod. xv. II. 349 = glaucus, *Wall.*
fluggeoides, Muell. Arg. in Linnaea, xxxii. (1863) 16 = Securinega fluggeoides.
fluitans, *Benth. ex Muell. Arg. l. c.* 36.—Bras.
foetidus, Pav. ex Baill. Adansonia, i. (1860–61) 34 = acuminatus.
Forskåhlii, Lepr. ex Baill. l. c. 59 = Euphorbia granulata, aegyptiaca.
Francavillanus, *Muell. Arg. in Linnaea*, xxxii. (1863) 20.—Am. austr.
fraxinifolius, Lodd. Bot. Cab. t. 839 = Glochidion lanceolarium?
frondosus, *Wall. Cat.* n. 7932.—Malaya.
fruticosus, Baill. Adansonia, v. (1864–65) 356, sphalm. = P. Conami.
fruticosus, Heyne, ex Wall. Cat. n. 7899 A = simplex.
Fuernrohrii, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 15.—Austral.
fulvirameus, Muell. Arg. in Flora, xlviii. (1865) 385 = Glochidion fulvirameum.
fusco-luridus, *Muell. Arg. in DC. Prod.* xv. II. 346.—Madag.
fuscus, *Muell. Arg. in Flora*, xlviii. (1865) 380.—Sumatra.
Galeottianus, *Baill. Adansonia*, i. (1860–61) 32.—Mexic.
Gardneri, Thw. Enum. Pl. Zeyh. 282 = simplex.
Gardnerianus, Baill. ex Muell. Arg. in Linnaea, xxxii. (1863) 33 = simplex.
garipensis, E. Mey. ex Muell. Arg. l. c. 7 = tenellus.
Gasstroemi, *Muell. Arg. in DC. Prod.* xv. II. 358.—Austral.
Gaudichaudi, Muell. Arg. in Flora, xlviii. (1865) 379 = Glochidion marianum.
genistoides, *Sond. in Linnaea*, xxiii. (1850) 134.—Afr. austr.
glabrescens, *Muell. Arg. l. c.* xxxii. (1863) 48.—Malaya.

PHYLLANTHUS:—
gladiatus, *Muell. Arg. in Linnaea*, xxxii. (1863) 52.—Bras.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* ii. 115.—Mexic.
glaucescens, Schlecht. & Cham. in Linnaea, vi. (1831) 364 = adenodiscus.
glaucifolius, Wall. Cat. n. 7923 = Glochidion glaucifolium.
glaucinus, *Muell. Arg. in Linnaea*, xxxii. (1863) 13.—Sumatra.
glaucogynus, Muell. Arg. in Flora, xlviii. (1865) 389 = Glochidion Moonii.
glaucophyllus, *Sond. in Linnaea*, xxiii. (1850) 133.—Afr. austr.
glaucus, Muell. Arg. in Flora, xlviii. (1865) 386 = Glochidion glaucum, *Muell. Arg.*
glaucus, *Wall. Cat.* n. 7927 A.—Reg. Himal.; China.
glaucus, Wall. l. c. n. 7927 B = Fluggea microcarpa, Andrachne cordifolia.
Glaziovii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 41.—Bras.
glomeratus, Wall. Cat. n. 7903 = P. Emblica.
glomerulatus, *Muell. Arg. in Flora*, xlviii. (1865) 375.—Sumatra.
Goudotianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 8.—Madag.
gracilentus, *Muell. Arg. in Flora*, xlviii. (1865) 380.—Java.
gracilipes, *Muell. Arg. in Linnaea*, xxxii. (1863) 47.—Java.
gracilis, *Baill. Etud. Gén. Euph.* 630. t. 25.—Java.
gracilis, Roxb. Hort. Beng. 69; Fl. Ind. iii. 654 = maderaspatensis.
gracilissimus, *Baill. Adansonia*, ii. (1861–62) 14.—Cuba.
gracillimus, Muell. Arg. in DC. Prod. xv. II. 408, in syn. = cyclanthera.
grandiflorus, Crantz, Inst. i. 144 = grandifolius.
grandiflorus, Spreng. Syst. iii. 22 = multiflorus.
grandifolius, *Linn. Sp. Pl.* 981.—Am. austr.
grandifolius, Spreng. Syst. iii. 22 = ovatus.
grandisepalus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 72.—Austral.
graveolens, *H. B. & K. Nov. Gen. et Sp.* ii. 112.—Am. austr.
Grayanus, *Muell. Arg. in Flora*, xlviii. (1865) 380.—Ins. Tahiti.
Griffithii, *Muell. Arg. in Linnaea*, xxxii. (1863) 27.—Reg. Himal.
Grisebachianus, *Muell. Arg. l. c.* 26.—Cuba.
griseus, Wall. Cat. n. 7918 = reticulatus, glaucus, *Wall.*, Fluggea microcarpa.
Gueinzii, *Muell. Arg. in Linnaea*, xxxii. (1863) 18.—Afr. austr.
guianensis, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 51 = Schomburgkianus.
Gunnii, *Hook. f. in Hook. l. c.* vi. (1847) 284.—Tasmania.
guyanensis, Muell. Arg. in DC. Prod. xv. II. 376 = attenuatus.
gymnanthus, Baill. Adansonia, ii. (1861–62) 240 = Bischofia javanica.
hakgalensis, *Thw. ex Trimen, Syst. Cat. Pl. Ceyl.* 80.—Zeylan.
Hamiltonianus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 75.—Reg. Himal.
Hamrur, *Forsk. Fl. Aegypt. Arab.* 159.—Arab.
Hasskarlianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 16.—Java.
hebecarpus, *Benth. Fl. Austral.* vi. 108.—Austral.
Helferi, *Muell. Arg. in Flora*, xlviii. (1865) 372.—Burma.
heliotropus, *Wright, ex Griseb. in Goett. Nachr.* (1865) 167.—Cuba.
heteradenius, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 63.—Bras.
heterodoxus, *Muell. Arg. in DC. Prod.* xv. II. 321.—Ins. Fiji.
heterophyllus, *E. Mey. ex Muell. Arg. in Linnaea*, xxxii. (1863) 43.—Afr. austr.
Heyneanus, Muell. Arg. l. c. 49 = longiflorus.
Heyneanus, Muell. Arg. in Flora, xlviii. (1865) 389 = Glochidion Heyneanum.
hirsutus, Muell. Arg. l. c. 371 = Glochidion hirsutum.
hirtellus, *Muell. Arg. in DC. Prod.* xv. II. 326.—Austral.

PHYLLANTHUS:—
hirtellus, F. Muell. ex Muell. Arg. in Linnaea, xxxii. (1863) 22 = thymoides.
Hoffmannseggii, Muell. Arg. l. c. 45 = diffusus.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 117 = Andrachne cordifolia.
Hohenackeri, Muell. Arg. in Flora, xlviii. (1865) 373 = Glochidion Hohenackeri.
hongkongensis, Muell. Arg. l. c. 371 = Glochidion hongkongense.
Hookeri, Muell. Arg. in Linnaea, xxxii. (1863) 19 = P. Urinaria.
humilis, Salisb. Prod. 391 = P. Niruri.
hypoleucus, *Muell. Arg. in Flora*, xlviii. (1865) 374.—Sumatra.
hypoleucus, Muell. Arg. in Linnaea, xxxii. (1863) 40 = lacteus.
hyssopifolioides, *H. B. & K. Nov. Gen. et Sp.* ii. 108.—Am. trop.
hyssopifolius, Muell. Arg. in DC. Prod. xv. II. 390 = praec.
hysteranthus, Muell. Arg. in Linnaea, xxxii. (1863) 52 = indicus.
incurvus, Sond. l. c. xxiii. (1850) 135 = heterophyllus.
incurvus, *Thunb. Prod. Pl. Cap.* 24.—Afr. austr.
indicus, *Muell. Arg. in Linnaea*, xxxii. (1863) 52.—Ind. or.
indigoferoides, *Benth. Fl. Austral.* vi. 110.—Austral.
indigoferoides, A. Cunn. ex Benth. l. c. 104 = Gasstroemi.
insignis, *Muell. Arg. in DC. Prod.* xv. II. 1271.—Java.
insulanus, Muell. Arg. in Flora, xlviii. (1865) 387 = Glochidion insulanum.
introductus, Steud. Nom. ed. II. ii. 327 = Andrachne fruticosa.
jacobinensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 6.—Bras.
jamaicensis, Griseb. Fl. Brit. W. Ind. 34 = reticulatus.
janeirensis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 45.—Bras.
japonicus, *Muell. Arg. in Linnaea*, xxxii. (1863) 52.—Japon.
Jardini, *Muell. Arg. l. c.* 21.—Ins. Pacif.
javanensis, Poir. ex Spreng. Syst. iii. 21 = maderaspatensis.
javanicus, Muell. Arg. in DC. Prod. xv. II. 363, in syn. = maderaspatensis.
javanicus, *Muell. Arg. in Linnaea*, xxxii. (1863) 50.—Java.
juglandifolius, Willd. Enum. Hort. Berol. Suppl. 64 = grandifolius.
junceus, *Muell. Arg. in DC. Prod.* xv. II. 411.—Cuba.
juniperinoides, Muell. Arg. in Linnaea, xxxii. (1863) 18 = Lawii.
juniperinus, Wall. Cat. n. 7901 = parvifolius.
Jussieuanus, Muell. Arg. in Flora, xlviii. (1865) 386 = Glochidion rigidum.
kanalensis, Baill. Adansonia, ii. (1861–62) 234 = Glochidion Lenormandi.
kanalophilus, Muell. Arg. in Flora, xlviii. (1865) 375 = Glochidion kanalense.
khasicus, *Muell. Arg. l. c.* 389.—Ind. or.
Kipareh, Muell. Arg. l. c. 373 = Glochidion glaucum, *Blume*.
Kirganelia, Blanco, Fl. Filip. ed. II. 480 = P. Niruri.
Kirganelia, Willd. Sp. Pl. iv. 587 = P. Casticum.
Kirkianus, *Muell. Arg. in Flora*, xlvii. (1864) 486.—Afr. trop.
Klotzschianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 53.—Bras.
Kollmannianus, *Muell. Arg. in Flora*, xlviii. (1865) 378.—Java.
Korthalsii, *Muell. Arg. l. c.* 377.—Borneo.
Kunthiana, Baill. Etud. Gén. Euph. 629 = salviaefolius.
Kurzianus, *Muell. Arg. in DC. Prod.* xv. II. 1272.—Java.
lacteus, *Muell. Arg. l. c.* 402.—Bras.
lacunarius, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 14.—Austral.
laevigatus, *Muell. Arg. in Flora*, xlviii. (1865) 374.—Burma; Malaya.
lagoënsis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 65.—Bras.
Lambertianus, Muell. Arg. in Flora, xlviii. (1865) 372. = Glochidion Lambertianum.
lamprophyllus, *Muell. Arg. in DC. Prod.* xv. II. 324.—Java.

PHYLLANTHUS:—
lanceolarius, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion lanceolarium.
lanceolatus, *Poir. Encyc.* v. 299.—Ins. Maurit.
lasiogynus, *Muell. Arg. in DC. Prod.* xv. II. 357.—Bras.
lathyroides, *H. B. & K. Nov. Gen. et Sp.* ii. 110.—Mexic.
latifolius, *Sw. Fl. Ind. Occ.* ii. 1109.—Jamaic.
laurifolius, A. Rich. Fl. Cub. Fanerog. ii. 216 = Savia sessiliflora.
Lawii, *J. Grah. Cat. Pl. Bomb.* 181.—Ind. or.
laxiflorus, *Benth. Pl. Hartw.* 90.—Guatemala.
ledifolius, A. Cunn. ex Benth. Fl. Austral. vi. 109 = thymoides.
lepidocarpus, Sieb. & Zucc. in Abh. Acad. Muench. iv. II. (1843) 143 = P. Urinaria.
leprocarpus, Wight, Ic. t. 1895[4] = P. Urinaria.
leptocaulos, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 47.—Bras.
leptoclados, *Benth. Fl. Hongk.* 312.—Ins. Hongkong.
leptogynus, Muell. Arg. in Flora, xlviii. (1865) 389 = Glochidion Gardneri.
leptophyllus, *Muell. Arg. in DC. Prod.* xv. II. 411.—Bras.
Leschenaultii, *Muell. Arg. in Linnaea*, xxxii. (1863) 37.—Ind. or.
leucogynus, *Muell. Arg. in Flora*, xlviii. (1865) 75.—Java.
leucophyllus, Strach. & Winterb. ex Baill. Etud. Gén. Euph. 593 = Fluggea microcarpa.
leucopyrus, Koen. ex Roxb. Hort. Beng. 69; Fl. Ind. iii. 658 = Fluggea Leucopyrus.
Lhotzkyanus, Hochst. ex Steud. Nom. ed. II. ii. 327 = Micrantheum ericoides.
Liebmannianus, *Muell. Arg. in DC. Prod.* xv. II. 366.—Mexic.
Lindbergii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 35.—Bras.
Lindenianus, Baill. Adansonia, ii. (1861–62) 13 = cyclanthera.
linearis, Mart. in Nov. Act. Nat. Cur. xi. (1823) 38 = flagelliformis.
linearis, *Sw. Fl. Ind. Occ.* ii. 1113.—Jamaic.
linearis, Wall. Cat. n. 7906 = maderaspatensis.
linifolius, Vahl, ex Baill. Adansonia, i. (1860–61) 84 = pentandrus.
linoides, Hochst. ex Baill. l. c. = pentandrus.
littoralis, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion littorale.
Llanosi, Muell. Arg. l. c. 387 = Glochidion Llanosi.
loandensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 329.—Afr. trop.
lobocarpus, *Benth. Fl. Austral.* vi. 97.—Austral.
longiflorus, *Heyne, ex Wall. Cat.* n. 7905.—Ind. or.
longifolius, Jacq. Hort. Schoenb. ii. 36 = distichus.
longifolius, *Lam. Illustr.* iii. 348.—Ins. Mascar.
longifolius, Sond. in Linnaea, xxiii. (1850) 135 = maderaspatensis.
longipes, *Muell. Arg. in Linnaea*, xxxii. (1863) 11.—Ind. or.
Lonphali, Herb. Madr. ex Wall. Cat. n. 7895 = P. Niruri.
loranthoides, *Baill. Adansonia*, ii. (1861–62) 238.—N. Caled.
Loureirii, Muell. Arg. in DC. Prod. xv. II. 435 = Briedelia tomentosa.
Lucena, Heyne, ex Roth, Nov. Pl. Sp. 185 = Fluggea Leucopyrus.
lucens, Poir. Encyc. v. 206 = Andrachne fruticosa.
lucidus, Hort. ex Steud. Nom. ed. II. ii. 327 = Fluggea microcarpa.
lucidus, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion lucidum.
lutescens, Muell. Arg. l. c. 377 = Glochidion lutescens.
lycioides, H. B. & K. Nov. Gen. et Sp. ii. 112 = acuminatus.
Macraei, *Muell. Arg. in Linnaea*, xxxii. (1863) 29.—Ind. or.
macrocalyx, *Muell. Arg. l. c.* 48.—Ind. or.
macrocarpus, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion macrocarpum.
macrochorion, Baill. Adansonia, ii. (1861–62) 232 = Glochidion macrochorion.
macrophyllus, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion macrophyllum, *Muell. Arg.*
macrostachyus, Muell. Arg. in Linnaea, xxxii. (1863) 16 = Picramnia Antidesma.

PHYLLANTHUS:—
madagascariensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 35.—Madag.
maderaspatensis, Forsk. Fl. Aegypt. Arab. 159 = tenellus.
maderaspatensis, *Linn. Sp. Pl.* 982.—Geront. trop.
maderaspatensis, Sesse & Moç. ex Baill. Adansonia, i. (1860–61) 25 = polygonoides.
maderaspatensis, Thou. ex Baill. Adansonia, ii. (1861–62) 46 = nummulariaefolius.
malabaricus, Muell. Arg. in Linnaea, xxxiv. (1865–66) 69 = Glochidion malabaricum.
Mannianus, *Muell. Arg. in Flora*, xlvii. (1864) 514.—Afr. trop.
Manono, *Muell. Arg. l. c.* xlviii. (1865) 377.—Ins. Tahiti.
marginatus, Herb. Heyne, ex Wall. Cat. n. 7899 A = simplex.
marianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 17.—Ins. Marian.
Martii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 27.—Bras.
Melleri, *Muell. Arg. in Flora*, xlvii. (1864) 514.—Madag.
Meyerianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 42.—Afr. austr.
micrandrus, *Muell. Arg. l. c.* 27.—Venezuela.
micrantheoides, *Baill. Adansonia*, ii. (1861–62) 238.—N. Caled.
micranthus, *A. Rich. Fl. Cub. Fanerog.* ii. 216.—Cuba.
microcarpus, Muell. Arg. in Linnaea, xxxii. (1863) 51 = reticulatus.
microcladus, *Muell. Arg. l. c.* xxxiv. (1865–66) 71.—Austral.
microdendron, *Muell. Arg. in Journ. Bot.* ii. (1864) 330.—Afr. trop.
microphyllinus, *Muell. Arg. l. c.* 331.—Afr. trop.
microphyllus, *H. B. & K. Nov. Gen. et Sp.* ii. 109.—Venezuela.
microphyllus, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 51 = diffusus.
microphyllus, Mart. Reise Bras. i. 285 = lathyroides.
mimosaefolius, Salisb. Prod. 391 = P. Emblica.
mimosoides, Lodd. Bot. Cab. t. 721 = lathyroides.
mimosoides, *Sw. Prod. Veg. Ind. Occ.* 27.—Ind. occ.
minimus, *Wright, in Sauv. Fl. Cub.* 122.—Cuba.
minutiflorus, *F. Muell. ex Muell. Arg. in Linnaea*, xxxiv. (1865–66) 75.—Austral.
minutulus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 54.—Bras.
Miquelianus, Muell. Arg. in Linnaea, xxxii. (1863) 33 = simplex.
mirabilis, *Muell. Arg. in Flora*, xlvii. (1864) 513.—Siam.
Mitchelli, *Benth. Fl. Austral.* vi. 103.—Austral.
Mocinianus, *Baill. Adansonia*, i. (1860–61) 35.—Mexic.
mollis, *Muell. Arg. in Flora*, xlviii. (1865) 387.—Java.
moluccanus, Muell. Arg. l. c. 380 = Glochidion moluccanum.
montanus, *Sw. Fl. Ind. Occ.* ii. 1117.—Jamaic.
montevidensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 37.—Reg. Argent.
Moonii, Muell. Arg. in DC. Prod. xv. II. 312 = Glochidion Moonii.
mucronatus, H. B. & K. Nov. Gen. et Sp. ii. 112 = acuminatus.
mucronatus, Wall. Cat. n. 7893 = P. Urinaria.
multicaulis, *Muell. Arg. in Linnaea*, xxxii. (1863) 18.—Afr. austr.
multiflorus, *Poir. Encyc.* v. 298.—Madag.
multiflorus, Willd. Sp. Pl. iv. 581 = reticulatus.
multiloculare, Muell. Arg. in Flora, xlviii. (1865) 370 = Glochidion multiloculare.
muricatus, Herb. Madr. ex Wall. Cat. n. 7893 D = P. Urinaria.
myrianthus, *Muell. Arg. in DC. Prod.* xv. II. 317.—N. Caled.
Myrsinites, *H. B. & K. Nov. Gen. et Sp.* ii. 111.—Am. austr.
myrtaceus, *Sond. in Linnaea*, xxiii. (1850) 134.—Afr. austr.
myrtifolius, *Moon, Cat. Pl. Ceyl.* 65.—Zeylan.
myrtifolius, Wall. Cat. n. 7940 = reticulatus.
myrtilloides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 158.—Cuba.

PHYLLANTHUS:—
Myrtillus, Wall. Cat. n. 7892 D = Sauropus quadrangularis.
nanogynus, *Muell. Arg. in Flora*, xlviii. (1865) 376.—Malaya.
Naviluri, Miq. ex Muell. Arg. in Linnaea, xxxii. (1863) 75 (Breyniae sp.).—Ind. or.
neilgherrensis, Muell. Arg. in Flora, xlviii. (1865) 385 = Glochidion neilgherrense.
nemoralis, Muell. Arg. l. c. 389 = Glochidion nemorale.
nemorum, *Russ. ex Wall. Cat.* n. 7897 B; *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 70.—Ind. or.
neo-granatensis, *Muell. Arg. l. c.* xxxii. (1863) 10.—N. Granat.
neopeltandrus, *Griseb. in Goett. Nachr.* (1865) 167.—Cuba.
nepalensis, Muell. Arg. in Flora, xlviii. (1865) 375 = Glochidion velutinum.
nephradenius, Muell. Arg. in DC. Prod. xv. II. 423 = longiflorus.
neurocarpus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 69.—Mexic.
nigrescens, *Muell. Arg. in DC. Prod.* xv. II. 348.—Ins. Philipp.
Niruri, Blanco, Fl. Filip. ed. I. 690 = Blancoanus.
Niruri, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 158 = pruinosus, carnosulus, tenuicaulis.
Niruri, *Linn. Sp. Pl.* 981.—Reg. trop.
Niruri, Schlecht. & Cham. in Linnaea, v. (1830) 87 = lathyroides.
Niruri, Thunb. Fl. Jap. 56 = P. Urinaria.
Niruri, Vell. Fl. Flum. x. t. 16 = acutifolius.
Niruri, Wall. Cat. n. 7894 = maderaspatensis.
Niruri, Wall. l. c. n. 7895 A = rotundifolius.
niruroides, *Muell. Arg. in Journ. Bot.* ii. (1864) 331.—Afr. trop.
nitidulus, *Muell. Arg. in DC. Prod.* xv. II. 1272.—Java.
nitidus, Herb. Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 110 = Eurya japonica.
nitidus, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion zeylanicum.
nivosus, *Bull, Cat.* (1873) 9; *W. G. Sm. Flor. Mag.* N. S. (1874) t. 120.—Ins. Pacif.
nobilis, *Muell. Arg. in DC. Prod.* xv. II. 414.—Guian.
Novae-Hollandiae, Baill. Adansonia, vi. (1866) 343 = baccatus.
Novae-Hollandiae, *Muell. Arg. in DC. Prod.* xv. II. 346.—Austral.
nummulariaefolius, *Poir. Encyc.* v. 302, partim.—Madag.
nummulariaefolius, Poir. l. c., partim = orbiculatus.
nummularioides, *Muell. Arg. in Linnaea*, xxxii. (1863) 5.—Ins. S. Doming.
nutans, *Sw. Prod. Veg. Ind. Occ.* 27.—Jamaic.
obcordatus, Hort. ex Willd. Enum. Hort. Berol. Suppl. 65 = maderaspatensis.
obliquus, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion zeylanicum.
obliquus, Wall. Cat. n. 7947 = Wightianus.
oblongifolius, Dennst. Schluess. Hort. Malab. 31 = Breynia rhamnoides.
obovatus, Mill. Arg. in Flora, xlviii. (1865) 387 = Glochidion obovatum.
obovatus, Muhl. ex Willd. Sp. Pl. iv. 574 = carolinensis.
obscurus, Willd. l. c. 581 = Glochidion obscurum.
obtusatus, *Muell. Arg. in DC. Prod.* xv. II. 433.—Bras.
obtusus, Schrank, in Syll. Ratisb. ii. (1828) 65 = Fluggea microcarpa.
ochrophyllus, *Benth. Fl. Austral.* vi. 99.—Austral.
octomerus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 30.—Bras.
odontadenius, *Muell. Arg. in Journ. Bot.* ii. (1864) 331.—Afr. trop.
oligotrichus, *Muell. Arg. in Flora*, xlviii. (1865) 379.—Java.
oppositifolius, *Baill. Adansonia*, ii. (1861–62) 53, nomen; *Muell. Arg. in Linnaea*, xxxii. (1863) 24.—Ins. Maurit.
orbicularis, *H. B. & K. Nov. Gen. et Sp.* ii. 111. t. 106.—Am. austr.
orbiculatus, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113.—Am. trop.
oreophilus, *Muell. Arg. in Linnaea*, xxxii. (1863) 49.—Zeylan.

PHYLLANTHUS:—
ornatus, Muell. Arg. in Flora, xlviii. (1865) 390 = Glochidion ornatum.
ovalifolius, *Forsk. Fl. Aegypt. Arab.* 159.—Arab.
ovalis, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 135 = Andrachne ovalis.
ovatus, Desf. ex Poir. Encyc. Suppl. iv. 404 (Quid?).—Hab.?
ovatus, *Poir. Encyc.* v. 297.—Ins. Martinic.
oxycarpus, *Muell. Arg. in DC. Prod.* xv. II. 1270.—Sumatra.
oxycladus, Muell. Arg. in Linnaea, xxxii. (1863) 26 = Pavonianus.
oxyphyllus, Miq. Fl. Ind. Bat. Suppl. 448 = frondosus.
oxyphyllus, Muell. Arg. in Linnaea, xxxii. (1863) 40 = acutifolius.
pachyphyllus, Muell. Arg. in DC. Prod. xv. II. 353 = coriaceus.
pacificus, *Muell. Arg. in Linnaea*, xxxii. (1863) 31.—Ins. Pacif.
pallidifolius, Muell. Arg. in DC. Prod. xv. II. 424 = pulcher.
pallidus, Muell. Arg. l. c. 283 = pulcher.
pallidus, *Wright, ex Griseb. in Goett. Nachr.* (1865) 168.—Cuba.
Pancherianus, Baill. Adansonia, ii. (1861–62) 235 = Glochidion Pancherianum.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 50.—Paraguay.
parvifolius, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 63.—Reg. Himal.
parvifolius, Steud. Nom. ed. II. ii. 327 = lathyroides.
parvulus, Sond. in Linnaea, xxiii. (1850) 132 = tenellus.
patens, Miq. ex Muell. Arg. in Linnaea, xxxii. (1863) 34 = simplex.
patens, Roxb. Hort. Beng. 69; Fl. Ind. iii. 667 (Breyniae sp.).—Ind. or.
Pavonianus, *Baill. Adansonia*, i. (1860–61) 30.—Peruv.
pedunculatus, Boiv. ex Baill. l. c. ii. (1861–62) 46 = nummulariaefolius.
pedunculatus, *Kostel. Allg. Med.-Pharm. Fl.* v. 1769.—Ind. or.
Peltandrus, Muell. Arg. in Linnaea, xxxii. (1863) 11 = Thwaitesianus.
penangensis, *Muell. Arg. in Flora*, xlviii. (1865) 388.—Ind. or.
pendulus, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 662.—Ind. or.
pentandrus, Roxb. ex Thw. Enum. Pl. Zeyl. 282 = reticulatus.
pentandrus, *Schum. & Thonn. Beskr. Guin. Pl.* 419.—Afr. trop.
pentaphyllus, Wright, ex Griseb. in Goett. Nachr. (1865) 167 = P. Niruri.
perpusillus, *Baill. Adansonia*, v. (1864–65) 358.—Bras.
Perrottetianus, Muell. Arg. in Flora, xlviii. (1865) 387 = Glochidion neilgherrense.
persimilis, *Muell. Arg. in Linnaea*, xxxii. (1863) 34.—N. Caled.
Pervilleanus, *Muell. Arg. l. c.* 13.—Afr. austr.
petiolaris, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 664.—Ind. or.
philippinensis, Muell. Arg. in Flora, xlviii. (1865) 376, 377 = Glochidion philippinense.
phillyreaefolius, *Poir. Encyc.* v. 299.—As. trop.; Ins. Mascar.
physocarpus, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. trop.
pilosus, *Muell. Arg. in DC. Prod.* xv. II. 432.—Cochinch.
piluliferus, Fenzl, in Flora, xxvii. (1844) 312 = pentandrus.
pimeleoides, A. DC. 9me Not. Pl. Rar. Jard. Genèv. 15 = calycinus.
pinifolius, *Baill. Adansonia*, v. (1864–65) 353.—Bras.
piscatorum, *H. B. & K. Nov. Gen. et Sp.* ii. 113.—Venezuela.
Placenta, Hassk. in Hoev. & De Vriese, Tijdschr. xi. (1844) 225 = seq.
placentatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = Sauropus rhamnoides?
platycalyx, *Muell. Arg. in DC. Prod.* xv. II. 318.—N. Caled.

PHYLLANTHUS :—

podocarpus, *Muell. Arg. in Flora*, xlviii. (1865) 388.—Ins. Fiji.
Poeppigianus, Muell. Arg. in DC. Prod. xv. II. 323 = Glochidion Poeppigianum.
Pohlianus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 49.—Bras.
Poiretianus, Muell. Arg. in Linnaea, xxxii. (1863) 39 = orbiculatus.
polycarpus, *Muell. Arg. in Flora*, xlviii. (1865) 387.—Hab.?
polygamus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 256 = Fluggea microcarpa.
polygonoides, *Spreng. Syst.* iii. 23.—Am. bor.
polymorphus, Muell. Arg. in Linnaea, xxxii. (1863) 24 = phillyreaefolius.
polyphyllus, Dalz. & Gibs. Bomb. Fl. 234 = Lawii.
polyphyllus, Herb. Wight, ex Wall. Cat. n. 7895 D = P. Niruri.
polyphyllus, Wall. Cat. n. 7893 D = P. Urinaria.
polyphyllus, *Willd. Sp. Pl.* iv. 586.—Ind. or.
polyspermus, Schum. & Thonn. Beskr. Guin. Pl. 416 = reticulatus.
pomaceus, Moon, Cat. Pl. Ceyl. 65 (Breyniae sp.).—Zeylan.
praetervisus, Muell. Arg. in Linnaea, xxxiv. (1865–66) 73 = parvifolius.
pratensis, Panch. ex Baill. Adansonia, ii. (1861–62) 237 = simplex.
Preissianus, Klotzsch, in Lehm. Pl. Preiss. i. 179 = calycinus.
Prieurianus, Muell. Arg. in Linnaea, xxxii. (1863) 12 = reticulatus.
procerus, *Wright, in Sauv. Fl. Cub.* 125.—Cuba.
prostratus, *Muell. Arg. in Journ. Bot.* ii. (1864) 330.—Afr. trop.
pruinosus, *Poepp. ex A. Rich. Fl. Cub. Fanerog.* ii. 216.—Cuba.
Pseudo-cicca, *Griseb. in Goett. Nachr.* (1865) 166.—Cuba.
Pseudo-conami, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 43.—Am. austr.
Pseudo-niruri, *Muell. Arg. in Flora*, xlvii. (1864) 539.—Afr. trop.
puberulus, Miq. ex Baill. Etud. Gén. Euph. 614 = reticulatus.
puberus, Muell. Arg. in Flora, xlviii. (1865) 387 = Glochidion sinicum.
pubescens, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 420 = Glochidion Llanosi.
pubescens, Moon, Cat. Pl. Ceyl. 65 = Glochidion Moonii.
pubescens, Wall. Cat. n. 7917 A, B = Breynia discigera.
pubigerus, A. Rich. Fl. Cub. Fanerog. ii. 216 = Savia sessiliflora.
pulchellus, Endl. in Enum. Pl. Hueg. 19 = calycinus.
pulcher, *Wall. Cat.* n. 7908.—Malaya.
pumilus, *Muell. Arg. in DC. Prod.* xv. II. 349.—Ins. Philipp.
purpurascens, H. B. & K. Nov. Gen. et Sp. ii. 110 = lathyroides.
purpureus, *Muell. Arg. in Journ. Bot.* ii. (1864) 329.—Afr. trop.
purpureus, Wright, ex Griseb. in Goett. Nachr. (1865) 168 = erythrinus.
pycnocarpus, Muell. Arg. in Flora, xlviii. (1865) 386 = Glochidion pycnocarpum.
pycnophyllus, Muell. Arg. in DC. Prod. xv. II. 322 = Glochidion microphyllum.
quadrangularis, Klein, ex Willd. Sp. Pl. iv. 585 = Sauropus quadrangularis.
quercinus, *Muell. Arg. in Flora*, xlviii. (1865) 386.—Java.
quinquefidus, Sesse & Moç. ex Baill. Adansonia, i. (1860–61) 39 = grandifolius.
racemifer, Steud. Nom. ed. II. ii. 327 = Breynia racemosa.
racemigerus, *Muell. Arg. in Linnaea*, xxxii. (1863) 23.—Bras.
racemosus, Heyne, ex Wall. Cat. n. 7902 = polyphyllus.
racemosus, Linn. f. Suppl. 415 = Agyneia bacciformis.
ramiflorus, Muell. Arg. in Flora, xlviii. (1865) 374 = Glochidion ramiflorum.
ramiflorus, Pers. Syn. ii. 591 = Securinega ramiflora.
ramillosus, *Muell. Arg. in Linnaea*, xxxii. (1863) 36.—Bras.

PHYLLANTHUS :—

ramosissimus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 70.—Austral.
ramosus, Vell. Fl. Flum. x. t. 17 = cladotrichus.
reclinatus, Roxb. Hort. Beng. 69; Fl. Ind. iii. 669 = Breynia racemosa.
Regnellianus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 58.—Bras.
Reichenbachianus, Sieber, ex Baill. Adansonia, i. (1860–61) 80 = Fluggea microcarpa.
Reinwardtii, *Muell. Arg. in Flora*, xlviii. (1865) 379.—Java.
reticulatus, Lodd. Bot. Cab. t. 116 = nutans.
reticulatus, *Poir. Encyc.* v. 298.—Geront. trop.
retusus, Dennst. Schluess. Hort. Malab. 31 (Breyniae sp.).—Ind. or.
retusus, Roxb. Hort. Beng. 69; Fl. Ind. iii. 657 = Fluggea microcarpa.
revolutus, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 135 = myrtaceus.
rhabdocarpus, *Muell. Arg. in DC. Prod.* xv. II. 1271.—Java.
rhamnoides, Boj. ex Muell. Arg. l. c. 435 = Breynia racemosa.
rhamnoides, Retz. Obs. v. 30; Willd. Sp. Pl. iv. 580 = Breynia rhamnoides.
rhamnoides, Roxb. Hort. Beng. 104; Fl. Ind. iii. 663 = Sauropus quadrangularis.
rhamnoides, Wall. Cat. n. 7911 B (= *Melanthesopsis patens.*).—Ind. or.
Rheedii, *Wight, Ic.* t. 1895.—Ind. or.
rhomboidalis, *Muell. Arg. in Linnaea*, xxxii. (1863) 52.—Madag.
rhytidospermum, *Muell. Arg. l. c.* xxxiv. (1865–66) 70.—Austral.
Riedelianus, *Muell. Arg. l. c.* xxxii. (1863) 16.—Bras.
rigens, *Muell. Arg. in Flora*, xlvii. (1864) 513, in obs.—Austral.
rigidulus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 72.—Austral.
rigidus, *Tate, in Trans. Roy. Soc. S. Austral.* vi. (1883) 101.—Austral.
Roemerianus, Scheele, in Linnaea, xxv. (1852) 583 = Andrachne Roemeriana.
Roeperianus, *Wall. Cat.* n. 7904; *Muell. Arg. in Linnaea*, xxxii. (1863) 28.—Reg. Himal.
rosellus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 53.—Bras.
rosmarinifolius, *Muell. Arg. l. c.* 60.—Bras.
rotundatus, Herb. Wight, ex Wall. Cat. n. 7928 D = Fluggea microcarpa.
rotundatus, *Poir. Encyc.* v. 297.—Ind. or.
rotundifolius, *Klein, ex Willd. Sp. Pl.* iv. 584.—Ind. or.; Afr. trop.
rotundifolius, Wall. Cat. n. 7892 D = Sauropus quadrangularis.
Roxburghii, *Muell. Arg. in Linnaea*, xxxii. (1863) 47.—Reg. Himal.
rubellus, *Muell. Arg. in DC. Prod.* xv. II. 379.—Peruv.
rubens, Boj. ex Baker, Fl. Maurit. 309 = P. Urinaria.
ruber, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 22 = Breynia rubra.
ruber, Spreng. Syst. iii. 22.—Cochinch.
rufidulus, *Muell. Arg. in Linnaea*, xxxii. (1863) 29.—N. Caled.
rufo-glaucus, *Muell. Arg. in Flora*, xlviii. (1865) 372.—Java.
rupestris, *H. B. & K. Nov. Gen. et Sp.* ii. 110.—Bras.
ruscifolius, *Muell. Arg. in DC. Prod.* xv. II. 358.—N. Granat.
ruscoides, H. B. K. Nov. Gen. et Sp. ii. 113 = acuminatus.
salicifolius, Baill. Adansonia, ii. (1861–62) 239 = Glochidion salicifolium.
salviaefolius, *H. B. & K. Nov. Gen. et Sp.* ii. 116.—Am. austr.
samarensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 73.—Ins. Philipp.
sandwicensis, *Muell. Arg. l. c.* xxxii. (1863) 31.—Ins. Sandvic.
saxosus, F. Muell. l. c. xxv. (1852) 441 = Gunnii.
scaber, *Klotzsch, in Lehm. Pl. Preiss.* i. 179.—Austral.
scabrellus, Webb, in Hook. Niger Fl. 175 = rotundifolius, P. Niruri.
scandens, Herb. Roxb. ex Thw. Enum. Pl. Zeyl. 282 = reticulatus.

PHYLLANTHUS :—

scandens, *Muell. Arg. in DC. Prod.* xv. II. 415.—Cuba.
Schomburgkianus, *Muell. Arg. l. c.* 387.—Guian.
scoparius, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 74.—Bras.
scoparius, Welw. Apont. 591 = pentandrus.
Seemannianus, *Muell. Arg. in Flora*, xlviii. (1865) 374.—Ins. Fiji.
Sellowianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 37.—Bras.
semicordatus, *Muell. Arg. in Flora*, xlviii. (1865) 376.—Java.
senensis, *Muell. Arg. in DC. Prod.* xv. II. 335.—Afr. trop.
sepialis, *Muell. Arg. in Bremen, Abh.* vii. (1880) 25.—Afr. trop.
sepiarius, Roxb. ex Wall. Cat. n. 7914 = Breynia rhamnoides.
sericeus, *Muell. Arg. in Flora*, xlviii. (1865) 390.—Java.
serissaefolius, Wall. Cat. n. 7900 = fasciculatus.
sikkimensis, Muell. Arg. in Linnaea, xxxii. (1863) 48 = Hamiltonianus.
silheticus, *Muell. Arg. in Flora*, xlviii. (1865) 378.—Ind. or.
similis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 71.—Austral.
simplex, *Retz. Obs.* v. 29.—As. trop.; Ins. Pacif.
simplicaulis, *Muell. Arg. in Linnaea*, xxxii. (1863) 58.—Bras.
Simsianus, Wall. Cat. n. 7920 = Andrachne fruticosa.
sinensis, Muell. Arg. in Linnaea, xxxii. (1863) 12 = reticulatus.
singalensis, *Muell. Arg. l. c.* 48.—Sumatra.
sinicus, Muell. Arg. l. c. 50 = nobilis.
smilacifolius, *Griseb. Fl. Brit. W. Ind.* 710. nomen.—Ins. Trinit.
societatis, *Muell. Arg. in DC. Prod.* xv. II. 364.—Ins. Pacif.
spathulifolius, *Griseb. in Goett. Nachr.* (1865) 169.—Cuba.
speciosus, *Jacq. Coll.* ii. 360.—Jamaic.
speciosus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 22.—Malaya.
spectabilis, Hueg. ex Vis. in Att. Ist. Venez. Sc. Ser. III. iv. (1858–59) 139 = bicolor.
sphaerogynus, Muell. Arg. in Flora, xlviii. (1865) 375 = Glochidion sphaerogynum.
spinescens, Wall. Cat. n. 7934 = reticulatus.
spinulosus, Herb. Heyne, ex Wall. Cat. n. 7897 A = Lawii.
sponiaefolius, *Muell. Arg. in Linnaea*, xxxii. (1863) 25.—Peruv.
Spruceanus, *Muell. Arg. l. c.* 40.—Bras.
squamatus, *Wright, in Sauv. Fl. Cub.* 123.—Cuba.
squamifolius, *Spreng. Syst.* iii. 21.—Cochinch.
stellatus, Retz. Obs. v. 29 = Glochidion rigidum.
stenocladus, Muell. Arg. in Flora, xlviii. (1864) 536 = Adami.
stenopterus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 74.—N. Granat.
stipulaceus, Boj. Hort. Maurit. 280 = maderaspatensis.
Stocksii, Muell. Arg. in Linnaea, xxxii. (1863) 51 = indicus.
strictus, Roxb. Hort. Beng. 69; Fl. Ind. iii. 670 = Sauropus albicans.
stylosus, Griff. Itin. Notes, 33, nomen = Griffithii.
subcarnosus, *Wright, ex Griseb. in Goett. Nachr.* (1865) 168; *et in Sauv. Fl. Cub.* 125.—Cuba.
subcordatus, Baill. Adansonia, ii. (1861–62) 53 = phillyreaefolius.
subcrenulatus, *F. Muell. Fragm.* i. 108.—Austral.
subemarginatus, *Muell. Arg. in Linnaea*, xxxii. (1863) 39.—Bras.
suberosus, *Wight, ex Wall. Cat.* n. 7910.—Ind. or.
subglomeratus, Poir. Encyc. v. 304 = acuminatus.
sublanatus, *Schum. & Thonn. Beskr. Guin. Pl.* 420.—Afr. trop.
subobscurus, *Muell. Arg. in DC. Prod.* xv. II. 1271.—Java.
subscandens, *Muell. Arg. in Flora*, xlviii. (1865) 372.—Java.
suffultus, Wall. Cat. n. 7939 (Breyniae sp.).—Ind. or.
sundaicus, *Muell. Arg. in DC. Prod.* xv. II. 1272.—Malaya.

PHYLLANTHUS:—
superbus, Muell. Arg. in Flora, xlviii. (1865) 375 = Glochidion superbum.
Swartzii, *Kostel. Allg. Med.-Pharm. Fl.* v. 1771.—Jamaic.
sylvaticus, Steud. Nom. ed. II. ii. 327 = Breynia virgata.
symphoricarpoides, *H. B. & K. Nov. Gen. et Sp.* ii. 114.—N. Granat.
symplocoides, Muell. Arg. in Flora, xlviii. (1865) 389 = Glochidion montanum.
taitensis, Muell. Arg. l. c. 380 = Glochidion taitense.
Tatei, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 55.—Austral.
taxifolius, D. Don, Prod. Fl. Nep. 63 = P. Emblica.
Telfairianus, Wall. Cat. n. 7935 = phillyreaefolius.
tenellus, Benth. Bot. Voy. Sulph. 165 = graveolens.
tenellus, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 668.—Ins. Maurit.
tenellus, Wall. Cat. n. 7892 = debilis, longiflorus, Sauropus quadrangularis.
tenuicaulis, *Muell. Arg. in Linnaea*, xxxii. (1863) 44.—Cuba.
tetrander, Blanco, Fl. Filip. ed. II. 480 = Blancoanus.
tetrandrus, Roxb. Hort. Beng. 69; Fl. Ind. iii. 674 = Roxburghii.
tetrandrus, Wall. Cat. n. 7930 = columnaris, distichus.
Teysmanni, *Muell. Arg. in DC. Prod.* xv. II. 1270.—Java.
thesioides, *Benth. Fl. Austral.* vi. 98.—Austral.
thesioides, Muell. Arg. in Linnaea, xxxii. (1863) 36 = dumetosus.
Thomsoni, *Muell. Arg. in Flora*, xlviii. (1865) 375.—Reg. Himal.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 418 = maderaspatensis.
Thwaitesianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 11.—Zeylan.
thymoides, *Sieber, ex Sond. in Linnaea*, xxviii. (1856) 566, nomen; *Muell. Arg. in DC. Prod.* xv. II. 372.—Austral.
timoriensis, Muell. Arg. in Linnaea, xxxii. (1863) 13 = P. Casticum.
tinctorius, Vahl, ex Baill. Etud. Gén. Euph. 633 = Breynia rhamnoides.
tomentosus, Muell. Arg. in Flora. xlviii. (1865) 371 = Glochidion tomentosum.
tomentosus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = Glochidion molle.
torrentium, *Muell. Arg. in DC. Prod.* xv. II. 316.—N. Caled.
trachygyne, *Benth. Fl. Austral.* vi. 103.—Austral.
trachyspermus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 14.—Austral.
tremulus, *Griseb. Fl. Brit. W. Ind.* 34.—Jamaic.
triandrus, Muell. Arg. in Flora, xlviii. (1865) 379 = Glochidion eleutherostylum.
trichogynus, Muell. Arg. l. c. 380 = Glochidion trichogynum.
trinervius, Wall. Cat. n. 7922 = Sauropus trinervius.
tristis, A. Juss. Euph. Tent. 108 = Breynia rhamnoides.
turbinatus, Koen. ex Roxb. Hort. Beng. 104; Fl. Ind. iii. 666 (Breyniae sp.).—Ind. or.
turbinatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = Breynia virgata.
turbinatus, Sims, Bot. Mag. t. 1862 = Andrachne fruticosa.
uakgalensis, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 242, sphalm. = hakgalensis.
uberiflorus, F. Muell. ex Baill. Adansonia, vi. (1866) 343 = Novae-Hollandiae.
umbratus, *Muell. Arg. in DC. Prod.* xv. II. 356.—Bras.
urceolatus, *Baill. Adansonia*, ii. (1861–62) 239.—N. Caled.
urceolatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 22 = Breynia rhamnoides.
Urinaria, *Linn. Sp. Pl.* 982.—Amphig. trop.
Urinaria, Wall. Cat. n. 7895 B, E = P. Niruri.
Urinaria, Willd. Sp. Pl. iv. 583 = rotundifolia.
ussuriensis, Rupr. & Maxim. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 210, 222 = simplex.
vacciniifolius, *Muell. Arg. in DC. Prod.* xv. II. 322.—Guian.
vaccinioides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 105.—Afr. trop.
vaccinioides, Scheele, in Linnaea, xxv. (1852) 585 = Cluytia heterophylla.

PHYLLANTHUS:—
varians, Muell. Arg. in Flora, xlviii. (1865) 380 = Glochidion varians.
velutinus, Muell. Arg. l. c. 387 = Glochidion velutinum.
venosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 254 = maderaspatensis.
venulosus, *Muell. Arg. in Flora*, xlviii. (1865) 374.—Ins. Fiji.
verrucosus, Thunb. Prod. Pl. Cap. 24 (Securinegae sp.).—Afr. austr.
Vespertilio, Baill. Adansonia, ii. (1861–62) 233 = Glochidion Vespertilio.
Vieillardi, *Baill. l. c.* 236.—N. Caled.
villosus, *Poir. Encyc.* v. 297.—China.
virens, *Muell. Arg. in DC. Prod.* xv. II. 415.—Cuba.
virens, Wall. Cat. n. 7928 = Fluggea microcarpa.
virgatus, Forst. f. Prod. 65 = simplex.
virgineus, Pers. Syn. ii. 591 = P. Casticum.
virgulatus, *Muell. Arg. in Journ. Bot.* ii. (1864) 330.—Afr. trop.
virosus, Roxb. ex Willd. Sp. Pl. iv. 578 = Fluggea microcarpa.
virosus, Wall. Cat. n. 7928 = reticulatus.
virosus, Wight, in Wall. Cat. n. 7939 C = Breynia rhamnoides.
vitiensis, *Muell. Arg. in Flora*, xlviii. (1865) 374.—Ins. Fiji.
Vitis-idea, Koen. ex Roxb. Hort. Beng. 69; Fl. Ind. iii. 665 = Breynia rhamnoides.
wagapensis, *Muell. Arg. in Flora*, xlviii. (1865) 372.—N. Caled.
Wallichianus, Muell. Arg. l. c. 387 = Glochidion Wallichianum.
Welwitschianus, *Muell. Arg. in Journ. Bot.* ii. (1864) 330.—Afr. trop.
Wightianus, *Muell. Arg. in Linnaea*, xxxii. (1863) 47.—Ind. or.
Wightianus, Muell. Arg. l. c. 6 = Fluggea microcarpa.
Wightianus, Wall. Cat. n. 7919 = reticulatus.
Wilkesianus, *Muell. Arg. in DC. Prod.* xv. II. 396.—Ins. Fiji.
williamsioides, *Griseb. in Goett. Nachr.* (1865) 169.—Cuba.
Wrightii, *Muell. Arg. in Flora*, xlviii. (1865) 378.—Ins. Hongkong.
zeylanicus, Muell. Arg. in DC. Prod. xv. II. 281 = Glochidion zeylanicum.
zeylanicus, Muell. Arg. in Linnaea, xxxii. (1863) 49 = anabaptizatus.
Zippelianus, *Muell. Arg. in DC. Prod.* xv. II. 433.—Ins. Timor.
ziziphoides, Baill. ex Gibert, Enum. Pl. Montev. 54, nomen = Sellowianus.
Zollingeri, Muell. Arg. in Flora, xlviii. (1865) 371 = Glochidion Zollingeri.
Zollingeri, Muell. Arg. in Linnaea, xxxii. (1863) 47 = pulcher.
zygophylloides, *Muell. Arg. in DC. Prod.* xv. II. 336.—Hab. ?

PHYLLARTHRON, DC. ex Meissn. Gen. 301, Comm. 210 (1840); et Prod. ix. 243 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1052.
ARTHROPHYLLUM, Boj. Hort. Maurit. 221 (1837).
Bernierianum, *Seem. in Bonplandia*, vii. (1859) 224.—Madag.
Bojeranum, *DC. Prod.* ix. 243.—Madag.
comorense, *DC. l. c.* 244.—Ins. Comoro.
Noronhianum, *DC. l. c.* 243.—Madag.
Poivreanum, *DC. l. c.*—Madag.
Thouarsianum, *DC. l. c.* 244.—Madag.

PHYLLARTHUS, Neck. Elem. ii. 85 (1790) = **Cereus**?, Haw. (Cact.).

PHYLLAUREA, Lour. Fl. Cochinch. 575 (1790) = **Codiaeum**, Rumph. (Euphorb.).
Codiaeum, Lour. l. c. = C. variegatum.

PHYLLEPIDUM, Rafin. in Med. Repos. N. York, v. (1808) 356. *AMARANTACEAE*.—Cf. Benth. & Hook. f. iii. 23.
scariosum, Schult. Syst. vi. 293 = squarrosum.
squamosum, Rafin. Fl. Tellur. iii. 41 = squarrosum.
squarrosum, *Rafin. in Med. Repos. N. York*, v. (1808) 356.—Am. bor.

PHYLLIMENA, Blume, ex DC. Prod. v. 636, in syn. (1836) = **Enhydra**, Lour. (Compos.).
longifolia, Blume, ex Miq. Fl. Ind. Bat. ii. 83 = E. longifolia.
paludosa, Blume, ex DC. Prod. v. 637 = E. paludosa.

PHYLLIREA, Tourn. ex Adans. Fam. ii. 223 (1763) = **Phillyrea**, Linn. (Oleac.).

PHYLLIS, Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 140.
NOBULA, Adans. Fam. ii. 145 (1763).
Galopina, Cruse, in Linnaea, vi. (1831) 20 = Galopina circaeoides.
indica, Linn. Sp. Pl. 232 (Quid?).—Ind.
Nobla, *Linn. l. c.*—Ins. Canar.
parviflora, Steud. Nom. ed. II. ii. 328, sphalm. = seq.
pauciflora, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 141 = P. Nobla.
viridis, Salisb. Prod. 61 = P. Nobla.

PHYLLOBAEA, Benth. in Benth. & Hook. f. Gen. ii. 1020 (1876). *GESNERACEAE*, Benth. & Hook. f. l. c.
amplexicaulis, *Benth. l. c.*; *et C. B. Clarke, in DC. Monog. Phan.* v. 140.—Burma.

PHYLLOBOTRYON, Muell. Arg. in Flora, xlvii. (1864) 534. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 325.
Soyauxianum, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 288.—Afr. trop.
spathulatum, *Muell. Arg. in Flora*, xlvii. (1864) 534.—Afr. trop.

PHYLLOBOTRYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = **Genista**, Linn. (Legum.).
anglica, Fourr. l. c. = G. anglica.

PHYLLOBOTRYUM, Muell. Arg. in DC. Prod. xv. II. 1231 (1866) = **Phyllobotryon**, Muell. Arg. (Euphorb.).

PHYLLOBRYON, Miq. Syst. Pip. 50 (1843) = **Peperomia**, Ruiz & Pav. (Piperac.).
Haenkeanum, Presl, Epim. Bot. 222 = Pep. Haenkeana.
Pavoni, Miq. Syst. Pip. 50 = Pep. foliiflora.

PHYLLOCACTUS, Link, Handb. ii. 10 (1831). *CACTEAE*, Benth. & Hook. f. i. 849.
DISISOCACTUS, G. K[unze,] in Bot. Zeit. iii. (1845) 533; Salm-Dyck, Cact. Hort. Dyck. ed. II. 57 (1850).
DISOCACTUS, Lindl. Bot. Reg. (1845) t. 9.
EPIPHYLLUM, Haw. Syn. Pl. Succ. 197 (1812), partim.
PHYLLOCEREUS, Miq. in Bull. Sc. Phys. et Nat. Brux. (1839) 112.
Ackermanni, *Walp. Rep.* ii. 341; *et Salm-Dyck, Cact. Hort. Dyck.* ed. II. 55.—Mexic.
acutifrons, *Hort. ex Salm-Dyck, l. c.* ed. I. 87, nomen.—Hab. ?
angularis, *Lem. ex Labour. Monog. Cact.* 671, nomen.—Hab. ?
angulifer, G. Don, in Loud. Encyc. Pl. Suppl. ii. 1380 = seq.
anguliger, *Lem. Jard. Fleur.* i. (1851) t. 92.—Mexic.
biformis, *Labour. Monog. Cact.* 418.—Honduras.
caulorrhizus, *Lem. Jard. Fleur.* i. (1851) *Misc.* 6.—Hab. ?
Cooperi ×, Hort. ex Regel, Gartenfl. (1883) 218.
crenatus, *Walp. Rep.* v. 820; *Foerst. Handb. Cact.* ed. I. 441.—Honduras.
grandis, *Lem. in Fl. des Serres*, iii. (1847) 255, verso.—Honduras.
guyanensis, *Brongn. ex Labour. Monog. Cact.* 416.—Guian.
Hookeri, *Walp. Rep.* ii. 341; *et Salm-Dyck, Cact. Hort. Dyck.* ed. II. 55, 223.—Bras.; Guian.
ignescens, *Hort. Dresd. ex Haage, Cact. Vers.* (1857) 10.—Hab. ?
Jenkinsoni, Hort. Angl. ex Haage, l. c. = phyllanthoides.
latifrons, *Walp. Rep.* ii. 341; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 55.—Mexic.
Londonii ×, Haage, ex Labour. Monog. Cact. 429, nomen.
longipes ×, Hort. Belg. ex Labour. l. c., nomen.
Lothii ×, Pfeiff. ex Labour. l. c., nomen.
Macquiana ×, Salm-Dyck. ex Labour. l. c., nomen.

PHYLLOCACTUS:—
macropterus, *Lem. Illustr. Hortic.* xi. (1864) *Misc.* 73.—Bras.
Maelenii ×, Salm-Dyck, ex Labour. Monog. Cact. 429, nomen
marginatus, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 37, nomen.—Hab.?
maurantianus ×, Hort. Berdl. ex Labour. Monog. Cact. 429, nomen.
mexicanus ×, Salm-Dyck, ex Labour. l. c. = Cereus mexicanus.
oxypetalus, *Link, ex Walp. Rep.* ii. 341.—Mexic.; Guatemala.
phyllanthoides, *Link, Handb.* ii. 11; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 55.—Mexic.
Phyllanthus, *Link, l. c.*; *Salm-Dyck, l. c.* 56.—Am. trop.
purpureus, *Hort. ex Haage, Verz. Cact.* 27.—Hab.?
Russelianus, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 37.—Bras.
Selloii ×, Hort. Sello, ex Labour. Monog. Cact. 429, nomen.
serratus, *Brongn. ex Labour. l. c.* 417.—Mexic.
stenopetalus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 228.—Hab.?
strictus, *Lem. Illustr. Hortic.* i. (1854) *Misc.* 107.—Cuba.
triumphans, *Hort. Angl. ex Haage, Cact. Verz.* (1857) 10, nomen.—Hab.?

PHYLLOCALYMMA, Benth. in Enum. Pl. Hueg. 61 (1837) = **Angianthus**, Wendl. (Compos.).
filaginoides, Steetz, in Lehm. Pl. Preiss. i. 437 = A. micropoides.
micropodioides, Benth. in Enum. Pl. Hueg. 62 = A. micropoides.

PHYLLOCALYX, Berg, in Linnaea, xxvii. (1854) 306 = **Eugenia**, Mich. (Myrtac.).
calycinus, Berg, l. c. 307 = E. calycina, *Cambess.*
calystegius, Berg, in Mart. Fl. Bras. xiv. 1. 331.—Bras.
cerasiflorus, Berg, in Linnaea, xxvii. (1854) 308.—Bras.
cupreus, Berg, in Mart. Fl. Bras. xiv. 1. 592.—Bras.
edulis, Berg, l. c. 327 = E. Selloi.
formosus, Berg, in Linnaea, xxvii. (1854) 307 = E. formosa, *Cambess.*
grandifolius, Berg, in Mart. Fl. Bras. xiv. 1. 333.—Bras.
herbaceus, Berg, l. c. 328.—Bras.
involucratus, Berg, in Linnaea, xxvii. (1854) 307 = E. involucrata.
laevigatus, Berg, in Mart. Fl. Bras. xiv. 1. 329.—Bras.
ligustrinus, Berg, l. c. 592.—Bras.
limbatus, Berg, l. c. 332.—Bras.
Luschnathianus, Berg, l. c. 333 = E. Luschnathiana.
macrosepalus, Berg, l. c. 332 = E. macrocalyx.
marginatus, Berg, l. c.—Bras.
membranaceus, Berg, l. c. 334.—Bras.
pubescens, Berg, l. c. 592.—Bras.
racemosus, Berg, l. c. 334.—Bras.
Regelianus, Berg, l. c. 591.—Bras.
retusus, Berg, l. c. 331.—Bras.
Riedelianus, Berg, l. c. 590.—Bras.
speciosus, Berg, in Linnaea, xxvii. (1854) 307 = E. speciosa.
strictus, Berg, in Mart. Fl. Bras. xiv. 1. 330.—Bras.
tomentosus, Berg, in Linnaea, xxvii. (1854) 307 = E. tomentosa.

PHYLLOCALYX, A. Rich. Tent. Fl. Abyss. i. 160. t. 34 (1847) = **Crotalaria**, Tourn. (Legumin.).
Quartinianus, A. Rich. l. c. = C. platycalyx.

PHYLLOCARPA, Nutt. ex Moq. in DC. Prod. xiii. 11. 108 (1849) = **Atriplex**, Tourn. (Chenop.).
spinosa, Nutt. l. c. = A. spinosa.

PHYLLOCARPUS. Ried. ex Endl. Gen. Suppl. ii. 97 (1842); Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 142. *LEGUMINOSAE*, Benth. & Hook. f. i. 564.
Pterocarpus, *Ried. ex Endl. Gen. Suppl.* ii. 97.—Bras.
Riedelii, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 143 = P. Pterocarpus.

PHYLLOCEPHALIUM, Miq. Fl. Ind. Bat. ii. 20 (1856) = seq.

PHYLLOCEPHALUM, Blume, Bijdr. 888 (1826) = **Centratherum**, Cass. (Compos.).
frutescens, Blume, l. c. 889 (= *Decaneurum frutescens*).—Java.

PHYLLOCEREUS, Miq. in Bull. Sc. Phys. et Nat. Brux. (1839) 112 = **Phyllocactus**, Link (Cact.).

PHYLLOCHLAMYS, Bur. in DC. Prod. xvii. 217 (1873). *URTICACEAE*, Benth. & Hook. f. iii. 359.
spinosa, *Bur. l. c.* 218.—Ind. or.; Malaya.

PHYLLOCLADUS, Rich. Conif. 129. t. 3 (1826). *CONIFERAE*, Benth. & Hook. f. iii. 432.
BROWNETERA, Rich. in Ann. Mus. Par. xvi. (1810) 299, nomen prius.
ROBERTIA, Rich. ex Carr. Conif. ed. I. 498 (1855).
THALAMIA, Spreng. Anleit. ii. 218 (1817).
alpinus, Hook. f. Fl. N. Zel. i. 235. t. 53 = trichomanoides.
aspleniifolia, Hook. f. in Hook. Lond. Journ. Bot. iv. (1845) 151 = rhomboidalis.
Billardieri, Mirb. in Mém. Mus. Par. xiii. (1825) 76 = rhomboidalis.
Cunninghami, Hort. ex Carr. Conif. ed. II. 707 = trichomanoides.
glauca, Hort. ex Carr. Conif. ed. I. 502 = trichomanoides.
glauca, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 149; ed. II. (1871) 94.—N. Zel.
hypophylla, *Hook. f. in Hook. Ic. Pl.* t. 889.—Borneo.
hypophylla, Hort. ex Carr. Conif. ed. II. 707 = trichomanoides.
rhomboidalis, *Rich. Conif.* 130. t. 3. f. 2.—Tasman.
rhomboidalis, A. Rich. Fl. N. Zél. 363 = trichomanoides.
serratifolia, Nois. ex Gord. Pinet. Suppl. 43 = rhomboidalis.
trichomanoides, *D. Don, in Lamb. Pin.* ed. II. App.—N. Zel.

PHYLLOCORYNE, Hook. f. in Trans. Linn. Soc. xxii. (1856) 51. t. 11 = **Scybalium**, Schott (Balanoph.).
jamaicensis, Hook. f. l. c. = S. jamaicense.

PHYLLOCOSMUS, Klotzsch, in Abh. Akad. Berl. 1856 (1857) 232. t. 1. *LINEAE*, Benth. & Hook. f. i. 245.
PENTACOCCA, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 600.
africanus, *Klotzsch, in Abh. Akad. Berl.* 1856 (1857) 233.—Afr. trop.
sessiliflorus, *Oliver, Fl. Trop. Afr.* i. 273.—Afr. trop.

PHYLLOCYCLUS, Kurz, in Journ. Soc. Beng. xlii. (1873) ii. 235 = **Canscora**, Lam. (Gentian.).
Helferiana, Kurz, l. c. 236 = C. Helferiana.
Parishii, Kurz, l. c. = C. Parishii.

PHYLLOCYTISUS, Fourn. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 358 = **Cytisus**, Linn. (Legumin.).
sessilifolius, Fourr. l. c. = C. sessilifolius.

PHYLLODES, Lour. Fl. Cochinch. 13 (1790) = **Phrynium**, Willd. (Scitam.).
placentaria, Lour. l. c. = P. capitatum.

PHYLLODIUM, Desv. Journ. Bot. i. (1813) 123. t. 5 = **Desmodium**, Desv. (Legumin.).
elegans, Desv. in Mém. Soc. Linn. Par. iv. (1826) 324 = D. elegans.
lutescens, Desv. Journ. Bot. i. (1813) 124 = D. lutescens.
pulchellum, Desv. in Mém. Soc. Linn. Par. iv. (1826) 324 = seq.
pulchrum, Desv. Journ. Bot. i. (1813) 124. t. 5. f. 24 = D. pulchellum.
vestitum, Benth. in Miq. Pl. Jungh. 217 = D. vestitum.

PHYLLODOCE, Link, Handb. ii. 132 (1831) = **Acacia**, Tourn. (Legumin.).
alata, Link, l. c. = A. alata.
angustifolia, Link, l. c. 133 = A. suaveolens.
armata, Link, l. c. = A. armata.
decipiens, Link, l. c. 132 = A. decipiens.
dodonaeifolia, Link, l. c. 133 = A. dodonaeifolia.
floribunda, Link, l. c. = A. longifolia.
genistaefolia, Link, l. c. = A. diffusa.

PHYLLODOCE:—
juniperina, Link, Handb. ii. 133 = Acacia juniperina.
linifolia, Link, l. c. = A. linifolia.
longifolia, Link, l. c. = A. longifolia.
myrtifolia, Link, l. c. = A. myrtifolia.
Sophora, Link, l. c. = A. longifolia?
stricta, Link, l. c. = A. stricta.
suaveolens, Link, l. c. = A. suaveolens.
undulata, Link, l. c. 132 = A. armata.
verticillata, Link, l. c. 133 = A. verticillata.

PHYLLODOCE, Salisb. Parad. Lond. t. 36 (1806) = **Bryanthus**, S. G. Gmel. (Ericac.).
caerulea, Bab. Man. Brit. Bot. ed. I. 194 = B. taxifolius.
empetriformis, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 160 = B. empetriformis.
Pallasiana, D. Don, l. c. 159 = B. aleuticus.
taxifolia, Salisb. Parad. Lond. t. 36 = B. taxifolius.

PHYLLODOLON, Salisb. Gen. Pl. Fragm. 90 (1866) = **Allium**, Tourn. (Liliac.).
fistulosum, Salisb. l. c. = A. fistulosum.

PHYLLOGLOTTIS, Salisb. Gen. Pl. Fragm. 15 (1866) = **Eriospermum**, Jacq. (Liliac.).
foliolifera, Salisb. l. c. = E. folioliferum.

PHYLLOLOBIUM, Fisch. in Spreng. Nov. Prov. 33 (1818) = **Astragalus**, Tourn. (Legumin.).
chinense, Fisch. in Spreng. l. c. = A. complanatus.
zanzibarense, Boj. ex Loud. Hort. Brit. 285, nomen (Quid?).—Afr. trop.

PHYLLOMA, Ker-Gawl. in Bot. Mag. t. 1585 (1813) = **Lomatophyllum**, Willd. (Liliac.).
aloiflorum, Ker-Gawl. l. c. = L. borbonicum.
aloifolium, Steud. Nom. ed. II. ii. 328, sphalm. = praec.
borbonicum, Schult. f. Syst. vii. 361 = L. borbonicum.
Jacquini, G. Don, in Sweet, Hort. Brit. ed. III. 685 = Aloë hexapetala.
macrum, Sweet, l. c. ed. I. 423 = L. macrum.
rufocinctum, Sweet, l. c. = L. rufocinctum.

PHYLLOMATIA, Benth. in Ann. Wien. Mus. ii. (1838) 113 = **Rhynchosia**, Lour. (Legumin.).

PHYLLOMELIA, Griseb. Cat. Pl. Cub. 139 (1866). *RUBIACEAE*, Benth. & Hook. f. ii. 116.
coronata, *Griseb. l. c.*—Cuba.

PHYLLOMERIA, Griseb. Ges. Abh. 282 (1880), sphalm. = praec.

PHYLLONOMA, Willd. ex Schult. Syst. vi. p. xx (1820). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
DULONGIA, H. B. & K. Nov. Gen. et Sp. vii. 76. t. 623 (1825).
ruscifolia, *Willd. ex Schult. Syst.* vi. 210.—Mexic.

PHYLLOPAPPUS, Walp. in Linnaea, xiv. (1840) 507 = **Microseris**, D. Don (Compos.).
lanceolatus, Walp. l. c. = M. Forsteri.

PHYLLOPODIUM, Benth. in Hook. Comp. Bot. Mag. i. (185) 372. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 944.
bracteatum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 373.—Afr. austr.
capitatum, *Benth. l. c.*—Afr. austr.
cuneifolium, *Benth. l. c.*—Afr. austr.
diffusum, *Benth. l. c.*—Afr. austr.
heterophyllum, *Benth. l. c.*—Afr. austr.
Krebsianum, *Benth. in DC. Prod.* x. 353.—Afr. austr.
pumilum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 373.—Afr. austr.

PHYLLOPUS, DC. Prod. iii. 177 (1828) = **Henriettea**, DC. (Melastom.).
Martii, Mart. Nov. Gen. et Sp. iii. 143. t. 275 = seq.
Martiusii, DC. Prod. iii. 178 = H. Martii.

PHYLLORHACHIS, Trimen, in Journ. Bot. xvii. (1879) 353. t. 205. *GRAMINEAE*, Benth. & Hook. f. iii. 1108.
sagittata, *Trimen, l. c.* 355.—Afr. trop.

PHYLLOSCHOENUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 171 = **Juncus**, Tourn.
acutiflorus, Fourr. l. c. 172 = J. acutiflorus.
alpinus, Fourr. l. c. = J. alpinus.
anceps, Fourr. l. c. = J. anceps.
lagenarius, Fourr. l. c. = J. Fontanesii.
lamprocarpus, Fourr. l. c. = J. lampocarpus.
obtusiflorus, Fourr. l. c. = J. obtusiflorus.
striatus, Fourr. l. c. = J. striatus.
supinus, Fourr. l. c. 171 = J. supinus.

PHYLLOSPADIX, Hook. Fl. Bor. Am. ii. 171. t. 186 (1838). *NAIADACEAE*, Benth. & Hook. f. iii. 1017.
Scouleri, *Hook. Fl. Bor. Am.* ii. 171. t. 186.—Am. bor. occ.
serrulatus, *Rupr. ex Aschers. in Linnaea*, xxxv. (1867–68) 169.—Alaska.
Torreyi, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 303. —Am. bor. occ.

PHYLLOSTACHYS, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 745. t. 5. *GRAMINEAE*, Benth. & Hook. f. iii. 1208.
aurea, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 716.—China; Japon.
bambusoides, *Sieb. & Zucc. in Abh. Akad. Muench.* iii. II. (1843) 745. t. 5. f. 3.—Japon.
edulis, A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 623 = Bambusa mitis.
flexuosa, *A. & C. Rivière, l. c.* 758.—China.
Fortunei, *A. & C. Rivière, l. c.* 301.—China.
Kumasaca, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 39.—Japon.
macrantha, Sieb. & Zucc. in Flora, xxix. (1846) 34 = bambusoides.
Mazeli, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 697 = P. Quilioi.
megastachya, Steud. in Flora, xxix. (1846) 21 = bambusoides.
mitis, A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 689 = Bambusa mitis.
Newinii, *Hance, in Journ. Bot.* xiv. (1876) 294.—China.
nidularia, *Munro, in Gard. Chron.* (1876) II. 773, 774. —Japon.
nigra, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 38.—China; Japon.
puberula, Munro, in Gard. Chron. (1876) II. 773, 774 = nigra.
Quilioi, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 697.—Japon.
reticulata, C. Koch, Dendrol. ii. II. 356 = bambusoides.
Stauntoni, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 37.—China.
sulphurea, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 773.—Hab.?
violascens, *A. & C. Rivière, l. c.* 770.—Hab.?
viridi-glaucescens, A. & C. Riviere, l. c. 700 = nana.

PHYLLOSTACHYS, Torr. in Ann. Lyc. N. York, iii. (1836) 404, in obs. = **Carex**, Linn. (Cyperac.).

PHYLLOSTEGIA, Benth. in Bot. Reg. sub t. 1292 (1830). *LABIATAE*, Benth. & Hook. f. ii. 1216.
brevidens, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 343.—Ins. Sandvic.
Chamissonis, Benth. in Linnaea, vi. (1831) 79 = glabra.
clavata, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
dentata, Benth. l. c. = vestita.
floribunda, *Benth. Lab. Gen. et Sp.* 653.—Ins. Sandvic.
glabra, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
grandiflora, *Benth. l. c.*—Ins. Sandvic.
Haliakalae, Wawra, in Flora, lv. (1872) 530 = mollis.
haplostachya, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 345.—Ins. Sandvic.
Hillebrandi, *H. Mann, in Mem. Bost. Soc. Nat. Hist.* i. (1866–69) 536.—Ins. Sandvic.
hirsuta, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
honolulensis, Wawra, in Flora, lv. (1872) 531 = parviflora.

PHYLLOSTEGIA:—
leptostachys, Benth. in Bot. Reg. sub t. 1292 = parviflora.
Macraei, Benth. in DC. Prod. xii. 554 = glabra.
macrophylla, Benth. in Bot. Reg. sub t. 1292 = parviflora.
microphylla, Benth. in Linnaea, vi. (1831) 570 (Verbenacea).—Unalask.
mollis, *Benth. l. c.* 79.—Ins. Sandvic.
parviflora, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
racemosa, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
stachyoides, *A. Gray, in Proc. Am. Acad.* (1861–62) 344.—Ins. Sandvic.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 56.—Ins. Tahiti.
truncata, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 345.—Ins. Sandvic.
vestita, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
Waimeae, *Wawra, in Flora*, lv. (1872) 531.—Ins. Sandvic.

PHYLLOSTEMA, Neck. Elem. ii. 301 (1790) = **Simaba**, Aubl. (Simarub.).

PHYLLOSTYLON, Capan. ex Benth. & Hook. f. Gen. iii. 352 (1880). *URTICACEAE*, Benth. & Hook. f. l. c.
brasiliensis, *Capan. l. c.*—Bras.

PHYLLOTA, Benth. in Enum. Pl. Hueg. 33 (1837). *LEGUMINOSAE*, Benth. & Hook. f. i. 470.
aspera, Benth. in Ann. Wien. Mus. ii. (1838) 77 = phylicoides.
barbata, *Benth. in Enum. Pl. Hueg.* 33.—Austral.
Baueri, Benth. in Ann. Wien. Mus. ii. (1838) 77 = phylicoides.
Billardieri, Benth. l. c. = phylicoides.
comosa, Benth. l. c. = phylicoides.
diffusa, F. Muell. Fragm. i. 8 = Pultenaea diffusa.
gracilis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) I. 267.—Austral.
grandiflora, Benth. in Ann. Wien. Mus. ii. (1838) 77 = phylicoides.
humifusa, *Benth. Fl. Austral.* ii. 95.—Austral.
Luehmanni, *F. Muell. Fragm.* x. 33.—Austral.
phylicoides, *Benth. in Ann. Wien. Mus.* ii. (1838) 77. —Austral.
pilosa, Benth. l. c. = phylicoides.
pleurandroides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 38.—Austral.
squarrosa, Benth. in Ann. Wien. Mus. ii. (1838) 77 = phylicoides,
Sturtii, *Benth. Fl. Austral.* ii. 95.—Austral.
Urodon, F. Muell. in Papers Roy. Soc. Tasm. (1877) 120 = Pultenaea Urodon.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 267 = barbata.

PHYLLOTAENIUM, André, in Illustr. Hortic. xix. (1872) t. 88 = **Xanthosoma**, Schott (Aroid.).
Lindeni, André, l. c. = X. Lindeni.
mirabile, Hort. ex Mast. in Gard. Chron. (1874) II. 258 = X. mirabile.

PHYLLOTHECA, Nutt. ex Moq. in DC. Prod. xiii. II. 98 (1849) = **Atriplex**, Tourn. (Chenop.).
parvifolia, Nutt. ex Moq. l. c. = A. californica.

PHYLLOXYLON, Baill. Adansonia, ii. (1861) 54. *LEGUMINOSAE*.
NEOBARONIA, Baker, in Journ. Linn. Soc. xxi. (1884) 336.
decipiens, *Baill. Adansonia*, ii. (1861) 54.—Ins. Maurit.

PHYLLYMENA, Blume, ex Miq. Fl. Ind. Bat. ii. 83 (1856) = PHYLLIMENA, Blume = **Enhydra**, Lour. (Compos.).

PHYLLYREA, G. Don, Gen. Syst. iv. 44 (1837), sphalm. = **Phillyrea**, Linn. (Oleac.).

PHYMASPERMUM, Less. Syn. Comp. 253 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 422.
ADENACHAENA, DC. Prod. vi. 49 (1837).
JACOSTA, DC. l. c. (1837).
OLIGOGLOSSA, DC. l. c. (1837).
aciculare, *Benth. & Hook. f. Gen.* ii. 423.—Afr. austr.
carnosulum, *Benth. & Hook. f. l. c.*—Afr. austr.
junceum, *Less. Syn. Comp.* 253.—Afr. austr.
leptophyllum, *Benth. & Hook. f. Gen.* ii. 423.—Afr. austr.
parvifolium, *Benth. & Hook. f. l. c.*—Afr. austr.

PHYMATANTHUS, Sweet, Geran. t. 43 (1820) = **Pelargonium**, L'Hérit. (Geran.).
elatus, Sweet, l. c. t. 96 = Pelarg. elatum.
grandiflorus ×, Sweet, Hort. Brit. ed. I. 75, nomen.
intertinctus ×, G. Don, in Sweet, l. c. ed. III. 108, nomen.
latifolius ×, Colville, ex Sweet, l. c. ed. I. 75, nomen.
Serra, Hoffmgg. Verz. Pfl. 99 = Pelarg. Serra.
tricolor, Sweet, Geran. t. 43 = Pelarg. violareum.
villosus ×, Sweet, Hort. Brit. ed. I. 76, nomen.

PHYMATIDIUM, Lindl. Gen. et Sp. Orch. 209 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 569.
delicatulum, *Lindl. l. c.* 210.—Bras.
denticulatum, Steud. Nom. ed. II. ii. 328, sphalm. = praec.
falcifolium, *Lindl. Gen. et Sp. Orch.* 210.—Bras.
hysteranthum, *Rodrig. Orch. Nov.* ii. 228.—Bras.
myrtophilum, *Rodrig. l. c.* 229.—Bras.
tillandsioides, *Rodrig. l. c.* 228.—Bras.

PHYMATIS, E. Mey. in Drège, Zwei Pfl. Docum. 211 (1843) = **Carum**, Rupp. (Umbell.).
cyminoides, E. Mey. l. c. = C. hispidum.

PHYMATOCARPUS, F. Muell. Fragm. iii. 120 (1862). *MYRTACEAE*, Benth. & Hook. f. i. 706.
Maxwelli, *F. Muell. l. c.* ix. 45.—Austral.
porphyrocephalus, *F. Muell. l. c.* iii. 121.—Austral.

PHYMOSIA, Desv. in Ham. Prod. Pl. Ind. Occ. 49 (1825) = **Sphaeralcea**, A. St. Hil. (Malvac.).
abutiloides, Desv. l. c. 50 = S. abutiloides.

PHYODINA, Rafin. Fl. Tellur. ii. 16 (1836) = **Tradescantia**, Linn. (Commel.).
gracilis, Rafin. l. c. = T. gracilis.

PHYRRHEIMA, Hassk. in Flora, liv. (1871) 257 = PYRRHEIMA, Hassk. = **Tradescantia**, Linn. (Commel.).

PHYSA, Thou. Gen. Nov. Madag. 20 (1806) = **Mollugo**, Linn. (Ficoid.).
madagascariensis, DC. Prod. i. 393.—Madag.

PHYSACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1085 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.

PHYSALIDIUM, Fenzl, in Tchihat. Asie Min. Bot. i. 327 (1860). *CRUCIFERAE*, Benth. & Hook. f. i. 88.
stylosum, *Fenzl, l. c.*—Persia.

PHYSALIS, Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 890.
ALICABON, Rafin. Sylva Tellur. 56 (1838).
ALKEKENGI, Tourn. ex Adans. Fam. ii. 218 (1763).
EPETORHIZA, Steud. Nom. ed. II. i. 556 (1840).
EPETETIORHIZA, Steud. l. c. ii. 328 (1841).
EXODECONUS, Rafin. Sylva Tellur. 57 (1838).
HERSCHELIA, Bowdich, Excurs. Mad. 159 (1825).
MEGISTA, Fourr. in Ann. Soc. Linn. N. S. xvii. (1869) 115.
PENTAPHILTRUM, Reichb. Nom. 121 (1841).
QUINCULA, Rafin. Atl. Journ. 145 (1832).
abyssinica, Hort. Berol. ex Nees, in Linnaea, vi. (1831) 448 = minima.
adulterina, *Fisch. ex Steud. Nom.* ed. I. 516, nomen.—Hab.?
aequata, *Jacq. f. ex Nees, in Linnaea*, vi. (1831) 470. —Am. bor.; Ind. or.
Alkekengi, *Linn. Sp. Pl.* 183.—Europ.; Japon.
Alkekengi, Lour. Fl. Cochinch. 133 = minima.
allogona, *Ind. Sem. Hort. Lips.* (1840) 3; *ex Dun. in DC. Prod.* xiii. I. 452.—Hab.?

PHYSALIS:—
Alpini, Jacq. Eclog. i. 39, in obs. = Withania somnifera.
angulata, Heyne, ex Wall. Cat. n. 2634 = minima.
angulata, *Linn. Sp. Pl.* 183.—Reg. calid.
angulata, Ruiz & Pav. Fl. Per. ii. 42 = lanceifolia.
angulata, Walt. Fl. Carol. 99 = obscura.
angustifolia, *Nutt. in Journ. Acad. Philad.* vii. (1834) 113.—Florida.
arborescens, *Linn. Sp. Pl.* ed. II. 261.—Mexic.
arborescens, Thunb. Prod. Pl. Cap. 37 = Withania somnifera.
arenaria, Hort. ex Nees, in Linnaea, vi. (1831) 471 = angulata.
arenaria, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 35.—Bras.
aristata, Ait. Hort. Kew. ed. I. i. 244 = Withania aristata.
atriplicifolia, Poir. Encyc. Suppl. ii. 348 = philadelphica.
barbadensis, Herb. Wight, ex Wall. Cat. n. 2634 = minima.
barbadensis, Jacq. Misc. ii. 359 = pubescens.
barbadensis, Lam. Encyc. ii. 102 = peruviana.
begonifolia, Roxb. in Beats. Tracts St. Helena, App. 317 = Mellissia begonifolia.
brasiliensis, Sendt. in Mart. Fl. Bras. x. 133 = obscura?
bulbosa, Hedw. ex Steud. Nom. ed. II. ii. 329 = chenopodifolia.
campechiana, Linn. Syst. ed. X. 933 = arborescens.
campechiensis, Hill, Veg. Syst. ix. 27 = praec.
capsicifolia, Dun. in DC. Prod. xiii. 1. 449 = angulata.
cardiophylla, Torr. in Bot. Mex. Bound. 153 = crassifolia.
Carpenterii, Riddell, Cat. Fl. Ludov.; N. Orl. Med. & Surg. Journ. viii. (1852) 758 = (Athenaeae sp.)—Am. bor.
chenopodifolia, *Lam. Illustr.* ii. 28.—Peruv.
chenopodiifolia, Willd. Sp. Pl. i. 1023 = philadelphica.
ciliata, Sieb. & Zucc. Fl. Jap. Fam. Nat. ii. 22 = angulata.
cordata, *Mill. Gard. Dict.* ed. VIII. n. 14.—Mexic.
cordifolia, *Dun. in DC. Prod.* xiii. 1. 441.—Hab.?
corymbosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = Cardiospermum Halicacabum.
Coztomatl, *Moç. & Sesse, ex Dun. in DC. Prod.* xiii. II. 450.—Mexic.
crassifolia, *Benth. Bot. Voy. Sulph.* 40.—Calif.
crenata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = minima.
curassavica, *Linn. Sp. Pl.* 182.—Venezuela.
curassavica, Willd. Sp. Pl. i. 1020 = viscosa.
daturaefolia, Lam. Encyc. ii. 102 = Nicandra physaloides.
decumbens, *Dun. in DC. Prod.* xiii. 1. 436.—Hab.?
dentata, *Pav. ex Dun. l. c.* 441.—Mexic.
diffusa, Willd. Herb. ex Nees, in Linnaea, vi. (1831) 480 = minima.
divaricata, D. Don, Prod. Fl. Nep. 97 = minima.
dubia, Link, Enum. Hort. Berol. i. 181 = angulata.
Dunali, *Montr. in Mém. Acad. Lyon*, x. (1860) 245.—N. Caled.
edulis, Sims, Bot. Mag. t. 1068 = peruviana.
Elliottii, Kunze, in Linnaea, xx. (1847) 33 = lanceolata.
esculenta, Salisb. Prod. 132 = peruviana.
Fendleri, *A. Gray, in Proc. Am. Acad.* x. (1874) 66.—Am. bor. occ.
flexuosa, Linn. Sp. Pl. 182 = Withania somnifera.
flexuosa, Russ. ex Wall. Cat. n. 2634 = angulata.
foetens, *Poir. Encyc. Suppl.* ii. 348.—Am. trop.
foetidissima, Hort. Madr. ex Dun. in DC. Prod. xiii. 1. 436 = viscosa.
foetidissima, Lag. Gen. et Sp. Nov. 10 = foetens.
frutescens, DC. Fl. Fr. iii. 611 = Withania frutescens.
fusco-maculata, *Dun. in DC. Prod.* xiii. 1. 437.—Reg. Argent.?
Galeottiana, *Walp. Rep.* vi. 574.—Mexic.
geniculata, *Miers, Illustr. S. Am. Pl.* ii. *App.* 56.—Mexic.
glabra, *Benth. Bot. Voy. Sulph.* 39.—Calif.
glabra, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 132 = Galeottiana.
glabriuscula, *Dun. in DC. Prod.* xiii. 1. 436.—Bolivia.
glutinosa, *Schlecht. in Linnaea*, xix. (1847) 310.—Mexic.
gracilis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 37.—Mexic.

PHYSALIS:—
grandiflora, *Hook. Fl. Bor. Am.* ii. 90.—Am. bor.
Halicacabum, Crantz, Inst. ii. 370; Scop. Fl. Carn. ed. II. i. 160 = P. Alkekengi.
Halicacabum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = Cardiospermum Halicacabum.
hederaefolia, *A. Gray, in Proc. Am. Acad.* x. (1875) 65.—Am. bor. occ.
Hermanni, Dun. in DC. Prod. xiii. 1. 444 = minima.
heterophylla, Nees, in Linnaea, vi. (1831) 463, partim = virginiana.
hirsuta, Dun. in DC. Prod. xiii. 1. 445 = pubescens.
hirsuta, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 132 = Athenaea hirsuta.
hyemalis, Salisb. Prod. 132 = P. Alkekengi.
hygrophila, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 86.—Bras.
incana, Desf. Tabl. ed. I. 68 = chenopodifolia.
incana, Hort. Par. ex Dun. in DC. Prod. xiii. 1. 440 = peruviana.
indica, Lam. Encyc. ii. 102 = minima.
integrifolia, Pav. ex Dun. in DC. Prod. xiii. 1. 438 = curassavica.
ixocarpa, Brot. ex DC. Cat. Hort. Monsp. 50, nomen; et ex Hornem. Hort. Hafn. Suppl. 26 = aequata.
ixocarpa, Hort. Bonn. ex Nees, in Linnaea, vi. (1831) 471 = angulata.
Jacquini, Link, Enum. Hort. Berol. i. 180 = viscosa.
laevigata, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 131.—Mexic.
Lagascae, Roem. & Schult. Syst. iv. 679 = minima.
lanceifolia, *Nees, in Linnaea*, vi. (1831) 473.—Am. austr.
lanceifolia, Rugel, ex Kunze, in Linnaea, xx. (1847) 33 = lanceolata.
lanceolata, Ell. Sketch, i. 278 = viscosa.
lanceolata, *Michx. Fl. Bor. Am.* i. 149.—Am. bor. occ.
latifolia, Lam. Illustr. ii. 29 = peruviana.
laurifolia, D. Dietr. Syn. Pl. i. 695 = lanceolata.
limensis, Retz. Obs. v. 22 = prostrata.
Linkiana, Nees, in Linnaea, vi. (1831) 471 = angulata.
lobata, *Torr. in Ann. Lyc. N. York*, ii. (1826) 226.—Am. bor. occ.
longifolia, Nutt. in Trans. Am. Phil. Soc. v. (1837) 193 = lanceolata.
lurida, Salisb. Prod. 132 = Scopola physaloides.
lutea, *Medic. in Act. Acad. Theod. Palat.* iv. *Phys.* (1780) 186.—Hab.?
maritima, M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 407 = viscosa.
maurelloides, *Schlecht. in Linnaea*, xxvii. (1854) 508.—N. Granat.
maxima, *Mill. Gard. Dict.* ed. VIII. n. 15.—Mexic.
megistocarpa, Zuccagni, in Roem. Collect. 130 = philadelphica.
mendocina, *Phil. in Anal. Univ. Chil.* (1862) II. 402.—Chili.
micrantha, Link, Enum. Hort. Berol. i. 181 = minima.
minima, *Linn. Sp. Pl.* 183.—Geront. trop.
minutiflora, *Moç. & Sesse, ex Dun. in DC. Prod.* xiii. 1. 451.—Mexic.
miraflorensis, *Dun. l. c.* 443.—Bolivia.
mollis, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 194.—Texas; N. Mexic.
mollis, Torrey, Bot. Mex. Bound. 153 = Fendleri, hederaefolia.
mucronata, *Willd. Enum. Hort. Berol. Suppl.* 11, nomen.—Hab.?
muricata, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 209.—Calif.
Neesiana, *Sendt. in Mart. Fl. Bras.* x. 131.—Bras.
nicandroides, *Schlecht. in Linnaea*, xix. (1847) 311.—Mexic.
nodosa, Lam. Illustr. ii. 29 = foetens.
nolanoides, *A. DC. in DC. Prod.* xiii. 1. 690.—Peruv.
nutans, J. F. Gmel. Syst. 382 = viscosa.
nyctaginea, Dun. in DC. Prod. xiii. 1. 440 = virginiana.
obscura, *Michx. Fl. Bor. Am.* i. 149.—Am. bor.
obscura, Torr. & Fras. ex Dun. in DC. Prod. xiii. 1. 440 = virginiana.
ochroleuca, Salisb. Prod. 132 = viscosa.
origanifolia, *Lam. Illustr.* ii. 28.—Bras.
orinocensis, H. B. & K. Nov. Gen. et Sp. iii. 12 = Withania orinocensis.
Orizabae, *Dun. in DC. Prod.* xiii. 1. 452.—Mexic.
ovata, Poir. Encyc. Suppl. ii. 347 = philadelphica.
Palmeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 235.—Calif.

PHYSALIS:—
parviflora, R. Br. Prod. 447 = minima.
parviflora, *Hort. ex Zuccagni, in Roem. Collect.* 130.—Hab.?
patula, *Mill. Gard. Dict.* ed. VIII. n. 12.—Mexic.
pedunculata, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 132.—Mexic.
pennsylvanica, Willd. ex Nees, in Linnaea, vi. (1831) 463 = virginiana.
pensylvanica, Linn. Sp. Pl. ed. II. 1670 = viscosa.
peruviana, *Linn. l. c.*—Reg. calid.
peruviana, Mill. Gard. Dict. ed. VIII. n. 16 = Nicandra physaloides.
philadelphica, *Lam. Encyc.* ii. 101.—Am. bor.
prostrata, *Domb. ex L'Hérit. Stirp. Nov.* 43. t. 22.—Peruv.
pruinosa, Ell. Sketch, i. 279 = obscura.
pruinosa, Linn. Sp. Pl. 184 = pubescens.
Pseudo-angulata, Blume, Bijdr. 706 = minima.
pubescens, R. Br. Prod. 447 = peruviana.
pubescens, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 227 = viscosa.
pubescens, Hort. ex Dun. in DC. Prod. xiii. 1. 439 = chenopodifolia.
pubescens, *Linn. Sp. Pl.* 183.—Am. bor.
pubescens, Willd. ex Nees, in Linnaea, vi. (1831) 479 = minima.
pumila, Nutt. in Trans. Am. Phil. Soc. v. (1837) 193 = lanceolata.
ramosa, *Forsk. Fl. Aegypt. Arab.* 204.—Arabia.
ramosa, Mill. Gard. Dict. ed. VIII. n. 9 = pubescens.
ramosa, Xaares, ex Zuccagni, in Roem. Collect. 129 = Saracha Zuccagniana.
ramosissima, Mill. ex Dun. in DC. Prod. xiii. 1. 449 = angulata.
Rothiana, Herb. Berol. ex Nees, in Linnaea, vi. (1831) 467 = pubescens.
Rothiana, Roem. & Schult. Syst. iv. 677 = minima.
sabeana, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 6 = lobata.
Sagunda, Walp. Rep. iii. 23, sphalm. = *P. Sugunda*.
Sancti-Josephi, *Dun. in DC. Prod.* xiii. 1. 451.—Mexic.
scariosa, Webb & Berth. Phyt. Canar. iii. 120 = Withania somnifera.
Schiedeana, *Dun. in DC. Prod.* xiii. 1. 452.—Mexic.
Schraderiana, Bernh. in Linnaea, xiii. (1839) 361 = Saracha viscosa.
Sendtneri, *Dun. in DC. Prod.* xiii. 1. 438.—Bosnia.
solanacea, Mert. ex Roth, Catalect. ii. 26 = Saracha procumbens.
somnifera, Linn. Sp. Pl. 182 = Withania somnifera.
spectabilis, Salisb. Prod. 133 = Scopola physaloides.
staminea, Muhl. ex Nees, in Linnaea, vi. (1831) 467 = pubescens.
stramonifolia, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 242 = Scopola lurida.
suberifera, *Dun. in DC. Prod.* xiii. 1. 442.—Bras.
suberosa, Cav. Ic. ii. 1 = Withania frutescens.
subtriflora, Ruiz & Pav. Fl. Per. ii. 42. t. 178 = Withania subtriflora.
Sugunda, Buch.-Ham. ex Wall. Cat. n. 2635 = Withania somnifera.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 741.—Guian.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 337 = minima.
tomentosa, Medic. in Act. Acad. Theod. Palat. iv. Phys. (1780) 184. t. 4 = peruviana.
tomentosa, Thunb. Prod. Pl. Cap. 37 = Withania somnifera.
tomentosa, Walt. Fl. Carol. 99 = viscosa.
tuberosa, *Cav. Desc.* 370.—Hispan.
tuberosa, Willd. Enum. Hort. Berol. 232 = chenopodifolia.
tuberosa, Zuccagni, in Roem. Collect. 130 = peruviana.
turbinata, Medic. in Act. Acad. Theod. Palat. iv. Phys. (1780) 189. t. 5 = pubescens.
urceolata, Ruiz & Pav. ex Dun. in DC. Prod. xiii. 1. 475 (= *Salpichroma urceolatum*).—Peruv.
villosa, *Mill. Gard. Dict.* ed. VIII. n. 13.—Mexic.
villosa, Moench, ex Steud. Nom. ed. II. ii. 329 = Withania somnifera.
villosa, Roth, Nov. Pl. Sp. 122 = minima.
violacea, *Carr. in Rev. Hortic.* (1882) 216.—Hab.?
virginiana, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. bor.
virginica, A. Gray, in Proc. Am. Acad. x. (1875) 65 = praec.

PHYSALIS:—
viscido-pubescens, Dun. in DC. Prod. xiii. 1. 442 = virginiana.
viscosa, Ell. Sketch, i. 279 = pubescens.
viscosa, A. Gray, Man. Bot. ed. I. 354 = virginiana.
viscosa, *Linn. Sp. Pl.* 183.—Am. calid.
Walteri, Nutt. in Journ. Acad. Philad. vii. (1834) 112 = viscosa.
Wrightii, *A. Gray, in Proc. Am. Acad.* x. (1875) 63. —Texas.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* iii. 13 = Athenaea xalapensis.

PHYSALOBIUM, Steud. Nom. ed. II. ii. 330 (1841) = PHYSOLOBIUM, Benth. = **Kennedya**, Vent. (Legumin.).

PHYSALOIDES, Moench, Meth. 473 (1794) = **Withania**, Pauq. (Solan.).
prostrata, Moench, Meth. Suppl. 178 = Physalis prostrata.
somnifera, Moench, Meth. 473 = W. somnifera.

PHYSANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 167. t. 29 (1861) = **Courbonia**, Brongn. (Cappar.).
glaucum, Klotzsch, l. c. = C. decumbens.

PHYSANTHERA, Bert. ex Steud. Nom. ed. II. ii. 330, 463 (1841) = **Rodriguezia**, Ruiz & Pav. (Orchid.).
callistachys, Bert. ex Steud. l. c. = R. stricta.

PHYSANTHYLLIS, Boiss. Voy. Bot. Espagne, ii. 162. t. 162 (1840) = **Anthyllis**, Riv. (Legumin.).
tetraphylla, Boiss. l. c. = A. tetraphylla.

PHYSARIA, A. Gray, Gen. Illustr. i. 162 (1848). *CRUCIFERAE*, [Benth. & Hook. f. i. 966].
didymocarpa, *A. Gray, l. c.*—Am. bor. occ.
Geyeri, *A. Gray, l. c.*—Am. bor. occ.
Newberryi, *A. Gray, in Ives, Rep. Colorado River, Bot.* 6.—Am. bor. occ.
oregona, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 363.—Am. bor. occ.

PHYSARUS, Steud. Nom. ed. II. ii. 330 (1841), sphalm. = **Physurus**, Rich. (Orchid.).

PHYSEDRA, Hook. f. in Benth. & Hook. f. Gen. i. 827 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
Barteri, *Cogn. in DC. Monog. Phan.* iii. 525.—Afr. trop.
heterophylla, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 553. —Afr. trop.
longipes, *Hook. f. in Oliver, l. c.*—Afr. trop.

PHYSENA, Noronha, ex Thou. Gen. Nov. Madag. 6 (1806). *PASSIFLOREAE*, Benth. & Hook. f. i. 815.
madagascariensis, *Steud. Nom.* ed. II. ii. 330, nomen; *Thou. ex Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 54.—Madag.
sessiliflora, *Tul. l. c.* 55.—Madag.

PHYSETOBASIS, Hassk. in Versl. en Med. Kon. Akad. Wetensch. v. (1857) 99; et in Flora, xl. (1857) 104 = **Holarrhena**, R. Br. (Apocyn.).
macrocarpa, Hassk. ll. cc. = H. macrocarpa.

PHYSIANTHUS, Mart. Nov. Gen. et Sp. i. 53. t. 32 (1824) = **Araujia**, Brot. (Asclep.).
albens, Mart. l. c. 54 = A. sericifera.
angustifolius, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 292 = A. angustifolia.
auricomus, R. Grah. in Bot. Mag. t. 3891 (= *Schubertia graveolens*).—Bras.
megapotamicus, Spreng. Syst. iv. Cur. Post. 112 = A. angustifolia.

PHYSICARPOS, DC. Prod. ii. 115 (1825) = PHUSICARPOS, Poir. = **Hovea**, R. Br. (Legumin.).

PHYSICHILUS, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 310 = **Hygrophila**, R. Br. (Acanth.).
barbatus, Nees, in DC. Prod. xi. 82 = H. barbata.
senegalensis, Nees, l. c. 81 = H. senegalensis.
Serpyllum, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 311 = H. Serpyllum.

PHYSIDIUM, Schrad. in Goett. Gel. Anz. i. (1821) 714 = **Angelonia**, Humb. & Bonpl. (Scrophul.).
procumbens, Schrad. l. c. = A. procumbens.

PHYSIGLOCHIS, Neck. Elem. iii. 245 (1790) = **Carex**, Dill. (Cyperac.).

PHYSINGA, Lindl. in Bot. Reg. (1838) Misc. 32 = **Epidendrum**, Linn. (Orchid.).
prostrata, Lindl. l. c.—Guian.

PHYSIPHORA, Soland. ex DC. Prod. i. 314 (1824) = **Alsodeia**, Thou. (Violar.).
laevigata, Soland. ex DC. l. c. = A. Physiphora.

PHYSKIUM, Lour. Fl. Cochinch. 662 (1790) = **Vallisneria**, Linn. (Hydroch.).
natans, Lour. l. c. 663 = V. Physkium.

PHYSOCALYCIUM, Vest, in Flora, iii. (1820) 409 = **Bryophyllum**, Salisb. (Crassul.).

PHYSOCALYMMA, Pohl, in Flora, x. (1827) 152; et Pl. Bras. Ic. i. 99. t. 82, 83 (1827). *LYTHRARIEAE*, Benth. & Hook. f. i. 781.
floridum, Pohl, Pl. Bras. Ic. i. 100. t. 82, 83 = scaberrimum.
scaberrimum, *Pohl, in Flora*, x. (1827) 153.—Bras.

PHYSOCALYMNA, DC. Prod. iii. 89 (1828) = praec.

PHYSOCALYX, Pohl, in Flora, x. (1827) 147; et Pl. Bras. Ic. i, 63. t. 53 (1827). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 966.
aurantiacus, *Pohl, in Flora*, x. (1827) 147; *et Pl. Bras. Ic.* i. 65.—Bras.
major, *Mart. Nov. Gen. et Sp.* iii. 2. t. 201.—Bras.
minor, Mart. l. c. 202 = aurantiacus.
rhinanthoides, Cham. in Linnaea, viii. (1833) 23 = Melasma rhinanthoides.

PHYSOCARPA, Rafin. New Fl. Am. iii. 73 (1836) = **Neillia**, D. Don (Rosac.).
amurensis, Maxim. in Act. Hort. Petrop. vi. (1879) 221.—As. or.
glabra, Rafin. New Fl. Am. iii. 73 = N. opulifolia.
opulifolia, Rafin. l. c. = N. opulifolia.
riparia, Rafin. l. c. = N. opulifolia.
tomentosa, Rafin. l. c. 74 = N. opulifolia.
Torreyi, Maxim. in Act. Hort. Petrop. vi. (1879) 221 = N. Torreyi.

PHYSOCARPON, Neck, Elem. ii. 164 (1790) = **Lychnis**, Tourn. (Caryoph.).

PHYSOCARPUM, Bercht. & Presl, Rostl. i. Ranunc. 14 (1823) = **Thalictrum**, Tourn. (Ranunc.).
clavatum, Bercht. & Presl, l. c. = T. clavatum.
podocarpum, Bercht. & Presl, l. c. 15 = T. podocarpum.

PHYSOCARPUS, Maxim. in Act. Hort. Petrop. vi. (1879) 219 = PHYSOCARPA, Rafin. = **Neillia**, D. Don (Rosac.).

PHYSOCAULIS, Tausch, in Flora, xvii. (1834) 342 = **Chaerophyllum**, Linn. (Umbell.).
nodosus, Koch, Syn. Fl. Germ. ed. II. 348 = C. nodosum.

PHYSOCHEILUS, Nutt. ex Benth. in DC. Prod. x. 536, in syn. (1846) = **Orthocarpus**, Nutt. (Labiat.).
variegatus, Nutt. ex Benth. l. c. = O. purpurascens.

PHYSOCHLAENA, C. Koch, in Linnaea, xxii. (1849) 737 = seq.

PHYSOCHLAINA, G. Don, Gen. Syst. iv. 470 (1837). *SOLANACEAE*, Benth. & Hook. f. ii. 902.
BELENIA, Decne. in Jacquem. Voy. Bot. 113. t. 120 (1844 ?).
dahurica, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. v. (1850) 471.—Dahuria.
grandiflora, Hook. Bot. Mag. t. 4600 = praealta.
orientalis, *G. Don, Gen. Syst.* iv. 470.—Oriens.
physaloides, G. Don, l. c. = Scopola physaloides.
praealta, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. v. (1850) 473.—Reg. Himal.
rubricaulis, *Miers, l. c.* 471.—Hab. ?
Semenowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) i. 95.—As. centr.

PHYSOCLAINA, Boiss. Fl. Orient. iv. 293 (1879) = praec.

PHYSOCODON, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 212 = **Melochia**, Linn. (Stercul.).
brachybotrys, Turcz. l. c. 213.—Bras.
macrobotrys, Turcz. l. c.—Ecuador.

PHYSODEIRA, Hanst. in Linnaea, xxvi. (1853) 207 = **Episcia**, Mart. (Gesner.).
bicolor, Hanst. l. c. = E. bicolor.

PHYSODIA, Salisb. Gen. Pl. Fragm. 37 (1866) = **Urginea**, Steinh. (Liliac.).
physodes, Salisb. l. c. = U. physodes.

PHYSODIUM, Presl, Rel. Haenk. ii. 150. t. 72 (1836). *STERCULIACEAE*, Benth. & Hook. f. i. 223.
corymbosum, *Presl, l. c.*—Mexic.
dubium, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.

PHYSOGETON, Jaub. & Spach, Illustr. ii. 48. t. 135 (1846) = **Halanthium**, C. Koch (Chenop.).
acanthophyllus, Jaub. & Spach, Illustr. ii. 48. t. 135 = H. purpureum.
purpureus, Moq. in DC. Prod. xiii. ii. 202 = H. purpureum.

PHYSOLEPIDION, Schrenk, Enum. Pl. Nov. 97 (1841) = **Lepidium**, Linn. (Crucif.).
repens, Schrenk, l. c.—Soongar.

PHYSOLEPIDIUM, Endl. Suppl. ii. 108 (1842) = praec.

PHYSOLEUCAS, Jaub. & Spach, Illustr. v. 48. t. 445 (1855) = **Leucas**, R. Br. (Labiat.).
acrodonta, Jaub. & Spach, l. c. = L. acrodonta.
arabica, Jaub. & Spach, l. c. 49 = L. inflata.
pachystachya, Jaub. & Spach, l. c. t. 445 = L. inflata.
Schimperi, Jaub. & Spach, l. c. = L. acrodonta.

PHYSOLOBIUM, Benth. in Hueg. Bot. Archiv, t. 2 (1837) = **Kennedya**, Vent. (Legumin.).
carinatum, Benth. in Enum, Pl. Hueg. 39 = K. parviflora.
elatum, Benth. in Hueg. Bot. Archiv, t. 2 = K. Stirlingii.
gracile, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 608 = K. parviflora.
macrophyllum, Meissn. in Lehm. Pl. Preiss. ii. 222 = K. macrophylla.
Stirlingii, Benth. in Hueg. Bot. Archiv, sub t. 2 = K. Stirlingii.

PHYSOLOPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) ii. 728 = **Angelica**, Riv. (Umbellif.).
saxatile, Turcz. l. c. 729 = A. saxatilis.

PHYSOLYCHNIS, Rupr. in Mém. Acad. Pétersb. Sér. VII. xiv. (1869) n. 4 (Sert. Tianschan.) 41 = **Lychnis**, Tourn. (Caryoph.).
altaica, Rupr. l. c.—Sibir. altaic.
apetala, Rupr. l. c. = L. apetala.
gonosperma, Rupr. l. c.—Turkestan.
uralensis, Rupr. l. c.—Ross.

PHYSONDRA, Rafin. Atl. Journ. 145 (1832) = **Astragalus**, Tourn. (Legumin.).
dispar, Rafin. l. c. = A. multiflorus.
longifolia, Rafin. l. c. = A. pictus.

PHYSOPHORA, Link, Enum. Hort. Berol. i. 278 (1821) = **Physospermum**, Cusson (Umbell.).
sylvatica, Link, l. c. = Physospermum commutatum.

PHYSOPLEXIS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = **Phyteuma**, Linn. (Campan.).
comosa, Schur, l. c. = Phyt. comosum.

PHYSOPODIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 403. Genus dubium *LYTHRARIARUM*, Benth. & Hook. f. i. 776.
volubile, *Desv. l. c.*—Ins. Borbon.

PHYSOPSIS, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 34. *VERBENACEAE*, Benth. & Hook. f. ii. 1139.
spicata, *Turcz. l. c.* 35.—Austral.

PHYSOPTYCHIS, Boiss. Fl. Orient. i. 260 (1867) = **Vesicaria**, Tourn. (Crucif.).
gnaphalodes, Boiss. l. c. = V. gnaphalodes.

PHYSORHYNCHUS, Hook. Ic. Pl. tt. 821, 822 (1852). *CRUCIFERAE*, Benth. & Hook. f. i. 100.
brahuicus, *Hook. l. c.*—Ind. bor. occ.
Chamaerapistrum, Boiss. Fl. Orient. i. 403 = brahuicus.

PHYSORHYNCUS, Hook. f. & T. Anders. in Hook. f. Fl. Brit. Ind. i. 165 (1872), sphalm. = praec.

PHYSOSIPHON, Lindl. Bot. Reg. sub t. 1797 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 490.
carinatus, *Lindl. Bot. Reg.* (1838) *Misc.* 72.—Mexic.
emarginatus, *Lindl. l. c.* sub t. 1797.—Am. austr.
Loddigesii, *Lindl. l. c.*—Mexic.
nicaraguensis, Liebm. Ind. Sem. Hort. Haun. (1853) 20 = Pleurothallis nicaraguensis.
ochraceus, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 17.—Mexic.
pubescens, *Rodrig. Orch. Nov.* i. 27.—Bras.
punctulatus, *Reichb. f. in Bot. Zeit.* xxiv. (1866) 385.—N. Granat.
serrulatus, *Rodrig. Orch. Nov.* ii. 78.—Bras.
spiralis, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
tubatus, Reichb. f. in Walp. Ann. vi. 188 = Loddigesii.

PHYSOSPERMUM, Cusson, in Mém. Soc. Méd. Par. (1782) 279. *UMBELLIFERAE*, Benth. & Hook. f. i. 882.
ALSCHINGERA, Vis. Fl. Dalm. iii. 69 (1849).
DANAA, All. Fl. Pedem. ii. 34. t. 63 (1785).
HAENSELERA, Lag. Gen. et Sp. Nov. 13 (1816).
PHYSOPHORA, Link, Enum. Hort. Berol. i. 278 (1821).
PSEUDOSPERMUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 517 (1821).
acteaefolium, Eichw. Casp. 32 = Eleutherospermum grandifolium.
acteaefolium, *Presl, Delic. Prag.* 128.—Sicil.; Ital.
aegopodioides, *Boiss. Fl. Orient.* ii. 923.—Macedon.
angelicaefolium, Guss. Ind. Sem. Hort. Bocc. (1825); ex ej. Fl. Sic. Prod. i. 356 = acteaefolium.
aquilegifolium, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 134 = commutatum.
cicutarium, Spreng. Sp. Umbellif. 23 = Eleutherospermum grandifolium.
commutatum, *Spreng. Umbellif. Prod.* 19.—Europ.; As. Min.
cornubiense, DC. Prod. iv. 246 = commutatum.
nudicaule, C. A. Mey. Verz. Pfl. Cauc. 132 = commutatum.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 40.—Kokan.
verticillatum, Vis. Fl. Dalm. iii. 358, in obs. = acteaefolium.

PHYSOSPERMUM, Lag. Amen. Nat. ii. 97 (1821) = **Pleurospermum**, Hoffm. (Umbellif.).
Gussoni, Lag. l. c. = Pleurosp. austriacum.

PHYSOSTEGIA, Benth. in Bot. Reg. sub t. 1289 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1204.
denticulata, Benth. in Bot. Reg. sub t. 1289 = virginiana.
imbricata, Benth. in DC. Prod. xii. 434 = parviflora.
imbricata, Hook. Bot. Mag. t. 3386 = virginiana.
intermedia, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 257.—Am. bor. occ.
parviflora, *Nutt. ex Benth. in DC. Prod.* xii. 434, in syn.; *et ex A. Gray, in Proc. Am. Acad.* viii. (1873) 371.—Am. bor.
speciosa, Sweet, Hort. Brit. ed. II. 406 = virginiana.
truncata, Benth. Lab. Gen. et Sp. 505 = Brazoria truncata.
truncata, Hook. Bot. Mag. t. 3494 = Brazoria scutellarioides.
variegata, Benth. in Bot. Reg. sub t. 1289 = virginiana.
virginiana, *Benth. l. c.*; *Lab. Gen. et Sp.* 504.—Am. bor.

PHYSOSTELMA, Wight, Contrib. 39 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 777.
CYSTIDIANTHUS, Hassk. Cat. Hort. Bog. Alt. 126 (1844).
campanulatum, Decne. in DC. Prod. viii. 633 = Wallichii.
laurifolium, *Walp. Ann.* iii. 63.—Sumatra.
Wallichii, *Wight, Contrib.* 40.—Singapore.

PHYSOSTEMON, Mart. Nov. Gen. et Sp. i. 72. t. 45 (1824) = **Cleome**, Linn. (Cappar.).
ambiguum, Bong. in Bull. Sc. Acad. Pétersb. i. (1836) 115 = C. stenophylla (Cf. Addend.).
intermedium, Moric. Pl. Nouv. Am. 62. t. 42 = C. guianensis.
lanceolatum, Mart. Nov. Gen. et Sp. i. 78. t. 45 = C. stenophylla (Cf. Addend.).
rotundifolium, Mart. l. c. 74. t. 47.—Bras.
tenuifolium, Mart. l. c. 73. t. 46.—Bras.

PHYSOSTIGMA, Balf. in Trans. Roy. Soc. Edinb. xxii. (1861) 310. t. 16. *LEGUMINOSAE*, Benth. & Hook. f. i. 538.
cylindrospermum, *Holmes, in Pharm. Journ.* Ser. III. ix. (1879) 913.—*Cf. Trimen, in Journ. Bot.* xvii. (1879) 185, 186.—Afr. trop.
venenosum, *Balf. in Trans. Roy. Soc. Edinb.* xxii. (1861) 310.—Afr. trop.

PHYSOTRICHIA, Hiern, in Journ. Bot. xi. (1873) 161. *UMBELLIFERAE*.
Buchananii, *Benth. ex Oliver, in Hook. Ic. Pl.* t. 1358.—Afr. trop.
Welwitschii, *Hiern, in Journ. Bot.* xi. (1873) 161.—Afr. trop.

PHYSURUS, Rich. in Mém. Mus. Par. iv. (1818) 55. *ORCHIDEAE*, Benth. & Hook. f. iii. 597.
ERYTHRODES, Blume, Bijdr. 410. t. 72 (1825).
MICROCHILUS, Presl, Rel. Haenk. i. 94 (1827).
ORCHIPEDUM, Breda, Kuhl. et Hasselt, Gen. et Sp. Orch. [t. 10] (1827).
QUETELETIA, Blume, Orch. Archip. Ind. 117. t. 37 (1858).
aratanhensis, *Rodrig. Orch. Nov.* ii. 290.—Bras.
argenteus, *Hort. ex C. Koch & Lauche, in Allg. Gartenz.* (1857) 5.—Hab.?
arietinus, *Reichb. f. & Warm. in Otia Bot. Hamb.* 82.—Bras.
bicolor, *Rodrig. Orch. Nov.* ii. 290.—Bras.
bifalcis, *Lindl. Gen. et Sp. Orch.* 502.—Peruv.
Blumei, *Lindl. in Wall. Cat.* n. 7397.—Ind. or.; Malaya.
brachyceras, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 33 = Goodyera striata.
brachyrrhynchus, *Reichb. f. Xenia Orch.* ii. 184.—Venezuela.
bracteatus, *Blume, Orch. Archip. Ind.* 97.—Sumatra.
calophyllus, *Reichb. f. Beitr. Orch. Centr. Am.* 64.—Costa Rica.
chiriquensis, *Reichb. f. ex Linden, Illustr. Hortic.* xxix. (1882) 116, nomen.—N. Granat.
claviger, *Reichb. f. in Bonplandia*, iv. (1856) 211.—Mexic.
commelinoides, *Rodrig. Orch. Nov.* i. 193.—Bras.
commelynifolius, *Reichb. f. in Flora*, xlviii. (1865) 274.—Cuba.
corniculatus, *Reichb. f. l. c.*—Cuba.
debilis, *Lindl. Gen. et Sp. Orch.* 503.—Bras.
decorus, *Reichb. f. in Gard. Chron.* (1873) 177.—Sumatra.
densiflorus, *Lindl. Gen. et Sp. Orch.* 502.—Bras.
elatior, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Ind. occ.
foliosus, *Lindl. Gen. et Sp. Orch.* 503.—Peruv.
glandulosus, Lindl. in Journ. Linn. Soc. i. (1857) 180 = Cystorchis glandulosus.
hirsutus, *Lindl. l. c.*—Reg. Himal.
hirtellus, *Lindl. Gen. et Sp. Orch.* 501.—Ind. occ.
humilis, *Blume, Orch. Archip. Ind.* 96.—Java.
hylibates, *Reichb. f. Otia Bot. Hamb.* 83.—Bras.
hyphaematicus, *Reichb. f. Xenia Orch.* ii. 184.—Venezuela.
lacteolus, *Rodrig. Orch. Nov.* ii. 292.—Bras.
lamprophyllus, *Linden & Reichb. f. in Hamb. Gartenz.* xvi. (1860) 15.—Bras.
latifolius, *Blume, Orch. Archip. Ind.* 95.—Java.
loxoglottis, *Reichb. f. Beitr. Orch. Centr. Am.* 64.—Guatemala.

PHYSURUS:—
maculatus, *Hook. Bot. Mag.* t. 5305.—Ecuador.
Maureverti, *Miq. Fl. Ind. Bat.* iii. 735.—Java.
metallescens, *Rodrig. Orch. Nov.* ii. 291.—Bras.
minor, *Lindl. Gen. et Sp. Orch.* 503.—Peruv.
mystacinus, *Reichb. f. Xenia Orch.* ii. 183.—Peruv.
nobilis, *Reichb. f. in Gard. Chron.* (1873) 177.—Bras.
Ortgiesii, *Reichb. f. l. c.*—N. Granat.
ovatus, *Lindl. Gen. et Sp. Orch.* 502.—Peruv.
pauciflorus, *Lindl. l. c.* 504.—Bras.
pictus, *Lindl. l. c.*—Bras.
plantagineus, *Lindl. l. c.* 503.—Ind. occ.
Preslei, *Lindl. l. c.* 501.—Peruv.
querceticola, *Lindl. l. c.* 505.—Mexic.?
rariflorus, *Lindl. Orch. Linden*, 28.—Venezuela.
repens, *Lindl. Gen. et Sp. Orch.* 502.—Peruv.
roseus, *Lindl. l. c.* 501.—Bras.
Sagraeanus, *A. Rich. Fl. Cub. Fanerog.* ii. 253.—Cuba.
stigmatopterus, *Reichb. f. Xenia Orch.* ii. 185.—Bras.
tridax, *Reichb. f. Beitr. Orch. Centr. Am.* 64.—Costa Rica.
vaginatus, *Hook. Ic. Pl.* t. 449.—Guatemala.
vasiflorus, Linden, Illustr. Hortic. xxix. (1882) 116, sphalm. = rariflorus.
vesicifer, *Reichb. f. Beitr. Orch. Centr. Am.* 63.—Costa Rica.
viridiflorus, *Lindl. in Journ. Linn. Soc.* i. (1857) 180.—Java.
xystophyllus, *Reichb. f. Xenia Orch.* ii. 183.—Venezuela.

PHYTARRHIZA, Vis. in Mem. Ist. Venet. v. (1855) 340 = **Tillandsia**, Linn. (Bromel.).
anceps, E. Morr. in Belg. Hortic. xxix. (1879) 368. tt. 20, 21 = T. anceps.
crocata, E. Morr. l. c. xxx. (1880) 87.—Bras.
Duratii, Vis. in Mem. Ist. Venet. v. (1855) 340 = T. Duratii.
Hamaleana, E. Morr. in Belg. Hortic. xx. (1870) 97 = T. Hamaleana.
Lindeni, E. Morr. in Gard. Chron. (1879) ii. 460 = T. Lindeni.
monadelpha, E. Morr. in Belg. Hortic. xxxii. (1882) 168. t. 7.—Am. austr.

PHYTELEPHAS, Ruiz & Pav. Syst. Veg. 299 (1798). *PALMAE*, Benth. & Hook. f. iii. 921.
ELEPHANTUSIA, Willd. Sp. Pl. iv. 1156 (1805).
aequatorialis, *Spruce, in Journ. Linn. Soc.* xi. (1871) 179, 180.—Ecuador.
aureo-costata, *Linden, in Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
Bonplandiana, *Gaudich. Bot. Voy. Bonite*, t. 30.—Hab.?
Endlicheriana, *Gaudich. l. c.*—Hab.?
Humboldtiana, *Gaudich. l. c.*—Hab.?
Kunthiana, *Gaudich. l. c.*—Hab.?
macrocarpa, *Ruiz & Pav. Syst. Veg.* 301.—Am. austr.
microcarpa, *Ruiz & Pav. l. c.* 302.—Peruv.
Orbignyana, *Gaudich. Bot. Voy. Bonite*, t. 29.—Hab.?
Pavonii, *Gaudich. l. c.* tt. 16, 29.—Hab.?
Persooniana, *Gaudich. l. c.* t. 30.—Hab.?
Poeppigii, *Gaudich. l. c.* t. 16.—Hab.?
Ruizii, *Gaudich. l. c.* t. 14.—Hab.?
Schottii, *H. Wendl. in Kerch. Palm.* 253, nomen.—Hab.?
Willdenowiana, *Gaudich. Bot. Voy. Bonite*, t. 30.—Hab.?

PHYTEUMA, Linn. Gen. ed. I. 46 (1737). *CAMPANULACEAE*, Benth. & Hook. f. ii. 561.
ASYNEUMA, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 335.
PETROMARULA, Vent. ex Hedw. f. Gen. 139 (1806).
PHYSOPLEXIS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47.
PODANTHUM, Boiss. Fl. Orient. iii. 945 (1875).
RAPUNCULUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
TRACANTHELIUM, Schur, Enum. Pl. Trans. 431 (1866).
adulterinum ×, Wallr. Beitr. Fl. Hercyn. 188.—German.
ambiguum, Schur, Enum. Pl. Trans. 945 (Index) = canescens.

PHYTEUMA:—
americanum, Hill, Veg. Syst. viii. 15 = Campanula americana.
amplexicaule, *Willd. Sp. Pl.* i. 925.—As. Min.
angustatum, Wender. ex Steud. Nom. ed. II. ii. 330, nomen (Quid?).—Hab.?
angustifolium, *Ledeb. Fl. Ross.* ii. 874.—Rossia.
anthericoides, *Nym. Consp.* 484.—Serbia.
argutum, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 183.—Turkestan.
armeriaefolium, Tausch, ex Reichb. Fl. Germ. Excurs. 857 = humile.
asperum, *Boiss. Diagn.* Ser. I. xi. 72.—Kurdistan.
atropurpureum, Hoppe, Taschenb. (1802) 27; et in Flora, ii. (1819) 68 = spicatum.
attenuatum, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 207.—Turkestan.
Aucheri, A. DC. in DC. Prod. vii. 456 = amplexicaule.
austriacum, G. Beck, in Verh. Zool. Bot. Ges. Wien, xxxii. (1883) 179 = orbiculare.
Balbisii, *A. DC. Monog. Camp.* 200.—Europ.
begonifolium, Roxb. Hort. Beng. 85; Fl. Ind. i. 505 = Pentaphragma begoniaefolium.
betonicaefolium, Losc. Pard. Ser. Inc. 66 = orbiculare.
betonicaefolium, Vill. Hist. Pl. Dauph. ii. 518. t. 12 = Michelii.
betonicifolium, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 190 = praec.
Bovelini, *Hegetschw. Fl. Schw.* 224.—Europ.
brevifolium, Schleich. Cat. Pl. Helvet. (1821) 25 = orbiculare.
campanuloides, *Bieb. Fl. Taur. Cauc.* i. 156.—Reg. Caucas.
canescens, *Waldst. & Kit. Pl. Rar. Hung.* i. 12. t. 14.—Hungar.; Reg. Caucas.
capensis, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
capituliforme, Rochel, Pl. Banat. Rar. 6 = orbiculare.
cappadocicum, *Boiss. Diagn.* Ser. I. xi. 73.—Cappadoc.
Carestiae, Biroli, in Brugnat. Giorn. Fis. II. i. (1818) 37 = humile.
Carestiae, Loisel, Fl. Gall. ed. II. i. 145 = serratum.
Charmelii, Sieber, ex Spreng. Pugill. i. 15 = Sieberi.
Charmelii, Vill. Hist. Pl. Dauph. ii. 516 = Scheuchzeri.
charmelioides, Biroli, in Mem. Ac. Torin. xxiv. (1820) 577 = Scheuchzeri.
cichoriforme, *Boiss. Diagn.* Ser. I. iv. 37.—As. Min.
collinum, Guss. Pl. Rar. 97. t. 19 = limoniifolium.
Columnae, Thomas, Cat. Pl. Suiss. (1818) 22; et ex Gaud. Fl. Helv. ii. 178 = Scheuchzeri.
comosum, Gouan, Fl. Monsp. 27 = orbiculare.
comosum, *Linn. Sp. Pl.* 171.—Europ. austr.
confusum, Kern. in Zeitschr. Ferdinand. Tirol. III. xv. (1870) 247 = orbiculare.
cordatum, Balb. Misc. ii. 10 = Balbisii.
cordatum, Reichb. Fl. Germ. Excurs. 297 = Sieberi.
cordatum, Sims, Bot. Mag. t. 1466 = orbiculare.
cordifolium, Lapeyr. Hist. Abr. Pl. Pyr. i. 10 = Michelii.
cordifolium, Vill. Hist. Pl. Dauph. ii. 517 = orbiculare.
corniculatum, [Clairv.] Man. Herb. 63 = Scheuchzeri.
corsicum, Sieber, ex Rochel, Bot. Reise Banat. 69 = orbiculare.
crispum, Pourr. in Mém. Acad. Toul. iii. (1788) 324 = Jasione humilis.
dubia, Frivald. in Flora, xx. (1837) 234 = Trachelium rumelianum.
elegans, *Hegetschw. Fl. Schw.* 228.—Europ.
ellipticifolium, Vill. Hist. Pl. Dauph. ii. 517 = orbiculare.
ellipticum, Auct. sphalm. = praec.
ellipticum, Poir. Encyc. vi. 73 = lanceolatum.
ellipticum, Sibth. & Sm. Fl. Graec. Prod. i. 143 = Sibthorpianum.
elongatum, *Hegetschw. Fl. Schw.* 227.—Europ.
fistulosum, *Reichb. in Flora*, v. (1822) 534.—Europ. austr.
foliosum, Kit. in Linnaea, xxxii. (1863) 426 = canescens.
giganteum, *Nym. Consp.* 483.—Ins. Rhodus.
glaciale, Hoppe, ex Nym. Consp. 485 = Sieberi.
globulariaefolium, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 100; Hegetschw. Reisen Gebirgsstock, 146 = pauciflorum.
gracile, Boiss. & Heldr. Diagn. Ser. I. xi. 74 = lobelioides.
graminifolium, Sieber, in Flora, v. (1822) 648 = hemisphaericum.

PHYTEUMA:—
Grisebachii, Pant. ex Nym. Consp. 484 = P. Michelii.
haemisphericum, Hill, Veg. Syst. viii. 15 = hemisphaericum.
Halleri, *All. Fl. Pedem.* i. 116.—Europ.
Halleri, Duby, Bot. Gall. ed. II. i. 312 = spicatum.
hemisphaericum, Geners. Fl. Scep. Elench. n. 195; ex Wahlenb. Fl. Carpat. 63 = orbiculare.
hemisphaericum, *Linn. Sp. Pl.* 170.—Europ.
Hegetschweileri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 104.—Europ.
hispidum, Hegetschw. Fl. Schw. 224 = orbiculare?
hispidum, *Schleich. Cat. Pl. Helvet.* ed. IV. 25, nomen. —Europ.
humile, Roth, Enum. Pl. Phan. Germ. i. 696 = Scheuchzeri.
humile, *Schleich. ex Murith, Guide Bot. Val.* 84—Europ.
inaequatum, Kit. ex Schult. Oesterr. Fl. ed. II. i. 398 = orbiculare.
intermedium, Hegetschw. Reisen Gebirgsstock, 147 = hemisphaericum.
intermedium, Jan, ex A. DC. Monog. Camp. 186 = Sieberi.
Jacquini, Sieber, Reise Kreta, i. 437; ii. 316 = Trachelium Jacquini.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 192.—Japon.
Kotschyi, Boiss. Diagn. Ser. I. xi. 71 = amplexicaule.
lanceolatum, Desf. in Ann. Mus. Par. xi. (1808) 55 = lobelioides.
lanceolatum, Vill. Hist. Pl. Dauph. ii. 517 = orbiculare.
lanceolatum, *Willd. Sp. Pl.* i. 924.—As. Min.; Syria.
latifolium, *Dalla Torre, Atl. Alpenfl.* 160.—Europ.
leianthum, *Trautv. in Act. Hort. Petrop.* iii. II. (1875) 274.—As. Min.
limonifolium, Ledeb. Fl. Ross. ii. 873 = P. Otites.
limonifolium, *Sibth. & Sm. Fl. Graec. Prod.* i. 144.—Europ. austr.; As. Min.
limoniifolium, C. A. Mey. Verz. Pfl. Cauc. 83 = pulchellum.
linifolium, *Boiss. & Heldr. Diagn.* Ser. I. xi. 75.—As. Min.
lobelioides, *Willd. Phyt.* 6. t. 4.—As. Min.
longifolium, *Hegetschw. Fl. Schw.* 225.—Europ.
lycium, *Boiss. Diagn.* Ser. I. xi. 76.—Lycia.
Michelii, *All. Fl. Pedem.* i. 115.—Europ.
Michelii, Hegetschw. Reisen Gebirgsstock, 148; Fl. Schw. 228 = orbiculare.
Michelii, Lapeyr. Hist. Abr. Pl. Pyr. 109 = hemisphaericum.
Michelii, F. W. Schmidt, Fl. Boëm. ii. 85 = Scheuchzeri.
minutum, Roem. & Schult. Syst. v. 86 = Jasione foliosa.
multicaule, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 208.—Turkestan.
nanum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = pauciflorum.
nigrum, *F. W. Schmidt, Fl. Boëm.* ii. 87.—Europ.
orbiculare, Jacq. Fl. Ausfr. t. 437 = Scheuchzeri.
orbiculare, *Linn. Sp. Pl.* 170.—Europ.
Otites, *Trautv. in Act. Hort. Petrop.* vi. (1879) 49.—As. Min.
ovale, Hoppe, Taschenb. (1794) 82 = nigrum.
ovatum, Hoffm. Deutschl. Fl. ed. II. i. 105 = nigrum.
ovatum, Lam. Illustr. ii. 68 = Scheuchzeri.
ovatum, F. W. Schmidt, Fl. Boëm. ii. 87 = Halleri.
paniculatum, Janka, ex Nym. Consp. 483 = canescens.
parviflorum, Mutel, Fl. Fr. t. 35. f. 269, sphalm. = pauciflorum.
pauciflorum, Johann. ex Roem. & Schult. Syst. v. 78 = sibiricum.
pauciflorum, *Linn. Sp. Pl.* 170.—Europ.
persicaefolium, Hoppe, ex A. DC. Monog. Camp. 196 = Michelii.
Pestalozzae, Boiss. Diagn. Ser. I. xi. 74 = lobelioides.
pilosum, Hegetschw. Reisen Gebirgsstock, 149 = orbiculare.
pinnatum, *Linn. Sp. Pl.* 171.—Ins. Creta.
Pseudoorbiculare, Pant. in Verh. Ver. Naturw. Heilk. Presb. ii. (1871–72) 53 = orbiculare.
pulchellum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 35.—As. Min.; Persia.
Rapunculus, Pers. in Linn. Syst. ed. XV. (1797) 220, in obs. = nigrum.

PHYTEUMA:—
Regelii, *Trautv. in Act. Hort. Petrop.* vi. (1879) 53.—As. centr.
reniforme, Dulac, Fl. Hautes-Pyr. 456 = Scheuchzeri.
repandum, Sibth. & Sm. Fl. Graec. Prod. i. 143 = limonifolium.
rigidifolium, Dufour, ex Roem. & Schult. Syst. v. 87 = Jasione foliosa.
rigidum, *Willd. Sp. Pl.* i. 925.—As. Min.; Syria.
rumelicum, Griseb. Spicil. Fl. Rumel. ii. 291 = Trachelium rumelianum.
salicifolium, Kit. ex Schult. Oesterr. Fl. ed. II. i. 400, in obs.; A. DC. Monog. Camp. 205 = canescens.
salignum, Waldst. & Kit. ex Bess. Prim. Fl. Galic. i. 368 = canescens.
Scheuchzeri, *Alt. Fl. Pedem.* i. 116. t. 39.—Europ. austr.
Scheuchzeri, F. W. Schmidt, Fl. Boëm. ii. 86 = orbiculare.
scorzonerifolium, Vill. Hist. Pl. Dauph. ii. 519. t. 12 = Michelii.
serratum, *Viv. App. Fl. Cors. Prod.* 1.—Ins. Corsica.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 184.—Turkestan.
sibiricum, *Vest, ex Roem. & Schult. Syst.* v. 77.—Sibir.
Sibthorpianum, *Roem. & Schult. l. c.* 84.—Graecia.
Sieberi, Schur, Enum. Pl. Transs. 429 = humile.
Sieberi, *Spreng. Pugill.* i. 15.—Europ.
spicatum, Lap. Hist. Abr. Pl. Pyr. 110 = Halleri.
spicatum, *Linn. Sp. Pl.* 171.—Europ.
spiciforme, Rochel, Bot. Reise Banat. 69 = spicatum.
Sinai, A. DC. in DC. Prod. vii. 455 = lanceolatum.
strictum, Sims, Bot. Mag. t. 2145 = limonifolium.
stylidioides, Boiss. Diagn. Ser. I. xi. 73 = lobelioides.
stylosum, Hort. Par. ex Schrank, Pl. Rar. Hort. Monac. t. 49 = limonifolium.
sublyratum, Panč. ex Nym. Consp. 484 = Halleri.
supinum, G. Don, Gen. Syst. iii. 749 = Jasione supina.
tenuifolium, *A. DC. Monog. Camp.* 201.—Byzant.
tetramericum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 105 = seq.
tetramerium, Schur, l. c. iv. (1853) 47 = spicatum.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 438.—Reg. Himal.
transsilvanicum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = hemisphaericum.
tricolor, *Molina, Sagg. Chile*, ed. II. 133.—Chile.
urticaefolium, [Clairv.] Man. Herb. 63 = Halleri.
veronicaefolium, Schrad. ex A. DC. Monog. Camp. 196 = Michelii.
virgatum, Jensch. ex Trautv. in Act. Hort. Petrop. vi. (1879) 49 = P. Otites.
virgatum, Kotschy, ex A. DC. in DC. Prod. vii. 455 = Sibthorpianum.
virgatum, Lodd. Bot. Cab. t. 667 = limonifolium.
virgatum, Willd. Sp. Pl. i. 914 = Campanula Rapunculus.

PHYTEUMA, Lour. Fl. Cochin. 138 (1790) = **Sambucus**, Tourn. (Caprifol.).
bipinnata, Lour. l. c. = S. ebuloides.
cochinchinensis, Lour. l. c. 139.—Cochinchin.

PHYTEUMOIDES, Smeathm. ex DC. Prod. iv. 414 (1830) = **Virecta**, Sm. (Rubiac.).
hirsutus, Smeathm. ex DC. l. c. = V. multiflora.

PHYTEUMOPSIS, Juss. ex Poir. Encyc. Suppl. iv. 405 (1816) = **Marshallia**, Schreb. (Compos.).
angustifolia, Poir. l. c. 406 = M. angustifolia.
lanceolata, Poir. l. c. 405 = M. lanceolata.
latifolia, Poir. l. c. 406 = M. latifolia.

PHYTHOLACCA, Brot. Fl. Lusit. ii. 224 (1804) sphalm. = **Phytolacca**, Tourn.

PHYTOCRENE, Wall. in Phil. Mag. iii. (1823) 223; et Pl. As. Rar. iii. 11. t. 216 (1832). *OLACINEAE*, Benth. & Hook. f. i. 354.
GYNOCEPHALUM, Blume, Bijdr. 483 (1825).
borneensis, *Becc. Malesia*, i. (1877) 128.—Borneo.
bracteata, *Wall. Pl. As. Rar.* iii. 12.—Malaya.
calicarpa, Griff. Notul. iv. 327 = gigantea.
crinipes, Baill. ex Bur. Mon. Bignon. 146, in obs. = hirsuta.
dasycarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 248.—Ins. Celebes.
gigantea, Blume, l. c. i. 41 = macrophylla.

PHYTOCRENE:—
gigantea, *Wall. Pl. As. Rar.* iii. 11. t. 216.—Burma.
hirsuta, *Blume, Rumphia*, iv. 37.—Ins. Celebes.
hirsuta, Teysm. & Binn. Cat. Hort. Bog. 207 = dasycarpa.
luzoniensis, *Baill. Adansonia*, x. (1872) 281.— ns. Philipp.
macrocarpa, Griff. Notul. iv. 322 = bracteata.
macrophylla, *Blume, Rumphia*, iv. 36.—Java.
oblonga, *Wall. Pl. As. Rar.* iii. 12.—Malaya.
palmata, *Wall. l. c.*—Malaya.
stylocarpa, Griff. Notul. iv. 320 = palmata.

PHYTOLACA, Hill, Hort. Kew. 215 (1768) sphalm. = seq.

PHYTOLACCA, Tourn. ex Linn. Syst. ed. I (1735). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 84.
PIRCUNIA, Bert. Merc. Chil. (1829); ex Am. Journ. Sc. Ser. I. xxiii. (1833) 264.
SARCOCA, Rafin. Fl. Tellur. iii. 55 (1836).
abyssinica, *Hoffm. in Comm. Gotting.* xii. (1796) 27.—Afr. trop. et austr.
abyssinica, Hook. & Arn. Bot. Beech. Voy. 94 = brachystachys.
acinosa, *Roxb. Hort. Beng.* 35; *Fl. Ind.* ii. 458.—Reg. Himal.; China.
acuminata, Hort. ex Moq. in DC. Prod. xiii. II. 33 = icosandra.
americana, Linn. Sp. Pl. 441 = decandra.
arborea, Hort. ex Moq. in DC. Prod. xiii. II. 31 = dioica.
asiatica, *Linn. Sp. Pl.* 441.—Ind. or.
australis, *Phil. in Anal. Univ. Chil.* (1873) 536.—Chili.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* ii. 183.—N. Granat.
bogotensis, H. Mann, in Proc. Am. Acad. vii. (1867) 198 = brachystachys.
bogotensis, Miq. Sert. Exot. 6. t. 3 = icosandra.
brachystachys, *Moq. in DC. Prod.* xiii. II. 31.—Ins. Sandvic.
chilensis, Bridges, ex Moq. l. c. 25 = Anisomeria littoralis.
chilensis, *Miers, Trav.* ii. 532, nomen.—Chili.
decandra, Descourt. Fl. Antill. v. t. 312 = octandra.
decandra, *Linn. Sp. Pl.* ed. II. 631.—Am. bor.
dioica, *Linn. l. c.* 632.—Am. austr.
dodecandra, L'Hérit. Stirp. Nov. 143 = abyssinica.
drastica, Poepp. & Endl. Nov. Gen. et Sp. i. 26. tt. 43, 44 = Anisomeria drastica.
elongata, Salisb. Prod. 345 = abyssinica.
esculenta, Van Houtte, in Fl. des Serres, iv. (1848) 398 B = acinosa.
gigantea, *Mart. Hort. Reg. Monac.* (1829) 57, nomen. (Quid?).—Hab.?
* *gracilis*, Herb. ex Moq. in DC. Prod. xiii. II. 34, 236 = Deeringia celosioides.
heptandra, Retz. Obs. vi. 29 = stricta.
icosandra, *Linn. Syst.* ed. X. 1040.—Ind. or.
icosandra, Sims, Bot. Mag. t. 2633 = mexicana.
Kaempferi, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 404 = acinosa.
Latbenia, Maxim. Ind. Sem. Hort. Petrop. (1869) 24 = Rivina laevis.
littoralis, *Poepp. & Endl. Nov. Gen. et Sp.* i. 27. t. 45.—Chili.
longespica, Moq. in DC. Prod. xiii. II. 33 = octandra.
lutea, Marsigl. ex Steud. Nom. ed. I. 618 = abyssinica.
macrostachya, Willd. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. II. 344 = icosandra.
malabarica, Crantz, Inst. ii. 484 = icosandra.
mexicana, Crantz, l. c. = octandra.
mexicana, Gaertn. Fruct. i. 377 = icosandra.
mexicana, *Sweet, Hort. Brit.* ed. I. 337.—Mexic.
octandra, *Linn. Sp. Pl.* ed. II. 631.—Japon.
oleracea, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 292.—Guiana.
pekinensis, *Hance, in Journ. Bot.* vii. (1869) 199.—China.
polystigma, Benth. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. II. 344 = icosandra.
polystyla, *Schomb. ex Moq. in DC. Prod.* xiii. II. 460, nomen.—Hab.?
populifolia, Salisb. Prod. 345 = dioica.
pruinosa, *Fenzl, Del. Sem. Hort. Vindob.* (1855) 6.—As. Min.
purpurascens, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1851); App. 13.—Cf. Linnaea, xxv. (1852) 297 = icosandra.

PHYTOLACCA:—
resedifolia, Hort. Berol. ex Moq. in DC. Prod. xiii. II. 30 = stricta.
resediformis, Hort. Berol. ex Moq. l. c. = stricta.
rivinoides, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 15.—Venezuela.
rugosa, A. Br. & Bouché, l. c. (1851) App. 13.—Cf. Linnaea, xxv. (1852) 297 = octandra.
scandens, Hilsenb. & Boj. ex Moq. in DC. Prod. xiii. II. 30 = abyssinica.
sessiliflora, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 15.—Mexic.
stricta, *Hoffm. in Comm. Gotting.* xii. (1796) 27.—Afr. austr.
thyrsiflora, *Fenzl, ex J. A. Schmidt, in Mart. Fl. Bras.* II. 343.—Bras.
triquetra, Moench, Meth. Suppl. 107 = icosandra.
vulgaris, Crantz, Inst. ii. 484 = decandra.

PHYTOXIS, Molina, Sagg. Chile, ed. II. 145, 290 (1810) = **Sphacele**, Benth. (Labiat.).
acidissima, Bert. ex Benth. in DC. Prod. xii. 258 = Cuminia fernandezia.
acidissima, Molina, Sagg. Chile, ed. II. 291 = S. campanulata.
sideritifolia, Molina, l. c. 145 = S. campanulata.

PHYTOXYS, Spreng. Syst. ii. 676 (1825) = praec.

PIARADENA, Rafin. Fl. Tellur. iii. 91 (1836) = **Salvia**, Tourn. (Labiat.).
fulgens, Rafin. l. c. = S. fulgens.

PIARANTHUS, R. Br. in Mem. Wern. Soc. i. (1809) 23. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 782.
OBESIA, Haw. Syn. Pl. Succ. 42 (1812).
aridus, *G. Don, Gen. Syst.* iv. 114.—Afr. austr.
decorus, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 163.—Afr. austr.
fasciculatus, Schult. Syst. vi. 10 = Stapelia fasciculata.
geminatus, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 163.—Afr. austr.
Gussoneanus, G. Don, Gen. Syst. iv. 113 = Stapelia europaea.
incarnatus, G. Don, l. c. 114 = Boucerosia incarnata.
mammillaris, G. Don, l. c. = Boucerosia mammillaris.
parviflorus, Spreng. Syst. i. 841 (Carallumatis sp.).—Afr. austr.
piliferus, Sweet, Hort. Brit. ed. II. 359 = Trichocaulon piliferum.
pullus, R. Br. in Mem. Wern. Soc. i. (1809) 23 = Boucerosia mammillaris.
punctatus, *R. Br. l. c.*—Afr. austr.
ramosus, Sweet, Hort. Brit. ed. II. 359 = Stapelia ramosa.
serrulatus, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 163.—Afr. austr.

PIARIMULA, Rafin. Fl. Tellur. ii. 102 (1836) = **Lippia**, Linn. (Verben.).
chinensis, Rafin. l. c. = L. nodiflora.

PIAROPHYLA, Rafin. Fl. Tellur. ii. 68 (1836) = **Saxifraga**, Tourn.
cordifolia, Rafin. l. c. = S. crassifolia.
elliptica, Rafin. l. c. = S. crassifolia.
sibirica, Rafin. l. c. = S. sibirica.

PIAROPUS, Rafin. Fl. Tellur. ii. 81 (1836) = **Eichhornia**, Kunth (Ponted.).
azureus, Rafin. l. c. 82 = E. azurea.
crassipes, Rafin. l. c. 81 = E. speciosa.
mesomelas, Rafin. l. c.—Hab.?
tricolor, Rafin. l. c. = E. azurea.
undulatus, Rafin. l. c. iv. 121 = E. azurea.

PICARDENIA, Steud. Nom. ed. II. ii. 332 (1841), sphalm. = PICRADENIA, Hook. = **Actinella**, Nutt. (Compos.).

PICCIA, Neck. Elem. ii. 436 (1790) = **Moronobea**, Aubl. (Guttif.).

PICCONIA, DC. Prod. viii. 288 (1844) = **Notelaea**, Vent. (Oleac.).
excelsa, DC. l. c. = N. excelsa.

PICEA, Link, in Abh. Akad. Berl. (1827) 179. *CONIFERAE*, Benth. & Hook. f. iii. 439.
ABIES, D. Don, in Loud. Arb. Brit. iv. 2329 (1838).
VEITCHIA, Lindl. in Gard. Chron. (1861) 265.
ajanensis, *Fisch. ex Trautv. & C. A. Mey. in Middend. Reise (Fl. Ochot.* 87).—Reg. Amur.; Japon.
alba, *Link, Handb.* ii. 478; *et in Linnaea*, xv. (1841) 519.—Am. bor.
Alcockiana, *Carr. Conif.* ed. II. 343.—Japon.
amabilis, Gord. Pinet. 154 = Abies subalpina.
amabilis, Hort. ex Carr. Conif. ed. II. 269 = Abies nobilis.
amabilis, Loud. Arb. Brit. iv. 2342 = Abies amabilis.
Apollinis, Rauch. ex Gord. Pinet. Suppl. 44 = Abies cephalonica.
aromatica, Carr. Conif. ed. II. 310 = Abies balsamea.
balsamea, Loud. Arb. Brit. vi. 2339 = Abies balsamea.
bifolia, A. Murr. in Gard. Chron. (1875) I. f. 97 = Abies lasiocarpa.
brachyphylla, Gord. Pinet. ed. II. 201 = Abies brachyphylla.
brachyphylla, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 518 = Abies homolepis.
bracteata, Loud. Arb. Brit. iv. 2348 = Abies bracteata.
Breweriana, *S. Wats. in Proc. Am. Acad.* xx. (1885) 378.—Calif.
Brunoniana, Spach, ex Gord. Pinet. Suppl. 9 = Tsuga Brunoniana.
californica, Carr. Conif. ed. I. 261 = Tsuga Pattoniana.
canadensis, Link, in Linnaea, xv. (1841) 524 = Tsuga canadensis.
cephalonica, Loud. Encyc. Trees, 1039. f. 1940–1946 = Abies cephalonica.
cilicica, Rauch. ex Gord. Pinet. Suppl. 50 = Abies cilicica.
cinerea, Baumann, ex Gord. Pinet. 153 = Abies pectinata.
coerulea, Link, in Linnaea, xv. (1841) 522 = alba.
concolor, Gord. Pinet. 155; A. Murr. in Gard. Chron. (1875) II. 135 = Abies concolor.
Davidiana, *Bertr. in Ann. Sc. Nat.* Sér. V. xx. (1874) 86.—Tibet.
Douglassi, Link, in Linnaea, xv. (1841) 524 = Pseudotsuga Douglasii.
Engelmanni, *Engelm. in Trans. Acad. Sc. St. Louis*, ii. (1863) 212.—Am. bor. occ.
excelsa, *Link, in Linnaea*, xv. (1841) 517.—Europ.
falcata, Carr. Conif. ed. II. 314 = Abies falcata.
firma, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 518 = Abies homolepis.
firma, Sieber, ex Gord. Pinet. 147 = Abies firma.
Fortunei, A. Murr. in Proc. Hort. Soc. (1862) 421 = Keteleeria Fortunei.
Fraseri, Emerson, Trees Shrubs Massach. ed. I. 88; ed. II. i. 104 = Abies balsamea.
Fraseri, Loud. Arb. Brit. iv. 2340 = Abies Fraseri.
glaucescens, Gord. Pinet. 149 = Abies religiosa.
Glehnii, *Mast. in Gard. Chron.* (1880) I. 300.—Ins. Sachalin.
grandis, Gord. Pinet. 155 = Abies falcata.
grandis, Loud. Arb. Brit. iv. 2341. f. 2245–2246 = Abies grandis.
grandis, Newberry, in Pacif. Rail. Rep. vi. 46. t. 6, partim = Abies concolor.
Herbertiana, Madden, ex Carr. Conif. ed. I. 222 = Abies Webbiana.
hirtella, Loud. Arb. Brit. iv. 2349 = Abies religiosa.
holophylla, Gord. Pinet. ed. II. 206 = Abies holophylla.
homolepis, Carr. Conif. ed. II. 290 = Abies homolepis.
Hudsonia, Hort. ex Gord. Pinet. 148 = Abies Fraseri.
japonica, *Regel, Ind. Sem. Hort. Petrop.* (1865) 33.—Japon.
jezoënsis, Carr. Conif. ed. I. 255 = Menziesii.
jezoënsis, Maxim. in Bull. Acad. Phys.-Math. Pétersb. xv. (1857) 235, nomen = ajanensis.
Khutrow, Carr. Conif. ed. I. 258 = P. Morinda.
kukunaria, Wender. Pflanz. Bot. Gaerten, (1832) 11 = Abies pectinata.
lasiocarpa, Hort. ex Gord. Pinet. Suppl. 53 = Abies grandis.
Lowi, Hort. ex Gord. l. c. ed. II. 218 = Abies Lowiana.
Lowiana, Gord. l. c. Suppl. = Abies grandis.
magnifica, Hort. ex Carr. Conif. ed. II. 269 = Abies magnifica.

PICEA:—
Maximowiczii, *Regel, Ind. Sem. Hort. Petrop.* (1865) 33, nomen; *Mast. in Gard. Chron.* (1880) i. 363.—Japon.
Menziesii, Carr. Conif. ed. I. 237 = sitchensis.
Menziesii, Engelm. in Trans. Acad. Sc. St. Louis, ii. (1863) 214 = pungens.
Mertensiana, Hort. ex Gord. Pinet. 18 = Tsuga Mertensiana.
metensis, Hort. ex Gord. l. c. 152 = Abies pectinata.
microphylla, Carr. Conif. ed. II. 311 = Abies microphylla.
microsperma, Carr. l. c. 339 = ajanensis.
montana, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 169 = excelsa.
Morinda, *Link, in Linnaea*, xv. (1841) 522.—Reg. Himal.
mucronata, Carr. Conif. ed. II. 312 = Pseudotsuga Douglasii.
Naphta, Knight, in Loud. Encyc. Trees, 1053 = Abies Webbiana.
nigra, *Link, Handb.* ii. 478; *et in Linnaea*, xv. (1841) 520.—Am. bor.
nobilis, Loud. Arb. Brit. iv. 2342 = Abies nobilis.
Nordmanniana, Loud. Encyc. Trees, 1042. f. 1950 = Abies Nordmanniana.
numidica, R. Smith, ex Gord. Pinet. ed. II. 220 = Abies numidica.
obovata, *Ledeb. Fl. Alt.* iv. 201.—Europ. or.; Sibir.
Omorika, *Mast. in Gard. Chron.* (1884) i. 308, 309. f. 58.—Serbia.
orientalis, *Carr. Conif.* ed. I. 244.—As. Min.
panachaica, Heldr. ex. Carr. l. c. ed. II. 283; et P. Laws. Pinet. Brit. ii. 175, 176 = Abies cephalonica.
Parryana, Hort. ex Parry, in Gard. Chron. (1883) ii. 725 = pungens.
Parsonsii, Fowler, l. c. (1872) 394 = Abies Lowiana.
Parsonsii, Hort. ex Gord. Pinet. Suppl. 52 = Abies grandis.
pectinata, Loud. Arb. Brit. iv. 2329 = Abies pectinata.
Pichta, Loud. l. c. 2338 = Abies sibirica.
Pindrow, Loud. l. c. 2346 = Abies Webbiana.
pinnosa, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 515 = Abies brachyphylla.
Pinsapo, Loud. Encyc. Trees, 1041. f. 1947 = Abies Pinsapo.
polita, *Carr. Conif.* ed. I. 256.—Japon.
pungens, *Engelm. in Gard. Chron.* (1879) i. 334; (1882) i. 145; (1883) ii. 725. f. 130.—Am. bor. occ.
pyramidalis, Hort. ex Gord. Pinet. Suppl. 49 = Abies pectinata.
religiosa, Loud. Arb. Brit. iv. 2349 = Abies religiosa.
Rinzi, Hort. ex Gord. Pinet. 152 = Abies pectinata.
rubra, A. Dietr. Fl. Berl. ii. 795; Link, Handb. ii. 478; et in Linnaea, xv. (1841) 521 = nigra.
Schrenkiana, Fisch. & Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 253 = obovata.
sibirica, Hort. ex Gord. Pinet. 156 = Abies sibirica.
sitchensis, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 87).—Am. bor. occ.
Smithiana, Boiss. Fl. Orient. v. 700 = P. Morinda.
subarctica, Schur, Sertum. (1853) 69. n. 2630 = excelsa.
tianschanica, *Rupr. Sert. Tiansch.* 72.—Turkestan.
Tschonofskiana, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 518 = Abies homolepis.
Tschugatskoyae, Hort. ex Carr. Conif. ed. II. 319 = alba.
Veitchii, Hort. ex Gord. Pinet. Suppl. 56 = Abies Veitchii.
Veitchii, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 515, 518 = Abies brachyphylla, homolepis.
vulgaris, Link, in Abh. Berl. Acad. (1827) 180 (nomen prius) = excelsa.
Webbiana, Gord. Pinet. 160 = Abies firma.
Webbiana, Loud. Arb. Brit. iv. 2344 = Abies Webbiana.
Withmanniana, Carr. Conif. ed. I. 260 = orientalis.
Wittmanniana, Fisch. ex Gord. Pinet. ed. II. 15 = orientalis.

PICKERINGIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 389 (1840). *LEGUMINOSAE*, Benth. & Hook. f. i. 466.
montana, *Nutt. ex Torr. & Gray, l. c.*—Calif.

PICKERINGIA, Nutt. in Journ. Acad. Philad. vii. (1834) 95 = **Ardisia**, Sw. (Myrsin.).
paniculata, Nutt. l. c. = A. Pickeringia.

PICNOCOMON, Wallr. ex DC. Prod. iv. 647 (1830) = **Cephalaria**, Schrad. (Dipsac.).
montanum, Pomel, Nouv. Mat. Fl. Atl. 64.—Algeria.

PICNOMON, Adans. Fam. ii. 116 (1763) = **Cnicus**, Linn. (Compos.).
Acarna, Cass. in Dict. Sc. Nat. xl. 188 = C. Acarna.
spinosum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = C. Acarna.

PICOTIA, Roem. & Schult. Syst. iv. p. x (1819) = **Omphalodes**, Moench (Borag.).
brassicaefolia, Roem. & Schult. l. c. 85 = O. amplexicaulis.
cappadocica, Roem. & Schult. l. c. = O. cornifolia.
Fortisii, Roem. & Schult. l. c. 87 = O. Fortisii.
linifolia, Roem. & Schult. l. c. 86 = O. linifolia.
lithospermifolia, Roem. & Schult. l. c. = Paracaryum rugulosum.
littoralis, Roem. & Schult. l. c. = O. littoralis.
nitida, Roem. & Schult. l. c. 84 = O. nitida.
scorpioides, Roem. & Schult. l. c. 87 = O. scorpioides.
verna, Roem. & Schult. l. c. 85 = O. verna.

PICRADENIA, Hook. Fl. Bor. Am. i. 317. t. 108 (1833) = **Actinella**, Nutt. (Compos.).
Richardsonii, Hook. l. c. = A. Richardsonii (cf. Addend.).

PICRAENA, Lindl. Fl. Med. 208 (1838). *SIMARUBEAE*, Benth. & Hook. f. i. 311.
Aeschryon, Vell. Fl. Flum. i. t. 152 (1827).
Muenteria, Walp. Rep. v. 398 (1846).
excelsa, *Lindl. Fl. Med.* 208.—Ind. occ.
Vellozii, *Engl. in Mart. Fl. Bras.* xii. ii. 227.—Bras.

PICRAENA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 263 = **Astragalus**, Tourn. (Legumin.).
amara, Stev. l. c. = A. amarus.

PICRAMNIA, Sw. Prod. Veg. Ind. Occ. 27 (1788). *SIMARUBEAE*, Benth. & Hook. f. i. 315.
Tariri, Aubl. Pl. Gui. Suppl. 37 (1775) nomen prius.
andicola, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 265.—Mexic.
Antidesma, Sieber, ex Steud. Nom. ed. II. ii. 332 = pentandra.
Antidesma, *Sw. Prod. Veg. Ind. Occ.* 27.—Am. trop.
antidesmoides, *Griseb. Fl. Brit. W. Ind.* 140.—Ins. Trinit.
apetala, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 264.—N. Granat.
bahiensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 446.—Bras.
Bonplandiana, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 266.—Mexic.
Camboita, *Engl. in Mart. Fl. Bras.* xii. ii. 239.—Bras.
caracasana, *Engl. l. c.* 236.—Venezuela.
Carpinterae, *Polak. in Linnaea*, xli. (1877) 553.—Costa Rica.
ciliata, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 20.—Bras.
connaroides, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 261.—N. Granat.
Corallodendron, Tul. l. c. 259 = sphaerocarpa.
cubensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 445.—Cuba.
Fessonia, *DC. Prod.* ii. 66.—Mexic.
filipetala, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 446 = Alvaradoa amorphoides.
Gardneri, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 578.—Bras.
Glazioviana, *Engl. in Mart. Fl. Bras.* xii. ii. 242.—Bras.
gracilis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 260.—N. Granat.
grandifolia, *Engl. in Mart. Fl. Bras.* xii. ii. 235.—Bras.
Kunthii, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 267 = sphaerocarpa?
latifolia, *Tul. l. c.* 258.—N. Granat.
Lindeniana, *Tul. l. c.* 266.—Mexic.
longissima, *Tul. l. c.* 257.—N. Granat.
macrostachys, *Engl. in Mart. Fl. Bras.* xii. ii. 238.—Guian.
Martiana, *Engl. l. c.* 237.—Bras.
micrantha, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 262 = pentandra.

PICRAMNIA:—
nitida, *Engl. in Mart. Fl. Bras.* xii. ii. 234.—Bras.
oblongifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 445.—Cuba.
parvifolia, *Engl. in Mart. Fl. Bras.* xii. ii. 242.—Bras.
pendula, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 262 = Sellowii.
pentandra, *Sw. Fl. Ind. Occ.* i. 220.—Ind. occ.
polyantha, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 577.—Mexic.
ramiflora, *Planch. l. c.* 578.—Bras.
Regnelli, *Engl. in Mart. Fl. Bras.* xii. ii. 241.—Bras.
reticulata, *Griseb. Cat. Pl. Cub.* 50.—Cuba.
Riedelii, *Regel & Rach, Ind. Sem. Hort. Petrop.* (1858) 50; *et in Gartenfl.* (1859) 2.—Bras.
Sellowii, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 578.—Bras.
sphaerocarpa, *Planch. l. c.*—N. Granat.
Spruceana, *Engl. in Mart. Fl. Bras.* xii. ii. 238.—Bras.
Tariri, *DC. Prod.* ii. 66.—Guian.
teapensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 265.—Mexic.
tetramera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 598.—Mexic.
triandra, Stokes, Bot. Mat. Med. iv. 535 = P. Antidesma.
umbrosa, Seem. Bot. Voy. Herald, 97. t. 24 = latifolia.
xalapensis, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 577.—Mexic.
Warmingiana, *Engl. in Mart. Fl. Bras.* xii. ii. 234.—Bras.

PICRANENA, Endl. Ench. 607 (1841) sphalm. = **Picraena**, Lindl. (Simarub.).

PICRASMA, Blume, Bijdr. 247 (1825). *SIMARUBEAE*, Benth. & Hook. f. i. 311.
Nima, Buch.-Ham. ex A. Juss. in Mém. Mus. Par. xii. (1825) 516.
ailanthoides, Planch. in Hook. Lond. Journ. Bot. v. (1846) 573 = quassioides.
andamanica, *Kurz, Rep. Andam. Veg. App.* 38; *et ex A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 520.—Ins. Andaman.
Denhamii, *Seem. Fl. Vit.* 33.—Ins. N. Hebrid.
excelsa, Planch. in Hook. Lond. Journ. Bot. v. (1846) 574 = Picraena excelsa.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 383 = quassioides.
javanica, *Blume, Bijdr.* 248.—Java.
napalensis, *Benn. Pl. Jav. Rar.* 201.—Reg. Himal.
quassioides, *Benn. l. c.* 198.—Reg. Himal.; China.
Vellozii, Planch. in Hook. Lond. Journ. Bot. v. (1846) 574 = Picraena Vellozii.

PICRELLA, Baill. Adansonia, x. (1871) 150. *SIMARUBEAE*.
trifoliata, *Baill. l. c.*—Mexic.

PICREUS, Juss. in Dict. Sc. Nat. xl. 194 (1826) = **Pycreus**, Beauv. (Cyperac.).

PICRIA, Benth. & Hook. f. Gen. ii. 812 (1876) = PICRIUM, Schreb. = **Coutoubea**, Aubl. (Gentian.).

PICRIA, Lour. Fl. Cochinch. 392 (1790) = **Curanga**, Juss. (Scrophul.).
Fel-terrae, Lour. l. c. 393 = C. amara?
surinamensis, Spreng. Syst. ii. 843.—Guian.

PICRICARYA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Olea**, Tourn.
oppositifolia, Dennst. l. c. = O. dioica.

PICRIDIUM, Desf. Fl. Atlant. ii. 220 (1799). *COMPOSITAE*, Benth. & Hook. f. ii. 527.
Reichardia, Roth, Bot. Abh. 35 (1787).
albidum, DC. Fl. Fr. iv. 16 = Crepis albida.
alpinum, Philippe, Fl. Pyrén. i. 576 = Lactuca alpina.
arabicum, Hochst. & Steud. ex DC. Prod. vii. 182 = tingitanum.
cernuum, Duf. ex DC. l. c. 183 (= *Zollikoferia pumila*).—Hispan.; Arab.?
crassifolium, Willk. in Bot. Zeit. v. (1847) 862 = vulgare.

PICRIDIUM:—
crystallinum, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 452.—Ins. Canar.
Derbesii, *Castagne, Suppl. Cat. Pl. Marseille*, 25.—Gall. austr.
dichotomum, *Fisch. & Mey. ex DC. Prod.* vii. 183.—Reg. Caucas.; As. Min.; Persia.
discolor, *Pomel, Nouv. Mat. Fl. Atl.* 6.—Alger.
gaditanum, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 232.—Hispan.
hispanicum, Poir. Encyc. Suppl. iv. 410 = tingitanum.
intermedium, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 451.—Reg. Mediterr.
ligulatum, Kunze, in Flora, xxix. (1846) 701 = intermedium.
ligulatum, Vent. Jard. Malm. 68. t. 68 = tingitanum.
lingulatum, Boiss. Voy. Bot. Espagne, ii. 390 = vulgare.
macrogaranum, Panč. ex Nym. Consp. 461 = vulgare.
macrophyllum, Vis. & Panč. in Mem. Ist. Venet. xv. (1870) 3 = vulgare.
maritimum, Reichb. f. Ic. Fl. Germ. xix. 26. t. 56 = vulgare.
orientale, DC. Fl. Fr. iv. 16 = tingitanum.
pinnatifidum, Lag. Gen. et Sp. Nov. 24 = tingitanum.
prenanthoides, *Rouy, in Bull. Soc. Bot. Fr.* xxxi. (1884) 274, nomen.—Hispan.
rupestre, *Pomel, Nouv. Mat. Fl. Atl.* 261.—Alger.
Saharae, *Pomel, l. c.* 262.—Alger.
Schimperi, *Presl, Bot. Bemerk.* 107.—Arab.
tingitanum, *Desf. Fl. Atlant.* ii. 220.—Reg. Mediterr.; Oriens.
vulgare, *Desf. l. c.* 221.—Europ.; Reg. Mediterr.

PICRINA, Reichb. ex Steud. Nom. ed. II. ii. 332 (1841) = seq.

PICRIS, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 511.
CHOEROSERIS, Link, Handb. i. 798 (1829).
DECKERA, Sch. Bip. in Flora, xvii. (1834) 479.
HAGIOSERIS, Boiss. Diagn. Ser. I. xi. 35 (1849).
HELMINTHIA, Juss. Gen. 170 (1789).
MEDICUSIA, Moench, Meth. 536 (1794).
MICRODERIS, DC. Prod. vii. 127 (1838).
PTILOSIA, Tausch, in Flora, xi. (1828) I. Erg. 78, in nota.
SPITZELIA, Sch. Bip. in Flora, xvi. (1833) 725.
VIGINEIXIA, Pomel, Nouv. Mat. Fl. Atl. 12 (1874).
abyssinica, *Sch. Bip. in Flora*, xxii. (1839) 20.—Abyssin.
acaulis, Bernh. Syst. Verz. Erf. 141 = Leontodon hispidus.
aculeata, *Vahl, Symb. Bot.* ii. 89.—Sicil.; Afr. bor.
albida, *Ball, in Journ. Bot.* xi. (1873) 370.—Marocc.
alpina, Lam. Fl. Fr. ii. 109 = Crepis alpina.
altissima, *Delile, Fl. Egypte*, 260. t. 141.—Egypt.
altissima, Desf. Cat. Hort. Par. ed. III. 149 = hieracioides.
altissima, Ledeb. ex Reichb. Fl. Germ. Excurs. 254 = Sprengeriana.
angustifolia, DC. Prod. vii. 130 = hieracioides.
arvalis, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 371 = hieracioides.
aspera, Gilib. Fl. Lituan. i. 227 = hieracioides.
aspera, Poir. Encyc. Suppl. v. 310 = Sprengeriana.
asperrima, Lindl. in Bot. Reg. (1838) Misc. 58 = hieracioides.
asplenioides, *Linn. Sp. Pl.* 793; *Desf. Fl. Atlant.* ii. 222.—Afr. bor.
atlantica, *Ball, in Journ. Linn. Soc.* xvi. (1878) 537.—Marocc.
attenuata, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 125 = hieracioides.
auriculata, Sch. Bip. Cich. Suppl. 24; ex Nym. Consp. 467 = hieracioides.
autumnalis, All. Fl. Pedem. i. 211 = Leontodon autumnalis.
Balansae, *Benth. & Hook. f. Gen.* ii. 513.—Oriens.
barbarorum, Lindl. Bot. Reg. (1838) Misc. 58 = hieracioides.
Blancheana, *Boiss. Fl. Orient.* iii. 741.—Syria.
campylocarpa, *Boiss. & Heldr. Diagn.* Ser. I. xi. 37.—Pamphylia.
camtschatica, Steud. Nom. ed. II. ii. 333, vide *kamtschatica*.
Chaixii, Poir. Encyc. Suppl. iv. 410 = pauciflora.
comosa, *Benth. & Hook. f. Gen.* ii. 513.—Europ.

PICRIS:—
coronopifolia, DC. Prod. vii. 131 = radicata.
corymbosa, Gren. et Godr. Fl. Fr. ii. 304 = spinulosa.
crepoides, Saut. in Flora, xiii. (1830) 409 = hieracioides.
crinita, Reut. in Compt. Rend. Soc. Hallér. (1854–56) 111 = hieracioides.
cupuligera, Walp. Ann. i. 461 = pilosa.
cyanocarpa, *Boiss. Diagn.* Ser. I. xi. 37.—Arab.
dahurica, DC. Prod. vii. 129 = hieracioides.
damascena, *Boiss. & Gaill. ex Boiss. Fl. Orient.* iii. 740.—Syria.
danubialis, All. Fl. Pedem. i. 211 = Leontodon hastilis.
daurica, Link, Enum. Hort. Berol. ii. 284 = seq.
davurica, Fisch. ex Hornem. Hort. Hafn. Suppl. 155 = hieracioides.
echioides, *Linn. Sp. Pl.* 792.—Europ.; Afr. bor.
filii, *Benth. & Hook. f. Gen.* ii. 512.—Ins. Azor.
flagellaris, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 120.—Gallia.
flexuosa, *Thunb. in Trans. Linn. Soc.* ii. (1794) 340.—Japon.
foetida, Lam. Fl. Fr. ii. 108 = Crepis foetida.
glaucescens, DC. Prod. vii. 130 = strigosa.
globulifera, Desf. Tabl. ed. I. 89, nomen; ed. II. 104 = Crepis Dioscoridis.
glomerata, *C. Koch, in Linnaea*, xxiii. (1850) 666.—Armenia.
goniocaula, *Boiss. Fl. Orient.* iii. 736.—Persia.
gracilis, Jord. Pugill. Pl. Nov. 113 = hieracioides.
grandiflora, Tenore, Ind. Sem. Hort. Neap. (1830) 13; Syll. Fl. Neap. 397 = hieracioides.
hamulosa, Wall. Cat. n. 3255 = hieracioides.
hieracioides, Boiss. Voy. Bot. Espagne, ii. 384 = longifolia, *Boiss. & Reut.*
hieracioides, *Linn. Sp. Pl.* 792.—Europ.; As. temp.; Austral.; N. Zel.
hieracioides, Sibth. & Sm. Fl. Graec. Prod. ii. 132 = spinulosa.
Hieracium, With. Bot. Arr. Brit. Pl. ed. I. ii. 482 = hieracioides.
hirta, All. Fl. Pedem. i. 210 = Leontodon hirtus.
hispida, All. l. c. = Leontodon hispidus.
hispidissima, Koch, Syn. Fl. Germ. ed. I. 421 = laciniata.
hispidissima, Lecoq & Lamotte, ex Boiss. Fl. Orient. iii. 735 = spinulosa.
humifusa, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 137 = echioides.
humilis, *DC. Prod.* vii. 130.—Afr. trop.
integrifolia, Desf. Tabl. ed. I. 89, nomen = Sprengeriana.
japonica, Thunb. Fl. Jap. 299 = hieracioides.
kamtschatica, Ledeb. in Mém. Acad. Pétersb. v. (1812) 557 = hieracioides.
Kotschyi, *Boiss. Fl. Orient.* iii. 738.—Syria.
lacera, Baill. Hist. Pl. viii. 294 = Crepis lacera.
laciniata, *Vis. in Flora*, xii. (1829) I. *Erg.* 24.—Dalmat.
laevis, *C. A. Mey. Verz. Pfl. Cauc.* 59.—Reg. Caucas.
lappacea, Reichb. in Flora, vii. (1824) 104 = hieracioides.
laxa, DC. Prod. vii. 129 = Sprengeriana.
longifolia, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 520, nomen.—Rossia.
longifolia, *Boiss. & Reut. Pugill. Pl. Nov.* 69.—Hispan.
longirostris, *Sch. Bip. in Mus. Senckenb.* iii. (1839) 60.—Abyss.
lyrata, Delile, Fl. Egypte, 259. t. 40. f. 3 = radicata.
macloviana, Urv. ex DC. Prod. vii. 59 = Leuceria gossypina.
macrantha, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 120.—Gallia.
macrorhiza, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. & Nat. Toul.* vi. (1883–84) 178.—Gallia.
monticola, Lamotte, Prod. Fl. Centr. Fr. ii. 453 = hieracioides.
nepalensis, Baxt. in Loud. Hort. Brit. Suppl. ii. 656 = hieracioides.
nigricans, *Grogn. in Mém. Hist. Nat. Soc. Eduenne*, i. (1865) 169.—Gallia.
nilotica, Sieber, ex Steud. Nom. ed. I. 619; et Reichb. in Flora, vii. (1824) 106 = sulphurea.
oligocephala, Schur, Enum. Pl. Transs. 361 = hieracioides.
olympica, *Boiss. Diagn.* Ser. I. iv. 26.—Bithynia.
orophila, Timb. in Bull. Soc. Sc. Phys. & Nat. Toul. i. (1872–73) 93 = hieracioides.

PICRIS:—
paleacea, Vest, in Syll. Ratisb. i. (1824) 78 = hieracioides.
pauciflora, Guss. ex Tenore, Ind. Sem. Hort. Neap. (1830) 13 = hieracioides.
pauciflora, *Willd. Sp. Pl.* iii. 1557.—Europ. austr.; Oriens.
pilosa, *Delile, Fl. Egypte*, 260. t. 41.—Afr. bor.
pinnatifida, Jord. Pugill. Pl. Nov. 112 = stricta.
prenanthoides, Schur, Enum. Pl. Transs. 360 = hieracioides.
pygmaea, *Hill, Veg. Syst.* iii. 167 (Quid?).—Europ.
pyrenaica, Gaertn. Fruct. ii. 366. t. 159. f. 2 = pauciflora.
pyrenaica, Linn. Sp. Pl. 792 = Crepis blattarioides.
pyrenaica, Vill. Hist. Pl. Dauph. iii. 148 = hieracioides.
radicata, *Less. Syn. Comp.* 134.—Afr. bor.; Arab.
repens, Lour. Fl. Cochinch. 478 (= *Barkhausia repens.*).—China.
rhagadioloides, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 218 = Sprengeriana.
Rhagadiolus, Pers. Syn. ii. 370 = Sprengeriana.
rigens, *Benth. & Hook. f. Gen.* ii. 512.—Ins. Azor.
rigida, Ledeb. in Spreng. Syst. iii. 663 = hieracioides.
rubra, Lam. Fl. Fr. ii. 109 = Crepis rubra.
ruderalis, Salisb. Prod. 179 = echioides.
ruderalis, F. W. Schmidt, ex Willd. Sp. Pl. iii. 1558 = hieracioides.
sancta, *F. W. Schmidt, Samml. Phys. Aufs.* i. (1795) 273.—Europ.
saxatilis, All. Fl. Pedem. i. 211 = Leontodon pyrenaicus.
scaberrima, Guss. ex Tenore, Syll. Fl. Neap. 396 = laciniata.
scabra, *Forsk. Fl. Aegypt. Arab.* 143.—Arabia.
sonchoides, Vest, in Flora, iii. (1820) 5 = hieracioides.
spinosa, Poir. Encyc. Suppl. iv. 408 = echioides.
spinosissima, *Geners. in Schult. Cat. Hort. Bot. Crac.* 58.—Hab. ?
spinulosa, *Bertol. ex Guss. Fl. Sic. Syn.* ii. 400.—Europ. austr.; As. Min.
Sprengeriana, Chaix, ex Lapeyr. Hist. Abr. Pl. Pyr. 468 = pauciflora.
Sprengeriana, *Poir. Encyc.* v. 310.—Europ. austr.; As. Min.; Aegypt.
squarrosa, Steetz, in Lehm. Pl. Preiss. i. 488 = hieracioides.
striata, Spreng. Syst. iii. 664 = Hieracium capense.
stricta, Jord. Cat. Jard. Dijon (1848), 29 = spinulosa.
strigosa, *Bieb. Fl. Taur. Cauc.* ii. 250.—Oriens.
strigosa, Hort. ex Reichb. Fl. Germ. Excurs. 254 = pauciflora.
strigosa, Wall. Cat. n. 3256 = hieracioides.
sulphurea, *Delile, Fl. Egypte*, 258. t. 40.—Aegypt.; Arab.
Taraxaci, All. Fl. Pedem. i. 211 = Leontodon Taraxaci.
taraxaciflora, *DC. Prod.* vii. 131.—Afr. bor.
transsilvanica, Schur, Enum. Pl. Transs. 360 = hieracioides.
tuberosa, All. Fl. Pedem. i. 210 = Leontodon tuberosus.
tuberosa, Lapeyr. ex Monn. Ess. Hierac. 49 = Hieracium pulmonarioides.
umbellata, Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ii. 273 = hieracioides.
undulata, Dulac, Fl. Hautes-Pyr. 496 = hieracioides.
Villarsii, Jord. Cat. Jard. Dijon (1848), 29 = hieracioides.
virens, Desf. Cat. Hort. Par. ed. III. 399 = hieracioides.
Willkommii, *Nym. Syll.* 53.—Hispan.

PICRIUM, Schreb. Gen. 791 (1791) = **Coutoubea**, Aubl. (Gentian.).
spicatum, Schreb. ex Progel, in Mart. Fl. Bras. vi. I. 210 = C. spicata.

PICROCOCCUS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 262 = **Vaccinium**, Linn.
elevatus, floridanus et *stamineus*, Nutt. l. c. = V. stamineum.

PICRODENDRON, Planch. in Hook. Lond. Journ. Bot. v. (1846) 579. *SIMARUBEAE*, Benth. & Hook. f. i. 315.
arboreum, *Planch, l. c.* 580.—Jamaic.
Calunga, Mart. ex Engl. in Mart. Fl. Bras. xii. II. 214 = Simaba ferruginea.
Juglans, Griseb. Fl. Brit. W. Ind. 177 = arboreum.

PICROLEMMA, Hook. f. in Benth. & Hook. f. Gen. i. 312 (1862). *SIMARUBEAE*, Benth. & Hook. f. l. c.
Sprucei, *Hook. f. l. c.*—Am. trop.

PICROPHLAEUS, Blume, Bijdr. 1019 (1826) = **Fagraea**, Thunb. (Logan.).
javanensis, Blume, l. c. 1020.—Malaya.

PICROPHYTA, F. Muell. in Linnaea, xxv. (1852) 421 = **Goodenia**, Sm.
albiflora, F. Muell. l. c. = G. albiflora.
calcarata, F. Muell. l. c. 423 = G. calcarata.

PICRORHIZA, Royle, ex Benth. Scroph. Ind. 47 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 962.
Kurroa, *Royle, ex Benth. l. c.*—Reg. Himal.
Kurrooa, Royle, Illustr. Bot. Himal. 291 = praec.
Lindleyana, Steud. Nom. ed. II. ii. 333 = P. Kurroa.

PICROSIA, D. Don, in Trans. Linn. Soc. xvi. (1833) 183. *COMPOSITAE*, Benth. & Hook. f. ii. 530.
PSILOPOGON, Phil. in Linnaea, xxxiii. (1864–65) 126.
australis, Orbign. Voy. Am. Mérid. vii. I. t. 10 = longifolia.
longifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 183.—Peruv.
runcinata, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 32 = longifolia.

PICROTHAMNUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 417 = **Artemisia**, Linn. (Compos.).
desertorum, Nutt. l. c. = A. spinescens.

PICTETIA, DC. in Bibl. Univ. Genèv. xxix. (1825) 40; et in Ann. Sc. Nat. Sér. I. ix. (1825) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 514.
angustifolia, *Griseb. Cat. Pl. Cub.* 73.—Cuba.
aristata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 94; *et Mém. Lég.* vii. 303. t. 47. f. 5.—Ind. occ.
Desvauxii, *DC. ll. cc.* 94; 304. t. 47. f. 4.—Ins. S. Doming.
Jussiaei, *DC. ll. cc.* 94; 304.—Hab.?
microphylla, *Benth. in Hemsl. Diagn. Pl. Nov.* 8.—Mexic.
obcordata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 94; *et Mém. Lég.* vii. 303. t. 47. f. 1.—Ins. S. Doming.
pubescens, Hochst. in Flora, xxix. (1846) 599 = Ormocarpum bibracteatum.
squamata, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 94 = Ormocarpum sennoides.
ternata, *DC. l. c.*; *et Mém. Lég.* vii. 304. t. 47. f. 2.—Ins. S. Doming.

PIDDINGTONIA, A. DC. in DC. Prod. vii. 341 (1839) = **Pratia**, Gaudich. (Campan.).
cyanocarpa, Hassk. in Bonplandia, vii. (1859) 179.—Java.
montana, Miq. Fl. Ind. Bat. ii. 573 = Pratia montana.
nummularia, DC. Prod. vii. 341 = Pratia begonifolia.
Palliardii, Lehm. in Hamb. Gartenz. vii. (1851) 337.—Hab.?
patens, Miq. Fl. Ind. Bat. ii. 573 = Pratia montana.

PIERARDIA, Rafin. Fl. Tellur. iv. 41 (1836) = **Dendrobium**, Sw. (Orchid.).
bicolor, Rafin. l. c. = D. Pierardi.

PIERARDIA, Roxb. Hort. Beng. 28 (1814); Fl. Ind. ii. 254 (1832) = **Baccaurea**, Lour. (Euphorb.).
Barteri, Baill. Adansonia, iv. (1863–64) 137 = B. Barteri.
costulata, Muell. Arg. in Flora, xlvii. (1864) 469 = B. costulata.
courtallensis, Wight, Ic. t. 1912 = B. courtallensis.
dasystachya, Miq. Fl. Ind. Bat. i. II. 358 = B. dasystachya.
dulcis, Jack, in Trans. Linn. Soc. xiv. (1823) 120 = B. dulcis.
dulcis, Wall. Cat. n. 8075 = B. bracteata.
flaccida, Wall. l. c. n. 8074 = B. flaccida.
Griffoniana, Baill. Adansonia, iv. (1863–64) 136 = B. Griffoniana.
macrocarpa, Miq. Fl. Ind. Bat. Suppl. 441 = B. macrocarpa.
macrophylla, Muell. Arg. in Flora, xlvii. (1864) 516 = B. macrophylla.

PIERARDIA:—
macrostachya, Wight & Arn. in Wight, Ic. tt. 1912–1913 = Baccaurea sapida.
Motleyana, Muell. Arg. in Flora, xlvii. (1864) 516 = B. Motleyana.
parviflora, Muell. Arg. in Linnaea, xxxii. (1863) 82 = B. parviflora.
pubera, Miq. Fl. Ind. Bat. Suppl. 442 = B. pubera.
pyrrhodasya, Miq. l. c. = B. pyrrhodasya.
racemosa, Blume, Bijdr. 579 = B. racemosa.
rhakodiscus, Hassk. in Bull. Soc. Bot. Fr. vi. (1859) 713 = Cyclostemon rhacodiscus.
sapida, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 254 = B. sapida.
Seemanni, Muell. Arg. in Flora, xlvii. (1864) 469 = B. Seemanni.

PIERCEA, Mill. Gard. Dict. ed. VII. (1759) = **Rivina**, Plum. (Phytolacc.).
acuminata, Rafin. New Fl. Am. iv. 13 = R. humilis.
glabra, Mill. Gard. Dict. ed. VIII. n. 1 = R. humilis.
obliquata, Rafin. New Fl. Am. iv. 13 = R. humilis.
tomentosa, Mill. Gard. Dict. ed. VIII. n. 2 = R. humilis.

PIERIS, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159. *ERICACEAE*, Benth. & Hook. f. ii. 588.
AEGIALEA, Klotzsch, in Linnaea, xxv. (1851) 17; et in Walp. Ann. ii. 1113 (1852).
PORTUNA, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 268.
floribunda, *Benth. & Hook. f. Gen.* ii. 588.—Am. bor.
formosa, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 159.—Ind. or.; China.
Griffithiana, Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 460 = Leucothoë Griffithiana.
japonica, D. Don, ex G. Don, Gen. Syst. iii. 832 = ovalifolia.
lanceolata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 159 = ovalifolia.
mariana, *Benth. & Hook. f. Gen.* ii. 588.—Am. bor.
nitida, *Benth. & Hook. f. l. c.*—Am. bor.
ovalifolia, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 159.—Reg. Himal.; Japon.
phillyreaefolia, *DC. Prod.* vii. 599.—Am. bor.
sinensis, Sweet, Hort. Brit. ed. III. 439 = Vaccinium bracteatum.
villosa, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 461.—Reg. Himal.

PIEROTIA, Blume, Mus. Bot. Lugd. Bat. i. 179 (1850) = **Ixonanthes**, Jack (Lineae).
lucida, Blume, l. c. 180 = I. icosandra.
reticulata, Blume, l. c. = I. petiolaris.

PIERREA, Hance, in Journ. Bot. xv. (1877) 339. *SAMYDACEAE*.
dictyoneura, *Hance, l. c.*—Cochinch.

PIGAFETTA, Adans. Fam. ii. 223 (1763) = **Eranthemum**, Linn. (Acanth.).

PIGAFETTA, Benth. & Hook. f. Gen. iii. 933 (1883), sphalm. = seq.

PIGAFETTIA, Becc. Malesia, i. (1877) 89. *PALMAE*, Benth. & Hook. f. iii. 933.
elata, *H. Wendl. in Kerch. Palm.* 253.—Ins. Celebes.
filaris, *Becc. Malesia*, i. (1877) 90, in obs.—Ins. Ceram.
papuana, *Becc. l. c.* 89.—N. Guin.

PIGEA, DC. Prod. i. 307 (1824) = **Ionidium**, Vent. (Violar.).
Banksiana, Ging. ex DC. l. c. = I. heterophyllum.
calycina, DC. l. c. = I. calycinum.
filiformis, DC. l. c. = I. filiforme.
floribunda, Lindl. in Mitch. Three Exped. ii. 165 = I. floribundum.
glauca, Endl. in Enum. Pl. Hueg. 5 = I. calycinum.
monopetala, Ging. ex DC. Prod. i. 307 = I. filiforme.

PIKRIA, G. Don, Gen. Syst. iv. 617 (1837) = PICRIA, Lour. = **Curanga**, Juss. (Scrophul.).

PILANTHUS, Poit. ex Endl. Gen. 1289 (1841) = **Centrosema**, DC. (Legumin.).

PILASIA, Rafin. Fl. Tellur. ii. 53 (1836) = **Urginea**, Steinh. (Liliac.).
filifolia, Rafin. l. c. = U. filifolia.

PILDERIA, Klotzsch, in Monatsb. Akad. Berl. (Maerz, 1854) 127; et in Abh. Akad. Berl. (1855) 66. t. 7 A = **Begonia**, Linn.
hirsuta, Klotzsch, in Monatsb. Akad. Berl. (Maerz, 1854) 127 = B. cordifolia.
urticaefolia, Klotzsch, l. c. = B. Pilderia.

PILEA, Lindl. Coll. Bot. t. 4 (1821). *URTICACEAE*, Benth. & Hook. f. iii. 384.
ADENIA, Torr. Fl. N. York, ii. t. 122 (1843).
ADIKE, Rafin. New Fl. Am. i. 63 (1836).
DUBREUILIA, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 489.
DUBRUEILIA, Gaudich. in Freyc. Voy. Bot. 495 (1826).
acuminata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 302.—Mexic.
alsinaefolia, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1846) 12 = serpyllifolia.
alsinifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 211.—N. Granat.
angulata, Blume, Mus. Bot. Lugd. Bat. ii. 55 = stipulosa.
anisophylla, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 193.—Reg. Himal.
anomala, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 21.—Bolivia.
approximata, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 123.—Reg. Himal.
arguta, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 218.—Ecuador.
articulata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 180.—Ins. Maurit.
astrogramma, *Miq. in Mart. Fl. Bras.* iv. I. 201.—Bras.
atroviridis, *Baker, Fl. Maurit.* 275.—Ins. Mascar.
auriculata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 299.—Costa Rica.
Balfourii, *Baker, Fl. Maurit.* 276.—Ins. Mascar.
betulaefolia, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 227 = hederacea.
boehmerioides, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 232.—Ins. Mascar.
Boiviniana, *Wedd. l. c.* 227.—Ins. Mascar.
bracteosa, *Wedd. l. c.* 245.—Reg. Himal.
callitrichoides, *Blume, Mus. Bot. Lugd. Bat.* ii. 44.—Venezuela.
callitrichoides, Kunth, Ind. Sem. Hort. Berol. (1846) 12 = serpyllifolia.
callitrichoides, Schlecht. in Linnaea, xxvii. (1854) 495 = muscosa.
capitata, *Baker, in Journ. Linn. Soc.* xxi. (1885) 446.—Madag.
capitellata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 220.—Bolivia.
carnosula, *Wedd. l. c.* 208.—N. Granat.
celebica, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 102, 105.—Ins. Celebes.
centradenioides, *Seem. Bot. Voy. Herald*, 194.—N. Granat.
cephalantha, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 209.—Ins. S. Doming.
ceratocalyx, *Wedd. in DC. Prod.* xvi. I. 148.—Peruv.
ceratomera, *Wedd. l. c.* 132.—Afr. trop.
Chamaedrys, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 229.—Ins. Porto Rico.
chilensis, Wedd. l. c. 221 = elliptica.
chrysosplenioides, *Wedd. l. c.* 231.—N. Granat.
ciliaris, Griseb. Cat. Pl. Cub. 60 = Lindeniana.
ciliaris, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 209.—Java.
ciliata, Blume, Mus. Bot. Lugd. Bat. ii. 46 = Swartzii.
citriodora, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 216.—Am. trop.
clandestina, *Wedd. in DC. Prod.* xvi. I. 120.—Jamaic.
Commersoniana, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Borbon.
consanguinea, *Wedd. in DC. Prod.* xvi. I. 112.—Ins. S. Doming.
corymbosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 50.—Ind. occ.
crassifolia, Blume, l. c. 52 = densiflora.
crassifolia, *Hance, in Journ. Bot.* xx. (1882) 294.—China.
Cruegeriana, *Wedd. in DC. Prod.* xvi. I. 122.—Venezuela.
cruzensis, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 228 = acuminata.

PILEA :—
cubensis, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 214 = serrulata.
cuneifolia, *Wedd. l. c.* 212.—Jamaic.
cuneiformis, *Wedd. l. c.* Sér. IV. i. (1854) 186.—Ins. Maurit.
dauciodora, *Wedd. l. c.* Sér. III. xviii. (1852) 223.—Peruv.
decora, Wedd. l. c. Sér. IV. i. (1854) 284 = elegans.
deltoidea, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 298.—Costa Rica.
dendrophila, *Miq. in Mart. Fl. Bras.* iv. i. 202.—Bras.
densiflora, *Kunth, Ind. Sem. Hort. Berol.* (1846) 12.—Peruv.
depressa, *Blume, Mus. Bot. Lugd. Bat.* ii. 46.—Ind. occ.
diffusa, Griseb. in Goett. Abh. vii. (1857) 193 = pubescens.
diffusa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 222.—Ind. occ.
diversifolia, *Wedd. l. c.* 212.—Peruv.
Dombeyana, *Wedd. l. c.* 221.—Peruv.
elegans, C. Gay, Fl. Chil. v. 364 = elliptica.
elegans, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 215.—Ind. occ.
elliptica, *Hook. f. Fl. Antarct.* 344.—Chili.
elongata, Blume, Mus. Bot. Lugd. Bot. ii. 55 = elegans.
elongata, Miq. in Zoll. Syst. Verz. Ind. Archip. 102 = trinervia.
falcata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 304.—Mexic.
fallax, *Wedd. in DC. Prod.* xvi. i. 120.—Am. trop.
fasciata, *Wedd. l. c.* 149.—Am. trop.
flexuosa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 218.—Am. austr.
Forsythiana, *Wedd. in DC. Prod.* xvi. i. 111.—Ind. occ.
Franquevilleana, *Wedd. l. c.* 108.—Ind. occ.
fuscata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 300 = pubescens.
Gaudichaudiana, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 226 = pubescens.
glaberrima, Blume, Mus. Bot. Lugd. Bat. ii. 54 = smilacifolia.
globosa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 208.—N. Granat.
glomerata, *Griseb. Cat. Pl. Cub.* 69.—Cuba.
gnidioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 173.—Ind. occ.
Goglado, Blume, Mus. Bot. Lugd. Bat. ii. 53 = smilacifolia.
Goglado, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 186 = scripta.
Goudotiana, *Wedd. l. c.* Sér. III. xviii. (1852) 216.—N. Granat.
grandifolia, Blume, Mus. Bot. Lugd. Bat. ii. 52 = grandis.
grandis, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 216.—Jamaic.
Griffithii, *Blume, Mus. Bot. Lugd. Bat.* ii. 55.—Malacca.
grossecrenata, Miq. in Mart. Fl. Bras. iv. i. 199 = pubescens.
guadalupensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 223.—Ins. Guadeloup.
guyanensis, Wedd. l. c. 221 = pubescens.
haitensis, Wedd. l. c. 219 = rugosa.
hederacea, *Wedd. l. c.* 222.—Ind. occ.
herniariifolia, Blume, Mus. Bot. Lugd. Bat. ii. 44 = muscosa.
herniarioides, Lindl. Coll. Bot. sub t. 4; Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 207 = muscosa.
heteronema, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 173.—Ind. occ.
Hilariana, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 210.—Bras.
hirsuta, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 220.—N. Granat.
hirtella, *Miq. in Mart. Fl. Bras.* iv. i. 202.—Bras.
Hookeriana, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 226.—Reg. Himal.
Huegelii, Blume, Mus. Bot. Lugd. Bat. ii. 53 = P. Wightii.
hyalina, *Fenzl, in Denkschr. Acad. Wien (Math.-Nat.)*, i. (1850) 250.—Bras.

PILEA :—
hygrophila, *Blume, Mus. Bot. Lugd. Bat.* ii. 56.—Ind. or.
hypnophila, *Baker, in Journ. Bot.* xx. (1882) 267.—Madag.
imparifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 212.—Guian.
inaequalis, *Wedd. l. c.* 229.—Ind. occ.
insignis, *Blume, Mus. Bot. Lugd. Bat.* ii. 56.—Ins. Borbon.
insolens, *Wedd. in DC. Prod.* xvi. i. 118.—Reg. Himal.
integrifolia, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 298.—Mexic.
Jamesoniana, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 224 = dauciodora.
javanica, Wedd. l. c. Sér. IV. i. (1854) 187 = Pellionia javanica.
Kakurang, *Blume, Mus. Bot. Lugd. Bat.* ii. 57.—Ins. Banda.
laevicaulis, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Borbon.
lamioides, *Wedd. l. c.* Sér. III. xviii. (1852) 213.—Peruv.
lanceolata, *Wedd. l. c.* 208.—Ind. occ.
latifolia, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 249.—Am. austr.
laxa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 230.—Jamaic.
leucophaea, *Blume, Mus. Bot. Lugd. Bat.* ii. 53.—Java.
Lindeniana, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 210.—N. Granat.
Lindenii, Blume, Mus. Bot. Lugd. Bat. ii. 49 = praec.
longifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 266.—Madag.
longipes, *Baker, l. c.* xxi. (1885) 446.—Madag.
longipes, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 301.—Mexic.
lucens, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Afr. trop.
lucida, *Blume, Mus. Bot. Lugd. Bat.* ii. 48.—Ind. occ.
Lundii, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 299.—Mexic.
lurida, *Wright, in Sauv. Fl. Cub.* 148.—Ind. occ.
macrodonta, *Baker, in Journ. Linn. Soc.* xx. (1883) 266.—Madag.
Manniana, *Wedd. in DC. Prod.* xvi. i. 117.—Afr. trop.
marginata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 238.—Peruv.
Maximiliani, *Miq. in Mart. Fl. Bras.* iv. i. 200.—Bras.
melastomoides, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 186 = trinervia.
mexicana, *Wedd. l. c.* Sér. III. xviii. (1852) 214.—Mexic.
miconiaefolia, Miq. in Zoll. Syst. Verz. Ind. Archip. 102, 106 = smilacifolia.
microphylla, Griseb. Cat. Pl. Cub. 59 = trianthemoides.
microphylla, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 302 = muscosa.
modesta, *Baker, in Journ. Linn. Soc.* xx. (1883) 265.—Madag.
mollis, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 251.—Am. austr.
mongolica, *Wedd. in DC. Prod.* xvi. i. 135.—Mongol.
montana, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 228 = pubescens.
Multiflora, *Wedd. l. c.* 218.—Am. trop.
muscosa, Hort. ex Wedd. in DC. Prod. xvi. i. 107 = serpyllifolia.
muscosa, *Lindl. Coll. Bot.* t. 4.—Am. trop.
Mutisiana, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 218.—N. Granat.
nana, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 299.—Mexic.
nepalensis, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 187.—Nepal.
nitida, *Wedd. l. c.* Sér. III. xviii. (1852) 211.—Peruv.
nudicaulis, *Wedd. l. c.* 208.—Ind. occ.
nummulariaefolia, *Wedd. l. c.* 225.—Ind. occ.
nummulariifolia, Griseb. in Goett. Abh. vii. (1857) 193 = repens.
nutans, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 196. t. 7. f. 11–13.—Am. trop.
obesa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Nepal.

PILEA :—
obtusata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 300.—Ins. Monserrat.
oreophila, Blume, Mus. Bot. Lugd. Bat. ii. 54 = trinervia.
ovalis, *Griseb. Fl. Brit. W. Ind.* 158.—Ind. occ.
oxyodon, Wedd. in Arch. Mus. Par. viii. (1855–56) 221 = bracteosa.
pachycarpa, *Wedd. l. c.* 225.—Reg. Himal.
Parietaria, *Blume, Mus. Bot. Lugd. Bat.* ii. 48.—Ind. occ.
Parietaria, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 173 = glomerata.
Pavonii, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 219.—Peruv.
pedunculata, Blume, Mus. Bot. Lugd. Bat. ii. 46 = elliptica.
pellucida, Blume, l. c. 54 = trinervia.
peltata, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 242.—China.
pendula, Blume, Mus. Bot. Lugd. Bat. ii. 47 = rupipendia.
pendula, *Griseb. Cat. Pl. Cub.* 59.—Ind occ.
peperomifolia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 297 = muscosa.
peplidifolia, Schlecht. in Linnaea, xxvii. (1854) 496 = serpyllifolia.
peploides, *Hook. & Arn. Bot. Beech. Voy.* 96.—As. trop.; Ins. Pacif.
petiolaris, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 187 = stipulosa.
Poeppigiana, *Wedd. l. c.* Sér. III. xviii. (1852) 225.—Peruv.
portoricensis, *Wedd. in DC. Prod.* xvi. i. 151.—Ins. Portorico.
Portula, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 297 = muscosa.
portulacina, Blume, Mus. Bot. Lugd. Bat. ii. 45 = muscosa.
producta, Blume, l. c. 56 = umbrosa.
propinqua, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 190.—Ind. occ.
pteropodon, *Wedd. in DC. Prod.* xvi. i. 144.—N. Granat.
pubescens, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 302.—Bras.
pulegifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 213.—Peruv.
pumila, *A. Gray, Man. Bot. N. U. St.* ed. I. 437.—Am. bor.
pumila, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 305.—Mexic.
punctata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 222.—Peruv.
puncticaulis, Wedd. in Arch. Mus. Par. viii. (1855–56) 181 = ciliaris.
pygmaea, Miq. in Zoll. Syst. Verz. Ind. Archip. 102, 106 = peploides.
pyrrhotricha, *Miq. Fl. Ind. Bat. Suppl.* 413.—Sumatra.
quadrifolia, *A. Rich. Tent. Fl. Abyss.* ii. 263.—Abyss.
radicans, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 223.—Jamaic.
radicans, Wight, Ic. t. 1974 = Wightii.
repanda, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Maurit.
repens, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 304.—Mexic.
reticulata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 215.—Jamaic.
rhexioides, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 304.—Mexic.
rhizobola, *Miq. in Mart. Fl. Bras.* iv. i. 202.—Bras.
rhombea, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 305.—N. Granat.
Riedlei, Blume, Mus. Bot. Lugd. Bat. ii. 55 = lucens.
Rivoriae, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 210 = ciliaris.
rivularis, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 266.—Ins. Mascar.
robusta, *Liebm. Ind. Sem. Hort. Haun.* (1852) 20.—*Cf. Ann. Sc. Nat.* Sér. III. xix. (1853) 366.—Mexic.
rotundata, *Griseb. Fl. Brit. W. Ind.* 157.—Ind. occ.
rubiaefolia, *Blume, Mus. Bot. Lugd. Bat.* ii. 49.—Guatemala.
rufa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 220.—Jamaic.

PILEA:—
rugosa, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 219.—Ind. occ.
rupicola, Wedd. l. c. 224 = pubescens.
rupipendia, *Wedd. l. c.* Sér. IV. i. (1854) 186.—Afr. trop.
salicifolia, Wedd. l. c. Sér. III. xviii. (1852) 209 = nudicaulis.
Sanctae-crucis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 302.—Ins. S. Crux.
sarcophylla, *Wedd. in DC. Prod.* xvi. I. 109.—Ind. occ.
scripta, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 187.—Reg. Himal.
scrobiculata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 300.—Mexic.
semidentata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 215.—Ind. occ.
serpyllacea, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 296 = serpyllifolia.
serpyllifolia, *Wedd. in DC. Prod.* xvi. I. 107.—Am. trop.
serratifolia, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 235.—Peruv.
serrulata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 213.—Ind. occ.
sessiliflora, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 242.—Ind. occ.
sessilifolia, *Wedd. l. c.* 204.—Ins. Mascar.
smilacifolia, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ind. or.; Java.
spathulata, *Griseb. Cat. Pl. Cub.* 59.—Ind. or.
Spruceana, *Wedd. in DC. Prod.* xvi. I. 161.—Peruv.
Stae-crucis, Auct. = P. Sanctae-crucis.
stelluligera, Griseb. Cat. Pl. Cub. 59 = Wrightiana.
stelluligera, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 184.—Ind. occ.
stipulosa, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 102.—Zeylan.
stolonifera, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 226.—Ind. occ.
strangulata, *Franch. & Sav. Enum. Pl. Jap.* i. 438.—Japon.
strigosa, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 225 = repens.
subcrenata, Blume, Mus. Bot. Lugd. Bat. ii. 54 = lucens.
subcrenata, *Wedd. in DC. Prod.* xvi. I. 148.—Am. trop.
sublucens, *Wedd. l. c.* 130.—Afr. trop.
submissa, *Wedd. l. c.* 151.—Peruv.
subpubera, Miq. in Zoll. Syst. Verz. Ind. Archip. 102, 105 = Achudemia javanica.
subserrata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 207.—N. Granat.
succulenta, Hook. f. in Trans. Linn. Soc. xx. (1847) 182 = microphylla.
succulenta, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 256.—Venezuela.
Swartzii, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 228.—Ind. occ.
Symmeria, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 246.—Reg. Himal.
tenerrima, Miq. in Linnaea, xxvi. (1853) 219 = muscosa.
tenuifolia, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 202.—Reg. Himal.
tetraphylla, *Blume, Mus. Bot. Lugd. Bat.* ii. 50.—Afr. trop.
Thouarsiana, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Mascar.
thymifolia, Blume, Mus. Bot. Lugd. Bat. ii. 44 = globosa.
Trianaeana, *Wedd. in DC. Prod.* xvi. I. 121.—N. Granat.
trianthemoides, Lindl. Coll. Bot. sub t. 4 = muscosa.
tribolata, Blume, Mus. Bot. Lugd. Bat. ii. 48, sphalm. = trilobata.
trichosanthes, *Wedd. in DC. Prod.* xvi. I. 120.—Am. trop.
trichotoma, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 303.—Mexic.
trilobata, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Maurit.
trinervia, *Wight, Ic.* t. 1973.—Ind. or.
triplinervia, Blume, Mus. Bot. Lugd. Bat. ii. 53 = lucens.
uliginosa, *Phil. in Linnaea*, xxx. (1859–60) 199.—Chili.

PILEA:—
umbellata, Wedd. in DC. Prod. xvi. I. 126 = insignis.
umbrosa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 187.—Reg. Himal.
uncidens, Wedd. l. c. Sér. III. xviii. (1852) 224 = dauciodora.
uninervis, Griseb. Cat. Pl. Cub. 59 = serrulata.
urticaefolia, Blume, Mus. Bot. Lugd. Bat. ii. 47 = rupipendia.
Urticella, *Wedd. in DC. Prod.* xvi. I. 157.—Venezuela.
variegata, *Seem. Bot. Voy. Herald*, 194.—N. Granat.
venulosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 57.—Ins. Celebes.
verbascifolia, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ins. Maurit.
virgata, *Wedd. in DC. Prod.* xvi. I. 112.—Jamaic.
vulcanica, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 303.—Mexic.
Wallichii, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 186 = umbrosa.
Wattersii, *Hance, in Journ. Bot.* xxiii. (1885) 327.—China.
Wightii, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 186.—Ind. or.; Malaya.
Wrightiana, *Wedd. in DC. Prod.* xvi. I. 113.—Cuba.
Wydlerii, Blume, Mus. Bot. Lugd. Bat. ii. 49 = ciliaris.
xalapensis, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1853) 227 = pubescens.

PILEANTHUS, Labill. Nov. Holl. Pl. ii. 11. t. 149 (1806). *MYRTACEAE*, Benth. & Hook. f. i. 698.
filifolius, *Meissn. in Journ. Linn. Soc.* i. (1857) 45.—Austral.
Limacis, *Labill. Nov. Holl. Pl.* ii. 11. t. 149.—Austral.
peduncularis, *Endl. in Ann. Wien. Mus.* ii. (1838) 196.—Austral.
vernicosus, F. Muell. Fragm. i. 255 = peduncularis.

PILEOCALYX, Gasparr. in Rendic. Acc. Sc. Nap. vi. (1847) 409 = **Cucurbita**, Tourn.
elegans, Gasparr. l. c. = C. maxima.

PILEOSTEGIA, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 57. *SAXIFRAGEAE*, Benth. & Hook. f. i. 641.
viburnoides, *Hook. f. & Thoms. l. c.* 76.—Reg. Himal.

PILEOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 276 = **Ilex**, Tourn.
mexicana, Turcz. l. c. = I. mexicana, *Black.*

PILEOSTIGMA, Hochst. in Flora, xxix. (1846) 598 = **Bauhinia**, Linn. (Legumin.).
acidum, Benth. in Miq. Pl. Jungh. 261 = B. malabarica.
malabaricum, Benth. l. c. 261 = B. malabarica.
pyrrhocarpum, Hochst. in Flora, xxix. (1846) 598 = B. reticulata.
racemosum, Benth. in Miq. Pl. Jungh. 262 = B. racemosa.
reticulatum, Hochst. ex Benth. l. c. 261 = B. reticulata.
rufescens, Benth. l. c. 262 = B. rufescens.

PILETOCARPUS, Hassk. in Flora, xlix. (1866) 212 = **Aneilema**, R. Br. (Commel.).
protensus, Hassk. l. c.; et Commel. Ind. 15 = A. protensum.
vitiensis, Hassk. ll. cc. 212; 18 = A. vitiense.

PILINOPHYTUM, Klotzsch, in Wiegm. Archiv, vii. (1841) 255 = **Croton**, Linn. (Euphorb.).
capitatum, Klotzsch, l. c. = C. capitatus.
Lindheimeri, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 232 = C. capitatus.

PILIOCALYX, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 185; et in Ann. Sc. Nat. Sér. V. iii. (1865) 225. *MYRTACEAE*, Benth. & Hook. f. Gen. i. 1006.
Boudonini, *Brongn. & Gris, ll. cc.* 186; 226.—N. Caled.
bullatus, *Brongn. & Gris, ll. cc.* xiii. (1866) 470; vi. (1866) 263.—N. Caled.
laurifolius, *Brongn. & Gris, ll. cc.* xii. (1865) 186; iii. (1865) 226.—N. Caled.
micranthus, *Brongn. & Gris, ll. cc.*—N. Caled.

PILIOCALYX:—
robustus, *Brong. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 185; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 224.—N. Caled.
wagapensis, *Brongn. & Gris, ll. cc.* xiii. (1866) 471; vi. (1866) 263.—N. Caled.

PILIOSANTHES, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 121, sphalm. = **Peliosanthes**, Andr. (Haemod.).

PILIOSTIGMA, Walp. Ann. i. 258 (1848–49), sphalm. = PILEOSTIGMA, Hochst. = **Bauhinia**, Linn. (Legumin.).

PILITIS, Lindl. Introd. Nat. Syst. ed. II. 443 (1836) = **Richea**, R. Br. (Epacrid.).
acerosa, Lindl. l. c. = R. acerosa.
Milligani, Hook. f. Fl. Tasm. i. 266, t. 83 = R. Milligani.

PILLERA, Endl. Prod. Fl. Norf. 91 (1833) = **Mucuna**, Adans. (Legumin.).
macrocarpa, Endl. l. c. = M. macrocarpa.

PILOBLEPHIS, Rafin. New Fl. Am. iii. 52 (1836) = **Satureia**, Tourn. (Labiat.).
ericoides, Rafin. l. c. 53 = S. rigida.
rigida, Rafin. l. c. = S. rigida.

PILOCARPUS, Vahl, Eclog. i. 29. t. 10 (1796). *RUTACEAE*, Benth. & Hook. f. i. 299.
amarus, Blanco, Fl. Filip. ed. II. 540 = Lunasia amara.
atropurpureus, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 152, 419 = pennatifolius.
fluminensis, Casar. ex Engl. in Mart. Fl. Bras. xii. II. 134 = pauciflorus.
giganteus, *Engl. l. c.* 136.—Bras.
Goudotianus, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 284.—N. Granat.
grandiflorus, *Engl. in Mart. Fl. Bras.* xii. II. 137.—Bras.
guyanensis, *Van Hall, in Fl. Jard. Pays-Bas*, iii. (1860) 116.—Guian.
heterophyllus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 170.—Cuba.
Humboldtii, *Spreng. Syst.* iv. *Cur. Post.* 126.—Mexic.
latifolius, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. III. vii. (1867) 285.—Guian.
longeracemosus, *Engl. in Mart. Fl. Bras.* xii. II. 135.—Bras.
macrocarpus, *Engl. l. c.* 138.—Bras.
officinalis, *Poehl, Unters. Blaett. Piloc. Off.* (1880) 18, sine diagn.—Am. austr.
parviflorus, Mart. in Nov. Act. Nat. Cur. xi. (1823) 177 = spicatus.
pauciflorus, *A. St. Hil. in Bull. Soc. Philom.* (1823) 131.—Bras.
pennatifolius, *Lem. in Jard. Fleur.* iii. (1852–53) t. 263.—Bras.
pinnatifidus, Carr. in Rev. Hortic. (1868) 60, sphalm. = praec.
pinnatifolius, Engl. in Mart. Fl. Bras. xii. II. 138, sphalm. = praec.
pinnatus, Mart. ex Engl. l. c. = praec.
racemosus, *Vahl, Eclog.* i. 29. t. 10.—Ind. occ.
Riedelianus, *Engl. in Mart. Fl. Bras.* xii. II. 135.—Bras.
Selloanus, *Engl. l. c.* 136.—Bras.
simplex, Baill. Adansonia, xi. (1875) 276 = pennatifolius.
spicatus, *A. St. Hil. in Bull. Soc. Philom.* (1823) 131. t. 10.—Bras.
subcoriaceus, *Engl. in Mart. Fl. Bras.* xii. II. 134.—Bras.
trijugatus, Lem. in Jard. Fleur. iii. (1852–53) sub t. 263 = pennatifolius.

PILOCEREUS, Lem. Cact. Gen. Nov. et Sp. 6 (1839) = **Cereus**, Mill. (Cact.).
Andryanus, Cels, ex Lem. in Rev. Hortic. (1862) 427 = C. caerulescens.
angulosus, Foerst. l. c. 428.—Hab. ?
Arrabidae, Lem. l. c. 429 = C. Arrabidae.
Celsianus, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 40, 185; Lem. in Rev. Hortic. (1862) 428.—Reg. Andin.
chrysomallus, Lem. in Fl. des Serres, iii. (1847) sub t. 242 = C. chrysomallus.

PILOCEREUS:—
coerulescens, Lem. in Rev. Hortic. (1862) 427 = Cereus caerulescens.
Columna, Lem. Cact. Gen. et Sp. Nov. 9 = C. Columna-Trajani.
Cometes, Mittl. ex Foerst. Handb. Cact. ed. I. 357 = C. Cometes.
Consolei, Lem. in Rev. Hortic. (1862) 427.—Hab.?
Curtisii, Salm-Dyck, Cact. Hort. Dyck. ed. I. 24 = C. Curtisi.
Dantwitzii, Hort. Haage.—Cf. Gard. Chron. (1873) 7. f. 1.—Peruv.
divaricatus, Lem. in Rev. Hortic. (1862) 427 = C. divaricatus.
Engelmanni, Lem. Illustr. Hortic. ix. (1862) Misc. 98 = C. Engelmanni.
fimbriatus, Lem. in Rev. Hortic. (1862) 427 = C. grandispinus.
floccosus, Lem. Illustr. Hortic. (1866) sub t. 470 = C. floccosus.
Forsterii, Sencke, ex Lem. in Rev. Hortic. (1862) 428 = *Houlletii*.
fossulatus, Croucher, in Gard. Chron. (1873) 983. f. 197 (= seq.?)—Hab.?
fossulatus, Labour. in Rev. Hortic. Sér. IV. iv. (1855) 25.—Bolivia.
foveolatus, Labour. ex Lem. l. c. (1862) 428 = *Celsianus*.
glaucescens, Labour. Cact. 279 = C. glaucescens.
grandispinus, Lem. in Rev. Hortic. (1862) 427 = C. grandispinus.
Haworthii, Console, l. c. 428 = C. Haworthii.
Hermentianus, Lem. Cact. 63, nomen = C. Hermentianus.
Houlletianum, Lem. in Rev. Hortic. (1862) 428, in syn. = C. virens.
Houlletii, Lem. l. c.—Mexic.
jubatus, Salm-Dyck, Cact. Hort. Dyck. ed. I. 24; Lem. l. c. 427 = C. Cometes.
leucocephalus, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 126.—Mexic.
militaris, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 40 = C. chrysomallus.
Moritzianus, Lem. Cact. 63, nomen = C. Moritzianus.
niger, Poit. in Rev. Hortic. Sér. II. iv. (1845–46) 289 = C. niger.
nigricans, Sencke, ex Lem. in Illustr. Hortic. (1866) Misc. 20 = C. nigricans.
oligogonus, Foerst. ex Lem. in Rev. Hortic. (1862) 428 = C. virens.
Plumieri, Lem. l. c. 427 = C. polygonus.
polyedrophorus, Lem. l. c. 428.—Bras.
polygonus, Salm-Dyck, Cact. Hort. Dyck. 24, in obs. = C. polygonus.
polylophus, Salm-Dyck, l. c. 24; ed. II. 184 = C. polylophus.
Robini, Lem. Illustr. Hort. xi. (1864) Misc. 74.—Cuba.
Schottii, Lem. l. c. 428 = C. Schottii.
scoparius, Poselg. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 126.—N. Mexic.
senilis, Lem. Cact. Gen. et Sp. Nov. 6 = C. senilis.
sublanatus, Foerst. ex Haage, Verz. Cact. 22 = C. sublanatus.
tilophorus, Lem. in Rev. Hortic. (1862) 428 = C. virens.
Vellozii, Lem. l. c. 427 = C. fluminensis.
virens, Lem. Illustr. Hortic. (1866) Misc. 20, in obs. = C. virens.
Williamsii, Lem. in Rev. Hortic. (1862) 428 (an seq.?).—Hab.?
Williamsii, Schelh. Cat. (1845); ex Foerst. Handb. Cact. ed. I. 354 = C. senilis.

PILOGYNE, Schrad. Ind. Sem. Hort. Gotting. (1835).—Cf. Linnaea, xx. (1837) Litt. 89 = **Melothria**, Linn. (Cucurb.).
affinis, Schrad. in Eckl. & Zeyh. Enum. 278 = M. punctata.
cuspidata, Schrad. l. c. = M. punctata.
dilatata, Schrad. l. c. = M. cordata.
Ecklonii, Schrad. l. c. = M. punctata.
Garcini, Harv. Thes. Cap. i. t. 96 = Trochomeria Hookeri.
Garcini, Wight, in Ann. & Mag. Nat. Hist. Ser. I. viii. (1843) 268 = Blastania Garcini.
lucida, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 36. t. 6 = M. lucida.
membranacea, Schrad. in Eckl. & Zeyh. Enum. 278 = M. punctata.

PILOGYNE:—
Peneyana, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 38 = Melothria Peneyana.
setosa, Regel, Cat. Pl. Hort. Aksakov. 111.—Hab.?
suavis, Schrad. Ind. Sem. Hort. Gotting. (1835).—Cf. Linnaea, xi. (1837) Litt. 89 = M. punctata.
tenuifolia, Schrad. in Linnaea, xii. (1838) 412 = M. punctata.
velutina, Schrad. in Eckl. & Zeyh. Enum. 278; et in Linnaea, xii. (1838) 412 = M. velutina.
Zehneria, Steud. Nom. ed. II. ii. 334 = M. Baueriana.

PILOISA, Rafin. Sylva Tellur. 43 (1838) = **Cordia**, Linn. (Borag.).
curassavica, Rafin. l. c. = C. curassavica.
globosa, Rafin. l. c. = C. globosa.

PILOPHORA, Jacq. Fragm. 32. t. 35 (1800) = **Manicaria**, Gaertn. (Palm.).
saccifera, H. Wendl. in Kerch. Palm. 253 = M. saccifera.
testicularis, Jacq. Fragm. 32 = M. saccifera.

PILOPUS, Rafin. Fl. Tellur. ii. 102 (1836) = **Grisebachia**, Klotzsch [1838] (Ericac.).
nodiflora, Rafin. l. c. = G. plumosa.

PILOREA, Rafin. Fl. Tellur. ii. 80 (1836) = **Wahlenbergia**, Schrad. (Campan.).
graminifolia, Rafin. l. c. = W. graminifolia.

PILOSANTHUS, Steud. Nom. ed. II. ii. 331 (1841) sphalm. = PSILOSANTHUS, Neck. = **Liatris**, Schreb. (Compos.).

PILOSELLA, Kostel. Hort. Prag. 104 (1844) = **Sisymbrium**, Tourn. (Crucif.).
Thaliana, Kostel. l. c. = S. Thalianum.

PILOSELLA, [Rupp.] Sch. Bip. fratt. in Flora, xlv. (1862) 417 = **Hieracium**, Tourn. (Compos.) (spp. Am. exceptis).
abscissa, Sch. Bip. fratt. in Flora, xlv. (1862) 436 = H. abscissum.
adenocephala, Sch. Bip. in Linnaea, xxxiii. (1864–65) 759.—Bolivia.
albiflora, Sch. Bip. fratt. in Flora, xlv. (1862) 438 = H. albiflorum.
albocinerea, Arv. in Bull. Soc. Dauph. (1880) 281.—Gallia.
alpicola, Sch. Bip. fratt. in Flora, xlv. (1862) 426 = H. alpicolum.
anchusoides, Arv. in Bull. Soc. Dauph. (1880) 281.—Gallia.
anthura, Sch. Bip. fratt. in Flora, xlv. (1862) 440 = H. anthurum.
arguta, Sch. Bip. fratt. l. c. 438 = H. argutum.
argyrocoma, Sch. Bip. fratt. l. c. 422 = H. Pilosella.
aurantiaca, Sch. Bip. fratt. l. c. 426 = H. aurantiacum.
Auricula, Sch. Bip. fratt. l. c. 425 = H. Auricula.
auriculaeformis, Sch. Bip. fratt. l. c. 424 = H. *bifurcum* ×, Bieb.
Avilae, Sch. Bip. fratt. l. c. 437 = H. Avilae.
Bauhini, Arv. in Bull. Soc. Dauph. (1880) 280 = H. Bauhini.
biflora, Arv. l. c. 279.—Gallia.
bifurca, Sch. Bip. fratt. in Flora, xlv. (1862) 423 = H. *bifurcum* ×, Bieb.
bihariensis, Arv. in Bull. Soc. Dauph. (1880) 280 = H. bihariense.
Blyttiana, Sch. Bip. fratt. in Flora, xlv. (1862) 425 = H. Blyttianum.
boliviensis, Sch. Bip. fratt. l. c. 436 = Crepis boliviensis.
brachiata, Sch. Bip. fratt. l. c. 424 = H. brachiatum.
brachyphylla, Sch. Bip. fratt. l. c. 426 = H. floribundum.
carcina, Arv. in Bull. Soc. Dauph. (1880) 282.—Gallia.
caroliniana, Sch. Bip. fratt. in Flora, xlv. (1862) 437 = H. marianum.
castellana, Sch. Bip. fratt. l. c. 425 = H. *bifurcum*, Bieb.
cernua, Sch. Bip. fratt. l. c. 423 = H. cernuum.
chilensis, Sch. Bip. fratt. l. c. 436 = H. chilense.
comata, Sch. Bip. fratt. l. c. 438 = H. comatum.
cymigera, Arv. in Bull. Soc. Dauph. (1880) 280 = H. cymosum.

PILOSELLA:—
cymosa, Sch. Bip. fratt. in Flora, xlv. (1862) 429 = Hieracium cymosum.
debilis, Sch. Bip. fratt. l. c. 438 = H. debile.
decolorans, Sch. Bip. fratt. 426 = H. decolorans.
dubia, Sch. Bip. fratt. l. c. 429 = H. dubium.
echioides, Sch. Bip. fratt. l. c. 431 = H. echioides.
fallax, Arv. in Bull. Soc. Dauph. (1880) 282 = H. praealtum.
Faurei, Arv. l. c. 279 = H. lycopsifolium.
flaccida, Sch. Bip. fratt. in Flora, xlv. (1862) 437 = H. flaccidum.
florentina, Sch. Bip. fratt. l. c. 427 = H. piloselloides.
floribunda, Arv. in Bull. Soc. Dauph. (1880) 280 = H. floribundum.
Friesii, Sch. Bip. fratt. in Flora, xlv. (1862) 436 = H. Schultzii.
frigida, Sch. Bip. fratt. l. c. 438 = H. frigidum.
fulvipes, Sch. Bip. fratt. l. c. 436 = H. fulvipes.
fusca, Arv. in Bull. Soc. Dauph. (1880) 280 = H. aurantiacum?
Fussiana, Sch. Bip. fratt. in Flora, xlv. (1862) 427 = H. piloselloides.
glacialis, Sch. Bip. fratt. l. c. 426 = H. glaciale.
glaucifolia, Sch. Bip. fratt. l. c. 436 = H. glaucifolium.
glomerata, Arv. in Bull. Soc. Dauph. (1880) 281 = H. glomeratum.
gracilis, Sch. Bip. fratt. in Flora, xlv. (1862) 439 = H. gracile.
Hoppeana, Sch. Bip. fratt. l. c. 422 = H. Pilosella.
horrida, Sch. Bip. fratt. l. c. 439 = H. horridum.
hybrida, Sch. Bip. fratt. 424 = H. hybridum.
hyperborea, Sch. Bip. fratt. l. c. 429 = H. hyperboreum.
incana, Sch. Bip. fratt. l. c. 431 = H. incanum.
jubata, Sch. Bip. fratt. l. c. 438 = H. jubatum.
juncea, Sch. Bip. fratt. l. c. 436 = H. junceum.
junciformis ×, Arv. in Bull. Soc. Bot. Fr. xxvi. (1879) Bibl. 39.—Gallia.
Laggeri, Sch. Bip. fratt. in Flora, xlv. (1862) 426 = H. tomentosum.
Lagopus, Sch. Bip. fratt. l. c. 436 = H. Lagopus.
leptocephala, Sch. Bip. fratt. l. c. 437 = H. leptocephalum.
leucotricha, Sch. Bip. fratt. l. c. = H. leucotrichum.
longipes, Sch. Bip. fratt. l. c. 423 = H. praealtum.
longipila, Sch. Bip. fratt. l. c. 439 = H. longipilum.
macrantha, Sch. Bip. fratt. l. 422 = H. Pilosella.
macrotrichia, Sch. Bip. fratt. l. c. 426 = H. macrotrichum.
magellanica, Sch. Bip. fratt. l. c. 438 = H. magellanicum.
Mandonii, Sch. Bip. in Linnaea, xxxiii. (1864–65) 760.—Bolivia.
mariana, Sch. Bip. fratt. in Flora, xlv. (1862) 438 = H. marianum.
mexicana, Sch. Bip. fratt. l. c. 436 = H. mexicanum.
micromegas, Sch. Bip. fratt. l. c. 426 = H. petraeum.
multiflora, Sch. Bip. fratt. l. c. 430 = H. sabinum.
myosotidifolia, Sch. Bip. fratt. l. c. 438 = H. mysotidifolium.
myriadenia, Sch. Bip. fratt. l. c. 425 = H. myriadenum.
Nestleri, Sch. Bip. fratt. l. c. 430 = H. cymosum.
niveopappa, Sch. Bip. fratt. l. c. 435 = H. niveopappum.
norvegica, Arv. in Bull. Soc. Dauph. (1880) 281 = H. norvegicum.
officinarum, Sch. Bip. fratt. in Flora, xlv. (1862) 421, 422 = H. Pilosella.
orthotricha, Sch. Bip. fratt. l. c. 436 = H. orthotrichum.
paniculata, Sch. Bip. fratt. l. c. 440 = H. paniculatum.
patagonica, Sch. Bip. fratt. l. c. 438 = H. patagonicum.
Peleteriana, Sch. Bip. fratt. l. c. 421 = H. Pilosella.
pennsilvanica, Sch. Bip. fratt. l. c. 439 = H. Gronovii.
persica, Sch. Bip. fratt. l. c. 431 = H. sabinum.
petraea, Sch. Bip. fratt. l. c. 426 = H. petraeum.
poliotricha, Arv. in Bull. Soc. Dauph. (1880) 281 = H. cymosum.
praealta, Sch. Bip. fratt. in Flora, xlv. (1862) 429, 431 = H. praealtum.
praemorsiformis, Sch. Bip. fratt. l. c. 435 = H. praemorsiforme.

PILOSELLA :—
pratensis, Sch. Bip. fratt. in Flora, xlv. (1862) 426 = Hieracium pratense.
primulaeformis, Arv. in Bull. Soc. Dauph. (1880) 280. —Gallia.
procera, Sch. Bip. fratt. in Flora, xlv. (1862) 431 = H. procerum.
Pseudo-Pelleteriana, Magnier.—Cf. Bull. Soc. Bot. Fr. xxvii. (1880) Bibl. 170 = H. Pilosella.
pumila, Sch. Bip. fratt. in Flora, xlv. (1862) 426 = H. glaciale.
Radula, Sch. Bip. fratt. l. c. 431 = H. Balansae.
relicina, Sch. Bip. fratt. l. c. 439 = H. relicinum.
rhodantha, Sch. Bip. fratt. l. c. 425 = H. rhodanthum.
rosea, Sch. Bip. fratt. l. c. 437 = H. roseum.
Rothiana, Sch. Bip. fratt. l. c. 431 = H. setigerum.
sabina, Sch. Bip. fratt. l. c. 430 = H. sabinum.
scabra, Sch. Bip. fratt. l. c. 439 = H. scabrum.
Schultesii, Sch. Bip. fratt. l. c. 425 = H. *bifurcum* ×, Bieb.
Scouleri, Sch. Bip. fratt. l. c. 439 = H. Scouleri.
semiglabrata, Sch. Bip. fratt. l. c. 436 = H. Commersonii.
serpyllifolia, Sch. Bip. fratt. l. c. 425 = H. serpyllifolium.
setigera, Arv. in Bull. Soc. Dauph. (1880) 281 = H. setigerum.
sparsa, Sch. Bip. fratt. in Flora, xlv. (1862) 428 = H. sparsum.
sparsipila, Magnier.—Cf. Bull. Soc. Bot. Fr. xxvii. (1880) Bibl. 170 (Quid ?).—Gallia.
spathulata, Sch. Bip. fratt. in Flora, xlv. (1862) 439 = H. compositum.
sphaerocephala, Sch. Bip. fratt. l. c. 423 = H. cernuum.
stolonifera, Sch. Bip. fratt. l. c. = H. stoloniflorum.
stoloniflora, Sch. & Sch. Bip. fratt. l. c. = H. stoloniflorum.
strigosa, Sch. Bip. fratt. l. c. 436 = H. strigosum.
stuposa, Sch. Bip. fratt. l. c. 435 = H. stuposum.
suecica, Sch. Bip. fratt. l. c. 425 = H. suecicum.
sulphurea, Sch. Bip. fratt. l. c. 425 = H. sulphureum.
Tenoreana, Sch. Bip. fratt. l. c. 422 = H. Pilosella.
thyrsoidea, Sch. Bip. fratt. l. c. 436 = H. thyrsoideum.
tovarense, Sch. Bip. fratt. l. c. 437 = H. tovarense.
trichodonta, Sch. Bip. in Linnaea, xxxiii. (1864–65) 760.—Bolivia.
tristis, Sch. Bip. fratt. in Flora, xlv. (1862) 439 = H. triste.
Urvillei, Sch. Bip. fratt. l. c. 437 = H. Urvillei.
Vahlii, Sch. Bip. fratt. l. c. 426 = H. Vahlii.
velutina, Sch. Bip. fratt. l. c. 422 = H. Pilosella.
venosa, Sch. Bip. fratt. l. c. 437 = H. venosum.
versicolor, Arv. in Bull. Soc. Dauph. (1880) 279 = H. floribundum ?
Villarsii, Sch. Bip. fratt. in Flora, xlv. (1862) 424 = H. stoloniflorum ?
Visianii, Sch. Bip. fratt. l. c. 424 = H. cernuum.
vulgaris, Arv. in Bull. Soc. Dauph. (1880) 278 = H. dubium.
Wolfgangiana, Arv. l. c. 281 = H. pratense.
Ziziana, Sch. Bip. fratt. in Flora, xlv. (1862) 430 = H. praealtum.

PILOSPERMA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 243. *GUTTIFERAE*, Benth. & Hook. f. i. 171.
caudatum, *Planch. & Triana, l. c.* 244.—N. Granat.

PILOSTACHYS, Rafin. New Fl. Am. iv. 89 (1836) = **Polygala**, Tourn.
hyemalis, Rafin. l. c. = Polyg. hyemalis.
nana, Rafin. l. c. = Polyg. nana.

PILOSTYLES, Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 21. t. 1 = **Apodanthes**, Poit. (Cytin.).
aethiopica, Welw. in Trans. Linn. Soc. xxvii. (1869) 67.—Afr. trop.
Berterii, Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 21. t. 1 = A. Berterii.
Blanchetii, R. Br. in Trans. Linn. Soc. xix. (1845) 247 = A. Blanchetii.
Calliandrae, R. Br. l. c. = A. Calliandrae.
Caulotreti, Hook. f. in DC. Prod. xvii. 116.—Venezuela.
Haussknechtii, Boiss. in Bibl. Univ. Genèv. xxv. (1866) 259.—Persia.
Ingae, Hook. f. in DC. Prod. xvii. 116.—N. Granat.
Thurberi, A. Gray, in Mem. Am. Acad. N. S. v. (1855) 326.—N. Mexic.

PILOTHECA, Mitch. Journ. Trop. Austral. 347 (1848), sphalm. = **Philotheca**, Rudge (Rutac.).

PILOTRICHUM, Hook. f. & T. Anders. in Hook. f. Fl. Brit. Ind. i. 141 (1872), sphalm. = PTILOTRICHUM, C. A. Mey. = **Alyssum**, Tourn. (Crucif.).

PILUMNA, Lindl. Bot. Reg. (1844) Misc. 73 = **Trichopilia**, Lindl. (Orchid.).
fragans, Lindl. l. c. 74 = T. fragrans.
laxa, Lindl. l. c. 73 = T. laxa.
nobilis, Reichb. f. in Linnaea, xxii. (1849) 843 = T. nobilis.
Wageneri, Reichb. f. in Bonplandia, ii. (1854) 15 = T. Wageneri.

PIMECARIA, Rafin. Alsog. Am. 64 (1838) = **Myrica**, Linn.
odorata, Rafin. l. c. = M. cerifera.

PIMELA, Lour. Fl. Cochinch. 407 (1790) = **Canarium**, Linn. (Burser.).
acutifolia, Blume, Mus. Bot. Lugd. Bat. i. 221 = C. rostratum.
alba, Lour. Fl. Cochinch. 408 = C. album.
altissima, Blume, Mus. Bot. Lugd. Bat. i. 225 = C. altissimum.
angustifolia, Blume, l. c. 226 = C. angustifolium.
caryophyllacea, Blume, l. c. 222.—China.
decumana, Blume, l. c. 223 = C. decumanum.
denticulata, Blume, l. c. 226 = C. denticulatum.
dichotoma, Blume, l. c. 222 = C. dichotomum.
glabra, Blume, l. c. = C. balsamiferum.
hirsuta, Blume, l. c. 223 = C. hirsutum.
hispida, Blume, l. c. 224 = C. hispidum.
Kipella, Blume, l. c. 220 = C. Kipella.
Kitengo, Blume, l. c. 221 = C. Kitenga.
laxiflora, Blume, l. c. 221 = C. longiflorum.
legitima, Blume, l. c. 222 = C. legitimum.
luzonica, Blume, l. c. 220 = C. luzonicum.
nigra, Lour. Fl. Cochinch. 407 = C. Pimela.
oleosa, Lour. l. c. = C. microcarpum.
paucijuga, Blume, Mus. Bot. Lugd. Bat. i. 226 = C. balsamiferum.
rigida, Blume, l. c. 222 = C. rigidum.
stricta, Blume, l. c. 226 = C. strictum.
villosa, Blume, l. c. 223 = C. Cumingii.

PIMELANDRA, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 88. *MYRSINEAE*, Benth. & Hook. f. ii. 646.
disticha, *Benth. & Hook. Gen.* ii. 647.—Ins. Philipp.
erecta, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 531.—Reg. Himal.
eugeniaefolia, *Benth. & Hook. f. Gen.* ii. 647.—Reg. Himal.
fuliginosa, *Benth. & Hook. f. l. c.*—Java.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 531.—Reg. Himal.
semidentata, *Benth. & Hook. f. Gen.* ii. 647.—Java.
Teysmanniana, Benth. & Hook. f. l. c. = Wallichii.
Wallichii, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 88 ; *et in DC. Prod.* viii. 106.—Malaya.

PIMELEA, Banks & Soland. ex Gaertn. Fruct. i. 186. t. 39 (1788). *THYMELAEACEAE*, Benth. & Hook. f. iii. 189.
BANKSIA, Forst. Char. Gen. 7. t. 4 (1776) (nomen prius).
CALYPTROSTEGIA, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 72.
COOKIA, J. F. Gmel. Syst. ii. 24 (1791).
GYMNOCOCCA, Fisch. & Mey. in Ind. Sem. Hort. Petrop. x. 46 (1845).
HETEROLAENA, Fisch. & Mey. l. c. 47 (1845).
MACROSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 177.
THECANTHES, Wikstr. in Vet. Akad. Handl. Stockh. (1818) 271.
aeruginosa, F. Muell. Fragm. vii. 2 = sylvestris.
affinis, Loud. Hort. Brit. Suppl. iii. 609 = rosea.
alpina, *F. Muell. ex Meissn. in DC. Prod.* xiv. 511.—Austral.
altior, *F. Muell. Fragm.* i. 84.—Austral.
ammocharis, *F. Muell. in Hook. Kew Journ.* ix (1857) 24.—Austral.
angustifolia, *R. Br. Prod.* 360.—Austral.
angustifolia, F. Muell. ex Meissn. in DC. Prod. xiv. 501 = stricta.

PIMELEA :—
arenaria, *A. Cunn. in Bot. Mag.* t. 3270.—N. Zel.
argentea, *R. Br. Prod.* 362.—Austral.
axiflora, *F. Muell. First Gen. Rep.* (1853) 17.—Austral.
axillaris, Banks & Soland. ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 280 = virgata.
Baxteri, A. Cunn. ex Meissn. in DC. Prod. xiv. 507 = imbricata.
Behrii, Schlecht. in Linnaea, xx. (1853–55) 583 = octophylla.
Bowmanni, *F. Muell. ex Benth. Fl. Austral.* vi. 30.—Austral.
brachyphylla, *Benth. l. c.* 11.—Austral.
brevifolia, *R. Br. Prod.* 359.—Austral.
brevifolia, Meissn. in DC. Prod. xiv. 497 = brachyphylla.
brevituba, *Fawc. in H. O. Forbes, Nat. Wand. East. Archip.* 516.—Ins. Timor.
buxifolia, *Hook. f. Handb. N. Zeal. Fl.* 243.—N. Zel.
campicola, A. Cunn. ex Benth. Fl. Austral. vi. 16 = glauca.
cernua, R. Br. Prod. 359 = spathulata.
cinerea, *R. Br. l. c.* 361.—Austral.
clavata, *Labill. Nov. Holl. Pl.* i. 11.—Austral.
cluytioides, Meissn. in Lehm. Pl. Preiss. ii. 271 = serpyllifolia.
collina, *R. Br. Prod.* 359.—Austral.
collina, A. Cunn. ex Benth. Fl. Austral. vi. 18 = linifolia.
colorans, *A. Cunn. ex Meissn. in DC. Prod.* xiv. 499. —Austral.
colorans, Lindl. in Mitch. Journ. Trop. Austral. 362 = collina.
concreta, *F. Muell. Fragm.* v. 78.—Austral.
congesta, A. Cunn. ex Benth. Fl. Austral. vi. 31 = curviflora.
congesta, R. Cunn. l. c. 32 = hirsuta.
congesta, F. Muell. Fragm. viii. 9 = longifolia.
Cornucopiae, *Vahl, Enum.* i. 305.—Austral.
crinita, Lindl. in Bot. Reg. (1838) Misc. 59 = imbricata.
Cunninghamii, Meissn. in DC. Prod. xiv. 498 = collina.
curviflora, *R. Br. Prod.* 362.—Austral.
decussata, R. Br. l. c. 360 = ferruginea.
denticulata, A. Cunn. ex Meissn. in DC. Prod. xiv. 498 = collina.
dichotoma, Schlecht. in Linnaea, xx. (1847) 581 = flava.
diosmaefolia, Lodd. Bot. Cab. t. 1708 = ferruginea.
diosmifolia, A. Cunn. ex Benth. Fl. Austral. vi. 29 = flava.
distinctissima, F. Muell. First Gen. Rep. (1853) 17 = microcephala.
drupacea, *Labill. Nov. Holl. Pl.* i. 10. t. 7.—Austral.
elachantha, *F. Muell. First Gen. Rep.* (1853) 17.—Austral.
elata, F. Muell. l. c. 17 = ligustrina.
elegans, F. Moore, in Illustr. Hortic. viii. (1861) t. 295 = ligustrina.
Eyrei, *F. Muell. Fragm.* v. 109.—Austral.
ferruginea, *Labill. Nov. Holl. Pl.* i. 10. t. 5.—Austral.
filamentosa, Rudge, in Trans. Linn. Soc. x. (1815) 287. t. 14 = linifolia.
filiformis, *Hook. f. in Hook. Lond. Journ.* vi. (1847) 280.—Tasman.
flava, *R. Br. Prod.* 361.—Austral.
flava, Meissn. in Lehm. Pl. Preiss. i. 605 = sulphurea.
floribunda, *Meissn. in DC. Prod.* xiv. 505.—Austral.
Forrestiana, *F. Muell. Fragm.* xi. 46.—Austral.
glauca, *R. Br. Prod.* 360.—Austral.
Gnidia, *Willd. Sp. Pl.* i. 50.—N. Zel.
graciliflora, Hook. Bot. Mag. t. 3288 = sylvestris.
gracilis, R. Br. Prod. 362 = curviflora.
grandiflora, *Donn, Hort. Cantab.* ed. VI. 5.—Austral.
Gunnii, Hook. f. Fl. Tasm. i. 332 = cinerea.
Haastii, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 396. —N. Zel.
haematostachya, *F. Muell. Fragm.* i. 84.—Austral.
Hendersonii, R. Grah. in Edinb. N. Phil. Journ. xxvi. (1838) 197 = rosea.
heterophylla, *F. Muell. Fragm.* i. 187.—Austral.
Hewardiana, Meissn. in Linnaea, xxvi. (1853–55) 346 = elachantha.
hirsuta, *Meissn. in DC. Prod.* xiv. 513.—Austral.
hispida, *R. Br. Prod.* 360.—Austral.

PIMELEA :—
hispida, A. Cunn. ex Meissn. in DC. Prod. xiv. 503 = rosea.
Holroydi, *F. Muell. Fragm.* vi. 159. t. 59.—Austral.
humilis, *R. Br. Prod.* 361.—Austral.
humilis, Lindl. Bot. Reg. t. 1268 = glauca.
hypericina, A. Cunn. ex Hook. Bot. Mag. t. 3330 = ligustrina.
imbricata, *R. Br. Prod.* 361.—Austral.
incana, R. Br. l. c. = nivea.
intermedia, Hort. ex R. Grah. in Edinb. N. Phil. Journ. xxvi. (1839) 421 = sylvestris.
intermedia, Lindl. Bot. Reg. t. 1439 = glauca.
involucrata, Banks & Soland. ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 274 = linifolia.
laevigata, *Gaertn. Fruct.* i. 186. t. 39. f. 1.—N. Zel.
lanata, R. Br. Prod. 360 = hispida.
lanata, Hensl. in Maund, Botanist, ii. t. 61 = sericea.
lanceolata, Hort. Nois. ex Meissn. in DC. Prod. xiv. 515 = pauciflora.
latifolia, *R. Br. Prod.* 362.—Austral.
Lehmanniana, *Meissn. in Lehm. Pl. Preiss.* i. 603.—Austral.
leptospermoides, *F. Muell. Fragm.* vii. 2.—Austral.
leptostachya, *Benth. Fl. Austral.* vi. 24.—Austral.
ligustrina, *Labill. Nov. Holl. Pl.* i. 9. t. 3.—Austral.
ligustrina, Sieber, ex Meissn. in DC. Prod. xiv. 498 = linifolia.
linariifolia, A. Cunn. ex Meissn. in DC. l. c. 515 = microcephala.
Lindleyana, Meissn. l. c. 499 = spathulata.
linifolia, A. Cunn. ex Meissn. l. c. 497 = spathulata.
linifolia, *Sm. Bot. Nov. Holl.* 31. t. 11.—Austral.
linoides, A. Cunn. in Field, N. S. Wales, 326 = linifolia.
longiflora, *R. Br. Prod.* 361.—Austral.
longifolia, *Banks & Soland. ex Wikstr. in Vet. Akad. Handl. Stockh.* (1818) 280.—N. Zel.
Lyallii, *Hook. f. Fl. N. Zel.* i. 222.—N. Zel.
macrocephala, Hook. Bot. Mag. t. 4543 = suaveolens.
marginata, Meissn. in DC. Prod. xiv. 497 = collina.
Maxwelli, *F. Muell. ex Benth. Fl. Austral.* vi. 12.—Austral.
Menkeana, Lehm. Pl. Preiss. i. 604 = suaveolens.
micrantha, F. Muell. ex Meissn. in Linnaea, xxvi. (1853–55) 351 = curviflora.
microcephala, *R. Br. Prod.* 361.—Austral.
microcephala, Meissn. in Lehm. Pl. Preiss. i. 606 = imbricata.
Milligani, *Meissn. in DC. Prod.* xiv. 509.—Tasman.
Mitchellii, Meissn. l. c. 506 = collina.
modesta, Meissn. in Lehm. Pl. Preiss. ii. 268 = brevifolia.
Muelleri, Meissn. in Linnaea, xxvi. (1853–55) 351 = curviflora.
myriantha, Meissn. in Lehm. Pl. Preiss. i. 607 = argentea.
myrtifolia, Schlecht. in Linnaea, xx. (1847) 582 = glauca.
nana, R. Grah. in Edinb. N. Phil. Journ. xxix. (1840) 174 = imbricata.
nervosa, *Meissn. in Lehm. Pl. Preiss.* ii. 269.—Austral.
Neypergiana, Hort. ex Decne. in Rev. Hortic. Sér. IV. i. (1852) 80 = Preissii.
Nieppergiana, [Carr.] l. c. (1881) 440 = praec.
nivea, *Labill. Nov. Holl. Pl.* i. 10. t. 6.—Tasman.
Novae-Zeylandiae, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 609 = laevigata.
nutans, Meissn. in Linnaea, xxvi. (1853–55) 348 = spathulata.
octophylla, *R. Br. Prod.* 361.—Austral.
ovalifolia, Meissn. in DC. Prod. xiv. 502 = hirsuta.
paludosa, R. Br. Prod. 360 = linifolia.
parviflora, Lodd. ex Meissn. in DC. Prod. xiv. 515 = pauciflora.
parvifolia, Meissn. in Linnaea, xxvi. (1853–55) 345 = flava.
pauciflora, *R. Br. Prod.* 360.—Austral.
Paxtoni, Hort. ex Regel, in Gartenfl. (1852) 104 = nivea ?
penicillaris, *F. Muell. in Melb. Chemist* (Octob. 1883).—Austral.
petraea, *Meissn. in Linnaea*, xxvi. (1853–55) 347.—Austral.
petrophila, *F. Muell. l. c.* xxv. (1852) 442.—Austral.
phylicoides, *Meissn. in Lehm. Pl. Preiss.* ii. 271.—Austral.

PIMELEA :—
physodes, *Hook. Ic. Pl.* t. 865.—Austral.
pilibunda, A. Cunn. ex Benth. Fl. Austral. vi. 21 = imbricata.
pilosa, Willd. Sp. Pl. i. 50 = virgata.
Preissii, *Meissn. in Lehm. Pl. Preiss.* i. 601.—Austral.
Preissii, Schlecht. in Linnaea, xx. (1847) 581 = glauca.
prinifolia, Nois. ex Steud. Nom. ed. II. ii. 334 = glauca.
propinqua, A. Cunn. ex Meissn. in DC. Prod. xiv. 512 = curviflora.
prostrata, Willd. Sp. Pl. i. 51 = laevigata.
punicea, *R. Br. Prod.* 359.—Austral.
pygmaea, *F. Muell. in Linnaea*, xxvi. (1853–55) 346.—Austral.
quadrifaria, Banks & Soland. ex Meissn. in DC. Prod. xiv. 517 = laevigata.
ramulosa, A. Cunn. ex Meissn. l. c. 515 = pauciflora.
rigida, Meissn. l. c. 502 = linifolia.
rosea, *R. Br. Prod.* 360.—Austral.
sanguinea, *F. Muell. Fragm.* i. 84.—Austral.
Schlechtendaliana, Meissn. in Bot. Zeit. vi. (1848) 394 = glauca.
sericea, *R. Br. Prod.* 361.—Austral.
sericea, Meissn. in DC. Prod. xiv. 517 = arenaria.
sericeo-villosa, *Hook. f. Handb. N. Zeal. Fl.* 245.—N. Zel.
sericostachya, *F. Muell. Fragm.* iv. 162.—Austral.
serpyllifolia, *R. Br. Prod.* 360.—Austral.
Shepherdii, *Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 609.—Austral. ?
Shuttleworthiana, Meissn. in DC. Prod. xiv. 518 = argentea.
silvestris, F. Muell. Fragm. vii. 5, *vide* sylvestris.
simplex, *F. Muell. in Linnaea*, xxv. (1852) 443.—Austral.
spathulata, *Labill. Nov. Holl. Pl.* i. 9. t. 4.—Austral.
speciosa, Regel, Cat. Pl. Hort. Aksakov. 111 (Quid ?).—Hab. ?
spectabilis, *Lindl. Swan River App.* 41.—Austral.
spicata, *R. Br. Prod.* 362.—Austral.
spiculigera, F. Muell. ex Benth. Fl. Austral. vi. 23 = spicata.
stricta, *Meissn. in Linnaea*, xxvi. (1853–55) 348.—Austral.
suaveolens, *Meissn. in Lehm. Pl. Preiss.* i. 603.—Austral.
sulphurea, *Meissn. in Bot. Zeit.* vi. (1848) 396.—Austral.
sylvestris, *R. Br. Prod.* 361.—Austral.
thymifolia, Presl, Bot. Bemerk. 107 = curviflora.
tinctoria, Meissn. in Lehm. Pl. Preiss. i. 603 = suaveolens.
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 243.—N. Zel.
trichostachya, *Lindl. in Mitch. Journ. Trop. Austral.* 355.—Austral.
trichostachya, F. Muell. Fragm. vii. 7 = simplex.
umbratica, A. Cunn. ex Meissn. in DC. Prod. xiv. 510 = drupacea.
uniflora, Regel, Cat. Pl. Hort. Aksakov. 111 (Quid ?).—Hab. ?
Urvilleana, *A. Rich. Fl. N. Zél.* 175.—N. Zel.
Verschaffeltii, C. Morr. in Ann. Soc. Bot. Gand, iii. (1847) 451. t. 166 = spectabilis.
vestita, Meissn. in DC. Prod. xiv. 513 = argentea.
villifera, A. Cunn. ex Benth. Fl. Austral. vi. 32 = hirsuta.
villifera, *Meissn. in Lehm. Pl. Preiss.* ii. 271.—Austral.
villosa, Banks & Soland. ex Meissn. in DC. Prod. xiv. 517 = arenaria.
villosa, Banks & Soland. ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 282 = sericea.
villosa, Meissn. in DC. Prod. xiv. 508 = longiflora.
viminea, Schlecht. in Linnaea, xx. (1847) 583 = octophylla.
virgata, *Vahl, Enum.* i. 306.—N. Zel.
viridula, Lindb. in Finsk. Vet. Soc. Forhandl. ix. (1867) 61, cum ic. = clavata.
Weippugiana, Paxt. Mag. Bot. xv. (1849) 191, sphalm. = *Neypergiana*.

PIMELEODENDRON, Muell. Arg. in DC. Prod. xv. II. 1143 (1866) = seq.

PIMELODENDRON, Hassk. in Versl. en Med. Kon. Akad. Wetensch. iv. (1855) 140. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 331.
STOMATOCALYX, Muell. Arg. in DC. Prod. xv. II. 1142 (1866).
amboinicum, *Hassk. in Versl. en Med. Kon. Akad. Wetensch.* iv. (1855) 140.—Ins. Molucc.

PIMENTA, Lindl. Coll. Bot. sub t. 19 (1821–25). *MYRTACEAE*, Benth. & Hook. f. i. 717.
AMOMIS, Berg, in Linnaea, xxvii. (1856) 416.
MYRCIA, Soland. ex Lindl. Coll. Bot. sub t. 19 (1821–25).
acris, *Kostel. Allg. Med. Pharm. Fl.* iv. 1526; *Wight, Illustr.* ii. 12.—Ind. or.
acuminata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 270.—Porto Rico.
aromatica, Kostel. Allg. Med. Pharm. Fl. iv. 1525 = officinalis.
citrifolia, Kostel. l. c. = acris.
communis, Benth. & Hook. f. Gen. i. 717 = officinalis.
fulvescens, Blume, Mus. Bot. Lugd. Bat. i. 80 = Eugenia fulvescens.
officinalis, *Lindl. Coll. Bot.* sub t. 19.—Ind. occ.
Pimento, *Griseb. Fl. Brit. W. Ind.* 241.—Ind. occ.
Pseudo-caryophyllus, Blume, Mus. Bot. Lugd. Bat. i. 80 = Eugenia Pseudocaryophyllus.
vulgaris, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 270 (= seq. ?).—Porto Rico.
vulgaris, Lindl. in Loud. Encyc. Pl. 418; Wright, Illustr. ii. 12 = officinalis.

PIMENTELIA, Wedd. Monog. Cinch. 94. t. 27 B (1849). *RUBIACEAE*, Benth. & Hook. f. ii. 33.
glomerata, *Wedd. Monog. Cinch.* 94.—Peruv.

PIMENTELLA, Walp. Ann. ii. 789 (1851–52) = praec.

PIMENTUS, Rafin. Sylva Tellur. 105 (1838) = **Pimenta**, Lindl. etc. (Myrtac.).
aromatica, Rafin. l. c. 158 = Eugenia Greggii.
cotinifolia, Rafin. l. c. 105 = Pimenta acris.
geminata, Rafin. l. c. (Quid ?).—Hab. ?
Gregia, Rafin. l. c. = Eugenia Greggii.
laurinus, Rafin. l. c. = Symplocos spicata.
saligna, Rafin. l. c. = Melaleuca Leucadendron.
triflora, Rafin. l. c. = Eugenia triflora.
vera, Rafin. l. c. = Pimenta officinalis.

PIMIA, Seem. in Bonplandia, x. (1862) 366. *STERCULIACEAE*, Benth. & Hook. f. i. 984.
rhamnoides, *Seem. l. c.*—Ins. Fiji.

PIMPINELE, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = **Pimpinella**, Riv. (Umbell.).
anisa, St. Lag. l. c. = Pimpinella Anisum.

PIMPINELLA, Tourn. ex Rupp. Fl. Jen. ed. Hall. 285 (1745); Adans. Fam. ii. 293 (1763) = **Poterium**, Linn. (Rosac.).
agrimonoides, Gaertn. Fruct. ii. 162 = Pot. hybridum.
minor, Lam. Fl. Fr. ii. 343 = Pot. Sanguisorba.
officinalis, Lam. l. c.; Gaertn. Fruct. ii. 161 = Pot. officinale.
Sanguisorba, Gaertn. l. c. 162 = Pot. Sanguisorba.
spinosa, Gaertn. l. c. = Pot. spinosum.

PIMPINELLA, [Riv.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 893.
ACRONEMA, Edgew. in Trans. Linn. Soc. xx. (1846) 51.
ANISOMETROS, Hassk. in Flora, xxx. (1847) 602.
ANISUM, Riv. ex Linn. Gen. ed. I. 83 (1737).
BUNIUM, Koch, Syn. Fl. Germ. ed. II. 315 (1843).
CHESNEYA, Bertol. in Nov. Comm. Bonon. v. (1842) 427 (Misc. Bot. i. 17. t. 2).
GAYTANIA, Muenst. in Bot. Zeit. i. (1843) 730.
GYMNOSCIADIUM, Hochst. in Flora, xxvii. (1844) 20.
HETERACHAENA, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1846) 577.
LEDEBURIA, Link, Enum. Hort. Berol. i. 286 (1821).
LERESCHIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 127.
MURRITHIA, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 576.
PANCICIA, Vis. Sem. Hort. Patav. (1857) 9.—Cf. Linnaea, xxix. (1857–58) 732.
PETROSCIADIUM, Edgew. in Trans. Linn. Soc. xx. (1846) 51.

PIMPINELLA :—
PLATYRHAPHE, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 56 (1867).
REUTERA, Boiss. Elench. 46 (1838).
SISARUM, [Tourn.] Adans. Fam. ii. 97 (1763).
TRAGIOPSIS, Pomel, Nouv. Mat. Fl. Atl. 139 (1874).
TRAGIUM, Spreng. Pl. Umb. Prod. 26 (1813).
TRAGOSELINUM, Tourn. ex Hall. Enum. Stirp. Helv. ii. 428 (1742).
acaulis, Krock. Fl. Siles. Suppl. i. 341 = P. Saxifraga ?
achilleifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 684.—Reg. Himal.
acronemaefolia, *C. B. Clarke, l. c.* 686.—Reg. Himal.
acuminata, *C. B. Clarke, l. c.*—Reg. Himal.
adscendens, *Dalz. in Hook. Kew Journ.* ii. (1850) 261. —Ind. or.
affinis, *Ledeb. Fl. Ross.* ii. 257.—Reg. Caucas.
alba, Gueldenst. Reisen, ii. 192, &c.—Cf. Ledeb. Fl. Ross. ii. 257 (Quid ?).—Rossia.
albescens, *Franch. Pl. Davidian.* i. 139.—Mongol.
alpestris, Schult. Syst. vi. 837 (Index) = seq.
alpina, Vest, ex Schult. l. c. 387, in obs.; Host, Fl. Austr. i. 399 = P. Saxifraga.
anagodendron, *C. Bolle, Ind. Sem. Hort. Berol.* (1861) *App.* 9.—Ins. Teneriff.
anethifolia, D. Don, Prod. Fl. Nep. 184 = Carum anethifolium.
angelicaefolia, Lam. Encyc. i. 451 = Aegopodium Podagraria.
angustifolia, Gilib. Fl. Lituan. ii. 42 = P. Saxifraga.
angustilobum, *Benth. & Hook. f. Gen.* i. 894.—Chili.
anisetum, *Boiss. & Bal. Diagn.* Ser. II. vi. 78.—Cappadoc.
anisoides, *Brign. in Mem. Acc. Sc. Nap.* i. *Fis.* (1819) 8. t. 1.—Ital.
anisoides, Guss. Cat. Hort. Boccad. (1821) 50 = Gussonii.
anisoides, Tenore, Viagg. in Calabria, 119 = Gussonii.
Anisum, Georgi, Beschr. Russ. Reich. iv. 860 = aromatica.
Anisum, S. G. Gmel. Iter, iv. 13 (Quid ?).—Sibir.
Anisum, *Linn. Sp. Pl.* 264.—Graec.; Aegypt.
anthriscoides, *Boiss. Fl. Orient.* ii. 874.—Syria; Persia.
apiifolia, Boiss. Diagn. Ser. I. x. 25 = Kotschyana.
apiodora, *A. Gray, in Proc. Am. Acad.* vii. (1868) 345. —Calif.
apula, Bernh. Sel. Sem. Hort. Erfurt. (1834).—Cf. in Linnaea, x. (1835) Litt. 79 = P. Tragium ?
arabica, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 130 = cretica.
aromatica, *Bieb. Fl. Taur. Cauc.* i. 241.—Reg. Caucas.
Asopii, Orph. ex Boiss. Fl. Orient. ii. 873 = P. Saxifraga.
aurea, *DC. Prod.* iv. 120.—Persia.
austriaca, Mill. Gard. Dict. ed. VIII. n. 5 = magna.
barbata, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 129. —Persia.
bella, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 11. 686.—Reg. Himal.
betonicaefolia, Willd. ex Ledeb. Fl. Ross. ii. 254 = rotundifolia.
betulaefolia, Adams, ex Ledeb. l. c. = rotundifolia.
bipinnata, Boeber, ex Georgi, Beschr. Russ. Reich. iv. 860 = magna.
bisecta, *Baker, in Journ. Linn. Soc.* xx. (1883) 152.— Madag.
bubonoides, Brot. Fl. Lusit. i. 463 = villosa.
bubonoides, DC. Prod. iv. 121 = Gussonii.
Buchii, *Webb & Berth. Phyt. Canar.* ii. 151.—Ins. Canar.
caespitosa, *Benth. ex Hook. f. Fl. Brit. Ind.* ii. 689.— Reg. Himal.
caffra, *D. Dietr. Syn. Pl.* ii. 947.—Afr. austr.
calicina, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 182.—Japon.
Calverti, Boiss. Diagn. Ser. II. ii. 73 = P. Saxifraga.
Candolleana, *Wight & Arn. Prod.* 369.—Ind. or.
canescens, Loisel. in Desv. Journ. Bot. ii. (1809) 271. t. 10 = P. Tragium.
capensis, *Thunb. Prod. Pl. Cap.* 51.—Afr. austr.
capillacea, Poit. ex Urban, in Mart. Fl. Bras. xi. 1. 342 = Apium Ammi.
capillifolia, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 29.—Turkestan.
cappadocica, *Boiss. & Bal. Diagn.* Ser. II. vi. 79.— Cappadoc.
carvifolia, Benth. & Hook. f. Gen. i. 894 = Carum flexuosum.

PIMPINELLA :—
chiliosciadia, Boiss. & Heldr. Diagn. Ser. II. ii. 86 = Athamanta chiliosciadia.
chrysantha, *Orph. ex Boiss. Diagn.* Ser. II. ii. 77. —Oriens.
cordata, E. Mey. in Drège, Zwei Pfl. Docum. 154, 155 = caffra.
corymbosa, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 131.—As. Min.; Syria.
cretica, *Poir. Encyc. Suppl.* i. 684.—Graec.; As. Min.; Arab.
cretica, Vis. Sem. Hort. Patav. (1844); Hampe, ex Boiss. Fl. Orient. ii. 875 (= *Bunium creticum*, Urv.). —Europ. austr.
crinita, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 131 = Psammogeton biternatum.
crispa, Hornem. Hort. Hafn. i. 289 = P. Saxifraga.
Cumbrae, *Link, in Buch, Beschr. Ins. Canar.* 152.— Ins. Canar.
dahurica, *Turcz. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 13.—Sibir.; Dahur.
Danaa, Bieb. Casp. 163 = Physospermum commutatum.
Dendroselinum, *Webb & Berth. Phyt. Canar.* ii. 152. —Ins. Canar.
Dendrotragium, Webb & Berth. l. c. t. 72 = praec.
depressa, DC. Prod. iv. 120 = P. Tragium.
deverroides, *Boiss. Fl. Orient.* ii. 873.—Persia.
dichotoma, Hort. ex Steud. Nom. ed. II. ii. 335 = Apium Ammi.
dichotoma, Linn. Mant. i. 58 = Carum dichotomum.
dichotoma, Spreng. Syst. i. 883 = Trinia Kitaibelii.
dioica, Bess. Enum. Pl. Volh. 13 = Trinia Henningii.
dioica, Bieb. Fl. Taur. Cauc. i. 242 = Trinia Hoffmannii.
dioica, Hort. ex Link, Enum. Hort. Berol. i. 285 = Apium Ammi.
dioica, Huds. Fl. Angl. ed. II. i. 128 = Trinia vulgaris.
dioica, Lapeyr. Hist. Abr. Pl. Pyr. i. 165 = Seselinanum.
dioica, Linn. Mant. ii. 357 = Trinia glauca.
dioica-rossica, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 44 = Trinia Hoffmanni.
dissecta, Pollini, Fl. Veron. i. 396 = P. Saxifraga.
dissecta, Retz. Obs. iii. 30 = magna.
diversifolia, *DC. Prod.* iv. 122.—Reg. Himal.; China.
domingensis, Willd. ex Schult. Syst. vi. 411 = Apium Ammi.
dumetorum, Boenn. ex Reichb. Fl. Germ. Excurs. 478 = magna.
ebracteata, *Baker, in Journ. Linn. Soc.* xx. (1883) 152. —Madag.
eriocarpos, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 249.—Syria; Assyria.
etbaica, *Schweinf. in Verh. Zool.-Bot. Ver. Wien*, xviii. (1868) 667.—Afr. trop.
ferulifolia, Stokes, Bot. Mat. Med. ii. 147 = Carum ferulaefolium.
flava, C. A. Mey. Verz. Pfl. Cauc. 122 = aurea.
flexuosa, Stokes, Bot. Mat. Med. ii. 145 = Conopodium denudatum.
genevensis, Vill. Hist. Pl. Dauph. ii. 604 = P. Saxifraga.
germanica, Vill. ex Spreng. in Schult. Syst. vi. 391 (Quid ?).—Europ.
glabra, Rostk. & Schmidt, Fl. Sedin. 133 = magna.
glauca, Georgi, Beschr. Russ. Reich. iv. 860 = Trinia Henningii.
glauca, Hoffm. Gen. Umbell. ed. I. 92 = Trinia Kitaibelii.
glauca, Linn. Sp. Pl. 264 = Trinia vulgaris.
glauca, Presl, Fl. Sicil. p. xxv = P. Tragium.
glaucescens, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 130 = barbata.
Gouani, *Benth. & Hook. f. Gen.* i. 894.—Europ.
gracilis, Bisch. Del. Sem. Hort. Heidelb. (1848) 4.— Cf. Linnaea, xxiv. (1851) 189 = affinis.
Griffithiana, *Boiss. Diagn.* Ser. II. ii. 74.—Afghan.
Gussonii, *Bertol. Fl. Ital.* iii. 270.—Ital.
Gymnosciadium, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 14.—Afr. trop.
hastata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 687. —Reg. Himal.
Henningii, Tenore, Syll. Fl. Neap. 153, in syn. = Carum dichotomum.
Heyneana, *Wall. Cat.* n. 566.—Ind. or.
hircina, Mill. Gard. Dict. ed. VIII. n. 3; Leers, Fl. Herborn. 79; Moench, Hass. Pl. Indig. 144 = P. Saxifraga.

PIMPINELLA :—
hirtella, A. Rich. Tent. Fl. Abyss. i. 323 = peregrina.
hispida, Loisel. in Desv. Journ. Bot. ii. (1809) 272 = peregrina.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 686.—Reg. Himal.
humilis, Willd. ex Ledeb. Fl. Ross. ii. 281 = Trinia Hoffmanni.
hyrcina, Krock. Fl. Siles. i. 480 = *hircina*.
integerrima, *Benth. & Hook. f. Gen.* i. 894.—Am. bor.
integrifolia, Wood, Am. Bot. & Flor. 139 = praec.
intermedia, Nees, in Flora, ii. (1819) 293 (Quid ?).— Europ.
involucrata, Jan, ex DC. Prod. iv. 123 = Carum Carvi.
involucrata, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 697 = Schultzia Benthami.
involucrata, Wight & Arn. Prod. 369 = Carum Roxburghianum.
javana, *DC. Prod.* iv. 122.—Java.
Kotschyana, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 133.—Persia.
Kraussiana, Meissn. ex Krauss, in Flora, xxvii. (1844) 431 = caffra.
laciniata, Gilib. Fl. Lituan. ii. 43 = P. Saxifraga.
laciniata, Thore, Chlor. Landes, 108 = magna.
lateriflora, *Dalz. & Gibs. Bomb. Fl.* 106.—Ind. or.
lateriflora, Link, Enum. Hort. Berol. i. 285 = Apium Ammi.
latifolia, Gilib. Fl. Lituan. ii. 42 = P. Saxifraga.
laxiflora, *Baker, in Journ. Linn. Soc.* xxi. (1884) 349. —Madag.
leptophylla, Pers. Syn. i. 324 = Apium leptophyllum.
Leschenaultii, *DC. Prod.* iv. 122.—Ind. or.
leucocarpa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = P. Tragium.
longiradiata, Boiss. & Heldr. in Boiss. Diagn. Ser. II. ii. 75 = P. Tragium.
lucida, Schur, Enum. Pl. Transs. 250 = P. Saxifraga.
lutea, *Desf. Atlant.* i. 255, t. 76.—Alger.
macedonica, *Boiss. & Orph. ex Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 135.—Macedon.
magna, Habl. Phys. Beschr. Taur. 239; et ex Bieb. Fl. Taur. Cauc. i. 240 = P. Saxifraga.
magna, *Linn. Mant.* ii. 217.—Europ.; Reg. Caucas.
major, *Huds. Fl. Angl.* ed. I. 110.—Europ.
major, Mill. Gard. Dict. ed. VIII. n. 1; Gouan, Illustr. 21 = magna.
major, Wallr. ex Steud. Nom. ed. II. ii. 335 = P. Saxifraga.
media, Geners. Elench. n. 273; ex Spreng. in Schult. Syst. vi. 387 = P. Saxifraga.
media, [Weber, in] Wigg. Prim. Fl. Holsat. 26 = magna.
minor, Ehrh. ex Spreng. in Schult. Syst. vi. 386 = P. Saxifraga.
monoica, *Dalz. in Hook. Kew Journ.* iii. (1851) 212.— Ind. or.
multicaulis, Poir. Encyc. Suppl. i. 684 = Trinia Henningii.
napiformis, Vis.—Cf. Linnaea, xxiv. (1851) 180 (= *Bunium creticum*, Urv.).—Europ. austr.
nemorosa, *Benth. & Hook. f. Gen.* i. 912.—Chili.
nervosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 684.—Reg. Himal.
nigra, Mill. Gard. Dict. ed. VIII. n. 4; Willd. Prod. Fl. Berl. 110 = P. Saxifraga.
nodiflora, Stokes, Bot. Mat. Med. ii. 149 = Apium nodiflorum.
nodosa, Urv. in Mém. Soc. Linn. Par. i. (1822) 289 = cretica.
nudicaulis, *Trautv. in Act. Hort. Petrop.* ii. (1873) 473. —Armenia.
olenea, Boiss. & Heldr. ex Boiss. Diagn. Ser. II. ii. 75 = P. Tragium.
olivieroides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 871.—Persia.
Olivieri, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 132.—Mesopot.
oreophila, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 195.—Afr. trop.
orientalis, Gouan, Illustr. 21 = magna.
ovata, Spreng. in Schult. Syst. vi. 837 (Index) = P. Saxifraga.
palmetorum, *Steud. Nom.* ed. II. ii. 335.—Arab.
Panatjan, Mirb. ex Rosenth. Pl. Diaph. 533 = P. Pruatjan.
Pansil, *Benth. & Hook. f. Gen.* i. 912.—Chili.

PIMPINELLA :—
Parishiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 309.—Burma.
parnassica, Boiss. & Heldr. ex Boiss. Diagn. Ser. II. ii. 75 = P. Tragium.
peregrina, Kotschy, ex Boiss. Diagn. Ser. I. x. 26 = affinis.
peregrina, Lej. Fl. Spa, i. 145 = magna.
peregrina, *Linn. Mant.* ii. 357.—Europ. austr.; As. Min.
petraea, Willd. ex Spreng. Sp. Umbell. 134 = P. Tragium.
peucedanifolia, *Fisch. ex Ledeb. Fl. Ross.* ii. 256.—Reg. Cauc.
podagrarioides, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 876.—As. Min.
polyclada, Boiss. & Heldr. Diagn. Ser. II. ii. 75 = P. Tragium.
polygama, Pourr. ex Nym. Consp. 308 = Trinia vulgaris.
poteriifolia, Schur, Enum. Pl. Transs. 249 = P. Saxifraga.
pratensis, Thuill. Fl. Par. ed. II. 154 = P. Saxifraga.
pretenderis, Orph. ex Boiss. Fl. Orient. ii. 872 = P. Tragium.
prolifera, *Krock. Fl. Siles. Suppl.* i. 342 (Quid?).—Europ.
Pruatjan, *Molkenb. in Miq. Pl. Jungh.* 97.—Java.
Pseudo-tragium, DC. Prod. iv. 122 = P. Tragium.
puberula, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 129.—Syria; Persia.
pubescens, Wall. Cat. n. 567 = diversifolia.
pumila, Jacq. Enum. Stirp. Vindob. 52 = Trinia vulgaris.
ramosissima, DC. Prod. iv. 122 = aurea.
ramosissima, Fisch. ex Trevir. Ind. Sem. Hort. Vratisl. (1819) 3 = Trinia Kitaibelii.
ramosissima, Hort. ex Link, Enum. Hort. Berol. i. 285 = Apium Ammi.
ranunculifolia, *Boiss. Fl. Orient.* ii. 865.—Beluchist.
Reuteriana, Boiss. Diagn. Ser. I. x. 26 = affinis.
rhodantha, *Boiss. in Tchihat. Asie Min. Bot.* i. 414.—Armenia.
rosea, Lindem. in Bull. Soc. Nat. Mosc. xxiii. (1850) II. 488 = *rubra*?
rotunda, Dulac, Fl. Hautes-Pyr. 349 = P. Saxifraga.
rotundifolia, *Bieb. Fl. Taur. Cauc.* i. 242.—Reg. Cauc.; Persia.
rotundifolia, Scop. Fl. Carn. ed. II. i. 208 = P. Saxifraga.
rubra, Hoppe & Schleich. ex Spreng. in Schult. Syst. vi. 384 = magna.
rugosa, Kunze, in Flora, xxix. (1846) 654 = magna.
rugulosa, *Schnizl. Obs. Hort. Erlang.* (1851).—Hab.?
rupestris, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) 12.—Hispan.
Sartorii, Heldr. in Boiss. Diagn. Ser. II. ii. 75 = P. Tragium.
Saxifraga, *Linn. Sp. Pl.* 163.—Europ.; As. bor.
Saxifraga, Scop. Fl. Carn. ed. II. i. 209 = magna.
Saxifraga, Ucria, Hort. Panorm. 139 = P. Tragium.
Schweinfurthii, *Aschers. in Verh. Bot. Ver. Brand.* xxi. (1880) 67.—Aegypt.
segetum, Spreng. Prod. Pl. Umb. 35 = Carum segetum.
Serra, *Franch. & Sav. Enum. Pl. Jap.* ii. 372.—Japon.
siifolia, *Leresche, in Journ. Bot.* xvii. (1879) 198.—Hispan.
sikkimensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 685.—Reg. Himal.
simensis, *Benth. & Hook. f. Gen.* i. 895.—Afr. trop.
sinica, Hance, in Journ. Bot. vi. (1868) 113 = diversifolia.
stadensis, *D. Dietr. Syn. Pl.* ii. 947.—Afr. austr.
Stocksii, *Boiss. Fl. Orient.* ii. 865.—Beluchist.
Stracheyi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 688.—Reg. Himal.
taurica, Hort. ex Steud. Nom. ed. II. ii. 336 = peregrina.
tenera, *Benth. ex Hook. f. Fl. Brit. Ind.* ii. 686.—Reg. Himal.
tenuicaulis, *Baker, in Journ. Linn. Soc.* xx. (1883) 153.—Madag.
tenuifolia, *DC. Prod.* iv. 123.—Persia.
tenuifolia, Schwaegr. & Koerte, ex Steud. Nom. ed. II. ii. 335 = magna.
tenuifolia, Spreng. in Schult. Syst. vi. 393 = P. Tragium.

PIMPINELLA :—
tenuis, Sieber, ex Spreng. in Schult. Syst. vi. 394 = cretica.
Thomasii, *Benth. & Hook. f. Gen.* i. 894, 896.—Calabr.
tomentosa, *Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 689.—Ind. or.
trachycarpum, Benth. & Hook. f. Gen. i. 895 = Aphanopleura trachysperma.
Tragium, *Vill. Prosp.* 24.—Europ. austr.; Oriens; Ind. or.
trifoliata, Wall. Cat. n. 565 = diversifolia.
triloba, *Hegetschw. Fl. Schw.* 264.—Helvet.
tripartita, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 164.—Afghan.
tripartita, Kalenicz. in Bull. Soc. Nat. Mosc. xviii. (1845) I. 231 = rotundifolia.
triradiata, Boiss. Diagn. Ser. I. x. 27 = P. Tragium.
variifolia, Salisb. Prod. 168 = P. Saxifraga.
villosa, *Schousb. Vextr. Marokk.* 139.—Reg. Mediterr. occ.
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 685.—Reg. Himal.
zagrosica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 872.—Persia.

PINALIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 31, in syn. (1825); Lindl. Orch. Scel. 21 (1826) = **Eria**, Lindl. (Orchid.).
alba, Buch.-Ham. l. c. = E. convallarioides.

PINANGA, Blume, Rumphia, ii. 76. t. 87 (1836). *PALMAE*, Benth. & Hook. f. iii. 884.
CLADOSPERMA, Griff. Notul. iii. 165 (1851).
OPHIRIA, Becc. in Ann. Jard. Buitenz. ii. (1885) 128.
bifida, Blume, Rumphia, ii. 92 = disticha.
borneensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 151.—Borneo.
caesia, *Blume, Rumphia*, ii. 84, in obs.—Ins. Celebes.
calapparia, H. Wendl. in Kerch. Palm. 253 = Actinorhytis calapparia.
celebica, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 175.—Ins. Celebes.
cochinchinensis, *Blume, Rumphia*, ii. 77, in obs.—Cochinch.
coronata, *Blume, l. c.* 83.—Ins. Celebes.
costata, Blume, l. c. 80 = Kuhlii.
Dicksonii, *Blume, l. c.* 77, in obs.—Ind. or.
disticha, *Blume, l. c.*—Malaya.
furfuracea, *Blume, l. c.* 88.—Ins. Celebes.
globulifera, *Blume, in Bull. Néerl.* (1839) 65.—Malaya.
gracilis, *Blume, Rumphia*, ii. 77, in obs.—Reg. Himal.; Burma.
hexasticha, *Kurz, For. Fl. Brit. Burma*, ii. 539.—Burma.
inaequalis, *Blume, Rumphia*, ii. 91.—Ins. Celebes.
javana, *Blume, l. c.* 85.—Java.
Junghuhnii, *Miq. Pl. Jungh.* 157.—Sumatra.
Kuhlii, *Blume, Rumphia*, ii. 82. t. 111.—Malaya.
latisecta, *Blume, l. c.* 79.—Sumatra.
maculata, *Porte, ex Lem. in Illustr. Hortic.* x. (1863) t. 361.—Ins. Philipp.
malaiana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 175.—Malaya.
minor, *Blume, Rumphia*, ii. 86.—Ins. Celebes.
Nenga, Blume, l. c. 77 = Nenga Wendlandiana.
nóxa, *Blume, l. c.* 81.—Java.
paradoxa, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 179.—Malaya.
patula, *Blume, Rumphia*, ii. 87 (sp. dub.).—Sumatra.
pisiformis, Teijsm. ex Becc. in Ann. Jard. Buitenz. ii. (1885) 136 = Gulubia costata.
pumila, Blume, Rumphia, ii. 77, in obs. = Nenga Wendlandiana?
purpurea, Hort. Bog. ex Miq. Fl. Ind. Bat. Suppl. 590 = Cyrtostachys Renda.
salicifolia, *Blume, Rumphia*, ii. 93.—Borneo.
Sanderiana, *Bull, Cat.* (1885) 15.—Malaya.
Smithii, Hort. Melb. ex Scheff. in Ann. Jard. Buitenz. i. (1876) 154 = Ptychosperma elegans.
tenella, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 179.—Borneo.
ternatensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 149.—Ins Ternate.

PINANGA :—
Veitchi, *H. Wendl.—Cf. Regel, Gartenfl.* (1880) 261.—Borneo.
vestiaria, Blume, Rumphia, ii. 77, in obs. = Areca vestiaria.

PINARDA, Vell. Fl. Flum. 23 (1825); i. t. 52 (1827) = **Micranthemum**, Michx. (Scrophul.).
repens, Vell. ll. cc. = M. orbiculatum.

PINARDIA, Cass. in Dict. Sc. Nat. xli. 38 (1826) = **Chrysanthemum**, Tourn. (Compos.).
anisocephala, Cass. l. c. = C. viscosum.
carinata, Vis. Fl. Dalm. ii. 90, in obs. = C. carinatum.
coronaria, Less. Syn. Comp. 255 = C. coronarium.
frutescens, Less. in Linnaea, vi. (1831) 168 = Phymaspermum carnosulum.
nodosa, Less. l. c. 169 = C. nodosum.
Roxburghii, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 44 = C. coronarium.
viscosa, Vis. Fl. Dalm. ii. 90, in obs. = C. viscosum.

PINARDIA, Neck. Elem. i. 5 (1790) = **Aster**, Tourn. (Compos.).

PINAROPAPPUS, Less. Syn. Comp. 143 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 529.
roseus, *Less. l. c.*—Mexic.
scapiger, *Walp. in Linnaea*, xiv. (1840) 508.—Mexic.

PINCECNITIA et PINCENECTIA, Lem. Illustr. Hortic. viii. (1861) Misc. 57 = seq.

PINCENECTITIA, Hort. ex Lem. Illustr. Hortic. viii. (1861) Misc. 48, 59 = **Nolina**, Michx. (Liliac.).
glauca, Hort. ex Lem. l. c. 61 = N. recurvata.
gracilis, Hort. ex Lem. l. c. = N. Hartwegiana.
tuberculata, Hort. ex Lem. l. c. 48, 59, 61 = N. recurvata.

PINCENICTITIA, Baker, in Journ. Linn. Soc. xviii. (1880) 234 = praec.

PINCINECTIA, Hort. ex Lem. Illustr. Hortic. viii. (1861) Misc. 57 = praec.

PINCKNEYA, Michx. Fl. Bor. Am. i. 103, t. 13 (1803). *RUBIACEAE*, Benth. & Hook. f. ii. 47.
PINKNEA, Pers. Syn. i. 197 (1805).
PINKNEYA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 81.
ionantha, *Planch. & Linden.—Cf. Bot. Zeit.* xii. (1854) 365.—N. Granat.
pubens, *Michx. Fl. Bor. Am.* i. 103. t. 13.—Am. bor.

PINEDA, Ruiz & Pav. Prod. 76. t. 14 (1794) = **Banara**, Aubl. (Samyd.).
incana, Ruiz & Pav. Syst. Veg. 133 = B. incana.

PINELIA, Lindl. Fol. Orchid. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 528.
hypolepta, *Lindl. l. c.*—Bras.

PINELLIA, Tenore, 'Sem. Hort. Nap. (1830)'; et in Att. Acc. Sc. Nap. iv. (1839) 57. *AROIDEAE*, Benth. & Hook. f. iii. 964.
ATHERURUS, Blume, Rumphia. i. 136 (1835).
HEMICARPURUS, Nees, Del. Sem. Hort. Vratisl. (1839) 4.
angustata, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 123 = tuberifera.
fornicata, Pritz. Ic. Ind. i. 846 = tubifera.
pedatisecta, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 341.—China.
tripartita, *Schott, Syn. Aroid.* i. 5.—Japon.
tuberifera, *Tenore, in Att. Acc. Sc. Nap.* iv. (1839) 57.—Japon.
Wawrae, *Engl. in DC. Monog. Phan.* ii. 567.—China.

PINGRAEA, Cass. in Dict. Sc. Nat. xli. 57 (1826) = **Baccharis**, Linn. (Compos.).
angustifolia, Cass. l. c. 58 = B. Pingraea.

PINGUICOLA, Zumag. Fl. Pedem. i. 17 (1847) = seq.

PINGUICULA, Tourn. ex Linn. Syst. ed. I (1735). *LENTIBULARIEAE*, Benth. & Hook. f. ii. 988.
BRANDONIA, Reichb. Consp. 127 (1828).
ISOLOBA, Rafin. Fl. Tellur. iv. 58 (1836).

PINGUICULA:—
acuminata, *Benth. Pl. Hartw.* 71.—Mexic.
acutifolia, Michx. Fl. Bor. Am. i. 11 = villosa.
alba, Kuehl, ex Hoppe, Taschenb. (1800) 225 = alpina.
albanica, Griseb. Spicil. Fl. Rumel. ii. 9 = hirtiflora.
albida, *Wright, ex Griseb. Cat. Pl Cub.* 162.—Cuba.
alpestris, Pers. Syn. i. 18 = alpina.
alpina, Berger. Fl. Bass. Pyr. i. 17; ex DC. Fl. Fr. Suppl. 405 = lusitanica.
alpina, *Linn. Sp. Pl.* 17.—Europ.
alpina, G. H. Weber, Spicil. Fl. Gotting. 1 = vulgaris.
antarctica, *Vahl, Enum.* i. 192.—Reg. Magellan.
australis, Chapm. Fl. S. U. St. 284 = elatior.
australis, Nutt. in Journ. Acad. Philad. vii. (1834) 103 = pumila.
Bakeriana, Hort. Sander.—Cf. Gard. Chron. (1881) i. 541. f. 102 = caudata.
borealis, Salisb. Prod. 90 = vulgaris.
brachyloba, Schult. Mant. i. 167 = alpina.
calyptrata, *H. B. & K. Nov. Gen. et Sp.* ii. 226.—N. Granat.
campanulata, Lam. in Journ. Hist. Nat. Par. i. (1792) 336. t. 18. f. 1 = lutea.
camtschatica, Willd. ex Link, Jahrb. i. III. (1820) 54 = vulgaris,
caudata, *Schlecht. in Linnaea*, vii. (1832) 393.—Mexic.
chilensis, *Clos, in C. Gay, Fl. Chil.* iv. 365.—Chili.
coerulea, Walt. Fl. Carol. 63 = elatior, pumila.
corsica, Bernard & Gren. ex Gren. & Godr. Fl. Fr. ii. 443 = leptoceras.
crenatiloba, *DC. Prod.* viii. 30.—Mexic.
crystallina, *Sibth. & Sm. Fl. Graec.* i. 11.—Ins. Cyprus.
daurica, Pall. ex Link, Jahrb. i. III. (1820) 54 = vulgaris.
davurica, Schult. Mant. i. 373 = praec.
edentula, Hook. Exot. Fl. t. 16 = lutea.
elatior, *Michx. Fl. Bor. Am.* i. 11.—Am. bor.
elongata, *Benj. in Linnaea*, xx. (1847) 318.—N. Granat.
filifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 162.—Cuba.
flavescens, Floerke, ex Hoppe, Taschenb. (1800) 14 = alpina.
floridensis, Chapm. Fl. S. U. St. Suppl. 635 = pumila.
Flos-mulionis, E. Morr. Belg. Hortic. xxii. (1872) 371. t. 27 = caudata.
gelida, Schur, Enum. Pl. Transs. 548 = alpina.
glandulosa, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 65).—Sibir.
grandiflora, Koch, Syn. Fl. Germ. ed. I. 579 = leptoceras.
grandiflora, *Lam. Encyc.* iii. 22.—Europ. occid.
grandiflora, Pollini, Fl. Veron. i. 24 = vulgaris.
Gypsophila, Wallr. in Linnaea, xiv. (1840) 533 = vulgaris.
heterophylla, *Benth. Pl. Hartw.* 70.—Mexic.
hirtiflora, *Tenore, Fl. Neap. Prod.* p. vi; *Fl. Nap.* iii. 18. t. 201.—Europ. austr.
involucrata, A. DC. in DC. Prod. viii. 30, sphalm. = *involuta*, Schrank.
involuta, *Ruiz & Pav. Fl. Per.* i. 20. t. 31. f. c.—Peruv.
involuta, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1818) 3 = villosa.
kamtschatica, Willd. ex Ledeb. Fl. Ross. iii. 3 = vulgaris.
laeta, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 80 = hirtiflora.
leptoceras, *Reichb. Pl. Crit.* i. 69.—Europ.
leptoceratia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 = praec.
lilacina, *Schlecht. & Cham. in Linnaea*, v. (1830) 94.—Mexic.
lilacina, Seem. Bot. Voy. Herald, 324 = crenatiloba.
longifolia, Gaud. Fl. Helv. i. 45 = leptoceras?
longifolia, Ram. ex DC. Fl. Fr. iii. 728 = grandiflora.
lusitanica, *Linn. Sp. Pl.* 17.—Europ. occ.
lutea, *Walt. Fl. Carol* 63.—Am. bor.
macroceras, Pall. ex Link, Jahrb. i. III. (1820) 54 = vulgaris.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 226 = caudata.
macrostyla, *Benj. in Linnaea*, xx. (1847) 317.—Peruv.
megaspilaea, Boiss. & Heldr. ex Boiss. Fl. Orient. iv. 2 = hirtiflora.
microceras, Cham. in Linnaea, vi. (1831) 568 = vulgaris.
moranensis, H. B. & K. Nov. Gen. et Sp. ii. 226 = caudata.
nana, Mart. & Gal. ex Hemsl. Biol. Centr. Am. Bot. ii. 471 = crenatiloba.
oblongiloba, A. DC. in DC. Prod. viii. 27 = caudata.
obtusiloba, A. DC. l. c. 30 = lilacina.
orchidioides, A. DC. l. c. 27 = caudata.
ovata, Stokes, Bot. Mat. Med. i. 37 = vulgaris.
oxyptera, Reichb. ex Benj. in Linnaea, xx. (1847) 318 (*occyptera*) = vulgaris?
pallida, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 288 = alpina.
pumila, *Michx. Fl. Bor. Am.* i. 11.—Am. bor.
purpurea, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 112. t. 5. f. 1 = alpina.
sibirica, *Vest, ex A. DC. in DC. Prod.* viii. 32.—Sibir.
sodalium, *F. Muell. ex Fourn. in Bull. Soc. Bot. Fr.* xx. (1873) p. lxvii.—Mexic.
spathulata, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 515.—Sibir.
subaequalis, Stokes, Bot. Comment. i. 123 = lusitanica.
vallisneriaefolia, *Webb, Otia Hisp.* 48. t. 44.—Hispan.
variegata, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 77.—Sibir.
villosa, Gunn. Fl. Norveg. ii. 71 = vulgaris.
villosa, Huds. Fl. Angl. 8 = lusitanica.
villosa, *Linn. Sp. Pl.* 17.—Reg. arct.
vulgaris, *Linn. l. c.*; ed. II. 25.—Europ.; Sibir.; Am. bor.
vulgaris, Petagn. Inst. Bot. ii. 38 = hirtiflora.

PINGUIN, [Dill.] Adans. Fam. ii. 67 (1763) = **Bromelia**, Linn.

PINILLOSIA, Ossa, in DC. Prod. v. 528 (1836).
COMPOSITAE, Benth. & Hook. f. ii. 342.
TETRACANTHUS, Wright, ex Griseb. Cat. Pl. Cub. 153 (1866).
bellioides, *Griseb. Cat. Pl. Cub.* 153.—Cuba.
tetranthoides, *DC. Prod.* v. 528.—Ind. occ.

PINKNEA, Pers. Syn. i. 197 (1805) = **Pinckneya**, Michx. (Rubiac.).
pubescens, Pers. l. c. = Pinckneya pubens.

PINKNEYA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 81 = **Pinckneya**, Michx. (Rubiac.).

PINTOA, C. Gay, Fl. Chil. i. 479, t. 16 (1846).
ZYGOPHYLLEAE, Benth. & Hook. f. i. 269.
chilensis, *C. Gay, l. c.* 480.—Chili.

PINUS, [Tourn.] Linn. Syst. ed. I (1735). *CONIFERAE*, Benth. & Hook. f. iii. 438.
APINUS, Neck. Elem. iii. 369 (1790).
CEMBRA, Opiz, Seznam, 27 (1852).
STROBUS, Opiz, in Lotos, iv. (1854) 94.
abasica, Hort. ex Carr. Conif. ed. I. 352 = halepensis.
abchasica, Hort. l. c. = halepensis.
Abies, Duroi, Obs. Bot. 39 = Abies pectinata.
Abies, Linn. Sp. Pl. 1002 = Picea excelsa.
Abies, Lour. Fl. Cochinch. 579 = Cunninghamia sinensis.
Abies, Pall. Fl. Ross. i. 6 = Picea obovata.
Abies, Siev. in Pall. N. Nord. Beitr. vii. 349 = Picea orientalis.
Abies, Thunb. Fl. Jap. 275 = Picea polita.
acapulcensis, G. Don, in Sweet, Hort. Brit. ed. III. 769 = apulcensis.
aculcensis, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
adunca, Bosc, ex Desf. Tabl. ed. II. 247; et ex Poir. Encyc. Suppl. iv. 418 = insignis.
africana, Hort. ex C. Koch, Dendrol. ii. II. 271 = P. Pinea.
alba, Ait. Hort. Kew. ed. I. iii. 371 = Picea alba.
albicaulis, *Engelm. in Trans. Acad. St. Louis*, ii. (1863) 209.—Am. bor.
Alcoquiana, Parl. in DC. Prod. xvi. II. 417 = Picea Alcockiana.
alepensis, Poir. Encyc. v. 338 = halepensis.
alopecuroides, Hort. ex Gord. Pinet. 209 = serotina.
altissima, Hort. ex Carr. Conif. ed. I. 385 = P. Laricio.
altissima, Ledeb. ex Gord. Pinet. 186 = sylvestris.
amabilis, Dougl. in Hook. Comp. Bot. Mag. ii. (1836) 93 = Abies amabilis.
amabilis, Parl. in DC. Prod. xvi. 426 = Abies lasiocarpa, magnifica.
amecaënsis, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
americana, Duroi, Obs. Bot. 39 = Tsuga canadensis.
americana, Gaertn. Fruct. ii. 60. t. 91 = Picea nigra.
americana, Steud. Nom. ed. I. 621 = Larix americana.
angulata, Roezl, Cat. Sem. Conif. Mexic. (1857) 27 = Montezumae.
Antoineana, Roezl, l. c. 14 = P. Pseudostrobus.
Apollinis, Antoine, Conif. 73 = Abies cephalonica.
apulcensis, *Lindl. Bot. Reg.* (1839) *Misc.* 63.—Mexic.
arabica, Sieber, ex Spreng. Syst. iii. 886 = halepensis.
aracanensis, Knight, ex Gord. Pinet. 179 = P. Pinea.
Araragi, Endl. Syn. Conif. 83 = Tsuga Sieboldii.
araucana, Molina, Sagg. Chile, ed. I. 182 = Araucaria imbricata.
arctica, Hort. ex Carr. Conif. ed. II. 456 = P. Pinea.
aristata, Engelm. in Am. Journ. Sc. Ser. II. xxxiv. (1862) 331; et in Trans. Acad. Sc. St. Louis, ii. (1863) 205 = Balfouriana, *S. Wats.*
arizonica, *Engelm. ex Rothr. in Wheeler, Bot. Rep. U. St. Geogr. Surv.* vi. 260.—Arizona.
Armandi, *Franch. Pl. Davidian.* i. (1884) 285.—Mongol.
armena, C. Koch, in Linnaea, xxii. (1849) 297 = sylvestris.
astecaënsis, Roezl, ex Gord. Pinet. ed. II. 290 = apulcensis.
atlantica, Endl. Syn. Conif. 137 = Cedrus Libani.
atrovirens, Roezl, ex Gord. Pinet. Suppl. 71 = Hartwegii.
Aucklandii, Lodd. ex Gord. Pinet. 195 = Gerardiana.
australis, Dum.-Cours. Bot. Cult. ed. II. vi. 460 (Quid?).—Austral.
australis, Hort. Berol. ex Steud. Nom. ed. II. ii. 237 = halepensis.
australis, Michx. f. Hist. Arbr. Am. i. 64. t. 6 = palustris.
austriaca, Link, in Flora, xi. (1828) 32; Hoess. Anleit. 6 = P. Laricio.
Ayacahuite, *Ehrenb. ex Schlecht. in Linnaea*, xii. (1838) 492.—Mexic.; Guatemal.
aztecaënsis, Roezl, Cat. Sem. Conif. Mexic. (1857) 26 = Montezumae.
baborensis, W. R. M'Nab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 697 = Abies numidica.
Backhousiana, Roezl, ex Gord. Pinet. Suppl. 72 = Montezumae.
bahamensis, *Griseb. Fl. Brit. W. Ind.* 503.—Ins. Baham.
baldensis, Zuccagni, in Roem. Collect. 158 = Abies pectinata.
Balfouriana, [*Jeffrey*,] *ex A. Murr. in Bot. Exped. Oregon.* 1, cum tab.—Calif.
Balfouriana, *S. Wats. in Bot. King's Exped.* v. 331.—Am. bor.
balsamea, Linn. Sp. Pl. 1002 = Abies balsamea.
Banksiana, *Lamb. Gen. Pin.* 21. ed. I. 7.—Am. bor. et arct.
Banksiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 218, partim = contorta.
Beardsleyi, A. Murr. in Edinb. N. Phil. Journ. N. S. i. (Jan.-Apr. 1855) 286 = ponderosa.
Benthamiana, Hartw. in Journ. Hort. Soc. ii. (1847) 189 = ponderosa.
Bessereriana, Roezl, Cat. Sem. Conif. Mexic. (1857) 20 = P. Teocote.
bicolor, Maxim. ex Parl. in DC. Prod. xvi. II. 418 = Picea Alcockiana.
bifida, Antoine, Conif. 79. t. 31. f. 2 = Abies firma.
binato-folio, Gilib. Exercit. ii. 414 = sylvestris.
Bolanderi, Parl. in DC. Prod. xvi. II. 379 = contorta.
Bonapartea, Roezl, in Gard. Chron. (1858) 358 = P. Ayacahuite.
Boothiana, Roezl, ex Gord. Pinet. Suppl. 72 = Montezumae.
borealis, Salisb. Prod. 398 = sylvestris.
Bothiana, Roezl, Cat. Conif. Mexic. (1857) 12 = *Boothiana*.
Boucheiana, Roezl, l. c. = Montezumae.
Bourcieri, Carr. in Rev. Hortic. Sér. IV. iii. (1854) 225 = seq.

PINUS:—
Boursieri, Carr. Conif. ed. I. 398 = contorta.
brachyphylla, Parl. in DC. Prod. xvi. II. 424 = Abies brachyphylla.
brachyptera, Engelm. in Wisliz. Tour N. Mexico, 89 = ponderosa.
bracteata, D. Don, in Trans. Linn. Soc. xvii. (1836) 443 = Abies bracteata.
Brunoniana, Wall. Pl. As. Rar. iii. 24. t. 247 = Tsuga Brunoniana.
bruttia, Tenore, Prod. Fl. Nap. p. lix = pyrenaica.
bullata, Roezl, Cat. Sem. Conif. Mexic. (1857) 16 = Montezumae.
Bungeana, *Zucc. in Endl. Syn. Conif.* 166.—China.
Buonapartea, Roezl, ex Gord. Pinet. 218 = P. Ayacahuite.
cairica, D. Don, ex Gord. l. c. 166 = halepensis.
calabrica, Hort. ex Gord. l. c. 168 = P. Laricio.
californica, Loisel. in Duham. Arb. ed. Nov. v. 243; Hartw. in Journ. Hort. Soc. ii. (1847) 189 = insignis.
Calocote, Roezl, ex Gord. Pinet. Suppl. 73 = P. Teocote.
canadensis, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 163 = Tsuga Mertensiana.
canadensis, Duroi, Obs. Bot. 38 = Picea alba.
canadensis, Linn. Sp. Pl. ed. II. 1421 = Tsuga canadensis.
canaliculata, *Miq. in Journ. Bot. Néerl.* i. (1861) 86.—China.
canariensis, *C. Sm. in Buch, Beschr. Canar. Ins.* 159.—Ins. Canar.
caramanica, Bosc, ex Loud. Encyc. Trees, 957 = P. Laricio.
caramaniensis, Bon Jardin. ed. (1837) 974; ex Loud. l. c. = praec.
carica, D. Don, in Fellows, Discov. Lycia, 294 = pyrenaica.
carpathica, Ungrisches Magaz. iii. 38; ex Carr. Conif. ed. II. 476 = P. Pumilio.
carpatica, Hort. ex Gord. Pinet. 180 = Picea excelsa.
Carrierei, Roezl, Cat. Sem. Conif. Mexic. (1857) 18 = Montezumae.
caucasica, Fisch. ex Gord. Pinet. 186 = sylvestris.
Cavendishiana, Hort. ex Carr. Conif. ed. II. 429 = sinensis.
Cavendishiana, Hort. ex Parl. in DC. Prod. xvi. II. 390 = P. Khasya.
cebenensis, Hort. ex Gord. Pinet. 168 = P. Laricio.
Cedrus, Linn. Sp. Pl. 1001 = Cedrus Libani.
Cedrus, Roezl, Cat. Sem. Conif. Mexic. (1857) 23 = leiophylla.
Cedrus, Uspensk. in Bull. Soc. Nat. Mosc. (1834) 389 = P. Cembra.
Cembra, *Linn. Sp. Pl.* 1000.—Europ. austr.; As. bor.
Cembra, Thunb. Fl. Jap. 274 = parviflora.
cembroides, Gord. in Journ. Hort. Soc. i. (1846) 236 = edulis.
cembroides, Newberry, in Pacif. Rail. Rep. vi. 44, 90 = albicaulis.
cembroides, *Zucc. in Abh. Akad. Muench.* i. (1832) 392.—Arizona; Mexic.
cephalonica, Endl. Cat. Hort. Vindob. i. 218 = Abies cephalonica.
chalmaënsis, Roezl, ex Gord. Pinet. Suppl. 73 = Montezumae.
Chihuahuana, *Engelm. in Wisliz. Tour N. Mexico*, 103.—Am. bor. occ.
Chilghoza, Elph. ex Knight, Syn. Conif. 30 = Gerardiana.
chinensis, Knight, ex Gord. Pinet. 176 = P. Pinaster.
Chylla, Lodd. Cat. (1836) 50 = excelsa.
cilicica, Antoine & Kotschy, in Oestr. Bot. Wochenbl. iii. (1853) 409 = Abies cilicica.
cinerea, Roehl. Deutschl. Fl. 376; ex Carr. Conif. ed. I. 246 = Picea excelsa.
clamaënsis, Carr. l. c. ed. II. 585 = Hartwegii.
Clanbrasiliana, Forbes, Hort. Woburn. 210 = Picea excelsa.
clausa, *Vasey, Cat. For. Trees*, 30; *et Sargent, Rep. For. N. Am.* (*U. St. 10th Census*, ix.) 199.—Florida.
coarctata, Roezl, Cat. Sem. Conif. Mexic. (1857) 16 = Montezumae.
coerulea, Lodd. ex Loud. Hort. Brit. 387 = Picea alba.
colchica, Booth, ex Gord. Pinet. 166 = halepensis.
Colorado, Parl. in DC. Prod. xvi. II. 407 = P. Ayacahuite.

PINUS:—
commutata, Parl. in DC. Prod. xvi. II. 417 = Picea Engelmanni.
Comonfortii, Roezl, Cat. Sem. Conif. Mexic. (1857) 22 = leiophylla.
concolor, Engelm. ex Gord. Pinet. 155 = Abies concolor.
conglomerata, Graefer, ex Tenore, Syll. Fl. Neap. 477 = pyrenaica?
contorta, Boland. in Proc. Calif. Acad. iii. (1866) 227 = muricata.
contorta, *Dougl. in Loud. Encyc. Trees*, 975. f. 915.—Am. bor. occ.
cornea, Roezl, ex Gord. Pinet. Suppl. 74 = Montezumae.
corrugata, Roezl, ex Gord. l. c. = Hartwegii.
corsicana, Loud. Arboret. Brit. iv. 2001 = P. Laricio.
Coulteri, *D. Don, in Trans. Linn. Soc.* xvii. (1837) 440.—Calif.
Craigiana, A. Murr. in Edinb. N. Phil. Journ. N. S. i. (Jan.–Apr. 1855) 288 = ponderosa.
cretacea, Kalenicz. in Bull. Soc. Nat. Mosc. (1849) I. 295 (Quid?).—Rossia.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 530.—Am. bor.; Ind. occ.
cubensis, Hort. ex Gord. Pinet. 234 = occidentalis.
cupressoides, Molini, Sagg. Chile, 168 = Araucaria imbricata?
dahurica, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) I. 101; et ex Endl. Syn. Conif. 128 = Larix dahurica.
dalmatica, Vis. Fl. Dalm. i. 199, in obs. = P. Laricio.
Damarra, Steud. Nom. ed. I. 622 = seq.
Dammara, Lamb. Pin. ed. I. i. 61. t. 38 = Agathis loranthifolia.
Davidiana, W. R. M'Nab, in Prod. Roy. Irish Acad. Ser. II. ii. (1877) 702 = Abies Davidiana.
davurica, Ledeb. Fl. Ross. iii. 673 = *dahurica*.
Decaisneana, Roezl, Cat. Sem. Conif. Mexic. (1857) 19 = Montezumae.
Decandolleana, Roezl, l. c. 22 = leiophylla.
decidua, Wall. ex Carr. Conif. ed. I. 188 = Tsuga Brunoniana.
deflexa, Torr. Bot. Mex. Bound. 209 = Jeffreyi.
densiflora, *Sieb. & Zucc. Fl. Jap.* ii. 22. t. 112.—Japon.
denticulata, Steud. Nom. ed. II. ii. 337 = Picea nigra.
Deodora, Roxb. Hort. Beng. 69; Fl. Ind. iii. 651 = Cedrus Libani.
depauperata, Roezl, ex Gord. Pinet. Suppl. 74; Carr. Conif. ed. II. 573 = Montezumae.
dependens, Roezl, Cat. Sem. Conif. Mexic. (1857) 23 = leiophylla.
detritis, Hort. Angl. ex Carr. Conif. ed. II. 467 = P. Pinaster.
devoniana, Lindl. Bot. Reg. (1839) Misc. 62 = Montezumae.
Dicksonii, Hort. ex Carr. Conif. ed. I. 300 = excelsa.
dioica, Vell. Fl. Flum. x. tt. 55, 56 = Araucaria brasiliana.
divaricata, Hort. ex Gord. Pinet. 163 = Banksiana.
Doelleriana, Roezl, Cat. Sem. Conif. Mexic. (1857) 14 = Montezumae.
Don-Pedrii, Roezl, l. c. 31 = P. Ayacahuite.
Douglasii, Lamb. Gen. Pin. ed. II. iii. t. 90 = Pseudotsuga Douglasii.
dumosa, D. Don, Prod. Fl. Nep. 55 = Tsuga Brunoniana.
Durangensis, Roezl, ex Gord. Pinet. 218 = P. Ayacahuite.
echinata, Hort. ex Carr. Conif. ed. I. 370 = P. Pumilio.
echinata, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. bor.
Edgariana, Hartw. in Journ. Hort. Soc. iii. (1848) 217 = muricata.
edulis, *Engelm. in Wisliz. Tour N. Mexico*, 88.—N. Mexic.
effusa, Salisb. Prod. 399 = Cedrus Libani.
Ehrenbergii, Endl. Syn. Conif. 151 = Hartwegii.
Ehrenbergii, Roezl, Cat. Sem. Conif. Mexic. (1857) 21 = leiophylla.
elegans, Roezl, l. c. 12 = Montezumae.
Elliottii, Engelm. in Sargent, Cat. For. Trees, 74; et in Trans. Acad. Sc. St. Louis, iv. (1880) 186 = cubensis.
Endlicheriana, Roezl, Cat. Sem. Conif. Mexic. (1857) 19 = Hartwegii.
Engelmanni, *Carr. in Rev. Hortic.* (1854) 227; *et Conif.* ed. I. 356.—Mexic.
Engelmanni, Torr. in Pacif. Rail. Rep. iv. 141 = ponderosa.

PINUS:—
ericetorum, Thore, Promen. Gascogne, 32 (Quid?).—Gallia.
erzeroomica, Calvert, ex Gord. Pinet. 186 = sylvestris.
escandoniana, Roezl, Cat. Sem. Hort. Conif. Mexic. (1857) 24 = P. Pseudo-strobus.
Escarena, Risso, Hist. Nat. Eur. ii. 340, 459 = P. Pinaster.
excelsa, Lam. Fl. Fr. ii. 202 = Picea excelsa.
excelsa, *Wall. ex Lamb. Pin.* ed. II. i. 40. t. 26; *Wall. Pl. As. Rar.* iii. 1. t. 201.—Reg. Himal.
excorticata, Hort. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 217 = Bungeana.
exserta, Roezl, Cat. Sem. Conif. Mexic. (1857) 27 = P. Pseudo-strobus.
falciformis, *Parl. in DC. Prod.* xvi. II. 685.—Borneo.
fastuosa, Salisb. Prod. 398 = P. Pinea.
Fenzleyi, Antoine & Kotschy, ex Carr. in Rev. Hortic. (1864) 259 = seq.
Fenzlii, Antoine & Kotschy, ex Carr. Conif. ed. II. 496 = P. Laricio.
fertilis, Roezl, ex Gord. Pinet. Suppl. 76 = cembroides.
filifolia, *Lindl. Bot. Reg.* (1840) *Misc.* 61.—Guatemala.
Finhonnoskiana, Parl. in DC. Prod. xvi. II. 431 = Abies homolepis.
Finlaysoniana, Wall. Cat. n. 6062 = Merkusii.
firma, Antoine, Conif. 70. t. 27/28 = Abies firma.
firma, W. R. M'Nab, in Proc. Roy. Irish Acad. 28 (1876) 686 = Abies brachyphylla.
Fischeri, Booth, ex Lodd. Cat. ed. (1836); P. Laws. Man. Conif. 333 = P. Pumilio.
flexilis, *James, in Long's Exped.* ii. 27, 35.—Am. bor. occ.
flexilis, A. Murr. Bot. Exped. Oregon, 1 = albicaulis.
Fortunei, Parl. in DC. Prod. xvi. II. 430 = Keteleeria Fortunei.
fragilis, Hort. ex Carr. Conif. ed. II. 457 = P. Pinea.
Fraseri, Lodd. Cat. (1836) 50 = rigida.
Fraseri, Pursh, Fl. Am. Sept. ii. 639 = Abies Fraseri.
Fremontiana, Endl. Syn. Conif. 183 = monophylla.
Fremontiana, Gord. in Journ. Hort. Soc. iv. (1849) 293 = edulis.
Frieseana, Wichura, in Flora, xlii. (1859) 409 = sylvestris.
frondosa, Roezl, ex Gord. Pinet. Suppl. 76 = Hartwegii.
futilis, Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 190, sphalm. = *fertilis*.
Galocote, Roezl, ex Gord. Pinet. Suppl. 76 = P. Teocote.
Geitneri, Roezl, l. c. = Hartwegii.
genevensis, Hort. ex Carr. Conif. ed. I. 372 = sylvestris.
genuensis, Cook, Sketch. Spain, ii. 236 = halepensis.
georgica, Hort. ex Gord. Pinet. 187 = palustris.
Gerardi, Forbes, Hort. Woburn. 210 = seq.
Gerardiana, *Wall. ex Lamb. Gen. Pin.* ed. II. iii. 151.—Reg. Himal.
Ghiesbrechtii, *Hort. ex Carr. Conif.* ed. II. 426.—Hab.?
glabra, *Walt. Fl. Carol.* 237.—Am. bor.
glauca, Moench, Verz. 73 = Picea alba.
glomerata, Salisb. Prod. 398 = P. Pinaster.
Gordoniana, Hartw. ex Gord. in Journ. Hort. Soc. ii. (1847) 79 = Montezumae.
gracilis, Roezl, Cat. Sem. Hort. Conif. Mexic. (1857) 21 = leiophylla.
grandis, Dougl. in Hook. Comp. Bot. Mag. ii. (1836) 147 = Abies grandis.
grandis, Roezl, Cat. Sem. Conif. Mexic. (1857) 11 = Montezumae.
Greggii, *Engelm. ex Parl. in DC. Prod.* xvi. II. 396.—Mexic.
Grenvilleae, Gord. in Journ. Hort. Soc. ii. (1847) 77 = Montezumae.
Griffithii, M'Clell. in Griff. Notul. iv. 17; Ic. Pl. As. iii. t. 365 (excl. anal.) = excelsa.
Griffithii, Parl. in DC. Prod. xvi. II. 411 = Larix Griffithii.
Grozelierii, Carr. in Rev. Hortic. (1869) 126 = monticola.
Haageana, Roezl, Cat. Sem. Conif. Mexic. (1857) 15 = Montezumae.
hagenaviensis, Hort. ex C. Koch, Dendrol. ii. II. 274 = sylvestris.
haguenoviensis, Hort. ex Parl. in DC. Prod. xvi. II. 385 = sylvestris.
halepensis, Bieb. Fl. Taur. Cauc. ii. 408 = P. Laricio.
halepensis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Reg. Mediterr.

PINUS:—
hamata, Roezl, Cat. Sem. Conif. Mexic. (1857) 32 = P. Ayacahuite.
Hamiltonii, Tenore, Cat. Ort. Nap. (1845) 90 = P. Pinaster.
Harryana, W. R. M'Nab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 689 = Abies homolepis.
Hartwegii, *Lindl. Bot. Reg.* (1839) *Misc.* 62.—Mexic.
Heldreichii, Christ, in Verh. Naturf. Ges. Basel, N. S. iii. (1863) 549 = P. Laricio.
helenica, Hort. ex C. Koch, Dendrol. ii. II. 292 = P. Pinaster.
Hendersonii, Roezl, Cat. Sem. Conif. Mexic. (1857) 15 = Montezumae.
heteromorpha, Roezl, l. c. 28 = P. Pseudo-strobus.
heterophylla, *C. Koch, in Linnaea*, xxii. (1849) 295.—Oriens.
heterophylla, Presl, Epim. Bot. 236 = parviflora.
hierosolimitana, Duham. Arbr. ii. 126 = halepensis.
hirtella, H. B. & K. Nov. Gen. et Sp. ii. 5 = Abies religiosa.
hispanica, Cook, Sketch. Spain, ii. 337 = pyrenaica.
holophylla, Parl. in DC. Prod. xvi. II. 424 = Abies firma.
homolepis, Antoine, Conif. 78. t. 31. f. 1 = Abies homolepis.
Hookeriana, W. R. M'Nab, in Proc. Irish Acad. Ser. II. ii. (1876) 212 = Tsuga Hookeriana.
horizontalis, Roezl, Cat. Sem. Conif. Mexic. (1857) 17 = Montezumae.
Hoseriana, Roezl, l. c. 24 = P. Pseudo-strobus.
hudsonica, Poir. Encyc. v. 339 = Banksiana.
Hugeli, Roezl, ex Gord. Pinet. Suppl. 77 = P. Teocote.
huisquilucaënsis, Roezl, Cat. Sem. Conif. Mexic. (1857) 22 = leiophylla.
humilis, Link, in Abh. Berl. Akad. 1827 (1830) 170 = sylvestris.
humistrata, Carr. Conif. ed. II. 388 = P. Cembra.
inflexus, Roezl, Cat. Sem. Conif. Mexic. (1857) 19 = Montezumae.
inops, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 163 = contorta.
inops, [*Soland. in*] *Ait. Hort. Kew.* ed. I. iii. 367.—Am. bor.
insignis, *Dougl. ex Loud. Arboret.* iv. 2243. f. 2132–2137.—Calif.
insularis, *Endl. Syn. Conif.* 157.—Ins. Philipp.
intermedia, Lodd. Cat. (1836); ex Parl. in DC. Prod. xvi. II. 411 = Larix leptolepis.
intermedia, Wangh. Beitr. Holz. Forst. Nord. Holz. 42. t. 16; Duroi, Baumz. ed. Pott, ii. 115 (err. typ. 114) = Larix americana.
interposita, Roezl, ex Gord. Pinet. Suppl. 78 = P. Teocote.
Iztacihuatlii, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
japonica, Forbes, Pinet. Woburn. 33 = densiflora.
japonica, Hort. ex Carr. Conif. ed. I. 365 = P. Pinaster.
japonica, Hort. ex Carr. l. c. ed. II. 457 = P. Pinea.
japonica, Parl. in DC. Prod. xvi. II. 431 = Picea japonica.
japonica, Thunb. Fl. Jap. 275 = Larix japonica.
Jeffreyana, Loud. Encyc. Pl. ii. 1544 = seq.
Jeffreyi, *A. Murr. Bot. Exped. Oreg.* 2, cum tab.; *et ex A. Murr. in Edinb. N. Phil. Journ.* xi. (1860) 224. —Calif.
jezoënsis, Antoine, Conif. 97. t. 37. f. 1 = Picea sitchensis.
Jostii, Roezl, Cat. Sem. Conif. Mexic. (1857) 15 = filifolia.
Kaempferi, Lamb. Gen. Pin. ed. III. 7 = Pseudolarix Kaempferi.
kamtschatica, Endl. Syn. Conif. 135 = Pseudolarix Kaempferi.
karamana, Mast. in Gard. Chron. (1884) I. 480 = P. Laricio.
Kegelii, Roezl, ex Gord. Pinet. Suppl. 78 = P. Teocote.
Kasya, Parl. in DC. Prod. xvi. II. 390 = P. Khasya.
Keteleerii, Roezl, Cat. Sem. Conif. Mexic. (1857) 13 = Montezumae.
khasia, Engelm. in Trans. Acad. Sc. St. Louis, iv. (1880) 179 = P. Khasya.
Khasya, *Royle, ex Gord. in Loud. Gard. Mag.* xvi. (1840) 8 (Kesuja).—Reg. Himal.; Burma.
khasyana, Griff. Notul. iv. 18 = P. Khasya.
Khutrow, Royle, Illustr. Bot. Himal. 350, 353. t. 84. f. 1 = Picea Morinda.

PINUS:—
Kochiana, Klotzsch, ex C. Koch, in Linnaea, xxii. (1849) 296 = sylvestris.
koraiensis, *Sieb. & Zucc. Fl. Jap.* ii. 28. t. 116.—Korea; Kamtchat.
Krelagi, Roezl, ex Gord. Pinet. Suppl. 78 = Hartwegii.
laeta, Salisb. Prod. 399 = Larix europaea.
Lambertiana, *Dougl. in Trans. Linn. Soc.* xv. (1827) 500.—Calif.
lanceolata, Lamb. Gen. Pin. ed. I. i. 52 = Cunninghamia sinensis.
laricina, Duroi, Obs. Bot. 49 = Larix americana.
Laricio, *Poir. Encyc.* v. 339.—Europ. austr.
Laricio, Santi, Viagg. Terz. 60. t. 1; Savi, Fl. Pisan. ii. 353 = P. Pinaster.
Larix, Dougl. in Hook. Comp. Bot. Mag. ii. (1836) 109 = Larix occidentalis.
Larix, Linn. Sp. Pl. 1001 = Larix europaea.
Larix, Siev. in Pall. N. Nord. Beitr. vii. 229 = Larix davurica.
Larix, Thunb. Fl. Jap. 275 = Larix leptolepis.
lasiocarpa, [Balf.] in A. Murr. Bot. Exped. Oregon, 1 = Abies concolor.
lasiocarpa, Hook. Fl. Bor. Am. ii. 163 = Abies subalpina.
latisquama, *Engelm. in Gard. Chron.* (1882) II. 712. —Mexic.
Latteri, Mason, in Journ. As. Soc. Beng. (1849) I. 74 = Merkusii.
Lawsoni, *Roezl, ex Gord. Pinet. Suppl.* 64.—Mexic.
laxa, Ehrh. Beitr. iii. 24 = Picea alba.
Ledebourii, Endl. Syn. Conif. 131 = Larix leptolepis.
leioclada, Stev. in Bull. Soc. Nat. Mosc. (1838) I. 44 = Abies pectinata.
leiophylla, *Schlecht. & Cham. in Linnaea*, vi. (1832) 354.—Mexic.
Lemoniana, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 512 = P. Pinaster.
leptolepis, Endl. Syn. Conif. 130 = Larix leptolepis.
Lerdoi, Roezl, Cat. Sem. Conif. Mexic. (1857) 23 = leiophylla.
Leroyi, Roezl, ex Gord. Pinet. Suppl. 79 = Montezumae.
leucodermis, Antoine, in Oestr. Bot. Zeitschr. xiv. (1864) 366 = P. Laricio.
leucosperma, *Maxim. in Bull. Acad. Pétersb.* xxvii (1881) 558.—China.
Lindleyana, Loud. ex Gord. in Journ. Hort. Soc. v. (1850) 215 = Montezumae.
Lindleyi, Carr. Man. Pl. iv. 349 = praec.
Llaveana, Schiede, in Linnaea, xii. (1838) 488 = cembroides.
Llaveana, Torr. Bot. Mex. Bound. 208. t. 55 = Parryana.
Loddigesii, Loud. Arboret. Brit. iv. 2269 = rigida.
Loiseleuriana, Carr. Conif. ed. I. 382 = pyrenaica.
longifolia, Roezl, ex Hemsl. Biol. Centr. Am. Bot. iii. 187 = Hartwegii.
longifolia, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 651.—Reg. Himal.
longifolia, Salisb. Prod. 398 = palustris.
lophosperma, Lindl. in Gard. Chron. (1860) 46 = Torreyana.
Loudoniana, Gord. Pinet. 230 = P. Ayacahuite.
Lowiana, W. R. McNab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 680. t. 46. f. 5 = Abies concolor.
Lowii, Roezl, Cat. Sem. Conif. Mexic. (1857) 18 = Hartwegii.
lucida, Salisb. Prod. 399 = Abies pectinata.
lutea, Lodd. ex Gord. Pinet. 170 = echinata.
lutea, Walt. Fl. Carol. 237 = palustris.
Lyallii, Parl. in DC. Prod. xvi. II. 412 = Larix Lyallii.
Mac-Intoshiana, Hort. ex P. Laws. Cat. (1855) 15 = contorta.
macrocarpa, Lindl. Bot. Reg. (1840) Misc. 61 = Coulteri.
macrophylla, Engelm. in Wisliz. Tour N. Mexico, 103 = Engelmanni.
macrophylla, Lindl. Bot. Reg. (1839) Misc. 63 = Montezumae.
macrophylla, Torr. in Sitgreave's Rep. 173 = ponderosa.
maderiensis, Tenore, Ind. Sem. Hort. Neap. (1854). —Cf. in Ann. Sc. Nat. Sér. IV. ii. (1854) 379 = P. Pinea.
magellensis, Schouw, in Ann. Sc. Nat. Sér. III. iii. (1845) 233 = P. Pumilio.

PINUS:—
magnifica, W. R. M'Nab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 700 = Abies magnifica.
magnifica, Roezl, Cat. Sem. Conif. Mexic. (1857) 28 = Montezumae.
mandshurica, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 382 = P. Cembra.
mariana, Duroi, Obs. Bot. 38 = Picea nigra.
maritima, R. Br. in Ait. Hort. Kew. ed. II. v. 315 = P. Laricio.
maritima, Lam. Fl. Fr. ii. 201 = P. Pinaster.
maritima, Mill. Gard. Dict. ed. VIII. n. 7 = halepensis.
marylandica, Hort. ex Antoine, Conif. 88 = Picea nigra.
Massoniana, *D. Don, in Lamb. Pin.* ed. II. ii. 16. t. 8. —China.
Massoniana, Gord. Pinet. 176 = P. Pinaster.
Massoniana, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 503 = densiflora.
Massoniana, Sieb. & Zucc. Fl. Jap. ii. 24. t. 113, 114 = Thunbergii.
Maximowiczii, Parl. in DC. Prod. xvi. II. 431 = Picea Maximowiczii.
Menziesii, Dougl. ex Lamb. Gen. Pin. ed. II. iii. 161. t. 71 = Picea sitchensis.
Merkiana, Gord. Pinet. ed. II. 448 (Index) = Merkusii.
Merkusii, *Jungh. & De Vriese, Pl. Nov. Ind. Bat. Or.* 5. t. 2.—Sumatra.
Mertensiana, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 163 = Tsuga Mertensiana.
michoacaënsis, Roezl, Cat. Sem. Conif. Mexic. (1857) 26 = Montezumae.
microcarpa, Lamb. Gen. Pin. ed. II. ii. 56. t. 37; ed. III. 87. t. 53 = Larix americana.
microcarpa, Roezl, Cat. Sem. Conif. Mexic. (1857) 21 = P. Teocote.
minor, Hort. ex Loud. Hort. Brit. Suppl. ii. 657 = halepensis.
minor, Loud. Encyc. Pl. 963. f. 1784 = P. Pinaster.
mitis, Michx. Fl. Bor. Am. ii. 204 = echinata.
monophylla, Hort. ex Parl. in DC. Prod. xvi. II. 385 = sylvestris.
monophylla, *Torr. & Frém. Rep.* 319. t. 4.—Am. bor. occ.
monspeliensis, Salzm. ex Dun. in Mém. Acad. Montp. (Sect. Sc.) ii. (1851) 82 = P. Pinaster.
monstrosa, Roezl, Cat. Sem. Conif. Mexic. (1857) 11 = Montezumae.
montana, Hoffm. Deutschl. Fl. ed. I. 340 = sylvestris.
montana, Lam. Fl. Fr. iii. 651 = P. Cembra.
montana, Mill. Gard. Dict. ed. VIII. n. 5 = P. Pumilio.
montana, Schur, Enum. Pl. Transs. 946 (Index) = Picea excelsa.
monteragensis, Godefr. ex Carr. Conif. ed. I. 353, sphalm. = seq.
montereyensis, Rauch, ex Gord. Pinet. 197 = insignis.
Montezumae, Gord. l. c. 232 = Hartwegii.
Montezumae, *Lamb. Gen. Pin.* ed. I. iii. 149. t. 64.—Mexic.
Monti-Allegri, Roezl, Cat. Sem. Conif. Mexic. (1857) 22 = leiophylla.
monticola, *Dougl. ex Lamb, l. c.* t. 87.—Am. bor. occ.
Morinda, Hort. ex Gord. Pinet. 12 = Picea Morinda.
Mugho, Poir. Encyc. v. 336 = P. Pumilio.
Mughus, Jacq. Ic. Pl. Rar. t. 193 = sylvestris.
Mughus, Scop. Fl. Carn. ed. II. ii. 247 = P. Pumilio.
Mugo, Turra, Fl. Ital. Prod. 67 = sylvestris.
Mulleriana, Roezl, Cat. Sem. Conif. Mexic. (1857) 21 = P. Teocote.
muricata, Boland. in Proc. Calif. Acad. iii. (1866) 227 = contorta.
muricata, *D. Don, in Trans. Linn. Soc.* xvii. (1837) 441.—Calif.
Murrayana, [Balf.] Bot. Exped. Oreg. 2, cum ic.; A. Murr. in Trans. Bot. Soc. Edinb. vi. (1860) 351 = contorta.
Naphta, Hort. ex Antoine, Conif. 80 = Abies Pindrow.
neglecta, Low, ex Gord. Pinet. 176 = P. Pinaster.
Neilreichiana ×, Reichardt, in Verh. Zool.-Bot. Ges. Wien (1876), 461.—Europ.
Neosa, Govan, ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 658 = Gerardiana.
nepalensis, Chambr. Trait. Arbr. Résin. Conif. 342 = excelsa.
nepalensis, Royle, ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 217 = P. Pinaster.

PINUS:—
Ne-plus-ultra, Roezl, ex Gord. Pinet. Suppl. 80 = Montezumae.
Nesselrodiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 16 = Montezumae.
nigra, Ait. Hort. Kew. ed. I. iii. 370 = Picea nigra.
nigra, Link, in Abh. Berl. Akad. 1827 (1830) 173 = P. Laricio.
nigrescens, Tenore, Fl. Nap. v. 366 = seq.
nigricans, Host, Fl. Austr. ii. 628 = P. Laricio.
nitida, Roezl, Cat. Sem. Conif. Mexic. (1857) 29 = Montezumae.
nivea, Booth, ex Carr. Conif. ed. I. 305 = P. Strobus.
nivea, Hort. ex Carr. l. c. ed. II. 401 = monticola.
nobilis, Dougl. ex Loud. Encyc. Pl. Suppl. i. 1276 = Abies nobilis.
nootkatensis, Manetti, ex Gord. Pinet. Suppl. 67 = ponderosa.
Nordmanniana, Stev. in Bull. Soc. Nat. Mosc. (1838) I. 45. t. 2 = Abies Nordmanniana.
Northumberlandiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 13 = Montezumae.
nova-hollandica, Lodd. ex Gord. Pinet. 176 = P. Pinaster.
nova-zelandica, Lodd. ex G. Don, in Loud. Hort. Brit. 387 = P. Pinaster.
Novae-Hollandiae, Lodd. ex Carr. Conif. ed. II. 466 = praec.
nummularia, Gord. Pinet. 128 = Larix leptolepis.
Nuttallii, Parl. in DC. Prod. xvi. II. 412 = Larix occidentalis.
obliqua, Saut. in Reichb. Fl. Germ. Excurs. 159 = P. Pumilio.
obovata, Antoine, Conif. 96. t. 37. f. 2 = Picea obovata.
obovata, Turcz. in Bull. Soc. Nat. Mosc. (1838) I. 101; xxvii. (1854) I. 412 = Picea orientalis.
Ocampii, Roezl, Cat. Sem. Conif. Mexic. (1857) 28 = Montezumae.
occidentalis, H. B. & K. Nov. Gen. et Sp. ii. 4 = Montezumae.
occidentalis, *Sw. Prod. Veg. Ind. Occ.* 103.—Ind. occ.
Ocote, Roezl, ex Gord. Pinet. Suppl. 81 = Montezumae.
Ocotechino, Roezl, ex Parl. in DC. Prod. xvi. II. 401 = leiophylla.
Omorika, Panč. Eine Oestl. Alp. (1876) 4; et in Gard. Chron. (1877) I. 620, 470 = Picea Omorika.
oocarpa, *Schiede, in Linnaea*, xii. (1838) 491.—Mexic.; Guatem.
oocarpoides, Lindl. ex Loud. Encyc. Trees, 1118 = praec.
orientalis, Falk, Topog. Beitr. ii. 260 = Haloxylon Ammodendron.
orientalis, Frivald. ex Carr. Conif. ed. I. 210 = Picea excelsa.
orientalis, Linn. Sp. Pl. ed. II. 1421 = Picea orientalis.
Orizabae, Gord. in Journ. Hort. Soc. i. (1846) 237 = P. Pseudo-strobus.
Ortgiesiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 25 = Montezumae.
osteosperma, Engelm. in Wisliz. Tour N. Mexico, 89 = cembroides.
Otteana, Roezl, ex Vilm. Cat. Conif. Mexiq. 18 = P. Teocote.
Padufia, Ledeb. ex Gord. Pinet. 186 = sylvestris.
Pallasiana, Lamb. Gen. Pin. ed. II. ii. t. 5 = P. Laricio.
Pallasii, Parol. Hort. Bot. Parol. (1841) 3 = pyrenaica.
palmiensis, Hort. ex Gord. Pinet. Suppl. 63 = seq.
Palmieri, Manetti, ex Gord. l. c. = palustris.
palustris, *Mill. Gard. Dict.* ed. VIII. n. 14.—Am. bor.
Papeleui, Roezl, Cat. Sem. Conif. Mexic. (1857) 31 = Hartwegii.
Paroliniana, Webb, ex Carr. Conif. ed. I. 391 = P. Laricio.
Parolinii, Vis. in Mem. Ist. Venet. vi. (1856) 243 = P. Laricio.
Parryana, *Engelm. in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 332, in nota.—Calif.
Parryana, Gord. Pinet. 202 = ponderosa.
parviflora, *Sieb. & Zucc. Fl. Jap.* ii. 27. t. 115.—Japon.
parvifolia, Hort. ex Carr. Conif. ed. II. 385 = parviflora.
Pattoniana, W. R. M'Nab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 211, 212. t. 23. f. 2 = Tsuga Mertensiana.
Pattoniana, Parl. in DC. Prod. xvi. II. 429 = Tsuga Pattoniana.

PINUS:—
patula, *Schiede & Deppe, ex Schlecht. in Linnaea*, xii. (1838) 488.—Mexic.
Pawlikowskiana, Roezl, ex Gord. Pinet. Suppl. 81 = Montezumae.
Paxtoni, Roezl, Cat. Sem. Conif. Mexic. (1857) 14 = Montezumae.
pectinata, Lam. Fl. Fr. ii. 202 = Abies pectinata.
pendula, Griff. in Journ. Trav. i. 287 = Picea Morinda.
pendula, Salisb. Prod. 399 = Larix americana.
pendula, [Soland. in] Ait. Hort. Kew. ed. I. iii. 369 = Larix americana.
Penicillus, Lapeyr. Hist. Pl. Pyr. 63 = pyrenaica.
pentaphylla, Carr. Conif. ed. II. 384 = parviflora.
persica, Strangw. in Loud. Gard. Mag. xv. (1839) 130 = halepensis.
Pescatorei, Roezl, ex Vilm. Cat. Conif. Mexiq. 18 (Quid?).—Mexic.
Peuce, *Griseb. Spicil. Fl. Rumel.* ii. 349.—Macedon.
Picea, Duroi, Obs. Bot. 37 = Picea excelsa.
Picea, Habl. Taur. 97 = P. Laricio.
Picea, Linn. Sp. Pl. 1001 = Abies pectinata.
Picea, Pall. Fl. Ross. i. I. 7 = Abies sibirica.
Pichta, Fisch. ex Lodd. Cat. (1836) 50 = Abies sibirica.
Pinaster, Bess. Fl. Galic. ii. 294 = P. Laricio.
Pinaster, Loud. Arboret. 2218 = Thunbergii.
Pinaster, [*Soland. in*] *Ait. Hort. Kew.* ed. I. iii. 367.—Reg. Mediterr.
Pinceana, *Gord. Pinet.* 204.—Mexic.
Pindrow, Royle, Illustr. Bot. Himal. 354. t. 86 = Abies Webbiana.
Pinea, Gord. Pinet. ed. I. 179, partim = densiflora.
Pinea, Habl. Taur. 97; ex Bieb. Fl. Taur. Cauc. ii. 408 = P. Laricio.
Pinea, *Linn. Sp. Pl.* 1000.—Reg. Mediterr.
Pinsapo, Boiss. Voy. Bot. Espagne, ii. 584 = Abies Pinsapo.
Pithyusa, Strangw. ex Gord. in Loud. Gard. Mag. xvi. (1840) 638 = seq.
Pityusa, Stev. in Bull. Soc. Nat. Mosc. (1838) I. 49 = halepensis.
Planchonii, Roezl, Cat. Sem. Conif. Mexic. (1857) 18 = Montezumae.
Poiretiana, Hort. ex Gord. Pinet. 168 = P. Laricio.
polita, Antoine, Conif. 95. t. 36. f. 1 = Picea polita.
ponderosa, *Dougl. in Loud. Arboret.* iv. 2243. f. 2132–2137.—Am. bor. occ.
pontica, C. Koch, in Linnaea, xxii. (1849) 297 = sylvestris.
Popocatepetlii, Roezl, Cat. Sem. Conif. Mexic. (1857) 38 = P. Ayacahuite.
porphyrocarpa, [A. Murr.] in P. Laws. Pinet. Brit. i. 83 = monticola.
prasina, Roezl, Cat. Sem. Conif. Mexic. (1857) 24 = P. Pseudo-strobus.
protuberans, Roezl, l. c. 27 = P. Pseudo-strobus.
Pseudo-halepensis, Denhardt, ex Carr. Conif. ed. I. 400 = pyrenaica.
Pseudo-larix, Steud. Nom. ed. II. ii. 338 = Larix leptolepis.
Pseudo-strobus, Gord. Pinet. 237 = Montezumae.
Pseudo-strobus, *Lindl. Bot. Reg.* (1839) Misc. 63.—Mexic.
Pseudo-taeda, *Tenore, Ind. Sem. Hort. Neap.* (1857); *ex Parl. in DC. Prod.* xvi. II. 398.—Hab.?
pumila, Hort. ex C. Koch, Dendrol. ii. II. 275 = sylvestris.
pumila, Regel, Ind. Sem. Hort. Petrop. (1858) 23; in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 211 = P. Cembra.
Pumilio, *Haenke, in Jirasek, Beob. Riesengeb.* (1791) 68.—Europ. austr.
pungens, *Lamb. ex Michx. Hist. Arb. Am.* i. 61. t. 5.—Am. bor.
pygmaea, Fisch. ex Endl. Syn. Conif. 142 = P. Cembra.
pygmaea, Hort. ex C. Koch, Dendrol. ii. II. 275 = sylvestris.
pyramidalis, Hort. ex Gord. Pinet. 168 = P. Laricio.
pyramidalis, Reum. ex Antoine, Conif. 13 = P. Pumilio.
pyramidalis, Salisb. Prod. 399 = Picea excelsa.
pyrenaica, *Lapeyr. Hist. Abr. Pl. Pyr.* 146.—Europ. austr.; Pyren.
quadrifolia, Parry, ex Parl. in DC. Prod. xvi. II. 402 = Parryana.
radiata, D. Don, in Trans. Linn. Soc. xvii. (1837) 442 = insignis.

PINUS:—
reflexa, *Engelm. in Coult. Bot. Gaz.* vii. (1882) 4.—Am. bor. occ.
Regeliana, Roezl, Cat. Sem. Conif. Mexic. (1857) 29 = P. Pseudo-strobus.
Reginae-Ameliae, C. Koch, Dendrol. ii. II. 391 (Index) = Abies Apollinis.
religiosa, H. B. & K. Nov. Gen. et Sp. ii. 5 = Abies religiosa.
resinosa, Loisel. Duham. Arb. ed. Nov. v. 237. f. 5 = pyrenaica.
resinosa, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
resinosa, Savi, Fl. Pis. ii. 354 = sylvestris.
resinosa, [*Soland. in*] *Ait. Hort. Kew.* ed. I. iii. 367.—Am. bor.
resinosa, Torr. in Ann. Lyc. N. York, ii. (1828) 249 = ponderosa.
retracta, Roezl, ex Gord. Pinet. Suppl. 82 = Montezumae.
rhaetica ×, Bruegg. in Jahrb. Ber. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 121.—Europ.
Richardiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 18 = Montezumae.
rigensis, Loud. Hort. Brit. 387, nomen = sylvestris.
rigida, Hook. & Arn. Bot. Beech. Voy. 160 = insignis.
rigida, *Mill. Gard. Dict.* ed. VIII. n. 10.—Am. bor.
rigida, Porcher, Resources S. St. 584 = echinata.
Rinzii, Roezl, Cat. Sem. Conif. Mexic. (1857) 25 = Montezumae.
robusta, Roezl, l. c. 19 = Hartwegii.
Roezlii, Carr. Conif. ed. II. 575 = Hartwegii.
Rohanii, Roezl, Cat. Sem. Conif. Mexic. (1857) 25.
romana, Hort. ex Loud. Arboret. iv. 2201 = seq.
romania, Hort. ex Steud. Nom. ed. II. ii. 338 = seq.
romeniaca, Cels, Cat. Arb. Pl. (1817) 28, nomen = P. Laricio.
rostrata, Hort. ex C. Koch, Dendrol. ii. II. 280 = P. Pumilio.
rotundata, Link, in Abh. Berl. Akad. 1827 (1830), 168 = P. Pumilio.
Royleana, Jamieson, ex Lindl. in Journ. Hort. Soc. ix. (1855) 52 = echinata.
Roylei, Lindl. ex Gord. Pinet. 170 = echinata.
rubescens, Roezl, Cat. Sem. Conif. Mexic. (1857) 17 = Montezumae.
rubra, Lamb. Pin. ed. II. ii. 43. t. 28 = Picea nigra.
rubra, Michx. f. Arb. Am. Sept. i. 45. t. 1 = resinosa.
rubra, Mill. Gard. Dict. ed. VIII. n. 3 = sylvestris.
rubra, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 166 = Massoniana.
rubra, Sieber, ex Gord. Pinet. Suppl. 58 = densiflora.
rubraeflora, Loud. ex Gord. Pinet. 172 = P. Pumilio.
rudis, Endl. Syn. Conif. 151 = Hartwegii.
rudis, Roezl, ex Gord. Pinet. Suppl. 70 = P. Pseudo-strobus.
Rumeliana, Roezl, Cat. Sem. Conif. Mexic. (1857) 12 = Montezumae.
rupestris, Michx. f. Arbr. Am. Sept. i. 49. t. 2 = Banksiana.
Russelliana, Lindl. Bot. Reg. (1839) Misc. 63 = Montezumae.
ruthenica, Hort. ex Carr. Conif. ed. II. 471 = inops.
Sabiniana, *Dougl. in Trans. Linn. Soc.* xvi. (1833) 749.—Calif.
Salzmanni, Dun. in Mém. Acad. Montp. (Sect. Sc.) ii. (1851) 82 = P. Laricio.
Sancta-helenica, Loud. ex Carr. Conif. ed. II. 466 = P. Pinaster.
sanguinea, Lapeyr. Hist. Abr. Pl. Pyr. 587 = P. Pumilio.
san-rafaeliana Roezl, Cat. Conif. Mexic. (1857) 13 = Montezumae.
saskatchawensis, Hook. ex Parl. in DC. Prod. xvi. II. 381 = contorta.
sativa, Lam. Fl. Fr. ii. 200 = Picea excelsa.
scarina, Hort. Cels, ex Steud. Nom. ed. II. ii. 338 = P. Pinaster.
scariosa, Lodd. Cat. (1836) 56; ex Loud. Encyc. Trees, 953 = sylvestris.
Schrenkiana, Antoine, Conif. 97 = Picea orientalis.
scoparia, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
scopifera, Miq. in Zoll. Syst. Verz. Ind. Archip. 82 = densiflora.
scotica, Willd. ex Endl. Syn. Conif. 172 = sylvestris.
selenolepis, Parl. in DC. Prod. xvi. II. 427 = Abies Veitchii.

PINUS:—
serenagensis, Madden, ex Gord. Pinet. 200 = longifolia.
serotina, Hort.—Cf. Bon Jard. (1837) 976; ex Antoine, Conif. 23 = palustris.
serotina, *Michx. Fl. Bor. Am.* ii. 205.—Am. bor.
Shasta, Carr. Conif. ed. II. 390 = albicaulis.
sibirica, Lodd. Cat. ex Carr. l. c. 352 = Larix dahurica.
sibirica, Turcz. in Bull. Soc. Nat. Mosq. (1838) I. 101; xxvii. (1854) I. 412 = Abies sibirica.
Sieboldii, W. R. M'Nab, in Proc. Roy. Irish Acad. Ser. II. ii. (1876) 213 = Tsuga Sieboldii.
Sinclairiana, Carr. Conif. ed. I. 355 = ponderosa.
Sinclairii, Hook. & Arn. Bot. Beech. Voy. 392 = insignis, ponderosa.
sinensis, Endl. Syn. Conif. 158, partim = P. Khasya.
sinensis, *Lamb. Gen. Pin.* iii. t. 2.—China.
sitchensis, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 164 = Picea sitchensis.
Skinneri, Hort. ex Carr. Conif. ed. I. 313 = oocarpa.
Skinnerii, Forbes, ex Gord. Pinet. 223 = filifolia.
Smithiana, D. Don, ex Lamb. Gen. Pin. iii. t. 88 = Picea Morinda.
Smithiana, Wall. Cat. n. 6063 = Picea Morinda.
Soulangeana, Roezl, Cat. Sem. Conif. Mexic. (1857) 11 = Montezumae.
spectabilis, Lamb. Gen. Pin. ed. II. ii. 3; D. Don, Prod. Fl. Nep. 55 = Abies Webbiana.
spinosa, Roezl, Cat. Sem. Conif. Mexic. (1857) 12 = Montezumae.
spinulosa, Griff. Notul. iv. 17 = Picea Morinda.
squamosa, Bosc, in Nouv. Cours Agric. ed. XI. 479; et Loud. Arboret. iv. 2292 = P. Pumilio.
squarrosa, Kalenicz. in Bull. Soc. Nat. Mosc. xxii. (1849) I. 295 (Quid?).—Rossia.
squarrosa, Walt. Fl. Carol. 237 (Quid?).—Carolin.
Stendishii, Roezl, Cat. Sem. Conif. Mexic. (1857) 30 = Hartwegii.
striata, Buch.-Ham. ex Gord. Pinet. 160 = Abies Webbiana.
strobiformis, Engelm. in Wisliz. Tour N. Mexico, 102 = P. Ayacahuite.
Strobus, Buch.-Ham. Acc. Nepal. 83 = excelsa.
Strobus, *Linn. Sp. Pl.* 1001.—Am. bor.
Strobus, Thunb. Fl. Jap. 275 = koraiensis.
subarctica, Schur, Enum. Pl. Transs. 946 (Index) = P. Pumilio.
subpatula, Roezl, ex Gord. Pinet. Suppl. 84 = patula.
sudeticus, Ungrisches Magaz. iii. 38; ex Carr. Conif. ed. II. 476 = P. Pumilio.
suffruticosa, Roezl, ex Gord. Pinet. Suppl. 84 = Hartwegii.
sumatrana, Hort. ex Mirb. in Mém. Mus. Par. xiii. (1825) 69 = Merkusii.
sylvestris, Baumg. Enum. Stirp. Transs. ii. 304 = P. Laricio.
sylvestris, Gouan, Fl. Monsp. 418 = halepensis.
sylvestris, Herb. Finlays. ex Lour. Fl. Cochinch. ed. Willd. 709 = Merkusii.
sylvestris, *Linn. Sp. Pl.* 1001.—Europ.; As. bor.
sylvestris, Lour. Fl. Cochinch. 579 = Massoniana.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 1 = P. Pinaster.
sylvestris, Thunb. Fl. Jap. 274 = Thunbergii.
syrtica, Thore, Promen. Gascogne, 161 = P. Pinaster.
tabulaeformis, Hort. ex Carr. Conif. ed. II. 510 = Thunbergii.
tabulaeformis, Hort. ex C. Koch, Dendrol. ii. II. 320 = P. Strobus.
Taeda, Blanco, Fl. Filip. ed. I. 767 = insularis.
Taeda, *Linn. Sp. Pl.* 1000.—Am. bor.
Tamrac, A. Murr. in Gard. Chron. (1869) 191 = contorta.
tatarica, Hort. ex Carr. Conif. ed. I. 389 = P. Laricio.
tatarica, Hort. ex C. Koch, Dendrol. ii. II. 289 = halepensis.
tatarica, Mill. Gard. Dict. ed. VIII. n. 4 = sylvestris.
taurica, Lodd. Cat. (1836); ex Loud. Arboret. iv. 2202 = P. Laricio.
taurina, Steud. Nom. ed. II. ii. 339, sphalm. = praec.
taxifolia, Lamb. Gen. Pin. ed. I. 27. t. 33 = Pseudotsuga Douglasii.
taxifolia, Salisb. Prod. 399 = Abies balsamea.
Tchugatskoi, Fisch. ex Henk. & Hochst. Syn. Nadelh. 157 = Abies cilicica.
tenangaënsis, Roezl, Cat. Sem. Conif. Mexic. (1857) 10 = Montezumae.

PINUS:—
tenuifolia, *Benth. Pl. Hartw.* 92.—Guatemala.
tenuifolia, Salisb. Prod. 399 = P. Strobus.
Teocote, *Cham. & Schlecht. in Linnaea*, v. (1830) 76.—Mexic.
tetragona, Moench, Meth. 364 = Picea alba.
Thibaudiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 15 = Montezumae.
Thilemannii, Roezl, l. c. 16 = Montezumae.
Thunbergii, *Parl. in DC. Prod.* xvi. II. 388.—Japon.
timoriensis, Loud. Arboret. Brit. iv. 2269 = insularis.
tinctoria, Wall. ex D. Don, Prod. Fl. Nep. 55 = Abies Webbiana.
tlamacaënsis, Roezl, ex Gord. Pinet. Suppl. 85 = Hartwegii.
tomacocaënsis, Roezl, l. c. = Montezumae.
Torreyana, *Parry, ex Torr. in Bot. Mex. Bound.* 210. t. 58.—Calif.
tortuosa, Steud. Nom. ed. II. ii. 339 = sylvestris.
Troubezkoiana, Roezl, Cat. Sem. Conif. Mexic. (1857) 11 = Montezumae.
Tschonoskiana, Parl. in DC. Prod. xvi. II. 431 = Abies homolepis.
Tschugatskoi, Hort. ex C. Koch, Dendrol. ii. II. 221 = Tchugatskoi.
Tsuga, Antoine, Conif. 83. t. 32. f. 2 = Tsuga Sieboldii.
tuberculata, D. Don, in Trans. Linn. Soc. xvii. (1836) 442 = insignis.
tuberculata, *Gord. in Journ. Hort. Soc.* iv. (1849) 218, cum tab.; *et Pinet.* 211.—Calif.
tumida, Roezl, ex Gord. Pinet. Suppl. 85 = P. Teocote.
turbinata, Bosc, ex Loud. Encyc. Trees, 975 = inops.
tzompoliana, Roezl, Cat. Sem. Conif. Mexic. (1857) 24 = P. Pseudo-strobus.
uliginosa, Neum. ex Wimm. in Uebers. Arb. Schles. Ges. 1837 (1838) 96 = P. Pumilio.
umbraculifera, Hort. ex C. Koch, Dendrol. ii. II. 320 = P. Strobus.
uncinata, Ram. ex DC. Fl. Fr. iii. 726 = P. Pumilio.
uncinata, Steud. Nom. ed. II. ii. 339 = sylvestris.
valida, Roezl, Cat. Sem. Conif. Mexic. (1857) 16 = Montezumae.
Van-Geertii, Roezl, l. c. 13 = Montezumae.
Van-Houttei, Roezl, l. c. 14 = Montezumae.
variabilis, Lamb. Gen. Pin. ed. II. i. 25. t. 16 = inops.
variabilis, Pursh, Fl. Am. Sept. ii. 643 = echinata.
variegata, Hort. ex C. Koch, Dendrol. ii. II. 290 = (Quid?).—Hab.?
Veitchii, W. R. M'Nab, in Proc. Roy. Irish Acad. (1876) 686 = Abies Veitchii.
Veitchii, Roezl, Cat. Sem. Conif. Mexic. (1857) 32 = P. Ayacahuite.
venusta, Dougl. in Hook. Comp. Bot. Mag. ii. (1836) 152 = Abies bracteata.
verrucosa, Roezl, Cat. Sem. Conif. Mexic. (1857) 22 = leiophylla.
Verschaffeltii, Roezl, l. c. 29 = Montezumae.
verticillata, Siebold, in Verh. Batav. Gen. xii. (1830) 12 = Sciadopitys verticillata.
Vilmoriniana, Roezl, Cat. Sem. Conif. Mexic. (1857) 20 = P. Teocote.
viminalis, Alstroem. in Vet. Acad. Handl. Stockh. (1777) 310 = Picea excelsa.
virescens, Neilr. Nachtr. 68 = Picea alba.
virginiana, Mill. Gard. Dict. ed. VIII. n. 9 = inops.
Webbiana, Wall. ex Lamb. Gen. Pin. ed. II. i. 77. t. 44 = Abies Webbiana.
Wilsonii, Roezl, Cat. Sem. Conif. Mexic. (1857) 19 = Montezumae.
Wincesteriana, Gord. in Journ. Hort. Soc. ii. (1847) 158 = Montezumae.
Wrightii, *Engelm. in Trans. Acad. Sc. St. Louis*, iv. (1880) 185.—Am. bor.
Zacatlanae, Roezl, Cat. Sem. Conif. Mexic. (1857) 26 = Montezumae.
zamoraënsis, Roezl, ex Gord. Pinet. Suppl. 86 = filifolia.
Zitacuaria, Gord. Pinet. 267 = seq.
Zitacuarii, Roezl, Cat. Sem. Conif. Mexic. (1857) 29 = Montezumae.

PINZONA, Mart. & Zucc. in Abh. Akad. Muench. i. (1832) 371 = **Curatella**, Linn. (Dillen.).
calineoides, Eichl. in Mart. Fl. Bras. xiii. I. 71.—Ins. Guadeloup.
coriacea, Mart. & Zucc. in Abh. Akad. Muench. i. (1837) 371 = C. americana.

PIOCTONON, Rafin. Sylva Tellur. 88 (1838) = **Heliotropium**, Tourn. (Borag.).
antillanum, Rafin. l. c.—Ind. occ.
persicum, Rafin. l. c. = H. persicum.
ternatum, Rafin. l. c. = H. fruticosum.

PIONANDRA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 353 = **Cyphomandra**, Mart. (Solan.).
allophylla, Miers, in Seem. Bot. Voy. Herald, 174 = C. allophylla.
betacea, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 358 = C. betacea.
cajanumensis, Miers, l. c. 363 = C. cajanumensis.
capsicoides, Miers, l. c. 360 = C. capsicoides.
ciliata, Miers, ex Sendt. in Mart. Fl. Bras. x. 123 = C. ciliata.
coriacea, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 363 = C. coriacea.
cornigera, Miers, Illustr. South Am. Pl. ii. 140 = C. cornigera.
crotonifolia, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 364 = Solanum crotonifolium.
cylindrica, Miers, Illustr. South Am. Pl. ii. 140 = C. cylindrica.
diploconos, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 357 = C. diploconos.
divaricata, Miers, l. c. 361 = C. divaricata.
elliptica, Miers, Illustr. South Am. Pl. ii. 140 = C. elliptica.
flagrans, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 364 = Solanum inaequale.
floribunda, Miers, l. c. 354 = C. diploconos.
fragrans, Miers, l. c. 357 = C. fragrans.
Fraxinella, Miers, Illustr. South Am. Pl. ii. 140 = C. Fraxinella.
Gardneri, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 358 = C. calycina.
Hartwegii, Miers, l. c. 363 = C. Hartwegi.
laxiflora, Miers, Illustr. South Am. Pl. ii. 140 = C. laxiflora.
narensis, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 364 = Solanum narense.
obliqua, Miers, l. c. 359 = C. obliqua.
oxyphylla, Miers, Illustr. South Am. Pl. ii. 140 = C. oxyphylla.
pendula, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 359 = C. pendula.
pinnata, Miers, in Ann. & Mag. Nat. Hist. Sér. II. xv. (1855) 199.—Bras.
premnaefolia, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 360 = C. Velloziana.
pubescens, Miers, l. c. 356 = C. sciadostylis.
Tegore, Miers, l. c. 362 = C. Tegore.
trachyphylla, Miers, l. c. 364 = Solanum trachyphyllum.
velutina, Miers, Illustr. South Am. Pl. ii. 140 = C. velutina.
viridiflora, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 359 = C. viridiflora.

PIPALIA, Stokes, Bot. Mat. Med. iv. 456 (1812) = **Litsea**, Lam. (Laurin.).
solitaria, Stokes, l. c. = L. Sebifera.

PIPAREA, Aubl. Pl. Gui. ii. App. 30. t. 386 (1775) = **Casearia**, Jacq. (Samyd.).
dentata, Aubl. l. c. 31. t. 386 = C. dentata.
multiflora, Gaertn. f. Fruct. iii. 231. t. 224. f. 1 = C. javitensis.

PIPER, Linn. Gen. ed. I. 333 (1737). *PIPERACEAE*, Benth. & Hook. f. iii. 129.
AMALAGO, Rafin. Sylva Tellur. 84 (1838).
ARTANTHE, Miq. Comm. Phyt. 40. t. 7, 8 (1840).
BETELA, Rafin. Sylva Tellur. 85 (1838).
CALLIANIRA, Miq. Syst. Pip. 344 (1843).
CARPUNYA, Presl, Epim. Bot. 228 (1850).
CARPUPICA, Rafin. Sylva Tellur. 85 (1838).
CAULOBRYON, Klotzsch, ex C. DC. in DC. Prod. xvi. I. 240 (1869).
CHAVICA, Miq. Syst. Pip. 222 (1843).
CHURUMAYA, Rafin. Sylva Tellur. 85 (1838).
COCCOBRYON, Klotzsch, in Miq. Syst. Pip. 343 (1843).
CUBEBA, Rafin. Sylva Tellur. 84 (1838).
ENCKEA, Kunth, in Linnaea, xiii. (1839) 590.
GONISTUM, Rafin. Sylva Tellur. 85 (1838).
HECKERIA, Kunth, in Linnaea, xiii. (1839) 564.
LEPIANTHES, Rafin. Sylva Tellur. 84 (1838).

PIPER:—
Macropiper, Miq. Comm. Phyt. 35 (1840).
Methysticum, Rafin. Sylva Tellur. 85 (1838).
Muldera, Miq. Comm. Phyt. 34 (1840).
Nematanthera, Miq. in Linnaea, xviii. (1844) 606. t. 11.
Ottonia, Spreng. Neue Entdeck. i. 255 (1820).
Oxodium, Rafin. Sylva Tellur. 85 (1838).
Peltobryon, Klotzsch, in Miq. Syst. Pip. 369 (1843).
Peperidia, Kostel. Allg. Med. Pharm. Fl. ii. 455 (1831).
Piperiphorum, Neck. Elem. iii. 294 (1790).
Pothomorphe, Miq. Comm. Phyt. 36 (1840).
Quebitea, Aubl. Pl. Gui. ii. 838 t. 327 (1775).
Rhyncholepis, Miq. Syst. Pip. 282 (1843).
Schilleria, Kunth, in Linnaea, xiii. (1839) 609, 726.
Schizonephos, Griff. Notul. iv. 383 (1854).
Serronia, Gaudich. in Deless. Ic. Sel. iii. 54. t. 90 (1837).
Sphaerostachys, Miq. Syst. Pip. 375 (1843).
Steffensia, Kunth, in Linnaea, xiii. (1839) 609, 726.
Suensonia, Gaudich. ex Miq. Syst. Pip. 535 (1843).
abbreviatum, Opiz, in Presl, Rel. Haenk. i. 157 = P. Chaba, *Blume*.
abutilifolium, *C. DC. in DC. Prod.* xvi. 1. 372.—Bras.
abutiloides, *Kunth, in Linnaea*, xiii. (1839) 721.—Bras.
acarouanyanum, *C. DC. in DC. Prod.* xvi. 1. 311.—Guian.
accrescens, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 118.—Ind. or.
acre, *Blume, in Verh. Batav. Gen.* xi. (1826) 185. f. 13.—Java.
aculeatum, *C. DC. in Journ. Bot.* iv. (1866) 216.—Bras.
acuminatum, Forst. f. Prod. 5 = Peperomia rhomboidea.
acuminatum, Linn. Sp. Pl. 30 = Peperomia obliqua.
acuminatum, Pers. Syn. i. 35 = Peperomia acuminata.
acutifolium, Hort. Berol. ex Steud. Nom. ed. II. ii. 339 = tuberculatum.
acutifolium, Poepp. ex Kunth, in Linnaea, xiii. (1839) 614 = mollicomum.
acutifolium, *Ruiz & Pav. Fl. Per.* 38. t. 64. f. a.—Peruv.
adenandrum, *C. DC. in DC. Prod.* xvi. 1. 273.—Guian.
adenophorum, *C. DC. l. c.* 274.—Guian.
adscendens, Endl. Prod. Fl. Norf. 36 = Peperomia Baueriana.
aduncum, *Linn. Sp. Pl.* 29.—Am. trop.
aduncum, Vell. Fl. Flum. 26; i. t. 60 = hirsutum.
aemulum, Endl. Prod. Fl. Norf. 36 = Peperomia reflexa.
aequale, *Vahl, Eclog.* i. 4. t. 3.—Am. trop.
affine, *C. DC. in DC. Prod.* xvi. 1. 286.—Guian.
affine, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 129 = Peperomia quadrifolia.
Afzelianum, Schult. Mant. i. 235 = subpeltatum.
aggregatum, Berg, in Act. Helv. vii. (1777) 181. t. 10 = Lacistema myricoides.
alare, *Ham. Prod. Pl. Ind. Occ.* 3.—Guian.
alatum, Vahl, Enum. i. 342 = Peperomia alata.
albido-punctatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 190. f. 16.—Java.
albidum, *H. B. & K. Nov. Gen. et Sp.* i. 55.—N. Granat.
albo-maculatum, D. Dietr. Syn. Pl. i. 117 = variegatum.
album, *Vahl, Enum.* i. 334.—Java.
alnoides, *Kunth, in Linnaea*, xiii. (1839) 643.—Bras.
alpinum, *C. DC. in DC. Prod.* xvi. 1. 328.—Ecuador.
alpinum, Sw. Prod. Veg. Ind. Occ. 15 = Peperomia alpina.
alsinoides, Lam. Illustr. i. 82 = Peperomia alsinoides.
alveolatum, *Opiz, in Presl, Rel. Haenk.* i. 155.—Peruv.
Amalago, *Linn. Sp. Pl.* 29.—Ind. occ.
amazonicum, *C. DC. in DC. Prod.* xvi. 1. 272.—Bras.
ambiguum, Salisb. Prod. 11 = medium.
amboinense, *C. DC. in DC. Prod.* xvi. 1. 347.—Malaya.
amplectens, *C. DC. l. c.* 293.—Bras.
amplexicaule, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia amplexicaulis.

PIPER:—
amplexifolium, Link, Enum. Hort. Berol. i. 38 = Peperomia amplexifolia.
amplifolium, *C. DC. in DC. Prod.* xvi. 1. 262.—Bras.
amplum, *Kunth, in Linnaea*, xiii. (1839) 618.—Bras.
andicola, *H. B. & K. Nov. Gen. et Sp.* i. 55.—Peruv.
androgynum, *C. DC. in Journ. Bot.* iv. (1866) 165.—N. Granat.
angulatum, Poir. Encyc. Suppl. iv. 468 = Peperomia angulata.
angustatum, Poir. l. c. 469 = Peperomia angustata.
angustifolium, Lam. Illustr. i. 81 = consanguineum.
angustifolium, Pav. ex C. DC. in DC. Prod. xvi. 1. 324 = Schlechtendaliana.
angustifolium, Roxb. Hort. Beng. 80; Fl. Ind. i. 161 = Peperomia Roxburghiana.
angustifolium, *Ruiz & Pav. Fl. Per.* i. 38.—Peruv.
angustum, Spreng. Syst. i. 108, sphalm. = augustum.
anisatum, *H. B. & K. Nov. Gen. et Sp.* i. 58.—Venezuela.
anisodorum, Blanco, Fl. Filip. ed. II. 16 = marginatum.
anisodorum, Naves, ex Villar, in Blanco, l. c. ed. III. Nov. App. 175 = P. Betle.
anisumolens, Blanco, l. c. ed. I. 23 = marginatum.
anonaefolium, *Kunth, in Linnaea*, xiii. (1839) 619.—Guian.
apiculatum, *C. DC. in DC. Prod.* xvi. 1. 273.—Bras.
appendiculatum, *C. DC. l. c.* 369.—Ecuador.
arborescens, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 438 = bantamense.
arborescens, *Miq. Syst. Pip.* 320; Wight, Ic. t. 1900, partim.—Ind. or.
arborescens, Roxb. Hort. Beng. 80; Fl. Ind. i. 159 = Thottea dependens.
arborescens, Thw. Enum. Pl. Zeyl. 293 = caninum.
arborescens, Wall. Cat. n. 6648 A = miniatum.
arboreum, Aubl. Pl. Gui. i. 23 = macrophyllum, verrucosum.
arcuatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 180. f. 11.—Java.
arcuatum, Opiz, in Presl, Rel. Haenk. i. 151 = Opizianum.
arcuatum, Thw. Enum. Pl. Zeyl. 293 = caninum, zeylanicum.
areolatum, *C. DC. in DC. Prod.* xvi. 1. 303.—Peruv.
argyrophyllum, *Miq. Syst. Pip.* 330.—Zeylan.
arifolium, Hort. ex Miq. in Mart. Fl. Bras. iv. 1. 9 = Peperomia arifolia.
aristolochioides, Lam. Illustr. i. 80 = pyrifolium.
Arnottianum, C. DC. in DC. Prod. xvi. 1. 352 = P. Chaba, *Hunter*.
aromaticum, Lam. Illustr. i. 79—nigrum.
aromaticum, Willd. Enum. Hort. Berol. Suppl. 3 = Peperomia inaequalifolia.
Artanthe, *C. DC. in DC. Prod.* xvi. 1. 271.—N. Granat.
artanthopse, *C. DC. in Mém. Soc. Phys. Genèv.* xviii. (1866) 245.—Costa Rica.
arthantopse, C. DC. in Journ. Bot. iv. (1866) 161 = praec.
articulatum, *A. Rich. Fl. Cub. Fanerog.* ii. 227.—Cuba.
articulosum, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 91; *Contrib.* 101.—Reg. Argent.
asarifolium, D. Dietr. Syn. Pl. i. 111 = Peperomia asarifolia.
asperifolium, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 105.—Guian.
asperifolium, *Ruiz & Pav. Fl. Per.* i. 37. t. 56. f. b.—Peruv.
asperiusculum, H. B. & K. Nov. Gen. et Sp. i. 51 = hirsutum.
asperum, Pers. Syn. i. 33 = scabrum.
atrosanguineum, *C. DC. in Linnaea*, xxxvii. (1871–73) 338.—Bras.
attenuatum, *Buch.-Ham. ex Wall. Cat.* n. 6642 B.—Ind. or.; Malaya.
attenuatum, Herb. ex Link, Jahrb. i. iii. (1820) 63 = laevigatum.
augustum, *Rudge, Ic. Pl. Gui.* 10. t. 7.—Guian.
aurantiacum, *Wall. Cat.* n. 6658 A.—Reg. Himal.
aurantiacum, *Wall. l. c.* 6658 B (sp. diversa).—Reg. Himal.
auriculatum, Blume, in Verh. Batav. Gen. xi. (1826) 171. t. 8 = miniatum.
auriculatum, *C. DC. in DC. Prod.* xvi. 1. 372.—Bras.

PIPER:—
auritum, *H. B. & K. Nov. Gen. et Sp.* i. 54.—Mexic.
auritum, Sieber, ex Kunth, in Linnaea, xiii. (1839) 698 = dilatatum.
australasicum, C. DC. in DC. Prod. xvi. 1. 353 = hederaceum.
australe, A. Cunn. ex Miq. in Hook. Lond. Journ. Bot. iv. (1845) 423 = Peperomia leptostachya.
austro-caledonicum, *C. DC. in DC. Prod.* xvi. 1. 346.—N. Caled.
avanum, *Wall. Cat.* n. 6661.—Burma.
avellanum, *C. DC. in DC. Prod.* xvi. 1. 302.—Guian.
avene, Hassk. Cat. Hort. Bog. Alt. 71 = Peperomia Reinwardtiana.
baccans, *C. DC. in DC. Prod.* xvi. 1. 279.—Bras.
baccatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 172. t. 3. f. 9.—Java.
baccatum, C. DC. in DC. Prod. xvi. 1. 242 = nigrum.
bahiense, *C. DC. l. c.* 371.—Bras.
Banksii, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 61.—Austral.
bantamense, *Blume, in Verh. Batav. Gen.* xi. (1826) 164. t. 3. f. 5 = caninum.
Baptisianum, *C. DC. in DC. Prod.* xvi. 1. 253.—Bras.
barbatum, *H. B. & K. Nov. Gen. et Sp.* i. 55.—Ecuador.
Barclayanum, *C. DC. in DC. Prod.* xvi. 1. 336.—Ins. Tobia.
Bartlingianum, *C. DC. l. c.* 257.—Guian.
Baryanum, *C. DC. l. c.* 272.—Bras.
basellaefolium, Poir. Encyc. Suppl. iv. 468 = Peperomia basellaefolia.
basellaefolium, Wall. Cat. n. 6665 = Peperomia Wallichii.
begoniaefolium, *Hook. & Arn. Bot. Beech. Voy.* 310.—Mexic.
Bennettianum, *C. DC. in DC. Prod.* xvi. 1. 287.—Bras.
Benthamianum, *C. DC. l. c.* 264.—Venezuela.
Berlandieri, *C. DC. l. c.* 295.—Mexic.
Berteroanum, *C. DC. l. c.* 249.—Mexic.
Betel, Blanco, Fl. Filip. ed. I. 22 = seq.
Betle, *Linn. Sp. Pl.* 28.—Ind. or.; Malaya.
bidentatum, Stokes, Bot. Mat. Med. i. 79 = P. Betle.
bilineatum, Blume, in Verh. Batav. Gen. xi. (1826) 231; Enum. Pl. Jav. fasc. i. 75 = Peperomia bilineata.
birmanicum, C. DC. in Journ. Bot. iv. (1866) 163 = Peperomia rostrata.
biunciale, Poir. Encyc. Suppl. iv. 467 = Peperomia biuncialis.
Blanchetii, *C. DC. in DC. Prod.* xvi. 1. 318.—Bras.
blandum, Jacq. Coll. iii. 210 = Peperomia blanda.
blattarum, *Spreng. Neue Entdeck.* ii. 100.—Portorico.
boehmeriaefolium, *Wall. Cat.* n. 6654 A.—Reg. Himal.; Burma.
bogotense, *C. DC. in Journ. Bot.* iv. (1866) 214.—N. Granat.
Boissierianum, *C. DC. in DC. Prod.* xvi. 1. 305.—N. Granat.
Boivinii, *C. DC. in Journ. Bot.* iv. (1866) 164.—Ins. Mascar.
bolivianum, *C. DC. in DC. Prod.* xvi. 1. 280.—Bolivia.
bombycatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 = P. Lowong.
Bontianum, *C. DC. in DC. Prod.* xvi. 1. 314.—Bras.
borbonense, C. DC. l. c. 339 = pyrifolium.
borneense, *N. E. Br. in Gard. Chron.* (1882) 1. 108.—Borneo.
botrytes, *Vahl, Enum.* i. 321.—Am. austr.
Boucheanum, *C. DC. in DC. Prod.* xvi. 1. 254.—Bras.
Bourgeaui, *C. DC. in Linnaea*, xxxvii. (1871–73) 358.—Mexic.
brachyphyllum, Willd. Enum. Hort. Berol. Suppl. 3 = Peperomia trinervis.
brachypodon, *C. DC. in DC. Prod.* xvi. 1. 327.—Ins. Gorgona.
brachystachyum, *C. DC. l. c.* 296.—Ecuador.
brachystachyum, Vahl, Enum. i. 354 = Peperomia brachystachya.
brachystachyum, Wall. Cat. n. 6656, partim = peepuloides.
brachystachyum, *Wall. l. c.* partim.—Reg. Himal.
bracteatum, Thomps. in Trans. Linn. Soc. ix. (1808) 203. t. 21. f. 2 = Peperomia scandens.
brasiliense, *C. DC. in DC. Prod.* xvi. 1. 259.—Bras.
Bredemeyeri, *Jacq. Eclog.* 125. f. 84.—Am. austr.
brevicaule, *Poir. Encyc. Suppl.* iv. 458.—Ind. occ.

PIPER:—
brevispicatum, *Opiz, in Presl, Rel. Haenk.* i. 151. t. 28. f. 1.—Hab. ?
brevistipatum, *C. DC. in Linnaea,* xxxvii. (1871–73) 338.—Bras.
Brongniartii, *C. DC. in DC. Prod.* xvi. i. 327.—Peruv.
bullatum, Vahl, Enum. i. 313 = rugosum.
bullosum, *C. DC. in Journ. Bot.* iv. (1866) 165.—N. Granat.
Burenii, *C. DC. in Linnaea,* xxxvii. (1871–73) 347.—Bras.
burmanum, C. DC. in DC. Prod. xvi. i. 356, sphalm. = seq.
buruanum, *Miq. Ann. Mus. Bot. Lngd. Bat.* i. 141.—Ins. Sunda.
cabellense, *C. DC. in DC. Prod.* xvi. i. 305.—Venezuela.
cabralanum, *C. DC. l. c.* 282.—Bras.
caladiifolium, *C. DC. l. c.* 330.—Mexic.
calceolarium, *C. DC. in Journ. Bot.* iv. (1866) 217.—N. Granat.
calceolatum, C. DC. in DC. Prod. xvi. i. 307 = calceolarium.
caldense, *C. DC. in Linnaea,* xxxvii. (1871–73) 343.—Bras.
californicum, *Presl, Rel. Haenk.* i. 160.—Calif.
callosum, Opiz, in Presl, l. c. i. 152 = P. Chaba, *Hunter.*
callosum, *Ruiz & Pav. Fl. Per.* i. 34. t. 53. f. a.—Peruv.
calocoma, *C. DC. in DC. Prod.* xvi. i. 264.—N. Granat.
Cambessedesii, *C. DC. l. c.* 271.—Bras.
canaliculatum, Klotzsch, ex Miq. Fl. Ind. Bat. i. II. 440 = miniatum.
canaliculatum, *Opiz, in Presl, Rel. Haenk.* i. 156.—Ins. Philipp.
candelarianum, *C. DC. in Linnaea,* xxxvii. (1871–73) 357.—Costa Rica.
candidum, Blume, Cat. Gew. Buitenz. 38 = Peperomia candida.
caninum, *Blume, in Verh. Batav. Gen.* xi. (1826) 214. f. 26.—Ind. or.; Malaya.
capense, *Linn. f. Suppl.* 90.—Afr. austr.
caraboboënse, *C. DC. in DC. Prod.* xvi. i. 259.—Venezuela.
caracasanum, *Bredem. ex Link, Jahrb.* i. III. (1820) 61.—Venezuela.
caraccasanum, Opiz, ex Presl, Rel. Haenk. i. 155 = controversum.
cardiophyllum, *C. DC. Prod.* xvi. i. 374.—Mexic.
caripense, *H. B. & K. Nov. Gen. et Sp.* i. 48.—Venezuela.
carpinifolium, *C. DC. in DC. Prod.* xvi. i. 255.—Bras.
Carpunya, *Ruiz & Pav. Fl. Per.* i. 37. t. 63. f. b.—Peruv.
Carpupija, Pers. Syn. i. 31 = praec.
cartagoanum, *C. DC. in Linnaea,* xxxvii. (1871–73) 350.—Costa Rica.
casapiense, *C. DC. in DC. Prod.* xvi. i. 367.—Peruv.
Casimirianum, *Hemsl. Biol. Centr. Am. Bot.* iii. 46.—Nicaragua.
cassinoides, D. Dietr. Syn. Pl. i. 111 = Peperomia alsinoides.
cassinoides, *Opiz, in Presl, Rel. Haenk.* i. 153.—Peruv.
catalpaefolium, H. B. K. Nov. Gen. et Sp. i. 58 = marginatum.
caudatum, Vahl, Eclog. Am. i. 3 = marginatum.
ceanothifolium, *H. B. & K. Nov. Gen. et Sp.* i. 56.—Venezuela.
celebicum, *Blume, in Verh. Batav. Gen.* xi. (1826) 157.—Ins. Celebes.
celtidifolium, Desf. Cat. Hort. Par. ed. III. 414 = unguiculatum.
celtidifolium, *Ham. Prod. Pl. Ind. Occ.* 3.—Ind. occ.
celtidifolium, H. B. & K. Nov. Gen. et Sp. i. 50 = aduncum.
celtidiforme, *Opiz, in Presl, Rel. Haenk.* i. 152.—Ins. Philipp.
ceramicum, *C. DC. in DC. Prod.* xvi. i. 350.—Malaya.
cernuum, Opiz, ex DC. l. c. 385 = dimidiatum.
cernuum, *Vell. Fl. Flum.* 25; i. t. 58.—Bras.
ceylanicum, C. DC. in DC. Prod. xvi. i. 242 = trineuron.
Chaba, *Blume, in Verh. Batav. Gen.* xi. (1826) 168.—Borneo.

PIPER:—
Chaba, *Hunter, in As. Res.* ix. (1809) 391.—Ind. or.; Malaya.
Chamissonis, *Steud. Nom.* ed. II. ii. 340.—Mexic.
chavicoides, *C. DC. in DC. Prod.* xvi. i. 307.—Bras.
Chawya, Buch.-Ham. in Wall. Cat. n. 6650 C = P. Betle.
chelseanum, Roem. & Schult. Syst. i. 337 = tomentosum.
chilense, *C. DC. in DC. Prod.* xvi. i. 367.—Chili.
chimonanthifolium, *Kunth, in Linnaea,* xiii. (1839) 628.—Bras.
chinantlense, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 131.—Mexic.
chinense, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 430.—China.
Churumayu, *Ruiz & Pav. Fl. Per.* i. 35. t. 58. f. b.—Peruv.
Chuvya, Miq. Syst. Pip. 267 = P. Betle.
Chuvya, *Roxb. Hort. Beng.* 4, nomen; *Hunter, ex C. DC. in DC. Prod.* xvi. i. 353 (sp. dub.).—Siam.
ciliatum, *C. DC. in Linnaea,* xxxvii. (1871–73) 347.—Bras.
ciliatum, Poir. Encyc. Suppl. iv. 469 = Peperomia ciliata.
ciliatum, Vent. ex Wawra, Bot. Ergeb. Maxim. Bras. 132 = Peperomia trinervis.
cinereum, *C. DC. in Journ. Bot.* iv. (1866) 214.—N. Granat.
circinnatum, Schult. Mant. i. 245 = Peperomia circinnata.
circulare, Desf. ex C. DC. in DC. Prod. xvi. i. 444 = Peperomia circinnata.
citrifolium, *Lam. Illustr.* i. 80.—Guian.
citrifolium, Link, Jahrb. i. III. (1820) 63 = Hoffmanseggianum.
citrifolium, Schott, ex Miq. in Mart. Fl. Bras. iv. i. 56 = colubrinum.
Claussenianum, *C. DC. in DC. Prod.* xvi. i. 275.—Bras.
clusiaefolium, Jacq. Coll. iii. 209 = Peperomia obtusifolia.
Clusii, *C. DC. in DC. Prod.* xvi. i. 340.—Afr. trop.
clypeatum, Wall. Cat. n. 6665 A et B (Fici sp.).—Ind. or.
cnernavacanum, C. DC. in Linnaea, xxxvii. (1871–72) 363, sphalm. = cuernavacanum.
coccoloboides, *Kunth, l. c.* xiii. (1839) 666.—Bras.
colipanum, *C. DC. l. c.* xxxvii. (1871–73) 349.—Mexic.
colonum, Presl, Bot. Bemerk. 112 = nigrum.
coloratum, Poir. Encyc. Suppl. iv. 467 = Peperomia colorata.
colubrinum, *Link, Jahrb.* i. III. (1820) 61.—Bras.
commutatum, Steud. Nom. ed. II. ii. 340 = plantagineum.
concavum, Vahl, Enum. i. 350 = Peperomia concava.
concinnum, *C. DC. in DC. Prod.* xvi. i. 328.—Bras.
concinnum, Haw. Rev. Pl. Succ. 198 = Peperomia concinna.
confusum, *C. DC. in Journ. Bot.* iv. (1866) 211.—Cuba.
congestum, Poir. Encyc. Suppl. iv. 469 = Peperomia congesta.
conjugatum, Poir. l. c. 466 = Peperomia conjugata.
consanguineum, *Kunth, in Linnaea,* xiii. (1839) 623.—Guian.
controversum, *Steud. Nom.* ed. II. ii. 340.—Mexic.
convexum, Blume, in Verh. Batav. Gen. xi. (1826) 236 = Peperomia convexa.
coralfalgense, *C. DC. in DC. Prod.* xvi. i. 266.—Bras.
corcovadense, *C. DC. l. c.* 255.—Bras.
cordatum, *C. DC. in Journ. Bot.* iv. (1866) 213.—Peruv.
cordifolium, Sw. Prod. Veg. Ind. Occ. 15 = Peperomia cordifolia.
cordillerianum, C. DC. in DC. Prod. xvi. i. 332 = umbellatum.
cordovanum, *C. DC. in Linnaea,* xxxvii. (1871–73) 352.—Mexic.
cordulatum, *C. DC. in Journ. Bot.* iv. (1866) 217.—Panama.
coriaceum, Vahl, Enum. i. 314 = Peperomia coriacea.
cornifolium, *H. B. & K. Nov. Gen. et Sp.* i. 52.—N. Granat.
coruscans, *H. B. & K. l. c.* 53.—N. Granat.
corylifolium, Kunth, in Linnaea, xiii. (1839) 699 = dilatatum.

PIPER:—
corylistachyon, *C. DC. in DC. Prod.* xvi. i. 346.—Ins. Philipp.
cosmianum, *C. DC. l. c.* 257.—Bras.
costaricense, *C. DC. l. c.* 328.—Costa Rica.
crassifolium, *C. DC. l. c.* 301.—Bras.
crassinervium, *H. B. & K. Nov. Gen. et Sp.* i. 48.—N. Granat.
crassipes, *Korth. ex C. DC. in DC. Prod.* xvi. i. 344.—Malaya.
crassispicatum, *Opiz, in Presl, Rel. Haenk.* i. 152.—Panama.
crassum, *Blume, in Verh. Batav. Gen.* xi. (1826) 189. f. 15.—Java.
crassum, Vell. Fl. Flum. 25; i. t. 59 = densespicatum.
crebrinerve, Willd. ex Link, Jahrb. i. III. (1820) 63 = tumidum.
crebrinode, Kunth, in Linnaea, xiii. (1839) 655 = tumidum.
crocatum, *Ruiz & Pav. Fl. Per.* i. 35. t. 55. f. a.—Peruv.
crystallinum, Vahl, Enum. i. 341 = Peperomia crystallina.
Cubeba, Boj. Hort. Maurit. 355 = pyrifolium.
Cubeba, *Linn. f. Suppl.* 90.—Malaya.
Cubeba, Vahl, Enum. i. 332 = caninum.
cubense, *C. DC. in Journ. Bot.* iv. (1866) 163.—Cuba.
cuernavacanum, *C. DC. in Linnaea,* xxxvii. (1871–73) 363 (cnernavacanum).—Mexic.
cujabanum, *Manso, ex Miq. in Mart. Fl. Bras.* iv. i. 54.—Bras.
cumanense, *H. B. & K. Nov. Gen. et Sp.* i. 47.—Venezuela.
cumaralense, *C. DC. in DC. Prod.* xvi. i. 262.—N. Granat.
cumbotianum, *C. DC. l. c.* 268.—Venezuela.
Cumingianum, *Miq. Syst. Pip.* 329.—Malaya.
cuneatum, *Hook. ex Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 429.—Ins. S. Vincent.
cuneifolium, D. Dietr. Syn. Pl. i. 119 = prunifolium.
cuneifolium, Jacq. Coll. iv. 127 = Peperomia obtusifolia.
Curumaya, D. Dietr. Syn. Pl. i. 117 = P. Churumayu.
curvatum, *Ruiz & Pav. Fl. Per.* i. 36. t. 54. f. a.—Peruv.
cuspidatum, *Ham. Prod. Pl. Ind. Occ.* 3.—Ins. S. Doming.
cyrtopodon, *C. DC. in DC. Prod.* xvi. i. 337.—Bras.
darienense, *C. DC. l. c.* 374.—Panama.
dasyoura, *C. DC. l. c.* 269.—Peruv.
dasypodum, *C. DC. l. c.* 304.—Peruv.
decipiens, *C. DC. l. c.* 273.—Mexic.
decrescens, *C. DC. l. c.* 251.—Mexic.
decumanum, Aubl. Pl. Gui. i. 21; Vahl, Enum. i. 331 = marginatum.
decumanum, *Linn. Diss. Herb. Amb. in Amoen. Acad.* v. 128.—Ind. occ.
decumanum, Mill. Gard. Dict. ed. VIII. n. 9 = Millii.
decurrens, *C. DC. in Journ. Bot.* iv. (1866) 215.—Costa Rica.
demeraranum, *C. DC. in DC. Prod.* xvi. i. 298.—Guian.
dendrophilum, *D. Dietr. Syn. Pl.* i. 108.—Mexic.
densespicatum, *Steud. Nom.* ed. II. ii. 340.—Bras.
densum, *Blume, in Verh. Batav. Gen.* xi. (1826) 193. f. 18.—Java.
denudans, *C. DC. in DC. Prod.* xvi. i. 324.—Bras.
denudatum, *Opiz, ex Presl, Rel. Haenk.* i. 158.—Ins. Philipp.
dependens, Vahl, Enum. i. 338 = Peperomia dependens.
Deppeanum, D. Dietr. Syn. Pl. i. 112 = Peperomia Deppeana.
Descourtilsianum, *C. DC. in DC. Prod.* xvi. i. 281.—Mexic.
diandrum, *C. DC. in Linnaea,* xxxvii. (1871–73) 364.—Mexic.
dichotomum, Hort. Par. ex Kunth, l. c. xiii. (1839) 594 = unguiculatum.
dichotomum, Poepp. ex Kunth, l. c. 647 = P. Pseudomollicoma.
dichotomum, Pohl, ex Miq. in Mart. Fl. Bras. iv. i. 70 = Pohlianum.
dichotomum, *Ruiz & Pav. Fl. Per.* i. 35. t. 60.—Peruv.
diffusum, Blume, ex Miq. in Linnaea, xx. (1847) 130 = sarmentosum.
diffusum, Vahl, Enum. i. 333 = attenuatum.

PIPER:—
dilatatum, *Pohl, ex Miq. in Mart. Fl. Bras.* iv. 1. 57. —Bras.
dilatatum, Reichb. ex Kunth, in Linnaea, xiii. (1839) 698 = ulmifolium.
dilatatum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 105.—Am. austr.
dimidiatum, *C. DC. in DC. Prod.* xvi. 1. 371.—Bras.
diospyrifolium, *Kunth, in Linnaea*, xiii. (1839) 627.—Bras.
discolor, Balb. ex Kunth, l. c. 592 = unguiculatum.
discolor, Schlecht. l. c. v. (1830) 74 = Schlechtendalii.
discolor, Sw. Prod. Veg. Ind. Occ. 15 = Peperomia discolor, *Loud.*
discolorum, *Desf. Cat. Hort. Par.* ed. III. 414.—Hab.?
disjunctum, *C. DC. in Linnaea*, xxxvii. (1871–73) 334. —Mexic.
dissimile, Poir. Encyc. Suppl. iv. 468 = Peperomia blanda.
distachyon, *Hill, Syst. Veg.* xxvi. 20. t. 20. f. 1.—Am. austr.
distachyon, Linn. Sp. Pl. 30 = Peperomia distachya.
divaricatum, *G. F. W. Mey. Prim. Fl. Esseq.* 15.—Guian.
divergens, *C. DC. in DC. Prod.* xvi. 1. 371.—Bras.
diversifolium, Poir. Encyc. Suppl. iv. 467 = Peperomia diversifolia.
dolabriforme, Poir. l. c. 464 = Peperomia dolabriformis.
Dombeyanum, C. DC. in DC. Prod. xvi. 1. 332 = umbellatum.
dubium, A. Dietr. Sp. Pl. i. 694 = celtidifolium.
dubium, Miq. Syst. Pip. 339 = Perrottetii.
Duchassaingii, *C. DC. in DC. Prod.* xvi. 1. 251.—Ins. Guadeloup.
dumosum, *Rudge, Pl. Gui.* 13. t. 14.—Guian.
Dupasquieri, *C. DC. in Linnaea*, xxxvii. (1871–72) 361.—Bras.
egense, *C. DC. in DC. Prod.* xvi. 1. 264.—Bras.
ellipticum, *Hook. & Arn. Bot. Beech. Voy.* 443.—Nicaragua.
ellipticum, Lam. Illustr. i. 83 = Peperomia elliptica.
ellipticum, Roem. & Schult. Syst. i. 328 = celebicum.
elongatum, *C. DC. in DC. Prod.* xvi. 1. 243.—Bras.
elongatum, Poir. Encyc. Suppl. iv. 466 = Peperomia elongata.
elongatum, Vahl, Enum. i. 312 = angustifolium.
emarginatum, Vahl, l. c. 339 = Peperomia emarginata.
emarginellum, Sw. ex Wikstr. in Vet. Akad. Handl. Stockh. (1827) 56; Adnot. Bot. 2 = Peperomia emarginella.
Enckea, *C. DC. in DC. Prod.* xvi. 1. 244.—Bras.
epigynium, *C. DC. in Linnaea*, xxxvii. (1871–73) 346. —Costa Rica.
epunctulatum, *C. DC. in DC. Prod.* xvi. 1. 338.—Bras.
eriopodon, *C. DC. l. c.* 284.—N. Granat.
estrellense, *C. DC. l. c.* 272.—Bras.
eucalyptifolium, *Rudge, Ic. Pl. Gui.* 10. t. 6.—Guian.
eucalyptiphyllum, *C. DC. in DC. Prod.* xvi. 1. (errat.) 492.—Bras.
eucalyptophyllum, C. DC. l. c. 252 = praec.
evonymifolium, Kunth, in Linnaea, xiii. (1839) 624 = lanceolatum.
exasperatum, *Vahl, Enum.* i. 332.—Ind. or.
excavatum, *Ruiz & Pav. Fl. Per.* i. 34. t. 54. f. b.—Peruv.
excelsum, *Forst. f. Prod.* 5.—Austral.; N. Zel.; Ins. Pacif.
exiguum, Blume, in Verh. Batav. Gen. xi. (1826) 232 = Peperomia exigua.
eximium, Kunth, in Linnaea, xiii. (1839) 665 = obliquum.
exsectum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 93.—Bras.
exserens, *C. DC. in DC. Prod.* xvi. 1. 261.—Bras.
extensum, Roem. & Schult. Syst. i. 325 = Peperomia elongata.
Fadyenii, *C. DC. in DC. Prod.* xvi. 1. 284.—Ind. occ.
falcatum, A. Dietr. Sp. Pl. i. 639 = Opizianum.
fallax, Vahl, Enum. i. 335 = nigrum.
fasciculare, Rudge, Ic. Pl. Gui. t. 4 = Lacistema myricoides.
fasciculatum, Ruiz & Pav. Syst. Veg. 362 = umbellatum.
Fendlerianum, *C. DC. in Journ. Bot.* iv. (1866) 213.—Venezuela.

PIPER:—
fernandezianum, Bert. ex Steud. Nom. ed. II. ii. 340 = Peperomia fernandeziana?
filiforme, Pers. Syn. i. 34 = Peperomia filiformis.
filiforme, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia obtusifolia.
Finlaysonianum, *Wall. Cat.* n. 6657.—Siam.
firmum, *C. DC. in DC. Prod.* xvi. 1. 242.—Malaya.
Fischerianum, *C. DC. l. c.* 277.—Mexic.
fistulosum, *C. DC. in Journ. Bot.* iv. (1866) 212.—N. Granat.
flagellare, H. B. & K. Nov. Gen. et Sp. i. 54 = obliquum.
flavicans, *C. DC. in DC. Prod.* xvi. 1. 293.—Bras.
flexicaule, *C. DC. l. c.* 308.—Guian.
flexuosum, *Jacq. Eclog. Am.* 139. t. 93.—Venezuela.
flexuosum, Rudge, Pl. Gui. t. 14 = flexicaule.
fluminense, *C. DC. in DC. Prod.* xvi. 1. 308.—Bras.
foetidum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 25 (Quid?).—Java.
foliiflorum, Vahl, Enum. i. 356 = Peperomia foliiflora.
foliosum, Poir. Encyc. Suppl. iv. 466 = Peperomia foliosa.
Fontanesianum, Heynh. Nom. ii. 530 = discolorum.
fontinale, Roem. & Schult. Syst. i. 335 = Peperomia fontinalis.
formosum, *C. DC. in DC. Prod.* xvi. 1. 286.—Peruv.
Forstenii, *C. DC. l. c.* 348.—Ins. Amboin.
foveolatum, C. DC. l. c. 329 = Poiteanum.
fragile, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 234.—N. Guin.
Francovilleanum, *C. DC. in Journ. Bot.* iv. (1866) 719.—Bras.
Fraseri, *C. DC. in DC. Prod.* xvi. 1. 269.—Ecuador.
freirifolium, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 274 = Peperomia exigua.
Friedrichsthalii, *C. DC. in DC. Prod.* xvi. 1. 327.—Guatemala.
frutescens, *C. DC. in Linnaea*, xxxvii. (1871–73) 337. —Bras.
fruticosum, Steud. Nom. ed. II. ii. 341 = asperifolium, *Ruiz & Pav.*
fuligineum, *Kunth, in Linnaea*, xiii. (1839) 655.—Bras.
Funckii, *C. DC. in DC. Prod.* xvi. 1. 310.—Venezuela.
furcatum, D. Dietr. Syn. Pl. i. 110 = Peperomia furcata.
fuscescens, *C. DC. in DC. Prod.* xvi. i. 291.—N. Granat.
fuscum, *C. DC. l. c.* 319.—Bras.
Futokadsura, *Sieb. Fl. Jap. Fam. Nat.* ii. 107.—Japon.
Gabrielianum, *C. DC. in DC. Prod.* xvi. 1. 268.—Guian.
galeatum, *C. DC. l. c.* 242.—Ind. or.
Galeottianum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 128 = Peperomia dendrophila.
galioides, Poir. Encyc. Suppl. iv. 470 = Peperomia galioides.
Gardneri, *C. DC. in DC. Prod.* xvi. 1. 254.—Bras.
Gaudichaudianum, *Kunth, in Linnaea*, xiii. (1839) 639. —Bras.
gemellum, Willd. ex Link, Jahrb. i. III. (1820) 62 = subpeltatum.
geniculatum, Hort. ex Link, l. c. = unguiculatum.
geniculatum, *Sw. Prod. Veg. Ind. Occ.* 15.—Ind. occ.
gigantifolium, *C. DC. in DC. Prod.* xvi. 1. 306.—Bras.
glabellum, *C. DC. l. c.* 312.—Guian.
glabellum, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia glabella.
glabratum, *Kunth, in Linnaea*, xiii. (1839) 633.—Bras.
glabrescens, *C. DC. in DC. Prod.* xvi. 1. 271.—Guian.
glabrum, *Mill. Gard. Dict.* ed. VIII. n. 12.—Honduras.
glanduligerum, *C. DC. in Journ. Bot.* iv. (1866) 215. —Venezuela.
glandulosum, Opiz, in Presl, Rel. Haenk. i. 158 = miniatum.
glaucescens, Jacq. Eclog. Am. t. 76 = unguiculatum.
globosum, *C. DC. in DC. Prod.* xvi. 1. 340.—Costa Rica.
globosum, Poir. Encyc. v. 475 = Peperomia umbilicata.
glyphicum, Hoffmgg. ex Kunth, in Linnaea, xiii. (1839) 573 = nigrum.
Godetii, *C. DC. in Linnaea*, xxxvii. (1871–72) 361.—Bras.
gracile, *Ruiz & Pav. Fl. Per.* i. 35. t. 59. f. b.—Peruv.

PIPER:—
granatense, *D. Dietr. Syn. Pl.* i. 117.—N. Granat.
grande, *Vahl, Eclog. Am.* ii. 3. t. 11.—N. Granat.
grandifolium, Afzel. Remed. Guineen. vi. 39 = subpeltatum.
grandifolium, *H. B. & K. Nov. Gen. et Sp.* i. 46.—N. Granat.
granulosum, Ruiz & Pav. Fl. Per. i. 38. t. 64. f. b = angustifolium.
Griffithii, *C. DC. in Journ. Bot.* iv. (1866) 163.—Reg. Himal.
griseum, *C. DC. in DC. Prod.* xvi. 1. 281.—Peruv.
guahamense, *C. DC. l. c.* 336.—Ins. Marian.
guanacostense, *C. DC. in Linnaea*, xxxvii. (1871–73) 356.—Nicaragua.
guianense, *C. DC. in DC. Prod.* xvi. 1. 301.—Guian.
Guigual, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 20 = brachystachyum.
Guildingianum, Spreng. Syst. iv. Cur. Post. 20 = Peperomia scandens.
guineense, *Schum. & Thonn. Beskr. Guin. Pl.* 19.—Afr. trop.
Haenkeanum, D. Dietr. Syn. Pl. i. 114 = Peperomia Haenkeana.
Haenkeanum, *Opiz, in Presl, Rel. Haenk.* i. 159.—Ins. Philipp.
Hahnii, *C. DC. in Linnaea*, xxxvii. (1871–72) 354.—Ins. Martinic.
halesiaefolium, *Kunth, ex C. DC. in DC. Prod.* xvi. 1. 309.—Bras.
Hamiltonii, *C. DC. l. c.* 360.—Reg. Himal.
Hapnium, *Buch.-Ham. in Wall. Cat.* n. 6650 D.—Ind. or.
Hartwegianum, *C. DC. in DC. Prod.* xvi. 1. 369.—N. Granat.
Haworthianum, Schult. Mant. i. 245 = Peperomia Haworthiana.
Hayneanum, *C. DC. in DC. Prod.* xvi. 1. 253.—Ins. Martinic.
hederaceum, *A. Cunn. ex Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 435.—Austral.
hemionitidifolium, D. Dietr. Syn. Pl. i. 114 = Peperomia hemionitidifolia.
Henshenii, *C. DC. in Linnaea*, xxxvii. (1871–73) 336. —Bras.
heptandrum, *C. DC. in DC. Prod.* xvi. 1. 244.—Peruv.
herbaceum, Miq. Syst. Pip. 96 = Peperomia obliqua.
hernandiaefolium, Vahl, Enum. i. 344 = Peperomia hernandiaefolia.
heteromallum, Pers. Syn. i. 33 = Peperomia secunda.
heterophyllum, *Ruiz & Pav. Fl. Per.* i. 34. t. 56. f. a. —Peruv.
heterostachyum, D. Dietr. Syn. Pl. i. 111 = Peperomia heterostachya.
Hilarianum, *Steud. Nom.* ed. II. ii. 341.—Bras.
Hillianum, *C. DC. in DC. Prod.* xvi. 1. 290.—Bras.
hirsutissimum, D. Dietr. Syn. Pl. i. 122 = Haenkeanum.
hirsutum, *Sw. Fl. Ind. Occ.* i. 60.—Am. trop.
hirtellum, *C. DC. in Journ. Bot.* iv. (1866) 218.—Bras.
hirtulum, *C. DC. in DC. Prod.* xvi. 1. 371.—Bras.
hispidulum, Sw. Prod. Veg. Ind. Occ. 15 = Peperomia hispidula.
hispidum, *H. B. & K. Nov. Gen. et Sp.* i. 50.—N. Granat.
hispidum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 131 = mexicanum.
hispidum, Sw. Prod. Veg. Ind. Occ. 15 = hirsutum.
Hoffmannseggianum, *Schult. Mant.* i. 242.—Bras.
Hohenackeri, *C. DC. in DC. Prod.* xvi. 1. 243.—Guian.
Holtonii, *C. DC. l. c.* 300.—N. Granat.
hongkongense, *C. DC. l. c.* 347.—Ins. Hongkong.
Hookeri, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 437.—Ind. or.
Hookerianum, *C. DC. in DC. Prod.* xvi. 1. 292.—Peruv.
Hostmannianum, *C. DC. l. c.* 287.—Am. trop.
Huegelianum, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 58.—Austral.
humile, Mill. Gard. Dict. ed. VIII. n. 4 = Milleri.
humile, Vahl, Enum. i. 349 = Peperomia humilis.
hyalinum, Wall. Cat. n. 6662 = Peperomia exigua.
hydrangeifolium, Kunth, in Linnaea, xiii. (1839) 659 = crassinervium.
hydrocotylefolium, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 129 = Peperomia umbilicata.

PIPER :—

hymenophyllum, *Miq. in Wight, Ic.* t. 1942.—Ind. or.
hypoglaucum, *Miq. in Ann. Bot. Mus. Lugd. Bat.* i. 139.—Sumatra.
imperiale, *C. DC. in DC. Prod.* xvi. 1. 339.—Darien.
impunctatum, Link, Jahrb. i. III. (1820) 61 = colubrinum.
inaequale, *C. DC. in Journ. Bot.* iv. (1866) 215.—Guian.
inaequalifolium, Vahl, Enum. i. 355 = Peperomia inaequalifolia.
incanum, Haw. Suppl. Pl. Succ. 2; Link & Otto, Ic. Pl. Sel. Rar. 17. t. 7 = Peperomia incana.
incisum, Rich. ex C. DC. in DC. Prod. xvi. 1. 455 = Peperomia quadrifolia.
incurvum, *Sieber, ex C. DC. l. c.* 294.—Ins. Martinic.
indecorum, *Kunth, in Linnaea*, xiii. (1839) 722.—Peruv.
indicum, C. DC. in DC. Prod. xvi. 1. 362 = leptostachyum, *Wall.*
insectifugum, *C. DC. ex Seem. Fl. Vit.* 262.—Ins. Fiji.
insigne, *Kunth, in Linnaea*, xiii. (1839) 667.—Guian.
insulare, C. DC. in DC. Prod. xvi. 1. 362 = trineuron.
interruptum, *Opiz, in Presl, Rel. Haenk.* i. 157.—Ins. Philipp.
inversum, *C. DC. in Linnaea*, xxxvii. (1871–73) 359.—Bras.
irazuanum, *C. DC. l. c.* 340.—Costa Rica.
Jaborandi, *Vell. Fl. Flum.* 24; i. t. 55.—Bras.
Jacquemontianum, Kunth, in Linnaea, xiii. (1839) 631 = citrifolium.
Jacquini, *C. DC. in DC. Prod.* xvi. 1. 280.—Venezuela?
Jacquinianum, *Schult. Mant.* i. 234.—Venezuela?
jalapense, *C. DC. in DC. Prod.* xvi. 1. 277.—Mexic.
jalapense, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 131 = lapathifolium.
janeiroënse, *C. DC. in Linnaea*, xxxvii. (1871–73) 339.—Bras.
japurense, *C. DC. in DC. Prod.* xvi. 1. 263.—Bras.
javanicum, C. DC. l. c. 343 = caninum.
javitense, *H. B. & K. Nov. Gen. et Sp.* i. 54.—Venezuela.
Jayeri, *C. DC. in DC. Prod.* xvi. 1. 358.—Ins. Philipp.
julifiorum, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 9.—Bras.
Jussiaeanum, *Kunth, in Linnaea*, xiii. (1839) 680.—Bras.
Kappleri, *C. DC. in Journ. Bot.* iv. (1866) 213.—Guian.
Karok, Blume, Cat. Gew. Buitenz. 33 = attenuatum.
Karwinskianum, *Kunth, in Linnaea*, xiii. (1839) 700.—Mexic.
Kegelianum, *C. DC. in DC. Prod.* xvi. 1. 372.—Bras.
Kegelii, *C. DC. l. c.* 323.—Guian.
kewense, *C. DC. l. c.* 327.—Hab.?
khasianum, *C. DC. l. c.* 349.—Reg. Himal.
Klotzschianum, *C. DC. l. c.* 257.—Bras.
Korthalsii, *Miq. Ann. Bot. Mus. Lugd. Bat.* i. 139.—Malaya.
Kunthianum, *Steud. Nom.* ed. II. ii. 341.—Bras.
Kunthii, *C. DC. in DC. Prod.* xvi. 1. 250.—Mexic.
Labillardieri, *C. DC. l. c.* 361.—Ins. Amboin.
lacunosum, *H. B. & K. Nov. Gen. et Sp.* i. 51.—Am. austr.
laetum, *C. DC. in DC. Prod.* xvi. 1. 256.—Bras.
laeve, *Vahl, Enum.* i. 332.—Ind. or.
laevifolium, Blume, Cat. Gew. Buitenz. 33 = Peperomia laevifolia.
laevigatum, *H. B. & K. Nov. Gen. et Sp.* i. 56.—N. Granat.
lagoaënse, *C. DC. in Linnaea*, xxxvii. (1871–73) 346.—Bras.
lanatum, Roxb. Hort. Beng. 80; Fl. Ind. i. 159 = caninum.
lanatum, *Wight, ex Miq. in Hook. Lond. Journ. Bot.* v. (1846) 553.—Ind. or.
lancaefolium, Schott, ex C. DC. in DC. Prod. xvi. 1. 324 = denudans.
lanceaefolium, *H. B. & K. Nov. Gen. et Sp.* i. 49.—Bras.
lanceatum, A. Dietr. Sp. Pl. i. 636 = lancifolium.
lanceolatum, Roxb. Hort. Beng. 80; Fl. Ind. i. 159 = miniatum.
lanceolatum, *Ruiz & Pav. Fl. Per.* i. 36.—Peruv.
lanceolatum, Salzm. ex Miq. in Mart. Fl. Bras. iv. 1. 47 = hirsutum.

PIPER :—

lancifolium, *Ham. Prod. Pl. Ind. Occ.* 3.—Jamaic.
Langsdorffianum, *C. DC. in DC. Prod.* xvi. 1. 290.—Bras.
Langsdorfii, *Schrank, in Flora*, vii. (1824) II. *Beibl.* 10.—Bras.
lapathifolium, *Steud. Nom.* ed. II. ii. 341.—Mexic.
latifolium, *Forst. f. Prod.* 5.—Ins. Pacif.
latifolium, Hunter, in As. Res. ix. (1809) 390 = sarmentosum.
latifolium, Lam. Illustr. i. 81 = subpeltatum.
latum, *H. B. & K. Nov. Gen. et Sp.* i. 57.—N. Granat.
laurifolium, H. B. & K. l. c. 47 = laurinum.
laurifolium, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.
laurinum, *Roem. & Schult. Syst.* i. 291.—Peruv.
laxiflorum, Poir. Encyc. Suppl. iv. 466 = Peperomia laxiflora.
laxum, *Vahl, Enum.* i. 326 —Ins. Philipp.
Lechlerianum, *C. DC. in DC. Prod.* xvi. 1. 269.—Peruv.
Ledebourii, *C. DC. l. c.* 281.—Bras.
Lehmannianum, *C. DC. l. c.* 368.—Peruv.
leiophyllum, *C. DC. l. c.* 297.—Bras.
Lenormandianum, *C. DC. l. c.* 299.—Guian.
lentaginoides, *Kunth, in Linnaea*, xiii. (1839) 683.—Bras.
lenticellosum, *C. DC. in Journ. Bot.* iv. (1866) 167.—N. Granat.
leonense, *C. DC. l. c.* 163.—Afr. trop.
lepidotum, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 45; *Contrib.* 97.—Reg. Argent.
leptophyllum, *C. DC. in DC. Prod.* xvi. 1. 279.—Venezuela.
leptostachyon, *Nutt. in Am. Journ. Sc.* Ser. I. v. (1822) 287.—Florida.
leptostachyum, *Wall. Cat.* n. 6649.—Burma.
leptostachyum, Willd. ex Kunth, in Linnaea, xiii. (1839) 604; A. Rich. Fl. Cub. Fanerog. ii. 228 = tenue.
lepturum, *Kunth, l. c.* 679.—Bras.
Lessertianum, *C. DC. in Journ. Bot.* iv. (1866) 164.—Ins. Philipp.
leucophyllum, C. DC. in DC. Prod. xvi. 1. 278 = verbascifolium.
Lhotzkyanum, *Kunth, in Linnaea*, xiii. (1839) 657.—Bras.
Liebmannii, *C. DC. l. c.* xxxvii. (1871–73) 344.—Mexic.
Lindbergii, *C. DC. in DC. Prod.* xvi. 1. 297.—Bras.
Lindenianum, *C. DC. in Journ. Bot.* iv. (1866) 218.—Cuba.
Lindenii, *C. DC. in DC. Prod.* xvi. 1. 248.—Mexic.
linearifolium, *C. DC. in Linnaea*, xxxvii. (1871–73) 355.—Costa Rica.
lineatum, *Ruiz & Pav. Fl. Per.* i. 35. t. 60. f. a.—Peruv.
linguaefolium, *Ham. Prod. Pl. Ind. Occ.* 3.—Jamaic.
Linkii, *A. Dietr. Sp. Pl.* i. 653.—Am. austr.
Linkii, *Miq. Syst. Pip.* 317.—Java.
Lonchites, *Schult. Mant.* i. 241.—Malaya.
Lonchites, Wall. Cat. n. 6644 A, B. = caninum, miniatum.
longepedunculatum, D. Dietr. Syn. Pl. i. 110 = Peperomia longepedunculata.
longicuspe, Willd. ex Kunth, in Linnaea, xiii. (1839) 629 = citrifolium.
longiflorum, Willd. ex D. Dietr. Syn. Pl. i. 112 = Peperomia longiflora.
longifolium, *Ruiz & Pav. Fl. Per.* i. 37. t. 61. f. a.—Peruv.
longum, Blume, in Verh. Batav. Gen. xi. (1826) 197 = P. Chaba, *Hunter.*
longum, *Linn. Sp. Pl.* 29.—As. trop.
Lowong, *Blume, in Verh. Batav. Gen.* xi. (1826) 161.—Java.
loxense, Poir. Encyc. Suppl. iv. 470 = Peperomia loxensis.
Lucaeanum, *Kunth, in Linnaea*, xiii. (1839) 678.—Bras.
lunare, D. Dietr. Syn. Pl. i. 111 = Peperomia lunaria.
Lundii, *C. DC. in Linnaea*, xxxvii. (1871–73) 339.—Bras.
Luschnathianum, *Kunth, l. c.* xiii. (1839) 617.—Bras.
luzonense, *C. DC. in DC. Prod.* xvi. 1. 350.—Ins. Philipp.
Macgillivrayi, *C. DC. ex Seem. Fl. Vit.* 262. t. 75.—Ins. Fiji.
machadoënse, *C. DC. in DC. Prod.* xvi. 1. 254.—Bras.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* i. 46.—Mexic.

PIPER :—

macrophyllum, Loud. Encyc. Pl. 28 = Rohrii.
macrophyllum, Schlecht. & Cham. in Linnaea, v. (1830) 73 = lapathifolium.
macrophyllum, Sw. Prod. Veg. Ind. Occ. 15 = geniculatum.
macrorhizum, Poir. Encyc. Suppl. iv. 471 = Peperomia macrorhiza.
macrorhynchon, *C. DC. in DC. Prod.* xvi. 1. 263.—Bras.
macrostachyon, Vahl, Enum. i. 341 = Peperomia macrostachya.
macrostachyum, *C. DC. in DC. Prod.* xvi. 1. 371.—Bras.
macrourum, H. B. & K. Nov. Gen. et Sp. i. 54 = tuberculatum.
maculatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 199. f. 20.—Java.
maculosum, *Hill, Veg. Syst.* xxvi. 19. t. 19. f. 3.—Am. austr.
maculosum, Linn. Sp. Pl. 30 = Peperomia maculosa.
madagascariense, *C. DC. in DC. Prod.* xvi. 1. 379.—Madag.
magicum, Reinw. ex Miq. in Linnaea, xxi. (1848) 484 = majusculum.
magnoliaefolium, Haw. Syn. Pl. Succ. 6; Miq. in Mart. Fl. Bras. iv. 1. 24 = Peperomia obtusifolia.
magnoliaefolium, Jacq. Coll. iii. 210 = Peperomia magnoliaefolia.
majusculum, *Blume, in Verh. Batav. Gen.* xi. (1826) 210.—Malaya.
malabarense, C. DC. in DC. Prod. xvi. 1. 242 = nigrum.
malacophyllum, *C. DC. l. c.* 337.—Bras.
Malamiri, *Blume, in Verh. Batav. Gen.* xi. (1826) 203. f. 22 (sp. dub.).—Java.
Malamiris, Linn. Sp. Pl. 29 (farrago spp.)
Malamiris, Roxb. Hort. Beng. 80; Fl. Ind. i. 160 = attenuatum.
Malmoris, Wall. Cat. n. 6642 A = sylvaticum, hymenophyllum, argyrophyllum.
mananthum, *Wright, in Sauv. Fl. Cub.* 144.—Cuba.
Mandoni, *C. DC. in DC. Prod.* xvi. 1. 275.—Peruv.
manillanum, Miq. Syst. Pip. 339 = *clypeatum* (Fici sp.).
marequitense, *C. DC. in DC. Prod.* xvi. 1. 290.—N. Granat.
margaritiferum, Bert. ex Steud. Nom. ed. II. ii. 342 = Peperomia magaritifera.
marginatum, Haw. Suppl. Pl. Succ. 3 = Peperomia obtusifolia.
marginatum, *Jacq. Ic. Pl. Rar.* ii. 2. t. 215.—Ind. occ.
marianum, Opiz, in Presl, Rel. Haenk. i. 159 = P. Betle.
maritimum, Blume ex C. DC. in DC. Prod. xvi. 1. 356 = P. Chaba, *Hunter.*
marowinianum, *C. DC. l. c.* 337.—Bras.
marowynense, *C. DC. l. c.* 268.—Guian.
Martensianum, *C. DC. l. c.* 251.—Mexic.
Martianum, *Kunth, in Linnaea*, xiii. (1839) 691.—Bras.
mascarense, D. Dietr. Syn. Pl. i. 120 = pyrifolium.
Mastersianum, *C. DC. in DC. Prod.* xvi. 1. 282.—Bras.
Matthewsii, *C. DC. l. c.* 261.—Peruv.
mauritianum, Schult. Mant. i. 245 = Peperomia elliptica.
maynasianum, *C. DC. in DC. Prod.* xvi. 1. 262.—Peruv.
maypurense, Poir. Encyc. Suppl. iv. 468 = Peperomia maypurensis.
medium, *Jacq. Ic. Pl. Rar.* i. 2. t. 8.—Mexic.
megalophyllum, *C. DC. in Linnaea*, xxxvii. (1871–73) 357.—Mexic.
Melamiris, C. DC. in DC. Prod. xvi. 1. 359, sphalm. = *P. Malamiris.*
melanostictum, *C. DC. l. c.* 274.—Mexic.
melastomoides, *Schlecht. & Cham. in Linnaea*, v. (1830) 74.—Mexic.
membranaceum, *C. DC. in DC. Prod.* xvi. 1. 366.—Mexic.
methysticum, *Forst. f. Pl. Escul.* 76.—Ins. Pacif.
methysticum, Linn. f. Suppl. 91 = latifolium.
methysticum, Roxb. Hort. Beng. 80; Fl. Ind. i. 159 = majusculum.
mexicanum, *C. DC. in DC. Prod.* xvi. 1. 276.—Mexic.
miconiaefolium, *C. DC. l. c.* 375.—Ecuador.
micranthera, *C. DC. in Linnaea*, xxxvii. (1871–73) 354.—Nicaragua.
microphyllum, Poir. Encyc. Suppl. iv. 469 = Peperomia microphylla.

PIPER:—
microstachyon, *Vahl, Eclog. Am.* ii. 3.—Guian.
microstigma, *C. DC. in DC. Prod.* xvi. 1. 351.—Malaya.
Miersinum, *C. DC. l. c.* 278.—Ecuador.
Mikanianum, *Steud. Nom.* ed. II. ii. 342.—Bras.
Milleri, *Roem. & Schult. Syst.* i. 337.—Ind. occ.
Millii, *C. DC. in DC. Prod.* xvi. 1. 380.—Hab.?
Minarum, *C. DC. l. c.* 304.—Bras.
miniatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 166.—Malaya.
Miquelianum, *C. DC. in DC. Prod.* xvi. 1. 254.—Bras.
miradorense, *C. DC. l. c.* 380.—Mexic.
misantlense, *C. DC. l. c.* 286.—Mexic.
mite, *Ruiz & Pav. Fl. Per.* i. 37. t. 58. f. a.—Peruv.
modestum, *C. DC. in DC. Prod.* xvi. 1. 275.—Guian.
molle, Poir. Encyc. Suppl. iv. 467 = Peperomia mollis.
mollicomum, *Kunth, in Linnaea*, xiii. (1839) 648.—Bras.
mollipilosum, *C. DC. in DC. Prod.* xvi. 1. 262.—Bras.
mollissimum, *Blume, in Verh. Batav. Gen.* xi. (1826) 223. f. 30.—Java.
moluccanum, Spreng. Syst. i. 112 = miniatum.
monostachion, Vell. Fl. Flum. 26; i. t. 63 = Peperomia hederacea.
monostachyos, Poir. Encyc. Suppl. iv. 465 = Peperomia pyrifolia.
monostachyum, Vahl, Enum. i. 343 = Peperomia monostachya.
montanum, *C. DC. in Journ. Bot.* iv. (1866) 214.—N. Granat.
Morisonianum, *DC. in DC. Prod.* xvi. 1. 311.—Bras.
Moritzianum, *C. DC. l. c.* 288.—N. Granat.
mucronatum, Rich. ex C. DC. l. c. 409 = Peperomia melanostigma.
mucronulatum, *Blume, in Verh. Batav. Gen.* xi. (1826) 218. f. 27.—Java.
Muelleri, *C. DC. in DC. Prod.* xvi. 1. 243.—Mexic.
Mullesua, Buch.-Ham. in D. Don, Prod. Fl. Nep. 20 = brachystachyum.
multinervium, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 130.—Mexic.
multiplinervium, *C. DC. in Journ. Bot.* iv. (1866) 214.—N. Granat.
munchanum, *C. DC. in DC. Prod.* xvi. 1. 258.—Peruv.
muricatum, *Blume, Cat. Gew. Buitenz.* 33.—Java.
murinum, Roem. & Schult. Syst. i. 325 = Peperomia pyrifolia.
Murrayanum, *C. DC. in DC. Prod.* xvi. 1. 280.—Jamaic.
muscosum, Schult. Mant. i. 245 = Peperomia muscosa.
myosuroides, Rudge, Pl. Rar. Gui. 4. t. 5 = Peperomia myosuroides.
Myosuros, Willd. ex D. Dietr. Syn. Pl. i. 110 = Peperomia scandens.
myrtifolium, Vahl, Enum. i. 341 = Peperomia myrtifolia.
Nagelii, *C. DC. in DC. Prod.* xvi. 1. 365.—Java.
naranjoanum, *C. DC. in Linnaea*, xxxvii. (1871–73) 363.—Costa Rica.
Neesianum, *C. DC. in DC. Prod.* xvi. 1. 256.—Mexic.
negroënse, *C. DC. l. c.* 297.—Bras.
Nematanthera, *C. DC. l. c.* 367.—Guian.
nematostachyum, Schult. Mant. i. 248 = Peperomia nematostachya.
nemorosum, Vahl, Enum. i. 341 = Peperomia acuminata.
nemostachyum, A. Dietr. Sp. Pl. i. 156, sphalm. = Peperomia nematostachya.
nepalense, *Miq. Syst. Pip.* 318.—Reg. Himal.
nervoso-venosum, Blume, in Verh. Batav. Gen. xi. (1826) 228 = Peperomia nervoso-venosa.
nervosum, *C. DC. in DC. Prod.* xvi. 1. 374.—Mexic.
nervulosum, *C. DC. in DC. Prod.* xvi. 1. 268.—Bras.
Nhandi, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = marginatum.
nigrescens, *Blume, in Verh. Batav. Gen.* xi. (1826) 161. f. 3.—Java.
nigrescens, Miq. Ann. Mus. Bot. Lugd. Bat. i. 137 = sulcatum.
nigricans, *Willd. ex Link, Jahrb.* i. III. (1820) 60.—Ins. Maurit.
nigrum, Beyr. ex Kunth, in Linnaea, xiii. (1839) 618 = amplum.
nigrum, Lam. ex Link, Jahrb. i. III. (1820) 60 = nigricans.
nigrum, *Linn. Sp. Pl.* 28.—Ind. or.; Malaya.

PIPER:—
nigrum, Wall. Cat. n. 4429 = Wightii.
nigrum, Wall. l. c. n. 6643, E, F = porphyrophyllum.
nilghirianum, C. DC. in DC. Prod. xvi. 1. 364 = hymenophyllum.
nitens, Desf. Tabl. ed. I. 211 = Peperomia trinervis?
nitidulum, *Opiz, in Presl, Rel. Haenk.* i. 154.—Mexic.
nitidum, Sw. Prod. Veg. Ind. Occ. 15 = geniculatum.
nitidum, *Vahl, Enum.* i. 314.—Am. trop.
nobile, *C. DC. in DC. Prod.* xvi. 1. 289.—Venezuela.
nodulosum, Link, Enum. Hort. Berol. i. 36 = unguiculatum.
Novae-Hispaniae, *C. DC. in DC. Prod.* xvi. 1. 301.—Mexic.
Novae-Hollandiae, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 60.—Austral.
novo-granatense, *C. DC. in DC. Prod.* xvi. 1. 313.—N. Granat.
nubigenum, Kunth, in Linnaea, xiii. (1839) 654 = crocatum.
nudum, *C. DC. in DC. Prod.* xvi. 1. 325.—Mexic.
nummulariaefolium, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia nummularifolia.
nummularifolium, Lam. & Willd. ex C. DC. in DC. Prod. xvi. 382 = Filix sterilis (Crypt.).
nummularifolium, Pohl, ex Miq. in Mart. Fl. Bras. iv. 1. 21 = Peperomia circinnata.
nummularium, Lam. Illustr. i. 82 = Peperomia serpyllifolia.
nutans, Opiz, in Presl, Rel. Haenk. i. 156. t. 28. f. 2 = tuberculatum.
oaxacanum, *C. DC. in DC. Prod.* xvi. 1. 274.—Mexic.
obcordatum, Stokes, Bot. Comment. i. 176 = Peperomia cordifolia.
obesum, Vahl, Enum. i. 321 = scabrum.
obliquum, Balb. ex Kunth, in Linnaea, xiii. (1839) 614 tuberculatum.
obliquum, Pers. Syn. i. 35 = Peperomia acuminata.
obliquum, *Ruiz & Pav. Fl. Per.* i. 37. t. 63. f. a.—Peruv.
oblongatum, *Opiz, in Presl, Rel. Haenk.* i. 153.—Peruv.
oblongifolium, *C. DC. in DC. Prod.* xvi. 1. 273.—Bras.
oblongum, *H. B. & K. Nov. Gen. et Sp.* i. 52.—Venezuela.
oblongum, Schlecht. & Cham. in Linnaea, v. (1830) 73 = Chamissonis.
oblongum, Wall. Cat. n. 6663 C = Peperomia Heyneana.
obovatum, *Ruiz & Pav. Fl. Per.* i. 36. t. 53. f. b.—Peruv.
obovatum, Vahl, Eclog. Am. i. 5 = Peperomia obversa.
obscurum, *C. DC. in Linnaea*, xxxvii. (1871–73) 348.—Bras.
obtusifolium, *Hill, Syst. Veg.* xxvi. 19. t. 19. f. 1.—Am. austr.
obtusifolium, Jacq. Coll. i. 141 = Peperomia polystachya.
obtusifolium, Linn. Sp. Pl. 30 = Peperomia obtusifolia.
obtusissimum, Miq. Syst. Pip. 337 = *clypeatum* (Fici sp.).
obtusum, *C. DC. in DC. Prod.* xvi. 1. 328.—Cuba.
obumbratum, *C. DC. l. c.* 265.—Bras.
obversum, Vahl, Enum. i. 354 = Peperomia obversa.
Oerstedii, *C. DC. in Linnaea*, xxxvii. (1871–73) 359.—Costa Rica.
officinarum, C. DC. in DC. Prod. xiv. 1. 356 = P. Chaba, *Hunter*.
Olfersianum, Kunth, in Linnaea, xiii. (1839) 646 = hirsutum.
olivaceum, *C. DC. in Linnaea*, xxxvii. (1871–73) 344.—Bras.
Opizianum, D. Dietr. Syn. Pl. i. 110 = Peperomia dependens.
Opizianum, [*Fuernr.*] *in Flora*, xii. (1829) 228.—Peruv.
Opisii, Steud. Nom. ed. II. ii. 342 = asperifolium.
orbiculatum, Poir. Encyc. v. 476 = Peperomia orbiculata.
orbiculatum, Spreng. Syst. i. 121 = Peperomia fimbriata.
ornatum, *N. E. Br. in Gard. Chron.* (1884) II. 424.—Ins. Celebes.
orthostachyum, *Kunth, in Linnaea*, xiii. (1839) 602.—Bras.

PIPER:—
ottoniaefolium, *C. DC. in Journ. Bot.* iv. (1866) 213.—N. Granat.
Ottonis, C. DC. in DC. Prod. xvi. 1. 332 = peltatum.
ovale, *C. DC. in Journ. Bot.* iv. (1866) 166.—Venezuela.
ovalifolium, *C. DC. in DC. Prod.* xvi. 1. 305.—Bras.
ovalifolium, D. Dietr. Syn. Pl. i. 112 = Peperomia ovalifolia.
ovalifolium, Heyne, in Wall. Cat. n. 6663 B = Peperomia dindigulensis.
ovatum, *Vahl, Eclog. Am.* i. 3. t. 1.—Ins. Trinit.
pachyphyllum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 436.—Madag.
pachystachyon, *C. DC. in Journ. Bot.* iv. (1866) 216.—N. Granat.
padifolium, *C. DC. in DC. Prod.* xvi. 1. 253.—Bras.
pallens, Poir. Encyc. Suppl. iv. 469 = Peperomia pallens.
pallescens, *C. DC. in Linnaea*, xxxvii. (1871–72) 362.—Bras.
pallidum, Forst. f. Prod. 5 = Peperomia pallida.
pallidum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 120.—Ind. or.
pallidum, Salisb. Prod. 11 = Peperomia polystachya.
paludosum, *C. DC. in DC. Prod.* xvi. 1. 283.—Bras.
palustre, *C. DC. l. c.* 293.—Bras.
panamense, *C. DC. in Journ. Bot.* iv. (1866) 216.—Panama.
panduratum, *C. DC. in DC. Prod.* xvi. 1. 250.—Jamaic.
pangoënse, *C. DC. l. c.* 368.—Peruv.
paniculatum, Willd. ex C. DC. l. c. 401 = Peperomia secunda.
papantlense, *C. DC. l. c.* 338.—Mexic.
paraënse, *C. DC. l. c.* 290.—Bras.
paraguayense, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 45; *Contrib.* 97 ('n. 2').—Paraguay.
paraguayense, *Parodi, l. c.* 90; *Contrib.* 99 ('n. 8') (sp. div.).—Paraguay.
paramaribense, *C. DC. in DC. Prod.* xvi. 1. 297.—Guian.
paramattense, C. DC. l. c. 353 = Novae-Hollandiae.
Parkerianum, *C. DC. l. c.* 373.—Guian.
Parthenium, *Mart. Syst. Mat. Med. Bras.* 100.—Bras.
parvifolium, *Blanco, Fl. Philip.* ed. I. 23.—Ins. Philipp.
parvulum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 130 = Peperomia mexicana.
patens, Hook. & Arn. Bot. Beech. Voy. 310 = nervosum.
patens, *Opiz, in Presl, Rel. Haenk.* i. 154.—Peruv.
patulum, *Bertol. Fl. Guatim.* 7. t. 1.—Guatemala.
patulum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 128 = miradorense.
paulianum, *C. DC. in DC. Prod.* xvi. 1. 294.—Bras.
Pavoni, *C. DC. l. c.* 294.—Peruv.
pedicellare, *C. DC. in Journ. Bot.* iv. (1866) 211.—N. Granat.
pedicellatum, C. DC. l. c. 164 = boehmeriaefolium.
pedicellosum, *Wall. Cat.* n. 6646 A.—Singapore.
pedicellosum, Wall. l. c. B = sumatranum.
pedunculare, *C. DC. in DC. Prod.* xvi. 1. 291.—Guian.
pedunculatum, *C. DC. in Journ. Bot.* iv. (1866) 162.—N. Granat.
pedunculatum, Reinw. ex Miq. in Linnaea, xxi. (1848) 484 = caninum.
peepuloides, *Roxb. Hort. Beng.* 4; *Fl. Ind.* i. 157.—Reg. Himal.
peepuloides, Wall. Cat. n. 6650 = nepalense, P. Chaba, *Hunter*, P. Betle.
pellucidum, Linn. Sp. Pl. 30 = Peperomia pellucida.
peltatum, *Linn. l. c.*—Am. trop.
peltatum, Ruiz & Pav. Fl. Per. i. 38. t. 59. f. a = umbellatum.
Peltobryon, *C. DC. in DC. Prod.* xvi. 1. 259.—Am. trop.
peltoideum, Poir. Encyc. Suppl. iv. 465 = Peperomia peltoidea.
penangense, *C. DC. in DC. Prod.* xvi. 1. 353.—Ins. Penang.
pendulum, Willd. ex Schult. Mant. i. 248 = Peperomia pendula.
peploides, Poir. Encyc. Suppl. iv. 469 = Peperomia peploides.

PIPER:—

pepuloides, D. Dietr. Syn. Pl. i. 108, sphalm. = peepuloides.
pereskiaefolium, Jacq. Coll. iv. 126 = Peperomia pereskiaefolia.
perforatum, D. Dietr. Syn. Pl. i. 108 = Peperomia perforata.
Perrottetii, *C. DC. in DC. Prod.* xvi. 1. 379.—Ins. Borbon.
persicariaefolium, *Kunth, in Linnaea*, xiii. (1839) 684.—Bras.
peruvianum, *C. DC. in DC. Prod.* xvi. 1. 283.—Peruv.
petiolare, *C. DC. in Journ. Bot.* iv. (1866) 212.—N. Granat.
petiolatum, *C. DC. l. c.* 161.—Reg. Himal.
philippinense, *C. DC. in DC. Prod.* xvi. 1. 353.—Ins. Philipp.
philippinum, *Miq. Syst. Pip.* 322.—Ins. Philipp.
phthinotrichon, *C. DC. in DC. Prod.* xvi. 1. 298.—Peruv.
Phyllanthus, Willd. ex A. Dietr. Sp. Pl. i. 183 = Peperomia foliiflora.
phyllostictum, *C. DC. in DC. Prod.* xvi. 1. 340.—Java.
phytolaccaefolium, *Opiz, in Presl, Rel. Haenk.* i. 151.—Ecuador.
pilosiusculum, *Opiz, l. c.* 154. t. 29.—Panama.
pilosum, *C. DC. in Linnaea*, xxxvii. (1871–73) 351.—Bras.
pilosum, Vahl, Enum. i. 338 = Peperomia pilosa.
piluliferum, *H. B. & K. Nov. Gen. et Sp.* i. 53.—Ecuador.
pinnatum, *Lour. Fl. Cochinch.* 31.—China.
piperoides, Schult. Mant. i. 240, sphalm. = peepuloides.
pisoënse, *C. DC. in DC. Prod.* xvi. 1. 278.—Bras.
planifolium, Poir. Encyc. v. 473 = Peperomia acuminata.
plantagineum, Buch.-Ham. ex Wall. Cat. n. 6659 = Hamiltonii.
plantagineum, Lam. Illustr. i. 80 = P. Amalago.
plantagineum, *Schlecht. in Linnaea*, vi. (1831) 353.—Mexic.
plantaginifolium, D. Dietr. Syn. Pl. i. 110 = chilense.
platyphyllum, *C. DC. in DC. Prod.* xvi. 1. 375.—Mexic.
plicatum, D. Dietr. Syn. Pl. i. 112 = Peperomia plicata.
Plukenetianum, *C. DC. in DC. Prod.* xvi. 1. 315.—Bras.
Poeppigii, *C. DC. l. c.* 260.—Bras.
Poeppigii, *Steud Nom.* ed. II. ii. 342.—Peruv.
Pohlianum, *C. DC. in DC. Prod.* xvi. 1. 256.—Bras.
Poiretianum, *C. DC. l. c.* 325.—Bras.
Poiteanum, *Kunth, in Linnaea*, xiii. (1839) 607.—Guian.
polybotrya, Poir. Encyc. Suppl. iv. 465 = Peperomia polybotrya.
polystachyon, Ait. Hort. Kew. ed. I. i. 49 = Peperomia polystachya.
populifolium, *Kunth, in Linnaea*, xiii. (1839) 710.—Bras.
populifolium, Opiz, in Presl, Rel. Haenk. i. 160 = cardiophyllum.
porphyrophyllum, *N. E. Br. in Gard. Chron.* (1884) II. 438.—Malaya.
portulacaefolium, Poir. Encyc. Suppl. iv. 470 = Peperomia portulacaefolia.
portulacoides, Lam. Illustr. i. 82 = Peperomia portulacoides.
potamogetonifolium, *Opiz, in Presl, Rel. Haenk.* i. 156.—Ins. Marian.
pothifolium, *Kunth, in Linnaea*, xiii. (1839) 671.—Bras.
pothiforme, *Wall. ex C. DC. in DC. Prod.* xvi. 1. 351.—Burma.
pothoides, *Wall. Cat.* n. 6639 A.—Burma.
pothomorphe, *C. DC. in DC. Prod.* xvi. 1. 308, 492.—Mexic.
praegnans, Rich. ex C. DC. l. c. 287 = Hostmannianum.
praemorsum, Vahl, Eclog. Am. i. 4 = geniculatum.
Preslii, *C. DC. in DC. Prod.* xvi. 1. 257.—N. Granat.?
prianamense, *C. DC. l. c.* 376.—Sumatra.
prismaticum, *C. DC. in Linnaea*, xxxvii. (1871–73) 342.—Costa Rica.
propinquum, *C. DC. in Journ. Bot.* iv. (1866) 167.—Peruv.
pruinosum, H. B. & K. Nov. Gen. et Sp. i. 59 = peltatum.

PIPER:—

prunifolium, *Jacq. Eclog. Am.* i. 140. t. 94.—Venezuela.
Pseudo-amalago, *C. DC. in Linnaea*, xxxvii. (1871–73) 353.—Bras.
Pseudo-asperifolium, *C. DC. in DC. Prod.* xvi. 1. 318.—Mexic.
Pseudo-blattarum, *C. DC. in Linnaea*, xxxvii. (1871–73) 362.—Bras.?
Pseudo-chavica, *C. DC. in DC. Prod.* xv. 1. 351.—Malaya.
Pseudo-churumayu, *Poepp. ex Steud. Nom.* ed. II. ii. 342; *C. DC. l. c.* 288.—Bras.
Pseudo-cubeba, Korth. ex Miq. Comm. Phyt. 22 = sumatranum.
Pseudo-fuligineum, *C. DC. in Linnaea*, xxxvii. (1871–73) 355.—Costa Rica.
Pseudo-Lindenii, *C. DC. l. c.* 335.—Nicaragua.
Pseudo-mollicomum, *C. DC. in DC. Prod.* xvi. 1. 278.—Peruv.
Pseudo-peruvianum, *C. DC. l. c.* 282.—Bras.
Pseudo-pothifolium, *C. DC. l. c.* 289.—Bras.
Pseudo-propinquum, *C. DC. in Linnaea*, xxxvii. (1871–73) 341.—Costa Rica.
Pseudo-velutinum, *C. DC. in DC. Prod.* xvi. 1. 282.—Bras.
psilostachyum, *H. B. & K. Nov. Gen. et Sp.* i. 50.—N. Granat.
psittacorum, *Endl. Prod. Fl. Norf.* 37.—Ins. Norfolk.
pteropodon, *C. DC. in DC. Prod.* xvi. 1. 255.—Bras.
pubescens, Haw. Syn. Pl. Succ. 7 = Peperomia Haworthiana.
pubescens, Pers. Syn. i. 34 = Peperomia pubescens.
pubescens, Soland. ex C. DC. in DC. Prod. xvi. 1. 342 = mollissimum.
pubescens, Vahl, Enum. i. 348 = Peperomia Vahlii.
pulchellum, Ait. Hort. Kew. ed. I. 50 = Peperomia pulchella.
pulchrum, *C. DC. in DC. Prod.* xvi. 1. 305.—N. Granat.
pulicare, D. Dietr. Syn. Pl. i. 110 = Peperomia scandens?
pumilum, D. Dietr. l. c. = Peperomia pumila.
punctatissimum, *C. DC. in DC. Prod.* xvi. 1. 255.—Bras.
punctatum, Poepp. ex Kunth, in Linnaea. xiii. (1839) 689 = heterophyllum.
punctatum, *Ruiz & Pav. Fl. Per.* i. 36. t. 62. f. a.—Peruv.
pupuloides, Spreng. Syst. i. 117, sphalm. = peepuloides.
Purdiei, *C. DC. in DC. Prod.* xvi. 1. 299.—N. Granat.
purpurascens, *Desf. Cat. Hort. Par.* ed. III. 414.—Hab.?
purpurascens, D. Dietr. Syn. Pl. i. 119 = angustifolium.
purpureum, Vahl, Enum. i. 343 = Peperomia purpurea.
pusillum, Blume, in Verh. Batav. Gen. xi. (1826) 232, 382 = Peperomia reflexa.
pyrifolium, Opiz, in Presl, Rel. Haenk. i. 158 = unguiculatum.
pyrifolium, *Vahl, Enum.* i. 327.—Ins. Mascar.
quadrangulare, Thomps. in Trans. Linn. Soc. ix. (1808) 202 = Peperomia quadrangularis.
quadrangulum, Willd. ex Miq. in Mart. Fl. Bras. iv. 1. 21 = Peperomia angulata.
quadrifolium, Linn. Sp. Pl. ed. II. 43 = Peperomia quadrifolia.
quadrifolium, Vell. Fl. Flum. 26; i. t. 62 = Peperomia reflexa.
quinqueangulatum, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 85.—Malaya.
quitense, *C. DC. in Journ. Bot.* iv. (1866) 163.—Ecuador.
racemosum, *Mill. Gard. Dict.* ed. VIII. n. 13.—Mexic.
racemosum, Vahl, Enum. i. 337 = Peperomia secunda.
radicans, *Vahl, l. c.* 333.—Ins. Maurit.
Radula, H. B. & K. Nov. Gen. et Sp. i. 48 = Bredemeyeri.
Rafflesii, *C. DC. in DC. Prod.* xvi. 1. 349.—Sumatra.
rarum, *C. DC. l. c.* 284.—Bras.
realgoanum, *C. DC. in Linnaea*, xxxvii. (1871–73) 335.—Nicaragua.
recurvatum, Blume, in Verh. Batav. Gen. xi. (1826) 233. f. 38 = Peperomia recurvata.
recurvum, *Blume, l. c.* 176. t. 4.—Java.
recurvum, Rich. ex C. DC. in DC. Prod. xvi. 1. 301 = inaequale.

PIPER:—

reflexum, Linn. f. Suppl. 91 = Peperomia reflexa.
Regelii, *C. DC. in DC. Prod.* xvi. 1. 309.—Bras.
Regnellii, *C. DC. l. c.* 307.—Bras.
Reinwardtianum, *C. DC. l. c.* 354.—Malaya.
reniforme, *Poir. Encyc.* v. 465.—Hab.?
reniforme, Willd. ex Link, Jahrb. i. III. (1820) 64 = Verhuellia elegans.
repens, Poir. Encyc. Suppl. iv. 467 = Peperomia scandens.
reticulatum, Hoffmgg. Verz. Pfl. 100 = plantagineum.
reticulatum, *Linn. Sp. Pl.* 29.—Ind. occ.
reticulatum, Vell. Fl. Flum. 261; i. t. 61 = ceanothifolium.
reticulosum, *Opiz, in Presl, Rel. Haenk.* i. 155.—Am. trop.
retrofractum, *Vahl, Enum.* i. 314.—Ind. or.
retusum, Linn. f. Suppl. 91 = Peperomia retusa.
revolutum, *Rich. ex H. B. & K. Nov. Gen. et Sp.* i. t. 3.—Hab.?
rhododendrifolium, *Kunth, in Linnaea*, xiii. (1839) 621.—Bras.
rhombeum, Vahl, Enum. i. 353 = Peperomia rhombea.
rhomboidale, *Dum. Cours. Bot. Cult.* ed. II. vi. 379; *et ex Desf. Cat. Hort. Par.* ed. III. 414.—Hab.?
rhombophyllum, *C. DC. in DC. Prod.* xvi. 1. 352.—Malaya.
Rhyncholepsis, *C. DC. l. c.* 344.—Malaya.
rhynchostachyum, *C. DC. l. c.* 326.—Guian.
ribesioides, *Wall. Cat.* n. 6637.—Burma; Malaya.
richardiaefolium, *Kunth, in Linnaea*, xiii. (1839) 669.—Bras.
Richardianum, *C. DC. in DC. Prod.* xvi. 1. 249.—Ind. occ.
Riedelii, *C. DC. l. c.* 325.—Bras.
rigidum, *C. DC. l. c.* 264.—Cuba.
riparium, H. B. & K. Nov. Gen. et Sp. i. 48 = grandifolium.
rivinoides, *Kunth, in Linnaea*, xiii. (1839) 682.—Bras.
rivulare, *C. DC. in Linnaea*, xxxvii. (1871–72) 360.—Bras.
Rohrii, *C. DC. in DC. Prod.* xvi. 1. 296.—Bras.
Rollingo, Schult. Mant. i. 247 = Peperomia Rollingo.
rostratum, Roxb. Hort. Beng. 80; Fl. Ind. i. 160 = Peperomia rostrata.
rotundatum, Poir. Encyc. Suppl. iv. 468 = Peperomia rotundata.
rotundifolium, *Hill, Veg. Syst.* xxvi. 19. t. 19. f. 2.—Jamaic.
rotundifolium, Linn. Sp. Pl. 30 = Peperomia cyclophylla.
rotundifolium, Sw. Obs. Bot. 20 = Peperomia circinata.
rotundum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = nigrum.
Roxburghianum, Schult. Mant. i. 242 = Peperomia Roxburghiana.
rubellum, Haw. ex Spreng. Syst. iv. Cur. Post. 20 = Peperomia rubella.
rubescens, *C. DC. in DC. Prod.* xvi. 1. 300.—Guian.
rubioides, Poir. Encyc. Suppl. iv. 470 = Peperomia rubioides.
rubricaule, Link, Enum. Hort. Berol. i. 39 = Peperomia pereskiaefolia.
rubricaule, Nees, in Hort. Phys. Berol. 48. t. 8 = Peperomia rubricaulis.
rubrum, Reinw. ex Miq. Fl. Ind. Bat. i. II. 440 = miniatum.
rude, *H. B. & K. Nov. Gen. et Sp.* i. 49.—N. Granat.
Rudgeanum, *C. DC. in DC. Prod.* xvi. 1. 373.—Guian.
ruficaule, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 92; *Contrib.* 102.—Reg. Argent.
rufinerve, *Opiz, in Presl, Rel. Haenk.* i. 159.—Ins. Philipp.
rugosum, Bredem. ex Willd. Enum. Hort. Berol. Suppl. 3 = Bredemeyeri.
rugosum, *Lam. Illustr.* i. 81.—Ins. S. Doming.
rugosum, Opiz, in Presl, Rel. Haenk. i. 152 = crocatum.
rugosum, Vahl, Eclog. Am. i. 4 = citrifolium.
rugulosum, *C. DC. in Journ. Bot.* iv. (1866) 211.—N. Granat.
Ruizianum, *C. DC. in DC. Prod.* xvi. 1. 308.—Mexic.
Ruizianum, D. Dietr. Syn. Pl. i. 118 = obliquum.
rumicifolium, *C. DC. in DC. Prod.* xvi. 1. 297.—Bras.
Rumphii, *C. DC. l. c.* 354.—Ins. Ternate.
rupestre, Poir. Encyc. Suppl. iv. 465 = Peperomia rupestris.

PIPER:—
ruscifolium, Poir. Encyc. Suppl. iv. 466 = Peperomia ruscifolia.
Sagoti, *C. DC. in Journ. Bot.* iv. (1866) 162.—Guian.
Sagranum, *C. DC. in DC. Prod.* xvi. 1. 292.—Cuba.
salicariaefolium, *Kunth, in Linnaea*, xiii. (1839) 636.—Bras.
salicifolium, *Vahl, Enum.* i. 312.—Guian.
salicinum, *Opiz, in Presl, Rel. Haenk.* i. 150.—Panama.
salignum, Poir. Encyc. Suppl. iv. 466 = Peperomia saligna.
saliuncum, A. Dietr.,Sp. Pl. i. 635, sphalm. = salicinum.
salviaefolium, *C. DC. in DC. Prod.* xvi. 1. 270.—Peruv.
Salzmanni, *C. DC. l. c.* 387, nomen.—Bras.
sancarlosianum, *C. DC. l. c.* 337.—Bras.
sanctum, *Schlecht. ex Miq. in Linnaea*, xviii. (1844) 714.—Mexic.
san-joseanum, *C. DC. in Linnaea*, xxxvii. (1871–73) 351.—Costa Rica.
santaremense, *C. DC. in DC. Prod.* xvi. 1. 275.—Bras.
sarmentosum, *Roxb. Hort. Beng.* 4; *Fl. Ind.* i. 160.—Ind. or.; Malaya.
sarmentosum, Wall. Cat. n. 6641 = longum.
saururoides, *Ham. Prod. Pl. Ind. Occ.* 3.—Ind. occ.
Saururus, Burm. Ind. Fl. Malab. 3 = P. Betle.
saxatilis, Wall. in Roxb. Fl. Ind. ed. Carey, i. 163 = Peperomia reflexa.
saxicolum, Vahl, Enum. 345 = Peperomia pubescens.
scabridum, Haenke, in Flora, xii. (1829) 228, nomen = scabriusculum.
scabrifolium, *Hook. & Arn. Bot. Beech. Voy.* 310.—Mexic.
scabriusculum, *A. Dietr. Sp. Pl.* i. 647.—Peruv.
scabrum, Lam. Illustr. i. 80 = aduncum.
scabrum, Opiz, in Presl, Rel. Haenk. i. 151 = scabriusculum.
scabrum, Poepp. ex Kunth, in Linnaea, xiii. (1839) 647 = Poeppigii, *Steud.*
scabrum, *Ruiz & Pav. Fl. Per.* i. 34. t. 55.—Peruv.
scabrum, Sw. Fl. Ind. Occ. i. 59 = hirsutum.
scabrum, Willd. ex Kunth, in Linnaea, xiii. (1839) 614 = tuberculatum.
scandens, Poepp. ex Miq. in Mart. Fl. Bras. iv. 1. 10 = Peperomia pellucida.
scandens, Sw. Prod. Veg. Ind. Occ. 15 = Peperomia glabella.
scandens, Vahl, Enum. i. 346 = Peperomia scandens.
Schachii, *C. DC. in DC. Prod.* xvi. 1. 288.—Ind. occ.
Schiedeanum, *Steud. Nom.* ed. II. ii. 343.—Mexic.
schizonephos, *C. DC, in DC. Prod.* xvi. 1. 241.—Malacca.
Schlechtendalianum, *C. DC. l. c.* 324.—Mexic.
Schlechtendalii, *Steud. Nom.* ed. II. ii. 343.—Mexic.
Schlimii, *C. DC. in DC. Prod.* xvi. 1. 274.—Venezuela.
Schottii, *C. DC. l. c.* 315.—Bras.
scopulorum, *Kunth, in Linnaea*, xiii. (1839) 669.—Bras.
scutatum, Willd. ex A. Dietr. Sp. Pl. i. 143 = peltatum.
scutellaefolium, Vahl, Enum. i. 356 = Peperomia scutellaefolia.
scutellatum, Pers. Syn. i. 34 = Peperomia scutellaefolia.
scutelliferum, *C. DC. in Linnaea*, xxxvii. (1871–73) 349.—Bras.
scutiphyllum, *Ham. Prod. Pl. Ind. Occ.* 3.—Ins. S. Doming.
sebastianopolitense, *C. DC. in DC. Prod.* xvi. 1. 252.—Bras.
secundum, Poepp. ex Kunth, in Linnaea, xiii. (1839) 612 = geniculatum.
secundum, Poir. Encyc. iv. 474 = Peperomia secunda.
secundum, *Ruiz & Pav. Fl. Per.* i. 36.—Peruv.
secundum, Willd. ex Cham. & Schlecht. in Linnaea, v. (1830) 73 = variegatum.
Seemannianum, *C. DC. in Journ. Bot.* iv. (1866) 164.—N. Hibern.
Selloi, *C. DC. in DC. Prod.* xvi. 1. 252.—Bras.
Sellowianum, *Kunth, in Linnaea*, xiii. (1839) 658.—Bras.
septemnerve, Vahl, Enum. i. 349 = Peperomia septemnervis.
septuplinervium, *C. DC. in DC. Prod.* xvi. 1. 313.—Panama.

PIPER:—
serpens, Sw. Prod. Veg. Ind. Occ. 16 = Verhuellia serpens.
serpentarioides, Nees, ex Miq. in Mart. Fl. Bras. iv. 1. 9 = Peperomia serpentarioides.
serpyllifolium, Vahl, Enum. i. 351 = Peperomia serpyllifera.
sessilifolium, Willd. ex Schult. Mant. i. 248 = Peperomia sessilifolia.
sidaefolium, Link & Otto, Ic. Pl. Sel. Rar. 15. t. 6 = subpeltatum.
Sieberi, *C. DC. in DC. Prod.* xvi. 1. 248.—Ind. occ.
silvestre, Vell. Fl. Flum. 25; i. t. 56 = amplum.
simplex, D. Dietr. Syn. Pl. i. 109 = Peperomia simplex.
simplex, *Endl. Prod. Fl. Ins. Norf.* 37.—Ins. Norfolk.
sinense, *C. DC. in DC. Prod.* xvi. 1. 361.—China.
singulare, *C. DC. in Journ. Bot.* iv. (1866) 161.—N. Granat.
Siriboa, Forst. f. Prod. 4 = austro-caledonicum.
Siriboa, Heyne, in Wall. Cat. n. 6651 B = P. Hapnium.
Siriboa, Linn. Sp. Pl. 29 = P. Betle.
Sirium, *C. DC. in DC. Prod.* xvi. 1. 361.—Malaya.
smilacifolium, *H. B. & K. Nov. Gen. et Sp.* i. 56.—Venezuela.
smilacifolium, Willd. ex Kunth, in Linnaea, xiii. (1839) 603 = laevigatum.
Solmsianum, *C. DC. in DC. Prod.* xvi. 1. 291.—Bras.
sororium, *C. DC. l. c.* 295.—Peruv.
spathulatum, D. Dietr. Syn. Pl. i. 109 = Peperomia spathulata.
speciosum, H. B. & K. Nov. Gen. et Sp. i. 59 = peltatum.
sphaerostachyum, *C. DC. in DC. Prod.* xvi. 1. 326.—Cuba.
Sprengelianum, *C. DC. l. c.* 283.—Bras.
Spruceanum, *C. DC. in Journ. Bot.* iv. (1866) 216.—Peruv.
spurium, Link, Enum. Hort. Berol. i. 37 = nigrum.
Staphyle, Roem. & Schult. Syst. i. 337 = racemosum.
stellatum, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia stellata.
stellatum, Vell. Fl. Flum. 26; i. t. 64 = Peperomia pereskiaefolia.
stellipilum, *C. DC. in DC. Prod.* xvi. 1. 262.—Bras.
stipulaceum, *Opiz, in Presl, Rel. Haenk.* i. 160.—Mexic.
stoloniferum, Poir. Encyc. Suppl. iv. 470 = Peperomia stolonifera.
striatum, *C. DC. in Linnaea*, xxxvii. (1871–73) 345.—Costa Rica.
striatum, Vahl, Enum. i. 343 = Peperomia striata.
stupposum, A. Dietr. Sp. Pl. i. 645 = crocatum.
stylosum, Miq. Ann. Bot. Mus. Lugd. Bat. i. 139 = Peperomia rostratum.
suaveolens, Desf. Cat. Hort. Par. ed. III. 414 = Peperomia suaveolens.
subaequale, *C. DC. in DC. Prod.* xvi. 1. 371.—Bras.
subandinum, *C. DC. l. c.* 275.—Peruv.
subauriculatum, *C. DC. l. c.* 323.—Bras.
subcordatum, *C. DC. l. c.* 353.—Malaya.
subflavum, *C. DC. in Journ. Bot.* iv. (1866) 210.—N. Granat.
subfuscum, *C. DC. l. c.* 217.—Costa Rica.
subglaucum, *C. DC. in Linnaea*, xxxvii. (1871–73) 337.—Bras.
subpedicellatum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 122.—Ind. or.
subpeltatum, *Willd. Sp. Pl.* i. 166.—Afr. trop.; Ind. or.; Malaya.
subrotundum, Blume, in Verh. Batav. Gen. xi. (1826) 235. f. 39 = Peperomia Reinwardtiana.
subrotundum, Haw. Suppl. Pl. Succ. 2 = Peperomia magnoliaefolia.
subscutatum, *C. DC. in DC. Prod.* xvi. 1. 321.—Peruv.
subseptemnervium, *C. DC. l. c.* 377.—Ins. Penang.
subulatum, *C. DC. in Journ. Bot.* iv. (1866) 165.—N. Granat.
Suipigua, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 20.—Nepal.
sulcatum, *Blume, Cat. Gew. Buitenz.* 34.—Java.
sulcatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 20 = Peperomia reflexa.
sumatranum, *C. DC. in DC. Prod.* xvi. 1. 343.—Burma; Malaya.

PIPER:—
sundaicum, *Blume, in Verh. Batav. Gen.* xi. (1826) 187 f. 14.—Java.
surinamense, *C. DC. in DC. Prod.* xvi. 1. 296.—Am. trop.
Swartzianum, *C. DC. l. c.* 265.—Ind. occ.
sylvaticum, *Roxb. Hort. Beng.* 4; *Fl. Ind.* i. 156.—Ind. or.; Burma.
sylvestre, *Lam. Illustr.* i. 79.—Ins. Mascar.
sylvestre, Lour. Fl. Cochinch. 30 = caninum?
syringaefolium, Balb. ex Kunth, in Linnaea, xiii. (1839) 599 = P. Amalago.
syringaefolium, *Vahl, Enum.* i. 328.—Ind. or.
talinifolium, Poir. Encyc. Suppl. iv. 465 = Peperomia talinifolia.
tarapotianum, *C. DC. in DC. Prod.* xvi. 1. 246.—Peruv.
teapense, *C. DC. l. c.* 260.—Mexic.
tectoniaefolium, *Kunth, in Linnaea*, xiii. (1839) 661.—Bras.
tejucense, *C. DC. in DC. Prod.* xvi. 1. 287.—Bras.
tenellum, Sw. Prod. Veg. Ind. Occ. 16 = Peperomia tenella.
tenue, *H. B. & K. Nov. Gen. et Sp.* i. 56.—N. Granat.
tenuicuspe, *C. DC. in DC. Prod.* xvi. 1. 304.—Peruv.
tenuiflorum, D. Dietr. Syn. Pl. i. 111 = Peperomia tenuiflora, *Opiz.*
tenuiflorum, *Vahl, in West, Bidr. Ste-Croix*, 268.—Ind. occ.
tenuifolium, *C. DC. in DC. Prod.* xvi. 1. 321.—Mexic.
tenuinerve, *C. DC. l. c.* 313.—Bras.
tenuispicatum, Steud. Nom. ed. II. ii. 343 = leptostachyum.
tenuistylum, *C. DC. in DC. Prod.* xvi. 1. 260.—Peruv.
tequendanense, *C. DC. in Journ. Bot.* iv. (1866) 215.—N. Granat.
terminale, D. Dietr. Syn. Pl. i. 120 = thermale.
terminale, H. B. & K. Nov. Gen. et Sp. i. 57 = unguiculatum.
tetragonum, Vahl, Enum. i. 349 = Peperomia tetragona.
tetraphyllum, Forst. f. Prod. 5 = Peperomia reflexa.
tetraphyllum, *Vell. Fl. Flum.* 26; i. t. 65.—Bras.
Teysmanii, *C. DC. in DC. Prod.* xvi. 1. 345.—Ins. Celebes.
thermale, *Vahl, Enum.* i. 329.—Ind. or.
Thwaitesii, C. DC. in DC. Prod. xvi. 1. 357 = caninum.
tigerianum, *C. DC. in Journ. Bot.* iv. (1866) 300.—Honduras.
tiguanum, C. DC. l. c. 218 = praec.
tiliaefolium, Desv. ex Ham. Prod. Pl. Ind. Occ. 4 = Bagassa Sagotiana.
tiliaefolium, *Schlecht. & Cham. in Linnaea*, vi. (1831) 352.—Mexic.
tithymaloides, Vahl, Enum. i. 340 = Peperomia tithymaloides.
tocacheanum, *C. DC. in DC. Prod.* xvi. 1. 264.—Darien.
tomentosum, *C. DC. l. c.* 372.—Bras.
tomentosum, *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
tomentosum, Vahl, Enum. i. 351 = Peperomia tomentosa.
Trianae, *C. DC. in Journ. Bot.* iv. (1866) 162.—N. Granat.
triandrum, *F. Muell. Fragm.* v. 197.—Austral.
tricarinatum, Haw. Suppl. Pl. Succ. 102 = Peperomia tricarinata.
trichocarpa, *C. DC. in Linnaea*, xxxvii. (1871–73) 350.—Bras.
trichoneuron, *C. DC. in DC. Prod.* xvi. 1. 281.—Guian.
trichostachyon, *C. DC. l. c.* 242.—Ind. or.
tricuspe, *C. DC. l. c.* 263.—Panama.
trifolium, Linn. Sp. Pl. 30 = Peperomia trifolia.
trifolium, Willd. ex C. DC. in DC. Prod. xvi. 1. 443 = Peperomia portulacoides.
trigonioideum, *C. DC. l. c.* 318.—Bras.
trigonum, *C. DC. in Journ. Bot.* iv. (1866) 212.—N. Granat.
trinerve, *C. DC. in DC. Prod.* xvi. 1. 292.—Panama.
trinerve, Vahl, Enum. i. 342 = Peperomia trinervis.
trineuron, *Miq. in Hook. Lond. Journ. Bot.* v. (1846) 555.—Zeylan.
trioicum, Roxb. Hort. Beng. 4; Fl. Ind. i. 151 = nigrum.
triquetrum, *Opiz, in Presl, Rel. Haenk.* i. 160.—Mexic.
tristachyon, *C. DC. in DC. Prod.* xvi. 1. 335.—Ins. Sandvic.

PIPER :—
tristachyum, Poir. Encyc. Suppl. iv. 465 = Peperomia tristachya.
tristemon, *C. DC. in DC. Prod.* xvi. 1. 328.—Venezuela.
truncatum, *Vell. Fl. Flum.* 25 ; i. t. 57.—Bras.
tuberculatum, *Jacq. Ic. Pl. Rar.* ii. 2. t. 210.—Am. austr.
tuberosum, D. Dietr. Syn. Pl. i. 114 = Peperomia tuberosa.
tumidum, *H. B. & K. Nov. Gen. et Sp.* i. 51.—Ecuador.
turbinatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = longum.
turrialvanum, *C. DC. in Linnaea*, xxxvii. (1871–73) 342.—Costa Rica.
Uhdei, *C. DC. in DC. Prod.* xvi. 1. 248.—Mexic.
ulmifolium, *Steud. Nom.* ed. II. ii. 344.—Ind. occ.
umbellatum, Jacq. Coll. ii. 335 = subpeltatum.
umbellatum, *Linn. Sp. Pl.* 30.—Amphig. trop.
umbellatum, Sieber, ex Kunth, in Linnaea, xiii. (1839) 566 = peltatum.
umbellatum, Vahl, Enum. i. 337 = Dombeyanum.
umbilicatum, Vahl, l. c. i. 356 = Peperomia umbilicata.
umbrosum, Blume, in Verh. Batav. Gen. xi. (1826) 201 = Peperomia nervoso-venosa.
umbrosum, *H. B. & K. Nov. Gen. et Sp.* i. 51.—N. Granat.
undeninervium, *C. DC. in DC. Prod.* xvi. 1. 295.—Bras.
ungaramense, *C. DC. l. c.* 243.—Malaya.
unguiculatum, Poepp. ex Kunth, in Linnaea, xiii. (1839) 603 = laevigatum.
unguiculatum, *Ruiz & Pav. Fl. Per.* i. 34. t. 37. f. B.—Peruv.
urostachyum, *Hemsl. Biol. Centr. Am. Bot.* iii. 57.—Nicaragua.
vaginans, *C. DC. in Linnaea*, xxxvii. (1871–73) 353.—Bras.
Vahlii, D. Dietr. Syn. Pl. i. 111 = Peperomia Vahlii.
variegatum, *H. B. & K. Nov. Gen. et Sp.* i. 47.—N. Granat.
variegatum, Pers. Syn. i. 35 = Peperomia maculosa.
variifolium, *C. DC. in DC. Prod.* xvi. 1. 308.—Mexic.
vasculosum, Wall. Cat. n. 6660 = brachystachyum.
velutinum, *H. B. & K. Nov. Gen. et Sp.* i. 52.—Peruv.
velutinum, Opiz, in Presl, Rel. Haenk. 157 = acutifolium.
venezuelense, *C. DC. in Journ. Bot.* iv. (1866) 216.—Venezuela.
venosum, *C. DC. in DC. Prod.* xvi. 1. 340.—Java.
venustum, Wall. Cat. n. 6666 (Pothos sp.).—Singapore.
veraguense, *C. DC. in DC. Prod.* xvi. 1. 294.—Costa Rica.
verbascifolium, *C. DC. l. c.* 258.—Am. trop.
verrucosum, Sieber, ex Steud. Nom. ed. II. ii. 344 = tuberculatum.
verrucosum, *Sw. Prod. Veg. Ind. Occ.* 15.—Jamaic.
verrucosum, Willd. Sp. Pl. i. 163 = dilatatum.
verruculosum, *C. DC. in Journ. Bot.* iv. (1866) 215.—Costa Rica.
verticillatum, Linn. Syst. ed. X. 856 = Peperomia pulchella.
vestitum, *C. DC. in Journ. Bot.* iv. (1866) 166.—Borneo.
Victorianum, *C. DC. l. c.* 218.—Venezuela.
villosulum, *C. DC. in Linnaea*, xxxvii. (1871–73) 352.—Bras.
villosum, Blume, ex Miq. Fl. Ind. Bat. i. II. 454 = muricatum.
villosum, *C. DC. in Journ. Bot.* iv. (1866) 212.—N. Granat.
villosum, Rich. ex C. DC. in DC. Prod. xvi. 1. 442 = Peperomia rotundata.
viminale, *Opiz, in Presl, Rel. Haenk.* i. 150. t. 26. f. 1.—Ins. Philipp.
Visiani, Hort. ex Buek, Ind. ad DC. Prod. iv. 296 = sphalm. = sylvestre.
Voigtii, *C. DC. in Linnaea*, xxxvii. (1871–72) 365.—Ind. or.
Wagneri, *C. DC. in DC. Prod.* xvi. 1. 302.—Panama.
Walkeri, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 438 = argyrophyllum.
Warakaboura, *C. DC. in DC. Prod.* xvi. 1. 257.—Am. trop.
Weddellii, *C. DC. l. c.* 326.—Bras.
Wightii, Miq. in Hook. Lond. Journ. Bot. v. (1846) 552 = hymenophyllum.
Willdenowii, *Kunth, ex Miq. in Mart. Fl. Bras.* iv. 1. 56.—Bras.
Wulfschlaegelii, *C. DC. in DC. Prod.* xvi. 1. 249.—Jamaic.
Wydlerianum, *C. DC. l. c.* 280.—Porto Rico.
xylopioides, *Kunth, in Linnaea*, xiii. (1839) 610.—Bras.
xylosteoides, *Steud. Nom.* ed. II. ii. 344.—Bras.
Yaguarundi, *Parodi, in Anal. Soc. Cient. Argent.* vi. (1878) 44 ; *Contrib.* 96.—Reg. Argent.
yucatanense, *C. DC. in Linnaea*, xxxvii. (1871–73) 334.—Mexic.
zacuapanum, C. DC. in DC. Prod. xvi. 1. 330 = tiliaefolium.
zeylanicum, *Miq. in Hook. Lond. Journ. Bot.* iv. (1845) 436.—Zeylan.
Zippelia, *C. DC. in DC. Prod.* xvi. 1. 256.—Java.
Zollingeri, *C. DC. l. c.* 364.—Malaya.
Zollingerianum, *C. DC. l. c.* 351.—Java.
Zuccarinii, C. DC. l. c. 365 (Pothos sp. ?).—Ind. or.

PIPERELLA, Presl, Fl. Sic. p. xxxvii (1826) = **Micromeria**, Benth. (Labiat.).
filiformis, Presl, l. c. = M. filiformis.

PIPERI, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = **Piper**, Linn.
betlum, St. Lag. l. c. = Piper Betle.
lonchitum, St. Lag. l. c. = Piper Lonchites.
mullesuanum, St. Lag. l. c. = Piper brachystachyum.
nematantherum, St. Lag. l. c. = Piper Nematanthera.
schizonephum, St. Lag. l. c. = Piper Schizonephos.
scopulosum, St. Lag. = Piper scopulosum.

PIPERIPHORUM, Neck. Elem. iii. 294 (1790) = **Piper**, Linn.

PIPERODENDRUM, Heist. Syst. 12 (1748) (Quid ?).

PIPEROMIA, Pritz. Ic. Ind. i. 217 (1855) sphalm. = seq.

PIPERONIA, Pritz. Ic. Ind. ii. 853 (1865) sphalm. = **Peperomia**, Ruiz & Pav. (Piper.).

PIPTADENIA, Benth. in Hook. Journ. Bot. iv. (1842) 334. *LEGUMINOSAE*, Benth. & Hook. f. i. 589. STACHYCHRYSUM, Boj. Hort. Maurit. 114 (1837).
africana, *Hook. f. in Hook. Niger Fl.* 320.—Afr. trop.
biuncifera, *Benth. in Hook. Journ. Bot.* iv. (1842) 337.—Bras.
Blancheti, *Benth. in Trans. Linn. Soc.* xxx. (1875) 371.—Bras.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 347.—Madag.
boliviana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 370.—Bolivia.
Cebil, *Griseb. in Goett. Abh.* xxiv. (1879) 121.—Reg. Argent.
chrysostachys, *Benth. in Trans. Linn. Soc.* xxx. (1875) 368.—Madag.
colubrina, *Benth. in Hook. Journ. Bot.* iv. (1842) 340.—Bras.
communis, *Benth. l. c.* 337.—Bras.
contorta, *Benth. in Trans. Linn. Soc.* xxx. (1875) 368.—Am. trop.
falcata, *Benth. in Hook. Journ. Bot.* iv. (1842) 341.—Bras.
filicicoma, Benth. l. c. 336 = contorta.
flabellata, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 353.—Madag.
flava, *Benth. in Trans. Linn. Soc.* xxx. (1875) 371.—Am. trop.
foetida, *Benth. l. c.* 366.—Mexic.
foliolosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 336.—Bras.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 347.—Madag.
Greveana, *Baill. l. c.* (1883) 353.—Madag.
guianensis, Benth. in Hook. Journ. Bot. iv. (1842) 335 = Stryphnodendron guianense.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 273.—Afr. trop.
inaequalis, *Benth. in Hook. Journ. Bot.* iv. (1842) 339.—Bras.
latifolia, *Benth. l. c.* 335.—Bras.

PIPTADENIA :—
laxa, *Benth. in Hook. Journ. Bot.* iv. (1842) 335.—Bras.
leptostachya, *Benth. l. c.* 339.—Bras.
macradenia, *Benth. l. c.* 335.—Bras.
macrocarpa, *Benth. l. c.* 341.—Bras.
Mannii, *Oliver, Fl. Trop. Afr.* ii. 329.—Afr. trop.
micracantha, *Benth. in Trans. Linn. Soc.* xxx. (1875) 369.—Bras.
microphylla, Benth. in Hook. Journ. Bot. iv. (1842) 341 = macrocarpa.
moniliformis, *Benth. l. c.* 339.—Bras.
nitida, *Benth. l. c.* 336.—Bras.
oudhensis, *Brand. Forest Fl.* 168.—Ind. or.
paniculata, *Benth. in Hook. Journ. Bot.* iv. (1842) 338.—Bras.
patens, *Benth. Bot. Voy. Sulph.* 89.—Nicarag.
peregrina, *Benth. in Hook. Journ. Bot.* iv. (1842) 340.—Bras.
Pervillei, *Vatke, in Linnaea*, xliii. (1880–82) 109.—Madag.
Poeppigii, *Klotzsch, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 367.—Bras.
polyptera, *Benth. l. c.* 370.—Bras.
polystachya, *Miq. in Linnaea*, xviii. (1844) 590.—Guian.
psilostachya, *Benth. in Hook. Journ. Bot.* iv. (1842) 336.—Guian.
pteroclada, *Benth. in Trans. Linn. Soc.* xxx. (1875) 370.—Peruv.
pterosperma, *Benth. in Hook. Journ. Bot.* iv. (1842) 338.—Bras.
ramosissima, *Benth. l. c.* 336.—Bras.
rigida, *Benth. l. c.* 338.—Bras.
stenadenia, Griseb. Cat. Pl. Cub. 81 = Dichrostachys nutans.
suaveolens, Griseb. Fl. Brit. W. Ind. 710 = flava.
suaveolens, *Miq. in Linnaea*, xviii. (1844) 589.—Ind. occ.
subtilifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 337.—N. Granat.
trisperma, *Benth. l. c.* 337.—Bras.
uaupensis, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 366.—Bras.
viridiflora, Benth. in Hook. Journ. Bot. iv. (1842) 337 = flava.

PIPTANDRA, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 323 = **Scholtzia**, Schau. (Myrtac.).
spathulata, Turcz. l. c. 324 = S. spathulata.

PIPTANTHUS, Sweet, Brit. Flow. Gard. t. 264 (1828). *LEGUMINOSAE*, Benth. & Hook. f. i. 465.
nepalensis, *Sweet, l. c.*—Reg. Himal.

PIPTATHERUM, Beauv. Agrost. 17. t. 5. f. 10, 11 (1812) = **Oryzopsis**, Michx. (Gramin.).
acuminatum, J. & C. Presl, Rel. Haenk. i. 221 = Eriochloa polystachya.
aequiglume, Munro, ex Duthie, in Atkins. Gaz. N. W. Ind. 626.—Ind. or.
angustifolium, Munro, ex Aitch. in Journ. Linn. Soc. xviii. (1880) 106.—Afghanist.
annulatum, Raddi, Agrost. Bras. 30 = Eriochloa polystachya.
Blancheanum, E. Desv. in Boiss. Diagn. Ser. II. iv. 127.—Oriens.
coerulescens, Beauv. Agrost. 18. t. 5 = O. coerulescens.
comosum, Roem. & Schult. Syst. ii. 330 (= *Urachne parviflora*).—Europ. austr. ; Sibir.
confine, Schult. Mant. ii. 184 = Eriochloa polystachya.
elegans, Beauv. Agrost. 173, nomen.—Hab. ?
flavescens, Sieber, ex Nees, Agrost. Bras. 16 = Eriochloa polystachya.
frutescens, Schult. Mant. ii. 183 (= *Urachne parviflora*).—Europ. austr. ; Sibir.
holciforme, Roem. & Schult. Syst. ii. 328 = O. holciformis.
laeve, [Nees, in] Meyen, Reise, i. 466, 484, nomen (= *Helopus Meyeni*).—Peruv.
laterale, Munro, ex Aitch. in Journ. Linn. Soc. xviii. (1880) 106 = *molinioides*.
Lindleyanum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 149.—Chili.
longearistatum, Boiss. & Haussk. ex Boiss. Fl. Orient. v. 509.—Persia.

PIPTATHERUM:—
macratherum, Nees, ex Steud. Syn. Pl. Gram. 123 (= *Urachne parviflora*).—Europ.
miliaceum, Coss. Notes Crit. 129 = *multiflorum*.
miliaceum, Fourn. in Bull. Soc. Bot. Fr. xiii. (1866) 119 (an praec. ?).—Reg. Mediterr.
molinioides, Boiss. Diagn. Ser. I. vii. 121.—Persia; Afghanist.
multiflorum, Beauv. Agrost. 18 (= *Urachne parviflora*).—Europ.; Sibir.
nigrum, Torr. Fl. U. St. i. 79 = Oryzopsis melanocarpa.
obtusum, [Nees, in] Meyen, Reise, ii. 32 (= *Helopus obtusus*).—Peruv.
paradoxum, Beauv. Agrost. 18 = O. paradoxa.
punctatum, Beauv. l. c. = Eriochloa polystachya.
ramosum, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 148 = *laeve*.
setosum, A. Rich. Fl. Cub. Fanerog. ii. 311.—Cuba.
soongaricum, Duthie, in Atkins. Gaz. N. W. Ind. 626.—Ind. or.
sphacelatum, Boiss. Fl. Orient. v. 508.—Persia.
Thomasii, Kunth, Enum. Pl. i. 177 (= *Urachne parviflora*).—Europ.; Sibir.
virescens, Boiss. Fl. Orient. v. 507.—Europ. or.; As. Min.

PIPTOCALYX, Oliver, ex Benth. Fl. Austral. v. 292 (1870). *MONIMIACEAE*, Benth. & Hook. f. iii. 143.
Moorei, *Oliver, l. c.*—Austral.

PIPTOCALYX, Torr. Bot. U. St. Expl. Exped. ii. 413. t. 12 (1874) = **Krynitzkia**, Fisch. & Mey. (Borag.).
circumscissus, Torr. l. c. 414. t. 12 = K. circumscissa.

PIPTOCARPHA, R. Br. in Trans. Linn. Soc. xii. (1816) 121. *COMPOSITAE*, Benth. & Hook. f. ii. 231.
CARPHOBOLUS, Schott, in Spreng. Syst. iv. Cur. Post. 409 (1827).
MONANTHEMUM, Griseb. Fl. Brit. W. Ind. 354 (1861).
VANILLOSMA, Spach, Hist. Vég. Phan. x. 39 (1841); xiv. 424 (1848).
asterotricha, *Baker, in Mart. Fl. Bras.* vi. II. 126, in obs.—Peruv.
axillaris, *Baker, l. c.* 122.—Bras.
brasiliana, *Cass. in Dict. Sc. Nat.* xli. 109.—Bras.
Brownii, DC. Prod. vii. 261 = brasiliana.
chontalensis, *Baker, in Mart. Fl. Bras.* vi. II. 132.—Nicaragua.
cinerea, *Baker, l. c.* 129.—Bras.
Cruegeri, *Baker, l. c.* 132.—Ins. Trinit.
elaeagnoides, *Baker, l. c.* 126.—N. Granat.
Lechleri, *Baker, l. c.* 127, in obs.—Peruv.
leprosa, *Baker, l. c.* 128.—Bras.
lucida, *Benn. ex Baker, l. c.* 121.—Bras.
Lundiana, *Baker, l. c.* 130.—Bras.
macropoda, *Baker, l. c.* 123.—Bras.
notata, *Baker, l. c.* 129.—Bras.
oblonga, *Baker, l. c.* 121.—Bras.
opaca, *Baker, l. c.* 124.—Bras.
oxyphylla, *Baker, l. c.* 120.—Bras.
pannosa, *Baker, in Journ. Bot.* xiii. (1875) 203.—Bras.
pellucida, *Baker, in Mart. Fl. Bras.* vi. II. 127.—Bras.
Poeppigiana, *Baker, l. c.* 131.—Peruv.
polycephala, *Baker, l. c.* 125.—Guian.
pyrifolia, *Baker, l. c.* 120.—Bras.
quadrangularis, *Baker, l. c.* 128.—Bras.
ramiflora, *Baker, l. c.* 127.—Bras.
Riedelii, *Baker, l. c.* 124.—Bras.
rotundifolia, *Baker, l. c.* 125.—Bras.
Sellowii, *Baker, l. c.* 130.—Bras.
senescens, *Baker, l. c.*—Bras.
Sprucei, *Baker, l. c.* 129, in obs.—Peruv.
tereticaulis, *Baker, l. c.* 132.—N. Granat.
tomentosa, *Baker, in Journ. Bot.* xiii. (1875) 203.—Bras.
triflora, *Benn. ex Baker, in Mart. Fl. Bras.* vi. II. 132.—Guian.
umbellulata, *Baker, l. c.* 126.—Bras.
Vauthieriana, *Baker, l. c.*—Bras.
venulosa, *Baker, l. c.* 131.—Bras.

PIPTOCARPHA, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 110 = **Chuquiraga**, Juss. (Compos.).
diacanthoides, Hook. & Arn. l. c. = C. leucoxylon.
excelsa, Hook. & Arn. l. c. = C. excelsa.

PIPTOCELUS, Presl, Bot. Bemerk. 36 (1844); in Abh. Boehm. Ges. V. iii. (1845) 466. Gen. dub. *CELASTRINEARUM*.—Cf. Benth. & Hook. f. Gen. i. 360.
Haenkeanus, *Presl, ll. cc.* 36; 466.—Ecuador.
myrianthus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 449 = Schinus dependens.

PIPTOCEPHALUM, Sch. Bip. in Bonplandia, viii. (1860) 369 = **Catananche**, Linn. (Compos.).
carpholepis, Sch. Bip. l. c. 369, 370 = C. lutea.
luteum, Sch. Bip. l. c. = C. lutea.
phaiolepis, Ball, in Journ. Linn. Soc. xvi. (1878) 534 = C. lutea.

PIPTOCERAS, Cass. in Dict. Sc. Nat. l. 469 (1828) = **Centaurea**, Linn. (Compos.).
babylonicum, Cass. l. c. liv. 489 = C. babylonica.
Behen, Cass. l. c. 487 = C. Behen.

PIPTOCHAETIUM, J. & C. Presl, Rel. Haenk. i. 222. t. 37 (1830) = **Oryzopsis**, Michx. (Gramin.).
bicolor, E. Desv. in C. Gay, Fl. Chil. vi. 273.—Chili.
chaetaphorum, Griseb. in Goett. Abh. xxiv. (1879) 298.—Reg. Argent.
erianthum, Bal. in Bull. Soc. Bot. Fr. xxxii. (1885) 244.—Reg. Argent.
hirtum, Phil. in Anal. Univ. Chil. (1873) 559.—Chili.
laevissimum, Phil. in Linnaea, xxxiii. (1864–65) 280.—Chili.
lasianthum, Griseb. in Goett. Abh. xxiv. (1879) 297.—Reg. Argent.
mucronatum, Griseb. l. c. 296.—Reg. Argent.
ovatum, E. Desv. in C. Gay, Fl. Chil. vi. 273.—Chili.
pallidum, Phil. ex Griseb. in Goett. Abh. xxiv. (1879) 297 = *laevissimum*.
panicoides, E. Desv. in C. Gay, Fl. Chil. vi. 270.—Chili.
purpuratum, Phil. in Linnaea, xxix. (1857–58) 86.—Chili.
setifolium, J. & C. Presl, Rel. Haenk. i. 222 (= *Urachne panicoides*).—Chili; Bras.
trichotomum, Griseb. in Goett. Abh. xxiv. (1879) 297 (= *Urachne trichotomum*).—Reg. Argent.
tuberculatum, E. Desv. in C. Gay, Fl. Chil. vi. 272.—Chili.
uruguense, Griseb. in Goett. Abh. xxiv. (1879) 297.—Reg. Argent.
verrucosum, Phil. in Linnaea, xxxiii. (1864–65) 280.—Chili.

PIPTOCHLAMYS, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 358 = **Thymelaea**, Endl.
hirsuta, C. A. Mey. l. c. = T. hirsuta.

PIPTOCLAINA, G. Don, Gen. Syst. iv. 364 (1837) = **Heliotropium**, Tourn. (Borag.).
malabarica, G. Don, l. c. = H. supinum.
supina, G. Don, l. c. = H. supinum.

PIPTOCOMA, Cass. in Bull. Soc. Philom. (1817) 10; et in Dict. Sc. Nat. xli. 111 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 233.
rufescens, *Cass. l. c.* 10; (1818) 58; *et in Dict. Sc. Nat.* xli. 111.—Ins. S. Doming.

PIPTOCOMA, Less. in Linnaea, iv. (1829) 315 = **Lychnophora**, Mart. (Compos.).
lychnophorioides, Less. l. c. = L. trichocarpha.

PIPTOLAENA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 25 = **Voacanga**, Thou. (Apocyn.).
Dregei, A. DC. in DC. Prod. viii. 358 = V. Dregei.

PIPTOLEPIS, Benth. Pl. Hartw. 29 (1840) = **Forestiera**, Poir. (Oleac.).
phillyreoides, Benth. l. c. = F. phillyreoides.

PIPTOLEPIS, Sch. Bip. in Pollichia, xx-xxi. (1863) 380. *COMPOSITAE*, Benth. & Hook. f. ii. 233.
buxoides, *Sch. Bip. l. c.* 383.—Bras.
ericoides, *Sch. Bip. l. c.*—Bras.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. II. 144.—Bras.

PIPTOLEPIS:—
imbricata, *Sch. Bip. in Pollichia*, xx-xxi. (1863) 383.—Bras.
leptospermoides, *Sch. Bip. l. c.* 382.—Bras.
Martiana, *Sch. Bip. l. c.* 385.—Bras.
Oleaster, *Sch. Bip. l. c.* 384.—Bras.
Pseudo-myrtus, *Sch. Bip. l. c.*—Bras.

PIPTOMERIS, Turcz. in Bull. Soc. Nat. Mosc. xxvi. I. (1853) 257 = **Jacksonia**, R. Br. (Legumin.).
aphylla, Turcz. l. c. 258 = J. Piptomeris.

PIPTOPOGON, Cass. in Dict. Sc. Nat. xlviii. 434, 507 (1827) = **Hypochoeris**, Linn. (Compos.).
decipiens, Cass. l. c. = H. Achyrophorus.
glaucum, Sch. Bip. in Nov. Act. Nat. Cur. xxi. I. (1845) 161 = H. Achyrophorus.
laevigatum, Sch. Bip. l. c. 165 = H. Achyrophorus.
macrospermum, C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 95 = Scorzonera macrosperma.

PIPTOPTERA, Bunge, in Act. Hort. Petrop. v. (1877) 644. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 75.
turkestana, *Bunge, l. c.* 645; vi. (1879) 449.—Turkestan.

PIPTOSACCOS, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 415. *INC. SEDIS*.
hypophyllantha, *Turcz. l. c.*—Malaya.

PIPTOSPATHA, N. E. Br. in Gard. Chron. (1879) I. 138. f. 20. *AROIDEAE*, Benth. & Hook. f. iii. 985.
insignis, *N. E. Br. l. c.*—Borneo.

PIPTOSTEGIA, Reichb. Nom. 113 (1841) = **Ipomoea**, Linn. (Convolv.).
Gomesii, Mart. Syst. Mat. Med. Bras. 78.—Bras.
operculata, Reichb. Nom. 113 = I. operculata.
Pisonis, Mart. Syst. Mat. Med. Bras. 78.—Bras.

PIPTOSTEMMA, Spach, Hist. Vég. Phan. x. 34 (1841) = **Nassauvia**, Comm. (Compos.).

PIPTOSTEMMA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 191 = **Angianthus**, Wendl. (Compos.).
carpesioides, Turcz. l. c, 192 = A. phyllocephalus.

PIPTOSTEMUM, Steud. Nom. ed. II. ii. 344 (1841) = PIPTOSTEMMA, Spach = **Nassauvia**, Comm. (Compos.).

PIPTOSTIGMA, Oliver, in Journ. Linn. Soc. viii. (1865) 158. t. 2. *ANONACEAE*, Benth. & Hook. f. i. 957.
glabrescens, *Oliver, l. c.* 159.—Afr. trop.
pilosum, *Oliver, l. c.*—Afr. trop.

PIPTOSTYLIS, Dalz. in Hook. Kew Journ. iii. (1851) 33. t. 2 = **Clausena**, Burm. f. (Rutac.).
indica, Dalz. l. c. = C. indica.

PIPTURUS, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 196. *URTICACEAE*, Benth. & Hook. f. iii. 390.
NOTHOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 137, in nota, t. 14 [1856].
PRAETORIA, Baill. Étud. Gén. Euph. 470. t. 11 (1858).
acuminatus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 304.—Malaya.
albidus, A. Gray, ex H. Mann, in Proc. Am. Acad. vii. (1867) 201 = Gaudichaudianus.
argenteus, *Wedd. in DC. Prod.* xvi. I. 235^{19}.—Java.
asper, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 197.—Ins. Philipp.
Candolleanus, Wedd. l. c. 196 = argenteus.
ceramicus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 304.—Ins. Ceram.
ellipticus, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 197.—Ins. Philipp.
Gaudichaudianus, *Wedd. l. c.*—Ins. Sandvic.
gracilipes, *A. Gray, ex Wedd. in DC. Prod.* xvi. I. 235^{30}.—Ins. Sandvic.
incanus, Wedd. in DC. l. c. 235^{18} = velutinus.

PIPTURUS:—
integrifolia, *Baker, in Journ. Bot.* xx. (1882) 267.—Madag.
mollissimus, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 449.—Malaya.
platyphyllus, *Wedd. in DC. Prod.* xvi. i. 235[19].—Ins. Pacif.
propinquus, Wedd. in Arch. Mus. Par. viii. (1855–56) 447. t. 15 D = argenteus.
repandus, *Wedd. l. c.* 448.—Java.
rotundifolius, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 197.—Ins. Maurit.
taitensis, Wedd. l. c. = Gaudichaudianus.
velutinus, *Wedd. l. c.* 196.—Ins. Malay. Mascar. Pacif.

PIQUERIA, Cav. Ic. iii. 18. t. 235 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 238.
PHALACRAEA, DC. Prod. v. 105 (1836).
ageratoides, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 430 = Alomia ageratoides.
angustata, Gardn. l. c. 432 = Alomia angustata.
artemisioides, *H. B. & K. Nov. Gen. et Sp.* iv. 153.—Ecuador.
attenuata, Gard. in Hook. Lond. Journ. Bot. vi. (1847) 430 = Gymnocoronis spilanthoides.
cinerea, Gardn. l. c. 432 = Alomia cinerea.
densiflora, *Benth. Bot. Voy. Sulph.* 110.—Ecuador.
eupatorioides, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 431 = Trichogonia salviaefolia.
Eupatorium, Gardn. l. c. 430 = Clibadium rotundifolium.
floribunda, *DC. Prod.* v. 105.—Peruv.; Chili.
fastigiata, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 431 = Alomia fastigiata.
foliosa, Gardn. l. c. 432 = Alomia foliosa.
galioides, *DC. Prod.* v. 105.—Peruv.
latifolia, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 430 = Ageratum conyzoides.
longipetiolata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 183 = Gymnocoronis spilanthoides.
myriadenia, Sch. Bip. ex Baker, l. c. 192 = Alomia myriadenia.
ovata, G. Don, ex Loud. Hort. Brit. 337 = trinervia.
pilosa, *H. B. & K. Nov. Gen. et Sp.* iv. 153.—Mexic.
polyphylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 192 = Alomia polyphylla.
pubescens, *Sm. in Rees, Cycl.* xxvii. n. 2.—Peruv.
quinqueflora, *Cass. in Dict. Sc. Nat.* xli. 116.—Peruv.
serrata, *A. Gray, in Proc. Am. Acad.* xv. (1880) 25.—Mexic.
subcordata, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 430 = Gymnocoronis spilanthoides.
triflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 77.—Mexic.
trinervia, *Cav. Ic.* iii. 19. t. 235.—Mexic.

PIRANHEA, Baill. Adansonia, vi. (1866) 235. t. 6. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 281.
trifoliata, *Baill. l. c.* 236. t. 6.—Bras.

PIRARDA, Adans. Fam. ii. 499 (1763) = **Ethulia**, Linn. (Compos.).

PIRATINERA, Aubl. Pl. Gui. ii. 888. t. 340 (1775) = **Brosimum**, Sw. (Urtic.).
guianensis, Aubl. l. c. = B. Aubletii.

PIRCUNIA, Bert. in Merc. Chil. (1829); ex Am. Journ. Sc. Ser. I. xxiii. (1833) 264 = **Phytolacca**, Linn.
abyssinica, Moq. in DC. Prod. xiii. II. 30 = Phyt. abyssinica.
chilensis, Moq. l. c. 29 = Phyt. chilensis.
dioica, Moq. l. c. 30 = Phyt. dioica.
drastica, Bert. in 'Merc. Chil. (1829)'; ex Am. Journ. Sc. Ser. I. xxiii. (1833) 264 = Anisomeria drastica.
esculenta, Moq. in Fl. des Serres, ix. (1853–54) 236 = Phyt. acinosa.
Latbenia, Moq. in DC. Prod. xiii. II. 29 = P. acinosa.
stricta, Moq. l. c. 30 = Phyt. stricta.

PIRENIA, C. Koch, Dendrol. i. 188 (1867) = PYRENIA, Clairv. = **Pyrus**, Tourn. (Rosac.).

PIRIGARA, Aubl. Pl. Gui. i. 487 (1775) = **Gustavia**, Linn. (Myrtac.).
alata, Span. in Linnaea, xv. (1841) 204 = G. alata.
globosa, Span. l. c. = G. globosa.

PIRIGARA:—
hexapetala, Aubl. Pl. Gui. i. 490 = Gustavia fastuosa.
insignis, Kunth, ex Hemsl. Biol. Centr. Am. Bot. i. 413 = G. superba.
speciosa, H. B. & K. Nov. Gen. et Sp. vii. 262 = G. latifolia.
superba, H. B. & K. l. c. 261 = G. superba.
tetrapetala, Aubl. Pl. Gui. i. 487 = Grias Aubletiana.
valida, Blume, Bijdr. 1096 = Planchonia valida.

PIRIGARDA, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 511 (1879), sphalm. = praec.

PIRINGA, Juss. in Mém. Mus. Par. vi. (1820) 399 = **Gardenia**, Ellis (Rubiac.).

PIRIPEA, Aubl. Pl. Gui. ii. 627. t. 253 (1775) = **Buchnera**, Linn. (Scrophul.).
palustris, Aubl. l. c. 628. t. 253 = B. palustris.

PIRIQUETA, Aubl. Pl. Gui. i. 298. t. 117 (1775). *TURNERACEAE* [Benth. & Hook. f. i. 807].
BURCARDIA, Schreb. Gen. i. 530 (1789).
BURGHARTIA, Scop. Introd. 229 (1777).
ERBLICHIA, Seem. Bot. Voy. Herald, 130 (1853).
assurensis, *Urban, in Eichl. Jahrb.* ii. (1883) 60.—Bras.
aurea, *Urban, l. c.* 69.—Bras.
Berneriana, *Urban, l. c.* 78.—Madag.
capensis, *Urban, l. c.*—Afr. austr.
caroliniana, Urban, l. c. 71 = fulva.
cistoides, G. F. W. Mey. ex Steud. Nom. ed. II. ii. 344 = villosa.
Duarteana, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 66.—Bras.
foliosa, Garcke, in Linnaea, xxii. (1849) 63 = villosa.
fulva, *Chapm. Fl. S. U. St.* 146.—Am. bor.
glabra, Chapm. l. c. 147 = fulva.
ionidioides, A. Rich. Ess. Fl. Cub. 605 = villosa.
lanceolata, Benth. in Hook. Journ. Bot. iv. (1842) 117 = villosa.
longifolia, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1880) 275 = villosa.
madagascariensis, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 79.—Madag.
nitida, *Urban, l. c.* 70.—Bras.
odorata, *Urban, l. c.* 80.—Am. centr.
ovata, *Urban, l. c.* 76.—Ins. S. Doming.
plicata, *Urban, l. c.* 65.—Bras.
racemosa, *Sweet, Hort. Brit.* ed. I. 154.—Bras.
rosea, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 64.—Bras.
Selloi, *Urban, l. c.* 63.—Bras.
sidifolia, *Urban, l. c.* 61.—Bras.
stenophylla, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1167, nomen = villosa.
sulfurea, *Urban & Rolfe, ex Urban, in Eichl. Jahrb. Berl.* ii. (1883) 62.—Bras.
Tamberlikii, *Urban, l. c.* 67.—Bras.
tomentosa, Chapm. Fl. S. U. St. 147 = fulva.
tomentosa, H. B. & K. Nov. Gen. et Sp. vi. 128 = villosa.
villosa, *Aubl. Pl. Gui.* i. 298. t. 117.—Guian.
viscosa, *Griseb. Cat. Pl. Cub.* 114.—Cuba; Bras.

PIRITANERA, R. H. Schomb. in Proc. Linn. Soc. ii. (1851) 161 = PIRATINERA, Aubl. = **Brosimum**, Sw. (Urtic.).

PIROLA, Neck. in Act. Acad. Theod. Palat. ii. (1770) 488 = **Pyrola**, Tourn. (Ericac.).

PIRONNEAVA, Gaudich. Bot. Voy. Bonite, tt. 63, 64 (1844–52) = **Aechmea**, Ruiz & Pav. (Bromel.).
glomerata, Gaudich. l. c. t. 63 = A. augusta.
Lueddemanniana, C. Koch, Wochenschr. ix. (1866) 182 = A. caerulescens.
Morreniana, Regel, Gartenfl. (1874) 257 = A. glomerata.
platynema, Gaudich. Bot. Voy. Bonite, t. 64 = A. platynema.
ramosa, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 148 = A. ramosa.
roseo-coerulea, C. Koch, Wochenschr. iv. (1861) 189 = A. glomerata.
spectabilis, C. Koch, l. c. iii. (1860) 338 = A. spectabilis.

PIROPHORUM, Neck. Elem. ii. 72 (1790) = **Pyrus**, Tourn. (Rosac.).

PIRRONEANA, Benth. & Hook. f. Gen. iii. 663 (1883), sphalm. = PIRONNEAVA, Gaudich. = **Aechmea**, Ruiz & Pav. (Bromel.).

PIRUS, Hall, Enum. Stirp. Helv. i. 351 (1742) = **Pyrus**, Tourn. (Rosac.).

PISAURA, Bonato, Monog. (1793); ex Endl. Gen. 1194 (1841) = **Lopezia**, Cav. (Onagr.).
automorpha, Bonato, l. c. = L. racemosa.

PISCIDIA, Linn. Syst. ed. X. 1155 (1759). *LEGUMINOSAE*, Benth. & Hook. f. i. 550.
acutata, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 106 = Pterocarpus ancylocalyx.
carthagenensis, Jacq. Enum. Pl. Carib. 27 = P. Erythrina.
Corallodendrum, Steud. Nom. ed. II. ii. 344 = Erythrina Corallodendron?
Erythrina, *Linn. Syst.* ed. X. 1155.—Am. trop.
florida, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 106 = Pterocarpus Rohrii.
inebrians, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 394 = P. Erythrina.
longifolia, *Willd. Sp. Pl.* iii. 920.—Mexic.
punicea, Cav. Ic. iv. 8 = Sesbania punicea.
toxicaria, Salisb. Prod. 336 = P. Erythrina.

PISONIA, Plum. ex Linn. Gen. ed. I. 42 (1737). *NYCTAGINEAE*, Benth. & Hook. f. iii. 9.
BESSERA, Vell. Fl. Flum. 147 (1825); iv. t. 12 (1827).
CALPIDIA, Thou. Hist. Vég. Isles Afr. Austr. (1804) 37. t. 10.
CEODES, Forst. Char. Gen. 141. t. 71 (1776).
COLUMELLA, Vell. Fl. Flum. 155 (1827); iv. t. 17 (1827).
PALLAVIA, Vell. l. c. 151 (1825); iv. t. 12 (1827).
TORRUBIA, Vell. l. c. 139 (1825); iii. t. 150 (1827).
TRAGULARIA, Koen. ex Roxb. Fl. Ind. ii. 217 (1832).
aculeata, *Linn. Sp. Pl.* 1026.—Cosmop. trop.
acuminata, *Mart. ex F. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 357.—Bras.
alba, *Span. in Linnaea*, xv. (1841) 342.—Ind. or.; Malaya.
anisophylla, *Hassk. Hort. Bog. Desc.* 85.—Malaya.
Brunoniana, A. Cunn. ex Choisy, in DC. Prod. xiii. II. 441 = excelsa.
Brunoniana, *Endl. Prod. Fl. Norf.* 43.—Austral.; N. Zel.; Ins. Norf.
buxifolia, Rottb. in Nye Samml. Danske Vidensk. Selsk. Skr. ii. (1783) 536. t. 4. f. 2 = Maba buxifolia.
Cafferana, *Casar. Nov. Stirp. Bras. Dec.* 68.—Bras.
Calpidia, *Steud. Nom.* ed. II. ii. 345.—Ins. Maurit.
campestris, *Netto, in Ann. Sc. Nat.* Sér. V. v. (1866) 83.—Bras.
Capparosa, Netto, l. c. 82 = Neea theifera.
cauliflora, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 417.—Malaya.
coccinea, Sw. Prod. Veg. Ind. occ. 60 = inermis.
combretiflora, *Mart. ex F. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 360.—Bras.
comosa, *Choisy, in DC. Prod.* xiii. II. 444.—Bras.
cordifolia, *Mart. ex F. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 359.—Bras.
costata, *Choisy, in DC. Prod.* xiii. II. 446.—Ins. Maurit.
crenulata, *F. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 356.—Bras.
cuneifolia, Schlecht. in Linnaea, xxiii. (1850) 571 = obtusata.
Darwinii, *Hemsl. Bot. Voy. Challenger*, i. II. 20.—Ins. Fernand.-Noronh.
discolor, *Spreng. Syst.* ii. 168.—Jamaic.
elliptica, *F. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 361.—Bras.
eucalyptifolia, H. B. & K. Nov. Gen. et Sp. vii. 197 = inermis.
excelsa, *Blume, Bijdr.* 735.—Malaya.
ferruginea, *Klotzsch, ex Choisy, in DC. Prod.* xiii. II. 445.—N. Granat.
floribunda, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 193.—Ins. Galap.
florida, *Choisy, in DC. Prod.* xiii. II. 443.—Bras.
Forsteriana, Endl. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 403 = grandis.

PISONIA:—
fragrans, Desf. Cat. Hort. Par. ed. III. 390 = obtusata ?
fragrans, *Dum.-Cours. Bot. Cult.* ed. II. vii. 114.—Hab. ?
fragrans, *Link, Enum. Hort. Berol.* i. 353.—Hab. ?
Georgina, Wall. Cat. n. 6768 = aculeata.
glaucescens, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. II. 360 = campestris.
graciliflora, *Mart. ex J. A. Schmidt, l. c.* 358.—Bras.
grandis, *R. Br. Prod.* 422.—Austral. ; Ins. Pacif.
grandis, A. Cunn. ex Hook. f. Fl. N. Zel. i. 209 = Brunoniana.
guianensis, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1131, nomen.—Guian.
heterophylla, *Choisy, in DC. Prod.* xiii. II. 444.—Bras.
hirsuta, *Choisy, l. c.* 445.—Bras.
hirtella, *H. B. & K. Nov. Gen. et Sp.* ii. 217.—Mexic.
inermis, Forst. f. Prod. 75 = grandis.
inermis, *Jacq. Select. Am.* 275.—Am. trop.
lanceolata, Choisy, in DC. Prod. xiii. II. 442 = P. Calpidia.
latifolia, *Desf. Cat. Hort. Par.* ed. III. 391.—Hab. ?
laurifolia, *J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 364.—Bras.
laxa, *Netto, in Ann. Sc. Nat.* Sér. V. v. (1866) 84.—Bras.
laxiflora, *Choisy, in DC. Prod.* xiii. II. 444.—Bras.
limonella, Blume, Bijdr. 735 = aculeata.
longirostris, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 40.—Malaya.
loranthoides, H. B. & K. Nov. Gen. et Sp. vii. 197 = aculeata.
macrocarpa, Presl, Symb. Bot. ii. 8. t. 56 = excelsa.
macrophylla, *Choisy, in DC. Prod.* xiii. II. 446.—Ins. Mascar.
macrophylla, *Link, Enum. Hort. Berol.* i. 354.—Hab. ?
major, *Baill. Adansonia*, x. (1872) 185.—Ins. Rawak.
malabarica, Choisy, in DC. Prod. xiii. II. 447 = alba ?
mexicana, Willd. Enum. Hort. Berol. Suppl. 20 = hirtella.
minor, *Choisy, in DC. Prod.* xiii. II. 443.—Bras.
mitis, Herb. Banks, ex Griseb. Fl. Brit. W. Ind. 71 = inermis.
mitis, *Linn. Sp. Pl.* 1026.—Ind. or.
mollis, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 136.—Guian.
monotaxadenia, *Wright, in Sauv. Fl. Cub.* 137.—Cuba.
Mooriana, F. Muell. Fragm. i. 20 = Brunoniana.
morindifolia, R. Br. in Wall. Cat. n. 7130 = alba.
myrtifrons, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. II. 360 = campestris.
nigricans, Choisy, in DC. Prod. xiii. II. 442 = inermis.
nigricans, *Sw. Prod. Veg. Ind. Occ.* 60.—Am. trop.
nitida, *Desf. Cat. Hort. Par.* ed. III. 390.—Hab. ?
nitida, *Mart. ex J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 356.—Bras.
nitida, *Willd. Enum. Hort. Berol. Suppl.* 20, nomen.—Madag.
noxia, Netto, in Ann. Sc. Nat. Sér. V. v. (1866) 80 = tomentosa.
obovata, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1075, nomen.—Guian.
obovata, *Link, Enum. Hort. Berol.* i. 354.—Hab. ?
obtusata, *Jacq. Hort. Schoenb.* iii. 35.—Ind. occ.
Olfersiana, *Link, Klotzsch & Otto, Ic. Pl. Rar.* 37. t. 15.—Bras.
olitoria, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 154.—Malaya.
ovalifolia, Choisy, in DC. Prod. xiii. II. 441 = P. Calpidia.
Pacurero, H. B. & K. Nov. Gen. et Sp. ii. 218 = inermis.
palicureoides, Casar. Nov. Stirp. Bras. Dec. 68 = Olfersiana.
Pallavia, Steud. Nom. ed. II. ii. 345 = aculeata.
pedicellaris, Griseb. ex Sauv. Fl. Cub. 137 = aculeata.
pernambucensis, Casar. Nov. Stirp. Bras. Dec. 69 = subcordata.
procera, Bert. ex Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 191 = Brunoniana.
psammophila, *Mart. ex J. A. Schmidt, in Mart. Fl. Bras.* xiv. II. 362.—Bras.
pubescens, *H. B. & K. Nov. Gen. et Sp.* ii. 218.—Venezuela.

PISONIA:—
racemosa, Wall. Cat. n. 7296 = Petunga longifolia.
rosea, *Linden, Cat.* n. 17 (1862) 17 ; *Teijsm. & Binn. Cat. Hort. Bog.* 92.—Hab. ?
rotundata, *Griseb. Cat. Pl. Cub.* 283.—Cuba.
Sinclairii, Hook. f. Fl. N. Zel. i. 209. t. 50 = Brunoniana.
subcordata, *Sw. Prod. Veg. Ind. Occ.* 60.—Ind. occ.
subferruginosa, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. xiv. II. 358, 411 = graciliflora.
subovata, Poir. Encyc. v. 347 = subcordata.
sylvestris. *Teijsm & Binn. in Tijdschr. Nederl. Ind.* ix. (1855) 349, 355 ; *et ex Hassk. Retzia*, 6.—Java.
tomentosa, *Casar. Nov. Stirp. Bras. Dec.* 69.—Bras.
tomentosa, Walp. Ann. i. 561, sphalm. = Plumbago tomentosa.
umbellifera, Seem. ex Nadeaud, Enum. Pl. Tahiti, 46 = excelsa.
uniseriata, Wright, ex Sauv. Fl. Cub. 138 = monotaxadenia.
venosa, *Choisy. in DC. Prod.* xiii. II. 444.—Bras.
villosa, Poir. Encyc. v. 347 = aculeata.
viscosa, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 19 ; *et in Baker, Fl. Maurit.* 263.—Ins. Rodriguez.
viscosa, *Seem. in Bonplandia*, ix. (1861) 258.—Ins. Fiji.
Yagua-pinda, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 211 ; *Contrib.* 61.—Paraguay.
Zapallo, *Griseb. in Goett. Abh.* xxiv. (1879) 39.—Reg. Argent.

PISOSPERMA, Sond. in Harv. & Sond. Fl. Cap. ii. 498 (1861–62). *CUCURBITACEAE*, Benth. & Hook. f. i. 831.
capense, *Sond. l. c.*—Afr. austr.

PISTACHIA, Salisb. Prod. 171 (1796) = seq.

PISTACIA, Linn. Hort. Cliff. 456 (1737). *ANACARDIACEAE*, Benth. & Hook. f. i. 419.
LENTISCUS, [Tourn.] Linn. Syst. ed. I (1735).
TEREBINTHUS, P. Br. Hist. Jamaic. 345 (1735).
acuminata, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 53 = P. Khinjuk.
americana, Mill. Gard. Dict. ed. VIII. n. 7 = Simaruba glauca ?
atlantica, DC. Prod. ii. 64 = mutica.
atlantica, *Desf. Fl. Atlant.* ii. 364.—Reg. Mediterr.
cabulica, Stocks, in Hook. Kew Journ. iv. (1852) 143 = mutica.
cappadocica ×, Planch. in Bull. Soc. Bot. Fr. xi. (1864) p. xlvi.—Europ.
chia, Desf. Tabl. ed. I. 199 = mutica.
chinensis, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 89.—China.
crassifolia, Salisb. Prod. 172 = P. Terebinthus.
fagaroides, *Willd. Enum. Hort. Berol. Suppl.* 66.—Hab. ?
gummifera, Salisb. Prod. 172 = P. Lentiscus.
hybrida ×, Gasparr. ex Parl. Fl. Ital. v. 383.—Europ.
integerrima, Stew. ex Brand. For. Fl. Ind. 122 = P. Khinjuk.
Khinjuk, *Stocks, in Hook. Kew Journ.* iv. (1852) 143.—Aegypt. ; Persia ; Reg. Himal.
Lentiscus, *Linn. Sp. Pl.* 1026.—Reg. Mediterr.
macrophylla, Pers. Syn. ii. 615 = vera.
massiliensis, Mill. Gard. Dict. ed. VIII. n. 6 = P. Lentiscus.
mexicana, *H. B. & K. Nov. Gen. et Sp.* vii. 22. t. 608.—Mexic.
mutica, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 338.—Reg. Mediterr. or. ; Persia ; Afghan.
mutica, Raul. ex Boiss. Fl. Orient. ii. 6 = P. Terebinthus.
narbonnensis ×, Linn. Sp. Pl. 1025.—Gall. austr.
nemausensis, Req. ex Planch. in Bull. Soc. Bot. Fr. xi. (1864) p. xlvi. = praec.
nigricans, Crantz, Inst. i. 184 = vera.
occidentalis, Baill. Adansonia, xi. (1874) 181 = Dacryodes hexandra.
officinarum, Ait. Hort. Kew. ed. I. iii. 398 = vera.
oleosa, *Lour. Fl. Cochinch*, 615.—Cochinch.
palaestina, Boiss. Diagn. Ser. I. ix. 1 = P. Terebinthus.
reticulata, Willd. Sp. Pl. iv. 751 = vera.
Simaruba, Linn. Sp. Pl. 1026 = Bursera gummifera.
terebinthina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Terebinthus.

PISTACIA:—
Terebinthus, Bieb. Fl. Taur. Cauc. ii. 418 = mutica.
Terebinthus, *Linn. Sp. Pl.* 1025.—Europ. austr. ; Reg. Mediterr.
Terebinthus, Mill. Gard. Dict. ed. VIII. n. 1 = vera.
Therebinthus, Scop. Fl. Carn. ed. II. ii. 262 = P. Terebinthus.
trifolia, Linn. Sp. Pl. 1025 = vera.
variifolia, Salisb. Prod. 171 = vera.
vera, *Linn. Sp. Pl.* 1025.—Reg. Mediterr. ; Oriens.
vera, Mill. Gard. Dict. ed. VIII. n. 4 = P. Terebinthus.
Vitex, Steud. Nom. ed. I. 627 = Vitex pubescens.

PISTIA, Linn. Gen. ed. I. 275 (1737). *AROIDEAE*, Benth. & Hook. f. iii. 964.
APIOSPERMUM, Klotzsch, in Abh. Akad. Berl. (1853) 351.
KODDA-PAIL, Adans. Fam. ii. 75 (1763).
LIMNONESIS, Klotzsch, in Abh. Akad. Berl. (1853) 352.
ZALA, Lour. Fl. Cochinch. 405 (1790).
aegyptiaca, Schleid. in Otto & Dietr. Allg. Gartenz. vi. (1838) 19 = P. Stratiotes.
aethiopica, Fenzl, ex Klotzsch, in Abh. Akad. Berl. (1853) 354 = P. Stratiotes.
africana, Presl, Epim. Bot. 240 = P. Stratiotes.
amazonica, Presl, l. c. = P. Stratiotes.
brasiliensis, Klotzsch, in Abh. Akad. Berl. (1853) 356 = P. Stratiotes.
commutata, Schleid. in Otto & Dietr. Allg. Gartenz. vi. (1838) 20 = P. Stratiotes.
crispata, Blume, Rumphia, i. 78 = P. Stratiotes.
Cumingii, Klotzsch, in Abh. Akad. Berl. (1853) 354 = P. Stratiotes.
Gardneri, Klotzsch, l. c. 356 = P. Stratiotes.
Horkeliana, Miq. in Linnaea, xviii. (1844) 81 = P. Stratiotes.
Leprieuri, Blume, Rumphia, i. 79 = P. Stratiotes.
linguaeformis, Blume, l. c. 79 = P. Stratiotes.
minor, Blume, l. c. 78 = P. Stratiotes.
natalensis, Klotzsch, in Abh. Akad. Berl. (1853) 354 = P. Stratiotes.
obcordata, Schleid. in Otto & Dietr. Allg. Gartenz. vi. (1838) 20 = P. Stratiotes.
occidentalis, Blume, Rumphia, i. 79 = P. Stratiotes.
Schleideniana, Klotzsch, in Abh. Akad. Berl. (1853) 356 = P. Stratiotes.
spathulata, Michx. Fl. Bor. Am. ii. 162 = P. Stratiotes.
Stratiotes, *Linn. Sp. Pl.* 963.—Reg. trop.
texensis, Klotzsch, in Abh. Akad. Berl. (1853) 356 = P. Stratiotes.
Turpini, C. Koch, in Bot. Zeit. x. (1852) 577 = P. Stratiotes.
Weigeltiana, Presl, Epim. Bot. 240 = P. Stratiotes.

PISTOLOCHIA, Bernh. Syst. Verz. Erf. 57 (1800) = **Corydalis**, Vent. (Papav.).
cava, Bernh. l. c. 74 = C. tuberosa.
intermedia, Bernh. l. c. = C. fabacea.
solida, Bernh. l. c. = C. bulbosa.

PISTOLOCHIA, Rafin. Fl. Tellur. iv. 98 (1836) = **Aristolochia**, Linn.
angustifolia, Rafin. l. c. = A. angustifolia, *Cham.*
latifolia, Rafin. l. c. = A. Pistolochia.
Serpentaria, Rafin. l. c. = A. Serpentaria.

PISTORINIA, DC. Prod. iii. 399 (1828) = **Cotyledon**, Tourn. (Crassul.).
brachyantha, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 59 = C. Cossoniana.
breviflora, Coss. & Bal. l. c. xx. (1873) 247 = C. Cossoniana.
breviflora, Boiss. Elench. 42 = C. hispanica.
hispanica, DC. Prod. iii. 399 = C. hispanica.
intermedia, Boiss. & Reut. Diagn. Ser. II. ii. 60 = C. hispanica.
Salzmanni, Boiss. Voy. Bot. Espagne, ii. 224 = C. hispanica.

PISUM, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
abyssinicum, A. Br. in Flora, xxiv. (1841) 268 = sativum.
americanum, Mill. Gard. Dict. ed. VIII. n. 5 = Lathyrus magellanicus.
Aphaca, Brot. Fl. Lusit. ii. 145 = Lathyrus Aphaca.
arvense, *Linn. Sp. Pl.* 727.—Europ. ; As. bor.
arvense, Moris, Fl. Sard. i. 576 = elatius.

PISUM:—
Aucheri, Jaub. & Spach, Illustr. i. 91. t. 46 = formosum.
baclium, Hort. ex Steud. Nom. ed. II. ii. 345 (*badium*?) = sativum.
biflorum, Rafin. Caratteri, 71 = elatius.
biflorum, Stokes, Bot. Mat. Med. iv. 30 = sativum.
borussicum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
chlorospermum, Hort. l. c. = sativum.
coccineum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 356 = sativum.
coerulescens, Hort. ex Steud. Nom. ed II. ii. 345 = sativum.
commune, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 572 = arvense, sativum.
dasigynum, Rafin. in Am. Monthly Mag. (1819) 194 (Quid?).—Am. bor.
decurrens, Stokes, Bot. Mat. Med. iv. 30 = Lathyrus Ochrus.
elatius, *Bieb. Fl. Taur. Cauc.* ii. 151.—Reg. Mediterr.; Oriens.
elatum, Ser. in DC. Prod. ii. 368 = praec.
excorticatum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
fertile, Hort. ex Steud. l. c. = sativum.
formosum, *Alef. in Bonplandia*, ix. (1861) 237; *et Boiss. Fl. Orient.* ii. 624.—Reg. Caucas.; Persia.
frigidum, Alef. in Bot. Zeit. xviii. (1860) 204 = formosum.
fulvum, *Sibth. & Sm. Fl. Graec.* vii. 79. t. 688.—As. Min.; Syria.
granulatum, Lloyd, Fl. Loire-Inf. 75 = elatius.
humile, *Boiss. & Noé, Diagn.* Sér. II. ii. 45.—Syria.
humile, Mill. Gard. Dict. ed. VIII. n. 2 = sativum.
Jomardi, *Schrank, in Flora*, iv. (1821) 309.—Aegypt.
leptolobum, Reichb. Fl. Germ. Excurs. 533 = arvense.
macrocarpum, Ser. ex Schur, Enum. Pl. Transs. 172 = sativum.
macrospermum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
maritimum, Linn. Sp. Pl. 727 = Lathyrus maritimus.
multiflorum, Stokes, Bot. Mat. Med. iv. 33 = Lathyrus maritimus.
Ochrus, Linn. Sp. Pl. 727 = Lathyrus Ochrus.
parviflorum, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
praecox, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
prolificum, Hort. ex Steud. l. c. = sativum.
quadratum, Hort. ex Steud. l. c. = sativum.
quadratum, Reichb. Fl. Germ. Excurs. 533 = arvense.
ramulare, Reichb. l. c. = sativum.
roseum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
rugosum, Hort. ex Steud. l. c. = sativum.
saccharatum, Hort. ex Reichb. Fl. Germ. Excurs. 533 = sativum.
sativum, *Linn. Sp. Pl.* 727.—Europ.; As. bor.
sibiricum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
smyrnense, Hort. ex Steud. l. c. = sativum.
tetragonum, Hort. ex Pasq. Cat. Orto Bot. Nap. 81 = sativum.
thebaicum, Willd. Enum. Hort. Berol. Suppl. 51 = sativum.
Tuffetii, Lesson, Fl. Rochefortine, 170 = elatius.
umbellatum, Mill. Gard. Dict. ed. VIII. n. 3 = sativum.
uniflorum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
uniflorum, Moench, Meth. 160 = arvense.
variegatum, Presl, Fl. Sic. p. xxiii = elatius.
viride, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.
vulgare, Jundz. Opis Rosl. w Litw. Rosn. 305 = sativum.
zeylanicum, Hort. ex Steud. Nom. ed. II. ii. 345 = sativum.

PITAVIA, Molina, Sagg. Chili, ed. II. 287 (1810).
RUTACEAE, Benth. & Hook. f. i. 297.
GALVEZIA, Ruiz & Pav. Prod. i. 56. t. 35 (1794).
punctata, *Molina, Sagg. Chile*, ed. II. 287.—Chili.
spicata, Steud. Nom. ed. II. ii. 345 = Ercilla volubilis.

PITAVIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 215 (1838) = **Cneoridium**, Hook. f. (Simarub.).
dumosa, Nutt. l. c. = C. dumosum.

PITCAIRINIA, Regel, in Act. Hort. Petrop. ii. (1873) 435, sphalm. = seq.

PITCAIRNIA, L'Hérit. Sert. Angl. 7. t. 11 (1788).
BROMELIACEAE, Benth. & Hook. f. iii. 665.
COCHLIOPETALUM, Beer, Bromel. 68 (1857).
CODONANTHES, Rafin. Fl. Tellur. iv. 24 (1836).
HEPETIS, Sw. Prod. Veg. Ind. Occ. 56 (1788).
LAMPROCONUS, Lem. Jard. Fleur. ii. (1852) sub t. 127.
MELINONIA, Brongn. in E. Morr. Cat. Brom. Jard. Liège (1873), 10.
NEUMANNIA, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 369.
ORTHOPETALUM, Beer, Bromel. 70 (1857).
ORTHOPHYTUM, Beer, in Flora, xxxvii. (1854) 347.
PEPINIA, A. Brongn. ex André, Illustr. Hortic. xvii. (1870) 32. t. 5.
PHLOMOSTACHYS, Beer, Bromel. 45 (1857).
SPIRASTIGMA, L'Hérit. ex Schult. f. Syst. vii. p. lxx (1830).
affinis, C. Koch, Ind. Sem. Hort. Berol. (1857) App. 5 = albucaefolia.
albiflora, Spreng. Syst, iv. Cur. Post. 131 = seq.
albiflos, *Herb. in Bot. Mag.* t. 2642.—Bras.
albucaefolia, *Schrad. Blumenb.* 24. t. 3.—Ind. occ.
alta, *Hassk. in Verh. Nat. Ver. Nederl. Ind.* i. (1856) 5.—Ind. occ.
Altensteinii, *Lem. in Fl. des Serres*, ii. (1846) 162.—Venezuela.
Andreana, *Linden, in Illustr. Hortic.* xx. (1873) 146. t. 139.—Venez.; N. Granat.
angustifolia, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 401.—Ins. S. Crux.
aphelandraeflora, *Lem. Illustr. Hortic.* xvi. (1869) *Misc.* 90.—Bras.
araneosa, *Baker, in Journ. Bot.* xix. (1881) 231.—N. Granat.
asterotricha, Poepp. & Endl. Nov. Gen. et Sp. ii. 42. t. 158 = ferruginea.
atrorubens, *Baker, in Journ. Bot.* xix. (1881) 307.—Panama.
aurantiaca, Tenore, in Ann. Sc. Nat. Sér. IV. ii. (1854) 378 = densiflora.
australis, *Hort. Par. ex C. Koch, Ind. Sem. Hort. Berol.* (1856) *App.* 4.—Bras.
Beycalema, Beer, Bromel. 63 = muscosa.
brachiata, *Cham. ex Link, Jahrb.* i. II. (1820) 192.—Ins. Philipp.?
bracteata, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 202.—Ind. occ.
Brochinia, D. Dietr. Syn. Pl. ii. 1062 = Brocchinia paniculata.
bromeliaefolia, *L'Hérit. Sert. Angl.* 7.—Ind. occ.
caerulea, *Benth. & Hook. f. Gen.* iii. 666.—Chili.
caricifolia, *Mart. in Schult. f. Syst.* vii. II. 1242.—Bras.
carnea, *Beer, in Oestr. Bot. Zeitschr.* viii. (1858) 182.—Panama.
cernua, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 12 = heterophylla.
chilensis, Lodd. Cat. ex Loud. Hort. Brit. 118, nomen = Puya chilensis.
chrysantha, *Phil. Fl. Atac.* 50.—Chili.
cinnabarina, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 202.—Bras.
coarctata, Pers. Syn. i. 344 = Puya chilensis.
commutata, Regel, Gartenfl. (1868) 8 = bracteata.
concolor, *Baker, in Journ. Bot.* xix. (1881) 269.—Peruv.
consimilis, *Baker, l. c.* 266.—Bolivia.
corallina, *Linden.—Cf. Belg. Hortic.* xxiii. (1873) 112; *et ex Carr. in Rev. Hortic.* (1875) 251.—N. Granat.
corcovadensis, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 384.—Bras.
crystallina, Pers. Syn. i. 344 = Puya lanuginosa.
Decaisnei, C. Koch, Ind. Sem. Hort. Berol. (1857) App. 5 = fulgens.
decora, A. Dietr. in Allg. Gartenz. xv. (1847) 353 = integrifolia.
densiflora, *Brongn. in Hortic. Univ.* vi. 228; *ex Walp. Ann.* vi. 79.—Mexic.
discolor, Loisel. Herb. Amat. v. t. 345 = Billbergia speciosa.
distacaia, Beer, Bromel. 58 = Billbergia speciosa?
echinata, *Hook. Bot. Mag.* t. 4709.—N. Granat.
elata, *Liebm. Ind. Sem. Hort. Haun.* (1849) 14.—Mexic.

PITCAIRNIA:—
elegans, *Regel, Cat. Pl. Hort. Aksakov.* 112, nomen.—Hab.?
excelsa, E. Morr. Belg. Hortic. xxv. (1875) 381 = pulverulenta.
exscapa, Liebm. Ind. Sem. Hort. Haun. (1848) 12; Hook. Bot. Mag. t. 4591 = heterophylla.
fastuosa, C. Morr. in Ann. Soc. Bot. Gand, iii. (1847) 411. t. 16 = Billbergia thyrsoidea.
ferruginea, *Ruiz & Pav. Fl. Per.* iii. 36.—Peruv.; Mexic.
firma, *Baker, in Journ. Bot.* xix. (1881) 268.—Hab.?
flammea, *Lindl. in Bot. Reg.* t. 1092.—Bras.
flavescens, Baker, in Bot. Mag. t. 6318 = xanthocalyx.
floccosa, *Regel, in Act. Hort. Petrop.* iii. (1875) 124.—Hab.?
fulgens, *Decne. in Cat. Linden.* (1850) 18; *et ex A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 137.—Guatemala.
fulgens, F. G. Dietr. Vollst. Lexik. Gaertn. II. Nachr. vii. 44 = *splendens*, Hort.
Funkiana, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 337; *Regel, Gartenfl.* (1855) 45.—Venezuela.
furfuracea, Jacq. Eclog. Am. i. 117. t. 79; Sims, Bot. Mag. t. 2657 = latifolia.
furfuracea, Willd. Enum. Hort. Berol. 346 = Puya Bonplandiana.
Gireoudiana, A. Dietr. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 105 = bracteata.
Glymiana, *C. Koch, Wochenschr.* xi. (1868) 89.—Hab.?
graminifolia, Hort. ex Schrad. Blumenb. 27 = integrifolia.
heterophylla, *Beer, Bromel.* 68.—Mexic.
humilis, *Lodd. ex Sweet, Hort. Brit.* ed. I. 425, nomen.—Am. austr.
humilis, Tenore, Ind. Sem. Hort. Neap. (1855); ex C. Muell. in Walp. Ann. vi. 83 = Jacksoni?
imbricata, *Regel, Gartenfl.* (1868) 135. t. 579; *Baker, in Journ. Bot.* xix. (1881) 307.—Mexic.
inermis, *D. Dietr. Syn. Pl.* ii. 1061.—Peruv.
integrifolia, *Ker-Gawl. in Bot. Mag.* t. 1462.—Ind. occ.
intermedia, Hort. ex Schult. f. Syst. vii. II. 1248 = latifolia.
iridiflora, *Hort. ex Beer, Bromel.* 51.—Hab.?
Jacksoni, *Hook. Bot. Mag.* t. 4540.—Guatemala.
Kalbreyeri, *Baker, in Journ. Bot.* xix. (1881) 273.—N. Grenat.
Karwinskyana, *Schult. f. Syst.* vii. II. 1239.—Mexic.
Kegeliana, *C. Koch, ex Baker, in Journ. Bot.* xix. (1881) 230.—Guian.
Kegeliana, *Schlecht. in Linnaea*, xxiv. (1851) 664.—Hab.?
laevis, Beer, Bromel. 60 = suaveolens.
laevis, Willd. ex Schult. f. Syst. vii. II. 1249 = pungens.
lanata, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 203 = Puya lanata.
lanuginosa, *Ruiz & Pav. Fl. Per.* iii. 35. t. 258.—Peruv.
latifolia, Andr. Bot. Rep. t. 322 = bromeliaefolia.
latifolia, [DC. in] Red. Lil. ii. t. 73–74 = bracteata.
latifolia, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 401.—Ind. occ.
Lechleri, *Baker, in Journ. Bot.* xix. (1881) 269.—Peruv.
Lehmanni, *Baker, l. c.* 273.—N. Granat.
lepidota, Regel, in Act. Hort. Petrop. ii. (1873) 435 = Andreana.
Liebmanni, C. Koch, Ind. Sem. Hort. Berol. (1857) App. 7 = heterophylla.
longifolia, Beer, Bromel. 67 = heterophylla.
longifolia, Hook. Bot. Mag. t. 4775 = pulverulenta.
macrocalyx, Hook. l. c. t. 4705 = Funkiana.
maidifolia, *Decne. ex Planch. in Fl. des Serres*, ix. (1853–54) 151.—Guatemala.
Maroni ×, André, in Rev. Hortic. (1885) 108.
media, *Lodd. ex Sweet, Hort. Brit.* ed. I. 425, nomen.—Am. austr.
megasepala, *Baker, in Journ. Bot.* xix. (1881) 229.—N. Granat.
micrantha, Lindl. Bot. Reg. (1843) Misc. 44 = suaveolens.
microcalyx, *Baker, in Journ. Bot.* xix. (1881) 228.—Venezuela.

PITCAIRNIA :—
monstrosa, *Beer, Bromel.* 68.—Hab. ?
montalbensis, *Linden, ex A. Dietr. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 138.—Mexic.
Moreliana, Hort. ex C. Koch, Ind. Sem. Hort. Berol. (1857) App. 8 = australis.
Moritziana, *C. Koch & Bouché, l. c.* (1856) *App.* 4.—Guatemala.
Morrenii, Lem. in Jard. Fleur. iii. (1853) t. 291 = heterophylla.
muscosa, *Mart. ex Schult. f. Syst.* vii. II. 1240.—Bras.
nubigena, *Planch. in Fl. des Serres*, viii. (1852–53) 265. t. 847.—Veneznela.
nuda, *Baker, in Journ. Bot.* xix. (1881) 269.—Guian.
ochroleuca, *Baker, l. c.* 306.—Guatemala.
odorata, Hort. ex Regel, Gartenfl. iv. (1855) 46 = albiflos.
Olfersi, Link, in Verhandl. Ver. Gartenb. vii. (1831) 363. t. 3 = flammea.
orgyalis, Baker, in Journ. Bot. xix. (1881) 273 = Lehmanni.
paniculata, Ruiz & Pav. Fl. Per. iii. 86. t. 260 = pulverulenta.
pauciflora, *Baker, in Journ. Bot.* xix. (1881) 230.—Guian.
penduliflora, *A. Rich. Fl. Cub. Fanerog.* ii. 262.—Cuba.
petiolata, *Baker, in Journ. Bot.* xix. (1881) 307.—Guatemala.
platyphylla, Schrad. Blumenb. 26 = bromeliaefolia.
polyanthoides, Brongn. in Rev. Hortic. Sér. IV. iv. (1855) 244 = recurvata.
pruinosa, *H. B. & K. Nov. Gen. et Sp.* i. 295.—Venezuela.
pulverulenta, *Ruiz & Pav. Fl. Per.* iii. 36. t. 259.—Peruv. ; Bolivia.
pungens, *H. B. & K. Nov. Gen. et Sp.* i. 294.—Am. austr.
punicea, *Lindl. ex Hassk. in Verh. Nat. Ver. Nederl. Ind.* i. (1856) 12 ; *Beer, Bromel.* 56.—Mexic.
punicea, *Scheidw. in Bull. Acad. Brux.* ix. I. (1842) 25.—Mexic.
pyramidata, Link, Enum. Hort. Berol. i. 308 = Puya Bonplandiana.
pyramidata, Pers. Syn. i. 344 = Puya pyramidata.
racemosa, Woodf. ex Schult. f. Syst. vii. II. 1245 = bracteata.
ramosa, Jacq. Eclog. i. 154, in obs. = latifolia.
recurvata, *C. Koch, Ind. Sem. Hort. Berol.* (1857) *App.* 4.—Bras.
Redouteana, Beer, Bromel. 57 = bromeliaefolia.
Redouteana, Schult. f. Syst. vii. II. 1243 = angustifolia.
rhodostachys, *Hassk. in Verh. Nat. Ver. Nederl. Ind.* i. (1856) 8.—Hab. ?
ringens, Link, Klotzsch & Otto, Ic. Pl. Rar. 63 ; Regel, Gartenfl. (1853) 162. t. 53 = Karwinskyana.
Roezli, *E. Morr. in Belg. Hortic.* xxxv. (1885) 285. t. 18–19.—Peruv.
rubicunda, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 3 = australis.
secunda, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 301 = Tillandsia secunda.
spathacea, *Griseb. in Goett. Abh.* xxiv. (1879) 329.—Reg. Argent.
speciosa, Hort. ex Schult. f. Syst. vii. II. 1250, nomen. (Quid ?).—Hab. ?
splendens, Hort.—Cf. Rev. Hortic. Sér. I. iii. (1836) 157 = cinnabarina ?
splendens, *Warsz. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 176.—Am. centr.
Sprucei, *Baker, in Journ. Bot.* xix. (1881) 303.—Bras.
staminea, *Lodd. Bot. Cab.* t. 722.—Bras.
suaveolens, *Lindl. Bot. Reg.* t. 1069.—Bras.
subpetiolata, *Baker, in Journ. Bot.* xix. (1881) 267.—Bras.
sulphurea, Andr. Bot. Rep. t. 249 = bracteata.
tabulaeformis, *Linden, Cat.* n. 17 (1862) 5 ; *et E. Morr. in Belg. Hortic.* xii. (1862) 257.—Mexic.
tomentosa, Hort. ex F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. vi. 305 = angustifolia.
undulata, Hort. ex Beer, Bromel. 65 = seq. ?
undulata, *Schiedw. in Otto & Dietr. Allg. Gartenz.* x. (1842) 275.—Bras.
undulatifolia, Hook. Bot. Mag. t. 4241 = Altensteinii.

PITCAIRNIA :—
vallisoletana, *Lex. in La Llave et Lex. Nov. Veg. Desc. fasc.* i. 19.—Mexic.
violacea, *Brongn. in Ann. Fl. Pom.* i. (1847) 116.—*Cf. Otto & Dietr. Allg. Gartenz.* xv. (1847) 299.—Mexic. ?
virescens, C. Koch, Ind. Sem. Hort. Berol. (1857) App. 4 (Caraguatae sp.).—Peruv.
viridiflora, *Regel, Ind. Sem. Hort. Petrop.* (1866) 81.—Hab. ?
Warszewicziana, *Klotzsch, ex Beer, Bromel.* 66 (Warszewitziana).—Peruv.
Wendlandi, *Baker, in Journ. Bot.* xix. (1881) 306.—Hab. ?
xanthocalyx, *Mart. Ind. Sem. Hort. Monac.* (1848) *Adn.* 4.—*Cf. Linnaea*, xxiv. (1851) 195.—Bras.
zeifolia, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 11 ; (1857) App. 4 = maidifolia.

PITCHERIA, Nutt. in Journ. Acad. Philad. vii. (1834) 93 = **Rhynchosia**, Lour. (Legumin.).
galactoides, Nutt. l. c. 94 = R. galactoides.

PITHECELLOBIUM, Mart. Hort. Monac. 188 (1829) = **Pithecolobium**, Mart. (1837) (Legumin.).
cyclocarpum, Mart. l. c. = Enterolobium cyclocarpum.
inundatum, Mart. l. c. (Quid ?).—Hab. ?
Unguis-cati, Mart. l. c. (nomen) = Pithecol. Unguis-cati.

PITHECOCTENIUM, Mart. ex DC. in Meissn. Gen. 300 ; Comm. 208 (1840) ; et Prod. ix. 193 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1038.
Aubletii, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 12.—Am. trop.
botryoides, *DC. Prod.* ix. 195.—Bras.
buccinatorium, DC. l. c. = Bignonia buccinatoria.
Catharinae, *DC. l. c.* 196.—Bras.
cinereum, *DC. l. c.* 195.—Mexic.
clematideum, *Griseb. in Goett. Abh.* xxiv. (1879) 257.—Reg. Argent.
cordifolium, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 48.—Bras.
cuneifolium, *DC. Prod.* ix. 196.—Bras.
cynanchoides, *DC. l. c.* 195.—Bras.
frutescens, *DC. l. c.* 196.—Bras.
granulosum, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1158, nomen.—Guian.
hexagonum, *DC. Prod.* ix. 195.—Mexic.
laxiflorum, *DC. l. c.*—Mexic.
Lundii, *DC. l. c.* 196.—Bras.
muricatum, *Moç. ex DC. l. c.* 194.—Mexic.
obovatum, *Mart. ex DC. l. c.* 196.—Guian.
panamense, Benth. Bot. Voy. Sulph. 129 = Anemopaegma orbiculatum.
parviflorum, *Mart. ex DC. Prod.* ix. 197.—Bras.
reticulare, *DC. l. c.*—Bras.
rufescens, *DC. l. c.* 174.—Hab. ?
scabriusculum, *Mart. ex DC. l. c.* 197.—Bras.
Squalus, *DC. l. c.* 194.—Bras.
stipulare, *Mart. ex DC. l. c.*—Bras.
tenuiflorum, *Mart. ex DC. l. c.* 197.—Bras.
Vitalbae, *DC. l. c.* 196.—Bras.

PITHECOLOBIUM, Mart. in Flora, xx. (1837) II. Beibl. 114 ; (Herb. Fl. Bras. 114) (Pithecollobium). *LEGUMINOSAE*, Benth. & Hook. f. i. 597.
CATHORMION, Hassk. Retzia, i. 231 (1855).
SPIROLOBA, Rafin. Sylva Tellur. 119 (1838).
ZYGIA, P. Br. Hist. Jamaic. t. 22. f. 3 (1756).
acatlense, *Benth. in Trans. Linn. Soc.* xxx. (1875) 593.—Mexic.
acutangulum, Miq. Fl. Ind. Bat. Suppl. 282 = angulatum.
adianthifolium, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 218.—Guian.
affine, *Baker, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 577.—Malacca.
albicans, *Benth. l. c.* 592.—Mexic.
altissimum, *Oliver, Fl. Trop. Afr.* ii. 364.—Afr. trop.
amplum, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 596 ; *Mart. Fl. Bras.* xv. II. 451.—Bras.
anamallayanum, Bedd. Fl. Sylv. i. t. 189 = subcoriaceum.
angulatum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 208.—As. trop.
asplenifolium, *Griseb. Cat. Pl. Cub.* 83.—Cuba.
Auaremotemo, *Mart. in Flora*, xx. (1837) II. *Beibl.* 115.—Bras.

PITHECOLOBIUM :—
auriculatum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 217.—Bras.
Benthamianum, *Miq. in Linnaea*, xviii. (1844) 592.—Guian.
Berterianum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 220 = fragrans.
Bertolonii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 588.—Guatemala.
bicolor, *Benth. in Mart. Fl. Bras.* xv. II. 451.—Bras.
bigeminum, Hassk. Retzia, i. 223 = Albizzia lucida.
bigeminum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 115, in obs.—Ind. or. ; Malaya.
Blancheti, *Benth. in Trans. Linn. Soc.* xxx. (1875) 587.—Bras.
borneense, *Benth. l. c.* 579.—Malaya.
brevifolium, *Benth. in A. Gray, Pl. Wright.* i. 67.—Am. bor. occ.
Brongniartii, Duchass. & Walp. in Flora, xxxvi. (1853) 232 = micradenium.
bubalinum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 576.—Malaya.
calliandriflorum, Wright, ex Griseb. Cat. Pl. Cub. 83 = P. Hystrix.
calliandrifolium, Benth. in Trans. Linn. Soc. xxx. (1875) 573, sphalm. = praec.
campestre, *Spruce, ex Benth. l. c.* 583 ; *et in Mart. Fl. Bras.* xv. II. 436.—Bras.
candidum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 201 = excelsum.
cataractae, Benth. l. c. 213 = glomeratum.
cauliflorum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 116.—Bras.
cinereum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 216 = P. Saman.
cinereum, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 413 = Mimosa sericantha.
circinale, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 201.—Ind. occ.
claviflorum, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 596 ; *et in Mart. Fl. Bras.* xv. II. 452.—Bras.
Clypearia, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 209.—As. trop.
coccineum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 594.—Peruv.
cochleatum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 115, in obs.—Bras.
cognatum, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 107.—Am. centr.
comosum, Benth. l. c. iii. (1844) 221 = Calliandra comosa.
confertum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 577.—Malacca.
contortum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 115, in obs.—Malaya.
corymbosum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 221.—Guian.
cyclocarpum, Mart. in Flora, xx. (1837) II. Beibl. 115, in obs. = Enterolobium cyclocarpum.
daulense, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 586.—Ecuador.
divaricatum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 213.—Bras.
diversifolium, *Benth. l. c.* 201.—Bras.
dulce, *Benth. l. c.* 199.—Am. trop.
dumosum, *Benth. l. c.* 223.—Bras.
elachistophyllum, *A. Gray. ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 352.—Mexic.
ellipticum, Hassk. Retzia, i. 225 = Inga elliptica.
excelsum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 115, in obs.—Ecuador.
fagifolium, *Blume, ex Miq. Fl. Bat.* i. I. 35.—Java.
falcifolium, Hassk. Pl. Jav. Rar. 418 = montanum.
fasciculatum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 208.—Malaya.
ferrugineum, *Benth. l. c.* 216.—Guian.
filamentosum, *Benth. l. c.* v. (1846) 106.—Am. trop.
filicifolium, *Benth. l. c.* iii. (1844) 205.—Mexic ; Jamaic.
filipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 588.—Am. trop.
floribundum, *Spruce, ex Benth. l. c.* 584 ; *et in Mart. Fl. Bras.* xv. II. 437.—Bras.
foliolosum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 223.—Bras.
Forfex, Benth. l. c. 199 = P. Unguis-cati.

PITHECOLOBIUM:—
Fournieri, *Benth. in Trans. Linn. Soc.* xxx. (1875) 574.—N. Caled.
fragrans, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 220.—Am. trop.
furcatum, *Benth. l. c.* v. (1846) 106.—Mexic.
geminatum, *Benth. l. c.* iii. (1844) 202.—Ind. or.
glabratum, Mart. in Flora, xx. (1837) II. Beibl. 116 = cauliflorum.
Glaziovii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 597; *et in Mart. Fl. Bras.* xv. II. 454.—Bras.
glomeratum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 213.—Guian.
glomeriflorum, *Kurz, For. Fl. Brit. Burma*, i. 430.—Burma.
gracile, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 44.—Ind. or.
grandiflorum, *Benth. Fl. Austral.* ii. 424.—Austral.
guadalupense, Chapm. Fl. S. U. St. 116 = P. Unguis-cati.
gummiferum, Mart. in Flora, xx. (1837) II. Beibl. 116 = Enterolobium ellipticum.
Hendersonii, *F. Muell. Fragm.* v. 191.—Austral.
hymeneaefolium, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 198.—Am. trop.
Hystrix, *Benth. in Hook. Ic. Pl.* t. 1168.—Cuba.
inaequale, *Benth. in Trans. Linn. Soc.* xxx. (1875) 596.—Am. trop.
incuriale, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 218.—Bras.
Junghuhnianum, *Benth. in Miq. Pl. Jungh.* 269.—Java.
Kegelii, *Meissn. in Linnaea*, xxi. (1848) 249.—Guian.
laetum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 203.—Bras.
lanceolatum, Benth. l. c. v. (1846) 105 = ligustrinum.
Langsdorfii, *Benth. l. c.* iii. (1844) 205.—Bras.
lasiogynum, Mart. ex Benth. in Trans. Linn. Soc. xxx. (1875) 413 = Mimosa pithecolobioides.
lasiopus, Benth. in Hook. Journ. Bot. ii. (1840) 141 = cauliflorum.
lateriflorum, Blume, ex Miq. Fl. Ind. Bat. i. I. 34 = lobatum?
latifolium, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 214.—Am. trop.
laxiflorum, *Benth. l. c.* 212.—Ins. Timor.
lentiscifolium, Wright, in Sauv. Fl. Cub. 36 = tortum.
leucophyllum, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 581; *et in Mart. Fl. Bras.* xv. II. 433.—Bras.
ligustrinum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 213 = Vahlianum.
ligustrinum, *Klotzsch, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 571.—Am. trop.
lindseaefolium, *Spruce, ex Benth. l. c.* 590; *et in Mart. Fl. Bras.* xv. II. 443.—Bras.
lobatum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 208.—Burma; Malaya.
longiflorum, *Benth. l. c.* v. (1846) 107.—Bras.
lucidum, *Benth. l. c.* iii. (1844) 207.—China.
Lucyi, F. Muell. Fragm. vi. 201 = Archidendron Lucyi.
lusorium, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 204.—Bras.
macrophyllum, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 593.—Peruv.
macrostachyum, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 105.—Am. trop.
macrostachyum, Benth. l. c. iii. (1844) 198 = ligustrinum.
marginatum, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 586; *et in Mart. Fl. Bras.* xv. II. 439.—Bras.
Martianum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 218 = incuriale.
Mathewsi, *Benth. l. c.* 222.—Peruv.
micradenium, *Benth. l. c.* 217.—Ind. occ.
micranthum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 589.—Ins. S. Doming.
microcalyx, *Spruce, ex Benth. l. c.* 582; *et in Mart. Fl. Bras.* xv. II. 434.—Bras.
microcarpum, *Benth. l. c.* 576.—Malaya.
microphyllum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 200 = P. Unguis-cati.
Minahassae, *Teijsm. & Binn. Cat. Hort. Bog.* 273, nomen.—Malaya.
Miquelianum, Meissn. in Linnaea, xxi. (1848) 250 = adiantifolium?

PITHECOLOBIUM:—
moniliferum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 211.—Malaya; Austral.
montanum, *Benth. l. c.* 209.—Reg. Himal.; Malaya.
Mottleyanum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 575.—Borneo.
multiflorum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 220.—Am. trop.
niopoides, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 591; *et in Mart. Fl. Bras.* xv. II. 447.—Bras.
nitidum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 202 = umbellatum.
oblongum, *Benth. l. c.* 198.—Am. trop.
obovale, *Wright, in Sauv. Fl. Cub.* 36; *Benth. in Trans. Linn. Soc.* xxx. (1875) 582.—Cuba.
oppositum, *Miq. Fl. Ind. Bat. Suppl.* 283.—Sumatra.
Palmeri, *Hemsl. Diagn. Pl. Nov.* 50.—Mexic.
panamense, Walp. & Duchass. in Linnaea, xxiii. (1850) 746 = hymeneaefolium.
panurense, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 586; *et in Mart. Fl. Bras.* xv. II. 440.—Bras.
papuanum, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 22.—N. Guin.
paraguense, *Benth. in Trans. Linn. Soc.* xxx. (1875) 574.—Paraguay.
parvifolium, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 223.—Am. trop.
pauciflorum, Benth. l. c. 212 = lobatum.
pedicellare, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 219.—Bras.
Pervilleanum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 586.—Madag.
polycarpum, Poepp. & Endl. Nov. Gen. et Sp. iii. 81 = laetum.
polycephalum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 219.—Am. trop.
prehensile, *Benth. in Trans. Linn. Soc.* xxx. (1875) 593.—Cuba.
pruinosum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 211.—Austral.
pubescens, *Benth. in Hook. Journ. Bot.* ii. (1840) 141.—Guian.
Quaremotemo, Walp. Rep. i. 932, sphalm. = P. Auaremotemo.
ramiflorum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 215.—Bras.
rhombeum, *Benth. l. c.* 217.—Bras.
roseum, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 595 = latifolium.
rostratum, *Miq. Fl. Ind. Bat. Suppl.* 282.—Java.
salutare, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 216.—N. Granat.
Saman, *Benth. l. c.*—Am. trop.
sanguineum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 594; *et in Mart. Fl. Bras.* xv. II. 447.—Bras.
scalare, *Griseb. in Goett. Abh.* xxiv. (1879) 123.—Reg. Argent.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 352.—Mexic.
Schomburgkii, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 219 = Enterolobium Schomburgkii.
scutiferum, Benth. l. c. 211 = lobatum.
Selloi, *Benth. l. c.* 214.—Bras.
sericiflorum, *Benth. l. c.* v. (1846) 105.—Venezuela.
sessile, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 22.—N. Guin.
simabaefolium, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 443.—Am. trop.
simabifolium, Spruce, ex Benth. in Trans. Linn. Soc. xxx. (1875) 589 = praec.
sophorocarpum, *Benth. in Benth. & Hook. f. Gen.* i. 598.—Am. trop.
spinifolium, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 200 = circinale.
Splitgerberianum, Miq. Stirp. Surin. Sel. 5 = Albizzia Lebbek.
Spruceanum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 590; *et in Mart. Fl. Bras.* xv. II. 443.—Bras.
stipellatum, Bernh. ex Benth. l. c. 590 = Kegelii.
stipulare, *Benth. l. c.* 596.—Bras.
subacutum, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 210.—Ins. Philipp.
subcoriaceum, *Thw. Enum. Pl. Zeyl.* 100.—Ind. or.
tenuiflorum, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 222 = tortum.

PITHECOLOBIUM:—
tortum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 114.—Bras.
Tozerii, F. Muell. Fragm. v. 10 = grandiflorum.
trapezifolium, *Benth. in Hook. Journ. Bot.* ii. (1840) 142.—Am. trop.
triflorum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 591.—Ecuador.
turbinatum, Benth. in Hook. Lond. Journ. Bot. v. (1846) 106 = filamentosum.
umbellatum, *Benth. l. c.* iii. (1844) 202.—Ind. or.; Malaya.
umbellatum, Miq. Fl. Ind. Bat. i. I. 37 = moniliferum.
Unguis-cati, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 200.—Am. trop.
unifoliolatum, *Benth. l. c.* 212.—Bras.
Vahlianum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 594.—Am. trop.
Vaillantii, F. Muell. Fragm. v. 9 = Archidendron Vaillantii.
variegatum, Blume, ex Miq. Fl. Ind. Bat. i. I. 37 = montanum.
Vincentis, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 222 = tortum.
zanzibaricum, *S. Moore, in Journ. Bot.* xv. (1877) 292.—Afr. trop.

PITHECOSERIS, Mart. ex DC. Prod. v. 84 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 236.
pacourinoides, *Mart. ex DC. l. c.*—Bras.

PITHECURUS, Willd. ex Kunth, Rév. Gram. ii. 571 (1829); Enum. Pl. i. 490 (1833) = **Andropogon**, Linn. (Gramin.).
canescens, Willd. ex Steud. Nom. ed. II. ii. 346 = Trachypogon polymorphus.
gracilis, Willd. ex Kunth, Rév. Gram. ii. 571 = A. Myosurus.

PITHOCARPA, Lindl. Swan River App. 23 (1839). *COMPOSITAE*, Benth. & Hook. f. ii. 318.
corymbulosa, *Lindl. l. c.*—Austral.
major, Steetz, in Lehm. Pl. Preiss. i. 446 = corymbulosa.
pulchella, Lindl. Swan River App. 23 = corymbulosa.

PITHOSILLUM, Cass. in Dict. Sc. Nat. xli. 164 (1826) = **Senecio**, Tourn. (Compos.).
lyratum, Cass. l. c. 165 = S. Pithosillum.

PITHURANTHOS, DC. Prod. iv. 144 (1830) = **Pituranthos**, Viv. (Umbellif.).

PITRAEA, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) II. 328 = **Priva**, Adans. (Verben.).
chilensis, Turcz. l. c. 329 = Priva laevis.

PITTONIA, Plum. ex Adans. Fam. ii. 177 (1763) = **Tournefortia**, Linn. (Borag.).
foetidissima, Pritz. Ic. Ind. i. 855, sphalm. = Passiflora foetida.
hirsuta, Pritz. l. c. = T. hirsutissima.

PITTONIOTIS, Griseb. in Bonplandia, vi. (1858) 8 = **Antirrhoea**, Comm. (Rubiac.).
protracta, Griseb. l. c. = A. protracta.
trichantha, Griseb. l. c. = A. trichantha.

PITTOSPOROIDES, Soland. ex Gaertn. Fruct. i. 286 (1788) = **Pittosporum**, Banks.
umbellatum, Soland. ex Gaertn. l. c. = Pittosporum umbellatum.

PITTOSPORUM, Banks, ex Gaertn. Fruct. i. 286. t. 59 (1788). *PITTOSPOREAE*, Benth. & Hook. f. i. 131.
CYLBANIDA, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 135.
GLYASPERMUM, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 11.
PITTOSPOROIDES, Soland. ex Gaertn. Fruct. i. 286 (1788).
PSEUDITEA, Hassk. in Flora, xxv. (1842) II. Beibl. 30; et Cat. Hort. Bog. Alt. 160 (1844).
QUINSONIA, Montr. in Mém. Acad. Lyon, x. (1860) 178.
SENACIA, Comm. ex DC. Prod. i. 347 (1824).

PITTOSPORUM:—
abyssinicum, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 89; *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 41.—Abyss.
acacioides, A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 109 = phillyraeoides.
acuminatum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 152.—Ins. Sandvic.
Andersonii, *Fisch. ex G. Don, Gen. Syst.* i. 374.—Hab.?
angustifolium, Lodd. Bot. Cab. t. 1859 = phillyraeoides.
arborescens, *Rich, ex A. Gray, Bot. U. St. Expl. Exped.* i. 223.—Ins. Fiji.
arbutifolium, Hort. ex DC. Prod. i. 346 = viridiflorum.
Baudouinii, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 189.—N. Caled.
bicolor, *Hook. Journ. Bot.* i. (1834) 249.—Austral.
brachyandrum, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 141.—Madag.
brachysepalum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 366.—Ins. Philipp.
Brackenridgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 225.—Ins. Fiji.
bracteolatum, *Endl. Prod. Fl. Norf.* 78.—Ins. Norf.
Buchanani, *Hook. f. Handb. N. Zeal. Fl.* 725.—N. Zel.
Campbelli, *F. Muell. Phyt. New Hebr.* 5.—N. Hebrid.
capense, Hort. ex DC. Prod. i. 346 = viridiflorum.
capitatum, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 188.—N. Caled.
cauliflorum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 151.—Ins. Sandvic.
ceylanicum, *Wight, Illustr.* i. 173.—Zeylan.
chelidospermum, *Blume, Mus. Bot. Lugd. Bat.* i. 161. N. Guin.
chinense, Donn, Hort. Cantabr. ed. IV. 48 = P. Tobira.
Colensoi, *Hook. f. Fl. N. Zel.* i. 22.—N. Zel.
Colensoi, T. Kirk, in Trans. N. Z. Inst. iv. (1872) 262 = tenuifolium.
commutatum, Putterl. Syn. Pittosp. 10 = viridiflorum.
confertiflorum, *A. Gray, Bot. U. St. Expl. Exped.* i. 232.—Ins. Sandvic.
coriaceum, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 448.—Ins. Madera.
cornifolium, *A. Cunn. in Bot. Mag.* t. 3161.—N. Zel.
crassifolium, *Soland. ex Putterl. Syn. Pittosp.* 12.—N. Zel.
crenulatum, Putterl. l. c. 15 = pimeleoides.
crispum, *Desf. Cat. Hort. Par.* ed. III. 410.—Hab.?
dasycaulon, *Miq. Anal. Bot. Ind.* iii. 5.—Ind. or.
densiflorum, Putterl. in Endl. Nov. Stirp. Dec. 44; et Syn. Pittosp. 9 = floribundum?
Deplanchei, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 187.—N. Caled.
discolor, Hort. Berol. ex Regel, Gartenfl. (1852) 103, 133 = bicolor.
echinatum, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 189.—N. Caled.
elegans, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 121 = eugenioides.
ellipticum, *T. Kirk, in Trans. N. Z. Inst.* i. ed. II. (1871) 84, 86.—N. Zel.
Enderi, *Regel, Gartenfl.* (1868) 226. t. 587; *et Ind. Sem. Hort. Petrop.* (1868) 90.—Austral.
enkianthoides, R. Cunn. & Hueg. ex Putterl. Syn. Pittosp. 7 = eugenioides.
eriocarpum, *Royle, Illustr. Bot. Himal.* 77.—Reg. Himal.
erioloma, *C. Moore & F. Muell. in Fragm.* vii. 139.—Ins. Dom. Howe.
eugenioides, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 106 (eugeinoides).—N. Zel.
expense, Nois. ex Steud. Nom. ed. I. 628, nomen (Quid?).—Hab.?
fasciculatum, *Hook. f. Fl. N. Zel.* i. 24.—N. Zel.
fasciculatum, T. Kirk, in Trans. N. Z. Inst. iv. (1872) 262 = tenuifolium.
Fernandezii, Vidal, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 13 = brachysepalum.
ferrugineum, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 27.—Malaya; Austral.
fissicalyx, *Miq. Illustr. Fl. Arch. Ind.* 81.—Ins. Buru.
flavum, Hook. Bot. Mag. t. 4799 = Hymenosporum flavum.
floribundum, *Wight & Arn. Prod.* 154.—Ind. or.

PITTOSPORUM:—
Fortunei, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 562 = glabratum.
fulvum, Rudge, in Trans. Linn. Soc. x. (1811) 298. t. 20 = revolutum.
Gilliesianum, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 143; ed. II. (1871) 87.—N. Zel.
glabratum, *Lindl. in Journ. Hort. Soc.* i. (1846) 230.—Reg. Himal.; China.
glabrum, *Hook. & Arn. Bot. Beech. Voy.* 110.—Ins. Sandvic.
gracile, *Panch. Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 187.—N. Caled.
hirsutum, Link, Enum. Hort. Berol. i. 233 = revolutum.
hirtum, Hort. ex Tenore, in Atti 3ª Riun. Sc. Ital. (1841) 502 = Syncarpia laurifolia.
hirtum, Willd. Enum. Hort. Berol. 261 = revolutum.
Huegelianum, Putterl. in Endl. Stirp. Nov. Dec. 43 = bicolor.
Humblotianum, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 471.—Madag.
humile, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 198.—Reg. Himal.
Huttonianum, *T. Kirk, in Trans. N. Z. Inst.* i. ed. II. (1871) 90.—N. Zel.
intermedium, *T. Kirk, l. c.* 84.—N. Zel.
ioides, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 139.—Madag.
japonicum, Hort. ex Presl, Symb. Bot. i. 40 = Sloanea monosperma.
javanicum, *Blume, Mus. Bot. Lugd. Bat.* i. 159.—Java.
Kirkii, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* i. ed. II. (1871) 91.—N. Zel.
lanceolatum, A. Cunn. in Mitch. Journ. Trop. Austral. 272 = phillyraeoides.
lanceolatum, Putterl. Syn. Pittosp. 17 = Celastrus lanceolatus.
laurifolium, *Brouss. ex Roem. & Schult. Syst.* v. 432.—Ins. Teneriff.
ledifolium, A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 109 = bicolor.
ligustrifolium, A. Cunn. ex Loud. Hort. Brit. Suppl. i. 597, nomen; et ex Putterl. Syn. Pittosp. 16 = phillyraeoides.
linifolium, A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 109 cfr. typ. = *tinifolium*.
longifolium, Putterl. Syn. Pittosp. 15 = phillyraeoides.
loniceroides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 186.—N. Caled.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 5.—Afr. trop.
mauritianum, Lodd. Cat. (1825); ex G. Don, Gen. Syst. i. 374 = P. Senacia.
Mayi, *Hort. ex Regel, Cat. Pl. Hort. Aksakov.* 112, nomen.—Hab.?
melanospermum, *F. Muell. Fragm.* i. 70.—Austral.
microcarpum, Putterl. Syn. Pittosp. 7 = eugenioides.
moluccanum, Miq. Illustr. Fl. Arch. Ind. 76 = ferrugineum.
monticolum, *Miq. l. c.* 83.—Java.
nanum, Hook. Comp. Bot. Mag. i. (1835) 275 = Marianthus procumbens.
neelgherrense, *Wight & Arn. Prod.* 154.—Ind. or.
novoguineense, *Miq. Illustr. Fl. Arch. Ind.* 79.—N. Guin.
obcordatum, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 121.—N. Zel.
ochrosiaefolium, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1842) 15.—Madag.
oleaefolium, A. Cunn. ex Putterl. Syn. Pittosp. 17 = phillyraeoides.
ovatifolium, F. Muell. Fragm. ii. 78 = ferrugineum.
ovatum, T. Kirk, in Trans. N. Z. Inst. iv. (1872) 267 = ellipticum.
pachylobum, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 140.—Madag.
Pancheri, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 188.—N. Caled.
paniculatum, *Brongn. & Gris, l. c.*—N. Caled.
parviflorum, *Putterl. in Lehm. Pl. Preiss.* i. 189.—Austral.
patulum, *Hook. f. Handb. N. Zeal. Fl.* 19, 725.—N. Zel.
pauciflorum, *Hook. & Arn. Bot. Beech. Voy.* 168. t. 32.—China.
Pervillei, *Blume, Mus. Bot. Lugd. Bat.* i. 159.—Madag.

PITTOSPORUM:—
phillyraeoides, *DC. Prod.* i. 347 (phylliraeoides).—Austral.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 227.—Ins. Fiji.
pimeleoides, *A. Cunn. ex Putterl. Syn. Pittosp.* 15.—N. Zel.
polyspermum, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 138.—Madag.
procumbens, Hook. Comp. Bot. Mag. i. (1835) 275 = Marianthus procumbens.
radicans, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 108 = reflexum.
Ralphii, *T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 161.—N. Zel.
reflexum, *R. Cunn. ex A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 108.—N. Zel.
reticulatum, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 138.—Ins. Borbon.
revolutum, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 27.—Austral.
rhombifolium, *A. Cunn. ex Hook. Ic. Pl.* t. 621.—Austral.
rhytidocarpum, *A. Gray, Bot. U. St. Expl. Exped.* i. 228. t. 18.—Ins. Fiji.
Richii, *A. Gray, l. c.* 224.—Ins. Fiji.
rigidum, *Hook. f. Fl. N. Zel.* i. 22. t. 10.—N. Zel.
Ročanum, Putterl. Syn. Pittosp. 16 = phillyraeoides.
rubiginosum, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 108.—Austral.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 367.—Java.
Rumphii, Putterl. Syn. Pittosp. 7 = ferrugineum.
salicinum, Lindl. in Mitch. Journ. Trop. Austral. 97 = phillyraeoides.
Senacia, *Putterl. Syn. Pittosp.* 7.—Ins. Mascar.
serrulatum, Jack, ex Griff. in Calc. Journ. Nat. Hist. iv. (1844) 195 = Celastrus pauciflorus.
Simsoni, *Montr. in Mém. Acad. Lyon*, x. (1860) 178.—N. Caled.
sinense, Desf. Tabl. ed. II. 231 = viridiflorum.
sinuatum, *Blume, Mus. Bot. Lugd. Bat.* i. 161.—N. Guin.
spathulatum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 151.—Ins. Sandvic.
stenopetalum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 320.—Madag.
taitense, *Putterl. Syn. Pittosp.* 6.—Ins. Tahiti.
tenuifolium, *Gaertn. Fruct.* i. 286. t. 59. f. 7.—N. Zel.
terminalioides, *Planch. ex A. Gray, Bot. U. St. Expl. Exped.* i. 231.—Ins. Sandvic.
tetraspermum, *Wight & Arn. Prod.* 154.—Ind. or.
timorense, *Blume, Mus. Bot. Lugd. Bat.* i. 160.—Ins. Timor.
tinifolium, A. Cunn. in Ann. Nat. Hist. Sér. I. iv. (1840) 109 = ferrugineum.
Tobira, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 27.—Japon.; China.
tobiroides, *A. Gray, Bot. U. St. Expl. Exped.* i. 226. t. 17.—Ins. Fiji.
tomentosum, Bonpl. Desc. Pl. Rar. Malm. 56 = revolutum.
translucens, Hook. ex Putterl. Syn. Pittosp. 13 = tenuifolium.
tuberculatum, *Zeyh. ex Steud. Nom.* ed. I. 628, nomen.—Hab.?
turbinatum, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 187.—N. Caled.
umbellatum, A. Cunn. ex Putterl. Syn. Pittosp. 7 = eugenioides.
umbellatum, *Gaertn. Fruct.* i. 286.—N. Zel.
undulatum, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 368 = taitense.
undulatum, *Vent. Jard. Cels*, t. 76.—Austral.
uniflorum, *Putterl. Syn. Pittosp.* 12.—Hab.?
venulosum, *F. Muell. Fragm.* vi. 186.—Austral.
verticillatum, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1842) 16.—Madag.
verticillatum, *Montr. in Mém. Acad. Lyon*, x. (1860) 178.—N. Caled.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 188.—N. Caled.
virgatum, *T. Kirk, in Trans. N. Z. Inst.* i. ed. II. (1871) 86, 90.—N. Zel.
viridiflorum, *Sims, Bot. Mag.* t. 1684.—Afr. austr.
Wallichii, Putterl. Syn. Pittosp. 17 = Celastrus pauciflorus.
zeylanicum, Walp. Rep. i. 252 = ceylanicum.

PITTUNIA, Miers, Trav. Chile, ii. 531 (1826), sphalm. = **Petunia**, Juss. (Solan.).

PITUMBA, Aubl. Pl. Gui. ii. App. 29. t. 385 = **Casearia**, Jacq. (Samyd.).
edulis, Rich. ex Eichl. in Mart. Fl. Bras. xiii. 1. 470 = C. macrophylla.
guyannensis, Aubl. Pl. Gui. t. 385 = C. macrophylla.

PITURANTHOS, Viv. Fl. Lib. Spec. 15. t. 7 (1824).
UMBELLIFERAE, Benth. & Hook. f. i. 890.
DEVERRA, DC. Mém. Omb. 45 (1829); Prod. iv. 143 (1830).
DEVERRARIA, DC. ex Steud. Nom. ed. II. i. 497 (1840).
ERIOCYCLA, Lindl. in Royle, Illustr. Bot. Himal. 232. t. 51 (1835).
HYMENOPHORA, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 281.
PITHURANTHOS, DC. Prod. iv. 144 (1830).
denudatus, *Viv. Fl. Lib. Spec.* 15. t. 7. f. 1–2.—Afr. bor.
nudus, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 680.—Reg. Himal.
Thomsoni, *C. B. Clarke, l. c.*—Reg. Himal.

PITYOPSIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 317 = **Chrysopsis**, Ell. (Compos.).
argentea, Nutt. l. c. = C. graminifolia.
falcata, Nutt. l. c. = C. villosa.
graminifolia, Nutt. l. c. = C. graminifolia.
pinifolia, Nutt. l. c. 317, 318 = C. pinifolia.

PITYRANTHE, Thw. Enum. Pl. Zeyl. 29 (1858).
TILIACEAE, Benth. & Hook. f. i. 232.
verrucosa, *Thw. l. c.*—Zeylan.

PITYRANTHUS, Mart. in Denkschr. Akad. Muench. v. (1817) 179 = **Alternanthera**, Forsk. (Amarant.).
crassifolius, Mart. l. c. = A. Achyrantha.

PITYRODIA, R. Br. Prod. 513 (1810). *VERBENACEAE*, Benth. & Hook. f. ii. 1140.
DASYMALLA, Endl. Nov. Stirp. Dec. 11 (1839).
QUOYA, Gaudich. in Freyc. Voy. Bot. 453. t. 66 (1826).
atriplicina, *F. Muell. ex Benth. Fl. Austral.* v. 52.—Austral.
Bartlingii, *Benth. l. c.* 49.—Austral.
cuneata, *Benth. l. c.* 51.—Austral.
dilatata, *F. Muell. ex Benth. l. c.*—Austral.
Drummondii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 213.—Austral.
exsuccosa, F. Muell. Fragm. i. 60 = Dicrastylis ochrotricha.
hemigenioides, *F. Muell. ex Benth. Fl. Austral.* v. 48.—Austral.
Lewellini, F. Muell. Fragm. viii. 50 = Chloanthes Lewellini.
loricata, F. Muell. l. c. x. 14 = Chloanthes loricata.
myriantha, F. Muell. l. c. i. 236 = Dicrastylis fulva.
Oldfieldii, *F. Muell. ex Benth. Fl. Austral.* v. 52.—Austral.
paniculata, *F. Muell. ex Benth. l. c.* 53.—Austral.
racemosa, *Benth. l. c.* 50.—Austral.
salvifolia; *R. Br. Prod.* 513.—Austral.
uncinata, *Benth. Fl. Austral.* v. 48.—Austral.
verbascina, *F. Muell. ex Benth. l. c.* 50.—Austral.

PITYROPHYLLUM, Beer, Bromel. 79 (1857) = **Tillandsia**, Linn. (Bromel.).
erubescens, Beer, l. c. = T. ionantha.
gracile, Beer, l. c. = T. ionantha.

PITYROSPERMA, Sieb. & Zucc. in Abh. Akad. Muench. iii. IV. (1843) 734 = **Cimicifuga**, Linn. (Ranunc.).
acerinum, Sieb. & Zucc. l. c. 735 = C. japonica.
biternatum, Sieb. & Zucc. l. c. 736 = C. biternata.
obtusilobum, Sieb. & Zucc. l. c. 735 = C. obtusiloba.

PIVONNEAVA, Hook. f. Bot. Mag. t. 5668 (1867) sphalm. = PIRONNEAVA, Gaudich. = **Aechmea**, Ruiz & Pav. (Bromel.).

PLACEA, Miers, Trav. Chile, ii. 529 (1826), nomen; et in Bot. Reg. (1841) t. 50. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 725.
Arzae, *Phil. in Anal. Univ. Chil.* (1873) 541.—Chili.

PLACEA:—
Germaini, *Phil. in Linnaea*, xxix. (1857–58) 67.—Chili.
grandiflora, *Lem. Illustr. Hortic.* xv. (1868) t. 574.—Chili.
lutea, *Phil. in Linnaea*, xxxiii. (1864–65) 259.—Chili.
ornata, *Miers, Trav. Chile*, ii. 529, nomen; *et in Bot. Reg.* (1841) t. 50.—Chili.
pratensis, Poepp. Fragm. Syn. Pl. Chile, 5; ex Phil. f. Cat. Pl. Vasc. Chil. 291 = Hippeastrum pratense.

PLACOCARPA, Hook. f. in Benth. & Hook. f. Gen. ii. 107 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
mexicana, *Hook. f. l. c.* 108.—Mexic.

PLACODISCUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 332. *SAPINDACEAE*.
leptostachys, *Radlk. l. c.* ix. (1879) 606.—Afr. trop.
turbinatus, *Radlk. l. c.* viii. (1878) 332.—Afr. trop.

PLACODIUM, Benth. & Hook. f. Gen. ii. 136 (1873), sphalm. = **Plocama**, Ait. (Rubiac.).

PLACOLOBIUM, Miq. Fl. Ind. Bat. i. 1. 1082 (1857). Gen. dub. *LEGUMINOSARUM*.—Cf. Benth. & Hook. f. i. 465.
sumatranum, *Miq. l. c.*—Sumatra.

PLACOMA, J. F. Gmel. Syst. ii. 296, 390 (1791) = **Plocama**, Ait. (Rubiac.).

PLACOPODA, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 835. *RUBIACEAE*.
virgata, *Balf. f. l. c.* 836.—Ins. Socotr.

PLACOSTIGMA, Blume, Fl. Jav. Praef. 6 (1828) = **Podochilus**, Blume (Orchid.).

PLACUS, Lour. Fl. Cochinch. 496 (1790) = **Blumea**, DC. (Compos.).
aromaticus, Baill. Hist. Pl. viii. 303 = B. aromatica.
balsamifer, Baill. l. c. = B. balsamifera.
chinensis, Baill. l. c. = B. chinensis.
foetidus, Baill. l. c. = Pluchea camphorata.
grandis, Baill. l. c. = B. densiflora.
indicus, Baill. l. c. = Pluchea indica.
laevis, Lour. Fl. Cochinch. 497 = Baccharis laevis.
suaveolens, Baill. Hist. Pl. viii. 303 = Pluchea Quitoc.
tomentosus, Lour. Fl. Cochinch. 497 = Baccharis cochinchinensis.

PLADAROXYLON, Hook. f. in Hook. Ic. Pl. t. 1055 (1870) = **Senecio**, Tourn. (Compos.).
Leucadendron, Hook. f. l. c. = S. Leucadendron.

PLADERA, Soland. ex Roxb. Hort. Beng. [10] (1814); Fl. Ind. i. 400 (1832) = **Canscora**, Lam. (Gentian.).
alata, Buch.-Ham. ex Wall. Cat. n. 4364 B = C. perfoliata.
decussata, Roxb. Hort. Beng. [10]; Fl. Ind. i. 402 = C. decussata.
decussata, Wall. Cat. n. 4364, partim = Exacum pedunculatum.
fastigiata, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 429 = Hoppea fastigiata.
perfoliata, Roxb. Hort. Beng. 83; Fl. Ind. i. 402 = C. perfoliata.
pulchella, Wall. Cat. n. 4368 = C. Schultesii.
pusilla, Roxb. Hort. Beng. [10]; Fl. Ind. i. 403 = Hoppea dichotoma.
pusilla, Thw. Enum. Pl. Zeyl. 204 = Hoppea fastigiata.
sessiliflora, Roxb. Hort. Beng. 83; Fl. Ind. i. 400 = C. sessiliflora.
virgata, Griff. Notul. iv. 92 = C. andrographioides.
virgata, Roxb. Hort. Beng. [10]; Fl. Ind. i. 401 = C. diffusa.

PLAEA, Pers. Syn. i. 451 (1805) = **Pleea**, Michx. (Liliac.).

PLAESIANTHA, Hook. f. in Benth. & Hook. f. Gen. i. 681 (1865). *RHIZOPHOREAE*, Benth. & Hook. f. l. c.
Lobbii, *Hook. f. l. c.*—Borneo.

PLAGIACANTHUS, Nees, in DC. Prod. xi. 335 (1847) = **Dianthera**, Linn. (Acanth.).
racemosus, Nees, l. c. = D. racemosa.

PLAGIANTHERA, Reichb. & Zoll. in Verh. Natuurk. Ver. Nederl. Ind. i. (1856) 19; et in Linnaea, xxviii. (1856) 321 = **Mallotus**, Lour. (Euphorb.).
affinis, Baill. Etud. Gén. Euph. 424 = M. claoxyloides.
oppositifolia, Reichb. & Zoll. in Verh. Natuurk. Ver. Nederl. Ind. i. (1856) 20; et in Linnaea, xxviii. (1856) 321 = M. Blumeanus.

PLAGIANTHUS, Forst. Char. Gen. 85. t. 43 (1776). *MALVACEAE*, Benth. & Hook. f. i. 202.
ASTEROTRICHION, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. 19. t. 8 (1840).
BLEPHARANTHEMUM, Klotzsch, l. c. (1840).
GYNATRIX, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 34.
HALOTHAMNUS, F. Muell. Pl. Vict. i. 158 (1862).
LAWRENCIA, Hook. Ic. Pl. tt. 261, 417 (1840).
PHILIPPODENDRUM, Poit. in Ann. Sc. Nat. Sér. II. viii. (1837) 183. t. 3.
WRENCIALA, A. Gray, Bot. U. St. Expl. Exped. i. 180, in obs. (1854).
Berthae, *F. Muell. Fragm.* v. 103.—Austral.
betulinus, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 25.—N. Zel.
Dampieri, Hort.—Cf. Gard. Chron. (1884) I. 670; II. 200 = sidoides.
diffusus, *Benth. Fl. Austral.* i. 190.—Austral.
discolor, Aschers. in Ind. Sem. Hort. Berol. (1861) App. 10 = sidoides.
divaricatus, *Forst. Char. Gen.* 86.—N. Zel.
glomeratus, *Benth. in Journ. Linn. Soc.* vi. (1862) 103.—Austral.
humilis, Blanco, Fl. Filip. ed. I. 526 = Turraea pumila.
Lampenii, Booth, ex Lindl. in Bot. Reg. (1838) Misc. 22 = sidoides.
linariifolia, *J. Buch. in Trans. N. Z. Inst.* xvi. (1884) 394.—N. Zel.
Lyallii, Hook. f. ex Benth. in Journ. Linn. Soc. vi. (1862) 103 (Gayae sp.).—N. Zel.
microphyllus, *F. Muell. Fragm.* i. 29.—Austral.
petiolaris, Backh. ex Benth. in Journ. Linn. Soc. vi. (1862) 102, nomen = sidoides.
pulchellus, *A. Gray, Bot. U. S. Expl. Exped.* i. 181.—Austral.
sidoides, *Hook. Bot. Mag.* t. 3396.—Tasmania.
spicatus, *Benth. in Journ. Linn. Soc.* vi. (1862) 103.—Austral.
squamatus, Benth. l. c. = microphyllus.
tasmanicus, A. Gray, Bot. U. St. Expl. Exped. i. 181; Benth. in Journ. Linn. Soc. vi. (1862) 103 = pulchellus.
urticinus, A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 25 = betulinus.

PLAGIDIA, Rafin. New Fl. Am. iv. 42 (1836) = **Anychia**? Rich. (Illecebr.).
rufa, Rafin. l. c. (Quid?).—Am. bor.

PLAGIOBASIS, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 108. *COMPOSITAE*, Benth. & Hook. ii. 1236.
centauroides, *Schrenk, l. c.* 109.—Soongar.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 361.—Turkestan.

PLAGIOBOTHRYS, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 46 (1835). *BORAGINEAE* [Benth. & Hook. f. ii. 851].
arizonicus, *Greene, ex A. Gray, in Proc. Am. Acad.* xx. (1885) 284.—Am. bor. occ.
canescens, *Benth. Pl. Hartw.* 326.—Calif.
Cooperi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 285.—Calif.
glomeratus, *A. Gray, l. c.* 286.—Am. bor. occ.
hispidus, *A. Gray, l. c.*—Calif.
Kingii, *A. Gray, l. c.* 281.—Am. bor. occ.
nothofulvus, *A. Gray, l. c.* 285.—Am. bor. occ.
procumbens, *A. Gray, l. c.* 283.—Chili.
rufescens, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 46.—Am. bor. occ.; Chili.

PLAGIOBOTHRYS :—
shastensis, *Greene, ex A. Gray, in Proc. Am. Acad.* xx. (1885) 284.—Calif.
tenellus, *A. Gray, l. c.* 283.—Am. bor. occ.
tinctorius, *A. Gray, l. c.*—Chili.
Torreyi, *A. Gray, l. c.* 284.—Calif.
ursinus, *A. Gray, l. c.* 285.—Calif.

PLAGIOCARPUS, Benth. in Hook. Ic. Pl. t. 1162 (1873). *LEGUMINOSAE.*
axillaris, *Benth. l. c.*—Austral.

PLAGIOCARYUM, Willd. ex Steud. Nom. ed. I. 628 (1821) (Quid ?).
scabrum, Willd. ex Steud. l. c.—Bras.

PLAGIOCHEILUS, Arn. ex DC. Prod. vi. 142 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 430.
HIPPIA, Linn. f. Suppl. 389 (1781) partim ; H. B. & K. Nov. Gen. et Sp. iv. 301 (1820).
POLYGYNE, Phil. in Linnaea, xxxiii. (1864–65) 170.
bogotensis, *Wedd. Chlor. And.* i. 62.—N. Granat.
ciliaris, *Wedd. l. c.* 227.—Bolivia.
frigidus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 48. t. 248.—Peruv.
peduncularis, *Wedd. Chlor. And.* i. 61.—Ecuador.
prostratus, *Benth. Pl. Hartw.* 136.—Peruv.
solivaeformis, *DC. Prod.* vi. 142.—Bolivia.
tanacetoides, *Haenke, ex DC. l. c.*—Am. austr.

PLAGIOCHILUS, Lindl. Veg. Kingd. 712 (1847) sphalm. = praec.

PLAGIOLIRION, Baker, in Gard. Chron. (1883) II. 38. *AMARYLLIDEAE.*
Horsmanni, *Baker, l. c.*—N. Granat.

PLAGIOLOBA, Reichb. Nom. 182 (1841) = **Hesperis**, Linn. (Crucif.).

PLAGIOLOBIUM, Sweet, Fl. Austral. t. 2 (1827) = **Hovea**, R. Br. (Legumin.).
chorizemaefolium, Sweet, l. c. = H. chorizemaefolia.
ilicifolium, Sweet, l. c. sub t. 2 = H. chorizemaefolia.
macrocalyx, Walp. Rep. v. 434, sphalm. = Platylobium obtusangulum.

PLAGIOLYTRUM, Nees, in Proc. Linn. Soc. i. (1841) 95 = **Tripogon**, Roth (Gramin.).
calycinum, Nees, l. c. = T. bromoides.
filiforme, Nees, l. c. = T. unidentatum.
unidentatum, Nees, l. c. = T. unidentatum.

PLAGION, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = PLAGIUS, L'Hérit. = **Chrysanthemum**, Tourn. (Compos.).

PLAGIOPHYLLUM, Schlecht. in Linnaea, xiii. (1839) 428 = **Centradenia**, G. Don (Melastom.).
grandifolium, Schlecht. l. c. 429 = C. grandifolia.
parviflorum, Triana, in Trans. Linn. Soc. xxviii. (1871) 32, sphalm. = seq.
parvifolium, Schlecht. in Linnaea, xiii. (1839) 428 = C. inaequilateralis.

PLAGIOPODA, Spach, Hist. Vég. Phan. x. 402 (1841) = **Grevillea**, R. Br. (Proteac.).

PLAGIOPTERON, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 244. t. 13. *TILIACEAE*, Benth. & Hook. f. i. 238.
fragrans, Griff. l. c. t. 13 = suaveolens.
suaveolens, *Griff. l. c.* 244.—Burma.

PLAGIORHEGMA, Maxim. Prim. Pl. Amur. 34. t. 1 (1859) = **Jeffersonia**, Barton (Berberid.).
dubium, Maxim. l. c. = J. dubia.

PLAGIOSCYPHUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 335. *SAPINDACEAE.*
cauliflorus, *Radlk. l. c.* 336.—Madag.

PLAGIOSETUM, Benth. in Hook. Ic. Pl. xiii. 33 (1877); et Fl. Austral. vii. 494 (1878). *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
refractum, *Benth. ll. cc.*—Austral.

PLAGIOSTEMON, Klotzsch, in Linnaea, xii. (1838) 232 = **Simocheilus**, Klotzsch (Ericac.).
bicolor, Klotzsch, l. c. = S. bicolor.
puberulus, Klotzsch, l. c. = S. Klotzschianus.
pubescens, Benth. in DC. Prod. vii. 703, sphalm. = praec.

PLAGIOSTIGMA, Presl, Bot. Bemerk. 127 (1844); in Abh. Boehm. Ges. Wiss. V. iii. (1845) 557 = **Aspalathus**, Linn. (Legumin.).
pineum, Presl, ll. cc. = A. pinea.

PLAGIOSTIGMA, Zucc. in Abh. Akad. Muench. iv. III. (1846) 154; (Fl. Jap. Fam. Nat. ii. 98) = **Ficus**, Tourn. (Urtic.).
pumila, Zucc. ll. cc. = F. pumila.
stipulata, Zucc. ll. cc. = F. pumila.

PLAGIOTAXIS, Wall. Cat. n. 1269, 1270 (1831) = **Chickrassia**, A. Juss. (Meliac.).
Chickrassa, Wall. Cat. n. 1269 = C. tabularis.
grandiflora, Wall. l. c. n. 1271 = Chisocheton grandiflorus.
velutina, Wall. l. c. n. 1270 = Chickrassia tabularis.

PLAGISTRA, Rafin. Fl. Tellur. iv. 98 (1836) = **Aristolochia**, Tourn.
cretica, Rafin. l. c. = A. cretica.

PLAGIUS, L'Hérit. ex DC. Prod. vi. 135 (1837) = **Chrysanthemum**, Tourn. (Compos.).
ageratifolius, L.Hérit. ex DC. l. c. = C. flosculosum.
Allionii, L'Hérit. ex DC. l. c. = C. flosculosum.
grandiflorus, L'Hérit. ex DC. l. c.—Afr. bor.
virgatus, DC. l. c. = C. flosculosum.

PLANARIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 416 = **Chaetocalyx**, DC. (Legumin.).
latisiliquum, Desv. l. c. 417 = C. latisiliqua.

PLANCHONIA, Blume, in Fl. des Serres, vii. (1851–52) 24. *MYRTACEAE*, Benth. & Hook. f. i. 721.
alata, Blume, l. c. 25 = timorensis.
crenata, *Miers, in Trans. Linn. Soc.* Ser. II. i. (1875) 91.—Austral.
elliptica, *Miers, l. c.* 93.—Borneo.
littoralis, Blume, in Fl. des Serres, vii. (1851–52) 25 = valida.
nitida, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 93, sphalm. = valida.
sumatrana, Blume, in Fl. des Serres, vii. (1851–52) 25 = valida.
sundaica, Miq. Fl. Ind. Bat. i. 1. 493 = valida.
tetraptera, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 93 = timorensis.
timorensis, *Blume, in Fl. des Serres*, vii. (1851–52) 25.—Ins. Timor. ; Java.
undulata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 256.—Java.
valida, *Blume, in Fl. des Serres*, vii. (1851–52) 24.—Malaya.

PLANCHONIA, Dun. in DC. Prod. xiii. 1. 471 (1852) = **Salpichroa**, Miers (Solanac.).
arbutifolia, Dun. l. c. 475 = S. rhomboidea.

PLANCHONIA, J. Gay, ex Benth. & Hook. f. Gen. i. 154 (1862) = **Polycarpaea**, Lam. (Caryoph.).

PLANCIA, Neck. Elem. i. 49 (1790) = **Leontodon**, Linn. (Compos.).

PLANERA, Giseke, Prael. Linn. 205 (1792) = **Costus**, Linn. (Scitam.).
speciosa, Giseke, l. c. 244 = C. speciosus.

PLANERA, J. F. Gmel. Syst. 150 (1791). *URTICACEAE*, Benth. & Hook. f. iii. 353.
Abelicea, Schult. Syst. vi. 304 = Zelkova cretica.
acuminata, Lindl. in Gard. Chron. (1861) 428 = Zelkova acuminata.
americana, Hort. ex Steud. Nom. ed. II. ii. 347 = aquatica.
aquatica, *J. F. Gmel. Syst.* 150.—Am. bor.
carpinifolia, Wats. Dendrol. t. 106 = Zelkova crenata.
crenata, Desf. Tabl. ed. II. 242 = Zelkova crenata.
Davidii, Hance, in Journ. Bot. vi. (1868) 333 (= *Hemiptelea Davidii*).—China.

PLANERA :—
Gmelini, Michx. Fl. Bor. Am. ii. 248 = aquatica.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 66 = Zelkova acuminata.
Kaki, Hort. ex Planch. in DC. Prod. xvii. 167 = Zelkova acuminata.
Kiaki, Hort. ex Planch. l. c. = Zelkova acuminata.
parvifolia, Sweet, Hort. Brit. ed. II. 464 = Ulmus parvifolia.
Richardi, Michx. Fl. Bor. Am. ii. 248 = Zelkova crenata.
Richardi, Spreng. Syst. i. 493, partim = aquatica.
ulmifolia, Michx. f. Hist. Arb. Am. iii. 283 = aquatica.

PLANOTIA, Munro, in Trans. Linn. Soc. xxvi. (1868) 70. *GRAMINEAE*, Benth. & Hook. f. iii. 1209.
PLATONIA, Kunth, Rév. Gram. i. 139. 327. t. 76 (1830).
acuminatissima, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 72.—N. Granat.
aperta, *Munro, l. c.* 73.—N. Granat.
elata, *Munro, l. c.* 71.—Ecuador.
nobilis, *Munro, l. c.* 72.—N. Granat.
virgata, *Munro, l. c.* 71.—Ins. Trinit.

PLANTAGINASTRUM, Heist. Syst. 7 (1748) (Quid ?).

PLANTAGINELLA, Dill. ex Moench, Meth. 427 (1794) = **Limosella**, Linn. (Scrophul.).
aquatica, Moench, l. c. = L. aquatica.

PLANTAGINELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 140 = **Plantago**, Tourn.
alpina, Fourr. l. c. = Plantago alpina.
carinata, Fourr. l. c. = Plantago carinata.
crassifolia, Fourr. l. c. = Plantago crassifolia.
graminea, Fourr. l. c. = Plantago maritima.
maritima, Fourr. l. c. = Plantago maritima.
subulata, Fourr. l. c. = Plantago subulata.

PLANTAGO, [Tourn.] Linn. Syst. ed. I (1735). *PLANTAGINEAE*, Benth. & Hook. f. ii. 1224.
ARNOGLOSSUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 292 (1821).
ASTEROGEUM, S. F. Gray, l. c. 294 (1821).
CORONOPUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 140.
LAGOPUS, Fourr. l. c. (1869).
PLANTAGINELLA, Fourr. l. c. (1869).
PSYLLIUM, Tourn. ex Juss. Gen. 90 (1789).
abyssinica, *Hochst. ex A. Br. in Flora*, xxiv. (1841) i. 279; *et ex Steud. in Flora*, xxxii. (1849) 413.—Abyss.
acanthophylla, *Decne. in DC. Prod.* xiii. 1. 730.—Europ. austr.
accedens, Rafin. Fl. Ludov. 33 = virginica.
adriatica, Campana, Cat. Hort. Ferrar. 22 = major.
adspersa, Bernh. Sel. Sem. Hort. Erf. (1836).—Cf. Linnaea, xii. (1838) Litt. 75 = P. Cynops.
aegyptica, Jacq. Coll. i. 45 = squarrosa.
affinis, *Decne. in DC. Prod.* xiii. 1. 725.—Reg. Andin.
afra, Linn. Sp. Pl. ed. II. 168 = P. Psyllium.
agrestis, Salzm. ex Steud. Nom. ed. II. ii. 347 = P. Psyllium.
agrostophylla, *Decne. in DC. Prod.* xiii. 1. 708.—N. Granat.
akkensis, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 66, nomen.—Marocc.
albicans, Hort. ex Roem. & Schult. Syst. iii. 127 = ovata.
albicans, *Linn. Sp. Pl.* 114.—Reg. Mediterr.
albicans, Pall. Reise, ii. 329 = minuta.
albicans, Schousb. ex Spreng. Syst. i. 439 = velutina.
albiflora, *Rafin. New Fl. Am.* iv. 11.—Am. bor.
aliena, Schrad. ex Roem. & Schult. Syst. iii. 119, in obs. = P. Lagopus.
alopecura, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Alopecurus.
alopecuroides, *Lam. Illustr.* i. 342.—Afr. bor.
Alopecurus, *Decne. in DC. Prod.* xiii. 1. 709.—Am. trop.
alpina, *Linn. Sp. Pl.* 114.—Europ. med.
alpina, Sieber, in Roehl. Deutschl. Fl. ed. III. i. 809 = maritima.
alpina, Vill. Hist. Pl. Dauph. ii. 302 = montana.
altissima, Linn. Sp. Pl. ed. II. 164 = lanceolata.
altissima, Loisel. Fl. Gall. ed. I. 88 = major.
ambigua, Guss. Fl. Sic. Prod. i. 185 = lanceolata.

PLANTAGO:—
amblodes, Rafin. New Fl. Am. iv. 12 (Quid?).—Am. bor.
amplexicaulis, *Cav. Ic.* ii. 22. t. 125.—Reg. Mediterr.; Oriens.; Ind. bor. occ.
andicola, Decne. ex Barn. Monog. Plantag. 17 = barbata.
angustata, Schur, Enum. Pl. Transs. 559 = major.
antarctica, *Decne. in DC. Prod.* xiii. 1. 703.—Austral.
aquatica, Gueldenst. Reisen, ii. 58 = Alisma Plantago.
aquatilis, *Griseb. in Goett. Abh.* xxiv. (1879) 221.—Reg. Argent.
arabica, *Boiss. Diagn.* Ser. I. xii. 94.—Arabia.
arachnoides, *Schrenk, Enum. Pl. Nov.* (1841) 16.—As. austr.
arborescens, *Poir. Encyc.* v. 389.—Ins. Canar.
Archeri, Hook. f. Fl. Tasman. i. 303 = tasmanica?
arctica, *Decne. in DC. Prod.* xiii. 1. 700.—Kamtschat.
arctica, Trev. Ind. Sem. (1829); ex Decne. l. c. 694 = major.
arenaria, Decne. in DC. Prod. xiii. 1. 735 = P. Psyllium.
arenaria, *Waldst. & Kit. Pl. Rar. Hung.* i. 51. t. 51.—Europ.; As. Min.
argentea, Brot. Fl. Lusit. i. 156 = lanceolata.
argentea, *Chaix, in Vill. Hist. Pl. Dauph.* i. 376; ii. 302.—Europ.
argentea, Desf. Fl. Atlant. i. 136 = ovata.
argentea, Lam. Illustr. i. 340 = monosperma.
argentea, Sieber, ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 464 = cylindrica.
argentea, Tenore, Fl. Nap. Suppl. i. 59 = albicans.
argentea, Webb, Iter Hisp. 17 = nivalis.
argyrophylla, *Decne. in DC. Prod.* xiii. 1. 711.—N. Granat.
aristata, Michx. Fl. Bor. Am. i. 95 = patagonica.
arvensis, Presl, Delic. Prag. 69 = P. Lagopus.
asiatica, Ledeb. Fl. Alt. i. 143 = sibirica.
asiatica, Linn. Sp. Pl. 113 = major.
asiatica, Turcz. in Bull. Soc. Nat. Mosc. (1838) 1. 99 = paludosa.
aspera, Gaud. Fl. Helv. i. 403 = maritima.
atrata, Hoppe, Cent. ii. Taschenb. (1799) 85; Hoffm. Deutschl. Fl. ed. II. i. 1. 76, in obs. = montana.
atrata, Tenore, Fl. Nap. Prod. App. iv. 10 = lanceolata.
atrofusca, *Rafin. Atl. Journ.* 150.—Hab.?
attenuata, James, in Long, Exped. i. 445; ex Torr. in Ann. Lyc. N. York, ii. (1828) 237 = eriopoda.
attenuata, Wall. in Roxb. Fl. Ind. ed. Carey, i. 422 = lanceolata.
aucklandica, *Hook. f. Fl. Antarct.* i. 64.—Ins. Auckland.
australis, *Lam. Illustr.* i. 339.—Reg. Argent.
azorica, Hochst. ex Seub. Fl. Azor. 30 = lanceolata.
barbata, *Forst. in Comm. Gotting.* ix. (1789) 25. t. 4.—Reg. Magellan.
Barneoudii, *Decne. in DC. Prod.* xiii. 1. 710.—N. Granat.
baufuloides, *Pomel, Nouv. Mat. Fl. Atl.* 311.—Algeria.
Bauphula, Edgew. in Hook. Journ. Bot. ii. (1840) 285 = amplexicaulis.
Bellardi, *All. Fl. Pedem.* i. 82. t. 85. f. 3.—Reg. Mediterr.; Oriens.
Bellardi, Drège, ex Decne. in DC. Prod. xiii. 1. 719 = caffra.
bellidifolia, Viv. Pl. Aegypt. Dec. iv. 23 = ciliata.
bellidioides, Decne. in DC. Prod. xiii. 1. 701 = varia?
Bernoulliana, *Vatke, in Verh. Bot. Ver. Brandenb.* (1874) 48.—Guatemala.
Berteroi, *Steinh. ex Decne. in DC. Prod.* xiii. 1. 726.—Chili.
Berteroniana, *Steud. in Flora*, xxxii. (1849) 407.—Chili.
Bertolonii, Godr. in Gren. & Godr. Fl. Fr. ii. 721 = media.
bicallosa, *Decne. in DC. Prod.* xiii. 1. 725.—Bras.
bicarinata, *G. F. W. Mey. Prim. Fl. Esseq.* 88.—Guian.
bidentata, Murith, Bot. Val. 85 = alpina.
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 117.—Am. bor. occ.
Bigelovii, S. Wats. in Bot. King's Exped. 212 = pusilla.
Bismarckii, *Nederl. in Monatschr. Bef. Gartenb.* (1881) 16. t. 1.—Patagon.
bonariensis, *Steud. in Flora*, xxxii. (1849) 408.—Reg. Argent.

PLANTAGO:—
borealis, Lange, in Fl. Dan. t. 2707 = maritima.
brachyantha, *Phil. Fl. Atac.* 47.—Chili.
brachypetala, Wallr. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 402 = brachystachys.
brachyphylla, *Edgew. ex Decne. in DC. Prod.* xiii. 1. 696.—Reg. Himal.
brachyphylla, Roem. & Schult. Syst. iii. 136 = subulata.
brachystachys, *Kunze, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 402.—Chili.
bracteata, Moench, Meth. 459 = major.
bracteosa, *Phil. in Anal. Univ. Chil.* (1873) 534.—Chili.
brasiliensis, Sims, Bot. Mag. t. 2616 = coriacea.
Bridgesii, *Decne. in DC. Prod.* xiii. 1. 725.—Chili.
Brongniartii, *Barn. Monog. Plantag.* 35.—Ins. Sandvic.
Browniana, Schult. Mant. iii. 86 = seq.
Brownii, *Rafin. in Mém. Soc. Linn. Par.* vi. (1827) 485.—Austral.; N. Zel.
brutia, *Tenore, Prod. Fl. Nap.* 59.—Ital.
bulbosa, Crantz, Inst. ii. 330 = P. Serraria.
bullata, Donn, Hort. Cantab. ed. I. 15 = crispa.
Bungei, *Steud. Nom.* ed. II. ii. 348.—Sibir. altaic.
Burchelli, *Decne. in DC. Prod.* xiii. 1. 720.—Afr. austr.
byzantina, C. Koch, in Linnaea, xxi. (1848) 712 = lanceolata.
caespitosa, *Phil. in Linnaea*, xxix. (1857-58) 37.—Chili.
cafra, *Decne. in DC. Prod.* xiii. 1. 719.—Afr. austr.
calabrica, Gaud. ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 452 = media.
californica, Greene, in Bull. Calif. Acad. i. III. (1885) 123 = heterophylla.
callosa, Colla, in Mem. Acc. Torin. xxxix. (1836) 6 = tumida.
camtchatica, Hook. & Arn. Bot. Beech. Voy. 156 = hirtella?
camtschatica, Link, Enum. Hort. Berol. i. 120 = major.
camtschatica, Rugel, ex Decne. in DC. Prod. xiii. 1. 700 = Rugelii.
canadensis, Hort. Par. ex Pursh, Fl. Am. Sept. i. 98 = cordata.
canaliculata, *Hoffmgg. Verz. Pfl. Nachtr.* iii. 51.—Hab.?
canaliculata, Pourr. Obs. 15; ex Decne. in DC. Prod. xiii. 1. 728 = maritima.
candicans, Auct. ex Decne. l. c. 713 = patagonica.
candicans, Schrad. ex Barn. Monog. Plantag. 43 = nivea.
Candollei, *Rafin. in Mém. Soc. Linn. Par.* vi. (1827) 453.—Chili.
canescens, *Adams, in Mém. Soc. Nat. Mosc.* ix. (1834) 233. t. 13. f. 1.—Sibir. or.
canescens, Schrad. Ind. Sem. Hort. Gotting.; ex Decne. in DC. Prod. xiii. 1. 713 = patagonica.
capensis, Boj. Hort. Maurit. 263 = lanceolata.
capensis, Thunb. Prod. Pl. Cap. 29 = remota.
capillaris, *E. Mey. ex Decne. in DC. Prod.* xiii. 1. 719.—Afr. austr.
capitata, Hoppe, ex Reichb. Fl. Germ. Excurs. 395 = argentea.
capitata, Tenore, Prod. Fl. Nap. 59 = lanceolata.
capitella, Ram. ex DC. Fl. Fr. iii. 414 = carinata.
capitellata, DC. l. c. = carinata.
capitellata, Soleir. ex Willk. & Lange, Prod. Fl. Hisp. ii. 357 = subulata.
caricina, *Decne. in DC. Prod.* xiii. 1. 708.—N. Granat.
carinata, Lange, in Kjoeb. Vidensk. Meddel. 1861 (1862) 57 = acanthophylla.
carinata, Moench, Meth. 460 = Loeflingii.
carinata, *Schrad. ex Mert. & Koch, Deutschl. Fl.* i. 810.—Europ. austr.; As. Min.
carnosa, R. Br. Prod. 425 = Brownii.
carnosa, *Lam. Illustr.* i. 342.—Afr. austr.
caroliniana, Pursh, Fl. Am. Sept. i. 98 = sparsiflora, heterophylla?
caroliniana, Walt. Fl. Carol. 85 = virginica.
caspica, Steud. Nom. ed. II. ii. 348 = Loeflingii.
Catananche, Clarke, Trav. iii. 336 = cretica.
catharinea, *Decne. in DC. Prod.* xiii. 1. 726.—Bras.
caulescens, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 141.—Mexic.
ceratophylla, Hoffmgg. & Link, Fl. Port. i. 431 = macrorhiza.
Chamissoi, Wallr. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 402 = P. Princeps.

PLANTAGO:—
chilensis, Desf. Cat. Hort. Par. ed. III. 390 = Candollei.
chilensis, *Rapin, in Mém. Soc. Linn. Par.* vi. (1827) 475.—Am. austr.
chottica, *Pomel, Nouv. Mat. Fl. Atl.* 124.—Alger.
ciliata, Boiss. Voy. Bot. Espagne, ii. 535 = albicans.
ciliata, *Desf. Fl. Atlant.* i. 137. t. 39.—Afr. bor.; Oriens; Ind. bor. occ.
cinerascens, Ser. ex Barn. Monog. Plantag. 25 = alpina.
cinerea, Domb. ex Decne. in DC. Prod. xiii. 1. 712 = tumida.
clausa, *Steud. in Flora*, xxxii. (1849) 407.—Chili.
Columnae, Gouan, Illustr. 6 = P. Coronopus.
Commersoniana, *Decne. ex Barn. Monog. Plantag.* 37.—Reg. Argent.
commutata, Guss. Suppl. Fl. Sic. Prod. i. 46 = P. Coronopus.
compressa, Rafin. in Am. Month. Mag. (1811) 356, nomen (Quid?).—Am. bor.
concinna, Salisb. Prod. 46 = media.
congesta, *Ruiz & Pav. Fl. Per.* i. 51. t. 79.—Peruv.
connivens, Moench, Meth. Suppl. 174 = virginica.
consanguinea, Decne. in DC. Prod. xiii. 1. 703 = varia.
contorta, Guss. Fl. Sic. Prod. i. 186 = lanceolata.
cordata, *Lam. Illustr.* i. 138.—Am. bor.
coriacea, *Cham. & Schlecht. in Linnaea*, i. (1826) 171.—Bras.
Cornuti, Gouan, Illustr. 6 = major.
Cornuti, Guebh. ex Decne. in DC. Prod. xiii. 1. 698 = sibirica.
Cornuti, Jacq. Misc. ii. 351 = P. Coronopus.
coronopifolia, Brot. Fl. Lusit. i. 157 = macrorhiza.
coronopifolia, Murr. Syst. ed. XIII. 132 = P. Coronopus.
coronopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Coronopus.
Coronopus, Bertol. Amoen. Ital. 129 = macrorhiza.
Coronopus, *Linn. Sp. Pl.* 115.—Europ.; As. bor.; Austral.
Coronopus, Sauss. Voy. Alp. 66 = maritima.
crassa, Willd. Phyt. 3 = crispa.
crassifolia, *Forsk. Fl. Aegypt. Arab.* 31.—Europ.
crassifolia, Roth, in Usteri, Ann. Bot. xiv. (1795) 24 = crispa.
crassinervis, *Hornem. Hort. Hafn. Suppl.* 18.—Hab.?
crassipes, *Borb. in Oestr. Bot. Zeitschr.* xxviii. (1878) 392.—Europ.
crenata, Blanco, Fl. Filip. ed. I. 56 = major.
cretica, Chaub. ex Decne. in DC. Prod. xiii. 1. 706 = Bellardi.
cretica, *Linn. Sp. Pl.* 114.—Reg. Mediterr. or.
crispa, *Jacq. Coll. Suppl.* 34. t. 16.—Hab.?
crispa, Savi, ex Decne. in DC. Prod. xiii. 1. 734 = P. Cynops.
crithmoides, Desf. Fl. Atlant. i. 140 = macrorhiza.
crypsoides, *Boiss. Fl. Orient.* iv. 888.—Aegypt.
cucullata, Lam. Illustr. i. 339 = maxima.
cucullata, Pursh, Fl. Am. Sept. i. 98 = eriopoda.
Cumingiana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 44.—Chili.
Cunninghamii, Decne. in DC. Prod. xiii. 1. 702 = debilis?
Cupani, Guss. Fl. Sic. Prod. i. 190 = P. Coronopus.
curta, Engelm. ex Decne. in DC. Prod. xiii. 1. 713 = patagonica.
cylindrica, *Forsk. Fl. Aegypt. Arab.* 31.—Aegypt.; Arab.
Cynops, *Linn. Sp. Pl.* 116.—Europ.; Sibir.
Cynops, Sibth. & Sm. Fl. Graec. Prod. i. 103 = arenaria.
cynopsidea, Schult. Obs. Bot. 25 = P. Psyllium.
cynopsis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Cynops.
dalmatica, Schott, ex Steud. Nom. ed. II. ii. 348 = lanceolata.
Daltoni, Decne. in DC. Prod. xiii. 1. 703 = tasmanica.
dasyurus, Rafin. Cat. 15 = P. Lagopus.
debilis, *R. Br. Prod.* 425.—Austral.
debilis, Nees, in Lehm. Pl. Preiss. i. 491 = varia.
Decaisnei, *Barn. Monog. Plantag.* 45.—Chili.
decipiens, *Barn. l. c.* 16.—Am. bor.
declinata, Hort. ex Roem. & Schult. Syst. iii. 127 = ovata.
decumbens, Bernh. ex Reichb. Fl. Germ. Excurs. 396 = lanceolata.

PLANTAGO:—
decumbens, Forsk. Fl. Aegypt. Arab. 30 = ovata.
decumbens, C. Gay, ex Barn. Monog. Plantag. 44 = tumida.
dentata, Fl. Corc. in Ionios Anthol. iv. 946 = major.
dentata, Roth, Tent. Fl. Germ. i. 61 = maritima.
denticulata, Gaud. Fl. Helvet. i. 403 = maritima.
denticulata, Link, ex Schrank, Pl. Rar. Ic. 25; ex Decne. in DC. Prod. xiii. 1. 716, cum cit. falsa = lusitanica.
Deppeana, *Vatke, in Verh. Bot. Ver. Brandenb.* xvi. (1874) 53.—Calif.
depressa, *Willd. Enum. Hort. Berol. Suppl.* 8.—Sibir.
deserticola, *Phil. Fl. Atac.* 46.—Chili.
diffusa, *Presl, ex Steud. Nom.* ed. II. ii. 348, nomen.—Hab.?
divaricata, Zuccagni, in Roem. Collect. 126 = P. Psyllium.
Dregeana, *Presl, Bot. Bemerk.* 105; *Decne. in DC. Prod.* xiii. 1. 695.—Afr. austr.
Drummondi, Decne. l. c. 701 = varia.
dubia, Linn. Fl. Suec. ed. II. p. xvi (cf. 46) = lanceolata.
Durandoi, *Pomel, Nouv. Mat. Fl. Atl.* 125.—Alger.
Durvillei, Delile, ex Steud. Nom. ed. II. ii. 348 = hirtella?
echioides, *Decne. in DC. Prod.* xiii. 1. 722.—Texas.
elata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 62 = lanceolata.
elata, *Steud. in Flora*, xxxii. (1849) 413.—Abyss.
elongata, Pursh, Fl. Am. Sept. ii. 729 = pusilla.
eriantha, Decne. in DC. Prod. xiii. 1. 707 = ciliata.
eriocarpa, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 279 = P. Psyllium.
eriophora, Hoffmgg. & Link, Fl. Port. i. 423 = lanceolata.
eriopoda, *Torr. in Ann. Lyc. N. York*, ii. (1828) 237.—Am. bor.
eriorrhiza, Willd. ex Spreng. Syst. i. 437 = linearis.
eriostachya, Tenore, Prod. Fl. Nap. 13 = P. Lagopus.
erosa, Wall. in Roxb. Fl. Ind. ed. Carey, i. 423 = major.
Eschscholtziana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 45.—Chili.
euphratica, *Decne. ex Barn. Monog. Plantag.* 51.—Cappadoc.
evacina, *Boiss. Fl. Orient.* iv. 887.—Afghan.
exaltata, Hornem. Hort. Hafn. 140 = major.
exigua, Murr. in Comm. Gotting. (1778) 94. t. 5 = pumila.
exilis, Decne. in DC. Prod. xiii. 1. 702 = varia.
falcifolia, Salisb. Prod. 47 = carnosa.
fernandezia, *Bert. ex Decne. in DC. Prod.* xiii. 1. 704.—Ins. J. Fernand.
filiformis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 77 = major.
filiformis, Decne. in DC. Prod. xiii. 1. 714 = patagonica.
filiformis, C. Koch, in Linnaea, xxi. (1848) 709 = P. Coronopus.
firma, *Kunze, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 402.—Chili.
flexuosa, Gaud. ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 456 = lanceolata.
floccosa, *Decne. in DC. Prod.* xiii. 1. 723.—Mexic.
fornicata, C. Koch, in Linnaea, xxi. (1848) 712 = P. Lagopus.
Frankii, *Steud. in Flora*, xxxii. (1849) 409.—Am. bor.
frigida, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 402 = macrantha.
frigida, Willd. ex Schult. Mant. iii. 76 = nubigena.
fruticosa, Hort. Matrit. ex Roem. & Schult. Syst. iii. 119, in obs. = P. Lagopus.
fruticosa, Hort. Par. ex Roem. & Schult. l. c. 121 = lusitanica.
fuscescens, *Jord. Obs. Pl. Crit.* iii. (1846) 231.—Gallia.
Galeottiana, *Decne. in DC. Prod.* xiii. 1. 726.—Mexic.
garganica, Hort. ex Decne. l. c. 735 = P. Psyllium.
garganica, Tenore, ex Baxt. in Loud. Hort. Brit. Suppl. ii. 659, nomen = praec.?
Gaudichaudii, Barn. Monog. Plantag. 15 = varia.
Gayana, *Decne. in DC. Prod.* xiii. 1. 709.—Chili.
genevensis, Poir. Encyc. v. 390 = P. Cynops.
gentianoides, Decne. in DC. Prod. xiii. 1. 721 = brachyphylla.

PLANTAGO:—
gentianoides, *Sm. in Rees, Cycl.* xxvii. [n. 12]—As. Min.
Gerardi, Pourr. in Mém. Acad. Toul. iii. (1788) 324 = lanceolata.
Gerardi, Schult. Oestr. Fl. ed. II. i. 298 = subulata.
gibbosa, Rafin. in Am. Monthly Mag. (1818) 344 = maritima.
gigantea, *Decne. in DC. Prod.* xiii. 1. 724.—Bras.
glabra, Nutt. Gen. Am. i. 100 = eriopoda?
glabrata, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 276 = tasmanica.
glareosa, *Kern. ex Meynh. Kalocsa Vidék. Növényt.* 150.—Hungar.
glauca, C. A. Mey. Verz. Pfl. Cauc. 115 = P. Lagopus.
globosa, *Rafin. New Fl. Am.* iv. 11.—Am. bor.
glomerata, *Poir. Encyc.* v. 385.—Ins. Teneriff.
gnaphaloides, Nutt. Gen. Am. i. 100 = patagonica.
gonophylla, Rafin. Atl. Journ. 149 (Quid?).—Am. bor.
Gouani, J. F. Gmel. Syst. 251 = major.
Goudotiana, *Decne. in DC. Prod.* xiii. 1. 699.—N. Granat.
gracilis, Hort. ex Schult. Mant. iii. 70 = major.
gracilis, Poir. Voy. Barb. ii. 115 = maritima.
graminea, Lam. Illustr. i. 343 = maritima.
graminea, Willd. ex Schult. Mant. iii. 79 = xorulensis.
graminiformis, Lam. Fl. Fr. ii. 311 = maritima.
grandiflora, Meyen, Reise, i. 348 = macrantha.
Griffithsii, *Decne. in DC. Prod.* xiii. 1. 700.—Afghan.
Grisebachii, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 52.—Reg. Argent.
Guilleminiana, *Decne. l. c.* 722.—Bras.
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* v. (1846) 446.—Tasman.
Halleri, Rapin, ex Decne, in DC. Prod. xiii. 1. 729 = maritima.
Halleri, *Schleich. Cat. Pl. Helv.* ed. IV. 26.—Europ.
Hamiltoni, *T. Kirk, in Trans. N. Z. Inst.* xi. (1879) 465.—N. Zel.
Hartwegii, Decne. in DC. Prod. xiii. 1. 724 = hirtella.
Hasskarlii, *Decne. l. c.* 700.—Java.
Hausknechtii, *Vatke, in Verh. Bot. Ver. Brandenb.* xvi. (1874) 53.—Syria; Peruv.
heterophylla, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 177.—Am. bor.
hirsuta, Fl. Corc. in Ionios Anthol. iv. 947 = lanceolata, P. Coronopus.
hirsuta, Gilib. Fl. Lituan. i. 17 = maritima.
hirsuta, Gill. ex Decne. in DC. Prod. xiii. 1. 725 = affinis.
hirsuta, Kit. ex Steud. Nom. ed. II. ii. 349 = P. Psyllium.
hirsuta, Pourr. Chlor. Narb. ex Barn. Monog. Plantag. 24 = subulata.
hirsuta, Ruiz & Pav. Fl. Per. i. 51 = limensis.
hirsuta, Thunb. Prod. Pl. Cap. 29 = carnosa.
hirtella, Barn. Monog. Plantag. 9 = Berteroi.
hirtella, *H. B. & K. Nov. Gen. et Sp.* ii. 229. t. 127.—Am. bor. et austr.
hispida, R. Br. Prod. 425 = varia.
hispidula, Bert. ex Steud. in Flora, xxxii. (1849) 405 = macrosperma.
hispidula, *Ruiz & Pav. Fl. Per.* i. 51. t. 78.—Chili.
holosericea, Gaud. ex Roem. & Schult. Syst. iii. 126 = argentea.
Holostea, Lam. Illustr. i. 340 = Bellardi.
Holosteum, Scop. Fl. Carn. ed II. i. 108 = subulata.
Hookeriana, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 39 = patagonica.
humifusa, Bernh. ex Reichb. Fl. Germ. Excurs. 863 = major.
humilis, *Decne. in DC. Prod.* xiii. 1. 724.—Peruv.
humilis, Jan, ex Decne. l. c. 730 = subulata.
hungarica, Waldst. & Kit. Pl. Rar. Hung. iii. 225 = lanceolata.
hybrida, Bart. Fl. N. Am. iii. 77; Comp. Fl. Philad. ii. 214 = pusilla.
hygrophila, *Steud. in Flora*, xxxii. (1849) 403.—Chili.
imberbis, Hook. f. Fl. Antarct. 66 = barbata.
incana, Ram. ex DC. Fl. Fr. iii. 414 = alpina.
incana, Roem. & Schult. Syst. iii. 134 = subulata.
incana, Stokes, Bot. Mat. Med. i. 212 = media.
incisa, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 23.—Java.
incomta, Willd. ex Ledeb. Fl. Ross. iii. 476 = tenuiflora.

PLANTAGO:—
incurvata, Murr. in Comm. Gotting. iii. 1780 (1781) 19. t. 6 = subulata.
indica, Linn. Syst. ed. X. 896 = pumila?
integralis, Gaud. Fl. Helvet. i. 403 = maritima.
intermedia, Gilib. Pl. Europ. i. 125 = major.
intermedia, C. Koch, in Linnaea, xxi. (1848) 711 = lanceolata.
intermedia, Lapeyr. Hist. Abr. Pl. Pyr. 69 = P. Lagopus.
intermedia, *Steud. in Flora*, xxxii. (1849) 414.—Austral.
interrupta, Poir. Encyc. v. 375 = sparsiflora.
irrigua, Fisch. ex Reichb. Fl. Germ. Excurs. 396 = lanceolata.
Ispaghula, Roxb. ex Flem. in As. Res. xi. (1810) 174 = ovata.
italica, S. G. Gmel. ex Steud. Nom. ed. I. 630 = arenaria.
Jacquini, Roem. & Schult. Syst. iii. 140 = P. Coronopus.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 469.—Japon.
juncoides, Decne. in DC. Prod. xiii. 1. 731, partim = decipiens.
juncoides, Lam. Illustr. i. 342 = maritima.
kamtschatica, vide *camtschatica*.
kamtschatika, Hook. Comp. Bot. Mag. ii. (1836) 60 = Rugelii.
karakabanica, Boiss. & Huet, ex Boiss. Fl. Orient. iv. 881 = saxatilis.
kentukensis, Michx. Fl. Bor. Am. i. 94 = cordata.
Kerstenii, Aschers. in Sitzb. Ges. Naturf. Fr. Berl. (1868) 23 = palmata.
kurdica, Vatke, in Verh. Bot. Ver. Brandenb. xvi. (1874) 52 = lanceolata.
lachnantha, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 446.—Turkestan.
laciniata, Willk. in Bot. Zeit. vi. (1848) 413 = P. Serraria.
Lagascae, Duf. ex Barn. Monog. Plantag. 24 = subulata.
lagocephala, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 445.—Turkestan.
lagopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Lagopus.
lagopodioides ×, Schrank, in Denkschr. Akad. Muench. 1813 (1814) 71 = lusitanica.
lagopoides, Desf. Fl. Atlant. i. 135 = amplexicaulis.
lagopoides, Viv. Fl. Lib. Spec. 7 = P. Lagopus.
Lagopus, *Linn. Sp. Pl.* 114.—Reg. Mediterr.; Oriens.
Lagopus, Pursh, Fl. Am. Sept. i. 99 = patagonica.
Lagurus, Roth, Nov. Pl. Sp. 87 = lusitanica.
Lamarkii, Jord. Obs. Pl. Nouv. iii. 228 = monosperma.
lanata, Hort. Monsp. ex Barn. Monog. Plantag. 43 = nivea.
lanata, Host, Fl. Austr. i. 210 = lanceolata.
lanata, Lag. & Rodr. in Anal. Cienc. Nat. v. (1802) 269 = nivalis.
lanata, Poir. Voy. Barb. ii. 115 = pilosa.
lanata, Raddi, Osserv. Isola Madera (1821), 11; ex Schult. Mant. iii. 78 = Raddiana.
lanata, Willd. ex Spreng. Syst. i. 439; Wight, in Wall. Cat. n. 6414 = ovata.
lanceaefolia, Salisb. Prod. 47 = lanceolata.
lanceolata, Hook. Fl. Bor. Am. ii. 123, partim = eriopoda.
lanceolata, *Linn. Sp. Pl.* 113.—Europ.; As. bor.
lanigera, *Hook. f. Handb. N. Zeal. Fl.* 227.—N. Zel.
lanuginosa, DC. Fl. Fr. Suppl. 377, in syn. = lanceolata.
lasiothrix, *Kunze, ex Decne. in DC. Prod.* xiii. 1. 709.—Chili.
latibracteata, *Trautv. Pl. Imag.* 20. t. 14.—Sibir.
latifolia, Fisch. Livl. Zust. 107; ex Ledeb. Fl. Ross. iii. 481 = media.
latifolia, Salisb. Prod. 46 = major.
laxiflora, *Decne. in DC. Prod.* xiii. 1. 699.—Afr. austr.
leiocephala, *Wallr. ex C. Koch, in Linnaea*, xxi. (1848) 713.—Reg. Caucas.
leiopetala, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1833) 17.—Ins. Canar.
leonensis, *Steud. in Flora*, xxxii. (1849) 404.—Chili.
leptophylla, *Decne. in DC. Prod.* xiii. 1. 723.—Bras.

PLANTAGO:—
leptostachys, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 277 = tasmanica.
leptostachys, *Ledeb. Fl. Ross.* iii. 479.—Sibir. altaic.
leptostachys, *E. Mey. ex Decne. in DC. Prod.* xiii. 1. 720.—Afr. austr.
Lessingii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 47.—Soongar.
leucophylla, *Decne. in DC. Prod.* xiii. 1. 708.—Venezuela.
limensis, *Pers. Syn.* i. 139.—Peruv.
limosa, Kit. ex Roem. & Schult. Syst. iii. 111 = major.
Lindeniana, Decne. ex Wedd. Chlor. And. ii. 164 = sericea.
linearifolia, Muhl. Cat. 15; ex Torr. Fl. U. St. 185 = pusilla.
linearis, *H. B. & K. Nov. Gen. et Sp.* ii. 229.—Ecuador.
linearis, Rafin. New Fl. Am. iv. 11 (Quid?).—Am. bor.
Linkii, Roem. & Schult. Syst. iii. 117, in obs. = lanceolata.
litorea, *Phil. Fl. Atac.* 46.—Chili.
littoralis, *Roxb. Hort. Beng.* [10], nomen.—Afr. austr.
Loeflingii, Bieb. Beschr. Casp. App. 208 = tenuiflora.
Loeflingii, Huds. Fl. Angl. ed. I. 52 = maritima.
Loeflingii, *Linn. Sp. Pl.* 115.—Reg. Mediterr.; Oriens.
Loeflingii, Thunb. Prod. Pl. Cap. 29 = cafra.
longestyla, *Presl, Bot. Bemerk.* 105.—Aegypt.
longifolia, Decne. in DC. Prod. xiii. 1. 717 = macrocarpa.
longiscapa, Fl. Corc. in Ionios Anthol. iv. 947 = lanceolata.
longiscapa, Jacquem. ex Decne. in DC. Prod. xiii. 1. 696 = major.
longissima, *Decne. l. c.* 720.—Afr. austr.
longistipes, Royle, ex Barn. Monog. Plantag. 28 = lanceolata.
Loscosii, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 358.—Hispan.
Loureiri, *Roem. & Schult. Syst.* iii. 112.—China; Japon.
ludoviciana, Rafin. Fl. Ludov. 33 = virginica.
lusitanica, *Linn. Sp. Pl.* ed. II. 1667.—Lusitan.
lusitanica, Willd. Sp. Pl. i. 644 = P. Lagopus.
lutulenta, Lamotte, Prod. Fl. Centr. Fr. ii. 618 = major.
luzuloidea, *Decne. in DC. Prod.* xiii. 1. 710.—N. Granat.
macrantha, *Decne. ex Barn. Monog. Plantag.* 45.—Chili.
macrocarpa, *Cham. & Schlecht. in Linnaea*, i. (1826) 166.—Am. bor. occ.
macrorhiza, *Poir. Voy. Barb.* ii. 114 (machroriza).—Reg. Mediterr.
macrosperma, *Steud. in Flora*, xxxii. (1849) 405.—Chili.
macrostachys, *Decne. in DC. Prod.* xiii. 1. 724.—Reg. Argent.
maderensis, *Decne. l. c.* 733.—Ins. Madera.
madritensis, Lag. ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 459 = P. Lagopus.
major, Bert. ex Barn. Monog. Plantag. 9 = Berteroi.
major, Ell. Sketch, i. 201 = Rugelii.
major, *Linn. Sp. Pl.* 112.—Europ.; As. et Am. bor.
major, Lour. Fl. Cochinch. 71 = Loureiri.
marginata, *Steud. in Flora*, xxxii. (1849) 407.—Chili.
marina, Woodw. in With. Bot. Arr. Brit. Pl. ed. II. i. 147 = maritima.
maritima, Auct. Plur. Am. = decipiens.
maritima, Barton, Fl. Philad. Prod. 26 = pusilla.
maritima, *Linn. Sp. Pl.* 114.—Europ.; Am. bor. et arct.
maritima, Willk. in Flora, xxxv. (1852) 291 = crassifolia.
matritensis, Lag. ex Decne. in DC. Prod. xiii. 1. 716 = lusitanica.
mauritanica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 105.—Afr. bor. occ.
maxima, *Jacq. Coll.* i. 82.—Sibir.
maxima, Ruching. Fl. Venet. 45 = major.
media, Blanco, Fl. Filip. ed. II. 38 = major.
media, Hook. & Arn. Bot. Beech. Voy. 43 = Candollii.
media, *Linn. Sp. Pl.* 113.—Europ.; As. bor.
mexicana, *Link, Enum. Hort. Berol.* i. 121.—Mexic.
micrantha, *Ledeb. ex Steud. Nom.* ed. II. ii. 349, nomen.—Sibir.
microcephala, Hort. ex Decne. in DC. Prod. xiii. 1. 719 = Loeflingii.

PLANTAGO:—
microcephala, Poir. Encyc. v. 378 = ovata.
microcephala, Royle, ex Barn. Monog. Plantag. 29 = lanceolata.
minima, DC. Fl. Fr. iii. 408 = major.
minima, Less. ex Barn. Monog. Plantag. 41 = Lessingii.
minor, Fries, Mant. ii. 12 = tenuiflora.
minor, Gilib. Fl. Lituan. i. 15 = major.
minuta, Link, in Schrad. Journ. Bot. (1800) 1. 59 = Bellardi.
minuta, C. A. Mey. Verz. Pfl. Cauc. 115 = Loeflingii.
minuta, *Pall. Reise*, iii. 716.—Reg. Caucas.
missouriensis, *Steud. in Flora*, xxxii. (1849) 409.—Am. bor.
Mitchellii, Decne. in DC. Prod. xiii. 1. 701 = varia.
Mohnikei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 119.—Japon.
mollis, *Barn. Monog. Plantag.* 45; *Decne. in DC. Prod.* xiii. 1. 713.—Chili.
mollis, Hook. & Arn. Bot. Beech. Voy. 43 = patagonica.
monanthos, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 606.—Reg. Magellan.
mongolica, *Decne. in DC. Prod.* xiii. 1. 707.—Mongolia.
monosperma, *Pourr. in Mém. Acad. Toul.* iii. (1788) 325.—Mont. Pyren.
monspeliensis, Willd. Enum. Hort. Berol. 160 = ovata.
montana, Griseb. Spicil. Fl. Rumel. ii. 303 = saxatilis.
montana, *Huds. Fl. Angl.* ed. I. 53; *Lam. Illustr.* i. 341.—Europ. med.
monticola, *Decne. in DC. Prod.* xiii. 1. 711.—Peruv.
multiceps, *H. B. & K. Nov. Gen. et Sp.* ii. 228.—Mexic.
multinervis, Rafin. New Fl. Am. iv. 10 (Quid?).—Am. bor.
multiplinervia, *Steud. in Flora*, xxxii. (1849) 410.—Am. bor.
Myosurus, *Lam. Illustr.* i. 342.—Reg. Argent.
nana, Bertol. Misc. Bot. i. 12 = Loeflingii.
nana, Tratt. Archiv, t. 42 = major.
neglecta, Guss. Fl. Sic. Prod. i. 191 = P. Coronopus.
nigricans, Herb. ex Roem. & Schult. Syst. iii. 117 = montana.
nigricans, Link, ex Reichb. Fl. Germ. Excurs. 396 = lanceolata.
nitida, *Hort. ex Roem. & Schult. Syst.* iii. 149.—Hab.?
nivalis, *Boiss. Voy. Bot. Espagne*, ii. 533.—Hispan.
nivalis, Jord. Obs. Pl. Nouv. iii. 232 = monosperma.
nivea, *H. B. & K. Nov. Gen. et Sp.* ii. 228.—Mexic.
notata, *Lag. Gen. et Sp. Nov.* 7.—Reg. Mediterr. et Caucas.
nubigena, *H. B. & K. Nov. Gen. et Sp.* ii. 227.—Ecuador.
nuda, *Luce, Topog. Nachr. Oesel*, 37 (Quid?).—Ins. Balt.
numidica, Poir. ex Decne. in DC. Prod. xiii. 1. 729 = maritima.
nutans, Poir. Encyc. v. 381 = lanceolata.
Nuttallii, Rapin, in Mém. Soc. Linn. Par. vi. (1827) 470 = patagonica.
oblongifolia, *Decne. in DC. Prod.* xiii. 1. 700.—Am. bor.
oblongifolia, Schur, in Oestr. Bot. Zeitschr. x. (1860) 249 = media.
obscura, *Steud. in Flora*, xxxii. (1849) 407.—Chili.
obtusata, *Decne. in DC. Prod.* xiii. 1. 711.—Chili.
occidentalis, Decne. l. c. 722 = virginica.
officinarum, Crantz, Inst. ii. 329 = major.
oliganthos, Roem. & Schult. Syst. iii. 122 = maritima.
Olivieri, Decne. ex Barn. Monog. Plantag. 37 = notata.
Orbignyana, Desf. ex Barn. l. c. 10, in obs. = seq.?
Orbignyana, *Steinh. ex Decne. in DC. Prod.* xiii. 1. 704.—Peruv.
Oreades, *Decne. l. c.* 699.—N. Granat.
orientalis, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 33.—Persia.
ovata, *Forsk. Fl. Aegypt. Arab.* 31.—Reg. Mediterr.; Oriens.
ovatifolia, *Steud. in Flora*, xxxii. (1849) 412.—Arabia.
ovina, Vill. Prosp. 19 = alpina.
pachyneura, *Steud. in Flora*, xxxii. (1849) 406.—Chili.

PLANTAGO:—
pachyphylla, *A. Gray, in Proc. Am. Acad.* vi. (1861) 54.—Ins. Sandvic.
pallida, Salisb. Prod. 47 = albicans.
palmata, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 19.—Afr. trop.
paludosa, *Turcz. ex Ledeb. Fl. Ross.* iii. 478.—Sibir. baical.
paradoxa, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 277 = Brownii.
Paralias, *Decne. in DC. Prod.* xiii. 1. 725.—Reg. Argent.
parviflora, Desf. Fl. Atlant. i. 141 = P. Psyllium.
parviflora, Hort. ex Roem. & Schult. Syst. iii. 147 = stricta.
parvula, Roem. & Schult. l. c. 125 = albicans.
parvula, Wedd. ex Reichb. Fl. Germ. Excurs. 396 = Weldenii.
patagonica, Bert. ex Steud. in Flora, xxxii. (1849) 405 = P. Rancaguae.
patagonica, *Jacq. Ic. Pl. Rar.* ii. 9. t. 306; *Coll. Suppl.* 35.—Am. bor. et austr.
pauciflora, Gilib. Fl. Lituan. i. 16 = major.
pauciflora, Lam. Illustr. i. 342 = barbata.
pauciflora, Pursh, Fl. Am. Sept. i. 99 = maritima, decipiens.
penicillata, Endl. & Fenzl, Sert. Cabul. 1 = ciliata.
pentasperma, *Hemsl. Bot. Voy. Challenger*, i. 11. 265.—Ins. Amsterd.
perpusilla, Decne. in DC. Prod. xiii. 1. 697 = heterophylla.
Perreymondii, *Barn. Monog. Plantag.* 46.—Venezuela.
phaeopus, Paine, in Proc. Palest. Exp. Soc. App. iii. 120 = Haussknechtii.
phaeostoma, *Boiss. & Heldr. Diagn.* Ser. II. iv. 71.—Aegypt.; Arab.
Philippiana, *Vatke, in Verh. Bot. Verh. Brandenb.* xvi. (1874) 49.—Chili.
philippica, *Cav. Ic.* iv. 35. t. 359.—Ins. Philipp.
pilosa, Kit. ex Spreng. Syst. i. 435 = argentea.
pilosa, Pourr. in Mém. Acad. Toul. iii. (1788) 324 = Bellardi.
platypetala, Wallr. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 402 = macrantha.
plicata, Schott, Nym. & Kotschy, Analect. Bot. 4 = brutia.
polymorpha, Banks & Soland. ex Hook. f. Fl. Antarct. 339 = barbata.
polysperma, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 431.—Sibir. altaic.
polystachia, Fl. Corc. in Ionios Anthol. iv. 946 = major.
polystachya, *Viv. Pl. Aegypt. Dec.* 22.—Aegypt.
praecox, C. A. Mey. Verz. Pl. Cauc. 115 = notata.
Preslii, Tenore, Prod. Fl. Nap. Suppl. v. p. vii = lanceolata.
Princeps, *Cham. & Schlecht. in Linnaea*, i. (1826) 167.—Ins. Sandvic.
prionota, *Pomel, Nouv. Mat. Fl. Atl.* 310.—Alger.
procera, Salisb. Prod. 46 = maxima.
prolifera, Webb, ex Barn. Monog. Plantag. 50 = Webbii.
Pseudo-lusitanica, Roem. & Schult. Syst. iii. 121 = P. Lagopus.
Pseudo-patagonica, *Vatke, in Verh. Bot. Ver. Bradenb.* xvi. (1874) 50.—Chili.
Pseudo-psyllium, Desf. Cat. Hort. Par. ed. III. 77 = P. Psyllium.
psyllia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Psyllium.
Psyllium, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 245 = arabica.
Psyllium, DC. Fl. Fr. iii. 416, in syn. = arenaria.
Psyllium, Herb. Aegypt. ex Presl, Bot. Bemerk. 105 = longestyla.
Psyllium, *Linn. Sp. Pl.* 115.—Reg. Mediterr.; Oriens; Ind. bor. occ.
pubescens, Lapeyr. Hist. Abr. Pl. Pyr. 71 = serpentina.
pulla, *Fl. Corc. in Ionios Anthol.* iv. 950 (Quid?).—Ins. Corcyra.
pumila, *Linn. f. Suppl.* 125.—Reg. Mediter.; Oriens; Ind. bor. occ.
pumila, Weld. ex Reichb. Fl. Germ. Excurs. 396 = Weldenii.
pungens, Lapeyr. Hist. Abr. Pl. Pyr. 71 = subulata.
purpurascens, Nutt. ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 454; et in Trans. Am. Phil. Soc. v. (1837) 177 = virginica.
purpurascens, *Willk. in Oestr. Bot. Zeitschr.* xxv. (1875) 110.—Ins. Balear.

PLANTAGO:—
Purshii, Roem. & Schult. Syst. iii. 120 = patagonica.
pusilla, Bunge, Verz. Suppl. Fl. Alt. 13 = Bungei.
pusilla, *Nutt. Gen. Am.* i. 100.—Am. bor.
pygmaea, Lam. Illustr. i. 341 = Bellardi.
Queleniana, Gaudich. in Freyc. Voy. Bot. 445 = Princeps.
Quillotae, *Vatke, in Verh. Bot. Ver. Brandenb.* xvi. (1874) 50.—Chili.
quinquenervia, Schleich. Cat. 38; ex Roem. & Schult. Syst. iii. 117 = montana.
quinquenervis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 25 = major.
Raddiana, *Schult. Mant.* iii. 78.—Ins. Madera.
radicata, Hoffmgg. & Link, Fl. Port. i. 428 = subulata.
ramosa, Aschers. Fl. Brandenb. 547 = arenaria.
Rancaguae, *Steud. in Flora*, xxxii. (1849) 405.—Chili.
Raoulii, *Decne. in DC. Prod.* xiii. 1. 703.—N. Zel.
rectirostris, Wallr. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 401 = tomentosa.
recurvata, Linn. Mant. ii. 198 = subulata.
recurvata, Willd. ex Spreng. Syst. i. 437 = serpentina.
remota, Bieb. ex Spreng. l. c. in syn. = tenuiflora.
remota, *Lam. Illustr.* i. 341.—Afr. austr.
remotiflora, Stocks, in Hook. Kew Journ. iv. (1852) 179 = Stocksii.
rhodosperma, *Decne. in DC. Prod.* xiii. 1. 722.—Texas.
Richardsoni, Decne. l. c. 698 = eriopoda.
rigens, Schult. Mant. iii. 83 = rigida.
rigida, Hort. ex Roem. & Schult. Syst. iii. 144 = P. Psyllium.
rigida, *H. B. & K. Nov. Gen. et Sp.* ii. 227.—Ecuador.
rigidiuscula, D. Dietr. Syn. Pl. i. 435 = rigida.
robusta, *Roxb. in Beats. Tracts St. Helen. App.* 317.—Ins. S. Helena.
Rocae, *Lorentz, in Roca, Exped. Rio-Negro*, ii. 250.—Patagon.
rosea, Hort. ex Reichb. Fl. Germ. Excurs. 394 = major.
rosetana, Poir. Encyc. Suppl. iv. 433 = pumila.
rubens, Hassk. Cat. Hort. Bogor. Alt. 95 = Hasskarlii.
rubens, Host, Fl. Austr. i. 211 = argentea.
Rugelii, *Decne. in DC. Prod.* xiii. 1. 700.—Am. bor.
rugosa, Hochst. in Flora, xxiv. (1841) 1. Intell. 22, nomen; et ex Steud. l. c. xxxii. (1849) 413 = stricta.
runcinata, Decne. in DC. Prod. xiii. 1. 702 = varia.
ruthenica, Adams, ex Decne. l. c. 735 = arenaria.
salicifolia, Decne. l. c. 729, in syn., sphalm., vide *falcifolia*.
salina, Decne. l. c. 720 = amplexicaulis.
salsa, Pall. Reise, i. 486 = maritima.
sarcophylla, Boiss. Fl. Orient. iv. 893 = squarrosa.
sarda, *Presl, Bot. Bemerk.* 105.—Sardinia.
saxatilis, *Bieb. Fl. Taur. Cauc.* i. 109.—Europ. or.; Persia.
scabra, Moench, Meth. 461 = pumila?
Schiedeana, *Decne. in DC. Prod.* xiii. 1. 723.—Mexic.
Schottii, Roem. & Schult. Syst. iii. 118 = lanceolata.
Schrenkii, *C. Koch, in Linnaea*, xxi. (1848) 710.—Ins. Waigatsch.
Schwarzenbergiana, *Schur, in Verh. Siebenb. Ver. Naturw.* vi. (1855) 3; *et in Oestr. Bot. Wochenbl.* vi. (1856) 282.—Europ.
scirpoides, Lam. Illustr. i. 342 = saxatilis.
scorzoneraefolia, Lam. l. c. = maritima.
Seetzenii, Hort. ex Decne. in DC. Prod. xiii. 1. 733 = Webbii.
Selloi, *J. A. Schmidt, in Mart. Fl. Bras.* vi. iv. 171.—Bras.
sempervirens, *Crantz, Inst.* ii. 331 (Quid?).—Europ.
sericea, Benth. Cat. Pl. Pyr. 112 = monosperma.
sericea, *Ruiz & Pav. Fl. Per.* i. 51.—Peruv.
sericea, Waldst. & Kit. Pl. Rar. Hung. ii. 163 = argentea.
sericophylla, Decne. in DC. Prod. xiii. 1. 702 = varia.
serpentina, Koch, Syn. Fl. Germ. ed. I. 599 = carinata.
serpentina, Lam. Illustr. i. 343 = subulata.
serpentina, Vill. Prosp. 19; All. Fl. Pedem. i. 82 = maritima.
Serraria, *Linn. Syst.* ed. X. 896.—Reg. Mediterr.
serrata, Hall. f. ex Gaud. Fl. Helv. i. 403 = subulata.

PLANTAGO:—
sessiliflora, Lapeyr. Hist. Abr. Pl. Pyr. 72 = subulata.
setosa, Bertol. Misc. Bot. i. 11 = cylindrica *vel* ciliata?
setosa, *Clarke, Trav.* ii. 643 = cylindrica.
sibirica, Fisch. ex Barn. Monog. Plantag. 13 = depressa.
sibirica, Georgi, Beschr. Russ. Reich. III. iv. 736 = media.
sibirica, *Hort. ex Poir. Encyc. Suppl.* iv. 433.—Sibir.
sicula, Presl, Delic. Prag. 70 = P. Psyllium.
sinaica, Decne. in DC. Prod. xiii. 1. 733 = arabica.
sinuata, Lam. Illustr. i. 338 = major.
Sorokini, *Bunge, in Act. Hort. Petrop.* vi. (1879) 393.—Turkestan.
sparsiflora, *Michx. Fl. Bor. Am.* i. 94.—Am. bor.
spathulata, *Hook. f. Fl. N. Zel.* i. 208.—N. Zel.
sphaerocephala, Poir. Encyc. v. 381 = montana.
sphaerostachys, *Hegetschw. Fl. Schw.* 116.—Europ.
spinulosa, Decne. in DC. Prod. xiii. 1. 713 = patagonica.
squalida, Salisb. Prod. 47 = P. Psyllium.
squamata, O. F. Muell. in Fl. Dan. t. 691 = maritima.
squamosa, Decne. in DC. Prod. xiii. 1. 714, sphalm., vide *squarrosa*, Bové.
squarrosa, Bové, ex Decne. l. c. 733 = arabica.
squarrosa, *Murr. in Comm. Gotting.* (1781) 38. t. 3.—Syria; Aegypt.
squarrosa, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 178 = patagonica.
Stauntoni, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxi. (1871) 33.—Ins. Amsterdam.
Steinheilii, Barn. Monog. Plantag. 38 = hispidula.
stellaris, F. Muell. Fragm. ii. 23 = Brownii.
stellaris, Salisb. Prod. 47 = P. Coronopus.
Stocksii, *Boiss. ex Decne. in DC. Prod.* xiii. 1. 720.—Persia; Afghan.
stricta, Boutelou, ex Willk. & Lange, Prod. Fl. Hisp. ii. 351 = P. Psyllium.
stricta, *Schousb. Vextr. Marokko*, 69.—Reg. Mediterr.; Persia.
stricta, Sieber, ex Decne. in DC. Prod. xiii. 1. 736 = pumila.
struthionis, A. Cunn. ex Decne. l. c. 702 = varia.
strutia, Walp. Rep. iv. 180, in syn., sphalm. = brutia.
stylosa, Bess. ex Decne. in DC. Prod. xiii. 1. 699 = major.
subsinuata, Hornem. Hort. Hafn. i. 139 = major.
subtrinervis, *Phil. in Linnaea*, xxxiii. (1864–65) 221.—Chili.
subulata, *Linn. Sp. Pl.* 115.—Europ. austr.
subulata, Roth, Tent. Fl. Germ. i. 62 = maritima.
subulata, Wulf. in Jacq. Coll. i. 204. t. 10 = carinata.
suffruticosa, Lam. Fl. Fr. ii. 313 = P. Cynops.
sylvatica, H. Mart. Fl. Mosq. 29 = lanceolata.
syrtica, Viv. Fl. Lib. Spec. 7. t. 3 = notata.
Tabernaemontani, Baumg. Enum. Stirp. Transs. i. 89 = major.
tasmanica, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 276.—Austral.
tatarica, *Decne. in DC. Prod.* xiii. 1. 696.—Reg. Himal.
tenuiflora, *Waldst. & Kit. Pl. Rar. Hung.* i. 37. t. 39.—Europ. or.; As. bor.
tenuis, Hoffmgg. & Link, Fl. Port. i. 426 = Bellardi.
teretifolia, Sieber, ex Rapin, in Mém. Soc. Linn. Par. vi. (1827) 473 = maritima.
texensis, Rafin. New. Fl. Am. iv. 11 (Quid?).—Texas.
Thunbergii, Poir. Encyc. Suppl. iv. 431 = carnosa.
tibetica, *Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind.* iv. 706.—Reg. Himal.
Timbali, Jard. Pugill. Pl. Nov. 138 = lanceolata.
tomentosa, Cham. & Schlecht. in Linnaea, i. (1826) 169 = Cumingiana, Eschscholtziana.
tomentosa, Gilib. Fl. Lituan. i. 17 = lanceolata.
tomentosa, Isabelle, ex Decne. in DC. Prod. xiii. 1. 725 = P. Paralias.
tomentosa, *Lam. Illustr.* i. 340.—Reg. Argent.
tomentosa, Schiede, ex Decne. in DC. Prod. xiii. 1. 723 = Schiedeana.
transsilvanica, Schur, in Oestr. Bot. Zeitschr. x. (1860) 249 = media.
triandra, *Bergqr. in Minneskr. Fisiog. Sällks. Lund*, (1877) n. 8. p. 16.—N. Zel.
triantha, Spreng. Syst. i. 439 = Brownii.
trinervis, Fl. Corc. in Ionios Anthol. iv. 947 = lanceolata.

PLANTAGO:—
triquetra, Pers. Syn. i. 139 = subulata.
truncata, Barn. Monog. Plantag. 14 = brachystachys.
truncata, *Cham. & Schlecht. in Linnaea*, i. (1826) 170.—Chili.
tubulosa, *Decne. in DC. Prod.* xiii. 1. 728.—Peruv.
tumida, *Link, Enum. Hort. Berol.* i. 121.—Chili.
uliginosa, Baumg. Enum. Stirp. Transs. i. 89 = gentianoides.
uliginosa, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 199 = major.
uncialis, *Decne. ex Barn. Monog. Plantag.* 42.—Chili.
uniflora, *Hook. f. Fl. N. Zel.* i. 207.—N. Zel.
uniflora, Linn. Sp. Pl. 115 = Littorella juncea.
uniglumis, *Wallr. ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 402.—Chili.
Urvillei, Decne. in DC. Prod. xiii. 1. 723 = hirtella.
vaginata, Vent. Jard. Cels, 29. t. 29 = P. Lagopus.
valparadisiaca, *Decne. in DC. Prod.* xiii. 1. 725.—Chili.
varia, *R. Br. Prod.* 424.—Austral.
varia, A. Cunn. ex Hook. f. Fl. N. Zel. i. 208 = Raoulii.
variabilis, Link, ex Barn. Monog. Plantag. 29 = lanceolata.
velutina, *Poir. Encyc.* v. 379.—Italia.
velutina, Tenore, Prod. Fl. Nap. 13 = albicans.
veratrifolia, *Decne. in DC. Prod.* xiii. 1. 721.—Mexic.
vestita, Roem. & Schult. Syst. iii. 149 = congesta.
Victorialis, Kotschy, ex Decne. in DC. Prod. xiii. 1 705 = cylindrica.
Victorialis, Poir. Encyc. v. 377 = argentea.
villifera, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 87.—Japon.
villosa, Lam. Illustr. i. 341 = Bellardi.
villosa, Moench, Meth. 459 = ovata.
virescens, Barn. Monog. Plantag. 33 = eriopoda.
virginica, Giseck. ex Roem. & Schult. Syst. iii. 122 = depressa.
virginica, *Linn. Sp. Pl.* 113.—Am. bor. et austr.
virginica, Riehl, ex Steud. in Flora, xxxii. (1849) 410 = missourensis.
virginica, Walt. Fl. Carol. 85 = sparsiflora?
Webbii, *Barn. Monog. Plantag.* 50.—Ins. Canar.
Weddelliana, *Decne. in DC. Prod.* xiii. 1. 712.—Bolivia.
Weldenii, *Reichb. Fl. Germ. Excurs.* 396.—Dalmatia.
Wrightiana, Decne. in DC. Prod. xiii. 1. 712 = patagonica.
Wulfenii, Bernh. ex Willd. Enum. Hort. Berol. i. 161 = maritima.
Wulfenii, Spreng. Fl. Hal. 54 = carinata.
Wulfenii, Sturm, Deutschl. Fl. Heft 24 = subulata.
xorulensis, Barn. Monog. Plantag. 44 = linearis.
xorulensis, *H. B. & K. Nov. Gen. et Sp.* ii. 228.—Mexic.

PLANTIA, Herb. in Lindl. Bot. Reg. (1844) Misc. 89 = **Hexaglottis**, Vent. (Irid.).
flava, Herb. l. c. = H. longifolia.

PLANTINIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317 = **Phleum**, Linn. (Gramin.).
aspera, Bubani, l. c. = Phleum asperum.

PLAPPERTIA, Reichb. Consp. 146 (1828) = **Chailletia**, DC.

PLASO, Adans. Fam. ii. 325 (1763) = **Butea**, Roxb. (Legumin.).

PLATANARIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 39 (1821) = **Sparganium**, Linn. (Typhac.).
natans, S. F. Gray, l. c. 40 = S. natans.

PLATANOCARPUM, Korth. Obs. Naucl. Ind. 18 (1839) = **Sarcocephalus**, Afzel. (Rubiac.).
africanum, Hook. f. in Hook. Ic. Pl. t. 787 = Mitragyna africana.
cordatum, Korth. Obs. Naucl. Ind. 19 = S. cordatus.
subditum, Korth. l. c. = S. subditus.

PLATANOCARPUS, Korth, Verh. Nat. Gesch. Bot. 152. t. 32 (1840?) = praec.

PLATANOCEPHALUS, Vaill. ex Crantz, Inst. i. 349 (1766) = **Anthocephalus**, A. Rich. (1834) (Rubiac.).
orientalis, Crantz, l. c. = A. morindaefolius.

PLATANOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = **Platanus**, Tourn.

PLATANTHERA, Rich. in Mém. Mus. Par. iv. 48 (1818) = **Habenaria**, Willd. (Orchid.).
acuifera, Lindl. in Wall. Cat. n. 7045 = H. acuifera.
acuminata, Lindl. l. c. n. 7040; Gen. et Sp. Orch. 289. —Ind. or.
affinis, Wight, Ic. t. 1693 = H. Galeandra.
albida, Lindl. Syn. Brit. Fl. ed. I. 261 = H. albida.
angustata, Lindl. Gen. et Sp. Orch. 290.—Java.
angustifolia, Reichb. f. Otia Bot. Hamb. 39 = Herminium angustifolium.
aphylla, Lindl. Gen. et Sp. Orch. = H. aphylla, *R. Br.*
arcuata, Lindl. l. c. 289.—Ind. or.
bifolia, Rich. in Mém. Mus. Par. iv. (1818) 57 = H. bifolia.
Bigelovii, Wood, Class-Book (1861), 685 = H. fissa?
Blephariglottis, Lindl. Gen. et Sp. Orch. 291 = H. Blephariglottis.
Blumii, Lindl. l. c. 290.—Java.
borealis, Reichb. f. Ic. Fl. Germ. xiii–xiv. 125 = H. dilatata.
brachyglossa, Reichb. Ic. Pl. Crit. ix. (1831) 19 = H. bifolia.
brachyphylla, Lindl. Gen. et Sp. Orch. 293 = H. crassifolia.
bracteata, Torr. Fl. N. York, ii. 279 = H. viridis.
canarensis, Lindl. ex Reichb. f. in Flora, xli. (1877) 100 [Pl. Hohen. n. 142] = H. crassifolia.
candida, Lindl. Gen. et Sp. Orch. 295 = H. triflora.
Carducciana, Goiran, in Nuov. Giorn. Bot. Ital. xv. (1883) 332.—Ital.
Championi, Lindl. in Hook. Kew Journ. vii. (1855) 37 = H. Galeandra.
chlorantha, Cust. ex Reichb. in Moessl. Handb. ed. II. ii. 1565 = H. bifolia.
Chorisiana, Reichb. f. Ic. Fl. Germ. xiii–xiv. 128 = H. Chorisiana.
ciliaris, Lindl. Gen. et Sp. Orch. 292 = H. ciliaris.
clavigera, Lindl. l. c. 289 = H. densa.
commelinifolia, Lindl. in Wall. Cat. n. 7037 = H. commelinifolia.
constricta, Lindl. l. c. n. 7043.—Ind. or.
convallariaefolia, Lindl. Gen. et Sp. Orch. 287.—Kamtschat.
cordifolia, Lindl. in Wall. Cat. n. 7049 = Hemipilia cordifolia.
crispa, Lindl. Gen. et Sp. Orch. 294.—Am. bor.
cristata, Lindl. l. c. 291 = H. cristata.
cubitalis, Lindl. l. c. 292 = H. cubitalis.
decipiens, Lindl. l. c. 290.—Kamtschat.; Sibir.
densa, Lindl. in Wall. Cat. n. 7046 = H. densa.
dentata, Lindl. Gen. et Sp. Orch. 296.—China.
dilatata, Lindl. ex Beck, Bot. N. & M. St. 347 = H. dilatata.
diphylla, Reichb. f. Ic. Fl. Germ. xiii–xiv. 128 = H. cordata.
dolichorrhiza, Reichb. f. l. c. 127 = H. hyperborea.
elegans, Lindl. Gen. et Sp. Orch. 285 = H. elegans.
Erdingeri ×, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 229.—Europ.
fimbriata, Lindl. Gen. et Sp. Orch. 293 = H. fissa?
fissa, Lindl. l. c. 294 = H. peramoena.
flava, Lindl. l. c. 293 = H. herbiola.
florentia, Franch. & Sav. Enum. Pl. Jap. ii. 514.—Japon.
foetida, Geyer, ex Hook. Kew Journ. vii. (1855) 376 = Herminium congestum.
foliosa, Brongn. in Duperr. Voy. Coq. Bot. 194. t. 38 = H. maxillaris.
Galeandra, Reichb. f. in Linnaea, xxv. (1852) 226 = H. Galeandra.
geniculata, Lindl. in Wall Cat. n. 7042 = H. geniculata.
Ghiesbreghtiana, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 30 = H. Ghiesbreghtiana.
gigantea, Lindl. in Wall. Cat. n. 7052 = H. Susannae.
gracilis, Lindl. Gen. et Sp. Orch. 288 = H. gracilis.
graminea, Lindl. l. c. 289 = *Lindleyi*.
graminea, Lindl. l. c. 292 = H. graminea, *Spreng.*
grandiflora, Lindl. l. c. 294 = H. grandiflora, *Torr.*
gratissima, La Pilaye, in Mém. Soc. Linn. Par. iv. (1826) 423.—Terra Nova.
herbiola, Lindl. Gen. et Sp. Orch. 287 = H. herbiola.
Heyneana, Lindl. in Wall. Cat. n. 7044 = H. Heyneana.
hologlottis, Maxim. Prim. Fl. Amur. 268.—Dahuria.

PLATANTHERA:—
holopetala, Lindl. Gen. et Sp. Orch. 291 = Habenaria Blephariglottis.
Hookeri, Lindl. l. c. 286 = H. Hookeriana.
huronensis, Lindl. l. c. 288 = H. hyperborea.
hybrida ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 118.—Europ.
hyperborea, Lindl. Gen. et Sp. Orch. 287 = H. hyperborea.
iantha, Wight, Ic. t. 1692.—Ind. or.
incisa, Lindl. Gen. et Sp. Orch. 293 = H. incisa.
japonica, Lindl. l. c. 290 = H. japonica.
Keiskei, Franch. & Sav. Enum. Pl. Jap. ii. 31 = H. Keiskei.
Koenigii, Lindl. Gen. et Sp. Orch. 286 = H. hyperborea.
lacera, G. Don, in Sweet, Hort. Brit. ed. III. 650 = H. lacera.
latilabris, Lindl. Gen. et Sp. Orch. 289.—Ind. or.
leucophaea, Lindl. l. c. 294 = H. leucophaea.
leucostachys, Lindl. l. c. 288 = H. leucostachys.
limosa, Lindl. in Ann. Nat. Hist. Ser. I. iv. (1840) 381 = H. limosa.
Lindleyi, Steud. Nom. ed. II. ii. 351.—Am. bor.
linifolia, Lindl. in Wall. Cat. n. 7041 = H. graminea, *Lindl.*
longebracteata, Lindl. l. c. n. 7048.—Ind. or.
lucida, Lindl. l. c. 7047 = H. lucida.
lutea, Wight, Ic. t. 919 = H. Perrottetiana.
Mandarinorum, Reichb. f. in Linnaea, xxv. (1852) 226.—China.
marginata, Lindl. in Wall. Cat. n. 7038 = H. marginata.
Menziesii, Lindl. Gen. et Sp. Orch. 286.—Am. bor.
Michauxii, Wood, Class Book (1861), 685 = H. Michauxii.
minor, Reichb. f. in Bot. Zeitschr. xxxvi. (1878) 75.—Japon.
montana, Reichb. f. Ic. Fl. Germ. xiii–xiv. 123 = H. nigra.
neuropetala, Franch. & Sav. Enum. Pl. Jap. ii. 33 = H. neuropetala.
obcordata, Lindl. in Wall. Cat. n. 7050 = H. Galeandra.
obtusata, Lindl. Gen. et Sp. Orch. 284 = H. obtusata.
obtusata, Schur, in Verh. Siebenb. Ver. Naturw. xix. (1868) 206 = H. bifolia.
oligantha, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 86.—Dahuria.
ophrydioides, F. Schmidt, Reisen Amurl. 182.—Ins. Sachalin.
orbiculata, Lindl. Gen. et Sp. Orch. 286 = H. orbiculata.
Orchidis, Lindl. in Wall. Cat. n. 7039.—Ind. or.
oreades, Franch. & Sav. Enum. Pl. Jap. ii. 514 = H. Keiskei.
peramoena, A. Gray, Man. Bot. N. U. St. ed. I. 473 = H. peramoena.
pervia, Peterm. Pflanzenschl. 591 = H. bifolia.
psychodes, Lindl. Gen. et Sp. Orch. 294 = H. lacera.
psycodes, A. Gray, Man. Bot. U. St. ed. I. 472 = H. fissa?
quercicola, Wood, Class-Book (1861), 685.—Am. bor.
radiata, Lindl. Gen. et Sp. Orch. 296 = H. radiata.
Reinii, Franch. & Sav. Enum. Pl. Jap. ii. 513.—Japon.
repens, Wood, Class-Book (1861), 685 = H. repens.
rhynchocarpa, Thw. Enum. Pl. Zeyl. 310 = H. rhynchocarpa.
robusta, Lindl. in Wall. Cat. n. 7036 = H. Susannae.
rostrata, Lindl. l. c. n. 7051 = H. rostrata.
rotundifolia, Lindl. Gen. et Sp. Orch. 292 = H. rotundifolia.
Rumphii, Brongn. in Duperr. Voy. Coq. Bot. 194. t. 38 a = H. Rumphii.
sachalinensis, F. Schmidt, Reisen Amurl. 181.—Ins. Sachalin.
satyrioides, Reichb. f. Ic. Fl. Germ. xiii–xiv. 131.—Europ.
Schischmareffiana, Lindl. Gen. et Sp. Orch. 286 = Herminium congestum.
Schuriana, Fuss, in Verh. Siebenb. Ver. Naturw. xix. (1868) 206 = H. bifolia.
solstitialis, Boenn. ex Reichb. Fl. Germ. Excurs. 120 = H. bifolia.
stenostachya, Lindl. in Hook. Kew Journ. vii. (1855) 37 = H. stenostachya.
stricta, Lindl. Gen. et Sp. Orch. 288.—Am. bor.

PLATANTHERA:—
Susannae, Lindl. Gen. et Sp. Orch. 295 = Habenaria Susannae.
tipuloides, Lindl. l. c. 285 = Tipularia camtschatica.
tricruris, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 265.—Abyss.
uniflora, Lindl. Gen. et Sp. Orch. 295 = H. triflora?
virescens, C. Koch, in Linnaea, xxii. (1849) 288 = H. bifolia.
viridis, Lindl. Syn. Brit. Fl. ed. I. 261 = H. viridis.
vulcanica, Lindl. in Ann. Nat. Hist. Ser. I. iv. (1840) 381 = H. vulcanica.
Wankelii, Reichb. f. in Reichb. Fl. Saxon. 89 = H. bifolia.

PLATANUS, [Tourn.] Linn. Hort. Cliff. 447 (1737). *PLATANACEAE*, Benth. & Hook. f. iii. 396.
acerifolia, Willd. Sp. Pl. iv. 474 = orientalis.
algeriensis, Hort. ex C. Koch, Dendrol. ii. 1. 468 = orientalis.
californica, Benth. Bot. Voy. Sulph. 54 = racemosa.
cuneata, Willd. Sp. Pl. iv. 473 = orientalis.
digitata, Hort. ex Steud. Nom. ed. II. ii. 352 = orientalis.
elongata, Steud. l. c. ed. I. 632 = orientalis.
excelsa, Salisb. Prod. 393 = occidentalis.
hispanica, Tenore, Cat. Ort. Nap. (1845) 91 = orientalis.
hybridus, Brot. Fl. Lusit. ii. 487 = occidentalis.
integrifolia, Hort. ex C. Koch, Dendrol. ii. 1. 469 = occidentalis.
laciniata, Hort. ex C. Koch, l. c. 468 = orientalis.
Lindeniana, *Mart. & Gal. in Bull. Brux.* x. (1843) II. 342.—Mexic.
lobata, Moench, Meth. 358 = occidentalis.
macrophylla, Cree, ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 659 = orientalis.
mexicana, Moric. Pl. Nouv. Am. 39. t. 26 = racemosa.
mexicana, Torr. in Emory, Notes Mil. Reconnois. 151 = Wrightii.
nana, Hort. ex C. Koch, Dendrol. ii. 1. 470 = orientalis.
nepalensis, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 612 = orientalis.
occidentalis, Hook. & Arn. Bot. Beech. Voy. 160 = racemosa.
occidentalis, *Linn. Sp. Pl.* 999.—Am. bor.
orientalis, *Linn. l. c.*—Europ. austr.; Oriens.
palmata, Moench, Meth. 358 = orientalis.
racemosa, *Nutt. ex Audubon, Birds Am.* iv. t. 362; *et N. Am. Sylva, Suppl.* i. 47. t. 15.—Am. bor. occ.
Reuteri, Hort. ex C. Koch, Dendrol. ii. 1. 468 = orientalis.
umbraculifera, Hort. ex C. Koch, l. c. = orientalis.
umbrosa, Salisb. Prod. 393 = orientalis.
undulata, Steud. Nom. ed. I. 632 = orientalis.
vulgaris, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 291 = orientalis.
Wrightii, *S. Wats. in Proc. Am. Acad.* x. (1875) 349. —Arizona; Mexic.

PLATEA, Blume, Bijdr. 646 (1825). *OLACINEAE*, Benth. & Hook. f. i. 353.
axillaris, Thw. Enum. Pl. Zeyl. 44 = Gomphandra axillaris.
coriacea, Thw. l. c. = Gomphandra polymorpha.
corniculata, *Becc. Malesia*, i. (1877) 117.—Malaya.
excelsa, *Blume, Bijdr.* 646.—Java.
Griffithiana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 110 = Phlebocalymna Griffithiana.
latifolia, *Blume, Bijdr.* 646.—Java.
laxiflora, Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 111 = Gomphandra laxiflora.
Lobbiana, Miers. l. c. = Phlebocalymna Lobbiana.
papuana, *Becc. Malesia*, i. (1883) 257.—N. Guin.
Riedeliana, *Becc. l. c.* (1877) 116.—Ins. Celebes.
sumatrana, *Blume, Mus. Bot. Lugd. Bat.* i. 249.—Sumatra.
Wightiana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 111 = Gomphandra polymorpha.

PLATEANA, Salisb. Gen. Pl. Fragm. 100 (1866) = **Narcissus**, Tourn. (Amaryll.).

PLATENIA, Karst. in Linnaea, xxviii. (1856) 250 = **Cocos**, Linn. (Palm.).
Chiragua, Karst. l. c. (= *Syagrus Chiragua.*)—N. Granat.

PLATHYMENIA, Benth. in Hook. Journ. Bot. iv. (1842) 333. *LEGUMINOSAE*, Benth. & Hook. f. i. 589.
CHRYSOXYLON, Casar. Nov. Stirp. Bras. Dec. 59 (1843).
foliolosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 334.—Bras.
reticulata, *Benth. l. c.*—Bras.

PLATOLARIA, Rafin. Sylva Tellur. 78 (1838) = **Tecoma**, Juss. (Bignon.).
flavescens, Rafin. l. c. = T. flavescens.

PLATONIA, Kunth, Rév. Gram. i. 139, 327. t. 76 (1830) = **Planotia**, Munro (Gramin.).
elata, Kunth, l. c. = Planotia elata.
virgata, Griseb. Fl. Brit. W. Ind. 530 = Planotia virgata.

PLATONIA, Mart. Nov. Gen. et Sp. iii. 168. t. 289 (1829). *GUTTIFERAE*, Benth. & Hook. f. i. 174.
grandiflora, *Planch & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 299.—Bras.
insignis, *Mart. Nov. Gen. et Sp.* iii. 169.—Bras.

PLATONIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Lippia**, Linn. (Verben.).
nudiflora, Rafin. l. c. = L. nodiflora.

PLATOSTOMA, Beauv. Fl. Owar. ii. 61. t. 95 (1805). *LABIATAE* [Benth. & Hook. f. ii. 1172].
PLATYSTOMA, Benth. & Hook. f. l. c., sphalm. (1876).
africanum, *Beauv. Fl. Owar.* ii. 61. t. 95.—Afr. trop.
flaccidum, *Benth. & Hook. f. Gen.* ii. 1173.—Ind. or.; Afr. trop.

PLATUNUM, A. Juss. in Ann. Mus. Par. vii. (1806) 76 = **Holmskioldia**, Retz. (Verben.).
rubrum, A. Juss. l. c. = H. sanguinea.

PLATYCAPNOS, Bernh. in Linnaea, viii. (1833) 471 = **Fumaria**, Linn. (Papav.).
saxicola, Willk. in Bot. Zeit. vi. (1848) 367 = F. saxicola.
spicatus, Bernh. in Linnaea, viii. (1833) 471 = F. spicata.
tenuilobus, Pomel, Nouv. Mat. Fl. Atl. 240.—Alger.

PLATYCARPHA, Less. in Linnaea, vi. (1831) 688. *COMPOSITAE*, Benth. & Hook. f. ii. 462.
carlinoides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 430.—Afr. trop.
glomerata, *Less. in Linnaea*, vi. (1831) 688; *DC. Prod.* v. 71.—Afr. austr.

PLATYCARPIDIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 309 = **Platysace**, Bunge (Umbell.).
validum, F. Muell. l. c. 310 = Platysace valida.

PLATYCARPUM, Humb. & Bonpl. Pl. Aequin. ii. 81. t. 104 (1809). *RUBIACEAE*, Benth. & Hook. f. ii. 44.
orinocense, *Humb. & Bonpl. l. c.*—Venezuela.

PLATYCARYA, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 741. t. 5. *JUGLANDEAE*, Benth. & Hook. f. iii. 400.
FORTUNAEA, Lindl. in Journ. Hort. Soc. i. (1846) 150.
FORTUNEA, Poit. in Rev. Hortic. Sér. II. v. (1846) 282.
strobilacea, *Sieb. & Zucc. in Abh. Akad. Muench.* iii. II. (1843) 743. t. 5.—Japon.

PLATYCENTRUM, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 123. t. 11 B = **Begonia**, Linn.
annulatum, C. Koch, in Berl. Gartenz. (1857); ex A. DC. in DC. Prod. xv. I. 350 = B. Griffithii.
Cathcartii, Klotzsch, in Abh. Akad. Berl. (1855) 125 = B. Cathcartii.
discolor, Miq. Fl. Ind. Bat. i. I. 694 = B. Evansiana.
erosum, Miq. l. c. = B. erosa.
Hamiltonianum, Miq. l. c. 695 = B. acuminata.
multangulum, Miq. l. c. = B. multangula.
Rex, Seem. in Harting. Parad. t. 81, 82 = B. Rex.
robustum, Miq. Fl. Ind. Bat. i. I. 694 = B. robusta.
rubro-venium, Klotzsch, in Abh. Akad. Berl. (1855) 124 = B. rubro-venia.

PLATYCENTRUM :—
rupicolum, Miq. Fl. Ind. Bat. i. I. 693 = Begonia rupicola.
tenuifolium, Miq. l. c. = B. tenuifolia.
Teysmannianum, Miq. l. c. 1092 = B. Teysmaniana (in Addend.).
xanthinum, Klotzsch, in Abh. Akad. Berl. (1855) 123 = B. xanthina.
Zollingerianum, Klotzsch, l. c. 125 = B. Zollingeriana.

PLATYCENTRUM, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 114. t. 5. *MELASTOMACEAE*, Benth. & Hook. f. i. 760.
clidemioides, *Naud. l. c.*—Guian.

PLATYCHAETA, Boiss. Diagn. Sér. I. xi. 5 (1849) = **Pulicaria**, Gaertn. (Compos.).
Aucheri, Boiss. l. c. = Pul. Aucheri.
carnosa, Boiss. l. c. Ser. II. iii. 10.—Afghan.
glaucescens, Boiss. l. c. Ser. I. xi. 5 = Pul. glaucescens.
glutinosa, Boiss. l. c. = Pul. glutinosa.
mucronifolia, Boiss. l. c.—Persia.
velutina, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 208.—Persia.
villosa, Boiss. Diagn. Ser. II. iii. 11 = Beluchist.; Sind.

PLATYCHEILUS, Cass. in Dict. Sc. Nat. xxxiv. 206, 212 (1825) = **Trixis**, P. Br. (Compos.).
ochroleucus, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 33, in syn. = T. ochroleuca.

PLATYCHILUM, Delaun. Herb. Amat. t. 187 (1815) = **Hovea**, R. Br. (Legumin.).
Celsianum, Delaun. l. c. = H. Celsi.

PLATYCLADUS, Spach, Hist. Vég. Phan. xi. 333 (1842) = **Thuya**, Linn. (Conif.).
dolabrata, Spach, l. c. 337 = T. dolabrata.
stricta, Spach, l. c. 335 = T. orientalis.

PLATYCLINIS, Benth. in Journ. Linn. Soc. xviii. (1881) 295. *ORCHIDEAE*, Benth. & Hook. f. iii. 496.
abbreviata, *Benth. ex Hemsl. in Gard. Chron.* (1881) II. 656.—Java.
brachyota, *Hemsl. l. c.*—Java.
Cobbiana, *Hemsl. l. c.*—Ins. Philipp.
commelynifolia, Benth. in Journ. Linn. Soc. xviii. (1881) 295 = Microstylis commelynifolia.
cornuta, *Benth. ex Hemsl. in Gard. Chron.* (1881) II. 656.—Java.
edentula, *Benth. ex Hemsl. l. c.*—Java.
filiformis, *Benth. ex Hemsl. l. c.*—Ins. Philipp.
glumacea, *Benth. ex Hemsl. l. c.*—Ins. Philipp.
latifolia, *Hemsl. l. c.*—Ins. Philipp.
longifolia, *Hemsl. l. c.*—Hab.?

PLATYCLINIUM, T. Moore, in Moore & Ayres, Gard. Mag. Bot. ii. (1850) 153 = **Begonia**, Linn.
cinnabarina, T. Moore, l. c. = B. cinnabarina.

PLATYCODON, A. DC. Monog. Camp. 125 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 556.
autumnale, Decne. in Rev. Hortic. Sér. III. ii. (1848) 361 = grandiflorum.
chinense, Lindl. & Paxt. Flow. Gard. ii. (1851-52) 121 = grandiflorum.
grandiflorum, *A. DC. Monog. Camp.* 125.—As. bor.; China; Japon.
homallanthium, A. DC. l. c. 126 = Wahlenbergia homallanthina.
sinensis, Lem. Jard. Fl. sub t. 250 = grandiflorum.

PLATYCODON, Reichb. Nom. 236 (1841) = **Daucus**, Tourn. (Umbell.).

PLATYCORYNE, Reichb. f. in Bonplandia, iii. (1855) 212. *ORCHIDEAE*, Benth. & Hook. f. iii. 632.
Pervillei, *Reichb. f. l. c.*—Madag.

PLATYCRATER, Sieb. & Zucc. Fl. Jap. 62. t. 27 (1835). *SAXIFRAGEAE*, Benth. & Hook. f. i. 642.
arguta, *Sieb. & Zucc. l. c.* 64.—Japon.

PLATYCYAMUS, Benth. in Mart. Fl. Bras. xv. I. 323 (1862). *LEGUMINOSAE*, Benth. & Hook. f. i. 531.
Regnellii, *Benth. in Mart. Fl. Bras.* xv. I. 323.—Bras.

PLATYDAUCON, Reichb. Nom. 205 (1841) = **Daucus**, Tourn. (Umbell.).

PLATYDESMA, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 317. *RUTACEAE*, Benth. & Hook. f. i. 991.
campanulata, *H. Mann, l. c.*—Ins. Sandvic.

PLATYESTES, Salisb. Gen. Pl. Fragm. 21 (1866) = **Lachenalia**, Jacq. (Liliac.).
nervosa, Salisb. l. c. = L. nervosa.
purpureo-coerulea, Salisb. l. c. = L. purpureo-coerulea.
racemosa, Salisb. l. c. = L. racemosa.

PLATYGONIA, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 33. t. 5 = **Trichosanthes**, Linn. (Cucurb.).
himalensis, Naud. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 608 = T. himalensis.
Kaempferi, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 33. t. 5 = T. cucumeroides.

PLATYGYNA, Mercier, in Ser. Bull. Bot. (1830) 168. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 328.
ACANTHOCAULON, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1850).
hexandra, Muell. Arg. in DC. Prod. xv. II. 914 = urens.
pruriens, Baill. Hist. Pl. v. 215 = urens.
urens, *Mercier, in Ser. Bull. Bot.* i. (1830) 168.—Cuba.

PLATYHYMENIA, Walp. Rep. i. 858 (1842), sphalm. = **Plathymenia**, Benth. (Legumin.).

PLATYLEPIS, Kunth, Enum. Pl. ii. 269 (1837) = **Ascolepis**, Nees (Cyperac.).
brasiliensis, Kunth, l. c.—Bras. et Afr. trop.
capensis, Kunth, l. c. = A. capensis.
dioica, Steud. Syn. Pl. Cyp. 131.—Afr. austr.
guianensis, Nees, in Mart. Fl. Bras. ii. I. 63.—Am. trop.
leococephala, Nees, l. c.—Bras.
xanthocephala, Nees, l. c. 62.—Bras.

PLATYLEPIS, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 34; Orch. Iles Fr. et Bourb. 39. t. 6. *ORCHIDEAE*, Benth. & Hook. f. iii. 604.
DIPLOGASTRA, Reichb. f. in Flora, xlviii. (1865) 183.
NOTIOPHRYS, Lindl. in Journ. Linn. Soc. i. (1857) 189; vi. (1862) 138.
Commelynae, *Reichb. f. in Linnaea*, xli. (1877) 63.—Ins. Tahiti.
glandulosa, *Reichb. f. l. c.* 62.—Afr. austr.
goodyeroides, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 34.—Ins. Maurit.
heteromorpha, *Reichb. f. Otia Bot. Hamb.* 52.—Ins. Samoa.
occulta, Reichb. f. in Linnaea, xli. (1877) 62 = goodyeroides.
polyadenia, *Reichb. f. in Flora*, lxviii. (1885) 537.—Ins. Comoro.
sechellarum, *S. Moore, in Baker, Fl. Maurit.* 340.—Ins. Seychelles.

PLATYLOBIUM, Sm. Spec. Bot. N. Holl. 17 (1793); et in Trans. Linn. Soc. ii. (1794) 350. *LEGUMINOSAE*, Benth. & Hook. f. i. 473.
CHEILOCOCCA, Salisb. ex Sm. Spec. Bot. N. Holl. 17 (1793), cum cit. falsa.
alternifolium, *F. Muell. in Wing, South. Sc. Record*, iii. (1883) 99.—Austral.
aphyllum, *Hort. ex Steud. Nom.* ed. II. ii. 352, nomen.—Austral.
formosum, *Sm. Spec. Bot. N. Holl.* 17. t. 6; *et in Trans. Linn. Soc.* ii. (1794) 350.—Austral.
gracile, *Hort. ex Steud. Nom.* ed. II. ii. 352, nomen.—Austral.
lanceolatum, Andr. Bot. Rep. t. 205 = Bossiaea heterophylla.
macrocalyx, Meissn. in Lehm. Pl. Preiss. i. 80 = obtusangulum.
microphyllum, Sims, Bot. Mag. t. 863 = Bossiaea microphylla.
Murrayanum, Hook. l. c. t. 3259 = triangulare.
obcordatum, DC. Prod. ii. 116 = seq.?
obcordatum, Vent. Jard. Malm. sub t. 31 = Bossiaea microphylla.
obtusangulum, *Hook. Bot. Mag.* t. 3258.—Austral.

PLATYLOBIUM:—
ovatum, Andr. Bot. Rep. t. 266 = Bossiaea heterophylla.
ovatum, Sieber, ex DC. Prod. ii. 116 = formosum.
parviflorum, Sm. Spec. Bot. N. Holl. 18 = formosum.
reticulatum, Sieber, ex Benth. in Ann. Wien. Mus. ii. (1838) 84 = Mirbelia grandiflora.
scolopendrium, Andr. Bot. Rep. t. 191; Vent. Jard. Malm. t. 55 = Bossiaea scolopendria.
spinosum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 284 = Bossiaea calycina.
triangulare, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 266. —Austral.
triangulare, Sims, Bot. Mag. t. 1508 = obtusangulum.

PLATYLOPHUS, Cass. in Dict. Sc. Nat. xliv. 36 (1826) = **Centaurea**, Linn. (Compos.).
nemoralis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 96 = C. nigra.
obscurus, Fourr. l. c. = C. nigra.

PLATYLOPHUS, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 92. *SAXIFRAGEAE*, Benth. & Hook. f. i. 652.
TRIMERISMA, Presl, Bot. Bemerk. 73 (1844).
trifoliatus, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 92.—Afr. austr.

PLATYMERIUM, Bartl. ex DC. Prod. iv. 619 (1834). Gen. dub. *RUBIACEARUM*.—Cf. Benth. & Hook. f. Gen. ii. 29.
glomeratum, *Bartl. ex DC. l. c.*—Ins. Philipp.

PLATYMETRA, Noronha, ex Salisb. Gen. Pl. Fragm. 9 (1866) = **Tupistra**, Ker-Gawl. (Liliac.).

PLATYMISCIUM, Vog. in Linnaea, xi. (1837) 198. *LEGUMINOSAE*, Benth. & Hook. f. i. 548.
HYMENOLOBIUM, Benth. in Flora, xx. (1837) II. Beibl. 122.
Blancheti, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 83; *et in Mart. Fl. Bras.* xv. i. 272.—Bras.
dichotomum, Benth. in Ann. Wien. Mus. ii. (1838) 104 = floribundum.
filipes, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 83.—Guian.
floribundum, *Vog. in Linnaea*, xi. (1837) 199.—Bras.
gracile, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 82.—Peruv.
hebestachyum, *Benth. l. c.* 81.—N. Granat.
latifolium, Benth. in Ann. Wien. Mus. ii. (1838) 104 = floribundum.
luteum, Benth. l. c. = nitens.
nitens, *Vog. in Linnaea*, xi. (1837) 417.—Bras.
parviflorum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 81.—Am. centr.
platystachyum, Griseb. Fl. Brit. W. Ind. 200, sphalm. = seq.
polystachyum, *Benth. in Seem. Bot. Voy. Herald*, 111. t. 21.—Am. trop.
praecox, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 104.—Bras.
pubescens, *Micheli, in Kjoeb. Vidensk. Meddel.* (1875) 101.—Bras.
speciosum, *Vog. in Linnaea*, xi. (1837) 200.—Bras.
stipulare, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 82.—Peruv.
trifoliolatum, *Benth. l. c.*—Mexic.
Trinitatis, *Benth. l. c.*—Ins. Trinitat.

PLATYNEMA, Schrad. Ind. Sem. Hort. Gotting. (1835). —Cf. Linnaea, xi. (1837) Litt. 89 = **Mertensia**, Roth (Borag.).
paniculata, Schrad. l. c. = M. paniculata.

PLATYNEMA, Wight & Arn. in Edinb. N. Phil. Journ. (July, 1833) 179 = **Tristellateia**, Thou. (Malpig.).
laurifolium, Wight & Arn. l. c. = T. australasica.
parvifolium, D. Dietr. Syn. Pl. ii. 1590 = Hiptage parvifolia.

PLATYPETALUM, R. Br. in Parry, 1st Voy. Suppl. 266 (1824) = **Braya**, Sternb. & Hoppe (Crucif.).
capitatum, Turcz. ex Ledeb. Fl. Ross. i. 195 = B. rosea.
dubium, R. Br. in Parry, 1st Voy. Suppl. 266 = B. rosea.

PLATYPETALUM:—
involucratum, Bunge, Verz. Suppl. Fl. Alt. 77 = Braya involucrata.
nanum, Turcz. ex Bunge, Ind. Sem. Hort. Dorpat. (1841) 8 = B. nana.
purpurascens, R. Br. in Parry, 1st Voy. Suppl. 267 = B. alpina.

PLATYPODIUM, Vog. in Linnaea, xi. (1837) 420. *LEGUMINOSAE*, Benth. & Hook. f. i. 546.
CALLISEMAEA, Benth. in Ann. Wien. Mus. ii. (1838) 105.
elegans, *Vog. in Linnaea*, xi. (1837) 420.—Bras.
grandiflorum, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 73.—Bras.
viride, Vog. in Linnaea, xi. (1837) 422 = elegans.

PLATYPTELEA, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 55 = **Aphanopetalum**, Endl. (Saxifrag.).
clematidea, J. Drumm. ex Harv. l. c. = A. occidentale.

PLATYPTERIS, H. B. & K. Nov. Gen. et Sp. iv. (1820) 200 = **Verbesina**, Linn. (Compos.).
crocata, H. B. & K. l. c. 201 = V. crocata.

PLATYRAPHIUM, Cass. in Dict. Sc. Nat. xxxv. 173 (1825) = **Cnicus**, Linn. (Compos.).
Billardieri, Cass. l. c. xli. 307 = C. Diacantha.
Jacquini, Cass. l. c. 306 = C. Diacantha.

PLATYRHAPHE, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 56 (1867) = **Pimpinella**, Riv. (Umbell.).
japonica, Miq. l. c. = Pimp. diversifolia.

PLATYRHIZA, Rodrig. Orch. Nov. ii. 230 (1881). *ORCHIDEAE*.
quadricolor, *Rodrig. l. c.* 231.—Bras.

PLATYSACE, Bunge, in Lehm. Pl. Preiss. i. 285 (1845). *UMBELLIFERAE* [Benth. & Hook. f. i. 874].
CATEPHA, Lechen. ex Reichb. Handb. 221 (1837).
FISCHERA, Spreng. Pl. Umb. Prod. 27 (1813).
PLATYCARPIDIUM, F. Muell. in Hook. Kew Journ. ix. (1857) 309.
SIEBERA, Reichb. Consp. 145 (1828).
TRACHYMENE, DC. Prod. iv. 72 (1830).
cirrosa, *Bunge, in Lehm. Pl. Preiss.* i. 285.—Austral.
flexuosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 29.—Austral.
trachymenoides, *F. Muell. Fragm.* i. 232.—Austral.
valida, *F. Muell. l. c.*—Austral.

PLATYSEMA, Benth. in Ann. Wien. Mus. ii. (1838) 122 = **Centrosema**, DC. (Legumin.).
triquetrum, Benth. l. c. = C. triquetrum (in Addend.).

PLATYSEPALUM, Welw. ex Baker, in Oliver, Fl. Trop. Afr. ii. 131 (1871). *LEGUMINOSAE*.
violaceum, *Welw. ex Baker, l. c.*—Afr. trop.

PLATYSMA, Blume, Bijdr. 295. t. 43 (1825) = **Podochilus**, Blume (Orchid.).
gracilis, Blume, l. c. = Pod. gracilis.

PLATYSPERMUM, Hoffm. Gen. Umbell. 64 (1814) = **Daucus**, Tourn. (Umbell.).
alatum, Schult. Syst. vi. 837 (Index) = D. alatus.
aureum, Pomel, Nouv. Mat. Fl. Atl. 148 = D. aureus.
grandiflorum, Pomel, l. c. = D. grandiflorus, *Desf.*
littorale, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 78 = D. bessarabicus.
muricatum, Hoffm. Gen. Umbell. 64 = D. muricatus.
orientale, Eichw. Pl. Casp. Cauc. 5 = D. pulcherrimus.
pubescens, Pomel, Nouv. Mat. Fl. Atl. 148 = D. littoralis.
pulcherrimum, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 78 = D. pulcherrimus.

PLATYSPERMUM, Hook. Fl. Bor. Am. i. 68. t. 18 B (1829). *CRUCIFERAE*, Benth. & Hook. f. i. 72.
scapigerum, *Hook. l. c.*—Am. bor.

PLATYSTACHYS, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 11; Beer, Bromel. 80 (1857) = **Tillandsia**, Linn. (Bromel.).
anceps, Beer, Bromel. 80 = T. anceps.
azurea, Beer, l. c. 90 = T. azurea.
bulbosa, Beer, l. c. 83 = T. bulbosa.
cinerascens, Beer, l. c. 263 = T. disticha.
complanata, E. Morr. in Belg. Hortic. (1872) 300 = T. complanata.
cyanea, C. Koch & Sello, Ind. Sem. Hort. Berol. (1854) App. 11 = T. cyanea.
digitata, Beer, Bromel. 84 = T. polystachya.
disticha, Beer, l. c. 264 = T. helicnioides.
Ehrenbergiana, C. Koch, ex Hemsl. Biol. Centr. Am. Bot. iii. 320 = T. Ehrenbergiana.
Ehrenbergii, Beer, Bromel. 264 = T. Ehrenbergii.
Ehrenbergii, C. Koch, Ind. Sem. Hort. Berol. (1873) App. iv. 5 = T. utriculata.
erythraea, Beer, Bromel. 83 = T. bulbosa.
filifolia, Beer, l. c. 264 = T. filifolia.
floribunda, Beer, l. c. = T. floribunda.
geniculata, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 345.—Bras.
glaucophylla, Beer, Bromel. 82 (= *Vriesia glaucophylla*).—N. Granat.
gracilis, Beer, l. c. 254 = Aechmea tillandsioides.
havanensis, Beer, l. c. 85 = T. fasciculata.
heptantha, Beer, l. c. 80 = T. heptantha.
inanis, Beer, l. c. 82 = T. bulbosa.
incarnata, Beer, l. c. 264 = T. incarnata.
juncea, Beer, l. c. 86 = T. tenuifolia.
Kunthiana, Beer, l. c. 90 = T. Kunthiana.
latifolia, C. Koch, Ind. Sem. Hort. Berol. (1873) App. iv. 5 = T. latifolia.
Moritziana, Beer, Bromel. 265 = T. Moritziana.
parviflora, Beer, l. c. 88 = T. parviflora.
patens, C. Koch, Ind. Sem. Hort. Berol. (1873) App. iv. 5 = T. flexuosa.
Plumierii, Beer, Bromel, 91 = T. serrata.
polystachia, Beer, l. c. 87 = T. polystachya.
pruinosa, Beer, l. c. 265 = T. pruinosa.
purpurea, Beer, l. c. 89 = T. purpurea.
scoparia, Beer, l. c. 265 = T. straminea.
setacea, Beer, l. c. 80 = T. compressa.
tortilis, Beer, l. c. 266 = T. pruinosa.
utriculata, Beer, l. c. = T. utriculata.
violacea, Beer, l. c. 264.—Mexic.
viridiflora, Beer, l. c. 81.—Mexic.

PLATYSTEMMA, Wall. Pl. As. Rar. ii. 41. t. 151 (1831). *GESNERACEAE*, Benth. & Hook. f. ii. 1020.
majus, Wall. l. c. 42 = violoides.
violoides, *Wall. l. c.* 42. t. 151.—Reg. Himal.

PLATYSTEMON, Benth. in Trans. Hort. Soc. Ser. II. i. (1834) 405. *PAPAVERACEAE*, Benth. & Hook. f. i. 51.
BOOTHIA, Dougl. ex Benth. in Trans. Hort. Soc. Ser. II. i. (1834) 405.
californicus, *Benth. l. c.*—Am. bor. occ.
leiocarpus, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 47 = californicus.

PLATYSTEPHIUM, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 80 = **Egletes**, Cass. (Compos.).
graveolens, Gardn. l. c. 81 = E. viscosa.

PLATYSTIGMA, Benth. in Trans. Hort. Soc. Ser. II. i. (1834) 406. *PAPAVERACEAE*, Benth. & Hook. f. i. 51.
MECONELLA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 64 (1838).
californicum, *Benth. & Hook. f. Gen.* i. 51.—Am. bor. occ.
lineare, *Benth. in Trans. Hort. Soc.* Ser. II. i. (1834) 407.—Am. bor. occ.
oreganum, *Benth. & Hook. f. ex S. Wats. Bibliog. Ind. N. Am. Bot.* 43.—Am. bor. occ.

PLATYSTIGMA, R. Br. in Wall. Cat. n. 7523 (1832). *EUPHORBIACEAE* [Benth. & Hook. f. iii. 283].
myristiceum, *R. Br. l. c.*—Ind. or.

PLATYSTOMA, Benth. & Hook. f. Gen. iii. 1172 (1876) sphalm. = **Platostoma**, Beauv. (Labiat.).

PLATYSTYLIS, Lindl. Gen. et Sp. Orch. 18 (1830) = **Liparis**, Rich. (Orchid.).
atropurpurea, Lindl. l. c.—Java.
decurrens, Lindl. l. c.—Java.

PLATYSTYLIS, Sweet, Brit. Flow. Gard. Ser. I. t. 239 (1828) = **Lathyrus**, Tourn. (Legumin.).
cyanea, Sweet, l. c. = L. cyaneus.
sessilifolia, Sweet, l. c. sub t. 239 = L. sessilifolius.
stipulacea, G. Don, Gen. Syst. ii. 341 = L. venosus.

PLATYTHECA, Steetz, in Lehm. Pl. Preiss. i. 220 (1845). *TREMANDREAE*, Benth. & Hook. f. i. 134.
crassifolia, Steetz, l. c. 222 = galioides.
Crucianella, Steetz, l. c. 221 = galioides.
galioides, *Steetz, l. c.* 220.—Austral.
verticillata, Baill. Hist. Pl. v. 67 = galioides.

PLATYZAMIA, Zucc. in Abh. Akad. Muench. iv. (1845) 23. t. 4 = **Dioon**, Lindl. (Cycad.).
rigida, Zucc. l. c. = D. edule.

PLATZCHAETA, Sch. Bip. in F. Schultz, Arch. Fl. 131 (1855) = PLATYCHAETA, Boiss. = **Pulicaria**, Gaertn. (Compos.).
uniflora, Sch. Bip. l. c. = Pul. glutinosa.

PLAZERIA, Steud. Nom. ed. II. ii. 353 (1841) = seq.

PLAZERIUM, Willd. ex Kunth, Enum. Pl. i. 474 (1833) = **Saccharum**, Linn. (Gramin.).
rufum, Willd. ex Kunth, l. c. = S. cayennense.

PLAZIA, Ruiz & Pav. Prod. 104 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 486.
AGLAODENDRON, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 175.
brasiliensis, Spreng. Neue Entdeck. ii. 137 = Stiftia chrysantha.
cheiranthifolia, *Wedd. Chlor. And.* i. 12.—Chili.
conferta, *Ruiz & Pav. Syst. Veg.* 187.—Peruv.
daphnoides, *Wedd. Chlor. And.* i. 13. t. 2.—Peruv.
parviflora, Spreng. Syst. iv. Cur. Post. 301 = Stiftia parviflora.

PLECOSPERMUM, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 124. t. 4. *URTICACEAE*, Benth. & Hook. f. iii. 361.
CARDIOGYNE, Bur. in DC. Prod. xvii. 232 (1873).
cuneifolium, Thw. in Hook. Kew Journ. vi. (1854) 303 = Cudrania javanensis.
spinosum, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 124.—Ind. or.

PLECOSTIGMA, Turcz. in Trautv. Pl. Imag. 8 t. 2 (1844) = **Gagea**, Salisb. (Liliac.).
pauciflorum, Turcz. l. c. = G. pauciflora.

PLECTANEIA, Thou. Gen. Nov. Madag. 11 (1806). *APOCYNACEAE*, Benth. & Hook. f. ii. 708.
Thouarsii, *Roem. & Schult. Syst.* iv. 420.—Madag.

PLECTANTHERA, Mart. Nov. Gen. et Sp. i. 39. t. 36 (1824) = **Luxemburgia**, A. St. Hil. (Ochnac.).
ciliosa, Mart. l. c. 40 = L. polyandra.
floribunda, Mart. l. c. = L. speciosa.

PLECTOCEPHALUS, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 51 (1830) = **Centaurea**, Linn. (Compos.).
abyssinicus, Boiss. Diagn. Ser. II. iii. 62 = C. abyssinica.
americanus, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 51 = C. americana.
chilensis, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1446 = C. chilensis.
cyanoides, Boiss. Diagn. Ser. II. iii. 62 = C. varians.

PLECTOCOMIA, Mart. & Blume, in Schult. f. Syst. vii. II. 1333 (1830). *PALMAE*, Benth. & Hook. f. iii. 934.
assamica, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 97.—Assam.
elongata, *Mart. & Blume, in Schult. f. Syst.* vii. 1333.—Java.
geminiflora, *H. Wendl. in Kerch. Palm.* 254.—Malaya.

PLECTOCOMIA:—
himalayana, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 100.—Reg. Himal.
Hystrix, *Linden, in Illustr. Hortic.* xxviii. (1881) 32.—Borneo.
icthyospinus, *Hort.—Cf. Gard. Chron.* v. (1876) 735.—Hab. ?
khasyana, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 98.—Reg. Himal.
macrostachya, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 207.—Burma.
montana, Griff. ex T. Anders. in Journ. Linn. Soc. xi. (1871) 12 = himalayana.
Mulleri, *Blume, Rumphia*, iii. 71.—Borneo.
rigida, H. Wendl. in Bot. Zeit. xvii. (1859) 165 = Mulleri.
spectabilis, *Hort. ex H. Wendl. in Kerch. Palm.* 254, nomen.—Hab. ?
sumatrana, Miq. Fl. Ind. Bat. Suppl. 592 = elongata ?

PLECTOGYNE, Link, in Otto & Dietr. Allg. Gartenz. ii. (1834) 265 = **Aspidistra**, Ker-Gawl. (Liliac.).
variegata, Link, l. c. = A. elatior.

PLECTOMA, Rafin. Fl. Tellur. iv. 110 (1836) = **Utricularia**, Linn. (Lentib.).
inflata, Rafin. l. c. = U. inflata.
stellata, Rafin. l. c. = U. stellaris.

PLECTOPOMA, Hanst. in Linnaea, xxvi. (1853) 201 = **Gloxinia**, L'Hérit. (Gesner.).
eucodonioides ×, Van Houtte, Fl. des Serres, xviii. (1869–70). t. 1858.
fimbriatum, Hanst. in Linnaea, xxvi. (1853) 201 = G. glabrata.
glabratum, Hanst. l. c. xxxiv. (1865–66) 434 = G. glabrata.
gloxiniflorum, Hanst. l. c. xxvii. (1854) 714 = G. glabrata.
myriostigma ×, Van Houtte, in Fl. des Serres, xix. (1873) t. 1969–70.
naegelioides ×, Van Houtte, l. c. xvii. (1867–68) 67.

PLECTRANTHERA, Benth. & Hook. f. Gen. i. 319 (1862), sphalm. = PLECTANTHERA, Mart. = **Luxemburgia**, A. St. Hil. (Ochnac.).

PLECTRANTHUS, L'Hérit. Stirp. Nov. 84, verso, t. 41, 42 (1785). *LABIATAE*, Benth. & Hook. f. ii. 1175.
GERMANEA, Lam. Encyc. ii. 690 (1786).
ISODON, Schrad. ex Benth. Lab. Gen. et Sp. 40 (1832).
RABDOSIA, Hassk. in Flora, xxv. (1842) II. Beibl. 25.
amboinensis, Spreng. Syst. ii. 690 = Coleus aromaticus.
amethystoides, *Benth. Lab. Gen. et Sp.* 45.—China.
amoenus, Wall. ex Hook. f. Fl. Brit. Ind. iv. 615 = Orthosiphon rubicundus.
aromaticus, Roxb. Hort. Beng. 45; Fl. Ind. iii. 22 = Coleus aromaticus.
asper, Spreng. Syst. ii. 690 = Coleus barbatus.
australis, R. Br. Prod. 506 = parviflorus.
barbatus, Andr. Bot. Rep. t. 594 = Coleus barbatus.
Barrelieri, Spreng. Syst. ii. 691 = Ocimum Basilicum.
Benthamianus, *Miq. Fl. Ind. Bat.* ii. 946.—Java.
bicolor, Blume, Bijdr. 837 = Coleus bicolor.
Buckia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 117 = ternifolius.
Buergeri, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 101 = glaucocalyx.
bullatus, Benth. in DC. Prod. xii. 56 = nilgherricus.
calycinus, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 230.—Afr. austr.
canescens, *Benth. Lab. Gen. et Sp.* 33.—Madag.
caninus, Roth, Nov. Pl. Sp. 279 = Coleus spicatus.
capillipes, *Benth. in DC. Prod.* xii. 57.—Zeylan.
carnosus, Sm. in Rees, Cycl. xxvii. n. 7 = Anisochilus carnosus.
ciliatus, *E. Mey. Comm. Pl. Afr. Austr.* 227.—Afr. austr.
Coetsa, Benth. in DC. Prod. xii. 58, partim = glaucocalyx.
Coetsa, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 117.—Reg. Himal.; Burma.
Coetsa, Thw. Enum. Pl. Zeyl. 237 = menthoides.
coleoides, *Benth. in DC. Prod.* xii. 64.—Ind. or.

PLECTRANTHUS:—
coloratus, D. Don, Prod. Fl. Nep. 116 = Geniosporum strobiliferum.
coloratus, *E. Mey. Comm. Pl. Afr. Austr.* 238.—Afr. austr.
comosus, Sims, Bot. Mag. t. 2318 = Coleus barbatus.
Concan, err. typ. in Hook. f. Fl. Brit. Ind. v. 768 = Stocksii.
concolor, Benth. in DC. Prod. xii. 75, in syn. = Coleus macrophyllus.
congestus, *R. Br. Prod.* 506.—Austral.
cordifolius, D. Don, Prod. Fl. Nep. 116 = incanus.
crassifolius, Hort. ex Benth. Lab. Gen. et Sp. 51, 60 = Coleus aromaticus, Anisochilus carnosus.
crassifolius, Vahl, Symb. Bot. i. 44 = Coleus spicatus ?
crenulatus, *Hook. f. Fl. Brit. Ind.* iv. 623.—Burma.
cylindraceus, *Hochst. ex Benth. in DC. Prod.* xii. 60.—Abyss.
cymosus, *Baker, in Journ. Linn. Soc.* xxi. (1885) 434.—Madag.
decumbens, *Hook. f. l. c.* vii. (1864) 210.—Afr. trop.
defoliatus, *Hochst. ex Benth. in DC. Prod.* xii. 60.—Abyss.
divaricatus, Weinm. in Syll. Ratisb. i. (1824) 68 = incanus.
dubius, Spreng. Syst. ii. 691 = Anisochilus carnosus.
dubius, *Vahl, ex Benth. Lab. Gen. et Sp.* 711.—Japon.
Eckloni, *Benth. in DC. Prod.* xii. 64.—Afr. austr.
excisus, *Maxim. Prim. Fl. Amur.* 213.—Manchur.
foetidus, *Benth. Lab. Gen. et Sp.* 35.—Austral.
Forskâlaei, Vahl, Symb. Bot. i. 44 = madagascariensis.
Forskâlaei, Willd. Sp. Pl. iii. 169 = Coleus barbatus.
Forsteri, *Benth. Lab. Gen. et Sp.* 38.—Ins. N. Hebrid.
fruticosus, *L'Hérit. Stirp. Nov.* 85. t. 41.—Afr. austr.
fruticosus, *Wight, ex Hook. f. Fl. Brit. Ind.* iv. 623.—Ind. or.
furcatus, Hort. ex Steud. Nom. ed. II. ii. 353 = Craniotome versicolor.
galeatus, Vahl, Symb. Bot. i. 43 = Coleus galeatus.
Gardneri, Thw. Enum. Pl. Zeyl. 237 = subincisus.
Gerardianus, *Benth. in Wall. Pl. As. Rar.* ii. 17.—Reg. Himal.
glandulosus, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 17.—Afr. trop.
glaucocalyx, *Maxim. Prim. Fl. Amur.* 212.—Amur.
graciliflorus, Benth. in DC. Prod. xii. 56 = Gerardianus.
graveolens, R. Br. Prod. 506 = parviflorus.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 623.—Reg. Himal.
Helferi, *Hook. f. l. c.*—Burma.
hexaphyllus, *Baker, in Journ. Linn. Soc.* xx. (1883) 231.—Madag.
hians, Benth. in DC. Prod. xii. 57 = nigrescens.
hirtus, *Benth. Lab. Gen. et Sp.* 38.—Afr. austr.
hispidus, *Benth. in Wall. Pl. As. Rar.* ii. 17.—Reg. Himal.
Hookeri, C. B. Clarke, ex Hook. f. Fl. Brit. Ind. iv. 620 = P. Coetsa.
incanus, *Link, Enum. Hort. Berol.* ii. 120.—Ind. or.
inconspicuus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 101 = inflexus.
indicus, Spreng. Syst. ii. 691 = Ocimum adscendens.
inflexus, *Vahl, ex Benth. Lab. Gen. et Sp.* 711.—Japon.
ingratus, Benth. in DC. Prod. xii. 74, in syn. = Coleus atropurpureus.
ingratus, Blume, Bijdr. 836 = Coleus ingratus.
insignis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 210.—Afr. trop.
intermedius, *Benth. in DC. Prod.* xii. 56.—Zeylan.
intermedius, *Zoll. & Mor. Syst. Verz. Zoll.* 55.—Java.
javanicus, *Benth. Lab. Gen. et Sp.* 45.—Java.
laciniatus, Blume, Bijdr. 838 = Coleus laciniatus.
lanceolatus, *Boj. ex Benth. Lab. Gen. et Sp.* 40.—Madag.
lavanduloides, *Baker, in Journ. Linn. Soc.* xx. (1883) 230.—Madag.
laxiflorus, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 228.—Afr. austr.
Leschenaultii, *Benth. Lab. Gen. et Sp.* 34.—Java.
longicornis, *F. Muell. Fragm.* v. 51.—Austral.
longiflorus, *Benth. Lab. Gen. et Sp.* 33.—Madag.
longitubus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 102.—Japon.
lucidus, Burch. ex Benth. in DC. Prod. xii. 68 = strigosus.

PLECTRANTHUS:—
lycopifolius, *A. Rich. Tent. Fl. Abyss.* ii. 181.—Abyss.
Macraei, Benth. Lab. Gen. et Sp. 42 = menthoides.
macranthus, *Hook. f. Fl. Brit. Ind.* iv. 616.—Reg. Himal.; Burma.
macrophyllus, Blume, Bijdr. 835 = Coleus macrophyllus.
maculatus, Boj. ex Benth. Lab. Gen. et Sp. 52 = Coleus Bojeri.
maculosus, Salisb. Prod. 88 = punctatus.
madagascariensis, *Benth. Lab. Gen. et Sp.* 37.—Madag.; Afr. austr.
Maddeni, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 620.—Reg. Himal.
malabaricus, Herb. Madr. ex Wall. Cat. n. 2735 = Coleus malabaricus.
marmoritis, *Hance, in Journ. Bot.* xii. (1874) 53.—China.
marrubioides, *Hochst. ex Benth. in DC. Prod.* xii. 60.—Abyss.
mauritianus, Boj. Hort. Maurit. 254 = madagascariensis.
Maximowiczii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 101 = glaucocalyx.
maypurensis, Spreng. Syst. ii. 691 = incanus.
melissoides, *Benth. Lab. Gen. et Sp.* 39.—Reg. Himal.
Melleri, *Baker, in Journ. Bot.* xx. (1882) 243.—Madag.
menthoides, *Benth. in Wall. Pl. As. Rar.* ii. 17.—Ind. or.
menthoides, Mor. Syst. Verz. Zoll. 55 = rufescens.
menthoides, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 181 = Ocimum adscendens, Orthosiphon tomentosus.
micranthus, Spreng. Syst. ii. 691 = Moschosma polystachyum.
mollis, Spreng. l. c. 690 = incanus.
mollissimus, Wall. ex Benth. Lab. Gen. et Sp. 26 = Orthosiphon rubicundus.
monachorum, Spreng. Syst. ii. 690 = Ocimum sanctum.
monadelphus, Buch.-Ham. ex Wall. Cat. n. 2729 = Coleus spicatus.
monadelphus, Roxb. Hort. Beng. 45; Fl. Ind. iii. 22 = Coleus barbatus.
montanus, *Benth. in Wall. Pl. As. Rar.* ii. 17.—Ind. or.
montanus, Wight, ex Hook. f. Fl. Brit. Ind. iv. 617 = nigrescens.
moschatus, R. Br. Prod. 506 = Moschosma australe.
multiflorus, Hochst. ex Benth. in DC. Prod. xii. 49 = Moschosma multiflora.
mysurensis, Heyne, ex Wall. Cat. n. 2728 = Coleus barbatus.
nankinensis, *Spreng. Syst.* ii. 691.—China.
nepetaefolius, *Benth. in DC. Prod.* xii. 57.—Ind. or.
nigrescens, *Benth. Lab. Gen. et Sp.* 710.—Zeylan.
nilgherricus, *Benth. in DC. Prod.* xii. 57.—Ind. or.
nudiflorus, Willd. Sp. Pl. iii. 168 = Coleus scutellarioides.
oblongifolius, Wall. Pl. As. Rar. ii. 16 = Gerardianus.
ocymoides, Herb. Madr. ex Wall. Cat. n. 2732 = Coleus scutellarioides.
ovatifolius, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 136.—Afr. trop.
ovatus, *Benth. Lab. Gen. et Sp.* 709.—Arab.
Palisoti, Benth. l. c. 39 = Solenostemon ocymoides.
paniculatus, Jacq. Fragm. 62 = parviflorus.
Parishii, *Hook. f. Fl. Brit. Ind.* iv. 622.—Burma.
parviflorus, R. Br. Prod. 506 = Moschosma polystachyum.
parviflorus, *Willd. Hort. Berol.* t. 65.—Austral.; Ins. Pacif.
Patchouli, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 624.—Reg. Himal.
paucicrenatus, *Franch. in Révoil, Çomali*, 56.—Afr. trop.
pekinensis, Maxim. Prim. Fl. Amur. 213, in nota = amethystoides.
petiolaris, *E. Mey. Comm. Pl. Afr. Austr.* 228.—Afr. austr.
pubescens, Willd. ex Benth. in Linnaea, xi. (1837) 344 = madagascariensis.
pulneyensis, *Hook. f. Fl. Brit. Ind.* iv. 617.—Ind. or.
punctatus, *L'Hérit. Stirp. Nov.* 87 t. 42.—Abyss.
purpuratus, *Harv. Thes. Cap.* i. 53. t. 83.—Afr. austr.
ramosissimus, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 17.—Afr. trop.

PLECTRANTHUS:—
rariflorus, Hochst. in Flora, xxviii. (1845) 67 = Ocimum obtusifolium.
repens, *Wall. Pl. As. Rar.* ii. 17.—Reg. Himal.
riparius, Hochst. in Flora, xxviii. (1845) 67 = Moschosma riparium.
rivularis, *Wight, ex Hook. f. Fl. Brit. Ind.* iv. 617.—Ind. or.
rotundifolius, *Spreng. Syst.* ii. 690.—Ins. Mascar.
rubicundus, D. Don, Prod. Fl. Nep. 116 = Orthosiphon rubicundus.
rufescens, *Benth. in DC. Prod.* xii. 59.—Java.
rugosus, *Wall. Pl. As. Rar.* ii. 17.—Reg. Himal.
Rutenbergianus, *Vatke, in Bremen, Abh.* ix. (1885) 135.—Madag.
saccatus, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 227.—Afr. austr.
salvioides, *Benth. Lab. Gen. et Sp.* 45.—Ind. or.
scabrellus, Benth. in DC. Prod. xii. 58 = nigrescens.
Schimperi, *Vatke, in Linnaea*, xxxvii. (1871–73) 317.—Abyss.
scrophularioides, *Wall. Pl. As. Rar.* ii. 16.—Reg. Himal.
scutellarioides, Blume, Bijdr. 837 = Coleus Blumei.
scutellarioides, R. Br. Prod. 506 = Coleus scutellarioides.
secundus, Roxb. Hort. Beng. 45; Fl. Ind. iii. 20 = incanus.
Serra, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 454.—Manchuria.
Sieberi, Benth. Lab. Gen. et Sp. 710 = parviflorus.
sinensis, *Miq. in Journ. Bot. Néerl.* i. (1861) 115.—China.
spicatus, *E. Mey. Comm. Pl. Afr. Austr.* 230.—Afr. austr.
spicatus, Wight, ex Wall. Cat. n. 2729 = Coleus scutellarioides.
stachyoides, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 136.—Afr. trop.
Stocksii, *Hook. f. Fl. Brit. Ind.* iv. 618.—Ind. or.
Stracheyi, *Benth. ex Hook. l. c.*—Reg. Himal.
striatus, *Benth. in Wall. Pl. As. Rar.* ii. 17.—Reg. Himal.
striatus, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 618 = Stocksii.
striatus, Wight, ex Hook. f. l. c. 617 = pulneyensis.
strigosus, *Benth. ex E. Mey. Comm. Pl. Afr. Austr.* 229.—Afr. austr.
strobiliferus, Roxb. Hort. Beng 45; Fl. Ind. iii. 23 = Anisochilus carnosus.
subincisus, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Ind. or.
subspicatus, Hochst. in Flora, xxviii. (1845) 67 = spicatus.
ternatus, *Sims, Bot. Mag.* t. 2460.—Madag.
ternifolius, *D. Don, Prod. Fl. Nep.* 117.—Reg. Himal.; China.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 944.—Java.
Thunbergii, *Benth. Lab. Gen. et Sp.* 37.—Afr. austr.
thymiflorus, Spreng. Syst. ii. 690 = Orthosiphon tomentosus.
tomentosus, *Benth. ex E. Mey. Comm. Pl. Afr. Austr.* 229.—Afr. austr.
tristis, Spreng. Syst. ii. 690 = Orthosiphon tomentosus.
tuberosus, Blume, Bijdr. 838 = Coleus parviflorus.
tuberosus, Roxb. ex Benth. Lab. Gen. et Sp. 26 = Orthosiphon rubicundus.
urticaefolius, Salisb. Prod. 88 = fruticosus, *L'Hérit.*
urticifolius, *Hook. f. Fl. Brit. Ind.* iv. 622.—Ind. or.
veronicifolius, *Hance, in Journ. Bot.* xxiii. (1885) 327.—China.
vestitus, *Benth. Lab. Gen. et Sp.* 32.—Madag.
villosus, Sieber, ex Benth. Lab. Gen. et Sp. 38 = madagascariensis.
virgatus, D. Don, Prod. Fl. Nep. 116 = Orthosiphon rubicundus.
viscosus, Spreng. Syst. ii. 691 = Ocimum viscosum.
Walkeri, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 354.—Ind. or.
Wightii, *Benth. Lab. Gen. et Sp.* 41.—Ind. or.
Wightii, J. Grah. Cat. Pl. Bomb. 148 = Stocksii.
zeylanicus, *Benth. Lab. Gen. et Sp.* 36.—Zeylan.

PLECTRECA, Rafin. Fl. Tellur. iv. 119 (1836) = **Vernonia**, Schreb. (Compos.).
corymbosa, Rafin. l. c. 120 = V. corymbosa.

PLECTRELMINTHUS, Rafin. Fl. Tellur. iv. 42 (1836) = **Angraecum**, Bory (Orchid.).
bicolor, Rafin. l. c. = A. caudatum.

PLECTRITES, DC. Mém. Valér. 13; Prod. iv. 651 (1830) = **Valerianella**, Tourn.
brachystemon, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 47 = V. congesta.
capitata, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 50 = V. aphanoptera.
congesta, DC. Prod. iv. 631 = V. congesta.
longiflora, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 51 = V. longiflora, *Walp.*
macrocera, Torr. & Gray, Fl. N. Am. ii. 50 = V. macrocera.
major, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 37 = V. samolifolia.
samolifolia, Hoeck, l. c. = V. samolifolia.
spathulata, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 51 = V. Nuttallii.

PLECTROCARPA, Gill. in Hook. Bot. Misc. iii. (1833) 166. *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 268.
tetracantha, *Gill. l. c.*—Chili.

PLECTRONEMA, Rafin. Fl. Tellur. iv. 10 (1836) = **Zephyranthes**, Herb. (Amaryll.).
candida, Rafin. l. c. = Z. candida.

PLECTRONIA, Buching. ex Krauss, in Flora, xxvii. (1844) 348 = **Olinia**, Thunb. (Lythrar.).
floribunda, Buching. l. c. = O. cymosa.

PLECTRONIA, Linn. Mant. i. 16 (1767). *RUBIACEAE*, Benth. & Hook. f. ii. 110.
CANTHIUM, Lam. Encyc. i. 602 (1783).
CLUSIOPHYLLEA, Baill. Adansonia, xii. (1878) 197.
DONDISIA, DC. Prod. iv. 469 (1830).
LYCIOSERISSA, Roem. & Schult. Syst. iv. p. xxiv. 353 (1819).
MITRASTIGMA, Harv. in Hook. Lond. Journ. i. (1842) 20.
PHALLARIA, Schum. & Thonn. Beskr. Guin. Pl. 112 (1827).
PSILOSTOMA, Klotzsch, in Eckl. & Zeyh. Enum. 361 (1837).
PSYDRAX, Gaertn. Fruct. i. 125. t. 26 (1788).
WEBERA, Cramer, Disp. Syst. 38 (1803).
acuminata, *Baker, Fl. Maurit.* 147.—Ins. Seychell
angustifolia, *Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 154.—Ind. or.
bibracteata, *Baker, Fl. Maurit.* 146.—Ins. Mascar.
Boiviniana, *Baker, in Journ. Linn. Soc.* xxi. (1885) 411.—Madag.
buxifolia, *Baker, l. c.*—Madag.
campanulata, *Bedd. Forest Man. Bot.* 134/6.—Zeylan.
carinata, *Baker, Fl. Maurit.* 147.—Ins. Seychell.
celastroides, *Baker, l. c.* 146.—Ins. Seychell.
ciliata, *D. Dietr. Syn. Pl.* i. 856.—Afr. austr.
citrifolia, *Eckl. & Zeyh. Enum.* 363.—Afr. austr.
corymbosa, Burm. f. Fl. Cap. Prod. 6 = ventosa, Celastrus acuminatus.
densiflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 167.—Madag.
didyma, *Bedd. Forest. Man. Bot.* 134/5.—Ind. or.
didyma, Brand. For. Fl. 276 = umbellata.
didyma, Kurz, For. Fl. Brit. Burma, ii. 35 = parvifolia, *Benth. & Hook. f.*
glabra, *Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 153.—Burma.
gracilipes, *Kurz, For. Fl. Brit. Burma*, ii. 36.—Ins. Andaman.
hirsuta, *DC. Prod.* iv. 476.—Afr. trop.
horrida, *Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 153.—Burma; Malaya.
Leschenaultii, Bedd. Forest. Man. Bot. 134/6 = angustifolia?
Macconneli, Horne, ex Baker, in Journ. Linn. Soc. xx. (1883) 363 = Pisonia grandis.
Macgregori, *Horne, l. c.*—Ins. Fiji.
macrocarpa, *Bedd. Forest. Man. Bot.* 134/6.—Ind. or.
madagascariensis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 189.—Madag.
monstrosa, *A. Rich. l. c.*—Ins. Philipp.
Mundtiana, Sond. in Harv. & Sond. Fl. Cap. iii. 17 = seq.
Mundtii, *Poepp. Sylva Cap.* ed. I. 19.—Afr. austr.
neilgherrensis, *Bedd. Forest. Man. Bot.* 134/6.—Ind. or.

PLECTRONIA:—
parviflora, *Bedd. Forest. Man. Bot.* 134/5.—Ind. or.
parviflora, Harv. & Sond. Fl. Cap. iii. 17 = pauciflora.
parvifolia, *Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 153.—Burma; Malaya.
pauciflora, *Klotzsch, ex Eckl. & Zeyh. Enum.* 363.—Afr. austr.
Rheedii, *Bedd. Forest. Man. Bot.* 134/5.—Ind. or.
Schimperi, *Hochst. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 136.—Abyss.
Schimperiana, Vatke, in Linnaea, xl. (1876) 195 = praec.
spinosa, *Klotzsch, ex Eckl. & Zeyh. Enum.* 362.—Afr. austr.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 58.—Ind. or.
trilocularis, *Baker, Fl. Maurit.* 147.—Ins. Rodriguez.
umbellata, *Baker, in Journ. Linn. Soc.* xx. (1883) 168.—Madag.
umbellata, *Benth. & Hook. f. Gen.* ii. 115.—Ins. Philipp.
venosa, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 85.—Afr. trop.
ventosa, *Linn. Mant.* i. 52.—Afr. austr.
viburnoides, *Baker, Fl. Maurit.* 146.—Ins. Maurit.
zanzibarica, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 231.—Afr. trop.

PLECTRONIA, Lour. Fl. Cochinch. 162 (1790) = **Acanthopanax**, Decne. & Planch. (Araliac.).
chinensis, Lour. l. c. = A. aculeatum.

PLECTROPHORA, H. C. Focke, in Tijdschr. Nat. Wetensch. i. (1848) 212 = **Comparettia**, Poepp. & Endl. (Orchid.).
iridifolia, H. C. Focke, l. c.—Guian.

PLECTROTROPIS, Schum. & Thonn. Beskr. Guin. Pl. 338 (1827) = **Vigna**, Savi (Legumin.).
angustifolia, Schum. & Thonn. l. c. = V. capensis.
hirsuta, Schum. & Thonn. l. c. 339 = V. capensis.
striata, Steud. Nom. ed. II. ii. 354 = V. capensis.

PLECTRURUS, Rafin. Neogenyt. 4 (1825) = **Tipularia**, Nutt. (Orchid.).
discolor, Rafin. l. c. = T. discolor.

PLEEA, Michx. Fl. Bor. Am. i. 247. t. 25 (1803). *LILIACEAE*, Benth. & Hook. f. iii. 829.
Sieberi, Reichb. ex Benth. Fl. Austral. vii. 30 = Anguillaria dioica.
tenuifolia, *Michx. Fl. Bor. Am.* i. 248.—Am. bor.

PLEGERINA, Arruda, ex Koster, Trav. Bras. 499 (1816). *INC. SED.*
PLERAGINA, Steud. Nom. ed. I. 632 (1821).
odorata, *Arruda, l. c.* 499, 500.—Bras.
rufa, *Arruda, l. c.* 499.—Bras.
umbrosissima, *Arruda, l. c.* 499, 500.—Bras.

PLEGORHIZA, Molina, Sagg. Chile, ed. I. 164, 351 (1782) = **Statice**, Linn. (Plumbag.).
adstringens, Willd. Sp. Pl. ii. 487 = S. chilensis.
Guaicuru, Molina, Sagg. Chile, ed. I. 164, 351 = S. chilensis.
Quicura, Steud. Nom. ed. II. ii. 354 = praec.

PLEGORRHIZA, Spreng. Anleit. ii. II. 884 (1817) = praec.

PLEIARINA, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
Humboldtiana, Rafin. l. c. = S. Humboldtiana.

PLEIENTA, Rafin. Fl. Tellur. iii. 30 (1836) = **Sabbatia**, Adans. (Gentian.).
capitata, Rafin. l. c. = Erythraea Centaurium?
dodecandra, Rafin. l. c. = S. chloroides.
fasciculata, Rafin. l. c.—Hab.?
flexuosa, Rafin. l. c.—Hab.?
leucantha, Rafin. l. c.; et New Fl. Am. iv. 92 = S. chloroides.
quinquenervia, Rafin. New Fl. Am. iv. 91.—Am. bor.
rigida, Rafin. Fl. Tellur. iii. 30.—Hab.?

PLEIMERIS, Rafin. Sylva Tellur 21 (1838) = **Thunbergia**, Retz. (Acanth.).
capensis, Rafin. l. c. = T. capensis.

PLEIOCARPA, Benth. in Benth. & Hook. f. Gen. ii. 699 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
mutica, *Benth. in Hook. Ic. Pl.* t. 1181.—Afr. trop.
rostrata, *Benth. l. c.* t. 1182.—Afr. trop.

PLEIOCHITON, Naud. ex A. Gray, Bot. U. St. Expl. Exped. i. 583. t. 73 (1854). *MELASTOMACEAE*, Benth. & Hook. f. i. 764.
crassifolium, *Naud. ex A. Gray, l. c.* 583. t. 73.—Am. austr.
ebracteatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 96.—Bras.

PLEIOCOCCA, F. Muell. Fragm. ix. 117 (1875). *RUTACEAE.*
Wilcoxiana, *F. Muell. l. c.*—Austral.

PLEIOGYNE, C. Koch, in Bot. Zeit. i. (1843) 40 = **Cotula**, Tourn. (Compos.).
anthemoides, C. Koch, l. c. = C. anthemoides.
australis, C. Koch, l. c. = C. australis.
cardiospermum, Edgew. in Trans. Linn. Soc. xx. (1846) 71 = C. anthemoides.
microcephala, C. Koch, in Bot. Zeit. i. (1843) 40 = C. microcephala.
multifida, Sond. in Linnaea, xxv. (1852) 484 = C. reptans.
nudicaulis, C. Koch, in Bot. Zeit. i. (1843) 40 = C. nudicaulis.
reptans, C. Koch, l. c. = C. reptans.
villosa, C. Koch, l. c. = C. villosa.

PLEIOGYNIUM, Engl. in DC. Monog. Phan. iv. 255 (1883). *ANACARDIACEAE.*
Solandri, *Engl. l. c.*—Austral.

PLEIOMERIS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 87 = **Myrsine**, Linn.
canariensis, A. DC. l. c. 88 = M. canariensis.

PLEIONE, D. Don, Prod. Fl. Nep. 36 (1825) = **Coelogyne**, Lindl. (Orchid.).
diphylla, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 66 = C. maculata?
humilis, D. Don, Prod. Fl. Nep. 37 = C. humilis.
Lagenaria, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 5 = C. Lagenaria.
maculata, Lindl. & Paxt. l. c. = C. maculata.
praecox, D. Don, Prod. Fl. Nep. 37 = C. praecox.
Wallichiana, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 66 = C. praecox.

PLEIOSMILAX, Seem. Journ. Bot. vi. (1868) 193. t. 81 = **Smilax**, Linn. (Liliac.).
Menziesii, Seem. l. c. 194. t. 81 = S. melastomifolia.
sandwicensis, Seem. l. c. 193 = S. sandwicensis.
vitiensis, Seem. l. c. = S. vitiensis.

PLEIOSPORA, Harv. Thes. Cap. i. 51. t. 81 (1859). *LEGUMINOSAE*, Benth. & Hook. f. i. 475.
cajanifolia, *Harv. l. c.* 51, 52.—Afr. austr.

PLEIOSTEMON, Sond. in Linnaea, xxiii. (1850) 135 = **Securinega**, Juss. (Euphorb.).
verrucosum, Sond. l. c. 136 (= *Phyllanthus verrucosus*).—Afr. austr.

PLEIOTAXIS, Steetz, in Peters, Reise Mossamb. Bot. 499. t. 51 (1863). *COMPOSITAE*, Benth. & Hook. f. ii. 487.
pulcherrima, *Steetz, l. c.* 500.—Afr. trop.

PLEISOLIRION, Rafin. Fl. Tellur. ii. 28 (1836) = **Paradisea**, Mazz. (Liliac.).
Liliastrum, Rafin. l. c. = Paradisea Liliastrum.

PLENCKIA, Moç. & Sesse, ex DC. Prod. i. 724 (1824) = **Choisya**, H. B. & K. (Rutac.).

PLENCKIA, Rafin. Specch. i. 194 (1814) = **Mollugo**, Linn. (Ficoid.).
setiflora, Rafin. l. c. = M. hirta.

PLENCKIA, Reiss. in Mart. Fl. Bras. xi. I. 30 (1861). *CELASTRINEAE*, Benth. & Hook. f. i. 368.
populnea, *Reiss. l. c.* 31.—Bras.

PLEOCARPHUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 228 = **Jungia**, Linn. f. (Compos.).
dentatus, Phil. in Anal. Univ. Chil. (1861) 51; et in Linnaea, xxxiii. (1864–65) 120.—Chili.
revolutus, D. Don, in Trans. Linn. Soc. xvi. (1830) 228.—Chili.

PLEOCARPUS, Walp. Ann. i. 997 (1848–49) = praec.

PLEOGYNE, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43. *MENISPERMACEAE*, Benth. & Hook. f. i. 963.
MICROCLISIA, Benth. in Benth. & Hook. f. Gen. i. Pars 1ma, Add. 435 (1862).
australis, *Benth. Fl. Austral.* i. 59.—Austral.
Cunninghami, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 324 = australis.

PLEOMELE, Salisb. Prod. 245 (1796) = **Dracaena**, Vand. (Liliac.).
aloifolia, Salisb. l. c. = Sanseviera guineensis.
fragrans, Salisb. l. c. = D. fragrans.

PLEONANTHUS, Ehrh. Beitr. iv. 148 (1789) = **Dianthus**, Linn. (Caryoph.).

PLEONOTOMA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 184 = **Bignonia**, Linn.
chondrogona, Miers, l. c. = B. chondrogona.
Clematis, Miers, l. c. = B. Clematis.
decomposita, Miers, l. c. = B. decomposita.
flava, Miers, l. c. 185 = B. flava.
flavida, Miers, l. c. = B. flavida.
glaberrima, Miers, l. c. 184 = B. glaberrima.
inaequalis, Miers, l. c. = B. inaequalis.
jasminifolia, Miers, l. c. = B. jasminifolia.
lanceolata, Miers, l. c. = Tabebuia lanceolata.
moringaefolia, Miers, l. c. = B. moringaefolia.
peregrina, Miers, l. c. 183.—Bras.
tetragonocaulis, Miers, l. c. 184 = B. tetragonocaulos.
triphylla, Miers, l. c. 185 = B. triternata.
triternata, Miers, l. c.—Bras.
variabilis, Miers, l. c. 184 = B. variabilis.

PLEOPOGON, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 189 = **Lycurus**, H. B. & K. (Gramin.).
setosum, Nutt. l. c.—Calif.

PLEOROTHYRIUM, Endl. Ench. 197 (1841) = **Pleurothyrium**, Nees (Laurin.).

PLEOTHECA, Wall. Cat. nn. 6215, 6216 (1830) = **Spiradiclis**, Blume (Rubiac.).
bifida, Wall. Cat. n. 6216 = S. bifida.
cylindrica, Wall. l. c. n. 6215 = S. cylindrica.

PLERAGINA, Steud. Nom. ed. I. 632 (1821) = **Plegerina**, Arruda (Inc. sed.).

PLERANDRA, A. Gray, Bot. U. St. Expl. Exped. i. 729. t. 95 (1854). *ARALIACEAE*, Benth. & Hook. f. i. 946.
BAKERIA, Seem. Journ. Bot. ii. (1864) 248.
NESOPANAX, Seem. l. c. 249 (1864).
Grayi, *Seem. Miss. Viti*, 437, nomen; *Journ. Bot.* ii. (1864) 242.—Ins. Fiji.
jatrophifolia, *Hance, in Journ. Bot.* xix. (1881) 275.—Hab.?
paucidentata, Baill. Adansonia, xii. (1878) 137 = Tetraplasandra paucidens.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 729. t. 95.—Ins. Fiji.
Vieillardi, *Baill. Adansonia*, xii. (1878) 136.—N. Caled.
vitiensis, *Baill. Hist. Pl.* vii. 169.—Ins. Fiji.

PLEROMA, D. Don, in Mem. Wern. Soc. iv. (1823) 293 = **Tibouchina**, Aubl. (Melastom.).
adenostemon, A. Gray, Bot. U. St. Expl. Exped. i. 603 = T. adenostemon.
adenostemon, Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. multiflora.
aegopogon, Triana, l. c. 45 = T. aegopogon.
aemulum, Triana, l. c. 41 = T. aemula, hygrophila.
albiflorum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 347 = T. hospita.

PLEROMA:—
angustifolium, Triana, in Trans. Linn. Soc. xxviii. (1871) 47 = Tibouchina angustifolia.
arboreum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 351 = T. arborea.
argenteum, Gardn. l. c. i. (1842) 172 = T. holosericea.
asperior, Triana, in Trans. Linn. Soc. xxviii. (1871) 44 = T. asperior.
australe, Triana, l. c. 47 = T. australis.
barbigerum, Triana, l. c. 45. t. 3 = T. barbigera.
Benthamianum, Gardn. in Hook. Bot. Mag. t. 4007 = T. Benthamiana.
bicolor, Triana, in Trans. Linn. Soc. xxviii. (1871) 47. —Bolivia.
bipenicillatum, Triana, l. c. 45 = T. bipenicillata.
Candolleanum, Triana, l. c. 44. t. 3 = T. Candolleana.
capitatum, Triana, l. c. 46.—Bolivia.
cardinale, Triana, l. c. 44 = T. cardinalis.
Chamaecistus, Triana, l. c. 47.—Ins. Martinic.
chironioides, Triana, l. c.—Ins. Dominic.
ciliare, Triana, l. c. 46.—N. Granat.
cistoides, Triana, l. c. 47.—Ins. S. Vincent.
citrinum, Triana, l. c. 46.—Bolivia.
Claussenii, Triana, 41 = T. Claussenii.
clidemioides, Berg, ex Triana, l. c. 43 = T. clidemioides.
clinopodifolium, Triana, l. c. 46 = T. clinopodifolia.
collinum, Triana, l. c. 42 = T. collina.
confertiflorum, Triana, l. c. 46.—Bolivia.
coronatum, Triana, l. c. 41 = T. coronata.
corymbosa, Benth. Pl. Hartw. 181 = T. Lindeniana.
cubensis, Wright, in Sauv. Fl. Cub. 52 = Chaetolepis saturaeioides, cubensis.
debile, Triana, in Trans. Linn. Soc. xxviii. (1871) 45 = T. debilis.
dendroides, Triana, l. c. 44 = T. dendroides.
diffusum, DC. Prod. iii. 152.—Porto Rico.
Diplostegium, Triana, in Trans. Linn. Soc. xxviii. (1871) 40 = T. canescens.
echinatum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 350 = T. Gardneriana.
echinatum, Triana, in Trans. Linn. Soc. xxviii. (1871) 47.—Peruv.
elegans, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 350 = T. elegans.
erigeron, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 45 = Pterolepis striphnocalyx.
eriocladum, Triana, l. c. 47.—N. Granat.
exappendiculatum, Triana, l. c. = T. mollis.
fissinervium, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 352 = T. fissinervia.
Fontanesiana, Gardn. l. c. i. (1842) 171 = T. granulosa.
Fothergilli, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. Fothergillae.
foveolatum, Triana, l. c. 41 = T. foveolata.
frigidulum, Triana, l. c. 42 = T. Martiusiana.
Galeottianum, Triana, l. c. 46.—Mexic.
Gardnerianum, Triana, l. c. 42 = T. Gardneriana.
Gaudichaudianum, A. Gray, Bot. U. St. Expl. Exped. i. 603 = T. Gaudichaudiana.
Gayanum, Triana, in Trans. Linn. Soc. xxviii. (1871) 46.—Peruv.
glandulosum, Triana, l. c. 48.—Guian.
glomeratum, G. Don, ex Loud. Hort. Brit. 174.—Ind. occ.
gracile, A. Gray, Bot. U. St. Expl. Exped. i. 604 = T. gracilis.
granulosum, D. Don, in Mem. Wern. Soc. iv. (1823) 295 = T. granulosa.
heteromallum, D. Don, l. c. = T. heteromalla.
hieracioides, Triana, in Trans. Linn. Soc. xxviii. (1871) 45 = T. hieracioides.
hispidum, Triana, l. c. 46 = T. longifolia.
holosericeum, D. Don, in Mem. Wern. Soc. iv. (1823) 295 = T. holosericea.
hospitum, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. dubia.
Kunthianum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 351 = T. Moricandiana.
Kunthianum, Hook. f. in Bot. Mag. t. 4412 = T. semidecandra.
Kunthianum, Paxt. Mag. Bot. xii. (1846) 125 = T. Benthamianum.
lanceolatum, Griseb. in Goett. Abh. xxiv. (1879) 128 = T. longifolia.
Langsdorfianum, Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. Gardneri, Langsdorffianum.
latifolium, Triana, l. c. 47.—Bolivia.

PLEROMA:—
laxum, DC. Prod iii. 151.—Peruv.
ledifolium, DC. l. c. = Brachyotum ledifolium.
lepidotum, Triana, in Trans. Linn. Soc. xxviii. (1871) 44. t. 3 = Tibouchina lepidota.
Lhotskyanum, Triana, l. c. 43 = T. Lhotzkyana.
longifolium, Triana, l. c. 45 = T. longifolia.
macranthum, Hook. f. Bot. Mag. t. 5721 = T. semidecandra.
macrochiton, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. macrochiton.
martiale, Triana, l. c. 44 = T. martialis.
maurocarpum, Triana, l. c. 47.—Peruv.
Maximilianum, Triana, l. c. 43 = T. Maximiliana.
melastomoides, Triana, l. c. 45 = T. melastomoides, exasperata.
mexicanum, DC. Prod. iii. 152.—Mexic.
micranthella, Wright, in Sauv. Fl. Cub. 51 = T. longifolia.
molle, Triana, in Trans. Linn. Soc. xxviii. (1871) 43 = T. Chamissoana.
Moricandianum, Triana, l. c. 42. t. 2 = T. Moricandiana.
multiflorum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 351 = T. multiflora.
mutabile, Triana, in Trans. Linn. Soc. xxviii. (1871) 40 = T. mutabilis.
Naudinianum, Triana, l. c. 46.—Mexic.
obscurum, Triana, l. c. 44 = T. corymbosa.
ochypetalum, D. Don, in Mem. Wern. Soc. iv. (1823) 295 = T. ochypetala.
Orbignyanum, Triana, in Trans. Linn. Soc. xxviii. (1871) 46.—Bolivia.
ornatum, Triana, l. c. 162. t. 3. f. 31 p. = T. ornata.
paniculare, Triana, l. c.—Bolivia.
papyriferum, Triana, l. c. 45 = T. papyrifera.
parasiticum, Berg, ex Triana, l. c. 46 = T. tuberosa.
paratropicum, Griseb. in Goett. Abh. xix. (1874) 140. —Reg. Argent.
petiolata, Paxt. Mag. Bot. xi. (1844) 267 = T. Gaudichaudiana.
pogonantherum, Triana, in Trans. Linn. Soc. xxviii. (1871) 45 = T. pogonanthera.
Pohlianum, Berg, ex Triana, l. c. 46 = T. canescens.
Preslianum, Triana, l. c. 43 = T. estrellensis.
pulchellum, Griseb. in Goett. Abh. xxiv. (1879) 128 = T. gracilis.
pulchrum, Cogn. in Mart. Fl. Bras. xiv. III. 387, sphalm. = praec.
pulchrum, Triana, in Trans. Linn. Soc. xxviii. (1871) 41 = T. pulchra, Saldanhaei.
Raddianum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 351 = T. Saldanhaei.
Raddianum, Triana, in Trans. Linn. Soc. xxviii. (1871) 41 = T. Raddiana, Sellowiana.
redivivum, Berg, ex Triana, l. c. 43 = T. rediviva.
Regnellii, Triana, l. c. 44 = T. martialis.
rufipile, Triana, l. c. 46.—Mexic.
salviaefolium, Triana, l. c. 44 = T. salviaefolia.
sarmentosum, Hook. f. Bot. Mag. t. 5629 = *laxum*.
scaberrimum, Triana, in Trans. Linn. Soc. xxviii. (1871) 47 = T. scaberrima.
scabriusculum, Triana, l. c. 46.—Mexic.
Schiedianum, Triana, l. c. 45.—Mexic.
semidecandrum, Triana, l. c. 41 = T. semidecandra.
sericans, Triana, l. c. 44 = T. stenocarpa.
setulosum, Triana, l. c. = T. setulosa.
spoliatum, A. Gray, Bot. U. St. Expl. Exped. i. 603 = T. Gaudichaudiana.
squamatum, Wright, in Sauv. Fl. Cub. 51 = Chaetolepis saturaeioides.
stenocarpum, Triana, in Trans. Linn. Soc. xxviii. (1871) 44 = T. stenocarpa.
stenopetalum, Triana, l. c. 46.—Peruv.
Steudelii, Triana, l. c. 48 = T. Lhotzkyana.
strigosum, Triana, l. c. 47 = T. ornata.
striphnocalyx, Triana, l. c. 45 = Pterolepis striphnocalyx.
sulcatum, Triana, l. c. 44 = T. sulcata.
Thereminianum, Triana, l. c. 41 = T. Thereminiana.
Tibouchinum, Triana, l. c. 45. t. 3 = T. aspera.
trichopodum, DC. Prod. iii. 151 = T. trichopoda.
trichopodum, Triana, in Trans. Linn. Soc. xxviii. (1871) 41 = T. multiceps.
trifolium, Triana, l. c. 42 = T. frigidula.
tuberosum, Gardn. ex Triana, l. c. 46 = T. tuberosa.
urceolare, Triana, l. c. 42 = T. urceolaris.
ursinum, Triana, l. c. 43 = T. ursina.

PLEROMA:—
velutinum, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = Tibouchina velutina, Blanchetiana.
villosissimum, Triana, l. c. 43 = T. villosissima.
villosum, DC. Prod. iii. 152 = Melastoma villosa.
vimineum, D. Don, in Mem. Soc. Wern. iv. (1823) 296 = T. viminea.
vimineum, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. Urvilleana.
virgatum, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 350 = T. virgata.
Weddelli, Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = T. Weddellii.

PLESIAGOPUS, Rafin. Fl. Tellur. iv. 78 (1836) = **Ipomoea**, Linn. (Convolv.).
brasiliensis, Rafin. l. c. = I. biloba.
cuneifolia, Rafin. l. c. = I. cuneifolia, *Mart.*?
maritima, Rafin. l. c. = I. teretistigma.
rotundifolia, Rafin. l. c. = I. biloba.
sovana, Rafin. l. c. = I. biloba.

PLESILIA, Rafin. New Fl. Am. iv. 55 (1836); Fl. Tellur. iv. 83 (1836) = **Evolvulus**, Linn. (Convolv.).
cuneifolia, Rafin. Fl. Tellur. iv. 83 = E. cuneifolius.

PLESISA, Rafin. Fl. Tellur. iv. 110 (1836) = **Utricularia**, Linn. (Lentib.).
bipartita, Rafin. l. c. = U. bipartita.

PLESMONIUM, Schott, Syn. Aroid. 34 (1856). *AROIDEAE*, Benth. & Hook. f. iii. 978.
dubium, Schott, Prod. Aroid. 125 = margaritiferum.
margaritiferum, *Schott, Syn. Aroid.* 34.—Ind. or.
nobile, *Schott, in Miq. Ann. Mus. Lugd. Bat.* i. 279 (sp. dub.).—Ins. Ceram.

PLETHIANDRA, Hook. f. in Benth. & Hook. f. Gen. i. 772 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
Motleyi, *Hook. f. l. c.*—Borneo.

PLETHIOSPHACE, Opiz, Seznam, 75 (1852) = **Salvia**, Tourn. (Labiat.).
Leistneriana, Opiz, l. c. = S. pratensis.
pratensis, Opiz, l. c. = S. pratensis.
stenantha, Opiz, l. c. = S. pratensis?
sylvestris, Opiz, l. c. = S. sylvestris.

PLETHOSTEPHIA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 32 = **Cordia**, Linn. (Borag.).
angiocarpa, Miers, l. c. = C. angiocarpa.

PLEUDIA, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
egyptiaca, Rafin. l. c. = S. aegyptiaca.

PLEURACHNE, Schrad. in Comm. Gotting. vii. (1832) 147. t. 4. f. 3; Anal. Fl. Cap. 47. t. 4 (1832) = **Ficinia**, Schrad. (Cyper.).
secunda, Schrad. ll. cc. 147; 47 = F. secunda.
Sieberi, Schrad. ll. cc. 148; 48 = = F. secunda.

PLEURADENIA, Rafin. Atl. Journ. 182 (1833) = **Euphorbia**, Linn.
coccinea, Rafin. l. c. = E. pulcherrima.

PLEURADENIA, Rafin. Neogenyt. 2 (1825) = **Collinsonia**, Linn. (Labiat.).
precox, Rafin. l. c. = C. scabriuscula.
scabra, Rafin. l. c. = C. scabriuscula.

PLEURANDRA, Labill. Nov. Holl. Pl. ii. 5. t. 143 (1806) = **Hibbertia**, Andr. (Dilleniac.).
acerosa, R. Br. ex DC. Syst. i. 422 = H. acerosa.
acicularis, Labill. Nov. Holl. Pl. ii. 6. t. 144 = H. acicularis.
astrotricha, Sieber, ex Spreng. Syst. iv. Cur. Post. 191 = H. Billardieri.
bracteata, R. Br. ex DC. Syst. i. 415 = H. bracteata, *Benth.*
calycina, DC. Syst. i. 422 = H. stricta.
camforosma, Sieber, ex Spreng. Syst. iv. Cur. Post. 191 = H. fasciculata.
candicans, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = nitida.
cinerea, R. Br. ex DC. Syst. i. 417 = H. sericea.

PLEURANDRA :—
cinerea, Sieber, ex Benth. Fl. Austral. i. 28 = Hibbertia Billardieri.
cistiflora, Reichb. Ic. Bot. Exot. i. 57.—Austral.
cistiflora, Sieber, ex Spreng. Syst. iv. Cur. Post. 191 = H. stricta.
cistoidea, Hook. in Mitch. Journ. Trop. Austral. 363 = H. stricta.
Cneorum, DC. Syst. i. 416 = H. nitida.
cognata, Steud. in Lehm. Pl. Preiss. i. 265 = H. acerosa.
crassifolia, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 5 = H. crassifolia.
densiflora, Hook. Journ. Bot. i. (1834) 245 = H. sericea.
diamesogenos, Steud. in Lehm. Pl. Preiss. i. 265 = H. lineata.
empetrifolia, DC. Syst. i. 420 = H. Billardieri.
enervia, DC. l. c. 421.—Austral.
ericifolia, DC. l. c. 420 = H. stricta.
fumana, Sieber, ex Benth. Fl. Austral. i. 27 = H. stricta.
furfuracea, R. Br. ex DC. Syst. i. 417 = H. furfuracea.
glaucophylla, Steud. in Lehm. Pl. Preiss. i. 264 = H. aurea ?
hemignosta, Steud. l. c. 265 = Candollea teretifolia.
hibbertioides, Steud. l. c. = Candollea teretifolia.
hirsuta, Hook. Comp. Bot. Mag. i. (1835) 273 = H. hirsuta.
hypericoides, DC. Syst. i. 421 = H. hypericoides.
incana, Lindl. in Mitch. Three Exped. ii. 156 = H. stricta.
intermedia, DC. Syst. i. 420 = H. pedunculata.
juniperina, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 6 = H. acerosa.
microphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 191 = H. stricta.
mucronata, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 139 = H. mucronata.
nitida, R. Br. ex DC. Syst. i. 416 = nitida.
obovata, R. Br. ex Benth. Fl. Austral. i. 28 = H. Billardieri.
ovata, Labill. Nov. Holl. Pl. ii. 5. t. 143 = H. Billardieri.
parviflora, R. Br. ex DC. Syst. i. 418 = H. Billardieri.
pedunculata, R. Br. l. c. 419 = H. gracilipes.
recurvifolia, Steud. in Lehm. Pl. Preiss. i. 264 = H. recurvifolia.
reticulata, Hook. Journ. Bot. i. (1834) 245 (Pulteneaeae sp. ?).—Tasmania.
riparia, R. Br. ex DC. Syst. i. 419 = H. stricta.
scabra, R. Br. l. c. 418 = H. Billardieri.
sericea, R. Br. l. c. 416 = H. sericea.
striata, G. Don, Gen. Syst. i. 74, sphalm. = *stricta*, Sieber.
stricta, DC. Syst. i. 422 = H. stricta.
stricta, Sieber, ex Spreng. Syst. iv. Cur. Post. 191 = praec. ?
triandra, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 280 = H. acicularis ?
verrucosa, Turcz. l. c. xxv. (1852) II. 139 = H. verrucosa.

PLEURANDRA, Rafin. Fl. Ludov. 95 (1817) = **Pleurostemon**, Rafin. (gen. dub.).—Cf. Benth. & Hook. f. i. 787.
alba, Rafin. l. c. 95 = Pleurostemon album.

PLEURANDROPSIS, Baill. Adansonia, x. (1872) 305 = **Asterolasia**, F. Muell. (Rutac.).
phebalioides, Baill. l. c. 306 = A. pleurandroides.

PLEURANDROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = praec.

PLEURANTHE, Salisb. in Knight, Prot. 49 (1809) = **Protea**, Linn.
amplexicaulis, Knight, l. c. 51 = Prot. amplexicaulis.
bupleurifolia, Knight, l. c.—Afr. austr.
cordifolia, Knight, l. c. = Prot. cordata.
ericaefolia, Knight, l. c. 50.—Afr. austr.
glastifolia, Knight, l. c. = Prot. humilis.
subulaefolia, Knight, l. c.—Afr. austr.

PLEURANTHIUM, Benth. in Journ. Linn. Soc. xviii. (1881) 312; Benth. & Hook. f. Gen. iii. 526 (1883). *ORCHIDEAE*, Benth. & Hook. f. l. c.
adendrobium, *Benth. & Hook. f. l. c.*—Cuba.

PLEURANTHUS, Rich. ex Pers. Syn. i. 65 (1805) = **Dulichium**, Pers. (Cyper.).

PLEURAPHIS, Torr. in Ann. Lyc. N. York, i. (1824) 148. t. 10 = **Hilaria**, H. B. & K. (Gramin.).
Jamesii, Torr. l. c. = H. Jamesii.
mutica, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 95 = H. mutica.
rigida, Thurber, in S. Wats. Bot. Calif. ii. 293 = H. rigida.
sericea, Nutt. ex Benth. in Journ. Linn. Soc. xix. (1881) 62 = H. sericea.

PLEURASTIS, Rafin. Fl. Tellur. iv. 12 (1836) = **Lycoris**, Herb. (Amaryll.).

PLEUREIA, Rafin. Sylva Tellur. 147 (1838) = **Psychotria**, Linn. (Rubiac.).
compressa, Rafin. l. c. = Ps. calycina.

PLEUREMIDIS, Rafin. Fl. Tellur. iv. 67 (1836) = **Thunbergia**, Retz. (Acanth.).
grandiflora, Rafin. l. c. = T. grandiflora.

PLEURENDOTRIA, Rafin. Fl. Tellur. ii. 73 (1836) = **Tellima**, R. Br. (Saxifr.).
parviflora, Rafin. l. c. = T. parviflora.
reniformis, Rafin. l. c. 74 = T. parviflora.

PLEURENODON, Rafin. Fl. Tellur. iii. 80 (1836) = **Hypericum**, Tourn.

PLEURICOSPORA, A. Gray, in Proc. Am. Acad. vii. (1868) 369. *MONOTROPEAE*, Benth. & Hook. f. ii. 608.
fimbriolata, *A. Gray, l. c.*—Calif.

PLEURIMARIA, Rafin. Fl. Tellur. iii. 31 (1836) = **Chlora**, Adans. (Gentian.).
mascariensis, Rafin. l. c. = C. perfoliata.
perfoliata, Rafin. l. c. = C. perfoliata.
quadrifolia, Rafin. l. c. = C. quadrifolia.

PLEURISANTHES, Baill. Adansonia, xi. (1874) 201. *OLACINEAE*.
Artocarpi, *Baill. l. c.*—Guian.

PLEUROBLEPHARON, Kunze, ex Reichb. Consp. 212[a] (1828). (Orchid.—Quid ?).

PLEUROBOTRYUM, Rodrig. Orch. Nov. i. 20 (1877). *ORCHIDEAE*.
atropurpureum, *Rodrig. l. c.*—Bras.

PLEUROCALYPTUS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xiv. (1867) 264. *MYRTACEAE*.
Deplanchei, *Brongn. & Gris, l. c.*—N. Caled.

PLEUROCARPAEA, Benth. Fl. Austral. iii. 460 (1866). *COMPOSITAE*, Benth. & Hook. f. ii. 227.
denticulata, *Benth. l. c.*—Austral.

PLEUROCARPUS, Klotzsch, in Bonplandia, vii. (1859) 4 = **Rhyssocarpus**, Endl. (Rubiac.).
decemfidus, Klotzsch, l. c.—Hab. ?

PLEUROCHAENIA, Griseb. in Goett. Abh. ix. (1861) 55 = **Miconia**, Ruiz & Pav. (Melastom.).
cubensis, Griseb. Cat. Pl. Cub. 101 = M. quadrangularis.
quadrangularis, Griseb. in Goett. Abh. ix. (1861) 55 = M. quadrangularis.
rigida, Griseb. l. c. = M. rigida.

PLEUROCOFFEA, Baill. in Bull. Soc. Linn. Par. i. (1880) 270. *RUBIACEAE*.
Boiviniana, *Baill. l. c.*—Madag.

PLEURODESMIA, Arn. in Edinb. N. Phil. Journ. xvi. (1834) 316 = **Schumacheria**, Vahl (Dilleniac.).
Grahamii, Arn. l. c. = S. castaneaefolia.

PLEUROGYNA, Eschsch. ex Cham. & Schlecht. in Linnaea, i. (1826) 187, 188. *GENTIANEAE*, Benth. & Hook. f. ii. 816.
LOMATOGONIUM, A. Br. in Flora, xiii. (1830) 221.
NARKETIS, Rafin. Fl. Tellur. iii. 26 (1836).

PLEUROGYNA:—
brachyanthera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 120.—Reg. Himal.
carinata, Edgew. in Trans. Linn. Soc. xx. (1846) 85 = carinthiaca.
carinthiaca, *G. Don, Gen. Syst.* iv. 188.—Europ.; As. bor.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 91 = carinthiaca.
Lubahniana, *Vatke, in Bremen, Abh.* ix. (1885) 127.—Madag.
minor, *Benth. & Hook. f. Gen.* ii. 816.—Ind. or.
Purshii, Steud. Nom. ed. II. ii. 355 = carinthiaca.
rotata, G. Don, Gen. Syst. iv. 188 = carinthiaca.
rotata, Franch. & Sav. Enum. Pl. Jap. ii. 450 = Swertia diluta.
rotata, *Griseb. Gen. et Sp. Gent.* 309.—Am. arct.
spathulata, *Kern, in Ber. Naturw. Ver. Innsbruck*, i. (1870) 104; *Nov. Pl. Sp. Dec.* ii. 8.—Reg. Himal.
Stelleriana, G. Don, Gen. Syst. iv. 188 = carinthiaca.
sulcata, G. Don. l. c. = rotata.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 120.—Reg. Himal.

PLEUROGYNE, Griseb. Gen. et Sp. Gent. 309 (1839) = praec.

PLEUROLOBUS, Jaume St. Hil. in Nouv. Bull. Soc. Philom. iii. (1812) 192 = **Desmodium**, Desv. (Legumin.).
brevifolia, Jaume St. Hil. in Desv. Journ. Bot. i. (1813) 61 = Aeschynomene brevifolia.
gangeticus, Jaume St. Hil. in Nouv. Bull. Soc. Philom. iii. (1812) 192 = E. gangeticum.
heterocarpus, Jaume St. Hil. in Desv. Journ. Bot. i. (1813) 61 = D. polycarpum.
hirtus, Jaume St. Hil. in Nouv. Bull. Soc. Philom. iii. (1812) 192 = Aeschynomene hirta.
maculatus, Jaume St. Hil. l. c. = D. gangeticum.
pubescens, Jaume St. Hil. in Desv. Journ. Bot. i. (1813) 61 = Aeschynomene pubescens.
remotus, Jaume St. Hil. in Nouv. Bull. Soc. Philom. iii. (1812) 192 = D. Scalpe.
spicatus, Jaume St. Hil. l. c. = D. incanum.
triflorus, Jaume St. Hil. l. c. = D. triflorum.

PLEUROMENES, Rafin. Sylva Tellur. 144 (1838) = **Prosopis**, Linn. (Legumin.).
heterocarpa, Rafin. l. c. 145 = Pros. Stephaniana.

PLEUROPAPPUS, F. Muell. in Trans. Vict. Inst. i. (1855) 37 = **Angianthus**, Wendl. (Compos.).
phyllocalymmeus, F. Muell. l. c. = A. Pleuropappus.

PLEUROPETALON, Blume, Mus. Bot. Lugd. Bat. i. 248 (1850) = **Villaresia**, Ruiz & Pav. (Olacin.).
samoënse, A. Gray, Bot. U. St. Expl. Exped. i. 299.—Ins. Pacif.
suaveolens, Blume, Mus. Bot. Lugd. Bat. i. 248.—Java.

PLEUROPETALUM, Benth. & Hook. f. Gen. i. 354 (1862) = praec.

PLEUROPETALUM, Hook. f. in Proc. Linn. Soc. i. (1845) 278; et in Trans. Linn. Soc. xx. (1847) 221. *PORTULACEAE*, Benth. & Hook. f. i. 157.
ALLOCHLAMYS, Moq. in DC. Prod. xiii. II. 463 (1849).
MELANOCARPUM, Hook. f. in Benth. & Hook. f. Gen. iii. 24 (1880).
costaricense, *Hort. Kew. ex Hemsl. Biol. Centr. Am. Bot.* iii. 12.—Am. trop.
Darwinii, *Hook. f. in Hook. Lond. Journ. Bot.* v. (1846) 109. t. 2.—Ins. Galapag.

PLEUROPHORA, D. Don, in Edinb. N. Phil. Journ. (Oct.-Dec. 1831) 112. *LYTHRARIEAE*, Benth. & Hook. f. i. 779.
anomala, *Koehne, in Mart. Fl. Bras.* xiii. II. 307.—Bras.
pilosiuscula, C. Gay, Fl. Chil. ii. 373 = polyandra.
polyandra, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 315.—Chili.
pungens, *D. Don, in Edinb. N. Phil. Journ.* (*Oct.-Dec.* 1831) 112.—Chili.
pusilla, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 315.—Chili.
saccocarpa, *Koehne, in Engl. Bot. Jahrb.* ii. (1882) 426.—Paraguay.
squarrosa, Walp. Rep. iv. 644 = polyandra.

PLEUROPHYLLUM, Hook. f. Fl. Antarct. i. 30. t. 22, 24 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 278.
ALBINEA, Homb. & Jacquinot, ex Decne. Bot. Voy. Astrol. & Zél. 37 (1853).
PACHYTHRIX, Hook. f. Fl. Antarct. i. 32, in syn. (1844).
criniferum, *Hook. f. Fl. Antarct.* i. 32. t. 24, 25.—Ins. Auckland.
Hombronii, *Decne. in Bot. Voy. Astrol. & Zél.* 36.—Ins. Auckland.
Hookeri, *J. Buch. in Trans. N. Z. Inst.* xvi. (1884) 395.—N. Zel.
oresigenesum, Decne. Bot. Voy. Astrol. & Zél. 37 = criniferum.
speciosum, *Hook. f. Fl. Antarct.* i. 31. tt. 22, 23.—Ins. Auckland.

PLEUROPLITIS, Trin. Fund. Agrost. 174 (1820) = **Arthraxon**, Beauv. (Gramin.).
caucasica, Rupr. ex Regel, in Bull. Acad. Pétersb. x. (1866) 373 = A. ciliaris.
centrasiatica, Griseb. in Ledeb. Fl. Ross. iv. 477 = A. ciliaris.
ciliata, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 152 = A. ciliaris.
Junghuhnii, Nees, ex Steud. Syn. Pl. Gram. 414 = A. Junghuhnii.
lancifolia, Regel, in Bull. Acad. Pétersb. x. (1866) 375 = A. microphyllus.
Langsdorffii, Trin. Fund. Agrost. 175. t. 16 = A. ciliaris.
major, Arn. ex Miq. Fl. Ind. Bat. iii. 481 = A. Junghuhnii.
major, Regel, in Bull. Acad. Pétersb. x. (1866) 370 = A. cuspidatus.
microphylla, Regel, l. c. = A. microphyllus.
plumbea, Nees, ex Steud. Syn. Pl. Gram. 414 = A. ciliaris.
producta, Griseb. in Ledeb. Fl. Ross. iv. 478 = Pollinia tenuis.
Quartiniana, Regel, in Bull. Acad. Pétersb. x. (1866) 376 = A. ciliaris.
Schimperi, Regel, l. c. 369 = A. microphyllus.
violacea, Nees, ex Steud. Syn. Pl. Gram. 414 = A. ciliaris.

PLEUROPOGON, R. Br. in Parry, 1st Voy. Suppl. 289 (err. typ. 189) (1824). *GRAMINEAE*, Benth. & Hook. f. iii. 1192.
LOPHOCHLAENA, Nees, in Ann. Nat. Hist. Ser. I. i. (1838) 283.
californicum, *Benth. ex Vasey, Grass. U. St.* 40.—Calif.
Douglasii, *Trin. ex Steud. Nom.* ed. II. ii. 355.—Am. bor.
refractum, *Benth. ex Vasey, Grass. U. St.* 40.—Calif.
Sabinii, *R. Br. in Parry, 1st Voy. Suppl.* 289 (err. tpy. 189).—Reg. arct.

PLEUROPTERANTHA, Franch. in Revoil, Çomali, 59. t. 5 (1882). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 1218.
Revoili, *Franch. l. c.* 60. t. 5.—Afr. trop.

PLEUROPTERUS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 587 = **Polygonum**, Tourn.
cordatus, Turcz. l. c. = Polyg. multiflorum, *Thunb.*

PLEUROSPA, Rafin. Fl. Tellur. iv. 8 (1836) = **Anthurium**, Schott (Aroid.).
cordifolia, Rafin. l. c. 9 = A. cordifolia.
reticulata, Rafin. l. c. 8.—Hab. ?

PLEUROSPERMUM, Hoffm. Gen. Umbell. Praemonend. p. viii (1814). *UMBELLIFERAE*, Benth. & Hook. f. i. 915.
AULACOSPERMUM, Ledeb. Fl. Alt. iv. 334 (1833).
ENYMONOSPERMUM, Spreng. ex DC. Prod. iv. 244, in syn. (1830).
GRAFIA, Reichb. Handb. 219 (1837).
HLADNIKIA, Koch, in Flora, xix. (1836) 166; et Syn. Fl. Germ. ed. I. 320 (1836).
HYMENIDIUM, Lindl. in Royle, Illustr. Bot. Himal. 233 (1833).
HYMENOLAENA, DC. Prod. iv. 244 (1830).
MALABAILA, Tausch, in Flora, xvii. (1834) 356.
PHYSOSPERMUM, Cusson, in Mém. Soc. Méd. Par. (1782) 279.

PLEUROSPERMUM:—
PTEROCYCLUS, Klotzsch, in Bot. Ergeb. Waldem. Reise, t. 47 (1862).
angelicoides, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 703.—Reg. Himal.
apiolens, *C. B. Clarke, l. c.* 705.—Reg. Himal.
Archangelica, *Ledeb. Fl. Alt.* i. 369.—Sibir. altaic.
austriacum, *Hoffm. Gen. Umbell. Praemonend.* p. x.—Europ. or.
Benthami, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 703.—Reg. Himal.
Brunonis, *Benth. ex C. B. Clarke, l. c.* 706.—Reg. Himal.
camtschaticum, *Hoffm. Gen. Umbell. Praemonend.* p. x.—Kamtschat.
Candollii, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 703.—Reg. Himal.
cicutarium, D. Dietr. Syn. Pl. ii. 988 = Molopospermum cicutarium.
cicutarium, Lindl. in Royle, Illustr. Bot. Himal. 233 = Selinum tenuifolium.
corydalifolium, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 62.—Afghan.
densiflorum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 706.—Reg. Himal.
dentatum, *Benth. ex C. B. Clarke, l. c.* 704.—Reg. Himal.
Gmelini, Bong. Vég. Ins. Sitcha, 141 = P. Archangelica.
Gmelini, *Steud. Nom.* ed. II. ii. 355.—Sibir.
Golaka, *Reichb. Fl. Germ. Excurs.* 441.—Europ.
Govanianum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 702.—Reg. Himal.
Hookeri, *C. B. Clarke, l. c.* 705.—Reg. Himal.
kamtschaticum, Auct. = camtschaticum.
lithuanicum, *Downar, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) i. 180.—Europ.
pulchrum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 63.—Afghan.
pumilum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 704.—Reg. Himal.
rotundatum, *Benth. ex C. B. Clarke, l. c.* 703.—Reg. Himal.
sikkimense, *C. B. Clarke, l. c.* 702.—Reg. Himal.
stellatum, *Benth. ex C. B. Clarke, l. c.* 704.—Reg. Himal.
stylosum, *C. B. Clarke, l. c.*—Reg. Himal.
toluccense, D. Dietr. Syn. Pl. ii. 988 = Arracacia toluccensis.
turkestanicum, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 295.—Turkestan.
uralense, *Hoffm. Gen. Umbell. Praemonend.* p. ix.—Sibir. ural.
uralense, Turcz. in Bull. Soc. Nat. Mosc. (1838) i. 93; xvii. (1844) 753 = austriacum.

PLEUROSTACHYS, Brongn. in Duperr. Voy. Coq. Bot. 172. t. 31 (1829). *CYPERACEAE*.
NEMOCHLOA, Nees, in Meyen, Reise, i. 108 (1834).
NOMOCHLOA, Nees, in Linnaea, ix. (1834) 299; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 110.
Beyrichii, *Steud. Syn. Pl. Cyp.* 139.—Bras.
elegans, *Kunth, Enum, Pl.* ii. 285.—Bras.
extenuata, *Steud. Syn. Pl. Cyp.* 139.—Bras.
floribunda, Kunth, Enum. Pl. ii. 285 = tenuiflora.
foliosa, *Kunth, l. c.* 284.—Bras.
Gaudichaudii, *Brogn. in Duperr. Voy. Coq. Bot.* 174.—Bras.
graminifolia, *Brogn. l. c.* 175.—Bras.
grandifolia, Boeck. in Kjoeb. Vidensk. Meddel. (1879–80) 25 = millegrana.
macrantha, *Kunth, Enum. Pl.* ii. 286.—Bras.
Martiana, *Steud. Syn. Pl. Cyp.* 139.—Bras.
millegrana, *Steud. l. c.*—Bras.
Orbigniana, *Brongn. in Duperr. Voy. Coq. Bot.* 175.—Bras.
puberula, *Boeck. in Flora*, lxiii. (1880) 453.—Am. austr.
Sellowii, *Kunth, Enum. Pl.* ii. 285.—Bras.
sparsiflora, *Kunth, l. c.* 286.—Bras.
stricta, *Kunth, l. c.*—Bras.
tenuiflora, *Brongn. in Duperr. Voy. Coq. Bot.* 175.—Bras.
Urvillii, *Brongn. l. c.* 173. t. 31.—Bras.

PLEUROSTEMON, Rafin. in Journ. Phys. lxxxix. (1819) 258. Genus dubium.—Cf. Benth. & Hook. f. i. 789.
PLEURANDRA, Rafin. Fl. Ludov. 95 (1817).
album, Rafin. Adn. (1820) ex DC. Prod. iii. 64.—Am. bor.

PLEUROSTENA, Rafin. Fl. Tellur. iii. 12 (1836) = **Polygonum**, Tourn.
serotina, Rafin. l. c.; et New Fl. Am. iii. 41 = Polyg. serotinum.

PLEUROSTIGMA, Hochst. in Flora, xxiv. (1841) i. Intell. 42 = **Bouchea**, Cham. (Verben.).
subrotundum, Hochst. l. c. = B. marrubiifolia.
sulphureum, Hochst. l. c. = B. pterygocarpa.

PLEUROSTIMA, Rafin. Fl. Tellur. ii. 97 (1836) = **Barbacenia**, Vand. (Amaryll.).
purpurea, Rafin. l. c. 98 = B. purpurea.

PLEUROSTYLIA, Wight & Arn. Prod. 157 (1834). *CELASTRINEAE*, Benth. & Hook. f. i. 363.
BOOTTIA, Ayres, ex Baker, Fl. Maurit. 49 (1877).
Heynei, Wight & Arn. Prod. 157 = Wightii.
leucocarpa, *Baker, Fl. Maurit.* 49.—Ins. Maurit.
pachyphloea, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 104.—Madag.
Wightiana, Tul. l. c. 105, sphalm. = seq.
Wightii, *Wight & Arn. Prod.* 157.—Ind. or.

PLEUROSTYLIS, Walp. Rep. i. 536 (1842) = praec.

PLEUROTAENIA, Hohen. ex Benth. & Hook. f. Gen. i. 920 (1865) = **Peucedanum**, Tourn. (Umbell.).

PLEUROTHALLIS, R. Br. in Ait. Hort. Kew. ed. II. v. 211 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 488.
ACIANTHERA, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 292.
ACRONIA, Presl, Rel. Haenk. i. 104 (1827).
ANATHALLIS, Barb. Rodrig. Orch. Nov. 72 (1877).
CENTRANTHERA, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 293.
CHAETOCEPHALA, Barb. Rodrig. Orch. Nov. 37 (1877).
CROCODEILANTHE, Reichb. f. Xenia Orch. i. 10. t. 6 (1854).
CRYPTOPHORANTHUS, Barb. Rodrig. Orch. Nov. 79 (1877).
DUBOISIA, Karst. in Otto & Dietr. Allg. Gartenz. xv. (1847) 394.
DUBOIS-REYMONDIA, Karst. in Bot. Zeit. vi. (1848) 397.
HUMBOLDTIA, Ruiz & Pav. Prod. 121. t. 27 (1794).
MYOXANTHUS, Poepp. & Endl. Nov. Gen. et Sp. i. 50. t. 88 (1835).
RHYNCHOPERA, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Rar. 103. t. 41 (1841).
SPECKLINIA, Lindl. Gen. et Sp. Orch. 8 (1830).
TALPINARIA, Karst. Fl. Columb. i. 153. t. 76 (1858–61).
Acjanthera, *Lindl. Fol. Orch. Pleuroth.* 23.—Bras.
acuminata, H. C. Focke, in Tijdschr. Nat. Wetensch. ii. (1849) 195 = semperflorens.
acuminata, *Lindl. Bot. Reg.* (1842) *Misc.* 70.—Guian.
acutissima, *Lindl. Fol. Orch. Pleuroth.* 43.—Bras.
affinis, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Peruv.
agathophylla, *Reichb. f. Xenia, Orch.* iii. 25.—Bolivia.
alata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 17 = corniculata.
albida, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 327 = obovata.
alligatorifera, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 299.—Bras.
alpestris, *Lindl. Gen. et Sp. Orch.* 7.—Jamaic.
amoena, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 16.—Mexic.
angustifolia, *Lindl. Bot. Reg.* sub t. 1797.—Mexic.
antennifera, *Lindl. Fol. Orch. Pleuroth.* 30.—Peruv.
anthoctenium, *Reichb. f. in Linnaea*, xli. (1877) 94.—Ins. S. Doming.
anthoxantha, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Bras.* 149.—Bras.
aphthosa, *Lindl. Bot. Reg.* (1838) *Misc.* 42.—Mexic.
apiculata, *Lindl. Fol. Orch. Pleuroth.* 17.—Venezuela.
arachnantha, *Reichb. f. Otia Bot. Hamb.* 18.—Ecuador.
Arbuscula, *Lindl. Bot. Reg.* (1842) *Misc.* 72.—Peruv.
Archidonae, *Lindl. Fol. Orch. Pleuroth.* 15.—Peruv.
arcuata, *Lindl. l. c.* 34.—Bras.

PLEUROTHALLIS:—
aristata, *Hook. in Ann. Nat. Hist.* Ser. I. ii. (1839) 329. t. 15.—Guian.
aristulata, *Lindl. Fol. Orch. Pleuroth.* 41.—Bras.
articulata, *Lindl. Bot. Reg.* (1842) *Misc.* 83.—Bras.
aspasicensis, *Reichb. f. in Bonplandia*, iii. (1855) 73.—Venezuela.
asperilinguis, *Reichb. f. & Warsz. l. c.* ii. (1854) 114.—Peruv.
atropurpurea, *Lindl. Bot. Reg.* (1842) *Misc.* 81.—Jamaic.
aurantiaca, *Rodrig. Orch. Nov.* i. 10.—Bras.
aurea, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 397.—Venezuela.
aurea, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 16 = chrysantha.
auriculata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 356.—Bras.
auriculigera, Reichb. f. in Gard. Chron. (1871) 1579 = Masdevallia auriculigera.
aviceps, Reichb. f. l. c. = Masdevallia aviceps.
barbacenensis, *Rodrig. Orch. Nov.* ii. 11.—Bras.
barbata, H. C. Focke, in Bot. Zeit. xi. (1853) 227 = barbulata.
barbata, *Westcott, in Phytologist*, i. (1841) 54.—Bras.
Barberiana, *Reichb. f. in Gard. Chron.* (1881) II. 6.—Am. trop.
barbulata, *Lindl. Fol. Orch. Pleuroth.* 40.—Guian.
Beyrichii, *Reichb. f. in Linnaea*, xxii. (1849) 829.—Bras.
bicarinata, *Lindl. Bot. Reg.* (1839) *Misc.* 14.—Bras.
bicolor, Lindl. l. c. (1842) Misc. 76 = strupifolia.
bicornis, *Lindl. Fol. Orch. Pleuroth.* 7.—Peruv.
bicruris, *Lindl. l. c.* 8.—Peruv.
bidentata, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
bidentula, *Rodrig. Orch. Nov.* ii. 20.—Bras.
biflora, H. C. Focke, in Tijdschr. Nat. Wetensch. ii. (1849) 197 = orbicularis.
bilamellata, *Reichb. f. in Linnaea*, xxii. (1849) 830.—Mexic.
Binoti, *Regel, in Act. Hort. Petrop.* vii. (1880) 388.—Bras.
biserrula, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.
bistuberculata, *Rodrig. Orch. Nov.* ii. 35.—Bras.
bivalvis, *Lindl. Orch. Linden.* 2.—Venezuela.
blepharophylla, *Griseb. Cat. Pl. Cub.* 260.—Cuba.
bogotensis, Lindl. Orch. Linden. 1 = Mathewsii.
boliviana, *Reichb. f. in Bonplandia*, iii. (1855) 224.—Bolivia.
Bowmanni, *Reichb. f. in Gard. Chron.* (1869) 989.—Bras.
brachyglottis, *Reichb. f. in Flora*, xlviii. (1865) 275.—Ind. occ.
brachypetala, *Griseb. Cat. Pl. Cub.* 257.—Cuba.
breviflora, *Lindl. Bot. Reg.* (1841) *Misc.* 59.—Mexic.
brevipes, *H. C. Focke, in Tijdschr. Nat. Wetensch.* ii. (1849) 198.—Guian.
Bufonis, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 225.—Venezuela.
cabellensis, *Reichb. f. in Linnaea*, xxii. (1849) 832.—Mexic.
caespitosa, *Rodrig. Orch. Nov.* i. 14.—Bras.
calceolaris, *Reichb. f. in Linnaea*, xli. (1877) 15.—N. Granat.
campestris, *Rodrig. Orch. Nov.* i. 8.—Bras.
canaligera, *Reichb. f. in Linnaea*, xli. (1877) 15.—N. Granat.
Capanèmae, *Rodrig. Orch. Nov.* ii. 16.—Bras.
capensis, Lindl. Fol. Orch. Pleuroth. 44, sphalm. = *Polystachya capensis*.
capillaris, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
cardiantha, *Reichb. f. in Linnaea*, xli. (1877) 15.—N. Granat.
cardiostola, *Reichb. f. in Bonplandia*, ii. (1854) 26.—Venezuela.
cardiothallis, *Reichb. f. in Bot. Zeit.* xv. (1857) 158.—Nicaragua.
Cardium, *Reichb. f. in Bonplandia*, ii. (1854) 26.—Venezuela.
casapensis, *Lindl. Bot. Reg.* (1842) *Misc.* 76.—Peruv.
Cassidis, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 384.—N. Granat.
caulescens, *Lindl. in Hook. Journ. Bot.* i. (1834) 9.—Peruv.
cauliflora, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355; *Hook. Ic. Pl.* t. 50.—Peruv.
Centranthera, *Lindl. Fol. Orch. Pleuroth.* 37.—Bras.

PLEUROTHALLIS:—
ceratathallis, *Reichb. f. in Bonplandia*, ii. (1854) 25.—Venezuela.
chamaelepanthes, *Reichb. f. l. c.* iii. (1855) 240.—Peruv.
chamaestelis, Reichb. f. in Linnaea, xxii. (1849) 825 = Stelis florea.
chamensis, *Lindl. Orch. Linden.* 2.—Venezuela.
chitonoides, *Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 613, nomen.—Ins. Trinit.
chloidophylla, *Reichb. f. in Linnaea*, xxii. (1849) 830.—Bras.
chloroleuca, *Lindl. Orch. Linden.* 2.—N. Granat.
chrysantha, *Lindl. Fol. Orch. Pleuroth.* 14.—Mexic.
ciliata, *Knowl. & Westc. Flor. Cab.* i. (1837) 39.—Guian.
circumplexa, *Lindl. Bot. Reg.* (1838) *Misc.* 24.—Mexic.
clandestina, *Lindl. Fol. Orch. Pleuroth.* 43.—Venezuela.
clausa, *De Puydt, Les Orchidées* (1880), 32–33. f. 44–53.—Hab. ?
coccinea, Hook. Exot. Fl. ii. t. 129 = Rodriguezia secunda.
cochlearis, *Reichb. f. in Bot. Zeit.* x. (1852) 764.—Mexic.
compressiflora, *Rodrig. Orch. Nov.* ii. 13.—Bras.
conanthera, *Reichb. f. Xenia Orch.* iii. 41. t. 217.—Mexic.
congesta, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 17 = octomerioides.
convexifolia, *Rodrig. Orch. Nov.* i. 17.—Bras.
convoluta, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 107.—N. Granat.
cordata, *Lindl. Gen. et Sp. Orch.* 5.—Peruv.
cordata, Lindl. Fol. Orch. Pleuroth. 10 = coriacardia.
cordifolia, *Reichb. f. in Bonplandia*, ii. (1854) 26.—Venezuela.
coriacardia, *Reichb. f. l. c.*—Venezuela.
coriacea, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 116.—Porto Rico.
corniculata, *Lindl. in Bot. Reg.* (1842) *Misc.* 83.—Mexic.
crassa, *Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 613, nomen.—Mexic.
crassifolia, *H. C. Focke, in Tijdschr. Nat. Wetensch.* ii. (1849) 196.—Guian.
crassifolia, Reichb. f. in Linnaea, xxii. (1849) 832 = minutalis.
crassipes, *Lindl. Fol. Orch. Pleuroth.* 29.—Venezuela.
crenata, *Lindl. in Gard. Chron.* (1846) 207; *Fol. Orch. Pleuroth.* 38.—Mexic.
crepidophylla, *Reichb. f. Otia Bot. Hamb.* 18.—Ecuador.
crinita, Rodrig. Orch. Nov. i. 16 = Lepanthes crinita.
crocea, *Rodrig. l. c.* ii. 23.—Bras.
Crocodilanthe, *Lindl. Fol. Orch. Pleuroth.* 4.—Peruv.
crocodiliceps, *Reichb. f. in Bonplandia*, iii. (1855) 72.—Venezuela.
cryptantha, *Rodrig. Orch. Nov.* i. 13.—Bras.
cubensis, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 328.—Cuba.
cultellifolia, *Rodrig. Orch. Nov.* i. 19.—Bras.
cyclochila, *Lindl. Fol. Orch. Pleuroth.* 8.—N. Granat.
declivis, *Lindl. l. c.* 36.—Peruv.
decurrens, *Poepp. & Endl. Nov. Gen. et Sp.* i. 48.—Peruv.
decurva, *Lindl. Fol. Orch. Pleuroth.* 8.—Peruv.
delicatula, *Lindl. l. c.* 38.—Jamaic.
dendrophila, *Reichb. f. in Linnaea*, xxii. (1849) 827.—Venezuela.
dichroa, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 356.—Bras.
diffusa, *Poepp. & Endl. Nov. Gen. et Sp.* i. 49. t. 86.—Peruv.
Dinotherii, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 114.—Peruv.
diptera, *Lindl. Fol. Orch. Pleuroth.* 44.—Peruv.
discoidea, *Lindl. in Bot. Reg.* sub t. 1797.—Ins. Trinit.
disticha, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 49 = Oberonia brevifolia.
disticha, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 16.—Mexic.
dubia, *A. Rich. & Gal. l. c.*—Mexic.
dura, *Lindl. Fol. Orch. Pleuroth.* 32.—Peruv.
ecallosa, *Rodrig. Orch. Nov.* ii. 24.—Bras.
echinantha, *Rodrig. l. c.* 22.—Bras.

PLEUROTHALLIS:—
elachopus, *Reichb. f. in Gard. Chron.* (1884) I. 109.—Venezuela.
elegans, *Lindl. Bot. Reg.* (1842) *Misc.* 70.—N. Granat.
elongata, *Hort. Hamb. ex Klotzsch, in Ind. Sem. Hort. Berol.* (1853) *App.* 1.—Hab. ?
emarginata, *Lindl. Fol. Orch. Pleuroth.* 25.—Jamaic.
emarginata, Lindl. Gen. et Sp. Orch. 6 = Physosiphon emarginatus.
endotrachys, *Reichb. f. in Linnaea*, xli. (1877) 95.—Costa Rica.
ephemera, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 356.—Bras.
erinacea, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
exarticulata, *Rodrig. Orch. Nov.* ii. 27.—Bras.
exasperata, *Lindl. Fol. Orch. Pleuroth.* 15.—Venezuela.
expansa, *Lindl. l. c.* 4.—Peruv.
fallax, Reichb. f. in Bonplandia, iii. (1855) 224 = tribuloides.
Felis-lingua, *Rodrig. Orch. Nov.* ii. 18.—Bras.
fenestrata, *Rodrig. l. c.* 12.—Bras.
filifera, *Lindl. Fol. Orch. Pleuroth.* 20.—Peruv.
fimbriata, *Lindl. l. c.* 17.—Bras.
flavida, *Lindl. l. c.* 44, 45; *et Stelis*, 19.—Guian.
flexuosa, *Lindl. Bot. Reg.* (1842) *Misc.* 69.—Peruv.
floribunda, *Lindl. l. c.* 73.—Ins. Martinic.
floribunda, *Poepp. & Endl. Nov. Gen. et Sp.* i. 48. t. 84.—Peruv.
floripecten, *Reichb. f. in Bonplandia*, ii. (1854) 25.—N. Granat.
floripicta, *Lindl. in Mem. Am. Acad.* viii. (1861) 219, nomen.—Cuba.
floruleuta, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 223.—N. Granat.
Fockei, *Lindl. Fol. Orch. Pleuroth.* 23.—Guian.
foetens, *Lindl. Bot. Reg.* (1843) *Misc.* 5.—Bras.
foliata, *Griseb. Fl. Brit. W. Ind.* 610.—Jamaic.
foliosa, Hook. Bot. Mag. t. 2746 = Gomesa foliosa.
Fons-florum, *Lindl. Fol. Orch. Pleuroth.* 5.—Peruv.
foveata, *Lindl. l. c.* 31.—Peruv.
fragilis, *Lindl. Bot. Reg.* (1841) *Misc.* 91.—Bras.
fraterna, *Lindl. Fol. Orch. Pleuroth.* 3.—Bolivia.
Fritillaria, *Reichb. f. in Bonplandia*, iii. (1855) 240.—N. Granat.
Fuegi, *Reichb. f. Beitr. Orch. Centr. Am.* 97. t. 10.—Guatemala.
fulgens, *Reichb. f. in Gard. Chron.* (1875) II. 516.—Costa Rica.
fusca, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Bras.
galeata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 107.—N. Granat.
Gardneri, Lindl. Bot. Reg. (1842) Misc. 83 = Restrepia Gardneri.
gelida, *Lindl. l. c.* (1841) *Misc.* 91.—Jamaic.
geniculata, *Lindl. Fol. Orch. Pleuroth.* 15.—Peruv.
Ghiesbreghtiana, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 16 = quadrifida.
gigantea, *Lindl. Fol. Orch. Pleuroth.* 28.—Peruv.
glanduligera, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Bras.
glossopogon, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.
glumacea, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Bras.
gracilis, *Rodrig. Orch. Nov.* i. 6.—Bras.
gracillima, *Lindl. Fol. Orch. Pleuroth.* 35.—Peruv.
grandiflora, *Lindl. Bot. Reg.* sub t. 1797.—Peruv.
gratiosa, *Reichb. f. in Bonplandia*, ii. (1854) 25.—Venezuela.
Grobyi, *Lindl. Bot. Reg.* t. 1797.—Guiana.
hamosa, *Rodrig. Orch. Nov.* ii. 31.—Bras.
harpophylla, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
Hartwegii, Lindl. Bot. Reg. (1842) Misc. 80 = geniculata.
Hartwegii, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 106.—N. Granat.
hastulata, *Reichb. f. & Warm. in Otia Bot. Hamb.* 92.—Bras.
hemirhoda, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 133.—N. Granat.
hians, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
Hoffmannseggiana, *Reichb. f. in Linnaea*, xxii. (1849) 827.—Bras.

PLEUROTHALLIS:—
Hookeri, Regel, in Act. Hort. Petrop. vii. (1880) 545 = strupifolia.
hygrophila, *Rodrig. Orch. Nov.* i. 7.—Bras.
hymenantha, *Lindl. Bot. Reg.* sub t. 1797.—Bras.; Cuba.
hypnicola, *Lindl. l. c.* (1842) *Misc.* 75.—Bras.
Hystrix, *Reichb. f. in Bonplandia*, ii. (1854) 26.—Venezuela.
imbricata, Rodrig. Orch. Nov. i. 15 = Lepanthes imbricata.
immersa, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 224.—N. Granat.
Imrayi, *Lindl. Fol. Orch. Pleuroth.* 9 (Imraei).—Ins. Dominic.
inaequalis, *Lindl. Bot. Reg.* (1842) *Misc.* 76.—Bras.
incompta, *Reichb. f. in Bonplandia*, ii. (1854) 24.—Venezuela.
incurva, *Lindl. Fol. Orch. Pleuroth.* 30.—Peruv.
intricata, *Lindl. Orchid. Linden.* 1.—N. Granat.
ionantha, *Reichb. f. in Linnaea*, xxii. (1849) 830.—Venezuela.
jacarepaguaënsis, *Rodrig. Orch. Nov.* ii. 29.—Bras.
Jamiesoni, *Lindl. Bot. Reg.* sub t. 1797.—N. Granat.
janeirensis, *Rodrig. Orch. Nov.* ii. 29.—Bras.
johannensis, *Rodrig. l. c.* 23.—Bras.
Kefersteiniana, *Reichb. f. in Bot. Zeit.* x. (1852) 673.—Venezuela.
Kegelii, *Reichb. f. in Linnaea*, xli. (1877) 132.—Guian.
Klotzscheana, *Reichb. f. l. c.* xxii. (1849) 828.—Bras.
Krameriana, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 294.—Am. centr.
laciniata, Rodrig. Orch. Nov. i. 14 = Lepanthes laciniata.
laevigata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 106.—N. Granat.
lamellaris, *Lindl. Fol. Orch. Pleuroth.* 8.—Bolivia.
Lanceana, *Lodd. Bot. Cab.* t. 1767.—Guian.
Lanceola, *Spreng. Syst.* iii. 731.—Jamaic.
lanceolata, *Lindl. Gen. et Sp. Orch.* 5.—Peruv.
Lansbergii, Regel, Ind. Sem. Hort. Petrop. (1856) 17; in Ann. Sc. Nat. Sér. IV. vi. (1856) 373 = P. Cardium.
lateritia, *Reichb. f. in Gard. Chron.* (1872) 731.—Costa Rica.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* i. 364.—N. Granat.
laurifolia, Reichb. f. in Bonplandia, ii. (1854) 23 = Roezlii.
laxa, *Lindl. Gen. et Sp. Orch.* 7.—Jamaic.
leontoglossa, *Reichb. f. in Linnaea*, xxii. (1849) 831.—Bras.
lepanthiformis, Reichb. f. l. c. xviii. (1844) 398 = villosa.
leptotifolia, *Rodrig. Orch. Nov.* i. 15.—Bras.
leucopyramis, *Reichb. f. in Linnaea*, xli. (1877) 47.—Costa Rica?
lichenicola, *Griseb. Cat. Pl. Cub.* 259.—Cuba.
ligulata, *Lindl. Fol. Orch. Pleuroth.* 29.—Peruv.
lilacina, *Rodrig. Orch. Nov.* i. 18.—Bras.
Lindenii, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 397.—Venezuela.
linearis, *Lindl. Bot. Reg.* (1842) *Misc.* 82.—Mexic.
Lingua, *Lindl. l. c.* 80.—Bras.
linguifera, *Lindl. Fol. Orch. Pleuroth.* 10.—Bolivia.
liparanges, *Reichb. f. in Gard. Chron.* (1885) i. 532.—Bras.
listrostachys, Reichb. f. in Bonplandia, iii. (1855) 71 = stenopetala.
longicaulis, *Lindl. Bot. Reg.* (1842) *Misc.* 72.—Bras.
longilabris, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 328.—Cuba.
longipes, *Koern. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 281.—Venezuela.
longirostris, *H. C. Focke, in Bot. Zeit.* xi. (1853) 340.—Guian.
longissima, Lindl. Fol. Orch. Pleuroth. 31 = racemiflora.
loranthophylia, *Reichb. f. in Bot. Zeit.* x. (1852) 674.—Am. austr.
luctuosa, *Reichb. f. in Linnaea*, xli. (1877) 48.—Costa Rica.
luteola, Lindl. Bot. Reg. (1841) Misc. 1 = fragilis.
macra, *Lindl. Fol. Orch. Pleuroth.* 10.—Peruv.
macroblepharis, *Reichb. f. in Gard. Chron.* (1874) ii. 772.—Peruv.?
macrocardia, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.

PLEUROTHALLIS:—
macrophylla, *H. B. & K. Nov. Gen. et Sp.* i. 365.—N. Granat.
macrophyta, *Rodrig. Orch. Nov.* i. 19.—Bras.
macropoda, *Rodrig. l. c.* ii. 25.—Bras.
macrorhiza, *Lindl. in Hook. Journ. Bot.* i. (1834) 9.—Peruv.
macuconensis, *Rodrig. Orch. Nov.* ii. 17.—Bras.
Magdalenae, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
malachantha, *Reichb. f. l. c.* 223.—Bras.
Mandonii, *Reichb. f. Xenia Orch.* iii. 24.—Bolivia.
mantiguyrana, *Rodrig. Orch. Nov.* ii. 14.—Bras.
marginalis, *Reichb. f. in Bonplandia*, iii. (1855) 224.—Bras.
marginata, *Lindl. Bot. Reg.* (1838) *Misc.* 42.—Guatemal.
Mathewsii, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Peruv.; N. Granat.
melachyla, *Rodrig. Orch. Nov.* ii. 19.—Bras.
melanantha, *Reichb. f. in Flora*, xlviii. (1865) 275.—Cuba.
Memor, *Reichb. f. in Bonplandia*, iv. (1856) 330.—Hab.?
meridana, *Reichb. f. in Linnaea*, xxii. (1849) 826.—Venezuela.
mesophylla, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 16 = circumplexa.
micrantha, *Rodrig. Orch. Nov.* ii. 21.—Bras.
microcardia, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
microlepanthes, *Griseb. Fl. Brit. W. Ind.* 610.—Jamaic.
microphylla, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 17.—Mexic.
Miersii, Lindl. Bot. Reg. (1842) Misc. 84 = Restrepia Miersii.
minax, Reichb. f. in Bonplandia, ii. (1854) 24 = plumosa.
minutalis, *Lindl. Fol. Orch. Pleuroth.* 40.—Mexic.
Miqueliana, *Lindl. l. c.* 17.—Guian.
misera, *Lindl. l. c.* 36.—Peruv.
modestissima, *Reichb. f. & Warm. in Otia Bot. Hamb.* 93.—Bras.
monocardia, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
montana, *Rodrig. Orch. Nov.* i. 5.—Bras.
Moritzii, *Reichb. f. in Linnaea*, xxii. (1849) 824.—Venezuela.
moschata, *Reichb. f. Xenia Orch.* iii. 42. t. 217.—Costa Rica.
Moseni, *Reichb. f. Otia Bot. Hamb.* 93.—Bras.
mucronata, *Lindl. in Mem. Am. Acad.* N. S. viii. (1861) 219, nomen.—Cuba.
multicaulis, Poepp. & Endl. Nov. Gen. et Sp. i. 47. t. 82 = ruscifolia.
multicuspidata, *Reichb. f. in Linnaea*, xli. (1877) 117.—N. Granat.
multirostris, *Reichb. f. l. c.* 49.—Jamaic.
Murex, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
muscicola, Rodrig. Orch. Nov. i. 9 = Lepanthes muscicola.
muscifera, Lindl. Bot. Reg. (1842) Misc. 79 = Restrepia muscifera.
muscoidea, *Lindl. l. c.* (1838) *Misc.* 89.—Bras.
muscosa, Rodrig. Orch. Nov. i. 7 = Lepanthes muscosa.
myrticola, *Rodrig. l. c.* 9.—Bras.
naraniensis, *Reichb. f. Beitr. Orch. Centr. Am.* 96.—Costa Rica.
navicularis, *Lindl. Fol. Orch. Pleuroth.* 6.—N. Granat.
navilinguis, *Reichb. f. in Bonplandia*, iii. (1855) 72.—N. Granat.
nemorosa, *Rodrig. Orch. Nov.* i. 11.—Bras.
nicaraguensis, *Reichb. f. in Walp. Ann.* vi. 171.—Nicaragua.
nittiorhyncha, *Lindl. Fol. Orch. Pleuroth.* 20.—N. Granat.
nubigena, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 326 (*rubigena*) = corniculata.
nuda, Reichb. f. in Walp. Ann. vi. 184 = hemirhoda.
nummularia, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
oblongifolia, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Cuba.
obovata, *Lindl. Bot. Reg.* (1842) *Misc.* 75.—Peruv.
obscura, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 17 = bilamellata.

PLEUROTHALLIS:—
occulta, *Knowles & Westc. ex G. Don, in Sweet, Hort. Brit.* ed. III. 636, nomen.—Bras.
ochreata, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
octomeriaeformis, Reichb. f. in Bonplandia, ii. (1854) 25 = obovata.
octomerioides, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Mexic.
octophrys, *Reichb. f. in Linnaea*, xli. (1877) 95.—Bras.
odontotepala, *Reichb. f. in Flora*, xlviii. (1865) 275.—Cuba.
oligantha, *Rodrig. Orch. Nov.* ii. 33.—Bras.
ophicephala, *Rodrig. l. c.* 294.—Bras.
ophiocephala, Lindl. Bot. Reg. (1838) Misc. 34 = Restrepia ophiocephala.
orbicularis, *Lindl. l. c.* (1842) *Misc.* 79.—Guian.
ovalifolia, *Reichb. f. in Walp. Ann.* vi. 188.—Guian.
pachyglossa, *Lindl. Bot. Reg.* (1840) *Misc.* 68.—Mexic.
pachyphylla, *Reichb. f. in Linnaea*, xxii. (1849) 824.—Bras.
pachyrachis, A. Rich. Fl. Cub. Fanerog. ii. 234 = Bulbophyllum pachyrrhachis.
pachyrhyncha, A. Rich. ex Benth. in Journ. Linn. Soc. xviii. (1881) 299 = Bulbophyllum bracteolatum.
palmiformis, *Lindl. Fol. Orch. Pleuroth.* 3.—Peruv.
pandurifera, *Lindl. Bot. Reg.* (1842) *Misc.* 70.—Bras.
panduripetala, *Rodrig. Orch. Nov.* ii. 12.—Bras.
Pantasmi, *Reichb. f. in Bonplandia*, iii. (1855) 224.—Nicaragua.
papillosa, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
pardipes, *Reichb. f. in Saund. Refug. Bot.* t. 119.—Bras.
parvifolia, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Bras.
pavimentata, *Reichb. f. in Linnaea*, xli. (1877) 48.—Bras.
pectinata, *Lindl. Bot. Reg.* (1839) *Misc.* 1.—Bras.
pedunculatis, Hook. Journ. Bot. iii. (1841) 308 = aphthosa.
peduncularis, *Lindl. Bot. Reg.* (1843) *Misc.* 47.—Bras.
pedunculata, *Reichb. f. in Linnaea*, xxii. (1849) 822.—Venezuela.
pelioxantha, *Rodrig. Orch. Nov.* i. 17.—Bras,
perplexa, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 15.—Hab.?
perpusilla, *Reichb. f. in Seem. Bot. Voy. Herald*, 215.—Panama.
phalangifera, Reichb. f. in Walp. Ann. vi. 168 = Mathewsii.
phyllocardia, *Reichb. f. Beitr. Orch. Centr. Am.* 97.—Costa Rica.
Pichinchae, *Reichb. f. in Linnaea*, xli. (1877) 47.—Ecuador.
picta, Hook. Bot. Mag. t. 3897 = strupifolia.
picta, *Lindl. Bot. Reg.* sub t. 1797.—Guian.
pilifera, *Lindl. Fol. Orch. Pleuroth.* 9.—Peruv.
pisifera, *Lindl. l. c.* 36.—Peruv.
plantaginea, *Lindl. Bot. Reg.* (1842) *Misc.* 82.—Peruv.
plumosa, *Lindl. l. c.* 72.—Costa Rica.
Poeppigii, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Peruv.
polygonoides, *Griseb. Fl. Brit. W. Ind.* 609.—Ind. occ.
polyliria, *Endres & Reichb. f. in Gard. Chron.* (1871) 1483.—Costa Rica.
polystachya, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 16.—Mexic.
pristeoglossa, *Reichb. f. & Warm. Otia Bot. Hamb.* 92.—Bras.
procumbens, *Lindl. Fol. Orch. Pleuroth.* 35.—Venezuela.
prolifera, *Herb. ex Lindl. Bot. Reg.* t. 1298.—Bras.
prostrata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 327.—Cuba.
pruinosa, *Lindl. Bot. Reg.* (1842) *Misc.* 75.—Guian.
puberula, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxii. (1854) 393 = Restrepia ophiocephala.
pubescens, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Mexic.
pulchella, *Lindl. in Hook. Exot. Fl.* ii. sub t. 123.—Ecuador.
punctata, Ker-Gawl. in Bot. Reg. t. 759 = Notylia punctata.
punctata, *Lindl. l. c.* sub t. 1797.—Bras.

PLEUROTHALLIS:—
punctata, Rodrig. Orch. Nov. i. 11 = serpentula.
purpurea, D. Don, Prod. Fl. Nep. 33 = Bulbophyllum Careyanum.
pusilla, *Lindl. Bot. Reg.* (1842) *Misc.* 82.—Peruv.
pyrsodes, *Reichb. f. in Gard. Chron.* (1876) II. 386.—Am. centr.
quadrifida, *Lindl. Bot. Reg.* (1842) *Misc.* 70.—Mexic.
quitensis, *Reichb. f. in Bonplandia*, iii. (1855) 240.—Ecuador.
racemiflora, Lindl. Fol. Orch. Pleuroth. 34 = oblongifolia.
racemiflora, *Lindl. in Hook. Exot. Fl.* ii. t. 123.—Am. centr.; Jamaic.
ramosa, *Rodrig. Orch. Nov.* ii. 17.—Bras.
ramulosa, *Lindl. Fol. Orch. Pleuroth.* 33.—Peruv.
Raymondi, *Reichb. f. in Bonplandia*, ii. (1854) 26.—Venezuela.
recurva, *Lindl. Bot. Reg.* (1841) *Misc.* 1.—Bras.
restrepioides, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 356.—Peruv.
retusa, *Lindl. Bot. Reg.* (1842) *Misc.* 81.—Mexic.
Reymondi, Reichb. f. in Walp. Ann. iii. 520 = Raymondi.
rhodotantha, *Reichb. f. in Linnaea*, xxii. (1849) 825.—N. Granat.
rhomboglossa, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
rictoria, *Reichb. f. in Linnaea*, xli. (1877) 14.—N. Granat.
rigidifolia, *Reichb. f. in Bonplandia*, iii. (1855) 224.—Peruv.
rio-grandensis, *Rodrig. Orch. Nov.* ii. 28.—Bras.
Roezlii, *Reichb. f. in Linnaea*, xli. (1877) 13.—N. Granat.
roseo-punctata, *Lindl. Orch. Linden.* 2.—Venezuela.
rostellata, *Rodrig. Orch. Nov.* i. 12.—Bras.
rostriflora, *Reichb. f. in Linnaea*, xxii. (1849) 827.—Bras.
rubens, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
ruberrima, *Lindl. Orch. Linden.* 1.—Venezuela.
rubigena, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 326, sphalm. = *nubigena*.
rubroviridis, *Lindl. l. c.* 327.—Cuba.
rufobrunnea, *Lindl. Fol. Orch. Pleuroth.* 36.—Mexic.
rupestris, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
ruscifolia, *R. Br. in Ait. Hort. Kew.* ed. II. v. 211.—Am. trop.
sagittata, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Reise*, 150.—Bras.
sagittifera, H. B. & K. Nov. Gen. et Sp. i. 365. t. 91 = Notylia multiflora, *Lindl.*
saltatoria, *Lindl. Fol. Orch. Pleuroth.* 30.—Bolivia.
sarcophylla, *Reichb. f. in Bonplandia*, iii. (1855) 224.—Venezuela.
saurocephala, *Lodd. Bot. Cab.* t. 1571.—Bras.
scabrata, *Lindl. Fol. Orch. Pleuroth.* 30.—Bolivia.
scabrilinguis, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Peruv.
scabripes, *Lindl. Bot. Reg.* (1839) *Misc.* 94.—Bras.
scalaris, Griseb. Cat. Pl. Cub. 259 = nummularia.
Scapha, *Reichb. f. in Gard. Chron.* (1874) II. 162.—Hab.?
scariosa, *Lindl. Bot. Reg.* (1842) *Misc.* 71.—Mexic.
Schiedei, *Reichb. f. in Linnaea*, xxii. (1849) 826.—Mexic.
Sclarea, *Reichb. f. l. c.* xli. (1877) 49.—N. Granat.
sclerophylla, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
secunda, *Poepp. & Endl. Nov. Gen. et Sp.* i. 49.—Peruv.
segoviensis, *Reichb. f. in Bonplandia*, iii. (1855) 223.—Nicaragua.
semipellucida, *Reichb. f. in Linnaea*, xxii. (1849) 823.—Venezuela.
semiscabra, *Lindl. Orch. Linden.* 2.—Venezuela.
semperflorens, *Lindl. Fol. Orch. Pleuroth.* 40.—Guian.
seriata, *Lindl. Bot. Reg.* (1840) *Misc.* 75.—Bras.
serpentula, *Rodrig. Orch. Nov.* ii. 20.—Bras.
serrifera, *Lindl. Fol. Orch. Pleuroth.* 34.—Bras.
serrulatipetala, *Rodrig. Orch. Nov.* ii. 33.—Bras.
sertularioides, *Spreng. Syst.* iii. 731.—Jamaic.
setigera, *Lindl. Fol. Orch. Pleuroth.* 38.—Peruv.
sicaria, *Lindl. Bot. Reg.* (1841) *Misc.* 91.—Am. trop.
Sirene, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.
Smithiana, *Lindl. Bot. Reg.* (1843) *Misc.* 57.—Bras.
Sonderana, *Reichb. f. in Linnaea*, xxii. (1849) 830.—Bras.

PLEUROTHALLIS:—
soratana, *Reichb. f. Xenia Orch.* iii. 25.—Bolivia.
spathulata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 17 = tribuloides.
spectrilinguis, *Reichb. f. in Gard. Chron.* (1882) II. 457.—Hab.?
sphenochila, *Lindl. Fol. Orch. Pleuroth.* 22.—N. Granat.
spiculifera, *Lindl. l. c.* 43.—Bras.
spilantha, *Rodrig. Orch. Nov.* ii. 32.—Bras.
spiralis, *Lindl. Gen. et Sp. Orch.* 7.—Peruv.
Sprucei, *Lindl. Fol. Orch. Pleuroth.* 35.—Bras.
stenopetala, *Lodd. ex Lindl. Bot. Reg.* (1838) *Misc.* 95.—Am. trop.
stenostachya, *Reichb. f. in Linnaea*, xviii. (1844) 399.—Mexic.
stigmatoglossa, Reichb. f. ex Lindl. Fol. Orch. Pleuroth. 18 = Restrepia ophiocephala.
striata, *H. C. Focke, in Tijdschr. Nat. Wetensch.* iv. (1851) 63.—Guian.
strupifolia, *Lindl. Bot. Reg.* (1839) *Misc.* 2.—Bras.
subpellucida, Klotzsch, Ind. Sem. Hort. Berol. (1852) App. 1; et in Ann. Sc. Nat. Sér. III. xix. (1853) 375 = loranthophylla.
subsinuata, *Lindl. Fol. Orch. Pleuroth.* 37.—Hab.?
succosa, Lindl. Gen. et Sp. Orch. 5 = ruscifolia.
sulphurea, *Rodrig. Orch. Nov.* ii. 15.—Bras.
supervacanea, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 180.—Hab.?
surinamensis, H. C. Focke, in Tijdschr. Nat. Wetensch. ii. (1849) 194 = picta.
tentaculata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 354.—Peruv.
tenuissima, *Reichb. f. in Linnaea*, xviii. (1844) 399.—Mexic.
teres, *Lindl. Bot. Reg.* sub t. 1797.—Bras.
testaefolia, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 328.—Ind. occ.; Venezuela.
translucida, *Rodrig. Orch. Nov.* ii. 19.—Bras.
triangularis, Klotzsch & Karst. in Otto & Dietr. Allg. Gartenz. xv. (1847) 329 = chamensis.
tribuloides, *Lindl. Gen. et Sp. Orch.* 6.—Ind. occ.
tricarinata, H. C. Focke, in Bot. Zeit. xi. (1853) 889 = Fockei.
tricarinata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 49. t. 87.—Peruv.
trichophora, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 326.—Cuba.
trichopoda, Reichb. f. in Bonplandia, iv. (1856) 330 = subsinuata.
trichopoda, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 17.—Mexic.
trichyphis, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
tridentata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 289.—Venezuela.
trifida, *Lindl. Bot. Reg.* (1842) *Misc.* 82.—Bras.
trigonopoda, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 361 = sicaria.
trilineata, *Lindl. Fol. Orch. Pleuroth.* 29.—Peruv.
trilineata, Rodrig. Orch. Nov. i. 6 = Lepanthes trilineata.
tripterantha, *Reichb. f. in Bonplandia*, ii. (1854) 24.—Venezuela.
tripteris, Reichb. f. in Linnaea, xxii. (1849) 829 = sicaria.
tripterygia, *Reichb. f. in Bonplandia*, ii. (1854) 24.—Venezuela.
triquetra, Klotzsch, ex Lindl. Fol. Orch. Pleuroth. 13 = chamensis.
tristis, *Rodrig. Orch. Nov.* ii. 293.—Bras.
Trulla, *Reichb. & Warsz. in Bonplandia*, ii. (1854) 114.—Peruv.
truncata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Ecuador.
truncicola, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Reise*, 149.—Bras.
truxillensis, *Reichb. f. in Bonplandia*, ii. (1854) 25.—Venezuela.
tubata, Lodd. ex Steud. Nom. ed. II. ii. 356 = Physosiphon Loddigesii.
tuberculata, Wright, in Sauv. Fl. Cub. 228 = Epidendrum Leochilus.
tubulosa, *Lindl. Fol. Orch. Pleuroth.* 19.—Peruv.
ujarensis, Lindl. l. c. = Restrepia ujarensis.
undulata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 48.—Peruv.
uniflora, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 355.—Bras.

PLEUROTHALLIS:—
univaginata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 326.—Ind. occ.
Urbaniana, *Reichb. f. in Ber. Deutsch. Bot. Ges.* iii. (1885) 279.—Porto Rico.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 234.—Cuba.
variegata, Rodrig. Orch. Nov. i. 10 = Lepanthes variegata.
velaticaulis, *Reichb. f. in Linnaea*, xxii. (1849) 824.—Venezuela.
velatipes, *Reichb. f. l. c.* 828.—Venezuela.
ventricosa, *Lindl. Fol. Orch. Pleuroth.* 16.—Peruv.
verrucosa, *Reichb. f. in Bonplandia*, ii. (1854) 24.—Venezuela.
vilipensa, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 3.—Am. centr.
villosa, *Knowles & Westc. Flor. Cab.* ii. (1838) 78.—Mexic.
violacea, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 16.—Mexic.
viridula, *Lindl. Fol. Orch. Pleuroth.* 19.—Venezuela.
vittata, *Lindl. Bot. Reg.* (1838) *Misc.* 73.—Mexic.; Venezuela.
Wageneriana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 274 = vittata.
Warmingii, *Reichb. f. Otia Bot. Hamb.* 93.—Bras.
Wawraeana, *Rodrig. Orch. Nov.* ii. 14.—Bras.
Weddelliana, Reichb. f. Xenia Orch. iii. 24.—Bolivia.
Wilsoni, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 326.—Ind. occ.
Wrightii, *Reichb. f. in Flora*, xlviii. (1865) 276.—Ind. occ.
xanthochlora, *Reichb. f. in Linnaea*, xxii. (1849) 823.—N. Granat.
xiphizusa, Reichb. f. in Walp. Ann. vi. 172 = P. Crocodilanthe.
xiphochila, *Reichb. f. in Linnaea*, xxii. (1849) 831.—Venezuela.
zephyrina, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.

PLEUROTHYRIUM, Nees, Syst. Laurin. 349 (1836). *LAURINEAE*, Benth. & Hook. f. iii. 159.
bifidum, *Nees, l. c.* 351.—Peruv.
chrysophyllum, *Nees, l. c.*—Peruv.
Chrysothyrsus, Meissn. in DC. Prod. xv. 1. 169 = Nectandra Arnottiana.
cuneifolium, *Nees, Syst. Laurin.* 352.—Bras.; Peruv.
ferrugineum, Meissn. in DC. Prod. xv. 1. 169 (= *Persea ferruginea*, Benth. & Hook. f.).—Peruv.
macranthum, Nees, in Linnaea, xxi. (1848) 512 = cuneifolium.
Poeppigii, *Nees, Syst. Laurin.* 349.—Peruv.
velutinum, Meissn. in DC. Prod. xv. 1. 170 (= *Persea velutina*).—N. Granat.

PLEUTERON, Rafin. Sylva Tellur. 109 (1838) = **Capparis**, Tourn.
Baduca, Rafin. l. c. = C. zeylanica.
Breynia, Rafin. l. c. = C. Breynia.
comosa, Rafin. l. c. = C. comosa.
frondosa, Rafin. l. c. = C. frondosa.
hastata, Rafin. l. c. = C. Cynophallophora.
linearis, Rafin. l. c. = C. linearis.
siliquosa, Rafin. l. c. = C. Breynia.
tenuis, Rafin. l. c. = C. tenuisiliqua.
torulosa, Rafin. l. c. = C. jamaicensis.

PLEXAURE, Endl. Prod. Fl. Norf. 30 (1833) = **Phreatia**, Lindl. (Orchid.).
cladophylax, Reichb. f. in Linnaea, xix. (1847) 370.—Ins. Carolin.
Dendrophylax, Reichb. f. l. c. 371.—Ins. Carolin.
limenophylax, Endl. Prod. Fl. Norf. 30 = Phreatia limenophylax.

PLEXINIUM, Rafin. Fl. Tellur. ii. 32 (1836) = **Androcymbium**, Willd. (Liliac.).
punctatum, Rafin. l. c. = A. leucanthum.

PLEXIPUS, Rafin. Fl. Tellur. ii. 104 (1836) = **Bouchea**, Cham. (Verben.).
cuneifolius, Rafin. l. c. = B. cuneifolia.

PLEXISTENA, Rafin. Fl. Tellur. ii. 20 (1836) = **Allium**, Tourn. (Liliac.).
margaritea, Rafin. l. c. = A. margaritaceum.

PLICULA, Rafin. Sylva Tellur. 55 (1838) = **Acnistus**, Schott (Solan.).
umbellata, Rafin. l. c. = A. umbellatus.

PLINIA, Blanco, Fl. Filip. ed. I. 423 (1837) = **Kayea**, Wall. (Guttif.).
paniculata, Blanco, l. c. = K. racemosa.

PLINIA, Plum. ex Linn. Syst. ed. I (1735) = **Eugenia**, Mich. (Myrtac.).
crocea, Linn. Mant. ii. 244, in obs.—Ind. occ.; Bras.
pedunculata, Linn. f. Suppl. 253 = E. uniflora, *Linn.*
pennata, Linn. Sp. Pl. 516 = *crocea.*
pentapetala, Linn. Mant. ii. 402 = *crocea.*
petiolata, Linn. Amoen. Acad. viii. 257 = E. uniflora, *Linn.*
rubra, Linn. Mant. ii. 243 = E. uniflora, *Linn.*
tetrapetala, Linn. l. c. 402 = E. uniflora, *Linn.*?

PLINTHANTHESIS, Steud. Syn. Pl. Gram. 14 (1855) = **Danthonia**, DC.? (Gramin.).
tenuior, Steud. l. c. 14.—Austral.
Urvillei, Steud. l. c.—Austral.

PLINTHINE, Reichb. Handb. 298 (1837) = **Arenaria**, Linn. (Caryoph.).
aggregata, Reichb. Ic. Fl. Germ. v. 33. t. 219 = A. tetraquetra.
tetraquetra, Reichb. l. c. = A. tetraquetra.

PLINTHOCROMA, Dulac, Fl. Hautes-Pyr. 419 (1867) = **Rhododendron**, Linn. (Ericac.).
ferrugineum, Dulac, l. c. = R. ferrugineum.

PLINTHUS, Fenzl, in Endl. Nov. Stirp. Dec. 51 (1839). *FICOIDEAE*, Benth. & Hook. f. i. 855.
cryptocarpus, *Fenzl, l. c.* 52.—Afr. austr.

PLOCAGLOTTIS, Steud. Nom. ed. II. ii. 356 (1841) = **Plocoglottis**, Blume (Orchid.).

PLOCAMA, Ait. Hort. Kew. ed. I. i. 292 (1789); iii. 508. *RUBIACEAE*, Benth. & Hook. f. ii. 136.
BARTLINGIA, Reichb. in Flora, vii. (1824) 241.
PLACODIUM, Benth. & Hook. f. Gen. ii. 136 (1873), sphalm.
PLACOMA, J. F. Gmel. Syst. 296, 390 (1791).
pendula, *Ait. Hort. Kew.* ed. I. i. 392.—Ins. Canar.

PLOCANDRA, E. Mey. Comm. Pl. Afr. Austr. 181 (1837) = **Chironia**, Linn. (Gentian.).
albens, E. Mey. l. c. 182.—Afr. austr.
palustris, Griseb. in DC. Prod. ix. 43 = C. palustris.
purpurascens, E. Mey. Comm. Pl. Afr. Austr. 182 = C. purpurascens (in Addend.).

PLOCOGLOTTIS, Blume, Bijdr. 380 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 541.
acuminata, *Blume, Mus. Bot. Lugd. Bat.* i. 46.—Malaya.
dilatata, *Blume, l. c.* 47.—Borneo.
fimbriata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 493.—Java.
javanica, *Blume, Bijdr.* 381.—Java.
latifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 47.—Java.
Lowii, *Reichb. f. in Gard. Chron.* (1865) 434.—Borneo.
moluccana, *Blume, Mus. Bot. Lugd. Bat.* i. 47.—Ins. Amboin.

PLOCOSPERMA, Benth. in Benth. & Hook. f. Gen. ii. 789 (1876). *LOGANIACEAE*, Benth. & Hook. f. l. c.
buxifolium, *Benth. in Hook. Ic. Pl.* t. 1195.—Guatemala.

PLOCOSTEMMA, Blume, Rumphia, iv. 30 (1848); et Mus. Bot. Lugd. Bat. i. 59. t. 14 (1849) = **Hoya**, R. Br. (Asclep.).
lasianthum, Blume, Rumphia, iv. 30 = H. lasiantha.
pallidum, Blume, Mus. Bot. Lugd. Bat. i. 60.—Sumatra.

PLOCOSTIGMA, Benth. err. typ. in Journ. Linn. Soc. xviii. (1881) 338, loco PLACOSTIGMA, Blume = **Podochilus**, Blume (Orchid.).

PLOESSLIA, Endl. Nov. Stirp. Dec. 39 (1839) = **Boswellia**, Roxb. (Burser.).
floribunda, Endl. l. c. 40 = B. papyrifera.

PLOIARUM, Korth. Verh. Nat. Gesch. Bot. 135. t. 25 (1842) = **Archytaea**, Mart. (Ternstr.).
elegans, Korth. l. c. = A. Vahlii.
oblongifolium, Miq. Fl. Ind. Bat. Suppl. 483.—Sumatra.

PLOKIOSTIGMA, Schuch. in Linnaea, xxvi. (1853) 39 = **Stackhousia**, Sm.
Lehmanni, Schuch. l. c. 40 = S. pubescens?

PLOSTAXIS, Rafin. New Fl. Am. iv. 87 (1836) = **Polygala**, Tourn.
lutea, Rafin. l. c. = Polyg. lutea.

PLOTEA, J. F. Gmel. Syst. 404 (1791) = seq.

PLOTIA, Adans. Fam. ii. 226 (1763) = **Myrsine**, Linn.
Adansonii, J. F. Gmel. Syst. 404.—Hab.?

PLOTIA, Neck. Elem. ii. 55 (1790) = **Embelia**, Burm. f. (Myrsin.).

PLOTIA, Schreb. ex Steud. Nom. ed. II. ii. 356 (1841) = **Poa**, Linn. (Gramin.).
brizoides, Schreb. ex Steud. l. c. = Poa obtusa.

PLOTTZIA, Arn. in Lindl. Introd. Nat. Syst. ed. II. 441 (1836) = **Paronychia**, Juss.
dichotoma, Arn. l. c. = Paron. dichotoma.

PLUCHEA, Cass. in Bull. Soc. Philom. (1817) 31; et in Dict. Sc. Nat. xlii. 1. (1826) *COMPOSITAE*, Benth. & Hook. f. ii. 290.
BERTHELOTIA, DC. Prod. v. 375 (1836).
EYREA, F. Muell. in Linnaea, xxv. (1852) 403.
GYMNOSTYLIS, Rafin. in Am. Monthly Mag. (1818) 268.
GYNEMA, Rafin. Fl. Ludov. 63 (1817).
KARELINIA, Less. in Linnaea, ix. (1834) 187; et in DC. Prod. v. 375 (1836).
LEPTOGYNE, Ell. Sketch, ii. 322 (1824).
PLUECHEA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 248.
SPIROPODIUM, F. Muell. Fragm. i. 33 (1858).
STYLIMNUS, Rafin. in Journ. Phys. lxxxix. (1819) 100.
TECMARSIS, DC. Prod. v. 93 (1836).
alopecuroides, Less. Syn. Comp. 207 = Pterocaulon virgatum.
amabilis, *Kunth, Ind. Sem. Hort. Berol.* (1845) 12.—Venezuela.
appendiculata, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 249.—Malaya.
arguta, *Boiss. Diagn.* Ser. II. iii. 5.—Ind. bor. occ.
aromatica, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 838.—Ins. Socotra.
auriculata, *Hemsl. Diagn. Pl. Nov.* 32.—Mexic.
baccharoides, *F. Muell. ex Benth. Fl. Austral.* iii. 528.—Austral.
balsamifera, Less. in Linnaea, vi. (1831) 150 = Blumea balsamifera.
basiflora, F. Muell. Rep. Babb. Exped. 12 = Thespidium basiflorum.
bifrons, *DC. Prod.* v. 451.—Am. bor.; Ind. occ.
bipinnata, Baill. Hist. Pl. viii. 300 = Garuleum bipinnatum.
Boivini, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 369.—Ins. Borbon.
borealis, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 212.—Am. bor. occ.
Bravae, *C. Bolle, in Bonplandia*, viii. (1860) 130.—Ins. Canar.
camphorata, *DC. Prod.* v. 452.—Am. bor.; Ind. occ.
carolinensis, G. Don, in Sweet, Hort. Brit. ed. III. 350 = odorata.
Chingoyo, *DC. Prod.* v. 452.—Peruv.
Cortesii, *DC. l. c.*—Mexic.
crassifolia, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 394.—Abyss.
crinita, Hook. & Arn. ex Baker, in Mart. Fl. Bras. vi. III. 105 = Stenachaenium megapotamicum.
dentex, *R. Br. ex Benth. Fl. Austral.* iii. 529.—Austral.

PLUCHEA:—
Dioscorides, *DC. Prod.* v. 450.—Afr. trop.
domingensis, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 327.—Ins. S. Doming.
Doniana, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 191.—Burma.
eupatorioides, *Kurz, For. Fl. Brit. Burma*, ii. 575.—Siam.
Eyrea, *F. Muell. Rep. Babb. Exp.* 11.—Austral.
fastigiata, *Griseb. in Goett. Abh.* xxiv. (1879) 184.—Reg. Argent.
filifolia, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 56 = Pterigeron filifolius.
floribunda, Hemsl. Diagn. Pl. Nov. 32 = Baccharis mucronata.
foetida, DC. Prod. v. 452 = camphorata.
foliolosa, DC. l. c. 451 = indica.
frutescens, Benth. in Hook. Ic. Pl. t. 1157 = arguta.
glabra, *Griseb. in Goett. Abh.* xxiv. (1879) 183.—Reg. Argent.
glabrata, DC. Prod. v. 452 = camphorata?
glutinosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 838.—Ins. Socotra.
heterophylla, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 325.—Afr. trop.
hirsuta, *Less. in Linnaea*, vi. (1831) 150.—China.
indica, *Less. l. c.*—As. et Austral. trop.
Kotschyi, Sch. Bip. in Flora, xxv. (1842) I. 134, nomen; et ex A. Rich. Tent. Fl. Abyss. i. 393 = Laggera aurita.
Kraussii, Sch. Bip. ex Steetz, in Peters, Reise Mossamb. Bot. 442 = P. Dioscoridis.
lanceolata, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 329.—Ind. bor. occ.; Afr. bor.
lanuginosa, Hook. f. Fl. Brit. Ind. iii. 266 = Blumea Malcolmii.
laxiflora, *Hook. & Arn. ex Baker, in Mart. Fl. Bras.* vi. III. 107.—Bras.
ligulata, F. Muell. Rep. Babb. Exped. 12 = Pterigeron liatroides.
linearifolia, *C. B. Clarke, Comp. Ind.* 95.—Reg. Himal.
longepedunculata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 12.—Venezuela.
macrocephala, DC. Prod. v. 450 = Stenachaenium megapotamicum.
macrocephala, F. Muell. Rep. Babb. Exped. 12 = Pterigeron macrocephalus.
macrophylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 107 = laxiflora.
marylandica, Cass. in Dict. Sc. Nat. xlii. 2 = camphorata.
montana, *Griseb. in Goett. Abh.* xxiv. (1879) 184.—Reg. Argent.
oblongifolia, *DC. Prod.* v. 451.—Bras.
obovata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 838.—Ins. Socotra.
odora, F. Muell. Rep. Babb. Exped. 12 = Pterigeron odorus.
odorata, *Cass. in Dict. Sc. Nat.* xlii. 3.—Am. austr.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 125 = oblongifolia.
ovalis, *DC. Prod.* v. 450.—Ind. bor. occ.; Afr. trop.
petiolata, Cass. in Dict. Sc. Nat. xlii. 2 = camphorata.
phagnaloides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 393 = Phagnalon scoparium.
pinnatifida, *Hook. f. in Hook. Ic. Pl.* t. 1156.—Afr. trop.
purpurascens, *DC. Prod.* v. 452.—Jamaic.
purpurascens, Torr. & Gray, Fl. N. Am. ii. 261 = camphorata.
pycnostachya, Less. Syn. Comp. 207 = Pterocaulon pycnostachyum.
Quitoc, *DC. Prod.* v. 450.—Bras.
rugosa, Less. Syn. Comp. 207 = Pterocaulon virgatum.
sagittalis, Less. l. c. = Baccharis sagittalis.
scabrida, *DC. Prod.* v. 453.—Ins. Philipp.
senegalensis, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 368.—Afr. trop.
Serra, *Franch. in Révoil, Çomali*, 35.—Afr. trop.
sessifolia, *DC. Prod.* v. 395.—Java.
sessilifolia, Steud. Nom. ed. II. ii. 356, sphalm. = praec.
sordida, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 328.—Afr. trop.
spicata, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 369.—Ins. Borbon.
spicata, Less. Syn. Comp. 207 = Pterocaulon virgatum.
squarrosa, *Benth. Fl. Austral.* iii. 529.—Austral.

PLUCHEA:—
subdecurrens, Benth. Pl. Hartw. 17 = auriculata.
subdecurrens, *Cass. in Dict. Sc. Nat.* xlii. 4.—Mexic.
subumbellata, Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 369 = Conyza Heudelotii.
tetranthera, *F. Muell. Rep. Babb. Exped.* 12.—Austral.
tomentosa, *DC. in Wight, Contrib.* 16.—Ind. or.
Tweediei, Hook. & Arn. ex Baker, in Mart. Fl. Bras. vi. III. 106 = P. Quitoc.
verrucosa, *Klatt, in Decken, Reise Ost-Afr.* iii. 76. —Afr. trop.
virgata, Less. in Linnaea, vi. (1831) 726 = Pterocaulon virgatum.
Wallichiana, *DC. Prod.* v. 451.—Ind. bor. occ.

PLUCHIA, Vell. Fl. Flum. 157 (1825); iv. t. 20 (1827) = **Diclidanthera**, Mart. (Styrac.).
curiosa, Vell. ll. cc.—Bras.

PLUECHEA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 248 = **Pluchea**, Cass. (Compos.).

PLUKENETIA, Linn. Sp. Pl. ed. I. 1192 (1753).
EUPHORBIACEAE, Benth. & Hook. f. iii. 327.
ACCIA, A. St. Hil. Morphol. 499 (1840).
ANABAENA, A. Juss. Euph. Tent. 46. t. 15 (1824).
BOTRYANTHE, Klotzsch, in Wiegm. Achiv, vii. (1841) 190. t. 9.
CERATOCOCCUS, Meissn. Gen. Comm. 369 (1843).
FRAGARIOPSIS, A. St. Hil. Morphol. 426, 499 (1840).
HEDRAIOSTYLUS, Hassk. Cat. Hort. Bog. Alt. 234 (1844).
PLUKNETIA, Linn. Gen. ed. I. 373 (1737).
PTEROCOCCUS, Hassk. in Flora, xxv. (1842) II. Beibl. 41.
SAJORIUM, Endl. Gen. Suppl. iii. 98 (1843).
VIGIA, Vell. Fl. Flum. Ic. ix. t. 128 (1827).
africana, *Sond. in Linnaea*, xxiii. (1850) 110.—Afr. austr.
brachybotrya, *Muell. Arg. l. c.* xxxiv. (1865–66) 158. —Peruv.
conophora, *Muell. Arg. in Flora*, xlvii. (1864) 530.— Afr. trop.
corniculata, *Sm. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 4.—Ind. or.; Malaya.
hastata, *Muell. Arg. in Flora*, xlvii. (1864) 469.—Afr. austr.
integrifolia, *Vahl, Eclog. Am.* iii. 43.—Guian.
occidentalis, Leand. ex Baill. Etud. Gén. Euph. 484 = tamnoides.
penninervia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 158.—Venezuela.
peruviana, *Muell. Arg. l. c.* 157.—Peruv.
polyadenia, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 335. —Bras.
tamnoides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 158.—Bras.
verrucosa, *Sm. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 4.—Guian.
volubilis, *Linn. Sp. Pl.* 1192.—Ind. occ.
volubilis, Linn. f. Suppl. 421 = verrucosa.

PLUKNETIA, Linn. Gen. ed. I. 373 (1737) = praec.

PLUMARIA, Opiz, Seznam, 75 (1852) = **Dianthus**, Linn. (Caryophyll.).
revoluta, Opiz, l. c. = D. superbus.
superba, Opiz, l. c. = D. superbus.
vulgaris, Opiz, l. c. = D. superbus.

PLUMBAGELLA, Spach, Hist. Vég. Phan. x. 333 (1841) = **Plumbago**, Tourn.
micrantha, Spach, l. c., in nota = Plumbago micrantha.

PLUMBAGIDIUM, Spach, Hist. Vég. Phan. x. 338 (1841) = **Plumbago**, Tourn.
auriculatum, Spach, l. c. 339 = Plumbago capensis.
roseum, Spach, l. c. = Plumbago rosea.
scandens, Spach, l. c. = Plumbago scandens.

PLUMBAGINELLA, Ledeb. Fl. Ross. iii. 471, in syn. (1846–51) = seq.

PLUMBAGO, Tourn. ex Linn. Syst. ed. I (1735).
PLUMBAGINEAE, Benth. & Hook. f. ii. 627.
MOLUBDA, Rafin. Sylva Tellur. 123 (1838).
PLUMBAGELLA, Spach, Hist. Vég. Phan. x. 333 (1841).

PLUMBAGO:—
PLUMBAGIDIUM, Spach, Hist. Vég. Phan. x. 338 (1841).
PLUMBAGINELLA, Ledeb. Fl. Ross. iii. 471, in syn. (1846–51).
THELA, Lour. Fl. Cochinch. 119 (1790).
alba, Hort. ex Pasq. Cat. Orto Bot. Nap. 82 = capensis.
amplexicaulis, *Oliver, in Journ. Linn. Soc.* xv. (1876) 96.—Afr. trop.
angustifolia, Spach, Hist. Vég. Phan. x. 337 = europaea.
aphylla, *Boj. ex Boiss. in DC. Prod.* xii. 694.— Madag.
auriculata, Blume, Bijdr. 736 = zeylanica.
auriculata, Lam. Encyc. ii. 270 = capensis.
caerulea, *H. B. & K. Nov. Gen. et Sp.* ii. 220.— Peruv.
capensis, *Thunb. Prod. Pl. Cap.* 33.—Afr. austr.
capensis, Willd. ex Boiss. in DC. Prod. xii. 694 = tristis.
coccinea, Salisb. Prod. 122; Boiss. l. c. = rosea.
coerulea, Auct. = caerulea.
eglandulosa, R. Br. in Salt, Abyss. App. 64, nomen = Ceratostigma abyssinicum.
europaea, *Linn. Sp. Pl.* 151.—Europ. austr.; Caucas.
flaccida, Moench, Meth. 429 = zeylanica.
floridana, Nutt. in Am. Journ. Sc. Sér. I. v. (1822) 290 = scandens.
glandulosa, Roem. & Schult. Syst. iv. 5 = Ceratostigma abyssinicum.
glandulosa, Willd. ex Roem. & Schult. l. c. 711 = caerulea.
grandiflora, Tenore, Cat. Orto Nap. (1845) 91 = capensis.
Humboldtiana, Roem. & Schult. Syst. iv. 711 = caerulea.
juncea, *Boj. Hort. Maurit.* 263.—Madag.
lactea, Salisb. Prod. 122 = zeylanica.
lapathifolia, Willd. Sp. Pl. i. 837 = europaea.
Larpentae, *Lindl. in Gard. Chron.* (1847) 732.— China.
mexicana, H. B. & K. Nov. Gen. et Sp. ii. 221 = scandens.
micrantha, *Ledeb. Fl. Alt.* i. 171.—Sibir. altaic.
occidentalis, Sweet, Hort. Brit. ed. I. 333 = scandens.
pulchella, *Boiss. in DC. Prod.* xii. 692.—Mexic.
purpurea, Salisb. Prod. 122 = europaea.
rhombifolia, Steud. Nom. ed. II. ii. 357, sphalm. = seq.
rhomboidea, Hook. Bot. Mag. t. 2917 = caerulea.
rhomboidea, Lodd. Bot. Cab. t. 1536 = pulchella.
rosea, *Linn. Sp. Pl.* ed. II. 215.—Ind. or.
sarmentosa, Lam. Illustr. i. 470 = scandens, zeylanica.
scandens, *Linn. Sp. Pl.* ed. II. 215.—Am. trop.
tomentosa, Hook. f. in Trans. Linn. Soc. xx. (1847) 194, err. typ. = Plantago tomentosa.
toxicaria, *Bertol. in Mem. Acc. Sc. Bolog.* ii. (1850) 564.—Afr. trop.
tristis, *Ait. Hort. Kew.* ed. II. i. 324.—Afr. austr.
undulata, Moench, Meth. Suppl. 153 = europaea.
virginica, Hook. f. in Trans. Linn. Soc. xx. (1847) 194, err. typ. = Plantago zeylanica.
viscosa, Blanco, Fl. Filip. ed. I. 78 = zeylanica.
vogeliaefolia, Eckl. & Zeyh. ex Boiss. in DC. Prod. xii. 694 = tristis.
zeylanica, *Linn. Sp. Pl.* 151.—Geront. trop.

PLUMEA, Lunan, Hort. Jamaic. ii. 77 (1814) = **Guarea**, Linn. (Meliac.).

PLUMERIA, Tourn. ex Linn. Syst. ed. I (1735).
APOCYNACEAE, Benth. & Hook. f. ii. 704.
CONOPHARYNGIA, G. Don, Gen. Syst. iv. 94 (1837).
HIMATANTHUS, Willd. in Roem. & Schult. Syst. v. p. xiii (1819).
PLUMIERA, Linn. Phil. Bot. 31 (1751).
acuminata, Ait. Hort. Kew. ed. II. ii. 70 = acutifolia.
acutifolia, *Poir. Encyc. Suppl.* ii. 667.—Mexic.
africana, *Mill. Gard. Dict.* ed. VIII. n. 5.—Afr. trop.
alba, Aubl. Pl. Gui. i. 259 = cuneata.
alba, Hort. ex A. DC. in DC. Prod. viii. 392 = acutifolia.
alba, H. B. & K. Nov. Gen. et Sp. iii. 230 = bicolor.
alba, *Linn. Sp. Pl.* 210.—Ind. occ.
ambigua, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 37.— Bras.

PLUMERIA:—
angustiflora, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. I. 42 = attenuata.
angustifolia, *A. DC. in DC. Prod.* viii. 394.—Guian.
arborea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = acutifolia.
arborescens, *G. Don, Gen. Syst.* iv. 93.—Jamaic.
articulata, *Vahl, Eclog. Am.* ii. 20.—Am. austr.
attenuata, *Benth. in Hook. Journ. Bot.* iii. (1841) 245. —Bras.
aurantia, Endl. in Harting. Parad. Vindob. t. 50 = rubra.
aurantia, Lodd. Cat. 15; ex G. Don, Gen. Syst. iv. 94 = lutea.
aurantiaca, Steud. Nom. ed. II. ii. 357 = praec.
Berterii, *A. DC. in DC. Prod.* viii. 393.—Cuba.
bicolor, *Ruiz & Pav. Fl. Per.* ii. 21. t. 141.—Am. austr.
bicolor, Seem. Bot. Voy. Herald, 168 = Tenorii.
Blandfordiana, Lodd. Cat. 15; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Am. austr.
bracteata, *A. DC. in DC. Prod.* viii. 394.—Bras.
carinata, *Ruiz & Pav. Fl. Per.* ii. 21. t. 140.— Peruv.
clusioides, *Griseb. Cat. Pl. Cub.* 171.—Cuba.
conspicua, *Hort. ex G. Don, in Loud. Hort. Brit.* 68. —Am. austr.
cuneata, *Sm. in Rees, Cycl.* xxvii. n. 6.—Guian.
drastica, *Mart. Reise Bras.* i. 547.—Bras.
dupliciflora, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 (Tabernaemontae sp.).—Java.
emarginata, *Griseb. Cat. Pl. Cub.* 171.—Cuba.
fallax, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 38.— Bras.
filifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 519.—Ind. occ.
floribunda, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 40.— Bras.
Gouani, D. Don, ex G. Don, Gen. Syst. iv. 94 = Lambertiana.
Hilariana, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 39.— Bras.
hypoleuca, Gasparr. in Ann. Civ. i. (1833) 122 = alba.
incarnata, *Mill. Gard. Dict.* ed. VIII. n. 2; *Ruiz & Pav. Fl. Per.* ii. 20. t. 138.—Peruv.
inodora, *Jacq. Enum. Pl. Carib.* 13.—N. Granat.
Jaegeri, *Muell. Arg. in Linnaea*, xxx. (1859–60) 397. —Am. trop.
jamaicensis, Lodd. Cat. 15; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Jamaic.
Jamesoni, *Hook. Bot. Mag.* t. 4751.—Ecuador.
Kerrii, G. Don, Gen. Syst. iv. 93 = tricolor.
Kunthiana, Kostel. Allg. Med.-Pharm. Flora, iii. 1067 = bicolor.
Lambertiana, *Lindl. Bot. Reg.* t. 1378.—Mexic.
lancifolia, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 41.— Bras.
laurifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 (Quid?).—Java.
leucantha, Lodd. Cat. 15; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Am. austr.
longifolia, *Boj. Hort. Maurit.* 210 (an seq.?).— Madag.
longifolia, *Lam. Encyc.* ii. 542.—Madag.
loranthifolia, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 42. —Bras.
lutea, *Ruiz & Pav. Fl. Per.* ii. 21. t. 142.—Peruv.
macrophylla, Lodd. Cat. 15; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Am. austr.
Malongo, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. I. 42 = attenuata.
Martii, *Muell. Arg. l. c.* 37.—Bras.
megaphylla, *A. DC. in DC. Prod.* viii. 391.—Mexic.
mexicana, Lodd. Bot. Cab. t. 1024 = Lambertiana.
Milleri, G. Don, Gen. Syst. iv. 93 = incarnata.
mollis, *H. B. & K. Nov. Gen. et Sp.* iii. 230.—Am. austr.
Mulongo, Benth. in Journ. Linn. Soc. v. (1861) 6 = attenuata.
nivea, *Lodd. ex Sweet, Hort. Brit.* ed. I. 275, nomen. —Am. austr.
nivea, Mill. Gard. Dict. ed. VIII. n. 4 = obtusa.
Northiana, Lodd. Cat. 16; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Am. austr.
obovata, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 40.— Bras.
obtusa, Bert. ex A. DC. in DC. Prod. viii. 393 = Berterii.

PLUMERIA:—
obtusa, *Linn. Sp. Pl.* 210.—Ind. occ.; Am. austr.
obtusa, Lour. Fl. Cochinch. 117 = acutifolia.
obtusifolia, Steud. Nom. ed. I. 633 = acutifolia.
papuana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 36.—N. Guin.
parvifolia, *Donn, Hort. Cantab.* ed. VI. 64; *Haw. Suppl. Pl. Succ.* 7.—Ind. occ.
phagedaenica, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 38 = fallax.
phagedaenica, *Mart. Reise Bras.* iii. 1128.—Bras.
puberula, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 39.—Bras.
pudica, *Jacq. Enum. Pl. Carib.* 13.—Ind. occ.
purpurea, *Ruiz & Pav. Fl. Per.* ii. 20. t. 187.—Peruv.
retusa, Bout. ex A. DC. in DC. Prod. viii. 392 = acutifolia.
retusa, *Lam. Encyc.* ii. 542.—Madag.
revolutifolia, Stokes, Bot. Mat. Med. i. 501 = alba.
rubra, *Linn. Sp. Pl.* 209.—Am. trop.
sericifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 171.—Cuba.
speciosa, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 36.—Bras.
subsessilis, *A. DC. in DC. Prod.* viii. 393.—Ind. occ.
Sucuuba, *Spruce, ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 40.—Bras.
Tenorii, *Gasparr. in Ann. Civ.* i. (1833) 123.—Jamaic.
tenuifolia, Lodd. Cat. 16; ex G. Don, Gen. Syst. iv. 94 (Quid?).—Am. austr.
tricolor, *Ruiz & Pav. Fl. Per.* ii. 20. t. 139.—Peruv.
tuberculata, *Lodd. Bot. Cab.* t. 681.—Ins. S. Doming.
velutina, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 38.—Bras.
versicolor, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Cuba?
Warmingii, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 99.—Bras.
xanthostoma, *Schlecht. in Linnaea*, viii. (1833) 523.—Mexic.

PLUMIERA, Linn. Phil. Bot. 31 (1751) = praec.

PLUMIERIA, Scop. Introd. 155 (1777) = praec.

PLUMMERA, A. Gray, in Proc. Am. Acad. xvii. (1881–82) 215. *COMPOSITAE.*
floribunda, *A. Gray, l. c.*—Arizona.

PLURIDENS, Neck. Elem. i. 86 (1790) = **Bidens**, Linn. (Compos.).

PLUTEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Lansium**, Rumph. (Meliac.).

PLUTONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Phaleria**, Jack (Thymel.).

PNEUMARIA, Hill, Veg. Syst. vii. 40 (1764) = **Mertensia**, Roth (1797) (Borag.).
maritima, Hill, l. c. = M. maritima.
sibirica, Hill, l. c. = M. sibirica.
virginica, Hill, l. c. = M. pulmonarioides.

PNEUMONANTHE, Gleditsch, in Mém. Acad. Berl. v. 1749 (1751) 130 = **Gentiana**, Tourn.
acaulis, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. acaulis.
adscendens, F. W. Schmidt, l. c. = G. decumbens.
algida, F. W. Schmidt, l. c. = G. frigida.
angustifolia, Gilib. Fl. Lituan. i. 34 = G. Pneumonanthe.
angustifolia, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10, 14 = G. acaulis.
asclepiadea, F. W. Schmidt, l. c. = G. asclepiadea.
bicuspidata, G. Don, Gen. Syst. iv. 194 = G. adsurgens.
campanulata, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 9 = G. punctata.
campestris, F. W. Schmidt, ex Steud. Nom. ed. II. ii. 357 = G. punctata.
Catesbaei, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. Saponaria.
depressa, G. Don, in Phil. Mag. (1836) 76; et in Trans. Linn. Soc. xvii. (1837) 510 = G. depressa.
exaltata, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = Eustoma silenifolium.
frigida, F. W. Schmidt, l. c. = G. frigida.

PNEUMONANTHE:—
gelida, G. Don, Gen. Syst. iv. 194 = Gentiana gelida.
glauca, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. glauca.
gracilis, Rafin. Fl. Tellur. iii. 28 = G. gracilis.
grandiflora, S. F. Gray, Nat. Arr. Brit. Pl. ii. 335 = G. acaulis.
incarnata, G. Don, Gen. Syst. iv. 195 = G. ochroleuca.
intermedia, G. Don, l. c. = G. ochroleuca.
Kurroo, G. Don, in Phil. Mag. (1836) 75; Royle, Illustr. Bot. Himal. 278 = G. Kurroo.
media, Rafin. Fl. Tellur. iii. 19, 28 = G. Pneumonanthe.
Michauxii, G. Don, Gen. Syst. iv. 194 = G. Saponaria.
minor, Rafin. Fl. Tellur. iii. 28 = G. Pneumonanthe?
montana, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. montana.
ochroleuca, G. Don, Gen. Syst. iv. 195 = G. ochroleuca.
ornata, G. Don, l. c. 194 = G. ornata.
pannonica, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 9 = G. pannonica.
plicata, F. W. Schmidt, l. c. 10, 13 = G. asclepiadea.
Pseudopneumonanthe, G. Don, in Sweet, Hort. Brit. ed. III. 471 = G. linearis.
punctata, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 9 = G. punctata.
purpurea, F. W. Schmidt, l. c. = G. purpurea.
rigida, Rafin. Fl. Tellur. iii. 28 = G. rigida.
Saponaria, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. Saponaria.
saxosa, F. W. Schmidt, l. c. = G. montana.
septemfida, F. W. Schmidt, l. c. 9 = G. septemfida.
Torreyana, Rafin. Fl. Tellur. iii. 28 = G. Torreyana.
triflora, F. W. Schmidt, in Roem. Arch. i. 1. (1796) 10 = G. triflora.
villosa, F. W. Schmidt, l. c. = G. ochroleuca.
viscosa, F. W. Schmidt, l. c. = Ixanthus viscosus.
vulgaris, F. W. Schmidt, l. c. = G. Pneumonanthe.

PNEUMONANTHOPSIS, Miq. Stirp. Surin. Sel. 150 (1850) = **Voyria**, Aubl. (Gentian.).
clavata, Miq. l. c. = V. clavata.
leucantha, Miq. l. c. = V. leucantha.

POA, Linn. Gen. ed. I. 20 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1196.
ALLAGOSTACHYUM, Nees, ex Steud. Nom. ed. II. i. 50 (1840).
ARCTOPHILA, Rupr. in Beitr. Pfl. Russ. Reich. ii. 62 (1845).
POIDIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836).
PLOTIA, Schreb. ex Steud. Nom. ed. II. ii. 356 (1841).
abbreviata, Blytt, ex Ledeb. Fl. Ross. iv. 373 = cenisia.
abbreviata, *R. Br. in Parry, 1st Voy. Suppl.* 287 (err. typ. 187).—Am. arct.
abietina, Schur, Enum. Pl. Trans. 771 = Leersia oryzoides.
abortiva, R. Rr. Prod. 181 = Heterachne Brownii.
abyssinica, *Jacq. Misc.* ii. 364.—Abyss.
acicularifolia, *J. Buch. Indig. Grasses N. Zeal.* t. 49.—N. Zel.
acicularis, Kunth, Enum. Pl. i. 342 = Eragrostis acicularis.
acinaciphylla, *E. Desv. in C. Gay, Fl. Chil.* vi. 412.—Chili.
acroleuca, *Steud. Syn. Pl. Gram.* 256.—Japon.
acutiflora, H. B. & K. Nov. Gen. et Sp. i. 161 = Eragrostis acutiflora.
adspersa, Drejer, Fl. Excurs. Hafn. 35 = serotina?
adusta, *J. & C. Presl, Rel. Haenk.* 271.—Peruv.
aegyptiaca, Willd. Enum. Hort. Berol. 107 = Eragrostis aegyptiaca.
aestivalis, *J. & C. Presl, Rel. Haenk.* i. 272.—Peruv.
aetnensis, *Guss. Fl. Sic. Prod.* i. 90.—Sicil.
affinis, R. Br. Prod. 179 = caespitosa.
affinis, Salzm. ex Steud. Nom. ed. II. ii. 357 = Eragrostis Vahlii.
agrostidea, DC. Syn. Gall. 132 = Airopsis agrostidea.
airoides, Koehl. Desc. Gram. 194 = Catabrosa aquatica.
airoides, Kunth, Enum. Pl. i. 360 = Eragrostis airoides.
airoides, Nutt. Gen. Am. i. 68 = Glyceria distans.
Aitchisoni, *Boiss. Fl. Orient.* v. 602.—Afghan.
alata, *Desv. Opusc.* 102.—Carolin.

POA:—
alba, Kunth, Enum. Pl. i. 343 = Eragrostis alba.
Alberti, *Regel, in Act. Hort. Petrop.* vii. (1880) 611.—Turkestan.
albida, Turcz. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 387 = Festuca sibirica.
algida, Rupr. Fl. Samojed. Cisural. 61. n. 311 = Phippsia algida.
algida, *Trin. in Linnaea*, x. (1836) 306.—Peruv.
Alopecurus, Kunth, Rév. Gram. i. 116 = Festuca Alopecuros.
alpina, Guss. Fl. Sic. Prod. i. 92 = insularis.
alpina, *Linn. Sp. Pl.* 67.—Reg. bor. et arct.
alpina, Pall. ex Roem. & Schult. Syst. ii. 536 = bulbosa.
alpina, S. Wats. in Bot. King's Exp. 386 = montana, *Vasey.*
alsodes, *A. Gray, Man. Bot. N. U. St.* ed. II. 562.—Am. bor.
altaica, C. A. Mey. Verz. Pfl. Cauc. 19 = cenisia.
altaica, *Trin. in Ledeb. Fl. Alt.* i. 97.—Reg. Caucas.; Sibir.
alternans, *Steud. Nom.* ed. II. ii. 358.—Afr. austr.
altissima, Gilib. Exercit. ii. 529; Moench, Meth. 185 = Glyceria aquatica.
amabilis, [Forsk.] Hort. ex Roem. & Schult. Syst. ii. 566 = Eragrostis aegyptiaca.
amabilis, Linn. Sp. Pl. 68 = Eragrostis amabilis.
amabilis, Walt. Fl. Carol. 79 = Eragrostis spectabilis.
ambigua, Ell. Sketch, i. 165 = Triodia ambigua.
amboinensis, Murr. Syst. ed. XIII. 98 = Eragrostis amboinensis.
amboinicia, Linn. Mant. ii. 557 = praec.
amoena, Kunth, Enum. Pl. i. 342 = Eragrostis amoena.
anceps, *Forst. f. Prod.* 8.—N. Zel.
anceps, Hegetschw. Fl. Schw. 81 = pratensis.
anceps, Presl, Cyp. & Gram. Sic. 43 = compressa.
anceps, Rehm. in Spraw. Kom. Fis. Krakow, vii. (1873) 5 = caesia.
andina, Nutt. ex S. Wats. in Bot. King's Exp. 388 (= *Atropis californica*).—Am. bor. occ.
andina, *Trin. in Linnaea*, x. (1835–36) 306.—Peruv.
angusta, *J. & C. Presl, Rel. Haenk.* i. 271.—Peruv.
angustata, R. Br. in Parry's 1st Voy. Suppl. 287 (err. typ. 187) = Glyceria angustata.
angustata, J. Vahl, in Bot. Notiser (1840) 155, nomen = Glyceria Vahliana.
angustifolia, Bast. Fl. Maine-et-Loire, 39 = nemoralis.
angustifolia, Linn. Sp. Pl. 67 = pratensis.
angustifolia, Reichb. Fl. Germ. Excurs. 47 = serotina.
angustifolia, Ucria, Hort. Reg. Panorm. 58 = trivialis.
angustifolia, Wahlenb. Fl. Suec. 58 = Leersia oryzoides.
annua, Cham. & Schlecht. in Linnaea, vi. (1831) 38 = infirma.
annua, *Linn. Sp. Pl.* 68.—Cosmop.
annua, Steud. Syn. Pl. Gram. 256 = puberula.
anomala, Kunth, Enum. Pl. i. 365 = Eragrostis Uniolae.
antennata, Delile, ex Poir. Encyc. Suppl. iv. 330 = Eragrostis aegyptiaca.
appennina, Jan, Elench. 2. n. 33; ex Bertol. Fl. Ital. i. 527 = alpina.
aquatica, Linn. Sp. Pl. 67 = Glyceria aquatica.
arachnifera, *Torr. in Marcy, Exp. Red Riv. Louis. Bot.* (1853) 301.—Am. bor.
araratica, Trautv. in Act. Hort. Petrop. ii. (1873) 486 = sterilis.
arctica, R. Br. in Parry's 1st Voy. Suppl. 288 (err. typ. 188) = cenisia.
arctica, C. Koch, in Linnaea, xxi. (1848) 403 = caucasica.
arenaria, Retz. Fl. Scand. ed. I. 16 = Glyceria maritima.
arenaria, Willd. ex Spreng. Syst. ii. 342 = Distichlis prostrata.
ariguensis, *Steud. Syn. Pl. Gram.* 259.—Chili.
aristata, Beauv. Agrost. 174 = Koeleria cristata.
aristata, *Phil. in Anal. Univ. Chil.* (1873) 574.—Chili.
articulata, Schrank, in Syll. Ratisb. i. (1824) 194 = Eragrostis articulata.
arundinacea, Bieb. Fl. Taur. Cauc. i. 60 = Glyceria remota.
arundinacea, Link, Hort. Berol. i. 176 = tatarica.
arundinacea, Moench, Meth. 186 = Glyceria maritima.
arundinacea, Poir. Encyc. Suppl. iv. 329 = Triodia cuprea.

POA:—
arvensis, *Gromow, ex Trautv. in Act. Hort. Petrop.* ix. (1884) 329, nomen.—Rossia.
aspera, Gaud. in Alpina, iii. (1808) 38 = caesia.
aspera, Jacq. Hort. Vindob. iii. 32 = Eragrostis aspera.
aspera, Schur, Enum. Pl. Transs. 771 = sterilis.
asperula, *Steud. Syn. Pl. Gram.* 252.—Groenl.
asthenes, Roem. & Schult. Syst. ii. 574 = Leptochloa capillacea.
atrata, Beauv. Agrost. 174 = Festuca sylvatica.
atrovirens, Desf. Fl. Atlant. i. 73. t. 17 = Eragrostis atrovirens.
atrovirens, Willd. ex Spreng. Syst. i. 340 = Eragrostis nigricans.
attenuata, 'Czetz. Erd. Mus. vi. (1872) 18' = alpina.
attenuata, Trin. in Bunge, Verz. Suppl. Fl. Alt. 9 = sterilis.
attica, Boiss. & Heldr. Diagn. Ser. I. xiii. 57 = pratensis.
aturensis, H. B. & K. Nov. Gen. et Sp. i. 161 = Eragrostis aturensis.
aulacosperma, Fresen. in Mus. Senckenb. ii. (1837) 144 = Eragrostis papposa.
aurata, Beauv. ex Steud. Nom. ed. II. i. 628 in syn. = Festuca spadicea.
aurea, Walp. Ann. i. 940 = Eragrostis plumosa.
auriculata, Hornem. ex Steud. Nom. ed. II. i. 563, in syn. = Eragrostis Neesii.
australis, R. Br. Prod. 179 = caespitosa.
australis, [Nees] Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 388 = caespitosa.
autumnalis, Muhl. in Ell. Sketch. i. 159 = flexuosa.
badensis, Haenke, ex Willd. Sp. Pl. i. 392 = alpina.
bahiensis, Schrad. in Schult. Mant. ii. 318 = Eragrostis bahiensis.
Balbisii, *Parl. Fl. Ital.* i. 360.—Sardin.
Balfourii, Parn. in Ann. & Mag. Nat. Hist. Ser. I. x. (1842) 122. t. 5 = caesia.
banatica, Kit. in Linnaea, xxxii. (1863) 311 = seq. ?
bannatica, Willd. ex Steud. Nom. ed. II. ii. 358 = Festuca montana.
barbarea, Spreng. ex Steud. l. c. = Festuca divaricata.
barbata, Thunb. Fl. Jap. 50. t. 10 = Eragrostis barbata.
Barrelieri, Bivona, Piante Ined. 5 = Glyceria fluitans.
Bergiana, Kunth, Rév. Gram. ii. 549 = Eragrostis Bergiana.
Bergii, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 374.—Reg. Argent.
Berteroniana, Kunth, Rév. Gram. i. 112 = Eragrostis Berteroniana.
bifaria, Vahl, Symb. Bot. ii. 19 = Eragrostis coromandelina.
biflora, Ferat, ex Nym. Consp. 834 = nemoralis.
biflora, Retz. Obs. v. 19 = Eragrostis minutiflora.
biformis, Kunth, Rév. Gram. ii. 471. t. 149 = Eragrostis biformis.
Bigelovii, *Vasey & Scribn. ex Vasey, Cat. Grass. U. St.* 81.—Am. bor. occ.
Billardieri, *Steud. Syn. Pl. Gram.* 262.—Austral.
binata, *Nees, Fl. Afr. Austr.* 379.—Afr. austr.
binervata, Beauv. Agrost. 174 = Chaixii.
binervata, Ehrh. ex Beauv. Agrost. 174 = Festuca sylvatica.
bipollicaris, *Hochst. in Flora*, xxxviii. (1855) 321.—Abyss.
Bivonae, Parl. in Guss. Fl. Sic. Syn. i. 99 = insularis.
blepharachne, *Steud. Syn. Pl. Gram.* 252.—Groenl.
blepharophora, *Trin. ex Steud. Nom.* ed. II. ii. 358.—Hab. ?
bohemica, Honck. Syn. Pl. Germ. i. 542; Mayer, ex Mert. & Koch, Deutschl. Fl. i. 621 (Quid ?).—Europ.
Bolanderi, *Vasey, in Coult. Bot. Gaz.* vii. (1882) 32.—Calif.
bonariensis, Kunth, Rév. Gram. i. 115 = pallens.
borbonica, *Kunth, l. c.* ii. t. 151 (non seq.).—Ins. Borbon.
borbonica, *Poir. Encyc. Suppl.* iv. 331.—Ins. Borbon.
borealis, *Kunth, Enum. Pl.* i. 326.—Am. bor. occ.
Borreri, Hook. & Arn. Brit. Fl. ed. VIII. 549 = Glyceria Borreri.
Boryana, Willd. Enum. Hort. Berol. 109 = Eragrostis ciliaris, *Link*.
botryoides, Trin. ex Bess. in Flora, xvii. (1834) 1. Beibl. 28, 30 = Leersia oryzoides.
Bourgaei, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 580.—Mexic.

POA:—
brachiata, Desv. Opusc. 100 = Panicum debile.
brachyphylla, Schult. Mant. ii. 304 = brevifolia.
brasiliensis, Raddi, Agrost. Bras. 51 = Eragrostis bahiensis.
brasiliensis, Spreng. Syst. i. 342 = Eragrostis elegans.
brevifolia, Gaud. in Alpina, iii. (1808) 34 = alpina.
brevifolia, Kunth, Rév. Gram. i. 111 = Aeluropus pubescens.
brevifolia, *Muhl. Desc. Gram.* 138.—Am. bor.
breviglumis, *Hook. f. Fl. Antarct.* 101.—N. Zel.
brizochloa, F. Muell. in Trans. Vict. Inst. i. (1855) 45 = nodosa.
brizoides, Linn. f. Suppl. 110 = Eragrostis brizoides, *Nees*.
brizoides, Wohlleb. Suppl. Fl. Halens. 5 = alpina.
bromoides, *Vahl, Symb. Bot.* iii. 10.—Peruv.
Brownei, Kunth, Rév. Gram. i. 112 = Eragrostis Brownei.
bryophila, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Sibir.
bulbosa, *Linn. Sp. Pl.* 70.—Europ.; As. or.; Afr. bor.
bulbosa, Steud. Syn. Pl. Gram. 256 = sinaica ?
cachectica, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 66. (Eragrostidis sp. ?).—Afr. trop.
caduca, *Kit. ex Rochel, Pl. Banat. Rar.* 26, nomen; *et in Linnaea*, xxxii. (1863) 311.—Banat.
caerulea, Bernh. Syst. Verz. Erf. 43 = Molinia caerulea.
caerulea, Knapp, Gram. Brit. t. 118 = pratensis.
caerulescens, Michx. ex Kunth, Enum. Pl. i. 319 = Triodia cuprea.
caesia, *Sm. Fl. Brit.* i. 103.—Reg. bor. temp.
caespitosa, *Forst. f. Prod.* 89.—Austral.; N. Zel.
caespitosa, Poir. Encyc. v. 73 = nemoralis.
californica, *Scribn. in Bull. Torrey Bot. Club*, x. (1883) 31.—Calif.
californica, Steud. Syn. Pl. Gram. 261 = Dissanthelium californicum.
Calotheca, Kunth, Enum. Pl. i. 340 = Briza Lamarckiana.
calycina, *Kunth, l. c.* 326.—Hab. ?
Cambessediana, Kunth, Rév. Gram. ii. 469. t. 148 = Eragrostis Cambessediana.
Campyle, Schult. Mant. ii. 304 = flexuosa.
canadensis, Beauv. Agrost. 155 = Briza canadensis.
canariensis, Willd. ex Spreng. Syst. i. 345 = Eragrostis minor.
canina, Hort. Gotting. ex Trin. in Mém. Acad. Sc. Pétersb. Sér. VI. i. (1831) 382, in obs. = cenisia.
capensis, Steud. in Flora, xii. (1829) 488 = Eragrostis Thunbergiana.
capillaris, Falk, Beitr. ii. 107 = Eragrostis pilosa.
capillaris, Link, Enum. Hort. Berol. i. 88 = Eragrostis mexicana.
capillaris, Linn. Sp. Pl. 68 = Eragrostis capillaris.
capillaris, Michx. Fl. Bor. Am. i. 67 = Eragrostis tenuis.
capillaris, Salzm. ex Steud. Nom. ed. II. ii. 358 = Eragrostis polytricha.
capillata, Mérat, Fl. Par. ii. 27 = Festuca ovina.
capitata, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 146 = Eragrostis reptans.
carinata, Poir. Encyc. v. 86 = Eragrostis reptans.
carniolica, Hladn. in Reichb. Ic. Fl. Germ. xi. 34 = concinna.
caroliniana, Spreng. Fl. Hal. Mant. 33 = Eragrostis Purshii.
catabrosodes, C. Koch, in Linnaea, xxi. (1848) 406 = Catabrosa Balansae.
Catharinae, Presl, Bot. Bemerk. 120 = sinaica.
caucasica, *Trin. in Mém. Acad. Sc. Pétersb.* Sér. VI. i. (1831) 378.—Reg. Caucas.
cenisia, *All. Auct. Fl. Pedem.* 40.—Europ.; As. bor.
cenisia, Schrad. Fl. Germ. i. 422 = alpina.
cerealis, Salisb. Prod. 20 = abyssinica.
cernua, Schleich. Cat. Pl. Helv. ed. IV. 26 = caesia.
cernua, *Willd. in Ges. Naturf. Fr. Berl. Neue Schr.* iv. (1803) 182.—Ind. or.
Chaixii, *Vill. Fl. Delph.* 7.—Europ.; As. Min.
chalcantha, Kunth, Enum. Pl. i. 339 = Eragrostis chalcantha.
Chapelieri, Kunth, Rév. Gram. ii. 543. t. 186 = Eragrostis elongata.
chariis, Schult. Mant. ii. 314 = Eragrostis Brownei.
chilensis, Hort. Par. ex Steud. Nom. ed. II. ii. 359 = Eragrostis gracilis.
chilensis, Moris, in Ann. Stor. Nat. Bologn. iv. (1830) 60; et in Mem. Acc. Torin. xxxvi. (1833) 196 = Eragrostis virescens.

POA:—
chilensis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Chili.
chiloënsis, *Phil. in Linnaea*, xxx. (1859–60) 206.—Chili.
chinensis, Burm. f. Fl. Ind. 27. t. 11; Koen. in Roth, Nov. Pl. Sp. 65 = Leptochloa capillacea.
chinensis, Hort. ex Presl, Repert. Bot. 252 = Eragrostis gracilis.
chinensis, Link, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 408 = Eragrostis pilosa.
chinensis, *Linn. Sp. Pl.* 69.—Ind. or.
chonotica, *Phil. in Linnaea*, xxix. (1857–58) 97.—Chili.
chorizanthe, *E. Desv. in C. Gay, Fl. Chil.* vi. 407.—Chili.
cilianensis, All. Fl. Pedem. ii. 246 = trivialis, Eragrostis cinnaeformis.
cilianensis, Bivona, ex Steud. Syn. Pl. Gram. 252 = insularis.
ciliaris, Linn. Syst. ed. X. 875 = Eragrostis ciliaris, *Link*.
ciliaris, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 67.—Afr. trop.
ciliata, Roxb. Hort. Beng. 82; Fl. Ind. i. 334 = Eragrostis ciliata.
cilicensis, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 234.—As. Min.
cinerea, Vill. Hist. Pl. Dauph, ii. 126 = pratensis.
cinnaeformis, Hornem. ex Schrad. in Linnaea, xii. (1838) 452 = Eragrostis cinnaeformis.
cinnoides, Hort. Hafn. ex Steud. Nom. ed. II. ii. 359 = praec.
clareosa, D. Dietr. Syn. Pl. i. 354, sphalm. = *glareosa*.
coarctata, Hall. f. ex Gaud. in Alpina, iii. (1808) 35 = nemoralis.
coarctata, Kit. ex Steud. Nom. ed. II. ii. 359 = laxa.
coerulea, Auct. vide caerulea.
cognata, Steud. Syn. Pl. Gram. 262 = nodosa.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 340.—N. Zel.
collina, Host, Gram. Austr. ii. t. 66 = alpina.
collina, C. Koch, ex Boiss. Fl. Orient. v. 610 = tatarica.
colorata, Sreud. Syn. Pl. Gram. 252 = cenisia ?
colpodea, T. Fries, in Ofver. Vetens. Akad. xxvi. (1869) 138 = stricta, *Lindeb*.
columbiensis, *Steud. Syn. Pl. Gram.* 261.—Am. bor.
commutata, *Roem. & Schult. Syst.* ii. 555.—Hispan.
compacta, *Salzm. ex Steud. Nom.* ed. II. ii. 359.—Bras.
complanata, Presl, ex Steud. l. c. (Quid ?).—Hab. ?
complanata, Schur, Enum. Pl. Transs. 770 = compressa.
compressa, *Linn. Sp. Pl.* 69.—Reg. bor. temp.
compressa, Schrenk, ex Ledeb. Fl. Ross. iv. 371 = sterilis.
compressa, Urv. in Mém. Soc. Linn. Par. iv. (1826) 600 = alpina.
compta, Kunth, Rév. Gram. i. 113 = Eragrostis compta.
Conceptionis, *Steud. Syn. Pl. Gram.* 259.—Chili.
concinna, R. Br. Prod. 180 = Eragrostis concinna.
concinna, *Gaud. Agrost. Helv.* 196.—Europ.
condensata, Kunth, Enum. Pl. i. 344 = Distichlis condensata.
conferta, Blytt, Norges Fl. i. 123 = caesia.
conferta, Ell. Sketch, i. 158 = Eragrostis conferta.
conglomerata, *Rupr. ex Galeotti, in Bull. Acad. Brux.* ix. 11. (1842) 235.—Mexic.
contracta, Retz. Obs. iii. 11 = Leptochloa capillacea, Eragrostis Thunbergiana.
contracta, Willd. ex Steud. Nom. ed. II. ii. 359 = Eragrostis pastoënsis.
controversa, Bal. in Bull. Soc. Bot. Fr. xxi. (1874) 16 = longifolia, *Trin*.
controversa, *Steud. Syn. Pl. Gram.* 260.—Ins. Falkl.
convoluta, Hartm. ex Reichb. Fl. Germ. Excurs. 45 = seq. ?
convoluta, Hornem. Hort. Hafn. ii. 953, add. = Glyceria distans.
Cookii, *Hook. f. in Phil. Trans.* clxviii. (1879) 14.—Ins. Kerguel.
coromandelina, Koen. ex Rottb. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 191; Roth, Nov. Pl. Sp. 70 = Eragrostis coromandelina.
coronensis, Schur, Enum. Pl. Transs. 773 = alpina.
costata, Schum. Enum. Pl. Saell. i. 28 = pratensis.
crispa, Link, Hort. Berol. i. 184 = laxa.
crispa, Thuill. Fl. Par. ed. II. 45 = bulbosa.

POA :—
cristata, Georgi, Beschr. Russ. Reich. iii. 693; Schreb. Spicil. Fl. Lips. 54 = Koeleria cristata.
cristata, Linn. Syst. ed. XII. 94 = Koeleria phleoides.
cristata, Savi, Bot. Etrusc. ii. 51 = Koeleria glauca.
cristata, Walt. Fl. Carol. 80.—Cf. Ell. Sketch, i. 164 = Festuca rigida.
crocata, Michx. Fl. Bor. Am. i. 68 = nemoralis.
Cummingii, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Chili.
curva, *Nees, ex Steud. Syn. Pl. Gram.* 259.—Chili.
curvata, Koel. Desc. Gram. 207 = Festuca elatior.
curvula, Schrad. in Goett. Gel. Anz. (1821) III. 2073 = Eragrostis Thunbergiana.
cuspidata, Baxt. ex Steud. Nom. ed. II. ii. 359 = flexuosa.
cuspidata, Nutt. in Barton, Comp. Fl. Philad. i. 61 = brevifolia.
cuspidata, Roth, Nov. Pl. Sp. 72 = Eragrostis Vahlii.
cylindracea, *Trin. ex Steud. Nom.* ed. II. ii. 359.—Hab.?
cylindrica, Roxb. Hort. Beng. 8; Fl. Ind. i. 334 = Eragrostis ciliata.
cynosuroides, Retz. Obs. iv. 20 = Eragrostis cynosuroides.
cyperoides, Pourr. ex Steud. Nom. ed. II. ii. 359 = Ammochloa subacaulis.
cyperoides, Thunb. Prod. Pl. Cap. 22 = Eragrostis cyperoides.
dacica, Porc. in Magyar Novényt. Lapok, ix. (1885) 132 = pumila.
dactyliformis, Steud. Syn. Pl. Gram. 426 = scaberula.
dactyloides, *Aubl. Pl. Gui.* i. 51.—Guian.
dactyloides, *H. B. & K. Nov. Gen. et Sp.* i. 163.—Ecuador.
dahurica, Hochst. & Steud. ex Boiss. Fl. Orient. v. 606 = sinaica.
dahurica, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Dahuria.
danthonioides, *Steud. Syn. Pl. Gram.* 258.—Chili.
debilis, Thuill. Fl. Par. ed. II. 43 = nemoralis.
debilis, *Torr. Fl. N. York*, ii. 459.—Am. bor.
decipiens, R. Br. Prod. 181 = Leptochloa capillacea.
decumbens, Scop. Fl. Carn. ed. II. i. 69 = Triodia decumbens.
decussata, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 79 = Glyceria procumbens.
deflexa, Rupr. Beitr. Pfl. Russ. Reich. ii. 62 = Laestadii.
delawarica, Balb. ex Spreng. f. Tent. Suppl. Syst. 6 = seq.?
delawarica, Link, Hort. Berol. i. 174 = Glyceria maritima.
delicatior, *Steud. Syn. Pl. Gram.* 256.—Java.
delicatula, [Fisch.] Hort. Petrop. ex Schrad. in Linnaea, xii. (1838) 448 = Eragrostis delicatula.
delicatula, Wilh. ex Steud. Syn. Pl. Gram. 250 = bulbosa.
densiflora, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 96 = arachnifera.
dentata, Torr. Fl. N. York, ii. t. 155 = Glyceria pallida.
denudata, *Steud. Syn. Pl. Gram.* 259.—Japon.
depauperata, G. Don pat. ex D. Don, in Mem. Wern. Soc. iii. (1821) 300 = leptostachya.
depauperata, *H. B. & K. Nov. Gen. et Sp.* i. 162.—Ecuador.
depauperata, Kit. ex Spreng. Pugill. i. 7 = nemoralis.
dependens, Hornem. Hort. Hafn. i. 92 = Eragrostis elongata.
depressa, J. & C. Presl, Fl. Čech. 20 = pratensis.
desertorum, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Desert. Kirgis.
despiciens, Link, Enum. Hort. Berol. i. 88 = Eragrostis plumosa.
diandra, R. Br. Prod. 180 = Eragrostis diandra.
diandra, Hort. ex Schrad. in Linnaea, xii. (1838) 451 = Eragrostis Purshii.
diandra, F. Muell. Rep. Babb. Exped. 21 = Eragrostis chaetophylla.
diandra, Roxb. Hort. Beng. 8; Fl. Ind. i. 336 = diarrhena.
diandra, Wall. ex Presl, Repert. Bot. 252 = Eragrostis interrupta, *Beauv.*
diantha, Steud. Syn. Pl. Gram. 256 = spondylodes.
diaphana, Boiss. Fl. Orient. v. 611 = soongarica.
diaphora, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 69 = soongarica.

POA :—
diarrhena, *Schult. Mant.* ii. 616.—Ind. or.
digitata, R. Br. Prod. 182 = Leptochloa subdigitata.
dinantha, *Wood, Class Book* (1861), 797 (Quid?).—Am. bor.
dioeca, Vent. ex Kunth, Enum. Pl. i. 336 = Eragrostis reptans.
dioica, Michx. ex Poir. Encyc. v. 87 = praec.
diomedarum, Kunth, Enum. Pl. i. 340 = Eragrostis diomedearum.
discolor, Hoppe, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 375 = alpina.
dissitiflora, *Roem. & Schult. Syst.* ii. 558.—Groenl.
distans, Desf. Cat. Hort. Par. ed. III. 21 = Eragrostis peruviana.
distans, Linn. Mant. i. 32 = Glyceria distans.
distans, Schultz, Prod. Fl. Starg. Suppl. i. 6 = serotina.
disticha, Wulf. in Jacq. Misc. ii. 74 = Oreochloa disticha.
distichophylla, R. Br. Prod. 182 = Distichlis maritima.
distichophylla, Gaud. in Alpina, iii. (1808) 39 = cenisia.
divaricata, Gouan, Illustr. 4 = Sphenopus Gouani.
divaricata, Vill. Hist. Pl. Dauph. ii. 127 = alpina.
diversifolia, *Hack. ex Boiss. Fl. Orient.* v. 600.—As. Min.; Syria.
dives, F. Muell. Census, 134 = Glyceria dives.
dolor, Boiss. & Heldr. Diagn. Ser. II. iv. 136 = cenisia.
dolosa, Janka, in Oestr. Bot. Zeitschr. xiv. (1864) 384 = laxa.
domingensis, Pers. Syn. i. 88 = Eragrostis gigantea, Leptochloa domingensis.
Douglasii, *Nees, in Ann. Nat. Hist.* Ser. I. i. (1838) 284.—Calif.
Drummondiana, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 418 = nodosa.
dubia, [Honck.] Verz. Aller Gew. Teutschl. 235 = pratensis.
dubia, Leers, Fl. Herborn. 28 = trivialis.
dubia, Suter, Fl. Helv. i. 49 = nemoralis.
dulcis, Salisb. Prod. 20 = Catabrosa aquatica.
dura, [Clairv.] Man. Herb. 21 = Glyceria procumbens.
dura, Scop. Fl. Carn. ed. II. i. 70 = Sclerochloa dura.
duriuscula, Willd. ex Spreng. Syst. i. 339 = annua.
Eatoni, *S. Wats. Bot. King's Exped.* 386.—Am. bor. occ.
effusa, Kit. in Schult. Oestr. Fl. ed. II. i. 227 = Leersia oryzoides.
effusa, Steud. Syn. Pl. Gram. 262 = caespitosa.
Ehrhartiana, E. Mey. Chlor. Hanov. i. 632 = Leersia oryzoides.
elatior, Moench, Enum. Pl. Hass. 37; Bernh. Syst. Verz. Erf. 44 = Festuca elatior.
elegans, R. Br. Prod. 181 = Eragrostis Brownei.
elegans, DC. Fl. Fr. iii. 62 = laxa.
elegans, Hall. f. ex Steud. Nom. ed. II. ii. 359 = serotina.
elegans, Poir. Encyc. v. 87 = Eragrostis ciliaris, *Link.*
elegans, Roxb. Hort. Beng. 82; Fl. Ind. i. 338 = Eragrostis elegantula.
elegans. Willd. ex Spreng. Syst. i. 343 = Eragrostis elegans.
elegantula, Kunth, Rév. Gram. i. 114 = Eragrostis elegantula.
Elliottii, Spreng. Syst. i. 338 = flexuosa.
elliptica, Willd. ex Steud. Nom. ed. II. ii. 359 = Eragrostis ciliaris, *Link.*
elongata, Torr. ex Spreng. Neue Entdeck. ii. 104 = Glyceria elongata.
elongata, Willd. Enum. Hort. Berol. i. 108 = Eragrostis elongata.
eminens, J. & C. Presl, Rel. Haenk. i. 272 = glumaris.
Enysii, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 500 = pusilla.
equitans, Kunth, Enum. Pl. i. 340 = Eragrostis equitans.
eragrostiformis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 88 = pratensis.
Eragrostis, Cav. Ic. i. 63 = Eragrostis major.
Eragrostis, Linn. Sp. Pl. 68 = Eragrostis minor.
Eragrostis, Vill. Hist. Pl. Dauph. ii. 135 = Eragrostis pilosa.
eremophila, *Phil. Fl. Atac.* 56.—Chili.
eriophora, *Steud. Syn. Pl. Gram.* 258.—Chili.
exigua, Dum. Obs. Gram. Belg. 113 = palustris.
exigua, *Hook. f. Handb. N. Zeal. Fl.* 338.—N. Zel.
expansa, J. F. Gmel. Syst. 181 = Sphenopus Gouani.

POA :—
expansa, Kunth, Rév. Gram. i. 113 = Eragrostis expansa.
falcata, Gaudich. in Freyc. Voy. Bot. 408. t. 25 = Eragrostis falcata.
familiaris, *Steud. Syn. Pl. Gram.* 426.—Japon.
fascicularis, Kunth, Rév. Enum. Pl. i. 339 = Eragrostis fascicularis.
fasciculata, Torr. Fl. N. Am. 107 = Glyceria distans.
Fenas, Steud. Syn. Pl. Gram. 252 = Festuca elatior.
Feratiana, Boiss. & Reut. Pugill. Pl. Nov. 128 = nemoralis.
ferruginea, Thunb. Fl. Jap. 50 = Eragrostis ferruginea.
fertilis, Host, Gram. Austr. iii. 10. t. 14 = Leersia oryzoides.
fertilis, Ledeb. Fl. Alt. i. 98 = sterilis.
fertilis, Reichb. Ic. Fl. Germ. xi. 37. t. 87 = serotina.
festucaeformis, Host, Gram. Austr. iii. 12. t. 17 = Glyceria distans.
festucoides, Lam. Illustr. i. 182 = spicata.
filifolia, Schur, Enum. Pl. Transs. 768 = pratensis.
filiformis, Link, ex Schrad. in Linnaea, xii. (1838) 451 = Eragrostis pilosa.
filiformis, Thunb. Prod. Pl. Cap. 21 = Eragrostis Thunbergiana.
filipes, *Lange, in Fl. Dan.* t. 2166.—Groenl.
firma, Wormsk. ex Roem. & Schult. ii. 546 = nemoralis.
firmula, S. F. Gray, Nat. Arr. Brit. Pl. ii. 103 = nemoralis.
flabellata, *Hook. f. in Phil. Trans.* clxviii. (1877) 22, in nota.—Reg. Magell.
flaccida, Moench, ex Steud. Nom. ed. II. i. 561, in syn. = abyssinica.
flaccidula, *Boiss. & Reut. Pugill. Pl. Nov.* 128.—Hispan.
flava, Linn. Sp. Pl. 68 = subverticillata, Triodia cuprea.
flava, Willd. ex Steud. Nom. ed. II. i. 563, in syn. = Eragrostis major.
flavescens, Kunth, Enum. Pl. i. 343 = Eragrostis minor.
flavescens, Thom. Cat. Pl. Suiss. (1818) 23, nomen; ex Hegetschw. Fl. Schw. 86 = alpina.
flavicans, *Ledeb. Fl. Ross.* iv. 373.—Ins. Unalaschk.
flexuosa, *Muhl. Desc. Gram.* 148.—Am. bor.
flexuosa, Roxb. Hort. Beng. 82; Fl. Ind. i. 339 = Roxburghiana.
flexuosa, Schleich. ex Gaud. in Alpina, iii. (1808) 40; Host, Gram. Austr. iv. 15. t. 26 = cenisia.
flexuosa, Sm. Eng. Bot. t. 1123 = laxa.
floccosa, Lehm. Pugill. iii. 39 = Eragrostis Bergiana.
floribunda, Schrad. ex Schult. Mant. ii. 319 = Eragrostis Vahlii.
floribunda, Willd. ex Spreng. Syst. i. 345 = Eragrostis minor.
fluitans, Scop. Fl. Carn. ed. II. i. 73 = Glyceria fluitans.
foliosa, *Hook. f. Handb. N. Zeal. Fl.* 338.—N. Zel.
Fonhi, *Phil. in Linnaea*, xxxiii. (1864–65) 294.—Chili.
Fordeana, F. Muell. Fragm. viii. 130 = Glyceria Fordeana.
formosissima, *Willd. ex Steud. Nom.* ed. II. ii. 360.—Am. austr.
Forsteri, *Phil. in Anal. Univ. Chil.* (1873) 573.—Chili.
Forsteri, *Steud. Syn. Pl. Gram.* 260.—Reg. Magell.
Frankii, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 35, in obs. = Eragrostis Frankii.
Freartitis, Orph. ex Boiss. Fl. Orient. v. 607 = caesia.
frigida, Gaud. in Alpina, iii. (1808) 32; Schleich. Cat. Pl. Nov. 20 = alpina.
fulva, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 378.—Am. arct.
fulvescens, *Trin. in Linnaea*, x. (1836) 306.—Peruv.
fusca, Desf. Cat. Hort. Par. ed. III. 387 = Eragrostis senegalensis.
gangetica, Roxb. Hort. Beng. 8; Fl. Ind. i. 340 = Eragrostis minor.
Gaudini, Roem. & Schult. Syst. ii. 548 = caesia.
Gayana, *E. Desv. in C. Gay, Fl. Chil.* vi. 416.—Chili.
Gebleri, *Trin. ex Steud. Nom.* ed. II. ii. 360.—Hab.?
gelida, *Roem. & Schult. Syst.* ii. 540.—Groenl.
gelida, Schur, Enum. Pl. Transs. 775 = alpina.
geniculata, *Dur. Expl. Algér.* t. 40. f. 4.—Alger.
Gerardi, All. Fl. Pedem. ii. 245 = Festuca spadicea.
glabra, Ehrh. Beitr. vi. 82 = pratensis.

POA :—

glabrata, Kunth, Enum. Pl. i. 344 = Eragrostis glabrata.
glareosa, Kunth, l. c. 339 = Eragrostis articulata.
glauca, Fellm. in Bull. Soc. Nat. Mosc. iii. (1831) 303 = caesia.
glauca, Poir. Encyc. v. 73 = gracilis.
glauca, Schk. Cat. Hort. Wittenberg. (1799) 49 = Koeleria glauca.
glauca, With. Bot. Arr. Brit. Pl. ed. III. ii. 148 = nemoralis.
glaucanthos, Schleich. ex Gaud. in Alpina, iii. (1808) 36 = nemoralis.
glaucescens, Roth, Catalect. fasc. iii. 15 = nemoralis.
glomerata, Desf. Tabl. ed. I. 218 = Eragrostis elongata, interrupta, *Beauv.*
glomerata, *Thunb. Prod. Pl. Cap.* 22.—Afr. austr.
glomerata, Walt. Fl. Carol. 80 = Eragrostis conferta.
glumaris, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 379.—Am. bor. occ.
glutinosa, Sw. Prod. Veg. Ind. Occ. 26 = Eragrostis glutinosa.
glycerioides, *Rupr. ex Galeotti, in Bull. Acad. Brux.* ix. II. (1842) 235, nomen.—Mexic.
Gmelini, [Honck.] Verz. Aller Gew. Teutschl. 242 (Quid ?).—Europ.
Gmelini, Koel. Desc. Gram. 161 = nemoralis.
gracilescens, Schrad. Hort. Gotting. fasc. 1 ; ex Roem. & Schult. Syst. ii. 547 = nemoralis.
gracilis, Bert. Misc. i. 420. t. 40. f. 1, 2 = sinaica.
gracilis, *Desf. Tabl.* ed. I. 17.—Hab. ?
gracilis, Hort. in Hafn. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 390, in obs. = Glyceria maritima ?
grandiflora, [Clairv.] Man. Herb. 21 = Festuca elatior ?
gregalis, Suter, Fl. Helv. i. 48 = pratensis.
groenlandica, *Steud. Syn. Pl. Gram.* 252.—Groenl.
Haenkei, Kunth, Enum. Pl. i. 343 = Eragrostis amoena.
Halleri, [Honck.] Verz. Aller Gew. Teutschl. 228 = laxa, nemoralis, Festuca elatior.
Halleri, Suter, Fl. Helv. i. 45 = cenisia.
Halleridis, Roem. & Schult. Syst. ii. 539 = cenisia.
halophila, Schiede, ex Steud. Nom. ed. II. ii. 360 = Glyceria distans.
hansiana, *Nees, in Royle, Illustr. Bot. Himal.* 46.—Ind. or.
havanensis, *Willd. ex Steud. l. c.*—Cuba.
hemipoa, Loret & Barr. Fl. Monsp. 749 = Festuca hemipoa.
heptantha, Steud. Syn. Pl. Gram. 255 = persica.
heteroclita, Desv. Opusc. 100 = tatarica.
heterophylla, Scheele, in Flora, xxvii. (1844) 58 = pratensis.
hexantha, *Wood, Class-Book* (1861), 797 (Quid ?).—Am. bor.
himalayana, *Nees, ex Steud. Syn. Pl. Gram.* 256.—Reg. Himal.
Hippuris, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 69 (Eragrostidis sp.).—Afr. trop.
hirsuta, Auct. Am. ex A. Gray, Man. Bot. N. U. St. ed. V. 632 = Eragrostis pectinacea.
hirsuta, Michx. Fl. Bor. Am. i. 68 = Eragrostis capillaris, *Nees.*
hirsutissima, *Link, Diss. Bot. Suerin* (1795), 44.—Hab. ?
hirta, *Thunb. Fl. Jap.* 49.—Japon.
Hoffmanniana, Opiz, Seznam, 76 = Leersia oryzoides.
Hohenackeri, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 69 = trivialis.
holciformis, *J. & C. Presl, Rel. Haenk.* i. 272.—Chili.
homomalla, *Nees, in Lehm. Pl. Preiss.* ii. 104.—Austral.
Hookeriana, F. Muell. Fragm. viii. 131 = Festuca Hookeriana.
Hosteana, Schur, Enum. Pl. Transs. 775 = alpina.
Howellii, *Vasey & Scribn. ex Vasey, Cat. Grass. U. St.* 82.—Am. bor. occ.
humilis, Ehrh. Beitr. vi. 84 = pratensis.
humilis, C. Koch, in Linnaea, xxi. (1848) 406 = Catabrosa humilis.
humilis, Lej. Fl. Spa, i. 49 = annua.
hybernica, Willd. ex Steud. Nom. ed. II. ii. 360 = Glyceria distans.
hybrida, *Gaud. in Alpina*, iii. (1808) 46.—Europ. ; Reg. Cauc.
hybrida, Mert. & Koch, Deutschl. Fl. i. 610, in obs. = Chaixii.
hydrophila, Pers. Syn. i. 89 = Leersia oryzoides.

POA :—

hydrophyla, Thuill. ex Steud. Nom. ed. I. 636, 638 = serotina
hypnoides, Lam. Illustr. i. 185 = Eragrostis reptans.
hypopsila, Steud. Syn. Pl. Gram. 263 = Eragrostis imbecilla.
iberica, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 15 = longifolia.
imbecilla, R. Br. Prod. 181 = Leptochloa capillacea.
imbecilla, Forst. f. Prod. 89 ; Spreng. Fl. Hal. Mant. i. 33 = Eragrostis imbecilla.
implexa, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 388 = caespitosa.
indica, Bory, ex Steud. Nom. ed. II. i. 564, in syn. = Eragrostis plumosa.
indica, *Koen. ex Rottb. in Ges. Naturf. Fr. Berl. Neue Schr.* iv. (1803) 194.—Ind. or.
infirma, *H. B. & K. Nov. Gen. et Sp.* i. 158.—Mexic.
insularis, *Parl. Fl. Ital.* i. 541.—Sicilia.
intercepta, Spreng. Syst. i. 347 = Eragrostis interrupta, *Beauv.*
intermedia, *J. Buch. Ind. Grass. N. Zeal.* t. 48.—N. Zel.
intermedia, Koel. Desc. Gram. 303 = Festuca elatior.
intermedia, *Steud. Syn. Pl. Gram.* 252.—Groenl.
interrupta, Lam. Illustr. i. 185 = Eragrostis interrupta, *Beauv.*
interrupta, Nutt. in Trans. Am. Phil. Soc. v. (1837) 146 = Eragrostis oxylepis.
interrupta, Sieber, ex Benth. Fl. Austral. vii. 646 = Eragrostis diandra.
japonica, Thunb. Fl. Jap. 31 = Eragrostis plumosa.
jubata, Kern. in Oestr. Bot. Zeitschr. xxiii. (1873) 6 = minor.
juldusicola, *Regel, in Act. Hort. Petrop.* vii. (1880) 612.—As. centr.
Julieti, *Phil. in Anal. Univ. Chil.* (1873) 575.—Chili.
juncea, Suter, Fl. Helv. i. 46 = nemoralis.
juncoides, Gaud. in Alpina, iii. (1808) 37 = nemoralis.
karatavica, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 525.—Turkest.
kerguelensis, Steud. Syn. Pl. Gram. 257 = Festuca scoparia.
Kingii, S. Wats. in Bot. King's Exp. 387 = glumaris.
Kirkii, *J. Buch. Indig. Gras. N. Zeal.* t. 51.—N. Zel.
Kitaibelii, Kunth, Rév. Gram. i. 117 = nemoralis.
Kitaibelii, Schult. Oesterr. Fl. ed. II. i. 225 = trivialis.
Koeleri, DC. Syn. Fl. Gall. 131 = alpina.
koelerioides, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Chili.
Koenigii, Kunth, Rév. Gram. i. 114 = Eragrostis interrupta, *Beauv.*
Labillardieri, *Steud. Syn. Pl. Gram.* 262.—Austral.
labradorica, *Steud. l. c.* 252.—Am. bor.
Laestadii, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 62.—Scandin.
laevis, *Borb. in Oestr. Bot. Zeitschr.* xxvii. (1877) 425.—Hungar.
laevis, R. Br. Prod. 179 = caespitosa.
lagopodioides, Kunth, Rév. Gram. i. 111 = Aeluropus pubescens.
lagopoides, Steud. Nom. ed. II. ii. 360 = Spartina cynosuroides.
Lamarckii, *Kunth, Enum. Pl.* i. 362.—Am. bor.
Langeana, Reichb. Fl. Germ. Excurs. 140* = compressa.
lanigera, *Nees, Agrost. Bras.* 491.—Bras.
lanuginosa, *Poir. Encyc.* v. 91.—Am. austr.
latiflora, Rupr. Beitr. Pfl. Russ. Reich. ii. 62. t. 4 = fulva.
latifolia, Forst. f. Prod. 8 = Centotheca lappacea.
latifolia, *Phil. in Linnaea*, xxix. (1857-58) 97.—Chili.
latifolia, Pohl, Tent. Fl. Bohem. i. 94 = Chaixii.
latispicea, F. Muell. Census, 134 = Glyceria latispicea.
laxa, *Haenke, in Jirasek, Beob. Riesengeb.* (1791) 118.—Europ. ; Am. bor.
laxa, Lam. Illustr. i. 183 = subverticillata.
laxa, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 374 = minor.
laxa, Wahlenb. Fl. Lapp. 40 ; R. Br. in Ross, Voy. ed. II. ii. 192 = cenisia.
laxiflora, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 96 = leptocoma.
laxiuscula, Lange, in Fl. Dan. t. 2946 = caesia.
leersioides, Guss. Fl. Sic. Prod. i. 98 = Eragrostis leersioides.
leersioides, *Kunth, Rév. Gram.* i. 116.—Sicilia.
Lejeunii, Dum. Obs. Gram. Belg. 112 = pratensis.

POA :—

Lemmoni, *Vasey, in Coult. Bot. Gaz.* iii. (1878) 13.—Calif.
lepida, *F. Muell. Fragm.* viii. 130.—Austral.
lepida, *Nees, ex Steud. Syn. Pl. Gram.* 257.—Chili.
lepida, *A. Rich. Tent. Fl. Abyss.* ii. 424.—Abyss.
leptantha, Kunth, Enum. Pl. i. 339 = Eragrostis leptantha.
leptoclada, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 422.—Abyss.
leptocoma, Bong. Vég. Ins. Sitcha, 54 = stenantha.
leptocoma, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 374.—Kamtschat.
leptostachya, R. Br. Prod. 180 = Eragrostis leptostachya.
leptostachya, *D. Don, in Mem. Wern. Soc.* iii. (1821) 299 (Quid ?).—Scotia.
ligularis, *Nees, ex Steud. Syn. Pl. Gram.* 257.—Bras.
ligulata, *Boiss. Voy. Bot. Espagne*, ii. 659. t. 178 a.—Hispan.
Lima, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 392 = Wangenheimia disticha.
limbata, *Link, Hort. Berol.* i. 179.—Nepal.
Lindsayi, *Hook. f. Handb. N. Zeal. Fl.* 340.—N. Zel.
linearis, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 67 = Eragrostis linearis, *Benth.*
linearis, *Trin. ex Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 145.—China.
lineata, Pers. Syn. ii. 89 = Glyceria nervata.
Linkii, Kunth, Rév. Gram. i. 113 = Eragrostis pilosa.
lithuanica, Gorski, in Eichw. Skizze, 117 = Glyceria remota.
littoralis, Gouan, Fl. Monsp. 470 = Aeluropus laevis.
littoralis, Hook. f. Handb. Fl. N. Zeal. 341, sphalm. = Festuca littoralis.
lobata, Kunth, Enum. Pl. i. 346 = Eragrostis geniculata ?
loliacea, Huds. Fl. Angl. ed. I. 35 = Festuca loliacea.
longifolia, *A. Rich. Tent. Fl. Abyss.* ii. 425.—Abyss.
longifolia, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 69.—Reg. Caucas.
lugens, Kunth, Enum. Pl. i. 331 = Eragrostis lugens.
lurida, Kunth, l. c. 342 = Eragrostis lurida.
macilenta, *A. Rich. Tent. Fl. Abyss.* ii. 428.—Abyss.
Mackayi, *J. Buch. Indig. Grass. N. Zeal.* t. 51.—N. Zel.
macrocalyx, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 103).—Sibir. or. ; Turkestan.
madagascariensis, *Lam. Illustr.* i. 185.—Madag.
malabarica, Klein, ex Steud. Nom. ed. II. ii. 30, 360 = Leptochloa capillacea.
malabarica, Linn. Sp. Pl. 69 = Centotheca lappacea.
malabarica, Retz. Obs. v. 19 ; ex Roem. & Schult. Syst. ii. 557 = chinensis.
malmundariensis, Lej. Fl. Spa, i. 50 = pratensis.
marginata, Willd. Enum. Hort. Berol. Suppl. 6 = nemoralis.
maritima, Cav. Ic. ii. t. 126 = Aeluropus laevis.
maritima, Hornem. in Fl. Dan. t. 1626 = Catabrosa aquatica.
maritima, Huds. Fl. Angl. ed. I. 35 = Glyceria maritima.
maritima, Oed. in Fl. Dan. t. 251 = Glyceria distans.
massauensis, Fresen. in Mus. Senckenb. ii. (1837) 143 = Aeluropus laevis.
maxima, Cav. Desc. 314 ; et in Anal. Cienc. Nat. vi. (1803) 149 = Briza maxima.
Maxwelli, *Benth. Fl. Austral.* vii. 653.—Austral.
maypurensis, H. B. & K. Nov. Gen. et Sp. i. 161 = Eragrostis maypurensis.
media, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 87 = alpina.
mediterranea, Nem. ex Chaub. in Act. Linn. Soc. Bord. xix. (1853) 45 = Glyceria distans.
megapotamica, Spreng. Syst. iv. Cur. Post. 35 = Eragrostis purpurascens.
megastachya, Koel. Desc. Gram. 181 = Eragrostis major.
melicoidea, *Link, Hort. Berol.* i. 186.—Hab. ?
melicoides, Fisch. ex Regel, in Act. Hort. Petrop. vii. (1880) 618 = stenantha.
melicoides, Nutt. Gen. Am. i. 68 = Graphephorum melicoideum.
Meratiana, *Kunth, Rév. Gram.* ii. 539. t. 184.—Bras.
mexicana, Lag. Gen. et Sp. Nov. 3 = Eragrostis mexicana.
mexicana, Link, Enum. Hort. Berol. i. 88 = Eragrostis pilosa.
Meyenii, *Nees, in Meyen, Reise* i. 484.—Peruv.
Michauxii, Kunth, Rév. Gram. i. 111 = Distichlis maritima.
micrantha, Schult. Mant. ii. 305 = Eragrostis capillaris.

POA:—
micrantha, Spreng. Syst. i. 344 = Eragrostis pilosa.
microstachya, *Desv. Opusc.* 100 (Quid?).—Afr.
microstachya, Link, Hort. Berol. i. 185 = Eragrostis arioides.
miliacea, Kit. in Schult. Oesterr. Fl. ed. II. i. 227 = Schultesii.
miliacea, *Link, Hort. Berol.* 174.—Gallia; Ital.
minima, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. i. 91 = Mibora verna.
minor, *Gaud. in Alpina*, iii. (1808) 44.—Europ.
minor, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 382 = cenisia.
minuta, Beauv. Agrost. 174 = Aira minuta.
minuta, Poir. Encyc. Suppl. iv. 327 = Festuca parvula.
modesta, *Phil. in Linnaea*, xxx. (1859–60) 205.—Chili.
modesta, Tuckerm. in Am. Journ. Sc. Ser. I. xlv. (1843) 45 = Sporobolus serotinus.
Molinerii, Balb. Elench. 85; Ser. Cat. 2; ex Gaud. Agrost. Helv. i. 191 = alpina.
Molinerii, Schleich. ex Gaud. Fl. Helv. i. 248 = concinna.
molinioides, *Trin. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 331.—Rossia.
montana, All. Fl. Pedem. ii. 245 = nemoralis.
montana, *Balansa, in Bull. Soc. Bot. Fr.* xxi. (1874) 17.—As. Min.
montana, Delarb. Fl. Auv. ii. 699 = Festuca spadicea.
montana, [Honck.] Verz. Aller Gew. Teutschl. 233 = pratensis.
montana, Parn. Grass. Scotl. 86 = caesia.
montana, *Vasey, in Monthly Rep. U. St. Dep. Agric.* (1874) 155.—Am. bor.
monticola, *Gaudich. in Freyc. Voy. Bot.* 408.—Ins. Sandvic.
monticola, *Phil. in Anal. Univ. Chil.* (1873) 576.—Chili.
Montufari, H. B. & K, Nov. Gen. et Sp. i. 159 = Eragrostis Montufari.
mucronata, Poir. Encyc. v. 91 = Eragrostis beninensis.
mucronata, Roth, Nov. Pl. Sp. 64 = Eragrostis mucronata.
mucronulata, Kunth, Rév. Gram. i. 114 = Eragrostis mucronata.
mulalensis, *H. B. & K. Nov. Gen. et Sp.* i. 162.—Ecuador.
multiflora, Forsk. Fl. Aegypt. Arab. 21 = Eragrostis major.
multiflora, Roxb. Hort. Beng. 8; Fl. Ind. i. 338 = Eragrostis multiflora.
multiradiata, *Regel, in Act. Hort. Petrop.* vii. (1880) 620.—Armenia.
muralis, [Honck.] Verz. Aller Gew. Teutschl. 226 = nemoralis.
muralis, Wibel, Prim. Fl. Werth. 114 = compressa.
nana, Savi, in Usteri, Ann. Bot. xxiv. (1800) 49 = Eragrostis peruviana.
naratavica, Anderss. in Walp. Ann. vi. 1013, sphalm. = karatavica.
naravetica, Anderss. l. c. 1258, sphalm. = praec.
nardoides, Kunth, Enum. Pl. i. 340 = Eragrostis coromandelina.
narifolia, Schur, Enum. Pl. Transs. 767 = pratensis.
Neesii, Kunth, l. c. 339 = Eragrostis Neesii.
neglecta, *Steud. Syn. Pl. Gram.* 253.—Dahuria.
nemoralis, Heyne, ex Wall. Cat. n. 3826 = Centotheca lappacea.
nemoralis, *Linn. Sp. Pl.* 69.—Europ.; As. bor.
nemoralis, Pursh, Fl. Am. Sept. i. 79 = serotina.
nemoralis, Torr. Fl. N. Am. i. 111 = alsodes.
nervata, Willd. Sp. Pl. i. 389 = Glyceria nervata.
nevadensis, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 66.—Am. bor. occ.
nigra, Clem. ex Willk. & Lange, Prod. Fl. Hisp. i. 83 = Eragrostis papposa.
nigricans, H. B. & K. Nov. Gen. et Sp. i. 159 = Eragrostis nigricans.
nitens, Kunth, Rév. Gram. i. 113 = Eragrostis nitida, *Link*.
nitens, [Weber] in Wigg. Prim. Fl. Holsat. Suppl. n. 6 = Hierochloë borealis.
nitida, Ell. Sketch, i. 162 = Eragrostis campestris.
nitida, Lam. Illustr. i. 182 = Koeleria cristata.
nitida, Savi, Bot. Etrusc. ii. 51 = Koeleria glauca.
nivalis, Schur, Enum. Pl. Transs. 776 = alpina.
nodosa, *Nees, in Lehm. Pl. Preiss.* ii. 105.—Austral.
Novarae, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxi. (1871) 31.—Ins. St. Paul (Mar. austr.).

POA:—
nutans, [Clairv.] Man. Herb. 21 = Melica nutans.
nutans, Gilib. Exercit. ii. 532 = nemoralis.
nutans, Klein, ex Steud. Nom. ed. II. ii. 359, 361 = cernua.
nutans, Link, Enum. Hort. Berol. i. 86 = Festuca nutans.
nutans, Retz. Obs. iv. 19 = Eragrostis Brownei.
nutkaensis, J. & C. Presl, Rel. Haenk. i. 272 = Glyceria distans.
Nuttalliana, Schult. Mant. ii. 303 = Glyceria distans.
Nuttallii, Kunth, Rév. Gram. i. 116 = Eragrostis Nuttalliana.
Nuttallii, Spreng. Syst. i. 344 = obtusa.
Nymanni, Tineo, in Guss. Fl. Sic. Syn. ii. 783 = pratensis.
oblonga, Baumg. Enum. Stirp. Transs. 238 = Eragrostis major.
obtusa, *Muhl. Desc. Gram.* 147; *Nutt. Gen. Am.* i. 67.—Am. bor.
obtusata, Beauv. Agrost. 175 = Eatonia obtusata.
obvallata, *Steud. Syn. Pl. Gram.* 258.—Chili.
ochotensis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 377.—Sibir.
oligantha, *Hochst. in Flora*, xxxviii. (1855) 322.—Abyss.
oligeria, *Steud. Syn. Pl. Gram.* 426.—Reg. Magellan.
Olmedi, H. B. & K. Nov. Gen. et Sp. i. 159 = Eragrostis Olmedi.
olympica, *Schott, Nym. & Kotschy, Analect. Bot.* 2.—As. Min.
oreophila, Boiss. & Heldr. Diagn. Ser. II. iv. 137, in obs. = cenisia.
oreophila, Schur, Enum. Pl. Transs. 773 = alpina.
ovalis, Tineo, Pl. Rar. Sicil. 21 = annua.
ovina, *A. Rich. Tent. Fl. Abyss.* ii. 428.—Abyss.
oxylepis, Torr. in Marcy, Expl. Red River Louis. Bot. (1853) 301 = Eragrostis oxylepis.
pachypogon, *Nees, ex Steud. Syn. Pl. Gram.* 258.—Chili.
pallens, Hall. f. ex Gaud. in Alpina, iii. (1808) 41 = cenisia.
pallens, *Poir. Encyc.* v. 91.—Reg. Argent.
pallescens, Koch, Syn. Fl. Germ. ed. I, Index 69 = praec.
pallida, Hort. ex Schrad. in Linnaea, xii. (1838) 450 = Eragrostis aegyptiaca.
pallida, Lag. Gen. et Sp. Nov. 3 = Eragrostis pilosa.
paludosa, [Honck.] Verz. Aller Gew. Teutschl. 226 = Leersia oryzoides?
palustris, Linn. Syst. ed. X. 874 = Leersia oryzoides.
palustris, H. Mart. Prod. Fl. Mosq. 19 = serotina.
palustris, O. F. Muell. in Fl. Dan. t. 750; DC. Fl. Fr. Suppl. 272 = nemoralis.
palustris, Wohlleb. Suppl. Fl. Halens. 5 = trivialis.
pamphylica, Boiss. Diagn. Ser. I. xiii. 58 = persica.
panicea, Retz. Obs. iii. 11 = Leptochloa capillacea.
panicea, Willd. ex Spreng. Syst. i. 338 = stolonifera.
paniciformis, A. Br. in Flora, xxiv. (1842) 274 = Eragrostis paniciformis.
panicoides, *Kunth, Enum. Pl.* i. 343.—Mexic.
paniculata, Roxb. Hort. Beng. 82; Fl. Ind. i. 340 = Eragrostis aspera.
pannonica, *Kern. in Oestr. Bot. Zeitschr.* xiv. (1864) 84.—Europ.
papillosa, Schrad. in Goett. Gel. Anz. (1821) III. 2074 = Eragrostis Uniolae.
paposana, *Phil. Fl. Atac.* 55.—Chili.
papposa, Dufour, ex Roem. & Schult. Syst. ii. 585 = Eragrostis papposa.
paradoxa, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 864 = soongarica.
paradoxa, Roem. & Schult. Syst. ii. 569 = Distichlis maritima.
parnassica, Boiss. & Heldr. in Boiss. Diagn. Ser. II. iv. 137, in obs. = pumila.
Parnellii, Bab. Man. Brit. Bot. ed. I. 368 = nemoralis.
parviflora, R. Br. Prod. 180 = Eragrostis pilosa.
parviflora, Pursh, Fl. Am. Sept. i. 80 = Glyceria nervata.
parvula, Roem. & Schult. Syst. ii. 571 = Festuca parvula.
parvula, Steud. Syn. Pl. Gram. 255 = persica.
paryadrica, *Boiss. in Tchihat. Asie Min. Bot.* ii. 608.—As. Min.
Pasqualii, Heldr. ex Parl. Fl. Ital. i. 343 = bulbosa.
pastoënsis, H. B. & K. Nov. Gen. et Sp. i. 160 = Eragrostis pastoënsis.

POA:—
patula, H. B. & K. Nov. Gen. et Sp. i. 158 = Eragrostis patula.
pauciflora, *Benth. ex Vasey, Grass. U. St.* 42.—Calif.
pauciflora, Roem. & Schult. Syst. ii. 549 = depauperata.
pectinacea, Geyer, ex Steud. Syn. Pl. Gram. 272 = Eragrostis Geyeri.
pectinacea, Kunth, Enum. Pl. i. 382 = Koeleria setacea.
pectinacea, Michx. Fl. Bor. Am. i. 69 = Eragrostis pectinacea.
pectinata, Lam. Illustr. i. 182 = Koeleria setacea.
pelligera, Rupr. Beitr. Pfl. Russl. ii. 64 = Graphephorum psilosanthum.
pellucida, R. Br. Prod. 181 = Eragrostis pilosa.
pendulina, J. Vahl, in Fl. Dan. t. 2343 = Laestadii.
Perouini, *Boiss. Fl. Orient.* v. 604.—Cilicia.
persica, *Trin. in Mém. Acad. Sc. Pétersb.* Sér. VI. i. (1831) 373.—Persia; Reg. Caucas.
peruviana, Jacq. Coll. i. 107 = Eragrostis peruviana.
phalaroides, *Nees, ex Steud. Syn. Pl. Gram.* 258.—Chili.
Philippii, *Steud. l. c.* 426.—Chili.
phleoides, Lam. Illustr. i. 182 = Koeleria phleoides.
phoenicoides, Koel. Desc. Gram. 201 = Brachypodium pinnatum.
Phoenix, Scop. Fl. Carn. ed. II. i. 74 = Graphephorum arundinaceum.
Phragmites, *Phil. in Anal. Univ. Chil.* (1873) 576.—Chili.
phryganoides, *Trin. in Mém. Acad. Sc. Pétersb.* Sér. VI. i. (1831) 389.—Kamtchat.
picta, *Trin. ex Steud. Nom.* ed. II. ii. 361.—Hab.?
pilifera, *Scheele, in Linnaea*, xxii. (1849) 344.—Texas.
pilosa, Linn. Sp. Pl. 68 = Eragrostis pilosa.
pilosissima, Kunth, Rév. Gram. i. 112 = Eragrostis articulata.
planicularis, Weber, in Usteri, Ann. Bot. i. (1791) 92 = compressa.
planiculmis, Weber, Suppl. Fl. Holsat. 3; ex Roem. & Schult. Syst. ii. 558 = praec.
plebeia, R. Br. Prod. 179 = caespitosa.
plumosa, Hort. Par. ex Schrad. in Linnaea, xii. (1838) 451 = Eragrostis gracilis.
plumosa, Retz. Obs. iv. 20 = Eragrostis amabilis.
poecila, *Phil. in Anal. Univ. Chil.* (1873) 573.—Chili.
poecilantha, Rupr. Beitr. Pfl. Russ. Reich. ii. 63. t. 5 = fulva.
Poidium, *Doell, in Mart. Fl. Bras.* ii. III. 119.—Bras.
Poiretii, Roem. & Schult. Syst. ii. 553 = Eragrostis pilosa.
Poiretii, *Steud. Syn. Pl. Gram.* 257.—Ins. Borbon.
polygama, *Steud. l. c.* 255.—Oriens.
polymorpha, Balb. ex Kunth, Enum. Pl. i. 332 = Eragrostis Berteroniana.
polymorpha, R. Br. Prod. 180 = Eragrostis Brownei.
polymorpha, Hort. Vindob. ex Schrad. in Linnaea, xii. (1838) 450 = Eragrostis Jacquini.
polymorpha, Koen. ex Rottb. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 194 = Eragrostis bahiensis.
polymorpha, Sieber, ex Kunth, Enum. Pl. i. 330 = Eragrostis tephrosanthos.
polymorpha, Wibel, Prim. Fl. Werth. 113 = nemoralis.
polynoda, Parn. Grass. Scotl. 84 = compressa.
polytricha, Kunth, Enum. Pl. i. 331 = Eragrostis polytricha.
porphyroclados, Nees, in Lehm. Pl. Preiss. ii. 105 = caespitosa.
porrantha, Steud. Syn. Pl. Gram. 262 = caespitosa.
praecox, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 135 = bulbosa.
pratensis, *Linn. Sp. Pl.* 67.—Reg. bor. temp.
pratensis, Pollich, Hist. Pl. Palat. i. 87 = trivialis.
Preslii, Kunth, Enum. Pl. i. 326 = Distichlis thalassica.
Pringlii, *Scribn. in Bull. Torrey Bot. Club*, x. (1883) 31.—Am. bor. occ.
procera, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 332.—Ind. or.
procumbens, Curt. Fl. Lond. fasc. vi. t. 11 = Glyceria procumbens.
prolifera, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 188 = bulbosa.

POA:—
prolifera, *Sw. Prod. Veg. Ind. Occ.* 27.—Ind. occ.
prostrata, H. B. & K. Nov. Gen. et Sp. i. 157 = Distichlis prostrata.
protuberata, Schur, Enum. Pl. Transs. 773 = bulbosa.
psammoides, Kunth, Enum. Pl. i. 328 = Eragrostis psammodes.
psammophila, Schur, Enum. Pl. Transs. 773 = bulbosa.
Pseudo-caesia, Schur, l. c. 772 = nemoralis.
Pseudo-concinna, Schur, l. c. 773 = bulbosa.
Pseudo-hybrida, Schur, l. c. 769 = trivialis.
Pseudo-laxa, Schur, l. c. 777 = laxa.
Pseudo-nemoralis, Schur, l. c. 771 = Leersia oryzoides.
psilocaulis, *Steud. Syn. Pl. Gram.* 256.—Japon.
psilophylla, *Hochst. in Flora*, xxxviii. (1855) 321.—Abyss.
psilosantha, Rupr. Beitr. Pfl. Russ. Reich. ii. 64 = Graphephorum psilosanthum.
psychrophila, Boiss. & Heldr. Diagn. Ser. I. xiii. 57 = cenisia.
puberula, *Steud. Syn. Pl. Gram.* 256.—Arab.
pubescens, R. Br. Prod. 181 = Eragrostis Brownei.
pubescens, Lej. Fl. Spa, i. 51 = pratensis.
pubiflora, *Benth. Pl. Hartw.* 261.—N. Granat.
pulchella, [Clairv.] Man. Herb. 20 = nemoralis?
pulchella, *Parl. in Atti 8va. Riun. Sc. Ital.* 586.—Ins. Cape Virid.
pulchella, *Salisb. Prod.* 21.—Ind. or.
pulchella, Stev. ex Bieb. Fl. Taur. Cauc. i. 417 = Festuca rigida.
pulchella, *Vasey, in Coult. Bot. Gaz.* vii. (1882) 32.—Am. bor.
pumila, *Host, Fl. Austr.* i. 146.—Europ. or.
pumila, C. Koch, in Linnaea, xxi. (1848) 403 = Catabrosa humilis.
Pumilio, *Hochst. in Flora*, xxxviii. (1855) 321.—Abyss.
punctata, Link, Enum. Hort. Berol. i. 87 = Eragrostis filiformis.
punctata, Linn. f. Suppl. 109 = Eragrostis pilosa.
pungens, Bieb. Fl. Taur. Cauc. i. 65 = Aeluropus pubescens.
pungens, Georgi, Beschr. Russ. Reich. iv. 693 = Aeluropus laevis.
pungens, Nutt. Gen. Am. i. 66 = brevifolia.
pungens, Torr. Fl. N. Am. 109 = flexuosa.
purpurascens, Spreng. Nov. Prov. 33 = Eragrostis purpurascens.
purpurascens, *Vasey, in Coult. Bot. Gaz.* vi. (1881) 297.—Am. bor.
purpurea, T. Kirk, in Trans. N. Z. Inst. ix. (1877) 500 = Kirkii.
Purshii, Hort. Erf. Sem. Coll. (1826);—Cf. Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 84 = Eragrostis Purshii.
pusilla, *Bergg. in Minneskr. Fisiog. Sällks. Lund*, (1877) 31. n. 8.—N. Zel.
pygmaea, *J. Buch. Indig. Grasses N. Zeal.* t. 50.—N. Zel.
pyramidalis, [Clairv.] Man. Herb. 19 = nemoralis?
pyramidata, Lam. Illustr. i. 183 = Koeleria cristata.
pyrenaica, Lange, ex Willk. & Lange, Prod. Fl. Hisp. i. 80 = minor.
quadriflora, Moench, ex Steud. Nom. ed. II. ii. 362 = Festuca sylvatica.
quadripedalis, Ehrenb. ex Koel. Desc. Gram. 160 = Chaixii.
quinquefida, Pursh, Fl. Am. Sept. i. 81 = Triodia cuprea.
racemosa, *Thunb. Prod. Pl. Cap.* 21.—Afr. austr.
racemosa, Vahl, Eclog. Am. i. 7 = Eragrostis Vahlii.
radicans, Moench, Meth. 186 = Eragrostis papposa.
Radula, *Franch. & Sav. Enum. Pl. Jap.* ii. 602.—Japon.
ramigera, F. Muell. in Trans. Vict. Inst. i. (1854) 45 = Glyceria ramigera.
ramosa, Savi, in Mem. Soc. Ital. Mod. ix. (1802) t. 10 = Aeluropus villosus.
ramosissima, *Hook. f. Fl. Antarct.* 101.—Ins. Dom. Auckland.
rariflora, Desf. Tabl. ed. II. 20 = nemoralis.
rariflora, Roem. & Schult. Syst. ii. 570 = Orthoclada rariflora.
rariflora, Steud. Nom. ed. I. 638 = Orthoclada laxa.
recta, Willd. ex Steud. l. c. ed. II. ii. 362 = nemoralis.
reflexa, *Vasey & Scribn. ex Vasey, Cat. Grass. U. St.* 83.—Am. bor. occ.

POA:—
refracta, Muhl. Desc. Gram. 146 = Eragrostis pectinacea.
remota, Forselles, in Act. Inst. Linn. Upsal. i. (1807) t. 1; (Tvenne Nya Vaext.) = Glyceria remota.
remota, Fries, ex Nym. Consp. 832 = Festuca sylvatica.
remota, Hartm. Handb. Skand. Fl. ed. II. 28 = Chaixii.
remota, *H. B. & K. Nov. Gen. et Sp.* i. 163.—Ecuador.
remotiflora, Rupr. Beitr. Pfl. Russ. Reich. ii. 63. t. 5 = Laestadii.
repens, Bieb. Fl. Taur. Cauc. iii. 69 = Aeluropus villosus.
repens, Willd. ex Steud. Nom. ed. II. ii. 362 = Distichlis prostrata.
reptans, Michx. Fl. Bor. Am. i. 69. t. 11 = Eragrostis reptans.
reptatrix, Cabr. ex Willk. & Lange, Prod. Fl. Hisp. i. 86 = Sphenopus Gouani.
resinulosa, *Nees, ex Steud. Syn. Pl. Gram.* 259.—Reg. Argent.
retroflexa, Curt. Fl. Lond. fasc. vi. t. 10 = Glyceria distans.
retrofracta, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. i. 91 = praec.
Reuteriana, *Boiss. & Buhse, in Mem. Soc. Nat. Mosc.* xii. (1860) 227.—Persia.
rhenana, Koel. Desc. Gram. 196 (Quid?).—Europ.
rhenana, Lej. Fl. Spa, i. 52 = nemoralis.
rigens, Hartm. Handb. Skand. Fl. ed. I. 448 = pratensis.
rigens, Trin. ex Ledeb. Fl. Ross. iv. 393 = Glyceria glumaris.
rigida, Huds. Fl. Angl. ed. I. 35 = Festuca rigida.
rigida, Linn. Cent. Pl. Rar. i. 5; Amoen. Acad. iv. 265 = praec.?
rigidifolia, *Steud. Syn. Pl. Gram.* 260.—Ins. Falkl.
riparia, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. iv. (1803) 185 = Eragrostis ciliaris, *Link*.
riparia, Wolff, in Hoffm. Deutschl. Fl. ed. II. i. 42 = Leersia oryzoides.
rivularis, Bernh. Syst. Verz. Erf. 41 = Catabrosa aquatica.
robusta, *Phil. in Anal. Univ. Chil.* (1873) 574.—Chili.
robusta, *Steud. Syn. Pl. Gram.* 426.—Reg. Magellan.
Rothii, Kunth, Rév. Gram. i. 114 = Eragrostis interrupta, *Beauv.*
rotundata, Trin. in Mém. Acad. Sc. Pétersb. Sér. VI. i. (1831) 378.—Ins. Unalaschk.
Roxburghiana, *Schult. Mant.* ii. 315.—Ind. or.
Royleana, *Nees, ex Steud. Syn. Pl. Gram.* 256.—Ind. or.
rubens, Lam. Illustr. i. 184 = Eragrostis amabilis.
rubens, Moench, Meth. 187 = Chaixii?
rubiginosa, Kunth, Enum. Pl. i. 339 = Eragrostis rubiginosa.
rubra, *Lepech. Reise*, i. 41.—Ross. med.
rubra, Willd. ex Spreng. Syst. i. 340 = Eragrostis capillaris.
rufescens, Schrad. ex Schult. Mant. ii. 319 = Eragrostis Vahlii.
rupestris, Bieb. ex C. A. Mey. Verz. Pfl. Cauc. 20 = sterilis.
rupestris, Roth, Nov. Pl. Sp. 71 = Eragrostis ciliata.
rupestris, With. Bot. Arr. Brit. Pl. ed. III. ii. 146. t. 24 = Glyceria procumbens.
Ruprechtii, *Peyr. in Linnaea*, xxx. (1859–60) 6.—Mexic.
sabauda, All. Fl. Pedem. ii. 245 = Festuca elatior.
sabauda, Suter, Fl. Helv. ii. 403 = Festuca sylvatica.
sabulosa, Turcz. ex Regel, in Act. Hort. Petrop. vii. (1880) 616 = caucasica.
salebrosa, Panz. ex Steud. Nom. ed. II. ii. 362 = nemoralis.
salina, Leyss. Fl. Hal. 21; et in Abh. Naturf. Ges. Halle, i. (1783) 366 = seq.?
salina, Pollich, Hist. Pl. Palat. i. 92 = Glyceria distans.
sarmentosa, Thunb. Prod. Pl. Cap. 21 = Eragrostis sarmentosa.
saxicola, *R. Br. Prod.* i. 180.—Tasmania.
scaberula, *Hook. f. Fl. Antarct.* 378.—Reg. Magellan.
scabra, Ehrh. Beitr. vi. 83 = trivialis.
scabra, Kit. ex Steud. Nom. ed. II. ii. 362; et in Linnaea, xxxii. (1863) 311 = sterilis.
scabrella, *Benth. ex Vasey, Grass. U. St.* 42.—Calif.

POA:—
scaligera, *Salzm. ex Steud. Nom.* ed. II. ii. 362.—Bras.
scariosa, Lag. Gen. et Sp. Nov. 3 = Festuca granatensis.
Scheuchzeri, Pourr. ex Steud. Nom. ed. II. ii. 362 = alpina.
Scheuchzeri, Suter, Fl. Helv. i. 50 = nemoralis.
Schiedeana, *Trin. ex Steud. Nom.* ed. II. ii. 362—Mexic.
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 423.—Abyss.
Schultesii, *Kunth, Rév. Gram.* i. 117.—Hungar.
scleroclada, Rupr. Beitr. Pfl. Russ. Reich. ii. 63. t. 6 = fulva.
sclerophylla, *Bergg. in Minneskr. Fisiog. Sällks. Lund*, (1877) 30. n. 8.—N. Zel.
sclerophylla, Kunth, Enum. Pl. i. 340 = Eragrostis sclerophylla.
sclerophylla, Spreng. ex Presl, Repert. Bot. 238 = Chaixii.
scoparia, *Kunth, Rév. Gram.* ii. 535. t. 182.—Reg. Argent.
secunda, *J. & C. Presl, Rel. Haenk.* i. 271.—Chili.
secunda, Zea, ex Roem. & Schult. Syst. ii. 697 = Glyceria procumbens.
secundiflora, Kunth, Enum. Pl. i. 342 = Eragrostis secundiflora.
Sellovii, *Nees, Agrost. Bras.* 491.—Reg. Argent.
semineutra, *Trin. Fund. Agrost.* 406.—Hungar.
seminuda, Kunth, Enum. Pl. i. 340 = Eragrostis seminuda.
semiverticillata, *Lag. ex Steud. Nom.* ed. II. ii. 362.—Hab.?
senegalensis, *Desv. Opusc.* 100.—Afr. trop.
senegalensis, Hort. Par. ex Desf. Cat. Hort. Par. ed. III. 21 = Eragrostis senegalensis.
serotina, *Ehrh. Beitr.* iv. 83.—Reg. bor. temp.
serotina, Gaud. Agrost. Helv. i. 208 = Leersia oryzoides.
serotina, Moerck, ex Bab. in Journ. Linn. Soc. xi. (1871) 343 = caesia.
serotina, Schrad. Fl. Germ. i. 292 = nemoralis.
serpentum, Nees, in Lehm. Pl. Preiss. ii. 106 = caespitosa.
seslerioides, All. Fl. Pedem. ii. 246 = Oreochloa pedemontana.
seslerioides, Michx. Fl. Bor. Am. i. 68 = Triodia cuprea.
sessiliflora, Kunth, Rév. Gram. i. 111 = Uniola sessiliflora.
sessilis, Lam. Illustr. i. 185 = Leptochloa capillacea.
sesslerioides, Loisel. Fl. Gall. ed. II. i. 82 = Oreochloa disticha.
setacea, Hoffm. Deutschl. Fl. ed. II. i. 44 = pratensis.
setacea, Huds. Fl. Angl. ed. I. 34 = trivialis.
setacea, Koel. Desc. Gram. 162 = Festuca ovina.
setifolia, *Benth. Pl. Hartw.* 262.—Peruv.
setifolia, Zett. Pl. Vasc. Pyr. 290 = violacea.
sicula, Jacq. Coll. ii. 304 = Demazeria sicula.
Sieberiana, *Kunth, Enum. Pl.* i. 357.—Austral.
Sieberiana, Spreng. Syst. iv. Cur. Post. 35 = caespitosa.
simensis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 422.—Abyss.
similis, Rupr. Beitr. Pfl. Russ. Reich. ii. 63. t. 6 = Laestadii.
simplex, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 86 = annua.
simplex, Wall. Fl. Carol. 79 = Eragrostis hirsuta.
simpliciflora, *Kunth, Enum. Pl.* i. 343.—Mexic.
sinaica, *Steud. Syn. Pl. Gram.* 256.—Arab.; Persia.
sinensis, J. F. Gmel. Syst. 181 = *chinensis*.
Smithii, Kunth, Rév. Gram. i. 111 = Aeluropus Smithii.
solida, Kunth, l. c. ii. 541 = Eragrostis Vahlii.
soongarica, *Boiss. Fl. Orient.* v. 611.—As. bor.
spadicea, Koel. Desc. Gram. 202 = Festuca spadicea.
speciosa, Roem. & Schult. Syst. ii. 573 = Eragrostis Brownei.
speciosa, Willd. ex Spreng. Syst. i. 344 = Eragrostis amabilis.
spectabilis, Pursh, Fl. Am. Sept. i. 81 = Eragrostis pectinacea.
sphondylodes, *Trin. ex Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 145.—China.
spicata, Bivona, Piante Ined. 2; ex Guss. Fl. Sic. Syn. ii. 784 = Glyceria fluitans.
spicata, *Linn. Mant.* i. 32.—Lusit.; Hispan.
spicata, Rainv. ex Reichb. Fl. Germ. Excurs. 140* = Glyceria maritima.

POA:—
spicata, Willd. ex Steud. Nom. ed. II. ii. 362 = Eleusine indica.
spinosa, Thunb. Prod. Pl. Cap. 22 = Eragrostis spinosa.
spondyloides, Steud. Syn. Pl. Gram. 256 = sphondylodes.
sporoboloides, A. Rich. Tent. Fl. Abyss. ii. 426 = Eragrostis interrupta, *Beauv.*
Sprengelii, Kunth, Rév. Gram. i. 116 = Eragrostis imbecilla.
squamata, Lam. Illustr. i. 185 (Eragrostidis sp.).—Afr. trop.
squamosa, Beauv. Agrost. 176 = praec.
squarrosa, *Licht. ex Roem. & Schult. Syst.* ii. 553.—Afr. austr.
squarrosa, Salzm. ex Steud. Syn. Pl. Gram. 274 = Eragrostis reptans.
stelligera, *Desv. Opusc.* 101.—Ind. occ.
stenantha, Janka, in Oestr. Bot. Zeitschr. xiv. (1864) 383 = alpina.
stenantha, *Trin. in Mém. Acad. Sc. Pétersb.* Sér. VI. i. (1831) 376.—As. bor. or.; Am. bor. occ.
stenantha, Turcz. ex Steud. Syn. Pl. Gram. 253 = neglecta.
stenoclada, *Kunth, Enum. Pl.* i. 343.—Peruv.
stenostachya, R. Br. Prod. 181 = Eragrostis stenostachya.
sterilis, *Bieb. Fl. Taur. Cauc.* i. 62.—As. Min.; Tauria.
stolonifera, Bell. in Mém. Acad. Turin, v. 1790–91 (1793) 215 = alpina.
stolonifera, *Muhl. Desc. Gram.* i. 39.—Am. bor.
straminea, *Steud. in Lechl. Berb. Am. Austr.* 52.—Chili.
striata, Lam. Illustr. i. 183 = Lamarckii.
striata, Michx. Fl. Bor. Am. i. 389 = Glyceria nervata.
striata, Thunb. Prod. Pl. Cap. 22 = Eragrostis Bergiana.
stricta, *D. Don, in Mem. Wern. Soc.* iii. (1821) 298 (Quid?).—Scotia.
stricta, Hegetschw. & Heer, Fl. Schw. 85 = alpina.
stricta, *Lindeb. in Bot. Notiser*, 1855 (1856), 10.—Scandinav.
stricta, Roth, Nov. Pl. Sp. 96 = Eragrostis Brownei.
stricta, Willd. ex Spreng. Syst. i. 343 = Eragrostis acutiflora.
strictula, Steud. Syn. Pl. Gram. 426 = sphondylodes.
strigosa, Hoffm. Deutschl. Fl. ed. II. i. 44 = pratense.
subalpina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 87 = alpina.
subcaerulea, Sm. Engl. Bot. t. 1004 = pratensis.
subcompressa, Parn. Grass. Brit. 84 = compressa.
subfastigiata, Trin. ex Ledeb. Fl. Alt. i. 96 = Glyceria subfastigiata.
subnuda, J. F. Gmel. Syst. 179 = Festuca sylvatica.
subsecunda, Lam. Illustr. i. 184 = Eragrostis reptans.
subspicata, *Kunth, Enum. Pl.* i. 326.—Peruv.
subtilis, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 532 = soongarica.
subtilis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 86 = alpina.
subulata, Bosc, ex Kunth, Enum. Pl. i. 325 = Distichlis maritima.
subulata, *Desv. Opusc.* 102.—Afr. trop.
subumbellata, Roem. & Schult. Syst. ii. 570 = Orthoclada rariflora.
subuniflora, *Kunth, Rév. Gram.* i. 115.—Mexic.
subuniflora, Schur, Enum. Pl. Transs. 948 (Index), sphalm. = nemoralis.
subverticillata, *Pers. Syn.* i. 92.—Am. bor.
sudetica, Haenke, in Jirasek, Beob. Riesengeb. (1791) 120 = Chaixii.
sudetica, C. A. Mey. Verz. Pfl. Cauc. 19, partim = longifolia.
sudetica, Schleich. Cat. Nov. 20; Gaud. Agrost. Helv. i. 215 = pratensis.
sulcata, Lag. Gen. et Sp. Nov. 3 = commutata.
sulcata, Schult. Mant. ii. 305 = Lamarckii.
supina, Baumg. Mant. 79 (1846) = laxa.
supina, Panz. in Sturm, Deutschl. Fl. Heft 34 = minor.
supina, Schrad. Fl. Germ. i. 289 = annua.
suzumeokatabica, Herb. Lugd. Batav. ex Steud. Syn. Pl. Gram. 287 = Glyceria airoides.
sylvatica, Chaix, in Vill. Hist. Pl. Dauph. ii. 128 = Chaixii.
sylvatica, Pollich, Hist. Pl. Palat. i. 83 = Festuca sylvatica.
sylvestris, *A. Gray, Man. Bot. N. U. St.* ed. I. 596.—Am. bor.
sylvicola, Guss. Enum. Pl. Inar. 371 = pratensis.

POA:—
syrtica, F. Muell. in Trans. Vict. Inst. i. (1854) 45 = Glyceria stricta.
tatarica, *Fisch. in Bess. Cat. Krzem. Suppl.* ii. 13.—Rossia; Persia.
tenax, H. B. & K. Nov. Gen. et Sp. i. 160 = Eragrostis tenax.
tenax, Link, Enum. Hort. Berol. i. 87 = Festuca elatior.
tenella, R. Br. Prod. 181; Link, Enum. Hort. Berol. i. 87 = Eragrostis plumosa.
tenella, Hoppe. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 374 = minor.
tenella, Linn. Sp. Pl. 69 = Eragrostis plumosa?
tenella, Nutt. Gen. Am. i. 67 = Eragrostis Nuttalliana.
tenella, Pall. Reise, iii. 537 = Eragrostis pilosa.
tenella, Pursh, Fl. Am. Sept. i. 80 = Eragrostis Purshii.
tenella, Unio It. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 84 = Eragrostis Frankii.
tenella, *Vahl, ex Hornem. Hort. Hafn.* i. 93.—Ind. or.
tenellula, Kunth, Rév. Gram. i. 113 = Eragrostis plumosa.
tenera, F. Muell. ex Hook. f. Fl. Tasm. ii. 124. t. 164 = caespitosa.
tenuiculmis, *Steud. Syn. Pl. Gram.* 259.—Chili.
tenuiflora, *Licht. ex Roem. & Schult. Syst.* ii. 567.—Afr. austr.
tenuiflora, Rafin. in Med. Repos. N. York, v. (1808) 353, nomen (Quid?).—Am. bor.
tenuifolia, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 96 (= *Atropis californica*).—Am. bor. occ.
tenuifolia, Nutt. ex S. Wats. in Bot. King's Exp. 387 (Glyceriae sp.).—Am. bor. occ.
tenuifolia, *A. Rich. Tent. Fl. Abyss.* ii. 425.—Abyss.
tenuis, [Clairv.] Man. Herb. 18 = nemoralis.
tenuis, Ell. Sketch, i. 156 = Eragrostis tenuis.
tenuissima, Schrad. in Linnaea, xii. (1838) 447 = Eragrostis plumosa.
tephrosanthos, Spreng. in Schult. Mant. ii. 316 = Eragrostis tephrosanthos.
ternitana, Spreng. ex Link, Handb. i. 62, nomen = Aeluropus laevis.
thalassica, H. B. & K. Nov. Gen. et Sp. i. 157; Kunth, Enum. Pl. i. 326 = Distichlis thalassica.
thermalis, Pers. Syn. i. 90 = alpina.
thessala, Boiss. & Orph. Diagn. Ser. II. iv. 135 = pumila.
Thouarsiana, *Kunth, Rév. Gram.* ii. 477. t. 152.—Madag.
thyphoides, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 75 = Koeleria phleoides.
Timoleontis, *Heldr. ex Boiss. Fl. Orient.* v. 607.—Graecia.
Torreyana, Spreng. Neue Entdeck. ii. 104 = Glyceria elongata.
tortuosa, Spreng. Syst. i. 345 = Roxburghiana.
transsilvanica, Schur, Enum. Pl. Transs. 776 = alpina.
tremula, Lam. Illustr. i. 185 = Eragrostis tremula.
tremula, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 86 = minor.
triangularis, Gilib. Exercit. ii. 531 = annua.
trichoclada, Rupr. Beitr. Pfl. Russ. Reich. ii. 62. t. 4 = Laestadii.
trichodes, Nutt. in Trans. Am. Phil. Soc. v. (1837) 146 = Eragrostis tenuis.
trichoides, *Hornem. ex Steud. Nom.* ed. II. ii. 363.—Hab.?
trichophylla, *Boiss. Fl. Orient.* v. 604.—Graecia.
trichopoda, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 136, err. typ. = praec.
tricolor, *Nees, ex Steud. Syn. Pl. Gram.* 259.—Chili.
triflora, Gilib. Exercit. 531 = serotina.
triflora, Moench, Meth. 187 = Festuca spadicea.
trinervata, DC. Fl. Fr. iii. 58 = Chaixii.
trinervata, Ehrh. Beitr. vi. 81, 131; Schrad. Spicil. Fl. Germ. 3 = Festuca sylvatica.
tristigmatica, *E. Desv. in C. Gay, Fl. Chil.* vi. 419.—Chili.
tristis, Trin. in Bunge, Verz. Suppl. Fl. Alt. 10 = laxa.
tristriata, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 252 = Catabrosa humilis.
triticea, Kunth, Rév. Gram. i. 116 = Eragrostis triticea.

POA:—
triticoides, *Decne. in Ann. Sc. Nat.* Sér. II. iv. (1835) 84.—Arab.
trivialis, Guss. Fl. Sic. Prod. i. 89 = pratensis.
trivialis, Leers, Fl. Herborn. t. 6. f. 2 = nemoralis.
trivialis, *Linn. Sp. Pl.* 67.—Reg. bor. temp.
trivialis, Pollich, Hist. Pl. Palat. i. 84; Koel. Desc. Gram. 57 = alpina.
tumidula, *Steud. Syn. Pl. Gram.* 258.—Chili.
tunetana, Spreng. Pugill. ii. 20 = Aeluropus villosus.
turgida, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 66 = Eragrostis rubiginosa?
typhoides, Willk. & Lange, Prod. Fl. Hisp. iii. 1078 = Koeleria phleoides.
udensis, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 103).—Sibir. or.
uliginosa, Willd. ex Spreng. Syst. i. 342 = remota.
umbrosa, *Trin. in Mém. Acad Pétersb.* Sér. VI. i. (1831) 386.—Bras.
uniflora, Muhl. Desc. Gram. 151 = Sporobolus serotinus.
uninervia, *Kunth, Enum. Pl.* i. 344.—Mexic.
Uniolae, Schrad. in Goett. Gel. Anz. (1821) III. 2074; Schult. Mant. ii. 312 = Eragrostis Uniolae.
unioloides, Retz. Obs. v. 19 = Eragrostis unioloides.
unioloides, Willd. ex Spreng. Syst. i. 344 = Eragrostis amabilis.
urssulensis, *Trin. in Bunge, Verz. Suppl. Fl. Alt.* 10.—Mont. Altai.
Urvillei, *Steud, Syn. Pl. Gram.* 262.—Reg. Magell.?
vaginaefolia, Steud. in Lechl. Berb. Am. Austr. 52 = seq.
vaginaeformis, Steud. ex Phil. i. Cat. Pl. Vasc. Chil. 332, sphalm. = vaginiflora.
vaginalis, *Benth. Pl. Hartw.* 261.—N. Granat.
vaginiflora, *Steud. Syn. Pl. Gram.* 259.—Chili.
Vahliana, Liebm. in Fl. Dan. t. 2401 = Glyceria Vahliana.
Vahlii, Roem. & Schult. Syst. ii. 563 = Eragrostis Vahlii.
valdiviana, *Phil. in Linnaea*, xxix. (1857–58) 97.—Chili.
vallesiana, [Honck.] Verz. Aller Gew. Teutschl. 222 (Quid?).—Helvet.
variabilis, Gaudich. in Freyc. Voy. Bot. 408 = Eragrostis wahowensis.
variegata, Hall. f. in Schleich. Pl. Helv. 38 = annua.
variegata, Host, ex Kunth, Enum. Pl. i. 354 = nemoralis.
variegata, Lam. Illustr. i. 181 = alpina.
venusta, Roem. & Schult. Syst. ii. 573 = concinna.
versicolor, Bess. Enum. Pl. Volh. 41; Trin. in Mém. Acad. Petersb. Sér. VI. iv. II. (1838) 62 = sterilis.
verticillata, Cav. Ic. i. 63. t. 93 = Eragrostis pilosa.
vestita, *Bosc, ex Steud. Nom.* ed. II. ii. 363.—Am. bor.
Villarsii, J. F. Gmel. Syst. 182 = pratensis.
villosa, Beyr. ex Schrad. in Linnaea, xii. (1838) 451 = Eragrostis velutina.
violacea, *Bell. in Mém. Acad. Turin*, v. (1792) 214.—Europ. austr.
violascens, *Phil. in Linnaea*, xxix. (1857–58) 100.—Chili.
virens, Jacq. ex Kunth, Enum. Pl. i. 353 = viridis.
virescens, Kunth, l. c. 329 = Eragrostis virescens.
virgata, *Poir. Encyc.* v. 78.—Ins. S. Doming.
virgata, Roth, Nov. Pl. Sp. 66 = Leptochloa filiformis.
virginica, *Zuccagni, in Roem. Collect.* 124.—Am. bor.
viridiflora, *Hochst. in Flora*, xxxviii. (1855) 323.—Abyss.
viridis, Gilib. Exercit. ii. 530 = pratensis.
viridis, *Schreb. ex Pursh, Fl. Am. Sept.* i. 79.—Am. bor.
viscosa, Retz. Obs. iv. 20 = Eragrostis amabilis.
vivipara, Fl. Corc. in Ionios Anthol. iii. (1834) 686 = bulbosa.
vivipara, Willd. Enum. Hort. Berol. 103 = alpina.
wahowensis, Kunth, Enum. Pl. i. 328 = Eragrostis wahowensis.
Walkeri, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 224.—N. Zel.
Walteri, Kunth, Rév. Gram. i. 116 = Eragrostis conferta.
Weigeltiana, Reichb. ex Linn. in Mém. Acad. Sc. Pétersb. Sér. VI. i. (1831) 410 = Eragrostis reptans.

POA:—
Wheeleri, *Vasey, in Rothr. Rep. Bot. U. St. Surv.* vi. 291.—Am. bor. occ.
Willemetiana, Godfr. in Willem. Fl. Lorrain. i. 86 = Chaixii.
Zeae, *Roem. & Schult. Syst.* ii. 558.—N. Granat.

POARCHON, Allem. Tract. Rio Jan. (1849) = **Trimezia**, Salisb. (Irid.).
fluminensis, Allem. l. c. = T. lurida.

POARCHON, Mart. ex Seub. in Mart. Fl. Bras. iii. 1. 223 (1855) = **Abolboda**, Humb. & Bonpl. (Xyrid.).
coeruleus, Mart. l. c. = A. Poarchon.

POARION, Reichb. Consp. 51 (1828) = **Koeleria**, Pers. (Gramin.).

POARIUM, Desv. in Ham. Prod. Pl. Ind. Occ. 46 (1825) = **Stemodia**, Linn. (Scrophul.).
veronicoides, Desv. l. c. = S. parviflora ?

POCILLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 129 = **Veronica**, Tourn. (Scroph.).
agrestis, Fourr. l. c. = V. agrestis.
Cymbalaria, Fourr. l. c. = V. Cymbalaria.
hederaefolia, Fourr. l. c. = V. hederaefolia.
persica, Fourr. l. c. = V. Tournefortii.
polita, Fourr. l. c. = V. agrestis.

POCOCKIA, Ser. in DC. Prod. ii. 185 (1825) = **Trigonella**, Linn. (Legumin.).
arabica, Boiss. Diagn. Ser. I. ix. 13 = T. arabica.
Aucheri, Boiss. l. c. 12 = T. brachycarpa.
biflora, Walp. Ann. ii. 347 = T. lunata.
cachemirica, Boiss. Diagn. Ser. I. ix. 12 = T. Emodi.
cilicica, Boiss. & Bal. l. c. Ser. II. v. 76 = T. brachycarpa.
cretica, Ser. in DC. Prod. ii. 185 = T. cretica.
elliptica, Boiss. Diagn. Ser. I. ix. 12 = T. elliptica.
glomerata, Boiss. l. c. = T. brachycarpa.
graeca, Boiss. & Sprun. l. c. ii. 16 = T. graeca.
Kotschyi, Boiss. l. c. ix. 12 = T. rytidocarpa.
lineata, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 54 = T. elliptica.
lunata, Boiss. Diagn. Ser. I. ix. 12 = T. lunata.
ovalis, Boiss. l. c. = Medicago ovalis.
platycarpos, Boiss. l. c. = Medicago platycarpa.
plicata, Boiss. & Bal. l. c. Ser. II. v. 76 = T. plicata.
radiata, Trautv. in Bull. Sc. Acad. Pétersb. viii. (1841) 271 = T. radiata.
rostrata, Boiss. & Bal. Diagn. Ser. II. vi. 43 = T. rostrata.
ruthenica, Boiss. l. c. Ser. I. ix. 12 = Medicago ruthenica.

POCOPHORUM, Neck. Elem. ii. 226 (1790) = **Rhus**, Linn. (Anacard.).

PODACHAENIUM, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 98. *COMPOSITAE*, Benth. & Hook. f. ii. 380.
COSMOPHYLLUM, C. Koch, Ind. Sem. Hort. Berol. (1854) 12.
DICALYMMA, Lem. Illustr. Hortic. ii. (1854) Misc. 37.
alatum, Walp. Ann. v. 230, sphalm. = paniculatum.
eminens, Baill. Hist. Pl. viii. 206 = paniculatum.
paniculatum, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 99.—Mexic.

PODADENIA, Thw. Enum. Pl. Zeyl. 273 (1861). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 318.
sapida, *Thw. l. c.* 274.—Zeylan.
Thwaitesii, Muell. Arg. in DC. Prod. xv. II. 791 = sapida.

PODAGRARIA, Hall. Enum. Stirp. Helv. ii. 427 (1742) = **Aegopodium**, Linn. (Umbell.).
Aegopodium, Moench, Meth. 90 = A. Podagraria.

PODALETRA, Rafin. Fl. Tellur. iv. 79 (1836) = **Convolvulus**, Tourn.
farinosa, Rafin. l. c. = C. farinosus.

PODALIRIA, Willd. Enum. Hort. Berol. 437 (1809) = seq.

PODALYRIA, Lam. Illustr. ii. 454. t. 327. f. 3, 4 (1793). *LEGUMINOSAE*, Benth. & Hook. f. i. 466.
alba, Sims, Bot. Mag. t. 1177 = Baptisia leucantha.
alba, Willd. Sp. Pl. ii. 503 = Baptisia alba.
albens, E. Mey. in Linnaea, vii. (1832) 146 = cuneifolia.
alpina, Willd. Enum. Hort. Berol. Suppl. 22 = Thermopsis corgonensis.
amoena, Eckl. & Zeyh. Enum. 155 = canescens.
angustifolia, Eckl. & Zeyh. l. c. 158 = argentea.
anomala, Lehm. ex Harv. & Sond. Fl. Cap. ii. 14 = sericea.
argentea, Eckl. & Zeyh. Enum. 157 = biflora.
argentea, *Salisb. Parad. Lond.* t. 7.—Afr. austr.
argentea, Willd. Sp. Pl. ii. 502 = Ammodendron Sieversii.
aurea, Willd. l. c. = Calpurnia lasiogyne.
australis, Willd. l. c. 503; Vent. Jard. Cels, t. 56 = Baptisia australis.
biflora, *Lam. Illustr.* ii. 471.—Afr. austr.
biflora, Sims, Bot. Mag. t. 753 = argentea.
bracteata, Muhl. Cat. ed. II. 42 = Baptisia leucophaea.
bracteata, Roxb. Hort. Beng. [31]; Fl. Ind. ii. 317 = Dalhousiea bracteata.
Burchellii. *DC. Prod.* ii. 101.—Afr. austr.
buxifolia, Eckl. & Zeyh. Enum. 157 = myrtillifolia.
buxifolia, Lam. Illustr. ii. 471, t. 327. f. 4 = glauca.
buxifolia, *Willd. Sp. Pl.* ii. 505.—Afr. austr.
calyptrata, *Willd. l. c.* 504.—Afr. austr.
cana, Willd. ex Walp. in Linnaea, xiii. (1839) 456 = Burchellii.
canescens, Eckl. & Zeyh. Enum. 158 = sericea.
canescens, *E. Mey. Comm. Pl. Afr. Austr.* 9.—Afr. austr.
capensis, Andr. Bot. Rep. t. 347 = Virgilia capensis.
coerulea, Michx. Fl. Bor. Am. i. 264 = Baptisia australis.
cordata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 8.—Afr. austr.
coruscans, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 171.—Afr. austr.
cuneifolia, Eckl. & Zeyh. Enum. 156, partim = argentea.
cuneifolia, *Vent. Hort. Cels*, t. 99.—Afr. austr.
genistoides, Willd. Sp. Pl. ii. 502 = Cyclopia galioides.
glauca, *DC. Prod.* ii. 102.—Afr. austr.
glauca, Eckl. & Zeyh. Enum. 157 = myrtillifolia.
Haematoxylon, Thonn. in Schum. & Beskr. Guin. Pl. 202 = Baphia nitida.
hamata, E. Mey. in Linnaea, vii. (1832) 146 = cuneifolia.
hirsuta, Willd. Sp. Pl. ii. 505 = cordata.
intermedia, Eckl. & Zeyh. Enum. 155 = canescens.
Kunthii, *Walp. in Linnaea*, xiii. (1839) 460.—Afr. austr.
laevigata, Willd. ex Walp. l. c. 453 = Cyclopia latifolia.
lanceolata, *Benth. in Ann. Wien. Mus.* ii. (1838) 68.—Afr. austr.
lanceolata, Poir. Encyc. x. 445 = Baptisia lanceolata.
lancifolia, Eckl. & Zeyh. Enum. 156 = Burchellii.
liparioides, DC. Prod. ii. 102 = biflora.
liparioides, Eckl. & Zeyh. Enum. 158 = argentea.
lupinoides, Willd. ex Cham. in Linnaea, vi. (1831) 545 = Thermopsis corgonensis.
lupinoides, Willd. Sp. Pl. ii. 504 = Thermopsis lanceolata.
Meyeriana, Eckl. & Zeyh. Enum. 155 = canescens.
microphylla, *E. Mey. in Linnaea*, vii. (1832) 147.—Afr. austr.
mollis, Michx. Fl. Bor. Am. i. 264 = Thermopsis mollis.
monosperma, Poir. Encyc. v. 440 = Ormosia dasycarpa.
Mundiana, Eckl. & Zeyh. Enum. 156 = glauca.
mutabilis, *Jacq. ex Steud.* Nom. ed. I. 639; ed. II. ii. 364.—Hab. ?
myrtifolia, D. Dietr. Syn. Pl. ii. 1498 = myrtillifolia.
myrtillifolia, Eckl. & Zeyh. Enum. 156 = calyptrata.
myrtillifolia, *Willd. Sp. Pl.* ii. 505.—Afr. austr.
obcordata, Poir. Encyc. v. 445 = Tephrosia obcordata.
oleaefolia, *Salisb. Parad. Lond.* t. 114.—Afr. austr.
orbicularis, *E. Mey. Comm. Pl. Afr. Austr.* 8; *Eckl. & Zeyh. Enum.* 156.—Afr. austr.
pallens, Eckl. & Zeyh. l. c. 158 = cuneifolia.
parvifolia, DC. Prod. ii. 102; Benth. in Ann. Wien Mus. ii. (1838) 68 = myrtillifolia.

PODALYRIA:—
patens, Eckl. & Zeyh. Enum. 159 = cuneifolia.
pedunculata, Eckl. & Zeyh. l. c. 158 = biflora.
perfoliata, Michx. Fl. Bor. Am. i. 263 = Baptisia perfoliata.
racemiflora, Harv. in Harv. & Sond. Fl. Cap. ii. 618 = glauca.
racemulosa, *DC. Prod.* ii. 102.—Afr. austr.
racemulosa, Eckl. & Zeyh. Enum. 157 = glauca.
reticulata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 11.—Afr. austr.
sericea, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Afr. austr.
sparsiflora, Eckl. & Zeyh. Enum. 157 = glauca.
speciosa, *Eckl. & Zeyh. l. c.*—Afr. austr.
splendens, Eckl. & Zeyh. l. c. 159 = cuneifolia.
styracifolia, Sims, Bot. Mag. t. 1580 = calyptrata.
subbiflora, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 439 = argentea.
subbiflora, DC. Prod. ii. 102 = biflora.
ternata, Willd. Sp. Pl. ii. 503 = Aspalathus ferruginea.
tetraptera, Hort. ex Steud. Nom. ed. I. 640 = Sophora tetraptera.
Thunbergiana, Eckl. & Zeyh. Enum. 155 = canescens.
tinctoria, Lam. Illustr. ii. 471 = Baptisia tinctoria.
trifoliata, Willd. Sp. Pl. ii. 504 = Argyrolobium sericeum.
uniflora, Michx. Fl. Bor. Am. i. 263 = Baptisia lanceolata.
velutina, *Burch. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 437.—Afr. austr.
villosa, Michx. Fl. Bor. Am. i. 264 = Baptisia villosa.
virgata, Willd. ex Walp. in Linnaea, xiii. (1839) 457 = myrtillifolia.

PODANTHERA, Wight, Ic. v. 22. t. 1759 (1851) = **Epipogum**, S. G. Gmel. (Orchid.).
pallida, Wight, l. c. = E. nutans.

PODANTHES, Haw. Syn. Pl. Succ. 32 (1812) = **Stapelia**, Linn. (Asclep.).
ciliata, Haw. l. c. 34 = Diplocyatha ciliata.
incarnata, Sweet, Hort. Brit. ed. II. 358 = Boucerosia incarnata.
irrorata, Haw. Syn. Pl. Succ. 33 = S. irrorata.
lepida, Haw. l. c. 34 = S. lepida.
pulchella, Haw. l. c. 33 = S. pulchella.
pulchra, Haw. l. c. 32 = S. roriflua.
roriflua, G. Don, Gen. Syst. iv. 118 = S. roriflua.
verrucosa, Haw. Syn. Pl. Succ. 33 = S. verrucosa.

PODANTHUM, Boiss. Fl. Orient. iii. 945 (1875) = **Phyteuma**, Linn. (Campan.).
amplexicaule, Boiss. l. c. 948 = Phyt. amplexicaule.
anthericoides, Janka, in Termész. Füzetek, ii. (1878) 30 = Phyt. lobelioides ?
asperum, Boiss. Fl. Orient. iii. 955 = Phyt. asperum.
brachylobum, Boiss. l. c. 947 = Campanula sessiliflora.
campanuloides, Boiss. l. c. 949 = Phyt. campanuloides.
canescens, Boiss. l. c. 950 = Phyt. canescens.
cappadocicum, Boiss. l. c. 946 = Phyt. cappadocicum.
cichoriforme, Boiss. l. c. 947 = Phyt. cichoriforme.
controversum, Boiss. l. c. 949 = Phyt. rigidum.
giganteum, Boiss. l. c. 946 = Phyt. giganteum.
lanceolatum, Boiss. l. c. 951 = Phyt. lanceolatum.
limonifolium, Boiss. l. c. = Phyt. limonifolium.
linifolium, Boiss. l. c. 954 = Phyt. linifolium.
lobelioides, Boiss. l. c. 953 = Phyt. lobelioides.
Lycium, Boiss. l. c. 954 = Phyt. Lycium.
macrodon, Boiss. & Haussk. ex Boiss. l. c. 956.—Persia.
Otites, Boiss. l. c. 952 = Phyt. Otites.
persicum, Boiss. l. c. 956.—Persia.
psilostachyum, Boiss. l. c. 948 = Campanula psilostachya.
pulchellum, Boiss. l. c. 947 = Phyt. pulchellum.
scoparium, Boiss. & Haussk. ex Boiss. l. c. 953.—Cataonia.
Sibthorpianum, Boiss. l. c. 950 = Phyt. Sibthorpianum.
strigillosum, Boiss. l. c. 957 = Campanula strigillosa.
tenuifolium, Boiss. l. c. 952 = Phyt. tenuifolium.
trichocalycinum, Boiss. l. c. 955 = Phyt. amplexicaule.
trichostegium, Boiss. l. c. 954.—Cappadoc.
virgatum, Boiss. l. c. 946 = Campanula Rapunculus.

PODANTHUS, Lag. Gen. et Sp. Nov. 24 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 356.
EUXENIA, Cham. in Nees, Hor. Phys. Berol. 75. t. 16 (1820).

PODANTHUS:—
gratus, D. Don, ex Sweet, Hort. Brit. ed. III. 353 = ovatifolius.
Mitiqui, *Lindl. in Loud. Hort. Brit.* 488.—Chili.
ovatifolius, *Lag. Gen. et Sp. Nov.* 24.—Bras.; Chili.

PODASAEMUM, Reichb. Consp. 50 (1828) = PODOSEMUM, Desv. = **Muehlenbergia**, Schreb. (Gramin.).

PODIA, Neck. Elem. i. 72 (1790) = **Centaurea**, Linn. (Compos.).

PODIANTHUS, Schnitzl. in Bot. Zeit. i. (1843) 739 = **Trichopus**, Gaertn. (Dioscor.).
arifolius, Schnitzl. l. c. = T. zeylanicus.

PODIONAPUS, Dulac, Fl. Hautes-Pyr. 82 (1867) = **Deschampsia**, Beauv. (Gramin.).
caespitosus, Dulac, l. c. 82 = D. caespitosa.
flexuosus, Dulac, l. c. 83 = D. flexuosa.

PODIOPETALUM, Hochst. in Flora, xxiv. (1841) 657 = **Dalbergia**, Linn. (Legumin.).
reticulatum, Hochst. l. c. 658 = D. obovata.

PODISONIA, Dum. ex Steud. Nom. ed. II. ii. 364 (1841), sphalm. = **Posidonia**, Koen. (Naiad.).

PODOCALLIS, Salisb. Gen. Pl. Fragm. 17 (1866) = **Massonia**, Thunb. (Liliac.).
nivea, Salisb. l. c. 17 = M. jasminiflora?

PODOCALYX, Klotzsch, in Wiegm. Archiv, vii. (1841) 202 = **Richeria**, Vahl (Euphorb.).
loranthoides, Klotzsch, l. c. = R. loranthoides.

PODOCARPUS, [L'Hérit.] Pers. Syn. ii. 580 (1807).
CONIFERAE, Benth. & Hook. f. iii. 435.
NAGEIA, Gaertn. Fruct. i. 191. t. 39 (1788), partim.
PRUMNOPITYS, Phil. in Linnaea, xxx. (1859–60) 731.
acicularis, Hort. ex Carr. Conif. ed. I. 456 = elata.
acutifolia, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 370.—N. Zel.
affinis, *Seem. Fl. Vit.* 266.—Ins. Fiji.
agathifolia, Blume, Rumphia, iii. 217 = latifolia.
alba, J. Baum, Cat. (1846); ex Henk. & Hochst. Syn. Nadelh. 404 (Quid?).—Hab.?
alpina, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Austral.
amara, *Blume, Enum. Pl. Jav.* 88.—Java.
andina, *Poepp. ex Endl. Syn. Conif.* 219.—Chili.
angustifolia, *Griseb. Cat. Pl. Cub.* 217.—Cuba.
angustifolia, *Parl. in DC. Prod.* xvi. II. 512.—Bolivia.
antarctica, Hort. ex Carr. Conif. ed. I. 450 = curvifolia.
antarctica, Hort. ex Carr. l. c. ed. II. 647 = Purdieana?
antillarum, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 75 = coriacea.
appressa, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 379.—Japon.
araucarioides, *Brongn. & Gris, ex Panch. & Sebert, Not. Bois N. Caléd.* 171.—N. Caled.
argotaenia, *Hance, in Journ. Bot.* xxi. (1883) 357.—China.
aristulata, *Parl. in DC. Prod.* xvi. II. 513.—Cuba.
aspleniifolia, Labill. Nov. Holl. Pl. ii. 71. t. 221 = Phyllocladus rhomboidalis.
Beccarii, *Parl. in DC. Prod.* xvi. II. 508.—Borneo.
Bidwilli, Hoibrenk, ex Endl. Conif. 213 = P. Totara.
biformis, Hook. Ic. Pl. t. 544 = Dacrydium Colensoi.
Blumei, *Endl. Syn. Conif.* 208.—Java.
bracteata, Blume, Enum. Pl. Jav. 88 = neriifolia.
bracteata, Hort. ex Carr. Conif. ed. II. 665 = macrophylla.
bracteata, Zoll. ex Parl. in DC. Prod. xvi. II. 516 = amara.
caesia, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 379.—Japon.
caespitosa, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xiii. (1866) 425 = alpina.
canaliculata, Hort. ex Carr. Conif. ed. II. 659 = macrophylla.

PODOCARPUS:—
chilena, Lechler, ex Parl. in DC. Prod. xvi. II. 510 = oleifolia.
chilina, *Rich. in Ann. Mus. Par.* xvi. (1810) 297, nomen; *et Conif.* 11.—Chili.
chinensis, Sweet, Hort. Brit. ed. I. 371; Wall. Cat. n. 6051 = macrophylla.
coreana, Hort. ex Carr. Conif. ed. II. 724 = Torreya nucifera.
coriacea, Hort. ex Carr. l. c. ed. I. 512 = Torreya nucifera.
coriacea, *Rich. Conif.* 14. t. 1. f. 3.—N. Granat.; Ind. occ.
corrugata, *Gord. Pinet. Suppl.* 88.—Japon.
costalis, *Presl, Epim. Bot.* 236.—Ins. Philipp.
Cumingii, *Parl. in DC. Prod.* xvi. II. 521.—Ins. Philipp.
cupressina, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Burma; Malaya.
curvifolia, *Carr. Conif.* ed. I. 450.—Hab.?
cuspidata, *Endl. Syn. Conif.* 207.—Japon.
cuspidata, Hort. ex Carr. Conif. ed. I. 465 = amara.
dacrydioides, *A. Rich. Ess. Fl. N. Zél.* 358. t. 39.—N. Zel.
Dieffenbachii, Hook. Ic. Pl. t. 547 = Veronica tetragona.
discolor, *Blume, Rumphia*, iii. 213.—Java.
Drouyniana, *F. Muell. Fragm.* iv. 86. t. 31.—Austral.
drupacea, Hort. ex Gord. Pinet. 45 = Cephalotaxus drupacea.
dulcamara, *Seem. in Bonplandia*, ix. (1861) 253.—Hab.?
elata, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Austral.
elata, Seem. ex Parl. in DC. Prod. xvi. II. 517 = affinis.
elongata, *L'Hérit. ex Pers. Syn.* ii. 580.—Afr. austr.
elongata, E. Mey. ex Endl. Syn. Conif. 218 = Meyeriana.
Endlicheriana, Carr. Conif. ed. I. 468 = neriifolia?
ensifolia, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 75 = elata.
eurhyncha, *Miq. Fl. Ind. Bat.* ii. 1074.—Sumatra.
excelsa, Lodd. Cat. (1836) 37; ex Endl. Syn. Conif. 213 = spinulosa.
falcata, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Afr. austr.
falcata, A. Cunn. ex Parl. in DC. Prod. xvi. II. 517 = elata.
falciformis, *Parl. l. c.* 685.—Borneo.
ferruginea, *G. Benn. ex D. Don, in Lamb. Gen. Pin.* ed. II. (1832) ii. [189].—N. Zel.
flagelliformis, *Hort. Belg. ex Carr. Conif.* ed. II. 666.—Japon.
glomerata, *D. Don, in Lamb. Gen. Pin.* ed. I. (1824) ii. 21; ed. Min. (1832) 141. n. 73.—Peruv.
gnidioides, *Carr. Conif.* ed. II. 656.—N. Caled.
grandifolia, *Endl. Syn. Conif.* 208.—Japon.
Horsfieldii, R. Br. in Wall. Cat. n. 6049 = cupressina.
Humboldti, Hort. ex Carr. Conif. ed. I. 451 = curvifolia.
Humboldti, Hort. ex Gord. Pinet. 288 = taxifolia.
imbricata, Blume, Enum. Pl. Jav. 89 = cupressina.
insignis, Hemsl. in Journ. Bot. xxiii. (1885) 287 = argotaenia.
jamaicaënsis, Hort. ex Gord. Pinet. 281 = Purdieana.
japonica, *Hort. Bogor. ex Siebold, in Ann. Soc. Hort. Pays-Bas* (1844), 35, 36.—Japon.
Junghuhniana, Miq. Pl. Jungh. 2 = neglecta.
koraiana, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 499 = Cephalotaxus pedunculata.
koraiana, C. Koch, in Berl. Allg. Gartenz. i. (1857) 228 = macrophylla.
koraiana, *Siebold, in Ann. Soc. Hort. Pays-Bas* (1844), 35.—Corea.
koreana, Hort. ex Parl. in DC. Prod. xvi. II. 516 = chinensis.
laeta, *Hoibrenk, ex Endl. Syn. Conif.* 214.—Austral.
Lamberti, *Klotzsch, ex Endl. l. c.* 211.—Bras.
lanceolata, Hort. ex Gord. Pinet. 275 = japonica.
latifolia, Blume, Enum. Pl. Jav. 89 = Blumei.
latifolia, R. Br. in Benn. Pl. Jav. Rar. 40 = Thunbergii.
latifolia, Hort. ex Carr. Conif. ed. I. 439 = cuspidata.
latifolia, *Wall. Pl. As. Rar.* i. 26. t. 30.—Ind. or.; Malaya.
Lawrencii, Hook. f. in Hook. Lond. Journ. Bot. iv. (1845) 151 = alpina.

PODOCARPUS:—
leptostachya, *Blume, Rumphia*, iii. 214.—Borneo.
linearis, Van Houtte, ex Gord. Pinet. 273 = elongata.
longifolia, Hort. ex Carr. Conif. ed. II. 665 = japonica.
longifolia, Hort. ex Gord. Pinet. 278 = macrophylla.
macrophylla, *D. Don, in Lamb. Gen. Pin.* ed. I. (1824) ii. 22; ed. Min. (1832) 143.—Japon.
macrophylla, Sweet, Hort. Brit. ed. I. 371 = Sweetii.
macrophylla, Wall. Cat. n. 6052 = neriifolia.
macrostachya, *Parl. in DC. Prod.* xvi. II. 510.—N. Granat.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 447.—Madag.
Maki, Sieb. & Zucc. Fl. Jap. Fam. Nat. ii. 108 = chinensis.
Makoya, Hort. ex Endl. Syn. Conif. 217 = chinensis.
Makoyae, Hort. ex Carr. Conif. ed. I. 457 = chinensis.
Makoyi, *Blume, Rumphia*, iii. 215.—Japon.
Makoyi, Hort. ex Parl. in DC. Prod. xvi. II. 516 = chinensis.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 218.—Afr. occ.
Matai, Lamb. ex Hook. f. Handb. N. Zeal. Fl. 741 = spicata.
Meyeriana, *Endl. Syn. Conif.* 218.—Afr. austr.
minor, *Parl. in DC. Prod.* xvi. II. 509.—N. Caled.
Miquelia, Hort. ex Parl. l. c. 516 = chinensis.
montana, Lodd. Cat. (1836) 37; ex Endl. Conif. 219 = taxifolia.
mucronata, Hort. ex Carr. Conif. ed. I. 446 = Purdieana.
mucronata, Hort. ex Carr. l. c. ed. II. 664 = macrophylla.
Nageia, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Japon.
neglecta, *Blume, Rumphia*, iii. 213.—Java.
neriifolia, *D. Don, in Lamb. Gen. Pin.* ed. I. (1824) ii. 21; ed. min. (1832) 142.—Reg. Himal.
nivalis, *Hook. Ic. Pl.* t. 582.—N. Zel.
nobilis, Hort. ex Carr. Conif. ed. I. 468 = neriifolia.
nobilis, Hort. ex Endl. Syn. Conif. 218 = Thunbergii.
Novae-Caledoniae, *Vieill. ex Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. vi. (1866) 242.—N. Caled.
nubicola, Makoy, ex Henk. & Hochst. Syn. Nadelh. 385 = seq.
nubigena, *Lindl. in Journ. Hort. Soc.* vi. (1851) 264.—Chili.
nucifera, Loud. Hort. Brit. 388 = Torreya nucifera.
oleifolia, *D. Don, in Lamb. Gen. Pin.* ed. I. (1824) ii. 20; ed. min. (1832) 140.—Peruv.
ovata, *Henk. & Hochst. Syn. Nadelh.* 381.—Japon.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 589.—Sumatra.
Pardii, J. Baum. Cat. (1846); ex Henk. & Hochst. Syn. Nadelh. 404, sphalm. = Purdieana.
parvifolia, *Parl. in DC. Prod.* xvi. II. 514.—Austral.
pectinata, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xvi. (1869) 330.—N. Caled.
philippeana, Benth. ex Parl. in DC. Prod. xvi. II. 375 = Agathis loranthifolia.
pinnata, Hort. ex Gord. Pinet. 138 = latifolia.
polystachya, R. Br. in Benn. Pl. Jav. Rar. 40 = neriifolia.
pruinosa, E. Mey. in Drège, Zwei Pfl. Docum. 212 = elongata.
pungens, Caley, ex D. Don, in Lamb. Gen. Pin. ed. I. (1824) ii. 23; ed. Min. (1832) 144 = spinulosa.
pungens, Van Houtte, ex Gord. Pinet. 285 = P. Totara.
Purdieana, *Hook. Ic. Pl.* t. 624.—Jamaic.
Purdieana, Hort. ex Carr. Conif. ed. I. 445 = salicifolia.
Purdieana, Hort. ex Carr. l. c. 451 = curvifolia.
rigida, Klotzsch, in Endl. Syn. Conif. 211 = glomerata.
rivularis, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xiii. (1866) 425 = Novae-Caledoniae.
Rumphii, *Blume, Rumphia*, iii. 214.—Ins. Molucc.
salicifolia, *Klotzsch & Karst. ex Endl. Syn. Conif.* 209.—N. Granat.
saligna, D. Don, in Lamb. Gen. Pin. ed. I. (1824) ii. 20; ed. Min. (1832) 139 = chilina.
Sellowii, *Klotzsch, ex Endl. Conif.* 209.—Bras.
sinensis, Teijsm. & Binn. Cat. Hort. Bog. 14 = chinensis.
spicata, *R. Br. in Benn. Pl. Jav. Rar.* 40.—N. Zel.
spicata, Poepp. & Endl. Nov. Gen. et Sp. iii. 18 = andina.

PODOCARPUS:—
spinulosa, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 75.—Austral.
spinulosa, Hort. ex Carr. Conif. ed. I. 463 = macrophylla.
spinulosa, Hort. ex Carr. l. c. ed. II. 652 = P. Totara.
Sprengelii, Blume, in Flora, vii. (1824) 292 = Rumphii.
Sprucei, *Parl. in DC. Prod.* xvi. II. 510.—Ecuador.
Sweetii, *Presl, Bot. Bemerk.* 110.—Afr. austr.
taxifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 2. t. 97.—Reg. Andin.
taxodioides, Carr. Conif. ed. II. 657 = Dacrydium taxoides.
tenuifolia, *Parl. in DC. Prod.* xvi. II. 521.—N. Caled.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 1072.—Sumatra.
thevetiaefolia, *Zipp. in Flora*, xii. (1829) 287, nomen; *Blume, Rumphia*, iii. 213.—N. Guin.
thujoides, R. Br. in Benn. Pl. Jav. Rar. 41 = dacrydioides.
Thunbergii, *Hook. in Lond. Journ. Bot.* i. (1842) 657. t. 22.—Afr. austr.
Thunbergii, Siebold, in Verh. Batav. Gen. xii. (1830) 384 = macrophylla.
Totara, *G. Benn. ex D. Don, in Lamb. Gen. Pin.* ed. II. (1832) [189].—N. Zel.
usta, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. vi. (1866) 243.—N. Caled.
valdiviana, [Nelson] Senilis, Pinaceae, 160 = *andina*.
variegatus, Hort.—Cf. Gard. Chron. (1861) 735 = P. Totara.
verticillata, Hort. ex G. Don, in Loud. Hort. Brit. 388 = macrophylla.
verticillata, Wall. ex Steud. Nom. ed. II. ii. 365 = Sciadopitys verticillata.
Vieillardii, *Parl. in DC. Prod.* xvi. II. 521.—N. Caled.
vitiensis, *Seem. in Bonplandia*, x. (1862) 366.—Ins. Fiji.
Vriesiana, Hort. ex Parl. in DC. Prod. xvi. II. 516 = macrophylla.
Wallichiana, Presl, Bot. Bemerk. 110 = latifolia.
Yacca, G. Don, in Loud. Hort. Brit. 388 = coriacea.
Yacca, G. Don, in Sweet, Hort. Brit. ed. III. 622 = taxifolia.
zamiaefolia, Hort. ex Carr. Conif. ed. I. 441 = latifolia.
zamiaefolia, A. Rich. Ess. Fl. N. Zél. 360 = Dammara australis.

PODOCENTRUM, Burch. ex Meissn. in Linnaea, xiv. (1840) 489 = **Emex**, Neck. (Polygon.).

PODOCHILUS, Blume, Bijdr. 295. t. 12 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 588.
APISTA, Blume, Bijdr. 296 (1825).
CRYPTOGLOTTIS, Blume, l. c. 296. t. 42 (1825).
HEXAMERIA, R. Br. in Benn. Pl. Jav. Rar. 26. t. 7 (1838).
PLACOSTIGMA, Blume, Fl. Jav. Praef. p. vi (1828).
PLATYSMA, Blume, Bijdr. 295. t. 43 (1825).
bicolor, Miq. Choix Pl. Buitenz. t. 22. f. 3 = Dendrobium subulatum.
cultratus, *Lindl. in Wall. Cat.* n. 7336.—Reg. Himal.; Burma.
densiflorus, *Blume, Rumphia*, iv. 44.—N. Guin.
falcatus, *Lindl. Gen. et Sp. Orch.* 234.—Zeylan.
gracilis, *Lindl. l. c.* 235.—Java.
lucescens, *Blume, Bijdr.* 295. t. 12.—Burma; Malaya.
malabaricus, *Wight, Ic.* v. t. 1748. f. 2.—Ind. or.
microphyllus, *Lindl. in Wall. Cat.* n. 7335.—Burma; Malaya.
muricatus, Teijsm. & Binn. Cat. Hort. Bog. 379 = scalpelliformis.
saxatilis, *Lindl. Gen. et Sp. Orch.* 235.—Zeylan.
scalpelliformis, *Blume, Rumphia*, iv. 45.—Ins. Molucc.; N. Guin.
sciuroides, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
similis, *Blume, Rumphia*, iv. 44.—Borneo.
tenuis, *Lindl. Gen. et Sp. Orch.* 235.—Java.
Zollingeri, *Miq. Choix Pl. Buitenz.* t. 25. f. 2 (an seq.?).—Java.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 42.—Java.

PODOCHREA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364 = **Astragalus**, Tourn. (Legumin.).
incana, Fourr. l. c. = A. incanus.
monspessulana, Fourr. l. c. = A. monspessulanus.

PODOCOCCUS, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 426. *PALMAE*, Benth. & Hook. f. iii. 915.
Barteri, *Mann & H. Wendl. l. c.*—Afr. trop.

PODOCOMA, Cass. in Bull. Soc. Philom. (1817) 137; et in Dict. Sc. Nat. xlii. 60 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 278.
ASTEROPSIS, Less. Syn. Comp. 188 (1832).
IXIOCHLAMYS, F. Muell. & Sond. in Linnaea, xxv. (1852) 466.
MORITZIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 279 (1873).
PODOPAPPUS, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 50.
bellidifolia, *Baker, in Mart. Fl. Bras.* vi. III. 16.—Bras.
Blanchetiana, *Baker, l. c.* 15.—Bras.
cuneifolia, *R. Br. in Sturt, Exped. Cent. Austral.* ii. *App.* 81.—Austral.
erigerifolia, Steud. Nom. ed. II. i. 584, in syn. sphalm. = primulaefolia.
glandulosa, *Baker, in Mart. Fl. Bras.* vi. III. 14, in nota.—N. Granat.
hieracifolia, *Cass. in Dict. Sc. Nat.* xlii. 60.—Am. austr.
hirsuta, *Baker, in Mart. Fl. Bras.* vi. III. 15.—Bras.
primulaefolia, *Cass. in Dict. Sc. Nat.* xlii. 61.—Am. austr.
Regnellii, *Baker, in Mart. Fl. Bras.* vi. III. 16.—Bras.

PODOCYTISUS, Boiss. & Heldr. Diagn. Ser. I. ix. 7 (1849) = **Laburnum**, Linn. (Legumin.).
caramanicus, Boiss. l. c. = L. caramanicum.

PODOGYNE, Hoffmgg. Verz. Pfl. 101, 185 (1824) = **Gynandropsis**, DC. (Capparid.).
pentaphylla, Hoffmgg. l. c. 185 = D. pentaphylla.

PODOLASIA, N. E. Br. in Gard. Chron. (1882) II. 70. *AROIDEAE*, Benth. & Hook. f. iii. 996.
stipitata, *N. E. Br. l. c.*—Borneo.

PODOLEPIS, Labill. Nov. Holl. Pl. ii. 56. t. 208 (1806). *COMPOSITAE*, Benth. & Hook. f. ii. 328.
PANAETIA, Cass. in Ann. Sc. Nat. Sér. I. xvii. (1829) 417; et in Dict. Sc. Nat. lx. 593 (1830).
RUTIDOCHLAMYS, Sond. in Linnaea, xxv. (1852) 497.
SCALIA, Sims, Bot. Mag. t. 956 (1806).
SCALIOPSIS, Walp. in Linnaea, xiv. (1840) 318.
SIEMSSENIA, Steetz, in Lehm. Pl. Preiss. i. 467 (1845).
STYLOLEPIS, Lehm. Sem. Hort. Hamb. (1828) 17.—Cf. Flora, xii. (1829) I. Erg. 55.
acuminata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 82.—Austral.
affinis, Sond. in Linnaea, xxv. (1852) 507 = canescens.
aristata, *Benth. in Enum. Pl. Hueg.* 64.—Austral.
auriculata, *DC. Prod.* vi. 162.—Austral.
canescens, *A. Cunn. ex DC. l. c.* 163.—Austral.
chrysantha, Endl. in Bot. Zeit. i. (1843) 458 = aristata.
contorta, *Lindl. in Bot. Reg.* (1838) *Misc.* 64.—Tasmania.
divaricata, A. Cunn. ex DC. Prod. vi. 151 = Gnephosis tenuissima.
ferruginea, DC. l. c. 163 = Helipterum citrinum.
filiformis, Steetz, in Lehm. Pl. Preiss. i. 465 = gracilis.
Gilberti, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 195 = Lessoni.
gracilis, *R. Grah. in Edinb. N. Phil. Journ.* (Jul.–Sept. 1828) 379.—Austral.
hieracioides, F. Muell. Fragm. i. 112 = longipedata.
inundata, A. Cunn. in DC. Prod. vi. 163 = canescens.
Kendallii, *F. Muell. in Wing, South. Sc. Record*, iii. (March 1883) 68.—Austral.
Lessoni, *Benth. Fl. Austral.* iii. 606.—Austral.
longipedata, *A. Cunn. ex DC. Prod.* vi. 163.—Austral.
Lucaeana, Walp. Rep. vi. 235 = longipedata.
microcephala, *Benth. Fl. Austral.* iii. 607.—Austral.
Mitchellii, Sond. in Linnaea. xxv. (1852) 508 = longipedata.
nutans, *Steetz, in Lehm. Pl. Preiss.* i. 464.—Austral.
pallida, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 78.—Austral.

PODOLEPIS:—
Rhytidochlamys, *F. Muell. Fragm.* iv. 79.—Austral.
rosea, *Steetz, in Lehm. Pl. Preiss.* i. 463.—Austral.
rosmarinifolia, DC. Prod. vi. 163 = Helichrysum diosmaefolium.
rugata, *Labill. Nov. Holl. Pl.* ii. 57. t. 28.—Austral.
Rutidochlamys, Benth. Fl. Austral. iii. 603 = P. Rhytidochlamys.
Siemssenia, *F. Muell. ex Benth. Fl. Austral.* iii. 607.—Austral.
subulata, Steetz, in Lehm. Pl. Preiss. i. 465 = aristata.
tetrachaeta, Walp. Ann. i. 422 = Leptorhynchos pulchellus.

PODOLOBIUM, R. Br. in Ait. Hort. Kew. ed. II. iii. 9 (1811) = **Oxylobium**, Andr. (Legumin.).
aciculare, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 98 = Chorizema Henchmanni.
aciculiferum, F. Muell. Fragm. i. 75 = O. aciculiferum.
Aquifolium, Lodd. ex G. Don, in Loud. Hort. Brit. 163 = O. trilobatum.
berberifolium, A. Cunn. ex Lindl. in Bot. Reg. (1841) Misc. 44.—Austral.
coriaceum, DC. Prod. ii. 103 = O. retusum.
heterophyllum, A. Cunn. ex Steud. Nom. ed. II. ii. 365 = O. heterophyllum.
humifusum, G. Don, Gen. Syst. ii. 116 = O. scandens.
obovatum, A. Gray, Bot. U. St. Expl. Exped. i. 379 = O. cuneatum.
procumbens, F. Muell. First Gen. Rep. 12 = O. procumbens.
scandens, DC. Prod. ii. 103 = O. scandens.
sericeum, DC. l. c. = Brachysema undulatum.
staurophyllum, Sieber, ex DC. l. c. = O. staurophyllum.
trilobatum, R. Br. in Ait. Hort. Kew. ed. II. iii. 9 = O. trilobatum.

PODOLOBUS, Rafin. in Am. Monthly Mag. (1819) 194 = **Stanleya**, Nutt. (Crucif.).

PODOLOPUS, Steud. Nom. ed. II. ii. 365 (1841) = praec.

PODOLOTUS, Benth. in Royle, Illustr. Bot. Himal. 198 (1835) = **Astragalus**, Tourn. (Legumin.).
hosackioides, Benth. l. c. = A. hosackioides.

PODONEPHELIUM, Baill. Adansonia, xi. (1874) 245. *SAPINDACEAE*.
concolor, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 120.—N. Caled.
Deplanchei, *Baill. Adansonia*, xi. (1874) 245.—N. Caled.
Homei, Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 120; et in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 540 = Deplanchei.
stipitatum, Baill. Hist. Pl. v. 396 = Deplanchei.

PODONIX, Rafin. Fl. Tellur. iv. 28 (1836) = **Tulipa**, Linn. (Liliac.).
albiflora, Rafin. l. c. = T. biflora.

PODONOSMA, Boiss. Diagn. Ser. I. xi. 113 (1849) = **Onosma**, Linn. (Borag.).
galalense, Schweinf. ex Boiss. Fl. Orient. iv. 1199.—Aegypt.; Arab.
syriacum, Boiss. Diagn. Ser. I. xi. 114 = O. syriacum.

PODOPAPPUS, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 50 = **Podocoma**, Cass. (Compos.).
hirsuta, Hook. & Arn. l. c. = Podocoma hirsuta.
pubescens, Hook. & Arn. l. c. = Podocoma hieracifolia.
tomentosus, Hook. & Arn. l. c. (= *Asteropsis macrocephala*).—Bras.

PODOPETALUM, Gaud. Fl. Helv. ii. 401 (1828) = **Trochiscanthes**, Koch (Umbell.).
nodiflorum, Gaud. l. c., in nota = T. nodiflorus.

PODOPETALUM, F. Muell. in Melb. Chemist (June 1882); ex ej. Census, 42. *LEGUMINOSAE*.
Ormondi, *F. Muell. Census*, 42, nomen.—Austral.

PODOPHANIA, Baill. in Bull. Soc. Linn. Par. i. (1880) 268. *COMPOSITAE.*
Ghiesbreghtiana, *Baill. l. c.*—Mexic.

PODOPHORUS, Phil. in Bot. Zeit. xiv. (1856) 648. *GRAMINEAE*, Benth. & Hook. f. iii. 1200.
bromoides, *Phil. l. c.* 649.—Ins. J. Fernand.

PODOPHYLLUM, Linn. Syst. ed. I (1735); Gen. ed. I. 151 (1737). *BERBERIDEAE*, Benth. & Hook. f. i. 45.
callicarpum, Rafin. Fl. Ludov. 14 = peltatum.
diphyllum, Linn. Sp. Pl. 505 = Jeffersonia binata.
Emodi, *Wall. Cat.* n. 814.—Reg. Himal.
hexandrum, Royle, Illustr. Bot. Himal. 64 = Emodi.
montanum, Rafin. Med. Fl. ii. 59 = peltatum.
peltatum, *Linn. Sp. Pl.* 505.—Am. bor.
pleianthum, *Hance, in Journ. Bot.* xxi. (1883) 175, 362.—China.
versipelle, *Hance, l. c.* 362.—China.

PODOPOGON, Ehrenb. ex Lindl. Veg. Kingd. 116 (1847), nomen (an seq.?).—Cf. Benth. & Hook. f. iii. 1096.

PODOPOGON, Rafin. Neogenyt. 4 (1825) = **Stipa**, Linn. (Gramin.).
avenaceus, Rafin. l. c. = S. avenacea.
barbatus, Rafin. l. c. = S. avenacea.
medius, Rupr. ex Steud. Syn. Pl. Gram. 118 = Beckera nubica.

PODOPTERUS, Humb. & Bonpl. Pl. Aequin. ii. 89. t. 107 (1809). *POLYGONACEAE*, Benth. & Hook. f. iii. 104.
mexicanus, *Humb. & Bonpl. l. c.*—Mexic.

PODORIA, Pers. Syn. ii. 5 (1807) = **Boscia**, Lam. (Cappar.).
angustifolia, Steud. Nom. ed. II. ii. 365; D. Dietr. Syn. Pl. iii. 232 = B. angustifolia.
senegalensis, Pers. Syn. ii. 5 = B. senegalensis.

PODOSAEMON, Spreng. Gen. i. 58 (1830) = PODOSEMUM, Desv. = **Muehlenbergia**, Schreb. (Gramin.).

PODOSAEMUM, Kunth, in Mém. Mus. Par. ii. (1815) 72 = praec.

PODOSCIADIUM, A. Gray, in Proc. Am. Acad. vii. (1868) 345. *UMBELLIFERAE.*
Bolanderi, *A. Gray, l. c.* 346.—Calif.
californicum, *A. Gray, l. c.*—Calif.

PODOSEMUM, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = **Muehlenbergia**, Schreb. (Gramin.).
agrostideum, Beauv. Agrost. 176 = M. capillaris.
alpestre, H. B. & K. Nov. Gen. et Sp. i. 131 = M. alpestris.
angustatum, J. & C. Presl, Rel. Haenk. i. 229 = M. angustata.
angustifolium, Steud. Syn. Pl. Gram. 170 = Agrostis Neesii.
angustum, Nees, Fl. Afr. Austr. 147 = Agrostis Neesii.
brachyphyllum, Nees, in Linnaea, xix. (1847) 690 = Polypogon brachyphyllus.
capillare, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = M. capillaris.
ciliatum, H. B. & K. Nov. Gen. et Sp. i. 128 = M. ciliata.
debile, H. B. & K. l. c. = M. debilis.
distichophyllum, J. & C. Presl, Rel. Haenk. i. 231 = M. distichophylla.
elegans, H. B. & K. Nov. Gen. et Sp. i. 130 = M. elegans.
foliosum, Link, Hort. Berol. i. 83 = M. mexicana.
glabratum, H. B. & K. Nov. Gen. et Sp. i. 130 = M. glabrata.
glaucum, Nees, in Linnaea, xix. (1847) 689 = Agrostis glauca.
glomeratum, Link, Hort. Berol. i. 84 = M. glomerata.
gracile, H. B. & K. Nov. Gen. et Sp. i. 131 = M. gracilis.
gymnostylum, Nees, Ind. Sem. Hort. Vratisl. (1850) 347; et in Linnaea, xxiv. (1851) 236 = M. gymnostyla.

PODOSEMUM:—
implicatum, H. B. & K. Nov. Gen. et Sp. i. 127 = Muehlenbergia implicata.
lachnanthum, Nees, Fl. Afr. Austr. 148 = Agrostis lachnantha.
mexicanum, Link, Hort. Berol. i. 84 = M. mexicana.
mucronatum, H. B. & K. Nov. Gen. et Sp. i. 129 = M. mucronata.
pungens, Link, Hort. Berol. i. 84 = Sporobolus pungens.
purpureum, Beauv. Agrost. 176 = Cinna arundinacea.
quadridentatum, H. B. & K. Nov. Gen. et Sp. i. 130 = M. quadridentata.
rigidum, H. B. & K. l. c. 129 = M. rigida.
setosum, H. B & K. l. c. = M. debilis.
soboliferum, Link, Hort. Berol. i. 83 = M. sobolifera.
stipoides, Cham. & Schlecht. in Linnaea, vi. (1831) 36 = Aristida Schiedeana.
stipoides, H. B. & K. Nov. Gen. et Sp. i. 131 = M. stipoides.
strictum, J. & C. Presl, Rel. Haenk. i. 230 = M. capillaris.
tenellum, H. B. & K. Nov. Gen. et Sp. i. 128 = M. tenella.
tenuiflorum, Link, Hort. Berol. i. 82 = M. Willdenowii.
tenuifolium, Nees, in Linnaea, xix. (1847) 690 = M. tenuifolia.
tenuissimum, J. & C. Presl, Rel. Haenk. i. 230 = M. tenuissima.
virens, Balb. ex Kunth, Enum. Pl. i. 202 = M. Berteroniana.
virescens, H. B. & K. Nov. Gen. et Sp. i. 132 = M. virescens.
virginicum, Link, Hort. Berol. i. 85 = Sporobolus virginicus.

PODOSPADIX, Rafin. Fl. Tellur. iv. 14 (1836) = **Pothos**, Linn. (Aroid.).
angustifolia, Rafin. l. c. 124 = Pothos angustifolius.
Harrisi, Rafin. l. c. = Anthurium Harrisii.
Humboldti, Rafin. l. c. = Caladium Humboldtii?
microphylla, Rafin. l. c. = Anthurium microphyllum.
reticulata, Rafin. l. c. 14 = Anthurium acaule.

PODOSPERMA, Labill. Nov. Holl. Pl. ii. 35. t. 177 (1806) = **Podotheca**, Cass. (1822) (Compos.).
angustifolium, Labill. l. c. = Podotheca angustifolia.
chrysanthum, F. Muell. Fragm. xii. 22 = Podotheca chrysantha.
gnaphaloides, F. Muell. l. c. = Podotheca gnaphaloides.
pedunculare, Reichb. ex DC. Prod. vi. 152 = Craspedia Richea.
Polakii, F. Muell. Census, 140 = seq.
Pollackii, F. Muell. Fragm. xii. 21 = Podotheca Pollackii.

PODOSPERMUM, DC. Fl. Fr. iv. 61 (1805) = **Scorzonera**, Linn. (Compos.).
angustifolium, Ledeb. ex Loud. Hort. Brit. Suppl. iii. 614 = S. angustifolia.
angustissimum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 43 = S. laciniata.
alpigenum, C. Koch, in Linnaea, xxiii. (1850) 658 = S. Jacquiniana.
armenaicum, Boiss. & Huet, Diagn. Ser. II. iii. 92 = S. armeniaca.
asperrimum, Sweet, Hort. Brit. ed. I. 206 = Leontodon asperrimus.
asperum, Sweet, l. c. 207 = Hymenonema Tournefortii.
Buxbaumii, C. Koch, in Linnaea, xvii. (1843) 274 = S. laciniata.
calcitrapaefolium, Boiss. Voy. Bot. Espagne, ii. 381 = S. intermedia?
calcitrapifolium, DC. Fl. Fr. Suppl. 455 = S. calcitrapifolia.
calcitrapifolium, C. A. Mey. Verz. Pfl. Cauc. 62 = S. intermedia.
callosum, DC. Prod. vii. 111 = S. villosa.
canum, C. A. Mey. Verz. Pfl. Cauc. 62 = S. Jacquiniana.
cariense, Boiss. Diagn. Ser. I. iv. 24 = S. floccosa.
cernuum, Unio It. ex Steud. Nom. ed. II. ii. 365 = S. Jacquiniana.
cespitosum, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 758 = S. Jacquiniana.
Columnae, DC. Prod. vii. 111 = S. villosa.

PODOSPERMUM:—
coronipifolium, Sch. Bip. ex Webb & Berth. Phyt. Canar. ii. 471 = Scorzonera laciniata?
decumbens, Gren. & Godr. Fl. Fr. ii. 310 = S. calcitrapifolia.
diversifolium, Schur, ex Nym. Consp. 465 = S. Jacquiniana.
eriospermum, Boiss. Diagn. Ser. I. vii. 5 = S. persepolitana.
heterophyllum, C. Koch, in Linnaea, xvii. (1843) 275 = S. laciniata.
heterophyllum, Schur, Bot. Rundsch. (1853) 7; et in Verh. Siebenb. Ver. Naturw. x. (1859) 61 = S. Jacquiniana.
integrifolium, Schur, Enum. Pl. Transs. 949 (Index) 364 = S. Jacquiniana.
intermedium, DC. Prod. vii. 110 = S. intermedia.
Jacquinianum, Koch, Syn. Fl. Germ. ed. I. 425 = S. Jacquiniana.
junceum, Steud. Nom. ed. II. ii. 366 = S. juncea.
laciniatum, DC. Fl. Fr. iv. 62 = S. laciniata.
loreum, Griseb. Spicil. Fl. Rumel. ii. 263 = S. Jacquiniana.
messeniacum, Steud. Nom. ed. II. ii. 366 = S. calcitrapifolia, Jacquiniana.
Meyeri, C. Koch, in Linnaea, xxiii. (1850) 659 = S. Jacquiniana.
molle, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 63 = S. mollis.
muricatum, DC. Syn. Fl. Gall. 265 = S. laciniata.
octangulare, Roth, ex Steud. Nom. ed. I. 755, in syn.; Link, Enum. Hort. Berol. ii. 279 = S. Jacquiniana.
pedunculare, Sieber, ex Benth. Fl. Austral. iii. 579 = Craspedia Richea.
phaeopappum, Boiss. Diagn. Ser. I. vii. 5 = S. multiscapa.
pinifolium, Hoffmgg. & Link, Fl. Port. ii. 127. t. 90ᵃ.—Lusit.
ponticum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 404, in textu.—As. Min.
pumilum, Sweet, Hort. Brit. ed. I. 206 (= *Zollikoferia pumila*).—Hispan.; Arab.
purpureum, Koch, ex Steud. Nom. ed. I. 755, in syn. = S. purpurea.
resedaefolium, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 470 = S. laciniata?
resedifolium, DC. Fl. Fr. iv. 61 = S. laciniata.
salinum, Schur, Enum. Pl. Transs. 364 = S. laciniata.
scorzoneroides, Turcz. ex DC. Prod. vii. 120 = S. humilis.
subulatum, DC. Fl. Fr. iv. 61 = S. laciniata.
taraxacifolium, Sweet, Hort. Brit. ed. I. 207 = Hypochoeris glabra.
Tenorii, DC. Prod. vii. 110 = S. laciniata.
tenuifolium, Hoffmgg. & Link, Fl. Port. ii. 129.—Hispan.
verrucosum, Boiss. & Bal. Diagn. Ser. II. v. 115 = S. verrucosa.
villosum, Stev. in DC. Prod. vii. 111 = S. mollis.
Willkommii, Sch. Bip. ex Willk. in Linnaea, xxv. (1852) 42 = S. intermedia.

PODOSTACHYS, Klotzsch, in Wiegm. Archiv, vii. (1841) 193 = **Croton**, Linn. (Euphorb.).
hirta, Klotzsch, l. c. 194 = C. glandulosus?
incana, Klotzsch, l. c. = C. incanus?
Lundianus, Didr. in Kjoeb. Vidensk. Meddel. (1857) 137 = C. Lundianus.
sclerocalyx, Didr. l. c. 134 = C. sclerocalyx.
Sellowiana, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. Lundianus.
serrata, Klotzsch, l. c. = C. serratus.
subfloccosa, Didr. in Kjoeb. Vidensk. Meddel. (1857) 135 = C. Lundianus.

PODOSTAURUS, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 45 = **Boenninghausenia**, Reichb. (Rutac.).
thalictroides, Jungh. l. c. 46 = B. albiflora.

PODOSTEMON, Michx. Fl. Bor. Am. ii. 164. t. 44 (1803). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 112.
DICRAEIA, Thou. Gen. Nov. Madag. 2 (1816).
POLYPLEVRUM, Tayl. ex Tul. in Arch. Mus. Par. vi. (1852) 118.
abrotanoides, *Nutt. in Journ. Acad. Philad.* vii. (1834) 105.—Am. bor.

PODOSTEMON:—
acuminatus, *Wedd. in DC. Prod.* xvii. 75.—Reg. Himal.
algaeformis, *Benth. & Hook. f. Gen.* iii. 112.—Ind. or.
Ceratophyllum, *Michx. Fl. Bor. Am.* ii. 165. t. 44.—Am. bor.
Chamissonis, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 103.—Bras.
dendroides, Thw. Enum. Pl. Zeyl. 222 = subulatus.
dichotomus, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 185.—Ind. or.
distichus, *Wedd. in DC. Prod.* xvii. 73.—Bras.
Dregeanus, Presl, Bot. Bemerk. 149 = Tristicha Dregeana.
ecostatus, Griff. ex Tul. in Ann. Sc. Nat. Sér. II. xi. (1849) 105 = Hookerianus.
elongatus, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 188.—Zeylan.
fruticulosus, *Wedd. in DC. Prod.* xvii. 74.—Bras.
Gardneri, *Harv. ex Thw. Enum. Pl. Zeyl.* 223.—Zeylan.
Griffithii, Wall. ex Gardn. in Flora, xxxiii. (1850) 40 = Hydrobryum Griffithii.
griseum, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 182 = olivaceus.
Hookerianus, *Wedd. in DC. Prod.* xvii. 74.—Ind. or.
Johnsonis, *Wedd. l. c.* 75.—Ind. or.
microcarpus, *Wedd. l. c.* 76.—Burma.
minor, *Benth. & Hook. f. Gen.* iii. 112.—Reg. Himal.
minutiflorus, *Benth. & Hook. f. l. c.*—Madag.
olivaceus, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 181.—Ind. or.
pterophyllus, *Benth. & Hook. f. Gen.* iii. 112.—Reg. Himal.; Burma.
repens, Rafin. in Am. Monthly Mag. (1811) 356, nomen (Quid?).—Am. bor.
rigidus, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 187 = dichotomus.
ruppioides, H. B. & K. Nov. Gen. et Sp. i. 246 = Apinagia ruppioides.
stylosus, *Benth. & Hook. f. Gen.* iii. 112.—Ind. or.
subulatus, *Gardn. in Calc. Journ. Nat. Hist.* vii. (1847) 184.—Zeylan.
Wallichiana, Griff. Notul. iv. 379 = seq.
Wallichii, *R. Br. in Wall. Cat.* n. 5225.—*Cf. Flora*, xxxiii. (1850) 40.—Reg. Himal.; Burma.
Wightii, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 186 = dichotomus.

PODOSTIGMA, Ell. Sketch, i. 326 (1817). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 755.
STYLANDRA, Nutt. Gen. Am. i. 170 (1818).
pubescens, *Ell. Sketch*, i. 326.—Am. bor.
viridis, Ell. l. c. 327 = Asclepiodora viridis.

PODOTHECA, Cass. in Dict. Sc. Nat. xxiii. 561 (1822). *COMPOSITAE*, Benth. & Hook. f. ii. 315.
LOPHOCLINIUM, Endl. in Bot. Zeit. i. (1843) 457
PHAENOPODA, Cass. in Dict. Sc. Nat. xlii. 84 (1826).
PODOSPERMA, Labill. Nov. Holl. Pl. ii. 35. t. 177 (1806), nomen prius.
angustifolia, *Less. Syn. Comp.* 273.—Austral.
chrysantha, *Benth. Fl. Austral.* iii. 602.—Austral.
fuscescens, *Benth. l. c.*—Austral.
gnaphalioides, *R. Grah. in Bot. Mag.* t. 3920.—Austral.
Pollackii, *F. Muell. Fragm.* xii. 21.—Austral.
pygmaea, *A. Gray, in Hook. Kew Journ.* iv. (1852) 227.—Austral.

POECHIA, Endl. Gen. Suppl. iv. 43 (1847) = **Psilostachys**, Hochst. (Amarant.).

POECILANDRA, Tul. in Ann. Sc. Nat. Sér. III. viii. (1847) 342. *OCHNACEAE*, Benth. & Hook. f. i. 320.
retusa, *Tul. l. c.*—Guian.

POECILANTHE, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 80. *LEGUMINOSAE*, Benth. & Hook. f. i. 547.
grandiflora, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 80.—Bras.
parviflora, *Benth. l. c.*—Bras.
subcordata, *Benth. l. c.*—Bras.

POECILOCHROMA, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 354. *SOLANACEAE*, Benth. & Hook. f. ii. 896.

POECILOCHROMA:—
Boissieri, Dun. & A. DC. in DC. Prod. xiii. 1. 687 (= *Lycioplesium Boissieri*).—Venezuela.
frondosum, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 355.—Peruv.
Funckianum, *Dun. & A. DC. in DC. Prod.* xiii. 1. 687.—Venezuela.
guttatum, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 356.—Peruv.
Lindenianum, *Miers, l. c.* 368.—N. Granat.
Lobbianum, *Miers, l. c.* 367.—N. Granat.
maculatum, *Miers, l. c.*—N. Granat.
punctatum, *Miers, l. c.* 354.—Peruv.
quitensis, *Miers, l. c.* (Acnisti sp.?).—Ecuador.

POECILOCNEMIS, Mart. ex Nees, in DC. Prod. xi. 286, 295 (1847) = **Geissomeria**, Lindl. etc. (Acanth.).
cincinnata, Mart. l. c. 287 = G. cincinnata.
macrostachya, Mart. l. c. 296 = Aphelandra macrostachya.
Minarum, Mart. l. c. 71 = Ebermaiera Minarum.
multiflora, Mart. l. c. 287 = G. cincinnata.

POECILODERMIS, Schott, Melet. 33 (1832) = **Sterculia**, Linn.
populnea, Schott, l. c. = S. diversifolia, *G. Don.*

POECILONEURON, Bedd. in Journ. Linn. Soc. viii. (1865) 267. t. 17. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 981.
indicum, *Bedd. l. c.*—Ind. or.
pauciflorum, *Bedd. Fl. Sylv.* i. t. 93.—Ind. or.

POECILOSTACHYS, Hack. in Sitzb. Acad. Wien, lxxxix. (1884) 131. *GRAMINEAE.*
geminata, Hack. l. c. 133 = Lophatherum geminatum.
Hildebrandtii, *Hack. l. c.* 132.—Madag.

POECILOSTEMON, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xiv. (1860) 256 = **Chrysochlamys**, Poepp. & Endl. (Guttif.).

POECILOTRICHE, Dulac, Fl. Hautes-Pyr. 528 (1867) = **Saussurea**, DC. (Compos.).
macrophylla, Dulac, l. c. = S. alpina.

POECKIA, Benth. & Hook. f. Gen. iii. 32 (1880), err. typ. = POECHIA, Endl. = **Psilostachys**, Hochst. (Amarant.).

POEDERIA, Reuss, Rep. Comm. ii. 55 (1802), sphalm. = **Paederia**, Linn. (Rubiac.).

POEONIA, Crantz, Stirp. Austr. ed. I. fasc. ii. 126 (1762); ed. II. 136 (1769) = **Paeonia**, Tourn. (Ranunc.).

POEPPIGIA, Bert. ex Férussac, Bull. Sc. Nat. xxiii. (1830) 109 = **Rhaphithamnus**, Miers (Verben.).
cyanocarpa, Bert. ex Hook. & Arn. Bot. Beech. Voy. 58 = R. cyanocarpus.

POEPPIGIA, Kunze, ex Reichb. Consp. 212[a] (1828) = **Tecophilaea**, Bert. (Haemod.).
chilensis, Kunze, ex Steud. Nom. ed. II. ii. 366 = T. violaeflora.

POEPPIGIA, Presl, Symb. Bot. i. 15. t. 8 (1830). *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
RAMIREZIA, A. Rich. Ess. Fl. Cub. t. 39 (1855).
densiflora, Tul. in Arch. Mus. Par. iv. (1844) 122 = procera.
excelsa, *A. Rich. Ess. Fl. Cub.* 484.—Cuba.
ferruginea, Tul. in Arch. Mus. Par. iv. (1844) 120 = procera.
procera, *Presl, Symb. Bot.* i. 16.—Am. centr.

POEVREA, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 76 = PEVRAEA, Comm. = **Combretum**, Linn.

POGADELPHA, Rafin. Fl. Tellur. iv. 29 (1836) = **Sisyrinchium**, Linn. (Irid.).
graminifolia, Rafin. l. c. 30 = S. graminifolia.
maculata, Rafin. l. c. 29 = S. graminifolia.

POGALIS, Rafin. Fl. Tellur. iii. 15 (1836) = **Polygonum**, Tourn.
barbata, Rafin. l. c. = Polyg. barbatum.
tinctoria, Rafin. l. c. = Polyg. tinctorium.
tomentosa, Rafin. l. c. = Polyg. tomentosum.

POGENDA, Rafin. Sylva Tellur. 10 (1838) = **Olea**, Tourn.
cernua, Rafin. l. c. = O. obtusifolia.

POGGENDORFFIA, Karst. in Linnaea, xxviii. (1856) 438; et Fl. Columb. i. 29. t. 15 (1858) = **Tacsonia**, Juss. (Passifl.).
rosea, Karst. ll. cc. 439; 29 = T. pinnatistipula.

POGOBLEPHIS, Rafin. Fl. Tellur. iii. 26 (1836) = **Gentiana**, Tourn.
missurica, Rafin. l. c. = G. Amarella.

POGOCHILUS, Falc. in Hook. Journ. Bot. iv. (1842) 73 = **Galeola**, Lour. (Orchid.).

POGOGYNE, Benth. Lab. Gen. et Sp. 414 (1834). *LABIATAE*, Benth. & Hook. f. ii. 1190.
Douglasii, *Benth. Lab. Gen. et Sp.* 414.—Calif.
multiflora, Benth. l. c. = Douglasii.
nudiuscula, *A. Gray, in Bot. Calif.* i. 597.—Calif.
parviflora, *Benth. Lab. Gen. et Sp.* 414.—Calif.
serpylloides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 386.—Calif.
tenuiflora, *A. Gray, l. c.* xi. (1876) 100.—Calif.
zizyphoroides, *Benth. Pl. Hartw.* 330.—Calif.

POGOMESIA, Rafin. Fl. Tellur. iii. 67 (1836) = **Tinantia**, Scheidw. (Commel.).
undata, Rafin. l. c. = T. fugax.

POGONANTHERA, Blume, in Flora, xiv. (1831) 520. *MELASTOMACEAE*, Benth. & Hook. f. i. 760.
corymbosa, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 240 = Geniostoma Lasiostemon.
pulverulenta, *Blume, in Flora*, xiv. (1831) 521.—Sumatra.
pulverulenta, Korth. Verh. Nat. Gesch. Bot. 247. t. 65 = reflexa.
reflexa, *Blume, in Flora*, xiv. (1831) 521.—Java; Amboina.
squamulata, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 24 = reflexa.

POGONANTHERA, Spach, Hist. Vég. Phan. ix. 583 (1840) = **Scaevola**, Linn. (Gooden.).

POGONANTHUS, Montr. in Mém. Acad. Lyon, x. (1860) 225. *INC. SED.*—Cf. Benth. & Hook. f. ii. 29.
Candollei, *Montr. l. c.*—N. Caled.

POGONATHERUM, Beauv. Agrost. 56. t. 11. f. 7 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1127.
HOMOPLITIS, Trin. Fund. Agrost. 166 (1820).
POGONOPSIS, J. & C. Presl, Rel. Haenk. i. 333. t. 46 (1830).
contortum, Brongn. in Duperr. Voy. Coq. Bot. 90. t. 17 = Pollinia articulata.
crinitum, Kunth, Enum. Pl. i. 478 = saccharoideum.
majus, Griseb. in Goett. Nachr. (1868) 92 = rufobarbatum.
polystachyum, Roem. & Schult. Syst. ii. 497 = saccharoideum.
refractum, Nees, ex Hook. & Arn. Bot. Beech. Voy. 239, in nota = saccharoideum.
rufobarbatum, *Griff. Notul.* iii. 81.—Ind. or.
saccharoideum, *Beauv. Agrost.* 176. t. 11. f. 7.—As. trop.
tenue, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 181 = Arthraxon microphyllus?

POGONATUM, Steud. Nom. ed. II. ii. 261, in syn. (1841) sphalm. = praec.

POGONELLA, Salisb. Gen. Pl. Fragm. 70 (1866) = **Simethis**, Kunth (Liliac.).
planifolium, Salisb. l. c. = S. bicolor.

POGONEMA, Rafin. Fl. Tellur. iv. 10 (1836) = **Zephyranthes**, Herb. (Amaryll.).
carinata, Rafin. l. c. = Z. carinata.

POGONETES, Lindl. Introd. Nat. Syst. ed. II. 443 (1836) = **Scaevola**, Linn. (Gooden.).
spinescens, Lindl. l. c. = S. spinescens.

POGONIA, Andr. Bot. Rep. t. 212 (1801). = **Myoporum**, Banks & Soland.
aspera, Hort. ex A. DC. in DC. Prod. xi. 710 = M. parvifolium.
debilis, Andr. Bot. Rep. t. 212 = M. debile.
glabra, Andr. l. c. t. 283 = M. acuminatum.
integerrima, Lag. Gen. et Sp. Nov. 10 = M. acuminatum.
scabra, Hort. ex A. DC. l. c. 710 = M. parvifolium.
tetrandra, Labill. Nov. Holl. Pl. i. 59. t. 83 = M. serratum.
tuberculata, Hort. ex A. DC. in DC. Prod. xi. 710 = M. parvifolium.
verrucosa, Desv. Journ. Bot. iv. (1814) 141 = M. verrucosum.

POGONIA, Juss. Gen. 65 (1789). *ORCHIDEAE*, Benth. & Hook. f. iii. 615.
APLOSTELLIS, Thou. Orch. Iles Austr. Afr. t. 24 (1822).
CLEISTES, Rich. in Mém. Mus. Par. iv. (1818) 31.
CODONORCHIS, Lindl. Gen. et Sp. Orch. 410 (1840).
CORDYLA, Blume, Bijdr. 416 (1825).
HAPLOSTELLIS, Endl. Gen. 219 (1837).
ISOTRIA, Rafin. in Desv. Journ. Bot. i. (1808) 220.
NERVILIA, Comm. ex Gaudich. in Freyc. Voy. Bot. 422. t. 35 (1826).
ODONECTIS, Rafin. in Med. Repos. N. York, v. (1808) 357.
ROPHOSTEMON, Endl. Gen. 216 (1837).
ROPTROSTEMON, Blume, Fl. Jav. Rar. Praef. p. vi (1828).
TRIPHORA, Nutt. Gen. Am. ii. 192 (1818).
affinis, *C. F. Austin, in A. Gray, Man. Bot. N. U. St.* ed. V. 507.—Am. bor.
aphylla, *Rodrig. Orch. Nov.* ii. 269.—Bras.
babianifolia, *Voigt, Hort. Suburb. Calc.* 632.—Reg. Himal.
bandana, *Blume, Orch. Archip. Ind.* 155.—Ins. Banda.
Barklyana, *Reichb. f. in Flora*, lxviii. (1885) 378; *et in Gard. Chron.* (1885) I. 726.—Ins. Mascar.
bella, *Reichb. f. & Warm. Otia Bot. Hamb.* 82.—Bras.
bicarinata, *Blume, Orch. Archip. Ind.* 152.—Ins. Mascar.
biflora, *Wight, Ic.* v. t. 1758.—Ind. or.
Bollei, *Reichb. f. Xenia Orch.* ii. 88.—As. or.
brasiliensis, *Rodrig. Orch. Nov.* i. 167.—Bras.
caloptera, *Reichb. f. & Warm. Otia Bot. Hamb.* 82.—Bras.
carinata, *Lindl. Gen. et Sp. Orch.* 414.—Ind. or.
carnosula, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 323.—Bras.
Commersonii, *Blume, Orch. Archip. Ind.* 152.—Madag.
concolor, *Blume, Mus. Bot. Lugd. Bat.* i. 32.—Java.
crispata, *Blume, l. c.*—Java.
cubensis, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 322 (rubensis).—Cuba.
Dallachyana, *F. Muell. ex Benth. Fl. Austral.* vi. 310.—Austral.
dilatata, Blume, Orch. Archip. Ind. t. 10 = punctata?
discolor, *Blume, Mus. Bot. Lugd. Bat.* i. 32.—Java.
divaricata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 203.—Am. bor.
flabelliformis, *Lindl. in Wall. Cat.* n. 7400.—Ind. or.; Malaya.
Fordii, *Hance, in Journ. Bot.* xxiii. (1885) 247.—China.
Gammieana, *Hook. f. Bot. Mag.* t. 6671.—Reg. Himal.
gentianoides, *Spreng. Syst.* iii. 706.—Jamaic.
gracilis, *Rodrig. in Rev. Engenh.* 74; *ex ej. Orch. Nov.* ii. 267.—Bras.
graminifolia, Voigt, Hort. Suburb. Calc. 632 = Spathoglottis pubescens.
grandiflora, *Reichb. f. Xenia Orch.* ii. 91.—Guian.
hirsuta, *Blume, Orch. Archip. Ind.* 152.—Madag.
holochila, *F. Muell. Fragm.* v. 200.—Austral.
japonica, Reichb. f. in Linnaea, xxv. (1852) 228 = similis.

POGONIA:—
Juliana, *Wall. Cat.* n. 7399.—Ind. or.
latipetala, *Rodrig. Orch. Nov.* ii. 265.—Bras.
lepida, *Reichb. f. Xenia Orch.* ii. 90.—Bras.
Lessonii, Reichb. f. l. c. 92 = tetraphylla.
Libonii, *Reichb. f. l. c.* 91.—Bras.
Liliastrum, *Reichb. f. l. c.* 89.—Bras.
macrantha, *Rodrig. in Rev. Engenh.* iii. 144; *ex ej. Orch. Nov.* ii. 265.—Bras.
macrophylla, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 335.—Cuba.
maculata, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 143.—Burma.
Mantiqueirae, *Reichb. f. & Warm. Otia Bot. Hamb.* 81.—Bras.
metallina, *Rodrig. in Rev. Engenh.* 74; *ex ej. Orch. Nov.* ii. 267.—Bras.
Miersii, *Reichb. f. Xenia Orch.* ii. 89.—Bras.
monantha, *Rodrig. Orch. Nov.* i. 167.—Bras.
montana, *Reichb. f. Xenia Orch.* ii. 91.—Bras.
Moritzii, Reichb. f. l. c. 89 = rosea.
Nervilia, Blume, Mus. Bot. Lugd. Bat. i. 32 = flabelliformis.
ophioglossoides, [*Ker-Gawl.*] *in Bot. Reg.* t. 148; *Nutt. Gen. Am.* ii. 192.—Am. bor.
ovata, *Lindl. Gen. et Sp. Orch.* 415.—Ins. Marian.
pachystomoides, *F. Muell. Fragm.* viii. 174.—Austral.
paludosa, *Reichb. f. Xenia Orch.* ii. 92.—Bras.
paranaënsis, *Rodrig. Orch. Nov.* ii. 268.—Bras.
parviflora, *Reichb. f. Xenia Orch.* ii. 90.—Guian.
pendula, *Lindl. in Bot. Reg.* t. 908.—Am. bor.
physurifolia, Reichb. f. in Nederl. Kruidk. Arch. iv. (1859) 324 = macrophylla.
plicata, *Lindl. Gen. et Sp. Orch.* 415.—Ind. or.
pluriflora, *Rodrig. Orch. Nov.* ii. 264.—Bras.
pulchella, Hook. f. Bot. Mag. t. 6851 = Fordii.
punctata, *Blume, Mus. Bot. Lugd. Bat.* i. 32.—Java.
purpurata, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. austr.
pusilla, *Reichb. f. & Warm. Otia Bot. Hamb.* 82.—Bras.
quadricallosa, *Rodrig. Orch. Nov.* ii. 263.—Bras.
Renschiana, *Reichb. f. Otia Bot. Hamb.* 73; *et in Bot. Zeit.* xxxix. (1881) 449.—Madag.
revoluta, *Rodrig. in Rev. Engenh.* 144; *ex ej. Orch. Nov.* ii. 266.—Bras.
Rodeiensis, *Rodrig. Orch. Nov.* ii. 269.—Bras.
rosea, *Hemsl. Biol. Centr. Am. Bot.* iii. 304.—Panama.
rubensis, Reichb. f. in Nederl. Kruidk. Arch. iv. (1859) 322, sphalm. = cubensis.
Scottii, *Reichb. f. in Flora*, lv. (1872) 276.—Reg. Himal.
similis, *Blume, Orch. Archip. Ind.* 148.—Japon.
simplex, Reichb. f. Xenia Orch. ii. 92 = Thouarsii.
speciosa, *Reichb. f. l. c.* 90.—Bras.
surinamensis, *Lindl. ex Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 674.—Guian.
tenuis, *Reichb. f. Xenia Orch.* ii. 91.—Ind. occ.
tetraphylla, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 16. t. 122.—Chili.
Thouarsii, *Blume, Orch. Archip. Ind.* 152.—Ins. Maurit.
trinervia, *Voigt, Hort. Suburb. Calc.* 632.—Ins. Molucc.
umbrosa, *Reichb. f. in Flora*, l. (1867) 102.—Afr. trop.
unguiculata, *Reichb. f. Xenia Orch.* ii. 91.—Bras.
uniflora, *F. Muell. Fragm.* v. 201.—Austral.
velutina, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 142.—Burma.
verticillata, *Nutt. Gen. Am.* ii. 192.—Am. bor.
vinosa, *Rodrig. Orch. Nov.* ii. 263.—Bras.
viridiflora, *Reichb. f. in Flora*, lxv. (1882) 532.—Java.
viridiflora, Wall. ex Voigt, Hort. Suburb. Calc. 632, nomen (Quid?).—Ind. or.?

POGONIOPSIS, Reichb. f. Otia Bot. Hamb. 82 (1881). *ORCHIDEAE*, Benth. & Hook. f. iii. 616.
Nidus-avis, *Reichb. f. l. c.*—Bras.

POGONITIS, Reichb. Handb. 226 (1837) = **Anthyllis**, Riv. (Legumin.).
montana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 196 = A. montana.

POGONOLEPIS, Steetz, in Lehm. Pl. Preiss. i. 440 (1845) = **Angianthus**, Wendl. (Compos.).
stricta, Steetz, l. c. = A. strictus.

POGONOLOBUS, F. Muell. Fragm. i. 55 (1858) = **Coelospermum**, Blume (Rubiac.).
reticulatus, F. Muell. l. c. 56 = C. reticulatum.

POGONOPHORA, Miers, ex Benth. in Hook. Kew Journ. vi. (1854) 372. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 288.
Cununri, Baill. Adansonia, iv. (1863–64) 288 = Cununria Spruceana.
Schomburgkiana, *Miers, ex Benth. in Hook. Kew Journ.* vi. (1854) 273.—Bras.
Trianae, *Muell. Arg. in Flora*, xlvii. (1864) 434.—N. Granat.

POGONOPHYLLUM, Didr. in Kjoeb. Vidensk. Meddel. (1857) 144 = **Micrandra**, Benth. (Euphorb.).
elatum, Didr. l. c. 145 = M. elata.

POGONOPSIS, J. & C. Presl, Rel. Haenk. i. 133. t. 46 (1830) = **Pogonatherum**, Beauv. (Gramin.).
tenera, J. & C. Presl, l. c. = Pogonath. saccharoideum.

POGONOPUS, Klotzsch, in Monatsb. Akad. Berl. (1853) 500. *RUBIACEAE*, Benth. & Hook. f. ii. 47.
CHRYSOXYLON, Wedd. Hist. Nat. Quinquin. 100 (1849).
HOWARDIA, Wedd. in Ann. Sc. Nat. Ser. IV. i. (1854) 66. t. 10.
exsertus, *Oerst. Am. Centr.* 17. t. 13.—Costa Rica.
Ottonis, *Klotzsch, in Monatsb. Akad. Berl.* (1853) 500.—Venezuela.

POGONORHYNCHUS, Crueg. in Linnaea, xx. (1847) 107 = **Miconia**, Ruiz & Pav. (Melastom.).
amplexans, Crueg. l. c. = M. umbrifera.
sessilifolius, Crueg. l. c. = M. tomentosa.

POGONOSPERMUM, Hochst. in Flora, xxvii. (1844) Beil. 5 = **Schwabea**, Endl. (Acanth.).
ciliare, Hochst. l. c. 6 = S. ciliaris.
hispidum, Hochst. l. c. = S. ciliaris.

POGONOSTEMON, Hassk. Cat. Hort. Bog. Alt. 130 (1844) = **Pogostemon**, Desf. (Labiat.).

POGONOSTIGMA, Boiss. Diagn. Ser. I. ii. 39 (1843) = **Tephrosia**, Pers. (Legumin.).
abyssinicum, Jaub. & Spach, Illustr. v. 88 = T. nubica.
arabicum, Boiss. Diagn. Ser. I. ii. 39 = T. Pogonostigma.
Boivini, Jaub. & Spach, Illustr. v. 87.—Arab.
nubicum, Boiss. Diagn. Ser. I. ii. 40 = T. nubica.

POGONOSTYLIS, Bertol. Fl. Ital. i. 312 (1833) = **Fimbristylis**, Vahl (Cyperac.).
squarrosus, Bertol. l. c. = F. squarrosa.

POGONOTROPHE, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 72. t. 2 = **Ficus**, Tourn. (Urtic.).
alnifolia, Miq. Pl. Jungh. 51 = F. obtusa.
assamica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 73 = F. laevis.
aurantiaca, Miq. in Zoll. Syst. Verz. Ind. Archip. 99 = F. glandulifera.
bornensis, Miq. Fl. Ind. Bat. i. II. 330 = F. obtusa.
ceylanica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 75 = F. laevis.
dasyphylla, Miq. l. c. 74 = F. laevis.
Emodi, Miq. l. c. 73 = F. laevis.
flavidula, Miq. Fl. Ind. Bat. Suppl. 435 = F. fulva, *Reinw.*
foveolata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 77 = F. foveolata.
glandulifera, Miq. l. c. = F. glandulifera.
javana, Miq. l. c. 75 = F. obtusa.
laevis, Miq. in Zoll. Syst. Verz. Ind. Archip. 93, 99 = F. laevis.
macrocarpa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 74 = F. macrocarpa.
phaeopoda, Miq. l. c. 76 = F. obtusa.
piperifolia, Miq. in Zoll. Syst. Verz. Ind. Archip. 93, 99 = F. obtusa.
pubigera, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 76 = F. foveolata.
pyrrhopoda, Miq. Fl. Ind. Bat. Suppl. 435 = F. obtusa.
reticulata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 76 = F. foveolata.

POGONOTROPHE :—
ribesioides, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 78 = Ficus recurva.
rigida, Miq. l. c. 74 = F. ramentacea.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 436 = F. glandulifera.
vagans, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 73 = F. laevis.
verrucosa, Miq. l. c. 77 = F. foveolata.
Wightiana, Miq. l. c. 74 = F. glabella.
zeylanica, Walp. Ann. i. 701 = F. laevis.

POGONURA, DC. ex Lindl. Introd. Nat. Syst. ed. II. 263 (1836) = **Perezia**, Lag. (Compos.).

POGOPETALUM, Benth. in Proc. Linn. Soc. i. (1840) 87 ; et in Trans. Linn. Soc. xviii. (1841) 684. t. 42 = **Emmotum**, Desv. (Olacin.).
acuminatum, Benth. ll. cc. 88 ; 685 = E. acuminatum.
acutum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 377 = E. fagifolium.
affine, Planch. ex Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 180 = E. acuminatum.
nitens, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 377 = E. nitens.
orbiculatum, Benth. in Proc. Linn. Soc. i. (1840) 88; et in Trans. Linn. Soc. xviii. (1841) 685. t. 42 = E. orbiculatum.

POGOSPERMUM, Brongn. in Ann. Sc. Nat. Sér. V. i. (1864) 327 = **Catopsis**, Griseb. (Bromel.).
Berteronianum, Brongn. l. c. 328 = C. nitida.
flavum, Brongn. l. c.—N. Granat.
floribundum, Brongn. l. c. 329 = C. nitida.
inconspicuum, Brongn. l. c.—Guian.
nitidum, Brongn. l. c. 328 = C. nitida.
nutans, Brongn. l. c. = C. nutans.
parviflorum, Brongn. l. c. 329 = Tillandsia parviflora.
Patschouly, Pritz. Ic. Ind. ii. 221, sphalm. = *Pogostemon Patchouly*.
sessiliflorum, Brongn. in Ann. Sc. Nat. Sér. V. i. (1864) 328 = C. nutans.

POGOSTEMON, Desf. in Mém. Mus. Par. ii. (1815) 154. t. 6. *LABIATAE*, Benth. & Hook. f. ii. 1179.
WENSEA, Wendl. Coll. Pl. iii. 24. t. 84 (1819).
amaranthoides, *Benth. in DC. Prod.* xii. 153 (amarantoides).—Reg. Himal.
atropurpureus, *Benth. l. c.* 154.—Ind. or.
Auricularia, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 127 = Dysophylla auricularia.
barbinervis, Benth. & Hook. f. Gen. ii. 1180 = Comanthosphace barbinervis.
brachystachyus, *Benth. in DC. Prod.* xii. 156.—Reg. Himal.
Cablin, *Benth. l. c.* 156.—Ins. Philipp.
comosus, *Miq. Fl. Ind. Bat.* ii. 963.—Java.
cristatus, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 127.—Java.
elsholtzioides, *Benth. in DC. Prod.* xii. 153.—Reg. Himal.
fraternus, *Miq. Fl. Ind. Bat.* ii. 963.—Reg. Himal. ; Java.
frutescens, J. Grah. Cat. Bomb. Pl. 149 = parviflorus.
Gardneri, *Hook. f. Fl. Brit. Ind.* iv. 632.—Ind. or.
glaber, *Benth. in Wall. Pl. As. Rar.* i. 31.—Reg. Himal.
gracilis, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 126.—Java.
Heyneanus, *Benth. in Wall. Pl. As. Rar.* i. 31.—Ind. or. ; Burma.
hirsutus, *Benth. Lab. Gen. et Sp.* 155.—Zeylan.
hirsutus, Wight, Ic. t. 1442 = Wightii.
imberbe, Wight, ex Hook. f. Fl. Brit. Ind. iv. 637 = atropurpureus.
intermedius, Benth. in Wall. Cat. n. 2327 = parviflorus.
japonicus, Benth. & Hook. f. Gen. ii. 1180 = Comanthosphace japonica.
longifolius, *F. Muell. Fragm.* v. 200.—Austral.
menthoides, *Blume. Bijdr.* 825.—Java.
mollis, *Benth. Lab. Gen. et Sp.* 155.—Ind. or.
mollis, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 126 = tomentosus.
paludosus, *Benth. in DC. Prod.* xii. 154.—Ind. or.
paniculatus, *Benth. in Wall. Pl. As. Rar.* i. 30.—Ind. or.
parviflorus, *Benth. l. c.* 31.—Ind. or. ; China.
Patchouly, Pellet. in Mém. Soc. Sc. Orléans, v. (1845) 277. t. 7 = Heyneanus.

POGOSTEMON :—
petiolaris, Benth. in DC. Prod. xii. 154 = Wightii.
plectrantoides, *Desf. in Ann. Mus. Par.* ii. (1803) 155. t. 6.—Ind. or.
pubescens, Benth. in DC. Prod. xii. 152 = parviflorus.
purpurascens. *Dalz. in Hook. Kew Journ.* ii. (1850) 337.—Ind. or.
purpuricaulis, Dalz. l. c. 336 = parviflorus.
quadrifolius, F. Muell. Fragm. v. 200, in obs. = Dysophylla quadrifolia.
reflexus, *Benth. in DC. Prod.* xii. 155.—Zeylan.
rotundatus, *Benth. in Wall. Pl. As. Rar.* i. 31.—Ind. or.
rotundatus, Wight, Ic. t. 1441 = mollis.
rupestris, *Benth. Lab. Gen. et Sp.* 156.—Zeylan.
speciosus, *Benth. in Wall. Pl. As. Rar.* i. 31.—Ind. or.
stellipila, Benth. & Hook. f. Gen. ii. 1180 = Comanthosphace stellipila.
strigosus, *Benth. in DC. Prod.* xii. 155.—Reg. Himal.
suavis, Tenore, in Parl. Giorn. Bot. ii. (1847) 56 = Heyneanus.
sublanceolatus, Benth. & Hook. f. Gen. ii. 1180 = Comanthosphace sublanceolata.
tomentosus, *Hassk. Cat. Hort. Bog. Alt.* 131.—Java.
travancoricus, *Bedd. Ic. Pl. Ind. Or.* i. 34.—Ind. or.
tuberculosus, *Benth. in Wall. Pl. As. Rar.* i. 31.—Reg. Himal.
velatus, *Benth. in DC. Prod.* xii. 155.—Ins. Philipp.
verticillatus, Miq. Fl. Ind. Bat. ii. 965 = Dysophylla verticillata.
vestitus, *Benth. in Wall. Pl. As. Rar.* i. 31.—Ind. or.
villosus, *Benth. Lab. Gen. et Sp.* 153.—Reg. Himal.
Wightii, *Benth. l. c.* 156.—Ind. or.

POGOSTOMA, Schrad. Ind. Sem. Hort. Gotting. (1831). —Cf. Linnaea, viii. (1833) Litt. 34 = **Capraria**, Tourn. (Scrophul.).
saxifragaefolia, Schrad. l. c. = C. saxifragaefolia.

POHLANA, Leand. Sacram. in Denkshr. Akad. Muench. (1819) 229 = **Zanthoxylum**, Linn. (Rutac.).
instrumentaria, Mart. Reise, ex Steud. Nom. ed. II. ii. 366, 796 = Z. rhoifolium.
Langsdorffii, Nees & Mart. in Nov. Act. Nat. Cur. xii. 1. (1824) 17 = Z. rhoifolium.

POICILLA, Griseb. Cat. Pl. Cub. 176 (1866). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 768.
ovatifolia, *Griseb. l. c.*—Cuba.
tamnifolia, *Griseb. l. c.* 177.—Cuba.

POIDIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836) = **Poa**, Linn. (Gramin.).
brasiliense, Nees, l. c.—Bras.

POIKILOSPERMUM, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 203 (1863–64). *URTICACEAE*, Benth. & Hook. f. iii. 389.
amboinense, *Zipp. & Miq. l. c.*—Ins. Amboin.

POINCETTIA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 252 = POINSETTIA, R. Grah. = **Euphorbia**, Linn.

POINCIA, Neck. Elem. ii. 449 (1790) = seq.

POINCIANA, Tourn. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 569.
DELONIX, Rafin. Fl. Tellur. ii. 92 (1836).
POINCIA, Neck. Elem. ii. 449 (1790).
alata, Burm. f. Fl. Ind. 333 = Caesalpinia pulcherrima.
bijuga, Burm. f. l. c. = Caesalpinia pulcherrima.
bijuga, Linn. Sp. Pl. ed. II. 709 = Caesalpinia bijuga.
bijugata, Jacq. Enum. Pl. Carib. 20 = praec.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 372.—Madag.
cassioides, Poir. Encyc. Suppl. iv. 448 = Caesalpinia cassioides.
compressa, *Sesse & Moç. ex G. Don, Gen. Syst.* ii. 433.—Mexic. ; Peruv.
coriaria, Jacq. Select. Am. 123. t. 175. f. 36 = Caesalpinia coriaria.
elata, *Linn. Cent. Pl.* ii. 16 ; *Amoen. Acad.* iv. 313.—Afr. trop.

POINCIANA :—
Gilliesii, Hook. Bot. Misc. i. (1830) 129 = Caesalpinia Gillesii.
hispida. G. Don, Gen. Syst. ii. 433 = Caesalpinia hispida ?
insignis, Kunth, Mim. t. 44 = Caesalpinia insignis.
lutea, *Regel, Cat. Pl. Hort. Aksakov.* 114.—Hab. ?
mucronata, Poir. Encyc. Suppl. iv. 448 = Caesalpinia mucronata.
Playfairii, *T. Anders. in Gard. Chron.* (1863) 388.—Arab.
pulcherrima, Linn. Sp. Pl. 380 = Caesalpinia pulcherrima.
pulchra, Salisb. Prod. 326 = praec.
punctata, Poir. Encyc. Suppl. iv. 449 = Caesalpinia punctata.
regia, *Boj. ex Hook. Bot. Mag.* t. 2884.—Madag.
Roxburghii, G. Don, Gen. Syst. ii. 433 = Peltophorum ferrugineum.
spinosa, Molina, Sagg. Chile, 158 = Caesalpinia tinctoria.
Tara, Ruiz & Pav. ex DC. Prod. ii. 481 = Caesalpinia tinctoria.

POINSETTIA, R. Grah. in Edinb. N. Phil. Journ. xx. (Jan. 1836) 412 = **Euphorbia**, Linn.
cyathophora, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253 = E. heterophylla.
dentata, Klotzsch & Garcke, l. c. = E. dentata.
Edwardsii, Klotzsch & Garcke, l. c. = E. heterophylla.
frangulaefolia, Klotzsch & Garcke, l. c. = E. geniculata.
geniculata, Klotzsch & Garcke, l. c. = E. geniculata.
heterophylla, Klotzsch & Garcke, l. c. = E. heterophylla.
insulana, Klotzsch & Garcke, l. c., nomen = E. insulana.
lancifolia, Klotzsch & Garcke, l. c. = E. lancifolia.
Morisoniana, Klotzsch & Garcke, l. c. = E. heterophylla, barbellata.
Oerstediana, Klotzsch & Garcke, l. c., nomen = E. Oerstediana.
pedunculata, Klotzsch & Garcke, in Seem. Bot. Voy. Herald, 277 = E. strigosa.
prunifolia, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253 = E. geniculata.
pulcherrima, R. Grah. in Edinb. N. Phil. Journ. xx. (Jan. 1836) 412 = E. pulcherrima.
punicea, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 253 = E. punicea.
radians, Klotzsch & Garcke, l. c. = E. radians.
Ruiziana, Klotzsch & Garcke, l. c., nomen = E. geniculata.
Schiedeana, Klotzsch & Garcke, l. c. = E. dentata.
xalapensis, Klotzsch & Garcke, l. c. = E. xalapensis.

POIRETIA, Cav. Ic. iv. 24 (1797) = **Sprengelia**, Sm. (Epacrid.).
cucullata, Cav. l. c. 25 = S. incarnata.

POIRETIA, J. F. Gmel. Syst. 263 (1791) = **Houstonia**, Gronov. (Rubiac.).
erecta, J. F. Gmel. l. c. = H. caerulea.
procumbens, J. F. Gmel. l. c. = Anotis rotundifolia.

POIRETIA, Sm. in Trans. Linn. Soc. ix. (1808) 304 = **Hovea**, R. Br. (Legumin.).
elliptica, Sm. l. c. 305 = H. Celsi.
linearis, Sm. l. c. 304 = H. linearis.

POIRETIA, Vent. Choix, t. 42 (1803). *LEGUMINOSAE*, Benth. & Hook. f. i. 513.
TURPINIA, Pers. Syn. ii. 314 (1807).
angustifolia, Vog. in Linnaea, xii. (1838) 53 = punctata.
aristata, Desv. in Journ. Bot. i. (1813) 122 = Pictetia aristata.
coriifolia, *Vog. in Linnaea*, xii. (1838) 55.—Bras.
densiflora, *Mart. in Flora*, xx. (1837) II. *Beibl.* 124.—Bras.
latifolia, *Vog. in Linnaea*, xii. (1838) 54.—Bras.
latisiliquosa, Desv. in Journ. Bot. i. (1813) 122 = Chaetocalyx latisiliqua.
multiflora, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 179 = scandens.
psoraloides, *DC. Prod.* ii. 315.—Reg. Argent.
pubescens, *Vog. in Linnaea*, xii. (1838) 52.—Bras.
punctata, *Desv. in Journ. Bot.* i. (1813) 122 ; *et in Ann. Sc. Nat.* Sér. I. ix. (1826) 411.—Bras.
refracta, *Griseb. Cat. Pl. Cub.* 72.—Cuba.
scandens, *Vent. Choix*, 42. t. 42.—Mexic.

POISSONIA, Baill. Adansonia, ix. (1870) 295 = **Coursetia**, DC. (Legumin.).
solanacea, Baill. l. c. 296 = C. orbicularis.

POITAEA, DC. Prod. ii. 263 (1825) = seq.

POITEA, Vent. Choix, 36. t. 36 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 500.
Campanilla, *DC. Prod.* ii. 263.—Ins. S. Doming.
galegiformis, Spreng. Syst. iii. 272 = seq.
galegoides, *Vent. Choix*, 36. t. 36.—Ins. S. Doming.
viciaefolia, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 99.—Ins. S. Doming.

POIVRAEA, A. Rich. Ess. Fl. Cub. 528 (1845) = seq.

POIVREA, Comm. ex Thou. Obs. Pl. Afr. Austr. 28 (1806) = **Combretum**, Linn.
aculeata, DC. Prod. iii. 18 = C. aculeatum.
aequinoctialis, Regel, Cat. Pl. Hort. Aksakov. 114.—Hab.?
alba, Rich. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 80 = *albiflora*.
albiflora, Tul. l. c.—Madag.
alternifolia, DC. Prod. iii. 17 = C. alternifolium.
barbata, Walp. Rep. ii. 64 = C. barbatum.
Berneriana, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 82.—Madag.
bracteosa, Hochst. in Flora, xxvii. (1844) 424.—Afr. austr.
coccinea, DC. Prod. iii. 18 = C. coccineum.
comosa, Sweet, Hort. Brit. ed. II. 196 = C. comosum.
conferta, Benth. in Hook. Niger Fl. 338 = C. confertum.
constricta, Benth. l. c. 337 = C. constrictum.
eriopetala, DC. Prod. iii. 18 = C. eriopetalum.
glutinosa, Klotzsch, in Peters, Reise Mossamb. Bot. 79 = C. Klotzschii.
grandiflora, Walp. Rep. ii. 64 = C. grandiflorum.
Hartmanniana, Schweinf. Pl. Nilot. 8. t. 2 = C. Hartmannianum.
intermedia, Walp. Rep. ii. 64 = C. comosum.
macrocalyx, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 81.—Madag.
mossambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. 78. t. 13 = C. constrictum.
nudiflora, Reichb. ex Walp. Rep. ii. 64 = C. aculeatum.
ovalis, Walp. l. c. = C. aculeatum.
Pavonii, Walp. l. e. = C. Pavonii.
pilosa, Wight & Arn. Prod. 317 = C. pilosum.
Roxburghii, DC. Prod. iii. 18 = C. decandrum.
rufipes, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 82.—Madag.
senensis, Klotzsch, in Peters, Reise Mossamb. Bot. 79.—Afr. trop.
sericea, Walp. Rep. ii. 64 = C. sericeum.
squamosa, Walp. l. c. = C. squamosum.
villosa, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 80 = C. villosum.
violacea, Tul. l. c. 79 = C. violaceum.

POKORNYA, Montr. in Mém. Acad. Lyon, x. (1860) 201 = **Lumnitzera**, Willd. (Combret.).
Ettingshauseni, Montr. l. c.—N. Caled.

POLAKIA, Stapf, in Denkschr. Acad. Wien, l. (1885) 43. *LABIATAE*.
paradoxa, *Stapf, l. c.*—Persia.

POLAMEIA, Reichb. Consp. 87 (1828) sphalm. = **Potameia**, Thou. (Laurin.).

POLANISIA, Rafin. in Am. Monthly Mag. (1818) 267; et in Journ. Phys. lxxxix. (1819) 98 = **Cleome**, Linn. (Capparid.).
angulata, DC. Prod. i. 242 = C. angulata.
brachiata, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 56.—Madag.
burtporensis, Munro, in Wight, Ic. t. 1072 = C. simplicifolia.
Chelidonii, DC. Prod. i. 242 = C. Chelidonii.
dianthera, DC. l. c. = C. Chelidonii.
dodecandra, DC. l. c. = C. Burmanni.
dumosa, Boj. ex Baker, in Journ. Bot. xx. (1882) 18 = C. dumosa (cf. Addend.).
felina, DC. Prod. i. 242 = C. felina.
graveolens, Rafin. in Am. Journ. Sc. Ser. I. i. (1819) 378 = C. graveolens.

POLANISIA:—
heterophylla, Wall. Cat. n. 6970 = Cleome Chelidonii.
icosandra, Wight & Arn. Prod. 22 = C. viscosa.
Leschenaultii, DC. Prod. i. 242 = C. Chelidonii.
lutea, Sond. in Harv. & Sond. Fl. Cap. i. 57 = C. lutea, *E. Mey.*
maculata, Sond. in Linnaea, xxiii. (1850) 6.—Afr. austr.
Maximiliani, Wawra, in Sitzb. Acad. Wien, Math Nat. xxxviii. (1860) 566 = C. foliosa.
micrantha, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 57 = C. micrantha, *Baker* (cf. Addend.).
microphylla, Eichl. in Mart. Fl. Bras. xiii. 1. 263.—Bras.
orthocarpa, Hochst. in Flora, xxiv. (1841) 1. Intell. 43, nomen; et Fenzl, ex Webb, Fragm. Fl. Aethiop. 23 = C. viscosa.
oxyphylla, DC. Prod. i. 242 = C. oxyphylla.
Schraderi, DC. l. c. = C. Chelidonii.
simplicifolia, Cambess. in Jacquem. Voy. Bot. 20. t. 20 = C. simplicifolia.
strigosa, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 56 = C. strigosa.
tenuifolia, Torr. & Gray, Fl. N. Am. i. 123 = C. tenuifolia.
trachysperma, Torr. & Gray, l. c. 669.—Texas.
uniglandulosa, DC. Prod. i. 242 = C. uniglandulosa.
viscosa, Blume, Bijdr. 52.—Java.
viscosa, DC. Prod. i. 242 = C. graveolens.

POLANYSIA, Rafin. in Am. Monthly Mag. (1819) 194 = praec.

POLATHERUS, Rafin. in Am. Monthly Mag. (1818) 268 = **Gaillardia**, Fouger. (Compos.).
scaber, Rafin. l. c. = G. aristata.

POLEMANNIA, Berg. ex Schlecht. in Linnaea, i. (1826) 250 = **Dipcadi**, Medic. (Liliac.).
hyacinthoides, Berg. l. c. = D. hyacinthoides.

POLEMANNIA, Eckl. & Zeyh. Enum. 347 (1837). *UMBELLIFERAE*, Benth. & Hook. f. i. 909.
grossulariaefolia, *Eckl. & Zeyh. Enum.* 348.—Afr. austr.
verticillata, *Sond. in Harv. & Sond. Fl. Cap.* ii. 551.—Afr. austr.

POLEMBRIUM, Steud. Nom. ed. II. ii. 367 (1841) = seq.

POLEMBRYON, Benth. & Hook. f. Gen. i. 299 (1862) = seq.

POLEMBRYUM, A. Juss. in Mém. Mus. Par. xii. (1825) 519. t. 28; Mem. Rut. 136. t. 49 = **Esenbeckia**, Kunth (Rutac.).
castaneaecarpum, A. Juss. ll. cc. t. 28. f. 49 = E. pilocarpoides.
Jussieui, Schott, Rutac. t. 6 = E. grandiflora.

POLEMONIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 46 (1737). *POLEMONIACEAE*, Benth. & Hook. f. ii. 823.
achilleaefolium, Willd. ex Roem. & Schult. Syst. iv. 793 = Phacelia pimpinelloides.
acutiflorum, Willd. l. c. 792 = caeruleum.
album, Fisch. Livl. Zust. 109 = caeruleum.
antarcticum, Griseb. in Goett. Abh. vi. (1854) 131 = micranthum.
arcticum, Nyl. & Saell. Herb. Mus. Fenn. 107 = humile.
boreale, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 92.—Sibir.
bursifolium, Willd. ex Roem. & Schult. Syst. iv. 793 (= *Eutoca bursifolia*).—Mexic.
caeruleum, *Linn. Sp. Pl.* 162.—Reg. bor. temp.
campanulatum, Fries, in Notis. ur Sällsk. Faun. Fl. Fenn. Förh. Haeft xiii. (1871-74) 258, 262 = caeruleum.
campanuloides, Thunb. Prod. Pl. Cap. 35 = Retzia campanuloides.
capitatum, Benth. in DC. Prod. ix. 317 = humile, caeruleum.
capitatum, Eschsch. in Mém. Acad. Pétersb. x. (1826) 282 = Gilia multicaulis.
carneum, *A. Gray, Syn. Fl. N. Am.* ii. 1. 151.—Am. bor. occ.

POLEMONIUM:—
ciliatum, Willd. ex Roem. & Schult. Syst. iv. 792 = Phacelia fimbriata.
confertum, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 73.—Am. bor. occ.
dissectum, Reichb. Ic. Fl. Germ. iv. 38 = lacteum.
dubium, Linn. Sp. Pl. 163 = Phacelia parviflora.
elatum, Salisb. Prod. 125 = caeruleum.
flavum, *Greene, in Coult. Bot. Gaz.* vi. (1881) 217.—N. Mexic.
foliosissimum, A. Gray, Syn. Fl. N. Am. ii. 1. 151 = mexicanum?
Frolovianum, Fisch. ex Herd. in Act. Hort. Petrop. i. (1874) 484 = caeruleum.
gracile, Fisch. Hort. Gorenk. ed. II. (1812) 28 = seq.?
gracile, *Willd. Enum. Hort. Berol. Suppl.* 10.—Dahuria.
grandiflorum, *Benth. in DC. Prod.* ix. 317.—Mexic.
humile, Salisb. Prod. 125 = reptans.
humile, Turcz. ex Ledeb. Fl. Ross. iii. 84 = caeruleum.
humile, *Willd. ex Roem. & Schult. Syst.* iv. 792.—Reg. arct. et Am. bor. occ.
lacteum, *Lehm. Sem. Hort. Hamb.* (1825) 17; *Pugill.* i. 27.—Sibir.
lanatum, Fisch. ex Hook. & Arn. Bot. Beech. Voy. 116 = humile.
lanatum, Pall. Reise, iii. 33 = caeruleum.
mexicanum, *Cerv. ex Lag. Gen. et Sp. Nov.* 10.—Mexic.
micranthum, *Benth. in DC. Prod.* ix. 318.—Am. bor. occ.; Chili.
moschatum, Wormsk. ex R. Grah. in Edinb. N. Phil. Journ. (Apr.–June 1829) 186 = humile.
muricatum, *Lag. ex Schrank, in Syll. Ratisb.* i. (1824) 199 (an Hydrophyllacea?).—Mexic.
Nyctelea, Linn. Sp. Pl. ed. II. 231 = Ellisia Nyctelea.
obscurum, Blanco, Fl. Filip. ed. I. 103 = Lepistemon flavescens.
pectinatum, *Greene, in Bull. Calif. Acad.* i. (1884) 10.—Calif.
pimpinelloides, Willd. ex Roem. & Schult. Syst. iv. 793 = Phacelia pimpinelloides.
prostratum, Rudolph, ex Georgi, Beschr. Russ. Reich. III. iv. 772 = humile.
pulchellum, Bunge, in Ledeb. Fl. Alt. i. 233 = humile.
pulcherrimum, Hook. Bot. Mag. t. 2979 = humile.
quadriflorum, Rafin. Atl. Journ. 177 (Quid?).—Am. bor.
reptans, *Linn. Syst.* ed. X. 925.—Am. bor.
Richardsonii, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1827) 175 = humile.
roëlloides, Thunb. Prod. Pl. Cap. 35 = Retzia roëlloides.
rubrum, Linn. Sp. Pl. 163 = Gilia coronopifolia.
sibiricum, D. Don, in Edinb. Phil. Journ. (Oct. 1822) 287 = caeruleum.
speciosum, Fisch. ex Steud. Nom. ed. II. ii. 367 = humile.
valerianaefolia, Gilib. Fl. Lituan. i. 45 = caeruleum.
villosum, Rudolph, ex Georgi, Beschr. Russ. Reich. III. iv. 771 = caeruleum.
viscosum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 154.—Am. bor. occ.
vulgare, S. F. Gray, Nat. Arr. Brit. Pl. ii. 341 = caeruleum.

POLIA, Lour. Fl. Cochinch. 164 (1790) = **Polycarpaea**, Lam. (Caryoph.).
arenaria, Lour. l. c. = Polycarpaea corymbosa.

POLIA, Tenore, Cat. Orto Bot. Nap. (1845) 92 = **Cypella**, Herb. (Irid.).
bonariensis, Tenore, l. c. = C. Herberti.
gracilis, Klatt, in Linnaea, xxxi. (1861-62) 545 = C. gracilis.
pusilla, Klatt, l. c. = C. pusilla.
ramosa, Klatt, l. c. 544 = C. Herberti.

POLIANTHES, Linn. Gen. ed. I. 97 (1737). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 737.
POLYANTHUS, Linn. Hort. Cliff. 126 (1737).
POTHOS, Adans. Fam. ii. 57 (1763).
TUBEROSA, Heist. Syst. 5 (1748).
ensifolia, Hort. ex Steud. Nom. ed. II. ii. 368 = Massonia ensifolia.
gracilis, Link, Enum. Hort. Berol. i. 330 = tuberosa.

POLIANTHES:—
maculata, *Mart. Ausw. Merkw. Pfl.* 19. t. 13.—Mexic.
mexicana, Zucc. in Abh. Akad. Muench. ii. (1837) 319 = tuberosa.
pygmaea, Jacq. Ic. Pl. Rar. t. 380 = Massonia ensifolia.
tuberosa, *Linn. Sp. Pl.* 316.—Mexic.

POLICARPEA, Lam. in Journ. Nat. Hist. Par. ii. (1792) 1 = **Polycarpaea**, Lam. (Caryoph.).

POLIGALA, Neck. Delic. Gallo-Belg. ii. 300 (1768) = **Polygala**, Tourn.

POLIGONUM, Neck. Delic. Gallo-Belg. i. 183 (1768) = **Polygonum**, Linn.

POLIODENDRON, Webb & Berth. Phyt. Canar. iii. 106. t. 173 (1836–50) = **Teucrium**, Tourn. (Labiat.).
heterophyllum, Webb & Berth. l. c. 107 = T. heterophyllum.

POLIOMINTHA, A. Gray, in Proc. Am. Acad. viii. (1873) 295, 365. *LABIATAE*, Benth. & Hook. f. ii. 1189.
glabrescens, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 549.—Texas; Mexic.
incana, *A. Gray, in Proc. Am. Acad.* viii. (1870) 296. —Am. bor. occ.
longiflora, *A. Gray, l. c.*—Mexic.
marifolia, *A. Gray, l. c.* (1872) 365.—Mexic.
mollis, *A. Gray, l. c.*—Texas.

POLIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Teucrium**, Tourn. (Labiat.).
album, Mill. Gard. Dict. ed. VIII. n. 4 = T. Polium.
angustifolium, Mill. l. c. n. 3 = T. Polium.
aransicum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
aureum, Moench, Meth. 385 = T. Polium.
brevicaule, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
canescens, Jord. & Fourr. l. c., nomen = T. Polium.
capitatum, Mill. Gard. Dict. ed. VIII. n. 5 = T. Polium.
chamaedryoides, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 475 = T. buxifolium.
collinum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
debile, Jord. & Fourr. l. c., nomen = T. Polium.
diffusum, Mill. Gard. Dict. ed. VIII. n. 13 = T. Polium.
druentinum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
drumense, Jord. & Fourr. l. c., nomen = T. Polium.
erectum, Mill. Gard. Dict. ed. VIII. n. 8 = T. Polium.
flexuosum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
fruticosum, Mill. Gard. Dict. ed. VIII. n. 11 = T. Polium.
graniticum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
integerrimum, Mill. Gard. Dict. ed. VIII. n. 14 = T. Polium.
lagopinum, Jord. & Fourr in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
latifolium, Mill. Gard. Dict. ed. VIII. n. 7 = T. Polium.
leptocladon, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
luteum, Mill. Gard. Dict. ed. VIII. n. 2 = T. Polium.
massiliense, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139 = T. aureum.
microphyllum, Jord. & Fourr. l. c. 138, nomen = T. Polium.
montanum, Jord. & Fourr. ex Nym. Consp. 566 = T. Polium.
montanum, Mill. Gard. Dict. ed. VIII. n. 1 = T. montanum.
perenne, Gueldenst. Reisen, ii. 178 = T. Polium.
pratense, Gueldenst. l. c. 163 = T. Polium.
pyrenaicum, Mill. Gard. Dict. ed. VIII. n. 6 = T. pyrenaicum.
ramosum, Mill. l. c. n. 9 = T. Polium.
rhodanicum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
saxicolum, Jord. & Fourr. l. c. = T. aureum.
serratum, Mill. Gard. Dict. ed. VIII. n. 12 = Teucrium Polium.
smyrnaeum, Mill. l. c. n. 15 = T. Polium.
spicatum, Mill. l. c. n. 10 = T. Polium.
supinum, Schur, Enum. Pl. Trans. 547 = T. montanum.
supravirens, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
tomentosum, Moench, Meth. Suppl. 133 = T. montanum.
vestitum, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 139, nomen = T. Polium.
vivariense, Jord. & Fourr. l. c., nomen = T. Polium.

POLLALESTA, H. B. & K. Nov. Gen. et Sp. iv. 46. t. 321 (1820) = **Oliganthes**, Cass. (Compos.).
vernonioides, H. B. & K. l. c. = O. triflora.

POLLIA, Thunb. Gen. Nov. i. 11 (1781). *COMMELINACEAE*, Benth. & Hook. f. iii. 846.
ACLISIA, E. Mey. in Presl, Rel. Haenk. i. 137. t. 25 (1827).
LAMPROCARPUS, Blume, ex Schult. f. Syst. vii. 1615, 1726 (1830).
Aclisia, *Hassk. Commel. Ind.* 55.—Reg. Himal.; Malaya.
condensata, *C. B. Clarke, in DC. Monog. Phan.* iii. 125.—Afr. trop.
crispata, *Benth. Fl. Austral.* vii. 90.—Austral.
cyanococca, F. Muell. Fragm. v. 40 = macrophylla, crispata.
didyma, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 451 = sorzogonensis.
elegans, *Hassk. in Miq. Pl. Jungh.* 149.—Java.
glaucescens, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1862) 305 = thyrsiflora.
gracilis, *C. B. Clarke, in DC. Monog. Phan.* iii. 124.—Ins. Johanna.
Horsfieldii, *C. B. Clarke, l. c.* 123.—Java.
indica, Thw. Enum. Pl. Zeyl. 323 = sorzogonensis.
japonica, Hance, in Journ. Bot. xvi. (1878) 233 = sorzogonensis.
japonica, *Thunb. Nov. Gen.* i. 12; *Fl. Jap.* 138.—Japon.
macrophylla, *Benth. Fl. Austral.* vii. 90.—Austral.
Mannii, *C. B. Clarke, in DC. Monog. Phan.* iii. 124.—Afr. trop.
pentasperma, *C. B. Clarke, l. c.* 129.—Reg. Himal.
purpurea, Hort. ex C. B. Clarke, l. c. 237 = Forrestia marginata.
sorzogonensis, Miq. Fl. Ind. Bat. iii. 541, partim = P. Aclisia.
sorzogonensis, *Steud. Nom.* ed. II. ii. 368.—Reg. Himal.; As. trop.; N. Caled.
subumbellata, *C. B. Clarke, in Journ. Linn. Soc.* xi. (1871) 451.—Reg. Himal.
sumatrana, *Hassk. Commel. Ind.* 56.—Ins. Philipp.?
thyrsiflora, *Endl. ex Hassk. in Miq. Pl. Jungh.* 150. —Ins. Andam.; Malaya.
Zollingeri, *C. B. Clarke, in DC. Monog. Phan.* iii. 127. —Java.

POLLICHIA, Medic. Bot. Beobacht. (1783) 247 = **Trichodesma**, R. Br. (Borag.).
africana, Medic. l. c. 248 = T. africanum.
indica, Medic. l. c. 247 = T. indicum.
latisepalea, F. Muell. Census, 100 = T. latisepaleum.
zeylanica, F. Muell. l. c. = T. zeylanicum.

POLLICHIA, Schrank, in Act. Erford. iii. (1782) 35 = **Lamium**, Tourn. (Labiat.).
amplexicaulis, Willd. Prod. Fl. Berol. 198 = L. amplexicaule.
Galeobdolon, Schrank, in Act. Erford. iii. (1782) 35 = L. Galeobdolon.
longicaulis, Krock. Fl. Siles. Suppl. ii. 150 = L. Galeobdolon?
montana, Pers. in Usteri, Ann. Bot. xiv. (1795) 39 = L. Galeobdolon.
vulgaris, Pers. l. c. = L. Galeobdolon.

POLLICHIA, [Soland. in] Ait. Hort. Kew. ed. I. i. 5 (1789). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 14.
MEERBURGIA, Moench, Meth. Suppl. 116 (1794).
NECKERIA, J. F. Gmel. Syst. 16 (1791).
campestris, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 5; iii. 505.—Afr. austr.

POLLINIA, Spreng. Pugill. ii. 10 (1815), farrago = seq. &c.

POLLINIA, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 304. *GRAMINEAE*, Benth. & Hook. f. iii. 1127.
DACTYLUS, Burm. f. Fl. Cap. Prod. 3 (1768).
EULALIA, Kunth, Rév. Gram. i. 160. t. 93 (1829).
LEPTATHERUM, Nees, in Proc. Linn. Soc. i. (1841) 92.
MICROSTEGIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836).
NEMASTACHYS, Steud. Syn. Pl. Gram. 357 (1855).
argentea, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 71.—Ind. or.; Austral.
articulata, *Trin. l. c.*—As. trop.; Austral.
aurea, *Benth. in Journ. Linn. Soc.* xix. (1881) 67.—Ins. Mascar.
brevifolia, Spreng. Syst. i. 288 = Andropogon brevifolius.
ciliaris, Spreng. l. c. 289 = Arthraxon ciliaris.
ciliata, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 186.—Ind. or.; China.
ciliata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 306.—Ind. or.
Cumingii, *Nees, in Hook. Kew Journ.* ii. (1850) 98.—As. trop.; Austral.
depauperata, Spreng. Pugill. ii. 11 = Andropogon depauperatus.
dispar, *Nees, ex Steud. Syn. Pl. Gram.* 410.—Java.
distachya, Spreng. Pugill. ii. 12 = Andropogon distachya.
dura, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 71 = Erianthus strictus.
eriopoda, Hance, in Journ. Bot. iv. (1866) 173 = Spodiopogon angustifolius.
eucnemis, *Nees, ex Steud. Syn. Pl. Gram.* 409.—Burma; Celebes.
fastigiata, Spreng. Pugill. ii. 13 = Andropogon fastigiatus.
filifolius, Benth. in Journ. Linn. Soc. xix. (1881) 67 = Erianthus filifolius.
filiformis, Spreng. Syst. i. 288 = Leptochloa cynosuroides.
fragilis, Spreng. l. c. 289 = Andropogon brevifolius.
fulta, Desv. Opusc. 69 = Andropogon Pseudograya.
fulva, Benth. Fl. Austral. vii. 526 = Cumingii.
fulva, Spreng. Neue Entdeck. ii. 93 = Chrysopogon montanus.
glabrata, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 70.—Ins. Societ.; N. Caled.
Gryllus, Spreng. Pugill. ii. 18 = Chrysopogon Gryllus.
imberbis, *Nees, ex Steud. Syn. Pl. Gram.* 410.—Ind. or.; Japon.
irritans, *Benth. Fl. Austral.* vii. 525.—Austral.
japonica, Franch. & Sav. Enum. Pl. Jap. ii. 608 = imberbis.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 290 = nuda.
lancea, Nees, ex Steud. Syn. Pl. Gram. 410 = ciliata, *Trin.*
laxa, Nees, ex Steud. l. c. = ciliata, *Trin.*
Lehmanni, *Arn. & Nees, ex Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 186.—Reg. Himal.
longisetus, Benth. in Journ. Linn. Soc. xix. (1881) 67 = Erianthus longisetosus.
Mackinlayi, *F. Muell. ex Benth. Fl. Austral.* vii. 527. —Austral.
microstachya, Desv. Opusc. 70 = Andropogon microstachyus.
monandra, Spreng. Syst. i. 288 = Pogonatherum saccharoideum.
monantha, *Nees, ex Steud. Syn. Pl. Gram.* 410.—Ind. or.
montana, Nees, ex Steud. l. c. 409 = ciliata, *Trin.*
nuda, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 307.—Ind. or.; Japon; Afr. austr.
pallida, Roem. & Schult. Syst. ii. 829 = Chrysopogon Gryllus.
polystachys, Spreng. Syst. i. 288 = Pogonatherum saccharoideum.
praemorsa, *Nees, ex Steud. Syn. Pl. Gram.* 409.—Java.
scoparia, Spreng. Pugill. ii. 13 = Andropogon scoparius.
setifolia, Nees, in Hook. Kew Journ. ii. (1850) 101 = articulata.
spectabilis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 305.—Ins. Carolin.

POLLINIA:—
striata, Spreng. Pugill. ii. 12 = Ischaemum laxum.
tenuis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 306.—Ins. Philipp.
tristachya, Thw. Enum. Pl. Zeyl. 368 = argentea.
undata, Spreng. Pugill. ii. 12 = Oplismenus compositus.
vagans, *Nees, ex Steud. Syn. Pl. Gram.* 410.—Reg. Himal.
vaginata, Spreng. Pugill. ii. 11 = Andropogon brevifolius.
versicolor, Benth. in Journ. Linn. Soc. xix. (1881) 67 = Erianthus versicolor.
villosa, *Munro, ex Benth. Fl. Hongk.* 420.—Reg. Himal.; China.
villosa, *Spreng. Syst.* i. 288.—Afr. austr.; Madag.
Wallichiana, Nees, ex Steud. Syn. Pl. Gram. 410 = ciliata, *Trin.*
Willdenowiana, Benth. in Journ. Linn. Soc. xix. (1881) 67 = imberbis.

POLLINIRHIZA, Dulac, Fl. Hautes-Pyr. 120 (1867) = **Listera**, Linn. (Orchid.).
cordata, Dulac, l. c. = L. cordata.
ovata, Dulac, l. c. = L. ovata.

POLOA, DC. in Guill. Arch. Bot. ii. (1833) 514 = **Pulicaria**, Gaertn. (Compos.).
Dalzellii, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 129 = Pul. angustifolia.
Wightiana, DC. in Guill. Arch. Bot. ii. (1833) 515; et in Wight, Contrib. 10 = Pul. Wightiana.

POLPODA, Presl, Polpoda, Nov. Pl. Gen. (1829); et Symb. Bot. i. t. 1 (1830). *FICOIDEAE*, Benth. & Hook. f. i. 858.
BLEPHAROLEPIS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 442 (1836).
capensis, *Presl, Polpoda, Nov. Pl. Gen.* (1829); *et Symb. Bot.* i. t. 1.—Afr. austr.

POLTOLOBIUM, Presl, Bot. Bemerk. 63 (1844); in Abh. Boehm. Ges. V. iii. (1845) 493 = **Andira**, Lam. (Legumin.).
hypoleucum, Presl, ll. cc. 63; 493.—Mexic.

POLYACANTHA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821) = **Centaurea**, Linn. (Compos.).
procumbens, S. F. Gray, l. c. = C. aspera.

POLYACANTHA, Hill, Herb. Brit. i. 69 (1769) = **Carduus**, Tourn. (Compos.).
acanthoides, Hill, l. c. = C. acanthoides.
acaulos, Hill, l. c. 71 = Cnicus acaulis.
crispa, Hill, l. c. = C. crispus.
palustris, Hill, l. c. 70 = Cnicus palustris.

POLYACANTHUS, Presl, Bot. Bemerk. 33 (1844); in Abh. Boehm. Ges. V. iii. (1845) 463 = **Gymnosporia**, Benth. & Hook. f. (Celastrin.).
angustifolius, Presl, ll. cc. 34; 464.—Afr. austr.
stenophyllus, Presl, ll. cc. 33; 463.—Afr. austr.

POLYACHYRUS, Lag. Amen. Nat. i. 37 (1811). *COMPOSITAE*, Benth. & Hook. f. ii. 499.
BRIDGESIA, Hook. Bot. Misc. ii. (1831) 222. t. 92.
DIAPHORANTHUS, Meyen, Reise, i. 406 (1834).
auritus, D. Don, in Phil. Mag. (1832) 390; et in Guill. Arch. Bot. ii. (1833) 466 = Poeppigii.
Bridgesii, Endl. ex Steud. Nom. ed. II. ii. 368 = sphaerocephalus.
carduoides, *Phil. Fl. Atac.* 28.—Chili.
decurrens, Lag. ex DC. Prod. vii. 53 = Poeppigii.
echinopsoides, DC. l. c. = sphaerocephalus.
fuscus, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 288.—Chili.
Gayi, *Remy, in C. Gay, Fl. Chil.* iii. 372.—Chili.
glandulosus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 423.—Peruv.
latifolius, *Phil. in Anal. Univ. Chil.* (1865) II. 348.—Chili.
litoralis, *Phil. in Linnaea*, xxviii. (1856) 714.—Chili.
macrotis, *Phil. l. c.* 713.—Chili.

POLYACHYRUS:—
multifidus, D. Don, in Phil. Mag. (1832) 390; et in Trans. Linn. Soc. xvi. (1833) 225 = niveus.
niveus, *Lag. ex DC. Prod.* vii. 53.—Chili; Peruv.
Poeppigii, *Kunze, ex Less. in Linnaea*, v. (1830) 5.—Chili.
roseus, *Phil. Fl. Atac.* 28.—Chili.
sphaerocephalus, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 330.—Peruv.
uniflorus, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 186.—Peruv.?
villosus, *Wedd. Chlor. And.* i. 56. t. 13.—Peruv.

POLYACTIDIUM, DC. Prod. v. 281 (1836) = **Erigeron**, Linn. etc. (Compos.).
delphinifolium, DC. l. c. 282 = E. delphinifolius.
Seemannii, Sch. Bip. in Seem. Bot. Voy. Herald, 301 = Achaetogeron Seemannii.
Sprengelii, DC. Prod. vii. 274 = Hysterionica montevidensis.

POLYACTIS, Less. Syn. Comp. 188 (1832) = **Erigeron**, Linn. (Compos.).
delphinifolia, Less. l. c. = E. delphinifolius.
Sprengelii, Schlecht. ex DC. Prod. vii. 274 = Hysterionica montevidensis.

POLYACTIUM, Eckl. & Zeyh. Enum. 65 (1835) = **Pelargonium**, L'Hérit. (Geran.).
aconitophyllum, Eckl. & Zeyh. l. c. 67 = Pel. aconitophyllum.
amatymbicum, Eckl. & Zeyh. l. c. 68 = Pel. amatymbicum.
anethifolium, Eckl. & Zeyh. l. c. 67 = Pel. anethifolium.
arenarium, Eckl. & Zeyh. l. c. 66 = Pel. arenarium.
caffrum, Eckl. & Zeyh. l. c. 68 = Pel. caffrum.
coniophyllum, Eckl. & Zeyh. l. c. 67 = Pel. flavum.
daucifolium, Eckl. & Zeyh. l. c. 68 = Pel. triste.
filipendulifolium, Eckl. & Zeyh. l. c. 66 = Pel. triste.
flavum, Eckl. & Zeyh. l. c. 67 = Pel. flavum.
fulgidum, Eckl. & Zeyh. l. c. = Pel. fulgidum.
gibbosum, Eckl. & Zeyh. l. c. = Pel. gibbosum.
hibridaefolium, Eckl. & Zeyh. l. c. 66 = Pel. heracleifolium.
lobatum, Eckl. & Zeyh. l. c. 65 = Pel. lobatum.
multiradiatum, Eckl. & Zeyh. l. c. 66 = Pel. radicatum.
papaverifolium, Eckl. & Zeyh. l. c. = Pel. triste.
peucedanifolium, Eckl. & Zeyh. l. c. 67 = Pel. anethifolium.
primulaeforme, Eckl. & Zeyh. l. c. 65 = Pel. pulverulentum.
radulaefolium, Eckl. & Zeyh. l. c. 66 = Pel. radulaefolium.
sphondylifolium, Eckl. & Zeyh. l. c = Pel. lobatum.
Sprengelii, Baker, in Mart. Fl. Bras. vi. III. 14, sphalm. = Hysterionica montevidensis.
triste, Eckl. & Zeyh. Enum. 67 = Pel. triste.
uitenhagense, Eckl. & Zeyh. l. c. 68 = Pel. schizopetalum.

POLYADENIA, Nees, Laurin. Expos. 18 (1833) = **Lindera**, Thunb. (Laurin.).
cuspidata, Miq. Fl. Ind. Bat. i. 1. 961.—Ins. Celebes.
gemmiflora, Nees, Syst. Laurin. 576 (= *Daphnidium gemmiflorum*).—Java.
grandifolia, Miq. Fl. Ind. Bat. i. 1. 964.—Java.
grandis, Hassk. Cat. Hort. Bog. Alt. 90 = Litsea tomentosa, *Blume*.
grandis, Nees, in Wall. Pl. As. Rar. ii. 62 (= *Tetranthera grandis*).—Ind. or.
insignis, Miq. Fl. Ind. Bat. i. 1. 962.—Sumatra.
lanceolata, Miq. l. c. 963.—Java.
lucida, Nees, Syst. Laurin. 575.—Java.
macrophylla, Miq. Fl. Ind. Bat. i. 1. 962.—Java.
Mandang, Nees, Syst. Laurin. 578.—Java.
pauciflora, Nees, l. c. 577.—Java.
pipericarpa, Miq. Fl. Ind. Bat. i. 1. 962.—Sumatra.
polyantha, Nees, Syst. Laurin. 577.—Java.
puberula, Miq. Fl. Ind. Bat. i. 1. 963.—Java.
reticulata, Nees, in Wall. Pl. As. Rar. ii. 62 = L. reticulata.
salicifolia, Miq. Fl. Ind. Bat. i. 1. 963.—Java.
subumbellifera, Steud. Nom. ed. II. ii. 369, in syn. = seq.
subumbelliflora, Nees, Syst. Laurin. 577.—Java.
umbellifera, D. Dietr. Syn. Pl. ii. 1362 = praec.

POLYALTHIA, Blume, Fl. Jav. Anonac. 68 (1829). *ANONACEAE*, Benth. & Hook. f. i. 25.
MONOON, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 15 (1865).
TRIVALVARIA, Miq. l. c. 19 (1865).
aberrans, *Maing. ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 67.—Malacca.
acuminata, *Oliver, Fl. Trop. Afr.* i. 26.—Afr. trop.
acuminata, *Thw. Enum. Pl. Zeyl.* 399.—Zeylan.
affinis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 37.—Java.
andamanica, *Kurz, Andam. Rep.* ed. II. 29.—Ins. Andam.
anomala, *Becc. in Nuov. Giorn. Bot. Ital.* iii. (1871) 188.—Borneo.
argentea, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 67.—Reg. Himal.
bifaria, *Benth. & Hook. f. ex Hook. f. l. c.* 62.—Burma.
biglandulosa, *Benth. & Hook. f. l. c.* 65.—Malaya.
cardiopetala, Dalz. in Hook. Kew Journ. ii. (1850) 39 = Goniothalamus cardiopetalus.
cauliflora, *Hook. f. & Thoms. Fl. Ind.* i. 138.—Singapore.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 14.—Ins. Celebes.
cerasoides, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 63.—Ind. or.
Chapelieri, *Baill. Adansonia*, viii. (1867–68) 349.—Ind. or.
cinnamomea, *Hook. f. & Thoms. Fl. Ind.* i. 138.—Penang.; Singapore.
coffeoides, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 62.—Ind. or.
costata, *Hook. f. & Thoms. in Hook. f. l. c.* 67.—Burma.
cuneiformis, *Blume, Fl. Jav. Anonac.* 75.—Java.
dubia, *Kurz, Andam. Rep.* ed. II. 29; *et in Journ. As. Soc. Beng.* xliii. (1874) II. 53.—Ins. Andam.
elliptica, *Blume, Fl. Jav. Anonac.* 73.—Java.
fragrans, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 63.—Ind. or.
fruticans, A. DC. in Mém. Soc. Phys. Génèv. v. (1832) 216 = Uvaria micrantha.
Holtzeana, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 230.—Austral.
hypoleuca, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 63.—Singapore.
Jenkinsii, *Benth. & Hook. f. ex Hook. f. l. c.* 64.—Reg. Himal.; Burma.
Jenkinsii, Benth. & Hook. f. l. c., partim = andamanica.
Kenthii, *Blume, Fl. Jav. Anonac.* 77.—Java.
Korinti, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 64.—Ind. or.
lanceolata, *Vidal, Phan. Cuming. Philipp.* 170.—Ins. Philipp.
lateriflora, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 52.—Burma.
longifolia, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 62.—Ind. or.
lucens, *Baker, in Journ. Linn. Soc.* xxi. (1884) 318.—Madag.
macrophylla, Blume, Fl. Jav. Anonac. 79. t. 39 = Goniothalamus macrophyllus.
macrophylla, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 66.—Burma; Malaya.
macrorhyncha, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 14.—Sumatra.
magnoliaeflora, *Maing. ex Hook. f. Fl. Brit. Ind.* i. 64.—Malacca.
maritima, *Baill. Adansonia*, viii. (1867–68) 348.—Madag.
microtus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 13.—Borneo.
miniata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 38.—Java.
Moonii, *Thw. Enum. Fl. Zeyl.* 9.—Zeylan.
nitida, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 64.—Singapore.
nitidissima, *Benth. Fl. Austral.* i. 51.—Austral.
obliqua, *Hook. f. & Thoms. Fl. Ind.* i. 138.—Malacca.
papuana, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 10.—N. Guin.
persicaefolia, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 66.—Zeylan.
Richardiana, *Baill. Adansonia*, viii. (1867–68) 350.—Madag.
rufescens, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 66.—Malabar.
sakumbangensis, *Miq. Mus. Bot. Lugd. Bat.* ii. 13.—Borneo.

POLYALTHIA:—
sclerophylla, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 65.—Malacca.
simiarum, *Benth. & Hook. f. ex Hook. f. l. c.* 63.—Reg. Himal.; Burma.
subcordata, *Blume, Fl. Jav. Anonac.* 71.—Java.
suberosa, *Benth. & Hook. f. ex Hook. f. Fl. Brit. Ind.* i. 65.—Ind. or.; Burma.
sumatrana, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 53.—Burma; Malaya.
vitiensis, *Seem. in Bonplandia*, ix. (1861) 254, nomen; Fl. Vit. 4. t. 3.—Ins. Fiji.

POLYANTHEMUM, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 228 = **Armeria**, Willd. (Plumbag.).

POLYANTHERIX, Nees, in Ann. Nat. Hist. Ser. I. i. (1838) 284 = **Elymus**, Linn. (Gramin.).
Hystrix, Nees, l. c. = E. Sitanion.

POLYANTHES, Jacq. Ic. Pl. Rar. t. 380 (1793); Coll. Suppl. 56 (1796) = **Polyxena**, Kunth (Liliac.).
pygmaea, Jacq. Il. cc.—Afr. austr.

POLYANTHES, Linn. Hort. Cliff. 126 (1737) = **Polianthes**, Linn. (Amaryll.).

POLYANTHUS, Auct. ex Benth. & Hook. f. Gen. iii. 737 (1883) = praec.

POLYARRHENA, Cass. in Dict. Sc. Nat. lvi. 172 (1826) = **Felicia**, Cass. (Compos.).
reflexa, Cass. l. c. 173 = F. reflexa.

POLYASTER, Hook. f. in Benth. & Hook. f. Gen. i. 299 (1862). *RUTACEAE*, Benth. & Hook. f. l. c.
boronioides, *Hook. f. l. c.*—Mexic.

POLYBAEA, Klotzsch, ex Benth. & Hook. f. Gen. ii. 570 (1876) = POLYBOEA, Klotzsch = **Cavendishia**, Lindl. (Vaccin.).

POLYBOEA, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1850) = **Bernardia**, Houst. (Euphorb.).
corensis, Klotzsch, l. c. = B. corensis.
Lorenzii, Griseb. in Goett. Abh. xix. (1874) 97 = B. Lorenzii.

POLYBOEA, Klotzsch, in Linnaea, xxiv. (1851) 30 = **Cavendishia**, Lindl. (Vaccin.).
crassifolia, Klotzsch, l. c. 31 = C. crassifolia.
laurifolia, Klotzsch, l. c. = C. laurifolia.
polyantha, Griseb. in Lechl. Berb. Am. Austr. 58 = Eurygania polyantha.
Quereme, Klotzsch, in Linnaea, xxiv. (1851) 31 = C. Querema.
velutina, Griseb. in Lechl. Berb. Am. Austr. 58.—Peruv.

POLYCALYMMA, F. Muell. & Sond. in Linnaea, xxv. (1852) 494 = **Myriocephalus**, Benth. (Compos.).
Stuartii, F. Muell. & Sond. l. c. = M. Stuartii.

POLYCANDIA, Steud. Nom. ed. II. ii. 369 (1841), sphalm. = **Polycardia**, Juss. (Celastr.).

POLYCANTHA, Hill, Veg. Syst. iv. 12 (1762) = **Cnicus**, Tourn. etc. (Compos.).
acanthoides, Hill, Hort. Kew. 59 = Carduus acanthoides.
acaulos, Hill, Herb. Brit. i. 71 = C. acaulis.
crispus, Hill, Hort. Kew. 59 = Carduus crispus.
palustris, Hill, l. c. = C. palustris.

POLYCARDIA, Juss. Gen. 377 (1789). *CELASTRINEAE*, Benth. & Hook. f. i. 365.
COMMERSONIA, Comm. ex Juss. Gen. 377 (1789).
FLORINDA, Noronha, ex Endl. Gen. 1086, in syn. (1841).
Aquifolium, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 103.—Madag.
epiphylla, Sm. in Rees, Cycl. xxviii. n. 1 = phyllanthoides.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 276.—Madag.
lateralis, *O. Hoffm. Sert. Madag.* 12.—Madag.
libera, *O. Hoffm. l. c.*—Madag.
madagascariensis, J. F. Gmel. Syst. 407 = phyllanthoides.
phyllanthoides, *Lam. Illustr.* ii. 100. t. 132.—Madag.

POLYCARENA, Benth. in Hook. Comp. Bot. Mag. i. (1835) 371. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 944.
aurea, *Benth. l. c.* 372.—Afr. austr.
capensis, *Benth. l. c.* 371.—Afr. austr.
capillaris, *Benth. l. c.* 372.—Afr. austr.
foliosa, *Benth. in DC. Prod.* x. 351.—Afr. austr.
gilioides, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 372.—Afr. austr.
intertexta, *Benth. l. c.*—Afr. austr.
plantaginea, *Benth. l. c.*—Afr. austr.
Plantaginis, Walp. Rep. iii. 309 = praec.
pubescens, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 372.—Afr. austr.
rariflora, *Benth. l. c.*—Afr. austr.
silenoides, *Harv. ex Benth. in DC. Prod.* x. 351.—Afr. austr.

POLYCARPA, Linden, ex Carr. in Rev. Hortic. (1868) 330 = **Idesia**, Maxim. (Bixin.).
Maximowiczii, Linden, ex Carr. l. c. = I. polycarpa.

POLYCARPAEA, Lam. in Journ. Hist. Nat. Par. ii. (1792) 8. t. 25. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 154.
ANTHYLLIS, Adans. Fam. ii. 271 (1763), partim.
AYLMERIA, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 276.
HAGAEA, Vent. Tabl. iii. 240 (1799).
HYALA, L'Hérit. ex DC. Prod. iii. 373 (1828).
LAHAYA, Roem. & Schult. Syst. v. p. xxx (1819).
MOLLIA, Willd. Hort. Berol. 11. t. 11. (1806).
PLANCHONIA, J. Gay, ex Benth. & Hook. f. Gen. i. 154 (1862).
POLIA, Lour. Fl. Cochinch. 164 (1790).
POLICARPEA, Lam. in Journ. Nat. Hist. Par. ii. (1792) 1.
ROBBAIREA, Boiss. Fl. Orient. i. 735 (1867).
aristata, *C. Sm. ex DC. Prod.* iii. 373.—Ins. Canar.
atherophora, *Steud. in Flora*, xxvi. (1843) 763.—Am. trop.
aurea, *Wight & Arn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 91.—Ind. or.
Benthamii, Wall. Cat. n. 1514 = Polycarpon Loeflingiae.
brasiliensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 183.—Bras.
breviflora, *F. Muell. in Rep. Babb. Exped.* 8.—Austral.
Burtoni, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 85.—Austral.
caespitosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 502.—Ins. Socotra.
candida, Webb & Berth. Phyt. Canar. i. 158. t. 21 = microphylla.
candidissima, Brunn. ex Bertol. Misc. Bot. iii. 9 = microphylla.
carnosa, *C. Sm. in Buch, Beschr. Canar. Ins.* 142.—Ins. Canar.
corymbosa, *Lam. Illustr.* ii. 129.—Amphig. trop.
cuspidata, *Schlecht. in Linnaea*, xiii. (1839) 408.—Mexic.
densiflora, Wall. Cat. n. 1513 = corymbosa.
depressa, DC. Prod. iii. 375 = Polycarpon Loeflingiae.
diffusa, *Wight, ex Arn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 91.—Ind. or.
divaricata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 502.—Ins. Socotra.
divaricata, Poir. ex Steud. Nom. ed. II. ii. 369 = Teneriffae.
eriantha, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 303 = corymbosa.
fallax, J. Gay, ex Oliver, Fl. Trop. Afr. i. 145 = corymbosa.
fragilis, *Delile, Fl. Egypte*, 209. t. 24. f. 1.—Aegypt.
frankenioides, *Presl, Rel. Haenk.* ii. 6.—Ins. Philipp.
Gayi, *Webb, in Hook. Niger. Fl.* 104.—Ins. Cap. Virid.
glabrifolia, *DC. Prod.* iii. 374.—Afr. trop.
glauca, C. Sm. in Tuckey, Congo, 250 = microphylla.
gnaphalodes, Poir. Encyc. Suppl. iv. 473 = microphylla.
humifusa, J. Gay, ex Oliver, Fl. Trop. Afr. i. 145 = corymbosa.
indica, Lam. in Journ. Hist. Nat. Par. ii. (1792) 8 = corymbosa.

POLYCARPAEA:—
involucrata, *F. Muell. in Rep. Babb. Exped.* 9.—Austral.
lanuginosa, Wall. Cat. n. 1515 = Polycarpon Loeflingiae.
latifolia, *Poir. Encyc. Suppl.* iv. 473.—Ins. Canar.
linearifolia, *DC. Prod.* iii. 374.—Afr. trop.
longiflora, *F. Muell. in Rep. Babb. Exped.* 8.—Austral.
marginata, *Presl, Bot. Bemerk.* 141.—Ind. or.
memphitica, Delile, Fl. Egypte, 211. t. 24. f. 2 = Polycarpon Loeflingiae.
microphylla, *Cav. in Anal. Cienc. Nat.* iii. (1801) 25.—Ins. Canar.
minuartioides, *DC. Prod.* iii. 375.—Hispan.
mozambica, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 15.—Afr. trop.
nivea, Webb, in Hook. Niger Fl. 104 = microphylla.
prostrata, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 263.—Afr. bor.; Syria; Arab.
rupicola, *Pomel, Nouv. Mat. Fl. Atl.* 202.—Alger.
serpyllifolia, Steud. Nom. ed. II. i. 803, in syn. = Paronychia capitata.
Smithii, *Link, in Buch, Beschr. Canar. Ins.* 142.—Ins. Canar.
spadicea, Lam. Illustr. ii. 129 = corymbosa.
spicata, *Wight, ex Arn. in Ann. Nat. Hist.* Ser. I. iii. (1839) 91.—Ind. or.; Afr. & Austral. trop.
spirostylis, *F. Muell. in Rep. Babb. Exped.* 9.—Austral.
staminodina, *F. Muell. l. c.* 8.—Austral.
staticaeformis, Hochst. & Steud. ex Fenzl, in Endl. Gen. 961 = spicata.
stellata, *DC. Prod.* iii. 374.—Afr. trop.
subulata, Wight & Arn. Prod. 358, sphalm. = corymbosa.
synandra, *F. Muell. in Rep. Babb. Exped.* 9.—Austral.
Teneriffae, *Lam. in Journ. Nat. Hist. Par.* ii. (1792) 8. t. 25.—Ins. Teneriff.
tenuifolia, *DC. Prod.* iii. 374.—Afr. trop.
violacea, *Benth. Fl. Austral.* i. 165.—Austral.
Wiestiana, *Presl, Bot. Bemerk.* 141.—Aegypt.

POLYCARPEA, Pomel, Nouv. Mat. Fl. Atl. 202 (1874) = praec.

POLYCARPIA, Webb & Berth. Phyt. Canar. i. 156 (1836) = praec.

POLYCARPOEA, Lam. in Journ. Hist. Nat. Par. ii. (1792) t. 25 = praec.

POLYCARPON, Loefl. It. 7 (1758); Linn. Syst. ed. X. 881 (1759). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 152.
ANTHYLLIS, Adans. Fam. ii. 271 (1763), partim.
ARVERSIA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 184. t. 112 (1829).
HAPALOSIA, Wight & Arn. Prod. 358 (1834).
TRICHLIS, Hall. Hort. Gotting. 26 (1743).
alsinefolium, *DC. Prod.* iii. 376.—Reg. Mediterr.; Afr. austr.; Austral.
alsinefolium, Webb & Berth. Phyt. Canar. ii. 155 = peploides.
apurense, *H. B. & K. Nov. Gen. et Sp.* vi. 40.—Am. austr.
arabicum, *Boiss. Diagn.* Ser. I. x. 13.—Aegypt.; Arab.
Benthamii, Hook. f. & Edgew. in Fl. Brit. Ind. i. 245, sphalm. = Loeflingiae.
Bivonae, J. Gay, in Duch. Rev. Bot. ii. (1847) 572 = peploides.
Cupani, Bubani, Dodecanthea, 14 = peploides.
depressum, Edgew. & Hook. f. in Fl. Brit. Ind. i. 245, sphalm. = Loeflingiae.
depressum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 174.—Calif.
depressum, Rohrb. in Mart. Fl. Bras. xiv. II. 257 = apurense.
diphyllum, Cav. Ic. ii. 40. t. 151. f. 1 = tetraphyllum.
floribundum, Willk. in Bot. Zeit. v. (1847) 430 = tetraphyllum.
Gmelini, Griseb. Spicil. Fl. Rumel. i. 214 = alsinefolium.
gnaphalodes, Steud. Nom. ed. II. i. 718, in syn., sphalm. = Polycarpaea microphylla.

POLYCARPON:—
herniariodes, Ball, in Journ. Bot. xi. (1873) 301 = peploides.
lanuginosum, Edgew. & Hook. f. in Fl. Brit. Ind. i. 245 = Loeflingiae.
latifolium, Bertol. in Nuov. Ann. Sc. Nat. Ser. II. ix. (1848) 138 = peploides.
Loeflingiae, *Benth. & Hook. f. Gen.* i. 153 (Loeflingii); *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 245.—Geront. trop.
magellanicum, Linn. f. Suppl. 115 = Donatia fascicularis.
peploides, *DC. Prod.* iii. 376.—Reg. Mediterr.
polyphyllum, *Blanco, Fl. Filip.* ed. I. 53.—Ins. Philipp.
pusillum, Roxb. ex Wight & Arn. Prod. 358 = Loeflingiae.
stipulifidum, Pers. Syn. i. 111 = Stipulicida setacea.
succulentum, *J. Gay, in Duch. Rev. Bot.* ii. (1847) 372.—Aegypt.; Arab.
succulentum, Webb & Berth. Phyt. Canar. i. 155 = tetraphyllum.
suffruticosum, *Griseb. in Goett. Abh.* xix. (1874) 77.—Reg. Argent.
Teneriffae, Steud. Nom. ed. II. i. 718, in syn., sphalm. = Polycarpaea Teneriffae.
tetraphyllum, *Linn. Syst.* ed. X. 881.—Amphig.
uniflorum, *Walt. Fl. Carol.* 83.—Am. bor.

POLYCENIA, Choisy, in Mém. Soc. Phys. Genèv. ii. II. (1823) 91 (Selag. 21. t. 2.) = **Hebenstretia**, Linn. (Selag.).
cordata, E. Mey. Comm. Pl. Afr. Austr. 245 = H. cordata.
dentata, Schimp. ex Choisy, in DC. Prod. xii. 4 = H. dentata.
fenestra, Steud. Nom. ed. II. ii. 370, sphalm. = seq.
fenestrata, E. Mey. Comm. Pl. Afr. Austr. 246.—Afr. austr.
fruticosa, E. Mey. l. c. 245 = H. fruticosa.
hebenstreitioides, Choisy, in Mém. Soc. Phys. Genèv. ii. II. (1823) 91 = H. repens.
lanceolata, E. Mey. Comm. Pl. Afr. Austr. 245.—Afr. austr.
tenera, Walp. Rep. iv. 143 = H. tenera.

POLYCEPHALOS, Forsk. Fl. Aegypt. Arab. 154 (1775) = **Sphaeranthus**, Vaill. (Compos.).
suaveolens, Forsk. l. c. = S. indicus.

POLYCHAETIA, Less. Syn. Comp. 371 (1832) = **Nestlera**, Spreng. (Compos.).
acerosa, DC. Prod. vi. 285 = N. acerosa.
brevifolia, DC. l. c. = Geigeria brevifolia.
Garnotii, Less. Syn. Comp. 372 = N. tenuifolia.
oppositifolia, DC. Prod. vi. 285 = N. prostrata.
passerinoides, DC. l. c. = Geigeria passerinoides.
pectidea, DC. l. c. = Geigeria pectidea.
relhanioides, Less. Syn. Comp. 371 = N. humilis.
tricephala, DC. Prod. vi. 285 = N. tricephala.

POLYCHAETIA, Tausch, ex Less. Syn. Comp. 129 (1832) = **Tolpis**, Adans. (Compos.).
coronopifolia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 404 = T. coronopifolia.

POLYCHILOS, Breda, Kuhl & Hasselt, Orch. ic. sine num. (1827) = **Phalaenopsis**, Blume (Orchid.).
Cornu-cervi, Breda, l. c. = Ph. Cornu-cervi.

POLYCHISMA, C. Muell. in Walp. Ann. vii. 489 (1868–71), sphalm. = POLYSCHISMA, Turcz. = **Pelargonium**, L'Hérit. (Geran.).

POLYCHLAENA, G. Don, Gen. Syst. i. 488 (1831) = **Melochia**, Linn. (Stercul.).
ramosa, G. Don, l. c. = M. corchorifolia.
simplex, G. Don, l. c. = M. corchorifolia.

POLYCHLAENA, Garcke, in Schweinf. Beitr. Fl. Aethiop. 56 (1867) = **Hibiscus**, Linn. (Malvac.).
adoënsis, Garcke, l. c. = H. adoënsis.

POLYCHNEMUM, Zumag. Fl. Pedem. i. 19, 401 (1849), sphalm. = **Polycnemum**, Linn. (Chenop.).

POLYCHROA, Lour. Fl. Cochinch. 559 (1790) = **Pellionia**, Gaudich. (Urtic.).
repens, Lour. l. c. = Pell. brevifolia?

POLYCLADOS, Phil. Fl. Atac. 34. t. 4. f. B (1860) = **Lepidophyllum**, Cass. (Compos.).
abietinus, Phil. in Anal. Univ. Chil. (1873) 492.—Chili.
cupressinus, Phil. Fl. Atac. 34. t. 4. f. B = L. cupressiforme?

POLYCLATHRA, Bertol. in Nov. Comm. Acad. Bonon. iv. (1840) 438. t. 46 (Fl. Guatim. 38. t. 12). Gen. dub. *CUCURBITACEARUM*.
PENTACLATHRA, Endl. Gen. Suppl. ii. 108 (1842).
cucumerina, *Bertol. ll. cc.*—Am. centr.

POLYCNEMON, F. Muell. Census, 29 (1882) = seq.

POLYCNEMUM, Linn. Gen. ed. II. 21 (1742). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 58.
SELAGO, Adans. Fam. ii. 268 (1763), partim.
acutum, Schrank, in Hoppe, Bot. Taschenb. (1798) 201 = Kochia arenaria.
alternifolium, Pall. ex Georgi, Beschr. Russ. Reich. III. iv. 665 = Petrosimonia crassifolia.
americanum, *Nutt. Gen. Am.* i. 30.—Am. bor.
arvense, *Linn. Sp. Pl.* 35.—Europ.; austr.
arvense, Sibth. & Sm. Fl. Graec. Prod. i. 23 = majus.
bonariense, *Willd. ex Link, Jahrb.* i. III. (1820) 67.—Reg. Argent.
brachiatum, Pall. Illustr. 62. t. 52; Bieb. in Mém. Soc. Nat. Mosc. i. (1806) 153 = Petrosimonia brachiata.
Camphorosma, Schrank, in Hoppe, Bot. Taschenb. (1798) 200 = Kochia arenaria.
camphorosmoides, Moq. in DC. Prod. xiii. II. 336 = Kochia arenaria.
conjugatum, Bieb. ex Ledeb. Fl. Ross. iii. 843 = Petrosimonia brachiata.
crassifolium, Pall. Illustr. 64. t. 55 = Petrosimonia crassifolia.
diandrum, *F. Muell. Native Pl. Vict.* i. 163.—Austral.
dichotomum, *Pall. Reise Suedl. Statth. Russ. Reich.* i. 287, 289, 511 (Quid?).—Rossia.
Echinus, Moq. in DC. Prod. xiii. II. 335 = Fontanesii.
erinaceum, Pall. Illustr. t. 48 = Nanophyton caspicum.
Fontanesii, *Dur. & Moq. in DC. Prod.* xiii. II. 335.—Afr. bor.
glaucum, Pall. Illustr. 63. tt. 53, 54 = Petrosimonia crassifolia.
Heuffelii, Láng, in Syll. Ratisb. ii. (1828) 219 = verrucosum.
hybridum, Andrz. ex Trautv. in Act. Hort. Petrop. ix. (1884) 140 = Chenopodium hybridum?
inundatum, Schrank, in Hoppe, Bot. Taschenb. (1798) 201 = arvense.
juniperinum, Bieb. in Mém. Soc. Nat. Mosc. i. (1806) 154 = Nanophyton caspicum.
majus, *A. Br. in Koch, Syn. Fl. Germ.* ed. II. 695.—Europ. austr.
malacophyllum, Bieb. in Mém. Soc. Nat. Mosc. i. (1806) 152 = Halimocnemis malacophylla.
mesembryanthema, F. Muell. Fragm. viii. 38 = Hemichroa mesembryanthema.
minus, Jord. ex Lloyd, Fl. L'Ouest Fr. ed. III. 261 = arvense.
monandrum, Pall. Reise, i. 483 = Petrosimonia monandra.
nuciferum, Pall. l. c. 635 = Halimocnemis sclerosperma.
oppositifolium, Pall. l. c. 484 = Petrosimonia crassifolia.
oppositifolium, Roem. & Schult. Syst. i. 525 = Petrosimonia brachiata.
pentandrum, *F. Muell. Native Pl. Vict.* i. 162.—Austral.
pumilum, Hoppe, ex Mert. & Koch, Deutschl. Fl. i. 404 = arvense.
recurvum, Loisel. Not. Fl. Fr. 151 = arvense.
salsum, Willd. Sp. Pl. i. 192 = Petrosimonia Volvox.
scleranthoides, Willd. ex Moq. in DC. Prod. xiii. II. 336 = bonariense.
sclerospermum, Pall. Reise, iii. 725 = Halimocnemis sclerosperma.
sedifolium, *Pall. Reise Suedl. Statth. Russ. Reich.* i. 258, 280 (Quid?).—Rossia.
sibiricum, Pall. Illustr. 61. t. 51 = Petrosimonia sibirica.
triandrum, Pall. Reise, i. 483 = Petrosimonia crassifolia, P. Volvox.
verrucosum, *Láng. in Syll. Ratisb.* i. (1824) 179.—Reg. Danub.; As. Min.
viaticum, Pall. Illustr. 57 = arvense.
Volvox, Pall. l. c. 60. t. 50 = Petrosimonia Volvox.
vulgare, Pall. Reise Suedl. Statth. Russ. Reich. i. 142, 143 = arvense?

POLYCODIUM, Rafin. in Am. Monthly Mag. (1818) 266 = **Vaccinium**, Linn.

POLYCOELIUM, A. DC. in DC. Prod. xi. 705 (1847) = **Myoporum**, Banks & Soland.
bontioides, A. DC. l. c. = M. bontioides.
chinense, A. DC. l. c. 706 = M. chinense.
sandwicense, A. DC. l. c. = M. sandwicense.

POLYCYCNIS, Reichb. f. in Bonplandia, iii. (1855) 218. *ORCHIDEAE*, Benth. & Hook. f. iii. 553.
barbata, *Reichb. f. l. c.*; iv. (1856) 323.—N. Granat.
gratiosa, *Endres & Reichb. f. in Gard. Chron.* (1871) 1451.—Costa Rica.
lepida, *Linden & Reichb. f. in Illustr. Hortic.* xvii. (1870) t. 19.—N. Granat.
muscifera, *Reichb. f. in Bonplandia*, iii. (1855) 218.—N. Granat.
vittata, *Reichb. f. in Walp. Ann.* vi. 618.—Guian.

POLYCYEMA, Voigt, Hort. Suburb. Calc. 141 (1845) = **Clausena**, Burm. f. (Rutac.).
pentaphyllum, Voigt, l. c. = C. pentaphylla.

POLYCYRTUS, Schlecht. in Linnaea, xvii. (1843) 126 = **Ferula**, Tourn. (Umbell.).
cachroides, Schlecht. l. c. = F. Schlechtendalii.

POLYDICLIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. 361 (1849) = **Nicotiana**, Linn. (Solan.).
multivalvis, Miers, l. c. 362 = N. quadrivalvis?
quadrivalvis, Miers, l. c. = N. quadrivalvis.

POLYDONTIA, Blume, Bijdr. 1104 (1826) = **Pygeum**, Gaertn. (Rosac.).
arborea, Blume, l. c. 1105 = Pyg. parviflorum.
ceylanica, Wight, Illustr. i. 203 = Pyg. Wightianum.
Walkerii, Wight, l. c. = Pyg. ceylanicum.

POLYDORA, Fenzl, in Flora, xxvii. (1844) 312 = **Vernonia**, Schreb. (Compos.).
stoechadifolia, Fenzl, l. c. = V. Perrottetii.

POLYECHMA, Hochst. in Flora, xxiv. (1841) 376 = **Hygrophila**, R. Br. (Acanth.).
abyssinicum, Hochst. ex Nees, in DC. Prod. xi. 83 = H. abyssinica.
coeruleum, Hochst. in Flora, xxiv. (1841) 377 = H. caerulea.
micranthum, Nees, in DC. Prod. xi. 83 = H. micrantha.
odorum, Nees, l. c. = H. odora.
radicans, Nees, l. c. = H. angustifolia.

POLYEMBRIUM, Schott, ex Steud. Nom. ed. II. ii. 367, in syn. (1841) = seq.

POLYEMBRYUM, Schott, ex Steud. Nom. ed. II. ii. 370 (1841) = POLEMBRYUM, A. Juss. = **Esenbeckia**, Kunth (Rutac.).

POLYGALA, [Tourn.] Linn. Syst. ed. I (1735). *POLYGALEAE*, Benth. & Hook. f. i. 136.
ANTHALOGEA, Rafin. New Fl. Am. iv. 88 (1836).
ASEMEIA, Rafin. l. c. (1836).
BADIERA, Hassk. Cat. Hort. Bog. Alt. 227 (1844).
BRACHYTROPIS, Reichb. Consp. 120 (1828).
CHAMAEBUXUS, [Tourn.] Spach, Hist. Vég. Phan. vii. 125 (1839).
CORYMBULA, Rafin. New Fl. Am. iv. 88 (1836).
ISOLOPHUS, Spach, Hist. Vég. Phan. vii. 112 (1839).
MICROLOPHIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195.
PHYLAX, Noronha, in Verh. Batav. Gen. v. (1790), ed. I. Art. IV. 3.
PILOSTACHYS, Rafin. New Fl. Am. iv. 89 (1836).
PLOSTAXIS, Rafin. l. c. 87 (1836).
POLYGALOIDES, Tourn. ex Hall. Enum. Stirp. Helvet. ii. 607 (1742).
PSYCHANTHUS, Spach, Hist. Vég. Phan. vii. 130 (1839).
SEMEIOCARDIUM, Zoll. in Tijdschr. Nederl. Ind. xvii. (1858) 245.
SENEGA, Spach, Hist. Vég. Phan. vii. 128 (1839).
SEXILIA, Rafin. New Fl. Am. iv. 87 (1836).
TRICHOLOPHUS, Spach, Hist. Vég. Phan. vii. 114 (1839).

POLYGALA :—
Triclisperma, Rafin. Specch. i. 117 (1814).
Zoroxus, Rafin. New Fl. Am. iv. 88 (1836).
abyssinica, *R. Br. in Salt, Abyss. App.* p. lxv, nomen; *Fresen. in Mus. Senckenb.* ii. (1837) 273.—Ind. or.; Afr. trop.
acanthoclada, *A. Gray, in Proc. Am. Acad.* xi. (1876) 73.—Am. bor. occ.
acerosa, E. Mey. in Drège, Zwei Pfl. Docum. 212 = ericaefolia.
acicularis, *Oliver, Fl. Trop. Afr.* i. 132.—Afr. trop.
aculeata, *Vell. Fl. Flum.* 291; vii. t. 62.—Bras.
acuminata, E. Mey. in Drège, Zwei Pfl. Docum. 212 = amatymbica.
acuminata, Willd. Sp. Pl. iii. 887 = Badiera acuminata.
acutifolia, Linn. Syst. ed. X. 1154 (Quid?).—Europ.?
acutifolia, Steud. Nom. ed. II. ii. 370 = lancifolia.
acutifolia, Torr. & Gray, Fl. N. Am. i. 128 = cymosa.
adenophora, *DC. Prod.* i. 327.—Guian.
adenophylla, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 29 (non al.).—Bras.
adenophylla, *A. St. Hil. Fl. Bras. Mer.* ii. 20.—Bras.
adenophylla, Steud. Nom. ed. II. ii. 370 = Monnina resedoides.
adoënsis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 37 = abyssinica.
adpressa, Steud. ex Stahl, Estud. Fl. Puerto-rico Foll. ii. 52 = longicaulis.
adscendens, Clarke, Trav. iii. 337 = venulosa.
aestuans, Linn. f. Suppl. 315 = Monnina aestuans.
affinis, *DC. Prod.* i. 322.—Afr. austr.
alba, *Nutt. Gen. Am.* ii. 87.—Am. bor.
albida, Steud. Nom. ed. I. 642 = P. Senega.
alopecuroides, Linn. Mant. ii. 260 = Muraltia alopecuroides.
alpestris, C. A. Mey. Verz. Pfl. Cauc. 103 = amara.
alpestris, Reichb. Pl. Crit. i. 25 = microcarpa.
Alphonsii, *Wall. Cat.* n. 4179.—Burma.
alpicola, *Rupr. Fl. Cauc.* 166, in textu.—Reg. Caucas.
alpina, Steud. Nom. ed. I. 642 = amara.
amara, *Linn. Syst.* ed. X. 1154.—Europ.
amarella, Coss. & Germ. Fl. Par. 56. t. 7 = calcarea.
amarella, Crantz, Stirp. Austr. ed. II. fasc. v. 438 = amara.
amatymbica, *Eckl. & Zeyh. Enum.* 24.—Afr. austr.
ambigua, Nutt. Gen. Am. ii. 89 = verticillata.
ambigua, Torr. & Gray, Fl. N. Am. i. 130 = Nuttallii.
amblyptera, Reichb. Pl. Crit. i. 91 = nicaeënsis.
amblyptera, F. Schultz, in Flora, xx. (1837) 752 = calcarea.
americana, *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
americana, Seem. Bot. Voy. Herald, 270 = puberula, *A. Gray*, ovalifolia.
amoena, Thunb. Prod. Pl. Cap. 120 = myrtifolia.
anatolica, Boiss. & Heldr. Diagn. Ser. II. i. 57 = major.
andrachnoides, Willd. Sp. Pl. iii. 875 = supina.
Angelisia, Tenore, Fl. Nap. v. 92 = microcarpa.
angulata, *DC. Prod.* i. 328.—Bras.
angustata, Schur, Enum. Pl. Transs. 89 = vulgaris.
angustifolia, A. W. Benn. in Mart. Fl. Bras. xiii. III. 14, partim = mollis.
angustifolia, Buch.-Ham. ex Wall. Cat. n. 4170 D = telephioides.
angustifolia, Gilib. Fl. Lituan. ii. 113 = vulgaris.
angustifolia, Heyne, ex Wight & Arn. Prod. 37 = chinensis.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 405. t. 511.—N. Mexic.
aparinoides, Hook. & Arn. Bot. Beech. Voy. 277 = verticillata.
aphylla, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 21. Bras.
appendiculata, Vell. Fl. Flum. 292; vii. t. 66 = paniculata.
appressa, *Benth. in Hook. Journ. Bot.* iv. (1842) 100.—Guian.
aquitanica, Clavaud, in Ann. Soc. Linn. Bord. xxxv. (1881) 362. t. 15. f. 5 = vulgaris.
arabica, Boiss. Diagn. Ser. I. i. 9 = irregularis.
arabica, Edgew. in Journ. As. Soc. Beng. xvi. (1847) II. 1213 = erioptera.
ardisioides, Wall. Cat. n. 4178 = crotalarioides.
areguensis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 201.—Am. austr.
arenaria, Oliver, Fl. Trop. Afr. i. 128 = persicariaefolia.
arenaria, *Willd. Sp. Pl.* iii. 880.—Afr. trop.

POLYGALA :—
arillata, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 199.—Ind. or.; Malaya.
Arnottiana, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 159 = Salomonia oblongifolia.
Arnottiana, Steud. Nom. ed. II. ii. 370 = rosmarinifolia.
arvensis, Roxb. Hort. Beng. 53; Fl. Ind. iii. 218 = Salomonia ciliata.
arvensis, Wall. Cat. n. 4172 = telephioides.
arvensis, Willd. Sp. Pl. iii. 876 = chinensis.
arvicola, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1842) 13.—Madag.
asbestina, *Burch. Trav.* i. 543.—Afr. austr.
ascendens, Auct. = *adscendens*.
aspalata, A. St. Hil. Fl. Bras. Mer. ii. 14, partim = sabulosa.
aspalatha, *Linn. Mant.* i. 99.—Bras.
asperuloides, DC. Prod. i. 329 = crucianelloides.
asperuloides, H. B. & K. Nov. Gen. et Sp. v. 403 = verticillata.
atropurpurea, *A. St. Hil. Fl. Bras. Mer.* ii. 10.—Bras.
attenuata, Hook. Journ. Bot. i. (1834) 195 = Hookeri.
attenuata, Lodd. Bot. Cab. t. 1000 = oppositifolia.
attenuata, Nutt. Gen. Am. ii. 90 = cymosa.
australis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 203.—Uruguay.
austriaca, Crantz, Stirp. Austr. ed. II. fasc. v. 439 = amara.
axillaris, Poir. Encyc. v. 489 = Floscopa axillaris.
badensis, Schimp. in Spenn. Fl. Frib. iii. 867 = serpyllacea.
baetica, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 559.—Hispan.
Balansae, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 240.—Marocc.
Baldwini, *Nutt. Gen. Am.* ii. 90.—Am. bor.
Ballii, Nym. Consp. 83 = vulgaris.
Banksii, Jacq. ex Steetz, in Lehm. Pl. Preiss. ii. 297 = Comesperma ericinum.
Barthiana, Fuss, in Verh. Siebenb. Ver. Naturw. xix. (1868) 210 = sibirica.
basaltica, Lamotte, Prod. Fl. Centr. Fr. i. 125 = vulgaris.
Beiliana, Eckl. & Zeyh. Enum. 23 = Muraltia Beiliana.
Berteriana, DC. Prod. i. 328 = hygrophila.
Beyrichii, Torr. & Gray, Fl. N. Am. i. 130 = alba.
bicolor, Hook. Journ. Bot. i. (1834) 194 = Boykini.
bicolor, H. B. & K. Nov. Gen. et Sp. v. 394 t. 507 = alba.
bicolor, Vell. Fl. Flum. 290; vii. t. 59 = Acanthocladus albicans.
Boissieri, *Coss. Notes Crit.* 100.—Hispan.
boliviensis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 171.—Bolivia.
Bonavisio, Vell. Fl. Flum. 292; vii. t. 64 = angulata.
borboniaefolia, Burch. ex DC. Prod. i. 322 = oppositifolia.
Bowkerae, *Harv. in Harv. & Sond. Fl. Cap.* i. 92.—Afr. austr.
Boykini, *Nutt. in Journ. Acad. Philad.* vii. (1834) 86.—Am. bor.
brachiata, Reichb. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 39 = appressa.
brachistachyos, Poir. Encyc. Suppl. iv. 477 = chinensis.
brachypetala, Wallr. Beitr. Fl. Hercyn. 246 (Quid?).—Europ.
brachypoda, *Tod. Nuov. Gen. e Sp.* 54.—Hab.?
brachyptera, *Griseb. Cat. Pl. Cub.* 12.—Cuba.
brachystachya, DC. Prod. i. 326; Blume, Bijdr. 59 = chinensis.
bracteata, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 32.—Bras.
bracteata, Thunb. Prod. Pl. Cap. 120 = bracteolata, *Linn.*
bracteolata, Burm. ex DC. Prod. i. 336 = Brachycarpaea varians.
bracteolata, *Forsk. Fl. Aegypt. Arab.* 213.—Arab.
bracteolata, *Linn. Sp. Pl.* 702.—Afr. austr.
brasiliensis, *Linn. Mant.* i. 99.—Bras.
brasiliensis, Mart. in Flora, xx. (1837) II. Beibl. 93 = paniculata.
brevifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 90.—Afr. austr.
brevifolia, *Nutt. Gen. Am.* ii. 89.—Am. bor.
brizoides, A. St. Hil. Fl. Bras. Mer. ii. 44. t. 88 = angustifolia.
bryoides, *A. St. Hil. l. c.* 39.—Bras.
Buchanani, D. Don, Prod. Fl. Nep. 199 = persicariaefolia.

POLYGALA :—
Buchenavii, *O. Hoffm. in Bremen, Abh.* vii. (1882) 240 (sp. dub.).—Madag.
Burmanni, *DC. Prod.* i. 322.—Afr. austr.
Burmanni, Eckl. & Zeyh. Enum. 21 = bracteolata, *Linn.*
buxifolia, Ball, ex Nym. Consp. 83 = vulgaris.
buxifolia, *H. B. & K. Nov. Gen. et Sp.* v. 407.—Mexic.
buxifolia, Reichb. Pl. Crit. i. 26 = calcarea.
buxifolia, St. Lag. in Ann. Soc. Bot. Lyon, vi. (1880) 132 = P. Chamaebuxus.
buxiformis, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 161 = telephioides.
calcarea, *Schultz, in Flora*, xx. (1837) 752.—Europ.
calcarea, Zett. ex Willk. & Lange, Prod. Fl. Hisp. iii. 557 = microcarpa.
californica, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 671.—Am. bor.
calvipes, Schlecht. in Linnaea, xiv. (1840) 160 = americana.
calycina, Presl, Bot. Bemerk. 15 = bracteolata, *Linn.*
campestris, Dalz. in Hook. Kew Journ. ii. (1850) 40 = elongata.
campestris, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 332.—Bras.
camporum, Benth. in Hook. l. c. iv. (1842) 100 = angustifolia.
canarana, A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 207 = Salomonia oblongifolia.
capillaris, *E. Mey. in Drège, Zwei Pfl. Docum.* 212.—Afr. trop. et austr.
capillifolia, *Desf. ex DC. Prod.* i. 327.—Hab.?
capitata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 199 = crotalarioides.
caracasana, *H. B. & K. Nov. Gen. et Sp.* v. 407.—Venezuela.
cardiocarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 293.—Burma.
caripensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 350.—Venezuela.
Carmichaelii, *Harv. in Harv. & Sond. Fl. Cap.* i. 19*.—Afr. austr.
carniolica, *A. Kern. Sched. Fl. Exsicc.* iii. 64.—Carniol.
caucasica, *Rupr. Fl. Cauc.* 165.—Reg. Caucas.
celosioides, *Mart. ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 35.—Bras.
cernua, Eckl. & Zeyh. Enum. 22 = abyssinica.
cernua, Thunb. Prod. Pl. Cap. 120 = virgata.
cestrifolia, A. St. Hil. Fl. Bras. Mer. ii. 51 = P. Laureola.
ceylanica, Heyne, ex Wall. Cat. n. 4183 = javana.
Chamaebuxus, *Linn. Sp. Pl.* 704.—Europ.
Chapmanii, *Torr. & Gray, Fl. N. Am.* i. 131.—Am. bor.
chatamensis, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 232.—Ins. Galap.
chinensis, *Linn. Sp. Pl.* 704.—China; Ind. or.
chloroneura, *Griseb. in Goett. Abh.* xix. (1874) 75.—Reg. Argent.
ciliata, Lebel, ex Gren. & Godr. Fl. Fr. i. 195 = vulgaris.
ciliata, Linn. Sp. Pl. 705 = Salomonia ciliata.
ciliata, Wight, ex A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 204 = chinensis.
ciliatifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 347.—Afr. austr.
ciliolata, Harv. in Harv. & Sond. Fl. Cap. i. 87 = ciliatifolia.
cineraria, Willd. Sp. Pl. iii. 880 (*cinearia*) = violacea, *Vahl.*
cinerea, DC. Prod. i. 330 = violacea, *Vahl.*
clavipes, Walp. Rep. iv. 648 = americana.
cluytioides, Burch. ex DC. Prod. i. 322 = myrtifolia.
Cneorum, *A. St. Hil. Fl. Bras. Mer.* ii. 38.—Bras.
coerulea, Miégev. ex Reichb. Pl. Crit. i. 25 = vulgaris.
comata, *Mart. ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 42.—Bras.
comata, Mart. ex A. W. Benn. l. c., partim = P. Timoutou.
commutata, Sweet, Hort. Brit. ed. II. 47 = paucifolia.
comosa, Schk. Handb. ii. 324. t. 194 = vulgaris.
conferta, *A. W. Benn. in Hemsl. Diagn. Pl. Nov.* 2.—Mexic.
confusa, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 165 = glaucoides.
coniicarpa, *Hance, in Walp. Ann.* ii. 80 (sp. dub.).—China.
conosperma, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 266.—Afr. trop.

POLYGALA:—
cordifolia, *Presl, Rel. Haenk.* v. 99.—Peruv.
cordifolia, Thunb. Prod. Pl. Cap. 120 = oppositifolia.
coriacea, *A. St. Hil. Fl. Bras. Mer.* ii. 19.—Am. austr.
corifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 129.—N. Granat.
corisoides, A. St. Hil. Fl. Bras. Mer. ii. 17. t. 83 = P. Cyparissias.
cornuta, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 63 = californica.
corsica, Gremli, Neue Beitr. Fl. Schweiz, iii. 4 = vulgaris.
corsica, Sieber, ex Steud. Nom. ed. II. ii. 371 = nicaeënsis.
coryfolia, C. Muell. in Walp. Ann. vii. 248 = corifolia?
corymbosa, Michx. Fl. Bor. Am. ii. 54, partim = cymosa.
corymbosa, Nutt. Gen. Am. ii. 89 = ramosa.
Coursiereana, Pomel, Nouv. Mat. Fl. Atl. 211 = nicaeënsis.
crotalarioides, *Buch.-Ham. ex DC. Prod.* i. 327.—Reg. Himal.
crotalarioides, Roxb. ex Steud. Nom. ed. II. ii. 371 = chinensis.
crotalloides, Roxb. ex Wight & Arn. Prod. 36, sphalm. = chinensis.
crucianelloides, *DC. Prod.* i. 329.—Ins. S. Doming.
cruciata, *Linn. Sp. Pl.* 706.—Am. bor.
cruciata, Torr. & Gray, Fl. N. Am. i. 127, partim = = brevifolia, *Nutt.*
cucullata, *Benth. Pl. Hartw.* 299.—Calif.
cucullata, Newberry, in Pacif. Rail. Rep. vi. 70 = californica.
Curtissii, *A. Gray, Man. Bot. N. U. St.* ed. V. 121.—Am. bor.
cuspidata, DC. Prod. i. 328 = P. Timoutou.
cuspidata, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 194 = cruciata.
cyanolopha, Hance, in Journ. Bot. iv. (1866) 117 = telephioides.
cyanophila, C. Muell. in Walp. Ann. vii. 245, sphalm. = seq.
cymosa, Poir. Encyc. v. 500 = ramosa.
cymosa, *Walt. Fl. Carol.* 179.—Am. bor.
Cyparissias, *A. St. Hil. Fl. Bras. Mer.* ii. 15.—Am. austr.
Darwiniana, *A. W. Benn. in Journ. Bot.* xvii. (1879) 203.—Patagon.
Decaisnei, Steud. Nom. ed. II. ii. 371 = spinescens.
decipiens, Bess. Enum. Pl. Volh. 73 = amara.
declinata, E. Mey. in Drège, Zwei Pfl. Docum. 212 = hispida.
decora, Sond. in Linnaea, xxiii. (1850) 14 = virgata.
decumbens, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 15.—Bras.
densiflora, Blume, Bijdr. 59 = glomerata.
densifolia, *A. St. Hil. Fl. Bras. Mer.* ii. 41.—Bras.
depauperata, Wall. Cat. n. 4180 = glomerata.
depressa, Wender. in Schrift. Marb. ii. (1831) 239; et in Berl. Jahrb. Pharm. xxxii. (1831) 109 = serpyllacea.
dichroma, Steud. Nom. ed. II. ii. 371 = Acanthocladus albicans.
dictyoptera, Boiss. Diagn. Ser. II. i. 59 = irregularis.
Dietrichiana, *D. Dietr. Syn. Pl.* iv. 905.—Afr. austr.
diffusa, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* ii. (1834) 114.—Afr. austr.
discolor, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 199 = leptalea.
disperma, *Vell. Fl. Flum.* 290; vii. t. 58.—Bras.
distans, *A. St. Hil. Fl. Bras. Mer.* ii. 24. t. 84.—Am. austr.
diversifolia, Linn. Sp. Pl. 703 = Badiera diversifolia.
diversifolia, Miq. in Linnaea, xix. (1847) 128 = longicaulis.
diversifolia, Pohl, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 7 = P. Laureola.
domingensis, Bert. ex Spreng. Syst. iii. 172 = Badiera Berteriana.
domingensis, *Jacq. Select. Am. ed. Min.* 252.—Ins. S. Doming.
Donii, Hook. f. in Hook. Niger Fl. 222 = tenuicaulis?
Dregeana, *Presl, Bot. Bemerk.* 14.—Afr. austr.
Duarteana, *A. St. Hil. Fl. Bras. Mer.* ii. 34.—Am. austr.
dubia, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 344 = vulgaris.

POLYGALA:—
dumosa, Poir. Encyc. v. 495 = Muraltia dumosa.
Dunaliana, A. St. Hil. Fl. Bras. Mer. ii. 32. t. 85 = brasiliensis.
dunensis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 344 = vulgaris.
dura, Wight, ex Wall. Cat. n. 4171 = glaucoides.
Eckloniana, Lehm. Del. Sem. Hort. Hamb. (1830) 8. —Cf. Linnaea, vi. (1831) Litt. 11 = ericaefolia.
Eckloniana, Presl, Bot. Bemerk. 15 = teretifolia.
elegans, Wall. Cat. n. 4186 = sibirica.
elongata, *Klein, ex Willd. Sp. Pl.* iii. 879.—Ind. or.
elongata, Presl, Fl. Sic. p. xvi, 136 = rosea.
emirnensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 89. —Madag.
empetrifolia, Houtt. Handleid. v. 433. t. 28. f. 1 = teretifolia.
ephedroides, Burch. Trav. i. 465 = virgata.
equisetoides, *A. St. Hil. Fl. Bras. Mer.* ii. 7.—Bras.
ericaefolia, *DC. Prod.* i. 323.—Afr. austr.
ericaefolia, Harv. in Harv. & Sond. Fl. Cap. i. 87, partim = macra.
ericoides, Burm. f. Fl. Cap. Prod. 19 = Muraltia Burmanni.
ericoides, Poir. Encyc. v. 497 = Muraltia Poiretii.
eriocephala, *F. Muell. ex Benth. Fl. Austral.* i. 139.—Austral.
erioptera, *DC. Prod.* i. 326.—As. et Afr. trop.
erubescens, E. Mey. in Drège, Zwei Pfl. Docum. 212 = hispida.
eumekes, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 172 = elongata.
eutaxioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 348.—Chili.
exigua, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 17.—Bras.
exigua, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 173 = chinensis.
exilis, *DC. Cat. Hort. Monsp.* 133.—Europ.
fasciculata, Poir. Encyc. v. 502 = Muraltia fasciculata.
fastigiata, *Nutt. Gen. Am.* ii. 89.—Am. bor.
fatua, Wallr. in Linnaea, xiv. (1840) 613; et Beitr. Fl. Hercyn. 243 (Quid?).—Europ.
Figariana, *Webb, Fragm. Fl. Aethiop.* 31.—Afr. trop.
filiformis, *A. St. Hil. Fl. Bras. Mer.* ii. 7.—Bras.
filiformis, Thunb. Prod. Pl. Cap. 121 = Muraltia filiformis.
fimbriata, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 13. —Bras.
Fimutu, J. F. Gmel. Syst. 1606 = P. Timoutou.
Fishiae, *Parry, in Proc. Davenp. Acad. Nat. Hist.* iv. (1884) 39 (sp. dub.).—Am. bor.
flabellata, Shuttlew. ex A. Gray, Pl. Wright. i. 41 = grandiflora.
flavescens, *DC. Cat. Hort. Monsp.* 134.—Europ. austr.
floribunda, *Benth. Pl. Hartw.* 58.—Am. austr.
floribunda, Boiss. & Huet, in Boiss. Diagn. Ser. II. v. 50 = major.
forojulensis, *Kern. in Oestr. Bot. Zeitschr.* xxiv. (1874) 102.—Europ.
fragilis, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 301 = gnidioides.
frutescens, Burm. f. ex DC. Prod. i. 308 = Ionidium heterophyllum.
fruticosa, Berg. Desc. Pl. Cap. 183 = oppositifolia.
furcata, Royle, Illustr. Bot. Himal. 76. t. 19 B = triphylla.
galapageia, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 233.—Ins. Galap.
galioides, Poir. Encyc. v. 503 = verticillata.
Garcini, *DC. Prod.* i. 323.—Afr. austr.
Gardneriana, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 19.—Bras.
Gariodiana, Jord. & Fourr. ex Verl. in Bull. Soc. Stat. Isère, xiv. (1872) 392 = nicaeënsis.
Gayii, *A. W. Benn. in Journ. Bot.* xvii. (1879) 168.—Am. austr.
genistoides, Poir. Encyc. v. 492 = virgata.
Gerardiana, Wall. Cat n. 4187 = abyssinica.
glabra, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 15.—Bras.
glabra, Heyne, ex A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 204 = chinensis.
glacialis, Bruegg. in Zeitschr. Ferdinand. Tirol. III. ix. (1860) 73 = alpina.
glandulosa, *H. B. & K. Nov. Gen. et Sp.* v. 404.—Mexic.
glandulosa, Vell. Fl. Flum. 293; vii. t. 70 = Monnina resedoides.

POLYGALA:—
glauca, E. Mey. in Drège, Zwei Pfl. Docum. 212, nomen = oppositifolia, myrtifolia.
glaucescens, Wall. Cat. n. 4182 = triphylla.
glaucoides, *Linn. Sp. Pl.* 705.—Zeylan.
glaucoides, Wight, in Wall. Cat. n. 4171 = chinensis.
globosa, Pohl, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 6 = violoides.
glochidiata, *H. B. & K. Nov. Gen. et Sp.* v. 400.—Venezuela.
glomerata, *Lour. Fl. Cochinch.* 426.—As. trop.
glomerata, Thw. Enum. Pl. Zeyl. 400 = sibirica.
glumacea, Sibth. & Sm. Fl. Graec. Prod. ii. 52 = monspeliaca.
gnidioides, *Willd. Sp. Pl.* iii. 878.—Chili; Peruv.
Gomesiana, *Welw. ex Oliver, Fl. Trop. Afr.* i. 126.—Afr. trop.
gracilipes, *Harv. in Harv. & Sond. Fl. Cap.* i. 90.—Afr. austr.
gracilis, *H. B. & K. Nov. Gen. et Sp.* v. 401.—Venezuela.
gracillima, Spruce, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 39 = appressa.
graminifolia, Poir. Encyc. v. 500 = cymosa.
grandiflora, Lodd. Bot. Cab. t. 1227 = myrtifolia.
grandiflora, *Walt. Fl. Carol.* 179.—Am. bor.
grandiflora, Wight, in Wall. Cat. n. 4175 = chinensis.
grandiflora, Willd. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 9 = seq.
grandifolia, *A. St. Hil. Fl. Bras. Mer.* ii. 52.—Bras.
grandis, Pall. Reise Suedl. Statth. Russ. Reich. ii. 190 = major?
granulata, Hochst. in Flora, xxiv. (1841) I. Intell. 30 = persicariaefolia.
Greggii, S. Wats. in Proc. Am. Acad. xvii. (1881-82) 325 = glandulosa.
guineensis, *Willd. Sp. Pl.* iii. 882.—Afr. trop.
Gundelsheimeri, C. Koch, in Linnaea, xix. (1847) 59 = supina.
gypsophiloides, Presl, ex Willk. & Lange, Prod. Fl. Hisp. iii. 558 = rosea.
hebantha, *Benth. Bot. Voy. Sulph.* 67.—Am. austr.
hebecarpa, *DC. Prod.* i. 330.—Ins. S. Doming.
hebeclada, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 10.—Bras.
hebeclada, *DC. Prod.* i. 331.—Bras.
Heisteria, Linn. Sp. Pl. 704 = Muraltia Heisteria.
hemipterocarpa, *A. Gray, Pl. Wright.* ii. 31.—N. Mexic.
herbiola, *A. St. Hil. Fl. Bras. Mer.* ii. 12.—Bras.
heterophylla, *Scheele, in Linnaea*, xvii. (1843) 336.—Bras.
heterophylla, F. Schultz, Fl. Pfalz (1846), 72, in syn. = calcarea.
heterotricha, Zipp. ex Span. in Linnaea, xv. (1841) 167 = humilis.
Heyneana, Wall. Cat. n. 4184 = sibirica.
Hilariana, Endl. in Linnaea, vii. (1832) 357 = grandifolia.
hirsuta, *A. St. Hil. Fl. Bras. Mer.* ii. 45.—Bras.
hirsutula, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 322 = glaucoides.
hispida, *Burch. ex DC. Prod.* i. 323.—Afr. austr.
hispida, E. Mey. in Drège, Zwei Pfl. Docum. 212 = affinis.
hispidula, Presl, Bot. Bemerk. 15 = peduncularis.
Hohenackeriana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 42.—Oriens.
Hookeri, *Torr. & Gray, Fl. N. Am.* i. 671.—Am. bor. occ.
hospita, Heuff. in Flora, xxxvi. (1853) 620 = supina.
Hostmanni, Miq. Stirp. Surin. Sel. 33 = trichosperma.
hottentotta, *Presl, Bot. Bemerk.* 15.—Afr. austr.
huillensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 133.—Afr. trop.
humilis, Lodd. Bot. Cab. t. 420 = Muraltia humilis.
humilis, *Span. in Linnaea*, xv. (1841) 167.—Ins. Timor.
humilis, Vell. Fl. Flum. 293; vii. t. 67 = paniculata.
hyalina, Wall. Cat. n. 4181 = triphylla.
hybrida ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 73.—Europ.
hybrida, DC. Prod. i. 325 = vulgaris.
hyemalis, Rafin. New Fl. Am. iv. 89 (Quid?).—Am. bor.
hygrophila, *H. B. & K. Nov. Gen. et Sp.* v. 395.—Venezuela.

POLYGALA:—
hypericoides, Webb, Fragm. Fl. Aethiop. 31 = persicariaefolia.
hypoglauca, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 169 = glaucoides.
hyssopifolia, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1842) 13; *et in Ann. Sc. Nat.* Sér. II. xx. (1843) 95.—Madag.
hyssopifolia, A. St. Hil. Consp. Polyg. Bras. Mer. 12; Fl. Bras. Mer. ii. 43 = hebeclada, *DC.*
ilheotica, *Wawra*, *Bot. Ergeb. Maxim. Bras.* 41. t. 54.—Bras.
illepida, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 92.—Afr. austr.
inaequiloba, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 349.—Mexic.
incarnata, Aubl. Pl. Gui. ii. 739 = adenophora.
incarnata, *Linn. Sp. Pl.* 701.—Am. bor.
insignis, *Klotzsch, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 8.—Bras.
insipida, Reichb. Fl. Germ. Excurs. 350 = amara.
insularis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 204.—Ins. Galapag.
intermedia, DC. Prod. i. 322 = peduncularis.
intermedia, Eckl. & Zeyh. Enum. 21 = bracteolata, *Linn.*
intermedia, Eichw. Skizze, 174 = vulgaris.
involutiflora, Lamotte, Prod. Centr. Fr. i. 126 = vulgaris.
irregularis, *Boiss. Diagn.* Ser. I. i. 8.—Oriens; Afr. trop.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxxii. (1882) 38.—Bras.
Jablotskiana, Fisch. & Mey. ex Boiss. Fl. Orient. i. 472 = Hohenackeriana.
Jamesi, Rafin. Atl. Journ. 146 (Quid?).—Am. bor.
japonica, Houtt. Handleid. x. 89. t. 62. f. 1 = sibirica.
javana, *DC. Prod.* i. 327.—Ind. or.; Malaya.
javanica, Steud. Nom. ed. II. ii. 371 = praec.
juncea, *A. St. Hil. Fl. Bras. Mer.* ii. 13.—Bras.
juniperifolia, Poir. Encyc. v. 496 = Muralтia juniperifolia.
juniperina, Cav. in Anal. Cienc. Nat. iv. (1801) 53 = microphylla.
karensium, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 392.—Burma.
khasiana, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 176 = sibirica.
Kleinii, Hassk. l. c. 158 = chinensis.
krumanina, *Burch. ex Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 16.—Afr. austr.
laevigata, E. Mey. in Drège, Zwei Pfl. Docum. 212 = serpentaria.
lagoana, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 22.—Bras.
lanata, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. i. (1842) 469 = hispida.
lanceolata, Poir. Encyc. v. 498 = Monnina lanceolata.
lanceolata, *Vell. Fl. Flum.* 291; vii. t. 63.—Bras.
lancifolia, *A. St. Hil. Fl. Bras. Mer.* ii. 42.—Bras.
lasiopoda, Presl, Bot. Bemerk. 15 = bracteolata, *Linn.*
latifolia, Ker-Gawl. in Bot. Reg. t. 645 = oppositifolia.
Laureola, *A. St. Hil. Fl. Bras. Mer.* ii. 50. t. 89.—Bras.
laxa, *Mart. ex Spreng. Syst.* iii. 170.—Bras.
laxa, Thunb. Prod. Pl. Cap. 121 = Muralтia laxa.
Lehmanniana, *Eckl. & Zeyh. Enum.* 23.—Afr. austr.
Lejeunii, Bor. Fl. Centr. Fr. ed. II. 71 = vulgaris.
Lensei, Bor. in Mém. Soc. Acad. Maine-et-Loire, xiv. (1863) 42 = vulgaris.
leptalea, *DC. Prod.* i. 325.—As. or.; Austral.
leptandroides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 349.—Mexic.
leptocaulis, Torr. & Gray, Fl. N. Am. i. 130 = paludosa.
leptoclada, *Boj. in Bouton*, 12me *Rapp. Ann. Maur.* (1842) 14.—Madag.
leptophylla, *Burch. Trav.* i. 400.—Afr. austr.
leptorhiza, DC. Prod. i. 326 = erioptera.
leptostachys, *Shuttlew. ex A. Gray, in Pl. Wright.* i. 41.—Am. bor.
leucantha, *A. W. Benn. in Journ. Bot.* xvii. (1879) 172.—Am. austr.
ligularis, Ker-Gawl. in Bot. Reg. t. 637 = myrtifolia.
ligularis, Spreng. ex Eckl. & Zeyh. Enum. 19 = virgata.
ligustroides, *A. St. Hil. Fl. Bras. Mer.* ii. 49.—Bras.

POLYGALA:—
linariaefolia, Poir. Encyc. v. 495 = chinensis.
linarifolia, Buch.-Ham. ex Wall. Cat. n. 4170 E. = telephioides.
linarifolia, Heyne, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 175 = rosmarinifolia.
linarifolia, Willd. Sp. Pl. iii. 877 = chinensis.
Lindheimeri, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 150.—Am. bor. occ.
linearis, R. Br. in Salt, Abyss. App. 65 = erioptera.
linearis, Heyne, ex Wight & Arn. Prod. 38 = chinensis.
linearis, Hook. f. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 37 = Duarteana.
linearis, Lag. Gen. et Sp. Nov. 22 = exilis.
linearis, E. Mey. in Drège, Zwei Pfl. Docum. 212 = rarifolia.
linifolia, Hook. f. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 37 = Duarteana.
linifolia, Scheele, in Linnaea, xvii. (1843) 335 = Nuttallii?
Linkiana, Sweet, Hort. Brit. ed. II. 46 = myrtifolia.
linoides, *Poir. Encyc.* v. 449.—Bras.
Liorani, Jord. ex Lamotte, Prod. Fl. Cent. Fr. i. 127 = serpyllacea.
longicaulis, *H. B. & K. Nov. Gen. et Sp.* v. 396.—Am. trop.
longifolia, A. Dietr. in Otto & Dietr. Allg. Gartenz. ii. (1834) 117 = macrophylla.
longifolia, Gilib. ex Steud. Nom. ed. II. ii. 372 = vulgaris.
longifolia, *Poir. Encyc.* v. 501.—Java.
longifolia, Presl, Bot. Bemerk. 15 = virgata.
Lourerii, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 242 = sibirica.
lucida, A. W. Benn. in Mart. Fl. Bras. xiii. III. 43 = P. Roubienna.
Ludwigiana, *Eckl. & Zeyh. Enum.* 23.—Afr. austr.
lupulina, Willd. Sp. Pl. iii. 880 = P. Timoutou.
lutea, *Linn. Sp. Pl.* 705.—Am. bor.
macra, *DC. Prod.* i. 323.—Afr. austr.
macra, Eckl. & Zeyh. Enum. 22 = ciliatifolia.
macra, Harv. in Harv. & Sond. Fl. Cap. i. 89, partim = refracta.
macra, E. Mey. in Drège, Zwei Pfl. Docum. 212 = virgata.
macradenia, *A. Gray, Pl. Wright.* i. 39.—Am. bor. occ.
macrantha, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 345 = oppositifolia.
macrolophos, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 167 = sibirica.
macropetala, Hassk. l. c. 168 = sibirica?
macrophylla, *D. Dietr. Syn. Pl.* iv. 905.—Afr. austr.
macrophylla, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 9 = grandifolia.
macroptera, *DC. Prod.* i. 332.—Madag.
macrostachya, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 171 = elongata.
madagascariensis, *O. Hoffm. in Bremen, Abh.* vii. (1882) 240.—Madag.
magna, Georgi, Beschr. Russ. Reich. III. iv. 1159 = major.
major, *Jacq. Fl. Austr.* v. 6.—Europ. austr.; Oriens.
major, Webb, Iter Hisp. 66 = Boissieri.
Mannii, *Oliver, Fl. Trop. Afr.* i. 130.—Afr. trop.
mariana, Mill. Gard. Dict. ed. VIII. n. 6 = incarnata.
maritima, *Vell. Fl. Flum.* 293; vii. t. 68.—Bras.
Marta, Steud. Nom. ed. II. ii. 372, sphalm. = Martii.
Martiana, A. W. Benn. in Mart. Fl. Bras. xiii. III. 13 = violacea.
Martii, *DC. Prod.* i. 332.—Bras.
mascatensis, *Boiss. Fl. Orient.* i. 470.—Arab.
mendocina, *Phil. Sert. Mend. Alt.* 4.—Chili.
mexicana, Moç. ex Cav. in Anal. Cienc. Nat. vii. (1804) 48 = scoparia.
Meyeriana, Presl, Bot. Bemerk. 14 = oppositifolia.
Michaleti, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 395 = vulgaris.
micrantha, Andr. Bot. Rep. t. 424 = Muralтia filiformis.
micrantha, *Guill. & Perr. Fl. Seneg. Tent.* i. 37.—Afr. trop.
micrantha, Thunb. Prod. Pl. Cap. 121 = Muralтia micrantha, filiformis.
micrantha, Willd. ex DC. Prod. i. 336 = Muralтia stipulacea.
microcarpa, *Gaud. Fl. Helv.* iv. 445.—Europ.; Reg. Caucas.

POLYGALA:—
microlopha, *Burch. ex DC. Prod.* i. 322.—Afr. austr.
microlopha, Eckl. & Zeyh. Enum. 22 = ericaefolia.
microphylla, *Linn. Sp. Pl.* ed. II. 989.—Lusitan.
microphylla, Thunb. Prod. Pl. Cap. 121 = Thunbergii.
microptera, A. W. Benn. in Hemsl. Diagn. Pl. Nov. 2 = incarnata.
minima, *Pohl, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 30.—Bras.
minutiflora, Presl, Rel. Haenk. ii. 100 = scoparia.
missurica, Rafin. New Fl. Am. iv. 89 (Quid?).—Am. bor.
mixta, Andr. Bot. Rep. t. 445 = Muralтia mixta.
mixta, Linn. f. Suppl. 316; Thunb. Prod. Pl. Cap. 121 = Muralтia Burmanni.
modesta, Miq. in Linnaea, xviii. (1844) 747 = tenuis.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 406.—Am. austr.
molluginifolia, A. St. Hil. Fl. Bras. Mer. ii. 25 = verticillata.
monninoides, H. B. & K. Nov. Gen. et Sp. v. 408 = angustifolia.
monopetala, Cambess. in Jacquem. Voy. Bot. 25. t. 27 = sibirica.
monosperma, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 19.—Bras.
monspeliaca, All. Fl. Pedem. ii. 301; Vill. Hist. Pl. Dauph. iii. 388 = vulgaris.
monspeliaca, Asso, Syn. Pl. Arag. 92 = rupestris.
monspeliaca, Blanco, Fl. Filip. ed. I. 557 = telephioides.
monspeliaca, *Linn. Sp. Pl.* 702.—Reg. Mediterr.
monspeliaca, Reichb. Fl. Germ. Excurs. 351 = exilis.
montana, Opiz, in Flora, v. (1822) 270 = vulgaris.
monticola, H. B. & K. Nov. Gen. et Sp. v. 405 = angustifolia.
Moquiniana, *A. St. Hil. Fl. Bras. Mer.* ii. 36. t. 86.—Bras.
Mori, Opiz, Natural. xi. (1826) 307, sphalm. = seq.
Moriana, Britting. in Flora, ix. (1826) 729 = vulgaris.
Morisiana, Reichb. f. Ic. Fl. Germ. xviii. 91 = nicaeënsis.
mucronata, *Baker, in Journ. Linn. Soc.* xx. (1883) 89.—Madag.
mucronata, A. W. Benn. in Mart. Fl. Bras. xiii. III. 8 = insignis.
mucronata, *Willd. Sp. Pl.* iii. 888.—Am. austr.
Muhlenbergii, G. Don, Gen. Syst. i. 358 = grandiflora.
multibracteata, Wall. Cat. n. 4174 = erioptera.
multicaulis, Tausch, in Flora (1821), 563 = vulgaris.
multicaulis, *Vell. Fl. Flum.* 294; vii. t. 71.—Bras.
multiceps, *Borb. ex Just, Jahresb.* x. 1882 (1885), 539.—Europ. austr.
multiceps, *Mart. ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 43.—Bras.
multiflora, *Poir. Encyc.* v. 497.—Afr. trop.
Munbyana, *Boiss. & Reut. Diagn.* Ser. II. v. 50.—Alger.
Mundiana, Eckl. & Zeyh. Enum. 22 = ericaefolia.
muraltioides, Eckl. & Zeyh. l. c. 24 = Muralтia Beiliana.
mutabilis, Dum. Fl. Belg. Prod. 31 = serpyllacea.
Myrsinites, Royle, Illustr. Bot. Himal. 76. t. 19 A = sibirica.
myrtifolia, Fries, Nov. Fl. Suec. ii. 227 = amara.
myrtifolia, *Linn. Sp. Pl.* 703.—Afr. austr.
myrtillifolia, *Nees, 'Samml.* t. 48'; *ex Pritz. Ic. Ind.* ii. 222 (Quid?).—Hab.?
myrtilloides, *Willd. Sp. Pl.* iii. 889.—Am. austr.
myrtillopsis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 130.—Afr. trop.
nana, *DC. Prod.* i. 328.—Am. bor.
Neaei, DC. l. c. 329 = aspalatha.
neglecta, Kern. in Oestr. Bot. Zeitschr. xviii. (1868) 37 = major.
nemoralis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 172.—Am. austr.
nemorivaga, *Pomel, Nouv. Mat. Fl. Atl.* 337.—Alger.
nemorosa, Willd. ex Steud. Nom. ed. II. ii. 372 = americana.
nicaeënsis, *Risso, ex Koch, Syn. Fl. Germ.* ed. I. 92.—Europ.; Reg. Mediterr.
nicaeënsis, Schur, Enum. Pl. Trans. 89 = major.
nicaeënsis, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 559 = baetica.

POLYGALA:—
nivea, Miégev. in Bull. Soc. Bot. Fr. xii. (1865) 341 = amara.
noucherensis, Cambess. in Jacquem. Voy. Bot. 24 = erioptera.
nova, Boissieu, Fl. Europ. i. t. 174. f. 1 = exilis.
nubica, Hochst. ex Oliver, Fl. Trop. Afr. i. 129 = erioptera.
nudicaulis, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 20.—Bras.
numidica, Pomel, Nouv. Mat. Fl. Atl. 211 = nicaeënsis.
nummularia, Burch. ex DC. Prod. i. 322 = oppositifolia.
nutans, Hook. f. in Hook. Niger Fl. 222 = arenaria.
nutkana, [Moç. & Sesse,] ex DC. Prod. i. 330 = ovalifolia.
nutkana, Torr. Bot. Mex. Bound. 49. t. 12 = cucullata.
Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 670.—Am. bor.
obovata, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 233. —Ins. Galap.
obovata, *A. St. Hil. Fl. Bras. Mer.* ii. 37.—Am. austr.
obscura, *Benth. Pl. Hartw.* 58.—Mexic.
obtusata, DC. Prod. i. 326 = erioptera.
obtusissima, Hochst. ex Oliver, Fl. Trop. Afr. i. 130 = senensis.
Ohlendorfiana, *Eckl. & Zeyh. Enum.* 22.—Afr. austr.
oleaefolia, *A. St. Hil. Fl. Bras. Mer.* ii. 49.—Bras.
oligantha, A. Rich. Tent. Fl. Abyss. i. 38 = erioptera.
oligophylla, DC. Prod. i. 325 = leptalea.
oppositifolia, *Linn. Mant.* ii. 259.—Afr. austr.
orbicularis, *Benth. Fl. Austral.* i. 140.—Austral.
ovalifolia, *DC. Prod.* i. 331.—Mexic.
ovalis, E. Mey. in Drège, Zwei Pfl. Docum. 212 = Ohlendorfiana.
ovata, Poir. Encyc. v. 498 = hebecarpa.
ovatifolia, A. Gray, Pl. Wright. i. 39, sphalm. = ovalifolia.
oxyococcoides, Desf. Fl. Atlant. ii. 127 = rupestris.
oxyphylla, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 8.—Bras.
oxyphylla, DC. Prod. i. 331 = P. Laureola.
oxyptera, Reichb. Pl. Crit. i. 25 = vulgaris.
pachyrrhiza, Casar. Nov. Stirp. Bras. Dec. 20 = violoides.
pallida, *E. Mey. in Drège, Zwei Pfl. Docum.* 212.—Afr. austr.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 325.—Am. bor. occ.
paludosa, Boenn. ex Reichb. Fl. Germ. Excurs. 351 = serpyllacea.
paludosa, *A. St. Hil. Fl. Bras. Mer.* ii. 8.—Bras.
paniculata, Forsk. Fl. Aegypt. Arab. p. cxvii = exilis.
paniculata, Le Conte, ex Torr. & Gray, Fl. N. Am. i. 129 = incarnata.
paniculata, *Linn. Syst.* ed. X. 1154.—Ind. occ.; Afr. trop.
papilionacea, *Boiss. Diagn.* Ser. I. i. 8.—Armen.
Pappeana, Eckl. & Zeyh. Enum. 23 = macra.
paraguayensis, A. W. Benn. in Journ. Bot. xvii. (1879) 173 = adenophylla.
Parryi, *A. W. Benn. in Hemsl. Diagn. Pl. Nov.* 21.—Mexic.
parviflora, Loisel. in Desv. Journ. Bot. ii. (1809) 360 = exilis.
parviflora, Poir. Encyc. v. 504 = Muraltia parviflora.
parviflora, Reichb. f. ex Nym. Consp. 83 = vulgaris.
pauciflora, Muhl. ex Willd. Sp. Pl. iii. 880 = paucifolia.
pauciflora, Thunb. Prod. Pl. Cap. 121 = Muraltia pauciflora.
paucifolia, *Willd. Sp. Pl.* iii. 880.—Am. bor.
Pearcei, *A. W. Benn. in Journ. Bot.* xvii. (1879) 201. —Bolivia.
pedemontana, Perrier, ex B. Verlot, in Rév. Hortic. (1863) 433 = vulgaris.
pedicellaris, A. St. Hil. Fl. Bras. Mer. ii. 47 = violoides.
peduncularis, *Burch. ex DC. Prod.* i. 323.—Afr. austr.
peduncularis, Eckl. & Zeyh. Enum. 20 = virgata.
peduncularis, A. Rich. Ess. Fl. Cub. 106 = violacea.
pedunculata, Harv. in Harv. & Sond. Fl. Cap. i. 86, in syn. = abyssinica.
pedunculosa, Thw. Enum. Pl. Zeyl. 400 = sibirica.
Penaea, Linn. Sp. Pl. 703 = Badiera Penaea.
perophylla, Pohl, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 18 = tenuis.
persicariaefolia, *DC. Prod.* i. 326.—As. Afr. et Austral. trop.

POLYGALA:—
persistens, *A. W. Benn. in Journ. Bot.* xvii. (1879) 170.—Chili.
peruviana, *A. W. Benn. l. c.* 173.—Peruv.
petalantha, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 5 = violoides.
Petitiana, *A. Rich. Tent. Fl. Abyss.* i. 37.—Abyss.
phrygia, Boiss. & Bal. ex Boiss. Fl. Orient. i. 473 = pruinosa.
phylicoides, Thunb. Prod. Pl. Cap. 121 = Muraltia phylicoides.
pilosa, *Baker, in Journ. Linn. Soc.* xxi. (1884) 321.—Madag.
pilosa, Poir. Encyc. v. 497 = Muraltia pilosa.
pinifolia, *Lam. Illustr.* iii. 159.—Afr. austr.
pinnata, Burm. f. Fl. Cap. Prod. 19 (Leguminosae sp.). —Afr. austr.
pisaurensis, *Caldesi, in Nuov. Giorn. Bot. Ital.* xi. (1879) 189.—Italia.
platycarpa, *Benth. Pl. Hartw.* 113.—Am. austr.
Poaya, Mart. Mat. Med. Bras. fasc. i. 13; Reise Bras. i. 285 = angulata.
podolica, DC. Prod. i. 325 = vulgaris.
Pohliana, A. St. Hil. Fl. Bras. Mer. ii. 45 = hirsuta.
polifolia, Presl, Rel. Haenk. ii. 101.—Bras.
polycephala, A. St. Hil. Fl. Bras. Mer. ii. 16 = aspalatha.
polygama, DC. Prod. i. 330 = grandiflora.
polygama, *Walt. Fl. Carol.* 179.—Am. bor.
polymorpha, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 865 = vulgaris, serpyllacea.
polyphylla, *DC. Prod.* i. 324.—Afr. austr.
polyphylla, Eckl. & Zeyh. Enum. 24 = Muraltia crassifolia.
pratensis, *A. W. Benn. in Journ. Bot.* xvii. (1879) 169. —Chili.
pratensis, *Phil. in Linnaea*, xxxiii. (1864) 18.—Chili.
Preslii, *D. Dietr. Syn. Pl.* iv. 904.—Peruv.
Preslii, Spreng. Syst. v. 551 = rosea.
procumbens, Roth, Nov. Pl. Sp. 329 = chinensis.
prostrata, F. Schultz, Fl. Pfalz (1846) 72, in syn. = serpyllacea.
prostrata, Willd. Sp. Pl. iii. 896 = chinensis.
pruinosa, *Boiss. Diagn.* Ser. I. i. 8.—Oriens.
pruinosa, Boiss. l. c., partim = nicaeënsis.
Pseudo-erica, *A. St. Hil. Fl. Bras. Mer.* ii. 40. t. 87.— Bras.
Pseudo-senega, Bertol. Misc. Bot. xv. 21 = lutea.
puberula, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 232.—Ins. Galapag.
puberula, *A. Gray, Pl. Wright.* i. 40.—Texas.
puberula, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 13 = violacea.
pubescens, Mart. in Denkschr. Bot. Ges. Regensb. i. (1815) 185 = violacea.
pubescens, Muhl. Cat. 66 = grandiflora.
pubescens, Rhode, ex Mutel, Fl. Fr. i. 129 = vulgaris.
pubescens, Schlecht. in Linnaea, xiv. (1840) 160 = puberula, *A. Gray*.
pubiflora, *Burch. ex DC. Prod.* i. 322.—Afr. austr.
pubiflora, Eckl. & Zeyh. Enum. 20 = bracteolata, *Linn.*
pulchella, *Cham. & Schlecht. in Linnaea*, v. (1830) 230.—Mexic.
pulchella, *A. St. Hil. Fl. Bras. Mer.* ii. 30.—Bras.
pulchella, Scheele, in Linnaea, xvii. (1843) 335 = glochidiata.
pulchra, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 32.— Java.
punctata, *A. W. Benn. in Journ. Bot.* xvii. (1879) 172. —Am. austr.
punctulata, Hochst. in Flora, xxix. (1841) I. Intell. 30, in obs. = persicariaefolia.
punctulata, *Vell. Fl. Flum.* 291; vii. t. 61.—Bras.
pungens, *Burch. Trav.* i. 304.—Afr. austr.
pungens, Drège, Zwei Pfl. Docum. 212 = ciliatifolia.
purpurea, Ait. Hort. Kew. ed. II. iv. 244 = paucifolia.
purpurea, Nutt. Gen. Am. ii. 88 = sanguinea.
pygmaea, Steud. Nom. ed. II. ii. 373 = crotalarioides.
pyxophylla, Reichb. Fl. Germ. Excurs. 350 = serpyllacea.
quadrangula, Presl, Rel. Haenk. ii. 100 = alba?
Quartiniana, Dillon, ex A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 263 = persicariaefolia.
quinqueangularis, Vell. Fl. Flum. 292; vii. t. 65 = angulata.

POLYGALA:—
quinqueflora, Buch.-Ham. ex Wall. Cat. n. 4170 C = telephioides.
quitensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 350.—Ecuador.
Raddiana, A. St. Hil. Fl. Bras. Mer. ii. 26 = exigua, glochidiata.
ramosa, *Ell. Sketch*, ii. 186.—Am. bor.
ramosa, Wight, in Wall. Cat. n. 4177 = chinensis.
ramosissima, *Cav. in Anal. Cienc. Nat.* iv. (1801) 53. —Ins. Teneriff.
ramulosa, *Boiss. & Bal. Diagn.* Ser. II. vi. 24.—As. Min.
rariflora, Steud. Nom. ed. II. ii. 372, sphalm. = seq.
rarifolia, *DC. Prod.* i. 332.—Afr. trop.
recta, Presl, Bot. Bemerk. 16 = leptophylla.
recurvata, Celak. in Oestr. Bot. Zeitschr. xxii. (1872) 109 = supina.
refracta, *Burch. ex DC. Prod.* i. 323.—Afr. austr.
refracta, Eckl. & Zeyh. Enum. 23 = illepida.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 292.—Japon.
remota, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 21. —Bras.
repens, Mérat, ex DC. Prod. i. 325 = amara.
reptans, Cheval. Fl. Par. iii. 850 = praec.?
resedoides, *A. St. Hil. Fl. Bras. Mer.* ii. 31.—Bras.
restiacea, E. Mey. in Drège, Zwei Pfl. Docum. 212 = macra.
reticulata, Presl, Delic. Prag. 228 = vulgaris.
retusa, Hochst. ex Oliver, Fl. Trop. Afr. i. 129 = erioptera.
revoluta, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 99.—Bras.
Reynoldsae, Chapm. Fl. S. U. St. Suppl. 613 = Rugelii.
rhinanthoides, *Soland. ex Benth. Fl. Austral.* i. 140.— Austral.
rhodoptera, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 11 = hebeclada.
rhombifolia, Eckl. & Zeyh. Enum. 18 = oppositifolia.
rigens, Burch. Trav. i. 465 = asbestina.
rigens, Eckl. & Zeyh. Enum. 22 = rarifolia.
rigens, E. Mey, in Drège, Zwei Docum. 212 = leptophylla.
rigida, *A. St. Hil. Fl. Bras. Mer.* ii. 18.—Bras.
rivinaefolia, H. B. & K. Nov. Gen. et Sp. v. 409 = americana.
Rogeri, Oertz. ex Tenore, Syll. Fl. Neap. 340 = flavescens.
rosea, Boiss. Voy. Bot. Espagne, ii. 81 = Boissieri.
rosea, *Desf. Fl. Atlant.* ii. 128. t. 176.—Reg. Mediterr.
rosea, Gren. & Godr. Fl. Fr. i. 194 = nicaeënsis.
rosea, Steud. Nom. ed. I. 643 = P. Senega.
rosmarinifolia, Eckl. & Zeyh. Enum. 20 = pedunculares.
rosmarinifolia, *Wight & Arn. Prod.* 37.—Ind. or.
Rothiana, Wight & Arn. l. c. = chinensis.
Rottleriana, Steud. Nom. ed. II. ii. 373 = violacea.
rotundifolia, Gilib. Fl. Lituan. ii. 113 = amara.
Roubienna, *A. St. Hil. Fl. Bras. Mer.* ii. 35.—Bras.
rubella, Willd. Sp. Pl. iii. 875 = polygama.
rufa, *Span. in Linnaea*, xv. (1841) 167.—Ins. Timor.
Rugelii, *Shuttlew. ex A. Gray, Pl. Wright.* i. 41.—Am. bor.
rugosa, Rafin. New Fl. Am. iv. 89 (Quid?).—Am. bor.
rupestris, *Pourr. in Mém. Acad. Toul.* iii. (1788) 325. —Reg. Mediterr. occ.
rupicola, *Hochst. & Steud. in Flora*, xxiv. (1841) I. *Intell.* 30.—Abyssin.
rupicola, Pomel, Nouv. Mat. Fl. Atl. 213 = rupestris.
Rusbyi, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 125.—Am. bor.
Rutenbergii, *O. Hoffm. in Bremen, Abh.* vii. (1882) 239.—Madag.
sabulosa, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 27. —Bras.
saginoides, *Griseb. Cat. Pl. Cub.* 13.—Cuba.
salasiana, *C. Gay, Fl. Chil.* i. 237.—Chili.
salicifolia, Poir. Encyc. v. 498 = Monnina longifolia.
Salviniana, *A. W. Benn. in Journ. Bot.* xvii. (1879) 203.—Am. centr.
sanguinea, *Linn. Sp. Pl.* 705.—Am. bor.
sanguinea, Nutt. Gen. Am. ii. 88 = Nuttallii.
sanguinea, Torr. & Gray, Fl. N. Am. i. 126 = fastigiata.

POLYGALA:—
saxatilis, Desf. Fl. Atlant. ii. 128. t. 175 = rupestris.
saxatilis, Portenschl. ex Willk. & Lange, Prod. Fl. Hisp. iii. 558 = rosea.
scabra, Edgew. in Trans. Linn. Soc. xx. (1846) 33 = abyssinica.
scabra, Linn. Sp. Pl. 703 = bracteolata, *Linn.*
scandens, *Vell. Fl. Flum.* 291; vii. t. 60.—Bras.
Schimperi, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 178 = erioptera.
Schimperi, Presl, Bot. Bemerk. 14 = rosea?
Schoenlankii, *O. Hoffm. & Hildebr. ex O. Hoffm. Sert. Madag.* 6.—Ins. Mascar.
Schreberi, Pall. ex Georgi, Beschr. Russ. Reich. Nachtr. 294 = supina.
sciophila, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 12 = ilheotica.
scoparia, Benth. Pl. Hartw. 8 = alba.
scoparia, *H. B. & K. Nov. Gen. et Sp.* v. 399.—Mexic.
sedoides, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 27.—Guian.
selaginoides, *A. W. Benn. l. c.* 38.—Bras.
semialata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 326.—Mexic.
seminuda, Harv. in Harv. & Sond. Fl. Cap. i. 86 = hottentotta.
senduaris, Buch.-Ham. ex Wall. Cat. n. 4170 B = telephioides.
Senega, *Linn. Sp. Pl.* 704.—Am. bor.
Seneka, Hill, Veg. Syst. xxii. 52 = praec.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 113.—Afr. trop.
sericea, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 38.—Bras.
serpentaria, *Eckl. & Zeyh. Enum.* 24.—Afr. austr.
serpillifolia, Poir. Encyc. v. 499 = erioptera, telephioides.
serpyllacea, *Weihe, in Flora*, ix. (1826) 745.—Europ.
serpyllifolia, Fischer-Oost. l. c. xxxvii. (1854) 98 = serpyllacea, amara.
serpyllifolia, Hose, in Usteri, Ann. Bot. xxi. (1797) 39 = serpyllacea?
serpyllifolia, Wight, in Wall. Cat. n. 4173 = erioptera.
setacea, *Michx. Fl. Bor. Am.* ii. 52.—Am. bor.
sherardioides, Pohl, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 29 = distans.
sibirica, *Linn. Sp. Pl.* 702.—As. temp. et trop.
Sieboldiana, Miq. in Versl. en Med. Kon. Akad. Wetensch. II. ii. (1868) 85 = sibirica.
Simassan, *Miq. Fl. Ind. Bat. Suppl.* 392.—Sumatra.
simplex, Burch. ex DC. Prod. i. 322 = virgata.
sinensis, J. F. Gmel. Syst. 1082 = chinensis.
Solierii, *C. Gay, Fl. Chil.* i. 238.—Chili.
sororia, Miq. Stirp. Surin. Sel. 32. t. 8 = variabilis.
sparsiflora, *Oliver, Fl. Trop. Afr.* i. 127.—Afr. trop.
spartioides, Burch. ex Eckl. & Zeyh. Enum. 20 = virgata.
spathulata, *Griseb. Cat. Pl. Cub.* 13.—Cuba.
spathulata, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid?).—Am. bor.
speciosa, Sims, Bot. Mag. t. 1780 = virgata.
spectabilis, *DC. Prod.* i. 331.—Am. austr.
spergulaefolia, A. St. Hil. Fl. Bras. Mer. ii. 28 = glochidiata.
sphenoptera, Fresen. in Mus. Senckenb. ii. (1837) 274 = persicariaefolia.
spilophylla, Steud. Nom. ed. II. ii. 373 = persicariaefolia.
spinescens, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 275.—Arab.
spinescens, Gill. in Hook. Bot. Misc. iii. (1833) 146 = scoparia.
spinosa, Linn. Sp. Pl. 704 = Mundtia spinosa.
Sprengeliana, Eckl. & Zeyh. Enum. 19 = virgata.
Spruceana, *A. W. Benn. in Journ. Bot.* xvii. (1879) 203.—Am. austr.
spuria, Stev. ex Ledeb. Fl. Ross. i. 271 = vulgaris.
squamifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 12.—Cuba.
squarrosa, Linn. f. Suppl. 315 = Muralтia squarrosa.
squarrosa, Soland. ex Benth. Fl. Austral. i. 141 = chinensis.
Stellera, DC. Prod. i. 327 = longicaulis.
stenoclada, *Benth. Fl. Austral.* i. 141.—Austral.
stenopetala, Klotzsch, in Peters, Reise Mossamb. Bot. 114 = rarifolia.
stenophylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 108.—Patagon.

POLYGALA:—
stenophylla, Klotzsch, in Peters, Reise Mossamb. Bot. t. 23 = rarifolia.
Steudeliana, Eckl. & Zeyh. Enum. 23 = refracta.
stipulacea, Burch. Trav. i. 19, 29 = Muralтia stipulacea.
stipulacea, Burm. f. Fl. Cap. Prod. 19 = Muralтia Heisteria.
stipulacea, Linn. Mant. ii. 260 = Muralтia stipulacea?
Stocksiana, Boiss. Diagn. Ser. II. i. 59 = Hohenackeriana.
straminea, Presl, Fl. Sicul. 136 = monspeliaca.
striata, Thunb. Prod. Pl. Cap. 121 = Muralтia striata.
stricta, A. W. Benn. in Mart. Fl. Bras. xiii. III. 13, partim = bryoides.
stricta, Endl. ex A. W. Benn. l. c. = violacea.
stricta, C. Gay, Fl. Chil. i. 236 = Gayii.
stricta, *A. St. Hil. Fl. Bras. Mer.* ii. 18.—Bras.
strigulosa, *Schlecht. in Linnaea*, xiv. (1840) 150.—Mexic.
subandina, *Phil. l. c.* xxxiii. (1864–65) 17.—Chili.
subspinosa, *S. Wats. in Am. Natural.* vii. (1873) 299.—Am. bor. occ.
subtilis, *H. B. & K. Nov. Gen. et Sp.* v. 393.—Am. austr.
subulata, E. Mey. in Drège, Zwei Pfl. Docum. 212 = bracteolata, *Linn.*
subuniflora, *Boiss. & Heldr. Diagn.* Ser. II. i. 59.—Graec.
sulcata, Willd. Sp. Pl. iii. 878 = variabilis.
sulphurea, A. W. Benn. in Mart. Fl. Bras. xiii. III. 17 = tenuis.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 392.—Sumatra.
supina, *Schreb. Ic. Desc. Pl. Dec.* 19. t. 10.—Oriens.
Szovitsiana, *Stev. ex Rupr. Fl. Cauc.* 295.—Reg. Caucas.
tamariscea, *Mart. ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 37.—Bras.
Tatarinowii, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) II. 523 = triphylla.
telephioides, Boiss. & Bal. Diagn. Ser. II. v. 49 = supina.
telephioides, *Willd. Sp. Pl.* iii. 876.—Ind. or.; Malaya.
tenella, Willd. l. c. 878 = paniculata.
tenuicaulis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 182.—Afr. trop.
tenuifolia, Link, Enum. Hort. Berol. ii. 220 = rarifolia.
tenuifolia, Poir. Encyc. v. 497 = Muralтia tenuifolia.
tenuifolia, Willd. Sp. Pl. iii. 879 = sibirica.
tenuis, *DC. Prod.* i. 329.—Bras.
tenuis, A. Dietr. in Otto & Dietr. Allg. Gartenz. ii. (1834) 119 = rarifolia.
tenuis, Torr. ex A. Gray, Pl. Wright. i. 41 = leptostachys.
teretifolia, *Linn. f. Suppl.* 316.—Afr. austr.
tetragona, Burch. ex DC. Prod. i. 322 = oppositifolia.
tetrasepala, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 37 = Petitiana.
Thea, Burm. f. Fl. Ind. 154 = seq.?
theezans, *Linn. Mant.* ii. 260.—Java; Japon.
thesioides, *Willd. Sp. Pl.* iii. 877.—Chili.; Peruv.
Thunbergii, *DC. Prod.* i. 333.—Afr. austr.
thuringiaca, Spreng. Syst. iii. 166 = nicaeënsis.
Thwaitesii, *Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 169.—Zeylan.
thymifolia, Thunb. Prod. Pl. Cap. 121 = Muralтia thymifolia.
Timbali, Le Grand, in Bull. Soc. Bot. Fr. xxviii. (1881) 54 = calcarea.
Timoutou, *Aubl. Pl. Gui.* ii. 737. t. 295.—Am. austr.
Timutu, Poir. Encyc. v. 491 = praec.
tinctoria, Vahl, Symb. Bot. i. 50 = javana.
tomentosa, Rottl. ex Mart. in Denkschr. Bot. Ges. Regensb. i. (1815) 185 = violacea.
tomentosa, *Thunb. Prod. Pl. Cap.* 120.—Afr. austr.
tomentosa, Vahl, ex DC. Prod. i. 326 = erioptera.
Torreyi, G. Don, Gen. Syst. i. 360 = alba.
toxoptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 348.—Malaya.
tranquebarica, Mart. in Denkschr. Bot. Ges. Regensb. i. (1815) 186 = chinensis.
tranquebarica, Wight, in Wall. Cat. n. 4169 = rosmarinifolia.
transsilvanica, Schur, Enum. Pl. Trans. 90 = vulgaris.
trichisperma, Jacq. Obs. Bot. iii. 16, sphalm. = seq.
trichosperma, *Linn. Syst.* ed. XII. iii. 232.—Bras.
triflora, Hassk. in Miq. Ann. Mus. Lugd. Bat. i. 175 = rosmarinifolia.

POLYGALA:—
triflora, Linn. Sp. Pl. 705 = chinensis.
triflora, Oliver, Fl. Trop. Afr. i. 128 = erioptera.
trinervata, Buch.-Ham. ex Wall. Cat. n. 4192 B = Salomonia cantoniensis.
trinervia, Thunb. Prod. Pl. Cap. 121 = Muralтia trinervia.
triphylla, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 200.—Ind. or.; Burma.
triphylla, Burm. f. Fl. Cap. Prod. 19 (Legumin. sp.).—Afr. austr.
triquetra, *Presl, Bot. Bemerk.* 15.—Afr. austr.
tutea, Crantz, Inst. ii. 71, err. typ. = lutea.
uliginosa, Pers. Syn. ii. 272 = P. Timoutou.
uliginosa, Weinm. Fl. Petrop. 70 = amara.
umbellata, Thunb. Prod. Pl. Cap. 120 = bracteolata, *Linn.*
umbrosa, *Mart. in Denkschr. Bot. Ges. Regensb.* i. (1815) 183.—Ind. or.
uncinata, Hochst. ex Harv. & Sond. Fl. Cap. i. 86 = abyssinica.
uncinata, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. i. (1842) 468 = rarifolia.
undulata, *Roxb. Hort. Beng.* [98]; *Fl. Ind.* iii. 219.—Ins. Penang.
unguiculata, Poir. Encyc. v. 494 = angustifolia.
uniflora, Michx. Fl. Bor. Am. ii. 53 = paucifolia.
Vahliana, DC. Prod. i. 326 = erioptera.
Vailliantii, Bess. Enum. Pl. Volh. 73 = vulgaris.
variabilis, *H. B. & K. Nov. Gen. et Sp.* v. 397.—Am. austr.
varians, Mart. in Denkschr. Bot. Ges. Regensb. i. (1815) 184 = chinensis.
Vayredae, *Costa, Suplem. Fl. Catal.* 10, 91.—Hispan.
velutina, *Presl, Rel. Haenk.* ii. 100.—Mexic.
venenosa, *Juss. ex Poir. Encyc.* v. 493.—Java.
venosa, Heyne, ex Wall. Cat. n. 4184 = sibirica.
venulosa, A. Dietr. in Otto. & Dietr. Allg. Gartenz. ii. (1834) 116 = Dietrichiana.
venulosa, *Sibth. & Sm. Fl. Graec. Prod.* ii. 52.—Graecia.
veronicea, F. Muell. in Trans. Vict. Inst. i. (1855) 117 = japonica.
versicolor, *Pomel, Nouv. Mat. Fl. Atl.* 336.—Alger.
versicolor, Willd. ex Steud. Nom. ed. II. ii. 373 = grandiflora.
verticillata, *Linn. Sp. Pl.* 706.—Am. bor.
verviana, Lej. Fl. Spa, ii. 92 = vulgaris.
viminea, Houtt. ex DC. Prod. i. 338 = Mundtia spinosa.
violacea, *Aubl. Pl. Gui.* ii. 735.—Guian.
violacea, Vahl, Symb. Bot. ii. 79 (= praec?).—Guian.
violoides, *A. St. Hil. Fl. Bras. Mer.* ii. 48.—Bras.
virgata, *Thunb. Prod. Pl. Cap.* 120.—Afr. austr.
virgata, Vell. Fl. Flum. 203; vii. t. 69 = lancifolia.
viridescens, Linn. Sp. Pl. 705 = sanguinea.
viridescens, Walt. Fl. Carol. 178 = nana.
viridis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 325.—Am. bor. occ.
volubilis, Boj. in Bouton, 12me Rapp. Ann. Maur. (1842) 14 = macroptera.
vulgaris, Asso, Syn. Stirp. Arag. 92 = calcarea.
vulgaris, Desf. Fl. Atlant. ii. 127 = rosea.
vulgaris, *Linn. Sp. Pl.* 702.—Europ.; As. Min.
vulgaris, Thunb. Fl. Jap. 277 = sibirica.
Wallichiana, Wight, Illustr. i. 49. t. 22 = persicariaefolia.
Warmingiana, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 6.—Bras.
Wattersii, *Hance, in Journ. Bot.* xix. (1881) 209.—China.
Webbiana, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 240.—Marocc.
Wightiana, Wall. Cat. n. 4190 = elongata.
Wolfgangiana, Bess. ex Ledeb. Fl. Ross. i. 271 = vulgaris.
Wrightii, Hemsl. Biol. Centr. Am. Bot. i. 62 = Monnina Wrightii.
Xanti, *A. Gray, in Proc. Am. Acad.* v. (1861) 153.—Calif.
Zablotskiana, *Fisch. & Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 347.—Persia.
Zeyheri, Spreng. ex Eckl. & Zeyh. Enum. 18 = oppositifolia.

POLYGALOIDES, Tourn. ex Hall. Enum. Stirp. Helv. ii. 607 (1742) = praec.

POLYGLOCHIN, Ehrh. Beitr. iv. 146 (1789) = **Carex**, Dill. (Cyperac.).

POLYGONARIA, Heist. Syst. 6 (1748) = **Corrigiola**, Linn. (Illecebr.).

POLYGONASTRUM, Moench, Meth. 637 (1794) = **Smilacina**, Desf. (Liliac.).
compressum, Moench, Meth. Suppl. 268 = Ophiopogon japonicus.
racemosum, Moench, Meth. 637 = S. racemosa.

POLYGONATUM, [Tourn.] Adans. Fam. ii. 54 (1763). *LILIACEAE*, Benth. & Hook. f. iii. 768.
AXILLARIA, Rafin. in Journ. Phys. lxxxix. (1819) 261.
CAMPYDORUM, Salisb. Gen. Pl. Fragm. 64 (1866).
EVALLARIA, Neck. Elem. iii. 189 (1790).
PERIBALLANTHUS, Franch. & Sav. Enum. Pl. Jap. ii. 524 (1879).
SIPHYALIS, Rafin. Fl. Tellur. iv. 17 (1836).
ambiguum, Link, ex Schult. f. Syst. vii. 299 = officinale.
anceps, Moench, Meth. 637 = officinale.
angustifolium, Pursh, Fl. Am. Sept. i. 234 = biflorum.
biflorum, *Ell. Sketch*, i. 393.—Am. bor.
bracteatum, G. Don, in Loud. Hort. Brit. 131; A. Dietr. in Otto & Dietr. Allg. Gartenz. iii. (1835) 223 = multiflorum.
brevistylum, *Baker, in Journ. Linn. Soc.* xiv. (1875) 556.—Reg. Himal.
canaliculatum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 148 = latifolium.
canaliculatum, Pursh, Fl. Am. Sept. i. 234 = giganteum.
Cathcartii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 559.—Reg. Himal.
chinense, Kunth, Enum. Pl. v. 146 = sibiricum.
ciliatum, Royle, Illustr. Bot. Himal. 380 = geminiflorum?
cilicicum, Schott, ex Kotschy, in Oestr. Bot. Wochenbl. vii. (1857) 205 = polyanthemum.
cirrhifolium, Royle, Illustr. Bot. Himal. 380 = sibiricum.
commutatum, A. Dietr. in Otto & Dietr. Allg. Gartenz. iii. (1835) 223 = giganteum.
falcatum, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858-59) 414.—Japon.
geminiflorum, *Decne. in Jacquem. Voy. Bot.* 170. t. 170.—Reg. Himal.
giganteum, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 222.—Am. bor.
glaberrimum, C. Koch, in Linnaea, xxii. (1849) 267 = officinale.
Govanianum, Royle, Illustr. Bot. Himal. 380 (sphalm. Govianum) = multiflorum.
graminifolium, *Hook. Ic. Pl.* t. 833.—Reg. Himal.
Griffithii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 558.—Reg. Himal.
Gussonii, Parl. Fl. Ital. iii. 44 = multiflorum.
hirtum, Pursh, Fl. Am. Sept. i. 234 = latifolium.
Hookeri, *Baker, in Journ. Linn. Soc.* xiv. (1875) 558.—Reg. Himal.
humile, Fisch. ex Maxim. Prim. Fl. Amur. 275 = officinale.
Inglesii, *Royle, Illustr. Himal. Bot.* 380, nomen.—Reg. Himal.
intermedium, Dum. Fl. Belg. Prod. 138; Bor. Fl. Cent. Fr. ed. III. ii. 615 = multiflorum.
involucratum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 205.—Japon.
Jacquemontianum, Kunth, Enum. Pl. v. 143 = verticillatum.
japonicum, *C. Morr. & Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 311.—Japon.
lasianthum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 209.—Japon.
latifolium, *Desf. in Ann. Mus. Par.* ix. (1807) 50.—Europ. austr.; As. et Am. bor.
latifolium, Pursh, Fl. Am. Sept. i. 235 = giganteum.
leptophyllum, Royle, Illustr. Bot. Himal. 380 = verticillatum.
macrophyllum, Sweet, Hort. Brit. ed. I. 415 = verticillatum.
macropodum, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 205; (1837) n. VII. 158.—China.
majale, All. Fl. Pedem. i. 130 = Convallaria majalis.

POLYGONATUM:—
Maximowiczii, *F. Schmidt, Reisen Amurl.* 185.—Ins. Sachalin.
multiflorum, *All. Fl. Pedem.* i. 131.—Europ.; As. bor.; Afghan.
multiflorum, Kunth, Enum. Pl. v. 138 = biflorum.
nervulosum, *Baker, in Journ. Linn. Soc.* xiv. (1875) 557.—Reg. Himal.
obtusifolium, Weinm. Elench. Hort. Pawlowsk. (1824) 175 = officinale.
officinale, *All. Fl. Pedem.* i. 131.—Europ.; Sibir.
oppositifolium, *Royle, Illustr. Bot. Himal.* 380.—Reg. Himal.
orientale, Desf. in Ann. Mus. Par. ix. (1807) 50. t. 7 = polyanthemum.
parviflorum, A. Dietr. in Otto & Dietr. Allg. Gartenz. (1835) 222 = biflorum.
polyanthemum, *Link, Handb.* i. 185; *A. Dietr. l. c.* 223.—Europ.; Reg. Cauc.; Syria; Persia.
Polygonatum, Jirasek, ex Schult. f. Syst. vii. 299 = officinale.
pruinosum, *Boiss. Fl. Orient.* v. 332.—Persia.
pubescens, Pursh, Fl. Am. Sept. i. 234 = biflorum.
punctatum, *Royle, Illustr. Bot. Himal.* 380.—Reg. Himal.
roseum, Hook. Bot. Mag. t. 5049 = verticillatum.
roseum, *Kunth, Enum. Pl.* v. 144.—Sibir.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) I. 436.—Turkestan.
sibiricum, [*Delar. in*] *Red. Lil.* vi. t. 315.—Reg. Himal.; As. bor.
stenophyllum, Maxim. Prim. Fl. Amur. 274 = verticillatum.
Thunbergii, C. Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 312 = latifolium.
umbellatum, *Baker, in Journ. Linn. Soc.* xiv. (1875) 553.—China.
uniflorum, Gilib. Exercit. ii. 463; Jaume St. Hil. Pl. Fr. t. 346 = officinale.
verticillatum, *All. Fl. Pedem.* i. 131.—Europ.; As. bor.
vulgare, Desf. in Ann. Mus. Par. ix. (1807) 49 = officinale.

POLYGONELLA, Michx. Fl. Bor. Am. ii. 240 (1803). *POLYGONACEAE*, Benth. & Hook. f. iii. 97.
GONOPYRUM, Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. vi. (1840) 144.
LYONIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
STOPINACA, Rafin. Fl. Tellur. iii. 11 (1836).
articulata, Meissn. Gen. Comm. 228 = Polygonum articulatum.
brevifolia, Meissn. in DC. Prod. xiv. 78 = Atraphaxis buxifolia.
brachystachya, *Meissn. l. c.* 80.—Florida.
buxifolia, Meissn. Gen. Comm. 228 = Atraphaxis buxifolia.
ciliata, *Meissn. in DC. Prod.* xiv. 81.—Florida.
Croomii, *Chapm. Fl. S. U. St.* 387.—Am. bor.
ericoides, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 231.—Texas.
gracilis, *Meissn. in DC. Prod.* xiv. 80.—Am. bor.
laetevirens, Meissn. Gen. Comm. 228 = Atraphaxis laetevirens.
lanceolata, Meissn. l. c. = Atraphaxis Laxmanni.
Meissneriana, *Shuttl. ex Meissn. in DC. Prod.* xiv. 81.—Am. bor.
parviflora, Steud. Nom. ed. I. 643, sphalm. = seq.
parvifolia, *Michx. Fl. Bor. Am.* ii. 241.—Am. bor.
polygama, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 231 = parvifolia.
pungens, Meissn. Gen. Comm. 228 = Atraphaxis pungens.

POLYGONIFOLIA, Adans. Fam. ii. 272 (1763) = **Corrigiola**, Linn. (Illecebr.).

POLYGONON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 132 = **Polygonum**, Tourn.
bistortum, St. Lag. l. c. = Polygonum Bistorta.
persicarium, St. Lag. l. c. 132 = Polygonum Persicaria.

POLYGONUM, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 116 (1737). *POLYGONACEAE*, Benth. & Hook. f. iii. 97.
AMPELYGONUM, Lindl. Bot. Reg. (1838) App. 63.
ANTENORON, Rafin. Fl. Ludov. 28 (1817).

POLYGONUM:—
ASICARIA, Neck. Elem. ii. 211 (1790).
BERGERIA, Koen. ex Steud. Nom. ed. II. i. 849, in syn. (1840).
BILDERDYKIA, Dum. Fl. Belg. 18 (1827).
BISTORTA, Tourn. ex Linn. Gen. ed. I. 115 (1737).
CHULUSIUM, Rafin. Ann. Nat. 12 (1820).
CHYLOCALYX, Hassk. in Flora, xxv. (1842) II. Beibl. 20.
CNOPOS, DIOCTIS, DISCOLENTA, GONONCUS, HEPTARINA, MITESIA, PLEUROSTENA, POGALIS, SPERMAULAXEN, SUNANIA, TASOBA, TRACAULON, Rafin. Fl. Tellur. iii. 12–16, 94, 95 (1836).
ECHINOCAULON, Spach, Hist. Vég. Phan. x. 521 (1841).
GONIATICUM, Stokes, Bot. Mat. Med. ii. 338 (1812).
HELXINE, Linn. Syst. ed. I. (1735).
HEPTARINIA, Rafin. New Fl. Am. iv. 48 (1836).
KOENIGIA, Linn. Mant. i. 3 (1767).
LAGUNEA, Lour. Fl. Cochinch. 220 (1790).
LYONELLA, Rafin. in Am. Monthly Mag. (1818) 266.
PERSICARIA, [Tourn.] Linn. Syst. ed. I. (1735).
PEUTALIS, Rafin. New Fl. Am. iv. 47 (1836); Fl. Tellur. iii. 14 (1836).
PLEUROPTERUS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 587.
TEPHIS, Adans. Fam. ii. 276 (1763).
THYSANELLA, A. Gray, in Bost. Journ. Nat. Hist. v. (1847) 232.
TINIARIA, Reichb. Handb. 236 (1837).
TOVARA, Adans. Fam. ii. 276 (1763).
TRACHOPYRON, Ger. ex Rafin. Fl. Tellur. iii. 10 (1836).
abyssinicum, *A. Rich. Tent. Fl. Abyss.* ii. 225.—Abyss.
acaule, *Boiss. Fl. Orient.* iv. 1043.—Persia.
acerosum, *Ledeb. ex Meissn. in DC. Prod.* xiv. 92.—Soongaria.
acetosaefolium, Vent. Jard. Cels, 88. t. 88 = Muehlenbeckia sagittifolia.
acetosum, *Bieb. Fl. Taur. Cauc.* i. 304.—Reg. Casp.
acetosum, Eichw. Pl. Casp. Cauc. 23 = cognatum.
acidulum, Willd. Enum. Hort. Berol. 429 = divaricatum.
acidum, Pall. Reise, ii. 25 = alpinum.
acidum, Willd. ex Meissn. in DC. Prod. xiv. 96 = acetosum.
acre, *H. B. & K. Nov. Gen. et Sp.* ii. 179.—Am. calid.
acre, Lam. Fl. Fr. iii. 234 = P. Hydropiper.
acre, Meissn. in Wall. Pl. As. Rar. iii. 57 = serrulatum.
acre, Sieber, ex Meissn. Monog. 79 = glabrum.
acuminatum, *H. B. & K. Nov. Gen. et Sp.* ii. 178.—Am. trop.
acuminatum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) n. IV. 12 = segetum.
acuminatum, Meissn. Monog. 78, partim. = glabrum.
acutatum, Lehm. Ind. Sem. Hort. Hamb. (1821) 6; Pugill. i. 28 = Fagopyrum cymosum.
acutifolium, Schur, Enum. Pl. Trans. 587 = aviculare.
adenophorum, Spreng. Syst. ii. 258 = lapathifolium.
adenophyllum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 48.—Afr. austr.
adenotrichum, Griff. Rep. Bot. Gard. Calc. (1843) 39 = chinense.
adpressum, R. Br. Prod. 420, partim = Muehlenbeckia gracillima.
adpressum, Labill. Nov. Holl. Pl. i. 99. t. 127 = Muehlenbeckia adpressa.
affine, *D. Don, Prod. Fl. Nep.* 70.—Reg. Himal.
affine, Steph. ex Spreng. Syst. ii. 256 = cognatum.
affine, Wall. Cat. n. 1683/3 = viviparum.
afghanicum, *Meissn. in DC. Prod.* xiv. 90.—Afghan.
agrestinum, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 559 = aviculare.
alatum, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 72.—As. et Afr. trop. et sub-trop.
alatum, Dulac, Fl. Hautes-Pyr. 169 = dumetorum.
aleppicum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1037.—Syria.
alopecuroides, *Turcz. ex Steud. Nom.* ed. II. ii. 374.—Sibir. baical.
alpestre, Heldr. ex Meissn. in DC. Prod. xiv. 694 = plebejum.
alpestre, C. A. Mey. Verz. Pfl. Cauc. 157 = cognatum.
alpestre, Schur, in Oestr. Bot. Zeitschr. i. (1861) 290 = P. Bistorta.

POLYGONUM:—
alpestre, Wall. Cat. n. 1725 = filicaule.
alpinum, *All. Fl. Pedem.* ii. 206. t. 68. f. 1.—Europ. austr.; As. bor.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 170 = mite.
altissimum, Moench, Meth. 630 = orientale.
ambiguum, Meissn. in Wall. Pl. As. Rar. iii. 54 = amplexicaule.
ammannoides, Jaub. & Spach, Illustr. ii. 28. t. 119 = cognatum.
amoenum, Blume, Bijdr. 532 = orientale.
amoenum, Salisb. Prod. 258 = P. Bistorta.
amphibium, *Linn. Sp. Pl.* 361.—Reg. bor. temp.
amplexicaule, *D. Don, Prod. Fl. Nep.* 70.—Reg. Himal.
angustifolium, D. Don, l. c. = viviparum.
angustifolium, Hort. ex Poir. Encyc. vi. 139 = Laxmanni.
angustifolium, Pall. Reise, iii. 230, 320 = alpinum.
angustifolium, Rafin. Fl. Tellur. iii. 16 (Quid?).—Hab.?
angustifolium, Roth, Tent. Fl. Germ. ii. 453 = minus.
angustissimum, F. Muell. Fragm. v. 92 = Muehlenbeckia polygonoides.
anomalum, *De Bruyn, in Miq. Pl. Jungh.* 311.—Java.
antihaemorrhoidale, Mart. Reise Bras. i. 550 = acre.
aphyllum, Krock. Fl. Siles. Suppl. ii. 75 = aviculare?
appressum, Steud. Nom. ed. II. ii. 374 = Muehlenbeckia adpressa.
aquaticum, A. Gray, ex Meissn. in DC. Prod. xiv. 116 = amphibium.
arachnoideum, Klotzsch, ex Meissn. l. c. 117 = lanigerum.
arborescens, Vell. Fl. Flum. 162; iv. t. 43 = Coccoloba nitida.
arcticum, Pall. ex Spreng. Syst. ii. 258 = sibiricum.
arenarium, Loisel. Nouv. Not. 18; Fl. Gall. ed. II. i. 284 = Bellardi.
arenarium, Rafin. ex Meissn. in DC. Prod. xiv. 98 = aviculare.
arenarium, *Waldst. & Kit. Pl. Rar. Hung.* i. 69. t. 67.—Europ. austr.
arenastrum, Bor. Fl. Cent. Fr. ed. III. ii. 559 = aviculare.
argenteum, Ehrenb. ex Meissn. in DC. Prod. xiv. 88 = maritimum.
argyrocoleon, *Steud. ex Kunze, in Linnaea*, xx. (1847) 17.—Persia.
aridum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1042.—Persia.
arifolium, *Linn. Sp. Pl.* 364.—Reg. Himal.; Am. bor.
arifolium, Spreng. Syst. ii. 258 = strigosum.
arifolium, Thunb. Fl. Jap. 168 = Thunbergii.
articulatum, R. Br. Prod. 420 = australe.
articulatum, *Linn. Sp. Pl.* 363.—Am. bor.
articulatum, Perr. ex Meissn. Monog. 54 = senegalense.
ascendens, Pritz. Ic. Ind. ii. 222, sphalm. = Polypogon littoralis.
asperulum, Wall. Cat. n. 1724 = muricatum.
assamicum, *Meissn. in DC. Prod.* xiv. 111.—Ind. or.; Burma.
Atraphaxis, Thunb. Fl. Cap. ed. Schult. 385 = seq.
atraphaxoides, *Thunb. Prod. Pl. Cap.* 77.—Afr. austr.
atriplicifolium, Wall. Cat. n. 1719 (= *Ceratogonum atriplicifolium*).—Ind. or.
attenuatum, *R. Br. Prod.* 420.—Austral.
auriculatum, Meissn. Monog. 59. t. 6 = chinense.
Austinae, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 212.—Calif.
australe, A. Rich. Ess. Fl. N. Zél. 178 = Muehlenbeckia australis.
australe, *Spreng. Syst.* ii. 258.—Austral.
australe, Thieb. ex Pers. Syn. i. 440 = hydropiperoides.
aviculare, Bert. ex Meissn. in DC. Prod. xiv. 88 = chilense.
aviculare, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1843) 407 = acetosum.
aviculare, *Linn. Sp. Pl.* 362.—Reg. bor. temp.
aviculare, Lour. Fl. Cochinch. 241 = plebejum.
aviculare, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 407 = Meyenii.
axillare, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 278 = Muehlenbeckia axillaris.

POLYGONUM:—
azoricum, H. C. Wats. ex Meissn. in DC. Prod. xiv. 111 = serrulatum.
Babingtonii, Endl. Gen. Suppl. iv. II. 47 = flaccidum.
Babingtonii, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 239 = senticosum.
baccatum, Soland. ex Meissn. in DC. Prod. xiv. 146 = Muehlenbeckia australis.
bacciferum, Boj. Hort. Maurit. 271 = praec.?
Balansae, *Boiss. Diagn.* Ser. II. iv. 78.—Algeria.
baldschuanicum, *Regel, in Act. Hort. Petrop.* viii. (1884) 684. t. 10.—Bokhara.
Barka, Buch.-Ham. ex Wall. Cat. n. 1722 = minus.
barbatum, Comm. ex Meissn. in DC. Prod. xiv. 115 = imberbe.
barbatum, *Linn. Sp. Pl.* 362.—Geront. trop.
barbatum, G. F. W. Mey. Prim. Fl. Esseq. 158 = acuminatum.
barbatum, Roxb. Hort. Beng. 29; Fl. Ind. ii. 289 = flaccidum.
barbatum, Thunb. Prod. Pl. Cap. 77 = lapathifolium.
barbatum, Wall. Cat. n. 1708 F = stagninum.
Baumgartenianum, Schur, Enum. Pl. Trans. 584 = lapathifolium.
Bellardi, *All. Fl. Pedem.* ii. 205. t. 90. f. 2.—Reg. Mediterr. et Cauc.
Bellardi, Blanco, Fl. Filip. ed. I. 314 = P. Persicaria.
Berteroi, *Phil. in Linnaea*, xxix. (1857–58) 38.—Chili.
Bertolonii, *Goiran & Lonivi, ex Goiran & Del Fabbro, Spec. Morph. Veg.—Cf. Nuov. Giorn. Bot. Ital.* ix. (1877) 170.—Hab.?
Beyrichianum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 42.—Bras.
biaristatum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 90.—Afghan.
bicolor ×, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 392.—Europ.
bicorne, Rafin. Fl. Ludov. 29 = pensylvanicum.
Bidwelliae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 294.—Calif.
biforme, Wahlenb. Fl. Suec. i. 242 = P. Persicaria.
biscaticlium, Buch.-Ham. ex Meissn. in DC. Prod. xiv. 106 = flaccidum.
biscatilium, Wall. Cat. p. 215. sub n. 1723 = praec.
Bishirae, Buch.-Ham. ex Meissn. l. c. 102 = stagninum.
Bistorta, Garcke, in Bot. Waldem. Reis. 135 = viviparum.
Bistorta, *Linn. Sp. Pl.* 360.—Reg. bor.
Bistorta, Walt. Fl. Carol. 131 = amphibium.
bistortoides, Pursh, Fl. Am. Sept. i. 271; Boiss. Diagn. Ser. I. v. 46 = P. Bistorta.
bistortum, Dulac, Fl. Hautes-Pyr. 168 = P. Bistorta.
Blumei, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 57.—Java.
Bocconi, DC. ex Meissn. in DC. Prod. xiv. 86 = scoparium.
Bolanderi, *Brew. ex A. Gray, in Proc. Am. Acad.* viii. (1873) 400.—Calif.
bonaërense, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 104.—Reg. Argent.
borneense, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 58.—Borneo.
Bowenkampi, *Phil. in Linnaea*, xxxiii. (1864–65) 225.—Chili.
brachiatum, Poir. Encyc. iv. 15 = chinense.
brachypodum, Baker, in Journ. Linn. Soc. xx. (1883) 239 = Meissnerianum.
bracteatum, Spreng. Syst. iv. Cur. Post. 154 = viviparum.
bracteosum, *Phil. Sert. Mend. Alt.* 42.—Chili.
brasiliense, *Koch, in Linnaea*, xxii. (1849) 206.—Bras.
Braunii, Bluff & Fingerh. Comp. Fl. Germ. i. 509 = mite.
brevisetum, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 61.—Guian.
Brittingeri, Opiz, Natural. viii. (1824) 74 = lapathifolium.
Brunonis, Wall. Cat. n. 1692 = affine.
bulbiferum, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1841) 94 = viviparum.
Bungeanum, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 77.—China.
buxifolium, Bieb. Fl. Taur. Cauc. i. 300 = Atraphaxis buxifolia.
buxifolium, Nutt. ex Bong. Vég. Isle Sitcha, 43 = aviculare.

POLYGONUM:—
caespitosum, Blume, Bijdr. 532 = P. Posumbu.
californicum, *Meissn. in DC. Prod.* xiv. 100.—Am. bor. occ.
çamporum, *Meissn. in Mart. Fl. Bras.* v. I. 21.—Bras.
Camptianum, Hort. ex Meissn. in DC. Prod. xiv. 147 = Muehlenbeckia complexa.
canescens, *Link, ex Dethard. Consp. Pl. Megalop.* 33 (Quid?).—Europ.
capitatum, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 73.—Reg. Himal.
capitatum, Korth. ex Meissn. in DC. Prod. xiv. 128 = alatum.
cappadocicum, *Boiss. & Bal. Diagn.* Ser. II. iv. 78.—Cappadoc.
Careyi, *Olney, Rhode Isl. Pl.* 29; *et ex A. Gray, Man. Bot. N. U. St.* ed. I. 386.—Am. bor.
carneum, C. Koch, in Linnaea, xxii. (1849) 197 = paleacum.
caucasicum, Hoffm. in Comm. Soc. Phys. Med. Mosq. i. I. (1808) 10 = Atraphaxis buxifolia.
cedrorum, Boiss. & Kotschy, Diagn. Ser. II. iv. 77 = Libani.
centinodium, Lam. Fl. Fr. iii. 237 = aviculare, Bellardi.
cereale, Salisb. Prod. 259 = Fagopyrum esculentum.
Chamissoanum, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 254 = seq.
Chamissonianum, Steud. Nom. ed. II. ii. 375 = Meissnerianum.
chilense, *C. Koch, in Linnaea*, xxii. (1849) 206.—Chili.
chinense, Houtt. Handleid. viii. 479 = multiflorum.
chinense, *Linn. Sp. Pl.* 363.—As. or.; Malaya.
chlorocoleum, Steud. ex Boiss. Fl. Orient. iv. 1034 = Bellardi.
ciliatum, Buch.-Ham. in D. Don, Prod. Fl. Nep. 73 = microcephalum.
ciliatum, Ehrenb. ex Meissn. in DC. Prod. xiv. 111 = serrulatum.
ciliatum, Kunze, ex Meissn. l. c. 142 = Paronychia chilensis.
ciliatum, *Lour. Fl. Cochinch.* 243.—China.
ciliatum, Ruiz, ex Meissn. in DC. Prod. xiv. 114 = acuminatum.
ciliatum, Willd. ex Spreng. Syst. ii. 255 = alpinum.
cilinode, *Michx. Fl. Bor. Am.* i. 241.—Am. bor.
ciliosum, Meissn. in DC. Prod. xiv. 95 = plebejum.
cirrhosum, Moç. & Sesse, ex Meissn. l. c. 184 = Antigonon leptopus.
claviculatum, Hort. Par. ex Meissn. l. c. 185 = Brunnichia cirrhosa.
cliffortioides, Meissn. in Wall. Pl. As. Rar. iii. 62 = plebejum.
coarctatum, *Dougl. ex Meissn. in DC. Prod.* xiv. 101.—Am. bor. occ.
coarctatum, Willd. ex Spreng. Syst. ii. 255 = sibiricum.
coccineum, Muhl. ex Willd. Enum. Hort. Berol. 428 = amphibium.
Coccoloba, Steud. Nom. ed. II. ii. 375 = Muehlenbeckia australis.
cochinchinense, Meissn. Monog. 55 = orientale.
cognatum, *Meissn. l. c.* 91.—Reg. Himal.
collare, Mart. ex Meissn. in DC. Prod. xiv. 119 = attenuatum.
compactum, *Hook. f. Bot. Mag.* t. 6476.—Japon.
complexum, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 455 = Muehlenbeckia complexa.
Comptonianum, Hort. ex Meissn. in DC. Prod. xiv. 147 = Muehlenbeckia complexa.
condensatum, Schultz, ex Nym. Consp. 628 = mite.
confertiflorum, Dougl. ex Meissn. in DC. Prod. xiv. 89 = P. Paronychia.
confertum, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1841) 116 = cognatum.
confusum, Meissn. in Wall. Pl. As. Rar. iii. 53 = paleaceum.
conspersum, Meissn. l. c. 56 = stagninum.
controversum, Guss. Fl. Sic. Prod. i. 472 = equisetiforme.
convolvulaceum, Lam. Fl. Fr. iii. 239 = P. Convolvulus.
convolvuliforme, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = seq.
Convolvulus. *Linn. Sp. Pl.* 364.—Reg. bor. temp.
cordatum, Hort. Berol. ex Meissn. in DC. Prod. xiv. 146 = Muehlenbeckia varians.

POLYGONUM:—
coronulatum, Steud. Nom. ed. II. ii. 375 = plebejum.
corrigioloides, *Jaub. & Spach, Illustr.* ii. 33. t. 124.—Persia.
corsicanum, *Link, Handb.* i. 300.—Ins. Corsic.
corymbosum, Willd. Sp. Pl. ii. 452 = chinense.
crassicaule, Rafin. Fl. Tellur. iii. 15 (Quid?).—Hab.?
crassifolium, Murr. Syst. Veg. ed. XIV. 378 = sibiricum.
crassinerve, Cesati, Stirp. Ital. fasc. 2, cum ic. = Roberti.
crenatum, Ehrenb. ex Meissn. in DC. Prod. xiv. 86 = equisetiforme.
criopolitanum, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 238.—China.
crispatum, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 138.—Ind. or.
crispulum, Sims, Bot. Mag. t. 1065 = Atraphaxis buxifolia.
crispum, Hornem. Hort. Hafn. Add. 962 = divaricatum.
cristatum, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 259.—Texas.
Cunninghamii, Meissn. in DC. Prod. xiv. 85 = Muehlenbeckia Cunninghami.
cuspidatum, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 84.—Japon.
cuspidatum, Willd. ex Spreng. Syst. ii. 256 = acuminatum.
cymosum, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 289 = chinense.
cymosum, Trevir. Del. Sem. Hort. Vrat. (1824); et in Nov. Act. Nat. Cur. xiii. (1826) I. 177 = Fagopyrum cymosum.
danubiale, *Kern. in Oestr. Bot. Zeitschr.* xxv. (1875) 255.—Europ.
daphnophyllum, Andrz. ex Meissn. in DC. Prod. xiv. 111 = minus.
Davisiae, *Brewer, ex A. Gray, in Proc. Am. Acad.* viii. (1873) 399.—Calif.
davuricum, Willd. ex Spreng. Syst. ii. 256 = Laxmanni.
Debeauxii, Legrand, in Bull. Soc. Bot. Fr. xxx. (1883) 71 = aviculare.
debile, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 63.—Ind. or.
debile, Steud. ex Meissn. in DC. Prod. xiv. 129 = alatum.
deciduum, Boiss. & Noé, in Bot. Zeit. xi. (1853) 734 = argyrocoleon.
decipiens, R. Br. Prod. 420 = minus.
declinatum, Vell. Fl. Flum. 162; iv. t. 41 = Coccoloba declinata.
decumbens, De Bruyn, in Miq. Pl. Jungh. 307 = P. Posumbu.
decumbens, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = lapathifolium.
delicatulum, *Meissn. in DC. Prod.* xiv. 127.—Reg. Himal.
densiflorum, Blume, Bijdr. 533 = chinense.
densiflorum, Meissn. in Mart. Fl. Bras. v. II. 13 = glabrum.
dentato-alatum, F. Schmidt, in Maxim. Prim. Fl. Amur. 232 = dumetorum, Anredera spicata.
denudatum, Bor. Fl. Centr. Fr. ed. III. ii. 559 = aviculare.
depressum, A. Cunn. ex Meissn. in DC. Prod. xiv. 147 = Muehlenbeckia axillaris.
dibotrys, D. Don, Prod. Fl. Nep. 73 = Fagopyrum cymosum.
dichotomum, Blume, Bijdr. 529 = pedunculare.
dichotomum, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
diclinum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 23 = Muehlenbeckia polygonoides.
diffusum, *Pall. ex Spreng. Syst.* ii. 255.—Sibir.
dioicum, Buch.-Ham. ex Meissn. in DC. Prod. xiv. 143 = Fagopyrum esculentum.
diospyrifolium, *Cham. & Schlecht. in Linnaea*, iii. (1828) 47.—Bras.
dissitiflorum, Guss. ex Nym. Consp. 639 = Bellardi.
distachyum, Mart. ex Meissn. in DC. Prod. xiv. 108 = acre.
divaricatum, Less. in Linnaea, ix. (1834) 204 = junceum.
divaricatum, *Linn. Sp. Pl.* 363.—Sibir.
divaricatum, Poir. Encyc. vi. 148 = scabrum.

POLYGONUM:—
divaricatum, Vill. Hist. Pl. Dauph. iii. 322 = alpinum.
Donianum, Spreng. Syst. iv. Cur. Post. 154 = affine.
Donii, Hort. ex Meissn. in DC. Prod. xiv. 119 = lapathifolium.
Donii, Meissn. Monog. 72 = serrulatum.
Donii, Wall. Cat. n. 1723 B = P. Posumbu.
Donii, Wight, Ic. t. 1801 = flaccidum.
Douglasii, *Greene, in Bull. Calif. Acad.* i. n. III. (1885) 125.—Am. bor. occ.
dracunculifolium, Boiss. Fl. Orient. iv. 1039 = luzuloides.
Dregeanum, Meissn. in Linnaea, xiv. (1840) 487 = aviculare.
Dregei, *Meissn. in DC. Prod.* xiv. 98.—Afr. austr.
Dryandri, Spreng. Syst. ii. 255 = plebejum.
dubium, Stein, ex A. Br. in Flora, vii. (1824) 357 = mite.
dumetorum, *Linn. Sp. Pl.* ed. II. 521.—Reg. bor. temp.
dumosum, *Boiss. Diagn.* Ser. I. vii. 83.—Persia.
effusum, Meissn. in DC. Prod. xiv. 93 = plebejum.
Ehrenbergii, Meissn. l. c. 86 = equisetiforme.
elatius, R. Br. Prod. 419 = lapathifolium.
elegans, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 419; Roxb. Hort. Beng. 29; Fl. Ind. ii. 291 = plebeium.
elegans, Hort. ex Meissn. in DC. Prod. xiv. 143 = Fagopyrum esculentum.
elegans, Tenore, Fl. Nap. Prod. Suppl. ii. p. lxvii; Fl. Nap. i. 207 = arenarium.
ellipticum, Willd. ex Spreng. Syst. ii. 253 = P. Bistorta.
emarginatum, Roth, Catalect. Bot. i. 48 = Fagopyrum esculentum.
emarginatum, Wall. Cat. n. 1688 = Fagopyrum cymosum.
Emodi, *Meissn. in Wall. Pl. As. Rar.* iii. 51, 54. t. 287.—Reg. Himal.
Engelmanni, *Greene, in Bull. Calif. Acad.* i. n. III. (1885) 126.—Am. bor.
ephedroides, *Hook. f. Fl. N. Zel.* i. 211.—N. Zel.
epilobioides, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 254.—Bras.
equisetiforme, Gren. & Godr. Fl. Fr. iii. 52 = scoparium.
equisetiforme, Mayer, ex Tenore, Syll. Fl. Neap. 195 = arenarium.
equisetiforme, *Sibth. & Sm. Fl. Graec. Prod.* i. 266; Fl. Graec. t. 364.—Reg. Mediterr.; Oriens.
erectum, Linn. Sp. Pl. 363 = aviculare.
erectum, *Vell. Fl. Flum.* 162; iv. t. 42.—Bras.
erythrodes, *Miq. Fl. Ind. Bat. Suppl.* 365.—Sumatra.
exiguum, *Meissn. Monog.* 90.—Afr. trop.
Fagopyrum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 73 = Fagopyrum cymosum.
Fagopyrum, Linn. Sp. Pl. 364 = Fagopyrum esculentum.
fastigiatum, Rafin. Fl. Tellur. iii. 15 (Quid?).—Hab.?
ferrugineum, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 252.—Bras.
filicaule, *Wall. Cat.* n. 1694.—Reg. Himal.
filiforme, Barton, Comp. Fl. Philad. i. 186 = tenue.
filiforme, Thunb. Fl. Jap. 163 = virginianum.
fimbriatum, *Ell. Sketch*, i. 588.—Am. bor.
fissum, Blume, Bijdr. 532 = barbatum.
flaccidum, *Meissn. in DC. Prod.* xiv. 107, partim.—Ind. or.; Malaya.
flaccidum, Roxb. Hort. Beng. 29; Fl. Ind. ii. 291 = serrulatum.
flagellare, Bertol. in Sebast. & Mauri, Prod. Fl. Rom. 142; Fl. Ital. iv. 382 = romanum.
flagellare, Spreng. Syst. ii. 255, iv. Cur. Post. 155 = Bellardi.
flagelliforme, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 409 = Bellardi.
flagelliforme, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 80.—Bras.
flavescens, Jord. ex Nym. Consp. 639 = Bellardi.
flexuosum, Benth. Pl. Hartw. 80 = Muehlenbeckia tamnifolia.
flexuosum, Heward, ex Meissn. in DC. Prod. xiv. 148 = Muehlenbeckia adpressa.
flexuosum, Pall. ex Ledeb. Fl. Ross. iii. 526 = divaricatum.
floribundum, *Schlecht. ex Spreng. Syst.* ii. 257.—Sibir.
floribundum, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 252 = acuminatum.

POLYGONUM:—
fluviatile, Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 47 = barbatum.
Forbesii, *Hance, in Journ. Bot.* xxi. (1883) 100.—China.
Forsterii, Hook. f. Handb. N. Zeal. Fl. 739, sphalm. = Muehlenbeckia adpressa?
frondosum, *Meissn. in DC. Prod.* xiv. 137.—Reg. Himal.
frutescens, Gueldenst. Reisen, i. 224 = Atraphaxis buxifolia.
frutescens, Hort. Par. ex Jaub. & Spach, Illustr. ii. 14 = Atraphaxis Laxmanni.
frutescens, Linn. Sp. Pl. 359 = Atraphaxis lanceolata.
frutescens, Vell. Fl. Flum. iv. t. 44 = Coccoloba Gardneri.
fruticosum, S. G. Gmel. ex Meissn. in DC. Prod. xiv. 78 = Atraphaxis lanceolata.
fruticulosum, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 407.—Bolivia.
geniculatum, Poir. Encyc. vi. 147 = aviculare.
geniculatum, Riddell, ex Meissn. in DC. Prod. xiv. 132 = Meissnerianum.
glabratum, Meissn. in Wall. Cat. n. 6285 = pedunculare.
glabrum, Hort. ex Reichb. Pl. Crit. v. 59 = lapathifolium.
glabrum, Roxb. ex D. Don, Prod. Fl. Nep. 71 = hydropiperoides.
glabrum, Roxb. Hort. Beng. 29; Fl. Ind. ii. 287 = serrulatum.
glabrum, *Willd. Sp. Pl.* ii. 447.—Reg. trop.
glandulosum, R. Br. Prod. 419 = lapathifolium.
glandulosum, Poir. Encyc. vi. 149 = P. Hydropiper.
glaucum, Nutt. Gen. Am. i. 254 = maritimum.
glutinosum, Wall. Cat. n. 1717 = lanigerum.
Gmelini, Hort. Par. ex Meissn. in DC. Prod. xiv. 99 = Bellardi.
Gmelini, *Steud. Nom.* ed. II. ii. 375.—Sibir.
Goyalparae, Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 47 = stagninum.
gracile, Guss. Enum. Pl. Inar. 285 = Bellardi.
gracile, Hort. ex Meissn. l. c. 143 = Fagopyrum esculentum.
gracile, Nutt. Gen. Am. i. 255 = Polygonella gracilis.
gracile, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 57 = P. Posumbu.
gracile, Salisb. Prod. 259; R. Br. Prod. 419 = P. Hydropiper.
gracilescens, Mart. ex Meissn. in DC. Prod. xiv. 108 = acre.
gracillimum, Spreng. Syst. iv. Cur. Post. 154 = macrophyllum.
gramineum, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 59.—Japon.
gramineum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = barbatum.
graminifolium, Hoffm. ex Meissn. in DC. Prod. xiv. 114 = acuminatum.
graminifolium, *Wierzb. ex Meissn. l. c.* xiv. 95.—Europ. austr.
grandiflorum, Bertol. Fl. Guatim. 12 = Antigonon guatemalense.
grandiflorum, Willd. Sp. Pl. ii. 440 = Atraphaxis grandiflora.
Greenei, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 295.—Calif.
gummifera, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 252.—Bras.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 278 = Muehlenbeckia appressa.
Gussonii, *Todaro, ex Cesati, Pass. & Gibell. Comp. Fl. Ital.* 262.—Europ. austr.
guttuliferum, Miq. ex Meissn. in DC. Prod. xiv. 129 = alatum.
gymnopus, *Franch. & Sav. Enum. Pl. Jap.* ii. 472.—Japon.
Hagei, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1838) 110 = polystachyum?
Hamiltonii, Meissn. in DC. Prod. xiv. I. 104 = stagninum.
Hamiltonii, Spreng. Syst. iv. Cur. Post. 155 = flaccidum.
Hartwrightii, *A. Gray, in Proc. Am. Acad.* viii. (1870) 234.—Am. bor. occ.
hastato-trilobum, Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 62 = Thunbergii.
hastatum, Boj. & Hilsenb. ex Baker, in Journ. Linn. Soc. xx. (1883) 239 = Meissnerianum.

POLYGONUM:—
hastatum, Murr. in Nov. Comm. Gotting. v. (1774) 37. t. 6 = sibiricum.
hastatum, *Zoll. & Mor. Verz. Ind. Archip.* ii. 110 (sp. dub).—Japon.
Hayii, Strach. & Winterb. ex Meissn. in DC. Prod. xiv. 137 = frondosum.
hecasanthum, Schur, Enum. Pl. Trans. 585 = P. Hydropiper.
Hermonis, Kotschy, ex Boiss. Fl. Orient. iv. 1040 = Libani.
herniarioides, Delile, Fl. Aegypt. Illustr. 61 = plebeium.
heterophyllum, Soland. ex Meissn. in DC. Prod. xiv. 147 = Muehlenbeckia complexa.
Hidropiper, Neck. Delic. Gallo Belg. i. 185 = P. Hydropiper.
hippopotami, *Ehrenb. ex Meissn. in DC. Prod.* xiv. 121.—Afr. trop.
hirsutum, Buch.-Ham. ex Meissn. in DC. Prod. xiv. 103 = viscosum.
hirsutum, *Walt. Fl. Carol.* 132.—Am. bor.
hirtum, Willd. ex Spreng. Syst. ii. 259 = orientale.
hispidulum, Blume, Bijdr. 535 = strigosum.
hispidum, Bab. in Trans. Linn. Soc. xviii. (1841) 100 = stagninum.
hispidum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 71 = flaccidum.
hispidum, H. B. & K. Nov. Gen. et Sp. ii. 178 = orientale.
Hookeri, *Meissn. in Ann. Sc. Nat.* Sér. V. vi. (1866) 352.—Reg. Himal.
Hornemanni, Meissn. in DC. Prod. xiv. 107 = stagninum.
horridum, Buch.-Ham. ex Roxb. Hort. Beng. 29; Fl. Ind. ii. 291 = strigosum.
humifusum, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 560 = aviculare.
humifusum, *Merk. ex Pall. N. Nord. Beitr.* vii. 289.—Sibir. altaic.
humile, *Meissn. in Wall. Pl. As. Rar.* iii. 59.—Reg. Himal.
hybridum, Chaub. ex St. Amans, Fl. Agen. 163 = mite.
Hydropiper, *Linn. Sp. Pl.* 361.—Reg. temp. bor. et austr.
Hydropiper, Lour. Fl. Cochinch. 240 = serrulatum.
Hydropiper, Michx. Fl. Bor. Am. i. 238 = acre.
hydropiperoides, *Michx. l. c.* 239.—Am. bor.
hydropiperoides, Pursh, Fl. Am. Sept. i. 270 = acre.
hypostictum, Miq. ex Meissn. in DC. Prod. xiv. 111 = minus.
hyssopifolium, Boj. Hort. Maurit. 271 = plebeium.
illecebroides, Meissn. in DC. Prod. xiv. 94 = plebeium.
imberbe, *Soland. ex Forst. f. Prod.* 90.—Ins. Tahiti.
imbricatum, *Nutt. ex S. Wats. in Am. Natural.* vii. (1873) 665.—Am. bor. occ.
imbricatum, Rafin. Fl. Tellur. iii. 16 (Quid?).—Hab.?
incanum, F. W. Schmidt, Fl. Boëm. iv. 90 = lapathifolium.
incarnatum, *Ell. Sketch*, i. 456.—Am. bor.
indicum, Heyne, in Roth, Nov. Pl. Sp. 208 = plebejum.
infestum, Salisb. Prod. 259 = P. Convolvulus.
injucundum, Lindl. Bot. Reg. t. 1250 = Muehlenbeckia chilensis.
intermedium, Ehrh. Beitr. v. 178, nomen; et vi. 143 = minus.
intermedium, Kit. ex Rochel, Bot. Reise Banat. 71; in Linnaea, xxxii. (1863) 363 = praec.?
intermedium, *Nutt. ex S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 378.—Am. bor.
intermedium, Rob. ex Loisel. Fl. Gall. ed. II. i. 52 = Roberti.
interruptum, *Bunge, Enum. Pl. Chin. Bor.* 58.—China.
inundatum, Mart. ex Meissn. in DC. Prod. xiv. 123 = orientale.
inundatum, Rafin. in Med. Repos. N. York, v. (1808) 359 (Quid?).—Am. bor.
japonicum, *Meissn. in DC. Prod.* xiv. 112.—Japon.
javanicum, *De Bruyn, in Miq. Pl. Jungh.* 303.—Java.
jucundum, *Meissn. Monog.* 71.—China.
junceum, *Ledeb. Fl. Ross.* iii. 529.—Sibir. ural.
junceum, Mitch. Journ. Trop. Austral. 85 = Muehlenbeckia Cunninghami.
Kitaibelianum, Sadler, Fl. Pesth. i. 1. 287 = Bellardi.

POLYGONUM:—
Kotschyanum, Boiss. Diagn. Ser. I. vii. 84 = thymifolium.
lacerum, *H. B. & K. Nov. Gen. et Sp.* ii. 179.—Am. trop.
lanatum, Roxb. Hort. Beng. 29; Fl. Ind. ii. 285 = lanigerum.
lanigerum, *R. Br. Prod.* 419.—Geront. trop.
lapathifolium, *Linn. Sp. Pl.* 360.—Reg. bor. et austr.
lapathifolium, Willd. Sp. Pl. ii. 446 = P. Persicaria.
latifolium, Gieseck. ex Lange, Consp. Fl. Groenl. i. 105 = aviculare.
latifolium, Kotschy, ex Boiss. Fl. Orient. iv. 1032 = alpinum.
laxiflorum, Pall. ex Meissn. in DC. Prod. xiv. 99 = Bellardi.
laxiflorum, Weihe, in Flora, ix. (1826) 746 = mite.
Laxmanni, *Lepech. in Nov. Act. Acad. Petrop.* x. (1797) 414.—Sibir.
laxum, Reichb. Pl. Crit. v. 56 = lapathifolium.
Lemanianum, *Meissn. in DC. Prod.* xiv. 91 (Lemannianum).—Afghan.
leptostachyum, *De Bruyn, in Miq. Pl. Jungh.* 307.—Java.
Libani, *Boiss. Diagn.* Ser. I. xii. 99.—Syria.
libanoticum, Ehrenb. ex Meissn. in DC. Prod. xiv. 96 = cognatum.
limbatum, *Meissn. l. c.* 123.—As. et Afr. trop.
linifolium, Heyne, in Roth, Nov. Pl. Sp. 207 = plebejum.
linifolium, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 168; et Cat. 40 = tenue.
linifolium, Perr. ex Meissn. Monog. 90 = exiguum.
littorale, Gren. & Godr. Fl. Fr. iii. 51 = Roberti.
littorale, Link, Enum. Hort. Berol. i. 385 = aviculare.
longisetum, De Bruyn, in Miq. Pl. Jungh. 307 = Blumei.
Lorinseri, Opiz, Seznam, 73, in syn. (Quid?).—Europ.
Loureirii, Poir. Encyc. Suppl. iv. 667 = ciliatum.
ludovicianum, *Meissn. in DC. Prod.* xiv. 116.—Am. bor.
luzuloides, *Jaub. & Spach, Illustr.* ii. 37. t. 126.—Persia.
Maackianum, Regel, Tent. Fl. Ussur. 127 = Thunbergii.
macranthum, *Meissn. in DC. Prod.* xiv. 107.—Reg. Himal.
macrochaeton, Fresen. in Flora, xxi. (1838) 601 = senegalense.
macrochaetum, Miq. in Linnaea, xviii. (1844) 242 = serrulatum.
macrohymenium, *Boiss. Fl. Orient.* iv. 1041.—Persia.
macrophyllum, *D. Don, Prod. Fl. Nep.* 70.—Reg. Himal.
macropterum, Hort. ex Meissn. in DC. Prod. xiv. 142 = Fagopyrum esculentum.
maculatum, Krock, Fl. Siles. Suppl. ii. 72 = lapathifolium.
maculatum, Rafin. Fl. Ludov. 29 = P. Persicaria.
madagascariense, *Meissn. in DC. Prod.* xiv. 113.—Madag.
Manoraene, Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 47 = barbatum.
marioticum, Ehrenb. ex Meissn. in DC. Prod. xiv. 86 = equisetiforme.
maritimum, Cham. & Schlecht. in Linnaea, iii. (1828) 51 = chilense.
maritimum, Eichw. Pl. Casp. Cauc. 23 = cognatum.
maritimum, *Linn. Sp. Pl.* 361.—Reg. bor. temp.
maritimum, Remy, in C. Gay, Fl. Chil. 269 = striatum.
maritimum, Vell. Fl. Flum. 161; iv. t. 39 = acre.
Maximowiczii, *Regel, Ind. Sem. Hort. Petrop.* (1864) 14.—Japon.
Meissneri, Wall. Cat. n. 1693 = Emodi.
Meissnerianum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 40.—Bras.
melastomaca, *Delile, Fl. Aegypt. Illustr.* 60 (sp. dub.).—Aegypt.
Metzianum, Miq. in Flora, xxxii. (1849) 557 = alatum.
Meyenii, *C. Koch, in Linnaea*, xxii. (1849) 205.—China.
Meyeri, Steud. Nom. ed. II. ii. 376 = paronychioides.
micranthum, Boiss. ex Meissn. in DC. Prod. xiv. 694 = corrigioloides.
micranthum, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 59.—Sumatra.

POLYGONUM:—
micranthum, *A. Rich. Fl. Cub. Fanerog.* ii. 183.—Cuba.
microcephalum, *D. Don, Prod. Fl. Nep.* 72.—Reg. Himal.
microcephalum, Hassk. in Flora, xiv. (1831) 561 = alatum.
microphyllum, Klotzsch, ex Meissn. in DC. Prod. xiv. 128 = filicaule.
microphyllum, F. Muell. l. c. 147 = Muehlenbeckia axillaris.
microphyllum, Ruiz, l. c. 98 = aviculare.
microspermum, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 560 = aviculare.
minimum, *S. Wats. in Bot. King's Exped.* 315.—Am. bor. occ.
minus, Ait. ex Meissn. in DC. Prod. xiv. 103 = hydropiperoides.
minus, *Huds. Fl. Angl.* ed. I. 148.—Reg. temp.
minus, Lag. ex Meissn. in DC. Prod. xiv. 118 = P. Persicaria.
minus, Reichb. Pl. Crit. v. t. 493 = mite.
minus, Tenore, Syll. Fl. Neap. 196 = serrulatum.
Miquelianum, Meissn. in DC. Prod. xiv. 92 = plebejum.
miserum, Willd. ex Spreng. Syst. ii. 256 = lacerum.
mite, Pers. Syn. i. 440 = hydropiperoides.
mite, *Schrank, Bair. Fl.* i. 668.—Europ.
mite, Wall. Cat. n. 1721. partim = P. Hydropiper, flaccidum.
mixtum, *Goiran & Lonivi, ex Goiran & Del Fabbro, Spec. Morph. Veg.—Cf. Nuov. Giorn. Bot. Ital.* ix. (1877) 170.—Hab.?
molle, *D. Don, Prod. Fl. Nep.* 72.—Reg. Himal.
molle, Wight, Ic. t. 1807 = polystachyum.
molliaeforme, *Boiss. Diagn.* Ser. I. vii. 84.—Persia; Tibet. occ.
monspeliense, Guss. Fl. Sic. Prod. i. 470 = Bellardi.
monspeliense, Thieb. ex Pers. Syn. i. 439 = aviculare.
monticolum, Steud. Nom. ed. II. ii. 376 = cognatum.
mucronatum, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1841) 115 = paronychioides.
Muhlenbergii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 295.—Am. bor.
multangulare, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 62 = Meissnerianum.
multiflorum, Gueldenst. Reisen, i. 421 (Quid?).—Reg. Caucas.
multiflorum, *Thunb. Fl. Jap.* i. 169.—China; Japon.
multisetum, *Delile, Fl. Aegypt. Illustr.* 60.—Aegypt.
muricatum, *Meissn. Monog.* 74.—Reg. Himal.
muricatum, Wall. Cat. n. 1697 = strigosum.
muticum, Moench, Meth. Suppl. 266 = virginianum.
myrianthum, Boiss. Fl. Orient. iv. 1034 = corrigioloides.
nanum, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 12 = aviculare.
natans, Gueldenst. Reisen, ii. 32 = amphibium.
neglectum, Bess. Enum. Pl. Volh. 45 = arenarium, aviculare.
nepalense, Hort. Elden. ex Fenzl, in Linnaea, xxiv. (1851) 318 = Fagopyrum esculentum.
nepalense, Meissn. Monog. 84; Wall. Cat. n. 1701 = alatum.
nervosum, Wallr. in Linnaea, xiv. (1840) 568; Beitr. Fl. Hercyn. 198 = aviculare.
niloticum, Lehm. ex Steud. Nom. ed. II. ii. 374 (Quid?).—Aegypt.
niloticum, Meissn. in DC. Prod. xiv. 119 = P. Persicaria.
nodosum, Pers. Syn. i. 440 = lapathifolium.
Nočanum, Boiss. in Bot. Zeit. xi. (1853) 734 = argyrocoleon.
nudiflorum, Rafin. New Fl. Am. iv. 51 (Quid?).—Am. bor.
nudum, Dulac, Fl. Hautes-Pyr. 169 = Bellardi.
nummularifolium, *Meissn. in DC. Prod.* xiv. 127.—Reg. Himal.
nutans, Roxb. Hort. Beng. 29; Fl. Ind. ii. 285 = lapathifolium.
obliquum, Kit. in Linnaea, xxxii. (1863) 364 (Quid?).—Hungar.
obtusatum, Steud. ex Meissn. in DC. Prod. xiv. 118 = lapathifolium.
obtusifolium, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = P. Hydropiper.
occidentale, Herb. Berol. ex Meissn. in DC. Prod. xiv. 144 = Fagopyrum tataricum.

POLYGONUM:—
ochreatum, S. G. Gmel. ex Meissn. in DC. Prod. xiv. 144 = divaricatum.
ochreatum, Houtt. Handleid. viii. 467. t. 49. f. 1 = tomentosum.
ochreatum, Willd. ex Meissn. in DC. Prod. xiv. 140 = alpinum.
ocreatum, *Linn. Sp. Pl.* 361.—Sibir.
odoratum, *Lour. Fl. Cochinch.* 243.—Cochinch.
oleraceum, Schur, Enum. Pl. Transs. 585 = P. Hydropiper.
oligophyllum, Boiss. Fl. Orient. iv. 1042 = salicornioides.
Olivieri, Jaub. & Spach, Illustr. ii. 31. t. 121 = polycnemoides.
Olivieri, Meissn. in DC. Prod. xiv. 92 = tubulosum.
orientale, Hort. ex Meissn. l. c. 119 = lapathifolium.
orientale, *Linn. Sp. Pl.* 362.—Geront. trop.
orientale, Wall. Cat. n. 1710 = tomentosum.
oryzetorum, *Blume, Bijdr.* 532 (oryzetum); *Meissn. in DC. Prod.* xiv. 106.—Java.
ovatum, Heyne, in Roth, Nov. Pl. Sp. 206 = chinense.
Owenii, Boj. in Ann. Sc. Nat. Sér. II. iv. (1835) 267. t. 9 (= *Ceratogonum atriplicifolium*).—Afr. trop.
oxyphyllum, Wall. Cat. n. 1715 = amplexicaule.
oxyspermum, C. A. Mey. & Bunge, in Ledeb. Cat. Hort. Dorpat. (1824) 5 = aviculare.
pachyrhizum, Trautv. in Act. Hort. Petrop. ii. (1873) 483 = luzuloides.
pachystachyum, *Rafin. Fl. Ludov.* 29; *et Meissn. Monog.* 80.—Am. bor.
paleaceum, *Wall. Cat.* n. 1684.—Reg. Himal.
pallens, Pers. Syn. i. 439 = lapathifolium.
pallidum, With. Bot. Arr. Brit. Pl. ed. III. ii. 381 = lapathifolium.
paludosum, *Griff. Rep. Bot. Gard. Calc.* 39.—Hab. ?
paludosum, Schur, Enum. Pl. Transs. 585 = mite.
palustre, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 158; *Contrib.* 51.—Reg. Argent.
panduraeforme, Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 48 = chinense, microcephalum.
paniculatum, Andr. ex Meissn. in DC. Prod. xiv. 118 = lapathifolium.
paniculatum, *Blume, Bijdr.* 533.—Reg. Himal.; Java.
pannonicum, *Simk. in Termész. Füzetek*, ii. (1878) 147. —Hungaria.
paraguayense, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 253.—Paraguay.
Paronychia, *Cham. & Schlecht. in Linnaea*, iii. (1828) 51.—Am. bor. occ.
Paronychia, C. A. Mey. Verz. Pfl. Cauc. 158 = paronychioides.
paronychioides, *C. A. Mey. in Bull. Soc. Nat. Mosc.* (1838) 356.—Reg. Cauc. et Himal.
Parryi, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 99.—Am. bor. occ.
parviflorum, Schott, in Oken, Isis, (1818) 822 = aviculare *vel* maritimum ?
parvifolium, Nutt. Gen. Am. i. 256 = Polygonella parvifolia.
patens, D. Don, Prod. Fl. Nep. 73 = chinense.
patulum, Bieb. Fl. Taur. Cauc. i. 304 = Bellardi.
patulum, Hort. ex Meissn. in DC. Prod. xiv. 119 = lapathifolium.
patulum, Karel. ex Meissn. l. c. 100 = arenarium.
pauciflorum, Endl. Gen. Suppl. iv. II. 49 = paronychioides.
pauciflorum, *Maxim. in Ind. Sem. Hort. Petrop.* (1866) 3.—Japon.
Pawlowskianum, *Glehn, in Act. Hort. Petrop.* iv. (1876) 77.—As. centr.
pedunculare, *Wall. Cat.* n. 1718.—As. et Austral. trop.
pensylvanicum, Bunge, Enum. Pl. Chin. Bor. 57 = Bungeanum.
pensylvanicum, Curt. Fl. Lond. fasc. t. 73 = lapathifolium.
pensylvanicum, *Linn. Sp. Pl.* 362.—Am. bor.
perfoliatum, *Linn. Syst.* ed. X. 1005.—Ind. or.; Malaya.
perforatum, Aitch. in Journ. Linn. Soc. xviii. (1880) 90 (non seq.).—Afghan.
perforatum, Meissn. Monog. 83 = alatum.
perforatum, Wall. Cat. n. 1700 = humile.
perpusillum, *Hook. f. in Hook. Ic. Pl.* t. 1490 A.—Reg. Himal.
Perrottetii, Meissn. in DC. Prod. xiv. 94 = exiguum.
Persicaria, Hook. & Arn. Bot. Beech. Voy. 68 = imberbe.

POLYGONUM:—
Persicaria, *Linn. Sp. Pl.* 361.—Reg. bor. temp.
Persicaria, Wahlenb. ex Nym. Consp. 637 = lapathifolium.
Persicaria, Wall. Cat. n. 1720 = glabrum.
persicarioides, *H. B. & K. Nov. Gen. et Sp.* ii. 179.—Mexic.
Persoonii, *Engelm. ex Meissn. in DC. Prod.* xiv. 103. —Texas.
peruvianum, *Meissn. l. c.* 122.—Peruv.
petiolatum, D. Don, Prod. Fl. Nep. 70 = amplexicaule.
pilosum, Roxb. Hort. Beng. 29; Fl. Ind. ii. 286 = orientale.
platycladum, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 73 (patycladium).—N. Caled.
plebeium, *R. Br. Prod.* 420.—Geront. trop. et temp.
pluriflorum, C. Koch, in Linnaea, xxii. (1849) 202 = cognatum.
podocephalum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 136; et ex Meissn. in DC. Prod. xiv. 129 = sphaerocephalum.
Poiretii, Meissn. Monog. 79 = glabrum.
polyanthos, *De Bruyn, in Miq. Pl. Jungh.* 304.—Java.
polybotryum, F. Muell. Fragm. iv. 130 = Muehlenbeckia polybotrya.
polycephalum, Wall. Cat. n. 1707 = chinense.
polycnemoides, *Jaub. & Spach, Illustr.* ii. 30. t. 120. —As. Min.; Persia.
polygaloides, *Meissn. in DC. Prod.* xiv. 101.—Am. bor. occ.
polygamum, Vent. Jard. Cels. 65 = Polygonella parvifolia.
polymorphum, Ledeb. Fl. Ross. iii. 524 = alpinum.
polyneuron, *Franch. & Sav. Enum. Pl. Jap.* ii. 471.—Japon.
polystachyum, *Wall. Cat.* n. 1686.—Reg. Himal.
portoricense, Bert. ex Endl. Gen. Suppl. iv. II. 47 = glabrum.
Posumbu, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 71. —As. or.
Posumbu, Thw. Enum. Pl. Zeyl. 245 = flaccidum.
Posumbu, Wall. Cat. n. 1722 = minus.
procumbens, De Bruyn, in Miq. Pl. Jungh. 307 = P. Posumbu.
procumbens, Gilib. Exercit. ii. 434 = aviculare.
propinquum, *Ledeb. Fl. Ross.* iii. 532.—Japon.
prostratum, *R. Br. Prod.* 419.—Austral.; N. Zel.
prostratum, Roxb. ex D. Don, Prod. Fl. Nep. 72 = plebejum.
provinciale, C. Koch, in Linnaea, xxii. (1849) 204 = aviculare.
Pseudo-hydropiper, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 653 = serrulatum.
Pseudo-lapathum, Schur, Enum. Pl. Transs. 584 = lapathifolium.
pterocarpum, *Wall. Cat.* n. 1690.—Reg. Himal.
Pteropus, Hance, in Journ. Bot. vii. (1869) 167 = Thunbergii.
pubescens, *Blume, Bijdr.* 532.—Java.
pulchellum, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 411 = Bellardi.
pulchrum, Blume, Bijdr. 530 = tomentosum.
punctatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 72 = alatum.
punctatum, Ell. Sketch, i. 455 = acre.
punctatum, Krock. Fl. Siles. Suppl. ii. 57 = lapathifolium ?
punctatum, Rafin. Ann. Nat. (1820) 12 (Quid ?).—Am. bor.
punctatum, Schwein. ex Meissn. in DC. Prod. xiv. 103 = hydropiperoides.
purpureum, Gilib. Exercit. ii. 433 = amphibium.
pusillum, Guss. ex Nym. Consp. 639 = aviculare.
pusillum, Krock. Fl. Siles. Suppl. ii. 64 = seq. ?
pusillum, Lam. Fl. Fr. iii. 235 = minus.
pyramidatum, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 411 = Fagopyrum esculentum.
quadrangulatum, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 353 = Muehlenbeckia tamnifolia.
quadrifidum, Buch.-Ham. ex Meissn. in DC. Prod. xiv. 122 = glabrum.
quadrifidum, Meissn. in Linnaea, xiv. (1840) 485 = lapathifolium.
radicosum, *Boiss. Diagn.* Ser. I. vii. 84.—Persia.
Raii, Bab. in Trans. Linn. Soc. xvii. (1836) 458 = Roberti.

POLYGONUM:—
ramiflorum, Janka, in Linnaea, xxx. (1859–60) 599 = Bellardi.
ramosissimum, *Michx. Fl. Bor. Am.* i. 237.—Am. bor.
ramoso-spicatum, Klotzsch, in Bot. Ergeb. Waldem. Reise, t. 87 = rumicifolium.
ramosum, Fisch. ex Hornem. Hort. Hafn. Suppl. 45 = Laxmanni.
Raptae, Herb. Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 47 = barbatum.
Rayi, Bab. Man. Brit. Bot. ed. II. 275 = *Raii*.
recumbens, *Royle, ex Bab. in Trans. Linn. Soc.* xviii. (1838) 116.—Reg. Himal.
refractum, Mart. ex Meissn. in DC. Prod. xiv. 132 = Meissnerianum.
repens, Meissn. in Wall. Cat. n. 1699 = capitatum.
restionoides, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 1039 = setosum.
reticulatum, De Bruyn, in Miq. Pl. Jungh. 309 = runcinatum.
reticulatum, C. Koch, in Linnaea, xxii. (1849) 200 = Bellardi.
rhaeticum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 115.—Europ.
rhizocaulon, De Bruyn, in Miq. Pl. Jungh. 306 = P. Posumbu.
riparium, Georgi, Bemerk. Reise Russ. Reich. i. 209 = Laxmanni.
riparium, Mart. ex Meissn. in DC. Prod. xiv. 108 = acre.
rivulare, Helfer, ex Meissn. l. c. 107 = flaccidum.
rivulare, Koen. ex Roxb. Hort. Beng. 29; Fl. Ind. ii. 290 = barbatum.
rivulare, *Phil. in Linnaea*, xxx. (1859–60) 199.—Chili.
rivulare, Roth, Nov. Pl. Sp. 206 = barbatum.
Roberti, *Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 409.—Europ.
robustum, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 60.—Java.
Roettleri, Roth, Nov. Pl. Sp. 205 = flaccidum.
romanum, *Jacq. Obs.* iii. 8. t. 58.—Europ. austr.
rottboellioides, Jaub. & Spach, Illustr. ii. 32. t. 122 = tubulosum.
Rottleri, Auct. = *Roettleri*.
Roxburghii, Meissn. in DC. Prod. xiv. 93 = plebejum.
Roylii, *Bab. in Trans. Linn. Soc.* xviii. (1841) 115.—Reg. Himal.
rubicundum, Soland. ex Meissn. in DC. Prod. xiv. 111 = serrulatum.
rubricaule, *Cham. in Linnaea*, viii. (1833) 130.—Bras.
rubricaule, Rafin. New Fl. Am. iv. 51 (Quid ?).—Am. bor.
rubriflorum, Frivald. ex Meissn. in DC. Prod. xiv. 99 = Bellardi.
rude, *Meissn. l. c.* 137.—Reg. Himal.
ruderale, Salisb. Prod. 259 = P. Persicaria.
rugosum, Buch.-Ham. ex Endl. Gen. Suppl. iv. II. 49 = Fagopyrum cymosum.
Ruizianum, *C. Koch, in Linnaea*, xxii. (1849) 207.—Hispan. ?
rumicifolium, Pall. ex Spreng. Syst. ii. 255 = sibiricum.
rumicifolium, *Royle, ex Bab. in Trans. Linn. Soc.* xviii. (1841) 112.—Reg. Himal.
runcinatum, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 73.—Reg. Himal.; Java.
rupestre, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 710 = cognatum.
rurivagum, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 560 = aviculare.
sachalinense, *F. Schmidt, ex Maxim. Prim. Fl. Amur.* 233.—Ins. Sachalin.
sagittatum, Boj. & Hilsenb. ex Baker, in Journ. Linn. Soc. xx. (1883) 239 = Meissnerianum.
sagittatum, D. Don, Prod. Fl. Nep. 73 = strigosum.
sagittatum, *Linn. Sp. Pl.* 363.—As. et. Am. bor.
salicifolium, Brouss. ex Willd. Enum. Hort. Berol. 428; Delile, Fl. Aegypt. 12 = serrulatum.
salicifolium, Schur, Enum. Pl. Transs. 583 = amphibium.
salicornioides, *Jaub. & Spach, Illustr.* ii. 32. t. 123.—Persia; Sind.
salignum, *Willd. Enum. Hort. Berol.* 430.—Sibir.
salsugineum, *Bieb. Beschr. Casp.* 169; *Fl. Taur. Cauc.* i. 304.—Reg. Caucas.
salsuginosum, Wallr. in Linnaea, xiv. (1840) 568; Beitr. Fl. Hercyn. 198 = Roberti.
Sanguinaria, *Remy, in C. Gay, Fl. Chil.* v. 270.—Chili.

POLYGONUM:—
scabridum, Willd. ex Meissn. in DC. Prod. xiv. 117 = persicarioides.
scabrinervium, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1841) 101 = glabrum.
scabrum, Moench, Meth. 629 = lapathifolium.
scabrum, Moench, Meth. Suppl. 267 = pensylvanicum.
scabrum, *Poir. Encyc.* vi. 148.—Afr. bor.
scandens, Linn. Sp. Pl. 364 = dumetorum, Anredera spicata.
scandens, *Vell. Fl. Flum.* 161 ; iv. t. 40.—Bras.
Schurii, Fuss, Fl. Transs. 560 = mite, P. Bistorta.
Schychowscii, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 150, nomen.—Rossia.
scoparium, *Req. ex Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 410.—Corsica.
segetum, *H. B. & K. Nov. Gen. et Sp.* ii. 177.—N. Granat.
senegalense, *Meissn. Monog.* 54.—Afr. bor. et trop.
senticosum, *Franch. & Sav. Enum. Pl. Jap.* i. 401.—Japon.
sericeum, Georgi, Bemerk. Reise Russ. Reich. i. 209 = seq. ?
sericeum, *Pall. Reise*, iii. 286, 320.—Sibir.
serotinum, *Rafin. Ann. Nat.* 12 (Quid ?).—Am. bor.
serotinum, Tenore, Syll. Fl. Neap. App. 566 = P. Persicaria.
serpyllaceum, Jaub. & Spach, Illustr. ii. 24. t. 117 = radicosum.
serratum, *Linn. Sp. Pl.* 360.—Afr. bor.
serratum, Poir. Encyc. vi. 144 = Poiretii.
serrulatum, Guss. ex Tenore, Fl. Nap. Prod. App. v. 12 = minus.
serrulatum, Hohen. ex Boiss. Fl. Orient. iv. 1029 = minus.
serrulatum, *Lag. Gen. et Sp. Nov.* 14.—Geront. temp. et trop.
serrulatum, Spreng. Syst. ii. 256 = glabrum.
setaceum, *Baldw. in Ell. Sketch*, i. 455.—Georgia.
setaceum, Nutt. ex Meissn. in DC. Prod. xiv. 80 = Polygonella gracilis.
setigerum, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 253 = acuminatum.
setosulum, *A. Rich. Tent. Fl. Abyss.* ii. 227.—Abyss.
setosum, *Jacq. Obs.* iii. 8. t. 57.—As. Min.
setosum, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid ?).—Am. bor.
shastense, *Brewer, ex A. Gray, in Proc. Am. Acad.* viii. (1873) 400.—Calif.
sibiricum, *Laxm. in Nov. Act. Petrop.* xviii. (1773) 531. t. 7. f. 2.—Reg. Himal. ; Sibir.
sibiricum, Linn. f. Suppl. 228 = alpinum.
Sieboldi, Hort. ex Meissn. in DC. Prod. xiv. 136 = cuspidatum.
Sieboldii, Meissn. l. c. 133 = sagittatum.
simlense, Royle, ex Bab. in Trans. Linn. Soc. xviii. (1841) 102 = lapathifolium.
sinarum, Desv. ex Meissn. Monog. 62 = Fagopyrum tataricum.
sinense, J. F. Gmel. Syst. 639 = chinense.
sinuatum, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyss.
sinuatum, *Royle, ex Bab. in Trans. Linn. Soc.* xviii. (1841) 108.—Reg. Himal.
somphocarpum, De Bruyn, in Nederl. Kruidk. Arch. i. (1848) 506 = lapathifolium.
songaricum, Schrenk, Enum. Pl. Nov. i. 8 = alpinum.
speciosum, Salisb. Prod. 259 = orientale.
speciosum, Wall. Cat. n. 1716 ; Meissn. Monog. 66 = amplexicaule.
spectabile, Lehm. Ind. Sem. Hort. Hamb. (1820) et (1825) ; ex Linnaea, iii. (1828) 50 = Bellardi.
spectabile, Mart. ex Meissn. in DC. Prod. xiv. 119 = attenuatum.
sphaerocephalum, *Wall. Cat.* n. 1703.—Ind. or.
sphaerostachyum, Meissn. Monog. 53 = macrophyllum.
spinulosum, *Boj. ex Meissn. in DC. Prod.* xiv. 113.—Ins. Maurit.
spirale, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 25 = oryzetorum.
splendens, Klotzsch, in Bot. Ergeb. Waldem. Reise, 136 = macrophyllum.
squamosum, *Rafin. Atl. Journ.* (1832) 16 (Quid ?).—Am. bor.
stagninum, *Buch.-Ham. ex Meissn. in Wall. Pl. As. Rar.* iii. 56.—Ind. or.
staticiflorum, Wall. Cat. n. 1704 = microcephalum.
stelligerum, *Cham. in Linnaea*, viii. (1833) 131.—Bras.

POLYGONUM:—
stenophyllum, Meissn. Monog. 52 = macrophyllum.
stoloniferum, Blanco, Fl. Filip. ed. I. 314 = barbatum.
stoloniferum, F. Schmidt, Reisen Amurl. 168 = Thunbergii.
striatum, Dulac, Fl. Hautes-Pyr. 169 = P. Convolvulus.
striatum, *C. Koch, in Linnaea*, xxii. (1849) 206.—Chili.
strictum, All. Fl. Pedem. ii. 207. t. 68. f. 2 = minus.
strictum, Ledeb. Fl. Alt. ii. 86 = Bellardi.
strictum, Meissn. in Linnaea, xiv. (1840) 485 = serrulatum.
strigillosum, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 57 = barbatum.
strigosum, *R. Br. Prod.* 420.—As. et Austral. trop.
strigosum, Buch.-Ham. ex Meissn. in DC. Prod. xiv. 129 = viscosum.
styligerum, *De Bruyn, in Miq. Pl. Jungh.* 311.—Java.
stypticum, *Cham. & Schlecht. in Linnaea*, iii. (1828) 53.—Bras.
subcordatum, Miq. in Journ. Bot. Néerl. i. (1861) 95 = orientale ?
subincanum, Mart. Hort. Erlang. (1814) 95 (Quid ?).—Hab. ?
subsessile, *R. Br. Prod.* 419.—Austral. ; N. Caled.
subspathaceum, D. Dietr. Syn. Pl. ii. 1320, sphalm. = Polypogon maritimus.
suffruticosum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 652 = equisetiforme.
suffultum, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 233.—Japon.
sumatranum, *De Bruyn, in Miq. Pl. Jungh.* 308.—Sumatra.
tamnifolium, H. B. & K. Nov. Gen. et Sp. ii. 180 = Muehlenbeckia tamnifolia.
tarmense, Ruiz, ex Meissn. in DC. Prod. xiv. 117 = persicarioides.
tataricum, Linn. Sp. Pl. 364 = Fagopyrum tataricum.
tataricum, Wall. ex D. Don, Prod. Fl. Nep. 72 = tortuosum.
tenellum, Blume, Bijdr. 530 ; et ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 57 = minus.
tenellum, Roxb. Hort. Beng. 29 ; Fl. Ind. ii. 289 = flaccidum.
tenerum, *Spreng. Syst.* ii. 254.—Georgia.
tenue, D. Don, Prod. Fl. Nep. 70 = macrophyllum.
tenue, *Michx. Fl. Bor. Am.* i. 238.—Am. bor.
tenuicaule, *Biss. & S. Moore, in Journ. Bot.* xvi. (1878) 135.—Japon.
tenuiflorum, Presl, Delic. Prag. 67 = lapathifolium.
tenuifolium, Steph. ex Ledeb. Fl. Ross. iii. 531 = salsugineum.
tephioides, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 158 ; *Contrib.* 52.—Reg. Argent.
terrestre, Hegetschw. Fl. Schw. 366 = amphibium.
tetragonum, Blume, Bijdr. 535 = pedunculare.
Thevenaei, *Lespin. in Bull. Soc. Bot. Fr.* vi. (1859) 657.—Gallia.
Thunbergii, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 84.—Japon.
thymifolium, *Jaub. & Spach, Illustr.* ii. 22. t. 116.—Persia.
tinctorium, *Ait. Hort. Kew.* ed. II. 31.—China.
tomentosum, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 113 = adenophyllum.
tomentosum, Schrank, Baier. Fl. i. 669 = lapathifolium.
tomentosum, Wall. Cat. n. 1709/4 = stagninum.
tomentosum, *Willd. Sp. Pl.* ii. 447.—Geront. trop.
torquatum, De Bruyn, in Miq. Pl. Jungh. 303 = orientale.
Torreyi, S. Wats. in Am. Natural. vii. (1873) 664 = minimum.
tortuosum, *D. Don, Prod. Fl. Nep.* 71.—Reg. Himal.
triandrum, Roland. ex Rottb. in Lit. Univ. Hafn. i. (1778) 274 = Triplaris americana.
triangulare, Wall. Cat. n. 1689 = Fagopyrum cymosum.
tripterocarpum, *A. Gray, ex Rothr. in Ann. Rep. Smithson. Inst.* 1867 (1868) 453.—Am. arct.
tristachyum, *Baker, in Journ. Bot.* xx. (1882) 244.—Madag.
tropicum, *C. Koch, in Linnaea*, xxii. (1849) 205.—Afr. trop.
truncatum, *A. Rich. Fl. Cub. Fanerog.* ii. 182.—Cuba.

POLYGONUM:—
truncatum, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 214 = runcinatum.
tubulosum, *Boiss. Diagn.* Ser. I. vii. 83.—Reg. Himal.
tumidum, Auch. ex Meissn. in DC. Prod. xiv. 86 = equisetiforme.
tumidum, *Delile, Fl. Aegypt. Illustr.* 60.—Aegypt.
turgidum, Thuill. Fl. Par. ed. II. i. 199 = lapathifolium.
typhoniifolium, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 239 = senticosum.
undulatum, Berg. Desc. Pl. Cap. 135 = atraphaxoides.
undulatum, Murr. in Comm. Gotting. v. 1774 (1775) 34. t. 5 = alpinum.
utriculatum, *Remy, in C. Gay, Fl. Chil.* iv. 267.—Chili.
uviferum, Linn. Sp. Pl. 365 = Coccoloba uvifera.
vaccinifolium, *Wall. Cat.* n. 1695.—Reg. Himal.
Venantianum, Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 319 ; Sert. 83. t. 3 = arenarium.
vernum, Rafin. Fl. Ludov. 30 = lapathifolium.
virgatum, Cham. & Schlecht. in Linnaea, iii. (1828) 45 = hydropiperoides.
virgatum, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 410 = Bellardi.
virgatum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 353 = arenarium.
virginianum, *Linn. Sp. Pl.* 360.—Reg. Himal. ; Am. bor.
viridiflorum, Poir. Encyc. vi. 140 = Cannabis sativa.
viscosum, Bast. ex Meissn. in DC. Prod. xiv. 119 = lapathifolium.
viscosum, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 71.—Reg. Himal.
viviparum, *Linn. Sp. Pl.* 360.—Reg. bor. et arct.
viviparum, Wall. ex D. Don, Prod. Fl. Nep. 70 = affine.
volcanicum, Benth. Pl. Hartw. 81 = Muehlenbeckia vulcanica.
volubile, Gilib. Exercit. ii. 435 = P. Convolvulus.
volubile, Turcz. in Bull. Soc. Nat. Mosc. (1840) 77 = Fagopyrum cymosum.
vulgare, Gueldenst. Reisen, ii. 44 = minus ?
Wallichii, *Meissn. Monog.* 83. t. 7.—Reg. Himal.
Wallichii, Wight, Ic. t. 1805 = sphaerocephalum.
Weyrichii, *F. Schmidt, in Maxim. Prim. Fl. Amur.* 234.—Ins. Sachalin.
Zippelii, *Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 64.—N. Guin.
zonulatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 = chinense.

POLYGYNE, Phil. in Linnaea, xxxiii. (1864–65) 170 = **Plagiocheilus**, Arn. (Compos.).
inconspicua, Phil. l. c. 171.—Chili.

POLYLEPIS, Ruiz & Pav. Prod. 80. t. 15 (1794).
ROSACEAE, Benth. & Hook. f. i. 623.
QUINASIS, Rafin. Sylva Tellur. 150 (1838).
hypoleuca, Wedd. Chlor. And. ii. t. 78 = racemosa.
incana, *H. B. & K. Nov. Gen. et Sp.* vi. 227.—Peruv.
lanuginosa, *H. B. & K. l. c.* 228.—Ecuador.
racemosa, *Ruiz & Pav. Syst. Veg.* i. 139.—Peruv.
sericea, *Wedd. Chlor. And.* ii. 238.—N. Granat. ; Venezuela.
tomentella, *Wedd. l. c.* 237. t. 78.—Peruv. ; Bolivia.
villosa, *H. B. & K. Nov. Gen. et Sp.* vi. 228.—Peruv.

POLYLOBIUM, Eckl. & Zeyh. Enum. 180 (1836) = **Lotononis**, DC. (Legumin.).
angustifolium, Eckl. & Zeyh. l. c. 182 = L. involucrata.
brachylobum, D. Dietr. Syn. Pl. iv. 962 = L. brachyloba.
calycinum, D. Dietr. l. c. = L. calycina.
carinatum, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. carinata.
corymbosum, Benth. l. c. = L. corymbosa.
debile, Eckl. & Zeyh. Enum. 181 = L. debilis.
erubescens, D. Dietr. Syn. Pl. iv. 962 = L. pumila.
falcatum, D. Dietr. l. c. = L. falcata.
fastigiatum, Eckl. & Zeyh. Enum. 183 = L. angustifolia.
filiforme, Eckl. & Zeyh. l. c. 181 = L. umbellata.
humifusum, D. Dietr. Syn. Pl. iv. 962 = L. humifusa.

POLYLOBIUM:—
intermedium, Eckl. & Zeyh. Enum. 181 = Lotononis trichopoda.
involucratum, Eckl. & Zeyh. l. c. 182 = L. involucrata.
micranthum, Presl, Bot. Bemerk. 123 = L. micrantha.
molle, D. Dietr. Syn. Pl. iv. 962 = L. mollis.
Mundianum, Eckl. & Zeyh. Enum. 183 = L. angustifolia.
oxypterum, Presl, Bot. Bemerk. 123 = L. oxyptera.
pallens, Eckl. & Zeyh. Enum. 182 = L. pallens.
pedunculare, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. peduncularis.
polycephalum, D. Dietr. Syn. Pl. iv. 962 = L. polycephala.
quinatum, Presl, Bot. Bemerk. 123 = L. quinata.
radulum, D. Dietr. Syn. Pl. iv. 962 = L. humifusa.
sparsiflorum, Eckl. & Zeyh. Enum. 181 = L. oxyptera.
tenellum, D. Dietr. Syn. Pl. iv. 962 = L. tenella.
tenuifolium, Eckl. & Zeyh. Enum. 182 = L. involucrata
trichopodum, Presl, Bot. Bemerk. 123 = L. trichopoda.
truncatum, Eckl. & Zeyh. Enum. 181 = L. umbellata.
typicum, Eckl. & Zeyh. l. c. 180 = L. trichopoda.
umbellatum, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. umbellata.
varium, Presl, Bot. Bemerk. 135 = L. varia.

POLYLOPHIUM, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 47. *UMBELLIFERAE*, Benth. & Hook. f. i. 929.
ACANTHOPLEURA, C. Koch, in Bot. Zeit. vii. (1849) 408.
GALBANON, Adans. Fam. ii. 499 (1763) nomen prius.
GALBANUM, D. Don, in Phil. Mag. ix. (1833) 46; et in Trans. Linn. Soc. xvi. (1833) 603.
involucratum, Boiss. Fl. Orient. ii. 1066 = orientale.
orientale, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 48.—Persia.
petrophilum, Boiss. Fl. Orient. ii. 1066 = thalictroides.
thalictroides, *Fenzl, in Tchihat. Asie Min. Bot.* i. 443.—As. Min.

POLYMERIA, R. Br. Prod. 488 (1810). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 875.
ambigua, *R. Br. l. c.*—Austral.
angusta, *F. Muell. Fragm.* vi. 100.—Austral.
calycina, *R. Br. Prod.* 488.—Austral.
distigma, *Benth. Fl. Austral.* iv. 433.—Austral.
lanata, R. Br. Prod. 488 = ambigua.
longifolia, *Lindl. in Mitch. Journ. Trop. Austral.* 398.—Austral.
marginata, *Benth. Fl. Austral.* iv. 432.—Austral.
occidentalis, *F. Muell. Census*, 95, nomen.—Austral.
pusilla, *R. Br. Prod.* 488.—Austral.
quadrivalvis, R. Br. l. c. = ambigua.

POLYMNIA, Linn. Diss. Chen. 22 (1751); Amoen. Acad. iii. 15 (1756). *COMPOSITAE*, Benth. & Hook. f. ii. 346.
ALYMNIA, Neck. Elem. i. 31 (1790).
POLYMNIASTRUM, Lam. Illustr. t. 712 (1798); iii. 287 (1823).
abyssinica, Linn. f. Suppl. 383 = Guizotia abyssinica.
africana, Poir. Encyc. Suppl. iv. 483 = Wedelia africana.
aspera, *Mart. ex DC. Prod.* v. 515.—Mexic.
australis, Spreng. ex DC. Prod. v. 542 = Wedelia latifolia.
benghalensis, Poir. ex Steud. Nom. ed. II. ii. 377 = Wedelia calendulacea.
calendulacea, Poir. Encyc. Suppl. iv. 482 = Wedelia calendulacea.
calicina, Poir. l. c. = Wedelia affinis, acapulcensis.
canadensis, *Linn. Sp. Pl.* 926.—Am. bor.
carnosa, Linn. f. Suppl. 384 = Didelta carnosa.
caroliniana, Poir. Encyc. v. 505 = Berlandiera tomentosa.
crenata, Poir. Encyc. Suppl. iv. 482 = Wedelia carnosa.
cruciana, Poir. l. c. = Wedelia cruciana.
dentata, *Moench, Meth. Suppl.* 242.—Hab.?
edulis, *Wedd. in Ann. Sc. Nat.* Sér. IV. vii. (1857) 114.—Am. austr.
frondosa, Bruce, Travels, vii. t. 52 = Guizotia abyssinica.
fruticosa, *Benth. Pl. Hartw.* 209.—N. Granat.
glabrata, *DC. Prod.* v. 515.—Chili.
gracilis, Poir. Encyc. Suppl. iv. 483 = Wedelia gracilis.
grandis, Hort. Berol. ex Kunth, Ind. Sem. Hort. Berol. (1847) 13 = Montañoa bipinnatifida.

POLYMNIA:—
graveolens, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 299.—Guian.
macrophylla, *Rafin. Fl. Ludov.* 70 (Quid?).—Am. bor.
macroscypha, *Baker, in Mart. Fl. Bras.* vi. III. 158.—Bras.
maculata, *Cav. Ic.* iii. 14. t. 227.—Mexic.
mollis, Poir. Encyc. Suppl. iv. 482 = Wedelia mollis.
parviflora, Poir. l. c. = Wedelia parviflora.
perfoliata, Poir. Encyc. v. 506 = Melampodium perfoliatum.
pyramidalis, *Triana, in Ann. Sc. Nat.* Sér. IV. ix. (1858) 40.—N. Granat.
revoluta, Auct. ex Steud. Nom. ed. II. ii. 377 = Melanopodium perfoliatum.
riparia, *H. B. & K. Nov. Gen. et Sp.* iv. 282.—N. Granat.
scabra, Zucc. in Abh. Akad. Muench. i. (1832) 313 = aspera.
serrata, Poir. Encyc. Suppl. iv. 482 = Wedelia serrata.
Siegesbeckia, *DC. Prod.* v. 516.—Bras.
silphioides, *DC. l. c.*—Bras.
sonchifolia, Griseb. in Goett. Abh. xxiv. (1879) 188 = silphioides.
sonchifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 47. t. 254.—Peruv.
spinosa, Linn. f. Suppl. 384 = Didelta spinosa.
Tetragonotheca, Linn. Syst. ed. XII. 576 = Tetragonotheca helianthoides.
Uvedalia, *Linn. Sp. Pl.* ed. II. 1303.—Am. bor.
variabilis, *Poir. Encyc.* v. 505.—Hab.?
Wedelia, Linn. Mant. i. 118 = Wedelia frutescens.
Wedelia, Mill. Gard. Dict. ed. VIII. n. 1 = P. Uvedalia.

POLYMNIASTRUM, Lam. Illustr. t. 712 (1798); iii. 287 (1823) = **Polymnia**, Linn. (Compos.).
urticaefolium, Cass. in Dict. Sc. Nat. lix. 246 = Polymnia variabilis.
variabile, Lam. Illustr. iii. 287 = Polymnia variabilis.

POLYNOME, Salisb. Gen. Pl. Fragm. 12 (1866) = **Dioscorea**, Plum.
alata, Salisb. l. c. = D. alata.
bulbifera, Salisb. l. c. = D. bulbifera.

POLYODON, H. B. & K. Nov. Gen. et Sp. i. 174. t. 55 (1815) = **Bouteloua**, Lag. (Gramin.).
distichus, H. B. & K. l. c. 175 = B. disticha.

POLYODONTIA, Meissn. Gen. 102 (1837) = POLYDONTIA, Blume = **Pygeum**, Gaertn. (Rosac.)

POLYOSMA, Blume, Bijdr. 658 (1825). *SAXIFRAGEAE*, Benth. & Hook. f. i. 647.
alangiacea, *F. Muell. Fragm.* viii. 8.—Austral.
chinensis, *Walp. Rep.* ii. 937.—China.
Cunninghamii, *Benn. Pl. Jav. Rar.* 196.—Austral.
discolor, *Baill. Adansonia*, x. (1871–72) 343.—N. Caled.
fragrans, *Benn. Pl. Jav. Rar.* 196.—Malaya.
fragrans, Wall. Cat. n. 8472, partim = integrifolia.
helicioides, *F. Muell. Papuan Pl.* ii. 8.—N. Guin.
ilicifolia, *Blume, Bijdr.* 658.—Java.
integrifolia, *Blume, l. c.* 659.—Ind. or.; Malaya.
laete-virens, Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 409 = mutabilis.
mutabilis, *Blume, Mus. Bot. Lugd. Bat.* i. 261.—Malaya.
Pancheri, *Baill. Adansonia*, x. (1871–72) 342.—N. Caled.
serrulata, *Blume, Bijdr.* 659.—Java.
spicata, *Baill. Adansonia*, x. (1871–72) 343.—N. Caled.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 261.—Java.
Wallichii, Benn. Pl. Jav. Rar. 196 = integrifolia.

POLYOSUS, Lour. Fl. Cochinch. 75 (1790).—Cf. **Polyozus**, Lour.

POLYOTUS, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 199 = **Gomphocarpus**, R. Br. (Asclep.).
angustifolius, Nutt. l. c. 201 = Asclepias stenophylla.
heterophyllus, Nutt. l. c. 199 = G. viridiflorus.
lanuginosus, Nutt. l. c. 200 (= *Acerates lanuginosa*).—Am. bor.
longifolius, Nutt. l. c. = Asclepias longifolia.
obovatus, Nutt. l. c. 201 = Asclepias obovata?

POLYOZUS, Blume, Bijdr. 947 (1826) = **Psychotria**, Linn. (Rubiac.).
acuminata, Blume, l. c. 948 = Ps. rostrata.
barbata, DC. Prod. iv. 495.—Ins. Hominoa.
latifolia, Blume, Bijdr. 948 = Ps. rostrata.
maderaspatana, DC. Prod. iv. 495 = Webera corymbosa.

POLYOZUS, Lour. Fl. Cochinch. 74 (1790), partim. Gen. dub. *RUBIACEARUM.*—Cf. Benth. & Hook. f. ii. 29.
POLYOSUS, Lour. l. c. 75 (1790).
bipinnata, Blanco, Fl. Filip. ed. II. 43 = Arthrophyllum bipinnatum.
bipinnata, *Lour. Fl. Cochinch.* 74.—Cochinch.
lanceolata, *Lour. l. c.* 75.—China.

POLYPAPPUS, Less. in Linnaea, iv. (1829) 314 = **Baccharis**, Linn. (Compos.).
discolor, DC. Prod. vii. 281 = Vanillosmopsis discolor.
parviflorus, DC. l. c. v. 398 = B. parviflora.
triplinervis, Less. Syn. Comp. 204 = B. triplinervis.

POLYPAPPUS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 178 = **Tessaria**, Ruiz & Pav. (Compos.).
sericeus, Nutt. l. c. = T. borealis.

POLYPARA, Lour. Fl. Cochinch. 61 (1790) = **Houttuyna**, Thunb. (Piper.).
cochinchinensis, Lour. l. c. = H. cordata.
cordata, Buek, Index ad DC. Prod. iv. 310 = praec.

POLYPHEMA, Lour. Fl. Cochinch. 546 (1790) = **Artocarpus**, Forst. (Urtic.).
Champeden, Lour. l. c. 547 = A. Polyphema.
Jaca, Lour. l. c. 546 = A. integrifolia.

POLYPHRAGMON, Desf. in Mém. Mus. Par. vi. (1820) 5. t. 2 = **Timonius**, Rumph. (Rubiac.).
amboinicum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 242.—Ins. Amboin.
anodon, Miq. l. c.—Ins. Amboin.
compressicaule, Miq. Fl. Ind. Bat. ii. 235.—Sumatra.
flavescens, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 134 = T. Jambosella.
hydrangeaefolium, Miq. Fl. Ind. Bat. ii. 235.—Sumatra.
minus, A. Rich. Diss. ex DC. Prod. iv. 445; et in Mém. Soc. Hist. Nat. Par. v. (1834) 231.—Hab.?
mutabile, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 240 = T. mutabilis.
ovale, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 216.—Borneo.
Pseudo-capitatum, Scheff. in Ann. Jard. Buitenz. i. (1876) 29.—Malaya.
rigidum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 243.—Hab.?
rufescens, Miq. l. c. 241.—Ins. Ternate.
sericanthum, Miq. l. c. 242 = Bobea sericantha, inaequisepala.
sericeum, Desf. in Mém. Mus. Par. vi. (1820) 6. t. 2 = T. Rumphii.
sessile, Scheff. in Ann. Jard. Buitenz. i. (1876) 67, nomen.—Malaya.
spiciflorum, Zipp. ex Span. in Linnaea, xv. (1841) 317 = *sericeum*.
splendens, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 241.—Sumatra.
stipulosum, Scheff. in Ann. Jard. Buitenz. i. (1876) 67, nomen.—Malaya.
trichocaulon, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 241.—Malaya.

POLYPLEVRUM, Tayl. ex Tul. in Arch. Mus. Par. vi. (1852) 118 = **Podostemon**, Michx.
orientale, Tayl. ex Tul. l. c. = Podost. Wallichii.

POLYPODIOPSIS, Carr. Conif. ed. II. 710 (1867). Gen. dub. *CONIFERARUM.*
Muellerii, Carr. l. c.—N. Caled.

POLYPODIUM *spinulosum*, Burm. f. Fl. Ind. 223. t. 67. f. 1 = Synaphea petiolaris, R. Br. (Proteac.), fide Meissn. in DC. Prod. xiv. 315.

POLYPOGON, Desf. Fl. Atlant. i. 66 (1798). *GRAMINEAE*, Benth. & Hook. f. iii. 1150.
NOWODWORSKYA, J. & C. Presl, Rel. Haenk. i. 238. t. 40, 351 (1830).
RASPAILIA, J. & C. Presl, l. c. (1830).
SANTIA, Savi, in Mem. Soc. Ital. Mod. viii. II. (1798) 479.

POLYPOGON :—
adscendens, Guss. in Bertol. Fl. Ital. ii. 777 = littoralis.
Adscensionis, *Trin. Agrost.* ii. 11.—Ins. Adscens.
affinis, Brongn. in Duperr. Voy. Coq. Bot. 19 = elongatus.
affinis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 141 = interruptus.
alopecuroides, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 88 = Agrostis exarata.
alpestris, Spreng. Syst. i. 243 = Muehlenbergia alpestris.
ascendens, Boiss. Fl. Orient. v. 521 = *adscendens*.
australis, Brongn. in Duperr. Voy. Coq. Bot. 21 = crinitus.
australis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 141 = interruptus.
brachyphyllus, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 552.—Mexic.
cachinalensis, *Phil. Fl. Atac.* 54.—Chili.
capensis, Hornem. ex Trin. in Mém. Acad. Sc. Pétersb. Sér. VI. i. (1831) 71 = Danthonia micrantha.
Chaetotropis, Trin. l. c. iv. II. (1845) 262 = Chaetotropis chilensis.
chonoticus, *Hook. f. Fl. Antarct.* 374.—Chili.
ciliatus, Spreng. Syst. i. 243 = Muehlenbergia ciliata.
crinitus, Nutt. Gen. Am. i. 50 = monspeliensis.
crinitus, *Trin. Gram. Unifl.* 171.—Chili.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 313.—Cuba.
demissus, *Steud. Syn. Pl. Gram.* 422.—Japon.
Drummondi, Steud. l. c. 184 = tenellus.
elongatus, *H. B. & K. Nov. Gen. et Sp.* i. 134.—Am. austr.
elongatus, Lag. Gen. et Sp. Nov. 3 = littoralis.
fasciculatus, Pers. Syn. i. 80 = Chaeturus fasciculatus.
filiformis, Spreng. ex Kunth, Enum. Pl. i. 39 = Reynaudia filiformis.
flavescens, *J. & C. Presl, Rel. Haenk.* i. 235.—Peruv.
fugax, *Nees, in Steud. Syn. Pl. Gram.* 184.—Austral.
glomeratus, Willd. Enum. Hort. Berol. 87 = Muehlenbergia glomerata.
gracilis, Spreng. Syst. v. 558 (Index) = Muehlenbergia tenella.
Higegaweri, *Steud. Syn. Pl. Gram.* 422.—Japon.
inaequalis, Trin. ex Steud. Nom. ed. II. ii. 378 = elongatus.
intermedius, *Carmich. in Trans. Linn. Soc.* xii. (1819) 504.—Ins. Trist. d'Acugn.
intermedius, Guss. ex Nym. Consp. 803 = monspeliensis.
interruptus, *H. B. & K. Nov. Gen. et Sp.* i. 134. t. 44.—Am. austr.
Lagascae, Roem. & Schult. Syst. ii. 326 = littoralis.
linearis, *Trin. in Linnaea*, x. (1836) 301.—Chili.
littoralis, Aitch. in Journ. Linn. Soc. xix. (1882) 193 = Agrostis verticillata.
littoralis, *Sm. Comp. Fl. Brit.* ed. II. 13.—Europ.
longiflorus, *Nees, ex Steud. Nom.* ed. II. ii. 378; *et ex Trin. in Mém. Acad. Pétersb.* Sér. VI. iv. II. (1845) 263.—Chili.
maritimus, Kunze, in Flora, xxix. (1846) 748 = monspeliensis.
maritimus, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 442.—Europ. austr.
mexicanus, *Schlecht. Hort. Hal.* (1842); *ex Steud. Syn. Pl. Gram.* 185.—Mexic.
mexicanus, Spreng. ex Steud. Nom. ed. II. i. 198, in syn. = Chaetium bromoides.
microstachys, *Phil. Fl. Atac.* 54.—Chili.
monspeliensis, *Desf. Fl. Atlant.* i. 66.—Amphig. trop. et temp.
nepalensis, *Nees, ex Steud. Nom.* ed. II. ii. 378; *et ex Trin. in Mém. Acad. Pétersb.* Sér. VI. iv. II. (1845) 255.—Nepal.
paniceus, Lag. Gen. et Sp. Nov. 3 = monspeliensis.
pauciflorus, Spreng. Syst. i. 243 = Agrostis pauciflora.
polysetus, Steud. in Flora, xii. (1829) 467 = monspeliensis.
pumilus, [Soland. in] Clarke, Trav. iii. 41 = maritimus.
racemosus, Nutt. Gen. Am. i. 51 = Muehlenbergia glomerata.
radicans, *Steud. Syn. Pl. Gram.* 184.—Chili.
sericeus, Spreng. Syst. i. 243 = Muehlenbergia capillaris.
setosus, Spreng. Fl. Hal. Mant. 31 = Muehlenbergia glomerata.
simplex, Spreng. Syst. iv. Cur. Post. 30 = Muehlenbergia maritima?
spathaceus, Steud. Nom. ed. II. ii. 378 = maritimus.

POLYPOGON :—
spicatus, Spreng. Syst. i. 243 = Muehlenbergia spicata.
strictus, *Nees, in Linnaea*, vii. (1832) 297.—Afr. austr.
subspathaceus, Req. in Ann. Sc. Nat. Sér. I. iv. (1825) 386 = maritimus.
subspicatus, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 443 = Chaeturus fasciculatus.
tataricus, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 3 = maritimus.
tenellus, *R. Br. Prod.* 173.—Austral.
tenellus, Spreng. Syst. i. 243 = Muehlenbergia tenella.
tenuis, Brongn. in Duperr. Voy. Coq. Bot. 22 = strictus.
vaginatus, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 44 = Alopecurus vaginatus.
zeylanicus, *Nees, ex Steud. Syn. Pl. Gram.* 183.—Zeylan.

POLYPOMPHOLYX, Lehm. Nov. Stirp. Pugill. viii. 48 (1844). *LENTIBULARIEAE*, Benth. & Hook. f. ii. 988.
Cosmiza, Rafin. Fl. Tellur. iv. 110 (1836).
Tetralobus, A. DC. in DC. Prod. viii. 667 (1844).
bicolor, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 160, nomen.—Guian.
Endlicheri, Lehm. Nov. Stirp. Pugill. viii. 48 = multifida.
exigua, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 50 = tenella.
laciniata, *Benj. in Linnaea*, xx. (1847) 496.—Bras.
latiloba, Lehm. Nov. Stirp. Pugill. viii. 49 = multifida.
multifida, *F. Muell. Fragm.* vi. 162.—Austral.
tenella, *Lehm. Nov. Stirp. Pugill.* viii. 50.—Austral.

POLYPORANDRA, Becc. Malesia, i. (1877) 125. *ICACINEAE*.
scandens, *Becc. l. c.*—Malaya.

POLYPREMUM, Adans, Fam. ii. 152 (1763) = **Valerianella**, Moench (Valer.).

POLYPREMUM, Linn. Syst. ed. I (1735). *LOGANIACEAE*, Benth. & Hook. f. ii. 791.
Cleyera, Adans. Fam. ii. 224 (1763).
Symphoranthus, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 219.
Linnaei, Michx. Fl. Bor. Am. i. 83 = procumbens.
procumbens, *Linn. Sp. Pl.* 111.—Am. bor.
Schlechtendahlii, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 350 = procumbens.

POLYPTERIS, Less. in Linnaea, vi. (1831) 518 = **Gaillardia**, Fouger. (Compos.).
brasiliensis, Less. l. c. = G. megapotamica.
integrifolia, DC. Prod. v. 659 = G. lanceolata.

POLYPTERIS, Nutt. Gen. Am. ii. 139 (1818). *COMPOSITAE* [Benth. & Hook. f. ii. 405].
callosa, *A. Gray, in Proc. Am. Acad.* xix. (1884) 30.—Am. bor. occ.
Hookeriana, *A. Gray, l. c.*—Texas; Mexic.
integrifolia, *Nutt. Gen. Am.* ii. 139.—Am. bor.
texana, *A. Gray, in Proc. Am. Acad.* xix. (1884) 30.—Texas; Mexic.

POLYRHAPHIS, Lindl. Veg. Kingd. 115 (1847) = **Pappophorum**, Schreb. (Gramin.).

POLYSCALIS, Wall. Cat. n. 6939 (1832) = **Cyathula**, Lour. (Amarant.).
capitata, Wall. l. c. n. 6940 = C. capitata.
sequax, Wall. l. c. n. 6939 = C. tomentosa.

POLYSCELIS, Hook. f. Fl. Brit. Ind. iv. 722, 723 (1885) = praec.

POLYSCHEMONE, Schott, Nym. & Kotschy, Analect. Bot. 55 (1854) = **Lychnis**, Tourn. (Caryoph.).
nivalis, Schott, Nym. & Kotschy, l. c. 56 = L. nivalis.

POLYSCHISMA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 269 = **Pelargonium**, L'Hérit. (Geran.).
anatymbicum, Turcz. l. c. 270 = Pel. amatymbicum.

POLYSCHISTIS, J. & C. Presl, Rel. Haenk. i. 294. t. 41. f. 12–18 (1830). Gen. dub. *GRAMINEARUM*.—Cf. Benth. & Hook. f. iii. 1096.
paupercula, *J. & C. Presl, l. c.*—Ins. Luzon.

POLYSCIAS, Forst. Char. Gen. 63. t. 32 (1775). *ARALIACEAE*, Benth. & Hook. f. i. 941.
Botryopanax, Miq. Ann. Mus. Bot. Lugd. Bat. i. 5 (1863).
Eupteron, Miq. Fl. Ind. Bat. i. I. 762 (1855).
Gastonia, Roep. in Bot. Zeit. vi. (1848) t. 1.
Grotefendia, Seem. Journ. Bot. ii. (1864) 247.
acuminata, *Seem. Journ. Bot.* iii. (1865) 181.—Ind. or.
australiana, F. Muell. Fragm. iv. 120 = Hedera australiana.
Ayresii, *Baker, Fl. Maurit.* 127.—Ins. Maurit.
Cumingiana, F. Villar, in Blanco, Fl. Phillip. ed. III. Nov. App. 102 = Panax Cumingianum.
cupularis, *Baker, Fl. Maurit.* 127.—Ins. Maurit.
cutispongia, *Baker, l. c.*—Ins. Mascar.
dichroostachya, *Baker, l. c.*—Ins. Maurit.
digitata, Blanco, Fl. Filip. ed. I. 224 = Heptapleurum rigidum.
disperma, Blanco, l. c. 226 = Arthrophyllum diversifolium.
nodosa, *Seem. Journ. Bot.* iii. (1865) 181.—Malaya.
paniculata, *Baker, Fl. Maurit.* 126.—Ins. Maurit.
papuana, *Seem. Journ. Bot.* iii. (1865) 181.—N. Guin.
pinnata, *Forst. Char. Gen.* 63. t. 32.—Ins. Pacif.
pinnata, Vitm. Summa Pl. ii. 448 = umbellata.
repanda, *Baker, Fl. Maurit.* 128.—Ins. Maurit. et Rodrig.
Rheedii, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 87 = Heptapleurum tomentosum.
sechellarum, *Baker, Fl. Maurit.* 128.—Ins. Seychelles.
umbellata, *Spreng. ex Steud. Nom.* ed. I. 646.—Ins. Pacif.

POLYSOLENIA, Hook. f. in Benth. & Hook. f. Gen. ii. 68 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Wallichii, *Hook. f. l. c.*—Reg. Himal.

POLYSPATHA, Benth. in Hook. Niger Fl. 543 (1849). *COMMELINACEAE*, Benth. & Hook. f. iii. 849.
Eckloniana, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 174 = Commelina Eckloniana.
paniculata, *Benth. in Hook. Niger Fl.* 543.—Afr. trop.

POLYSPHAERIA, Hook. f. in Benth. & Hook. f. Gen. ii. 108 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
lanceolata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 128.—Afr. trop.
ligustriflora, Vatke, in Linnaea, xl. (1876) 193 = Olinia cymosa.
multiflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 127.—Afr. trop.
parvifolia, *Hiern, l. c.* 128.—Afr. trop.
Schweinfurthii, *Hiern, l. c.*—Afr. trop.

POLYSPORA, Sweet, Hort. Brit. ed. I. 61 (1826) = **Gordonia**, Ellis (Ternstr.).
axillaris, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 143, 181 (Mém. Ternstr. 91) = Adinandra acuminata.
axillaris, Sweet, Hort. Brit. ed. I. 61; D. Don, in Hook. Bot. Mag. t. 4019 = G. anomala.
Lessertii, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 143 = G. anomala.

POLYSTACHYA, Hook. Exot. Fl. t. 103 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 540.
Encyclia, Poepp. & Endl. Nov. Gen. et Sp. ii. 10 (1838).
Epiphora, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 201.
abbreviata, *Reichb. f. in Gard. Chron.* (1874) II. 291.—Bras.
Adansoniae, *Reichb. f. in Flora*, xlviii. (1865) 185.—Afr. trop.
affinis, *Lindl. Gen. et Sp. Orch.* 73.—Afr. trop.
alpina, *Lindl. in Journ. Linn. Soc.* vi. (1862) 131.—Afr. trop.
anceps, *Ridl. l. c.* xxi. (1885) 473.—Madag.
angularis, *Reichb. f. in Flora*, l. (1867) 114.—Afr. trop.
Bennettiana, *Reichb. f. Otia Bot. Hamb.* 113.—Abyss.

POLYSTACHYA:—
bifida, *Lindl. in Journ. Linn. Soc.* vi. (1862) 129.—Afr. trop.
bracteosa, *Lindl. Bot. Reg.* (1840) *Misc.* 48.—Afr. trop.
caduca, *Reichb. f. Otia Bot. Hamb.* 110.—Afr. trop.
caespitosa, *Rodrig. Orch. Nov.* ii. 168.—Bras.
caloglossa, *Reichb. f. Otia Bot. Hamb.* 111.—Afr. trop.
capensis, Sond. ex Harv. Thes. Cap. ii. 51 = Ottoniana.
caracasana, *Reichb. f. in Bonplandia*, ii. (1854) 15.—Venezuela.
carnea, *A. Br. in Fl. des Serres*, xv. (1862-63) 45.—Afr. trop.
cerea, *Lindl. Bot. Reg.* (1840) *Misc.* 86.—Mexic.
clavata, *Lindl. l. c.* (1842) *Misc.* 61.—Honduras.
coriscensis, *Reichb f. Otia Bot. Hamb.* 112.—Afr. trop.
cultrata, *Lindl. Bot. Reg.* sub t. 851.—Ins. Maurit.
cultriformis, *Lindl. ex Spreng. Syst.* ii. 742.—Ins. Mascar.
dendrobiiflora, *Reichb. f. Otia Bot. Hamb.* 76.—Afr. trop.
dixantha, *Reichb. f. in Gard. Chron.* (1882) i. 294.—Afr. trop.
elastica, *Lindl. in Journ. Linn. Soc.* vi. (1862) 131.—Afr. trop.
elegans, *Reichb. f. Otia Bot. Hamb.* 113.—Afr. trop.
ensifolia, *Lindl. in Journ. Linn. Soc.* vi. (1862) 129.—Afr. trop.
estrellensis, *Reichb. f. in Linnaea*, xxv. (1852) 231.—Bras.
extinctoria, *Reichb. f. in Walp. Ann.* vi. 638.—Jamaic.
foliosa, *Reichb. f. l. c.* 640.—Ind. occ.
fusiformis, *Lindl. in Bot. Reg.* sub t. 851.—Ins. Mascar.
galeata, Reichb. f. in Walp. Ann. vi. 637 = grandiflora.
galericulata, *Reichb. f. Otia Bot. Hamb.* 111.—Afr. trop.
geraënsis, *Rodrig. Orch. Nov.* ii. 168.—Bras.
Gerrardi, *Harv. Thes. Cap.* ii. 49.—Afr. austr.
golungensis, *Reichb. f. in Flora*, xlviii. (1865) 185.—Afr. trop.
grandiflora, *Lindl. in Bot. Mag.* t. 3707.—Afr. trop.
hypocrita, *Reichb. f. in Gard. Chron.* (1881) ii. 685.—Afr. trop.
Jussieuana, *Reichb. f. in Walp. Ann.* vi. 641.—Madag.
laxiflora, *Lindl. in Journ. Linn. Soc.* vi. (1862) 129.—Afr. trop.
leonensis, *Reichb. f. Otia Bot. Hamb.* 112.—Afr. trop.
Lindleyana, Harv. Thes. Cap. ii. 50 = pubescens.
lineata, *Reichb. f. in Saund. Refug. Bot.* ii. tt. 80, 81.—Guatemala.
luteola, *Hook. Exot. Fl.* t. 103.—Am. trop.
luteola, Wight, Ic. t. 1678 = Wightii.
masasayensis, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Nicaragua.
mauritiana, *Spreng. Syst.* iii. 742.—Ins. Maurit.
membranacea, A. Rich. Fl. Cub. Fanerog. ii. 248 (= *Chloidia vernalis*).—Ind. occ.
minutiflora, Ridl. in Journ. Linn. Soc. xx. (1883) 330 = fusiformis.
modesta, *Reichb. f. in Flora*, l. (1867) 114.—Afr. trop.
multiflora, Ridl. in Journ. Linn. Soc. xxi. (1885) 475 err. typ. = *minutiflora*.
nana, Klotzsch, Ind. Sem. Hort. Berol. (1853) 12 = caracasana.
nana, *Reichb. f. in Walp. Ann.* vi. 638.—Peruv.
nitidula, *Reichb. f. in Flora*, l. (1867) 114.—Afr. trop.
odorata, *Lindl. in Journ. Linn. Soc.* vi. (1862) 130.—Afr. trop.
Ottoniana, *Reichb. f. in Hamb. Gartenz.* xi. (1855) 249.—Afr. austr.
pachyglossa, *Reichb. f. in Linnaea*, xli. (1877) 73.—Afr. trop.
paulensis, *Reichb. f. in Walp. Ann.* vi. 642.—Bras.
pinicola, *Rodrig. Orch. Nov.* ii. 169.—Bras.
puberula, *Lindl. in Bot. Reg.* sub t. 851.—Afr. trop.
pubescens, *Reichb. f. in Walp. Ann.* vi. 643.—Afr. austr.
purpurea, *Wight, Ic.* t. 1679.—Ind. or.
pyramidalis, *Lindl. in Journ. Linn. Soc.* vi. (1862) 130.—Afr. trop.
ramosa, Gardn. ex Thw. Enum. Pl. Zeyl. 307 = Josephia lanceolata.

POLYSTACHYA:—
ramulosa, *Lindl. Bot. Reg.* (1838) *Misc.* 76.—Afr. trop.
reflexa, *Lindl. l. c.* (1841) *Misc.* 18.—Afr. trop.
rhodoptera, *Reichb. f. in Hamb. Gartenz.* xiv. (1858) 214.—Afr. trop.
rigidula, *Reichb. f. in Flora*, l. (1867) 117.—Afr. austr.
rosea, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 474.—Madag.
rosellata, *Ridl. l. c.* xx. (1883) 330.—Madag.
rufinula, *Reichb. f. in Gard. Chron.* (1879) i. 41.—Afr. trop.
Sandersoni, *Harv. Thes. Cap.* ii. 49.—Afr. austr.
setifera, *Lindl. in Journ. Linn. Soc.* vi. (1862) 130.—Afr. trop.
shirensis, *Reichb. f. Otia Bot. Hamb.* 112.—Afr. trop.
similis, *Reichb. f. l. c.*—Afr. austr.
Steudneri, *Reichb. f. l. c.* 113.—Afr. trop.
sulfurea, *A. Br. in Fl. des Serres*, xv. (1862-63) 45.—Afr. trop.
superposita, *Reichb. f. Otia Bot. Hamb.* 111.—Afr. trop.
tessellata, *Lindl. in Journ. Linn. Soc.* vi. (1862) 130.—Afr. austr.
tricruris, *Reichb. f. in Flora*, l. (1867) 118.—Afr. austr.
virescens, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 474.—Madag.
Weigelti, Reichb. f. in Linnaea, xxv. (1852) 230 = foliosa.
Wightii, *Reichb. f. in Walp. Ann.* vi. 640.—Ind. or.
zeylanica, *Lindl. Bot. Reg.* (1838) *Misc.* 78.—Ins. Mascar. et Zeylan.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 39.—Java.

POLYSTEMMA, Decne. in DC. Prod. viii. 602 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 768.
viridiflora, *Decne. l. c.*—Mexic.

POLYSTEMON, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 95 = **Belangeria**, Cambess. (Saxifr.).
pentaphyllus, D. Don, l. c. = B. speciosa ?
triphyllus, D. Don, l. c. = B. tomentosa.

POLYSTIGMA, Meissn. Gen. 252; Comm. 161 (1839) = **Byronia**, Endl. (Ilicin.).
Hookeri, Meissn. l. c. Comm. 161 = B. sandwicensis.

POLYSTORTHIA, Blume, Fl. Jav. Praef. p. viii (1828) = **Pygeum**, Gaertn. (Rosac.).

POLYSTYLUS, Hasselt, ex Hassk. Retzia, 3 (1855) = **Phalaenopsis**, Blume (Orchid.).
Cornu-cervi, Hasselt, ex Hassk. l. c. = Ph. Cornu-cervi.

POLYTAENIA, DC. Coll. Mém. v. 53. t. 13 (1829); et Prod. iv. 196 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 922.
Pachiloma, Rafin. New Fl. Am. iv. 33 (1836).
Phaiosperma, Rafin. l. c. 32 (1836).
Nuttallii, *DC. Coll. Mém.* v. 53. t. 13; *et Prod.* iv. 196.—Am. bor.

POLYTAXIS, Bunge, Del. Sem. Hort. Dorpat. (1843) 8.—Cf. Linnaea, xviii. (1844) 156 = **Jurinea**, Cass. (Compos.).
Lehmanni, Bunge, ll. cc.—As. centr.

POLYTENIA, Rafin. New Fl. Am. iv. 33 (1836) = **Polytaenia**, DC. (Umbell.).

POLYTHECANDRA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 314 = **Clusia**, Linn. (Guttif.).
Schomburgkiana, Planch. & Triana, l. c. xiv. (1860) 229.—Guian.
Spruceana, Planch. & Triana, l. c.—Bras.

POLYTHRIX, Nees, in DC. Prod. xi. 285 (1847) = **Crossandra**, Salisb. (Acanth.).
stenandrium, Nees, l. c. 286 = C. madagascariensis.

POLYTHYSANIA, Hanst. in Linnaea, xxiv. (1853) 209 = **Alloplectus**, Mart. (Gesner.).
parviflora, Hanst. l. c. = A. parviflorus.

POLYTOCA, R. Br. in Benn. Pl. Jav. Rar. 20. t. 5 (1838). *GRAMINEAE*, Benth. & Hook. f. iii. 1112.
Cyathorhachis, Nees, ex Steud. Syn. Pl. Gram. 403 (1855).
bracteata, *R. Br. in Benn. Pl. Jav. Rar.* 20. t. 5.—Java.
macrophylla, *Benth. in Journ. Linn. Soc.* xix. (1881) 52.—Archip. Louisiade.
semiteres, *Benth. in Benth. & Hook. f. Gen.* iii. 1113.—Ind. or.
Wallichiana, *Benth. in Journ. Linn. Soc.* xix. (1881) 52.—Ind. or.

POLYTROPIS, Presl, Symb. Bot. 21. t. 13 (1830) = **Rhynchosia**, Lour. (Legumin.).
ferulaefolia, Presl, l. c. 22 = R. ferulaefolia.
pinnata, Eckl. & Zeyh. Enum. 246 = R. pinnata.
umbellata, Eckl. & Zeyh. l. c. = R. ferulaefolia.

POLYURA, Hook. f. in Hook. Ic. Pl. t. 1049 (1868). *RUBIACEAE*, Benth. & Hook. f. ii. 62.
geminata, *Hook. f. in Hook. Ic. Pl.* t. 1049.—Reg. Himal.

POLYXENA, Kunth, Enum. Pl. iv. 294 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 807.
Manlilia, Salisb. Gen. Pl. Fragm. 17 (1866).
Polyanthes, Jacq. Ic. Pl. Rar. t. 380 (1793); Coll. Suppl. 56 (1796).
pygmaea, *Kunth, l. c.*—Afr. austr.

POLYZONE, Endl. in Ann. Wien. Mus. ii. (1839) 191 = **Darwinia**, Rudge (Myrtac.).
purpurea, Endl. l. c. = D. purpurea.

POLYZYGUS, Dalz. in Hook. Kew Journ. ii. (1850) 260. *UMBELLIFERAE*, Benth. & Hook. f. i. 910.
tuberosus, *Walp. Ann.* ii. 715.—Ind. or.

POMADERRIS, Labill. Nov. Holl. Pl. i. 61. tt. 86, 87 (1804). *RHAMNEAE*, Benth. & Hook. f. i. 381.
Ledelia, Rafin. Sylva Tellur. 154 (1838).
Pomatiderris, Roem. & Schult. Syst. v. 622 (1819); H. B. & K. Nov. Gen. et Sp. vii. 60 (1825).
Pomatoderris, Roem. & Schult. l. c. p. xxxii. 414 (1819).
acuminata, Link, Enum. Hort. Berol. i. 235 = elliptica.
aemula, Steud. in Lehm. Pl. Preiss. i. 182 = Spyridium globulosum.
albicans, Steud. l. c. 184 = Trymalium albicans.
andromedaefolia, A. Cunn. in Field, N. S. Wales, 351 = phylliraeoides.
androsaemifolia, Heynh. Nom. i. 646, sphalm. = praec. ?
apetala, *Labill. Nov. Holl. Pl.* i. 52. t. 87.—Austral.
aspera, Sieber, ex DC. Prod. ii. 33 = apetala.
betulina, *A. Cunn. in Bot. Mag.* t. 3212.—Austral.
biaurita, F. Muell. Fragm. iii. 73 = obcordata.
Calvertiana, *F. Muell. Fragm.* ix. 138.—Austral.
capsularis, G. Don, ex Loud. Hort. Brit. 84 = Colubrina asiatica.
capsularis, *Montr. in Mém. Acad. Lyon*, x. (1860) 193.—N. Caled.
cinerea, *Benth. Fl. Austral.* i. 420.—Austral.
coactilifolia, F. Muell. ex Reiss. in Linnaea, xxix. (1857-58) 291 = Spyridium coactilifolium.
commixta, Steud. in Lehm. Pl. Preiss. i. 184 = Spyridium spadiceum.
discolor, DC. Prod. ii. 33 = elliptica.
Edgerleyi, *Hook. f. Handb. N. Zeal. Fl.* 43.—N. Zel.
elachophylla, *F. Muell. Fragm.* ii. 131.—Austral.
elliptica, *Labill. Nov. Holl. Pl.* i. 61. t. 86.—Austral.; N. Zel.
ericifolia, Hook. Journ. Bot. i. (1834) 257 = phylicaefolia.
ferruginea, *Sieber, ex DC. Prod.* ii. 33.—Austral.
Forrestiana, *F. Muell. Fragm.* ix. 139.—Austral.
globulosa, G. Don, ex Loud. Hort. Brit. 84 = Spyridium globulosum.
grandis, *F. Muell. Fragm.* iii. 68.—Austral.
hirsuta, Steud. in Lehm. Pl. Preiss. i. 184 = Spyridium spadiceum.
hirta, Reiss. in Endl. Nov. Stirp. Dec. 31 = ferruginea.
intangenda, *F. Muell. Fragm.* x. 52.—Austral.
intermedia, Sieber, ex DC. Prod. ii. 33 = elliptica.
Kumeraho, A. Cunn. ex Fenzl, in Enum. Pl. Hueg. 21; et in Ann. Nat. Hist. Ser. I. iii. (1839) 248 = elliptica.
lanigera, *Sims, Bot. Mag.* t. 1823.—Austral.

POMADERRIS:—
ledifolia, *A. Cunn. in Field, N. S. Wales*, 351.—Austral.
ligustrina, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 271 = ledifolia.
ligustrina, *Sieber, ex DC. Prod.* ii. 34.—Austral.
malifolia, Sieber, ex Benth. Fl. Austral. i. 418 = elliptica.
multiflora, Sieber, ex DC. Prod. ii. 33 = elliptica.
myrtilloides, *Fenzl, in Enum. Pl. Hueg.* 22.—Austral.
obcordata, *Fenzl, l. c.* 23.—Austral.
obovata, Hook. Comp. Bot. Mag. i. (1835) 277 = Spyridium obovatum.
obscura, Sieber, ex Fenzl, Enum. Pl. Hueg. 21 = lanigera.
ovaria, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 268 = racemosa.
ovata, Nois. ex Steud. Nom. ed. I. 646 = Ceanothus ovatus.
paniculosa, F. Muell. & Reiss. in Linnaea, xxix. (1857–58) 269 = racemosa.
parvifolia, Hook. Journ. Bot. i. (1834) 257 = Spyridium parvifolium.
phillyraefolia, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 266 = vacciniifolia.
phillyraefolia, Steud. in Lehm. Pl. Preiss. i. 183 = Spyridium globulosum.
philyraefolia, Fenzl, in Enum. Pl. Hueg. 22 = phylliraeoides.
phylicaefolia, *Lodd. Bot. Cab.* t. 120.—Austral.; N. Zel.
phylliraeoides, *Sieber, ex DC. Prod.* ii. 33.—Austral.
polifolia, Reiss. & F. Muell. in Linnaea, xxix. (1857–58) 269 = phylicaefolia.
polyantha, Steud. in Lehm. Pl. Preiss. i. 182 = Spyridium globulosum.
prostrata, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 284 = Spyridium eriocephalum.
prunifolia, *A. Cunn. ex Fenzl, in Enum. Pl. Hueg.* 22.—Austral.
pyrrhophylla, Steud. in Lehm. Pl. Preiss. i. 183 = Spyridium globulosum.
racemosa, *Hook. Journ. Bot.* i. (1834) 256.—Austral.
rosmarinifolia, Steud. in Lehm. Pl. Preiss. i. 184 = Trymalium ledifolium.
scabra, Hort. ex Steud. Nom. ed. II. ii. 379 = apetala.
Simsii, Sweet, Hort. Brit. ed. II. 114 = lanigera.
spatulata, G. Don, in Loud. Hort. Brit. 84 = Trymalium Billardieri.
stenopetala, F. Muell. Fragm. iii. 69 = myrtilloides.
subochreata, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 287 = Spyridium subochreatum.
subrepanda, *F. Muell. ex Reiss. l. c.* 267.—Austral.
subretusa, Steud. in Lehm. Pl. Preiss. i. 183 = Spyridium spadiceum.
Tainui, *Hector, in Trans. N. Z. Inst.* xi. (1879) 429.—N. Zel.
vacciniifolia, *Reiss. in Linnaea*, xxix. (1857–58) 266.—Austral.
viridirufa, Sieber, ex Fenzl, Enum. Pl. Hueg. 21 = ferruginea.
Wendlandiana, G. Don, in Loud. Hort. Brit. 84 = ferruginea.
westringiaefolia, Steud. in Lehm. Pl. Preiss. i. 185 = Spyridium westringiaefolium.
zizyphoides, Hook. & Arn. Bot. Beech. Voy. 61 = Alphitonia excelsa.

POMANGIUM, Reinw. in Syll. Ratisb. ii. (1828) 10 = **Argostemma**, Wall. (Rubiac.).
montanum, Blume, ex DC. Prod. iv. 418 = A. montanum.

POMARIA, Cav. Ic. v. 1. t. 402 (1799) = **Caesalpinia**, Linn. (Legumin.).
glandulosa, Cav. l. c. 2. t. 402 = C. bijuga.
glauca, Cav. Desc. 466.—Mexic.?
humilis, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 303 = Hoffmanseggia humilis.
Jamesii, Walp. Rep. i. 811 = C. bijuga.
melanosticta, Schau. in Linnaea, xx. (1847) 748 = Hoffmanseggia melanosticta.

POMASTERION, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 80 (1865–66) = **Actinostemma**, Griff. (Cucurb.).
japonicum, Miq. l. c. = A. racemosum.

POMATIDERRIS, Roem. & Schult. Syst. v. 622 (1819); H. B. & K. Nov. Gen. et Sp. vii. 60 (1825) = **Pomaderris**, Labill. (Rhamn.).

POMATIUM, Gaertn. f. Fruct. iii. 252. t. 225 (1805) = **Bertiera**, Aubl. (Rubiac.).
dubium, G. Don, Gen. Syst. iii. 505.—Afr. trop.
spicatum, Gaertn. f. Fruct. iii. 252 = B. Pomatium.

POMATIUM, Nees, ex Meissn. in Mart. Fl. Bras. v. II. 252 (1866) = **Nectandra**, Roland. (Laurin.).
canescens, Nees, l. c. = N. canescens.

POMATOCALPA, Breda, Kuhl & Hass. Orch. t. [15] (1827) = **Cleisostoma**, Blume (Orchid.).
spicatum, Breda, l. c. = C. callosum.

POMATODERRIS, Roem. & Schult. Syst. v. p. xxxii. 414 (1819) = **Pomaderris**, Labill. (Rhamn.).

POMATOSACE, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 499. *PRIMULACEAE.*
Filicula, *Maxim. l. c.* 500.—China.

POMATOTHECA, F. Muell. Fragm. x. 72 (1876). *FICOIDEAE.*
humillima, *F. Muell. l. c.*—Austral.

POMAX, Soland. ex Gaertn. Fruct. i. 112. t. 24 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 141.
glabra, DC. Prod. iv. 615 = umbellata.
hirta, DC. l. c. = umbellata.
rupestris, F. Muell. in Linnaea, xxv. (1852) 395 = umbellata.
umbellata, *Soland. ex Gaertn. Fruct.* i. 112.—Austral.

POMBALIA, Vand. Fasc. Pl. 7 (1771) = **Ionidium**, Vent. (Violar.).
Ipecacuanha, Vand. l. c. = I. Ipecacuanha.
Itouba, Ging. ex DC. Prod. i. 307 = I. Ipecacuanha.

POMBEA, Mutis, ex Caldas, in Seman. Nuev. Gran. ii. 27 (1810), nomen.

POMELIA, Durando, ex Pomel, Mat. Fl. Atl. 7 (1860) = **Daucus**, Tourn. (Umbell.).
setifolia, Durando, ex Pomel, l. c. 8 = D. setifolius.

POMEREULA, Domb. ex DC. Prod. iii. 190 (1828) = **Miconia**, Ruiz & Pav. (Melastom.).
tomentosa, Domb. ex DC. l. c. = M. pulverulenta.

POMETIA, Forst. Char. Gen. 109. t. 55 (1776), partim; Forst. f. Prod. 74 (1786), partim. *SAPINDACEAE*, Benth. & Hook. f. i. 407.
CNESMOCARPUS, Zipp. ex Blume, Rumphia, iii. 115 (1847).
DIPLOCARDIA, Zipp. ex Blume, l. c. (1847).
ECCREMANTHUS, Thw. in Hook. Kew Journ. vii. (1856) 272. t. 9.
IRINA, Blume, Bijdr. 230 (1825).
PROSTEA, Cambess. in Mém. Mus. Par. xviii. (1829) 25. t. 1 C.
eximia, *Bedd. Fl. Sylvat.* t. 157.—Ind. or.; Malaya.
macrocarpa, *Kurz, in Journ. As. Soc. Beng.* xliv. II. (1875) 205.—Malacca.
pinnata, *Forst. Char. Gen.* 110; *Forst. f. Prod.* 74.—Ins. Pacif.
ternata, Forst. f. Prod. 74 = Allophylus Cobbe.
ternata, Willd. Sp. Pl. iii. 398 = Schmidelia Cobbe.
tomentosa, Kurz, Andam. Rep. ed. II. 34 = eximia.

POMETIA, Vell. Fl. Flum. 80 (1825); ii. t. 87 (1827) = **Lucuma**, Molina (Sapot.).
lactescens, Vell. l. c. 81; ii. t. 87 = L. glycyphloea.
lutescens, Steud. Nom. ed. II. ii. 379, sphalm. = praec.

POMMEREULLA, Linn. f. Nov. Gram. 31 (1779). *GRAMINEAE*, Benth. & Hook. f. iii. 1174.
GRACILEA, Koen. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218.
ROYLEA, Nees, ex Steud. Nom. ed. II. ii. 475 (1841).
Cornucopiae, *Linn. f. Nov. Gram.* 31.—Ind. or.
elongata, Wight & Arn. ex Steud. Nom. ed. II. ii. 379 = P. Cornucopiae?
monoeca, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218 = Melanocenchris Rothiana.
Royleana, Steud. Nom. ed. II. ii. 379 = Melanocenchris Royleana.

POMPADOURA, Buchoz, ex DC. Prod. iii. 2 (1828) = **Calycanthus**, Linn.

POMPHIDEA, Miers, Apocyn. S. Am. 18 (1878). *APOCYNEAE.*
Swartziana, *Miers, l. c.* 19.—Ind. occ.

POMPILA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Sterculia**, Linn.

PONAEA, Schreb. Gen. i. 266 (1789) = **Toulicia**, Aubl. (Sapind.).
guianensis, J. F. Gmel. Syst. 640 = T. guianensis.
saponarioides, Willd. Sp. Pl. ii. 470 = T. guianensis.

PONARIA, Rafin. Med. Fl. ii. 109 (1830) = **Veronica**, Tourn. (Scrophul.).
laciniata, Rafin. l. c. = V. spuria *vel* austriaca.
latifolia, Rafin. l. c. = V. latifolia.
Pona, Rafin. l. c. = V. Ponae.

PONCELETIA, R. Br. Prod. 554 (1810) = **Sprengelia**, Sm. (Epacrid.).
monticola, A. Cunn. ex DC. Prod. vii. 768 = S. ponceletioides.
sprengelioides, R. Br. Prod. 554 = S. Ponceletia.

PONCELETIA, Thou. Fl. Trist. Acugn. 36 (1811) = **Spartina**, Schreb. (Gramin.).
arundinacea, Thou. l. c. = S. arundinacea.

PONCIRUS, Rafin. Sylva Tellur. 143 (1838) = **Citrus**, Tourn. (Rutac.).
trifoliatus, Rafin. l. c. = C. trifoliata.

PONERA, Lindl. Gen. et Sp. Orch. 113 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 527.
NEMACONIA, Knowles & Westc. Flor. Cab. ii. (1838) 127.
TETRAGAMESTUS, Reichb. f. in Bonplandia, ii. (1854) 21.
Adendrobium, Reichb. f. in Flora, xlviii. (1865) 278 = Pleuranthium Adendrobium.
albida, *Reichb. f. Beitr. Orch. Centr. Am.* 103.—Costa Rica.
amethystina, *Reichb. f. in Saund. Refug. Bot.* ii. t. 93.—Guatemala.
Behrii, Reichb. f. in Bonplandia, iii. (1855) 220 = Scaphyglottis Behrii.
bilineata, *Reichb. f. Beitr. Orch. Centr. Am.* 88.—Costa Rica.
conferta, Reichb. f. in Bonplandia, ii. (1854) 22 = Scaphyglottis conferta.
dubia, Reichb. f. l. c. iv. (1856) 327 = Scaphyglottis dubia.
esuriens, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 98.—N. Granat.
Felskyi, *Reichb. f. in Linnaea*, xli. (1877) 85.—Guian.
graminifolia, *Lindl. Bot. Reg.* (1839) *Misc.* 17.—Mexic.
inconspicua, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 616 (Quid?).—Guatemala.
juncifolia, *Lindl. Gen. et Sp. Orch.* 114.—Mexic.
Kienastii, Reichb. f. in Gard. Chron. (1877) I. 810 = Scaphyglottis Kienastii.
leucantha, Reichb. f. in Bonplandia, ii. (1854) 22 = Scaphyglottis leucantha.
macroglossa, *Reichb. f. in Bot. Zeit.* x. (1852) 639.—Guatemala.
mesocopis, Endres & Reichb. f. Xenia Orch. ii. 222. t. 200 = Scaphyglottis mesocopis.
modesta, *Reichb. f. in Linnaea*, xli. (1877) 85, in obs.—Am. trop.
pellita, *Reichb. f. in Gard. Chron.* (1880) II. 8.—Hab.?
pleurostachys, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 282.—N. Granat.
prolifera, *Reichb. f. l. c.* 22.—Ind. occ.
punctulata, *Reichb. f. l. c.* iii. (1855) 220.—Venezuela.
rosea, Reichb. f. l. c. ii. (1854) 22 = Scaphyglottis violacea.
stellata, Reichb. f. in Walp. Ann. vi. 454 = Scaphyglottis stellata.
striata, *Lindl. Bot. Reg.* (1842) *Misc.* 18.—Guatemala.
striolata, *Reichb. f. in Linnaea*, xli. (1877) 39, in obs.—Mexic.
violacea, *Reichb. f. in Bonplandia*, ii. (1854) 22.—Hab.?

PONERORCHIS, Reichb. f. in Linnaea, xxv. (1852) 227 = **Habenaria**, Willd. (Orchid.).
graminifolia, Reichb. f. l. c. 228.—Japon.

PONGAM, Adans. Fam. ii. 322 (1763) = seq.

PONGAMIA, Vent. Jard. Malm. 28. t. 28 (1803). *LEGUMINOSAE*, Benth. & Hook. f. i. 549.
GALEDUPA, Lam. Encyc. ii. 594 (1786).
acuminata, R. Grah. in Wall. Cat. n. 5901 = Derris acuminata.
amoena, Wall. l. c. n. 5912 = Derris amoena.
angustifolia, *Buch.-Ham. ex Wall. l. c.* n. 5917.—Nepal.
atropurpurea, Wall. l. c. n. 5910 = Millettia atropurpurea.
caerulea, R. Grah. in Wall. l. c. n. 5894 = Millettia caerulea.
cana, R. Grah. l. c. n. 5903 = Millettia cana.
canarensis, Dalz. in Hook. Kew Journ. ii. (1850) 37 = Derris canarensis.
cassioides, Wall. Cat. n. 5918 = Millettia pulchra.
caudata, R. Grah. in Wall. l. c. n. 5895 = Millettia caudata.
chinensis, *DC. Prod.* ii. 416.—China.
cinerea, R. Grah. in Wall. Cat. n. 5888 = Millettia cinerea.
Corcor, R. Grah. l. c. n. 5891 = Millettia racemosa.
coriacea, R. Grah. l. c. n. 5905 = Derris scandens.
Corollaria, *Miq. Fl. Ind. Bat.* i. 1. 149.—Ins. Amboin.
crassifolia, R. Grah. in Wall. Cat. n. 5913 = Spatholobus crassifolius.
cuneifolia, R. Grah. l. c. n. 5887 = Derris cuneifolia.
dubia, R. Grah. l. c. n. 5899 = Derris elliptica.
elegans, R. Grah. l. c. n. 5883 = Derris elegans.
elliptica, Sweet, Hort. Brit. ed. I. 131 = Derris elliptica.
elongata, R. Grah. in Wall. Cat. n. 5886 = Derris acuminata, oblonga.
emarginata, Wall. l. c. n. 5909 = Derris marginata.
extensa, Wall. l. c. n. 5900 = Millettia extensa.
ferruginea, R. Grah. in Wall. l. c. n. 5885 = Derris ferruginea.
floribunda, R. Grah. l. c. n. 5884 = Derris elegans.
fruticosa, R. Grah. l. c. n. 909 = Millettia fruticosa.
Galedupa, Herb. Madr. ex Wall. l. c. n. 5878 = glabra.
glabra, *Vent. Jard. Malm.* 28. t. 28.—As. trop.; Austral.; Ins. Pacif.
glandulosa, Griff. Notul. iv. 443 (an Derris?).—Ind. or.
grandiflora, *Vent. Jard. Malm.* sub t. 28.—Ind. or.
grandifolia, R. Grah. in Wall. Cat. n. 5882 = Derris sinuata.
grandifolia, Zoll. & Mor. Syst. Verz. Zoll. 2 = glabra.
heterocarpa, Wall. ex Voigt, Hort. Suburb. Calc. 240, nomen; Baker, in Hook. f. Fl. Brit. Ind. ii. 106 = Millettia cinerea.
Heyneana, R. Grah. in Wall. Cat. n. 5916 = Derris Heyneana.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 149 = Derris elliptica.
hypoleuca, Miq. l. c. 148 = Derris elliptica.
lucida, R. Grah. in Wall. Cat. n. 5880 (Quid?).—Ind. or.
macrophylla, R. Grah. l. c. n. 5892 = Millettia auriculata.
madagascariensis, Boj. ex Oliver, Fl. Trop. Afr. ii. 246 = Derris uliginosa.
marginata, R. Grah. in Wall. Cat. n. 5896 = Derris cuneifolia.
maritima, Wight, ex Wall. l. c. n. 5879 = Derris uliginosa.
mitis, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 128.—Ins. Nicobar.
monadelpha, R. Grah. in Wall. Cat. n. 5906 = Derris cuneifolia.
oblonga, R. Grah. l. c. n. 5893 = Millettia cinerea.
obovata, R. Grah. l. c. n. 5897 = Derris cuneifolia.
ovalifolia, Wight & Arn. Prod. 262 = Derris oblonga.
palustris, R. Grah. in Wall. Cat. n. 5889 = Millettia cinerea.
paniculata, R. Grah. l. c. n. 5904 = Millettia cinerea.
paniculata, Wight, ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 110 = Derris Heyneana.
pendula, R. Grah. in Wall. Cat. n. 5902 = Millettia pendula.
PONGAMIA:—
piscatoria, Seem. in Bonplandia, ix. (1861) 255 = Derris uliginosa.
Piscidia, *Steud. Nom.* ed. II. ii. 380.—Ind. or.
racemosa, R. Grah. in Wall. Cat. n. 5914 = Millettia racemosa.
reflexa, R. Grah. l. c. n. 5898 = Derris marginata.
religiosa, Wight, in Hook. Bot. Misc. iii. (1833) 301 = Derris uliginosa.
rigida, R. Grah. in Wall. Cat. n. 5908 = Spatholobus acuminatus.
rosea, R. Grah. l. c. n. 5907 = Spatholobus acuminatus.
rostrata, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, iii. (1846) 65, nomen.—Java.
secunda, R. Grah. in Wall. Cat. n. 5890 = Derris secunda.
sericea, Vent. Jard. Malm. sub t. 28 = Millettia sericea.
sinuata, Wall. Cat. n. 5911 = Derris sinuata.
tetraptera, Hort. Calc. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 106 = Millettia tetraptera.
triphylla, Wight, in Hook. Bot. Misc. iii. (1833) t. 41 = Derris uliginosa.
uliginosa, DC. Prod. ii. 416 = Derris uliginosa.
volubilis, Zoll. & Mor. Syst. Verz. Zoll. 3 = Derris elliptica.

PONGATIUM, Juss. Gen. 423 (1789) = **Sphenoclea**, Gaertn. (Campan.).
indicum, Lam. Illustr. ii. 44 = S. zeylanica.
spongiosum, Blanco, Fl. Filip. ed. I. 86 = S. zeylanica.

PONGELIA, Rafin. Sylva Tellur. 78 (1838) = **Bignonia**, Linn.
longiflora, Rafin. l. c. 79 = B. longiflora.

PONGELION, Adans. Fam. ii. 319 (1763) = **Adenanthera**, Linn. (Legumin.).

PONISTA, Rafin. Fl. Tellur. ii. 65 (1836) = **Saxifraga**, Tourn.
oregonensis, Rafin. l. c. 66 = S. petraea.
petrea, Rafin. l. c. = S. petraea.

PONTALETSJE, Adans. Fam. ii. 159 (1763) = **Lawsonia**, Linn. (Lythrar.).

PONTEDERAS, Hoffmgg. Verz. Pfl. 137 (1824) = seq.

PONTEDERIA, Linn. Syst. ed. I (1735). *PONTEDERIACEAE*, Benth. & Hook. f. iii. 837.
CARIGOLA, Rafin. Fl. Tellur. ii. 10 (1836).
KADAKIA, Rafin. l. c. 9 (1836).
NARUKILA, Adans. Fam. ii. 54 (1763).
UMSEMA, Rafin. in Med. Repos. N. York, v. (1808) 352.
alba, Buch.-Ham. ex Wall. Cat. n. 5095 D = Monochoria vaginalis.
angustifolia, Pursh, Fl. Am. Sept. i. 224 = cordata.
aquatica, Vell. Fl. Flum. 144; iii. t. 164 = Eichhornia azurea.
azurea, Herb. Cub. ex Schult. f. Syst. vii. 1148 = Heteranthera reniformis?
azurea, Hook. Bot. Mag. t. 2932 = Eichhornia speciosa.
azurea, Poepp. ex Kunth, Enum. Pl. iv. 122 = Heteranthera spicata.
azurea, Sw. Prod. Veg. Ind. Occ. 57 = Eichhornia azurea.
brasiliensis, *Willd. ex Schult. f. Syst.* vii. 1145.—Bras.
coerulea, *Maund, ex Steud. Nom.* ed. II. ii. 380 (Quid?).—Am. bor.
cordata, *Linn. Sp. Pl.* 288.—Am. calid.
cordata, Lour. Fl. Cochinch. 245 = Monochoria vaginalis.
cordifolia, Mart. in Schult. f. Syst. vii. 1142 = rotundifolia.
crassicaulis, Schlecht. in Abh. Nat. Ges. Halle, vi. (1862) 177, sphalm. = seq.
crassipes, Mart. Nov. Gen. et Sp. i. 9. t. 4 = Eichhornia speciosa.
dilatata, [Buch.-Ham. in] Symes, Embassy. ed. II. iii. 309 = Monochoria hastaefolia.
dubia, Blume, Enum. Pl. Jav. i. 33 = Monochoria vaginalis.
elongata, Balf. in Proc. Bot. Soc. Edinb. (1855) 50 = Eichhornia speciosa.
PONTEDERIA:—
eriantha, Miq. in Linnaea, xvii. (1843) 60 = rotundifolia.
falciloba, Solms, in DC. Monog. Phan. iv. 534 = rotundifolia.
hastata, Linn. Sp. Pl. 288 = Monochoria hastaefolia.
lagoënsis, Warm. in Kjoeb. Vidensk. Meddel. (1871) 323 = Reussia subovata.
lanceolata, Nutt. Gen. Am. i. 216 = cordata.
lanceolata, Wall. ex Kunth, Enum. Pl. iv. 135 = Monochoria vaginalis.
lancifolia, Muhl. ex Ell. Sketch, i. 382 = cordata.
limosa, Sw. Prod. Veg. Ind. Occ. 57 = Heteranthera limosa.
linearis, Hassk. in Flora, xxv. (1842) II. Beibl. 4 = Monochoria vaginalis.
Loureiriana, Schult. f. Syst. vii. 1145 = Monochoria vaginalis.
maculata, Donn, Hort. Cantab. ed. VI. 81 (Quid?).—Am. bor.
Martiana, Seub. in Mart. Fl. Bras. iii. 1. 92, sphalm. = seq.
Martiusiana, Schult. f. Syst. vii. 1143 = Eichhornia Martiana.
mucronata, Rafin. ex Torr. Fl. U. St. i. 343 = cordata.
natans, Beauv. Fl. Owar. ii. 18. t. 68. f. 2 = Eichhornia natans.
nymphaefolia, *Kunth, Enum. Pl.* iv. 126.—Bras.
obtusifolia, A. DC. Monog. Phan. iv. 567 (Index) = cordata.
ovalis, *Mart. ex Schult. f. Syst.* vii. 1140.—Bras.
ovata, *Hook. & Arn. Bot. Beech. Voy.* 218.—China.
ovata, Linn. Sp. Pl. 288 = Phrynium capitatum.
paniculata, Spreng. Neue Entdeck. iii. 18 = Eichhornia martiana.
paradoxa, Mart. in Schult. f. Syst. vii. 1144 = Eichhornia Schultesiana.
pauciflora, Blume, Enum. Pl. Jav. i. 32 = Monochoria vaginalis.
plantaginea, Roxb. Hort. Beng. [23]; Fl. Ind. ii. 123 = Monochoria vaginalis.
racemosa, Buch.-Ham. ex Wall. Cat. n. 5095 C = Monochoria vaginalis.
rotundifolia, *Linn. f. Suppl.* 192.—Guian.
sagittata, Presl, Rel. Haenk. ii. 116 = cordata.
sagittata, Roxb. Hort. Beng. [23]; Fl. Ind. ii. 124 = Monochoria hastaefolia.
sagittifolia, Heyne, ex Wall. Cat. n. 5093 = Monochoria hastaefolia.
Schomburgkiana, Klotzsch, in Schomb. Faun. & Fl. Brit. Gui. 1118 = Reussia subovata.
stricta, Burm. f. Fl. Cap. Prod. 9 (an Aponogetonis sp.?).—Afr. austr.
triandra, Banks, ex Schult. f. Syst. vii. 1148, in obs. = Heteranthera limosa.
tricolor, Mart. ex Seub. in Mart. Fl. Bras. iii. 1. 92 = Eichhornia Martiana.
tumida, Willd. ex Kunth, Enum. Pl. iv. 129 = Eichhornia azurea.
vaginalis, Blanco, Fl. Filip. ed. I. 255 = Monochoria hastaefolia.
vaginalis, Burm. f. Fl. Ind. 80 = Monochoria vaginalis.

PONTESIA, Vell. Fl. Flum. viii. t. 147 (1827) = **Riencourtia**, Cass. (Compos.).
errata, Steud. Nom. ed. II. ii. 380, sphalm. = seq.
serrata, Vell. Fl. Flum. viii. t. 147 = R. latifolia?

PONTHIEVA, R. Br. in Ait. Hort. Kew. ed. II. v. 199 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 593.
NERISSA, Rafin. Fl. Tellur. ii. 89 (1836).
SCHOENLEINIA, Klotzsch, ex Lindl. Veg. Kingd. 182 (1847).
andicola, *Reichb. f. in Linnaea*, xli. (1877) 52.—Ecuador.
cornuta, *Reichb. f. l. c.* 18.—Bolivia.
diptera, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.
Ephippium, *Reichb. f. in Linnaea*, xxviii. (1856) 382.—Mexic.
glandulosa, *R. Br. in Ait. Hort. Kew.* ed. II. v. 200.—Ind. occ.
guatemalensis, *Reichb. f. Beitr. Orch. Centr. Am.* 63.—Guatemala.
inaudita, *Reichb. f. in Linnaea*, xli. (1877) 18.—Peruv.

PONTHIEVA:—
lancifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 251.—Cuba.
maculata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 385.—N. Granat.
Mandonii, *Reichb. f. Xenia Orch.* iii. 18.—Bolivia.
montana, *Lindl. in Benth. Pl. Hartw.* 155.—Peruv.
multiflora, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 16. t. 123.—Chili; Peruv.
oblongifolia, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic.
petiolata, *Lindl. Bot. Reg.* t. 760.—Ind. occ.
rostrata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 385.—Mexic.
sylvicola, *Reichb. f. in Linnaea*, xli. (1877) 52.—Ecuador.
villosa, *Lindl. in Benth. Pl. Hartw.* 155.—N. Granat.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 101.—N. Granat.

PONTIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 318 = **Chrysanthemum**, Tourn. (Compos.).
Bauhini, Bubani, l. c. = C. coronopifolium, *Vill.*

PONTINIA, Fries, in Lindbl. Bot. Notiser (1843), 141 = **Lychnis**, Tourn. (Caryoph.).
Coeli-rosa, Fries, l. c. = L. Coeli-rosa.

PONTOPIDANA, Scop. Introd. 195 (1777) = **Couroupita**, Aubl. (Myrtac.).

PONTOPPIDANA, Steud. Nom. ed. II. ii. 381 (1841) = praec.

POOTIA, Dennst. Schluess. Hort. Malab. 37 (1818) = **Canscora**, Lam. (Gentian.).
triflora, Dennst. l. c. = C. perfoliata.

POOTIA, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 192; Fl. Ind. Bat. ii. 417 ('1856') = **Voacanga**, Thou. (Apocyn.).
cereopsifolia, Miq. ex Benth. in Journ. Linn. Soc. iii. (1858) 74 = Carallia integerrima.
exauriculata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 404.—Ins. Molucc.
grandifolia, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 193 = V. grandifolia.

POPONAX, Rafin. Sylva Tellur. 118 (1838) = **Acacia**, Willd. (Legumin.).
Farnesiana, Rafin. l. c. = A. Farnesiana.
tortuosa, Rafin. l. c. = A. tortuosa.

POPOWIA, Endl. Gen. 831 (1839). *ANONACEAE*, Benth. & Hook. f. i. 25.
affinis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 20 = ramosissima.
australis, *Benth. Fl. Austral.* i. 52.—Austral.
bancana, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 17.—N. Guin.
Barteri, *Baill. Adansonia*, viii. (1867–68) 324.—Afr. trop.
Beccarii, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 16.—N. Guin.
Beddomeana, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 68. Ind. or.
Boivini, *Baill. Adansonia*, viii. (1867-68) 317.—Madag.
caffra, *Hook. f. & Thoms. ex Benth. in Trans. Linn. Soc.* xxiii. (1862) 470.—Afr. austr.
cuspidata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 21.—Java.
foetida, *Maing. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 69.—Malacca.
fornicata, *Baill. Adansonia*, viii. (1867–68) 318.—Afr. trop.
glaucocarpa, *Baill. l. c.* 319.—Madag.
Helferi, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 69.—Burma.
Heudeloti, *Baill. l. c.* 320.—Afr. trop.
hirta, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 21.—Sumatra.
Kirkii, Benth. in Trans. Linn. Soc. xxiii. (1862) 470 = Cleistochlamys Kirkii.
macrocarpa, *Baill. Adansonia*, viii. (1867–68) 316.—Madag.
Mannii, *Baill. l. c.* 320.—Afr. trop.
nervifolia, *Maing. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 69.—Malacca.

POPOWIA:—
novo-guineensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 21.—N. Guin.
papuana, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 15.—N. Guin.
parvifolia, *Kurz, in Journ. Bot.* xiii. (1875) 324.—Ins. Nicobar.
parvifolia, *Scheff. in Ann. Jard. Buitenz*, ii. (1885) 16.—N. Guin.
pauciflora, *Maing. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 69.—Malacca.
Pervillei, *Baill. Adansonia*, viii. (1867–68) 317.—Madag.
pilosa, *Baill. l. c.* 322.—Madag.
pisocarpa, *Endl. in Walp. Rep.* i. 74.—Java.
ramosissima, Bedd. Ic. Pl. Ind. Or. t. 75 = Beddomeana.
ramosissima, *Hook. f. & Thoms. Fl. Ind.* i. 105.—Ind. or.
rufula, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 20 = ramosissima.
tomentosa, *Maing. ex Hook. f. & Thoms. Fl. Brit. Ind.* i. 70.—Malacca.
Vogelii, *Baill. Adansonia*, viii. (1867-68) 324; *Hist. Pl.* i. 221.—Afr. trop.

POPPIA, Hort. ex Carr. in Vilm. Fl. Pl. Terre, ed. III. 649 (1870) = POPPYA, Neck. = **Luffa**, Tourn. (Cucurb.).

POPPIGIA, Hook. & Arn. Bot. Beech. Voy. 58 (1832) = POEPPIGIA, Bert. = **Rhaphithamnus**, Miers (Verben.).

POPPYA, Neck. Elem. i. 241 (1790) = **Luffa**, Tourn. (Cucurb.).
Fabiana, C. Koch, in Allg. Gartenz. (1857) 60 = L. aegyptiaca.
operculata, M. Roem. Syn. Pepon. 59 = L. purgans.

POPULAGO, Tourn. ex Hall. Enum. Stirp. Helv. i. 319 (1742) = **Caltha**, Linn. (Ranunc.).
palustris, Scop. Fl. Carn. ed. II. i. 404 = C. palustris.

POPULUS, Linn. Syst. ed. I (1735); Gen. ed. I. 307 (1737). *SALICINEAE*, Benth. & Hook. f. iii. 412.
AIGIROS, Rafin. Alsog. Am. 42 (1838).
LEUCE, Opiz, Seznam, 59 (1852).
MONILISTUS, Rafin. Alsog. Am. 42 (1838).
OCTIMA, Rafin. l. c. (1838).
TACAMAHACA, Mill. Gard. Dict. ed. VI. 175. App. (1752).
TREMULA, Dum. in Hall, Bijdr. Nat. Wet. i. (1826) 46.
acerifolia, Lodd. ex Loud. Arboret. iii. 1640 = alba.
Acladesca, Hort. ex C. Koch, Wochenschr. viii. (1865) 238 = balsamifera.
Acladesca, Lindl. in Loud. Encyc. Pl. 840 = monilifera.
adenopoda, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 50.—China.
aegyptiaca, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 660 = alba.
alba, *Linn. Sp. Pl.* 1034.—Europ.; As. bor.
angulata, Ait. Hort. Kew. ed. I. iii. 407 = monilifera.
angulosa, Michx. Fl. Bor. Am. ii. 243 = praec.
angustifolia, *James, in Long, Exped.* i. 497; *ex Torr. in Ann. Lyc. N. York*, ii. (1828) 249.—Am. bor.
angustifolia, Newberry, in Pacif. Rail. Rep. vi. Bot. 89 = trichocarpa.
angustifolia, Weinm. Elench. Hort. Pawlowsk. 451, nomen; et ex Ledeb. Fl. Ross. iii. 629 = balsamifera.
arembergica, Lodd. ex Loud. Arboret. iii. 1640 = alba.
argentea, *Fouger. in Mém. Agric. Par.* (Print. 1786) 102.—Hab.?
argentea, Michx. f. Hist. Arb. Am. iii. 290. t. 9 = heterophylla.
Athemsay, Hort. ex Fouger. in Mém. Agric. Par. (Print. 1786) 96 = seq.
Athemse, Hort. ex Fouger. l. c. = heterophylla.
atheniensis, Lodd. ex C. F. Ludwig, Neue. Wilde Baumz. (1783) 35 = tremuloides.

POPULUS:—
australis, Tenore, Ind. Sem. Hort. Neap. (1830) 15; Fl. Nap. iv. Syll. 138 = tremula.
Bachofenii, Wierzb. ex Reichb. Ic. Fl. Germ. xi. 29. t. 616 = alba.
balsamifera, *Linn. Sp. Pl.* 1034.—As. et Am. temp.
balsamifera, Lyall, in Journ. Linn. Soc. vii. (1864) 134 = trichocarpa.
balsamifera, Mill. Gard. Dict. ed. VIII. n. 5 = monilifera, heterophylla.
belgica, Lodd. ex Loud. Arboret. iii. 1640 = alba.
benzoifera, Tausch, in Flora, xxi. (1838) 753 = tremuloides.
berolinensis ×, C. Koch, Wochenschr. viii. (1865) 239.
betulifolia, Pursh, Fl. Am. Sept. ii. 619 = nigra.
bicolor, Rafin. Alsog. Am. 42 (Quid?).—Am. bor.
biformis, Rafin. l. c. 43 (Quid?).—Am. bor.
Bolleana, Mast. in Gard. Chron. (1882) II. f. 96 = alba.
canadensis, Fouger. in Mém. Agric. Par. (Print. 1786) 94 = nigra.
canadensis, Michx. f. Hist. Arb. Am. iii. 302. f. 12 = monilifera.
canadensis, Moench, Verz. 81 = balsamifera.
candicans, Ait. Hort. Kew. ed. I. iii. 406 = balsamifera.
candicans, Lodd. ex Loud. Arboret. iii. 1640 = alba.
canescens, Sm. Fl. Brit. iii. 1080 = alba.
carolinensis, Fouger. in Mém. Agric. Par. (Print. 1786) 90 = heterophylla.
carolinensis, Moench, Verz. 81 = monilifera.
caudina, Tenore, Fl. Nap. v. 280 = nigra.
ciliata, Schur, Enum. Pl. Transs. 625 = monilifera.
ciliata, *Wall. Cat.* n. 2796.—Reg. Himal.
cilicica, Kotschy, ex Wesmael, in DC. Prod. xvii. II. 325 = tremula.
ciliolata, Rafin. Alsog. Am. 43 (Quid?).—Am. bor.
cinerea, Rafin. l. c. 42 (Quid?).—Am. bor.
cordata, Hort. ex Poir. Encyc. v. 236 = tremuloides.
cordifolia, Burgsdorf, Anleit. Erzieh. Holzart. ed. I. 177 = heterophylla.
croatica, Waldst. & Kit. ex Bess. in Flora, xv. (1832), II. Beibl. I. 14 = nigra.
deglubata, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 83 = Eucalyptus deglupta.
deltoides, Marsh. Arbust. 106 = monilifera.
deltoides, Marsh. l. c. 194 = grandidentata.
dilatata, Ait. Hort. Kew. ed. I. iii. 406 = pyramidalis.
diversifolia, Schrenk, Enum. Pl. Nov. 15 = euphratica.
Eugenei, Hort. ex C. Koch, Dendrol. ii. I. 493 = monilifera.
euphratensis, [Lindl.?] in Gard. Chron. (1849) 806 = seq.
euphratica, *Oliv. Voy. Atlas*, iii. 449. f. 45–46.—Oriens; As. centr.
excelsa, Salisb. Prod. 395 = alba.
fastigiata, Fouger. in Mém. Agric. Par. (Print. 1786) 82; Desf. Tabl. ed. I. 213 = nigra.
fastigiata, Poir. Encyc. v. 235 = pyramidalis.
flexibilis, Rozier, Cours Compl. Agric. vii. 618; Poir. Encyc. v. 234 = nigra.
flexilis, C. Koch, Dendrol. ii. I. 491 = praec.
Fremonti, *S. Wats. in Proc. Am. Acad.* x. (1875) 350.—Am. bor. occ.
glandulosa, Moench, Meth. 339 = monilifera.
glandulosa, Willd. ex Steud. Nom. ed. II. ii. 381 = tremuloides.
graeca, Ait. Hort. Kew. ed. I. iii. 407 = tremuloides.
grandidentata, *Michx. Fl. Bor. Am.* ii. 243.—Am. bor.
grisea, Lodd. ex Loud. Arboret. iii. 1640 = alba.
heterophilla, Fouger. in Mém. Agric. Par. (Print. 1786) 96 = heterophylla.
heterophylla, Duroi, Harbk. ii. 150 = monilifera.
heterophylla, Hort. ex C. Koch, Wochenschr. viii. (1865) 238 = balsamifera.
heterophylla, *Linn. Sp. Pl.* 1034.—Am. bor.
hudsoniana, Desf. Tabl. ed. II. 243 = seq.
hudsonica, Michx. f. Hist. Arb. Am. ii. 619 = nigra.
hybrida, *Bieb. Fl. Taur. Cauc.* ii. 423.—Europ.; Reg. Caucas.
hybrida, Reichb. Ic. Fl. Germ. xxi. 29 = alba.
intermedia, Mert. ex Loud. Arboret. iii. 1640 = alba.
italica, Duroi, Harbk. ii. 141 = pyramidalis.
laevigata, Ait. Hort. Kew. ed. I. iii. 406 = monilifera.
latifolia, Moench, Meth. 338 = balsamifera.

POPULUS:—
laurifolia, Ledeb. Fl. Alt. iv. 297 = balsamifera.
leucophylla, Schur, Enum. Pl. Transs. 623 = hybrida.
Lindleyana, Booth, ex Loud. Arboret. iv. 2651 = balsamifera.
Lindleyana, Hort. ex Steud. Nom. ed. II. ii. 381 = monilifera.
lombardica, Hort. ex Link, Handb. ii. 456 = nigra.
longifolia, Fisch. ex Loud. Arboret. iii. 1673 = balsamifera.
macrocarpa, Schrenk, Enum. Pl. Nov. ii. 16 = balsamifera.
macrophylla, Lindl. in Loud. Encyc. Pl. 840 = balsamifera.
macrophylla, Lodd. ex Loud. Arboret. iii. 1671 = monilifera.
major, Mill. Gard. Dict. ed. VIII. n. 4 = alba.
marylandica, Bosc, Nouv. Dict. xi. 409 = monilifera.
Medusæ, Booth, in Loud. Arboret. iv. 2651 = monilifera.
mexicana, *Wesmael, in DC. Prod.* xvi. II. 328.—Mexic.
monilifera, *Ait. Hort. Kew.* ed. I. iii. 406.—Am. bor.
monilifera, Newberry, in Pacif. Rail. Rep. vi. Bot. 89 = Fremonti.
monticola, Mert. ex Loud. Arboret. iii. 1645 = tremula.
moscoviensis ×, Schroeder, in Gard. Chron. (1882) II. 108.
nana, Royle, Illustr. Bot. Himal. 345 = Betula nana.
neapolitana, Tenore, Fl. Nap. v. 279 = nigra.
neglecta, Hort. ex Wesmael, in DC. Prod. xvi. II. 329 = monilifera.
nigra, *Linn. Sp. Pl.* 1034.—Europ.
nigra, Mill. Gard. Dict. ed. VIII. n. 3 = alba.
nivea, Willd. Berl. Baumz. 227 = alba.
nova, Aud. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 660 = monilifera.
odorata ×, Schroeder, in Gard. Chron. (1882) II. 108.
ontariensis, Lodd. ex Loud. Arboret. iii. 1676 = balsamifera.
oxyodonta, Mert. ex Loud. l. c. 1645 = tremula.
palmata, Hort. ex Loud. l. c. 1640 = alba.
pannonica, Kit. ex Bess. Enum. Pl. Volh. 38 = nigra.
pendula, Hort. ex Steud. Nom. ed. II. ii. 381 = alba.
pendula, Hort. ex Tausch, in Flora, xxi. (1838) 754 = tremuloides.
pendula, Lodd. ex Loud. Arboret. iii. 1646 = tremula.
polonica, Hort. ex Loud. Encyc. Trees, 824 = nigra.
pruinosa, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 210.—Sibir.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 540.—Mongol.
Pseudo-balsamifera, Turcz. in Bull. Soc. Nat. Mosc. (1838) I. 101; xxvii. (1854) I. 399 = balsamifera.
Pseudo-tremula, Schur, Enum. Pl. Transs. 624 = tremula.
pyramidalis, *Salisb. Prod.* 395; *Rozier, ex Lam. Encyc.* v. 235.—Hispan.
pyramidata, Moench, Meth. 339 = nigra.
pyriformis, Royle, Illustr. Bot. Himal. 344 = ciliata.
quercifolia, Hort. ex Loud. Arboret. iii. 1640 = alba.
repanda, Baumg. Enum. Stirp. Transs. i. 349 = tremula.
reticulata, Rafin. Alsog. Am. 43 (Quid ?).—Am. bor.
rotundifolia, Griff. Notul. iv. 382 = ciliata.
rubra, Hort. ex Poir. Encyc. v. 239 = nigra.
salicifolia, Hort. ex Loud. Arboret. iii. 1673 = balsamifera.
salicifolia, Lodd. ex Loud. l. c. 1652 = nigra.
salicifolia, Rafin. Alsog. Am. 43 (= praec. ?).—Am. bor.
sibirica, Hort. ex Tausch, in Flora, xxi. (1838) 754 = tremuloides.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 29.—Japon.
Simonii, Carr. in Rev. Hortic. (1867) 360 = balsamifera.
stricta, Mert. ex Loud. Arboret. iii. 1645 = tremula.
suaveolens, Fisch. ex Loud. l. c. 1674; et ex Ledeb. Fl. Ross. iii. 629 = balsamifera.
supina, Lodd. ex G. Don, in Loud. Hort. Brit. 401 = tremula.
Tacamahacca, Mill. Gard. Dict. ed. VIII. n. 6 = balsamifera.
tomentosa, Hort. ex Loud. Arboret. iii. 1640; Carr. in Rev. Hortic. (1867) 340 = alba.
tremula, *Linn. Sp. Pl.* 1034.—Europ.
tremuliformis, Emerson, Rep. Trees Shrubs Massach. ed. I. 243; ed. II. 279 = seq.

POPULUS:—
tremuloides, *Michx. Fl. Bor. Am.* ii. 243.—Am. bor.
trepida, Willd. Sp. Pl. iv. 803 = tremuloides.
trichocarpa, *Torr. & Gray, ex Hook. Ic. Pl.* t. 878.—Calif.
versicolor, Salisb. Prod. 395 = nigra.
villosa, Láng, ex Reichb. Fl. Germ. Excurs. 173 = tremula.
viminalis, Hort. Par. ex Poir. Encyc. v. 238 = alba.
viminalis, Lodd. ex Loud. Arboret. iii. 1673 = balsamifera.
viminea, Bon Jardin. (1845) 565; ex Wesmael, in DC. Prod. xvi. II. 330 = balsamifera.
viminea, Dum.-Cours. Bot. Cult. ed. II. vi. 401 = nigra.
virginiana, Dum.-Cours. l. c. 400 = monilifera.
virginiana, *Fouger. in Mém. Agric. Par.* (Print. 1786) 87.—Am. bor.
viridis, Lodd. ex Loud. Arboret. iii. 1652 = nigra.
vistulensis, Hort. ex Loud. Encyc. Trees, 824 = nigra.

PORANA, Burm. f. Fl. Ind. 51. t. 21. f. 1 (1768).
CONVULVULACEAE, Benth. & Hook. f. ii. 876.
DINETUS, Buch.-Ham. ex D. Don, in Sweet, Brit. Flow. Gard. t. 127 (1825).
DUPERREYA, Gaudich. in Freyc. Voy. Bot. 452. t. 63 (1826).
acuminata, Beauv. Fl. Owar. i. 65. t. 49 = Neuropeltis acuminata.
cordifolia, Ledeb. Ind. Hort. Dorpat. (1824) Suppl. 6 = racemosa.
dichotoma, Buch.-Ham. ex D. Don, Fl. Nep. 99 = racemosa.
elegans, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 571.—Java.
grandiflora, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 43; *Cat.* n. 1324.—Reg. Himal.
malabarica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 223.—Ind. or.
obtusa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1883) 83.—Ins. Socotra.
paniculata, *Roxb. Hort. Beng.* [13]; *Pl. Corom.* iii. 31. t. 235; *Fl. Ind.* i. 464.—Ind. or.; Malaya.
racemosa, Dalz. & Gibs. Bomb. Fl. 162 = malabarica.
racemosa, *Roxb. Hort. Beng.* [13]; *Fl. Ind.* i. 466.—Ind. or.; Burma.
sericea, *F. Muell. Fragm.* vi. 100.—Austral.
speciosa, Benth. & Hook. f. Gen. ii. 876, sphalm. = seq.
spectabilis, *Kurz, in Journ. Bot.* xi. (1873) 136.—Burma.
stenoloba, Kurz, l. c. = grandiflora.
tomentosa, Leschen. ex Choisy, in DC. Prod. ix. 436 = paniculata.
truncata, *Kurz, in Journ. Bot.* xi. (1873) 136.—Reg. Himal.
volubilis, *Burm. f. Fl. Ind.* 51. t. 21. f. 1; *Linn. Mant.* ii. 560.—Burma; Malaya.

PORANTHERA, Rafin. in Ser. Bull. Bot. i. (1830) 221 = **Chrysopogon**, Trin. (Gramin.).
ciliata, Rafin. l. c. = C. avenaceus.
nutans, Rafin. l. c. = C. nutans.

PORANTHERA, Rudge, in Trans. Linn. Soc. x. (1811) 302. t. 22. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 262.
alpina, *Cheesem. ex Hook. Ic. Pl.* t. 1366.—N. Zel.
Arbuscula, Sieber, ex Benth. Fl. Austral. vi. 56 = corymbosa.
Arbuscula, Sond. in Linnaea, xxviii. (1856) 567 = ericoides.
cicatricosa, F. Muell. ex Benth. Fl. Austral. vi. 55 = ericoides.
corymbosa, *A. Brongn. in Duperr. Voy. Coq. Bot.* 219. t. 50. f. A.—Austral.
Drummondii, Klotzsch, in Lehm. Pl. Preiss. ii. 231 = microphylla.
ericifolia, Hueg. Bot. Archiv, t. 8 = corymbosa.
ericifolia, *Rudge, in Trans. Linn. Soc.* x. (1811) 302. t. 22.—Austral.
ericoides, *Klotzsch, in Lehm. Pl. Preiss.* ii. 232.—Austral.
glauca, Klotzsch, l. c. 231 = ericoides.
Huegelii, *Klotzsch, l. c.*—Austral.
linarioides, Sieber, ex Benth. Fl. Austral. vi. 56 = corymbosa.
microphylla, *Brongn. in Duperr. Voy. Coq. Bot.* 218. t. 50 B.—Austral.
piceoides, Klotzsch, in Lehm. Pl. Preiss. ii. 232 = ericoides.

PORAQUEIBA, Aubl. Pl. Gui. 123. t. 47 (1775).
OLACINEAE, Benth. & Hook. f. i. 352.
BARRERIA, Willd. Sp. Pl. i. 1145 (1798).
MEISTERIA, Scop. Introd. 124 (1777).
acuminata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iv. (1859) 365 = sericea.
cubensis, *Wright, ex Griseb. Cat. Pl. Cub.* 119.—Cuba.
guianensis, *Aubl. Pl. Gui.* i. 123. t. 47.—Am. trop.
sericea, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 172.—Bras.
surinamensis, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 483 = guianensis.

PORCELIA, Ruiz & Pav. Prod. 84. t. 16 (1794).
ANONACEAE, Benth. & Hook. f. i. 23.
SAPRANTHUS, Seem. Journ. Bot. iv. (1866) 369. t. 54.
cinnamomea, *Ruiz & Pav. ex G. Don, Gen. Syst.* i. 92.—Hab. ?
grandiflora, Pers. Syn. ii. 95 = Asimina grandiflora.
nitidifolia, *Ruiz & Pav. Syst.* i. 144.—Peruv.
parviflora, Pers. Syn. ii. 95 = Asimina parviflora.
pygmaea, Pers. l. c. = Asimina pygmaea.
triloba, Pers. l. c. = Asimina triloba.

PORCELLITES, Cass. in Dict. Sc. Nat. xxv. 64 (1822); xliii. 42 (1826), partim = **Hypochoeris**, Linn. (Compos.).
apargioides, Less. in Linnaea, vi. (1831) 102 = H. apargioides.
brasiliensis, Less. l. c. 103 = H. brasiliensis.
cretensis, Cass. in Dict. Sc. Nat. xliii. 45 = H. cretensis.
helveticus, Cass. l. c. 43 = H. uniflora.
maculata, Cass. l. c. = H. maculata.
radicata, Cass. l. c. = H. radicata.

PORLIERA, Pers. Syn. i. 445 (1805) = seq.

PORLIERIA, Ruiz & Pav. Prod. 55. t. 9 (1794).
ZYGOPHYLLEAE, Benth. & Hook. f. i. 268.
angustifolia, *A. Gray, Pl. Wright.* i. 28.—N. Mexic.
hygrometra, *Ruiz & Pav. Fl. Per.* iv. t. 343; *Syst. Veg.* 94.—Peruv.; Chili.
hygrometrica, Pav. ex A. Gray, Pl. Wright. i. 28, in obs. = hygrometra.

POROCARPUS, Gaertn. Fruct. ii. 473. t. 178 (1791) = **Timonius**, Rumph. (Rubiac.).
Helminthotheca, Gaertn. l. c.—Zeylan.

POROCYSTIS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 353. *SAPINDACEAE*.
toulicioides, *Radlk. l. c.* 354.—Bras.

POROPHYLLUM, Vaill. ex Linn. Coroll. Gen. 14. (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
HUNTERIA, Moç. & Sesse, ex DC. Prod. v. 649 (1836).
KLEINIA, Jacq. Select. Stirp. Am. 215. t. 127 (1763).
amplexicaule, *Engelm. ex A. Gray, Pl. Wright.* i. 120.—Mexic.
angustissimum, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 410.—Bras.
arenosum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 285 = angustissimum.
cacalioides, *DC. Prod.* v. 650.—Ins. Philipp.
caesium, *Casar. Nov. Stirp. Bras. Dec.* 86.—Bras.
coloratum, *DC. Prod.* v. 650.—Mexic.
Cusimbua, DC. l. c. = Gynura angulosa.
decumbens, *DC. l. c.*—Mexic.
ellipticum, Cass. in Dict. Sc. Nat. xliii. 56 = ruderale.
Ervendbergii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 35.—Mexic.
exsertum, *Baker, in Mart. Fl. Bras.* vi. III. 284.—Reg. Argent.
filifolium, DC. Prod. v. 651 = suffruticosum.
filifolium, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 107, nomen.—Mexic.
gracile, *Benth. Bot. Voy. Sulph.* 29.—Am. bor. occ.
Greggii, A. Gray, Pl. Wright. i. 120 = gracile, scoparium.
Haenkei, *DC. Prod.* v. 649.—Chili.
hieracioides, DC. l. c. 650 = Gynura auriculata.
japonicum, DC. l. c. = Gynura pinnatifida.
jorullense, *Cass. in Dict. Sc. Nat.* xliii. 57.—Mexic.
lanceolatum, *DC. Prod.* v. 649.—Bras.

POROPHYLLUM :—
latifolium, *Benth. in Hook. Journ. Bot.* ii. (1840) 44.—Guian.
Linaria, *DC. Prod.* v. 649.—Mexic.
linariaefolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = praec.
linarifolium, St. Lag. l. c. viii. (1861) 203 = praec.
Lindenii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 308.—Mexic.
lineare, *DC. Prod.* v. 649.—Chili.
lineare, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 284 = Riedelii.
linifolium, DC. Prod. v. 649 = suffruticosum.
macrocephalum, *DC. l. c.* 648.—Mexic.
Martii, *Baker, in Mart. Fl. Bras.* vi. III. 284.—Bras.
nummularium, *DC. Prod.* v. 649.—Mexic.
obscurum, DC. l. c. 651 = lineare.
obtusifolium, *DC. l. c.* 650.—Mexic.
oppositifolium, *DC. l. c.* 651.—Reg. Argent.
prenanthoides, *DC. l. c.* 649.—Bras.
Riedelii, *Baker, in Mart. Fl. Bras.* vi. III. 284.—Bras.
ruderale, *Cass. in Dict. Sc. Nat.* xliii. 56.—Am. trop.
scoparium, *A. Gray, Pl. Wright.* i. 119.—Texas; Mexic.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 308.—Mexic.
Selloi, *DC. Prod.* v. 651.—Reg. Argent.
suffruticosum, *Cass. in Dict. Sc. Nat.* xliii. 57.—Am. austr.
tagetoides, *DC. Prod.* v. 650.—Mexic.
tridentatum, *Benth. Bot. Voy. Sulph.* 30.—Calif.
viridiflorum, *DC. Prod.* v. 648.—Mexic.

POROSPERMUM, F. Muell. Fragm. vii. 94 (1870). *ARALIACEAE.*
Michieanum, *F. Muell. l. c.* 95.—Austral.

POROSTEMA, Schreb. Gen. 517 (1791) partim = **Nectandra**, Roland. (Laurin.).

POROTHRINAX, H. Wendl. ex Griseb. Cat. Pl. Cub. 221 (1866) = **Thrinax**, Linn. (Palm.).
Pumilio, H. Wendl. ex Griseb. l. c. = T. Pumilio.

PORPA, Blume, Bijdr. 117 (1825) = **Triumfetta**, Linn. (Tiliac.).
repens, Blume, l. c.—Java.

PORPAX, Lindl. Bot. Reg. (1845) Misc. 62 = **Eria**, Lindl. (Orchid.).
reticulata, Lindl. l. c. = E. reticulata.

PORPAX, Salisb. Gen. Pl. Fragm. 9 (1866) = **Aspidistra**, Ker-Gawl. (Liliac.).

PORPHYRA, Lour. Fl. Cochinch. 69 (1790) = **Callicarpa**, Linn. (Verben.).
dichotoma, Lour. l. c. 70 = C. purpurea.

PORPHYROCODON, Hook. f. in Benth. & Hook. f. Gen. i. 79 (1862). *CRUCIFERAE*, Benth. & Hook. f. l. c.
pictum, *Hook. f. l. c.*—N. Granat.

PORPHYROCOMA, Scheidw. in Journ. Hortic. Prat. iii. 72.—Cf. Rev. Hortic. Sér. II. iv. (1849) 104; Hook. Bot. Mag. t. 4176 (1845) = **Dianthera**, Linn. (Acanth.).
lanceolata, Scheidw. ll. cc. (= *Orthotactus Pohlianus*).—Bras.

PORPHYROSCIAS, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 62 (1867) = **Peucedanum**, Tourn. (Umbell.).
decursiva, Miq. l. c.—Japon.

PORPHYROSPATHA, Engl. in DC. Monog. Phan. ii. 289 (1879). *AROIDEAE*, Benth. & Hook. f. iii. 980.
Hoffmannii, *Engl. l. c.* 291.—Costa Rica.
Schottiana, *Engl. l. c.* 290.—Costa Rica.

PORPHYROSTACHYS, Reichb. f. Xenia Orch. i. 18 (1858) = **Stenoptera**, Presl (Orchid.).
pilifera, Reichb. l. c.—Ecuador.

PORPHYROSTEMMA, Benth. ex Oliver, in Trans. Linn. Soc. xxix. (1873) 96. *COMPOSITAE*, Benth. & Hook. f. ii. 336.
Grantii, *Benth. ex Oliver, l. c.*—Afr. trop.

PORROTERANTHE, Steud. Syn. Pl. Gram. 287 (1855) = **Glyceria**, R. Br. (Gramin.).
Drummondii, Steud. l. c. = G. fluitans.

PORRUM, [Tourn.] Linn. Syst. ed. I (1735); Moench, Meth. Suppl. 263 (1794) = **Allium**, Tourn. (Liliac.).
amethystinum, Reichb. Fl. Germ. Excurs. 140[16] = A. Ampeloprasum.
Ampeloprasum, Mill. Gard. Dict. ed. VIII. n. 2 = A. Ampeloprasum.
arenarium, Reichb. Fl. Germ. Excurs. 110 = A. vineale.
ascalonicum, Reichb. l. c. = A. ascalonicum.
capitatum, Renault, Fl. Dép. Orne, 33 = A. vineale?
Cepa, Reichb. Fl. Germ. Excurs. 110 = A. Cepa.
commune, Reichb. l. c. 111 = A. Ampeloprasum.
crinitum, Reichb. l. c. 140[16] = A. sphaerocephalum.
descendens, Reichb. l. c. 110 = A. sphaerocephalum.
fistulosum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 76 = A. fistulosum.
lineare, Rafin. Fl. Tellur. ii. 20 = A. lineare.
nutans, Rafin. l. c. = A. nutans.
oleraceum, Moench, Meth. Suppl. 264 = A. oleraceum.
Ophioscorodon, Reichb. Fl. Germ. Excurs. 110 = A. sativum.
paniculatum, Moench, Meth. Suppl. 264 = A. paniculatum.
polyanthum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. rotundum.
rotundum, Reichb. Fl. Germ. Excurs. 111 = A. rotundum.
sativum, Mill. Gard. Dict. ed. VIII. n. 1 = A. Porrum.
sativum, Reichb. Fl. Germ. Excurs. 110 = A. sativum.
Schoenoprasum, Schur, Enum. Pl. Transs. 670 = A. Schoenoprasum.
Scorodoprasum, Reichb. Fl. Germ. Excurs. 110 = A. Scorodoprasum.
sectile, Schult. f. Syst. vii. 1009, in syn. sphalm. = A. Porrum.
sibiricum, Schur, Enum. Pl. Transs. 670 = A. sibiricum.
sphaerocephalum, Reichb. Fl. Germ. Excurs. 110 = A. sphaerocephalum.
strictum, Reichb. l. c. 111 = A. strictum.
vineale, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 76 = A. vineale.

PORTAEA, Tenore, in Atti 7[ma] Adun. Sc. Ital. 902 (1846) = **Juanulloa**, Ruiz & Pav. (Solan.).
aurantiaca, Tenore, l. c. 903 = J. parasitica.

PORTALESIA, Meyen, Reise, i. 316 (1834) = **Nassauvia**, Juss. (Compos.).
procumbens, Meyen, l. c. = N. Lagascae?

PORTEA, C. Koch, Ind. Sem. Hort. Berol. (1856) 7. *BROMELIACEAE*, Benth. & Hook. f. iii. 662.
kermesina, *C. Koch, l. c.*; *et in Ann. Sc. Nat.* Sér. IV. vi. (1856) 368.—Am. trop.

PORTENSCHLAGIA, Tratt. Arch. (1812) t. 250 = **Elaeodendron**, Jacq. (Tiliac.).
australis, Tratt. l. c. = E. australe.
integrifolia, Tratt. l. c. t. 284 = E. australe.
trifoliata, Pohl, ex Engl. in Mart. Fl. Bras. xii. II. 100 = Galipea bracteata.

PORTENSCHLAGIA, Vis. Fl. Dalm. iii. 45 (1849). *UMBELLIFERAE*, Benth. & Hook. f. i. 901.
ramosissima, *Vis. l. c.*—Dalmat.

PORTERELLA, Torr. in Hayd. Rep. Geol. Surv. Montana, 488 (1872) = **Laurentia**, Neck. (Campan.).
carnosula, Torr. l. c. = L. carnosula.

PORTERIA, Hook. Ic. Pl. t. 864 (1852) = **Phyllactis**, Pers. (Rubiac.).
bractescens, Hook. Ic. Pl. 864 = Phyll. bracteata.
ebractescens, Pritz. Ic. Ind. i. 882, sphalm. = praec.
oblongifolia, Karst. Fl. Columb. ii. 99 = Phyll. Mutisiana.
parviflora, Trev. in Bot. Zeit. xi. (1853) 354.—Venezuela.
pubescens, Karst. Fl. Columb. ii. 99.—Venezuela.
rotundifolia, Karst. l. c. = Phyll. cordifolia.

PORTESIA, Cav. Diss. vii. 369. t. 215, 216 (1789) = **Trichilia**, P. Br. (Meliac.).
diversifolia, M. Roem. Syn. Hesper. 116 = T. simplicifolia.

PORTESIA :—
echinocarpa, De Vriese, in Nederl. Kruidk. Arch. i. (1848) 251 = T. echinocarpa.
glabra, Griseb. Fl. Brit. W. Ind. 130 = T. glabra.
montana, M. Roem. Syn. Hesper. 115 = T. montana.
mucronata, Cav. Diss. vii. 370 = T. heterophylla.
ovata, Cav. l. c. 369. t. 215 = T. pallida.
rimosa, Blanco, Fl. Filip. ed. II. 250 = Amoora canarana.
simplicifolia, M. Roem. Syn. Hesper. 115 = T. simplicifolia.
trinitensis, M. Roem. l. c. 116 = T. trinitensis.

PORTLANDIA, P. Br. Hist. Jamaic. 164. t. 11 (1756). *RUBIACEAE*, Benth. & Hook. f. ii. 45.
acuminata, *Willd. ex Roem. & Schult. Syst.* v. 23.—Venezuela.
coccinea, *Sw. Fl. Ind. Occ.* i. 384.—Jamaic.
coriacea, Spreng. Syst. i. 708 = coccinea.
corymbosa, Ruiz & Pav. Fl. Per. ii. 49 = Erostemma corymbosum.
daphnoides, *R. Grah. in Edinb. N. Phil. Journ.* (1840–41) 206.—Cuba.
Ghiesbreghtiana, *Baill. Adansonia*, xii. (1879) 300; *Hist. Pl.* vii. 332.—Mexic.
grandiflora, *Linn. Syst.* ed. X. 928; *Jacq. Enum. Pl. Carib.* 16.—Ind. occ.
grandiflora, Zuccagni, in Roem. Collect. 129 = Solandra grandiflora.
gypsophila, *Macfad. ex Griseb. Fl. Brit. W. Ind.* 324.—Ind. occ.
hexandra, Jacq. Enum. Pl. Carib. 16 = Coutarea speciosa.
hexandra, Vell. ex Steud. Nom. ed. II. i. 429; ii. 383 = Coutarea pubescens.
longiflora, *Meissn. ex Griseb. Cat. Pl. Cub.* 126.—Cuba.
mexicana, *Hemsl. Diagn. Pl. Nov.* 31.—Mexic.
pendula, *Wright, ex Griseb. Cat. Pl. Cub.* 126.—Cuba.
platantha, *Hook. Bot. Mag.* t. 4534.—Am. trop.
speciosa, Baill. Hist. Pl. vii. 381 = Coutarea speciosa.
tetrandra, Forst. f. Prod. 15 = Bikkia grandiflora.

PORTULA, Dill. ex Moench, Meth. 661 (1794) = **Peplis**, Linn. (Lythrar.).
diffusa, Moench, l. c. = Peplis Portula.
nummulariaefolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 381 = Peplis erecta.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 553 = Peplis Portula.
Timeroyi, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 381 = Peplis erecta.

PORTULACA, Linn. Syst. ed. I. (1735). *PORTULACEAE*, Benth. & Hook. f. i. 157.
LEMIA, Vand. Fl. Lusit. 35. t. 2. f. 15 (1788); et in Roem. Script. 116. t. 7. f. 15 (1796).
MERIDA, Neck. Elem. ii. 382 (1790).
MERIDIANA, Linn. f. Suppl. 248 (1781).
adenostemon, *Vest, in Flora*, xii. (1829) I. *Erg.* 63.—Hab.?
Anacampseros, Linn. Sp. Pl. 445 = Anacampseros Telephiastrum.
anceps, A. Rich. Tent. Fl. Abyss. i. 301 = quadrifida.
arabica, *DC. Prod.* iii. 355 (Quid?).—Arab.
arachnoides, Haw. Misc. 142 = Anacampseros arachnoides.
Armitii, *F. Muell. Fragm.* x. 97.—Austral.
aurea, Hort. ex DC. Prod. iii. 353 = oleracea.
australis, *Endl. Atakta*, 7. t. 6.—Austral.
axilliflora, *Blanco, Fl. Filip.* ed. II. 285.—Ins. Philipp.
axilliflora, *Pers. Syn.* ii. 6.—Hab.?
Bareschta, Forsk. Fl. Aegypt. Arab. p. cxii (Quid?).—Arab.
bicolor, *F. Muell. Fragm.* i. 171.—Austral.
brasiliensis, *Vahl, in West, Bidr. Ste-Croix*, 287, nomen.—Am. trop.
caffra, *Thunb. Prod. Pl. Cap.* 85.—Afr. austr.
carnosa, Domb. ex DC. Prod. iii. 358 = Calandrinia paniculata.
caryophylloides, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 722 = grandiflora.
cincta, *Fenzl, in Tijdschr. Nederl. Ind.* xiv. (1857) 162.—Malaya.
consanguinea, Schlecht. in Linnaea, xxiv. (1851) 693 = oleracea?
crassicaulis, Jacq. Hort. Vindob. iii. 29 = Talinum triangulare.

PORTULACA:—
crassifolia, Jacq. Hort. Vindob. iii. t. 52 = Talinum triangulare.
cristata, Buch.-Ham. in Wall. Cat. n. 6844 = tuberosa.
cuneifolia, Vahl, Symb. Bot. i. 33 = Talinum cuneifolium.
decumbens, Vahl, l. c. = Orygia decumbens.
dichotoma, Gueldenst. Reisen, i. 192 = oleracea.
digyna, *F. Muell. Fragm.* i. 170.—Austral.
diptera, Zipp. ex Span. in Linnaea, xv. (1841) 207 = quadrifida.
elatior, *Mart. ex Rohrb. in Mart. Fl. Bras.* xiv. II. 302.—Bras.
eriophora, Casar. Nov. Stirp. Bras. Dec. ii. 23 = pilosa.
filamentosa, Haw. Misc. 142 = Anacampseros filamentosa.
filifolia, *F. Muell. Fragm.* i. 169.—Austral.
flava, DC. Prod. iii. 355, sphalm. = lutea.
foliosa, DC. Calq. Dess. Fl. Mex. 389 = pilosa.
foliosa, *Ker-Gawl. in Bot. Reg.* t. 793.—Afr. trop.
fruticosa, Linn. Syst. ed. X. 1045 = Talinum triangulare.
fruticosa, Thunb. Prod. Pl. Cap. 85 = Portulacaria afra.
fulgens, *Griseb. in Goett. Abh.* xxiv. (1879) 30.—Reg. Argent.
geniculata, Royle, Illustr. Bot. Himal. 221 = quadrifida.
Gilliesii, *Hook. Bot. Mag.* t. 3064.—Chili.
grandiflora, *Hook. l. c.* t. 2885; *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 192.—Bras.
guianensis, Steud. Nom. ed. II. ii. 382, sphalm. = seq.
guineensis, *Lindl. ex DC. Prod.* iii. 353.—Afr. trop.
halimoides, *Linn. Sp. Pl.* ed. II. 639.—Ind. occ.
helianthemoides, *Zipp. ex Span. in Linnaea*, xv. (1841) 207, nomen.—Ins. Timor.
Hilaireana, *G. Don, Gen. Syst.* iii. 74.—Bras.
hirsutissima, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 191.—Bras.
hortensis, Rupr. Fl. Ingr. 388 = oleracea.
imbricata, *Forsk. Fl. Aegypt. Arab.* 92.—Arab.
involucrata, Haw. Rev. Pl. Succ. 59 = suffruticosa.
Karwinskii, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 63.—Mexic.
laevis, Buch.-Ham. in Wall. Cat. n. 6841 = oleracea.
lanata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = pilosa.
lanceolata, *Engelm. in A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 154.—Texas.
lanceolata, Haw. Misc. 142 = Anacampseros lanceolata.
lanuginosa, Crantz, Inst. ii. 428; H. B. & K. Nov. Gen. et Sp. vi. 74 = pilosa.
Laruotteana, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 190 = mucronata.
latifolia, Hornem. Hort. Hafn. ii. 491 = oleracea.
linearis, Domb. ex DC. Prod. iii. 358 = Calandrinia lingulata.
linifolia, Forsk. Fl. Aegypt. Arab. 92 = quadrifida.
linifolia, *Roxb. Hort. Beng.* [36] nomen.—Ind. or.
lutea, *Soland. ex Forst. f. Pl. Escul.* 72; *Seem. Fl. Vit.* 9.—Ins. Societ.
macrorhiza, *Zipp. ex Span. in Linnaea*, xv. (1841) 207, nomen.—Ins. Timor.
marginata, *H. B. & K. Nov. Gen. et Sp.* vi. 72.—Venezuela.
maritima, S. G. Gmel. Reise, iv. 40 = oleracea.
megalantha, Steud. Nom. ed. II. ii. 383 = grandiflora.
mendocinensis, Gill. ex Rohrb. in Mart. Fl. Bras. xiv. II. 303 = grandiflora.
meridiana, Linn. f. Suppl. 248 = quadrifida.
microphylla, *A. Rich. Ess. Fl. Cub.* 620.—Cuba.
mucronata, *Hort. Vindob. ex Link, Enum. Hort. Berol.* ii. 2.—Bras.
napiformis, *F. Muell. ex Benth. Fl. Austral.* i. 169.—Austral.
officinarum, Crantz, Inst. ii. 428 = oleracea.
oleracea, *Linn. Sp. Pl.* 445.—Reg. calid.
oligosperma, *F. Muell. Fragm.* i. 170.—Austral.
olitoria, Pall. Reise, iii. 594 = oleracea.
paniculata, Jacq. Enum. Pl. Carib. 22 = Talinum patens.
paniculata, Linn. Sp. Pl. ed. II. 640 = Talinum triangulare.
parviflora, Griseb. Fl. Brit. W. Ind. 57, sphalm. = seq.
parvifolia, Haw. Syn. Pl. Succ. 122 = oleracea.
patens, Jacq. Hort. Vindob. ii. 71 = Talinum patens.
patens, *Linn. Mant.* ii. 242 (an praec. ?).—Am. trop.

PORTULACA:—
pilosa, Herb. Madr. ex Wall. Cat. n. 6844 = tuberosa.
pilosa, *Linn. Sp. Pl.* 445.—Am. bor. et austr.
pilosissima, *Hook. Bot. Misc.* ii. (1831) 220.—Peruv.
Portulacastrum, Linn. Sp. Pl. 446 = Sesuvium Portulacastrum.
prolifera, *Schum. & Thonn. Beskr. Guin. Pl.* 289.—Afr. trop.
prostrata, Domb. ex Lem. in Fl. des Serres, ii. (Avril, 1846) sub t. 5 = Calandrinia umbellata.
psammotropha, *Hance, in Walp. Ann.* ii. 660.—China.
pusilla, *H. B. & K. Nov. Gen. et Sp.* vi. 72.—Venezuela.
quadrifida, *Linn. Mant.* i. 73.—Geront. trop.
racemosa, Linn. Sp. Pl. ed. II. 640 = Talinum triangulare.
rediviva, *Wawra, in Sitzb. Acad. Wien, Math. Nat.* xxxviii. (1860) 564.—Afr. trop.
reflexa, Haw. Misc. 141 = Talinum patens.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = oleracea.
repens, Roxb. ex Wight & Arn. Prod. 356 = quadrifida.
retusa, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 154.—Am. bor. occ.
rostellata, Brign. in Linnaea, xviii. (1844) 509; Aless. in Nuov. Ann. Sc. Nat. Bologna, Ser. II. ii. (1844) 207 = marginata.
rubens, Haw. Misc. 142 = Anacampseros arachnoides.
rubricaulis, *H. B. & K. Nov. Gen. et Sp.* vi. 73.—Am. austr.
sativa, Haw. Misc. 36 = oleracea.
saxea, *Vell. Fl. Flum.* 202; v. t. 13.—Bras.
saxifragoides, *Welw. in Oliver, Fl. Trop. Afr.* i. 149.—Afr. trop.
sclerocarpa, *A. Gray, Bot. U. St. Expl. Exped.* i. 141.—Ins. Sandvic.
sedoides, Spruce, ex Rohrb. in Mart. Fl. Bras. xiv. II. 304 = pilosa.
sedoides, *Welw. in Oliver, Fl. trop. Afr.* i. 149.—Afr. trop.
setacea, Haw. Misc. 139 = pilosa.
simpliciuscula, *Mart. ex Rohrb. in Mart. Fl. Bras.* xiv. II. 301.—Bras.
splendens, Lindl. Bot. Reg. (1843) t. 34 = Thellusonii ?
stelliformis, [Moç. & Sesse,] ex DC. Prod. iii. 353 = foliosa.
stipularis, Dryand. ex DC. l. c. 356 = Anacampseros filamentosa.
suffrutescens, *Engelm. in Coult. Bot. Gaz.* vi. (1881) 236.—Arizona.
suffruticosa, Thw. Enum. Pl. Zeyl. 24 = oleracea.
suffruticosa, *Wight, ex Wall. Cat.* n. 6842.—Ind. or.
teretifolia, H. B. & K. Nov. Gen. et Sp. vi. 72 = pilosa ?
Thellusonii, *Lindl. Bot. Reg.* (1840) t. 31.—Hab. ?
Toston, Blanco, Fl. Filip. ed. I. 408 = Trianthema monogyna.
triangularis, Jacq. Enum. Pl. Carib. 22 = Talinum triangulare.
trigona, *Thunb. Prod. Pl. Cap.* 85.—Afr. austr.
tuberosa, *Roxb. Hort. Beng.* [91]; *Fl. Ind.* ii. 464.—Ind. or.
umbraticola, *H. B. & K. Nov. Gen. et Sp.* vi. 72.—N. Granat.
villosa, *Cham. in Linnaea*, vi. (1831) 565.—Ins. Sandvic.
viridis, Hort. ex DC. Prod. iii. 353 = oleracea.
Wightiana, *Wall. Cat.* n. 6845.—Ind. or.

PORTULACARIA, Jacq. Coll. i. 160. t. 22 (1786). *PORTULACEAE*, Benth. & Hook. f. i. 157.
afra, *Jacq. Coll.* i. 160. t. 22.—Afr. austr.
namaquensis, *Sond. in Harv. & Sond. Fl. Cap.* ii. 386—Afr. austr.

PORTULACASTRUM, Juss. ex Medic. Phil. Bot. i. 99 (1789) = **Trianthema**, Sauv. (Ficoid.).
monogynum, Medic. l. c. = T. monogyna.

PORTULACCA, Haw. Syn. Pl. Succ. 121 (1812) = **Portulaca**, Linn.

PORTUNA, Nutt. in Trans. Am. Phil. Soc. viii. (1843) 268 = **Pieris**, D. Don (Ericac.).
floribunda, Nutt. l. c. = Pieris floribunda.

POSIDONIA, Kon. in Kon. & Sims, Ann. Bot. ii. (1805) 95. t. 6. *NAIADACEAE*, Benth. & Hook. f. iii. 1015.
AEGLE, Dulac, Fl. Hautes-Pyr. 43 (1867).
CAULINIA, DC. Fl. Fr. iii. 156 (1805).
KERNERA, Willd. Sp. Pl. iv. 947 (1805).
TAENIDIUM, Targ. Tozz. Cat. Veg. Mar. 80. t. 1 (1826).
antarctica, Spreng. Syst. i. 181 = Cymodocea antarctica.
australis, *Hook. f. Fl. Tasman.* ii. 43.—Austral.
Caulini, *Kon. in Kon. & Sims, Ann. Bot.* ii. (1805) 96.—Reg. Mediterr.
oceanica, Delile, Fl. Aegypt. 30 = Caulini.
serrulata, Spreng. Syst. i. 181 = Cymodocea serrulata.

POSIDONION, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = praec.

POSKEA, Vatke, in Linnaea, xliii. (1880–82) 321. *BORAGINEAE*.
africana, *Vatke, l. c.*—Afr. trop.

POSOQUERIA, Aubl. Pl. Gui. i. 133. t. 51 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 83.
CYRTANTHUS, Schreb. Gen. 122 (1789).
PASORIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 80.
RAMSPEKIA, Scop. Introd. 145 (1777).
SOLENA, Willd. Sp. Pl. i. 961 (1797).
STANNIA, Karst. Ausw. Neue Gew. 27. t. 9 (1848).
acuminata, Mart. in Flora, xxiv. (1841) II. Beibl. 79 = longiflora.
acutifolia, *Mart. l. c.* 77.—Bras.
coriacea, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 240.—Mexic.
decora, DC. Prod. iv. 375 = latifolia.
drupacea, Gaertn. f. Fruct. iii. 77. t. 195. f. 1 = Randia drupacea.
dumetorum, Roxb. Fl. Ind. ed. Carey, ii. 564 = Randia dumetorum.
fasciculata, Roxb. l. c. 568 = Randia fasciculata.
floribunda, Roxb. l. c. 569 = Randia dumetorum.
formosa, *Planch. in Fl. des Serres*, vi. (1850–51) 169.—Venezuela.
fragrans, Roxb. Fl. Ind. ed. Carey, ii. 567 = Randia malabarica.
fragrantissima, *Linden & André, Illustr. Hortic.* xvii. (1870) t. 27.—Bras.
gracilis, *Roem. & Schult. Syst.* v. 227.—Guian.
havanensis, *DC. Prod.* iv. 375.—Cuba.
insignis, Nees, in Flora, iv. (1821) 301 = latifolia.
intermedia, Miq. in Linnaea, xix. (1847) 135 = latifolia.
latifolia, *Roem. & Schult. Syst.* v. 227.—Am. trop.
laurifolia, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 78.—Bras.
longiflora, *Aubl. Pl. Gui.* i. 134. t. 51.—Guian.
longiflora, Roxb. Fl. Ind. ed. Carey, ii. 569 = Randia longiflora.
longispina, Roxb. l. c. 566 = Randia dumetorum.
lucida, Mart. in Flora, xxiv. (1841) II. Beibl. 77 = latifolia.
macrophylla, Hemsl. Biol. Centr. Am. Bot. ii. 89 = latifolia.
macropus, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 79.—Bras.
montana, Mart. l. c. 80 = Hillia parasitica.
multiflora, Blume, Bijdr. 980 = Randia longiflora.
multiflora, *Lem. Illustr. Hortic.* xvi. (1869) t. 597.—Bras.
nutans, Roxb. Fl. Ind. ed. Carey, ii. 565 = Randia dumetorum.
palustris, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 78.—Bras.
panamensis, Walp. Ann. ii. 797 = latifolia.
revoluta, Schrad. in Goett. Gel. Anz. i. (1821) 714 = latifolia.
rigida, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 570 = Randia fasciculata.
Trinitatis, *DC. Prod.* iv. 375.—Ins. Trinit.
uliginosa, Roxb. Fl. Ind. ed. Carey, ii. 563 = Randia uliginosa.
versicolor, *Lindl. Bot. Reg.* (1841) t. 26.—Cuba.

POSORIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 80 = **Posoqueria**, Aubl. (Rubiac.).

POSSIRA, Aubl. Pl. Gui. ii. 934. t. 355 (1775) = **Swartzia**, Schreb. (Legumin.).
arborescens, Aubl. l. c. = S. triphylla.
dodecandra, Poir. Encyc. v. 577 = S. dodecandra.
grandiflora, Poir. Encyc. Suppl. iv. 532 = S. grandiflora.
pinnata, Poir. l. c. 533 = S. pinnata.
simplex, Sw. Prod. Veg. Ind. Occ. 82; Poir. Encyc. v. 577 = S. grandiflora.
triphylla, Sw. l. c. = S. triphylla.

POSSIRIA, Steud. Nom. ed. II. ii. 653, in syn. (1841), sphalm. = praec.

POSSURA, Aubl. ex Steud. Nom. ed. II. ii. 383 (1841) = praec.
guianensis, Aubl. ex Steud. l. c. = S. triphylla.

POSTIA, Boiss. & Blanche, in Boiss. Fl. Orient. iii. 182 (1875). *COMPOSITAE*, Benth. & Hook. f. ii. 1233.
bombycina, *Boiss. & Haussk. in Boiss. l. c.* 183.—Persia.
lanuginosa, *Boiss. l. c.* 182.—Syria.
microcephala, *Boiss. l. c.* 183.—Persia.
puberula, *Boiss. & Haussk. l. c.*—Persia.

POSTUERA, Rafin. Sylva Tellur. 10 (1838) = **Notelaea**, Vent. (Oleac.).
longifolia, Rafin. l. c. = N. longifolia.

POTALIA, Aubl. Pl. Gui. i. 394. t. 151 (1775). *LOGANIACEAE*, Benth. & Hook. f. ii. 795.
NICANDRA, Schreb. Gen. 283 (1789).
amara, *Aubl. Pl. Gui.* i. 394. t. 151.—Guian.
amara, Desf. Cat. Hort. Par. ed. III. 127 = Clavija lancifolia.
resinifera, Mart. Nov. Gen. et Sp. ii. 89 = amara.

POTAMEIA, Thou. Gen. Nov. Madag. 5 (1806). *LAURINEAE*, Benth. & Hook. f. iii. 154.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 446.—Madag.
Thouarsii, *Roem. & Schult. Syst.* iii. 476.—Madag.

POTAMICA, Poir. in Dict. Sc. Nat. xliii. 94 (1826) = praec.

POTAMOBRYON, Liebm. in Forh. Skand. Naturf. Kjoeb. (1847) 513. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 1219.
concinnum, *Liebm. l. c.*—Mexic.
laxum, *Liebm. l. c.* 514.—Mexic.
patulum, *Liebm. l. c.* 515.—Mexic.

POTAMOCHARIS, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 296 = **Mammea**, Linn. (Guttifer.).
Mamei, Rottb. l. c. = M. americana.

POTAMOCHLOA, Griff. in Journ. As. Soc. Beng. v. (1836) 571 = **Hygroryza**, Nees (Gramin.).
Retzii, Griff. l. c. 571. t. 24 = H. aristata.

POTAMOGETON, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 33 (1737). *NAIADACEAE*, Benth. & Hook. f. iii. 1014.
BUCCAFERREA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 316.
GROELANDIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 169.
GROENLANDIA, J. Gay, in Comp. Rend. Acad. Sc. xxxviii. (1854) 703.
PELTOPSIS, Rafin. in Journ. Phys. lxxxix. (1819) 101.
SPIRILLUS, J. Gay, in Comp. Rend. Acad. Sc. xxxviii. (1854) 703.
acuminatus, Schum. Enum. Pl. Saell. i. 49 = lucens.
acuminatus, Trevir. ex Roem. & Schult. Syst. iii. 509, in obs. = lucens?
acuminatus, Wahlenb. Fl. Upsal. 58 = praelongus.
acutifolius, Hornem. in Fl. Dan. t. 2046 = compressus.
acutifolius, *Link, in Roem. & Schult. Syst.* iii. 513.—Europ.; Reg. Caucas.; Austral.
acutifolius, J. & C. Presl, Fl. Cech. 37 = Friesii.
affinis, Boenn. ex Cham. & Schlecht. in Linnaea, ii. (1827) 216 = natans.
alpino-natans, *F. Schultz, in Pollichia*, xx-xxi. (1863) 228.—Europ. centr.
alpinus, Balb. Misc. 13 = rufescens.
amblyophyllus, C. A. Mey. Beitr. Pfl. Russ. Reich. vi. 10 = pectinatus.

POTAMOGETON:—
americanus, Cham. & Schlecht. in Linnaea, ii. (1827) 226 = lucens?
americanus, Roem. & Schult. Syst. iii. 509 = lucens.
amplexicaulis, *Kar. in Bull. Soc. Nat. Mosc.* (1839) 173.—Turkestan.
amplifolius, *Tuckerm. in Am. Journ. Sc.* Ser. I. xlv. (1843) 38; II. vi. (1848) 225.—Am. bor.
angustatus, Wallm. Fl. Ostrog. ed. I. (1812) (ex Ar. Benn. MS.) = acutifolius.
angustifolius, Bercht. & Presl, ex Opiz, Boehm. Gew. 23 = lucens?
angustifolius, Bercht. & Presl, Rostl. i. Alismac. 19 = pectinatus.
angustissimus, *H. B. & K. Nov. Gen. et Sp.* i. 370.—Mexic.
annulatus, Bell. in Mém. Acad. Turin, x, xi. (1802–3) 447 = rufescens.
Augustani, Bell. l. c. 448, in nota = seq.?
augustanus, Balb. Misc. 14. t. 3 = gramineus.
Berchtoldi, Fieb. in Berch. & Fieb. Potamog. 40. t. 4 = pusillus.
Berteroanus, *Phil. in Linnaea*, xxx. (1859–60) 200.—Chili.
Besseri, Steud. Nom. ed. II. ii. 384 = natans.
bifolius, Lapeyr. Hist. Abr. Pl. Pyren. Suppl. 27 = Vicia Faba (fide Benth. Cat. Pl. Pyr. 113).
Billotii, F. Schultz, Arch. Fl. Fr. et Allem. 61 = fluitans.
borealis, Rafin. in Med. Repos. N. York, v. (1808) 354 = marinus.
caespitosus, *Humnicki, Cat. Pl. Luxeuil*, 60; *Suppl.* 90.—Europ.
caespitosus, Nolte, in Fl. Dan. t. 2889 = rutilus.
canariensis, Link, in Buch, Beschr. Canar. Ins. 138 = fluitans.
capillaceus, Poir. Encyc. Suppl. iv. 535 = hybridus.
caricifolius, Willd. ex Schult. Mant. iii. 367 = tenuifolius.
Casparii, Weyl, in Oestr. Bot. Zeitschr. xx. (1870) 321 = rufescens.
Casparyi, Kohts, l. c. 291 = praec.
caudatus, Seidl, ex Opiz, Boehm. Gew. 23 = lucens.
cespitosus, Reichb. f. Skizze von Nolte, 21 = rutilus.
Cheesemanii, *Ar. Benn. in Journ. Bot.* xxi. (1883) 66.—N. Zel.
Claytonii, *Tuckerm. in Am. Journ. Sc.* Ser. I. xlv. (1843) 38.—Am. bor.
coloratus, *Hornem. in Fl. Dan.* t. 1449.—Europ.
complanatus, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 297 = compressus.
compressus, All. Fl. Pedem. i. 240 = acutifolius.
compressus, *Linn. Sp. Pl.* 127.—Reg. bor. temp.
compressus, Reichb. Fl. Germ. Excurs. 8 = Friesii.
compressus, Sm. Fl. Brit. i. 856; Mert. & Koch, Deutschl. Fl. i. 856 = pusillus.
compressus, Wahlenb. Fl. Suec. 110 = obtusifolius.
condylocarpus, Tausch, in Flora, xix. (1836) 423 = densus.
confervoides, Reichb. Ic. Fl. Germ. vii. 13 = Tuckermani.
contortus, Desf. Fl. Atlant. i. 150 = Zannichellia contorta.
corniculatus, Schur, Enum. Pl. Transs. 632 = lucens.
cornutus, J. & C. Presl, Fl. Čech. 37 = lucens.
crassipes, Kit. ex Cham. & Schlecht. in Linnaea, ii. (1827) 204 = gramineus.
crenulatus, D. Don, Prod. Fl. Nep. 22 = crispus.
crispatus, Wallm. ex Reichb. Fl. Germ. Excurs. 8 = crispus.
crispus, Darl. Fl. Cestr. 23 = perfoliatus.
crispus, Laterr. Fl. Bordel. (1811) 71, partim = heterophyllus.
crispus, *Linn. Sp. Pl.* 126.—Reg. temp. bor. et austr.
cristatus, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 139.—Reg. Amur.
curvifolius, Hartm. Fl. Scand. ed. I. 78 = nitens.
cuspidatus, Schrad. ex Mert. & Koch, Deutschl. Fl. i. 854 = acutifolius.
cuspidatus, Schrad. in Sm. Engl. Fl. i. 234 = compressus.
decipiens, *Nolte, ex Koch, Syn. Fl. Germ.* ed. II. 779.—Europ.
delicatulus, Bertol. Misc. Bot. xv. 17 = hybridus.
densus, *Linn. Sp. Pl.* 126.—Reg. bor. temp.
densus, Schwein. ex Beck, Bot. U. St. 368 = Elodea canadensis.

POTAMOGETON:—
denticulatus, Link, in Buch, Beschr. Canar. Ins. 138 = pusillus.
denticulatus, Mohl & Schlecht. in Bot. Zeit. xiv. (1856) 783, sphalm. = *delicatulus*.
digynus, *Wall. Cat.* n. 5177.—Ind. or.
dimorphus, Rafin. in Am. Monthly Mag. (1817) 358 = P. Spirillus.
distachyus, Bell. in Mém. Acad. Turin, x, xi. (1802–3) 447, in nota = gramineus.
divaricatus, *Wolfg. ex Schult. Mant.* iii. 355.—Hab.?
diversifolius, Barton, Fl. Philad. Prod. 27 = P. Spirillus.
diversifolius, Rafin. in Med. Repos. N. York, v. (1808) 354 = hybridus.
Drummondii, *Benth. Fl. Austral.* vii. 171.—Austral.
drupaceus, Láng, in Flora, xxix. (1846) 472 = pectinatus.
elegans, Wall. Cat. n. 5178 = polygonifolius.
epihydrum, Rafin. in Med. Repos. N. York, v. (1808) 354 = natans?
exstipulatus, Muhl. Cat. 18 = pauciflorus.
exstipulatus, Willd. ex Schult. Mant. iii. 365 = angustissimus, pauciflorus.
fasciculatus, Wolfg. in Schult. l. c. = marinus.
filicaulis, Schur, Enum. Pl. Transs. 633 = pectinatus.
filifolius, Phil. Fl. Atac. 50 = Ruppia maritima.
filiformis, Nolte, Nov. Fl. Holsat. 20 = marinus.
filiformis, Pers. Syn. i. 152 = pectinatus.
filiformis, Pursh, ex Tuckerm. in Am. Journ. Sc. Ser. II. vi. (1848) 230 = hybridus.
filiformis, *Rafin. in Med. Repos. N. York*, Ser. III. ii. (1811) 409.—Am. bor.
flabellatus, Bab. Man. Brit. Bot. ed. III. 343 = pectinatus.
flexicaulis, Dethard. [in Strelitz. Anzeig. (1809) n. 50] ex Koch, Syn. Fl. Germ. ed. I. 676 = praelongus.
flexuosus, Schleich. Cat. Pl. Helv. ed. III. 23; iv. 27 = praelongus.
fluitans, Liljebl. Fl. Suec. ed. II. 71 = sparganifolius.
fluitans, Pursh, Fl. Am. Sept. i. 120 = Claytonii.
fluitans, *Roth, Tent. Fl. Germ.* i. 72; ii. 202.—Reg. temp. et trop.
fluitans, Sm. Fl. Brit. iii. 1391 = rufescens.
fluviatilis, Roem. & Schult. Syst. iii. 570 (Index) = lucens.
foliorum, Rafin. in Med. Repos. N. York, Ser. III. ii. (1811) 409 = gramineus?
foliosus, Rafin. l. c. Ser. II. v. (1808) 354 = pauciflorus.
Friesii, *Rupr. Beitr. Pfl. Russl.* iv. 43.—Europ.
Gaudichaudii, Cham. & Schlecht. in Linnaea, ii. (1827) 197 = mucronatus?
gemmiparus, Robbins, ex Morong, in Coult. Bot. Gaz. v. (1880) 51 = pusillus?
glaucus, Gueldenst. Reisen, i. 49 (Quid?).—Ross. med.
gracilis, *Fries, Nov. Fl. Suec.* ed. II. 50.—Suecia.
gracilis, *Wolfg. in Schult. Mant.* iii. 355.—Sibir.
gramineus, R. Br. Prod. 343 = ochreatus.
gramineus, *Linn. Sp. Pl.* 127.—Europ.
gramineus, Michx. Fl. Bor. Am. i. 102 = pauciflorus.
gramineus, Savi, Fl. Pis. i. 194 = pusillus.
gramineus, Sm. Fl. Brit. i. 196 = obtusifolius.
gramineus, Wahlenb. Fl. Upsal. 57 = nitens.
graminifolius, Benkö, Transs. i. 123 = gramineus.
Griffithii, *Ar. Benn. in Journ. Bot.* xxi. (1883) 65.—Cambria.
Grisebachii, Heuff. in Verh. Zool. Bot. Ges. Wien, viii. (1858) 200 = rutilus.
helodes, Dum. Fl. Belg. 163 = coloratus.
heterophyllus, Auct. ex Fries, Nov. Fl. Suec. ed. II. 35; Sm. ex Nym. Consp. 682 = nitens.
heterophyllus, Biroli, Fl. Acon. i. 53 = polygonifolius?
heterophyllus, Buch.-Ham. ex Wall. Cat. n. 5181 = javanicus.
heterophyllus, Ell. Sketch, i. 222 = P. Spirillus?
heterophyllus, Schreb. Spicil. Fl. Lips. 21 = gramineus.
heterophyllus, Torr. Comp. Fl. N. St. 88 = hybridus.
Hillii, *Morong, in Coult. Bot. Gaz.* vi. (1881) 290.—Am. bor.
Hornemanni, E. Mey. Chlor. Hanov. 521 = coloratus.
hybridus, *Michx. Fl. Bor. Am.* i. 101.—Am. bor.
hybridus, Petagna, Inst. Bot. ii. 289; Thuill. Fl. Par. ed. II. 86 = gramineus.
illinoënsis, *Morong, in Coult. Bot. Gaz.* v. (1880) 50.—Am. bor.
indicus, Roth, ex Roem. & Schult. Syst. iii. 516 = Aponogeton monostachyon.
indicus, *Roxb. Hort. Beng.* 12; *Fl. Ind.* i. 452.—Ind. or.

POTAMOGETON :—
intermedius, *Cham. & Schlecht. in Linnaea*, ii. (1827) 221.—Europ.
interruptus, Kit. in Schult. Oestr. Fl. ed. II. i. 328 = pectinatus.
japonicus, Franch. & Sav. Enum. Pl. Jap. ii. 15, nomen = mucronatus.
javanicus, *Hassk. in Verh. Natuurk. Ver. Nederl. Ind.* i. (1856) 26.—As. trop.
Kirkii, Syme, in Engl. Bot. ed. III. ix. 31 = natans.
Kochii, F. Schultz, in Arch. Fl. Fr. et Allem. 72 = spathulatus.
lanceolatus, Eichw. Skizze, 126 = lucens.
lanceolatus, Fl. Corc. in Ionios Anthol. iv. 954 (err. typ. 854) = pectinatus.
lanceolatus, Poir. Encyc. Suppl. iv. 536 = lanciformis.
lanceolatus, Reichb. Ic. ex Nym. Consp. 681 = salicifolius.
lanceolatus, *Sm. Engl. Bot.* t. 1985.—Anglia.
lancifolius, Roth, ex Steud. Nom. ed. II. ii. 384 = densus.
lanciformis, *Roem. & Schult. Syst.* iii. 512.—Carinthia.
lateralis, *Morong, in Coult. Bot. Gaz.* v. (1880) 51.—Am. bor.
laticaulis, Wahlenb. Fl. Suec. 111 = acutifolius.
latifolius, Cham. & Schlecht. in Linnaea, ii. (1827) 731 (Index) = praec ?
Leschenaultii, Cham. & Schlecht. l. c. 222. t. 6. f. 23 = fluitans.
Liebmanni, Buchen. in Bremen, Abh. iii. (1872) 349 = Heteranthera graminea.
lithuanicus, Gorski, ex Reichb. Ic. Fl. Germ. vii. 19 = salicifolius.
Loeselii, [Honck.] Verz. Aller Gew. Teutschl. 487; Roem. & Schult. Syst. iii. 508 = perfoliatus.
lonchites, Tuckerm. in Am. Journ. Sc. Ser. II. vi. (1848) 226 = lucens.
longepedunculatus, Mérat, Rev. Fl. Par. 494 = gramineus.
longifolius, Bab. in Engl. Bot. Suppl. t. 2847; et Man. Brit. Bot. ed. I. 323 = decipiens.
longifolius, J. Gay, ex Poir. Encyc. Suppl. iv. 535 = lucens.
lucens, Geners. Fl. Scepus. (1798) 146 = gramineus.
lucens, *Linn. Sp. Pl.* 126.—Geront. trop.
lucens, [Weber] Wigg. Prim. Fl. Holsat. 15 = praelongus.
lucidus, Gueldenst. Reisen, i. 76; Salisb. Prod. 68 = lucens.
macrophyllus, Wolfg. ex Schult. Mant. iii. 358 = lucens.
malaianus, Miq. Illustr. Fl. Arch. Ind. 146 = mucronatus.
marianensis, *Cham. & Schlecht. in Linnaea*, ii. (1827) 228.—Ins. Marian.
marinus, Huds. Fl. Angl. ed. II. 77; Desf. Fl. Atlant. i. 150 = pectinatus.
marinus, *Linn. Sp. Pl.* 127.—Europ.
marinus, Pall. Reise, ii. 393 = pusillus.
maritimus, Pohl, Tent. Fl. Bohem. i. 159, in syn. = marinus.
mascarensis, *Cham. & Schlecht. in Linnaea*, ii. (1827) 228.—Ins. Mascar.
maximus, Roehl. ex Steud. Nom. ed. I. 648, 649 = densus.
microcarpus, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 24 = polygonifolius.
microstachys, Wolfg. ex Schult. Mant. iii. 360 = rufescens.
monogynus, J. Gay, ex Webb & Berth. Phyt. Canar. iii. 300 = trichoides.
monogynus, *J. Gay, in Coss. & Germ. Suppl. Fl. Par.* 89; *Fl. Par.* 572.—Europ.
montanus, *Presl, Rel. Haenk.* ii. 85.—Am. austr.
mucronatus, Nym. Syll. 387 = pusillus.
mucronatus, *Presl, Epim. Bot.* 245.—Ins. Philipp.
mucronatus, Schrad. ex Roem. & Schult. Syst. iii. 517 = Friesii *vel* lucens.
mysticus, *Morong, in Coult. Bot. Gaz.* v. (1880) 50.—Am. bor.
natans, *Linn. Sp. Pl.* 126.—Reg. temp. et subtrop.
natans, Sturm, Deutschl. Fl. Heft 9 = polygonifolius.
natans, Thunb. Prod. Pl. Cap. 32 = Thunbergii.
nerviger, Wolfg. ex Schult. Mant. iii. 359 = rufescens.
nervigerus, Eichw. Skizze, 125 = praec.
niagarensis, *Tuckerm. in Am. Journ. Sc.* Ser. II. vii. (1849) 354.—Am. bor.
nigrescens, Fries, Mant. iii. 17 = rufescens.
nitens, *Weber, Fl. Holsat. Suppl.* n. 11.—Europ.

POTAMOGETON :—
nitens, Willd. ex Cham. & Schlecht. in Linnaea, ii. (1827) 202 = lucens.
nodosus, Poir. Encyc. Suppl. iv. 535 = fluitans.
Nuttallii, Cham. & Schlecht. in Linnaea, ii. (1827) 226 = Claytonii ?
Oakesianus, *Robbins, ex A. Gray, Man. Bot. U. St.* ed. V. 485.—Am. bor.
oblongifolius, Hook. f. Handb. N. Zeal. Fl. 742 = gramineus.
oblongo-rufescens, Schultz, in Flora, xii. (1829) 230 = spathulatus.
oblongus, E. Mey. Chlor. Hanov. 519 = fluitans.
oblongus, Viv. Fragm. Fl. Ital. i. 1. t. 2 = polygonifolius.
obrutus, Wood, Class-Book (1845), 178; (1861) 675 = rufescens.
obscurus, DC. Fl. Fr. Suppl. 311 = rufescens.
obtusifolius, *Mert. & Koch, Deutschl. Fl.* i. 855.—Reg. temp. bor. et austr.
obtusus, Ducros, ex Gaud. Fl. Helv. i. 546 = rufescens.
occidentalis, *Sieber, ex Cham. & Schlecht. in Linnaea*, ii. (1827) 224.—Ins. Martinic.
ochreatus, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 117.—N. Zel.
octandrus, *Poir. Encyc. Suppl.* iv. 534.
Oederi, G. F. W. Mey. Fl. Hanov. Excurs. 536 = Friesii.
oppositifolius, DC. Bot. Gall. iii. 186 = densus.
owaihiensis, *Cham. & Schlecht. in Linnaea*, ii. (1827) 228.—Ins. Sandvic.
oxyphyllus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 161.—Japon.
paludosus, Bory, ex Cham. & Schlecht. in Linnaea, ii. (1827) 216 = natans.
palustris, Tees̩d. in Trans. Linn. Soc. v. (1800) 43 = gramineus.
panormitanus, Bivona, in Bivona f. Ined. 6; ex Guss. Fl. Sic. Syn. i. 207 = pusillus.
parnassifolius, Schrad. ex Mert. & Koch, Deutschl. Fl. i. 839 = natans.
parvifolius, Buchen. in Bremen, Abh. vii. (1880) 32 = javanicus.
pauciflorus, Lam. Fl. Fr. iii. 209 = densus.
pauciflorus, *Pursh, Fl. Am. Sept.* i. 121.—Am. bor.
paucifolius, Opiz, in Flora, v. (1822) 266 = gramineus.
pectinatus, *Linn. Sp. Pl.* 127.—Reg. temp. bor. et austr.
pensylvanicus, Willd. ex Cham. & Schlecht. in Linnaea, ii. (1827) 227 = Claytonii.
perfoliatus, *Linn. Sp. Pl.* 126.—Reg. temp. bor. et austr.
petiolaris, Presl, Delic. Prag. 151 = natans.
petiolaris, *Rafin. in Med. Repos. N. York*, Ser. III. ii. (1811) 409 (Quid ?).—Am. bor.
petiolatus, Wolfg. ex Schult. Mant. iii. 352 = natans.
plantagineus, Du Croz, ex Roem. & Schult. Syst. iii. 504 = coloratus.
Plantago, Batard, Fl. Maine-et-Loire, 64 = natans.
polygonifolius, *Pourr. in Mém. Acad. Toul.* iii. (1788) 325.—Europ.
polygonus, *Cham. in Linnaea*, ii. (1827) 184. t. 4. f. 11.—Bras.
porcatus, Muhl. Cat. 18 = P. Spirillus.
praelongus, Bréb. Fl. Normand. ed. II. 250 = nitens.
praelongus, F. Muell. Fragm. viii. 217 = perfoliatus ?
praelongus, *Wulf. in Roem. Arch.* iii. III. (1805) 331.—Europ.
Proteus, Cham. & Schlecht. in Linnaea, ii. (1827) 196. t. 5. f. 16, a, b = lucens.
pulcher, *Tuckerm. in Am. Journ. Sc.* Ser. I. xlv. (1843) 38.—Am. bor.
pumilus, Wolfg. ex Schult. Mant. iii. 354 = Claytonii.
purpurascens, Seidl, ex J. & C. Presl, Fl. Čech. 37 = rufescens.
Purshii, Tuckerm. in Am. Journ. Sc. Ser. II. vi. (1848) 228 = Claytonii ?
pusillus, *Linn. Sp. Pl.* 127.—Reg. temp. bor. et austr.
reptans, *Humnicki, Cat. Pl. Luxeuil* (1876), 61.—Europ.
rigidus, *Wolfg. in Schult. Mant.* iii. 359.—Europ.
Robbinsii, *Oakes, in Hovey, Mag. Hortic.* vii. (1841) 180.—Am. bor.
rotundifolius, Schultz, Prod. Fl. Starg. Suppl. i. 9 = lucens.
Roxburghianus, Schult. Mant. iii. 367 = indicus.
rufescens, *Schrad. in Cham. Adn. ad Fl. Berol.* (1815) 4.—Reg. bor. temp.
rutilus, *Wolfg. in Schult. Mant.* iii. 362.—Lithuan.
salicifolius, *Wolfg. l. c.* 355.—Lithuan.
semipellucidus, Koch & Ziz, Cat. Pl. Palat. 5, 18 = rufescens.

POTAMOGETON :—
serotinus, Schrad. ex Schult. Mant. iii. 351 = natans.
serratus, *Huds. Fl. Angl.* ed. I. 61 = crispus.
serratus, Linn. Sp. Pl. 126 = densus.
serratus, Roth, Tent. Fl. Germ. i. 73 = rufescens.
serratus, Scop. Fl. Carn. ed. II. i. 117 = praelongus.
serratus, [Weber,] Wigg. Prim. Fl. Holsat. 16 = lucens.
serrulatus, Opiz, in Flora, v. (1822) 267 = crispus.
serrulatus, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 139.—Reg. Amur.
setaceus, Gilib. ex Schult. Mant. iii. 355 = divaricatus.
setaceus, Linn. Sp. Pl. 127 = densus.
setaceus, Pursh, Fl. Am. Sept. i. 120 = hybridus.
setaceus, Schum. Enum. Fl. Saell. i. 51 = pectinatus.
setaceus, Tausch, in Flora, xix. (1836) 423 = densus.
siculus, *Bercht. & Presl, Rostl.* i. *Alismac.* 21 (Quid ?).—Sicilia.
siculus, Tineo, ex Guss. Syn. Fl. Sic. ii. 790 = coloratus.
sparganiifolius, Bab. Man. Brit. Bot. ed. VI. 363 = natans.
sparganiifolius, Laestad. ex Fries, Nov. Fl. Suec. Mant. i. 9 = lucens.
spathaeformis, Tuckerm. ex A. Gray, Man. Bot. U. St. ed. V. 487 = gramineus ?
spathulatus, Auct. ex Nym. Consp. 681 = fluitans.
spathulatus, Schrad. ex Koch & Ziz, Cat. Pl. Palat. 5, 18.—German.
Spirillus, *Tuckerm. in Am. Journ. Sc.* Ser. II. vi. (1848) 228.—Am. bor.
striatus, *Ruiz & Pav. Fl. Per.* i. 70. t. 136. f. b.—Peruv.
strictus, Phil. Fl. Atac. 50 = pectinatus.
Sturrockii, Ar. Benn. in Scott. Natural. N. S. i. (1883) 27 = pusillus.
subflavus, *Loret & Barr. Fl. Montp.* 671.—Gallia.
subtrichodes, Schur, Enum. Pl. Transs. 633 = pusillus.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* 597.—Sumatra.
syriacus, *Cham. & Schlecht. in Linnaea*, ii. (1827) 227.—Syria.
tataricus, Less. l. c. ix. (1834) 206 = obtusifolius.
tenuicaulis, F. Muell. Fragm. i. 90, 244 = javanicus.
tenuifolius, *H. B. & K. Nov. Gen. et Sp.* i. 296.—N. Granat.
tenuifolius, Rafin. in Med. Repos. N. York, Ser. III. ii. (1811) 409 (Quid ?).—Am. bor.
tenuissimus, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 342 = pusillus.
Thunbergii, *Cham. & Schlecht. in Linnaea*, ii. (1827) 221.—Afr. austr.
trichoides, Benth. Pl. Hartw. 341; Schur, in Oestr. Bot. Zeitschr. xx. (1870) 281 = pusillus.
trichoides, *Cham. & Schlecht. in Linnaea*, ii. (1827) 175.—Europ.
trichoides, Tuckerm. in Am. Journ. Sc. Ser. II. vii. (1849) 358 = Tuckermani.
tuberculatus, Tenore & Guss. Fl. Neap. Syll. App. v. 4 = trichoides.
tuberosus, Roxb. Hort. Beng. 12; Fl. Ind. i. 452 = crispus.
Tuckermani, *Robbins, ex A. Gray, Fl. U. St.* ed. II. 434.—Am. bor.
uliginosus, Boenn. ex Cham. & Schlecht. in Linnaea, ii. (1827) 216 = natans.
undulatus, *Wolfg. in Schult. Mant.* iii. 360.—Lithuan.
upsaliensis, *Tiselius, in Bot. Notiser* (1884), 15.—Europ.
vaginalis, J. Fisch. Verz. Gefaesspfl. Neu-Vorpom. (1861) 40 = seq.
vaginatus, Turcz. in Bull. Soc. Nat. Mosc. (1838) I. 102; xxviii. (1854) II. 66 – pectinatus.
Vaillantii, Roem. & Schult. Syst. iii. 514 = pectinatus.
variifolius, Thore, Chlor. Land. 47 = natans.
Vaseyi, *Robbins, ex A. Gray, Fl. N. U. St.* ed. V. 485.—Am. bor.
volhynicus, Bess. ex Roem. & Schult. Syst. iii. 509, in obs. = lucens ?
Zetterstedtii, *Wallm. in Bot. Zeit.* i. (1843) 256.—Am. bor.
Zizii, Koch, ex Roth, Enum. Pl. Germ. i. 531 = lucens.
zosteraceus, Bab. Man. Brit. Bot. ed. I. 325 = pectinatus.
zosteraceus, *Fries, Nov. Fl. Suec.* ed. II. 51.—Europ. bor.
zosterifolius, *Schum. Enum. Pl. Saell.* i. 50.—Reg. bor. temp.
zosterophyllus, Dum. Fl. Belg. 164 = praec.

POTAMOGETON, Walt. Fl. Carol. 90 (1788) = **Myriophyllum**, Ponted. (Halor.).
pinnatum, Walt. l. c. = M. scabratum.
verticillatum, Walt. l. c. = M. heterophyllum.

POTAMOGETUM, [Clairv.] Man. Herb. 34, 44 (1811) = **Potamogeton**, Tourn. (Naiad.).

POTAMOGITON, Rafin. in Med. Repos. N. York, v. (1808) 354 = **Potamogeton**, Tourn. (Naiad.).

POTAMOPHILA, R. Br. Prod. 211 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1116.
MALTEBRUNIA, Kunth, Rév. Gram. i. 183. t. 3 (1830).
leersioides, *Benth. in Journ. Linn. Soc.* xix. (1881) 55.—Madag.
parviflora, *R. Br. Prod.* 211.—Austral.
prehensilis, *Benth. in Journ. Linn. Soc.* xix. (1881) 55.—Madag.

POTAMOPHILA, Schrank, Pl. Rar. Hort. Monac. ii. t. 63 (1819) = **Microtea**, Sw. (Phytolacc.).
parviflora, Schrank, l. c. = M. maypurensis.

POTAMOPITYS, Buxb. ex Adans. Fam. ii. 256 (1763) = **Elatine**, Linn.
Alsinastrum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = E. Alsinastrum.

POTAMOXYLON, Rafin. Sylva Tellur. 78 (1838) = **Tabebuia**, Gomez (Bignon.).
alba, Rafin. l. c. = T. fluviatilis.

POTANINIA, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 465. *ROSACEAE*.
mongolica, *Maxim. l. c.* 466.—As. or.

POTANISIA, Wawra in Sitzb. Akad. Wien, Math.-Nat. xxxviii. (1860) 566 (err. typ.) = POLANISIA, Rafin. = **Cleome**, Linn. (Cappar.).

POTENTILLA, Linn. Syst. ed. I (1735); Gen. ed. I. 147 (1737). *ROSACEAE*, Benth. & Hook. f. i. 620.
BOOTIA, Bigel. Fl. Bost. ed. II. 351 (1824).
CHAMAEPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371.
COELAS, Dulac, Fl. Hautes-Pyr. 303 (1867).
COMARUM, Linn. Gen. ed. I. 148 (1737).
DACTYLOPHYLLUM, Spenn. Fl. Frib. iii. 1034 (1829).
DRYADANTHE, Endl. Gen. 1242 (1840).
DRYMOCALLIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371.
DYNAMIDIUM, Fourr. l. c. (1868).
FRAGA, Lapeyr. Hist. Abr. Pl. Pyr. 287 (1813).
FRAGARIASTRUM, Schur, Enum. Pl. Tranns. 187 (1866).
HORKELIA, Cham. & Schlecht. in Linnaea, ii. (1827) 26.
HYPARGYRIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371.
IVESIA, Torr. & Gray, ex Torr. in Pacif. Rail. Rep. vi. 72 (1857).
JUSSIEA, Linn. Laches. Lapp. i. 284 (1811).
LEHMANNIA, Tratt. Ros. Monog. iv. 144 (1824).
PANCOVIA, Heist. ex Adans. Fam. ii. 294 (1763).
PENTAPHYLLUM, Gaertn. Fruct. i. 349. t. 73 (1788).
POTENTILLOPSIS, Opiz, in Lotos, vii. (1857) 30.
QUINQUEFOLIUM, Tourn. ex Adans. Fam. ii. 295 (1763).
SIBBALDIA, Linn. Sp. Pl. ed. I. 284 (1753).
TORMENTILLA, Linn. Gen. ed. I. 147 (1737).
TRICHOTHALAMUS, Lehm. in Nov. Act. Nat. Cur. x. (1821) 585. t. 49.
TRIDOPHYLLUM, Neck. Elem. ii. 93 (1790).
abbreviata, *Zimm. Gatt. Potentill.* 19.—Europ.
abnormis, *Lehm. Del. Sem. Hort. Hamb.* (1849) 11; *et in Pugill.* ix. 19.—Oriens.
absinthiifolia, Dougl. ex Hook. Fl. Bor. Am. i. 188 = pensylvanica.
abyssinica, A. Rich. Tent. Fl. Abyss. i. 257 = reptans.
acaulis, Linn. Sp. Pl. 500 = subacaulis.
acutifolia, Gilib. Fl. Lituan. ii. 253 = recta.

POTENTILLA:—
adenophylla, *Boiss. & Hohen. Diagn.* Ser. I. iii. 5.—Armenia.
adnata, Wall. ex Lehm. Del. Sem. Hort. Hamb. (1849) 7; et in Pugill. ix. 9 = Geum elatum.
adpressa, Opiz, Seznam, 79 = collina.
adscendens, Baumg. Enum. Stirp. Transs. ii. 62 = heptaphylla.
adscendens, Lapeyr. Hist. Abr. Pl. Pyr. 289 = aurea.
adscendens, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 554 = inclinata.
adscendens, Wolf. ex Reichb. Fl. Germ. Excurs. 593 = chrysantha.
adscendens, Zimm. Gatt. Potentill. 6 = *ascendens*.
affinis, Host, Fl. Austr. ii. 41 = aurea.
africarum, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid ?).—Hab.?
agrimonoides, Bieb. Fl. Taur. Cauc. i. 403 = sericea.
agrivaga, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1870) 46 = verna.
alba, Cav. & Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 228 = caulescens.
alba, *Linn. Sp. Pl.* 498.—Europ.; Reg. Caucas.
albescens, Opiz, Seznam, 79; et ex Zimm. Gatt. Potentill. 19 = collina.
albifolia, Wall. Cat. n. 1018 = micropetala.
albo-villosa, Schur, Enum. Pl. Transs. 191 = argentea.
alchemilloides, *Lapeyr. in Mém. Acad. Toul.* i. (1782) 210. t. 17.—Mont. Pyren.
alchemilloides, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 291 = heptaphylla.
alpestris, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 83; Boiss. Fl. Orient. ii. 719 = Bungei.
alpestris, Hall. f. in Ser. Mus. Helvet. i. (1818) 53 = aurea.
alpestris, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. ii. 357 = gelida.
alpicola, La Soie, in Bull. Soc. Murith. v. (1875) 18 = aurea ?
alpigena, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
alpina, Dalla Torre, Atl. Alpenfl. 91 = aurea ?
altaica, Bunge, in Ledeb. Fl. Atl. ii. 252 = multifida.
alterniflora, Rich. ex Nestl. Monog. Pot. 41 = recta.
alternifolia, Steud. Nom. ed. II. ii. 386, sphalm. = praec.
Amansiana, F. Schultz, Arch. Fl. 219, 228, 377 = verna.
ambigua, *Cambess. in Jacquem. Voy. Bot.* 51. t. 62.—Reg. Himal.
ambigua, Gaud. Fl. Helv. iii. 391 = aurea.
amphibola, Schur, Enum. Pl. Transs. 198 = heptaphylla.
amurensis, Maxim. Prim. Fl. Amur. 98 = supina.
ancistrifolia, *Bunge, in Ledeb. Fl. Alt.* ii. 99.—China.
ancistrifolia, Galeotti, ex Lehm. Del. Sem. Hort. Hamb. (1849) 6; et in Pugill. ix. 1 = Richardii.
andicola, *Benth. Pl. Hartw.* 172.—N. Granat.
anemonefolia, *Lehm. Ind. Sem. Hort. Hamb.* (1853) 9; *et in Hamb. Gartenz.* ix. (1853) 505.—Japon.
angustifolia, DC. Fl. Fr. Suppl. 540 = hirta.
angustifolia, Rafin. Fl. Tellur. ii. 56 = palustris.
angustifolia, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 296 = nivea.
anomala, Ledeb. Fl. Ross. ii. 53 = reptans.
Anserina, *Linn. Sp. Pl.* 495.—Reg. temp. bor. et austr.
anserinoides, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 123 = praec.
apennina, *Tenore, Cat. Hort. Neap.* (1815) 50; *et Fl. Nap.* i. 291. t. 46.—Mont. Apennin.
approximata, *Bunge, in Ledeb. Fl. Alt.* ii. 241.—Reg. Caucas.; Sibir.
arachnoidea, Dougl. ex Steud. Nom. ed. II. ii. 386 = pensylvanica.
Arbuscula, D. Don, Prod. Fl. Nep. 256 = fruticosa.
arctica, Hort. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 50 = arguta.
arenaria, Borck. ex Gaertn. Mey. & Scherb. Fl. Wett. ii. 248 = subacaulis.
argaea, *Boiss. Diagn.* Ser. II. vi. 70.—As. Min.
argentata, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = seq.
argentea, *Linn. Sp. Pl.* 497.—Reg. bor. temp.
argentea, Willd. ex Ledeb. Fl. Ross. ii. 48 = dealbata.
argenteaeformis, *Kauffm. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 313.—Ross. med.

POTENTILLA:—
argentina, Huds. Fl. Angl. ed. I. 195 = P. Anserina.
arguta, Lehm. Pugill. iii. 25 = pensylvanica, glandulosa.
arguta, Nutt. in Journ. Acad. Philad. vii. (1834) 21 = glandulosa.
arguta, *Pursh, Fl. Am. Sept.* ii. 636.—Am. bor.
argyroloma, *Boiss. & Hohen. Diagn.* Ser. I. x. 6.—Persia.
argyrophylla, *Wall. Cat.* n. 1020.—Reg. Himal.
ascendens ×, Gremli, Beitr. Fl. Schweiz, 68.—Europ.
asperrima, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 609.—Sibir. altaic.
assurgens, Vill. Fl. Delph. 51 = inclinata.
astracanica, Jacq. Misc. ii. 349 = recta.
astragalifolia, Bunge, in Ledeb. Fl. Alt. ii. 246 = bifurca.
Atropos, Hoffmeist. Briefe aus Indien, App. ii. 364, nomen (sp. dub.).—Reg. Himal.
atropurpurea, De Vriese, Tuinb. Fl. i. (1855) 268, sphalm. = seq.
atrosanguinea, Lodd. Bot. Cab. t. 786 = argyrophylla.
aurea, *Linn. Cent. Pl.* ii. 18; *Amoen. Acad.* iv. 316.—Europ.
aurigena, Kern. ex Zimm. Gatt. Potentill. 22 = verna.
aurulenta, *Gremli, Fl. Schweiz*, ed. I. p. xii.—Helvet.
australis, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = hirta.
australis, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1870) 47 = verna.
autaretica, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = heptaphylla.
autumnalis, Opiz, ex Zimm. Gatt. Potentill. 19 (Quid ?).—Europ.
axilliflora, *Hook. f. Fl. Brit. Ind.* ii. 346.—Reg. Himal.
baldensis, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 220, nomen = aurea.
bannehalensis, Cambess. in Jacquem. Voy. Bot. 52 = Leschenaultiana.
barbata, Wall. Cat. n. 1030 = Mooniana.
Bauhini, Hort. ex Schlecht. in Linnaea, ii. (1827) 26 = heptaphylla.
Baumgarteniana, Schur, Enum. Pl. Transs. 191 = inclinata.
bellunensis, Huter & Porta, ex Zimm. Gatt. Potentill. 23 = aurea.
Benicskyi, Frivald. in Flora, xix. (1836) 437 = rupestris.
Benyctskyi, Steud. Nom. ed. II. ii. 386 = praec.
betonicaefolia, Poir. Encyc. v. 601 = nivea.
bicolor, Lindl. Bot. Reg. (1845) t. 62 = nepalensis.
bidens, Bertol. Misc. Bot. xxiv. 16 = bifurca.
Biebersteiniana, Tratt. Ros. Monog. iv. n. 24 = pensylvanica.
bifida, Pall. Reise, i. 215; ii. 523 = bifurca.
biflora, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 297.—Reg. bor. temp. et arct.
bifolia, Walp. Ann. ii. 470, sphalm. = praec.
bifurca, *Linn. Sp. Pl.* 497.—Reg. Caucas.; As. bor.
bifurcata, Poir. Encyc. v. 587 = praec.
bifurcata, Wall. ex Hook. f. Fl. Brit. Ind. ii. 355 = nepalensis.
Bigeloviana, Wender. in Linnaea, xvi. (1842) Litt. 112 = arguta.
Billotii, Boulay, in Billotia, i. (1869) 111 = verna.
bipinnatifida, Dougl. ex Hook. Fl. Bor. Am. i. 188 = pensylvanica.
biserrata, Schur, Enum. Pl. Transs. 198 = heptaphylla.
bithynica, Hornem. Hort. Hafn. Suppl. 55 = aurea.
Blaschkeana, Turcz. ex Lehm. Ind. Sem. Hort. Hamb. (1853) 9; et in Hamb. Gartenz. ix. (1853) 506 = gracilis.
Bocconi, Nestl. Monog. Pot. 73. t. 10. f. 2 = apennina.
Bogenhardiana, F. Schultz, in Flora, xlv. (1862) 461 = P. Fragariastrum.
bolzanensis, Zimm. Gatt. Potentill. 21 = verna.
Bouquojana, Nym. Consp. 225 (*Buquoyana*) = heptaphylla.
brachyloba, Borb. ex Zimm. Gatt. Potentill. 11 = collina.
brachypetala, *Fisch. & Mey. ex Lehm. Del. Sem. Hort. Hamb.* (1849) 9.—Reg. Caucas.
Brauniana, Hoppe, Bot. Taschenb. (1800) 137 = minima.
Breisnii, Hort. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 49 = glandulosa.

POTENTILLA:—
brevifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 442.—Am. bor.
breviscapa, Vest, in Flora, iv. (1821) 158 = micrantha.
breviscissa, Bertol. Misc. Bot. xxiv. 17 = multifida.
Breweri, *S. Wats. in Proc. Am. Acad.* viii. (1873) 555. —Calif.
brutia, Tenore, ex Nym. Consp. 224 = argentea.
Buccoana, *Clem. in Mem. Acc. Sc. Torin*, Ser. II. xvi. (1857) 275.—Bithynia.
Buguoyana, Knaf, in Flora, xxx. (1847) 181 = heptaphylla.
Bungei, *Boiss. Fl. Orient.* ii. 718.—Persia.
caespitosa, Lehm. Del. Sem. Hort. Hamb. (1849) 10; et in Pugill. ix. 17 = nivea.
calabra, Tenore, Prod. Fl. Nap. Suppl. ii. 47 = argentea.
calycina, *Boiss. & Bal. Diagn.* Ser. II. v. 94.—Kurdist.
calycina, Ehrenb. ex Boiss. Fl. Orient. ii. 704 = libanotica.
camonia, Rota, Prosp. Fl. Bergamo, 99 (Quid?).—Europ. austr.
campestris, Nutt. ex Torr. & Gray, Fl. N. Am. i. 439 = dissecta.
campicola, Jord. ex Verlot, Cat. Pl. Jard. Grenoble, (1856) 28 = aurea.
cana, Jord. l. c. = argentea.
cana, Wall. ex Hook. f. Fl. Brit. Ind. ii. 359 = supina.
canadensis, *Linn. Sp. Pl.* 498.—Am. bor.
candicans, Fisch. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 60 = pensylvanica.
candicans, *H. B. & K. Nov. Gen. et Sp.* vi. 216.—Am. austr.
candicans, Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 285 = P. Anserina?
Candolleana, Royle, Illustr. Bot. Himal. 203 = argyrophylla.
Candollei, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* iii. (1875–76) 563.—Hab.?
canescens, Bess. Fl. Galic. i. 330 = inclinata.
canescens, Hartm. Scand. Fl. ed. I. 200 = collina.
canescens, Spreng. Syst. ii. 537, ex parte = humifusa.
cappadocica, *Boiss. Diagn.* Ser. I. iii. 6.—As. Min.
cardiopetala, Bess. ex DC. Prod. ii. 579 = recta.
carniolica, *Kern. in Oestr. Bot. Zeitschr.* xx. (1870) 44.—Austria.
caroliniana, Poir. Encyc. v. 595 = canadensis.
cataclines, Lehm. Pugill. iii. 37 = argyrophylla.
caulescens, *Linn. Cent. Pl.* ii. 19; *Amoen. Acad.* iv. 317.—Europ. austr.
caulescens, Scop. Fl. Carn. ed. II. i. 361 = Clusiana.
Cautleyana, Royle, Illustr. Bot. Himal. 207 = argyrophylla.
centigrana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1874) 163.—Japon.
Charpentieri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 60.—Europ.
Chaubardiana, Timb. in Mém. Acad. Toul. Sér. IV. vi. (1856) 141 = verna.
chinensis, *Ser. in DC. Prod.* ii. 581.—China.
chrysantha, Griseb. Spicil. Fl. Rumel. i. 98 = hirta.
chrysantha, Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 78 = gracilis?
chrysantha, Reichb. Fl. Germ. Excurs. 593 = heptaphylla.
chrysantha, *Trevir. Ind. Sem. Hort. Vratisl.* (1818) 5; *Reichb. Pl. Crit.* vi. 21. f. 808.—Europ.
chrysocraspeda, Lehm. Del. Sem. Hort. Hamb. (1849) 10; *Pugill.* ix. 15 = aurea.
chrysopetala, Bess. ex DC. Prod. ii. 578 = hirta.
cicutariaefolia, Willd. Sp. Pl. ii. 1098 = supina.
cilicia, Boiss. Diagn. Ser. I. iii. 7 = Kotschyana.
cinerascens, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iii. (1863) 421 = sericea.
cinerea, Chaix, ex Vill. Prosp. 46; Hist. Pl. Dauph. iii. 567 = subacaulis.
cinerea, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 296; Rafin. ex DC. Prod. ii. 577 = argentea.
Clarkei, *Hook. f. Fl. Brit. Ind.* ii. 351.—Reg. Himal.
Clarkiana, Kellogg, in Proc. Calif. Acad. vii. (1877) 94 = Grayi?
Clementi, Jord. Pugill. 70 = subacaulis.
clivicola, Jord. Cat. Jard. Grenoble (1856) = aurea.
Clusiana, Geners. Elench. Scepus. n. 277; ex Wahlenb. Fl. Carpat. 157 = alba.
Clusiana, *Jacq. Fl. Austr.* ii. 10. t. 116.—Austria.

POTENTILLA:—
coccinea, Hoffmeist. Briefe aus Indien, App. ii. 364, nomen = nepalensis.
Collettiana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 53.—Afghan.
collina, Gren. & Godr. Fl. Fr. i. 533 = argentea.
collina, *Wibel, Prim. Fl. Werth.* ii. 267.—Europ.; As. Min.
colorata, Lehm. Sem. Hort. Hamb. (1821) 8 = nepalensis.
comariformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = palustris.
comaroides, *Humb. & Bonpl. ex Nestl. Monog. Pot.* 62. t. 4. f. 3.—Mexic.
comaroides, Steph. ex Ledeb. Fl. Ross. ii. 50 = stipularis.
Comarum, Nestl. Monog. Pot. 36 = palustris.
commutata, Lehm. Pugill. iii. 16 = microphylla.
concinna, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 411 = gracilis.
concinna, Richards. in Frankl. Journ. ed. II. App. 20 = humifusa.
conferta, Bunge, in Ledeb. Fl. Alt. ii. 240 = pensylvanica.
confertiflora, Torr. Fl. U. St. i. 499 = arguta.
confinis, Jord. Pugill. 68 = argentea.
congesta, *Baill. Hist. Pl.* i. 369.—Calif.
cordata, Schrank, Prim. Fl. Salisb. 130 = alba.
coriandrifolia, *D. Don, Prod. Fl. Nep.* 232.—Reg. Himal.
corsica, *Sieber, ex Lehm. Del. Sem. Hort. Hamb.* (1849) 9 = crassinervia.
corsica, *Soleir. ex Lehm. l. c.* 7; *et Pugill.* ix. 8.—Corsica.
corymbosa, Moench, Meth. Suppl. 279 = recta.
corymbosa, Pourr. in Mém. Acad. Toul. iii. (1788) 325 = caulescens.
coryphea, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
crassa, Tausch, ex Opiz, Boehm. Gew. 63, nomen = recta.
crassa, Uechtr. ex Zimm. Gatt. Potentill. 9 = incrassata.
crassinervia, *Vis. App. Fl. Cors. Prod.* 2.—Corsic.; Sardin.
crinita, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 41.—Am. bor. occ.
crocea, Hall. f. in Schleich. Cat. (1807) 20 = aurea.
cryptantha, Klotzsch, in Bot. Ergeb. Waldem. Reise, 156 = monanthes.
Cryptotaeniae, *Maxim. in Bull. Acad. Pétersb.* xviii. (1874) 162.—Japon.
cuneata, Baill. Hist. Pl. i. 369 = multijuga.
cuneata, Wall. Cat. n. 1015 = ambigua.
cuneifolia, Bertol. Misc. Bot. xxiv. 15 = ambigua.
curvidens, Schur, Enum. Pl. Transs. 190 = inclinata.
curviseta, *Hook. f. Fl. Brit. Ind.* ii. 358.—Reg. Himal.
dacica, Borb. ex Zimm. Gatt. Potentill. 5 = P. Tormentilla.
dahurica, G. Don, Gen. Syst. ii. 561 = *davurica*.
dasyantha, Hort. Genev. (1848); ex Lehm. Del. Sem. Hort. Hamb. (1849) 12 = aurea.
dasyphylla, Bunge, in Ledeb. Fl. Alt. ii. 243 = sericea.
davurica, Nestl. Monog. Pot. 31. t. 1 [bis] = fruticosa.
dealbata, *Bunge, in Ledeb. Fl. Alt.* ii. 250.—Sibir.
dealbata, Dougl. ex Hook. Fl. Bor. Am. i. 188 = Hippiana.
debilis, Schleich. Cat. Pl. Helv. ed. III. 23 = aurea.
decipiens, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = argentea.
declinata, *Hornem. Hort. Hafn. Suppl.* 56.—Hab.?
decumbens, Jord. Cat. Jard. Grenoble (1849), 22 = argentea.
delphinensis, *Gren. & Godr. Fl. Fr.* i. 530.—Gallia.
demissa, Jord. Cat. Jard. Grenoble (1849), 22 = argentea.
dentata, Forsk. Fl. Aegypt. Arab. 98 = viscosa.
denticulata, Wall. Cat. n. 1029 = supina.
denticulosa, Ser. in DC. Prod. ii. 573 = Fragaria indica.
deorum, *Boiss. & Heldr. Diagn.* Ser. II. ii. 51.—Thessal.
depauperata, *Engelm. ex A. Gray, in Proc. Am. Acad.* vii. (1868) 399.—Am. bor.
depressa, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 289 = aurea.
desertorum, *Bunge, in Ledeb. Fl. Alt.* ii. 257.—Reg. Himal.; Soongar.

POTENTILLA:—
Dethomasii, Reichb. Fl. Germ. Excurs. 595 = Detommasii.
Detommasii, Schur, Enum. Pl. Transs. 191 = argentea.
Detommasii, *Tenore, Prod. Fl. Nap. Suppl.* i. 61.—Europ. austr.
dichotoma, *Galeotti, ex Lehm. in Nov. Act. Nat. Cur.* xxiii. *Suppl.* (1856) 204 (sp. dub.).—Mexic.
dichotoma, Moench, Meth. 659 = norvegica.
Dickinsii, *Franch. & Sav. Enum. Pl. Jap.* ii. 337.—Japon.
diffusa, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 41 = Hippiana.
diffusa, Willd. Enum. Hort. Berol. 555 = norvegica.
digitata, Rafin. Fl. Tellur. ii. 55 = palustris.
digitato-flabellata, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1851) 14 = gracilis.
digitato-flabellata, Heidenr. in Oestr. Bot. Zeitschr. xxi. (1871) 169 = norvegica.
discolor, *Bunge, in Ledeb. Fl. Alt.* ii. 99.—China.
discolor, Cambess. in Jacquem. Voy. Bot. 53 = Salesoviana.
dissecta, *Pursh, Fl. Am. Sept.* i. 355.—Am. bor.
divaricata, DC. Fl. Fr. Suppl. 541 = recta.
divergens, Nym. Consp. 227 = P. Tormentilla.
diversifolia, Lehm. Pugill. ii. 9 = dissecta.
Dombeyi, *Nestl. Monog. Pot.* 38. t. 5. f. 2.—Chili.
Doubjouneana, *Cambess. in Jacquem. Voy. Bot.* 50.—Reg. Himal.
Drummondii, Lehm. Pugill. ii. 9 = dissecta.
Drusiana, De Vriese, Tuinb. Fl. i. (1855) 268. t. 12 = argyrophylla.
dryophylla, Pall. ex Ledeb. Fl. Ross. ii. 58 = nivea.
dubia, Moench, Enum. Pl. Hass. 263; Suter, Fl. Helv. i. 308 = aurea.
Durandii, Torr. & Gray, Fl. N. Am. i. 444 = Fragaria indica.
effusa, *Dougl. ex Lehm. Pugill.* ii. 8.—Am. bor. occ.
Egedii, Wormsk. in Fl. Dan. t. 1578 = P. Anserina.
Ehrenbergiana, *Schlecht. in Linnaea*, xiii. (1839) 261. —Mexic.
elata, Salisb. Prod. 362 = fruticosa.
elatior, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 295.—As. Min.; Reg. Caucas.
elegans, *Schlecht. in Linnaea*, ii. (1827) 22.—Kamtchat.
ellwindensis, Boiss. Diagn. Ser. I. vi. 51 = elvendensis.
elongata, Bess. ex Spreng. Syst. iv. Cur. Post. 199 = virgata.
elongata, Goldb. ex DC. Prod. ii. 577 = intermedia.
elvendensis, *Boiss. Fl. Orient.* ii. 709.—Persia.
emarginata, Desf. Tabl. ed. I. 177 = P. Fagiastrum.
emarginata, *Pursh, Fl. Am. Sept.* i. 353.—Am. bor. et arct.
engadinensis ×, Bruegg. in Cat. Bot. Gart. Zuerich (1868), 4.—Europ.
erecta, Malmberg, Förteckn. Karelska näs Kärlv. 314 = seq.?
erecta, Uspenski, ex Ledeb. Fl. Ross. ii. 45 = recta.
eriocarpa, *Wall. Cat.* n. 1012.—Reg. Himal.
erodora, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
erythrocaulon, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid?).—Hab.?
Ewersmanniana, *Fisch. ex Claus, Ind. Goebel, Reise*, ii. 272.—Ross.
exaltata, Bunge, Enum. Pl. Chin. Bor. 24 = chinensis.
explanata, Zimm. Gatt. Potentill. 20 = aurea.
fagineicola, Lamotte, Prod. Fl. Centr. Fr. i. 241 = aurea.
fallax, Rochel, Bot. Reise Banat. 71 = chrysantha.
fallax, Mor. ex Zimm. Gatt. Potentill. 5 = P. Tormentilla.
fastigiata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 440 = gracilis.
Fenzlii, Lehm. Del. Sem. Hort. Hamb. (1849) 8; et in Pugill. ix. 12 = Meyeri.
ferruginea, Dougl. ex Hook. Fl. Bor. Am. i. 186; Paxt. Mag. Bot. v. (1838) t. 223 = arguta.
filiformis, Vill. Hist. Pl. Dauph. iii. 564 = aurea.
filipendula, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 616 = tanacetifolia.
filipendula, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 286.—Dahur.
Fintelmanni ×, E. Otto, Hamb. Gartenzeit. v. (1849) 52.
fissa, Nutt. in Torr. & Gray, Fl. N. Am. i. 446 = glandulosa.

POTENTILLA:—
fissidens, *Borb. ex Zimm. Gatt. Potentill.* 9.—Europ.
flabellata, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 24.—Turkestan.
flabellifolia, Hook. ex Torr. & Gray, Fl. N. Am. i. 442 = gelida.
flabelliformis, Lehm. Pugill. ii. 12 = gracilis.
flagellaris, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 291.—Sibir.
floribunda, Pursh, Fl. Am. Sept. i. 355 = fruticosa.
Fontolii, Colla, Herb. Pedem. ii. 326 = grammopetala.
formosa, D. Don, Prod. Fl. Nep. 232 = nepalensis.
formosana, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 212 = discolor.
Fragaria, Láng, ex Tenore, Syll. Fl. Neap. 249; Hohen. Enum. Elisabethpol. 236 = micrantha.
Fragaria, Poir. Encyc. v. 599 = P. Fragariastrum.
fragariaefolia, C. C, Gmel. Fl. Bad. ii. 452 = P. Fragariastrum.
fragariaefolia, Hoppe, ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 196 = norvegica.
fragariaefolia, Klotzsch, ex Lehm. l. c. 176 = Fragaria indica.
fragariaefolia, Less. ex Ledeb. Fl. Ross. ii. 57 = nivea.
Fragariastrum, *Ehrh. ex Hall. f. in Ser. Mus. Helv.* i. (1818) 49.—Europ.
fragarifolia, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 203 = *fragariaefolia*.
fragariformis, *A. Gray, in Perry, Narr. Exp. Jap.* ii. 310.—Japon.
fragarioides, Habl. Desc. Phys. Taur. 146 = geoides.
fragarioides, *Linn. Sp. Pl.* 496.—As. bor.; Reg. Himal.
fragarioides, Poir. Encyc. v. 586 = rupestris.
fragarioides, Schlecht. & Cham. in Linnaea, ii. (1827) 25 = japonica.
fragarioides, Vill. Hist. Pl. Dauph. iii. 561 = P. Fragariastrum.
fragarioides, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 285 = Sprengeliana.
fragiformis, Willd. ex Schlecht. l. c. 294 = gelida.
fraterna, Wallr. in Linnaea, xiv. (1840) 578 = P. Fragariastrum.
Friesiana, *Lange, in Fl. Dan.* t. 2965.—Groenl.
frigida, Grev. in Mem. Wern. Soc. iii. (1821) 430 = emarginata.
frigida, Schiede, ex Steud. Nom. ed. II. ii. 387 = leptopetala.
frigida, *Vill. Hist. Pl. Dauph.* iii. 563.—Europ.; Am. bor.
fructicosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 580 = fruticosa.
frutescens, *Radde, Ber. Reis. Sued-ost. Sibir.* 94.—Sibir.
fruticosa, *Linn. Sp. Pl.* 495.—Reg. bor. temp.
fulgens, *Wall. ex Hook. Bot. Mag.* t. 2700.—Reg. Himal.
fusca, Schlecht. in Linnaea, xiii. (1839) 262 = haematochrus.
galliciana, Bess. Prim. Fl. Gall. i. 333 = aurea.
garipensis, E. Mey. in Drège, Zwei Pfl. Docum. 213; et ex Boiss. Fl. Orient. ii. 726 = supina.
Gaudini, Gremli, Excurs. Schw. (1874) 171 = subacaulis.
gelida, Fries, ex Nym. Consp. 225 = aurea.
gelida, *C. A. Mey. Verz. Pfl. Cauc.* 167.—Europ. et As. bor.
geminiflora, Schrank, in Syll. Ratisb. i. (1824) 216 = norvegica.
gentilis, Jord. Pugill. Pl. Nov. 69 = heptaphylla.
geoides, *Bieb. Fl. Taur. Cauc.* i. 404.—Tauria.
geranioides, *Willd. Sp. Pl.* ii. 1101.—Armen.; Syria.
geranoides, Schleich. Cat. Helv. (1807) 20 = aurea.
Gerardiana, Lindl. in Wall. Cat. n. 1023 = fragarioides.
glabra, Lodd. Bot. Cab. t. 914 = fruticosa.
glabrata, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 285 = fruticosa.
glacialis, Hall. f. in Ser. Mus. Helv. i. (1818) 51, 69 = frigida.
glacialis, Pourr. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 137 = nivalis.
glandulifera, Kras. ex Nym. Consp. 226 = aurea.
glandulosa, Boulay, in Billotia, i. (1864) 12 = verna.
glandulosa, Kras. in Oestr. Bot. Zeitschr. xvii. (1867) 303 = aurea.
glandulosa, *Lindl. Bot. Reg.* t. 1583.—Am. bor. occ.

POTENTILLA:—
glauca, Cambess. in Jacquem. Voy. Bot. 54 = bifurca.
glauca, Moris, Stirp. Sard. Elench. 18 = crassinervia.
glaucescens, Willd. ex Schlecht. Ges. Naturf. Fr. Berl. Mag. vii. (1813) 289 = subacaulis.
glaucophylla, Lehm. Del. Sem. Hort. Hamb. (1836) 7; Cf. Linnaea, xii. (1838) Litt. 83 = dissecta.
glutinosa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 446 = glandulosa.
glutinosa, Pursh, ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 50 = arguta.
Goldbachii, *Rupr. Fl. Ingr.* 319.—Reg. Caucas.
Gordoni, *Baill. Hist. Pl.* i. 369.—Am. bor.
gossypina, Nutt. ex Hook. Lond. Journ. Bot. vi. (1847) 219 = effusa.
gracilis, *Dougl. in Hook. Bot. Mag.* t. 2984.—Am. bor. occ.
gracilis, Wall. ex Hook. f. Fl. Brit. Ind. ii. 359 = Kleiniana.
Grahamiana, Wight & Arn. Prod. 301 = Leschenaultiana.
grammopetala, *Moretti, in Brugnat. Giorn. Fis.* ii. IX. (1826) 67.—Italia.
grandiflora, Bieb. Fl. Taur. Cauc. i. 409 = gelida.
grandiflora, *Linn. Sp. Pl.* 499.—Europ.; As. bor.
grandiflora, Scop. Fl. Carn. ed. II. i. 363 = aurea.
grandiflora, Thunb. Fl. Jap. 219 = japonica.
Grayi, *S. Wats. in Proc. Am. Acad.* viii. (1873) 557.—Calif.
Gremlii, Zimm. Gat. Potentill. 6 = *ascendens*.
Grenieri, F. Schultz, ex Nym. Consp. 224 = argentea.
Grenieriana, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = heptaphylla.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 351.—Reg. Himal.
groenlandica, R. Br. in Ross's Voy. ed. II. ii. 193 = emarginata.
grossa, Dougl. ex Hook. Fl. Bor. Am. i. 193 = norvegica.
Guentheri, Pohl, Tent. Fl. Bohem. ii. 185 = collina.
Gulielmi-Waldemarii, Klotzsch, Bot. Ergeb. Waldem. Reise, t. 13 = nepalensis.
Gunneri, *Hartm. Skand. Fl.* ed. I. 201.—Norveg.
haematochrus, *Lehm. Ind. Sem. Hort. Hamb.* (1836). —*Cf. Linnaea*, xii. (1838) *Litt.* 83.—Mexic.
Halleri, Ser. in Mus. Helv. i. (1818) 75 = aurea.
Halleriana, Tratt. Ros. Monog. iv. n. 86, in obs. = heptaphylla.
Haynaldiana, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 176.—Macedon.
Heerii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 61.—Europ.
Hegetschweileri ×, Bruegg. l. c. 59.—Europ.
Heidenreichii, Zimm. Gatt. Potentill. 10 = norvegica.
helvetica, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = heptaphylla.
helvetica, Schleich. Cat. Helv. (1797) 20 = frigida.
heptaphylla, Linn. Cent. Pl. ii. 13 = aurea.
heptaphylla, *Mill. Gard. Dict.* ed. VIII. n. 9.—Europ.; Oriens.
heterophylla, Lapeyr. Hist. Abr. Pl. Pyr. 289 = aurea.
Heuffeliana, Steud. Nom. ed. II. ii. 387 = heptaphylla.
Heynii, Roth, Nov. Pl. Sp. 235 = supina.
hiemalis, Cham. & Schlecht. in Linnaea, v. (1830) 572 = leptopetala.
Hippiana, *Lehm. Pugill.* ii. 7.—Am. bor. occ.
hirsuta, Michx. Fl. Bor. Am. i. 303 = norvegica.
hirsuta, Vahl, ex Hornem. in Fl. Dan. t. 1390 = nivea.
hirta, All. Fl. Pedem. iii. t. 71 = recta.
hirta, Haenke, in Abh. Boehm. Ges. Wiss. (1787) 112 = inclinata.
hirta, *Linn. Sp. Pl.* 497.—Europ. austr.; Oriens.
hirta, Schur, Enum. Pl. Transs. 192 = patula.
hispanica, *Zimm. Gatt. Potentill.* 7.—Hispan.
hispida, Nestl. Monog. Pot. 36 = viscosa.
hispida, Willd. Enum. Hort. Berol. 553 = pensylvanica.
hololeuca, Boiss. ex Lehm. Del. Sem. Hort. Hamb. (1849) 8; Pugill. ix. 9 = multifida, sericea.
holopetala, Walp. Ann. ii. 485 = *olopetala*.
holosericea, *Griseb. Spicil. Fl. Rumel.* i. 99.—Macedon.
holosericea, Nutt. ex Torr. & Gray, Fl. N. Am. i. 438 = sericea.
Hookeriana, *Lehm. Del. Sem. Hort. Hamb.* (1849) 10; *Pugill.* ix. 18.—Am. bor.
Hopwoodiana ×, Sweet, Flow. Gard. Ser. II. t. 61.
Humboldtiana, Tratt. Ros. Monog. iv. 41 = candicans.

POTENTILLA:—
humifusa, *Nutt. Gen. Am.* i. 310.—Am. bor. occ.
humifusa, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 289 = aurea.
hungarica, Schlecht. l. c. = inclinata.
hybrida, Wallr. Sched. Crit. 247 = P. Fragariastrum.
hypoleuca, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 619.—Mongolia.
ignescens, Hort. ex Steud. Nom. ed. V. ii. 388 = argyrophylla.
imbricata, Kar. & Kir. in Bull. Soc. Nat. Mosc. xiv. (1841) 416 = bifurca.
impolita, Wahlenb. Fl. Carpat. 155 = argentea.
inaperta, Jord. Cat. Jard. Grenoble (1849), 23 = collina.
incana, Gaertn. Mey. & Scherb. Fl. Wett. ii. 248 = subacaulis.
incanescens, Bess. ex Lindem. in Bull. Soc. Nat. Mosc. lv. (1880) i. 298 = inclinata.
incanescens, Opiz, Natural. ix. (1825) 136 = argentea.
incerta, Vill. Prosp. 46 (Quid ?).—Gallia.
incisa, Desf. Tabl. ed. I. 177; Nestl. Monog. Pot. 51 = aurea.
incisa, Lag. Gen. et Sp. Nov. 16 = Lagascana.
incisa, Lindl. ex Loud. Hort. Brit. Suppl. ii. 660 = glandulosa.
inclinata, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 69 = aurea.
inclinata, J. & C. Presl, Fl. Čech. 106 = collina.
inclinata, *Vill. Hist. Pl. Delph.* iii. 567. t. 45.—Europ.; As. bor.
incrassata, *Zimm. Gatt. Potentill.* 9.—Europ.
Inglisii, Royle, Illustr. Bot. Himal. 207. t. 41. f. 3 = fruticosa.
inquinans, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 624 = rupestris.
insignis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 285 = mollissima.
insignis, Royle, ex Lindl. Bot. Reg. (1841) t. 37 = argyrophylla.
integrifolia, Lapeyr. Hist. Abr. Pl. Pyr. 291 = nivalis.
intermedia, *Linn. Mant.* i. 76.—Europ.
intermedia, Nestl. Monog. Pot. 50 = heptaphylla.
intermedia, Poir. Encyc. v. 170 = aurea.
intermedia, Roth, Tent. Fl. Germ. i. 223 = recta.
intermedia ×, Tratt. Ros. Monog. iv. n. 58.
intermedia, Wahlenb. Fl. Carpat. i. 154 = inclinata.
italica, Lehm. Del. Sem. Hort. Hamb. (1849) 11; Pugill. ix. 20 = procumbens.
Jacquemontiana, Cambess. in Jacquem. Voy. Bot. 51 = argyrophylla.
Jamesoniana, Grev. in Mem. Wern. Soc. iii. (1821) 416 = nivea.
Jankeana, *Pantoč. in Oestr. Bot. Zeitschr.* xxiii. (1873) 5.—Montenegr.
japonica, *Blume, Bijdr.* 1105.—Japon.
Johanniniana, Goiran, Morph. Veg. Veron. 45. t. 3 = argentea.
jurana, Reut. in Comp.-Rend. Soc. Hallér. (1854-56) 108 = aurea.
juratensis, Ser. ex Nestl. Monog. Pot. 52 = aurea.
kashmirica, *Hook. f. Fl. Brit. Ind.* ii. 355.—Reg. Himal.
Keilhavii, Sommerf. in Mag. Naturvid. II. i. (1832) 244 = pulchella.
Kerneri ×, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 391.—Europ.
Kleiniana, *Wight & Arn. Prod.* 300.—Ind. or.
Kotschyana, *Fenzl, Pugill. Pl. Nov. Syr.* 6.—As. Min.
Kristofiana, *Zimm. Gatt. Potentill.* 29.—Austria.
kurdica, *Boiss. & Hohen. Diagn.* Ser. I. iii. 4.—Kurdist.
labradorica, Lehm. Del. Sem. Hort. Hamb. (1849) 12; Pugill. ix. 21 = norvegica.
laciniosa, Waldst. & Kit. ex Nestl. Monog. Pot. 45 = recta.
laeta, Reichb. Fl. Germ. Excurs. 595 = hirta.
laeta, Salisb. Prod. 362 = P. Tormentilla.
laevigata, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
Lagascana, *Ser. in DC. Prod.* ii. 586.—Hab. ?
lanata, Lam. Fl. Fr. iii. 646 = nivalis.
lanuginosa, Boulay, in Billotia, i. (1864) 12 = verna.
lanuginosa, *Fisch. & Sweet, ex Steud. Nom.* ed. II. ii. 388.—Reg. Caucas.

POTENTILLA :—

laxa, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 289 = aurea.
lazica, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 716.—As. Min.
Lehmanniana, *Rupr. Sert. Tiansch.* 45.—Turkestan.
lejocarpa, *Vis. & Panč. in Mem. Ist. Venet.* x. (1861) 431. t. 2.—Serbia.
lepida, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
leptopetala, *Lehm. Del. Sem. Hort. Hamb.* (1830) 8 ; *Pugill.* iii. 32.—Mexic.
Leschenaultiana, *Ser. in DC. Prod.* ii. 584.—Ind. or.
leucantha, Gueldenst. Reisen, i. 192 = rupestris.
leucochroa, Lindl. in Wall. Cat. n. 1019 = argyrophylla.
leuconota, *D. Don, Prod. Fl. Nep.* 230.—Reg. Himal.
leuconota, Wall. Cat. n. 1021, partim = pedunculares.
leucophylla, Pall. Reise, ii. 194 = nivea.
leucophylla, Torr. in Ann. Lyc. N. York, ii. (1828) 197 = Hippiana.
leucopolitana, P. J. Muell. ex F. Schultz, Arch. Fl. 272 = collina.
libanotica, *Boiss. & Sprun. Diagn.* Ser. I. iii. 4.—Syria.
lignosa, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 293.—Persia.
limosa, Boenn. ex Schur, Enum. Pl. Transs. 189 = supina.
Lindackeri, Tausch, in Flora, ii. (1819) 466 = collina.
Lindenbergii, Lehm. in Hamb. Gartenz. vii. (1851) 339 = fruticosa.
linealoba, [Moç. & Sesse,] ex Ser. in DC. Prod. ii. 582 = candicans.
lineata, Trevir. in Spreng. Syst. ii. 534 = fulgens.
Loddigesii, Spreng. Syst. iv. Cur. Post. 199 = inclinata.
longifolia, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 287 = viscosa.
longipes, *Ledeb. Fl. Ross.* ii. 50.—Rossia.
Loureironis, Tratt. Ros. Monog. iv. n. 4 = fruticosa.
lucida, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 296 = gelida.
lupinoides, Willd. Sp. Pl. ii. 1107 = nivalis.
lycopodioides, Baill. ex S. Wats. in Bibl. Ind. N. Am. Bot. 288 = Gordoni.
Mackayana ×, Sweet, Brit. Flow. Gard. Ser. II. t. 43.
Macnabiana ×, Lem. in Fl. des Serres, ii. (1846) t. 149.
macrantha, Ledeb. in Mém. Acad. Pétersb. v. (1812) 541 = nivea.
macrocalyx, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 252 = rupestris.
macrophylla, Schur, Enum. Pl. Transs. 190 = recta.
macrorrhiza, Juss. ex Lehm. Monog. Pot. 168 = speciosa.
macrorrhiza, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 292 = ranunculoides.
maculata, Fries, ex Lange, Consp. Fl. Groenl. i. 9 = Frieseana.
maculata, Gilib. Fl. Lituan. ii. 252 = verna.
maculata, Pourr. in Mém. Acad. Toul. iii. (1788) 326 = aurea.
magna, Jeanb. & Timb. in Bull. Soc. Phys. et Nat. Toul. iii. (1875-76) 564 = verna.
Mathoneti, Jord. Cat. Jard. Grenoble (1849), 23 = heptaphylla.
Mayana, Hort. ex Schlecht. in Linnaea, xviii. (1844) 508 = nepalensis.
medioxenia, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = heptaphylla.
megalontodon, Lehm. Del. Sem. Hort. Hamb. (1839) 8 = inclinata.
meifolia Wall. Cat. n. 1011 = coriandrifolia.
Menziesii ×, Paxt. Mag. Bot. xv. (1849) 247.
Meyeri, *Boiss. Diagn.* Ser. I. iii. 7.—As. Min. ; Persia.
micrantha, Ledeb. ex Hook. Fl. Bor. Am. i. 195 = nivea.
micrantha, *Ram. ex DC. Fl. Fr.* iv. 468.—Europ. austr. ; Oriens.
microdons, Schur, Enum. Pl. Transs. 192 = collina.
micropetala, *D. Don, Prod. Fl. Nep.* 231.—Reg. Himal.
micropetala, Reichb. Fl. Germ. Excurs. 593 = heptaphylla.
microphylla, *D. Don, Prod. Fl. Nep.* 231.—Reg. Himal.

POTENTILLA :—

microphylla, Zimm. Gatt. Potentill. 6 = reptans.
millegrana, Dougl. ex Hook. Fl. Bor. Am. i. 193 = norvegica.
millegrana, Engelm. ex Lehm. Del. Sem. Hort. Hamb. (1849) 11 ; Pugill. ix. 22 = rivalis.
minima, *Hall. f. in Schleich. Cat.* (1815) 28 ; *et in Ser. Mus. Helv.* i. (1818) 51, 67.—Europ.
minima, Lodd. Bot. Cab. t. 480 ; A. Gray, Man. Bot. N. U. St. ed. I. 122 = frigida.
minima, Neilr. ex Nym. Consp. 225 = aurea.
minor, Gilib. Fl. Lituan. ii. 251 = verna.
missurica, Schrad. Ind. Sem. Hort. Gotting.—Cf. Linnaea, viii. (1833) 26 = pensylvanica.
mixta, Nolte, ex Koch, Syn. Pl. Germ. ed. II. 239 = reptans.
mixta, Rochel, Bot. Reise Banat. 71 = hirta.
mollis, *Panč. ex Nym. Consp.* 223.—Serbia.
mollissima, *Lehm. Ind. Sem. Hort. Hamb.* (1831) 6 ; *Pugill.* ix. 11.—Turkestan.
monanthes, *Lindl. in Wall. Cat.* n. 1025.—Reg. Himal.
monspeliensis, Linn. Sp. Pl. 499 = norvegica.
montana, Brot. Fl. Lusit. ii. 350 = splendens.
montana, Schur, ex Nym. Consp. 227 = Sibbaldi.
montenegrina, *Pantoč. in Oestr. Bot. Zeitschr.* xxiii. (1873) 6.—Montenegr.
montivaga, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1870) 47 (Quid ?).—Gallia.
Mooniana, *Wight, Ic.* i. t. 233.—Reg. Himal. ; Zeylan.
Moorcroftii, Wall. Cat. n. 1014 = bifurca.
Morisoni, DC. Cat. Hort. Monsp. 135 = norvegica.
Mulleri ×, Hort. ex Lem. Illustr. Hortic. iv. (1857) t. 143.
multicaulis, Bunge, Enum. Pl. Chin. Bor. 25 = sericea.
multidentata, *Ser. ex G. Don, in Loud. Hort. Brit.* 214.—Austria.
multifida, Baker & S. Moore, in Journ. Linn. Soc. xvii. (1879) 381 = chinensis.
multifida, *Linn. Sp. Pl.* 496.—Europ. ; As. bor.
multifida, Pall. Reise, iii. 194 = sericea ?
multifida, Willd. ex Ledeb. Fl. Ross. ii. 41 = approximata.
multijuga, *Lehm. Del. Ind. Sem. Hamb.* (1849) 6 ; *Pugill.* ix. 2.—Calif.
Munroana, Lehm. ll. cc. 7 ; 6 = fragarioides.
Murrii, *Zimm. Gatt. Potentill.* 21.—Europ.
nana, H. B. & K. ex Nestl. Monog. Pot. 34. t. 17 = candicans.
nana, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 296.—Ins. Aleut.
Naspata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 230 = fulgens.
nebrodensis, Strobl, ex Zimm. Gatt. Potentill. 29 = caulescens.
neglecta, Baumg. Enum. Stirp. Transs. ii. 63 = argentea.
nemoralis, Bunge, in Ledeb. Fl. Alt. ii. 256 = flagellaris.
nemoralis, Gaud. Fl. Helv. iii. 383 = *ascendens*.
nemoralis, Lehm. Monog. Pot. 147. t. 13 = reptans.
nemoralis, Nestl. Monog. Pot. 65 = procumbens.
nepalensis, *Hook. Exot. Fl.* t. 88.—Reg. Himal.
Nestleri, *Fries, Nov. Fl. Suec.* 45.—Suecia.
Nestleriana, Tratt. Ros. Monog. iv. 91 = intermedia.
Neumanni, Opiz, Seznam, 79 = aurea.
Neumanniana, Reichb. Fl. Germ. Excurs. 592 = aurea.
Neumayeriana, Tratt. Ros. Monog. iv. 75 = P. Tormentilla.
nevadensis, *Boiss. Elench.* 40.—Hispan.
Neymeyeriana, Reichb. Fl. Germ. Excurs. 589, sphalm. = P. Tormentilla.
Newberryi, *A. Gray, in Proc. Am. Acad.* vi. (1866) 532.—Am. bor.
nitida, *Linn. Cent. Pl.* ii. 18 ; *Amoen. Acad.* iv. 316. —Europ. ; Sibir.
nitida, Scop. Fl. Carn. ed. II. i. 362 = alba.
nitida, Tenore, Fl. Nap. ii. 291 = apennina.
nitida, Thuill. Fl. Par. ed. II. 257 = splendens.
nivalis, Csató, in Oestr. Bot. Zeitschr. xxiii. (1873) 100 = Haynaldiana.
nivalis, *Lapeyr. in Mém. Acad. Toul.* i. (1782) 210. t. 16.—Mont. Pyren.
nivalis, Torr. in Ann. Lyc. N. York, i. (1824) 32. t. 3. f. 2 ; et ii. (1828) 195 = Geum Rossii.
nivea, Gunn. Fl. Norveg. ii. 46. t. 3 = Gunneri.
nivea, *Linn. Sp. Pl.* 499.—Reg. bor. temp. et arct.
Nordmanniana, Ledeb. Fl. Ross. ii. 53 = umbrosa.

POTENTILLA :—

normalis, *Bess. ex Spreng. Syst.* iv. *Cur. Post.* 199. —Syria.
norvegica, All. Fl. Pedem. ii. 58 = frigida.
norvegica, Geners. ex Wahlenb. Fl. Carpat. 155 = argentea.
norvegica, *Linn. Sp. Pl.* 499.—Reg. bor. temp. et arct.
norvegica, Schur, Enum. Pl. Transs. 189 (Agrimoniae sp.).—Hungar.
nuda, *Boiss. Diagn.* Ser. I. vi. 49.—Persia.
nudicaulis, *Boiss. & Bal. Diagn.* Ser. II. vi. 69.—Cappadoc.
nudicaulis, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 287 = viscosa.
nurensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 720.—Persia.
nutans, Gueldenst. Reisen, ii. 79 = Geum rivale ?
Nuttallii, Lehm. Pugill. ix. 44 = gracilis.
obliqua, Dougl. ex Hook. Fl. Bor. Am. i. 186 = arguta.
obovata, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iii. (1863) 422 = supina.
obscura, Willd. Sp. Pl. ii. 1100 = recta.
ochreata, Lindl. in Wall. Cat. n. 1028 = fruticosa.
odora, Mart. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 91 = umbrosa.
officinalis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 583 = P. Tormentilla.
olopetala, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 612 = gracilis.
ontopoda, Dougl. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 617 (Quid ?).—Am. bor.
opaca, Gaertn. Mey. & Scherb. Fl. Wett. ii. 247 = verna.
opaca, Krock. Fl. Siles. ii. 171 = inclinata.
opaca, Linn. Amoen. Acad. iv. 274 = aurea.
opaca, Sm. Engl. Bot. t. 2449 = intermedia.
opaca, Vill. Prosp. 47 ; Hist. Pl. Dauph. 566 = subacaulis.
opacata, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28, nomen ; et in Lamotte, Prod. Fl. Centr. Fr. i. 242 = aurea.
oregana, Nutt. ex Torr. & Gray, Fl. N. Am. i. 446 = glandulosa.
oreites, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
ornithopoda, Tausch, Hort. Canal. i. t. 10 = multifida.
ornithopodioides, Sweet, Hort. Brit. ed. II. 191 = praec. ?
oronoma, Jord. ex Verlot, Cat. Jard. Grenoble (1856), 28 = aurea.
ovalis, *Lehm. Del. Sem. Hort. Hamb.* (1849) 9 ; *Pugill.* ix. 14.—Mexic.
ovata, Poir. Encyc. v. 601 = subacaulis.
Oweriniana, *Rupr. ex Boiss. Fl. Orient.* ii. 705.—Reg. Caucas.
pallens, Moench, Meth. 658 = recta.
pallida, Lag. ex Bess. Enum. Pl. Volh. 69 = recta.
palustris, *Scop. Fl. Carn.* ed. II. i. 359.—Reg. bor. et arct.
pannosa, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 719.—Persia.
paradoxa, Nutt. in Torr. & Gray, Fl. N. Am. i. 437 = supina.
paradoxa, Schur, ex Nym. Consp. 224 = heptaphylla.
parnassica, Boiss. & Orph. Diagn. Ser. II. v. 95 = hirta.
parva, Frivald. ex Hampe, in Flora, xx. (1837) 231 = hirta.
parviflora, Desf. Tabl. ed. I. 177 ; [Clairv.] Man. Herb. 166 = micrantha.
parviflora, Gaud. Fl. Helvet. vi. 388 = heptaphylla.
parvifolia, *Fisch. in Lehm. Pugill.* iii. 6.—Soongar.
parvula, Schur, Enum. Pl. Transs. 194 = subacaulis.
patens, Herbich, Stirp. Rar. Bucov. 41 = delphinensis.
patula, *Waldst. & Kit. Pl. Rar. Hung.* ii. 218. t. 199. —Mont. Carpath.
pectinata, Fisch. ex DC. Prod. ii. 581 = pensylvanica.
pedata, Willd. Enum. Hort. Berol. Suppl. 38 = hirta.
pedemontana, Boiss. & Reut. Ind. Sem. Hort. Genev. (1861) 4 = grandiflora.
pedunculares, *D. Don, Prod. Fl. Nep.* 230.—Reg. Himal.
pennina ×, Gremli, Neue Beitr. Fl. Schweiz, Heft 1. (1880) 45.—Europ.
pensylvanica, *Linn. Mant.* i. 76.—Am. et As. bor.
pensylvanica, Reichb. Fl. Germ. Excurs. 596 = sanguisorbifolia.
pensylvanica, Schur, Enum. Pl. Transs. 189 = tanacetifolia.

POTENTILLA :—
pensylvanica, Willd. Enum. Hort. Berol. 553, partim = viscosa.
pensylvanica, Willk. & Lange, Prod. Fl. Hisp. iii. 236 = hispanica.
pentandra, Engelm. in Torr. & Gray, Fl. N. Am. i. 447 = rivalis.
pentaphylla, Richter, ex Pohl, Tent. Fl. Bohem. ii. 181 = recta.
permiata, Schur, Enum. Pl. Transs. 951 (Index), sphalm. = seq.
permixta, Schur, l. c. 197 = heptaphylla.
perpusilla, *Hook. f. Fl. Brit. Ind.* ii. 346.—Reg. Himal.
persica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 710.—Persia.
petiolulata, Gaud. Fl. Helv. iii. 374 = caulescens.
petraea, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 292.—Persia.
petrophila, Boiss. Voy. Bot. Espagne, ii. 728 = caulescens.
Pickeringii, *S. Wats. Bibl. Ind. N. Am. Bot.* 289.—Am. bor.
pigmaea, Jord. Obs. Pl. Crit. vii. (1849) 25 = rupestris.
pilosa, Kitt. Taschenb. ed. II. 1081 = verna.
pilosa, Vill. Prosp. 47 ; Hist. Pl. Dauph. iii. 570 = hirta.
pilosa, Willd. Sp. Pl. ii. 1100 = recta.
pilosissima, Schur, Enum. Pl. Transs. 195 = aurea.
pimpinelloides, Falk, Beitr. ii. 193 = bifurca.
pimpinelloides, *Linn. Sp. Pl.* 497.—Armen.
pimpinelloides, Panč. in Verh. Zool. Bot. Ges. Wien, vi. (1856) 487 = Visianii.
pinnatifida, Presl, Delic. Prag. 54 = reptans.
plattensis, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 439.—Am. bor. occ.
poëtarum, Boiss. & Sprun. Diagn. Ser. I. iii. 3 = speciosa.
polymorpha, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 753 = verna.
polyodonta, Borb. ex Zimm. Gatt. Potentill. 9 = inclinata.
polyphylla, Schur, Enum. Pl. Transs. 198 = heptaphylla.
polyphylla, Wall. Cat. n. 1026 = Mooniana.
polyschista, Boiss. & Hohen. Diagn. Ser. I. x. 6 = sericea.
polytricha, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
pontica, C. Koch, in Linnaea, xix. (1847) 44 = heptaphylla.
poteriifolia, *Boiss. Diagn.* Ser. I. vi. 50.—Persia.
poteriifolia, Vis. in Mem. Ist. Venet. x. (1861) 433 = Visianii.
potterioides, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 286 = fragarioides.
praecox, F. Schultz, in Pollichia, xvi-xvii. (1859) 5 = collina.
praeruptorum, F. Schultz, l. c. xviii–xix. (1861) 105 ; xxii–xxiv. (1866) 154 = aurea.
pratensis, Herbich, in Flora, xxxviii. (1855) 641 = delphinensis.
pratensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 23 ; x. (1859) 98 = patula.
procumbens, [Clairv.] Man. Herb. 166 = Sibbaldi.
procumbens, *Sibth. Fl. Oxon.* 162.—Europ.
properiflora, Jord. ex Verlot, Cat. Pl. Grenoble (1856), 28 = aurea.
prostrata, Baumg. Enum. Stirp. Transs. ii. 68 = micrantha.
prostrata, Haenke, ex Pohl, Tent. Fl. Bohem. ii. 181 = supina.
prostrata, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 76 = fruticosa.
prostrata, Moench, Meth. 659 = P. Fragariastrum.
prostrata, Rottb. in Skrift. Kiobenh. Selsk. Vidensk. x. (1765–69) 453 = nivea.
Pseudo-alpestris, Schur, ex Nym. Consp. 225 = aurea.
Pseudo-anserina, Bertol. Misc. Bot. xxiv. 15. t. 3 = P. Anserina.
Pseudo-frigida, Schur, Enum. Pl. Transs. 195 = aurea.
Pseudo-opaca, Gremli, Excurs. Schw. ed. I. p. xii = aurulenta.
pteropoda, *Royle, Illustr. Bot. Himal.* 207. t. 40. f. 2.—Reg. Himal.
puberula, *Krasan, in Oestr. Bot. Zeitschr.* xvii. (1867) 304.—Europ.
pubescens, Moench, Meth. 657 = viscosa.
pulchella, *R. Br. in Ross's Voy.* ed. II. ii. 193.—Ins. Melvil.
pulchella, Spreng. Syst. iv. Cur. Post. 199 = humifusa.
pulcherrima, Lehm. Pugill. ii. 11 = Hippiana.

POTENTILLA :—
pulchra, *Langsd. Bemerk. Reise Welt*, i. 45 ; *Voy. & Trav.* (1814) ii. 51, nomen.—Unalaschk.
pulchra, Salisb. Prod. 362 = P. Anserina.
Pulsatilla, C. Muell. in Walp. Ann. vii. 867 (Anemones sp.).—Cf. Regel, in Bull. Soc. Nat. Mosc. (1859) i. 225.—As. bor.
pulvinaris, *Fenzl, Pugill. Pl. Nov. Syr.* 6.—As. Min.
pumila, Poir. Encyc. v. 594 = canadensis.
purpurea, *Hook. f. Fl. Brit. Ind.* ii. 347.—Reg. Himal.
pusilla, Host, Fl. Austr. ii. 39 = verna.
pygmaea, C. Muell. in Walp. Ann. ii. 478 = rupestris.
pyrenaica, Ram. ex DC. Fl. Fr. iv. 459 = aurea.
pyrenaica, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 292 = alchemilloides.
radiata, *Lehm. Del. Ind. Sem. Hamb.* (1849) 9 ; *Pugill.* ix. 13.—Persia.
radicosa, Boiss. Diagn. Ser. I. vi. 49 = nuda.
ranunculoides, *Humb. & Bonpl. ex Nestl. Monog. Pot.* 56.—Mexic.
Ranunculus, *Lange, in Fl. Dan.* t. 2964.—Groenl.
reclinis, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = argentea.
recta, Jacq. Fl. Austr. iv. t. 383 = inclinata.
recta, *Linn. Sp. Pl.* 499.—Europ. ; As. bor.
recta, Nutt. Gen. Am. i. 310 = gracilis.
reptans, Georgi, Reise, i. 219 = flagellaris.
reptans, *Linn. Sp. Pl.* 499.—Europ. ; As. et Am. bor.
reptans, Willd. ex Ledeb. Fl. Ross. ii. 52 = procumbens.
reticulata, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iii. (1863) 420 = fruticosa.
retusa, O. F. Muell. in Fl. Dan. t. 799 = tridentata.
Reuteri, *Boiss. Diagn.* Ser. II. ii. 51.—Hispan.
rhaetica × , Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 58.—Europ.
rhenana, P. J. Muell. ex F. Schultz, Arch. Fl. 272 = collina.
Richardii, *Lehm. Del. Ind. Sem. Hamb.* (1849) 6 ; *Pugill.* ix. 1.—Mexic.
rigida, Nutt. in Journ. Acad. Philad. vii. 1. (1834) 20 = gracilis.
rigida, Wall. Cat. n. 1009 = fruticosa.
rivalis, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 437.—Am. bor. occ.
rivalis, Torr. in Pacif. Rail. Rep. iv. 84 = glandulosa.
Robbinsiana, Oakes, ex Torr. & Gray, Fl. N. Am. i. 441 = minima.
rotundifolia, Vill. Hist. Pl. Dauph. iii. 565 = aurea.
rubens, Vill. Prosp. 46 ; Moench, Meth. 657 = aurea.
rubescens, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
rubifolia, Boiss. Fl. Orient. ii. 723 = seq.
rubiformis, Czern. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 152 = elatior.
rubra, Hall. f. in Ser. Mus. Helv. i. (1818) 56 = palustris.
rubra, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 292 = comaroides.
rubricaulis, Bourg. ex S. Wats. Bibl. Ind. N. Am. Bot. 298 = Hippiana.
rubricaulis, Lehm. Pugill. ii. 11 = dissecta.
rubricaulis, Pall. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 94 = petraea.
ruderalis, Kitt. Taschenb. ed. II. 1081 = argentea.
rufescens, *Franch. & Sav. Enum. Pl. Jap.* ii. 340.—Japon.
rupestris, Falk, Beitr. ii. 193. t. 9 = recta.
rupestris, *Linn. Sp. Pl.* 496.—Europ. ; As. bor.
rupestris, Pall. ex Ledeb. Fl. Ross. ii. 41 = viscosa.
rupestris, Presl, Epim. Bot. 198 = glandulosa.
Ruprechti, *Boiss. Fl. Orient.* ii. 716.—Reg. Caucas.
Russelliana × , Hort. ex Sweet, Flow. Gard. t. 279.
ruthenia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. ii. 359 = supina.
ruthenica, Willd. Sp. Pl. ii. 1079 = norvegica.
sabauda, DC. Fl. Fr. iv. 468 = rupestris.
Sadleri, Reichb. Fl. Germ. Excurs. 594 = inclinata.
Sadleri, Schloss. ex Nym. Consp. 224 = recta.
Sadleri, Schur, Enum. Pl. Transs. 198 = heptaphylla.
Salesoviana, *Steph. in Mém. Soc. Nat. Mosc.* ii. (1809) 6. t. 3.—Reg. Himal. ; As. centr.
Salesovii, Steph. ex Willd. Enum. Hort. Berol. 552 = praec.
salisburgensis, Haenke, in Jacq. Coll. ii. 68 = aurea.
Salisii, Bor. in Mém. Soc. Acad. Maine-et-Loire, xiv. (1863) 41 = Sibbaldi.
Sanguisorba, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 286.—Sibir.

POTENTILLA :—
sanguisorbifolia, *F. O. Wolf, in Bull. Soc. Murith.* xi. (1883) 9.—Europ.
santolinoides, *Baill. ex S. Wats. Bibl. Ind. N. Am. Bot.* 289.—Am. bor.
sarmentosa, Bigel. Fl. Bost. ed. II. 204 = canadensis.
sarmentosa, Willd. Enum. Hort. Berol. 554 = canadensis.
Saundersiana, Royle, Illustr. Bot. Himal. 207. t. 41. f. 1 = multifida.
saxatilis, Boulay, in Billotia, i. (1869) 109 = aurea.
Saxifraga, *Ardoin, ex De Not. Ind. Sem. Genuen.* (1848) ; *ex Lehm. in Nov. Act. Nat. Cur.* xxiii. *Suppl.* (1856) 134.—Europ.
scepusiensis, *Kit. in Linnaea*, xxxii. (1863) 597.—Europ.
Schleicheri, Zimm. Gatt. Potentill. 26 = aurea.
Schrenkiana, *Regel, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 379.—As. centr.
Schultzii, P. J. Muell. ex F. Schultz, Arch. Fl. 306 = collina.
Schurii, Fuss, ex Zimm. Gatt. Potentill. 17 = patula.
sciaphila, *Zimm. l. c.* 5.—Europ.
Seidlitziana, *Bienert, in Seidlitz, Bot. Ergeb.* 96.—Armenia.
semiargentea × , Borb. in Oestr. Bot. Zeitschr. xxxii. (1882) 170.—Europ.
semidentata, Juss. ex Lehm. Monog. Pot. 175 = splendens.
semiloba, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = argentea.
septemsecta, *G. F. W. Mey. Fl. Hannov. Excurs.* 178.—Europ.
sericea, Dulac, Fl. Hautes-Pyr. 308 = alchemilloides.
sericea, Grev. in Mem. Wern. Soc. iii. (1821) 430 = pulchella.
sericea, *Linn. Sp. Pl.* 495.—Reg. Caucas. ; As. bor.
serotina, Vill. Hist. Pl. Delph. iii. 564 = verna.
Serpentini, *Borb. ex Zimm. Gatt. Potentill.* 22.—Europ.
sessilis, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 197 = reptans.
Sibbaldi, *Hall. f. in Ser. Mus. Helv.* i. (1818) 51.—Reg. bor. et austr.
Sibbaldia, Griessel. Kl. Bot. Schrift. i. 239 = praec.
Sibbaldiana, Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 203 = praec.
sibirica, Patr. ex Lehm. Monog. Pot. 88 = aurea ?
Siemersiana, Lehm. Sem. Hort. Hamb. (1821) 8 ; Pugill. i. 31 = fulgens.
Sieversiana, *Otto, ex Sweet, Hort. Brit.* ed. II. 191.—Hab. ?
silesiaca, Uechtr. in Jahresb. Schles. Ges. Vaterl. Cult. (1867) 82 = collina.
silvestris, Neck. Delic. Gallo-Belg. i. 222 = *sylvestris*.
similata, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid ?).—Hab. ?
simplex, Michx. Fl. Bor. Am. i. 303 = canadensis.
sischanensis, *Bunge, ex Lehm. Pugill.* ix. 3.—China.
Smoutii × , Lem. Fl. des Serres, iv. (1848) t. 373.
Sommerfeltii, *Lehm. Del. Ind. Sem. Hamb.* (1849) 6 ; *Pugill.* ix. 4.—Ins. Spitzberg.
songarica, Bunge, in Ledeb. Fl. Alt. ii. 244 = sericea.
sordida, Fries, Summa Veg. Scand. 44 = collina.
sordida, Klotzsch, ex Lehm. Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 53 = Mooniana.
sororia, Wender. Fl. Hass. 158 = caulescens.
speciosa, *Willd. Sp. Pl.* ii. 1110.—Graec. ; As. Min.
splendens, Buch.-Ham. ex Trevir. Ind. Sem. Hort. Wratisl. (1823) 3 = fulgens.
splendens × , Koch, Syn. Fl. Germ. ed. II. 243.—Europ.
splendens, *Ram. in DC. Fl. Fr.* iv. 467.—Gallia.
Sprengeliana, Hassk. Cat. Hort. Bog. Alt. 266 = japonica.
Sprengeliana, *Lehm. Monog. Pot.* 49.—Sibir.
spuria × , Kern. in Oestr. Bot. Zeitschr. xx. (1870) 41, 45.—Europ.
ssawellanica, Bienert, in Seidlitz, Bot. Ergeb. 93 = argaea.
stellulata, Rochel, Pl. Banat. Rar. [9] Index ; Bot. Reise Banat. 72 = subacaulis.
stenantha, Lehm. Del. Ind. Sem. Hamb. (1849) 7 ; Pugill. ix. 7 = Agrimonia Eupatoria.
sterilis, Garcke, Fl. Deutschl. ed. IV. 112 = P. Fragariastrum.
stipulaceum, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 566 = stipularis.
stipularis, Jeanb. & Timb. l. c. = seq. ?

POTENTILLA:—
stipularis, Lapeyr. in Mém. Acad. Toul. i. (1782) 212 = nivalis.
stipularis, *Linn. Sp. Pl.* 498.—Sibir.
stolonifera, Lehm. Ind. Sem. Hort. Hamb. (1831) = japonica.
stricta, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid?).—Hab.?
strictissima, *Zimm. Gatt. Potentill.* 5.—Europ.
strigosa, Pall. ex Pursh, Fl. Am. Sept. i. 356 = pensylvanica.
subacaulis, Lapeyr. Hist. Abr. Pl. Pyrén. 290 = frigida, verna.
subacaulis, *Linn. Syst.* ed. X. 1065.—As. bor.
subacaulis, Scop. Fl. Carn. ed. II. i. 364 = nitida.
subalpina, Zimm. Gatt. Potentill. 14 = heptaphylla.
subarenaria, Zimm. l. c. 21 = verna.
subargentea, Krock. Fl. Siles. iv. 513 = collina.
subcinerea, Borb. ex Zimm. Gatt. Potentill. 21 = verna.
suberecta ×, Zimm. l. c. 5.—Europ.
subpalmata, *Ledeb. Fl. Ross.* ii. 56.—Armenia.
subpedata, C. Koch, in Linnaea, xvi. (1842) 349 = reptans.
subpinnata, Ledeb. Ind. Sem. Hort. Dorpat. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 35 = multifida.
subrecta, Jord. Pugill. Pl. Nov. 65 = inclinata.
subrubens ×, Borb. ex Zimm. Gatt. Potentill. 21.—Europ.
subsalsa, Schur, Enum. Pl. Transs. 189 = supina.
subviscosa, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 97.—N. Mexic.
suffruticosa, Pourr. ex Lehm. Monog. Pot. 177 = frigida.
sulphurea, Lam. Fl. Fr. iii. 114 = recta.
supina, *Linn. Sp. Pl.* 497.—Geront. temp. et trop.
syenitea, Jord. Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
sylvestris, Neck. Delic. Gallo-Belg. i. 222 (*silvestris*) = P. Tormentilla.
sylvestris, Renault, Fl. Dép. Orne, 148 = procumbens.
tanacetifolia, Schur, Enum. Pl. Transs. 189 (Quid?).—Hungar.
tanacetifolia, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 286.—Sibir.
tanaitica, *Zinger, in Bull. Soc. Nat. Mosc.* lvii. (1882) II. 69.—Rossia.
taurica, Schur, Enum. Pl. Transs. 198 = heptaphylla.
taurica, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 291 = hirta.
Tellusonii ×, Hort. ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 207.
tenella, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 620 = multifida.
tenuifolia, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 285 = fruticosa.
tenuiloba, Jord. Pugill. Pl. Nov. 67 = argentea.
tenuirugis, *Pomel, Nouv. Mat. Fl. Atl.* 158.—Alger.
tergloviensis, Hacq. Pl. Alp. Carn. ii. 15. t. 4. f. 2 = nitida.
ternata ×, Bruegg. in Cat. Hort. Turic. (1870) Suppl. 3.—Europ.
ternata, C. Koch, in Linnaea, xix. (1847) 45 = aurea.
tetrandra, *Hook. f. Fl. Brit. Ind.* ii. 346.—Reg. Himal. et Altaic.
tetrapetala, Hall. f. in Ser. Mus. Helv. i. (1818) 51 = P. Tormentilla.
Thellusonii, Hort. Ind. Sem. Hort. Basil. (1848); ex Walp. Ann. ii. 519 = *Tellusoni*.
Thomasii, C. A. Mey. Verz. Pfl. Cauc. 168 = Meyeri.
Thomasii, Ser. in DC. Prod. ii. 576 = Detommasii.
Thurberi, *A. Gray, ex Lehm. Ind. Sem. Hort. Hamb.* (1854) 10; *et in Mem. Am. Acad.* N. S. v. (1855) 318.—N. Mexic.
thuringiaca, Bernh. ex Link, Enum. Hort. Berol. ii. 64 = heptaphylla.
thyrsiflora, Kern. Sched. Fl. Exsicc. ii. 21 = collina.
tiroliensis, Zimm. ex Kern. l. c. iii. 22 = verna.
Tomasii, C. Muell. in Walp. Ann. ii. 490, sphalm. = *Thomasii*.
tomentosa, Gilib. Fl. Lituan. ii. 254 = argentea.
tomentosa, Tenore, ex Nym. Syll. Suppl. 49 = collina.
Tommasii, Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 95, sphalm. = *Thomasii*.
Tommasiniana, F. Schultz, Arch. Fl. (1858) 264; et in Pollichia, xvi–xvii. (1859) 7 = arenaria.
Tonguei, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. II. 660 = nepalensis.

POTENTILLA:—
Tormentilla, *Neck. in Act. Acad. Theod. Palat.* ii. (1770) 491; *Schrank, Bair. Fl.* ii. 53.—Europ.; As. bor.
Tormentilla-elatior, Tratt. Ros. Monog. iv. 70 = praec.
Tormentilla-erecta, Scop. Fl. Carn. ed. II. i. 360 = praec.
Tormentilla-formosa, Tongue, ex Loud. Hort. Brit. Suppl. ii. 660 = nepalensis.
tormentilloides, Lange, ex Nym. Consp. 226 = aurea.
tormentilloides, Mayer, ex Tausch, in Flora, ii. (1819) 467 = procumbens.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 23 = aurea.
tridentata, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 216. t. 9.—Am. bor. et arct.
trifida, Lehm. in Hamb. Gartenz. vii. (1851) 263 = Fragaria indica.
trifida, Pall. Reise, i. 215 = bifurca.
trifolia, Gilib. Fl. Lituan. ii. 249 = norvegica.
trilobata, Poir. Encyc. Suppl. iv. 536 = aurea.
trullifolia, *Hook. f. Fl. Brit. Ind.* ii. 345.—Reg. Himal.
Turczaninowiana, *Stschégl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 164.—As. centr.
turicinensis, Siegf. ex Zimm. Gatt. Potentill. 20 = aurea.
tyrolensis, *Scop. Ann.* II. *Hist. Nat.* (1769) 52 (Quid?).—Europ.
ucranica, Lodd. Cat. (1818); ex Lehm. Monog. Pot. 195 = argentea.
Uechtritzii, Zimm. Gatt. Potentill. 9 = canescens.
umbrosa, *Stev. in Bess. Cat. Hort. Crem.* (1816) 110; *ex Bieb. Fl. Taur. Cauc.* iii. 357.—Tauria; Reg. Caucas.
uniflora, Ledeb. in Hook. Fl. Bor. Am. i. 195 = nivea.
uniflora, Ledeb. in Mém. Acad. Pétersb. v. (1812) 543 = gelida.
Vahliana, Lehm. Monog. Pot. 172 = nivea.
Vaillantii, Nestl. Monog. Pot. 75 = splendens.
valdepubens, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid?).—Hab.?
Valderia, *Linn. Syst.* ed. X. 1064.—Pedem.
Valderia, Vill. Hist. Pl. Dauph. iii. 572 = nivalis.
vallesiaca, *Huet, in Compt. Rend. Soc. Hallér.* (1853–54) 32.—Europ.
varia, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 755 = argentea.
variabilis, Klotzsch, ex Lehm. in Nov. Act. Nat. Cur. xxiii. Suppl. (1856) 42 = fragarioides.
varians, Moench, Meth. 658 = intermedia.
velutina, Lehm. Monog. Pot. 170 = subacaulis.
velutina, Wall. Cat. n. 1016 = peduncularis.
Verloti, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
verna, Hook. in Scoresby, Greenland, 431 = emarginata.
verna, *Linn. Sp. Pl.* 498.—Reg. bor. temp.
vernalis, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
verticillaris, Steph. ex Willd. Sp. Pl. ii. 1096 = multifida, sericea.
vesca, Scop. Fl. Carn. ed. II. i. 363 = Fragaria vesca.
vestita, Jord. in Cat. Pl. Jard. Grenoble (1849), 23 = subacaulis.
vestita, Klotzsch, in Bot. Ergeb. Waldem. Reise, 157 = argyrophylla.
vilifera, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 (Quid?).—Hab.?
villosa, Dulac, Fl. Hautes-Pyr. 308 = P. Fragariastrum.
villosa, Hall. f. ex Lehm. Monog. Pot. 100 = intermedia.
villosa, Pall. ex Pursh, Fl. Am. Sept. i. 353 = gelida.
viminea, Schrad. ex Lehm. Monog. Pot. 92 = aurea.
vindobonensis, Zimm. ex Kern. Sched. Fl. Exsicc. iii. 23 = aurea.
vinosa, Lodd. ex Spreng. Syst. iv. Cur. Post. 199 = inclinata.
virgata, *Lehm. Monog. Pot.* 75.—Podolia.
viscosa, *Donn, Hort. Cantab.* ed. II. 68.—Reg. Casp.
viscosa, Steud. Nom. ed. I. 653 = hirta.
Visianii, *Panč. in Mem. Ist. Venet.* x. (1861) 433; xii. (1864) 480.—Serbia.
visurgina, Weihe, ex Reichb. Fl. Germ. Excurs. 871 = norvegica.
vitodurinensis, Siegf. ex Zimm. Gatt. Potentill. 19 = aurea.

POTENTILLA:—
vivariensis, Jord. Cat. Jard. Grenoble (1849), 24 = verna.
Vockei, P. J. Muell. ex F. Schultz, Arch. Fl. 272 = collina.
vulcanica, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 717 = Kotschyana.
vulgaris, Kitt. Taschenb. ed. II. 1081 = verna.
Wallichiana, DC. ex Wall. Cat. n. 1022 = Kleiniana.
Wallichiana, Govan, ex Wall. Cat. n. 1013 = argyrophylla.
Wallichiana, Ser. in DC. Prod. ii. 574 = Fragaria indica.
Weimannii, Lodd. Bot. Cab. t. 706 = collina.
Wheeleri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 148.—Calif.
Wiemanniana, [Guenth. Grab. & Wimm.] Enum. Phan. Siles. 88 = collina.
Wolnyi, Kit. ex Rochel, Bot. Reise Banat. 71 = P. Fragariastrum.
Wrangeliana, Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop. vii. 54 = glandulosa.
xerophila, Jord. ex Verlot, Cat. Pl. Jard. Grenoble (1856), 28 = aurea.
xylorrhiza, Boiss. & Hohen. Diagn. Ser. I. x. 5 = lignosa.
Zichyi, *Szont. in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 1095.—Hungar.
Zimmeteri, Borb. in Oestr. Bot. Zeitschr. xxxii. (1882) 180 = verna.

POTENTILLOPSIS, Opiz, in Lotos, vii. (1857) 30 = **Potentilla**, Linn. (Rosac.).
Engelmanniana, Opiz, l. c. = Potentilla rivalis.

POTERANTHERA, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1835) Bot. 137. t. 8. *MELASTOMACEAE*, Benth. & Hook. f. i. 739.
ONOCTONIA, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 276.
calcarata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 33.—Bras.
crassipes, *Triana, l. c.* 33. t. 1. f. 17 c.—Bras.
pauciflora, *Triana, l. c.* t. 1. f. 17 b.—Bras.
pusilla, *Bong. in Mém. Acad. Pétersb.* Sér. VI. ii. (1835) *Bot.* 137. t. 8. f. 1.—Bras.

POTERIDIUM, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 43 = **Poterium**, Linn. (Rosac.).
annuum, Spach, l. c. = Poterium annuum.

POTERION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = **Poterium**, Linn. (Rosac.).
sanguisorbens, St. Lag. l. c. = Poterium Sanguisorba.

POTERIUM, Linn. Gen. ed. I. 289 (1737). *ROSACEAE*, Benth. & Hook. f. i. 624.
PIMPINELLA, Tourn. ex Rupp. Fl. Jen. ed. Hall. 285 (1745); Adans. Fam. ii. 293 (1763).
POTERIDIUM, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 43.
SANGUISORBA, Rupp. ex Linn. Syst. ed. I (1735).
SARCOPOTERIUM, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 43.
agrimonifolium, Cav. Elench. Hort. Madr. (1803); ex DC. Prod. ii. 595 = hybridum.
agrimonifolium, Link, in Webb & Berth. Phyt. Canar. ii. 10 = verrucosum.
agrimonoides, Linn. Sp. Pl. 994, in syn. = hybridum.
alophum, Spach, ex Nym. Consp. 240 = Spachianum.
alveolatum, Pomel, Nouv. Mat. Fl. Atl. 162 = seq.
alveolosum, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 38.—Afr. bor.
anceps, *Ball, in Journ. Bot.* xi. (1873) 332.—Marocc.
ancistroides, *Desf. Fl. Atlant.* ii. 346. t. 251.—Reg. Mediterr. occ.
ancistroides, Guir. ex Nym. Consp. 239 = rupicolum.
annuum, *Nutt. ex Hook. Fl. Bor. Am.* i. 198.—Am. bor. occ.
australe, Salisb. Prod. 360 = Acaena sericea.
balearicum, Bourg. ex Nym. Consp. 240 = Spachianum.
boreale, Salisb. Prod. 360 = officinale.
canadense, *A. Gray, Man. Bot. N. U. St.* ed. V. 150.—Am. bor.; Japon.
caudatum, *Ait. Hort. Kew.* ed. I. iii. 354.—Ins. Canar.

POTERIUM:—
collinum, Salisb. Prod. 361 = P. Sanguisorba.
compactum, *Boiss. Diagn.* Ser. I. x. 7.—Syria.
crispum, Pomel. Nouv. Mat. Fl. Atl. 162 = alveolosum.
Delorti, Jord. Obs. Pl. Crit. vii. (1849) 21 = P. Sanguisorba.
diandrum, *Hook. f. Fl. Brit. Ind.* ii. 362.—Reg. Himal.
dictyocarpum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 206 = ancistroides.
dictyocarpum, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 34 = P. Sanguisorba.
Duriaei, *Spach, l. c.* 33.—Alger.
Duriaei, Willk. in Bot. Zeit. v. (1847) 430 = verrucosum.
eriocarpum, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 36.—Italia.
filiforme, *Hook. f. Fl. Brit. Ind.* ii. 362.—Reg. Himal.
Fontanesii, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 41 = verrucosum.
frutescens, *Bory, Ess. Isles Fortun.* 346 (Quid?).—Ins. Canar.
fruticosum, Link, in Buch, Beschr. Canar. Ins. 155 = praec.
Gaillardoti, *Boiss. Diagn.* Ser. II. ii. 52.—Syria; Mesopot.
garganicum, *Tenore, Syll. Fl. Neap.* 261.—Italia.
glaucescens, Reichb. Fl. Germ. Excurs. 610 = P. Sanguisorba.
glaucum, Noé, ex Nym. Consp. 239 = villosum.
guestphalicum, Boenn. ex Bor. Fl. Centr. Fr. ed. III. ii. 213 = P. Sanguisorba.
humile, Lam. ex Steud. Nom. ed. I. 653 = Acaena magellanica.
hybridum, *Linn. Sp. Pl.* 994.—Europ. austr.
hybridum, T. Nees, Gen. Fl. Germ. Monoc. n. 79 = polyganum.
indicum, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 10.—Zeylan.
Jaubertii, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 33 = villosum.
lasiocarpum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 733.—Persia.
lateriflorum, *Coss. Notes Crit.* 107.—Hispan.
longifolium, *Hook. f. Fl. Brit. Ind.* ii. 363.—Reg. Himal.
lucidum, Salisb. Prod. 360 = Acaena lucida.
Magnolii, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 38 = verrucosum.
Magnolii, Willk. in Flora, xxxiv. (1851) 633 = P. Sanguisorba.
mauritanicum, Boiss. Voy. Bot. Espagne, ii. 205 = verrucosum.
megacarpon, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimp. 22 = verrucosum.
microphyllum, Jord. Obs. Pl. Crit. vii. (1849) 20 = Spachianum.
minus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 575 = P. Sanguisorba.
multicaule, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 44.—Hispan.; Marocc.
muricatum, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 36 = polygamum.
obscurum, Jord. Obs. Pl. Crit. vii. (1849) 23 = P. Sanguisorba.
officinale, *A. Gray, in Proc. Am. Acad.* vii. (1868) 340.—Europ.: As. bor.; Am. bor.
platylophum, Jord. Obs. Pl. Crit. vii. (1849) 22 = polygamum.
polygamum, Desf. Cat. Hort. Par. ed. III. 292 = verrucosum.
polygamum, Lej. Rev. Fl. Spa, 199 = P. Sanguisorba.
polygamum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 217. t. 198.—Europ.; Oriens.
rupicolum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 45.—Hispan.
Sanguisorba, *Linn. Sp. Pl.* 994.—Reg. bor. temp.
sitchense, *S. Wats. Bibliog. Ind. N. Am. Bot.* i. 303.—Am. bor. occ.
Spachianum, *Coss. Notes Crit.* 108.—Hispan.
spinosum, *Linn. Sp. Pl.* 994.—Syria.
stenolophum, Jord. Obs. Pl. Crit. vii. (1849) 22 = polygamum.
tenuifolium, *Franch. & Sav. Enum. Pl. Jap.* i. 133.—As. or.

POTERIUM:—
verrucosum, *Ehrenb. in Ind. Sem. Hort. Berol.* (1829); *ex Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 263.—Reg. Mediterr.; Arab.
verrucosum, Link, ex G. Don, Gen. Syst. ii. 595 = P. Sanguisorba?
vestitum, Pomel, Nouv. Mat. Fl. Atl. 161 = alveolosum.
villosum, *Sibth. & Sm. Fl. Graec. Prod.* ii. 238.—As. Min.

POTHA, Burm. Thes. Zeyl. 197 (1737); Linn. Fl. Zeyl. 156 (1747) = **Pothos**, Linn. (Aroideae).

POTHOIDIUM, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 70; Aroid. 26. t. 57 (1857). *AROIDEAE*, Benth. & Hook. f. iii. 999.
Lobbianum, *Schott, ll. cc.*—Ins. Philipp.

POTHOMORPHE, Miq. 'in Bull. Néerl. (1840)'; Comm. Phyt. 36 (1840) = **Piper**, Linn.
Dombeyana, Miq. Syst. Pip. 211 = Piper umbellatum.
hirsuta, Bartl. Ind. Sem. Hort. Gotting. (1863) 6.—Venezuela.
Lessertiana, Miq. Syst. Pip. 210 = Piper umbellatum.
mollissima, Miq. Comm. Phyt. 46 = Piper mollissimum.
Ottonis, Miq. Syst. Pip. 207 = Piper peltatum.
peltata, Miq. Comm. Phyt. 37, 45 = Piper peltatum.
pruinosa, Miq. l. c. 37 = Piper peltatum.
scutata, Miq. l. c. = Piper peltatum.
sidaefolia, Miq. l. c. 36 = Piper subpeltatum.
speciosa, Miq. l. c. 37 = Piper peltatum.
subpeltata, Miq. l. c. = Piper subpeltatum.
umbellata, Miq. l. c. 36 = Piper umbellatum.

POTHOS, Adans. Fam. ii. 57 (1763) = **Polianthes**, Linn. (Amaryll.).

POTHOS, Linn. Diss. Dass. 26 (1747); Amoen. Acad. i. 410 (1749). *AROIDEAE*, Benth. & Hook. f. iii. 999.
GONIURUS, Presl, Epim. Bot. 244 (1849).
PODOSPADIX, Rafin. Fl. Tellur. iv. 14 (1836).
TAPANAVA, Adans. Fam. ii. 470 (1763).
acaulis, Hook. Exot. Fl. ii. t. 122 = Anthurium Hookeri.
acaulis, Hort. Berol. ex Tausch, in Flora, xix. (1836) 424 = brachypodos.
acaulis, Jacq. Enum. Pl. Carib. 31 = Anthurium acaule.
aduncus, *Vell. Fl. Flum.* ix. t. 124.—Bras.
Albertisii, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 267.—N. Guin.
anceps, Spreng. ex Schult. Mant. iii. 301 = Gymnostachys anceps.
angustatus, H. B. & K. Nov. Gen. et Sp. i. 77 = Anthurium angustatum.
angustifolius, *Presl, Epim. Bot.* 243.—Ind. or.
angustifolius, Reinw. ex Miq. in Bot. Zeit. xiv. (1856) 561 = longifolius.
argyraeus, Hort. ex Engl. in DC. Monog. Phan. ii. 255 = Scindapsus pictus.
aureus, *Linden, ex André, in Illustr. Hortic.* xxvii. (1880) 69. t. 381.—Ins. Salomon.
auritus, Willd. ex Schult. Mant. iii. 301 = Syngonium Vellozianum.
australasicus, F. Muell. Fragm. i. 62 = longipes.
Barberianus, *Schott, Aroid.* 24. t. 53.—Malacca; Borneo.
Beccarianus, *Engl. in DC. Monog. Phan.* ii. 92.—Borneo.
bifarius, *Wall. Cat.* n. 4445 (Quid?).—Singapore.
brachypodos, *Tausch, in Flora*, xix. (1836) 424.—Ins. Martinic.
breviscapus, Poepp. ex Kunth, Enum. Pl. iii. 78 = Anthurium breviscapum.
brevistylus, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 267.—Borneo.
camtschaticus, Spreng. Syst. iii. 767 = Lysichitum camtschatcense.
cannaefolius, Dryand. in Sims, Bot. Mag. t. 603 = Spathiphyllum cannaeforme.
cannaefolius, Rudge, Pl. Gui. t. 33 = Philodendron Rudgeanum.
cannaeformis, H. B. & K. Nov. Gen. et Sp. i. 76 = Spathiphyllum cannaeforme.
cartilagineus, Desf. Cat. Hort. Par. ed. III. 8, 386 = Anthurium cartilagineum.

POTHOS:—
Cathcarti, *Schott, Aroid.* 22. t. 44.—Reg. Himal.
caudatus, Roxb. Hort. Beng. 83; Fl. Ind. i. 436 = Epipremnum mirabile.
celatocaulis, *N. E. Br. in Gard. Chron.* (1880) i. 200.—Borneo.
Chapelieri, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 19.—Madag.
clavatus, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 266.—N. Guin.
cognatus, Schott, Aroid. 22. t. 48 = scandens.
cordatus, Ait. Hort. Kew. ed. I. iii. 319 = macrophyllus?
cordatus, Linn. Sp. Pl. ed. II. 1373 = Anthurium Guildingii.
cordatus, Willd. Enum. Hort. Berol. 168 = Anthurium cordifolium.
coriaceus, R. Grah. in Edinb. Phil. Journ. (Apr. 1826) 353 = Anthurium coriaceum.
coriaceus, Salisb. Prod. 262 = macrophyllus?
crassinervius, Hook. Bot. Mag. t. 2987 = Anthurium Hookeri.
crassinervius, Jacq. Coll. iv. 122 = Anthurium crassinervium.
crenatus, Linn. Sp. Pl. ed. II. 1373 = Anthurium acaule.
Cumingianus, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 19 (Cummingianus).—Ins. Philipp.
Cuscuaria, Aubl. Pl. Gui. ii. 840 = Scindapsus Cuscuaria.
cylindricus, Presl, Epim. Bot. 243 = Loureirii.
decipiens, Schott, in Bonplandia, vii. (1859) 165 = scandens.
decurrens, Wall. Cat. p. 215. n. 4437, sphalm. = *decursivus*.
decursivus, Roxb. Hort. Beng. 11; Fl. Ind. i. 436 = Rhaphidophora decursiva.
decursivus, Wall. Pl. As. Rar. ii. 83 = Epipremnum mirabile.
digitatus, Jacq. Ic. iii. 18; Coll. iv. 119 = Anthurium digitatum.
drupaceus, *Dennst. Schluess. Hort. Malab.* 33.—Ind. or.
elegans, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 267.—N. Guin.
elegans, Wall. Cat. n. 4444 = Anadendrum montanum.
ellipticus, Moon, Cat. Pl. Ceyl. 11, nomen; et ex Miq. Fl. Ind. Bat. iii. 132 = remotiflorus.
elongellus, Hort. ex Kunth, Enum. Pl. iii. 69 = Anthurium longifolium.
Enderianus, *Hort.—Cf. Gard. Chron.* (1884) i. 711.—Borneo.
erneaphyllus, Vell. Fl. Flum. ix. t. 126 = Anthurium variabile.
erythrurus, *Tausch, in Flora*, xix. (1836) 425.—Hab.?
exiguiflorus, Schott, Aroid. 21. t. 41; Prod. Aroid. 561 = scandens.
fallax, Schott, Prod. Aroid. 560 = scandens.
flexuosus, H. B. & K. Nov. Gen. et Sp. vii. 151 = Anthurium flexuosum.
foetidus, Ait. Hort. Kew. ed. I. iii. 319 = Symplocarpus foetidus.
giganteus, Roxb. Hort. Beng. 11; Fl. Ind. i. 434 = Epipremnum giganteum.
glaucus, *Link & Otto, ex Steud. Nom.* ed. II. ii. 391, nomen.—Mexic.
glaucus, Schott, ex Kunth, Enum. Pl. iii. 73 = Anthurium coriaceum.
glaucus, Wall. Pl. As. Rar. ii. 45. t. 156 = Rhaphidophora glauca.
gracilis, *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 433.—Ins. Amboin.
gracilis, Rudge, Pl. Gui. i. 23. t. 32 = Anthurium Rudgeanum.
grandiflorus, Steud. Nom. ed. II. ii. 391, sphalm. = seq.
grandifolius, Jacq. Coll. iv. 121 = Anthurium grandifolium.
Harrisii, R. Grah. in Edinb. Phil. Journ. (Apr. 1826) 353 = Anthurium Harrisii.
hederaceus, *Aubl. Pl. Gui.* ii. 840.—Guian.
hederaceus, Zoll. & Mor. Syst. Verz. Zoll. 83 = Scindapsus hederaceus.
heptaphyllus, Vell. Fl. Flum. ix. t. 125 = Anthurium variabile.
heterophyllus, Roxb. Hort. Beng. 11; Fl. Ind. i. 437 = Lasia aculeata.

POTHOS:—
Hookeri, *Schott, Aroid.* 23. t. 46.—Zeylan.
Horsfieldii, Miq. Fl. Ind. Bat. iii. 178 = longifolius.
inaequilaterus, Engl. in DC. Monog. Phan. ii. 88 = Cumingianus.
insignis, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 267. —Borneo.
Junghuhnianus, Schott, Prod. Aroid. 567 = macrophyllus.
Junghuhnii, De Vriese, in Miq. Pl. Jungh. 104 = macrophyllus.
Korthalsianus, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 284 = Cumingianus.
laciniatus, Mart. & Gal. in Bull. Acad. Brux. x. 1. (1843) 122 = Anthurium podophyllum.
lanceolatus, Linn. Sp. Pl. ed. II. 1373 = Anthurium lanceolatum.
Lasia, Roxb. Hort. Beng. 11; Fl. Ind. i. 438 = Lasia aculeata.
latifolius, Linn. Syst. ed. X. 1252 (Quid?).—Ins. Amboin.
leptospadix, De Vriese, in Miq. Pl. Jungh. 105 = Zollingerianus.
leptostachyus, *Schott, Prod. Aroid.* 571.—Borneo.
leucocarpus, Hort. ex Kunth, Enum. Pl. iii. 68 = Anthurium scandens.
leucophaeus, Poepp. ex Engl. in Mart. Fl. Bras. iii. 11. 104 = Spathiphyllum cannaeforme.
longifolius, Hoffm. Verz. Pfl. 102; Nachtr. iii. 53 = Anthurium longifolium.
longifolius, *Link & Otto, ex Steud. Nom.* ed. II. ii. 391.—Mexic.
longifolius, *Presl, Epim. Bot.* 242.—Ins. Philipp.
longipes, *Schott, Aroid.* 23. t. 47.—Austral.
Loureirii, *Hook. & Arn. Bot. Beech. Voy.* 220.—China; Ins. Philipp.; Austral.
lucidus, Hort. ex Kunth, Enum. Pl. iii. 73 = Anthurium lucidum.
luzonensis, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 19.—Ins. Philipp.
macrophyllus, *De Vriese, in Miq. Pl. Jungh.* 104.—Java; Sumatra.
macrophyllus, Sw. Prod. Veg. Ind. Occ. 33 = Anthurium Selloum.
macrophyllus, Willd. Sp. Pl. i. 686 = Anthurium grandifolium.
macrostachyus, *Zoll. & Mor. Syst. Verz. Zoll.* 83 (sp. dub).—Java.
malaianus, Miq. Fl. Ind. Bat. Suppl. 596 = Anadendrum montanum.
marginatus, Wall. Cat. n. 4443 = Anadendrum marginatum.
maximus, Desf. Cat. Hort. Par. ed. III. 8, 386 = Anthurium Fontanesii.
membranuliferus, De Vriese, ex Engl. in DC. Monog. Phan. ii. 196 = Anthurium podophyllum.
membranuliferus, *Hort. Spaarn-Bergens.* 41; *ex Walp. Ann.* i. 762.—Hab.?
microphyllus, Hook. Bot. Mag. t. 2953 = Anthurium microphyllum.
microphyllus, *Presl, Epim. Bot.* 243.—Burma; China.
microstachyus, H. B. & K. Nov. Gen. et Sp. i. 75 = Anthurium microstachyum.
miniatus, Zipp. ex Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 129 = Epipremnum asperatum.
monopetalus, Reinw. ex Miq. Fl. Ind. Bat. iii. 188 = Anadendrum affine.
Mottleyanus, *Schott, Aroid.* 25. t. 55.—Borneo.
myosuroides, H. B. & K. Nov. Gen. et Sp. i. 75 = Anthurium myosuroides.
nervosus, Willd. ex Kunth, Enum. Pl. iii. 51 = Philodendron Ochrostemon.
nigrescens, Zipp. ex Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 131 = Zippelii.
obliquus, *Wall. Cat.* n. 4446 (Quid?).—Ind. or.
oblongifolius, *Tausch, in Flora*, xix. (1836) 425.—Hab.?
obtusifolius, Ait. Hort. Kew. ed. II. i. 269 = Anthurium obtusifolium.
odoratus, Anders. ex Sims, Bot. Mag. t. 603 = Spathiphyllum cannaeforme.
officinalis, Roxb. Hort. Beng. 11; Fl. Ind. i. 431 = Scindapsus officinalis.
ovatus, Walt. Fl. Carol. 224 = Orontium aquaticum.
oxyphyllus, *Miq. Fl. Ind. Bat.* iii. 181.—Java.
palmatus, Linn. Sp. Pl. ed. II. 1374 = Anthurium palmatum.

POTHOS:—
panduraeformis, H. B. & K. Nov. Gen. et Sp. i. 78 = Philodendron panduraeforme.
papuanus, *Becc. ex Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 267.—N. Guin.
parasiticus, Vell. Fl. Flum. ix. t. 121 = Anthurium Miquelianum.
pedatus, H. B. & K. Nov. Gen. et Sp. i. 78. t. 20 = Anthurium pedatum.
Peepla, Roxb. Hort. Beng. 11; Fl. Ind. i. 433 = Rhaphidophora Peepla.
pentandrus, *Dennst. Schluess. Hort. Malab.* 33.—Ind. or.
pentaphyllus, Willd. Sp. Pl. i. 687 = Anthurium pentaphyllum.
pertusus, Roxb. Hort. Beng. 83; Fl. Ind. i. 434 = Rhaphidophora pertusa.
pinnatifidus, Roxb. Hort. Beng. 83; Fl. Ind. i. 437 = Epipremnum mirabile.
pinnatus, Linn. Sp. Pl. ed. II. 1324 = Epipremnum mirabile.
pinnatus, Wall. Cat. n. 4438; Blanco, Fl. Filip. ed. I. 646 = Epipremnum medium.
Pipla, Schult. Mant. iii. 295 = Rhaphidophora Peepla.
platynevron, Desf. Cat. Hort. Par. ed. III. 8, 386 = Caladium platynevron.
podophyllus, Cham. & Schlecht. in Linnaea, vi. (1831) 22 = Anthurium podophyllum.
Putorii, Bartram, Fl. Virg. 60 = Symplocarpus foetidus.
quinquenervius, *H. B. & K. Nov. Gen. et Sp.* i. 76.—Am. austr.
reflexus, Hoffm. ex Kunth, Enum. Pl. iii. 73; Link & Otto, ex Steud. Nom. ed. II. ii. 391 = Anthurium coriaceum.
remotiflorus, *Hook. Ic. Pl.* t. 133.—Zeylan.
reptans, Link & Otto, ex Steud. Nom. ed. II. ii. 391 = Anthurium reptans.
rigidus, *Aubl. Pl. Gui.* ii. 839.—Guian.
Roxburghii, De Vriese, in Miq. Pl. Jungh. 103 = scandens.
Roxburghii, Schott, Aroid. 22. t. 37 = Vriesianus.
rubescens, Link & Otto, ex Steud. Nom. ed. II. ii. 391 = Anthurium rubescens.
rubricaulis, Hort. ex Kunth, Enum. Pl. iii. 69 = Anthurium Harrisii.
rubrinervius, Link, Enum. Hort. Berol. i. 109 = Anthurium rubrinervium.
Rumphii, *Schott, Melet.* 21.—Java.
sagittatus, Sims, Bot. Mag. t. 1584 = Anthurium rubrinervium.
sagittifolius, Rudge, Pl. Gui. 24. t. 34 = Urospatha sagittaefolia.
scandens, D. Don, Prod. Fl. Nep. 21 = Cathcarti.
scandens, Hook. Ic. Pl. t. 175 = Hookeri.
scandens, *Linn. Sp. Pl.* 968.—Ind. or.; Malaya; China.
scandens, Wall. Cat. n. 4435 = Vriesianus.
Schlechtendalii, Mart. & Gal. in Bull. Acad. Brux. x. 1. (1843) 122 = Anthurium Schlechtendalii.
scolopendrinus, Ham. Prod. Pl. Ind. Occ. 16 = Anthurium gracile.
scolopendrioides, *Desf. Tabl.* ed. I. 8.—Hab.?
Scolopendrium, Spreng. Nov. Prov. 34 = Anthurium acaule.
Seemanni, *Schott, in Bonplandia*, v. (1857) 44.—China.
solitarius, Vell. Fl. Flum. iv. t. 123 = Anthurium affine.
spectabilis, *Hort.—Cf. Gard. Chron.* (1869) 587.—Bras.
spinosus, Buch.-Ham. ex Wall. Cat. n. 4447 C = Lasia aculeata.
suaveolens, Desf. Cat. Hort. Par. ed. III. 8, 386 = Anthurium rubrinervium.
subcaulescens, Vell. Fl. Flum. ix. t. 122 = Anthurium coriaceum.
subsagittatus, H. B. & K. Nov. Gen. et Sp. i. 77 = Anthurium subsagittatum.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* 596.—Sumatra.
tener, *Schott, Aroid.* 24; *Prod. Aroid.* 572.—Ins. Amboin.?
tener, Wall. in Roxb. Fl. Ind. ed. Carey, i. 454 = gracilis.
terminalis, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 247 = Loureirii.
Thomsonianus, *Schott, Aroid.* 24. t. 51.—Ind. or.
triphyllus, *Willd. ex Schult. Mant.* iii. 301.—Am. austr.
undatus, Hort. ex Kunth, Enum. Pl. iii. 82 = Anthurium undatum.

POTHOS:—
undulatus, *Hort. ex Steud. Nom.* ed. II. ii. 391, nomen.—Hab.?
venosus, *Willd. ex Schult. Mant.* iii. 300.—Mexic.?
ventricosus, *Hoffmgg. Verz. Pfl.* 102 (Quid?).—Bras.
violaceus, Sw. Prod. Veg. Ind. Occ. 32 = Anthurium scandens.
viridis, *Parment. ex Steud. Nom.* ed. II. ii. 391, nomen.—Hab.?
Vrieseanus, *Schott, Aroid.* 22. t. 36.—Sumatra.
Wallichii, Steud. Nom. ed. II. ii. 391 = Rhaphidophora glauca.
Zippelii, *Schott, in Miq. Ann. Mus. Lugd. Bat.* i. 131.—Malaya; N. Guin.
Zollingeri, Engl. in DC. Monog. Phan. ii. 85, sphalm. = seq.
Zollingerianus, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 19.—Java; Sumatra.

POTHUAVA, Gaudich. Bot. Voy. Bonite, tt. 116, 117 (1844–52) = **Aechmea**, Ruiz & Pav. (Bromel.).
comata, Gaudich. l. c. t. 116 = A. comata.
nudicaulis, Regel, Gartenfl. (1882) 291. t. 1096 = A. nudicaulis.
Skinneri, C. Koch, Wochenschr. iv. (1861) 190.—Guatemala.
spicata, Gaudich. Bot. Voy. Bonite, t. 117 = A. nudicaulis.

POTIMA, R. Hedw. Gen. 143 (1806) = **Faramea**, Aubl. (Rubiac.).

POTTSIA, Hook. & Arn. Bot. Beech. Voy. 198. t. 43 (1836). *APOCYNACEAE*, Benth. & Hook. f. ii. 712.
EUTHODON, Griff. Notul. iv. 84; Ic. Pl. As. t. 458 (1854).
PARAPOTTSIA, Miq. Fl. Ind. Bat. ii. 1080 (1856).
TEYSMANNIA, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 193.
cantonensis, *Hook. & Arn. Bot. Beech. Voy.* 199.—Ind. or.; Java; China.
Hookeriana, Wight, Ic. t. 1306 = cantonensis.
ovata, A. DC. in DC. Prod. viii. 442 = cantonensis.

POUCHETIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 251. *RUBIACEAE*, Benth. & Hook. f. ii. 92.
africana, *DC. Prod.* iv. 393.—Afr. trop.
parviflora, *Benth. in Hook. Niger Fl.* 395.—Afr. trop.
Saxifraga, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 355.—Abyss.

POUPARTIA, Comm. ex Juss. Gen. 372 (1789) = **Spondias**, Linn. (Anacard.).
borbonica, J. F. Gmel. Syst. 728 = S. borbonica.
castanea, Engl. in DC. Monog. Phan. iv. 261 = Sclerocarya castanea.
dulcis, Blume, Bijdr. 1161 = S. dulcis.
excelsa, March. Rév. Anacard. 157 = Sclerocarya caffra.
guianensis, J. F. Gmel. Syst. 1609, sphalm. = *borbonica*.
mangifera, Blume, Bijdr. 1160 = Dracontomelon mangiferum.
minor, March. Rév. Anacard. 157.—Madag.
pinnata, Blanco, Fl. Filip. ed. I. 393 = Dracontomelon mangiferum.
pubescens, March. Rév. Anacard. 157 = S. pubescens.
triphylla, March. l. c. = Lanneoma velutina.

POUROUMA, Aubl. Pl. Gui. ii. 891. t. 341 (1775). *URTICACEAE*, Benth. & Hook. f. iii. 380.
PURUMA, Jaume St. Hil. Expos. ii. 313 (1805).
acuminata, *Mart. Syst. Mat. Med. Bras.* 34, nomen; *et ex Miq. in Mart. Fl. Bras.* iv. 1. 130.—Bras.
acutifolia, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 105.—Bras.
aspera, *Tréc. l. c.* 102.—Guian.
bicolor, *Mart. Syst. Mat. Med. Bras.* 34.—Bras.
cecropiaefolia, *Mart. Reise Bras.* iii. 1130; *et ex Miq. in Mart. Fl. Bras.* iv. 1. 123.—Bras.
cinerascens, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 125.—Bras.
digitata, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 106.—Guian.
edulis, *Dufresne, in Illustr. Hortic.* xx. (1873) 70.—N. Granat.
fuliginea, *Miq. in Mart. Fl. Bras.* iv. 1. 129.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 892. t. 341.—Guian.
heterophylla, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 125.—Bras.

POUROUMA :—
Jussiaeana, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 106.—Peruv.
mollis, *Tréc. l. c.* 102.—Am. trop.
multifida, Tréc. l. c. 107 = cecropiaefolia.
ovata, *Tréc. l. c.* 101.—Bras.
palmata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 29. t. 141. —Peruv.
retusa, Benth. ex Miq. in Mart. Fl. Bras. iv. 1. 128 = tomentosa.
sapida, *Karst. Fl. Columb.* ii. 19. t. 110.—N. Granat.
tomentosa, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 128. —Bras.
triloba, *Klotzsch, in Linnaea*, xx. (1847) 526; *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 104.—Peruv.; Bras.?
velutina, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 130. —Bras.
villosa, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 103.—Guian.

POURRETIA, Ruiz & Pav. Prod. 46. t. 7 (1794) = **Puya**, Molina (Bromel.).
Achupalla, Linden, Cat. n. viii. (1853) 31, nomen = Puya Bonplandiana.
aëranthos, Loisel. Herb. Amat. v. t. 304 = Tillandsia dianthoidea.
alpestris, Poepp. & Endl. Nov. Gen. et Sp. ii. 41. t. 156 = Pitcairnia caerulea.
auruacensis, Linden, Cat. n. viii. (1853) 31.—N. Granat.
Bonplandiana, D. Dietr. Syn. Pl. ii. 1060 = Puya Bonplandiana.
caerulea, Miers, Trav. ii. 531, nomen = Pitcairnia caerulea.
coarctata, Ruiz & Pav. Fl. Per. iii. 34 = Puya chilensis.
ferruginea, Spreng. Syst. ii. 23 = Pitcairnia ferruginea.
floccosa, C. Koch, Wochenschr. iii. (1860) 251 = Pitcairnia floccosa.
floccosa, Linden, Cat. n. viii. (1853) 31.—Hab.?
frigida, Linden, l. c. = Dyckia frigida.
inermis, E. Mey. ex Presl, Rel. Haenk. i. 123 = Pitcairnia inermis.
lanata, H. B. & K. Nov. Gen. et Sp. i. 296 = Puya lanata.
lanuginosa, Hort. ex Regel, in Act. Hort. Petrop. iii. (1875) 125 = Pitcairnia floccosa.
lanuginosa, Ruiz & Pav. Fl. Per. iii. 33 = Puya lanuginosa.
lepidota, Linden, Cat. n. viii. (1853) 31.—Hab.?
magnispatha, Colla, Hort. Ripul. App. ii. 334. t. 19 = Bilbergia speciosa.
paniculata, Ruiz & Pav. Syst. Veg. 81 = Pitcairnia pulverulenta.
pyramidata, H. B. & K. Nov. Gen. et Sp. i. 296 = Puya Bonplandiana.
pyramidata, Ruiz & Pav. Fl. Per. iii. 34 = Puya pyramidata.
rubricaulis, Miers, Trav. ii. 531 = Pitcairnia caerulea?
speciosa, Linden, Cat. n. viii. (1853) 31.—Hab.?
sympaganthera, Ruiz & Pav. Syst. Veg. 82 = Guzmania tricolor.
violacea, Hort. Laurent. ex Regel, in Act. Hort. Petrop. iii. (1875) 125 = Pitcairnia floccosa.
violacea, Linden, Cat. n. viii. (1853) 31.—Hab.?
yuccoides, Linden, l. c. n. lxxxvii. (1871) 25.—Hab.?

POURRETIA, Willd. Sp. Pl. iii. 844 (1801) = **Cavanillesia**, Ruiz & Pav. (Malvac.).
arborea, Willd. l. c. = C. umbellata.
platanifolia, Humb. & Bonpl. Pl. Aequin. ii. 162. t. 133 = C. platanifolia.
tuberculata, Mart. Reise Bras. ii. 582 = C. platanifolia?

POURTHIAEA. Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 146. *ROSACEAE*.
arguta, *Decne. l. c.* 147.—Reg. Himal.
Calleryana, *Decne. l. c.*—China.
coreana, *Decne. l. c. c.* 148.—Japon; Corea.
Cotoneaster, *Decne. l. c.* 149.—Japon.
Hookeri, Decne. l. c. = arguta.
lucida, *Decne. l. c.* 148.—Ins. Formosa.
Oldhamii, *Decne. l. c.* 149.—Japon.
salicifolia, Decne. l. c. 148 = arguta.
Thunbergii, *Decne. l. c.* 149.—Japon.
villosa, *Decne. l. c.* 147.—Japon.
Zollingeri, *Decne. l. c.* 149.—Japon.

POUSLOWIA, Wight, in Hook. Journ. Bot. iii. (1841) 185, 192 (Quid?).

POUTERIA, Aubl. Pl. Gui. i. 85 (1775). *SAPOTACEAE*, Benth. & Hook. f. ii. 633.
CHAETOCARPUS, Schreb. Gen. 75 (1789).
PROZETIA, Neck. Elem. i. 225 (1790).
amazonica, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* xii. (1882) 333.—Bras.
Caimito, Radlk. l. c. 312 = Lucuma Caimito.
chrysophylloides, Radlk. l. c. 333 = Lucuma chrysophylloides.
crassifolia, *Radlk. l. c.* 334.—Bras.
dictyoneura, *Radlk. l. c.* xiv. [1884] 464.—Cuba.
Gardneriana, Radlk. l. c. xii. (1882) 333 = Lucuma Gardneriana.
glomerata, Radlk. l. c. = Lucuma glomerata.
gomphiaefolia, Radlk. l. c. = Lucuma gomphiaefolia.
guianensis, *Aubl. Pl. Gui.* i. 86. t. 33 (= Lucuma psammophila, excl. fruct.).—Guian.
guianensis, Griseb. Fl. Brit. W. Ind. 403 = Sloanea Purdiaei.
lactescens, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 294 = Lucuma glycyphloea.
laevigata, *Radlk. l. c.* xiv. [1884] 457.—Bras.
lasiocarpa, Radlk. l. c. xii. (1882) 333 = Lucuma lasiocarpa.
lateriflora, Radlk. l. c. = Lucuma lateriflora.
laurifolia, Radlk. l. c. = Lucuma laurifolia.
lucens, Radlk. l. c. = Lucuma lucens.
macrocarpa, D. Dietr. Syn. Pl. i. 431 = Labatia macrocarpa.
neriifolia, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 333 = Lucuma neriifolia.
nitida, Radlk. l. c. = Lucuma nitida.
ochrosperma, *Radlk. l. c.* 335.—Bras.
parviflora, Radlk. l. c. 333 = Lucuma parviflora.
psammophila, Radlk. l. c. = Lucuma psammophila.
ramiflora, Radlk. l. c. = Lucuma ramiflora.
rigida, Radlk. l. c. = Lucuma rigida.
salicifolia, Radlk. l. c. = Lucuma Sellowii.
sessiliflora, Poir. Encyc. Suppl. iv. 546 (= *Labatia sessiliflora*).—Ind. occ.
torta, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 333 = Lucuma torta.

POUZOLZIA, Gaudich. in Freyc. Voy. Bot. 503 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 387.
ELKANIA, Schlecht. ex Wedd. in DC. Prod. xvi. 1. 227, in syn. (1869).
GONOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 509.
HYRTANANDRA, Miq. Pl. Jungh. 25 (1851).
LEPTOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 193. t. 57 (1856).
LEUCOCOCCUS, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 311.
MARGAROCARPUS, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 203.
MEMORIALIS, Buch.-Ham. in Wall. Cat. nn. 4598–4601 (1831).
PENTOCNIDE, Rafin. Fl. Tellur. iii. 48 (1836).
STACHYOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 227. t. 55 (1856).
abyssinica, *Blume, Mus. Bot. Lugd. Bat.* ii. 236.—Abyss.
acalyphoides, *Wedd. in DC. Prod.* xvi. 1. 234.—Venezuela.
acuta, Wight, Ic. vi. 39 = Wightii.
alienata, Gaudich. in Freyc. Voy. Bot. 503 = indica.
ambigua, Wight, Ic. vi. 39 = Wightii.
angustifolia, Wight, l. c. 43 = indica.
arachnoidea, Wedd. in DC. Prod. xvi. 1. 228 = macrura.
arnhemica, F. Muell. Fragm. iv. 87 = indica.
aspera, *Wedd. in DC. Prod.* xvi. 1. 233.—Peruv.; Bolivia.
aspera, Wight, Ic. vi. 40. t. 2095. f. 18 = scabra.
auriculata, *Wight, Ic.* vi. 42.—Ind. or.
Bennettiana, *Wight, l. c.* 40.—Ind. or.
bicuspidata, Wight, l. c. 42 = auriculata.
Boiviniana, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 405.—Madag.
borbonica, Wight, Ic. vi. 43 = viminea.
bornensis, Blume, Mus. Bot. Lugd. Bat. ii. 234 = indica.
bracteata, Benn. Pl. Jav. Rar. 67 = caudata.
canescens, Gaudich. in Freyc. Voy. Bot. 503 = laevigata.

POUZOLZIA :—
caudata, *Benn. Pl. Jav. Rar.* 66.—Ind. or.
ceramica, *Blume, Mus. Bot. Lugd. Bat.* ii. 234.—Ins. Molucc.
cinerascens, *Wedd. in DC. Prod.* xvi. 1. 230.—Sumatra.
cochinchinensis, Blume, Mus. Bot. Lugd. Bat. ii. 245 = indica.
concinna, Wight, Ic. vi. 40 = Wightii.
confinis, *Blume, Mus. Bot. Lugd. Bat.* ii. 230.—Ind. or.
conglobata, *Miq. Fl. Ind. Bat.* i. II. 260.—Java.
cordata, *Benn. Pl. Jav. Rar.* 67.—Java.
courtallensis, Wight, Ic. vi. 38. t. 2093. f. 10 = caudata.
cymosa, Wight, l. c. 42 = auriculata.
Dalzellii, Wight, l. c. 41 = pentandra.
diffusa, Wight, l. c. 38. t. 2099. f. 36 = indica.
discolor, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 408.—Peruv.
elegans, *Wedd. in DC. Prod.* xvi. 1. 230.—Ins. Formosa.
elliptica, Wight, Ic. vi. 42 = auriculata.
flaccida, A. Rich. Tent. Fl. Abyss. ii. 259 = Australina Schimperiana.
formicaria, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 407.—Peruv.
Forsteniana, Blume, Mus. Bot. Lugd. Bat. ii. 23 = indica.
fulgens, Wedd. in DC. Prod. xvi. 1. 229 = viminea.
Gardneri, Wight, Ic. t. 2092. f. 5 = Bennettiana.
Gardneriana, Wight, l. c. vi. 39 = Bennettiana.
glabra, Wight, l. c. 41 = pentandra.
glabrata, Wedd. in Arch. Mus. Par. viii. (1855–56) 406 = ovata.
glochidiata, Blume, Mus. Bot. Lugd. Bat. ii. 245 = indica.
glomerata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 492, in obs.; Hassk. Cat. Hort. Bog. Alt. 80 = indica.
guatemalana, *Wedd. in DC. Prod.* xvi. 1. 233.—Guatemala.
guineensis, *Benth. in Hook. Niger Fl.* 518.—Afr. trop.
heterocarpa, Wight, Ic. vi. 40 = Bennettiana.
heterophylla, Blume, Mus. Bot. Lugd. Bat. ii. 237 = indica.
heterophylla, *Wedd. in DC. Prod.* xvi. 1. 231.—Ins. Celebes.
hirta, *Hassk. Cat. Hort. Bog. Alt.* 80.—As. et Austral. trop.
hispida, Benn. Pl. Jav. Rar. 66 = hirta.
Huegeliana, Blume, Mus. Bot. Lugd. Bat. ii. 241 = Bennettiana.
hypericifolia, Blume, l. c. 242 = pentandra.
hypoleuca, *Wedd. in DC. Prod.* xvi. 1. 227.—Afr. trop.
indica, *Gaudich. in Freyc. Voy. Bot.* 503.—As. et Austral. trop.
integrifolia, *Dalz. in Hook. Kew Journ.* iii. (1851) 134.—Ind. or.
Johnsoniana, Wight, Ic. vi. 44. t. 2101. t. 47 = indica.
laevigata, *Gaudich. in Freyc. Voy. Bot.* 503.—Ins. Mascar.
laevis, Wedd. in DC. Prod. xvi. 1. 233 = virgata.
latifolia, *Wedd. l. c.* 234.—Mexic.
longifolia, Wight, Ic. vi. 40 = Wightii.
lucida, *Hassk. Cat. Hort. Bog. Alt.* 80.—Java.
lythroides, F. Muell. Fragm. v. 194 = hirta.
macrura, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 395.—Ins. Philipp.
microceltis, Miq. Fl. Ind. Bat. i. II. 260 = viminea.
microphylla, Wight, Ic. vi. 42 = indica.
minor, Wight, l. c. 43 = indica.
mixta, *Solms, in Sitzb. Ges. Naturf. Fr. Berl.* (1864) 21.—Afr. trop.
mollis, Blume, Mus. Bot. Lugd. Bat. ii. 230 = laevigata.
montana, Miq. Pl. Jungh. 23 = viminea.
muralis, Miq. l. c. 24 = indica.
mysorensis, Wight, Ic. vi. 39 = Bennettiana.
nana, Blume, Mus. Bot. Lugd. Bat. ii. 244 = indica.
neilgherrensis, Wight, Ic. vi. 42 = Wightii.
obliqua, *Wedd. in Arch. Mus. Par.* viii. (1855-56) 405. —Panama.
oblongifolia, Wight, Ic. vi. 42 = Wightii.
obscura, Blume, Mus. Bot. Lugd. Bat. ii. 241 = Bennettiana.

POUZOLZIA :—
occidentalis, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 410.—Nicaragua.
orientalis, *Wedd. l. c.* 413.—Ind. or.
ovalifolia, Wight, Ic. vi. 39 = Bennettiana.
ovalis, Miq. Pl. Jungh. 24 = viminea.
ovata, *Wedd. in DC. Prod.* xvi. 1. 232.—N. Granat.
ovata, Wight, Ic. vi. 41 = Wightii.
parietarioides, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 492 = indica.
parva, Blume, Mus. Bot. Lugd. Bat. ii. 234 = indica.
parvifolia, *Wight, Ic.* vi. 39. t. 2092. f. 1.—Zeylan; Java.
pauciflora, Benn. Pl. Jav. Rar. 67 = indica.
pauciflora, A. Rich. Tent. Fl. Abyss. ii. 259 = Droguetia diffusa.
pentandra, *Benn. Pl. Jav. Rar.* 64.—As. trop.
pilosa, Wight, Ic. vi. 43 = indica.
procridioides, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 412.—Afr. austr.
procumbens, Wight, Ic. vi. 42 = indica.
propinqua, Blume, Mus. Bot. Lugd. Bat. ii. 230 = auriculata.
prostrata, Benn. Pl. Jav. Rar. 67 = indica.
pyramidata, Wight, Ic. vi. 44 = indica.
quadrialata, Wight, l. c. 37. t. 2094. f. 12 = Bennettiana.
quinquenervis, Benn. Pl. Jav. Rar. 66 = hirta.
ramosissima, Wight, Ic. vi. 41 = pentandra.
repens, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 174 = Boehmeria repens.
revoluta, *Blume, Mus. Bot. Lugd. Bat.* ii. 240.—Sumatra.
Rheedii, Wight, Ic. vi. 43 = auriculata.
rostrata, Wight, l. c. 42 = auriculata.
Rothiana, *Blume, Mus. Bot. Lugd. Bat.* ii. 231.—Ind. or.
rotundifolia, Wight, Ic. vi. 42 = auriculata.
rubricaulis, *Wedd. in DC. Prod.* xvi. 1. 229.—Java.
scabra, *Wight, Ic.* vi. 41.—Ind. or.
scabrida, Wight, l. c. 43 = auriculata.
Sonnerati, Gaudich. in Freyc. Voy. Bot. 503 = Phenax vulgaris.
sponiaefolia, Schlecht. ex Blume, Mus. Bot. Lugd. Bat. ii. 244 = Bennettiana.
Stocksii, Wight, Ic. vi. 41 = pentandra.
strigosa, Hassk. Cat. Hort. Bog. Alt. 80 = indica.
suffruticosa, Wight, Ic. vi. 43 = indica.
tenuis, Blume, Mus. Bot. Lugd. Bat. ii. 235 = indica.
ternata, Benn. Pl. Jav. Rar. 66 = Wightii.
tetraptera, Wight, Ic. vi. 44 = indica.
tomentosa, Wight, l. c. 40 = Bennettiana.
trialata, Wight, l. c. 42 = Wightii.
triandra, Blume, Mus. Bot. Lugd. Bat. ii. 241 = parvifolia.
trichophora, Hohen. ex Wedd. in DC. Prod. xvi. 1. 222 = indica.
tuberosa, *Wight, Ic.* vi. 43.—Ind. or.
vesicaria *Wight, l. c.*—Ind. or.
viminea, *Wedd. in DC. Prod.* xvi. 1. 228.—Ind. or.
virgata, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 409.—Bras.
Waitziana, *Blume, Mus. Bot. Lugd. Bat.* ii. 239.—Java.
Walkeriana, Wight, Ic. vi. 41. t. 2095 f. 16 = pentandra.
Wallichiana, Wight, l. c. 41 = Wightii.
Wightii, *Benn. Pl. Jav. Rar.* 66.—Ind. or.
zeylanica, Benn. l. c. 67 = indica.

POZOA, Lag. Gen. et Sp. Nov. 13 (1816) = **Azorella**, Lam. (Umbell.).
coriacea, Lag. l. c.—Reg. Andin.
cuneifolia, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 103 = A. cuneifolia.
denticulata, Meyen, Reise, i. 314 = *coriacea*.
exigua, Hook. f. Handb. N. Zeal. Fl. 87.—N. Zel.
Fragosea, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 102 = A. Muelleri.
Haastii, Hook. f. Handb. N. Zeal. Fl. 88.—N. Zel.
hydrocotylifolia, Bridges, ex Gardn. in Field. Sert. Pl. t. 40.—Chili.
hydrocotyloides, Hook. f. Handb. N. Zeal. Fl. 88.—N. Zel.
incisa, Meyen, Reise, i. 314 = A. incisa.
pallida, T. Kirk, in Trans. N. Z. Inst. x. (1878) 419.—N. Zel.
Ranunculus, Hook. f. Fl. N. Zel. 85, in obs. = A. Ranunculus.

POZOA :—
reniformis, Hook. f. Fl. Antarct. 15. t. 11 = Azorella reniformis.
Roughii, Hook. f. Handb. N. Zeal. Fl. 89.—N. Zel.
trifoliata, Hook. f. Fl. N. Zel. 85 = A. trifoliolata.

POZOPSIS, Hook. Ic. Pl. t. 859 (1852) = **Huanaca**, Cav. (Umbell.).
cordifolia, Hook. l. c. = H. cordifolia.

PRADOSIA, Liais, Brés. 614 (1872) = **Lucuma**, Molina (Sapot.).
glycyphloea, Liais, l. c. = L. glycyphloea.

PRAEALSTONIA, Miers, in Journ. Linn. Soc. xvii. (1879) 291 = **Symplocos**, Linn. (Styrac.).
Arechea, Miers, l. c. = S. Arechea.
cernua, Miers, l. c. 292 = S. cernua.
coccinea, Miers, l. c. 291 = S. coccinea.
coriacea, Miers, l. c. 292 = S. coriacea.
Matthewsii, Miers, l. c. = S. Matthewsii.
mucronata, Miers, l. c. = S. mucronata.
nuda, Miers, l. c. = S. nuda.
octopetala, Miers, l. c. 291 = S. octopetala.
parvifolia, Miers, l. c. 292 = S. parvifolia.
reflexa, Miers, l. c. = S. reflexa.
rufescens, Miers, l. c. = S. rufescens.
serrulata, Miers, l. c. = S. serrulata.
theaeformis, Miers, l. c. 291 = S. Alstonia.
tomentosa, Miers, l. c. 292 = S. tomentosa.

PRAENANTHES, Hook. Bot. Misc. ii. (1831) 221, err. typ. = **Prenanthes**, Vaill. (Compos.).

PRAESEPIUM, Spreng. ex Reichb. Consp. 212[b] (1828), nomen. *ROSACEAE* (Quid?).

PRAETORIA, Baill. Etud. Gén. Euph. 470. t. 11 (1858) = **Pipturus**, Wedd. (Euphorb.).

PRANGOS, Lindl. in Quart. Journ. Sc. xix. (1825) 7.
UMBELLIFERAE, Benth. & Hook. f. i. 904.
ANISOPLEURA, Fenzl, in Flora, xxvi. (1843) 459; et in Endl. Gen. Suppl. iii. 82 (1843).
COLLADONIA, DC. Prod. iv. 240 (1830).
HEPTAPTERA, Marg. & Reut. in Mém. Soc. Phys. Genève. viii. (1838) 302 t. 5.
HETEROPTERA, Steud. Nom. ed. II. i. 756 (1840).
MELIOCARPUS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 84.
PERLEBIA, DC. Mém. Omb. 67 (1829).
PTEROMATHRUM, Koch, in DC. Prod. iv. 239 (1830).
angustifolia, *Nym. Syll.* 165.—Ital.
anisopetala, *DC. Prod.* iv. 240.—Syria.
Arcis-romanae, *Boiss. & Huet, Diagn.* Ser. II. ii. 105.—Armenia.
armena, Kotschy & Boiss. ex Boiss. Fl. Orient. ii. 938 = platychlaena.
asperula, *Boiss. Diagn.* Ser. I. x. 54.—Syria.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 83.—Persia.
cheilanthifolia, Boiss. l. c. 79 = Cachrys cheilanthifolia.
cinerea, Boiss. l. c. 80 = Szowitsii.
colladonioides, *Nym. Syll.* 165.—Graecia.
corymbosa, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 81.—Cappadoc.
cylindracea, *DC. Prod.* iv. 239.—Calabria.
denticulata, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 35.—Galatia.
ferulacea, *Lindl. in Quart. Journ. Sc.* xix. (1825) 7.—Europ. austr.; Oriens.
foeniculacea, C. A. Mey. Verz. Pfl. Cauc. 131 = ferulacea.
Haussknechtii, *Boiss. Fl. Orient.* ii. 940.—Persia.
Hermonis, *Boiss. l. c.* 943.—Syria.
humilis, Fisch. in Ledeb. Fl. Ross. ii. 359 = Szowitsii.
lophoptera, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 82.—As. Min.
macrocarpa, *Boiss. l. c.* 77.—Persia.
meliocarpoides, *Boiss. l. c.* 81.—As. Min.
microcarpa, *Boiss. l. c.* 83.—Persia.
odontoptera, *Boiss. l. c.* 78.—Persia.
ovatifolia, Boiss. l. c. 79 = odontoptera.
pabularia, *Lindl. in Quart. Journ. Sc.* xix. (1825) 7.—Reg. Himal.
pauciradiata, Boiss. & Hohen. Diagn. Ser. I. x. 55 = Trachydium elbrusense.

PRANGOS :—
Pestalozzae, Boiss. Diagn. Ser. I. x. 55 = meliocarpoides.
peucedanifolia, *Fenzl, in Flora*, xxvi. (1843) 463.—Mesopot.
platychlaena, *Boiss. in Tchihat. Asie Min. Bot.* i. 457.—As. Min.
pumila, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 77.—As. Min.
stenoptera, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 104 = ferulacea.
Szowitsii, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 78.—Persia.
thapsioides, DC. Prod. iv. 240 = Elaeoselinum foetidum.
triquetra, *Nym. Syll.* 313.—Europ.
tuberculata, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 943.—Persia.
Uechtritzii, *Boiss. & Haussk. l. c.* 938.—As. Min.
uloptera, *DC. Prod.* iv. 239.—Persia.

PRARAVINIA, Korth. Verh. Nat. Gesch. Bot. 189. t. 41 (1842?). *RUBIACEAE*, Benth. & Hook. f. ii. 75.
PARAVINIA, Hassk. in Flora, xxxi. (1848) 580.
densiflora, *Korth. Verh. Nat. Gesch. Bot.* 190. t. 41.—Borneo.

PRASANTHEA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 467 = **Paliavana**, Vand. (Gesner.).
prasinata, Decne. l. c. = Paliavana sericiflora.

PRASITELES, Salisb. Gen. Pl. Fragm. 102 (1866) = **Narcissus**, Linn. (Amaryll.).
viridiflorus, Salisb. l. c. = N. viridiflorus.

PRASIUM, Linn. Gen. ed. I. 345 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1217.
LEVINA, Adans. Fam. ii. 190 (1763).
coccineum, Walt. Fl. Carol. 166 = Physostegia virginiana.
creticum, Reichb. ex Nym. Consp. 564 = majus.
glabrum, Gaudich. in Freyc. Voy. Bot. 452. t. 64 = Phyllostegia glabra.
grandiflorum, Gaudich. l. c. 453. t. 65. f. 2 = Phyllostegia grandiflora.
hirsutum, Poir. Encyc. v. 611 = Stachys Swainsoni, aethiopica.
incarnatum, Walt. Fl. Carol. 165 = Macbridea pulchra.
javanicum, Blume, Bijdr. 840 = Gomphostemma javanicum.
laetum, Salisb. Prod. 89 = majus.
macrophyllum, Gaudich. in Freyc. Voy. Bot. 453. t. 65. f. 3 = Phyllostegia parviflora.
majus, *Linn. Sp. Pl.* 601.—Reg. Mediterr.
medium, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838), reimpr. 13.—Ins. Mader.
melissaefolium, Roxb. Hort. Beng. 45; Fl. Ind. iii. 26 = Gomphostemma melissaefolium.
minus, Linn. Sp. Pl. 601 = majus.
minus, Vahl, ex Walp. Rep. iii. 819 = Stachys Swainsoni.
parviflorum, Gaudich. in Freyc. Voy. Bot. 453. t. 65. f. 1 = Phyllostegia parviflorum.
phlomoides, Reinw. ex Blume, Bijdr. 840 = Gomphostemma phlomoides.
purpureum, Walt. Fl. Carol. 166 = Physostegia virginiana.

PRASKOINON, Rafin. Fl. Tellur. iv. 29 (1836) = **Allium**, Tourn. (Liliac.).
longifolium, Rafin. l. c. = A. longifolium.

PRASOPEPON, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 26 = **Cucurbitella**, Walp.
cucumifolius, Griseb. in Goett. Abh. xix. (1874) 146 = C. cucumifolia.
Duriaei, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 27. t. 2 = C. Duriaei.

PRASOPHYLLUM, R. Br. Prod. 317 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 608.
GENOPLESIUM, R. Br. Prod. 319 (1810).
affine, Lindl. Gen. et Sp. Orch. 516 = fuscum.
alpinum, R. Br. Prod. 318 = fuscum.
ansatum, *Fitzg. in Journ. Bot.* xxiii. (1885) 135.—Austral.

PRASOPHYLLUM :—
Archeri, *Hook. f. Fl. Tasman.* ii. 14. t. 113 B.—Austral.
attenuatum, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
australe, *R. Br. Prod.* 318.—Austral.
australe, Lindl. Gen. et Sp. Orch. 514 = elatum.
Baueri, Poir. Encyc. Suppl. iv. 547 (*Baveri*) = rufum.
brachystachyum, *Lindl. Gen. et Sp. Orch.* 513.—Tasman.
brevilabre, *Hook. f. Fl. Tasman.* ii. 11. t. 110 A.—Austral.
Brounii, Reichb. f. Beitr. Syst. Pfl. 16 = elatum ?
calopterum, *Reichb. f. in Linnaea*, xli. (1877) 57.—N. Caled.
Colensoi, *Hook. f. Fl. N. Zel.* i. 241.—N. Zel.
cucullatum, *Reichb. f. Beitr. Syst. Pfl.* 59.—Austral.
cyphochilum, *Benth. Fl. Austral.* vi. 340.—Austral.
densum, *Fitzg. in Journ. Bot.* xxiii. (1885) 135.—Austral.
despectans, *Hook. f. Fl. Tasman.* ii. 13. t. 113 A.—Austral.
Drummondii, Reichb. f. Beitr. Syst. Pfl. 60 = elatum ?
elatum, *R. Br. Prod.* 318.—Austral.
eriochilum, *Fitzg. in Journ. Bot.* xxiii. (1885) 135.—Austral.
filiforme, *Fitzg. l. c.* 137.—Austral.
Fimbria, *Reichb. f. Beitr. Syst. Pfl.* 60.—Austral.
fimbriatum, *R. Br. Prod.* 319.—Austral.
flavum, *R. Br. l. c.* 318.—Austral.
fuscum, *R. Br. l. c.*—Austral.
gibbosum, *R. Br. l. c.*—Austral.
giganteum, Endl. in Lehm. Pl. Preiss. ii. 12 = P. Fimbria.
giganteum, Lindl. Swan River App. 54 = elatum.
gracile, Lindl. l. c. = macrostachyum.
hians, *Reichb. f. Beitr. Syst. Pfl.* 61.—Austral.
intricatum, *C. Stuart, ex Benth. Fl. Austral.* vi. 346.—Austral.
laminatum, *Fitzg. in Journ. Bot.* xxiii. (1885) 136.—Austral.
Lindleyanum, Reichb. f. Beitr. Syst. Pfl. 58 = fuscum ?
longisepalum, *Fitzg. in Journ. Bot.* xxiii. (1885) 136.—Austral.
lutescens, Lindl. Gen. et Sp. Orch. 514 = australe.
macrostachyum, *R. Br. Prod.* 318.—Austral.
macrotys, Endl. in Lehm. Pl. Preiss. ii. 12 = seq. ?
macrotys, Lindl. Swan River App. 54 = elatum.
nigricans, *R. Br. Prod.* 319.—Austral.
nigricans, Endl. in Lehm. Pl. Preiss. ii. 12 = macrostachyum.
nudiscapum, Hook. f. Fl. Tasman. ii. 13 = brachystachyum.
nudum, Hook. f. Fl. N. Zel. i. 242 = rufum.
nudum, *Hook. f. Fl. Tasman.* ii. t. 113.—Austral.
ovale, *Lindl. Swan River App.* 54.—Austral.
parvifolium, *Lindl. l. c.*—Austral.
patens, *R. Br. Prod.* 318.—Austral.
plumaeforme, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
pumilum, *Hook. f. Fl. N. Zel.* i. 242.—N. Zel.
reflexum, *Fitzg. in Journ. Bot.* xxiii. (1885) 137.—Austral.
rostratum, Lindl. Gen. et Sp. Orch. 516 = fuscum.
rufum, *R. Br. Prod.* 319.—Austral.
striatum, *R. Br. l. c.* 318.—Austral.
Tepperi, *F. Muell. in Tepp. Pl. Ardross.* (Oct. 1880); *ex ej. Census*, 140.—Austral.
triangulare, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
truncatum, Lindl. Gen. et Sp. Orch. 513 = patens.
tunicatum, Hook. f. Fl. N. Zel. i. 242 = nudum.
viride, *Fitzg. in Journ. Bot.* xxiii. (1885) 135.—Austral.
Woollsii, *F. Muell. Fragm.* v. 100.—Austral.

PRASOXYLON, M. Roem. Syn. Hesper. 83, 101 (1846) = **Dysoxylum**, Blume (Meliac.).
alliaceum, M. Roem. l. c. 101 = D. alliaceum.

PRASUM, Linn. Syst. ed. I (1735), nomen = **Prasium**, Linn.

PRATIA, G. Don, Gen. Syst. iii. 699 (1834), partim = **Lobelia**, Plum. (Campanul.).

PRATIA, Gaudich. in Freyc. Voy. Bot. 456. t. 79 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 550.
Colensoa, Hook. f. Fl. N. Zel. i. 156 (1853).

PRATIA :—
Piddingtonia, A. DC. in DC. Prod. vii. 341 (1839).
Speirema, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 27.
angulata, *Hook. f. Fl. Antarct.* i. 43.—N. Zel.
arenaria, *Hook. f. l. c.* 41. t. 29.—Ins. Auckl.
atacamensis, *Phil. Fl. Atac.* 34.—Chili.
begoniaefolia, G. Don, Gen. Syst. iii. 699 = seq.
begonifolia, *Lindl. Bot. Reg.* t. 1373.—Reg. Himal.; Malaya.
boliviensis, *A. DC. in DC. Prod.* vii. 340.—Bolivia.
corymbosa, G. Don, Gen. Syst. iii. 699 = Lobelia corymbosa.
Cunninghamii, Hook. f. Fl. Antarct. i. 42 = erecta.
elliptica, *Hook. f. l. c.* 43.—Reg. Argent.
erecta, *Gaudich. in Freyc. Voy. Bot.* 456. t. 79.—Austral.
gelida, *Benth. Fl. Austral.* iv. 132.—Austral.
glandulifera, *Wedd. Chlor. And.* ii. 11. t. 45.—Peruv.
hederacea, *Cham. in Linnaea*, viii. (1833) 212.—Bras.
irrigua, *Benth. Fl. Austral.* iv. 132.—Austral.
linnaeoides, *Hook. f. Handb. N. Zeal. Fl.* 172.—N. Zel.
longiflora, *Hook. f. Fl. Antarct.* ii. 325.—Chili.
macrodon, *Hook. f. Handb. N. Zeal. Fl.* 172.—N. Zel.
montana, *Hassk. Cat. Hort. Bog. Alt.* 106.—Java.
nummularia, Benth. ex Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 210 = begonifolia.
nummularia, *A. Br. & Aschers. Ind. Sem. Hort. Berol.* (1861) *App.* 6.—Chili.
oligophylla, *Wedd. Chlor. And.* ii. 10. t. 45.—Peruv.; Bolivia.
pedunculata, *Benth. Fl. Austral.* iv. 133.—Austral.
pencana, *Phil. in Anal. Univ. Chil.* (1861) 53.—Chili.
perpusilla, *Hook. f. Handb. N. Zeal. Fl.* 172.—N. Zel.
platycalyx, *Benth. Fl. Austral.* iv. 132.—Austral.
puberula, *Benth. l. c.* 133.—Austral.
radicans, G. Don, Gen. Syst. iii. 700 = Lobelia radicans.
reniformis, *Kanitz, in Mart. Fl. Bras.* vi. iv. 136.—Bras.
repens, *Gaudich. in Freyc. Voy. Bot.* 456. t. 79.—Reg. Magell.
repens, Wedd. Chlor. And. ii. 9 = Hypsela reniformis.
rugulosa, Don, ex Steud. Nom. ed. II. ii. 392 = angulata.
serpyllacea, Presl, Prod. Monog. Lobel. 46 = hederacea.
subsessilis, *Wedd. Chlor. And.* ii. 10.—Bolivia.
Thunbergii, G. Don, Gen. Syst. iii. 700 = Lobelia radicans.
zeylanica, Hassk. Cat. Hort. Bog. Alt. 106 = begonifolia.

PRATICOLA, Ehrh. Beitr. iv. 146 (1789) = **Thalictrum**, Tourn. (Ranunc.).

PRAXELIS, Cass. in Dict. Sc. Nat. xliii. 261 (1826) = **Eupatorium**, Tourn. (Compos.).
capillaris, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 254 = E. capillare.
grandifolium, Sch. l. c. = E. kleinioides.
hecatantha, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 343 = E. kleinioides.
kleinioides, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 254 = E. kleinioides.
paucidentata, Sch. Bip. l. c. = E. kleinioides.
pedunculares, Sch. Bip. l. c. = E. kleinioides.
Riedelii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 342 = E. capillare.
sideritis, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 254 = E. kleinioides.
villosa, Cass. in Dict. Sc. Nat. xliii. 261 = E. urticaefolium.

PREAUXIA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 250 (1836–46) = **Chrysanthemum**, Tourn. (Compos.).
canariensis, Sch. Bip. l. c. 252.—Ins. Canar.
Dugourii, C. Bolle, in Bonplandia, vii. (1859) 297.—Ins. Teneriff.
jacobaeifolia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 251. t. 90.—Ins. Canar.

PREISSIA, Opiz, Seznam, 79 (1852) = **Avena**, Linn. (Gramin.).
strigosa, Opiz, l. c. = A. strigosa.

PREMNA, Linn. Mant. ii. 154 (1771). *VERBENACEAE*, Benth. & Hook. f. ii. 1152.
Baldingera, Dennst. Schluess. Hort. Malab. 31 (1818).
Gumira, Rumph. ex Hassk. in Flora, xxv. (1842) ii. Beibl. 26.
Holochiloma, Hochst. l. c. xxiv. (1841) 371.
Phaenicanthus, Thw. Enum. Pl. Zeyl. 242, in syn. (1861).
Scrophularioides, Forst. f. Prod. 91 (1786).
abbreviata, *Miq. Fl. Ind. Bat.* ii. 892.—Java.
acaulis, Wall. Cat. n. 1776 = herbacea.
acuminata, *R. Br. Prod.* 512.—Austral.
adenosticta, *Schau. in DC. Prod.* xi. 630.—Ins. Philipp.
amplectans, *Wall. Cat.* n. 1762.—Ind. or.
angustata, Wall. l. c. 1763 = amplectens.
arborea, Roth, Nov. Pl. Sp. 287 = Gmelina arborea.
asperrima, *Wall. ex Steud. Nom.* ed. II. ii. 392, nomen (Quid ?).—Ind. or.
attenuata, R. Br. Prod. 512 = obtusifolia.
barbata, *Wall. Cat.* n. 1768.—Reg. Himal.
bengalensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 577.—Ind. or.
bracteata, *Wall. Cat.* n. 1172.—Reg. Himal.
cana, Wall. l. c. 1782 = barbata.
cardiophylla, *Schau. in DC. Prod.* xi. 638.—Ins. Philipp.
ceramensis, *Miq. Fl. Ind. Bat.* ii. 893.—Ins. Ceram.
cordata, Blanco, Fl. Filip. ed. I. 489 = tomentosa.
cordata, R. Br. Prod. 512 = acuminata.
cordifolia, Brand. For. Fl. 367 = scandens.
cordifolia, J. Grah. Cat. Pl. Bomb. 155 = corymbosa.
cordifolia, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 78.—Malaya.
cordifolia, Wight, Ic. t. 1483 = coriacea.
coriacea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 573.—Ind. or.
corymbosa, *Rottl. & Willd. in Ges. Naturf. Fr. Neue Schrift.* iv. (1803) 187.—Ind. or.
Cumingiana, *Schau. in DC. Prod.* xi. 634.—Ins. Philipp.
cyclophylla, *Miq. Fl. Ind. Bat.* ii. 899.—Java.
Dallachyana, *Benth. Fl. Austral.* v. 59.—Austral.
densiflora, Wall. Cat. n. 1773 = corymbosa.
divaricata, *Wall. l. c.* n. 1781.—Burma; Malaya.
Doncaria, Buch.-Ham. ex Wall. l. c. 2647 = cordifolia.
esculenta, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 81.—Ind. or.
ferruginea, *A. Rich. Fl. Tent. Fl. Abyss.* ii. 172.—Abyss.
flavescens, *Buch.-Ham.. ex Wall. Cat.* n. 2649.—Ind. or.; Malaya.
flavescens, Juss. in Ann. Mus. Par. vii. (1806) 77 = tomentosa.
flavida, *Miq. Fl. Ind. Bat. Suppl.* 570.—Sumatra.
foetida, *Reinw. ex Blume, Bijdr.* 816.—Java.
Gaudichaudii, *Schau. in DC. Prod.* xi. 631.—Ins. Mariann.
glaberrima, *Wight, Ic.* t. 1484.—Ind. or.
glycycocca, F. Muell. Fragm. iii. 36 = obtusifolia.
Goeringii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxiv. (1863) ii. 216.—Java *vel* Japon. ?
grossa, *Wall. Cat.* n. 1764.—Burma.
herbacea, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 80.—Ind. or.
hispida, *Benth. in Hook. Niger Fl.* 485.—Afr. trop.
integerrima, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 577, sphalm. = glaberrima.
integrifolia, *Linn. Mant.* ii. 252.—Ind. or.; Malaya.
integrifolia, Willd. in Ges. Naturf. Fr. Neue Schrift. iv. (1803) 187 = divaricata.
interrupta, *Wall. Cat.* n. 1778.—Reg. Himal.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 97.—Japon.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 575.—Reg. Himal.
laevigata, *Miq. Fl. Ind. Bat.* ii. 895.—Sumatra.
lanata, Steud. Nom. ed. I. 137, in syn. = tomentosa.
latifolia, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 76.—Ind. or.
latifolia, Thw. Enum. Pl. Zeyl. 242 = tomentosa.
leucostoma, *Miq. Fl. Ind. Bat.* ii. 899.—Java.

PREMNA:—
leucostoma, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 159 = nauseosa.
limbata, *Benth. Fl. Austral.* v. 59.—Austral.
longifolia, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 79.—Ind. or.
longifolia, Wall. Cat. n. 1771 = bengalensis.
lucidula, Kurz, For. Fl. Brit. Burma. ii. 263 = divaricata.
lucidula, *Miq. Fl. Ind. Bat.* ii. 898.—Java.
macrophylla, *Wall. Cat.* n. 1765.—Burma.
mariannarum, *Schau. in DC. Prod.* xi. 632.—Ins. Mariann.
media, R. Br. Prod. 512 = integrifolia.
micrantha, *Schau. in DC. Prod.* xi. 635.—Reg. Himal.
micrantha, Thw. Enum. Pl. Zeyl. 242 = Thwaitesii.
microphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 217.—China.
milleflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 576.—Assam.
mollissima, Roth, Nov. Pl. Sp. 286 = latifolia.
mucronata, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 80 = latifolia.
multiflora, Zipp. ex Span. in Linnaea, xv. (1841) 330, nomen (Quid ?).—Ins. Timor.
nauseosa, *Blanco, Fl. Filip.* ed. I. 489.—Ins. Philipp.
oblongata, *Miq. Fl. Ind. Bat.* ii. 893.—Java.
obtusifolia, *R. Br. Prod.* 512.—Austral.
odorata, Blanco, Fl. Filip. ed. I. 488 = pubescens.
opulifolia, *Miq. Fl. Ind. Bat.* ii. 898.—Java.
ovalifolia, Wall. Cat. n. 1782 = integrifolia.
ovata, R. Br. Prod. 512 = integrifolia.
parasitica, *Blume, Bijdr.* 816.—Burma; Malaya.
parvifolia, *Roth, Nov. Sp. Pl.* 288.—Ind. or.
philippinensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxiv. (1863) II. 215.—Ins. Philipp.
pinguis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 579.—Reg. Himal.
populifolia, *Miq. Fl. Ind. Bat. Suppl.* 570.—Sumatra.
procumbens, *Moon, ex Wall. Cat.* n. 1780; *et Cat. Pl. Ceyl.* 45.—Zeylan.
pubescens, *Blume, Bijdr.* 816.—Java.
punctulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 575.—Malacca.
punduana, *Wall. Cat.* n. 2651.—Reg. Himal.
purpurascens, *Thw. Enum. Pl. Zeyl.* 242.—Zeylan.
pygmaea, Wall. Cat. n. 1776 = herbacea.
pyramidata, *Wall. l. c.* n. 1779.—Burma.
quadrifolia, *Schum. & Thonn. Beskr. Guin. Pl.* 275.—Afr. trop.
racemosa, Gamble, Darjeeling List Trees, 61 = bracteata.
racemosa, *Wall. Cat.* n. 1777.—Reg. Himal.; Burma.
ramosa, Wall. l. c. 1774 = scandens.
resinosa, *Schau. in DC. Prod.* xi. 637.—Afr. trop.
reticulata, Juss. in Ann. Mus. Par. vii. (1806) 68 = Petitia domingensis.
Roxburghiana, *Wall. Cat.* n. 2648.—Ind. or.
rufidula, *Miq. Fl. Ind. Bat. Suppl.* 569.—Sumatra.
salvifolia, Spreng. Syst. ii. 755 = Pityrodia salvifolia.
sambucina, Wall. Cat. n. 1775 = integrifolia.
scandens, Boj. Hort. Maurit. 257 = serratifolia.
scandens, Dalz. & Gibs. Bomb. Fl. 199 = coriacea.
scandens, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 82.—Ind. or.; Burma.
scandens, Wall. Cat. n. 1774, partim = micrantha.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 263.—Afr. trop.
serratifolia, Blanco, Fl. Filip. ed. II. 342 = foetida.
serratifolia, Linn. Mant. ii. 253 = integrifolia.
spinosa, Roxb. Hort. Beng. 46; Fl. Ind. iii. 77 = integrifolia.
subcordata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 216 = integrifolia.
syringaefolia, Zipp. ex Span. in Linnaea, xv. (1841) 330, nomen (Quid ?).—Ins. Timor.
taitensis, *Schau. in DC. Prod.* xi. 638.—Ins. Tahiti.
thrysiflora, Heyne, ex Wall. Cat. n. 1770 = Wightiana.
Thwaitesii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 579.—Zeylan.
thyrsoidea, *Wight, Ic.* t. 1485.—Ind. or.
tiliaefolia, Zipp. ex Span. in Linnaea, xv. (1841) 330 = timoriana.
timoriana, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 402.—Ins. Timor.
tomentosa, Kurz, For. Fl. Brit. Burma, ii. 260 = pyramidata.
tomentosa, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 581 = Gmelina arborea.
tomentosa, *Willd. Sp. Pl.* iii. 314.—Ind. or.

PREMNA:—
Tracyana, F. Muell. Fragm. v. 61 = Clerodendron Tracyanum.
trichostoma, Miq. Fl. Ind. Bat. ii. 892 = parasitica.
truncata, Turcz. in Bull. Soc. Nat. Mosc. xxxiv. (1863) II. 215 = corymbosa.
verticillata, *Poir. Encyc. Suppl.* i. 348.—Ind. or. ?
vestita, *Schau. in DC. Prod.* xi. 631.—Ins. Philipp.
viburnoides, Kurz, For. Fl. Brit. Burma, ii. 261 = latifolia.
viburnoides, *A. Rich. Tent. Fl. Abyss.* ii. 171.—Abyss.
viburnoides, Wall. Cat. n. 2646 C = integrifolia.
villosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 573.—Ind. or.
villosa, Moon, Cat. Pl. Ceyl. 45 = procumbens.
Wightiana, *Schau. in DC. Prod.* xi. 635.—Ind. or.
zanzibarensis, *Vatke, in Linnaea*, xlviii. (1880–82) 531.—Afr. trop.

PRENANTHES, Vaill. ex Linn. Gen. ed. I. 235 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 527.
ESOPON, Rafin. Fl. Ludov. 143 (1817).
HARPALYCE, D. Don, in Edinb. N. Phil. Journ. (Jan.-Mar. 1829) 308.
HYLETHALE, Link, Handb. i. 788 (1829).
NABALUS, Cass. in Dict. Sc. Nat. xxxiv. 94 (1825).
NARBALIA, Rafin. New Fl. Am. iv. 34 (1836).
acanthifolia, Willd. Sp. Pl. iii. 1542 = Lactuca acanthifolia.
acaulis, Roxb. Hort. Beng. 60; Fl. Ind. iii. 403 = Crepis acaulis.
affinis, *Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 42.—Java.
alata, *D. Dietr. Syn. Pl.* iv. 1309; *A. Gray, Syn. Fl. N. Am.* i. II. 435.—Am. bor.
alata, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 274 = Hookeri.
alba, Bigel. ex DC. Prod. vii. 242 = altissima.
alba, *Linn. Sp. Pl.* 798.—Am. bor.
alba, Walt. Fl. Carol. 193 = P. Serpentaria.
alliariaefolia, DC. Prod. vii. 195 = Brunoniana.
altissima, *Linn. Sp. Pl.* 797.—Am. bor.
altissima, Pall. Reise, iii. 655 = Lactuca sagittata.
amabilis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 842.—Ins. Socotra.
amplexicaulis, Mill. Gard. Dict. ed. VIII. n. 4 = purpurea.
aphylla, Nutt. Gen. Am. ii. 123 = Lygodesmia aphylla.
arborea, Brouss. Elench. Hort. Monsp. (1805) 47 = Sonchus arboreus.
aspera, *Michx. Fl. Bor. Am.* ii. 83.—Am. bor.
aspera, Schrad. ex Willd. Sp. Pl. iii. 1539; et ex DC. Prod. vii. 141 = Chondrilla aspera.
asplenifolia, Willd. Sp. Pl. iii. 1540 = Launaea asplenifolia.
attica, Spreng. Syst. iii. 654 = Chondrilla ramosissima.
autumnalis, Walt. Fl. Carol. 193 = virgata.
Boottii, *D. Dietr. Syn. Pl.* iv. 1309; *A. Gray, Syn. Fl. N. Am.* i. II. 435.—Am. bor.
Brunoniana, *Wall. Cat.* n. 3278.—Reg. Himal.
bulbosa, DC. Fl. Fr. iv. 7 = Crepis bulbosa.
callosa, *Klatt, in Sitzb. Akad. Muench.* (1878) 98.—Tibet.
Candolleana, Wall. Cat. n. 3280 = Ainsliaea pteropoda, Crepis silhetensis.
cervicornis, Boiss. Voy. Bot. Espagne, ii. 744 = Launaea spinosa.
chinensis, Thunb. Fl. Jap. 301 = Lactuca Fischeriana.
chondrilloides, Linn. Mant. i. 107 = Chondrilla prenanthoides.
chondrilloides, Pall. Reise, iii. 537 = Chondrilla graminea.
cordata, Willd. Hort. Berol. t. 25 = altissima.
crepidinea, Ell. Sketch, ii. 259 = P. Serpentaria.
crepidinea, *Michx. Fl. Bor. Am.* ii. 84.—Am. bor.
cymosa, *Spreng. Syst.* iii. 657.—Ins. Canar.
debilis, Thunb. Fl. Jap. 300 = Lactuca debilis.
decurrens, Schur, Enum. Pl. Transs. 953 (Index) = Lactuca viminea.
deltoidea, Bieb. Fl. Taur. Cauc. iii. 529 = Lactuca deltoidea.
deltoidea, Ell. Sketch, ii. 257 = altissima.
dentata, Thunb. Fl. Jap. 301 = Lactuca Thunbergii.
denticulata, Houtt. Handleid. x. 385 = Lactuca denticulata.
dichotoma, Wall. Cat. n. 3275 = Launaea aspleniifolia, nudicaulis.

PRENANTHES:—
diversifolia, Ledeb. Ic. Pl. Fl. Ross. ii. 17 = Crepis tenuifolia.
erucaefolia, Willd. ex Boiss. Fl. Orient. iii. 801 (= *Mulgedium prenanthoides*).—Reg. Caucas.
erysanifolia, Steud. Nom. ed. II. ii. 393, sphalm. = *erysimifolia*.
erysimifolia, Willd. Sp. Pl. iii. 1543 = Lactuca muralis.
exigua, A. Gray, Pl. Wright. ii. 105 = Lygodesmia exigua.
fastigiata, Blume, Bijdr. 836 = Crepis japonica.
Fraseri, D. Dietr. Syn. Pl. iv. 1309 = P. Serpentaria.
fruticosa, Willd. ex DC. Prod. vii. 69 = Trixis frutescens.
gigantea, Rafin. ex DC. l. c. 242 = crepidinea.
glabra, Heyne, in Wall. Cat. n. 3264 = Crepis acaulis.
glauca, Rafin. Fl. Ludov. 57 (nomen delend.).
glomerata, Decne. in Jacquem. Voy. Bot. 99. t. 107 = Crepis glomerata.
glomerata, Reichb. ex DC. Prod. vi. 393 = Senecio pubigerus.
graciliflora, Wall. Cat. n. 3260 = Lactuca graciliflora.
graminea, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 67 = Lactuca Fischeriana.
grandiflora, D. Dietr. Syn. Pl. iv. 1309 = Lygodesmia grandiflora.
hastata, Thunb. Fl. Jap. 301 = Lactuca denticulata.
hieracifolia, Willd. Sp. Pl. iii. 1541 = Phaecasium lampsanoides.
hispida, Bieb. Fl. Taur. Cauc. ii. 245 = Lactuca microcephala.
hispida, Pall. Reise, i. 501 = Chondrilla brevirostris.
hispidula, DC. Prod. vii. 195 = Brunoniana.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 412.—Reg. Himal.; Burma.
Hothae, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 207.—Burma.
humilis, Thunb. Fl. Jap. 302 (= *Youngia humilis*).—Japon.
illinoënsis, Pers. Syn. ii. 366 = aspera.
indica, Klein, ex Poir. Encyc. Suppl. ii. 330 = Lactuca Heyneana.
integra, Thunb. Fl. Jap. 300 = Crepis integra.
japonica, Linn. Mant. i. 107 = Crepis japonica.
javanica, *Willd. Sp. Pl.* iii. 1534.—Java.
juncea, Pursh, Fl. Am. Sept. ii. 498 = Lygodesmia juncea.
khasiana, *C. B. Clarke, Comp. Ind.* 273.—Reg. Himal.
Kotschyi, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 223.—China.
laciniata, Houtt. Handleid. x. 381 = Lactuca brevirostris.
laevigata, Blume, Bijdr. 836 = Lactuca laevigata.
laevigata, Wall. Cat. n. 3259 = Lactuca macrorhiza.
lanceolata, Houtt. Handleid. x. 383 = Crepis integra.
linifolia, *Roxb. Hort. Beng.* 60, nomen.—Ind. or. ?
longifolia, *Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 43.—Java.
lutea, Vell. Fl. Flum. 350; viii. t. 91 = Hypochoeris brasiliensis.
lyrata, Thunb. Fl. Jap. 303 = Crepis japonica.
mainensis, *A. Gray, Syn. Fl. N. Am.* i. II. 433.—Am. bor.
miamiensis, Riddell, Syn. Fl. W. St. 50 = alba.
minor, Steud. Nom. ed. II. ii. 393 = Stephanomeria minor.
multiflora, Thunb. Fl. Jap. 303 = Crepis japonica.
multiflosculosa, Nees, in Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 265 = Lactuca muralis.
muralis, Linn. Sp. Pl. 797 = Lactuca muralis.
nana, Torr. ex DC. Prod. vii. 241 = P. Serpentaria.
napifolia, Wall. Cat. n. 3277 = Crepis japonica.
Negretii, Req. ex DC. Prod. vii. 156 = Crepis leontodontoides.
nitissima, Hill, Veg. Syst. iii. 172, sphalm. = altissima.
obtusa, Buch.-Ham. in Wall. Cat. n. 3276 = Launaea nudicaulis.
ovata, Riddell, Syn. Fl. W. St. 50 = alba.
paniculata, Moench, Meth. 534 = Phaecasium lampsanoides.
Paosia, Buch.-Ham. ex DC. Prod. vii. 193, vide *Poosia*.
parviflora, Gilib. Fl. Lituan. i. 233 = Lactuca muralis.
parviflora, *Riddell, Syn. Fl. W. St.* 50.—Am. bor.
patens, Wall. Cat. n. 3258 = Lactuca Heyneana, Launaea nudicaulis.

PRENANTHES:—
pauciflora, Torr. in Ann. Lyc. N. York, ii. (1828) 210 = Stephanomeria runcinata.
pendula, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 421.—Ins. Canar.
pinnata, Linn. f. Suppl. 347 = Sonchus leptocephalus.
polymorpha, Ledeb. Fl. Alt. iv. 144 = Crepis nana.
Poosia, Buch.-Ham. ex Wall. Cat. n. 3265 = Crepis japonica.
procumbens, Roxb. Hort. Beng. 60; Fl. Ind. iii. 404 = Launaea nudicaulis.
proteophylla, Riddell, Syn. Fl. W. St. 50 = alba.
pulchra, DC. Fl. Fr. iv. 7 = Phaecasium lampsanoides.
pumila, Baldw. ex Ell. Sketch, ii. 262, in textu = Lygodesmia aphylla.
purpurea, *Linn. Sp. Pl.* 797.—Europ.
pygmaea, Ledeb. in Mém. Acad. Petersb. v. (1812) 553 = Crepis nana.
quinqueloba, Wall. Cat. n. 3261 = Senecio quinquelobus.
racemosa, *Michx. Fl. Bor. Am.* ii. 84.—Am. bor.
racemosa, Roxb. Hort. Beng. 60; Fl. Ind. iii. 405 = Lactuca Heyneana.
ramosissima, All. Fl. Pedem. i. 226. t. 33. = Lactuca viminea.
raphanifolia, DC. Prod. vii. 195 = Brunoniana.
repens, Linn. Sp. Pl. ed. I., errata = Lactuca repens.
rhombifolia, *Humb. ex Willd. Sp. Pl.* iii. 1535.—Venezuela.
rigida, Buch.-Ham. in Wall. Cat. n. 3263 = Crepis acaulis.
rostrata, Blume, Bijdr. 836 = Lactuca graciliflora.
ruanensis, Chick. in Coult. Bot. Gaz. vi. (1881) 191 = P. Serpentaria.
rubicunda, Willd. Sp. Pl. iii. 1537, excl. syn. = alba.
rubida, *Rafin. Atl. Journ.* 17 (Quid?).—Am. bor.
runcinata, James, in Long's Exped. ex Torr. & Gray, Fl. N. Am. ii. 472 = Stephanomeria runcinata.
runcinata, Lag. Gen. et Sp. Nov. 24 = Launaea nudicaulis.
sarmentosa, Wall. Cat. n. 3262 E = Emilia sonchifolia.
sarmentosa, Willd. Phyt. 10 = Launaea pinnatifida.
scandens, *Hook. f. ex Benth. & Hook. f. Gen.* ii. 527; *C. B. Clarke, Comp. Ind.* 274.—Reg. Himal.
Serpentaria, *Pursh, Fl. Am. Sept.* ii. 499. t. 24.—Am. bor.
sikkimensis, *Hook. f. Fl. Brit. Ind.* iii. 412.—Reg. Himal.
simplex, Pursh, Fl. Am. Sept. ii. 498 = virgata.
sinensis, J. F. Gmel. Syst. 1173 = *chinensis*.
sonchifolia, Bunge, Enum. Pl. Chin. Bor. 40 = Lactuca denticulata.
sonchifolia, Willd. Sp. Pl. iii. 1541 = Lactuca Heyneana.
sonchoides, DC. Cat. Fl. Monsp. 53 = Lactuca acuminata.
spathulata, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 192 = Crepis stenoma.
spicata, *Rafin. Atl. Journ.* 150 (Quid?).—Am. bor.
spinosa, Forsk. Fl. Aegypt. Arab. 144 = Launaea spinosa.
squarrosa, Thunb. Fl. Jap. 303 = Lactuca brevirostris.
striata, Blume, Bijdr. 885 = Crepis japonica.
suavis, Salisb. Parad. Lond. t. 85 = alba.
subdentata, Hook. Bot. Misc. ii. (1831) 221 = Picrosia longifolia.
taraxacoides, Wall. Cat. n. 3279 = Lactuca Heyneana.
Tatarinowii, *Maxim. Prim. Fl. Amur.* 474.—Manchuria.
tenuifolia, Linn. Sp. Pl. 797 = purpurea.
tenuifolia, Torr. in Ann. Lyc. N. York, ii. (1828) 210 = Stephanomeria minor.
transalpina, Vell. Fl. Flum. 350; viii. t. 92 = Hieracium Commersonii.
triquetra, Labill. Ic. Pl. Syr. Dec. iii. 4 = Lactuca triquetra.
tuberosa, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 99 = Lactuca microcephala.
versicolor, Fisch. ex Bunge, Enum. Pl. Chin. Bor. 40 = Lactuca Fischeriana.
viminea, Linn. Sp. Pl. 797 = Lactuca viminea.
violaefolia, *Decne. in Jacquem. Voy. Bot.* 100. t. 108.—Reg. Himal.
virgata, *Michx. Fl. Bor. Am.* ii. 83.—Am. bor.
viscosa, Baumg. Enum. Stirp. Transs. iii. 44 = Phaecasium lampsanoides.
vulgaris, Gueldenst. Reisen, i. 113 = Lactuca muralis.
Wolfiana, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 43 = purpurea.

PRENANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 43, 56 = praec.

PRENANTUS, Rafin. Atl. Journ. 150 (1832) = praec.

PREONANTHUS, Ehrh. Beitr. iv. 149 (1789) = **Anemone**, Tourn. (Ranunc.).
alpinus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 11 = A. alpina.
sulfureus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323 = A. alpina.

PREPTANTHE, Reichb. f. in Fl. des Serres, viii. (1852–53) 245 = **Calanthe**, R. Br. (Orchid.).
Sedeni ×, Reichb. f. in Gard. Chron. (1878) i. 168 = *C. Sedeni*.
vestita, Reichb. f. in Fl. des Serres, viii. (1852–53) 245 = C. vestita.
villosa, Reichb. f. in Bonplandia, v. (1857) 42 = C. pilosa.

PREPUSA, Mart. Nov. Gen. et Sp. ii. 120. t. 190 (1827). *GENTIANEAE*, Benth. & Hook. f. ii. 813.
campanulata, Griseb. ex Prog. in Mart. Fl. Bras. vi. i. 244 = connata.
connata, *Gardn. in Hook. Ic. Pl.* iii. t. 225–6.—Bras.
Hookeriana, *Gardn. in Bot. Mag.* t. 3909.—Bras.
montana, *Mart. Nov. Gen. et Sp.* ii. 121. t. 190.—Bras.

PRESCOTTIA, Lindl. in Hook. Exot. Fl. t. 115 (1825) (Prescotia); Bot. Reg. t. 1915, textu, n. 1916 (1836). *ORCHIDEAE*, Benth. & Hook. f. iii. 593.
DECAISNEA, Brongn. in Duperr. Voy. Coq. Bot. 192. t. 39 (1829).
GALEOGLOSSUM, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 31.
colorans, Lindl. Bot. Reg. t. 1915, textu, n. 1916 = stachyodes.
corcovadensis, *Reichb. f. in Linnaea*, xxii. (1849) 814.—Bras.
densiflora, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. vi. (1841) 53.—Bras.
epiphyta, *Rodrig. Orch. Nov.* i. 179.—Bras.
Galeottii, *Reichb. f. in Linnaea*, xix. (1847) 377.—Mexic.
gigantea, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 618, nomen.—Bras.
lanceaefolia, *Link & Otto, ex Steud. Nom.* ed. II. ii. 393, nomen.—Bras.
lancifolia, *Lindl. Gen. et Sp. Orch.* 453.—Bras.
leptostachya, *Lindl. in Bot. Reg.* sub t. 1915 (err. typ. t. 1916).—Bras.
Lindeniana, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 31.—Mexic.
longipetiolata, *Rodrig. Orch. Nov.* i. 177.—Bras.
micrantha, *Lindl. Bot. Reg.* sub t. 1915 (err. typ. t. 1916).—Bras.
microrhiza, *Rodrig. Orch. Nov.* i. 179.—Bras.
montana, *Rodrig. l. c.* 178.—Bras.
Myosurus, *Reichb. f. ex Griseb. Fl. Brit. W. Ind.* 640.—Ind. occ.
nivalis, *Rodrig. Orch. Nov.* ii. 278.—Bras.
octopollinica, *Rodrig. l. c.* 277.—Bras.
oligantha, *Lindl. Gen. et Sp. Orch.* 454.—Jamaic.
ophioglossoides, Spreng. Syst. iii. 706 = Microstylis ophioglossoides.
orchioides, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 386.—Mexic.
pachyrhiza, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 31.—Mexic.
pellucida, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 335.—Cuba.
petiolaris, *Lindl. Bot. Reg.* sub t. 1915 (err. typ. t. 1916).—Peruv.
phleoides, *Lindl. Gen. et Sp. Orch.* 453.—Bras.
plantaginea, Lindl. in Hook. Exot. Fl. t. 115 = seq.
plantaginifolia, *Lindl. l. c.* t. 115, in textu.—Bras.
pubescens, *Rodrig. Orch. Nov.* i. 178.—Bras.
rodeiensis, *Rodrig. l. c.* ii. 278.—Bras.
spiranthophylla, *Rodrig. l. c.* i. 177.—Bras.
stachyodes, *Lindl. Bot. Reg.* sub t. 1915 (err. typ. t. 1916).—Bras.
tennis, *Lindl. Gen. et Sp. Orch.* 454.—Peruv.
viacola, *Rodrig. Orch. Nov.* ii. 279.—Bras.

PRESLAEA, Mart. Nov. Gen. et Sp. ii. 75. t. 164 (1827) = **Heliotropium**, Tourn. (Borag.).
linifolia, A. St. Hil. Voy. Distr. Diam. ii. 433.—Bras.
paradoxa, Mart. Nov. Gen. et Sp. ii. 76.—Bras.
parciflora, Mart. in Flora, xxii. (1839) i. Beibl. 7.—Bras.
stenostachya, A. St. Hil. Voy. Distr. Diam. ii. 434.—Bras.

PRESLEA, G. Don, Gen. Syst. iv. 667 (1837) = **Preslia**, Opiz (Labiat.).

PRESLEA, Spreng. Syst. iv. Cur. Post. 338 (1827) = PRESLAEA, Mart. = **Heliotropium**, Tourn. (Borag.).

PRESLIA, Opiz, Natural. viii. (1824) 86; et in Flora, vii. (1824) 322. *LABIATAE*, Benth. & Hook. f. ii. 1183.
cervina, *Fresen. in Syll. Ratisb.* ii. (1828) 238.—Europ. austr.
glabriflora, Opiz, Natural. viii. (1824) 88; et in Flora, vii. (1824) 323 = cervina.
villiflora, Opiz, ll. cc. = cervina.

PRESTELIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 236 (1873) = **Eremanthus**, Linn. (Compos.).
eriopus, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 169 = E. eriopus.

PRESTINARIA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) i. Intell. 27; Steud. Nom. ed. II. ii. 393 (1841) = **Coreopsis**, Linn. (Compos.).
abyssinica, Sch. Bip. ex Hochst. l. c.; et ex Steud. l. c. = C. abyssinica.
bidentoides, Sch. Bip. ex Hochst. l. c.; et ex Steud. l. c. = C. Prestinaria.
macrantha, Sch. Bip. ex Walp. Rep. vi. 163 = C. macrantha.

PRESTOËA, Hook. f. in Benth. & Hook. f. Gen. iii. 899 (1883). *PALMAE*, Benth. & Hook. f. l. c.
pubigera, *Hook. f. l. c.*—Ins. Trinit.

PRESTONIA, R. Br. in Mem. Wern. Soc. i. (1809) 69. *APOCYNACEAE*, Benth. & Hook. f. ii. 709.
EXOTHOSTEMON, G. Don, Gen. Syst. iv. 82 (1837).
GUACHAMACA, De Gross, in Med. Bot. Criollo, i. 295; ex Ernst, in Trans. Bot. Soc. Edinb. x. (1870) 450.
HAEMADICTYON, Lindl. in Trans. Hort. Soc. vi. (1825) 70.
PENTALINON, Voigt, Hort. Suburb. Calc. 523 (1845).
annularis, *G. Don, Gen. Syst.* iv. 84.—Guian.
bahiensis, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 164.—Bras.
calycina, *Muell. Arg. l. c.* 162.—Bras.
cearensis, *Miers, Apocyn. S. Am.* 148.—Bras.
contorta, *Hemsl. Biol. Centr. Am. Bot.* ii. 311.—Mexic.
glabrata, *H. B. & K. Nov. Gen. et Sp.* iii. 222.—Ecuador.
hirsuta, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 163.—Bras.
hirsuta, Spreng. Syst. i. 637 = Echites Pavonii.
ipomaeifolia, *A. DC. in DC. Prod.* viii. 429.—Panama.
laeta, *Miers, Apocyn. S. Am.* 149.—Bras.
lanata, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 164.—Bras.
latifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 250—Guian.
lutescens, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 164.—Bras.
macrocarpa, *Hemsl. Biol. Centr. Am. Bot.* ii. 311.—Guatemala.
megalagrion, *Miers, Apocyn. S. Am.* 149.—Bras.
mexicana, *A. DC. in DC. Prod.* viii. 429.—Mexic.
mollis, *H. B. & K. Nov. Gen. et Sp.* iii. 221. t. 242.—Am. austr.
parviflora, *Benth. & Hook. f. Gen.* ii. 709.—N. Granat.
peruviana, *Spreng. Syst.* i. 637.—Peruv.
quinquangularis, *Spreng. l. c.*—Am. austr.
schizadenia, *Hemsl. Biol. Centr. Am. Bot.* ii. 312.—Mexic.
Seemanni, Miers, Apocyn. S. Am. 146 = ipomaeifolia.
sericea, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 360.—Mexic.

PRESTONIA:—
surinamensis, *Muell. Arg. in Linnaea*, xxx. (1859–60) 433.—Guian.
tomentosa, *R. Br. in Mem. Wern. Soc.* i. (1809) 70.—Bras.
tomentosa, Seem. Bot. Voy. Herald, 168 = ipomaeifolia.

PRESTONIA, Scop. Introd. 281 (1777) = **Abutilon**, Tourn. (Malvac.).

PRESTONIOPSIS, Muell. Arg. in Linnaea, xxx. (1859–60) 439; et in Bot. Zeit. xviii. (1860) 22 = **Dipladenia**, DC. (Apocyn.).
Fendleri, Miers, Apocyn. S. Am. 168 = D. Fendleri.
hirsuta, Miers, l. c. 167.—Venezuela.
pubescens, Muell. Arg. in Linnaea, xxx. (1859–60) 439; et in Bot. Zeit. xviii. (1860) 22.—Am. austr.
venosa, Miers, Apocyn. S. Am. 167.—Venezuela.

PRETERA, Steud. Nom. ed. II. ii. 394 (1831) sphalm. = seq.

PRETREA, J. Gay, in Ann. Sc. Nat. Sér. I. (1824) 457. *PEDALINEAE*, Benth. & Hook. f. ii. 1059.
DICEROCARYUM, Boj. in Ann. Sc. Nat. Sér. II. iv. (1835) 268. t. 10.
artemisiaefolia, Klotzsch, in Peters, Reise Mossamb. Bot. 189 = zanguebarica.
Bojeriana, Decne. in Duch. Rev. Bot. i. (1846) 517 = zanguebarica.
eriocarpa, *Decne. l. c.*—Afr. austr.
Forbesii, *Decne. in Ann. Sc. Nat.* Sér. V. iii. (1865) 334.—Afr. austr.
loasaefolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 188.—Afr. trop.
senecioides, Klotzsch, l. c. 189 = zanguebarica.
zanguebarica, *J. Gay, in Ann. Sc. Nat.* Sér. I. i. (1824) 457.—Afr. trop.

PREVOITA, Steud. Nom. ed. II. ii. 394 (1841), sphalm. = PREVOTIA, Adans. = **Cerastium**, Dill. (Caryoph.).

PREVOSTEA, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 497 = **Breweria**, R. Br. (Convolv.).
africana, Benth. in Hook. Niger Fl. 469 = B. africana.
amazonica, Choisy, in DC. Prod. ix. 437 (= *Reinwardtia glabra*).—Bras.
ferruginea, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 498 = B. ferruginea.
glabra, Choisy, l. c. (= *Reinwardtia glabra*).—N. Granat.
heterantha, G. Don, Gen. Syst. iv. 299 (= *Aniseia heterantha*).—Bras.
mossambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. 244.—Afr. trop.
sericea, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 498 (= *Reinwardtia sericea*).—N. Granat.
spectabilis, Meissn. in Mart. Fl. Bras. vii. 325.—Bras.
umbellata, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 497 (= *Reinwardtia umbellata*).—Bras.
velutina, Walp. Rep. vi. 542.—Mexic.

PREVOTIA, Adans. Fam. ii. 256 (1763) = **Cerastium**, Dill. (Caryoph.).

PRIANTHES, Pritz. Ic. Ind. ii. 232 (1865) = PRIONANTHES, Schrank = **Trixis**, P. Br. (Compos.).

PRIESTLEYA, DC. Mém. Lég. 190 (1825); et in Ann. Sc. Nat. Sér. I. iv. (1825) 90. *LEGUMINOSAE*, Benth. & Hook. f. i. 472.
ACHYRONIA, Wendl. Bot. Beobacht. 39. t. 12 (1798).
XIPHOTHECA, Eckl. & Zeyh. Enum. 166 (1836).
angustifolia, *Eckl. & Zeyh. Enum.* 165.—Afr. austr.
axillaris, DC. Mém. Lég. 197 = Amphithalea densa.
axillaris, E. Mey. Comm. Pl. Afr. Austr. 20 = Amphithalea cuneifolia.
capitata, *DC. Prod.* ii. 121.—Afr. austr.
Cephalotes, E. Mey. Comm. Pl. Afr. Austr. 18 = umbellifera.
elliptica, *DC. Mém. Lég.* 198.—Afr. austr.
elliptica, E. Mey. in Linnaea, vii. (1832) 150 = Amphithalea densa.
ericaefolia, DC. Mém. Lég. 195. t. 30 = Amphithalea ericaefolia.
graminifolia, *DC. Prod.* ii. 122.—Afr. austr.

PRIESTLEYA:—
hirsuta, *DC. Mém. Lég.* 193.—Afr. austr.
laevigata, DC. l. c. 195 = teres.
laevigata, Eckl. & Zeyh. Enum. 165 = angustifolia.
laevigata, E. Mey. Comm. Pl. Afr. Austr. 18 = capitata.
lanceolata, E. Mey. in Linnaea, vii. (1832) 150 = sericea?
lancifolia, C. B. Presl, ex J. S. Presl, Rostl. iii. 76 = hirsuta?
latifolia, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 447.—Afr. austr.
leiocarpa, *Eckl. & Zeyh. Enum.* 165.—Afr. austr.
Meyeri, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 65 = Amphithalea densa.
myrtifolia, *DC. Mém. Lég.* 194.—Afr. austr.
polycarpa, Steud. Nom. ed. II. ii. 394 = tecta.
rotundifolia, Walp. in Linnaea, xiii. (1839) 469 = tecta.
sericea, DC. Prod. ii. 122 = Amphithalea densa.
sericea, *E. Mey. Comm. Pl. Afr. Austr.* 19.—Afr. austr.
Sieberi, *Tausch, in Flora*, xxv. (1842) i. 281.—Afr. austr.
tecta, *DC. Prod.* ii. 122.—Afr. austr.
teres, *DC. l. c.*—Afr. austr.
Thunbergii, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 446.—Afr. austr.
tomentosa, DC. Prod. ii. 122 = villosa?
umbellifera, *DC. l. c.*—Afr. austr.
umbellifera, Eckl. & Zeyh. Enum. 165 = Thunbergii.
umbellifera, E. Mey. Comm. Pl. Afr. Austr. 17 = angustifolia.
vestita, *DC. Prod.* ii. 122.—Afr. austr.
villosa, *DC. l. c.*—Afr. austr.

PRIESTLEYA, [Moç. & Sesse,] ex DC. Prod. v. 564 (1836) = **Montañoa**, Cerv. (Compos.).
corymbosa, [Moç. & Sesse,] ex DC. l. c. 565 = M. tomentosa.
longifolia, [Moç. & Sesse,] ex DC. l. c. = M. grandiflora.
squarrosa, [Moç. & Sesse,] ex DC. l. c. = M. frutescens.

PRIEUREA, DC. Prod. iii. 58 (1828) = **Ludwigia**, Linn. (Onagr.).
senegalensis, DC. l. c.—Afr. trop.

PRIEURIA, Benth. & Hook. f. Gen. i. 789 (1867) = praec.

PRIMULA, Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 631.
ALEURITIA, Spach, Hist. Vég. Phan. ix. 360 (1840).
ARETIA, Link, Handb. ii. 411 (1829).
AUGANTHUS, Link, l. c. 414 (1829).
CANKRIENIA, De Vriese, in Miq. Pl. Jungh. 86 (1851).
EVOTROCHIS, Rafin. Fl. Tellur. ii. 76 (1836).
KABLIKIA, Opiz, Seznam, 55 (1852).
OSCARIA, Lilja, in Lindbl. Bot. Notiser (1839), 39.
PRIMULIDIUM, Spach, Hist. Vég. Phan. ix. 354 (1840).
acaulis, Hill, Veg. Syst. viii. 25; Jacq. Misc. i. 158 = vulgaris.
admontensis ×, Gusmus, ex Church. in Gard. Chron. (1884) II. 808.—Europ.
alba, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 185.—Europ. med.
algida, *Adam, ex Weber & Mohr, Beitr.* i. 46.—As. Min.; As. bor.
algida, Janka, ex Boiss. Fl. Orient. iv. 30 = frondosa.
Allionii, Hausm. Fl. Tirol. ii. 719 = tyrolensis.
Allionii, *Loisel. in Desv. Journ. Bot.* ii. (1809) 262. t. 11. f. 1.—Europ. austr.
alpestris, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 89 = elatior.
alpina, Loisel. Fl. Gall. ed. II. i. 160 = viscosa.
alpina, Salisb. Prod. 118 = P. Auricula.
alpina, Schleich. ex Reichb. Pl. Crit. ix. 11 = *pubescens*.
altaica, Lehm. Monog. Prim. 59. t. 5 = farinosa.
altaica, Pall. ex Ledeb. Fl. Alt. i. 209 = elatior.
altissima, D. Don, Prod. Fl. Nep. 78 = reticulata.
ambigua, Salisb. Prod. 117 = elatior.
amethystina, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 268.—China.

PRIMULA:—
amoena, *Bieb. Fl. Taur. Cauc.* i. 138.—Reg. Caucas.; Persia.
androsacea, Georgi, ex Ledeb. Fl. Ross. iii. 18 = Androsace Chamaejasme.
angustifolia, *Torr. in Ann. Lyc. N. York*, i. (1824) 34. t. 3. f. 3.—Am. bor. occ.
Arctotis ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 124.—Europ.
aretioides, Lehm. Monog. Prim. 90. t. 9 = Dionysia aretioides.
atroviolacea, Jacquem. ex Duby, in DC. Prod. viii. 41 = nivalis.
Aucheri, *Jaub. & Spach, Illustr.* i. 97. t. 49.—Arab.
Auricula, Hill, Veg. Syst. viii. 25 = farinosa.
Auricula, *Linn. Sp. Pl.* 143.—Europ.
Auricula, Vill. Hist. Pl. Dauph. ii. 469 = marginata.
auriculata, *Lam. Illustr.* ii. 429.—As. Min.; Persia.
auriculata, Ledeb. Fl. Alt. i. 211 = algida.
Balbisii, Lehm. Monog. Prim. 45 = ciliata.
Bayerni, Rupr. in Bull. Acad. Pétersb. vi. (1863) 238, in textu = nivalis.
bella, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 268.—China.
bellidifolia, *King, ex Hook. f. Fl. Brit. Ind.* iii. 486.—Reg. Himal.
Berninae ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 153.—Europ.
bicolor, Rafin. Caratteri, 78 = vulgaris.
biflora ×, Huter, ex Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 158.—Europ.
Bonjeani, Huegen, ex Nym. Consp. 605 = pedemontana.
borealis, Duby, in DC. Prod. viii. 43 = farinosa.
Boveana, Decne. ex Duby l. c. 35 = verticillata.
bracteata, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 266.—China.
Brandisii, Wiesb. in Oestr. Bot. Zeitschr. xxxii. (1882) 282 = officinalis.
breviscapa, [Clairv.] Man. Herb. 58 = vulgaris.
brevistyla ×, DC. Fl. Fr. Suppl. 383.—Europ.
bullata, *Franch. in Bull. Soc. Bot. Fr*, xxxii. (1885) 365.—China.
Bungeana, C. A. Mey. in Beitr. Pfl. Russ. Reich. vi. 22 = algida.
cachemeriana, W. M[unro], in Garden, xvi. (1879) 535 = denticulata.
caespitosa, Steud. Nom. ed. I. 655, vide *cespitosa*.
calliantha, *Franch. in Bull. Soc. Fr.* xxxii. (1885) 368.—China.
calycantha, Retz. Obs. ii. 10 = vulgaris.
calycina, Duby, ex Gaud. Fl. Helv. ii. 94 = glaucescens.
calycina, Reichb. Pl. Crit. i. 32. f. 70 = spectablis.
Candolleana, Reichb. l. c. vi. 18 = integrifolia.
capitata, *Hook. Bot. Mag.* t. 4550.—Reg. Himal.
capitellata, *Boiss. Diagn.* Ser. I. vii. 64.—As. Min.; Persia; Afghan.
carniolica, *Jacq. Fl. Austr.* v. 28.—Austria.
carniolica, Poll. Fl. Veron. i. 231 = spectabilis.
carpathica, Fuss, in Verh. Siebenb. Ver. Naturw. x. (1859) 149 = elatior.
cashmeriana, Carr. in Rev. Hortic. (1880) 330 = denticulata.
caucasica, C. Koch, in Linnaea, xxiii. (1850) 614 = algida.
cernua, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 271.—China.
cespitosa, Willd. ex Roem. & Schult. Syst. iv. 785 = Dionysia aretioides.
chinensis, Auct. = sinensis.
ciliata, *Moretti, in Brugnat. Giorn. Fis.* viii. (1815) 127.—Alp. Europ. austr.
ciliata, Schrank, Prim. Fl. Salisb. 64 = hirsuta.
Clarkei, *Watt, in Journ. Linn. Soc.* xx. (1882) 4.—Reg. Himal.
Clusiana, *Tausch, in Flora*, iv. (1821) 364.—Europ.
Clusii, Wiest, ex Nym. Consp. 605 = Clusiana.
Columnae, Tenore, Prod. Fl. Nap. 14 = officinalis.
commutata, Schott, in Oestr. Bot. Wochenbl. ii. (1852) 35 = viscosa.
concinna, Watt, in Journ. Linn. Soc. xx. (1882) 5 = farinosa.
confinis, Schott, ex Reichb. Ic. Fl. Germ. xvii. 40 = hirsuta.
cordifolia, *Kit. in Linnaea*, xxxii. (1863) 451.—Hungar.
cordifolia, Rupr. in Bull. Acad. Pétersb. vi. (1863) 225 = elatior.

PRIMULA :—
cordifolia, Schur, Enum. Pl. Transs. 554 = officinalis.
coronaria, Salisb. Prod. 117 = officinalis.
cortusoides, *Linn. Sp. Pl.* 144.—Sibir. ; Japon.
crassifolia, Lehm. Monog. Prim. 91. t. 9 = nivalis.
crenata, Lam. Illustr. i. 429 = marginata.
crenata, Schur, Enum. Pl. Transs. 553 = elatior.
cuneifolia, Duby, in DC. Prod. viii. 39 = seq. ?
cuneifolia, *Ledeb. in Mém. Acad. Pétersb.* v. (1814) 522.—Reg. arct.
Cushia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 78 = petiolaris.
daonensis, *Leyb. in Oestr. Bot. Wochenbl.* iv. (1854) 10.—Europ. austr.
darialica, *Rupr. in Bull. Acad. Pétersb.* vi. (1863) 236. —Reg. Caucas.
davurica, Lehm. Monog. Prim. 58 = farinosa.
davurica, *Spreng. Gartenz.* (1804) 113, cum fig.—Dahuria.
decipiens, Duby, in DC. Prod. viii. 44 = farinosa.
decora, Sims, Bot. Mag. t. 1922 = hirsuta.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 272.—China.
dentata, Donn, ex Roem. & Schult. Syst. iv. 133 = cortusoides ?
denticulata, *Sm. Exot. Bot.* ii. 109. t. 114.—Reg. Himal.
denticulata, Wight, Ic. t. 2000 = elliptica.
dentiflora, Andr. Bot. Rep. vi. t. 405 = cortusoides.
deusta, Banks, ex Roem. & Schult. Syst. iv. 152 (Quid ?). —Hab. ?
Dickieana, *Watt, in Journ. Linn. Soc.* xx. (1882) 9.—Reg. Himal.
digenea ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 79.—Europ.
Dinyana ×, Lagger, in Flora, xxii. (1839) 670.—Helvet.
discolor ×, Leyb. in Flora, xxxviii. (1855) 344.—Europ.
discolor, Schur, Enum. Pl. Transs. 554 = officinalis.
dolomitis, *Hort. Llewelyn, ex Baker, in Gard. Chron.* (1884) i. 577.—Europ. austr.
domestica, Hoffmgg. Verz. Pfl. Nachtr. ii. 185 = elatior.
dryadifolia, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 270.—China.
egaliksensis, *Wormsk. ex Hornem. in Fl. Dan.* t. 1511. —Am. arct.
egalliccensis, Lehm. Monog. Prim. 64. t. 7 = praec.
elata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 79 = denticulata.
elatior, *Hill, Veg. Syst.* viii. 25 ; *Mill. Gard.* ed. VIII. n. 2 ; *Jacq. Misc.* i. 158.—Europ.
elegans, Duby, in DC. Prod. viii. 42 = rosea.
elliptica, *Royle, Illustr. Bot. Himal.* 311. t. 76.—Reg. Himal.
elongata, *Watt, in Journ. Linn. Soc.* xx. (1882) 8.—Reg. Himal.
Elwesiana, *King, ex Watt, l. c.* 13.—Reg. Himal.
erosa, *Wall. Cat.* n. 611.—Reg. Himal.
Escheri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
exaltata, Lehm. Monog. Prim. 57 = farinosa.
exscapa, Hegetschw. Fl. Schw. 195 = viscosa.
Facchinii ×, Schott, Wilde Blendl. Prim. 18. t. 3.—Europ.
farinifolia, *Rupr. in Bull. Acad. Pétersb.* vi. (1863) 236.—Reg. Caucas.
farinosa, Bieb. Fl. Taur. Cauc. i. 139 = algida.
farinosa, Griseb. Spicil. Fl. Rumel. ii. 2 = frondosa.
farinosa, *Linn. Sp. Pl.* 143.—Reg. bor.
farinosa, Schangin, in Pall. N. Nord. Beitr. vi. 47 = auriculata.
Fedschenkoi, *Regel, in Act. Hort. Petrop.* iii. (1875) 133.—Turkestan.
filipes, *Watt, in Journ. Linn. Soc.* xx. (1882) 5.—Reg. Himal.
fimbriata, *Wall. ex Duby, in DC. Prod.* viii. 45.—Reg. Himal.
finmarchica, Jacq. Misc. i. 160 = sibirica.
finno-marchica, Georgi, Beschr. Russ. Reich. III. iv. 764 = praec.
flagellicaulis ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 79.—Europ.
flava, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 497.—China.
Floerkeana, *Schrad. in Kruenitz, Oecon. Encyc.* cxvii. (1811) 393.—Alp. Europ. austr.

PRIMULA :—
floribunda, *Wall. Tent. Fl. Nap.* 43. t. 33.—Reg. Himal.
Flueggeana, Lehm. Monog. Prim. 36. t. 2 = elatior.
Forsteri ×, Stein, in Gartenfl. (1879) 322.—Europ.
Freyeri, Hoppe, ex Reichb. Fl. Germ. Excurs. 403 = carniolica.
frondosa, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 204.—Thracia.
Gambeliana, *Watt, in Journ. Linn. Soc.* xx. (1882) 3. —Reg. Himal.
geraniifolia, *Hook. f. Fl. Brit. Ind.* iii. 484.—Reg. Himal.
gigantea, *Jacq. Misc.* i. 160.—Sibir.
gigantea, Lehm. Monog. Prim. 61. t. 6 = farinosa.
glabra, *Klatt, in Linnaea*, xxxvii. (1871–73) 500.—Reg. Himal.
glabrescens, F. Nyl. ex W. Nyl. & Saell. Herb. Mus. Fenn. 32, in nota = stricta.
glacialis, Adam, ex Rupr. in Bull. Acad. Pétersb. vi. (1863) 230 = auriculata.
glacialis, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 270.—China.
glacialis, Willd. ex Roem. & Schult. Syst. iv. 785 = algida.
glandulosa, Schur, Enum. Pl. Transs. 552 = elatior.
glaucescens, *Moretti, in Brugnat. Giorn. Fis.* II. v. (1822) 249.—Europ. austr.
glaucescens, Reichb. Fl. Germ. Excurs. 403 = spectabilis.
globifera, Griff. Itin. Notes, 148, nomen = capitata.
glutinosa, All. Auct. Pedem. 6 = Allionii.
glutinosa, Lapeyr. Hist. Abr. Pl. Pyr. 96 = viscosa.
glutinosa, *Wulf. in Jacq. Fl. Austr.* v. 41.—Europ. austr.
Goebelii ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 82.—Europ.
grandiflora, Bast. Suppl. Fl. Maine-et-Loire, 26 = carniolica.
grandiflora, Lam. Fl. Fr. ii. 248 = vulgaris.
grandis, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 395. —Reg. Caucas.
graveolens, Hegetschw. Fl. Schw. 194. t. 6 = viscosa.
Halleri, [Honck.] Verz. Aller Gew. Teutschl. 576 = longiflora.
Hampeana, *Wallr. Beitr. Fl. Hercyn.* 333 (Quid ?).—Europ.
Heerii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
helvetica, Donn, Hort. Cantab. ed. II. 20, nomen ; Lodd. Bot. Cab. t. 348 = *pubescens*.
Henryi ×, Hort.—Cf. Gard. Chron. (1881) i. 404.
heterochroma, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 70.—Persia.
Heydei, *Watt, in Journ. Linn. Soc.* xx. (1882) 5.—Reg. Himal.
hirsuta, *All. Fl. Pedem.* i. 93.—Europ.
hirsuta, Reichb. ex Nym. Consp. 604 = daonensis.
hirsuta, Vill. Hist. Pl. Dauph. ii. 469 = viscosa.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 97 = denticulata.
Hookeri, *Watt, in Journ. Linn. Soc.* xx. (1882) 14.—Reg. Himal.
Hornemanniana, Lehm. Monog. Prim. 55. t. 4 = farinosa.
Huguenini ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
humilis, Steud. Nom. ed. II. ii. 395 = pusilla.
Huteri ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 159.—Europ.
hybrida ×, Knaf, in Lotos, xiv. (1864) 84.—Europ.
hybrida, Schrank, Baier. Fl. i. 449 = vulgaris.
hyperborea, Spreng. Syst. i. 577 = cuneifolia.
imperialis, *Jungh. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 298.—Java.
incisa, Lam. Fl. Fr. ii. 250 = integrifolia.
inflata, Lehm. Monog. Prim. 26. t. 2. f. 1 = officinalis.
inodora, Hill, Herb. Brit. ii. 193 ; Gilib. Fl. Lituan. i. 32 = vulgaris.
integrifolia, Jacq. Fl. Austr. iv. 14. t. 327 = spectabilis.
integrifolia, *Linn. Sp. Pl.* 144.—Europ. austr.
integrifolia, Oed. in Fl. Dan. t. 188 = sibirica.
integrifolia, Scop. Fl. Carn. ed. II. i. 133 = carniolica.
integrifolia, Vis. Fl. Dalm. ii. 149 = Kitaibeliana.
intermedia ×, Facc. in Zeitschr. Ferdinand. Tirol. III. iv. (1854) 19 = *brevistyla*.
intermedia, Ledeb. in Mém. Acad. Pétersb. v. (1814) 519 = sibirica.

PRIMULA :—
intermedia ×, Portenschl. in Tratt. Arch. t. 387.—Europ.
intermedia, Sims, Bot. Mag. t. 1219 = farinosa.
intricata, Gren. & Godr. Fl. Fr. ii. 449 = elatior.
intrusa, Reichb. Pl. Crit. vii. 15. t. 631. f. 852 = sibirica.
involucrata, Sw. Cat. 562 ; ex Duby, in DC. Prod. viii. 35 ; Otto, ex Hook. Bot. Mag. t. 2842 = verticillata.
involucrata, *Wall. Cat.* n. 7107 ; *Naud. in Rev. Hortic.* Sér. II. v. (1846) 283.—Enrop. ; As. bor.
Jaeschkeana, Kern. Nov. Pl. Sp. Dec. ii. 1 = nivalis.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 400.—Japon.
Jellenkiana, Freyer, ex Nym. Consp. 604 = carniolica.
jesoana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 119.—Japon.
Jiraseckiana, Tratt. ex Reichb. Fl. Germ. Excurs. 402 = minima.
Kaufmanniana, *Regel, in Act. Hort. Petrop.* iii. (1875) 131.—As. med.
Kerneri ×, Goebl & Stein, in Oestr. Bot. Zeitschr. xxviii. (1878) 188.—Europ.
Kingii, *Watt, in Journ. Linn. Soc.* xx. (1882) 9.—Reg. Himal.
kisoana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 119.—Japon.
Kitaibeliana, *Schott, in Oestr. Bot. Wochenbl.* ii. (1852) 268.—Europ.
Kraettliana ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 99.—Europ.
lactea, Lam. Fl. Fr. ii. 250 = Androsace lactea.
laevigata, Duby, ex Reichb. Pl. Crit. i. 33 = glaucescens.
lasiopetala, *Wallr. Beitr. Fl. Hercyn.* 333 (Quid ?).—Europ.
lateriflora, Goupil, in Mém. Soc. Linn. Par. iii. (1825) 250. t. 10 = elatior.
latifolia, Koch, Syn. Fl. Germ. ed. I. 588 = hirsuta.
latifolia, Lapeyr. Hist. Abr. Pl. Pyr. 97 = viscosa.
lepida, Duby, in DC. Prod. viii. 44 = farinosa.
Listeri, *King, ex Hook. f. Fl. Brit. Ind.* iii. 485.—Reg. Himal.
longiflora, *All. Fl. Pedem.* i. 92. t. 39.—Europ.
longiflora, Auch. ex Duby, in DC. Prod. viii. 46 = Dionysia cespitosa.
longifolia, Curt. Bot. Mag. t. 392 = auriculata.
longiscapa, Ledeb. in Mém. Acad. Pétersb. v. (1814) 520 = farinosa.
lutea, Vill. Hist. Pl. Dauph. ii. 469 = P. Auricula.
luteola, *Rupr. in Bull. Acad. Pétersb.* vi. (1863) 233. —Reg. Caucas.
macrocalyx, Bunge, in Ledeb. Fl. Alt. i. 209 = officinalis.
macrocarpa, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 68.—Japon.
macrophylla, D. Don, Prod. Fl. Nep. 80 = nivalis.
macrophylla, C. Koch, in Linnaea, xxiii. (1850) 615 = auriculata.
magellanica, *Lehm. Monog. Prim.* 62. t. 6.—Reg. Magell.
Mandarina, Hoffm. in Otto & Dietr. Allg. Gartenz. iii. (1835) 194 = sinensis.
marginata, *Curt. Bot. Mag.* t. 191.—Europ.
marginata, Georgi, Beschr. Russ. Reich. III. iv. 763 (Quid ?).—Sibir.
maxima, *Hort. ex Steud. Nom.* ed. II. ii. 395, nomen. —Hab. ?
Maximowiczii, *Regel, in Act. Hort. Petrop.* iii. (1875) 139.—China bor.
media ×, Peterm. Fl. Lips. (1838) 77.—Europ.
megaseaefolia, *Boiss. Fl. Orient.* iv. 26.—As. Min.
Meyeri, Rupr. in Bull. Acad. Pétersb. vi. (1863) 224 = amoena.
microcalyx, Lehm. Monog. Prim. 46. t. 4 = marginata.
minima, Habl. Taur. 117 = vulgaris.
minima, *Linn. Sp. Pl.* 143.—Europ. med.
minuta, Banks, ex Roem. & Schult. Syst. iv. 152 (Quid ?).—Hab. ?
minutissima, *Jacquem. ex DC. Prod.* viii. 42.—Reg. Himal.
mistassinica, Michx. Fl. Bor. Am. i. 124 = farinosa.
modesta, Bisset & S. Moore, in Journ. Bot. xvi. (1878) 134 = farinosa.
mollis, *Nutt. ex Hook. Bot. Mag.* t. 4798.—Reg. Himal.
montana, Opiz, Natural. x. (1825) 225 = elatior.
montana, Reut. ex Nym. Consp. 603 = officinalis.
Moorcroftiana, Wall. Cat. n. 4988 = nivalis.

PRIMULA:—
moschata, *Salisb. Prod.* 118.—Hab.?
multiceps, Freyer, ex Duby, in DC. Prod. viii. 38 = carniolica.
Munroi, Lindl. Bot. Reg. (1847) t. 15 = involucrata.
Mureti, Charpentier, ex Koch, Syn. Fl. Germ. ed. II. 678 = seq.
Muretiana ×, Moritz. Pfl. Graub. 111.—Helvet.
muscoides, *Hook. f. ex Watt, in Journ. Linn. Soc.* xx. (1882) 15; *et Fl. Brit. Ind.* iii. 494.—Reg. Himal.
mutabilis, Lour. Fl. Cochinch. 104 = Hydrangea Hortensia.
nana, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 23 = petiolaris.
nivalis, Don, ex Reichb. Fl. Germ. Excurs. 405 = ciliata.
nivalis, Hort. ex Hoffmgg. Verz. Pfl. Nachtr. ii. 185 = alba.
nivalis, *Pall. Reise*, iii. 723.—As. et Am. bor.
nivalis, Turcz. ex Steud. Nom. ed. II. ii. 395 = farinosa.
nivea, Hort. ex Steud. l. c. = ciliata, villosa.
norwegica, Retz. Fl. Scand. ed. II. 55 = sibirica.
nutans, Georgi, Bemerk. einer Reise Russ. Reich. i. 200 = sibirica.
obconica, *Hance, in Journ. Bot.* xviii. (1880) 234.—China.
oblongifolia ×, Schur, Enum. Pl. Transs. 555.—Europ.
obovata ×, Huter, in Oestr. Bot. Zeitschr. xxiii. (1873) 125.—Europ.
obovata, Wall. Cat. n. 610 = floribunda.
obtusifolia, *Royle, Illustr. Bot. Himal.* 311. t. 77. f. 1.—Reg. Himal.
odontophylla, Wall. Cat. n. 7016 = rotundifolia.
odorata, Gilib. Fl. Lituan. i. 32 = officinalis.
oenensis, Thom. ex Schott, in Oestr. Bot. Wochenbl. ii. (1852) 36 = daonensis.
officinalis, Hill, Veg. Syst. viii. 25 = elatior?
officinalis, *Jacq. Misc.* i. 159.—Europ.; As. Min.
Olgae, *Regel, in Act. Hort. Petrop.* iii. (1875) 151.—Turkestan.
oreocharis, Hance, in Journ. Bot. xiii. (1875) 133 = Maximowiczii.
orientalis, Willd. ex Roem. & Schult. Syst. iv. 785 = = nivalis.
ornata, Banks, ex Roem. & Schult. l. c. 152, nomen (Quid?).—Hab.?
pachyscapa, *Goiran & Del Fabbro, Spec. Morph. Veg. ex Nuov. Giorn. Bot. Ital.* ix. (1877) 170.—Hab.?
Palinuri, *Petagn. Inst. Bot.* ii. 332.—Ital.
Pallasii, Lehm. Monog. Prim. 38. t. 3 = elatior.
pallida, Schott, in Oestr. Bot. Wochenbl. ii. (1852) 35 = hirsuta.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 257.—Am. bor. occ.
parvifolia, Duby, in DC. Prod. viii. 42 = farinosa.
patens, Turcz. in Bull. Soc. Nat. Mosc. (1838) 99 = cortusoides.
pedemontana, *Thom. Cat. Pl. Suiss.* (1818) 24, nomen; *et ex Reichb. Fl. Germ. Excurs.* 403.—Europ. austr.
Perreiniana, Flugg. in Ann. Mus. Par. xii. (1808) 420. t. 27 = elatior.
petiolaris, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 22.—Reg. Himal.
pinnatifida, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 271.—China.
pistiifolia, Griseb. in Goett. Abh. vi. (1854) 127 = Samolus spathulatus.
pistillaris, Hoffmgg. Verz. Pfl. Nachtr. ii. 187 = officinalis?
Plantae ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
poculiformis, Hook. f. Bot. Mag. t. 6582 = obconica.
Polliniana, Moretti, Primul. Ital. 12 = spectabilis.
polyantha, Mill. Gard. Dict. ed. VIII. n. 4 = elatior.
Portae ×, Huter, ex Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 81.—Europ.
praenitens, [Ker-Gawl. in] Bot. Reg. t. 539 = sinensis.
prolifera, *Wall. in As. Res.* xiii. (1820) 372.—Reg. Himal.
Pseudo-acaulis, Schur, Enum. Pl. Transs. 552 = vulgaris.
puberula ×, Schott, Wilde Blendl. Prim. 10. t. 2.—Europ.
pubescens ×, Jacq. Misc. i. 159.—Alp. Europ. med.
pubescens, Loisel. Fl. Gall. ed. I. 114 = viscosa.
pulchra, *Watt, in Journ. Linn. Soc.* xx. (1882) 3.—Reg. Himal.

PRIMULA:—
pumila ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 156.—Europ.
Pumilio, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 498.—Tibet. or.
purpurea, Royle, Illustr. Bot. Himal. 311. t. 77. f. 2 = nivalis.
pusilla, Goldie, in Edinb. Phil. Journ. vi. (1822) 322 = farinosa.
pusilla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 22.—Reg. Himal.
pycnorhiza, Ledeb. Fl. Ross. iii. 11 = auriculata.
pyrenaica, Miégev. in Bull. Soc. Bot. Fr. x. (1863) 28 = officinalis.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 428.—Japon.
reptans, *Hook. f. ex Watt, in Journ. Linn. Soc.* xx. (1882) 14; *et Fl. Brit. Ind.* iii. 494.—Reg. Himal.
reticulata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 21.—Reg. Himal.
rhaetica, Gaud. Fl. Helv. ii. 91 = *pubescens*, Jacq.
rosea, *Royle, Illustr. Bot. Himal.* 311. t. 75. f. 1.—Reg. Himal.
rotundifolia, Pall. Reise, iii. 223 = sibirica.
rotundifolia, *Wall. ex Roxb. Fl. Ind. ed. Carey*, ii. 18.—Reg. Himal.
rubra, [Honck.] Verz. Aller Gew. Teutschl. 575 = viscosa.
Rusbyi, *Greene, in Bull. Torrey Bot. Club.* viii. (1881) 122.—N. Mexic.
salisburgensis, Willd. ex Schott, Wilde Blendl. Prim. 8. t. 1 = Floerkeana.
Salisii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
sapphirina, *Hook. f. & Thoms. ex Watt, in Journ. Linn. Soc.* xx. (1882) 10; *et ex Hook. f. Fl. Brit. Ind.* iii. 492.—Reg. Himal.
Saundersiana, Royle, ex Hook. f. Fl. Brit. Ind. iii. 494 = minutissima.
Sauteri, Schultz, in Flora, xix. (1836) 123 = minima.
saxifragifolia, Lehm. Monog. Prim. 89. t. 9 = cuneifolia.
scotica, Hook. Fl. Lond. t. 133 = farinosa.
secundiflora, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 267.—China.
sedifolia, Salisb. Parad. Lond. t. 107 = Douglasia Vitaliana.
semperflorens, Loisel. ex Steud. Nom. ed. II. ii. 396 = sinensis.
septemloba, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 265.—China.
sericea, Krock. Fl. Siles. Suppl. i. 249 (Quid?).—Europ.
serrata, Georgi, Bemerk. einer Reise Russ. Reich. i. 201 = sibirica.
serratifolia, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 267.—China.
sertulosa, Kickx f. in Mém. Soc. Linn. Par. iv. (1826) 31. t. 3 = sinensis.
sessilis, Roxb. ex Steud. Nom. ed. II. ii. 396; Royle, ex Hook. f. Fl. Brit. Ind. iii. 493 = petiolaris.
sibirica, *Jacq. Misc.* i. 160.—As. et Am. arct.
Sibthorpi, *Hoffmgg. Verz. Pfl.* 189.—Iber.; Graec.
Sibthorpii, Hort. ex Reichb. Fl. Germ. Excurs. 402 = vulgaris.
Sieboldi, *E. Morr. Belg. Hortic.* xxiii. (1873) 97. t. 6.—Japon.
sikkimensis, *Hook. Bot. Mag.* t. 4597.—Reg. Himal.
simensis, Hochst. in Gardn. & Field. Sert. Pl. t. 48 = verticillata.
simplicissima, *Ledeb. ex Steud. Nom.* ed. II. ii. 396, nomen.—Hab.?
Simsii, Sweet, Hort. Brit. ed. I. 330 = villosa.
sinensis, Lour. Fl. Cochinch. 105 (sp. dub.).—China.
sinensis, *Sabine, ex Lindl. Coll. Bot.* t. 7.—China.
soldanelloides, *Watt, in Journ. Linn. Soc.* xx. (1882) 10; *et ex Hook. f. Fl. Brit. Ind.* iii. 493.—Reg. Himal.
sonchifolia, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 266.—China.
spathulacea, Jacquem. ex Duby, in DC. Prod. viii. 44 = elliptica.
spathulata, Royle, Illustr. Bot. Himal. 310, nomen = nivalis.
speciosa, D. Don, Prod. Fl. Nep. 78 = reticulata.
speciosa, Gmel. ex Steud. Nom. ed. II. ii. 396 = nivalis.

PRIMULA:—
spectabilis, Koch, Syn. Fl. Germ. ed. I. 589, partim = Clusiana.
spectabilis, *Tratt. Tabul.* i. 426.—Alp. Europ. austr.
spicata, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 269.—China.
Steinii ×, Obrist, ex Stein, in Gartenfl. (1879) 322. t. 991.
Stelviana, Vulp. in Flora, xli. (1858) 549 (err. typ. 249) = daonensis.
stenocalyx, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 498.—China.
Stirtoniana, *Watt, in Journ. Linn. Soc.* xx. (1882) 15; *et ex Hook. f. Fl. Brit. Ind.* iii. 495.—Reg. Himal.
Stracheyi, Hook. f. & Thoms. ex Watt, l. c. 14, partim = reptans.
Stracheyi, Hook. f. ex W. M[unro,] in Garden (1879) 535 = minutissima.
stricta, *Hornem. in Fl. Dan.* t. 1385.—Europ. bor.; Island.
Stuartii, Hook. f. Fl. Brit. Ind. iii. 490, partim = nivalis.
Stuartii, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 20.—Reg. Himal.
suaveolens, Bertol. in Desv. Journ. Bot. ii. (1813) 76 = officinalis.
suaveolens ×, Caruel, in Atti Soc. Ital. Sc. Nat. viii. (1865) 467.—Europ.
subarctica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 61 = elatior.
suffrutescens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 371.—Calif.
sylvestris, Scop. Fl. Carn. ed. II. i. 132 = vulgaris.
Telemachica, Klatt, in Journ. Bot. vi. (1868) 121 = denticulata.
tenella, *King, ex Watt, in Journ. Linn. Soc.* xx. (1882) 13; *et Hook. f. Fl. Brit. Ind.* iii. 492.—Reg. Himal.
ternovania ×, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 77.—Europ.
Thomasinii, Gren. & Godr. Fl. Fr. ii. 449 = officinalis.
tibetica, Watt, in Journ. Linn. Soc. xx. (1882) 6; et ex Hook. Fl. Brit. Ind. iii. 488 = P. Pumilio.
Tommasinii, Nym. Consp. 603 = officinalis.
Tournefortii, Rupr. in Bull. Acad. Pétersb. vi. (1863) 230 = auriculata.
travnicensis ×, Wiesb. in Oestr. Bot. Zeitschr. xxxii. (1882) 282.—Europ.
tridentata, D. Don, Prod. Fl. Nep. 77 = petiolaris.
truncata, Lehm. Monog. Prim. 87. t. 8 = minima.
tschuktschorum, Kjellm. in Nordensk. Vega-Exped. Vetensk. Iaktt. i. 516 = nivalis.
turkestanica, Hort.—Cf. Gard. Chron. (1877) II. 809 = nivalis.
tyrolensis, *H. W. Schott, Sippen Primeln*, 13.—Europ. med.
undulata, Fisch. ex Reichb. Pl. Crit. v. 41. t. 468. f. 660 = farinosa.
unicolor, Lange, Dansk. Fl. ed. III. 181 = officinalis.
uniflora, C. C. Gmel. Fl. Bad. i. 442 = vulgaris.
uniflora, *Klatt, in Linnaea*, xxxvii. (1871–73) 500.—Reg. Himal.
uralensis, Fisch. ex Reichb. Pl. Crit. vii. 18. t. 638 = officinalis.
urticifolia, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 497.—China.
vaginata, *Watt, in Journ. Linn. Soc.* xx. (1882) 4.—Reg. Himal.
variabilis, Bast. Suppl. Fl. Maine-et-Loire, 26 = vulgaris.
variabilis ×, Goupil, in Mém. Soc. Linn. Par. iii. (1825) 248.—Gallia.
venusta ×, Host, Fl. Austr. i. 248.—Alp. austr.
Venzoi ×, Huter, ex Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 155.—Europ.
veris, Lehm. Monog. Prim. 27 = officinalis.
veris, Linn. Sp. Pl. 142 = elatior, vulgaris, officinalis.
veris, Mill. Gard. Dict. ed. VIII. n. 1; Hill, Herb. Crit. ii. 193 = vulgaris.
vernalis, Salisb. Prod. 117 = vulgaris.
verticillata, *Forsk. Fl. Aegypt. Arab.* 42.—Arab.
villosa, Curt. Bot. Mag. t. 14 = *pubescens*.
villosa, Lam. Fl. Fr. ii. 250 = Androsace villosa.
villosa, Lapeyr. Hist. Abr. Pl. Pyr. 96 = viscosa.
villosa, Lehm. Monog. Prim. 77 = hirsuta.
villosa, *Wulf. in Jacq. Fl. Austr.* v. 41.—Europ. med.
viscosa, *All. Fl. Pedem.* i. 93.—Europ.
viscosa, Gaud. Fl. Helv. ii. 89 = villosa.

PRIMULA:—
viscosa, Vill. Prosp. 21; Hist. Pl Dauph. ii. 467 = hirsuta.
viscosa, Waldst. & Kit. Pl. Rar. Hung. ii. p. xxviii = Kitaibeliana.
Vitaliana, Linn. Sp. Pl. 143 = Douglasia Vitaliana.
vulgaris, Hill, Herb. Brit. ii. 193 = officinalis.
vulgaris, *Huds. Fl. Angl.* ed. I. 70.—Europ.
Wattii, *King, ex Watt, in Journ. Linn. Soc.* xx. (1882) 10.—Reg. Himal.
Wulfeniana, *Schott, Wilde Blendl. Prim.* 17. t. 6.—Europ.
yunnanensis, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 269.—China.

PRIMULIDIUM, Spach, Hist. Vég. Phan. ix. 354 (1840) = **Primula**, Linn.
sinense, Spach, l. c. 355 = Primula sinensis.

PRIMULINA, Hance, in Journ. Bot. xxi. (1883) 169. *GESNERACEAE.*
Tabacum, *Hance, l. c.*—China.

PRINGLEA, Anders. ex Hook. f. Fl. Antarct. ii. 238. t. 90, 91 (1845). *CRUCIFERAE*, Benth. & Hook. f. i. 76.
DIAPHORANTHUS, Anders. ex Hook. f. Fl. Antarct. 238 (1845).
antiscorbutica, *R. Br. ex Hook. f. l. c.* 239. t. 90, 91.—Reg. Antarct.

PRINGLEOPHYTUM, A. Gray, in Proc. Am. Acad. xx. (1885) 292. *ACANTHACEAE.*
lanceolatum, *A. Gray, l. c.* 293.—Calif.

PRINODIA, Griseb. in Goett. Abh. vii. (1857) 224, in obs. = **Ilex**, Tourn.
bumelioides, Griseb. l. c. = I. bumelioides.
sideroxyloides, Griseb. l. c. = I. sideroxyloides.

PRINOS, Gronov. ex Linn. Coroll. Gen. 6 (1737) = **Ilex**, Tourn.
ambiguus, Michx. Fl. Bor. Am. ii. 236 = I. ambigua.
ambiguus, Nutt. Gen. Am. i. 213 = Nemopanthes fascicularis.
ambiguus, Pursh, Fl. Am. Sept. i. 220 = I. mollis.
asprellus, Hook. & Arn. Bot. Beech. Voy. 176. t. 36 = I. asprella.
atomarius, Nutt. Gen. Am. i. 213 = I. lucida.
canadensis, Lodd. ex Steud. Nom. ed. II. ii. 396 = Nemopanthus fascicularis?
cassinoides, Hort. ex Steud. l. c. = I. Cassine?
confertus, Moench, Meth. 481 = I. verticillata.
coriaceus, Benth. Pl. Hartw. 60.—Mexic.
coriaceus, Pursh, Fl. Am. Sept. i. 221 = I. lucida.
crenatus, Regel, Ind. Sem. Hort. Petrop. (1866) 103 = I. crenata?
cymosa, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 140 = I. cymosa.
deciduus, DC. Prod. ii. 16 = I. decidua.
dioicus, Vahl, Eclog. Am. ii. 25 = I. dioica.
dubius, G. Don, Gen. Syst. ii. 20 = I. mollis.
glaber, Linn. Sp. Pl. 330 = I. glaber.
glaber, Wats. Dendrol. i. t. 27 = I. lucida.
Godajam, Colebr. ex Wall. Pl. As. Rar. iii. 38 = I. Godajam.
Gronovii, Michx. Fl. Bor. Am. ii. 236 = I. verticillata.
hypoglaucus, Miq. Fl. Ind. Bat. Suppl. 514.—Ins. Banca.
integer, Hook. & Arn. Bot. Beech. Voy. 261 = I. integra?
integerrimus, Vell. Fl. Flum. 145; iii. t. 167 = I. integerrima.
integrifolius, Ell. Sketch, ii. 706 = Nemopanthus fascicularis.
lacunosa, Miq. Fl. Ind. Bat. Suppl. 513.—Sumatra.
laevigatus, Pursh, Fl. Am. Sept. i. 220 = I. laevigata.
lanceolatus, Hill, Veg. Syst. xvi. 57; Pursh, l. c. = I. lanceolata.
lanceolatus, Macfad. Fl. Jamaic. i. 206 = I. montana.
laurinus, Thury, ex Choisy, Mém. Ternstr. 44. t. 3, in obs.—Afr. austr.
longifolius, Desf. Tabl. ed. II. 272, nomen.—Am. bor.
longipes, Rafin. Sylva Tellur. 50 = Nemopanthus fascicularis.
lucidus, Ait. Hort. Kew. ed. I. i. 478 = I. lucida.
Macfadyenii, Walp. Rep. i. 541 = I. montana.

PRINOS:—
montanus, Sw. Prod. Veg. Ind. Occ. 58 = Ilex montana.
myricoides, Willd. ex D. Dietr. Syn. Pl. ii. 1050 = I. myricoides?
nitidus, Hort. ex C. Koch, Dendrol. ii. 1. 227 = I. laevigata.
nitidus, Vahl, Eclog. Am. ii. 26 = I. nitida.
obtusatus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 457 = I. obtusata.
obtusatus, Willd. ex D. Dietr. Syn. Pl. ii. 1050.—Am. austr.
padifolius, Willd. Enum. Hort. Berol. 394 = I. verticillata.
parvifolius, Rafin. Sylva Tellur. 50.—Hab.?
prunifolius, Desf. Tabl. ed. II. 272 = I. verticillata.
punctatus, Rafin. Sylva Tellur. 50.—Hab.?
reticulatus, Rafin. l. c.—Hab.?
rugosus, Rafin. l. c.—Hab.?
serratus, Vell. Fl. Flum. 145; iii. t. 166 = I. acrodonta.
sideroxyloides, Sw. Prod. Veg. Ind. Occ. 58 = I. montana.
spicatus, Miq. Fl. Ind. Bat. i. II. 594 = I. spicata.
undulatus, Hort. ex Steud. Nom. ed. I. 657, nomen.—Hab.?
verrucosus, Rafin. Sylva Tellur. 50.—Hab.?
verticillatus, Linn. Sp. Pl. 330 = I. verticillata.

PRINSEPIA, Royle, Illustr. Bot. Himal. 206. t. 38 (1834). *ROSACEAE*, Benth. & Hook. f. i. 611.
CYCNIA, Lindl. Veg. Kingd. 543 (1847).
utilis, *Royle, Illustr. Bot. Himal.* 206.—Reg. Himal.

PRINTZIA, Cass. in Dict. Sc. Nat. xxxvii. 463 (1825); xliii. 324 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 333.
LIOYDIA, Neck. Elem. i. 4 (1790).
LLOYDIA, Benth. & Hook. f. Gen.. ii. 333 (1873).
aromatica, *Less. in Linnaea*, v. (1830) 275.—Afr. austr.
auriculata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 514.—Afr. austr.
Bergii, *Cass. in Dict. Sc. Nat.* xliii. 324.—Afr. austr.
Huttoni, *Harv. Thes. Cap.* ii. 37.—Afr. austr.
pyrifolia, *Less. Syn. Comp.* 108.—Afr. austr.

PRIONACHNE, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836). *GRAMINEAE*, Benth. & Hook. f. iii. 1155.
CHONDROLAENA, Nees, Fl. Afr. Austr. 133 (1841).
KTENOSACHNE, Steud. Syn. Pl. Gram. 150 (1855).
dentata, Nees, Fl. Afr. Austr. 134, in syn. = seq.
Ecklonii, *Nees, in Lindl. Introd. Nat. Syst.* ed. II. 448.—Afr. austr.

PRIONANTHES, Schrank, Pl. Rar. Hort. Monac. ii. t. 51 (1819) = **Trixis**, P. Br. (Compos.).
antimenorrhoea, Schrank, l. c. = T. divaricata.

PRIONANTHIUM, Desv. Opusc. 64 (1831). *GRAMINEAE.*
rigidum, *Desv. l. c.* 65. t. 4.—Ind. or.

PRIONANTIUM, Desv. Opusc. t. 4 (1831) = praec.

PRIONITES, Pritz. Ic. Ind. i. 896 (1855), sphalm. = **Prionotes**, R. Br. (Epacrid.).

PRIONITIS, Adans. Fam. ii. 499 (1763) = **Falcaria**, Riv. (Umbell.).
Falcaria, Dum. Fl. Belg. 77 = F. vulgaris.
falcata, Delarb. Fl. Auv. ed. II. 421 = F. vulgaris.

PRIONITIS, Linn. Hort. Cliff. 487 (1737); Oerst. in Kjoeb. Vidensk. Meddel. (1854) 137 = **Barleria**, Linn. (Acanth.).
Hystrix, Miq. Fl. Ind. Bat. ii. 809 = B. Prionitis.
pubiflora, Miq. l. c. 810 = B. Prionitis.

PRIONIUM, E. Mey. in Linnaea, vii. (1832) 130. *JUNCACEAE*, Benth. & Hook. f. iii. 869.
Palmita, *E. Mey. l. c.* 131.—Afr. austr.
serrata, Buchen. in Bremen, Abh. iv. (1875) 408 = P. Palmita.

PRIONOLEPIS, Poepp. & Endl. Nov. Gen. et Sp. iii. 55. t. 261 (1842) = **Liabum**, Adans. (Compos.).
silphioides, Poepp. & Endl. l. c.—Peruv.

PRIONOPHYLLUM, C. Koch, Ind. Sem. Hort. Berol. App. iv. 3 (1873) = **Dyckia**, Schult. f. (Bromel.).
Selloum, C. Koch, l. c. 7.—Bras.

PRIONOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 118 = **Alloplectus**, Mart. (Gesner.).
Forseithii, Oerst. l. c. 119 = A. Forseithii.

PRIONOPSIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 329 = **Aplopappus**, Cass. (Compos.).
Chapmanii, Torr. & Gray, Fl. N. Am. ii. 245 = Aster eryngiifolius.
ciliata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 329 = A. ciliatus.

PRIONOSEPALUM, Steud. Syn. Pl. Cyp. 266 (1855) = **Chaetanthus**, R. Br. (Restiac.).
Gilberti, Steud. l. c. = C. leptocarpoides.

PRIONOSTACHYS, Hassk. in Flora, xlix. (1866) 212 = **Aneilema**, R. Br. (Commel.).
ensifolia, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 213 = A. giganteum.
herbacea, Hassk. ex C. B. Clarke, l. c. 205 = A. herbaceum.
terminalis, Hassk. ex C. B. Clarke, l. c. 211 = A. nudiflorum.

PRIONOSTEMMA, Miers, in Trans. Linn. Soc. xxviii. (1872) 354 = **Hippocratea**, Linn. (Celastrin.).
aspera, Miers, l. c. 355 = H. aspera.
Kunthiana, Miers, l. c. 359 = H. decussata.
malpighiaefolia, Miers, l. c. 356 = H. malpighiaefolia.
scabridula, Miers, l. c. 357 = H. Vahliana, *Peyr.*
setulifera, Miers, l. c. 359 (Hippocrateae sp.).—Guatemala.
velutina, Miers, l. c. 358 = Salacia velutina.

PRIONOTES, R. Br. Prod. 552 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 616.
americana, Hook. Ic. Pl. t. 30 = Lebetanthus americanus.
cerinthoides, *R. Br. Prod.* 553.—Tasman.
secunda, Spreng. Syst. i. 631 = Dracophyllum secundum.

PRIONOTIS, Benth. & Hook. f. Gen. ii. 616 (1876) = praec.

PRIOPETALON, Rafin. Fl. Tellur. iv. 34 (1836) = **Alstroemeria**, Linn. (Amaryll.).
pallidum, Rafin. l. c. 35 = A. Ligtu.

PRIORIA, Griseb. Fl. Brit. W. Ind. 215 (1864). *LEGUMINOSAE*, Benth. & Hook. f. i. 585.
copaifera, *Griseb. l. c.*—Jamaic.; Panama.

PRIOTROPIS, Wight & Arn. Prod. 180 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 480.
cytisoides, *Wight & Arn. l. c.*—Reg. Himal.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 509.—Ins. Socotra.

PRISCIANA, Rafin. Sylva Tellur. 114 (1838) = **Heliophila**, Burm. f. (Crucif.).
capensis, Rafin. l. c. = H. callosa.

PRISMATANTHUS, Hook. & Arn. Bot. Beech. Voy. 203 (1836) = **Siphonostegia**, Benth. (Scrophul.).
bartsioides, Hook. & Arn. l. c. = S. chinensis.

PRISMATOCARPUS, L'Herit. Sert. Angl. 1. t. 3 (1788), partim. *CAMPANULACEAE*, Benth. & Hook. f. ii. 559.
CODIPHUS, Rafin. Fl. Tellur. ii. 77 (1836).
IREON, Scop. Introd. 229 (1777), nomen prius.
ROELLOIDES, Banks, ex A. DC. Monog. Camp. 164 (1830).
altiflorus, *L'Hérit. Sert. Angl.* 2.—Afr. austr.
Bergianus, Cham. in Linnaea, viii. (1833) 199 = subulatus.
brevilobus, *A. DC. in DC. Prod.* vii. 443.—Afr. austr.
Burchellii, *Vatke, in Linnaea*, xxxviii. (1874) 701.—Afr. austr.
campanuloides, Sond. in Harv. & Sond. Fl. Cap. iii. 589 = strictus.

PRISMATOCARPUS:—
Candolleanus, *Cham. in Linnaea*, viii. (1833) 197.—Afr. austr.
confertus, Moench, Meth. 496 = Specularia hybrida.
cordatus, Reichb. Fl. Germ. Excurs. 858 = Specularia hybrida.
crispus, *L'Hérit. Sert. Angl.* 2.—Afr. austr.
diffusus, *A. DC. Monog. Camp.* 164.—Afr. austr.
Ecklonii, *A. DC. l. c.* 168.—Afr. austr.
falcatus, Tenore, Prod. Fl. Nap. 16 = Specularia falcata.
fastigiatus, *Presl, ex A. DC. in DC. Prod.* vii. 442.—Afr. austr.
fruticosus, Eckl. ex A. DC. Monog. Camp. 168 = Ecklonii.
fruticosus, *L'Hérit. Sert. Angl.* 2.—Afr. austr.
grandiflorus, E. Mey. ex A. DC. in DC. Prod. vii. 443 = paniculatus.
Hildebrandtii, *Vatke, in Linnaea*, xxxviii. (1874) 701.—Afr. austr.
hirsutus, G. Don, in Loud. Hort. Brit. 77 = seq.
hirtus, Tenore, Prod. Fl. Nap. 16 = Specularia Speculum.
hirtus, Urv. ex A. DC. Monog. Camp. 350 = Specularia coa.
Hookeri, Sweet, Hort. Brit. ed. II. 328 = nitidus.
hybridus, L'Hérit. Sert. Angl. 3 = Specularia hybrida.
interruptus, Eckl. ex A. DC. Monog. Camp. 165 = paniculatus.
interruptus, *L'Hérit. Sert. Angl.* 2.—Afr. austr.
junceus, Buek, in Eckl. & Zeyh. Enum. 383 = Lightfootia juncea.
laricinus, Presl, Prod. Monog. Lobel. 52 = diffusus.
linariaefolius, A. DC. Monog. Camp. 169 = strictus.
lycopodioides, A. DC. in DC. Prod. vii. 443 = subulatus.
nitidus, *L'Hérit. Sert. Angl.* 2. t. 3.—Afr. austr.
paniculatus, *L'Hérit. l. c.* 2.—Afr. austr.
pedunculatus, A. DC. in DC. Prod. vii. 443 = paniculatus.
pentagonius, L'Hérit. Sert. Angl. 3 = Specularia pentagonia.
perfoliatus, Sweet, Hort. Brit. ed. I. 251 = Specularia perfoliata.
roelloides, Sond. in Harv. & Sond. Fl. Cap. iii. 586 = paniculatus.
scaber, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimp. 16 = Specularia falcata.
sessilis, *Eckl. ex A. DC. Monog. Camp.* 171.—Afr. austr.
Speculum, L'Hérit. Sert. Angl. 3 = Specularia Speculum.
strictus, *A. DC. Monog. Camp.* 169.—Afr. austr.
subulatus, *A. DC. l. c.* 166.—Afr. austr.
tenellus, *Oliver, in Hook. Ic. Pl.* t. 1460.—Afr. austr.
tenerrimus, *Buek, in Eckl. & Zeyh. Enum.* 384.—Afr. austr.

PRISMATOMERIS, Thw. in Hook. Kew Journ. viii. (1856) 268. t. 7 A. *RUBIACEAE*, Benth. & Hook. f. ii. 118.
albidiflora, *Thw. l. c.* 269.—Ind. or.; Malaya.
Fergusonii, *Thw. ex Bedd. Forest Man. Bot.* 134/10.—Zeylan.

PRISTIDIA, Thw. Enum. Pl. Zeyl. 149 (1859) = **Gaertnera**, Lam. (Logan.).
divaricata, Thw. l. c. = G. Koenigii.

PRISTIMERA, Miers, in Trans. Linn. Soc. xxviii. (1872) 360 = **Hippocratea**, Linn. (Celastrin.).
andina, Miers, l. c. 364.—Peruv.
apiculata, Miers, l. c. = H. integrifolia, *Seem.*
coriacea, Miers, l. c. 365 = H. coriacea.
granulosa, Miers, l. c. 363 = H. malpighiaefolia.
lepida, Miers, l. c. 361.—Venezuela.
oblongata, Miers, l. c. 362 = H. oblongata.
tenella, Miers, l. c. 365 = H. acapulcensis?
verrucosa, Miers, l. c. 364 = H. verrucosa.
Wrightiana, Miers, l. c. 363 = H. malpighiaefolia.

PRISTOCARPHA, E. Mey. ex DC. Prod. vi. 86 (1837) = **Athanasia**, Linn. (Compos.).
capitata, E. Mey. ex DC. l. c. = A. capitata.

PRITCHARDIA, Seem. & H. Wendl. in Bonplandia, ix. (1861) 260; x. (1862) 197. *PALMAE*, Benth. & Hook. f. iii. 928.
filifera, *Linden, ex André, in Illustr. Hortic.* xxiv. (1877) 32, 105.—Am. bor. occ.

PRITCHARDIA:—
Gaudichaudii, *H. Wendl. in Bonplandia*, x. (1862) 199.—Ins. Pacif.
grandis, Hort. Bull.—Cf. Gard. Chron. (1874) i. 415 = Licuala grandis.
macrocarpa, *Linden, ex André, in Illustr. Hortic.* xxvi. (1879) 105 t. 352.—Ins. Sandvic.
Martii, *H. Wendl. in Bonplandia*, x. (1862) 199.—Ins. Pacif.
pacifica, *Seem. & H. Wendl. l. c.* ix. (1861) 260; x. (1862) 197.—Ins. Fiji.
Vuylstekeana, *Hort.—Cf. Gard. Chron.* (1883) i. 693. f. 114.—Ins. Paumotu.

PRITZELIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126; Abh. Akad. Berl. (1855) 107. t. 10 B = **Begonia**, Linn.
angulata, Klotzsch, ex Wawra, Bot. Ergeb. Maxim. Bras. 51 = B. angulata.
coccinea, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126; Abh. (1855) 108 = B. coccinea.
deflexa, A. DC. in DC. Prod. xv. i. 362, in syn. = B. scandens.
Fischeri, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126; Abh. (1855) 108 = B. Fischeri.
glabra, A. DC. in DC. Prod. xv. i. 362, in syn. = B. scandens.
glauca, Klotzsch, in Monatsb. Berl. Akad. (Maerz. 1854) 126; Abh. (1855) 109 = B. glauca.
lucida, A. DC. in DC. Prod. xv. i. 362, in syn. = B. scandens.
montana, A. DC. l. c. in syn. = B. scandens.
Princeps, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126; Abh. (1855) 111 = B. Princeps.
ramentacea, Klotzsch, ll. cc. = B. ramentacea.
sanguinea, Klotzsch, ll. cc. 126; 109 = B. sanguinea.
zebrina, Klotzsch, ll. cc. 126; 110 = B. angularis.

PRITZELIA, F. Muell. Desc. Papuan Pl. i. 13 (1875).
PHILYDRACEAE, Benth. & Hook. f. iii. 840.
HETAERIA, Endl. Gen. 133 (1837).
PHILYDRELLA, Caruel, in Nuov. Giorn. Bot. Ital. x. (1878) 91.
pygmaea, *F. Muell. Desc. Papuan Pl.* i. 13.—N. Guin.

PRITZELIA, Schau. in Flora, xxvi. (1843) 405, 407, nomen = **Scholtzia**, Schau. (Myrtac.).
obovata, Schau. ex Walp. Rep. ii. 922 = S. obovata.

PRITZELIA, Walp. Rep. ii. 428 (1843) = **Trachymene**, Rudge (Umbell.).
didiscoides, Walp. l. c. = T. pilosa.

PRIVA, Adans. Fam. ii. 505 (1763). *VERBENACEAE*, Benth. & Hook. f. ii. 1145.
BLAIRIA, Houst. ex Linn. Gen. ed. I. 334, in syn. (1737).
CASTELIA, Cav. in Anal. Cienc. Nat. iii. (1801) 134.
PHELLODERMA, Miers, in Trans. Linn. Soc. xxvii. (1870) 100. t. 27.
PHRYMA, Forsk. Fl. Aegypt. Arab. p. cxv (1775).
PITRAEA, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) ii. 328.
STREPTIUM, Roxb. Pl. Corom. ii. 25. t. 146 (1798).
TORTULA, Roxb. ex Willd. Sp. Pl. iii. 359 (1800).
abyssinica, Jaub. & Spach, Illustr. v. 58. t. 453 = leptostachya.
aspera, *H. B. & K. Nov. Gen. et Sp.* ii. 278.—Mexic.
bahiensis, *DC. Prod.* xi. 533.—Bras.
crenata, Schrad. Ind. Sem. Hort. Gotting. (1831).—Cf. Linnaea, viii. (1833) Litt. 24 = aspera.
dentata, Juss. in Ann. Mus. Par. vii. (1806) 70; Pers. Syn. ii. 139 = leptostachya.
echinata, *Juss. l. c.* 69.—Am. trop.
Forskâlii, Jaub. & Spach, Illustr. v. 59. t. 455 = leptostachya.
Forskaolii, E. Mey. Comm. Pl. Afr. Austr. 275 = leptostachya.
glaberrima, Bocq. Rev. Verb. 116. t. 18 = Dipyrena glaberrima.
hispida, *Juss. in Ann. Mus. Par.* vii. (1806) 69.—Mexic.
laevis, *Juss. l. c.* 70.—Am. austr.
lamiifolia, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 325 = echinata.

PRIVA:—
lappulacea, Pers. Syn. ii. 139 = echinata.
leptostachya, *Juss. in Ann. Mus. Par.* vii. (1806) 70.—Ind. or.; Afr. trop.
mexicana, Pers. Syn. ii. 139 = hispida.
mexicana, Sieber, ex Steud. Nom. ed. II. ii. 397 = echinata.
Meyeri, Jaub. & Spach, Illustr. v. 57 = dentata.
orchidioides, Walp. Rep. iv. 36 = laevis.
trachelioides, Mart. & Gal. in Bull. Acad. Brux. xi. ii. (1844) 324 = aspera.
tuberosa, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 135.—Mexic.
virgata, Spreng. Syst. ii. 753 = Lippia urticoides.

PROBLASTES, Reinw. in Syll. Ratisb. ii. (1828) 10 = **Lumnitzera**, Willd. (Combret.).
cuneifolia, Reinw. l. c.—Java.

PROBOSCIDEA, Schmid. Anal. 49. t. 12, 13 (1747) = **Martynia**, Houst. (Pedalin.).
althaeifolia, Decne. in Ann. Sc. Nat. Sér. V. iii. (1865) 324 = M. altheaefolia.
arenaria, Decne. l. c. 326 = M. altheaefolia.
Botterii, Decne. l. c. 327 = M. Botterii.
cordifolia, Moench, Meth. 466 = M. Proboscidea.
fallax, Decne. in Ann. Sc. Nat. Sér. V. iii. (1865) 325 = Craniolaria fallax.
fragrans, Decne. l. c. 326 = M. fragrans.
Jussieui, Steud. Nom. ed. II. ii. 397 = M. Proboscidea.
triloba, Decne. in Ann. Sc. Nat. Sér. V. iii. (1865) 326 = M. triloba.
unibracteata, Decne. l. c. 325.—Bras.
violacea, Decne. l. c. = M. fragrans.

PROBOSCIDIA, Rich. ex DC. Prod. iii. 106 (1828) = **Rhynchanthera**, DC. (Melastom.).

PROBOSCIPHORA, Neck. Elem. i. 336 (1790) = **Rhynchocorys**, Griseb. (Scrophul.).

PROCKIA, P. Br. ex Linn. Syst. ed. X. 1074 (1759).
TILIACEAE, Benth. & Hook. f. i. 237.
JACQUINIA, Mutis, ex Linn. Mant. ii. 247 (1771).
KELLETTIA, Seem. Bot. Voy. Herald, 85, 254 (1853).
TINEA, Spreng. Neue Entdeck. ii. 165 (1821).
TRILIX, Linn. Mant. ii. 153 (1771).
acuta, [Moç. & Sesse,] ex DC. Prod. i. 260 = P. Crucis.
albicans, Blanco, Fl. Filip. ed. I. 430 = Alchornea Blumeana.
bahiensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 335.—Bras.
completa, Hook. Ic. Pl. t. 94 = septemnervia.
crucifolia, C. Muell. in Walp. Ann. vii. 451 = morifolia.
Crucis, *Linn. Syst.* ed. X. 1074.—Am. trop.
deltoides, *Lam. Illustr.* iii. 22.—Ins. Mascar.
flava, *Karst. Fl. Columb.* ii. 79. t. 140.—N. Granat.
integrifolia, *Willd. Sp. Pl.* ii. 1214.—Ins. Mascar.
laciniata, Poir. Encyc. v. 627 = theaeformis.
lobata, *Poir. l. c.*—Hab.?
lucida, *Wall. Cat.* n. 7245; *Steud. Nom.* ed. II. ii. 398.—Ind. or.
luzoniensis, *Presl, Rel. Haenk.* ii. 94.—Ins. Philipp.
macrostachya, [*Moç. & Sesse,*] *ex DC. Prod.* i. 261.—Mexic.
mexicana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 334.—Mexic.
morifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 357.—N. Granat.
obovata, Presl, Rel. Haenk. ii. 94 = Xylosma nitidum.
odorata, Domb. ex Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 288 = Banara mollis.
ovata, Lam. Illustr. iii. 22 = integrifolia.
racemosa, *Hort. Lecchi, ex Colla, Hort. Ripul.* 113.—Hab.?
rotundifolia, Eckl. & Zeyh. Enum. 16 = Dovyalis celastrioides.
septemnervia, *Spreng. Syst.* ii. 609.—Bras.
serrata, Willd. Sp. Pl. ii. 1213 = Thiodia serrata.
subcordata, [Moç. & Sesse,] ex DC. Prod. i. 260 = P. Crucis.
theaeformis, *Willd. Sp. Pl.* ii. 1214.—Ins. Mascar.
tomentosa, A. Rich. Ess. Fl. Cub. 218 = P. Crucis.
villosa, *Moric. Pl. Nouv. Am.* i. 67. t. 97.—Bras.

PROCLESIA, Klotzsch, in Linnaea, xxiv. (1851) 32 = **Cavendishia**, Lindl. (Vaccin.).
acuminata, Klotzsch, l. c. 34 = C. acuminata.
alata, Klotzsch, l. c.—Peruv.
alnifolia, Klotzsch, l. c. 35.—Venezuela.
Benthamiana, Klotzsch, l. c. 33 = C. acuminata.
bracteata, Klotzsch, l. c. 34.—Peruv.
capitata, Klotzsch, l. c. = C. nobilis.
cordata, Klotzsch, l. c. 33.—Venezuela.
cordifolia, Klotzsch, l. c. 32.—N. Granat.
Hartwegiana, Klotzsch, l. c. 35.—Ecuador.
melastomoides, Klotzsch, l. c. 32 = C. melastomoides.
Pseudo-pubescens, Klotzsch, l. c. 33.—N. Granat.
pubescens, Klotzsch, l. c. = C. pubescens.
scabriuscula, Klotzsch, l. c.—Ecuador.
splendens, Klotzsch, l. c. 34.—N. Granat.
strobolifera, Klotzsch, l. c. 32.—N. Granat.
veraguensis, Klotzsch, l. c. 35 = C. veraguensis.
Warszewiczii, Klotzsch, l. c. = C. Warscewiczii.

PROCRASSULA, Griseb. Spicil. Fl. Rumel. i. 323 (1843) = **Sedum**, Tourn. (Crassul.).
caespitosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 384 = S. caespitosum.
Magnolii, Griseb. Spicil. Fl. Rumel. i. 323 = S. caespitosum.
mediterranea, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 16 = S. caespitosum.
pallidiflora, Jord. & Fourr. l. c. = S. rubens.

PROCRIS, Comm. ex Juss. Gen. 403 (1789). *URTICACEAE*, Benth. & Hook. f. iii. 386.
SCIOBIA, Reichb. Nom. 66 (1841).
SCIOPHILA, Gaudich. in Freyc. Voy. Bot. 493. t. 120 (1826).
acuminata, Poir. Encyc. v. 629 = Elatostema acuminatum.
approximata, Wall. Cat. n. 4630 = Elatostema cuneatum.
australis, Spreng. Syst. iii. 846 = Elatostema sessile.
axillaris, *J. F. Gmel. Syst.* 267 (Quid ?).—Hab. ?
cantoniensis, Spreng. Syst. iii. 846 (alius ordinis).—China.
caudata, Poir. Encyc. v. 629 = Boehmeria caudata.
caulescens, Poir. Encyc. Suppl. iv. 554, in obs. (alius ordinis).—Hab. ?
cephalida, *Comm. ex Poir. Encyc.* v. 629.—Malaya; Ins. Mascar.
chinensis, Spreng. Syst. iii. 846 (alius ordinis).—China.
cochinchinensis, Spreng. l. c. (alius ordinis).—Cochinch.
cuneata, Blume, Bijdr. 509 = laevigata.
cyrtandraefolia, Zoll. & Mor. Syst. Verz. Zoll. 74 = Elatostema nigrescens.
divaricata, Steud. Nom. ed. II. ii. 398 = Pellionia divaricata.
diversifolia, Wall. Cat. n. 4631 = Elatostema surculosum.
elatostemoides, Steud. Nom. ed. II. ii. 398 = Pellionia elatostemoides.
elegans, Wall. Cat. n. 4632 = Elatostema surculosum.
erecta, Blanco, Fl. Filip. ed. I. 707 = Conocephalus erectus.
excelsa, Steud. in Flora, xxxiii. (1850) 261 = Boehmeria excelsa.
fagifolia, Poir. Encyc. v. 629 = Elatostema fagifolium.
ficoidea, Wall. Cat. n. 4635 = Elatostema ficoides.
formidata, Span. in Linnaea, xv. (1841) 344 = Laportea peltata.
frutescens, *Blume, Bijdr.* 510.—Malaya.
gibbosa, Wall. Cat. n. 7273 = Pellionia Daveauana.
glabra, Hook. & Arn. Bot. Beech. Voy. 96 = Urera sandwicensis.
gracilis, Blume, Bijdr. 513 = Elatostema gracile.
grandis, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 337.—Ins. Philipp.
Heyneana, Wall. Cat. n. 7272 = Pellionia Heyneana.
hirsuta, Blume, Bijdr. 509 = Elatostema hirsutum.
hirta, Poir. Encyc. v. 629 = Phenax urticaefolius.
hypocrypta, Comm. ex Poir. Encyc. v. 629 = Elatostema fagifolium.
hypoleuca, Hochst. ex Steud. in Flora, xxxiii. (1850) 261 = Debregeasia hypoleuca.
integrifolia, Boj. Hort. Maurit. 299 = cephalida.
integrifolia, D. Don, Prod. Fl. Nep. 61 = Elatostema sesquifolium.
laeta, Wall. Cat. n. 4637 = Elatostema surculosum.

PROCRIS:—
laevigata, *Blume, Bijdr.* 508.—As. et Afr. trop.
lateriflora, Poir. Encyc. Suppl. iv. 554 = Boehmeria cylindrica.
latifolia, *Blume, Bijdr.* 509.—Java.
littoralis, Poir. Encyc. v. 628 = Boehmeria cylindrica.
longifolia, Blume, Bijdr. 508 = cephalida.
longifolia, Poepp. ex Wedd. in DC. Prod. xvi. 1. 194 = Phenax angustifolius.
longifolia, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 167 = Villebrunea scabra.
lucida, Spreng. Syst. iii. 846 = cephalida.
macrophylla, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 222 = Boehmeria rugosissima.
maculata, Poir. Encyc. v. 628 = Elatostema maculatum.
melastomoides, *Spreng. Syst.* iii. 847.—Ins. S. Doming.
membranacea, Blume, Bijdr. 512 = Elatostema acuminatum.
mollis, Wall. Cat. n. 4633 = Elatostema molle.
monandra, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 61 = Elatostema surculosum.
montana, *Steud. Nom.* ed. II. ii. 398.—Ins. Norfolk.
nivea, Wedd. in DC. Prod. xvi. 1. 206, in syn. = Boehmeria nivea.
nudiflora, Poir. Encyc. Suppl. iv. 454 = Boehmeria nudiflora.
obtusa, Royle, Illustr. Bot. Himal. t. 83. f. 3 = Lecanthus Wightii.
obtusa, Wall. Cat. n. 4636 = Elatostema obtusum.
paludosa, Blume, Bijdr. 510 = Elatostema macrophyllum.
parva, Blume, l. c. 512 = Elatostema parvum.
pedunculares, Wall. Cat. n. 4634 = Lecanthus Wightii, Elatostema papillosum.
pedunculata, Wedd. in DC. Prod. xvi. 1. 191 = cephalida.
piperifolia, Wall. Cat. n. 4628 = Elatostema sesquifolium.
polycephala, Wall. l. c. n. 4629 = Elatostema sessile.
punctata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 61 = Elatostema sessile.
punctata, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 167 = Villebrunea scabra.
punctata, Wall. Cat. n. 4627 = Elatostema rupestre, lineolatum.
quinquenervia, Spreng. Syst. iii. 846 (alius ordinis).—Ins. S. Doming.
racemosa, Royle, Illustr. Bot. Himal. t. 83. f. 1 = Pilea Wightii.
radicans, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 218 = Pellionia radicans.
ramiflora, Poir. Encyc. v. 630 = Boehmeria ramiflora.
rostrata, Reinw. ex Blume, Bijdr. 510 = Pellionia elatostemoides.
rugosa, Poir. Encyc. v. 628 = Phenax ballotaefolius.
rupestris, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 60 = Elatostema rupestre.
rupicola, *Zipp. ex Span. in Linnaea*, xv. (1841) 344, nomen.—Ins. Timor.
sesquifolia, Reinw. ex Blume, Bijdr. 511 = Elatostema sesquifolium.
sessilis, Hook. & Arn. Bot. Beech. Voy. 70 = Elatostema sessile.
Sieberi, Steud. Nom. ed. II. ii. 398 = Chlorophora tinctoria.
sinuata, Blume, Bijdr. 511 = Elatostema sinuatum.
Sonnerati, Spreng. Syst. iii. 847 = Phenax vulgaris.
splendens, Lindl. Veg. Kingd. ed. III. 260. f. 175 = Boehmeria australis.
strigosa, Blume, Bijdr. 512 = Elatostema strigosum.
torresiana, Steud. Nom. ed. II. ii. 398 = cephalida.
trinervata, Poir. Encyc. v. 628 = Gyrotaenia trinervata.
umbellata, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 217 = Pellionia umbellata.
urticaefolia, Poir. Encyc. v. 628 = Phenax urticaefolius.
violacea, Blanco, Fl. Filip. ed. I. 706 = Conocephalus suaveolens.
Wightiana, Wall. Cat. n. 4638 = laevigata.

PROINEIA, Ehrh. Beit. iv. 149 (1789) = **Aira**, Linn. (Gramin.).

PROIPHYS, Herb. App. [Bot. Reg.] 42 (1821) = **Eurycles**, Salisb. (Amaryll.).
amboinensis, Herb. l. c. = E. sylvestris.

PROLONGOA, Boiss. Voy. Bot. Espagne, ii. 320 (1840) = **Chrysanthemum**, Tourn. (Compos.).
pectinata, Boiss. l. c. t. 93 = C. pectinatum.
Pseudanthemis, Kunze, in Flora, xxix. (1846) 699.—Hispan.
setabensis, Nym. Syll. 10 = C. paludosum.

PROMENAEA, Lindl. Bot. Reg. (1843) Misc. 13 = **Zygopetalum**, Hook. (Orchid.).
citrina, [P. N. Don,] Donn, Hort. Cantab. ed. XIII. 720 = Z. xanthinum.
florida, Reichb. f. in Hamb. Gartenz. xix. (1863) 12.—Bras.
graminea, Lindl. Bot. Reg. (1843) Misc. 13 (Aganosmatis sp.).—Guian.
guttata, Reichb. f. in Bonplandia, iv. (1856) 323 = Z. guttatum.
lentiginosa, Lindl. Bot. Reg. (1843) Misc. 13 = Z. lentiginosum.
microptera, Reichb. f. in Gard. Chron. (1881) II. 134 = Z. micropterum.
Rollissonii, Lindl. Bot. Reg. (1843) Misc. 13 = Z. Rollissonii.
stapelioides, Lindl. l. c. = Z. stapelioides.
xanthina, Lindl. l. c. = Z. xanthinum.

PRONACRON, Cass. in Dict. Sc. Nat. xliii. 370 (1826) = **Melampodium**, Linn. (Compos.).
ramosissima, Cass. l. c. = M. camphoratum.

PRONAYA, Hueg. Bot. Archiv, t. 6 (1837). *PITTOSPOREAE*, Benth. & Hook. f. i. 133.
CAMPYLANTHERA, Hook. Ic. Pl. t. 82 (1837).
SPIRANTHERA, Hook. Bot. Mag. sub t. 3523 (1836), nomen prius.
angustifolia, Lehm. Del. Sem. Hort. Hamb. (1845).—Cf. Linnaea, xix. (1847) 403 = Billardiera Lehmanniana.
elegans, *Hueg. Bot. Archiv*, t. 6.—Austral.
ericoides, Walp. Rep. ii. 770 = Marianthus procumbens.
Huegeliana, Putterl. in Lehm. Pl. Preiss. i. 204 = Billardiera variifolia.
lanceolata, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 364 = Billardiera variifolia ?
latifolia, Turcz. l. c. 363 = Billardiera coriacea.
Muelleriana, *Turcz. l. c.* xxxvi. (1863) 1. 561.—Austral.
pedunculosa, *Turcz. l. c.* 560.—Austral.
sericea, Turcz. l. c. xxvii. (1854) II. 363 = Billardiera variifolia.
speciosa, Endl. in Enum. Pl. Hueg. 9 = elegans.

PROSARTES, D. Don, in Proc. Linn. Soc. i. (1839) 48; et Trans. Linn. Soc. xviii. (1841) 531 = **Disporum**, Salisb. (Liliac.).
Hookeri, Torr. in Pacif. Rail. Rep. iv. 144 = D. Hookeri.
lanuginosa, D. Don, in Proc. Linn. Soc. i. (1839) 48; et Trans. Linn. Soc. xviii. (1841) 532 = D. lanuginosum.
Menziesii, D. Don, ll. cc. 48; 533 = D. Menziesii.
oregana, S. Wats. in Proc. Am. Acad. xiv. (1879) 271.—Am. bor. occ.
parvifolia, S. Wats. in Bot. Calif. ii. 179.—Calif.
trachyandra, Torr. in Pacif. Rail. Rep. iv. 144 = D. lanuginosum.
trachycarpa, S. Wats. Bot. King's Exped. 344 = D. trachycarpum.
viridescens, Regel, Tent. Fl. Ussur. 148.—Mandchuria.

PROSCEPHALEIUM, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 248. *RUBIACEAE*, Benth. & Hook. f. ii. 122.
javanicum, *Korth. l. c.*—Java.

PROSCEPHALIUM, Benth. & Hook. f. Gen. ii. 122 (1873) = praec.

PROSELIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 234 = **Chaetanthera**, Ruiz & Pav. (Compos.).
serrata, D. Don, l. c. 235 = C. chilensis.

PROSELIAS, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = **Astragalus**, Tourn. (Legumin.).
globosus, Stev. l. c. = A. globosus.
incanus, Stev. l. c. = A. incanus.
monspessulanus, Stev. l. c. = A. monspessulanus.
reduncus, Stev. l. c. = A. reduncus.
resupinatus, Stev. l. c. = A. fragrans.
sanguinolentus, Stev. l. c. = A. sanguinolentus.

PROSERPINACA, Linn. in Act. Soc. Sc. Upsal. (1741) 81. *HALORAGEAE*, Benth. & Hook. f. i. 675.
TRIXIS, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748), App 220.
palustris, *Linn. Sp. Pl.* 88.—Am. bor.
pectinata, *Lam. Illustr.* i. 214.—Am. bor.

PROSOPANCHE, De Bary, in Abh. Naturf. Ges. Halle, x. (1868) 267. tt. 1, 2. *CYTINACEAE*, Benth. & Hook. f. iii. 120.
Burmeisteri, *De Bary, l. c.*—Reg. Argent.

PROSOPIA, Reichb. Consp. 121 (1828)=**Pedicularis**, Tourn. (Scrophul.).

PROSOPIS, Linn. Mant. i. 10 (1767). *LEGUMINOSAE*, Benth. & Hook. f. i. 591.
ALGAROBIA, Benth. Pl. Hartw. 13 (1839).
LAGONYCHIUM, Bieb. Fl. Taur. Cauc. iii. 288 (1819).
MITOSTAX, Rafin. Sylva Tellur. 120 (1838).
NELTUMA, Rafin. l. c. 119 (1838).
PLEUROMENES, Rafin. l. c. 144 (1838).
SPIROLOBIUM, Orbign. Sert. Patag. t. 13 (1839).
STROMBOCARPA, A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 243.
abbreviata, *Benth. in Hook. Journ. Bot.* iv. (1842) 352.—Am. austr.
adesmioides, *Griseb. in Goett. Abh.* xix. (1874) 132.—Reg. Argent.
adstringens, Steud. Nom. ed. II. ii. 398 = *astringens*.
affinis, Spreng. Syst. ii. 326 = juliflora.
alba, *Griseb. in Goett. Abh.* xix. (1874) 131.—Reg. Argent.
Algarobilla, *Griseb. l. c.* (Abgarobilla).—Reg. Argent.
Alpataco, *Phil. in Anal. Univ. Chil.* ii. (1862) 394; *Linnaea*, xxxiii. (1864–65) 62.—Chili.
astringens, Gill. ex Hook. Bot. Misc. iii. (1833) 204 = Acacia atramentaria.
Bonplandiana, Spreng. Syst. iv. Cur. Post. 165 = Entada polystachya.
bracteolata, DC. Prod. ii. 447 = juliflora.
campestris, *Griseb. in Goett. Abh.* xix. (1874) 132.—Reg. Argent.
cinerascens, *A. Gray, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 381.—Texas.
cumanensis, H. B. & K. Nov. Gen. et Sp. vi. 310 = juliflora.
denudans, *Benth. in Hook. Journ. Bot.* iv. (1842) 351.—Am. austr.
domingensis, DC. Prod. ii. 447 = juliflora.
dubia, Guill. & Perr. Fl. Seneg. Tent. i. 242 = Acacia Sieberiana.
dubia, H. B. & K. Nov. Gen. et Sp. vi. 309 = Enterolobium cyclocarpum?
dulcis, Gill. ex Hook. Bot. Misc. iii. (1833) 203 = flexuosa.
dulcis, *Kunth, Mim.* 110. t. 34 = juliflora.
elegans, Spreng. Syst. iv. Cur. Post. 165 = Piptadenia rigida *vel* trisperma.
elephantina, E. Mey. Comm. Pl. Afr. Austr. 165 = Elephantorrhiza Burchellii.
elephantorrhiza, Spreng. Syst. iv. Cur. Post. 165 = Elephantorrhiza Burchellii.
Emoryi, Torr. in Emory, Notes Mil. Reconnois. 139 = pubescens.
ephedrioides, Gill. ex Hook. Bot. Misc. iii. (1833) 204 = Mimosa ephedroides.
faeculifera, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 426 = Parkia africana.
ferox, *Griseb. in Goett. Abh.* xxiv. (1879) 118.—Reg. Argent.
flexuosa, DC. Prod. ii. 447 = juliflora.
fruticosa, Meyen, Reise, i. 376 = juliflora.
glandulosa, Torr. in Ann. Lyc. N. York, ii. (1828) 192. t. 2 = juliflora.
globosa, Gill. ex Hook. Bot. Misc. iii. (1833) 205 = Mimosa Gilliesii.
heterophylla, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 82.—Mexic.
horrida, Kunth, Mim. 106. t. 33 = juliflora.
humilis, *Gill. ex Hook. Bot. Misc.* iii. (1833) 204.—Reg. Argent.
inermis, Gill. ex Hook. l. c. 203 = dulcis.
inermis, H. B. & K. Nov. Gen. et Sp. vi. 307 = juliflora.
juliflora, *DC. Prod.* ii. 447.—Am. trop.
Kirkii, Oliver, Fl. Trop. Afr. ii. 332 = Acacia albida.

PROSOPIS:—
lanceolata, Benth. in Hook. Journ. Bot. iv. (1842) 347 = oblonga.
limensis, *Benth. l. c.* 350.—Peruv.
microphylla, H. B. & K. Nov. Gen. et Sp. vi. 308 = Acacia tortuosa.
Nandubey, *Lorentz, ex Griseb. in Goett. Abh.* xxiv. (1879) 117.—Reg. Argent.
nigra, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 283.—Reg. Argent.
oblonga, *Benth. in Hook. Journ. Bot.* iv. (1842) 348.—Afr. trop.
odorata, Torr. & Frémont, Rep. 313. t. 1 = juliflora, pubescens.
pallida, H. B. & K. Nov. Gen. et Sp. vi. 309 = juliflora.
Panta, *Hieron. in Bol. Acad. Nac. Córdora*, iv. (1881) 284.—Reg. Argent.
pubescens, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 82.—Texas; Mexic.; Calif.
reptans, *Benth. in Hook. Journ. Bot.* iv. (1842) 352.—Am. austr.
ruscifolia, *Griseb. in Goett. Abh.* xix. (1874) 130.—Reg. Argent.
scandens, Stokes, Bot. Mat. Med. ii. 468 = Entada scandens.
sericantha, *Gill. ex Hook. Bot. Misc.* iii. (1833) 204.—Am. austr.
Siliquastrum, DC. Prod. ii. 447 = juliflora.
siliquosa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = P. Siliquastrum.
spicata, Burm. f. Fl. Ind. 102 = spicigera.
spicigera, *Linn. Mant.* i. 68.—Persia; Ind. or.
Stephaniana, *Kunth, ex Spreng. Syst.* ii. 326.—Reg. Caucas.; Ind. bor. occ.
striata, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 83.—Patagon.
strombulifera, *Benth. in Hook. Journ. Bot.* iv. (1842) 352.—Am. austr.
tamariscina, Spreng. Syst. iv. Cur. Post. 165 = Gagnebina tamariscina.
torquata, *DC. Prod.* ii. 448.—Am. austr.

PROSORUS, Dalz. in Hook. Kew Journ. iv. (1852) 345 = **Phyllanthus**, Linn. (Euphorb.).
cyanospermus, Thw. Enum. Pl. Zeyl. 281 = Ph. cyanospermus.
Gaertneri, Thw. in Hook. Kew Journ. viii. (1856) 272 = Ph. cyanospermus.
indicus, Dalz. in Hook. l. c. iv. (1852) 346 = Ph. indicus.
indicus, Thw. in Hook. l. c. vi. (1854) 299, partim = Ph. cyanospermus.

PROSPERO, Salisb. Gen. Pl. Fragm. 28 (1866) = **Scilla**, Linn. (Liliac.).
autumnale, Salisb. l. c. = S. autumnalis.
hyacinthoideum, Salisb. l. c. = S. hyacinthoides.
lingulatum, Salisb. l. c. = S. lingulata.
obtusifolium, Salisb. l. c. = S. obtusifolia.
parviflorum, Salisb. l. c. = S. numidica.
undulatum, Salisb. l. c. = Urginea undulata.

PROSPHYSIS, Dulac, Fl. Hautes-Pyr. 67 (1867) = **Festuca**, Linn. (Gramin.).
tenella, Dulac, l. c. 68 = F. montana?

PROSTANTHERA, Labill. Nov. Holl. Pl. ii. 18. t. 157 (1806). *LABIATAE*, Benth. & Hook. f. ii. 1217.
CHILODIA, R. Br. Prod. 507 (1810).
CRYPHIA, R. Br. l. c. 508 (1810).
KLANDERIA, F. Muell. in Linnaea, xxv. (1852) 426.
aspalathoides, *A. Cunn. ex Benth. Lab. Gen. et Sp.* 453.—Austral.
atriplicifolia, A. Cunn. l. c. 451 = ovalifolia.
Baxteri, *A. Cunn. l. c.* 452.—Austral.
Behriana, *Schlecht. in Linnaea*, xx. (1847) 610.—Austral.
caerulea, *R. Br. Prod.* 508.—Austral.
Caleyi, Benth. Lab. Gen. et. Sp. 454 = chlorantha?
calycina, *F. Muell. ex Benth. Fl. Austral.* v. 107.—Austral.
Campbellii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 252.—Austral.
canaliculata, *F. Muell. Fragm.* vi. 105.—Austral.
chlorantha, *F. Muell. ex Benth. Fl. Austral.* v. 108.—Austral.

PROSTANTHERA:—
coccinea, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 48 = microphylla.
cotinifolia, A. Cunn. ex Benth. Lab. Gen. et Sp. 452 = rotundifolia.
cryptandroides, *A. Cunn. l. c.* 453.—Austral.
cuneata, *Benth. in DC. Prod.* xii. 560.—Austral.
debilis, *F. Muell. Fragm.* viii. 147.—Austral.
decussata, *F. Muell. l. c.* i. 125.—Austral.
denticulata, *R. Br. Prod.* 509.—Austral.
Eckersleyana, *F. Muell. Fragm.* x. 17.—Austral.
empetrifolia, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 226.—Austral.
euphrasioides, *Benth. in Mitch. Journ. Trop. Austral.* 360.—Austral.
eurybioides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 48.—Austral.
Grylloana, *F. Muell. Fragm.* x. 17.—Austral.
hirtula, *F. Muell. ex Benth. Fl. Austral.* v. 97.—Austral.
incana, *A. Cunn. ex Benth. Lab. Gen. et Sp.* 455.—Austral.
incisa, *R. Br. Prod.* 509.—Austral.
incisa, Sieber, ex Benth. Lab. Gen. et Sp. 451 = Sieberi.
lasianthos, *Labill. Nov. Holl. Pl.* ii. 18. t. 157.—Austral.
Leichhardtii, *Benth. Fl. Austral.* v. 106.—Austral.
linearis, *R. Br. Prod.* 509.—Austral.
linearis, Sieber, ex Benth. Lab. Gen. et Sp. 460 = Westringia longifolia.
lithospermoides, *F. Muell. Fragm.* vi. 107.—Austral.
marifolia, *R. Br. Prod.* 509.—Austral.
melissifolia, *F. Muell. Fragm.* i. 19.—Austral.
microphylla, *A. Cunn. ex Benth. Lab. Gen. et Sp.* 454.—Austral.
nivea, *A. Cunn. l. c.* 452.—Austral.
odoratissima, *Benth. in Mitch. Journ. Trop. Austral.* 291.—Austral.
ovalifolia, *R. Br. Prod.* 509.—Austral.
phylicifolia, *F. Muell. Fragm.* i. 19.—Austral.
pimeloides, F. Muell. l. c. vi. 107 = saxicola.
prunelloides, *R. Br. Prod.* 508.—Austral.
retusa, R. Br. l. c. 509 = rotundifolia.
retusa, Sieber, ex Benth. Lab. Gen. et Sp. 455 = violacea.
rhombea, *R. Br. Prod.* 509.—Austral.
ringens, *Benth. in Mitch. Journ. Trop. Austral.* 363.—Austral.
rotundifolia, *R. Br. Prod.* 509.—Austral.
rugosa, *A. Cunn. ex Benth. Lab. Gen. et Sp.* 456.—Austral.
saxicola, *R. Br. Prod.* 509.—Austral.
Sieberi, *Benth. Lab. Gen. et Sp.* 451.—Austral.
spinosa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 48.—Austral.
staurophylla, *F. Muell. Fragm.* ix. 73.—Austral.
striatiflora, *F. Muell. in Linnaea*, xxv. (1852) 425.—Austral.
Sullivaniae, F. Muell. ex Benth. Fl. Austral. v. 104 = striatiflora.
thymifolia, A. Cunn. ex Benth. Lab. Gen. et Sp. 455 = violacea.
violacea, *R. Br. Prod.* 509.—Austral.
Walteri, *F. Muell. Fragm.* vii. 108.—Austral.
Wilkieana, *F. Muell. l. c.* viii. 230.—Austral.

PROSTEA, Cambess. in Mém. Mus. Par. xviii. (1829) 25. t. 1 C = **Pometia**, Forst. (Sapind.).
oblongifolia, Presl, Bot. Bemerk. 40 = Sapindus oblongifolius.
pinnata, Cambess. in Mém. Mus. Par. xviii. (1829) 25 = Pometia pinnata.

PROSTHECHEA, Knowles & Westc. Flor. Cab. ii. (1838) 111 = **Epidendrum**, Linn. (Orchid.).
glauca, Knowles & Westc. l. c. = E. glaucum.

PROSTHESIA, Blume, Bijdr. 866 (1826) = **Alsodeia**, Thou. (Violar.).
javanica, Blume, l. c. 867 = A. javanica.

PROTANTHERA, Rafin. in Med. Repos. N. York, II. iii. (1806) 422, nomen. *LILIACEAE*?
phalangioides, Rafin. l. c.—Am. bor.
tenue, Rafin. l. c.—Am. bor.

PROTEA, Linn. Gen. ed. II. 38 (1742); ed. V. 41 (1754) = **Leucadendron**, R. Br. (Proteac.).

PROTEA, Linn. Syst. ed. I (1735); Gen. ed. I. 22 (1737). *PROTEACEAE*, Benth. & Hook. f. iii. 169.
CHRYSODENDRON, Vaill. ex Meissn. in DC. Prod. xiv. 230 (1856-57).
ERODENDRUM, Salisb. Parad. Lond. t. 67 (1807).
LEPIDOCARPUS, Adans. Fam. ii. 284 (1763).
LEUCADENDRON, Linn. Syst. ed. II. 18 (1740).
PLEURANTHE, Salisb. in Knight, Prot. 49 (1810).
abietina, Buek, ex Meissn. in DC. Prod. xiv. 246 = acerosa.
abietina, Licht. ex Meissn. l. c. 248 = Spatalla pedunculata.
abietina, Poir. Encyc. Suppl. iv. 559 = Leucadendron aemulum.
abrotanifolia, Andr. Bot. Rep. t. 507 = Serruria phylicoides, odorata.
abrotanifolia, Salisb. Prod. 48 = Serruria Burmanni.
abyssinica, *Willd. Sp. Pl.* i. 522.—Abyss.
acaulis, *Thunb. Diss. Prot.* 40.—Afr. austr.
acerosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 95.—Afr. austr.
acicularis, Poir. Encyc. Suppl. iv. 559 = Petrophila acicularis.
acrocarpa, Poir. l. c. 570 = Serruria acrocarpa.
acufera, Cav. in Anal. Hist. Nat. i. (1799) 236 = Isopogon anethifolius.
acuminata, *Sims, Bot. Mag.* t. 1694.—Afr. austr.
acutifolia, Salisb. Parad. Lond. t. 2 = rosacea, *Thunb.*
adscendens, Steud. Nom. ed. II. ii. 399 = Serruria adscendens.
aemula, Poir. Encyc. Suppl. iv. 558, 573 = Serruria aemula.
Aitoni, Poir. l. c. 570 = Serruria Aitoni.
alba, Thunb. Diss. Prot. 31 = Leucadendron cinereum.
alopecuroides, Lam. Illustr. i. 240 = Nivenia Sceptrum.
alpina, *Knight, Prot.* 27.—Afr. austr.
amplexicaulis, *R. Br. in Trans. Linn. Soc.* x. (1810) 95.—Afr. austr.
anemonefolia, Salisb. Prod. 48 = Isopogon anemonefolius.
anethifolia, Salisb. l. c. = Isopogon anethifolius.
angolensis, *Welw. Apont.* 586.—Afr. trop.
angustata, *R. Br. in Trans. Linn. Soc.* x. (1810) 90.—Afr. austr.
angustata, Drège, ex Meissn. in DC. Prod. xiv. 245 = acaulis.
angustata, Poir. Encyc. Suppl. iv. 556 = Leucadendron angustatum.
angustifolia, Salisb. Prod. 49 = P. Scolymus.
apifolia, Sm. ex Meissn. in DC. Prod. xiv. 279 = Isopogon anemonefolius.
arborea, *Schult. Mant.* iii. 266.—Afr. austr.
arborescens, *Hort. ex Steud. Nom.* ed. II. ii. 399.—Afr. austr.
arcuata, Lam. Illustr. i. 234 = Leucadendron plumosum.
arenaria, Buek, ex Meissn. in DC. Prod. xiv. 245 = acaulis.
arenaria, Poir. Encyc. Suppl. iv. 571 = Serruria arenaria.
argentea, Linn. Sp. Pl. 94 = Leucadendron argenteum.
argentiflora, Andr. Bot. Rep. t. 447 = Serruria triternata.
ascendens, Lam. Illustr. i. 239 = Serruria adscendens.
asplenifolia, *Link, Enum. Hort. Berol.* i. 113.—Afr. austr.
asterifolia, *Knight, Prot.* 30.—Afr. austr.
attenuata, Poir. Encyc. Suppl. iv. 566 = Leucospermum attenuatum.
aulacea, Thunb. Diss. Prot. 31 = Aulax cneorifolia.
Aulax, Hibbert, ex Meissn. in DC. Prod. xiv. 242 = lorea.
auriculata, *Tausch, in Flora*, xxv. (1842) 1. 285.—Afr. austr.
australis, Poir. Encyc. Suppl. iv. 579 = Spatalla propinqua.
Banksii, Klotzsch, ex Meissn. in DC. Prod. xiv. 236 = Buekiana.
barbata, Lam. Illustr. i. 236 = speciosa.
barbigera, *Meissn. in DC. Prod.* x. 233.—Afr. austr.
bracteata, Thunb. Diss. Prot. 27 = Aulax pinifolia.
bracteolaris, Poir. Encyc. Suppl. iv. 578 = Spatalla parilis.
brassicaefolia, *Hort. ex Steud. Nom.* ed. II. ii. 399.—Afr. austr.
brevifolia, Poir. Encyc. Suppl. iv. 579 = Spatalla brevifolia.
Brownii, Poir. l. c. 570 = Serruria diffusa.
bruniades, Linn. f. Suppl. 117 = Leucadendron corymbosum.

PROTEA:—
bruniaefolia, *Knight, Prot.* 32.—Afr. austr.
Buekiana, *Meissn. in DC. Prod.* xiv. 236.—Afr. austr.
buxifolia, Poir. Encyc. Suppl. iv. 566 = Leucospermum buxifolium.
caespitosa, Andr. Bot. Rep. t. 526 = turbiniflora.
caffra, *Meissn. in DC. Prod.* xiv. 237.—Afr. austr.
callosa, Wendl. ex Steud. Nom. ed. II. ii. 34, 399 = Leucadendron callosum.
calocephala, *Meissn. in DC. Prod.* xiv. 239.—Afr. austr.
calycina, *Schneev. Ic. Pl. Rar.* t. 48.—Afr. austr.
canaliculata, *Andr. Bot. Rep.* t. 437.—Afr. austr.
canaliculata, Herb. Linn. Soc. Lond. ex Meissn. in DC. Prod. xiv. 242 = repens.
candicans, Andr. Bot. Rep. t. 294 = Leucospermum tomentosum.
candicans, Thunb. Prod. Pl. Cap. 186 = Serruria albicans.
capitata, Poir. Encyc. Suppl. iv. 575 = Nivenia capitata.
capitulata, Poir. l. c. 568 = Mimetes capitulata.
carinata, *Hort. ex Schult. Mant.* iii. 266.—Afr. austr.
carlescens, E. Mey. ex Meissn. in DC. Prod. xiv. 238 = punctata.
carnosa, *Hort. ex Roem. & Schult. Syst.* iii. 356 (Quid?).—Afr. austr.
cartilaginea, Poir. Encyc. Suppl. iv. 557 = Leucadendron cartilagineum.
caudata, Thunb. Diss. Prot. 26 = Spatalla caudata, Thunbergii.
caulescens, *E. Mey. ex Meissn. in DC. Prod.* xiv. 244.—Afr. austr.
cerifera, *Hort. ex Roem. & Schult. Syst.* iii. 356 (Quid?).—Afr. austr.
chamaelea, Lam. Illustr. i. 237 = Leucadendron decurrens.
ciliaris, Hort. ex Roem. & Schult. Syst. iii. 356 = Leucadendron marginatum.
ciliata, Breit. Hort. Breit. 380; Schult. Mant. iii. 266 = Leucadendron marginatum.
ciliata, Desf. Tabl. ed. II. 45 = Leucadendron grandiflorum.
ciliata, Poir. Encyc. Suppl. iv. 572 = Serruria ciliata.
cinerea, Ait. Hort. Kew. ed. I. i. 127 = Leucadendron cinereum.
cinerea, Willd. ex Meissn. in DC. Prod. xiv. 215 = Leucadendron imbricatum.
cinerea, Willd. Sp. Pl. i. 521 = Leucospermum tortum.
cneorifolia, Salisb. Prod. 49 = Aulax cneorifolia.
coarctata, Thunb. Fl. Cap. 122 = Serruria scariosa.
coccinea, *R. Br. in Trans. Linn. Soc.* x. (1810) 77.—Afr. austr.
comosa, Thunb. Diss. Prot. 28 = Leucadendron comosum.
compacta, *R. Br. in Trans. Linn. Soc.* x. (1810) 76.—Afr. austr.
compar, Poir. Encyc. Suppl. iv. 574 = Serruria compar.
concava, Lam. Illustr. i. 234 = Nivenia concava.
concinna, Poir. Encyc. Suppl. iv. 556 = Leucadendron concinnum.
concinna, Salisb. Prod. 50 = Leucadendron uliginosum.
concolor, *Hort. ex Steud. Nom.* ed. II. ii. 399.—Afr. austr.
congesta, Poir. Encyc. Suppl. iv. 573 = Serruria congesta.
conica, Lam. Illustr. i. 237 = Leucadendron adscendens, strictum.
conifera, Andr. Bot. Rep. t. 541 = Leucadendron strictum.
conifera, Linn. Mant. ii. 193 = Leucadendron adscendens, salignum.
conifera, Thunb. Diss. Prot. 37 = Leucadendron coniferum.
conocarpa, Poir. Encyc. v. 656 = Leucospermum ellipticum.
conocarpa, Thunb. Diss. Prot. 22 = Leucospermum conocarpum, ellipticum.
cordata, *Thunb. l. c.* 45.—Afr. austr.
cordifolia, Sims, Bot. Mag. t. 649 = cordata.
coriacea, Buek, ex Meissn. in DC. Prod. xiv. 238 = punctata.
coriacea, Licht. ex Roem. & Schult. Syst. iii. 384 = Mimetes coriacea.
coriacea, Willd. ex Meissn. in DC. Prod. xiv. 220 = Leucadendron venosum.
coronata, Andr. Bot. Rep. t. 469 = formosa.

PROTEA:—
coronata, Curt. ex Steud. Nom. ed. I. 658 = lorea.
coronata, Lam. Illustr. i. 236 = speciosa.
corymbosa, Thunb. Diss. Prot. 29 = Leucadendron corymbosum.
crassifolia, Poir. Encyc. Suppl. iv. 557 = Leucadendron crassifolium.
crassulaefolia, *Knight, Prot.* 28.—Afr. austr.
criniflora, Linn. f. Suppl. 117 = Leucospermum oleaefolium.
crinita, Jacq. ex Spreng. Syst. i. 465 = Leucospermum oleaefolium.
crinita, Thunb. Diss. Prot. 21 = Leucospermum crinitum.
cristata, *Lam. Illustr.* i. 235.—Afr. austr.
crithmifolia, Poir. Encyc. Suppl. iv. 574 = Nivenia crithmifolia.
cucullata, Lam. Illustr. i. 239 = Mimetes Hartogii.
cucullata, Linn. Mant. ii. 189 = Mimetes lyrigera.
cyanoides, Linn. l. c. 188 = Serruria cyanoides.
cyanoides, Thunb. Diss. Prot. 15 = Serruria nitida.
cygnea, Poir. Encyc. Suppl. iv. 569 = Serruria cygnea.
cynaroides, *Linn. Mant.* ii. 190.—Afr. austr.
daphnoides, Thunb. Fl. Cap. i. 490 = Leucadendron daphnoides.
decipiens, Poir. Encyc. Suppl. iv. 573 = Serruria decipiens.
declinata, *Hort. ex Meissn. in DC. Prod.* xiv. 247.—Afr. austr.
decora, Salisb. Prod. 50 = Leucadendron grandiflorum.
decumbens, Andr. Bot. Rep. t. 349 = Serruria Niveni.
decumbens, Thunb. Diss. Prot. 14 = Serruria decumbens.
decumbens, Willd. ex Meissn. in DC. Prod. xiv. 284 = Serruria glaberrima.
densa, *Willd. Enum. Hort. Berol. Suppl.* 7, nomen.—Afr. austr.
dichotoma, Cav. in Anal. Hist. Nat. i. (1799) 239 = Petrophila pulchella.
dichotoma, Lam. Illustr. i. 235 = Mimetes dichotoma, divaricata.
divaricata, Linn. Mant. ii. 194 = Isopogon anethifolius.
divaricata, Willd. ex Meissn. in DC. Prod. xiv. 265 = Mimetes thymelaeoides.
diversifolia, Poir. Encyc. Suppl. iv. 559, 577 = Sorocephalus diversifolius.
diversifolia, Willd. Enum. Hort. Berol. 139 = Leucadendron salignum.
dodonaeifolia, Buek, ex Meissn. in DC. Prod. xiv. 239 = longifolia.
echinulata, *Meissn. l. c.* 242.—Afr. austr.
effusa, *E. Mey. ex Meissn. l. c.* 240.—Afr. austr.
elevata, Poir. Encyc. Suppl. iv. 570 = Serruria elevata.
elliptica, Thunb. Diss. Prot. 22 = Leucospermum ellipticum.
elongata, *R. Br. in Trans. Linn. Soc.* x. (1810) 90.—Afr. austr.
erecta, Thunb. Fl. Cap. i. 454 = Serruria hirsuta.
ericaefolia, Willd. ex Meissn. in DC. Prod. xiv. 218 = Leucadendron corymbosum.
ericifolia, Poir. Encyc. Suppl. iv. 557 = Leucadendron ericifolium.
ericoides, *Hort. ex Schult. Mant.* iii. 266.—Afr. austr.
eriocephala, Licht. ex Roem. & Schult. Syst. ii. 379 = Serruria phylicoides, squarrosa.
erosa, Licht. ex Roem. & Schult. l. c. 360 = Leucospermum grandiflorum.
erosa, Willd. ex Meissn. in DC. Prod. xiv. 260 = Leucospermum crinitum.
falcata, A. Cunn. ex Meissn. l. c. 267 = Petrophila linearis.
falcifolia, *Knight, Prot.* 29.—Afr. austr.
fastigiata, Poir. Encyc. Suppl. iv. 559 = Petrophila fastigiata.
feniculacea, Poir. l. c. 572 = Serruria foeniculacea.
filifolia, Poir. l. c. 559 = Petrophila acicularis.
flagellaris, Poir. l. c. 573 = Serruria flagellaris.
flavescens, Willd. Enum. Hort. Berol. Suppl. 7 = Leucadendron flavescens.
florida, Thunb. Diss. Prot. 15 = Serruria florida.
foeniculacea, Auct. vide *feniculacea*.
formosa, Andr. Bot. Rep. t. 17 = Leucospermum medium.
formosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 79.—Afr. austr.
fucifolia, Salisb. Prod. 48 = Petrophila pulchella.
fulva, *Tausch, in Flora*, xxv. (1842) 285.—Afr. austr.

PROTEA:—
furcellata, Poir. Encyc. Suppl. iv. 571 = Serruria furcellata.
furcellata, Sm. ex Meissn. in DC. Prod. xiv. 277 = Isopogon anethifolius.
fusca, Linn. Sp. Pl. 95 = Leucadendron Levisanus.
fusciflora, Jacq. Hort. Schoenb. i. 11 = Leucadendron fusciflorum.
Gaguedi, J. F. Gmel. Syst. 225 = abyssinica.
glaberrima, Poir. Encyc. Suppl. iv. 569 = Serruria glaberrima.
glabra, *Thunb. Diss. Prot.* 42.—Afr. austr.
glauca, *Brouss. ex Roem. & Schult. Syst.* iii. 356.—Afr. austr.
glauca, Buek, ex Meissn. in DC. Prod. xiv. 233 = neriifolia.
glauca, Drège, ex Meissn. l. c. = melaleuca.
glaucophylla, Salisb. Parad. Lond. t. 11 = acaulis.
globosa, Andr. Bot. Rep. t. 307 = Leucadendron concolor.
Globularia, Lam. Illustr. i. 236 = Leucadendron Globularia.
globulariaefolia, *Knight, Prot.* 30.—Afr. austr.
glomerata, Andr. Bot. Rep. t. 264 = Serruria pedunculata.
glomerata, Linn. Mant. ii. 187 = Serruria glomerata.
glomerata, Thib. ex Meissn. in DC. Prod. xiv. 292 = Serruria phylicoides.
glomerata, Thunb. Diss. Prot. 18 = Serruria elongata.
glomerata, Willd. ex Meissn. in DC. Prod. xiv. 293 = ciliata.
glomerata, Willd. ex Spreng. Syst. i. 468 = Serruria adscendens.
glutinosa, *Knight, Prot.* 27.—Afr. austr.
gnidiaefolia, *Knight, l. c.* 31.—Afr. austr.
graminea, *Hort. ex Link, Enum. Hort. Berol.* i. 113.—Afr. austr.
grandiceps, Tratt. Thesaur. Bot. 5, in syn. = speciosa.
grandiflora, *Thunb. Diss. Prot.* 41.—Afr. austr.
grandis, *Hort. ex Meissn. in DC. Prod.* xiv. 247, nomen.—Afr. austr.
Gustaviana, Poir. Encyc. v. 663 = Nivenia parvifolia.
helvola, Licht. ex Roem. & Schult. Syst. iii. 379 = Serruria cygnea.
helvola, Willd. ex Meissn. in DC. Prod. xiv. 297 = Serruria elongata.
heterophylla, Thunb. Diss. Prot. 24 = Leucospermum diffusum.
Hibbertii, Poir. Encyc. Suppl. iv. 568 = Mimetes splendida.
hirsuta, Poir. l. c. 572 = Serruria hirsuta.
hirsuta, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 12 = Leucadendron hirsutum.
hirsuta, Willd. Enum. Hort. Berol. 139 = Leucadendron Levisanus.
hirta, Hort. ex Meissn. in DC. Prod. xiv. 216 = Leucadendron tortum.
hirta, *Klotzsch, ex Krauss, in Flora*, xxviii. (1845) 76.—Afr. austr.
hirta, Linn. Mant. ii. 188 = Mimetes hirta.
hirtella, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
humiflora, Andr. Bot. Rep. t. 532 = seq.
humiflorens, Willd. Enum. Hort. Berol. Suppl. 7 = humilis.
humifusa, *Hort. ex Meissn. in DC. Prod.* xiv. 246.—Afr. austr.
humilis, *R. Br. in Trans. Linn. Soc.* x. (1810) 95.—Afr. austr.
hypophylla, Thunb. Diss. Prot. 23 = Leucospermum Hypophyllum.
Hypophyllocarpodendrum, Linn. Mant. ii. 191 = Leucospermum Hypophyllum.
imberbis, Poir. Encyc. Suppl. iv. 576 = Sorocephalus imberbis.
imbricata, Poir. l. c. 557 = Leucadendron imbricatum.
imbricata, Thunb. Diss. Prot. 38 = Sorocephalus imbricatus.
imbricata, Wendl. Hort. Herrenh. 4. t. 14 = Leucadendron buxifolium.
incana, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
incompta, *R. Br. in Trans. Linn. Soc.* x. (1810) 83.—Afr. austr.
incorrupta, Krauss, in Flora, xxviii. (1845) 75, sphalm. = praec.
incurva, Andr. Bot. Rep. t. 429 = Leucadendron aemulum.

PROTEA:—
incurva, Thunb. Diss. Prot. 26 = Spatalla procera.
inflexa, Willd. Enum. Hort. Berol. Suppl. 7 = Leucadendron inflexum.
intermedia, Poir. Encyc. Suppl. iv. 575 = Nivenia media.
involucrata, Licht. ex Spreng. Syst. i. 457 = Leucospermum involucratum.
involucrata, Willd. ex Meissn. in DC. Prod. xiv. 222 = Leucadendron adscendens.
lacticolor, Salisb. Parad. Lond. t. 27 = longiflora.
laevis, *R. Br. in Trans. Linn. Soc.* x. (1810) 91.—Afr. austr.
laevis, Thunb. Fl. Cap. ed. Schult. 133 = Nivenia Lagopus.
Lagopus, Andr. Bot. Rep. t. 243 = Nivenia crithmifolia.
Lagopus, Thunb. Diss. Prot. 19 = Nivenia Lagopus.
Lagopus, Willd. Sp. Pl. i. 510 = Nivenia spicata.
lanata, Thunb. Diss. Prot. 30 = Sorocephalus lanatus.
lanceolata, DC. ex Meissn. in DC. Prod. xiv. 212 = Aulax cneorifolia.
lanceolata, Hort. ex Meissn. l. c. 248, nomen = seq.?
lanceolata, *E. Mey. ex Meissn. l. c.* 240.—Afr. austr.
latericolor, Meissn. l. c. 248, sphalm. = *lacticolor*.
latifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 75.—Afr. austr.
Laureola, Lam. Illustr. i. 234 = Leucadendron decorum.
laurifolia, Buek, ex Meissn. in DC. Prod. xiv. 236 = grandiflora.
laxa, Poir. Encyc. Suppl. iv. 578 = Spatalla gracilis.
lenta, Salisb. Prod. 50 = Leucadendron ovale?
Lepidocarpodendron, Linn. Mant. ii. 190 = cristata, longifolia, speciosa.
Lepidocarpon, *R. Br. in Trans. Linn. Soc.* x. (1810) 80.—Afr. austr.
Lepidocarpon, Sims, Bot. Mag. t. 674 = melaleuce.
leucantha, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
Levisana, Meissn. l. c. 214 = Leucadendron imbricatum.
Levisanus, Linn. Mant. ii. 194 = Leucadendron Levisanus, imbricatum.
ligulaefolia, Sweet, Hort. Brit. ed. I. 346 = longifolia.
linearis, Houtt. Handleid. iv. 116. t. 19. f. 2 = Leucadendron angustatum.
linearis, Thunb. Diss. Prot. 33 = Leucospermum lineare.
linguiformis, *Hort. ex Meissn. in DC. Prod.* xiv. 248, nomen.—Afr. austr.
linifolia, Jacq. Hort. Schoenb. i. 11 = Leucodendron linifolium.
longicaulis, *Knight, Prot.* 29.—Afr. austr.
longiflora, *Lam. Illustr.* i. 234.—Afr. austr.
longifolia, *Andr. Bot. Rep.* tt. 132, 133, 144.—Afr. austr.
longifolia, Salisb. Parad. Lond. t. 37 = laevis.
loranthifolia, *Knight, Prot.* 28.—Afr. austr.
lorea, *R. Br. in Trans. Linn. Soc.* x. (1810) 93.—Afr. austr.
lorea, Drège, ex Meissn. in DC. Prod. xiv. 242 = tenuifolia.
macrocephala, *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 13.—Afr. austr.
macrophylla, *R. Br. in Trans. Linn. Soc.* x. (1810) 78.—Afr. austr.
macrophylla, Buek, ex Meissn. in DC. Prod. xiv. 233 = barbigera.
madiensis, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 143.—Afr. trop.
magnifica, *Andr. Bot. Rep.* t. 438.—Afr. austr.
magnoliaefola, Buek, ex Meissn. in DC. Prod. xiv. 244 = tenax.
marginata, Lam. Illustr. i. 235 = grandiflora.
marginata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 15 = melaleuca.
marginata, Willd. Enum. Hort. Berol. Suppl. 7 = Leucadendron marginatum.
marginata, Willd. ex Meissn. in DC. Prod. xiv. 218 = Leucadendron decorum.
Massonii, Poir. Encyc. Suppl. iv. 568 = Mimetes Massoni.
media, Poir. l. c. 566 = Leucospermum medium.
melaleuca, *R. Br. in Trans. Linn. Soc.* x. (1810) 79.—Afr. austr.
mellifera, *Thunb. Diss. Prot.* 34.—Afr. austr.

PROTEA:—
micans, *Welw. Apont.* 586.—Afr. trop.
micrantha, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
mollis, *Hort. l. c.*—Afr. austr.
mollis, Poir. Encyc. Suppl. iv. 567, 577 = Leucospermum molle.
mollissima, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
mollissima, Poir. Encyc. Suppl. iv. 575 = Nivenia mollisima.
montana, *E. Mey. ex Meissn. in DC. Prod.* xiv. 240.—Afr. austr.
mucronata, Hort. l. c. = mucronifolia.
mucronata, Nois. ex Schult. Mant. iii. 266, nomen = praec.?
mucronifolia, *Salisb. Parad. Lond.* t. 24.—Afr. austr.
multifida, Nois. ex Schult. Mant. iii. 266 (Quid?).—Afr. austr.
Mundi, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* vi. (1838) 113.—Afr. austr.
mutica, Poir. Encyc. Suppl. iv. 455 = Leucadendron spathulatum.
myrtifolia, Thunb. Diss. Prot. 41 = Mimetes myrtifolia.
nana, Lam. Illustr. i. 233 = acaulis.
nana, Thunb. Diss. Prot. 30 = rosacea.
nectarina, Schrad. Sert. Hannov. 5. t. 21 = Lambertia formosa.
neriifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 81.—Afr. austr.
nigra, *Regel, Cat. Pl. Hort. Aksakov.* 116 (Quid?).—Afr. austr.
nigrita, DC. ex Meissn. in DC. Prod. xiv. 233 = melaleuca.
nitens, Thunb. Fl. Cap. i. 514 = Mimetes nitens.
nitida, Mill. Gard. Dict. ed. VIII. n. 3 = grandiflora.
nivea, Poir. Encyc. Suppl. iv. 578 = Spatalla nivea.
Niveni, Poir. l. c. 572 = Serruria Niveni.
nutans, Poir. ex Steud. Nom. ed. ii. 400 = Leucospermum nutans.
obliqua, Hort. Bollw. ex Meissn. in DC. Prod. xiv. 235 = longiflora.
obliqua, Lam. Encyc. v. 642 = Leucadendron concolor.
obliqua, Linn. f. Suppl. 117 = Leucadendron squarrosum.
obliqua, Thunb. Diss. Prot. 35 = Leucadendron plumosum.
obliqua, Willd. ex Meissn. in DC. Prod. xiv. 222 = Serruria adscendens.
oblongifolia, *Presl, Bot. Bemerk.* 40.—Afr. austr.
oborata, Roem. & Schult. Syst. iii. 378, in obs., sphalm. = seq.
obovata, Andr. ex Meissn. in DC. Prod. xiv. 25 = Serruria abrotanifolia.
obtusa, Sweet, Hort. Brit. ed. I. 346 = speciosa.
obtusata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 15 = Leucadendron obtusatum.
obtusifolia, *Buek, ex Meissn. in DC. Prod.* xiv. 235.—Afr. austr.
ochroleuca, Sm. Exot. Bot. ii. 43 = longiflora.
odorata, Thunb. Prod. Pl. Cap. 187 = Leucadendron odoratum.
odoratissima, Mass. ex R. Br. in Trans. Linn. Soc. x. (1810) 87 = mucronifolia.
ovalis, Buek, ex Meissn. in DC. Prod. xiv. 235 = Mundi.
ovata, *Thunb. in Mém. Acad. Pétersb.* iv. (1813–14) 548. t. 17.—Afr. austr.
pallens, Linn. Mant. ii. 193 = Leucadendron adscendens.
pallens, Thunb. Diss. Prot. 36 = Leucadendron decurrens.
pallens, Willd. ex Meissn. in DC. Prod. xiv. 222 = Leucadendron virgatum.
pallida, Salisb. Prod. 49 = Leucadendron decurrens.
parilis, Buek, Ind. ad DC. Prod. iv. 317 = Serruria phylicoides.
parviflora, Linn. Mant. ii. 195 = Leucadendron plumosum.
Passerina, Desf. Tabl. ed. II. 45, nomen = seq.?
Passerina, Hort. ex Meissn. in DC. Prod. xiv. 216, 251 = Leucadendron tortum.
patens, *R. Br. in Trans. Linn. Soc.* x. (1810) 82.—Afr. austr.
patens, Poir. Encyc. Suppl. iv. 567 = Leucospermum puberum.
patula, Thunb. Diss. Prot. 16 = Serruria glomerata.
patula, Willd. ex Meissn. in DC. Prod. xiv. 296 = Serruria Burmanni.
pauciflora, Poir. Encyc. Suppl. iv. 568 = Mimetes pauciflora.

PROTEA :—
Paulina, *Endl. Gen. Suppl.* iv. II. 77, nomen.—Nubia.
pedunculata, Lam. Illustr. i. 240 = Serruria pedunculata.
pedunculata, Poir. Encyc. Suppl. iv. 559, 578 = Spatalla pedunculata.
pendula, *R. Br. in Trans. Linn. Soc.* x. (1810) 87.—Afr. austr.
penicillata, E. Mey. ex Meissn. in DC. Prod. xiv. 235 = Mundi.
petiolata, Buek, ex Meissn. l. c. 245 = cynaroides.
phylicoides, Poir. Encyc. v. 659 = Serruria villosa.
phylicoides, F. W. Schmidt, ex Tausch, in Flora, xiii. (1830) 555 = Serruria congesta.
phylicoides, Thunb. Diss. Prot. 19 = Serruria hirsuta.
phylicoides, Willd. Sp. Pl. i. 510 = Serruria arenaria.
Pinaster, Hort. ex Steud. Nom. ed. II. ii. 400 = pinastrifolia?
pinastrifolia, *Hort. Loud. ex Dum.-Cours. Bot. Cult.* ed. II. i. 591.—Afr. austr.
pinifolia, Linn. ex Meissn. in DC. Prod. xiv. 212 = Aulax cneorifolia.
pinifolia, Linn. Mant. ii. 187 = Aulax pinifolia.
pinnata, Andr. Bot. Rep. t. 512 = Serruria pinnata.
pinnata, Cav. ex Meissn. in DC. Prod. xiv. 248 (Quid?).—Afr. austr.
plumigera, Thunb. Fl. Cap. 121 = Serruria simplicifolia.
plumosa, Ait. Hort. Kew. ed. I. i. 127 = Leucadendron plumosum.
polygaloides, Willd. ex Meissn. in DC. Prod. xiv. 215 = Leucadendron imbricatum.
polygaloides, Willd. Hort. Berol. ii. 100 = canaliculata, Leucadendron imbricatum.
polysperma, Poir. Encyc. Suppl. iv. 556 = Leucadendron platyspermum.
polystachia, Poir. l. c. 578 = Spatalla polystachya.
procumbens. Linn. f. Suppl. 116 = Serruria decumbens.
prolifera, Thunb. Diss. Prot. 29 = Spatalla prolifera.
prostrata, Thunb. Prod. Pl. Cap. 27 = Leucadendron prostratum.
pubera, Linn. Mant. ii. 192 = Leucospermum puberum.
pubescens, Poir. Encyc. Suppl. iv. 559 = Leucadendron pubescens.
pulchella, *Andr. Bot. Rep.* t. 270.—Afr. austr.
pulchella, Schrad. Sert. Hannov. 15 = Petrophila pulchella.
punctata, *Meissn. in DC. Prod.* xiv. 238.—Afr. austr.
purpurea, Linn. Mant. ii. 195 = Mimetes purpurea.
pyramidalis, Poir. Encyc. Suppl. iv. 578 = Spatalla parilis.
pyramidalis, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 11 = Leucadendron pyramidale.
pyrifolia, Buek, ex Meissn. in DC. Prod. xiv. 236 = Buekiana.
racemosa, Licht. ex Spreng. Syst. i. 471 = Spatalla gracilis.
racemosa, Linn. Mant. ii. 187 = Spatalla bracteata.
racemosa, Thunb. Diss. Prot. 25 = Spatalla ramulosa.
radiata, Andr. Bot. Rep. t. 646 = latifolia.
radiata, Bonpl. ex Reichb. Fl. Exot. iv. 35, cum ic. = longiflora.
Radula, *Hort. ex Link, Enum. Hort. Berol.* i. 113.—Afr. austr.
radulifolia, Donn, Cat. Hort. Cantab. ed. IV. 25 = P. Radula.
rangiferina, Hort. ex Schult. Mant. iii. 263 = coccinea.
rangiferina, Lee, ex Meissn. in DC. Prod. xiv. 277 = Isopogon anethifolius.
recondita, *Buek, ex Meissn. l. c.* 237.—Afr. austr.
reflexa, *Hort. ex Meissn. l. c.* 248 (Quid?).—Afr. austr.
repens, Andr. Bot. Rep. t. 453 = amplexicaulis.
repens, Linn. Mant. ii. 189 = mellifera.
repens, *Thunb. Diss. Prot.* 34.—Afr. austr.
reticulata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 13 = Leucadendron reticulatum.
retroflexa, *F. G. Dietr. Vollst. Lexik. Gaertn.* vii. 556.—Afr. austr.
retusa, Poir. Encyc. Suppl. iv. 555 = Leucadendron retusum.
revoluta, *R. Br. in Trans. Linn. Soc.* x. (1810) 90.—Afr. austr.
revoluta, Buek, ex Meissn. in DC. Prod. xiv. 242 = tenuifolia.
rigida, Poir. Encyc. Suppl. iv. 559 = Petrophila rigida.
rosacea, *Linn. Mant.* ii. 189.—Afr. austr.

PROTEA :—
Rouppelliae, *Meissn. in DC. Prod.* xiv. 237.—Afr. austr.
Roxburghii, Poir. Encyc. Suppl. iv. 574 = Serruria Roxburghii.
rugosa, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 11 = Leucadendron rugosum.
rupicola, *Mund, ex Meissn. in DC. Prod.* xiv. 236.—Afr. austr.
saligna, Andr. Bot. Rep. t. 572 = Leucadendron floridum.
saligna, Linn. Mant. ii. 194 = Leucadendron adscendens, salignum.
saligna, Thunb. Diss. Prot. 39 = Leucadendron uliginosum.
salsoloides, Poir. Encyc. Suppl. iv. 576 = Sorocephalus salsoloides.
salsoloides, Thunb. ex Meissn. in DC. Prod. xiv. 266 = Mimetes purpurea.
scabra, *R. Br. in Trans. Linn. Soc.* x. (1810) 91.—Afr. austr.
scabra, Poir. Encyc. Suppl. iv. 559 = Leucadendron scabrum.
scabrida, *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 14.—Aft. austr.
Sceptrum, Lam. Illustr. i. 241 = Nivenia parvifolia.
Sceptrum, Thunb. Diss. Prot. 21 = Nivenia Sceptrum.
Sceptrum-Gustavianum, Sparm. in Vet. Akad. Handl. Stockh. (1777) 55. t. 1 = Nivenia Sceptrum.
scolopendrina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = seq.
Scolopendrium, *R. Br. in Trans. Linn. Soc.* x. (1810) 94.—Afr. austr.
scolymocephala, Reichard, Syst. Pl. i. 271 = P. Scolymus.
Scolymus, *Thunb. Diss. Prot.* 33.—Afr. austr.
scoparia, Poir. Encyc. Suppl. iv. 571 = Serruria scoparia.
sericea, Linn. f. Suppl. 118 = Leucadendron sericeum.
sericea, Thunb. Diss. Prot. 39 = Leucadendron salignum.
sericifolia, Poir. Encyc. Suppl. iv. 578 = Spatalla sericea.
serraria, Thunb. Diss. Prot. 17 = Serruria Burmanni.
serrarioides, Soland. ex Meissn. in DC. Prod. xiv. 291 = Serruria hirsuta.
Serruria, Linn. Mant. ii. 188 = Serruria Burmanni.
sessile, Poir. Encyc. Suppl. iv. 557 = Leucadendron sessile.
setacea, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
setacea, Poir. Encyc. Suppl. iv. 576 = Sorocephalus setaceus.
simplicifolia, Poir. l. c. 570 = Serruria simplicifolia.
spatalloides, Poir. l. c. 577 = Sorocephalus spatalloides.
spathulata, Poir. l. c. 567 = Leucospermum spathulatum.
spathulata, Thunb. Diss. Prot. 44 = Nivenia spathulata.
spathulata, Willd. ex Spreng. Syst. i. 469 = Nivenia parvifolia.
spatulaefolia, *Knight, Prot.* 31.—Afr. austr.
speciosa, *Linn. Mant.* ii. 191; *Sims, Bot. Mag.* t. 1183.—Afr. austr.
speciosa, Thunb. Diss. Prot. 42 = cristata.
spectabilis, Licht. ex Spreng. Syst. i. 461 = latifolia.
spectabilis, Willd. ex Meissn. in DC. Prod. xiv. 238 = compacta.
sphaerocephala, Houtt. Handleid. iv. 99. t. 19. f. 1 = Serruria hirsuta, pedunculata, scariosa.
sphaerocephala, Linn. Mant. ii. 188 = Serruria phylicoides.
sphaerocephala, Thib. ex Meissn. in DC. Prod. xiv. 293 = Serruria ciliata.
sphaerocephala, Thunb. Diss. Prot. 16 = Serruria rubricaulis.
sphaerocephala, Willd. ex Spreng. Syst. i. 467 = Serruria arenaria.
spicata, Andr. Bot. Rep. t. 234 = Nivenia media.
spicata, Linn. Mant. ii. 187 = Nivenia spicata.
spicata, Willd. ex Meissn. in DC. Prod. xiv. 302 = Nivenia Lagopus.
spiralis, *Hort. ex Meissn. l. c.* 248.—Afr. austr.
squamata, Poir. Encyc. Suppl. iv. 560 = Petrophila squamata.
squarrosa, Knight, Prot. 127 = Leucodendron fusciflorum?
squarrosa, Poir. Encyc. Suppl. iv. 573 = Serruria squarrosa.
staticefolia, *Hort. ex Meissn. in DC. Prod.* xiv. 248—Afr. austr.

PROTEA :—
stellaris, Sims, Bot. Mag. t. 881 = Leucadendron fusciflorum.
Stilbe, Poir. Encyc. Suppl. iv. 571 = Serruria Stilbe.
stricta, Don, ex Steud. Nom. ed. II. ii. 401 = Leucadendron adscendens.
strobilina, Don, ex Steud. l. c. = Leucadendron concolor.
strobilina, Linn. Mant. ii. 192 = Leucadendron squarrosum.
strobilina, Thunb. Diss. Prot. 38 = Leucadendron ovale.
Strobus, Meissn. in DC. Prod. xiv. 242 = repens.
subacaulis, *Hort. ex Meissn. l. c.* 248.—Afr. austr.
tenax, *R. Br. in Trans. Linn. Soc.* x. (1810) 88.—Afr. austr.
tenella, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
tenera, *Hort. ex Steud. Nom.* ed. II. ii. 401 (Quid?).—Afr. austr.
tenuifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 90.—Afr. austr.
tenuifolia, Poir. Encyc. Suppl. iv. 577 = Sorocephalus tenuifolius.
tenuifolia, Salisb. Prod. 49 = Leucadendron Levisanus.
tenuifolia, Soland. ex Meissn. in DC. Prod. xiv. 296 = Serruria Burmanni.
tenuifolia, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 9 = Thunbergii.
teretifolia, Andr. Bot. Rep. t. 161 = Leucadendron abietinum.
teretifolia, Poir. Encyc. Suppl. iv. 559 = Petrophila teretifolia.
thesiifolia, *Knight, Prot.* 28.—Afr. austr.
Thunbergii, *Steud. Nom.* ed. II. ii. 401.—Afr. austr.
thymelaeoides, Poir. Encyc. Suppl. iv. 568 = Mimetes thymelaeoides.
thymifolia, *Knight, Prot.* 31.—Afr. austr.
thyoides, Sm. ex Meissn. in DC. Prod. xiv. 225 = Leucadendron scabrum.
thyrsoides, Lam. Illustr. i. 240 = Serruria elongata.
tomentosa, Thunb. Diss. Prot. 24 = Leucospermum tomentosum.
tomentosa, Willd. ex Meissn. in DC. Prod. xiv. 258 = Leucospermum spathulatum.
torta, Thunb. Diss. Prot. 31 = Leucadendron tortum.
tortuosa, Salisb. Prod. 48 = Leucospermum conocarpum.
totta, Linn. Mant. ii. 191 = Leucospermum tottum.
tridactylites, Cav. in Anal. Hist. Nat. i. (1799) 236 = Isopogon anemonefolius.
tridentata, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
trifida, Poir. Encyc. Suppl. iv. 560 = Petrophila trifida.
triternata, Andr. Bot. Rep. t. 337 = Serruria triplicato-ternata.
triternata, Thib. ex Meissn. in DC. Prod. xiv. 295 = Serruria Roxburghii.
triternata, Thunb. Diss. Prot. 18 = Serruria triternata.
truncata, *Thunb. Fl. Cap.* i. 493.—Afr. austr.
turbinata, Steud. Nom. ed. I. 661 = longifolia.
turbiniflora, *R. Br. in Trans. Linn. Soc.* x. (1810) 93.—Afr. austr.
uliginosa, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
umbellata, Thunb. Diss. Prot. 32 = Aulax cneorifolia.
umbonalis, Sweet, Hort. Brit. ed. I. 346 = longifolia.
venosa, *Lam. Illustr.* i. 234.—Afr. austr.
venosa, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 10 = venulosa.
venulosa, *Steud. Nom.* ed. II. ii. 401.—Afr. austr.
verticillata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 12 = Sorocephalus verticillatus.
vestita, Lam. Illustr. i. 239 = Leucospermum ellipticum.
vidua, [Ker-Gawl.] Andr. Bot. Rep. Rec. Pl. 39 = longifolia.
villifera, *Lindl. in Bot. Reg.* t. 1023.—Afr. austr.
villosa, Hort. Bollw. ex Meissn. in DC. Prod. xiv. 233 = melaleuca.
villosa, Jacq. ex Meissn. in DC. l. c. 265 = Mimetes thymelaeoides.
villosa, Lam. Illustr. i. 240 = Serruria villosa.
villosa, Poir. Encyc. Suppl. iv. 566 = Leucospermum grandiflorum.
villosa, Thunb. Prod. Pl. Cap. 187 = Serruria pilosa.

PROTEA:—
villosa, Willd. ex Spreng. Syst. i. 464 = Leucospermum buxifolium.
villosiuscula, Banks, ex Poir. Encyc. Suppl. iv. 566 = Leucospermum grandiflorum.
virgata, Andr. Bot. Rep. t. 577 = acerosa.
virgata, Poir. Encyc. Suppl. iv. 556 = Leucadendron virgatum.
virgata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 10 = Leucadendron acutum.
viscosa, *Hort. ex Meissn. in DC. Prod.* xiv. 248.—Afr. austr.
Wendlandi, Poir. Encyc. Suppl. iv. 556 = Leucadendron buxifolium.
Woodwardii, *Endl. Gen. Suppl.* iv. II. 77.—Afr. austr.

PROTEINIA, Reichb. Nom. 205 (1841) = **Saponaria**, Linn. (Carophyll.).

PROTEINOPHALLUS, Hook. f. Bot. Mag. t. 6195 (1875) = **Amorphophallus**, Blume (Aroid.).
Rivieri, Hook. f. l. c. = A. Rivieri.

PROTEOPSIS, Mart. & Zucc. ex DC. Prod. v. 16 (1836); Sch. Bip. in Pollichia, xx–xxi. (1863) 378, 433. *COMPOSITAE*, Benth. & Hook. f. ii. 234.
argentea, *Mart. & Zucc. ex DC. Prod.* v. 16.—Bras.
glauca, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 20 = Vernonia pedunculata.
scapigera, Mart. l. c. 55 = Vernonia scapigera.
Sellovii, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 434.—Bras.

PROTERPIA, Rafin. Sylva Tellur. 80 (1838) (Bignoniacea).
obtusifolia, Rafin. l. c. = 'Spathodea obtusifolia *seu* Tabebuia uliginosa,' sec. A. DC.

PROTIONOPSIS, Blume, Mus. Bot. Lugd. Bat. i. 229 (1850) = **Commiphora**, Jacq. (Burser.).
caudata, Blume, l. c. = C. caudata.
pubescens, Blume, l. c. = C. pubescens.
Roxburghiana, Blume, l. c. = C. caudata.

PROTIUM, Burm. f. Fl. Ind. 88 (1768). *BURSERACEAE* [Benth. & Hook. f. i. 325].
DAMMARA, Gaertn. Fruct. ii. 100 (1791).
ICICA, Aubl. Pl. Gui. i. 337 (1775).
ICICOPSIS, Engl. in Mart. Fl. Bras. xii. II. 255 (1874).
MARIGNIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 351.
Almecago, *March. in Baill. Adansonia*, viii. (1867–68) 43.—Bras.
altissimum, *March. l. c.* 51.—Am. trop.
Aracouchili, *March. l. c.*—Bras.
aromaticum, Engl. in Mart. Fl. Bras. xii. II. 273 = heptaphyllum.
Beandon, *March. ex Engl. in DC. Monog. Phan.* iv. 92.—Madag.
Blanchetii, *Engl. l. c.* 79.—Bras.
brasiliense, *Engl. in Mart. Fl. Bras.* xii. II. 268.—Bras.
Carana, *March. in Baill. Adansonia*, viii. (1867–68) 51.—Bras.
Copal, *Engl. in DC. Monog. Phan.* iv. 83.—Mexic.
coriaceum, *Engl. in Mart. Fl. Bras.* xii. II. 281.—Am. trop.
crassifolium, *Engl. l. c.* 270.—Am. trop.
decandrum, *March. in Baill. Adansonia*, viii. (1867–68) 51.—Guian.
divaricatum, *Engl. in Mart. Fl. Bras.* xii. II. 279. t. 55.—Bras.
elegans, *Engl. l. c.* 273.—Bras.
ferrugineum, *Engl. in DC. Monog. Phan.* iv. 79.—Bras.
giganteum, *Engl. in Mart. Fl. Bras.* xii. II. 277.—Bras.
grandifolium, *Engl. l. c.* 269.—Am. trop.
guianense, *March. in Baill. Adansonia*, viii. (1867–68) 52.—Am. trop.
heptaphyllum, *March. in Kjoeb. Vidensk. Meddel.* (1873) 55.—Am. trop.
Hostmanni, *Engl. in Mart. Fl. Bras.* xii. II. 266.—Am. trop.
Icicariba, *March. in Baill. Adansonia*, viii. (1867–68) 52.—Bras.
insigne, *Engl. in DC. Monog. Phan.* iv. 77.—N. Granat.
javanicum, *Burm. f. Fl. Ind.* 88.—Java.
laxiflorum, *Engl. in Mart. Fl. Bras.* xii. II. 275.—Bras.
macrophyllum, *Engl. l. c.* 277.—Am. trop.
madagascariense, *Engl. in DC. Monog. Phan.* iv. 91.—Madag.
Martianum, *Engl. in Mart. Fl. Bras.* xii. II. 266.—Bras.
Melinonis, *Engl. in DC. Monog. Phan.* iv. 68.—Guian.
multiflorum, *Engl. in Mart. Fl. Bras.* xii. II. 273.—Bras.
nitidum, *Engl. l. c.* 271.—Bras.
obtusifolium, *March. in Baill. Adansonia*, viii. (1867–68) 62.—Ins. Maurit.
ovatum, *Engl. in Mart. Fl. Bras.* xii. II. 264.—Bras.
paniculatum, *Engl. l. c.* 270.—Am. trop.
pilosissimum, *Engl. in DC. Monog. Phan.* iv. 80.—Bras.
polybotryum, *Engl. in Mart. Fl. Bras.* xii. II. 278.—Guian.
pubescens, *Engl. l. c.* 265.—Bras.
reticulatum, *Endl. in DC. Monog. Phan.* iv. 79.—Bras.
Riedelianum, *Engl. in Mart. Fl. Bras.* xii. II. 279. t. 56.—Bras.
Sagotianum, *March. in Adansonia*, viii. (1867–68) 69. f. 1.—Guian.
Schomburgkianum, Engl. in Mart. Fl. Bras. xii. II. 276 = decandrum.
serratum, *Engl. in DC. Monog. Phan.* iv. 88.—Ind. or.
Spruceanum, *Engl. in Mart. Fl. Bras.* xii. II. 276.—Am. trop.
subserratum, *Engl. in DC. Monog. Phan.* iv. 89.—Bras.
Tacamahaca, March. in Baill. Adansonia, viii. (1867–68) 52 = heptaphyllum.
tenuifolium, *Engl. in DC. Monog. Phan.* iv. 76.—Peruv.
trifoliolatum, *Engl. in Mart. Fl. Bras.* xii. II. 266.—Am. trop.
unifoliolatum, *Engl. l. c.* 262.—Am. trop.
venosum, *Engl. l. c.* 264.—Am. trop.
Warmingianum, *March. in Kjoeb. Vidensk. Meddel.* (1873) 55.—Bras.
Widgrenii, *Engl. in Mart. Fl. Bras.* xii. II. 272.—Bras.
Zollingeri, *Engl. in DC. Monog. Phan.* iv. 71.—Java.

PROTIUM, Wight & Arn. Prod. 176 (1834) = **Commiphora**, Jacq. (Burser.).
africanum, Harv. in Harv. & Sond. Fl. Cap. ii. 592 = C. Harveyi.
caudatum, Wight & Arn. Prod. 176 = C. caudata.
gileadense, Wight & Arn. l. c. 177 = C. Berryi.
mossambicense, Oliver, Fl. Trop. Afr. i. 329 = C. mossambicense.
pubescens, Wight & Arn. Prod. 176 = C. pubescens.
Roxburghiana, Wight & Arn. l. c. = C. caudatum.

PROTOHOPEA, Miers, in Journ. Linn. Soc. xvii. (1879) 289 = **Symplocos**, Linn. (Styrac.).
cernua, Miers, l. c. = S. Hartwegi.
tinctoria, Miers, l. c. = S. tinctoria.

PROTOLEPIS, Steud. Nom. ed. II. ii. 401 (1841), sphalm. = **Proteopsis**, Mart. & Zucc. (Compos.)

PROTORHUS, Engl. in Engl. Bot. Jahrb. i. (1881) 420. *ANACARDIACEAE*.
fulva, *Engl. l. c.* 421.—Madag.
Grandidieri, *Engl. l. c.* 422.—Madag.
latifolia, *Engl. l. c.* 421.—Madag.
longifolia, *Engl. l. c.* 422.—Afr. austr.
nitida, *Engl. l. c.* 421.—Madag.
oblongifolia, *Engl. l. c.*—Madag.
sericea, *Engl. l. c.*—Madag.
Thouarsii, *Engl. l. c.* 422.—Madag.

PROUSTIA, Lag. Amen. Nat. i. 33 (1811). *COMPOSITAE*, Benth. & Hook. f. ii. 500.
HARMODIA, D. Don, ex Steud. Nom. ed. II. i. 722 (1840).
angustifolia, *Wedd. Chlor. And.* i. 24.—Bolivia.
baccharoides, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 106.—Chili.
cinerea, *Phil. in Linnaea*, xxix. (1857–58) 109.—Chili.
cuneifolia, D. Don, in Trans. Linn. Soc. xvi. (1830) 202 = pungens.
domingensis, Spreng. ex DC. Prod. v. 30 = Vernonia buxifolia.
glandulosa, *DC. Prod.* vii. 27.—Chili.
ilicifolia, Hook. & Arn. Bot. Beech. Voy. 28 = pungens.
mendocina, *Phil. Sert. Mend. Alt.* 18.—Chili.
mexicana, Lag. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 201 = Perezia Humboldtii.
oblongifolia, *D. Don, l. c.* 199.—Peruv.
pungens, *Poepp. ex Less. Syn. Comp.* 110.—Chili.
pyrifolia, *Lag. Amen. Nat.* i. 33; *DC. in Ann. Mus. Par.* xix. (1812) 67. t. 4.—Chili.
reticulata, Lag. ex D. Don, in Trans. Linn. Soc. xvi. (1830) 200 = Perezia reticulata.
reticulata, *Phil. in Anal. Univ. Chil.* (1861) 50.—Chili.
Tipia, *Phil. Fl. Atac.* 28.—Chili.
Vanillosma, *Wright, in Sauv. Fl. Cub.* 83.—Cuba.

PROVENZALIA, Adans. Fam. ii. 469 (1763) = **Calla**, Linn. (Aroid.).
bispatha, Rafin. New Fl. Am. ii. 90; Fl. Tellur. iii. 67 = C. palustris.
brevis, Rafin. New Fl. Am. ii. 90 = C. palustris.
heterophyla, Rafin. l. c. = C. palustris.
palustris, Rafin. l. c. = C. palustris.

PROZETIA, Neck. Elem. i. 225 (1790) = **Pouteria**, Aubl. (Sapot.).

PROZOPSIS, C. Muell. in Walp. Ann. v. 62 (1858), err. typ. = POZOPSIS, Hook. = **Huanaca**, Cav. (Umbell.).

PRUMNOPITYS, Phil. in Linnaea, xxx. (1859–60) 731 **Podocarpus**, L'Hérit. (Conif.).
elegans, Phil. l. c. 732 = P. affinis.

PRUNELLA, Linn. Sp. Pl. ed. I. 600 (1753). *LABIATAE* [Benth. & Hook. f. ii. 1203].
BRUNELLA, Tourn. ex Linn. Syst. ed. I (1735).
aequinoctialis, H. B. & K. Nov. Gen. et Sp. ii. 323 = vulgaris.
alba, Pall. ex Bieb. Fl. Taur. Cauc. ii. 67 = laciniata.
algeriensis, Noé, in Bull. Soc. Bot. Fr. ii. (1855) 582 = vulgaris.
alpina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 59 = vulgaris.
alpina, Timb. in Bull. Soc. Bot. Fr. xiii. (1866), Sess. extr. p. cliii = grandiflora.
angustifolia, Wender. Schrift. Marb. ii. (1831) 258 = vulgaris.
australasica, Moric. ex Buch-Ham. in Ser. Bull. Bot. i. (1830) 162 = vulgaris.
australis, Sweet, Hort. Brit. ed. II. 410 = vulgaris.
bicolor ×, Beck, in Verh. Zool.-Bot. Ges. Wien, xxxii. (1883) 185.—Europ.
bicolor, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 8 (Quid?).—Java.
Browniana, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 597 = vulgaris.
caerulea, Gueldenst. ex Ledeb. Fl. Ross. iii. 393 = vulgaris.
canadensis, Mill. Gard. Dict. ed. VIII. n. 4 = vulgaris.
caroliniana, Mill. l. c. n. 6 = vulgaris.
cinerea, Rafin. New Fl. Am. ii. 30 = vulgaris.
cordata, Rafin. l. c. 31 = vulgaris.
dissecta, Wender. in Schrift. Marb. ii. (1831) 257 = vulgaris.
elongata, Dougl. ex Benth. Lab. Gen. et Sp. 417 = vulgaris.
Fischeriana, Buch.-Ham. in Ser. Bull. Bot. i. (1830) 164 = vulgaris.
grandiflora, *Jacq. Fl. Austr.* iv. 40. t. 377.—Europ.
hastaefolia, Brot. Fl. Lusit. i. 181 = grandiflora.
hastata, Spreng. Syst. ii. 692 = grandiflora.
heterophyla, Rafin. New Fl. Am. ii. 31 = vulgaris.
hirsuta, Wender. in Schrift. Marb. ii. (1831) 257 = vulgaris.
hirta, Bernh. ex Steud. Nom. ed. I. 661 = vulgaris.

PRUNELLA:—
hispida, Benth. in Wall. Pl. As. Rar. i. 66 = vulgaris.
hybrida, 'Kn.' ex Nym. Consp. 574 = laciniata.
hyssopifolia, *Linn. Sp. Pl.* 600.—Europ.
incisa, Link, Enum. Hort. Berol. ii. 121 = vulgaris.
indica, Burm. f. Fl. Ind. 130 = Acrocephalus capitatus.
intermedia, Brot. Fl. Lusit. i. 180 = laciniata.
intermedia, Link, Handb. i. 490 = vulgaris.
intermedia, Reichb. Pl. Crit. iii. 4 = Cleonia lusitanica.
laciniata, Leyss. Fl. Halens. 148 = grandiflora.
laciniata, *Linn. Sp. Pl.* ed. II. 837.—Europ.; Afr. bor.; Oriens.
laciniata, Walt. Fl. Carol. 163 = vulgaris.
latifolia, Donn, Hort. Cantab. ed. II. 80 = vulgaris.
longifolia, Pers. Syn. ii. 137 = vulgaris.
lusitanica, Linn. Sp. Pl. 601 = Cleonia lusitanica.
macrophylla, Steud. Nom. ed. II. ii. 402 = vulgaris.
mariquitensis, Willd. ex Benth. in Linnaea, xi. (1837) 345 = vulgaris.
microphylla, Rafin. New Fl. Am. ii. 29 = vulgaris.
Novae-Angliae, Mill. Gard. Dict. ed. VIII. n. 7 = vulgaris.
obtusifolia, Rafin. New Fl. Am. ii. 32 = vulgaris.
odorata, Lam. Fl. Fr. ii. 367 = Cleonia lusitanica.
officinalis, Crantz, Stirp. Austr. ed. II. fasc. iv. 279 = vulgaris.
ovata, Pers. Syn. ii. 137 = vulgaris.
parviflora, Gilib. Fl. Lituan. i. 88; Poir. Voy. Barb. ii. 188 = vulgaris.
pensylvanica, Willd. Hort. Berol. i. 9 = vulgaris.
petiolaris, Rafin. New Fl. Am. ii. 30 = vulgaris.
pinnatifida, Pers. Syn. ii. 137 = vulgaris.
pratensis, Schur, Enum. Pl. Transs. 543 = vulgaris.
purpurea, Gueldenst. ex Ledeb. Fl. Ross. iii. 393 = vulgaris.
pyrenaica, Philippe, Fl. Pyrén. ii. 173 = grandiflora.
reptans, Dum. Fl. Belg. 44 = vulgaris.
reticulata, Rafin. New Fl. Am. ii. 31 = vulgaris.
rosea, Rafin. l. c. = vulgaris.
sessilifolia, Rafin. l. c. 30 = vulgaris.
speciosa, Wender. in Schrift. Marb. ii. (1831) 257 = grandiflora.
sulphurea, Mill. Gard. Dict. ed. VIII. n. 5 = laciniata.
surrecta, Dum. Fl. Belg. 44 = vulgaris.
Tournefortii, Timb. in Bull. Soc. Bot. Fr. xiii. (1866), Sess. extr. p. cliv = grandiflora.
transsilvanica, Schur, Enum. Pl. Transs. 953 (Index) = grandiflora.
variabilis ×, Beck, in Verh. Zool.-Bot. Ges. Wien, xxxii. (1883) 186.—Europ.
violacea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 8 (Quid?).—Java.
violacea, Nym. Consp. 574 = laciniata.
vulgaris, *Linn. Sp. Pl.* 600.—Reg. temp.

PRUNOPHORA, Neck. Elem. ii. 71 (1790) = **Prunus**, Tourn. (Rosac.).

PRUNOPSIS, André, in Rev. Hortic. (1883) 967 = **Prunus**, Tourn. (Rosac.).
Lindleyi, André, l. c. 67 = Prunus triloba, *Lindl.*

PRUNUS, [Tourn.] Linn. Syst. ed. I (1735). *ROSACEAE*, Benth. & Hook. f. i. 609.
Amygdalopsis, Carr. in Rev. Hortic. (1862) 91.
Amygdalus, [Tourn.] Linn. Syst. ed. I (1735).
Armeniaca, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
Ceraseidos, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1843) 743. t. 5.
Cerasophora, Neck. Elem. ii. 71 (1790).
Cerasus, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 141 (1737).
Chimanthus, Rafin. Fl. Ludov. 26 (1817).
Druparia, [Clairv.] Man. Herb. 159 (1811).
Hagidryas, Griseb. Reise Rumel. i. 311 (1841).
Laurocerasus, [Tourn.] M. Roem. Fam. Rosifl. 5, 89 (1847).
Padus, Linn. Syst. ed. I (1735); Gen. ed. I. 142 (1737).
Persica, [Tourn.] Mill. Gard. Dict. ed. VI. (1752).
Prunophora, Neck. Elem. ii. 71 (1790).
Prunopsis, André, in Rev. Hortic. (1883) 367.

PRUNUS:—
Trichocarpus, Neck. Elem. ii. 70 (1790).
Tubopadus, Pomel, Mat. Fl. Atl. 8 (1860).
Acacia, Crantz, ex Poir. Encyc. v. 679 = seq.
Acacia-germanica, Crantz, Stirp. Austr. ed. I. fasc. ii. 58; ed. II. 93 = spinosa.
acida, Ehrh. Beitr. v. 162 = P. Cerasus.
acinaria, Desf. Tabl. ed. I. 179 = americana.
acinaria, Poir. Encyc. v. 677 = domestica.
acuminata, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 599 = *pubescens*, Clav.
acuminata, *Hook. f. Fl. Brit. Ind.* ii. 317.—Reg. Himal.
acuminata, Hort. ex C. Koch, Dendrol. i. 101 = americana.
acuminata, Michx. Fl. Bor. Am. 284 = maritima.
adenophylla, Wall. Cat. sub n. 718 = undulata.
aestiva, Salisb. Prod. 356 = P. Cerasus.
agrestis, Jord. & Fourr. Brev. Pl. Nov. ii. 37; Ic. Fl. Europ. i. 56. t. 155 = insititia.
alleghaniensis, *Porter, in Coult. Bot. Gaz.* ii. (1877) 85.—Am. bor.
Aloocha, Royle, Illustr. Bot. Himal. 205, nomen = insititia?
Amarella, *Reichb. Fl. Germ. Excurs.* 646.—Europ. austr.
ambigua, Salisb. Prod. 356 = insititia.
amelanchieriflora, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. x. (1876) 490 = insititia?
americana, *Marsh. Arbust.* 111.—Am. bor.
amygdalina, Poir. Encyc. v. 677 = domestica.
amygdalina, Webb, Iter Hisp. 47 = Ramburei.
Amygdalus, *Stokes, Bot. Mat. Med.* iii. 101.—Afr. bor.; Oriens.
Andersoni, *A. Gray, in Proc. Am. Acad.* vii. (1868) 337.—Am. bor. occ.
angustifolia, Marsh. Arbust. 111 = P. Chicasa.
arbascensis, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. ii. (1874) 451 = spinosa *vel* insititia.
arguta, Bigel. ex M. Roem. Syn. Rosifl. 86 = virginiana.
Armeniaca, Blanco, Fl. Filip. ed. II. 296 = triflora.
Armeniaca, *Linn. Sp. Pl.* i. 474.—Reg. Caucas.
Armeniaca, Thunb. Fl. Jap. 200 = P. Mume.
armenioides, Lieg. Syst. Pflaum. ii. 162; ex M. Roem. Syn. Rosifl. 52 = domestica.
aspera, *Thunb. Fl. Jap.* 201.—Japon.
Aubertiana, Ser. ex Steud. Nom. ed. II. ii. 402 = domestica.
aucubaefolia, Hort. ex C. Koch, Dendrol. i. 120 = P. Padus.
augustana, Poir. Encyc. v. 677 = domestica.
aureliana, Poir. l. c. 678 = domestica.
austera, Ehrh. Beitr. v. 160; vii. 50 = P. Cerasus.
austriaca, Lieg. Syst. Pflaum. ii. 10; ex M. Roem. Syn. Rosifl. 37 = domestica.
autumnalis, Lieg. l. c. 23; ex M. Roem. l. c. 44 = domestica.
avium, *Linn. Fl. Suec.* ed. II. 474.—Europ.
baccata, Linn. Amoen. Acad. vii. 456 (Quid?).—Rossia.
Ballota, Lamotte, Prod. Fl. Centr. Fr. i. 232 = domestica.
biferum, Carr. in Rev. Hortic. (1875) 416 (Quid?).—Hab.?
Blumii, *D. Dietr. Syn. Pl.* iii. 45.—China.
bokhariensis, Royle, Illustr. Bot. Himal. 205, nomen = insititia.
borealis, Poir. Encyc. v. 674 = pensylvanica.
borealis, Salisb. Prod. 356 = P. Padus.
brachybotrya, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 348.—Mexic.
brachypetala, *Walp. Ann.* i. 272.—Persia.
bracteata, *Franch. & Sav. Enum. Pl. Jap.* ii. 329.—Japon.
brasiliensis, Schott, ex Spreng. Syst. iv. 406 = sphaerocarpa.
brigantiaca, *Vill. Fl. Delphin.* 49; *Hist. Pl. Dauph.* i. 380.—Gallia.
brigantina, Nym. Consp. 213 = praec.
brignola, Poir. Encyc. v. 478 = domestica.
brignola, Reichb. Fl. Germ. Excurs. 645 = insititia.
brignoniensis, *Ehrh. Beitr.* vii. 50, nomen.—Europ.
Buergeriana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 92.—Japon.
Bungei, Walp. Rep. ii. 9 = humilis.
campanulata, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 103.—China.

PRUNUS:—
canadensis, Linn. Sp. Pl. ed. II. 678 = americana?
canadensis, Marsh. Arbust. 113 = virginiana.
canadensis, [Moç. & Sesse,] ex DC. Prod. ii. 539 = P. Capollin.
candicans, Bess. Enum. Pl. Volh. 58 = spinosa.
candicans, Willd. Enum. Hort. Berol. Suppl. 32; Balb. Cat. Taur. (1813) 62 = insititia?
Capollin, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 355. t. 8.—Mexic.
capricida, Wall. in DC. Prod. ii. 540 = undulata.
Caproniana, Hort. ex Reichb. Fl. Germ. Excurs. 644 in syn. = P. Cerasus.
Capuli, Cav. ex Spreng. Syst. ii. 477 = P. Capollin.
carolina, Duroi, Harbk. Baumz. i. 198 = seq.
caroliniana, *Ait. Hort. Kew.* ed. I. ii. 163.—Am. bor.
cartilaginea, Lehm. Ind. Sem. Hort. Hamb. (1833).—Cf. Linnaea, x. (1836) Lit. 76 = serotina.
catharinea, Steud. Nom. ed. II. ii. 402 = domestica.
caucasica, *Gueldenst. ex Pall. Fl. Ross.* i. 17.—Reg. Caucas.
cerasa, Crantz, Stirp. Austr. ed. I. fasc. ii. 58; ed. II. 94 = pumila.
Ceraseidos, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 103.—Japon.
cerasifera, Ehrh. Beitr. iv. 17 = divaricata.
cerasiflora, Paill. in Bull. Soc. Bot. Fr. xvi. (1869) p. xvi = insititia.
cerasifolia, S. Wats. Bibliog. Ind. N. Am. Bot. 306 = pensylvanica.
cerasoides, D. Don, Prod. Fl. Nep. 239 = P. Puddum.
Cerasus, *Linn. Sp. Pl.* 474.—Europ.; Oriens.
Cerasus, Scop. Fl. Carn. ed. II. i. 343 = avium.
cerea, Ehrh. Beitr. vii. 50 = domestica.
cereola, Poir. Encyc. v. 677 = domestica.
ceylanica, Miq. Fl. Ind. Bat. i. 366 = Pygeum Wightianum.
Chamaecerasus, *Jacq. Coll.* i. 133.—Europ. austr.; As. bor.
Chapronii, *Carr. in Rev. Hortic.* (1881) 467.—Roumania.
Chicasa, *Michx. Fl. Bor. Am.* i. 284.—Am. bor.
chinensis, Blume, Bijdr. 1104 = japonica.
cinerascens, Lamotte, Prod. Fl. Centr. Fr. i. 231 = insititia.
claudiana, Poir. Encyc. v. 677 = domestica.
coccinea, Rafin. Fl. Ludov. 135 = americana.
Cocomilia, *Tenore, Fl. Nap. Prod.* p. lxviii.—Italia.
colchica, Hort. ex C. Koch, Dendrol. i. 125 = P. Laurocerasus.
communis, Huds. Fl. Angl. ed. II. i. 212 = spinosa, insititia.
compressa, Lieg. Syst. Pflaum. ii. 148; ex M. Roem. Syn. Rosifl. 44 = domestica.
congestiflora, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 604 = spinosa?
cornuta, Wall. Cat. n. 716 = P. Padus.
coronata, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 597 (Quid?).—Gallia.
cuneata, Rafin. Ann. Nat. 11 = pumila.
cuneifolia, *Kit. in Linnaea*, xxxii. (1863) 603.—Europ. austr.
Cupaniana, Guss. ex Nym. Consp. 213 = P. Mahaleb.
damascena, Ehrh. Beitr. vii. 51 = domestica.
Damsonia, M. Roem. Syn. Rosifl. 54 = domestica.
dasycarpa, *Ehrh. Beitr.* vi. 90.—Oriens?
Davidiana, Franch. Pl. Davidian. i. 103 = P. Persica.
declinata, Marsh. Arbust. 112 = maritima.
demissa, D. Dietr. Syn. Pl. iii. 43 = virginiana.
densa, Martr. Pl. Crit. Tarn, 22 = spinosa.
densiflora, Jord. & Fourr. Brev. Pl. Nov. ii. 24 = insititia.
densiflora, Steud. Nom. ed. II. ii. 403 = virginiana.
depressa, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 600 (Quid?).—Gallia.
depressa, Lieg. Syst. Pflaum. ii. 198; ex M. Roem. Syn. Rosifl. 50 = domestica.
depressa, Pursh, Fl. Am. Sept. i. 332 = pumila.
Desvauxii, Bor. Fl. Centr. Fr. ed. III. ii. 182 = insititia.
diapraea, Poir. Encyc. v. 678 = domestica.
divaricata, *Ledeb. Ind. Hort. Dorp. Suppl.* (1824) 6.—Oriens.
domestica, *Linn. Sp. Pl.* 475.—Europ.; Reg. Cauc.
domestica, Thunb. Fl. Jap. 203 = japonica.
donarium, Siebold, in Verh. Batav. Gen. xii. (1830) 68 = paniculata.

PRUNUS:—
Duerinckii, Walp. Rep. ii. 10 = virginiana.
dulcis, Mill. ex Reichb. Fl. Germ. Excurs. 644 = avium.
dumetorum, Jord. & Fourr. Brev. Pl. Nov. ii. 26; Lamotte, Prod. Fl. Centr. Fr. i. 231 = insititia.
dumosa, Salisb. Prod. 356 = virginiana.
duracina, *Sweet, Hort. Brit.* ed. I. 134.—Europ. austr.
eburnea, *Aitch. in Journ. Linn. Soc.* xviii. (1880) 50.—Persia; Afghan.
elegans, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 601 (Quid ?).—Gallia.
elliptica, *Thunb. Fl. Jap.* 199.—Japon.
emarginata, *Walp. Rep.* ii. 9.—Am. bor. occ.
enucleata, Poir. Encyc. v. 678 = domestica.
erecta, Walp. Ann. iii. 854 = emarginata.
erubescens, Paill. in Bull. Soc. Bot. Fr. xvi. (1869) p. xv = insititia.
erythrocalyx, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 596 (Quid ?).—Gallia.
exigua, Bechst. Forstb. ed. V. 426 = insititia.
fallax, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xx. (1866) 29, nomen.—Europ.
fasciculata, *A. Gray, in Proc. Am. Acad.* x. (1874) 70.—Am. bor. occ.
fastigiata, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. x. (1876) 489 = spinosa.
ferruginea, *Steud. Nom.* ed. II. ii. 400.—Mexic.
ferruginea, Wall. Cat. sub n. 721 = rufa.
fimbriata, Steud. Nom. ed. II. ii. 403 = virginiana.
flava, Kit. in Linnaea, xxxii. (1863) 603 = cuneifolia.
floribunda, Hort. ex C. Koch, Dendrol. i. 119 = pumila.
floribunda, Weihe, in Flora, ix. (1826) 747 = *vinaria*.
foecundissima, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 595 = spinosa.
Fremonti, *S. Wats. in Bot. Calif.* ii. 442.—Calif.
fruticans, Weihe, in Flora, ix. (1826) 748 = spinosa.
fruticosa, Pall. Fl. Ross. i. 19 = P. Chamaecerasus.
galatensis, Poir. Encyc. v. 677 = domestica.
gallica, *Ehrh. Beitr.* vii. 51.—Europ.
glandulifolia, *Rupr. ex Maxim. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1856) 130.—Manchur.
glandulosa, Thunb. Fl Jap. 202 = japonica.
glandulosa, *Torr. & Gray, Fl. N. Am.* i. 408.—Am. bor. occ.
glaucifolia, Wall. Cat. n. 1008 = nepaulensis.
gracilis, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 243.—Texas.
graeca, *Desf. ex Steud. Nom.* ed. II. ii. 403.—Graecia.
grandifolia, Salisb. Prod. 356 = P. Laurocerasus.
grandissima, Lieg. Syst. Pflaum. ii. 62; ex M. Roem. Syn. Rosifl. 42 = domestica.
Grayana, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 107.—Japon.
gymnoclada, Clav. in Actes Linn. Soc. Bord. xxxviii. (1884) 602 (Quid ?).—Gallia.
Hausmanni, Nym. Consp. 214 = spinosa.
hirsuta, Ell. Sketch, i. 541 = virginiana ?
Hixa, Brouss. ex Willd. Enum. Hort. Berol. 517 = lusitanica.
hortensis, Pers. Syn. ii. 34 = P. Cerasus.
humilis, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 97.—China.
humilis, Colla, Herb. Pedem. ii. 293 = prostrata.
hungarica, Poir. Encyc. v. 676 = domestica.
Husmanni, Boechel, in Bot. Zeit. xi. (1853) 827 = spinosa.
hyemalis, Michx. Fl. Bor. Am. i. 284 = americana.
ilicifolia, *Walp. Rep.* ii. 10.—Am. bor. occ.
imperialis, Poir. Encyc. v. 676 = domestica.
incana, *Decne. in Rev. Hortic.* Sér. IV. ii. (1853) 281. t. 15.—Reg. Caucas.
incana, Schwein. in Long, Exped. Winnep. ii. 387 = pumila.
incana, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 263 = prostrata.
incisa, *Thunb. Fl. Jap.* 202.—Japon.
inermis, Gilib. Fl. Lituan. ii. 227 = domestica.
insignis, Carr. in Rev. Hortic. (1871) 534 = spinosa.
insititia, Fries, ex M. Roem. Syn. Rosifl. 35 = spinosa.
insititia, *Linn. Cent. Pl.* i. 12; *Amoen. Acad.* iv. 273.—Europ.; As. Min.; Reg. Himal.
insititia, Poir. Encyc. v. 678 = domestica.
insititia, Walt. Fl. Carol. 146 = P. Chicasa.
integerrima, Steud. Nom. ed. II. ii. 403 = acuminata.
integrifolia, *Walp. Ann.* iii. 854.—Peruv.
intermedia, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 603 = spinosa ?

PRUNUS:—
intermedia, Poir. Encyc. v. 674 = P. Chamaecerasus.
italica, Borck. Forstb. ii. 1409 = insititia.
italica, Lieg. Syst. Pflaum. ii. 75; ex M. Roem. Syn. Rosifl. 40 = domestica.
Itosakura, Siebold, in Verh. Batav. Gen. xii. (1830) 68 = incisa.
Jacquemontii, *Hook. f. Fl. Brit. Ind.* ii. 314.—Reg. Himal.
Jamasakura, Siebold, in Verh. Batav. Gen. xii. (1830) 68 = paniculata.
japonica, Carr. Rev. Hortic. (1873) 457 = humilis.
japonica, *Thunb. Fl. Jap.* 201.—Japon.; China.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 365; *Blume, Mélang. Bot.* (1855).—*Cf. Flora*, xli. (1858) 255.—Java.
Jenkinsii, *Hook. f. Fl. Brit. Ind.* ii. 317.—Assam.
Juliana, Poir. Encyc. v. 676 = domestica.
Juliana, Reichb. Fl. Germ. Excurs. 643 = P. Cerasus.
Junghuhniana, *Miq. Fl. Ind. Bat.* i. 1. 366.—Java.
Kerii, *Steud. Nom.* ed. II. ii. 403.—Japon.
Kolomikta, Maxim. in Bull. Phys.-Math. Pétersb. xv. (1856) 129 = Actinidia callosa.
lanceolata, Willd. Berl. Baumz. ed. I. 240 = pensylvanica.
lancifolia, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 600 = maritima.
latiflora, Clav. l. c. = insititia.
latifolia, Jord. & Fourr. Brev. Pl. Nov. ii. 27 = insititia.
latifolia, *Moench, Verz. Ausl. Baeume*, 85.—Am. bor.
laurifolia, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 458.—Ins. Timor.
laurifolia, Schlecht. in Linnaea, xiii. (1839) 404 = Schiedeana.
Laurocerasus, *Linn. Sp. Pl.* 474.—Oriens.
laxiflora, Kit. in Linnaea, xxxii. (1863) 602 = P. Cerasus ?
leucocarpa, M. Roem. Syn. Rosifl. 47 = domestica.
ligerina, Lloyd, Fl. L'ouest Fr. ed. III. 102 = spinosa.
littoralis, Bigel. Fl. Bost. ed. II. 193 = maritima.
lucida, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 597 = *erythrocalyx*.
lusitanica, Gueldenst. ex Ledeb. Fl. Ross. ii. 9 = P. Laurocerasus.
lusitanica, *Linn. Sp. Pl.* 473.—Europ. austr.
lusitanica, Walt. Fl. Carol. 146 = caroliniana.
lutea, Bechst. Forstb. ed. V. 425 = insititia.
Maackii, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 361.—Manchuria.
macrophylla, Poir. Encyc. Suppl. iv. 584 = duracina.
macrophylla, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. ii. (1843) 122.—Japon.
Mahaleb, *Linn. Sp. Pl.* 474.—Europ. austr.; Oriens.
maliformis, Poir. Encyc. v. 677 = domestica.
Marasca, Reichb. Fl. Germ. Excurs. 644 = P. Cerasus.
maritima, *Wangenh. Am.* 103.—Am. bor.
martabanica, *Kurz, For. Fl. Brit. Burma*, i. 434.—Burma; Ins. Andam.
martabanica, Steud. Nom. ed. II. ii. 403 (gen. dub.).—Ind. or.
Martrini, Genev. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 29, nomen; et in Bull. Soc. Bot. Fr. xiv. (1867) Bibl. 43 = spinosa.
Maximowiczii, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1856) 131.—Manchuria.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 353.—Mexic.
Meyeri, Boeckel, in Bot. Zeit. xi. (1853) 827 = spinosa.
micrantha, Steud. Nom. ed. II. ii. 403 = virginiana.
microcarpa, *C. A. Mey. Verz. Pfl. Cauc.* 166.—Reg. Cauc.; Persia.
micropetala, *C. Koch, Dendrol.* i. 118.—Calif.
microphylla, *Hemsl. Biol. Centr. Am. Bot.* i. 368.—Mexic.
microstigma, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. x. (1876) 490 = spinosa.
minutiflora, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 185.—Am. bor. occ.
minutiflora, S. Wats. in Bot. King's Exp. 420 = fasciculata.
Miqueliana, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 98.—Japon.
mirobolana, DC. Prod. ii. 533 = domestica.
Mississipi, Marsh. Arbust. 112 = americana.
mollis, Torr. Fl. N. & M. U. St. 470 = americana.
mollis, Walp. Rep. ii. 9 = emarginata.
mongolica, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 16.—Mongolia.
montana, Hort. ex C. Koch, Dendrol. i. 122 = virginiana.

PRUNUS:—
montana, Marsh. Arbust. 113 = pensylvanica ?
montana Schur, Enum. Pl. Transs. 179 = spinosa.
monticola, *C. Koch, in Ind. Sem. Hort. Berol.* (1854) *App.* 15.—As. Min.
multiglandulosa, Cav. in Anal. Cienc. Nat. iii. (1801) 59 = lusitanica.
Mume, *Sieb. & Zucc. Fl. Jap.* i. 29.—Japon.
myrobalana, Loisel. in Duham. Arb. ed. Nov. v. 184 = divaricata.
myrtellina, Poir. Encyc. v. 678 = domestica.
nana, Duroi, Obs. Bot. 12 = virginiana.
nana, *Stokes, Bot. Mat. Med.* iii. 103.—Europ.; Reg. Cauc.; Sibir.
napaulensis, C. Koch, Dendrol. i. 89 = seq.
nepalensis, Hort. ex C. Koch, l. c. = P. Armeniaca.
nepaulensis, *Steud. Nom.* ed. II. ii. 403.—Reg. Himal.
nicotianaefolia, Loisel. ex Steud. l. c. = duracina.
nigra, Ait. Hort. Kew. ed. I. ii. 165 = americana.
nigra, Ehrh. Beitr. v. 161.—Europ.
nigra, Reichb. Fl. Germ. Excurs. 645 = insititia.
nigricans, Ehrh. Beitr. vii. 126 = avium.
nitida, Salisb. Prod. 356 = caroliniana.
nitidissima, Hassk. Retzia, i. 175 = spinulosa.
obovata, Bigel. Fl. Bost. ed. II. 192 = virginiana.
occidentalis, *Sw. Prod. Veg. Ind. Occ.* 80.—Ind. occ.
odorata, Lam. Fl. Fr. iii. 108 = P. Mahaleb.
oeconomica, Borck. Forstb. ii. 1401 = domestica.
opaca, *Walp. Rep.* v. 648.—N. Granat.
orientalis, Walp. l. c. ii. 911 = microcarpa.
ovalifolia, Salisb. Prod. 356 = domestica.
oviformis, Jord. & Fourr. Brev. Pl. Nov. ii. 25 = spinosa.
oxycarpa, Bechst. Forstb. ed. V. 424 = P. Cerasus.
oxycarpa, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 111 = macrophylla.
oxypyrena, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 586, 595 = P. Padus.
Padus, *Linn. Sp. Pl.* 473,—Europ.; As. bor.
paniculata, *Thunb. Fl. Jap.* 200.—Japon.
parviflora, Paill. in Bull. Soc. Bot. Fr. (1869) p. xvi = spinosa.
pauciflora, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 97.—China bor.
pectinata, Walp. Rep. ii. 911 = P. Puddum.
pedunculares, Jord. & Fourr. Brev. Pl. Nov. ii. 25 = spinosa.
pedunculata, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 78.—Sibir. baical.
pendula, *Desf. Tabl.* ed. II. 272, nomen.—Hab. ?
Perdigona, Poir. Encyc. v. 676 = domestica.
pensylvanica, *Linn. f. Suppl.* 252.—Am. bor.
Persica, *Stokes, Bot. Mat. Med.* iii. 100.—As. temp.
persicifolia, Desf. Hist. Arbr. ii. 205 = pensylvanica
Pertigona, Hort. ex Steud. Nom. ed. II. ii. 404, sphalm. = domestica.
petraea, Tausch, in Flora, xxi. (1838) 719 = P. Padus.
Petzoldi, C. Koch, Dendrol. i. 92 = triloba.
phaeosticta, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 110 = punctata.
Pissardi, Carr. in Rev. Hortic. (1881) 190 = divaricata.
plena, Poir. Encyc. v. 671 = P. Cerasus.
pleuradenia, Griseb. Fl. Brit. W. Ind. 231 = sphaerocarpa.
podolica, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 301 (Quid ?).—Rossia.
pogonostyla, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 11.—China.
Poiretiana, Heynh. Nom. ii. 564 = maritima.
polymorpha, Spenn. Fl. Friburg. iii. 732 = spinosa, insititia, domestica.
porcorum, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 605 (Quid ?).—Gallia.
praecox, Poir. Encyc. v. 677 = domestica.
praecox, Salisb. Prod. 356 = spinosa.
prostrata, *Labill. Dec. Pl. Syr.* i. t. 6.—Reg. Mediterr.; Oriens.
Pruna, Crantz, Stirp. Austr. ed. I. fasc. ii. 59 = divaricata.
Pruneauliana, Steud. Nom. ed. II. ii. 404 = domestica.
Pseudo-armeniaca, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. v. 96.—Graecia.
Pseudo-cerasus, Lindl. in Trans. Hort. Soc. vi. (1826) 91 = paniculata.
pubescens, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 599 (Quid ?).—Gallia.
pubescens, Pursh, Fl. Am. Sept. i. 331 = maritima.
pubigera, Steud. Nom. ed. II. ii. 404 = maritima.

PRUNUS:—
Puddum, Hort. ex Hook. Exot. Fl. ii. sub t. 148 = Sarcococca pruniformis.
Puddum, *Roxb. ex Wall. Pl. As. Rar.* ii. 37.—Reg. Himal.
pumila, *Linn. Mant.* i. 75 (excl. syn.).—Am. bor.
pumila, Lumn. Fl. Poson. 201; Georgi, Beschr. Russ. Reich. III. iv. 1004 = P. Chamaecerasus.
pumila, Walt. Fl. Carol. 146 = umbellata.
punctata, *Hook. f. Fl. Brit. Ind.* ii. 317.—Reg. Himal.
pyaeidicalyx, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 596 = *erythrocalyx*.
pygmaea, Willd. Berl. Baumz. ed. I. 248 = maritima.
pyramidalis, DC. Fl. Fr. iv. 485 = domestica.
racemosa, Lam. Fl. Fr. iii. 107 = P. Padus.
Ramburei, *Boiss. Elench.* 39.—Hispan.
reclinata, Bosc, ex Spach, Hist. Vég. Phan. i. 398 = maritima.
reflexa, *Walp. Rep.* ii. 910.—Bras.
regia, Poir. Encyc. v. 678 = domestica.
rivularis, *Scheele, in Linnaea*, xxi. (1848) 594.—Texas.
rosea, Poir. Encyc. v. 671 = P. Cerasus.
rosiflora, Paill. in Bull. Soc. Bot. Fr. xvi. (1869) p. xvi = insititia.
rotundifolia, Rafin. Atl. Journ. 78 (Quid?).—Am. bor.
rubella, Bechst. Forstb. ed. V. 428 = insititia.
rubella, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 597 = *erythrocalyx*.
rubella, *Ehrh. Beitr.* v. 162.—Germania.
rubescens, Lamotte, Prod. Fl. Centr. Fr. i. 234 = domestica.
rubicunda, Bechst. Forstb. ed. V. 226, 492 = avium.
rubra, Ait. Hort. Kew. ed. I. ii. 163 = virginiana.
rubra, Willd. Berl. Baumz. ed. I. 238 = P. Padus.
rufa, *Steud. Nom.* ed. II. ii. 402.—Reg. Himal.
rustica, Lamotte, Prod. Fl. Centr. Fr. i. 233 = domestica.
rusticana, Jord. & Fourr. Brev. Pl. Nov. ii. 24; Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 594 = spinosa.
salicifolia, H. B. & K. Nov. Gen. et Sp. iv. 241. t. 563 = serotina.
salicina, *Lindl. in Trans. Hort. Soc.* vii. (1830) 239.—China.
saligna, Royle, Illustr. Bot. Himal. 204, nomen = P. Persica.
samydoides, *Schlecht. in Linnaea*, xiii. (1839) 92, 404.—Mexic.
Sanctae-Catharinae, Lamotte, Prod. Fl. Centr. Fr. i. 235 = domestica.
saxicola, Bor. Fl. Centr. Fr. ed. III. ii. 183 = domestica.
Schiedeana, *Steud. Nom.* ed. II. ii. 404.—Mexic.
semperflorens, Ehrh. Beitr. vii. 132 = P. Cerasus.
sempervirens, Salisb. Prod. 356 = lusitanica.
sempervirens, Willd. Enum. Hort. Berol. Suppl. 33 = caroliniana.
sepivaga, Jord. & Fourr. Brev. Pl. Nov. ii. 26 = insititia.
serotina, *Ehrh. Beitr.* iii. 20.—Am. bor.
serotina, Poir. Encyc. v. 665 = virginiana.
serotina, Roth, Catalect. fasc. i. 58 = P. Cerasus.
serotina, Schur, Enum. Pl. Transs. 178 = spinosa.
serratifolia, Marsh. Arbust. 114 = caroliniana.
serrulata, Lindl. in Trans. Hort. Soc. vii. (1830) 238 = paniculata.
sibirica, *Linn. Sp. Pl.* 474.—Sibir.
silvatica, Roxb. Fl. Ind. ii. 501 = P. Puddum.
Simonii, *Carr. in Rev. Hortic.* (1872) 111.—China.
sinensis, Hort. ex Carr. l. c. (1869) 300 = seq.?
sinensis, Pers. Syn. ii. 36 = japonica.
sphaerica, Willd. Berl. Baumz. ed. II. 315 = maritima.
sphaerocarpa, Michx. Fl. Bor. Am. i. 274 = maritima.
sphaerocarpa, *Sw. Prod. Veg. Ind. Occ.* 80.—Am. trop.
spinosa, *Linn. Sp. Pl.* 475.—Europ.; Afr. bor.; Oriens.
spinosa, Walt. Fl. Carol. 146 = americana.
spinosissima, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 281.—Turkestan.
spinulosa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 122.—Japon.

PRUNUS:—
Ssiori, *F. Schmidt, Reisen Amurl.* 124.—Ins. Sachalin.
stenopetala, Manceau, Pl. Phan. Maine, 7 = spinosa.
stenophyllus, Rafin. Fl. Ludov. 98 (Quid?).—Am. bor.
stipulacea, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 97.—China.
suaveolens, Kit. in Linnaea, xxxii. (1863) 602 = insititia.
subcordata, *Benth. Pl. Hartw.* 308.—Calif.
subhirtella, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 91.—Japon.
subinermis, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 598 (Quid?).—Gallia.
subrotunda, Bechst. Forstb. ed. V. 426 = insititia.
sundaica, Miq. Fl. Ind. Bat. i. 1. 365 = spinulosa.
Susquehanae, Willd. Enum. Hort. Berol. 519 = pumila.
sylvatica, Hook. f. Fl. Brit. Ind. ii. 314 = P. Puddum.
sylvestris, Habl. Taur. 108 = insititia.
sylvestris, Mill. ex Steud. Nom. ed. I. 663 = spinosa.
sylvestris, Pers. Syn. ii. 35 = avium.
syriaca, Borckh. Forstb. ii. 1406 = divaricata.
Tawakonia, Lindh. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 186 = rivularis.
tenerrima, *Carr. in Rev. Hortic.* (1870–71) 473.—Hab.?
texana, D. Dietr. Syn. Pl. iii. 45 = glandulosa.
texana, Scheele, in Linnaea, xxi. (1848) 593 = rivularis.
tiliaefolia, Salisb. Prod. 356 = P. Armeniaca.
tinctoria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 23 = Symplocos fasciculata.
tomentosa, *Thunb. Fl. Jap.* 203.—Japon.
trichocarpa, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 96 = tomentosa.
triflora, Rafin. Ann. Nat. 11 (Quid?).—Am. bor.
triflora, *Roxb. Hort. Beng.* 38.—Burma; China.
trifolia, Roxb. Fl. Ind. ii. 501, lapsu = praec.
triloba, *Lindl. in Gard. Chron.* (1857) 268.—China.
turonensis, Steud. Nom. ed. II. ii. 404 = domestica.
ulmifolia, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 281.—Turkestan.
umbellata, *Ell. Sketch*, i. 541.—Am. bor.
undulata, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 239.—Reg. Himal.
ursina, *Kotschy, ex Boiss. Fl. Orient.* ii. 652.—Syria.
vapincensis, Jord. & Fourr. Brev. Pl. Nov. ii. 26 = insititia.
varactensis, Bor. Fl. Centr. Fr. ed. III. ii. 183 = domestica.
varia, Ehrh. Beitr. vii. 127 = avium.
verrucosa, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 280.—Turkestan.
vinaria, Bechst. Forstb. ed. V. 427 (Quid?).—Europ.
violacea, M. Roem. Syn. Rosifl. 52 = domestica.
virescens, Paill. in Bull. Soc. Bot. Fr. xvi. (1869) p. xvi. = insititia.
virgata, Martr. Pl. Crit. Tarn, 21 = spinosa.
virginalis, Wender. in Schrift. Marb. ii. (1830) 253 = virginiana.
virginea, M. Roem. Syn. Rosifl. 50 = domestica.
virginiana, Duroi, Obs. Bot. 12 = serotina.
virginiana, *Linn. Sp. Pl.* 473.—Am. bor.
virginica, Steud. Nom. ed. II. ii. 246, in syn. sphalm. = praec.
vulgaris, Schur, Enum. Pl. Transs. 954 (Index) = P. Cerasus.
vulgatior, Clav. in Act. Linn. Soc. Bord. xxxviii. (1884) 594 = spinosa.
Wallichii, Steud. Nom. ed. II. ii. 404 = acuminata.
xanthocarpos, Hort. ex C. Koch, Dendrol. i. 89 = P. Armeniaca.
Zippeliana, *Miq. Fl. Ind. Bat.* i. 1. 367.—Java.

PRYONA, Miq. Fl. Ind. Bat. i. 1. 1081 (1855) = **Crudia**, Schreb. (Legumin.).
bantamensis, Miq. l. c.—Java.

PRZEWALSKIA, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 507. *SOLANACEAE*.
tangutica, *Maxim. l. c.* 508.—Tibet.

PSACALIUM, Cass. in Dict. Sc. Nat. xliii. 461 (1826) = **Senecio**, Tourn. (Compos.).
arboreum, DC. Prod. vi. 335.—Am. austr.
glabratum, DC. l. c.—Am. austr.

PSACALIUM:—
kleinioides, DC. Prod. vi. 335.—N. Granat.
laxiflorum, Benth. Pl. Hartw. 41 = S. Moreliae.
peltatum, Cass. in Dict. Sc. Nat. xliii. 461 = S. peltiferus.
thyrsoideum, DC. Prod. vi. 335 = S. insignis.
vaccinioides, DC. l. c.—N. Granat.

PSALINA, Rafin. Fl. Tellur. iii. 25 (1836) = **Gentiana**, Tourn.
exacoides, Rafin. l. c. = Belmontia cordata.
utriculosa, Rafin. l. c. = G. utriculosa.

PSAMMA, Beauv. Agrost. 143. t. 6. f. 1 (1812) = **Ammophila**, Host (Gramin.).
arenaria, Roem. & Schult. Syst. ii. 845 = A. arundinacea.
australis, Mabille, Rech. Pl. Corse, fasc. i. 33 = A. arundinacea.
baltica, Roem. & Schult. Syst. ii. 845 = A. baltica.
littoralis, Beauv. Agrost. 143. t. 6. f. 1 = A. arundinacea.
pallida, Presl, Cyp. et Gram. Sic. 24 = A. arundinacea.

PSAMMANTHE, Hance, in Walp. Ann. ii. 659 (1852) = **Sesuvium**, Linn. (Ficoid.).
marina, Hance, l. c. 660 = S. Portulacastrum.

PSAMMANTHE, Reichb. Nom. 205 (1841) (Caryophyll.?).

PSAMMISIA, Klotzsch, in Linnaea, xxiv. (1851) 42. *VACCINIACEAE*, Benth. & Hook. f. ii. 566.
alpicola, *Klotzsch, l. c.* 45.—N. Granat.
bicolor, *Klotzsch, l. c.* 44.—Peruv.
breviflora, *Klotzsch, l. c.*—Peruv.
costaricensis, *Klotzsch, l. c.* 47.—Am. centr.
cyathifera, *Klotzsch, l. c.* 43.—N. Granat.
falcata, *Klotzsch, l. c.* 44.—N. Granat.
formosa, *Klotzsch, l. c.* 46.—Guian.
glabra, *Klotzsch. l. c.*—Am. centr.
guianensis, *Klotzsch, l. c.* 43.—Guian.
Hookeriana, *Klotzsch, l. c.* 46.—N. Granat.
leucostoma, *Benth. ex Meissn. in Mart. Fl. Bras.* vii. 127.—Bras.
longicolla, *Hook. Bot. Mag.* t. 5526.—Am. austr.
longifolia, *Klotzsch, in Linnaea*, xxiv. (1851) 43.—Peruv.
macrophylla, *Klotzsch, l. c.* 45.—N. Granat.
nitida, *Klotzsch, l. c.*—N. Granat.
oblongifolia, *Regel, Ind. Sem. Hort. Petrop.* (1866) *Suppl.*; (1867) 33.—Hab.?
pauciflora, Griseb. in Lechl. Berb. Am. Austr. 58 = Macleania pauciflora.
penduliflora, *Klotzsch, in Linnaea*, xxiv. (1851) 43.—N. Granat.
Planchoniana, *Decne. in Rev. Hortic.* Sér. IV. iii. (1854) 182.—N. Granat.
ramiflora, *Klotzsch, in Linnaea*, xxiv. (1851) 44.—Am. centr.
rupestris, *Klotzsch, l. c.* 45.—N. Granat.
sarcantha, *Decne. in Rev. Hortic.* Sér. IV. iii. (1854) 181.—N. Granat.
sclerophylla, *Planch. & Linden, Fl. des Serres*, viii. (1852–53) 205. t. 825.—N. Granat.
tovarensis, *Klotzsch, in Linnaea*, xxiv. (1851) 44.—N. Granat.

PSAMMOGETON, Edgew. in Proc. Linn. Soc. i. (1845) 253; et in Trans. Linn. Soc. xx. (1846) 57. *UMBELLIFERAE*, Benth. & Hook. f. i. 929.
biternatum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 57.—Persia; Ind. bor. occ.
brevisetum, *Boiss. Fl. Orient.* ii. 1079.—Persia.
crinitum, Boiss. l. c. 1078 = biternatum.
setifolium, *Boiss. l. c.* 1079.—Persia; Afghan.
triternatum, Benth. & Hook. f. Gen. i. 929, sphalm. = biternatum.

PSAMMOPHILA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 345 = **Gypsophila**, Linn. (Caryophyll.).
muralis, Fourr. l. c. = G. muralis.

PSAMMOSERIS, Boiss. & Reut. Diagn. Ser. I. xi. 52 (1849) = **Crepis**, Vaill. (Compos.).
arabica, Boiss. & Reut. l. c. = C. arabica.
senecioides, Boiss. & Reut. l. c. = C. senecioides.

PSAMMOSTACHYS, Presl (Bot. Bemerk. 91 (1844)), in Abh. Boehm. Ges. V. iii. (1845) 521 = **Striga**, Lour. (Scrophul.).
varia, Presl, ll. cc. = Harveya varia.

PSAMMOTROPHA, Eckl. & Zeyh. Enum. 286 (1836). *FICOIDEAE*, Benth. & Hook. f. i. 858.
MALLOGONUM, Reichb. Handb. 236 (1837).
androsacea, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 265. —Afr. austr.
mucronata, *Fenzl, l. c.* 267.—Afr. austr.
myriantha, *Sond. in Harv. & Sond. Fl. Cap.* i. 147.—Afr. trop.
parvifolia, *Eckl. & Zeyh. Enum.* 287.—Afr. austr.
quadrangularis, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 264.—Afr. austr.
rigida, *Fenzl, l. c.* 266.—Afr. austr.

PSAMMOTROPHE, Benth. & Hook. f. Gen. ii. 1023 (Index) (1867) = praec.

PSANACETUM, Neck. Elem. i. 89 (1790) = **Tanacetum**, Tourn. (Compos.).
annuum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 89 = T. annuum.

PSANCHUM, Neck. Elem. i. 254 (1790) = **Cynanchum**, Linn. (Asclep.).

PSATHERIPS, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
fragilis, Rafin. l. c. = S. fragilis.

PSATHURA, Comm. ex Juss. Gen. 206 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 132.
CHICOINAEA, Comm. ex DC. Prod. iv. 462 (1830).
borbonica, *J. F. Gmel. Syst.* 577.—Ins. Borbon.
corymbosa, Gaertn. f. Fruct. iii. 82. t. 194. f. 4 = borbonica.
myrtifolia, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 214.—Ins. Maurit.
obovata, *Tussac, Fl. Antill.* ii. 78. t. 18.—Ins. S. Doming.
sechellarum, *Baker, Fl. Maurit.* 157.—Ins. Seychelles.
terniflora, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 214.—Ins. Maurit.

PSATHUROCHAETA, DC. Prod. v. 609 (1836) = **Melanthera**, Rohr (Compos.).
Dregei, DC. l. c.—Afr. austr.

PSATHYRA, Spreng. Anleit. ii. II. 614 (1818) = **Psathura**, Comm. (Rubiac.).

PSATHYROTES, A. Gray, Pl. Wright. ii. 100. t. 13; in Smithson. Contrib. v. (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 415.
annua, *A. Gray, Pl. Wright.* ii. 100. t. 13.—Mexic.
incisa, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 322.—Calif.
pilifera, *A. Gray, in Proc. Am. Acad.* xix. (1883) 50.—Utah.
ramosissima, *A. Gray, l. c.* vii. (1868) 363.—Calif.
scaposa, *A. Gray, Pl. Wright.* ii. 100.—Mexic.
Schottii, A. Gray, in Proc. Am. Acad. ix. (1874) 206 = Peucephyllum Schottii.

PSATURA, Poir. Encyc. vi. 582 (1804) = **Psathura**, Comm. (Rubiac.).

PSEDERA, Neck. Elem. i. 158 (1790) = **Hedera**, Tourn. (Araliac.).

PSEDOMELIA, Neck. Elem. iii. 150 (1790) = **Bromelia**, Linn.

PSELIUM, Lour. Fl. Cochinch. 622 (1790), partim = **Pericampylus**, Miers (Menisp.).
ambiguum, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 372.—Cochinch.
heterophyllum, Lour. Fl. Cochinch. 621.—Cochinch.

PSEPHELLUS, Cass. in Dict. Sc. Nat. xliii. 488 (1826) = **Centaurea**, Linn. (Compos.).
Aucherianus, Boiss. Fl. Orient. iii. 611 = C. Aucheriana.
calocephalus, Cass. in Dict. Sc. Nat. xliii. 488 = C. dealbata.
carthamoides, Fisch. & Mey. ex Boiss. Diagn. Ser. I. vi. 125 = C. carthamoides.

PSEPHELLUS :—
dealbatus, C. Koch, in Linnaea, xxiv. (1851) 438 = Centaurea dealbata.
declinatus, C. Koch, l. c. = C. leucophylla.
decumbens, C. Koch, l. c. = C. leucophylla.
heterophyllus, Boiss. & Haussk. Diagn. Ser. II. iii. 65 = C. Spruneri ?
hypoleucus, Boiss. Fl. Orient. iii. 610 = C. hypoleuca.
incanescens, Boiss. l. c. = C. integrifolia.
integrifolius, C. Koch, in Linnaea, xxiv. (1851) 438 = C. integrifolia.
kacheticus, Rehm. ex Boiss. Fl. Orient. iii. 608.—Reg. Caucas.
lazicus, Boiss. l. c. = C. dealbata.
leptophyllus, Boiss. l. c. 611 = C. leptophylla.
leucophyllus, C. A. Mey. Verz. Pfl. Cauc. 66 = C. leucophylla.
Marschallianus, C. Koch, in Linnaea, xxiv. (1851) 438 = C. Marschalliana.
pauciloba, Boiss. Fl. Orient. iii. 609 = C. pauciloba.
phaeopappus, Fisch. & Mey. ex DC. Prod. vi. 560 = C. leuzeoides.
salviaefolius, Boiss. Fl. Orient. iii. 609.—Reg. Caucas.
spectabilis, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 43 (= *Amberboa spectabilis*) = C. Balsamita ?
ustulatus, Fisch. & Mey. ex DC. Prod. vi. 576 = C. ustulata.
xanthocephalus, Fisch. & Mey. ex Boiss. Fl. Orient. iii. 610 = C. xanthocephala.

PSEUDACACIA, Moench, Meth. 145 (1794) = PSEUDOACACIA, Tourn. = **Robinia**, Linn. (Legumin.).

PSEUDAEGLE, Miq. Ann. Mus. Lugd. Bat. ii. 83 (1865–66) = **Citrus**, Linn. (Rutac.).
sepiaria, Miq. l. c. = C. trifoliata.

PSEUDAIS, Decne. in Ann. Sc. Nat. Sér. II. xix. (1843) 40. *THYMELAEACEAE*, Benth. & Hook. f. iii. 200.
coccinea, *Decne. l. c.*—Ins. Molucc.

PSEUDALANGIUM, F. Muell. Fragm. ii. 84 (1860) = **Marlea**, Roxb. (Cornac.).
polyosmoides, F. Muell. l. c. = M. vitiensis.

PSEUDALEIA, Thou. Gen. Nov. Madag. 15 (1806) = **Olax**, Linn.
madagascariensis, DC. Prod. i. 533 = O. Pseudaleia.
Thouarsii, Steud. Nom. ed. II. ii. 208, in syn. = O. Pseudaleioides.

PSEUDALEIOIDES, Thou. Gen. Nov. Madag. 15 (1806) = **Olax**, Linn.
Thouarsii, DC. Prod. i. 533 = O. Pseudaleioides.

PSEUDALOMIA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, i. (1844) 484, nomen (Compos.).

PSEUDANTHUS, Sieber, ex Spreng. Syst. iv. Cur. Post. 25 (1827). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 262.
CHORIZOTHECA, Muell. Arg. in Linnaea, xxxii. (1863) 76.
CHRYSOSTEMON, Klotzsch, in Lehm. Pl. Preiss. ii. 232 (1847).
brachyphyllus, *F. Muell. in Trans. Roy. Soc. N. S. Wales* (1881), 11.—Austral.
chrysus, Muell. Arg. in Flora, xlvii. (1864) 486 = Stachystemon polyandrus.
divaricatissimus, *Benth. Fl. Austral.* vi. 60.—Austral.
micranthus, *Benth. l. c.* 59.—Austral.
nematophorus, *F. Muell. Fragm.* ii. 14.—Austral.
nitidus, Muell. Arg. in DC. Prod. xv. II. 197 = virgatus.
occidentalis, F. Muell. Fragm. i. 107 = virgatus.
orientalis, *F. Muell. l. c.* ii. 14.—Austral.
ovalifolius, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 66.—Austral.
pimeleoides, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 25. —Austral.
polyandrus, F. Muell. Fragm. ii. 153 = Stachystemon polyandrus.
vermicularis, *F. Muell. in Trans. Roy. Soc. N. S. Wales* (1881), 11.—Austral.
virgatus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 56.—Austral.

PSEUDANTHUS, Wight, Ic. v. 3; vi. t. 1776 bis (1852) = **Nothosaerva**, Wight (Amarant.).
brachiatus, Wight, l. c. = N. brachiata.

PSEUDARTHRIA, Wight & Arn. Prod. 209 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 521.
ANARTHROSYNE, E. Mey. Comm. Pl. Afr. Austr. 124 (1835).
capitata, Hassk. Cat. Hort. Bog. Alt. 281; Pl. Jav. Rar. 390 = Desmodium capitatum.
confertiflora, *Baker, in Oliver, Fl. Trop. Afr.* ii. 167. —Afr. trop.
cordata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 39. t. 7.—Afr. trop.
densiflora, *Klotzsch, l. c.* 41.—Afr. trop.
fagifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 167.—Afr. trop.
gracilis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 41. —Afr. trop.
gyrans, Hassk. in Hoev. & De Vriese, Tijdschr. xi. (1844) 104 = Desmodium gyrans.
gyroides, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, iii. (1846) 63, nomen.—Java.
Hookeri, *Wight & Arn. Prod.* 209.—Ins. Maurit.
macrophylla, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 168.—Afr. trop.
pilosa, Wight & Arn. Prod. 209 = Rhynchosia pilosa, *Wall.*
polycarpa, Hassk. Cat. Hort. Bog. Alt. 282; Pl. Jav. Rar. 393 = Desmodium gyroides.
timoriensis, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, iii. (1846) 63, nomen.—Java.
viscida, *Wight & Arn. Prod.* 209.—Ind. or.; Java.

PSEUDATALAYA, Baill. Hist. Pl. v. 419 (1874) = **Atalaya**, Blume (Sapindac.).
multiflora, Baill. l. c. = A. multiflora.

PSEUDEHRETIA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 607 = **Ilex**, Tourn.
paniculata, Turcz. l. c.—Java.
umbellata, Turcz. l. c. = I. Godajam.

PSEUDELEPHANTOPUS, Rohr, in Skrift. Nat. Selsk. Kiobenh. ii. (1792) 214 = **Elephantopus**, Linn. (Compos.).
spicatus, Rohr, l. c. 216 = E. spicatus.

PSEUDEPIDENDRUM, Reichb. f. in Bot. Zeit. x. (1852) 733 = **Epidendrum**, Linn. (Orchid.).
spectabile, Reichb. f. l. c. = E. Pseudepidendrum.

PSEUDERIOPSIS, Reichb. f. in Linnaea, xxii. (1849) 852 = **Eriopsis**, Lindl. (Orchid.).
Schomburgkii, Reichb. f. l. c. 853 = E. Schomburgkii.

PSEUDEVAX, DC. ex Steud. Nom. ed. II. i. 620, in syn. (1840); ii. 405 (1841) = **Evax**, Gaertn. (Compos.).

PSEUDIMA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 300, 309; ix. (1879) 495, 515 = **Sapindus**, Tourn.
frutescens, Radlk. ll. cc. = S. frutescens.

PSEUDIOSMA, A. Juss. in Mém. Mus. Par. xii. (1825) 519 = **Zanthoxylum**, Linn. *vel* **Evodia**, Forst. (Rutac.).
asiatica, G. Don, Gen. Syst. i. 807.—Cochinch.

PSEUDITEA, Hassk. in Flora, xxv. (1842) II. Beibl. 30; et Cat. Hort. Bog. Alt. 160 (1844) = **Pittosporum**, Banks.
javanica, Hassk. ll. cc. = Pittosp. floribundum ?

PSEUDIXORA, Miq. Fl. Ind. Bat. ii. 209 (1856) = **Randia**, Houst. (Rubiac.).
auriculata, Miq. l. c. 210 = Anomanthodia auriculata.
javanica, Miq. l. c. = R. longiflora.
sumatrana, Miq. l. c. 209 = R. sumatrana.
truncata, Miq. l. c. 210 = Trisciadia truncata.
Zollingeriana, Miq. l. c. = R. longiflora.

PSEUDO-ACACIA, [Tourn.] Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 364 = **Robinia**, Linn. (Legumin.).
Halodendron, Moench, Meth. 145 = Halimodendron argenteum.
hispida, Moench, l. c. = R. hispida.
odorata, Moench, l. c. = R. Pseud-acacia.
vulgaris, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 364 = R. Pseud-acacia.

PSEUDOBARLERIA, T. Anders. in Journ. Linn. Soc. vii. (1864) 26 = **Petalidium**, Nees (Acanth.).
hirsuta, T. Anders. l. c. = Pet. Currori (in Addend.).

PSEUDOBARLERIA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 135 = **Barleria**, Linn. (Acanth.).
coerulea, Oerst. l. c. = B. strigosa.
hirsuta, Oerst. l. c. = B. polytricha.
polystachya, Oerst. l. c. = B. strigosa.
polytricha, Oerst. l. c. = B. polytricha.
Richardiana, Oerst. l. c. = B. Richardiana.

PSEUDOBASILICUM, Steud. Nom. ed. II. ii. 405 (1841) = PSEUDOBRASILIUM, Plum.

PSEUDOBESLERIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 128 (Gesn. Centr. Am. 54) = **Besleria**, Linn. (Acanth.).
umbrosa, Oerst. ll. cc. = B. umbrosa.

PSEUDOBRASILIUM, Plum. ex Adans. Fam. ii. 341 (1763).—Cf. PSEUDOBASILICUM, Steud. Nom. ed. II. ii. 405 (1841) (Quid ?).
glabrum, Lam. in Mém. Acad. Sc. Par. (1784) 345.—Hab. ?
hirsutum, Lam. l. c. 349.—Hab. ?

PSEUDOCALYX, Radlk. in Bremen, Abh. viii. (1883) 416. *ACANTHACEAE.*
saccatus, *Radlk. l. c.*—Afr. trop.

PSEUDOCAPSICUM, Medic. Phil. Bot. i. 122 (1789) = **Solanum**, Tourn.
diphyllum, Medic. l. c. = S. diphyllum.
lancifolium, Moench, Meth. Suppl. 180 = S. lanceolatum.
torulosum, Moench, Meth. 477 = S. aethiopicum.
undulatum, Moench, l. c. = S. Pseudocapsicum.

PSEUDOCARAPA, Hemsl. in Hook. Ic. Pl. t. 1458 (1884). *MELIACEAE.*
Championii, *Hemsl. l. c.*—Zeylan.

PSEUDOCAREX, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 146 (1865–66) = **Carex**, Dill. (Cyperac.).
plantaginea, Miq. l. c.—Japon.

PSEUDOCARYOPHYLLUS, Berg, in Linnaea, xxvii. (1854) 415 = **Myrtus**, Tourn.
coccolobaefolius, Berg, l. c. 416 = Myrcia coccolobaefolia.
costatus, Berg, in Mart. Fl. Bras. xiv. i. 607 = Myrcia costata.
fulvescens, Berg, in Linnaea, xxvii. (1854) 416 = Eugenia fulvescens.
heteroclitus, Berg, l. c. = Eugenia heteroclita.
Seemanni, Triana, ex Hemsl. Biol. Centr. Am. Bot. i. 407 = Myrtus Arayan.
sericeus, Berg, in Linnaea, xxvii. (1854) 415 = Eugenia Pseudocaryophyllus.
velutinum, Berg, in Mart. Fl. Bras. xiv. i. 607.—Bras.

PSEUDOCENTRUM, Lindl. in Journ. Linn. Soc. iii. (1859) 63, 64. *ORCHIDEAE*, Benth. & Hook. f. iii. 594.
bursarium, *Reichb. f. in Linnaea*, xli. (1877) 53.—Peruv.
Hoffmanni, *Reichb. f. l. c.*—Costa Rica.
macrostachyum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 64.—Peruv.
minus, *Benth. in Hook. Ic. Pl.* t. 1382.—Jamaic.

PSEUDO-CHAENOMELES, Carr. in Rev. Hortic. (1882) 238 = **Pyrus**, Tourn. (Rosac.).
Maulei, Carr. l. c. = Pyrus japonica.

PSEUDOCHROSIA, Blume, Mus. Bot. Lugd. Bat. i. 158 (1850). *APOCYNACEAE*, Benth. & Hook. f. ii. 700.
glomerata, *Blume, l. c.*—N. Guin.

PSEUDOCROTON, Muell. Arg. in Flora, lv. (1872) 24. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 304.
tinctorius, *Muell. Arg. l. c.*—Guatemala.

PSEUDOCYPERUS, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Cyp. 108 (1855) = **Fimbristylis**, Vahl (Cyperac.).
scirpinus, Steud. ll. cc. = F. bahiensis.

PSEUDODICTAMNUS, Tourn. ex Adans. Fam. ii. 191 (1763) = **Ballota**, Linn. (Labiat.).
acutus, Moench, Meth. 400 = B. hirsuta.
emarginatus, Moench, Meth. Suppl. 139 = B. africana.

PSEUDODRACONTIUM, N. E. Br. in Journ. Bot. xx. (1881) 193. t. 231. f. 2. *AROIDEAE*, Benth. & Hook. f. iii. 971.
anomalum, *N. E. Br. l. c.*—Cochinch.
Lacourii, *N. E. Br. l. c.* 194.—Cochinch.

PSEUDOELEPHANTOPUS, Steud. Nom. ed. II. ii. 405 = PSEUDELEPHANTOPUS, Rohr = **Elephantopus**, Linn. (Compos.).

PSEUDOEUGENIA, Scortech. in Journ. Bot. xxiii. (1885) 153. *MYRTACEAE.*
perakiana, *Scortech. l. c.*—Penins. Malay.

PSEUDOFUMARIA, Medic. Phil. Bot. i. 110 (1789) = **Corydalis**, Vent. (Papaver.).
capnoides, Borck. in Roem. Arch. i. ii. (1797) 45 = C. lutea.
intermedia, Borck. l. c. = C. fabacea.
lutea, Medic. Phil. Bot. i. 110 = C. lutea.
major, Borck. in Roem. Arch. i. ii. (1797) 45 = C. bulbosa.
minor, Borck. l. c. = C. bulbosa.
Schangini, Borck. l. c. = C. longiflora.
sibirica, Borck. l. c. = C. sibirica.

PSEUDO-GUNNERA, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 193 = **Gunnera**, Linn. (Halor.).
macrophylla, Oerst. l. c. = G. macrophylla.

PSEUDO-IRIS, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 417 = **Iris**, Tourn.
palustris, Medic. l. c. = I. Pseudacorus.

PSEUDOLARIX, Gord. Pinet. 292 (1858). *CONIFERAE* [Benth. & Hook. f. iii. 442].
Kaempferi, *Gord. l. c.*—China.

PSEUDOLMEDIA, Karst. Fl. Columb. ii. 21. t. 111 (1862 ?) = **Olmedia**, Ruiz & Pav. (Urtic.).
coriacea, Karst. l. c.—N. Granat.

PSEUDOLMEDIA, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 129. t. 5. *URTICACEAE*, Benth. & Hook. f. iii. 372.
OLMEDIOPSIS, Karst. Fl. Columb. ii. 17. t. 109 (1862 ?).
bucidaefolia, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 109.—Porto Rico.
ferruginea, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 131.—Peruv.
havanensis, *Tréc. l. c.* 130.—Cuba.
hirsuta, *Baill. Adansonia*, xi. (1873–76) 295.—Borneo.
laevigata, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 131.—Peruv.
macrophylla, *Tréc. l. c.* 132.—Peruv.
spuria, Griseb. Fl. Brit. W. Ind. 152 = havanensis.

PSEUDOLYSIMACHION, Opiz, Seznam, 80 (1852) = **Veronica**, Tourn. (Scrophul.).
commutatum, Opiz, l. c. = V. commutata.
cristatum, Opiz, in Lotos, iv. (1854) 45 = V. orchidea.
Kosteleckii, Opiz, Seznam, 80 = V. longifolia.
longifolium, Opiz, l. c. = V. longifolia.
Nenningii, Opiz, l. c. = V. longifolia.
Pettersii, Opiz, l. c. = V. longifolia.
riparium, Opiz, l. c. = V. longifolia.
spicatum, Opiz, l. c. = V. spicata.

PSEUDOMORUS, Bur. in Ann. Sc. Nat. Sér. V. xi. (1869) 371. *URTICACEAE*, Benth. & Hook. f. iii. 364.
Brunoniana, *Bur. l. c.* 372.—N. Caled.

PSEUDONEPHELIUM, Radlk. Sapind. Holland-Ind. Nachtr. 71 (1879) = **Nephelium**, Linn. (Sapind.).
fumatum, Radlk. l. c. = N. fumatum.

PSEUDOPANAX, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 366, partim. *ARALIACEAE*, Benth. & Hook. f. i. 941.
anomalum, C. Koch, l. c. = Panax anomalum.
arboreum, C. Koch, l. c. = Panax arboreum.
Colensoi, C. Koch, l. c. = Panax Colensoi.
crassifolium, *C. Koch, l. c.*—N. Zel.
Edgerleyi, C. Koch, l. c. = Panax Edgerleyi.
Gaudichaudii, C. Koch, l. c. = Panax Gaudichaudii.
Gunnii, C. Koch, l. c. = Panax Gunnii.
Lessonii, *C. Koch, l. c.*—N. Zel.
lineare, C. Koch, l. c. = Panax lineare.
ovatum, C. Koch, l. c. = Panax Gaudichaudii.
platyphyllum, C. Koch, l. c. = Panax platyphyllum.
simplex, C. Koch, l. c. = Panax simplex.

PSEUDOPETALON, Rafin. Fl. Ludov. 107 (1817) = **Zanthoxylum**, Linn. (Rutac.).
glandulosum, Rafin. l. c. 108 = Z. Clava-Herculis.
tricarpum, Rafin. Med. Fl. ii. 114 = Z. Clava-Herculis.

PSEUDOPHOLIDIA, A. DC. in DC. Prod. xi. 704 (1847) = **Pholidia**, R. Br. (Myopor.).
brevifolia, A. DC. l. c. = Phol. brevifolia.

PSEUDOPTERIS, Baill. Adansonia, xi. (1874) 243. *SAPINDACEAE.*
decipiens, *Baill. l. c.*—Madag.

PSEUDOPYXIS, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 189 (1867). *RUBIACEAE*, Benth. & Hook. f. ii. 135.
depressa, *Miq. l. c.*—Japon.
heterophylla, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 175.—Japon.
longituba, *Franch. & Sav. Enum. Pl. Jap.* ii. 391.—Japon.

PSEUDORAPHIS, Griff. Notul. iii. 29 (1851) = **Setaria**, Beauv. (Gramin.).
Brunoniana, Griff. l. c. = S. glauca.

PSEUDORCHIS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 213 (1821) = **Liparis**, Rich. (Orchid.).
Loeselii, S. F. Gray, l. c. = L. Loeselii.

PSEUDORYZA, Griff. Ic. Pl. As. Rar. t. 144 (1847) = **Leersia**, Sw. (Gramin.).
ciliata, Griff. l. c.; et Notul. iii. 1 = L. hexandra.

PSEUDORHACHICALLIS, Benth. & Hook. f. Gen. ii. 60 (1873) = PSEUDRACHICALLIS, Karst. = **Mallostoma**, Karst. (Rubiac.).

PSEUDOSANTALUM, Mill. Gard. Dict. ed. VIII. in syn. (1768) = **Caesalpinia**, Linn. (Legumin.).

PSEUDOSCIADIUM, Baill. Adansonia, xii. (1878) 130. *ARALIACEAE.*
Balansae, *Baill. l. c.*—N. Caled.

PSEUDOSCORDUM, Herb. Amaryll. 11 (1837) = **Nothoscordum**, Kunth (Liliac.).

PSEUDOSERIS, Baill. in Bull. Soc. Linn. Par. i. (1881) 282. *COMPOSITAE.*
Grandidieri, *Baill. l. c.*—Madag.
Rutenbergi, *Baill. l. c.*—Madag.

PSEUDOSMODINGIUM, Engl. Bot. Jahrb. i. (1881) 419. *ANACARDIACEAE.*
Andrieuxii, *Engl. l. c.* 420.—Mexic.
perniciosum, *Engl. l. c.*—Mexic.
Virletii, *Engl. l. c.*—Mexic.

PSEUDOSOPHORA, Sweet, Hort. Brit. ed. II. 122 (1830) = **Sophora**, Linn. (Legumin.).
alopecuroides, Sweet, l. c. = S. alopecuroides.
sericea, Sweet, l. c. = S. sericea.

PSEUDOSOROCEA, Baill. Adansonia, xi. (1875) 296 = **Sorocea**, A. St. Hil. (Urtic.).
Bonplandi, Baill. l. c.—Guian.

PSEUDOSOROCEA :—
muriculata, Benth. & Hook. f. Gen. iii. 378 = Sorocea muriculata.
Poeppigii, Baill. l. c. 297.—Bras.
Sprucei, Baill. l. c. 296.—Peruv.
uaupensis, Baill. l. c. 297.—Bras.

PSEUDOSPERMUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 517 (1821) = **Physospermum**, Cuss. (Umbell.).
commutatum, S. F. Gray, l. c. = Phys. commutatum.

PSEUDOSPONDIAS, Engl. in DC. Monog. Phan. iv. (1883) 258. *ANACARDIACEAE.*
microcarpa, *Engl. l. c.* 259.—Afr. trop.

PSEUDOSTACHYUM, Munro, in Trans. Linn. Soc. xxvi. (1868) 141. t. 4. *GRAMINEAE*, Benth. & Hook. f. iii. 1212.
compactiflorum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 252 = Melocalamus compactiflorus.
Helferi, *Kurz, l. c.* 253.—Burma.
polymorphum, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 142.—Ind. or.

PSEUDOSTREBLUS, Bur. in DC. Prod. xvii. 219 (1873). *URTICACEAE*, Benth. & Hook. f. iii. 357.
indica, *Bur. l. c.* 220.—Ind. or.

PSEUDOTREWIA, Miq. Fl. Ind. Bat. i. II. 414 (1859) = **Alchornea**, Sw. (Euphorb.).
cuneifolia, Miq. l. c. Suppl. 462 = A. cuneifolia.
macrophylla, Miq. l. c. i. II. 414 = Trewia macrophylla.

PSEUDOTSUGA, Carr. Conif. ed. II. 256 (1867). *CONIFERAE*, Benth. & Hook. f. iii. 441.
Davidiana, Carr. in Rev. Hortic. (1873) 38 = Abies Davidiana.
Douglasii, *Carr. Conif.* ed. II. 256.—Am. bor. occ.
Fortunei, W. R. McNab, in Proc. Roy. Irish Acad. Ser. II. ii. (1877) sub t. 49 = Keteleeria Fortunei.
glaucescens, Carr. Conif. ed. II. 274 = Abies religiosa.
jezoënsis, W. R. McNab, in Prod. Roy. Irish Acad. Ser. II. ii. (1877) sub t. 49 = Keteleeria Fortunei.
Lindleyana, Carr. in Rev. Hortic. (1868) 152 = Douglasii.
magnifica, W. R. McNab, in Proc. Roy. Irish Acad. Ser. II. ii. (1877) sub t. 49 = Abies magnifica.
nobilis, W. R. McNab, l. c. = Abies nobilis.

PSEUDRACHICALLIS, Karst. Fl. Columb. ii. 10 (1862) = **Mallostoma**, Karst. (Rubiac.).
floribunda, Karst. l. c. = M. caracasanum?

PSEUDUVARIA, Miq. Fl. Ind. Bat. i. II. 82 (1859) = **Mitrephora**, Blume (Anonac.).
Diepenhorstii, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 38 = M. Diepenhorstii.
reticulata, Miq. Fl. Ind. Bat. i. II. 32 = M. reticulata.

PSEVA, Rafin. in Journ. Phys. lxxxix. (1819) 261 = **Chimaphila**, Pursh (Ericac.).

PSIADIA, Jacq. Hort. Schoenb. ii. 13. t. 152 (1797). *COMPOSITAE*, Benth. & Hook. f. ii. 284.
ALIX, Comm. ex DC. Prod. v. 318 (1836).
ELPHEGEA, Cass. in Bull. Soc. Philom. (1818) 31; et in Dict. Sc. Nat. xiv. 361 (1819).
FRAPPIERIA, Cordem. in Adansonia, x. (1871) 24.
GLUTINARIA, Comm. ex DC. Prod. v. 318 (1836).
SARCANTHEMUM, Cass. in Bull. Soc. Philom. (1818) 74.
THOUARSIA, Vent. ex DC. Prod. v. 318 (1836).
altissima, *Benth. & Hook. f. Gen.* ii. 285.—Madag.
arabica, *Jaub. & Spach, Illustr.* iv. 85. tt. 352, 353.—Arab.; Afr. trop.
arguta, Voigt, Hort. Suburb. Calc. 408 = trinervia.
auriculata, *Baker, in Journ. Bot.* xx. (1882) 169.—Madag.
balsamica, DC. in Wight, Contrib. 10 = trinervia.
Coronopus, *Hook. f. & Benth. ex Balf. f. in Phil. Trans.* clxviii. (1879) 350.—Ins. Rodrig.
dentata, DC. Prod. v. 319 = trinervia.
dodoneaefolia, *Steetz, in Peters, Reise Mossamb. Bot.* 385.—Afr. trop.
Francavillea, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 365.—Ins. Borbon.
glauca, Ayres, ex Baker, Fl. Maurit. 171 = trinervia.
glutinosa, *Jacq. Hort. Schoenb.* ii. t. 152.—Ins. Maurit.
grandidentata, *Steetz, in Peters, Reise Mossamb. Bot.* 387.—Afr. trop.
hispida, *Benth. & Hook. f. Gen.* ii. 285.—Madag.
incana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 320.—Afr. trop.
integerrima, *DC. in Wight, Contrib.* 10.—Ins. Maurit.
integrifolia, Steud. Nom. ed. II. i. 179 = praec.
linearifolia, DC. Prod. v. 319 = trinervia.
madagascariensis, *DC. l. c.* 320.—Madag.
minor, *Steetz, in Peters, Reise Mossamb. Bot.* 391.—Afr. trop.
penninervia, *DC. Prod.* v. 319.—Ins. Maurit.
punctulata, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 196 = arabica.
quinquenervia, Steetz, in Peters, Reise Mossamb. Bot. 392 = trinervia.
resiniflora, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 286 = arabica.
retusa, *DC. Prod.* v. 319.—Ins. Borbon.
rodriguesiana, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 15; *et in Phil. Trans.* clxviii. (1879) 350.—Ins. Rodrig.
rotundifolia, *Hook. f. in Melliss, S. Helena*, 228, 286. t. 41.—Ins. S. Helena.
salviaefolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 181.—Madag.
Schweinfurthii, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 837.—Ins. Socotra.
trinervia, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 139.—Afr. trop.; Ins. Mascar.
urticaefolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 181.—Madag.
vernicosa, *Steetz, in Peters, Reise Mossamb. Bot.* 390.—Afr. trop.
volubilis, Baill. Hist. Pl. viii. 300 = Microglossa volubilis.

PSIDIASTRUM, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 272. *MYRTACEAE.*
dubium, *Bello, l. c.*—Porto Rico.

PSIDIOPSIS, Berg, in Linnaea, xxvii. (1854) 350. *MYRTACEAE*, Benth. & Hook. f. i. 713.
Moritzianum, *Berg, l. c.* 351.—Venezuela.

PSIDIUM, Linn. Gen. ed. I (1737) 140. *MYRTACEAE*, Benth. & Hook. f. i. 713.
ACCA, Berg, in Linnaea, xxvii. (1853–55) 138.
CALYTROPSIDIUM, Berg, l. c. (1853–55) 349.
GUAIAVA, Tourn. ex Adans. Fam. ii. 88 (1763).
GUAYABA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 16.
acidum, Mart. ex Berg, in Linnaea, xxvii. (1854) 436 = acutangulum.
acre, *Tenore, Ind. Sem. Hort. Neap.* (1829) 17.—Hab.?
acutangulum, *DC. Prod.* iii. 233.—Bras.
adamantium, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 292 = Campomanesia adamantium.
aerugineum, *Berg, in Mart. Fl. Bras.* xiv. I. 391.—Am. trop.
alatum, *Berg, l. c.* 604.—Bras.
albidum, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 283.—Bras.
albidum, Miq. in Linnaea, xxii. (1849) 532 = grandifolium.
amplexicaule, Pers. Syn. ii. 27 = montanum.
amygdalinum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 317.—Reg. Argent.
anceps, *Berg, in Mart. Fl. Bras.* xiv. I. 395.—Bras.
angustifolium, Lam. Encyc. iii. 16 = P. Guajava.
anthomega, *Vell. Fl. Flum.* 212; v. t. 51.—Bras.
apricum, Vell. ll. cc. 213; v. t. 58 = Campomanesia aprica.
Apysa, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 66; *Contrib.* 116.—Reg. Argent.
aquaticum, *Benth. in Hook. Journ. Bot.* ii. (1840) 318.—Guian.
Araça, *Raddi, in Opusc. Sc.* iv. (1823) 252.—Bras.
Arasa-hu, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 65; *Contrib.* 115.—Reg. Argent.
Arasa-pe, *Parodi, ll. cc.*—Reg. Argent.
Arasope-mi, *Parodi, ll. cc.*—Reg. Argent.
arboreum, *Vell. Fl. Flum.* 211; v. t. 50.—Bras.
argentum, *Berg, in Mart. Fl. Bras.* xiv. I. 388.—Bras.
aromaticum, Aubl. Pl. Gui. i. 485. t. 191 = Campomanesia aromatica.
aromaticum, Blanco, Fl. Filip. ed. I. 417 = P. Guajava.
aromaticum, Descourt. Fl. Med. Antill. v. 229. t. 364 = montanum.
aromaticum, D. Don, ex Berg, in Mart. Fl. Bras. xiv. I. 522 = Donianum.
australe, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 283.—Bras.
basanthum, *Berg, in Mart. Fl. Bras.* xiv. I. 601.—Bras.
Benthamianum, *Berg, in Linnaea*, xxvii. (1854) 362.—Guian.
Berterianum, *Berg, l. c.* 374.—Porto Rico.
Brownianum, *Mart. ex DC. Prod.* iii. 236.—Bras.
buxifolium, *Nutt. N. Am. Sylva*, i. 115. t. 25.—Am. bor.
calycolpoides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 183.—Cuba.
campestre, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 289 = Campomanesia Cambessedeana.
campomanisioides, *Berg, in Mart. Fl. Bras.* xiv. I. 520.—Bras,
caninum, *Lour. Fl. Cochinch.* 310 (sp. dub.).—China.
caninum, Mart. ex DC. Prod. iii. 235 = Campomanesia Guaviroba.
Cattleianum, [*Sabine,*] *in Trans. Hort. Soc.* iv. (1822) 317. t. 11.—Bras.
Cattleyanum, Weinw. in Syll. Ratisb. ii. (1828) 166 = humile.
cerasoides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 290.—Bras.
chinense, Lodd. ex Loud. Hort. Brit. 197, nomen (Quid?).—China.
chrysophyllum, *F. Muell. Sel. Pl. Indust. Cult.* 188.—Bras.
ciliatum, *Benth. in Hook. Journ. Bot.* ii. (1840) 318.—Guian.
cinereum, *Mart. ex DC. Prod.* iii. 234.—Bras.
cordatum, Sims, Bot. Mag. t. 1779 = montanum.
coriaceum, Mart. ex DC. Prod. iii. 236 = humile.
corymbosum, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 286 = Campomanesia corymbosa.
costa-ricense, *Berg, in Linnaea*, xxvii. (1854) 368.—Costa Rica.
crenatum, *Berg, l. c.* 373.—Am. austr.
Cujavillus, Burm. f. Fl. Ind. 114 = P. Guajava.
Cujavus, Linn. Syst. ed. X. 1055 = P. Guajava.
cuneatum, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 285.—Bras.
cuneifolium, *Tenore, Ind. Sem. Hort. Neap.* (1833) 14.—Hab.?
cupreum, *Berg, in Mart. Fl. Bras.* xiv. I. 393.—Bras.
decaspermum, Linn. f. Suppl. 252 = Timonius Jambosella.
decussatum, *DC. Prod.* iii. 235.—Bras.
densicomum, *Mart. ex DC. l. c.*—Bras.
desertorum, Mart. ex DC. l. c. 236 = Campomanesia repanda.
dichotomum, Weinm. in Syll. Ratisb. ii. (1828) 166 = P. Araça.
Donianum, *Berg, in Mart. Fl. Bras.* xiv. I. 521.—Bras.
dubium, *H. B. & K. Nov. Gen. et Sp.* vi. 152.—Venezuela.
dulce, *Vell. Fl. Flum.* 213; v. t. 56.—Bras.
elegans, Mart. ex DC. Prod. iii. 240 = Myrtus elegans.
elegans, *Miq. Fl. Ind. Bat.* i. I. 470.—Ins. Molucc.
emarginatum, *Ruiz & Pav. Fl. Per.* iv. t. 419.—Peruv.
erianthum, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 279.—Bras.
erosum, Miq. in Linnaea, xxii. (1849) 532 = Campomanesia fusca.
eugenioides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 290.—Bras.
eugenioides, Miq. in Linnaea, xxii. (1849) 533 = Campomanesia xanthocarpa.
ferrugineum, *Presl, in Oken, Isis*, xxi. (1828) 274.—Ins. Maurit.
firmum, *Berg, in Mart. Fl. Bras.* xiv. I. 390.—Bras.
fluviatile, *Rich. ex DC. Prod.* iii. 235.—Guian.
fragrans, Macfad. Fl. Jamaic. ii. 108 = P. Guajava.
fruticosum, *Vell. Fl. Flum.* 213; v. t. 57.—Bras.
galapagaeum, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 224.—Ins. Galapag.

PSIDIUM:—
Gardnerianum, *Berg, in Mart. Fl. Bras.* xiv. i. 389.—Bras.
glaucescens, *Berg, l. c.* 600.—Bras.
grandiflorum, Aubl. Pl. Gui. i. 483 = Campomanesia aromatica.
grandiflorum, *Ruiz & Pav. Fl. Per.* iv. t. 421.—Peruv.
grandifolium, *Mart. ex DC. Prod.* iii. 234.—Bras.
Guajava, *Linn. Sp. Pl.* 470.—Am. trop.
Guaviroba, DC. Prod. iii. 235 = Campomanesia Guaviroba.
Guayabita, *A. Rich. Ess. Fl. Cub.* 581.—Cuba.
guazumaefolium, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 280 = Campomanesia guazumaefolia.
guianense, Pers. Syn. ii. 27 = fluviatile.
Guildingianum, *Griseb. Fl. Brit. W. Ind.* 242.—Ind. occ.
guineense, Sw. Prod. Veg. Ind. Occ. 77 = P. Araça.
herbaceum, *Berg, in Mart. Fl. Bras.* xiv. i. 410.—Bras.
hians, *Mart. ex DC. Prod.* iii. 234.—Bras.
hians, Miq. in Linnaea, xxii. (1849) 533 = Campomanesia fusca.
humile, *Vell. Fl. Flum.* 211; v. t. 49.—Bras.
inaequilaterum, *Berg, in Mart. Fl. Bras.* xiv. i. 399.—Bras.
incanescens, *Mart. ex DC. Prod.* iii. 234.—Bras.
indicum, Boj. Hort. Maurit. 139 = Cattleianum.
indicum, *Raddi, Mem.* (1821) 6; *ex DC. Prod.* iii. 237 (err. typ. 337).—Bras.
intermedium, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 72 = P. Guajava.
itanareense, *Berg, in Mart. Fl. Bras.* xiv. i. 402.—Bras.
Itatiaiae, *Wawra, Itin. Princ. Coburgi*, i. 20.—Bras.
lacteum, *Berg, in Mart. Fl. Bras.* xiv. i. 403.—Bras.
lanceolatum, *Berg, in Linnaea*, xxx. (1859–60) 704.—Bras.
Langsdorffii, *Berg, in Mart. Fl. Bras.* xiv. i. 599.—Bras.
lanuginosum, *Ruiz & Pav. Fl. Per.* iv. t. 421.—Peruv.
Laruotteanum, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 282.—Bras.
latifolium, *Link, Enum. Hort. Berol.* ii. 27.—Am. austr.
laurifolium, *Berg, in Linnaea*, xxvii. (1854) 364.—Nicaragua.
leptocladum, *Berg, in Mart. Fl. Bras.* xiv. i. 409.—Bras.
lineatifolium, Pers. Syn. ii. 27 = Campomanesia lineatifolia.
littorale, *Raddi, in Opusc. Sc.* iv. (1823) 254. t. 7. f. 2.—Bras.
longifolium, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 229.—Afr. trop.
macahense, *Berg, in Mart. Fl. Bras.* xiv. i. 605.—Bras.
macrospermum, *Berg, in Mart. Fl. Bras.* xiv. i. 392.—Bras.
macrostemum, *Ruiz & Pav. Fl. Per.* iv. 420.—Peruv.
malifolium, F. Muell. Sel. Pl. Indust. Cult. 189 = Campomanesia malifolia.
maranhense, *Berg, in Mart. Fl. Bras.* xiv. i. 386.—Bras.
maribense, *Mart. ex DC. Prod.* iii. 233.—Bras.
mediterraneum, *Vell. Fl. Flum.* 212; v. t. 52.—Bras.
mengahiense, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 286.—Bras.
microcarpum, *Cambess. l. c.* 284.—Bras.
minus, Mart. ex DC. Prod. iii. 235 = P. Araça.
molle, *Bertol. Fl. Guatim.* 22. t. 9.—Guatemala.
montanum, *Sw. Prod. Veg. Ind. Occ.* 77.—Ind. occ.
monticolum, Berg, in Kjoeb. Vidensk. Meddel. (1855) 11 = molle.
Moritzianum, *Berg, in Linnaea*, xxvii. (1854) 359.—N. Granat.
multiflorum, Cambess. in A. St. Hil. Fl. Bras. ii. 281, 287 = Campomanesia multiflora.
Myrsinites, *DC. Prod.* iii. 236.—Bras.
myrsinoides, *Berg, in Mart. Fl. Bras.* xiv. i. 384.—Bras.
myrtifolium, *Lodd. ex Loud. Hort. Brit.* 197, nomen.—Hab.?
myrtoides, *Berg, in Mart. Fl. Bras.* xiv. i. 384.—Bras.

PSIDIUM:—
nigrum, *Lour. Fl. Cochinch.* 311.—Cochinch.
nitidum, *Wright, in Sauv. Fl. Cub.* 44.—Cuba.
nummularia, *Wright, l. c.*—Cuba.
nutans, *Berg, in Mart. Fl. Bras.* xiv. i. 394.—Bras.
oblongatum, *Berg, l. c.* 392.—Bras.
oblongifolium, *Berg, l. c.* 602.—Bras.
obovatum, Mart. ex DC. Prod. iii. 236 = Cattleianum.
obversum, Miq. in Linnaea, xxii. (1849) 532 = Campomanesia obversa.
Oerstedianum, *Berg, l. c.* xxvii. (1854) 360.—Guatemala.
oligospermum, *Link, Enum. Hort. Berol.* ii. 27 (= seq. ?).—Hab. ?
oligospermum, *Mart. ex DC. Prod.* iii. 236.—Bras.
ooideum, *Berg, in Mart. Fl. Bras.* xiv. i. 398.—Bras.
ovatifolium, *Berg, l. c.* 385.—Bras.
paraënse, *Berg, l. c.* 386.—Bras.
paraibicum, *Wawra, Itin. Princ. Coburgi*, i. 19. t. 1. f. B.—Bras.
paranense, *Berg, in Mart. Fl. Bras.* xiv. i. 604.—Bras.
parviflorum, *Benth. in Hook. Journ. Bot.* ii. (1840) 318.—Guian.
parvifolium, *Griseb. Cat. Pl. Cub.* 91.—Cuba.
persicifolium, *Berg, in Mart. Fl. Bras.* xiv. i. 407.—Bras.
pigmeum, *Arruda, in Koster, Trav. Braz.* 492.—Bras.
pilosum, Vell. Fl. Flum. 212; v. t. 54 = rufum.
Pohliana, *Berg, in Mart. Fl. Bras.* xiv. i. 390.—Bras.
polycarpon, *Lamb. in Trans. Linn. Soc.* xi. (1813) 231.—Am. trop.
pomiferum, Linn. Sp. Pl. ed. II. 672 = P. Guajava.
pratense, Poepp. ex Berg, in Linnaea, xxvii. (1854) 365 = rypdocarpum.
prostratum, Berg, l. c. 364 = P. Guajava.
pubescens, *Mart. ex DC. Prod.* iii. 234.—Bras.
pumilum, Vahl, Symb. Bot. ii. 56 = P. Guajava.
punctulatum, *DC. Prod.* iii. 233.—Bras.
punctulatum, Miq. in Linnaea, xxii. (1849) 513 = Campomanesia xanthocarpa.
pygmaeum, Vell. ex Steud. Nom. ed. II. ii. 406 = pigmaeum.
pyriferum, Linn. Sp. Pl. ed. II. 672 = P. Guajava.
pyriforme, Griseb. Fl. Brit. W. Ind. 241, sphalm. = praec.
racemosum, *Vell. Fl. Flum.* 212; v. t. 55.—Bras.
radicans, *Berg, in Mart. Fl. Bras.* xiv. i. 402.—Bras.
refractum, *Berg, l. c.* 394.—Bras.
rhombeum, *Berg, in Mart. Fl. Bras.* xiv. i. 383.—Bras.
Richardianum, *Berg, in Linnaea*, xxx. (1859–60) 705.—Ind. occ.
Riedelianum, *Berg, in Mart. Fl. Bras.* xiv. i. 603.—Bras.
riparium, *Mart. ex DC. Prod.* iii. 235.—Bras.
rivulare, Mart. ex DC. l. c. 233 = Campomanesia lineatifolia.
robustum, *Berg, in Mart. Fl. Bras.* xiv. i. 400.—Bras.
roseum, Mart. ex DC. Prod. iii. 264 = Eugenia rosea.
rotundatum, *Griseb. Cat. Pl. Cub.* 92.—Cuba.
rubescens, *Berg, in Mart. Fl. Bras.* xiv. i. 395.—Bras.
rubrum, *Lour. Fl. Cochinch.* 311.—Cochinch.
rufum, *Mart. ex. DC. Prod.* iii. 234.—Bras.
Ruizianum, *Berg, in Linnaea*, xxvii. (1854) 365.—Peruv.
rutidocarpum, G. Don, Gen. Syst. ii. 833, lapsu = seq.
rypdocarpum, *Ruiz & Pav. Fl. Per.* iv. t. 420.—Peruv.
salutare, *Berg, in Linnaea*, xxvii. (1854) 356.—N. Granat.
sapidissimum, Jacq. Hort. Schoenb. iii. 62 = P. Guajava.
Schiedeanum, *Berg, in Linnaea*, xxvii. (1854) 368.—Mexic.
Sellowianum, Berg, in Mart. Fl. Bras. xiv. i. 400 = arboreum.
sericeum, *Berg, l. c.* 389. = Bras.
sericiflorum, *Benth. Pl. Hartw.* 176.—N. Granat.
Sieberianum, *Berg, in Mart. Fl. Bras.* xiv. i. 387.—Bras.
sorocabense, *Berg, l. c.* 398.—Bras.

PSIDIUM:—
Sprucei, *Berg, in Mart. Fl. Bras.* xiv. i. 396.—Bras.
striatulum, *DC. Prod.* iii. 233.—Bras.
suaveolens, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 291 = Campomanesia suaveolens.
suffruticosum, *Berg, in Mart. Fl. Bras.* xiv. i. 387.—Bras.
tenuifolium, Mart. ex. DC. Prod. iii. 236 = Campomanesia tenuifolia.
terminale, *Vell. Fl. Flum.* 212; v. t. 53.—Bras.
ternatifolium, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 278 = grandifolium.
Thea, *Griseb. in Goett. Abh.* xix. (1874) 139.—Reg. Argent.
tomentosum, *Parodi, in Anal. Soc. Cient. Argent.* vii. (1879) 64; *Contrib.* 114.—Reg. Argent.
transalpinum, Vell. Fl. Flum. 213; v. t. 59 = Campomanesia transalpina.
turbiniflorum, *Mart. ex DC. Prod.* iii. 234. = Bras.
umbrosum, *Berg, in Mart. Fl. Bras.* xiv. i. 599.—Bras.
variabile, Berg, l. c. 400 = Cattleianum.
velutinum, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 288 = Campomanesia velutina.
vulgare, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = P. Guajava.
Widgrenianum, *Berg, in Mart. Fl. Bras.* xiv. i. 392.—Bras.
Willemetianum, *DC. Prod.* iii. 237 (err. typ. 337).—Ins. Maurit.
xidocarpum, Ruiz, ex Berg, in Linnaea, xxvii. (1854) 365 = rypdocarpum.
ypanemense, *Berg, in Mart. Fl. Bras.* xiv. i. 395.—Bras.

PSIGURIA, Neck. Elem. i. 137 (1790) = **Anguria**, Tourn. (Cucurb.).

PSILACHENIA, Benth. in Journ. Linn. Soc. xiii. (1873) 475, sphalm. = PSILOCHENIA, Nutt. = **Crepis**, Vaill. (Compos.).

PSILACTIS, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 71. *COMPOSITAE*, Benth. & Hook. f. ii. 269.
asteroides, *A. Gray, l. c.*—N. Mexic.
brevilingulata, *Sch. Bip. ex Hemsl. Diagn. Pl. Nov.* 34.—Mexic.
Coulteri, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 71.—Mexic.

PSILAEA, Miq. Fl. Ind. Bat. Suppl. 355 (1860) = **Linostoma**, Wall. (Thymel.).
dalbergioides, Miq. l. c. = L. pauciflorum.

PSILANTHUS, Hook. f. in Hook. Ic. Pl. t. 1129 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 115.
ebracteolatus, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 186.—Afr. trop.
Mannii, *Hook. f. in Hook. Ic. Pl. t.* 1129.—Afr. trop.
tetramerus, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 187.—Afr. trop.

PSILANTHUS, Juss. in Ann. Mus. Par. vi. (1805) 396 = **Passiflora**, Linn.
trinervia, Juss. l. c. = Pass. trinervia.
viridiflora, M. Roem. Syn. Pepon. 198 = Pass. viridiflora.

PSILARABIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = **Arabis**, Linn. (Crucif.).
glabra, Fourr. l. c. = A. perfoliata.

PSILATHERA, Link, Hort. Berol. i. 121 (1827) = **Sesleria**, Scop. (Gramin.).
tenella, Link, l. c. 122 = S. tenella.

PSILOBIUM, Jack, in Malay. Misc. ii. (1822) n. VII. 84. *RUBIACEAE*, Benth. & Hook. f. ii. 75.
capillare, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 313 = Morindopsis capillaris.
nutans, *Jack, in Malay. Misc.* ii. (1822) n. VII. 85.—Sumatra.
tomentosum, *Jack, l. c.* p. iii.—Malaya.

PSILOCARPHUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 340. *COMPOSITAE*, Benth. & Hook. f. ii. 297.
BENZANILLA, Remy, in C. Gay, Fl. Chil. iv. 109. t. 46. (1849).

PSILOCARPHUS:—
brevissimus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 340.—Am. bor. occ.
caulescens, *Benth. Pl. Hartw.* 319.—Calif.
globiferus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 340. = Calif.
oreganus, *A. Gray, in Proc. Am. Acad.* viii. (1873) 390 (non seq.).—Am. bor. occ.
oregonus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 341.—Am. bor. occ.
tenellus, *Nutt. l. c.*—Calif.

PSILOCARPUS, Pritz. Ic. Ind. ii. 233 (1865) = **Psophocarpus**, Neck. (Legumin.).

PSILOCARYA, Torr. in Ann. Lyc. N. York, iii. (1836) 359 = **Rynchospora**, Vahl (Cyperac.).
candida, Nees, in Mart. Fl. Bras. ii. 1. 117 = R. candida.
conferta, Nees, l. c. 116 = R. velutina.
corymbifera, Benth. in Hook. Ic. Pl. xv. 25. t. 1431 = seq.
corymbiformis, Benth. & Hook. f. Gen. iii. 1048.—Cuba.
divergens, Nees, in Mart. Fl. Bras. ii. 1. 116 = R. velutina.
mexicana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 248 = R. rufa.
nitens, Wood, Am. Bot. & Flor. 364 = R. nitens.
Pohlii, Nees, in Mart. Fl. Bras. ii. 1. 115 = R. velutina.
rhynchosporoides, Torr. in Ann. Lyc. N. York, iii. (1836) 361 = R. nitens.
robusta, Nees, in Mart. Fl. Bras. ii. 1. 116 = R. robusta.
rufa, Nees, l. c. 117 = R. rufa.
Schiedeana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 248 = R. eximia.
scirpoides, Torr. in Ann. Lyc. N. York, iii. (1836) 360 = R. scirpoides.
Sellowiana, Nees, in Mart. Fl. Bras. ii. 1. 117 = R. velutina.
Teneriffae, Torr. in Ann. Lyc. N. York, iii. (1836) 362, in adn. = R. micrantha.
texensis, Torr. l. c. 361 = R. nitens.
velutina, Nees, in Mart. Fl. Bras. ii. 1. 115. t. 7 = R. velutina.

PSILOCHLAENA, Walp. Rep. ii. 697 (1843) = seq.

PSILOCHENIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 437 = **Crepis**, Vaill. (Compos.).
occidentalis, Nutt. l. c. = C. occidentalis.

PSILOCHILUS, Rodrig. Orch. Nov. ii. 272. *ORCHIDEAE.*
modestus, *Rodrig. l. c.* 273.—Bras.

PSILOGYNE, DC. in Bibl. Univ. Genèv. xvii. (1838) 132 (Rev. Bign. 16) = **Vitex**, Linn. (Verben.).
viticifolia, DC. ll. cc. = V. multinervis.

PSILOLEPUS, Presl, Bot. Bemerk. 128 (1844) = **Aspalathus**, Linn. (Legumin.).
bracteatus, Presl, l. c. 129 = A. bracteata.
lanatus, Presl, l. c. = A. lanata.
pedunculatus, Presl, l. c. = A. pedunculata.

PSILONEMA, C. A. Mey. in Ledeb. Fl. Alt. iii. 50 (1831) = **Alyssum**, Linn. (Crucif.).
calycinum, C. A. Mey. in Bull. Sc. Acad. Pétersb. vii. (1840) 132 = A. calycinum.
campestre, Schur, Enum. Pl. Transs. 62 = A. campestre.
dasycarpum, C. A. Mey. in Ledeb. Fl. Alt. iii. 51 = A. dasycarpum.
hirsutum, Schur, Enum. Pl. Transs. 62 = A. hirsutum.
homalocarpum, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 63 = A. homalocarpum.
minimum, Schur, Enum. Pl. Transs. 62 = A. minimum.
ruderale, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 334 = A. calycinum.
sabulosum, Fourr. l. c. = A. calycinum.
vagum, Fourr. l. c. = A. calycinum.

PSILOPOGON, Hochst. in Flora, xxiv. (1841) 1. Intell. 20 = **Arthraxon**, Beauv. (Gramin.).
capensis, Hochst. l. c. xxix. (1846) 117 = Pollinia nuda.

PSILOPOGON:—
Figarei, De Not. Ind. Sem. Hort. Genuens. (1852) 25; et in Ann. Sc. Nat. Sér. III. xix. (1853) 370 = Arthraxon microphyllus.
Schimperi, Hochst. in Flora, xxix. (1846) 117, in nota = A. microphyllus.

PSILOPOGON, Phil. in Linnaea, xxxiii. (1864–65) 126 = **Picrosia**, D. Don (Compos.).
albiflorus, Phil. l. c. = Picr. longifolia.

PSILOPSIS, Neck. Elem. i. 319 (1790) = **Lamium**, Tourn. (Labiat.).

PSILOSANTHUS, Neck. Elem. i. 69 (1790) = **Liatris**, Schreb. (Compos.).

PSILOSIPHON, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 272 = **Lapeyrousia**, Pourr. (Irid.).
cyanescens, Welw. l. c. = L. cyanescens.
fragrans, Welw. l. c. = L. fragrans.
odoratissimus, Welw. l. c. 273 = L. odoratissima.

PSILOSOLENA, Presl, Bot. Bemerk. 102 (1844) = **Peddiea**, Harv. (Thymel.).
umbellata, Presl, l. c. = Peddiea Dregei.

PSILOSTACHYS, Hochst. in Flora, xxvii. (1844) 1. Beil. 6. t. 4. *AMARANTACEAE*, Benth. & Hook. f. iii. 32.
POECHIA, Endl. Gen. Suppl. iv. 43 (1847).
POECKIA, Benth. & Hook. f. Gen. iii. 32 (1880), err. typ.
gnaphalobrya, *Hochst. in Flora*, xxvii. (1844) 1. *Beil.* 6.—Afr. bor.
sericea, *Hook. f. in Benth. & Hook. f. Gen.* iii. 32.—Ind. or.

PSILOSTACHYS, Steud. Syn. Pl. Gram. 413 (1855) = **Dimeria**, R. Br. (Gramin.).
Hohenackeri, Steud. l. c. = D. Hohenackeri.

PSILOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 58 = **Cleidion**, Blume (Euphorb.).
axillaris, Turcz. l. c. = C. tricoccum.

PSILOSTEMON, DC. Prod. x. 35 (1846) = **Trachystemon**, D. Don (Borag.).
creticum, DC. l. c. 36 = T. creticum.
orientale, DC. l. c. = T. orientale.

PSILOSTOMA, Klotzsch, in Eckl. & Zeyh. Enum. 361 (1837) = **Plectronia**, Linn. (Rubiac.).
ciliata, Klotzsch, l. c. 362 = Plec. ciliata.

PSILOSTROPHE, DC. Prod. vii. 261 (1838) = **Riddellia**, Nutt. (Compos.).
gnaphalodes, DC. l. c. = R. arachnoidea.

PSILOTHAMNUS, DC. Prod. vi. 41 (1837) = **Gamolepis**, Less. (Compos.).
adpressifolius, DC. l. c. = G. ericoides.
ericaefolius, DC. l. c. = G. ericoides.

PSILOTHONNA, E. Mey. ex DC. Prod. vi. 40 (1837) = **Gamolepis**, Less. (Compos.).
Tagetes, E. Mey. ex DC. l. c. = G. annua.

PSILOTRICHUM, Blume, Bijdr. 544 (1825). *AMARANTACEAE*, Benth. & Hook. f. iii. 32.
LEIOSPERMUM, Wall. Cat. n. 6923 (1831).
calceolatum, *Moq. in DC. Prod.* xiii. 11. 280.—Ind. or.
capitatum, *F. Muell. Fragm.* i. 238.—Austral.
cordatum, Moq. in DC. Prod. xiii. 11. 280 = Psilostachys gnaphalobrya.
ferrugineum, *Moq. l. c.* 279.—Ind. or.
helichrysoides, F. Muell. Fragm. i. 237 = Trichinium helichrysoides.
nudum, *Moq. in DC. Prod.* xiii. 11. 280.—Ind. or.
nudum, Wight, Ic. t. 1775 = calceolatum.
ovatum, Moq. in DC. Prod. xiii. 11. 281 = calceolatum.
sandwichense, *Seem. Fl. Vit.* 198.—Ins. Sandvic.
scleranthum, *Thw. Enum. Pl. Zeyl.* 248.—Zeylan.
sericeum, Dalz. in Dalz. & Gibs. Bomb. Fl. 216 = Psilostachys sericea.
trichotomum, *Blume, Bijdr.* 545.—Ind. or.; Malaya.

PSILOXYLON, Thou. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 138 = **Fropiera**, Bouton (Myrtac.).
mauritianum, Baill. Adansonia, x. (1871) 41 = F. mauritiana.
mauritianum, Thou. ex Benth. in DC. Prod. x. 434 = Nuxia verticillata.

PSILURUS, Trin. Fund. Agrost. 93 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1205.
ASPRELLA, Host, Gram. Austr. iv. 17. t. 29 (1809).
MONERMA, Beauv. Agrost. 116 (1812), pro parte.
OPHIURINELLA, Desv. Opusc. 75. t. 5. f. 4 (1831).
aristatus, Duval-Jouve, in Bull. Soc. Bot. Fr. xiii. (1866) 132 = nardoides.
nardoides, *Trin. Fund. Agrost.* 93.—Europ. austr.
rottboellioides, *Griff. Notul.* iii. 95.—Afghan.

PSISTINA, Rafin. Sylva Tellur. 133 (1838) = **Helianthemum**, Tourn. (Cistin.).

PSISTUS, Neck. Elem. ii. 273 (1790) = **Helianthemum**, Tourn. (Cistin.).

PSITHYRISMA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85 = **Symphyostemon**, Miers, etc. (Irid.).
flexuosum, Herb. l. c., nomen = Sisyrinchium cuspidatum.
narcissoides, Herb. l. c., nomen = Symphyostemon narcissoides.
odoratissimum, Herb. l. c., nomen = Solenomelus biflorus.

PSITTACANTHUS, Mart. in Flora, xiii. (1830) 106 = **Loranthus**, Linn.
acinarius, Mart. l. c. 108 = L. acinarius.
acutifolius, G. Don, Gen. Syst. iii. 417 = L. acutifolius.
americanus, Mart. in Flora, xiii. (1830) 108 = L. americanus, *Linn.*
auriculatus, Eichl. in Mart. Fl. Bras. v. 11. 25 = L. auriculatus.
Berteroi, G. Don, Gen. Syst. iii. 418 = L. Berteroi.
bicalyculatus, Mart. in Flora, xiii. (1830) 108 = L. bicalyculatus.
biternatus, Blume, in Schult. f. Syst. vii. 1730 = L. biternatus.
brachynema, Eichl. in Mart. Fl. Bras. v. 11. 29.—Am. trop.
brasiliensis, Blume, in Schult. f. Syst. vii. 1730 = L. brasiliensis.
calyculatus, G. Don, Gen. Syst. iii. 415 = L. calyculatus.
cardiphyllus, Eichl. in Mart. Fl. Bras. v. 11. 27 = L. cardiphyllus.
chanduyensis, Eichl. l. c.—Ecuador.
cinctus, Mart. in Flora, xiii. (1830) 108 = L. cinctus.
claviceps, Eichl. in Mart. Fl. Bras. v. 11. 26 = L. claviceps.
clusiaefolius, Eichl. l. c. 30 = L. dichotomus.
Collum-cygni, Eichl. l. c. 35.—Am. trop.
cordatus, Blume, in Schult. f. Syst. vii. 1730 = L. cordatus.
corymbosus, Blume, l. c. = L. corymbosus.
corynocephalus, Eichl. in Mart. Fl. Bras. v. 11. 36.—Bras.
crassifolius, Mart. in Flora, xiii. (1830) 108 = L. crassifolius.
cucullaris, Blume, in Schult. f. Syst. vii. 1730 = L. cucullaris.
cuneifolius, Blume, l. c. = L. cuneifolius.
cupulifer, Blume, l. c. = L. cupulifer.
cupulifer, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 1162 = L. cucullaris.
decipiens, Eichl. in Mart. Fl. Bras. v. 11. 35.—Bras.
Destructor, G. Don, Gen. Syst. iii. 416 = L. Destructor.
dichotomus, Blume, in Schult. f. Syst. vii. 1730 = L. dichotomus.
dichroos, Mart. in Flora, xiii. (1830) 108 = L. dichrous.
divaricatus, Blume, in Schult. f. Syst. vii. 1730 = L. divaricatus.
drepanophyllus, Eichl. in Mart. Fl. Bras. v. 11. 40.—Bras.
Eschscholzianus, G. Don, Gen. Syst. iii. 417 = L. Eschscholzianus.
eucalyptifolius, Blume, in Schult. Syst. vii. 1730 = L. eucalyptifolius.
falcifrons, Mart. in Flora, xiii. (1830) 107 = L. falcifrons.
flagellaris, Mart. l. c. 108 = L. flagellaris.

PSITTACANTHUS :—
flavo-viridis, Eichl. in Mart. Fl. Bras. v. II. 41.—Bras.
formosus, G. Don, Gen. Syst. iii. 416 = Loranthus robustus.
furcatus, Mart. in Flora, xiii. (1830) 108 = L. furcatus.
glaucocoma, Eichl. in Mart. Fl. Bras. v. II. 32.—Bras.
grandiflorus, Blume, in Schult. f. Syst. vii. 1730 = L. grandiflorus, *Ruiz & Pav.*
grandiflorus, G. Don, Gen. Syst. iii. 415 = L. grandifolius.
grandifolius, Mart. in Flora, xiii. (1830) 108 = L. grandifolius.
guianensis, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 1161 = L. cordatus.
Hoffmannseggianus, Blume, in Schult. f. Syst. vii. 1730 = L. Hoffmannseggianus.
Holtoni, Eichl. in Mart. Fl. Bras. v. II. 24.—N. Granat.
Jacquini, G. Don, Gen. Syst. iii. 415 = L. americanus, *Linn.*
Karwinskianus, Eichl. in Mart. Fl. Bras. v. II. 26 = L. Karwinskyanus.
lamprophyllus, Eichl. l. c. 28.—Bras.
longibracteatus, Blume, in Schult. f. Syst. vii. 1730 = L. longibracteatus.
macrantherus, Eichl. in Mart. Fl. Bras. v. II. 26 = L. macrantherus.
martinicensis, Eichl. l. c. = L. martinicensis.
mexicanus, Blume, in Schult. f. Syst. vii. 1730 = L. mexicanus.
Mutisii, Blume, l. c. = L. Mutisii.
nodosus, Blume, l. c. = L. nodosus.
obmunitus, Blume, l. c. = L. obmunitus.
obovatus, Eichl. in Mart. Fl. Bras. v. II. 27 = L. obovatus, *Benth.*
peronopetalus, Eichl. l. c. 31.—Bras.
pinguis, Eichl. l. c. 37.—Bras.
plagiophyllus, Eichl. l. c.—Bras.
Plumieri, G. Don, Gen. Syst. iii. 417 = L. Plumieri.
ramiflorus, G. Don, l. c. 415 = L. ramiflorus.
robustus, Mart. in Flora, xiii. (1830) 108 = L. robustus.
scalpratus, Vahl, ex Eichl. in Mart. Fl. Bras. v. II. 25.—Ind. occ.
Schiedeanus, Blume, in Schult. f. Syst. vii. 1730 = L. Schiedeanus.
Siphon, Eichl. in Mart. Fl. Bras. v. II. 28.—Bras.
Smithii, Klotzsch, in Schomb. Fauna & Fl. Brit. Gui. 1162 = L. cucullaris.
speciosus, G. Don, Gen. Syst. iii. 415 = L. speciosus.
Sternbergianus, Blume, in Schult. f. Syst. vii. 1730 = L. Sternbergianus.
thyrsiflorus, G. Don, Gen. Syst. iii. 417 = L. thyrsiflorus.
Velloxianus, Mart. in Flora, xxiv. (1841) II. Beibl. 33 = L. dichrous.
Warmingii, Eichl. in Mart. Fl. Bras. v. II. 36.—Bras.

PSITTACOGLOSSUM, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 29 (1825) = **Maxillaria**, Ruiz & Pav. (Orchid.).
atratum, La Llave & Lex. l. c. 30 = M. atrata.

PSITTACOSCHOENUS, Nees, in Lehm. Pl. Preiss. ii. 87 (1846), in nota = **Gahnia**, Forst. (Cyperac.).
erythrocarpus, F. Muell. ex Boeck. in Linnaea, xxxviii. (1874) 341 = G. psittacorum.

PSOLANUM, Neck. Elem. ii. 60 (1790) = **Solanum**, Tourn.

PSOPHIZA, Rafin. Fl. Tellur. iv. 99 (1836) = **Aristolochia**, Tourn.
undulata, Rafin. l. c. = A. sempervirens?

PSOPHOCARPUS, Neck. Elem. iii. 45 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 540.
Botor, Adans. Fam. ii. 326 (1763).
Diesingia, Endl. in Flora, xv. (1832) 113.
Psilocarpus, Pritz. Ic. Ind. ii. 233 (1865).
comorensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 380.—Ins. Mascar.
longepedunculatus, Hassk. in Flora, xxv. (1842) II. Beibl. 75 = palustris.

PSOPHOCARPUS :—
Mabala, *Welw. Apont.* 589.—Afr. trop.
palmettorum, Guill. & Perr. Fl. Seneg. Tent. i. 222 = palustris.
palustris, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 420.—Afr. trop.
tetragonolobus, *DC. Prod.* ii. 403.—Ins. Maurit.; As. trop. (cult.).

PSORA, Hill, Veg. Syst. iv. 30 (1762) = **Centaurea**, Linn. (Compos.).
orientalis, Hill, Hort. Kew. 65 = C. orientalis.

PSORALEA, Linn. Gen. ed. II. 358 (1742). *LEGUMINOSAE*, Benth. & Hook. f. i. 491.
Asphalthium, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 380.
Cullen, Medic. l. c. 381 (1787).
Meladenia, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 576.
Orbexilum, Rafin. Atl. Journ. 145 (1832).
Rhynchodium, Presl, in Abh. Boehm. Ges. V. iii. (1845) 484; Bot. Bemerk. 54 (1844).
Ruteria, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 380.
acanthocarpa, F. Muell. Fragm. iii. 45 = Glycyrrhiza psoraleoides.
acaulis, *Stev. ex Bieb. Fl. Taur. Cauc.* ii. 206.—Reg. Caucas.
aculeata, *Linn. Sp. Pl.* 763.—Afr. austr.
aculeata, Thunb. ex Harv. & Sond. Fl. Cap. ii. 154 = bracteata.
acuminata, Eckl. & Zeyh. Enum. 229 = bracteata.
acuminata, Lam. Illustr. iii. 173 = aculeata.
adscendens, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 40.—Austral.
affinis, Eckl. & Zeyh. Enum. 224 = pinnata.
albicans, Eckl. & Zeyh. l. c. 231 = striata.
algoënsis, Eckl. & Zeyh. l. c. 229 = bracteata.
alnifolia, Bertol. Misc. Bot. ix. 10 = Rhynchosia tomentosa.
alopecurina, Bertol. l. c. 11 = Rhynchosia tomentosa.
alopecuroides, Poir. Encyc. v. 695 = Dalea alopecuroides.
altissima, *Bertol. in Nov. Comm. Bonon.* iii. (1839) 16. t. 2. f. 1.—Afr. austr.
americana, Linn. Sp. Pl. 763 = dentata.
andongensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 65.—Afr. trop.
angustifolia, *L'Hérit. ex Ait. Hort. Kew.* ed. I. iii. 79.—Afr. austr.
angustifolia, Willd. in Roem. & Usteri, Mag. Bot. iv. 11tes St. (1790) 33 = bituminosa.
aphylla, Jacq. Hort. Schoenb. ii. 51. t. 223 = hirta.
aphylla, *Linn. Pl. Afr. Rar.* 15; *Amoen. Acad.* vi. 93.—Afr. austr.
arabica, Hochst. & Steud. ex Boiss. Fl. Orient. ii. 193 = Tephrosia Pogonostigma.
arborea, Sims, Bot. Mag. t. 2090 = pinnata.
Archeri, *F. Muell. Fragm.* iv. 21.—Austral.
arenaria, Nutt. Gen. Am. ii. 103 = lanceolata.
argentea, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
argophylla, *Pursh, Fl. Am. Sept.* ii. 475.—Am. bor.
ascendens, C. Muell. in Walp. Ann. vii. 707 = adscendens.
astragalifolia, Poir. Encyc. Suppl. iv. 587 = capitata.
aurea, Poir. l. c. 590 = Dalea aurea.
australasica, Schlecht. in Linnaea, xx. (1847) 668 = patens.
axillaris, *Linn. f. Suppl.* 338.—Afr. austr.
axillaris, Poir. Encyc. v. 687 = angustifolia.
azurea, *Phil. Fl. Atac.* 14.—Chili.
badocana, *Blanco, Fl. Filip.* ed. II. 416; *Benth. Fl. Austral.* ii. 190.—Ins. Philipp.; Austral.
balsamica, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 55.—Austral.
biflora, *Harv. in Harv. & Sond. Fl. Cap.* ii. 157.—Afr. austr.
biflora, *Rudolph, in Schrad. Journ.* ii. (1799) 291.—Hab.?
bipedunculata, *Jacq. Coll.* iv. 146.—Hab.?
bituminosa, *Linn. Sp. Pl.* 763.—Reg. Mediter.; Arab.
Bowieana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 154.—Afr. austr.
brachiata, Dougl. ex Hook. Fl. Bor. Am. i. 137. t. 53 = esculenta.
bracteata, *Berg. Desc. Pl. Cap.* 224; *Linn. Mant.* ii. 264.—Afr. austr.

PSORALEA :—
bracteolata, Eckl. & Zeyh. Enum. 229 = bracteata.
caffra, *Eckl. & Zeyh. l. c.* 230.—Afr. austr.
cajanifolia, Benth. ex Harv. Thes. Cap. i. 52 = Pleiospora cajanifolia.
californica, *S. Wats. in Proc. Am. Acad.* xii. (1877) 251.—Calif.
campestris, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 301.—Am. bor.
candicans, *Eckl. & Zeyh. Enum.* 228.—Afr. austr.
candida, Poir. Encyc. v. 694 = Petalostemon candidus.
canescens, *Michx. Fl. Bor. Am.* ii. 57.—Am. bor.
capitata, *Linn. f. Suppl.* 339.—Afr. austr.
carnea, *E. Mey. Comm. Pl. Afr. Austr.* 85.—Afr. austr.
carnea, Poir. Encyc. v. 694 = Petalostemon carneus.
carthaginensis, Jacq. Enum. Pl. Carib. 27 = Dalea enneaphylla.
castorea, *S. Wats. in Am. Natural.* xii. (1878) 601; *et in Proc. Am. Acad.* xiv. (1879) 291.—Calif.
cephalantha, F. Muell. Fragm. iv. 35 = badocana.
Cephalotes, E. Mey. Comm. Pl. Afr. Austr. 87 = uncinata.
Cephalotes, C. Presl, ex J. Presl, Rostl. iii. 182; Eckl. & Zeyh. Enum. 230 = bracteata.
cinerea, *Lindl. in Mitch. Three Exped.* ii. 65.—Austral.
citriodora, Cav. Ic. iii. 36 = Dalea citriodora.
conferta, *Bertol. in Nov. Comm. Bonon.* iii. (1839) 17. t. 2. f. 2.—Afr. austr.
corylifolia, *Linn. Sp. Pl.* 764.—Ind. or.; Arab.
corymbosa, Poir. Encyc. v. 694 = Petalostemon corymbosus.
cryptocarpa, Torr. & Gray, Fl. N. Am. i. 301 = cuspidata.
Cullen, Poir. Encyc. v. 685 = glandulosa.
cuneifolia, Dum.-Cours. Bot. Cult. ed. II. vi. 101 = bracteata.
cuspidata, *Pursh, Fl. Am. Sept.* ii. 741.—Am. bor.
cyphocalyx, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 172.—Texas.
cytisoides, Linn. Sp. Pl. ed. II. 1075 = Indigofera cytisoides.
Dalea, Linn. l. c. ed. I. 764 = Dalea alopecuroides.
decidua, Berg. Desc. Pl. Cap. 220 = aphylla.
decidua, Sieber, ex Walp. in Linnaea, xiii. (1839) 512 = pinnata.
decumbens, *Ait. Hort. Kew.* ed. I. iii. 80.—Afr. austr.
decumbens, Willd. ex Walp. in Linnaea, xiii. (1839) 514 = Mundiana.
densa, E. Mey. Comm. Pl. Afr. Austr. 86 = bracteata.
dentata, *DC. Prod.* ii. 221.—Reg. Mediter. occ.
denudata, Hoffmgg. Verz. Pfl. 103 = aphylla.
diffusa, Eckl. & Zeyh. Enum. 225 = repens.
digitata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 300.—Am. bor.
divaricata, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 788.—Peruv.
Dregeana, Steud. Nom. ed. II. ii. 407 = uncinata.
Drummondii, Meissn. in Bot. Zeit. xiii. (1855) 31 = cinerea.
drupacea, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 221.—Persia; Turkest.
Eckloniana, Otto, ex Harv. & Sond. Fl. Cap. ii. 153 = striata.
Ecklonis, Hort. Viadr. ex Walp. in Linnaea, xiii. (1839) 515 = striata.
eglandulosa, Ell. Sketch, ii. 198 = melilotoides.
elata, Salisb. Prod. 332 = bituminosa.
elliptica, Pursh, Fl. Am. Sept. ii. 741 = lanceolata.
emphysodes, Jacq. Coll. iv. 144 = Dalea phymatodes.
enneaphylla, Bert. ex Steud. Nom. ed. II. ii. 407 = Dalea domingensis.
enneaphylla, Linn. Sp. Pl. ed. II. 1076 = Dalea enneaphylla.
eriantha, *Benth. in Mitch. Journ. Trop. Austral.* 131.—Austral.
esculenta, *Pursh, Fl. Am. Sept.* ii. 475. t. 22.—Am. bor.
exigua, Eckl. & Zeyh. Enum. 232 = obtusifolia.
fascicularis, *DC. Prod.* ii. 217.—Afr. austr.
fasciculata, Steud. Nom. ed. II. ii. 75, in syn., sphalm. = praec.
filifolia, Eckl. & Zeyh. Enum. 226 = aphylla.
filiformis, *Poir. Encyc.* v. 682.—Afr. austr.
floribunda, Nutt. ex Torr. & Gray, Fl. N. Am. i. 300 = tenuiflora.

PSORALEA:—
floridana, Shuttlew. ex A. Gray, Pl. Wright, i. 45 = P. Lupinellus.
foetida, *Presl, Delic. Prag.* 52.—Sicilia.
foliolosa, Ait. Hort. Kew. ed. I. iii. 82 = Dalea citriodora.
foliosa, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 399.—Afr. trop.
frutescens, Poir. Encyc. v. 684 = bituminosa, pubescens.
fruticosa, Kellogg, in Proc. Calif. Acad. vii. (1877) 91 = bracteata.
Gaudichaudiana, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 467.—Ins. Timor.
glabra, E. Mey. Comm. Pl. Afr. Austr. 83 = pinnata.
glandulosa, *Linn. Sp. Pl.* ed. II. 1075.—Chili.
glaucescens, Eckl. & Zeyh. Enum. 227 = oligophylla.
glaucina, *Harv. in Harv. & Sond. Fl. Cap.* ii. 147.—Afr. austr.
gracilis, Chapm. ex Torr. & Gray, Fl. N. Am. i. 303 = melilotoides.
Gueinzii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 145.—Afr. austr.
Gunnii, Hook. f. Fl. Tasm. i. 99 = adscendens.
hamata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 152.—Afr. austr.
Harveyana, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 82.—Afr. austr.
herbacea, Sieber, ex DC. Prod. ii. 168 = Lotononis umbellata.
Higuera, *Griseb. in Goett. Abh.* xxiv. (1879) 99.—Reg. Argent.
Higuerilla, *Gill. ex Hook. Bot. Misc.* iii. (1833) 181.—Reg. Argent.
hilaris, *Eckl. & Zeyh. Enum.* 230.—Afr. austr.
hirta, *Linn. Pl. Afr. Rar.* 15; *Amoen. Acad.* vi. 93.—Afr. austr.
hirta, Thunb. Prod. Pl. Cap. 136 = candicans.
hirta, Willd. ex Walp. in Linnaea, xiii. (1839) 515 = striata.
humilis, *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
hypogaea, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 302.—Am. bor.
incana, Nutt. in Fras. Cat. (1813); ex ej. Gen. Am. ii. 102 = argophylla.
intermedia, Bess. Cat. Hort. Crem. 111 = angustifolia.
involucrata, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
Jacquiniana, Eckl. & Zeyh. Enum. 226 = aphylla.
Jacquiniana, Sweet, Hort. Brit. ed. II. 138 = hirta.
Jamesii, Torr. in Ann. Lyc. N. York, ii. (1828) 175 = Dalea Jamesii.
Jaubertiana, *Fenzl, in Flora*, xxvi. (1843) 392.—Syria.
Kraussiana, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 81.—Afr. austr.
Krebsii, Vog. ex Walp. in Linnaea, xiii. (1839) 512 = pinnata.
lachnostachys, *F. Muell. Fragm.* iii. 105.—Austral.
laevigata, Eckl. & Zeyh. Enum. 224 = pinnata.
laevigata, Linn. f. Suppl. 339 = odoratissima?
Lagopus, Cav. Ic. i. 59 = Dalea Lagopus.
lanceolata, *Pursh, Fl. Am. Sept.* ii. 475.—Am. bor.
lasiostachys, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 13.—Peruv.
lateralis, Salisb. Prod. 332 = dentata.
lathyrifolia, *Balb. in Mém. Acad. Turin*, x. 1. (1810) 360. t. 8.—Hab.?
latifolia, Torr. in Ann. Lyc. N. York, ii. (1828) 176 = P. Onobrychis.
laxiflora, Nutt. in Torr. & Gray, Fl. N. Am. i. 299 = lanceolata.
laxiflora, Poir. Encyc. Suppl. iv. 590 = Dalea laxiflora.
Leichhardtii, F. Muell. Fragm. iv. 22 = Indigofera glandulosa.
leporina, Ait. Hort. Kew. ed. I. iii. 81 = Dalea Lagopus.
leucantha, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 54.—Austral.
linearifolia, *Torr. & Gray, Fl. N. Am.* i. 300.—Am. bor.
linearis, *Burm. f. Fl. Cap. Prod.* 22 (Quid?).—Afr. austr.
linearis, Thunb. Prod. Pl. Cap. 135 = axillaris.
longifolia, Pursh, Fl. Am. Sept. ii. 741 = Astragalus pictus, *A. Gray*.
Lupinellus, *Michx. Fl. Bor. Am.* ii. 58.—Am. bor.
lutea, Cav. Ic. iv. 12 = Dalea lutea.

PSORALEA:—
lutea, Molina, Sagg. Chile, 163 = glandulosa.
macradenia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 148.—Afr. austr.
macrorhiza, Fras. Cat. (1813); ex Pursh, Fl. Am. Sept. ii. 741 = cuspidata.
macrostachya, *DC. Prod.* ii. 220.—Calif.
marginata, *Meyen, Reise*, i. 436.—Peruv.
Martini, *F. Muell. Fragm.* v. 11.—Austral.
melanocarpa, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 234.—Mexic.
melilotoides, *Michx. Fl. Bor. Am.* ii. 58.—Am. bor.
Melilotus, Pers. Syn. ii. 347 = melilotoides.
mephitica, *S. Wats. in Am. Natural.* xii. (1878) 601; *et in Proc. Am. Acad.* xiv. (1879) 291.—Calif.
micrantha, *A. Gray, in Pacif. Rail. Rep.* iv. 77.—Am. bor. occ.
microcephala, *Reichb. ex Kunze, in Linnaea*, xx. (1847) 62.—Hab.?
mucronata, Thunb. Prod. Pl. Cap. 35 = aculeata.
multicaulis, Jacq. Hort. Schoenb. ii. 55. t. 230 = capitata.
multifoliolata, Clos, in C. Gay, Fl. Chil. ii. 87 = Dalea multifoliata.
multijuga, *Ell. Sketch*, ii. 198.—Am. bor.
Mundiana, *Eckl. & Zeyh. Enum.* 228.—Afr. austr.
mutabilis, Cav. Ic. iv. 65 = Dalea mùtabilis.
Mutisii, *Kunth, Mim.* 191. t. 54.—N. Granat.
nutans, Cav. Ic. iii. 1 = Dalea nutans.
obliqua, *E. Mey. in Linnaea*, vii. (1832) 164.—Afr. austr.
obovata, Roxb. Hort. Beng. 99; Fl. Ind. iii. 388 = plicata.
obtusifolia, *DC. Prod.* ii. 221.—Afr. austr.
obtusiloba, Torr. & Gray, Fl. N. Am. i. 300 = tenuiflora.
odoratissima, *Jacq. Hort. Schoenb.* ii. 54. t. 229.—Afr. austr.
oligophylla, *Eckl. & Zeyh. Enum.* 227.—Afr. austr.
Onobrychis, *Nutt. Gen. Am.* ii. 104.—Am. bor.
ononides, *Burm. f. Fl. Cap. Prod.* 22.—Afr. austr.
ononoides, Poir. Encyc. v. 688 = decumbens.
orbicularis, *Lindl. Bot. Reg.* t. 1971.—Calif.
palaestina, Gouan, Illustr. 51; Murr. Syst. ed. XIII. 570 = bituminosa.
palmata, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 451 = cyphocalyx.
parva, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 40 = adscendens.
parviflora, E. Mey. Comm. Pl. Afr. Austr. i. 86 = bracteata.
parviflora, Poir. Encyc. Suppl. iv. 590 = Astragalus gracilis.
parviflora, Tausch, in Flora, xix. (1836) 420 = Tauschiana.
patens, *Lindl. in Mitch. Three Exped.* ii. 9.—Austral.
pedunculata, [Ker.-Gawl.] in Bot. Reg. t. 223 = tomentosa.
pedunculata, Poir. Encyc. Suppl. iv. 590 = Dalea alopecuroides.
pentaphylla, *Linn. Sp. Pl.* 764.—Mexic.
phymatodes, Jacq. Ic. Pl. Rar. iii. t. 563 = Dalea phymatodes.
physodes, *Hook. Fl. Bor. Am.* i. 136.—Am. bor. occ.
pinnata, Andr. Bot. Rep. t. 474 = odoratissima.
pinnata, *Linn. Sp. Pl.* 762.—Afr. austr.
pinnata, Sieber, ex Walp. in Linnaea, xiii. (1839) 512 = filiformis.
plicaefolia, Eckl. & Zeyh. Enum. 231 = obtusifolia.
plicata, *Delile, Fl. Egypte*, 252. t. 37. f. 3.—Afr. trop.; Arab.; Ind. or.
plumosa, *F. Muell. Fragm.* iv. 22.—Austral.
plumosa, Reichb. Fl. Germ. Excurs. 869 = bituminosa.
polyphylla, *Eckl. & Zeyh. Enum.* 227.—Afr. austr.
polystachya, Poir. Encyc. Suppl. iv. 587 = dentata.
polysticta, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 150.—Afr. austr.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 3 (Quid?).—Ind. or.
prostrata, Linn. Pl. Afr. Rar. 15; Amoen. Acad. vi. 93 = Rhynchosia ferulaefolia.
pubescens, *Balb. ex Pers. Syn.* ii. 347.—Mexic.?
pubescens, *Poir. Encyc.* v. 686.—Peruv.
punctata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = Poiretia punctata.
purpurea, Poir. Encyc. v. 694 = Petalostemon violaceus.

PSORALEA:—
pustulata, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 54.—Austral.
racemosa, E. Mey. Comm. Pl. Afr. Austr. 87 = hamata.
racemosa, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
reclinata, Cav. Ic. i. 60 = Dalea reclinata.
repens, *Berg. Desc. Pl. Cap.* 223; *Linn. Mant.* ii. 265.—Afr. austr.
restioides, *Eckl. & Zeyh. Enum.* 226.—Afr. austr.
rhombifolia, *Torr. & Gray, Fl. N. Am.* i. 303.—Texas.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 463 = cuspidata.
rotundifolia, *Linn. f. Suppl.* 338.—Afr. austr.
rubescens, *Lour. Fl. Cochinch.* 444.—Cochinch.
rupicola, Eckl. & Zeyh. Enum. 230 = striata.
scabra, Nutt. in Torr. & Gray, Fl. N. Am. i. 300 = lanceolata.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 4.—Guatemala.
Schultzii, *F. Muell. Fragm.* ix. 155.—Austral.
scutellata, *Lour. Fl. Cochinch.* 443.—Cochinch.
sericea, Poir. Encyc. v. 687 = tomentosa.
sibirica, *Poir. Encyc. Suppl.* iv. 587.—Sibir.
simplex, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 303.—Am. bor.
simplicifolia, Baldw. ex Torr. & Gray. l. c. = virgata.
sinuata, Sieber, ex Walp. in Linnaea, xiii. (1839) 512 = filiformis.
spathulata, *E. Mey. Comm. Pl. Afr. Austr.* 85.—Afr. austr.
speciosa, Eckl. & Zeyh. Enum. 224 = pinnata.
spicata, Hort. ex Tausch, in Flora, xix. (1836) 421 = Tauschiana.
spicata, *Linn. Mant.* ii. 264.—Afr. austr.
spicata, Willd. ex Walp. in Linnaea, xiii. (1839) 514 = bracteata.
Stachydis, *Linn. f. Suppl.* 338.—Afr. austr.
Stachydis, Willd. ex Walp. in Linnaea, xiii. (1839) 515 = Indigofera cytisoides.
stachyera, Eckl. & Zeyh. Enum. 230 = bracteata.
Stachyos, Eckl. & Zeyh. l. c. 231 = hirta.
Stachyos, Thunb. Fl. Cap. ed. Schult. 608 = P. Stachydis.
stipulacea, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 466.—Ins. Timor.
stipulata, *Torr. & Gray, Fl. N. Am.* i. 688.—Am. bor.
striata, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
striata, Willd. ex Walp. in Linnaea, xiii. (1839) 515 = racemosa.
strobilina, *Hook. & Arn. Bot. Beech. Voy.* 332.—Calif.
subacaulis, *Torr. & Gray, Fl. N. Am.* i. 302.—Am. bor.
tamborensis, *Miq. Fl. Ind. Bat.* i. 1. 318.—Ins. Sunda.
taurica, Ledeb. ex Nym. Consp. 188 = bituminosa.
Tauschiana, *Steud. Nom.* ed. II. ii. 408.—Afr. austr.
tenax, *Lindl. in Mitch. Three Exped.* ii. 10.—Austral.
tenuiflora, *Pursh, Fl. Am. Sept.* ii. 475.—Am. bor.
tenuifolia, Eckl. & Zeyh. Enum. 225 = pinnata.
tenuifolia, *Linn. Sp. Pl.* 763.—Afr. austr.
tenuifolia, Thunb. Prod. Pl. Cap. 135 = fascicularis.
tenuifolia, Willd. ex Walp. in Linnaea, xiii. (1839) 513 = angustifolia.
tenuissima, *E. Mey. Comm. Pl. Afr. Austr.* 84.—Afr. austr.
Testariae, *F. Muell. Fragm.* v. 45.—Austral.
tetragonoloba, Linn. Mant. i. 104 = Cyamopsis psoralioides.
tetraphylla, Poir. Encyc. Suppl. iv. 589 = Poiretia psoralioides.
Thomii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 155.—Afr. austr.
Thunbergiana, Eckl. & Zeyh. Enum. 225 = fascicularis.
tomentosa, Cav. Ic. iii. 21 = Dalea verbenacea.
tomentosa, *Thunb. Prod. Pl. Cap.* 135.—Afr. austr.
triantha, *E. Mey. Comm. Pl. Afr. Austr.* 88.—Afr. austr.
triflora, DC. Prod. ii. 217 = axillaris.
triflora, Poir. Encyc. v. 683 = triantha.
triflora, Thunb. Prod. Pl. Cap. 135 = angustifolia.
tuberculata, Steud. Nom. ed. II. ii. 408, sphalm. = *Parosela tuberculata*.
uncinata, *Eckl. & Zeyh. Enum.* 231.—Afr. austr.
velutina, *E. Mey. Comm. Pl. Afr. Austr.* 89.—Afr. austr.

PSORALEA:—
venusta, *Eckl. & Zeyh. Enum.* 231.—Afr. austr.
verrucosa, Willd. Sp. Pl. iii. 1343 = angustifolia.
violacea, DC. Prod. ii. 243, in syn. = Petalostemon violaceus.
virgata, *Nutt. Gen. Am.* ii. 104.—Am. bor.
Weddelliana, *Baill. Adansonia*, ix. (1870) 292.—Bras.
Zeyheri, *Harv. Thes. Cap.* i. 51. t. 80.—Afr. austr.

PSOROPHYTUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 360 = **Hypericum**, Linn.
undulatum, Spach, l. c. = H. balearicum.

PSOROSPERMUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 157. *HYPERICINEAE*, Benth. & Hook. f. i. 167.
Afzelii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 579.—Afr. trop.
alternifolium, *Hook. f. in Hook. Niger Fl.* 243.—Afr. trop.
androsaemifolium, *Baker, in Journ. Bot.* xx. (1882) 19.—Madag.
axillare, *Benth. & Hook. f. Gen.* i. 167, in nota sub Haronga.—Madag.
brachypodum, *Baker, in Journ. Linn. Soc.* xx. (1883) 93.—Madag.
cerasifolium, *Baker, l. c.* xxi. (1884) 324.—Madag.
chionanthifolium, *Spach, in Ann. Sc. Nat.* Sér. II. v. (1836) 161.—Madag.
citrifolium, *Spach, l. c.* 159.—Madag.
cordiaefolium, *Spach, l. c.* 162.—Madag.
cornifolium, *Spach, l. c.* 161.—Madag.
cuspidatum, *Spach, l. c.* 162.—Madag.
discolor, *Baker, in Journ. Linn. Soc.* xxi. (1884) 323.—Madag.
discolor, *Spach, in Ann. Sc. Nat.* Sér. II. v. (1836) 160.—Madag.
Fanerana, *Baker, in Journ. Bot.* xx. (1882) 19.—Madag.
febrifugum, *Spach, in Ann. Sc. Nat.* Sér. II. v. (1836) 163.—Afr. trop.
ferrovestitum, *Baker, in Journ. Linn. Soc.* xx. (1883) 93.—Madag.
ferrugineum, Hook. f. in Hook. Niger Fl. 241 = febrifugum.
Forbesii, *Baker, in Journ. Linn. Soc.* xx. (1883) 94.—Madag.
leonense, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 578 = Haronga madagascariensis.
leptophyllum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 323.—Madag.
microcarpum, *Baker, l. c.* xx. (1883) 95.—Madag.
niloticum, *Kotschy, in Schweinf. Beitr. Fl. Aethiop.* 285.—Afr. trop.
pauciflorum, *Baker, in Journ. Linn. Soc.* xx. (1883) 94.—Madag.
senegalense, *Spach, in Ann. Sc. Nat.* Sér. II. v. (1836) 164.—Afr. trop.
tenuifolium, *Hook. f. in Hook. Ic. Pl.* t. 771.—Afr. trop.
tenuifolium, Kotschy, in Oestr. Bot. Zeitschr. xiv. (1864) 338 = niloticum.
trichophyllum, *Baker, in Journ. Linn. Soc.* xxi. (1884) 323.—Madag.
venulosum, *Baker, l. c.* xx. (1883) 93.—Madag.

PSYCHANTHUS, Spach, Hist. Vég. Phan. vii. 130 (1839) = **Polygala**, Tourn.
bracteolatus, Spach, l. c. 140 = Polyg. bracteolata, *Linn.*
grandiflorus, Spach, l. c. 135 = Polyg. myrtifolia.
myrtifolius, Spach, l. c. 137 = Polyg. myrtifolia.
oppositifolius, Spach, l. c. 133 = Polyg. oppositifolia.
speciosus, Spach, l. c. 138 = Polyg. virgata.

PSYCHECHILUS, Breda, Orch. Kuhl. et Hass. (1827) [t. 9] = **Zeuxine**, Lindl. (Orchid.).
gracilis, Breda, l. c. = Z. gracilis.
purpurascens, Blume, Orch. Arch. Ind. 105. t. 9. f. 3 = Hetaeria purpurascens.
violaceus, Hass. ex Blume, l. c. 105 = Hetaeria purpurascens.

PSYCHILIS, Rafin. Fl. Tellur. iv. 40 (1836) = **Epidendrum**, Linn. (Orchid.).
amena, Rafin. l. c. = E. bifidum, *Aubl.*

PSYCHINE, Desf. Fl. Atlant. ii. 68. t. 148 (1800). *CRUCIFERAE*, Benth. & Hook. f. i. 89.
arabica, Spreng. Syst. ii. 880 = Schouwia arabica.
numidica, Spreng. l. c. = stylosa.
stylosa, *Desf. Fl. Atlant.* ii. 69. t. 148.—Afr. bor.

PSYCHOPSIS, Rafin. Fl. Tellur. iv. 40 (1836) = **Oncidium**, Sw. (Orchid.).
picta, Rafin. l. c. = O. Papilio.

PSYCHOSPERMA, Dum. Anal. Fam. 56 (1829) = **Ptychosperma**, Labill. (Palm.).

PSYCHOTHRIA, Linn. Amoen. Acad. v. 395 (1760) = seq.

PSYCHOTRIA, Linn. Syst. ed. X. 929 (1759). *RUBIACEAE*, Benth. & Hook. f. ii. 123.
GALVANIA, Vand. Fl. Lusit. 15. t. 1. f. 7 (1788).
GRUMILEA, Gaertn. Fruct. i. 138. t. 28. f. 2 (1788).
GRUNDLEA, Steud. Nom. ed. II. i. 708 (1840).
HYLACIUM, Beauv. Fl. Owar. ii. 83. t. 113 (1807).
IPECACUANHA, Arruda, Diss. 44 (1810).
MAPOURIA, Aubl. Pl. Gui. i. 175. t. 67 (1775).
MYRSTIPHYLLUM, P. Br. Hist. Jamaic. 152 (1756).
PATABEA, Aubl. Pl. Gui. i. 110. t. 43 (1775).
PLEUREIA, Rafin. Sylva Tellur. 147 (1838).
POLYOZUS, Blume, Bijdr. 947 (1826).
PSYCHOTROPHUM, P. Br. Hist. Jamaic. 160. t. 13. f. 1, 2 (1756).
RONABEA, Aubl. Pl. Gui. i. 154. t. 59 (1775).
SIMIRA, Aubl. l. c. 170. t. 65 (1775).
STREBLOSA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 245.
STREMPELIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 180. t. 18. f. 3.
SULCANUX, Rafin. Sylva Tellur. 148 (1838).
VISCOIDES, Jacq. Select. Am. 73. t. 51 (1763).
ZWAARDEKRONIA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 252.
abrupta, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 205.—Afr. trop.
acuminata, *Benth. Bot. Voy. Sulph.* 107.—N. Granat.
acutiflora, *DC. Prod.* iv. 506.—Am. austr.
adenophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 166.—Ind. or.; Burma.
aeneofusca, *Muell. Arg. in Mart. Fl. Bras*, vi. v. 253.—Bras.
affinis, *Baker, Fl. Maurit.* 155.—Ins. Seychelles.
Afzelii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 205.—Afr. trop.
agnata, *DC. Prod.* iv. 505.—Ins. S. Doming.
alba, *Ruiz & Pav. Fl. Per.* ii. 58. t. 205. f. a.—Am. austr.
alpina, Poir. Encyc. v. 702 = Palicourea eriantha.
alpina, Sw. Prod. Veg. Ind. Occ. 44 = Palicourea alpina.
amazonica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 225.—Bras.
ambigua, Wight & Arn. Prod. 433 = Chasalia curviflora.
amethystina, Ruiz & Pav. Fl. Per. ii. 57 = Palicourea amethystina.
ampla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 462.—Bras.
ampla, Wall. Cat. n. 8333 (= Pavettae sp.).—Ind. or.
amplectens, *Benth. in Hook. Journ. Bot.* iii. (1841) 230.—Bras.
amplifolia, Hort. ex Steud. Nom. ed. I. 664 = Geophila violacea.
anacardifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = Palicourea macrocarpa.
anamallayana, *Bedd. Ic. Pl. Ind. Or.* 57. t. 235.—Ind. or.
anceps, *H. B. & K. Nov. Gen. et Sp.* iii. 360.—N. Granat.
andamanica, *Kurz, in Journ. Bot.* xiii. (1875) 328.—Ins. Andaman. et Nicob.
androsaemifolia, *Griseb. Fl. Brit. W. Ind.* 341.—Ind. occ.
anetoclada, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 206.—Afr. trop.
angulata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 243.—Burma; Malaya.
angustata, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 193.—Ins. Galapag.
angustifolia, *G. Don, Gen. Syst.* iii. 595 (sp. dub).—Afr. trop.

PSYCHOTRIA:—
angustifolia, *Poir. Encyc.* v. 703.—Ins. S. Doming.
anisocephala, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 332.—Bras.
anisoloba, *Muell. Arg. l. c.* 237.—Bras.
Ansellii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 214.—Afr. trop.
Antherura, Roem. & Schult. Syst. v. 188 = Antherura rubra.
apiculata, *Muell. Arg. in Flora*, lix. (1876) 549.—Bras.
apocynifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 47.—Bras.
apodantha, *A. Gray, l. c.* 46.—Ins. Samoa.
appendiculata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 350.—Bras.
arabica, *Klotzsch, in Sitzb. Acad. Wien, Math. Nat.* lii. *Abth.* i. (1866) 257.—Arabia.
arborea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 202.—Afr. trop.
arcuata, Benth. in Hook. Journ. Bot. iii. (1841) 229 = inundata.
ardisiaefolia, H. B. & K. Nov. Gen. et Sp. iii. 359 = alba.
arenosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 309.—Bras.
argentata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 243.—Java.
aristata, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 191.—Am. austr.
Arnottiana, Steud. Nom. ed. II. ii. 408 = connata.
Arrabidae, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 381.—Bras.
asiatica, Linn. Syst. ed. X. 929; Lam. Illustr. ii. 243. = Brownei.
asiatica, Poir. Encyc. v. 696 = glabrata.
asiatica, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 160 = calocarpa.
attenuata, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 192.—Am. austr.
aurantiaca, Muell. Arg. in Mart. Fl. Bras. vi. v. 244, in obs. = Palicourea crocea.
aurantiaca, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 165 = malayana.
Aureola, *Bartl. ex DC. Prod.* iv. 513.—Mexic.
auriculata, *Wright, ex Griseb. Cat. Pl. Cub.* 136.—Cuba.
axillaris, *Vell. Fl. Flum.* 67; ii. t. 32.—Bras.
axillaris, *Willd. Sp. Pl.* i. 962.—Guian.
bahiensis, DC. Prod. iv. 509 = cuspidata.
bahiensis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 338.—Bras.
Balbisiana, *DC. Prod.* iv. 517.—Jamaic.
bantamensis, *Miq. Fl. Ind. Bat.* ii. 288.—Java.
barbata, *Poir. Encyc.* v. 704.—Ins. Martinic.
barbata, *Span. in Linnaea*, xv. (1841) 319.—Ins. Timor.
barbiflora, *DC. Prod.* iv. 509.—Bras.
barcellana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 369.—Bras.
barraënsis, *Muell. Arg. l. c.* 254.—Bras.
Benthamiana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 204.—Afr. trop.
Berteriana, *DC. Prod.* iv. 515.—Ind. occ.
Beskeana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 360.—Bras.
Beyrichiana, *Muell. Arg. in Flora*, lix. (1876) 542, 545.—Bras.
bialata, *Wright, ex Griseb. Cat. Pl. Cub.* 138.—Cuba.
biaristata, *Bartl. ex DC. Prod.* iv. 513.—Mexic.
biattenuata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 465.—Bras.
bicolor, Bredem. ex Roem. & Schult. Syst. v. 191 = Palicourea crocea.
bicornis, *Wall. ex Voigt, Hort. Suburb. Calc.* 393, nomen.—Reg. Himal.
bidentata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 209.—Afr. trop.
bifaria, *Hiern, l. c.* 198.—Afr. trop.
bisulcata, *Wight & Arn. Prod.* 434.—Ind. or.
biternata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 366.—Bras.
Blanchetiana, *Muell. Arg. l. c.* 238.—Bras.
borjensis, *H. B. & K. Nov. Gen. et Sp.* iii. 357.—Venezuela.
brachiata, D. Dietr. Syn. Pl. i. 770, in syn., sphalm. = *bracteata*, F. G. Dietr.

PSYCHOTRIA:—
brachiata, Spreng. ex DC. Prod. iv. 516 = Rudgea caribaea.
brachiata, *Sw. Prod. Veg. Ind. Occ.* 45.—Ind. occ.
brachyandra, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 241.—Bras.
brachyantha, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 196.—Afr. trop.
brachybotrya, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 327.—Bras.
brachyceras, *Muell. Arg. in Flora*, lix. (1876) 544, 546.—Bras.
brachygyne, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 287.—Bras.
brachyloba, *Muell. Arg. l. c.* 243.—Bras.
brachypodum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = brevipes.
Brackenridgii, *A. Gray, in Proc. Am. Acad.* iv. (1860) 44.—Ins. Fiji.
bracteata, *DC. Prod.* iv. 510.—Guian.
bracteata, D. Dietr. Syn. Pl. i. 765 = Rudgea caribaea.
bracteata, F. G. Dietr. Vollst. Lexik. Gaertn. vii. 617 = Palicourea laxa.
bracteata, Wight & Arn. Prod. 434 = Dalzellii.
bracteocardia, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 362.—Bras.
bracteolata, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 228.—Mexic.
bracteosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 207.—Afr. trop.
brasiliensis, *Vell. Fl. Flum.* 66; ii. t. 29.—Bras.
Brassii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 204.—Afr. trop.
brevicollis, *Muell. Arg. in Flora*, lix. (1876) 548, 552.—Bras.
breviflora, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 263.—Bras.
brevipedunculata, *Muell. Arg. l. c.* 258.—Bras.
brevipes, *DC. Prod.* iv. 511.—Guian.
Browerii, *Seem. Fl. Vit.* 135.—Ins. Fiji.
Brownei, *Spreng. Syst.* i. 742.—Ind. occ.
brunnea, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 201.—Afr. trop.
bullata, *Seem. Fl. Vit.* 137.—Ins. Fiji.
Burchelliana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 364.—Bras.
byrsophylla, Spreng. Syst. i. 747 = rigida.
caerulea, Ruiz & Pav. Fl. Per. ii. 62 = Palicourea coerulea.
cajannensis, Desv. ex Ham. Prod. Fl. Ind. Occ. 29 = Palicouria pedunculosa.
calocarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 315.—Reg. Himal.; Burma.
caloneura, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 359.—Bras.
calva, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 199.—Afr. trop.
calycina, *H. B. & K. Nov. Gen. et Sp.* iii. 356.—N. Granat.
calycosa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Fiji.
campyloneura, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 302.—Bras.
capensis, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 230.—Afr. austr.
capitata, *Ruiz & Pav. Fl. Per.* ii. 59. t. 206. f. a.—Peruv.
capitata, Sieber, ex DC. Prod. iv. 531 = Chasalia capitata.
capitellata, DC. l. c. 514 = brachiata?
Carronis, *C. Moore & F. Muell. Fragm.* vii. 49.—Austral.
carthagenensis, *Jacq. Enum. Pl. Carib.* 16.—Am. austr.
casiquiaria, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 324.—Bras.
castellana, *Muell. Arg. l. c.* 261.—Bras.
cataractarum, *Muell. Arg. l. c.* 226.—Bras.
celastroides, *Griseb. Fl. Brit. W. Ind.* 341.—Ind. occ.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 210.—Ins. Celebes.
cernua, *Nadeaud, Enum. Pl. Tahiti*, 51.—Ins. Tahiti.
cernua, Steud. Nom. ed. II. ii. 408 = barbiflora.
chaenotricha, *DC. Prod.* iv. 509.—Bras.
chimarrhoides, DC. l. c. 514 = lanceolata, *Nutt.*
chlorantha, *Benth. in Hook. Journ. Bot.* iii. (1841) 225.—Guian.
chlorantha, Buch.-Ham. ex Wall. Cat. n. 8336 B = denticulata.

PSYCHOTRIA:—
chlorophylla, *Muell. Arg. in Flora*, lix. (1876) 543.—Bras.
chlorotica, *Muell. Arg. l. c.* 542, 545.—Bras.
chontalensis, Seem. in Bull, Cat. (1870) 5.—Cf. Journ. Bot. xix. (1881) 188 = pilosa.
chrysorhiza, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 111 = Morinda citrifolia.
ciliolata, *Schlecht. in Linnaea*, xxviii. (1856) 519.—Bras.
circinata, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = Palicourea elliptica.
citrifolia, Sw. Prod. Veg. Ind. occ. 43 = Rudgea caribaea.
clavipes, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 464.—Bras.
closterocarpa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 44.—Ins. Samoa.
coccinea, Poit. ex DC. Prod. iv. 529 = Palicourea coccinea.
coerulea, Steud. Nom. ed. II. ii. 409 = *caerulea*.
collina, Labill. Sert. Austro-Caled. 47. t. 47 = Brownei.
colorata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 372.—Bras.
comitis, *Muell. Arg. l. c.* 247.—Bras.
commutata, *DC. Prod.* iv. 518.—Am. austr.
concolor, *Benth. & Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 31.—Nicaragua.
conferta, *Benth. Bot. Voy. Sulph.* 107.—N. Granat.
confertiflora, Miq. Fl. Ind. Bat. ii. 285 = Korthalsiana.
congesta, *Hook. f. Fl. Brit. Ind.* iii. 162.—Ind. or.
congesta, *Spreng. ex DC. Prod.* iv. 515.—Jamaic.
conjungens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 264.—Bras.
connata, Kurz, For. Fl. Brit. Burm. ii. 10 = adenophylla.
connata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 163.—Ind. or.
consanguinea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 252.—Bras.
constricta, *Muell. Arg. in Flora*, lix. (1876) 544, 546.—Bras.
contracta, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 330.—Bras.
cordata, *A. Gray, in Proc. Am. Acad.* iv. (1860) 46.—Ins. Fiji.
cordifolia, F. G. Dietr. Vollst. Lexik. Gaertn. vii. 618 = Geophila macropoda.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 365.—Bras.
coriacea, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 242.—Borneo.
coriacea, *Poir. Encyc.* v. 703.—Am. austr.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 362.—Venezuela.
cornifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 191 = orinocensis.
cornigera, *Benth. in Hook. Journ. Bot.* iii. (1841) 227.—Ins. Trinit.; Guian.
cornuta, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 198.—Afr. trop.
coronata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 508.—Ind. occ.
coronata, Vell. Fl. Flum. 65; ii. t. 25 = Rudgea coronata.
corymbifera, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 247.—Bras.
corymbosa, Sieber, ex DC. Prod. iv. 523 (non Rubiacea).—Hab.?
corymbosa, *Sw. Prod. Veg. Ind. Occ.* 44.—Ind. occ.
costaricensis, *Polak. in Linnaea*, xli. (1877) 571.—Costa Rica.
costato-venosa, *Schlecht. l. c.* xxviii. (1856) 497.—Peruv.
costivenia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 508.—Cuba.
crassa, *Benth. in Hook. Journ. Bot.* iii. (1841) 227.—Guian.
crassifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 205.—Borneo.
crispa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 212.—Afr. trop.
cristata, *Hiern, l. c.* 205.—Afr. trop.
crocea, Sw. Prod. Veg. Ind. Occ. 44 = Palicourea crocea.
croceoides, Desv. ex Ham. Prod. Pl. Ind. Occ. 29 = Palicourea croceoides.
cujabensis, *Schlecht. in Linnaea*, xxviii. (1856) 507.—Bras.

PSYCHOTRIA:—
cumanensis, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 191.—Venezuela.
cuneifolia, *DC. Prod.* iv. 507.—Peruv.
curviflora, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 167 = Chasalia curviflora.
cuspidata, *Bredem. ex Roem. & Schult. Syst.* v. 192.—Venezuela.
cuspidella, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 209.—Sumatra; Borneo.
cyanea, Vell. Fl. Flum. 68; ii. t. 36 = Faramea salicifolia.
cyanocarpa, [*Seem*].—*Cf. Gard. Chron.* (1869) 1236.—Nicaragua.
cyanococca, [Seem. in] Floral Mag. (1870) t. 479 = praec.
cymosa, *Ruiz & Pav. Fl. Per.* ii. 59. t. 206. f. b.—Peruv.
Dallachiana, *Benth. Fl. Austral.* iii. 427.—Austral.
Dalzellii, *Hook. f. Fl. Brit. Ind.* iii. 163.—Ind. or.
daphnoides, *A. Cunn. ex Hook. Bot. Mag.* t. 3228.—Austral.
dasyophthalma, *Griseb. Fl. Brit. W. Ind.* 341.—Jamaic.
debilis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 346.—Bras.
decidua, Vell. Fl. Flum. 67; ii. t. 34 = alba.
decipiens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 241.—Bras.
deflexa, *DC. Prod.* iv. 510.—Guian.
densecostata, *Muell. Arg. in Flora*, lix. (1876) 544, 546.—Bras.
densevenosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 283.—Bras.
densiflora, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 189 = alba.
denticulata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 166.—Reg. Himal.; Burma.
dichotoma, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 190.—Am. austr.
diervilloides, *H. B. & K. Nov. Gen. et Sp.* iii. 358.—Am. austr.
diplosphaerica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 465.—Bras.
divaricata, Blume, Cat. Gew. Buitenz. 53 = divergens, *Blume*.
divaricata, *H. B. & K. Nov. Gen. et Sp.* iii. 362.—Ecuador.
divaricata, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 191 = glaucescens.
divergens, *Blume, Bijdr.* 959.—Java.
divergens, *Kurz, For. Fl. Brit. Burm.* ii. 13 (= sulcata?)—Burma.
diversicolor, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 252.—Bras.
Dodda-Jorrane, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 171 = bisulcata.
domingensis, Jacq. Enum. Pl. Carib. 16 = Palicourea domingensis.
Doniana, *Benth. in Hook. Niger Fl.* 419.—Afr. trop.
dura, *Sw. ex. Griseb. Fl. Brit. W. Ind.* 340.—Jamaic.
Eckloniana, *F. Muell. Extra-trop. Pl. Ind.* ed. [III.] (1880) 264.—Afr. austr.
egensis, *Muell. Arg. in Flora*, lix. (1876) 542, 545.—Bras.
elliptica, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 189 = commutata.
elliptica, *Ker-Gawl. in Bot. Reg. t.* 607.—Bras.
elongata, Bedd. Fl. Sylv. Anal. Gen. t. 17 = congesta, *Hook. f.*
elongata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 32.—Nicaragua.
elongata, *Hook. f. Fl. Brit. Ind.* iii. 163.—Ind. or.
emetica, *Linn. f. Suppl.* 144.—N. Granat.
emetica, Vell. Fl. Flum. 64; ii. t. 22 = P. Ipecacuanha.
erratica, *Hook. f. Fl. Brit. Ind.* iii. 168.—Reg. Himal.
erythrocarpa, *Schlecht. in Linnaea*, ix. (1834) 595.—Mexic.
erythrophylla, *Muell. Arg. in Flora*, lix. (1876) 542, 546.—Bras.
estrellana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 382.—Bras.
euneura, *Miq. Fl. Ind. Bat.* ii. 284.—Java.
evenia, *Wright, ex Griseb. Cat. Pl. Cub.* 136.—Cuba.
exannulata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 257.—Bras.
excelsa, *H. B. & K. Nov. Gen. et Sp.* iii. 355.—Mexic.

PSYCHOTRIA:—
expansa, Blume, Bijdr. 963 = malayana.
exserta, *DC. Prod.* iv. 517.—Ins. S. Doming.
extensa, *Miq. Fl. Ind. Bat.* ii. 288.—Java.
extratropica, *Muell. Arg. in Flora*, lix. (1876) 544.—Bras.
fastigiata, Spreng. Syst. i. 742 = Palicourea fastigiata.
febrifuga, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 32. t. 238.—Bras.
ferruginea, *Baker, Fl. Maurit.* 156.—Ins. Seychelles.
fetens, DC. Prod. iv. 516, sphalm. = *foetens*.
ficigemma, *DC. l. c.* 510.—Guian.
filipes, *A. Gray, in Proc. Am. Acad.* iv. (1860) 46.—Ins. Fiji.
filipes, *Hook. f. Fl. Brit. Ind.* iii. 170.—Zeylan.
fimbriata, Benth. in Hook. Journ. Bot. iii. (1841) 226 = Rudgea micrantha.
fissistipula, *Muell. Arg. in Flora*, lix. (1876) 547, 551.—Bras.
Fitzalani, *Benth. Fl. Austral.* iii. 428.—Austral.
flavescens, Spreng. Syst. i. 744 = Palicourea flavescens.
flavicans, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 339.—Bras.
flexuosa, *Willd. Sp. Pl.* i. 966.—Guian.; Bras.
floribunda, *H. B. & K. Nov. Gen. et. Sp.* iii. 359.—Am. austr.
fluminensis, *Vell. Fl. Flum.* 66; ii. t. 30.—Bras.
Fockeana, *Miq. in Linnaea*, xviii. (1844) 296.—Guian.
foetens, Sw. Prod. Veg. Ind. Occ. 43 = Brownei, foetida.
foetida, *Griseb. Fl. Brit. W. Ind.* 342.—Ind. occ.
foliosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 197.—Afr. trop.
formosa, *Cham. & Schlecht. in Linnaea*, iv. (1829) 24.—Bras.
Forsteriana, *A. Gray, in Proc. Am. Acad.* iv. (1860) 44.—Ins. Pacif.
forsteronioides, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
foveolata, *Ruiz & Pav. Fl. Per.* ii. 59. t. 207. f. b.—Peruv.
foveolata, Spreng. Syst. i. 746 = Coffea foveolata.
Franquevilleana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 325.—Bras.
fraterna, *Muell. Arg. l. c.* 251.—Bras.
fulgens, *Muell. Arg. l. c.* 257.—Bras.
fulva, *Buch.-Ham. ex Wall. Cat.* n. 8336.—Reg. Himal.
fulva, Wall. l. c. n. 8336. C = silhetensis.
furcata, *DC. Prod.* iv. 512.—Panama.
furcellata, *Vatke, in Bremen, Abh.* ix. (1885) 118.—Madag.
gabonica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 201.—Afr. trop.
gabrielis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 335.—Bras.
gandarussaefolia, DC. Prod. iv. 521, sphalm. = gendarussaefolia.
Gardneri, *Hook. f. Fl. Brit. Ind.* iii. 161.—Zeylan.
Gardneriana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 379.—Bras.
gendarussaefolia, *Blume, Bijdr.* 960.—Java.
glabrata, Bartl. ex DC. Prod. iv. 507 = cuneifolia.
glabrata, Sieber, ex DC. l. c. 514 = tenuifolia.
glabrata, *Sw. Prod. Veg. Ind. Occ.* 43.—Jamaic.
glabrescens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 363.—Bras.
glandulifera, *Thw. ex Hook. f. Fl. Brit. Ind.* iii. 161.—Zeylan.
glauca, *Polak. in Linnaea*, xli. (1877) 569.—Costa Rica.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* iii. 358.—Venezuela.
Glaziovii, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 267.—Bras.
globosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 208.—Afr. trop.
glomerata, *H. B. & K. Nov. Gen. et Sp.* iii. 362.—Am. austr.
goyazensis, *Muell. Arg. in Flora*, lix. (1876) 548, 551.—Bras.
gracilenta, *Muell. Arg. l. c.* 542, 545.—Bras.
graciliflora, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 35.—Costa Rica.
gracilis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Fiji.
gracilis, Ruiz & Pav. Fl. Per. ii. 63 = Geophila gracilis.

PSYCHOTRIA:—
granadensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 34.—Nicaragua.
grandiflora, *H. Mann, in Proc. Am. Acad.* vii. (1867) 170.—Ins. Sandvic.
grandifolia, *Hoffmgg. ex Roem. & Schult. Syst.* v. 190.—Bras.
grandifolia, Sieber, ex DC. Prod. iv. 532 = Chasalia grandifolia.
grandis, *Sw. Prod. Veg. Ind. Occ.* 43.—Am. trop.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 171.—Malacca.
guianensis, Raeusch. Nom. ed. III. 56 = Palicourea guianensis.
Haenkeana, *DC. Prod.* iv. 507.—Am. austr.
hancorniaefolia, *Benth. in Linnaea*, xxiii. (1850) 463.—Bras.
hastisepala, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 350.—Bras.
hebeclada, *DC. Prod.* iv. 513.—Mexic.
hebecladoides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 509.—Cuba.
hederacea, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 191 = Geophila violaefolia.
Helferiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 314.—Burma; Malaya.
herbacea, Jacq. Enum. Pl. Carib. 16; Linn. Sp. Pl. ed. II. 245 = Geophila reniformis.
herbacea, *Vell. Fl. Flum.* 64; ii. t. 21.—Bras.
heterocephala, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 333.—Bras.
heteromorpha, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 244.—Sumatra.
hexandra, *H. Mann, in Proc. Am. Acad.* vii. (1867) 170.—Ins. Sandvic.
Heyneana, Wall. Cat. n. 8327 = Thwaitesii, truncata.
hirsuta, *Sw. Prod. Veg. Ind. Occ.* 42.—Jamaic.
hirta, H. B. & K. Nov. Gen. et Sp. iii. 364 = rufescens.
hirta, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 191.—N. Granat.
hirtinervia, *Wawra, in Oestr. Bot. Zeitschr.* xxix. (1879) 216.—Bras.
hirtipes, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 379.—Bras.
Hoffmannseggiana, *Muell. Arg. l. c.* 336.—Bras.
homalosperma, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 393.—Ins. Bonin.
Hookeri, G. Don, Gen. Syst. iii. 585 = Coprosma triflora.
Hookeriana, Steud. Nom. ed. II. ii. 409 = praec.
horizontalis, Spreng. ex DC. Prod. iv. 514 = oligotricha.
horizontalis, *Sw. Prod. Veg. Ind. Occ.* 44.—Ind. occ.
horridula, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 344.—Bras.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 285.—Java.
Humboldtiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 333.—Bras.
humilis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 198.—Afr. trop.
hundensis, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 189 = alba.
hyacinthiflora, Ruiz & Pav. Fl. Per. ii. 62 = Palicourea hyacinthiflora.
hyacinthoides, Pers. Syn. i. 208 = praec.
hygrophiloides, *Benth. in Linnaea*, xxiii. (1850) 465.—Bras.
hypargyraea, *A. Gray, in Proc. Am. Acad.* iv. (1860) 46.—Ins. Fiji.
hyptoides, *Benth. in Hook. Journ. Bot.* iii. (1841) 231.—Guian.
ignea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 244.—Bras.
inaequifolia, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
infundibularis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 209.—Afr. trop.
insidens, *Hiern, l. c.* 208.—Afr. trop.
insularum, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Pacif.
interjecta, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 357.—Bras.
intermedia, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 175.—Bras.
intrudens, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 206.—Java.

PSYCHOTRIA:—
inundata, *Benth. in Hook. Journ. Bot.* iii. (1841) 229.—Bras.; Guian.
involucellaris, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 380.—Bras.
involucrans, *Muell. Arg. l. c.* 365.—Bras.
involucrata, *A. Rich. Fl. Cub. Fanerog.* ii. 27.—Cuba.
involucrata, Sw. Prod. Veg. Ind. Occ. 45 = officinalis.
involucrata, Willd. ex DC. Prod. iv. 508 = flexuosa.
iodotricha, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 375.—Bras.
Ipecacuanha, *Stokes, Bot. Mat. Med.* i. 365.—Bras.
ixoroides, *Bartl. ex DC. Prod.* iv. 522.—Ins. Philipp.
Jackii, *Hook. f. Fl. Brit. Ind.* iii. 167.—Malaya.
jambosioides, *Schlecht. in Linnaea*, xxviii. (1856) 518.—Bras.
janeirensis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 357.—Bras.
japurensis, *Muell. Arg. l. c.* 328.—Bras.
japurensis, *Muell. Arg. l. c.* 344.—Reg. Amazon.
jasminiflora, *Mast. in Gard. Chron.* (1879) II. 200; *Hook. f. Bot. Mag.* t. 6454.—Bras.
Johnsoni, *Hook. f. Fl. Brit. Ind.* iii. 175.—Ind. or.
justicoides, *Schlecht. in Linnaea*, ix. (1834) 596.—Mexic.
Kirkii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 206.—Afr. trop.
Kolly, *Schum. & Thonn. Beskr. Guin. Pl.* 110.—Afr. trop.
konguensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 200.—Afr. trop.
Korthalsiana, *Miq. Fl. Ind. Bat.* ii. 293.—Java; Sumatra.
laciniata, *Vell. Fl. Flum.* 66; ii. t. 28.—Bras.
laevigata, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = Palicourea laevigata.
laevigata, Wight & Arn. Prod. 433 = connata.
laevis, *DC. Prod.* iv. 505.—Porto Rico.
lagoënsis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 278.—Bras.
lamprophylla, *Spreng. Syst.* i. 747.—Am. austr.
lanata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 249.—Bras.
lanceolata, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 15; *et in Baker, Fl. Maurit.* 156.—Ins. Rodrigues.
lanceolata, *Nutt. in Am. Journ. Sc.* Ser. I. v. (1822) 290.—Florida.
lanceolata, Vell. Fl. Flum. 65; ii. t. 27 = Coussarea lanceolata.
Langsdorffiana, *Muell. Arg. in Flora*, lix. (1876) 541, 545.—Bras.
lasiophthalma, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 508.—Cuba.
lasiopus, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 297.—Bras.
lasiostylis, *Muell. Arg. in Flora*, lix. (1876) 548, 551.—Bras.
latifolia, Blume, Cat. Gew. Buitenz. 54 = robusta.
latifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 189.—Am. austr.
latistipula, *Benth. in Hook. Niger Fl.* 419.—Afr. trop.
laurifolia, Bert. ex DC. Prod. iv. 514 = tenuifolia.
laurifolia, Spreng. ex DC. l. c. 516 = polyantha.
laurifolia, Sw. Prod. Veg. Ind. Occ. 43 = glabrata.
laxa, Ruiz & Pav. Fl. Per. ii. 61 = Palicourea laxa.
laxa, *Sw. Fl. Ind. Occ.* i. 407.—Jamaic.
laxiflora, *Blume, Bijdr.* 964.—Java.
leiocarpa, *Cham. & Schlecht. in Linnaea*, iv. (1829) 22.—Bras.
leiocarpa, Mart. in Flora, xxi. (1838) II. Beibl. 61 = Rudgea leiocarpoides.
lepidocarpa, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 242.—Sumatra.
leptophylla, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 200.—Afr. trop.
leptothyrsa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 208.—Ins. Ceram.
leucocarpa, *Blume, Bijdr.* 964.—Java.
leucocephala, Brongn. in Lem. Hort. Univ. vi. (1845) 257 = Rudgea macrophylla.
leucophaea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 33.—Bras.
Lhotzkyana, Schlecht. in Linnaea, xxviii. (1856) 505 = flexuosa.

PSYCHOTRIA:—
ligustrina, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 188 = borjensis.
Lindleyana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 265. —Bras.
linearis, *Bartl. ex DC. Prod.* iv. 522.—Ins. Philipp.
lineata, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 191 = Palicourea apicata.
lineata, Spreng. Syst. i. 746 = oligantha, *DC.*
longepedunculata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 237.—Bras.
longevaginalis, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 201.—Afr. trop.
longicollis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 33 = Costa Rica.
longicuspis, *Muell. Arg. in Flora*, lix. (1876) 549, 552. —Bras.
longiflora, Poir. Encyc. v. 104 = Palicourea longiflora.
longiflora, *Willd. Sp. Pl.* i. 971.—Guian.
longifolia, Dalz. in Hook. Kew Journ. ii. (1850) 133 = Chasalia curviflora.
longifolia, *Hoffmgg. ex Roem. & Schult. Syst.* v. 190. —Bras.
longifolia, Spreng. Syst. i. 745 = Palicourea longifolia.
longipes, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 354.—Bras.
longipetiolata, *Thw. Enum. Pl. Zeyl.* 149.—Zeylan.
longistipula, *Benth. in Hook. Journ. Bot.* iii. (1841) 227.—Bras.
longistipulata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 248.—Bras.
longistylis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 209.—Afr. trop.
loniceroides, *Sieber, ex DC. Prod.* iv. 523.—Austral.
lophoclada, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 197.—Afr. trop.
lucens, *Hiern, l. c.* iii. 211.—Afr. trop.
lucida, *H. B. & K. Nov. Gen. et Sp.* iii. 361. t. 288.—Am. austr.
lucida, Spreng. ex DC. Prod. iv. 517 = Balbisiana.
lucida, Willd. ex Spreng. Syst. i. 740 = splendens.
lucidula, *Baker, in Journ. Linn. Soc.* xxi. (1885) 413. —Madag.
lupulina, *Benth. in Hook. Journ. Bot.* iii. (1841) 230. —Bras.; Guian.
lurida, Blume, Bijdr. 959 = Chasalia curviflora.
Luschnathiana, Schlecht. in Linnaea, xxviii. (1856) 512 = seq.
Luschnathii, *Klotzsch, ex Mart. in Flora*, xxiv. (1841) II. *Beibl.* 71.—Bras.
lutea, Sieber, ex DC. Prod. iv. 524, 622 (Apocynacea).
lutea, Spreng. ex DC. l. c. 530 = Palicourea barbinervia.
lutea, Willd. Sp. Pl. i. 971 (= *Nonatelia lutea*).—Guian.
luteo-virescens, Pers. Syn. i. 208 = Palicourea laxa.
luzoniensis, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 112.—Ins. Philipp.
macrantha, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 377. —Bras.
macrobotrys, Ruiz & Pav. Fl. Per. ii. 57 = Palicourea macrobotrys.
macrocalyx, *A. Gray, in Proc. Am. Acad.* iv. (1860) 46.—Ins. Fiji.
macrocarpa, *Hook. f. Fl. Brit. Ind.* iii. 162.—Ind. or.
macrophylla, *Ruiz & Pav. Fl. Per.* ii. 56. t. 202. f. a. —Peruv.
macrophylla, Vent. ex DC. Prod. iv. 517 = platyphylla.
macropoda, Ruiz & Pav. Fl. Per. ii. 63 = Geophila macropoda.
macrothyrsa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 225. —Bras.
madraspatana, *Hook. f. Fl. Brit. Ind.* iii. 167.—Ind. or.
magnoliaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 360.—Ecuador.
magnoliaefolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = Willdenowii.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 166.—Singapore.
malaneoides, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
malayana, *Jack, in Malay. Misc.* i. (1820) I. 3.—Malaya.
malayana, F. Villar, ex Vidal, Sinops. t. 57. f. F = P. Tacpo.
mamillaris, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 299. —Bras.
mandiocana, *Muell. Arg. in Flora*, lix. (1876) 545, 546.—Bras.
manillensis, *Bartl. ex DC. Prod.* iv. 522.—Ins. Philipp.
Mannii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 197.—Afr. trop.

PSYCHOTRIA:—
Mansoana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 243. —Bras.
mapourioides, *DC. Prod.* iv. 509.—Guian.
Mapuria, Roem. & Schult. Syst. v. 187 = nitida.
maranhana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 361. —Bras.
Marcgravii, *Spreng. Syst.* iv. *Cur. Post.* 79.—Bras.
marginata, Schlecht. in Linnaea, xxviii. (1856) 505 = Rudgea leiocarpoides.
marginata, *Sw. Prod. Veg. Ind. Occ.* 43.—Am. trop.
mariana, *Bartl. ex DC. Prod.* iv. 522.—Ins. Marian.
Martiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 339.—Bras.
Martiusii, *Muell. Arg. l. c.* 240.—Bras.
maypurensis, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 190.—Venezuela.
medica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 232.—Bras.
Medusula, *Muell. Arg. in Flora*, lix. (1876) 541, 545. —Bras.
megacoma, *Miq. Fl. Ind. Bat.* ii. 284.—Java.
megalocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 268.—Bras.
megalophylla, *Muell. Arg. l. c.* 224.—Bras.
megalosperma, Vahl, Eclog. Am. iii. 3 = Chione glabra.
megapontica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 361.—Bras.
melanotricha, *Muell. Arg. l. c.* 345.—Bras.
membranacea, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 189.—Am. austr.
membranifolia, *Bartl. ex DC. Prod.* iv. 522.—Ins. Philipp.
mesentericarpa, *Baker, in Journ. Linn. Soc.* xxi. (1885) 412.—Madag.
mesomorpha, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 308.—Bras.
mesotropa, *Muell. Arg. in Flora*, lix. (1876) 550, 554. —Bras.
mexicana, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 192.—Mexic.
micrantha, *H. B. & K. Nov. Gen. et Sp.* iii. 363. t. 284. —Am. austr.
microcarpa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 282. —Bras.
microcarpa, Wall. Cat. n. 8344 = tortilis.
microcephala, *Miq. Fl. Ind. Bat.* ii. 293.—Java?
microcephala, Miq. Stirp. Surin. Sel. 180 = Hoffmannseggiana.
microcephala, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 351.—Bras.
microdesma, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 36.—Costa Rica.
minutiflora, *Muell. Arg. in Flora*, lix. (1876) 548, 552.—Bras.
miradorensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 50.—Mexic.
mitis, Ruiz & Pav. Fl. Per. ii. 60 = Palicourea mitis.
mollis, *Poir. Encyc.* v. 701.—Mexic.
monocephala, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
montana, *Blume, Bijdr.* 961.—Java.
monticola, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 199.—Afr. trop.
monticola, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 315 = fulva.
Moonii, *Hook. f. Fl. Brit. Ind.* iii. 170.—Zeylan.
Morae, *Polak. in Linnaea*, xli. (1877) 570.—Costa Rica.
morindaeflora, *Wall. Cat.* n. 8438 A, B, C, partim.—Malaya.
mucronata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 211.—Afr. trop.
Muelleriana, *Wawra, Itin. Princ. Coburgi*, i. 124.—Bras.
multicolor, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 378. —Bras.
multiflora, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 109.—Afr. trop.
multiplex, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 319. —Bras.
myriantha, *Muell. Arg. in Flora*, lix. (1876) 549, 552. —Bras.
myrtifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = seq.
Myrtiphyllum, *Sw. Prod. Veg. Ind. Occ.* 44.—Jamaic.

PSYCHOTRIA:—
nematopoda, *F. Muell. Fragm.* vii. 48.—Austral.
nemorosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 109.—Bras.
nervosa, *Benth. in Hook. Journ. Bot.* iii. (1841) 229. —Guian.
nervosa, D. Don, Prod. Fl. Nep. 137 = denticulata?
nervosa, Sw. Prod. Veg. Ind. Occ. 43 = undata.
nesophila, *F. Muell. Fragm.* ii. 135.—Austral.
neurothrix, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 241. —Bras.
neurotricha, *DC. Prod.* iv. 505.—Porto Rico.
neurotricha, Schlecht. in Linnaea, xxviii. (1856) 520 = neurothrix.
nicaraguensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 34.—Nicaragua.
nicobarica, *Kurz, in Journ. Bot.* xiii. (1875) 328.—Ins. Nicobar.
nicotianaefolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 229.—Mexic.
nigropunctata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 207. —Afr. trop.
nitida, *Willd. Sp. Pl.* i. 963.—Guian.
nitidella, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 250.—Bras.
nitidula, *Cham. & Schlecht. in Linnaea*, iv. (1829) 25. —Bras.
nivosa, Linden, ex Berkeley, in Journ. Hort. Soc. N. S. i. (1866) 156 = Rudgea nivosa.
nonatelioides, Cham. & Schlecht. in Linnaea, iv. (1829) 26 = flexuosa.
noxia, *A. St. Hil. Pl. Rem. Brés.* 234. t. 21. f. A.—Bras.
nubica, *Delile, Cent. Pl. Méroé*, 66.—Nubia.
nuda, Wawra, Itin. Princ. Coburgi, i. 128 = Gardneriana.
nudiflora, *Wight & Arn. Prod.* 434.—Ind. or.
nutans, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Ocotea guianensis.
nutans, *Sw. Prod. Veg. Ind. Occ.* 43.—Ind. occ.
obconica, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
obesa, Wall. Cat. n. 8328 = Gaertnera obesa.
obfuscata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 377. —Bras.
obliquinervia, *Muell. Arg. in Flora*, lix. (1876) 549, 552.—Bras.
oblongifolia, E. Mey. in Drège, Zwei Pfl. Docum. 214 = Kraussia floribunda.
obovalis, *A. Rich. Fl. Cub. Fanerog.* ii. 27.—Cuba.
obovata, *Hemsl. Biol. Centr. Am. Bot.* ii. 50.—Mexic.
obovata, Ruiz & Pav. Fl. Per. ii. 58 = Palicourea obovata.
obovata, Wall. Cat. n. 8380 A = polycarpa, sarmentosa.
obscura, *Benth. in Hook. Niger Fl.* 419.—Afr. trop.
obscura, *Zoll. & Mor. Syst. Verz. Zoll.* 65.—Java.
obscurata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 253.—Bras.
obtegens, *Muell. Arg. l. c.* 277.—Bras.
obtusifolia, Lam. Illustr. ii. 243; Poir. Encyc. v. 697. —Madag.
obtusifolia, E. Mey. in Drège, Zwei Pfl. Docum. 214 (= *Canthium obovatum*).—Afr. austr.
obtusifolia, Sieber, ex DC. Prod. iv. 524 = Gaertnera edentata.
obvallata, Schum. & Thonn. Beskr. Guin. Pl. 111 = Geophila obvallata.
octocuspis, *Muell. Arg. in Flora*, lix. (1876) 548, 551. —Bras.
odorata, Blume, ex Miq. Fl. Ind. Bat. ii. 296 = malayana.
odorata, *Wright, ex Griseb. Cat. Pl. Cub.* 138.—Cuba.
officinalis, *Raeusch. Nom.* ed. III. 55.—Guian.
oligantha, *DC. Prod.* iv. 514.—Ins. S. Doming.
oligantha, *Miq. Fl. Ind. Bat.* ii. 288.—Java.
oligotricha, *DC. Prod.* iv. 514.—Am. austr.
opaca, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 263.—Bras.
ophioxyloides, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 168 = Chasalia curviflora.
orinocensis, *DC. Prod.* iv. 512.—Venezuela.
ornata, Spruce, ex Benth. & Hook. f. Gen. ii. 128 = Cephaëlis ornata.
ovoidea, *Wall. Cat.* n. 8383.—Malaya.
owariensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 204.—Afr. trop.

PSYCHOTRIA :—
oxyphylla, Wall. Cat. n. 8374 = Gaertnera Koenigii.
pachyneura, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 359. —Bras.
pacimonica, *Muell. Arg. l. c.* 337.—Bras.
padifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 189.—Mexic.
Palicourea, Sw. Fl. Ind. Occ. i. 433 = Palicourea guianensis.
palicoureoides, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 305.—Bras.
Palicurea, Willd. ex Steud. Nom. ed. II. ii. 410 = Palicourea petiolaris.
Palicurea, Willd. Sp. Pl. i. 971 = Palicourea guianensis.
palicureoides, Mart. in Flora, xxiv. (1841) II. Beibl. 64 = Rudgea palicoureoides.
pallens, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 108.—Bras.
pallida, Willd. ex Steud. Nom. ed. II. ii. 410 = Palicourea crocea.
paludosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 367.—Bras.
paniculata, *Willd. Sp. Pl.* i. 970.—Am. trop.
papantlensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 50.—Mexic.
paradoxa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 313. —Bras.
paraënsis, *Muell. Arg. l. c.* 245.—Bras.
parasitica, *Sw. Prod. Veg. Ind. Occ.* 44.—Ind. occ.
pardina, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 371.—Bras.
parviflora, *Span. in Linnaea*, xv. (1841) 319.—Ins. Timor.
parviflora, *Willd. Sp. Pl.* i. 962.—Guian.
parvifolia, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 35.—Nicaragua.
parvula, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Fiji.
patens, D. Dietr. Syn. Pl. i. 765 = Brownei, foetida.
patens, *Sw. Prod. Veg. Ind. Occ.* 45.—Ind. occ.
patentinervia, *Muell. Arg. in Flora*, lix. (1876) 543, 546.—Bras.
patentivenia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 208. —Ins. Celebes.
Patrisii, *DC. Prod.* iv. 510.—Guian.
patula, Willd. ex Roem. & Schult. Syst. v. 189 = alba.
pauciflora, *Bartl. ex DC. Prod.* iv. 522.—Ins. Philipp.
paucifolia, Willd. ex Spreng. Syst. i. 751 = longiflora.
pauridiantha, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 196. —Afr. trop.
Pavetta, Sw. Prod. Veg. Ind. Occ. 45 = Palicourea Pavetta.
Peckoltiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 366.—Bras.
pedunculata, Bartl. ex DC. Prod. iv. 507 = Haenkeana.
pedunculata, *Sw. Prod. Veg. Ind. Occ.* 44.—Jamaic.
pedunculosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = Palicourea pedunculosa.
pelagica, *Seem. Fl. Vit.* 136.—Ins. Fiji.
penangensis, *Hook. f. Fl. Brit. Ind.* iii. 175.—Ins. Penang.
pendula, *Hook. f. l. c.* 164.—Ins. Andaman.
pentaphthosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 284.—Bras.
pergamena, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 241.—Sumatra.
persimilis, *Muell. Arg. in Flora*, lix. (1876) 541, 545. —Bras.
Pervillei, *Baker, Fl. Maurit.* 155.—Ins. Seychelles.
petiolaris, Spreng. Syst. i. 745 = Palicourea petiolaris.
philippensis, *Cham. & Schlecht. in Linnaea*, iv. (1829) 21.—Ins. Philipp.
phyllocalymma, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 373.—Bras.
phyllocalymmoides, *Muell. Arg. l. c.* 374.—Bras.
Phytolacca, Spreng. ex DC. Prod. iv. 505 = laevis.
phytolaccea, *Poir. Encyc.* v. 704.—Am. austr.
Pickeringii, *A. Gray, in Proc. Am. Acad.* iv. (1860) 47.—Ins. Fiji.
picta, Wall. Cat. n. 8353 = calocarpa.
pilosa, *Ruiz & Pav. Fl. Per.* ii. 60. t. 208. f. a.—Peruv.
pilosiuscula, *Griseb. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
platycocca, *A. Gray, in Proc. Am. Acad.* iv. (1860) 47.—Ins. Fiji.

PSYCHOTRIA :—
platyneura, *Kurz, in Journ. Bot.* xiii. (1875) 327.—Ins. Andaman.
platyphylla, *DC. Prod.* iv. 517.—Porto Rico.
platypoda, *DC. l. c.* 510.—Guian.
platypodina, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 463.—Bras.
pleiocephala, *Muell. Arg. in Flora*, lix. (1876) 549, 552.—Bras.
plurivenia, *Thw. Enum. Pl. Zeyl.* 149.—Zeylan.
Poeppigiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 370.—Bras.
poliostemma, *Benth. Fl. Austral.* iii. 428.—Austral.
polyantha, *DC. Prod.* iv. 516.—Ins. S. Doming.
polycarpa, *Hook. f. Fl. Brit. Ind.* iii. 165.—Malaya.
polycephala, *Benth. in Hook. Journ. Bot.* iii. (1841) 231.—Guian.
polyneura, DC. Prod. iv. 520 = denticulata.
polyneura, *Kurz, in Journ. Bot.* xiii. (1875) 327.—Ins. Andaman.
polyodonta, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 286. —Bras.
polytricha, *Miq. Fl. Ind. Bat.* ii. 287.—Sumatra.
porophylla, Vell. Fl. Flum. 68; ii. t. 38 = Faramea porophylla.
portoriccensis, DC. Prod. iv. 515 = rufescens.
Pritchardii, *Seem. Fl. Vit.* 135.—Ins. Fiji.
procumbens, Vell. Fl. Flum. 64; ii. t. 23 = Coussarea procumbens.
psathuroides, *DC. Prod.* iv. 518.—Ins. Maurit.
Pseudo-pavetta, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 281.—Porto Rico.
psilogyne, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 279.—Bras.
puberula, *Wright, ex Griseb. Cat. Pl. Cub.* 135.—Cuba.
pubescens, Bartl. ex DC. Prod. iv. 513 = hebeclada.
pubescens, *Sw. Prod. Veg. Ind. Occ.* 44.—Am. trop.
pubigera, *Schlecht. in Linnaea*, xxviii. (1856) 514.—Bras.
pumila, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 207.—Afr. trop.
punctata, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 230.—Afr. trop.
punicea, Ruiz & Pav. Fl. Per. ii. 62 = Palicourea punicea.
purpurascens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 354.—Bras.
pyramidalis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 509.—Cuba.
pyrifolia, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 360.—Ins. J. Fernand.
quadrifolia, Rudge, Pl. Gui. 27. t. 42 = Palicourea quadrifolia.
quinquecuspis, *Muell. Arg. in Flora*, lix. (1876) 549, 552.—Bras.
quinqueradiata, *Polak. in Linnaea*, xlī. (1877) 570.—Costa Rica.
Rabeniana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 264. —Bras.
racemosa, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
racemosa, *Willd. Sp. Pl.* i. 966.—Bras.; Guian.
radians, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 242.—Bras.
recurva, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 206.—Afr. trop.
Reevesii, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 104 = elliptica.
refracta, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 262.—Bras.
Regina, *Muell. Arg. l. c.* 318.—Bras.
Regnellii, *Muell. Arg. in Flora*, lix. (1876) 548, 552.—Bras.
remota, *Benth. in Hook. Journ. Bot.* iii. (1841) 225.—Guian.
repanda, *Ruiz & Pav. Fl. Per.* ii. 61.—Peruv.
reptans, *Benth. in Hook. Niger Fl.* 418.—Afr. trop.
reticulata, *Ruiz & Pav. Fl. Per.* ii. 56. t. 212. f. b.—Peruv.
revoluta, *DC. Prod.* iv. 517.—Ins. S. Doming.
Rhinocerotis, *Reinw. ex Blume, Bijdr.* 961.—Java.
rhodocarpa, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 30.—Sumatra.
rhodoleuca, *Muell. Arg. in Flora*, lix. (1876) 541, 545. —Bras.
rhytidocarpa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 289.—Bras.

PSYCHOTRIA :—
Riedeliana, *Muell. Arg. in Flora*, lix. (1876) 548, 551.—Bras.
rigida, *Bredem. ex Roem. & Schult. Syst.* v. 192.—Bras.
robusta, *Blume, Bijdr.* 962.—Java.
rosea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 360.—Bras.
rostrata, Blume, Bijdr. 961 = Chasalia rostrata.
Roxburghii, *DC. Prod.* iv. 519.—Ins. Molucc.
rubra, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 336.—Bras.
rubra, Poir. Encyc. Suppl. iv. 597 = Antherura rubra.
rudgeoides, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 463. —Bras.
ruelliaefolia, *Muell. Arg. l. c.* 364.—Bras.
rufescens, *H. B. & K. Nov. Gen. et Sp.* iii. 364.—Am. trop.
rufescens, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 192 = micrantha.
rufipes, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 220.—Ins. Galap.
rufovaginata, *Griseb. Cat. Pl. Cub.* 136.—Cuba.
rugulosa, *H. B. & K. Nov. Gen. et Sp.* iii. 356.—Peruv.
rupestris, *Muell. Arg. in Flora*, lix. (1876) 542, 546.—Bras.
salicifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 51.—Mexic.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 356.—N. Granat.
salicifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = lamprophylla.
saligna, E. Mey. in Drège, Zwei Pfl. Docum. 214 = Kraussia lanceolata.
Salzmanniana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 362.—Bras.
sambucina, *Link, ex Roem. & Schult. Syst.* v. 188.—Bras.
santaremica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 328.—Bras.
sarmentosa, *Blume, Bijdr.* 964.—Ind. or.; Malaya.
scabriuscula, *Bartl. ex DC. Prod.* iv. 513.—Am. austr.
scandens, Hook. & Arn. Bot. Beech. Voy. 193 = serpens.
scandens, Moon, Cat. Pl. Ceyl. 14 = sarmentosa.
scandens, Reinw. ex Miq. Fl. Ind. Bat. ii. 287 = laxiflora.
Schlechtendaliana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 259.—Bras.
Schomburgkii, *Benth. in Hook. Journ. Bot.* iii. (1841) 231.—Guian.
Schottiana, *Muell. Arg. in Flora*, lix. (1876) 547, 551. —Bras.
Schuechiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 348.—Bras.
Schweinfurthii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 210.—Afr. trop.
sciadephora, *Hiern, l. c.* 202.—Afr. trop.
sclerophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 239.—Bras.
scrobiculata, Spreng. Syst. v. 572 (Index) = Coffea foveolata.
segregata, *Muell. Arg. in Flora*, lix. (1876) 549, 552.—Bras.
Sellowiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 239. —Bras.
semifissa, *Muell. Arg. l. c.* 340.—Bras.
serpens, *Linn. Mant.* ii. 204.—China.
sessiliflora, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 191.—Venezuela.
sessilifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. I. (1844) 228.—Mexic.
sessilis, Muell. Arg. in Mart. Fl. Bras. vi. v. 358 = axillaris.
sessilis, Vell. Fl. Flum. 65; ii. t. 26 = Rudgea sessilis.
setacea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 197.—Afr. trop.
setifera, *Benth. in Hook. Journ. Bot.* iii. (1841) 228.—Guian.
silhetensis, *Hook. f. Fl. Brit. Ind.* iii. 174.—Reg. Himal.
silvicola, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 262.—Bras.
Simira, Roem. & Schult. Syst. v. 187 = parviflora, *Willd.*

PSYCHOTRIA:—
simulans, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 363.—Bras.
singalensis, *Miq. Fl. Ind. Bat.* ii. 294.—Sumatra.
sonans, Muell. Arg. in Mart. Fl. Bras. vi. v. 231, sphalm. = rigida.
sordida, *Thw. Enum. Pl. Zeyl.* 149.—Zeylan.
sororia, *DC. Prod.* iv. 510.—Guian.
sororiella, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 375.—Bras.
soteropolitana, *Muell. Arg. l. c.* 260.—Bras.
Soyauxii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 213.—Afr. trop.
spathicalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 319.—Bras.
speciosa, Forst. f. Prod. 16 = Ixora vitiensis.
speciosa, Spreng. Syst. i. 746 = Palicourea speciosa.
sphaerocarpa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 161.—Reg. Himal.
sphaerocephala, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
spicata, *Benth. in Hook. Journ. Bot.* iii. (1841) 231.—Guian.
spicata, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
splendens, *Spreng. Syst.* i. 740.—Am. austr.
Sprucei, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 226.—Bras.
squarrosa, *Muell. Arg. l. c.* 231.—Bras.
stachyoides, *Benth. in Linnaea*, xxiii. (1850) 464.—Bras.
stellaris, *Muell. Arg. in Flora*, lix. (1876) 547, 551.—Bras.
stenocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 268.—Bras.
stenoclada, *Muell. Arg. l. c.* 254.—Bras.
stenophylla, *Hook. f. Fl. Brit. Ind.* iii. 161.—Zeylan.
stenophylla, Schott, in Spreng. Syst. iv. App. 404 = noxia.
stenophylla, Spreng. Syst. i. 744 = Palicourea angustifolia.
stictophylla, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 212.—Afr. trop.
stipulacea, Sw. ex DC. Prod. iv. 514 = nervosa.
stipulacea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 164.—Ins. Penang.
stipularis, Reinw. ex Miq. Fl. Ind. Bat. ii. 280 = robusta.
stipulata, *Hemsl. Biol. Centr. Am. Bot.* ii. 51.—Mexic.
stipulata, Vell. Fl. Flum. 66; ii. t. 31 = Ixora stipulata.
stipulosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 334.—Bras.
Storckii, *Seem. Fl. Vit.* 135.—Ins. Fiji.
strepens, DC. Prod. iv. 531, in syn. = Palicourea rigida.
strigosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 373.—Bras.
subaeneo-fusca, *Muell. Arg. l. c.* 464.—Bras.
subalata, *Wright, ex Griseb. Cat. Pl. Cub.* 137.—Cuba.
subcacuminalis, *Muell. Arg. l. c.* 277.—Bras.
subcrocea, *Muell. Arg. l. c.* 244.—Bras.
subcuspidata, *Muell. Arg. l. c.* 261.—Bras.
subfusca, *Muell. Arg. l. c.* 297.—Bras.
subherbacea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 208.—Afr. trop.
subintegra, *Hook. f. Fl. Brit. Ind.* iii. 162.—Ind. or.
subnuda, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 209.—Afr. trop.
subobliqua, *Hiern, l. c.* 206.—Afr. trop.
subobovata, *Miq. Fl. Ind. Bat.* ii. 287.—Java?
subpunctata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 201.—Afr. trop.
subremota, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 315.—Bras.
subrufa, Miq. Fl. Ind. Bat. ii. 292 = P. Rhinocerotis.
subscandens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 246.—Bras.
subsessilis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 32.—Costa Rica.
subspathacea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 266.—Bras.
subtomentosa, *Ruiz & Pav. Fl. Per.* ii. 61. t. 210. f. a.—Peruv.
subtriflora, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.

PSYCHOTRIA:—
subundulata, *Benth. in Hook. Journ. Bot.* iii. (1841) 227.—Guian.
succulenta, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 216.—Afr. trop.
suffruticosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 249.—Bras.
sulcata, *Wall. Cat.* n. 8370.—Burma.
sulphurea, Ruiz & Pav. Fl. Per. ii. 58 = Palicourea sulphurea.
sulphurea, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 218.—Afr. trop.
sulphurea, *Seem. Fl. Vit.* 134.—Ins. Fiji.
sumbavana, *Miq. Fl. Ind. Bat.* ii. 285.—Malaya.
suterella, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 380.—Bras.
sylvana, Pers. Syn. i. 207 = Geophila reniformis.
sylvatica, *Blume, Bijdr.* 963.—Java.
symplocifolia, *Kurz, For. Fl. Brit. Burma*, ii. 11.—Reg. Himal.; Burma.
syringaeflora, Schlecht. in Linnaea, xxviii. (1856) 516 = Ixora syringaeflora.
tabacifolia, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 236.—Bras.
tabacifolia, Wall. Cat. n. 8334 = montana.
tabernaefolia, Poir. Encyc. v. 704 = Palicourea tabernaefolia.
Tacpo, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 312.—Ins. Philipp.
tenella, *Muell. Arg. in Flora*, lix. (1876) 544, 546.—Bras.
tenerior, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 331.—Bras.
tenuifolia, *Sw. Prod. Veg. Ind. Occ.* 43.—Ins. S. Doming.
tenuinervis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 282.—Bras.
tenuiramea, Muell. Arg. in Flora, lix. (1876) 542 = Hoffmannseggiana.
tephrosantha, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Fiji.
terminalis, *Vell. Fl. Flum.* 65; ii. t. 24.—Bras.
ternifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 169.—Madag.
tetragona, *Seem. Fl. Vit.* 137.—Ins. Fiji.
tetrandra, Blume, Bijdr. 961 = Chasalia curviflora.
tetraphylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 234.—Bras.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iii. 173.—Reg. Himal.
Thwaitesii, *Hook. f. l. c.* 162.—Ind. or.
thyrsiflora, Ruiz & Pav. Fl. Per. ii. 57 = Palicourea thyrsiflora.
tinctoria, Ruiz & Pav. l. c. 62 = Palicourea tinctoria.
tinifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 190 = Palicourea tinifolia.
tomentosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 51.—Mexic.
tomentosa, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 370.—Bras.
tortilis, *Blume, Bijdr.* 958.—Malaya.
trachystyla, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 213.—Afr. trop.
triantha, *Muell. Arg. in Flora*, lix. (1876) 550, 553.—Bras.
tribracteata, *Wright, ex Griseb. Cat. Pl. Cub.* 137.—Cuba.
trichantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 169.—Madag.
trichocephala, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 32. t. 238.—Peruv.
trichoclada, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 368.—Bras.
tricholoba, *Muell. Arg. l. c.* 343.—Bras.
trichoneura, *Muell. Arg. l. c.* 367.—Bras.
trichophora, *Muell. Arg. in Flora*, lix. (1876) 541, 545.—Bras.
trichophoroides, *Muell. Arg. l. c.*—Bras.
trichostyla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 289.—Bras.
trichotoma, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 227.—Mexic.
trichotoma, *Tenore, in Parl. Giorn. Bot. Ital.* ii. (1847) vii–viii. 202.—Hab.?
trifida, *Ruiz & Pav. Fl. Per.* ii. 60. t. 209. f. b.—Peruv.
triflora, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 359 = Coprosma triflora.

PSYCHOTRIA:—
triflora, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 108 = Cremaspora africana.
trifolia, Sieber, ex DC. Prod. iv. 622 = Rauwolfia Lamarkii.
triphylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 233.—Bras.
trispicata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 509—Ind. occ.
truncata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 162.—Ind. or.
tubulosa, *A. Rich. Fl. Cub. Fanerog.* ii. 27.—Cuba.
tubulosa, Vell. Fl. Flum. 68; ii. t. 37 = Rudgea eriantha.
turbinata, *A. Gray, in Proc. Am. Acad.* iv. (1860) 45.—Ins. Fiji.
turbinella, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 374.—Bras.
tylophora, *Kurz, in Journ. Bot.* xiii. (1875) 328.—Ins. Nicobar.
uberabana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 371.—Bras.
uliginosa, *Sw. Prod. Veg. Ind. Occ.* 43.—Jamaic.
umbellata, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 109.—Afr. trop.
umbellata, Vell. Fl. Flum. 67; ii. t. 35 = Arrabideae.
umbraticola, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 214.—Afr. trop.
undata, *Jacq. Hort. Schoenb.* iii. 5. t. 260.—Ins. Baham.; Jamaic.
undulata, *Miq. Fl. Ind. Bat.* ii. 294.—Borneo.
undulata, Poir. Encyc. Suppl. iv. 591 = undata.
uniflora, Reinw. ex Miq. Fl. Ind. Bat. ii. 304 = Amaracarpus pubescens.
urophylla, *Schlecht. in Linnaea*, xxviii. (1856) 522.—Bras.
vaginalis, G. Don, Gen. Syst. iii. 596 = seq.
vaginans, DC. Prod. iv. 520 = Gaertnera Koenigii.
vaginans, Hook. f. Fl. Brit. Ind. iii. 176 = truncata?
Vauthieri, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 295.—Bras.
vellerea, *Muell. Arg. l. c.* 305.—Bras.
Vellosiana, Bentl. in Linnaea, xxiii. (1850) 464 = axillaris.
velutipes, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 356.—Bras.
venulosa, *Muell. Arg. l. c.* 294.—Bras.
verrucosa, *Muell. Arg. l. c.* 246.—Bras.
verticillata, *Muell. Arg. l. c.* 233.—Bras.
vestita, *Presl, Bot. Bemerk.* 77.—Bras.
viburnoides, *H. B. & K. Nov. Gen. et Sp.* iii. 361.—Venezuela.
viburnoides, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 189 = lucida.
villosa, *Ruiz & Pav. Fl. Per.* ii. 59. t. 207. f. a.—Peruv.
villosa, Vell. Fl. Flum. 67; ii. t. 33 = barbiflora.
viminea, Wall. Cat. n. 8354 = Gaertnera viminea.
violacea, Aubl. Pl. Gui. i. 145. t. 55 = Geophila violacea.
violacea, Willd. Sp. Pl. i. 966 = Palicourea violacea.
virens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 213.—Afr. trop.
virens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 33.—Bras.
virgata, *Ruiz & Pav. Fl. Per.* ii. 60. t. 209. f. a.—Peruv.
viridiflora, *Reinw. ex Blume, Bijdr.* 963.—Java.
viridiflora, Reinw. ex Kurz, For. Fl. Brit. Burm. ii. 12 = calocarpa.
viridiflora, Zoll. ex Miq. Fl. Ind. Bat. ii. 298 = sylvatica.
viridis, *Ruiz & Pav. Fl. Per.* ii. 61. t. 210. f. b.—Peruv.
viridissima, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 315 = montana.
viticina, Steud. Nom. ed. II. ii. 410 = connata.
vitiensis, Seem. in Bonplandia, ix. (1861) 257; x. (1862) 36 = calycosa.
vittoriensis, *Muell. Arg. in Flora*, lix. (1876) 548, 551.—Bras.
Vogeliana, *Benth. in Hook. Niger Fl.* 420.—Afr. trop.
volubilis, Roxb. ex Wight & Arn. Prod. 424 = Paederia foetida.
Wallichiana, Spreng. Syst. iv. Cur. Post. 79 = denticulata?
Wallisiana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 224.—Bras.

PSYCHOTRIA:—
Warmingii, *Muell. Arg. in Flora*, lix. (1876) 544, 546. —Bras.
Wawrana, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 263. —Bras.
Weddelliana, *Muell. Arg. l. c.* 234.—Bras.
Westii, *DC. Prod.* iv. 516.—Porto Rico.
Wightiana, *Hook. f. Fl. Brit. Ind.* iii. 167.—Ind. or.
Willdenowii, *DC. Prod.* iv. 505.—Am. austr.
Wrightii, *Baker, Fl. Maurit.* 156.—Ins. Seychelles.
xanthocephala, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 351.—Bras.
xantholoba, *Muell. Arg. in Flora*, lix. (1876) 542, 546.—Bras.
xanthophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 255.—Bras.
zambesiana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 203. —Afr. trop.
zanguebarica, *Hiern, l. c.* 214.—Afr. trop.

PSYCHOTRION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = praec.

PSYCHOTROPHUM, P. Br. Hist. Jamaic. 160. t. 13. f. 1, 2 (1756) = **Psychotria**, Linn. (Rubiac.).
asiaticum, Crantz, Inst. ii. 259 = Psychotria asiatica.
herbaceum, Crantz, l. c. = Geophila reniformis.

PSYCHRIDIUM, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 263 = **Astragalus**, Tourn. (Legumin.).
brachytropis, Stev. l. c. = A. brachytropis.
oroboides, Stev. l. c. = A. oroboides.

PSYCHROGETON, Boiss. Fl. Orient. iii. 156 (1875) = **Aster**, Tourn. (Compos.).
cabulicum, Boiss. l. c.—Afghan.

PSYCHROPHILA, Bercht. & Presl, Rostl. i. Ranunc. 80 (1823) = **Caltha**, Rupp. (Ranunc.).
andicola, C. Gay, Fl. Chil. i. 49 = C. andicola.
appendiculata, Bercht. & Presl, Rostl. i. Ranunc. 80 = C. appendiculata.
dioneaefolia, C. Gay, Fl. Chil. i. 51 = C. dioneaefolia.
sagittata, Bercht. & Presl, Rostl. i. Ranunc. 80 = C. sagittata.

PSYCOTHRIA, Linn. Pl. Afr. Rar. 9 (1759), sphalm. = **Psychotria**, Linn. (Rubiac.).

PSYCROPHILA, Rafin. Atl. Journ. 144 (1832) = **Caltha**, Rupp. (Ranunc.).
auriculata, Rafin. l. c. = C. sagittata.
sagittata, Rafin. l. c. = C. sagittata.

PSYDARANTA, Neck. Elem. iii. 145 (1790) = **Calathea**, G. F. W. Mey. (Scitam.).
comosa, Rafin. Fl. Tellur. iv. 53 = C. comosa.

PSYDARANTHA, Steud. Nom. ed. II. ii. 411 (1841) = praec.

PSYDAX, Steud. Nom. ed. II. ii. 411 (1841), sphalm. = seq.

PSYDRAX, Gaertn. Fruct. i. 125. t. 26 (1788) = **Plectronia**, Linn. (Rubiac.).
angustifolia, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 191.—Madag.
dicoccos, Gaertn. Fruct. i. 125. t. 26. f. 2 = Plect. didyma.
major, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 191.—Madag.
media, A. Rich. l. c.—Madag.
minor, Steud. Nom. ed. II. ii. 411, sphalm. = praec.

PSYLLIUM, Tourn. ex Juss. Gen. 90 (1789) = **Plantago**, Tourn.
aegyptiacum, Mirb. Hist. Pl. ed. II. xiv. [1800] 333 = Plant. squarrosa.
afrum, Mirb. l. c. 334 = Plant. Psyllium.
annuum, Mirb. l. c. 332, 333 = Plant. Psyllium.
annuum, Thuill. Fl. Par. ed. II. 81 = Plant. arenaria.
arborescens, Mirb. Hist. Pl. ed. II. xiv. [1800] 334 = Plant. arborescens.
arenarium, Mirb. l. c. 333 = Plant. arenaria.
Cynops, Mirb. l. c. 332, 334 = Plant. Cynops.

PSYLLIUM:—
erectum, Jaume St. Hil. Expos. i. II. 209 = Plantago Psyllium.
genevense, Mirb. Hist. Pl. ed. II. xiv. [1800] 332, 334 = Plant. Cynops.
indicum, Dum.-Cours. Bot. Cult. ed. II. ii. 492 = seq.?
indicum, Mirb. Hist. Pl. ed. II. xiv. [1800] 334 = Plant. pumila.
parviflorum, Mirb. l. c. = Plant. Psyllium.
pumilum, Mirb. l. c. = Plant. pumila.
ramosum, Gilib. Fl. Lituan. i. 17 = Plant. arenaria.
squarrosum, Dum.-Cours. Bot. Cult. ed. II. ii. 492 = Plant. squarrosa.
strictum, Mirb. Hist. Pl. ed. II. xiv. [1800] 333 = Plant. stricta.
suffruticosum, Jaume St. Hil. Expos. i. II. 209 = Plant. Cynops.

PSYLLOCARPUS, Mart. Nov. Gen. et Sp. i. 44. t. 28 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 146.
DIODOIS, Pohl, in Flora, viii. (1825) 183, nomen.
asparagoides, *Mart. Nov. Gen. et Sp.* i. 46.—Bras.
ericoides, *Mart. l. c.* 45. t. 28. f. 1.—Bras.
laricinus, Spreng. Syst. iv. Cur. Post. 39 = seq.
laricoides, *Mart. Nov. Gen. et Sp.* i. 45. t. 28. f. 2.—Bras.
thymbroides, *Mart. l. c.* 46.—Bras.

PSYLLOCARPUS, Pohl, ex DC. Prod. iv. 479 (1830) = **Declieuxia**, H. B. & K. (Rubiac.).
cordifolius, Pohl, l. c. 480 = D. cordigera.
divergentiflorus, Pohl, l. c. = D. divergentiflora.
foliosus, Pohl, l. c. = D. foliosa.
galioides, Pohl, l. c. 479 = D. galioides.
glaber, Pohl, l. c. 481 = D. chiococcoides.
imbricatus, Pohl, l. c. 480 = D. daphnoides.
prunosus, Pohl, l. c. 481 = D. glauca.
trichotomus, Pohl, l. c. = D. mucronulata.

PSYLLOPHORA, Ehrh. Beitr. iv. 146 (1789) = **Carex**, Dill. (Cyperac.).
capitata, Schur, Enum. Pl. Transs. 697 = C. capitata.
Davalliana, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 = C. Davalliana.
dioica, Schur, l. c. = C. dioica.
obtusata, Schur, l. c. = C. obtusata.
pauciflora, Schur, l. c. x. (1859) 201 = C. pauciflora.
pulicaris, Schur, l. c. iv. (1853) 78 = C. pulicaris.
pyrenaica, Schur, l. c. = C. pyrenaica.
Sieberiana, Opiz, Seznam, 80 = C. Davalliana.
vulgaris, Heuff. in Flora, xxvii. (1844) 528 = C. pulicaris.

PSYLLOTHAMNUS, Oliver, in Hook. Ic. Pl. t. 1499 (1885). *ILLECEBRACEAE*.
Beevori, *Oliver, l. c.*—Arab.

PSYLOXYLON, Thou. ex Gaudich. in Freyc. Voy. Bot. 30 (1826) nomen. Genus dubium *LYTHRARIEARUM*.

PSYTHIRHISMA, Herb. ex Lindl. Veg. Kingd. 161 (1847) = PSITHYRISMA, Herb. = **Symphyostemon**, Miers (Irid.).

PTACOSEIA, Ehrh. Beitr. iv. 147 (1789) = **Carex**, Dill. (Cyperac.).

PTAEROXYLON, Eckl. & Zeyh. Enum. 54 (1835). *SAPINDACEAE*, Benth. & Hook. f. i. 411.
utile, *Eckl. & Zeyh. l. c.*—Afr. austr.

PTARMICA, [Tourn.] Neck. Elem. i. 14 (1790) = **Achillea**, Linn. (Compos.).
abrotanoides, Vis. Fl. Dalm. ii. 81. t. 10 = A. abrotanoides.
acuminata, Ledeb. Fl. Ross. ii. 529 = A. Ptarmica.
ageratifolia, Nym. Consp. 364 = A. ageratifolia.
Aizoon, Griseb. Reise Rumel. ii. 304 = A. ageratifolia.
alpina, DC. Prod. vi. 22 = A. commutata, impatiens.
ambrosiaca, Boiss. & Heldr. Diagn. Ser. II. iii. 19 = A. ambrosiaca.
atrata, DC. Prod. vi. 20 = A. atrata.
Barrelieri, DC. l. c. 19 = A. Barrelierii.
biserrata, DC. l. c. 23 = A. biserrata.

PTARMICA:—
Bocconi, Steud. Nom. ed. II. ii. 411 = Achillea alpina.
borealis, DC. Prod. vi. 21 = A. sibirica.
cartilaginea, Ledeb. Fl. Ross. ii. 530 = A. Ptarmica.
cartilaginea, Miq. ex Heimerl, in Denkschr. Akad. Wien, xlviii. (1884) 188 = A. sibirica.
Clavennae, DC. Prod. vi. 21 = A. Clavennae.
Clusiana, Steud. Nom. ed. II. ii. 411 = A. atrata.
cristata, DC. Prod. vi. 22 = A. falcata?
Fraasii, Nym. Consp. 365 = A. Fraasii.
grandiflora, DC. Prod. vi. 23 = A. Ptarmica.
Haussknechtiana, Nym. Consp. 364 = A. Haussknechtii.
Herba-rota, DC. Prod. vi. 22 = A. Herba-rota.
hybrida, Nym. Consp. 365 = A. hybrida.
impatiens, DC. Prod. vi. 22 = A. impatiens.
impunctata, Nym. Consp. 365 = A. Kraettliana.
integrifolia, Gilib. Fl. Lituan. i. 216 = A. Ptarmica.
ircutiana, DC. Prod. vi. 22 = A. Ptarmica.
Jaborneggi, Halacsy, ex Heimerl, in Denkschr. Akad. Wien, xlviii. (1884) 155 = A. Jaborneggi.
Laggeri, Nym. Consp. 365 = A. Laggeri.
lenensis, Heimerl, in Denkschr. Akad. Wien, xlviii. (1884) 175 = A. Ptarmica.
Lereschii, Nym. Consp. 366 = A. obscura.
lingulata, DC. Prod. vi. 24 = A. lingulata.
macrophylla, DC. l. c. 21 = A. macrophylla.
mongolica, DC. l. c. 22 = A. sibirica.
Morisiana, Nym. Consp. 364 = A. Herba-rota.
moschata, DC. Prod. vi. 20 = A. moschata.
mucronulata, DC. l. c. = A. Barrelierii.
multifida, DC. Prod. vii. 295 = A. atrata.
multiflora, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 331 = A. alpina.
nana, DC. Prod. vi. 21 = A. nana.
oxyloba, DC. l. c. 20 = A. oxyloba.
pectinata, Schur, Enum. Pl. Transs. 326 = A. pectinata.
Pseudo-atrata, Schur, l. c. 327 = A. oxyloba.
pyrenaica, Nym. Consp. 364 = A. Ptarmica.
scardica, Griseb. Reise Rumel. ii. 304 = A. atrata.
Schouwii, Nym. Consp. 364 = A. Barrelierii.
serbica, Nym. l. c. = A. ageratifolia.
serrata, DC. Prod. vi. 22 = A. serrata.
sibirica, Ledeb. Fl. Ross. ii. 528 = A. sibirica.
speciosa, DC. Prod. vi. 23 = A. Ptarmica.
spinosa, Schur, Enum. Pl. Transs. 325 = A. impatiens.
tenuifolia, Schur, l. c. 327 = A. oxyloba.
Thomasiana, DC. Prod. vi. 21 = A. montana.
umbellata, DC. l. c. 22 = A. umbellata.
valesiaca, DC. l. c. 21 = A. valesiaca.
vulgaris, DC. l. c. 23 = A. Ptarmica.

PTELAEA, Moench, Meth. 55 (1794) = seq.

PTELEA, Linn. Syst. ed. I (1735). *RUTACEAE*, Benth. & Hook. f. i. 301.
BELLUCIA, Adans. Fam. ii. 344 (1763).
angustifolia, *Benth. Pl. Hartw.* 9.—Am. bor. occ.
aptera, *Parry, in Proc. Davenp. Acad.* iv. (1884) 39.—Calif.
arborea, Blanco, Fl. Filip. ed. I. 63 = Harpullia cupanioides.
Baldwinii, *Torr. & Gray, Fl. N. Am.* i. 215.—Am. bor.
Baldwinii, Young, Fl. Texas, 196 = angustifolia.
caroliniana, Hort. ex Steud. Nom. ed. II. ii. 412 = trifoliata.
microphylla, D. Dietr. Syn. Pl. i. 497 = trifoliata.
mollis, M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 406 = trifoliata.
monophylla, Lam. Illustr. i. 336 = trifoliata.
ovata, Lour. Fl. Cochinch. 82 = Ptelidium ovatum?
parvifolia, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* i. 170.—Mexic.
pentandra, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 83.—Mexic.
pentaphylla, Munchh. Hausv. v. 242 = trifoliata.
pinnata, Linn. f. Suppl. 126 = Zanthoxylum Blackburnia.
podocarpa, *DC. Prod.* ii. 83.—Mexic.
tomentosa, Rafin. Fl. Ludov. 108 = trifoliata.
trifoliata, Boland. Cat. 7 = angustifolia.
trifoliata, *Linn. Sp. Pl.* 118.—Am. bor.
trifoliolata, [Moç. & Sesse,] ex DC. Prod. ii. 83 = podocarpa.
viscosa, Linn. Sp. Pl. 118 = Dodonaea viscosa.
viticifolia, Salisb. Prod. 68 = trifoliata.

PTELEOCARPA, Oliver, in Trans. Linn. Soc. xxviii. (1873) 515. *OLACINEAE.*
longistyla, *Becc. Malesia*, i. (1877) 130.—Borneo.
malaccensis, *Oliver, in Trans. Linn. Soc.* xxviii. (1873) 515.—Malacca.

PTELIDIUM, Thou. Vég. Iles Afr. i. 11, 29 (1805); Gen. Nov. Madag. 24 (1806). *CELASTRINEAE*, Benth. & Hook. f. i. 363.
SERINGIA, Spreng. Anleit. ii. 694 (1818); Syst. i. 441 (1825).
ovatum, *Poir. Encyc. Suppl.* iv. 597.—Madag.

PTERACHNE, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 62, in nota (1842) = **Ascolepis**, Nees (Cyperac.).

PTERANDRA, A. Juss. in A. St. Hil. Fl. Bras. Mér. iii. 72. t. 179 A (1833). *MALPIGHIACEAE*, Benth. & Hook. f. i. 253.
latifolia, A. Juss. in Deless. Ic. Sel. iii. 19. t. 30 = Acmanthera latifolia.
psidiaefolia, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 73. t. 179 B = pyroidea.
pyroidea, *A. Juss. l. c.* 74.—Bras.

PTERANTHERA, Blume, Mus. Bot. Lugd. Bat. ii. 30. 1852 [1856] = **Vatica**, Linn. (Dipteroc.).
Mangachapoi, Blume, l. c. = Shorea Mangachapoi.
sinensis, Blume, l. c. = V. chinensis.

PTERANTHUS, Forsk. Fl. Aegypt. Arab. 36 (1775). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 18.
LOUICHEA, L'Hérit. Stirp. Nov. i. 135. t. 65 (1785).
echinatus, *Desf. Fl. Atlant.* i. 144.—Afr. bor.; Arab.
Forskåhlei, Mirb. in Hist. Nat. Pl. ed. II. x. 130 (*Forskaolii*) = echinatus.

PTERICHIS, Lindl. Gen. et Sp. Orch. 444 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 592.
ACRAEA, Lindl. in Benth. Pl. Hartw. 155 (1845).
Diuris, *Reichb. f. in Bonplandia*, ii. (1854) 10.—Venezuela.
galeata, *Lindl. Gen. et Sp. Orch.* 445.—Peruv.

PTERIDOPHYLLUM, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 719. t. 5. *PAPAVERACEAE*, Benth. & Hook. f. i. 54.
racemosum, *Sieb. & Zucc. l. c.* 720.—Japon.

PTERIDOPHYLLUM, Thw. in Hook. Kew Journ. vi. (1854) 65. t. 1 = **Filicium**, Thw. (Burserac.).
decipiens, Thw. l. c. 66 = F. decipiens.

PTERIGERON, A. Gray, Pl. Wright. i. 90, in nota (1852). *COMPOSITAE*, Benth. & Hook. f. ii. 292.
OLIGANTHEMUM, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 56.
STREPTOGLOSSA, Steetz, ex F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 491.
adscendens, *Benth. Fl. Austral.* iii. 533.—Austral.
decurrens, *Benth. l. c.* 531.—Austral.
dentatifolius, *F. Muell. Fragm.* ix. 119.—Austral.
filifolius, *Benth. Fl. Austral.* iii. 533.—Austral.
liatroides, *Benth. l. c.* 532.—Austral.
macrocephalus, *Benth. l. c.*—Austral.
microglossus, *Benth. l. c.*—Austral.
odorus, *Benth. l. c.*—Austral.

PTERIGIUM, Correa, in Ann. Mus. Par. viii. (1806) 397 = **Dipterocarpus**, Gaertn.
costatum, Correa, l. c. = D. alatus.
teres, Correa, l. c. x. (1807) 159 = Dryobalanops aromatica.

PTERIGOSTACHYUM, Nees, ex Steud. Nom. ed. II. ii. 412 (1841), nomen. *GRAMINEAE.*
Lehmanni, *Nees, ex Steud. l. c.*—Hab.?

PTERILEMA, Reinw. in Syll. Ratisb. ii. (1828) 13 = **Engelhardtia**, Leschen. (Jugland.).
aceriflorum, Reinw. l. c. = E. aceriflora.
acerifolia, Steud. Nom. ed. II. i. 412 = praec.

PTERIPHIS, Rafin. Fl. Tellur. iv. 99 (1836) = **Aristolochia**, Tourn.
tripteris, Rafin. l. c = A. tripteris.

PTERISANTHES, Blume, Bijdr. 192 (1825). *AMPELIDEAE*, Benth. & Hook. f. i. 387.
EMBAMMA, Griff. Notul. iv. 694 (1854).
cissioides, *Blume, Bijdr.* 193.—Malaya.
coriacea, Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 95 = polita.
heterantha, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 664.—Malacca.
involucrata, G. Don, Gen. Syst. i. 694, sphalm. = cissioides.
ovata, Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 95 = Vitis eriopoda.
pedata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 664.—Malacca.
polita, *M. Laws. l. c.* 663.—Burma; Malaya.

PTERIUM, Desv. in Journ. Bot. i. (1813) 75 = **Lamarkia**, Moench (Gramin.).
elegans, Desv. l. c. 76 = L. aurea.

PTERNANDRA, Jack, in Malay. Misc. ii. (1822) VII. 60. *MELASTOMACEAE*, Benth. & Hook. f. i. 771.
APTEUXIS, Griff. Notul. iv. 672 (1854).
EWYCKIA, Blume, in Flora, xiv. (1831) 524.
appendiculata, Walp. Rep. ii. 148, in syn., sphalm. = capitellata.
caerulescens, *Jack, in Malay. Misc.* ii. (1822) VII. 61.—Ind. or.
capitellata, *Jack, l. c.* p. iii.—Ind. or.
cordata, Baill. Hist. Pl. vii. 23 = Kibessia cordata.
cyanea, Triana, in Trans. Linn. Soc. xxviii. (1871) 153 = caerulescens.
echinata, Jack, in Malay. Misc. ii. (1822) VII. p. iii = Kibessia simplex.
echinata, Wall. Cat. n. 4078 A = Kibessia pubescens.
Korthalsiana, Triana, in Trans. Linn. Soc. xxviii. (1871) 153 = capitellata.
latifolia, Triana, l. c. = paniculata.
paniculata, *Benth. in Wall. Cat.* n. 4080; *Triana, l. c.*—Malaya.
parasitica, Finlays. & Wall. l. c. n. 4086 B = Pogonanthera reflexa.
simplex, Baill. Hist. Pl. vii. 24 = Kibessia pubescens?

PTERNIX, Hill, Veg. Syst. iv. 16 (1762) = **Carduus**, Tourn. (Compos.).
deflorata, Hill, Hort. Kew. 60 = C. defloratus.

PTEROCALYMMA, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 508 = **Lagerstroemia**, Linn. (Lythrar.).
calycina, Koehne, in Engl. Bot. Jahrb. iv. (1883) 25, sphalm. = seq.
paniculata, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 508 = L. calycina.

PTEROCALYMNA, Benth. & Hook. f. Gen. i. 783 (1867), err. typ. = praec.

PTEROCALYX, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 361 = **Alexandra**, Bunge (Chenop.).
strictissimus, Schrenk, l. c. = A. Lehmanni.

PTEROCARPOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = **Pterocarpus**, Linn. (Legumin.).
amphimenius, St. Lag. l. c. = Pteroc. Amphimenium.
ateleius, St. Lag. l. c. = Ateleia pterocarpa.
marsupius, St. Lag. l. c. = Pteroc. Marsupium.
peltarius, St. Lag. l. c. = Pteroc. Peltaria.

PTEROCARPUS, Berg. in Vet. Akad. Handl. Stockh. (1769) 116 = **Ecastaphyllum**, P. Br. (Legumin.).
Ecastaphyllum, Berg. l. c. = E. Brownei.

PTEROCARPUS, Linn. Fl. Zeyl. 196 (1747). *LEGUMINOSAE*, Benth. & Hook. f. i. 547.
AMPHYMENIUM, H. B. & K. Nov. Gen. et Sp. vi. 380 (1823).
ANCYLOCALYX, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 136. t. 2.
ECHINODISCUS, Benth. in Ann. Wien. Mus. ii. (1838) 94.
GRISELINIA, Scop. Introd. 302 (1777).
LINGOUM, Adans. Fam. ii. 319 (1763).
MOUTOUCHI, Aubl. Pl. Gui. ii. 748. t. 299 (1775).
NEPHRAEA, Noronha, ex Hassk. Cat. Hort. Bog. Alt. 283 (1844).

PTEROCARPUS:—
PHELLOCARPUS, Benth. in Ann. Wien. Mus. ii. (1838) 106.
VATAIREA, Aubl. Pl. Gui. ii. 755. t. 302 (1775).
WEINREICHIA, Reichb. Consp. 152 (1828).
abyssinicus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 233 = lucens.
Adansonii, *DC. Prod.* ii. 419.—Afr. trop.
advena, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 439.—Madag.
africanus, Hook. in W. Gray, Trav. West. Afr. 395. t. D = erinaceus.
Amerimnon, Poir. Encyc. v. 729 = Dalbergia Amerimnum.
Amphymenium, *DC. Prod.* ii. 418.—Mexic.
ancylocalyx, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 79.—Bras.
angolensis, *DC. Prod.* ii. 419.—Afr. trop.
anterus, J. F. Gmel. Syst. 1087 = Drepanocarpus lunatus.
Apalatoa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 111 (Apataloa) = Crudia Schreberi.
apterus, Gaertn. Fruct. ii. 351. t. 156 = Drepanocarpus lunatus.
Ateleia, DC. Prod. ii. 419 = Ateleia pterocarpa.
australis, *Endl. Prod. Fl. Norf.* 152.—Ins. Norfolk.
Berterii, Spreng. ex DC. Prod. ii. 421 = Ecastaphyllum Monetaria.
bilobus, Roxb. ex G. Don, Gen. Syst. ii. 376 = P. Marsupium.
buxifolius, Murr. Syst. ed. XIV. 642 = Brya Ebenus.
caucasicus, Jensch. ex Trautv. in Act. Hort. Petrop. viii. (1883) 192, sphalm. = Pterocarya caucasica.
corylifolia, Royle, Illustr. Bot. Himal. 461 (Index), sphalm. = Psoralea corylifolia.
crispatus, [Moç. & Sesse,] ex DC. Prod. ii. 418 = P. Draco.
cultratus, Vell. Fl. Flum. 301; vii. t. 95 = violaceus.
cyathiformis, [Moç. & Sesse,] ex DC. Prod. ii. 420 = Drepanocarpus cyathiformis.
dalbergioides, Roxb. Hort. Beng. 53; Fl. Ind. iii. 236 = indicus.
diadelphus, Blanco, Fl. Filip. ed. I. 563 = Derris thyrsiflora.
Draco, Lam. Tabl. Encyc. iii. 161 = indicus.
Draco, *Linn. Sp. Pl.* ed. II. 1662.—Am. austr.
dubius, Spreng. Syst. iv. Cur. Post. 268 = Ecastaphyllum dubium.
Ebenus, Pers. Syn. ii. 277 = Brya Ebenus.
echinatus, *Pers. l. c.*—Afr. trop.?
erinaceus, *Lam. Encyc.* v. 728.—Afr. trop.
erinaceus, F. Villar, in Blanco, Fl. Philip. ed. III. Nov. App. 68 = Vidalianus.
esculentus, *Schum. & Thonn. Beskr. Guin. Pl.* 330.—Afr. trop.
falcatus, Vell. Fl. Flum. 301; vii. t. 93 = Poecilanthe grandiflora.
flavus, Lour. Fl. Cochinch. 431 = indicus?
floribundus, *Wall. Cat.* n. 5846.—Ind. or.
frutescens, Blanco, Fl. Filip. ed. I. 562 = Derris uliginosa.
frutescens, Vell. Fl. Flum. 302; vii. t. 96 = Dalbergia variabilis.
glaber, Reich. ex DC. Prod. ii. 421 = Brya Ebenus.
gummifer, *Bert. ex DC. l. c.* 419.—Ins. S. Doming.
Hayesii, *Hemsl. Diagn. Pl. Nov.* 8.—Panama.
hemipterus, Gaertn. Fruct. ii. 351. t. 156. f. 2 = P. Draco.
hypostictus, *Miq. Fl. Ind. Bat. Suppl.* 296.—Sumatra.
indicus, Herb. Madr. ex Wall. Cat. n. 5915 = Derris oblonga.
indicus, *Willd. Sp. Pl.* iii. 904.—As. trop.
latifolius, Poir. Encyc. Suppl. iv. 610 = Lonchocarpus latifolius.
lucens, *Lepr. ex Guill. & Perr. Fl. Seneg. Tent.* i. 228.—Afr. trop.
lunatus, Linn. f. Suppl. 317 = Drepanocarpus lunatus.
lunulatus, Sieber, ex Steud. Nom. ed. II. ii. 412 = Machaerium robiniaefolium.
luteus, Poir. Encyc. Suppl. iv. 610 = indicus?
luteus, Vell. Fl. Flum. 301; vii. t. 92 = Platymiscium floribundum.
macrocarpus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 187.—Burma.
Marsupium, *Roxb. Pl. Corom.* ii. 9. t. 116.—Ind. or.
melliferus, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 239.—Afr. trop.
microcarpus, *Pers. Syn.* ii. 277.—Am. austr.
Moutouchi, Lam. Illustr. iii. 161 (*Montouchi*) = P. Draco.

PTEROCARPUS:—
niger, Vell. Fl. Flum. 300; vii. t. 91 = Dalbergia nigra.
obtusatus, *Miq. Fl. Ind. Bat.* i. 1. 136.—Ins. Sunda.
officinalis, Jacq. Select. Am. 283. t. 183. f. 92 = P. Draco.
orbiculatus, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 418.—Mexic.
ovalis, Linn. Amoen. Acad. viii. 260 = Ecastaphyllum Monetaria.
pallidus, Blanco, Fl. Filip. ed. I. 560 = indicus.
paniculatus, *Spreng. Syst.* iii. 192.—Bras.
Peltaria, DC. Prod. ii. 419 = Wiborgia fusca.
Plumieri, Poir. Encyc. Suppl. iv. 610 = Ecastaphyllum Monetaria.
polyspermus, Vell. Fl. Flum. 301; vii. t. 94 = Moldenhawera floribunda.
pubescens, Poir. Encyc. v. 730 = Lonchocarpus pubescens.
pubescens, *Spreng. Syst.* iv. *Cur. Post.* 268.—Mexic.
quercinus, Vell. Fl. Flum. 299; vii. t. 90 = Ecastaphyllum Monetaria.
Richardi, Poir. Encyc. Suppl. iv. 611 = Ecastaphyllum Monetaria.
Rohrii, *Vahl, Symb. Bot.* ii. 79.—Bras.
Roxburghii, *G. Don, Gen. Syst.* ii. 377.—Ind. or.
rufescens, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 79.—Peruv.
Sanguis-draconis, Crantz, Inst. ii. 77 = P. Draco.
santalinoides, L'Hérit. ex DC. Prod. ii. 419 = esculentus.
santalinus, Blanco, Fl. Filip. ed. I. 561 = indicus.
santalinus, Buch.-Ham. ex Wall. Cat. n. 5842 = P. Marsupium.
santalinus, *Linn. f. Suppl.* 318.—Ind. or.
sapindoides, DC. Prod. ii. 419 = Andira sapindoides.
scandens, *Poir. Encyc.* v. 730.—Venezuela.
senegalensis, Vahl, ex DC. Prod. ii. 419 = erinaceus.
sericeus, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 75.—Afr. austr.
simplicifolius, *Baker, in Oliver, Fl. Trop. Afr.* ii. 238.—Afr. trop.
Sissoo, Roxb. ex Wight & Arn. Prod. 264 = Dalbergia Sissoo.
suberosus, Pers. Syn. ii. 277 = P. Draco.
ternatus, Poir. Encyc. v. 727 = Ecastaphyllum Monetaria?
tinctorius, *Welw. Apont.* 584; *et ex Baker, in Oliver, Fl. Trop. Afr.* ii. 239.—Afr. trop.
uliginosus, Roxb. ex G. Don, Gen. Syst. ii. 377 = Derris uliginosa.
Vidalianus, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 309.—Ins. Philipp.
villosus, *Mart. ex Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 79.—Bras.
violaceus, *Vog. in Linnaea*, xi. (1837) 416.—Bras.
Wallichii, Wight & Arn. Prod. 267 = indicus.
Zollingeri, Miq. Fl. Ind. Bat. i. 1. 136 = indicus.

PTEROCARYA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 345. *JUGLANDEAE*, Benth. & Hook. f. iii. 399.
caucasica, *C. A. Mey. Verz. Pfl. Cauc.* 134.—Oriens.
chinensis, Hort. ex Lavall. Arbor. Segrez. 65 = stenoptera.
fraxinifolia, Spach, Hist. Vég. Phan. ii. 180 = caucasica.
laevigata, Hort. ex Lavall. Arbor. Segrez. 65 = stenoptera.
rhoifolia, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* i. 33.—Japon.
sorbifolia, *Sieb. & Zucc. l. c.*—Japon.
Spachiana, Lavall. Arbor. Segrez. 69 = caucasica.
stenoptera, *C. DC. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 34.—China.

PTEROCARYA, Nutt. ex Moq. in DC. Prod. xiii. 11. 106 (1849) = **Atriplex**, Tourn. (Chenop.).
spinosa, Nutt. ex Moq. l. c. 113 = A. canescens.

PTEROCAULON, Ell. Sketch, ii. 323 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 294.
CHLAENOBOLUS, Cass. in Dict. Sc. Nat. xlix. 337 (1827).
MONENTELES, Labill. Sert. Austro-Caled. 42. t. 43, 44 (1825).
SPHAERANTHOIDES, A. Cunn. ex DC. Prod. v. 456 (1836).

PTEROCAULON:—
alopecuroideum, DC. Prod. v. 454 = virgatum.
angustifolium, DC. l. c. = virgatum.
Billardieri, F. Muell. Papuan Pl. 43 = cylindrostachyum.
Bojeri, *Baker, Fl. Maurit.* 164.—Madag.
cylindrostachyum, *C. B. Clarke, Comp. Ind.* 98.—Burma; Malaya; Austral.
glandulosus, *F. Muell. Census*, 79.—Austral.
interruptum, DC. Prod. v. 454 = virgatum.
polystachyum, *DC. l. c.*—Bras.
pycnostachyum, *Ell. Sketch*, ii. 324.—Am. bor.
sphacelatum, *Benth. & Hook. f. Gen.* ii. 295.—Austral.
sphaeranthoides, *F. Muell. Census*, 79.—Austral.
spicatum, DC. Prod. v. 454 = virgatum.
verbascifolium, *F. Muell. Census*, 79.—Austral.
virgatum, *DC. Prod.* v. 454.—Am. trop.

PTEROCELASTRUS, Meissn. Gen. 68 (1837). *CELASTRINEAE*, Benth. & Hook. f. i. 365.
ASTEROCARPUS, Eckl. & Zeyh. Enum. 122 (1835), nomen prius.
arboreus, Walp. Rep. i. 535 = variabilis.
Burmanni, Walp. l. c. = tricuspidatus.
Dregeanus, *Sond. in Harv. & Sond. Fl. Cap.* i. 462.—Afr. austr.
litoralis, *Walp. Rep.* i. 535.—Afr. austr.
marginatus, *Baill. Adansonia*, xi. (1874) 266.—N. Caled.
nervosus, Walp. Rep. i. 535 = variabilis.
rostratus, *Walp. l. c.*—Afr. austr.
stenopterus, *Walp. l. c.*—Afr. austr.
tetrapterus, Walp. l. c. = variabilis.
tricuspidatus, *Walp. l. c.*—Afr. austr.
typicus, Walp. l. c. = tricuspidatus.
variabilis, *Sond. in Harv. & Sond. Fl. Cap.* i. 462.—Afr. austr.

PTEROCELTIS, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 292. *URTICACEAE*, Benth. & Hook. f. iii. 354.
Tatarinowii, *Maxim. l. c.* 293.—Mongolia.

PTEROCEPHALUS, Vaill. ex Adans. Fam. ii. 152 (1763) = **Scabiosa**, Tourn. (Dipsac.).
arabicus, Boiss. Diagn. Ser. I. x. 78.—Arab.
bellidifolius, Boiss. l. c. ii. 109 = S. pterocephala.
brachiatus, Spreng. Syst. i. 384 = S. brachiata.
brevis, Coult. Mém. Dipsac. 32. t. 1. f. 16 = S. papposa.
Broussonetii, Coult. ex DC. Prod. iv. 653 = S. Broussonetii.
canus, Coult. ex DC. l. c. = S. cana.
caramanicus, Boiss. & Heldr. Diagn. Ser. I. x. 77.—As. Min.
Coulteri, Boiss. Diagn. l. c. = S. papposa.
cyprius, Boiss. l. c. ii. 110 = *multiflorus*.
depressus, Coss. & Bal. in Bull. Soc. Bot. Fr. xx. (1873) 250.—Marocc.
diandrus, Lag. Gen. et Sp. Nov. 9 = S. papposa.
dumetorum, Coult. Mém. Dipsac. 32 = S. dumetorum.
frutescens, Hochst. in Flora, xxiv. (1841) 1. Intell. 27, nomen; et ex A. Rich. Tent. Fl. Abyss. i. 369 = S. frutescens.
incanus, Rafin. Fl. Tellur. iv. 94 (Quid?).—Hab.?
involucratus, Spreng. Syst. i. 384 = S. papposa.
kurdicus, Vatke, in Linnaea, xxxviii. (1874) 734.—Kurdistan.
lasiospermus, Link, in Buch, Beschr. Canar. Ins. 150 = S. fruticulosa.
lusitanicus, DC. Prod. iv. 653 = S. lusitanica.
marcescens, Moench, Meth. 491 = S. plumosa.
multiflorus, Poech, Enum. Pl. Ins. Cyp. 16.—Ins. Cyprus.
niveus, Coult. Mém. Dipsac. 32 = S. nivea.
palaestinus, Coult. l. c. 31. t. 1. f. 14 = S. brachiata.
papposus, Coult. l. c. 32. t. 1. f. 17 = S. papposa.
Parnassi, Spreng. Syst. i. 384 = S. pterocephala.
parviflorus, Fisch. & Mey. in Tchihat. Asie Min. Bot. ii. 223 = S. brachiata.
perennis, DC. Prod. iv. 653 = S. pterocephala.
persicus, Boiss. Diagn. Ser. I. vi. 72.—Persia.
Pinardi, Boiss. l. c. v. 88.—As. Min.
plumosus, Coult. Mém. Dipsac. 31. t. 1. f. 15 = S. plumosa.
plumosus, Heldr. ex Boiss. Diagn. Ser. I. x. 77 = *caramanicus*.

PTEROCEPHALUS:—
pulverulentus, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 152.—Syria.
Putkianus, Boiss. & Kotschy, ex Boiss. l. c. 150.—Persia.
pyrethrifolius, Boiss. & Hohen. Diagn. Ser. I. ii. 109.—Persia.
Quartinianus, A. Rich. Tent. Fl. Abyss. i. 369 = Scabiosa frutescens.
sanctus, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 266.—Arab.
Sibthorpianus, Steud. Nom. ed. II. ii. 413 = S. brachiata.
spathulatus, Coult. Mém. Dipsac. 32 = S. spathulata, *D. Dietr.*
strictus, Boiss. & Hohen. Diagn. Ser. I. ii. 110.—Persia.
Szovitsii, Boiss. Fl. Orient. iii. 150.—Persia.
tomentosus, Coult. ex DC. Prod. iv. 653 = S. sphaciotica.
Vaillanti, Lag. Gen. et Sp. Nov. 9 = S. papposa.
virens, Webb & Berth. Phyt. Canar. ii. 202. t. 81.—Ins. Canar.

PTEROCERAS, Hasselt, ex Hassk. in Flora, xxv. (1842) II. Beibl. 6 = **Sarcochilus**, R. Br. (Orchid.).
radicans, Hassk, l. c.; et in Hoev. & De Vriese, Tijdschr. ix. (1842) 142 = S. teres.

PTEROCHAETA, Steetz, in Lehm. Pl. Preiss. i. 455 (1845) = **Waitzia**, Wendl. (Compos.).
paniculata, Steetz, l. c. 456 = W. paniculata.

PTEROCHAETE, Arn. ex Boeck. in Linnaea, xxxvii. (1871–73) 600 = **Rynchospora**, Vahl (Cyperac.).
Drummondii, Arn. ex Boeck. l. c. = R. rariflora.

PTEROCHAETE, Boiss. Diagn. Ser. I. vi. 76 (1845) = **Pulicaria**, Gaertn. (Compos.).
Aucheri, Boiss. l. c. 77 = Pulic. Aucheri.
glaucescens, Boiss. l. c. 78 = Pulic. glaucescens
glutinosa, Boiss. l. c. 77 = Pulic. glutinosa.
mucronifolia, Boiss. l. c. 78.—Persia.

PTEROCHILUS, Hook. & Arn. Bot. Beech. Voy. t. 17 (1832) = **Microstylis**, Lindl. (Orchid.).
plantaginea, Hook. & Arn. l. c. = M. Rheedii.

PTEROCHITON, Torr. in Frém. Rep. 318 (1845) = **Atriplex**, Tourn. (Chenop.).
canescens, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 184 = A. canescens.
occidentale, Torr. in Frém. Rep. 318 = A. canescens.

PTEROCHLAMYS, Fisch. ex Endl. Gen. 294 (1837) = **Panderia**, Fisch. & Mey. (Chenop.).
pilosa, Fisch. & Mey. ex Ledeb. Fl. Ross. iii. 741 = Pand. pilosa.

PTEROCLADIS, Lamb. ex G. Don, in Sweet, Hort. Brit. ed. III. 349 (1839) = **Baccharis**, Linn. (Compos.).
genistelloides, Lamb. ex G. Don, l. c. = B. genistelloides.

PTEROCLADON, Hook. f. in Benth. & Hook. f. Gen. i. 763 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
Sprucei, *Hook. f. l. c.*—Peruv.

PTEROCLADUM, Triana, in Trans. Linn. Soc. xxviii. (1871) 97 = praec.

PTEROCOCCUS, Hassk. in Flora, xxv. (1842) II. Beibl. 41 = **Plukenetia**, Linn. (Euphorb.).
glaberrimus, Hassk. l. c. = Pluk. corniculata.

PTEROCOCCUS, Pall. Reise, App. 738. t. 5 (1776); Reise Russ. Reich. ii. App. 43 (1777) = **Calligonum**, Linn. (Polygon.).
aphyllus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 443 = C. leucocladum.
aphyllus, Pall. Reise, App. ii. 738 = C. Pallasia.
crispus, C. A. Mey. in Bull. Sc. Acad. Pétersb. viii. (1841) 340 = C. crispum.
leucocladus, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 211 = C. leucocladum.

PTEROCOCCUS:—
persicus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 191 = Calligonum persicum.
songaricus, C. A. Mey. in Bull. Sc. Acad. Pétersb. viii. (1841) 340 = C. songaricum.
tetrapterus, Meissn. in DC. Prod. xiv. 29 = C. tetrapterum.

PTEROCOELION, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 572. Gen. dub. *TILIACEARUM*. —Cf. Benth. & Hook. f. i. 985.
javanicum, Turcz. l. c.—Java.

PTEROCYCLUS, Klotzsch, in Bot. Ergeb. Waldem. Reise, 150. t. 47 (1862) = **Pleurospermum**, Hoffm. (Umbell.).
angelicoides, Klotzsch, ll. cc. = Pleurosp. angelicoides.

PTEROCYMBIUM, R. Br. in Benn. Pl. Jav. Rar. 219. t. 45 (1844) = **Sterculia**, Linn.
javanicum, R. Br. l. c. = S. campanulata.
nicobaricum, Didr. in Kjoeb. Vidensk. Meddel. (1854) 199.—Ins. Nicobar.

PTEROCYPERUS, Opiz, Seznam, 80 (1852) = **Cyperus** Mich.
esculentus, Opiz, l. c. = C. esculentus.

PTERODISCUS, Hook. Bot. Mag. t. 4117 (1844). *PEDALINEAE*, Benth. & Hook. f. ii. 1057.
aurantiacus, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 53.—Afr. trop.
Gayi, *Decne. in Ann. Sc. Nat.* Sér. V. iii. (1865) 336. —Afr. trop.
luridus, *Hook. f. Bot. Mag.* t. 5784.—Afr. austr.
speciosus, *Hook. l. c.* t. 4117.—Afr. austr.

PTERODON, Vog. in Linnaea, xi. (1837) 384. *LEGUMINOSAE*, Benth. & Hook. f. i. 551.
COMMILOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 110.
abruptus, *Benth. in Hook. Kew Journ.* ii. (1850) 236. —Bras.
emarginatus, *Vog. in Linnaea*, xi. (1837) 384.—Bras.
macrophyllus, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1103, nomen.—Guian.
polygalaeflorus, *Benth. in Hook. Kew Journ.* ii. (1850) 236.—Bras.
pubescens, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 127; *et in Mart. Fl. Bras.* xv. I. 306.—Bras.

PTEROGASTRA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 32. *MELASTOMACEAE*, Benth. & Hook. f. i. 742.
cupheoides, *Seem. Bot. Voy. Herald*, 122.—Am. centr.
divaricata, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 33.—Am. trop.
major, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 40. t. 2. f. 30 b.—Am. trop.
minor, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 33.—Venezuela.

PTEROGLOSSASPIS, Reichb. f. Otia Bot. Hamb. 67 (1878). *ORCHIDEAE*, Benth. & Hook. f. iii. 542.
eustachya, *Reichb. f. l. c.*—Abyss.

PTEROGLOSSIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 32 = **Reyesia**, Clos (Solan.).
laxa, Miers, l. c. 33 = R. chilensis.

PTEROGYNE, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 62, in nota (1842) = **Ascolepis**, Nees (Cyperac.).

PTEROGYNE, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 140. *LEGUMINOSAE*, Benth. & Hook. f. i. 586.
nitens, *Tul. l. c.*—Bras.

PTEROLEPIS, Endl. Gen. 1214 (1840) = **Osbeckia**, Linn. (Melastom.).

PTEROLEPIS, Miq. Comm. Phyt. 72 (1840). *MELASTOMACEAE*, Benth. & Hook. f. i. 742.
ARTHROSTEMMA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 355.
BRACHYANDRA, Naud l. c. ii. (1844) 143.

PTEROLEPIS:—
alpestris, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 39.—Bras.
Arstedii, Cogn. in Mart. Fl. Bras. xiv. III. 297, sphalm. = *Oerstedii*.
Balansaei, *Cogn. l. c.* 267.—Bras.
Buraeavi, *Cogn. l. c.* 268.—Bras.
capitata, Miq. Comm. Phyt. 79 = glomerata.
cataphracta, *Triana, in Trans. Linn. Soc.* xxviii (1871) 38.—Bras.
cerastifolia, Triana, l. c. 40 = Tibouchina cerastifolia.
clidemioides, *Triana, l. c.*—Bras.
exigua, *Triana, l. c.* 39.—Mexic.
filiformis, *Triana, l. c.* 39.—Bras.
glomerata, *Miq. Comm. Phyt.* 78.—Am. trop.; Ind. occ.
glomerata, Triana, in Trans. Linn. Soc. xxviii. (1871) 39 = Salzmannii, maritima.
Goudotiana, Triana, l. c. = pusilla.
herbacea, Triana, l. c. = Tibouchina herbacea.
Herincquiana, *Cogn. in Mart. Fl. Bras.* xiv. III. 268.—Paraguay.
heterostemon, Triana, in Trans. Linn. Soc. xxviii. (1871) 39. t. 2 = filiformis.
hirsutissima, Triana, l. c. = pauciflora.
Hostmanni, Steud. in Flora, xxvii. (1844) 720 = glomerata.
ladanoides, Triana, in Trans. Linn. Soc. xxviii. (1871) 39 = pumila.
lanceolata, *Cogn. in Mart. Fl. Bras.* xiv. III. 263.—Bras.
lasiophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 40 = Tibouchina lasiophylla.
longistyla, *Cogn. in Mart. Fl. Bras.* xiv. III. 283.—Bras.
macranthera, Triana, in Trans. Linn. Soc. xxviii. (1871) 40 = Tibouchina sebastianopolitana.
maritima, *Cogn. in Mart. Fl. Bras.* xiv. III. 273.—Bras.
Martiana, Triana, in Trans. Linn. Soc. xxviii (1871) 39 = Tibouchina sebastianopolitana.
nitida, Triana, l. c. = Tibouchina nitida.
Oerstedii, Triana, l. c. 40 = Tibouchina Oerstedii.
paludosa, *Cogn. in Mart. Fl. Bras.* xiv. III. 272.—Bras.
parnassiaefolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 38.—Bras.
pauciflora, *Triana, l. c.* 40.—Bras.
perpusilla, *Cogn. in Mart. Fl. Bras.* xiv. III. 280.—Bras.
Pohliana, *Cogn. l. c.* 279.—Bras.
polygonoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 39.—Bras.
pumila, Cogn. in Mart. Fl. Bras. xiv. III. 263 = pusilla.
pumila, *Crueg. in Linnaea*, xx. (1847) 102.—Am. trop.
pusilla, *Miq. l. c.* xviii. (1844) 619.—Am. trop.
repanda, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 39.—Bras.
Riedeliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 265.—Bras.
Salzmannii, *Cogn. l. c.* 271.—Bras.
saturejaeformis, *Cogn. l. c.* 277.—Bras.
simplicicaulis, Triana, in Trans. Linn. Soc. xxviii. (1871) 40 = Tibouchina simplicicaulis.
sipaneoides, *Cogn. in Mart. Fl. Bras.* xiv. III. 270.—Bras.
sipanioides, Triana, in Trans. Linn. Soc. xxviii. (1871) 39 = Trianaei.
striphnocalyx, *Cogn. in Mart. Fl. Bras.* xiv. III. 286.—Bras.
Trianaei, *Cogn. l. c.* 269.—Bras.
trichotoma, Cogn. l. c. 261 = pumila.
versicolor, Triana, in Trans. Linn. Soc. xxviii (1871) 40 = Tibouchina versicolor.
Weddelliana, *Triana, l. c.* 39.—Bolivia.

PTEROLEPIS, Schrad. in Goett. Gel. Anz. (1821) III. 2071 = **Scirpus**, Tourn. (Cyperac.).
scirpoides, Schrad. l. c. = S. littoralis.

PTEROLOBIUM, Andrz. ex DC. Syst. ii. 374 (1821) = **Thlaspi**, Tourn. (Crucif.).
Biebersteinii, Andrz. l. c. = T. latifolia.
macrophyllum, Rupr. Fl. Cauc. 125 = T. latifolia.

PTEROLOBIUM, R. Br. in Salt, Abyss. App. 64 (1814). *LEGUMINOSAE*, Benth. & Hook. f. i. 567.
CANTUFFA, J. F. Gmel. Syst. 677 (1791).

PTEROLOBIUM:—
KANTUFFA, Bruce, Trav. t. 14 (1790).
QUARTINIA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 259.
REICHARDIA, Roth, Nov. Pl. Sp. 210 (1821).
abyssinicum, A. Rich. Tent. Fl. Abyss. i. 246 = lacerans.
Brucei, Benth. & Hook. f. Gen. i. 588 = lacerans.
indicum, Hance, in Journ. Bot. xxi. (1883) 297 = subvestitum?
indicum, *A. Rich. Tent. Fl. Abyss.* i. 247, in obs.—Ind. or.
insigne, Blume, ex Miq. Fl. Ind. Bat. i. I. 106 = indicum.
Kantuffa, Wight & Arn. ex Steud. Nom. ed. II. ii. 413 = lacerans.
lacerans, *R. Br. in Salt, Abyss. App.* 64.—Abyssin.
lacerans, Wall. Cat. n. 5841 = indicum, Caesalpinia digyna.
macropterum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 71 = indicum.
microphyllum, Miq. Fl. Ind. Bat. i. I. 106 = indicum.
nitens, *F. Muell. ex Benth. Fl. Austral.* ii. 279.—Austral.
subvestitum, *Hance, in Journ. Bot.* xxii. (1884) 365.—China.

PTEROLOMA, DC. Prod. ii. 326, in textu (1825); Benth. in Miq. Pl. Jungh. 219 (1852) = **Desmodium**, Desv. (Legumin.).
auriculatum, DC. l. c. = D. triquetrum.
triquetrum, Benth. in Miq. Pl. Jungh. 217, 220 = D. triquetrum.

PTEROLOMA, Hochst. & Steud. in Flora, xxiv. (1841) I. Intell. 42; et ex Steud. Nom. ed. II. ii. 413 (1841) = **Dipterygium**, Decne. (Crucif.).
arabicum, Hochst. & Steud. ll. cc. = D. glaucum.

PTEROLOPHUS, Cass. in Dict. Sc. Nat. xliv. 34 (1826) = **Centaurea**, Linn. (Compos.).
lanceolatus, Cass. l. c. l. 249 = C. pterolopha.
pinnatifidus, Cass. l. c. 250 = C. alba.

PTEROMARATHRUM, Koch, ex DC. Prod. iv. 239 (1830) = **Prangos**, Lindl. (Umbell.).

PTERONEURON, Meissn. Gen. 10 (1837) = seq.

PTERONEURUM, DC. Syst. ii. 269 (1821) = **Cardamine**, Tourn. (Crucif.).
bipinnatum, Reichb. Fl. Germ. Excurs. 676 = C. thalictroides.
carnosum, DC. Syst. ii. 270 = C. carnosa.
corsicum, Jord. Diagn. i. 131 = C. graeca.
creticum, Jord. l. c. 133 = C. graeca.
Cupanii, Jord. l. c. = C. graeca.
dalmaticum, Vis. in Flora, xii. (1829) I. Erg. 17 = C. maritima.
decurrens, Blume, Bijdr. 51 = C. decurrens.
graecum, DC. Syst. ii. 270 = C. graeca.
javanicum, Blume, Bijdr. 51 = C. javanica.
maritimum, Reichb. Fl. Germ. Excurs. 676 = C. maritima.
microphyllum, J. & C. Presl, Delic. Prag. 237 = C. maritima.
Rochelianum, Reichb. ex Nym. Consp. 38 = C. graeca.
trichocarpum, Jord. Diagn. i. 132 = C. graeca.

PTERONIA, Linn. Sp. Pl. ed. II. 1176 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 259.
HENANTHUS, Less. Syn. Comp. 196 (1832).
PACHYDERIS, Cass. in Dict. Sc. Nat. lvi. 170. (1828).
PETRONIA, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 52.
PTEROPHORA, Neck. Elem. i. 78 (1790).
PTEROPHORUS, Vaill. ex Adans. Fam. ii. 118 (1763).
SCEPINIA, Neck. Elem. i. 78 (1790).
acerosa, DC. Prod. v. 357 = baccharoides.
acuminata, *DC. l. c.* 361.—Afr. austr.
adenocarpa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 109. —Afr. austr.
aromatica, Salisb. Prod. 188 = camphorata.
aspalatha, *DC. Prod.* v. 356.—Afr. austr.
aspera, Thunb. Prod. Pl. Cap. 143 = camphorata.
baccharoides, *Less. Syn. Comp.* 196.—Afr. austr.

PTERONIA:—
beckeoides, *DC. Prod.* v. 359.—Afr. austr.
callosa, *DC. l. c.* 362.—Afr. austr.
calvescens, Drège, ex Harv. & Sond. Fl. Cap. iii. 101 = gymnocline.
camphorata, *Linn. Sp. Pl.* ed. II. 1176.—Afr. austr.
Candollei, Harv. in Harv. & Sond. Fl. Cap. iii. 105 = glauca.
canescens, DC. Prod. v. 358 = cinerea.
caroliniana, Walt. Fl. Carol. 202 = Liatris squarrosa.
centauroides, *DC. Prod.* v. 362.—Afr. austr.
Cephalotes, Linn. f. Suppl. 358 = hirsuta.
Chamaepeuce, Spreng. Syst. iii. 440 = Cnicus Chamaepeuce.
Chamaepeuce, Tenore, Syll. Fl. Neap. 417 = Cnicus gnaphalodes.
chrysocomifolia, *Poir. Encyc.* v. 732.—Afr. austr.
ciliata, *Thunb. Prod. Pl. Cap.* 144; *DC. Prod.* v. 359.—Afr. austr.
cinerea, *Linn. f. Suppl.* 356.—Afr. austr.
connata, DC. Prod. v. 360 = flexicaulis.
crassifolia, *Poir. Encyc.* v. 731.—Afr. austr.
cylindracea, *DC. Prod.* v. 363.—Afr. austr.
dentata, Spreng. ex DC. l. c. 482 = Iphiona baccharidifolia.
divaricata, *Less. Syn. Comp.* 196.—Afr. austr.
echinata, Thunb. Prod. Pl. Cap. 143 = Felicia echinata.
elegans, *Sch. Bip. ex Walp. Rep.* ii. 970.—Afr. austr.
elongata, *Thunb. Fl. Cap. ed. Schult.* 631.—Afr. austr.
empetrifolia, *DC. Prod.* v. 363.—Afr. austr.
erythrochaeta, *DC. l. c.* 358.—Afr. austr.
fasciculata, *Linn. f. Suppl.* 357.—Afr. austr.
fasciculata, Willd. ex Less. Syn. Comp. 262 = Athanasia fasciculata.
fastigiata, Thunb. Prod. Pl. Cap. 143 = paniculata.
flexicaulis, DC. Prod. v. 360 = paniculata.
flexicaulis, *Linn. f. Suppl.* 355.—Afr. austr.
glabrata, DC. Prod. v. 362 = glauca.
glabrata, *Linn. f. Suppl.* 358.—Afr. austr.
glauca, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
glaucescens, *DC. Prod.* v. 357.—Afr. austr.
glomerata, *Linn. f. Suppl.* 356.—Afr. austr.
gymnocline, *DC. Prod.* v. 359.—Afr. austr.
heterocarpa, *DC. l. c.* 364.—Afr. austr.
hirsuta, *Linn. f. Suppl.* 356.—Afr. austr.
incana, *DC. Prod.* v. 358.—Afr. austr.
inflexa, *Linn. f. Suppl.* 356.—Afr. austr.
laricina, Houtt. ex DC. Prod. v. 364 = camphorata.
latisquama, DC. l. c. 363 = glauca.
leptolepis, *DC. l. c.*—Afr. austr.
leptospermoides, *DC. l. c.*—Afr. austr.
leucoclada, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 65.—Afr. austr.
leucoloma, *DC. Prod.* v. 362.—Afr. austr.
lucilioides, *DC. l. c.* 358.—Afr. austr.
lupulina, DC. l. c. 357 = inflexa.
marginata, Herb. Buitenz. ex C. B. Clarke, Comp. Ind. 124 = Inula Cappa.
marginata, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 52 = praec.?
membranacea, *Linn. f. Suppl.* 357.—Afr. austr.
microphylla, DC. Prod. v. 359 = glomerata.
minuta, Linn. f. Suppl. 357 = Nestlera humilis.
mucronata, *DC. Prod.* v. 362.—Afr. austr.
onobromoides, *DC. l. c.* 364.—Afr. austr.
oppositifolia, *Linn. Syst.* ed. XII. 538.—Afr. austr.
oppositifolia, E. Mey. ex DC. Prod. v. 363 = leptolepis.
ovalifolia, *DC. l. c.* 361.—Afr. austr.
pallens, *Linn. f. Suppl.* 357.—Afr. austr.
paniculata, *Thunb. Prod. Pl. Cap.* 143.—Afr. austr.
pauciflora, Sims, Bot. Mag. t. 1697 = Helipterum virgatum.
Porophyllum, Cav. Ic. iii. 13. t. 225 = Adenophyllum Porophyllum.
quinqueflora, *DC. Prod.* v. 363.—Afr. austr.
retorta, Linn. f. Suppl. 356 = hirsuta.
rigida, Berg. Desc. Pl. Cap. 231 = incana.
scabra, *Harv. in Harv. & Sond. Fl. Cap.* iii. 109.—Afr. austr.
scariosa, *Linn. f. Suppl.* 356.—Afr. austr.
sesuviifolia, DC. Prod. v. 360 = glabrata.
spinosa, Linn. f. Suppl. 357 = Hoplophyllum spinosum.
stoehelinoides, *DC. Prod.* v. 364.—Afr. austr.
stricta, Ait. Hort. Kew. ed. I. iii. 162 = camphorata.
succulenta, *Thunb. Prod. Pl. Cap.* 143.—Afr. austr.

PTERONIA:—
tenuifolia, *DC. Prod.* v. 363.—Afr. austr.
tomentosa, Lour. Fl. Cochinch. 489 (Quid?).—China.
tricephala, *DC. Prod.* v. 359.—Afr. austr.
turbinata, *DC. l. c.* 362.—Afr. austr.
uncinata, *DC. l. c.* 357.—Afr. austr.
undulata, *DC. l. c.* 361.—Afr. austr.
uniflora, Poir. Encyc. v. 733 = hirsuta?
verticillata, DC. Prod. v. 357 = uncinata?
villosa, *Linn. f. Suppl.* 356.—Afr. austr.
viscosa, DC. Prod. v. 364 = adenocarpa.
viscosa, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
xantholepis, DC. Prod. v. 358 = incana.

PTEROPAPPUS, Pritz. Ic. Ind. ii. 235 (1865), sphalm. = **Pterygopappus**, Hook. f. (Compos.).

PTEROPHORA, Harv. Gen. S. Afr. Pl. 223 (1838) = **Dregea**, E. Mey. (Asclep.).
Dregea, Harv. l. c. = D. floribunda.

PTEROPHORA, Neck. Elem. i. 78 (1790) = **Pteronia**, Linn. (Compos.).

PTEROPHORUS, Vaill. ex Adans. Fam. ii. 118 (1763) = **Pteronia**, Linn. (Compos.).
asper, Less. Syn. Comp. 195 = Pteronia camphorata.
camphoratus, Cass. in Dict. Sc. Nat. xxxvii. 474 = Pteronia camphorata.
strictus, Less. Syn. Comp. 195 = Pteronia camphorata.

PTEROPHYLLA, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 93 = **Weinmannia**, Linn. (Saxifr.).
fraxinea, D. Don, l. c. = W. fraxinea.
sessilis, Salm-Dyck, Hort. Dyck. 220, nomen (Quid?).—Hab.?

PTEROPHYLLUS, [Nelson] Senilis, Pinaceae, 163 (1866) = **Ginkgo**, Linn. (Conif.).
Gingko, C. Koch, Dendrol. ii. II. 107, sphalm. = seq.
salisburiensis, [Nelson] Senilis, Pinaceae, 163 = G. biloba.

PTEROPHYTON, Cass. in Bull. Soc. Philom. (1818) 76 = **Actinomeris**, Nutt. (Compos.).
alatum, Cass. in Dict. Sc. Nat. xliv. 49 = A. tetraptera.
alternifolium, Cass. l. c. = A. squarrosa.
ovatum, Cass. l. c. = A. ovata.
procerum, Cass. l. c. 50 = A. squarrosa.

PTEROPODIUM, DC. in Meissn. Gen. 300; Comm. 209 (1840) = **Jacaranda**, Juss. (Bignon.).
glabrum, DC. Prod. ix. 239.—Bolivia.
hirsutum, DC. l. c.—Bolivia.

PTEROPODIUM, Willd. ex Steud. Nom. ed. II. ii. 414 (1841) = **Calamagrostis**, Adans., etc. (Gramin.).
arundinaceum, Willd. ex Steud. l. c. = Deyeuxia triflora.
effusum, Willd. ex Steud. l. c. = Deyeuxia effusa.
heterophyllum, Willd. ex Steud. l. c. = Deyeuxia erianthra.
junceum, Willd. ex Steud. l. c. = Deyeuxia effusa.
planifolium, Willd. ex Steud. l. c. = Deyeuxia planifolia.
rigidum, Willd. ex Steud. l. c. = Deyeuxia rigida.
strictum, Willd. ex Steud. l. c. = C. Humboldtiana.

PTEROPOGON, DC. Prod. vi. 245 (1837) = **Helipterum**, DC. (Compos.).
australis, Nees, in Linnaea, xvi. (1842) 223 = H. pygmaeum.
Cassinianus, F. Muell. in Linnaea, xxv. (1852) 415 = Schoenia Cassiniana.
corymbosus, A. Gray, in Hook. Kew Journ. iv. (1852) 268 = H. corymbosum.
demissus, A. Gray, l. c. 269 = H. exiguum.
Drummondii, A. Gray, l. c. 267 = H. pygmaeum.
gracilis, A. Gray, l. c. 269 = H. gracile.
Humboldtianus, F. Muell. in Linnaea, xxv. (1852) 415 = H. Humboldtianum.
intermedius, F. Muell. l. c. 411.—Austral.
laevis, A. Gray, in Hook. Kew Journ. iv. (1852) 269 = H. laeve.

PTEROPOGON:—
oppositifolius, F. Muell. in Linnaea, xxv. (1852) 415 = Schoenia Cassiniana.
platyphyllus, F. Muell. l. c. 413 = Helipterum strictum.
polycephalus, A. Gray, in Hook. Kew Journ. iv. (1852) 268 = H. polycephalum.
pygmaeus, DC. Prod. vi. 245 = H. pygmaeum.
ramosissimus, F. Muell. in Linnaea, xxv. (1852) 412 = Helichrysum semifertile.
ramosus, A. Gray, in Hook. Kew Journ. iv. (1852) 270 = H. tenellum.
ruthenicus, C. A. Mey. in Goebel, Reise, ii. 284 = Tragopogon floccosus.
spicatus, Steetz, in Lehm. Pl. Preiss. i. 479 = H spicatum.

PTEROPOGON, Fenzl, in Flora, xxii. (1839) II. 727 = **Facelis**, Cass. (Compos.).
andicola, Nees, in Ind. Sem. Hort. Vratisl. (1841); ex eod. in Linnaea, xvi. (1842) 223 = seq.
chilensis, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 64 = F. apiculata.

PTEROPOGON, Neck. Elem. i. 109 (1840) = **Scabiosa**, Tourn. (Dipsac.).

PTEROPYRUM, Jaub. & Spach, Illustr. ii. 7. tt. 107–109 (1846). *POLYGONACEAE*, Benth. & Hook. f. iii. 85.
Aucherii, *Jaub. & Spach, l. c.* ii. 8. t. 107.—Persia.
ericoides, *Boiss. Fl. Orient.* iv. 1002.—Beluch.; Persia.
gracile, Boiss. Diagn. Ser. I. xii. 102 = Olivierii.
Griffithii, Meissn. in DC. Prod. xiv. 31 = Olivierii.
Noëanum, *Boiss. in Bot. Zeit.* xi. (1853) 734, nomen; *et ex Meissn. in DC. Prod.* xiv. 31.—Assyria.
Olivierii, *Jaub. & Spach, Illustr.* ii. 9. t. 108, (Oliverii).—Persia; Afghan.
scoparium, *Jaub. & Spach, l. c.* 10. t. 109.—Arabia.

PTEROSCLERIA, Nees, in Mart. Fl. Bras. ii. I. 196 (1842). *CYPERACEAE*, Benth. & Hook. f. iii. 1068.
guianensis, *Nees, l. c.*—Guian.
longifolia, *Griseb. Fl. Brit. W. Ind.* 579.—Ins. Trinit.; Guian.

PTEROSELINUM, Reichb. Fl. Germ. Excurs. 453 (1832) = **Peucedanum**, Tourn. (Umbell.).
alsaticum, Reichb. l. c. 454 = Peuced. alsaticum.
austriacum, Reichb. l. c. = Peuced. austriacum.
Chabraei, Reichb. l. c. 453 = Peuced. carvifolium.
glaucum, Reichb. l. c. = Peuced. Schottii.
intermedium, Schur, Enum. Pl. Transs. 265 = Peuced. austriacum.
montanum, Reichb. Fl. Germ. Excurs. 454 = Peuced. austriacum.
peucedanoides, Reichb. l. c. = Peuced. involucratum.
rablense, Reichb. l. c. = Peuced. rablense.
sibiricum, Reichb. l. c. = Peuced. baicalense.
transsilvanicum, Schur, Enum. Pl. Transs. 265 = Peuced. transsilvanicum.

PTEROSENECIO, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 300 (1884) = **Senecio**, Tourn. (Compos.).
grandis, Sch. Bip. ex Baker, l. c. 304 = S. grandis.

PTEROSIPHON, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 589 = **Cedrela**, Linn. (Meliac.).
multivenius, Turcz. l. c. 590 = C. angustifolia?

PTEROSPARTUM, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 440 (1880) = **Genista**, Linn. (Legumin.).
cantabricum, Willk. l. c. 441 = G. tridentata.
lasianthum, Willk. l. c. 440 = G. tridentata.
sagittale, Willk. l. c. = G. sagittalis.
Scolopendrinum, Willk. l. c. 469 = G. tridentata.
stenopterum, Willk. l. c. 441 = G. tridentata.
tridentatum, Willk. l. c. = G. tridentata.

PTEROSPERMUM, Schreb. Gen. 461 (1791). *STERCULIACEAE*, Benth. & Hook. f. i. 220.
SCZEGLEEWIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 233.
VELAGA, Adans. Fam. ii. 398 (1763); Gaertn. Fruct. ii. 245. t. 133 (1791).

PTEROSPERMUM:—
acerifolium, Benth. Fl. Hongk. 39 = heterophyllum.
acerifolium, *Willd. Sp. Pl.* iii. 729.—Ind. or.; Java.
acerifolium, Zoll. & Mor. Syst. Verz. Zoll. 27 = diversifolium.
aceroides, Wall. Cat. n. 1171 = acerifolium.
Blumeanum, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 311.—Sumatra; Java.
canescens, Roxb. Hort. Beng. 50; Fl. Ind. iii. 162 = suberifolium.
celebicum, *Miq. Illustr. Fl. Arch. Ind.* 87.—Ins. Celebes.
cinnamomeum, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 120, nomen; *et For. Fl. Brit. Burma*, i. 147.—Burma.
diversifolium, *Blume, Bijdr.* 88.—Ind. or.; Malaya.
Donianum, *Walp. Rep.* i. 350.—Ind. or.
elongatum, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 312.—Borneo.
fuscum, *Korth. l. c.*—Borneo.
glabrescens, *Wight & Arn. Prod.* 69.—Ind. or.
hastatum, Blanco, Fl. Filip. ed. I. 528 = diversifolium.
heterophyllum, *Hance, in Journ. Bot.* vi. (1868) 112.—China.
Heyneanum, G. Don, Gen. Syst. i. 538 = Donianum.
Heyneanum, *Wall. Cat.* n. 1169.—Ind. or.
Jackianum, *Wall. l. c.* n. 1164.—Ins. Penang.
javanicum, *Jungh. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 306.—Java.
lanceaefolium, Blume, Bijdr. 87 = Blumeanum.
lanceaefolium, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 163.—Ind. or.
Lawianum, Nimmo, in J. Grah. Cat. Pl. Bomb. 246 = Heyneanum.
obliquum, *Blanco, Fl. Filip.* ed. I. 529.—Ins. Philipp.
oblongum, Wall. Cat. n. 1165 = suberifolium.
obtusifolium, *Wight, ex Mast. in Hook. f. Fl. Brit. Ind.* i. 369.—Ind. or.
paniculatum, Roxb. Fl. Ind. ii. 642 = Brownlowia elata.
parvifolium, *Miq. Fl. Ind. Bat. Suppl.* 403.—Sumatra.
Pierrei, *Hance, in Journ. Bot.* xv. (1877) 329.—Cambodia.
plantanifolium, *Lodd. ex G. Don, in Loud. Hort. Brit.* 285, nomen.—Ind. or.
reticulatum, *Wight & Arn. Prod.* 69.—Ind. or.
rubiginosum, *Heyne, in Wall. Cat.* n. 1168.—Ind. or.
rubiginosum, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 28 = obliquum.
Sczegleewia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 573 = obliquum.
semisagittatum, Blanco, Fl. Filip. ed. II. 368 = obliquum.
semisagittatum, *Buch.-Ham. ex Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 160.—Ind. or.
suberifolium, *Lam. Illustr.* iii. 136.—Ind. or.
suberifolium, Willd. Sp. Pl. iii. 728 = Heyneanum.
subinaequale, *Miq. Fl. Ind. Bat. Suppl.* 404.—Sumatra.
subsessile, *Miq. l. c.* 403.—Sumatra.
sumatranum, *Miq. Illustr. Fl. Arch. Ind.* 87.—Sumatra.

PTEROSPORA, Nutt. Gen. Am. i. 269 (1818). *MONOTROPEAE*, Benth. & Hook. f. ii. 605.
andromedea, *Nutt. l. c.*—Am. bor.

PTEROSPOROPSIS, Kellogg, fide A. Gray, ex Benth. & Hook. f. Gen. ii. 606 (1876) = **Sarcodes**, Torr. (Monotr.).
sonoraënsis, Kellogg, ex Curran, in Bull. Calif. Acad. i. n. III. (1885) 141 = S. sanguinea.

PTEROSTEGIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 48 (1835). *POLYGONACEAE*, Benth. & Hook. f. iii. 94.
diphylla, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 168.—Calif.
drymarioides, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 48.—Calif.
fruticosa, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 212.—Calif.
galioides, *Greene, l. c.* 213.—Calif.
macroptera, *Benth. Bot. Voy. Sulph.* 44.—Calif.
microphylla, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 168 = drymarioides.

PTEROSTELMA, Wight, Contrib. 39 (1834) = **Hoya**, R. Br. (Asclep.).
acuminatum, Wight, l. c. = H. acuminata.
albiflora, Blume, Rumphia, iv. 33 = H. albiflora.

PTEROSTEMON, Schau. in Linnaea, xx. (1847) 736. *ROSACEAE*, Benth. & Hook. f. i. 615.
mexicanus, *Schau. l. c.*—Mexic.

PTEROSTEPHANUS, Kellogg, in Proc. Calif. Acad. iii. (1863) 20. f. 4 = **Anisocoma**, Torr. (Compos.).
runcinatus, Kellogg, l. c. 21 = A. acaulis.

PTEROSTIGMA, Benth. Scroph. Ind. 20 (1835) = **Adenosma**, R. Br. (Scrophul.).
capitatum, Benth. l. c. 21 = A. capitatum.
grandiflorum, Benth. l. c. = A. grandiflorum.
hirsutum, Miq. Fl. Ind. Bat. Suppl. 562.—Sumatra.
macrophyllum, Benth. Scroph. Ind. 21 = A. macrophyllum.
ovatum, Benth. in DC. Prod. x. 380 = A. ovatum.
rubiginosum, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 393 = A. grandiflorum.
spicatum, Benth. Scroph. Ind. 21 = A. capitatum.
stricta, Griff. Notul. iv. 96 = A. caeruleum.
subrepens, Thw. Enum. Pl. Zeyl. 426 = A. subrepens.
villosum, Benth. Scroph. Ind. 21 = A. caeruleum.
villosum, Thw. Enum. Pl. Zeyl. 218 = A. camphoratum.

PTEROSTYLIS, R. Br. Prod. 326 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 610.
DIPLODIUM, Sw. in Ges. Naturf. Fr. Berl. Mag. iv. (1810) 84.
acuminata, *R. Br. Prod.* 326.—Austral.
acuminata, Sieber, ex Lindl. Gen. et Sp. Orch. 391 = concinna.
alata, Reichb. f. Beitr. Syst. Pfl. 70 = praecox.
aphylla, *Lindl. Gen. et Sp. Orch.* 392.—Austral.
australis, *Hook. f. Fl. N. Zel.* i. 248.—N. Zel.
Banksii, *R. Br. ex A. Cunn. in Bot. Mag.* t. 3172.—N. Zel.
Baptistii, *Fitzg. Austral. Orch.* i. t. 2.—Austral.
barbata, Endl. in Lehm. Pl. Preiss. ii. 5 = pyramidalis.
barbata, *Lindl. Swan River App.* 53.—Austral.
clavigera, *Fitzg. in Journ. Bot.* xxiii. (1885) 138.—Austral.
coccina, *Fitzg. Austral. Orch.* i. t. 3.—Austral.
concinna, *R. Br. Prod.* 326.—Austral.
cucullata, *R. Br. l. c.* 327.—Austral.
curta, *R. Br. l. c.* 326.—Austral.
cycnocephala, *Fitzg. Austral. Orch.* i. t. 7.—Austral.
Daintreana, *F. Muell. ex Benth. Fl. Austral.* vi. 360.—Austral.
dubia, R. Br. Prod. 328 = cucullata.
dubia, Hook. f. Fl. Tasman. ii. 20. t. 115 = furcata.
emarginata, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 328.—N. Zel.
foliata, *Hook. f. Fl. N. Zel.* i. 249.—N. Zel.
furcata, *Lindl. Gen. et Sp. Orch.* 390.—Austral.
gibbosa, R. Br. Prod. 328 = rufa.
graminea, *Hook. f. Fl. N. Zel.* i. 248.—N. Zel.
grandiflora, *R. Br. Prod.* 327.—Austral.
hirsuta, Ait. ex Steud. Nom. ed. I. 667, nomen = obtusa?
hispidula, *Fitzg. Austral. Orch.* i. t. 5.—Austral.
longifolia, *R. Br. Prod.* 327.—Austral.
micromega, *Hook. f. Fl. N. Zel.* i. 248.—N. Zel.
Mitchelli, Lindl. in Mitch. Journ. Trop. Austral. 365 = rufa?
mutica, *R. Br. Prod.* 328.—Austral.
nana, *R. Br. l. c.* 327.—Austral.
nutans, *R. Br. l. c.*—Austral.
obtusa, *R. Br. l. c.*—Austral.
ophioglossa, *R. Br. l. c.* 326.—Austral.
parviflora, *R. Br. l. c.* 327.—Austral.
pedoglossa, *Fitzg. Austral. Orch.* i. t. 5.—Austral.
pedunculata, *R. Br. Prod.* 327.—Austral.; Tasman.
praecocissima, F. Muell. ex Benth. Fl. Austral. vi. 365 = vittata.
praecox, *Lindl. Gen. et Sp. Orch.* 388.—Austral.
puberula, *Hook. f. Fl. N. Zel.* i. 249.—N. Zel.
pyramidalis, Endl. in Lehm. Pl. Preiss. ii. 5 = reflexa.
pyramidalis, *Lindl. Swan River App.* 53.—Austral.
recurva, *Benth. Fl. Austral.* vi. 360.—Austral.
reflexa, *R. Br. Prod.* 327.—Austral.
revoluta, R. Br. l. c. = reflexa.

PTEROSTYLIS:—
rufa, *R. Br. Prod.* 327.—Austral.
rupestris, *Drumm. in Hook. Journ. Bot.* ii. (1840) 367.—Austral.
scabra, Lindl. Swan River App. 53 = reflexa.
scabrida, Lindl. Gen. et Sp. Orch. 389 = cucullata.
squamata, R. Br. Prod. 327 = rufa.
squamata, Lindl. Gen. et Sp. Orch. 388 = barbata.
striata, *Fitzg. Austral. Orch.* i. t. 5.—Austral.
trullifolia, *Hook. f. Fl. N. Zel.* i. 249.—N. Zel.
truncata, *Fitzg. Austral. Orch.* i. t. 3.—Austral.
turfosa, *Endl. in Lehm. Pl. Preiss.* ii. 5.—Austral.
vittata, *Lindl. Swan River App.* 53.—Austral.
Woollsii, *Fitzg. Austral. Orch.* i. t. 8.—Austral.

PTEROSTYRAX, Sieb. & Zucc. Fl. Jap. i. 94, t. 47 (1835) = **Halesia**, Linn. (Styrac.).
corymbosum, Sieb. & Zucc. l. c. 96 = H. corymbosa.
hispidum, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 132 = H. hispida.
micranthum, Sieb. & Zucc. l. c.—Japon.

PTEROTA, P. Br. Hist. Jamaic. 146. t. 5 (1756) = **Zanthoxylum**, Linn. (Rutac.).
Fagara, Crantz, Inst. ii. 417 = Z. Pterota.
piperita, Crantz, l. c. 418 = Z. pimpinelloides.
tragodes, Crantz, l. c. 417 = Z. tragodes.

PTEROTHECA, Cass. in Bull. Soc. Philom. (1816) 200. *COMPOSITAE*, Benth. & Hook. f. ii. 516.
CREPINIA, Reichb. in Moessl. Handb. ii. 1415 (1828).
MYOSERIS, Link, Enum. Hort. Berol. ii. 291 (1822).
TRICHOCREPIS, Vis. Stirp. Dalm. 19, t. 7 (1826).
TRICHOSERIS, Sch. Bip. in Flora, xxii. (1839) 19.
aralensis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 383.—Turkestan.
aspera, Reichb. f. Ic. Fl. Germ. xix. 37 = Crepis aspera.
bifida, C. B. Clarke, Comp. Ind. 256 = Falconeri.
bifida, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 43.—Oriens.
Dioscoridis, Reichb. f. Ic. Fl. Germ. xix. 37 = Crepis Dioscoridis.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 399.—Ind. or.
glareosa, *Trautv. in Act. Hort. Petrop.* iv. (1876) 386, in textu.—As. Min.
griselica, Serres, in Bull. Soc. Bot. Fr. iii. (1856) 275 = nemausensis.
macrantha, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 384, in obs.—As. centr.
nemausensis, *Cass. in Bull. Soc. Philom.* (1816) 200.—Europ. austr.
nemausensis, Reichb. Fl. Germ. Excurs. 269 = bifida.
nudicaulis, Dulac, Fl. Hautes-Pyr. 493 = nemausensis.
obovata, Boiss. Diagn. Ser. II. iii. 98 (= *Lagoseris orientalis*).—As. occ. et bor.
purpurea, C. B. Clarke, Comp. Ind. 256; Trautv. in Act. Hort. Petrop. iv. (1876) 386, in textu = bifida.
runcinata, *Trautv. l. c.* 385.—Armenia.
Ruppellii, Steud. Nom. ed. II. ii. 415 = bifida.
sancta, C. Koch, in Linnaea, xxiii. (1850) 692 = nemausensis.

PTEROTHECA, Presl, Symb. Bot. i. 55. t. 35 (1831) = **Rynchospora**, Vahl (Cyperac.).
Vahlii, Presl, l. c. = R. pterocarpa.

PTEROTHRIX, DC. Prod. vi. 279 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 324.
cymbaefolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 275.—Afr. austr.
perotrichoides, *Harv. l. c.*—Afr. austr.
spinescens, *DC. Prod.* vi. 280.—Afr. austr.
tomentosa, DC. l. c. = Amphiglossa tomentosa.

PTEROTRICHIS, Reichb. Nom. 129 (1841) = PHEROTRICHIS, Decne. = **Lachnostoma**, H. B. & K. (Asclep.).

PTEROTROPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 337 = **Thlaspi**, Tourn. (Crucif.).
beugesiaca, Fourr. l. c. = T. alpestre.
erratica, Fourr. l. c. = T. perfoliatum.
Gaudiniana, Fourr. l. c. = T. alpestre.
impropera, Fourr. l. c. = T. perfoliatum.
Lereschii, Fourr. l. c. = T. virgatum.
martialis, Jord. ex Fourr. l. c. = T. perfoliatum.
sylvestris, Fourr. l. c. = T. alpestre.
virens, Fourr. l. c. = T. alpestre.

PTEROTUM, Lour. Fl. Cochinch. 292 (1790). *INC. SEDIS.*
procumbens, *Lour. l. c.* 293.—Cochinch.

PTERYGIUM, Endl. Gen. 1013 (1841) = PTERIGIUM, Correa = **Dipterocarpus**, Gaertn.

PTERYGOCALYX, Maxim. Prim. Fl. Amur. 198, t. 9 (1858) = **Crawfurdia**, Wall. (Gentian.).
volubilis, Maxim. l. c.—Reg. Amur.

PTERYGOCARPUS, Hochst. in Flora, xxvi. (1843) 78 = **Dregea**, E. Mey. (Asclep.).
abyssinicus, Hochst. l. c. (= *Hoya africana*).—Afr. trop.

PTERYGODIUM, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 217. t. 3. *ORCHIDEAE*, Benth. & Hook. f. iii. 632.
OMMATODIUM, Lindl. Gen. et Sp. Orch. 365 (1838).
acutifolium, *Lindl. Gen. et Sp. Orch.* 366.—Afr. austr.
alatum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 218.—Afr. austr.
atratum, Sw. l. c. = Ceratandra chloroleuca.
caffrum, *Sw. l. c.*—Afr. austr.
carnosum, *Lindl. Gen. et Sp. Orch.* 367.—Afr. austr.
catholicum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 218.—Afr. austr.
cruciferum, *Sond. in Linnaea*, xix. (1847) 109.—Afr. austr.
inversum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 218.—Afr. austr.
magnum, *Reichb. f. in Flora*, l. (1867) 117.—Afr. austr.
platypetalum, *Lindl. Gen. et Sp. Orch.* 366.—Afr. austr.
rubiginosum, *Sond. ex Drège, in Linnaea*, xx. (1847) 220, nomen; *Bolus, in Journ. Linn. Soc.* xx. (1884) 486.—Afr. austr.
sulcata, Roxb. Hort. Beng. 63; Fl. Ind. iii. 452 = Zeuxine sulcata.
venosum, *Lindl. Gen. et Sp. Orch.* 367.—Afr. austr.
Volucris, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 218.—Afr. austr.

PTERYGOLEPIS, Reichb. Nom. 236 (1841) = PTEROLEPIS, Schrad. = **Scirpus**, Tourn. (Cyperac.).

PTERYGOLOMA, Hanst. in Linnaea, xxvi. (1853) 211 = **Alloplectus**, Mart. (Gesner.).
cristatum, Griseb. Fl. Brit. W. Ind. 464.—Ind. occ.
pubescens, Griseb. l. c.—Ind. occ.
repens, Hanst. in Linnaea, xxvi. (1853) 211 = A. repens.

PTERYGOPAPPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 120. *COMPOSITAE*, Benth. & Hook. f. ii. 303.
MAJA, Wedd. Chlor. And. i. 228, t. 27 D (1855).
Lawrencii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 120.—Tasman.

PTERYGOSTACHYUM, Nees, ex Steud. Syn. Pl. Gram. 413 (1855) = **Dimeria**, R. Br. (Gramin.).
Lehmanni, Nees, l. c.—Zeylan.

PTERYGOTA, Schott & Endl. Melet. 32 (1832) = **Sterculia**, Linn.
alata, R. Br. in Benn. Pl. Jav. Rar. 234 = S. alata.
alata, Thw. Enum. Pl. Zeyl. 29 = S. Thwaitesii.
brasiliensis, F. Allem. in Trab. Comm. Sc. Expl. Bot. i. (1862) 7.—Bras.
Roxburghii, Schott & Endl. Melet. 32 = S. alata.

PTERYPODIUM, Reichb. f. in Flora, l. (1867) 117, sphalm. = **Pterygodium**, Sw. (Orchid.).

PTERYXIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 624 (1840) = **Cymopterus**, Rafin. (Umbell.).
albiflora, Nutt. l. c. 625 = C. terebinthinus.
foeniculacea, Nutt. l. c. 624 = C. terebinthinus.
terebinthacea, Nutt. l. c. = C. terebinthinus.
thapsoides, Nutt. l. c. 625 = C. terebinthinus.

PTICHOCHILUS, Benth. in Journ. Linn. Soc. xviii. (1881) 341 = PTYCHOCHILUS, Schau. = **Tropidia**, Lindl. (Orchid.).

PTILAGROSTIS, Griseb. in Ledeb. Fl. Ross. iv. 447 (1853) = **Stipa**, Linn. (Gramin.).
mongholica, Griseb. l. c. = S. mongolica.

PTILANTHELIUM, Steud. Syn. Pl. Cyp. 166 (1855) = **Mesomelaena**, Nees (Cyperac.).
Chauvinii, Steud. l. c. 167 = M. deusta.

PTILEPIDA, Rafin. in Am. Monthly Mag. (1818) 268 = **Cephalophora**, Cav. (Compos.).

PTILERIS, Rafin. in Am. Monthly Mag. (1818) 268 = **Erechtites**, Rafin. (Compos.).
hieracifolius, Rafin. l. c. = E. praealta.

PTILIMNIUM, Rafin. in Am. Monthly Mag. (1819) 192 = **Discopleura**, DC. (Umbell.).
capillaceum, Rafin. l. c. = D. capillacea.
costatum, Rafin. Neogenyt. 2 = D. capillacea.
junceum, Rafin. l. c. = D. Nuttallii.

PTILINA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 479 (1840) = **Didiplis**, Rafin. (Lythrar.).
aquatica, Nutt. ex Torr. & Gray, l. c. 480 = D. linearis.

PTILIUM, Pers. Syn. i. 359, in syn. (1805) = PETILIUM, Linn. = **Fritillaria**, Tourn. (Liliac.).

PTILOCALYX, Torr. & Gray, in Pacif. Rail. Rep. ii. 170 (1855) = **Coldenia**, Linn. (Borag.).
Greggii, Torr. & Gray, l. c. = C. Greggii.

PTILOCHAETA, Nees, in Mart. Fl. Bras. ii. 1. 147. t. 8 (1842) = **Rynchospora**, Vahl (Cyperac.).
diodon, Nees, l. c. 148 = R. lunata.

PTILOCHAETA, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 52. *MALPIGHIACEAE*, Benth. & Hook. f. i. 258.
bahiensis, *Turcz. l. c.*—Bras.
nudipes, *Griseb. in Goett. Abh.* xxiv. (1879) 66.—Reg. Argent.

PTILOCNEMA, D. Don, Prod. Fl. Nep. 33 (1825) = **Pholidota**, Lindl. (Orchid.).
bracteata, D. Don, l. c. = Pholidota imbricata.

PTILOMERIS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 381 = **Baeria**, Fisch. & Mey. (Compos.).
affinis, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 174 = B. affinis.
anthemoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 382 = B. anthemoides.
aristata, Nutt. l. c. = B. coronaria.
coronaria, Nutt. l. c. = B. coronaria.
mutica, Nutt. l. c. = B. mutica.
tenella, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 173 = B. tenella.

PTILONEILEMA, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Gram. 201 (1855) = **Melanocenchris**, Nees (Gramin.).
plumosum, Steud. ll. cc. = M. plumosa.

PTILONELLA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 385 = **Blepharipappus**, Hook. (Compos.).
scabra, Nutt. l. c. 386 = B. scaber.

PTILOPHORA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 112 = **Microseris**, D. Don (Compos.).
major, A. Gray, l. c. 113 = M. major.
nutans, A. Gray, l. c. = M. nutans.

PTILORIA, Rafin. Atl. Journ. 145 (1832) = **Stephanomeria**, Nutt. (1841) (Compos.).
pauciflora, Rafin. l. c. = S. runcinata.
tenuifolia, Rafin. l. c. = S. minor.

PTILOSCIADIUM, Steud. Syn. Pl. Cyp. 149 (1855) = **Rynchospora**, Vahl (Cyperac.).
Lindenii, Steud. l. c. 150 = R. hirta.

PTILOSIA, Tausch, in Flora, xi. (1828) 1. Erg. 78, in nota = **Picris**, Linn. (Compos.).
apargioides, Tausch, l. c. = P. sulphurea.

PTILOSTEMON, Cass. in Bull. Soc. Philom. (1816) 200 = **Cnicus**, Linn. (Compos.).
appendiculatum, Cass. in Dict. Sc. Nat. xliv. 59 = C. fruticosus.
muticum, Cass. l. c. = C. Chamaepeuce.

PTILOSTEMUM, Steud. Nom. ed. II. ii. 415 (1841) = praec.

PTILOSTEPHIUM, H. B. & K. Nov. Gen. et Sp. iv. 253. tt. 387, 388 (1820) = **Tridax**, Linn. (Compos.).
balbisioides, Spreng. Syst. iii. 570 = T. balbisioides.
coronopifolium, H. B. & K. Nov. Gen. et Sp. iv. 255. t. 387 = T. coronopifolia.
trifidum, H. B. & K. l. c. 255. t. 388 = T. trifida.

PTILOTRICHUM, C. A. Mey. in Ledeb. Fl. Alt. iii. 64 (1831) = **Alyssum**, Tourn. (Crucif.).
Candolleanum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 13 = A. macrocarpum.
canescens, C. A. Mey. in Ledeb. Fl. Alt. iii. 66 = A. canescens.
cappadocicum, Boiss. Fl. Orient. i. 288.—As. Min.
cretaceum, Ledeb. Fl. Ross. i. 143 = A. cretaceum.
cyclocarpum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 159 (= *Koniga scardica*).—Macedon.; As. Min.
elongatum, C. A. Mey. in Ledeb. Fl. Alt. iii. 66 = A. canescens.
emarginatum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 160 (= *Koniga emarginata*).—Graecia.
glabrescens, Boiss. Fl. Orient. i. 289.—Cappadoc.
longicaule, Boiss. Elench. 13 = A. longicaule.
Peyrousianum, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 835 = A. Lapeyrousianum.
purpureum, Boiss. Elench. 12 = A. Lagascae.
pyrenaicum, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 835 = A. pyrenaicum.
rupestre, Boiss. Fl. Orient. i. 288, in obs. = A. rupestre.
saxigenum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 13 = A. macrocarpum.
spinosum, Boiss. Voy. Bot. Espagne, ii. 46 = A. spinosum.
strigulosum, Kunze, in Flora, xxix. (1846) 683 = A. maritimum.
tenuifolium, C. A. Mey. in Ledeb. Fl. Alt. iii. 67, in obs. = A. tenuifolium.
tortuosum, Willk. in Bot. Zeit. v. (1847) 234 = A. Lapeyrousianum.

PTILOTUM, Dulac, Fl. Hautes-Pyr. 313 (1867) = **Dryas**, Linn. (Rosac.).
octopetalum, Dulac, l. c. = D. octopetala.

PTILOTUS, R. Br. Prod. 415 (1810). *AMARANTACEAE*, Benth. & Hook. f. iii. 32.
aervoides, F. Muell. Fragm. vi. 231 = Trichinium aervoides.
alopecuroideus, F. Muell. l. c. 227 = Trichinium alopecuroideum.
amabilis, Span. in Linnaea, xv. (1841) 345 = corymbosus?
arthrolasius, F. Muell. Fragm. vi. 232 = Trichinium arthrolasium.
astrolasius, F. Muell. l. c. 233 = Trichinium astrolasium.
auriculifolius, F. Muell. Census, 28 = Trichinium auriculaefolium.
axillaris, F. Muell. l. c. = Trichinium axillare.
Beckeri, F. Muell. Fragm. vi. 233 = Trichinium Beckerianum.
brachyanthus, F. Muell. Census, 29 = Trichinium brachyanthum.
caespitulosus, F. Muell. Fragm. vi. 232 = Trichinium caespitulosum.
calostachyus, F. Muell. l. c. 231 = Trichinium calostachyum.
conicus, *R. Br. Prod.* 415.—Austral.
corymbosus, *R. Br. l. c.*—Austral.
declinatus, Nees, in Lehm. Pl. Preiss. i. 631 = Trichinium declinatum.
dissitiflorus, F. Muell. Census, 28 = Trichinium dissitiflorum.
distans, Poir. Encyc. Suppl. iv. 620 = Trichinium distans.
divaricatus, F. Muell. Fragm. vi. 229 = Trichinium divaricatum.

PTILOTUS:—
Drummondi, F. Muell. Fragm. vi. 229 = Trichinium Drummondii.
erubescens, Schlecht. in Linnaea, xx. (1847) 575 = Trichinium erubescens.
esquamatus, F. Muell. Census, 28 = Trichinium esquamatum.
exaltatus, Nees, in Lehm. Pl. Preiss. i. 630 = Trichinium exaltatum.
floribundus, F. Muell. Fragm. vi. 233 = Trichinium sericostachyum.
Forrestii, *F. Muell. Pl. N. W. Austral.* v. (1870) 240.—Austral.
Fraseri, F. Muell. Census, 29 = Trichinium Fraseri.
fusiformis, Poir. Encyc. Suppl. iv. 619 = Trichinium fusiforme.
gomphrenoides, *F. Muell. Fragm.* vi. 233, nomen; *et ex Benth. Fl. Austral.* v. 244.—Austral.
gracilis, Poir. Encyc. Suppl. iv. 620 = Trichinium gracile.
grandiflorus, *F. Muell. Fragm.* i. 237.—Austral.
helichrysoides, F. Muell. l. c. vi. 231 = Trichinium helichrysoides.
helipteroides, F. Muell. l. c. = Trichinium helipteroides.
hemisteirus, F. Muell. l. c. iv. 90 = Trichinium corymbosum.
holosericeus, F. Muell. l. c. vi. 229 = Trichinium holosericeum.
Hoodii, *F. Muell. l. c.* viii. 232.—Austral.
humilis, *F. Muell. l. c.* vi. 229.—Austral.
incanus, Poir. Encyc. Suppl. iv. 620 = Trichinium incanum.
lanatus, A. Cunn. ex Moq. in DC. Prod. xiii. II. 281 = Trichinium Cunninghamii.
latifolius, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 88.—Austral.
laxus, F. Muell. Census, 28 = Trichinium laxum.
lepidus, F. Muell. Fragm. iv. 89 = grandiflorus.
leucocomus, F. Muell. Census, 29 = Trichinium leucocomum.
Lindleyi, F. Muell. Fragm. vi. 233 = Trichinium obovatum.
macrocephalus, Poir. Encyc. Suppl. iv. 620 = Trichinium macrocephalum.
macrotrichus, *F. Muell. Fragm.* iv. 90.—Austral.
Manglesii, F. Muell. l. c. vi. 230 = Trichinium Manglesii.
Murrayi, *F. Muell. l. c.* iii. 145.—Austral.
nobilis, F. Muell. l. c. vi. 227 = Trichinium nobile, exaltatum, semilanatum.
obovatus, F. Muell. l. c. 228 = Trichinium obovatum.
ovatus, Moq. in DC. Prod. xiii. II. 281 = Psilotrichum calceolatum.
pachocephalus, F. Muell. Fragm. vi. 228 = Trichinium macrocephalum.
parviflorus, F. Muell. Census, 28 = Trichinium parviflorum.
parvifolius, F. Muell. Fragm. vi. 229 = Trichinium parvifolium.
Polakii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 274.—Austral.
polystachyus, F. Muell. Fragm. vi. 230 = Trichinium polystachyum, Stirlingii, roseum, laxum.
psilotrichoides, F. Muell. Census, 29 = Trichinium corymbosum.
pyramidatus, F. Muell. Fragm. vi. 230 = Trichinium pyramidatum.
Roei, F. Muell. Census, 29 = Trichinium Roei.
roseus, F. Muell. l. c. 28 = Trichinium roseum.
rotundifolius, F. Muell. Fragm. vi. 230 = Trichinium rotundifolium.
sandwicensis, *A. Gray, ex H. Mann, in Proc. Am. Acad.* vii. (1867) 200.—Ins. Sandvic.
sericostachyus, F. Muell. Fragm. vi. 230 = Trichinium sericostachyum.
Sewelliae, *F. Muell.* '*in Vict. Naturalist* (Jan. 1885)' (nobis latet).—Austral.
spathulatus, Poir. Encyc. Suppl. iv. 620 = Trichinium spathulatum.
spicatus, *F. Muell. ex Benth. Fl. Austral.* v. 243.—Austral.
Stirlingi, F. Muell. Census, 28 = Trichinium Stirlingii.
striatus, F. Muell. l. c. = Trichinium striatum.
trichotoma, Miq. Fl. Ind. Bat. i. 1. 1037 = Psilotrichum trichotomum.
villosiflorus, *F. Muell. Fragm.* iii. 125.—Austral.

PTILURUS, D. Don, in Trans. Linn. Soc. xvi. (1830) 218 = **Leuceria**, Lag. (Compos.).
daucifolius, D. Don, l. c. 219.—Peruv.

PTOSIMOPAPPUS, Boiss. Voy. Bot. Espagne, ii. 739 (1845) = **Centaurea**, Linn. (Compos.).
arboreus, Walp. Ann. ii. 930 = C. arborea.
argutus, Boiss. Diagn. Ser. II. iii. 60 = C. arguta.
bracteatus, Boiss. l. c. Ser. I. x. 104 = C. bracteata.
canariensis, Boiss. Voy. Bot. Espagne, ii. 740 = C. canariensis.
intybaceus, Boiss. l. c. = C. intybacea.
sempervirens, Boiss. l. c. = C. sempervirens.
uliginosus, Boiss. Diagn. Ser. II. iii. 60 = C. uliginosa.
Webbianus, Boiss. l. c. 61 = C. Webbiana.

PTYAS, Salisb. Gen. Pl. Fragm. 76 (1866) = **Aloë**, Tourn. (Liliac.).
disticha, Salisb. l. c. = A. disticha.

PTYCANTHERA, Decne. in DC. Prod. viii. 606 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 768.
Berterii, *Decne. l. c.*—Ins. S. Doming.

PTYCHANDRA, Scheff. in Ann. Jard. Buitenz. i. (1876) 140. *PALMAE*, Benth. & Hook. f. iii. 893.
glauca, *Scheff. l. c.* 160.—Ins. Batjan.
Musschenbroekiana, *Becc. Malesia*, i. (1877) 100, in obs.—Ins. Ternate.

PTYCHOCARPA, Spach, Hist. Vég. Phan. x. 402 (1841) = **Grevillea**, R. Br. (Proteac.).

PTYCHOCARYA, R. Br. in Wall. Cat. n. 3538 (1830) = **Scirpodendron**, Zipp. (Cyperac.).
macrocarpa, R. Br. ex Steud. Nom. ed. II. ii. 416 = S. costatum.

PTYCHOCENTRUM, Benth. in Ann. Wien. Mus. ii. (1838) 113 = **Rhynchosia**, Lour. (Legumin.).

PTYCHOCHILUS, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) (Pl. Meyen.) 431. t. 12. B = **Tropidia**, Lindl. (Orchid.).
septemnervis, Schau. l. c. = T. curculigoides.

PTYCHOCOCCUS, Becc. in Ann. Jard. Butenz. ii. (1885) 90. *PALMAE*.
arecinus, *Becc. l. c.* 99.—N. Guin.
paradoxus, *Becc. l. c.* 96.—N. Guin.

PTYCHODEA, Willd. ex Cham. & Schlecht. in Linnaea, iv. (1829) 168 = **Sipanea**, Aubl. (Rubiac.).
pedunculata, Willd. ex Cham. & Schlecht. l. c. = S. dichotoma.
sessiliflora, Willd. ex Cham. & Schlecht. l. c. = S. glomerata.

PTYCHODON, Klotzsch, ex Reichb. Nom. 172 (1841) = **Lafoensia**, Vand. (Lythrar.).
nummularifolius, Klotzsch, ex Koehne, in Mart. Fl. Bras. xiii. II. 347 = L. nummularifolia.

PTYCHOLEPIS, Griseb. in Lechl. Berb. Am. Austr. 58 (1857) = **Blepharodon**, Decne. (Asclep.).
ditassoides, Griseb. l. c.—Peruv.

PTYCHOMERIA, Benth. in Hook. Kew Journ. vii. (1855) 14 = **Gymnosiphon**, Blume (Burmann.).
capitata, Benth. l. c. 15.—Bras.
cornuta, Benth. l. c. 16 = G. cornutum.
cymosa, Benth. l. c. 15 = G. cymosum.
divaricata, Benth. l. c. 16 = G. divaricatum.
fimbriata, Benth. l. c. 14.—Bras.
mutica, Benth. l. c. 16.—Bras.
nivea, Griseb. Cat. Pl. Cub. 257.—Cuba.
tenella, Benth. in Hook. Kew Journ. vii. (1855) 17.—Bras.

PTYCHOPETALUM, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 376. *OLACINEAE*, Benth. & Hook. f. i. 346.
ATHESIANDRA, Miers, ex Benth. & Hook. f. Gen. i. 346 (1862).
anceps, *Oliver, Fl. Trop. Afr.* i. 347.—Afr. trop.
olacoides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 377.—Guian.
petiolatum, *Oliver, Fl. Trop. Afr.* i. 347.—Afr. trop.

PTYCHOPYXIS, Miq. Fl. Ind. Bat. Suppl. 402 (1860). *INC. SEDIS.*—Cf. Benth. & Hook. f. Gen. i. 217.
costata, *Miq. l. c.*—Sumatra.

PTYCHORAPHIS, Becc. in Ann. Jard. Bot. Buitenz. ii. (1885) 90. *PALMAE*.
augusta, *Becc. l. c.*—Ins. Nicobar.
philippinensis, *Becc. l. c.*—Ins. Philipp.
singaporensis, *Becc. l. c.*—Malaya.

PTYCHOSEMA, Benth. in Lindl. Swan River App. 16 (1839). *LEGUMINOSAE*, Benth. & Hook. f. i. 496.
anomalum, *F. Muell. Fragm.* ix. 62.—Austral.
pusillum, *Benth. in Lindl. Swan River App.* 16.—Austral.
trifoliolatum, *F. Muell, in Wing, South. Sc. Record*, ii. (1882) 72.—Austral.

PTYCHOSPERMA, Labill. in Mém. Inst. Par. ix. 1808 (1809), 252. *PALMAE*, Benth. & Hook. f. iii. 891.
NORMANBYA, F. Muell. Fragm. xi. 57, in obs. (1878); Becc. in Ann. Jard. Buitenz. ii. (1885) 91.
PSYCHOSPERMA, Dum. Anal. Fam. 56 (1829).
SEAFORTHIA, R. Br. Prod. 267 (1810).
alba, Scheff. in Tijdschr. Nederl. Ind. xxvii. (1864) 26 = Dictyosperma album.
Alexandrae, F. Muell. Fragm. v. 47 = Archontophoenix Alexandrae.
angustifolia, Blume, Rumphia, ii. 122 = Coleospadix oninensis.
appendiculata, Blume, l. c. = Drymophloeus appendiculatus.
arecina, Becc. Malesia, i. (1877) 58 = Ptychococcus arecinus.
arfakiana, Becc. l. c. 57 = Rhopaloblaste hexandra.
Beatricae, *F. Muell. in Melb. Chemist* (Feb. 1882).—Austral.
caesia, Miq. Fl. Ind. Bat. iii. 20 = Pinanga caesia.
calapparia, Miq. l. c. = Actinorhytis calapparia.
Capitis-Yorkii, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 217.—Austral.
caudata, Becc. Malesia, i. (1877) 55 = Pinanga caudata (in Addend.).
coccinea, Hort. ex Teijsm. & Binn. Cat. Hort. Bog. 69 = Cyrtostachys Renda.
cochinchinensis, Miq. Fl. Ind. Bat. iii. 23 = Pinanga cochinchinensis.
communis, Miq. l. c. 31 = Coleospadix oninensis.
coronata, Miq. l. c. 24 = Pinanga coronata.
costata, Miq. l. c. 25 = Pinanga Kuhlii.
Cunninghamiana, H. Wendl. in Bot. Zeit. xvi. (1858) 346 = Archontophoenix Cunninghamii.
Dicksoni, Miq. Fl. Ind. Bat. iii. 23 = Pinanga Dicksonii.
disticha, Miq. l. c. 28 = Pinanga disticha.
Drudei, *H. Wendl. in Benth. & Hook. f. Gen.* iii. 892.—Austral.
elegans, *Blume, Rumphia*, ii. 118.—Austral.
elegans, Hort.—Cf. Gard. Chron. (1870) 419 = Archontophoenix Cunninghamii.
filifera, H. Wendl. in Bonplandia, x. (1862) 195 = Vitiphoenix filifera.
furfuracea, Miq. Fl. Ind. Bat. iii. 26 = Pinanga furfuracea.
gracilis, *Labill. in Mém. Inst. Par.* ix. 1808 (1809), 253.—Nov. Hibernia.
inaequalis, Miq. Fl. Ind. Bat. iii. 27 = Pinanga inaequalis.
Junghuhnii, Miq. l. c. 28 = Pinanga Junghuhnii.
Kuhlii, Miq. l. c. 21 = Pinanga Kuhlii.
Laccospadix, Benth. Fl. Austral. vii. 140 = Calyptrocalyx australasicus.
lacerata, *Hort.—Cf. Gard. Chron.* (1868) 1317.—Am. centr.
latisecta, Miq. Fl. Ind. Bat. iii. 20 = Pinanga latisecta.
litigiosa, Becc. Malesia, i. (1877) 50 = Coleospadix litigiosa.
Macarthurii, *H. Wendl. ex Kew Report*, 1882 (1884), 55.—Austral.
maculata, Seem. in Gard. Chron. (1870) 697 = Pinanga maculata.
malaiana, Miq. Fl. Ind. Bat. iii. 25 = Pinanga malaiana.
micrantha, Becc. Malesia, i. (1877) 52 = Rhopaloblaste micrantha.
minor, Miq. Fl. Ind. Bat. iii. 23 = Pinanga minor.
Musschenbroekiana, Becc. Malesia, i. (1877) 53 = Ptychandra Musschenbroekiana.

PTYCHOSPERMA:—
Nenga, Teijsm. & Binn. Cat. Hort. Bog. 69 = Nenga Wendlandiana.
nobilis, *Seem. in Gard. Chron.* (1870) 697.—Hab. ?
Normanbyi, F. Muell. Fragm. xi. 56 = Areca Normanbyi.
noxa, Miq. Fl. Ind. Bat. iii. 23 = Pinanga noxa.
paniculata, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 3 = Mischophloeus paniculatus.
paradoxa, Scheff. in Ann. Jard. Buitenz. i. (1876) 83 = Ptychococcus paradoxus.
patula, Miq. Fl. Ind. Bat. iii. 26 = Pinanga patula.
pauciflora, *H. Wendl. in Bonplandia*, x. (1862) 193. —Ins. Fiji.
perbrevis, H. Wendl. l. c. 193 = Balaka perbrevis.
Pikeringii, *H. Wendl. l. c.* 194.—Ins. Fiji.
polystachya, Miq. Fl. Ind. Bat. Suppl. 590 = Areca triandra.
punicea, Miq. Fl. Ind. Bat. iii. 31 = Pinanga ternatensis.
Rumphii, Blume, Rumphia, ii. 119 = Drymophloeus olivaeformis.
rupicola, Thw. Enum. Pl. Zeyl. 328 = Loxococcus rupicola.
salicifolia, Miq. Fl. Ind. Bat. iii. 28 = Pinanga salicifolia.
saxatilis, Blume, Rumphia, ii. 121 = Drymophloeus saxatilis.
Seaforthii, Miq. Fl. Ind. Bat. iii. 21 = elegans, Archontophoenix Cunninghamii.
Seemannii, H. Wendl. in Bonplandia, ix. (1861) 261; x. (1862) 192 = Balaka Seemanni.
simplicifrons, Miq. Fl. Ind. Bat. Suppl. 590 = Pinanga disticha.
singaporensis, Becc. Malesia, i. (1877) 61 = Ptychoraphis singaporensis.
sylvestris, Miq. Fl. Ind. Bat. iii. 22 = Pinanga javana.
tahitensis, *H. Wendl. in Bonplandia*, x. (1862) 196.—Ins. Tahiti.
tenella, *H. Wendl. in Bot. Zeit.* xvii. (1859) 63.—Borneo.
Veitchii, H. Wendl. in Linnaea, xxxix. (1875) 213 = Archontophoenix Veitchii.
vestiaria, Miq. Fl. Ind. Bat. iii. 31 = Areca vestiaria.
vitiensis, *H. Wendl. in Bonplandia*, ix. (1861) 260; x. (1862) 195.—Ins. Fiji.

PTYCHOSTIGMA, Hochst. in Flora, xxvii. (1844) 23 = **Galiniera**, Delile (Rubiac.).
saxifragum, Hochst. l. c. = G. coffeioides.

PTYCHOTIS, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 124 = **Carum**, Rupp. (Umbell.).
achilleifolia, DC. Prod. iv. 109 = Pimpinella achilleifolia.
Ajowan, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 497 = C. copticum.
amoides, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 125 = C. ammoides.
anethifolia, DC. Prod. iv. 108 = C. anethifolium.
arabica, T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 21.—Aden.
aspera, Pomel, Nouv. Mat. Fl. Atl. 139 = C. ammoides.
atlantica, Coss. & Dur. in Bull. Soc. Bot. Fr. ix. (1862) 296.—Alger.
barbata, DC. Prod. iv. 109 = Pimpinella barbata.
Bunius, Reichb. Fl. Germ. Excurs. 474 = C. Bunius.
carvifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 537 = C. carvifolium.
coptica, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 496 = C. copticum.
dichotoma, Coss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 469 = C. dichotomum.
didyma, Sond. in Harv. & Sond. Fl. Cap. ii. 538 = C. didymum.
foeniculifolia, Steud. Nom. ed. II. ii. 572, in syn. = C. copticum.
heterophylla, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) 125 = C. Bunius.
hispida, Sond. in Harv. & Sond. Fl. Cap. ii. 537 = C. hispidum.
involucrata, Royle, Illustr. Bot. Himal. 229 = C. Roxburghianum.
Meisneri, Sond. in Harv. & Sond. Fl. Cap. ii. 536 = Apium humile.
puberula, DC. Prod. iv. 109 = Pimpinella puberula.
rigens, Harv. Gen. S. Afr. Pl. 134 (= *Bunium rigens*). —Afr. austr.
Roxburghiana, DC. Prod. iv. 109 = C. Roxburghianum.

PTYCHOTIS:—
sylvestris, Royle, Illustr. Bot. Himal. 228, 229 = Carum sylvestre.
tenuis, Sond. in Harv. & Sond. Fl. Cap. ii. 537. Afr. austr.
Thorei, Gren. & Godr. Fl. Fr. i. 735 = C. inundatum.
Timbali, Jord. Cat. Gr. Réc. Grenoble (1850), 15.—Gallia.
trachysperma, Boiss. Voy. Bot. Espagne, ii. 731 = ammoides.
Vargasiana, DC. Prod. iv. 109.—Venezuela.
verticillata, Duby, Bot. Gall. ed. II. i. 235 = C. ammoides.

PTYSSIGLOTTIS, T. Anders. in Thw. Enum. Pl. Zeyl. 235 (1860). *ACANTHACEAE*, Benth. & Hook. f. ii. 1117.
radicosa, *T. Anders. l. c.*—Zeylan.

PTYXOSTOMA, Vahl, in Danske Nat. Selsk. Skriv. vi. (1810) 95 = **Lonchostoma**, Wikstr. (1818) (Bruniac.).
monogyna, Vahl, l. c.—Afr. austr.
myrtoides, Vahl, l. c. 97 = L. acutiflorum.

PUBETA, Linn. Diss. Pl. Surin. 16 (1775); Amoen. Acad. viii. 264 (1785) = **Duroia**, Linn. f. (Rubiac.).

PUBILARIA, Rafin. Fl. Tellur. ii. 27 (1836) = **Simethis**, Kunth (1843) (Liliac.).
bicolor, Rafin. l. c. = S. bicolor.

PUCCINELLIA, Parl. Fl. Ital. i. 366 (1848) = **Glyceria**, R. Br. (Gramin.).
convoluta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 184 = G. distans.
distans, Parl. Fl. Ital. i. 367 = G. distans.
festucaeformis, Parl. l. c. 368 = G. distans.
Gussonii, Parl. l. c. 369 = G. distans.
maritima, Parl. l. c. 370 = G. maritima.
permixta, Parl. l. c. = G. Borreri.

PUCEDANUM, Hill, Hort. Kew. 94 (1768), err. typ. = **Peucedanum**, Tourn. (Umbell.).

PUERARIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 97. *LEGUMINOSAE*, Benth. & Hook. f. i. 537.
NEUSTANTHUS, Benth. in Miq. Pl. Jungh. i. 234 (1852).
anabaptis, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 253 = Shuteria hirsuta.
brachycarpa, *Kurz, l. c.* xlii. (1873) II. 232.—Burma.
Candollii, *Wall. Cat.* n. 5355.—Ind. or.
composita, R. Grah. in Wall. l. c. 5570 = Wallichii.
ferruginea, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 232 = Shuteria ferruginea.
hirsuta, *Kurz, l. c.* 254.—Pegu.
javanica, Benth. in Journ. Linn. Soc. ix. (1867) 125 = phaseoloides.
peduncularis, *R. Grah. in Wall. Cat.* n. 5354.—Reg. Himal.
phaseoloides, *Benth. in Journ. Linn. Soc.* ix. (1867) 125.—Ind. or.; Malaya; China.
Stracheyi, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 198.—Ind. or.
stricta, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 254.—Burma.
subspicata, Benth. in Journ. Linn. Soc. ix. (1867) 125 = phaseoloides.
Thomsoni, *Benth. l. c.* 122.—Reg. Himal.
Thunbergiana, *Benth. l. c.*—China; Japon.
tuberosa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 97.—Reg. Himal.
Wallichii, *DC. l. c.*—Reg. Himal.

PUGIONELLA, Salisb. Gen. Pl. Fragm. 128 (1866) = **Strumaria**, Jacq. (Amaryll.).
angustifolia, Salisb. l. c. = S. angustifolia.

PUGIONIUM, Gaertn. Fruct. ii. 291. t. 142 (1791). *CRUCIFERAE*, Benth. & Hook. f. i. 96.
cornutum, *Gaertn. l. c.*—Oriens.
dolabratum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 426.—Mongolia.

PUGIOPAPPUS, A. Gray, in Pacif. Rail. Rep. iv. 104 (1857) = **Leptosyne**, DC. (Compos.).
Bigelovii, A. Gray, l. c. = L. Bigelovii.
Breweri, A. Gray, in Proc. Am. Acad. viii. (1873) 660 = L. Bigelovii.
calliopsideus, A. Gray, l. c. 660 = L. calliopsidea.

PUJA, Molina, Sagg. Chile, ed. II. 153 (1810) = **Puya**, Molina (Orchid.).

PUKANTHUS, Rafin. Sylva Tellur. 53 (1838) = **Grabowskia**, Schlecht. (Solan.).
odoratus, Rafin. l. c. = G. boerhaaviaefolia.

PUKATERIA, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 120 = **Griselinia**, Forst. (Cornac.).
littoralis, Raoul, l. c. = G. lucida.

PULCHERIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Kadsura**, Kaempf. (Magnol.).

PULCHIA, Steud. Nom. ed. II. ii. 416 (1841), sphalm. = PLUCHIA, Vell. = **Diclidanthera**, Mart. (Styrac.).

PULEGIUM, Ray, ex Mill. Gard. Dict. ed. VI (1752) = **Mentha**, Tourn. (Labiat.).
aromaticum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 367 = M. Pulegium.
cervinum, Mill. Gard. Dict. ed. VIII. n. 3 = Preslia cervina.
erectum, Mill. l. c. n. 2 = M. Pulegium.
heterophyllum, Opiz, ex Boenn. Prod. Fl. Monast. 174 = M. Pulegium.
micranthum, Claus, in Beitr. Pfl. Russ. Reich. viii. 244 = M. Pulegium.
pubescens, Opiz, ex Boenn. Prod. Fl. Monast. 174 = M. Pulegium.
tomentella, Presl, Fl. Sic. p. xxxvi = M. Pulegium.
vulgare, Mill. Gard. Dict. ed. VIII. n. 1 = M. Pulegium.

PULICARIA, Gaertn. Fruct. ii. 461. t. 173 (1719). *COMPOSITAE*, Benth. & Hook. f. ii. 335.
DUCHESNIA, Cass. in Bull. Soc. Philom. (1817) 153.
FRANCOEURIA, Cass. in Dict. Sc. Nat. xxxviii. 374 (1825).
FRANKOERIA, Steud. Nom. ed. II. i. 646 (1840).
KILIANA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) I. Intell. 26, nomen.
PLATYCHAETA, Boiss. Diagn. Ser. I. xi. 5 (1849).
PLATZCHAETE, Sch. Bip. in F. Schultz, Arch. Fl. 131 (1855).
POLOA, DC. in Guill. Arch. Bot. ii. (1833) 514.
PTEROCHAETA, Boiss. Diagn. Ser. I. vi. 76 (1845).
STRABONIA, DC. Prod. v. 481 (1836).
TUBILIUM, Cass. in Bull. Soc. Philom. (1817) 153.
adenophora, *Franch. in Révoil, Çomali*, 37.—Afr. trop.
angustifolia, *DC. Prod.* v. 479.—Ind. or.
annua, Gaertn. Fruct. ii. 462 = Erigeron annuus.
arabica, Bourg. ex Nym. Consp. 394 = hispanica.
arabica, *Cass. in Dict. Sc. Nat.* xliv. 94.—Oriens.
argyrophylla, *Franch. in Révoil, Çomali*, 38.—Afr. trop.
armena, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 204.—Armenia.
aromatica, R. Br. in Salt, Abyss. App. p. 65 = undulata.
aspera, Pomel, Nouv. Mat. Fl. Atl. 35 = longifolia.
Aucheri, *Jaub. & Spach, Illustr.* iv. 78. t. 346.—Persia.
bellidiflora, Wallr. Sched. Crit. 483 = Erigeron annuus.
Boissieri, *Hook. f. Fl. Brit. Ind.* iii. 300.—Ind. or.
calycina, Presl, Delic. Prag. 96 = Inula montana.
canariensis, *C. Bolle, in Bonplandia*, vii. (1859) 295.—Ins. Canar.
capensis, *DC. Prod.* v. 479.—Afr. austr.
chrysopsidoides, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 153, nomen = petiolaris.
Clausonis, Pomel, Nouv. Mat. Fl. Atl. 36 = vulgaris.
congesta, C. Koch, in Linnaea, xxiii. (1850) 711 = odora.
crispa, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 223, in syn.; *Benth. & Hook. f. Gen.* ii. 336.—Afr. bor. et trop.; Arab.; Ind. or.
demissa, *Boiss. Fl. Orient.* iii. 205.—Afghan.
dentata, DC. Prod. v. 480 = dysenterica.
dentata, Guss. Fl. Sic. Syn. ii. 502 = vulgaris.
desertorum, *DC. ex Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 263.—Arab.
diversifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 840.—Ins. Socotra.
dumetorum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 501 = odora.
dysenterica, *Gaertn. Fruct.* ii. 462.—Europ.

PULICARIA:—
elata, Boiss. Diagn. Ser. I. xi. 6 = dysenterica.
erigeroides, DC. Prod. v. 480 = capensis.
fetida, *Rafin. Fl. Tellur.* ii. 48.—Hab. ?
filaginoides, *Pomel, Nouv. Mat. Fl. Atl.* 36.—Alger.
foliolosa, *DC. Prod.* v. 480.—Ind. or.
germanica, J. & C. Presl, Fl. Čech. 178 = vulgaris.
glaucescens, *Jaub. & Spach, Illustr.* iv. 79. t. 349.—Ind. or.; Persia.
glutinosa, *Jaub. & Spach, l. c.* 77. t. 348.—Arab.
gnaphalodes, *Boiss. Diagn.* Ser. I. vi. 76.—Persia; Turkestan; Afghan.
gracilis, *Heimerl, in Denkschr. Acad. Wien*, l. (1885) 56.—Persia.
gracilis, *Nym. Syll.* 19.—Lusit.
graeca, Sch. Bip. ex Nym. Consp. 394 = vulgaris.
grandidentata, *Jaub. & Spach, Illustr.* iv. 71. t. 345.—Arab.
Grantii, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 96.—Afr. trop.
graveolens, Nym. Syll. 19 = Inula graveolens.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 326.—Afr. trop.
hirta, J. & C. Presl, Fl. Čech. 178 = Inula hirta.
hispanica, *Boiss. Fl. Orient.* iii. 205, in obs.—Lusit.; Hispan.
hydrabadensis, Lindl. ex DC. Prod. v. 480 = Wightiana.
incisa, DC. l. c. 479 = undulata.
indica, Jaub. & Spach, Illustr. iv. 67. t. 343 = angustifolia.
indica, Rafin. Fl. Tellur. ii. 48 (Quid ?).—Hab. ?
inuloides, *DC. Prod.* v. 480.—Afr. bor.
inuloides, Schweinf. ex Ball, in Journ. Linn. Soc. xvi. (1878) 501 = longifolia.
involucrata, R. Br. in Salt, Abyss. App. 65, nomen = Ruppellii.
Kotschyi, Sch. Bip. ex Boiss. Fl. Orient. iii. 202 = dysenterica.
Kurtziana, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 327.—Afr. trop.
longifolia, *Boiss. Diagn.* Ser. II. iii. 16.—Afr. bor.
mauritanica, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 61, nomen.—Marocc.
monocephala, *Franch. in Révoil, Çomali*, 36.—Afr. trop.
odora, *Reichb. Fl. Germ. Excurs.* 239.—Europ. austr.
orientalis, *Jaub. & Spach, Illustr.* iv. 65. t. 342.—Arab.
paludosa, *Steud. Nom.* ed. II. ii. 416.—Lusit.
paludosa, Welw. ex Nym. Consp. 394 = hispanica.
palustris, *Hoffmgg. & Link, Fl. Portug.* ii. 293.—Lusit.
persica, Jaub. & Spach, Illustr. iv. 75. t. 347 = Aucheri.
petiolaris, *Jaub. & Spach, l. c.* 69. t. 344.—Arab.
planifolia, *Steud. Nom.* ed. II. ii. 416.—Lusit.
pratensis, Scheele, in Linnaea, xviii. (1844) 460 = vulgaris.
prostrata, Aschers. Fl. Brandenb. 304 = vulgaris.
Renschiana, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 197.—Afr. trop.
revoluta, *Nym. Syll.* 19.—Lusit.
rufescens, *Nym. l. c.*—Lusit.
Ruppellii, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 401.—Afr. trop.
sakhiana, *Klatt, in Sitzb. Akad. Muench.* (1878) 1. 87.—Reg. Himal.
salicina, J. & C. Presl, Fl. Cech. 178 = Inula salicina.
salviaefolia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 331.—As. centr.
saxicola, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 174 = angustifolia.
Schimperi, *DC. Prod.* vii. 285.—Arab.
sicula, *Moris, Fl. Sard.* ii. 363.—Europ. austr.
stephanocarpa, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 840.—Ins. Socotra.
Steveni, Nym. Syll. 19 = dysenterica.
Stocksii, Boiss. Diagn. Ser. II. iii. 16 = angustifolia.
Stocksii, *Hook. f. Fl. Brit. Ind.* iii. 300.—Ind. or.
trichocephala, DC. Prod. v. 478 = arabica.
uliginosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 463 = vulgaris.
uliginosa, *Hoffmgg. & Link, Fl. Portug.* ii. 294.—Lusit.
uliginosa, Stev. ex DC. Prod. v. 478 = dysenterica.
undulata, *Kostel. Allg. Med.-Pharm. Fl.* ii. 660; *DC. Prod.* v. 479.—Aegypt.; Arab.
undulata, C. A. Mey. Verz. Pfl. Cauc. 79 = vulgaris.

PULICARIA:—
vieraeoides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 840.—Ins. Socotra.
villosa, Link, Enum. Hort. Berol. ii. 336 = arabica.
viscida, R. Br. in Salt, Abyss. App. p. lxv = Felicia abyssinica.
viscosa, Koch, Syn. Fl. Germ. ed. I. 361; ed. II. 395 = Inula viscosa.
vulgaris, *Gaertn. Fruct.* ii. 461. t. 173.—Geront. temp.
Wightiana, *C. B. Clarke, Comp. Ind.* 128.—Ind. or.

PULMONARIA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 37 (1737). *BORAGINEAE*, Benth. & Hook. f. ii. 857.
affinis, Jord. Cat. Jard. Dijon (1848), 13 = saccharata.
alpestris, Lamotte, Prod. Fl. Centr. Fr. ii. 535 = officinalis.
alpina, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
alpina, Torr. in Ann. Lyc. N. York, ii. (1828) 224 = Mertensia alpina.
amoena, Stev. ex Lehm. Asperif. 296 = Mertensia dahurica.
angustata, Schrad. ex Roem. & Schult. iv. 54 = angustifolia.
angustifolia, Bess. Prim. Fl. Galic. i. 150 = mollis.
angustifolia, *Linn. Sp. Pl.* 135.—Europ.
armena, Schrank, in Nov. Act. Nat. Cur. ix. (1818) 99 = Mertensia dahurica.
azurea, Bess. Prim. Fl. Galic. i. 150 = angustifolia.
bracteata, Willd. ex Roem. & Schult. Syst. iv. 747 = Mertensia sibirica.
ciliata, James, ex Torr. in Ann. Lyc. N. York, ii. (1828) 224 = Mertensia sibirica.
Clusii, Baumg. Enum. Stirp. Transs. i. 123 = angustifolia.
cordato-folia, Gilib. Fl. Lituan. i. 23 = officinalis.
daurica, Cat. Hort. Viln. (1824); ex Roem. & Schult. Syst. iv. 57 = seq. ?
dauurica, Schrank, in Nov. Act. Nat. Cur. ix. (1818) 103 = seq.
davurica, Sims, Bot. Mag. t. 1743 = Mertensia dahurica.
denticulata, Roem. & Schult. Syst. iv. 746 = Mertensia sibirica.
digenea ×, Kern. in Oestr. Bot. Zeitschr. xxiii. (1873) 180.—Austria.
elegans, Hort. ex Roem. & Schult. Syst. iv. 745 = Mertensia paniculata.
elliptica, Rafin. New Fl. Am. iv. 17 (Mertensiae sp. ?).—Am. bor.
Gallorum, Gueldenst. Reisen, ii. 262 = officinalis.
glabra, Stokes, Bot. Mat. Med. i. 282 = Mertensia pulmonarioides.
gracilis, Willd. ex Roem. & Schult. Syst. iv. 747 = Mertensia dahurica.
grandiflora, DC. Cat. Hort. Monsp. (1813) 135 = saccharata.
hirta, Linn. Sp. Pl. ed. II. 1667 = Hyoscyamus physaloides.
hybrida ×, Kern. Monog. Pulmonar. 31.—Europ.
Kochii, Kern. in Oestr. Bot. Zeitschr. xxiii. (1873) 181 = *oblongata*.
lanceolata, Pursh, Fl. Am. Sept. ii. 729 = Mertensia lanceolata.
longifolia, Bast. ex Bor. Fl. Centr. Fr. ed. III. ii. 460 = angustifolia.
maculata, F. G. Dietr. Vollst. Lexic. Gaertn. v. 657 = officinalis.
maculata, Stokes, Bot. Mat. Med. i. 278 = angustifolia.
maculosa, Georgi, Reise, i. 2 = mollis.
maculosa, Reichb. Fl. Germ. Excurs. 338, in syn. = officinalis.
marginata, Nutt. Gen. Am. i. 115 = Mertensia lanceolata.
maritima, Linn. Sp. Pl. 136 = Mertensia maritima.
media, Host, Fl. Austr. i. 235 = angustifolia.
mollis, Guép. Fl. Maine-et-Loire, ed. III. 162, partim = *ovalis*.
mollis, Tenore, Syll. Fl. Neap. 84 = officinalis.
mollis, *Wolff, ex Heller, Fl. Wirceb. Suppl.* 13.—Europ. centr.
mollissima, Kern. Monog. Pulmonar. 47 = mollis.
montana, Lej. Fl. Spa, i. 98 = mollis.
montana, Wulf. ex Schrank, in Nov. Act. Nat. Cur. ix. (1818) 98 = angustifolia.
notha ×, Kern. Monog. Pulmonar. 32.—Europ.
oblongata, Schrad. in Roem. & Schult. Syst. iv. 744 (farrago spp.).—Europ.

PULMONARIA:—
oblongifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 43 = Mertensia oblongifolia.
obscura, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 341 = officinalis.
obscura, Rehm. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 495 = mollis.
officinalis, *Linn. Sp. Pl.* 135.—Europ.
officinalis, Oed. in Fl. Dan. t. 482 = saccharata.
orientalis, Mill. Gard. Dict. ed. VIII. n. 5 = Nonnea rosea.
ovalis ×, Bast. Fl. Maine-et-Loire, Suppl. 44.—Europ.
paniculata, Ait. Hort. Kew. ed. I. i. 181 = Mertensia paniculata.
parviflora, Knapp, in Flora, xlv. (1862) 273 = angustifolia.
parviflora, Michx. Fl. Bor. Am. i. 131 = Mertensia maritima.
parviflora, Schrank, in Nov. Act. Nat. Cur. ix. (1818) 103 = Mertensia paniculata.
parviflora, Schur, Enum. Pl. Transs. 472 = mollissima.
pauciflora, Gilib. Fl. Lituan. i. 23 = officinalis.
pilosa, Cham. in Linnaea, iv. (1829) 449 = Mertensia paniculata.
procumbens, Stokes, Bot. Mat. Med. i. 281 = Mertensia maritima.
pubescens, Willd. ex Roem. & Schult. Syst. iv. 744 = Mertensia paniculata.
pumila, Schrank, in Nov. Act. Nat. Cur. ix. (1818) 101 = Mertensia paniculata.
rubra, Schott, in Bot. Zeit. ix. (1851) 395 = mollis.
saccharata, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ.
sibirica, Linn. Sp. Pl. 135 = Mertensia sibirica.
simplicissima, Ledeb. in Mém. Acad. Pétersb. v. (1812) 518 = Mertensia simplicissima.
stiriaca, Kern. Monog. Pulmonar. 36 = angustifolia.
stylosa, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 62 = Mertensia stylosa.
suffruticosa, Linn. Sp. Pl. ed. II. 1667 = Lithospermum graminifolium.
tenella, Salisb. Prod. 113 = Mertensia pulmonarioides.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = mollis.
tuberosa, Martr. Fl. Tarn, i. 487, partim = *ovalis*.
tuberosa, Schrank, in Nov. Act. Nat. Cur. ix. (1818) 97 = angustifolia.
undulata, *Lavy, Vég. Orig.* 12 (sp. obsc.).—Hab. ?
Vallarsae, Kern. Monog. Pulmonar. 33. t. 11 = officinalis.
variabilis, Godr. Fl. Lorr. ii. 122 = angustifolia.
villosula, Roem. & Schult. Syst. iv. 745 = Mertensia villosula.
virginica, Linn. Sp. Pl. 135 = Mertensia pulmonarioides.
vulgaris, Holand, Fl. Moselle, i. 90 = officinalis.
vulgaris, Mérat, Fl. Par. ed. I. 70; II. ii. 164 (farrago spp.).—Europ.

PULSATILLA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 163 (1737) = **Anemone**, Tourn. (Ranunc.).
affinis, Lasch, in Linnaea, iii. (1828) 166 = A. pratensis.
africana, Spreng. Syst. ii. 664 = A. capensis.
ajanensis, Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 28 = A. vernalis.
alba, Reichb. Fl. Germ. Excurs. 732 = A. alpina.
albana, Bercht. & Presl, Rostl. i. Ranunculac. 22; Spreng. Syst. ii. 663 = A. albana.
alpina, G. Laws. in Proc. & Trans. Nova Scotian Inst. Nat. Sc. (1870) 23 = A. occidentalis.
alpina, Schrank, Baier. Fl. ii. 81 = A. alpina.
ambigua, Turcz. ex Pritz. in Linnaea, xv. (1841) 601 = A. albana.
amoena, Jord. Diagn. i. 54 = A. Pulsatilla.
angustifolia, Turcz. in Bull. Soc. Nat. Mosc. xiii. (1840) 61.—Sibir.
aperta, Schur, Enum. Pl. Transs. 4 = A. Pulsatilla.
apiifolia, Reichb. Ic. Fl. Germ. iv. t. 51 = A. alpina.
apiifolia, Sweet, Hort. Brit. ed. I. 3 = A. alpina.
baldensis, Delarb. Fl. Auv. ed. II. 553 = A. baldensis.
Bauhini, Tausch, ex Pritz. in Linnaea, xv. (1841) 580 = A. patens.
Bogenhardiana, Reichb. Ic. Fl. Germ. iv. 18 = A. Bogenhardiana.
Breynii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xii. (1854) 218.—Rossia.

PULSATILLA:—
Bungeana, C. A. Mey. ex Ledeb. Fl. Alt. ii. 371 = Anemone Bungeana.
Burseriana, Reichb. Fl. Germ. Excurs. 732 = A. alpina.
caffra, Eckl. & Zeyh. Enum. Pl. Afr. Austr. 1 = A. caffra.
campanella, Fisch. ex Regel, in Bull. Soc. Nat. Mosc. (1861) II. 31 = A. albana.
capensis, Steud. Nom. ed. II. i. 379, in syn. = A. capensis.
cernua, Bercht. & Presl, Rostl. i. Ranunculac. 22; Spreng. Syst. ii. 664 = A. cernua.
chinensis, C. Muell. in Walp. Ann. vii. 8 = A. chinensis.
coronaria, Borkh. in Rhein. Mag. i. (1793) 420 = A. coronaria.
davurica, Spreng. Syst. ii. 663 = A. dahurica.
georgica, Rupr. Fl. Cauc. 9, in textu = violacea?
grandiflora, Reichb. ex Walp. Rep. i. 19 = A. alpina.
grandis, Wender. in Schrift. Marb. ii. (1830) 257 = A. Pulsatilla.
Hackelii, Pohl, ex Maly, Fl. Styr. 2; Ledeb. Fl. Ross. i. 20 = A. Pulsatilla.
Hackelii, Reichb. Fl. Germ. Excurs. 733 = A. Halleri.
Halleri, Willd. Enum. Hort. Berol. 580 = A. Halleri.
hybrida, Mikan & Presl, in Sturm, Deutschl. Fl. Heft 46 = A. Halleri.
intermedia, Fisch. ex Schrank, in Syll. Ratisb. i. (1824) 221 (Quid?).—Sibir.
intermedia, Hoppe & Hornsch. Reise, 183 = A. montana.
intermedia, Lasch, in Linnaea, iii. (1828) 164 = patens.
intermedia, Sweet, Hort. Brit. ed. II. 3 = A. Pulsatilla.
Jankae, Nym. Syll. Suppl. 28 = A. pratensis.
Jankae, Schur, Enum. Pl. Transs. 4 = A. Jankae.
latifolia, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xii. (1854) 218.—Rossia.
lutea, Reichb. ex Walp. Rep. i. 19 = A. alpina.
media, Bogenh. in Flora, xxiii. (1840) 74 = A. Pulsatilla.
micrantha, Sweet, Hort. Brit. ed. II. 3 = A. alpina.
millefoliata, Ser. ex Nym. Consp. 2 = A. alpina.
montana, Reichb. Fl. Germ. Excurs. 733 = A. montana.
narcissiflora, Schrank, Baier. Fl. ii. 85 = A. narcissiflora.
nemorosa, Schrank, l. c. 82 = A. nemorosa.
Nigella, Jord. Diagn. i. 55 = A. montana.
nigricans, Stoerck, ex DC. Syst. i. 193 = A. pratensis.
nuda, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) II. 19.—Sibir.
Nuttalliana, Spreng. Syst. ii. 663 = A. patens.
obsoleta, Sweet, Hort. Brit. ed. II. 3 = A. pratensis.
patens, Mill, Gard. Dict. ed. VIII. n. 4 = A. patens.
pratensis, Mill. l. c. n. 2 = A. pratensis.
propera, Jord. Diagn. i. 54 = A. Pulsatilla.
propinqua, Lasch, in Linnaea, iii. (1828) 165 = A. vernalis.
Pseudo-patens, Schur, Enum. Pl. Transs. 4 = A. patens.
ranunculoides, Schrank, Baier. Fl. ii. 84 = A. ranunculoides.
recta, Gilib. Fl. Lituan. ii. 278 = A. Pulsatilla.
reflexa, Gilib. l. c. = A. pratensis.
rubra, Delarb. Fl. Auv. ed. II. 553 = A. pratensis.
slavica, Reuss, Květ. Slov. (1853) 5 = A. slavica.
sulphurea, Sweet, Hort. Brit. ed. II. 3 = A. vernalis.
tenuifolia, Spreng. Syst. ii. 664 = A. tenuifolia.
transsilvanica, Schur, Enum. Pl. Transs. 5 = A. Pulsatilla.
vernalis, Mill. Gard. Dict. ed. VIII. n. 3 = A. vernalis.
versicolor, Andrz. ex Rupr. in Bull. Phys.-Math. Acad. Pétersb. xii. (1854) 217 = A. stellata?
violacea, Rupr. Fl. Cauc. 6 = A. albana.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = A. Pulsatilla.
Wolfgangiana, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xii. (1854) 218 = A. Wolfgangiana.
Zichyi, Schur, in Oestr. Bot. Zeitschr. xiii. (1863) 317 = A. pratensis.

PULTENAEA, Sm. Bot. Nov. Holl. 35. t. 12 (1793). *LEGUMINOSAE*, Benth. & Hook. f. i. 470.
BARTLINGIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1848) 373.

PULTENAEA:—
EUCHILUS, R. Br. in Ait. Hort. Kew. ed. II. iii. 17 (1811).
SPADOSTYLES, Benth. in Ann. Wien. Mus. ii. (1838) 80.
URODON, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 17.
acerosa, *R. Br. ex Benth. Fl. Austral.* ii. 131.—Austral.
aciphylla, Benth. Enum. Pl. Hueg. 35; et in Ann. Wien. Mus. ii. (1838) 81 = reticulata.
adunca, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) I. 279.—Austral.
altissima, *F. Muell. ex Benth. Fl. Austral.* ii. 123.—Austral.
amoena, Sieber, ex Steud. Nom. ed. II. ii. 418 = linophylla.
argentea, A. Cunn. in Field, N. S. Wales, 347 = dentata.
aristata, *Sieber, ex DC. Prod.* ii. 112.—Austral.
aspalathoides, *Meissn. in Lehm. Pl. Preiss.* i. 73.—Austral.
aspera, Sieber, ex DC. Prod. ii. 113 = Phyllota phylicoides.
Ausfeldi, *Regel, Gartenfl.* (1865) 193. t. 475. f. 1–11.—Austral.
baeckeoides, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 83 = Gunnii.
Benthami, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 38.—Austral.
biloba, R. Br. in Bot. Mag. t. 2091 = scabra.
brachyphylla, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 279 = verruculosa.
brachytropis, *Benth. ex Lindl. in Bot. Reg.* (1841) *Misc.* 40.—Austral.
bracteata, Schrad. Sert. Hannov. t. 18 = linophylla.
Brunonis, Benth. in Ann. Wien. Mus. ii. (1838) 81 = Latrobea Brunonis.
calycina, *Benth. Fl. Austral.* ii. 121.—Austral.
canaliculata, *F. Muell. in Trans. Vict. Inst.* i. (1855) 119.—Austral.
candida, Lodd. Bot. Cab. t. 1236 = tenuifolia.
canescens, A. Cunn. in Field, N. S. Wales, 346 = plumosa.
caniculata, Walp. Ann. vii. 662, sphalm. = canaliculata.
capitellata, Sieber, ex DC. Prod. ii. 112 = stricta.
clutiaefolia, *Regel, Ind. Sem. Hort. Petrop.* (1860) 108.—Hab.?
comosa, Sieber, ex DC. Prod. ii. 113 = Phyllota phylicoides.
conferta, *Benth. Fl. Austral.* ii. 118.—Austral.
cordata, R. Grah. in Hook. Bot. Mag. t. 3443 = juniperina.
cordifolia, Benth. in Ann. Wien. Mus. ii. (1838) 82 = praec.
crassifolia, *Lodd. Cat. ex Loud. Hort. Brit.* 164, nomen.—Austral.
cuneata, Benth. in Ann. Wien. Mus. ii. (1838) 83 = microphylla.
daphnoides, *Wendl. Bot. Beobacht.* 49.—Austral.
deltoidea, Sieber, ex DC. Prod. ii. 110 = scabra.
densifolia, *F. Muell. in Trans. Vict. Inst.* i. (1855) 119.—Austral.
dentata, *Labill. Nov. Holl. Pl.* i. 103. t. 131.—Austral.
diemenica, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 277 = pedunculata.
diffusa, *Hook. Fl. Tasm.* i. 91. t. 14.—Tasman.
Drummondii, *Meissn. in Lehm. Pl. Preiss.* ii. 221.—Austral.
echinata, Spreng. Syst. iv. Cur. Post. 173, sphalm. = seq.
echinula, *Sieber, ex DC. Prod.* ii. 112.—Austral.
elliptica, *Sm. in Trans. Linn. Soc.* ix. (1808) 246.—Austral.
empetrifolia, *Meissn. in Lehm. Pl. Preiss.* i. 76.—Austral.
epacridea, F. Muell. Fragm. iv. 21 = staphylloides.
ericaefolia, *Benth. in Lindl. Swan River App.* 13.—Austral.
ericoides, Vent. Jard. Malm. t. 35 = Aotus villosa.
eriophora, *Lodd. ex Steud. Nom.* ed. II. ii. 418.—Austral.
euchila, *DC. Prod.* ii. 112.—Austral.
fasciculata, *Benth. in Ann. Wien. Mus.* ii. (1838) 82.—Austral.
ferruginea, Rudge, in Trans. Linn. Soc. xi. (1815) 300. t. 23 = villosa.

PULTENAEA:—
filifolia, F. Muell. Fragm. i. 9 = tenuifolia.
flexilis, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 502.—Austral.
foliolosa, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 83.—Austral.
fuscata, F. Muell. in Trans. Vict. Inst. i. (1855) 119 = vestita.
genistoides, Meissn. in Lehm. Pl. Preiss. i. 73 = Latrobea genistoides.
glabra, *Benth. Fl. Austral.* ii. 125.—Austral.
glaucescens, Sieber, ex DC. Prod. ii. 112 = linophylla.
graveolens, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1885) 68.—Austral.
Gunnii, *Benth. in Ann. Wien. Mus.* ii. (1838) 82.—Austral.
Hartmanni, *F. Muell. Fragm.* viii. 166.—Austral.
hibbertioides, *Hook. f. Fl. Tasman.* i. 89.—Austral.
hispidula, *R. Br. ex Benth. Fl. Austral.* ii. 133.—Austral.
humilis, *Benth. in Hook. f. Fl. Tasman.* i. 91.—Austral.
hypolampra, Sieber, ex DC. Prod. ii. 111 = elliptica.
ilicifolia, Andr. Bot. Rep. t. 320 = Oxylobium trilobatum.
incarnata, *Sweet, Hort. Brit.* ed. I. 108 (Quid?).—Austral.
incurvata, *A. Cunn. in Field, N. S. Wales*, 346.—Austral.
involucrata, *Benth. Fl. Austral.* ii. 138.—Austral.
juncea, Willd. Sp. Pl. ii. 506 = Viminaria denudata.
juniperina, *Labill. Nov. Holl. Pl.* i. 103. t. 130.—Austral.
lanata, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 83 = villosa.
largiflorens, *Benth. Fl. Austral.* ii. 134.—Austral.
laxa, *Kunze, Pugill. Pl. Nov.* 11.—Austral.
laxiflora, *Benth. Fl. Austral.* ii. 133.—Austral.
Lindleyana, *Regel, Ind. Sem. Hort. Petrop.* (1856) 39.—Austral.
linophylla, *Schrad. Sert. Hannov.* 28.—Austral.
microphylla, *Sieber, ex DC. Prod.* ii. 112.—Austral.
mollis, *Lindl. in Mitch. Three Exped.* ii. 260.—Austral.
montana, Lindl. l. c. 178 = scabra.
mucronata, Lodd. Bot. Cab. t. 1711 = polifolia.
mucronata, *F. Muell. Fragm.* i. 8.—Austral.
Muelleri, *Benth. Fl. Austral.* ii. 138.—Austral.
myrtoides, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 81.—Austral.
nana, Andr. Bot. Rep. t. 434 = Chorizema ilicifolium.
neurocalyx, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) I. 281.—Austral.
obcordata, Andr. Bot. Rep. t. 574 = daphnoides.
obcordata, *Benth. Fl. Austral.* ii. 120.—Austral.
obovata, *Benth. l. c.* 123.—Austral.
ochreata, *Meissn. in Lehm. Pl. Preiss.* i. 75.—Austral.
Ottonis, *Regel, Gartenfl.* (1857) 343.—Austral.
oxalidifolia, A. Cunn. ex Steud. Nom. ed. II. ii. 418 = ternata.
paleacea, *Willd. Sp. Pl.* ii. 506.—Austral.
parviflora, *Sieber, ex DC. Prod.* ii. 111.—Austral.
pedunculares, Benth. in Ann. Wien. Mus. ii. (1838) 83 = seq.
pedunculata, *Hook. Bot. Mag.* t. 2859.—Austral.
petiolaris, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 82.—Austral.
phylicoides, Sieber, ex DC. Prod. ii. 113 = Phyllota phylicoides.
pimeleoides, Hook. f. Fl. Tasman. i. 88 = dentata.
pinifolia, *Meissn. in Lehm. Pl. Preiss.* ii. 220.—Austral.
plumosa, *Sieber, ex DC. Prod.* ii. 111.—Austral.
polifolia, *A. Cunn. in Field, N. S. Wales*, 346.—Austral.
polygalifolia, Rudge, in Trans. Linn. Soc. xi. (1815) 303. t. 25 = villosa.
procumbens, *A. Cunn. in Field, N. S. Wales*, 347.—Austral.
prostrata, *Benth. in Hook. f. Fl. Tasman.* i. 89.—Austral.
protaeoides, Sieber, ex DC. Prod. ii. 112 = stipularis.
psoraleoides, Sieber, ex Steud. Nom. ed. II. ii. 418 = stipularis.
pteronioides, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 280 = strobilifera.
pycnocephala, *F. Muell. ex Benth. Fl. Austral.* ii. 114.—Austral.

PULTENAEA:—
racemulosa, Sieber, ex DC. Prod. ii. 111 = villosa ?
reticulata, *Benth. Fl. Austral.* ii. 118.—Austral.
retorta, Wendl. Hort. Herrenh. ii. 13. t. 9 = Dillwynia ericifolia.
retusa, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 502 = linophylla.
rigida, *R. Br. ex Benth. Fl. Austral.* ii. 130.—Austral.
rosea, *F. Muell. Fragm.* ii. 15.—Austral.
rosmarinacea, Sieber, ex Steud. Nom. ed. II. ii. 418 = Aotus villosa.
rosmarinifolia, Lindl. Bot. Reg. t. 1584 = polifolia.
rosmarinifolia, Sieber, ex DC. Prod. ii 108 = Aotus villosa.
rotundifolia, *Benth. Fl. Austral.* ii. 121.—Austral.
rubiaefolia, Andr. Bot. Rep. t. 351 = Mirbelia reticulata.
rupestris, *Baxt. in Loud. Hort. Brit. Suppl.* iii. 619, nomen (Quid ?).—Austral.
scabra, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 18.—Austral.
selaginoides, *Hook. f. Fl. Tasman.* i. 87.—Tasman.
setigera, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 82 = procumbens.
setulosa, *Benth. Fl. Austral.* ii. 132.—Austral.
Skinneri, *F. Muell. Fragm.* viii. 166.—Austral.
spinulosa, *Benth. Fl. Austral.* ii. 121.—Austral.
squarrosa, Sieber, ex DC. Prod. ii. 113 = Phyllota phylicoides.
staphyleoides, *A. Cunn. ex G. Don, Gen. Syst.* ii. 124.—Austral.
stenophylla, A. Cunn. ex G. Don, l. c. = microphylla.
stipularis, *Sm. Bot. Nov. Holl.* 35. t. 12; *et in Trans. Linn. Soc.* ix. (1808) 245.—Austral.
stricta, *Sims, Bot. Mag.* t. 1588.—Austral.
strobolifera, *Meissn. in Lehm. Pl. Preiss.* i. 76.—Austral.
styphelioides, Steud. Nom. ed. II. ii. 418, sphalm. = staphyleoides.
subspicata, *Benth. Fl. Austral.* ii. 137.—Austral.
subumbellata, *Hook. Bot. Mag.* t. 1584.—Austral.
Sweetii, G. Don, in Sweet, Hort. Brit. ed. III. 150 = flexilis.
sylvatica, Sieber, ex DC. Prod. ii. 104 = Oxylobium Pultenaeae.
tenella, *Benth. Fl. Austral.* ii. 122.—Austral.
tenuifolia, *R. Br. in Bot. Mag.* t. 2086.—Austral.
ternata, *F. Muell. Fragm.* i. 8.—Austral.
thymifolia, Sieber, ex DC. Prod. ii. 111. = elliptica.
tuberculata, Pers. Syn. i. 454 = elliptica.
uncinata, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 83 = microphylla.
Urodon, *Benth. Fl. Austral.* ii. 124.—Austral.
verruculosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 278.—Austral.
verticillata, Turcz. l. c. 279 = empetrifolia.
vestita, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 19.—Austral.
villifera, *Sieber, ex DC. Prod.* ii. 111.—Austral.
villosa, Andr. Bot. Rep. t. 309 = Aotus villosa.
villosa, *Willd. Sp. Pl.* ii. 507.—Austral.
virgata, Sieber, ex DC. Prod. ii. 108 = Aotus villosa.
viscidula, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1885) 69.—Austral.
viscosa, *R. Br. ex Benth. Fl. Austral.* ii. 127.—Austral.

PULTENEYA, Hoffmgg. Verz. Pfl. 191 (1824) = praec.

PULTNAEA, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1836) 195, sphalm. = praec.

PULVINARIA, Fourn. in Mart. Fl. Bras. vi. iv. 214 (1885). *ASCLEPIADEAE.*
Lhotzkyana, *Fourn. l. c.* 215.—Bras.

PUMILEA, P. Br. Hist. Jamaic. 188 (1756) = **Turnera**, Plum

PUMILO, Schlecht. in Linnaea, xxi. (1848) 448 = **Rutidosis**, DC. (Compos.).
argyrolepis, Schlecht. l. c. = R. Pumilo.
Preissii, Sond. in Linnaea, xxv. (1852) 487 = R. Pumilo.

PUNDUANA, Steetz, in Peters, Reise Mossamb. Bot. 345 (1863) = **Vernonia**, Schreb. (Compos.).
volkameriaefolia, Steetz, l. c. 346 = V. volkameriaefolia.
vulgaris, Steetz, l. c. = V. volkameriaefolia.

PUNEERIA, Stocks, in Journ. As. Soc. Bombay, iii. (1849) 55 = **Withania**, Pauq. (Solanac.).
coagulans, Stocks, l. c. 56 = W. coagulans.

PUNGAMIA, Lam. Illustr. t. 603 (1792 ?) = **Pongamia**, Vent. (Legumin.).

PUNICA, [Tourn.] Linn. Syst. ed. I (1735). *LYTHRARIEAE*, Benth. & Hook. f. i. 784.
GRANATUM, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132.
RHOEA, St. Lag. l. c. 133 (1880).
florida, Salisb. Prod. 354 = P. Granatum.
Granatum, *Linn. Sp. Pl.* 472.—Europ. austr.; Maurit.
grandiflora, Hort. ex Steud. Nom. ed. I. 669 = P. Granatum.
nana, Linn. Sp. Pl. ed. II. 676 = P. Granatum.
protopunica, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 512.—Ins. Socotra.
spinosa, Lam. Fl. Fr. iii. 483 = P. Granatum.

PUNICELLA, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 333 = **Balaustion**, Hook. (Myrtac.).
carinata, Turcz. l. c. = B. pulcherrimum.

PUPAL, Adans. Fam. ii. 268 (1763) = seq.

PUPALIA, Juss. in Ann. Mus. Par. ii. (1803) 132. *AMARANTACEAE*, Benth. & Hook. f. iii. 31.
CODIVALIA, Rafin. Fl. Tellur. iii. 40 (1836).
DESMOCHAETA, DC. Cat. Hort. Monsp. 101 (1813).
KOMMIA, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 181 (1867).
SYAMA, Jones, in As. Res. iv. (1795) 261.
Alopecurus, Fenzl, ex Moq. in DC. Prod. xiii. ii. 328 = Cyathula cylindrica.
atropurpurea, *Moq. l. c.* 331.—As. et Afr. trop.
densiflora, Mart. Nov. Gen. et Sp. ii. tt. 156, 158 = Cyathula achyranthoides.
distantiflora, *A. Rich. Tent. Fl. Abyss.* ii. 217.—Abyss.
echinata, Juss. ex Steud. Nom. ed. I. 269, 669 = lappacea.
Figarei, *Tod. Ind. Sem. Hort. Panorum.* (1866) 34.—Aegypt.
globosa, Hochst. & Steud. in Flora, xxiv. (1841) i. Intell. 22, nomen = Cyathula globulifera.
holosericea, Fenzl, ex Moq. in DC. Prod. xiii. ii. 329, 333 = Cyathula globulifera.
lappacea, *Juss. in Ann. Mus. Par.* ii. (1803) 132; *Moq. l. c.* 331.—As. et Afr. trop.
mollis, *Moq. l. c.* 333.—Afr. trop.
natalensis, Sond. in Linnaea, xxiii. (1850) 97 = Cyathula natalensis.
orbiculata, *Wight, Ic.* v. t. 1783.—Ind. or.
ortacantha, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 181.—Abyss.
patula, Juss. ex Steud. Nom. ed. I. 269, 669 = lappacea.
prostrata, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 321 = Cyathula geniculata.
remotiflora, *Moq. in DC. Prod.* xiii. ii. 333.—Afr. austr.
Schimperiana, Hochst. in Flora, xxiv. (1841) i. Intell. 22, nomen = Cyathula cylindrica.
sordida, *Moq. in DC. Prod.* xiii. ii. 333.—Reg. Argent.
styracifolia, Juss. ex Steud. Nom. ed. I. 269, 669 = lappacea.
subfusca, *Moq. in DC. Prod.* xiii. ii. 332.—Afr. trop.
Thonningii, *Moq. l. c.* 333.—Afr. trop.
velutina, Moq. l. c. 332 = lappacea.

PURARIA, Wall. Cat. n. 5352–55, 5570 (1831 ?) = **Pueraria**, DC. (Legumin.).

PURCHIA, Dum. Anal. Fam. 24 (1829) = PURSHIA, Spreng. = **Onosmodium**, Michx. (Borag.).

PURDIAEA, Planch. in Hook. Lond. Journ. Bot. v. (1846) 250. t. 9. *CYRILLEAE* [Benth. & Hook. f. ii. 1226].
COSTAEA, A. Rich. Fl. Cub. Fanerog. ii. 75 (1853).
angustifolia, Wright, ex Griseb. Cat. Pl. Cub. 53 = stenopetala.
monodynama, *Griseb. l. c.*—Cuba.
nutans, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 251. t. 9.—N. Granat.
stenopetala, *Griseb. in Goett. Abh.* ix. (1861) 45.—Cuba.

PURGOSEA, Haw. in Phil. Mag. iii. (1828) 184 = **Crassula**, Linn.
aloides, Sweet, Hort. Brit. ed. II. 223 = C. alooides.
alpestris, G. Don, Gen. Syst. iii. 104 = C. alpestris.
barbata, G. Don, l. c. = C. barbata.
capitellata, Sweet, Hort. Brit. ed. II. 223 = C. capitella.
cephalophora, G. Don, Gen. Syst. iii. 103 = C. cephalophora.
ciliata, Sweet, Hort. Brit. ed. II. 223 = C. ciliata.
concinna, Sweet, l. c. = C. concinna.
concinnella, Sweet, l. c. = C. concinnella.
conspicua, Sweet, l. c. = C. conspicua (in Addend.).
corymbosula, Loud. ex Steud. Nom. ed. II. ii. 421 = seq.
corymbulosa, Sweet, Hort. Brit. ed. II. 223 = C. corymbulosa.
Cotyledonis, Sweet, l. c. = C. Cotyledonis.
crenulata, G. Don, Gen. Syst. iii. 103 = C. crenulata.
debilis, G. Don, l. c. 104 = C. debilis.
dentata, G. Don, l. c. = C. dentata.
hemisphaerica, G. Don, l. c. = C. hemisphaerica.
hirta, G. Don, l. c. 103 = C. hirta.
ligulifolia, Sweet, Hort. Brit. ed. II. 223 = C. ligulifolia.
linguaefolia, Sweet, l. c. = C. linguaefolia.
minima, G. Don, Gen. Syst. iii. 105 = C. minima.
montana, G. Don, l. c. 104 = C. montana.
obovata, Sweet, Hort. Brit. ed. II. 223 = C. obovata.
pertusa, Sweet, l. c. = C. corymbulosa.
pertusula, Haw. in Phil. Mag. iii. (1828) 184.—Afr. austr.
punctata, Sweet, Hort. Brit. ed. II. 223 = C. punctata.
pyramidalis, G. Don, Gen. Syst. iii. 104 = C. pyramidalis.
ramuliflora, Sweet, Hort. Brit. ed. II. 223 = C. ramuliflora.
rupestris, G. Don, Gen. Syst. iii. 105 = C. rupestris.
sediflora, Steud. Nom. ed. II. ii. 421 = C. sediflora.
spicata, G. Don, Gen. Syst. iii. 105 = C. spicata.
tecta, G. Don, l. c. = C. tecta.
thyrsiflora, Sweet, Hort. Brit. ed. II. 223 = C. thyrsiflora.
tomentosa, Sweet, l. c. = C. tomentosa.
turrita, Sweet, l. c. = C. turrita.

PURGOSIA, G. Don, Gen. Syst. iii. 103 (1834) = praec.

PURKINJIA, Presl, Symb. Bot. ii. 17. t. 64 (1833) = **Ardisia**, Sw. (Myrsin.).
nodosa, Presl, l. c.—Mexic.

PURPURELLA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 301 = **Tibouchina**, Aubl. (Melastom.).
albiflora, Triana, in Trans. Linn. Soc. xxviii. (1871) 35 = T. hospita.
grossa, Triana, l. c. = T. grossa.
Itatiaiae, Wawra, in Oestr. Bot. Zeitschr. xxxii. (1882) 38 = T. Itatiaiae.
muricata, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 301 = T. grossa.
reticulata, Naud. l. c. 302 = T. reticulata.

PURSHIA, DC. in Trans. Linn. Soc. xii. (1817) 157. *ROSACEAE*, Benth. & Hook. f. i. 617.
KUNZIA, Spreng. Anleit. ii. ii. 869 (1818).
TIGAREA, Pursh, Fl.-Am. Sept. i. 333. t. 15 (1814).
glandulosa, *Curran, in Bull. Calif. Acad.* i. n. iii. (1885) 153.—Calif.
tridentata, *DC. in Trans. Linn. Soc.* xii. (1817) 158.—Am. bor. occ.

PURSHIA, Dennst. Schluess. Hort. Malab. 35 (1818) = **Centranthera**, R. Br. (Scrophul.).
ciliata, Dennst. l. c. = C. procumbens.

PURSHIA, Rafin. in Am. Monthly Mag. (1819) 191, sub n. 33 = BURSHIA, Rafin. = **Myriophyllum**, Ponted. (Halorag.).

PURSHIA, Spreng. Anleit. ii. 1. 450 (1817) = **Onosmodium**, Michx. (Borag.).
hispida, Lehm. Asperif. ii. 382 = O. hispidum.
mollis, Lehm. l. c. 383 = O. molle.

PURUMA, Jaume St. Hil. Expos. ii. 313 (1805) = **Pourouma**, Aubl. (Urtic.).

PUSCHKINIA, Adams, in Nov. Act. Acad. Petrop. xiv. (1805) 164. t. B. *LILIACEAE*, Benth. & Hook. f. iii. 813.
ADAMSIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 16.
dubia, Kunth, Enum. Pl. iv. 338 = Hyacinthus spicatus.
hyacinthoides, *Baker, in Linn. Soc. Journ.* xi. (1871) 435.—Kurdistan.
libanotica, Zucc. in Abh. Akad. Muench. iii. (1837–43) 238 = scilloides.
scilloides, *Adams, in Nov. Act. Acad. Petrop.* xiv. (1805) 164.—As. Min.; Syria.
scilloides, Sieber, Reise Kreta, ii. 319 = Chionodoxa nana.
sicula, Van Houtte, in Fl. des Serres, xxi. (1875) 117 = scilloides.

PUTORIA, Pers. Syn. i. 524 (1805). *RUBIACEAE*, Benth. & Hook. f. ii. 136.
balearica, Steud. Nom. ed. II. ii. 67, in syn. = calabrica.
brevifolia, *Coss. & Dur. in Ann. Sc. Nat.* Sér. IV. i. (1854) 224, nomen.—Alger.
calabrica, *Pers. Syn.* i. 524.—Reg. Mediterr.
cymosa, Pomel, Nouv. Mat. Fl. Atl. 80 = calabrica.
hispanica, Boiss. & Reut. Diagn. Ser. II. ii. 108 = calabrica.
indica, DC. Prod. iv. 577 = Anotis Leschenaultiana?
microphylla, Pomel, Nouv. Mat. Fl. Atl. 79 = brevifolia.
tenella, Pomel, l. c. 71 = brevifolia.

PUTRANJIVA, Wall. Tent. Fl. Nap. 61 (1826). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 277.
NAGEIA, Roxb. Hort. Beng. 71 (1814); Fl. Ind. iii. 766 (1832).
PALENGA, Thw. in Hook. Kew Journ. viii. (1856) 270. t. 7.
amblyocarpa, Muell. Arg. in DC. Prod. xv. II. 444 = Roxburghii.
Roxburghii, *Wall. Tent. Fl. Nap.* 61.—Ind. or.; Burma.
sphaerocarpa, Muell. Arg. in DC. Prod. xv. II. 443 = Roxburghii.
zeylanica, *Muell. Arg. l. c.* 444.—Zeylan.

PUTTERLICKIA, Endl. Gen. 1086 (1840). *CELASTRINEAE*, Benth. & Hook. f. i. 366.
pyracantha, *Endl. l. c.*—Afr. austr.

PUTZEYSIA, Klotzsch, in Abh. Akad. Berl. (1855) 134 = **Begonia**, Linn.
gemmipara, Klotzsch, l. c. 135 = B. gemmipara.

PUTZEYSIA, Planch. & Linden, in Linden, Cat. n. 22 (1857) 3 = **Aesculus**, Linn. (Sapind.).
rosea, Planch. & Linden, l. c.; et in Belg. Hortic. vii. (1857) 198 (= *Billia columbiana*).—N. Granat.

PUYA, Molina, Sagg. Chile, 160, 351 (1782). *BROMELIACEAE*, Benth. & Hook. f. iii. 666.
POURRETIA, Ruiz & Pav. Prod. 46. t. 7 (1794).
PUJA, Molina, Sagg. Chile, ed. II. 153 (1810).
alpestris, C. Gay, Fl. Chil. vi. 12 = Pitcairnia caerulea.
Altensteinii, Link, Klotzsch & Otto, Ic. Sel. i. 1. t. 1 = Pitcairnia Altensteinii.
Bonplandiana, *Schult. f. Syst.* vii. 1236.—N. Granat.
caerulea, Auct. = *coerulea*.
carnea, Regel, Cat. Pl. Hort. Aksakov. 117 = Pitcairnia carnea?
chilensis, *Molina, Sagg. Chile*, 160, 351.—Chili.
coarctata, Fisch. Sert. Petrop. t. 19 = chilensis.
coerulea, Miers, in Lindl. Bot. Reg. (1840) t. 11 = Pitcairnia caerulea.

PUYA:—
edulis, *E. Morr. in Belg. Hortic.* xxviii. (1878) 354.—Bras.
floccosa, *E. Morr. l. c.* xxxv. (1885) 81.—Venezuela.
gigantea, Phil. in Linnaea, xxxiii. (1864–65) 246 = chilensis.
Gigas, *E. Andr. in Rev. Hortic.* (1881) 314.—N. Granat.
grandiflora, Hook. Bot. Mag. t. 5234 = Pitcairnia ferruginea.
guyanensis, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1068, nomen.—Guian.
heterophylla, Lindl. Bot. Reg. (1840) t. 71 = Pitcairnia heterophylla.
lanata, *Schult. f. Syst.* vii. 1233.—Peruv.
lanuginosa, *Schult. f. l. c.* 1234.—Peruv.
longifolia, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) 483. t. 101 = Pitcairnia heterophylla.
macrostachya, *A. Dietr. Allg. Gartenz.* xvi. (1848) 141, 146.—Guian.
maidifolia, C. Morr. in Ann. Soc. Bot. Gand, v. (1849) 453. t. 289 = Pitcairnia maidifolia.
maydifolia, Decne. in Rev. Hortic. Sér. III. v. (1851) 347 = *maidifolia*.
meridensis, E. Morr. in Belg. Hortic. xxxv. (1885) 81 = floccosa.
paniculata, Phil. in Linnaea, xxxiii. (1864–65) 247 (Pitcairniae sp.).—Chili.
pyramidata, *Schult. f. Syst.* vii. 1235.—Peruv.
recurvata, Schiedw. in Otto & Dietr. Allg. Gartenz. x. (1842) 275 = Pitcairnia recurvata.
Roezli, E. Morr. in Belg. Hortic. xxxv. (1885) 80 (Pitcairniae sp.).—Peruv.
rubricaulis, Steud. Nom. ed. II. ii. 419 = Pitcairnia caerulea?
saxatilis, Mart. Reise Bras. ii. 757 (= *Encholirium spectabile*).—Bras.
suberosa, Molina, Sagg. Chile, ed. II. 153 = chilensis.
sulphurea, Hook. Bot. Mag. t. 4696 = Pitcairnia Wendlandi.
virescens, Hook. l. c. t. 4991 (Caraguatae sp.).—Peruv.
Warszewiczii, H. Wendl. l. c. t. 5225 = Pitcairnia atrorubens.
Whytei, Hook. f. l. c. t. 5732 = Pitcairnia caerulea.

PYCHNANTHEMUM, G. Don, Gen. Syst. iv. 667 (1837), sphalm. = **Pycnanthemum**, Michx. (Labiat.).

PYCHNOSTACHYS, G. Don, Gen. Syst. iv. 686 (1837), sphalm. = **Pycnostachys**, Hook. (Labiat.).

PYCNANDRA, Benth. in Benth. & Hook. f. Gen. ii. 658 (1876). *SAPOTACEAE*, Benth. & Hook. f. l. c.

PYCNANTHEMUM, Michx. Fl. Bor. Am. ii. 7. t. 33 (1803). *LABIATAE*, Benth. & Hook. f. ii. 1184.
BRACHYSTEMUM, Michx. Fl. Bor. Am. ii. 5. tt. 31, 32 (1803).
KOELLIA, Moench, Meth. 407 (1794).
TULLIA, Leavenw. in Am. Journ. Sc. Ser. I. xx. (1830) 343. t. 5.
alatum, Rafin. Fl. Ludov. 40 (Quid?).—Am. bor.
albescens, *Torr. & Gray, ex A. Gray, in Am. Journ. Sc.* Ser. I. xlii. (1842) 45.—Am. bor.
aristatum, *Michx. Fl. Bor. Am.* ii. 8. t. 33.—Am. bor.
arkansanum, Fresen. in Flora, xxv. (1842) 325 = muticum.
californicum, *Torr. ex Dur. in Journ. Acad. Philad.* Ser. II. iii. (1855) 99.—Calif.
clinopodioides, *Torr. & Gray, ex A. Gray, in Am. Journ. Sc.* Ser. I. xlii. (1842) 45.—Am. bor.
dubium, A. Gray, l. c. = P. Tullia.
elongatum, Blanco, Fl. Filip. ed. II. 333 = Hyptis spicigera.
hyssopifolium, Benth. Lab. Gen. et Sp. 329 = aristatum.
incanum, *Michx. Fl. Bor. Am.* ii. 7.—Am. bor.
lanceolatum, *Pursh, Fl. Am. Sept.* ii. 410.—Am. bor.
leptodon, *A. Gray, Syn. Fl. N. Am.* ii. 1. 355.—Am. bor.
linifolium, *Pursh, Fl. Am. Sept.* ii. 409.—Am. bor.
Loomisii, Nutt. in Journ. Acad. Philad. vii. (1834) 100 = incanum.
Monardella, Michx. Fl. Bor. Am. ii. 8. t. 34 = Monarda clinopodia.
montana, *Michx. l. c.* 8.—Am. bor.
muticum, *Pers. Syn.* ii. 128.—Am. bor.
nudum, *Nutt. Gen. Am.* ii. 34.—Am. bor.
ovatum, Nutt. ex Benth. in DC. Prod. xii. 188 = muticum.

PYCNANTHEMUM:—
pilosum, Nutt. Gen. Am. ii. 33 = muticum.
rosmarinoides, Rafin. Fl. Ludov. 40 (Quid?).—Am. bor.
setosum, Nutt. in Journ. Acad. Philad. vii. (1834) 100 = aristatum.
subulatum, Blanco, Fl. Filip. ed. II. 333 = Hyptis brevipes.
tenuifolium, Schrad. Hort. Gotting. 10 = linifolium.
Torrei, *Benth. Lab. Gen. et Sp.* 329.—Am. bor.
Tullia, *Benth. l. c.* 328.—Am. bor.
verticillatum, Pers. Syn. ii. 128 = lanceolatum.
verticillatum, Pursh, Fl. Am. Sept. ii. 410 = aristatum.
virginicum, Nutt. Gen. Am. ii. 33 = Torrei.
virginicum, Pers. Syn. ii. 128 = lanceolatum.

PYCNANTHES, Rafin. New Fl. Am. iv. 95 (1836) = **Pycnanthemum**? Michx. (Labiat.).
ciliata, Rafin. l. c.—Am. bor.
parvifolia, Rafin. l. c.—Am. bor.
saxatilis, Rafin. l. c.—Am. bor.

PYCNARRHENA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44. *MENISPERMACEAE*, Benth. & Hook. f. i. 38.
longifolia, *Becc. Malesia*, i. (1877) 160.—Ins. Timor.
lucida, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 87.—Java.
mecistophylla, *Miers, in Ann. Nat. Hist.* Ser. III. xx. (1867) 12; *Contrib.* iii. 353 (an gen.?).—Reg. Himal.
novoguineensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 87.—N. Guin.
planiflora, Hook. f. & Thoms. Fl. Ind. 206, sphalm. = pleniflora.
planifolius, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44, sphalm. = seq.
pleniflora, *Miers, l. c.* Ser. III. xx. (1867) 12; *Contrib.* iii. 353. t. 141.—Reg. Himal.
tumefacta, *Miers, ll. cc.* 12; 353.—Borneo.

PYCNOBOTRYA, Benth. in Benth. & Hook. f. Gen. ii. 715 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
nitida, *Benth. in Hook. Ic. Pl.* t. 1183.—Afr. trop.

PYCNOCEPHALUM, DC. Prod. v. 83 (1836) = **Eremanthus**, Less. (Compos.).
plantaginifolium, DC. l. c. = E. plantaginifolius.
scapigerum, DC. l. c. = E. scapigerus.
spathulaefolium, DC. l. c. = E. plantaginifolius.

PYCNOCOMA, Benth. in Hook. Niger Fl. 508 (1849). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 326.
cornuta, *Muell. Arg. in Flora*, xlvii. (1864) 483.—Afr. trop.
gigantea, *Baill. Etud. Gén. Euph.* 411.—Ins. Comor.
macrophylla, *Benth. in Hook. Niger Fl.* 508.—Afr. trop.
minor, *Muell. Arg, in Flora*, xlvii. (1864) 483.—Afr. trop.
reticulata, *Baill. Adansonia*, i. (1860–61) 259.—Madag.
rigidifolia, *Baill. Etud. Gén. Euph.* 411.—Madag.
trewioides, *Baill. l. c.*—Ins. Comor.

PYCNOCOMON, Hoffmgg. & Link, Fl. Port. ii. 93. t. 88 (1820) = **Scabiosa**, Tourn.; et **Cephalaria**, Schrad. (Dipsac.).
albescens, Wallr. Sched. Crit. 46, in nota = C. leucantha.
alpina, Wallr. l. c. = C. alpina.
attenuata, Wallr. l. c. = C. attenuata.
boetica, Boiss. Voy. Bot. Espagne, ii. 788 = C. leucantha.
centauroides, Wallr. Sched. Crit. 46, in nota = C. corniculata.
corniculata, Wallr. l. c. = C. corniculata.
cretacea, Wallr. l. c. = C. cretacea.
leucantha, Wallr. l. c. = C. leucantha.
rigida, Wallr. l. c. = C. rigida.
rutaefolia, Hoffmgg. & Link, Fl. Port. ii. 94. t. 88 = Scabiosa rutaefolia.
scabra, Wallr. Sched. Crit. 46, in nota = C. scabra.
syriaca, Wallr. l. c. = C. syriaca.
tatarica, Wallr. l. c. = C. tatarica.
transsylvanica, Wallr. l. c. = C. transsylvanica.
uralensis, Wallr. l. c. = C. corniculata.
ustulata, Wallr. l. c. = C. ustulata.

PYCNOCOMON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 132 = PICNOMON, Adans. = **Cnicus**, Linn. (Compos.).

PYCNOCOMUS, Hill, Veg. Syst. iv. 38 (1762) = **Centaurea**, Linn. (Compos.).
sonchifolius, Hill, Hort. Kew. 67 = C. sonchifolia.

PYCNOCYCLA, Lindl. in Royle, Illustr. Bot. Himal. 232. t. 51 (1834). *UMBELLIFERAE*, Benth. & Hook. f. i. 881.
DICYCLOPHORA, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 89.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 333 = glauca.
Aucheriana, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 88.—Persia; Beluchist.
cespitosa, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 953.—Persia.
flabellifolia, *Boiss. Diagn.* Ser. II. ii. 105.—Persia.
glauca, *Lindl. in Royle, Illustr. Bot. Himal.* 232. t. 51.—Ind. or.; Abyss.
macrocarpa, Decne. ex Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 75 = Cachrys eriantha.
nodiflora, *Decne. ex Boiss. l. c.* 88.—Persia.
spinosa, *Decne. l. c.* Sér. II. iii. (1835) 259.—Persia; Afghan.
tomentosa, *Decne. l. c.* 258.—Arab.

PYCNOLACHNE, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 215 = **Lachnostachys**, Hook. (Verben.).
ledifolia, Turcz. l. c. = L. Walcottii.

PYCNONEURUM, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 340. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 761.
junciforme, *Decne. l. c.*—Madag.
sessiliflorum, *Decne. l. c.* 341 (sessiflorum).—Madag.

PYCNOPHYLLUM, Remy, in Ann. Sc. Nat. Sér. III. vi. (1846) 355. t. 20. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
STICHOPHYLLUM, Phil. Fl. Atac. 19. t. 1 D (1860).
bryoides, *Rohrb. in Linnaea*, xxxvi. (1869–70) 662.—Chili.
convexum, *Griseb. in Goett. Abh.* xix. (1874) 75.—Reg. Argent.
Lechlerianum, *Rohrb. in Linnaea*, xxxvi. (1869–70) 664.—Peruv.
molle, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 355. t. 20. f. 2–8.—N. Granat.
sulcatum, *Griseb. in Goett. Abh.* xix. (1874) 75.—Am. austr.
tetrastichum, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 356. t. 20. f. 1.—Bolivia.

PYCNORHACHIS, Benth. in Benth. & Hook. f. Gen. ii. 776 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
Maingayi, *Hook. f. Fl. Brit. Ind.* iv. 63.—Malacca.

PYCNOSANDRA, Blume, Mus. Bot. Lugd. Bat. ii. 191 (1856) = **Cyclostemon**, Blume (Euphorb.).
serrata, Blume, l. c. 192 = C. serratus.
timorensis, Blume, l. c. = C. Teysmanni.

PYCNOSORUS, Benth. in Enum. Pl. Hueg. 63 (1837) = **Craspedia**, Forst. (Compos.).
chrysanthus, Sond. in Linnaea, xxv. (1852) 492 = C. chrysantha.
globosus, Benth. in Enum. Pl. Hueg. 63 = C. globosa.
globosus, Lindl. in Mitch. Journ. Trop. Austral. 353 = C. chrysantha.

PYCNOSPORA, R. Br. ex Wight & Arn. Prod. 197 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 521.
hedysaroides, *R. Br. l. c.*—Austral.
nervosa, Wight & Arn. l. c. = hedysaroides.

PYCNOSTACHYS, Hook. Exot. Fl. iii. t. 202 (1827). *LABIATAE*, Benth. & Hook. f. ii. 1177.
ECHINOSTACHYS, E. Mey. Comm. Pl. Afr. Austr. 243 (1837).
abyssinica, *Fresen. in Flora*, ii. (1838) 608.—Abyss.
coerulea, *Hook. Exot. Fl.* iii. t. 202.—Madag.
reticulata, *Benth. in DC. Prod.* xii. 83.—Afr. austr.
urticifolia, *Hook. Bot. Mag.* t. 5365.—Afr. trop.

PYCNOSTELMA, Bunge, ex Decne. in DC. Prod. viii. 512 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 759.
chinensis, *Bunge, ex Decne. l. c.*—China; Japon.

PYCREUS, Beauv. Fl. Owar. ii. 48. t. 86 (1807). *CYPERACEAE* [Benth. & Hook. f. iii. 1044].
DISTIMUS, Rafin. in Journ. Phys. lxxxix. (1819) 105.
albo-marginatus, *Nees, in Mart. Fl. Bras.* ii. 1. 9.—Amphig. trop.
angulatus, *Nees, in Linnaea*, ix. (1834) 283.—Amphig. calid.
Baccha, *Nees, l. c.*—Ind. or.
Bachu, Nees, ex Boeck. in Linnaea, xxxv. (1867–68) 473, in syn. = praec.
capillaris, Nees, l. c. = globosus.
densiflorus, Steud. Nom. ed. II. ii. 419, sphalm. = seq.
densifolius, *Nees, in Linnaea*, ix. (1834) 283, nomen. —Hab. ?
Elliottianus, Nees, l. c. = polystachyos.
flavescens, *Beauv. ex Reichb. Fl. Germ. Excurs.* 72.—Cosmop.
globosus, *Reichb. l. c.* 140[10].—Cosmop.
infirmus, Nees, in Mart. Fl. Bras. ii. 1. 8 = propinquus.
inundatus, Nees, l. c. 10, in obs. = P. Baccha.
laevigatus, Nees, in Linnaea, x. (1836) 130 = Cyperus laevigatus.
lateralis, Nees, l. c. ix. (1834) 283 = Cyperus laevigatus.
Linnaeanus, *Nees, l. c.*, nomen.—Hab. ?
Maximiliani, Nees, in Mart. Fl. Bras. ii. 1. 7 = propinquus.
megapotamicus, *Nees, l. c.* 6.—Bras.
Monti, Beauv. ex Reichb. Fl. Germ. Excurs. 72 = Cyperus Monti.
mucronatus, Nees, in Linnaea, ix. (1834) 283 = Cyperus laevigatus.
Mundtii, *Nees, l. c.*—Afr. trop.; Madag.
nitens, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 53.—Geront. trop.
Olfersianus, Nees, in Mart. Fl. Bras. ii. 1. 8 = polystachyos.
paniculatus, Nees, in Linnaea, ix. (1834) 283 = polystachyos.
pannonicus, Beauv. ex Reichb. Fl. Germ. Excurs. 72 = Cyperus pannonicus.
paraënsis, Nees, in Mart. Fl. Bras. ii. 1. 8 = flavescens.
polystachyos, *Beauv. Fl. Owar.* ii. 48. t. 86.—Cosmop.
propinquus, *Nees, in Mart. Fl. Bras.* ii. 1. 7.—Am. trop.
pulvinatus, Nees, ex Steud. Nom. ed. II. i. 472; ii. 419 = nitens.
pumilus, *Nees, in Linnaea*, ix. (1834) 283.—Ind. or.
puncticulatus, Nees, in Mart. Fl. Bras. ii. 1. 10, in obs. = P. Baccha.
pygmaeus, Nees, in Linnaea, ix. (1834) 283 = Cyperus pygmaeus.
retusus, *Nees, ex C. B. Clarke, in Journ. Linn. Soc.* xxi. (1884) 71.—Ins. Maurit.
sabulosus, *Nees, in Mart. Fl. Bras.* ii. 1. 10.—Bras.
sanguinolentus, *Nees, in Linnaea*, ix. (1834) 283.—Geront. trop.
squarrosus, Nees, l. c. = Cyperus squarrosus.
umbrosus, *Nees, l. c.* x. (1835–36) 130.—Afr. austr.

PYGEUM, Gaertn. Fruct. i. 218. t. 46 (1788). *ROSACEAE*, Benth. & Hook. f. i. 610.
DIGASTER, Miq. Fl. Ind. Bat. Suppl. 329 (1860).
GERMARIA, Presl, Epim. Bot. 221 (1850).
POLYDONTIA, Blume, Bijdr. 1104 (1826).
POLYSTORTHIA, Blume, Fl. Jav. Praef. viii (1828).
acuminatum, *Colebr. in Trans. Linn. Soc.* xii. (1819) 360. t. 18.—Reg. Himal.
acuminatum, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. ii. 319 = glaberrimum.
acuminatum, Wight, Ic. t. 993 = Gardneri.
africanum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 191.—Afr. trop.
Andersoni, *Hook. f. Fl. Brit. Ind.* ii. 320.—Ind. or.
arboreum, *Endl. ex Kurz, For. Fl. Brit. Burma*, i. 435.—Burma.
arboreum, Endl. Gen. 1250 = parviflorum.
Blumei, Teijsm. & Binn. Cat. Hort. Bog. 252 = parviflorum.
brevifolium, *Hook. f. Fl. Brit. Ind.* ii. 321.—Malacca.
capitellatum, *Hook. f. l. c.*—Burma.

PYGEUM:—
celebicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 212.—Ins. Celebes.
ceylanicum, Auct. = zeylanicum.
ceylanicum, Bedd. Fl. Sylv. t. 59 = Wightianum.
confusum, *Blume, Mél. Bot.* (1855); *et in Flora*, xli. (1858) 256.—Java.
denudatum, Buch.-Ham. ex Wall. Cat. n. 7190 = acuminatum.
Gardneri, *Hook. f. Fl. Brit. Ind.* ii. 321.—Ind. or.
glaberrimum, *Hook. f. l. c.* 319.—Reg. Himal.
Griffithii, *Hook. f. l. c.* 322.—Malacca.
griseum, *Blume, Mél. Bot.* (1855); *et in Flora*, xli. (1858) 256.—Java.
lampongum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 212.—Sumatra.
lanceolatum, *Hook. f. Fl. Brit. Ind.* ii. 319.—Singapore.
latifolium, *Miq. Fl. Ind. Bat.* i. 1. 361.—Sumatra.
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 319.—Malacca.
montanum, *Hook. f. l. c.* 321.—Reg. Himal.
ovatum, *Teijsm. & Binn. Cat. Hort. Bog.* 252.—Java.
oxycarpum, Hance, in Journ. Bot. viii. (1870) 242 = Prunus macrophylla.
parviflorum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 412.—Java.
parvifolium, Gardn. ex Thw. Enum. Pl. Zeyl. 103 = Wightianum.
persimile, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 306.—Burma.
phaeostictum, Hance, in Journ. Bot. viii. (1870) 72 = Prunus punctata.
polystachyum, *Hook. f. Fl. Brit. Ind.* ii. 320.—Malacca.
sumatranum, *Miq. Fl. Ind. Bat. Suppl.* 619.—Sumatra.
Walkerii, Blume, Mél. Bot. (1855); et in Flora, xli. (1858) 256 = zeylanicum.
Wightianum, *Blume, ll. cc.*—Ind. or.
zeylanicum, Dalz. & Gibs. Bomb. Fl. 89 = Gardneri.
zeylanicum, *Gaertn. Fruct.* i. 218. t. 46.—Zeylan.

PYGMAEA, Hook. f. Handb. N. Zeal. Fl. 217 (1864) = **Veronica**, Tourn. (Scrophul.).
ciliolata, Hook. f. l. c.—N. Zel.
pulvinaris, Hook. f. l. c.—N. Zel.
Thomsoni, J. Buch. in Trans. N. Z. Inst. xiv. (1882) 353.—N. Zel.

PYGMEA, J. Buch. in Trans. N. Z. Inst. xiv. (1882) 353, err. typ. = praec.

PYRACANTHA, M. Roem. Syn. fasc. iii. (Rosifl.) 104, 219 (1847) = **Crataegus**, Tourn. (Rosac.).
chinensis, M. Roem. l. c. 220 = C. Pyracantha.
coccinea, M. Roem. l. c. 219 = C. Pyracantha.
crenulata, M. Roem. l. c. 220 = C. crenulata.
pauciflora, M. Roem. l. c. 220 = C. Pyracantha.

PYRAGMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Uvaria**, Linn. (Anonac.).

PYRAMIA, Cham. in Linnaea, ix. (1834) 458. *MELASTOMACEAE*, Benth. & Hook. f. i. 737.
ACIPETALUM, Turcz. in Bull. Soc. Nat. Mosc. (1848) I. 577, partim.
Lychnitis, Klotzsch, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 29 = Trembleya Lychnitis.
pityrophylla, *Cham. in Linnaea*, ix. (1834) 459.—Bras.
salviaefolia, *Cham. l. c.*—Bras.
striatella, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 60.—Bras.

PYRAMIDANTHE, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 39 (1865–66) = **Melodorum**, Hook. f. & Thoms. (Anonac.).
rufa, Miq. l. c. = M. prismaticum.

PYRAMIDIUM, Boiss. Diagn. Ser. II. i. 47 (1853). *CRUCIFERAE*, Benth. & Hook. f. i. 97.
Griffithianum, *Boiss. l. c.*—Afghan.

PYRAMIDOCARPUS, Oliver, in Journ. Linn. Soc. ix. (1867) 171. *SAMYDACEAE*, Benth. & Hook. f. i. 798.
Blackii, *Oliver, l. c.*—Afr. trop.

PYRAMIDOPTERA, Boiss. Diagn. Ser. II. ii. 106 (1856). *UMBELLIFERAE*, Benth. & Hook. f. i. 903.
cabulica, *Boiss. l. c.*—Afghan.

PYRAMIDOSTYLIUM, Mart. ex Peyr. in Mart. Fl. Bras. xi. i. 155 (1878) = **Salacia**, Linn. (Celastr.).
Bacupari, Mart. ex Peyr. l. c. 160 = S. crassifolia.
banisterioides, Mart. ex Peyr. l. c. 155 = S. paniculata.
ciliatum, Mart. ex Peyr. l. c. 156 = S. undulata.
fimbriatum, Mart. ex Peyr. l. c. = S. undulata.
grandifolium, Mart. ex Peyr. l. c. = S. grandifolia.
parviflorum, Mart. ex Peyr. l. c. 160 = S. glomerata.

PYRARDA, Cass. in Dict. Sc. Nat. xli. 120 (1836) = **Grangea**, Adans. (Compos.).
ceruanoides, Cass. l. c. 121 = G. maderaspatana.

PYRECNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 (Quid ?).

PYRENACANTHA, Wight, in Hook. Bot. Misc. ii. (1831) 107. *OLACINEAE*, Benth. & Hook. f. i. 345 ; iii. 220.
ADELANTHUS, Endl. Gen. 1327. n. 6839 ; Suppl. ii. 80 (1841).
CAVANILLA, Thunb. Nov. Gen. 105 (1792).
MOLDENHAUERA, Spreng. Syst. i. 373 (1825).
SADRUM, Soland. ex Baill. in DC. Prod. xvii. 18 (1873).
chlorantha, *Baker, in Journ. Linn. Soc.* xxi. (1884) 331.—Madag.
grandiflora, *Baill. Adansonia*, x. (1872) 270.—Afr. austr.
kamassana, *Baill. l. c.* 272.—Afr. trop.
Kirkii, *Baill. l. c.*—Afr. trop.
scandens, *Planch. ex Harv. Thes. Cap.* i. 14. t. 23.—Afr. austr.
Vogeliana, *Baill. Adansonia*, x. (1872) 271.—Afr. trop.
volubilis, *Wight, in Hook. Bot. Misc.* ii. (1831) 107.—Ind. or.

PYRENARIA, Blume, Bijdr. 1119 (1826). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 185.
EUSYNAXIS, Griff. Notul. iv. 560 (1854).
acuminata, *Planch. ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 174.—Malaya.
attenuata, *Seem. in Bonplandia*, vii. (1859) 49.—Burma.
barringtoniaefolia, *Seem. l. c.*—Ind. or.
camelliaeflora, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 46.—Burma.
diospyricarpa, *Kurz, l. c.* xlii. (1873) II. 60.—Burma.
lanceolata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 40.—Java.
lasiocarpa, *Korth. Verh. Nat. Gesch. Bot.* 147.—Java.
masocarpa, *Korth. l. c.*—Borneo.
oidocarpa, *Korth. l. c.*—Java.
serrata, *Blume, Bijdr.* 1120.—Java.
villosula, *Miq. Fl. Ind. Bat. Suppl.* 484.—Sumatra.

PYRENIA, [Clairv.] Man. Herb. 161 (1811) = **Pyrus**, Tourn. (Rosac.).
Amelanchier, [Clairv.] l. c. 162 = Amelanchier vulgaris.
Aria, [Clairv.]. l. c. = Pyrus Aria.
Aucuparia, [Clairv.] l. c. = Pyrus Aucuparia.
Chamaemespylus, [Clairv.] l. c. = Pyrus Chamaemespilus.
Malus, [Clairv.] l. c. = Pyrus Malus.
Pyrus, [Clairv.] l. c. 161 = Pyrus communis.
Sorbus, [Clairv.] l. c. 162 = Pyrus Sorbus.
torminalis, [Clairv.] l. c. = Pyrus torminalis.

PYRENOGLYPHIS, Karst. Fl. Columb. ii. 141. t. 174 (1869) = **Bactris**, Jacq. (Palm.).
balanoidea, Karst. l. c. 142, in obs. = B. balanoides.
Cruegeriana, Karst. l. c., in obs. = B. Cruegeriana.
major, Karst. l. c. 141. t. 174 = B. major.
ovata, Karst. l. c. 142, in obs. = B. ovata.

PYRETHRARIA, Pers. ex Steud. Nom. ed. II. ii. 420 (1841) = **Cotula**, Tourn. (Compos.).
dichotoma, Pers. ex Steud. l. c. = C. Pyrethraria.

PYRETHRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.

PYRETHRUM, Hall. Enum. Stirp. Helv. ii. 720 (1742) = **Chrysanthemum**, Tourn. (Compos.).
abrotanifolium, Bunge, ex Ledeb. Fl. Ross. ii. 549.—Sibir. altaic.
abrotanifolium, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 86 = Anthemis montana.
Achilleae, DC. Prod. vi. 57 = C. Achilleae.
Achilleae, Wendl. in Flora, xii. (1829) I. Erg. 29 = C. macrophyllum.
achilleifolium, Bieb. Fl. Taur. Cauc. ii. 327 = C. achilleaefolium.
adauctum, Link, in Buch, Beschr. Canar. Ins. 149 = C. pinnatifidum.
alpestre, [Clairv.] Man. Herb. 247 = C. coronopifolium.
alpinum, Baumg. Enum. Stirp. Transs. iii. 108 = Anthemis carpathica.
alpinum, Schrank, Prim. Fl. Salisb. 215 = C. alpinum.
ambiguum, Ledeb. Fl. Alt. iv. 118 = C. montanum.
anethifolium, Willd. Enum. Hort. Berol. 904 = C. anethifolium.
anomalum, Lange, in Kjoeb. Vidensk. Meddel. (1861) 75 = C. oppositifolium.
argenteum, Adam, ex Ledeb. Fl. Ross. ii. 552 = C. praealtum.
argenteum, Boiss. Fl. Orient. iii. 354 = C. argenteum.
argyranthemoides, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 343 = Tanacetum argyranthemoides.
armenum, C. Koch, in Linnaea, xvii. (1843) 47 = C. millefoliatum.
arundanum, Boiss. Voy. Bot. Espagne, ii. 317.—Hispan.
arvense, Salzm. ex DC. Prod. vi. 54 = C. tingitanum ; Matricaria glabra.
Athanasia, Boiss. Fl. Orient. iii. 353 = Tanacetum fruticulosum.
atratum, Less. Syn. Comp. 254 = C. atratum.
Aucheri, Heimerl, ex Stapf, in Denkschr. Akad. Wien, l. (1885) 112.—Caria.
Aucherianum, DC. Prod. vii. 297 = Tanacetum Aucherianum.
auriculatum, Boiss. ex Boiss. Fl. Orient. iii. 327 = C. auriculatum.
balsaminatum, M. Ward, in Journ. Linn. Soc. xvii. (1880) 532, sphalm. = seq.
Balsamita, Willd. Sp. Pl. iii. 2153 = C. Balsamita.
Barrelieri, Dufour, ex DC. Prod. vi. 55 = C. coronopifolium.
Baumanni, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 283.—Tauria.
bipinnatifidum, Link, in Buch, Beschr. Canar. Ins. 149 = C. anethifolium.
bipinnatum, Willd. Sp. Pl. iii. 2160 = C. bipinnatum.
Bocconi, Willd. l. c. 2159 = C. Bocconi.
Bonplandianum, H. B. & K. Nov. Gen. et Sp. iv. 300 = Xanthocephalum centauroides.
Bourgaei, Boiss. Fl. Orient. iii. 356.—Armenia.
breviradiatum, Ledeb. in Mém. Acad. Pétersb. v. (1812) 577 = C. coronarium.
breviradiatum, Schweinitz, ex Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 397 = Matricaria discoidea.
Broussonetii, Choisy, in Buch, Beschr. Canar. Ins. 149 = C. Broussonetii.
cadmeum, Boiss. Diagn. Ser. I. iv. 10.—Mesopot.
canescens, Boiss. Fl. Orient. iii. 353 = Tanacetum canescens.
cappadocicum, DC. Prod. vii. 297 = Tanacetum cappadocicum.
carneum, Bieb. Fl. Taur. Cauc. ii. 325 = C. coronopifolium, *Willd.*
carniolicum, Dollin. ex Boiss. Fl. Orient. iii. 342 = C. corymbosum.
cassium, Boiss. Diagn. Ser. I. xi. 23.—Syria.
Catananche, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61 = C. Catananche.
caucasicum, Willd. Sp. Pl. iii. 2156 = C. caucasicum.
cedrorum, Schott & Kotschy, in Boiss. Fl. Orient. iii. 347 = *cappadocicum*.
ceratophylloides, Tenore, Fl. Nap. ii. 233 = C. coronopifolium.
chiliophyllum, Boiss. Diagn. Ser. I. xi. 26, in syn.—Persia.
chiliophyllum, Fisch. & Mey. ex DC. Prod. vi. 59 = C. millefoliatum.
chrysanthemifolium, Hort. Angl. ex DC. Prod. vi. 7 = Anthemis parthenioides.
cilicium, Boiss. Diagn. Ser. II. iii. 28.—Syria.

PYRETHRUM :—
cinerarifolium, Trev. Ind. Sem. Hort. Vratisl. App. ii. (1820) 2 ; et in Nov. Act. Nat. Cur. xiii. (1826) 204 = Chrysanthemum cinerariaefolium.
cinereum, Griseb. Spicil. Fl. Rumel. ii. 202.—Balkan.
Clausonis, Battand. in Bull. Soc. Bot. Fr. xxxii. (1885) 340 = C. paludosum.
Clusii, Fisch. ex Reichb. Fl. Germ. Excurs. 232 ; et ex Koch, Syn. Fl. Germ. ed. I. 379 = C. corymbosum.
conicum, Less. in Linnaea, ix. (1834) 189.—Rossia.
coronopifolium, Willd. Enum. Hort. Berol. 904 = C. grandiflorum.
corymbiferum, S. G. Gmel. Reise durch Russl. i. 115 = C. corymbosum ?
corymbiferum, Schrank, Baier. Fl. ii. 405 = C. corymbiferum.
corymbosum, Kral. ex Ball, in Journ. Linn. Soc. xvi. (1878) 509 = C. Webbianum.
corymbosum, Link, in Buch, Beschr. Canar. Ins. 149 = C. anethifolium.
corymbosum, Scop. Fl. Carn. ed. II. ii. 148 ; Moench, Meth. 597 = C. corymbosum.
Cota, Eichw. Skizze, 149 = Anthemis Cota ?
crassipes, Stschegl. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 172.—Soongar.
crithmifolium, Willd. Enum. Hort. Berol. 903 = C. anethifolium.
daucifolium, Ledeb. Fl. Ross. ii. 549 = C. daucifolium.
Decaisneanum, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 426.—Japon.
decipiens, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 48 = Matricaria decipiens.
defloratum, Hort. ex A. Br. in Bot. Zeit. x. (1852) 650 = Matricaria discoidea.
delicatulum, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 31 = Matricaria oreades.
densum, Labill. Ic. Pl. Syr. Dec. iii. 12. t. 8 = C. densum.
discoideum, Bess. ex DC. Prod. vi. 131 = C. argenteum.
discoideum, Ledeb. Ic. Fl. Ross. t. 153 = Tanacetum Ledebourii.
diversifolium, R. Grah. in Edinb. N. Phil. Journ. (Apr.–Jul. 1826) 174 ; et in Hook. Exot. Fl. iii. t. 215 = Brachycome diversifolia.
duderanum, Boiss. Diagn. Ser. I. xi. 26 = *myriophyllum*.
dumosum, Boiss. Fl. Orient. iii. 349 = Tanacetum dumosum.
elegans, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 101 = C. maritimum.
elongatum, Duch. in Journ. Soc. Hortic. (1858) 658 = C. cinerariaefolium.
elongatum, Fisch. & Mey. ex DC. Prod. vi. 56 = Matricaria oreades.
elongatum, Link, in Buch, Beschr. Canar. Ins. 149 = = C. Broussonetii.
exaltatum, Griseb. Spicil. Fl. Rumel. ii. 201 = C. trichophyllum.
ferulaceum, Webb & Berth. Phyt. Canar. ii. 281 = Tanacetum ferulaceum.
filipendulinum, Boiss. Diagn. Ser. I. xi. 24 = Tanacetum polycephalum.
Fischeri, Willd. ex Nym. Consp. 372 = C. corymbosum.
flabellifolium, Boiss. Fl. Orient. iii. 355 = Tanacetum flabellifolium.
flaveolum, Hoffmgg. & Link, Fl. Port. ii. 341. t. 104.—Hispan.
flavo-virens, Boiss. Diagn. Ser. I. xi. 25.—Persia.
foeniculaceum, Link, in Buch, Beschr. Canar. Ins. 149 = seq. ?
foeniculaceum, Willd. Enum. Hort. Berol. 903 = C. anethifolium.
frutescens, Willd. Sp. Pl. iii. 2150 = C. frutescens.
fruticosum, Spreng. ex DC. Prod. vi. 45 = C. praealtum.
fruticulosum, Fenzl, ex Boiss. Fl. Orient. iii. 338.—As. Min.
fruticulosum, Spreng. Cent. 37 ; Fl. Hal. Mant. 51 = C. praealtum.
fuscatum, Willd. Sp. Pl. iii. 2156 = Matricaria fuscata.
Gayanum, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 15 ; xxii. (1875) 61 = C. Mawii.
glabratum, Less. Syn. Comp. 254 = Matricaria glabrata.
glabrum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 48 = seq. ?
glabrum, Lag. Gen. et Sp. Nov. 30 = C. paludosum.

PYRETHRUM:—
graminifolium, Guss. Pl. Rar. 349 = Chrysanthemum graminifolium.
grandiflorum, Holl. & Reichb. in Flora, xiii. (1830) 382 = C. pinnatifidum.
grandiflorum, Willd. Enum. Hort. Berol. 904 = C. grandiflorum.
Halleri, Willd. Sp. Pl. iii. 2152 = C. coronopifolium.
Heldreichianum, Fenzl, in Tchihat. Asie Min. Bot. ii. 273 = *fruticulosum*, Fenzl.
heterophyllum, Baumg. ex Fuss, in Verh. Siebenb. Ver. Naturw. x. (1859) 104 = C. heterophyllum.
hirtum, Less. Syn. Comp. 254 = Matricaria hirta.
hispanicum, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 316 = Matricaria Chamomilla.
hispanicum, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 98 = C. pulverulentum.
hispidum, Link, in Buch, Beschr. Canar. Ins. 149 = C. frutescens.
hybridum, Wender. Ind. Sem. Hort. Marb. App. 9 = C. macrophyllum.
incanum, Ledeb. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 620 = Tanacetum Meyerianum.
indicum, Cass. in Dict. Sc. Nat. xliv. 149 = C. indicum.
indicum, Roxb. Hort. Beng. 62; Fl. Ind. iii. 436 = C. coronarium.
inodorum, Moench, Meth. 597 = Matricaria inodora.
jacobaeaeforme, C. Koch, in Linnaea, xxiv. (1851) 331 = C. palustre.
Kirilowii, Turcz. in DC. Prod. vi. 61.—Sibir. baical.
Kittaryanum, C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vii. (1850) 4.—Rossia.
Kotschyi, Boiss. Diagn. Ser. I. vi. 88.—Persia.
lanatum, Spreng. ex DC. Prod. vi. 42 = Egletes bellidiflora.
larvatum, Nym. Consp. 371, in syn. = C. coronopifolium.
latifolium, Willd. Enum. Hort. Berol. 904 = C. lacustre.
lavandulaefolium, Fisch. ex Trautv. in Act. Hort. Petrop. i. (1871–72) 181.—Mongol.
leptophyllum, Stev. ex Bieb. Fl. Taur. Cauc. iii. 580 = Tanacetum leptophyllum.
leucanthemiflorum, Boiss. & Huet, Diagn. Ser. II. iii. 28 = Tanacetum Aucherianum.
leucanthemoides, Tausch, in Flora, xii. (1829) i. Erg. 61 = C. atratum.
Leucanthemum, Franch. Fl. Cher-et-Loir, 307 = C. Leucanthemum.
leucophyllum, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 355.—Syria.
lingulatum, Boiss. l. c. 357.—Persia.
macrocarpum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61, nomen.—Algeria.
macrocephalum, Coss. & Dur. ex Coss. l. c. 48 = C. trifurcatum.
macrophyllum, Willd. Sp. Pl. iii. 2154 = C. macrophyllum.
macrotum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61 = C. macrotum.
Mandianum, Hort. ex G. Don, in Loud. Hort. Brit. 353.—Ins. Maurit.
Maresii, Coss. in Bull. Soc. Bot. Fr. iv. (1857) 16 = C. Maresii.
marginatum, Miq. Fl. Ind. Bat. ii. 85 = Tanacetum marginatum.
maritimum, Sm. Fl. Brit. ii. 901 = C. maritimum.
matricarioides, Paxt. ex Vilm. Fl. Pl. Terre, ed. III. 264 = C. indicum.
maximum, Hoffmgg. Verz. Pfl. Nachtr. ii. 189 = C. lacustre.
microcephalum, Hassk. Cat. Hort. Bog. Alt. 101.—Japon.
millefoliatum, Ledeb. Fl. Alt. iv. 120 = Tanacetum Meyerianum.
millefoliatum, Willd. Sp. Pl. iii. 2160 = C. millefoliatum.
minimum, DC. Fl. Fr. iv. 924 = C. tomentosum.
modestum, Heimerl, ex Stapf, in Denkschr. Akad. Wien, l. (1885) 59.—Persia.
montanum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 89 = C. montanum?
mucronatum, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 619.—Turkestan.
mucronulatum, Hoffmgg. & Link, Fl. Port. ii. 337.—Lusit.
multicaule, Willd. Sp. Pl. iii. 2158 = C. multicaule.

PYRETHRUM:—
multifidum, DC. Prod. vi. 63 = Chrysanthemum multifidum.
Myconis, Moench, Meth. Suppl. 247 = C. Myconis.
myriophyllum, C. A. Mey. Verz. Pfl. Cauc. 74 = Tanacetum myriophyllum.
nitens, Boiss. & Noé, Diagn. Ser. II. iii. 30.—Armenia.
nivale, Adam, ex Spreng. Syst. iii. 585 = Anthemis iberica.
nivale, Sch. Bip. in Boiss. Fl. Orient. iii. 339 = Tanacetum nivale.
niveum, Lag. Elench. Hort. Madr. (1805); Gen. et Sp. Nov. 30 = C. praealtum.
oligocephalum, DC. Prod. vii. 297 = C. millefoliatum.
oppositifolium, Spreng. Syst. iii. 585 = C. oppositifolium.
orientale, Willd. Sp. Pl. iii. 2159 = Anthemis Bieberstieniana.
Pallasianum, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 423.—As. bor.
palmatifidum, DC. Prod. vi. 54 = Tanacetum palmatifidum.
palustre, Willd. Sp. Pl. iii. 2154 = C. palustre.
parthenifolium, Willd. l. c. 2156 = C. praealtum.
Parthenium, Sm. Fl. Brit. ii. 900 = C. Parthenium.
parviflorum, Willd. Sp. Pl. iii. 2158 = Matricaria inodora.
paucifolium, C. A. Mey. ex Boiss. Fl. Orient. iii. 340 = C. coccineum.
pectinatum, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 180 = Tanacetum pulchellum.
pectinatum, Hoffmgg. & Link, Fl. Port. ii. t. 103 = C. pectinatum.
persicum, Boiss. Diagn. Ser. I. vi. 89.—Persia.
pinnatifidum, Link, in Buch, Beschr. Canar. Ins. 149 = C. grandiflorum.
pinnatifidum, Willd. Sp. Pl. iii. 2154 = C. multifidum.
pinnatum, Boiss. Fl. Orient. iii. 355 = Tanacetum pinnatum.
platyrachis, Boiss. l. c. 356.—Persia.
polycephalum, Sch. Bip. ex Boiss. l. c. 351.—Mesopot.
polyphyllum, Boiss. l. c. = Tanacetum millefoliatum.
poteriifolium, Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 312 = C. corymbosum.
Pourretii, Timb. ex Nym. Consp. 372 = C. corymbosum.
praecox, Bieb. Fl. Taur. Cauc. ii. 324 = Matricaria praecox.
procumbens, Kostel. Allg. Med.-Pharm. Fl. ii. 691 = C. indicum.
Pseud-anthemis, Boiss. & Huet, Diagn. Ser. II. iii. 28 = Tanacetum Aucherianum.
Pseudo-parthenium, Boiss. & Kotschy, Diagn. Ser. II. iii. 28 = Tanacetum nivale.
ptarmicaeflorum, Webb & Berth. Phyt. Canar. ii. 282.—Ins. Canar.
ptarmicaefolium, Willd. Sp. Pl. iii. 2151 = Achillea grandiflora.
pulchellum, Turcz. ex DC. Prod. vi. 55 = Tanacetum pulchellum.
pulchrum, Ledeb. Ic. Pl. Alt. i. 20. t. 84 = Tanacetum pulchrum.
Pulicaria, Willd. ex Spreng. Syst. iii. 589 = Lidbeckia americana.
pulverulentum, Lag. in Varied. Cienc. ii. iv. (1805) 40 = C. pulverulentum.
pulverulentum, Willd. Enum. Hort. Berol. 906 = C. praealtum.
pumilum, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 40 = Matricaria praecox.
pusillum, Willd. ex Spreng. Syst. iii. 587 = Matricaria praecox.
radicans, Cav. Desc. 199 = C. radicans.
roseum, Bieb. Fl. Taur. Cauc. ii. 324 = C. coccineum.
roseum, Lindl. Bot. Reg. t. 1084 = C. coronopifolium.
Roylei, DC. Prod. vi. 56 = C. tibeticum.
santolinoides, DC. in Ann. Sc. Nat. Sér. II. ii. (1834) 264 = Tanacetum sinaicum.
saxatile, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 382 = C. millefoliatum.
saxatile, Lodd. Bot. Cab. t. 126 = C. alpinum.
scariosum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 61.—Algeria.
segetum, Moench, Meth. 597 = C. segetum.
sericeum, Bieb. Fl. Taur. Cauc. ii. 323 = C. argenteum.

PYRETHRUM:—
serotinum, Hort.—Cf. Gard. Chron. (1878) ii. 492 = Chrysanthemum uliginosum.
serotinum, Willd. Sp. Pl. iii. 2151 = C. serotinum.
setaceum, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 44.—Kokan.
seticuspe, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 423.—Japon.
setifolium, Regel, Ind. Sem. Hort. Petrop. (1876) 36, nomen.—Hab.?
silaifolium, Stev. in DC. Prod. vi. 57 = Matricaria grandiflora.
simplicifolium, Willd. Sp. Pl. iii. 2151 = Egletes domingensis.
sinense, DC. Prod. vi. 62 = C. sinense.
sorbifolium, Boiss. ex Boiss. Fl. Orient. iii. 343.—Armenia.
spathulaefolium, J. Gay, ex Nym. Consp. 373.—Oriens.
speciosum, Willd. Enum. Hort. Berol. Suppl. 60 = C. grandiflorum.
subcorymbosum, Schur, Bot. Rundreise (1853), 80; et in Verh. Siebenb. Ver. Naturw. (1859) 146 = C. subcorymbosum.
sulphureum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 17 = Tanacetum sulphureum.
syriacum, Boiss. Diagn. Ser. I. xi. 24 = C. densum.
Szowitsii, Boiss. l. c. 25 = Tanacetum myriophyllum.
tabrisianum, Boiss. Fl. Orient. iii. 356.—Persia.
tanacetifolium, Boiss. l. c. 341 = Tanacetum Aucherianum.
tanacetoides, DC. Prod. vi. 59 = Tanacetum Meyerianum.
Tanacetum, DC. l. c. 63 = C. Balsamita.
Tchihatchewii, Hort. ex Clémenc. in Rev. Hortic. (1869) 380.—Oriens?
tenuifolium, Willd. Enum. Hort. Berol. 906 = C. corymbosum.
tenuilobum, Boiss. Fl. Orient. iii. 352 = Tanacetum Aucheri.
tenuisectum, Boiss. l. c. 338.—Persia.
tenuissimum, Trautv. in Act. Hort. Petrop. ix. (1884) 392.—Turkest.
tibeticum, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 147 = C. tibeticum.
tingitanum, DC. Prod. vi. 54 = C. tingitanum.
tomentellum, Boiss. Fl. Orient. iii. 354 = Tanacetum tomentellum.
tomentosum, [Clairv.]. Man. Herb. 247 = C. alpinum.
tomentosum, DC. Fl. Fr. Suppl. 477 = C. tomentosum.
transiliense, Regel, in Act. Hort. Petrop. v. (1877) 618.—Turkestan.
trichophyllum, Griseb. Spicil. Fl. Rumel. ii. 201 = C. trichophyllum?
trifurcatum, Willd. Sp. Pl. iii. 2158 = C. trifurcatum.
Turreyanum, Less. Syn. Comp. 254 = C. cinerariaefolium.
uliginosum, Waldst. & Kit. in Willd. Sp. Pl. iii. 2152 = C. uliginosum.
umbelliferum, Boiss. Fl. Orient. iii. 352 = Tanacetum umbelliferum.
umbrosum, Boiss. l. c. 357 = C. umbrosum.
uniflorum, Fisch. & Mey. in DC. Prod. vi. 60 = Tanacetum uniflorum.
Vahlii, Boiss. & Reut. Diagn. Ser. II. iii. 29 = Tanacetum Vahlii.
velutinum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) i. 419 = C. bipinnatum.
versicolor, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 98 = C. pulverulentum.
vulgare, Boiss. Fl. Orient. iii. 352 = Tanacetum vulgare.
Waldsteinii, Janka, in Just, Jahresb. iv. (1878) 1062 = C. rotundifolium.
Webbianum, Coss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 509 = C. Webbianum.
Willemoti, Duch. in Journ. Soc. Hortic. (1859) 210 = C. cinerariaefolium.
Willkommii, Sch. Bip. ex Nym. Syll. 11 = Tanacetum Vahlii.
Zawadskii, Nym. l. c. = C. Zawadskii.

PYRETHRUM, Medic. in Act. Acad. Theod. Palat. iii. (1775) 237. t. 18 = **Spilanthes**, Jacq. (Compos.).
Acmella, Medic. l. c. 243. t. 19 = S. Acmella.
bidens, Medic. l. c. 241. t. 18 = seq.?
Spilanthus, Medic. l. c. 242 = S. oleracea.

PYRGOSEA, Eckl. & Zeyh. Enum. 298 (1837) = PURGOSEA, Haw. = **Crassula**, Linn.

PYRGUS, Lour. Fl. Cochinch. 120 (1790) = **Ardisia**, Sw. (Myrsin.).
racemosa, Lour. l. c. 121 = A. Pyrgus.

PYROLA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 123 (1737). *ERICACEAE*, Benth. & Hook. f. ii. 602.
ACTINOCYCLUS, Klotzsch, in Monatsb. Akad. Berl. (1857) 14.
AMELIA, Alefeld, in Linnaea, xxviii. (1856) 25.
ERXLEBENIA, Opiz, Seznam, 41 (1852).
RAMISCHIA, Opiz, l. c. 82 (1852).
THELAIA, Alefeld, in Linnaea, xxviii. (1856) 33.
americana, G. Don, in Sweet, Hort. Brit. ed. II. 341 = rotundifolia.
angustifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 283.—Mexic.
aphylla, *Sm. in Rees, Cycl.* xxix. n. 7.—Am. bor.
asarifolia, Michx. Fl. Bor. Am. i. 251 = rotundifolia.
asarifolia, Radius, Diss. 23. t. 4 = chlorantha.
bracteata, Hook. Fl. Bor. Am. ii. 47 = rotundifolia.
chlorantha, *Sw. in Vet. Acad. Handl. Stockh.* (1810) 190. t. 5.—Europ.; Am. bor.
conferta, Fisch. ex Ledeb. Fl. Ross. ii. 930 = minor.
conferta, Willd. ex DC. Prod. vii. 773 = rotundifolia.
convoluta, Barton, Fl. Philad. Prod. 50 = chlorantha.
corymbosa, Bertol. Misc. Bot. iii. 12 = Chimaphila umbellata.
declinata, Moench, Meth. 71 = rotundifolia.
dentata, Sm. in Rees, Cycl. xxix. n. 6 = picta.
Durandi, Rafin. Atl. Journ. 119 (Quid?).—Am. bor.
elata, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 270 = rotundifolia.
elliptica, *Nutt. Gen. Am.* i. 273.—Am. bor.
flexuosa, Rafin. Atl. Journ. 151 (Quid?).—Am. bor.
frutescens, Gilib. Fl. Lituan. ii. 195 = Chimaphila umbellata.
fruticosa, Eschsch. ex Ledeb. Fl. Ross. ii. 924 = Cladothamnus pyrolaeflorus.
grandiflora, Radius, Diss. 27. t. 3. f. 2 = rotundifolia.
groenlandica, Hornem. in Fl. Dan. t. 1817 = rotundifolia.
Halleri, Steud. Nom. ed. II. ii. 422 = Moneses grandiflora.
hybrida, Vill. Hist. Pl. Dauph. iii. 588 = secunda.
incarnata, Fisch. ex DC. Prod. vii. 773 = rotundifolia.
intermedia, Schleich. Cat. ed. III. 23 = minor.
Liebmannii, *Lange, in Kjoeb. Vidensk. Meddel.* (1867) 113. t. 1.—Mexic.
maculata, Linn. Sp. Pl. 396 = Chimaphila maculata.
magna, Gueldenst. ex Ledeb. Fl. Ross. ii. 928 = rotundifolia.
major, Lam. Fl. Fr. ii. 530 = rotundifolia.
maritima, Kenyon, in Phytol. ii. (1847) 727 = rotundifolia.
media, Hayne, Arzneigew. iv. 22 = chlorantha.
media, *Sw. in Vet. Acad. Handl. Stockh.* (1804) 257.—Europ.; Reg. Caucas.
Menziesii, R. Br. ex D. Don, in Mem. Wern. Soc. v. (1824) 245 = Chimaphila Menziesii.
minor, Gilib. ex DC. Prod. vii. 773 = rotundifolia.
minor, *Linn. Sp. Pl.* 396.—Reg. bor. temp.
minor, Pursh, Fl. Am. Sept. i. 299 = chlorantha.
noveboracensis, Colden, ex Torr. Fl. N. U. St. i. 432 = rotundifolia.
obovata, Bertol. Misc. Bot. iii. 11 = rotundifolia.
occidentalis, R. Br. ex D. Don, in Mem. Wern. Soc. v. (1824) 232 = chlorantha.
ovalifolia, Schur, Enum. Pl. Transs. 449 = rotundifolia.
oxypetala, *C. F. Austin, ex A. Gray, Man. Bot. N. U. St.* ed. V. 302.—Am. bor.
picta, *Sm. in Rees, Cycl.* xxix. n. 8.—Am. bor.
pumila, Hornem. ex Cham. & Schlecht. in Linnaea, i. (1826) 514 = rotundifolia.
renifolia, *Maxim. Prim. Fl. Amur.* 190.—Manchuria.
reticulata, Rafin. ex DC. Prod. vii. 773 = elliptica.
rosea, Sm. Engl. Bot. t. 2543 = minor.
rotundifolia, Benth. Pl. Hartw. 66 = angustifolia.
rotundifolia, *Linn. Sp. Pl.* 396.—Reg. bor. temp.
rotundifolia, Michx. Fl. Bor. Am. i. 251, partim = elliptica.
rotundifolia, Oed. in Fl. Dan. t. 110 = media.
rotundifolia, Pall. ex DC. Prod. vii. 774 = minor.
rotundifolia, Sturm, Deutschl. Fl. Heft 13 = chlorantha.

PYROLA:—
Sartorii, *Hemsl. Biol. Centr. Am. Bot.* ii. 283.—Mexic.
secunda, *Linn. Sp. Pl.* 396.—Reg. bor. temp.
serotina, Melicocq, in Ann. Pas-de-Calais (1848-49), 223.—Cf. Bull. Soc. Bot. Fr. i. (1854) 162 = rotundifolia.
subaphylla, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 433.—Japon.
uliginosa, Torr. & Gray, ex Torr. Fl. N. York, i. 453 = rotundifolia.
umbellata, Linn. Sp. Pl. 396 = Chimaphila umbellata.
uniflora, Linn. l. c. 397 = Moneses grandiflora.
urceolata, Poir. Encyc. v. 743 = Galax aphylla.
verticillata, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = Moneses grandiflora.
virens, Schweig. & Koerte, Fl. Erlang. Add. 154 = chlorantha.

PYROLIRION, Herb. App. [Bot. Reg.] 37 (1821) = **Zephyranthes**, Herb. (Amaryll.).
albicans, Herb. Amaryll. 184.—Peruv.
aurantiacum, Lem. Jard. Fleur. iv. (1854) t. 377. Misc. 27.—Peruv.
aureum, Herb. App. [Bot. Reg.] 37 = Z. aurea.
flammeum, Herb. l. c.—Peruv.
flavum, Herb. l. c.—Peruv.
tubiflorum, M. Roem. Syn. Ensat. 127 = Z. aurea.

PYROPHORUM, DC. Prod. ii. 633 (1825) = PIROPHORUM, Neck. = **Pyrus**, Tourn. (Rosac.).

PYROPSIS, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 22 (1840) = **Madia**, Molina (Compos.).
elegans, Hort. ex Fisch. Mey. & Avé-Lall. l. c. = M. elegans.

PYROSPERMUM, Miq. Fl. Ind. Bat. Suppl. 402 (1860) = **Kurrimia**, Wall. (Celastr.).
calophyllum, Miq. l. c. = K. paniculata.

PYROSTEGIA, Presl, (Bot. Bemerk. 93 (1844)); in Abh. Boehm. Ges. V. iii. (1845) 523 = **Bignonia**, Tourn.
acuminata, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 188, nomen.—Bras.
amabilis, Miers, l. c.—Guian.
dichotoma, Miers, l. c. = B. dichotoma, *Vell.*
ignea, Presl, Bot. Bemerk. 93 = B. ignea.
intaminata, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 188, nomen.—Bras.
lauta, Miers, l. c.—Bras.
longiflora, Miers, l. c. = B. longiflora.
ornata, Miers, l. c. = B. ornata.
pallida, Miers, l. c. = B. pallida.
parvifolia, Miers, l. c.—Bras.
puberula, Miers, l. c.—Bras.
reticulata, Miers, l. c.—Paraguay.
venusta, Miers, l. c. = B. venusta.

PYROSTOMA, G. F. W. Mey. Prim. Fl. Esseq. 219 (1818) = **Vitex**, Linn. (Verbenac.).
ternatum, G. F. W. Mey. l. c. 220 = V. triflora.

PYROSTRIA, Comm. ex Juss. Gen. 206 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 111.
Commersonii, *J. F. Gmel. Syst.* 246.—Ins. Maurit.
comorensis, Boj. ex Baker, Fl. Maurit. 146 = Plectronia bibracteata.
cordifolia, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 216 = macrophylla.
fasciculata, *Boj. ex Baker, Fl. Maurit.* 148.—Ins. Mascar.
lancifolia, *Boj. ex Baker, l. c.* 149.—Ins. Mascar.
macrophylla, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 217.—Ins. Maurit.
obliqua, Boj. & Baker, Fl. Maurit. 149 = fasciculata.
oleoides, Boj. Hort. Maurit. 170 = seq. ?
oleoides, Lam. Illustr. i. 289 = Commersonii.
orbicularis, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 216 = macrophylla.
polymorpha, *A. Rich. l. c.* 217.—Ins. Maurit.
salicifolia, Willd. Sp. Pl. i. 614; Pers. Syn. i. 131 = Commersonii.
spinosa, Miq. Fl. Ind. Bat. ii. 313 = Vangueria spinosa.
trilocularis, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 14.—Ins. Rodrig.
umbellata, Boj. Hort. Maurit. 170 = Plectronia umbellata.

PYROSTRIA, Roxb. Hort. Beng. 83 (1814); Fl. Ind. i. 403 (1832) = **Timonius**, Rumph. (Rubiac.).
hexasperma, Roxb. ll. cc. = T. Rumphii?

PYROTHECA, Steud. Nom. ed. II. ii. 422 (1841), sphalm. = GYROTHECA, Salisb. = **Lachnanthes**, Ell. (Haemod.).
tinctoria, Steud. l. c. = L. tinctoria.

PYRRANTHUS, A. Gray, Bot. U. St. Expl. Exped. i. 615, in syn. (1854) = seq.

PYRRHANTHUS, Jack, in Malay. Misc. ii. (1822) n. VII. 57 = **Lumnitzera**, Willd. (Combret.).
albus, Wall. Cat. n. 4019 = L. racemosa.
littoreus, Jack, in Malay. Misc. ii. (1822) n. VII. 57 = L. coccinea.

PYRRHEIMA, Hassk. in Flora, lii. (1869) 366 = **Tradescantia**, Linn. (Commel.).
Loddigesii, Hassk. l. c. 367 = T. fuscata.
minus, Hassk. l. c. 368 = T. fuscata.

PYRRHOCOMA, Walp. Rep. ii. 593 (1843) = PYRROCOMA, Hook. = **Aplopappus**, Cass. (Compos.).

PYRRHOPAPPUS, DC. Prod. vii. 144 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 523.
SITILIAS, Rafin. New Fl. Am. iv. 85 (1836).
carolinianus, *DC. Prod.* vii. 144.—Am. bor.
grandiflorus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 430 = scaposus.
multicaulis, Curtiss, ex A. Gray, Syn. Fl. N. Am. i. II. 441 = carolinianus.
multicaulis, *DC. Prod.* vii. 144.—Texas; Mexic.
pauciflorus, *DC. l. c.*—Mexic.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xi. (1876) 80.—Arizona.
scaposus, *DC. Prod.* vii. 144.—Am. bor. occ.
Sessaeanus, *DC. l. c.*—Mexic.
taraxacoides, DC. l. c. 145 (al. gen.).—Afr. austr.

PYRRHOPAPPUS, A. Rich. Tent. Fl. Abyss. i. 463 (1847) = **Lactuca**, Tourn. (Compos.).
Hochstetteri, A. Rich. l. c. = L. capensis.
humilis, A. Rich. l. c. = L. humilis.

PYRRHOSA, Endl. Gen. 830 (1839) = **Myristica**, Linn.
Horsfieldii, Wight, Ic. t. 1857 = M. Horsfieldia.

PYRRHOTRICHIA, Wight & Arn. Prod. 238 (1834) = **Eriosema**, DC. (Legumin.).
tuberosa, Wight & Arn. l. c. = E. chinense.

PYRROCOMA, Hook. Fl. Bor. Am. i. 306. t. 107 (1833) = **Aplopappus**, Cass. (Compos.).
angustifolia, DC. Prod. v. 351.—Chili.
arguta, Torr. & Gray, Fl. N. Am. ii. 244 = A. racemosus.
aurea, Phil. in Linnaea, xxviii. (1856) 733.—Chili.
Bezanillana, Remy, in C. Gay, Fl. Chil. iv. 65.—Chili.
canescens, Phil. in Linnaea, xxviii. (1856) 730.—Chili.
carthamoides, Hook. Fl. Bor. Am. i. 307 = A. carthamoides.
Cerberoana, Remy, in C. Gay, Fl. Chil. iv. 66.—Chili.
denticulata, Phil. in Linnaea, xxviii. (1856) 730.—Chili.
foliosa, Phil. l. c. 732.—Chili.
foliosa, Torr. & Gray, ex Duch. Rev. Bot. i. (1845-46) 81 = A. Fremonti.
glomerata, Torr. & Gray, Fl. N. Am. ii. 244 = A. racemosus.
grindelioides, DC. Prod. v. 350 = A. squarrosus.
ilicifolia, Remy, in C. Gay, Fl. Chil. iv. 67 = A. Remyanus.
linifolia, Phil. in Linnaea, xxviii. (1856) 730.—Chili.
longipes, Phil. l. c. 733.—Chili.
Macraeana, Remy, in C. Gay, Fl. Chil. iv. 64.—Chili.
Menziesii, Hook. & Arn. Bot. Beech. Voy. 351 = Bigelovia veneta.
Nuttallii, Remy, in C. Gay, Fl. Chil. iv. 67 = A. pinnatifidus.

PYRROCOMA :—
paniculata, Torr. & Gray, Fl. N. Am. ii. 244 = Aplopappus racemosus.
parvifolia, DC. Prod. v. 351.—Chili.
pinea, Phil. in Linnaea, xxviii. (1856) 731.—Chili.
pristiphylla, Remy, in C. Gay, Fl. Chil. iv. 65.—Chili.
racemosa, Torr. & Gray, Fl. N. Am. ii. 244 = A. racemosus.
radiata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 333 = A. carthamoides.
reticulata, Phil. in Linnaea, xxviii. (1856) 731.—Chili.
rigida, Phil. l. c. 732.—Chili.
saxatilis, Remy, in C. Gay, Fl. Chil. iv. 64.—Chili.
setigera, Phil. in Linnaea, xxviii. (1856) 733.—Chili.

PYRROTRICHIA, Baker, in Hook. f. Fl. Brit. Ind. ii. 219 (1876) = praec.

PYRULARIA, Michx. Fl. Bor. Am. ii. 231 (1803). *SANTALACEAE*, Benth. & Hook. f. iii. 223.
CALINUX, Rafin. in Med. Repos. N. York, v. (1808) 352.
HAMILTONIA, Muhl. ex Willd. Sp. Pl. iv. 1114 (1805).
SPHAEROCARYA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 371 (1824).
ceylanica, A. DC. in DC. Prod. xiv. 629 = Scleropyrum Wallichianum.
edulis, *A. DC. l. c.* 628.—Reg. Himal.
moschifera, *A. DC. l. c.* 629.—Java.
oleifera, A. Gray, ex Chapm. Fl. S. U. St. 396 = seq.
pubera, *Michx. Fl. Bor. Am.* ii. 231.—Am. bor.
Wallichiana, A. DC. in DC. Prod. xiv. 629 = Scleropyrum Wallichianum.

PYRUS, [Tourn.] Linn. Syst. ed. I (1735). *ROSACEAE*, Benth. & Hook. f. i. 626.
APIROPHORUM, Neck. Elem. ii. 72 (1790).
ARIA, M. Roem. Fam. Rosifl. iii. 101, 124 (1847).
AZAROLUS, Borckh. Handb. Forstb. ii. 1224 (1803).
CHAENOMELES, Lindl. in Trans. Linn. Soc. xiii. (1822) 97.
CHAMAEMESPILUS, Medic. Phil. Bot. i. 138, 135 (1789).
CYDONIA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
ERIOLOBUS, M. Roem. Fam. Rosifl. iii. 104, 216 (1847).
HAHNIA, Medic. Gesch. 81 (1793).
LAZAROLUS, Medic. Phil. Bot. i. 15 (1789).
MALUS, Tourn. ex Linn. Gen. ed. I. 145, in syn. (1737); Mill. Gard. Dict. ed. VI (1752).
MESPILOPHORA, Neck. Elem. ii. 73 (1790).
MESPILUS, [Tourn.] Linn. Syst. ed. I. (1735).
MICROMELES, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 168.
PIROPHORUM, Neck. Elem. ii. 72 (1790).
PYRENIA, [Clairv.] Man. Herb. 161 (1811).
SORBUS, [Tourn.] Linn. Syst. ed. I (1735).
TORMINALIS, Medic. Phil. Bot. i. 134, 155 (1789).
TORMINARIA, M. Roem. Fam. Rosifl. iii. 101, 130 (1847).
acerba, DC. Prod. ii. 635 = P. Malus.
Achras, Gaertn. Fruct. ii. 44. t. 87 = communis.
aestivalis, Lamotte, Prod. Fl. Centr. Fr. i. 279 = communis.
albicans, Kit. in Linnaea, xxxii. (1863) 583 = slavonica.
alnifolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 350.—Japon.
alnifolia, Lindl. in Trans. Linn. Soc. xiii. (1821) 98 = Amelanchier alnifolia.
alpina, Willd. Enum. Hort. Berol. 527 = P. Chamaemespilus?
Amelanchier, Du Roi, Harbk. Baumz. ii. 219; Lam. Fl. Fr. iii. 492; Linn. f. Suppl. 256 = Amelanchier vulgaris.
Amelanchier, Hort. ex C. Koch, Dendrol. i. 181 = Amelanchier canadensis.
americana, Cooper, in Am. Natural. iii. (1870) 406 = sambucifolia.
americana, *DC. Prod.* ii. 637; *Spreng. Syst.* ii. 511.—Am. bor.
amygdaliformis, *Vill. Cat. Hort. Argent.* 323.—Europ. austr.; As. Min.
amygdaloides, Link, Handb. ii. 80, sphalm. = praec.
angulata, *Kit. ex Rochel, Bot. Reise Banat.* 73; *et in Linnaea*, xxxii. (1863) 583.—Europ.

PYRUS:—
angustifolia, *Ait. Hort. Kew.* ed. I. ii. 176.—Am. bor.
angustifolia, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 359 = Bovei.
anthyllidifolia, Sm. in Rees, Cycl. xxix. n. 29 = Osteomeles anthyllidifolia.
apetala, Munchh. Hausv. v. 247 = P. Malus.
apiosa, Hort. ex Steud. Nom. ed. I. 670 = P. Malus.
arbutifolia, *Linn. f. Suppl.* 256.—Am. bor.
arctica, Hort. ex C. Koch, Dendrol. i. 192 = P. Aria.
arguta, Tausch, in Flora, xvii. (1834) 490 = intermedia.
Aria, Brandis, For. Fl. 206 = lanata.
Aria, *Ehrh. Beitr.* iv. 20.—Europ.; As. bor.
Aria, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. ii. 375 = vestita.
Aria, Wimm. & Grab. Fl. Siles. (1829) ii. 21 = sudetica.
armeniacaefolia, Dochn. Obstk. i. 16 = P. Malus.
astracanica, DC. Prod. ii. 635 = P. Malus.
atrovirens, Hort. ex C. Koch, Dendrol. i. 194 = rotundifolia.
Aucuparia, *Ehrh. Plantag.* 20; *ex ej. Beitr.* vi. 94; *Gaertn. Fruct.* ii. 45. t. 87.—Europ.; As. bor.
Aucuparia, E. Mey. Pl. Labrad. 81 = americana, sambucifolia.
auricularis, Knoop, Pomol. ii. 38. t. 4 = P. Pollveria.
auriculata, DC. Prod. ii. 636 = pinnatifida?
Azarolus, Scop. Fl. Carn. ed. II. i. 347 = Crataegus Azarolus.
baccata, Hort. ex Hoffmgg. Verz. Pfl. 104 = rubicunda.
baccata, *Linn. Mant.* i. 75.—Reg. Himal.; As. bor.
badensis, Hort. ex C. Koch, Dendrol. i. 192 = P. Aria.
Balansae, Decne. Jard. Fruit. Mus. i. t. 6 = communis.
barauyensis, *Kit. in Linnaea*, xxxii. (1863) 582 (Quid?). —Europ. austr.
Bartramiana, Tausch, in Flora, xxi. (1838) 715 = Amelanchier canadensis.
betulaefolia, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 101.—China.
Boissieriana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 87 = cordata.
Bollwylleriana, DC. Fl. Fr. Suppl. 530 = P. Pollveria.
Botryapium, Linn. f. Suppl. 255 = Amelanchier canadensis.
Bourgaeana, Decne. Jard. Fruit. Mus. i. t. 2 = communis.
Boveana, Decne. l. c. t. 10 = Bovei.
Bovei, *Steud. Nom.* ed. II. ii. 423.—Syria.
brachypoda, Kern. ex Wenzig, in Linnaea, xxxviii. (1874) 17 = communis.
Calleryana, *Decne. Jard. Fruit. Mus.* i. sub t. 8.—Japon.
canadensis, Billb. in Linneska Samfund, i. 1832 (1833) 24 = arbutifolia, Amelanchier canadensis.
canescens, Spach, Hist. Vég. Phan. ii. 129 = communis.
canoplia, Guss. ex Decne. Jard. Fruit. Mus. i. sub t. 13 = amygdaliformis.
capitata, Hort. ex Lindl. in Trans. Hort. Soc. vii. (1828) 232 = arbutifolia.
castanea, Steud. Nom. ed. I. 670, sphalm. = *cestiana*.
castaneifolia, *Hook. f. Fl. Brit. Ind.* ii. 379.—Reg. Himal.
Caucasi, Gueldenst. Reisen, i. 423 (Quid?).—Reg. Cauc.
cavillea, Hort. ex Steud. Nom. ed. I. 670 = P. Malus.
cerasifera, *Tausch, in Flora*, xxi. (1838) 714.—Sibir.
ceratocarpa, Wender. in Schrift. Marb. ii. (1830) 254 = prunifolia.
cestiana, Hort. ex Steud. Nom. ed. I. 670, in syn. = P. Malus.
Chamaemespilus, *Ehrh. Beitr.* iv. 19.—Europ.
chinensis, Hort. ex Steud. Nom. ed. I. 670, in syn. = communis.
chinensis, Roxb. Hort. Beng. 38; Fl. Ind. ii. 511 = sinensis, *Lindl.*
chinensis, Spreng. Syst. ii. 510 = sinensis, *Poir.*
ciliata, *Kit. in Linnaea*, xxxii. (1863) 582 (Quid?).—Europ. austr.
communis, Dur. in Bull. Soc. Bot. Fr. vii. (1860) 31, partim = cordata.
communis, Gouan, Hort. Monsp. 242 = amygdaliformis.
communis, *Linn. Sp. Pl.* 479.—Europ.; As. bor.; Reg. Himal.
communis, Thunb. Fl. Jap. 207 = sinensis, *Lindl.*

PYRUS:—
cordata, *Desv. Obs. Pl. Anj.* 152.—Gallia; Persia.
coronaria, *Linn. Sp. Pl.* 480.—Am. bor.
coronaria, Wangenh. Nord-Am. Holzart. 61. t. 21. f. 47 = angustifolia.
corymbosus, Gilib. Fl. Lituan. ii. 234 = communis.
Cotoneaster, Moench, Meth. 681 = Cotoneaster integerrima.
crataegifolia, *Targ.-Tozz. ex Savi, Tratt. Alb. Tosc.* ed. II. i. 169.—Ital.
crenata, D. Don, Prod. Fl. Nep. 237 = communis?
crenata, Hort. ex C. Koch, Dendrol. i. 192 = P. Aria.
crenata, Lindl. Bot. Reg. t. 1655 = vestita.
cretica, Hort. ex C. Koch, l. c. = P. Aria.
cretica, Willd. Sp. Pl. ii. 1015 = Amelanchier vulgaris.
csiklovensis, *Kit. in Linnaea*, xxxii. (1863) 583 (Quid?). —Europ. austr.
cuneifolia, Guss. Pl. Rar. 202 = amygdaliformis.
cuspidata, *Bertol. in Mem. Acad. Sc. Bolog.* Ser. II. iv. (1864) 311.—Reg. Himal.
Cydonia, *Linn. Sp. Pl.* 480.—Reg. Mediterr. et Cauc.
dasyphylla, Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 604; Borckh. Handb. Forstb. ii. 1269 = P. Malus.
decipiens, Bechst. Forstb. ed. V. (1843) 152 = rotundifolia.
densiflora, *Steud. Nom.* ed. II. ii. 423.—Hab.?
denticulata, *Hort. Angl. ex Dum.-Cours. Bot. Cult.* ed. II. v. 435.—China.
depressa, Lindl. in Trans. Hort. Soc. vii. (1828) 230 = arbutifolia.
dioica, Moench, Verz. 87 = P. Malus.
discolor, Maxim. Prim. Fl. Amur. 103 = P. Aucuparia.
diversifolia, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 133 = rivularis.
divisa, Link, Enum. Hort. Berol. ii. 38 = P. Malus.
domestica, Ehrh. Plantag. p. 20; ex ej. Beitr. vi. 95 = P. Sorbus.
domestica, Medic. Gesch. 87; Borckh. in Roem. Arch. i. III. 90 = communis.
edulis, Willd. Enum. Hort. Berol. 527; Wats. Dendrol. Brit. i. 52 = P. Aria.
elaeagnifolia, Steud. Nom. ed. I. 670 = seq.
elaeagrifolia, *Pall. in Nov. Act. Petrop.* vii. (1789) 355. t. 10.—As. Min.
epirotica, Hort. ex Steud. Nom. ed. I. 670 = P. Malus.
eriopleura, Reichb. Fl. Germ. Excurs. 630 = amygdaliformis.
eriostyla, Guss. Fl. Sic. Prod. Suppl. 154 = P. Malus.
falerna, Hort. ex Steud. Nom. ed. I. 670 = communis.
favonia, Hort. ex Steud. l. c. = communis.
fennica, Bab. Man. Brit. Bot. ed. III. 111 = pinnatifida.
ferruginea, *Hook. f. Fl. Brit. Ind.* ii. 379.—Reg. Himal.
florentina, Targ.-Tozz. in Mem. Soc. Mod. xx. II. (1829) 302. t. 5 = crataegifolia.
floribunda, Lindl. Bot. Reg. t. 1006 = arbutifolia.
foliolosa, Wall. Cat. n. 677, partim = Wallichii.
foliolosa, *Wall. Pl. As. Rar.* ii. 81. t. 189.—Reg. Himal.
Fontanesiana, Steud. Nom. ed. II. ii. 423 = P. Malus.
fraxinifolia, Cooper, in Am. Natural. iii. (1870) 406 = sambucifolia?
fusca, Rafin. Med. Fl. ii. 254 = rivularis.
germanica, *Hook. f. Stud. Fl.* ed. I. 127.—Europ. or.; Oriens.
glabra, *Boiss. Diagn.* Ser. I. vi. 53.—Persia.
glandulosa, Moench, Meth. 680 = Crataegus coccinea.
glutinosa, Madiot. ex Steud. Nom. ed. II. ii. 423 (Quid?).—Hab.?
gracilis, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 131 = sambucifolia.
graeca, Hort. ex Steud. Nom. ed. II. ii. 423 = P. Aria.
grandifolia, Lindl. in Trans. Hort. Soc. vii. (1828) 233; Bot. Reg. t. 1154 = arbutifolia.
granulosa, *Bertol. in Mem. Acad. Sc. Bolog.* Ser. II. iv. (1864) 312.—Reg. Himal.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 377.—Reg. Himal.
hadiensis, *Forsk. Fl. Aegypt. Arab.* 212.—Arabia.
heterophylla, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 581.—Turkestan.
heterophylla, *Steud. Nom.* ed. II. ii. 423.—Hab.?
Hoggii, Franch. & Sav. Enum. Pl. Jap. ii. 349 = Tschonoskii.

PYRUS :—
hortensis, Mill. ex Trautv. in Act. Hort. Petrop. viii. (1883) 339, nomen = communis?
Hostii, Hort. ex C. Koch, Dendrol. i. 192 = P. Aria.
hyberna, *Pall. Reise in Suedl. Statth. Russ. Reich.* i. 349 (Quid?).—Rossia.
hybrida, Moench, Verz. 90. t. 6 = spuria.
hybrida, Sm. Fl. Brit. ii. 534 = pinnatifida.
hybrida, Steud. Nom. ed. I. 670 = prunifolia.
hyemalis, *Hort. ex C. Koch, Hort. Dendrol.* 183, in adnot.—Hab.?
incana, *C. Koch, Dendrol.* i. 217.—Armenia.
indica, Coleb. in Wall. Pl. As. Rar. ii. 56 = Docynia indica.
indica, Roxb. Hort. Beng. 38; Fl. Ind. ii. 511 = seq.
insignis, *Hook. f. Fl. Brit. Ind.* ii. 377.—Reg. Himal.
integerrima, Wall. ex D. Don, Prod. Fl. Nep. 237 = Photinia integrifolia.
intermedia, Bechst. Diana, iv. 110 = rotundifolia.
intermedia, *Ehrh. Beitr.* iv. 20.—Europ.
intermedia, Soyer-Willem. ex Nym. Consp. 242 = latifolia.
intermedia, Tenore, Fl. Nap. iv. 279 = Crataegus tomentosa?
irregularis, Du Roi, Harbk. Baumz. ii. 216 = P. Pollveria.
Jacquemontiana, *Decne. Jard. Fruit. Mus.* i. t. 8.—Reg. Himal.
japonica, *Thunb. Fl. Jap.* 207.—Japon.
Kaido, Siebold, ex C. Koch, Dendrol. i. 209 = spectabilis.
kamounensis, Wall. Cat. n. 678 = lanata.
karensium, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 306 = granulosa.
khasiana, *Hook. f. Fl. Brit. Ind.* ii. 378.—Reg. Himal.
Kotschyana, Boiss. ex Decne. Jard. Fruit. Mus. i. t. 18 = elaeagrifolia.
kumaonensis, Hook. f. Fl. Brit. Ind. ii. 375, sphalm. = kamounensis.
Kumaoni, *Decne. Jard. Fruit. Mus.* sub t. 7.—Reg. Himal.
lanata, *D. Don, Prod. Fl. Nep.* 237.—Reg. Himal.
lanuginosa, *DC. Prod.* ii. 637.—Hungar.
latifolia, Lindl. ex Steud. Nom. ed. II. ii. 424 = intermedia.
latifolia, Syme, in Bot. Exch. Club Rep. 1872–4 (1875) 19 = rotundifolia.
liquescens, Hort. ex Steud. Nom. ed. I. 670 = communis.
lobata, C. Koch, Hort. Dendrol. i. 182 = Docynia indica.
longipes, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 310.—Alger.
ludoviciana, Rafin. Fl. Ludov. 158 (Quid?).—Am. bor.
lusitanica, Hort. ex Steud. Nom. ed. I. 670 = P. Cydonia.
malifolia, Spach, Hist. Vég. Phan. ii. 131 = P. Cydonia.
maliformis, Hort. ex Steud. Nom. ed. I. 670 = P. Cydonia.
Malus, *Linn. Sp. Pl.* 479.—Europ.; As. bor.; Reg. Himal.
marginata, Kit. in Linnaea, xxxii. (1863) 582 = ciliata.
matrensis, Kit. l. c. (Quid?).—Hungar.
Maulei, Mast. in Gard. Chron. i. (1874) 756 = japonica.
melanocarpa, Willd. Enum. Hort. Berol. 525 = arbutifolia.
Mengo, Siebold, ex C. Koch, Dendrol. i. 213 = Sieboldi.
meridionalis, Guss. Fl. Sic. Syn. ii. II. 831 = P. Aria.
Michauxii, Bosc, ex Poir. Encyc. Suppl. iv. 452 = amygdaliformis.
Michauxii, Hort. ex C. Koch, Dendrol. i. 216 = communis.
micrantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 351.—Japon.
microcarpa, DC. Prod. ii. 636; Spreng. Syst. ii. 511 = americana.
microcarpa, Wendl. ex C. Koch, Dendrol. i. 211 = baccata.
microphylla, *Wall. Cat.* n. 676.—Reg. Himal.
mitis, Syme, Engl. Bot. ed. III. iii. 256 = P. Malus.
montana, *Hort. ex C. Koch, Hort. Dendrol.* 183, in adnot.—Hab.?
Montis-Sinai, Hort. ex Steud. Nom. ed. I. 670 = amygdaliformis.

PYRUS :—
nebrodensis, Guss. Fl. Sic. Prod. i. 569 = Cotoneaster nebrodensis.
nepalensis, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. ii. 374 = P. Pashia.
nepalensis, Lodd. ex G. Don, in Loud. Hort. Brit. 202 = P. Aria.
Neumanniana, Tausch, in Flora, xxi. (1838) I. Beibl. 76 = Amelanchier canadensis?
nivalis, Jacq. Fl. Austr. ii. 4 = communis.
nivalis, Lindl. Bot. Reg. t. 1484 = amygdaliformis.
nivalis, Pall. ex Georgi, Beschr. Russ. Reich. iv. 1014 = elaeagrifolia.
Nussia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 237 = Stranvaesia glaucescens.
oblonga, Hort. ex Steud. Nom. ed. I. 670 = P. Cydonia.
oblongifolia, Spach, Hist. Vég. Phan. ii. 128 = amygdaliformis.
obtusata, *Steud. Nom.* ed. II. ii. 424.—Hab.?
odorata, Hort. ex C. Koch, Dendrol. i. 192 = P. Aria.
oleaefolia, Hort. l. c. 216 = communis.
olivaefolia, Hort. l. c. = communis.
orientalis, Hort. ex Nois. Cat. (1812); ex Thouin, in Mém. Mus. Par. i. (1815) 171 = amygdaliformis.
orientalis, Pall. Reise Suedl. Statth. Russ. Reich. ii. 88, 188, 222, 236, 453 (Quid?).—Rossia.
ovalis, Willd. Berl. Baumz. ed. I. 259 = Amelanchier canadensis.
paradisiaca, Hort. ex Steud. Nom. ed. I. 670 = P. Malus.
parasomila, Hort. l. c. 671 = P. Malus.
parviflora, Desf. in Ann. Mus. Par. xii. (1808) 52 = amygdaliformis.
Pashia, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 236.—Reg. Himal.
pensylvanica, Fisch. ex Steud. Nom. ed. I. 671 (Quid?).—Am. bor.
pensylvanica, Hort. ex Lindl. in Trans. Hort. Soc. vii. (1828) 231 = arbutifolia.
persica, Pers. Syn. ii. 40 = elaeagrifolia.
pinnatifida, *Ehrh. Plantag.* p. 22; *ex ej. Beitr.* vi. 93.—Europ. bor.
pinnatifida, Sm. Engl. Bot. t. 2331, partim = P. Aria.
Piraster, Borkh. Handb. Forstbot. ii. 1287 = communis.
pohuashanensis, Hance, in Journ. Bot. xiii. (1875) 132 = P. Aucuparia.
Pollveria, Lej. Fl. Spa, ii. 311; Rev. Fl. Spa, 94 = P. Malus.
Pollveria, *Linn. Mant.* ii. 244.—Europ.
Pollvilleriana, Borkh. Handb. Forstb. ii. 1251 = praec.
Pollweriana, Rev. Hortic. (1885) 416. f. 76, 78 = praec.
Pollwilleriana, Decne. Jard. Fruit. Mus. i. t. 31 = praec.
Pollwylleriana, Link, Handb. ii. 80; Wallr. Beitr. Fl. Hercyn. 205 = praec.
polonica, Hort. ex C. Koch, Dendrol. i. 208 = prunifolia.
polycarpa, *Hook. f. Fl. Brit. Ind.* ii. 378.—Reg. Himal.
pompejana, Hort. ex Steud. Nom. ed. I. 671 = communis.
praecox, Hort. ex Lindl. Bot. Reg. t. 1484 = amygdaliformis.
praecox, Pall. Fl. Ross. i. 22 = P. Malus.
praemorsa, Guss. Fl. Sic. Syn. i. 560 = P. Aucuparia.
prasomila, Hort. ex Steud. Nom. ed. I. 670, in syn. = P. Malus.
prunifolia, Hort. ex Steud. l. c. ed. II. ii. 424 = arbutifolia.
prunifolia, *Willd. Phyt.* 8.—Sibir.
pubens, Lindl. in Trans. Hort. Soc. vii. (1828) 232 = arbutifolia.
pulverulenta, Hort. ex C. Koch, Dendrol. i. 208 = P. Malus.
pumila, Hort. l. c. 203 = P. Malus.
pumila, Neum. ex Tausch, in Flora, xxi. (1838) I. Beibl. 77 = arbutifolia.
pyraina, Rafin. ex Decne. Jard. Fruit. Mus. i. sub t. 13 = amygdaliformis.
pyramidalis, Hort. ex Steud. Nom. ed. I. 671 = communis.
Pyraster, Medic. Gesch. 87; Borck. in Roem. Arch. i. III. (1798) 90 = communis.
quinqueflora, *Buch.-Ham. ex G. Don, Gen. Syst.* ii. 647.—Ind. or.

PYRUS :—
rhamnoides, *Hook. f. Fl. Brit. Ind.* ii. 377.—Reg. Himal.
Ringo, *Wenzig, in Linnaea*, xxxviii. (1874) 37.—Japon.
Riversii, Hort. ex C. Koch, Dendrol. i. 209 = spectabilis.
rivularis, *Dougl. in Hook. Fl. Bor. Am.* i. 203. t. 68.—Am. bor. occ.
rotundifolia, Auct. ex Steud. Nom. ed. I. 671 = Amelanchier vulgaris.
rotundifolia, *Moench, Meth.* 680; *Bechstt. Forstbot.* ed. V. 152.—Europ.
rubelliana, Hort. ex Steud. Nom. ed. I. 671 = P. Malus.
rubicunda, *Hoffmgg. Verz. Pfl.* 192.—Sibir.
rufescens, Hort. ex Steud. Nom. ed. I. 671 = communis.
rupicola, Syme, Engl. Bot. ed. III. iii. 244 = P. Aria.
salicifolia, Balb. in Mém. Acad. Turin, 1805–8 (1809), 216; Sibth. & Sm. Fl. Graec. Prod. i. 344 (1809) = amygdaliformis.
salicifolia, Habl. Taur. 107; ex Bieb. Fl. Taur. Cauc. i. 390 = elaeagrifolia.
salicifolia, *Pall. Reise*, iii. 734.—Reg. Caucas.
salviaefolia, Petter, Bot. Wegw. Spalato, 103 = amygdaliformis.
salvifolia, *DC. Prod.* ii. 634.—Gallia.
sambucifolia, *Cham. & Schlecht. in Linnaea*, ii. (1827) 36.—As. bor. or.; Am. bor.
sanguinea, Pursh, Fl. Am. Sept. i. 340 = Amelanchier alnifolia, canadensis.
sanguisorbifolia, Stokes, Bot. Mat. Med. iii. 120 = Amelanchier vulgaris.
saracenia, Hort. ex C. Koch, Dendrol. i. 219 = persica.
sativa, C. Koch, l. c. 215 = communis.
sativa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 (Quid?).—Java.
saxatilis, Schlecht. ex Ledeb. Fl. Ross. ii. 97 = Sieversii.
scabiosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 23 = Pygeum parviflorum?
scandica, Aschers. Fl. Brandenb. 207 = intermedia.
scandica, Bab. Man. Brit. Bot. ed. III. 111 = rotundifolia.
semilobata, Bechst. Forstbot. ed. V. 152 = rotundifolia.
semipinnata, Bechst. l. c. 153 = intermedia.
semipinnata, *Roth, Enum. Pl. Phan. Germ.* ii. 438.—Europ.
sempervirens, Willd. Enum. Hort. Berol. Suppl. 35 = angustifolia.
septiloba, Stokes, Bot. Mat. Med. iii. 121 = torminalis.
sibirica, Hort. ex C. Koch, Dendrol. i. 208 = prunifolia.
sicula, Tineo, Cat. Hort. Panorm. (1827) 208 = amygdaliformis.
Sieboldi, *Regel, Ind. Sem. Hort. Petrop.* (1858) 51.—Japon.
Sieversii, Hort. ex C. Koch, Dendrol. i. 208 = prunifolia.
Sieversii, *Ledeb. Fl. Alt.* ii. 222.—Sibir. altaic.; Reg. Himal.
sikkimensis, *Hook. f. Fl. Brit. Ind.* ii. 373.—Reg. Himal.
sikkimensis, Hook. f. l. c. 377, 378, in syn. = granulosa, Griffithii.
Simonii, *Carr. in Rev. Hortic.* (1872) 28.—China.
Sinai, Desf. Hist. Arb. ii. 144 = amygdaliformis.
sinaica, Dum.-Cours. Bot. Cult. ed. II. v. 435; Thouin, in Mém. Mus. Par. i. (1815) 170. t. 9 = amygdaliformis.
sinensis, Dum.-Cours. l. c. 428 = spectabilis.
sinensis, Hort. ex C. Koch, Dendrol. i. 215 = communis.
sinensis, *Lindl. in Trans. Hort. Soc.* vi. (1826) 396.—China.
sinensis, *Poir. Encyc. Suppl.* iv. 452 (sp. div.).—China.
sinica, Royle, Illustr. Bot. Himal. 206 = sinensis, *Lindl.*
slavonica, *Kit. in Linnaea*, xxxii. (1863) 583.—Europ. austr.
sorbifolia, Bosc, ex Wats. Dendrol. i. t. 53 = pinnatifida.
sorbifolia, *Cham. ex Spreng. Syst.* iv. *Cur. Post.* 343.—Kamtschat.
Sorbus, *Gaertn. Fruct.* ii. 45. t. 87.—Europ.; Afr. bor.

PYRUS:—
spectabilis, *Ait. Hort. Kew.* ed. I. ii. 175.—China.
sphaerocarpa, Wender. Ind. Sem. Hort. Marb. (1835). —Cf. Linnaea, xi. (1837) Litt. 92 = baccata.
spinosa, *Forsk. Fl. Aegypt. Arab.* 211.—Anatolia.
spuria, *DC. Prod.* ii. 637.—Hab.?
stipulacea, Hort. ex Baxt. in Loud. Hort. Brit. Suppl. ii. 662 (Quid?).—Reg. Himal.
suaveolens, Wender. in Linnaea, v. (1830) Litt. 55 = coronaria.
subcordata, Ledeb. Fl. Ross. ii. 95 = rivularis.
sudetica, *Tausch, in Flora*, xvii. (1834) 76.—Suecia.
subfusca, *Regel, in Act. Hort. Petrop.* i. (1870) 126.—Rossia.
suecica, Roehl. ex Steud. Nom. ed. I. 671; Garcke, Fl. Deutschl. ed. IX. 140 = intermedia.
Sumboshia, Spreng. Syst. iv. Cur. Post. 196 = P. Cydonia.
sylvestris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 562 = P. Malus.
sylvestris, Moench, Meth. 679 = communis.
syriaca, *Boiss. Diagn.* Ser. I. x. 1.—As. Min.; Syria.
syrmiensis, *Kit. in Linnaea*, xxxii. (1863) 581.—Europ. austr.
Tardevieta, Lamotte, Prod. Fl. Centr. Fr. i. 280 = communis.
tenuifolia, Stokes, Bot. Mat. Med. iii. 120 = Amelanchier canadensis.
Thomsoni, *King, ex Hook. f. Fl. Brit. Ind.* ii. 379.—Reg. Himal.
thuringiaca, Ilse, ex Nym. Consp. 241 = pinnatifida.
tianschanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 287.—Turkestan.
Tinei, Todaro, ex Decne. Jard. Fruit. Mus. i. sub t. 13 = amygdaliformis.
tomentosa, *DC. Prod.* ii. 637.—Sibir.
tomentosa, Moench, Meth. 680 = P. Pollveria.
tomentosa, *Roxb. Fl. Ind.* ii. 512.—Persia.
Toringo, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 41 = Sieboldi.
torminalis, *Ehrh. Beitr.* vi. 92.—Europ.
trilobata, *DC. Prod.* ii. 636.—Syria.
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 169.—Japon.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 288.—Turkestan.
umbellata, *Dum.-Cours. Bot. Cult.* ed. II. v. 429.—China.
upsaliensis, Hort. Lodd. ex G. Don, in Loud. Hort. Brit. 201 = P. Malus.
ursina, Herb. Strach. & Winterb. ex Hook. f. Fl. Brit. Ind. ii. 376 = P. Aucuparia.
ursina, Wall. Cat. n. 675 = foliolosa.
ussuriensis, Maxim. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 132, 237; et in Prim. Fl. Amur. 102 = sinensis, *Lindl.*
Valerna, Hort. ex Steud. Nom. ed. I. 671 = communis.
variolosa, Wall. Cat. n. 680 = P. Pashia.
verruculosa, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iv. (1864) 312 = P. Pashia.
vestita, *Wall. Cat.* n. 679.—Reg. Himal.
Wallichii, *Hook. f. Fl. Brit. Ind.* ii. 376.—Reg. Himal.
Wangenheimiana, Tausch, in Flora, xxi. (1838) 714 = Amelanchier canadensis.

PYTHAGOREA, Lour. Fl. Cochinch. 243 (1790) = **Homalium**, Jacq. (Samyd.).
africana, E. Mey. in Drège, Zwei Pfl. Docum. 214 = H. africanum.
cochinchinensis, Lour. Fl. Cochinch. 243 = H. fagifolium.
rufescens, E. Mey. ex Presl, Bot. Bemerk. 42 = H. rufescens.

PYTHAGOREA, Rafin. in Journ. Phys. lxxxix. (1819) 96 = **Lythrum**, Linn.
alata, Rafin. l. c. = L. alatum.
linearis, Rafin. l. c. = L. lineare.
virgata, Rafin. l. c. = L. virgatum.

PYTHION, Mart. in Flora, xiv. (1831) 459 = **Amorphophallus**, Blume (Aroid.).
campanulatum, Mart. l. c. = A. campanulatus.

PYTHIUS, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
dulcis, Rafin. l. c. = E. dulcis.
epithymoides, Rafin. l. c. = E. epithymoides.
fuscata, Rafin. l. c. = E. Pithyum.
obtusata, Rafin. l. c. = E. obtusata.

PYTHONIUM, Schott, Melet. i. 17 (1832) = **Thomsonia**, Wall. (Aroid.).
bulbiferum, Schott, l. c. 18 = Amorphophallus bulbifer.
guineense, G. Don, in Sweet, Hort. Brit. ed. III. 631, nomen.—Afr. trop.
Hookeri, Kunth, Enum. Pl. iii. 31 = Anchomanes Hookeri.
sylvaticum, Wight, Ic. t. 802 = Synantherias sylvatica.
Wallichianum, Schott, Melet. i. 17 = T. napalensis.

PYXA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Costus**, Linn. (Scitam.).

PYXIDANTHERA, Michx. Fl. Bor. Am. i. 152. t. 17 (1803). *DIAPENSIACEAE*, Benth. & Hook. f. ii. 619.
barbulata, *Michx. l. c.*—Am. bor.
granulata, Pritz. Ic. Ind. i. 913, sphalm. = praec.
spathulata, Muhl. Cat. 25 = Lepuropetalum spathulatum.

PYXIDANTHUS, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 150. t. 6 = **Blakea**, P. Br. (Melastom.).
latifolius, Naud. l. c. 152 = B. Pyxidanthus.
Lindenianus, Naud. l. c. 151 = B. Lindeniana.
Schlimii, Naud. l. c. = B. Schlimii.

PYXIDIUM, Moq. in DC. Prod. xiii. II. 262 (1849) (vox fructus desc. lapsu pro genere habita) = **Amaranthus**, Tourn.
gangeticum, Moq. l. c. 261 = A. gangeticus.
graecizans, Moq. l. c. 263 = A. Blitum.
melancholicum, Moq. l. c. 262 = A. gangeticus.
oleraceum, Moq. l. c. 273 = A. oleraceus, lividus.
viride, Moq. l. c. 274 = A. viridis.

PYXIPOMA, Fenzl, in Ann. Wien. Mus. ii. (1839) 293 = **Sesuvium**, Linn. (Ficoid.).
polyandrum, Fenzl, l. c. = S. Portulacastrum.

Q

QUADRANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 (Quid?).

QUADRELLA, J. S. Presl, Rostl. ii. 260 (1825) = **Capparis**, Tourn.
Breynia, J. S. Presl, l. c. 261 = C. Breynia.
crotonoides, J. S. Presl, l. c. 260 = C. crotonoides.
furfuracea, J. S. Presl, l. c. = C. furfuracea.
intermedia, J. S. Presl, l. c. 261 = C. odoratissima.
jamaicensis, J. S. Presl, l. c. = C. jamaicensis.
sidaefolia, J. S. Presl, l. c. 260 = C. crotonoides.
torulosa, J. S. Presl, l. c. 261 = C. jamaicensis.

QUADRIA, Ruiz & Pav. Prod. 16. t. 33 (1794) = **Gevuina**, Molina (Proteac.).
Avellana, Gaertn. f. Fruct. iii. 220 = G. Avellana.
heterophylla, Ruiz & Pav. Fl. Per. i. 63. t. 99 = G. Avellana.

QUADRIALA, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 194 (Fam. Nat. Fl. Jap. i. 86. t. 2) = **Buckleya**, Torr. (Santal.).
lanceolata, Sieb. & Zucc. ll. cc. 195 (87) = B. lanceolata (in Addend.).

QUADRICOSTA, Dulac, Fl. Hautes-Pyr. 329 (1867) = **Ludwigia**, Linn. (Onagrar.).
palustris, Dulac, l. c. = L. palustris.

QUADRIFARIA, Manetti, ex Gord. Pinet. Suppl. 14 (1862) = **Araucaria**, Juss. (Conif.).
imbricata, Manetti, ex Gord. l. c. = A. imbricata.

QUAIACUM, Scop. Introd. 212 (1777), sphalm. = **Guaiacum**, Plum. (Zygophyll.).

QUALEA, Aubl. Pl. Gui. i. 5. t. 1, 2 (1775). *VOCHYSIACEAE*, Benth. & Hook. f. i. 976.
AGARDHIA, Spreng. Syst. i. 4 (1825).
AMPHILOCHIA, Mart. Nov. Gen. et Sp. i. 127. t. 77 (1824).
LOZANIA, Mutis, ex Caldas, in Seman. Nuev. Gran. (1810) III. 20.
SCHUECHIA, Endl. Gen. 1178 (1840).
acuminata, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 40.—Bras.
albiflora, *Warm. l. c.* 36.—Am. trop.
calcarata, Link, Jahrb. i. III. (1820) 24 = Erisma violaceum.
cassiquiarensis, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 34.—Bras.
coerulea, *Aubl. Pl. Gui.* i. 7. t. 2.—Guian.
cordata, *Spreng. Syst.* i. 17.—Bras.
cryptantha, *Spreng. ex Warm. in Mart. Fl. Bras.* xiii. II. 53.—Bras.
densiflora, *Warm. l. c.* 50.—Am. trop.
dichotoma, *Warm. ex Wille, in Overs. Danske Vidensk. Selsk. Forh.* (1882) 181.—Am. trop.
disticha, Spreng. Syst. iv. Cur. Post. 10 = Callisthene major.
ecalcarata, *Mart. Nov. Gen. et Sp.* iii. 130. t. 78.—Bras.
elongata, *Warm. in Mart. Fl. Bras.* xiii. II. 48.—Am. trop.
fasciculata, Spreng. Syst. iv. Cur. Post. 10 = Callisthene fasciculata.
Gardneriana, *Warm. in Mart. Fl. Bras.* xiii. II. 35.—Am. trop.
Gestasiana, *A. St. Hil. in Mém. Mus. Par.* vi. (1820) 254.—Bras.
glauca, *Warm. in Mart. Fl. Bras.* xiii. II. 49.—Am. trop.
Glaziovii, *Warm. l. c.* 53.—Am. trop.
grandiflora, *Mart. Nov. Gen. et Sp.* iii. 133. t. 79.—Bras.
ingens, *Warm. in Mart. Fl. Bras.* xiii. II. 40.—Bras.
intermedia, *Warm. l. c.* 50.—Am. trop.
Jundiahy, *Warm. in Kjoeb. Vidensk. Meddel.* (1867) 31.—Bras.
laurifolia, Spruce, ex Warm. in Mart. Fl. Bras. xiii. II. 109 = Erisma laurifolium.
Lundii, *Warm. l. c.* 47.—Am. trop.
lutea, Desf. ex Link, Jahrb. i. III. (1820) 24 = Erisma nitidum.
macropetala, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 41.—Bras.
macroptera, *Link, ex A. Dietr. Sp. Pl.* i. 101.—Bras.
maliformis, Link, ex Warm. in Mart. Fl. Bras. xiii. II. 24 = Callisthene fasciculata.
marginata, *Warm. l. c.* 52.—Am. trop.
microphylla, *Warm. l. c.* 37.—Am. trop.
minor, Spreng. Syst. iv. Cur. Post. 10 = Callisthene minor.
Muelleriana, *Schomb. Reise in Brit. Guian.* iii. (1848) 1099.—Guian.
multiflora, *Mart. Nov. Gen. et Sp.* i. 134. t. 80.—Bras.
parviflora, *Mart. l. c.* 135. t. 81.—Bras.
Perrini, *Spreng. Syst.* i. 17.—Guian.
pilosa, *Warm. in Mart. Fl. Bras.* xiii. II. 45.—Am. trop.
psidiifolia, *Spruce, ex Warm. l. c.* 46.—Am. trop.
pulcherrima, *Spruce. ex Warm. l. c.* 37.—Am. trop.
retusa, *Spruce, ex Warm. l. c.* 34.—Bras.
rosea, *Aubl. Pl. Gui.* i. 5. t. 1.—Guian.
Schomburgkiana, *Warm. in Mart. Fl. Bras.* xiii. II. 39.—Guian.
Selloi, *Warm. l. c.* 49.—Am. trop.
Sprucei, *Warm. l. c.* 38.—Am. trop.
subvillosa, *Link, ex A. Dietr. Sp. Pl.* i. 101.—Bras.
trichanthera, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 35.—Bras.
verticillata, Spreng. Syst. i. 17 = Vochysia verticillata.
violacea, Mart. & Zucc. ex Schult. Mant. i. Add. II. 99 = rosea.

QUAMASIA, Rafin. in Am. Monthly Mag. (1818) 265 = **Camassia**, Lindl. (Liliac.).
esculenta, Rafin. l. c. = C. esculenta.
scilloides, Rafin. Cat. 15 = C. esculenta.

QUAMASSIA, Rafin. Fl. Tellur. ii. 25 (1836) = praec.

QUAMOCLIDION, Choisy, in DC. Prod. xiii. II. 429 (1849) = **Mirabilis**, Riv. (Nyctag.)
angulatum, Choisy, l. c.—Mexic.
multiflorum, Torr. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 321 = M. multiflora.
nyctagineum, Choisy, in DC. Prod. xiii. II. 429 = M. triflora.
oxybaphoides, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 320 = Oxybaphus Wrightii.

QUAMOCLIT, Tourn. ex Moench, Meth. 453 (1794) = **Ipomoea**, Linn. (Convolv.).
acuta, Gard. Chron. (1880) I. 588, sphalm. = Quercus acuta.
acutangula, Choisy, in DC. Prod. ix. 335 = I. coccinea.
angulata, Boj. Hort. Maurit. 224 = I. angulata.
cholulensis, G. Don, Gen. Syst. iv. 259 = I. coccinea.
coccinea, Moench, Meth. 453 = I. coccinea.
dichotoma, G. Don, Gen. Syst. iv. 259 = I. coccinea.
digitata, G. Don, l. c. 260 = I. digitata, *Linn.*
eustachiana, G. Don, l. c. 259 = I. triloba.
globosa, Benth. Pl. Hartw. 89 = I. Hartwegi.
globosa, G. Don, Gen. Syst. iv. 259 = I. peduncularis.
grandiflora, G. Don, l. c. = I. Funis.
hastigera, G. Don, l. c. = I. hederifolia.
hederifolia, G. Don, l. c. = I. hederifolia.
Kerberi, Fourn. in Bull. Soc. Bot. Fr. xxx. (1883) 187. —Mexic.
leucantha, G. Don, Gen. Syst. iv. 258 = I. leucantha.
longiflora, G. Don, l. c. 259 = I. Bona-nox.
luteola, G. Don, l. c. 258 = I. coccinea.
Mina, G. Don, l. c. 259 = I. versicolor.
mutica, Choisy, in DC. Prod. ix. 335 = I. serotina.
Nationis, Hook. Bot. Mag. t. 5432 = I. Nationis.
pallescens, Ann. Flor. & Pom. (1839) 85 = I. hederifolia.
pedata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 270 = I. armata.
pennata, Voigt, Hort. Suburb. Calc. 353 = I. Quamoclit.
phoenicea, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 433 = I. coccinea.
phoenicea, Wight, ex Meissn. in Mart. Fl. Bras. vii. 218 = I. hederifolia.
pinnata, Boj. Hort. Maurit. 224 = I. Quamoclit.
Ruiziana, G. Don, Gen. Syst. iv. 258 = I. coccinea.
ruselliaeflora, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 271 = I. hederifolia.
sagittaefolia, Choisy, in DC. Prod. ix. 335 = I. sagittaefolia.
sanguinea, G. Don, Gen. Syst. iv. 259 = I. hederifolia.
serotina, G. Don, l. c. = I. serotina.
solanifolia, Choisy, in DC. Prod. ix. 335 = I. solanifolia.
triloba, G. Don, Gen. Syst. iv. 259 = I. triloba.
tubulosa, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 270 = I. tubulosa.
vitifolia, G. Don, Gen. Syst. iv. 259 = I. peduncularis.
vulgaris, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 434 = I. Quamoclit.

QUAMOCLITA, Rafin. Fl. Tellur. iv. 74 (1836) = QUAMOCLIT, Tourn. = **Ipomoea**, Linn. (Convolv.).
bipinnata, Rafin. l. c. (Quid?).—Hab.?
coccinea, Rafin. l. c. = I. coccinea.
cordata, Rafin. l. c. (Quid?).—Hab.?
denticulata, Rafin. l. c. = I. denticulata.
hybrida, Rafin. l. c. (Quid?).—Hab.?
lacunosa, Rafin. l. c. = I. lacunosa.
litoralis, Rafin. l. c. = I. littoralis.
multifida, Rafin. New Fl. Am. iv. 57 (Quid?).—Am. bor.
rubra, Rafin. Fl. Tellur. iv. 74 = Gilia coronopifolia.
solanifolia, Rafin. l. c. = I. solanifolia.

QUAMOCLITIA, Lowe, Man. Fl. Mad. ii. 51 (1868) = QUAMOCLIT, Tourn. = **Ipomoea**, Linn. (Convolv.).

QUAPIRA, Auct. ex Pfeiff. Nom. ii. 900 (1874) = **Guapira**, Aubl. (Inc. Sed.).

QUAPOJA, Batsch, Tab. Aff. 89 (1800) = seq.

QUAPOYA, Aubl. Pl. Gui. ii. 897 (1775) = **Clusia**, Linn. (Guttif.).
acuminata, Walp. Rep. i. 393 = C. acuminata, *Spreng.*
comans, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 239 = Renggeria comata.
laxiflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 11 = C. laxiflora.
littoralis, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 239 = Renggeria littoralis.
Pana-panari, Aubl. Pl. Gui. ii. 900. t. 344 = C. macrocarpa.
Pana-Pary, Steud. Nom. ed. II. ii. 425 = praec.
scandens, Aubl. Pl. Gui. ii. 898 = C. microcarpa.
sulphurea, Poepp. & Endl. Nov. Gen. et Sp. iii. 12 = C. sulphurea.
surinamensis, Miq. Stirp. Surin. Sel. 92 = C. macrocarpa.

QUAQUA, N. E. Br. in Gard. Chron. (1879) II. 8. *ASCLEPIADEAE.*
hottentotorum, *N. E. Br. l. c.*—Afr. austr.

QUARARIBEA, Aubl. Pl. Gui. ii. 691. t. 278 (1775). *MALVACEAE*, Benth. & Hook. f. i. 212.
GERBERIA, Scop. Introd. 286 (1777).
guianensis, *Aubl. Pl. Gui.* ii. 692. t. 278.—Bras.; Guian.
Martini, *Baill. Adansonia*, x. (1872) 180.—Guian.
pterocalyx, *Hemsl. Diagn. Pl. Nov.* 4.—Panama.
turbinata, *Poir. Encyc. Suppl.* iv. 636.—Bras.

QUARENA, Rafin. Sylva Tellur. 39 (1838) = **Cordia**, Linn. (Borag.).
indica, Rafin. l. c. = C. Myxa.
sinensis, Rafin. l. c. = C. sinensis.
spinescens, Rafin. l. c. = C. spinescens.

QUARTINIA, Endl. Gen. Suppl. ii. 94 (1842). *LYTHRARIEAE*, Benth. & Hook. f. i. 777.
RHYACOPHILA, Hochst. in Flora, xxiv. (1841) 659.
repens, *Endl. ex Walp. Rep.* v. 673.—Abyss.
turfosa, A. Rich. Tent. Fl. Abyss. i. 277. t. 51 = repens.

QUARTINIA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 259 = **Pterolobium**, R. Br. (Legumin.).
abyssinica, A. Rich. l. c. 260 = P. lacerans.

QUASSIA, Linn. Sp. Pl. ed. II. 553 (1762). *SIMARUBEAE*, Benth. & Hook. f. i. 308.
africana, *Baill. Adansonia*, viii. (1867–68) 89.—Afr. trop.
alatifolia, Stokes, Bot. Mat. Med. ii. 491 = amara.
amara, *Linn. Sp. Pl.* ed. II. 553.—Guian.
Aruba, D. Dietr. Syn. Pl. ii. 1416 = Simaba Aruba.
Cedron, Baill. Hist. Pl. iv. 406 = Simaba Cedron.
crocea, Vahl, Eclog. Am. iii. 12 = amara.
dioica, O. Berg, Mat. Med. 355 = Simaruba glauca.
excelsa, Sw. Prod. Veg. Ind. Occ. 67 = Simaruba excelsa.
ferruginea, D. Dietr. Syn. Pl. ii. 1416 = Simaba ferruginea.
floribunda, D. Dietr. l. c. = Simaba floribunda.
glauca, Spreng. Syst. ii. 319 = Simaruba glauca.
guianensis, D. Dietr. Syn. Pl. ii. 1416 = amara.
officinalis, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guian.
orinocensis, D. Dietr. Syn. Pl. ii. 1416 = Simaba orinoccensis.
pentandra, Stokes, Bot. Mat. Med. ii. 489 = Simaruba excelsa.
polygama, Lindsay, in Trans. Roy. Soc. Edinb. iii. (1794) 205. t. [6] = Simaruba excelsa.
Simaruba, Blanco, Fl. Filip. ed. II. 247 = Cupania glabrata.
Simaruba, Linn. f. Suppl. 234 = Simaruba amara.
suaveolens, D. Dietr. Syn. Pl. ii. 1416 = Simaba suaveolens.
tricarpa, Blanco, Fl. Filip. ed. I. 351 = Cupania pleuropteris.
trichilioides, D. Dietr. Syn. Pl. ii. 1416 = Simaba trichilioides.
undulata, D. Dietr. l. c. = Zwingera undulata.
versicolor, Spreng. Syst. iv. Cur. Post. 163 = Simaruba versicolor.

QUATERNELLA, Ehrh. Beitr. iv. 149 (1789) = **Cerastium**, Dill. (Caryoph.).

QUEBITEA, Aubl. Pl. Gui. ii. 838. t. 327 (1775) = **Piper**, Linn.
guianensis, Aubl. l. c. (Quid?).—Guian.

QUEBRACHIA, Griseb. in Goett. Abh. xix. (1874) 51. 114; xxiv. (1879) 95. *ANACARDIACEAE.*
Lorentzii, *Griseb. l. c.*—Reg. Argent.

QUEKETTIA, Lindl. Bot. Reg. (1839) Misc. 3. *ORCHIDEAE*, Benth. & Hook. f. iii. 569.
microscopica, *Lindl. l. c.*—Bras.

QUELTIA, Salisb. in Trans. Hort. Soc. i. (1812) 351 = **Narcissus**, Tourn. (Amaryll.).
alba, Haw. Suppl. Pl. Succ. 127 = N. incomparabilis.
ampla, Salisb. in Trans. Hort. Soc. i. (1812) 351 = N. incomparabilis.
apodantha, Kunth, Enum. Pl. v. 856 = N. juncifolius.
Assoana, Kunth, l. c. 730 = N. juncifolius.
aurantia, Haw. Suppl. Pl. Succ. 123 = N. incomparabilis.
aurantiaca, Salisb. Gen. Pl. Fragm. 100, sphalm. = praec.
Bernardi, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 164 = N. Macleayi.
biternaceus, Salisb. Gen. Pl. Fragm. 100 = N. incomparabilis?
capax, Salisb. in Trans. Hort. Soc. i. (1812) 353 = N. capax.
cernua, M. Roem. Syn. Ensat. 203 = N. cernuus.
concolor, Haw. Suppl. Pl. Succ. 126 = N. incomparabilis.
Coornei, M. Roem. Syn. Ensat. 204 = N. triandrus.
coronata, M. Roem. l. c. 203 = N. triandrus.
foetida, Herb. Amaryll. 311 = N. incomparabilis.
galanthifolia, Haw. Suppl. Pl. Succ. 128 = N. galanthifolius.
Gouani, Haw. Monog. Narciss. 5. n. 2 = N. incomparabilis.
incomparabilis, Haw. Suppl. Pl. Succ. 125 = N. incomparabilis.
Jonquilla, Herb. Amaryll. 315. t. 38. f. 24 = N. Jonquilla.
juncifolia, Herb. l. c. 314. t. 43. f. 1 = N. juncifolius.
Maclayi, M. Roem. Syn. Ensat. 204 = N. Macleayi.
Macleana, Herb. Amaryll. 310. t. 39. f. 1 = N. Macleayi.
montana, Herb. l. c. f. 4 = N. poculiformis.
montana, Rafin. Fl. Tellur. iv. 123 = praec.?
nivea, Haw. Monog. Narciss. 6. n. 7 = N. incomparabilis.
odora, Herb. Amaryll. 313 = N. odorus.
orientalis, Herb. l. c. 312. t. 38. f. 28 = N. Tazetta.
pallida, Salisb. Gen. Pl. Fragm. 100 = N. Tazetta.
Parkinsonii, M. Roem. Syn. Ensat. 205 = N. Macleayi.
poculiformis, Haw. Suppl. Pl. Succ. 127 = N. poculiformis.
poculigera, Salisb. in Trans. Hort. Soc. i. (1812) 352 = N. poculiformis.
pumila, Herb. Amaryll. 316. t. 41. f. 26 = N. pumilus.
pusilla, Herb. l. c. 315. t. 43. f. 2 = N. juncifolius.
Sabini, M. Roem. Syn. Ensat. 205 = N. Macleayi.
semipartita, Haw. Suppl. Pl. Succ. 126 = N. incomparabilis.
triandra, M. Roem. Syn. Ensat. 203 = N. triandrus.

QUELUSIA, Vand. Fl. Lusit. 23. t. 10. f. 10 (1788); et in Roem. Script. Hisp. Lusit. 101 (1796) = **Fuchsia**, Linn. (Onagrar.).
regia, Vand. ex Vell. Fl. Flum. 149 (1825); et iv. t. 6 (1827) (Quid?).—Bras.

QUERCUS, [Tourn.] Linn. Syst. ed. I (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 407.
ARCAULA, Rafin. Alsog. Am. 30 (1838).
BALANAULAX, Rafin. l. c. 28 (1838).
BALANOPLIS, Rafin. l. c. 29 (1838).
CERRIS, Rafin. l. c. (1838).
CYCLOBALANOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 77.
CYCLOBALANUS, Oerst. l. c. (1866) 80.
DRYOPSILA, Rafin. Alsog. Am. 25 (1838).
ERIODRYS, Rafin. l. c. 30 (1838).
LITHOCARPUS, Blume, Bijdr. 526 (1825).
PASANIA, Oerst. in Kjoeb. Vidensk. Meddel. (1866) 81.
PERYTIS, Rafin. Alsog. Am. 29 (1838).
SCOLODRYS, Rafin. l. c. (1838).
SYNAEDRYS, Lindl. Introd. Nat. Syst. ed. II. 441 (1836).

QUERCUS:—
Abelicea, Lam. Encyc. i. 725 = Zelkova cretica.
abietum, Kotschy, ex A. DC. in DC. Prod. xvi. II. 8 = Q. Robur.
acerifolia, Hort. ex Petz. & Kirchn. Arb. Muscav. 652 = rubra.
acroglandis, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 23 = agrifolia.
acuminata, Hort. ex Endl. Gen. Suppl. iv. II. 28 = annulata.
acuminata, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 217 = calophylla.
acuminata, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 636. —Ind. or.
acuminatissima, A. DC. in DC. Prod. xvi. II. 102 = Junghuhnii.
acuta, Buch.-Ham. ex Wall. Cat. n. 3735 = Castanopsis tribuloides.
acuta, Rafin. Alsog. Am. 22 (Quid ?).—Am. bor.
acuta, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 290 = thalassica.
acuta, *Thunb. Fl. Jap.* 175.—Japon.
acutidens, Torr. Bot. Mex. Bound. 207 = dumosa.
acutifolia, *Née, in Anal. Cienc. Nat.* iii. (1801) 267.—Mexic.
acutiglandis, Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 146 = agrifolia.
acutissima, *Carruth. in Journ. Linn. Soc.* vi. (1862) 33.—China.
aegilopifolia, Boiss. ex Endl. Gen. Suppl. iv. II. 25 = lusitanica.
Aegilops, Griseb. Spicil. Fl. Rumel. ii. 44 = macedonica.
Aegilops, *Linn. Sp. Pl.* 996.—Europ. austr.; Oriens.
Aegilops, Mill. Gard. Dict. ed. VIII. n. 7 = lanuginosa.
Aegilops, Scop. Fl. Carn. ed. II. ii. 241 = Q. Cerris.
aegylopifolia, Boiss. ex A. DC. in DC. Prod. xvi. II. 17 = lusitanica.
aegylopifolia, Pers. Syn. ii. 570 = Q. Pseudo-suber.
Aesculus, Boiss. Fl. Orient. iv. 1164 = Q. Robur.
aestivalis, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 385, in textu = Q. Robur.
Afares, Pomel, Nouv. Mat. Fl. Atl. 391 = castaneaefolia.
affinis, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 222 = obtusata.
affinis, [Scheidw.] in Hortic. Belg. iv. (1837) 321. t. 17 = nitens.
afghanistanensis, Hort. ex C. Koch, Dendrol. ii. II. 33 = Q. Robur.
africana, Barrow, Trav. S. Afr. i. 83 = Ocotea bullata.
agnostifolia, Hort. ex C. Koch, l. c. 62 = nigra.
agrifolia, *Née, in Anal. Cienc. Nat.* iii. (1801) 271.—Calif.
ajudaghiensis, *Stev. ex Petz. & Kirchn. Arb. Muscav.* 614.—Rossia.
Alams, Benth. Pl. Hartw. 55 = calophylla.
alba, *Linn. Sp. Pl.* 996.—Am. bor.
aliena, *Blume, Mus. Bot. Lugd. Bat.* i. 298.—China; Japon.
almaguerensis, *Humb. & Bonpl. Pl. Aequin.* ii. 157. t. 131.—N. Granat.
alnifolia, *Poech, Enum. Pl. Ins. Cyp.* 12.—Ins. Cyprus.
alpestris, Boiss. Elench. 83 = lusitanica.
alpina, Endl. Gen. Suppl. iv. II. 25, sphalm. = *Alzina*.
altissima, Hort. ex Petz. & Kirchn. Arb. Muscav. 634 = Q. Robur.
Alzina, Lapeyr. Hist. Arbr. Pl. Pyr. ii. 584 = Q. Ilex.
ambigua, Humb. & Bonpl. Pl. Aequin. ii. 51. t. 93 = obtusata.
ambigua, *Kit. ex Rochel, Pl. Banat. Rar.* 26, nomen; *et in Linnaea*, xxxii. (1863) 354.—Europ.
ambigua, Michx. f. Hist. Arb. Am. ii. 120. t. 24 = rubra.
Amherstiana, *Wall. Cat.* n. 2783.—Burma.
amplifolia, Guss. Fl. Sic. Syn. ii. 607 = Q. Robur.
anceps, Korth. Verh. Nat. Gesch. Bot. 204 = spicata.
Andersoni, Hook. f. ex C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 127 = pachyphylla.
angulizans, Rafin. Alsog. Am. 22 (Quid ?).—Am. bor.
angustata, Blume, in Verh. Batav. Gen. ix. (1825) 212; Fl. Jav. Cup. 15. t. 72. t. 7 = Q. Pseudo-molucca.
Annenkowi, *Kauffm. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 168.—Rossia.
annulata, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 445 = Durandii.

QUERCUS:—
annulata, Hook. f. ex A. DC. in DC. Prod. xvi. II. 99 = semiserrata.
annulata, Korth. Verh. Nat. Gesch. Bot. 213 = Teysmannii.
annulata, Sm. in Rees, Cycl. xxix. n. 22 = glauca.
apennina, Lam. Encyc. i. 725 = Q. Robur.
aprica, Rafin. Alsog. Am. 26 (Quid ?).—Am. bor.
aquatica, *Walt. Fl. Carol.* 234.—Am. bor.
aquifolia, Kotschy, ex A. DC. in DC. Prod. xvi. II. 108 = coccifera.
Arcaula, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 56 = spicata.
arcoglandis, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 23, sphalm. = *acroglandis*.
Arcola, Buch.-Ham. ex Wall. Cat. p. 159. n. 2781 = *Arcaula*.
arcuata, Kotschy, ex A. DC. in DC. Prod. xvi. II. 56 = coccifera.
argentata, *Korth. Verh. Nat. Gesch. Bot.* 215. t. 47. f. 1–17.—Sumatra.
argyrophylla, Wall. Cat. n. 2788 = Castanopsis argentea.
aristata, *Hook. & Arn. Bot. Beech. Voy.* 444.—Mexic.
armata, D. Don, Prod. Fl. Nep. 56 = Castanopsis tribuloides.
armata, Roxb. Hort. Beng. 104; Pl. Corom. iii. 92. t. 296; Fl. Ind. iii. 640 = Castanopsis armata.
armeniaca, *Kotschy, Eichen*, t. 25.—Armenia.
aspera, Bosc, in Mém. Inst. Par. [viii.] (1807) 328 = Q. Robur.
asperata, Pers. Syn. ii. 571 = Q. Robur.
asplenifolia, Hort. ex A. DC. in DC. Prod. xvi. II. 42 = Q. Cerris.
atropurpurea, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
atrosanguinea, Hort. l. c. = Q. Robur.
atro-virens, Sm. in Rees, Cycl. xxix. n. 74 = Q. Robur.
Aucherii, *Jaub. & Spach, Illustr.* i. 113. t. 58.—Archip. Graec.
aurea, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
aurea, Kit. ex Reichb. Ic. Fl. Germ. xii. 8; Wierzb. ex Kotschy, Eichen, t. 4 = Q. Robur.
aurea, Rafin. Alsog. Am. 20 (Quid ?).—Am. bor.
australis, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 233 = Q. Robur.
australis, Link, Handb. ii. 466 = lusitanica.
austriaca, Willd. Sp. Pl. iv. 454 = Q. Cerris.
Auzandri, Gren. & Godr. Fl. Fr. iii. 119 = Q. Pseudo-coccifera.
Auzin, Secondat, ex Bosc, in Mém. Inst. Par. [viii.] (1807) 320 = Q. Robur.
avellanaeformis, *Colm. & Boutel. Exám. Árb. Pen.* 9.—Hispan.
axillaris, Schur, in Oestr. Bot. Wochenbl. vii. (1857) 18 = Q. Robur.
baetica, Buek, in DC. Ind. Prod. iv. 322 = lusitanica.
Ballota, Desf. in Mém. Acad. Sc. Par. 1790 (1797) 395. t. 6; Fl. Atlant. ii. 350 = Q. Ilex.
Baloot, Griff. Itin. Notes, 328 = Q. Ilex.
bambusaefolia, *Fortune, in Gard. Chron.* (1860) 170.—China.
bambusaefolia, *Hance, ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 116.—China.
bancana, Kurz, For. Fl. Brit. Burma, ii. 485 = polystachya.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 361; *et in Flora*, liii. (1870) 251.—Ins. Banca.
Banga, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 57 = lanuginosa.
Banisteri, Michx. Hist. Chênes Am. t. 27 = ilicifolia.
Banja, Endl. Gen. Suppl. iv. II. 27 = Q. Banga.
barbinervis, *Benth. Pl. Hartw.* 56.—Mexic.
Beccariana, *Benth. in Hook. Ic. Pl.* t. 1315.—Penang; Borneo.
Bennettii, *Miq. Fl. Ind. Bat.* i. I. 857.—Malaya.
Benthami, *A. DC. in DC. Prod.* xvi. II. 29.—Mexic.
berberidifolia, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 172 = dumosa.
bicolor, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 396.—Am. bor.
Bivoniana, Guss. Fl. Sic. Syn. ii. 604 = Q. Ilex.
Blancoi, *A. DC. in DC. Prod.* xvi. II. 97.—Ins. Philipp.
Blumeana, *Korth. Verh. Nat. Gesch. Bot.* 208. t. 44.—Malaya.

QUERCUS:—
Boissieri, Reut. in Boiss. Diagn. Ser. I. xii. 119 = lusitanica.
Bombyx, Hort. Leroy, ex C. Koch, Dendrol. ii. II. 72 = serrata.
Bonplandiana, Sweet, Hort. Brit. ed. I. 370 = obtusata.
borealis, Michx. f. N. Am. Sylv. i. 98. t. 26 = coccinea.
Bourgaei, *Oerst. ex Hemsl. Biol. Centr. Am. Bot.* iii. 168.—Mexic.
brachybalanos, Kotschy, ex A. DC. in DC. Prod. xvi. II. 54 = coccifera.
brachycarpa, Guss. ex Parl. Fl. Ital. iv. 180 = Q. Robur.
brachycarpa, Kotschy, ex A. DC. in DC. Prod. xvi. II. 18 = lusitanica.
brachyphylla, Kotschy, Eichen, t. 9 = Q. Robur.
brachystachys, *Benth. Pl. Hartw.* 91.—Mexic.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 108.—Burma.
Brantii, *Lindl. Bot. Reg.* (1840) *Misc.* 41.—Kurdistan.
brevifolia, Kotschy, ex A. DC. in DC. Prod. xvi. II. 11 Q. Robur.
brevipes, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 232 = Q. Robur.
brevipetiolata, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 359 = lamponga.
Breweri, Engelm. in Bot. Calif. ii. 96 = Garryana.
Brossa, Bosc, in Mém. Inst. Par. [viii.] (1807) 319 = Q. Toza.
brutia, Tenore, Ind. Sem. Hort. Neap. (1825) 11. n. 20 = Q. Robur.
Budayana, Haberle, ex Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 195 = lanuginosa.
Buergerii, *Blume, Mus. Bot. Lugd. Bat.* i. 299.—Japon.
bumelioides, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 188.—Am. centr.
Bungeana, *F. B. Forbes, in Journ. Bot.* xxii. (1884) 83, 85.—China.
calicina, Poir. Encyc. Suppl. ii. 216 = Q. Ilex.
californica, Cooper, in Smithson. Rep. (1858) 261 = Kelloggii.
callicarpifolia, Griff. Itin. Notes, ii. 87, partim = fenestrata.
calliprinos, Webb, Iter Hisp. 15 = coccifera.
callosa, Benth. Pl. Hartw. 91 = tomentosa.
callosa, Mart. ex A. DC. in DC. Prod. xvi. II. 28 = laxa.
calophylla, *Schlecht. & Cham. in Linnaea*, v. (1830) 79.—Mexic.
camuta, Hort. ex Petz. & Kirchn. Arb. Muscav. 628 = Q. Toza.
cana, Hort. ex Steud. Nom. ed. II. ii. 426 = Q. Cerris.
canariensis, Willd. Enum. Hort. Berol. 975 = lusitanica.
candicans, *Née, in Anal. Cienc. Nat.* iii. (1801) 277.—Hispan.
candida, *Hort. ex Steud. Nom.* ed. II. ii. 426, nomen.—Am. bor.
canescens, *Blume, Mus. Bot. Lugd.-Bat.* i. 296.—Japon.
caraballoana, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 209.—Ins. Philipp.
carduchorum, *C. Koch, in Linnaea*, xxii. (1849) 320.—Kurdistan.
carduchorum, Kotschy, ex A. DC. in DC. Prod. xvi. II. 49 = Libani.
carpatica, *Kit. in Linnaea*, xxxii. (1863) 353.—Hungaria.
carpinea, *Kotschy, ex A. DC. in DC. Prod.* xvi. II. 108.—Syria.
carpinifolia, Rafin. Alsog. Am. 21 (Quid ?).—Am. bor.
Cassura, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 57 = semecarpifolia.
castanaeifolia, Pantoč. Adnot. Fl. Hercegov. 31 = macedonica.
Castanea, Emerson, Trees Massach. ed. I. 137. t. 5 = Q. Prinus.
Castanea, Muhl. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 396 = prinoides.
Castanea, *Née, in Anal. Cienc. Nat.* iii. (1801) 276.—Mexic.
castaneaefolia, Coss. ex J. Gay, in Bull. Soc. Bot. Fr. viii. (1861) 421 = Q. Pseudo-suber.

QUERCUS:—
castaneaefolia, *C. A. Mey. Verz. Pfl. Cauc.* 44.—Reg. Caucas.
castanicarpa, Roxb. Hort. Beng. 68; Fl. Ind. ii. 640 = Castanopsis castanicarpa.
castellana, Bosc, ex Pers. Syn. ii. 571 = Q. Toza.
castellana, Poir. Encyc. Suppl. ii. 226 = Q. Ilex.
Catesbaei, *Michx. Hist. Chênes Am.* tt. 29, 30.—Am. bor.
Catungea, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 56 = Castanopis armata.
caudata, Lindl. in Wall. Cat. n. 2787 = Castanopsis tribuloides?
cedrorum, Kotschy, Eichen, t. 37 = Q. Robur.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 110.—Ins. Celebes.
cenomanensis, Desp. ex Endl. Gen. Suppl. iv. II. 24 = Q. Toza.
cerrioides, Willk. & Costa, in Linnaea, xxx. (1859–60) 123 = Q. Robur.
Cerris, Blanco, Fl. Filip. ed. I. 727 = Castanopsis javanica.
Cerris, DC. Fl. Fr. iii. 311 = Q. Toza.
Cerris, *Linn. Sp. Pl.* 997.—Europ.; Oriens.
Cerris, Pall. Fl. Ross. ii. 4 = Q. Robur.
chainolepis, Kotschy, ex A. DC. in DC. Prod. xvi. II. 55 = coccifera.
Championi, *Benth. in Hook. Kew Journ.* vi. (1854) 113.—China.
chinantlensis, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 179 = cuneifolia.
Chincapin, Rafin. Alsog. Am. 23 = Q. Prinus.
chinensis, *Abel, Narr. Journ. China*, 165, cum ic. 363.—China.
chinensis, Bunge, Enum. Pl. Chin. Bor. 61 = Bungeana.
Chinquapin, Pursh, Fl. Am. Sept. ii. 634 = Q. Prinus.
chrysolepis, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 173.—Calif.
chrysophylla, *Humb. & Bonpl. Pl. Aequin.* ii. 42. t. 87.—Mexic.
cinerea, *Michx. Hist. Chênes Am.* t. 14.—Am. bor.
cinerea, Rafin. Alsog. Am. 27 (an praec.?).—Am. bor.
cintrana, Welw. ex Nym. Consp. 662 = occidentalis.
circinata, *Née, in Anal. Cienc. Nat.* iii. (1801) 272.—Mexic.
citrifolia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 187.—Costa Rica.
classinervia, A. DC. in DC. Prod. xvi. II. 87, sphalm. = crassinervia.
Clivia, *Dehnh. ex Pasq. Cat. Orto Bot. Nap.* (1867) 87.—Ital.
coccifera, *Linn. Sp. Pl.* 995.—Reg. Mediterr.
coccinea, *Wangenh. Beitr. Anpfl. Holz.* 44. f. 9.—Am. bor.
collina, Schleich. ex Endl. Gen. Suppl. iv. II. 24 = Q. Robur.
commutata, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 177 = nitens.
Companoana, *Vidal, Sinops.* 41. t. 92. f. D.—Ins. Philipp.
comptoniaefolia, Hort. ex C. Koch, Dendrol. ii. II. 27 = Q. Robur.
concentrica, Blanco, Fl. Filip. ed. II. 502 = Llanosii.
concentrica, *Lour. Fl. Cochinch.* 572.—Cochinch.
Concordia, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
condensata, Schur, in Oestr. Bot. Wochenbl. vii. (1857) 18 = Q. Robur.
conferta, Kit. in Schult. Oesterr. Fl. ed. II. i. 619 = Q. Farnetto.
confertifolia, *Humb. & Bonpl. Pl. Aequin.* ii. 53. t. 94.—Mexic.
confertifolia, Torr. Bot. Mex. Bound. 207 = hypoleuca.
congesta, Presl, Delic. Prag. 32 = Q. Robur.
conglomerata, Pers. Syn. ii. 570 = Q. Robur.
conocarpa, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 207 = Reinwardtii.
conocarpa, *Oudem. in Versl. en Med. Kon. Akad. Wetensch.* xii. (1861) 206.—Malaya.
consobrina, Kotschy, ex A. DC. in DC. Prod. xvi. II. 54 = coccifera.
conspersa, Benth. Pl. Hartw. 91 = acutifolia.
Cookii, Loud. Arb. Brit. iii. 1926 = lusitanica.
cooperta, *Blanco, Fl. Filip.* ed. II. 503.—Ins. Philipp.
cordata, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 211.—Mexic.

QUERCUS:—
coriacea, Bechst. Forstbot. ed. V. 211 = Q. Robur.
cornea, *Lour. Fl. Cochinch.* 572.—China.
coronensis, Schur, Enum. Pl. Transs. 607 = Q. Robur.
corrugata, *Hook. Ic. Pl.* t. 403.—Guatemala.
Cortesii, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 175.—Mexic.
corticosa, Rafin. Alsog. Am. 24 (Quid?).—Am. bor.
corymbifolia, Ehrenb. ex Boiss. Fl. Orient. iv. 1167 = lusitanica.
costaricensis, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 184.—Costa Rica.
costata, *Blume, Bijdr.* 522.—Malaya.
costata, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 207 = sundaica.
crassifolia, Benth. Pl. Hartw. 56 = splendens.
crassifolia, *Humb. & Bonpl. Pl. Aequin.* ii. 49. t. 91.—Mexic.
crassinervia, *Blume, Mus. Bot. Lugd. Bat.* i. 292.—Java.
crassipes, *Humb. & Bonpl. Pl. Aequin.* ii. 37. t. 83.—Mexic.
crassipocula, Torr. in Pacif. Rail. Rep. iv. 137; v. 365 = chrysolepis.
crenata, Lam. Encyc. i. 724 = Q. Pseudo-suber.
cretensis, Lam. ex Steud. Nom. ed. I. 673 = Zelkova cretica.
cretica, Raulin, ex A. DC. in DC. Prod. xvi. II. 54 = coccifera.
crinita, Lam. Encyc. i. 718 = Q. Cerris.
crispa, Hort. ex C. Koch, Dendrol. ii. II. 56 = Q. Ilex.
crispa, Hort. ex Steud. Nom. ed. II. ii. 426 = Q. Cerris.
crispa, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 18 = Q. Robur.
crispata, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 386 = Q. Robur.
crispula, *Blume, Mus. Bot. Lugd. Bat.* i. 298.—Japon.
crispula, Vukot. in Rad Jugos. Akad. Zagreb. xxii. 1872 (1873) 14 (Quid?).—Europ.
cubana, A. Rich. Fl. Cub. Fanerog. ii. 230 = virginiana.
cuneata, Roxb. ex Wall. Cat. n. 3732 = lucida.
cuneata, Wangenh. Beitr. Teutsch. Holzg. Forstw. Nordam. Holz. 78. t. 5 = falcata.
cuneifolia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 182.—Mexic.
cuneifolia, Rafin. Alsog. Am. 27 (Quid?).—Am. bor.
Cupaniana, Guss. Fl. Sic. Syn. ii. 606 = Q. Robur.
cuprea, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
cupressoides, Hort. l. c. 29 = Q. Robur.
cupulatus, Gilib. Exercit. ii. 398 = Q. Robur.
cuspidata, *Thunb. Fl. Jap.* 176.—Japon.
cyclophora, *Endl. Gen. Suppl.* iv. II. 28.—Malaya.
cyclophylla, Welw. ex Nym. Consp. 662 = Q. Ilex.
cylindracea, Guss. ex Parl. Fl. Ital. iv. 178 = Q. Robur.
Cypri, Kotschy, ex A. DC. in DC. Prod. xvi. II. 19 = lusitanica.
Cypria, Jaub. & Spach. Illustr. i. 110 = alnifolia.
cyrtopoda, *Miq. Fl. Ind. Bat.* i. I. 869 (sp. dub.).—Sumatra.
cyrtorhyncha, *Miq. l. c. Suppl.* 350.—Malaya.
Daimio, Hort. ex C. Koch, Dendrol. ii. II. 45 = dentata.
Dalbertisii, *F. Muell. in Vict. Natural.* (Dec. 1884) 124.—N. Guin.
Dalechampii, Tenore, Ind. Sem. Hort. Neap. (1830) 15; et ej. Syll. 469 = Q. Robur.
daphnoidea, *Blume, Fl. Jav. Cupul.* 28. t. 16.—Java; Sumatra.
dasystachya, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 221.—Borneo.
dchorochensis, Boiss. Fl. Orient. iv. 1165, sphalm. = *dshorochensis*.
dealbata, *Hook. f. & Thoms. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 107.—Reg. Himal.
dealbata, Royle, Illustr. Bot. Himal. 346, in nota = dilatata.
dealbata, Wall. Cat. n. 2769 = incana.
decipiens, Bechst. Forstbot. ed. V. 211 = Q. Robur.
decipiens, Mart. ex Gal. in Bull. Acad. Brux. x. II. (1843) 214 = reticulata.
densiflora, *Hook. & Arn. Bot. Beech. Voy.* 391.—Calif.
densifolia, *Abel, Narr. Journ. China*, 165, 363.—China.

QUERCUS:—
dentata, Bartr. Trav. ed. II. (1794) 28 = aquatica.
dentata, *Thunb. Fl. Jap.* 177.—Japon.
dentata, Wats. Dendrol. t. 93 = Q. Cerris.
dentosa, Lindl. in Wall. Cat. n. 2775 = glauca.
depressa, Blume, in Verh. Batav. Gen. ix. (1825) 209 = spicata.
depressa, *Humb. & Bonpl. Pl. Aequin.* ii. 50. t. 92.—Mexic.
depressa, Roxb. Hort. Beng. 104; Fl. Ind. iii. 640 = cyclophora.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 349.—Sumatra.
dilatata, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 188 = Q. Robur.
dilatata, *Lindl. in Wall. Cat.* n. 2785.—Reg. Himal.
dilatata, Rafin. Alsog. Am. 24 (Quid?).—Am. bor.
dipsacina, Kotschy, ex A. DC. in DC. Prod. xvi. II. 55 = coccifera.
discocarpa, *Hance, in Journ. Bot.* xii. (1874) 242.—Malaya.
discolor, *Ait. Hort. Kew.* ed. I. iii. 358.—Am. bor.
dispar, Kotschy, ex A. DC. in DC. Prod. xvi. II. 55 = coccifera.
dispar, Rafin. Alsog. Am. 19 (Quid?).—Am. bor.
dissecta, Hort. ex C. Koch, Dendrol. ii. II. 27 = Q. Robur.
divaricata, *Lindl. in Wall. Cat.* n. 2790 (sp. dub.).—Burma.
diversifolia, Née, in Anal. Cienc. Nat. iii. (1801) 270 = tomentosa.
Douglasii, *Hook. & Arn. Bot. Beech. Voy.* 391.—Calif.
Drummondii, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 170.—Texas.
dschorochensis, Kotschy, Eichen, t. 39 = seq.
dshorochensis, C. Koch, in Linnaea, xxii. (1849) 328 = Q. Robur.
dubia, Lindl. in Wall. Cat. n. 2786 = Castanopsis indica.
dubia, Linn. Pl. Surin. n. 127; Amoen. Acad. viii. 263 (Quid?).—Guian.
dumosa, *Nutt. N. Am. Sylva*, i. 18.—Calif.
Dunnii, Kellogg, in Pacif. Rural Press (June 1879); ex Curran, in Bull. Calif. Acad. i. n. III. (1885) 147 = Palmeri.
Durandii, *Buckl. in Proc. Acad. Sc. Philad.* 1860 (1861) 445.—Am. bor.
Durinus, Rafin. Alsog. Am. 19 (Quid?).—Am. bor.
dysophylla, *Benth. Pl. Hartw.* 55.—Mexic.
echinacea, Torr. in Pacif. Rail. Rep. iv. 137 = densiflora.
echinata, A. DC. in DC. Prod. xvi. II. 683, sphalm. = *echinacea*.
echinata, Kotschy, ex A. DC. l. c. 55 = coccifera.
echinata, Lam. Fl. Fr. ii. 207 = Q. Aegilops.
echinata, Salisb. Prod. 393 = Q. Cerris.
echinoides, R. Br. Campst. in Ann. & Mag. Nat. Hist. Ser. IV. vii. (1871) 251 = densiflora.
Ehrenbergii, *Kotschy, Eichen*, t. 15.—Cilicia; Syria.
elegans, Blume, in Verh. Batav. Gen. ix. (1825) 208 = spicata.
elephantum, *Hance, in Journ. Bot.* xiii. (1875) 365.—Cambodia.
elliptica, Liebm. ex A. DC. in DC. Prod. xvi. II. 31 = glaucoides.
elliptica, *Née, in Anal. Cienc. Nat.* iii. (1801) 278.—Mexic.
elongata, Willd. Sp. Pl. 444 = falcata.
Emoryi, Porter & Coulter, Fl. Colorado, 127 = undulata?
Emoryi, *Torr. in Emory, Not. Mil. Reconnoiss.* 151. t. 9.—Am. bor. occ.
encleisacarpa, *Korth. Verh. Nat. Gesch. Bot.* 209. t. 45.—Malaya.
erucaefolia, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 388 = Q. Robur.
esculenta, Hort. ex C. Koch, Dendrol. ii. II. 80 = lusitanica.
Esculus, Linn. Mant. ii. 496 = Q. Cerris.
Esculus, Linn. Sp. Pl. 996 = Q. Robur.
Esculus, Pollini, Fl. Veron. iii. 125 = Q. Farnetto.
eugeniaefolia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 185.—Guatemala.
eumorpha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 107.—Burma.
Ewyckii, *Korth. Verh. Nat. Gesch. Bot.* 212. t. 46.—Malaya.

QUERCUS:—
excelsa, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 174.—Mexic.
exoniana ×, Poir. Encyc. Suppl. ii. 225 = seq.
exoniensis ×, Lodd. ex Sweet, Hort. Brit. ed. I. 370; et ex Webb, Iter Hisp. 13.
expansa, Poir. Encyc. Suppl. ii. 216 = Q. Ilex.
extensa, Schur, Enum. Pl. Transs. 609 = Q. Robur.
Eyrei, *Champ. ex Benth. in Hook. Kew Journ.* vi. (1854) 114.—Ins. Hongkong.
Fabri, *Hance, in Journ. Linn. Soc.* x. (1869) 202.—China.
fagifolia, Hort. ex C. Koch, Dendrol. ii. II. 56 = Q. Ilex.
fagiformis, Jungh. in Bonplandia, vi. (1858) 83 = Junghuhnii.
faginea, Lam. Encyc. i. 725 = lusitanica.
faginea, Tenore, Syll. Fl. Neap. 469 = Q. Robur.
falcata, *Michx. Hist. Chênes Am.* t. 28.—Am. bor.
Falconeri, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 197.—Burma.
falkenbergensis, Booth, ex Loud. Arb. Brit. iii. 1926 = Q. Robur.
farinulenta, *Hance, in Journ. Bot.* xiii. (1875) 365.—Cambodia.
Farnetto, *Tenore, Syll. Fl. Neap.* 470.—Europ. austr.
fastigiata, Lam. Encyc. i. 725; DC. Fl. Fr. Suppl. 351 = Q. Robur.
Fendleri, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 170.—Am. bor. occ.
fenestrata, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 633.—Reg. Himal.
fenestrata, Wall. Cat. n. 2784, partim = lanceaefolia.
Fennessi, Hort. ex A. DC. in DC. Prod. xvi. II. 6 = Q. Robur.
Fenzlii, Kotschy, Eichen, t. 24 = coccifera.
Fernandezii, *Vidal, Sinops.* 41. t. 92. f. E.—Ins. Philipp.
ferox, Roxb. Hort. Beng. 104; Fl. Ind. iii. 639 = Castanopsis tribuloides.
ferruginea, Michx. f. Hist. Arb. Am. ii. 92. t. 18 = nigra.
filicifolia, Hort. ex A. DC. in DC. Prod. xvi. II. 6 = Q. Robur.
filiformis, Muhl. ex A. DC. l. c. 21 = bicolor.
filipendula, Vukot. in Flora, li. (1868) 330 = Q. Robur.
fissa, *Champ. ex Benth. in Hook. Kew Journ.* vi. (1854) 114.—Ins. Hongkong.
flava, Née, ex Spreng. Syst. iii. 857 = magnoliaefolia.
flavida, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 178.—Mexic.
floccosa, *Liebm. l. c.*—Mexic.
floribunda, Wall. Cat. n. 2773 = dilatata.
floridana, Shuttlew. ex A. DC. in DC. Prod. xvi. II. 22 = obtusiloba.
foeminea, Mill. Gard. Dict. ed. VIII. n. 2 = Q. Robur.
Fontanesii, Guss. Ind. Sem. Hort. Boccad. (1825); ex ej. Fl. Sic. Syn. ii. 605 = Q. Pseudo-suber.
Fordii, Carr. in Rev. Hortic. (1861) 114. ff. 24, 25 = Q. Ilex.
Frainetto, Tenore, Cat. Hort. Nap. (1819) 65, sphalm. = Q. Farnetto.
frondosa, Steud. Nom. ed. II. ii. 426 = Q. Cerris.
fructipendula, Schrank, Baier. Fl. i. 660 = Q. Robur.
fruticosa, Brot. Fl. Lusit. ii. 31 = humilis.
fulhamensis ×, Hort. ex Steud. Nom. ed. II. ii. 426.
fulva, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 183.—Mexic.
fulvescens, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 70 = chrysolepis.
furfuracea, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 189 = acutifolia.
fusca, Rafin. Alsog. Am. 19 (Quid ?).—Am. bor.
Galeotti, *Martens, in Bull. Acad. Brux.* x. II. (1843) 220.—Mexic.
Gambelii, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 179 = undulata.
Gambellii, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 169 = Douglasii.
Garryana, *Dougl. ex Hook. Fl. Bor. Am.* ii. 159.—Am. bor. occ.
geltowiensis, Hort. ex C. Koch, Dendrol. ii. II. 32 = Q. Robur.
gemelliflora, *Blume, in Verh. Batav. Gen.* ix. (1825) 222.—Java.

QUERCUS:—
gemelliflora, Herb. Buerger, ex Blume, Mus. Bot. Lugd. Bat. i. 299 = Buergerii.
Genabii, Hort. ex C. Koch, Dendrol. ii. II. 62 = nigra.
georgiana, *M. A. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 408.—Am. bor.
germana, *Schlecht. & Cham. in Linnaea*, v. (1830) 78.—Mexic.
germanica, Lasch, in Bot. Zeit. xv. (1857) 413 = Q. Robur.
Ghiesbreghtii, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 212.—Mexic.
gibraltarica, C. Koch, Dendrol. ii. II. 81 = lusitanica.
gilva, *Blume, Mus. Bot. Lugd. Bat.* i. 306.—Japon.
glaberrima, Blume, in Verh. Batav. Gen. ix. (1825) 210 = spicata.
glabra, Blanco, Fl. Filip. ed. I. 727 = ovalis.
glabra, Buch.-Ham. ex Wall. Cat. n. 3731 = acuminata.
glabra, Buerger, ex Blume, Mus. Bot. Lugd. Bat. i. 290, 304 = thalassica, marginata.
glabra, *Thunb. Fl. Jap.* 175.—Japon.
glabrescens, *Benth. Pl. Hartw.* 56, 348.—Mexic.
glabrescens, Kern. in Oestr. Bot. Zeitschr. xxvi. (1876) 230 = *Kernerii*.
glabrescens, Seem. Bot. Voy. Herald, 252 = Warszewiczii.
glandulifera, *Blume, Mus. Bot. Lugd. Bat.* i. 295.—Japon.
glauca, Bosc, Mém. Chênes, 26; ex Loisel. in Duham. Arb. ed. Nov. vii. 161 = humilis.
glauca, Buerger, ex Blume, Mus. Bot. Lugd. Bat. i. 290 = thalassica.
glauca, Martr. & Timb. in Bull. Soc. Bot. Fr. xi. (1864) p. xvi = Q. Ilex.
glauca, *Thunb. Fl. Jap.* 175.—Reg. Himal.; Japon.
glaucescens, *Humb. & Bonpl. Pl. Aequin.* ii. 29. t. 78.—Mexic.
glaucoides, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 209.—Mexic.
glomerata, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 640.—Ins. Penang.
glomerata, Wall. Cat. n. 2791 = Castanopsis sumatrana.
glutinosa, *Blume, Mus. Bot. Lugd. Bat.* i. 304.—Java.
Goedelii, Balansa & Kotschy, ex A. DC. in DC. Prod. xvi. II. 46 = Q. Aegilops.
gonoloba, Rafin. Alsog. Am. 25 (Quid ?).—Am. bor.
gracilipes, Miq. Fl. Ind. Bat. Suppl. 347 = spicata.
gracilis, *Korth. Verh. Nat. Gesch. Bot.* 207.—Borneo.
gracilis, Lange, in Kjoeb. Vidensk. Meddel. (1861) 36 = Q. Ilex.
graeca, Kotschy, Eichen, t. 30 = Q. Aegilops.
Grahami, *Benth. Pl. Hartw.* 57.—Mexic.
Grahami, Hort. Hafn. ex A. DC. in DC. Prod. xvi. II. 66 = acutifolia.
gramuntia, Linn. Sp. Pl. 995 = Q. Ilex.
grandifolia, D. Don, in Lamb. Gen. Pin. ii. 27. t. 8 = spicata.
grandis, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 183.—Guatemala.
granulata, *Liebm. l. c.* 186.—Costa Rica.
granulata, Rafin. Alsog. Am. 21 (Quid ?).—Am. bor.
Griffithii, *Hook. f. & Thoms. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 104.—Reg. Himal.
grisea, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 171.—Am. bor. occ.
grosseserrata, *Blume, Mus. Bot. Lugd. Bat.* i. 306.—Japon.
guatimalensis, *A. DC. in DC. Prod.* xvi. II. 78.—Guatemala.
Gulliveri, *F. Muell. in Vict. Natural.* i. (1885) 167.—N. Guin.
Guppyi, *F. Muell. l. c.* (Dec. 1884) 123, in textu.—Ins. Salomon.
Haas, Kotschy, Eichen, t. 2 = Q. Robur.
haliphlaeos, Lam. Encyc. i. 718 = Q. Cerris.
haliphleos, Guss. Fl. Sic. Syn. ii. 605 = Q. Pseudo-suber.
Hancei, *Benth. Fl. Hongk.* 322.—China.
Harlandi, *Hance, ex Walp. Ann.* iii. 382.—China.
Hartwegi, Benth. Pl. Hartw. 56 = obtusata.
Hartwissiana, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 387 = Q. Robur.
Hasseltii, Korth. ex Miq. Fl. Ind. Bat. i. I. 861 = omalokos.
hastata, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 171 = Emoryi.

QUERCUS:—
Haynaldiana, *Simk. in Magyar Növényt. Lapok*, vii. (1883) 63. t. 3.—Hungar.
Helferiana, *A. DC. in DC. Prod.* xvi. II. 101.—Burma.
hemisphaerica, Bartr. Trav. 320 = nigra.
hemisphaerica, Hort. ex Endl. Gen. Suppl. iv. II. 25 = virginiana.
Hentzei, Petz. Arb. Muscav. 627 = Q. Robur.
heteroloba, Rafin. Alsog. Am. 25 (Quid ?).—Am. bor.
heterophyla, Rafin. l. c. 24 (Quid ?).—Am. bor.
heterophylla, Hort. Parment. ex A. DC. in DC. Prod. xvi. II. 42 = Q. Cerris.
heterophylla, *Michx. f. Hist. Arbr. Am.* ii. 87.—Am. bor.
Hindsii, Benth. Bot. Voy. Sulph. 55 = lobata.
hirsuta, Lindl. in Wall. Cat. n. 3734 = lappacea.
hirsutula, Blume, Mus. Bot. Lugd. Bat. i. 298 = aliena.
hispanica, Lam. Encyc. i. 723 = Q. Pseudo-suber.
hispanica, Willk. in Flora, xxxv. (1852) 315 = Q. Robur.
Hodginsii, Lodd. Cat. (1836) ex Steud. Nom. ed. II. ii. 427 = Q. Robur.
holosericea, Blume, Mus. Bot. Lugd. Bat. i. 298 = aliena.
Horsfieldii, Miq. Fl. Ind. Bat. i. I. 856 = semiserrata.
Humboldtii, *Bonpl. in Humb. & Bonpl. Pl. Aequin.* ii. 155. t. 130.—N. Granat.
Humboldtii, Kotschy, ex A. DC. in DC. Prod. xvi. II. 32 = Lindeni.
humilis, DC. Fl. Fr. iii. 312 = Q. Toza.
humilis, *Lam. Encyc.* i. 719.—Reg. Mediterr. occ.
humilis, Mill. Gard. Dict. ed. VIII. n. 4 = Q. Robur ?
humilis, Walt. Fl. Carol. 234 = cinerea.
humosa, Blume, Mus. Bot. Lugd. Bat. i. 298 = aliena.
hungarica, Hubeny, ex Rochel, Bot. Reise Banat. 73; Kit. in Linnaea, xxxii. (1863) 353 = Q. Farnetto.
hybrida, Bechst. Forstbot. ed. V. 211 = Q. Robur.
hybrida, Brot. Fl. Lusit. ii. 31 = lusitanica.
hyemalis, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 385, in textu = Q. Robur.
hypochrysa, Stev. l. c. 388 = Q. Robur.
hypoleuca, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 384.—N. Mexic.
hypoleuca, Kotschy, ex A. DC. in DC. Prod. xvi. II. 45 = lusitanica.
hypoleuca, Miq. Fl. Ind. Bat. i. I. 869 = Korthalsii.
hypophlaeos, Petz. Arb. Muscav. 656 = falcata.
Hystrix, *Korth. Verh. Nat. Gesch. Bot.* 201.—Malaya.
iberica, Bieb. Fl. Taur. Cauc. ii. 402 = Q. Robur.
Ibicis, Kotschy, ex A. DC. in DC. Prod. xvi. II. 9 = Q. Robur.
Ijam-ahoso, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 93 = Meliosma myriantha.
Ilex, *Linn. Sp. Pl.* 995.—Reg. Mediterr.; Oriens.
Ilex, Lour. Fl. Cochinch. 571 = Helferiana.
ilexoides, Rafin. Alsog. Am. 23. (Quid ?).—Am. bor.
ilicifolia, Salisb. Prod. 392 = Q. Ilex.
ilicifolia, *Wangenh. Forstw.* 79. *f.* 17.—Am. centr.
imbricaria, A. Gray, ex A. DC. in DC. Prod. xvi. II. 63 = Q. Phellos.
imbricaria, *Michx. Hist. Chênes Am.* tt. 15, 16.—Am. bor.
imbricata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 57 = lamellosa.
incana, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 642.—Reg. Himal.
incisa, Hort. ex Steud. Nom. ed. II. ii. 427 = Q. Cerris.
induta, *Blume, in Verh. Batav. Gen.* ix. (1825) 220.—Java.
inermis, Ehrenb. ex A. DC. in DC. Prod. xvi. II. 18; et ex Unger & Kotschy, Insel Cypern, 216 = lusitanica.
infectoria, Oliv. Voy. Atlas, tt. 14, 15 = lusitanica.
inops, Kotschy, ex A. DC. in DC. Prod. xvi. II. 54 = coccifera.
insignis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 219.—Mexic.
integrifolia, Hort. ex Steud. Nom. ed. I. 673 = Q. Ilex.
intermedia, Boenn. ex Reichb. Fl. Germ. Excurs. 177; D. Don, ex Leighton, Fl. Shrop. 473 = Q. Robur.
intermedia, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 223 = calophylla.
inversa, Lindl. & Paxt. Flow. Gard. i. (1850–51) 58. f. 36 = thalassica.

QUERCUS:—
Irwinii, *Hance, in Ann. Sc. Nat. Sér.* IV. xviii. (1862) 229.—Ins. Hongkong.
iteaphylla, *Hance, in Journ. Bot.* xxii. (1884) 229.—Ins. Hongkong.
ithaburensis, Decne. l. c. Sér. II. iv. (1835) 348 = Q. Aegilops.
Jacobi, R. Br. Campst. in Ann. & Mag. Nat. Hist. Ser. IV. vii. (1871) 255 = Garryana.
Jama-buwa, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 93 = Meliosma pungens.
javensis, *Miq. l. c.* i. 117.—Java.
Jenkinsii, *Benth. in Hook. Ic. Pl.* t. 1312–13.—Reg. Himal.
Jordanae, *Villanueva, Apuntes Un Nuev. Roble*, (1875) 7.—Ins. Philipp.
Juergensenii, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 188.—Mexic.
Junghuhnii, *Miq. Fl. Ind. Bat.* i. 1. 853.—Java.
Kajan, Miq. in Zoll. Syst. Verz. Ind. Archip. 87 = Q. Hystrix.
Kamroopii, D. Don, Prod. Fl. Nep. 57 = lineata?
Kelloggii, *Newberry, in Pacif. Rail. Rep.* vi. *Bot.* 28, 89. f. 6.—Calif.
Kernerii ×, Simk. in Magyar Növényt. Lapok, vii. (1883) 69.—Europ.
Korthalsii, *Blume, Mus. Bot. Lugd. Bat.* i. 292.—Java.
Korthalsii, Endl. Gen. Suppl. iv. II. 28 = Teysmannii.
Kurzii, Hance, in Journ. Bot. xvi. (1878) 328 = lanceaefolia.
lacera, *Blume, Mus. Bot. Lugd. Bat.* i. 306.—Japon.
laciniata, *Vukot. in Rad Jugos. Akad. Zagreb.* xxii. 1872 (1873) 19.—Europ.
laciniosa, Bor. Fl. Centr. Fr. ed. III. ii. 588, in obs. = Q. Robur.
laeta, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 179.—Mexic.
laevigata, *Blume, Mus. Bot. Lugd. Bat.* i. 304.—Japon.
laevis, Walt. Fl. Carol. 234 = Catesbaei?
lamellata, Roxb. Hort. Beng. 104; Fl. Ind. iii. 641 = lamellosa.
lamellosa, *Sm. in Rees, Cycl.* xxix. n. 23.—Reg. Himal.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 348.—Malaya.
lamprophyllos, C. Koch, in Linnaea, xxii. (1849) 327 = Q. Robur.
lanata, Sm. in Rees, Cycl. xxix. n. 27 = incana.
lanceaefolia, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 634.—Reg. Himal.
lanceolata, *Humb. & Bonpl. Pl. Aequin.* ii. 34. t. 81.—Mexic.
lanceolata, Mart. & Gal. ex A. DC. in DC. Prod. xvi. II. 79 = oajacana.
lancifolia, Benth. Pl. Hartw. 55 = nitens.
lancifolia, Liebm. ex A. DC. in DC. Prod. xvi. II. 72 = leiophylla.
lancifolia, *Schlecht. & Cham. in Linnaea*, v. (1830) 78.—Mexic.
lanigera, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 215.—Mexic.
lanuginosa, *D. Don, Prod. Fl. Nep.* 57.—Reg. Himal.
lanuginosa, Thuill. Fl. Par. ed. II. 502 = Q. Robur.
lappacea, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 637.—Reg. Himal.; Malaya.
Lasistan, Kotschy, ex A. DC. Prod. xvi. II. 5 = Q. Robur.
latifolia, Hort. ex Steud. Nom. ed. II. ii. 427 = imbricaria.
laurifolia, *Michx. Hist. Chênes Am.* t. 17.—Am. bor.
laurifolia, Miq. Pl. Jungh. 11 = Teysmannii.
laurina, *Humb. & Bonpl. Pl. Aequin.* ii. 32. t. 80.—Mexic.
laurina, Liebm. ex A. DC. in DC. Prod. xvi. II. 70 = lanceolata.
laurina, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 210 = depressa.
laxa, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 181.—Mexic.
laxiflora, Lindl. ex Wall. Cat. n. 2774 = glauca.
Leana, *Nutt. N. Am. Sylv.* i. 25. t. 5 bis.—Am. bor.
leiophylla, *A. DC. in DC. Prod.* xvi. II. 71.—Mexic.
leptobalana, Guss. Fl. Sic. Syn. ii. 608 = Q. Toza.
leptocarpa, *Vukot.—Cf. Oestr. Bot. Zeitschr.* xxx. (1880) 163.—Europ.
leptocarpos, Kotschy, ex Boiss. Fl. Orient. iv. 1167 = lusitanica.

QUERCUS:—
leptogyne, *Korth. Verh. Nat. Gesch. Bot.* 206 (sp. dub.).—Borneo.
Lezermiana, *Bosc, ex Poir. Encyc. Suppl.* ii. 226.—Europ. austr.
Libani, *Oliv. Voy. Atlas*, t. 32.—Syria.
Lindeni, *A. DC. in DC. Prod.* xvi. II. 32.—N. Granat.
Lindleyana, *Wall. Cat.* n. 2782.—Burma.
lineata, *Blume, Bijdr.* 523.—Reg. Himal.; Java.
lineata, Miq. Pl. Jungh. 10 = Junghuhnii.
linguaefolia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 180.—Mexic.
litoralis, *Blume, Mus. Bot. Lugd. Bat.* i. 303 (sp. dub.).—Java.
litseifolia, *Hance, in Journ. Bot.* xxii. (1884) 228.—China.
Llanosii, *A. DC. in DC. Prod.* xvi. II. 97.—Ins. Philipp.
lobata, *Née, in Anal. Cienc. Nat.* iii. (1801) 277.—Calif.
lobulata, Soland. ex Sm. in Abbot, Insects Georgia, i. 93. t. 47 = obtusiloba.
longaeva, Salisb. Prod. 392 = Q. Robur.
longepedunculata, Willk. Forstl. Fl. 864 (Index) = *longipetiolata*.
longifolia, C. Koch, in Linnaea, xxii. (1849) 327 = Q. Robur.
longifolia, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 185 = acutifolia.
longifolia, Rafin. Alsog. Am. 21 (Quid?).—Am. bor.
longiglanda, Torr. & Frém. in Frém. Geol. Mem. Upper Calif. 15, 17 = lobata.
longipes, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 387 = Q. Robur.
longipetiolata, Schur, Enum. Pl. Transs. 607 = Q. Robur.
Look, *Kotschy, Eichen*, t. 21.—Syria.
Lorentii, *Hochst. in Lorent, Wanderung.* 327.—As. Min.
Loureirii, Hance, in Journ. Linn. Soc. x. (1869) 201 = Castanopsis tribuloides.
lucida, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 635.—Malaya.
Lucombeana ×, Sweet, Hort. Brit. ed. I. 370.
lusitanica, *Lam. Encyc.* i. 719.—Reg. Mediterr.; Oriens.
lusitanica, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 666 = humilis.
lutea, *Née, in Anal. Cienc. Nat.* iii. (1801) 269.—Mexic.
lutescens, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 219.—Mexic.
lyrata, Spreng. Syst. iii. 864, partim = lobata.
lyrata, *Walt. Fl. Carol.* 235.—Am. bor.
McCormickii, *Carruth. in Journ. Linn. Soc.* vi. (1862) 32.—China.
macedonica, *A. DC. in DC. Prod.* xvi. II. 50.—Macedon.
Mackiana, Hook. Ic. Pl. t. 224 = lappacea.
macranthera, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) II. 259 = castaneaefolia.
macrocarpa, Endl. Gen. Suppl. iv. II. 24, sphalm. = *microphylla*.
macrocarpa, *Michx. Hist. Chênes Am.* tt. 2, 3.—Am. bor.
macrolepis, Kotschy, Eichen, t. 16 = Q. Aegilops.
macrophylla, Miq. Fl. Ind. Bat. Suppl. 351 = sundaica.
macrophylla, Née, in Anal. Cienc. Nat. iii. (1801) 274 = magnoliaefolia.
macrostipulata, Guss. ex Parl. Fl. Ital. iv. 178 = Q. Robur.
magnoliaefolia, *Née, in Anal. Cienc. Nat.* iii. (1801) 268.—Mexic.
Maingayi, *Benth. in Hook. Ic. Pl.* t. 1314.—Ins. Penang.
malacophylla, Schur, in Oestr. Bot. Zeitschr. x. (1860) 181 = Q. Robur.
mannifera, Lindl. Bot. Reg. (1840) Misc. 40 = Q. Robur.
mappacea, Korth. Verh. Nat. Gesch. Bot. 202 = sundaica.
marginata, *Blume, Mus. Bot. Lugd. Bat.* i. 304.—Japon.
maritima, Rafin. Alsog. Am. 27 (Quid?).—Am. bor.
maritima, Willd. Sp. Pl. iv. 424 = aquatica.
marylandica, Du Roi, Harbk. Baumz. ii. 274 = nigra.

QUERCUS:—
Matasii, Siebold, in Verh. Batav. Gen. xii. (1830) 26 = glauca.
menesiensis, *Kit. in Linnaea*, xxxii. (1863) 353.—Hungar.
Merkusii, Endl. Gen. Suppl. iv. II. 28 = lineata.
mespilifolia, *Wall. Cat.* n. 2766.—Burma.
mespilifolia, Wallr. Sched. Crit. 494 = Q. Robur.
Mesto, Boiss. Voy. Bot. Espagne, ii. 579. t. 166 = coccifera.
mexicana, Benth. Pl. Hartw. 56 = crassipes.
mexicana, Humb. & Bonpl. Pl. Aequin. ii. 35. t. 82 = Q. Castanea.
Michauxii, *Nutt. Gen. Am.* ii. 215.—Am. bor.
microbalanos, Bor. Fl. Centr. Fr. ed. III. ii. 588, in obs. = Q. Robur.
microcalyx, Korth. Verh. Nat. Gesch. Bot. 204 = spicata.
microcarpa, Lapeyr. Hist. Abr. Pl. Pyr. 582 = Q. Robur.
microcarpa, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 184 = elliptica.
microphylla, *Née, in Anal. Cienc. Nat.* iii. (1801) 264.—Mexic.
Miqueliana, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 360 = Bennettii.
Mirbeckii, Dur. in Duch. Rev. Bot. ii. (1846) 426 = lusitanica.
mitis, *Banks, ex Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 15 (sp. dub.).—Ins. Mader.
mixta, A. DC. in DC. Prod. xvi. II. 83 = spicata.
mollis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 216.—Mexic.
mollis, Nutt. Gen. Am. ii. 215 = bicolor.
molucca, Blanco, Fl. Filip. ed. I. 726 = Llanosii.
molucca, *Linn. Sp. Pl.* 1199.—Ins. Molucc.
mongolica, *Fisch. ex Turcz. in Bull. Soc. Nat. Mosc.* (1838) 101, nomen; xxvii. (1854) I. 409.—As. or.
montana, Willd. Sp. Pl. iv. 440 = Q. Prinus.
monticola, Petz. & Kirchn. Arb. Muscav. 644 = praec?
Morehus, Kellogg, in Proc. Calif. Acad. ii. (1863) 36 = Wislizeni.
Morisii, *Borzi, in Nuov. Giorn. Bot. Ital.* xiii. (1881) 10. t 1.—Ins. Sardoa.
Moulei, *Hance, in Journ. Bot.* xiii. (1875) 363.—China.
mucronata, Willd. Sp. Pl. iv. 436 = Q. Castanea.
Muehlenbergii, Engelm. in Trans. Acad. Sc. St. Louis, iii. (1878) 391 = Q. Prinus.
muricata, Pallas, Pract. Bot. vii. 172; ex Willk. & Lange, Prod. Fl. Hisp. i. 240 = lusitanica.
muricata, Roxb. Hort. Beng. 104; Fl. Ind. iii. 635 = sundaica.
myrsinaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 305.—Japon.
myrtifolia, Willd. Sp. Pl. iv. 424 = aquatica.
naiadarum, *Hance, in Journ. Bot.* xxii. (1884) 227.—China.
nana, Willd. Sp. Pl. iv. 443 = aquatica.
Nara, Hort. ex A. DC. in DC. Prod. xvi. II. 14 = aliena.
Neaei, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 173 = Garryana.
nectandrifolia, *Liebm. l. c.* 175.—Mexic.
nepaulensis, Desf. Cat. Hort. Par. ed. III. 414 = incana.
neurophylla, Miq. Fl. Ind. Bat. Suppl. 351 = sundaica?
nigra, *Linn. Sp. Pl.* 995.—Am. bor.
nigra, Marsh. Arbust. 120 = discolor.
nigra, Rugel, ex A. DC. in DC. Prod. xvi. II. 60 = rubra.
nigra, Thore, Chlor. Land. 381 = Q. Toza.
nigrescens, Rafin. Alsog. Am. 19 (Quid?).—Am. bor.
nigricans, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
nitens, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 217.—Mexic.
nitida, *Blume, Mus. Bot. Lugd. Bat.* i. 294 (sp. dub.).—Sumatra.
nitida, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 210 = acutifolia.
nitida, Rafin. Alsog. Am. 20 (Quid?).—Am. bor.
nobilis, Hort. ex C. Koch, Dendrol. ii. II. 65 = falcata.
nobilis, Hort. Ottoland. ex Mast. in Gard. Chron. (1875) II. 816 = nigra.

QUERCUS:—
Novi-Orleani, Petz. & Kirchn. Arb. Muscav. 656 = aquatica.
nudinervis, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 182 = obtusata.
oajacana, *Liebm. l. c.* 178.—Mexic.
oblonga, Hort. ex C. Koch, Dendrol. ii. II. 56 = Q. Ilex.
oblongata, D. Don, Prod. Fl. Nep. 57 = incana?
oblongifolia, R. Br. Campst. in Ann. & Mag. Nat. Hist. Ser. IV. vii. (1871) 252 = chrysolepis?
oblongifolia, *Torr. in Sitgreaves, Rep.* 173. t. 19.—Calif.; N. Mexic.
obovata, *Bunge, Enum. Pl. Chin. Bor.* 62.—China.
obtecta, *Poir. Encyc. Suppl.* ii. 182—China.
obtusa, Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 152, in syn. = laurifolia.
obtusata, *Humb. & Bonpl. Pl. Aequin.* ii. 26. t. 76.—Mexic.
obtusifolia, D. Don, Prod. Fl. Nep. 56 = semecarpifolia.
obtusiloba, *Michx. Hist. Chênes Am.* t. 1.—Am. bor.
occidentalis, *J. Gay, in Ann. Sc. Nat.* Sér. IV. vi. (1856) 243.—Europ. austr.
ocoteaefolia, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 176 = nitens.
ocreata, Jungh. ex Miq. Fl. Ind. Bat. i. I. 937 (= *Lepidadenia sumatrana*).—Sumatra.
Oerstediana, R. Br. Campst. in Ann. & Mag. Nat. Hist. Ser. IV. vii. (1871) 250 = Garryana.
oidocarpa, *Korth. Verh. Nat. Gesch. Bot.* 216. t. 47. f. 18.—Malaya.
oleoides, Schlecht. & Cham. in Linnaea, v. (1830) 79 = virginiana.
oligodes, Rafin. Alsog. Am. 28 (Quid?).—Am. bor.
oligoneura, *Korth. Verh. Nat. Gesch. Bot.* 203 (sp. dub.).—Sumatra.
olivaeformis, Michx. f. Hist. Arb. Am. ii. 32. t. 2 = macrocarpa.
olla, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 197 (sp. dub.).—Assam.
omalokos, *Korth. Verh. Nat. Gesch. Bot.* 214.—Malaya.
omissa, *A. DC. in DC. Prod.* xvi. II. 28.—Mexic.
oocarpa, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 184 = Warszewiczii.
oogyne, *Miq. Fl. Ind. Bat. Suppl.* 351.—Sumatra.
oophora, Kotschy, Eichen, t. 26 = Brantii.
Orizabae, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 189.—Mexic.
ovalifolia, Bosc, ex Pers. Syn. ii. 570 = lusitanica.
ovalis, *Blanco, Fl. Filip.* ed. II. 502.—Ins. Philipp.
oxyadenia, Torr. in Sitgreaves, Rep. 172. t. 17 = agrifolia.
oxycarpa, *Vukot. in Rad Jugos. Akad. Zagreb.* xxii. 1872 (1873) 13.—Europ.
oxyodon, Miq. Ann. Mus. Bot. Lugd. Bat. i. 114 = lineata.
oxyrhyncha, Miq. Fl. Ind. Bat. Suppl. 347 = lineata.
pachyphylla, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 197.—Reg. Himal.
Pagoda, Rafin. Alsog. Am. 23 (Quid?).—Am. bor.
palaestina, Kotschy, Eichen, t. 19 = coccifera.
paleacea, Desv. Journ. Bot. i. (1813) 48 = Q. Robur.
pallida, *Blume, Bijdr.* 524.—Java.
pallida, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 28; et in Verh. Zool.-Bot. Ges. Wien (1858), 195 = Q. Robur.
palmata, *Vukot. in Rad Jugos. Akad. Zagreb.* xxii. (1872) 11.—Europ.
Palmeri, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1878) 393.—Calif.
paludosa, Hort. ex Petz. Arb. Muscav. 646 = bicolor.
palustris, *Du Roi, Harbk. Baumz.* ii. 268. t. 5. f. 4.—Am. bor.
palustris, Rugel, ex A. DC. in DC. Prod. xvi. II. 62 = coccinea.
palustris, Torr. & Gray, in Pacif. Rail. Rep. ii. 175 = rubra.
pandurata, Humb. & Bonpl. Pl. Aequin. ii. 28 = obtusata.
pandurata, *Rafin. Alsog. Am.* 21.—Am. bor.
pannonica, Hort. ex Endl. Gen. Suppl. iv. II. 24 = Q. Farnetto.
pannosa, Bosc, ex A. DC. in DC. Prod. xvi. II. 21 = bicolor.
Parmenteria, Hort. Soulang. ex Mutel, Fl. Fr. iii. 204 = Q. Robur.

QUERCUS:—
paucilamellosa, A. DC. in DC. Prod. xvi. II. 101 = lamellosa.
pectinata, Hort. ex C. Koch, Dendrol. ii. II. 27 = Q. Robur.
pedemontana, Colla, Herb. Pedem. v. 185 = Q. Robur.
pedunculariis, Née, in Anal. Cienc. Nat. iii. (1801) 270 = tomentosa.
pedunculata, Ehrh. Beitr. v. 161 = Q. Robur.
pedunculata, Née, ex A. DC. in DC. Prod. xvi. II. 33 = tomentosa.
pedunculiflora, C. Koch, in Linnaea, xxii. (1849) 324 = Q. Robur.
penangensis, Miq. Fl. Ind. Bat. i. I. 859 = cyclophora.
pendula, Lodd. Cat. (1836) ex Loud. Arboret. iii. 1732 = Q. Robur.
pendulina, Kit. in Schult. Oester. Fl. ed. II. i. 620 = Q. Robur.
perseaefolia, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 188 = elliptica.
persica, *Jaub. & Spach, Illustr.* i. 109. t. 55.—Persia.
petiolaris, Benth. Pl. Hartw. 55 = polymorpha.
petiolaris, Boiss. & Reut. Diagn. Ser. I. xii. 120 = lusitanica.
petiolata, Schur, in Oestr. Bot. Wochenbl. vii. (1857) 18 = Q. Robur.
Pfaeffingeri, Kotschy, Eichen, t. 23 = lusitanica.
phellodrys, *Link, ex Steud. Nom.* ed. I. 674; II. ii. 427, nomen.—Hab.?
Phellos, *Linn. Sp. Pl.* 994.—Am. bor.
philippinensis, A. DC. in DC. Prod. xvi. II. 97 = lineata?
phillyraeoides, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 406.—Japon.
Phullata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 57 = glauca.
Pinanga, *Blume, Mus. Bot. Lugd. Bat.* i. 303 (sp. dub.).—Java.
pinnatifida, *Franch. & Sav. Enum. Pl. Jap.* ii. 497.—Japon.
pinnatifida, C. C. Gmel. Fl. Bad. iv. 673 = Q. Robur.
pinnatifida, Vukot. in Rad Jugos. Acad. Zagreb. xxii. 1872 (1873) 12 = Q. Robur?
pinnatiloba, C. Koch, in Linnaea, xxii. (1849) 326 = Q. Robur.
placentaria, Blume, Bijdr. 518 = spicata.
placentaria, Wall. Cat. n. 2779 = cyclophora.
platycarpa, *Blume, Fl. Jav. Cupulif.* 27. t. 15.—Java.
plumbea, *Blume, Mus. Bot. Lugd. Bat.* i. 293 (sp. dub.).—Sumatra.
polyantha, Lindl. in Wall. Cat. n. 2771 = serrata.
polycarpa, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 174 = Q. Robur.
polycarpos, Kotschy, ex A. DC. in DC. Prod. xvi. II. 18 = lusitanica.
polymorpha, *Schlecht. & Cham. in Linnaea*, v. (1830) 78.—Mexic.
polyneura, Miq. Pl. Jungh. 11 = lineata.
polystachya, *Wall. Cat.* n. 2789.—Burma.
pontica, *C. Koch, in Linnaea*, xxii. (1849) 319.—As. Min.
prasina, Bosc, ex Endl. Suppl. iv. II. 25 = humilis.
prasina, Pers. Syn. ii. 568 = Q. Ilex.
prinodes, Voigt, Hort. Suburb. Calc. 277 = Castanopsis indica.
prinoides, Rafin. Alsog. Am. 22 (Quid?).—Am. bor.
prinoides, Willd. Sp. Pl. iv. 440 = Q. Prinus.
Prinus, M. A. Curt. in Rep. Geol. Surv. N. Carol. (1880) 33, partim = Michauxii.
Prinus, *Linn. Sp. Pl.* 995.—Am. bor.
pruinosa, *Blume, in Verh. Batav. Gen.* ix. (1825) 217.—Java.
pruinosa, Mor. ex A. DC. in DC. Prod. xvi. II. 86 = Q. Pseudo-molucca.
Pseud-ilex, *Chatin, in Bull. Soc. Bot. Fr.* xvi. (1869) 22.—Europ.
Pseudo-aegilops, Petz. Arb. Muscav. 627 = Q. Robur.
Pseudo-annulata, Blume, Mus. Bot. Lugd. Bat. i. 299 = Teysmannii.
Pseudo-cerris, Boiss. Diagn. Ser. I. xii. 118 = Q. Cerris.
Pseudo-coccifera, *Desf. Fl. Atlant.* ii. 349.—Reg. Mediterr.
Pseudo-coccifera, Labill. Ic. Pl. Syr. Dec. v. 9. t. 6 = coccifera.
Pseudo-dchorochensis, Boiss. Fl. Orient. iv. 1165 = Q. Robur.
Pseudo-molucca, *Blume, in Verh. Batav. Gen.* ix. (1825) 214.—Java.

QUERCUS:—
Pseudo-molucca, Mor. ex A. DC. in DC. Prod. xvi. II. 85 = spicata.
Pseudo-pedunculata, Vukot. in Rad Jugos. Akad. Zagreb. xliv. (1878) 209 – Q. Robur.
Pseudo-sessilis, Schur, Enum. Pl. Transs. 608 = Q. Robur.
Pseudo-suber, *Santi, Viag. Tosc.* i. 156. t. 3; *Desf. Fl. Atlant.* ii. 348.—Europ.
Pseudo-tscharakensis, Kotschy, ex A. DC. in DC. Prod. xvi. II. 8 = Q. Robur.
pubescens, Brot. Fl. Lusit. ii. 31 = Q. Toza.
pubescens, Szovits, ex A. DC. in DC. Prod. xvi. II. 13 = castaneaefolia.
pubescens, Willd. Berl. Baumz. 279 = Q. Robur.
pubinervis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 211.—Mexic.
pulchella, *Humb. & Bonpl. Pl. Aequin.* ii. 44. t. 88.—Mexic.
pulverulenta, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
pumila, Michx. Fl. Bor. Am. ii. 196 = Q. Prinus.
pumila, Walt. Fl. Carol. 234 = Q. Phellos.
pungens, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 171 = grisea.
purpurea, Lodd. ex Loud. Arboret. iii. 1735 = Q. Robur.
Pyrami, Kotschy, Eichen, t. 3 = Q. Aegilops.
pyramidalis, C. C. Gmel. Fl. Bad. iii. 699 = Q. Robur.
pyrenaica, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 70 = Q. Robur.
pyrenaica, Willd. Sp. Pl. iv. 451 = Q. Toza.
pyrifolia, Blume, Mus. Bot. Lugd. Bat. i. 304 = spicata.
Quexigo, Cook, ex Willk. & Lange, Prod. Fl. Hisp. i. 240 = lusitanica.
quinqueloba, Engelm. in Trans. Acad. Sc. St. Louis, iii. (1878) 542 = nigra.
racemosa, Jack, in Malay. Misc. ii. (1822) n. VII. 86 = spicata.
racemosa, Lam. Encyc. i. 715 = Q. Robur.
Ragnal, Lodd. Cat. ex Loud. Arboret. iii. 1849 = Q. Cerris.
Raineri, *Bèreng. Guir. Forest.* t. 1; *ex Pritz. Ic. Ind.* ii. 238.—Hab.?
Rajah, *Hance, in Journ. Bot.* xvi. (1878) 198.—Malaya.
Ransomi, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 23 = lobata.
Rassa, *Miq. Fl. Ind. Bat. Suppl.* 350.—Malaya.
Ravenscroftiana, A. Murr. in Gard. Chron. (1866) 1018 (sp. dub.).—Am. austr.
Raynal, C. Koch, Dendrol. ii. II. 77 = Q. Cerris.
recurvans, Kotschy, ex A. DC. in DC. Prod. xvi. II. 56 = coccifera.
regalis, Burnet, ex Endl. Gen. Suppl. iv. II. 24 = Q. Robur.
regia, Lindl. Bot. Reg. (1840) Misc. 41 = Libani.
Reinwardtii, *Korth. Verh. Nat. Gesch. Bot.* 211.—Sumatra.
repanda, Benth. Pl. Hartw. 55 = microphylla.
repanda, *Humb. & Bonpl. Pl. Aequin.* ii. 31. t. 79.—Mexic.
repanda, Michx. Hist. Chênes Am. t. 5 = alba.
resinosa, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 182 = magnoliaefolia.
reticulata, *Humb. & Bonpl. Pl. Aequin.* ii. 40. t. 86.—Am. bor. occ.
retusa, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 187 = virginiana.
retusa, Rafin. Alsog. Am. 22 (Quid?).—Am. bor.
reversa, Benth. in Hook. Kew Journ. vi. (1854) 112 = thalassica.
rhamnifolia, Miq. Fl. Ind. Bat. i. I. 853 = Castanopsis rhamnifolia.
rhioënsis, Hance, in Journ. Bot. xvi. (1878) 198 = spicata.
Richardii, *Bosc, in Mém. Inst. Par.* [viii.] (1807) 326.—Hispan.
rigida, C. Koch, ex A. DC. in DC. Prod. xvi. II. 18 = lusitanica.
rigida, Willd. Sp. Pl. iv. 434 = coccifera.
Robur, Asso, Syn. Stirp. Arag. 136 = Q. Toza.
Robur, *Linn. Sp. Pl.* 996.—Europ.; As. occ.
Robur, Pall. Fl. Ross. ii. 3 = mongolica.
rosacea, Bechst. Forstbot. ed. V. 210 = Q. Robur.
rostrata, Vukot. in Rad Jugos. Akad. Zagreb. li. 1880) 17 (Quid?).—Europ.

QUERCUS:—
rotundata, *Blume, in Verh. Batav. Gen.* ix. (1825) 219. —Java.
rotundifolia, Lam. Encyc. i. 723 = Q. Ilex.
Roxburghii, Endl. Gen. Suppl. iv. II. 28 = serrata.
rubens, Petz. & Kirchn. Arb. Muscav. 629 = Q. Robur.
rubra, Benth. Pl. Hartw. 337 = Kelloggii.
rubra, *Linn. Sp. Pl.* 996.—Am. bor.
rubra, Rugel, ex A. DC. in DC. Prod. xvi. II. 63 = nigra.
Rudkini ×, Britton, in Bull. Torrey Bot. Club, ix. (1882) tt. 10–12.—Am. bor.
rugosa, Née, in Anal. Cienc. Nat. iii. (1801) 275 = crassifolia.
rugulosa, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 209.—Mexic.
rumelica, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 353, in obs. = Q. Robur.
runcinata, Engelm. in Trans. Acad. Sc. St. Louis, iii. (1878) 542 = rubra.
Sada, [Mast. in] Gard. Chron. (1875) II. 685 = rubra?
Sadleriana, *R. Br. Campst. in Ann. & Mag. Nat. Hist.* Ser. IV. vii. (1871) 249.—Am. bor. occ.
salicifolia, Benth. Pl. Hartw. 56 = tlapuxahuensis.
salicifolia, Hort. ex Steud. Nom. ed. II. ii. 428 = Q. Robur.
salicifolia, *Née, in Anal. Cienc. Nat.* iii. (1801) 265.—Mexic.
salicina, *Blume, Mus. Bot. Lugd. Bat.* i. 305.—Japon.
Salzmanniana, Webb, Iter Hisp. 12 = lusitanica.
San-Sabeana, Buckley, in Young, Bot. Texas, 507; ex Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 145 = Durandii.
sapotaefolia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 185.—Guatemala.
Sartorii, Botteri, ex A. DC. in DC. Prod. xvi. II. 66 = acutifolia.
Sartorii, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 177.—Mexic.
sclerophylla, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 59. t. 37.—China.
scolopendrifolia, Hort. ex C. Koch, Dendrol. ii. II. 28 = Q. Robur.
scyphigera, *Hance, in Journ. Bot.* xvi. (1878) 199.—Ins. Banca.
scytophylla, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 180.—Mexic.
Secondati, Steud. Nom. ed. I. 674 = Q. Cerris.
Seemanni, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 188.—Am. austr.
segoviensis, *Liebm. l. c.* 186.—Nicaragua.
semecarpifolia, *Sm. in Rees, Cycl.* xxix. n. 20.—Reg. Himal.
semiserrata, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 641. —Reg. Himal.
sempervirens, Mill. Gard. Dict. ed. VIII. n. 3 = Q. Ilex.
sempervirens, Walt. Fl. Carol. 234 = virginiana.
sericea, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 361.—Ins. Banca.
sericea, Willd. Sp. Pl. iv. 424 = Q. Phellos.
Serra, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 174.—Mexic.
serrata, Roxb. Hort. Beng. 68; Fl. Ind. iii. 641 = Castanopsis indica.
serrata, *Thunb. Fl. Jap.* 176.—Reg. Himal.; China; Japon.
serratifolia, Benth. ex Petz. & Kirch. Arb. Muscav. 632 = Libani?
sessiliflora, Salisb. Prod. 392 = Q. Robur.
sessilifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 305 (sessifolia).—Japon.
sessilis, Ehrh. Beitr. v. 161 = Q. Robur.
setifera, *Hort.—Cf. Gard. Chron.* (1874) I. 726; *et ex T. Moore, in Flor. & Pomol.* (1875) 199. f. 8.—Japon.
Shumardii, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 444 = discolor?
Sibthorpii, Kotschy, ex Boiss. Fl. Orient. iv. 1169 = coccifera.
sideroxyla, *Humb. & Bonpl. Pl. Aequin.* ii. 39. t. 85.—Mexic.
Sieboldiana, Blume, Mus. Bot. Lugd. Bat. i. 290 = thalassica.
silvicolarum, *Hance, in Journ. Bot.* xxii. (1884) 229. —China.

QUERCUS:—
sinuata, Martr. & Timb. in Bull. Soc. Bot. Fr. xi. (1864) p. xi = Q. Ilex.
sinuata, Walt. Fl. Carol. 235 = alba?
Sirokasi, *Siebold, in Verh. Batav. Gen.* xii. (1830) 26. —Japon.
Skinneri, *Benth. Pl. Hartw.* 90.—Mexic.
Smilax, Linn. Sp. Pl. 994 = Q. Ilex.
sonchifolia, Booth, ex Petz. & Kirch. Arb. Muscav. 658 (Quid?).—Hab.?
sonomensis, Benth. ex A. DC. in DC. Prod. xvi. II. 62 = Kelloggii.
sororia, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 175.—Mexic.
sphacelata, *Blume, Mus. Bot. Lugd. Bat.* i. 304 (sp. dub.).—Java.
sphaerocarpa, Vukot. in Schloss. & Vukot. Bilinar, 472 = Q. Robur.
spicata, Humb. & Bonpl. Pl. Aequin. ii. 46. t. 89 = reticulata.
spicata, *Sm. in Rees, Cycl.* xxix. n. 12.—Reg. Himal.; Malaya.
spinulosa, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 218 = crassifolia.
splendens, *Née, in Anal. Cienc. Nat.* iii. (1801) 275.—Mexic.
squamata, Roxb. Hort. Beng. 68; Fl. Ind. iii. 638 = spicata.
squarrosa, *Kotschy, ex A. DC. in DC. Prod.* xvi. II. 108.—Syria.
stellata, Wangenh. Forstw. 78. t. 6. f. 15 = obtusiloba.
stipularis, Humb. & Bonpl. Pl. Aequin. ii. 47. t. 90 = splendens.
stolonifera, Lapeyr. Hist. Abr. Pl. Pyr. 582 = Q. Toza.
Streimeri, Heuff. ex Freyn, in Verh. Zool.-Bot. Ges. Wien, xxvii. (1878) 422 = Q. Robur.
striata, *Siebold, ex André, in Illustr. Hortic.* xvii. (1870) 242. t. 44.—Japon.
strigosa, Wierz. ex Rochel, Bot. Reise Banat. 73 (Quid?). —Europ.
strombocarpa, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 176.—Mexic.
subalpina, Kotschy, ex A. DC. in DC. Prod. xvi. II. 9 = Q. Robur.
Suber, Kotschy, Eichen, t. 33 = Q. Pseudo-suber.
Suber, Linn. Sp. Pl. 995 = Q. Ilex.
subera, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = praec.
suberoides, *Rafin. in Med. Repos. N. York*, v. (1808) 360.—Sicilia.
suberosa, Salisb. Prod. 392 = Q. Ilex.
sublobata, Kit. in Schult. Oester. Fl. ed. II. i. 619 = Q. Robur.
subvelutina, Schur, Enum. Pl. Transs. 606 = Q. Robur.
sundaica, *Blume, in Verh. Batav. Gen.* ix. (1825) 216. —Malaya.
sundaica, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 207 (excl. syn. Naves) = Companoana.
susedana, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 15 = lanuginosa.
synbalanos, *Hance, in Journ. Bot.* xxii. (1884) 228.—Ins. Hongkong.
syriaca, Kotschy, Eichen, t. 1 = lusitanica.
syspirensis, C. Koch, in Linnaea, xxii. (1849) 328 = Q. Robur.
tardiflora, Czern. ex Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) I. 385 = Q. Robur.
tauricola, Kotschy, Eichen, t. 10 = lusitanica.
Tauza, Desf. Tabl. ed. II. 245 = Q. Toza.
Tauzin, Pers. Syn. ii. 571 = Q. Toza.
Tchihatcheffii, Kotschy, ex A. DC. in DC. Prod. xvi. II. 48 = seq.
Tchihatchewi, Kotschy, in Tchihat. Asie Min. Bot. ii. 468 = Libani.
Tema, Buch.-Ham. ex Wall. Cat. n. 3733 = lanceaefolia.
Tennesi, Hort. ex Wesmael, in Bull. Congr. Bot. Brux. (1864) 275 = Q. Robur.
tenuicola, Boiss. Fl. Orient. iv. 1167, in syn. sphalm. = *tauricola*.
texana, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 444 = rubra.
Teysmannii, *Blume, Mus. Bot. Lugd. Bat.* i. 300 (Tysmannii).—Java; Sumatra.
thalassica, *Hance, in Hook. Kew Journ.* i. (1849) 176. —China.
thelecarpa, Miq. Pl. Jungh 9 = Q. Pseudo-molucca.

QUERCUS:—
Thirkeana, C. Koch, in Linnaea, xix. (1847) 14 = lusitanica.
Thomasii, Tenore, Ind. Sem. Hort. Neap. (1825) 11. n. 21 = Q. Robur.
Thomsoni, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 109.—Reg. Himal.
Thomsoniana, A. DC. in DC. Prod. xvi. II. 98 = lineata.
tinctoria, Bartr. Trav. ed. II. (1794) 37; Michx. Hist. Chênes Am. tt. 24, 25 = discolor.
tlapuxahuensis, *A. DC. in DC. Prod.* xvi. II. 29.—Mexic.
tolimensis, *Humb. & Bonpl. Pl. Aequin.* ii. 153. t. 129. —N. Granat.
tomentella, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1878) 393.—Calif.
tomentosa, Ehrh. ex A. DC. in DC. Prod. xvi. II. 6 = Q. Robur.
tomentosa, *Willd. Sp. Pl.* iv. 437.—Mexic.
Tommasini, Kotschy, ex Vis. Fl. Dalm. Suppl. 45 = Q. Robur.
torulosa, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 14 (Quid?).—Europ.
Tosa, Loisel. in Duham. Arb. ed. Nov. vii. 179 = Q. Toza.
totutlensis, *A. DC. in DC. Prod.* xvi. II. 62.—Mexic.
Tournefortii, Willd. Sp. Pl. iv. 453 = Q. Cerris.
Toza, *Gillet, ex Bosc, in Lam. Journ. Hist. Nat.* ii. (1792) 115. t. 2. f. 3.—Europ. austr.
Toza, Griseb. Spicil. Fl. Rumel. ii. 337 = Q. Robur.
tribuloides, Sm. in Rees, Cycl. xxix. n. 13 = Castanopsi tribuloides.
tricolor, Hort. ex Petz. & Kirch. Arb. Muscav. 624 = Q. Robur.
tridens, Humb. & Bonpl. Pl. Aequin. ii. 35. t. 82 = Q. Castanea.
tridentata, Engelm. ex A. DC. in DC. Prod. xvi. II. 64 = nigra.
triloba, Michx. Hist. Chênes Am. t. 26 = falcata.
tristis, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 174 = Q. Castanea.
trojana, Kotschy, ex A. DC. in DC. Prod. xvi. II. 46 = seq.?
trojana, Webb, in Loud. Gard. Mag. xv. (1839) 590; et Jaub. & Spach, Illustr. i. t. 57 A = Q. Aegilops.
tuberculata, Liebm. in. Overs. Danske Vidensk. Selsk. Forh. (1854) 181 = polymorpha.
turbinata, Blume, Bijdr. 523 = lineata.
turbinata, Kit. in Linnaea, xxxii. (1863) 353 = Q. Robur.
turbinata, Liebm. in Overs. Danske Vidensk. Selsk. Forh. (1854) 186 = guatimalensis.
turbinata, Roxb. Hort. Beng. 68; Fl. Ind. iii. 636 = Thomsoni.
Turneri, Hort. ex A. DC. in DC. Prod. xvi. II. 43 = Q. Pseudo-suber.
Turneri, Willd. Enum. Hort. Berol. 975 = lusitanica.
Tysmannii, Blume, Mus. Bot. Lugd. Bat. i. 300, sphalm. = Teysmannii.
uliginosa, Wangenh. Forstw. 80. t. 6 = nigra.
umbonata, Hance, in Journ. Bot. xii. (1874) 241 = cyclophora.
umbrosa, Endl. Gen. Suppl. iv. II. 26 = calophylla.
undulata, Benth. Pl. Hartw. 81 = Benthami.
undulata, Engelm. in Trans. Acad. Sc. St. Louis, iii. (1878) 392, partim = Durandii.
undulata, Kit. in Linnaea, xxxii. (1863) 354 (Reliq. Kitaib. 110) = Q. Robur.
undulata, C. Koch, l. c. xix. (1847) 14 = lusitanica.
undulata, *Torr. in Ann. Lyc. N. York*, ii. (1828) 248. t. 4.—Am. bor. occ.
Ungeri, Kotschy, Eichen, t. 13 = Q. Aegilops.
urceolaris, *Jack, in Malay. Misc.* ii. (1822) VII. 87.—Sumatra.
urticaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 296.—China; Japon.
uvariifolia, *Hance, in Journ. Bot.* xxii. (1884) 227.—China.
vaccinifolia, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 106 = chrysolepis.
valentina, Cav. Ic. ii. 25 = lusitanica.
valentina, Lam. ex Steud. Nom. ed. I. 675 = discolor.
valida, Kotschy, ex A. DC. in DC. Prod. xvi. II. 55 = coccifera.
Vallonea, A. DC. l. c. 45 = Q. Aegilops.
Vallonea, Kotschy, Eichen, t. 7 = Q. Cerris.
variabilis, *Blume, Mus. Bot. Lugd. Bat.* i. 297.—Japon.

QUERCUS:—
varians, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 214 = polymorpha.
variegata, Lodd. ex Steud. Nom. ed. II. ii. 428 = Q. Cerris.
variifolia, Sweet, Hort. Brit. ed. II. 465 = Q. Ilex.
varingaefolia, Miq. Pl. Jungh. 12 = javensis.
Vaseyana, *Buckl. in Bull. Torrey Bot. Club*, x. (1883) 91.—Texas.
velutina, Lam. Encyc. i. 721 = discolor?
velutina, L'Herit. ex A. DC. in DC. Prod. xvi. II. 21 = bicolor.
velutina, *Lindl. in Wall. Pl. As. Rar.* ii. 41. t. 150.—Burma.
verrucosa, Rafin. Alsog. Am. 26 (Quid?).—Am. bor.
versicolor, Rafin. l. c. 20 (Quid?).—Am. bor.
vertesiensis, *Kit. in Linnaea*, xxxii. (1863) 354, nomen.—Hungar.
vesca, Kotschy, Eichen, t. 36 = Libani.
Vibrayeana, *Franch. & Sav. Enum. Pl. Jap.* ii. 498.—Japon.
Vidalii, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 209.—Ins. Philipp.
villosa, Walt. Fl. Carol. 235 = obtusiloba?
viminalis, Bosc, in Mém. Inst. Par. [viii.] (1807) 316 = Q. Robur.
virens, Ait. Hort. Kew. ed. I. iii. 356 = virginiana.
virgata, Martr. Fl. Tarn, 641 = Q. Robur.
Virgiliana, Tenore, Fl. Nap. v. 262 = Q. Robur.
virginiana, *Mill. Gard. Dict.* ed. VIII. n. 16.—Am. bor.
vulcanica, Boiss. ex Kotschy, Eichen, t. 18 = Q. Robur.
Wallichiana, *Lindl. in Wall. Cat.* n. 2778.—Malaya.
Warszewiczii, *Liebm. in Overs. Danske Vidensk. Selsk. Forh.* (1854) 187.—Guatemala.
Welandii, Simk. in Magyar Növényt. Lapok, vii. (1883) 67 = Q. Robur.
Wislizeni, *A. DC. in DC. Prod.* xvi. II. 67.—Calif.
Woodii, *Hance, in Journ. Bot.* xii. (1874) 240.—Ins. Philipp.
xalapensis, *Humb. & Bonpl. Pl. Aequin.* ii. 24. t. 75.—Mexic.
xylina, [Scheidw.] Hortic. Belge, iv. (1837) 321. t. 18 = reticulata.
xylocarpa, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 196.—Assam; Burma.

QUERIA, Loefl. It. Hisp. 48 (1752). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 152.
argyrocoma, Eaton, N. Am. Bot. ed. VII.; ex Eaton & Wright, N. Am. Bot. ed. VIII. 343 = Paronychia argyrocoma.
canadensis, Linn. Sp. Pl. 90 = Anychia dichotoma.
capillacea, Nutt. Gen. Am. i. 159 = Anychia dichotoma.
chilensis, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 428 = Donatia fascicularis?
dichotoma, Maxim. ex Franch. & Sav. Enum. Pl. Jap. i. 405 = *trichotoma*.
dichotoma, Moench, Meth. 351 = Anychia dichotoma.
hispanica, *Linn. Sp. Pl.* 90.—Reg. Mediterr. & Caucas.
multiflora, Hort. ex Steud. Nom. ed. I. 675 = Anychia dichotoma.
sessiliflora, Eaton, N. Am. Bot. ed. VII.; ex Eaton & Wright, N. Am. Bot. ed. VIII. 343 = Paronychia sessiliflora.
trichotoma, Thunb. in Trans. Linn. Soc. ii. (1794) 329 = Wikstroemia japonica.

QUESNELIA, Gaudich. Bot. Voy. Bonite, t. 54 (1844–49). *BROMELIACEAE*, Benth. & Hook. f. iii. 665.
GUESMELIA, Walp. Ann. i. 841 (1849).
LIEVENA, Regel, Gartenfl. (1880) 289.
Augustocoburgi, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 150.—Bras.
centralis, Wawra, l. c. = lateralis.
lateralis, *Wawra, l. c.* 149.—Bras.
roseo-marginata, Carr. in Rev. Hortic. (1880) 70; et ex E. Morr. in Belg. Hortic. xxxi. (1881) 82. t. 4 = rufa, *Gaudich.*
rufa, *Gaudich. Bot. Voy. Bonite*, t. 54.—Bras.
rufa, *E. Morr. in Belg. Hortic.* xxxii. (1882) 115. t. 4–6.—Guian.
Skinneri, Hort. ex E. Morr. l. c. xxxi. (1881) 82 = rufa, *Gaudich.*
strobilospica, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 149.—Bras.

QUESNELIA:—
Van Houttei, *E. Morr. in Belg. Hortic.* xxxi. (1881) 163, 350. t. 18.—Bras.

QUETELETIA, Blume, Orch. Archip. Ind. 117. t. 37 (1858) = **Physurus**, Rich. (Orchid.).
plantaginifolia, Blume, l. c. 118 (= *Orchipedum plantaginifolium*).—Java.

QUIINA, Aubl. Pl. Gui. Suppl. 19. t. 379 (1775). *GUTTIFERAE*, Benth. & Hook. f. i. 176.
GUIINA, Crueg. in Linnaea, xx. (1847) 115.
ROBINSONIA, Scop. Introd. 218 (1777).
TOUROULIA, Aubl. Pl. Gui. 492. t. 194 (1775).
crenata, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 163.—Guian.
Cruegeriana, *Griseb. Fl. Brit. W. Ind.* 106.—Ins. Trinit.
Decaisneana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 315.—Guian.
florida, *Tul. l. c.* Sér. III. xi. (1849) 167.—Bras.
guianensis, *Aubl. Pl. Gui. Suppl.* 19.—Guian.
guianensis, Crueg. ex Griseb. Fl. Brit. W. Ind. 106 = Cruegeriana.
jamaicensis, *Griseb. l. c.* 105.—Jamaic.
leptoclada, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 159.—Guian.
longifolia, *Spruce, ex Planch. & Triana, l. c.* Sér. IV. xv. (1861) 314.—Bras.
macrophylla, *Tul. l. c.* Sér. III. xi. (1849) 164.—Venezuela.
macrostachya, *Tul. l. c.* 162.—Peruv.
micrantha, *Tul. l. c.* 160.—Bras.
obovata, *Tul. l. c.* 157.—Guian.
Poeppigiana, *Tul. l. c.* 161.—Bras.
rhytidopus, *Tul. l. c.* 166.—Bras.
sessilis, *Choisy, Planch. & Triana, l. c.* Sér. IV. xv. (1861) 312.—Guian.
ternatiflora, *Wright, in Sauv. Fl. Cub.* 9.—Cuba.
tinifolia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 311.—Bras.

QUILAMUM, Blanco, Fl. Filip. ed. I. 851 (1837) = **Crypteronia**, Blume (Lythrar.).
luteum, Blanco, l. c. = C. pubescens.

QUILESIA, Blanco, Fl. Filip. ed. I. 176 (1837) = **Chailletia**, DC.
sericea, Blanco, l. c. 177 = C. Griffithii.

QUILIUSA, Hook. f. in Journ. Linn. Soc. x. (1869) 460, sphalm. = QUELUSIA, Vand. = **Fuchsia**, Linn. (Onagrar.)

QUILLAIA, Molina = seq.

QUILLAJA, Molina, Sagg. Chile, 354 (1782). *ROSACEAE*, Benth. & Hook. f. i. 614.
CULLAY, Molina, ex Steud. Nom. ed. II. i. 452 (1840).
FONTENELLEA, A. St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 141. t. 7.
SMEGMADERMOS, Ruiz & Pav. Prod. 133. t. 31 (1794).
brasiliensis, *Mart. Syst. Mat. Med. Bras.* 127.—Bras.
lanceolata, D. Dietr. Syn. Pl. ii. 1616 = seq.
lancifolia, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March 1831) 231.—Bras.
Molinae, DC. Prod. ii. 547 = Q. Saponaria.
petiolaris, D. Don, in Edinb. N. Phil. Journ. (Oct.–Dec. 1831) 110 = Azara Gilliesii.
Poeppigii, *Walp. Rep.* ii. 52.—Chili.
Saponaria, *Molina, Sagg. Chile*, 175, 354.—Chili.
Saponaria, Poir. Encyc. vi. 33 = Q. Smegmadermos.
Sellowiana, *Walp. Rep.* ii. 52.—Bras.
smegmadermis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.
Smegmadermos, *DC. Prod.* ii. 547.—Chili.
Smegmadermos, Poepp. ex Walp. Rep. ii. 52 = Poeppigii.

QUINARIA, Lour. Fl. Cochinch. 272 (1790) = **Clausena**, Burm. f. (Rutac.).
Lansium, Lour. l. c. = C. Wampi.

QUINARIA, Rafin. Med. Fl. ii. 122 (1830) = **Vitis**, Tourn. (Ampel.).
hederacea, Rafin. l. c. = V. hederacea.
hirsuta, Rafin. l. c. = V. hederacea.
japonica, Rafin. Sylva Tellur. 87 = V. heterophylla.

QUINASIS, Rafin. Sylva Tellur. 150 (1838) = **Polylepis**, Ruiz & Pav. (Rosac.).
villosa, Rafin. l. c. = villosa.

QUINCHAMALA, Willd. Sp. Pl. i. 1217 (1797) = seq.

QUINCHAMALIUM, Molina, Sagg. Chile, ed. I. 151 (1782). *SANTALACEAE*, Benth. & Hook. f. iii. 220.
andinum, *Phil. in Bot. Zeit.* xv. (1857) 747.—Chili.
Berteroanum, *Phil. l. c.* 748.—Chili.
bracteosum, *Phil. l. c.* 747.—Chili.
breviflorum, Presl, Epim. Bot. 246 = majus.
carnosum, *Phil. Fl. Atac.* 48.—Chili.
chilense, Lam. Illustr. ii. 125. t. 142 = procumbens.
chilense, *Molina, Sagg. Chile*, 151.—Chili.
chilense, Phil. ex A. DC. in DC. Prod. xiv. 625 = majus.
chilense, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 412 = gracile.
chilense, Willd. Sp. Pl. i. 1217 = linifolium.
Dombeyi, Brongn. in Duperr. Voy. Coq. Bot. t. 52 = procumbens.
elegans, *Presl, Epim. Bot.* 246.—Chili.
ericoides, Brongn. in Duperr. Voy. Coq. Bot. t. 52 = gracile.
excrescens, *Phil. in Bot. Zeit.* xv. (1857) 748.—Chili.
fruticulosum, *Steud. Nom.* ed. II. ii. 429.—*Cf. Miers, in Journ. Linn. Soc.* xvii. (1878) 138.—Chili.
gracile, Brongn. in Duperr. Voy. Coq. Bot. 231. t. 52.—Chili.
linarioides, *Phil. in Bot. Zeit.* xv. (1857) 746.—Chili.
linifolium, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 412 = procumbens.
linifolium, *Miers, in Journ. Linn. Soc.* xvii. (1878) 135.—Chili.
litorale, *Phil. in Linnaea*, xxxiii. (1864–65) 234.—Chili.
majus, *Brongn. in Duperr. Voy. Coq. Bot.* 229. t. 51.—Chili.
minutum, Phil. in Linnaea, xxxiii. (1864–65) 233 = gracile.
parviflorum, *Phil. in Bot. Zeit.* xv. (1857) 747.—Chili.
patagonicum, *Phil. f. in Anal. Soc. Cient. Argent.* x. (1880) 143.—Patagon.
patagonicum, Spreng. Syst. i. 537 = Arjona tuberosa.
pratense, *Phil. in Bot. Zeit.* xv. (1857) 747.—Chili.
procumbens, *Ruiz & Pav. Fl. Per.* ii. 1.—Peruv.
purpureum, *Phil. in Bot. Zeit.* xv. (1857) 747.—Chili.
rugosum, Phil. in Linnaea, xxxiii. (1864–65) 233 = majus.
tenue, Steud. Nom. ed. II. ii. 429 = gracile.
thesioides, *Phil. Fl. Atac.* 48.—Chili.

QUINCULA, Rafin. Atl. Journ. 145 (1832) = **Physalis**, Linn. (Solanac.).
lobata, Rafin. l. c. = P. lobata.

QUINETIA, Cass. in Dict. Sc. Nat. lx. 590 (1830). *COMPOSITAE*, Benth. & Hook. f. ii. 316.
Urvillei, *Cass. l. c.* 591.—Austral.

QUINIO, Schlecht. in Linnaea, xxvi. (1854) 732 = **Cocculus**, DC. (Menisp.).
cocculoides, Schlecht. l. c. = C. macrocarpus.

QUINQUEDULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Litsea**, Lam. (Laurin.).

QUINQUEFOLIUM, Tourn. ex Adans. Fam. ii. 295 (1763) = **Potentilla**, Linn. (Rosac.).

QUINQUELOBUS, Benj. in Linnaea, xx. (1847) 316. '*GENUS SPURIUM*.'—Cf. Benth. & Hook. f. ii. 987.
BENJAMINIA, Mart. Fl. Bras. x. 255 (1847).
glaber, Benj. in Linnaea, xx. (1847) 316 = Limnophila gratioloides.
minor, Benj. l. c. 317.—Asia.
splendens, Benj. l. c. 316.—Asia.
utricula riaeformis, Benj. in Mart. Fl. Bras. x. 255 = seq.
utricula riaeoides, Benj. in Linnaea, xx. (1847) 316.—Bras.

QUINQUELOCULARIA, C. Koch, in Linnaea, xxiii. (1850) 630 = **Campanula**, Tourn.
crispa, C. Koch, l. c. 631 = C. crispa.

QUINQUINA, Condam. in ' Act. Acad. Par. (1738) 226' = **Cinchona**, Linn. (Rubiac.).

QUINSONIA, Montr. in Mém. Acad. Lyon, x. (1860) 178 = **Pittosporum**, Banks.
coccinea, Montr. l. c. 179.—N. Caled.

QUINTILIA, Endl. Gen. 1408 (1841) = **Stauranthera**, Benth. (Gesner.).
coerulea, DC. Prod. ix. 278 = S. ecalcarata.

QUINTINIA, A. DC. Monog. Camp. 92, in adnot. (1830). *SAXIFRAGEAE*, Benth. & Hook. f. i. 645.
elliptica, *Hook. f. Fl. N. Zel.* i. 78.—N. Zel.
Fawkneri, *F. Muell. Fragm.* vi. 92.—Austral.
serrata, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 356.—N. Zel.
Sieberi, *A. DC. Monog. Camp.* 92, in adnot.—Austral.
Verdonii, *F. Muell. Fragm.* ii. 125.—Austral.

QUIRINA, Rafin. Sylva Tellur. 102 (1838) = **Cuphea**, P. Br. (Lythrar.).
microphyla, Rafin. l. c. = C. microphylla.

QUIRIVELIA, Poir. Encyc. vi. 42 (1804) = **Ichnocarpus**, R. Br. (1809) (Apocyn.).
zeylanica, Poir. l. c. = I. frutescens.

QUIROSIA, Blanco, Fl. Filip. ed. II. 398 (1845) = **Crotalaria**, Dill. (Legumin.).
anceps, Blanco, l. c. = C. verrucosa.
secunda, Blanco, l. c. = C. linifolia.

QUISQUALIS, Linn. Sp. Pl. ed. II. 556 (1762) = *COMBRETACEAE*, Benth. & Hook. f. i. 689.
KLEINIA, Crantz, Inst. ii. 488 (1766).
SPHALANTHUS, Jack, in Malay. Misc. ii. (1822) n. VII. 55 (1822).
bibracteata, *Regel, Cat. Pl. Hort. Aksakov.* 118, nomen. —Hab.?
densiflora, *Wall. Cat.* n. 4011.—Burma; Malaya.
ebracteata, Beauv. Fl. Owar. i. 57. t. 5 = indica.
glabra, Burm. f. Fl. Ind. 104. t. 28. f. 2 = indica.
grandiflora, Miq. in Journ. Bot. Néerl. i. (1861) 119 = indica.
indica, Blanco, Fl. Filip. ed. I. 361 = malabarica.
indica, *Linn. Sp. Pl.* ed. II. 556.—As. trop.
longiflora, Presl, Epim. Bot. 216 = indica.
Loureiri, G. Don, Gen. Syst. ii. 667 = indica.
madagascariensis, *Boj. Hort. Maurit.* 134, nomen.—Madag.
malabarica, *Bedd. Ic. Pl. Ind. Or.* i. 33. t. 155.—Ind. or.
obovata, Schum. & Thonn. Beskr. Guin. Pl. 218 = indica.
parviflora, *Gerr. ex Harv. & Sond. Fl. Cap.* ii. 512. —Afr. austr.
pubescens, Burm. f. Fl. Ind. 104. t. 35. f. 2 = indica.
sinensis, Lindl. Bot. Reg. (1844) t. 15 = indica.
spinosa, Blanco, Fl. Filip. ed. II. 254 = malabarica.
spinosa, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 81 = indica.
villosa, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 426 = indica.

QUIVISIA, Comm. ex Juss. Gen. 264 (1789). *MELIACEAE*, Benth. & Hook. f. i. 330.
CALODRYUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 401. t. 57.
GILIBERTIA, J. F. Gmel. Syst. 682 (1791).
GINNANIA, M. Roem. Syn. Hesper. 79, 90 (1846).
PAYERIA, Baill. Adansonia, i. (1860–61) 50. t. 3.
anomala, *O. Hoffm. Sert. Madag.* 10.—Madag.
chilosantha, *Boj. Hort. Maurit.* 58.—Ins. Maurit.
decandra, *Cav. Diss.* vii. 367. t. 211.—Ins. Mascar.
filipes, Baker, Fl. Maurit. 46 = trichopoda?
fruticosa, J. F. Gmel. Syst. 633 = oppositifolia?
heterophylla, *Cav. Diss.* vii. 368. t. 213.—Ins. Mascar.
heterophylla, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = ovata.
laciniata, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 12; *et in Baker, Fl. Maurit.* 46.—Ins. Rodrig.
laciniosa, C. DC. in DC. Monog. Phan. i. 431, sphalm. = praec.
lanceolata, C. DC. l. c. 433.—Madag.
mauritiana, Baker, Fl. Maurit. 45 = ovata, heterophylla, decandra, oppositifolia.

QUIVISIA:—
oppositifolia, *Cav. Diss.* vii. 368. t. 214.—Ins. Borbon.
ovata, *Cav. l. c.* 368. t. 212.—Ins. Borbon.
racemosa, Pers. Syn. i. 468 = decandra.
rigida, C. DC. in DC. Monog. Phan. i. 433 = chilosantha.
Sieberi, *C. DC. l. c.* 430.—Ins. Maurit.
tetramera, *C. DC. l. c.* 432.—Madag.
trichopoda, *Baill. Adansonia*, xi. (1874) 255.—Ins. Maurit.

QUOYA, Gaudich. in Freyc. Voy. Bot. 453. t. 66 (1826) = **Pityrodia**, R. Br. (Verben.).
atriplicina, F. Muell. Fragm. i. 235 = P. atriplicina.
cuneata, Gaudich. in Freyc. Voy. Bot. 454 = P. cuneata.
dilatata, F. Muell. Fragm. vi. 157 = P. dilatata.
hemigenoides, F. Muell. l. c. 156 = P. hemigenioides.
loxocarpa, F. Muell. l. c. iv. 80 = P. Drummondii.
Oldfieldii, F. Muell. l. c. = P. Oldfieldii.
paniculata, F. Muell. l. c. = P. paniculata.
racemosa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 194 = P. racemosa.
stachyodes, F. Muell. Fragm. v. 50; vi. 158 = P. racemosa.
verbascina, F. Muell. l. c. iv. 80 = P. verbascina.

R

RABDOCHLOA, Beauv. Agrost. 84 (1812) = **Leptochloa**, Beauv. (Gramin.).
cruciata, Beauv. l. c. = Chloris cruciata.
domingensis, Beauv. l. c. = L. domingensis.
monostachya, Beauv. l. c. = L. monostachya.
mucronata, Beauv. l. c. = Eleusine aegyptiaca.
virgata, Beauv. l. c. = Chloris virgata.
Vulpiastrum, De Not. Ind. Sem. Hort. Genuens. (1852) 26; et in Ann. Sc. Nat. Sér. III. xix. (1853) 372.—Nubia.

RABDOSIA, Hassk. in Flora, xxv. (1842) II. Beibl. 25 = **Plectranthus**, L'Hérit. (Labiat.).
javanica, Hassk. l. c. = P. javanicus.

RABELAISIA, Planch. in Hook. Lond. Journ. Bot. iv. (1845) 519. t. 18 = **Lunasia**, Blanco (Rutac.).
parvifolia, Planch. l. c. = L. parvifolia.
philippinensis, Planch. l. c. = L. amara.

RABENHORSTIA, Reichb. Nom. 159 (1841) = **Berzelia**, Brongn. (Bruniac.).

RACAPA, M. Roem. Syn. Hesper. 123 (1846) = **Carapa**, Aubl. (Meliac.).
procera, M. Roem. l. c. = C. procera.

RACARIA, Aubl. Pl. Gui. Suppl. 24. t. 382 (1775). Gen. dub. *SAPINDACEARUM*.—Cf. Benth. & Hook. f. i. 392.
sylvatica, *Aubl. l. c.*—Guian.
sylvestris, Miers, in Journ. Linn. Soc. xvii. (1879) 342, sphalm. = praec.

RACHIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 124; et in Abh. Akad. Berl. (1855) 68. t. 6 B = **Begonia**, Linn.
auriformis, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 124 = B. incana.
incana, Klotzsch, in Abh. Akad. Berl. (1855) 69 = B. incana.
Meyeri, Klotzsch, l. c. = B. tomentosa.
peltata, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 124 = B. incana.
polygonata, Klotzsch, ex A. DC. in DC. Prod. xv. 1. 327 = B. polygonata.

RACHICALLIS, DC. Prod. iv. 433 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 49.
RHACHICALLIS, Spach, Hist. Vég. Phan. viii. 374 (1839).
capitata, Benth. Pl. Hartw. 195 = Hedyotis Hartwegiana.

RACHICALLIS:—
caracasana, DC. Prod. iv. 434 = Mallostoma caracasanum.
nitida, DC. l. c. 433 = Mallostoma nitidum.
rupestris, *DC. l. c.* 434.—Ind. occ.
setosa, Steud. Nom. ed. II. ii. 429 = Mallostoma setosum.

RACKA, J. F. Gmel. Syst. 245 (1791) = **Avicennia**, Linn. (Verben.).
ovata, Roem. & Schult. Syst. iii. 207 = A. officinalis.
torrida, J. F. Gmel. Syst. 245 = A. officinalis.

RACLATHRIS, Rafin. Sylva Tellur. 167 (1838) = **Rochelia**, Reichb. (Borag.).
cerinthoides, Rafin. l. c. = Rochelia stellulata.

RACLETIA, Adans. Fam. ii. 449 (1763) (Quid?).

RACOMA, Willd. ex Steud. Nom. ed. II. ii. 429 (1841), sphalm. = ROCAMA, Forsk. = **Trianthema**, Sauv. (Ficoid.).

RACOUBEA, Aubl. Pl. Gui. i. 589. t. 236 (1775) = **Homalium**, Jacq. (Samyd.).
guianensis, Aubl. l. c. = H. Racoubea.

RACUA, J. F. Gmel. Syst. 1612 (Index) = RACKA, J. F. Gmel. = **Avicennia**, Linn. (Verben.).

RADACKIA, Cham. & Endl. in Ann. Wien. Mus. i. (1836) 186. Genus indesc. *LEGUMINOSARUM*. Cf. Benth. & Hook. f. i. 465.
amicorum, Cham. & Endl. l. c.—Ins. Pacif.

RADAMAEA, Benth. in DC. Prod. x. 509 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 973.
montana, *Benth. in DC. Prod.* x. 509.—Madag.
prostrata, *Benth. l. c.*—Ins. Galega.

RADDIA, Bertol. in Bologn. Opusc. Sc. iii. (1819) 410 = **Olyra**, Linn. (Gramin.).
brasiliensis, Bertol. l. c. = O. floribunda.

RADDIA, Fl. Corc. in Ionios Anthol. ii. (1834) 448 = **Crypsis**, Ait. (Gramin.).
aculeata, Fl. Corc. l. c. 450 = C. aculeata.

RADDIA, Spreng. Syst. i. 177, in syn. (1825); Miers, in Trans. Linn. Soc. xxviii. (1872) 388 = **Salacia**, Linn. (Celastrin.).
arborea, Miers, l. c. 389 = S. undulata.
calypsoides, Miers, l. c. 396 = S. calypsoides.
conferta, Miers, l. c. 394 = S. conferta.
crassifolia, Miers, l. c. 396 = S. crassifolia.
elliptica, Miers, l. c. 395 = S. elliptica.
fasciculata, Miers, l. c. 391.—Bras.
firmifolia, Miers, l. c. 393 = S. elliptica.
glomerata, Miers, l. c. 397 = S. glomerata.
grandiflora, Miers, l. c. 391 = S. grandiflora, *Peyr.*
impressifolia, Miers, l. c. 392 = S. grandiflora, *Peyr.*
lacunosa, Miers, l. c. = S. lacunosa.
oblongifolia, Miers, l. c. 393 = S. elliptica.
pachyphylla, Miers, l. c. 396 = S. pachyphylla.
verrucosa, Miers, l. c. 395 = S. verrucosa, *Wright.*

RADDISIA, Leandro Sacram. in Denkschr. Akad. Muench. vii. (1821) 237. t. 15 = **Salacia**, Linn. (Celastrin.).
arborea, Schult. Mant. i. 347 = S. undulata.

RADEMACHIA, Steud. Nom. ed. I. 676 (1821) err. typ. = seq.

RADERMACHIA, Thunb. in Vet. Akad. Handl. Stockh. (1776) 251 = **Artocarpus**, Forst. (Urtic.).
arborea, Steud. Nom. ed. II. ii. 429, sphalm. = *Raddisia arborea*.
incisa, Thunb. in Vet. Akad. Handl. Stockh. (1776) 253 = A. incisa.
integra, Thunb. l. c. 254 = A. integrifolia.

RADERMACHIA, Zoll. & Mor. in Zoll. Syst. Verz. Ind. Archip. 53 (1854) = **Stereospermum**, Cham. (Bignon.).
amoena, Seem. in Journ. Bot. viii. (1870) 146 = S. hypostictum.
Banaibana, Bur. in Baill. Adansonia, ii. (1861–62) 194 = S. Banaibanai.

gigantea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 250 = S. hypostictum.
glandulosa, Miq. l. c. = S. glandulosum.
Lobbii, Miq. l. c. = S. hypostictum.
pinnata, Seem. in Journ. Bot. viii. (1870) 147 = S. pinnatum.
quadripinna, Seem. l. c. = S. Seemannii.
stricta, Zoll. & Mor. in Zoll. Syst. Verz. Ind. Archip. 53 = S. glandulosum.

RADIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Mimusops**, Linn. (Sapot.).

RADIA, A. Rich. in Kunth, Syn. Pl. i. 300 (1822) = **Barbacenia**, Vand. (Amaryll.).
tubiflora, A. Rich. l. c. = B. tubiflora (in Addend.).

RADIANA, Rafin. Specch. i. 88 (1814); ex DC. Prod. iii. 353 = **Cypselea**, Turp. (Ficoid.).
petiolata, Rafin. l. c. = C. humifusa.

RADIATA, Medic. Phil. Bot. i. 208 (1789) = **Medicago**, Tourn. (Legumin.).

RADICULA, Dill. ex Moench, Meth. 262 (1794) = **Nasturtium**, Linn. (Crucif.).
lancifolia, Moench, l. c. = N. amphibium.
palustris, Moench, l. c. 263 = N. palustre.
pinnata, Moench, l. c. = N. sylvestre.
pyrenaica, Cav. Desc. 423 = N. pyrenaicum.

RADIOLA, [Dill.] Roth, Tent. Fl. Germ. i. 71 (1788).
LINEAE, Benth. & Hook. f. i. 242.
LINOIDES, Rupp. Fl. Jen. ed. Hall. 91 (1745).
MILLEGRANA, Adans. Fam. ii. 269 (1763).
dichotoma, Moench, Meth. 288 = R. Linoides.
filiformis, Dulac, Fl. Hautes-Pyr. 226 = R. Linoides.
linoidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.
Linoides, *Roth, Tent. Fl. Germ.* i. 71.—Europ.
Millegrana, Sm. Engl. Bot. t. 895 = R. Linoides.
multiflora, Aschers. Fl. Brandenb. 106 = R. Linoides.
tetrapetala, Steud. Nom. ed. II. ii. 429, sphalm. = R. Linoides.

RADIUSIA, Reichb. Consp. 148 (1828) = **Sophora**, Linn. (Legumin.).
alopecuroides, Reichb. ex Steud. Nom. ed. II. ii. 429 = S. alopecuroides.
chinensis, Heynh. Nom. ii. 582 = S. chinensis.
flavescens, Endl. ex Heynh. l. c. = S. flavescens.
galegoides, Heynh. l. c. = S. flavescens.
sericea, Heynh. l. c. = S. sericea.

RADOJITSKYA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 176 = **Lachnaea**, Linn. (Thymel.).
capensis, Turcz. l. c. 177 = L. axillaris.

RAFFENALDIA, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 413. *CRUCIFERAE*, Benth. & Hook. f. i. 101.
COSSONIA, Dur. in Ann. Sc. Nat. Sér. III. xx. (1853) 82. t. 6.
primuloides, *Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1853) 413.—Alger.

RAFFLESIA, R. Br. in Trans. Linn. Soc. xiii. (1821) 201. *CYTINACEAE*, Benth. & Hook. f. iii. 119.
Arnoldi, *R. Br. l. c.* 207.—Sumatra.
Cumingii, R. Br. l. c. xix. (1844) 243 = manillana.
Hasseltii, *Suring. in Veth, Midden-Sumatra*, iv. II. *Flora*, 26, cum ic.—Sumatra.
Horsfieldii, *R. Br. in Trans. Linn. Soc.* xix. (1844) 242.—Java.
Lagascae, Blanco, Fl. Filip. ed. II. 595 = R. Patma.
manillana, *Teschem. in Bost. Journ. Nat. Hist.* iv. (1844) 63, 65, t. 6.—Ins. Philipp.
Patma, *Blume, Korte Beschr. Patma* (1825), 21.—Java.
philippensis, Blanco, Fl. Filip. ed. II. 565 = manillana.
Rochussenii, *Teysm. & Binn. in Nat. Tijdschr. Nederl. Ind.* (1850) 425.—Java.
Schadenbergiana, *Goepp. ex Hieron. in Gartenfl.* xxxiv. (1885) 3. t. 1177.—Ins. Philipp.
Titan, Jack, Desc. Malay. Plants, iii. [1823 ?], 1 = Arnoldi.
Tuan-Mudae, *Becc. in Atti Soc. Ital. Sc. Nat.* xi. (1868) 197.—Borneo.

RAFINESQUIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 429. *COMPOSITAE* [Benth. & Hook. f. ii. 533].
californica, *Nutt. l. c.*—Calif.
neo-mexicana, *A. Gray, Pl. Wright.* ii. 103.—Am. bor. occ.

RAFINESQUIA, Rafin. Fl. Tellur. ii. 96 (1836) = **Hosackia**, Dougl. (Legumin.).
comosa, Rafin. l. c. = H. bicolor.
obovata, Rafin. l. c. (Quid. ?).—Am. bor.

RAFINESQUIA, Rafin. New Fl. Am. iii. 51 (1836) = **Calamintha**, Tourn. (Labiat.).
angustifolia, Rafin. l. c. 52 = C. Nuttallii ?
coccinea, Rafin. Fl. Tellur. iii. 83 = C. coccinea.

RAFINESQUIA, Rafin. Sylva Tellur. 79 (1838) = **Jacaranda**, Juss. (Bignon.).
caerulea, Rafin. l. c. = J. caerulea.

RAFNIA, Thunb. Nov. Gen. 144 (1800); Prod. Pl. Cap. 123 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 475.
HYBOTROPIS, E. Mey. ex Steud. Nom. ed. II. i. 780 (1840).
OEDMANNIA, Thunb. Nov. Gen. 136 (1800).
PELECYNTHIS, E. Mey. Comm. Pl. Afr. Austr. 13 (1835).
VASCOA, DC. Mém. Leg. 186 (1825); Prod. ii. 119 (1825).
affinis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 36.—Afr. austr.
alpina, Eckl. & Zeyh. Enum. 160 = triflora.
amplexicaulis, *Thunb. Nov. Gen.* 144.—Afr. austr.
angulata, *Thunb. l. c.* 147.—Afr. austr.
angustifolia, Thunb. l. c. = angulata.
axillaris, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 468 = affinis.
axillaris, Eckl. & Zeyh. Enum. 161 = crassifolia.
axillaris, *Thunb. Nov. Gen.* 146.—Afr. austr.
axillaris, Willd. ex Walp. in Linnaea, xiii. (1839) 505 = Euchlora serpens.
cordata, Eckl. & Zeyh. Enum. 159 = ovata.
cordata, *Mart. in Denkschr. Akad. Muench.* vi. 1816–1817 (1820) 189.—Afr. austr.
corymbosa, Walp. in Linnaea, xiii. (1839) 464 = cuneifolia.
crassifolia, *Harv. Thes. Cap.* i. 45. t. 71.—Afr. austr.
cuneifolia, E. Mey. Comm. Pl. Afr. Austr. 12 = elliptica.
cuneifolia, *Thunb. Nov. Gen.* 145.—Afr. austr.
dichotoma, *Eckl. & Zeyh. Enum.* 161.—Afr. austr.
diffusa, Eckl. & Zeyh. l. c. 160 = triflora.
diffusa, *Thunb. Nov. Gen.* 149.—Afr. austr.
Ecklonis, *E. Mey. Comm. Pl. Afr. Austr.* 13.—Afr. austr.
elliptica, Sieber, ex Presl, Bot. Bemerk. 44 = myrtifolia.
elliptica, *Thunb. Nov. Gen.* 145.—Afr. austr.
erecta, Eckl. & Zeyh. Enum. 160 = elliptica.
erecta, *Thunb. Nov. Gen.* 149.—Afr. austr.
fastigiata, *Eckl. & Zeyh. Enum.* 160.—Afr. austr.
filifolia, Eckl. ex Walp. Ann. i. 208 = Aspalathus corymbosa.
filifolia, Thunb. Nov. Gen. 148 = angulata.
humilis, *Eckl. & Zeyh. Enum.* 162.—Afr. austr.
intermedia, Vog. ex Walp. in Linnaea, xiii. (1839) 463 = elliptica.
lancea, *DC. Prod.* ii. 119.—Afr. austr.
lancea, E. Mey. Comm. Pl. Afr. Austr. 12 = lancifolia.
lancifolia, *Presl, Bot. Bemerk.* 44.—Afr. austr.
Meyeriana, Presl, l. c. = elliptica.
myrtifolia, *Presl, l. c.*—Afr. austr.
opposita, *Thunb. Nov. Gen.* 146.—Afr. austr.
ovata, *E. Mey. Comm. Pl. Afr. Austr.* 12.—Afr austr.
pauciflora, Eckl. &. Zeyh. Enum. 162 = opposita.
perfoliata, *E. Mey. Comm. Pl. Afr. Austr.* 12.—Afr. austr.
perfoliata, Willd. Sp. Pl. iii. 949 = Baptisia perfoliata.
racemosa, *Eckl. & Zeyh. Enum.* 161.—Afr. austr.
reflexa, Walp. Rep. ii. 833, in syn. = elliptica.
retroflexa, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 469 = diffusa.
retroflexa, Eckl. & Zeyh. Enum. 160 = elliptica.
retroflexa, *Thunb. Nov. Gen.* 148.—Afr. austr.
retusa, Vent. Jard. Malm. 53 = Templetonia retusa.

RAFNIA:—
rhomboidea, Walp. in Linnaea, xiii. (1839) 464 = cuneifolia.
spicata, *Thunb. Nov. Gen.* 147.—Afr. austr.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 36.—Afr. austr.
triflora, *Thunb. Nov. Gen.* 145.—Afr. austr.
virens, *E. Mey. Comm. Pl. Afr. Austr.* 11.—Afr. austr.

RAILLARDELLA, Benth. in Benth. & Hook. f. Gen. ii. 442 (1876). *COMPOSITAE*, Benth. & Hook. f. l. c.
argentea, *A. Gray, in Bot. Calif.* i. 417.—Calif.
Eiseni, Kellogg, ex A. Gray, Syn. Fl. N. Am. i. II. 380 = scaposa.
Muirii, *A. Gray, in Bot. Calif.* i. 618.—Calif.
Pringlei, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 17.—Calif.
scaposa, *A. Gray, in Bot. Calif.* i. 417.—Calif.

RAIANIA, Scop. Introd. 460 (1838) = **Rajania**, Linn. (Diosc.).

RAILLARDA, Endl. Gen. 460 (1838) = seq.

RAILLARDIA, Spreng. Gen. ii. 796 (1831) = seq.

RAILLIARDA, DC. Prod. vi. 440 (1837) = seq.

RAILLIARDIA, Gaudich. in Freyc. Voy. Bot. 469. t. 83 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 442.
arborea, *A. Gray, in Proc. Am. Acad.* v. (1861) 134.—Ins. Sandvic.
argentea, A. Gray, l. c. vi. (1865) 550 = Raillardella argentea.
ciliolata, *DC. Prod.* vi. 441.—Ins. Sandvic.
Hillebrandi, *H. Mann, in Proc. Am. Acad.* vii. (1867) 175.—Ins. Sandvic.
latifolia, *A. Gray, l. c.* v. (1861) 132.—Ins. Sandvic.
laxiflora, *DC. Prod.* vi. 441.—Ins. Sandvic.
linearis, *Gaudich. in Freyc. Voy. Bot.* 469. t. 83.—Ins. Sandvic.
Menziesii, *A. Gray, in Proc. Am. Acad.* v. (1861) 133.—Ins. Sandvic.
montana, *H. Mann, l. c.* vii. (1867) 176.—Ins. Sandvic.
platyphylla, *A. Gray, l. c.* v. (1861) 134.—Ins. Sandvic.
scabra, *DC. Prod.* vi. 441.—Ins. Sandvic.
scaposa, A. Gray, in Proc. Am. Acad. vi. (1865) 551 = Raillardella scaposa.
struthioloides, *A. Gray, l. c.* (1861) 134.—Ins. Sandvic.

RAJANIA, Linn. Gen. ed. I. 306 (1737) = *DIOSCOREACEAE*, Benth. & Hook. f. iii. 744.
JANRAIA, Adans. Fam. ii. 76 (1763).
angustifolia, Griseb. ex Kunth, Enum. Pl. v. 446 = cubensis.
angustifolia, *Sw. Prod. Veg. Ind. Occ.* 59.—Ins. S. Doming.
brasiliensis, *Griseb. in Mart. Fl. Bras.* iii. I. 48.—Bras.
cirrhata, *Billb. ex Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 111.—Costa Rica.
cordata, *Linn. Sp. Pl.* 1032.—Ind. occ.
cubensis, *Kunth, Enum. Pl.* v. 446.—Cuba.
flexuosa, Bello, in Anal. Soc. Esp. Hist. Nat. xii. (1883) 124 = hastata ?
flexuosa, Poir. Encyc. vi. 59 = Dioscorea Bridgesii.
hastata, Griseb. in Mart. Fl. Bras. iii. I. 47, excl. syn. = Dioscorea trachyandra.
hastata, *Linn. Sp. Pl.* 1032.—Ind. occ.
hastata, Poepp. ex Kunth, Enum. Pl. v. 452 = microphylla.
hexaphylla, Thunb. Fl. Jap. 149 = Stauntonia hexaphylla.
lobata, *Poir. Encyc.* vi. 58.—Peruv.
microphylla, *Kunth, Enum. Pl.* v. 451.—Cuba.
mucronata, *Willd. Sp. Pl.* iv. 787.—Ins. S. Doming.
ovata, *Sw. Prod. Veg. Ind. Occ.* 59.—Ind. occ.
pleioneura, *Griseb. Fl. Brit. W. Ind.* 588.—Ind. occ.
quinata, Thunb. Fl. Jap. 148 = Akebia quinata.
quinquefolia, *Linn. Sp. Pl.* 1032.—Ind. occ.
verticillata, Vell. Fl. Flum. x. t. 115 = Herreria Salsaparilha.

RAJANIA, Walt. Fl. Carol. 53, 247 (1788) = **Brunnichia**, Banks (Polygon).
caroliniana, J. F. Gmel. Syst. 581 = seq.
ovata, Walt. Fl. Carol. 247 = B. cirrhosa.

RALEIGHIA, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 97 = **Abatia**, Ruiz & Pav. (Samyd.).
americana, Gardn. l. c. 98 = A. americana.

RAMATUELA, H. B. & K. Nov. Gen. et Sp. vii. 253. t. 656 (1825). *COMBRETACEAE*, Benth. & Hook. f. i. 686.
argentea, *H. B. & K. l. c.* 254.—Am. trop.
virens, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. II. 100.—Bras.

RAMATUELLA, Poir. in Dict. Sc. Nat. xliv. 426 (1826) = praec.

RAMELIA, Baill, Adansonia, xi. (1874) 132. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 326.
codonostylis, *Baill. l. c.*—N. Caled.

RAMEYA, Baill. Adansonia, ix. (1870) 315. *MENISPERMACEAE*.
calopicrosia, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 458.—Madag.
capitata, *Baill. Adansonia*, ix. (1870) 315.—Ins. Comor.
loucoubensis, Baill. l. c. = Triclisia loucoubensis.
macrocarpa, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 458.—Madag.

RAMIREZIA, A. Rich. Fl. Cub. t. 39 (1855) = **Poeppigia**, Presl (Legumin.).
cubensis, A. Rich. l. c. = P. excelsa.

RAMISCHIA, Opiz, Seznam, 82 (1852) = **Pyrola**, Tourn. (Ericac.).
secunda, Garcke, Fl. Deutschl. ed. IV. 222 = P. secunda.
secundiflora, Opiz, Seznam, 82 = P. secunda.

RAMONDIA, Rich. in Pers. Syn. i. 216 (Ramonda) (1805); Jaume St. Hil. Expos. i. 280 (1805). *GESNERACEAE*, Benth. & Hook. f. ii. 1024.
CHAIXIA, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 38 (1818).
JANCAEA, Boiss. Pl. Or. Nov. Dec. i. 4 (1875).
JANKAEA, Boiss. Fl. Orient. iv. 82 (1879).
LOBIROTA, Dulac, Fl. Hautes-Pyr. 430 (1867).
MYCONIA, Lapeyr. Hist. Abr. Pl. Pyr. 115 (1813).
Heldreichii, *Janka, ex Boiss. Pl. Or. Nov. Dec.* i. 5.—Thessalia.
Myconi, Reichb. Fl. Germ. Excurs. 388 = pyrenaica.
Nathaliae, Panč. & Petrov. Fl. Agr. Nyssani, 574; ex Just, Bot. Jahresb. xi. 850 = serbica.
pyrenaica, *Rich. in Pers. Syn.* i. 216.—Mont. Pyren.
scapiflora, Jaume St. Hil. Expos. i. 280 = pyrenaica.
scapigera, DC. Fl. Fr. Suppl. 417, sphalm. = praec.
serbica, *Panč. ex C. B. Clarke, in DC. Monog. Phan.* v. 168.—Servia.

RAMOTHA, Rafin. Fl. Tellur. ii. 15 (1836) = **Xyris**, (Gronov.).
africana, Rafin. l. c. (Quid?).—Afr.
floridana, Rafin. l. c. = X. flexuosa?
operculata, Rafin. l. c. = X. operculata.
pubescens, Rafin. l. c. = X. pubescens.
vera, Rafin. l. c. = X. indica?

RAMPHICARPA, Reichb. Nom. 119 (1841) = **Rhamphicarpa**, Benth. (Scrophul.).

RAMPHIDIA, Miq. Fl. Ind. Bat. iii. 730 (1858) = RHAMPHIDIA, Lindl. = **Hetaeria**, Blume (Orchid.).

RAMPHOCARPUS, Neck. Elem. ii. 438 (1790) = **Erodium**?, L'Hérit. (Geran.).

RAMPHOSPERMUM, Andrz. ex Reichb. in Moessl. Handb. ed. II. i. p. lvii (1827) = RHAMPHOSPERMUM, Reichb. = **Brassica**, Tourn. (Crucif.).

RAMPINIA, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 129 = **Herpetospermum**, Wall. (Cucurb.).
herpetospermoides, C. B. Clarke, l. c. 130 = H. caudigerum.

RAMSPEKIA, Scop. Introd. 145 (1777) = **Posoqueria**, Aubl. (Rubiac.).

RAMTILLA, DC. in Wight, Contrib. 18 (1834) = **Guizotia**, Cass. (Compos.).
oleifera, DC. l. c. = G. abyssinica.

RAMUSIA, E. Mey. in Drège, Zwei Pfl. Docum. 215 (1843) = **Asystasia**, Blume (Acanth.).
nyctaginea, E. Mey. l. c.; et ex Nees, in DC. Prod. xi. 167 = A. coromandeliana.

RAMUSIA, Nees, in DC. Prod. xi. 309 (1847) = **Peristrophe**, Nees (Acanth.).
tridentata, Nees, l. c.—Afr. austr.

RANAPALUS, Kellogg, in Proc. Calif. Acad. vii. (1877) 113 = **Herpestis**, Gaertn. f. (Scrophul.).
Eisenii, Kellogg, l. c. = H. rotundifolia.

RANARIA, Cham. in Linnaea, viii. (1825) 30 = **Herpestis**, Gaertn. f. (Scrophul.).
monnierioides, Cham. l. c. 31 = H. Ranaria.

RANCAGUA, Poepp. & Endl. Nov. Gen. et Sp. i. 15. tt. 24, 25 (1835) = **Lasthenia**, Cass. (Compos.).
Bridgesii, Poepp. & Endl. l. c. t. 25 = L. obtusifolia.
Fevillei, Poepp. & Endl. l. c. t. 24 = L. obtusifolia.
glaberrima, Endl. ex Walp. Rep. vi. 180 = L. glabrata.

RANDALIA, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 47. t. 5 = **Eriocaulon**, Linn.
decangularis, Beauv. l. c. = E. decangulare.
densa, Beauv. l. c. = E. deustum.
depressa, Beauv. l. c. = E. depressum.
fistulosa, Beauv. l. c. = E. fistulosum.
scariosa, Beauv. l. c. = E. scariosum.

RANDIA, Houst. ex Linn. Gen. ed. I. 376 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 88.
CANTHIOPSIS, Seem. Fl. Vit. 166. t. 46 (1866).
CATU-NAREGAM, Adans. Fam. ii. 85 (1763).
CERISCUS, Gaertn. f. Fruct. iii. 140. t. 28 (1797).
CUPI, Adans. Fam. ii. 158 (1763).
CUPIA, DC. Prod. iv. 393 (1830).
FOSCARENIA, Vell. ex Vand. Fl. Lusit. 7 (1788); et in Roem. Script. 78 (1796).
GRIFFITHIA, Wight & Arn. Prod. 399 (1834).
GYNOPACHIS, Blume, Cat. Gew. Buitenz. 48 (1823).
LACHNOSIPHONIUM, Hochst. in Flora, xxv. (1842) 236.
NAREGA, Rafin. Sylva Tellur. 98 (1838).
OXYCEROS, Lour. Fl. Cochinch. 150 (1790).
PSEUDIXORA, Miq. Fl. Ind. Bat. ii. 209 (1856).
STYLOCORYNA, Cav. Ic. iv. 45. t. 368 (1797).
TALANGNINIA, Chapel. ex DC. Prod. iv. 389 (1830).
THIOLLIERA, Montr. in Mém. Acad. Lyon, x. (1860) 217.
XEROMPHIS, Rafin. Sylva Tellur. 21 (1838).
accedens, *Hance, in Ann. Sc. Nat.* Ser. V. v. (1866) 218.—China.
aculeata, *Linn. Sp. Pl.* 1192.—Ind. occ.
acuminata, *Benth. in Hook. Niger Fl.* 385.—Afr. trop.
africana, *G. Don, Gen. Syst.* iii. 502 (Quid?).—Afr. trop.
angatensis, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 108.—Ins. Philipp.
anisophylla, *Hook. f. Fl. Brit. Ind.* iii. 114.—Malaya.
armata, DC. Prod. iv. 387 = Basanacantha armata.
auriculata, Steud. Nom. ed. II. ii. 430 = Anomanthodia auriculata.
Beccariana, *Baill. Adansonia*, xii. (1879) 246.—Borneo.
Benthamiana, *F. Muell. Fragm.* ix. 180.—Austral.
Bojeriana, *Baill. Adansonia*, xii. (1879) 245.—Madag.
Bowieana, A. Cunn. ex Hook. Bot. Mag. t. 3409 = macrantha.
Brandzeana, *Baill. Adansonia*, xii. (1879) 244.—N. Caled.
Buchananii, *Oliver, in Hook. Ic. Pl.* t. 1356.—Afr. trop.
calophylla, Griseb. Cat. Pl. Cub. 123 = Casasia calophylla.
calycina, Cham. in Linnaea, ix. (1834) 246 (= Basanacanthae sp.).—Bras.
Candolleana, *Wight & Arn. Prod.* 399.—Ind. or.
canthioides, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 194.—China.
capitata, *DC. Prod.* iv. 387.—Mexic.
caudata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 96.—Afr. trop.
chartacea, *F. Muell. Fragm.* ix. 180, in obs.—Austral.
chinensis, DC. Prod. iv. 388, in syn. = sinensis.
ciliolata, *Wright, in Sauv. Fl. Cub.* 60.—Cuba.
clusiaefolia, Chapm. Fl. S. U. St. 179 = Genipa clusiifolia.
coriacea, Benth. in Hook. Niger Fl. 387 = Tricalysia coriacea.
corymbosa, Wight & Arn. Prod. 399 = Candolleana.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 179.—Ins. Philipp.
Cupia, Steud. Nom. ed. II. ii. 430 = longiflora.
deccanensis, Bedd. Ic. Pl. Ind. Or. i. 57 = Candolleana.
decussata, Wall. ex Voigt, Hort. Suburb. Calc. 381 = Hedyotis macrophylla.
densiflora, *Benth. Fl. Hongk.* 155.—As. et Austral. trop.
Devoniana, *Benth. & Hook. f. Gen.* ii. 89.—Afr. trop.
dilleniacea, *Baill. Adansonia*, xii. (1879) 245.—Borneo.
dioica, Karst. in Linnaea, xxx. (1859–60) 153 = Basanacantha dioica.
Doniana, Benth. in Hook. Niger Fl. 385 = Sherbournia foliosa.
drupacea, *DC. Prod.* iv. 389.—Java.
dumetorum, *Lam. Illustr.* ii. 227.—Geront. trop.
echinocarpa, [*Moç. & Sessé,*] *ex DC. Prod.* iv. 385.—Mexic.
edulis, *Kostel. Allg. Med.-Pharm. Fl.* ii. 579.—Cochinch.
exaltata, *Griff. Notul.* iv. 262.—Burma.
fasciculata, *DC. Prod.* iv. 386.—Reg. Himal.; Burma.
ferox, *DC. l. c.* 387.—Mexic.
Fitzalani, *F. Muell. ex Benth. Fl. Austral.* iii. 411.—Austral.
floribunda, DC. Prod. iv. 386 = dumetorum.
foliolosa, Wight & Arn. Prod. 398 = aculeata.
gambica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 96.—Afr. trop.
Gardneri, *Hook. f. Fl. Brit. Ind.* iii. 112.—Zeylan.
genipaeflora, *DC. Prod.* iv. 389.—Afr. trop.
glomerata, *Benth. Bot. Voy. Sulph.* 103.—Ecuador.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 112.—Reg. Himal.
hebecarpa, Benth. in Hook. Journ. Bot. iii. (1841) 213 = Basanacantha hebecarpa.
heterophylla, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 14; *et in Baker, Fl. Maurit.* 141.—Ins. Rodrig.
hirta, F. Muell. Census, 74 = Gardenia hirta.
hondensis, *Karst. in Linnaea*, xxx. (1859–60) 154.—N. Granat.
horrida, *Roem. & Schult. Syst.* v. 248.—Cochinch.
Humboldtiana, DC. Prod. iv. 387 = Basanacantha Humboldtiana.
hygrophyla, *Hook. f. Fl. Brit. Ind.* iii. 115.—Burma.
Karstenii, *Polak. in Linnaea*, xli. (1877) 568.—Costa Rica.
Kraussii, Harv. Thes. Cap. i. 22 = dumetorum.
Lachnosiphonium, *Hochst. in Flora*, xxv. (1842) 237.—Afr. austr.
latifolia, Lam. Encyc. iii. 24 = aculeata.
laurifolia, Benth. & Hook. f. Gen. ii. 88 = Gardneri.
leucocarpa, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 194.—Ins. Hongkong.
longiflora, *Lam. Encyc.* iii. 26.—Reg. Himal.; Malaya.
longiflora, Salisb. Parad. Lond. t. 93 = macrantha.
longispina, DC. Prod. iv. 386 = dumetorum.
longistyla, DC. l. c. 388 = Macrosphyra longistyla.
Macarthuri, *F. Muell. Papuan Pl.* i. 68.—N. Guin.
macracantha, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 234.—Java.
macrantha, *DC. Prod.* iv. 388.—Afr. trop.
macrocarpa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 97.—Afr. trop.
macrophylla, *Benth. & Hook. f. Gen.* ii. 89; *et ex Hook. f. Fl. Brit. Ind.* iii. 114.—Malaya.
maculata, *DC. Prod.* iv. 388.—Afr. trop.
maculata, *Span. in Linnaea*, xv. (1841) 315.—Ins. Timor.
madagascariensis, *DC. Prod.* iv. 389.—Madag.
malabarica, *Lam. Encyc.* iii. 25.—Ind. or.

RANDIA:—
malabarica, Wall. Cat. n. 8255 = fasciculata, dumetorum.
malleifera, *Benth. & Hook. f. Gen.* ii. 89.—Afr. trop.
malleiflora, Walp. Ann. ii. 795, sphalm. = praec.
mitis, Linn. Sp. Pl. 1192 = aculeata.
monantha, Benth. Pl. Hartw. 84 = Basanacantha monantha.
Moorei, *F. Muell. ex Benth. Fl. Austral.* iii. 411.—Austral.
Munsae, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 99.—Afr. trop.
Mussaenda, *DC. Prod.* iv. 388.—Am. austr.
nigrescens, *Griseb. Cat. Pl. Cub.* 123.—Cuba.
nitida, *DC. Prod.* iv. 387.—N. Granat.
nutans, DC. l. c. 386 = dumetorum.
obovata, *H. B. & K. Nov. Gen. et Sp.* iii. 409.—N. Granat.
obovata, Ruiz & Pav. Fl. Per. ii. 68 = pubescens.
obscura, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 108 = Gardenia obscura.
octomera, *Benth. & Hook. f. Gen.* ii. 89.—Afr. trop.
ovata, *Duchass. ex Griseb. in Bonplandia*, vi. (1858) 8.—Panama.
oxypetala, *Lindl. Bot. Reg.* (1843) *Misc.* 63.—Reg. Himal.
pallens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 96.—Afr. trop.
parviflora, *Lam. Encyc.* iii. 25.—Ind. or.
parvifolia, *Harv. Thes. Cap.* i. 23.—Afr. austr.
parvifolia, Lam. Encyc. iii. 25 = Catesbaea parvifolia.
patula, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 235.—Java.
polysperma, Roxb. Hort. Beng. 15; Fl. Ind. i. 527 = Petunga Roxburghii.
propinqua, G. Don, Gen. Syst. iii. 502 = Gardenia propinqua.
pubescens, *Ruiz & Pav. Fl. Per.* ii. t. 220. f. b.—Peruv.
racemosa, *Maxim. in Bull. Acad. Pétersb.* xxix. (1884) 167.—Ins. Philipp.; Archip. Luchu.
racemosa, Roxb. Hort. Beng. 15; Fl. Ind. i. 525 = Petunga Roxburghii.
reticulata, Benth. in Hook. Niger Fl. 386 = Tricalysia reticulata.
rigida, DC. Prod. iv. 386 = fasciculata.
Rottleri, Wight & Arn. Prod. 397 = dumetorum.
rotundifolia, *Ruiz & Pav. Fl. Per.* ii. 68.—Peruv.
rubens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 95.—Afr. trop.
rudis, *E. Mey. ex Harv. Thes. Cap.* i. 22.—Afr. austr.
rugulosa, *Hook. f. Fl. Brit. Ind.* iii. 113.—Ind. or.
Ruiziana, *DC. Prod.* iv. 388.—Peruv.
Sagraeana, Griseb. Cat. Pl. Cub. 122 = Scolosanthus crucifer.
scandens, DC. Prod. iv. 387 = longiflora.
scandens, Lam. Encyc. iii. 26 = Gardenia scandens.
sessilis, *F. Muell. Fragm.* vii. 47.—Austral.
Sherborniae, Hook. Niger Fl. 385 = Sherbournia foliosa.
sikkimensis, *Hook. f. Fl. Brit. Ind.* iii. 114.—Reg. Himal.
sinensis, *Roem. & Schult. Syst.* v. 248.—China.
speciosa, *Bedd. Forest Man. Bot.* 133.—Ind. or.
speciosa, *DC. Prod.* iv. 388.—Guian.
speciosa, Hook. Ic. Pl. t. 824 = Gardenia speciosa.
spinosa, Karst. Fl. Columb. ii. 128, adnot. = Basanacantha armata.
spinosa, Poir. Encyc. Suppl. ii. 829; Blume, Bijdr. 981 = dumetorum.
Stanleyana, *Walp. Ann.* ii. 794.—Afr. trop.
stipularis, F. Muell. Papuan Pl. i. 69 = stipulosa.
stipulosa, Miq. Fl. Ind. Bat. ii. 228 = dumetorum.
stipulosa, *Moore & F. Muell. ex F. Muell. Fragm.* vii. 47.—Austral.
stricta, Roxb. Hort. Beng. 15; Fl. Ind. i. 526 = Hyptianthera stricta.
sumatrana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 235.—Sumatra.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 54.—Ins. Tahiti.
Talangninia, *DC. Prod.* iv. 389.—Madag.
tetracantha, DC. l. c. 387 = Basanacantha tetracantha.
tetrandra, *DC. l. c.*—N. Granat.
tetrasperma, *Benth. & Hook. f. ex Brand. For. Fl.* 272.—Reg. Himal.
tomentosa, *Hook. f. Fl. Brit. Ind.* iii. 110.—Burma; Malaya.

RANDIA:—
tomentosa, Wight & Arn. Prod. 398 = dumetorum.
triflora, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 138.—Nepal.
uliginosa, *Poir. Encyc. Suppl.* ii. 829; *DC. Prod.* iv. 386.—Ind. or.; Burma.
urcelliformis, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 104 = Gardenia urcelliformis.
venulosa, Wall. Cat. n. 8301 = Petunga venulosa.
Vieillardi, *Baill. Adansonia*, xii. (1879) 295.—N. Caled.
virosa, *Buch.-Ham. in Trans. Linn. Soc.* xvii. (1835) 165.—Ind. or.
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 113.—Reg. Himal.; Burma.
Wrightiana, Griseb. Cat. Pl. Cub. 122 = Scolosanthus Wrightii.
xalapensis, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 239.—Mexic.
Zollingeri, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1866) 248.—Ins. Sunda.
Zollingeriana, Teijsm. & Binn. Cat. Hort. Bog. 384 = praec.

RANDONIA, Coss. in Bull. Soc. Bot. Fr. vi. (1859) 391. *RESEDACEAE*, Benth. & Hook. f. i. 111.
africana, *Coss. l. c.* 392.—Alger.

RANGIA, Griseb. Cat. Pl. Cub. 122 (1866), err. typ. = **Randia**, Houst. (Rubiac.).

RANGIUM, Juss. in Dict. Sc. Nat. xxiv. 200 (1822) = **Forsythia**, Vahl (Oleac.).

RANISIA, Salisb. Gen. Pl. Fragm. 143 (1866) = **Gladiolus**, Linn. (Irid.).
concolor, Salisb. l. c. = G. tristis.
gracilis, Salisb. l. c. = G. gracilis.
hirsuta, Salisb. l. c. = G. hirsutus.
recurva, Salisb. l. c. = G. recurvus.
trichonemifolia, Salisb. l. c. = G. trichonemifolius.
tristis, Salisb. l. c. = G. tristis.
versicolor, Salisb. l. c. = G. grandis.

RANULA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 324 = **Ranunculus**, Tourn.
aconitifolia, Fourr. l. c. = Ranunc. aconitifolius.
platanifolia, Fourr. l. c. = Ranunc. aconitifolius.

RANUNCULASTRUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 324 = **Ranunculus**, Tourn.
acre, Fourr. l. c. = Ranunculus acris.
albonaevum, Fourr. l. c. xvii. (1869) 194 = Ranunculus bulbosus.
auricomum, Fourr. l. c. xvi. (1868) 324 = Ranunculus auricomus.
Boraeanum, Fourr. l. c. = Ranunculus acris.
bulbiferum, Fourr. l. c. xvii. (1869) 194 = Ranunculus bulbosus.
bulbosum, Fourr. l. c. xvi. (1868) 324 = Ranunculus bulbosus.
gracile, Fourr. l. c. = Ranunculus montanus.
lanuginosum, Fourr. l. c. = Ranunculus lanuginosus.
mixtum, Fourr. l. c. = Ranunculus nemorosus.
nemorivagum, Fourr. l. c. = Ranunculus acris.
nemorosum, Fourr. l. c. = Ranunculus nemorosus.
radicescens, Fourr. l. c. = Ranunculus nemorosus.
repens, Fourr. l. c. = Ranunculus repens.
reptabundum, Fourr. l. c. = Ranunculus repens.
sparsipilum, Fourr. l. c. = Ranunculus bulbosus.
valdepubens, Fourr. l. c. xvii. (1869) 194 = Ranunculus bulbosus.
vulgatum, Fourr. l. c. xvi. (1868) 324 = Ranunculus acris.

RANUNCULUS, [Tourn.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
APHANOSTEMMA, A. St. Hil. Fl. Bras. Mer. i. 12 (1825).
BATRACHIUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 720 (1821).
CASALEA, A. St. Hil. Fl. Bras. Mer. i. 8. t. 1 (1825).
CERATOCEPHALA, Moench, Meth. 218 (1794).
CYNOMORBIUM, Opiz, Fl. Časlav. Cent. iii. 261, 255 (1804).—Cf. Lotos, iv. (1854) 43.
CYPRIANTHE, Spach, Hist. Vég. Phan. vii. 220 (1839).

RANUNCULUS:—
CYRTORHYNCHA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 26 (1838).
FICARIA, [Dill.] Hall. Enum. Stirp. Helv. i. 321 (1742).
FLAMMULA, Dod. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 324.
GAMPSOCERAS, Stev. in Bull. Soc. Nat. Mosc. xxv. (1852) 1. 541, in textu.
GLOSSOPHYLLUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325.
HECATONIA, Lour. Fl. Cochinch. 302 (1790); DC. Syst. i. 227 (1818).
HERICINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325.
IONOSMANTHUS, Jord. & Fourr. Ic. Pl. Europ. ii. 13 (1870).
KRAPFIA, DC. Syst. i. 228 (1818).
LEUCONOË, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323.
NOTOPHILUS, Fourr. l. c. 325 (1868).
PACHYLOMA, Spach, Hist. Vég. Phan. vii. 194 (1839).
PFUNDIA, Opiz, Seznam, 73 (1852).
RANULA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 324.
RANUNCULASTRUM, Fourr. l. c. 324 (1868).
SCOTANUM, Adans. Fam. ii. 459 (1763).
STYLURUS, Rafin. Fl. Ludov. 27 (1817).
THORA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325.
XERODERA, Fourr. l. c. 324 (1868).
XIPHOCOMA, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 270, in textu; et xxv. (1852) I. 537.
abnormis, *Cutanda & Willk. in Linnaea*, xxx. (1859–60) 83.—Hispan.
abortivus, *Linn. Sp. Pl.* 551.—Am. bor.
abrotanifolius, Auct. ex Steud. Nom. ed. I. 676; ed. II. ii. 431 = divaricatus.
acaulis, *Banks & Soland. in DC. Syst.* i. 270.—N. Zel.
accessivus, Jord. Diagn. i. 70 = Villarsii.
acetosellaefolius, *Boiss. Elench.* 5.—Hispan.
aconitifolius, *Linn. Sp. Pl.* 551.—Europ.
acris, *Linn. l. c.* 554.—Europ.; As. bor.
acris, Willk. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 935 = nemorosus.
acutidentatus, Rupr. Fl. Cauc. 286 = seq.
acutilobus, *Ledeb. Fl. Ross.* i. 40; *Rupr. l. c.* 22.—Reg. Caucas.
adoneus, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 56.—Am. bor.
adscendens, *Brot. Fl. Lusit.* ii. 370.—Hispan.; Lusit.
adscendens, Brot. Phyt. Lusit. ii. 229 = Broteri.
aduncus, Gren. & Godr. Fl. Fr. i. 32 = Villarsii.
affinis, *R. Br. in Parry*, 1*st. Voy. App.* 265.—Reg. arct.
affinis, F. Schultz, in Flora, xxiv. (1841) 558 = trichophyllus.
afghanicus, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 148.—Afghan.
Agerii, Bertol. in Opusc. Sc. Bologn. iii. (1819) 182. t. 6 = chaerophyllos.
agrarius, All. Auct. Pedem. 27; Poir. Encyc. vi. 117 = sardous.
alatus, Poir. l. c. 127 = Soliva sessilis.
Alberti, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 223.—Turkest.
albicans, Jord. Obs. Pl. Crit. vi. (1847) 10 = monspeliacus.
albonaevus, Jord. Diagn. i. 81 = bulbosus.
Aleae, *Willk. in Linnaea*, xxx. (1859–60) 84.—Hispan.
aleuticus, *Mert. in Linnaea*, v. (1830) 65, nomen.—Ins. Aleut.
algidus ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 78.—Europ.
alismaefolius, *Geyer, ex Benth. Pl. Hartw.* 295.—Am. bor. occ.
alismoides, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 13 = angustifolius.
alpestris, *Linn. Sp. Pl.* 553.—Europ.
alpestris, Pall. Reise, ii. 568 = nivalis.
alpicola, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 177 = nemorosus.
alpicola, Timb. in Bull. Soc. Bot. Fr. xv. (1868), Sess. Extr. p. xc = Villarsii.
altaicus, Laxm. in Nov. Comm. Acad. Petrop. xviii. (1774) 533 = nivalis.

RANUNCULUS:—
Amansii, Jord. Diagn. i. 79 = nemorosus.
Amarillo, *Bertol. Fl. Guatimal.* 24.—Guatemala.
ambigens, S. Wats. Bibliog. Ind. N. Am. Bot. i. 16 = obtusiusculus.
ambiguus, Jord. Diagn. i. 76 = polyanthemos.
ambiguus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 252 = cassubicus.
amblyolobus, *Boiss. & Hohen. Diagn.* Ser. I. viii. 5.—Persia.
Ammanni, Gunn. Fl. Norveg. ii. 103 = hyperboreus.
ammoides, Pall. ex DC. Prod. i. 42 = trilobus.
amoenus, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 241 = adoneus ?
amoenus, Ledeb. Fl. Alt. ii. 320 = affinis.
amphibius, James, in Long, Exped. Rocky Mount. ii. 342 = multifidus.
amphitricha, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 237.—N. Zel.
amplexicaulis, *Linn. Sp. Pl.* 549.—Europ. austr.
amurensis, *Maxim. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 41, nomen.—Reg. Amur.
Andersonii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 327.—Am. bor. occ.
andinus, *Phil. in Linnaea*, xxxiii. (1864–65) 2.—Chili.
anemonaefolius, *DC. ex Poir. Encyc. Suppl.* iv. 663.—Turcia.
anemonefolius, *DC. Syst.* i. 282.—As. Min.
anemoneus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 97.—Austral.
anemonoides, Sievers, ex Steud. Nom. ed. I. 677, 679 = Callianthemum rutaefolium.
anemonoides, *Zahlbr. in Flora*, vi. (1823) 220.—Styria.
angulatus, Presl, Delic. Prag. 7 = sardous.
angustifolius, *DC. Rapp. Voy.* i. 78; *ex ej. Fl. Fr. Suppl.* 636.—Mont. Pyren.
apiifolius, Desf. in Pict. Hort. Par. 37. t. 73; ex DC. Syst. i. 260 = monspeliacus.
apiifolius, *Pers. Syn.* ii. 105.—Reg. Argent.
apiifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = sceleratus.
aquaticus, Neck. Delic. Gallo-Belg. i. 242; Lam. Fl. Fr. iii. 184 = aquatilis.
aquatilis, *Linn. Sp. Pl.* 556, partim.—Reg. temp.
aquatilis, Linn. l. c., partim = peltatus.
arachnoideus, *C. A. Mey. Verz. Pfl. Cauc.* 201.—Reg. Caucas.
arcticus, Richards. in Frankl. 1st Journ. ed. I. App. 741 = affinis.
argemonifolius, *Griseb. in Goett. Abh.* xix. (1874) 70.—Reg. Argent.
argyreus, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 352.—As. Min.
armeniacus, Boiss. & Huet, Diagn. Ser. II. v. 9 = anemonefolius.
arvensis, *Linn. Sp. Pl.* 555.—Europ.; As. bor.
ascendens, Steud. Nom. ed. II. ii. 432 = adscendens.
Aschenbornianus, *Schau. in Linnaea*, xx. (1847) 719.—Mexic.
Aschersonii, *Freyn, in Bot. Centralb.* vi. (1881) *Beil.* [iii.] 13.—Afr. bor.
asiaticus, Bory & Chaub. Nouv. Fl. Pélop. 35 = chaerophyllos.
asiaticus, Gueldenst. Reisen, i. 193 = ophioglossifolius.
asiaticus, Lepech. Reisen, iii. 40 = Trollius asiaticus.
asiaticus, *Linn. Sp. Pl.* 552.—Oriens.
asiaticus, Thunb. Fl. Jap. 241 = acris.
astrantiaefolius, Boiss. ex Rupr. Fl. Cauc. 25 = caucasicus.
astrantiaefolius, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 29 = nemorosus.
atlanticus, Ball, in Journ. Bot. xi. (1873) 296 = acris.
atlanticus, Pomel, Nouv. Mat. Fl. Atl. 248 = aquatilis.
attenuatus, Royle, Illustr. Bot. Himal. 53 = hirtellus.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 351.—Persia.
Aucheri, Clem. ex Boiss. Fl. Orient. i. 43 = Sibthorpii.
aucklandicus, *A. Gray, Bot. U. St. Expl. Exped.* i. 8.—Ins. Auckland.
aurasiacus, *Pomel, Nouv. Mat. Fl. Atl.* 379.—Alger.
aureus, Schleich. Cat. Pl. Helv. ed. III. 24 = nemorosus.
auricomus, *Linn. Sp. Pl.* 551.—Reg. bor. temp.

RANUNCULUS:—
auricomus, Schlecht. in Linnaea, vi. (1831) 577 = cassubicus.
Bachii, *Wirtg. in Verhandl. Preuss. Rheinl.* ii. (1846) 22.—Europ.
Baerianus, Bogusl. in Erman, Archiv, vi. (1848) 60 = pygmaeus.
Baidarae, *Rupr. Fl. Cauc.* 22.—Reg. Caucas.
Balbisii, Moris, Fl. Sard. i. 40 = bulbosus.
balearicus, Freyn, in Oestr. Bot. Zeitschr. xxvi. (1876) 158 = palustris.
Barthii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 41 = acris.
batrachioides, *Pomel, Nouv. Mat. Fl. Atl.* 249.—Alger.
Baudotii, *Godr. in Mém. Soc. Roy. Nancy*, 1839 (1840), 21.—Europ. occ.; Marocc.
Bauhini, Tausch, in Flora, xvii. (1834) 525 = aquatilis.
Baurii, *Mac Owan, in Journ. Linn. Soc.* xviii. (1881) 390.—Afr. austr.
Beckii, G. Don, Gen. Syst. i. 39 = multifidus.
belgicus, *Dum. Comm. Bot.* 52, in nota.—Belgium.
Belvisii, DC. Syst. i. 291 = repens.
Berardi, Vill. ex Steud. Nom. ed. II. ii. 432 = rutaefolius.
Berteroanus, *Phil. in Linnaea*, xxviii. (1856) 661.—Chili.
Berteroanus, Presl, Bot. Bemerk. 7 = saxifragaefolius.
Berteronianus, Steud. Nom. ed. II. ii. 432 = trisepalus.
Bertolonii, Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 378, in textu = bilobus.
bilobus, *Bertol. Misc. Bot.* xix. 5.—Europ. austr.
binatus, Kit. ex Reichb. Fl. Germ. Excurs. 723 = auricomus.
bipartitus, *Gromow, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 41, nomen.—Rossia.
biternatus, *Sm. in Rees, Cyclop.* xxix. n. 48.—Reg. Magell.
blepharicarpos, Boiss. Elench. 5 = spicatus.
blepharicarpos, Hack. & Winkl. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 921 = suborbiculatus.
blepharicarpos, Lange, in Kjoeb. Vidensk. Meddel. (1865) 57 = escurialensis.
Bloomeri, *S. Wats. in Bot. Calif.* ii. 426.—Calif.
Blumei, Steud. Nom. ed. II. ii. 432 = diffusus.
bonariensis, *Poir. Encyc.* vi. 102.—Reg. Argent.
Bonplandianus, *H. B. & K. Nov. Gen. et Sp.* v. 46.—N. Granat.
Boraeanus, Jord. Obs. Pl. Crit. vi. (1847) 19 = acris.
borealis, Trautv. in Bull. Soc. Nat. Mosc. xxxiii. (1860) i. 72 = acris.
Bourgaei, *Boiss. Fl. Orient.* i. 43.—Armenia.
brachiatus, Schleich. Cat. Pl. Helv. ed. III. 24 = bulbosus.
brachylobus, *Boiss. & Hohen. Diagn.* Ser. I. viii. 6.—As. Min.; Persia.
brachyphyllus, Tenore, Fl. Nap. t. 149 = brevifolius.
bracteatus, Schleich. ex DC. Prod. i. 41 = bulbosus.
brevicaulis, Hook. Fl. Bor. Am. i. 13 = ovalis.
brevifolius, *Tenore, Prod. Fl. Nap.* 68.—Ital.; Graec.
brevipes, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 14.—N. Granat.
brevirostris, Edgew. in Trans. Linn. Soc. xx. (1846) 28 = laetus.
breviscapus, *DC. Syst.* i. 253.—Peruv.
Breynii, J. F. Gmel. Syst. 880 = seq.
Breyninus, Crantz, Stirp. Austr. ed. I. fasc. ii. 91; ed. II. 115 = nemorosus.
Broteri, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 930.—Hispan.
brutius, *Tenore, Prod. Fl. Nap.* 61.—Ital.; Graec.; As. Min.
Buchanani, *Hook. f. Handb. N. Zeal. Fl.* 5.—N. Zel.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 5.—Japon.
Buhsei, *Boiss. Fl. Orient.* i. 45.—Persia.
bulbifer, Jord. Diagn. i. 80 = bulbosus.
bulbilliferus, *Boiss. & Hohen. Diagn.* Ser. I. viii. 3.—Persia.
bulbosus, Costa, Fl. Catal. 7 = Aleae.
bulbosus, *Linn. Sp. Pl.* 554.—Europ.; Persia.
bulbosus, Plan. Fl. Gallega, 58 = occidentalis, *Freyn.*
bulbosus, Willk. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 932 = castellanus.
bullatus, *Linn. Sp. Pl.* 550.—Reg. Mediterr.
Bungei, Steud. Nom. ed. II. ii. 432 = aquatilis.

RANUNCULUS:—
bupleurifolius, DC. Syst. i. 243, in syn. = pyrenaeus.
bupleuroides, *Brot. Fl. Lusit.* ii. 365.—Lusitan.
bupleuroides, Lapeyr. ex Nym. Consp. 10 = pyrenaeus.
cabulicus, Boiss. Diagn. Ser. II. i. 9 = muricatus.
cadmicus, *Boiss. l. c.* Ser. I. i. 65.—Caria; Cyprus.
caespititius, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 359 = R. Flammula.
caespitosus, Thuill. Fl. Par. ed. II. i. 279 = divaricatus.
caespitosus, Wall. Cat. n. 4701 = affinis.
Calcarae, *Tineo, ex Guss. Fl. Sic. Syn.* ii. 889.—Sicilia.
Caleyanus, *G. Don, Gen. Syst.* i. 37 (Quid ?).—Anglia.
californicus, *Benth. Pl. Hartw.* 295.—Am. bor. occ.
calthaefolius, Jord. Obs. Pl. Crit. vi. (1847) 2 = R. Ficaria.
Calverti, Boiss. & Huet, Diagn. Ser. II. v. 8 = anemonefolius.
Camissonis, Auct. vide *Chamissonis*.
cammarifolius, Arv. Ess. Pl. Dauph. 21 = acris.
Camozzianus, Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 242 = Sibthorpii.
camtschaticus, Steud. Nom. ed. II. ii. 432 = *kamchaticus*.
canadensis, Jacq. Misc. ii. 343 = pensylvanicus.
cantoniensis, DC. Prod. i. 43 = pensylvanicus.
canus, Benth. Pl. Hartw. 294 = occidentalis, *Nutt.*
canus, Torr. in Pacif. Rail. Rep. iv. 62 = californicus.
Canuti, *Coss. ex Ardoino, Fl. Alp. Marit.* 16.—Liguria.
capensis, *Thunb. Prod. Pl. Cap.* 94.—Afr. austr.
capillaceus, Thuill. Fl. Par. ed. II. i. 278 = aquatilis.
capillaris, Gater. Pl. Montaub. 102 = aquatilis.
Capillus-naidis, Arv. Ess. Pl. Dauph. 18 = lutulentus.
cappadocicus, *Willd. Sp. Pl.* ii. 1326.—Cappadoc.
cardiophyllus, Hook. Fl. Bor. Am. i. 14 = affinis.
carinatus, Freyn, in Kern. Sched. Fl. Exsicc. i. 21 = confusus.
carinthiacus, Hoppe, in Sturm, Deutschl. Fl. Heft 46 = montanus.
carnosus, Wall. Cat. n. 4699 = sceleratus.
carolinianus, DC. Syst. i. 292 = repens.
carolinianus, Torr. & Gray, Fl. N. Am. i. 23 = palmatus.
carpaticus, Herbich, Sel. Pl. Rar. Gallic. 15 = montanus.
carpaticus, Wahlenb. ex Nym. Consp. 9 = R. Thora.
carpetanus, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 3.—Hispan.
carpetanus, Bourg. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 921 = suborbiculatus.
cassius, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 354.—As. Min.; Syria.
cassubicus, *Linn. Sp. Pl.* 551.—Europ.; Sibir.
castellanus, *Boiss. & Reut. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 932.—Hispan.
caucasicus, *Bieb. Fl. Taur. Cauc.* ii. 27.—Reg. Caucas.
caucasicus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 5 = Buhsei.
cervicornis, Kittel, Taschenb. ed. II. 773 = auricomus.
Cesatianus, *Caldes. in Nuov. Giorn. Bot. Ital.* xi. (1879) 324.—Ital.
Cesatii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 79.—Europ.
chaerophylloides, Jord. Obs. Pl. Crit. vi. (1847) 5 = chaerophyllos.
chaerophyllos, *Linn. Sp. Pl.* 555.—Reg. Mediterr.; Oriens.
Chamissoi, Spreng. Syst. ii. 650 = seq.
Chamissonis, Schlecht. Animad. Ran. i. 12. t. 1 (*Camissonis*) = glacialis.
chiclensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 28.—Peruv.
chilensis, *DC. Syst.* i. 286.—Chili.
chinensis, Bunge, Enum. Pl. Chin. Bor. 3 = pensylvanicus.
chionophilus, *Boiss. Fl. Orient.* i. 38.—Syria.
Chius, *DC. Syst.* i. 299.—Europ. austr.; As. Min.
chondrodes, Pomel, Nouv. Mat. Fl. Atl. 251 = chaerophyllos.
choorensis, Royle, Illustr. Bot. Himal. 53 = hirtellus.
chordorhizos, *Hook. f. Handb. N. Zeal. Fl.* 723.—N. Zel.
chrysanthus ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 80.—Europ.
cibiniensis, Schur, Enum. Pl. Transs. 22 = bulbosus.

RANUNCULUS:—
cicutarius, *Schlecht. Animad. Ran.* i. 25. t. 4. f. 2.—Reg. Casp.
circinatus, *Sibth. Fl. Oxon.* 175.—Europ.
circinnatoides, Arv. Ess. Pl. Dauph. 18 = divaricatus.
circinnatus, Winkler, ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 912 = fucoides.
claviformis, Dulac, Fl. Hautes-Pyr. 221 = R. Ficaria.
Clintonii, Beck, Bot. N. & M. St. 9 = repens.
cochlearifolius, Hornem. Hort. Hafn. ii. 529 = ophioglossifolius.
cochlearifolius, Ruiz & Pav. ex DC. Prod. i. 29 = macropetalus.
coenosus, Godr. in Gren. & Godr. Fl. Fr. i. 19 = Lenormandi.
coenosus, Guss. Suppl. Fl. Sic. Prod. 187 = omiophyllus.
coiliguensis, *Steud. in Flora*, xxxix. (1856) 404.—Chili.
collinus, R. Br. ex DC. Syst. i. 271 = parviflorus.
colocensis, *Menyh. Kalocsa Videk. Növényt.* 26.—Europ.
colonorum, Endl. in Enum. Pl. Hueg. 1 = lappaceus.
Columnae, All. Fl. Pedem. ii. 50 = Seguieri.
comatus, Link, ex Schlecht. Animad. Ran. i. 24, in obs. = chaerophyllos.
Commersonii, DC. ex Poir. Encyc. Suppl. iv. 662 = biternatus.
commutatus, *Badaro, in Brugnat. Giorn. Fis.* II. ix. (1826) 81.—Ital.
complicatus, *Kit. in Linnaea*, xxxii. (1863) 485.—Hungar.
concinnatus, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 182 = montanus.
confervoides, Fries, Summa Veg. Scand. 139 = omiophyllus.
confusus, *Gren. & Godr. Fl. Fr.* i. 22.—Europ.
confusus, [Clairv.] Man. Herb. 177 = parnassifolius.
cordifolius, Bast. Fl. Maine-et-Loire, 207 = ophioglossifolius.
cordifolius, *Spreng. Syst.* iv. *Cur. Post.* 218.—Reg. Argent.
cordiformis, Dulac, Fl. Hautes-Pyr. 219 = parnassifolius.
cordigerus, *Viv. Fl. Cors. Sp. Nov.* 8.—Ins. Corsica.
cornutus, Boiss. Diagn. Ser. II. v. 10 = Pinardi.
cornutus, *DC. Syst.* i. 300.—Syria.
coronatus, Ehrenb. ex Boiss. Fl. Orient. i. 56 = muricatus.
corsicus, DC. Fl. Fr. Suppl. 637 = palustris.
cortusaefolius, *Willd. Enum. Hort. Berol.* 588.—Ins. Fortunat.
Crantzii, Hedw. in Roem. Coll. Bot. 38 (*Cranzii*) = nemorosus.
crassicaulis, *Hort. Chels. ex G. Don, in Loud. Hort. Brit.* 231, nomen.—Europ.
crassipes, *Hook. f. Fl. Antarct.* 224.—Ins. Kerguel.
crenatus, Bertol. Fl. Ital. v. 514, excl. syn. = bilobus.
crenatus, *Waldst. & Kit. Pl. Rar. Hung.* i. 9. t. 10.—Europ. austr.
creticus, *Linn. Sp. Pl.* 550.—Ins. Creta.
creticus, Sieber, ex Steud. Nom. ed. II. ii. 432 = nemorosus.
crithmifolius, *Hook. f. Handb. N. Zeal. Fl.* 6.—N. Zel.
crithmifolius, Nym. Consp. 10 = glacialis.
croaticus, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 182 = Gouani.
crymophilus, *Boiss. & Hohen. Diagn.* Ser. I. viii. 6.—Persia.
Csatoi, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 46 = acris.
cubensis, *Griseb. Cat. Pl. Cub.* 1.—Cuba.
cuneatus, *Boiss. Diagn.* Ser. I. viii. 2.—As. Min.; Syria.
cuneatus, Hook. Journ. Bot. i. (1834) 242 = lappaceus.
cuneatus, Sommerf. in Mag. Naturvid. II. i. (1832) 238 = nivalis.

RANUNCULUS:—
cuneifolius, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 306.—Mongolia.
cuneilobus, A. Rich. Tent. Fl. Abyss. i. 7. t. 3 = distrias.
cupreus, *Boiss. & Heldr. Diagn.* Ser. I. viii. 3.—Ins. Creta.
curvirostratus, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iv. (1864) 313 = arvensis.
curvirostris, *Freyn, in Bot. Centralb.* vi. (1881) *Beil.* [iii.] 15.—Chili.
cyclophyllus, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 19 = monspeliacus.
cylindricus, Jord. Diagn. i. 61 = monspeliacus.
Cymbalaria, *Pursh, Fl. Am. Sept.* ii. 392.—Reg. bor. temp.
cymbalarifolius, Balb. ex Moris, Stirp. Sard. Elench. 2 = bulbosus.
dahuricus, Turcz. ex Ledeb. Fl. Ross. i. 732 = affinis.
damascenus, *Boiss. & Gaill. Diagn.* Ser. II. vi. 5.—Syria.
dasycarpus, *Boiss. Fl. Orient.* i. 28.—Persia.
daucifolius, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 270 = cicutarius.
dealbatus, Lapeyr. Hist. Abr. Pl. Pyr. 315 = aconitifolius.
debilis, *Rafin. in Med. Repos. N. York*, v. (1808) 359.—Pensylvan.
decipiens, Pomel, Nouv. Mat. Fl. Atl. 381 = palustris.
declinatus, Gilib. Fl. Lituan. ii. 270 = R. Flammula.
decumbens, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. v. (1865) 375 = diffusus.
Delacouri, Gaudefr. & Mabille, Note sur Renonc. (1874) 7; ex Bull. Soc. Dauph. fasc. iii. (1876) 69 = nemorosus.
delphinifolius, Fries, ex Steud. Nom. ed. II. ii. 432 = polyanthemos.
delphinifolius, *H. B. & K. Nov. Gen. et Sp.* v. 48.—Mexic.
delphinifolius, Torr. in Eaton, Man. Bot. ed. IV. 424 = multifidus.
delphinifolius, Torr. & Gray, Fl. N. Am. i. 659 = californicus.
demissus, *DC. Syst.* i. 275.—Hispan.; Oriens.
demissus, Orphan. ex Nym. Consp. 11 = Villarsii.
depressus, Fisch. ex Steud. Nom. ed. I. 678, nomen = praec.?
depressus, Georgi, Beschr. Russ. Reich. III. iv. 1070 = ophioglossifolius.
depressus, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 393.—N. Zel.
dertropòdius, Steud. ex A. Rich. Tent. Fl. Abyss. i. 5 = tembensis.
dichotomiflorus, *Lag. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 927.—Hispan.
dichotomus, Moç. & Sessé, ex DC. Syst. i. 288 = orthorhynchus?
diffusus, *DC. Prod.* i. 38.—Ind. or.; Malaya.
digitatus, Gilib. Fl. Lituan. ii. 266 = sceleratus.
digitatus, *Hook. Kew Journ.* iii. (1851) 124. t. 4.—Am. bor.
digitatus, Willd. ex Ledeb. Fl. Ross. i. 30 = pedatus.
dimorphorhizus, Brot. Phyt. Lusit. ii. 227 = chaerophyllos.
discolor, Steud. in Lehm. Pl. Preiss. i. 263 = leptaceus.
dissectifolius, *F. Mueli. ex Benth. Fl. Austral.* i. 11.—Austral.
dissectus, Auch. ex Tchihat. Asie Min. Bot. i. 378 = Sibthorpii.
dissectus, *Bieb. Fl. Taur. Cauc.* ii. 25.—Tauria.
dissectus, Hook. & Arn. Bot. Beech. Voy. 316 = californicus.
dissectus, Portenschl. ex Steud. Nom. ed. II. ii. 433 = alpestris.
distans, Royle, Illustr. Bot. Himal. 53 = laetus.
distrias, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 7.—Abyss.
divaricatus, A. Gray, Pl. Wright. ii. 8 = aquatilis.
divaricatus, Koch, Syn. Fl. Germ. ed. I. 12 = circinatus.
divaricatus, *Schrank, Baier. Fl.* ii. 103; *Moench, Meth.* 214.—Europ.
divergens, Jord. Obs. Pl. Crit. vi. (1847) 14 = Sprunerianus.
diversifolius, *Boiss. & Kotschy, in Boiss. Fl. Orient.* i. 45.—Armenia.

RANUNCULUS:—
diversifolius, Gilib. Fl. Lituan. ii. 262 = aquatilis.
dodecapetalus, Steph. ex Ledeb. Fl. Ross. i. 47 = Oxygraphis glacialis.
dolosus, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 384.—Reg. Casp.
Donianus, *Pritz. in Walp. Rep.* ii. 740.—Mexic.
Drouetii, F. Schultz, Arch. Fl. Fr. et Allem. 10 = trichophyllus.
dubius, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 909.—Hispan.
echinatus, Crantz, Stirp. Austr. ed. I. fasc. ii. 95; ed. II. 118 = arvensis.
echinatus, Vent. Jard. Cels, 73 = muricatus.
edulis, *Boiss. & Hohen. Diagn.* Ser. I. viii. 4.—As. Min.; Persia.
Eisenii, Kellogg, in Proc. Calif. Acad. vii. (1877) 115 = Nelsonii?
elatus, *Steud. in Flora*, xxxix. (1856) 405.—Chili.
elbrusensis, *Boiss. Fl. Orient.* i. 34.—Persia.
elegans, C. Koch, in Linnaea, xv. (1841) 248 = anemonefolius.
eleophilus, Arv. Ess. Pl. Dauph. 19 (Quid?).—Gallia.
elongatus, *F. Schultz, in Billotia*, i. (1869) 113.—Europ.
Enysii, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 394.—N. Zel.
ericetorum, Pourr. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 922 = escurialensis.
erinifolius, *Tenore, Syll. Fl. Neap.* 273.—Ital.
eriocalyx, *Scheele, in Flora*, xxvi. (1843) 301.—Helvet.
eriocarpus, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 351.—Cappadoc.
eriophyllus, C. Koch, in Linnaea, xix. (1847) 46 = palustris.
eriorrhizus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 5.—Persia.
eriotorus, Jord. Diagn. i. 69 = Villarsii.
erodiifolius, *C. Gay, Fl. Chil.* i. 45.—Chili.
Eschscholtzii, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 241, partim = affinis.
Eschscholtzii, Schlecht. Animad. Ran. ii. 16. t. 1 = nivalis.
escurialensis, *Boiss. & Reut. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 922.—Hispan.
eustrigulosus, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 44 = acris.
exiguus, Urv. in Mém. Soc. Linn. Par. iv. (1826) 615 = biternatus.
exilis, *Phil. Fl. Atac.* 7.—Chili.
extorris, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 204 = ternatus.
falcatus, *Linn. Sp. Pl.* 556.—Europ.; Reg. Himal.
fallax, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 37 = cassubicus.
fascicularis, Barton, Comp. Fl. Philad. ii. 25 = repens.
fascicularis, *Muhl. ex Bigel. Fl. Bost.* ed. I. 137.—Am. bor.
fascicularis, Spreng. Neue Entdeck. i. 228 = recurvatus.
fascicularis, S. Wats. Bot. King's Exp. 9 = californicus.
feniculaceus, Gilib. Exercit. i. 370 = aquatilis.
Fenzlii, *Boiss. Fl. Orient.* i. 44.—Cilicia.
fibrillosus, *C. Koch, in Linnaea*, xix. (1847) 47.—Bithynia.
fibrosus, Pomel, Nouv. Mat. Fl. Atl. 251 = chaerophyllos.
fibrosus, Wall. Cat. n. 4706 = pensylvanicus.
Ficaria, *Linn. Sp. Pl.* 550.—Europ.; Reg. Caucas.
Ficaria, Walt. Fl. Carol. 159 = Caltha palustris.
ficariaeformis, F. Schultz, Arch. Fl. 123 = R. Ficaria.
ficarioides, *Bory & Chaub. Nouv. Fl. Pélop.* 34.—Graecia.
filamentosus, *Wedd. Chlor. And.* ii. 301.—Bolivia.
filiformis, Michx. Fl. Bor. Am. i. 320 = R. Flammula.
fistulosus, Brign. Fasc. Pl. Forojul. 25 = ophioglossifolius.
fistulosus, Pursh, ex Torr. in Ann. Lyc. N. York, ii. (1828) 163 = multifidus.
flabellaris, Rafin. in Am. Monthly Mag. (1818) 344 = multifidus.
flabellatus, Bivona, Sic. Pl. Cent. i. 68 = bulbosus.
flabellatus, Desf. Fl. Atlant. i. 438. t. 114 = chaerophyllos.
flabellatus, Heuff. ex Kittel, Taschenb. ed. II. 773 = auricomus.

RANUNCULUS:—
flabellifolius, Heuff. ex Reichb. Fl. Germ. Excurs. 723 = cassubicus.
flaccidus, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 224 = biternatus.
flaccidus, *Hook. f. & Thoms. Fl. Ind.* i. 38.—Reg. Himal.
flaccidus, Pers. in Usteri, Ann. Bot. xiv. (1795) 39 = aquatilis.
flagelliformis, *Sm. in Rees, Cycl.* xxix. n. 13.—Am. austr.
Flamma, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 389 = R. Flammula.
flammeus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 719 = R. Flammula.
Flammula, D. Don, in Royle, Illustr. Bot. Himal. 53 = pulchellus.
Flammula, *Linn. Sp. Pl.* 548.—Reg. bor. temp.
Flammula, Michx. Fl. Bor. Am. i. 321 = oblongifolius.
Flammula, Pursh, Fl. Am. Sept. ii. 391 = obtusiusculus.
Flammula, Walt. Fl. Carol. 159 = pusillus.
Flammulae, Krock. Fl. Siles. ii. 1. 248 = R. Flammula.
flammuloides, Rafin. ex DC. Prod. i. 32 = R. Flammula.
floribundus, Bab. in Trans. Bot. Soc. Edinb. v. (1858) 77 = peltatus.
fluitans, *Lam. Fl. Fr.* iii. 184.—Europ.
fluviatilis, Bigel. Fl. Bost. ed. I. 139 = multifidus.
fluviatilis, Pursh, Fl. Am. Sept. ii. 395 = aquatilis.
fluviatilis, [Weber, ex] Wigg. Prim. Fl. Holsat. 42 = fluitans.
fluviatilis, Willd. Sp. Pl. ii. 1333 = peucedanifolius.
foeniculaceus, Gilib. Fl. Lituan. ii. 261 = divaricatus.
fontanus, *Presl, Delic. Prag.* 6.—Sicilia.
Forskoehlii, *DC. Syst.* i. 303.—Arab.
Franchetianus, Bor. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 8 = acris.
fraternus, *Schrenk, Enum. Pl. Nov.* (1841) 103.—Soongaria.
Friesianus, Jord. Obs. Pl. Crit. vi. (1847) 17 = acris.
Friesii, *Gren. & Godr. Fl. Fr.* i. 34.—Gallia.
frigidus, Schrank, Pl. Rar. Hort. Monac. t. 57 = lacerus.
frigidus, Willd. Sp. Pl. ii. 1312; DC. Prod. i. 35 = nivalis.
fucoides, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 912.—Hispan.
fumariaefolius, Desf. ex DC. Syst. i. 256 = millefoliatus.
furcatus, Berger. Fl. Bass. Pyr. ii. 409 = Gouani.
Galeottii, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 276.—Mexic.
gallecicus, *Freyn, ex Willk. Illustr. Fl. Hisp. Balear.* i. 100.—Hispan.
garganicus, Tenore, Ind. Sem. Hort. Neap. (1830) 13 = millefoliatus.
gelidus, Hoffmgg. ex Reichb. Fl. Germ. Excurs. 720 = glacialis.
gelidus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 133.—As. centr.
gelidus, Schur, Enum. Pl. Transs. 18 = montanus.
geoides, *H. B. & K. Nov. Gen. et Sp.* v. 47.—Mexic.
geoides, Siev. in Pall. N. Nord. Beitr. vii. 201 = ophioglossifolius.
geraniifolius, *Hook. f. Fl. N. Zel.* i. 9.—N. Zel.
geraniifolius, Pourr. in Mém. Acad. Toul. iii. (1788) 326 = montanus.
geranioides, Blume, Bijdr. 2 = diffusus.
geranioides, *H. B. & K. Nov. Gen. et Sp.* v. 44.—N. Granat.
giganteus, *Wedd. Chlor. And.* ii. 304.—Peruv.
Gilliesii, *Steud. in Flora*, xxxix. (1856) 404.—Chili.
glabellus, Royle, Illustr. Bot. Himal. 53 = hirtellus.
glaberrimus, *Hook. Fl. Bor. Am.* i. 12.—Am. bor. occ.
glabratus, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 18 = hirtellus.
glabrifolius, Hook. Journ. Bot. i. (1834) 243 = rivularis.
glabriusculus, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 19.—Terr. Samoyed.
glacialis, *Linn. Sp. Pl.* 553.—Reg. bor. et arct.
glandulifer, *Poepp. Fragm. Syn. Pl. Chile*, 28.—Chili.
Gmelini, *DC. Syst.* i. 303.—Sibir.
Gmelini, Schlecht. Animad. Ranunc. ii. 35 = multifidus.
Godleyanus, *Hook. f. Handb. N. Zeal. Fl.* 723.—N. Zel.

RANUNCULUS:—
Goldei, *Meinsh. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 44, nomen.—Rossia.
Gonnetii, Jord. Diagn. i. 65 = monspeliacus.
Gouani, Hegetschw. Fl. Schw. i. 402 = Villarsii.
Gouani, Lecoq. & Lamotte, Cat. Pl. Vasc. Fr. 52 = nemorosus.
Gouani, Sm. in Rees, Cycl. xxix. n. 41 = montanus.
Gouani, *Willd. Sp. Pl.* ii. 1322.—Mont. Pyren.
gracilipes, *Hook. f. Handb. N. Zeal. Fl.* 8.—N. Zel.
gracilis, Bory & Chaub. Nouv. Fl. Pélop. 35 = chaerophyllos.
gracilis, *Clarke, Trav.* iii. 336; *DC. Syst.* i. 256.—Ins. Cos.
gracilis, Schleich. Cat. Pl. Helv. ed. III. 24 = montanus.
graecus, Griseb. Spicil. Fl. Rumel. i. 314 = muricatus.
gramineus, Habl. Phys. Beschr. Taur. 149 = illyricus.
gramineus, *Linn. Sp. Pl.* 549.—Europ.; Marocc.
gramineus, Spreng. Syst. ii. 648 = longicaulis.
graminifolius, Salisb. Prod. 372 = gramineus.
granatensis, Boiss. Diagn. Ser. II. i. 8 = acris.
grandiflorus, *Linn. Sp. Pl.* 555.—Reg. Caucas.; Cappadoc.
grandiflorus, Payot, Fl. Mont-Blanc, 13 (Quid?).—Europ.
grandifolius, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 330.—Sibir. altaic.
grandifolius, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 38 = megaphyllus.
granulatus, Griseb. Spicil. Fl. Rumel. i. 306; Pomel, Nouv. Mat. Fl. Atl. 380 = chaerophyllos.
gregarius, Brot. Fl. Lusit. ii. 369 = chaerophyllos.
Grenieranus, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 304 = Villarsii.
gruinalis, Schott, in Nym. & Kotschy, Analect. Bot. 33 = Gouani.
Guilelmi-Jordani, *Aschers. in Verh. Bot. Ver. Brand.* xxi. (1880) 64.—Aegypt.
Gunnae, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 666.—Abyss.
Gunnianus, *Hook. Journ. Bot.* i. (1834) 244. t. 133.—Austral.
Guzmanni, *Humb. Tabl.* 69; *ex DC. Syst.* i. 263.—Reg. Andin.
Haarbachii, *De Not. ex Cesati, in Bibl. Ital.* xci. (1838) 347.—Ital.
Haastii, *Hook. f. Handb. N. Zeal. Fl.* 6.—N. Zel.
Halleri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 78.—Europ.
halophilus, Schlecht. Animad. Ran. i. 23. t. 4. f. 1 = R. Cymbalaria.
hastatus, Walker, ex Wight, Illustr. i. 5 = sagittifolius.
hawaiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 10.—Ins. Sandvic.
Haynaldi, *Menyh. Kalocsa Videk. Növényt.* 28.—Europ.
hebecarpus, *Hook. & Arn. Bot. Beech. Voy.* 316.—Am. bor. occ.
hederaceus, Bourg. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 908 = ololeucos.
hederaceus, *Linn. Sp. Pl.* 556.—Europ. occ.
hederaceus, Pall. Reise, ii. 511 = hyperboreus.
hederaefolius, Salisb. Prod. 373 = hederaceus.
Hegetschweileri, *Ducommun, Taschenb.* 18.—Helvet.
Heldreichanus, Jord. Obs. Pl. Crit. vi. (1847) 14 = Sprunerianus.
hemignostus, *Steud. in Flora*, xxxix. (1856) 404.—Peruv.
Henriquesii, *Freyn, in Bot. Centralb.* vi. (1881) *Beil.* [iii.] 21.—Europ.
heterophyllus, Lapeyr. Hist. Abr. Pl. Pyr. 316 = aconitifolius.
heterophyllus, Sm. in Rees, Cycl. xxix. n. 12 = bonariensis.
heterophyllus, [Weber, in] Wigg. Prim. Fl. Holsat. 42 = aquatilis.
heterorrhizus, *Boiss. & Bal. Diagn.* Ser. II. vi. 6.—Phrygia.
heucheraefolius, Presl, Fl. Sicul. 15 = palustris.
hierosolymitanus, *Boiss. Fl. Orient.* i. 36.—Syria.
Hilarii, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 209.—Reg. Argent.
hirsutus, Curt. Fl. Lond. Fasc. ii. t. 40 = sardous.
hirtellus, *Royle, Illustr. Bot. Himal.* 53.—Reg. Himal.
hirtus, Banks & Soland. ex DC. Syst. i. 289 = plebeius.

RANUNCULUS:—
hirtus, Hook. f. Fl. Tasman. i. 6 = lappaceus.
hirtus, Muhl. ex Steud. Nom. ed. II. ii. 433 = recurvatus.
hispidus, Hook. Fl. Bor. Am. i. 19, partim = occidentalis, *Nutt.*
hispidus, Michx. l. c. 321 = repens.
hispidus, Pursh, Fl. Am. Sept. ii. 395 = pensylvanicus.
Hohenackeri, *Presl, Bot. Bemerk.* 7, nomen.—Reg. Cauc.
Hollianus, *Reichb. in Flora*, xiii. (1830) 130.—Lusitania.
hololeucus Auct. = ololeucos.
holophyllus, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 220 = sceleratus.
homoeophyllus, Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 906 = omiophyllus.
Hookeri, *Regel, Fl. Ost-Sibir.* i. 47.—As. bor. or.; Am. bor. occ.
Hookeri, Schlecht. in Linnaea, ix. (1834) 610 = repens?
Hornemanni, *Schlecht. Animad. Ran.* ii. 36.—Am. bor.
Hornschuchii, Hoppe, in Sturm, Deutschl. Fl. Heft 46 = Villarsii.
hortensis, Pers. Syn. ii. 104 = asiaticus.
hortensis, Schur, Enum. Pl. Transs. 21 = repens.
Huetii, *Boiss. Diagn.* Ser. II. v. 7.—Armenia.
humilis, Collie, ex Hook. Bot. Beech. Voy. 4. t. 2 = obtusatus.
humilis, D. Don, ex G. Don, Gen. Syst. i. 34 = Donianus.
humilis, Huet, ex Nym. Consp. 14, in obs. = parviflorus?
humilis, Pers. Syn. ii. 102 = pusillus.
hybridus, *Biria, Hist. Nat. Renoncul.* 38.—Europ. austr.
hybridus ×, Payot, Fl. Mont-Blanc, 12.—Europ.
hydrocharis, Spenn. Fl. Friburg. iii. 1007 = aquatilis.
hydrocharoides, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 306.—Mexic.
Hydrocotile, Domb. ex DC. Syst. i. 251 = flagelliformis.
hydrocotyloides, Wall. Cat. n. 4703 = diffusus.
hydrophilus, Bunge, Enum. Pl. Chin. Bor. 2 = aquatilis.
hydrophilus, *Gaudich. ex Mirb. in Ann. Sc. Nat.* Sér. I. v. (1825) 105.—Ins. Falkland.
hyperboreus, *Rottb. in Skrift. Kjoeb. Selsk.* x. (1770) 458.—Reg. bor. et arct.
hypotrichus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 275 [bis].—Rumel.
hystriculus, A. Gray, in Proc. Am. Acad. vii. (1868) 328 = Kumlienia hystricula.
illyricus, Gouan, Fl. Monsp. 269 = monspeliacus.
illyricus, *Linn. Sp. Pl.* 552.—Europ. austr.; Reg. Caucas.
illyricus, Poir. Encyc. vi. 121 = chaerophyllos.
incisifolius, Reichb. ex Kittel, Taschenb. ed. II. 773 = auricomus.
incisus, Hook. f. Fl. N. Zel. i. 10 = rivularis.
incomparabilis, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 174.—Thracia.
inconspicuus, Hook. f. Fl. Tasman. i. 9. t. 2 = rivularis.
incrassatus, Guss. Fl. Sic. Syn. ii. 50 = chius.
indicus, Roxb. Hort. Beng. 43; Fl. Ind. ii. 671 = sceleratus.
infestus, Salisb. Prod. 373 = repens.
inops, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 183 = Villarsii.
insignis, *Hook. f. Fl. N. Zel.* i. 8. t. 2.—N. Zel.
insubricus, *Cesati, in Bibl. Ital.* xci. (1838) 346.—Ital.
insularis, Viviani, Fl. Cors. Prod. App. i. 2 = chaerophyllos.
intermedius, Eaton, Man. Bot. ed. III. 424 = repens.
intermedius, Knaf, in Flora, xxix. (1846) 289 = peltatus.
intermedius, Poir. Encyc. vi. 116 = sardous.
intermedius, Schur, Enum. Pl. Transs. 14 = R. Ficaria.
inundatus, R. Br. ex DC. Syst. i. 269 = rivularis.
involucratus, Lapeyr. Hist. Abr. Pl. Pyr. 318 = Gouani.
isopyroides, DC. Syst. i. 238 = Callianthemum rutaefolium.
isthmicus, *Boiss. Diagn.* Ser. I. vi. 4.—Graecia.
Jacquini, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 1020 = nivalis.
japonicus, Langsd. ex DC. Prod. i. 38 = Langsdorfii.

RANUNCULUS :—

japonicus, Thunb. in Trans. Linn. Soc. ii. (1794) 337 = acris.
javanicus, *Blume*, *Bijdr.* 3.—Java.
kamchaticus, DC. Syst. i. 302 = Oxygraphis glacialis.
Kaufmanni, *Clerc*, *ex Trautv. in Act. Hort. Petrop.* viii. (1883) 45, nomen.—Ross.
Kayserii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 43 = acris.
Kladnii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 42 = acris.
Kochii, Beurl. in Bot. Notiser (1852), 156 = seq.?
Kochii, *Ledeb. Fl. Ross.* i. 731.—Reg. Caucas.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 5.—Persia.
Krapfia, *DC. in Deless. Ic. Sel. Pl.* i. 9. t. 35.—Chili.
Kunthii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 12.—N. Granat.
lacerus, *Bell. in Mém. Acad. Turin*, v. (1790–91) 233. t. 8.—Gallia.
lacerus, Mutel, Fl. Fr. i. 18 = pyrenaeus.
laciniatus, Baumg. Enum. Stirp. Transs. ii. 131 = sardous.
laciniatus, [Clairv.] Man. Herb. 177 = lacerus.
lacustris, Beck & Tracy, in Trans. Albany Inst. i. (1830) 148 ; Bot. N. & Mid. St. 9 = multifidus.
laetus, Salisb. Prod. 373 = bulbosus.
laetus, *Wall. Cat.* n. 4702.—Reg. Himal.
Lagascanus, *DC. Prod.* i. 43.—Hispan.
Lambertianus, *D. Dietr. Syn. Pl.* iii. 316.—Mexic.
lanceolatus, [Clairv.] Man. Herb. 177 = pyrenaeus.
lanceolatus, Pall. ex DC. Prod. i. 32 = R. Flammula.
lancifolius, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iii. (1863) 423 = R. Cymbalaria.
lancipetalus, *Griseb. in Goett. Abh.* xix. (1874) 69.—Reg. Argent.
Langsdorfii, DC. Prod. i. 34 = multifidus.
Langsdorfii, *Spreng. Syst.* ii. 652.—Japon.
lanuginosaeformis, Selin. ex Fellm. Pl. Lapp. Or. 3 = acris.
lanuginosus, Amo, Fl. Iber. vi. 720 = nemorosus.
lanuginosus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 6 = constantinopolitanus.
lanuginosus, Costa, Introd. Fl. Catal. 6 = sardous.
lanuginosus, Ledeb. Fl. Ross. i. 42 = acris.
lanuginosus, *Linn. Sp. Pl.* 554.—Europ. ; Reg. Caucas.
lanuginosus, Pursh, Fl. Am. Sept. ii. 394 = repens.
lanuginosus, Walt. Fl. Carol. 159 = recurvatus.
lappaceus, *Sm. in Rees, Cyclop.* xxix. n. 61.—Austral.
lapponicus, *Linn. Sp. Pl.* 553.—Reg. bor. et arct.
lapponicus, Oed. Fl. Dan. t. 144 = nivalis.
lapponicus, Soleir. ex Nym. Consp. 10 = demissus.
lapponicus, Vill. Hist. Pl. Dauph. iii. 743 = Villarsii.
lasiocarpus, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 323.—Sibir. altaic.
lasiostemon, Fenzl, Pugill. Pl. Nov. Syr. 15 = Fenzlii.
lateriflorus, *DC. Syst.* i. 251.—Europ. austr. ; Oriens.
latilobus, Kit. in Linnaea, xxxii. (1863) 485 = complicatus.
laxicaulis, Darby, Bot. S. St. 204 = obtusiusculus.
Lechleri, *Schlecht. in Linnaea*, xxvii. (1854) 557.—Peruv.
Lecokii, Bor. Fl. Centr. Fr. ed. III. ii. 17 = nemorosus.
leiodiscus, Boiss. & Huet, ex Boiss. Fl. Orient. i. 55 = marginatus.
Lemmoni, *A. Gray*, *in Proc. Am. Acad.* x. (1874) 68.—Am. bor. occ.
Lenormandi, *F. Schultz*, *in Flora*, xx. (1837) 726.—Europ.
leontinensis, *Freyn*, *in Willk. & Lange*, *Prod. Fl. Hisp.* iii. 910.—Hispan.
leptaleus, *DC. Syst.* i. 258.—Ins. Cyprus.
leptocaulis, Hook. Journ. Bot. i. (1834) 244 = parviflorus.
leptopetalus, Rafin. Fl. Ludov. 83 = recurvatus.
Lerchenfeldianus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 14 = montanus.
leucothrix, Ball, in Journ. Bot. xi. (1873) 296 = chaerophyllos.
Limosella, *F. Muell. ex T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 177.—N. Zel.
limoselloides, *F. Muell. ex Hook. f. Ic. Pl.* t. 1081.—N. Zel.
limoselloides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 275 [bis].—Venezuela.
limosus, Nutt. in Torr. & Gray, Fl. N. Am. i. 20 = multifidus.
linearilobus, *Bunge*, *in Arb. Naturf. Ver. Riga*, i. (1847) 122.—Turkestan.

RANUNCULUS :—

Lingua, *Linn. Sp. Pl.* 549.—Europ. ; Sibir.
Lingua, Pursh, Fl. Am. Sept. ii. 391 = obtusiusculus.
litoralis, *Phil. in Linnaea*, xxviii. (1856) 663.—Chili.
litoralis, *Schlecht. in Lechl. Berb. Am. Austr.* 54, nomen.—Chili.
Llaveanus, *Schlecht. in Linnaea*, x. (1836) 233.—Mexic.
lobatus, [Clairv.] Herb. Man. 177 = aconitifolius.
lobatus, *Jacquem. Voy. Bot.* 4.—Reg. Himal.
lobatus, Moench, Meth. 214 = muricatus.
lomatocarpus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 36.—Oriens.
longepedunculatus, *Scheidw. in Hortic. Belge*, v. (1838) 163. t. 9.—Mexic.
longicaulis, *Ledeb. ex Spreng. f. Tent. Suppl. Syst.* 15.—Europ. austr.
longicaulis, C. A. Mey. in Ledeb. Fl. Alt. ii. 308 = pulchellus.
longifolius, Lam. Fl. Fr. iii. 189 = R. Lingua.
longipes, Lange, ex Cut. Fl. Madr. 103 = pedunculatus.
longirostris, *Godr. in Mém. Soc. Roy. Nancy*, 1839 (1840) 39.—Am. bor.
Lorentii, Hochst. in Lorent, Wanderungen, 338 = lomatocarpus.
lucidus, Poir. Encyc. vi. 113 = repens.
lugdunensis, Jord. Diagn. i. 63 = monspeliacus.
lusitanicus, *Freyn*, *in Flora*, lxiii. (1880) 24.—Lusitan.
lutulentus, *Songeon & Perrier*, *in Billot*, *Annot. Fl. Fr. et Allem.* 181.—Europ.
luzulaefolius, Amo, Fl. Iber. vi. 714 = gramineus.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 4.—N. Zel.
lycoctonifolius, Hegetschw. Fl. Schw. 544 = montanus.
Macauleyi, *A. Gray*, *in Proc. Am. Acad.* xv. (1880) 45.—Am. bor.
maclovianus, *Urv. in Mém. Soc. Linn. Par.* iv. (18 6) 615.—Ins. Falkland.
macranthus, *Scheele*, *in Linnaea*, xxi. (1848) 585.—Texas ; Mexic.
macrocarpus, Pritz. Ic. Ind. ii. 239, sphalm. = macropus, *Boiss. & Buhse.*
macropetalus, *DC. Prod.* i. 29.—Peruv.
macrophyllus, Desf. Fl. Atlant. i. 437 = palustris.
macrophyllus, Ledeb. Fl. Ross. i. 42 = grandiflorus.
macrophyllus, Schur, Enum. Pl. Transs. 17 = acris.
macropus, *Boiss. & Buhse*, *in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 5.—Persia.
macropus, *Hook. f. in Hook. Ic. Pl.* t. 634.—N. Zel.
macrorhizus, *Schnizl. Ic.* t. 178. f. 1.—Hab. ?
macrorrhynchus, *Boiss. Diagn.* Ser. I. vi. 5.—Mesopot.
madagascariensis, *Freyn*, *in Bremen*, *Abh.* vii. (1880) 7.—Madag.
magellanicus, Comm. ex DC. Syst. i. 294 = petiolaris.
magellensis, Tenore, Prod. Fl. Nap. 69 = crenatus.
major, Dum. Fl. Belg. Prod. 128 = acris.
Malabailae, *Boiss. Fl. Orient.* i. 37.—Cilicia.
malacophyllus, Schur, Enum. Pl. Transs. 16 = acris.
Mandonianus, *Wedd. Chlor. And.* ii. 299.—Bolivia.
marginatus, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 318.—Graec.
marilandicus, Poir. Encyc. vi. 126 = repens.
marinus, Fries, Summa Veg. Scand. 141 = Baudotii.
maritimus, *Phil. in Anal. Univ. Chil.* (1872) 663.—Chili.
Marschlinsii, Steud. Nom. ed. II. ii. 434 = demissus.
marsicus, Guss. & Tenore, in Fl. Neap. Syll. App. iv. p. ix = auricomus.
Martini, Lamotte, Prod. Fl. Centr. Fr. i. 43 = trichophyllus.
marylandicus, Poir. Encyc. vi. 126 = repens.
mauiensis, *A. Gray*, *Bot. U. St. Expl. Exped.* i. 11.—Ins. Sandvic.
mauritanicus, Pomel, Nouv. Mat. Fl. Atl. 249 = omiophyllus.
mediterraneus, Griseb. ex Schur, Enum. Pl. Transs. 22 = sardous.
medius, Bernh. Cat. Pl. Hort. Erfurt. ; ex Schlecht. Animad. Ran. i. 14 = aconitifolius.
meganthus, Rafin. Fl. Ludov. 82 (Quid ?).—Am. bor.
megaphyllus, *Steud. Nom.* ed. II. ii. 434.—Ins. Mader.
meifolius, Pomel, Nouv. Mat. Fl. Atl. 380 = millefoliatus.
Meinshauseni, *Schrenk*, *in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 309.—Soongar.
membranaceus, Fresen. in Mus. Senckenb. ii. (1837) 270 = pinnatus.

RANUNCULUS :—

membranaceus, Royle, Illustr. Bot. Himal. 53 = pulchellus.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* i. 7.—Afr. austr.
Meyerianus, *Rupr. Fl. Cauc.* 25.—Reg. Caucas.
micranthus, *Bruegg. in Zeitschr. Ferdinand. Tirol.* III. ix. (1860) 7.—Europ.
micranthus, Nutt. in Torr. & Gray, Fl. N. Am. i. 18 = abortivus.
microcarpus, Bertol. Misc. Bot. xxii. 17. t. 6 = muricatus.
microcarpus, *Presl*, *Rel. Haenk.* ii. 70.—Chili.
Millani, *F. Muell. in Hook. Kew Journ.* vii. (1855) 358.—Austral.
millefoliatus, *Vahl*, *Symb. Bot.* ii. 63.—Reg. Mediterr.
millefolius, [Soland. in] Russ. Aleppo, ed. II. ii. 254 = myriophyllus.
Millii, *Boiss. & Heldr. ex Boiss. Fl. Orient.* i. 35.—Graec.
minimus, Hort. ex Schlecht. Animad. Ran. i. 19 = ophioglossifolius.
minutiflorus, *Bert. ex Phil. in Linnaea*, xxviii. (1856) 661.—Chili.
minutus, *C. Gay*, *Fl. Chil.* i. 38.—Chili.
minutus, Leyb. in Flora, xxxvii. (1854) 369 = montanus.
miser, *Phil. in Linnaea*, xxviii. (1856) 662.—Chili.
mitis, Gilib. Fl. Lituan. ii. 268 = auricomus.
mixtus, Jord. Obs. Pl. Crit. vii. (1849) 1 = nemorosus.
Moellendorffii, *Hance*, *in Journ. Bot.* xvii. (1879) 7.—China.
mollis, Wall. Cat. n. 4704 = diffusus.
monanthos, *Phil. in Linnaea*, xxviii. (1856) 662.—Chili.
Monroi, *Hook. f. Fl. N. Zel.* ii. 323.—N. Zel.
monspeliacus, Georgi, Beschr. Russ. Reich. III. iv. 1067 = illyricus.
monspeliacus, *Linn. Sp. Pl.* 553.—Reg. Mediterr.
monspeliacus, Winkl. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 922 = Winkleri.
monspessulanus, Jord. Diagn. i. 64 = monspeliacus.
montanus, Orsini, ex Tenore, Syll. 270 = Villarsii.
montanus, Presl, ex Nym. Consp. 12 = acris.
montanus, Sibth. & Sm. Fl. Graec. Prod. ii. 359 = Sibthorpii.
montanus, *Willd. Sp. Pl.* ii. 1321.—Europ.
montteanus, *Phil. in Linnaea*, xxx. (1859–60) 185.—Chili.
Moorcroftianus, D. Don, in Royle, Illustr. Bot. Himal. 53 = pulchellus.
moschatus, *Uspensky*, *in Bull. Soc. Nat. Mosc.* vii. (1834) 375, nomen.—Rossia.
Moseleyi, *Hook. f. in Phil. Trans.* clxviii. (1879) 10.—Ins. Kerguel.
mucronatus, Bertol. in Nov. Comm. Acad. Bonon. vi. (1843) 224. t. 8. f. 2 = oxyspermus.
Muelleri, *Benth. Fl. Austral.* i. 13.—Austral.
mugellensis, Steud. Nom. ed. I. 679, sphalm. = *magellensis.*
multicaulis, *D. Don*, *ex G. Don*, *Gen. Syst.* i. 34.—Mexic.
multifidus, Forsk. Fl. Aegypt. Arab. 102 = Forskoehlii.
multifidus, *Pursh*, *Fl. Am. Sept.* ii. 736.—Am. bor.
multiflorus, Dulac, Fl. Hautes-Pyr. 218 = aconitifolius.
multiflorus, Gilib. Fl. Lituan. ii. 263 = polyanthemos.
multiscapus, Hook. f. Fl. N. Zel. i. 9. t. 5 = lappaceus.
muricatus, *Linn. Sp. Pl.* 555.—Europ. ; Oriens.
muricatus, Moench, Meth. 215 = parviflorus.
myosuroides, *Boiss. & Kotschy*, *Diagn.* Ser. II. v. 6.—Syria.
Myosurus, Afzel. in Liljebl. Svensk Fl. 230 = Myosurus minimus.
myriophyllus, *DC. Syst.* i. 257.—As. Min. ; Mesopot.
nanus, Fisch. ex DC. Prod. i. 33 = R. Cymbalaria.
nanus, Hook. Journ. Bot. i. (1834) 242 = lappaceus.
napaulensis, DC. Prod. i. 39 = diffusus.
napellifolius, Crantz, Stirp. Austr. ed. I. fasc. ii. 90 ; ed. II. 114 = acris, nemorosus, polyanthemos.
napellifolius, *DC. Syst.* i. 282.—Graec.
natans, C. A. Mey. in Ledeb. Fl. Alt. ii. 315 = hyperboreus.
natans, *Nees*, *ex G. Don*, *Gen. Syst.* i. 31.—Mexic.
neapolitanus, Bourg. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 930, 931 = Broteri, Aleae.

RANUNCULUS:—
neapolitanus, Tenore, Ind. Sem. Hort. Neap. (1825) 11. n. 22 = bulbosus.
neapolitanus, Tod. ex Freyn, Kennt. Ranunc. 20 = palustris.
Nelsonii, *A. Gray, in Proc. Am. Acad.* viii. (1872) 374. —Am. bor. occ.
nemorivagus, Jord. Diagn. i. 74 = acris.
nemorosus, *DC. Syst.* i. 280.—Europ. austr.
Nendtwichii, Frivald. in Flora, xix. (1836) 437 = lateriflorus.
nepalensis, Spreng. Syst. ii. 655 = diffusus.
nepalinus, Jacquem. ex Aitch. in Journ. Linn. Soc. xviii. (1880) 30 = hirtellus.
nephelogenes, Edgew. in Trans. Linn. Soc. xx. (1846) 28 = pulchellus.
nervosus, Royle, Illustr. Bot. Himal. 53 = hirtellus.
nevadensis, *Willk. in Linnaea*, xxx. (1859–60) 85.—Hispan.
nidulans, Pomel, Nouv. Mat. Fl. Atl. 252 = chaerophyllos.
nigrescens, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 921.—Hispan.
nitidus, Muhl. Cat. ed. II. 56 = repens.
nitidus, Walt. Fl. Carol. 159 = abortivus?
nivalis, Lag. in Anal. Cienc. Nat. v. (1804) 280 = demissus.
nivalis, Lerchew. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 14 = montanus.
nivalis, *Linn. Sp. Pl.* 553.—Reg. bor. et arct.
nivalis, Pall. Reise, iii. 653 = polyrhizos.
nivalis, Sibth. & Sm. Fl. Graec. Prod. i. 382 = Sibthorpii.
nivicola, *Hook. Ic. Pl.* t. 571.—N. Zel.
nodiflorus, *Linn. Sp. Pl.* 549.—Europ.
nubigenus, *H. B. & K. Nov. Gen. et Sp.* v. 42.—Peruv.
Nuttallii, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 56.—Am. bor. occ.
obesus, *Trautv. in Act. Hort. Petrop.* iii. 11. (1875) 275 —Reg. Caucas.
oblongifolius, *Ell. Sketch*, ii. 58.—Am. bor. occ.
obtectus, Wall. Cat. n. 4705 = diffusus.
obtusatus, *Poepp. Fragm. Syn. Pl. Chil.* 28.—Chili.
obtusifolius, *Hornem. Hort. Hafn.* ii. 528.—Hispan.
obtusiusculus, *Rafin. in Med. Repos. N. York*, v. (1808) 359; *et in Desv. Journ. Bot.* i. (1808) 225. —Am. bor.
occidentalis, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 929.—Hispan.
occidentalis, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 22, excl. syn.—Am. bor. occ.
occidentalis, Torr. & Gray, l. c., partim = Nelsonii.
oenanthefolius, Tenore & Guss. in Atti Acc. Nap. v. 1. (1843) 318 = repens.
Olgae, *Regel, Pl. Nov. Fedsch.* 1.—Turkestan.
oligocarpos, *Höchst. ex A. Rich. Tent. Fl. Abyss.* i. 5. —Abyss.
oligocarpus, *Bruegg. in Zeitschr. Ferdinand. Tirol.* III. ix. (1860) 7.—Europ.
ololeucos, *Lloyd, Fl. Loire-Inf.* 3.—Europ.
olyssiponensis, Pers. Syn. ii. 106 = spicatus.
omiophyllus, *Tenore, Fl. Nap.* iv. 338.—Europ.
ophioglossifolius, *Vill. Hist. Pl. Dauph.* iii. 731.—Europ. occ.; Reg. Mediterr.
ophioglossoides, Willd. Sp. Pl. ii. 1310 = ophioglossifolius.
oppositifolius, *Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 513.—Hungar.
orbicularis, *Steud. in Flora*, xxxix. (1856) 406.—Chili.
oreophilus, Bieb. Fl. Taur. Cauc. iii. 383 = Villarsii.
oreophilus, Walp. Ann. iii. 811, sphalm. = seq.
oreophytus, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 89.—Afr. trop.
orientalis, *Linn. Sp. Pl.* 555.—Afr. bor.; Oriens.
ornithopodioides, Stev. ex Ledeb. Fl. Ross. i. 30 = pedatus.
ornithorhynchus, Walp. Rep. i. 43, sphalm. = orthorhynchus.
orthophytus, Walp. l. c. v. 5, sphalm. = oreophytus.
orthorhynchus, *Hook. Fl. Bor. Am.* i. 21. t. 9.—Am. bor. occ.
oryzetorum, Bunge, Enum. Pl. Chin. Bor. 2 = sceleratus.
ovalifolius, Lamb. ex G. Don, Gen. Syst. i. 33 = rhomboideus.
ovalis, *Rafin. in Desv. Journ. Bot.* iv. (1814) 268. —Am. bor.

RANUNCULUS:—
ovatifolius, Steud. Nom. ed. II. i. 434 = praec.
ovatus, DC. Syst. i. 247, in syn. = R. Flammula.
oxynotus, *A. Gray, in Proc. Am. Acad.* x. (1874) 68.—Calif.
oxyrhynchus, Griseb. Spicil. Fl. Rumel. i. 312 = Sprunerianus.
oxyspermus, *Willd. Sp. Pl.* ii. 1328.—Ross.; As. Min.
Pacheri, *Dalla Torre, Atl. Alpenfl.* 53.—Europ.
pachyrrhizus, *Hook. f. Handb. N. Zeal. Fl.* 8.—N. Zel.
palaestinus, Boiss. Diagn. Ser. I. viii. 4 = constantinopolitanus.
palensis, Berger. Fl. Bass. Pyr. ii. 405 = sardous.
palimbifolius, *Wedd. Chlor. And.* ii. 303.—Ecuador; Peruv.
Pallasii, *Schlecht. Animad. Ran.* i. 15. t. 2.—Reg. arct.
pallidior, [Chaix, in] Vill. Hist. Pl. Dauph. i. 335; iii. 751 = sardous.
pallidus, [Soland. in] Russ. Aleppo, ed. II. ii. 254 = sardous.
palmatus, [Clairv.] Man. Herb. 177 = aconitifolius.
palmatus, *Elt. Sketch*, ii. 61.—Am. bor.
paludosus, *Poir. Voy. Barb.* ii. 184.—Afr. bor.
paludosus, Willk. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 924 = chaerophyllos.
palustris, Amo, Fl. Iber. vi. 720 = Broteri.
palustris, Hegetschw. ex Kittel, Taschenb. ed. II. 773 = auricomus.
palustris, Lange, ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 930 = adscendens.
palustris, *Linn. ex Sm. in Rees, Cyclop.* xxix. n. 52.—Reg. Mediterr.; Oriens.
palustris, Willk. ex Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 931 = Aleae.
pangiensis, *Watt, in Journ. Linn. Soc.* xviii. (1881) 377.—Reg. Himal.
panormitanus, Tod. in Atti Acc. Sc. Palermo, N. S. i. (1845) 15 = bulbosus.
pantothrix, Brot. Fl. Lusit. ii. 375 = trichophyllus.
pantothrix, Ell. Sketch, ii. 56 = aquatilis.
papillatus, Dulac, Fl. Hautes-Pyr. 218 = hederaceus.
paradoxus, Weihe, in Linnaea, x. (1835–36) 418. t. 2 = acris.
Pardalianches, Steud. Nom. ed. I. 680 = R. Thora.
parnassifolius, *Linn. Sp. Pl.* 549.—Europ.
parviflorus, Gouan, Fl. Monsp. 270 = sardous.
parviflorus, *Linn. Syst.* ed. X. 1087.—Reg. Mediterr.
parviflorus, Steud. Nom. ed. II. ii. 434 = parnassifolius.
parvulus, [Clairv.] Man. Herb. 179 = Villarsii.
parvulus, J. F. Gmel. Syst. 879 = falcatus.
parvulus, Linn. Mant. i. 79 = sardous.
pascuicolus, Jord. Diagn. i. 73 = acris.
patagonicus, *Poepp. Fragm. Syn. Pl. Chil.* 29.—Chili.
paucicarpus, Arv. Ess. Pl. Dauph. 17 = ololeucos.
paucidentatus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 309.—Soongar.
pauciflorus, Spreng. Syst. ii. 657 = parviflorus.
paucistamineus, Tausch, in Flora, xvii. (1834) 525 = trichophyllus.
pectinatus, Dubois, Méthod. 454 = aquatilis.
pedatifidus, Hook. Fl. Bor. Am. i. 18 = Hookeri.
pedatifidus, Schlecht. Animad. Ran. ii. 18 = affinis.
pedatifidus, *Sm. in Rees, Cyclop.* xxix. n. 72.—Dahuria.
pedatus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 112. t. 108. —Hungar.
pedunculatus, *Sm. in Rees, Cyclop.* xxix. n. 49.—Reg. Magellan.
pedunculatus, Viv. ex Nym. Consp. 8 = chaerophyllos.
pedunculatus, *Lange, Desc. Pl. Nov. Hisp.* i. 4.—Hispan.
pedunculatus, Viv. Fl. Cors. Sp. Nov. 8 = sardous.
peloponnesiacus, Boiss. Diagn. Ser. I. i. 63 = chaerophyllos.
peltatus, Moench, Meth. 214 = aquatilis.
peltatus, *Schrank, Baier. Fl.* ii. 103.—Reg. temp. bor.
pensylvanicus, *Linn. f. Suppl.* 272.—Am. bor.
pensylvanicus, Pursh, Fl. Am. Sept. ii. 393 = repens.
persicus, *DC. Syst.* i. 288.—Persia.
Perusianus, Jeanb. & Timb. in Bull. Soc. Phys. Nat. Toulouse, iii. (1875–76) 330 = nemorosus.
peruvianus, *Pers. Syn.* ii. 103.—Peruv.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* v. 45.—Mexic.

RANUNCULUS:—
Petiveri, Amo, Fl. Iber. vi. 702 = Baudotii.
Petiveri, Coss. & Germ. Fl. Par. i. 10 = ololeucos.
Petiveri, Koch, in Sturm, Deutschl. Fl. Heft 82; et Syn. Fl. Germ. ed. II. 13 = confusus.
petroselinus, *Biria, Hist. Nat. Renoncul.* 43. t. 2.—Ins. Borbon.
peucedanifolius, *All. Fl. Pedem.* ii. 53.—Reg. bor. temp.
peucedanifolius, Gilib. Exercit. i. 369 = fluitans.
peucedanoides, Desf. Fl. Atlant. i. 444 = peucedanifolius.
Peyrousianus, Freyn, in Willk. & Lange, Prod. Fl. Hisp. iii. 936 = tuberosus.
philonotis, Ehrh. in Hannov. Mag. (1783) 17tes St. 270 = sardous.
philonotis, Pursh, Fl. Am. Sept. ii. 393 = repens.
phrygius, *Boiss. Fl. Orient.* i. 29.—Phrygia.
Phthora, Steud. Nom. ed. II. ii. 435 = *R. Pthora*.
phyteumaefolius, Spreng. Syst. iv. Cur. Post. 218 = bonariensis.
pilosus, Dulac, Fl. Hautes-Pyr. 220 = Gouani.
pilosus, *H. B. & K. Nov. Gen. et Sp.* v. 45.—Am. austr.
pilosus, C. Muell. in Walp. Ann. vii. 21, sphalm. = setoso-pilosus.
pilulifer, Hook. Ic. Pl. t. 600 = parviflorus.
pimpinellifolius, Hook. Journ. Bot. i. (1834) 243 = lappaceus.
pimpinelloides, D. Don, in Royle, Illustr. Bot. Himal. 53 = Callianthemum cachemirianum.
Pinardi, *Boiss. Diagn.* Ser. II. v. 10.—Syria.
pinguis, *Hook. f. Fl. Antarct.* 3. t. 1.—Ins. Auckland.
pinnatus, *Poir. Encyc.* vi. 126.—Afr. trop.
pinnatus, Wight, Illustr. i. 6 = Wallichianus.
plantagineus, All. Fl. Pedem. ii. 48 = pyrenaeus.
plantagineus, Pers. Syn. ii. 102 = bupleuroides.
plantaginifolius, *Murr. in Comm. Gotting.* viii. (1777) 39. t. 2.—Sibir.
plantaginifolius, Salisb. Prod. 372 = amplexicaulis.
platanifolius, Linn. Mant. i. 79 = aconitifolius.
platensis, *Spreng. Syst.* v. 586 (Index).—Reg. Argent.
platyspermus, *Fisch. Hort. Gorenk.* ed. II. (1812) 48.—Reg. Khirgis.; Turkestan.
plebeius, *R. Br. ex DC. Syst.* i. 288.—Austral.; N. Zel.
pleurocarpus, *Maxim. Prim. Fl. Amur.* 21.—Reg. Amur.
polianthos, Neck. Delic. Gallo-Belg. i. 240 = polyanthemos.
polyanthemoides, Bor. Fl. Centr. Fr. ed. III. ii. 16 = nemorosus.
polyanthemoides, Schur, Enum. Pl. Transs. 20 = polyanthemos.
polyanthemos, *Linn. Sp. Pl.* 554.—Europ.; Reg. Caucas.
polymorphus, All. Fl. Pedem. ii. 49 = auricomus.
polymorphus, Rochel, Pl. Banat. Rar. in Index, nomen; Bot. Reise Banat. 74 = montanus.
polypetalus, Gill. ex Hook. Bot. Misc. iii. (1833) 134 = peduncularis.
polypetalus, Rafin. Fl. Ludov. 83 (Quid?).—Am. bor.
polypetalus, Royle, Illustr. Bot. Himal. 54. t. 11. f. 2 = Oxygraphis polypetala.
polyphyllus, *Waldst. & Kit. ex Willd. Sp. Pl.* ii. 1331. —Hungar.
polyrhizos, *Steph. ex Willd. Sp. Pl.* ii. 1324.—Ross.; Sibir.
polyrhizus, Soleir. ex Nym. Consp. 10 = demissus.
ponticus, C. Koch, in Linnaea, xix. (1847) 46 = Sibthorpii.
ponticus, Willd. Enum. Hort. Berol. Suppl. 40 = nemorosus.
pozoaefolius, *C. Gay, Fl. Chil.* i. 34.—Chili.
praecox, Salisb. Prod. 372 = R. Ficaria.
praemorsus, *H. B. & K. ex DC. Syst.* i. 292.—Am. austr.
pratensis, Phil. in Linnaea, xxxiii. (1864–65) 2 = erodiifolius.
pratensis, Presl, Delic. Prag. 9 = palustris.
Preslii, *Weihe, ex Opiz, Natural.* xi. (1826) 346.—Europ.
procerus, Moris, Fl. Sard. i. 45 = palustris.
procumbens, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 355.—Persia.
prolifer, *Reinw. ex De Vriese, in Miq. Pl. Jungh.* 77. —Java.
propinquus, C. A. Mey. in Ledeb. Fl. Alt. ii. 332 = acris.

RANUNCULUS:—
prostratus, Poir. Encyc. vi. 113 = repens.
protensus, *Kit. in Linnaea*, xxxii. (1863) 484.—Hungar.
protractus, *Steud. Nom.* ed. II. ii. 435; *et in Flora*, xxxix. (1856) 406.—Chili.
Pseudo-bulbosus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 84 = sardous.
Pseudo-cassubicus, Schur, Enum. Pl. Transs. 23 = auricomus.
Pseudo-fluitans, Newbould, ex Syme, Engl. Bot. ed. III. i. 20 = aquatilis.
Pseudo-hirsutus, Schur, Enum. Pl. Transs. 22 = sardous.
Pseudo-montanus, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 42 = montanus.
Pseudo-philonotis, *Griseb. in Goett. Abh.* xix. (1874) 69.—Reg. Argent.
Pseudopsis, Jord. Diagn. i. 68 = demissus.
Pseudo-thora, Host, Fl. Austr. ii. 108 = hybridus.
Pseudo-Villarsii, Schur, Enum. Pl. Transs. 19 = Villarsii.
psilostachys, *Griseb. Spicil. Fl. Rumel.* i. 304.—Macedon.
psychrophilus, *Wedd. Chlor. And.* ii. 300.—Bolivia.
Pthora, Crantz, Stirp. Austr. ed. I. fasc. ii. 98; ed. II. 119 = hybridus.
pubescens, Bert. ex Steud. Nom. ed. II. ii. 435 = protractus.
pubescens, Lag. Gen. et Sp. Nov. 19 = Lagascanus.
pubescens, *Thunb. Prod. Pl. Cap.* 94.—Afr. austr.
pulchellus, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 333.—Sibir.; Reg. Himal.
Pumilio, R. Br. ex DC. Syst. i. 271 = parviflorus.
pumilus, Poir. Encyc. vi. 133 = divaricatus.
pumilus, Thuill. Fl. Par. ed. II. i. 277 = sardous.
Purshii, Richards. in Frankl. 1st Journ. ed. II. App. 23 = multifidus.
pusillus, Ledeb. in Mém. Acad. Pétersb. v. (1812) 546 = multifidus.
pusillus, *Poir. Encyc.* vi. 99.—Am. bor.
pusillus, Pomel, Nouv. Mat. Fl. Atl. 249 = batrachioides.
pygmaeus, Gueldenst. ex Ledeb. Fl. Ross. i. 27 = falcatus.
pygmaeus, *Wahlenb. Fl. Lapp.* 157. t. 8. f. 1.—Reg. bor. et arct.
pyrenaeus, Gouan, Illustr. 33 = Gouani.
pyrenaeus, *Linn. Mant.* ii. 248.—Europ. austr.
Questieri, Billot, Annot. Fl. Fr. et Allem. 122 = nemorosus.
Raddeanus, *Regel, Ind. Sem. Hort. Petrop.* (1865) 39.—Reg. Caucas.
radians, Revel, in Act. Linn. Soc. Bord. xix. (1853) 120 = trichophyllus.
radicans, C. A. Mey. in Ledeb. Fl. Alt. ii. 316 = multifidus.
radicescens, Jord. Pugill. Pl. Nov. 2 = nemorosus.
Raimondii, *Wedd. Chlor. And.* ii. 305.—Peruv.
rectirostris, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 303.—Alger.
rectus, Bor. Fl. Centr. Fr. ed. III. ii. 15 = acris.
rectus, Déségl.—Cf. Bull. Soc. Bot. Fr. xxv. (1878) Bibl. 107 = praec.?
rectus, Ravin, in Bull. Soc. Sc. L'Yonne, xiv. II. (1860) 92 (Quid?).—Gallia.
recurvatus, Bong. Veg. Sitch. 5 = Nelsonii.
recurvatus, *Poir. Encyc.* vi. 125.—As. et Am. bor.
renifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Thora.
reniformis, Gilib. Fl. Lituan. ii. 269 = cassubicus.
reniformis, Kittel, Taschenb. ed. II. 773 = auricomus.
reniformis, *Wall. Cat.* n. 4709.—Ind. or.
repens, *Linn. Sp. Pl.* 554.—Reg. bor. temp.
reptabundus, Jord. Diagn. i. 83 = repens.
reptabundus, Rupr. in Beitr. Pfl. Russ. Reich. ii. 10 = sceleratus?
reptans, Linn. Sp. Pl. 549 = R. Flammula.
reticulatus, Regel & Schmitz, ex Seub. in Flora, xxviii. (1845) 334 = arvensis.
Reuterianus, *Boiss. Diagn.* Ser. I. i. 65.—As. Min.
Revellierii, Bor. in Mém. Soc. Acad. Maine-et-Loire, i. (1857) 85; iv. (1858) 117 = pedunculatus.
rhaeadifolius, Poir. Encyc. Suppl. iv. 664 = trilobus?
rhipiphyllus, Bast. ex Bor. Fl. Centr. Fr. ed. III. ii. 11 = aquatilis.
rhoeadifolius, DC. Syst. i. 284 = trilobus.
rhombifolius, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 1 = bullatus.

RANUNCULUS:—
rhomboideus, Goldie, in Edinb. Phil. Journ. vi. (April 1822) 329. t. 11. f. 1 = ovalis.
rhynchocarpus, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 355 = lomatocarpus.
rigescens, *Turcz. ex Osten-Sack. & Rupr. Sert. Tiansch.* 37.—Turkestan.
rigidus, *Godr. in Mém. Soc. Roy. Nancy*, 1839 (1840) 41.—Afr. austr.
rigidus, Pers. in Usteri, Ann. Bot. xiv. (1795) 39 = divaricatus.
Rionii, *Lagger, in Flora*, xxxi. (1848) 49.—Helvet.
riparius, Edgew. in Trans. Linn. Soc. xx. (1846) 28 = pensylvanicus.
rivularis, *Banks & Soland. ex DC. Syst.* i. 270.—Austral.; N. Zel.
Robertsoni, *Benth. Fl. Austral.* i. 10.—Austral.
Robini, Rafin. Fl. Ludov. 82 = obtusiusculus.
robustus, Pomel, Nouv. Mat. Fl. Atl. 250 = chaerophyllos.
Rosani, Tenore, Prod. Fl. Neap. p. xxxii = trilobus.
roseus, Hegetschw. Fl. Schw. 540 = glacialis.
rotundifolius, Jord. Diagn. i. 66 = monspeliacus.
rufosepalus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 217.—Turkestan.
rufulus, Brot. Fl. Lusit. ii. 367 = chaerophyllos.
rumelicus, *Griseb. Spicil. Fl. Rumel.* i. 305.—Graec.
rupestris, *Guss. Ind. Sem. Boccad.* (1826) 8.—*Cf. Willk. & Lange, Prod. Fl. Hisp.* iii. 981.—Hispan.
rutaefolius, *Linn. Sp. Pl.* 552.—Europ.; Sibir.
rutaefolius, Mill. Gard. Dict. ed. VIII. n. 6 = millefoliatus.
Rutenbergii, *Freyn, in Bremen, Abh.* vii. (1880) 9.—Madag.
ruthenicus, Jacq. Hort. Vindob. iii. 19 = plantaginifolius.
Sabinii, R. Br. in Parry, 1st Voy. App. 264 = pygmaeus.
sagittifolius, *Hook. Ic. Pl.* t. 173.—Zeylan.
sahendicus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 6.—Persia.
salicifolius, Stokes, Bot. Mat. Med. iii. 259 = R. Lingua.
salsugineus, Link, Handb. ii. 392, sphalm. = *salsuginosus*, Pall.
salsuginosus, DC. Prod. i. 33 = plantaginifolius.
salsuginosus, D. Don, in Royle, Illustr. Bot. Himal. 53 = lobatus.
salsuginosus, *Georgi, Bemerk. Russ. Reich.* i. 222 (Quid?).—Rossia.
salsuginosus, Pall. Reise, iii. 213 = R. Cymbalaria.
salsuginosus, Wall. Cat. n. 4708 = pulchellus.
salsus, *Pall. Reise Prov. Russ. Reich.* iii. 318.—Sibir. or.
samojedorum, *Rupr. Beitr. Pfl. Russ. Reich.* ii. 18.—Ross. arct.
sanguineus, Mill. Gard. Dict. ed. VIII. n. 10 = asiaticus.
saniculaefolius, Schur, Enum. Pl. Transs. 21 = nemorosus.
saniculaefolius, *Viv. Fl. Lib. Spec.* 29. t. 11. f. 2.—Afr. bor.
saniculaeformis, Muhl. ex DC. Prod. i. 40 = recurvatus.
sardonicus, Gueldenst. ex Ledeb. Fl. Ross. i. 47 = ophioglossifolius.
sardous, *Crantz, Stirp. Austr.* ed. I. fasc. ii. 84; ed. II. 111.—Europ.; Afr. bor.; As. Min.
sarmentoso-repens, *Steud. in Flora*, xxxix. (1856) 403.—Peruv.
sarmentosus, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 244 = R. Cymbalaria.
sarmentosus, *Griseb. in Goett. Abh.* xxiv. (1879) 13.—Reg. Argent.
Sartorianus, Boiss. & Heldr. Diagn. Ser. II. i. 8 = Villarsii.
saxatilis, Balb. Misc. Bot. i. 27 = monspeliacus.
saxatilis, Vill. Prosp. 50 = praec.?
saxifragaefolius, Steph. ex Steud. Nom. ed. II. ii. 435 = R. Cymbalaria.
saxifragaefolius, *Steud. Nom.* ed. II. ii. 435; *et in Flora*, xxxix. (1856) 406.—Chili.
scaber, *Presl, Fl. Sicul.* i. 13.—Sicilia.
scandens, *Bogusl. in Erman, Archiv*, vi. (1848) 60.—Ross.
scapigerus, Hook. Journ. Bot. i. (1834) 244 = lappaceus.
sceleratus, *Linn. Sp. Pl.* 551.—Reg. bor. temp.

RANUNCULUS:—
Schimperianus, Hochst. & A. Rich. Tent. Fl. Abyss. i. 4 = simensis.
Schlechtendalii, Hook. Fl. Bor. Am. i. 21. t. 11 = repens.
Schmidtii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 34, in obs. = R. Lingua.
Schottii, *Dalla Torre, Atl. Alpenfl.* 54.—Europ.
Schowii, *Tin. in Guss. Fl. Sic. Syn.* ii. 889.—Sicil.
Schraderianus, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 44 = chius.
Schrenkianus, *Fisch. & Mey. ex Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) I. 64.—Soongar.
Schurii, Fuss, ex Schur, Enum. Pl. Transs. 16 = montanus.
scutatus, Schott, ex Dalla Torre, Atl. Alpenfl. 54 = Schottii.
scutatus, Waldst. & Kit. Pl. Rar. Hung. ii. 205 = R. Thora.
sedunensis, Rion, ex Wolf. in Bull. Soc. Murith. vii–viii. (1879) 28 = Rionii.
segetalis, Kit. ex Uechtr. in Flora, iv. (1821) 40 = arvensis.
Seguieri, *Vill. Prosp.* 50; *Hist. Pl. Dauph.* iii. 737.—Europ.
semicalvus, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 2, in textu = bulbatus.
semidecurrens, Opiz, in Flora, vii. (1824) I. Beil. 84 = repens.
semiverticillatus, *Phil. in Anal. Univ. Chil.* i. (1861) 60.—Chili.
septentrionalis, *De Bray, ex Fleisch. & Lindem. Fl. Deutsch. Ostseeprov.* 195.—Prov. Balt.
septentrionalis, Poir. Encyc. vi. 125 = repens?
serbicus, *Vis. Cat. Sem. Hort. Patav. Suppl.* (1858); *et in Mem. Ist. Venet.* ix. (1860) 168.—Banat.
sericans, *Steud. in Flora*, xxxix. (1856) 405.—Chili.
sericeus, *Poir. Encyc.* vi. 109.—Ins. Maurit.
sericeus, Willd. Enum. Hort. Berol. 589 = illyricus.
sericocephalus, *Hook. f. Fl. Antarct.* 225.—Ins. Falkland.
sericophyllus, *Hook. f. Handb. N. Zeal. Fl.* 6.—N. Zel.
sessiliflorus, R. Br. ex DC. Syst. i. 302 = parviflorus.
sessiliflorus, Spreng. Syst. iv. Cur. Post. 219 = platensis.
sessiliflorus, Wedd. ex Griseb. in Goett. Abh. xxiv. (1879) 13 = Hilarii.
setoso-pilosus, *Steud. in Flora*, xxxix. (1856) 404.—Peruv.
Sewerzowi, *Regel, in Act. Hort. Petrop.* v. (1877) 221.—Turkestan.
sibbaldioides, *H. B. & K. ex DC. Syst.* i. 293.—Am. austr.
sibiricus, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 246 = nivalis.
sibiricus, Spreng. Syst. ii. 652 = multifidus.
sibiricus, Schraub, ex Schlecht. Animad. Ran. ii. 38, nomen (Quid?).—Hab.?
Sibthorpii, *Boiss. Diagn.* Ser. II. i. 7.—Graec.
siculus, Presl, Delic. Prag. 8 = acris.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 5.—Japon.
silvaticus, Vill. Prosp. 51 = acris?
simensis, *Fresen, in Mus. Senckenb.* ii. (1837) 269.—Abyss.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 6.—N. Zel.
songaricus, *Schrenk. in Fisch. & Mey. Enum. Pl. Nov. Schrenk.* ii. 67.—Soongar; Turkest.
sparsipilus, Jord. Diagn. i. 80 = bulbosus.
speciosus, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 722 = bulbosus.
sphaerospermus, Boiss. & Blanche, Diagn. Ser. II. v. 6 = aquatilis.
spicatus, *Desf. Fl. Atlant.* i. 438.—Afr. bor.
spicatus, Plan. Fl. Gallega, 54 = nigrescens.
spicatus, Presl, Fl. Sicul. 14 = rupestris.
Spreitzenhoferi, *Heldr. in Oestr. Bot. Zeitschr.* xxviii. (1878) 50.—Ins. Ionian.
spretus, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 17 = nemorosus.
Sprunerianus, *Boiss. Diagn.* Ser. I. i. 64 (Sprunnerianus).—Graec.; Syria.
squamosus, Dulac, Fl. Hautes-Pyr. 218 = R. Thora.
squarrosus, Laest. ex Nym. Consp. 12 = acris?
squarrosus, Pomel, Nouv. Mat. Fl. Atl. 381 = orientalis.

RANUNCULUS:—
stagnalis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 5 = simensis.
stagnatilis, Wallr. Sched. Crit. 285 = aquaticus.
stenocarpus *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 6. —Abyss.
stenopetalus, *Hook. Ic. Pl.* t. 677.—Chili.
Steveni, Andrz. ex Bess. Enum. Pl. Volh. 22 = acris.
stipatus, Jord. Diagn. i. 72 = acris.
stoloniferus, *Hemsl. Diagn. Pl. Nov.* 17.—Mexic.
striatus, Hochst. in Flora, xxiv. (1841) I. Intell. 28, nomen; et ex A. Rich. Tent. Fl. Abyss. i. 8 = pinnatus.
strigillosus, *Boiss. et Huet, Diagn.* Ser. II. v. 7.—As. Min.; Persia.
strigulosus, Schur, Enum. Pl. Transs. 17 = acris.
suaneticus, *Rupr. Fl. Cauc.* 21.—Reg. Caucas.
subacri-bulbosus, *Wesm. in Bull. Acad. Brux.* (1862) 14. t. 298.—Armenia.
subalpinus, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 40, in textu = acris.
subapetalus, Vict. Aug. in Ann. Linn. Soc. i. 193; ex Duby, Bot. Gall. i. 13 = parviflorus.
submersus, Godr. ex Carion, in Bull. Soc. Bot. Fr. vi. (1859) 498 = aquatilis.
suborbiculatus, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 921.—Hispan.
subpinnatus, Wight & Arn. Prod. 4 = diffusus.
subscaposus, *Hook. f. Fl. Antarct.* 5.—N. Zel.
subtilis, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 41 = acris.
subtilis, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 393. —Reg. Cauc.
subtrifolius, Schur, Enum Pl. Transs. 22 = sardous.
subtuberosus, Gilib. Fl. Lituan. ii. 262 = acris.
sulphureus, [Soland. in] Phipps, Voy. 202 = nivalis.
supranudus, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 1 = bullatus.
sylvaticus, Gren. et Godr. Fl. Fr. i. 33 = nemorosus.
sylvaticus, Thuill. Fl. Par. ed. II. 276 = acris.
Szovitzianus, *Boiss. Fl. Orient.* i. 42.—Persia.
szurulensis, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 14 = montanus.
Tachireoi, *Franch. & Sav. Enum. Pl. Jap.* ii. 267.—Japon.
Tappeineri, Bamberg. in Flora, xxxv. (1852) 625 = pygmaeus.
tauricus, Pall. Tabl. Taur. 52 = ophioglossifolius.
tauriensis, All. ex Steud. Nom. i. 681 = pyrenaeus.
telonensis, Robert, ex Biria, Hist. Nat. Renoncul. 37 = R. Flammula.
tembensis, *Fresen. in Mus. Senckenb.* ii. (1837) 271.—Abyss.
tenellus, Marschl. in Flora, xvii. (1834) II. Beibl. 84 = demissus.
tenellus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 23 = Nelsonii.
tenellus, *Viv. Pl. Aegypt. Dec.* iv. 6.—Aegypt.
Teneriffa, Pers. Syn. ii. 103 = cortusaefolius.
Tenorii, Jord. Diagn. i. 62 = illyricus.
tenuicaulis, *Cheesem. in Trans. N. Z. Inst.* xvii. (1885) 235.—N. Zel.
tenuifolius, Schleich. Cat. Pl. Helv. ed. IV. 28 = montanus.
tenuirostris, *Pomel, Nouv. Mat. Fl. Atl.* 250.—Alger.
tenuirostris, Steud. ex Hochst. in Flora, xxvii. (1844) 97 = oreophytus.
ternatus, DC. Syst. i. 242 = acris.
ternatus, *Thunb. Fl. Jap.* 241.—China; Japon.
testiculatus, Crantz, Stirp. Austr. ed. I. fasc ii. 97; ed. II. 119; Bieb. Fl. Taur. Cauc. iii. 386 = falcatus.
texensis, Engelm. ex Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1847) 210 = oblongifolius.
Thomasii, Tenore, Ind. Sem. Hort. Neap. (1827) 25 = millefoliatus.
Thomasii, *Tenore, Prod. Fl. Nap.* 69.—Italia.
Thora, *Linn. Sp. Pl.* 550.—Europ. austr.
Thora, Sibth. & Sm. Fl. Graec. Prod. i. 380 = brevifolius.
Thora, Sturm, Deutschl. Fl. Heft 19 = hybridus.
Thora, Weigel. ex Steud. Nom. ed. I 681 = auricomus.
Timbali, Mabille & Gaudf. Not. sur Renonc. (1874).—Cf. Bull. Soc. Bot. Fr. xxi. (1874) Bibl. 216 = nemorosus.
tomentosus, Moench, Meth. 212 = illyricus.
tomentosus, Poir. Encyc. vi. 127 = repens.
Tommasinii, Reichb. ex Freyn, in Oestr. Bot. Zeitschr. xxv. (1875) 116 = velutinus.

RANUNCULUS:—
tomophyllus, Jord. Diagn. i. 71 = acris.
trachycarpus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 46.—Oriens.
trachyspermus, Ell. Sketch, ii. 65 = parviflorus.
trachyspermus, *Engelm. ex Engelm. & Gray, in Rost. Journ. Nat. Hist.* v. (1847) 211.—Am. bor. occ.
trachyticus, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 41 = acris.
transsilvanicus, Schur, l. c. 46 = acris.
Traunfellneri, Hoppe, in Sturm, Deutschl. Fl. Heft 46 = alpestris.
Traunfellneri, Schott, Skizz. Ranunk. Alloph. 8 = praec.?
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 4.—N. Zel.
trichocarpus, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* i. 47.—Armen.; Persia.
trichophyllos, Losc. Ser. Inconf. Pl. Arag. 2 = confusus.
trichophyllus, *Chaix, in Vill. Hist. Pl. Dauph.* i. 335. —Europ.; Oriens.
tricrenatus, *Rupr. in Beitr. Pfl. Russ. Reichs*, ii. 19.—Reg. Samojed.
tridentatus, H. B. & K. ex DC. Syst. i. 252 = R. Cymbalaria.
trifoliatus, Muhl. ex Steud. Nom. ed. II. ii. 436 = fascicularis.
trifolius, Moench, Meth. Suppl. 70 = pensylvanicus.
trigonocarpus, *Boiss. Diagn.* Ser. I. viii. 2.—As. Min.
trilobatus, D. Don, Prod. Fl. Nep. 194 = diffusus.
trilobatus, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 547.—N. Zel.
trilobatus, Kit. ex. Schlecht. Animad. Ran. ii. 21 = montanus.
trilobus, *Desf. Fl. Atlant.* i. 437. t. 113.—Reg. Mediterr.
tripartitus, *DC. Ic. Gall. Rar.* i. 15.—Gall. occ.
tripartitus, Dubourg, ex Steud. Nom. ed. II. ii. 436 = Lenormandi.
tripartitus, Nolte, Novit. Fl. Holsat. 51 = aquatilis.
triphyllos, Wallr. Beitr. Fl. Hercyn. 214; et in Linnaea, xiv. (1840) 584 = confusus.
trisepalus, *Gill. ex Hook. Bot. Misc.* iii. (1833) 133.—Am. austr.
triternatus, Poir. Encyc. Suppl. iv. 662, in syn. = biternatus.
trullifolius, *Hook. f. Fl. Antarct.* 226.—Ins. Falkland.
Tschuktschorum, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 19, in textu.—Terr. Tschuktsch.
tuberculatus, DC. Syst. i. 297 = arvensis.
tuberculatus, C. A. Mey. Verz. Pfl. Cauc. 202 = lomatocarpus.
tuberosus, Hornem. Hort. Hafn. ii. 527 = Hornemanni.
tuberosus, Lapeyr. Hist. Abr. Pl. Pyr. i. 320 = nemorosus.
tuberosus, Schur, in Oesterr. Bot. Zeitschr. xi. (1861) 82 = montanus.
tuberosus, Stur, ex Nym. Consp. 12 = acris.
turkestanicus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 217.—Turkestan.
udus, *Freyn, in Bremen, Abh.* vii. (1880) 10.—Madag.
uliginosus, Willd. Enum. Hort. Berol. 586 = ophioglossifolius.
umbellatus, Roxb. ex Willd. l. c. 588 = sceleratus.
umbrosus, Tenore & Guss. Fl. Neap. Syll. App. v. 15 = lanuginosus.
unalaschcensis, Bess. ex Ledeb. Fl. Ross. i. 32 = R. Flammula.
uncinatus, *D. Don, ex G. Don, Gen. Syst.* i. 35.—Mexic.
vaginalis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 13.—N. Granat.
vaginalis, Portenschl. ex Freyn, Kennt. Ranunc. 26 = palustris.
vaginatus, Sommerauer, in Flora, xvi. (1833) 182 = crenatus.
valdepubens, Jord. Diagn. i. 82 = bulbosus.
valesiacus, Suter, Fl. Helvet. i. 325 = lacerus.
variifolius, Salisb. Prod. 373 = auricomus.
velutinus, Schur, Enum. Pl. Transs. 19 = lanuginosus.
velutinus, *Tenore, Ind. Sem. Hort. Neap.* (1825) 11. n. 23.—Europ. austr.
ventricosus, Vent. Jard. Cels, sub. t. 73 = muricatus.
vernus, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 1018 = auricomus, cassubicus.
Vernyi, *Franch. & Sav. Enum. Pl. Jap.* i. 8.—Japon.
verrucosulus, Poir. ex Steud. Nom. ed. II. ii. 436 = sardous, trilobus.

RANUNCULUS:—
verrucosus, Bercht. & Presl, Rostl. i. Ranunculac. 61, nomen (Quid?).—Europ.
vestitus, *Wall. Cat.* n. 4707.—Burma.
Villarsii, *DC. Fl. Fr.* iv. 896.—Europ. austr; Reg. Caucas.
villiferus, Jord. Diagn. i. 82 = bulbosus.
villosus, DC. Syst. i. 283 = constantinopolitanus.
villosus, St. Amans, Fl. Agen. 227. t. 5 = nemorosus.
villosus, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1877) 306 = palustris.
vitifolius, Royle, Illustr. Bot. Himal. 53 = diffusus.
vulgatus, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 15 = acris.
Wallichianus, *Wight & Arn. Prod.* 4.—Ind. or.
Warionii, Freyn, in Flora, lxiii. (1880) 25 = spicatus.
Weyleri, *Marès, in Bull. Soc. Bot. Fr.* xii. (1865) 232, in textu; *et ex Willk. in Linnaea*, xl. (1876) 131.—Ins. Balear.
Winkleri, *Freyn, in Willk. & Lange, Prod. Fl. Hisp.* iii. 922.—Hispan.
xantholeucos, Coss. & Dur. in Bull. Soc. Bot. Fr. xxvii. (1880) 67 = batrachioides.
Xatardi, Lapeyr. Suppl. Hist. Pl. Pyr. 77 = parviflorus.
yunnanensis, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 5.—China.
Zuccarinii, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 5 = ternatus.

RAOULIA, Hook. f. Fl. N. Zel. i. 134. tt. 36, 37 (1846). *COMPOSITAE*, Benth. & Hook. f. ii. 307.
apice-nigra, *T. Kirk, in Trans. N. Z. Inst.* xi. (1879) 464.—N. Zel.
australis, *Hook. f. ex Raoul, Choix Pl. Nouv. Zél.* 20. t. 15.—N. Zel.
bryoides, *Hook. f. Fl. N. Zel.* ii. 322.—N. Zel.
catipes, *Hook. f. Fl. Tasman.* i. 206.—Austral.
eximia, *Hook. f. Handb. N. Zeal. Fl.* 149.—N. Zel.
glabra, *Hook. f. Fl. N. Zel.* i. 135.—N. Zel.
Goyeni, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 373.—N. Zel.
grandiflora, *Hook. f. Fl. N. Zel.* i. 136. t. 37 A.—N. Zel.
Haastii, *Hook. f. Handb. N. Zeal. Fl.* 148.—N. Zel.
Hectori, *Hook. f. l. c.*—N. Zel.
M'Kayi, *F. Buch. in Trans. N. Z. Inst.* xiv. (1882) 354. t. 34. f. 3.—N. Zel.
mammillaria, *Hook. f. Handb. N. Zeal. Fl.* 150.—N. Zel.
Munroi, *Hook. f. l. c.* 148.—N. Zel.
Parkii, *F. Buch. in Trans. N. Z. Inst.* xiv. (1882) 355.—N. Zel.
Petriensis, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 549.—N. Zel.
Planchoni, *Hook. f. ex Benth. Fl. Austral.* iii. 651.—Tasman.
rubra, *F. Buch. in Trans. N. Z. Inst.* xiv. (1882) 349.—N. Zel.
subsericea, *Hook. f. Fl. N. Zel.* i. 136.—N. Zel.
subulata, *Hook. f. Handb. N. Zeal. Fl.* 149.—N. Zel.
tasmanica, Hook. f. Fl. Tasman. t. 58 = catipes.
tenuicaulis, *Hook. f. Fl. N. Zel.* i. 135.—N. Zel.

RAPA, Tourn. ex Linn. Syst. ed. I (1735) = **Brassica**, Tourn. (Crucif.).
Napus, Mill. Gard. Dict. ed. VIII. n. 3 = B. campestris.
oblonga, Mill. l. c. n. 2 = B. campestris.
rotunda, Mill. l. c. n. 1 = B. oleracea.

RAPANEA, Aubl. Pl. Gui. i. 121. t. 46 (1775) = **Myrsine**, Linn.
guianensis, Aubl. l. c. = M. floribunda.
oblonga, Pohl, ex Miq. in Mart. Fl. Bras. x. 308 = M. floribunda.
oblongifolia, Pohl, l. c. 310 = M. leuconeura.
villosissima, Miq. l. c. 322 (Index) = M. villosissima.

RAPATEA, Aubl. Pl. Gui. i. 305. t. 118 (1775). *RAPATEACEAE*, Benth. & Hook. f. iii. 858.
MNASIUM, Schreb. Gen. 214 (1789).
angustifolia, *Spruce, ex Koern. in Linnaea*, xxxvii. (1871–73) 469.—Venezuela.
arthrophylla, Mart. ex Seub. in Mart. Fl. Bras. iii. 1. 132 = Schoenocephalium arthrophyllum.
flava, Kunth, Enum. Pl. iii. 367 (= *Monotrema flavum*). —Venezuela.

RAPATEA:—
Friderici-Augusti, *Rob. Schomb. Rapat. Frid.-Aug. & Saxo-Frid.* 11. t. 1.—Guian.
gracilis, Poepp. & Endl. ex Endl. Gen. 131 = Cephalostemon gracilis.
gracilis, Seub. in Mart. Fl. Bras. iii. 1. 128 = Cephalostemon Riedelianus.
longipes, *Spruce, ex Koern. in Linnaea*, xxxvii. (1871–73) 472.—Bras.
Martiana, Nees, ex Seub. in Mart. Fl. Bras. iii. 1. 130 = Schoenocephalium Martianum.
paludosa, *Aubl. Pl. Gui.* i. 305. t. 118.—Am. trop.
pandanoides, *Linden & André, Illustr. Hortic.* xx. (1873) 213. t. 153–4.—Bras.
parviflora, Nees, ex Seub. in Mart. Fl. Bras. iii. 1. 132 = Schoenocephalium arthrophyllum.
pycnocephala, *Seub. in Mart. Fl. Bras.* iii. 1. 128.—Bras.
sphaerocephala, Schult. f. Syst. vii. 1149 = Thurnia sphaerocephala.
Spruceana, *Koern. in Linnaea*, xxxvii. (1871–73) 470.—Venezuela.
squarrosa, Kunth, Enum. Pl. iii. 367 = Cephalostemon squarrosus.
squarrosa, Spruce, ex Koern. in Linnaea, xxxvii. (1871–73) 447 = Cephalostemon affinis.
unilateralis, Schult. f. Syst. vii. 1149 = Spathanthus unilateralis.

RAPHANIS, Dod. ex Moench, Meth. 267 (1794) = **Cochlearia**, Tourn. (Crucif.).
magna, Moench, l. c. = C. Armoracia.

RAPHANISTRUM, Tourn. ex Adans. Fam. ii. 424 (1763) = **Raphanus**, Tourn. (Crucif.).
arvense, Wallr. Sched. Crit. 336 = Raphanus Raphanistrum.
fugax, Nym. Consp. 29 = Raphanus Raphanistrum.
Gayanum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 44 = Raphanus sativus.
innocuum, Moench, Meth. 217; Medic. in Usteri, Ann. Bot. viii. (1794) 39 = Raphanus Raphanistrum.
Lampsana, Gaertn. Fruct. ii. 300. t. 143 = Raphanus Raphanistrum.
Landra, Reichb. Fl. Germ. Excurs. 656 = Raphanus Landra.
maritimum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 688 = Raphanus maritimus.
odessanum, Andrz. ex Bess. Enum. Pl. Volh. 82 = Raphanus Raphanistrum.
pterocarpum, Juss. ex DC. Syst. ii. 661 = Enarthrocarpus pterocarpus.
recurvatum, Juss. l. c. = Enarthrocarpus lyratus.
rostratum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 22 = Raphanus rostratus.
segetum, Baumg. Enum. Stirp. Transs. ii. 280 = Raphanus Raphanistrum.
silvestre, Aschers. Fl. Brandenb. 65 = Raphanus Raphanistrum.
vulgare, S. F. Gray, Nat. Arr. Brit. Pl. ii. 687 = Raphanus Raphanistrum.

RAPHANOCARPUS, Hook. f. in Hook. Ic. Pl. t. 1084 (1871). *CUCURBITACEAE*.
Kirkii, *Hook. f. l. c.*—Afr. trop.
Welwitschii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 541.—Afr. trop.

RAPHANOPSIS, Welw. Apont. 547, nomen (1858) (Quid?).

RAPHANUS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 196 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 101.
DONDISIA, Scop. Introd. 316 (1777).
DURANDEA, Delarb. Fl. Auv. ed. II. 365 (1800).
ORMYCARPUS, Neck. Elem. iii. 82 (1790).
RAPHANISTRUM, Tourn. ex Adans. Fam. ii. 424 (1763).
aegyptius, L'Hérit. ex DC. Syst. ii. 661 = Enarthrocarpus pterocarpus.
albiflorus, *Presl, Fl. Sic.* i. 109.—Sicilia.
albus, Boenn. ex Reichb. Fl. Germ. Excurs. 656 = R. Raphanistrum.
amplexicaulis, Viv. Fl. Lib. Spec. 37 = Brassica cyrenaica.
arcuatus, Willd. Sp. Pl. iii. 562 = Chorispora tenella.

RAPHANUS:—
articulatus, Stokes, Bot. Mat. Med. iii. 482 = R. Raphanistrum.
arvense, Wallr. ex Aschers. Fl. Brandenb. 65 = R. Raphanistrum.
Aucheri, *Boiss. Diagn.* Ser. I. viii. 45.—Syria; Persia.
biarticulatus, *Willd. Enum. Hort. Berol. Suppl.* 46.—China.
Blaisii, Godr. & Gren. in Godr. Migr. Vég. ed. II. 37; et in Mém. Acad. Stanislas, Sér. III. [xvii.] 1853 (1854) 361 = Rapistrum Blaisii.
cakiloideus, DC. Syst. ii. 669 = Erucaria Oliverii.
caspicus, Willd. ex Ledeb. Fl. Ross. i. 215 = Goldbachia laevigata.
caudatus, Linn. Mant. i. 95 = sativus.
cheiranthifolius, Willd. Hort. Berol. i. 19 = Brassica Cheiranthos.
chinensis, Mill. Gard. Dict. ed. VIII. n. 5 = sativus.
cornutus, Gilib. Fl. Lituan. ii. 57 = R. Raphanistrum.
Erucaria, J. F. Gmel. Syst. 991 = Brassica adpressa.
Erucastrum, Crantz, ex Steud. Nom. ed. II. ii. 437 = Brassica Erucastrum.
erucoides, Linn. f. Suppl. 299 = Brassica oleracea.
fugax, *Presl, Fl. Sic.* i. 109.—Sicilia.
Gayanus, G. Don, in Sweet, Hort. Brit. ed. III. 44 = sativus.
ibericus, Bieb. Fl. Taur. Cauc. ii. 129 = Chorispora iberica.
infestus, Salisb. Prod. 273 = R. Raphanistrum.
laevigatus, Bieb. Fl. Taur. Cauc. ii. 129 = Goldbachia laevigata.
lanceolatus, Willd. Sp. Pl. iii. 562 = Brassica juncea.
Landra, *Moretti, ex DC. Syst.* ii. 668.—Italia.
longistylus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 65, 133 = R. Raphanistrum.
lyratus, Forsk. Fl. Aegypt. Arab. 119 = Enarthrocarpus lyratus.
maritimus, *Sm. Engl. Bot.* t. 1643.—Europ. occ.
microcarpus, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 750.—Hispan.
niger, Mill. Gard. Dict. ed. VIII. n. 4 = sativus.
odessanus, Spreng. Syst. ii. 855 = R. Raphanistrum.
oleifer, Hort. ex Steud. Nom. ed. II. ii. 437 = sativus.
orbicularis, Mill. Gard. Dict. ed. VIII. n. 3 = sativus.
orientalis, Willd. ex Steud. Nom. ed. II. ii. 437 = Erysimum brachycarpum.
pilosus, Willd. Sp. Pl. iii. 562 = Wormskioldia heterophylla.
pinnatus, Viv. Fl. Lib. Spec. 37. t. 16. f. 4 = Rapistrum bipinnatum.
primuloides, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 413 = Raffenaldia primuloides.
pterocarpus, Pers. Syn. ii. 209 = Enarthrocarpus pterocarpus.
pugioniformis, *Boiss. Diagn.* Ser. I. viii. 46.—Syria.
Radicula, Pers. Syn. ii. 208 = sativus.
Raphanistrum, Cav. Desc. 425 = microcarpus.
Raphanistrum, *Linn. Sp. Pl.* 669.—Europ.
recurvatus, Pers. Syn. ii. 209 = Enarthrocarpus lyratus.
rostratus, *DC. Syst.* ii. 666.—Reg. Casp.
rotundifolius, Bieb. Fl. Taur. Cauc. ii. 130 = Erysimum brachycarpum.
rotundus, Mill. Gard. Dict. ed. VIII. n. 2 = sativus.
sativus, *Linn. Sp. Pl.* 669.—Europ.
segetum, Clavaud, in Act. Linn. Soc. Bord. xxxv. (1881) 288 = R. Raphanistrum.
sibiricus, Linn. Sp. Pl. 669 = Chorispora sibirica.
sinensis, Thunb. ex Pritz. Ic. Ind. ii. 239 = sativus?
strictus, Fisch. ex Bieb. Fl. Taur. Cauc. iii. 452 = Diptychocarpus strictus.
stylosus, Dulac, Fl. Hautes-Pyr. 194 = R. Raphanistrum.
sylvestris, Lam. Fl. Fr. ii. 495 = R. Raphanistrum.
tataricus, Falk, Beitr. ii. 218. t. 13 = Chorispora tenella.
tauricus, Adam, in Weber & Mohr, Beitr. i. 62 = Erysimum brachycarpum.
tauricus, Hoffm. ex Steud. Nom. ed. II. ii. 437 = Sobolewskia lithophila.
tenellus, Pall. Reise, iii. 741 = Chorispora tenella.
turgidus, Pers. Syn. ii. 209 (= *Sinapis turgida*).—Aegypt.

RAPHELINGIA, Dum. Anal. Fam. 60 (1829) = **Ornithogalum**?, Tourn. (Liliac.).

RAPHIA, Beauv. Fl. Owar. i. 75 (1804). *PALMAE*, Benth. & Hook. f. iii. 935.
METROXYLON, Spreng. Gen. 283 (1830).
Gaertneri, *G. Mann & H. Wendl. in Trans. Linn. Soc.* xxiv. (1864) 437.—Afr. trop.
Hookeri, *G. Mann & H. Wendl. l. c.* 438.—Afr. trop.
longiflora, *G. Mann & H. Wendl. l. c.*—Afr. trop.
lyciosa, Comm. ex Kunth, Enum. Pl. iii. 217 = pedunculata.
nicaraguensis, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 52, in obs. = vinifera.
pedunculata, *Beauv. Fl. Owar.* i. 78.—Madag.
polymita, Comm. ex Kunth, Enum. Pl. iii. 217 = pedunculata.
Ruffia, Mart. Hist. Nat. Palm. iii. 217 = pedunculata.
taedigera, Mart. l. c. 216 = vinifera.
textilis, *Welw. Apont.* 584. n. 2.—Afr. trop.
vinifera, *Beauv. Fl. Owar.* i. 77.—Afr. trop.
Welwitschii, *H. Wendl. in Trans. Linn. Soc.* xxiv. (1864) 439. t. 42.—Afr. trop.

RAPHIDIOCYSTIS, Hook. f. in Benth. & Hook. f. Gen. i. 828. *CUCURBITACEAE*, Benth. & Hook. f. l. c.
brachypoda, *Baker, in Journ. Bot.* xx. (1882) 113.—Madag.
Mannii, *Hook. f. in Benth. & Hook. f. Gen.* i. 828.—Afr. trop.
Welwitschii, *Hook. f. l. c.*—Afr. trop.

RAPHIDOPHORA, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 168 = RHAPHIDOPHORA, Schott.

RAPHIDOPHYLLUM, Hochst. in Flora, xxiv. (1841) 666 = **Sopubia**, Buch.-Ham. (Scrophul.).
ramosum, Hochst. l. c. 668 = S. ramosa.
simplex, Hochst. l. c. 667 = S. Dregeana.

RAPHIDOSPORA, Reichb. Handb. 190 (1837) = RHAPHIDOSPORA, Nees = **Justicia**, Linn. (Acanth.).

RAPHIODON, Benth. in DC. Prod. xii. 96 (1848) = RHAPHIODON, Schau. = **Hyptis**, Jacq. (Labiat.).

RAPHIOLEPIS, Lindl. in Bot. Reg. t. 468 (1820). *ROSACEAE*, Benth. & Hook. f. i. 627.
OPA, Lour. Fl. Cochinch. 308 (1790), partim.
RHAPIOLEPIS, Poir. in Dict. Sc. Nat. xv. 314 (1827).
crataegoides, M. Roem. Syn. Rosifl. 113 = indica.
indica, *Lindl. in Bot. Reg.* t. 468.—China.
integerrima, Hook. & Arn. Bot. Beech. Voy. 263 = japonica.
japonica, *Sieb. & Zucc. Fl. Jap.* i. 162.—Japon.
laevis, Lodd. ex G. Don, Gen. Syst. ii. 602 = Photinia laevis.
latifolia, *Lodd. ex G. Don, in Loud. Hort. Brit.* 202, nomen (Quid?).—China.
Loureiri, Spreng. Syst. ii. 508 = indica?
Mertensii, Sieb. & Zucc. Fl. Jap. i. 164 = japonica.
ovata, *Briot, in Rev. Hortic.* (1870–71) 348.—Japon.
Pheostemon, Lindl. Coll. Bot. sub t. 3 = indica.
pheostemonia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = praec.
rubra, Lindl. Coll. Bot. t. 3 = indica.
salicifolia, Lindl. l. c. sub t. 3 = indica.
Sieboldii, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 146 = japonica.
sinensis, M. Roem. Syn. Rosifl. 114 = indica.
spiralis, G. Don, Gen. Syst. ii. 602 = indica.

RAPHIONACME, Harv. in Hook. Lond. Journ. Bot. i. (1842) 22. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 745.
APOXYANTHERA, Hochst. in Flora, xxv. (1842) 18.
RHAPHIONACME, Walp. Rep. vi. 480 (1847).
ZUCCHELLIA, Decne. in DC. Prod. viii. 492 (1844).
divaricata, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 23.—Afr. austr.
obovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 250.—Afr. austr.
pubescens, *Hochst. in Flora*, xxvii. (1844) 827.—Afr. austr.
purpurea, *Harv. Thes. Cap.* i. 41. t. 66.—Afr. austr.
Zeyheri, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 23.—Afr. austr.

RAPHIONE, Salisb. Gen. Pl. Fragm. 89 (1866) = **Allium**, Tourn. (Liliac.).
carinata, Salisb. l. c. = A. carinatum.
oleracea, Salisb. l. c. = A. oleraceum.
pallens, Salisb. l. c. = A. paniculatum.
paniculata, Salisb. l. c. = A. paniculatum.

RAPHIOSTYLES, Benth. & Hook. f. Gen. i. 351 (1862) = RHAPHIOSTYLIS, Planch. = **Apodytes**, E. Mey. (Olacin.).

RAPHIS, Beauv. Agrost. 120 (1812) = RHAPHIS, Lour. = **Chrysopogon**, Trin. (Gramin.).

RAPHISANTHE, Lilja, in Linnaea, xv. (1841) 263 = **Blumenbachia**, Schreb. (Loasac.).
lateritia, Lilja, l. c. = Loasa lateritia?

RAPHISTEMMA, Wall. Pl. As. Rar. ii. 50. t. 163 (1831). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 758.
ciliatum, Hook. f. Bot. Mag. t. 5704 = Daemia extensa.
Hooperianum, *Decne. in DC. Prod.* viii. 516.—Siam; Java.
pulchellum, *Wall. Pl. As. Rar.* ii. 50. t. 163.—Reg. Himal.; Burma.

RAPINIA, Lour. Fl. Cochinch. 127 (1790) = **Sphenoclea**, Gaertn. (Campan.).
herbacea, Lour. l. c. = S. zeylanica.

RAPINIA, Montr. in Mém. Acad. Lyon, x. (1860) 243. *VERBENACEAE*, Benth. & Hook. f. ii. 1152.
collina, *Montr. l. c.*—N. Caled.

RAPISTRELLA, Pomel, Mat. Fl. Atl. 11 (1860) = **Rapistrum**, Tourn. (Crucif.).
ramosissima, Pomel, l. c.—Afr. bor.

RAPISTRUM, Tourn. ex Medic. Phil. Bot. i. 190 (1789). *CRUCIFERAE*, Benth. & Hook. f. i. 99.
ARTHROLOBUS, Andrz. ex DC. Syst. ii. 430 (1821).
CORDYLOCARPUS, Desf. Fl. Atlant. ii. 79. t. 152 (1800).
DIDESMUS, Desv. Journ. Bot. iii. (1814) 160. t. 25.
OTOCARPUS, Dur. in Duch. Rev. Bot. ii. (1847) 435.
RAPISTRELLA, Pomel, Mat. Fl. Atl. 11 (1860).
SCHRANKIA, Medic. Pflanzeng. 42 (1792).
SPHAERULARIA, Heist. Syst. 9 (1748).
aegyptiacum, R. Br. in Ait. Hort. Kew. ed. II. iv. 74 = Ochthodium aegyptiacum.
aegyptium, *Baill. Hist. Pl.* iii. 197; *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 46.—Afr. bor.
arvense, All. Fl. Pedem. i. 258 = Raphanus Raphanistrum.
bipinnatum, *Coss. & Kral. in Bull. Soc. Bot. Fr.* iv. (1857) 395.—Alger.
Blaisii, *Gren. in Mém. Soc. Emul. Doubs*, Sér. III. ii. (1858) 404.—Hab. ?
bursaefolium, Berger. Phyt. iii. 165; ex DC. Syst. ii. 649 = Calepina Corvini.
Cakile, Berger. l. c. 163; ex DC. l. c. 429 = Cakile maritima.
clavatum, DC. l. c. 433 = rugosum.
confusum, Pomel, Nouv. Mat. Fl. Atl. 356 = rugosum.
conoideum, *Pomel, l. c.* 356.—Alger.
Cordylocarpus, Pomel, l. c. = Moricaria Cordylocarpus.
costatum, *DC. Syst.* ii. 434.—Syria.
diffusum, Crantz, Class. Crucif. 105 = perenne.
filiforme, Moench, Meth. Suppl. 69 = Crambe filiformis.
glabrum, *Host, Fl. Austr.* ii. 220.—Austr.
glandulosum, Berger. Phyt. iii. 163; ex DC. Syst. ii. 672 = Bunias orientalis.
hirsutum, Host, Fl. Austr. ii. 220 = rugosum.
hirtum, Host, l. c. = rugosum.
hispanicum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 6 = Linnaeanum.
hispanicum, Medic. Pflanzeng. 19 = Crambe hispanica.
hispidum, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 416.—Hab. ?
hypogaeum, Duby, Bot. Gall. ed. II. i. 54 = Morisia hypogaea.
Linnaeanum, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 5.—Europ. austr.
maritimum, Berger. Phyt. iii. 173; ex DC. Syst. ii. 428 = Cakile maritima.

RAPISTRUM:—
microcarpum, Jord. ex Loret, in Bull. Soc. Bot. Fr. vi. (1859) 93 = Linnaeanum.
orientale, *DC. Syst.* ii. 443.—Graecia.
paniculatum, Gaertn. Fruct. ii. 285 = Neslia paniculata.
perenne, *All. Fl. Pedem.* i. 258.—Europ.
perfoliatum, Berger. Phyt. iii. t. 167; ex DC. Syst. ii. 575 = Myagrum perfoliatum.
Raphanistrum, Crantz, Class. Crucif. 107 = Raphanus Raphanistrum.
rugosum, *All. Fl. Pedem.* i. 257.—Europ.; Oriens.
scabrum, Moench, Meth. 211 = Crambe hispanica.
strictissimum, Pomel, Nouv. Mat. Fl. Atl. 356 = rugosum.
venosum, Nym. Consp. 29 = Linnaeanum.
virgatum, *Pomel, Nouv. Mat. Fl. Atl.* 357.—Alger.

RAPOLOCARPUS, Boj. Hort. Maurit. 44 (1837) err. typ. = **Ropalocarpus**, Boj. (Tiliac.).

RAPOUREA, Reichb. Consp. 118 (1828), sphalm. = **Ropourea**, Aubl. (Verben. ?).

RAPUNCULUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111 = **Campanula**, Tourn.
patulus, Fourr. l. c. = C. patula.
persicaefolius, Fourr. l. c. = C. persicifolia.
verus, Fourr. l. c. = C. Rapunculus.

RAPUNCULUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Phyteuma**, Linn. (Campan.).
comosus, Mill. Gard. Dict. ed. VIII. n. 2 = P. comosum.
esculentus, Fl. Dan. ex Steud. Nom. ed. II. ii. 437 = Campanula rotundifolia.
hemisphaericus, Mill. Gard. Dict. ed. VIII. n. 3 = P. hemisphaericum.
orbicularis, Mill. l. c. n. 5 = P. orbiculare.
pauciflorus, Mill. l. c. n. 4 = P. pauciflorum.
spicatus, Mill. l. c. n. 1 = P. spicatum.

RAPUNTIUM, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768) = **Lobelia**, Linn. (Campan.).
acuminatum, Presl, Prod. Monog. Lobel. 24 = L. acuminata.
acutangulum, Presl, l. c. 16 = L. triquetra.
affine, Presl, l. c. 13 = L. affinis.
affine, Presl, l. c. 25 = L. Cliffortiana.
alatum, Presl, l. c. 14 = L. anceps.
alsinoides, Presl, l. c. 22 = L. alsinoides.
amoenum, Presl, l. c. 23 = L. amoena.
amygdalinum, Presl, l. c. 27 = L. amygdalina.
anceps, Presl, l. c. 14 = L. anceps.
angulatum, Presl, l. c. 30 = Pratia angulata.
aquaticum, Presl, l. c. 21 = L. aquatica.
arabideum, Presl, l. c. 18 = Laurentia arabidea.
arenarioides, Presl, l. c. 17 = L. trigona.
bellidifolium, Presl, l. c. 15 = L. Erinus.
Besserianum, Presl, l. c. 28 = L. polyphylla.
bicolor, Presl, 17 = L. Erinus.
bifidum, Presl, l. c. 30 = Laurentia Dregeana.
blandum, Presl, l. c. 27 (= *Tupa blanda*).—Chili.
Boryanum, l. c. 24 = L. serpens.
bracteosum, Presl, l. c. 29 = L. polyphylla.
brevifolium, Presl, l. c. 23 = L. brevifolia.
Breynii, Presl, l. c. 19 = L. Breynii.
Bridgesii, Presl, l. c. 28 = L. Bridgesii.
Brownianum, Presl, l. c. 30 = L. gibbosa.
Burmannianum, Presl, l. c. 31 = L. Burmanniana.
caeruleum, Presl, l. c. 20 = L. coronopifolia.
caespitosum, Presl, l. c. 13 = L. radicans.
campanuloides, Presl, l. c. 14 = L. radicans.
camporum, Presl, l. c. 15 = L. camporum.
capillifolium, Presl, l. c. 19 = L. capillifolia.
capitatum, Presl, l. c. 11 = L. lutea.
cardinale, Mill. Gard. Dict. ed. VIII. n. 1 = L. cardinalis.
Cavanillesianum, Presl, Prod. Monog. Lobel. 27 = L. laxiflora.
ceratophyllum, Presl, in Eckl. & Zeyh. Enum. 399 = L. tomentosa.
chamaedrifolium, Presl, Prod. Monog. Lobel. 21 = L. chamaedrifolia.
Chamaepitys, Presl, l. c. 20 = L. tomentosa.
cheiranthifolium, Presl, l. c. 18 = L. serpens.
chenopodifolium, Presl, l. c. 31 = L. Cliffortiana.
chinense, Presl, l. c. 13 = L. chinensis.

RAPUNTIUM:—
cinereum, Presl, Prod. Monog. Lobel. 30 = Lightfootia albicaulis.
circaeoides, Presl, l. c. 25 = Lobelia circaeoides.
cirsiifolium, Presl, l. c. 27 = L. cirsiifolia.
Claytonianum, Presl, l. c. 23 = L. spicata.
Cliffortianum, Mill. Gard. Dict. ed. VIII. n. 3 = L. Cliffortiana.
coccineum, Moench, Meth. Suppl. 277 = L. cardinalis.
coeruleum, Auct. = *caeruleum*.
collinum, Presl, Prod. Monog. Lobel. 23 = L. collina.
coloratum, Presl, l. c. 24 = L. colorata.
coloratum, Presl, l. c. 30 = L. amoena.
cordatum, Presl, in E. Mey. Comm. Pl. Afr. Austr. 285 = L. Preslii.
cordifolium, Moench, Meth. Suppl. 277 = L. Cliffortiana.
cordigerum, Presl, Prod. Monog. Lobel. 26 = L. cordigera.
coronopifolium, Presl, l. c. 20 = L. coronopifolia.
corymbosum, Presl, in E. Mey. Comm. Pl. Afr. Austr. 287 = L. corymbosa.
cuneiforme, Presl, Prod. Monog. Lobel. 14 = L. anceps.
decurrens, Presl, l. c. 24 = L. decurrens.
dentatum, Presl, l. c. 21 = L. dentata.
Dortmanna, Presl, l. c. 18 = L. Dortmanna.
Dregeanum, Presl, l. c. 22 = L. Dregeana.
Ecklonianum, Presl, in Eckl. & Zeyh. Enum. 396 = L. Eckloniana.
ericoides, Presl, Prod. Monog. Lobel. 12 = L. ericoides.
erinoides, Mill. Gard. Dict. ed. VIII. n. 9 = Laurentia repens.
Erinus, Mill. l. c. n. 8 = L. Erinus.
euphrasioides, Presl, Prod. Monog. Lobel. 12 = L. euphrasioides.
exaltatum, Presl, l. c. 24 = Haynaldia exaltata.
excelsum, Presl, l. c. 29 = L. salicifolia.
fastigiatum, Presl, l. c. 15 = L. fastigiata.
fenestrale, Presl, l. c. 13 = L. fenestralis.
fervens, Presl, l. c. 30 = L. fervens.
filiforme, Presl, l. c. 18 = L. serpens.
flaccidum, Presl, l. c. 13 = L. Erinus.
flavum, Presl, in Eckl. & Zeyh. Enum. 394 = L. flava.
flexuosum, Presl, Prod. Monog. Lobel. 16 = L. natalensis.
flexuosum, Presl, l. c. 23 = L. flexuosa.
fluviatilis, Presl, l. c. 30 = Isotoma fluviatilis.
foliosum, Presl, l. c. 24 = L. decurrens.
fulgens, Presl, l. c. 26 = L. fulgens.
genistoides, Presl, in E. Mey. Comm. Pl. Afr. Austr. 285 = L. linearis.
gibbosum, Presl, Prod. Monog. Lobel. 13 = L. gibbosa.
glabrifolium, Presl, l. c. 12 = L. leptocarpa.
glandulosum, Presl, l. c. 21 = L. glandulosa.
gracile, Presl, l. c. = L. gracilis, *Andr.*
gramineum, Presl, l. c. 26 = L. graminea.
gruinum, Presl, l. c. 25 = L. gruina.
Haenkeanum, Presl, l. c. 26 = L. Haenkeana.
heteromallum, Presl, l. c. 15 = L. triquetra.
heterophyllum, Presl, l. c. 13 = L. heterophylla.
hirsutum, Moench, Meth. 656 = L. hirsuta.
hirtum, Mill. Gard. Dict. ed. VIII. n. 6 = L. hirsuta.
hyssopifolium, Presl, Prod. Monog. Lobel. 29 = L. hyssopifolia.
incisum, Presl, in Eckl. & Zeyh. Enum. 397 = L. pubescens.
inflatum, Mill. Gard. Dict. ed. VIII. n. 5 = L. inflata.
inundatum, Presl, Prod. Monog. Lobel. 30 = Isotoma fluviatilis.
irriguum, Presl, l. c. = Pratia irrigua.
Kalmii, Presl, l. c. 23 = L. Kalmii.
kamtschaticum, Presl, l. c. 24 = L. sessilifolia.
Kohautianum, Presl, l. c. 15 = L. senegalensis.
Kraussii, Presl, l. c. 27 = L. Kraussii.
Krebsianum, Presl, l. c. 15 = L. Erinus.
Kunthianum, Presl, l. c. 27 = L. laxiflora.
laciniatum, Presl, l. c. = Siphocampylus Lamarckii.
lasianthum, Presl, in E. Mey. Comm. Pl. Afr. Austr. 288 = L. lasiantha.
laxiflorum, Presl, Prod. Monog. Lobel. 26 = L. laxiflora.
Leschenaultianum, Presl, l. c. 24 = L. excelsa.
linarioides, Presl, l. c. 22 = L. linarioides.
longiflorum, Mill. Gard. Dict. ed. VIII. n. 7 = Isotoma longiflora.
longifolium, Presl, Prod. Monog. Lobel. 26 = L. longifolia.

RAPUNTIUM :—
Lucaeanum, Presl, Prod. Monog. Lobel. 27 = Lobelia Lucaeana.
luteum, Presl, l. c. 11 = L. lutea.
maculare, Presl, l. c. 20 = L. coronopifolia.
membranaceum, Presl, l. c. 30 = L. membranacea.
Michauxii, Presl, l. c. 25 = L. Cliffortiana?
micranthum, Presl, l. c. = L. micrantha.
microdon, Presl, l. c. 15 = L. Erinus.
molle, Presl, l. c. 30 = L. mollis.
monticolum, Presl, l. c. 25 = L. monticola.
mucronatum, Presl, l. c. 29 = L. mucronata, *Cav.*
multicaule, Presl, l. c. 31 = L. multicaule.
Mundianum, Presl, in Eckl. & Zeyh. Enum. 395 = L. leptocarpa.
muscoides, Presl, Prod. Monog. Lobel. 22 = L. muscoides.
nanum, Presl, l. c. = L. nana.
nicotianaefolium, Presl, l. c. 24 = L. nicotianaefolia.
nummularioides, Presl, l. c. 21 = L. nummularioides.
nummularium, Presl, l. c. 30 = Pratia begonifolia.
Nuttallianum, Presl, l. c. 23 = L. Nuttallii.
obovatum, Presl, l. c. 25 = L. incana.
Ottonianum, l. c. 14 = L. Ottoniana.
ovatum, Presl, l. c. 16 = L. pubescens.
ovatum, Presl, l. c. 29 (= *Tupa ovata*, G. Don).—Peruv.
paludosum, Presl, l. c. 18 = L. paludosa.
parvifolium, Presl, l. c. 31 = L. parvifolium.
patulum, Presl, l. c. 30 = L. patula.
paucidentatum, Presl, l. c. 20 = L. tomentosa.
pauciflorum, Presl, l. c. 15 = L. pauciflora.
pedunculare, Presl, l. c 19 = L. hirsuta.
pedunculatum, Presl, l. c. 30 = Pratia pedunculata.
persicifolium, Presl, l. c. 27 = L. persicaefolia.
pinifolium, Presl, l. c. 19 = L. pinifolia.
platycaulum, Presl, l. c. 14 = L. madagascariensis.
Poeppigianum, Presl, l. c. 29 = L. polyphylla.
polyphyllum, Presl, l. c. 28 = L. polyphylla.
procumbens, Presl, l. c. 17 = L. procumbens.
pteropodum, Presl, l. c. 21 = L. patula.
puberulum, Presl, l. c. 23 = L. puberula.
pubescens, Presl, l. c. 17 = P. pubescens.
purpurascens, Presl, l. c. = L. purpurascens.
purpureum, Presl, l. c. 29 = L. purpurea, *Lindl.*
pusillum, Presl, l. c. 22 = L. pusilla.
pygmaeum, Presl, l. c. 22 = Laurentia micrantha.
pyramidale, Presl, l. c. 23 = L. pyramidalis.
quadrangulare, Presl, l. c. 16 = L. quadrangularis.
racemosum, Presl, l. c. 25 = L. racemosa, *Sims.*
radicans, Presl, l. c. 14 = L. radicans.
rapunculoides, Presl, l. c. 23 = L. rapunculoides.
Reinwardtianum, Presl, l. c. 14 = L. trigona.
reniforme, Presl, l. c. 15 = L. reniformis.
Richardianum, Presl, l. c. 27 = L. decumbens.
rigidulum, Presl, l. c. 26 = L. laxiflora.
roseum, Presl, l. c. 24 = L. rosea.
rugulosum, Presl, l. c. 30 = Pratia angulata.
rupestre, Presl, l. c. 15 = L. rupestris.
scabripes, Presl, l. c. 17 = L. Erinus.
secundum, Presl, l. c. 29 = L. secunda.
serpens, Presl, l. c. 15 = L. serpens.
serrulatum, Presl, l. c. 14 = L. urens.
setaceum, Presl, l. c. 19 = L. setacea.
simplex, Presl, l. c. 17 = L. debilis.
simplicicaule, Presl, l. c. 30 = L. gibbosa.
Simsii, Presl, l. c. 20 = L. Simsii.
siphiliticum, Mill. Gard. Dict. ed. VIII. n. 2 = L. syphilitica.
sonchifolium, Presl, Prod. Monog. Lobel. 27 = L. sonchifolia.
spartioides, Presl, in Eckl. & Zeyh. Enum. 395 = L. spartioides.
speciosum, Presl, Prod. Monog. Lobel. 19 = L. syphilitica.
spicatum, Presl, l. c. 31 = L. spicata, *Ruiz & Pav.*
splendens, Presl, l. c. 26 = L. fulgens.
strictum, Presl, l. c. 27 = L. stricta.
subdentatum, Presl, l. c. 28 = L. polyphylla.
subtile, Presl, l. c. 25 = L. trigona.
succulentum, Presl. l. c. 13 = L. succulenta.
syphiliticum, Moench, Meth. 655 = L. syphilitica.
telephioides, Presl, Prod. Monog. Lobel. 18 = L. serpens.
tenerum, Presl, l. c. 15 = L. tenera.
tenuifolium, Presl, in Eckl. & Zeyh. Enum. 395 = L. ericetorum.
tenuius, Presl, Prod. Monog. Lobel. 30 = L. tenuior.
thapsoideum, Presl, l. c. 24 = Haynaldia thapsoidea.
thermale, Presl, l. c. 12 = L. leptocarpa.

RAPUNTIUM :—
Thunbergii, Presl, Prod. Monog. Lobel. 20 = Lobelia coronopifolia.
tomentosum, Presl, l. c. 19 = L. tomentosa.
trialatum, Presl, l. c. 13 = L. trigona.
triquetrum, Presl, l. c. 16 = L. triquetra.
Tupa, Presl, l. c. 28 = L. Tupa.
umbellatum, Presl, l. c. 31 = L. umbellata.
uranocomum, Presl, l. c. 24 = Haynaldia uranocoma.
urens, Mill. Gard. Dict. ed. VIII. n. 4 = L. urens.
variifolium, Presl, Prod. Monog. Lobel. 12 = L. variifolia.
veronicaefolium, Presl, l. c. 23 = L. veronicaefolia.
Wallichianum, Presl, l. c. 24 = L. pyramidalis.
xalapense, Presl, l. c. 25 = L. Cliffortiana.
zeylanicum, Presl, l. c. 13 = L. zeylanica.

RAPUTIA, Aubl. Pl. Gui. ii. 670. t. 272 (1775). *RUTACEAE*, [Benth. & Hook. f. i. 285].
PHOLIDANDRA, Neck. Elem. i. 348 (1790).
SCIURIS, Schreb. Gen. 24 (1789).
alba, *Engl. in Mart. Fl. Bras.* xii. II. 102.—Bras.
aromatica, *Aubl. Pl. Gui.* ii. 671. t. 272.—Guian.
Conchocarpus, Schult. Mant. i. 126 = Cusparia macrophylla.
heterophylla. *DC. in Mém. Mus. Par.* ix. (1822) 153.—Porto Rico.
magnifica, *Engl. in Mart. Fl. Bras.* xii. II. 102.—Bras.
Ossana, *Engl. l. c.* 104.—Cuba.
trifoliata, *Engl. l. c.* 103.—Bras.

RARAM, Adans. Fam. ii. 35 (1763) = **Cenchrus**, Linn. (Gramin.).

RASPAILIA, J. & C. Presl, Rel. Haenk. i. 238. t. 40 (1830) = **Polypogon**, Desf. (Gramin.).
agrostoides, J. & C. Presl, l. c. 239 = P. elongatus.

RASPALIA, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 377. t. 37. *BRUNIACEAE*, Benth. & Hook. f. i. 672.
angulata, E. Mey. in Drège, Zwei Pfl. Docum. 215 = Erica modesta.
aspera, *E. Mey. l. c.*, nomen; *et ex Harv. & Sond. Fl. Cap.* ii. 321.—Afr. austr.
capitella, Presl, Bot. Bemerk. 39 = Brunia staavioides.
microphylla, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 378.—Afr. austr.
passerinoides, *Presl, Bot. Bemerk.* 39.—Afr. austr.
phylicoides, *Presl, l. c.*—Afr. austr.
struthioloides, *Presl, l. c.* 40.—Afr. austr.
teres, E. Mey. in Drège, Zwei Pfl. Docum. 215 = microphylla.
villosa, Presl, Bot. Bemerk. 39 = Brunia villosa.

RASSIA, Neck. Elem. ii. 12 (1790) = **Gentiana**, Linn.

RATABIDA, Loud. Hort. Brit. Suppl. ii. 665 (1839), sphalm. = RATIBIDA, Rafin. = **Lepachys**, Rafin. (Compos.).

RATHEA, Karst. Fl. Columb. i. 77. t. 38 (1859?) = **Tacsonia**, Juss. (Passifl.).
floribunda, Karst. l. c. = T. floribunda.

RATHEA, Karst. in C. Koch & Fint. Wochenschr. i. (1858) 377 = **Synechanthus**, H. Wendl. (Palm.).
fibrosa, Karst. l. c. = S. fibrosus.

RATHKEA, Schum. & Thonn. Beskr. Guin. Pl. 355 (1827) = **Ormocarpum**, Beauv. (Legumin.).
glabra, Schum. & Thonn. l. c. = O. sennoides.
pubescens, Walp. Ann. i. 1105 = O. bibracteatum.

RATIBIDA, Rafin. in Am. Monthly Mag. (1818) 268 = **Lepachys**, Rafin. (Compos.).
columnaris, Rafin. l. c.; D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 361 = L. columnaris.
pulcherrima, Steud. Nom. ed. II. ii. 439 = L. columnaris.
sulcata, Rafin. in Journ. Phys. lxxxix. (1819) 100 = L. columnaris.

RATONIA, DC. Prod. i. 618 (1824). *SAPINDACEAE*, Benth. & Hook. f. i. 399.
ARYTERA, Blume, Rumphia, iii. 170 (1847).
EPHIELIS, Schreb. Gen. 253 (1789).
ERNSTINGIA, Scop. Introd. 322 (1777).
GELONIUM, Gaertn. Fruct. ii. 271. t. 139 (1791).

RATONIA:—
LACHNOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 571.
LEPIDOPETALUM, Blume, Rumphia, iii. 171 (1847).
MATAYBA, Aubl. Pl. Gui. i. 331. t. 128 (1775), nomen prius.
MISCHOCARPUS, Blume, Bijdr. 238 (1825); Rumphia, iii. 166 (1847).
PEDICELLIA, Lour. Fl. Cochinch. 655 (1790).
SARCOTOECHIA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 501, 659.
TINA, Roem. & Schult. Syst. v. 414 (1819).
ZYGOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 574.
Alphandi, F. Muell. Fragm. iv. 158 = Castanospora Alphandi.
anodonta, *Benth. & Hook. f. Gen.* i. 400; *Benth. Fl. Austral.* i. 461.—Austral.
apetala, *Griseb. Fl. Brit. W. Ind.* 126.—Ind. occ.
apetala, Seem. Fl. Vit. 47 = Elattostachys apetala.
Cordierii, F. Muell. Fragm. ix. 93 = Synima Cordierii.
Daemeliana, F. Muell. l. c. 96 = Toechima Daemeliana.
diphyllostegia, F. Muell. l. c. v. 145 = Diploglottis Cunninghamii.
distylis, *F. Muell. ex Benth. Fl. Austral.* i. 462.—Austral.
domingensis, *DC. Prod.* i. 618.—Ins. S. Doming.
exangulata, *F. Muell. Fragm.* iv. 156.—Austral.
falcata, *Seem. Fl. Vit.* 47.—Ins. Fiji.
grandissima, *F. Muell. Fragm.* iv. 156.—Austral.
Hornei, *Seem. Fl. Vit.* 47.—N. Caled.
lachnocarpa, *F. Muell. Fragm.* iv. 157.—Austral.
lachnopetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 586.—Ins. Philipp.
Lessertiana, Benth. & Hook. f. Gen. i. 400 = Cupania Lessertiana.
Martyana, F. Muell. Fragm. v. 6 = Sarcopteryx Martyana.
O'Shanesiana, *F. Muell. Fragm.* ix. 96.—Austral.
pyriformis, *Benth. & Hook. f. Gen.* i. 400; *Benth. Fl. Austral.* i. 461.—Austral.
spathulata, *Griseb. Cat. Pl. Cub.* 45.—Cuba.
stipitata, *Benth. Fl. Austral.* i. 461.—Austral.
Storckii, Seem. Fl. Vit. 47 = Cupaniopsis Storckii.
tenax, *Benth. Fl. Austral.* i. 461.—Austral.
Zygolepis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 586.—Ins. Philipp.

RATOPITYS, Carr. Conif. ed. II. 227 (1867) = RAXOPITYS, Nelson = **Cunninghamia**, R. Br. (Conif.).

RATZEBURGIA, Kunth, Rév. Gram. ii. 487. t. 158 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1130.
AIKINIA, Wall. Pl. As. Rar. iii. 46. t. 273 (1832).
pulcherrima, *Kunth, Rév. Gram.* ii. 487.—Ind. or.
Schimperi, Steud. Nom. ed. II. ii. 439 = Elionurus Royleanus.

RAUIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 151, 167. *RUTACEAE*, [Benth. & Hook. f. i. 285].
racemosa, Nees & Mart. l. c. 169 = Cusparia Fontanesiana.
resinosa, *Nees & Mart. l. c.*—Bras.

RAUKANA, Seem. in Journ. Bot. iv. (1866) 352 = **Panax**, Linn. (Araliac.).
Edgerleyi, Seem. l. c. = P. Edgerleyi.

RAUVOLFIA, Plum. ex Linn. Gen. ed. I. 62 (1737) = **Rauwolfia**, Plum. (Apocyn.).

RAUWENHOFFIA, Scheff. in Ann. Jard. Buitenz. ii. (1885) 21. *ANONACEAE*.
siamensis, *Scheff. l. c.* 23.—Siam.
uvarioides, *Scheff. l. c.*—Ins. Aru.

RAUWOLFIA, Plum. ex Linn. Gen. ed. I. 62 (1737) (Rauvolfia). *APOCYNACEAE*, Benth. & Hook. f. ii. 697.
CYRTOSIPHONIA, Miq. Fl. Ind. Bat. ii. 401 (1856).
DISSOLAENA, Lour. Fl. Cochinch. 137 (1790).
HEURCKIA, Muell. Arg. in Flora, liii. (1870) 168.
OPHIOXYLON, Linn. Fl. Zeyl. 239 (1747).
affinis, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 34.—Bras.
Alphonsiana, *Muell. Arg. in Linnaea*, xxx. (1859-60) 394.—Ind. occ.

RAUWOLFIA:—
amsoniaefolia, *A. DC. in DC. Prod.* viii. 338.—Ins. Philipp.
angustifolia, Salisb. Prod. 146 = nitida.
bahiensis, *A. DC. in DC. Prod.* viii. 339.—Bras.
Beddomei, *Hook. f. Fl. Brit. Ind.* iii. 634.—Ind. or.
biauriculata, *Muell. Arg. in Linnaea*, xxx. (1859-60) 396.—Ind. occ.
Blanchetii, *A. DC. in DC. Prod.* viii. 340.—Bras.
brasiliensis, *Spreng. Syst.* i. 834.—Bras.
caffra, *Sond. in Linnaea*, xxiii. (1850) 77.—Afr. austr.
canescens, *Linn. Sp. Pl.* ed. II. 303.—Ind. occ.
cubana, *A. DC. in DC. Prod.* viii. 339.—Cuba.
decurva, *Hook. f. Fl. Brit. Ind.* iii. 633.—Ind. or.
densiflora, *Benth. ex Hook. f. l. c.*—Ind. or.
glabra, Cav. Ic. iii. 50 = Vallesia dichotoma.
grandiflora, *Mart. ex A. DC. in DC. Prod.* viii. 341.—Bras.
heterophylla, *Willd. ex Roem. & Schult. Syst.* iv. 805.—Am. austr.
hirsuta, Jacq. Enum. Pl. Carib. 14 = canescens.
laevigata, Willd. ex Roem. & Schult. Syst. iv. 805 = Malouetia jasminoides.
Lamarkii, *A. DC. in DC. Prod.* viii. 337.—Am. trop.
lanceolata, *A. DC. l. c.*—Porto Rico.
lanceolata, Griseb. Fl. Brit. W. Ind. 408, in syn. = Lamarkii.
latifolia, A. DC. in DC. Prod. viii. 339 = Lamarkii.
ligustrina, Willd. ex Roem. & Schult. Syst. iv. 805 = ternifolia.
longifolia, *A. DC. in DC. Prod.* viii. 338.—Mexic.
lycioides, *Cav. in Anal. Cienc. Nat.* v. (1802) 69.—Hab.?
micrantha, *Hook. f. Fl. Brit. Ind.* iii. 633.—Ind. or.
microcarpa, *Hook. f. l. c.*—Burma.
Moricandii, *A. DC. in DC. Prod.* viii. 340.—Bras.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 78.—Afr. austr.
nitida, *Jacq. Enum. Pl. Carib.* 14; *Select. Am.* 47.—Ind. occ.
nitida, Lam. Illustr. ii. 304 = Lamarkii.
nitidissima, Steud. Nom. ed. I. 682 = nitida.
obtusiflora, *A. DC. in DC. Prod.* viii. 341.—Ins. Nossibé.
odontophora, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 150.—Ecuador.
oppositifolia, Spreng. Neue Entdeck. iii. 33 = Tabernaemontana Berterii?
parvifolia, Bert. ex Spreng. Syst. i. 834 = ternifolia.
paucifolia, *A. DC. in DC. Prod.* viii. 340.—Bras.
peguana, *Hook. f. Fl. Brit. Ind.* iii. 632.—Burma.
polyphylla, *Benth. in Hook. Journ. Bot.* iii. (1841) 241.—Guiana.
psychotrioides, *H. B. & K. Nov. Gen. et Sp.* iii. 231.—Am. austr.
Pulaparia, *Roxb. ex Sm. in Rees, Cycl.* xxix. n. 5.—Hort. Calcutt.
reflexa, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 405.—Java.
salicifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 519.—Cuba.
sandwicensis, *A. DC. in DC. Prod.* viii. 339.—Ins. Sandvic.
Sellowii, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 33.—Bras.
Senegambiae, *A. DC. in DC. Prod.* viii. 340—Afr. trop.
serpentina, *Benth. ex Kurz, For. Fl. Brit. Burma*, ii. 171.—Ind. or.; Java.
Sprucei, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 34.—Bras.
strempelioides, *Griseb. Cat. Pl. Cub.* 170.—Cuba.
striata, Poir. Encyc. vi. 83 = Ochrosia borbonica.
subpubescens, Linn. Mant. ii. 345 = canescens.
sumatrana, *Jack, in Malay. Misc.* i. (1820) v. 22.—Sumatra.
ternifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 232.—Am. trop.
tetraphylla, Linn. Sp. Pl. 208 = nitida, canescens.
tomentosa, Jacq. Enum. Pl. Carib. 14 = canescens.
viridis, Willd. ex Roem. & Schult. Syst. iv. 805 = psychotrioides.
vomitoria, *Afzel. Stirp. Guin. Med. Sp. Nov.* 1.—Afr. trop.
Weddeliana, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 32.—Bras.

RAUWOLFIA, Ruiz & Pav. Fl. Per. et Chil. ii. 26. t. 152 (1799) = **Citharexylum**, Mill. (Verben.).
dentata, Tafalla, ex D. Don, in Edinb. N. Phil. Journ. (Jan.-Mar. 1831) 237 = C. dentatum.
flexuosa, Ruiz & Pav. Fl. Per. ii. 16 = C. spinosum.
macrophylla, Ruiz & Pav. l. c. = C. spinosum.
pubescens, Willd. ex Roem. & Schult. Syst. iv. 805 = C. molle.
spinosa, Cav. Ic. vi. 16.—Peruv.

RAVENALA, Adans. Fam. ii. 67 (1763). *SCITAMINEAE*, Benth. & Hook. f. iii. 656.
PHENAKOSPERUM, Endl. Prod. Fl. Norf. 35, in nota (1833).
URANIA, Schreb. Gen. 212 (1789).
guyanensis, *Steud. Nom.* ed. II. ii. 439.—Guian.
madagascariensis, *J. F. Gmel. Syst.* 567; *Poir. Encyc. Suppl.* iv. 656; *Lam. Illustr.* ii. 371.—Madag.

RAVENEA, H. Wendl. ex Bouché, in Monatschr. Ver. Befoerd. Gartenb. (1878) 197, 323, 324. Gen. dub. *PALMARUM*.—Cf. Benth. & Hook. f. iii. 883.
Hildebrandti, H. Wendl. ex Bouché, l. c.—Ins. Comor.

RAVENIA, Benth. & Hook. f. Gen. iii. 883 (1883), sphalm. = praec.

RAVENIA, Vell. Fl. Flum. 20 (1825); i. t. 49 (1827). *RUTACEAE*, Benth. & Hook. f. i. 286.
LEMONIA, Lindl. Bot. Reg. (1840) t. 59.
infelix, *Vell. Fl. Flum.* 21; i. t. 49.—Bras.
spectabilis, *Engl. in Mart. Fl. Bras.* xii. II. 126.—Cuba; Bras.

RAVENSARA, Sonner. Voy. Ind. Or. iii. 248. t. 127 (1782). *LAURINEAE*, Benth. & Hook. f. iii. 151.
AGATHOPHYLLUM, Juss. Gen. 431 (1789).
EVODIA, Gaertn. Fruct. ii. 100 [t. 103] (1791).
acuminata, *Baill. Adansonia*, ix. (1868-70) 303.—Madag.
aromatica, *J. F. Gmel. Syst.* 754.—Madag.
floribunda, *Baill. Adansonia*, ix. (1868-70) 304.—Madag.
Lastellii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 447.—Madag.
retusa, *Baill. Adansonia*, ix. (1868-70) 303.—Madag.
Topak, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 448.—Madag.

RAVIA, Schult. Addit. ii. ad Mant. ii. 215, in syn. (1824), err. typ. = **Rauia**, Nees & Mart. (Rutac.).

RAVNIA, Oerst. in Kjoeb. Vidensk. Meddel. (1852) 49 (Centr. Am. Rub.). *RUBIACEAE*, Benth. & Hook. f. ii. 40.
triflora, *Oerst. l. c.* 50.—Costa Rica.

RAWSONIA, Harv. & Sond. Fl. Cap. i. 67 (1859). *BIXINEAE*, Benth. & Hook. f. i. 127.
lucida, *Harv. & Sond. l. c.*—Afr. austr.

RAXAMARIS, Rafin. Fl. Tellur. iii. 54 (1836) = **Ornithogalum**, Tourn. (Liliac.).
parviflora, Rafin. l. c. 55 = O. exscapum.

RAXOPITYS, [Nelson], Senilis, Pinaceae, 97 (1866) = **Cunninghamia**, R. Br. (Conif.).
Cunninghamii, [Nelson], l. c. = C. sinensis.

RAYANIA, Meissn. in DC. Prod. xiv. 185 (1856) = RAJANIA, Walt. = **Brunnichia**, Banks (Polygon.).

RAYERIA, Gaudich. Bot. Voy. Bonite, t. 108 (1844-52) = **Alona**, Lindl. (Convolv.).
teretifolia, Gaudich. l. c.—Hab.?

RAZISEA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 142. *ACANTHACEAE*, Benth. & Hook. f. ii. 1106.
spicata, *Oerst. l. c.*—Costa Rica.

RAZOUMOWSKIA, Hoffm. Hort. Mosq. (1808); ex Bieb. Fl. Taur. Cauc. iii. 629 (1819) = **Arceuthobium**, Bieb. (Loranth.).
capensis, Hoffm. l. c.; ex J. S. Presl, Rostlin. ii. 28 = Viscum capense.
caucasica, Hoffm. l. c. Introd. n. 1. f. 1; ex Bieb. Fl. Taur. Cauc. iii. 629 = A. Oxycedri.
jamaicensis, Hoffm. l. c.; ex J. S. Presl, Rostlin. ii. 28 = Dendrophthora opuntioides.
mexicana, Hoffm. l. c.; ex J. S. Presl, l. c. = A. vaginatum.
Oxycedri, F. Schultz, ex Nym. Consp. 320 = A. Oxycedri.

RAZUMOVIA, Spreng. ex Juss. in Dict. Sc. Nat. xliv. 526 (1826) = **Humea**, Sm. (Compos.).
paniculata, Spreng. ex DC. Prod. vi. 158 = H. elegans.

RAZUMOVIA, Spreng. Fl. Hal. Mant. 45 (1807) = **Centranthera**, R. Br. (Scrophul.).
tranquebarica, Spreng. l. c. = C. humifusa.

REA, Bert. ex Decne. in Guill. Arch. Bot. i. (1833) 513. tt. 9, 10 = **Dendroseris**, D. Don (Compos.).
Berteriana, Decne. l. c. 515 = D. Berteriana.
lacerata, Phil. in Bot. Zeit. xxviii. (1870) 499 = D. lacerata.
leucantha, Bert. ex DC. Prod. vii. 243 = D. neriifolia.
macrantha, Bert. ex Decne. in Guill. Arch. Bot. i. (1833) 514. t. 9 = D. macrophylla.
marginata, Bert. ex Decne. l. c. 518. t. 9 = D. macrophylla.
micrantha, Bert. ex Decne. l. c. 517 = D. micrantha.
mollis, Bert. ex Decne. l. c. 519 = D. micrantha.
neriifolia, Decne. l. c. 517 = D. neriifolia.
pinnata, Bert. ex Decne. l. c. 516 = D. Berteriana.

REANA, Brign. Ind. Sem. Hort. Moden. (1849).—Cf. Ann. Sc. Nat. Sér. III. xii. (1849) 365 = **Euchlaena**, Schrad. (Gramin.).
Giovannini, Brign. ll. cc. = E. mexicana.
luxurians, Dur. in Bull. Soc. Acclim. Sér. II. ix. (1872) 581 = E. mexicana.
luxurians, Hort.—Cf. Gard. Chron. (1875) II. 174; (1876) I. 566 = Tripsacum dactyloides.

REAUMUREA, Steud. Nom. ed. I. 682 (1821) = seq.

REAUMURIA, Linn. Syst. ed. X. 1081 (1759). *TAMARISCINEAE*, Benth. & Hook. f. i. 161.
EICHWALDIA, Ledeb. in Eichw. Pl. Casp. 38. t. 34 (1831).
Billardieri, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. III. viii. (1847) 381.—Syria.
cistoides, Adam, in Weber & Mohr, Beitr. i. 61 = hypericoides.
filifolia, *Jaub. & Spach, Illustr.* iii. 54, 58.—Persia.
floribunda, *Regel, Cat. Pl. Hort. Aksakov.* 118.—Hab.?
Floyeri, *S. Moore, in Journ. Bot.* xv. (1877) 289.—Beluchist.
fruticosa, *Boiss. Fl. Orient.* i. 759.—Persia.
hirtella, *Jaub. & Spach, Illustr.* iii. 54, 55.—Egypt.; Arab.
hypericoides, *Willd. Sp. Pl.* ii. 1250.—Reg. Caucas.; Persia.
hyrcanica, Jaub. & Spach, Illustr. iii. 55, 60 = hypericoides.
kaschgarica, *Rupr. Sert. Tiansch.* 42.—Turkestan.
linifolia, Salisb. Parad. Lond. t. 18 = hypericoides.
mucronata, *Jaub. & Spach, Illustr.* iii. 54, 57.—Egypt.
oxiana, *Boiss. Fl. Orient.* i. 759.—Reg. Casp.; Persia.
palaestina, *Boiss. Diagn.* Ser. I. x. 10.—Palaestin.
persica, Boiss. Fl. Orient. i. 761 = filifolia.
squarrosa, *Jaub. & Spach, Illustr.* iii. 59.—Persia; Afghan.
stenophylla, Jaub. & Spach, l. c. 54 = mucronata.
Stocksii, *Boiss. Fl. Orient.* i. 761.—Beluchist.
tatarica, Jaub. & Spach, Illustr. iii. 55 = hypericoides.
trigyna, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 425.—Mongol.
vermiculata, Bieb. Fl. Taur. Cauc. ii. 17 = hypericoides.
vermiculata, *Linn. Syst.* ed. X. 1081.—Ital.; Afr. bor.
vermiculata, Linn. l. c., partim = hirtella.

REBENTISCHIA, Opiz, in Lotos, jv. (1854) 104 = **Trisetum**, Pers. (Gramin.).
flavescens, Opiz, l. c. = T. flavescens.

REBIS, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 26 = **Ribes**, Linn. (Saxifr.).
alpinoides, Steud. Nom. ed. II. ii. 439 = Ribes alpinoides.
discolor, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 28 = Ribes albifolium.
ebracteolata, Spach, l. c. 31 = Ribes alpinoides.
frigida, Spach, l. c. 29 = Ribes hirtum.
Gayana, Spach, l. c. = Ribes Gayanum.
macrobotrys, Spach, l. c. 27 = Ribes macrobotrys.
punctata, Spach, l c. 30 = Ribes punctata.

REBOUDIA, Coss. & Dur. in Bull. Soc. Bot. Fr. iii. (1856) 704 = **Erucaria**, Gaertn. (Crucif.).
erucarioides, Coss. & Kralik, l. c. 705 = E. erucarioides.

REBOULEA, Kunth, Rév. Gram. 341. t. 84 (1830) = **Eatonia**, Rafin. (Gramin.).
gracilis, Kunth, l. c. = E. obtusata.
obtusata, A. Gray, Man. Bot. N. U. St. ed. I. 591 = E. obtusata.
pennsylvanica, A. Gray, l. c. = E. purpurascens.
truncata, Torr. ex Munro, in Journ. Linn. Soc. vi. (1862) 43 = E. obtusata.

RECCHIA, Moç. & Sesse, ex DC. Syst. i. 411 (1818) = **Rigiostachys**, Planch. (Simarub.).
mexicana, Moç. & Sesse, l. c. = Rigiostachys bracteata.

RECEVEURA, Vell. Fl. Flum. 237 (1825); v. tt. 119, 120 (1827) = **Hypericum**, Linn.
cordata, Vell. Fl. Flum. 237; v. t. 119.—Bras.
graveolens, Vell. l. c. 237; v. t. 120 = H. brasiliense.

RECHSTEINERIA, Regel, in Flora, xxi. (1848) 247 = **Gesneria**, Linn.
allagophylla, Regel, l. c. = G. allagophylla.
spicata, Kumm. ex Hanst. in Mart. Fl. Bras. viii. 357 = G. tribracteata.
stricta, Hanst. in Linnaea, xxvi. (1853) 214. t. 1. f. 19 = G. stricta.
tribracteata, Klotzsch, ex Hanst. in Mart. Fl. Bras. viii. 356 = G. tribracteata.

RECTOMITRA, Blume, Mus. Bot. Lugd. Bat. i. 6 (1849) = **Kibessia**, DC. (Melastom.).
cordata, Triana, in Trans. Linn. Soc. xxviii. (1871) 153 = K. cordata.
galeata, Blume, Mus. Bot. Lugd. Bat. i. 6.—Borneo.
tuberculata, Blume, l. c. 7 = K. tuberculata.

REDIA, Casar. Nov. Stirp. Bras. Dec. 51 (1842) = **Cleidion**, Blume (Euphorb.).
tricocca, Casar. l. c. 52 = C. tricoccum.

REDOUTEA, Vent. Jard. Cels, t. 11 (1800) = **Fugosia**, Juss. (Malvac.).
heterophylla, Vent. l. c. = F. heterophylla.
tripartita, H. B. & K. Nov. Gen. et Sp. v. 293 = F. tripartita.

REDOWSKIA, Cham. & Schlecht. in Linnaea, i. (1826) 33. t. 2. *CRUCIFERAE*, Benth. & Hook. f. i. 92.
sophiaefolia, *Cham. & Schlecht. l. c.*—As. bor.

REDUTEA, Pers. Syn. ii. 254 (1807) = REDOUTEA, Vent. = **Fugosia**, Juss. (Malvac.).

REEDIA, F. Muell. Fragm. i. 239. t. 10 (1859). *CYPERACEAE*, Benth. & Hook. f. iii. 1067.
spathacea, *F. Muell. l. c.* i. 240. t. 10; ii. 182.—Austral.

REEVESIA, Lindl. in Brande, Journ. [II. ii.] (1827) 109, 112; Bot. Reg. t. 1236 (1829). *STERCULIACEAE*, Benth. & Hook. f. i. 219.
pubescens, *Mast. in Hook. f. Fl. Brit. Ind.* i. 364.—Reg. Himal.
thyrsoidea, *Lindl. in Brande, Journ.* [II. ii.] (1827) 112; *Bot. Reg.* t. 1236.—China.
Wallichii, *R. Br. in Benn. Pl. Jav. Rar.* 231, in adnot.—Reg. Himal.

REGELIA, Hort. ex H. Wendl. in Illustr. Hortic. xii. (1865) Misc. 6 = **Verschaffeltia**, H. Wendl. (Palm.).
magnifica, Hort. ex H. Wendl. l. c. = V. splendida.
majestica, Hort. ex Gard. Chron. (1865) 292 = V. splendida.
Princeps, Hort. ex Balf. f. in Baker, Fl. Maurit. 388 = V. splendida.

REGELIA, Lem. Illustr. Hortic. vii. (1860) sub t. 245, in nota = **Karatas**, Adans. (Bromel.).
caeruleum, Lem. l. c. (= *Nidularium caeruleum*).—Hab. ?
cruentum, Lem. l. c. = K. cruenta.
Innocentii, Lem. l. c. = K. Innocentii.
Meyendorffii, Lem. l. c. = K. Carolinae.

REGELIA, Schau. in Linnaea, xvii. (1843) 243. *MYRTACEAE*, Benth. & Hook. f. i. 706.
adpressa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 25 = Beaufortia micrantha.
ciliata, *Schau. in Linnaea*, xvii. (1843) 244.—Austral.
gibbosa, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 168 = inops.
globosa, Benth. Fl. Austral. iii. 171, sphalm. = praec.
grandiflora, *Benth. l. c.* 170.—Austral.
inops, *Schau. in Lehm. Pl. Preiss.* ii. 224.—Austral.

REGIA, Loud. ex C. DC. in DC. Prod. xvi. II. 136 (1864) = **Juglans**, Linn.
maxima, Loud. l. c. = J. regia.

REGMUS, Dulac, Fl. Hautes-Pyr. 328 (1867) = **Circaea**, Tourn. (Onagrar.).
alpinus, Dulac, l. c. = C. alpina.
lutetianus, Dulac, l. c. = C. lutetiana.

REGNALDIA, Baill. Adansonia, i. (1860–61) 187 = **Chaetocarpus**, Thw. (Euphorb.).
cluytioides, Baill. l. c. 188 = C. castanocarpus.

REGNELLIA, Rodrig. Orch. Nov. i. 81 (1877) = *ORCHIDEAE*.
purpurea, *Rodrig. l. c.* 79.—Bras.

REHMANNIA, Libosch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 36 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 960.
SPARMANNIA, Buc'hoz, Pl. Nouv. Découv. t. 1 (1779), nomen prius.
chinensis, *Libosch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 36.—China.
glutinosa, Steud. Nom. ed. II. ii. 440 = chinensis.
lutea, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 538.—Japon.
Piasezkii, *Maxim. l. c.* xxvi. (1880) 502.—China.

REICHARDIA, Dennst. Schluess. Hort. Malab. 32 (1818) = **Tabernaemontana**, Linn. (Apocyn.).
grandiflora, Dennst. l. c.—Ind. or.
jasminoides, Dennst. l. c.—Ind. or.

REICHARDIA, Roth, Bot. Abh. 35 (1787) = **Picridium**, Desf. (1799) (Compos.).
integrifolia, Moench, Meth. 546 = P. vulgare.
picroides, Roth, Bot. Abh. 35 = P. vulgare.
runcinata, Moench, Meth. 546 = P. tingitanum.
tingitana, Roth, Bot. Abh. 35 = P. tingitanum.

REICHARDIA, Roth, Catalect. ii. 64 (1800) = **Maurandia**, Orteg. (Scrophul.).
scandens, Roth, l. c. 65 = M. semperflorens.

REICHARDIA, Roth, Nov. Pl. Sp. 210 (1821) = **Pterolobium**, R. Br. (Legumin.).
decapetala, Roth, l. c. 212 = Caesalpinia sepiaria.
decapetala, Wight, ex Steud. Nom. ed. II. ii. 440 = P. lacerans.
hexapetala, Roth, Nov. Pl. Sp. 210 = P. indicum.
pentapetala, Blanco, Fl. Filip. ed. II. 233 = P. indicum.

REICHELEA, A. W. Benn. in Journ. Linn. Soc. xi. (1871) 271, sphalm. = seq.

REICHELIA, Schreb. Gen. 200 (1789) = **Hydrolea**, Linn. (Hydroph.).
angustifolia, J. F. Gmel. Syst. 1613, sphalm. = Phyllanthus angustifolius.

REICHELIA:—
guianensis, Spreng. Syst. i. 940 = Hydrolea caroliniana.
montevidensis, Spreng. l. c. = H. montevidensis.
palustris, Bilb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 141 = H. spinosa.
palustris, Blanco, Fl. Filip. ed. I. 220 = Sphenoclea zeylanica.
palustris, J. F. Gmel. Syst. 499 = H. caroliniana.

REICHEMBACHANTHUS, Rodrig. Orch. Nov. ii. 164 (1881). *ORCHIDEAE*.
modestus, *Rodrig. l. c.* 165.—Bras.

REICHENBACHIA, Spreng. in Bull. Soc. Philom. (1823) 54. t. 1. *NYCTAGINEAE*, Benth. & Hook. f. iii. 8.
caniflora, Mart. in Flora, xx. (1837) II. Beibl. 90 = Leucaster caniflorus.
hirsuta, *Spreng. in Bull. Soc. Philom.* (1823) 54. t. 1.—Bras.

REICHENHEIMIA, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 54. t. 4A = **Begonia**, Tourn.
Stelsneri, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiv. (1856) 204.—Hab. ?
subpeltata, Klotzsch, in Abh. Akad. Berl. (1855) 55 = B. subpeltata.
Thwaitesii, Klotzsch, l. c. = B. tenera.
zeylanica, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiv. (1856) 204 = B. zeylanica (in Addend.).

REICHERTIA, Karst. in Bot. Zeit. vi. (1848) 397 = **Schultesia**, Mart. (Gentian.).
gracilis, Karst. Fl. Columb. i. 60, in obs. = S. gracilis ?
linarifolia, Karst. in Bot. Zeit. vi. (1848) 397; Fl. Columb. i. 59 = S. stenophylla.
rosea, Karst. ll. cc. = S. stenophylla.

REIDIA, Wight, Ic. v. 27. tt. 1903, 1904 (1852) = **Phyllanthus**, Linn. (Euphorb.).
fimbriata, Wight, l. c. t. 1904 = P. fimbriatus.
floribunda, Wight, l. c. t. 1903 = P. Wightianus.
glabrescens, Miq. Fl. Ind. Bat. i. II. 374 = P. glabrescens.
glaucescens, Miq. l. c. = P. pulcher.
gracilipes, Miq. l. c. = P. gracilipes.
gracilis, Miq. l. c. 373 = P. gracilis.
latifolia, Wight, Ic. t. 1904 = P. macrocalyx.
ovalifolia, Wight, l. c. = P. longiflorus ?
polyphylla, Wight, l. c. = P. anabaptizatus.
singalensis, Miq. Fl. Ind. Bat. Suppl. 449 = P. singalensis.

REIFFERSCHEIDIA, Presl, Rel. Haenk. ii. 74. t. 62 (1836). *DILLENIACEAE*, Benth. & Hook. f. i. 13.
speciosa, *Presl, l. c.*—Am. austr.

REIFFERSCHIEDIA, Spach, Hist. Vég. Phan. vii. 413 (1839), sphalm. = praec.

REIGERA, Opiz, Seznam, 83 (1852) = **Scirpus**, Tourn. (Cyperac.).
maritima, Opiz, l. c. = S. maritimus.

REILIA, Steud. Syn. Pl. Cyp. 311 (1855) = **Eryngium** ?, Tourn. (Umbell.).—Cf. Benth. & Hook. iii. 863.
eryngioides, Steud. l. c.—Reg. Argent.

REIMARIA, Fluegge, Gram. Monog. 213 (1810), partim. *GRAMINEAE*, Benth. & Hook. f. iii. 1097.
aberrans, *Doell, in Mart. Fl. Bras.* ii. II. 38.—Am. trop.
acuta, *Fluegge, Gram. Monog.* 217.—Am. trop.
brasiliensis, *Schlecht. in Bot. Zeit.* x. (1852) 17.—Am. trop.
conferta, Nees, Agrost. Bras. 15; et in Mart. Fl. Bras. ii. II. 37 = acuta.
diffusa, Spreng. Neue Entdeck. iii. 14 = Zizania miliacea.
elegans, Fluegge, Gram. Monog. 216 = Paspalum pulchellum.
fascicularis, Link, Hort. Berol. ii. 203 = Panicum Schraderi.
laxa, Reichb. ex Spreng. f. Tent. Suppl. Syst. 2 = Panicum lanatum.

REIMARIA:—
oligostachya, *Munro, ex Benth. in Journ. Linn. Soc.* xix. (1881) 34.—Cuba; Florida.
puberula, Link, Hort. Berol. ii. 203 = Paspalum ischnocaulon.
purpurascens, Link, l. c. 203 = Panicum violascens?

REINECKIA, Karst. in C. Koch, Wochenschr. i. (1858) 349 = **Synechanthus**, H. Wendl. (Palm.).
triandra, Karst. l. c. = S. Warszewiczianus.

REINECKIA, Kunth, in Abh. Akad. Berl. 1842 (1844) 29. *LILIACEAE*, Benth. & Hook. f. iii. 771.
LIRIOPE, Salisb. Gen. Pl. Fragm. 74 (1866).
SANSEVIELLA, Reichb. Consp. 44 (1828), nomen prius.
carnea, *Kunth, in Abh. Akad. Berl.* 1842 (1844) 29.—China; Japon.

REINERA, Dennst. Schluess. Hort. Malab. 33 (1818) = **Curinila**, Roem. & Schult. (Asclep.?).
baypinensis, Dennst. l. c. = C. Rheedi.

REINERIA, Moench, Meth. Suppl. 44 (1802) = **Tephrosia**, Pers. (Legumin.).
reflexa, Moench, l. c. 45 = T. reflexa.

REINHARDTIA, Liebm. in Overs. Vidensk. Selsk. Forh. 1845 (1846) 9. *PALMAE*, Benth. & Hook. f. iii. 906.
elegans, *Liebm. l. c.*; *et ex Mart. Hist. Nat. Palm.* iii. 311.—Mexic.

REINIA, Franch. & Sav. Enum. Pl. Jap. ii. (1876) 314 = **Itea**, Gronov. (Saxifrag.).
racemosa, Franch. & Sav. l. c. = I. japonica.

REINWARDTIA, Blume, ex Nees, in Syll. Ratisb. i. (1824) 196 = **Saurauja**, Willd. (Ternstr.).
javanica, Blume, l. c. = S. Blumia.

REINWARDTIA, Dum. Comm. Bot. 19 (1822). *LINEAE*, Benth. & Hook. f. i. 243.
KITTELOCHARIS, Alef. in Bot. Zeit. xxi. (1863) 282.
MACROLINUM, Reichb. Handb. 306 (1827).
indica, Dum. Comm. Bot. 19 = trigyna.
repens, Planch. in Hook. Lond. Journ. Bot. vii. (1848) 523 = trigyna.
tetragyna, *Planch. l. c.*—Ind. or.
trigyna, Dalz. & Gibs. Bomb. Fl. 16 = tetragyna.
trigyna, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 522.—Ind. or.

REINWARDTIA, Korth. Verh. Nat. Gesch. Bot. 101. t. 12 (1840?) = **Ternstroemia**, Mutis.
elongata, Korth. l. c. 103 = T. micrantha.
patens, Korth. l. c. 102. t. 12 = T. micrantha.

REINWARDTIA, Spreng. Syst. i. 527 (1825) = **Breweria**, R. Br. (Convolv.).
ferruginea, Steud. Nom. ed. II. ii. 440 = B. ferruginea.
glabra, Spreng. Syst. i. 863.—N. Granat.
heterantha, Steud. Nom. ed. II. ii. 440.—Bras.
sericea, Spreng. Syst. i. 863.—N. Granat.
umbellata, Steud. Nom. ed. II. ii. 440.—Bras.

REISSEKIA, Endl. Gen. 1103 (1840). *RHAMNEAE*, Benth. & Hook. f. i. 386.
cordifolia, *Endl. l. c.*; *A. Gray, Bot. U. St. Expl. Exped.* i. 285.—Bras.
smilacina, Endl. l. c. = cordifolia.

REISSIPA, Steud. ex Klotzsch, in Lehm. Pl. Preiss. ii. 230 (1847) = **Monotaxis**, Brongn. (Euphorb.).
pleurandroides, Steud. ex Klotzsch, l. c. = M. grandiflora.

REJOUA, Gaudich. in Freyc. Voy. Bot. 450. t. 61 (1826) = **Tabernaemontana**, Plum. (Asclep.).
aurantiaca, Gaudich. l. c. 451. t. 61 = T. aurantiaca.
scandens, Seem. in Bonplandia, x. (1862) 296 = Carruthersia scandens.

RELBUNIUM, Benth. & Hook. f. Gen. ii. 149 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Aschenbornii, *Hemsl. Biol. Centr. Am. Bot.* ii. 62.—Mexic.
ciliatum, *Hemsl. l. c.*—Mexic.
hypocarpium, *Hemsl. l. c.* 63.—Mexic.
laevigatum, *Hemsl. l. c.*—Mexic.
microphyllum, *Hemsl. l. c.*—Mexic.
polyplocum, *Hemsl. Diagn. Pl. Nov.* 53. = Mexic.
tuberculosum, *Ball, in Journ. Linn. Soc.* xxii. (1885) 42.—Peruv.

RELCHELA, Steud. Syn. Pl. Gram. 101 (1855) = **Deyeuxia**, Clar. (Gramin.).
panicoides, Steud. l. c.—Chili.

RELHAMIA, J. F. Gmel. Syst. 247 (1791) = **Curtisia**, Ait. (Cornac.).
faginea, J. F. Gmel. l. c. = C. faginea.

RELHANIA, L'Hérit. Sert. Angl. 22. t. 29 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 326.
ECLOPES, Gaertn. Fruct. ii. 440. t. 169 (1791).
MICHAUXIA, Neck. Elem. i. 12 (1790).
RHYNCHOCARPUS, Less. Syn. Comp. 382 (1832).
RHYNCHOPSIDIUM, DC. in Mém. Soc. Phys. Genèv. vii. II. (1836) 283, 286.
acicularis, Desf. ex Steud. Nom. ed. II. ii. 440 = pungens.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 300.—Afr. austr.
apiculata, Harv. l. c. 303 = quinquenervis.
Bellidiastrum, *L'Hérit. Sert. Angl.* 24.—Afr. austr.
Bellidiastrum, Poir. ex DC. Prod. vi. 291 = Osmites Bellidiastrum.
calycina, L'Hérit. Sert. Angl. 24 = Peyrousea calycina.
centauroides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 301.—Afr. austr.
cuneata, *L'Hérit. Sert. Angl.* 23.—Afr. austr.
decussata, L'Hérit. l. c. 24 = cuneata.
dichotoma, Willd. ex Less. Syn. Comp. 372 = Nestlera humilis.
dumosa, E. Mey. ex DC. Prod. vi. 284 = Nestlera conferta.
ericoides, *Cass. in Dict. Sc. Nat.* xlv. 30.—Afr. austr.
faginea, Steud. Nom. ed. II. ii. 440, sphalm. = Curtisia faginea.
genistaefolia, *L'Hérit. Sert. Angl.* 22.—Afr. austr.
lateriflora, L'Hérit. l. c. 23 = sessiliflora.
lateriflora, Link, Enum. Hort. Berol. ii. 340 = quinquenervis.
laxa, *L'Hérit. Sert. Angl.* 23.—Afr. austr.
linifolia, Willd. ex Less. Syn. Comp. 371 = Nestlera humilis.
microphylla, *L'Hérit. Sert. Angl.* 22.—Afr. austr.
multipunctata, *DC. Prod.* vi. 286.—Afr. austr.
paleacea, L'Hérit. Sert. Angl. 24 = ericoides.
passerinoides, L'Hérit. l. c. 23 = Geigeria passerinoides.
pedunculata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 302.—Afr. austr.
pedunculata, *L'Hérit. Sert. Angl.* 23.—Afr. austr.
pinnata, Thunb. Prod. Pl. Cap. 146 = Ursinia filiformis.
pumila, Thunb. l. c. = pedunculata, *L'Hérit.*
pungens, *L'Hérit. Sert. Angl.* 24.—Afr. austr.
quinquenervis, *Thunb. Prod. Pl. Cap.* 146.—Afr. austr.
recurva, DC. Prod. vi. 286 = squarrosa.
recurva, *Less. Syn. Comp.* 376.—Afr. austr.
reflexa, Thunb. Prod. Pl. Cap. 146 = Nestlera biennis.
rotundifolia, *Less. Syn. Comp.* 375.—Afr. austr.
santolinoides, L'Hérit. Sert. Angl. 24 = ericoides.
sedifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 301.—Afr. austr.
sessiliflora, *Thunb. Prod. Pl. Cap.* 146.—Afr. austr.
speciosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 302.—Afr. austr.
squarrosa, *L'Hérit. Sert. Angl.* 22.—Afr. austr.
styphelioides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 303.—Afr. austr.
tomentosa, Thunb. Prod. Pl. Cap. 146 = Amphiglossa tomentosa.
triflora, E. Mey. ex DC. Prod. vi. 285 = Nestlera tricephala.
trinervis, *Thunb. Prod. Pl. Cap.* 146.—Afr. austr.
virgata, *L'Hérit. Sert. Angl.* 23.—Afr. austr.
viscosa, *L'Hérit. l. c.* 22.—Afr. austr.

RELLESTA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 337 = **Swertia**, Linn. (Gentian.).
cyanea, Turcz. l. c. = S. tetrapetala.

REMACLEA, C. Morr. Belg. Hortic. iii. (1853) 1. t. 1 = **Trimezia**, Salisb. (Irid.).
funebris, C. Morr. l. c. 3. f. 1 = T. lurida.

REMBERTIA, Adans. Fam. ii. 226 (1763) = **Diapensia**, Linn.

REME, Adans. Fam. ii. 245 (1763) = **Trianthema**, Sauv. (Ficoid.).

REMIJIA, DC. in Bibl. Univ. Genèv. (1829) ii. 185. *RUBIACEAE*, Benth. & Hook. f. ii. 33.
MACROCNEUMUM, Vand. Fl. Lusit. et Bras. 14 (1788).
angatensis, Blanco, Fl. Filip. ed. II. 115 = Randia angatensis.
Bergeniana, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 13.—Bras.
cujabensis, *Wedd. l. c.*—Bras.
densiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 215.—Guian.
ferruginea, *DC. Prod.* iv. 357.—Bras.
firmula, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 13.—Bras.
Hilarii, *DC. Prod.* iv. 357.—Bras.
lacunosa, *Spruce, in Journ. Linn. Soc.* v. (1861) 12.—Bras.
Lambertiana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 13 = Cascarilla Lambertiana.
Macrocnemia, *Wedd. l. c.*—Bras.
macrophylla, *Benth. & Hook. f. ex Flueck. Chinarind.* 16.—N. Granat.
obscura, Blanco, Fl. Filip. ed. II. 116 (*oscura*) = Gardenia obscura.
odorata, Blanco, l. c. 115 (*olorata*) = Randia densiflora.
paniculata, *DC. Prod.* iv. 357.—Bras.
pedunculata, *Flueck. Chinarind.* 17. t. 6.—N. Granat.
prismatostylis, *Benth. & Hook. f. ex Flueck. l. c.* 16.—N. Granat.
Purdieana, *Wedd. in Ann. Sc. Nat.* Sér. III. ix. (1849) 272.—N. Granat.
Riveroana, Flueck. Chinarind. 17 = Cascarilla Riveroana.
tenuiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 214.—Guian.
Vellozii, *DC. Prod.* iv. 357.—Bras.

REMIREA, Aubl. Pl. Gui. i. 44. t. 16 (1775). *CYPERACEAE*, Benth. & Hook. f. iii. 1058.
MEGIA, Schreb. Gen. 786 (1791).
diffusa, Sieber, ex Steud. Syn. Pl. Gram. 194 = Arundo occidentalis.
distichophylla, Boeck. in Flora, xli. (1858) 410 = maritima.
maritima, *Aubl. Pl. Gui.* i. 45. t. 16.—Reg. trop.
pedunculata, R. Br. Prod. 236 = maritima.
rigidissima, Steud. Syn. Pl. Cyp. 317 = maritima.
Wightiana, Nees, in Wight, Contrib. 92 = maritima.

REMUSATIA, Schott, Melet. i. 18 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 974.
Hookeriana, *Schott, in Oestr. Bot. Wochenbl.* viii. (1858) 133.—Reg. Himal.
vivipara, *Schott, Melet.* i. 18.—Ind. or.; Burma.

REMYA, Hillebr. ex Benth. & Hook. f. Gen. ii. pars 1ma, Add. 536 (1873); 1231 (1876). *COMPOSITAE*.
mauiensis, *Hillebr.* (*in Herb. Kew.*).—Ins. Sandvic.

RENANTHERA, Lour. Fl. Cochinch. 521 (1790). *ORCHIDEAE*, Benth. & Hook. f. iii. 577.
NEPHRANTHERA, Hassk. Cat. Hort. Bog. Alt. 44 (1844).
amabilis, Boxall, ex Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 241 (Quid?).—Ins. Philipp.
Arachnites, Lindl. Gen. et Sp. Orch. 217 = Arachnanthe moschifera.
bilinguis, Reichb. f. Xenia Orch. i. 7. t. 4 = Arachnanthe bilinguis.
coccinea, *Lour. Fl. Cochinch.* 521.—China.
elongata, *Lindl. Gen. et Sp. Orch.* 218.—Malaya.
Flosaëris, Reichb. f. Xenia Orch. i. 88 = Arachnanthe moschifera.

RENANTHERA:—
histrionica, *Reichb. f. in Gard. Chron.* (1878) II. 74.—Malaya.
Hookeriana, *Reichb. f. Xenia Orch.* ii. 42.—Borneo.
labrosa, Reichb. f. l. c. i. 88 = Arachnanthe bilinguis.
Lowei, *Reichb. f. l. c.* 89.—Borneo.
matutina, Lindl. Bot. Reg. (1843) t. 41 = elongata.
matutina, *Lindl. Gen. et Sp. Orch.* 218; *Blume, Mus. Bot. Lugd. Bat.* i. 61.—Java.
micrantha, Blume, Rumphia, iv. 54; Mus. Bot. Lugd. Bat. i. 60 = elongata.
moluccana, *Blume, Rumphia*, iv. 54.—Ins. Amboin.
Rohaniana, Reichb. f. Xenia Orch. i. 89 = Hookeriana.
Storiei, *Reichb. f. in Gard. Chron.* (1880) II. 296.—Ins. Philipp.
Sulingi, *Lindl. Gen. et Sp. Orch.* 217.—Java.

RENARDA, Regel, in Trautv. Regel, Maxim. & Winkl. Decas Pl. Nov. 5 (1882); et in Act. Hort. Petrop. viii. (1883) 277. *UMBELLIFERAE.*
siifolia, *Regel, ll. cc.* 6; 278.—Turkestan.

RENARDIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 466 = **Trimeria**, Harv. (Bixin.).
leiocarpa, Turcz. l. c.—Afr. austr.

RENEALMIA, R. Br. Prod. Add. [591] (1810) = **Libertia**, Spreng. (Irid.).
grandiflora, R. Br. l. c. [592] = L. grandiflora.
ixioides, Ker-Gawl. Gen. Irid. 27 = L. ixioides.
paniculata, R. Br. Prod. Add. [592] = L. paniculata.
pulchella, R. Br. l. c. = L. pulchella.

RENEALMIA, Houtt. Handleid. viii. 335. t. 47. f. i. (1777) = **Villarsia**, Vent. (Gentian.).
capensis, Houtt. l. c. = V. ovata.

RENEALMIA, Linn. Gen. ed. I. 340 (1737) = **Tillandsia**, Linn. (Bromel.).
disticha, Linn. Syst. ed. X. 974 = T. heliconioides.
monostachia, Linn. Sp. Pl. 287 = T. tenuifolia.
paniculata, Linn. l. c. 286 = T. paniculata.
pendula, Gaertn. f. Fruct. iii. 13 t. 102 = T. pendula.
polystachia, Linn. Sp. Pl. 286 = T. polystachia.
recurvata, Linn. l. c. 287 = T. recurvata.
usneoides, Linn. l. c. = T. usneoides.

RENEALMIA, Linn. f. Suppl. 7 (1781). *SCITAMINEAE*, Benth. & Hook. f. iii. 648.
ETHANIUM, Salisb. in Trans. Hort. Soc. i. (1812) 281, 282.
GETHYRA, Salisb. l. c. (1812).
PEPERIDIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
SIPHOTRIA, Rafin. Fl. Tellur. iv. 50 (1836).
africana, *Benth. in Hook. Ic. Pl.* t. 1430.—Afr. trop.
amoena, *A. Rich. Fl. Cub. Fanerog.* ii. 254.—Cuba.
aromatica, Griseb. Fl. Brit. W. Ind. 601 = domingensis.
bracteosa, *Griseb. l. c.*—Ind. occ.
breviscapa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 25. t. 135.—Peruv.
calcarata, Andr. Bot. Rep. t. 421 = Alpinia calcarata.
Cardamomum-medium, Sweet, Hort. Brit. ed. II. 493, nomen = Amomum costatum.
caribaea, Griseb. Fl. Brit. W. Ind. 601 = racemosa.
concolor, D. Dietr. Syn. Pl. i. 14, sphalm. = Thalia concolor.
domingensis, Horan. Prod. Scitam. 32 = occidentalis.
erecta, D. Dietr. Syn. Pl. i. 14, sphalm. = Thalia erecta.
exaltata, Blanco, Fl. Filip. ed. II. 1 = Alpinia gracilis.
exaltata, *Linn. f. Suppl.* 7.—Ind. occ.
fasciculata, Rosc. Scit. 14. t. 109 = fasciculatum.
geniculata, D. Dietr. Syn. Pl. i. 14, sphalm. = Thalia geniculata.
gracilis, Blanco, Fl. Filip. ed. I. 1 = Alpinia gracilis.
jamaicensis, Horan. Prod. Scitam. 32 = occidentalis.
macrantha, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 25. t. 134.—Peruv.
marantaefolia, D. Dietr. Syn. Pl. i. 14, sphalm. = Thalia marantaefolia.
minor, [Desf.] Hort. Par. ex Roem. & Schult. Syst. i. 32 = Alpinia calcarata.
monosperma, *Miq. in Linnaea*, xxii. (1849) 80.—Guian.

RENEALMIA:—
mutica, Salisb. in Trans. Hort. Soc. i. (1812) 280 = Alpinia mutica.
nutans, Andr. Bot. Rep. t. 360 = Alpinia nutans.
occidentalis, *Sweet, Hort. Brit.* ed. II. 493; *Griseb. in Goett. Abh.* vii. (1857) 274.—Ind. occ.
Paco-Secora, *Horan. Prod. Scitam.* 32.—Ins. S. Vincent.
pauciflora, Griseb. Cat. Pl. Cub. 256, in syn. = amoena.
polystachia, *Jacq. Select. Am.* 93.—Cuba.
punicea, *Sweet, Hort. Brit* ed. II. 493.—Ind. or.
racemosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 26 = breviscapa.
racemosa, *A. Rich. Fl. Cub. Fanerog.* ii. 254; *Griseb. Fl. Brit. W. Ind.* 601.—Am. trop.
Ruiziana, Horan. Prod. Scitam. 33 = breviscapa.
strobilifera, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 25. t. 135.—Peruv.
sylvestris, *Horan. Prod. Scitam.* 32.—Ind. occ.
thyrsoidea, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 25. t. 134.—Peruv.
tuberosa, D. Dietr. Syn. Pl. i. 14 = Calathea tuberosa.
ventricosa, *Griseb. Cat. Pl. Cub.* 256.—Cuba.

RENGGERIA, Meissn. Gen. 42; Comm. 31 (1837). *GUTTIFERAE*, Benth. & Hook. f. i. 171.
SCHWEIGGERA, Mart. Nov. Gen. et Sp. iii. 166. t. 297 (1829).
acuminata, Seem. Bot. Voy. Herald, 88 = Clusia acuminata.
comata, *Meissn. ex Walp. Rep.* i. 393.—Bras.
guianensis, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 104.—Guian.
littoralis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 11.—Bras.
montana, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1093, nomen.—Guian.

RENGIFA, Poepp. & Endl. Nov. Gen. et Sp. iii. 12. t. 210 (1842). *GUTTIFERAE*, Benth. & Hook. f. i. 171.
acuminata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 243.—Guian.
peruviana, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 12.—Peruv.
scandens, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 241.—Guian.

RENIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Inocarpus**, Forst. (Legumin.).

RENNELLIA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 255. *RUBIACEAE*, Benth. & Hook. f. ii. 118.
elliptica, *Korth. l. c.* 257.—Sumatra.
ovalis, *Korth. l. c.*—Sumatra.
speciosa, *Benth. & Hook. f. Gen.* ii. 118.—Burma.

RENSCHIA, Vatke, in Linnaea, xliii. (1880-82) 94 = **Tinnea**, Kotschy & Peyr. (Labiat.).
heterotypica, Vatke, l. c. = T. heterotypica.

RENSSELAERIA, Beck, Bot. N. Mid. St. 382 (1833) = **Peltandra**, Rafin. (Aroid.).
virginica, Beck, l. c. = P. virginica.

REPANDRA, Lindl. Orch. Scel. 12 (1826) = **Disa**, Berg. (Orchid.).

REPHESIS, Rafin. Sylva Tellur. 59 (1838) = **Ficus**, Tourn. (Urtic.).
calyptrata, Rafin. l. c. = F. calyptrata.
ovata, Rafin. l. c. = F. ovata.

REPTONIA, A. DC. in DC. Prod. viii. 153 (1844). *MYRSINEAE*, Benth. & Hook. f. ii. 648.
EDGEWORTHIA, Falc. in Proc. Linn. Soc. i. (1842) 129.
MONOTHECA, A. DC. in DC. Prod. viii. 152 (1844).
buxifolia, *A. DC. in DC. Prod.* viii. 153.—Ind. or.
laurina, Benth. Fl. Hongk. 208 = Sarcosperma laurina.

REQUIENIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 91 = **Tephrosia**, Pers. (Legumin.).
obcordata, DC. l. c.; Lég. Mém. vi. t. 37 = T. obcordata.
sphaerosperma, DC. ll. cc. 91; . 38 = T. sphaerosperma.

RESEDA, Tourn. ex Linn. Syst. ed. I (1735). *RESEDACEAE*, Benth. & Hook. f. i. 112.
ARKOPODA, Rafin. Fl. Tellur. iii. 73 (1836).
ERESEDA, Spach, Hist. Vég. Phan. vii. 101 (1839).
HEXASTYLIS, Rafin. Fl. Tellur. iii. 73 (1836).
PECTANISIA, Rafin. l. c. 72 (1836).
TEREIANTHES, Rafin. l. c. (1836).
abyssinica, Fresen. in Mus. Senckenb. ii. (1837) 106 = Caylusea abyssinica.
affinis, Kotschy, ex Muell. Arg. Monog. Resed. 124 = arabica.
alba, Delile, Fl. Aegypt. 15 = decursiva.
alba, *Linn. Sp. Pl.* 449.—Europ.; Oriens.
alba, St. Amans, Fl. Agen. 187 = Jacquinii.
alba, Ruiz, ex Muell. Arg. Monog. Resed. 107 = bipinnata.
alopecuros, *Boiss. Diagn.* Ser. I. viii. 55.—Palaestina.
Alphonsi, *Muell. Arg. in Bot. Zeit.* xiv. (1856) 35.—Algeria.
amblyocarpa, *Fresen. in Mus. Senckenb.* ii. (1837) 108.—Abyssin.
anomala, J. Gay, ex Muell. Arg. Monog. Resed. 124 = arabica.
arabica, *Boiss. Diagn.* Ser. I. i. 6.—Aegypt.; Arab.
aragonensis, Loscos & Pardo, Ser. Imperf. 51 = R. Phyteuma.
arborea, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 722 = odorata.
armena, *Boiss. Diagn.* Ser. I. i. 5.—Armenia.
atriplicifolia, J. Gay, ex Muell. Arg. Monog. Resed. 146 = Aucheri.
attenuata, *Ball, in Journ. Linn. Soc.* xvi. (1877) 338.—Marocc.
Aucheri, *Boiss. Diagn.* Ser. I. i. 5.—Mesopot.; Persia.
baetica, J. Gay, ex Muell. Monog. Resed. 109 = bipinnata.
Balansae, *Muell. Arg. l. c.* 119.—Cilicia.
Barrelieri, *Bertol. Fl. Ital.* v. 32.—Ital.; Hispan.
bipinnata, Loscos & Pardo, Ser. Imperf. 51 = macrostachya.
bipinnata, *Willd. Enum. Hort. Berol.* i. 499.—Hispan.
Boissieri, *Muell. Arg. in Bot. Zeit.* xiv. (1856) 37.—Aegypt.
bracteata, Boiss. Diagn. Ser. I. vi. 22 = pruinosa.
Buhseana, *Muell. Arg. in Bot. Zeit.* xiv. (1856) 36.—Beluchist.
Bungei, *Boiss. Fl. Orient.* i. 433.—Persia.
cahirana, Muell. Arg. Monog. Resed. 176 = Boissieri.
calicinalis, Lam. Fl. Fr. iii. 204 = R. Phyteuma.
canescens, Linn. Sp. Pl. 448 = Astrocarpus sesamoides.
canescens, *A. St. Hil. in Ann. Sc. Nat.* Sér. II. vii. (1837) 375.—Arab.
capensis, Burm. ex Muell. Arg. Monog. Resed. 222 = Astrocarpus Clusii.
capensis, Burm. f. Fl. Cap. Prod. 13 = Oligomeris capensis.
capensis, Willd. ex Muell. Arg. Monog. Resed. 214 = Oligomeris glaucescens.
caudata, DC. ex Muell. Arg. l. c. 154 = pruinosa.
chinensis, *Lour. Fl. Cochinch.* 299.—China.
clausa, Reichb. ex Muell. Arg. in Bot. Zeit. xiv. (1856) 39 = lutea.
Clusii, Spreng. Fl. Hal. Mant. 41 = Astrocarpus sesamoides.
cochinchinensis, *Lour. Fl. Cochinch.* 299.—Cochinch.
collina, J. Gay, in Expl. Sc. Algér. t. 71. f. 2 = R. Phyteuma.
complicata, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) 13.—Hispan.
confusa, Pomel, Nouv. Mat. Fl. Atl. 224 = R. Phyteuma.
constricta, Lange, Pl. Nov. Hisp. fasc. i. 6. t. 8 = lanceolata.
crispa, Mill. Gard. Dict. ed. VIII. n. 2 = lutea.
crispa, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 892 = alba.
crispata, *Bertol. Fl. Ital.* v. 26.—Sicil.
crispata, Link, Enum. Hort. Berol. ii. 8 = R. Luteola.
crystallina, *Webb & Berth. Phytog. Canar.* i. 102. t. 9.—Ins. Canar.
decursiva, *Forsk. Fl. Aegypt. Arab.* 67.—Arab.
difformis, Moench, Meth. Suppl. 22 = Caylusea canescens.
diffusa, *Ball, in Journ. Linn. Soc.* xvi. (1877) 339.—Marocc.
dipetala, Ait. Hort. Kew. ed. I. ii. 132 = Oligomeris capensis.
dipetala, Drège, ex Presl, Bot. Bemerk. 8 = Oligomeris Dregeana.

RESEDA:—
Dregeana, Presl, Bot. Bemerk. 8—Oligomeris Dregeana.
Duriaeana, *J. Gay, in Expl. Sc. Algér.* t. 71. f. 1; *et ex Muell. Arg. in Bot. Zeit.* xiv. (1856) 38.—Afr. bor.
elata, *Coss. & Bal. ex Muell. Arg. in DC. Prod.* xvi. II. 578; *et in Bull. Soc. Bot. Fr.* xx. (1873) 242.—Marocc.
erecta, Lag. Gen. et Sp. Nov. 17 = stricta.
eremophila, Boiss. Diagn. Ser. I. viii. 54 = decursiva.
filiformis, Pourr. ex Muell. Arg. Monog. Resed. 200 = virgata.
fruticosa, Ehrenb. ex Muell. Arg. l. c. 229 = Caylusea canescens.
fruticulosa, Linn. Syst. ed. X. 1046 = alba.
Gayana, Boiss. Voy. Bot. Espagne, ii. 76. t. 21 = alba.
gigantea, Pourr. ex Willk. & Lange, in Prod. Fl. Hisp. iii. 890 = suffruticosa.
glauca, Brot. Fl. Lusit. ii. 307 = virgata.
glauca, Eichw. Casp. Cauc. 24 = R. Luteola.
glauca, *Linn. Sp. Pl.* 449.—Hispan.
glauca, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Oligomeris capensis.
globulosa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 45.—Reg. Caucas.
gracilis, Tenore, Syll. 232 = lutea.
gredensis, *Muell. Arg. in DC. Prod.* xvi. II. 582; *Cutanda, Apend. Fl. Matrit.* 748.—Hispan.
Gussoni, Boiss. & Reut. Diagn. Ser. II. i. 49 = R. Luteola.
Haussknechtii, *Muell. Arg. in DC. Prod.* xvi. II. 574.—Persia.
hexagyna, Forsk. Fl. Aegypt. Arab. 92 = canescens.
Hookeri, Guss. Enum. Pl. Inar. 28 = alba.
incisa, Tenore, Prod. Fl. Nap. p. xxviii = alba.
inflata, Ehrenb. ex Muell. Arg. Monog. Resed. 189 = lutea.
inodora, *Reichb. Ic. Fl. Germ.* ii. 22. t. 99 f. 4445.—Europ. occ.
Jacquinii, *Reichb. l. c.*—Europ. austr.
kahirina, Boiss. Fl. Orient. i. 430 = Boissieri.
kurdica, *Boiss. & Noé, in Boiss. l. c.* 426.—Kurdist.
laciniata, Dulac, Fl. Hautes-Pyr. 183 = R. Phyteuma.
laevigata, *G. Don, Gen. Syst.* i. 289.—Aegypt.
Lamottei, Jord. ex Nym. Consp. 69 = lutea.
lanceolata, *Lag. Gen. et Sp. Nov.* 17.—Hispan.
Lancerotae, Webb, ex Delile, Ind. Sem. Hort. Monsp. (1836) 27 = crystallina.
leucantha, *Hegelm. ex Willk. & Lange, Prod. Fl. Hisp.* iii. 892.—Hispan.
linifolia, *Hornem. Hort. Hafn.* ii. 501.—Europ. austr.; Aegypt.
litorea, J. Gay, ex Nym. Consp. 69 = Jacquinii.
littoralis, Muell. Arg. Monog. Resed. 141 = Jacquinii.
longicalycina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Phyteuma.
lurida, *Muell. Arg. Monog. Resed.* 152.—Afr. or.
lusitanica, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 897 = alba.
lutea, *Linn. Sp. Pl.* 449.—Europ.; Oriens.
Luteola, *Linn. l. c.* 448.—Europ.; Oriens.
macrobotrys, *Boiss. Fl. Orient.* i. 433.—Persia.
macrocarpa, Ehrenb. ex Muell. Arg. Monog. Resed. 156 = stenostachya.
macrosperma, Reichb. in Flora, xiii. (1830) 130 = media.
macrostachya, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 890.—Hispan.
media, *Lag. Gen. et Sp. Nov.* 17.—Lusitan.
mediterranea, Bess. ex Boiss. Fl. Orient. i. 428 = inodora.
mediterranea, Jacq. Coll. i. 147 = Jacquinii.
mediterranea, Linn. ex Muell. Arg. Monog. Resed. 227 = Caylusea canescens.
mediterranea, *Linn. Sp. Pl.* ed. II. 564.—Afr. bor.
mediterranea, Sadler, Verz. 71 = inodora.
mediterranea, Schoen, ex Muell. Arg. Monog. Resed. 124 = arabica.
mediterranea, Willd. ex Muell. Arg. l. c. 136 = R. Phyteuma.
microcarpa, *Muell. Arg. in Bot. Zeit.* xiv. (1856) 35.—Persia.
microphylla, Presl, Bot. Bemerk. 8 = Oligomeris capensis.
mucronata, Tineo, Cat. Pl. Hort. Panorm. (1827) 280 = lutea.
mucronulata, Guss. Fl. Sic. Prod. Suppl. 147 = lutea.
muricata, *Presl, Bot. Bemerk.* 8.—Palaest.; Arab.
myriophylla, Tenore, Fl. Neap. App. i. 28 = alba.
neglecta, Muell. Arg. Monog. Resed. 178 = lutea.

RESEDA:—
nilgherrensis, Muell. Arg. in Bot. Zeit. xiv. (1856) 34 = odorata.
ochracea, Moench, Meth. 58 = alba.
Ochradeni, *Boiss. Diagn.* Ser. II. i. 49.—Persia.
odorata, Gueldenst. ex Ledeb. Fl. Ross. i. 236 = R. Phyteuma.
odorata, *Linn. Syst.* ed. X. 1046.—Afr. bor.?
oligandra, Edgew. in Journ. As. Soc. Beng. vii. (1838) II. 764 = Oligomeris glaucescens.
orientalis, *Boiss. Fl. Orient.* i. 427.—Syria.
orthostyla, C. Koch, in Linnaea, xv. (1845) 705 = lutea.
papillosa, Muell. Arg. in Bot. Zeit. xiv. (1856) 38 = Duriaeana.
parviflora, DC. ex Muell. Arg. Monog. Resed. 114 = alba.
pedunculata, R. Br. in Salt, Abyss. App. 64 = Caylusea abyssinica.
Phiteuma, Neck. Delic. Gallo-Belg. i. 210 = lutea.
Phyteuma, Brot. Fl. Lusit. ii. 305 = media.
Phyteuma, Forsk. Fl. Aegypt. Arab. p. lxvii = pruinosa.
Phyteuma, *Linn. Sp. Pl.* 449.—Reg. Mediterr.; As. Min.
Phyteuma, Kralik, ex Muell. Arg. Monog. Resed. 232 = arabica.
pilosa, Link, ex Muell. Arg. l. c. 135 = media.
podocarpos, Viv. Pl. Aegypt. Dec. 7 = Caylusea canescens.
podolica, *Rehm. Notiz. Veg. Schwarz. Meer.* 55.—Podolia.
praetervisa, Muell. Arg. Monog. Resed. 232 = arabica.
propinqua, R. Br. in Denh. & Clapp. Trav. App. 227 = decursiva.
pruinosa, Bové, ex Muell. Arg. Monog. Resed. 156 = stenostachya.
pruinosa, *Delile, Fl. Aegypt. Illustr.* 63.—Aegypt.
pruinosa, Schimp. ex Muell. Arg. Monog. Resed. 159 = muricata.
Pseudo-phyteuma, Brot. ex Nym. Consp. 68 = R. Phyteuma.
Pseudo-virens, Frivald. in Flora, xx. (1837) 232 = R. Luteola.
purpurascens, Linn. Sp. Pl. 449 = Astrocarpus Clusii.
pygmaea, Scheele, in Flora, xxvi. (1843) 426 = Astrocarpus sesamoides.
quadragyna, Walp. Rep. ii. 464 = *tetragyna*.
Quartiniana, Buching. ex Muell. Arg. Monog. Resed. 152 = lurida.
Quartiniana, A. Rich. Tent. Fl. Abyss. i. 13 = pruinosa.
ramosissima, G. Don, ex Kunze, in Linnaea, xxiv. (1851) 163 = stricta.
ramosissima, *Pourr. ex Willd. Enum. Hort. Berol.* 499; *Spreng. Syst.* ii. 464.—Hispan.
Reuteriana, *Muell. Arg. in Bot. Zeit.* xiv. (1856) 37.—Alger.
rotundifolia, Kotschy, ex Muell. Arg. Monog. Resed. 147 = Aucheri.
salicifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 666 = R. Luteola.
saxatilis, Bal. ex Muell. Arg. Monog. Resed. 165 = Reuteriana.
saxatilis, Pourr. ex Willd. Enum. Hort. Berol. 500 = stricta.
Schimperi, Presl, Bot. Bemerk. 8 = Oligomeris glaucescens.
Schraderi, *Boiss. Diagn.* Ser. II. i. 49.—Persia.
scoparia, *Brouss. ex Willd. Enum. Hort. Berol.* i. 499.—Ins. Mader.
sesamoides, Linn. Sp. Pl. 449 = Astrocarpus sesamoides.
sesamoides, Sieber, ex Muell. Arg. Monog. Resed. 221. = Astrocarpus sesamoides.
spartioides, Sieber, ex Muell. Arg. l. c. 94 = Ochradenus baccatus.
spathulata, *E. Mey. ex Muell. Arg. l. c.* 211 = Oligomeris spathulata.
spicata, Humb. ex Muell. Arg. l. c. 136 = R. Phyteuma.
stellata, Lam. Fl. Fr. iii. 204 = Astrocarpus sesamoides.
stenostachya, *Boiss. Diagn.* Ser. I. i. 5.—Anatolia.
stipulacea, DC. ex Muell. Arg. Monog. Resed. 200 = virgata.
stricta, Kralik, ex Muell. Arg. l. c. 187 = lutea.
stricta, *Pers. Syn.* ii. 10.—Algeria.
subulata, Delile, Fl. Aegypt. 15 = Oligomeris glaucescens.
suffruticosa, *Loefl. It.* ed. Germ. 79; *ex Willk. & Lange, Prod. Fl. Hisp.* iii. 890 = alba.
suffruticulosa, Link, ex Muell. Arg. Monog. Resed. 189 = lutea.
suffruticulosa, Linn. Sp. Pl. ed. II. 645 = alba.

RESEDA:—
tenuifolia, Wallr. Beitr. Fl. Hercyn. 204; et in Linnaea, xiv. (1840) 574 = lutea.
tetragyna, Forsk. Fl. Aegypt. Arab. 92 = arabica.
tinctoria, Salisb. Prod. 345 = R. Luteola.
tomentosa, *Boiss. Diagn.* Ser. I. i. 4.—Cappadoc.
tomentosa, Coss. in Bull. Soc. Bot. Fr. vi. (1859) 392 = villosa.
Tournefortii, Schult. Obs. Bot. 89 = R. Phyteuma.
tricuspis, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 241.—Marocc.
tridens, Viv. Pl. Aegypt. Dec. 6 = Oligomeris glaucescens.
tripartita, Buhse, ex Muell. Arg. Monog. Resed. 161 = Buhseana.
truncata, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 45.—Macedon.; Anatol.
truncula, Ehrenb. ex Muell. Arg. Monog. Resed. 191 = lutea.
undata, Amo, Fl. Iber. vi. 322 = suffruticosa.
undata, Boiss. Voy. Bot. Espagne, ii. 75. t. 20 = Barrelieri.
undata, Ehrenb. ex Muell. Arg. Monog. Resed. 109 = decursiva.
undata, Habl. Phys. Desc. Taur. 146 = lutea.
undata, Linn. Syst. ed. X. 1046 = alba.
undulata, Gilib. Fl. Lituan. ii. 210 = R. Luteola.
undulata, Link, Handb. ii. 518, sphalm. = *undata*.
urnigera, *Webb, Fragm. Fl. Aethiop.* 25.—Afr. bor.
villosa, *Coss. in Bull. Soc. Bot. Fr.* vi. (1859) 392.—Algeria.
virescens, *Hornem. Hort. Hafn.* ii. 501.—Hispan.
virgata, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 6.—Lusitan.
viridis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 501.—Ins. Socotr.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = lutea.
Yannesii, Cutanda, Fl. Comp. Madr. 151 = lutea.

RESEDELLA, Webb & Berth. Phyt. Canar. i. 106 (1836) = **Oligomeris**, Cambess. (Resed.).
dipetala, Webb & Berth. l. c. 107 = O. glaucescens.
Dregeana, Muell. Arg. in Bot. Zeit. xiv. (1856) 39 = O. Dregeana.
subulata, Webb & Berth. Phyt. Canar. i. 107. t. 11 = O. glaucescens.

RESINARIA, Comm. ex Lam. Encyc. i. 349 (1784) = **Terminalia**, Linn. (Combret.).

RESTIARIA, Lour. Fl. Cochinch. 639 (1790). *INC. SEDIS.*
cordata, *Lour. l. c.*—China.

RESTIO, Linn. Syst. ed. XII. 725 (1767). *RESTIACEAE*, Benth. & Hook. f. iii. 1031.
BALOSKION, Rafin. Fl. Tellur. iv. 32 (1836).
CHONDROPETALON, Rafin. l. c. 33 (1836).
CRASPEDOLEPIS, Steud. Syn. Pl. Cyp. 264 (1855).
ISCHYROLEPIS, Steud. in Flora, xxiii. (1850) 229, nomen; Syn. Pl. Cyp. 249 (1855).
LEIENA, Rafin. Fl. Tellur. iv. 33 (1836).
MEGALOTHECA, F. Muell. Fragm. viii. 98 (1873).
RHODOCOMA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836).
abortivus, *Nees, in Lehm. Pl. Preiss.* ii. 60.—Austral.
acuminatus, Thunb. Diss. Rest. 13 = Cannamois simplex.
alboaristatus, Nees, in Linnaea, v. (1830) 635 = Hypodiscus duplicatus.
amblycoleus, *F. Muell. Fragm.* viii. 65.—Austral.
ameles, Steud. Syn. Pl. Cyp. 252 = ferruginosus.
anceps, Burch. ex Mast. in Journ. Linn. Soc. x. (1869) 267 = Hypolaena anceps.
applanatus, *Spreng. Syst.* i. 185.—Austral.
arcuatus, *Mast. in DC. Monog. Phan.* i. 247.—Afr. austr.
argenteus, Nees, in Linnaea, v. (1830) 656 = Ceratocaryum argenteum.
argenteus, Thunb. Diss. Rest. 14 = Thamnochortus argenteus.
aristatus, Thunb. l. c. 10. f. 4 = Hypodiscus aristatus.
articulatus, Retz. Obs. iv. 14 = Lepironia mucronata.
asperiflorus, Nees, in Linnaea, v. (1830) 656 = Elegia asperiflora.
australis, *R. Br. Prod.* 245.—Austral.
bifarius, *Mast. in Journ. Linn. Soc.* x. (1869) 278.—Afr. austr.

RESTIO:—
bifidus, Auct. ex Mast. in DC. Monog. Phan. i. 280 = Garnotianus.
bifidus, *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 7.—Afr. austr.
bifurcus, Mast. in Journ. Linn. Soc. viii. (1865) 247, partim = furcatus.
bifurcus, *Nees, ex Mast. l. c.*—Afr. austr.
bigeminus, *Nees, ex Mast. l. c.* 246.—Afr. austr.
brizoides, Steud. Syn. Pl. Cyp. 255 = tremulus.
Brownii, Kunth, Enum. Pl. iii. 417 = Leptocarpus aristatus.
callistachyus, Kunth, l. c. 400 = Mastersi.
capillaris, *Kunth, l. c.* 405.—Afr. austr.
caricinus, Mass. ex Mast. in DC. Monog. Phan. i. 327 = Thamnochortus caricinus.
casuarinaeformis, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 227 = subverticillatus.
Cephalotes, Kunth, Enum. Pl. iii. 448, sphalm. = Cannomois Cephalotes.
cernuus, Linn. f. Suppl. 425; Thunb. Diss. Rest. 10. f. 2 = Thamnochortus cernuus.
chasmatocoleus, F. Muell. Fragm. viii. 71 = laxus.
chaunocoleus, *F. Muell. l. c.* 64.—Austral.
Chondropetalum, Nees, in Linnaea, v. (1830) 652 = Elegia deusta.
cincinnatus, *Mast. in Journ. Linn. Soc.* viii. (1865) 240.—Afr. austr.
cinerascens, *R. Br. Prod.* 246.—Austral.
cinerascens, Nees, ex Mast. in DC. Monog. Phan. i. 335 = Leptocarpus tenax.
clavatus, R. Br. 246 = Hypolaena fastigiata.
clavatus, Nees, in Lehm. Pl. Preiss. ii. 58 = Hypolaena exsulca.
complanatus, *R. Br. Prod.* 245.—Austral.
compressus, R. Br. l. c. = applanatus.
compressus, *Rottb. Desc. et Ic.* 6. t. 2 f. 4.—Afr. austr.
concolor, *Steud. Syn. Pl. Cyp.* 251.—Afr. austr.
confertospicatus, *Steud. l. c.* 256.—Austral.
confinis, Steud. l. c. = ornatus.
crinalis, *Mast. in Journ. Linn. Soc.* viii. (1865) 229.—Afr. austr.
crispatus, *R. Br. Prod.* 246.—Austral.
crispatus, Nees, ex Benth. Fl. Austral. vii. 421 = Caustis flexuosa.
crispatus, Nees, ex Benth. l. c. 238 = Hypolaena lateriflora.
curviramis, *Kunth, Enum. Pl.* iii. 395.—Afr. austr.
curvulus, *Nees, in Ann. & Mag. Nat. Hist.* Ser. I. vi. (1841) 50.—Austral.
cuspidatus, *Thunb. in Hoffm. Phytog. Blaetter*, i. 8.—Afr. austr.
debilis, *Nees, in Linnaea*, v. (1830) 641.—Afr. austr.
deformis, *R. Br. Prod.* 245.—Austral.
deformis, Steud. Syn. Pl. Cyp. 255 = leptocarpoides.
depauperatus, *Kunth, Enum. Pl.* iii. 405.—Afr. austr.
dichotomus, Gaertn. Fruct. ii. 12 = Willdenowia teres.
dichotomus, Linn. Syst. ed. XII. 735 = Thamnochortus fruticosus.
dichotomus, Rottb. Desc. 2; Progr. 10 = Thamnochortus dichotomus.
dichotomus, Thunb. Diss. Rest. 21 = triflorus, triticeus.
dichotomus, Willd. ex Kunth, Enum. Pl. iii. 406 = perplexus.
diffusus, Spreng. Syst. i. 185 = Leptocarpus tenax.
digitatus, Nees, in Linnaea, v. (1830) 638 = obtusissimus.
digitatus, *Thunb. Diss. Rest.* 18.—Afr. austr.
digitatus, Thunb, ex Mast. in Journ. Linn. Soc. xiv. (1875) 417 = Cannamois Cephalotes.
digitatus, Zeyh. ex Kunth, Enum. Pl. iii. 456 = Willdenowia striata.
digynus, Link, ex Mast. in Journ. Linn. Soc. viii. (1865) 250 = Mastersi.
dimorphus, *R. Br. Prod.* 246.—Austral.
dispar, *Mast. in Journ. Linn. Soc.* viii. (1865) 246.—Afr austr.
distachyos, Rottb. Desc. 8; Progr. 11 = Thamnochortus umbellatus.
distichus, Rottb. ll. cc. 6; 11 = Thamnochortus distichus.
divaricatus, Mast. in Journ. Linn. Soc. viii. (1865) 236 = macer.
dubius, Klotzsch, ex Mast. l. c. 250 = Mastersi.
echinatus, Kunth, Enum. Pl. iii. 384 = squarrosus.
Ecklonii, *Mast. in Journ. Linn. Soc.* viii. (1865) 236.—Afr. austr.
egregius, *Hochst. in Flora*, xxviii. (1845) 337.—Afr. austr.

RESTIO:—
elatus, *Mast. in Journ. Linn. Soc.* viii. (1865) 226.—Afr. austr.
elegans, Poir. Encyc. vi. 171 = Cannomois Cephalotes.
Elegia, Murr. Syst. ed. XIII. 738 = Elegia juncea.
Eleocharis, *Nees, ex Mast. in Journ. Linn. Soc.* viii. (1865) 238.—Afr. austr.
elongatus, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 7 = Thamnochortus elongatus.
equisetaceus, Reichb. ex Mast. in Journ. Linn. Soc. x. (1869) 250 = Dovea macrocarpa.
erectus, Nees, ex Mast. l. c. 243 = Elegia juncea.
erectus, Nees, in Linnaea, v. (1830) 655 = Elegia propinqua.
erectus, Thunb. Diss. Rest. 14 = Thamnochortus erectus.
eriophorus, Reichb. ex Mast. in Journ. Linn. Soc. x. (1869) 229 = Thamnochortus fruticosus.
fasciculatus, R. Br. Prod. 247 = Hypolaena fasciculata.
fastigiatus, *R. Br. Prod.* 246.—Austral.
fastigiatus, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 250 = Mastersi.
ferruginosus, Kunth, ex Mast. in DC. Monog. Phan. i. 248 = subverticillatus.
ferruginosus, *Link, ex Kunth, Enum. Pl.* iii. 393.—Afr. austr.
festucaeformis, *Nees, ex Mast. in Journ. Linn. Soc.* viii. (1865) 248.—Afr. austr.
filiformis, *Poir. Encyc.* vi. 173.—Afr. austr.
flexuosus, R. Br. Prod. 247 = Hypolaena flexuosa.
fraternus, *Kunth, Enum. Pl.* iii. 386.—Afr. austr.
fruticosus, Thunb. Diss. Rest. 16 = Leptocarpus paniculatus.
fruticosus, Thunb. ex Mast. in Journ. Linn. Soc. xiv. (1875) 417 = Leptocarpus modestus.
fuirenoides, Kunth, Enum. Pl. iii. 386 = setiger.
furcatus, *Nees, ex Mast. in Journ. Linn. Soc.* viii. (1865) 242.—Afr. austr.
fuscus, Nees, ex Mast. l. c. x. (1869) 250 = Dovea macrocarpa.
Garnotianus, *Kunth, Enum. Pl.* iii. 392.—Afr. austr.
Gaudichaudianus, *Kunth, l. c.* 387.—Afr. austr.
glaucus, Behr, ex Mast. in DC. Monog. Phan. i. 301.—Hab?
glomeratus, *Thunb. Diss. Rest.* 18.—Afr. austr.
glumaceus, Klotzsch, ex Krauss, in Flora, xxviii. (1845) 339 = triticeus.
gracilior, *F. Muell. ex Benth. Fl. Austral.* vii. 226.—Austral.
gracilis, *R. Br. Prod.* 245.—Austral.
gracilis, Nees, ex Benth. Fl. Austral. vii. 233 = Leptocarpus tenax.
graminifolius, *Kunth, Enum. Pl.* iii. 407—Afr. austr.
grandis, Spreng. ex Nees, in Linnaea, v. (1830) 660 = Lamprocaulos grandis.
Harveii, *Mast. in Journ. Linn. Soc.* viii. (1865) 253. t. 15.—Afr. austr.
Helenae, *Mast. l. c.* 233.—Afr. austr.
hirtellus, Kunth, Enum. Pl. iii. 426 = Leptocarpus peronatus.
Hystrix, *Mast. in Journ. Linn. Soc.* x. (1869) 276.—Afr. austr.
imbricatus, Thunb. Diss. Rest. 9 = Thamnochortus imbricatus.
impolitus, Kunth, Enum. Pl. iii. 404 = Hypolaena impolita.
incurvatus, Thunb. Diss. Rest. 18 = Leptocarpus incurvatus.
intermedius, *Kunth, Enum. Pl.* iii. 388.—Afr. austr.
intermedius, Steud. in Flora, xii. (1829) 132 = Elegia juncea.
ischaemoides, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 248 = festucaeformis.
junceus, Zeyh. & Nees, ex Nees, in Linnaea, v. (1830) 658 = Elegia juncea.
Kunthii, Steud. Syn. Pl. Cyp. 251 = triflorus.
laniger, *Kunth, Enum. Pl.* iii. 386.—Afr. austr.
lateriflorus, R. Br. Prod. 247 = Hypolaena lateriflora, elongata.
lateriflorus, Nees, ex Benth. Fl. Austral. vii. 222 = fastigiatus.
laxus, *R. Br. Prod.* 245. n. 3.—Austral.
laxus, R. Br. l. c. 246. n. 12 = Leptocarpus tenax.
Lehmanni, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 225 = ocreatus.
leptocarpoides, *Benth. Fl. Austral.* vii. 229.—Austral.
leptoclados, *Mast. in Journ. Linn Soc.* viii. (1865) 241.—Afr. austr.

RESTIO:—
leptostachyus, *Kunth, Enum. Pl.* iii. 407.—Afr. austr.
limbatus, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 235 = macer.
Loxocarya, Mast. in DC. Monog. Phan. i. 272 = Loxocarya cinerea.
lucaeanus, Kunth, Enum. Pl. iii. 385 = cuspidatus.
lucens, *Poir. Encyc.* vi. 169.—Afr. austr.
Ludwigii, *Steud. Syn. Pl. Cyp.* 254.—Afr. austr.
macer, *Kunth, Enum. Pl.* iii. 390.—Afr. austr.
macrogynus, Nees, ex Mast. in DC. Monog. Phan. i. 288 = egregius.
Mastersi, *F. Muell. Fragm.* viii. 68, in textu.—Afr. austr.
Megalotheca, Benth. Fl. Austr. viii. 222 = seq.
megalothecus, *F. Muell. l. c.* 99.—Austral.
membranaceus, Nees, in Linnaea, v. (1830) 657 = Elegia juncea.
micans, Nees, l. c. 649 = Thamnochortus micans.
microstachys, R. Br. Prod. 246 = Leptocarpus tenax.
microstachys, Nees, ex Mast. in DC. Monog. Phan. i. 247 = Gaudichaudianus.
miser, *Kunth, Enum. Pl.* iii. 392.—Afr. austr.
monanthos, *Mast. in Journ. Linn. Soc.* viii. (1865) 238.—Afr. austr.
monocephalus, *R. Br. Prod.* 245.—Austral.
monostachyus, Steud. in Flora, xii. (1829) 133 = bifidus.
mucronatus, Nees, in Linnaea, v. (1830) 660 = Dovea mucronata.
multiflorus, *Spreng. Syst.* i. 187.—Afr. austr.
Neesii, *Mast. in Journ. Linn. Soc.* viii. (1865) 232.—Afr. austr.
neglectus, Klotzsch, ex Hochst. in Flora, xxviii (1845) 338 = Leptocarpus neglectus.
nitens, *Nees, in Lehm. Pl. Preiss.* ii. 59.—Austral.
nudus, Nees, in Linnaea, v. (1830) 651 = Elegia nuda.
nutans, R. Br. Prod. 245 = Leptocarpus aristatus.
nutans, Steud. in Flora, xii. (1829) 104 = Ludwigii.
nutans, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 7 = Thamnochortus elongatus.
oblongus, *Mast. in Journ. Linn. Soc.* viii. (1865) 229.—Afr. austr.
obtusissimus, *Steud. Syn. Pl. Cyp.* 252.—Afr. austr.
ocreatus, *Kunth, Enum. Pl.* iii. 385.—Afr. austr.
oligocephalus, F. Muell. Fragm. viii. 68 = monocephalus.
oligostachyus, Kunth, Enum. Pl. iii. 399 = bifidus.
ornatus, *Steud. Syn. Pl. Cyp.* 256.—Austral.
oxylepis, Kunth, Enum. Pl. iii. 427 = Leptocarpus oxylepis.
pachystachyus, *Kunth*, l. c. 399.—Afr. austr.
paleaceus, Nees, ex Mast. in Journ. Linn. Soc. x. (1869) 225, 227, 230 = Leptocarpus modestus, Thamnochortus elongatus, dichotomus.
pallens, R. Br. Prod. 245 = gracilis.
pallens, Nees, ex Benth. Fl. Austral. vii. 233 = Leptocarpus tenax.
paniculatus, Link, Handb. i. 134 = multiflorus.
paniculatus, Rottb. Desc. 4; Progr. 10 = Leptocarpus paniculatus.
pannosus, *Mast. in Journ. Linn. Soc.* viii. (1865) 244.—Afr. austr.
parviflorus, Thunb. Diss. Rest. 13 = Elegia parviflora.
pauciflorus, Poir. Encyc. vi. 168.—Afr. austr.
pedicellatus, Mast. in Journ. Linn. Soc. viii. (1865) 252.—Afr. austr.
peronatus, Kunth, Enum. Pl. iii. 426 = ocreatus.
perplexus, *Kunth, Enum. Pl.* iii. 406.—Afr. austr.
pilisepalus, Steud. Syn. Pl. Cyp. 256 (Cyperacea).—Ins. Waigiou.
polystachyus, Kunth, Enum. Pl. iii. 402 = Mastersi.
propinquus, Nees, in Linnaea, v. (1830) 653 = Elegia propinqua.
Pseudoleptocarpus, Kunth, Enum. Pl. iii. 399 = bifidus.
pubescens, R. Br. Prod. 247 = Hypolaena pubescens.
punctulatus, Nees, ex Mast. in Journ. Linn. Soc. viii. (1865) 242 = Thamnochortus distichus.
purpurascens, *Nees, ex Mast. l. c.* 249.—Afr. austr.
pycnostachyus, *Mast. l. c.* 232.—Afr. austr.
quadratus, *Mast. l. c.* x. (1869) 277. = Afr. austr.
quinquefarius, *Nees, in Linnaea*, v. (1830) 639.—Afr. austr.
racemosus, Lam. Illustr. iii. 400 = Dovea tectorum.
ramiflorus, Nees, in Linnaea, v. (1830) 644 = Leptocarpus paniculatus.
Rhodocoma, *Mast. in Journ. Linn. Soc.* x. (1869) 275.—Afr. austr.
rottboellioides, *Kunth, Enum. Pl.* iii. 394.—Afr. austr.

RESTIO:—
saroclados, *Mast. in DC. Monog. Phan.* i. 291.—Afr. austr.
scariosus, Sieber, ex Kunth, Enum. Pl. iii. 430 = Thamnochortus dichotomus.
scariosus, Thunb. Diss. Rest. 15 = Thamnochortus fruticosus.
schoenoides, *Kunth, Enum. Pl.* iii. 391.—Afr. austr.
Scopa, Thunb. Diss. Rest. 19 = Cannomois Cephalotes.
scoparius, *Kunth, Enum. Pl.* iii. 390.—Afr. austr.
setiger, *Kunth, l. c.* 385.—Afr. austr.
setuligerus, Nees, in Lehm. Pl. Preiss. ii. 58 = Leptocarpus tenax.
Sieberi, *Kunth, Enum. Pl.* iii. 287.—Afr. austr.
simplex, Forst. ex Steud. in Flora, xii. (1829) 134 = Thamnochortus elongatus.
simplex, Linn. ex Mast. in Journ. Linn. Soc. x. (1869) 231 = Thamnochortus imbricatus.
simplex, Murr. Syst. ed. XIII. 738 = Leptocarpus simplex.
simplex, *Thunb. Diss. Rest.* 16.—Afr. austr.
Sonderianus, *Mast. in Journ. Linn. Soc.* viii. (1865) 252.—Afr. austr.
sphacelatus, *R. Br. Prod.* 246.—Austral.
spicifer, Lam. Illustr. iii. 399 = Thamnochortus cernuus.
spicigerus, Kunth, Enum. Pl. iii. 440, sphalm. = Thamnochortus cernuus.
spicigerus, Thunb. Diss. Rest. 11 = Thamnochortus spicigerus.
spinulosus, *Kunth, Enum. Pl.* iii. 402.—Afr. austr.
Sprengelii, Mast. in Journ. Linn. Soc. viii. (1865) 224 = cuspidatus.
squamosus, *Thunb. in Hoffm. Phytog. Blatter*, i. (1803) 8.—Afr. austr.
squarrosus, *Poir. in Lam. Encyc.* vi. 174.—Afr. austr.
squarrosus, Spreng. Syst. i. 186 = cuspidatus.
Steudelii, F. Muell. Fragm. viii. 70 = ornatus.
stipularis, Banks, ex Mast. in Journ. Linn. Soc. xxi. (1885) 588 = Elegia stipularis.
strobilifer, *Kunth, Enum. Pl.* iii. 398.—Afr. austr.
subfalcatus, *Nees, ex Mast. in Journ. Linn. Soc.* viii. (1865) 231.—Afr. austr.
subtilis, *Nees, ex Mast. l. c.* 251.—Afr. austr.
subulatus, *Mast. l. c.* 248.—Afr. austr.
subverticillatus, *Mast. l. c.* 227.—Afr. austr.
sulcatus, Kunth, Enum. Pl. iii. 404 = Hypodiscus Willdenovia.
synchroolepis, Steud. Syn. Pl. Cyp. 250 = Hypodiscus synchroolepis.
tectorum, Linn. f. Suppl. 425 = Dovea tectorum.
tectorum, Sieber, ex Kunth, Enum. Pl. iii. 462 = Elegia nuda.
tectorum, Thunb. Fl. Cap. i. 85; Steud. in Flora, xii. (1829) 132 = Elegia deusta.
tenellus, Nees, in Lehm. Pl. Preiss. ii. 57 = Leptocarpus aristatus.
tenuis, Burch. ex Mast. in Journ. Linn. Soc. x. (1869) 265 = Hypolaena tenuis.
tenuis, Krauss, in Flora, xxviii. (1845) 337 = curviramis.
tenuissimus, *Kunth, Enum. Pl.* iii. 394.—Afr. austr.
tetragonus, *Thunb. Diss. Rest.* 17.—Afr. austr.
tetraphyllus, *Labill. Nov. Holl. Pl.* ii. 77. tt. 226, 227.—Austral.
tetrasepalus, Steud. Syn. Pl. Cyp. 251 = Thamnochortus imbricatus.
Thamnochortus, Thunb. Diss. Rest. 15 = Thamnochortus dichotomus.
thyrsifer, Lam. Illustr. iii. 399 = Elegia propinqua.
thyrsifer, Rottb. Desc. 8; Progr. 11 = Elegia juncea.
tremulus, *R. Br. Prod.* 245.—Austral.
triflorus, *Rottb. Desc. et Ic.* 3. t. 2. f. 2.—Afr. austr.
triflorus, Sieber, ex Mast. in Journ. Linn. Soc. viii. (1865) 235 = Sieberi.
triflorus, Steud. in Flora, xii. (1829) 133 = bifidus.
trisepalus, *Nees, in Lehm. Pl. Preiss.* ii. 58.—Austral.
triticeus, *Rottb. Desc.* 7; *Progr.* 11.—Afr. austr.
triticeus, Thunb. ex Mast. in Journ. Linn. Soc. xiv. (1875) 417 = pannosus.
tropicus, *R. Br. Prod.* 246.—Austral.
umbellatus, Thunb. Diss. Rest. 11 = Thamnochortus umbellatus.
umbellatus, Thunb. ex Mast. in Journ. Linn. Soc. xiv. (1875) 415 = Thamnochortus cernuus, imbricatus.
uncinatus, Nees, ex Benth. Fl. Austral. vii. 421 = Caustis recurvata.
vacillans, Steud. Syn. Pl. Cyp. 255 = Hypolaena fastigiata.

RESTIO:—
vaginatus, Thunb. Diss. Rest. 10 = Thamnochortus imbricatus.
vaginatus, Willd. Sp. Pl. iv. 719 = bifidus.
venustulus, *Kunth, Enum. Pl.* iii. 388.—Afr. austr.
verticillaris, Linn. f. Suppl. 425; Thunb. Diss. Rest. 19 = Elegia verticillaris.
verticillatus, Spreng. Syst. i. 186 = Elegia verticillaris.
vilis, *Kunth, Enum. Pl.* iii. 389.—Afr. austr.
vimineus, Murr. Syst. ed. XIII. 738; Linn. Herb. ex Mast. in Journ. Linn. Soc. x. (1869) 231 = Thamnochortus dichotomus.
vimineus, Rottb. Desc. et Ic. 4. t. 2. f. 1 = Leptocarpus incurvatus.
vimineus, Thunb. ex Mast. in Journ. Linn. Soc. xiv. (1875) 420 = perplexus.
virgatus, Rottb. Desc. 5; Progr. 10 = Cannomois Cephalotes.
Wallichii, *Mast. in Journ. Linn. Soc.* viii. (1865) 234.—Afr. austr.
xyridioides, Kunth, Enum. Pl. iii. 397 = quinquefarius.

RESTREPIA, H. B. & K. Nov. Gen. et Sp. i. 366. t. 94 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 491.
antennifera, *H. B. & K. Nov. Gen. et Sp.* i. 367. t. 9.—N. Granat.
antennifera, Lindl. Fol. Orch. Restrep. 1, partim = maculata.
aspasicensium, *Reichb. f. in Bonplandia*, iii. (1855) 70.—Venezuela.
biflora, Regel, in Ann. Sc. Nat. Sér. IV. vi. (1856) 373 = Pleurothallis hemirhoda.
cucullata, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 108.—N. Granat.
Dayana, *Reichb. f. in Gard. Chron.* (1875) II. 257.—Costa Rica.
elegans, *Karst. in Otto & Dietr. Allg. Gartenz.* xv. (1847) 202; *Ausw. Neuer Gew. Venez.* 8. t. 2.—Venezuela.
erythroxantha, *Reichb. f. in Linnaea*, xxii. (1849) 817.—N. Granat.
Falkenbergii, *Reichb. f. in Gard. Chron.* (1880) I. 232.—N. Granat.
Gardneri, *Benth. in Benth. & Hook. f. Gen.* iii. 491.—Bras.
guttulata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 357.—Ecuador.
hypolepta, *Reichb. f. in Walp. Ann.* vi. 204.—Bras.
Kegelii, *Reichb. f. in Linnaea*, xli. (1877) 133.—Guian.
Lansbergii, Hook. Bot. Mag. t. 5257 = xanthophthalma.
Lansbergii, *Reichb. f. & Wagen. in Bonplandia*, ii. (1854) 23.—Venezuela.
lonchophylla, *Rodrig. Orch. Nov.* i. 35.—Bras.
maculata, *Lindl. Orchid. Linden.* 4.—N. Granat.
microphylla, *Rodrig. Rev. de Engenh.* iii. (1881) 110; *ex ej. Orch. Nov.* ii. 115.—Bras.
Miersii, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Bras.* 150.—Bras.
muscifera, *Reichb. f. ex Lindl. Fol. Orchid. Restrep.* 2.—Guatemala.
nuda, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 275 = Pleurothallis hemirhoda.
ophiocephala, *Reichb. f. in Bonplandia*, ii. (1854) 88.—Mexic.
parvifolia, Lindl. Orchid. Linden. 4 = Brachionidium parvifolium.
prorepens, *Reichb. f. in Gard. Chron.* (1877) I. 810.—Costa Rica.
punctulata, Lindl. Fol. Orchid. Restrep. 2 = elegans.
Reichenbachiana, *Endres, in Gard. Chron.* (1875) II. 356.—Costa Rica.
rhyneantha, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 114.—Peruv.
Schlimii, *Reichb. f. in Linnaea*, xli. (1877) 45.—N. Granat.
tentaculata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 34. t. 59.—Peruv.
ujarensis, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Costa Rica.
varicosa, *Lindl. Fol. Orchid. Restrep.* 3.—Peruv.
vittata, Lindl. in Journ. Hort. Soc. iii. (1848) 314 = Pleurothallis hemirhoda.
Wageneri, *Reichb. f. in Bonplandia*, ii. (1854) 23.—Venezuela.
xanthophthalma, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 300.—Guatemala.

RESUPINARIA, Rafin. Sylva Tellur. 115 (1838) = **Sesbania**, Scop. (Legumin.).
grandiflora, Rafin. l. c. 116 = S. grandiflora.

RETAMA, Rafin. Sylva Tellur. 22 (1838); Boiss. Voy. Bot. Espagne, ii. 143 (1839) = **Genista**, Tourn. (Legumin.).
albiflora, Rafin. l. c. 22 = Cytisus albus.
angulata, Griseb. Spicil. Fl. Rumel. i. 5.— Europ. austr.
atlantica, Pomel, Nouv. Mat. Fl. Atl. 172 = G. atlantica.
Bovei, Webb, Otia Hisp. 24 = G. monosperma.
dasycarpa, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 57, nomen = G. dasycarpa.
Duriaei, Webb, in Ann. Sc. Nat. Sér. II. xx. (1843) 279 = G. Duriaei.
Gussonei, Webb, l. c. 280 = G. monosperma.
hipponensis, Webb, Otia Hisp. 25.—Hispan.
lutea, Rafin. Sylva Tellur. 22 = G. sphaerocarpa?
microcarpa, Webb, in Ann. Sc. Nat. Sér. II. xx. (1843) 280.—Ins. Canar.
monosperma, Boiss. Voy. Bot. Espagne, ii. 144 = G. monosperma.
parviflora, Webb, Otia Hisp. 23 = G. parviflora?
Raetam, Webb & Berth. Phyt. Canar. ii. 56 = G. Raetam.
rhodorhizoides, Webb & Berth. l. c. 54 = G. rhodorhizoides.
Spachii, Webb, in Ann. Sc. Nat. Sér. II. xx. (1843) 281 = G. rhodorhizoides.
sphaerocarpa, Boiss. Voy. Bot. Espagne, ii. 144 = G. sphaerocarpa.
Webbii, Webb, Otia Hisp. 24; Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 57 = G. monosperma.

RETAMILIA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 483; Contrib. i. 285 = **Retanilla**, Brongn. (Rhamn.).

RETANILLA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 364. t. 2. *RHAMNEAE*, Benth. & Hook. f. i. 384.
MOLINAEA, Comm. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 364.
RETAMILIA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 483.
affinis, C. Gay, Fl. Chil. ii. 26 = stricta.
articulata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 486.—Chili.
crassa, *Phil. in Linnaea*, xxxiii. (1864–65) 38.—Chili.
Ephedra, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 365.—Peruv.
glauca, *Phil. in Linnaea*, xxxiii. (1864–65) 37.—Chili.
obcordata, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 365.—Peruv.
spinifer, *C. Gay, Fl. Chil.* ii. 27.—Chili.
stricta, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 173.—Chili.
trinervia, Hook. & Arn. l. c. 174 = Trevoa trinervis.

RETINARIA, Gaertn. Fruct. ii. 187. t. 120. f. 4 (1791) = **Gouania**, Linn. (Rhamn.).
scandens, Gaertn. l. c. = G. tiliaefolia.
volubilis, Gaertn. l. c. t. 120. f. 4 = G. tiliaefolia.

RETINIPHYLLUM, Humb. & Bonpl. Pl. Aequin. i. 86. t. 25 (1808). *RUBIACEAE*, Benth. & Hook. f. ii. 98.
AMMIANTHUS, Spruce, ex Benth. & Hook. f. Gen. ii. 98 (1873).
COMMIANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 223.
concolor, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 8.—Bras.
discolor, *Muell. Arg. l. c.* 11.—Bras.
Martianum, *Muell. Arg. l. c.* 9.—Bras.
pallidum, *Muell. Arg. l. c.* 12.—Bras.
pilosum, *Muell. Arg. l. c.* 7.—Bras.
rhabdocalyx, *Muell. Arg. l. c.* 10.—Bras.
scabrum, *Benth. in Hook. Journ. Bot.* iii. (1841) 222.—Venezuela.
Schomburgkii, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 12.—Bras.
secundiflorum, *Humb. & Bonpl. Pl. Aequin.* i. 86. t. 25.—Venezuela.
speciosum, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 10.—Bras.
truncatum, *Muell. Arg. l. c.* 11.—Bras.

RETINISPORA, Sieb. & Zucc. Fl. Jap. ii. 36. tt. 121–123 (1842) = **Thuya**, Linn. (Conif.).
andelyensis, Carr. in Rev. Hortic. (1880) 178 = Cupressus thyoides (cf. Addend.).
decurvata, Hort. ex Carr. Conif. ed. II. 141 = T. orientalis.
decussata, Hort. ex Gord. Pinet. ed. II. 364 = T. orientalis.
Devriesiana, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 495 = T. occidentalis.
dubia, Carr. Conif. ed. II. 141 = T. occidentalis.
Ellwangeriana, Carr. in Rev. Hortic. (1869) 349 = T. occidentalis.
ericoides, Zucc. ex Gord. Pinet. Suppl. 91 = T. orientalis, occidentalis.
filicoides, Hort. Syme, in Gard. Chron. (1876) i. 235. f. 40 = Cupressus obtusa (cf. Addend.).
filifera, Fowler, in Gard. Chron. (1872) 1490 ; Stand. ex Gord. Pinet. ed. II. 364 = Cupressus obtusa (cf. Addend.).
Fusinoki, Zucc. ex Gord. Pinet. Suppl. 95 = Cupressus obtusa (cf. Addend.).
glauca, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 495 (Quid ?).—Hab. ?
japonica, Hort. l. c. (Quid ?).—Hab. ?
juniperoides, Carr. Conif. ed. II. 140 = T. orientalis.
leptoclada, Zucc. ex Gord. Pinet. Suppl. 91 = Cupressus thyoides (cf. Addend.).
lycopodioides, Gord. l. c. 92 = Cupressus pisifera (cf. Addend.).
meldensis, Carr. Conif. ed. II. 103 = T. occidentalis.
monstrosa, Hort. ex Carr. l. c. 132 = Cupressus obtusa (cf. Addend.).
obtusa, Sieb. & Zucc. Fl. Jap. ii. 38. t. 121 = Cupressus obtusa (cf. Addend.).
pisifera, Sieb. & Zucc. l. c. 39. t. 122 = Cupressus pisifera (cf. Addend.).
plumosa, Hort. ex Carr. Conif. ed. II. 791 ; Veitch, ex C. Koch, Dendrol. ii. II. 170 = Cupressus pisifera (cf. Addend.).
Pseudo-squarrosa, Carr. Conif. ed. II. 140 = Cupressus thyoides (cf. Addend.).
recurvata, Hort. ex Mast. in Journ. Linn. Soc. xviii. (1881) 495 = T. orientalis.
rigida, Carr. Conif. ed. II. 141 = T. orientalis.
squarrosa, Hort. ex Carr. l. c. 140, 141 = Cupressus pisifera (cf. Addend.).
squarrosa, Sieb. & Zucc. Fl. Jap. ii. 40. t. 123 = Cupressus pisifera (cf. Addend.).
stricta, Hort. ex Gord. Pinet. ed. II. 428 = Cupressus pisifera (cf. Addend.).
tetragona, Barron, ex Gord. l. c. 429 = Cupressus obtusa (cf. Addend.).

RETINODENDRON, Korth. Verh. Nat. Gesch. Bot. 55. t. 8 (1839) = **Vatica**, Linn. (Dipteroc.).
lanceaefolium, Korth. l. c. 56 = V. lanceaefolia.
pauciflorum, Korth. l. c. 58 = V. pauciflora.
Rassak, Korth. l. c. 56. t. 8 = V. Rassak.

RETINOPHLEUM, Benth. & Hook. f. Gen. i. 570 (1865), sphalm. = RHETINOPHLOEUM, Karst. = **Cercidium**, Tul. (Legumin.).

RETINOSPORA, Carr. Conif. ed. I. 132 (1855) = RETINISPORA, Sieb. & Zucc. = **Thuya**, Linn. (Conif.).

RETTBERGIA, Raddi, Agrost. Bras. 17 (1823) = **Chusquea**, Kunth (Gramin.).
bambusaeoides, Raddi, l. c. = C. Gaudichaudii.
capitata, Nees, in Linnaea, ix. (1834) 489 = C. capitata.

RETZIA, Thunb. in Phys. Sällsk. Handl. Stockh. i. (1776) i. 55. t. 1. *SOLANACEAE*, Benth. & Hook. f. ii. 905.
SOLENOSTIGMA, Klotzsch, ex Walp. Rep. iii. 125 (1844–45).
campanuloides, *Spreng. Syst.* i. 589.—Afr. austr.
capensis, *Thunb. in Phys. Sällsk. Handl. Stockh.* i. (1776) i. 55. t. 1.—Afr. austr.
pilosa, Rottl. ex Choisy, in DC. Prod. ix. 403 = Convolvulus Rottlerianus.
roëlloides, *Spreng. Syst.* i. 589.—Afr. austr.
spicata, J. F. Gmel. Syst. 333 = capensis.

REUSSIA, Dennst. Schluess. Hort. Malab. 33 (1818) = **Paederia**, Linn. (Rubiac.).
sarmentosa, Dennst. l. c. = P. tomentosa.

REUSSIA, Endl. Gen. 139 (1836). *PONTEDERIACEAE*.
subovata, *Solms, in DC. Monog. Phan.* iv. 534.—Am. austr.
triflora, *Seub. in Mart. Fl. Bras.* iii. i. 96.—Bras.

REUTERA, Boiss. Elench. 46 (1838) = **Pimpinella**, Riv. (Umbell.).
acuminata, Edgew. in Trans. Linn. Soc. xx. (1846) 52 = P. acuminata.
albiflora, Costa, Supl. Fl. Catal. 304.—Hispan.
aurea, Boiss. Fl. Orient. ii. 863 = P. aurea.
catalaunica, Costa, ex Nym. Consp. 309 = *gracilis*.
cervariaefolia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 134 = P. aurea ?
deverroides, Boiss. l. c. 135 = P. deverroides.
dichotoma, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 863.—Persia.
flabellifolia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 135.—Cappadoc.
flava, Boiss. l. c. 135 = P. aurea.
Fontanesii, Boiss. Voy. Bot. Espagne, ii. 244 = P. lutea.
gracilis, Boiss. Elench. 46.—Hispan.
papillaris, Boiss. Diagn. Ser. II. ii. 76.—Afghan.
pastinacaefolia, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 134.—Persia.
procumbens, Boiss. Elench. 47.—Hispan.
puberula, Loscos & Pardo, Ser. Inconf. Arag. 44 = *gracilis*.
rigidula, Boiss. & Orph. Diagn. Ser. II. ii. 77 = P. chrysantha.
tenuis, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 862.—Cataonia.
tragiodes, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 135.—Persia.

REVEESIA, Walp. Ann. i. 105 (1848-49), sphalm. = **Reevesia**, Lindl. (Stercul.).

REVERCHONIA, A. Gray, in Proc. Am. Acad. xvi. (1881) 107. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 1222.
arenaria, *A. Gray, l. c.*—Texas.

REYESIA, Clos, in C. Gay, Fl. Chil. iv. 418. t. 52 (1849). *SOLANACEAE*, Benth. & Hook. f. ii. 909.
PTEROGLOSSIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 32.
chilensis, *Clos, in C. Gay, Fl. Chil.* iv. 419.—Chili.

REYNANDIA, Kunth, Rév. Gram. 195. t. 9 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1120.
filiformis, *Kunth, l. c.*—Ins. S. Doming.

REYNOLDSIA, A. Gray, Bot. U. St. Expl. Exped. i. 723. t. 92, 93 (1854) = **Trevesia**, Vis. (Araliac.).
pleiosperma, A. Gray, l. c. 725. t. 93.—Ins. Samoa.
sandwicensis, A. Gray, l. c. 724. t. 92.—Ins. Sandvic.
tahitensis, Nadeaud, Enum. Pl. Tahiti, 63.—Ins. Tahiti.
verrucosa, Seem. in Journ. Bot. ii. (1864) 245.—Ins. Tahiti.

REYNOSIA, Griseb. Cat. Pl. Cub. 33 (1866). *RHAMNEAE*.
latifolia, *Griseb. l. c.* 34.—Cuba.
mucronata, *Griseb. l. c.* 34, 283.—Cuba.
retusa, *Griseb. l. c.* 34.—Cuba.

REYNOUTHRIA, Steud. Nom. ed. I. 684 (1821) = seq.

REYNOUTRIA, Houtt. Handleid. viii. 639. t. 51 (1777). *INC. SEDIS.*
japonica, *Houtt. l. c.* 640.—Japon.

RHABARBARUM, Tourn. ex Adans. Fam. ii. 277 (1763) = **Rheum**, Linn. (Polygon.).
palmatum, Moench, Meth. 628 = Rheum palmatum.
Rhaponticum, Moench, l. c. = Rheum Rhaponticum.
Ribes, Moench, l. c. = Rheum Ribes.

RHABDADENIA, Muell. Arg. in Mart. Fl. Bras. vi. i. 173. t. 52 (1860). *APOCYNACEAE*, Benth. & Hook. f. ii. 727.
barbata, *Miers, Apocyn. S. Am.* 123.—Ind. occ.
Berteri, *Muell. Arg. in Linnaea*, xxx. (1859–60) 435.—Ins. S. Doming.
biflora, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 175.—Am. austr.

RHABDADENIA:—
campestris, *Miers, Apocyn. S. Am.* 121.—Bras.
cordata, *Miers, l. c.* 122.—Mexic.
cubensis, *Muell. Arg. in Linnaea*, xxx. (1859–60) 435.—Ind. occ.
Ehrenbergii, Muell. Arg. ex Griseb. Fl. Brit. W. Ind. 415 = Echites paludosa.
laxiflora, *Miers, Apocyn. S. Am.* 120.—Ind. occ.
Lindeniana, *Muell. Arg. in Linnaea*, xxx. (1859–60) 437.—Cuba.
lucida, *Miers, Apocyn. S. Am.* 123.—Venezuela.
macrostoma, *Muell. Arg. in Linnaea*, xxx. (1859–60) 435.—Guian.
madida, *Miers, Apocyn. S. Am.* 121.—Bras.
nervosa, *Miers, l. c.* 122.—Ind. occ.
paludosa, *Miers, l. c.* 119.—Bras.
Pohlii, *Muell. Arg. in Mart. Fl. Bras.* vi. i. 174.—Bras.
Sagraei, Muell. Arg. in Griseb. Fl. Brit. W. Ind. 415 = Echites Sagraei.
Wrighteana, Muell. Arg. in Linnaea, xxx. (1859–60) 435 = Mandevilla Wrighteana (in Addend.).

RHABDIA, Mart. Nov. Gen. et Sp. ii. 136, t. 195 (1827). *BORAGINEAE*, Benth. & Hook. f. ii. 842.
ROTULA, Lour. Fl. Cochinch. 121 (1790).
crebrifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 435 ; Contrib. ii. 222. t. 84 = lycioides.
fluvialis, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 175 = lycioides.
lycioides, *Mart. Nov. Gen. et Sp.* ii. 136. t. 195.—As. Afr. et Am. trop.
sericea, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 176 = lycioides.
viminea, Dalz. ex Hook. Ic. Pl. t. 823 = lycioides.

RHABDOCALYX, Lindl. Veg. Kingd. 629 (1847) = **Cordia**, Linn. (Borag.).

RHABDOCHLOA, Kunth, in Mém. Mus. Par. ii. (1815) 73 = RABDOCHLOA, Beauv. = **Leptochloa**, Beauv. (Gramin.).

RHABDOCRINUM, Reichb. Consp. 65 (1828) = **Lloydia**, Salisb. (Liliac.).
graecum, Reichb. l. c. = L. graeca.
serotinum, Reichb. l. c. = L. alpina.

RHABDOSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 68. *UMBELLIFERAE*, Benth. & Hook. f. i. 898.
Aucheri, *Boiss. l. c.* 69.—Persia.
petiolare, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 899.—Persia.
stenophyllum, *Boiss. & Haussk. l. c.* 898.—Persia.

RHABDOSTIGMA, Hook. f. in Benth. & Hook. f. Gen. ii. 109 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Kirkii, *Hook. f. l. c.*—Afr. trop.

RHABDOTHAMNUS, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 460. *GESNERACEAE*, Benth. & Hook. f. ii. 1011.
negriana, F. Muell. Fragm. vii. 152 = Negria rhabdothamnoides.
scabrosus, Steud. Nom. ed. II. ii. 443 = seq.
Solandri, *A. Cunn. in Ann. Nat. Hist.* Ser. I. i. (1838) 460.—N. Zel.

RHABDOTHECA, Cass. in Dict. Sc. Nat. xlviii. 424 (1827) = **Launaea**, Cass. (Compos.).
angustifolia, Pomel, Nouv. Mat. Fl. Atl. 263 (= *Zollikoferia arabica*.).—Alger.
asplenifolia, Webb, in Hook. Niger Fl. 146 = L. aspleniifolia.
Brunnera, Webb, l. c. 147 = Sonchus Brunneri.
Candolleana, Pomel, Nouv. Mat. Fl. Atl. 263 (= *Zollikoferia Candolleana*).—Alger.
Cassiniana, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 286 (= *Zollikoferia Cassiniana*).—Abyss.
chondrilloides, Webb, in Hook. Niger Fl. 146 = L. nudicaulis.
divaricata, Webb, l. c. = L. nudicaulis.
longiloba, Pomel, Nouv. Mat. Fl. Atl. 263 (= *Zollikoferia longituba*).—Alger.
nudicaulis, Pomel, l. c. 262 = L. nudicaulis.
picridioides, Webb, in Hook. Niger Fl. 146.—Ins. Cap. Virid.
prenanthoides, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 286 = Sonchus prenanthoides.

RHABDOTHECA :—
pumila, Pomel, Nouv. Mat. Fl. Atl. 263 (=*Zollikoferia pumila*).—Alger.
quercifolia, Pomel. l. c. (=*Zollikoferia quercifolia*).—Alger.
resedifolia, Pomel, l. c. = Launaea nudicaulis.
Rueppellii, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 286.—Afr. trop.
sonchoides, Cass. in Dict. Sc. Nat. xlviii. 425 = L. nudicaulis.
spinosa, Webb, in Hook. Niger Fl. 147 = L. spinosa.
squarrosa, Pomel, Nouv. Mat. Fl. Atl. 263 (=*Zollikoferia arabica*).—Alger.
tenuiloba, Pomel, l. c. (=*Zollikoferia tenuiloba*).—Alger.

RHACHICALLIS, Spach, Hist. Veg. Phan. viii. 374 (1839) = **Rachicallis**, DC. (Rubiac.).

RHACOMA, Adans. Fam. ii. 117 (1763) = **Leuzea**, DC. (Compos.).

RHACOMA, Linn. Syst. ed. X. 896 (1759) = **Myginda**, Linn. (Celastr.).
Crossopetalum, Linn. l. c. = M. Rhacoma.
Uragoga, Baill. Hist. Pl. vi. 27 = M. Uragoga.

RHADAMANTHUS, Salisb. Gen. Pl. Fragm. 37 (1866). *LILIACEAE*, Benth. & Hook. f. iii. 808.
convallarioides, *Salisb. l. c.*—Afr. austr.

RHADINOCARPUS, Vog. in Linnaea, xii. (1838) 108 = **Chaetocalyx**, DC. (Legumin.).
acutifolius, Vog. l. c. 111 = C. acutifolia (cf. Addend.).
brasiliensis, Vog. l. c. 110 = C. brasiliensis.
multiflorus, Vog. l. c. 108.—N. Granat.

RHADIOLA, Savi, in Giorn. Tosc. Sc. Med. Fis. Nat. i. (1840) 194, err. typ. = **Radiola**, Dill. (Lineae).

RHAEO, C. B. Clarke, in DC. Monog. Phan. iii. 316 (1881), err. typ. = **Rhoeo**, Hance (Commel.).

RHAGADIOLUS, Tourn. ex Scop. Introd. 122 (1777). *COMPOSITAE*, Benth & Hook. f. ii. 510.
GARHADIOLUS, Jaub. & Spach, Illustr. iii. 119. tt. 284, 285 (1849).
HEDYPNOIS, Schreb. Gen. 552 (1791).
arenarius, *Benth. & Hook. f. Gen.* ii. 510.—Marocc.
creticus, All. Fl. Pedem. i. 226 = R. Hedypnois.
edulis, *Gaertn. Fruct.* ii. 354.—Reg. Mediterr.; Oriens.
Hedypnois, *All. Fl. Pedem.* i. 226 = Reg. Mediterr.
Hedypnois, *Fisch. & Mey. ex Fisch. Mey. & Trautv. Ind. Sem. Hort. Petrop.* iv. 46.—Armenia; Persia.
intermedius, Tenore, Fl. Med. ii. 25 = edulis.
Koelpinia, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 265 = Koelpinia linearis.
Kolpinia, Lam. Illustr. iii. 234 = praec.
lampsanaefolius, Mirb. Hist. Nat. Pl. xi. [1805] 328 = edulis.
lampsanoides, Desf. Fl. Atlant. ii. 234 = edulis.
lyratus, Lam. Illustr. iii. 234 = edulis.
pendulus, *Benth. & Hook. f. Gen.* ii. 510.—Reg. Mediterr.
radiatus, All. Fl. Pedem. i. 227 = Hyoseris radiata.
rigidus, Pomel, Nouv. Mat. Fl. Atl. 17 = edulis.
scaber, All. Fl. Pedem. i. 226 = Hyoseris scabra.
stellatus, DC. Prod. vii. 80, in syn. = Hyoseris radiata.
stellatus, Gaertn. Fruct. ii. 354 = edulis.
taraxacoides, All. Fl. Pedem. i. 227 = Leontodon hirtus, hispidus.
Zacintha, All. l. c. = Zacintha verrucosa.

RHAGANUS, E. Mey. in Drège, Zwei Pfl. Docum. 216 (1843) = **Bersama**, Fresen. (Sapind.).
lucidus, E. Mey. l. c.; et in Flora, xxxii. (1849) 67.—Afr. austr.

RHAGODIA, R. Br. Prod. 408 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 49.
baccata, Moq. in DC. Prod. xiii. II. 50 = Billardieri.
Billardieri, *R. Br. Prod.* 408.—Austral.
Candolleana, Moq. Chenop. Enum. 10 = Billardieri.
chenopodioides, Moq. l. c. 11 = Billardieri?
congesta, Moq. in DC. Prod. xiii. II. 51 = Billardieri.
crassifolia, *R. Br. Prod.* 408.—Austral.
dioica, *Nees, in Lehm. Pl. Preiss.* i. 636.—Austral.
Drummondii, Moq. in DC. Prod. xiii. II. 52 = crassifolia.
Eschscholtziana, *Fenzl, in Denkschr. Akad. Wien, (Math.-Nat.)* i. (1850) 257.—Ins. Sandvic.

RHAGODIA:—
Gaudichaudiana, *Moq. Chenop. Enum.* 11.—Austral.
hastata, *R. Br. Prod.* 408.—Austral.
linifolia, *R. Br. l. c.* 408.—Austral.
linifolia, Nees, in Lehm. Pl. Preiss. i. 637 = Preissii.
nitrariacea, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 73 = Chenopodium nitrariaceum
nutans, *R. Br. Prod.* 408.—Austral.
obovata, *Moq. Chenop. Enum.* 10.—Austral.
parabolica, *R. Br. Prod.* 408.—Austral.
parvifolia, Moq. in DC. Prod. xiii. II. 52 = crassifolia.
Preissii, *Moq. l. c.* 49.—Austral.
prostrata, A. Cunn. ex Benth. Fl. Austral. v. 156 = spinescens.
radiata, Nees, in Lehm. Pl. Preiss. i. 637 = Billardieri.
reclinata, A. Cunn. ex Benth. Fl. Austral. v. 153 = parabolica.
spinescens, *R. Br. Prod.* 408.—Austral.

RHAMINDIUM, Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 39 (1884), err. typ. = **Rhamnidium**, Reiss. (Rhamn.).

RHAMNELLA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 30 (1867) = **Microrhamnus**, A. Gray (Rhamn.).
japonica, Miq. l. c. = M. franguloides.

RHAMNIDIUM, Reiss. in Mart. Fl. Bras. xi. I. 94. t. 31 (1861). *RHAMNEAE*, Benth. & Hook. f. i. 378.
cognatum, *Reiss. l. c.* 95.—Bras.
elaeocarpum, *Reiss. l. c.* 94.—Bras.
glabrum, *Reiss. l. c.* 95.—Bras.
molle, *Reiss. l. c.*—Bras.
reticulatum, *Griseb. Cat. Pl. Cub.* 33.—Cuba.
retusum, *Griseb. l. c.* 32.—Cuba.
revolutum, Chapm. Fl. S. St. Suppl. 612 = Reynosia latifolia.
revolutum, *Griseb. Cat. Pl. Cub.* 32.—Cuba.

RHAMNOIDES, Tourn. ex Moench, Meth. 343 (1794) = **Hippophaë**, Linn. (Elaeagn.).
Hippophaë, Moench, l. c. = H. Rhamnoides.

RHAMNOPSIS, Reichb. Consp. 188 (1828) = **Flacourtia**, Comm. (Bixin.).
sepiaria, Reichb. l. c. = F. sepiaria.

RHAMNOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.

RHAMNUS, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 58 (1737). *RHAMNEAE*, Benth. & Hook. f. i. 377.
ALATERNUS, Mill. Gard. Dict. ed. VI (1752).
AMPELOPLIS, Rafin. Sylva Tellur. 33 (1838).
ATADINUS, Rafin. l. c. 30 (1838).
ATULANDRA, Rafin. l. c. 31 (1838).
CERVISPINA, [Dill.] Moench, Meth. 686 (1794).
ENDOTROPIS, Rafin. Sylva Tellur. 31 (1838).
FORGEROUXA, Neck. Elem. ii. 124 (1790).
FRANGULA, Tourn. ex Hall. Enum. Helv. i. 164 (1742); et Mill. Gard. Dict. ed. VI (1752).
GIRTANNERIA, Neck. Elem. ii. 121 (1790).
HETTLINGERIA, Neck. l. c. 124 (1790).
LITHOPLIS, Rafin. Sylva Tellur. 32 (1838).
MARCORELLA, Neck. Elem. ii. 122 (1790).
PALIURUS, Tourn. ex Mill. Gard. Dict. ed. VI (1752).
PERFONON, Rafin. Sylva Tellur. 29 (1838).
SARCOMPHALUS, Rafin. l. c. (1838).
SCIADOPHILA, Phil. in Linnaea, xxviii. (1856) 618.
VERLANGIA, Neck. Elem. ii. 125 (1790).
acuminata, Colebr. in Roxb. Fl. Ind. i. 615 = Colubrina asiatica.
adriatica, Jord. ex Nym. Consp. 145 = intermedia.
agrestis, Lour. Fl. Cochinch. 197 = Zizyphus agrestis.
affinis, *Blume, Bijdr.* 1140.—Java.
africana, Mill. Gard. Dict. ed. VIII. n. 4 = Noltea africana.
Alaternus, *Linn. Sp. Pl.* 193.—Reg. Mediterr.; As. Min.
alnifolia, *L'Hérit. Sert. Angl.* 5.—Am. bor.
alnifolia, Pursh, Fl. Am. Sept. i. 166 = Purshiana.
alnoides, S. F. Gray, Nat. Arr. Brit. Pl. ii. 621 = R. Frangula.

RHAMNUS:—
alpina, *Linn. Sp. Pl.* 193.—Europ.; Afr. bor.
alpina, Pall. Fl. Ross. ii. 23 = grandifolia.
alpina, Richards. in Frankl. Journ. App. 734 = alnifolia.
alpina, Sibth. & Sm. Fl. Graec. Prod. i. 158 = fallax.
amygdalina, Desf. Fl. Atlant. i. 198 = oleoides.
amygdalina, Moris, Stirp. Sard. Elench. fasc. i. 11 = persicaefolia.
angulata, Pers. Syn. i. 240 = Zizyphus angulata.
angustifolia, Hort. ex C. Koch, Dendrol. i. 17 = R. Alaternus.
aphylla, Gmel. ex Steud. Nom. ed. II. ii. 443 = Noaea spinosissima.
aragonensis, [Asso,] Syn. Arag. 27 = lycioides.
arenicola, Casar. Nov. Stirp. Bras. Dec. 76 = Scutia arenicola.
arguta, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. (1866) n. XI. 6.—China.
Arnottiana, *Gardn. ex Thw. Enum. Pl. Zeyl.* 74.—Zeylan.
asiatica, Lam. Encyc. iv. 474 = Colubrina asiatica.
asplenifolia, Carr. in Rev. Hortic. (1872) 195 = incana, *Carr.*?
australis, Pers. Syn. i. 241 = Paliurus aculeatus.
balearica, Hort. ex Steud. Nom. ed. II. ii. 443 = R. Alaternus.
balearica, Hort. Par. ex Link, Handb. ii. 121 = R. Frangula?
balearica, *Willk. in Oestr. Bot. Zeitschr.* xxv. (1875) 112.—Hispan.
baptica, Decne. ex Rondot, Vert de Chine, 58, nomen (Quid?).—China.
biflora, D. Dietr. Syn. Pl. i. 807, sphalm. = seq.
biniflora, [Moç. & Sesse,] ex DC. Prod. ii. 26 = Karwinskia Humboldtiana.
Boissieri, Heldr. ex Boiss. Fl. Orient. ii. 18 = prunifolia.
brasiliensis, Spreng. Syst. i. 768 = Ilex pubiflora.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 31.—Japon.
burgundiaca, Desf. Tabl. ed. II. 202 = *hybrida*.
buxifolia, *Poir. Voy. Barb.* ii. 127.—Afr. bor.
californica, *Eschsch. in Mém. Acad. Pétersb.* x. (1823) 285.—Calif.; Mexic.
canadensis, Hort. ex Koch, Dendrol. i. 610 = caroliniana.
capensis, Spreng. Syst. i. 767 = Elaeodendron croceum.
capensis, Thunb. Prod. Pl. Cap. 44 = Scutia Commersonii.
capillaris, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 483 = lycioides.
capreaefolia, *Schlecht. in Linnaea*, xx. (1841) 464.—Mexic.
cardiocarpa, Willd. ex Roem. & Schult. Syst. v. 276 = tinctoria.
cardiosperma, Willd. ex DC. Prod. i. 24 = tinctoria.
carniolica, Kern. in Zeitschr. Ferdinand. Tirol. III. xv. (1870) 282 = saxatilis.
caroliniana, *Walt. Fl. Carol.* 101.—Am. bor.
carpinifolia, Pall. Fl. Ross. i. II. 24. t. 10 = Zelkova crenata.
cassinoides, *Lam. Encyc.* iv. 474.—Ins. S. Doming.
castaneifolia, *Hort. ex Lebas, in Rev. Hortic.* (1867) 306.—Hab.?
cathartica, *Linn. Sp. Pl.* 193.—Europ. As. et Afr. bor.
cathartica, Pall. Reise, iii. 423 = dahurica.
celtifolia, *Thunb. Prod. Pl. Cap.* 44.—Afr. austr.
Chacaye, Domb. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Discaria serratifolia.
chlorophora, Decne. in Compt. Rend. Par. xliv. (1857) 1141 = tinctoria.
cinerascens, *Blume, Bijdr.* 1141.—Java.
circumcissa, Linn. f. Suppl. 152 = Scutia Commersonii.
circumcissa, Russ. ex Wall. Cat. n. 4245 C = Zizyphus Jujuba.
Clusii, Willd. Enum. Hort. Berol. 250 = R. Alaternus.
collina, Salisb. Prod. 139 = R. Alaternus.
colubrina, Jacq. Enum. Pl. Carib. 16; Linn. Sp. Pl. ed. II. 280 = Colubrina ferruginosa.
coriacea, Brouss. in Roem. & Schult. Syst. v. 291; Nees, Hor. Phys. Berol. 114. t. 22 = integrifolia.
coriacea, Buch, in Abh. Berl. Akad. 1816–17 (1819), 382 = racemosa.
coriacea, Willd. ex Spreng. Syst. i. 768 (=*Mespilodaphne glauca*).—Bras.

RHAMNUS:—
cornifolia, *Boiss. & Hohen. Diagn.* Ser. I. ix. 1.—Kurdist.; Persia.
corymbosa, Vahl, ex Roem. & Schult. Syst. v. 294 = Sageretia corymbosa.
costata, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. x. (1866) n. XI. 17.—Japon.
crenata, Baker & S. Moore, in Journ. Linn. Soc. xvii. (1879) 380 = parvifolia.
crenata, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 146.—Japon.
crenulata, *Ait. Hort. Kew.* ed. I. i. 263.—Ins. Teneriff.
Crini, Orph. ex Boiss. Fl. Orient. ii. 20 = fallax.
crocea, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 261.—Am. bor. occ.
cubensis, Jacq. Enum. Pl. Carib. 16 = Ceanothus cubensis.
cubensis, Pers. Syn. i. 240 = Rhus lineatifolia.
cumanensis, *Loefl. It. Hisp.* 182.—Venezuela.
cuneata, Dougl. in Hook. Fl. Bor. Am. i. 124 = Ceanothus cuneatus.
cuneifolia, Poir. Encyc. Suppl. iv. 89 = oleoides.
dahurica, *Pall. Reise*, iii. 721(dauurica).—Ind. or.; As. bor.
Dalanta, Blanco, Fl. Filip. ed. II. 121 = Zizyphus Dalanta.
diffusa, *Clos, in C. Gay, Fl. Chil.* ii. 18.—Chili.
divaricata, Forsk. Fl. Aegypt. Arab. 204 = Zizyphus Spina-Christi.
domingensis, Jacq. Enum. Pl. Carib. 17 = Ceanothus reclinatus.
Drummondii, Young, Fl. Texas, 204 = Colubrina texensis.
elegans, H. B. & K. Nov. Gen. et Sp. vii. 53 = Sageretia elegans.
elliptica, Sw. Prod. Veg. Ind. Occ. 50 = Ceanothus reclinatus.
Ephedra, Domb. ex DC. Prod. ii. 29 = Retanilla Ephedra.
erythroxyla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Erythroxylum.
Erythroxylon, Georgi, Beschr. Russ. Reich. Nachtr. 262; Bieb. Fl. Taur. Cauc. i. 168 = Pallasii.
Erythroxylum, *Pall. Reise*, iii. 722.—Sibir.
fallax, *Boiss. Diagn.* Ser. II. v. 73.—Graecia.
ferrea, Vahl, Symb. Bot. iii. 41 = Scutia ferrea.
ferruginea, Nutt. in Journ. Acad. Philad. vii. (1834) 90 = Colubrina ferruginosa.
filiformis, Roth, Nov. Pl. Sp. 153 = Sageretia oppositifolia.
Frangula, *Linn. Sp. Pl.* 193.—Europ.; As. et Afr. bor.
franguloides, Michx. Fl. Bor. Am. i. 153 = alnifolia.
franguloides, Willd. ex Roem. & Schult. Syst. v. 295 = californica.
glabrata, Heyne, in Wall. Cat. n. 7479 = Zizyphus rugosa.
glandulosa, *Ait. Hort. Kew.* ed. I. i. 265.—Ins. Canar.
globosa, Bunge, Enum. Pl. Chin. Bor. 14 = tinctoria.
globosa, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 146 = japonica.
glomerata, Hook. ex Benth. Pl. Hartw. 9 = Colubrina glomerata.
Goudotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 379.—N. Granat.
graeca, *Boiss. & Reut. Diagn.* Ser. II. v. 74.—Graecia.
graeca, Kotschy, ex Boiss. Fl. Orient. ii. 14 = punctata.
grandifolia, *Fisch. & Mey. ex Ledeb. Fl. Ross.* i. 504.—Reg. Caucas.
guayaquilensis, *H. B. & K. Nov. Gen. et Sp.* vii. 54.—Ecuador.
Guicciardii, Heldr. & Sart. ex Boiss. Fl. Orient. ii. 20 = Sibthorpiana.
hederacea, Delort. ex Nym. Consp. 145 = R. Alaternus.
Heldreichii, *Boiss. Diagn.* Ser. II. i. 120.—As. Min.
heterogenea, Burm. f. Fl. Ind. 61 = Zizyphus Spina-Christi.
hirsuta, Wight & Arn. Prod. 165 = dahurica.
hirtella, *Boiss. Fl. Orient.* ii. 15.—As. Min.
hispanica, Hort. ex C. Koch, Dendrol. i. 617 = R. Alaternus.
horrida, Salisb. Prod. 139 = Paliurus aculeatus.
Humboldtiana, Roem. & Schult. Syst. v. 295 = californica.
humifusa, Schur, Enum. Pl. Transs. 142 = saxatilis.
hybrida ×, L'Hérit. Sert. Angl. 5.

RHAMNUS:—
hydriensis, Hacq. Pl. Alp. t. 2 = pumila.
iguanaea, Jacq. Enum. Pl. Carib. 16; Linn. Sp. Pl. ed. II. 282 = Celtis aculeata.
iguanea, *Vell. Fl. Flum.* 96; ii. t. 147.—Bras.
ilicifolia, Kellogg, in Proc. Calif. Acad. ii. (1863) 37 = crocea.
illyrica, Griseb. ex Pant. in Verh. Ver. Nat.-Heilk. Presb. ii. (1871–72) 110 = graeca.
incana, *Carr. in Rev. Hortic.* Sér. IV. vii. (1858) 657.—Gall. austr.
incana, *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 603.—Ins. Molucc.
inebrians, R. Br. in Salt, Abyss. App. 64 = R. Staddo?
inermis, Gilib. Fl. Lituan. ii. 133 = alpina.
infectoria, Koch, Syn. Fl. Germ. ed. I. 148 = intermedia.
infectoria, *Linn. Mant.* i. 40.—Europ. austr.
infectoria, Tchihat. Asie Min. Bot. Atl. t. 9 = petiolaris.
infusionum, Delile, in Ferr. & Galin. Voy. Abyss. iii. 111 = R. Staddo.
insularis, Kellogg, in Proc. Calif. Acad. ii. (1863) 20 (*insulus*) = crocea.
integrifolia, *DC. Cat. Hort. Monsp.* 136.—Ins. Canar.
integrifolia, Spreng. ex Meissn. in DC. Prod. xv. i. 97 (= *Mespilodaphne glauca*).—Bras.
intermedia, *Steud. & Hochst. in Flora*, x. (1827) 74.—Europ.
japonica, *Maxim. in Mem. Acad. Pétersb.* Sér. VII. x. (1806) n. XI. 11.—Japon.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 647.—Java.
Jujuba, Linn. Sp. Pl. 194 = Zizyphus Jujuba.
kurdica, *Boiss. & Hohen. Diagn.* Ser. I. ii. 3.—Mesopot.; Persia.
laevigata, Vahl, Symb. Bot. iii. 41 = Reynosia latifolia.
lanceolata, *Pursh, Fl. Am. Sept.* i. 166.—Am. bor.
Lando, *Llanos, Fragm.* 57 (Quid?).—Ins. Philipp.
latifolia, *L'Hérit. Sert. Angl.* 5. t. 8.—Ins. Azor.
laurifolia, Nutt. in Torr. & Gray, Fl. N. Am. i. 260 = californica.
leprosa, *Blume, Bijdr.* 1139.—Java.
libanotica, *Boiss. Diagn.* Ser. II. i. 119.—As. Min.; Syria.
Lindeniana, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 379.—N. Granat.
linearis, *Clos, in C. Gay, Fl. Chil.* ii. 19.—Chili.
linearis, Sessé & Moç. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 367 = Adolphia infesta.
lineata, Linn. Cent. Pl. ii. 11; Amoen. Acad. iv. 308 = Berchemia lineata.
lineata, Lour. Fl. Cochinch. 159 = Berchemia Loureiriana.
lineatifolia, Roem. & Schult. Syst. v. 294 = Rhus lineatifolia.
littoralis, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 30 = Paliurus ramosissimus.
longifolia, [*Desf.*] *Dum.-Cours. Bot. Cult.* ed. II. vi. 260.—Am. bor.
longifolia, Mill. Gard. Dict. ed. VIII. n. 3 = saxatilis.
Lotus, Linn. Sp. Pl. 194 = Zizyphus Lotus.
lucida, Roxb. Hort. Beng. 86; Fl. Ind. i. 605 = Scutia Commersonii.
lucida, Salisb. Prod. 139 = Zizyphus sativa.
lycioides, Brot. Fl. Lusit. i. 300 = oleoides.
lycioides, *Linn. Sp. Pl.* ed. II. 279.—Reg. Mediterr. occ.
lycioides, Pall. Fl. Ross. ii. 26 = Pallasii.
Lycium, Scop. Fl. Carn. ed. II. i. 163 = infectoria, saxatilis.
macrophylla, *Carr. in Rev. Hortic.* (1876) 159.—China.
madagascariensis, Pourr. ex Baill. Adansonia, i. (1860–61) 252 = Gelonium adenophorum.
major, Carr. in Rev. Hortic. (1874) 355 = californica?
mauritiana, Soyer-Willem. in Usteri, Ann. Bot. xviii. (1796) 20 = Zizyphus Jujuba.
micrantha, Linn. Syst. ed. X. 937 = Trema micrantha.
microcarpa, *Boiss. Fl. Orient.* ii. 20.—Reg. Caucas.
microphylla, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 295.—Mexic.
microphylla, Roxb. Fl. Ind. i. 613 = Zizyphus rotundifolia.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = infectoria.
minutiflora, Michx. Fl. Bor. Am. i. 154 = Sageretia Michauxii.
montana, Rosten. in Feuill. Jeunes Nat. x. (1879–80) 132, nomen = alpina?

RHAMNUS:—
mucronata, *Schlecht. in Linnaea*, xv. (1841) 465.—Mexic.
myrtifolia, *Willk. in Linnaea*, xxv. (1852) 18.—Hispan.
myrtina, Burm. f. Fl. Ind. 60 = Scutia Commersonii.
mystacina, Ait. Hort. Kew. ed. I. i. 266 = Helinus mystacinus.
Nabeca, Forsk. Fl. Aegypt. Arab. 204 = Zizyphus Spina-Christi.
nana, Carr. in Rev. Hortic. (1874) 355 = californica.
Napeca, Linn. Sp. Pl. 194 = Zizyphus Napeca.
neapolitana, Link, Handb. ii. 120 (Quid?).—Italia.
nemoralis, Salisb. Prod. 139 = R. Frangula.
Nerija, Spreng. Syst. iv. Cur. Post. 86 = Elaeodendron glaucum.
nipalensis, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 640.—Reg. Himal.
nummularia, Burm. f. Fl. Ind. 61 = Zizyphus rotundifolia.
obscura, *Schrank, in Syll. Ratisb.* i. (1824) 202.—Ind. occ.
obtusifolia, Hook. ex Torr. & Gray, Fl. N. Am. i. 685 = Zizyphus obtusifolia.
Oenoplia, Linn. Sp. Pl. ed. II. 279 = seq.
Oenopolia, Linn. l. c. ed. I. 194 = Zizyphus Oenoplia.
oleifolia, Hook. Fl. Bor. Am. i. 123. t. 44 = californica.
oleoides, Lam. Fl. Fr. ii. 545 = infectoria.
oleoides, *Linn. Sp. Pl.* ed. II. 279.—Reg. Mediterr.
oppositifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = Luvunga eleutherandra.
oreigenes, Hance, in Journ. Bot. vii. (1869) 114; viii. (1870) 312 = crenata.
palaestina, *Boiss. Diagn.* Ser. II. i. 119.—Syria; Arab.
Paliurus, Linn. Sp. Pl. 194 = Paliurus aculeatus.
Pallasii, *Fisch. & Mey. in Ind. Sem. Hort.* iv. 46.—Reg. Caucas.
paniculata, Thonn. ex DC. Prod. ii. 30 = Chailletia toxicaria?
parviflora, Klein & Willd. ex Roem. & Schult. Syst. v. 295 = Sageretia oppositifolia.
parvifolia, *Bunge, Enum. Pl. Chin. Bor.* 14.—China.
parvifolia, Torr. & Gray, Fl. N. Am. i. 262 = lanceolata.
parvifolia, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 150 = dahurica.
pauciflora, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 137.—Abyss.
pentapetala, Gilib. Fl. Lituan. ii. 131 = R. Frangula.
pentaphylla, Jacq. Obs. Bot. ii. 17 = Rhus pentaphylla.
persica, *Boiss. Diagn.* Ser. I. ii. 4.—Persia; Reg. Himal.
persicaefolia, *Moris, Stirp. Sard. Elench.* fasc. ii. 2.—Ins. Sardoa.
peruviana, Hort. ex Lam. Encyc. iii. 320 = Maytenus octogona.
petiolaris, *Boiss. & Bal. Diagn.* Ser. II. v. 75.—As. Min.; Syria.
picenensis, *Duval-Jouve, in Mém. Acad. Montp.* ix. (1878), reimpr. 12.—Gallia.
polifolia, Vahl, Symb. Bot. iii. 41 = Phylica ramosissima.
polygama, *Jacq. Enum. Pl. Carib.* 17.—Ind. occ.
polymorpha, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 713 = dahurica.
pontica, *Boiss. Fl. Orient.* ii. 21.—As. Min.
prinifolia, Salisb. Prod. 139 = prinoides.
prinoides, *L'Hérit. Sert. Angl.* vi. t. 9.—Afr. austr.
procumbens, *Edgew. in Trans. Linn. Soc.* xx. (1846) 43.—Reg. Himal.
prunifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 157.—Graecia.
prunifolia, Sieber, ex Steud. Nom. ed. II. ii. 444 = oleoides.
pubescens, Poir. Encyc. iv. 464 = infectoria.
pubescens, Sibth. & Sm. Fl. Graec. iii. 33 = Sibthorpiana.
pubescens, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvi. (1872) 379 = Ceanothus pubescens.
pumila, *Linn. Mant.* i. 49.—Europ.
pumila, Wulf. in Jacq. Coll. ii. 141. t. 11 = rupestris.
punctata, *Boiss. Diagn.* Ser. I. ii. 4.—Syria; Cilicia.
purpurea, *Edgew. in Trans. Linn. Soc.* xx. (1846) 44.—Reg. Himal.
Purshiana, *DC. Prod.* ii. 25.—Am. bor.
pusilla, Tenore, Prod. Fl. Neap. 16 = pumila.

RHAMNUS:—
quitensis, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 295.—Ecuador.
quitensis, Spreng. Syst. i. 769 = Ilex paraguensis.
racemosa, *Duham. Arb. ed. Nov.* iii. 48.—Hab. ?
ramiflora, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
ramosissima, Auct. ex Steud. Nom. ed. I. 686; ed. II. ii. 444 = Paliurus ramosissimus.
recta, Forsk. Fl. Aegypt. Arab. 204 = Zizyphus Spina-Christi.
repens, Page, Prod. Southampt. Bot. Gard. 23; Hort. ex C. Koch, Dendrol. i. 614 = pumila.
Retamilla, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 483 = seq.
Retanilla, Domb. ex DC. Prod. ii. 29 = Retanilla obcordata.
robusta, *Hort.—Cf. Rev. Hortic.* (1879) 100.—Hab. ?
rotundifolia, Dum.-Cours. Bot. Cult. ed. II. iv. 260 = R. Alaternus.
rotundifolia, Pers. Syn. i. 240 = Zizyphus Xylopyrus.
ruffa, *Vell. Fl. Flum.* 96; ii. t. 148.—Bras.
rumeliacea, Frivald. in Flora, xviii. (1835) 332 = rupestris.
rupestris, C. A. Mey. Verz. Pfl. Cauc. 133 = microcarpa.
rupestris, Royle, Illustr. Bot. Himal. 169, nomen = procumbens.
rupestris, *Scop. Fl. Carn.* ed. II. i. 164. t. 5.—Europ. or.
rupestris, Vill. Hist. Pl. Dauph. ii. 538 = pumila.
sanguinea, Pers. Syn. i. 239 = seq.
Sanguino, Orteg. Hort. Matr. Dec. 87 = R. Frangula.
Saquil, Domb. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Discaria serratifolia.
Sarcomphalus, Linn. Syst. ed. X. 937 = Sarcomphalus laurinus.
saxatilis, Host, Syn. 119 = infectoria.
saxatilis, *Jacq. Enum. Stirp. Vindob.* 39, 212.—Europ. austr.
saxatilis, Sibth. & Sm. Fl. Graec. Prod. i. 158 = graeca.
scandens, Hill, Hort. Kew. 453 = Berchemia volubilis.
scandens, Spreng. Syst. i. 768 = Tralliana scandens.
sectipetala, *Mart. Herb. Fl. Bras.* n. 1085.—Bras.
sempervirens, Hort. ex Dum.-Cours. Bot. Cult. ed. II. iv. 258 = *hybrida*.
senticosa, *H. B. & K. Nov. Gen. et Sp.* vii. 54.—Peruv.
serrata, *Willd. ex Roem. & Schult. Syst.* v. 295.—Mexic.
serrulata, H. B. & K. Nov. Gen. et Sp. vii. 51. t. 617 = praec.
Shortii, Nutt. in Journ. Acad. Philad. vii. (1834) 91 = lanceolata.
Sibthorpiana, *Roem. & Schult. Syst.* v. 286.—Graecia.
sicula, Linn. Syst. ed. XII. iii. App. 229 = Argania Sideroxylon.
sinensis, Ser. ex Rondot, Vert de Chine, 53 = dahurica.
Sladdo, Walp. Ann. ii. 267, sphalm. = R. Staddo.
soporifera, Lour. Fl. Cochinch. 158 = Zizyphus sativa.
Spartium, Domb. ex Roem. & Schult. Syst. v. 512 = Discaria serratifolia.
spathulaefolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 46.—Reg. Caucas.
sphaerosperma, *Sw. Prod. Veg. Ind. Occ.* 50.—Ind. occ.
spiciflora, *A. Rich. Tent. Fl. Abyss.* i. 138.—Abyss.
Spina-Christi, Linn. Sp. Pl. 195 = Zizyphus Spina-Christi.
spinosa, Gilib. Fl. Lituan. ii. 132 = cathartica.
spinosa, *Hemsl. Diagn. Pl. Nov.* 6.—Panama.
splendens, *Blume, Bijdr.* 1140.—Java.
Staddo, *A. Rich. Tent. Fl. Abyss.* i. 138.—Abyss.
subsempervirens, Hort. ex C. Koch, Dendrol. i. 614 = pumila.
surinamensis, *Scop. Delic. Pl. Insubr.* iii. 10. t. 5.—Guian.
sylvaticus, Serres, in Bull. Soc. Bot. Fr. iv. (1857) 436 = cathartica.
Talanai, *Blanco, Fl. Filip.* ed. I. 170.—Ins. Philipp.
Teidis, Brouss. ex Roem. & Schult. Syst. v. 291, in obs. = integrifolia.
terminalis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 190 = Sageretia oppositifolia.
terniflora, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 26.—Mexic.
tetragona, Linn. f. Suppl. 153; Thunb. Prod. Pl. Cap. 42 = Cassine scandens.

RHAMNUS:—
texensis, Torr. & Gray, Fl. N. Am. i. 263 = Colubrina texensis.
Thea, Osbeck, It. 232 = Sageretia theezans.
theezans, Linn. Mant. ii. 207 = Sageretia theezans.
tinctoria, Mutel, Fl. Fr. i. 218 = infectoria.
tinctoria, *Waldst. & Kit. Pl. Rar. Hung.* iii. 283. t. 255.—Europ. or.; As. Min.
tomentella, Benth. Pl. Hartw. 303 = californica.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 21 = tinctoria.
trigyna, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 190 = Sageretia oppositifolia.
trinervis, Cav. Ic. vi. 3 = Zizyphus exserta.
tripartita, Ucria, in Roem. Arch. i. 1. (1796) 68 = Rhus albida.
triquetra, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 376.—Reg. Himal.
ulmoides, Gueldenst. ex Ledeb. Fl. Ross iii. 645 = Zelkova crenata.
umbellata, Cav. Ic. vi. 2. t. 504 = californica.
uninervis, Dulac, Fl. Hautes-Pyr. 333 = oleoides.
utilis, Decne. in Compt. Rend. Par. xliv. (1857) 1141 = dahurica.
valentina, Willd. Sp. Pl. i. 1096 = pumila.
velutina, Boiss. Elench. Pl. Nov. Hisp. 29 = lycioides.
venosa, Lam. ex Poir. Encyc. iv. 474 = Ceanothus reclinatus.
Villarsi, Link, Handb. ii. 120 = pumila.
Villarsii, Jord. Obs. Pl. Crit. vii. (1849) 18 = infectoria.
virgata, Maxim. in Mém. Acad. Pétersb. Sér. VII. x. (1866) n. xi. 13 = parvifolia.
virgata, Roxb. Hort. Beng. 17; Fl. Ind. i. 604 = dauurica.
virginicus, Page, Prod. Bot. Southamp. Gard. 23 (Quid ?).—Am. bor.
vitiensis, Benth. Fl. Austral. i. 413 = Dallachya vitiensis.
Vitis-idaea, Burm. f. Fl. Ind. 61 = Breynea rhamnoides.
volubilis, Linn. f. Suppl. 132 = Berchemia volubilis.
vulgaris, Pers. Syn. i. 240 = Zizyphus sativa.
Wichellii, Hort. ex C. Koch, Dendrol. i. 604 = cathartica.
Wicklia, Hort. ex C. Koch, l. c. = cathartica.
Wightii, *Wight & Arn. Prod.* 164.—Ind. or.
Wihhor, Luce, in Ind. Hort. Dorpat. (1824) 6; et in Topog. Nachr. Oesel, 62 = cathartica.
Willdenowiana, Roem. & Schult. Syst. v. 295 = longifolia.
Wulfeni, [Hoppe, in] Flora, vi. (1823) 173 = rupestris.
Wulfenii, Roth, Enum. Pl. Phan. Germ. i. 745 = pumila.
xanthocarpa, Hort. ex C. Koch, Dendrol. i. 605 = cathartica.
xylopyra, Retz. Obs. ii. 11 = Zizyphus Xylopyrus.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 477.—Afr. austr.
Zeyheri, Spreng. ex Sond. l. c. 230, Add. 20 = Elaeodendron croceum.
zizyphoides, Spreng. Fl. Hal. Mant. 37 = Alphitonia excelsa.
Zizyphus, Linn. Pl. Pl. 194 = Zizyphus sativa.
zonulata, Blanco, Fl. Filip. ed. I. 172 = Zizyphus Xylopyrus.

RHAMPHICARPA, Benth. in Hook. Comp. Bot. Mag. i. (1835) 368. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 969.
MACROSIPHON, Hochst. in Flora, xxiv. (1841) 373.
Cameroniana, *Oliver, in Journ. Linn. Soc.* xiv. (1876) 95.—Afr. trop.
curviflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 368.—Madag.
fistulosa, Benth. in DC. Prod. x. 504 = longiflora.
Herzfeldiana, *Vatke, in Linnaea*, xliii. (1880–82) 311.—Afr. trop.
Heuglinii, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 100.—Afr. trop.
humilis, Hochst. in Flora, xxvii. (1844) 101 = Cycnium humifusum.
longiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 368.—Ind. or.; Austral.
recurva, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 122.—Afr. trop.
serrata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 228.—Afr. trop.
tubulosa, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 368.—Afr. austr.
veronicaefolia, *Vatke, in Linnaea*, xliii. (1880–82) 312.—Afr. trop.

RHAMPHIDIA, Lindl. in Journ. Linn. Soc. i. (1857) 181 = **Hetaeria**, Blume (Orchid.).
alsinefolia, Lindl. l. c. 182 = Myrmechis gracilis.
discoidea, Reichb. f. in Linnaea, xli. (1877) 59.—N. Caled.
elongata, Lindl. in Journ. Linn. Soc. i. (1857) 181 = H. elongata.
Gardneri, Thw. Enum. Pl. Zeyl. 313 = H. Gardneri.
grandiflora, Lindl. in Journ. Linn. Soc. i. (1857) 182 = Myrmechis glabra.
Mannii, Reichb. f. Otia Bot. Hamb. 109.—Afr. trop.
ovalifolia, Lindl. in Journ. Linn. Soc. i. (1857) 181 = H. ovalifolia.
rubens, Lindl. l. c. 182 = H. rubens.
rubicunda, Reichb. f. in Seem. Fl. Vit. 294 = Goodyera rubicunda ?
scripta, Reichb. f. in Linnaea, xli. (1877) 59.—N. Caled.
tenuis, Lindl. in Journ. Linn. Soc. i. (1857) 182 = H. tenuis.
Vieillardi, Reichb. f. in Linnaea, xli. (1877) 58.—N. Caled.

RHAMPHOSPERMUM, Andrz. ex Bess. Enum. Pl. Volh. 83 (1822) = **Brassica**, Tourn. (Crucif.).
album, Andrz. ex Reichb. in Moessl. Handb. ed. II. ii. 1186 = B. alba.
arvense, Andrz. ex Bess. Enum. Pl. Volh. 83 = B. Sinapis.
orientale, Andrz. l. c. = B. Sinapis.
volgense, Andrz. ex Rupr. Fl. Ingr. 97 = B. juncea.

RHANTERIUM, Desf. Fl. Atlant. ii. 291. t. 240 (1799). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
adpressum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 252.—Alger.
epapposum, *Oliver, in Hook. Ic. Pl.* t. 1367.—Beluchist.
intermedium, Pomel, Nouv. Mat. Fl. Atl. 37 = adpressum.
suaveolens, *Desf. Fl. Atlant.* ii. 291. t. 240.—Afr. bor.

RHAPHANOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 65, 133 = **Raphanus**, Tourn. (Crucif.).
landrus, St. Lag. l. c. 133 = Raphanus Landra.

RHAPHEDOSPERA, Wight, Ic. t. 1554 (1850), sphalm. = RHAPHIDOSPORA, Nees = **Justicia**, Houst. (Acanth.).

RHAPHIA, Auct. ex Steud. Nom. ed. II. ii. 445 (1841) = **Raphia**, Beauv. (Palm.).

RHAPHIDIOCYSTIS, Hook. f. in Oliver, Fl. Trop. Afr. ii. 553 (1871) = **Raphidiocystis**, Hook. f. (Cucurb.).

RHAPHIDOPHORA, Hassk. in Flora, xxv. (1842) II. Beibl. 11, partim; Schott, Gen. Aroid. t. 77 (1858). *AROIDEAE*, Benth & Hook. f. iii. 992.
RAPHIDOPHORA, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 168.
affinis, *Schott, Prod. Aroid.* 385.—Reg. Himal.
amplissima, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 129 = Epipremnum amplissimum.
angulata, Schott, Prod. Aroid. 379 = montana.
angustata, Engl. in DC. Monog. Phan. ii. 241 = montana.
angustata, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 128.—Sumatra.
angustifolia, *Schott, in Bonplandia*, v. (1857) 45.—Malaya; N. Guin.
Beccarii, *Engl. Bot. Jahrb.* i. (1881) 181.—Borneo.
Calophyllum, Schott, Prod. Aroid. 380 = R. Peepla.
caudata, Schott, l. c. 382 = Epipremnum mirabile.
conica, *Engl. Bot. Jahrb.* i. (1881) 181.—N. Guin.
Cunninghamii, Schott, in Bonplandia, ix. (1861) 367 = Epipremnum mirabile ?
decursiva, *Scott, l. c.* v. (1857) 45.—Ind. or.
dilacerata, Koch, in Regel, Gartenfl. (1864) 5. t. 423 = decursiva.
eximia, *Schott, in Bonplandia*, v. (1857) 45; *Prod. Aroid.* 387.—Reg Himal.
fallax, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 128 = montana.
glauca, *Schott, in Bonplandia*, v. (1857) 45.—Reg. Himal.
grandis, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 349 = eximia.

RHAPHIDOPHORA:—
honkongensis, Schott, Prod. Aroid. 378 = angustifolia.
Hookeri, *Schott, l. c.* 381.—Reg. Himal.
Huegelii, Schott, l. c. 384 = Epipremnum medium.
insignis, Schott, Gen. Aroid. t. 77 = eximia?
Korthalsiana, Herb. Lugd. Bat. ex Engl. in DC. Monog. Phan. ii. 250 = Epipremnum medium.
Korthalsii, *Schott, in Miq. Ann. Mus. Lugd. Bat.* i. 129.—Malaya.
lacera, Hassk. in Flora, xxv. (1842) II. Beibl. 11 = Epipremnum mirabile.
lancifolia, *Schott, Prod. Aroid.* 380.—Reg. Himal.
lingulata, *Schott, in Bonplandia*, v. (1857) 45.—Java.
Lobbii, *Schott, Prod. Aroid.* 379.—Malaya.
maxima, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 269.—Borneo.
megastigma, *Engl. Bot. Jahrb.* i. (1881) 180.—Sumatra.
montana, *Schott, in Miq. Ann. Mus. Lugd. Bat.* i. 128.—Java.
oblongifolia, Schott, l. c. 128 = spathacea.
Peepla, *Schott, in Bonplandia*, v. (1857) 45.—Ind or.; Malaya.
Peepla, Thw. Enum. Pl. Zeyl. 336 = pertusa.
pertusa, *Schott, Prod. Aroid.* 382.—As. trop. [Austral.; Ins. Pacif.].
pinnata, Schott, in Bonplandia, v. (1857) 45 = Epipremnum mirabile.
pinnatifida, Schott, l. c = Epipremnum mirabile?
pteropoda, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 268.—N. Guin.
puberula, *Engl. Bot. Jahrb.* i. (1881) 180.—Sumatra.
spathacea, *Schott, in Miq. Ann. Mus. Lugd. Bat.* i. 129.—Java.
Storckiana, *Schott, in Bonplandia*, x. (1862) 346.—Ins. Fiji.
sylvestris, Engl. in DC. Monog. Phan. ii. 239 = angustifolia.
tenuis, *Engl. Bot. Jahrb.* i. (1881) 181.—Borneo.
vitiensis, Schott, in Bonplandia, ix. (1861) 367 = Epipremnum mirabile.
Wallichii, Schott, Prod. Aroid. 383 = Epipremnum mirabile.
Zippeliana, Schott, in Miq. Ann. Mus. Lug. Bat. i. 129 = Epipremnum asperatum.

RHAPHIDOPHYLLUM, Benth. in DC. Prod. x. 521 (1846) = RAPHIDOPHYLLUM, Hochst. = **Sopubia**, Buch.-Ham. (Scrophul.).

RHAPHIDOSPERMA, G. Don, in Sweet, Hort. Brit. ed. III. 559 (1839), sphalm. = seq.

RHAPHIDOSPORA, Nees, in Wall. Pl. As. Rar. iii. 115 (1832) = **Justicia**, Houst. (Acanth.).
abyssinica, Nees, in DC. Prod. xi. 500 = J. glabra.
ciliata, Nees, l. c. = J. pubigera.
cordata, Nees, l. c. 499 = J. cordata.
dichotoma, Nees, l. c. 500 = Dianthera dichotoma.
glabra, Nees, in Wall. Pl. As. Rar. iii. 115 = J. glabra.
leptántha, Nees, in DC. Prod. xi. 501 = J. leptantha.
membranifolia, Miq. Fl. Ind. Bat. ii. 850.—Java.
tenella, T. Anders. in Journ. Linn. Soc. ix. (1867) 515 = Dianthera collina.
tenella, Nees, in DC. Prod. xi. 501.—Java.
vestita, G. Don, in Sweet, Hort. Brit. ed. III. 559 = Nelsonia campestris.

RHAPHIODON, Schau. in Flora, xxvii. (1844) 345 = **Hyptis**, Jacq. (Labiat.).
Echinus, Schau. l. c. = H. Sideritis.

RHAPHIOLEPIS, Poir. in Dict. Sc. Nat. xv. 314 (1827) = **Raphiolepis**, Lindl. (Rosac.).

RHAPHIONACME, C. Muell. in Walp. Rep. vi. 480 (1846–47) sphalm. = **Raphionacme**, Harv. (Asclep.).

RHAPHIOPHALLUS, Schott, Gen. Aroid. t. 27 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 972.
Hohenackeri, *Schott, l. c.*—Ind. or.

RHAPHIOSTYLIS, Planch. ex Benth. in Hook. Niger Fl. 259 (1849) = **Apodytes**, E. Mey. (Olacin.).
beninensis, Planch. ex Benth. l. c. 259 = A. beninensis.
Heudelotii, Planch. ex Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 391 = A. beninensis.

RHAPHIS, Lour. Fl. Cochinch. 552 (1790) = **Chrysopogon**, Trin. (Gramin.).
acicularis, Desv. Opusc. 69 = C. aciculatus.
arguta, Nees, ex Steud. Syn. Pl. Gram. 391 = C. argutus.
arundinacea, Desv. Opusc. 69 = Sorghum halepense.
arundinacea, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 499.—Ins. Philipp.
ciliolata, Nees, ex Steud. Syn. Pl. Gram. 396 = C. serrulatus.
coerulescens, Desv. Opusc. 69 = C. violascens.
coeruleus, Nees, in Royle, Illustr. Bot. Himal. 417, nomen; et ex Steud. Syn. Pl. Gram. 395 = C. serrulatus.
echinulata, Nees, ll. cc. 417; 397 = C. Gryllus.
glauca, Nees, ex Steud. l. c. = C. montanus.
Gryllus, Desv. Opusc. 69 = C. Gryllus.
javanica, Nees, ex Steud. Syn. Pl. Gram. 396 = C. aciculatus.
microstachya, Nees, in Royle, Illustr. Bot. Himal. 417; et ex Steud. Syn. Pl. Gram. 397 = C. violascens.
muricata, Nees, ll. cc. 416; 394 = Andropogon squarrosus.
orientalis, Desv. Opusc. 69 = C. Wightianus.
repens, Nees, ex Steud. Syn. Pl. Gram. 397 = C. montanus.
Royleana, Nees, l. c. = C. Gryllus.
stricta, Nees, in Hook. Kew Journ. ii. (1850) 99 = Andropogon intermedius.
trivalvis, Kunth, Enum. Pl. i. 505, sphalm. = seq.
trivialis, Lour. Fl. Cochinch. 553 = C. aciculatus.
villosula, Nees, ex Steud. Syn. Pl. Gram. 397 = C. violascens.
Wightiana, Nees, ex Steud. l. c. 395 = C. Wightianus.
zeylanica, Nees, ex Steud. l. c. 397 = C. Arnottianus.

RHAPHIS, Walp. Ann. iii. 471 (1852), sphalm. = **Rhapis**, Linn. f. (Palm.).

RHAPHISPERMUM, Benth. in DC. Prod. x. 509 (1848). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 973.
gerardioides, *Benth. in DC. Prod.* x. 509.—Madag.

RHAPHISTEMMA, Meissn. Gen. 268 (1840) = **Raphistemma**, Walt. (Asclep.).

RHAPHISTEMUM, Walp. Ann. i. 506 (1848–49) sphalm. = praec.

RHAPHITAMNUS, Miers, in Trans. Linn. Soc. xxvii. (1870) 95. t. 26. *VERBENACEAE*, Benth. & Hook. f. ii. 1149.
POEPPIGIA, Bert. ex Bull. Férussac, xxiii. (1830) 109.
amoenus, *Miers, in Trans. Linn. Soc.* xxvii. (1870) 97.—Chili.
buxifolius, *Miers, l. c.* 98.—Chili.
cyanocarpus, *Miers, l. c.* 96. t. 26.—Chili.
longiflorus, *Miers, l. c.* 98.—Chili.
parvifolius, *Miers. l. c.* 99.—Chili.
serratifolius, *Miers, l. c.* 99.—Chili.

RHAPIDOPHYLLUM, H. Wendl. & Drude, in Bot. Zeit. xxxiv. (1876) 803. *PALMAE*, Benth. & Hook. f. iii. 925.
Hystrix, *H. Wendl. & Drude, l. c.*—Florida.

RHAPIDOSPORA, Reichb. Nom. 110 (1841), sphalm. = RHAPHIDOSPORA, Nees = **Justicia**, Houst. (Acanth.).

RHAPIS, Linn. f. in Ait. Hort. Kew. ed. I. iii. 473 (1789). *PALMAE*, Benth. & Hook. f. iii. 930.
RHAPHIS, Walp. Ann. iii. 471 (1852), sphalm.
acaulis, Willd. Sp. Pl. iv. 1093 = Sabal Adansoni.
arundinacea, Ait. Hort. Kew. ed. I. iii. 474; ed. II. v. 474 = Rhapidophyllum Hystrix.
aspera, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 624 (Quid?).—Hab.?
caroliniana, Hort. ex Kunth, Enum. Pl. iii. 246 = Rhapidophyllum Hystrix.
cochinchinensis, *Mart. Hist. Nat. Palm.* iii. 254.—Cochinch.
cordata, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 624 (Quid?).—Hab.?
flabelliformis, *L'Hérit. ex Ait. Hort. Kew.* ed. I. iii. 473.—China; Japon.

RHAPIA:—
humilis, *Blume, Rumphia*, ii. 54.—China.
javanica, *Blume, l. c.* 56.—Java.
Kwanwortsik, H. Wendl. Ind. Palm. (1854) 34 = flabelliformis.
major, *Blume, Rumphia*, ii. 55.—China.
Sirotsik, Hort. ex H. Wendl. in Kerch. Palm. 255 = humilis.

RHAPONTICA, Hill, Veg. Syst. iv. 47 (1762) = seq.

RHAPONTICUM, [Vaill.] Hall. Enum. Stirp. Helv. ii. 687 (1742) = **Centaurea**, Linn. (Compos.).
acaule, DC. Prod. vi. 664 = C. Chamaerhaponticum.
argentatum, Moench, Meth. 562 = C. glastifolia.
atriplicifolium, DC. Prod. vi. 663 = Serratula atriplicifolia.
Behen, Kostel. Allg. Med. Pharm. Fl. ii. 606 = C. Behen.
Calcitrapa, Scop. Fl. Carn. ed. II. ii. 136 = C. Calcitrapa.
canariense, DC. Prod. vi. 664 = C. cynaroides.
caspium, Karel. ex Ledeb. Fl. Ross. ii. 751 = *nitidum*.
caulescens, Coss. & Bal. in Bull. Soc. Bot. Fr. xx. (1873) 251; xxii. (1875) 62 = C. Cossoniana.
centauroides, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1440 = C. centauroides.
ciliatum, Lam. Fl. Fr. ii. 39 = C. nigra.
cynaroides, Less. Syn. Comp. 6 = Cnicus cynaroides.
dahuricum, Turcz. in Bull. Soc. Nat. Mosc. xi. (1838) 96 = C. monantha.
davuricum, Ledeb. Fl. Ross. ii. 751 = praec.
deltoidum, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1440 = Serratula atriplicifolia.
doronicaefolium, Fisch. ex Sweet, Hort. Brit. ed. II. 287.—Sibir.
eriophorum, Scop. Fl. Carn. ed. II. ii. 135 = C. eriophora.
heleniifolium, Gren. & Godr. Fl. Fr. ii. 239 = C. Rhaponticum.
Jacea, Scop. Fl. Carn. ed. II. ii. 136 = C. Jacea.
karatavicum, Regel & Schmalh. in Act. Hort. Petrop. vi. (1879) 320.—Turkestan.
lyratum, Nym. Consp. 416 = C. Rhaponticum.
nitidum, Fisch. in DC. Prod. vi. 664.—Reg. Casp.
Pallasii, G. Don, ex Loud. Hort. Brit. 360, nomen = C. Rhaponticum.
paniculatum, Scop. Fl. Carn. ed. II. ii. 136 = C. paniculata.
pulchrum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 36 = C. pulchra.
pungens, Franch. & Sav. Enum. Pl. Jap. ii. 415 = Serratula atriplicifolia.
pusillum, Boiss. Fl. Orient. iii. 592 = C. pygmaea.
pygmaeum, DC. Prod. vi. 665 = C. pygmaea.
salinum, Less. in Linnaea, ix. (1835) 179 = C. salina.
scariosum, Lam. Fl. Fr. ii. 38 = C. Rhaponticum.
uniflorum, DC. Diss. Comp. 33 = C. monantha.

RHAPTOCARPUS, Miers, Apocyn. S. Am. 151 (1878) = **Echites**, P. Br. (Apocyn.).
apiculatus, Miers, l. c. 153 = E. coalita.
coalitus, Miers, l. c. 152 = E. coalita.
didymus, Miers, l. c. = E. didyma.
Martii, Miers, l. c. = E. Martii.
odoriferus, Miers, l. c. 151 = E. odorifera.

RHAPTOMERIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 41 = **Cyclea**, Arn. (Menisp.).
Burmanni, Miers, l. c. = C. Burmannii, peltata.

RHAPTONEMA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xx. (1867) 261. *MENISPERMACEAE*.
cancellata, *Miers, l. c.* 262.—Madag.

RHAPTOPETALUM, Oliver, in Journ. Linn. Soc. viii. (1865) 159. t. 12. *OLACINEAE*, Benth. & Hook. f. i. 995.
coriaceum, *Oliver, l. c.* 160.—Afr. trop.
Soyauxii, *Oliver, in Hook. Ic. Pl.* t. 1405.—Afr. trop.

RHAPTOSTYLUM, Humb. & Bonpl. Pl. Aequin. ii. 139. t. 125 (1809) = **Heisteria**, Jacq. (Olacin.).
acuminatum, Humb. & Bonpl. l. c.—N. Granat.
pentandrum, Benth. ex Reiss. in Mart. Fl. Bras. xi. 1. 77 = H. pentandra.

RHAZYA, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 80. *APOCYNACEAE*, Benth. & Hook. f. ii. 703.
orientalis, *A. DC. in DC. Prod.* viii. 386.—Graecia.
stricta, *Decne. in Ann. Sc. Nat.* Sér. II. iv. (1835) 80. —Arabia; Afghan.

RHEA, Endl. Ench. 241, in syn. (1841) = REA, Bert. = **Dendroseris**, D. Don (Compos.).

RHEEDEA, Linn. Gen. ed. I. 364 (1737) = seq.

RHEEDIA, Linn. Sp. Pl. ed. I. 1193 (1753). *GUTTIFERAE*, Benth. & Hook. f. i. 175.
CHLOROMYRON, Pers. Syn. ii. 73 (1807).
LAMPROPHYLLUM, Miers, in Trans. Linn. Soc. xxi. (1853) 249. t. 26.
VERTICILLARIA, Ruiz & Pav. Prod. Fl. Per. 81. t. 15 (1794).
acuminata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 314.—Peruv.
americana, Hort. ex Steud. Nom. ed. II. ii. 446 = lateriflora.
aristata, *Griseb. Cat. Pl. Cub.* 38.—Ind. occ.
Benthamiana, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 320 = macrophylla.
brasiliensis, *Planch. & Triana, l. c.* 310.—Bras.
calyptrata, *Planch. & Triana, l. c.* 311.—Bras.
Commersonii, *Planch. & Triana, l. c.* 312.—Madag.
Cupi, *Planch. & Triana, l. c.* 323.—Am. austr.
edulis, *Planch. & Triana, l. c.* 310.—Panama.
elliptica, *Wright, in Sauv. Fl. Cub.* 9.—Cuba.
floribunda, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 319.—Bras.
fruticosa, *Wright, ex Griseb. Cat. Pl. Cub.* 38.—Cuba.
Gardneriana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 321.—Bras.
javanica, Burm. f. Fl. Ind. 118 (Quid?).—Java.
lateriflora, Bert. ex DC. Prod. i. 516 = Sloanea Berteriana.
lateriflora, *Linn. Sp. Pl.* 1193.—Am. trop.
longifolia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 322.—Bras.
macrophylla, *Planch. & Triana, l. c.* 309.—Bras.
Madruno, *Planch. & Triana, l. c.* 315.—N. Granat.
parviflora, *Planch. & Triana, l. c.* 323.—Guian.
Pervillei, *Planch. & Triana, l. c.* 312.—Madag.
pulvinata, *Planch. & Triana, l. c.* 320.—N. Granat.
ruscifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 167.—Ind. occ.
Sieberi, *Choisy, in Mém. Soc. Phys. Genèv.* xii. (1851) 427.—Ins. Trinit.
Smeathmanii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 312.—Afr. trop.
virens, *Planch. & Triana, l. c.* 317.—Guian.

RHEITHROPHYLLUM, Hassk. Cat. Hort. Bog. Alt. 123 (1844) = seq.

RHEITROPHYLLUM, Hassk. in Flora, xxv. (1842) II. Beibl. 56 = **Aeschynanthus**, Jack (Gesner.).
subverticillatum, Hassk. l. c. = A. angustifolia.

RHEKTOPHYLLUM, N. E. Br. in Journ. Bot. xx. (1882) 194. t. 230. *AROIDEAE*, Benth. & Hook. f. iii. 981.
mirabile, *N. E. Br. l. c.* 195.—Afr. trop.

RHESA, Walp. Rep. i. 538 (1842), sphalm. = BHESA, Buch.-Ham. = **Kurrimia**, Wall. (Celastr.).

RHETINODENDRON, Meissn. Gen. 216; Comm. 136 (1839). *COMPOSITAE* [Benth. & Hook. f. ii. 442].
BALBISIA, DC. in Guill. Arch. Bot. ii. (1833) 333.
INGENHOUZIA, Bert. ex DC. Prod. vi. 447 (1837).
VENDREDIA, Baill. Hist. Pl. viii. 264 (1882).
Berteroi, *Hemsl. Bot. Voy. Challenger*, i. III. 39.—Ins. J. Fernand.

RHETINOLEPIS, Coss. in Bull. Soc. Bot. Fr. iii. (1856) 707 = **Anthemis**, Mich. (Compos.).
lonadioides, Coss. l. c. 708.—Alger.

RHETINOPHLOEUM, Karst. Fl. Columb. ii. 25. t. 113 (1862?) = **Cercidium**, Tul. (Legumin.).
viride, Karst. l. c.—N. Granat.

RHEUM, Linn. Syst. ed. I (1735). *POLYGONACEAE*, Benth. & Hook. f. iii. 100.
acuminatum, *Hook. f. & Thoms. in Bot. Mag.* t. 4877. —Reg. Himal.
aplostachyum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 442 = R. Rhaponticum.
australe, D. Don, Prod. Fl. Nep. 75 = Emodi.
austriacum, Hort. ex G. Don, in Loud. Hort. Brit. 161, nomen (Quid?).—Europ.
caspicum, Pall. in Nov. Act. Acad. Petrop. x. (1797) 382 = tataricum.
Collinianum, *Baill. in Rev. Hortic.* (1883) 420.—China.
compactum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 442 = R. Rhaponticum.
compactum, *Linn. Sp. Pl.* ed. II. 531.—Mongol.
cordatum, Hort. ex Steud. Nom. ed. II. ii. 446 (Quid?). —Hab.?
crassinervium, *Fisch. ex Lindl. Med. Fl.* 950.—Hab.?
crispum, Hort. ex G. Don. in Loud. Hort. Brit. 161, nomen = Rumex crispus?
cruentum, *Siev. in Pall. N. Nord. Beitr.* vii. 294.—Sibir. altaic.
cuneatum, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxvii. (1864) I. 387, nomen.—Turkestan.
dentatum, Mart. Ind. Sem. Hort. Monac. (1839).—Cf. Linnaea, xiv. (1840) Litt. 135 = hybridum.
digynum, Wahlenb. Fl. Lapp. 101. t. 9. f. 2 = Oxyria digyna.
Emodi, *Wall. Cat.* n. 1727.—Reg. Himal.
Emodi, Wall. l. c. 1727 C = Webbianum.
Emodium, Wall. ex Nees & Eberm. Med.-Pharm. Bot. ii. 455 = Emodi.
esculentum, Salisb. Prod. 257 = R. Rhaponticum.
Fedtschenkoi, *Maxim. ex Regel, Pl. Nov. Fedsch.* 77. —Turkestan.
fenestratum, Loud. Hort. Brit. 161 (Quid?).—Hab.?
Franzenbachii, *Muent. in Act. Congr. Bot. Amst.* 1877 (1879) 212.—As. temp.
hybridum, *Murr. in Nov. Comm. Gotting.* ii. v. 1774 (1775) 50. t. 11.—Mongolia.
leucorrhizum, *Pall. in Nov. Act. Acad. Petrop.* x. (1797) 381.—Sibir.
macropterum, *Mart. Ind. Sem. Hort. Monac.* (1848); *ex Meissn. in DC. Prod.* xiv. 33.—Hab.?
microcarpum, *Meissn. l. c.*—Hab.?
Moorcroftianum, Meissn. l. c. 36 = spiciforme.
Moorcroftianum, *Royle, Illustr. Bot. Himal.* 315, 318. —Reg. Himal.
muricatum, Blanco, Fl. Filip. ed. I. 327 = undulatum.
nanum, Sievers, ex Pall. in Nov. Act. Acad. Petrop. x. (1797) 381 = leucorrhizum.
nobile, *Hook. f. & Thoms. Illustr. Himal. Pl.* t. 19.—Reg. Himal.
nutans, Pall. Fl. Ross. ii. I. (Index) = compactum.
officinale, *Baill. Adansonia*, x. (1871–72) 246; xi. (1873–76) 229.—Tibet.
palmatum, *Linn. Syst.* ed. X. 1010.—Mongol.
pumilum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 503.—China.
racemiferum, *Maxim. l. c.*—Mongol.
Rhabarbarum, Linn. Sp. Pl. 372 = undulatum.
Rhaponticum, Delarb. Fl. Auv. ed. II. 527 = Rumex alpinus.
Rhaponticum, S. G. Gmel. ex Ledeb. Fl. Ross. iii. 497 = tataricum.
Rhaponticum, *Linn. Sp. Pl.* 371.—Sibir.
rhizostachyum, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 254.—Soongaria.
Ribes, *Linn. Sp. Pl.* 372.—Syria; Persia; Afghan.
rotundatum, Stokes, Bot. Mat. Med. ii. 429 = R. Rhaponticum.
rotundifolium, Fisch. ex Sweet, Hort. Brit. ed. II. 440 (Quid?).—Hab.?
rugosum, *Desf. Cat. Hort. Par.* ed. III. 389.—Hab.?
sanguineum, Hort. Halens. ex Meissn. in DC. Prod. xiv. 33 = macropterum.
sibiricum, Pall. N. Nord. Beitr. vii. 264; Fl. Ross. ii. I. (Index) = R. Rhaponticum.
songaricum, *Schrenk. in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 114.—Soongaria.
spiciforme, *Royle, Illustr. Bot. Himal.* 318. t. 78.—Reg. Himal.
tataricum, *Linn. f. Suppl.* 229.—Tartaria.
tetragonopus, *Mart. Ind. Sem. Hort. Monac.* (1839). —*Cf. Linnaea*, xiv. (1840) *Litt.* 135.—Hab.?
undulatum, *Linn. Sp. Pl.* ed. II. 531.—As. med.

RHEUM:—
undulatum, Pall. Reise, iii. 317, 322 = R. Rhaponticum.
uninerve, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 503.—Mongol.
Webbiana, *Royle, Illustr. Bot. Himal.* 318. t. 78 A.—Reg. Himal.

RHEXIA, Linn. Coroll. Gen. 6. (1737). *MELASTOMACEAE*, Benth. & Hook. f. i. 747.
acerosa, Schrank & Mart. ex. DC. Prod. iii. 125 = Marcetia acerosa.
Acisanthera, Linn. Syst. ed. X. 998 = Acisanthera quadrata.
Acisanthera, Willd. Sp. Pl. ii. 303 = Acisanthera recurva.
acuminata, Schrank, ex DC. Prod. iii. 139 = Pyramia pityrophylla.
adamantium, A. St. Hil. ex Humb. & Bonpl. Rhex. 157. t. 60 = Cambessedesia Hilariana.
agrestis, Mart. & Schrank, ex DC. Prod. iii. 126 = Microlicia agrestis.
Agrostemma, Mart. & Schrank, l. c. 133 = Tibouchina gracilis.
alata, Raddi, in Mem. Soc. Ital. Mod. xviii. Fis. (1820) 387 = Tibouchina granulosa.
alba, Mart. & Schrank, ex DC. Prod. iii. 104 = Lavoisiera alba.
Alifanus, Walt. Fl. Carol. 130 = glabella.
alpestris, Mart. ex DC. Prod. iii. 140 = Pterolepis alpestris.
alsinefolia, Schrank & Mart. l. c. 118 = Acisanthera alsinaefolia.
alternans, Schrank & Mart. l. c. 120 = Microlicia serpyllifolia.
amazonum, Schrank, ex DC. l. c. 123 (= *Pachyloma coriaceum*).—Bras.
angustifolia, Nutt. Gen. Am. i. 244 = lanceolata.
angusturensis, Humb. & Bonpl. Rhex. 77. t. 29 = Pterolepis glomerata.
anomala, Schrank & Mart. ex DC. Prod. iii. 109 = Macairea rufescens.
aquatica, Sw. Fl. Ind. Occ. ii. 650; Humb. & Bonpl. Rhex. 104. t. 40 = Nepsera aquatica.
arcuata, Cogn. in Mart. Fl. Bras. xiv. III. 502 (Index), lapsu = *armata*.
arenaria, Schrank & Mart. ex DC. Prod. iii. 114 = Siphanthera arenaria.
arenariaefolia, Mart. & Schrank, l. c. 120 = Microlicia arenariaefolia.
armata, Spreng. Syst. ii. 308 = Chaetostoma dismoides.
aspera, *Voigt, in Syll. Ratisb.* ii. (1828) 55 (sp. dub.). —Hab.?
aspera, Willd. Sp. Pl. ii. 304 = Tibouchina aspera.
aurea, Desv. ex Ham. Prod. Pl. Ind. Occ. 35 = lutea.
avicularis, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia cinerea.
axillaris, Pav. ex DC. l. c. 101 = Meriana prunifolia.
baccharoides, Schrank & Mart. ex DC. l. c. 119 = Microlicia baccharoides.
balsamifera, Mart. & Schrank, l. c. 110 = Microlicia balsamifera.
berberifolia, Humb. & Bonpl. Rhex. 110. t. 42 = Comolia berberifolia.
bicolor, Andr. ex D. Don, in Mem. Wern. Soc. iv. (1822) 301 = Aciotis discolor.
bicornis, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 278 = Acisanthera recurva.
bicuspis, Schrank, ex DC. Prod. iii. 123 (= *Pachyloma coriaceum*).—Bras.
bidentata, Nees & Mart. in Nov. Act. Nat. Cur. xii. I. (1824) 53 = Cambessedesia Hilariana.
bilocularis, Schrank & Mart. ex DC. Prod. iii. 114 = Piptanthera paludosa.
bivalvis, Vahl, Eclog. Am. i. 38 = Acisanthera brevifolia.
blepharocentra, Mart. ex DC. Prod. iii. 103 = Lavoisiera imbricata.
Bonplandii, Humb. & Bonpl. Rhex. 159 (Index) = Monochaetum Bonplandii.
bracteolaris, Schrank & Mart. in DC. Prod. iii. 141 = Marcetia bracteolaris.
brasiliana, *Vand. Fl. Lusit.* 23.—Bras.
brasiliensis, Hoffmgg. Verz. Pfl. 75 = Luehea speciosa.
brevifolia, Rich. ex DC. Prod. iii. 117 = Acisanthera trivalvis.

RHEXIA :—
Bucardia, Schrank, ex DC. Prod. iii. 110 = Cambessedesia latevenosa.
calcarata, Schlecht. in Linnaea, xiii. (1839) 431 = Monochaetum Candolleanum.
campanularis, Humb. & Bonpl. Rhex. 35. t. 14 = Brachyotum campanulare.
Candolleana, Mart. ex DC. Prod. iii. 129 = Tibouchina Candolleana.
canescens, Humb. & Bonpl. Rhex. 14. t. 6 = Brachyotum canescens.
canescens, Humb. & Bonpl. l. c. 47. t. 18 = Monochaetum Bonplandii.
capitata, Rich. in Humb. & Bonpl. l. c. 84. t. 32 = Pterolepis glomerata.
cardinalis, Humb. & Bonpl. l. c. 96. t. 37 = Tibouchina cardinalis.
cataphracta, Schrank & Mart. ex DC. Prod. iii. 102 = Lavoisiera imbricata.
caudata, Schrank & Mart. l. c. 186 = Miconia urophylla.
cerastiifolia, Schrank & Mart. l. c. 137 = Tibouchina cerastifolia.
cernua, Humb. & Bonpl. Rhex. 32. t. 13 = Brachyotum cernuum.
Chamaecistus, Sieber, ex DC. Prod. iii. 134 (= *Pleroma Chamaecistus*).—Bras.
chamaedrifolia, Schrank & Mart. ex DC. l. c. 111 = Cambessedesia Espora.
Christophoriana, Schrank & Mart. l. c. 107 = Rhynchanthera dichotoma.
ciliata, Mart. l. c. 103 = Lavoisiera imbricata.
ciliosa, *Michx. Fl. Bor. Am.* i. 221.—Am. bor.
cinerea, Mart. ex DC. Prod. iii. 124 = Marcetia cinerea.
circaeifolia, Humb. & Bonpl. Rhex. 101. t. 39 = Aciotis circaeifolia.
Cochobambae, Schrank & Mart. ex DC. iii. 120 = Microlicia oligantha.
cognata, *Weinm. in Syll. Ratisb.* ii. (1828) 20.—Hab.?
collina, Salzm. ex Triana, in Trans. Linn. Soc. xxviii. (1873) 43 = Tibouchina Lhotzkyana.
compta, Mart. & Schrank, ex DC. Prod. iii. 103 = Lavoisiera imbricata.
condylocarpa, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 124 = Pterogastra cupheoides.
conferta, Humb. & Bonpl. Rhex. 53. t. 20 = Brachyotum confertum.
cordata, Spreng. Neue Entdeck. i. 301 = Microlicia cordata.
cordigera, Mart. & Schrank, ex DC. Prod. iii. 124 = Marcetia cordigera.
corymbosa, Mart. & Schrank, l. c. 110 = Cambessedesia corymbosa.
corymbosa, Raddi, in Mem. Soc. Ital. Sc. Moden. xx. (1829) 124. t. 2. f. 1 = Tibouchina corymbosa.
crassifolia, Mart. & Schrank, ex DC. Prod. iii. 104 = Lavoisiera crassifolia.
crenulata, Mart. & Schrank, l. c. 111 = Microlicia crenulata.
cubensis, Griseb. Cat. Pl. Cub. 104 = mariana.
dasystaminea, Schrank, ex DC. Prod. iii. 130 = Tibouchina granulosa.
decipiens, Salzm. ex Triana, in Trans. Linn. Soc. xxviii (1873) 44 = Tibouchina setulosa.
decumbens, Poir. Encyc. Suppl. iv. 628 = Dissotis decumbens.
decussata, Mart. & Schrank, ex DC. Prod. iii. 124 = Marcetia decussata.
Deppeana, Schlecht. & Cham. in Linnaea, v. (1830) 566 (= Arthrostema Schiedeanum).
dichotoma, Sieber, ex Griseb. Fl. Brit. W. Ind. 267 = Pterolepis glomerata.
dichrananthera, Ruiz & Pav. Fl. Per. iii. 84 = Monochaetum dicranantherum.
diffusa, Spreng. Syst. ii. 307 = Ossaea microphylla, *Triana*.
divaricata, Humb. & Bonpl. Rhex. 59. t. 22 = Pterogastra divaricata.
diversifolia, Humb. & Bonpl. l. c. 119. t. 45 = Arthrostemma ciliatum.
Drummondii, Hort. Angl. ex Heynh. Nom. i. 683 (Quid?).—Am. bor.
echinata, Ruiz & Pav. Fl. Per. iii. 85 (= *Pleroma echinatum*).—Peruv.
elachistophylla, Schrank & Mart. in DC. Prod. iii. 120 = Microlicia variolosa.
elegans, *Bonpl. ex Spach, Hist. Vég. Phan. Atl.* 16. t. 30.—Hab.?
elegans, *Schlecht. in Linnaea*, xiii. (1839) 432.—Mexic.

RHEXIA :—
elliptica, Raddi, in Mem. Soc. Ital. Sc. Moden. xx. (1829) 116. t. 1. f. 1 = Tibouchina Fothergillae.
enervia, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia euphorbioides.
ericoides, Spreng. Neue Entdeck. i. 302 = Marcetia carinata.
Espora, A. St. Hil. ex Humb. & Bonpl. Rhex. 152. t. 58 = Cambessedesia Espora.
estrellensis, Raddi, in Mem. Soc. Ital. Moden. xviii. Fis. (1820) 388 = Tibouchina estrellensis.
euphorbioides, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia euphorbioides.
excelsa, Humb. & Bonpl. Rhex. 90. t. 34 = Centronia excelsa.
excoriata, Mart. & Schrank, ex DC. Prod. iii. 124 = Marcetia excoriata.
Exul, Schrank & Mart. ex DC. l. c. 119 = Microlicia graveolens, variolosa.
fasciculata, Schrank & Mart. l. c. = Microlicia fasciculata.
fasciculata, Spreng. Neue Entdeck. iii. 62 = Cambessedesia Hilariana.
Fasikan, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 25 = Bredia hirsuta.
firmula, Mart. & Schrank, ex DC. Prod. iii. 104 = Lavoisiera firmula.
fissinervia, Mart. & Schrank, l. c. = Tibouchina fissinervia.
flexuosa, Ruiz & Pav. Fl. Per. iii. 85. t. 320 = Tibouchina longifolia.
floribunda, Mart. ex DC. Prod. iii. 103 = Lavoisiera imbricata.
floribunda, Schlecht. in Linnaea, xiii. (1839) 431 = Monochaetum rivulare.
Fontanesii, Humb. & Bonpl. Rhex. 93. t. 36 = Tibouchina granulosa.
formosissima, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 121 = Tibouchina granulosa.
Fothergillae, Schrank & Mart. ex DC. Prod. iii. 108 = Tibouchina Fothergillae.
fragilis, *Bertol. Fl. Guatimal.* 16.—Guatemala.
frigidula, Schrank & Mart. ex DC. Prod. iii. 127 = Tibouchina frigidula.
fulva, Spreng. Neue Entdeck. i. 301; Syst. ii. 308 = Microlicia serpyllifolia.
galioides, Schrank & Mart. ex DC. Prod. iii. 111 = Cambessedesia Hilariana.
gentianoides, Mart. & Schrank, l. c. 104 = Lavoisiera gentianoides.
glabella, *Michx. Fl. Bor. Am.* i. 222.—Am. bor.
glandulosa, *Bertol. Fl. Guatimal.* 15.—Guatemala.
glandulosa, Bonpl. Desc. Pl. Malm. 70. t. 27 (= *Pleroma glandulosa*).—Guian.
glomerata, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 276 = Pterolepis glomerata.
glutinosa, Linn. f. Suppl. 216 = Bucquetia glutinosa.
gracilis, Humb. & Bonpl. Rhex. 138. t. 52 = Tibouchina gracilis.
grandiflora, Vahl, Symb. Bot. iii. [106] Corrig.; Humb. & Bonpl. l. c. 25. t. 11 = Rhynchanthera grandiflora.
grandifolia, *Schlecht. in Linnaea*, xiii. (1839) 429.—Mexic.
graveolens, Mart. & Schrank, ex DC. Prod. iii. 119 = Microlicia graveolens.
helvola, Spreng. Syst. ii. 309 = Microlicia hypericoides.
herbacea, Raddi, in Mem. Soc. Ital. Moden. xviii. Fis. (1820) 389 = Aciotis paludosa.
herbacea, Schrank & Mart. ex DC. Prod. iii. 137 = Tibouchina herbacea.
heteranthera, Humb. & Bonpl. Rhex. 87. t. 33 = Melastoma repens.
heterophylla, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 316 = Chaetolepis heterophylla.
heterostemon, Mart. & Schrank, ex DC. Prod. iii. 126 = Trembleya parviflora.
hexandra, Schrank, l. c. 115 = Aciotis paludosa.
hieracioides, Mart. & Schrank, l. c. 133 = Tibouchina hieracioides.
Hilariana, Humb. & Bonpl. Rhex. 147. t. 56 = Cambessedesia Hilariana.
hirsuta, Mart. & Schrank, ex DC. Prod. iii. 119 = Microlicia graveolens, variolosa.
hirsuta, Salisb. Prod. 365 = virginica.
hispida, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Pterolepis pumila.
holosericea, Humb. & Bonpl. Rhex. 29. t. 12 = Tibouchina holosericea.

RHEXIA :—
hospita, Schrank & Mart. ex DC. Prod. iii. 129 = Tibouchina hospita.
hygrophila, Mart. ex DC. l. c. 128 = Tibouchina hygrophila.
hypericoides, Willd. Sp. Pl. ii. 303 = Desmoscelis villosa.
ilicifolia, Schrank & Mart. ex DC. Prod. iii. 111 = Cambessedesia ilicifolia.
imbricata, Thunb. Pl. Bras. Dec. i. 10. t. 2. f. 2 = Lavoisiera imbricata.
inaequalifolia, [Moç. & Sesse,] ex DC. Prod. iii. 122 = Arthrostema ciliatum.
inaequilateralis, Schlecht. & Cham. in Linnaea, v. (1830) 567 = Centradenia inaequilateralis.
incana, Spreng. Syst. ii. 309 = Monochaetum Bonplandii.
inconstans, Vahl, Eclog. Am. i. 37 = Tibouchina ornata.
indecora, Humb. & Bonpl. Rhex. 131. t. 49 = Aciotis indecora.
inundata, Schrank & Mart. ex DC. Prod. iii. 117 = Acisanthera inundata.
isophylla, Schrank & Mart. l. c. 120 = Microlicia isophylla.
itambana, Mart. & Schrank, l. c. 104 = Lavoisiera itambana.
jucunda, Schrank & Mart. l. c. 139 = Miconia staminea.
jungermannioides, Mart. & Schrank, l. c. 119 = Microlicia jungermannioides.
juniperina, Humb. & Bonpl. ex Spreng. Syst. ii. 310 = Marcetia juniperina.
jussiaeoides, *Linn. f. Suppl.* 215.—Guian.
ladanoides, Rich. in Humb. & Bonpl. Rhex. 72. t. 27 = Pterolepis pumila.
lanceaeflora, Mart. & Schrank, ex DC. l. c. 118 = Comolia lanceaeflora.
lanceolata, Humb. & Bonpl. Rhex. 56. t. 21 = Tibouchina longifolia.
lanceolata, *Walt. Fl. Carol.* 129.—Am. bor.
Langsdorffiana, Humb. & Bonpl. Rhex. 135. t. 51 = Tibouchina Langsdorffiana.
Langsdorffiana, Kunth, ex DC. Prod. iii. 124 = Marcetia taxifolia.
lasiandra, Schrank & Mart. ex DC. l. c. 127 = Tibouchina Martinsiana.
latevenosa, Mart. l. c. 110 = Cambessedesia latevenosa.
latifolia, Aubl. Pl. Gui. i. 336. t. 129. f. 2 = Comolia Aubletii.
lepidota, Humb. & Bonpl. Rhex. 38. t. 15 = Tibouchina lepidota.
leptophylla, Humb. & Bonpl. l. c. 64. t. 24 = Comolia leptophylla.
leucantha, Sw. Prod. Veg. Ind. Occ. 61 = Meriania leucantha.
Leuzeana, Humb. & Bonpl. Rhex. 144. tt. 54, 55 = Bertolonia Leuzeana.
limnobios, Schrank & Mart. ex DC. Prod. iii. 117 = Acisanthera limnobios.
limosa, Mart. ex DC. l. c. 108 = Rhynchanthera limosa.
linearifolia, Poir. Encyc. vi. 2 = lutea.
linearifolia, Poir. ex DC. Prod. iii. 122 = Ludwigia alternifolia.
linifolia, Mart. ex DC. l. c. 105 = Microlicia linifolia.
longifolia, Vahl, Eclog. Am. i. 39 = Tibouchina longifolia.
ludoviciana, Rafin. Fl. Ludov. 92 = mariana.
lutea, *Walt. Fl. Carol.* 130.—Am. bor.
lutescens, Ruiz & Pav. Fl. Per. iii. 84 = Brachyotum lutescens.
Lychnitis, Schrank & Mart. ex DC. Prod. iii. 126 = Trembleya Lychnitis.
lychnitoides, Schrank & Mart. l. c. 132 = Desmoscelis villosa.
macrochiton, Mart. l. c. 129 = Tibouchina macrochiton.
mariana, *Linn. Sp. Pl.* 346.—Am. bor.
Martiusiana, Ser. in DC. Prod. iii. 137 = Tibouchina sebastianopolitana.
microphylla, Humb. & Bonpl. Rhex. 4. t. 2 = Chaetolepis microphylla.
mollis, Humb. & Bonpl. l. c. 50. t. 19 = Tibouchina mollis.
Moricandiana, Mart. ex DC. Prod. iii. 128 = Tibouchina Moricandiana.
mucorifera, Mart. & Schrank, ex DC. l. c. 103 = Lavoisiera mucorifera.

RHEXIA:—
multiflora, Humb. & Bonpl. Rhex. 41. t. 16 = Monochaetum multiflorum.
muricata, Humb. & Bonpl. l. c. 1. t. 1 = Tibouchina grossa.
myrtoidea, Humb. & Bonpl. l. c. 7. t. 3 = Monochaetum myrtoideum.
napaulensis, Ser. ex DC. Prod. iii. 142 = Osbeckia chinensis.
Nianga, Mart. & Schrank, ex DC. l. c. 163 = Oxymeris Nianga.
nummularioides, Humb. & Bonpl. Rhex. 62. t. 23 = Comolia nummularioides.
nymphaeifolia, Humb. & Bonpl. l. c. t. 53 = Bertolonia nymphaeaefolia.
obtusa, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
obvallata, Sieber, ex Oken, Isis (1822) 453; Miq. Comm. Phyt. 79 = Pterolepis glomerata.
ochypetala, Ruiz & Pav. Fl. Per. iii. 86. t. 321. f. a = Tibouchina ochypetala.
oleaefolia, Schrank & Mart. ex DC. Prod. iii. 139 = Microlepis oleaefolia.
ornata, Rich. in Humb. & Bonpl. Rhex. 69. t. 26 = Tibouchina ornata.
oxypetala, Spreng. Syst. ii. 306 = Tibouchina oxypetala.
Pachyloma, Mart. ex DC. Prod. iii. 123 (= *Pachyloma coriaceum*).—Bras.
paludosa, Salzm. ex Griseb. Fl. Brit. W. Ind. 269 = Acisanthera brevifolia.
paniculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 222 = Oxyspora paniculata.
paniculata, Mart. & Schrank, ex DC. Prod. iii. 137 = Tibouchina sebastianopolitana.
parnassifolia, Mart. & Schrank, l. c. 140 = Pterolepis parnassiaefolia.
parvifolia, Schlecht. in Linnaea, xiii. (1839) 428 = Centradenia inaequilateralis.
pendulifolia, Bonpl. Desc. Pl. Malm. 68. t. 26 = Aciotis pendulifolia.
pentanthera, Ser. ex DC. Prod. iii. 108 = Rhynchanthera Schrankiana.
petiolata, Walt. Fl. Carol. 130 = ciliosa?
phlogiformis, Mart. & Schrank, ex DC. Prod. iii. 126 = Trembleya phlogiformis.
piloselloides, Humb. & Bonpl. Rhex. 12. t. 5 = Castratella piloselloides.
pityrophylla, Mart. ex DC. Prod. iii. 139 = Pyramia pityrophylla.
polygonoides, Mart. l. c. 137 = Pterolepis polygonoides.
polystachya, Humb. & Bonpl. Rhex. 98. t. 38 = Aciotis polystachya.
pratensis, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 276 = Pterolepis glomerata.
Princeps, Humb. & Bonpl. Rhex. 122. t. 46 = Dissotis eximia.
proteaeformis, Schrank & Mart. ex DC. Prod. iii. 130 = Tibouchina holosericea.
psammophila, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 37 = Comolia denudata.
pulcherrima, Mart. & Schrank, ex DC. Prod. iii. 104 = Lavoisiera pulcherrima.
pumila, Humb. & Bonpl. Rhex. t. 35 = Pterolepis pumila.
punctata, Mart. & Schrank, ex DC. Prod. iii. 104 = Lavoisiera punctata.
punctatissima, Mart. & Schrank, l. c. 118 = Acisanthera punctatissima.
pungens, Mart. & Schrank, l. c. 112 = Chaetostoma pungens.
purpurata, Schrank & Mart. l. c. 110 = Cambessedesia purpurata.
purpurea, Sw. Prod. Veg. Ind. Occ. 64 = Meriania purpurea.
Purshii, Spreng. Syst. v. 590 (Index) = virginica.
quadrangularis, Salzm. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 42 = Tibouchina urceolaris.
quadrivalvis, Mart. ex DC. Prod. iii. 124 = Marcetia taxifolia.
quaternifolia, Schrank & Mart. ex DC. l. c. 140 = Microlepis quaternifolia.
quinquenervis, Ruiz & Pav. Fl. Per. iii. 83 = Brachyotum quinquenerve.
Raddiana, Moric. ex DC. Prod. iii. 129 = Tibouchina Raddiana.
Radula, Humb. & Bonpl. Rhex. 107. t. 41 = Macairea Radula.
recurva, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Acisanthera recurva.

RHEXIA:—
regalis, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 277 = Pterolepis pumila.
repanda, Mart. & Schrank, ex DC. Prod. iii. 141 = Pterolepis repanda.
reptans, Mart. ex Steud. Nom. ed. II. ii. 447 = Tibouchina clinopodifolia.
resinosa, Schrank, ex DC. Prod. iii. 111 = Microlicia sincorensis.
reticulata, Humb. & Bonpl. Rhex. 21. t. 9 = Tibouchina reticulata.
rosea, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen (Quid?).—Am. bor.
rosmarinifolia, Ruiz & Pav. Fl. Per. iii. 84 = Brachyotum rosmarinifolium.
rosmarinoides, Schrank & Mart. ex DC. Prod. iii. 126 = Trembleya rosmarinoides.
rostrata, Schrank & Mart. l. c. 108 = Rhynchanthera rostrata.
rotundifolia, Humb. & Bonpl. Rhex. 66. t. 25 = Graffenrieda rotundifolia.
rufipilis, Schlecht. in Linnaea, xiii. (1839) 430 (= *Pleroma rufipile*).—Mexic.
sarmentosa, Humb. & Bonpl. Rhex. 23. t. 10 (= *Pleroma laxum*).—Peruv.
scabriuscula, Schlecht. in Linnaea, xiii. (1839) 431 (= *Pleroma scabriusculum*).—Mexic.
scandens, Rich. ex DC. Prod. iii. 127 = Adelobotrys scandens.
Schiedeana, Schlecht. & Cham. in Linnaea, v. (1830) 565 (= *Pleroma rufipile*).—Mexic.
Schrankiana, Mart. ex DC. Prod. iii. 107 = Rhynchanthera Schrankiana.
scoparia, Mart. ex DC. Prod. iii. 112 = Marcetia carinata.
scoparia, A. St. Hil. ex Humb. & Bonpl. Rhex. 154. t. 59 = Microlicia scoparia.
sebastianopolitana, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 126. t. 2 = Tibouchina sebastianopolitana.
semiadnata, Schrank, ex DC. Prod. iii. 140 = Pterolepis alpestris.
semidecandra, Schrank & Mart. ex DC. l. c. 129 = Tibouchina semidecandra.
semiserrata, Schrank & Mart. l. c. 167 = Huberia semiserrata.
septemnervia, Walt. Fl. Carol. 130 = virginica.
Seringeana, Mart. ex DC. Prod. iii. 119 = Microlicia graveolens.
serrulata, *Nutt. Gen. Am.* i. 243.—Am. bor.
serrulata, Rich. ex Humb. & Bonpl. Rhex. 74. t. 28 = Rhynchanthera serrulata.
Sertularia, Schrank & Mart. ex DC. Prod. iii. 125 = Comolia Sertularia.
setosa, Schrank & Mart. l. c. 119 = Microlicia subsetosa.
setosa, Spreng. Neue Entdeck. i. 301 = Microlicia setosa.
sherardioides, Schrank, ex DC. Prod. iii. 133 = Pterolepis glomerata.
sileniflora, Humb. & Bonpl. Rhex. 129. t. 48 = Aciotis sileniflora.
sincorensis, Mart. ex DC. Prod. iii. 111 = Microlicia sincorensis.
sipaneoides, Mart. l. c. 141 = Pterolepis sipaneoides.
speciosa, Bert. ex DC. Prod. iii. 101 = Meriana leucantha.
speciosa, Humb. & Bonpl. Rhex. 9. t. 4 = Meriania speciosa.
spicata, Schrank & Mart. ex DC. Prod. iii. 15 = Ossaea brachystachya.
stachyoides, Humb. & Bonpl. Rhex. 113. t. 43 = Desmoscelis villosa.
stachyoides, Schrank & Mart. ex DC. iii. 108 = Rhynchanthera rostrata.
stenocarpa, Schrank & Mart. l. c. 130 = Tibouchina stenocarpa.
stricta, Humb. & Bonpl. Rhex. 19. t. 8 = Brachyotum strigosum.
stricta, *Pursh, Fl. Bor. Am.* i. 258.—Am. bor.
strigillosa, Schrank & Mart. ex DC. Prod. iii. 133 = Tibouchina gracilis.
strigosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Tibouchina ornata.
strigosa, Willd. ex Spreng. Syst. ii. 309 = Pterolepis pumila.
striphnocalyx, Mart. ex DC. Prod. iii. 140 = Pterolepis striphnocalyx.
suberosa, Spreng. Neue Entdeck. i. 302 = Cambessedesia Hilariana.

RHEXIA:—
subtriplinervia, Hort. Berol. ex Heynh. Nom. i. 683 (Quid?).—Hab.?
superba, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 118. t. 1. f. 4 = Tibouchina Raddiana.
tamariscina, Schrank & Mart. ex DC. Prod. iii. 125 = Marcetia tamariscina.
taxifolia, A. St. Hil. in Humb. & Bonpl. Rhex. 150. t. 57 = Marcetia taxifolia.
tenella, Humb. & Bonpl. l. c. 79. t. 30 = Ernestia tenella.
tenuifolia, Mart. ex DC. Prod. iii. 125 = Marcetia tenuifolia.
tetragona, Mart. & Schrank, ex DC. l. c. 103 = Lavoisiera tetragona.
tetraptera, Desv. ex Ham. Prod. Pl. Ind. Occ. 35 = virginica.
tetrasticha, Mart. & Schrank, ex DC. Prod. iii. 112 = Marcetia tetrasticha.
thymifolia, Herb. Monac. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 26 = Microlicia serpyllifolia.
thymifolia, Humb. & Bonpl. Rhex. 133. t. 50 = Appendicularia thymifolia.
thymifolia, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia variolosa, serpyllifolia.
tibouchinoides, Schrank & Mart. l. c. 129 = Tibouchina trichopoda.
tortuosa, Humb. & Bonpl. Rhex. 17. t. 7 (= *Pleroma mexicanum*).—Mexic.
trichocalycina, Mart. & Schrank, ex DC. Prod. iii. 120 = Microlicia trichocalycina.
trichotoma, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 277 = Pterolepis pumila.
trichotoma, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = Pterolepis glomerata.
triflora, Mart. & Schrank, ex DC. Prod. iii. 126 = Trembleya parviflora.
triflora, Raddi, in Mem. Soc. Ital. Moden. xx. (1829) 122. t. 1. f. 2 = Tibouchina Moricandiana.
trifolia, Schrank & Mart. ex DC. Prod. iii. 127 = Tibouchina frigidula.
trivalvis, Vahl, Eclog. Am. i. 39 = Acisanthera trivalvis.
uda, Mart. ex DC. Prod. iii. 136 = Pterolepis pumila.
uncata, Schrank & Mart. ex DC. l. c. 120 = Microlicia uncata.
uniflora, Raddi, in Mem. Soc. Ital. Moden. xviii. Fis. (1820) 388 = Tibouchina Raddiana.
uniflora, Vahl, Symb. Bot. ii. 48 = Acisanthera recurva.
urceolaris, Schrank & Mart. ex DC. Prod. iii. 139 = Tibouchina urceolaris.
variabilis, Mart. & Schrank, l. c. 118 = Acisanthera variabilis.
variolosa, Schrank & Mart. ex DC. Prod. iii. 120 = Microlicia variolosa. cordata.
venusta, Schrank, & Mart. l. c. 119 = Microlicia serpyllifolia.
veronicaefolia, Schrank & Mart. l. c. 121 = Siphanthera cordata.
versicolor, Lindl. Bot. Reg. t. 1066 = Tibouchina versicolor.
vestita, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia vestita.
villosa, Aubl. Pl. Gui. i. 334. t. 129. f. 1 = Comolia villosa.
villosissima, Rich. in Humb. & Bonpl. Rhex. 81. t. 31 = Desmoscelis villosa.
viminalis, Mart. & Schrank, ex DC. Prod. iii. 103 = Microlicia viminalis.
viminea, D. Don, in Bot. Reg. t. 664 = Tibouchina viminea.
violacea, Mart. ex DC. Prod. iii. 118 = Comolia violacea.
virginica, *Linn. Sp. Pl.* 346.—Am. bor.
virgulata, Schrank & Mart. ex DC. Prod. iii. 119 = Microlicia serpyllifolia.
viscidula, Mart. ex Cham. in Linnaea, ix. (1834) 385 = Comolia sessilis.
viscidula, Schrank & Mart. ex DC. Prod. iii. 118 = Comolia violacea.
zeylanica, Link, Enum. Hort. Berol. i. 380 = Tibouchina holosericea.

RHIGIOCARYA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 100 = **Chasmanthera**, Hochst. (Menisp.).
racemifera, Miers, l. c. 101.—Afr. trop.

RHIGIOPHYLLUM, Hochst. in Flora, xxv. (1842) 232. *CAMPANULACEAE*, Benth. & Hook. f. ii. 559.
squarrosum, *Hochst. l. c.*—Afr. austr.

RHIGIOTHAMNUS, Spach, Hist. Vég. Phan. x. 9 (1841) = **Dicoma**, Cass. (Compos.).

RHIGOSPIRA, Miers, Apocyn. S. Am. 67 (1878) = **Tabernaemontana**, Linn. (Apocyn.).
paucifolia, Miers, l. c. 69 = T. paucifolia.
quadrangularis, Miers, l. c. 68 = Ambellania quadrangularis.
reticulata, Miers, l. c. 69 = T. reticulata.
sinuosa, Miers, l. c. 70 = T. Sprucei.
Sprucei, Miers, l. c. = T. Sprucei.
ternstroemiacea, Miers, l. c. 71 = T. ternstroemiacea.
venulosa, Miers, l. c. 68 = Ambellania macrophylla.

RHIGOZUM, Burch. Trav. i. 299 (1822). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1048.
RHIZOGUM, Reichb. Consp. 125 (1828).
brachiatum, E. Mey. in Drège, Zwei Pfl. Docum. 264 = trichotomum.
obovatum, *Burch. Trav.* i. 389.—Afr. austr.
trichotomum, *Burch. l. c.* 299.—Afr. austr.

RHINACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 76 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1112.
calcaratus, *Nees, in DC. Prod.* xi. 444.—Ind. or.
communis, *Nees, in Wall. Pl. As. Rar.* iii. 109.—Ind. or.; Malaya.
gracilis, Boj. ex Nees, in DC. Prod. xi. 506 = Hypoëstes gracilis.
gracilis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 218.—Afr. trop.
macilentus, Presl, Bot. Bemerk. 95 = communis.
Nasuta, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 79 = communis.
oblongus, *Nees, in DC. Prod.* xi. 444.—Afr. trop.
osmospermus, T. Anders. in Journ. Linn. Soc. vii. (1863) 51; ix. (1868) 522 = communis.
osmospermus, *Boj. ex Nees, in DC. Prod.* xi. 443.—Madag.
Rottlerianus, Nees, in Wall. Pl. As. Rar. iii. 109 = communis.
scoparius, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 88.—Ins. Socotra.
tubulosus, Presl, Bot. Bemerk. 95 = Justicia tubulosa.

RHINACTINA, Less. in Linnaea, vi. (1831) 119 = **Aster**, Tourn. (Compos.).
limonifolia, Less. l. c. = A. obovatus.
uniflora, Bunge, ex DC. Prod. v. 279 = A. eremophilus.

RHINACTINA, Willd. in Ges. Nat. Fr. Berl. Mag. i. (1807) 139 = **Jungia**, Linn. f. (Compos.).
cinerarioides, Willd. ex DC. Prod. vii. 55 = Jungia ferruginea.
purpurea, Willd. ex Steud. Nom. ed. I. 687 = Leuceria purpurea.

RHINANTHERA, Blume, Bijdr. 1121 (1826) = **Scolopia**, Schreb. (Bixin.).
Blumei, Steud. Nom. ed. II. ii. 448.—Java.
odoratissima, Blume, Bijdr. 1121.—Java.

RHINANTHUS, Linn. Sp. Pl. ed. I. 603 (1753), partim; Bieb. Fl. Taur. Cauc. ii. 68 (1808) = **Rhynchocorys**, Griseb. (Scrophul.).
Columnae, Jaub. & Spach, Illustr. iv. 138 = Rhynch. Elephas.
Elephas, Linn. Sp. Pl. 603 = Rhynch. Elephas.
orientalis, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 12 = Rhynch. Elephas.
orientalis, Linn. Sp. Pl. 603 = Rhynch. orientalis.
strictus, C. Koch, in Linnaea, xxii. (1849) 684 = Rhynch. Elephas.

RHINANTHUS, Linn. Syst. ed. I (1735); Gen. ed. I. 184 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 979.
ALECTOROLOPHUS, Hall. Enum. Stirp. Helv. ii. 623 (1742); Moench, Meth. 440 (1794).
CRISTA-GALLI, Rupp. Fl. Jen. ed. Hall. 240 (1745).
ELEPHAS, Tourn. ex Adans. Fam. ii. 211 (1763).
PROBOSCIPHORA, Neck. Elem. i. 336 (1790).

RHINANTHUS:—
acuminatus, Poir. Encyc. Suppl. v. 624 = Castilleja pallida.
adulterinus, Wallr. Beitr. Fl. Hercyn. 226 (Quid?).—Europ.
Alectorolophus, Pollich, Hist. Pl. Palat. ii. 177 = major.
alpestris, Steud. Nom. ed. II. ii. 448 = major.
alpinus, Baumg. Enum. Stirp. Transs. ii. 194 = major.
alpinus, Koch, Syn. Fl. Germ. ed. I. 627 = angustifolius.
alpinus, Lam. Fl. Fr. ii. 354 = Bartsia alpina.
angustifolius, *C. C. Gmel. Fl. Bad.* ii. 669.—Europ.
aristatus, Čelak. in Oestr. Bot. Zeitschr. xx. (1870) 132 = angustifolius.
bicolor, Poir. Encyc. Suppl. v. 623 = Bartsia Trixago.
bifidus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 94 = Pedicularis carnosa.
buccalis, Wallr. in Flora, xxv. (1842) 504 = major.
capensis, Linn. Syst. ed. XII. 405; Thunb. Prod. Pl. Cap. 98 = Bartsia Trixago.
coccineus, Lam. Encyc. ii. 60 = Castilleja coccinea.
creticus, F. G. Dietr. Vollst. Gaertn. Lexik. viii. 137 = Bartsia Trixago.
Crista-galli, Linn. Sp. Pl. 603 = major, minor.
Crista-galli, Vill. Fl. Delph. 62 = major.
cristatus, Stokes, Bot. Mat. Med. iii. 378 = major.
glaber, Lam. Fl. Fr. ii. 352 = minor.
glacialis, Personn. in Bull. Soc. Bot. Fr. x. (1863) 745 = major?
glauca, Poir. Encyc. Suppl. ii. 309 = Lagotis glauca.
grandiflorus, Bluff, Fingerh. Comp. Fl. Germ. ed. I. ii. 61 = major?
hirsutus, Lam. Fl. Fr. ii. 353 = major.
indicus, Linn. Sp. Pl. 603 = Geniosporum elongatum.
inflatus, Salisb. Prod. 96 = major, minor.
major, *Ehrh. Beitr.* vi. 144.—Europ.
maritimus, Lam. Fl. Fr. ii. 353 = Bartsia Trixago.
maximus, Lam. Encyc. ii. 61 = Bartsia viscosa.
maximus, Willd. Sp. Pl. iii. 189 = Bartsia Trixago.
minor, *Ehrh. Beitr.* vi. 144.—Europ.
montanus, Sauter, in Flora, xl. (1857) 361 = Bartsia alpina.
nigricans, Meinsh. Fl. Ingr. 259 = minor?
obscurus, Steph. Nom. Pl. Agric. Mosq. (1804) 35 (Quid?).—Rossia.
pallens, Wibel, Prim. Fl. Werth. 142 = seq.?
pallidus, Lam. Encyc. ii. 60 = Castilleja pallida.
parviflorus, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 61 = minor.
pratensis, Vill. Fl. Delph. 62 = minor.
pubescens, Boiss. & Heldr. ex Boiss. Fl. Orient. iv. 479 = major.
pulcher, Schummel, ex Opiz, Natural. xi. (1826) 449 = Bartsia alpina.
ramosus, Schur, Enum. Pl. Transs. 511 = minor.
Reichenbachii, Benth. in DC. Prod. x. 558 = major.
riphaeus, Krock. Fl. Siles. Suppl. ii. 160 = major, minor.
scaber, Thunb. Prod. Pl. Cap. 98 = Alectra melampyroides.
spicatus, Poir. Encyc. Suppl. v. 623 = Bartsia aspera.
stenophyllus, Schur, Enum. Pl. Transs. 512 = minor.
tenuifolius, Poir. Encyc. Suppl. v. 624 = Orthocarpus tenuifolius.
Tournefortii, *Jaub. & Spach, Illustr.* iv. 137. t. 393.—Oriens.
trifidus, Vahl, Symb. Bot. i. 44 = Bungea trifida.
Trixago, Linn. Syst. ed. XII. 1107; Gouan, Hort. Monsp. 296 = Bartsia Trixago.
Trixago, Thuill. Fl. Par. ed. II. 304 = minor.
versicolor, Lam. Encyc. ii. 61 = Bartsia Trixago.
villosus, Pers. Syn. ii. 151 = major.
virginicus, Linn. Herb. ex Benth. in DC. Prod. x. 558 = Lamourouxia serratifolia.
virginicus, Linn. Sp. Pl. 603 = Gerardia quercifolia.
viscosus, Lam. Fl. Fr. ii. 354; Brot. Fl. Lusit. i. 187 = Bartsia viscosa.
vulgaris, Gueldenst. ex Ledeb. Fl. Ross. iii. 266 = major, minor.

RHINANTUS, Gilib. Exercit. i. 132 (1792) = praec.

RHINCHOGLOSSUM, Blume, Bijdr. 741 (1826) = **Rhynchoglossum**, Blume (Gesnerac.).

RHINCHOSIA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 78 = **Rhynchosia**, Lour. (Legumin.).

RHINCOSPORA, Gaudich. in Freyc. Voy. Bot. 415 (1826) = **Rynchospora**, Vahl (Cyperac.).

RHINIACHNE, Hochst. ex Steud. Syn. Pl. Gram. 360, in syn. (1855) = **Thelepogon**, Roth (Gramin.).
Princeps, Hochst. ex Steud. l. c. = T. elegans.

RHINIUM, Schreb. Gen. 701 (1791) = **Doliocarpus**, Roland. (Dillen.).

RHINOCARPUS, Bert. ex H. B. & K. Nov. Gen. et Sp. viii. 5. t. 601 (1825) = **Anacardium**, Linn.
excelsa, Bert. ex H. B. & K. l. c. = A. Rhinocarpus.

RHINOGLOSSUM, Pritz. Ic. Ind. i. 933 (1855), sphalm. = **Rhynchoglossum**, Blume (Gesner.).

RHINOLOBIUM, Arn. in Mag. Zool. & Bot. ii. (1838) 420 = **Schizoglossum**, E. Mey. (Asclep.).
lineare, Decne. in DC. Prod. viii. 556, in syn. (= *Lagarinthus gracilis*).—Afr. austr.
tenue, Arn. in Mag. Zool. & Bot. ii. (1838) 421.—Afr. austr.

RHINOPETALUM, Fisch. ex Alex. in Edinb. N. Phil. Journ. [viii]. (Oct.–Dec. 1829) 19 = **Fritillaria**, Tourn. (Liliac.).
Boissieri, Klatt, in Hamb. Gartenz. xvi. (1860) 439 = F. gibbosa.
Carelini, Steud. Nom. ed. II. ii. 448, sphalm. = seq.
Karelini, Fisch. ex Sweet, Brit. Flow. Gard. Ser. II. t. 283 = F. gibbosa.
stenantherum, Regel, in Act. Hort. Petrop. iii. II. (1875) 294.—Turkestan.
triste, Eversm. ex Ledeb. Fl. Ross. iv 149 = F. gibbosa.

RHINOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 56 = **Thesium**, Linn. (Santal.).
brevifolia, Turcz. l. c. 57 = T. leptocaule.
longifolia, Turcz. l. c. = T. Turczaninowii.

RHINOSTIGMA, Miq. Fl. Ind. Bat. Suppl. 495 (1860) = **Garcinia**, Linn. (Guttif.).
macrophyllum, Miq. l. c. = G. macrophylla.
parvifolium, Miq. l. c. = G. dioica?

RHIPIDODENDRON, Spreng. Anleit. ii. I. 243 (1817) = seq.

RHIPIDODENDRUM, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 164 = **Aloë**, Tourn. (Liliac.).
dichotomum, Willd. l. c. 166 = A. dichotoma.
distichum, Willd. l. c. 165 = A. plicatilis.
plicatile, Haw. Rev. Pl. Succ. 45 = A. plicatilis.

RHIPIDOSTIGMA, Hassk. Retzia, 103 (1855) = **Maba**, Forst. (Ebenac.).
Teijsmanni, Hassk. l. c. 106 = M. Teijsmanni.
Zollingeri, Hassk. l. c. 104 = M. hermaphroditica.

RHIPOGONUM, Forst. Char. Gen. 49. t. 25 (1776) (Ripogonum). *LILIACEAE*, Benth. & Hook. f. iii. 764.
album, *R. Br. Prod.* 293.—Austral.
album, Sieber, ex Kunth, Enum. Pl. v. 161 = Smilax glycyphylla.
discolor, *F. Muell. Fragm.* vii. 78.—Austral.
dubium, *Endl. Prod. Fl. Norfolk.* 30.—Ins. Norfolk.
Elseyanum, *F. Muell. Fragm.* i. 44.—Austral.
Fawcettianum, *F. Muell. ex Benth. Fl. Austral.* vii. 9.—Austral.
Moorianum, F. Muell. Fragm. i. 44 = album.
parviflorum, R. Br. Prod. 293 = scandens.
scandens, *Forst. Char. Gen.* 50.—N. Zel.

RHIPSALIS, Gaertn. Fruct. i. 137. t. 28 (1788). *CACTEAE*, Benth. & Hook. f. i. 850.
CASSYTHA, Mill. Gard. Dict. ed. VIII. (1768).
HARIOTA, Adans. Fam. ii. 243 (1763).
LEPISMIUM, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 315, 380.
PFEIFFERA, Salm-Dyck, Cact. Hort. Dyck. ed. I. 40 (1845).
aethiopica, *Welw. in Journ. Linn. Soc.* iii. (1859) 152, nomen.—Afr. trop.
alata, Steud. Nom. ed. II. i. 333 = pachyptera.
alternata, *Lem. Cact.* 80, nomen.—Am. trop.
brachiata, Hook. Bot. Mag. t. 4039 = R. Saglionis.

RHIPSALIS:—
calamiformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 135 = funalis.
Cassytha, *Gaertn. Fruct.* i. 137. t. 28 (Cassutha).—Am. trop.
cassythoides, *G. Don, Gen. Syst.* iii. 176.—Mexic.
cereiformis, *Foerst. Handb. Cact.* ed I. 454.—Hab.?
cereuscula, Haw. ex Phil. Mag. (1830) 112 = R. Saglionis.
conferta, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 229.—Hab.?
coriacea, *Polak. in Linnaea*, xli. (1877) 562.—Costa Rica.
cribrata, Pfersd. ex N. E. Br. in Gard. Chron. (1877) i. 716 = penduliflora.
crispa, Hort. ex Foerst. Handb. Cact. ed. I. 450 = crispata.
crispata, *Pfeiff. Enum. Diagn. Cact.* 130.—Hab.?
cylindrica, *Steud. Nom.* ed. II. ii. 448.—Bras.
dichotoma, *G. Don, Gen. Syst.* iii. 176.—N. Granat.
echinata, Hort. ex Pfeiff. Enum. Diagn. Cact. 136 = mesembryanthoides.
fasciculata, *Haw. Suppl. Pl. Succ.* 83.—Ind. occ.
floccosa, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 134.—Hab.?
funalis, *Salm-Dyck, ex DC. Prod.* iii. 476.—Am. austr.
grandiflora, Haw. Suppl. Pl. Succ. 83 = funalis.
Hookeriana, G. Don, Gen. Syst. iii. 176 = R. Cassutha.
horrida, *Baker, in Journ. Linn. Soc.* xxi. (1884) 347.—Madag.
Houlletiana, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 64.—Bras.
Houlletii, Hook. f. Bot. Mag. t. 6089 = praec.
Knightii, *Foerst. Handb. Cact.* ed. I. 456.—Bras.
Lorentziana, *Griseb. in Goett. Abh.* xxiv. (1879) 139.—Reg. Argent.
lumbricoides, Lem. in Salm-Dyck, Hort. Dyck. ed. (1841) 1842; et Rev. Hortic. (1865) 219 = sarmentacea.
macrocarpa, *Miq. in Bull. Sc. Nat. Phys. Néerl.* (1838) 49.—Hab.?
mesembryanthemoides, Steud. Nom. ed. II. ii. 449 = seq.
mesembryanthoides, *Haw. Rev. Pl. Succ.* 71.—Am. austr.
micrantha, *DC. Prod.* iii. 476.—Ecuador.
Miquelii, *Lem. Cact.* 80, nomen.—Am. trop.
Mittleri, *Foerst. Handb. Cact.* ed. I. 455.—Bras.
monacantha, *Griseb. in Goett. Abh.* xxiv. (1879) 140.—Reg. Argent.
Myosurus, *Foerst. Handb. Cact.* ed. I. 455.—Bras.
pachyptera, *Pfeiff. Enum. Diagn. Cact.* 132.—Ind. occ.
paradoxa, *Salm-Dyck, Cact. Hort. Dyck.* ed. I. 39; ed. II. 59, 228.—Bras.
parasitica, DC. Prod. iii. 476 = undulata.
parasitica, Haw. Syn. Pl. Succ. 187 = fasciculata.
pendula, Hort. ex Pfeiff. Enum. Diagn. Cact. 133 = R. Cassytha.
penduliflora, *N. E. Br. in Gard. Chron.* (1877) i. 716.—Bras.?
pentagona, Hort. ex Foerst. Handb. Cact. ed. I. 453 = seq.
pentaptera, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 105.—Bras.
platycarpa, Lem. in Rev. Hortic. Sér. IV. ix. (1860) 502 = robusta.
platycarpa, *Pfeiff. Enum. Diagn. Cact.* 131.—Bras.
pterocaulis, Hort. ex Foerst. Handb. Cact. ed. I. 453 = paradoxa.
ramulosa, *Pfeiff. Enum. Diagn. Cact.* 130.—Hab?
rhombea, *Pfeiff. l. c.*—Hab?
Riedeliana, *Regel, Ind. Sem. Hort. Petrop.* (1860) 49.—Bras.
robusta, *Lem. in Rev. Hortic.* Sér. IV. ix. (1860) 502.—Bras.?
rugulosa, *Lem. Illust. Hortic.* viii. (1861) sub t. 293.—Ind. occ.
Saglionis, *Otto, ex Walp. Rep.* ii. 936.—Reg. Argent.
salicornioides, *Haw. Suppl. Pl. Succ.* 83.—Bras.
sarmentacea, *Otto & Dietr. Allg. Gartenz.* ix. (1841) 98.—Reg. Argent.
spatulata, *Otto, ex Sweet, Hort. Brit.* ed. III. 288.—Hab.?
Swartziana, *Pfeiff. Enum. Diagn. Cact.* 131.—Ind. occ.
teres, *Steud. Nom.* ed. II. ii. 449.—Bras.
trigona, *Pfeiff. Enum. Diagn. Cact.* 133.—Bras.
Turpinii, *Lem. Cact.* 80, nomen.—Am. trop.
undulata, *Pfeiff. Enum. Diagn. Cact.* 136.—Ind. occ.

RHIZAERIS, Rafin. Sylva Tellur. 90 (1838) = **Lagunclaria**, Gaertn. (Combret.).
alba, Rafin. l. c. = L. racemosa.

RHIZEMYS, Rafin. Fl. Tellur. iv. 26 (1836) = **Testudinaria**, Salisb. (Dioscor.).
elephantipes, Rafin. l. c. = T. elephantipes.
montana, Rafin. l. c.—Hab.?

RHIZIRIDEUM, Fourr. in Ann. Soc. Linn. Lyon, N.S. (1869) 160 = **Allium**, Tourn. (Liliac.).
acutangulum, Fourr. l. c. = A. senescens.
fallax, Fourr. l. c. = A. senescens.

RHIZIUM, Dulac, Fl. Hautes-Pyr. 243 (1867) = **Elatine**, Linn.
curvum, Dulac, l. c. = E. Hydropiper.
verticillatum, Dulac, l. c. = E. Alsinastrum.

RHIZOBOLUS, Gaertn. ex Schreb. Gen. 369 (1789); et Fruct. ii. 93. t. 98 (1791) = **Caryocar**, Linn. (Ternstr.).
butyrosus, J. F. Gmel. Syst. 840 = C. butyrosum.
glaber, Steud. Nom. ed. II. ii. 449 = C. glabrum.
Pekea, Gaertn. Fruct. ii. 93. t. 98. f. 1 = C. tomentosum.
Saouvari, Correa, in Ann. Mus. Par. viii. (1806) 394. t. 5. f. 2 = C. glabrum.
Souari, Steud. Nom. ed. I. 688, sphalm. = praec.
tuberculosum, J. F. Gmel. Syst. 840 = C. tomentosum.

RHIZOBOTRYA, Tausch, in Flora, xix. (1836) 33 = **Cochlearia**, Tourn. (Crucif.).
alpina, Tausch, l. c. 34 = C. brevicaulis.

RHIZOCEPHALUM, Wedd. Chlor. And. ii. 11. t. 46 (1857). *CAMPANULACEAE*, Benth. & Hook. f. ii. 549.
Candollei, *Wedd. l. c.* 12. t. 46.—N. Granat.
pumilum, *Wedd. l. c.* 13.—Bolivia.
subulatum, *Benth. & Hook. f. Gen.* ii. 549.—Ecuador.

RHIZOCEPHALUS, Boiss, Diagn. Ser. I. v. 68 (1844); xiii. 43 (1853) = **Heleochloa**, Host (Gramin.).
orientalis, Boiss. l. c. 69.—Mesopot.

RHIZOCORALLON, Hall. in Rupp. Fl. Jen. 301 (1745) = **Corallorhiza**, Hall. (Orchid.).

RHIZOGUM, Reichb. Consp. 125 (1828) = **Rhigozum**, Burch. (Bignon.).

RHIZOPHORA, Linn. Gen. ed. I. 137 (1737). *RHIZOPHOREAE*, Benth. & Hook. f. i. 679.
AZOPHORA, Neck. Elem. ii. 361 (1790).
MANGIUM, Rumph. ex Scop. Introd. 218 (1777).
MANGLE, Adans. Fam. ii. 445 (1763).
MANGLES, Plum. ex DC. Prod. iii. 32 (1828).
Aegiceras, Gaertn. f. Fruct. iii. t. 45. f. 1 = Aegiceras minus.
Aegiceras, J. F. Gmel. Syst. 747 = Aegiceras majus.
americana, Nutt. N. Am. Sylv. i. 112. t. 24 = R. Mangle.
apiculata, Blume, Fl. Jav. i. 91 = Bruguiera gymnorrhiza.
australis, Steud. Nom. ed. II. ii. 449 = Bruguiera Rheedii.
Candel, Linn. Sp. Pl. 443 = Kandelia Rheedii.
Candelaria, DC. Prod. i. 32 = Brugueria gymnorrhiza.
Candelaria, Macrae, ex Blume, Mus. Bot. Lugd. Bat. i. 137 = Bruguiera caryophylloides, malabarica.
Candelaria, Wall. Cat. n. 4878 = mucronata.
caryophylloides, *Burm. f. Fl. Ind.* 109.—As. trop.
caryophylloides, Jack, in Malay. Misc. i. (1820) n. v. 34 = Bruguiera caryophylloides.
caseolaris, Linn. Amoen. Acad. iv. 123 = Sonneratia acida.
ceratophylloides, Rausch. Nom. ed. III. 136, sphalm. = caryophylloides.
conjugata, Linn. Sp. Pl. 443 = Bruguiera gymnorrhiza.
corniculata, Linn. Amoen. Acad. iv. 123 = Aegiceras majus.
cylindrica, Linn. Sp. Pl. 443 = Bruguiera caryophylloides.
decandra, Roxb. Hort. Beng. [36], nomen = Ceriops Roxburghiana.

RHIZOPHORA:—
eriopetala, Steud. Nom. ed. II. ii. 449 = Bruguiera eriopetala.
glomerulata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 143 = Ceriops Zippeliana.
gymnorrhiza, Linn. Sp. Pl. 443 = Bruguiera gymnorrhiza.
Lamarckii, *Montr. in Mém. Acad. Lyon*, x. (1860) 201.—N. Caled.
latifolia, *Miq. Fl. Ind. Bat. Suppl.* 324.—Sumatra.
macrorhiza, Griff. Notul. iv. 664 (*macrohiza*) = mucronata.
Mangle, *Linn. Sp. Pl.* 443.—Littora trop.
Mangle, Roxb. Hort. Beng. [36]; Fl. Ind. ii. [459] = mucronata.
mucronata, *Lam. Encyc.* vi. 169.—Geront. trop.
mucronata, Naves, in Blanco, Fl. Philipp. ed. III. t. 135 = Bruguiera gymnorrhiza.
obtusa, Dennst. Schluess. Hort. Malab. 15, 32 (= *Problastes cuneifolia*).—Java.
pachypoda, *Baill. Adansonia*, xi. (1873-76) 309.—N. Caled.
Palun, *DC. Prod.* iii. 33.—Ins. Molucc.
parviflora, Roxb. Fl. Ind. ii. 461 = Bruguiera parviflora.
pauciflora, Griff. Notul. iv. 664, sphalm. ? = praec.
polandra, Blanco, Fl. Filip. ed. I. 396 = Bruguiera eriopetala.
racemosa, G. F. W. Mey. Prim. Fl. Esseq. 185 = R. Mangle.
Rheedii, Steud. Nom. ed. II. ii. 449 = Bruguiera gymnorrhiza.
rugens, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 233 = mucronata.
sexangula, *Lour. Fl. Cochinch.* 297.—Cochinch.
stylosa, Griff. Notul. iv. 665 = Kandelia Rheedii.
Tagal, Perr. in Mém. Soc. Linn. Par. iii. (1825) 138 (Quid?).—Guian.
timoriensis, DC. Prod. iii. 32 = Ceriops Candolleana.
tinctoria, Blanco, Fl. Filip. ed. I. 394 = Bruguiera gymnorrhiza.

RHODACTINEA, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 449 = **Barnadesia**, Mutis (Compos.).
rosea, Gardn. l. c. 450 = B. rosea.

RHODACTINIA, Benth. & Hook. f. Gen. ii. 484 (1873), sphalm. = praec.

RHODALSINE, J. Gay, in Ann. Sc. Nat. Sér. III. iv. (1845) 25, in adnot. = **Arenaria**, Rupp. (Caryophyll.).
procumbens, J. Gay, l. c. = A. geniculata.

RHODAMNIA, Jack, in Malay. Misc. ii. (1822) n. VII. 48. *MYRTACEAE*, Benth. & Hook. f. i. 715.
MONOXORA, Wight, Illustr. ii. 12. t. 97 (1841).
argentea, *Benth. Fl. Austral.* iii. 278.—Austral.
Blairiana, *F. Muell. Fragm.* ix. 141.—Austral.
cinerea, Jack, in Malay. Misc. ii. (1822) n. VII. 48 = trinervia.
concolor, Miq. Fl. Ind. Bat. Suppl. 315 = trinervia.
glauca, *Blume, Mus. Bot. Lugd. Bat.* i. 79.—N. Guin.
globosa, *Blume, l. c.*—Borneo.
latifolia, *Miq. Fl. Ind. Bat.* i. 1. 480.—Malaya.
Mulleri, Blume, Mus. Bot. Lugd. Bat. i. 79 = trinervia.
Nageli, Miq. Fl. Ind. Bat. i. 1. 478 = trinervia.
sessiliflora, *Benth. Fl. Austral.* iii. 277.—Austral.
spectabilis, Blume, Mus. Bot. Lugd. Bat. i. 78 = trinervia.
subtriflora, Blume, l. c. 79 = trinervia.
trinervia, *Blume, Mus. Bot. Lugd. Bat.* i. 79.—Burma; Malaya; Austral.

RHODANTHE, Lindl. Bot. Reg. t. 1703 (1834) = **Helipterum**, DC. (Compos.).
atropurpurea, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 776 = *atrosanguinea*.
atrosanguinea, J. Drumm. in Gard. Chron. (1861) 622.—Austral.
immaculata, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 967 = H. Manglesii.
maculata, Hort. cf. Gard. Chron. (1861) 622 = H. Manglesii?
Manglesii, Lindl. Bot. Reg. t. 1703 = H. Manglesii.

RHODAX, Spach, in Ann, Sc. Nat. Sér. II. vi. (1836) 363 = **Helianthemum**, Tourn. (Cistin.).
alpestris, Fuss, Fl. Transs. 78 = H. cinereum.
ambiguus, Fuss, l. c. 79.—Transs.
canus, Fuss. l. c. 79 = H. canum.
Chamaecistus, Spach, Hist. Veg. Phan. vi. 40 = H. cinereum.
dichotomus, Spach, l. c. 44 = H. dichotomum.
marifolius, Fuss, Fl. Transs. 79 = H. vineale.
montanus, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 364 = H. montanum.
oelandicus, Fuss, Fl. Transs. 78 = H. cinereum.
origanifolius, Spach, in Ann. Sc. Nat. Sér. II. iv. (1836) 364 = H. origanifolium.
penicillatus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 340 = H. cinereum.
polyanthus, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 364 = H. polyanthum.
prostratus, Pomel, Nouv. Mat. Fl. Atl. 354 = H. prostratum.
vinealis, Fuss, Fl. Transs. 79 = H. vineale.

RHODEA, Endl. Gen. 155 (1837) = **Rohdea**, Roth (Liliac.).

RHODIA, Linn. Syst. ed. I (1735) = **Sedum**, Tourn. (Crassul.).
officinarum, Crantz, Inst. i. 191 = Sedum roseum, *Scop.*

RHODIOLA, Linn. Gen. ed. I. 318 (1737) = **Sedum**, Tourn. (Crassul.)
algida, Fisch. & Mey. in Schrank, Enum. Pl. Nov. 70 = S. algidum.
asiatica, D. Don, Prod. Fl. Nep. 213 = S. asiaticum.
atropurpurea, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 39) = S. atropurpureum.
elongata, Fisch. & Mey. in Schrank. Enum. Pl. Nov. 68 = S. roseum, *Scop.*
gelida, Schrenk, Enum. Pl. Nov. 67 = S. gelidum.
imbricatum, Edgew. in Trans. Linn. Soc. xx. (1846) 47 = S. roseum, *Scop.*
integrifolia, Rafin. Atl. Journ. 146 = S. roseum, *Scop.*
Kirilowi, Regel, ex Maxim. Prim. Fl. Amur. 472 = S. Kirilowii.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = S. roseum, *Scop.*
odora, Salisb. Prod. 308 = S. roseum, *Scop.*
odorata, Lam. Fl. Fr. iii. 647 = S. roseum, *Scop.*
quadrifida, Fisch. & Mey. in Schrank, Enum. Pl. Nov. 69 = S. quadrifidum.
rosea, Linn. Sp. Pl. 1035 = S. roseum, *Scop.*
sibirica, Sweet, Hort. Brit. ed. III. 268 = S. roseum, *Scop.*
Stephani, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 39) = S. Stephani.

RHODIOLA, Lour. Fl. Cochinch. 627 (1790) = **Cardiospermum**, Linn. (Sapind.).
biternata, Lour. l. c. = C. Halicacabum (Masc.).

RHODOCALYX, Muell. Arg. in Mart. Fl. Bras. vi. 1. 172. t. 51 (1860). *APOCYNACEAE*, Benth. & Hook. f. ii. 725.
calycosus, Miers, Apocyn. S. Am. 140 = Echites calycosa.
cinereus, Miers, l. c. 141 = Haplophyton cimicidum.
coccineus, Miers, l. c. = Dipladenia coccinea.
crassifolius, Miers, l. c. 139 = Echites crassifolia.
crassipes, Miers, l. c. 140 = Echites crassipes.
cuneifolius, *Miers, l. c.* 142.—Bras.
hypoleucus, Miers, l. c. 140 = Macrosiphonia hypoleuca.
lanuginosus, Miers, l. c. 139 = Macrosiphonia lanuginosa.
ovatus, Miers, l. c. 141 = Dipladenia coccinea.
rotundifolius, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 173.—Bras.
suaveolens, Miers, Apocyn. S. Am. 139 = Macrosiphonia hypoleuca.
Tweedianus, *Miers, l. c.* 142.—Bras.

RHODOCHITON, Zucc. in Abh. Akad. Muench. i. (1832) 306, in syn. t. 13. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 935.
volubile, *Zucc. l. c.*—Mexico.

RHODOCHLAENA, Spreng. Syst. ii. 595 (1825) = **Rhodolaena**, Thou. (Chlaenac.).

RHODOCHLAMYS, Schau. in Linnaea, xx. (1847) 706 = **Salvia**, Tourn. (Labiat.).
speciosa, Schau. l. c. 707 = S. Sessei.

RHODOCISTUS, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 367 = **Cistus**, Tourn.
Berthelotianus, Spach, l. c. = C. vaginatus.

RHODOCLADA, Baker, in Journ. Linn. Soc. xxi. (1884) 327. *LINACEAE.*
rhopaloides, *Baker, l. c.* 328.—Madag.

RHODOCODON, Baker, in Journ. Linn. Soc. xviii. (1881) 280. t. 8. *LILIACEAE*, Benth. & Hook. f. iii. 811.
madagascariensis, *Baker, l. c.*—Madag.

RHODOCOMA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836) = **Restio**, Linn.
capense, Nees, ex Steud. Syn. Pl. Cyp. 249 = Restio Rhodocoma?
Equisetum, Nees, ex Mast. in Journ. Linn. Soc. x. (1869) 275 = Restio Rhodocoma.

RHODODENDRON, Linn. Sp. Pl. ed. I. 392 (1753).
ERICACEAE, Benth. & Hook. f. ii. 599.
ANTHODENDRON, Reichb. in Moessl. Handb. i. p. xl. 244, 308 (1827).
AZALEA, Linn. Syst. ed. I. (1735).
BRACHYCALYX, Sweet, ex Steud. Nom. ed. II. i. 220 (1840).
CHAMAECISTUS, Regel, Gartenfl. (1874) 60.
HOCHENWARTIA, Crantz, Inst. ii. 468 (1766).
HYMENANTHES, Blume, Bijdr. 862 (1826).
LOISELERIA, Reichb. Fl. Germ. Excurs. 416, in nota (1831).
LOISELEURIA, Reichb. Nom. 127 (1841).
OSMOTHAMNUS, DC. Prod. vii. 715 (1839).
PLINTHOCHROMA, Dulac, Fl. Hautes-Pyr. 419 (1867).
RHODORA, Linn. Sp. Pl. ed. II. 561 (1762).
RHODOTHAMNUS, Lindl. & Paxt. Flow. Gard. i. (1850–51) t. 22.
STEMOTIS, Rafin. Sylva Tellur. 152 (1838).
THEIS, Salisb. ex DC. Prod. vii. 715 (1835).
TSUTSUSI, Adans. Fam. ii. 164 (1763).
VIREYA, Blume, Bijdr. 854 (1826).
acuminatum, *Hook. f. in Hook. Ic. Pl.* t. 886.—Borneo.
Adansoni, *Pépin, Ann. Fl. et Pom.* (1839) 304.—Hab.?
aeruginosum, Hook. f. Rhod. Sikkim-Himal. t. 22 = campanulatum.
afghanicum, *Aitch. & Hemsl. in Journ. Linn. Soc* xviii. (1880) 75.—Afghan.
albiflorum, *Hook. Fl. Bor. Am.* ii. 43. t. 33; *ex R. Grah. in Edinb. N. Phil. Journ.* xxiv. (1838) 422.—Am. bor. occ.
Albrechtii, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 227.—Japon.
album, *Blume, Cat. Gew. Buitenz.* 72.—Java.
album, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 154 = arboreum.
album, Hoffmgg. Verz. Pfl. 192 = albiflorum.
algarvense, Page, Prod. Southhampt. Gard. 38 = baeticum?
alstroemerioides ×, Lem. Jard. Fleur. iv. (1854) t. 384.
altaclerense, Lindl. Bot. Reg. t. 1414 = arboreum.
amoenum, Planch. in Fl. des Serres, ix. (1853–54) 80; et in Rev. Hort. Sér. IV. iii. (1854) 64 = indicum.
angustifolium, Hoppe, in Flora, xx. (1837) 185, 205 = hirsutum.
Anthopogon, *D. Don, in Mem. Wern. Soc.* iii. (1821) 409.—As. bor.; Reg. Himal.
Anthopogon, Wall. Cat. n. 759, partim = setosum.
anthopogonoides, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 350.—China.
apoanum, *Stein, Gartenfl.* (1885) 194. t. 1196.—Ins. Philipp.
arborescens, *Torr. Fl. U. St.* i. 425.—Am. bor.
arboreum, *Sm. Exot. Bot.* i. 9.—Ind. or.; Reg. Himal.; Zeylan.
arfakianum, *Becc. Malesia*, i. (1878) 201.—N. Guin.
argenteum, Hook. f. Rhod. Sikkim-Himal. t. 9 = grande.
aristatum, Royle, Illustr. Bot. Himal. 42, nomen = barbatum?

RHODODENDRON:—
aromaticum, Wall. Cat. n. 1520 = R. Anthopogon.
assamicum, Hort.—Cf. Gard. Chron. (1881) i. 8 (Quid?).—Assam.
Aucklandii, Hook. f. Rhod. Sikkim-Himal. t. 11 = Griffithianum.
aureum, Georgi, Reise Russ. Reich. i. 214 = chrysanthum.
azaleoides, Desf. Tabl. ed. II. 272 = ponticum.
baeticum, *Boiss. & Reut. Diagn.* Ser. II. iii. 118.—Hispan.; Lusit.
balsaminaeflorum ×, T. Moore, Fl. & Pomol. (1883) 81. t. 587.
barbatum, *G. Don, Gen. Syst.* iii. 844.—Reg. Himal.
Batemani, Hook. f. Bot. Mag. t. 5387 = campanulatum.
bhotanicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 475.—Reg. Himal.
bicolor, Sweet, Hort. Brit. ed. II. 344 = nudiflorum.
bigener, Hort. ex Steud. Nom. ed. II. ii. 450 = maximum.
blandfordiaeflorum, Hook. Bot. Mag. t. 4930 = cinnabarinum.
Blandinianum, *Martr. in Hérincq, Hortic. Franç.* (1859) 152. t. 13.—Am. bor.
Blumei, Nutt. in Hook. Kew Journ. v. (1853) 366 = niveum.
Boothii, *Nutt. l. c.* 356.—Reg. Himal.
brachycarpum, *D. Don, ex G. Don, Gen. Syst.* iii. 843.—Japon.
brachycarpum, Zucc. ex Maxim. in Mém. Acad. Pétersb. Sér. VII. xvi. (1871) 39 = indicum.
Breynii, Planch. in Fl. des Sérres, ix. (1853–54) 77; et in Rev. Hortic. Sér. IV. iii. (1854) 46 = indicum.
Brookeanum, *Low, ex Lindl. in Journ. Hort. Soc.* iii. (1848) 82.—Borneo.
Buergeri, Miq. Ann. Mus. Bot. Lugd. Bat. i. 34 = indicum.
Burmanni, G. Don, Gen. Syst. iii. 846 = ledifolium.
buxifolium, *Low, ex Hook. f. in Hook. Ic. Pl.* t. 890.—Borneo.
calendulaceum, Hook. & Arn. Bot. Beech. Voy. 362 = occidentale.
calendulaceum, *Torr. Fl. U. St.* i. 425.—Am. bor.
californicum, *Hook. Bot. Mag.* t. 4863.—Am. bor. occ.
Calleryi, Planch. in Fl. des Sérres, ix. (1853–54) 81; in Rev. Hortic. Sér. IV. iii. (1854) 66 = indicum.
calophyllum, Nutt. in Hook. Kew Journ. v. (1853) 362 = Maddeni.
calycinum, Planch. in Fl. des Serres, ix. (1853–54) 81; et in Rev. Hortic. Sér. IV. iii. (1854) 65 = ledifolium.
camelliaeflorum, *Hook. f. Rhod. Sikkim-Himal.* t. 28.—Reg. Himal.
campanulatum, *D. Don. in Mem. Wern. Soc.* iii. (1821) 410.—Reg. Himal.
Campbelliae, Hook. f. Rhod. Sikkim-Himal. t. 6 = arboreum.
campylocarpum, *Hook. f. l. c.* t. 30.—Reg. Himal.
campylogynum, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 10.—China.
camtschaticum, *Pall. Fl. Ross.* i. 48. t. 33.—Kamtschat.; Ins. Aleut.
Candelabrum, Hook. f. Rhod. Sikkim-Himal. t. 29 = Thomsonii.
canescens, G. Don, Gen. Syst. iii. 848 = nudiflorum.
capitatum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 351.—China.
Carnarvoni ×, Sweet, Hort. Brit. ed. II. 342, nomen.
Cartonianum ×, DC. Prod. vii. 727.
catawbiense, Hoffmgg. Verz. Pfl. Nachtr. ii. 194 = ellipticum.
catawbiense, *Michx. Fl. Bor. Am.* i. 258.—Am. bor.
Catesbaei, Forbes, Hort. Woburn. 90 = seq.
Catesbaeum, Dum.-Cours. Bot. Cult. ed. II. vii. 165 = ponticum?
caucaseum, Sims, Bot. Mag. t. 1145 = seq.
caucasicum, *Pall. Fl. Ross.* i. 46. t. 31.—Reg. Caucas.
Cavroni ×, André, in Rev. Hortic. (1885) 60, cum ic.
celebicum, DC. Prod. vii. 725 = malayanum.
cephalanthum, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 9.—China.
Chamaecistus, Cham. & Schlecht. in Linnaea, i. (1826) 513 = Redowskianum.
Chamaecistus, Linn. Sp. Pl. 392 = Rodothamnus Chamaecistus.
Championae, *Hook. Bot. Mag.* t. 4609.—China.
Chapmanii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 61.—Am. bor.

RHODODENDRON :—
chrysanthum, *Pall. Reise*, iii. 729.—Davuria.
ciliatum, *Hook. f. Rhod. Sikkim-Himal.* t. 24.—Reg. Himal.
cinnabarinum, *Hook. f. l. c.* t. 8.—Reg. Himal.
cinnamomeum, Wall. Cat. n. 760 = arboreum.
cistiflorum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Chamaecistus.
citrinum, *Hassk. Cat. Pl. Hort. Bog. Alt.* 161.—Java.
Clivianum ×, J. Sm. in Bot. Mag. t. 4478.
Collettianum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 75.—Afghan.
cordifolium, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 23.—Hab. ?
crispiflorum, Planch. in Fl. des Serres, ix. (1853–54) 83 = indicum.
Cunninghami, Hort. ex C. Koch, Dendrol. ii. 1. 166 = ponticum.
Curtisii ×, T. Moore, Flor. & Pomol. (1884) 113. t. 615.
Dalhousiae, *Hook. f. Rhod. Sikkim-Himal.* 2. t. 2.—Reg. Himal.
Danielsianum, Planch. in Fl. des Serres, ix. (1853–54) 79 = indicum.
dauricum, *Linn. Sp. Pl.* 392.—Sibir.
deciduum, Andr. ex Steud. Nom. ed. II. ii. 450 = ponticum.
decumbens, D. Don, ex G. Don, Gen. Syst. iii. 846 = indicum.
deflexum, Griff. Notul. iv. 303; Ic. Pl. As. t. 519 = triflorum.
Degronianum, *Carr. in Rev. Hortic.* (1869) 368.—Japon.
dilatatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 34.—Japon.
durionifolium, *Becc. Malesia*, i. (1878) 202.—Borneo.
Edgarii, Gamble, Darjeeling List Trees, 52 = campanulatum.
Edgeworthii, *Hook. f. Rhod. Sikkim-Himal.* 2. t. 21.—Reg. Himal.
elaeagnoides, Hook. f. l. c. t. 23 B = lepidotum.
ellipticum, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 194.—Am. bor.
enneandrum ×, DC. Prod. vii. 728.
ericoides, *Low, ex Hook. f. in Hook. Ic. Pl.* t. 887.—Borneo.
eximium, Nutt. in Hook. Kew Journ. v. (1853) 364 = Falconeri.
Falconeri, *Hook. f. Rhod. Sikkim-Himal.* t. 10.—Reg. Himal.
Farrerae, *Tate, ex Sweet, Brit. Flow. Gard.* Ser. II. t. 95.—China.
ferrugineum, *Linn. Sp. Pl.* 392.—Europ.; As. bor.
flavum, G. Don, Gen. Syst. iii. 847 = ponticum.
flavum, Pall. Reise, iii. 320 = chrysanthum.
formosum, *Wall. Pl. As. Rar.* iii. 3. t. 207.—Reg. Himal.
Fortuni, [*T. Moore,*] *Lindl. in Gard. Chron.* (1859) 868; *Hook. f. Bot. Mag.* t. 5595 (Fortunei).—China.
fragrans, Hort. ex DC. Prod. vii. 722 = maximum.
fragrans, Maxim. Rhod. As. Or. 16 = R. Anthopogon.
fulgens, *Hook. f. Rhod. Sikkim-Himal.* t. 25.—Reg. Himal.
gaultherioides, Boiss. & Bal. ex Boiss. Pl. Or. Nov. Dec. i. 4 = Orphanidesia gaultherioides.
germanicum, Hoppe, in Flora, xx. (1837) 187 = hirsutum.
Gibsonii, Paxt. Mag. Bot. viii. (1841) t. 217 = formosum.
glabratum, Hoppe, in Flora, xx. (1837) 185 = hirsutum.
glaucum, G. Don, Gen. Syst. iii. 848 = viscosum.
glaucum, *Hook. f. Rhod. Sikkim-Himal.* t. 17.—Reg. Himal.
Gowenianum ×, Sweet, Brit. Flow. Gard. Ser. I. t. 263.
gracile, *Low, ex Lindl. in Journ. Hort. Soc.* iii. (1848) 84.—Borneo.
grande, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 176.—Reg. Himal.
Griffithianum, *Wight, Ic. Pl.* t. 1203.—Reg. Himal.
Griffithii, Hook. f. Rhod. Sikkim-Himal. 2, sphalm. = praec.
halense ×, Gremb. in Oestr. Bot. Zeitschr. xxiv. (1874) 375.—Europ.
hatamense, *Becc. Malesia*, i. (1878) 202.—N. Guin.
Henryi, *Hance, in Journ. Bot.* xix. (1881) 241; xxii. (1884) 22.—China.

RHODODENDRON :—
hirsutiformum ×, Gremb. in Oestr. Bot. Zeitschr. xxiv. (1874) 376.—Europ.
hirsutum, *Linn. Sp. Pl.* 392.—Europ. med.
hispidum, *D. Don, in Edinb. Phil. Journ.* vi. (1822) 48.—Japon.
hispidum, Torr. Fl. U. St. i. 425 = viscosum.
Hodgsonii, *Hook. f. Rhod. Sikkim-Himal.* t. 15.—Reg. Himal.
Hookeri, *Nutt. in Hook. Kew Journ.* v. (1853) 358.—Reg. Himal.
hyacinthiflorum, Hort. ex Steud. Nom. ed. II. ii. 450 = ponticum.
hybridum ×, Ker-Gawl. in Bot. Reg. t. 195.
indicum, *Sweet, Hort. Brit.* ed. II. 343; *Brit. Flow. Gard.* Ser. II. t. 128.—China.
intermedium, Schur, Enum. Pl. Transs. 448 = hirsutum.
intermedium, Tausch, in Flora, xix. (1836) 36 = ferrugineum.
intermedium, Wender. l. c. ix. (1826) 357 = praec. ?
jasminiflorum, *Hook. Bot. Mag.* t. 4524.—Java.
javanicum, *Benn. Pl. Jav. Rar.* 85. t. 19.—Java.
Jenkinsii, *Nutt. in Hook. Kew Journ.* v. (1853) 361.—Reg. Himal.
Jodogawa, Hort. Bog. ex Hassk. Cat. Hort. Bog. Alt. 162 = Keiskei.
Kaempferi, Planch. in Fl. des Serres, ix. (1853–54) 77 = indicum.
kamtschaticum, *Pall. Fl. Ross.* i. 48. t. 33.—As. bor. or.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 163.—Japon.
Kendrickii, *Nutt. in Ann. & Mag. Nat. Hist.* Ser. II. xii. (1853) 10.—Reg. Himal.
Knightianum ×, Forbes, Hort. Woburn. 90.
Keysii, *Nutt. in Hook. Kew Journ.* v. (1853) 353.—Reg. Himal.
Kochii, *Stein, in Gartenfl.* xxxii. (1885) 193. t. 1195.—Ins. Philipp.
Koempferi, Planch. in Rev. Hortic. Sér. IV. iii. (1854) 46, sphalm. = *Kaempferi*.
Konori, *Becc. Malesia*, i. (1878) 200.—N. Guin.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 43.—Sumatra.
lampongum, Miq. Fl. Ind. Bat. Suppl. 585 = malayanum.
lanatum, *Hook. f. Rhod. Sikkim-Himal.* t. 16.—Reg. Himal.
lancifolium, Hook. f. l. c. t. 4 = barbatum.
lancifolium, Moench, Meth. 45 = ponticum.
lapponicum, *Wahlenb. Fl. Suec.* 249.—Reg. bor. et arct.
lateritium, Planch. in Fl. des Serres, ix. (1853–54) 80; et in Rev. Hortic. Sér. IV. iii. (1854) 63 = indicum.
latifolium, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 195.—Am. bor. ?
latifolium, Hoppe, in Flora, xxiii. (1840) 525 = hirsutum.
ledifolium, *G. Don, Gen. Syst.* iii. 846.—China.
lepidotum, *Wall. Cat.* n. 758.—Reg. Himal.
leptocarpum, *Nutt. in Hook. Kew Journ.* vi. (1854) 256.—Reg. Himal.
leucanthum, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 115 = ledifolium.
Lindleyi, *T. Moore, in Gard. Chron.* (1864) 364.—Ind. or. ?
linearifolium, *Poir. Encyc.* vi. 267.—Ind. or. ?
linearifolium, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 131.—Japon.
Lobbii, *Hort.—Cf. Gard. Chron.* (1871) 1323.—Hab. ?
longiflorum, *Lindl. in Journ. Hort. Soc.* iii. (1848) 88.—Borneo.
longifolium, Nutt. in Hook. Kew Journ. v. (1853) 366 = grande.
Loureiriana, *G. Don, Gen. Syst.* iii. 846.—Cochinch.
Lowei, *Hook. f. in Hook. Ic. Pl.* t. 883.—Borneo.
Lowei, Loud. Gard. Mag. xi. (1835) 79, 190 = ponticum.
lucidum, *Nutt. in Hook. Kew Journ.* v. (1853) 363.—Reg. Himal.
luteum, *Sweet, Hort. Brit.* ed. II. 343.—Turcia.
macranthum, G. Don, Gen. Syst. iii. 846 = indicum.
macranthum, Griff. Notul. iv. 303; Ic. Pl. As. t. 520 = Maddeni.
macrocarpos, Griff. Itin. Notes, 138 = Dalhousiae.

RHODODENDRON :—
macrocarpum, Van Houtte, in Fl. des Serres, xi. (1857) 83 (Quid ?).—Ind. or.
macrophyllum, *D. Don, ex G. Don, Gen. Syst.* iii. 843.—Am. bor. occ.
macrosepalum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 228.—Japon.
macrostemon, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. ix. 41.—Japon.
Maddeni, *Hook. f. Rhod. Sikkim-Himal.* t. 18.—Reg. Himal.
malayanum, *Jack, in Malay. Misc.* ii. (1822) n. vii. 17.—Sumatra.
Manglesi ×, Hort. Veitch.—Cf. Gard. Chron. (1885) ii. 48.
Mariae, *Hance, in Journ. Bot.* xx. (1882) 230.—China.
maximum, Hook. Fl. Bor. Am. ii. 43 (excl. syn.) = macrophyllum.
maximum, *Linn. Sp. Pl.* 392.—Am. bor.
maximum, Thunb. Fl. Jap. 181 = Metternichii.
Metternichii, *Sieb. & Zucc. Fl. Jap.* i. 23. t. 9.—Japon.
micranthum, Maxim. in Mém. Acad. Pétersb. Sér. VII. xvi. (1870) n. ix. 18 = R. Anthopogon.
micranthum, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 586.—China.
minus, Michx. Fl. Bor. Am. i. 258 = punctatum.
molle, *G. Don, Gen. Syst.* iii. 846.—China.
molle, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 131 = sinense.
Morterii ×, Sweet, Brit. Flow. Gard. Ser. II. i. t. 10.
moulmainense, *Hook. Bot. Mag.* t. 4904.—Burma.
mucronatum, G. Don, Gen. Syst. iii. 846 = ledifolium.
mucronulatum, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. vii. 155; xxi. (1848) i. 585 = dauricum.
multicolor, *Miq. Fl. Ind. Bat. Suppl.* 586.—Sumatra.
mutabile, Webb, ex Royle, Illustr. Bot. Himal. 259 = campanulatum.
myrtifolium, Lodd. Bot. Cab. t. 908 = ponticum.
myrtifolium, Schott & Kotschy, ex Schott, in Bot. Zeit. ix. (1851) 17 = ferrugineum.
narcissiflorum, *Planch. in Fl. des Serres*, ix. (1853–54) 82; et in Rev. Hortic. Sér. IV. iii. (1854) 67 (sp. dub.).—China.
nilagiricum, Zenk. in Ann. Sc. Nat. Sér. II. vi. (1836) 150; Pl. Ind. t. 15 = arboreum.
nitidum, Torr. Fl. U. St. i. 425 = viscosum.
nivale, *Hook. f. Rhod. Sikkim-Himal.* t. 26.—Reg. Himal.
niveum, *Hook. f. in Journ. Hort. Soc.* vii. (1852) 78.—Reg. Himal.
nobile, Wall. Cat. n. 1521 = campanulatum, barbatum.
Nobleanum, Hort. ex Lindl. Bot. Reg. t. 1820. f. 1 = caucasicum.
nudiflorum, *Torr. Fl. U. St.* i. 424.—Am. bor.
Nuttallii, *Booth, ex Nutt. in Hook. Kew Journ.* v. (1853) 355.—Reg. Himal.
oblongum, Griff. Notul. iv. 303; Ic. Pl. As. t. 523 = Griffithianum.
obovatum, Hook. f. Rhod. Sikkim-Himal. Consp. 6 = lepidotum.
obtusum, Hort. ex Wats. Dendrol. ii. t. 162 = ponticum.
obtusum, Planch. in Fl. des Serres, ix. (1853–54) 80 et in Rev. Hortic. Sér. IV. iii. (1854) 64 = indicum.
occidentale, *A. Gray, in Bot. Calif.* i. 458.—Am. bor. occ.
odoratum, Lodd. ex Steud. Nom. ed. II. ii. 450 = ponticum.
officinale, Salisb. Parad. Lond. t. 80 = chrysanthum.
Oldhami, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. ix. 34.—Ins. Formosa.
orbiculare, *Decne. in Fl. des Serres*, xxii. (1877) 169.—Tibet.
ornatissimum ×, Versch. in Illustr. Hortic. xiv. (1867) t. 530.
ornatum, Sweet, Brit. Flow. Gard. Ser. II. t. 137 = viscosum.
ovatum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 230.—China.
palustre, Turcz. ex DC. Prod. vii. 724 = R. Anthopogon.
papuanum, *Becc. Malesia*, i. (1878) 201.—N. Guin.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 475.—Burma.
parviflorum, Dum.-Cours. Bot. Cult. ed. I. 253 = ponticum.
parviflorum, Hort. ex DC. Prod. vii. 723 = punctatum

RHODODENDRON:—
parviflorum, F. Schmidt, in Mém. Acad. Pétersb. Sér. VII. xii. (1868) n. II. (Reisen Amurl.) 158, sphalm. = *parvifolium*.
parvifolium, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 237 = R. Anthopogon.
pelargoniaeflorum ×, Van Houtte, in Fl. des Serres, x. (1854–55) 225. t. 1063.
pendulum, *Hook. f. Rhod. Sikkim-Himal.* t. 13.—Reg. Himal.
phoeniceum, G. Don, Gen. Syst. iii. 846 = ledifolium.
pictum ×, Forbes, Hort. Woburn. 232.
planifolium, Nutt. in Hook. Kew Journ. v. (1853) 365 = campanulatum.
polifolium, Scop. Fl. Carn. ed. II. i. 287 = Andromeda Polifolia.
ponticum, *Linn. Sp. Pl.* ed. II. 562.—As. Min.; Syria.
praecox ×, Davis, ex Gard. Chron. (1878) I. 335.
procerum, Salisb. Prod. 287 = maximum.
prunifolium, *Forbes, Hort. Woburn.* 91 (Quid?).—Ind. or.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 350.—China.
pulchellum, Salisb. Prod. 287 = R. Rhodora.
pulcherrimum, Lindl. Bot. Reg. t. 1820 = caucasicum.
pulchrum, Sweet, Brit. Flow. Gard. Ser. II. t. 117 = ledifolium.
pumilum, *Hook. f. Rhod. Sikkim-Himal.* t. 14.—Reg. Himal.
pumilum, Nutt. in Hook. Kew Journ. v. (1853) 354 = leptocarpum.
punctatum, *Andr. Bot. Rep.* t. 36.—Am. bor.
puniceum ×, Planch. in Fl. des Serres, ix. (1853–54) 79.
puniceum, Roxb. Hort. Beng. 33; Fl. Ind. ii. 409 = arboreum.
purpureum, G. Don, Gen. Syst. iii. 843 = maximum.
purpureum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 154 = arboreum.
Purshii, G. Don, l. c. = maximum.
quinquefolium, *Bisset & S. Moore, in Journ. Bot.* xv. (1877) 292.—Japon.
ramentaceum, Planch. in Fl. des Serres, ix. (1853–54) 81; et in Rev. Hortic. Sér. IV. iii. (1854) 65 = indicum.
Redowskianum, *Maxim. Prim. Fl. Amur.* 189.—Reg. Amur.
reticulatum, *D. Don, ex G. Don, Gen. Syst.* iii. 846.—Japon.
retusum, *Benn. Pl. Jav. Rar.* 88. t. 20.—Java; Sumatr.
Rhodora, *J. F. Gmel. Syst.* 694.—Am. bor.
rhombicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 164.—Japon.
Rollissonii, Lindl. Bot. Reg. (1813) t. 25 = arboreum.
Rovellii ×, Leroy, in Rev. Hortic. (1867) 159.
Roylei, Hook. f. Rhod. Sikkim-Himal. t. 7 = cinnabarinum.
rugosum, *Low, ex Hook. f. in Hook. Ic. Pl.* t. 885.—Borneo.
Russelianum ×, Sweet, Brit. Flow. Gard. Ser. II. t. 9 = catawbiense.
salicifolium, *Becc. Malesia*, i. (1878) 202.—Borneo.
salignum, Hook. f. Rhod. Sikkim-Himal. Consp. 6. t. 23 = lepidotum.
scabrum, G. Don, Gen. Syst. iii. 846 = Metternichii.
Schlippenbachii, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 226.—Korea.
semibarbatum, *Maxim. l. c.* 229.—Japon.
Seniavini, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. IX. 38.—China.
serpyllifolium, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 165.—Japon.
setosum, *D. Don, in Trans. Wern. Soc.* iii. (1821) 408.—Reg. Himal.
Seymouri ×, Herb. ex Otto & Dietr. Allg. Gartenz. v. (1837) 312.
Shepherdii, *Nutt. in Hook. Kew Journ.* v. (1853) 360.—Reg. Himal.
Sieboldii, Miq. Ann. Mus. Bot. Lugd. Bat. i. 33 = indicum.
simiarum, *Hance, in Journ. Bot.* xix. (1881) 243; xxii. (1884) 22.—China.
Simsii, Planch. in Fl. des Serres, ix. (1853–54) 78 = indicum.
sinense, *Sweet, Brit. Flow. Gard.* Ser. I. t. 290.—Japon.
Smirnowi, *Trautv. ex Regel, in Gartenfl.* (1885) 335.—Reg. Caucas.
Smithii, Nutt. in Hook. Bot. Mag. t. 5120 = barbatum.
Smithii ×, Sweet, Hort. Brit. ed. II. 342.

RHODODENDRON:—
sparsiflorum, Nutt. in Hook. Kew Journ. v. (1853) 363 = camelliaeflorum.
speciosum, Salisb. Prod. 287 = ponticum.
speciosum, Sweet, Hort. Brit. ed. II. 343 = indicum?
Standishii ×, Paxt. Mag. Bot. xv. (1849) 210.
subcordatum, *Becc. Malesia*, i. (1878) 203.—Borneo.
sublanceolatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 163.—Japon.
Taylori ×, Veitch, in Flor. Mag. N. S. (1877) t. 242.
Teysmannii, *Miq. Fl. Ind. Bat. Suppl.* 585.—Sumatra.
thibaudiense, Hort. ex Dombr. Fl. Mag. iv. (1865) t. 253 = cinnabarinum.
Thomsonii, *Hook. f. Rhod. Sikkim-Himal.* t. 12.—Reg. Himal.
Thunbergii, Planch. in Fl. des Serres, ix. (1853–54) 78 = indicum.
thymifolium, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 351.—China.
tigrinum, Hort. ex Steud. Nom. ed. II. ii. 451 = maximum.
Toverenae, *F. Muell. in Vict. Natural.* i. (1884) 101.—N. Guin.
tricolor ×, Sweet, Hort. Brit. ed. II. 343.
triflorum, *Hook. f. Rhod. Sikkim-Himal.* t. 19.—Reg. Himal.
trispidum, Torr. Fl. North & Middle St. i. 184, sphalm. = *hispidum*, Torr.
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 230.—Japon.
tubiflorum, DC. Prod. vii. 725 = malayanum.
undulatum, Steud. Nom. ed. II. ii. 451 = arboreum.
Ungerni, *Trautv. ex Regel, in Gartenfl.* (1885) 335.—Reg. Caucas.
vaccinioides, *Hook. f. Rhod. Sikkim-Himal.* t. 3.—Reg. Himal.
variolosum, *Becc. Malesia*, i. (1878) 206.—Borneo.
Vaseyi, *A. Gray, in Proc. Am. Acad.* xv. (1880) 48.—Am. bor.
Veitchianum, *Hook. Bot. Mag.* t. 4992.—Burma.
velutinum, *Becc. Malesia*, i. (1878) 204.—Borneo.
venosum, Nutt. in Hook. Kew Journ. v. (1853) 365 = Falconeri.
venustum, Salisb. Prod. 287 = arboreum.
verticillatum, *Low, ex Lindl. in Journ. Hort. Soc.* iii. (1848) 86.—Borneo.
virgatum, *Hook. f. Rhod. Sikkim-Himal.* t. 26.—Reg. Himal.
viscosum, Fisch. ex Herd. in Act. Hort. Petrop. i. (1870) 350 = Redowskianum.
viscosum, *Torr. Fl. U. St.* i. 424.—Am. bor.
vittatum, *Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 66.—China.
Wallichii, Hook. f. Rhod. Sikkim-Himal. t. 6 = campanulatum.
Weyrichii, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xvi. (1871) n. IX. 26.—Japon.
Wightii, *Hook. f. Rhod. Sikkim-Himal.* t. 27.—Reg. Himal.
Wilsoni, Nutt. ex Hook. Bot. Mag. t. 5116 = grande.
Windsorii, Nutt. in Hook. Kew Journ. v. (1853) 357 = arboreum.
zeylanicum, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1364 = arboreum.

RHODOGERON, Griseb. Cat. Pl. Cub. 151 (1866). *COMPOSITAE*, Benth. & Hook. f. ii. 291.
coronopifolius, *Griseb. l. c.* 152.—Cuba.

RHODOLAENA, Thou. Hist. Vég. Isles. Afr. 47. t. 13 (1806). *CHLAENACEAE*, Benth. & Hook. f. i. 195.
PANDORA, Noronha, ex Thou. l. c. 48 (1806).
RHODOCHLAENA, Spreng. Syst. ii. 595 (1825).
acutifolia, *Baker, in Journ. Linn. Soc.* xxi. (1884) 322.—Madag.
altivola, *Thou. Hist. Vég. Isles. Afr.* 48. t. 13.—Madag.

RHODOLEIA, Champ. ex Hook. Bot. Mag. t. 4509 (1850). *HAMAMELIDEAE*, Benth. & Hook. f. i. 668.
Championi, *Hook. l. c.*—China.
formosa, Champ. ex Hance, in Walp. Ann. iii. 842 = Championi.
Teysmanni, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* vi. (1857) 123.—Sumatra.

RHODOLIRION, Phil. in Linnaea, xxix. (1857–58) 65 = **Hippeastrum**, Herb. (Amaryll.).
andinum, Phil. l. c. 66 = H. Rhodolirion.
montanum, Phil. l. c. 65 = H. uniflorum.

RHODOMYRTUS, Reichb. Nom. 117 (1841). *MYRTACEAE*, Benth. & Hook. f. i. 713.
cymiflora, *F. Muell. ex Benth. Fl. Austral.* iii. 273.—Austral.
macrocarpa, *Benth. Fl. Austral.* iii. 273.—Austral.
psidioides, *Benth. l. c.* 272.—Austral.
tomentosa, *Wight, Spicil. Neilgherr.* i. 60. t. 71.—Ind. or.; Malaya.
trineura, *F. Muell. ex Benth. Fl. Austral.* iii. 272.—Austral.

RHODOPHIALA, Presl, Bot. Bemerk. 115 (1844) **Hippeastrum**, Herb. (Amaryll.)
amarylloides, Presl, l. c. = H. pratense.
andina, Phil. in Anal. Univ. Chil. (1873) 543 = H. Herbertianum.
biflora, Phil. in Linnaea, xxix. (1857–58) 66.—Chili.
laeta, Phil. Fl. Atac. 51 = H. pratense.
modesta, Phil. in Anal. Univ. Chil. (1873) 544 = H. modestum.
uniflora, Phil. Fl. Atac. 51 = H. uniflorum.
Volckmanni, Phil. in Linnaea, xxxiii. (1864–65) 259 = H. pratense.

RHODOPHORA, Neck. Elem. ii. 91 (1790) = **Rosa**, Tourn.

RHODOPSIS, Lilja, Fl. Sverig. Suppl. i. 42 (1840) = **Calandrinia**, H. B. & K. (Portulac).
discolor, Lilja, l. c. = C. discolor.
speciosa, Lilja, l. c. = C. Menziesii?

RHODOPSIS, Reichb. Nom. 168 (1841) = **Rosa**, Tourn.

RHODOPTERA, Rafin. Fl. Tellur. iii. 45 (1836) = **Rumex**, Tourn. (Polygon.).
roseus, Rafin. l. c. 46 = Rumex roseus.

RHODORA, Linn. Sp. Pl. ed. II. 561 (1763) = **Rhododendron**, Linn. (Ericac.).
canadensis, Linn. l. c. = Rhodod. Rhodora.
congesta, Moench, Meth. 68 = Rhodod. Rhodora.

RHODORHIZA, Webb, in Bot. Reg. (1841) Misc. 69 = **Convolvulus**, Linn.
florida, Webb, l. c. 70 = C. floridus.
fruticulosa, Webb & Berth. Phyt. Canar. iii. 33 = C. fruticulosus.
glandulosa, Webb, in Bot. Reg. (1841) Misc. 70 = C. fruticulosus.
Perraudieri, C. Bolle, in Bonplandia, ix. (1861) 54 = C. Perraudieri (in Addend.).
scoparia, Webb, in Bot. Reg. (1841) Misc. 70 = C. scoparius.
virgata, Webb & Berth. Phyt. Canar. iii. 30 = C. scoparius.
volubilis, C. Bolle, in Bonplandia, ix. (1861) 54 = C. Massoni.

RHODORMIS, Rafin. Fl. Tellur. iii. 93 (1836) = **Salvia**, Tourn. (Labiat.).
rosaefolia, Rafin. l. c. 94 = S. rosaefolia.

RHODORRHIZA, Webb & Berth. Phyt. Canar. iii. 28 (1841–60) = RHODORIZA, Webb = **Convolvulus**, Linn.

RHODOSERIS, Turcz. in Bull Soc. Nat. Mosc. xxiv. (1851) II. 94 = **Onoseris**, DC. (Compos.).
conspicua, Turcz. l. c. 95 = O. Isotypus.

RHODOSPATHA, Poepp. & Endl. Nov. Gen. et Sp. iii. 91. t. 300 (1845). *AROIDEAE*, Benth. & Hook. f. iii. 990.
ANEPSIAS, Schott, Gen. Aroid. t. 73 (1858).
ATIMETA, Schott, l. c. t. 71 (1858).
blanda, *Schott, in Journ. Bot.* ii. (1864) 53.—Bras.
heliconiifolia, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 317.—Bras.
latifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 91. t. 300.—Peruv.
Lechleriana, Schott, Prod. Aroid. 351 = latifolia.
Melinoni, Engl. Bot. Jahrb. i. (1881) 483 = blanda.

RHODOSPATHA:—
Moritziana, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 109.—Venezuela.
oblongata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 91.—Bras.; Peruv.
Poeppigii, Schott, Gen. Aroid. 72 = latifolia.
surinamensis, Schott, in Bonplandia, ix. (1861) 368 = oblongata.
Wendlandii, *Schott, in Journ. Bot.* ii. (1864) 52.—Am. centr.

RHODOSPHAERA, Engl. Bot. Jahrb. i. (1881) 423. *ANACARDIACEAE.*
rhodanthema, *Engl. l. c.* 424.—Austral.

RHODOSTACHYS, Phil. in Linnaea, xxix. (1857–58) 57. *BROMELIACEAE*, Benth. & Hook. f. iii. 662.
RUCKIA, Regel, Gartenfl. (1868) 69. t. 571.
andina, *Phil. in Linnaea*, xxix. (1857–58) 58.—Chili.
bicolor, *Benth. & Hook. f. Gen.* iii. 662.—Chili.
grandiflora, *Phil. in Linnaea*, xxx. (1859–60) 202.—Chili.
Joinvillei, Benth. & Hook. f. Gen. iii. 662 = pitcairniaefolia.
litoralis, *Phil. in Linnaea*, xxx. (1859–60) 201.—Chili.
pitcairniaefolia, *Benth. & Hook. f. Gen.* iii. 662.—Chili.

RHODOSTOMA, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 286 = **Palicourea**, Aubl. (Rubiac.).
gardenioides, Scheidw. l. c. = P. gardenioides.

RHODOTHAMNUS, Lindl. & Paxt. Flow. Gard. i. (1850–51) t. 22 = **Rhododendron**, Linn. (Ericac.).
kamtchaticus, Lindl. & Paxt. l. c. = Rhodod. kamtchaticum.

RHODOTHAMNUS, Reichb. in Moessl. Handb. 688 (1827). *ERICACEAE*, Benth. & Hook. f. ii. 596.
Chamaecistus, *Reichb. l. c.*—Europ.

RHODOTYPOS, Sieb. & Zucc. Fl. Jap. 187. t. 99 (1835). *ROSACEAE*, Benth. & Hook. f. i. 613.
kerrioides, *Sieb. & Zucc. l. c.*—Japon.

RHODOXYLON, Rafin. Fl. Tellur. iv. 79 (1836) = **Convolvulus**, Tourn.
floridum, Rafin. l. c. 80 = C. floridus.
scoparium, Rafin. l. c. = C. scoparius.

RHOEA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = **Punica**, Tourn. (Lythrar.).
Punica, St. Lag. l. c. = P. Granatum.

RHOEO, Hance, in Walp. Ann. iii. 659 (1853). *COMMELINACEAE*, Benth. & Hook. f. iii. 855.
discolor, *Hance, l. c.* 660.—Mexic.

RHOIACARPOS, A. DC. in DC. Prod. xiv. 634 (1857) = **Colpoon**, Berg. (Santal.).
capensis, A. DC. l. c. 635.—Afr. austr.

RHOICARPOS, Ind. Kew. supra, i. 586, in syn., sphalm. = praec.

RHOMBIFOLIUM, Rich. ex DC. Prod. ii. 235 (1825) = **Clitoria**, Linn. (Legumin).
canescens, Rich. ex DC. l. c. 236 = C. guianensis.

RHOMBODA, Lindl. in Journ. Linn. Soc. i. (1857) 181 = **Hetaeria**, Blume (Orchid.).
lamellata, Miq. Fl. Ind. Bat. iii. 747 = H. lamellata.
longifolia, Lindl. in Journ. Linn. Soc. i. (1857) 181 = H. longifolia.

RHOMBOLYTHRUM, Link, Hort. Berol. ii. 96 (1833) = **Triodia**, R. Br. (Gramin.).
Berteroanum, E. Desv. in C. Gay, Fl. Chil. vi. 387.—Chili.
quinquenervium, Phil. in Linnaea, xxxiii. (1864–65) 293.—Chili.
rhomboideum, Link, Hort. Berol. ii. 296 = T. filiformis.
valdivianum, Phil. in Linnaea, xxxiii. (1864–65) 294 = Poa valdiviana.

RHOMBOSPORA, Korth. in Nederl. Kruidk. Arch. ii. (1850) 114 = **Greenia**, Wight & Arn. (Rubiac.).
Commersonii, Korth. l. c.—Sumatra.
sumatrana, Miq. Fl. Ind. Bat. ii. 345 = Wendlandia glabrata.

RHOPALA, Schreb. Gen. i. 62 (1789) = **Roupala**, Aubl. (Proteac.).

RHOPALEPHORA, Hassk. in Bot. Zeit. xxii. (1864) 58 = **Aneilema**, R. Br. (Commel.).
Blumei, Hassk. l. c. 59 = A. monadelphum.

RHOPALOBLASTE, Scheff. in Ann. Jard. Buitenz. i. (1876) 137. *PALMAE*, Benth. & Hook. f. iii. 891.
hexandra, *Scheff. l. c.* 156. t. 26–27 f. 1.—Ins. Molucc.
micrantha, *Benth. & Hook. f. Gen.* iii. 892.—N. Guin.
Princeps, *Hort. Bull.—Cf. Gard. Chron.* (1880) i. 759.—Hab.?
singaporensis, Benth. & Hook. f. Gen. iii. 892 = Ptychoraphis singaporensis.

RHOPALOCARPUS, Teijsm. & Binn. ex Miq. Ann. Mus. Bat. Lugd. Bat. ii. 22 (1865) = **Anaxagorea**, A. St. Hil. (Anonac.).
fruticosus, Teijsm. & Binn. ex Miq. l. c. = A. fruticosa.

RHOPALOCNEMIS, Jungh. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 213. *BALANOPHOREAE*, Benth. & Hook. f. iii. 238.
PHAEOCORDYLIS, Griff. in Trans. Linn. Soc. xx. (1846) 100. t. 8.
phalloides, *Jungh. in Nov. Act. Nat. Cur.* xviii. Suppl. i. (1841) 215.—Reg. Himal.; Java.

RHOPALOSTIGMA, Phil. Fl. Atac. 42. t. 6 (1860) = **Phrodus**, Miers (Solanac.).
microphyllum, Phil. l. c.—Chili.
pendulum, Phil. l. c.—Chili.
thymifolia, Phil. in Linnaea, xxix. (1857–58) 24 = P. Bridgesii.

RHOPALOSTIGMA, Schott, in Oestr. Bot. Zeit. ix. (1859) 39 = **Staurostigma**, Scheidw. (Aroid.).
Riedelianum, Schott, l. c. = S. Riedelianum.

RHOPALOSTYLIS, Klotzsch, ex Baill. Adansonia, v. (1864–65) 317 = **Dalechampia**, Linn. (Euphorb.).
buettnerioides, Klotzsch, l. c. = D. micrantha.

RHOPALOSTYLIS, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 180. *PALMAE*, Benth. & Hook. f. iii. 890.
Baueri, *H. Wendl. & Drude, in Bot. Zeit.* xxxv. (1877) 638.—Ins. Norfolk et Chatham.
sapida, *H. Wendl. & Drude, in Kerch. Palm.* 255.—N. Zel.

RHOPIUM, Schreb. Gen. 608 (1791) = **Phyllanthus**, Linn. (Euphorb.).
citrifolium, Willd. Sp. Pl. iv. 150 = P. attenuatus.

RHOPHOSTEMON, Wittst. Etymol. Handw. 763 (1856) = ROTROSTEMON, Blume = **Pogonia**, Juss (Orch.).

RHORADENDRON, Griseb. in Lechl. Berb. Am. Austr. 59 (1857), sphalm. = **Phoradendron**, Nutt. (Loranthac.).

RHUACOPHILA, Blume, Enum. Pl. Jav. 13 (1827) = **Dianella**, Lam. (Liliac.).
celebica, Blume, l. c. 14 = D. celebica.
javanica, Blume, l. c. = D. nemorosa.

RHUS, [Tourn.] Linn. Gen. ed. I. 84 (1737). *ANACARDIACEAE*, Benth. & Hook. f. i. 418.
COTINUS, [Tourn.] Linn. Syst. ed. I (1735).
LOBADIUM, Rafin. in Am. Monthly Mag. (1819) 357; et in Journ. Phys. lxxxix. (1819) 98.
MALOSMA, Engl. in DC. Monog. Phan. iv. 393 (1883).
METOPIUM, P. Br. Hist. Jamaic. 177 (1756).
PHILOSTEMON, Rafin. Fl. Ludov. 107 (1817).
POCOPHORUM, Neck. Elem. ii. 226 (1790).
SCHMALZIA, Desv. ex DC. Prod. ii. 72 (1825).
STYPHONIA, Nutt. in Torr. & Gray, Fl. N. Am. i. 220 (1838).
TOXICODENDRON, [Tourn.] Linn. Syst. ed. I (1735).
TURPINIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
VERNIX, Adans. Fam. ii. 342 (1763).

RHUS:—
abyssinica, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 31, nomen; *& ex Oliver, Fl. Trop. Afr.* i. 438.—Abyss.
acuminata, DC. Prod. ii. 68 = succedanea.
acuminata, E. Mey. in Drège, Zwei Pfl. Docum. 216 = laevigata.
acutidens, *Engl. in DC. Monog. Phan.* iv. 423.—Afr. austr.
acutiloba, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 612 = R. Toxicodendron.
aequalis, *Pers. Syn.* i. 326.—Hab.?
affinis, Wall. Cat. n. 995 = semialata.
africana, *Mill. Gard. Dict.* ed. VIII. n 11.—Afr. austr.
aglaophylla, Eckl. & Zeyh Enum. 148 = undulata.
ailanthoides, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 89 = Picrasma quassioides.
alata, Thunb. Prod. Pl. Cap. 52 = Hippobromus alatus.
albicans, Willd. Enum. Hort. Berol. 325 = albida.
albida, *Schousb. Vextr. Marokko*, 142.—Afr. bor.; Arab.; Syria.
albomarginata, *Sond. in Harv. & Sond. Fl. Cap.* i. 519.—Afr. austr.
amazonica, Poepp. ex Engl. in DC. Monog. Phan. iv. 278 = Tapirira guianensis.
Amela, D. Don, Prod. Fl. Nep. 248 = semialata.
amoena, Salisb. Prod. 169 = R. Coriaria.
ampla, *Engl. in DC. Monog. Phan.* iv. 429.—Afr. trop.
Andrieuxii, *Engl. l. c.* 389.—Mexic.
angolensis, *Engl. l. c.* 448.—Afr. trop.
angustifolia, *Linn. Sp. Pl.* 267.—Afr. austr.
angustifolia, E. Mey. ex Harv. & Sond. Fl. Cap. i. 512 = puberula.
antillana, *Egg. Fl. St. Croix*, 41.—Ins. S. Crucis.
arborea, DC. Prod. ii. 73 = Schmidelia Cominia.
arborea, Macfad. ex Griseb. Fl. Brit. W. Ind. 177 = Picrodendron arboreum.
arborescens, *DC. Prod.* ii. 73.—N. Granat.
arenaria, Wierzb. ex Engl. in DC. Monog. Phan. iv. 351 = R. Cotinus.
argentea, Eckl. & Zeyh. Enum. 149 = dissecta.
argentea, Mill. Gard. Dict. ed. VIII. n. 12 = angustifolia.
argyrophylla, Presl, Bot. Bemerk. 42 = Anaphrenium argenteum.
aromatica, *Ait. Hort. Kew.* ed. I. i. 367.—Am. bor.
atomaria, Jacq. Hort. Schoenb. iii. 51. t. 343 = villosa.
atra, Forst. f. Prod. 23 = Oncocarpus vitiensis.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 5.—Arab.
bahamensis, G. Don, Gen. Syst. ii. 72 (Quid?).—Ins. Bahama.
bicolor, Licht. ex Schult. Syst. vi. 661 = tomentosa.
Bolusii, *Sond. ex Engl. in DC. Monog. Phan.* iv. 436.—Afr. austr.
brasiliensis, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 198.—Bras.
Bucki-Amela, Roxb. Fl. Ind. ii. 99 = semialata.
Bucki-Amelam, Roxb. Hort. Beng. 22 = praec.
Burchelli, *Sond. ex Engl. in DC. Monog. Phan.* iv. 412.—Afr. austr.
Burkeana, *Sond. in Harv. & Sond. Fl. Cap.* i. 514.—Afr. austr.
Burmanni, DC. Prod. ii. 69 = mucronata.
Cacodendron, Ehrh. in Hann. Mag. (1783) 227; et ej. Beitr. iii. 20 = Ailanthus glandulosa.
calycina, Steud. Nom. ed. II. ii. 451 = Loxostylis alata.
canadensis, Marsh. Arbust. 129 = aromatica.
canadensis, Mill. Gard. Dict. ed. VIII. n. 5 = typhina.
carolinense, Marsh. Arbust. 129 = glabra?
carolinina, Mill. Gard. Dict. ed. VIII. n. 4 = glabra.
caustica, Hook. & Arn. Bot. Beech. Voy. 15 = Lithraea venenosa.
Cavanillesii, DC. Prod. ii. 69 = lucida.
celastroides, *Sond. in Harv. & Sond. Fl. Cap.* i. 519.—Afr. austr.
chinensis, Mill. Gard. Dict. ed. VIII. n. 7 = semialata.
ciliata, *Licht. ex Schult. Syst.* vi. 661.—Afr. austr.
ciliolata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 470.—Mexic.
cirrhiflora, Linn. f. Suppl. 184 (= *Cissus pauciflora*).—Afr. austr.
Clausseniana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 469 = Lithraea Aroeirinha.
Cobbe, Linn. Sp. Pl. 267 = Allophylus Cobbe.
Cominia, Linn. Syst. ed. X. 964 = Schmidelia Cominia.

RHUS:—
Commersonii, *Poir. Encyc. Suppl.* v. 264.—Bras.
concinna, *Burch. Trav.* i. 360.—Afr. austr.
concolor, Presl, Bot. Bemerk. 42 = Anaphrenium concolor.
copallina, *Linn. Sp. Pl.* 266.—Am. bor.
corallina, J. F. Gmel. Syst. 1615 (Index), sphalm. = praec.
coriacea, *Engl. in DC. Monog. Phan.* iv. 418.—Afr. austr.
Coriaria, *Linn. Sp. Pl.* 265.—Reg. Mediterr.; Persia.
cotina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Cotinus.
cotinoides, *Nutt. ex Torr. & Gray. Fl. N. Am.* i. 217, in syn.; *et ex Chapm. Fl. S U. St.* 70.—Am. bor.
Cotinus, *Linn. Sp. Pl.* 267.—Reg. Medit.; Oriens; Himal.; China.
Cotinus, Torr. & Gray, Fl. N. Am. i. 216 = cotinoides.
crassinervia, Presl, Bot. Bemerk. 42 = laevigata.
crataegiformis, Pers. Syn. i. 326 = albida.
crenata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 28.—Afr. austr.
crenatifolia, *Schlecht. in Linnaea*, xvi. (1842) 483.—Mexic.
crenulata, A. Rich. Tent. Fl. Abyss. i. 142 = glaucescens.
crispa, Harv. ex Engl. in DC. Monog. Phan. iv. 443 = Gueinzii.
cuneifolia, *Linn. f. Suppl.* 183.—Afr. austr.
cuneifolia, E. Mey. ex Harv. & Sond. Fl. Cap. i. 513 = dentata.
decipiens, E. Mey. in Drège, Zwei Pfl. Docum. 216 = Schmidelia decipiens.
decipiens, Wight & Arn. Prod. 172 = Filicium decipiens.
dentata, *Thunb. Prod. Pl. Cap.* 52.—Afr. austr.
denudata, Eckl. & Zeyh. Enum. 143 = viminalis.
denudata, Licht. ex Schult. Syst. vi. 661 = lancea.
Dhuna, *Buch.-Ham. in Wall. Cat.* n. 8502 (sp. dub.).—Nepal.
digitata, Linn. f. Suppl. 184 (= *Cissus Thunbergii*, Eckl. & Zeyh.).—Afr. austr.
dimidiata, Thunb. in Hoffm. Phytog. Blaetter, i. (1803) 28 (= *Cissus dimidiata*).—Afr. austr.
dioica, Brouss. ex Willd. Enum. Hort. Berol. 325 = albida.
discolor, *E. Mey. in Drège, Zwei Pfl. Docum.* 216, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 507.—Afr. austr.
discolor, Schrad. ex Engl. in DC. Monog. Phan. iv. 408 = tomentosa.
dispar, Presl, Bot. Bemerk. 42 = Anaphrenium dispar.
dissecta *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 29.—Afr. austr.
divaricata, *Eckl. & Zeyh. Enum.* 146.—Afr. austr.
diversifolia, Engl. in DC. Monog. Phan. iv. 395, sphalm. = seq.
diversiloba, Torr. & Gray, Fl. N. Am. i. 218 = R. Toxicodendron.
Dregeana, *Sond. in Harv. & Sond. Fl. Cap.* i. 516.—Afr. austr.
Eckloniana, Presl, Bot. Bemerk. 41 = mucronata.
Eckloniana, *Sond. in Harv. & Sond. Fl. Cap.* i. 515.—Afr. austr.
Ecklonis, Schrad. ex Harv. & Sond. l. c. 509 = tomentosa.
elegans, Ait. Hort. Kew. ed. I. i. 366 = glabra.
elliptica, Thunb. Fl. Cap. ii. 214 = tomentosa.
elongata, Eckl. & Zeyh. Enum. 144 = viminalis.
elongata, Jacq. Hort. Schoenb. iii. 52 = mucronata.
erosa, Drège, ex Presl, Bot. Bemerk. 41 = Schmidelia erosa.
erosa, *Thunb. Fl. Cap.* ii. 212.—Afr. austr.
excisa, E. Mey. ex Harv. & Sond. Fl. Cap. i. 518 = undulata.
excisa, *Thunb. Prod. Pl. Cap.* 187.—Afr. austr.
fastigiata, Eckl. & Zeyh. Enum. 146 = puberula.
fastuosa, Salisb. Prod. 170 = semialata.
ferruginea, Teijsm. & Binn. ex Engl. in DC. Monog. Phan. iv. 450 = rufa.
filicina, [Moç. & Sesse,] ex DC. Prod. ii. 67 = Bursera bipinnata.
foliosa, *A. Rich. Tent. Fl. Abyss.* i. 143.—Abyss.
fragrans, Licht. ex Schult. Syst. vi. 661 = viminalis.
fraxinifolia, D. Don, Prod. Fl. Nep. 248 = Evodia fraxinifolia.
fraxinifolia, Salisb. Prod. 169 = succedanea.
Gerrardi, Harv. ex Engl. in DC. Monog. Phan. iv. 442 = viminalis.

RHUS:—
glabra, *Linn. Sp. Pl.* 265.—Am. bor.
glauca, Hochst. ex Oliver, Fl. Trop. Afr. i. 438 = glaucescens?
glauca, *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 27; *Pers. Syn.* i. 326; *Desf. Hist. Arb.* ii. 326.—Afr. austr.
glaucescens, *A. Rich. Tent. Fl. Abyss.* i. 143.—Abyss.
glaucovirens, Engl. in DC. Monog. Phan. iv. 432 = Zeyheri.
glutinosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 144.—Abyss.
gracillima, *Engl. in DC. Monog. Phan.* iv. 445.—Afr. austr.
gracilis, Hort. ex Engl. l. c. iv. 377 = typhina.
grandidens, *Harv. ex Engl. l. c.* 440.—Afr. austr.
grandidentata, DC. Prod. ii. 72 = dentata.
grandifolia, *Engl. in DC. Monog. Phan.* iv. 434.—Afr. austr.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 12.—Reg. Himal.
Gueinzii, *Sond. in Harv. & Sond. Fl. Cap.* i. 515.—Afr. austr.
heterophylla, *DC. Cat. Hort. Monsp.* (1813) 136.—Hab.?
Hindsiana, *Engl. in DC. Monog. Phan.* iv. 388.—Calif.
hirsuta, Salisb. Prod. 170 = villosa.
hirta, Harv. ex Engl. in DC. Monog. Phan. iv. 425 = tridentata.
horrida, *Eckl. & Zeyh. Enum.* 151.—Afr. austr.
humilis, Eckl. & Zeyh. l. c. 146 = puberula.
humilis, Salisb. Prod. 170 = R. Toxicodendron.
hypoleuca, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 43.—China.
Hypselodendrum, Moench, Meth. 73 = typhina.
incana, *Engl. in DC. Monog. Phan.* iv. 428.—Afr. austr.
incana, Mill. Gard. Dict. ed. VIII. n. 8 = villosa.
incisa, *Linn. f. Suppl.* 183.—Afr. austr.
indica, Herb. Madr. ex Wight & Arn. Prod. 172 = mysurensis.
insignis, *Hook. f. Fl. Brit. Ind.* ii. 11.—Reg. Himal.
insignis, Oliver, Fl. Trop. Afr. i. 437 = Anaphrenium abyssinicum.
integerrima, Wall. Cat. n. 8474 = Pistacia Khinjuk.
integrifolia, *Benth. & Hook. f. ex S. Wats. in Wheeler, Rep. Bot.* 84.—Calif.
integrifolia, Engl. in DC. Monog. Phan. iv. 388, in syn. sphalm. = Hindsiana.
japonica, Buch.-Ham ex D. Don, Prod. Fl. Nep. 248 = seq.
javanica, Linn. Sp. Pl. 265 = semialata.
juglandifolia, Wall. ex D. Don, Prod. Fl. Nep. 248; Cat. n. 996 = Wallichii.
juglandifolia, *Willd. ex Schult. Syst.* vi. 649; *H. B. & K. Nov. Gen. et Sp.* vii. 8. t. 603.—Am. trop.
Kaempferi, Sweet, Hort. Brit. ed. I. 97 = vernicifera.
Kakrasingee, Royle, Illustr. Bot. Himal. 175 = Pistacia Khinjuk.
khasiana, *Hook. f. Fl. Brit. Ind.* ii. 10.—Reg. Himal.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 439.—Afr. trop.
Kotschyana, Fenzl, in Flora, xxvii. (1844) 312 = glaucescens.
Krebsiana, *Presl, Bot. Bemerk.* 41.—Afr. austr.
laevicaulis, Torr. ex A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 28 = glabra.
laevigata, *Linn. Sp. Pl.* ed. II. 1672.—Afr. austr.
laevis, Wall. ex G. Don, Gen. Syst. ii. 69 = R. Cotinus.
lancea, *Linn. f. Suppl.* 184.—Afr. austr.
lancea, E. Mey. ex Harv. & Sond. Fl. Cap. i. 516 = Dregeana.
lanceolata, A. Gray, ex Engl. in DC. Monog. Phan. iv. 384 = copallina.
laurina, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 219 = Calif.
lavandulaefolia, Presl, Bot. Bemerk. 42 = stenophylla.
Lentii, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 16.—Calif.
lentiscifolia, Stokes, Bot. Mat. Med. ii. 164 = copallina.
leucantha, Jacq. Hort. Schoenb. iii. 50 = copallina.
leucocarpa, E. Mey. in Drège, Zwei Pfl. Docum. 216 = Schmidelia leucocarpa.
Lindeniana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 468 = juglandifolia.
lineata, Spreng. Syst. i. 936 = lineatifolia.

RHUS:—
lineatifolia, *Orteg. Hort. Matr. Dec.* 89.—Cuba.
livida, Salisb. Prod. 170 = laevigata.
lobata, Hook. Fl. Bor. Am. i. 127 = R. Toxicodendron.
lobata, *Poir. Encyc. Suppl.* v. 264.—Ins. Teneriff.
longifolia, Sond. in Harv. & Sond. Fl. Cap. i. 522 = Protorhus longifolia.
longipes, *Engl. in DC. Monog. Phan.* iv. 431.—Afr. trop.
longispina, *Eckl. & Zeyh. Enum.* 148.—Afr. austr.
lucida, Ait. Hort. Kew. ed. II. ii. 166 = mucronata.
lucida, *Linn. Sp. Pl.* 267.—Afr. austr.
lucida, E. Mey. ex Engl. in DC. Monog. Phan. iv. 412 = glauca.
macrocarpa, *Engl. l. c.* 449.—Afr. austr.
macrophylla, *Hook. & Arn. Bot. Beech. Voy.* 413.—Mexic.
macrophylla, A. Rich. ex Engl. in DC. Monog. Phan. iv. 426 = abyssinica.
magalismontana, *Sond. in Harv. & Sond. Fl. Cap.* i. 510.—Afr. austr.
marginata, *E. Mey. in Drège, Zwei Pfl. Docum.* 216, nomen.—Afr. austr.
Meerburgiana, *Schult. Syst.* vi. 663.—Hab.?
melanocarpa, E. Mey. in Drège, Zwei Pfl. Docum. 216 = Schmidelia melanocarpa.
melintangensis, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 219 = Canarium glaucum.
meridionalis, Spreng. Syst. i. 936 = Commersonii.
metopia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Metopium.
metopioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 468.—Cuba.
Metopium, *Linn. Syst.* ed. X. 964.—Am. bor.; Ind. occ.
Meyeriana, Presl, Bot. Bemerk. 41 = puberula.
micrantha, Eckl. & Zeyh. Enum. 149 = undulata.
micrantha, Thunb. Fl. Cap. ii. 221 = dentata.
microcarpa, Steud. Nom. ed. I. 689 = R. Toxicodendron.
microlepis, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 468 = hypoleuca.
microphylla, *Engelm. in A. Gray, Pl. Wright.* i. 31.—Texas.
mollis, *H. B. & K. Nov. Gen. et Sp.* vii. 10. t. 602.—Mexic.
mollis, *Jacq. Fragm.* 74.—Afr. austr.
mollis, E. Mey. in Drège, Zwei Pfl. Docum. 216 = parvifolia.
monophylla, E. Mey. l. c. = Schmidelia monophylla.
mucronata, Eckl. & Zeyh. Enum. 150 = africana.
mucronata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 519 = excisa.
mucronata, *Thunb. in Hoffm. Phytog. Blaetter*, i. (1803) 27.—Afr. austr.
mucronifolia, Sond. in Harv. & Sond. Fl. Cap. i. 521 = Anaphrenium abyssinicum.
mysurensis, *Heyne, ex Wight & Arn. Prod.* 172.—Ind. or.
natalensis, *Bernh. ex Krauss, in Flora*, xxvii. (1844) 349.—Afr. austr.
nepalensis, Hoffmgg. Verz. Pfl. Nachtr. ii. 35 = pruinosa.
nervosa, Eckl. & Zeyh. Enum. 147 = excisa.
nervosa, Poir. Encyc. Suppl. v. 264 = undulata.
nitida, *Engl. in DC. Monog. Phan.* iv. 434.—Afr. trop.
nodosa, *Blume, Bijdr.* 1164.—Java.
obliqua, E. Mey. ex Harv. & Sond. Fl. Cap. i. 444 = Clausena inaequalis.
obliqua, *Thunb. Fl. Cap.* ii. 224.—Afr. austr.
oblongifolia, E. Mey. in Drège, Zwei Pfl. Docum. 216 = Sapindus oblongifolius.
obovata, *Sond. in Harv. & Sond. Fl. Cap.* i. 508.—Afr. austr.
obovatifolia, Stokes, Bot. Mat. Med. ii. 159 = R. Cotinus.
obscura, Bieb. Fl. Taur. Cauc. i. 243 = Pterocarya caucasica.
Odina, Buch.-Ham. ex Wall. Cat. n. 8475 = Odina Wodier.
odorata, Nois. ex Steud. Nom. ed. I. 689, nomen (Quid?).—Hab.?
ornifolia, Pall. in Gueldenst. Reisen, i. 282 = R. Coriaria.
Osbeckii, *Carr. in Rev. Hortic.* (1887) 111.—China.
Osbeckii, Steud. Nom. ed. II. ii. 452 = semialata.

RHUS:—
ovata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 358.—Calif.
Oxyacantha, Schousb. ex Cav. in Anal. Cienc. Nat. iii. (1801) 36 = albida.
Oxyacanthoides, Dum.-Cours. Bot. Cult. vi. 662; ed. II. vi. 214 = albida.
Oxymetopium, Griseb. Cat. Pl. Cub. 67 = R. Metopium.
pachyrrhachis, Hemsl. Biol. Centr. Am. Bot. i. 218 = sempervirens.
pallens, Eckl. & Zeyh Enum. 147 = excisa.
pallida, E. Mey. in Drège, Zwei Pfl. Docum. 216 = viminalis.
panaciformis, F. Muell. Fragm. vii. 22 = rufa.
paniculata, *Wall. Cat.* n. 993.—Reg. Himal.; Burma.
paniculosa, Sond. in Harv. & Sond. Fl. Cap. i. 522 = Anaphrenium paniculosum.
parviflora, *Roxb. Hort. Beng.* 22; *Fl. Ind.* ii. 100.—Reg. Himal.
parvifolia, Engl. in DC. Monog. Phan. iv. 419, sphalm. = praec.
parvifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 510.—Afr. austr.
pauciflora, Linn. f. Suppl. 183 = Hippobromus alatus.
pendulina, Eckl. & Zeyh. Enum. 145 = mucronata.
pendulina, Jacq. ex Willd. Enum. Hort. Berol. 324 = viminalis.
pentaphylla, *Desf. Fl. Atlant.* i. 267. t. 77.—Reg. Mediterr.; Afr. trop.
perniciosa, H. B. & K. Nov. Gen. et Sp. vii. 10 = Pseudosmodingium perniciosum.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 144 = glutinosa.
pilipes, Presl, Bot. Bemerk. 41 = mucronata.
pistachiaefolia, Salisb. Prod. 169 = copallina.
platypoda, E. Mey. in Drège, Zwei Pfl. Docum. 216 = horrida.
plicaefolia, Eckl. & Zeyh. Enum. 148 = excisa.
Plukenetiana, *Eckl. & Zeyh. l. c.* 147.—Afr. austr.
polyantha, Benth. Pl. Hartw. 60 = Picramnia polyantha.
populifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 216 nomen; *et ex Harv. & Sond. Fl. Cap.* i. 508.—Afr. austr.
potentillaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 469.—Mexic.
pruinosa, *Hoffmgg. Verz. Pfl. Nachtr.* II. 198.—Ind. or.
pterocarpa, [Moç. & Sesse,] ex DC. Prod. ii. 84 = Spathelia rhoifolia.
pterota, Presl, Bot. Bemerk. 41 = longispina.
puberula, *Eckl. & Zeyh. Enum.* 146.—Afr. austr.
pubescens, Eckl. & Zeyh. l. c. 145 = pyroides.
pubescens, *Thunb. Prod. Pl. Cap.* 52.—Afr. austr.
pubigera, Blume, Bijdr. 1165 = succedanea.
pulcherrima, Oliver, Fl. Trop. Afr. i. 436 = Anaphrenium pulcherrimum.
pumila, Meerb. Pl. Sel. Ic. t. 14 = Meerburgiana.
pumila, *Michx. Fl. Bor. Am.* i. 182.—Am. bor.
punjabensis, *J. L. Stew. ex Brand. For. Fl.* 120.—Reg. Himal.
pyroides, *Burch. Trav.* i. 340.—Afr. austr.
pyroides, A. Rich. Tent. Fl. Abyss. i. 145 = villosa.
Quartiniana, A. Rich. l. c. 141 = glaucescens.
quercifolia, Steud. Nom. ed. I. 689 = R. Toxicodendron.
quinquefolia, Stokes, Bot. Mat. Med. ii. 161 = R. Metopium.
Radaelijawei, Mill. Gard. Dict. ed. VIII. n. 13 = Connarus monocarpus.
radicans, Linn. Sp. Pl. 266 = R. Toxicodendron.
refracta, *Eckl. & Zeyh. Enum.* 145.—Afr. austr.
Rehmanniana, *Engl. in DC. Monog. Phan.* iv. 422.—Afr. austr.
retinorrhaea, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 142.—Abyss.
retusa, Zoll. ex Teijsm. & Binn. Cat. Hort. Bog. 230 = rufa.
rhodanthema, F. Muell. in Journ. Pharm. Soc. Vict. i. (1858) 43; et ex Benth. Fl. Austral. i. 489 = Rhodosphaera rhodanthema.
rigida, *Mill. Gard. Dict.* ed. VIII. n. 14.—Afr. austr.
rimosa, Eckl. & Zeyh. Enum. 150 = rigida.
rosaefolia, *Hoffmgg. Verz. Pfl. Nachtr.* II. 199.—Bras.
rosmarinifolia, Thunb. Fl. Cap. ed. Schult. 262 = stenophylla.

RHUS:—
rosmarinifolia, *Vahl, Symb. Bot.* iii. 50.—Afr. austr.
Roxburghii, Steud. Nom. ed. II. ii. 452 = semialata.
rubifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 470.—Mexic.
rufa, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* xxvii. (1863) 52.—Java.
rufescens, *Desv. in Ham. Prod. Fl. Ind. Occ.* 32 (an Anacard.?).—Jamaic.
rufescens, Eckl. & Zeyh. Enum. 144 = discolor.
Saeneb, Forsk. Fl. Aegypt. Arab. 206 (non Anacard.).—Arabia.
salicifolia, Presl, Bot. Bemerk. 42 = Anaphrenium abyssinicum.
salicina, Sond. in Harv. & Sond. Fl. Cap. i. 522 = Anaphrenium abyssinicum.
Samo, Tul. in Ann. Sc. Nat. Sér. III. vi. (1846) 367 = juglandifolia.
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 369.—Ins. Sandvic.
saxatilis, [Moç. & Sesse,] ex DC. Prod. ii. 71 (non Anacard.).—Mexic.
scandens, Salisb. Prod. 170 = R. Toxicodendron.
Schiedeana, *Schlecht. in Linnaea*, xvi. (1842) 480.—Mexic.
Schimperi, Hochst. ex Engl. in DC. Monog. Phan. iv. 441 = glaucescens.
schinoides, Willd. ex Schult. Syst. vi. 649 = Schinus terebinthifolius?
schmideloides, Schlecht. in Linnaea, xvi. (1842) 482 = aromatica.
scoparia, *Eckl. & Zeyh. Enum.* 148.—Afr. austr.
scytophylla, *Eckl. & Zeyh. l. c.* 150.—Afr. austr.
semialata, *Murr. in Comm. Gotting.* vi. (1784) 27. t. 3.—Reg. Himal.; China; Ins. Sandvic.
sempervirens, *Scheele, in Linnaea*, xxiii. (1850) 566.—Texas.
sericea, Eckl. & Zeyh. Enum. 146 = puberula.
serraefolia, Burch. Trav. ii. 100 = erosa.
simarubaefolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 367. t. 44.—Ins. Fiji.
Simonii, *Carr. in Rev. Hortic.* (1866) 339.—China.
simplicifolia, Salisb. Prod. 170 = R. Cotinus.
sinuata, Eckl. & Zeyh. Enum. 147 = obovata.
sinuata, *Thunb. Prod. Pl. Cap.* 52.—Afr. austr.
Sonderi, *Engl. in DC. Monog. Phan.* iv. 435.—Afr. austr.
spathulata, Eckl. & Zeyh. Enum. 148 = undulata.
spicata, Thunb. Fl. Cap. ii. 217 = Schmidelia decipiens.
stenophylla, *Eckl. & Zeyh. Enum.* 144.—Afr. austr.
striata, Ruiz & Pav. Fl. Per. iii. 29. t. 252 = juglandifolia.
suaveolens, Ait. Hort. Kew. ed. I. i. 368 = aromatica.
subcordata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 470 = rubifolia.
subferruginata, Presl, Bot. Bemerk. 41 = divaricata.
succedanea, *Linn. Mant.* ii. 221.—Reg. Himal; China; Japon.
Sumac, Targ. Tozz. Osserv. Bot. Dec. vi. 3 = R. Coriaria.
sylvestris, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 140.—Japon.
syriaca, Boiss. & Bal. ex Boiss. Fl. Orient. ii. 5 = albida.
taitensis, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 361 = simarubaefolia.
tenuiflora, Presl, Bot. Bemerk. 41 = mucronata.
tenuinervis, *Engl. in DC. Monog. Phan.* iv. 423.—Afr. trop.
terebinthifolia, *Schlecht. & Cham. in Linnaea*, v. (1830) 600.—Mexic.
theesans, Hort. ex G. Don, in Loud. Hort. Brit. 110 = undulata.
Theezera, Pers. Syn. i. 325 = pentaphylla.
Thunbergiana, Schult. Syst. vi. 651 = glauca.
Thunbergii, Hook. Ic. Pl. t. 595 = Anaphrenium argenteum.
thyrsiflora, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 567.—Ins. Socotra.
tomentosa, *Linn. Sp. Pl.* 266.—Afr. austr.
toxicarium, Salisb. Prod. 170 = R. Toxicodendron.
toxicodendra, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.
Toxicodendron, *Linn. Sp. Pl.* 266.—Am. bor.
transvaalensis, *Engl. in DC. Monog. Phan.* iv. 440.—Afr. austr.
triceps, E. Mey. in Drège, Zwei Pfl. Docum. 216 = rigida.

RHUS:—
trichocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 84.—Japon.
tridactyla, *Burch. Trav.* i. 340.—Afr. austr.
tridactyla, Eckl. & Zeyh. Enum. 144 = Eckloniana.
tridentata, Linn. f. Suppl. 184 (= *Cissus pauciflora*).—Afr. austr.
tridentata, *Sond. in Harv. & Sond. Fl. Cap.* i. 511.—Afr. austr.
tridentata, Willd. ex Engl. in DC. Monog. Phan. iv. 438 = pentaphylla.
trifoliata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = Evodia glabra.
trifoliata, Willd. Enum. Hort. Berol. Suppl. 15 = laevigata.
trijuga, Poir. Encyc. vii. 506 = Schinus terebinthifolius?
trilobata, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 219.—Am. bor. occ.; Mexic.
tripartita, DC. Prod. ii. 72, in syn. = albida.
triphylla, *Medic. Pfl. Anat.* 106, nomen.—Hab.?
typhia, Crantz, Inst. ii. 275 = seq.
typhina, *Linn. Cent. Pl.* ii. 14.—Am. bor.
undulata, *Jacq. Hort. Schoenb.* iii. 52. t. 346.—Afr. austr.
undulata, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 239 = Schmidelia decipiens.
undulata, A. Rich. Tent. Fl. Abyss. i. 145 = glaucescens.
varielobata, Steud. Nom. ed. II. ii. 452 = R. Toxicodendron.
variifolia, [Moç. & Sesse,] ex DC. Prod. ii. 73 = heterophylla?
Veatchiana, Kellogg, in Proc. Calif. Acad. ii. (1863) 24 = Veatchia cedrosensis.
Veitchiana, Engl. in DC. Monog. Phan. iv. 452, sphalm. = praec.
velutina, Wall. Cat. n. 998; et ex G. Don, Gen. Syst. ii. 69 = R. Cotinus.
venenata, *DC. Prod.* ii. 68.—Am. bor.
venosa, *Griseb. Cat. Pl. Cub.* 67.—Cuba.
vernicifera, *DC. Prod.* ii. 68, partim.—Tibet. or.; Japon.
vernicifera, DC. l. c., partim = Wallichii.
vernicifera, Salisb. Prod. 169 = venenata.
verniciflua, Stokes, Bot. Mat. Med. ii. 164 = vernicifera.
Vernix, Linn. Sp. Pl. 265, partim = venenata, vernicifera.
Vernix, Thunb. Fl. Jap. 121 = vernicifera.
verrucosa, Scheele, in Linnaea, xxi. (1848) 592 = R. Toxicodendron.
villosa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 145 = pyroides.
villosa, *Linn. f. Suppl.* 183.—Afr. trop. et austr.
villosa, E. Mey. ex Engl. in DC. Monog. Phan. iv. 427 = refracta.
villosissima, *Engl. l. c.* 447.—Afr. austr.
viminalis, *Ait. Hort. Kew.* ed. I. i. 368; *Vahl, Symb. Bot.* iii. 50.—Afr. austr.
viminalis, Jacq. Hort. Schoenb. iii. 51. t. 344 = lancea.
viminalis, A. Rich. Tent. Fl. Abyss. i. 142 = retinorrhaea.
virens, Lindheim. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 159 = sempervirens.
viridiflora, Duham. Arb. ed. Nov. ii. 163 = typhina.
viridiflora, Hort. ex Engl. in DC. Monog. Phan. iv. 376 = glabra.
viticifolia, *F. Muell. ex Benth. Fl. Austral.* i. 489.—Austral.
Wallichii, *Hook. f. Fl. Brit. Ind.* ii. 11.—Reg. Himal.
Wallichii, Sweet, Hort. Brit. ed. II. 118 = an praec.?
Welwitschii, *Engl. in DC. Monog. Phan.* iv. 428.—Afr. trop.
Wildingii, Dehnh. Revis. Napol. i. III. 172; ex Walp. Rep. i. 551 = viminalis.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 514.—Afr. austr.
ziziphina, Tineo, Pl. Rar. Sic. Pugill. i. 8 = albida.

RHYACOPHILA, Hochst. in Flora, xxiv. (1841) 659 = **Quartinia**, Endl. (Lythrar.).
repens, Hochst. l. c. = Q. repens.

RHYDITOSPERMUM, Walp. Rep. vi. 196 (1846–47), sphalm. = RHYTIDOSPERMUM, Sch. Bip. = **Matricaria**, Tourn. (Compos.).

RHYNCHADENIA, A. Rich. Fl. Cub. Fanerog. ii. 248. t. 85 (1853) = **Macradenia**, R. Br. (Orchid.).
cubensis, A. Rich. l. c. = M. lutescens.

RHYNCHANDRA, Reichb. Nom. 56 (1841) = seq.

RHYNCHANTHERA, Blume, Bijdr. Tabell. 78 (1826) = **Corymbis**, Thou. (Orchid.).
paniculata, Blume, l. c.—Java.

RHYNCHANTHERA, DC. Prod. iii. 106 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 738.
PROBOSCIDIA, Rich. ex DC. Prod. iii. 106 (1828).
THENARDIA, [Moç. & Sesse,] ex DC. l. c. 108 (1828).
acuminata, *Benth. in Hook. Journ. Bot.* ii. (1840) 299.—Guian.
adenophora, Miq. Comm. Phyt. 73. t. 10 = Acisanthera recurva.
ambigua, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 210 = grandiflora.
betulifolia, *Cogn. in Mart. Fl. Bras.* xiv. III. 171.—Guian.
brachyrhyncha, *Cham. in Linnaea*, ix. (1834) 379.—Bras.
bracteata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 31.—N. Granat.
collina, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 208.—Bolivia.
cordata, *DC. Prod.* iii. 107.—Bras.
cordifolia, Walp. Rep. ii. 120 = acuminata.
dichotoma, *DC. Prod.* iii. 107.—Bras.
Fothergillae, DC. l. c. 108 = Tibouchina Fothergillae.
Gardneri, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 207.—Bras.
Glazioviana, *Cogn. in Mart. Fl. Bras.* xiv. III. 189.—Bras.
grandiflora, *DC. Prod.* iii. 107.—Guian.; Bras.
Haenkeana, *DC. l. c.*—Peruv.
hispida, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 212.—Bras.
Hookeri, *Naud. l. c.*—Peruv.
insignis, Naud. l. c. 210 = mexicana.
intermedia, Naud. l. c. 211 = hispida.
latifolia, *Cogn. in Mart. Fl. Bras.* xiv. III. 167.—Bras.
laxa, *Cogn. l. c.* 176.—Bras.
limosa, *DC. Prod.* iii. 108.—Bras.
Matthaei, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 209 = Haenkeana.
Maximowiczii, *Cogn. in Mart. Fl. Bras.* xiv. III. 185.—Bras.
mexicana, *DC. Prod.* iii. 108.—Mexic.
modesta, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 212 = serrulata.
monodynama, DC. Prod. iii. 107 = grandiflora.
monodynama, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 209 = cordata.
novemnervia, *DC. Prod.* iii. 107.—Bras.
ovalifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 208.—Bras.
parviflora, *Naud. l. c.* 213.—Guian.
pentanthera, DC. Prod. iii. 108 = Schrankiana.
Regnellii, *Cogn. in Mart. Fl. Bras.* xiv. III. 184.—Bras.
Riedeliana, *Cogn. l. c.*—Bras.
rosea, *Cogn. l. c.* 181.—Paraguay.
rostrata, *DC. Prod.* iii. 108.—Bras.
salicifolia, Mart. ex DC. l. c. 109 = serrulata.
Schrankiana, *DC. l. c.* 107.—Bras.
Schrankiana, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 214, partim = dichotoma.
secundiflora, *Naud. l. c.* 207.—Paraguay.
serrulata, *DC. Prod.* iii. 108.—Am. trop.
simplicicaulis, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849) 208 = ovalifolia.
stachydimorpha, DC. Prod. iii. 108 = rostrata.
stricta, *Cogn. in Mart. Fl. Bras.* xiv. III. 168.—Bras.
ursina, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 207.—Bras.
verbenoides, *Cham. in Linnaea*, ix. (1834) 378.—Bras.
villosissima, *Cogn. in Mart. Fl. Bras.* xiv. III. 187.—Bras.
Weddellii, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 213.—Bras.

RHYNCHARRHENA, F. Muell. Fragm. i. 128 (1859) = **Pentatropis**, R. Br. (Asclep.).
atropurpurea, F. Muell. l. c. = P. atropurpurea.
quinquepartita, F. Muell. l. c. = P. quinquepartita.

RHYNCHELYTHRUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Panicum**, Linn. (Gramin.).
Dregeanum, Nees, l. c. 447 = P. roseum.

RHYNCHELYTRUM, Endl. Gen. 83 (1837) = praec.

RHYNCHELYTRUM, Hochst. in Flora, xxvii. (1844) 249. *GRAMINEAE*, Benth. & Hook. f. iii. 1120.
grandiflorum, Hochst. l. c. (Panici sp.).—Afr. trop.
microstachyum, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 97 = *grandiflorum*.
ruficomum, *Hochst. ex Steud. Syn. Pl. Gram.* 120.—Afr. trop.

RHYNCHIUM, Dulac, Fl. Hautes-Pyr. 290 (1867) = **Vicia**, Tourn. (Legumin.).
plicatum, Dulac, l. c. 291 = V. Ervilia.

RHYNCHOCARPA, Schrad. in Linnaea, xii. (1838) 403 = **Kedrostis**, Medic. (Cucurbit.).
africana, Aschers. in Schweinf. Beitr. Fl. Aethiop. 268 = K. africana.
Bainesii, Hook. f. in Oliver, Fl. Trop. Afr. ii. 564 = K. Bainesii.
corallina, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 180 = Corallocarpus Fenzlii.
Courbonii, Naud. l. c. xviii. (1862) 197 = Corallocarpus velutinus.
deltoidea, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 105 (sp. dub.).—Burma.
dissecta, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 179 = K. africana.
Ehrenbergii, Schweinf. Beitr. Fl. Aethiop. 251 = Corallocarpus Ehrenbergii.
epigaea, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 178 = Corallocarpus epigaeus.
erostris, Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 672 = Corallocarpus erostris.
etbaica, Aschers. ex Cogn. in DC. Monog. Phan. iii. 655 = Corallocarpus velutinus.
foetida, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 627 = K. rostrata.
foetida, Schrad. in Linnaea, xii. (1838) 403 = K. foetidissima.
foetidissima, Walp. Rep. ii. 197 = K. foetidissima.
Gijef, Schweinf. ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 566 = Corallocarpus Gijef.
glomerata, Hort.—Cf. Gard. Chron. (1880) II. 819.—Bras.
hirtella, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 181 = K. hirtella.
odorata, Hance, in Journ. Linn. Soc. xiii. (1873) 104 = Melothria odorata.
pedunculosa, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 182 = Corallocarpus pedunculosus.
rostrata, Naud. l. c. 177 = K. rostrata.
Schimperi, Naud. l. c. 180 = Corallocarpus Schimperi.
Welwitschii, Naud. l. c. xviii. (1862) 198. t. 10 = Corallocarpus Welwitschii.

RHYNCHOCARPUS, Less. Syn. Comp. 382 (1832) = **Relhania**, L'Hérit. (Compos.).
lateriflorus, Less. l. c. 383 = Relh. sessiliflora.

RHYNCHOCARPUS, Reinw. ex Blume, Cat. Gew. Buitenz. 84 (1823) = **Cyrtandra**, Forst. (Gesner.).
coccineus, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 12 = Cyrtandra coccinea.
cordifolius, Reinw. ex Blume, Cat. Gew. Buitenz. 84 = C. cordifolia.
glaber, Reinw. l. c. = C. longe-petiolata.
longirostris, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 12 = C. longirostris.
mollis, Reinw. ex De Vriese, l. c. 16 = C. mollis.
nemorosus, Reinw. ex Blume, Cat. Gew. Buitenz. 84 = C. longe-petiolata.
repens, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 14 = C. repens.
spicatus, Reinw. ex De Vriese, l. c. = C. spicata.
umbellatus, Reinw. ex De Vriese, l. c. 13 = C. cuneata.

RHYNCHOCORYS, Griseb. Spicil. Fl. Rumel. ii. 12 (1844). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 978.
ELEPHANTINA, Bertol. Fl. Ital. vi. 279 (1844).
ELEPHAS, Tourn. ex Adans. Fam. ii. 211 (1763).
PROBOSCIPHORA, Neck. Elem. i. 336 (1790).
RHINANTHUS, Linn. Sp. Pl. 603 (1753), partim.

RHYNCHOCORYS:—
Elephas, *Griseb. Spicil. Fl. Rumel.* ii. 12.—Europ. austr.; Oriens.
maxima, *Richter, in Denkschr. Akad. Wien*, l. (1885) 25.—Persia.
orientalis, *Benth. in DC. Prod.* x. 559.—Reg. Caucas.
strictus, Boiss. Fl. Orient. iv. 478 = R. Elephas.

RHYNCHODIA, Benth. in Benth. & Hook. f. Gen. ii. 719 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
CERCOCOMA, Miq. Fl. Ind. Bat. ii. 445 (1859?).
RHYNCHOSPERMUM, A. DC. in DC. Prod. viii. 431 (1844).
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 667.—Ind. or.

RHYNCHODIUM, Presl (Bot. Bemerk. 54 (1844)), in Abh. Boehm. Ges. V. iii. (1845) 484 = **Psoralea**, Linn. (Legumin.).
acaule, Presl, Bot. Bemerk. 54 = P. acaulis.
bituminosum, Presl, l. c. = P. bituminosa.
foetidum, Presl, l. c. = P. foetida.
hirtum, Presl, l. c. = P. hirta.
palestinum, Presl, l. c. = P. bituminosa.
sericeum, Presl, l. c. = P. tomentosa.

RHYNCHOGLOSSUM, Blume, Bijdr. 741 (1826) (Rhinchoglossum). *GESNERACEAE*, Benth. & Hook. f. ii. 1019.
ANTONIA, R. Br. in Wall. Pl. As. Rar. iii. 65 (1832).
LOXOTIS, R. Br. ex Benth. Scroph. Ind. 57 (1835).
KNAPPIA, Ferd. Bauer, ex Steud. Nom. ed. II. i. 846 (1840); ii. 75 (1841).
RHINOGLOSSUM, Pritz. Ic. Ind. i. 933 (1855).
Blumei, DC. Prod. ix. 274 = obliquum.
klugioides, *C. B. Clarke, in DC. Monog. Phan.* v. 163.—Ins. Philipp.
obliquum, *Blume, Bijdr.* 741.—Ind. or.; Malaya.
obliquum, Hohen. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 366 = Klugia Notoniana.
Rheedii, DC. Prod. ix. 274 = obliquum.
scabrum, Dalz. in Hook. Kew Journ. ii. (1850) 140 = Klugia Notoniana.
zeylanicum, Hook. Bot. Mag. t. 4198 = obliquum.

RHYNCHOLACIS, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 95. *PODOSTOMACEAE*, Benth. & Hook. f. iii. 110.
crassipes, *Spruce, ex Wedd. in DC. Prod.* xvii. 56.—Bras.
Hydrocichorium, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 95.—Guian.
linearis, *Tul. in Mart. Fl. Bras.* iv. 1. 241. t. 74. f. 5.—Bras.
macrocarpa, *Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 95.—Guian.
oligandra, *Wedd. in DC. Prod.* xvii. 57.—Bras.
tenuifolia, *Spruce, ex Wedd. l. c.*—Bras.
varians, *Spruce, ex Wedd. l. c.* 56.—Bras.

RHYNCHOLEPIS, Miq. Syst. Pip. 282 (1843) = **Piper**, Linn.
brevicuspis, Miq. l. c. 283 = P. Rhyncholepsis.
Cumingiana, Miq. l. c. 282 = P. Rhyncholepsis.
Haeankeana, Miq. l. c. 284 = P. denudatum.
Roxburghii, Miq. l. c. = Peperomia rostrata.

RHYNCHOLEPSIS, C. DC. in DC. Prod. xvi. 1. 240 (1869) = praec.

RHYNCHOPAPPUS, Dulac, Fl. Hautes-Pyr. 493 (1867) = **Crepis**, Vaill. (Compos.).
albidus, Dulac, l. c. = C. alpina.
cinereus, Dulac, l. c. = C. taraxacifolia.
foetidus, Dulac, l. c. = C. foetida.

RHYNCHOPERA, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Rar. 103. t. 41 (1850) = **Pleurothallis**, R. Br. (Orchid.).
pedunculata, Klotzsch, l. c. = pedunculata.
punctata, Karst. Ausw. Neuer Gew. Venez. 21. t. 7.—Venezuela.

RHYNCHOPETALUM, Fresen. in Flora, xxi. (1838) II. 603 = **Lobelia**, Linn. (Campan.).
montanum, Fresen. l. c. 604 = L. Rhynchopetalum.

RHYNCHOPSIDIUM, DC. in Mém. Soc. Phys. Genèv. vii. II. (1836) 283, 286 = **Relhania**, L'Hérit. (Compos.).
pedunculatum, DC. Prod. vi. 290 = Relh. pedunculata, *Harv.*
pumilum, DC. l. c. = Relh. pedunculata, *L'Hérit.*
sessiliflorum, DC. in Mém. Soc. Phys. Genèv. vii. II. (1836) 283 = Relh. sessiliflora.

RHYNCHOPYLE, Engl. Bot. Jahrb. i. (1881) 183.
AROIDEAE, Benth. & Hook. f. iii. 985.
elongata, *Engl. l. c.* 184.—Borneo.
marginata, *Engl. l. c.*—Borneo.

RHYNCHOSIA, Lour. Fl. Cochinch. 460 (1790).
LEGUMINOSAE, Benth. & Hook. f. i. 542.
ARCYPHYLLUM, Ell. in Journ. Acad. Philad. i. (1818) 371.
CHRYSOSCIAS, E. Mey. Comm. Pl. Afr. Austr. 139 (1835).
COPISMA, E. Mey. l. c. 132 (1835).
CYANOSPERMUM, Wight & Arn. Prod. 259 (1834).
DOLICHOLUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 354.
ERIOSEMA, Reichb. Consp. 150 (1828).
EURIOSMA, Desv. in Ann. Sc. Nat. Ser. I. ix. (1826) 421.
HIDROSIA, E. Mey. Comm. Pl. Afr. Austr. 89 (1835).
NOMISMIA, Wight & Arn. Prod. 236 (1834).
ORTHODANUM, E. Mey. Comm. Pl. Afr. Austr. 131 (1835).
PHYLLOMATIA, Benth. in Ann. Wien. Mus. ii. (1838) 113.
PITCHERIA, Nutt. in Journ. Acad. Philad. vii. (1834) 93.
POLYTROPIA, Presl, Symb. Bot. i. 21. t. 13 (1830).
PTYCHOCENTRUM, Benth. in Ann. Wien. Mus. ii. (1838) 113.
SIGMODOSTYLES, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 93.
acuminata, Eckl. & Zeyh. Enum. 253 = caribaea.
acuminatissima, *Miq. Fl. Ind. Bat.* i. I. 171.—Java.
acutifolia, *Benth. Fl. Austral.* ii. 266.—Austral.
acutissima, *Thw. Enum. Pl. Zeyl.* 413.—Ind. or.
adenantha, *Miq. Fl. Ind. Bat.* i. I. 170.—Java.
adenodes, *Eckl. & Zeyh. Enum.* 254.—Afr. austr.
aequinoctialis, Duchass. & Walp. in Linnaea, xxiii. (1850) 743 = reticulata.
Afzelii, R. Br. ex Baker, in Oliver, Fl. Trop. Afr. ii. 225 = Eriosema Afzelii.
albo-nitens, Lem. Illustr. Hortic. viii. (1861) t. 290 = Desmodium scutatum.
amatymbica, Eckl. & Zeyh. Enum. 254 = adenodes.
angustifolia, *DC. Prod.* ii. 388.—Afr. austr.
angustifolia, Engelm. ex A. Gray, Pl. Wright. i. 44 = texana.
antennulifera, *Baker, in Oliver, Fl. Trop. Afr.* ii. 223.—Afr. trop.
argentea, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 419.—Afr. trop.
argentea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 253.—Afr. austr.
aurea, *DC. Prod.* iii. 386.—Ind. or.
aureo-guttata, *Andersson. in Vet. Akad. Handl. Stockh.* 1853 (1855) 252.—Ins. Galapag.
australis, *Benth. Fl. Austral.* ii. 267.—Austral.
avensis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 222.—Burma.
Balansae, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 31.—Paraguay.
barbata, *Steud. Nom.* ed. II. ii. 453.—Peruv.
Beddomei, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 222.—Ind. or.
biflora, DC. Prod. ii. 387 = Atylosia scarabaeoides.
Bocandeana, *Webb, in Hook. Kew Journ.* ii. (1850) 345.—Ins. Cap. Virid.
Borianii, Schweinf. Reliq. Kotschy, 30 = Glycine Borianii.
bracteata, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 225.—Ind. or.
bullata, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 257.—Afr. austr.
cajanoides, Guill. & Perr. Fl. Seneg. Tent. i. 215 = Eriosema cajanoides.
calycina, *Guill. & Perr. l. c.* 214.—Afr. trop.
calycosa, *Hemsl. Diagn. Pl. Nov.* 48.—Panama.
cana, *DC. Prod.* ii. 386.—Ind. or.
candicans, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 258.—Burma.

RHYNCHOSIA:—
Candollei, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 473 = minima.
capitata, DC. Prod. ii. 386 = aurea.
caribaea, Auct. Plur. ex Benth. in Mart. Fl. Bras. xv. I. 205 = minima.
caribaea, *DC. Prod.* ii. 384.—Am. bor.; Ind. occ.; Afr. trop. et austr.
Cephalotes, Steud. Nom. ed. II. ii. 453 = Psoralea tomentosa.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 387.—Madag.
chrysoposta, *Steud. Nom.* ed. II. ii. 453.—Afr. austr.
Chrysoscias, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 248.—Afr. austr.
Cienkowskii, Schweinf. Reliq. Kotschy. 31 = Tephrosia lupinifolia.
Clausseni, *Benth. in Mart. Fl. Bras.* xv. I. 201.—Bras.
clivorum, *S. Moore, in Journ. Bot.* xvi. (1878) 131.—Afr. austr.
codoorensis, Bedd. Ic. Pl. Ind. Or. i. 70 = Heynei.
confertiflora, A. Rich. Tent. Fl. Abyss. i. 231 = Pseudarthria confertiflora.
congensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 217.—Afr. trop.
coodoorensis, Baker, in Hook. f. Fl. Brit. Ind. ii. 220 = *codoorensis.*
cordata, Steud. Nom. ed. II. ii. 453 = Eriosema cordatum.
corylifolia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. I. 202.—Bras.
crassifolia, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 252.—Afr. austr.
crinita, DC. Prod. ii. 389 = Eriosema crinitum.
crotalarioides, DC. l. c. 387 = Flemingia congesta.
Cunninghamii, *Benth. Fl. Austral.* ii. 266.—Austral.
cyanosperma, *Benth. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 218.—As. et Afr. trop.
debilis, *G. Don, Gen. Syst.* iii. 347.—Afr. trop.
densiflora, *DC. Prod.* ii. 386.—Ind. or.; Afr. trop.
densiflora, Wall. Cat. n. 5492 D = aurea.
difformis, DC. Prod. ii. 384 = tomentosa.
diffusa, *DC. l. c.* 388.—Mexic.
discolor, Klotzsch, in Peters, Reise Mossamb. Bot. 30 = R. Memnonia.
discolor, *Mart. & Gal. Bull. Acad. Brux.* x. II. (1843) 199.—Mexic.
diversifolia, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 33.—Paraguay.
Dregei, Steud. Nom. ed. II. ii. 453 = Eriosema squarrosum.
Ecklonei, *Steud. l. c.*—Afr. austr.
edulis, *Griseb. in Goett. Abh.* xix. (1874) 123.—Reg. Argent.
elegans, *A. Rich. Tent. Fl. Abyss.* i. 228.—Abyss.
elongata, DC. Prod. ii. 386 = aurea.
erecta, DC. l. c. 384 = tomentosa.
ervoidea, DC. Mém. Lég. 368; Prod. ii. 386 = minima.
erythrinoides, Schlecht. & Cham. in Linnaea, v. (1830) 587 = phaseoloides.
exigua, *Andersson. in Vet. Acad. Handl. Stockh.* 1853 (1855) 252.—Ins. Galapag.
faginea, Guill. & Perr. Fl. Seneg. Tent. i. 212 = Flemingia faginea.
falcata, E. Mey. Comm. Pl. Afr. Austr. 136 = minima.
Falconeri, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 224.—Reg. Himal.
ferruginea, *A. Rich. Tent. Fl. Abyss.* i. 229.—Abyss.
ferulaefolia, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 249.—Afr. austr.
ficifolia, *Benth. l. c.* 251.—Afr. austr.
filicaulis, R. Grah. in Wall. Cat. n. 5502 = minima.
filicaulis, *Welw. ex. Baker, in Oliver, Fl. Trop. Afr.* ii. 221.—Afr. trop.
filipes, *Benth. in Linnaea*, xxiv. (1851) 644.—Ind. or.
flagellaris, Fenzl, in Flora, xxvii. (1844) 312, nomen = minima.
flavissima, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 219.—Afr. trop.
Fridericiana, *DC. Prod.* ii. 387.—Ins. Philipp.
frutescens, Steud. Nom. ed. II. i. 459, sphalm. = rufescens.
galactioides, DC. Prod. ii. 387 = Calopogonium galactoides.
galactioides, *Endl. ex Walp. Rep.* i. 790.—Am. bor.
Garcini, *DC. Prod.* ii. 388.—Persia.

RHYNCHOSIA:—
gibba, E. Mey. in Linnaea, vii. (1832) 170 = caribaea.
glabra, DC. Prod. ii. 387 = glandulosa.
glandulosa, *DC. l. c.* 388.—Afr. austr.
glomerata, Guill. & Perr. Fl. Seneg. Tent. i. 216 = Eriosema glomeratum.
Grahami, Wall. Cat. n. 5504 = Eriosema chinense.
grandiflora, Schlecht. & Cham. in Linnaea, v. (1830) 588 = Eriosema grandiflorum.
grandifolia, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.
Grantii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 223.—Afr. trop.
Heynei, *Wight & Arn. Prod.* 240.—Ind. or.
himalensis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 225.—Reg. Himal.
hirsuta, *Eckl. & Zeyh. Enum.* 256.—Afr. austr.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 198.—Mexic.
Hookeri, *G. Don, Gen. Syst.* ii. 347.—Ind. occ.
humilis, Eckl. & Zeyh. Enum. 253 = totta.
incana, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 223 = R. Pseudo-cajan.
inflata, Boj. Hort. Maurit. 104 = caribaea.
intermedia, Kotschy & Peyr. Pl. Tinn. 7. n. 10 = caribaea?
Lablab, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 386.—Madag.
latifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 285.—Am bor.
latissima, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 221.—Afr. trop.
laxiflora, Cambess. in Jacquem. Voy. Bot. 44 = minima.
leucophylla, *Benth. in Mart. Fl. Bras.* xv. I. 202.—Bras.
leucoscias, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 249.—Afr. austr.
lineata, *Benth. in Mart. Fl. Bras.* xv. I. 202.—Bras.
lobata, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 419.—Bras.
longepedunculata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 226 = Eriosema longepedunculatum.
longeracemosa, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 198.—Mexic.; Panama.
longeramosa, Griseb. Fl. Brit. W. Ind. 191, in syn., sphalm. = praec.
lucida, DC. Prod. ii. 387 = Strongylodon ruber.
Luschnathiana, *Walp. in Linnaea*, xiv. (1840) 295.—Bras.
macrocarpa, *Benth. Pl. Hartw.* 11.—Mexic.
macrophylla, DC. Prod. ii. 385 = reticulata.
malacophylla, Boj. Hort. Maurit. 104 = caribaea.
Mannii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 217.—Afr. trop.
maxima, Boj. Hort. Maurit. 105 = Baukea insignis.
medicaginea, DC. Prod. ii. 386 = minima.
melanosperma, Klotzsch, in Peters, Reise Mossamb. Bot. 30 = caribaea.
melanosticta, *Griseb. in Goett. Abh.* xix. (1874) 124.—Reg. Argent.
Memnonia, *DC. Prod.* ii. 386.—Afr. trop.; Arab.; Sind.
mendacinensis, *Gill. ex Hook. & Arn. in Bot. Misc.* iii. (1833) 199.—Chili.
mendozinensis, Walp. Rep. i. 788 = praec.
menispermoidea, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 102.—Am. bor.
mexicana, Hook. & Arn. Bot. Beech. Voy. 287 = minima.
mexicana, Steud. Nom. ed. II. ii. 454 = Eriosema grandiflorum.
Meyeri, *Steud. l. c.*—Afr. austr.
microphylla, Wall. Cat. n. 5497 = minima.
microscias, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 249.—Afr. austr.
minima, *DC. Prod.* ii. 385.—Cosmop. trop. et subtrop.
mollis, *DC. l. c.* 388.—Afr. trop.
mollissima, *Dalz. in Journ. Linn. Soc.* xiii. (1873). 186.—Ind. or.
mollissima, *G. Don, Gen. Syst.* ii. 347.—Afr. trop.
mollissima, *Shuttl. ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 256.—Am. bor.
mollissima, *Zoll. & Mor. in Nat. en Geneesk. Arch. Nêerl. Indie*, iii. (1846) 78.—Java.
monosperma, *Griseb. in Goett. Abh.* xix. (1874) 123.—Reg. Argent.
nervosa, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 253.—Afr. austr.
nitens, *Benth. ex Harv. l. c.* 257.—Afr. austr.
nitida, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.

RHYNCHOSIA :—
nuda, DC. Mém. Légum. 366; Prod. ii. 385 = minima.
nummularia, *DC. Prod.* ii. 386.—Ind. or.
Orthodanum, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 257. —Afr. austr.
pallida, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 32.—Paraguay.
paniculata, Steud. Nom. ed. II. ii. 454 = totta.
parviflora, Steud. l. c. = Eriosema parviflorum.
parvifolia, *DC. Mém. Légum.* 367; *Prod.* ii. 385.—Ins. S. Doming.
phaseoloides, *DC. Prod.* ii. 385.—Am. trop.
picta, Steud. Nom. ed. II. i. 416, in syn., sphalm. = secunda.
pilosa, *Steud. l. c.* ii. 454.—Afr. austr.
pilosa, *Wall. Cat.* n. 5499.—Burma.
pinnata, *Harv. Thes. Cap.* i. 50.—Afr. austr.
platyphylla, *Benth. in Mart. Fl. Bras.* xv. I. 201.—Bras.
polystachya, A. Rich. Tent. Fl. Abyss. i. 231 = Eriosema polystachyum.
polystachya, Steud. Nom. ed. II. ii. 454 = Eriosema cajanoides.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 121.—Panama.
precatoria, DC. Prod. ii. 385 = phaseoloides.
prostrata, R. Grah. in Wall. Cat. n. 5496 = minima.
Pseudo-cajan, *Cambess. in Jacquem, Voy. Bot.* 45. t. 55.—Reg. Himal.
psoraloides, *DC. Prod.* ii. 389.—Madag.
puberula, *Steud. Nom.* ed. II. ii. 454; *Harv. in Harv. & Sond. Fl. Cap.* ii. 255.—Afr. austr.
pubescens, *DC. Prod.* ii. 386.—Abyss.
pulchella, *DC. l. c.* 389.—Am. austr.
pulverulenta, Stocks, in Hook. Kew Journ. iv. (1852) 147 = R. Memnonia.
punctata, DC. Mém. Légum. 365; Prod. ii. 385 = minima.
quadrata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 254.—Afr. austr.
reniformis, DC. Prod. ii. 384 = tomentosa.
resinosa, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 218.—Afr. trop.
reticulata, *DC. Prod.* ii. 385.—Am. trop.
rhodophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 133.—Madag.
rhombifolia, DC. Prod. ii. 386 = minima.
rhomboidea, *F. Muell. ex Benth. Fl. Austral.* ii. 265. —Austral.
rigidula, *DC. Prod.* ii. 388.—Afr. austr.
rigidula, Eckl. & Zeyh. Enum. 255 = totta.
riparia, Eckl. & Zeyh. l. c. = glandulosa.
rosea, DC. Prod. ii. 387 = Canavalia obtusifolia.
rostrata, *Benth. Fl. Austral.* ii. 266.—Austral.
rotundifolia, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.
rufa, DC. Prod. ii. 388 = Eriosema rufum.
rufescens, *DC. l. c.* 387.—Ind. or.; Malaya.
saligna, Steud. Nom. ed. II. ii. 454 = Eriosema salignum.
scarabaeoides, DC. Prod. ii. 387 = Atylosia scarabaeoides.
Schimperi, *Hochst. ex Boiss. Fl. Orient.* ii. 626.—Oriens.
secunda, *Eckl. & Zeyh. Enum.* 252.—Afr. austr.
Senna, *Gill. ex Hook. & Arn. in Bot. Misc.* iii. (1833) 199.—Reg. Argent.
sennaarensis, Hochst. ex Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 655 = flavissima.
sericea, *Gill. ex Hook. & Arn. in Bot. Misc.* iii. (1833) 199.—Chili.
sericea, *Span. in Linnaea*, xv. (1841) 195.—Ind. or.; Malaya.
sericea, Vidal, Sinops. t. 40. f. D = Flemingia congesta.
sericophylla, *Kunze, in Linnaea*, xx. (1847) 63.—Java.
sessiliflora, DC. Prod. ii. 389 = Tephrosia brevipes.
sessilifolia, Benth. in Mart. Fl. Bras. xv. I. 46, sphalm. = praec.
sigmodes, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 251.—Afr. austr.
simplicifolia, DC. Prod. ii. 389 = Eriosema simplicifolium.
simplicifolia, Walp. Rep. i. 787 = singulifolia.
simplicifolia, Wood, Class-Book (1861), 321 = tomentosa.
singulifolia, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.
splendens, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 654.—Afr. trop.
stipulosa, A. Rich. Tent. Fl. Abyss. i. 229 = viscosa.

RHYNCHOSIA :—
striata, *G. Don, Gen. Syst.* ii. 347.—Afr. trop.
suaveolens, *DC. Prod.* ii. 387.—Ind. or.
subsericea, Walp. Ann. ii. 434 = Galactia dubia.
tenuicaulis, Wall. Cat. n. 5495 = minima.
tenuis, Steud. Nom. ed. II. ii. 454 = minima.
texana, Torr. & Gray, Fl. N. Am. i. 687 = R. Senna.
tomentosa, Baill. in Bull. Soc. Linn. Par. i. (1883) 387 = cyanosperma.
tomentosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 23.—Am. bor.
tomentosa, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 186 = sericea.
totta, *DC. Prod.* ii. 388.—Afr. austr.
trichodes, Harv. in Harv. & Sond. Fl. Cap. ii. 258 = totta.
trinervis, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.
tuberosa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 227 = Eriosema tuberosum.
uniflora, Harv. in Harv. & Sond. Fl. Cap. ii. 249 = angustifolia.
velutina, R. Grah. in Wall. Cat. n. 5501 = Atylosia rugosa.
velutina, *Wight & Arn. Prod.* 238.—Ind. or.
versicolor, *Baker, in Journ. Linn. Soc.* xx. (1883) 132. —Madag.
vestita, Wall. Cat. n. 5505 = Atylosia barbata.
villosula, Thw. Enum. Pl. Zeyl. 412 = viscosa.
violacea, DC. Prod. ii. 388 = Eriosema violaceum.
virgata, R. Grah. in Wall. Cat. n. 5503 = Eriosema chinense.
viscida, DC. Prod. ii. 387 = Pseudarthria viscida.
viscidula, *Steud. Nom.* ed. II. ii. 454.—Afr. austr.
viscosa, *DC. Prod.* ii. 387.—As. et Afr. trop.
volubilis, *Lour. Fl. Cochinch.* 460.—China; Japon.
volubilis, Wood, Class-Book (1861), 321; Bot. & Fl. 96 = tomentosa.
Wightiana, R. Grah. in Wall. Cat. n. 5500 = Atylosia Candollei.
Zeyheri, Steud. Nom. ed. II. ii. 454 = Eriosema squarrosum.

RHYNCHOSPERMUM, A. DC. in DC. Prod. viii. 431 (1844) = **Rhynchodia**, Benth. (Scrophul.).
ellipticum, A. DC. l. c. = Rhynchodia Wallichii.
Wallichii, A. DC. l. c. = Rhynchodia Wallichii.

RHYNCHOSPERMUM, Lindl. in Journ. Hort. Soc. i. (1846) 74 = **Trachelospermum**, Lem. (Apocyn.).
jasminoides, Lindl. l. c. = T. jasminoides.

RHYNCHOSPERMUM, Reinw. in Syll. Ratisb. ii. (1828) 7. *COMPOSITAE*, Benth. & Hook. f. ii. 263.
LEPTOCOMA, Less. in Linnaea, vi. (1831) 130.
ZOLLINGERIA, Sch. Bip. in Flora, xxxvii. (1854) 273.
verticillatum, *Reinw. in Syll. Ratisb.* ii. (1828) 8.—Java.

RHYNCHOSPORA, Willd. Enum. Hort. Berol. 71 (1809) = **Rynchospora**, Vahl (Cyperac.).

RHYNCHOSTELIS, Reichb. f. in Bot. Zeit. x. (1852) 770 = **Leochilus**, Knowl. & Westc. (Orchid.).
pygmaea, Reichb. f. l. c. = L. pygmaeus (in Addend.).

RHYNCHOSTEMON, Steetz, in Lehm. Pl. Preiss. ii. 333 (1847) = **Thomasia**, J. Gay (Stercul.).
canescens, Steetz, l. c. 335 = T. glutinosa.
glutinosum, Steetz, l. c. 334 = T. glutinosa.

RHYNCHOSTIGMA, Benth. in Benth. & Hook. f. Gen. ii. 771 (1876) = **Toxocarpus**, Wight & Arn. (Asclep.).
brevipes, Benth. in Hook. Ic. Pl. sub t. 1189.—Afr. trop.
parviflorum, Benth. l. c.—Afr. trop.
racemosum, Benth. l. c. t. 1189.—Afr. trop.

RHYNCHOSTYLIS, Blume, Bijdr. 285. t. 49 (1825) (Rynchostylis). *ORCHIDEAE*, Benth. & Hook. f. iii. 574.
gurwalica, Reichb. f. in Walp. Ann. vi. 888 = retusa.
guttata, Reichb. f. in Bonplandia, ii. (1854) 93 = retusa.
papillosa, Heynh. Nom. ii. 594 = Saccolabium papillosum.
praemorsa, Blume, Bijdr. 286 = retusa.
retusa, *Blume, l. c.*—Ind. or.; Malaya.
violacea, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Ins. Philipp.

RHYNCHOSTYLIS, Tausch, in Flora, xvii. (1834) 343 = **Chaerophyllum**, Linn. (Umbell.).
hirsutus, Tausch, l. c. = C. hirsutum.

RHYNCHOTECHUM, Blume, Bijdr. 775 (1825). *GESNERACEAE*, Benth. & Hook. f. ii. 1016.
CHEILOSANDRA, Griff. in Lindl. Veg. Kingd. 672 (1847).
CHILIANDRA, Griff. Notul. iv. 150 (1854).
CORYSANTHERA, Wall. Cat. n. 6411 (1831?).
alternifolium, *C. B. Clarke, in DC. Monog. Phan.* v. 198.—Reg. Himal.
Brandisii, *C. B. Clarke, l. c.* 199.—Burma.
calycinum, *C. B. Clarke, l. c.*—Reg. Himal.
ellipticum, *DC. Prod.* ix. 285.—Reg. Himal.
hispidum, *C. B. Clarke, in DC. Monog. Phan.* v. 197. —Java.
latifolium, *Hook. f. & Thoms. ex C. B. Clarke, Comm. et Cyrt. Beng.* t. 93.—Reg. Himal.; Burma.
parviflorum, *Blume, Bijdr.* 775.—Penang; Java.
tenue, *C. B. Clarke, in DC. Monog. Phan.* v. 199.—Ins. Philipp.
vestitum, *Wall. ex C. B. Clarke, Comm. et Cyrt. Beng.* t. 92.—Reg. Himal.

RHYNCHOTHECA, Ruiz & Pav. Prod. 82. t. 15 (1794). *GERANIACEAE*, Benth. & Hook. f. i. 275.
AULACOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 149.
RHYNCOTHELIA, Pers. Syn. ii. 92 (1807).
diversifolia, H. B. & K. Nov. Gen. et Sp. v. 233. t. 465 = spinosa.
integrifolia, H. B. & K. l. c. 232. t. 464 = spinosa.
spinosa, *Ruiz & Pav. Syst. Veg.* 142.—Ecuador.

RHYNCOSIA, Webb, in Hook. Kew Journ. ii. (1850) 345, sphalm. = **Rhynchosia**, Lour. (Legumin.).

RHYNCOSPERMUM, A. DC. in DC. Prod. viii. 431 (1844), sphalm. = RHYNCHOSPERMUM, Lindl.

RHYNCOSTYLIS, Steud. Nom. ed. II. ii. 457 (1841), sphalm. = **Rhynchostylis**, Blume (Orchid.).

RHYNCOTHECUM, A. DC. in DC. Prod. ix. 285 (1845), sphalm. = **Rhynchotechum**, Blume (Gesner.).

RHYNCOTHELIA, Pers. Syn. ii. 92 (1807) = **Rhynchotheca**, Ruiz & Pav. (Geran.).

RHYNEA, DC. Prod. vi. 154 (1837); in Deless. Ic. Sel. iv. t. 52 (1839) = **Cassinia**, R. Br. (Compos.).
phylicaefolia, DC. ll. cc.—Afr. austr.

RHYNEA, Scop. Introd. 262 (1777) = **Mesua**, Linn. (Guttif.).

RHYNOSPERMUM, Walp. Ann. iii. 920 (1852-53), sphalm. = RHYNCHOSPERMUM, DC. = **Rhynchodia**, Benth. (Scrophul.).

RHYPARIA, Hassk. Cat. Hort. Bog. Alt. 239 (1844) = **Ryparosa**, Blume (Euphorb.).

RHYSOSPERMUM, Gaertn. f. Fruct. iii. 232. t. 224 (1805) = **Notelaea**, Vent. (Oleac.).
ellipticum, Gaertn. f. l. c. t. 224. f. 2 = N. nervosa.
nervosa, Gaertn. f. l. c. 232 = N. nervosa.

RHYSOTOECHIA, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 498, 656. *SAPINDACEAE*.
bifoliolata, *Radlk. l. c.* 656.—Austral.
flavescens, *Radlk. l. c.* 658.—Austral.
grandifolia, *Radlk. l. c.* 657.—Borneo.
Mortoniana, *Radlk. l. c.* 656.—Austral.
ramiflora, *Radlk. l. c.* 657.—Ins. Celebes.
Robertsoni, *Radlk. l. c.* 657.—Austral.

RHYSSOCARPUS, Endl. in Bot. Zeit. i. (1843) 459. *RUBIACEAE*, Benth. & Hook. f. ii. 81.
PLEUROCARPUS, Klotzsch, in Bonplandia, vii. (1859) 3.
pubescens, *Endl. in Bot. Zeit.* i. (1843) 459.—Am. trop.

RHYSSOLOBIUM, E. Mey. Comm. Pl. Afr. Austr. 217 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 769.
dumosum, *E. Mey. l. c.*—Afr. austr.

RHYSSOSTELMA, Decne. in DC. Prod. viii. 590 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 751.
nigricans, *Decne. l. c.*—Reg. Argent.

RHYTACHNE, Desv. in Ham. Prod. Pl. Ind. Occ. 11 (1825). Gen. dub. *GRAMINEARUM*, Benth. & Hook. f. iii. 1096.
rottboellioides, *Desv. l. c.* 12.—Afr. trop.; Ind. occ.

RHYTICARPUS, Sond. in Harv. & Sond. Fl. Cap. ii. 540 (1862). *UMBELLIFERAE*, Benth. & Hook. f. i. 887.
difformis, *Benth. & Hook. f. Gen.* ii. 887.—Afr. austr.
Ecklonis, *Sond. in Harv. & Sond. Fl. Cap.* ii. 540.—Afr. austr.
rugosus, *Sond. l. c.*—Afr. austr.

RHYTICARYUM, Becc. Malesia, i. (1878) 256 = **Ryticaryum**, Becc. (Icacin.).

RHYTIDANDRA, A. Gray, Bot. U. St. Expl. Exped. i. 302 (1854) = **Marlea**, Roxb. (Cornac.).
polyosmoides, F. Muell. Fragm. ii. 176 = M. vitiensis.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 303. t. 28 = M. vitiensis.

RHYTIDANTHE, Benth. in Enum. Pl. Hueg. 63 (1837) = **Leptorhynchos**, Less. (Compos.).
scabra, Benth. l. c. = L. elongatus.

RHYTIDOLOBUS, Dulac, Fl. Hautes-Pyr. 113 (1867) = **Hyacinthus**, Tourn. (Liliac.).
appendiculatus, Dulac, l. c. = H. romanus.

RHYTIDOPHYLLUM, Mart. Nov. Gen. et Sp. iii. 38 (1829) (Rytidophyllum). *GESNERACEAE*, Benth. & Hook. f. ii. 1006.
auriculatum, *Hook. Bot. Mag.* t. 3562.—Hab. ?
Berteroanum, *Mart. Nov. Gen. et Sp.* iii. 39.—Ind. occ.
crenulatum, *DC. Prod.* vii. 524.—Cuba.
exsertum, *Griseb. Cat. Pl. Cub.* 198.—Cuba.
floribundum, *Lem. Fl. des Serres*, ii. (1846) t. 178.—Cuba.
grande, Linden, ex Hanst. in Linnaea, xxxiv. (1865–66) 311 = tomentosum.
grande, *Mart. Nov. Gen. et Sp.* iii. 39.—Ind. occ.
Humboldtii, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 18 = Campanea Humboldtii.
leucomallon, *Hanst. in Linnaea*, xxxiv. (1865–66) 312.—Ind. occ.
melastoma, Poepp. & Endl. Nov. Gen. et Sp. iii. 7 (= *Moussonia melastoma*).—Peruv.
Oerstedii, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 17 = Campanea Oerstedii.
petiolare, *DC. Prod.* vii. 524.—Ins. S. Doming.
Plumerianum, *DC. l. c.*—Ins. S. Doming.
prasinatum, *Mart. ex Steud. Nom.* ed. II. ii. 448.—Bras.
purpureum, Poepp. & Endl. Nov. Gen. et Sp. iii. 6. t. 206 = Isoloma purpurea.
Tigridia, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 18 = Campanea grandiflora.
tomentosum, Hort. Berol. ex Hanst. in Linnaea, xxxiv. (1865–66) 310 = petiolare.
tomentosum, *Mart. Nov. Gen. et Sp.* iii. 39.—Ind. occ.
Wrightianum, *Griseb. Cat. Pl. Cub.* 198.—Cuba.

RHYTIDOSPERMUM, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 277 (1836–50) = **Matricaria**, Tourn. (Compos.).
inodorum, Sch. Bip. l. c. = M. inodora.

RHYTIDOSPORUM, F. Muell. ex Hook. f. Fl. Tasm. (1855) = **Marianthus**, Hueg. (Pittosp.).
procumbens, F. Muell. l. c. = M. procumbens.
Stuartianum, F. Muell. l. c. = M. procumbens.

RHYTIDOSTYLIS, Reichb. Nom. 101 (1841) = RYTIDOSTYLIS, Hook. & Arn. = **Elaterium**, Jacq. (Cucurb.).

RHYTIDOTUS, Hook. f. in Benth. & Hook. f. Gen. ii. 101 (1873) = **Rytidotus**, Hook. f. (Rubiac.).

RHYTIGLOSSA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 285, 444 (1836) = **Dianthera**, Gronov. (Acanth.).
acuminata, Nees, in DC. Prod. xi. 354 = D. comata.
acuminatissima, Miq. Stirp. Surin. Sel. 129.—Guian.

RHYTIGLOSSA:—
amazonica, Nees, in Mart. Fl. Bras. ix. 127.—Bras.
Anagallis, Mart. & Nees, in Mart. l. c. 119 = Dianthera Anagallis.
androsaemifolia, Nees, in DC. Prod. xi. 352 = D. androsaemifolia.
angustifolia, Nees, in Mart. Fl. Bras. ix. 121 = D. angustifolia.
Arnottiana, Nees, in DC. Prod. xi. 343.—Bras.
axillaris, Nees, in Mart. Fl. Bras. ix. 122.—Bras.
barbata, Nees, in DC. Prod. xi. 355.—Ind. occ.
breviflora, Nees, l. c. 352 = D. breviflora.
campestris, Nees, in Mart. Fl. Bras. ix. 118 = D. campestris.
Candelariae, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 158 = D. Candelariae.
caracasana, Nees, in Mart. Fl. Bras. ix. 127.—Am. trop.
cayennensis, Nees, in DC. Prod. xi. 346 = D. cayennensis.
chamaedryoides, Nees, in Mart. Fl. Bras. ix. 123.—Bras.
ciliata, Nees, in Linnaea, xv. (1841) 364.—Afr. austr.
cordifolia, Nees, in Mart. Fl. Bras. ix. 121.—Bras.
corymbulosa, Nees, in DC. Prod. xi. 355.—Guatemala.
cristata, Regel, Gartenfl. (1852) 132.—Am. austr. ?
cuspidulata, Nees, in DC. Prod. xi. 348.—Peruv.
dasyclados, Mart. ex Nees, in Mart. Fl. Bras. ix. 126 = D. dasyclados.
distorta, Nees, l. c. 125 = D. distorta.
divergens, Nees, l. c. 128.—Bras.
Ecklonіana, Nees, in DC. Prod. xi. 336.—Afr. austr.
ensiformis, Wood, Class-Book (1861), 536 = D. americana ?
geniculata, Nees, in DC. Prod. xi. 341 = D. secunda.
genuflexa, Nees, in Mart. Fl. Bras. ix. 121.—Bras.
glandulifera, Presl, Bot. Bemerk. 95 = *origanoides*.
glandulosa, Hochst. in Flora, xxviii. (1845) 71 = Blechum laxiflorum.
havanensis, Nees, in DC. Prod. xi. 344 = D. comata.
hirsuta, Nees, l. c. 342.—Peruv.
Hookeriana, Nees, l. c. 348 = D. Hookeriana.
humilis, Nees, l. c. 340 = D. ovata.
Hyssopus, Nees, l. c. 345.—Mexic.
ilhensis, Moric. ex Nees, l. c. 342.—Bras.
inaequalis, Nees, l. c. 349 = D. inaequalis.
indica, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 281 = D. indica.
laeta, Nees, in Mart. Fl. Bras. ix. 126 = D. laeta.
laevilinguis, Nees, l. c. 120.—Bras.
lanceolata, Buek, Index in DC. Prod. iii. 379 = *lineolata*.
latifolia, Nees, in DC. l. c. xi. 353 = D. latifolia.
lavandulaefolia, Nees, in Mart. Fl. Bras. ix. 124.—Bras.
leucophloea, Nees, l. c. 129.—Bras.
Lindeniana, Nees, in DC. Prod. xi. 349 = D. Lindeniana.
linearis, Nees, in Mart. Fl. Bras. ix. 125.—Bras.
lineolata, Nees, in DC. Prod. xi. 341.—Peruv.
longiflora, Nees, in Mart. Fl. Bras. ix. 128 = D. longiflora.
lucida, Nees, in DC. Prod. xi. 341 = D. secunda.
maynensis, Nees, l. c. 346.—Peruv.
menthoides, Nees, in Mart. Fl. Bras. ix. 122.—Bras.
mexicana, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 157 = D. mexicana.
Meyeniana, Nees, in Mart. Fl. Bras. ix. 121 = Beloperone Meyeniana.
microphylla, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 159 = D. Candelariae.
microstachya, Nees, in DC. Prod. xi. 342.—Peruv.
Moricandiana, Nees, l. c. 341.—Guian.
nana, Nees, l. c. 346.—Mexic.
nemoralis, Nees, l. c. 159, in syn., nomen.—Hab. ?
oblonga, Nees, l. c. 343.—N. Granat.
oblonga, Nees, in Mart. Fl. Bras. ix. 124 = D. oblonga.
obtusifolia, Nees, l. c. 120 = D. ovata.
origanoides, Nees, in Linnaea, xv. (1841) 365.—Afr. austr.
ovata, Nees, in DC. Prod. xi. 336.—Afr. austr.
ovatifolia, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 156.—Mexic.
paniculata, Hort. Borbon. ex T. Anders. in Journ. Agri.-Hort. Soc. Ind. N. S. i. (1868) 282 (Justiciae sp.).—Hab. ?
paniculata, Nees, in Mart. Fl. Bras. ix. 129.—Bras.
pauciflora, Nees, l. c. 123.—Bras.
pectoralis, Nees, l. c. 128 = D. pectoralis.

RHYTIGLOSSA:—
pedunculosa, Nees, in DC. Prod. xi. 339 = Dianthera americana.
piauhiensis, Nees, in Mart. Fl. Bras. ix. 124.—Bras.
Poeppigiana, Nees, l. c. 123.—Bras.
polygonoides, Nees, in DC. Prod. xi. 353.—N. Granat.
prolixa, Nees, l. c. 336.—Afr. austr.
ptychostoma, Nees, l. c. 337 = Justicia ptychostoma.
punctata, Nees, l. c. 355 = D. punctata.
radicosa, Nees, l. c. 344 = Ptyssiglottis radicosa.
repens, Nees, in Mart. Fl. Bras. ix. 119.—Bras.
reptans, Nees, in DC. Prod. xi. 343 = D. reptans.
rubicunda, Hochst. in Flora, xxviii. (1845) 71.—Afr. austr.
Ruiziana, Nees, in DC. Prod. xi. 345 = D. lineolata.
Sagraeana, A. Rich. Fl. Cub. Fanerog. ii. 163 = D. Sagraeana.
sarmentosa, Nees, in Mart. Fl. Bras. ix. 119.—Bras.
scabra, Nees, in DC. Prod. xi. 354.—Hab. ?
secunda, Nees, l. c. 340 = D. secunda.
sessilis, Nees, l. c. 345 = D. sessilis.
speciosa, Nees, l. c. 351 = D. speciosa.
symphyantha, Nees, in Mart. Fl. Bras. ix. 125.—Bras.
tenuiflora, Nees, in DC. Prod. xi. 340.—Peruv.
verticillaris, Nees, l. c. 354.—Porto Rico.
viridiflora, Nees, l. c. 346 = Gatesia laete-virens.

RHYTILEUCOMA, F. Muell. Fragm. ii. 90 (1860) = **Chilocarpus**, Blume (Apocyn.).
chilocarpoides, F. Muell. l. c. = C. australis.

RHYTIS, Lour. Fl. Cochinch. 660 (1790) = **Antidesma**, Burm. (Euphorb.).
fruticosa, Lour. l. c. = A. fruticosum.

RHYTISPERMUM, Link, Handb. i. 579 (1829) = **Lithospermum**, Linn. (Borag.).
apulum, Link, l. c.; Reichb. f. Ic. Fl. Germ. xviii. 67 = L. apulum.
apulum, Webb & Berth. Phyt. Canar. iii. 51 = L. microspermum.
arvense, Link, Handb. i. 579 = L. arvense.
dispermum, Link, l. c. 580 = Rochelia stellulata.
fruticosum, Link, l. c. 579 = L. fruticosum.
incrassatum, Reichb. f. Ic. Fl. Germ. xviii. 67 = L. incrassatum.
medium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 122 = L. arvense.
minimum, Reichb. f. Ic. Fl. Germ. xviii. 68 = L. minimum.
patulum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 122.—Europ.
prostratum, Link, Handb. i. 579 = L. fruticosum.
purpuro-coeruleum, Link, l. c. = L. purpuro-caeruleum.
tenuiflorum, Reichb. f. Ic. Fl. Germ. xviii. 67 = L. tenuiflorum.
tinctorium, Link, Handb. i. 579 = Alkanna tinctoria.
villosum, Link, l. c. 580 = L. villosum.

RHYTTIGLOSSA, T. Anders. in Journ. Linn. Soc. vii. (1864) 118, sphalm. = RHYTIGLOSSA, Nees = **Dianthera**, Linn. (Acanth.).

RIANA, Aubl. Pl. Gui. i. 237. t. 94 (1775) = **Alsodeia**, Thou. (Violar.).
Conohoria, Pritz. Ic. Ind. 939, sphalm. = A. flavescens.
guianensis, Aubl. Pl. Gui. i. 237 = A. prunifolia.
tricapsularis, Blanco, Fl. Filip. ed. I. 850 = Chailletia Helferiana.

RIBEIREA, Arruda, ex Koster, Trav. Braz. 499 (1816). *ROSACEAE* ?
sorbilis, *Arruda, ex Koster, l. c.*—Bras.

RIBES, Linn. Gen. ed. I. 68 (1737). *SAXIFRAGEAE*, Benth. & Hook. f. i. 654.
BOTRYCARPUM, A. Rich. Bot. Méd. ii. 487 (1823), in adnot.
CALOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 21.
CEROPHYLLUM, Spach, Hist. Vég. Phan. vi. 152 (1838).
CHRYSOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 18.
COREOSMA, Spach, l. c. 22 (1835).
GROSSULARIA, Tourn. ex Adans. Fam. ii. 243 (1763).
LIEBICHIA, Opiz, Seznam, 59 (1852).

RIBES:—
Rebis, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 26.
Ribesium, Medic. Phil. Bot. i. 120 (1789).
Robsonia, Reichb. Handb. 233 (1837).
acerifolium, Hort. ex C. Koch, Dendrol. i. 649 = rubrum.
acerrimum, Rochel, ex Schult. Oesterr. Fl. ed. II. i. 432 = petraeum.
aciculare, *Sm. in Rees, Cycl.* xxx. n. 25.—Sibir.
acidum, *Ehrh. Beitr.* v. 161, nomen.—Europ.
acidum, Turcz. ex Steud. Nom. ed. II. ii. 457 = rubrum.
aconitifolium, Hort. ex C. Koch, Dendrol. i. 661 = nigrum.
aculeatum, Salisb. Prod. 355 = R. Grossularia.
acuminatum, Wall. Cat. n. 6834 = glaciale.
affine, Dougl. ex Bong. in Mém. Acad. Pétersb. ii. (1833) 138 = laxiflorum.
affine, H. B. & K. Nov. Gen. et Sp. vi. 60 = campanulatum.
albidum, Paxt. Mag. Bot. x. (1843) 55 = sanguineum.
albiflorum, A. Gray, Bot. U. St. Expl. Exped. i. 662, sphalm. = seq.
albifolium, *Ruiz & Pav. Fl. Per.* iii. 12. t. 132. f. 6.—Peruv.
albinervium, Michx. Fl. Bor. Am. i. 110 = rubrum.
album, Gilib. Fl. Lituan. ii. 137 = alpinum.
alceaefolium, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 343 = sanguineum.
alpestre, Wall. ex Decne. in Jacquem. Voy. Bot. 64. t. 75 = R. Grossularia.
alpinoides, *Domb. ex Spach, in Ann. Sc. Nat.* Sér. II. iv. (1835) 31.—Chili.
alpinum, *Linn. Sp. Pl.* 200.—Europ.; As. bor.
alpinum, C. A. Mey. Verz. Pfl. Cauc. 154 = orientale.
alpinum, Siev. in Pall. N. Nord. Beitr. vii. 345 = saxatile.
altaicum, Hort. ex C. Koch, Dendrol. i. 660 = nigrum.
altaicum, Lodd. ex Baxt. in Loud. Hort. Brit. Suppl. ii. 667 = triste.
altissimum, Turcz. ex Ledeb. Fl. Ross. ii. 199 = petraeum.
ambiguum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 251.—Japon.
americanum, Hort. ex C. Koch, Dendrol. i. 649 = rubrum.
americanum, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. bor.
americanum, Pall. Fl. Ross. i. II. 34 = laxiflorum.
apiifolium, Hort. ex C. Koch, Dendrol. i. 661 = nigrum.
atropurpureum, C. A. Mey. in Ledeb. Fl. Alt. i. 268 = petraeum.
augustum, Dougl. ex Baxt. in Loud. Hort. Brit. Suppl. ii. 667 = sanguineum.
aureum, *Pursh, Fl. Am. Sept.* i. 164.—Am. bor. occ.
bacciferum, Hort. ex C. Koch, Dendrol. i. 657 = alpinum.
baicalense, Turcz. ex Steud. Nom. ed. II. ii. 457 = rubrum.
balsamiferum, Kellogg, in Proc. Calif. Acad. ii. (1863) 94 = cereum.
Beatonii ×.—Cf. Loud. Gardn. Mag. xix. (1843) 269.
Berteroanum, *Phil. in Linnaea*, xxviii. (1856) 648.—Chili.
bicolor, Less. ex Ledeb. Fl. Ross. ii. 197 = fragrans.
bicolor, *Phil. in Linnaea*, xxviii. (1856) 646.—Chili.
Biebersteinii, Berland. in Mém. Soc. Phys. Genèv. iii. II. (1826) 60; et in DC. Prod. iii. 482 = petraeum.
Billenei, Medic. Pfl. Anat. 106 = americanum.
Billiardii, *Carr. in Rev. Hortic.* (1867) 140.—Hab.?
boreale, Turcz. ex Steud. Nom. ed. II. ii. 457 = nigrum.
brachystachyum, *Phil. in Anal. Univ. Chil.* (1872) 723.—Chili.
bracteosum, *Dougl. ex Hook. Fl. Bor. Am.* i. 233.—Am. bor. occ.
bracteosum, S. Wats. Bot. King's Exped. 99 = Hudsonianum.
bullatum, *Otto & Dietr. Allg. Gartenz.* x. (1842) 267.—Hab.?
burejense, *F. Schmidt, Reisen Amurl.* 42.—Reg. Amur.
californicum, Hook. & Arn. Bot. Beech. Voy. 346 = Menziesii.
callibotrys, *Wender. Analect. Krit. Bemerk.* (1852) n. 12.—Am. bor.?

RIBES:—
callibotrys, Wendl. ex Steud. Nom. ed. II. ii. 457; Walp. Ann. v. 23 (Quid?).—Hab.?
campanulatum, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 500.—Mexic.
campanulatum, Moench, Meth. 683 = americanum.
carpathicum, Kit. ex Schult. Oesterr. Fl. ed. II. i. 432 = petraeum.
caucasicum, Adams, ex Roem. & Schult. Syst. v. 507 = R. Grossularia.
caucasicum, Bieb. Fl. Taur. Cauc. iii. 160 = petraeum.
cereum, *Dougl. in Trans. Hort. Soc.* vii. (1830) 512.—Am. bor. occ.
chifuense, Hance, in Journ. Bot. xiii. (1875) 36 = fasciculatum.
chilense, C. Koch, Wochenschr. ii. (1859) 138 = villosum.
ciliatum, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 500.—Mexic.
ciliatum, C. Koch, in Linnaea, xvi. (1842) 355 = petraeum.
collinum, *Phil. in Linnaea*, xxviii. (1856) 647.—Chili.
crispum, Dulac, Fl. Hautes-Pyr. 264 = R. Grossularia.
crispum, Hort. ex C. Koch, Dendrol. i. 661 = nigrum.
cucullatum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 340.—Chili.
cuneatum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 426.—Sibir. altaic.
cuneifolium, *Ruiz & Pav. Fl. Per.* iii. 13.—Peruv.
Cynosbati, *Linn. Sp. Pl.* 202.—Am. bor.; Japon.
dauricum, Georgi, Reise, i. 381; ex Ledeb. Fl. Ross. ii. 196 = R. Diacantha.
decumbens, Ledeb. ex Steud. Nom. ed. II. ii. 457, nomen (Quid?).—Sibir.
densiflorum, *Phil. in Linnaea*, xxviii. (1856) 649.—Chili.
dependens, Domb. ex Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 27 = macrobotrys.
desmocarpum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 87.—Reg. Himal.
Diacantha, *Pall. Reise*, iii. 722.—Sibir.
Dikuscha, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) I. 254 = nigrum.
Dillenii, Medic. Beobacht. (1782) 326 = americanum.
dioicum, Moench, Meth. 683 = alpinum.
divaricatum, *Dougl. in Trans. Hort. Soc.* vii. (1830) 515.—Am. bor. occ.
dubium, Jacques, in Rev. Hortic. Sér. IV. iii. (1854) 279 = R. Grossularia.
ebracteatum, Berland. in Mém. Soc. Phys. Genèv. iii. II. (1826) 60 = aureum.
ebracteolatum, C. Gay, Fl. Chil. iii. 34 = alpinoides.
echinatum, Dougl. ex Lindl. Bot. Reg. sub t. 1349 = lacustre.
fasciculatum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 189.—China; Japon.
ferox, Sm. in Rees, Cycl. xxx. n. 26 = Menziesii.
flavum, *Berland. in Mém. Soc. Phys. Genèv.* iii. II. (1826) 60. t. 2.—Hab.?
Fleischmanni, Reichb. ex C. Koch, Dendrol. i. 657 = rubrum.
floridum, Georgi, Beschr. Russ. Reich. III. iv. Nachtr. 262 = laxiflorum.
floridum, L'Hérit, Stirp. Nov. 4 = americanum.
Fontainesii, Colla, Hort. Ripul. App. iii. 2 = flavum.
fragrans, Lodd. Bot. Cab. t. 1533 = aureum.
fragrans, *Pall. in Nov. Act. Acad. Petrop.* x. (1797) 377.—Sibir.
frigidum, H. B. & K. Nov. Gen. et Sp. vi. 62 = hirtum.
fuchsioides, Moç. & Sesse, ex Berland. in Mém. Soc. Phys. Genèv. iii. II. (1826) 58 = speciosum.
Gayanum, *Steud. Nom.* ed. II. ii. 439, in syn.; *Walp. Rep.* ii. 360.—Chili.
glaciale, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 513.—Reg. Himal.
glandulosum, Ait. Hort. Kew. ed. I. i. 279 = prostratum.
glandulosum, *Grauer, Dec.* 2.—Hab.?
glandulosum, *Ruiz & Pav. Fl. Per.* iii. 13.—Chili.
glutinosum, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 476 = sanguineum.
glutinosum, Jacquem. Voy. Bot. 65 = orientale.
Gordonianum ×, Lem. in Fl. des Serres, ii. (1846) t. 165.
gracile, *Michx. Fl. Bor. Am.* i. 111.—Am. bor.

RIBES:—
gracile, Pursh, Fl. Am. Sept. i. 165 = rotundifolium.
gracile, Torr. Fl. N. U. St. i. 269 = R. Cynosbati.
graveolens, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 535 = fragrans.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 88.—Reg. Himal.
Grossularia, Georgi, Beschr. Russ. Reich. III. iv. 799 = aciculare.
Grossularia, *Linn. Sp. Pl.* 201.—Europ.; Afr. bor.; Reg. Himal.
grossulariaefolium, Reichb. ex C. Koch, Dendrol. i. 657 = alpinum.
grossularioides, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 250.—Japon.
grossularioides, Michx. ex Steud. Nom. ed. I. 691 = oxyacanthoides.
grossularium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = R. Grossularia.
heterotrichum, Hort. ex C. Koch, Dendrol. i. 662 = americanum.
heterotrichum, *C. A. Mey. in Ledeb. Fl. Alt.* i. 270.—Sibir. altaic.
himalense, Decne. in Jacquem. Voy. Bot. 66. t. 67 = = rubrum.
himalense, Royle, Illustr. Bot. Himal. 225 = R. Grossularia.
hirtellum, Michx. Fl. Bor. Am. i. 111 = oxyacanthoides.
hirtum, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 501.—Am. austr.
Hladnickianum, Freyer, ex Steud. Nom. ed. II. ii. 457 = rubrum.
Hladnikii, Reichb. ex Nym. Consp. 267 = alpinum.
holosericeum, *Hort. Berol. ex Otto & Dietr. Allg. Gartenz.* x. (1842) 266.—Hab.?
Hookerianum, A. Murr. Oreg. Circ. 1 = Lobbii?
horridum, *Rupr. ex Maxim. Prim. Fl. Amur.* 117.—Reg. Amur.
hortense, Lam. ex Steud. Nom. ed. I. 691 = rubrum.
hudsonianum, Bourg. ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 336 = prostratum.
hudsonianum, *Richards. in Frankl. 2nd Journ. App.* 6.—Am. bor.
hybridum, Bess. Prim. Fl. Galic. i. 186 = R. Grossularia.
incarnatum, *Wedd. Chlor. And.* ii. 215.—Peruv.
inebrians, Lindl. Bot. Reg. t. 1471 = cereum.
inodorum, Link, Handb. ii. 7 = aureum.
integrifolium, *Phil. in Gartenfl.* (1881) 195.—Chili.
intermedium, Bercht. & Presl, Rostl. i. Grossular. 3 = nigrum.
intermedium, *Tausch, in Flora*, xxi. (1838) 720.—Hab.?
irriguum, Dougl. in Trans. Hort. Soc. vii. (1830) 516 = divaricatum.
irriguum, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 53 = oxyacanthoides.
japonicum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 253.—Japon.
jorullense, H. B. & K. Nov. Gen. et Sp. vi. 61 = ciliatum.
Kunthii, *Berland. in Mém. Soc. Phys. Genèv.* iii. II. (1826) 60.—Mexic.
laciniatum, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 87 = glaciale.
lacustre, *Poir. Encyc. Suppl.* ii. 856.—Am. bor.
lathraeum, Hort. ex C. Koch, Dendrol. i. 641 = rotundifolium.
laxiflorum, *Pursh, Fl. Am. Sept.* ii. 731.—Am. bor.
laxiflorum, Richards. in Frankl. 1st Journ. App. ed. II. 7 = prostratum.
leptanthum, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 53.—Am. bor. occ.
leptostachyum, *Benth. Pl. Hartw.* 186.—N. Granat.
leptostachyum, Decne. in Jacquem. Voy. Bot. 65. t. 76 = orientale.
leucocladon, Reichb. ex C. Koch, Dendrol. i. 657 = alpinum.
Lobbii, *A. Gray, in Am. Natural.* x. (1876) 274.—Calif.
longiflorum, [Nutt.] in Fras. Cat. (1813) n. 74, nomen; et ex Torr. & Gray, Fl. N. Am. i. 552 = aureum.
lucidum, *Kit. in Linnaea*, xxxii. (1863) 481.—Hungar.
luridum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 87—Reg. Himal.

RIBES:—
macracanthum, Lodd. ex G. Don, in Loud. Hort. Brit. 90, nomen (Quid?).—Hab.?
macrobotrys, Hort. ex C. Koch, Dendrol. i. 653 = petraeum.
macrobotrys, *Ruiz & Pav. Fl. Per.* iii. 12.—Peruv.
macrocalyx, Hance, in Journ. Bot. xiii. (1875) 35 = grossularioides.
magellanicum, *Poir. Encyc. Suppl.* ii. 856.—Reg. Magellan.
malvaceum, Sm. in Rees, Cycl. xxx. n. 13 = sanguineum.
melancholicum, Siev. ex Pall. in Nov. Act. Acad. Petrop. x. (1797) 378 = triste.
melanthum, Boiss. & Hohen. Diagn. Ser. I. x. 19 = orientale.
Menziesianum, Roem. & Schult. Syst. v. 507 = seq.
Menziesii, *Pursh, Fl. Am. Sept.* ii. 732.—Am. bor. occ.
mexicanum, Spreng. Syst. iv. Cur. Post. 100 = Kunthii.
Meyeri, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 260.—Mont. Altai.
microphyllum, *H. B. & K. Nov. Gen. et Sp.* vi. 62.—Mexic.
missouriense, Hort. ex C. Koch, Dendrol. i. 662 = americanum.
missouriense, Nutt. ex Torr. & Gray, Fl. N. Am. i. 548 = gracile.
missurense, Lodd. ex Wendl. f. in Bartl. & Wendl. f. Beitr. ii. 17 = aureum.
mogollonicum, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 121.—N. Mexic.
molle, *Poepp. ex Rach, in Ind. Sem. Hort. Petrop.* (1858) 42.—Chili.
montanum, *Phil. in Linnaea*, xxx. (1859–60) 210.—Chili.
multiflorum, H. B. & K. Nov. Gen. et Sp. vi. 60 = Kunthii.
multiflorum, *Kit. in Roem. & Schult. Syst.* v. 493.—Europ. austr.
multiflorum, Willd. Enum. Hort. Berol. Suppl. 12 = alpinum.
nemorosum, *Phil. in Linnaea*, xxviii. (1856) 645.—Chili.
nevadense, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 65 = sanguineum.
nigrum, *Linn. Sp. Pl.* 201.—Europ.; As. bor.
nigrum, Richards. in Frankl. Narr. 1st Journ. App. 734 = hudsonianum.
niveum, Lindl. Bot. Reg. t. 1692 = gracile.
nubigenum, *Phil. in Linnaea*, xxviii. (1856) 646.—Chili.
occidentale, Hook. & Arn. Bot. Beech. Voy. 346 = Menziesii.
odoratum, Schlecht. ex Hemsl. Biol. Centr. Am. Bot. i. 386 = ciliatum.
odoratum, Wendl. f. in Bartl. & Wendl. f. Beitr. ii. 15 = aureum.
odoriferum, *C. Koch, in C. Koch & Fint. Wochenschr.* ii. (1859) 137.—Syria.
officinale, Steud. Nom. ed. I. 691, sphalm. = seq.
officinarum, Dum.-Cours. Bot. Cult. ed. II. v. 303 = rubrum.
olidum, Moench, Meth. 683 = nigrum.
opulifolium, Hort. ex C. Koch, Dendrol. i. 657 = alpinum.
opulifolium, Kit. in Linnaea, xxxii. (1863) 483 = multiflorum.
Oregoni, Hérincq, Hortic. Franç. (1872) t. 8 = aureum.
orientale, *Desf. Hist. Arb.* ii. 88; *Poir. Encyc. Suppl.* ii. 856.—Oriens; Reg. Himal.
Ovallei, *Phil. in Linnaea*, xxxiii. (1864–65) 83.—Chili.
oxyacanthoides, Georgi, Beschr. Russ. Reich. III. iv. 799 = R. Grossularia.
oxyacanthoides, *Linn. Sp. Pl.* 201.—Am. bor.
oxyacanthoides, Michx. Fl. Bor. Am. i. 111 = lacustre.
pallidum, *Otto & Dietr. Allg. Gartenz.* x. (1842) 268.—Hab.?
palmatum, Desf. Cat. Hort. Par. ed. III. 274 = aureum.
parviflorum, *Phil. in Linnaea*, xxviii. (1856) 648.—Chili.
parviflorum, *Wedd. Chlor. And.* ii. 215.—Ecuador.
parvifolium, *Phil. in Linnaea*, xxxiii. (1864–65) 84.—Chili.
pauciflorum, Turcz. ex Ledeb. Fl. Ross. ii. 200 = nigrum.
pendulum, Salisb. Prod. 355 = rubrum.
pensylvanicum, Lam. Encyc. iii. 49 = americanum.

RIBES:—
petiolare, Dougl. in Trans. Hort. Soc. vii. (1830) 514 = hudsonianum.
petraeum, Hort. Prag. ex Tausch, in Flora, xxi. (1838) 720 = urceolatum.
petraeum, Sm. Engl. Bot. t. 705 = rubrum.
petraeum, *Wulf. in Jacq. Misc.* ii. 36.—Europ.; As. bor.
pilosum, Reichb. ex C. Koch, Dendrol. i. 657 = alpinum.
pinetorum, *Greene, in Coult. Bot. Gaz.* v. (1880) 157.—N. Mexic.
pneobalsamum, Siev. in Pall. N. Nord. Beitr. vii. 218 = fragrans.
polyanthes, *Phil. in Linnaea*, xxviii. (1856) 649.—Chili.
polycarpum, J. F. Gmel. Syst. 419 = procumbens.
procumbens, *Pall. Fl. Ross.* ii. 35. t. 65.—Dahur.
propinquum, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 70.—Sibir. or.
prostratum, *L'Hérit. Stirp. Nov.* 3. t. 2.—Am. bor.
pulchellum, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 191.—Sibir.
pumilum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 551 = cereum.
punctatum, Lindl. Bot. Reg. t. 1278 = orientale.
punctatum, *Ruiz & Pav. Fl. Per.* iii. 12. t. 233. f. a.—Chili.
quercetorum, *Greene, in Bull. Calif. Acad.* i. n. III. (1885) 83.—Calif.
reclinatum, Linn. Sp. Pl. 201 = R. Grossularia.
recurvatum, Michx. Fl. Bor. Am. i. 109 = americanum.
reniforme, Nutt. in Journ. Acad. Philad. vii. (1834) 25 = cereum.
resinosum, James, Cat. 177; ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 332 = cereum.
resinosum, *Pursh, Fl. Am. Sept.* i. 163.—Am. bor.
resinosum, Sims, Bot. Mag. t. 1583 = orientale.
rigens, *Hort. ex Regel, Gartenfl.* (1866) 50.—Hab.?
rigens, Michx. Fl. Bor. Am. i. 110 = prostratum.
ringens, Hort. ex C. Koch, Dendrol. i. 654 = petraeum.
Roeseri, Orph. ex Boiss. Fl. Orient. ii. 816 = multiflorum.
Roezli, *Regel, Gartenfl.* (1879) 226.—Am. bor. occ.
rotundifolium, Engelm. in Trans. Am. Phil. Soc. N. S. xii. (1861) 193 = oxyacanthoides?
rotundifolium, *Michx. Fl. Bor. Am.* i. 110.—Am. bor.
rubrum, *Linn. Sp. Pl.* 200.—Europ.; As. et Am. bor.
rupicola, *Phil. in Linnaea*, xxx. (1859–60) 210.—Chili.
sanguineum, *Pursh, Fl. Am. Sept.* i. 164.—Am. bor. occ.
sativum, Syme, Eng. Bot. ed. III. iv. 42 = rubrum.
saxatile, *Pall. in Nov. Act. Acad. Petrop.* x. (1797) 376.—Sibir.
saxosum, Hook. Fl. Bor. Am. i. 231 = oxyacanthoides.
scabrum, Domb. ex Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 25 = viscosum.
Schlechtendalii, Lange, Ind. Sem. Hort. Haun. (1870) 31 = rubrum.
Schmidtianum, Tausch, in Flora, xxi. (1838) 719 = americanum.
Scopolii, Reichb. ex C. Koch, Dendrol. i. 657 = alpinum.
setosum, A. Gray, in Proc. Am. Acad. viii. (1873) 383 = lacustre.
setosum, *Lindl. Bot. Reg.* t. 1237; *et in Trans. Hort. Soc.* vii. (1830) 243.—Am. bor.
sibiricum, Hort. ex C. Koch, Dendrol. i. 649 = rubrum.
speciosum, *Pursh, Fl. Am. Sept.* ii. 731.—Am. bor. occ.
spicatum, Robs. in Trans. Linn. Soc. iii. (1797) 240. t. 21 = rubrum.
spicatum, Schult. Oesterr. Fl. ed. II. i. 432 = multiflorum.
spinosum, Lam. Fl. Fr. iii. 470 = R. Grossularia.
Stakori, *Fisch. ex Sweet, Hort. Brit.* ed. II. 238, nomen.—Sibir.
stamineum, Hornem. Hort. Hafn. 237 = rotundifolium.
stamineum, Sm. in Rees, Cycl. xxx. n. 30 = speciosum.
stenocarpum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 475.—China.
sterile, Hort. ex C. Koch, Dendrol. i. 657 = alpinum.
stramineum, Berland. in Mém. Soc. Phys. Genèv. iii. II. (1856) 58. t. 3, sphalm. = *stamineum*.
suaveolens, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 256 = fragrans.
sublobatum, *Phil. in Linnaea*, xxviii. (1856) 647.—Chili.

RIBES:—
subvestitum, Hook. Bot. Mag. t. 4931 = Lobbii.
subvestitum, Hook. & Arn. Bot. Beech. Voy. 346 = Menziesii.
subvestitum, Hort. ex C. Koch, Dendrol. i. 645 = oxyacanthoides.
sylvestre, Syme, Eng. Bot. ed. III. iv. 43 = rubrum.
Takare, *D. Don, Prod. Fl. Nep.* 208.—Reg. Himal.
tenuiflorum, Lindl. in Trans. Hort. Soc. vii. (1830) 242 = aureum.
tomentosum, C. Koch, in Wochenschr. ii. (1859) 138 = divaricatum.
tortuosum, *Benth. Bot. Voy. Sulph.* 17.—Calif.
tortuosum, Hort. ex C. Koch, Dendrol. i. 657 = alpinum.
trifidum, Hort. ex C. Koch, l. c. 662 = americanum.
trifidum, Michx. Fl. Bor. Am. i. 110 = prostratum.
triflorum, Bigel. Fl. Bost. ed. II. 90 = oxyacanthoides.
triflorum, Hook. & Arn. Bot. Beech. Voy. 346 = divaricatum.
triflorum, Willd. Hort. Berol. t. 61 = rotundifolium.
trilobum, Meyen, Reise, i. 314 = sanguineum.
triste, Bunge, Verz. Suppl. Fl. Alt. 21 = petraeum.
triste, *Pall. in Nov. Act. Acad. Petrop.* x. (1797) 378.—Sibir.
tubiflorum, C. A. Mey. in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 140 = sanguineum.
tubulosum, Eschsch. in Mém. Acad. Pétersb. x. (1821–22) 283 = sanguineum.
urceolatum, *Tausch, in Flora*, xxi. (1838) 720.—Hab.?
Uva-crispa, Linn. Sp. Pl. 201 = R. Grossularia.
valdivianum, *Phil. in Linnaea*, xxviii. (1856) 650.—Chili.
velutinum, *Greene, in Bull. Calif. Acad.* i. n. III. (1885) 83.—Calif.
viburnifolium, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 202.—Calif.
villosum, *C. Gay, Fl. Chil.* iii. 35.—Chili.
villosum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 547 = divaricatum.
villosum, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 514 = orientale.
vinosum, Dum.-Cours. Bot. Cult. ed. II. vii. App. 57 = rubrum.
viridissimum, Reichb. ex C. Koch, Dendrol. i. 657 = alpinum.
viscosissimum, *Pursh, Fl. Am. Sept.* i. 163.—Am. bor. occ.
viscosum, Cels, ex Steud. Nom. ed. I. 691 = resinosum.
viscosum, *Ruiz & Pav. Fl. Per.* iii. 13.—Peruv.
vitifolium, Host, Fl. Austr. i. 308 = multiflorum.
vulgare, C. Koch, Fl. Jen. 36 = R. Grossularia.
vulgare, Lam. Encyc. iii. 47 = rubrum.
Wolfii, Rothr. in Am. Natural. viii. (1874) 358 = sanguineum.
xorullense, Spreng. Syst. iv. Cur. Post. 100 = ciliatum.

RIBESIOIDES, Linn. Fl. Zeyl. 190 (1747) = **Embelia**, Burm. f. (Myrsin.).

RIBESIUM, Medic. Phil. Bot. i. 120 (1789) = **Ribes**, Linn. (Saxifr.).
campanulatum, Medic. Pfl. Anat. 106 = Ribes americanum.
Dillenii, Medic. Phil. Bot. i. 120 = Ribes americanum.
nigrum, Medic. l. c. = Ribes nigrum.
rubrum, Medic. l. c. = Ribes rubrum.

RICARDIA, Adans. Fam. ii. 158 (1763) = RICHARDIA, Houst. = **Richardsonia**, Kunth (Rubiac.).

RICAURTEA, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 47 = **Doliocarpus**, Roland. (Dillen.).
congestiflora, Triana, l. c. 48 = D. Rolandri.
nitida, Triana, l. c. 47 = D. nitida.

RICHAEIA, Thou. Gen. Nov. Madag. 25 (1806) = **Weihea**, Spreng. (Rhizoph.).
madagascariensis, Poir. in Dict. Sc. Nat. xlv. 446 = Cassipourea madagascariensis.

RICHARDIA, Houst. ex Linn. Gen. ed. I. 100 (1737) = **Richardsonia**, Kunth (Rubiac.).
adscendens, Steud. Nom. ed. II. ii. 458 = Richardsonia adscendens.
brasiliensis, Gomez, Mem. sobre Ipecac. 31 (1801) = Richardsonia pilosa.

RICHARDIA :—
divergens, Steud. Nom. ed. II. ii. 459 = Richardsonia grandiflora.
emetica, Schult. f. Syst. vii. 87 = Richardsonia rosea.
grandiflora, Steud. Nom. ed. II. ii. 459 = Richardsonia grandiflora.
Haenkeana, Steud. l. c. = Richardsonia Haenkeana.
humistrata, Steud. l. c. = Richardsonia humistrata.
lateralis, Steud. l. c. = Richardsonia grandiflora.
pilosa, Ruiz & Pav. Fl. Per. iii. 50 = Richardsonia pilosa.
rosea, Schult. f. Syst. vii. 88 = Richardsonia rosea.
scabra, Linn. Sp. Pl. 330 = Richardsonia pilosa.
sparsa, Steud. Nom. ed. II. ii. 459 = Richardsonia grandiflora.
stellaris, Steud. l. c. = Richardsonia stellaris.
villosa, [Moç. & Sesse,] ex DC. Prod. iv. 569 = Richardsonia adscendens.

RICHARDIA, Kunth, in Mém. Mus. Par. iv. (1818) 437. t. 20. *AROIDEAE*, Benth. & Hook. f. iii. 982.
COLOCASIA, Link, Diss. Bot. Suerin (1795) 77; et Handb. i. 267 (1829).
OTOSMA, Rafin. Fl. Tellur. iv. 8. (1836); New Fl. Am. ii. 90 (1836).
ZANTEDESCHIA, Spreng. Syst. iii. 765 (1826).
aethiopica, Spreng. Syst. iii. 765; v. 594, sphalm. = seq.
africana, *Kunth, in Mém. Mus. Par.* iv. (1818) 433.—Afr. austr.
albo-maculata, *Hook. Bot. Mag.* t. 5140.—Afr. austr.
angustiloba, *Schott, in Journ. Bot.* iii. (1865) 35.—Afr. trop.
hastata, *Hook. Bot. Mag.* t. 5176.—Afr. austr.
melanoleuca, *Hook. f. l. c.* t. 5765.—Afr. austr.

RICHARDIA, Lindl. Veg. Kingd. 715 (1847), sphalm. = REICHARDIA, Roth = **Picridium**, Desf. (Compos.).

RICHARDSONIA, Kunth, in Mém. Mus. Par. iv. (1818) 430. *RUBIACEAE*, Benth. & Hook. f. ii. 147.
RICARDIA, Adans. Fam. ii. 158 (1763).
RICHARDIA, Houst. ex Linn. Gen. ed. I. 100 (1737).
SCHIEDEA, Bartl. Ord. 210 (1830).
adscendens, *DC. Prod.* iv. 569.—Mexic.
brasiliensis, Hayne, Arzn. Gew. viii. t. 21 = pilosa.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 31.—Cuba.
divergens, DC. Prod. iv. 568 = grandiflora.
emetica, Mart. Spec. Mat. Med. Bras. 11 = rosea.
grandiflora, *Cham. & Schlecht. in Linnaea*, iii. (1828) 351.—Bras.
Haenkeana, *DC. Prod.* iv. 569.—Mexic.
humistrata, *Cham. & Schlecht. in Linnaea*, iii. (1828) 353.—Bras.
lateralis, DC. Prod. iv. 568 = grandiflora.
muricata, *Griseb. Cat. Pl. Cub.* 143.—Cuba.
pilosa, *H. B. & K. Nov. Gen. et Sp.* iii. 350.—Am. trop.
rosea, *A. St. Hil. Pl. Us. Brés.* n. VII. t. 7.—Bras.
scabra, A. St. Hil. l. c. t. 8; Mart. Spec. Mat. Med. Bras. 10 = pilosa.
sericea, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 354.—Bras.
sparsa, DC. Prod. iv. 568 = grandiflora.
stellaris, *Cham. & Schlecht. in Linnaea*, iii. (1828) 352.—Bras.

RICHEA, R. Br. Prod. 555 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
CYSTANTHE, R. Br. Prod. 555 (1810).
PILITIS, Lindl. Introd. Nat. Syst. ed. II. 443 (1836).
acerosa, *F. Muell. Fragm.* vi. 69.—Tasman.
Desgrassii, *Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. et Zél.* 85—Tasman.
dracophylla, *R. Br. Prod.* 555, partim.—Tasman.
dracophylla, R. Br. l. c., partim = scoparia.
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 273.—Austral.
Milligani, *F. Muell. Fragm.* vi. 69.—Tasman.
pandanifolia, *Hook. f. Fl. Antarct.* i. 50.—Tasman.
procera, *F. Muell. Fragm.* vi. 68.—Tasman.
scoparia, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 273.—Tasman.
sprengelioides, *F. Muell. Fragm.* vi. 68.—Tasman.

RICHEA, Labill. Voy. i. 186. t. 16 (1798) = **Craspedia**, Forst. (Compos.).
glauca, Labill. l. c. t. 16 = C. Richea.

RICHEIA, Steud. Nom. ed. I. 691 (1821) = RICHAEIA, Thou. = **Weihea**, Spreng. (Rhizoph.).

RICHELLA, A. Gray, in Proc. Am. Acad. ii. (1852) 325. *ANONACEAE*, Benth. & Hook. f. i. 26.
monosperma, *A. Gray, l. c.*—Ins. Fiji.

RICHERIA, Vahl, Eclog. Am. i. 30. t. 4 (1796). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 286.
GUARANIA, Baill. Etud. Gén. Euph. 598 (1858).
PODOCALYX, Klotzsch, in Wiegm. Archiv, vii. (1841) 202.
australis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 17.—Bras.
Gardneriana, Baill. ex Muell. Arg. in DC. Prod. xv. II. 468 = grandis.
grandis, *Vahl, Eclog. Am.* i. 30. t. 4.—Am. trop.
laurifolia, Baill. ex Muell. Arg. in DC. Prod. xv. II. 468 = grandis.
longifolia, Baill. ex Muell. Arg. l. c. 467 = grandis.
loranthoides, *Muell. Arg. l. c.* 469.—Am. trop.
purpurascens, Baill. ex Muell. Arg. l. c. 467 = grandis.
Spruceana, Baill. ex Muell. Arg. l. c. 468 = grandis.

RICHIAEA, Benth. & Hook. f. Gen. i. 681 (1867), sphalm. = RICHAEIA, Thou. = **Weihea**, Spreng. (Rhizoph.).

RICHIEA, G. Don, Gen. Syst. i. 276 (1831) = **Ritchiea**, R. Br. (Capparid.).

RICHTERIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 126 = **Chrysanthemum**, Tourn. (Compos.).
pyrethroides, Kar. & Kir. l. c. 127 = C. Richteria.

RICINELLA, Muell. Arg. in Linnaea, xxxiv. (1865–66) 153 = **Adelia**, Linn. (Euphorb.).
barbinervis, Muell. Arg. l. c. 154 = A. barbinervis.
haemiolandra, Muell. Arg. in DC. Prod. xv. II. 730.—Ind. occ.
membranifolia, Muell. Arg. in Mart. Fl. Bras. xi. II. 306.—Bras.
myricaefolia, Muell. Arg. in Linnaea, xxxiv. (1865–66) 154 = Bernardia myricaefolia.
oaxacana, Muell. Arg. l. c. = A. oaxacana.
pedunculosa, Muell. Arg. l. c. 153 = A. pedunculosa.
sylvestris, Muell. Arg. in DC. Prod. xv. II. 730 = A. sylvestris.
triloba, Muell. Arg. in Linnaea, xxxiv. (1865–66) 153 = triloba.

RICINOCARPOS, Desf. in Mém. Mus. Par. iii. (1817) 459. t. 22. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 263.
ECHINOSPHAERA, Sieber, ex Steud. Nom. ed. II. i. 538 (1840).
ROEPERIA, Spreng. Syst. iii. 13, 147 (1826).
Bowmani, *F. Muell. Fragm.* i. 181.—Austral.
cyanescens, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 60.—Austral.
glaucus, *Endl. in Enum. Pl. Hueg.* 18.—Austral.
ledifolius, *F. Muell. Fragm.* i. 76.—Austral.
major, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 59.—Tasman.
marginatus, *Benth. Fl. Austral.* vi. 73.—Austral.
Mitchelli, Sond. in Linnaea, xxviii. (1856) 563 = Bertya Mitchellii.
muricatus, *Muell. Arg. l. c.* xxxiv. (1865–66) 61.—Austral.
pinifolius, *Desf. in Mém. Mus. Par.* iii. (1817) 459. t. 22.—Austral.
psilocladus, *Benth. Fl. Austral.* vi. 71.—Austral.
puberulus, Baill. Etud. Gén. Euph. 344 = Bowmani.
rosmarinifolius, *Benth. Fl. Austral.* vi. 72.—Austral.
sidaefolius, Baill. Etud. Gén. Euph. 344 = pinifolius.
sidaeformis, F. Muell. ex Benth. Fl. Austral. vi. 70 = pinifolius.
speciosus, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Austral.
tasmanicus, Sond. & F. Muell. in Linnaea, xxviii. (1856) 562 = Bertya rosmarinifolia.
trichophorus, *Muell. Arg. l. c.* xxxiv. (1865–66) 60.—Austral.
tuberculatus, *Muell. Arg. l. c.*—Austral.
undulatus, Lehm. Pl. Preiss. ii. 370 = glaucus.
velutinus, *F. Muell. Fragm.* ix. 2.—Austral.

RICINOCARPUS, A. Juss. Euph. Tent. 36 (1824) = praec.

RICINODENDRON, Muell. Arg. in Flora, xlvii. (1864–65) 533. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 297.
africanus, *Muell. Arg. l. c.*—Afr. trop.

RICINOIDES, Tourn. ex Moench, Meth. 286 (1794) = **Chrozophora**, Neck. (Euphorb.).
tinctoria, Moench, l. c. = C. tinctoria.

RICINOPHYLLUM, Pall. ex Ledeb. Fl. Ross. ii. 375 (1844) = **Fatsia**, Decne. & Planch. (Araliac.).
americanum, Pall. ex Ledeb. l. c. = F. horrida.

RICINUS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 295 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 321.
CATAPUTIA, Ludw. Gen. 81 (1760).
africanus, Mill. Gard. Dict. ed. VIII. n. 5 = communis.
americanus, Mill. l. c. n. 2 = Jatropha Curcas.
angulatus, *Thunb. Diss. Ricin.* 6.—Ind. or.
Apelta, Lour. Fl. Cochinch. 585 = Mallotus Apelta.
armatus, Andr. Bot. Rep. t. 430 = communis.
badius, Reichb. Ic. Bot. Exot. ii. 21. t. 154 = communis.
chinensis, Thunb. Diss. Ricin. 5 = communis?
communis, *Linn. Sp. Pl.* 1007.—Reg. trop.
dicoceus, Roxb. Fl. Ind. iii. 690 = Mallotus moluccanus.
digitatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = communis.
dioicus, Cheval. ex Steud. Nom. ed. II. ii. 459 = Mallotus integrifolius.
dioicus, Forst. f. Prod. 67 = Macaranga dioica.
dioicus, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 690.—Ins. Amboin.
dioicus, Wall. Cat. n. 7828 = Coelodiscus montanus.
europaeus, T. F. Nees, Gen. Pl. Monochl. t. 38 = communis.
floribundus, Reinw. ex Muell. Arg. in DC. Prod. xv. II. 962 = Mallotus floribundus.
furfuraceus, Wall. Cat. n. 7805 = Daphniphyllum majus.
Gibsoni.—Cf. Gard. Chron. (1877) II. 692 = communis?
glaucus, Hoffmgg. Verz. Pfl. Nachtr. ii. 199 = communis.
globosus, Willd. Sp. Pl. iv. 567 = Mettenia globosa.
hybridus, Bess. Cat. Hort. Crem. (1816) Suppl. ii. 13 = communis.
inermis, Mill. Gard. Dict. ed. VIII. n. 6; Jacq. Misc. ii. 362 = communis.
inermis, Wall. Cat. n. 7806 = Macaranga hypoleuca.
integrifolius, Willd. Sp. Pl. iv. 567 = Mallotus integrifolius.
japonicus, Thunb. Diss. Ricin. 4 = communis?
Jarak, Thunb. Fl. Javan. 23 = Jatropha Curcas.
Krappa, Hort. Berol. ex Steud. Nom. ed. II. ii. 459 = communis.
laevis, DC. Not. Jard. Genèv. i. 31 = communis.
lanceolatus, Thou. ex Baill. Adansonia, i. (1860–61) 255 = Mallotus integrifolius.
leucocarpus, Bertol. in Giorn. Arcad. xxi. (1824) 192 [bis] = communis.
lividus, Jacq. Misc. ii. 360 = communis.
macrocarpus, Hort. ex Steud. Nom. ed. II. ii. 459 = communis.
macrophyllus, *Bertol. in Giorn. Arcad.* xxi. (1824) 192 [bis].—Hab.?
Mappa, Linn. Amoen. Acad. iv. 124 = Macaranga Mappa.
Mappa, Roxb. Hort. Beng. 69; Fl. Ind. iii. 690 = Macaranga Tanarius.
Mappa, Wall. Cat. n. 7810 = Macaranga Roxburghii.
medicus, Forsk. Fl. Aegypt. Arab. 164 = communis.
medius, J. F. Gmel. Syst. 1615 = praec.
megalosperma, Delile, Cent. Pl. Méroé, 89 = communis.
minor, Mill. Gard. Dict. ed. VIII. n. 7 = communis.
montanus, Wall. Cat. n. 7727 = Baliospermum axillare.
morifolius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 (Quid?).—Java.
nanus, Balb. Cat. Hort. Bot. Taurin. (1813) 65 = communis.
Obermanni, Hort. ex Groenland, in Rev. Hortic. Sér. IV. vii. (1858) 601 = sanguineus.
odoratus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = Commersonia echinata.

RICINUS:—
paniculatus, *Link & Otto, ex Steud. Nom.* ed. II. ii. 459, nomen.—Hab.?
peltatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = communis.
perennis, Hort. ex Steud. Nom. ed. II. ii. 459 = communis.
pictus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = Codiaeum variegatum.
portoricensis, Juss. ex Baill. Etud. Gén. Euph. 314 = Jatropha hernandiaefolia.
pulchellus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 (Quid?).—Java.
purpurascens, Bertol. Misc. Bot. ix. 4 = communis.
rugosus, Mill. Gard. Dict. ed. VIII. n. 4 = communis.
rutilans, Muell. Arg. in DC. Prod. xv. II. 1017, in syn. = communis.
salicinus, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 142 = Homonoia riparia.
sanguineus, *Hort. ex Groenland, in Rev. Hortic.* Sér. IV. vii. (1858) 601.—Hab.?
scaber, Bertol. ex Moris, Stirp. Sard. Elench. 41 = communis.
speciosus, Burm. f. Fl. Ind. 207 (err. typ. 307) = communis.
spectabilis, Blume, Bijdr. 623 = communis.
Tanarius, Linn. Diss. Herb. Amb. in Amoen. Acad. iv. 125 = Macaranga Tanarius.
Tanarius, Lour. Fl. Cochinch. 584 = Mallotus floribundus.
tomentosus, Gaud. in Freyc. Voy. Bot. 489. t. 116 = Adriana tomentosa.
tomentosus, Thunb. Diss. Ricin. 6 (Quid?).—Java.
trilobus, Reinw. ex Blume, Cat. Gew. Buitenz. 108 = Macaranga triloba.
trilobus, Thunb. Diss. Ricin. 6 (= praec.?).—Zeylan.
tunisensis, Desf. Cat. Hort. Par. ed. III. 340 = communis.
undulatus, Bess. Cat. Hort. Bot. Krzem. (1811) 90 = communis.
urens, Mill. Gard. Dict. ed. VIII. n. 3 = communis.
viridis, Willd. Sp. Pl. iv. 564 = communis.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = communis.

RICOILA, Renealm. ex Rafin. Fl. Tellur. iii. 23 (1836) = **Gentiana**, Tourn.
bavarica, Rafin. l. c. = G. bavarica.
pumila, Rafin. l. c. = G. pumila.
verna, Rafin. l. c. = G. verna.

RICOPHORA, Mill. Gard. Dict. ed. VI. 175 App. (1752) = **Dioscorea**, Plum.

RICOTIA, Linn. Sp. Pl. ed. II. 912 (1763). *CRUCIFERAE*, Benth. & Hook. f. i. 72.
aegyptiaca, Linn. l. c. = R. Lunaria.
cantoniensis, *Lour. Fl. Cochinch.* 397.—China.
carnosula, *Boiss. & Heldr. Diagn.* Ser. I. viii. 30.—As. Min.
cretica, *Boiss. & Heldr. l. c.* 29.—Ins. Creta.
Lunaria, *DC. Syst.* ii. 284.—Syria.
multifida, Herb. Banks, ex DC. l. c. 685 = Heliophila diffusa.
Pestalotiana, *Cesati, in Bot. Zeit.* xiv. (1856) 532. t. 9.—As. Min.
pinnata, Moench, Meth. 272 = R. Lunaria.
sinuata, *Boiss. & Heldr. Diagn.* Ser. I. viii. 30.—As. Min.
tenera, Herb. Banks, ex DC. Syst. ii. 685 = Heliophila Peltaria.
tenuifolia, *Sibth. & Sm. Fl. Graec. Prod.* ii. 17.—As. Min.

RIDAN, Adans. Fam. ii. 130 (1763) = **Actinomeris**, Nutt. (Compos.).

RIDDELIA, Rafin. New Fl. Am. iv. 15. (1836) = **Melochia**, Dill. (Stercul.).
antiphyla, Rafin. l. c. 16 = M. hirsuta, *Cav.*

RIDDELLIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 371. *COMPOSITAE*, Benth. & Hook. f. ii. 401.
PSILOSTROPHE, DC. Prod. vii. 261 (1838).
arachnoidea, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 94.—Mexic.
Cooperi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358.—Am. bor. occ.
tagetina, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 371 (Tagetinae).—Am. bor. occ.

RIDELIA, Spach, Hist. Nat. Vég. Phan. ix. 227 (1840), sphalm. = RIEDELIA, Cham. = **Lantana**, Linn. (Verben.).

RIDLEIA, Endl. Gen. 1000 (1840) = RIEDLEA, Vent. = **Melochia**, Dill. (Stercul.).

RIDOLFIA, Moris, Ind. Sem. Hort. Taur. (1841); Fl. Sard. ii. 212. t. 75 (1842) = **Carum**, Rupp. (Umbell.).
segetum, Moris, ll. cc = C. Ridolfia.

RIEDELIA, Cham. in Linnaea, vii. (1832) 240 = **Lantana**, Linn. (Verben.).
lippioides, Cham. l. c. = L. Chamissonis.
officinalis, Lacerda, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 320 = Vandellia diffusa.

RIEDELIA, Meissn. in Mart. Fl. Bras. vii. 171 (1863) = **Satyria**, Klotzsch (Vaccin.).
bahiensis, Mart. l. c. 172 = S. panurensis.

RIEDELIA, Oliver, in Hook. Ic. Pl. t. 1419 (1883). *SCITAMINEAE.*
curviflora, *Oliver, l. c.*—Malaya.

RIEDELIA, Trin. ex Kunth, Enum. Pl. i. 515 (1833) = **Arundinella**, Raddi (Gramin.).
Mikani, Trin. ex Kunth. l. c. = A. brasiliensis.

RIEDLEA, Vent. in Mém. Inst. Sc. Phys. Par. i. (1807) 2 = **Melochia**, Dill. (Stercul.).
Aleurodendron, Steud. Nom. ed. II. ii. 459.—Java.
Berteriana, DC. Prod. i. 492 = Waltheria glabra.
borbonica, DC. l. c. 491 = M. borbonica.
capitata, Boj. Hort. Maurit. 38 = M. corchorifolia.
caracasana, DC. Prod. i. 492 = M. caracassana.
Chamaedrys, G. Don, Gen. Syst. i. 529 = M. Chamaedrys.
concatenata, DC. Prod. i. 492 = M. corchorifolia.
concinna, Walp. Ann. ii. 166 = M. concinna.
corchorifolia, DC. Prod. i. 491 = M. corchorifolia.
cordata, Steud. Nom. ed. II. ii. 459 = M. arborea.
corymbosa, [Moç. & Sesse,] ex DC. Prod. i. 491 = M. corymbosa (in Addend.).
crenata, DC. Prod. i. 491 = M. tomentosa.
cubensis, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 209.—Cuba.
depressa, DC. Prod. i. 491 = M. depressa.
dichotoma, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 208.—Guian.
elongata, Presl, Rel. Haenk. ii. 148.—Calif.
eriantha, DC. Prod. i. 492.—Guian.
graminifolia, Steud. Nom. ed. II. ii. 460 = M. graminifolia.
heterotricha, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 211 = M. hirsuta, *Cav.*
hirsuta, DC. Prod. i. 492 = M. hirsuta, *Cav.*
hypoleuca, Walp. Ann. ii. 166 = M. hypoleuca, *Miq.*
hyptoides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 210.—Bras.
inflata, DC. Prod. i. 491 = M. lupulina.
interrupta, Schlecht. in Linnaea, xi. (1837) 375 = M. interrupta.
jamaicensis, DC. Prod. i. 492 = M. venosa.
Jurgensenii, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 211 = M. Jurgensenii.
littoralis, Poepp. & Endl. Nov. Gen. et Sp. iii. 73.—Bras.
lupulina, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856), 115 = M. lupulina.
melissaefolia, Presl, Rel. Haenk. ii. 149.—Mexic.
mollis, DC. Prod. i. 491 = M. mollis.
multiflora, Presl, Rel. Haenk. ii. 147.—Peruv.
nodiflora, DC. Prod. i. 491 = M. nodiflora.
odorata, DC. l. c. = M. odorata.
peruviana, DC. l. c. = M. peruviana.
polystachya, DC. l. c. 490 = M. venosa.
pteronema, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 208.—Hab.?
quinquealata, Steud. Nom. ed. II. ii. 460 = Columbia floribunda.
radiata, Blume, Bijdr. 86 = M. corchorifolia.
ramosissima, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 571, in obs.—Ind. or.
ramuliflora, Miq. in Linnaea, xix. (1847) 132.—Guian.
scutellarioides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 210 = M. scutellarioides.
sericea, Steud. Nom. ed. II. ii. 460 = M. venosa.

RIEDLEA:—
serrata, Vent. Choix, t. 37 = Melochia hirsuta, *Cav.*
setulosa, Miq. in Hoev. & De Vriese, Tijdschr. x. (1843) 80.—Guian.
siphonandra, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858). I. 208.—Ecuador.
sparsiflora, Klotzsch, ex Walp. Ann. ii. 166.—Guian.
supina, DC. Prod. i. 491 = M. corchorifolia.
tenella, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 212 = M. tenella.
thymifolia, Presl, Rel. Haenk. ii. 149.—Peruv.
tiliaefolia, DC. Prod. i. 491 = M. arborea.
tomentella, Presl, Rel. Haenk. ii. 148 = M. tomentella.
truncata, DC. Prod. i. 491 = M. corchorifolia.
Ulmariae, Steud. Nom. ed. II. ii. 460 = M. ulmarioides.
urticaefolia, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 209 = M. nodiflora.
velutina, DC. Prod. i. 491.—Ind. or.
venosa, DC. Prod. i. 492 = M. venosa.
villosissima, Presl, Rel. Haenk. ii. 146.—Ins. Marian.
virgultosa, Macfad. Fl. Jamaic. i. 104 = M. venosa.
Wallichiana, Steud. Nom. ed. II. ii. 460 = M. arborea.

RIEDLEIA, DC. Prod. i. 490 (1824) = praec.

RIEDLEJA, Hassk. Cat. Hort. Bog. Alt. 205 (1844) = praec.

RIEDLIA, Dum. Anal. Fam. 46 (1829) = praec.

RIENCOURTIA, Cass. in Dict. Sc. Nat. xlv. 466 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 345.
PONTESIA, Vell. Fl. Flum. Ic. viii. t. 147 (1827).
RIENCURTIA, Cass. in Bull. Soc. Philom. (1818) 26.
TETRANTHA, Poit. ex DC. Prod. v. 696, 503 (sphalm. PETRANTHA) (1836).
angustifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 287 = oblongifolia.
glomerata, *Cass. in Dict. Sc. Nat.* xlv. 468.—Guian.
latifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 286.—Bras.
longifolia, *Baker, in Mart. Fl. Bras.* vi. III. 144.—Bras.
oblongifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 287.—Bras.
spiculifera, *Cass. in Dict. Sc. Nat.* xlv. 467.—Am. austr.?
tenuifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 287.—Bras.
tenuifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 145 = oblongifolia.

RIENCURTIA, Cass. in Bull. Soc. Philom. (1818) 26 = praec.

RIESENBACHIA, Presl, Rel. Haenk. ii. 36. t. 54 (1835). *ONAGRARIEAE*, Benth. & Hook. f. i. 792.
racemosa, *Presl, l. c.*—Mexic.

RIESSIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 123; et in Abh. Akad. Berl. (1855) 64. t. 5 C = **Begonia**, Linn.
ferruginea, Klotzsch, l. c. 124 = B. Hookeriana.
pulchella, Klotzsch, l. c. = B. pulchella.

RIGIDELLA, Lindl. Bot. Reg. (1840) t. 16. *IRIDEAE*, Benth. & Hook. f. iii. 691.
flammea, *Lindl. l. c.*—Mexic.
immaculata, *Herb. in Lindl. Bot. Reg.* (1841) *Misc.* 62. t. 68.—Guatemala.
maculata, Walp. Ann. iii. 612, sphalm. = praec.
orthantha, *Lem. in Fl. des Serres*, Sér. I. i. (1845) 107. t. 45.—Mexic.

RIGIOLEPIS, Hook. f. in Hook. Ic. Pl. t. 1160 (1876). *VACCINIACEAE*, Benth. & Hook. f. ii. 572.
borneensis, *Hook. f. in Hook. Ic. Pl.* t. 1160.—Borneo.

RIGIOPAPPUS, A. Gray, in Proc. Am. Acad. vi. (1865) 548. *COMPOSITAE*, Benth. & Hook. f. ii. 406.
leptocladus, *A. Gray, l. c.*—Calif.

RIGIOSTACHYS, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 29. *SIMARUBEAE*, Benth. & Hook. f. i. 309.
RECCHIA, Moç. & Sesse ex DC. Syst. i. 411 (1818).
bracteata, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 30.—Mexic.

RIMA, Sonner. Voy. Nouv. Guin. 99. tt. 57–60 (1776) = **Artocarpus**, Forst. (Urtic.).

RINANTHUS, Gilib. Fl. Lituan. i. 307 (1781) = **Rhinanthus**, Linn. (Scrophul.).

RINDERA, Pall. Reise, i. 486 (1771). *BORAGINEAE*, Benth. & Hook. f. ii. 849.
CYPHOMATTIA, Boiss. Fl. Orient. iv. 272 (1879).
MATTIA, Schult. Obs. Bot. 30, 32 (1809).
angustifolia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 415.—Armenia.
Aucheri, Bunge, l. c. = Paracaryum Aucheri.
caespitosa, *Bunge, l. c.*—As. Min.
canescens, *DC. ex A. DC. in DC. Prod.* x. 168, in obs. —As. Min.; Persia.
Columnae, Roem. & Schult. Syst. iv. 762 = Cynoglossum Columnae.
corymbiformis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 415 = Paracaryum corymbiforme.
corymbulosa, C. Koch, in Linnaea, xxii. (1849) 648 = tetraspis.
cristata, Roem. & Schult. Syst. iv. 761 = Paracaryum cristatum.
cyclodonta, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 415.—Turkest.
echinata, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) i. 92.—Soongar.
emarginata, Roem. & Schult. Syst. iv. 83 = Paracaryum angustifolium.
eriantha, *Ledeb. ex Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 415 = lanata.
glastifolia, Roem. & Schult. Syst. iv. 83 = Paracaryum glastifolium.
glochidiata, Wall. Cat. n. 926 = Paracaryum glochidiatum.
graeca, *Boiss. & Heldr. Diagn.* Ser. I. vii. 30.—Graecia.
incana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 415 = Paracaryum incanum.
laevigata, Roem. & Schult. Syst. iv. 83, 765 = tetraspis.
lanata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 415.—As. Min.; Persia.
lateriflora, Roem. & Schult. Syst. iv. 762 = Pectocarya lateriflora.
leptophylla, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 415 = Paracaryum leptophyllum.
magellensis, Roem. & Schult. Syst. iv. 763 = Cynoglossum magellense.
ochroleuca, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 408.—Sibir. altaic.
pilosa, Roem. & Schult. Syst. iv. 762 = Pectocarya lateriflora.
pubescens, *C. Koch, in Linnaea*, xxii. (1849) 648.—Pontus.
punctata, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 415 = lanata.
tetraspis, L'Hérit. ex DC. & A. DC. in DC. Prod x. 168 = canescens.
tetraspis, *Pall. Reise*, i. 486.—Ross.; Sibir.
umbellata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 415.—Austr.; Sibir.

RINOPODIUM, Salisb. Gen. Pl. Fragm. 28 (1866) = **Scilla**, Linn. (Liliac.).
bifolia, Salisb. l. c. = S. bifolia.

RINOREA, Aubl. Pl. Gui. i. 235. t. 93 (1775) = **Alsodeia**, Thou. (Violar.).
guianensis, Aubl. l. c. = A. paniculata.
integrifolia, Ging. in DC. Prod. i. 313 = A. integrifolia.

RINXOSTYLIS, Rafin. Sylva Tellur. 87 (1838) = **Vitis**, Tourn. (Ampel.).
repanda, Rafin. l. c. = V. repanda.

RINZIA, Schau. in Linnaea, xvii. (1843) 239 = **Baeckea**, Linn. (Myrtac.).
crassifolia, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 331 = B. platystemona.

RINZIA:—
fumana, Schau. in Linnaea, xvii. (1843) 239 = Baeckea fumana.
longifolia, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 331 = B. schelleraefolia.
oxycoccoides, Turcz. l. c. = B. oxycoccoides.

RIOCREUXIA, Decne. in DC. Prod. viii. 640 (1848). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 780.
torulosa, *Decne. l. c.*—Afr. austr.

RIPASIA, Rafin. New Fl. Am. ii. 48 (1836) = **Baptisia**, Vent. (Legumin.).
cerulea, Rafin. l. c. = B. australis.

RIPIDIUM, Trin. Fund. Agrost. 169 (1820) = **Erianthus**, Michx. (Gramin.).
japonicum, Trin. l. c. = Miscanthus sinensis.
Ravennae, Trin. l. c. = E. Ravennae.
strictum, Trin. l. c. = E. Hostii.

RIPOGONUM, Forst. Char. Gen. 49. t. 25 (1776) = **Rhipogonum**, Forst. (Liliac.).

RIPSELAXIS, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
herbacea, Rafin. l. c. = S. herbacea.

RIPSOCTIS, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
egyptiaca, Rafin. l. c. = S. Safsaf?.
indica, Rafin. l. c. = S. tetrasperma.

RIQUERIA, Pers. Syn. i. 152 (1805) = seq.

RIQUEURIA, Ruiz & Pav. Prod. 18 (1794). Gen. dub. *RUBIACEARUM.*
avenia, *Ruiz & Pav. Fl. Per.* i. 70.—Peruv.

RISSOA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 324 = **Atalantia**, Correa (Rutac.).
ceylanica, Arn. l. c. = A. ceylanica.

RITCHIEA, R. Br. in Denh. & Clapp. Voy. App. 225 (1826). *CAPPARIDEAE*, Benth. & Hook. f. i. 110.
erecta, *Hook. f. in Hook. Ic. Pl.* t. 769.—Afr. trop.
fragrans, *R. Br. in Denh. & Claph. Voy. App.* 225.—Afr. trop.
polypetala, *Hook. ex Hook. f. in Bot. Mag.* t. 5344.—Afr. trop.

RITINOPHORA, Neck. Elem. ii. 229 (1790) = **Amyris**, P. Br. (Burser.).

RITTERA, Schreb. Gen. 364 (1789) = **Swartzia**, Schreb. (Legumin.).
dodecandra, Vahl, Symb. Bot. ii. 60 = S. dodecandra.
grandiflora, Vahl, Eclog. Am. ii. 37 = S. grandiflora.
pinnata, Vahl, l. c. 38 = S. pinnata.
simplex, Vahl, Symb. Bot. ii. 60 = S. grandiflora.
triphylla, Sw. ex Steud. Nom. ed. II. ii. 460 = S. triphylla.

RIVEA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 407. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 868.
Argyreia, *Mart. ex Choisy, in DC. Prod.* ix. 327.—Bras.
Bona-nox, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 409. (Conv. Or. 27) = hypocrateriformis.
brasiliana, *Mart. ex Choisy, in DC. Prod.* ix. 326.—Bras.
cordata, *Choisy, l. c.*—Bras.
cuneata, Wight, in Hook. Journ. Bot. iii. (1841) 199 = Argyreia cuneata.
exaltata, *Klotzsch, in Schomb. Fauna et Fl. Brit. Gui.* 1084, nomen.—Guian.
fragrans, Nimmo, in J. Grah. Cat. Pl. Bomb. 127 = hypocrateriformis.
hastata, *Klotzsch, in Schomb. Fauna et Fl. Brit. Gui.* 1152, nomen.—Guian.
hirsuta, Wight, Ic. t. 891 = Argyreia hirsuta.
hypocrateriformis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 408 (*Conv. Or.* 26).—Ind. or.
obtecta, Choisy, l. c. 410 (Conv. Or. 28) = Argyreia obtecta.
ornata, Aitch. Cat. Pl. Punj. 100 = hypocrateriformis.
ornata, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 409 (Conv. Or. 27).—Ind. or.
pomacea, Wight, Ic. t. 888 = Argyreia pomacea.

RIVEA:—
subincana, *Choisy, in DC. Prod.* ix. 325.—Bras.
tiliaefolia, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 407 (Conv. Or. 25) = Argyreia tiliaefolia.
zeylanica, Thw. Enum. Pl. Zeyl. 209 = Argyreia populifolia.

RIVERIA, H. B. & K. Nov. Gen. et Sp. vii. 266. t. 659 (1825) = **Swartzia**, Schreb. (Legumin.).
nitens, H. B. & K. l. c. 267 = S. nitens.

RIVINA, Plum. ex Linn. Syst. ed. I (1735). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 80.
DOBERA?, Rafin. Fl. Tellur. iii. 55 (1836).
PIERCEA, Mill. Gard. Dict. ed. VII. (1759).
RIVINIA, Linn. Hort. Cliff. 35 (1737).
SOLANOIDES, Tourn. ex Moench, Meth. 307 (1794).
TITHONA, Linn. Syst. ed. I. (1735).
acuminata, H. B. & K. Nov. Gen. et Sp. ii. 184 = Mohlana secunda.
acuminata, Rafin. New Fl. Am. iv. 13 = humilis.
affinis, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 30 = Mohlana nemoralis.
apetala, Schum. & Thonn. Beskr. Guin. Pl. 84 = Mohlana guineensis.
aurantiaca, *Warsz. ex Schenk, Ind. Sem. Werceburg.* (1861); *ex Urban, Add. Ind. Sem. Hort. Berol.* (1880) 46.—Am. austr.?
brasiliensis, Nocca, in Usteri, Ann. Bot. vi. (1793) 63 = humilis.
canescens, D. Don, ex Steud. Nom. ed. II. ii. 460 = humilis.
dodecandra, Jacq. Obs. Bot. i. 6 = Villamilla octandra.
Ehrenbergiana, Klotzsch, ex Moq. in DC. Prod. xiii. II. 11 = Villamilla octandra.
glabrata, H. B. & K. Nov. Gen. et Sp. ii. 184 = humilis.
gracilis, Salisb. Prod. 67 = humilis.
humilis, *Linn. Sp. Pl.* 121.—Am. trop.
inaequalis, Hook. Ic. Pl. t. 130 = Mohlana nemoralis.
laevis, Linn. Mant. i. 41 = humilis.
lanceolata, Willd. Enum. Hort. Berol. Suppl. 8 = humilis.
Latbenia, Buch.-Ham. ex Wall. Cat. n. 6959 = humilis?
latifolia, Lam. Illustr. i. 324 = Mohlana latifolia.
mexicana, Moç. & Sesse, ex Moq. in DC. Prod. xiii. II. 14 = humilis?
Moritziana, Klotzsch, ex Moq. l. c. 11 = Villamilla octandra,
Mutisii, Willd. ex Schult. Mart. iii. 305, in obs. = Villamilla octandra.
obliquata, *Rafin. New Fl. Am.* iv. 13.—Am. bor.
octandra, Linn. Cent. Pl. ii. 9 = Villamilla octandra.
orientalis, Moq. in DC. Prod. xiii. II. 12 = humilis.
pallida, Salisb. Prod. 67 = humilis.
paniculata, Linn. Cent. Pl. ii. 9 = Salvadora persica.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1868) 206; *Contrib.* 55.—Paraguay.
peruviana, Moq. in DC. Prod. xiii. II. 10 = Villamilla peruviana.
plumbaginifolia, Willd. ex Moq. l. c. 13 = humilis.
portulaccoides, Nutt. in Trans. Am. Phil. Soc. v. (1837) 167 = humilis.
procumbens, Ruiz, ex Moq. in DC. Prod. xiii. II. 13 = humilis.
puberula, H. B. & K. Nov. Gen. et Sp. ii. 184 = humilis.
purpurascens, Schrad. Gen. Pl. Illustr. 17. t. 5 = humilis.
scandens, Mill. Gard. Dict. ed. VIII. n. 2 = Villamilla octandra.
secunda, Ruiz & Pav. Fl. Per. i. 65 = Mohlana secunda.
seguierioides, Klotzsch, ex Moq. in DC. Prod. xiii. II. 14 = Ledenbergia seguierioides.
tetrandra, Desf. Tabl. ed. II. 49 = humilis.
tinctoria, Ham. ex G. Don, in Loud. Hort. Brit. Suppl. i. 598, nomen (Quid?).—Venezuela.
tinctoria, Moq. in DC. Prod. xiii. II. 10 = Villamilla peruviana.
viridiflora, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 105.—Porto Rico.
viridis, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 185 = humilis.

RIVINIA, Linn. Hort. Cliff. 35 (1737) = praec.

RIXEA, C. Morr. in Ann. Soc. Bot. Gand, i. (1845) 225, t. 22 = **Tropaeolum**, Linn. (Geran.).
azurea, C. Morr. l. c. = T. azureum.

RIXIA, Lindl. Veg. Kingd. 367 (1847) = praec.

RIZOA, Cav. in Anal. Cienc. Nat. iii. (1801) 132 = **Gardoquia**, Ruiz & Pav. (Labiat.).
ovatifolia, Cav. l. c. 133 = G. multiflora.

ROBBAIREA, Boiss. Fl. Orient. i. 735 (1867) = **Polycarpaea**, Lam. (Caryophyll.).
prostrata, Boiss. l. c. = P. prostrata.

ROBBIA, A. DC. in DC. Prod. viii. 444 (1844) = **Malouetia**, DC. (Apocyn.).
cestroides, A. DC. l. c. 445 = M. cestroides.
gossypina, Miers, Apocyn. S. Am. 108.—Bras.
macrocarpa, Miers, l. c. = Echites macrocarpa.

ROBERGIA, Schreb. Gen. 309 (1789) = **Rourea**, Aubl. (Connar.).
frutescens, J. F. Gmel. Syst. 729; Willd. Sp. Pl. ii. 752 = Rourea frutescens.
glabra, Spreng. Syst. iv. Cur. Post. 188 = Rourea glabra.
hirsuta, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 455 = Tapiria hirsuta.

ROBERTA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.

ROBERTIA, DC. Fl. Fr. Suppl. 453 (1815) = **Hypochoeris**, Linn. (Compos.).
pinnatifida, Spreng. Syst. iii. 661 = H. pinnatifida.
taraxacoides, DC. Fl. Fr. Suppl. 453 = H. taraxacoides.
tenuiflora, Boiss. Diagn. Ser. I. xi. 35 = H. tenuiflora.

ROBERTIA, Mérat, Fl. Par. 211 (1812) = **Eranthis**, Salisb. (Ranunc.).
hiemalis, Mérat, l. c. = E. hyemalis.

ROBERTIA, Rich. ex Carr. Conif. ed. I. 498 (1855) = **Phyllocladus**, Rich. (Conif.).

ROBERTIA, Scop. Introd. 154 (1777) = **Bumelia**, Sw. (Sapot.).
decandra, Scop. l. c. = B. lycioides.

ROBERTIUM, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1837) 99 = **Geranium**, Tourn.
anemonaefolium, Picard, l. c. 134 = G. anemonaefolium.
lucidum, Picard, l. c. = G. lucidum.
macrorrhizum, Picard, l. c. = G. macrorrhizum.
minutiflorum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 351 = G. Robertianum.
modestum, Fourr. l. c. = G. Robertianum.
purpureum, Fourr. l. c. = G. Robertianum.
semiglabrum, Fourr. l. c. = G. Robertianum.
simile, Fourr. l. c. = G. Robertianum.
vulgare, Picard, in Mém. Soc. Agric. Boulogne, Sér. II. i. (1837) 134 = G. Robertianum.

ROBERTSIA, Endl. Gen. 740 (1839) = **Sideroxylon**, Linn. (Sapot.).

ROBERTSONIA, Haw. Syn. Pl. Succ. 321 (1812) = **Saxifraga**, Linn.
crenata, Haw. l. c. 322 = S. umbrosa.
cuneifolia, Haw. Saxif. Enum. 56 = S. cuneifolia.
daurica, Haw. l. c. = S. davurica.
dentata, Haw. Syn. Pl. Succ. 323 = S. umbrosa.
Geum, Haw. Saxif. Enum. 53 = S. Geum.
hibernica, Bree, ex Sternb. Rev. Saxif. Suppl. ii. 5 = S. umbrosa.
hieracifolia, Link, Handb. ii. 40 = S. hieracifolia.
hirsuta, Haw. Saxif. Enum. 54 = S. umbrosa.
hybrida, Link, Handb. ii. 39 = S. hybrida.
micranthifolia, Haw. Syn. Pl. Succ. 322 = S. erosa.
nivalis, Link, Handb. ii. 40 = S. nivalis.
pensylvanica, Link, l. c. = S. pensylvanica.
polita, Haw. Syn. Pl. Succ. 322 = S. umbrosa.
punctata, Haw. Saxif. Enum. 55 = S. punctata.
sarmentosa, Link, Handb. ii. 40 = S. sarmentosa.
semipubescens, Link, l. c. = S. pensylvanica.
serrata, Haw. Syn. Pl. Succ. 322 = S. umbrosa.
sphaeroidea, Haw. l. c. = S. umbrosa.
stellaris, Link, Handb. ii. 40 = S. stellaris.
umbrosa, Haw. Saxif. Enum. 55; S. F. Gray, Nat. Arr. Brit. Pl. ii. 529 = S. umbrosa.

ROBESCHIA, Hochst. ex Fourn. Thèse Crucif. 146 (1865) = **Sisymbrium**, Tourn. (Crucif.).
sinaica, Hochst. ex Fourn. l. c. = S. Sophia.

ROBINA, Aubl. Pl. Gui. ii. 768 (1775), err. typ. = seq.

ROBINIA, Linn. Gen. ed. I. 220 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 499.
PSEUDO-ACACIA, [Tourn.] Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 364.
Acacia, Linn. Syst. ed. X. 1161 = R. Pseud-acacia.
aculeata, Juss. ex DC. Prod. ii. 314 = Pictetia Jussiaei.
aculeata, *Vahl, in West, Bidr. Ste Croix*, 300.—Ind. occ.
acuminata, *Schlecht. in Linnaea*, xii. (1838) 306.—Mexic.
alata, Mill. Gard. Dict. ed. VIII. n. 6 = Piscidia Erythrina.
altagana, Pall. Fl. Ross. i. 68 = Caragana microphylla.
amara, *Lour. Fl. Cochinch.* 455.—China; Cochinch.
ambigua, Poir. Encyc. Suppl. iv. 690 = dubia.
amoena, Hort. ex C. Koch, Dendrol. i. 60 = viscosa.
amorphaefolia, Hort. ex C. Koch, l. c. 56 = R. Pseud-acacia.
angulata, Hort. ex C. Koch, l. c. = R. Pseud-acacia.
argentea, Siev. in Pall. N. Nord. Beitr. vii. 326 = Ammodendrum Sieversii.
argentiflora, Schum. & Thonn. Beskr. Guin. Pl. 352 = Lonchocarpus sericeus.
aturensis, *Humb. ex Spreng. Syst.* iii. 246.—N. Granat.
aurea, Hort. ex C. Koch, Dendrol. i. 56 = R. Pseud-acacia.
Bessoniana, Hort. ex C. Koch, l. c. 58 = R. Pseud-acacia.
bullata, Hort. ex C. Koch, l. c. 56 = R. Pseud-acacia.
Camusetii, Leroy, ex C. Koch, l. c. = R. Pseud-acacia.
candida, Roxb. Hort. Beng. 56; Fl. Ind. iii. 327 = Tephrosia candida.
capensis, Burm. f. Fl. Cap. Prod. 22 = Calpurnia robinioides.
Caragana, Linn. Sp. Pl. 722 = Caragana arborescens.
Chamlagu, L'Hérit. Stirp. Nov. 161. t. 77 = Caragana Chamlagu.
chinensis, Pers. Syn. ii. 310 = Caragana Chamlagu.
coccinea, Aubl. Pl. Gui. ii. 773 = Ormosia coccinea.
coluteoides, Hort. ex C. Koch, Dendrol. i. 56 = R. Pseud-acacia.
complexa, Hort. ex C. Koch, l. c. 61 = hispida.
cornigera, Hort. ex C. Koch, l. c. 56 = R. Pseud-acacia.
crispa, Hort. ex C. Koch, l. c. = R. Pseud-acacia.
cubensis, *H. B. & K. Nov. Gen. et Sp.* vi. 392.—Cuba.
cyanescens, Schum. & Thonn. Beskr. Guin. Pl. 351 = Lonchocarpus cyanescens.
cylindrica, Hort. ex C. Koch, Dendrol. i. 56 = R. Pseud-acacia.
davurica, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 304.—Dahuria.
Decaisneana, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
dissecta, Verlot, l. c. = R. Pseud-acacia.
domingensis, Spreng. ex DC. Prod. ii. 168 = Corynella pauciflora.
dubia, Blume, ex Miq. Fl. Ind. Bat. i. 1. 154 = Millettia sericea.
dubia, *Fouc. in Desv. Journ. Bot.* ii. (1813) 204.—Am. bor.
dubia, Poir. Encyc. vi. 227 = Sabinea dubia.
echinata, Mill. Gard. Dict. ed. VIII. n. 2 = R. Pseud-acacia.
edwardsiaefolia, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
Ehrenbergia, *Schlecht. in Linnaea*, xii. (1838) 303.—Mexic.
fastigiata, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
ferox, Pall. Reise, iii. 752 = Caragana spinosa.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* vi. 395.—Venezuela.
ferruginea, Roxb. Hort. Beng. [98]; Fl. Ind. iii. 329 = Derris ferruginea.
filiformis, Roxb. ex Wight & Arn. Prod. 206 = Galactia dubia.
flava, *Lour. Fl. Cochinch.* 556.—China bor.
flexilis, Salisb. Prod. 336 = Caragana microphylla.
florida, Vahl, in West, Bidr. Ste Croix, 300 = Sabinea florida.
fragilis, Salisb. Prod. 336 = R. Pseud-acacia.

ROBINIA :—
frutescens, Linn. Sp. Pl. ed. II. 1044 = Caragana frutescens.
Frutex, Linn. l. c. ed. I. 723 = Caragana arborescens.
fruticosa, Georgi, Reise, ii. 715 = Caragana frutescens.
fruticosa, Roxb. Hort. Beng. 56; Fl. Ind. iii. 328 = Millettia fruticosa.
Galega, Walp. Rep. i. 677 = seq.
Galuga, Roxb. ex Wight & Arn. Prod. 211 = Millettia racemosa.
Garagana, Crantz, Inst. ii. 60, sphalm. = *R. Caragana*.
glabra, Mill. Gard. Dict. ed. VIII. n. 5 = Dalbergia campechiana.
glabrescens, Hoffmgg. Verz. Pfl. Nachtr. ii. 57 = hispida.
glandulosa, Spreng. Syst. iii. 248 = Poiretia psoraloides.
glutinosa, Sims, Bot. Mag. t. 560 = viscosa.
glyciphylla, *Poir. Encyc.* vi. 227.—Ins. Martinic.
grandiflora, Bieb. Fl. Taur. Cauc. ii. 168 = Caragana grandiflora.
grandiflora, Linn. Sp. Pl. 722 = Sesbania grandiflora.
guineensis, *Willd. Enum. Hort. Berol.* 769.—Afr. trop.
Halimodendron, Link, Handb. ii. 201 = seq.
Halodendrum, Pall. Reise, ii. 741 = Halimodendrum argenteum.
Heynei, Wall. Cat. n. 5653 = Calpurnia lasiogyne.
hirsuta, Lindem. in Bull. Soc. Nat. Mosc. lv. (1880) 1. 296 = hispida?
hispida, *Linn. Mant.* i. 101.—Am. bor.
hybrida, Audib. ex DC. Prod. ii. 262 = dubia.
inermis, Jacq. Select. Am. 210 = R. Pseud-acacia.
intermedia, Soulange, in Ann. Soc. Hortic. Par. ii. (1828) 42 = viscosa.
jaspidea, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
javanica, Burm. f. Fl. Ind. 163 = Cassia glauca.
javanica, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 291 = Tephrosia candida.
jubata, Pall. in Nov. Act. Acad. Petrop. x. (1797) 370. t. 6 = Caragana jubata.
latifolia, Mill. Gard. Dict. ed. VIII. n. 9 = Lonchocarpus violaceus.
latifolia, Poir. Encyc. vi. 224 = Poitea Campanilla.
latisiliqua, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
leucantha, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 291 = Tephrosia candida.
linearis, Hort. ex C. Koch, Dendrol. i. 56 = R. Pseud-acacia.
lucida, Salisb. Prod. 337 = Caragana Chamlagu.
ludoviciana, Rafin. Fl. Ludov. 164 (Quid?).—Am. bor.
macrophylla, Roxb. Hort. Beng. 56; Fl. Ind. iii. 329 = Millettia auriculata.
macrophylla, Schrad. ex G. Don, Gen. Syst. ii. 238 = hispida.
macrophylla, *Spreng. Syst.* iii. 246.—Bras.
maculata, *H. B. & K. Nov. Gen. et Sp.* vi. 393.—Mexic.
martinicensis, Pers. Syn. ii. 312 = Sabinea dubia.
melanocarpa, *Schlecht. in Linnaea*, xii. (1838) 305.—Mexic.
microphylla, Hort. ex Loud. Hort. Brit. Suppl. ii. 667 = R. Pseud-acacia.
microphylla, Pall. Astragal. 116 = Caragana microphylla.
mitis, Linn. Sp. Pl. ed. II. 1044 = Pongamia glabra.
mitis, Lour. Fl. Cochinch. 455 = Pongamia chinensis.
mollis, Bieb. Fl. Taur. Cauc. iii. 477 = Caragana frutescens.
mollis, Vahl, ex DC. Prod. ii. 260 = Lonchocarpus sericeus.
mollissima, Vahl, ex Poir. Encyc. vi. 227 = Cupania Poiretii.
mongolica, Sw. ex Bess. Cat. Hort. Crem. (1816) 116 = Caragana arborescens.
monophyllus, Hort. ex C. Koch, Dendrol. i. 56 = R. Pseud-acacia.
montana, Bartr. Cat. ex Pursh, Fl. Am. Sept. ii. 488 = hispida.
multiflora, Schum. & Thonn. Beskr. Guin. Pl. 350 (Millettiae sp.?).—Afr. trop.
nana, Ell. Sketch, ii. 243 = hispida.
neo-mexicana, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 314.—Mexic.
Nicou, Aubl. Pl. Gui. ii. 771 = Lonchocarpus rufescens.
Panacoco, Aubl. l. c. 768 = Swartzia tomentosa.
patula, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
pendula, Orteg. Hort. Matr. Dec. 26 = R. Pseud-acacia.
pinnata, Hort. ex Steud. Nom. ed. I. 693; ed. II. ii. 461 = R. Pseud-acacia?

ROBINIA:—
polyanthos, Sw. Fl. Ind. Occ. iii. 1260 = Corynella polyantha.
procera, Lodd. Cat. ex G. Don, in Loud. Hort. Brit. 304 = R. Pseud-acacia.
Pseud-acacia, *Linn. Sp. Pl.* 722.—Am. bor.
purpurea, *Hort. ex Link, Enum. Hort. Berol.* ii. 242.—Am. bor.
pygmaea, Linn. Sp. Pl. 723 = Caragana pygmaea.
pyramidalis, Verlot, in Rev. Hortic. (1873) 155 = R. Pseud-acacia.
pyramidata, *Mill. Gard. Dict.* ed. VIII. n. 7.—Am. centr.
racemosa, Roxb. Hort. Beng. 56; Fl. Ind. iii. 329 = Millettia racemosa.
radicans, Sw. ex Steud. Nom. ed. I. 693 (Quid?).—Hab.?
rosea, Loisel. in Duham. Arb. ed. Nov. t. 17 = hispida.
rosea, Mill. Gard. Dict. ed. VIII. n. 4.—Am. bor.
rubiginosa, Banks, ex Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 92 = Lonchocarpus rubiginosus.
rubiginosa, *Nees, ex Mart. in Nov. Act. Nat. Cur.* xii. 1. (1824) 31.—Bras.
rubiginosa, Poir. Encyc. vi. 227 = Cupania Poiretii.
scandens, Willd. Sp. Pl. iii. 1134 = Lonchocarpus Nicou.
Schiedeana, *Schlecht. in Linnaea*, xii. (1838) 306.—Mexic.
sennoides, Roxb. Hort. Beng. [98]; Fl. Ind iii. 328 = Mundulea suberosa.
sepium, Jacq. Enum. Pl. Carib. 28 = Lonchocarpus sepium.
sepium, Sw. Fl. Ind. Occ. iii. 1258 = Lonchocarpus violaceus.
sericea, Poir. Encyc. vi. 226 = Lonchocarpus sericeus.
sericea, Sieber, ex Presl, Symb. Bot. i. 14 = Tephrosia candida.
sophoraefolia, Hort. Vindob. ex Steud. Nom. ed. I. 693 = R. Pseud-acacia.
speciosa, Sw. ex Bess. Cat. Hort. Crem. (1816) 116 = Caragana arborescens.
spectabilis, Dum.-Cours. Bot. Cult ed. II. vi. 140 = R. Pseud-acacia.
spinifolia, Desv. Journ. Bot. (1814) 78 = Pictetia Desvauxii.
spinosa, Linn. Mant. ii. 269 = Caragana spinosa.
spinosissima, Laxm. in Nov. Comm. Acad. Petrop. xv. (1771) 558 = Caragana spinosa.
squamata, Hort. Berol. (1844) ex Walp Ann. i. 231 = Brongniartia robinioides.
squamata, Vahl, Symb. Bot. iii. 88 = Ormocarpum sennoides.
squamosa, Poir. Encyc. vi. 224 = Ormocarpum sennoides.
striata, Willd. Sp. Pl. iii. 1132 = Geoffraea superba.
stricta, Hoffmgg. Verz. Pfl. 194 = R. Pseud-acacia.
subdecandra, L'Hérit. Stirp. Nov. 157. t. 75 = Calpurnia lasiogyne.
suberosa, Roxb. Hort. Beng. 56; Fl. Ind. iii. 327 = Mundulea suberosa.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 349 = Millettia Thonningii.
tomentosa, Fisch. Cat. Hort. Gorenk. (1812) 71 = Caragana frutescens.
tomentosa, Willd. Sp. Pl. iii. 1134 = Swartzia tomentosa.
tortuosa, Hoffmgg. Verz. Pfl. 193 = R. Pseud-acacia.
tragacanthoides, Pall. in Nov. Act. Acad. Petrop. x. (1797) 371. t. 7 = Caragana tragacanthoides.
triflora, L'Hérit. Stirp. Nov. 162 = Halimodendron argenteum.
uliginosa, Willd. Sp. Pl. iii. 1133 = Derris uliginosa.
umbraculifera, Loud. Hort. Brit. Suppl. ii. 667 = R. Pseud-acacia.
undulata, Hort. ex C Koch, Dendrol. i. 56 = R. Pseud-acacia.
Uterhartii, Poit. in Rev. Hortic. Sér. II. ii. (1843–44) 447 = R. Pseud-acacia.
variegata, *Schlecht. in Linnaea*, xii. (1838) 301.—Mexico.
vesicaria, Jacq. Coll. i. 105 = Sesbania platycarpa.
viciaefolia, Bert. ex DC. Prod. ii. 263 = Poitea viciaefolia?
violacea, Beauv. Fl. Owar. ii. 28. t. 76 = Lonchocarpus sericeus.
violacea, Jacq. Enum. Pl. Carib. 28 = Lonchocarpus violaceus.
viscosa, *Vent. Jard. Cels*, t. 4.—Am. bor.

ROBINSONIA, DC. in Guill. Arch. Bot. ii. (1833) 333. *COMPOSITAE*, Benth. & Hook. f. ii. 441.
corrugata, Phil. in Bot. Zeit. xiv. (1856) 645 = Gayana.
evenia, *Phil. l. c.* 644.—Ins. J. Fernand.
Gayana, *Decne. in Ann. Sc. Nat.* Sér. II. i. (1834) 28. t. 1.—Ins. J. Fernand.
gracilis, *Decne. l. c.* 29—Ins. J. Fernand.
longifolia, *Phil. in Bot. Zeit.* xiv. (1856) 644.—Ins. J. Fernand.
macrocephala, *Decne. in Ann. Sc. Nat.* Sér. II. i. (1834) 28. t. 1.—Chili; Ins. J. Fernand.
micrantha, Phil. ex Hemsl. Bot. Voy. Challenger, i. III. 40 = gracilis.
nervosa, Phil. in Bot. Zeit. xiv. (1856) 645 = Plantago fernandezia.
thurifera, *Decne. in Ann. Sc. Nat.* Sér. II. i. (1834) 28.—Ins. J. Fernand.

ROBINSONIA, Scop. Introd. 218 (1777) = **Quiina** Aubl. (Guttif.).
guianensis, J. F. Gmel. Syst. 796 = Q. guianensis.
melianthifolia, Willd. Sp. Pl. ii. 999 = Q. guianensis.

ROBIQUETIA, Gaudich. in Freyc. Voy. Bot. 426. t. 34 (1826) = **Saccolabium**, Blume (Orchid.).
ascendens, Gaudich. l. c. = S. ascendens.

ROBSONIA, Reichb. Handb. 233 (1837) = **Ribes**, Linn. (Saxifr.).
speciosa, Spach, Hist. Vég. Phan. vi. 181 = Ribes speciosum.

ROBYNSIA, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 193 = **Pachyrhizus**, Rich. (Legumin.).
lobata, Mart. & Gal. l. c. 194 = P. palmatilobus.
macrophylla, Mart. & Gal. l. c. 193 = P. angulatus.

ROCAMA, Forsk. Fl. Aegypt. Arab. p. cviii. 71 (1775) = **Trianthema** Linn. (Ficoid.).
arabica, J. F. Gmel. Syst. 455 = T. pentandra.
digyna, DC. Prod. iii. 352 = T. pentandra.
prostrata, Forsk. Fl. Aegypt. Arab. p. cviii = T. pentandra.

ROCCARDIA, Neck. Elem. i. 74 (1790) = **Helipterum**, DC. (Compos.).

ROCCARDIA, Rafin. Fl. Tellur. iv. 119 (1836) = **Staehelina**, DC. (Compos.).
purpurea, Rafin. l. c. = S. dubia.

ROCHEA, DC. Pl. Grass. t. 103 (1810?). *CRASSULACEAE*, Benth. & Hook. f. i. 658.
DIETRICHIA, Tratt. Arch. Gew. n. 449 (1812).
KALOSANTHES, Haw. Rev. Succ. 6 (1821).
albiflora, DC. Prod. iii. 393 = Crassula albiflora.
bicolor, Steud. Nom. ed. II. ii. 462 = odoratissima.
biconvexa, *DC. Prod.* iii. 394.—Afr. austr.
Burmanniana, Steud. Nom. ed. II. ii. 462 = Crassula Burmanniana.
coccinea, *DC. Pl. Grass.* t. 1.—Afr. austr.
cymosa, DC. Prod. iii. 394 = Crassula cymosa.
dejecta, Steud. Nom. ed. II. ii. 462 = Crassula dejecta.
dichotoma, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 309 = Crassula abyssinica.
falcata, DC. Pl. Grass. t. 103 = Crassula falcata.
fascicularis, *DC. Prod.* iii. 394.—Afr. austr.
flava, DC. l. c. = Crassula flava.
jasminea, *DC. l. c.*—Afr. austr.
media, DC. l. c. = versicolor.
microphylla, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 369 = jasminea.
odoratissima, *DC. Prod.* iii. 394.—Afr. austr.
perfoliata, DC. l. c. 393 = Crassula perfoliata.
tiniflora, *Lem. Illustr.* Hortic. viii. (1861) sub t. 269.—Afr. austr.
undosa, *Steud. Nom. ed.* II. ii. 462.—Afr. austr.
undulata, Steud. l. c. = Crassula undulata.
vaginata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 309 = Crassula abyssinica.
versicolor, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. ii. 369 = odoratissima.
versicolor, *Link, Enum. Hort. Berol.* i. 299; *DC. Prod.* iii. 394.—Afr. austr.

ROCHEA, Salisb. in Trans. Hort. Soc. i. (1812) 322 = **Geissorhiza**, Ker-Gawl. (Irid.).
venusta, Salisb. l. c. = G. Rochensis.

ROCHEA, Scop. Introd. 296 (1777) = **Aeschynomene**, Linn. (Legumin.).

ROCHEFORTIA, Sw. Prod. Veg. Ind. Occ. 53 (1788); Fl. Ind. Occ. i. 551. t. 11 (1797). *BORAGINEAE*, Benth. & Hook. f. ii. 841.
DIPLOSTYLIS, Karst. & Triana, in Linnaea, xxviii. (1856) 433.
acanthophora, *Griseb. Fl. Brit. W. Ind.* 482.—Ind. occ.
brasiliensis, *Willd. ex Schult. Syst.* vi. 210.—Bras.
cuneata, *Sw. Prod. Veg. Ind. Occ.* 54.—Jamaic.
Jacquini, Griseb. in Goett. Abh. vii. (1857) 253 = Ehretia spinosa.
ovata, *Sw. Prod. Veg. Ind. Occ.* 54.—Jamaic.

ROCHELIA, Reichb. in Flora, vii. (1824) 243. *BORAGINEAE*, Benth. & Hook. f. ii. 852.
MACCOYA, F. Muell. Fragm. i. 127 (1859).
RACLATHRIS, Rafin. Sylva Tellur. 167 (1838).
cancellata, *Boiss. & Bal. Diagn.* Ser. II. iii. 136.—Cilicia.
cardiosepala, *Bunge, in Mém Sav. Etr. Pétersb.* vii. (1851) 420.—Turkest.
glomerata, Torr. in Ann. Lyc. N. York, ii. (1828) 226 = Krynitzkia glomerata.
incana, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xiv. (1841) 710.—Turkest.
leiocarpa, *Ledeb. Fl. Alt.* i. 172.—Turkest.
leiosperma, Steud. Nom. ed II. ii. 56, in syn., sphalm. = praec.
Maccoya, *F. Muell. ex Benth. Fl. Austral.* iv. 408.—Austral.
macrocalyx, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 419.—Persia; Afghan.
mollis, *Bunge, ex Boiss. Fl. Orient.* iv. 245.—Persia.
peduncularis, *Boiss. Diagn.* Ser. I. vii. 35.—Persia.
persica, *Bunge, ex Boiss. Fl. Orient.* iv. 244.—Persia.
rectipes, Stocks, in Hook. Kew Journ. iv. (1852) 176 = macrocalyx.
saccharata, Reichb. in Flora, vii. (1824) 243 = stellulata.
stellulata, Aitch. in Journ. Linn. Soc. xix. (1882) 178 = stylaris.
stellulata, *Reichb. Ic. Fl. Germ.* ii. 13. t. 123.—Europ.
stylaris, *Boiss. Fl. Orient.* iv. 245.—Afghan.

ROCHELIA, Roem. & Schult. Syst. iv. 11 (1819) = **Echinospermum**, Sw. (Borag.).
asperata, Boj. ex DC. Prod. x. 152, in nota = Cynoglossum Rochelia.
barbata, Roem. & Schult. Syst. iv. 110 = E. barbatum.
Borboniae, Roem. & Schult. l. c. 108, 781 = Cynoglossum Rochelia.
condylophora, Roem. & Schult. l. c. 782 = E. condylophorum.
cynoglossoides, Roem. & Schult. l. c. 111, 783 = E. cynoglossoides.
deflexa, Roem. & Schult. l. c. 109 = deflexum.
echinophora, Roem. & Schult. l. c. 110 = E. minimum.
gracilis, Roem. & Schult. l. c. 111 = Pectocarya linearis.
javanica, Roem. & Schult. l. c. 108, 780 = E. javanicum.
Lappula, Roem. & Schult. l. c. 109 = E. Lappula.
marginata, Roem. & Schult. l. c. 110 = E. marginatum.
patens, Nutt. in Journ. Acad. Philad. vii. (1834) 44 = E. patens.
patula, Roem. & Schult. Syst. iv. 781 = E. patulum.
Redowskii, Roem. & Schult. l. c. 110, 782 = E. Redowskii.
spinocarpos, Roem. & Schult. l. c. 111, 783 = E Vahlianum.
squarrosa, Roem. & Schult. l. c. 109 = E. patulum.
strigosa, Roem. & Schult. l. c. 782 = Cynoglossum borbonicum.
virginica, Roem. & Schult. l. c. 108 = E. virginicum.
zeylanica, Roem. & Schult. l. c. 108, 780 = Cynoglossum denticulatum.

ROCHETIA, Delile, in Rochet, Sec. Voy. Choa, 344 (1846) = **Trichilia**, P. Br. (Meliac.).
choënsis, Delile, l. c. = T. emetica.

ROCHONIA, DC. Prod. v. 345 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 259.
BREPHOCTON, Rafin. Fl. Tellur. ii. 49 (1836).
cinerarioides *DC. Prod.* v. 345.—Madag.
cuneata, *DC. l. c.*—Madag.

RODETIA, Moq. in DC. Prod. xiii. II. 323 (1849) = **Bosea**, Linn. (Amarant.).
Amherstiana, Moq. l. c. = B. Amherstiana.

RODGERSIA, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 389. *SAXIFRAGEAE*, Benth. & Hook. f. i. 635.
japonica, Regel, Gartenfl. (1871) t. 708 = seq.
podophylla, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 389.—Japon.

RODIGIA, Spreng. Neue Entdeck. i. 275 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 511.
PACHYLEPIS, Less. Syn. Comp. 139 (1832).
SCLEROLEPIS, Monn. Ess. Hierac. 81 (1829).
alliata, G. Don, ex Loud. Hort. Brit. 328 = Hypochoeris Achyrophorus.
commutata, *Spreng. Neue Entdeck.* i. 275.—Graecia.
interrupta, Sch. Bip. Cich. Suppl. ii. 152 = commutata.
laevigata, Spreng. Neue Entdeck. i. 276 = Hypochoeris Achyrophorus.

RODRIGUEZIA, Ruiz & Pav. Prod. 115. t. 25 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 559.
BURLINGTONIA, Lindl. Bot. Reg. t. 1927 (1837).
PHYSANTHERA, Bert. ex Steud. Nom. ed. II. ii. 330, 463 (1841).
bahiensis, *Reichb. f. in Bonplandia*, ii. (1854) 90.—Bras.
Barkeri, Hook. Bot. Mag. t. 2746 (Gomesae sp.).—Bras.
Batemani, *Lindl. in Poepp. & Endl. Nov. Ge Sp.* i. 41. t. 70.—Peruv.
bifolia, *Rodrig. Orch. Nov.* ii. 239.—Bras.
brachystachys, *Reichb. f. & Warm. in Otia Bot. Hamb.* 87.—Bras.
caloplectron, *Reichb. f. in Gard. Chron.* (1883) I. 368.—N. Granat.
candida, *Batem. in Bot. Reg.* (1837) sub t. 1927.—Guian.
carnea, *Lindl. l. c.* (1843) *Misc.* 72.—N. Granat.
cochlearis, Lindl. l. c. (1842) Misc. 22 = Leochilus cochlearis.
crispa, Lindl. l. c. (1839) Misc. 86 = Gomesa crispa.
decora, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.
eleutherosepala, *Rodrig. Orch. Nov.* ii. 240.—Bras.
ensifolia, Reichb. f. in Walp. Ann. vi. 695, sphalm. = seq.
ensiformis, *Ruiz & Pav. Syst. Veg.* 218.—Peruv.
fragrans, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.
granadensis, *Reichb. f. l. c.*—N. Granat.
lanceolata, Lodd. Bot. Cab. t. 676 = secunda.
lanceolata, *Ruiz & Pav. Syst. Veg.* 219.—Peruv.
laxiflora, Lindl. Bot. Reg. (1837) Misc. 86 = Gomesa chrysostoma.
Leeana, *Reichb. f. in Gard. Chron.* (1883) II. 38.—Hab.?
Lehmanni, *Reichb. f. l. c.* (1883) I. 403.—N. Granat.
leochilina, *Reichb. f. l. c.* (1871) 970.—Costa Rica.
leucantha, *Rodrig. Orch. Nov.* i. 146.—Bras.
luteola, *N. E. Br. in Gard. Chron.* (1883) I. 688.—Hab.?
macrostachya, Hort. Petrop. ex Regel, Ind. Sem. Hort. Petrop. (1856) 21 = Gomesa Fischeri.
maculata, Lindl. Bot. Reg. (1840) Misc. 89 = Leochilus oncidioides.
maculata, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.
mascarenensis, *Spreng. Syst.* iii. 719.—Ins. Mascar.
mascarensis, Steud. Nom. ed. II. ii. 463, sphalm. = praec.
microphyta, *Rodrig. Rev. de Engenh.* III. 144; *ex ej. Orch. Nov.* ii. 241.—Bras.
obtusifolia, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.
pardina, *Reichb. f. l. c.* xx. (1862) 428.—Bras.
planifolia, Lindl. in Trans. Hort. Soc. vii. (1830) 67 = Gomesa planifolia.
pubescens, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.
recurva, Lindl. in Trans. Hort. Soc. vii. (1830) 67 = Gomesa recurva.
refracta, *Reichb. f. in Bonplandia*, ii. (1854) 90, in syn.—Bras.
rigida, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.

RODRIGUEZIA:—
secunda, *H. B. & K. Nov. Gen. et Sp.* i. 367. t. 92.—Am. trop.
Stangeana, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 314.—Hab.?
stenochila, Lindl. Orchid. Linden. 23 = Scelochilus stenochila.
sticta, *Steud. Nom.* ed. II. ii. 463.—Jamaic.
suaveolens, Lindl. Gen. et Sp. Orch. 195 = Gomesa foliosa.
venusta, *Reichb. f. in Bot. Zeit.* x. (1852) 771.—Bras.

RODSCHIEDIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Croton**? Linn. (Euphorb.).
serrata, Dennst. l. c.—Ind. or.

RODSCHIEDIA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 413 (1800) = **Capsella**, Medic. (Crucif.).
Bursa-pastoris, Gaertn. Mey. & Scherb. l. c. 435 = C. Bursa-pastoris.

RODSCHIEDIA, Miq. in Linnaea, xviii. (1844) 585 = **Securidaca**, Linn. (Polygal.).
surinamensis, Miq. l. c.—Guian.

ROËA, Hueg. ex Benth. in Enum. Pl. Hueg. 34 (1837) = **Sphaerolobium**, Sm. (Legumin.).
linophylla, Hueg. l. c. = S. linophyllum.
nudiflora, Meissn. in Lehm. Pl. Preiss. i. 59 = S. nudiflorum.

ROEBELIA, Engel, in Linnaea, xxxiii. (1864–65) 680. t. 3. f. 5. *PALMAE*, Benth. & Hook. f. iii. 883.
solitaria, *Engel, l. c.*—N. Granat.

ROEGNERIA, C. Koch, in Linnaea, xxi. (1848) 413 = **Agropyron**, J. Gaertn. (Gramin.).
caucasica, C. Koch, l. c. = A. Raegnerii.

ROEHLINGIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Tetracera**, Linn. (Dillen.).
suaveolens, Dennst. l. c. = T. laevis.

ROELA, Scop. Introd. 149 (1777), sphalm. = **Roëlla**, Linn. (Campan.).

ROELANA, Comm. ex DC. Prod. i. 575 (1824) = **Erythroxylum**, P. Br. (Lineae).
laurifolia, Comm. ex DC. l. c. = E. laurifolium.

ROËLLA, Linn. Coroll. Gen. 4 (1737). *CAMPANULACEAE*, Benth. & Hook. f. ii. 558.
angustifolia, Roxb. in Beats. Tracts St. Helena, 320 = Wahlenbergia clivosa.
bracteata, A. DC. Monog. Camp. 179 = spicata.
bryoides, *Buek, in Eckl. & Zeyh. Enum.* 386.—Afr. austr.
caespitosa, Banks, ex A. DC. in DC. Prod. vii. 447 = muscosa.
campestris, A. DC. l. c. = spicata.
cernua, Brouss. Elench. Hort. Monsp. (1805) 49 = Wahlenbergia lobelioides.
ciliata, Drège, ex A. DC. in DC. Prod. vii. 446 = Merciera brevifolia.
ciliata, *Linn. Sp. Pl.* 170.—Afr. austr.
cinerea, A. DC. Monog. Camp. 175 = Lightfootia grisea.
compacta, Thunb. ex A. DC. in DC. Prod. vii. 448, 496 = Merciera tenuifolia.
decumbens, Loud. Hort. Brit. 77 = Wahlenbergia capensis.
decurrens, Andr. Bot. Rep. t. 238 = Wahlenbergia capensis.
decurrens, *L'Hérit. Sert. Angl.* 4.—Afr. austr.
Dregeana, A. DC. in DC. Prod. vii. 446 = ciliata.
Dunantii, A. DC. Monog. Camp. 175 = reticulata.
Eckloniana, *Buek, in Eckl. & Zeyh. Enum.* 385.—Afr. austr.
elegans, *Paxt. Mag. Bot.* vi. (1839) 27.—Afr. austr.
erecta, Banks, ex A. DC. in DC. Prod. vii. 444 = Prismatocarpus altiflorus.
ericoides, Buek, in Eckl. & Zeyh. Enum. 385 = spicata.
ericoides, Spreng. Syst. i. 723 = Prismatocarpus paniculatus.
filiformis, Lam. Illustr. ii. 66 = squarrosa.
filiformis, Poir. Encyc. vi. 232 = Lightfootia fasciculata.

ROËLLA:—
fruticosa, L'Hérit. ex A. DC. in DC. Prod. vii. 444 = Prismatocarpus fruticosus.
glabra, Poir. Encyc. vi. 232 = Lightfootia fasciculata.
glabrella, Banks, ex A. DC. in DC. Prod. vii. 446 = Lightfootia grisea.
glauca, Buek, in Eckl. & Zeyh. Enum. 387 = spicata.
glomerata, *A. DC. in DC. Prod.* vii. 447.—Afr. austr.
gracilis, Eckl. & Zeyh. Enum. 385 = ciliata.
incurva, Banks, ex A. DC. Monog. Camp. 172 = ciliata.
interrupta, Banks, ex A. DC. in DC. Prod. vii. 443 = Prismatocarpus interruptus.
lateriflora, Banks, ex A. DC. l. c. 496 = Merciera brevifolia.
latiloba, *A. DC. l. c.* 447.—Afr. austr.
leptosepala, *Sond. in Harv. & Sond. Fl. Cap.* iii. 593.—Afr. austr.
linifolia, Roxb. in Beats. Tracts St. Helena, 321 = Wahlenbergia linifolia.
minor, Buek, in Eckl. & Zeyh. Enum. 385 = ciliata.
muscosa, *Linn. f. Suppl.* 143.—Afr. austr.
nitida, Banks, ex A. DC. in DC. Prod. vii. 445 = Prismatocarpus nitidus.
paniculata, Roxb. in Beats. Tracts St. Helena, 320 = Wahlenbergia Roxburghii.
pedunculata, Berg. Desc. Pl. Cap. 42 = Prismatocarpus paniculatus.
pinifolia, Thunb. ex A. DC. in DC. Prod. vii. 448, 496 = Merciera tenuifolia.
prostrata, Eckl. & Zeyh. Enum. 387 = spicata.
prostrata, E. Mey. ex A. DC. in DC. Prod. vii. 447 = reticulata.
recurva, Banks, ex A. DC. l. c. 418 = Lightfootia paniculata.
recurvata, *A. DC. l. c.* 447.—Afr. austr.
reflexa, Banks, ex A. DC. l. c. 418 = Lightfootia fasciculata.
reticulata, Lam. Encyc. vi. 232; Loisel. Herb. Amat. t. 332 = ciliata.
reticulata, *Linn. Sp. Pl.* 170.—Afr. austr.
rigidula, Banks, ex A. DC. in DC. Prod. vii. 447 = squarrosa.
secunda, Buek, in Eckl. & Zeyh. Enum. 386 = spicata.
spicata, *Linn. f. Suppl.* 143.—Afr. austr.
squarrosa, *Berg. Desc. Pl. Cap.* 42.—Afr. austr.
tenuifolia, Thunb. Fl. Cap. ii. 28 = Merciera tenuifolia.
ternifolia, Willd. ex Harv. & Sond. Fl. Cap. iii. 592 = reticulata.
Thunbergii, A. DC. Monog. Camp. 174 = Lightfootia tenella.

ROËLLOIDES, Banks, ex A. DC. Monog. Camp. 164 (1830) = **Prismatocarpus**, L'Hérit. (Campan.).

ROELPINIA, Scop. Introd. 231 (1777), err. typ. = KOELPINIA, Scop. = **Acronychia**, Forst. (Rutac.).

ROEMERA, Tratt. Gen. Pl. Disp. 88 (1802) = **Steriphoma**, Spreng. (1827) (Cappar.).
paradoxa, Tratt. l. c. = S. desmoides.

ROEMERIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Scaevola**, Linn. (Gooden.).
Lobelia, Dennst. l. c. = S. Koenigii.

ROEMERIA, Medic. in Usteri, Ann. Bot. iii. (1792) 15; DC. Syst. ii. 92 (1821). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
argemonoides, *Pomel, Nouv. Mat. Fl. Atl.* 247.—Afr. bor.
bicolor, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) I. 249.—Turkestan.
bivalvis, *DC. Syst.* ii. 93.—Mesopot.
hybrida, DC. l. c. 92 = violacea.
hybrida, Unio It. ex Presl, Bot. Bemerk. 8 = Schimperi.
orientalis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 374.—Persia; Mesopot.; Afghan.
pinnatifida, *Bélanger, Voy. Ind.*, absque num.—Ind. or.
refracta, *DC. Syst.* ii. 93.—Tauria.
rhoeadiflora, *Boiss. Diagn.* Ser. I. vi. 7.—Persia.
Schimperi, *Presl, Bot. Bemerk.* 8.—Syria.
vermiculata, *Lehm. ex Sweet, Hort. Brit.* ed. II. 585.—Persia.
violacea, *Medic. in Usteri, Ann. Bot.* iii. (1792) 15.—Europ.; Afr. bor.

ROEMERIA, Moench, Meth. 341 (1794) = **Amaranthus**, Linn.
polygonoides, Moench, l. c. = A. polygonoides.

ROEMERIA, Thunb. Nov. Gen. ix. 130 (1798), genus spurium.
argentea, Thunb. l. c. 131 = Anaphrenium argenteum.
inermis, Thunb. l. c. = Sideroxylon inerme.
melanophleos, Thunb. l. c. = Myrsine melanophlœos.

ROEMERIA, Tratt. ex DC. Syst. ii. 92 (1821) = **Steriphoma**, Spreng. (1827) (Cappar.).

ROEMERIA, Zea, ex Roem. & Schult. Syst. i. 61, 287 (1817) = **Diarrhena**, Rafin. (Gramin.).
Zeae, Roem. & Schult. l. c. 287 = D. americana.

ROEPERA, A. Juss. in Mém. Mus. Par. xii. (1825) 454. t. 15. f. 3 = **Zygophyllum**, Linn.
aurantiaca, Lindl. in Mitch. Three Exped. ii. 70 = Z. fruticosum.
Billardieri, A. Juss. in Mém. Mus. Par. xii. (1825) 434. t. 5. f. 3 = Z. Billardieri.
buxifolia, F. Muell. ex Benth. Fl. Austral. vi. 117 = Neoroepera buxifolia.
fabagifolia, A. Juss. in Mém. Mus. Par. xii. (1825) 525. t. 15 = Z. fruticulosum.
fruticulosa, A. Juss. l. c. 454. t. 5. f. 3 = Z. fruticulosum.
latifolia, Hook. f. Fl. Tasman. i. 60 = Z. apiculatum.

ROEPERIA, F. Muell. in Hook. Kew Journ. ix. (1857) 15 = **Gynandropsis**, DC. (Cappar.).
cleomoides, F. Muell. l. c. = G. Muelleri.

ROEPERIA, Spreng. Syst. iii. 13, 147 (1826) = **Ricinocarpus**, Desf. (Euphorb.).
pinifolia, Spreng. l. c. 147 = Ricinocarpus pinifolius.

ROEPEROCHARIS, Reichb. f. Otia Bot. Hamb. 104 (1881) = **Habenaria**, Willd. (Orchid.).
Bennettiana, Reichb. f. l. c. = H. Bennettiana (in Addend.).
platyanthera, Reichb. f. l. c. = H. platyanthera.

ROESLINIA, Moench, Meth. Suppl. 211 (1802) = **Chironia**, Linn. (Gentian.).
angustifolia, G. Don, Gen. Syst. iv. 203 = C. frutescens.
baccifera, G. Don, l. c. = C. baccifera.
decussata, G. Don, in Sweet, Hort. Brit. ed. III. 472 = C. frutescens.
frutescens, G. Don, Gen. Syst. iv. 203 = C. frutescens.
tetragona, Moench, Meth. Suppl. 212 = C. baccifera.

ROETTLERA, Vahl, Enum. i. 88 (1805) = ROTTLERA, Vahl = **Didymocarpus**, Wall. (Gesner.).

ROEZLIA, Regel, in Act. Hort. Petrop. i. (1871) 98 = **Monothactum**, Naud. (Melast.).
granadensis, Regel, l. c.; et Gartenfl. (1871) 353. = M. quadrangulare.

ROGERIA, J. Gay, in Ann. Sc. Nat. Sér. I. i. (1824) 457. *PEDALINEAE*, Benth. & Hook. f. ii. 1057.
adenophylla, *J. Gay, l. c.*—Afr. trop.
brasiliensis, *J. Gay, l. c.*—Bras.
longiflora, *J. Gay, l. c.*—Afr. austr.
microcarpa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 190.—Afr. trop.

ROGIERA, Planch. in Fl. des Serres, v. (1849) t. 442 = **Rondeletia**, Linn. (Rubiac.).
amoena, Planch. l. c. = Rond. amoena.
cordata, Planch. l. c. sub t. 442 = Rond. cordata.
elegans, Planch. l. c. sub t. 442 = Rond. Roezlii.
elegantissima, Regel, Gartenfl. (1865) t. 490 = Rond. gratissima.
gratissima, Linden, ex Planch. in Fl. des Serres, xv. (1862-65) 133 = Rond. gratissima.
latifolia, Decne. in Rev. Hortic. Sér. IV. ii. (1853) 121. t. 7 = Rond. amoena.
menechma, Planch. in Fl. des Serres, v. (1849) sub t. 442 = Rond. amoena.
Roezlii, Planch. l. c. = Rond. Roezlii.
rugosa, Hemsl. Biol. Centr. Am. Bot. ii. 17, in syn. = Rond. amoena.
urophylla, Hort.—Cf. Gard. Chron. (1857) 756.—Mexic.
versicolor, Lindl. & Paxt. Flow. Gard. ii. (1851-52) 69; Lem. Jard. Fleur. ii. (1852) t. 112 = Rond. amoena.

ROHDEA, Roth, Nov. Pl. Sp. 196 (1821). *LILIACEAE*, Benth. & Hook. f. iii. 772.
RHODEA, Endl. Gen. 155 (1837), sphalm.
TITRAGYNE, Salisb. Gen. Pl. Fragm. 9 (1866).
japonica, *Roth, Nov. Pl. Sp.* 197.—Japon.
nutans, Schult. f. Syst. vii. 1653 = Tupistra nutans.
Tupistra, Schult. f. l. c. 173 = Tupistra squalida.

ROHRIA, Schreb. Gen. 30 (1789) = **Tapura**, Aubl. (Chaillet.).
petioliflora, Willd. Sp. Pl. i. 186 = T. guianensis.
Schreberi, J. F. Gmel. Syst. 93 = T. guianensis?
Tapura, J. F. Gmel. l. c. = T. guianensis.

ROHRIA, Vahl, in Danske Nat. Selsk. Skriv. i. II. (1791) 15 = **Berkheya**, Ehrh. (Compos.).
armata, Vahl, l. c. ii. II. (1793) 39 = B. carthamoides.
bisulca, Thunb. Nov. Gen. 124 = Cullumia bisulca.
carlinoides, Vahl, in Danske Nat. Selsk. Skriv. i. II. (1791) 17 = B. carlinoides.
carthamoides, Thunb. l. c. iii. (1793) 103. t. 8 = B. carthamoides.
ciliaris, Thunb. l. c. 99 = Cullumia ciliaris.
cruciata, Thunb. l. c. 104 = B. cruciata.
cuneata, Thunb. l. c. 105 = B. cuneata.
cynaroides, Vahl, l. c. i. II. (1791) 16 = B. cynaroides.
decurrens, Thunb. l. c. iii. (1793) 104 = Stephanocoma berkheyoides.
grandiflora, Thunb. l. c. 107 = B. grandiflora.
hispida, Thunb. l. c. 101 = Cullumia hispida.
ilicifolia, Vahl, l. c. ii. II. (1793) 40. t. 7 = B. grandiflora.
incana, Thunb. l. c. iii. (1793) 106 = B. fruticosa.
lanceolata, Thunb. l. c. 98 = B. lanceolata.
monanthos, Thunb. l. c. 102 = B. uniflora.
obovata, Thunb. l. c. 106 = B. obovata.
palmata, Thunb. l. c. 108 = B. palmata.
patula, Thunb. Nov. Gen. 124 = Cullumia patula.
pectinata, Thunb. l. c. = Cullumia pectinata.
pungens, Thunb. l. c. 125 = B. pungens.
revoluta, Vahl, in Danske Nat. Selsk. Skriv. ii. II. (1793) 40 = B. lanceolata.
setosa, Thunb. l. c. iii. (1793) 101 = Cullumia setosa, decurrens.
spinosissima, Thunb. l. c. 108 = B. spinosissima.
squarrosa, Thunb. l. c. 100 = Cullumia squarrosa.
sulcata, Thunb. Nov. Gen. 123 = Cullumia sulcata.

ROIA, Scop. Introd. 226 (1777) = **Swietenia**, Jacq. (Meliac.).

ROJOC, Adans. Fam. ii. 146 (1763) = **Morinda**, Linn. (Rubiac.).

ROKEJEKA, Forsk. Fl. Aegypt. Arab. 90 (1775) = **Gypsophila**, Linn. (Caryoph.).
capillaris, Forsk. l. c. p. lxvi = G. Rokejeka.
deserti, J. F. Gmel. Syst. 703 = G. Rokejeka.

ROLANDRA, Rottb. in Soc. Med. Havn. Collect. ii. (1775) 256. *COMPOSITAE*, Benth. & Hook. f. ii. 237.
argentea, *Rottb. l. c.* 258.—Am. austr.
diacantha, Cass. in Dict. Sc. Nat. xlvi. 170 = argentea.
fruticosa, Rottb. in Soc. Med. Havn. Collect. ii. (1775) 268 = argentea.
monacantha, Cass. l. c. = argentea.
reptans, Willd. ex DC. Prod. v. 91 = Trichospira menthoides.
terminalis, Spreng. Syst. iii. 673 = Ichthyothere Cunabi.

ROLDANA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 10 (1825) = **Senecio**, Tourn. (Compos.)
lobata, La Llave, l. c. = S. Roldana.

ROLFINKIA, Zenk. Pl. Ind. 13. t. 14 (1837) = **Centratherum**, Cass. (Compos.).
centaurioides, Zenk. l. c. 14. t. 14 = C. reticulatum.

ROLLANDIA, Gaudich. in Freyc. Voy. Bot. 458. t. 74 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 546.
ambigua, *G. Don, Gen. Syst.* iii. 698.—Ins. Sandvic.
calycina, *G. Don. l. c.* 699.—Ins. Sandvic.
crispa, Gaudich. in Freyc. Voy. Bot. 459 = lanceolata.
Delessertiana, Gaudich. Bot. Voy. Bonite, t. 75 = lanceolata.

ROLLANDIA :—
Humboldtiana, *Gaudich. Bot. Voy. Bonite*, t. 76.—Ins. Sandvic.
Kaalae, Wawra, in Flora, lvi. (1873) 45 = Humboldtiana.
lanceolata, *Gaudich. in Freyc. Voy. Bot.* 458.—Ins. Sandvic.
longiflora, *Wawra, in Flora*, lvi. (1873) 44.—Ins. Sandvic.
montana, Gaudich. in Freyc. Voy. Bot. 13. t. 74 = lanceolata.
pedunculosa, *Wawra, in Flora*, lvi. (1873) 46.—Ins. Sandvic.
pinnatifida, G. Don, Gen. Syst. iii. 698 = Delissea pinnatifida.
scabra, Wawra, in Flora, lvi. (1873) 46 = lanceolata.

ROLLINIA, A. St. Hil. Fl. Bras. Mer. i. 28. t. 5 (1825). *ANONACEAE*, Benth. & Hook. f. i. 27.
biflora, *G. Don, Gen. Syst.* i. 91.—Peruv.
cuspidata, *Mart. Fl. Bras.* xiii. i. 20.—Bras.
dolabripetala, A. St. Hil. Fl. Bras. Mer. i. 29 = longifolia.
edulis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 30.—N. Granat.
emarginata, *Schlecht. in Linnaea*, ix. (1835) 318.—Bras.
exalbida, *Mart. Fl. Bras.* xiii. i. 19.—Bras.
exsucca, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 199.—Guian.
fagifolia, *A. St. Hil. Fl. Bras. Mer.* i. 29.—Bras.
fagifolia, *Schlecht. in Linnaea*, ix. (1835) 316.—Bras.
glaucescens, Miq. Stirp. Surin. Sel. 108 = resinosa.
glaucescens, *Sond. in Linnaea*, xxii. (1849) 557.—Bras.
laurifolia, *Schlecht. l. c.* ix. (1835) 319.—Bras.
longifolia, *A. St. Hil. Fl. Bras. Mer.* i. 29.—Bras.
membranacea, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 30.—N. Granat.
micrantha, *G. Don, Gen. Syst.* i. 91.—Peruv.
mucosa, Baill. Adansonia, viii. (1868) 268 = Sieberi.
multiflora, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 97.—Guian.
orthopetala, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 200.—Guian.
parviflora, *A. St. Hil. Pl. Us. Brés.* t. 30.—Bras.
Poeppigii, *Mart. Fl. Bras.* xiii. i. 47.—Bras.
pterocarpa, *G. Don, Gen. Syst.* i. 91.—Peruv.
puberula, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 200.—Guian.
pulchrinervia, *A. DC. l. c.* 201.—Guian.
resinosa, *Spruce, ex Benth. in Hook. Kew Journ.* v. (1853) 9.—Guian.
rufinervis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 30.—N. Granat.
rugulosa, *Schlecht. in Linnaea*, ix. (1835) 317.—Bras.
salicifolia, *Schlecht. l. c.*—Bras.
Sieberi, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 199.—Mexic.
silvatica, *Mart. Fl. Bras.* xiii. i. 18.—Bras.
Sonderiana, Walp. Ann. iii. 813 = glaucescens.
sylvatica, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 159.—Bras.
tenuifolia, *DC. ex Steud. Nom.* ed. II. ii. 464.—Guian.
tinifolia, *Klotzsch, in Schomb. Faun. & Fl. Brit. Gui.* 1163, nomen.—Guian.

ROLOFA, Adans. Fam. ii. 256 (1763) = **Mollugo**, Linn. (Ficoid.).

ROMANA, Vell. Fl. Flum. 54 (1825), partim = **Buddleia**, Linn. (Logan.).
campestris, Vell. Fl. Flum. 55; i. t. 146 = B. elegans.
crenata, Vell. l. c. t. 147.—Bras.
integerrima, l. c. t. 148.—Bras.

ROMANZOFFIA, Cham. in Nees, Hor. Phys. Berol. 71. t. 14 (1820). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 830.
altera, Cham. in Linnaea, ii. (1827) 732, 609 = sitchensis.
sitchensis, *Bong. in Mém. Acad. Pétersb.* ii. (1833) 156.—Ins. Sitcha.
unalaschcensis, *Cham. in Nees, Hor. Phys. Berol.* 72. t. 14.—Ins. Aleut.

ROMANZOVIA, Spreng. Syst. i. 584 (1825) = praec.

ROMANZOWIA, DC. Prod. i. 319 (1824) = praec.

ROMBUT, Rumph. ex Adans. Fam. ii. 284 (1763) = **Cassytha**, Linn. (Laurin.).

ROMNEYA, Harv. in Hook. Lond. Journ. Bot. iv. (1845) 74. t. 3. *PAPAVERACEAE*, Benth. & Hook. f. i. 51.
Coulteri, *Harv. l. c.* 75.—Calif.

ROMOVIA, Muell. Arg. in DC. Prod. xv. II. 1134, in syn. (1866), sphalm. = RONNOWIA, Buc'hoz = **Omphalea**, Linn. (Euphorb.).

ROMUALDEA, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvi. (1872) 370 = **Hippocratea**, Linn. (Celastr.).
apiculata, Triana & Planch. l. c., in nota = H. megacarpa.
integrifolia, Triana & Planch. l. c. = H. integrifolia.
lancifolia, Triana & Planch. l. c. = H. lancifolia (in Addend.).
macrophylla, Triana & Planch. l. c. = H. macrophylla.
malpighifolia, Triana & Planch. l. c. 372 = H. integrifolia.
Wrightiana, Triana & Planch. l. c. = H. malpighiaefolia ?

ROMULEA, Maratti, Diss. Romul. et Saturn. Romae, 13. t. 1 (1772). *IRIDEAE*, Benth. & Hook. f. iii. 694.
ILMU, Adans. Fam. ii. 497 (1763), nomen prius.
SPATALANTHUS, Sweet, Brit. Flow. Gard. Ser. I. t. 300 (1829).
TRICHOMEMA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 195 (1821).
TRICHONEMA, [Ker-Gawl.] in Bot. Mag. t. 575 (1802).
arenaria, *Eckl. Top. Verz.* 18.—Afr. austr.
aurea, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 399 = sublutea.
barbata, *Baker, in Journ. Bot.* xiv. (1876) 236.—Afr. austr.
bulbocodioides, *Eckl. Top. Verz.* 19.—Afr. austr.
Bulbocodium, Kunze, in Flora, xxix. (1846) 689 = Clusiana.
Bulbocodium, Parl. Fl. Ital. iii. 246 = Linaresii.
Bulbocodium, *Sebast. & Mauri, Fl. Roman.* 17.—Reg. Mediterr.
caelestina, *Vis. Cat. Hort. Patav.* (1842) 145.—Hab. ?
camerooniana, *Baker, in Journ. Bot.* xiv. (1876) 236.—Afr. trop.
candida, *Tenore, Cat. Ort. Nap.* (1845) 95.—Afr. austr.
caulescens, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 399 = bulbocodioides.
chloroleuca, *Baker, in Journ. Linn. Soc.* xvi. (1877) 89.—Afr. austr.
chloroleuca, Eckl. Top. Verz. 20 = bulbocodiodes.
Clusiana, *Nym. Syll. Suppl.* 62; *Baker, in Journ. Linn. Soc.* xvi. (1877) 87.—Hispan.; Lusit.
Columnae, *Sebast. & Mauri, Fl. Roman.* 18.—Reg. Mediterr.
corsica, *Jord. & Fourr. Brev. Pl. Nov. Fasc.* ii. 107.—Ins. Corsic.
crocea, Boiss. & Heldr. Diagn. Ser. I. xiii. 18 = R. Bulbocodium.
crocifolia, *Vis. Fl. Dalm. Suppl.* 30. t. 2. f. 2.—Dalmat.
cruciata, *Eckl. Top. Verz.* 19; *Baker, in Journ. Linn. Soc.* xvi. (1877) 89.—Afr. austr.
cuprea, *Baker, in Journ. Bot.* xiv. (1876) 236.—Afr. austr.
dichotoma, Baker, in Journ. Linn. Soc. xvi. (1877) 89 = rosea.
elegans, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 400.—Afr. austr.
elongata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 87.—Ins. Sardoa.
filifolia, *Eckl. Top. Verz.* 20.—Afr. austr.
flaveola, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 106 = R. Bulbocodium.
flexuosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 400.—Afr. austr.
fragrans, Eckl. Top. Verz. 19 = Ixia fragrans.
grandiflora, Tin. in Todaro, Ind. Sem. Hort. Panorm. (1857) 44 = R. Bulbocodium.
grandiscapa, *J. Gay, ex Baker, in Journ. Linn. Soc.* xvi. (1877) 87.—Ins. Canar.
Hartungii, Parl. ex G. Hartung, Geogn. Verh. Ins. Lanzarote, Fuertav. 143 = grandiscapa.
hirsuta, *Eckl. Top. Verz.* 19.—Afr. austr.

ROMULEA:—
latifolia, *Baker, in Journ. Bot.* xiv. (1876) 237.—Afr. austr.
ligustica, *Parl. Fl. Ital.* iii. 249.—Italia.
Linaresii, De Not. & Caldesi, ex Parl. Fl. Ital. iii. 249 = ligustica.
Linaresii, *Parl. Fl. Panorm.* i. 38.—Europ.; As. Min.
Linaresii, Rolli, ex Parl. Fl. Ital. iii. 251 = ramiflora.
longifolia, *Baker, in Journ. Linn. Soc.* xvi. (1877) 89.—Afr. austr.
Macowani, *Baker, in Journ. Bot.* xiv. (1876) 236.—Afr. austr.
minima, *Tenore, in Mem. Acc. Sc. Borb. Nap.* iii. II. (1832) 117.—Italia.
minuta, Eckl. Top. Verz. 19 = Pauridia hypoxioides.
minutiflora, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 399.—Afr. austr.
modesta, Jord. Ic. Fl. Europ. t. 110 = R. Columnae.
multiflora, Req. ex Parl. Fl. Ital. iii. 243 = R. Bulbocodium.
neglecta, *Jord. & Fourr. Brev. Pl. Nov. Fasc.* ii. 107.—Italia.
nivalis, Boiss. Fl. Orient. v. 116 = R. Bulbocodium.
nivalis, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 400.—Afr. austr.
numidica, *Jord. & Fourr. Brev. Pl. Nov. Fasc.* i. 49.—Algeria.
obscura, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 399.—Afr. austr.
Parlatorii, Todaro, Ind. Sem. Hort. Panorm. (1857) 45;—Cf. Linnaea, xxix. (1857–58) 749 = ramiflora.
parviflora, *Eckl. Top. Verz.* 19.—Afr. austr.
pudica, Baker, in Journ. Linn. Soc. xvi. (1877) 89 = rosea.
pulchella, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 106 = R. Bulbocodium.
purpurascens, Parl. Fl. Panorm. i. 40 = ramiflora.
purpurascens, *Tenore, in Mem. Acc. Sc. Borb. Nap.* iii. II. (1832) 117.—Reg. Mediterr.
pylia, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 398.—Afr. austr.
pylia, Nym. Consp. 704 = R. Bulbocodium.
ramiflora, *Tenore, Ind. Sem. Hort. Neap.* (1826) 3; *et in Mem. Acc. Sc. Borb. Nap.* iii. II. (1832) 117. t. 7.—Reg. Mediterr.
ramosa, Eckl. Top. Verz. 19 = arenaria ?
recurva, Eckl. l. c. 20 = bulbocodiodes.
reflexa, *Eckl. l. c.* 18.—Afr. austr.
Requienii, *Parl. Fl. Ital.* iii. 248.—Ins. Corsic.
Revelieri, Jord. & Fourr. Brev. Pl. Nov. Fasc. i. 49 = bulbocodiodes.
Rollii, Parl. Fl. Ital. iii. 251 = ramiflora.
rosea, *Eckl. Top. Verz.* 19.—Afr. austr.
similis, *Eckl. l. c.*—Afr. austr.
speciosa, Baker, in Journ. Linn. Soc. xvi. (1877) 89 = rosea.
spiralis, *Baker, l. c.* 90.—Afr. austr.
subalbida, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 108 = R. Columnae.
sublutea, *Baker, in Journ. Linn. Soc.* xvi. (1877) 88.—Afr. austr.
subpalustris, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 398.—Afr. austr.
subpalustris, Nym. Consp. 704 = purpurascens.
syrtica, Jord. & Fourr. Brev. Pl. Nov. Fasc. i. 48 = R. Bulbocodium.
tabularis, Eckl. & Zeyh. 18, nomen = speciosa.
tortuosa, *Baker, in Journ. Linn. Soc.* xvi. (1877) 88.—Afr. austr.
tridentifera, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 398.—Afr. austr.
tubata, *Klatt, l. c.* 401.—Afr. austr.
uliginosa, Kunze, in Flora, xxix. (1846) 690 = R. Bulbocodium.
umbellata, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 402 = R. Bulbocodium.
uncinata, Klatt, l. c. 401 = hirsuta.
vulgaris, Eckl. Top. Verz. 18 = rosea.
Zeyheri, *Eckl. l. c.* 19.—Afr. austr.

RONABEA, Aubl. Pl. Gui. i. 154. t. 59 (1775) = **Psychotria**, Linn. (Rubiac.).
arborea, Blanco, Fl. Filip. ed. II. 114 (= *Canthium mite*).—Ins. Philipp.
australis, A. Rich. Voy. Astrol. Bot. i. 265 = Coprosma foetidissima ?
bipinnata, Blanco, Fl. Filip. ed. I. 162 (= *Canthium mite*).—Ins. Philipp.

RONABEA:—
didymocarpos, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 170.—Guian.
emetica, A. Rich. l. c. = Psychotria emetica.
erecta, Aubl. Pl. Gui. i. 156 = P. axillaris.
latifolia, Aubl. l. c. 154. t. 59 = P. axillaris.
morindoides, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 170.—Guian.
myodendron, A. Rich. l. c. = P. leiocarpa.
viscoides, A. Rich. l. c. = P. parasitica.

RONABIA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = praec.
myodendra, St. Lag. l. c. = P. leiocarpa.

RONCELIA, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 289 (1870), sphalm. = ROUCELA, Dum. = **Campanula**, Tourn.

RONDACHINE, Bosc, Encyc. Agric. vi. (1816) 180 = **Brasenia**, Schreb. (Nymph.).

RONDELETIA, Linn. Gen. ed. I. 59 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 48.
ARACHNIMORPHA, Desv. in Ham. Prod. Pl. Ind. Occ. 28 (1825).
ARACHNOTHRYX, Planch. in Fl. des Serres, v. (1849) sub t. 442.
LIGHTFOOTIA, Schreb. Gen. 122 (1789).
PETESIA, P. Br. Hist. Jamaic. 143. tt. 2, 3 (1756).
ROGIERA, Planch. in Fl. des Serres, v. (1849) t. 442.
STEVENSIA, Poit. in Ann. Mus. Par. iv. (1804) 235. t. 60.
WILLDENOVIA, J. F. Gmel. Syst. 362 (1791).
affinis, *Hemsl. Diagn. Pl. Nov.* 28.—Mexic.
africana, T. Winterb. Account Sierra Leone, ii. 46 = Crossopteryx Kotschyana.
alaternoides, *A. Rich. Fl. Cub. Fanerog.* ii. 13.—Cuba.
americana, *Linn. Sp. Pl.* 172.—Ind. occ.; Am. austr.
amoena, *Hemsl. Diagn. Pl. Nov.* 26.—Mexic.
arborescens, *Griseb. Fl. Brit. W. Ind.* 327.—Ind. occ.
asiatica, Blanco, Fl. Filip. ed. I. 146 = Wendlandia paniculata.
asiatica, Linn. Sp. Pl. 172 = Tarenna zeylanica.
avenia, *Wright, in Sauv. Fl. Cub.* 63.—Cuba.
Backhousii, *Hook. f. Bot. Mag.* t. 6290.—Am. trop.
Berteriana, *DC. Prod.* iv. 409.—Ins. S. Doming.
biflora, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 276 = Sipanea biflora.
brachyacantha, *DC. Prod.* iv. 410.—Bras.
brevipes, *Benth. Pl. Hartw.* 191.—N. Granat.
Budda, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 138 = Wendlandia coriacea.
buddleioides, *Benth. Pl. Hartw.* 69.—Mexic.
buxifolia, *Vahl, Eclog. Am.* ii. 11. t. 12.—Ind. occ.
byrsophylla, Spreng. Syst. iv. Cur. Post. 74 = Wendlandia coriacea.
camarioca, *Wright, in Sauv. Fl. Cub.* 62.—Cuba.
capitata, *Benth. in Hook. Journ. Bot.* iii. (1841) 217.—Guian.
capitellata, *Hemsl. Diagn. Pl. Nov.* 28.—Mexic.
chamaebuxifolia, *Griseb. Cat. Pl. Cub.* 128.—Cuba.
cincta, *Griseb. Fl. Brit. W. Ind.* 329.—Ind. or.
cinerea, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 141 = Wendlandia exserta.
coccinea, [Moç. & Sesse,] ex DC. Prod. iv. 408 = odorata.
cordata, *Benth. Pl. Hartw.* 85.—Guatemala.
coriacea, Spreng. Syst. i. 707 = Rudgea eriantha.
coriacea, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 142 = Wendlandia coriacea.
cornifolia, Benth. Pl. Hartw. 39 = Deppea cornifolia.
correifolia, *Griseb. Cat. Pl. Cub.* 129.—Cuba.
corymbosa, Jack, in Malay. Misc. i. (1820) n. I. 4 = Greenea Jackii.
cumanensis, *H. B. & K. Nov. Gen. et Sp.* iii. 394.—Venezuela.
cymosa, Poir. Encyc. vi. 256 = Plectronia didyma.
daphnoides, *Griseb. Fl. Brit. W. Ind.* 327.—Jamaic.
densiflora, Blume, Bijdr. 974 = Wendlandia densiflora.
dichotoma, Pers. Syn. i. 203 = americana.
dilatata, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 286.—Guian.
discolor, *H. B. & K. Nov. Gen. et Sp.* iii. 396.—N. Granat.
disperma, *Jacq. Enum. Pl. Carib.* 16.—Am. austr.
dubia, *Hemsl. Diagn. Pl. Nov.* 28.—Mexic.

RONDELETIA:—
elongata, *Bartl. ex DC. Prod.* iv. 409.—Mexic.
eriantha, *Benth. Pl. Hartw.* 192.—N. Granat.
eriocarpa, *Karst. in Linnaea*, xxx. (1859-60) 147.—N. Granat.
erythroneura, *Karst. l. c.* 148.—N. Granat.
exserta, Roxb. Hort. Beng. 14; Fl. Ind. i. 523 = Wendlandia exserta.
febrifuga, Afzel. ex G. Don, Gen. Syst. iii. 516 = Crossopteryx Kotschyana.
floribunda, G. Don, Gen. Syst. iii. 516 = Holarrhena africana.
glauca, *Griseb. Fl. Brit. W. Ind.* 329.—Ind. occ.
glomerulata, *Spreng. Syst.* i. 707.—Guian.
gracilis, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
gratissima, *Hemsl. l. c.* 25.—Am. trop.
Heynii, Roem. & Schult. Syst. v. 234 = Wendlandia exserta.
hirsuta, Spreng. ex DC. Prod. iv. 409 = Berteriana.
hirsuta, *Sw. Prod. Veg. Ind. Occ.* 41.—Jamaic.
hirta, Bartl. ex DC. Prod. iv. 408 = panamensis.
hirta, *Sw. Prod. Veg. Ind. Occ.* 41.—Jamaic.
hypoleuca, *Griseb. Cat. Pl. Cub.* 128.—Cuba.
incana, Spreng. Syst. i. 707 = royenaefolia.
incana, *Sw. Prod. Veg. Ind. Occ.* 41.—Jamaic.
intermedia, *Hemsl. Diagn. Pl. Nov.* 26.—Mexic.
Jurgensenii, *Hemsl. l. c.* 29.—Mexic.
laevigata, *Ait. Hort. Kew.* ed. II. i. 366.—Ind. occ.
laevigata, *DC. Prod.* iv. 408 (sp. dub.).—Ins. Trinit.
lanceolata, Spreng. Syst. i. 707 = Mussaenda luteola.
Landia, Spreng. l. c. = Mussaenda Landia.
laniflora, *Benth. Pl. Hartw.* 85.—Guatemala.
latifolia, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 43.—Am. centr.
laurifolia, *Sw. Fl. Ind. Occ.* i. 363.—Jamaic.
leptacantha, *DC. Prod.* iv. 410.—Cuba.
leucophylla, *H. B. & K. Nov. Gen. et Sp.* iii. 395.—Mexic.
ligustroides, *Hemsl. Diagn. Pl. Nov.* 26.—Mexic.
Lindeniana, *A. Rich. Fl. Cub. Fanerog.* ii. 13.—Cuba.
linguiformis, *Hemsl. Diagn. Pl. Nov.* 29.—Mexic.
longiflora, Cham. in Linnaea, ix. (1834) 240 = Hindsia longiflora.
longifolia, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 137 = Adenosacme longifolia.
loniceroides, G. Don, Gen. Syst. iii. 516 = Usteria guineensis.
lucida, Wall. Cat. n. 8453 (= *Webera fragrans*).—Malaya.
macrophylla, Hort. ex Hemsl. Biol. Centr. Am. Bot. ii. 17 = amoena.
microdon, *DC. Prod.* iv. 408.—Cuba.
microdonta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = praec.
microphylla, *Wright, ex Griseb. Cat. Pl. Cub.* 127.—Cuba.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 43.—Nicarag.
nitida, *Hemsl. Diagn. Pl. Nov.* 29.—Mexic.
obovata, Linn. Syst. ed. XII. 163 = odorata.
odorata, *Jacq. Enum. Pl. Carib.* 16.—Cuba; Mexic.
orissensis, Roth, Nov. Pl. Sp. 142 = Wendlandia exserta.
panamensis, *DC. Prod.* iv. 408.—Panama.
paniculata, Roxb. Hort. Beng. 15; Fl. Ind. i. 521 = Wendlandia paniculata.
parviflora, *Poir. Encyc.* vi. 252.—Ins. Martinic.
pedicellaris, *Wright, in Sauv. Fl. Cub.* 63.—Cuba.
peduncularis, *A. Rich. Fl. Cub. Fanerog.* ii. 14.—Cuba.
pendula, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 140 = Wendlandia pendula.
phialanthoides, *Griseb. Cat. Pl. Cub.* 129.—Ind. occ.
pilosa, *Sw. Prod. Veg. Ind. Occ.* 41.—Ind. occ.
Poitaei, *Griseb. Fl. Brit. W. Ind.* 328; *et in Mem. Am. Acad.* N. S. viii. (1863) 505.—Ind. occ.
polita, *Griseb. l. c.* 326.—Jamaic.
proxima, D. Don, Prod. Fl. Nep. 139 = Wendlandia tinctoria.
pubescens, *H. B. & K. Nov. Gen. et Sp.* iii. 396.—N. Granat.
Purdiei, *Hook. f. Bot. Mag.* t. 5669.—N. Granat.
racemosa, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 133 = Palicourea laevigata.
racemosa, Hort. ex Tenore, in Parl. Giorn. Bot. Ital. ii. (1847) VII-VIII. 203 = Psychotria trichotoma.
racemosa, *Sw. Fl. Ind. Occ.* i. 360.—Jamaic.

RONDELETIA:—
reflexa, *Benth. Pl. Hartw.* 192.—N. Granat.
repens, *Linn. Syst.* ed. X. 928.—Jamaic.?
rhynchospora, *H. B. & K. Nov. Gen. et Sp.* iii. 395.—N. Granat.
rigida, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 505.—Ind. occ.
Roezlii, *Hemsl. Biol. Centr. Am. Bot.* ii. 23.—Guatemala.
royenaefolia, *DC. Prod.* iv. 410.—Ins. S. Doming.
rugosa, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 43.—Am. centr.
scabra, *Hemsl. Diagn. Pl. Nov.* 29.—Mexic.
scandens, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 524.—Ins. Maurit.
speciosa, Lodd. Bot. Cab. t. 1893; Paxt. Mag. Bot. ii. (1836) 242 = odorata.
spicata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 139 = Greenea Jackii.
spinescens, Spreng. Syst. i. 708 = Randia nitida.
splendens, Regel, Cat. Pl. Hort. Aksakov. 124 (Quid?).—Hab.?
stenosiphon, *Hemsl. Diagn. Pl. Nov.* 26.—Mexic.
stereocarpa, *Griseb. Fl. Brit. W. Ind.* 327.—Ins. Dominic.
stricta, Roth, Nov. Pl. Sp. 140 = Hyptianthera stricta.
strigosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 24.—Guatemala.
superba, Regel, Cat. Pl. Hort. Askakov. 124 (Quid?).—Hab.?
tetrandra, Roxb. Hort. Beng. 85; Fl. Ind. i. 524 = Hedyotis macrophylla.
thyrsiflora, Hort. ex Planch. in Fl. des Serres, viii. (1852-53) 13, sub t. 754 = cordata.
thyrsiflora, Roth, Nov. Pl. Sp. 141 = Wendlandia exserta.
thyrsoidea, *Sw. Prod. Veg. Ind. Occ.* 41.—Jamaic.
tinctoria, Blume, Bijdr. 974 = Wendlandia glabrata.
tinctoria, Roxb. Hort. Beng. 15; Fl. Ind. i. 522 = Wendlandia tinctoria.
tinifolia, *Griseb. Cat. Pl. Cub.* 129.—Cuba.
tomentosa, *Sw. Prod. Veg. Ind. Occ.* 41.—Jamaic.
triflora, Vahl, Symb. Bot. iii. 34 = pilosa.
trifolia, *Jacq. Select. Am.* 60. t. 43.—Jamaic.
trifoliata, Linn. Sp. Pl. ed. II. 1671 = praec.
umbellata, J. F. Gmel. Syst. 1616 = *umbellulata*, Sw.
umbellulata, Macfad. ex Griseb. Fl. Brit. W. Ind. 328 = hirta.
umbellulata, Sw. Prod. Veg. Ind. Occ. 41 = buxifolia.
venosa, *Wright, ex Griseb. Cat. Pl. Cub.* 128.—Cuba.
verbenacea, Griseb. l. c. 130 = Ceratopyxis verbenacea.
versicolor, J. Sm. in Bot. Mag. t. 4579 = amoena.
villosa, *Hemsl. Diagn. Pl. Nov.* 27.—Mexic.
virgata, *Sw. Prod. Veg. Ind. Occ.* 41.—Ins. S. Doming.
volubilis, [Moç. & Sesse,] ex DC. Prod. iv. 470 = Lygodisodea mexicana.

RONNBERGIA, E. Morr. & André, Illustr. Hortic. (1874) 120. t. 177. *BROMELIACEAE*, Benth. & Hook. f. iii. 661.
columbiana, *E. Morr. in Belg. Hortic.* (1885) 82.—N. Granat.
Morreniana, *Linden & André, in Illustr. Hortic.* (1874) 120. t. 177.—N. Granat.

RONNOWIA, Buc'hoz, Pl. Nouv. Découv. 6. t. 4 (1779) = **Omphalea**, Linn. (Euphorb.).
domingensis, Buc'hoz, l. c. = O. commutata?

ROPALA, J. F. Gmel. Syst. 225 (1791) = **Roupala**, Aubl. (Proteac.).

ROPALOCARPUS, Boj. Hort. Maurit. 44 (1837). *TILIACEAE*, Benth. & Hook. f. i. 238.
RAPOLOCARPUS Boj. (err. typ. in loc. cit.).
lucidus, *Boj. l. c.*—Madag.
Thouarsianus, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 393.—Madag.
triplinervius, *Baill. Adansonia*, x. (1871) 106.—Madag.

ROPALON, Rafin. New Fl. Am. ii. 17 (1836) = **Nuphar**, Sm. (Nymph.).
sagittatum, Rafin. l. c. = N. sagittaefolium.

ROPALOPETALUM, Griff. Notul. iv. 716 (1854) = **Artabotrys**, R. Br. (Anonac.).
uniflorum, Griff. l. c. 717 = A. burmanicus.

ROPHOSTEMON, Endl. Gen. 21 (1837), sphalm. = ROTROSTEMON, Blume = **Pogonia**, Juss. (Orchid.).

ROPHOSTEMUM, Reichb. Nom. 55 (1841) = praec.

ROPOUREA, Aubl. Pl. Gui. i. 198. t. 78 (1775). *VERBENACEAE*?
CAMAX, Schreb. Gen. i. 144 (1789).
IDESIA, Scop. Introd. 199 (1777).
PLOTIA, Neck. Elem. ii. 55 (1790).
guianensis, *Aubl. Pl. Gui.* i. 198.—Guian.

ROPTROSTEMON, Blume, Fl. Jav. Praef. p. vi (1828) = **Pogonia**, Juss. (Orchid.).
concolor, Lindl. Gen. et Sp. Orch. 453 = P. concolor.
discolor, Lindl. l. c. = P. discolor.

RORAM, Endl. Gen. 85, in syn. (1836), sphalm. = RARAM, Adans. = **Cenchrus**, Linn. (Gramin.).

RORELLA, Hall. ex All. Fl. Pedem. ii. 88 (1785) = **Drosera**, Linn.
longifolia, All. l. c. = Drosera longifolia.
lusitanica, Rafin. Fl. Tellur. iii. 36 = Drosophyllum lusitanicum.
rotundifolia, All. Fl. Pedem. ii. 88 = Drosera rotundifolia.

RORIDA, J. F. Gmel. Syst. 260 (1791) = **Cleome**, Linn. (Cappar.).
arabica, J. F. Gmel. l. c. = C. droserifolia.

RORIDULA, Forsk. Fl. Aegypt. Arab. 35 (1775) = **Cleome**, Linn. (Cappar.).
arabica, Roem. & Schult. Syst. iii. 13 = C. droserifolia.
droserifolia, Forsk. Fl. Aegypt. Arab. p. lxii = C. droserifolia.
tetrandra, Vitm. Summa Pl. ii. 68 = C. droserifolia.

RORIDULA, Linn. Gen. ed. VI. 567 (1764). *DROSERACEAE*, Benth. & Hook. f. i. 664.
IREON, Burm. f. Prod. Fl. Cap. 6 (1768).
IRIDION, Roem. & Schult. Syst. v. 436, in syn. (1819).
dentata, *Linn. Gen.* ed. VI. 567.—Afr. austr.
gorgonias, *Planch. in Ann. Sc. Nat.* Sér. III. ix. (1848) 307.—Afr. austr.
muscicarpa, Gaertn. Fruct. i. 298. t. 62 = dentata.
verticillata, Pers. Syn. i. 253 = dentata.

RORIPA, Scop. Fl. Carn. ed. I. 520 (1760) = **Nasturtium**, R. Br. (Crucif.).
amphibia, Bess. Enum. Pl. Volh. 27 = N. amphibium.
anceps, Fuss, Fl. Transs. 47 = N. anceps.
armoracioides, Fuss, l. c. = N. armoracioides.
auriculata ×, Menyh. Kalocsa Vidék. Növényt. 39.—Hungar.
austriaca, Spach, Hist. Vég. Phan. vi. 513 = N. austriacum.
barbareoides, Čelak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 458 = N. sylvestre.
Borbasii, Menyh. in Oestr. Bot. Zeitschr. xxix. (1879) 173 = *auriculata*.
Camelinae, Spach, Hist, Vég. Phan. vi. 514 = N. austriacum.
hispanica, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 845 = N. hispanicum.
Kerneri, Menyh. Kalocsa Vidék. Növényt. 39 = N. brachycarpum.
lippizensis, Reichb. Ic. Fl. Germ. xii. 15 = N. lippizense.
Menyharthiana ×, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 246.—Hungar.
nasturtioides, Spach, Hist. Vég. Phan. vi. 506 = N. palustre.
palustris, Bess. Enum. Pl. Volh. 27; Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 844 = N. palustre.
pyrenaica, Spach, Hist. Vég. Phan. vi. 506 = N. pyrenaicum.
riparia, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 71 = N. amphibium.
rivularis, Reichb. Ic. Fl. Germ. xii. 15 = N. sylvestre.
rusticana, Gren. & Godr. Fl. Fr. i. 127 = Cochlearia Armoracia.
silvestris, Fuss, Fl. Transs. 47 = *sylvestris*.
stolonifera, Nym. Consp. 35 = N. amphibium.
sylvestris, Bess. Enum. Pl. Volh. 27 = N. sylvestre.
terrestis, Fuss, Fl. Transs. 47 = N. anceps?

ROSA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 146 (1737). *ROSACEAE*, Benth. & Hook. f. i. 625.
HULTENIA, Reichb. Handb. 243 (1837).
HULTHEMIA, Dum. Not. Nov. Gen. Hulth. (1824); Anal. Fam. 39 (1829).
LOWEA, Lindl. Bot. Reg. t. 1261 (1829).
RHODOPHORA, Neck. Elem. ii. 91 (1790).
RHODOPSIS, Reichb. Nom. 168 (1841).
abietina, Gren. Fl. Jurass. 134 = tomentosa.
abscondita, Christ, in Gremli, Excurs. Schw. ed. IV. 167 = sepium.
abstenta, *Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 347.—Persia.
abyssinica, *R. Br. in Salt, Abyss. App.* 64.—Abyss.
acantha, *Waitz, ex Link, Enum. Hort. Berol.* ii. 60, nomen.—Hab.?
Acharii, Billb. in Svensk Bot. ix. t. 577 = canina.
acicularis, *Lindl. Ros. Monog.* 44. t. 8.—As. bor.; Am. bor.
acicularis, Rupr. ex Nym. Consp. 236 = carelica.
aciphylla, Rau, Enum. Ros. 69 = canina.
aculeatissima, Dum.-Cours. Bot. Cult. ed. I. iii. 340 = spinosissima.
acuminata, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 216 = humilis.
acuminata, Sw. ex Reichb. Fl. Germ. Excurs. 614 = cinnamomea.
acuta, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 270 = canina.
acuta, *Fisch. ex Sweet, Hort. Brit.* ed. II. 180.—Reg. Caucas.
addita, *Déségl. in Journ. Bot.* xi. (1874) 168.—Persia.
adenocalyx, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 434 = canina.
adenophora, *Kit. in Schult. Oestr. Fl.* ed. II. ii. 69.—Hungaria.
adenophylla, *Willd. Enum. Hort. Berol.* 546.—Hab.?
adjecta, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 105 = alpina.
adscita, Déségl. in Billotia, i. (1864) 34 = canina.
aeduensis, Déségl. & Gillot, in Bull. Soc. Bot. Belg. xix. (1880) 37 = sepium.
affinis, Rau, Enum. Ros. 79 = canina.
affinis, Sternb. in Flora, ix. (1826) 1. Beil. 80 = spinosissima.
Afzeliana, Fries, Fl. Halland. 87 = canina.
aginensis, Ripart, ex Déségl. in Journ. Bot. xii. (1874) 169 = Amansii.
agraria, Ripart, ex Déségl. l. c. 350 = canina.
agrestina, Ripart, ex Déségl. l. c. = canina.
agrestis, C. C. Gmel. Fl. Bad. ii. 416 = gallica.
agrestis, *Savi, Fl. Pis.* i. 475.—Europ.; Afr. bor.
agrestis, Sw. Summa Pl. 18; Adnot. Bot. 39 = canina.
alba, All. Fl. Pedem. ii. 139 = sempervirens.
alba, Gaertn. Mey. & Scherb. Fl. Wett. ii. 232 = tomentosa.
alba, *Linn. Sp. Pl.* 492.—Reg. Caucas.
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1883) 278.—Turkestan.
albicans, *Godet, ex Boiss. Fl. Orient.* ii. 675.—Persia.
albiflora, Opiz, in Flora, v. (1822) 268 = agrestis.
albiflora, *Steud. Nom.* ed. II. ii. 465.—Tauria.
albo-lutescens, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 319, in obs. 1.—Gallia.
aleutensis, Crép. l. c. xiv. (1875) 47; xvi. (1876) 25 = nutkana?
Aliothi, Christ, in Gremli, Excurs. Schw. ed. II. 177 = trachyphylla.
Allionii, *Burnat & Gremli, Suppl. Mon. Roses Alpes-Marit.* 16.—Europ.
alpestris, Déségl. Ess. Monog. Obs. 55 = alpina.
alpestris, Rapin, ex Rent. Cat. Pl. Vasc. Genèv. ed. II. 68 = glauca.
alpicola, *Rouy, in Bull. Soc. Bot. Fr.* xxii. (1875) 295.—Gallia.
alpina, *Linn. Sp. Pl.* ed. II. 703.—Europ.
alpina, Pall. Fl. Ross. ii. 61 = Gmelini.
alpina, Pall. Reise, ii. 518 = spinosissima.
alpinoides, *Déségl. in Bull. Soc. Sc. Angers* (1878).—*Cf. Bull. Soc. Bot. Fr.* xxv. (1878) *Bibl.* 107.—Gallia.
alpino-minuta, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 295 = alpicola.
alpiphila, Arv. Ess. Pl. Dauph. (1871) 27 = glauca.
altaica, Willd. Enum. Hort. Berol. 543 = spinosissima.
amabilis, Borck. Forstbot. iii. 1833 = seq.?
amabilis, Wrede, ex Steud. Nom. ed. I. 695 = perfecta.
Amansii, *Déségl. & Ripart, ex Déségl. in Journ. Bot.* xi. (1874) 169.—Europ.

ROSA:—
ambigens, Gremli, Excurs. Schw. ed. II. 178 = canina.
ambigua, Lej. Rev. Fl. Spa, 98 = canina.
ambigua, *Léman, in Journ. Phys.* lxxxvii. (1818) 366.—Gallia.
ambigua, *Waitz, ex Link, Enum. Hort. Berol.* ii. 60, nomen; *Spreng. Syst.* ii. 548.—Hab.?
amblyotis, C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. vi. (1847); (Zimmtr.) 30 = nutkana.
americana, [Waitz] Breit. Hort. Breit. 416; Link, Enum. Hort. Berol. ii. 56 = blanda.
amoena, Dum.-Cours. Bot. Cult. ed. II. v. 477 = centifolia.
amoena, Waitz, ex Link, Enum. Hort. Berol. ii. 60, nomen = praec.?
amoyensis, Hance, in Journ. Bot. vi. (1868) 297 = microcarpa.
amphora, Gren. ex Déségl. in Billotia, i. (1864) 121 = canina.
amygdalifolia, Ser. in DC. Prod. ii. 601 = laevigata.
analoga, Déségl. in Bull. Soc. Bot. Belg. xix. (1880) 32 = canina.
anceps, *Bouteill. in Bull. Soc. Bot. Fr.* xxvii. (1880) 298.—Gallia.
Ancistrum, *Léman, in Journ. Phys.* lxxxvii. (1818) 367.—Gallia.
andegavensis, Bast. Fl. Maine-et-Loire, 189 = canina.
Andorae, *Burnat & Gremli, Suppl. Mon. Roses Alpes-Marit.* 45.—Europ.
Andreae, *Lange, Ind. Sem. Hort. Haun.* (1874) 23.—Ins. Sachalin.
Andrewsii, Tratt. Ros. ii. 205 = alpina.
Andrzeiouskii, Bor. Fl. Centr. Fr. ed. II. n. 692 = mollissima.
Andrzeiovii, Bess. Cat. Hort. Crem. Suppl. iv. 19; Enum. Pl. Volh. 19 = canina.
Andrzeiowskii, Nym. Consp. 232 = canina.
Andrzejowsciana, Koch, Syn. Fl. Germ. ed. I. 227 = seq.
Andrzejowscii, Stev. ex Bieb. Fl. Taur. Cauc. iii. 339, in nota = canina.
Anemone, Hort. ex Red. Ros. ii. 116 = centifolia.
anemoneflora, *Andr. Ros.* t. 32.—Gallia.
anemoneflora, *Fortune, ex Lindl. in Journ. Hort. Soc.* ii. (1847) 316.—China.
anglica, Hort. ex Steud. Nom. ed. I. 695 = centifolia.
anisopoda, Christ, in Gremli, Excurs. Schw. ed. IV. 165 = ferox.
anisopoda, Vukot. in Kern. Sched. Fl. Excurs. iii. 55 = zagrabiensis.
annesiensis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 75 = tomentosa.
anomala, Ripart, ex Déségl. & Gillot, in Bull. Soc. Bot. Belg. xix. (1880) 37 = tomentosa.
anoplantha, Christ, in Gremli, Excurs. Schw. ed. IV. 178 = pomifera.
anserinaefolia, Boiss. Diagn. Ser. I. vi. 51 = Beggeriana.
apennina, Woods, Tour. Fl. 123 = agrestis.
apiifolia, Willd. Enum. Hort. Berol. Suppl. 38 = centifolia.
approximata, Déségl. in Billotia, i. (1864) 37 = gallica.
apricorum, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 245 = rubiginosa.
arabica, Crép. l. c. 344 = rubiginosa.
arborea, Pers. Syn. ii. 50 = moschata.
arcana, *Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 385.—Europ.
arduennensis, Crép. in Bull. Acad. Brux. Sér. II. xiv. (1862) 102 = tomentosa.
arduennensis, Lej. Fl. Spa, ii. 340, sphalm. = Rubus arduennensis.
arenaria, Bieb. ex Spreng. Syst. ii. 547 = Willdenowii.
arenivaga, Déségl. ex Jullien, in Mém. Soc. Acad. Maine-et-Loire, xii. (1862) 69 = gallica.
arguta, *Muss.-Pursch. ex Crép. in Bull. Soc. Bot. Belg.* xi. (1872) 78.—Reg. Caucas.
arguta, Stev. ex Bieb. Fl. Taur. Cauc. iii. 348 = tomentosa.
argute-serrata, Opiz, in Lotos, iv. (1854) 46 (Quid?).—Europ.
aristata, Lapeyr. Fl. Pyr. t. 105 = cinnamomea.
arkansana, *Port. & Coult. Fl. Colorado*, 38.—Am. bor. occ.
armata, Stev. ex Bess. Enum. Pl. Volh. 60 = canina.
armatissima, *Déségl. & Ripart, in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 114.—Gallia.
armena, *Boiss. Fl. Orient.* ii. 674.—Armenia.
Armidae, Steud. Nom. ed. II. ii. 465 = canina.

ROSA:—
armoricana, *Boullu, in Ann. Soc. Bot. Lyon*, ix. (1882) 328.—Gallia.
arvatica, Puget, ex Baker, in Naturalist, i. (1864) 101 = canina.
arvensis, *Huds. Fl. Angl.* ed. I. 192.—Europ.
arvensis, Lucé, Fl. Osil. 169 = tomentosa.
arvensis, Roth, Tent. Fl. Germ. ii. 554 = canina.
arvina, *Krock. Fl. Siles.* ii. 1. 150 (Quid?).—Europ.
arvina, Lloyd, Fl. Ouest Fr. ed. II. 180 = gallica.
aspera, *Schleich. Cat. Pl. Helv.* ed. IV. 29.—Helvet.
aspernata, Déségl. in Journ. Bot. xii. (1874) 171 = canina.
asperrima, *Godet, ex Boiss. Fl. Orient.* ii. 678.—Persia.
aspratilis, Crép. ex Déségl. in Journ. Bot. xii. (1874) 171 = canina.
aspreiicola, Gremli, Excurs. Schw. ed. II. 177 = canina?
assimilis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 103 = gallica.
atrichostyla, *Gren. in Mém. Soc. Emul. Doubs*, Sér. IV. ix. (1875) 426.—Gallia.
atropurpurea, *Boullu, in Ann. Soc. Bot. Lyon*, ix. (1882) 274.—Gallia.
atropurpurea, *Brot. Fl. Lusit.* ii. 488.—Lusitan.
atrovirens, Viv. Fl. Ital. Fragm. i. 4. t. 6 = sempervirens.
Aucheri, *Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 344.—Persia.
Aucheriana, Crép. in Heurck, Obs. Bot. 30 = praec.
Aunieri, Cariot, Etr. Fl. ed. IV. ii. 180 = canina.
aurantiaca, Hort. ex Steud. Nom. ed. I. 695 = R. Eglanteria.
aurelianensis, Red. ex Link, Handb. ii. 93 = gallica.
australis, Kern. ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 244 = pomifera.
austriaca, Crantz, Stirp. Austr. ed. I. fasc. ii. 33 (err. typ. 36); ed. II. 86 = gallica.
austriaca, Hort. ex C. Koch, Dendrol. i. 225 = R. Eglanteria.
Axmanni, C. C. Gmel. Fl. Bad. iv. 357 = gallica.
ayreshirea, Hort. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 211, in obs. = arvensis.
baicalensis, *Turcz. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 12.—Sibir. baical.
Bakeri, Déségl. in Journ. Bot. ii. (1864) 267 = canina.
Balansaea, *Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 553.—As. Min.
baldensis, Kern. ex Déségl. l. c. 217, in obs. = arvensis.
balearica, Desf. Tabl. ed. I. 175; Pers. Syn. ii. 49 = sempervirens.
ballonianum, Herm. in Kirsch. Fl. Alsat. i. 245 = spinosissima.
balsamea, Bess. Cat. Hort. Crem. (1811) Suppl. iv. 18 = sepium.
balsamica, Willd. Enum. Hort. Berol. Suppl. 38 = canina.
baltica, *Roth, Nov. Pl. Sp.* 234.—Europ.
Banksiae, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iii. 256.—China.
Banksiana, Abel, Narr. Journ. China, 160 = praec.
barbulata, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 430 = canina.
basilica, Roess. Ros. i. 184 = gallica.
batavica, Breit. Hort. Breit. 416 (Quid?).—Hab.?
Beatricis, *Burn. & Gremli, Suppl. Mon. Roses Alpes-Marit.* 14.—Gallia.
Beggeriana, *Schrenk, Enum. Pl. Nov.* 73.—As. temp.
belgica, Brot. Fl. Lusit. i. 338 = gallica.
belgica, Mill. Gard. Dict. ed. VIII. n. 17 = damascena.
Belladonna, Hort. ex Andr. Ros. iii. 46, in obs. = gallica.
bellevallis, Puget, ex Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 426 = canina.
bengalensis, Pers. Syn. ii. 50 = indica.
berberifolia, *Pall. in Nov. Act. Acad. Petrop.* x. (1792) 379 t. 10. f. 5.—Persia.
Berneti, *Schmidely, in Ann. Soc. Bot. Lyon*, vii. (1880) 180.—Europ.
Besseri, Tratt. Ros. ii. 128 = spinosissima.
bibracteata, Bast. in DC. Fl. Fr. v. Suppl. 537 = arvensis.
bicolor, Jacq. Hort. Vindob. i. 1 = R. Eglanteria.
bicolor, Opiz, in Lotos, iv. (1854) 70 (= praec.?).—Europ.
Biebersteiniana, Tratt. Ros. ii. 5 = canina.
Biebersteinii, Lindl. in Loud. Encyc. Pl. 444 = ferox.

ROSA :—
bifera, Poir. Encyc. vi. 276 = damascena.
biflora, Aubl. Pl. Gui. i. 514 = centifolia ?
biflora, Krock. Fl. Siles. ii. 1. 151 = alpina.
Billietii, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 337 = agrestis.
bipinnata, Dum.-Cours. Bot. Cult. ed. I. iii. 351 = centifolia.
bipinnata, Hort. ex Steud. Nom. ed. I. 695 = arvina.
biserrata, Mérat, Fl. Par. 190 = canina.
biturigensis, Bor. Fl. Centr. Fr. ed. II. ii. 630 = spinosissima.
blanda, *Ait. Hort. Kew.* ed. I. ii. 202.—Am. bor.
blanda, Pursh, Fl. Am. Sept. i. 344 = nitida.
Blondaeana, Ripart, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 133 = canina.
Boissieri, *Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 340. —As. Min.
Bonnieri, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 295 = alpicola.
Boraeana, Béraud, in Mém. Soc. Agric. Angers, v. (1842) 353 = gallica.
Borbasiana, *H. Braun, in Flora*, lxviii. (1885) 114.—Europ.
Borboniana, Desp. Ros. Gall. 106 = indica.
Borckhausenii, Steud. Nom. ed. I. 695 = canina.
Boraeeana, Béraud, ex Bor. Fl. Centr. Fr. ed. III. ii. 216 = gallica.
borealis, Tratt. Ros. ii. 141 = spinosissima.
Boreykiana, Bess. Enum. Pl. Volh. 61 = alba.
Borkhausenii, Tratt. Ros. i. 114 = tomentosa.
Borkhausii, Gaud. ex Ser. Mél. Bot. i. 41 = canina.
Borreri, Woods, in Trans. Linn. Soc. xii. (1817) 210 = canina.
Bourbonia, Roess. Rosen, ii. 28 = gallica.
Bourgeauiana, Crép. Prim. Mon. Bor. 303 = acicularis.
Boursoultii ×, Sweet, Hort. Brit. ed. I. 138, nomen.
Bovernieriana, Lag. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 392 = canina.
brachiata, *Déségl. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 103.—Europ.
brachyantha, Opiz, in Lotos, iv. (1854) 46 (Quid ?).—Europ.
brachypoda, *Déségl. & Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 339.—Gallia.
bracteata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = involucrata.
bracteata, *Wendl. Bot. Beobacht.* 50.—China.
bractescens, Woods, in Trans. Linn. Soc. xii. (1817) 216 = canina.
Brandisii, Keller, in Oestr. Bot. Zeitschr. xxxiv. (1884) 12, 172 = Sabini ?
Braniciana, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 327, nomen.—Rossia.
Braunii, *Keller, in Oestr. Bot. Zeitschr.* xxxii. (1882) 39.—Austria.
brevipedunculata, Opiz, in Lotos, ix. (1854) 46 (Quid ?).—Europ.
brevistyla, DC. Fl. Fr. Suppl. 537 = stylosa.
britannica, Déségl. ex Baker, in Journ. Linn. Soc. xi. (1871) 218 = tomentosa.
Bronneri, *Dierb. in Flora*, xi. (1828) 1. *Erg.* 108.—Europ.
Broteri, Tratt. Ros. ii. 98 = moschata.
Brotheri, *Scheutz, in Bot. Centralb.* ii. (1880) 1245.—Reg. Caucas.
Brownii, Tratt. Ros. ii. 96 = moschata.
brunica, *Bouv. in Bull. Inst. Nat. Genèv.* xix. (1875) 92.—Europ.
Brunneri, Dierb. in Flora, xi. (1828) Index, Erg. 124, sphalm. = Bronneri.
Brunonii, Lindl. Ros. Monog. 120. t. 14 = moschata.
Brunonis, Wall. Cat. n. 689 = praec.
bullata, Savi, Fl. Ital. i. t. 34 = centifolia.
Bungeana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 84 = hemisphaerica, *Herrm.*
burgundiaca, Ehrh. Beitr. vi. 97; Pers. Syn. ii. 48 = centifolia.
burgundica, Gmel. ex Steud. Nom. ed. II. ii. 466 = parvifolia.
Burnati, Burnat. & Gremli, Roses Alpes-Marit. 92 = canina.
caballicensis, Puget, ex Déségl. in Billotia, i. (1864) 35 = montana.
cabulica, *Boiss. Fl. Orient.* ii. 678.—Afghan.
Caesarea, Boiss. l. c. 672 = hemisphaerica, *Herrm.*
caesia, Sm. Engl. Bot. t. 2367 = canina.

ROSA :—
calabrica, Huter, ex Burnat. & Gremli, Roses Alpes-Marit. 79 = rubiginosa.
Caledoniae, Borrer, ex Baker, in Journ. Linn. Soc. xi. (1871) 234 = canina.
calendarum, Borck. Forstbot. 330 = damascena.
californica, *Schlecht. in Linnaea*, ii. (1827) 35.—Calif.
calvatostyla, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 429 = canina.
calycina, Bieb. Fl. Taur. Cauc. iii. 349 = canina.
calycina, Moench, Verz. 124 (Quid ?).—Europ.
camberiensis, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 342 = rubiginosa.
campaniaca, *Roess. Rosen*, ii. 24.—Hab. ?
campanulata, Ehrh. Beitr. vi. 97 = turbinata.
campestris, Dum. Fl. Belg. Prod. 93 = trachyphylla.
campestris, Sw. Summa Pl. 18 = canina.
campestris, Wallr. Ros. Pl. 110 = spinosissima.
camtschatica, Erman, Reise, iii. 303 = nutkana.
camtschatica, Steud. Nom. ed. I. 696 = *kamtschatica*.
canariensis, *Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 535, in obs.—Ins. Canar.
candida, Scop. Fl. Carn. ed. II. i. 354 = arvensis.
Candolleana, Thory, Ros. Candoll. 6 = spinosissima.
canescens, Baker, in Naturalist, i. (1864) 97 = canina.
canescens, *Krock. Fl. Siles.* ii. 1. 153.—Europ.
canina, *Linn. Sp. Pl.* 492.—Europ.; As. temp.
canina, Siev. ex Ledeb. Fl. Alt. ii. 229 = Gmelini.
canina, Soland. ex Bab. in Journ. Linn. Soc. xi. (1871) 304 = spinosissima.
cantabrica, Crép. in Willk. & Lange, Prod. Fl. Hisp. iii. 210 = stylosa.
capnoides, Kern. ex Déségl in Bull. Soc. Bot. Belg. xv. (1876) 565 = tomentosa.
capreolata, Neill, in Edinb. Phil. Journ. (1820) 104 = arvensis.
carbonariensis, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxvii.—Gallia.
cardinalis, Lawr. Ros. t. 59 = gallica.
carelica, Fries, in Fl. Dan. Suppl. ii. 6. t. 76 = acicularis.
carelica, *Fries, Summa Veg. Scand.* i. 171, 556.—Europ.
Carionii, Déségl. & Gillot, in Bull. Soc. Bot. Belg. xix. (1880) 34 = canina.
Carioti, Chabert, in Cariot, Etud. Fleurs, ii. (1865) 677 = canina.
carissima, *Roess. Rosen*, 68.—Hab. ?
carnea, Dum.-Cours. Bot. Cult. ed. II. v. 478 = gallica.
carnea, Hort. ex Léman, in Journ. Phys. lxxxvii. (1818) 366 = centifolia.
Carolinà, Duroi, Harbk. Baumz. ii. 355; Walt. Fl. Carol. 149 = humilis.
Carolina, *Linn. Sp. Pl.* 492.—Am. bor.
carolinensis, Marsh, Arbust. 135 = R. Carolina.
caroliniana, Michx. Fl. Bor. Am. i. 295 = humilis.
carpatica, *Kit. in Linnaea*, xxxii. (1863) 592.—Europ.
caryophyllacea, Bess. Enum. Pl. Volh. 19 = tomentosa.
caryophyllacea, Christ, in Gremli, Excurs. Schw. ed. IV. 164 = rubiginosa.
caryophyllata, Hort. ex Red. Ros. i. 114, in obs. = centifolia.
caryophyllea, Poir. Encyc. vi. 276 = centifolia.
casta, Tratt. ex Link, Handb. ii. 85 = suavis.
catalaunica, Costa, Supl. Fl. Catal. 26 = canina.
caucasea, *Lindl. Ros. Monog.* 97. t. 11.—Reg. Caucas.
caucasica, Pall. Pl. Ross. i. 11. 62 = canina.
celerata, Baker, in Naturalist, i. (1864) 99 = canina.
Celsiana, Hort. ex C. Koch, Dendrol. i. 252 = damascena.
celsicola, Arv. Ess. Pl. Dauph. (1871) 27 = glauca.
Celsii, *Léman, in Journ. Phys.* lxxxvii. (1818) 367 (nomen).—Gallia.
centifolia, *Linn. Sp. Pl.* 491.—Reg. Caucas.
centifolia, Lour. Fl. Cochinch. 323 (Quid ?).—China.
centipetala, Stokes, Bot. Mat. Med. iii. 133 = praec.
cerasifera, Timb. in Bull. Soc. Bot. Fr. xi. (1864) 146 = collina.
cerea, Roess. ex Red. Ros. i. 69 = R. Eglanteria.
Chaberti, *Déségl. in Cariot, Etud. Fleurs*, ii. (1865) 180.—Europ.
Chaboissaei, Gren. Fl. Jurass. 241 = canina.
Chailletii, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 339 = rubiginosa.
Chamaerhodon, Vill. Hist. Pl. Dauph. iii. 555 = gallica.
Chapusii, Godet, Fl. Jura, Suppl. 73 = canina.

ROSA :—
Chavini, Rapin, ex Reut. Cat. Pl. Vasc. Genèv. ed II. 69 = canina.
cheriensis, Déségl. in Billotia, i. (1864) 45 = agrestis.
cherokeensis, Donn, Hort. Cantab. ed. VI. 144 = laevigata.
chinensis, Jacq. Obs. Bot. iii. 7 = indica.
chlorophylla, Ehrh. Beitr. ii. 69 = R. Eglanteria.
Christiana ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.–xxiv. (1880) 62.—Europ.
Christii ×, Wiesb. in Oestr. Bot. Zeitschr. xxix. (1879) 146.—Europ.
ciliaris, Wils. ex Baker, in Journ. Linn. Soc. xi. (1871) 229 = canina.
ciliata, *Link, Handb.* ii. 90.—Europ.
ciliato-petala, Bess. Enum. Pl. Volh. 66 = pomifera.
ciliato-petala, Koch, Syn. Fl. Germ. ed II. 253 = mollissima.
cinerascens, Cariot, Etud. Fleurs, ii. (1865) 677 = canina.
cinerascens, Dum. Fl. Belg. Prod. 93 = tomentosa.
cinerea, Rapin, in Gren. Fl. Jurass. 238 = canina.
cinerea, Sw. in Svensk Bot viii. t. 553 = cinnamomea.
cinerosa, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 380 = canina.
cinnabarina, Wrede, ex Breit. Hort. Breit. 419 (Quid ?).—Hab. ?
cinnamomea, Kar. & Kir. ex Déségl. in Journ. Bot. xii. (1874) 168 = dissimilis.
cinnamomea, Herrm. Diss. 7 = alpina.
cinnamomea, *Linn. Sp. Pl.* 491.—Europ.; As. bor.
cinnamomea, Roth, Tent. Fl. Germ. i. 214 = R. Eglanteria.
ciphiana, Sm. Fl. Brit. ii. 537 = spinosissima.
cladoleia, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 265 = canina.
clinophylla, *Thory, in Red. Ros.* i. 43.—Ind. or.
clivorum, Scheutz, Stud. Skand. Ros. (1872) 28 = tomentosa.
clotildea, Timb. ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 260 = stylosa.
Clusiana, Bouv. in Bull. Soc. Bot. Fr. xiii. (1866) p. xxiv = pomifera, alpina.
Clusiana, *Waitz, ex Tratt. Ros.* i. 119.—Europ.
coccinea, Wrede, ex Steud. Nom. ed I. 697 (Quid ?).—Hab. ?
coerulescens, Kern. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 293 = ferruginea.
coerulescens, Link, Handb. ii. 83 (Quid ?).—Hab. ?
cochinchinensis, *G. Don, Gen. Syst.* ii. 585.—Cochinch.
collina, Jacq. Fl. Austr. ii. 58 = canina.
collina, Sm. Engl. Bot. t. 1895 = stylosa.
collincola, Ehrh. Beitr. ii. 70 = cinnamomea.
collivaga, Cottet, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 247, 297 = tomentosa.
columnifera, Borck. ex Reichb. Fl. Germ. Excurs. 621 = sylvestris.
commutata, Scheutz, Stud. Skand. Ros. (1872) 41 = flexuosa.
comosa, Ripart, in F. Schultz, Arch. Fl. Fr. et Allem. 254 = rubiginosa.
complicata, Gren. Fl. Jurass. 239 = glauca.
concavifolia, *Waitz, ex Tratt. Ros.* ii. 66.—Hab. ?
concinna, Hort. Humb. (1834) ex Steud. Nom. ed II. ii. 467 (Quid ?).—Hab. ?
concinna, Lagg. & Puget, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 494 = canina.
condensata, Puget, ex Déségl. in Billotia, i. (1864) 118 = canina.
conferta, Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 28 = tomentosa.
confusa, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 246 = tomentosa.
conica, Chabert, ex Cariot, Etud. Fleurs, ii. (1865) 171 = arvina.
consanguinea, Gren. Fl. Jurass. 225 = trachyphylla.
consimilis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 90 = spinosissima.
conspicua, Bor. l. c. xii. (1862) 55 = arvensis.
controversa, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 499 = canina.
corallina, Willd. Enum. Hort. Berol. Suppl. 38 = humilis.
corbariensis, Debeaux, in Act. Linn. Soc. Bord. xxx. (1875) p. clxi = sepium.
cordata, *Cariot, Etud. Fleurs*, ed. VII. ii. (1884) 228.—Gallia.
cordifolia, Baker, in Naturalist, i. (1864) 24 = glauca.

ROSA:—
cordifolia, Host, Fl. Austr. ii. 23 = gallica.
coriacea, Crép. in Heurck, Obs. Bot. 30 = canina.
coriacea, *Opiz, in Flora*, v. (1822) 268.—Europ.
coriifolia, Fries, Novit. Fl. Suec. ed. I. (1814) 33; ed. II. 147 = canina.
coronata, Crép. Man. Fl. Belg. 52, in nota = involuta.
corrugata, *Caldesi, in Nuov. Giorn. Bot. Ital.* xii. (1880) 100.—Europ.
coruscans, Waitz, ex Link, Enum. Hort. Berol. ii. 57 = rugosa.
corylifolia, *Vukot. ex Borb. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 16.—Croatia.
corymbifera, Borck. Holzart. 319 = canina.
corymbosa, Bast. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 213 = arvensis.
corymbosa, Bosc, ex Desf. Cat. Hort. Par. ed. III. 272 = blanda.
corymbosa, Breit. Hort. Breit. 418 = perfecta.
corymbosa, Ehrh. Beitr. iv. 21 = R. Carolina.
corymbosa, *Moench, Verz.* 119.—Europ.
Cotteti, Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 31 = trachyphylla.
Crantzii, Schultz, Obs. Bot. 94 = rubiginosa.
crassifolia, Wallm. in Liljebl. Svensk Fl. ed. III. 268 = canina.
cremsensis, Kern. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 578 = mollissima.
Crepini, Miègev. ex Crép. l. c. viii. (1869) 274 = glauca.
Crepiniana, Déségl. ex Baker, in Naturalist, i. (1864) 97 = canina.
cretacea, *Kalenicz. in Bull. Soc. Nat. Mosc.* xxii. (1849) i. 297, 301, nomen.—Rossia.
cretica, Bosc, Encyc. Agric. vii. 705 (Quid?).—Ins. Creta.
cretica, Thory, Ros. (1820) 110 = rubiginosa.
cretica, Vest, ex Tratt. Ros. ii. 83 = glutinosa.
cretica, Wallr. Ros. 144, partim = alpina.
crispa, Hort. ex Red. Ros. ii. 12 = arvina.
croatica, Kit. in Linnaea, xxxii. (1863) 589 = gallica.
cryptopoda, Baker, in Naturalist, i. (1864) 66 = agrestis.
cryptopoda, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 428 = canina.
cucumerina, Tratt. Ros. ii. 181 = laevigata.
cuprea, Jacq. Fragm. 31 = gallica.
Cursor, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 217 = setigera.
curticola, *Puget, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 114.—Europ.
curvula, Timb. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 367 = canina.
cuspidata, Bieb. Fl. Taur. Cauc. i. 396 = tomentosa.
cuspidatoides, Crép. in Scheutz, Stud. Skand. Ros. (1872) 37 = tomentosa.
cymbifolia, *Léman, in Journ. Phys.* lxxxvii. (1818) 366.—Gallia.
cymosa, Tratt. Ros. i. 87 = microcarpa.
Cynorrhodon, Wallr. Ros. 206; ex Red. Ros. ii. 51 = canina.
Czackiana, Bess. Enum. Pl. Volh. 61 = gallica.
daënensis, Boiss. Diagn. Ser. I. vii. 51; Fl. Orient. ii. 677 = Beggeriana.
dahurica, Auct. = davurica.
dalmatica, *Kern. in Oestr. Bot. Zeitschr.* xx. (1870) 8.—Europ.
damascena, *Mill. Gard. Dict.* ed. VIII. n. 15.—Oriens.
dasistema, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 219 = blanda.
Davidi, *Crép. in Bull. Soc. Bot. Belg.* xiii. (1874) 253.—China.
davurica, *Pall. Fl. Ross.* i. II. 61.—Dahuria.
decipiens, Bor. Fl. Centr. Fr. ed. III. ii. 218 = gallica.
decora, *Kern. ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 512 = flexuosa.
De-la-Soiei, Lagg. & Puget, in Bull. Soc. Murith. ii. (1873) 43 = glauca.
Dematranea, *Lagg. & Puget, ex Cottet, l. c.* iv. (1875) 32.—Helvet.
densa, Timb. in Bull. Soc. Bot. Fr. xi. (1864) 141 = rubiginosa.
densiflora, Tausch, ex Opiz, Boehm. Gew. (1823) 61, nomen (Quid?).—Bohem.
depressa, Gremli, Excurs. Schw. ed. II. 178 = trachyphylla.
Deseglisei, Bor. Fl. Centr. Fr. ed. III. ii. 224 = canina.
Desvauxii, *Desp. Ros. Gall.* 107.—Hab.?

ROSA:—
devoniensis ×, Hort.—Cf. Gard. Chron. (1841) 470.
dianthiflora, *Andr. Ros.* t. 31.—Europ.
dibracteata, Thory, Ros. (1820) 135 = arvensis.
dichroa, Lerch, in Oestr. Bot. Zeitschr. xxii. (1872) 145 = spinosissima.
Dicksoni, *Lindl. in Trans. Hort. Soc.* vii. (1830) 224.—Hibernia.
didoënsis, *Boiss. Fl. Orient.* ii. 685.—Reg. Caucas.
diffusa, Roxb. Hort. Beng. 92; Fl. Ind. ii. 515 = multiflora.
digitaria, Bosc, ex DC. Prod. ii. 622 = alba.
dijonensis, *Roess. Ros.* i. 93.—Gallia.
Dilleniana, *Baker, ex Déségl. in Journ. Bot.* v. (1867) 42.—Europ.
diminuta, Bor. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 546 = rubiginosa.
dimorpha, Bess. Cat. Hort. Crem. (1811) Suppl. iii. 19 = tomentosa.
dimorphacantha, Martin, ex March. & Hardy, in Bull. Soc. Bot. Belg. vii. (1868) 248 = rubiginosa.
Diomedis, Gren. ex Déségl. in Billotia, i. (1864) 121 = canina.
discreta, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 237 = glauca.
dissimilis, *Déségl. in Journ. Bot.* xii. (1874) 168.—Rossia.
distans, Christ, in Gremli, Excurs. Schw. ed. IV. 170 = tomentosa.
diversifolia, Vent. Jard. Cels, 35 = indica.
diversisepala, *H. Braun, in Beck, Fl. Hernstein*, 353.—Europ.
djimilensis, *Boiss. Fl. Orient.* ii. 673.—As. Min.
dolosa, Godet, Fl. Jura, Suppl. 72 = canina.
Doniana, Woods, in Trans. Linn. Soc. xii. (1817) 185 = involuta.
drosophora, *H. Braun, ex Kern. Sched. Fl. Exsicc.* iii. 56.—Tyrol.
druentica, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 344.—Gallia.
dryadea, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 502 = trachyphylla.
dubia, Bastard, ex Déségl. l. c. 353 = canina.
dubia, Carr. in Rev. Hortic. (1868) 269 = thyrsiflora.
dubia, Wibel, Prim. Fl. Werth. 263 = tomentosa.
Dufftii ×, Schultz, ex Haussk. in Mitth. Geogr. Ges. Jena, ii. (1885) 24.—Europ.
dumalis, Bechst. Forstbot. ed. V. (1843) 582 = canina.
dumalis, Tratt. ex Link, Handb. ii. 94 = gallica.
dumetorum, Lange, in Fl. Dan. t. 2718 = seq.?
dumetorum, Sm. Engl. Bot. t. 2579 = inodora.
dumetorum, Thuill. Fl. Par. ed. II. 250 = canina.
dumosa, Puget, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 101 = tomentosa.
dumosa, Salisb. Prod. 359 = canina.
Dupontii, *Déségl. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 58.—Europ.
Durandii, Crép. in Bull. Soc. Bot. Fr. xxii. (1875) 19 = nutkana.
Ecae, Aitch. in Journ. Linn. Soc. xviii. (1880) 54 = xanthina.
echinata, Dupont, ex DC. Prod. ii. 607 = cinnamomea.
echinata, *Waitz, ex Link, Enum. Hort. Berol.* ii. 60, nomen.—Hab.?
echinocarpa, Ripart, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 150 = macrantha.
edita, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 347 = canina.
editorum, Ripart, ex Déségl. l. c. 286 = alpina.
Eglanteria, *Linn. Sp. Pl.* 491.—Europ.; Oriens.
Eglanteria, Mill. Gard. Dict. ed. VIII. n. 4 = rubiginosa.
elasmacantha, *Trautv. Ind. Sem. Hort. Petrop.* (1868) 25.—As. bor.
elata, Christ, in Gremli, Excurs. Schw. ed. IV. 163 = gallica.
elatior, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 297 = sepium.
elegans, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 214 = R. Carolina.
elegans, Salisb. Prod. 358 = damascena.
elliptica, *Tausch, ex Tratt. Ros.* ii. 69; *et in Opiz, Boehm. Gew.* (1823) 60.—Europ.
elongata, Dum.-Cours. Bot. Cult. ed. II. v. 484 (err. typ. 448) = multiflora.
elongata, Roess. ex Steud. Nom. ed. II. ii. 468 = R. Carolina.

ROSA:—
elymaitica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 675.—Persia.
engolismensis, *Déségl. & Guillon, in Ann. Soc. Bot. Lyon*, ix. (1882) 6.—Europ.
enneaphylla, Rafin. Préc. Découv. 35 = R. Carolina.
episcopalis, Hort. ex Roess. Rosen, 239 = gallica.
erinacea, *Desp. Ros. Gall.* 18.—Gallia.
erinacea, Hardy, ex Nym. Consp. 236 = involuta.
eriocarpa, *Léman, in Journ. Phys.* lxxxvii. (1818) 367, nomen.—Hab.?
eriosa, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 570 = tomentosa.
eriostyla, Ripart & Déségl. l. c. 334 = canina.
erratica, Ripart, in Soc. Dauph. Ech. Pl. iv. (1877) 109 = stylosa.
erronea, Ripart, ex Crép. Bull. Soc. Bot. Belg. viii. (1869) 257 = arvensis.
erubescens, *Andr. Ros.* t. 18.—Hab.?
erythrantha, Bor. Fl. Centr. Fr. ed. III. ii. 224 = canina.
erythrella, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 330 = canina.
Euratiana, Bosc, ex Link, Handb. ii. 87 = seq.?
Evratina, Bosc, ex Poir. Encyc. Suppl. iv. 714 = R. Carolina, rubiginosa, villosa.
exilis, [*Crép.*] *ex Boutt. in Bull. Soc. Bot. Belg.* vii. (1868) 220.—Europ.
falcata, Puget, ex Crép. l. c. viii. (1869) 237 = montana.
fallaciosa, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 391 = canina.
fallax, Gren. ex Gren. & Godr. Fl. Fr. i. 561 = graveolens.
fallax, Puget, ex Baker, in Journ. Linn. Soc. xi. (1869) 226 = canina.
fallens, Déségl. in Lamotte, Prod. Fl. Centr. Fr. i. 266 = canina.
farinosa, Bechst. Forstbot. 643; Rau, Enum. Ros. 147 = tomentosa.
farinulenta, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 246 = tomentosa.
fastigiata, Bast. Fl. Maine-et-Loire, Suppl. 30 = stylosa.
fastigiata, Salisb. Prod. 358 = arvensis.
fastuosa, Hort. ex C. Koch, Dendrol. i. 239 = rugosa.
Favrati, Christ, in Gremli, Excurs. Schw. ed. IV. 170 = tomentosa.
Fedtschenkoana, *Regel, in Del. Sem. Hort. Petrop.* (1876) 36, nomen.—Turkestan.
Fendleri, *Crép. in Bull. Soc. Bot. Belg.* xv. (1876) 91.—Calif.
fenestralis, Hort. ex Red. Ros. iii. 72 = ferruginea.
fenestrata, Donn, Hort. Cantab. ed. VIII. 170 = setigera.
ferox, Ait. Hort. Kew. ed. II. iii. 262 = rugosa.
ferox, *Bieb. Fl. Taur. Cauc.* iii. 339.—As. Min.; Persia.
ferox, Lawr. Ros. t. 42 = cinnamomea.
ferruginea, *Vill. Prosp.* 46.—Europ.
fimbriata, Gremli, Excurs. Schw. ed. II. 178 = tomentosa.
finitima, Déségl. ex Cottet, in Bull. Soc. Murith. iii. (1874) 39 = canina.
firma, Puget, ex Déségl. in Billotia, i. (1869) 118 = canina.
firmula, Godet, Fl. Jura, Suppl. (1874) 38 = canina.
Fischeri, *Link, Handb.* ii. 88 (= seq.?).—Sibir.
Fischeriana, *Bess. ex Link, Enum. Hort. Berol.* ii. 60, nomen.—Rossia.
fissispina, Wierzb. ex Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 102 = canina.
flagellaris, Christ, in Gremli, Excurs. Schw. ed. II. 176 = ferox.
flava, *Donn, Hort. Cantabr.* ed. IV. 121 (Quid?).—China.
flava, *Wickstr. in Vet. Akad. Handl. Stockh.* (1820) 273.—Tauria.
flavescens, Hassk. Cat. Pl. Hort. Bog. Alt. 265 = arvina.
flexibilis, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 317 = canina.
flexicaulis, Godet, ex Boiss. Fl. Orient. ii. 675 = albicans.
flexuosa, Rafin. Préc. Decouv. 35 = canina.
flexuosa, *Rau, Enum. Ros.* 127.—Europ.
floccida, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 567 = tomentosa.

ROSA:—
floribunda, Bess. Cat. Hort. Crem. (1816) Suppl. iv. 19 = rubiginosa.
floribunda, Hort. ex Andr. Ros. t. 106 = multiflora.
florida, Donn, Hort. Cantab. ed. VIII. 159 = R. Carolina.
florida, Poir. Encyc. Suppl. iv. 715 = multiflora.
fluvialis, O. F. Muell. in Fl. Dan. t. 868 = cinnamomea.
foecundissima, Muench. Hausv. v. (1770) 279 = cinnamomea.
foetida, Bast. Fl. Maine-et-Loire, Suppl. 29 = tomentosa.
foetida, Herrm. Diss. 18 = R. Eglanteria.
foliolosa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 460.—Am. bor.
foliosa, *Léman, in Journ. Phys.* lxxxvii. (1818) 366.—Gallia.
foliolosa, Nutt. l. c. partim = carolina.
Fontanesii, Pomel, Nouv. Mat. Fl. Atl. 157 = agrestis.
formosa, Roess. ex Red. Ros. i. 74 = gallica.
formosula, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 437 = sepium.
Forsteri, Sm. Engl. Fl. ii. 392 = canina.
Fortuneana, Lem. Jard. Fleur. ii. (1852) Misc. 26 = seq.
Fortuniana ×, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 71.—China.
Fourroei, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 98 = gallica.
fragariaeflora, Ser. ex DC. Prod. ii. 601 = Banksiae ?
fragariaefolia, Steud. Nom. ed. II. ii. 468, sphalm. = praec.
fragrans, *Desp. Ros. Gall.* 101.—Hab. ?
fragrans, Red. Ros. i. t. 19 = indica.
fragrans, Salisb. Prod. 358 = R. Carolina.
francofurtana, Muench, Hausv. v. (1770) 288; Borck. Forst. Beschr. Holzart. 312 = turbinata.
francofurtensis, Desf. Tabl. ed. I. 175 = praec.
Franzonii, Christ, Rosen Schweiz (1875), 174 = ferruginea.
Fraseriana ×, Hort. ex G. Don, in Loud. Hort. Brit. 211.
fraxinea, Willd. Enum. Hort. Berol. Suppl. 37 = blanda.
fraxinellaefolia, *Andr. Ros.* t. 100.—Hab. ?
fraxinifolia, Borck. Vers. Forstw. Beschr. Holz. 301 = cinnamomea.
fraxinifolia, Dur. & Hilg. in Pacif. Rail. Rep. v. III. 7 = californica.
fraxinifolia, C. C. Gmel. Fl. Bad. ii. 413 = blanda.
fraxinifolia, Hook. Fl. Bor. Am. i. 199 = nutkana.
fraxinifolia, Roess. Ros. i. 219 = R. Carolina.
friburgensis, Lagg. & Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 248 = pomifera.
Friedlaenderi, Nym. Consp. 235 = seq.
Friedlanderiana, Bess. Enum. Pl. Volh. 46 = canina.
Friesii, *Lagg. & Puget, ex Cottet, in Bull. Soc. Murith.* iv. (1875) 27.—Europ.
Friesii, Scheutz, Stud. Ros. Skand. (1872) 39 = canina.
frondosa, Stev. ex Spreng. Syst. ii. 554 = canina.
frutetorum, Bess. Cat. Hort. Crem. (1811) Suppl. iii. 20; Enum. Pl. Volh. 18 = canina.
fugax, Gren. Fl. Jura (1864) 239 = glauca.
fulgens, Andamow. ex Opiz, Natural. xi. (1826) 449 (Quid ?).—Europ.
fulgens, Christ, in Gremli, Excurs. Schw. ed. IV. 160 = cinnnamomea.
fusca, Moench, Meth. 688 = arvensis.
fusiformis, Crép. in Willk. & Lange, Prod. Fl. Hisp. iii. 214 = canina.
gallica, *Linn. Sp. Pl.* 492.—Europ.; Reg. Caucas.
gallicoides, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 218 = arvensis.
Gallinariae, Burnat & Gremli, Suppl. Mon. Ros. Alpes-Marit. 25 = canina ?
Gandogeriana, Debeaux, in Bull. Soc. Bot. Fr. xxi. (1874) 9 = sempervirens.
Gaudini, Puget, in Déségl. Rev. Sect. Toment. Rosa (1866), 47 = pomifera.
Gebleriana, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 80.—Sibir. altaic.
gemella, Willd. Enum. Hort. Berol. 544 = blanda.
gemina, *Opiz, in Lotos*, iv. (1854) 46.—Europ.
geminata, Rau, Enum. Ros. 98 = gallica.
genevensis, Puget, in Bull. Soc. Bot. Belg. xv. (1876) 567 = tomentosa.

ROSA:—
Gennarii, Huet, ex Gennari, in Mem. Acc. Torin. xxv. (1858) 463 = canina.
gentilis, Sternb. in Flora, ix. (1826) Beil. 79 = spinosissima.
georgina, Hort. Hamb. ex Steud. Nom. ed. II. ii. 469 (Quid ?).—Hab. ?
germanica, Hort. ex Steud. l. c. ed. I. 699 = turbinata.
germanica, Maerkl. ex Steud. l. c. = gallica.
Gillotii, *Déségl. & Lucand. in Ann. Soc. Bot. Lyon*, ix. (1882) 15.—Gallia.
Gisleri, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 247 = tomentosa.
glaberrima, Dum. Fl. Belg. Prod. 94 = canina.
glabra, *Andr. Ros.* t. 40.—Hab. ?
glabra, Baker, in Naturalist, i. (1864) 24 = glauca.
glabra, Meyen, Reise, i. 303 = Meyeniana.
glabrata, *Kit. in Linnaea*, xxxii. (1863) 588.—Europ.
glabrata, Vest, ex Tratt. Ros. ii. 220 = ferruginea.
glabrescens, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 340 = rubiginosa.
glabrifolia, Rupr. Stirp. Fl. Petrop. Diatr. 65 = cinnamomea.
glandulifera, Roxb. Hort. Beng. 38; Fl. Ind. ii. 514 = moschata.
glandulosa, Bell. in Mém. Acad. Turin, v. 1790–1791 (1793) 230 = alpina.
glandulosa, Bess. Cat. Hort. Crem. (1816) 118; Enum. Pl. Volh. 20 = canina.
glandulosa, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 91 = montana.
glandulosa, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 214 = R. Carolina ?
glauca, Dierb. Fl. Heidelb. ii. 140 = arvensis.
glauca, Pourr. in Mém. Acad. Toul. iii. (1788) 326 = ferruginea.
glauca, Schott, in Bess. Enum. Pl. Volh. 60 = canina.
glauca, *Vill. ex Loisel. in Desv. Journ. Bot.* ii. (1809) 336.—Europ.
glaucescens, Desv. in Mérat, Fl. Par. 192 = canina.
glaucescens, Wulf. in Roem. Arch. iii. III. (1805) 376 = ferruginea.
glaucifolia, *Opiz, in Flora*, v. (1822) 268.—Europ.
glaucina, Ripart, ex T. Durand, in Bull. Soc. Bot. Belg. xix. II. (1880) 60 = canina.
glaucophylla, Ehrh. Beitr. ii. 69 = hemisphaerica, *Herrm.*
glaucophylla, Winch, Ess. Geogr. Distr. Northumb. ed. I. 45 = canina.
glauco-pubescens, *Opiz, in Lotos*, iv. (1854) 69.—Europ.
globata, Déségl. ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 279 = canina.
globosa, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 215 = humilis.
globularis, Franch. ex Bor. Fl. Centr. Fr. ed. III. ii. 221 = canina.
glutinosa, Dematra, Ess. Monog. 6 = pomifera.
glutinosa, Schultz, Prod. Fl. Starg. Suppl. i. 27 = agrestis.
glutinosa, *Sibth. & Sm. Fl. Graec. Prod.* i. 348.—Europ.; As. Min.
Gmelini, *Bunge, in Ledeb. Fl. Alt.* ii. 228.—Sibir.
Godeti, Gren. in Godet, Fl. Jura, Suppl. 73 = trachyphylla.
gombensis, Lagg. & Puget, in Bull. Soc. Murith. iii. (1875) 54 = pomifera.
gorenkensis, Fisch. ex Spreng. Syst. iv. Cur. Post. 200 = cinnamomea.
gracilens, Crép. in Willk. & Lange, Prod. Fl. Hisp. iii. 212 = montana.
gracilescens, Gren. ex Déségl. in Billotia, i. (1869) 121 = canina.
gracilis, Woods, in Trans. Linn. Soc. xii. (1817) 186 = involuta.
granata, Red. ex Link, Handb. ii. 93 = gallica.
granatensis, Willk. in Linnaea, xxv. (1852) 24 = spinosissima.
grandiflora, Lindl. Ros. Monog. 55 = spinosissima.
grandiflora, Salisb. Prod. 358 = gallica.
grandiflora, Wallr. Ann. Bot. 66 = pulverulenta.
grandifolia, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 339 = rubiginosa ?
granensis, *Kmet. ex Kern. Sched. Fl. Exsicc.* ii. 34.—Hungar.
grata, Salisb. Prod. 358 = gallica.
graveolens, *Gren. & Godr. Fl. Fr.* i. 560.—Europ.
Gremlii, Christ, in Gremli, Excurs. Schw. ed. II. 176 = rubiginosa.

ROSA:—
Grenierii, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 168 = pomifera.
Grevillei, Sweet, Hort. Brit. ed. I. 138 = multiflora.
grisea, Wrede, ex Steud. Nom. ed. I. 699 (Quid ?).—Hab. ?
grossularifolia, Andr. Ros. t. 30 = gallica.
Guepini, Desv. Fl. Anjou, 325 = canina.
Guilelmi-Waldemarii, Klotzsch, in Bot. Ergeb. Waldem. Reise, 153 = macrophylla.
Guineti, *Schmidely, in Ann. Soc. Bot. Lyon*, vii. (1880) 177.—Gallia.
guttensteinensis, Jacq. f. ex Spreng. Syst. iv. Cur. Post. 200 = ferruginea.
gymnocarpa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 461.—Am. bor.
haberiana, Puget, ex Déségl. in Billotia, i. (1864) 37 = canina.
Hackeliana, Nym. Consp. 233 = Heckeliana.
haematodes, *Boiss. Fl. Orient.* ii. 684.—Reg. Caucas.
Hailstoni, Baker, in Lond. Bot. Exch. Club Rep. (1868) 7 = canina.
Halácsyi, *H. Braun, in Beck, Fl. Hernstein*, 356.—Europ.
hallandica, Scheutz, in Ofvers. Vet. Akad. Foerh. Stockh. (1873) n. II. 46 = canina.
Halleri, Krock. Fl. Siles. ii. I. 150 = arvensis.
Hampeana, *Griseb. ex Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 329.—Hab. ?
Hardii ×, Paxt. Mag. Bot. x. (1843) 195 = involucrata.
hawrana, *Kmet, ex Kern. Sched. Fl. Exsicc.* ii. 38.—Hungar.
Heckeliana, *Tratt. Ros.* ii. 85.—Ital.; Graecia.
Hecleliana, Boiss. Fl. Orient. ii. 680, sphalm. = praec.
Heldreichii, *Boiss. & Reut. Diagn.* Ser. II. ii. 49.—Thessalia.
helvetica, Hall. f. in Roem. Arch. i. II. (1797) 6 = villosa.
hemisphaerica, *Herrm. Diss.* 18.—Phrygia.
hemisphaerica, *Spreng. Syst.* iv. *Cur. Post.* 200.—Hab. ?
hemitricha, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 373 = canina.
Heriteriana, Link, Handb. ii. 84 = alpina.
Herporhodon, Ehrh. Beitr. ii. 71 = arvensis.
heteracantha, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 350.—Sibir.
heterophylla, Timb. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 527 = sepium.
heterophylla, Woods, in Trans. Linn. Soc. xii. (1817) 195 = mollissima.
hetrostyla, Griff. Notul. iv. 454; Ic. Pl. As. iv. t. 574 = moschata.
hibernica, Hook. ex Lindl. Ros. 51 = spinosissima.
hibernica, Sm. Engl. Fl. ii. 393 = glauca.
Hillebrandtii, *Weitenw. in Opiz, Natural.* xi. (1826) 450.—Europ.
Hirciana, *H. Braun, in Oestr. Bot. Zeitschr.* xxxii. (1882) 6.—Europ. austr.
hirta, *H. Braun, Beitr. Kenntn. Gatt. Rosa* (1885), 108.—Europ.
hispanica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 44 = canina.
hispanica, *Mill. Gard. Dict.* ed. VIII. n. 7.—Hispan.
hispida, Hort. ex Poir. Encyc. vi. 286 = villosa.
hispida, Krock. Fl. Siles. ii. I. 152 = alpina.
hispida, Muench. Hausv. v. (1770) 281; Borck. Forstbot. ii. 1332 = tomentosa.
hispida, Red. Ros. 159 = spinosissima.
hispida, Sims, in Bot. Mag. t. 1570 = R. Eglanteria.
hispidocarpa, Chab. in Cariot, Etud. Fleurs, ii. (1865) 677 = R. Carolina.
hispidula, *Ripart, in Bull. Soc. Bot. Belg.* xv. (1876) 386.—Europ.
Histrix, Léman, in Bull. Soc. Philom. (1818) 94 = micrantha.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 153 = macrophylla.
holikensis, *Kmet, in Oestr. Bot. Zeitschr.* xxxiv. (1884) 19.—Austria.
hollandica, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 484 (err. typ. 448) = multiflora.
hollandica, Pers. ex Steud. Nom. ed. I. 699 = centifolia.
hololeia, Ripart, ex Cottet, in Bull. Soc. Murith. iii. (1874) 38 = canina.
holosericea, *Borck. Forstbot.* iii. 1815.—Europ.
holosericea, Duroi, Obs. Bot. 20 = gallica.
holostrica, *Roess. Rosen*, 188.—Europ.

ROSA:—
homolaga, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 426 = canina.
Hookeriana, Wall. Cat. n. 691; Bertol. Misc. Bot. xxiv. 14 = macrophylla.
horrida, Bess. ex Fisch. Cat. Hort. Gorenk. (1812) 66 = ferox.
horrida, Spreng. Syst. ii. 549 = viminea.
horridula, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 323 = canina.
hudsoniana, Thory, in Red. Ros. i. 95 = R. Carolina.
humifusa, Red. ex Reichb. Fl. Germ. Excurs. 623 = gallica.
humifusa, Tratt. Ros. ii. 102 = sempervirens.
humilis, Bess. Cat. Hort. Crem. (1811) Suppl. iv. 19; Enum. Pl. Volh. 20 = canina.
humilis, *Marsh. Arbust. Am.* 136.—Am. bor.
humilis, Tausch, in Flora, ii. (1819) 465 = gallica.
hungarica, *Kern. in Oestr. Bot. Zeitschr.* xix. (1869) 234.—Europ.
hyacinthina ×, Sweet, Hort. Brit. ed. I. 138, nomen.—Hab.?
hybernica, Thory, Ros. 141 = glauca.
hybrida, Schleich. Cat. Pl. Helv. ed. III. 24 = arvina.
hybrida, Vill. Hist. Pl. Dauph. iii. 554 = alpina.
Hystrix, Lindl. Ros. Monog. 129. t. 17 = laevigata.
Ibara, C. Koch, Ind. Sem. Hort. Berol. (1867) App. i. 9, sphalm. = R. Iwara.
iberica, Stev. in Bieb. Fl. Taur. Cauc. iii. 539 = rubiginosa.
Ilseana, *Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 334.—Europ.
imitata, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 120 = canina.
immitis, Déségl. l. c. 97 = stylosa.
imperialis, Hort. ex C. Koch, Dendrol. i. 252 = damascena.
implexa, Gren. Fl. Jurass. 238 = canina.
imponens, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 237 = montana.
incana, Falk, ex Wikstr. in Vet. Akad. Handl. Stockh. (1820) 275 = laxa.
incana, *Kit. ex Tratt. Ros.* i. 135.—Hungar.
incarnata, Mill. Gard. Dict. ed. VIII. n. 19 = gallica.
incerta, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 384 = canina.
inclinata, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 326 = glauca.
incomparabilis, Chab. in Cariot, Etud. Fleurs, ii. (1865) 170 = gallica.
inconsiderata, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 117 = canina.
inconspicua, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 357 = canina.
indica, *Linn. Sp. Pl.* 492.—Ind. or.; China.
inerma, Bertol. in Mem. Acc. Sc. Bologn. Ser. II. iii. (1863) 419 = seq.?
inermis, *Bosc, Encyc. Agric.* vii. 705.—Ind. or.; China.
inermis, Delaun. ex Mill. Gard. Dict. ed. VIII. n. 6 = alpina.
inermis, *Luce, Topog. Nachr. Oesel*, 170.—Ins. Balt.
inermis, Roxb. Hort. Beng. 38; Fl. Ind. ii. 516 = Banksiae.
infesta, *Kmet, ex Kern. Sched. Fl. Exsicc.* ii. 32.—Hungar.
inodora, Fries, Nov. Fl. Suec. 9 = agrestis.
inodora, Hook. in Curt. Fl. Lond. ed. II. t. 117 = canina.
insidiosa, Gren. Fl. Jurass. 233 = tomentosa.
insidiosa, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 501 = trachyphylla.
insignis, Déségl. & Ripart, in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 112 = canina.
intercalaris, Déségl. l. c. 104 = alpina.
intermedia, *Bosc, Encyc. Agric.* vii. 705.—Europ.
intermedia, Carr. Rev. Hortic. xvii. (1868) 270 = thyrsiflora.
intermedia, Chab. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 240 = arvina.
intermedia, Desv. ex Link, Handb. ii. 91; Kit. in Linnaea, xxxii. (1863) 591 = canina.
intermissa, Nym. Consp. 232, sphalm. = *intromissa*.
interveniens, *Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 354.—Europ.
intricata, Crép. l. c. viii. (1869) 298 = tomentosa.
intricata, Gren. Fl. Jurass. (1864) 239 = glauca.
intromissa, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 298 = tomentosa.

ROSA:—
involucrata, Braam, Ic. Pl. in Chin. Nasc. t. 17 = bracteata.
involucrata, *Roxb. Hort. Beng.* 38; *Fl. Ind.* ii. 513.—Ind. or.; China.
involuta, *Sm. Fl. Brit.* iii. 1398; *Engl. Fl.* ii. 377.—Europ.
irregularis, *Déségl. & Guillon, in Ann. Soc. Bot. Lyon*, ix. (1882) 7.—Europ.
Isaurae, Tratt. Ros. ii. 72 = pulverulenta.
Iwara, *Sieber, ex Regel, Ind. Sem. Hort. Petrop.* (1861) 53.—Japon.
Jacquemontii, *Crép. ex Hook. f. Fl. Brit. Ind.* ii. 368.—Reg. Himal.
jactata, Déségl. in Journ. Bot. xii. (1874) 169 = canina.
japonica, *Waitz, ex Steud. Nom.* ed. I. 699.—Japon.
Jordani, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 146 = sepium.
Jundzilli, Bess. Cat. Pl. Hort. Crem. (1816) 117 = canina?
Jundzilliana, Baker, in Naturalist, i. (1864) 65 = tomentosa.
Jundzilliana, Bess. ex Enum. Pl. Volh. 61, 67 = canina.
jurana, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 344 = rubiginosa.
kalksburgensis ×, Wiesb. in Oestr. Bot. Zeitschr. xxix. (1879) 144.—Europ.
kamtchatica, Vent, Jard. Cels, t. 67 = cinnamomea.
kaschgarica, *Rupr. Sert. Tiansch.* 46.—Turkest.
kentuckensis, Rafin. Ann. Gén. Sc. Phys. v. (1820) 213 = setigera.
Kingstonia, Hort. ex Dum.-Cours. Bot. Cult. ed II. v. 476 = centifolia.
Kluckii, Auct. = seq.
Klukii, Bess. Cat. Hort. Crem. (1816) 118 = rubiginosa.
Kmetiana, *Borb. Magyar Akad. Temész. Koxlem.* xvi. (1880) 437, 454.—Hungar.
Kosciuskiana, *Bess. ex Link, Enum. Hort. Berol.* ii. 60, nomen = montana.
Kosingiana, G. Don, Gen. Syst. ii. 565 = seq.
Kosinsciana, Bess. Enum. Pl. Volh. 64 = canina.
Kotschyana, *Boiss. Diagn.* Ser. I. x. 3.—Persia.
Krockeri, *Tratt. Ros.* ii. 231.—Europ.
labradora, *Hort. Hamb. ex Steud. Nom.* ed. II. ii. 469.—Labrador.
laccea, *Roess. Rosen*, ii. 56.—Hab.?
lacerans, *Boiss. & Buhse, in Nouv. Mém Soc. Nat. Mosc.* xii. (1860) 85.—Persia.
laciniata, Lej. ex Reichb. Fl. Germ. Excurs. 620 = canina?
lactea, Loisel. in Desv. Journ. Bot. ii. (1809) 337 = stylosa.
lacteola, Waitz, ex Steud. Nom. ed. I. 700 = centifolia.
lactiflora, Déségl. in Fourr. Cat. Cours du Rhone (1869), 76 = rubiginosa.
ladanifera, Timb. in Bull. Soc. Hist. Nat. Toul. iv. (1871) 173 = agrestis.
laevigata, *Michx. Fl. Bor. Am.* i. 295.—Am. bor.
laevigata, *Tausch, ex Opiz, Boehm. Gew.* (1823) 61.—Europ.
laevipes, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 298 = sepium.
laevis, Boullay, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 244 = gallica.
lagenaria, Vill. Hist. Pl. Dauph. iii. 553 = alpina.
Laggeri, *Puget, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 118.—Europ.
lanceolata, Opiz, in Flora, v. (1822) 268 = canina.
latebrosa, Déségl. in Journ. Bot. xii. (1874) 170 = canina.
latispina, Boiss. Diagn. Ser. II. ii. 49 = cabulica.
Lattuga, Savi, Fl. Ital. i. 99 = centifolia.
Laurentiae, Andr. Ros. t. 75 = seq.
Lawranceana, *Sweet, Hort. Suburb. Lond.* (1818) 119.—China.
laxa, Lindl. Ros. Monog. t. 3 = lucida.
laxa, *Retz. in Hoffm. Phytog. Blaetter*, i. (1803) 39.—Sibir.
Ledebourii, Spreng. Syst. ii. 551 = tomentosa.
Leeana, *Sweet, Guide*, 195; *Hort. Brit.* ed. III. 214.—Hab.?
Lehmanniana, *Bunge, in Mém. Sav. Etr. Pétersb.* xii. (1851) 287.—Turkestan.
Lemaitrei, Ripart, in Verlot, Cat. Pl. Dauph. 394 = canina.

ROSA:—
Lemanii, Bor. Fl. Centr. Fr. ed. III. ii. 230 = micrantha.
lepida, *Waitz, ex Tratt. Ros.* i. 22.—Hab.?
Lereschii, *Rapin, in Bull. Soc. Bot. Belg.* xiv. (1875) 237.—Europ.
Leschenaultiana, *Red. & Thory, ex Wight & Arn. Prod.* 301.—Ind. or.
leucantha, Loisel. in Desv. Journ. Bot. ii. (1809) 338; Bieb. Fl. Taur. Cauc. iii. 352 = canina.
leucochroa, Desv. Journ. Bot. ii. (1809) 316 = stylosa.
Leveillei, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxv.—Gallia.
Lheritieranea, Red. Ros. iii. 21 = alpina.
libanotica, Boiss. Diagn. Ser. I. x. 4 = glutinosa.
Libertiana, Tratt. Ros. ii. 80 = micrantha.
Likana, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 31 = gallica.
Lindleyana, Tratt. Ros. ii. 190 = involucrata.
Lindleyi, Spreng. Syst. ii. 647 = lucida.
Lindleyi, Wall. ex Hook. f. Fl. Brit. Ind. ii. 367 = moschata.
Linkii, Dehnh. Revista Napolit. i. III. 163; ex Walp. Rep. ii. 12 = multiflora.
lionensis, *Roess. Rosen*, i. 90.—Hab.?
litigiosa, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 239, 267 = canina.
livescens, Bess. Cat. Hort. Crem. Suppl. iv. 19; et Enum. Pl. Volh. 29 = canina?
livida, Host, Fl. Austr. ii. 25 = ferruginea.
Lloydi, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 391 = canina.
longepedunculata, La Soie, in Christ, Rosen Schw. (1873) 180 = montana.
longicuspis, Bertol. Misc. Bot. xxi. 15 = macrophylla.
longifolia, Willd. Sp. Pl. ii. 1079 = indica.
Longii ×, Sweet, ex Steud. Nom. ed. II. ii. 470.
Loureiriana, *G. Don, Gen. Syst.* ii. 585.—China; Cochinch.
Lucandiana, *Déségl. & Gillot, in Bull. Soc. Bot. Belg.* xix. (1880) 36.—Gallia.
Luciae, *Franch. & Rochebr. l. c.* x. (1871) 324.—Japon.
lucida, Auct. ex Meehan, Nat. Flow. ii. 33. t. 9 = humilis.
lucida, *Andr. Ros.* t. 78.—China.
lucida, *Ehrh. Beitr.* iv. 22.—Am. bor.
lucida, Koch, Syn. Fl. Germ. ed. II. 444 = baltica.
lucida, Lawr. Ros. t. 84 = bracteata.
ludebunda, Gren. & Paill. in Gren. Rev. 69 = canina.
lugdunensis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 141 = agrestis.
lugdunensis, Wrede, ex Breit. Hort. Breit. 425 = lionensis.
lurida, Andr. Ros. t. 92 = ferruginea.
lusitanica, *Roess. Rosen*, iii. 219.—Lusit.
Lusseri, Lagg. & Puget, in Bull. Soc. Murith. ii. (1873) 42 = ferox.
lutea, Brot. Fl. Lusit. ii. 337 = hemisphaerica, *Herrm.*
lutea, Mill. Gard. Dict. ed. VIII. n. 11; Lam. Fl. Fr. iii. 132 = R. Eglanteria.
lutescens, Pursh, Fl. Am. Sept. ii. 735 = blanda.
lutetiana, Léman, in Bull. Soc. Philom. (1818) 93 = canina.
Lyellii, Lindl. Ros. Monog. 12. t. 1 = involucrata.
lyonensis, Steud. Nom. ed. I. 700 = lionensis.
Lyonii, Pursh, Fl. Am. Sept. i. 345 = humilis.
Macartnea, Dum.-Cours. Bot. Cult. ed. I. i. 351 = laevigata.
Macartnea, Dum.-Cours. l. c. ed. II. v. 483 = bracteata.
macrantha, Desf. Fl. Sarthe, 77 = canina.
macroacantha, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 314 = canina.
macrocarpa, *Boiss. Fl. Orient.* ii. 684.—As. Min.
macrocarpa, Mérat, Fl. Par. 190 = canina.
macrocarpa, Nutt. ex Rafin. Med. Bot. ii. 258 = nutkana.
macrodonta, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxv.—Gallia.
macrophylla, *Lindl. Ros. Monog.* 35. t. 6.—Reg. Himal.; China.
macropoda, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 265 = spinosissima.
macrostylis, Stokes, Bot. Mat. Med. iii. 132 = arvensis.
Maerklini, *Dierb. in Flora*, xi. (1828) I. *Erg.* 108.—Austria.

ROSA :—
maialis, *Desf. Fl. Atlant.* i. 400.—Afr. bor.
majalis, Borrer, in Hook. Fl. Bor. Am. i. 201 = acicularis?
majalis, Herrm. Diss. 8 = cinnamomea.
majalis, Wallr. Ros. 164 = ferruginea.
Makeka, Dupont, ex Reichb. Fl. Germ. Excurs. 622 = gallica.
maldensis, *Roess. Ros.* ii. 3.—Hab.?
malmundariensis, Lej. Fl. Spa, i. 231 = canina.
Malyi, *Kern. in Oestr. Bot. Zeitschr.* xix. (1869) 325. —Europ.
Mandonii, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 111 = canina.
Manettii, *Hort.—Cf. Gard. Chron.* (1849) 724, 742.—Hab.?
maracandica, *Bunge, in Mém. Sav. Etr. Pétersb.* xii. (1851) n. VII. 286.—Turkestan.
Marcyana, Boullu, in Bull. Soc. Bot. Fr. xxiii. (1876) p. lxvi = tomentosa.
Mareyana, Boullu, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 556, sphalm. = praec.
marginata, Wallr. Ann. Bot. 68 = canina.
marisensis, *Simck. & H. Braun, in Termész. Füzetek*, ix. (1885) 40.—Europ.
maritima, *Wrede, ex Breit. Hort. Breit.* 425.—Europ.
marmorea, *Roess. Ros.* i. 186.—Hab.?
marsica, *Godet, in Nouv. Giorn. Bot. Ital.* ix. (1877) 162.—Italia.
Martini, Gren. Fl. Jurass. (1864) 242 = canina.
massiliensis, *Breit. ex Steud. Nom.* ed. I. 700; ed. II. ii. 470.—Hab.?
Mathonneti, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 263 = spinosissima.
matraënsis ×, Borb.—Cf. Oestr. Bot. Zeitschr. xxxiv. (1884) 261.—Europ.
Maukschii, Kit. in Linnaea, xxxii. (1863) 588 = scepusiensis.
Maukschii, *Schult. Oestr. Fl.* ed. II. ii. 69.—Hungar.
maxima, Desf. Tabl. ed. I. 174 = centifolia.
Maximiliani, Nees, in Neuwied, Reise Nord-Am. ii. 434 = Woodsii.
Maximowicziana, Regel, in Act. Hort. Petrop. v. (1877) 378 = multiflora?
media, Hort. ex DC. Prod. ii. 620 = centifolia.
media, *B. Martin.—Cf. Bull. Soc. Bot. Fr.* xxx. (1883) *Bibl.* 186.—Gallia.
medioxima, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 110 = canina.
megacarpa, Nutt. ex Torr. & Gray, Fl. N. Am. i. 460 = nutkana?
megalocarpa, *Déségl. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 117.—Europ.
melanocarpa, Link, Enum. Hort. Berol. ii. 57 = spinosissima.
meldensis, Dum.-Cours. Bot. Cult. ed. II. v. 474 = centifolia.
menstrua, Andr. Roses, t. 50 = damascena.
menstrualis, Dum.-Cours. Bot. Cult. ed. II. v. 486 = praec.
mentita, Déségl. in Billotia, i. (1864) 43 = agrestis.
meridionalis, Burm. & Gremli, Roses Alpes-Marit. 75 = canina.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 354.—Mexic.
mexicana, *Willd. ex Spreng. Syst.* ii. 555.—Mexic.
Meyeniana, *Steud. Nom.* ed. II. ii. 470.—Chili.
micans, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 555 = tomentosa.
micrantha, Bor. Fl. Centr. Fr. ed. II. ii. 182, in obs. = rubiginosa.
micrantha, DC. Fl. Fr. vi. 539 = canina.
micrantha, *Sm. Engl. Bot.* t. 2490.—Europ.; As. Min.
microcarpa, Bess. Cat. Hort. Crem. (1811) Suppl. iv. 20; Enum. Pl. Volh. 18, 60, 62, 81 = spinosissima.
microcarpa, *Lindl. Ros. Monog.* 130. t. 18.—China.
microcarpa, *Retz. in Hoffmgg. Phytog. Blaetter*, i. (1803) 40.—Sibir.
microphylla, Desf. Fl. Atlant. i. 401 = sempervirens.
microphylla, *Roxb. Hort. Beng.* 38; *Fl. Ind.* ii. 515.—China.
microphylla, Willd. ex Spreng. Syst. ii. 547 = Willdenowii.
microtypos ×, Borb. & Vukot.—Cf. Oestr. Bot. Zeitschr. xxxiv. (1884) 260.—Europ.
millesia, Linn. Amoen. Acad. iv. 484 (Quid?).—Gallia.

ROSA :—
miniata, *Wrede, ex Steud. Nom.* ed. I. 700; ed. II. ii. 470.—Europ. austr.
minima, *Hort. Cantab.—Cf. Gard. Chron.* (1881) i. 242.—Hab.?
minima, *Roess. Rosen*, 139.—Hab.?
minuta, Bor. ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 167 = pomifera.
minuta, *Hort. ex Steud. Nom.* ed. II. ii. 470.—Hab.?
minutifolia, *Parry, in Bull. Torrey Bot. Club*, ix. (1882) 97.—Calif.
mirabilis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 101 = gallica.
mirogojana, Vukot. & H. Braun, ex Kern. Sched. Fl. Exsicc. iii. 47 = gallica.
Missonia, *Hort. ex Steud. Nom.* ed. I. 700; ed. II. ii. 470.—Hab.?
missourensis, *Hort. ex Steud. l. c.* II. ii. 470.—Am. bor.?
mitis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 84 = laceraas.
mitissima, C. C. Gmel. Fl. Bad. iv. 358 = spinosissima.
mixta, Chab. in Cariot, Etud. Fleurs, ii. (1865) 677 = gallica.
mixta, Tratt. Ros. i. 136 = alpina.
modesta, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 260 = stylosa.
mollis, Ledeb. ex Spreng. Syst. ii. 551 = tomentosa.
mollis, *J. C. Presl, Fl. Cech.* 102.—Europ.
mollis, Sm. Engl. Bot. t. 2459 = mollissima.
mollissima, *Willd. Prod. Fl. Berl.* 437.—Europ.; As. Min.
monophylla, Hort. ex Steud. Nom. ed. II. ii. 470 = berberifolia.
monspeliaca, Gouan, Fl. Monsp. 255 = alpina.
monstrosa, Hort. ex Breit. Hort. Breit. 425 = centifolia.
montana, *Chaix, in Vill. Hist. Pl. Dauph.* i. 346; iii. 547.—Europ.
montana, DC. Fl. Fr. Suppl. 532 = tomentosa.
montana, Murith, Bot. Val. 91 = alpina.
montana, Stev. ex Spreng. Syst. ii. 552 = canina.
Montezumae, *Humb. & Bonpl. ex Red. & Thory, Ros.* i. 55.—Mexic.
monticola, Rapin, Cant. Vaud. ed. II. 194 = canina.
montivaga, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 107 = canina.
Moorcroftii, Wall. Cat. n. 693 = R. Eglanteria?
moschata, Benth. Fl. Hongk. 106 = Luciae.
moschata, *Herrm. Diss.* 15.—Oriens.
mosquensis, Spreng. Syst. iv. Cur. Post. 200 = cinnamomea.
mucronulata, Déségl. in Godet, Fl. Jura, Suppl. (1869) 71 = canina.
multiflora, Andr. Ros. t. 83 = japonica.
multiflora, Benth. Pl. Hongk. 107 = microcarpa.
multiflora, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. ii. 365 = involucrata.
multiflora, *Dum.-Cours. Bot. Cult.* ed. I. iii. 346.—Hab.?
multiflora, Reyn. ex Reichb. Fl. Germ. Excurs. 620 = ferruginea.
multiflora, *Thunb. Fl. Jap.* 214.—Japon; China.
multiflora, Wrede, ex Roess. Ros. 68 = centifolia.
multivaga, Déségl. in Bull. Soc. Bot. Belg. xiv. (1875) 341 = rubiginosa.
Mureti ×, Rapin, l. c. 236.—Europ.
muricata, *Waitz, ex Link, Enum. Hort. Berol.* ii. 56.—Hab.?
Murithii, Puget, in Bull. Soc. Murith. iii. (1875) 55 = pomifera.
muscosa, Mill. Gard. Dict. ed. VIII. n. 22 = gallica.
mutabilis, Bradbury, ex James, in Long's Exped. i. 69 = setigera.
mutabilis, Dum.-Cours. Bot. Cult. ed. II. v. 475 = gallica.
mutica, O. F. Muell. in Fl. Dan. t. 688 = blanda, cinnamomea.
myriacantha, DC. Fl. Fr. iv. 439 = spinosissima.
myriantha, *Carr. in Rev. Hortic.* (1865) 448.—Calif.
myropolarum, Steud. Nom. ed. I. 700 = damascena.
myrtifolia, Hall. f. ex Steud. l. c. = agrestis.
nana, Thory, Ros. 80 = spinosissima.
nankinensis, Lour. Fl. Cochinch. 324 = indica.
nanula, *Hoffmgg. Verz. Pfl.* 108.—China.
napaulensis, Andr. Ros. t. 82 = moschata.
nebrodensis, Guss. Fl. Sic. Syn. i. 563 = canina.

ROSA :—
neglecta, *Léman, in Journ. Phys.* lxxxvii. (1818) 367, nomen.—Gallia.
neglecta, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 524 = agrestis.
nemoralis, *Léman, in Bull. Soc. Philom.* (1818) 94.—Gallia.
nemorivaga, Déségl. in Billotia, i. (1864) 40 = flexuosa.
nemorosa, Libert, in Lej. Fl. Spa, ii. 311 = micrantha.
nemorum, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 245 = gallica.
nepalensis, Lindl. ex Steud. Nom. ed. II. ii. 470 = moschata.
nevadensis, Crép. in Willk. & Lange, Prod. Fl. Hisp. iii. 210 = stylosa.
nitens, Desv. in Mérat, Fl. Par. (1812) 192 = canina.
nitida, *Willd. Enum. Hort. Berol.* 544.—Am. bor.
nitidula, Bess. Enum. Pl. Volh. 20, 61, 67 = canina.
nivalis, Donn, Cat. Hort. Cantab. ed. VIII. 170 = involuta.
nivea, DC. Cat. Hort. Monsp. 137 = laevigata.
nivea, Dupont. ex Lindl. in Bot. Reg. sub t. 861 = moschata.
nivea, Hort. ex Steud. Nom. ed. I. 700 = centifolia.
nivea, Rafin. in Ann. Gén. Sc. Phys. v. (1870) 218 = blanda.
Noisettana, Bosc, Encyc. Agric. vii. 705 = seq.
Noisettiana, Red. Ros. ii. 77 = indica.
normandica, Hort. ex Dum.-Cours. Bot. Cult. ed. II. v. 476 = centifolia.
nova, Dum.-Cours. l. c. ed. I. iii. 351 = *oblongifolia*.
nuda, Woods, in Trans. Linn. Soc. xii. (1817) 205 = canina.
numidica, *Gren. ex Déségl. in Journ. Bot.* xii. (1874) 171.—Afr. bor.
nutkana, *Presl, Epim. Bot.* 203.—Am. bor. occ.
oblonga, Déségl. & Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 332 = canina.
oblongata, *Opiz, in Lotos*, iv. (1854) 69.—Europ.
oblongifolia, Dum.-Cours. Bot. Cult. ed. I. iii. 351 (Quid?).—Hab.?
obovata, Lagg. & Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 23 = canina.
obovata, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 217 = humilis.
obscura, Puget, in Fourr. Cat. Pl. Rhone (1869), 75 = canina.
obtusa, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 352.—Gallia.
obtusifolia, Christ & Gremli, in Gremli, Excurs. Schw. ed. IV. 174 = scaphusiensis.
obtusifolia, Desv. Journ. Bot. ii. (1809) 317 = canina.
occulta, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 273 = canina.
ochroleuca, *Sw. ex Thunb. in Mus. Nat. Acad. Ups. App.* xxiii. (1816) 26, nomen.—Sibir.
ochroleuca, Waitz, ex Wikstr. in Vet. Acad. Handl. Stockh. (1820) 271, 272 = lutescens.
odorata, Sweet, Hort. Brit. ed. I. 138 = indica.
odoratissima, Hort. ex Steud. Nom. ed. II. ii. 470 = indica.
odoratissima. Scop. Fl. Carn. ed. II. i. 354 = alpina.
Oederiana, Tratt. Ros. ii. 133 = spinosissima.
oenensis, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 328 = canina.
oleicarpa, Thory, Ros. 114 = sepium.
olympica, Donn, Hort. Cantab. ed. VI. 144 = gallica.
omissa, Déségl. in Billotia, i. (1864) 47 = mollissima.
opaca, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 332 = canina.
opacifolia, Chab. in Cariot, Etud. Fleurs, ii. (1865) 677 = gallica.
operta, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 287 = micrantha.
oplisthes, *Boiss. Fl. Orient.* ii. 674.—Reg. Caucas.
Opsostemma, Ehrh. Beitr. ii. 72 = moschata.
Orbessanea, Red. Ros. ii. 21 = turbinata.
oreinosa, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 284 = alpina.
orientalis, *Dupont, ex DC. Prod.* ii. 607.—As. Min.; Persia.
Orphanidis, Boiss. & Reut. Diagn. Ser. II. ii. 50 = Heckeliana.
orthacantha, Kern. in Christ, Rosen Schw. (1873) 168 = canina.
orthosepala, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 768 = cinnamomea, ferruginea.

ROSA:—
ostensa, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 426 = canina.
ovata, Lej. Fl. Spa, ii. 312 = arvensis.
oxyacantha, Bieb. Fl. Taur. Cauc. iii. 338 = spinosissima.
oxyodon, *Boiss. Fl. Orient.* ii. 674.—Reg. Caucas.
oxyphylla, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 315, in obs. = canina.
Ozanonii, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 88 = spinosissima.
pallens, *Retz. ex Spreng. Syst. App.* 200.—Suecia.
pallida, Tratt. ex Link, Handb. ii. 85 = ochroleuca.
palustris, Buch.-Ham. ex Lindl. Ros. Monog. 8 = involucrata.
palustris, Marsh. Arbust. 135; Roess. Rosen, i. 240 = R. Carolina.
pampinula, Roess. ex Steud. Nom. ed. I. 700 = R. Eglanteria.
paniculata, Hort. Genev. ex DC. Prod. ii. 600 = indica.
pannonica, Wiesb. in Oestr. Bot. Zeitschr. xxix. (1879) 143 = gallica.
pannosa, Link, Handb. ii. 83 = indica?
Pantocekii, *Holuby, ex Pant. in Verh. Ver. Natur- u. Heilk. Presb.* ii. (1871–72) 117.—Europ.
papaverina, Borck. Forstbot. iii. 1814 = seq.?
papaverina, Moench, ex Roess. Rosen, i. 182 = gallica.
paradoxa, Burnat & Gremli, Roses Alpes-Marit. 126 = arvensis.
parisiensis, *Wrede, ex Breit. Hort. Breit.* 426.—Hab.?
parviflora, Ehrh. Beitr. iv. 21 = humilis.
parvifolia, *Andr. Ros.* t. 56.—Hab.?
parvifolia, Ehrh. Beitr. vi. 97 = centifolia.
parvifolia, Lindl. Ros. Monog. 70 = humilis.
parvifolia, Pall. Fl. Ross. i. II. 62 = spinosissima.
parvula, *Gren. ex Crép. in Bull. Soc. Bot. Belg.* xv. (1869) 287.—Europ.
parvula, Guss. ex Nym. Consp. 233 = rubiginosa.
parvula, Sauzé & Maill. Cat. Pl. Deux-Sèvres (1864), 27 = stylosa.
patenti-ramea, Debeaux, ex Crép. in Bull. Soc. Bot. Belg. xxi. (1882) 56 = ferruginea.
pauciflora, *Lindl. ex Sweet, Hort. Brit.* ed. II. 179.—Sibir.
pellita, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 555 = tomentosa.
pellucina, Arv. ex Déségl. l. c. 259 = spinosissima.
pendula, Salisb. Prod. 359 = spinosissima.
pendulina, Ait. Hort. Kew. ed. I. ii. 208 = alpina.
pendulina, Linn. Sp. Pl. 492 = spinosissima.
pennina, La Soie, ex Cottet, in Bull. Soc. Murith. iii. (1874) 39 = glauca.
pennsylvanica, Andr. Ros. t. 101 = gallica *vel* R. Carolina.
pensylvanica, Michx. Fl. Bor. Am. i. 296 = R. Carolina.
pensylvanica, Wangenh. Beytr. Forstwiss. 113 = humilis.
pentaphylla, *Schrank, in Syll. Ratisb.* i. (1824) 215.—China.
perfecta, *Wrede, ex Steud. Nom.* ed. I. 701.—Hab.?
permixta, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 147 = ferox.
permutata, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 575.—Europ.
Perrieri, Long, ex Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 425 = montana.
persica, J. F. Gmel. Syst. 855 (nomen prius) = berberifolia.
persicifolia, Hort. ex Hoffmgg. Verz. Pfl. 108 = indica.
personata, Gremli, Excurs. Schw. ed. IV. 178 = pomifera.
perusiana, Timb. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 582 = pomifera.
pervirens, *Gren. ex Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 257.—Hab.?
petiolaris, Kit. in Linnaea, xxxii. (1863) 592 = carpatica.
petrogenes, Ozan. ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 261 = spinosissima.
phoenicia, *Boiss. Diagn.* Sér. I. x. 4.—Cilicia; Syria.
phrygia, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 249 = rubiginosa.
picta, *Wrede, ex Steud. Nom.* ed. I. 701.—Hab.?
pilosa, *Hort. ex Steud. l. c.* ed. II. ii. 471.—Oriens.
pilosa, *Opiz, in Flora*, v. (1822) 268.—Europ.
pilosiuscula, *Opiz, l. c.*—Europ.

ROSA:—
pimpinellifolia, Herb. Hook. f. & Thoms. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 262 = Webbiana.
pimpinellifolia, Linn. Syst. ed. X. 1062 = spinosissima.
pimpinelloides, Pall. Reise Suedl. Statth. Russ. Reich. i. 349 (Quid?).—Rossia.
pinnatifida, *Andr. Ros.* t. 41.—Europ.
pisocarpa, *A. Gray, in Proc. Am. Acad.* viii. (1872) 382.—Am. bor. occ.
Pissardi, *Carr. in Rev. Hortic.* (1880) 314.—Reg. Casp.
Pitcairnea, Sabine, ex Link, Handb. ii. 85 = spinosissima.
platyacantha, Schrenk, in Bull. Sc. Acad. Pétersb. x. (1842) 254 = xanthina.
platyphylla, Rau, Enum. Ros. 82 = canina.
platyphylloides, Déségl. & Ripart, in Fourr. Cat. Pl. Cours Rhone (1869), 75 = canina.
plicata, *Bechst. Forstbot.* ed. V (1843), 583.—Europ.
plicata, Roess. Rosen, ii. 58 = praec.?
podolica, Tratt. ex Link, Handb. ii. 91 = tomentosa.
Pokornyana ×, Kmet.—Cf. Oestr. Bot. Zeitschr. xxxiii. (1833) 225.—Europ.
Polliniana, Spreng. Pugill. ii. 66 = arvina.
polyadena, *Burnat & Gremli, Suppl. Mon. Roses Alpes-Marit.* 26.—Europ.
polyantha, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 128 = multiflora.
polyanthos, Roess. Rosen, t. 35 = centifolia.
polymorpha, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 769 = canina.
polyphylla, Willd. Enum. Hort. Berol. Suppl. 37 = acicularis, spinosissima.
pomifera, *Herrm. Diss.* 16.—Graecia; Reg. Caucas.
pomifera, Lamotte, Cat. Pl. Centr. Fr. 38; Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 217 = mollissima.
Pommaretii, *Puget, ex Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 286.—Gallia.
pomponia, DC. Fl. Fr. iv. 443 = centifolia.
porrigens, Gremli, Excurs. Schw. ed. IV. 163 = gallica.
Portenschlagiana, *Tratt. Ros.* ii. 203.—Hab.?
Portlandia, Hort. ex Andr. Ros. t. 19 = ruficaulis.
portlandica, Roess. Rosen, i. 240 = praec.?
portugalensis, *Roess. l. c.* ii. 45.—Hab.?
poteriifolia, Bess. Enum. Pl. Volh. 62 = spinosissima.
poteriifolia, Schott & Kotschy, ex Boiss. Fl. Orient. ii. 679 = glutinosa.
Poterium, *Léman, in Journ. Phys.* lxxxvii. (1818) 366.—Gallia.
Pougeti, *Bor. ex Déségl. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 136.—Gallia.
Pouzini, Tratt. Ros. ii. 112 = canina.
praecox, Boullu, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 571 = tomentosa.
praecox, *Hort. ex Steud. Nom.* ed. II. ii. 471.—Hab.?
praenestina, Mill. ex Thory, Ros. 92 = gallica.
praeterita, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 499 = canina.
pratensis, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 215 = humilis.
pratensis, Sw. ex Steud. Nom. ed. II. ii. 471 = canina.
Presliana, Opiz, in Seits, Rosen (1819), 143 = mollis.
primoriensis, *Vukot. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 44.—Europ.
procera, *Hort. ex Steud. Nom.* ed. II. ii. 471.—Sibir.
procera, Salisb. Prod. 359 = alba.
procumbens, *Roxb. Hort. Beng.* 38, nomen.—Ind. or.
procumbens, Ser. Mél. Bot. i. 5 = arvensis.
prolifera, Hort. ex Steud. Nom. ed. I. 701; ed. II. ii. 471 = multiflora.
properata, Boullu, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 571 = tomentosa.
propinqua, Déségl. l. c. xiv. (1875) 343 = rubiginosa.
prostrata, DC. Cat. Hort. Monsp. 138 = sempervirens.
protea, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 503 = trachyphylla.
provincialis, Bieb. Fl. Taur. Cauc. i. 396 = ferox.
provincialis, Herrm. Diss. 9 = gallica.
proxima, Cottet, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 248 = pomifera.
pruinosa, Baker, in Naturalist, i. (1864) 96 = canina.
pruinosa, *Lindl. in Loud. Encyc. Pl.* 448.—Sibir.
Pseudo-flexuosa, Ozan. ex Déségl. in Billotia, i. (1864) 42 = flexuosa.

ROSA:—
Pseudo-indica, Lindl. Ros. Monog. 132 = indica.
Pseudo-lucida, Bess. ex DC. Prod. ii. 604 = baltica.
pseudopsis, Gremli, Excurs. Schw. ed. IV. 175 = canina.
Pseudo-rubiginosa, Lej. Fl. Spa, i. 229 = mollissima.
Pseudo-sepium, Callay, ex Cottet, in Bull. Soc. Murith. iii. (1874) 42 = agrestis.
Pseudo-vestita, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxvii.—Gallia.
psilophilla, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 501 = trachyphylla.
psilophylla, Rau, Enum. Ros. 101 = canina.
puberula, Desv. ex Ser. Mél. Bot. i. 41 = canina.
pubescens, *Léman, in Bull. Soc. Philom.* (1818) 94.—Gallia.
pubescens, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 514 = moschata.
pubescens, *Schleich. Cat. Pl. Helv.* ed. III. 24.—Europ.
Pugeti, Bor. ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 136 = flexuosa.
pulchella, Bor. Fl. Centr. Fr. ed. II. n. 662 = gallica.
pulchella, Salisb. Prod. 358 = spinosissima.
pulchella, Schott & Kotschy, ex Boiss. Fl. Orient. ii. 680 = orientalis.
pulchella, *Willd. Enum. Hort. Berol.* 545.—Hab.?
pulchella, Woods, in Trans. Linn. Soc. xii. (1817) 196 = tomentosa.
pulverulenta, Baker, in Journ. Linn. Soc. xi. (1869) 223 = agrestis.
pulverulenta, *Bieb. Fl. Taur. Cauc.* i. 399.—Reg. Caucas.
pulverulenta, Guss. ex Nym. Consp. 232 = glutinosa.
pumila, Scop. Fl. Carn. ed. II. i. 353 = gallica.
punicea, Mill. Gard. Dict. ed. VIII. n. 12 = R. Eglanteria.
purpurascens, *Fisch. ex Sweet, Hort. Brit.* ed. II. 180.—Sibir.
purpurascens, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 351.—Gallia.
purpurea ×, Hort. ex G. Don, in Loud. Hort. Brit. 211.
pusilla, Rafin. in Ann. Nat. 11; et in Ann. Gén. Sc. Phys. v. (1820) 218 = humilis.
pustulosa, Bertol. Fl. Ital. v. 195 = glutinosa.
Puymaurea, Gren. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 549 = rubiginosa.
pycnacantha, *Borb. in Magyar Akad. Természes. Kozlem.* xvi. (1880) 438, 448.—Hungar.
pycnocephala, Christ, in Gremli, Excurs. Schw. ed. IV. 169 = tomentosa.
pygmaea, Bieb. Fl. Taur. Cauc. i. 397 = gallica.
pygmaea, Krock. Fl. Siles. ii. 154 = Krockeri.
pyramidalis, *Wrede, ex Roess. Rosen*, i. 67.—Hab.?
pyrenaica, Gouan, Illustr. 31 = alpina.
pyriformis, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 387 = canina.
quassata, Gren. in Mém. Soc. Emul. Doubs, Ser. IV. ix. (1875) 439 = sepium.
racemosa, *Andr. Ros.* t. 49.—Gallia.
Rafinesqueiana, Tratt. Ros. ii. 234 = R. Carolina.
Rafinesquii, Ser. in DC. Prod. ii. 611 = blanda.
ramealis, Puget, ex Baker, in Journ. Linn. Soc. xi. (1869) 228 = canina.
ramosissima, Rau, Enum. Ros. 74 = canina.
ramulosa, Godr. Fl. Lorr. ed. II. i. 231 = canina.
ranunculiflora, *Andr. Ros.* t. 33.—Hab.?
ranunculiformis, Steud. Nom. ed. I. 701; Link, Handb. ii. 92, 93 = praec.
Rapa, Bosc, Encyc. Agric. vii. 704, 706 = lucida.
Rapini, Boiss. & Bal. Diagn. Ser. II. vi. 72 = hemisphaerica, *Herrm.*
Ratomsciana, Bess. Enum. Pl. Volh. 65 = canina.
Raui, Tratt. Ros. ii. 35 = canina.
Ravaudi, Boullu, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 295 = montana.
Ravellae, Christ, in Gremli, Excurs. Schw. ed. IV. 162 = involuta.
reclinata ×, Red. & Thory, Ros. iii. 79.
recognita, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 296 = canina.
recondita, Puget, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 107 = pomifera.
recurva, Roxb. Hort. Beng. 38; Fl. Ind. ii. 514 = moschata?
Redoutea, Thory, Ros. 43 = *Reduteana*.
Redowskiana, *Bess. ex Sweet, Hort. Brit.* ed. II. 180.—Hab.?
Reduteana, Thory, ex Red. Ros. i. 101 = nitida.

ROSA :—

Regeliana, Linden & André, Illustr. Hortic. xviii. (1871) t. 47 = rugosa.
Regelii, *Reut. Ind. Sem. Hort. Genev.* (1867) 4.—China.
Regina, Roess. Rosen, i. 47 = centifolia ?
Reginae, Hort. ex Steud. Nom. ed. I. 702 = praec. ?
Reichenbachiana, *Waitz, in Flora*, vii. (1824) 417.—Hab. ?
remensis, DC. Fl. Fr. iv. 443 = gallica.
remensis, Dum.-Cours. Bot. Cult. ed. II. v. 474 = centifolia.
repanda, Hort ex Steud. Nom. ed. II. ii. 471 (Quid ?).—Scotia.
repens, Muench, Hausv. v. (1770) 281 = gallica.
repens, Scop. Fl. Carn. ed. II. i. 355 = arvensis.
reptans, Crép. ex Bout. in Bull. Soc. Bot. Fr. xxvii. (1880) 298 = arvensis.
resinosa, Bor. Fl. Centr. Fr. ed. II. ii. 233 = mollissima.
resinosa, Lej. Rev. Fl. Spa, 96 = rubiginosa.
resinosa, Sternb. in Reichb. Fl. Germ. Excurs. 616 = pomifera.
resinosoides, Crép. in Cottet, 'in Bull. Soc. Murith. iii. (1874)' 44 = mollissima.
reticulata, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 332 = canina.
Reuteri, Godet, Fl. Jura, 208, in obs. = glauca.
reversa, J. & C. Presl, Fl. Cech. 103 = alpina.
reversa, Waldst. & Kit. Pl. Rar. Hung. iii. 293. t. 264 = spinosissima.
Reynieri, Hall. f. in Roem. Arch. i. II. (1797) 7 = montana.
rhaetica, Gremli, Excurs. Schw. ed. IV. 164 = rubiginosa.
rhemensis, *Roess. Rosen*, ii. 64.—Hab. ?
Rhodani, Chab. ex Boullu, in Bull. Soc. Bot. Fr. xxiii. (1876) p. lxiv = arvina.
rhoetica, Kern. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 304 = glauca.
rhombifolia, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxiii.—Gallia.
rhynchocarpa, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 331, in obs. = canina.
rigida, Willd. ex Déségl. l. c. xi. (1872) 84 = ferox.
rigidula, Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 23 = canina.
riparia, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 216 = humilis.
Ripartii, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 87 = spinosissima.
Robertsoni, Baker, ex Syme, Engl. Bot. ed. III. iii. 207 = involuta.
Rosenbergiana, Red. & Thory, Ros. ii. 113 = turbinata.
rossica, Pall. ex Spreng. Neue Entdeck. iii. 254 = spinosissima.
rothomagensis, Rouy, in Bull. Soc. Bot. Fr. xxii. (1875) 297 = agrestis.
rotundifolia, Reichb. Fl. Germ. Excurs. 617 = rubiginosa.
Rousselii, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 353 = canina.
Roxbourgii, Tratt. Ros. ii. 233 = microphylla.
Roxburghii, Sweet, Hort. Brit. ed. I. 138 = multiflora.
rubella, Sm. Engl. Bot. t. 2521 = spinosissima.
rubella, Winch, ex J. Hogg, in Journ. Linn. Soc. iv. (1860) 198 = spinosissima.
rubelliflora, *Ripart, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 109 = canina.
rubeoides, Andr. Ros. t. 84 = macrophylla ?
rubescens, Ripart, ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 110 = canina.
rubicunda, Hall, f. in Roem. Arch. i. II. (1797) 6 = ferruginea.
rubifolia, R. Br. in Ait. Hort. Kew. ed. II. iii. 260 = setigera.
rubifolia, Vukot. in Rad Jugos. Akad. Zagreb. lxix. (1884) 21 = subalbida.
rubiginosa, Coss. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 344 = canina.
rubiginosa, *Linn. Mant.* ii. 564.—Europ.; Reg. Caucas.
rubiginosa, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 446 = micrantha.
rubra, Lam. Fl. Fr. iii. 130 = gallica.
rubricans, Roess. ex Steud. Nom. ed. I. 702 = centifolia ?
rubrifolia, Rupr. ex Boiss. Fl. Orient. ii. 684 = haematodes.
rubrifolia, Vill. ist. Pl. Dauph. iii. 549 = ferruginea.

ROSA :—

rubripetala, *Roem. Hort. Turic. ex Steud. Nom.* ed. II. ii. 472.—Hab. ?
rubrispina, Bosc, ex Poir. Encyc. Suppl. iv. 715 = nitida.
ruficaulis, *Ehrh. Beitr.* vii. 138.—Am. bor.
Ruga ×, Lindl. in Bot. Reg. t. 1389.
rugibinosa, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 251, sphalm. = rubiginosa.
rugosa, Dematra, Monog. Ros. (1818) 4 = Friesii.
rugosa, *Thunb. Fl. Jap.* 213.—Japon.
rupestris, Crantz, Stirp. Austr. ed. I. fasc. ii. 32; ed. II. 85 = alpina.
rupincola, *Fisch. ex Sweet, Hort. Brit.* ed. II. 180.—Sibir. altaic.
Ruprechtii, *Boiss. Fl. Orient.* ii. 682.—Reg. Caucas.
ruralis, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 79 = gallica.
ruscinonensis, Gren. & Déségl. ex Déségl. in Billotia, i. (1864) 33 = moschata.
Russelliana, *Sincl. ex Forbes, Hort. Woburn.* 117.—Hab. ?
rustica, Léman, in Bull. Soc. Philom. (1818) 93 = stylosa.
rusticana, Déségl. in Billotia, i. (1864) 34 = arvensis.
sabauda, Rapin, in Bull. Soc. Hallér. (1852–53) 178 = involuta.
Sabina, Dupont. ex Red. Ros. i. 94 = rubiginosa.
Sabini, Woods, in Trans. Linn. Soc. xii. (1817) 188 = involuta.
salaevensis, Christ, ex Nym. Consp. 237 = alpina.
salaevensis, Rapin, in Bull. Soc. Hallér. (1852–53) 178 = montana.
salaevensis, Verlot, Cat. Pl. Dauph. 115 = montana.
salicifolia, Red. Ros. iii. 121 (Index) = R. Carolina.
Salisii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 63.—Europ.
salvanensis, La Soie, ex Cottet, in Bull. Soc. Murith. iii. (1874) 43 = ferox.
Sambrancheriana, La Soie, ex Cottet, l. c. 39 = glauca.
sancta, *Andr. Ros.* t. 98.—Europ.
sancta, *A. Rich. Tent. Fl. Abyss.* i. 262.—Abyssin.
sanguinea, Roess. ex Red. Ros. i. 76 = gallica.
sanguisorbaefolia, Nois. ex Red. l. c. ii. 8 = turbinata.
sanguisorbifolia, Donn, Cat. Hort. Cantab. ed. VIII. 169 = spinosissima.
sanguisorbifolia, La Soie, in Christ, Rosen Schw. (1873) 181 = montana.
sarmentacea, Woods, in Trans. Linn. Soc. xii. (1817) 213 = canina.
saxatilis, Hort. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 276 = Fischeriana.
saxatilis, Stev. ex Bieb. Fl. Taur. Cauc. 349 = canina.
saxigena, H. Braun.—Cf. Oestr. Bot. Zeitschr. xxxii. (1882) 410 = canina.
Sayi, *Schwein. in Long, Exped. Winnep.* ii. 388.—Am. bor.
scabrata, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 241 = canina.
scabrata, *Henning, in Mém. Soc. Nat. Mosc.* vi. (1823) 88.—Reg. Caucas.
scabriuscula, Winch, ex Sm. in Engl. Bot. t. 1896 = tomentosa.
scandens, Mill. Gard. Dict. ed. VIII. n. 8 = sempervirens.
scandens, Moench, Verz. Ausl. Baeume, 118 = arvensis.
scaphusiensis, *Christ, in Gremli, Excurs. Schw.* ed. IV. 175.—Helvet.
scepusiensis, *Kit. in Linnaea*, xxxii. (1863) 588.—Hungar.
schergiana, *Boiss. Fl. Orient.* ii. 686.—Syria.
Scheutzii, Christ, in Flora, lvii. (1874) 511 = mollissima.
Schimperiana, Hochst. l. c. xxiv. (1841) 1. Intell. 31 = abyssinica.
Schmidtii, *H. Braun, Beitr. Kenntn. Gatt. Rosa* (1885), 14 [72].—Europ.
Schottiana, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 358 = canina.
Schrenkiana, *Crép. in Bull. Soc. Bot. Belg.* xiv. (1875) 31.—As. centr.
Schultzii, Ripart, in F. Schultz, Fl. Fr. et Allem. 254 = glauca.
sciaphila, Schur, Enum. Pl. Transs. 202 = rubiginosa.
sclerophylla, Scheutz, Stud. Skand. Rosa (1872), 20 = canina.
Scopoliana, Tratt. Ros. [i.] ii. 229 = cinnamomea.

ROSA :—

scopulorum, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 545.—Gallia.
scotica, Mill. Gard. Dict. ed. VIII. n. 5 = spinosissima.
scotophylla, *Boullu, in Cariot, Etude Fleurs*, ed. VII. ii. (1884) 242.—Gallia.
semiglabra, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 240 = canina.
semiglandulosa, Ripart, l. c. xv. (1876) 500 = canina.
semperflorens, *Curt. Bot. Mag.* t. 284.—China.
semperflorens, Desf. Tabl. ed. I. 175 = damascena.
semperflorens, Desv. Obs. (1818) 154 = gallica.
sempervirens, Bast. Ess. Fl. Maine-et-Loire, 188 = canina.
sempervirens, Dup. ex Steud. Nom. ed. I. 702 = moschata.
sempervirens, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. ii. 368 = Leschenaultiana.
sempervirens, *Linn. Sp. Pl.* 492.—Europ.; Oriens; Ind. or.
sempervirens, Roth, Tent. Fl. Germ. i. 218 = rubiginosa.
senticosa, Achar. in Vet. Akad. Handl. Stockh. (1813) 91. t. 13 = canina.
separabilis, *Déségl. in Bull. Soc. Bot. Belg.* xix. (1880) 30.—Europ.
seperina, Sauzé, ex Déségl. l. c. xv. (1876) 217 = arvensis.
sepincola, Sw. ex Fries, Novit. ed. II. 147 = canina.
sepium, Lam. Fl. Fr. iii. 129; Borck. ex Rau, Enum. 90 = canina.
sepium, Thuill. Fl. Par. ed. II. 252 = agrestis.
seposita, *Déségl. in Bull. Soc. Bot. Belg.* xix. (1880) 28.—Gallia.
septicola, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 149 = micrantha.
Serafini, Viv. Fl. Lib. Spec. 67 = seq.
Seraphini, Viv. Fl. Cors. Sp. Nov. 8 = agrestis.
sericea, *Lindl. Ros. Monog.* 105. t. 12.—Reg. Himal.
Seringeana, Godr. Fl. Lorr. ed. II. 255 = tomentosa.
serotina, *Donn, ex Thory, Ros.* 142.—Hab. ?
serpens, Gilib. Fl. Lituan. ii. 238 = canina.
serpens, Wibel, Prim. Fl. Werth. 266 = arvensis.
Serrae, Dehnh. Rev. Napol. i. III. 163; ex Walp. Rep. ii. 11 = bracteata.
serrulata, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 218 = humilis.
sertorum, Wrede, ex Steud. Nom. ed. I. 703 = perfecta.
setigera, *Michx. Fl. Bor. Am.* i. 295.—Am. bor.
setosa, *Bess. ex Link, Enum. Hort. Berol.* ii. 60, nomen.—Podolia.
setulifera, Timb. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 511 = flexuosa.
Sherardi, Davies, Welsh Botanol. 49 = tomentosa.
sibirica, Steph. ex Ledeb. Fl. Ross. ii. 78 = canina.
sibirica, Tratt. Ros. ii. 230 = spinosissima.
sicula, Tratt. l. c. 68 = Heckeliana.
Sieboldi, *Crép. ex Franch. & Sav. Enum. Pl. Jap.* i. 136.—Japon.
Silverhielmii, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 195.—Rossia.
similata, Puget, ex Déségl. in Billotia, i. (1864) 38 = canina.
Simkovicsii, *Kmet, in Oestr. Bot. Zeitschr.* xxxiv. (1884) 18.—Europ.
simplex, Scop. Fl. Carn. ed. II. i. 353 = cinnamomea.
simplicifolia, Salisb. Prod. 359; Pers. Syn. ii. 47 = berberifolia.
sinica, Ait. Hort. Kew. ed. II. iii. 261 = laevigata.
sinica, Desf. ex DC. Prod. ii. 606 = blanda.
sinica, Hook. Bot. Mag. t. 2847 = laevigata.
sinica, Murr. Syst. ed. XIII. 394 = indica.
sinica, Tratt. Ros. i. 89 = semperflorens.
sinuatidens, Christ, in Gremli, Excurs. Schw. ed. IV. 168 = canina.
Sirei, Christ, l. c. 162 = spinosissima.
Solandri, Tratt. Ros. ii. 150 = blanda.
solstitialis, Bess. Prim. Fl. Galic. i. 324 = canina.
soongarica, Bunge, in Ledeb. Fl. Alt. ii. 226 = cinnamomea.
sorbifolia, Hort. ex Steud. Nom. ed. I. 703; ed. II. ii. 473 = spinosissima.
spathulaefolia, *Vukot. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 48.—Europ.
speciosa, *Andr. Ros.* t. 26.—Hab. ?
speciosa, Déségl. in Billotia, i. (1864) 39 = flexuosa.

ROSA:—
sphaerocarpa, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 279 = canina.
sphaeroidea, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 338.—Europ.
sphaerophora, Ripart, ex Nym. Consp. 233 = micrantha.
spherica, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 333 = canina.
spinatifolia, Steud. Nom. ed. II. ii. 473, sphalm. = spinulifolia.
spinosissima, *Linn. Sp. Pl.* 491.—Europ.; As. Min.
spinosissima, Pesneau, Cat. Pl. Loire-Inf. (1841) 181 = baltica.
spinosissima, Wahlenb. Fl. Upsal. 171 = cinnamomea.
spinulifolia, Dematra, Monog. Ros. 8 = rubiginosa.
spithamea, A. Gray, in Bot. Calif. ii. 444 = californica.
Sprengeliana, Tratt. Ros. ii. 163 = R. Carolina.
spreta, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 89 = spinosissima.
spuria, Puget, ex Déségl. l. c. xxviii. (1872) 109 = canina.
squarrosa, Rau, Enum. Ros. 77 = canina.
stebonensis, Roess. Rosen, i. 238 = canina.
stenocarpa, Désegl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 113 = micrantha.
stephanocarpa, Déségl. & Ripart, l. c. 115 = montana.
stipulacantha, Bast. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 356 = canina.
stipularis, Mérat, Fl. Par. 192 = canina.
striata, Wrede, ex Steud. Nom. ed. I. 703 = gallica.
stricta, Macoun & Gibs. in Trans. Bot. Soc. Edinb. xii. (1876) 324 = blanda.
stricta, Muhl. Cat. 50 = spinosissima.
stupenda, Crép. ex Boiss. Fl. Orient. ii. 684 = macrocarpa.
stylosa, *Desv. Journ. Bot.* ii. (1809) 317.—Europ.
stylosa, Pers. Syn. ii. 47 = arvensis.
suaveolens, Pursh, Fl. Am. Sept. i. 346 = rubiginosa.
suavifolia, Lightf. Fl. Scot. 262 = rubiginosa.
suavis, *Willd. Enum. Hort. Berol. Suppl.* 37.—Hab.?
subalba, *Hort. ex Steud. Nom.* ed. II. ii. 472.—China.
subalbida, *Vukot. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 21.—Europ.
subcinnamomea, Borb. in Oestr. Bot. Zeitschr. xxxiv. (1884) 261 = canina.
subcristata, Baker, in Naturalist, i. (1864) 97 = canina.
subdola, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 528 = agrestis.
subdola, Kmet, ex H. Braun, in Flora, lxviii. (1885) 114 = Borbasiana.
Suberti, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 352 = canina.
subglobosa, Sm. Engl. Fl. ii. 38 = tomentosa.
subhispida, *Vukot. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 39.—Europ.
subhybrida, *Borb. ex Vukot. l. c.* 20.—Europ.
subinermis, *Bess. ex Déségl. in Bull. Soc. Sc. Angers*, (1878).—*Cf. Bull. Soc. Bot. Fr.* xxv. (1878) *Bibl.* 107.—Europ.
subinermis, Chab. in Cariot, Etud. Fleurs, ii. (1865) 173 = gallica.
subintrans, Gren. ex Déségl. in Billotia, i. (1869) 121 = canina.
sublaevis, *Boullu, in Bull. Soc. Bot. Fr.* xxiii. (1876) p. lxi.—Gallia.
submirra ×, Vukot.—Cf. Oestr. Bot. Zeit. xxxiv. (1884) 260.—Europ.
submissa, *Vukot. in Rad Jugos. Akad. Zagreb.* lxix. (1884) 17.—Europ.
submitis, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 332 = canina.
subolida, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 124 = flexuosa.
subrubra, Thory & Red. ex Steud. Nom. ed. II. ii. 472 = R. Eglanteria.
subsessiliflora, *Boullu, in Feuill. Jeunes Nat.* x. (1879–80) 89.—Europ.
subumbellata, Schur, Enum. Pl. Transs. 203 = indica?
subvillosa, *Léman, in Journ. Phys.* lxxxvii. (1818) 366.—Gallia.
Sufferti, Kirsch. Fl. Alsat. i. 247 = tomentosa.
sulfurea, Thory, Ros. 123 = R. Eglanteria.
sulphurea, Ait. Hort. Kew. ed. I. ii. 201 = hemisphaerica, *Herrm.*
superbissima, *Waitz, ex Steud. Nom.* ed. I. 703; ed. II. ii. 472.—Hab.?

ROSA:—
surculosa, Woods, in Trans. Linn. Soc. xii. (1817) 228 = canina.
Swartziana, Fries, Fl. Halland. 87 = canina.
Swartzii, *Billb. in Svensk Fl.* t. 541 = praec.
sylvatica, Gater. Pl. Montaub. 94 = gallica.
sylvestris, *Crantz, Stirp. Austr.* ed. I. fasc. ii. 30; ed. II. 84.—Europ.
sylvestris, Herrm. Diss. 10 = arvensis.
sylvestris, Lindl. Syn. Brit. Fl. 101 = tomentosa.
sylvestris, Schrank, Baier. Fl. ii. 23 = canina.
sylvicola, Déségl. & Ripart, in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 122 = rubiginosa.
sylvularum, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 333.—Europ.
syntrichostyla, Ripart, l. c. 312 = canina.
systyla, Bast. Fl. Maine-et-Loire, Suppl. 31 = stylosa.
systylomorpha, Gren. in Mém. Soc. Emul. Doubs, Sér. IV. ix. (1875) 229 = canina.
sytnensis, *Kmet, ex Kern. Sched. Fl. Exsicc.* ii. 28.—Hungar.
Szaboi, *Borb.—Cf. Oestr. Bot. Zeitschr.* xxxii. (1882) 338.—Europ.
tartarica, *Roess. Rosen*, ii. 52.—Hab.?
taurica, Bieb. Fl. Taur. Cauc. i. 394 = canina.
teneriffensis, Donn, Cat. Hort. Cantab. ed. VI. 143 (Quid?).—Ins. Teneriff.
tenuiglandulosa, Mérat, Fl. Par. 189 = rubiginosa.
tenuis, Beck, Fl. Frankf. i. 285 = spinosissima.
terebinthacea, Bess. Enum. Pl. Volh. 21, 61, 66 = tomentosa.
terebinthacea, *Bess. ex Spreng. Syst.* ii. 551.—Galicia.
terebinthinacea, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 159 = flexuosa.
terebinthinacea, Gren. ex Baker, in Journ. Linn. Soc. xi. (1871) 235 = canina.
ternata, Poir. Encyc. vi. 284 = laevigata.
tetrapetala, Royle, Illustr. Bot. Himal. 208. t. 42 = sericea.
Thea, Savi, Fl. Ital. ii. 17. t. 47 = indica.
Thedenii, *Scheutz, Stud. Skand. Ros.* (1872) 27.—Europ. bor.
Thomasii, Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 24 = flexuosa.
Thoryi, Tratt. Ros. ii. 85 = multiflora?
Thunbergii, Tratt. l. c. i. 86 = multiflora.
thuringiaca, *Crép. in Bull. Soc. Bot. Belg.* viii. (1869) 319.—Europ.
thuringiaca, *Link, Handb.* ii. 94.—Europ. med.
thyrsiflora, *Leroy, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 204.—Japon.
Timeroyi, Chab. in Cariot, Etud. Fleurs, ed. V. ii. 189 = canina.
tirolensis, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 331 = canina.
tolosana, *Timb. in Bull. Soc. Hist. Nat. Toul.* iv. (1870) 173.—Europ.
tomentella, Léman, in Bull. Soc. Philom. (1818) 94 = canina.
tomentosa, *Sm. Fl. Brit.* ii. 539.—Europ.; As. Min.
tortuosa, Wierzb. ex Baker, in Journ. Linn. Soc. xi. (1871) 232 = canina.
torulosa, Wall. ex Hook. f. Fl. Brit. Ind. ii. 366 = macrophylla.
Touranginiana, Déségl. & Ripart, in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 102 = canina.
trachyphylla, Bor. Fl. Centr. Fr. ed. III. ii. 228 = canina.
trachyphylla, *Rau, Enum. Ros.* 124.—Europ.
transalpina, Christ, in Gremli, Excurs. Schw. ed. IV. 163 = arvensis.
transiens, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 8 = glauca.
transmota, Crép. in Bull. Soc. Bot. Belg. viii. (1876) 239 = canina.
transsilvanica, Schur, Enum. Pl. Transs. 202 = canina.
trichocarpa, Waitz, ex Tratt. Ros. ii. 45 = alba?
trichoidea, *Ripart, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 386.—Gallia.
trichoneura, Ripart, ex Crép. l. c. viii. (1869) 240 = canina.
tricolor, *Hort. ex Steud. Nom.* ed. I. 703; ed. II. ii. 473.—Hab.?
triflora, Chab. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 245 = gallica.
trifoliata, Bosc, Dict. Agric. ed. II. xiii. 280 = laevigata.

ROSA:—
trifoliolata, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 213 = setigera.
triphylla, Roxb. Hort. Beng. 38; Fl. Ind. ii. 515 = anemoneflora, *Fortune.*
tristis, Kern. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 568 = tomentosa.
tuguriorum, Willd. Enum. Hort. Berol. 544 = arvensis.
tunoniensis, Déségl. in Billotia, i. (1864) 46 = tomentosa.
turbinata, *Ait. Hort. Kew.* ed. I. ii. 206.—Europ.
turbinata, Vill. Hist. Pl. Dauph. iii. 550 = alpina.
turbinella, Sw. ex Spreng. Syst. ii. 554 = cinnamomea.
turgida, Pers. Syn. ii. 49 = blanda.
turkestanica, *Regel, in Act. Hort. Petrop.* v. (1877) 349.—Turkestan.
tuschetica, *Boiss. Fl. Orient.* ii. 673.—Reg. Caucas.
tyroliensis, Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 495 = *tirolensis.*
ucenensis, Boullu, in Cariot, Etud. Fleurs, ed. VII. ii. (1884) 256 = rubiginosa.
uliginosa, *Bray, ex Fleisch. & Lindem. Fl. Deutsch. Ostseeprov.* 179 (Quid?).—Prov. balt.
uliginosa, *Gilib. Fl. Lithuan. Inchoat. Collect.* v. 240. —Hab.?
umbellata, Leers, Fl. Herborn. 117, 286 = rubiginosa.
umbellata, Leyss. Fl. Hal. 92; Libert, in Lej. Fl. Spa, ii. 313 = canina.
umbellifera, Sw. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 570 = tomentosa.
Uncinella, Bess. Cat. Pl. Hort. Crem. (1811) Suppl. iv. 21 = canina.
uncinelloides, Puget, ex Nym. Consp. 235 = canina.
undulaeflora, *Andr. Ros.* t. 108.—Am. bor.
unguicularis, Bertol. Misc. Bot. xxii. 15. t. 3 = Webbiana.
unguiculata, Desf. Tabl. ed. I. 175 = centifolia.
unica, Hort. ex Hoffmgg. Verz. Pfl. 109 = centifolia.
urbica, Léman, in Bull. Soc. Philom. (1818) 93 = canina.
uriensis, Lagg. & Puget, ex Cottet, in Bull. Soc. Murith. iv. (1875) 28 = tomentosa.
usitatissima, Gater. Pl. Montaub. 94 = alba.
ustalica, Vukot. in Rad Jugos. Akad. Zagreb. lxix. (1884) 14 = alpina.
Vaillantiana, Bor. ex Baker, in Journ. Linn. Soc. xi. (1871) 224 = agrestis.
valesiaca, Lagg. & Puget, ex Cottet, in Bull. Soc. Murith. ii. (1873) 43 = canina.
Vanheurckiana, *Crép. ex Boiss. Fl. Orient.* ii. 683.—Armenia.
varia, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 775 = gallica.
varians, Pohl, ex Link, Handb. ii. 93 (Quid?).—Europ.
Vayredae, *Costa, ex Vayr. in Anal. Soc. Esp. Hist. Nat.* xi. (1882) 83.—Hab.?
velutina, [Clairv.] Man. Herb. 163 = tomentosa.
velutinaeflora, Déségl. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 100 = gallica.
venosa, Sw. ex Spreng. Syst. ii. 554 = glauca.
Ventenatiana ×, Red. & Thory, Ros. iii. 83.
venusta, Scheutz, Stud. Skand. Rosa (1872), 36 = mollissima.
venusta, Waitz, ex Reichb. Fl. Germ. Excurs. 613 = alpina.
vepretorum, Link, Handb. ii. 91 = canina.
verecunda, *Waitz, ex Tratt. Ros.* i. 21.—Hab.?
Verloti, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 274 = canina.
versicolor, Hort. ex Steud. Nom. ed. I. 703 = gallica.
versicolor, Steud. l. c. = spinosissima.
versicolor, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. ii. (1874) 63 = agrestis.
verticillacantha, Mérat, Fl. Par. 190 = canina.
vestita, Godet, Fl. Jura, 210 = tomentosa.
vestita, Sternb. in Flora, ix. (1826) 1. Beil. 77 = ferruginea.
Vetteri, Favr. in Gremli, Neue Beitr. Fl. Schw. i. 10 = cinnamomea.
Viburnum, Red. Ros. ii. 58 = centifolia.
vicina, Crép. in Bull. Soc. Bot. Belg. viii. (1869) 241 = canina.
Villarsiana, Sieber, ex Reichb. Fl. Germ. Excurs. 612 = spinosissima.
Villarsii, Tratt. ex Link, Handb. ii. 86 = alpina.

ROSA:—
villosa, Bieb. Fl. Taur. Cauc. i. 395 = mollissima.
villosa, Falk, Beitr. ii. 191 = tomentosa.
villosa, *Linn. Sp. Pl.* 491.—Europ.; As. med.
villosa, Pall. Fl. Ross. ii. 63 = pomifera.
villosa, Sm. Engl. Bot. t. 583 = involuta, pomifera.
villosiuscula, Ripart, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 266 = canina.
villosula, Paill. Rev. Fr. Conté, 362; ex Billotia, i. (1869) 119 = canina.
Vilmosina, Hort. ex Steud. Nom. ed. I. 704; ed. II. ii. 473 = gallica.
viminea, *Lindl. Ros. Monog.* 49.—Rossia.
vinacea, Baker, in Naturalist, i. (1864) 101 = canina.
vinealis, Ripart, ex Déségl. in Billotia, i. (1864) 36 = canina.
vinetorum, Ripart, ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 499 = canina.
vinodora, Kern. in Oestr. Bot. Zeitschr. xix. (1869) 329 = agrestis.
violacea, Hort. ex Roess. Rosen, i. 200 = gallica.
virescens, Déségl. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 73 = gallica.
virgata, Gremli, Excurs. Schw. ed. II. 177 = canina.
virginalis, Wrede, ex Steud. Nom. ed. I. 704 = centifolia?
virginea, Ripart, ex Déségl. in Journ. Bot. xii. (1874) 167 = stylosa.
virginiana, Mill. Gard. Dict. ed. VIII. n. 10; Duroi, Harbk. Baumz. ii. 353 = R. Carolina.
virginica, Roess. Rosen, t. 13; Spreng. Nov. Prov. 36 = praec.
virgultorum, Ripart, ex Déségl. in Billotia, i. (1864) 44 = agrestis.
viridicata, Puget, ex Crép. in Bull. Soc. Bot. Belg. viii. (1869) 239, 266 = canina.
viscida, Puget, ex Crép. l. c. 241 = canina.
viscida, *Schultz, in Flora*, viii. (1825) II. *Beibl.* 78.—Europ.
viscida, *Tausch, ex Opiz, Boehm. Gew.* (1823) 61.—Bohemia.
viscidula, *Gren. ex Bouv. in Bull. Inst. Nat. Genèv.* xix. (1875) 107.—Europ.
viscosa, Jan, Cat. 8 = agrestis.
vogesiaca, Desport. Ros. Gall. (1828) 88 = glauca.
Vukotinovicii, Borb. ex Vukot. in Rad Jugos. Akad. Zagreb. lxix. (1884) 23 = alpina.
vulpina, Wallr. Rosae Hist. (1828) 201 = R. Eglanteria.
Waiziana, Reichb. in Moessl. Handb. ed. II. ii. 870 = Reichenbachiana.
Wallichii, Sabine, in Wall. Cat. n. 687 = Leschenaultiana.
Wallichii, Tratt. Ros. ii. 193 = sericea.
wasserburgensis, Kirsch. Fl. Alsat. i. (1852) 247 = flexuosa.
Watsoni, Baker, in Naturalist, i. (1864) 98 = canina.
Webbiana, *Wall. Cat.* n. 683.—Reg. Himal.
Wettsteinii, *H. Braun, in Oestr. Bot. Zeitschr.* xxxv. (1885) 303.—Europ.
Wichurae, C. Koch, Wochenschr. xii. (1869) 202 = multiflora.
Wichuraiana, Crép. ex Déségl. in Bull. Soc. Bot. Belg. xv. (1876) 204 = Luciae.
Wietziana, Auct. ex Reichb. in Moessl. Handb. ed. II. ii. 871 = Reichenbachiana.
Willdenowii, *Spreng. Syst.* ii. 547.—Reg. Casp.; Sibir.
Willdenowii, Tratt. Ros. ii. 81 = pulchella.
Wilsoni, Borr. in Hook. Brit. Fl. ed. I. 228 = involuta.
Wolfgangiana, Bess. Enum. Pl. Volh. 61, 67 = gallica.
Wolfii, *La Soie, ex Déségl. in Bull. Soc. Bot. Belg.* xv. (1876) 345.—Europ.
Woodsii, A. Gray, in Mem. Am. Acad. iv. (1849) 42 = californica.
Woodsii, *Lindl. Ros. Monog.* 21.—Am. bor.
Woodsii, Regel, in Act. Hort. Petrop. (1877) 299 = nutkana.
Wulfenii, *Tratt. Ros.* i. 200.—Europ.
xanthina, *Lindl. Ros. Monog.* 132.—China.
xerampelina, *Hort. ex Breit. Hort. Breit.* 433.—Europ.
Ywara, *Carr. in Rev. Hort.* (1876) 357.—Japon.
zagrabiensis, *Vukot. & H. Braun, ex Kern. Sched. Fl. Exsicc.* iii. 55.—Croatia.
zalana, *Wiesb. in Oestr. Bot. Zeitschr.* xxix. (1879) 142.—Europ.

ROSA:—
zamensis, *Simk. & H. Braun, in Termész. Füzetek*, ix. (1885) 41.—Hungar.
Zeytheriana, *Dehn. Revista Napolit.* i. III. 165; *ex Walp. Rep.* ii. 12.—Hab.?

ROSALESIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 9 (1824) = **Brickellia**, Ell. (Compos.).
glandulosa, La Llave, l. c. = B. Rosalesia.

ROSANOVIA, Benth. & Hook. f. Gen. ii. 1004, sphalm. (1876) = seq.

ROSANOWIA, Regel, Gartenfl. xxi. (1872) 33. t. 712 = **Sinningia**, Nees (Gesner.).
conspicua, Regel, l. c.—Bras.
hybrida ×, Nagy, in Illustr. Wien. Gartenfl. Zeitschr. x. (1885) 126.—Bras.
ornata ×, Van Houtte, in Fl. de Serres, xxiii. (1882) 149.

ROSAURA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Wedelia**?, Jacq. (Compos.).

ROSCHERIA, H. Wendl. ex Baker, Fl. Maurit. 386 (1877). *PALMAE*, Benth. & Hook. f. iii. 913.
melanochoetes, *H. Wendl. l. c.* 387.—Ins. Mascar.

ROSCIA, D. Dietr. Syn. Pl. i. 552 (1839), sphalm. = BOSCIA, Thunb. = **Toddalia**, Juss. (Rutac.).

ROSCOEA, Roxb. Hort. Beng. 46 (1814); Fl. Ind. iii. 54 (1832) = **Sphenodesma**, Jack (Verbenac.).
pentandra, Roxb. ll. cc. 46; 54 = S. pentandra.
tomentosa, Roxb. ll. cc. [95]; 56 = Conyza tomentosa.
villosa, Roxb. l. c. [95]; 55 = Laggera aurita.

ROSCOEA, Sm. Exot. Bot. ii. 97. t. 108 (1804). *SCITAMINEAE*, Benth. & Hook. f. iii. 641.
CAUTLEA, Royle, Illustr. Bot. Himal. 361 (1839).
alpina, *Royle, Illustr. Bot. Himal.* 361.—Reg. Himal.
capitata, *Sm. in Trans. Linn. Soc.* xiii. (1822) 461.—Nepal.
elatior, *Sm. l. c.* 460.—Reg. Himal.
exilis, Sm. ex Horan. Prod. Scitam. 21 = purpurea.
gracilis, Sm. in Trans. Linn. Soc. xiii. (1822) 460 = elatior.
lutea, *Hassk. in Flora*, xlvii. (1864) 21.—Ins. Philipp.
lutea, Royle, Illustr. Bot. Himal. 361. t. 89 = elatior.
nigro-ciliata, *Hassk. in Flora*, xlvii. (1864) 21.—Ins. Philipp.
procera, Wall. Pl. As. Rar. iii. t. 242 = purpurea.
purpurea, *Sm. Exot. Bot.* ii. 97. t. 108.—Reg. Himal.; Burma.
spicata, *Sm. in Trans. Linn. Soc.* xiii. (1822) 460.—Reg. Himal.

ROSCYNA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 364 = **Hypericum**, Linn.
americana, Spach, l. c. = H. pyramidatum.
Gebleri, Spach, l. c. = H. Gebleri.
Gmelini, Spach, l. c. = H. Ascyron.
japonica, Blume, Mus. Bot. Lugd. Bat. ii. 21 = H. japonicum.

ROSEA, Klotzsch, in Monatsb. Akad. Berl. (1853) 501 = **Tricalysia**, A. Rich. (Rubiac.).
crassifolia, Klotzsch, l. c. = T. crassifolia.
jasminiflora, Klotzsch, l. c. = T. jasminiflora.

ROSEA, Mart. Nov. Gen. et Sp. ii. 58 (1826) = **Iresine**, P. Br. (Amarant.).
elatior, Mart. l. c. 59. t. 155 = I. elatior.
Poeppigiana, Klotzsch, ex Seub. in Mart. Fl. Bras. v. I. 226 = I. elatior.
virgata, Klotzsch, ex Seub. l. c. = I. elatior.

ROSENBERGIA, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 30 = **Cobaea**, Cav. (Polemon.).
gracilis, Oerst. l. c. 31 = C. gracilis.
penduliflora, Karst. Fl. Columb. i. 27. t. 14 = C. penduliflora.

ROSENIA, Thunb. Nov. Gen. 161 (1801). *COMPOSITAE*, Benth. & Hook. f. ii. 327.
glandulosa, *Thunb. l. c.* 162.—Afr. austr.
spinescens, *DC. Prod.* vi. 280.—Afr. austr.

ROSILLA, Less. Syn. Comp. 245 (1832). *COMPOSITAE*, [Benth. & Hook. f. ii. 397].
lutea, *Less. l. c.*—Mexic.

ROSLINIA, G. Don, Gen. Syst. iv. 203 (1837) = ROESLINIA, Moench = **Chironia**, Linn. (Gentian.).

ROSLINIA, Neck. Elem. i. 357 (1790) = **Justicia**, Houst. (Acanth.).

ROSMARINUS, [Tourn.] Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1197.
angustifolius, Mill. Gard. Dict. ed. VIII. n. 1 = officinalis.
chilensis, Molina, Sagg. Chile, 158 = Sphacele campanulata.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 25 = officinalis?
eriocalix, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 44 = officinalis.
flexuosus, Jord. & Fourr. l. c. = officinalis.
latifolius, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
laxiflorus, Noé, ex Lange, in Kjoeb. Vidensk. Meddel. (1863) 12 = officinalis.
officinalis, *Linn. Sp. Pl.* 23.—Reg. Mediterr.
prostratus, Fl. Corc. in Ionios Anthol. ii. (1834) 446 = officinalis.
rigidus, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 43 = officinalis.
tenuifolius, Jord. & Fourr. l. c. = officinalis.

ROSPIDIOS, A. DC. in DC. Prod. viii. 220 (1844) = **Diospyros**, Linn. (Ebenac.).
vaccinioides, A. DC. l. c. = D. vaccinioides.

ROSSENIA, Vell. Fl. Flum. 31 (1825); i. t. 77 (1827) = **Cusparia**, Humb. (Rutac.).
pentaphylla, Vell. ll. cc.—Bras.

ROSSINA, Steud. Nom. ed. II. ii. 652 (1841), sphalm. = POSSIRA, Aubl. = **Swartzia**, Schreb. (Legumin.).

ROSSMAESSLERA, Reichb. Nom. 113 (1841) = **Gilia**, Ruiz & Pav. (Polemon.).

ROSSMANNIA, Klotzsch, in Monatsb. Akad. Berl. (Maerz 1854) 125; et in Abh. Akad. Berl. 1854 (1855) 99. t. 9 A = **Begonia**, Linn.
repens, Klotzsch, ll. cc. = B. repens.

ROSSOLIS, Adans. Fam. ii. 245 (1763) = **Drosera**, Linn.
rotundifolia, Moench, Meth. 57 = D. rotundifolia.
septemtrionalis, Scop. Fl. Carn. ed. II. i. 232 = D. rotundifolia.

ROSTELLARIA, Gaertn. f. Fruct. iii. 135. t. 207 (1805). 'Genus valde dubium,' Benth. & Hook. f. ii. 653.
SARCORHYNA, Presl, Bot. Bemerk. 103 (1844).
Lessertiana, *Gaertn. f. l. c.*—Hab.?

ROSTELLARIA, Nees, in Wall. Pl. As. Rar. iii. 76 (1832) = **Justicia**, Houst. (Acanth.).
adenostachya, Nees, l. c. 101 = J. procumbens.
crinita, Nees, l. c. = J. micrantha.
diffusa, Nees, l. c. 100 = J. diffusa.
haplostachya, Benth. & Hook. f. Gen. ii. 1110 = J. haplostachya.
hedyotidifolia, Nees, in Wall. Pl. As. Rar. iii. 100 = J. diffusa.
japonica, Carr. in Rev. Hortic. (1866) 400 = J. procumbens?
micrantha, Steud. Nom. ed. II. i. 838, in syn., sphalm. = seq.
mollissima, Nees, in Wall. Pl. As. Rar. iii. 101 = J. simplex.
parviflora, Benth. in Hook. Niger Fl. 481 = J. tenella.
peploides, Nees, in Wall. Pl. As. Rar. iii. 101 = J. quinqueangularis.
procumbens, Nees, l. c. = J. procumbens.
quinquangularis, Nees, l. c. = J. quinqueangularis.
reptans, Nees, ex Benth. & Hook. f. Gen. ii. 1110 = J. trifolioides.
rotundifolia, Nees, in Wall. Pl. As. Rar. iii. 100 = J. diffusa.
Schimperi, Hochst. in Flora, xxiv. (1841) I. Intell. 24 = Rhinacanthus calcaratus?

ROSTELLARIA :—
sexangularis, Hochst. in Flora, xxv. (1842) Beil. 1. 144 = Dicliptera sexangularis.
tenella, Nees, ex Benth. & Hook. f. Gen. ii. 1110 = Justicia tenella.
Vahlii, Nees, in Wall. Pl. As. Rar. iii. 102 = J. diffusa.

ROSTELLULARIA, Reichb. Handb. 190 (1837) = **Justicia**, Houst. (Acanth.).
abyssinica, Brongn. ex Nees, in DC. Prod. xi. 372 = J. simplex.
adenostachya, Nees, l. c. 373 = J. procumbens.
Blancoi, Hassk. in Flora, xlvii. (1864) 54 = Diantherа americana.
brachystachya, Nees, in DC. Prod. xi. 369 = J. brachystachya.
crenulata, Nees, l. c. = J. tenella.
crinita, Nees, l. c. 373 = J. micrantha.
diffusa, Nees, l. c. 371 = J. diffusa.
glandulosa, Nees, l. c. 373 = J. simplex.
gracilis, Wight, Ic. t. 1541 = Rungia linifolia.
haplostachya, Nees, in DC. Prod. xi. 369 = J. haplostachya.
hedyotidifolia, Nees, l. c. 370 = J. diffusa.
juncea, Nees, l. c. 376 = J. procumbens.
media, Nees, l. c. 374 = J. procumbens.
mollissima, Nees, l. c. 373 = J. procumbens.
obtusa, Nees, l. c. 374 = J. Blumeana.
peploides, Nees, l. c. 375 = J. quinqueangularis.
pogonanthera, F. Muell. in Linnaea, xxv. (1852) 431 = J. procumbens.
procumbens, Nees, in DC. Prod. xi. 371 = J. procumbens.
psychotrioides, Nees, l. c. 370 = J. psychotrioides.
pumila, Hochst. ex Nees, l. c. 372 = J. simplex.
quinquangularis, Nees, l. c. 375 = J. quinqueangularis.
reptans, Nees, l. c. 368 = J. trifolioides.
rotundifolia, Nees, l. c. 370 = J. simplex, diffusa.
Royeniana, Nees, l. c. 373 = J. Royeniana.
sarmentosa, Zoll. ex Nees, l. c. 370 = J. procumbens.
simplex, Wight, Ic. iv. t. 1542 = J. notha.
tenella, Nees, in DC. Prod. xi. 369 = J. tenella.
trichochila, Miq. in Journ. Bot. Néerl. i. (1861) 118.—China.
Vahlii, Nees, in DC. Prod. xi. 376 = J. diffusa.

ROSTKOVIA, Desv. Journ. Bot. i. (1808) 324. t. 12. *JUNCACEAE*, Benth. & Hook. f. iii. 867.
MARSIPPOSPERMUM, Desv. Journ. Bot. i. (1808) 328. t. 12.
brevifolia, Phil. in Linnaea, xxix. (1857–58) 76 = Distichia brevifolia.
clandestina, Phil. l. c. = Distichia clandestina.
gracilis, *Hook. f. Fl. Antarct.* 83.—Ins. Antarct.
gracilis, *Phil. in Linnaea*, xxix. (1857–58) 75.—Chili.
grandiflora, *Hook. f. Fl. Antarct.* 81.—Ins. Antarct.
magellanica, Hook. f. l. c. = sphaerocarpa.
Novae-Zelandiae, J. Buch. in Trans. N. Z. Inst. iv. (1872) 227 = gracilis, *Hook. f.*
sphaerocarpa, *Desv. Journ. Bot.* i. (1808) 327.—Reg. Magellan.

ROSTRARIA, Trin. Fund. Agrost. 149 (1820) = **Trisetum**, Pers. (Gramin.).
laevis, Trin. l. c. 150 = T. paniceum.
pubescens, Trin. l. c. = T. phleoides.

ROTALA, Linn. Mant. ii. 143 (1771). *LYTHRARIAE*, [Benth. & Hook. f. i. 776].
alata, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 171.—Ind. or.
apetala, F. Muell. Fragm. iii. 108 = Ammannia Rotala.
cordata, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 172.—Ind. or.
decussata, DC. Prod. iii. 76 = Ammannia pentandra.
decussata, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 467.—Afr. trop.
decussata, Hiern, l. c., partim = illecebroides.
densiflora, Koehne, in Engl. Bot. Jahrb. i. (1881) 164 = Ammania pentandra.
dentifera, *Koehne, l. c.* 161.—Mexic.
diandra, F. Muell. in Trans. Phil. Soc. N. S. W. xiv. (1881) 89 = Ammannia diandra.
elatinoides, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 466.—Afr. trop.
filiformis, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 468.—Afr. trop.

ROTALA :—
fimbriata, *Wight, Ic. t.* 217.—Ind. or.
floribunda, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 156.—Ind. or.
fontinalis, *Hiern, in Oliver, Trop. Afr.* ii. 468.—Afr. trop.
hexandra, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 167.—Ind or.
illecebroides, *Koehne, l. c.* 161.—Ind. or.
indica, Koehne, l. c. 172 = Ammania peploides.
leptopetala, *Koehne, l. c.* 162.—As. or.
macrandra, *Koehne, l. c.* 176.—Ind. or.
mexicana, *Cham. & Schlecht. in Linnaea*, v. (1830) 567.—Mexic.
myriophylloides, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 469 = Hydrolythrum Wallichii.
nummularia, *Welw. ex Hiern, l. c.* 468.—Afr. trop.
occultiflora, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 152.—Ind. or.
pentandra, Wight, Ic. i. sub t. 217 = Ammannia pentandra.
petaloidea, Wight, ex Steud. Nom. ed. II. ii. 474 = Ammannia Rotala.
pusilla, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 128 = mexicana.
ramosior, *Koehne, in Mart. Fl. Bras.* xiii. II. 194.—Bras.
repens, Koehne.—Cf. Sitzb. Bot. Ver. Prov. Brandenb. (1877) 48 = Quartinia repens.
Ritchiei, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 386.—Ind. or.
rotundifolia, Koehne, l. c. i. (1881) 175 = Ammannia rotundifolia.
Roxburghiana, Wight, Ic. t. 260. f. B. = Ammannia pentandra.
serpiculoides, *Welw. ex Hiern, in Oliver, Fl. Trop. Afr.* ii. 469.—Afr. trop.
simpliciuscula, *Koehne, in Engl. Bot. Jahrb.* i. (1882) 159.—Ind. or.
stagnina, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 467.—Afr. trop.
subrotunda, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 174.—Ind. or.
tenella, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 467.—Afr. trop.
tenuis, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 177.—Ind. or.
verticillaris, Hiern, in Oliver, Fl. Trop. Afr. ii. 467 = mexicana.
verticillaris, Linn. Mant. ii. 175 = Ammannia Rotala.
Wallichii, Koehne, in Engl. Bot. Jahrb. i. (1881) 154 = Hydrolythrum Wallichii.

ROTANG, Adans. Fam. ii. 24 (1763) = **Calamus**, Linn. (Palm.).

ROTANGA, Boerh. ex Crantz, Inst. i. 127 (1766) = **Calamus**, Linn. (Palm.).
Calamus, Crantz, l. c. = C. Rotang.

ROTBOLLA, Zumag. Fl. Pedem. i. 142 (1849), sphalm. = **Rottboellia**, Linn. f. (Gramin.).

ROTERBE, Klatt, in Mart. Fl. Bras. iii. 1. 543. t. 71 (1871) = BOTHERBE, Steud. = **Calydorea**, Herb. (Irid.).
bulbosa, Steud. ex Baker, in Journ. Linn. Soc. xvi. (1877) 134 = Libertia elegans.
campestris, Klatt, in Mart. Fl. Bras. iii. 1. 543 = C. campestris.
elegans, Steud. ex Baker, in Journ. Linn. Soc. xvi. (1877) 153 = Libertia elegans.
gracilis, Klatt, in Mart. Fl. Bras. iii. 1. 544 = C. nuda.
luteola, Klatt, l. c. = C. luteola.

ROTHECA, Rafin. Fl. Tellur. iv. 69 (1836) = **Clerodendron**, Linn. (Verben.).
bicolor, Rafin. l. c. = C. serratum.
ternifolia, Rafin. l. c. = C. serratum.

ROTHERIA, Meyen, Reise, i. 402 (1834) = **Cruckshanksia**, Hook. & Arn. (Rubiac.).
lanceolata, Meyen, l. c. = C. hymenodon.

ROTHIA, Lam. Journ. Hist. Nat. i. (1792) 16. t. 1 = **Hymenopappus**, L'Hérit. (Compos.).
carolinensis, Lam. l. c. = H. scabiosus.

ROTHIA, Pers. Syn. ii. 688 (1807). *LEGUMINOSAE*, Benth. & Hook. f. i. 477.
GOETZEA, Reichb. Consp. 204 (1828).
HARPELEMA, Jacq. f. Eclog. Pl. ii. 6 (1844).
WESTONIA, Spreng. Syst. iii. 230 (1826).
XEROCARPUS, Guill. & Perr. Fl. Seneg. Tent. 169. t. 44 (1832).
hirsuta, *Baker, in Oliver, Fl. Trop. Afr.* ii. 7.—Afr. trop.
trifoliata, *DC. Prod.* ii. 382.—Ind. or.; Austral.

ROTHIA, Schreb. Gen. 531 (1791) = **Andryala**, Linn. (Compos.).
andryaloides, Gaertn. Fruct. ii. 371. t. 174. f. 9 = A. Rothia.
argentea, Lapeyr. Hist. Abr. Pl. Pyr. 485 = A. ragusina.
cheiranthifolia, Roth, Catalect. fasc. i. 105 = A. integrifolia.
floccosa, Pomel, Nouv. Mat. Fl. Atl. 3 = A. floccosa.
laxiflora, Salzm. ex DC. Prod. vii. 247 = A. laxiflora.
runcinata, Roth, Catalect. fasc. i. 107 = A. integrifolia.
tenuifolia, Tineo, Cat. Pl. Hort. Panorm. (1827) 280 = A. dentata.

ROTHMANNIA, Thunb. in Vet. Acad. Handl. Stockh. (1776) 63 = **Gardenia**, Ellis (Rubiac.).
Bowieana, Benth. in Hook. Niger Fl. 384 = Randia macrantha.
capensis, Thunb. in Vet. Acad. Handl. Stockh. (1776) 65. f. 2 = G. Rothmannia.
longiflora, Salisb. Parad. Lond. t. 65 = Randia maculata.
macrophylla, R. Br. in Wall. Cat. n. 8304 = Randia macrophylla.
malleifera, Hook. Niger Fl. 383 = Randia malleifera.
Stanleyana, Hook. l. c. = Randia maculata.

ROTHROCKIA, A. Gray, in Proc. Am. Acad. xx. (1885) 295. *ASCLEPIADEAE*.
cordifolia, *A. Gray, l. c.*—Am. bor.

ROTMANNIA, Neck. Elem. ii. 450 (1790) = **Eperua**, Aubl. (Legumin.).

ROTTBOELIA, Scop. Introd. 233 (1777) = **Ximenia**, Linn. (Olacin.).

ROTTBOELLA, Murr. Syst. 124 (1784) = **Rottboellia**, Linn. f. (Gramin.).

ROTTBOELLIA, Host, Gram. Austr. i. 19. t. 24 (1801) = **Lepturus**, R. Br. (Gramin.).

ROTTBOELLIA, Linn. f. Nov. Gram. Gen. in Amoen. Acad. x. 22 (1779), partim. *GRAMINEAE*, Benth. & Hook. f. iii. 1129.
APOGONIA, Fourn. ex Benth. & Hook. f. Gen. iii. 1130 (1883).
COELORACHIS, Brongn. in Duperr. Voy. Coq. Bot. 64. t. 141 (1829).
CYMBACHNE, Retz. Obs. vi. 36 (1791).
LASIURUS, Boiss. Diagn. Ser. II. iv. 145 (1859).
MANISURIS, Linn. Mant. ii. 164 (1767).
PELTOPHORUS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188.
PHACELURUS, Griseb. Spicil. Fl. Rumel. ii. 423 (1844).
PHOLIURUS, Trin. in Spreng. Neue Entdeck. ii. 67 (1821).
STEGOSIA, Lour. Fl. Cochinch. 51 (1790).
adscendens, Brot. Fl. Lusit. i. 84 = Lepturus cylindricus.
agropyroides, *Hack. in Bol. Soc. Broter.* iii. (1885) 135.—Afr. trop.
altissima, Poir. Voy. Barb. ii. 105 = compressa.
anthephoroides, Steud. in Flora, xxix. (1846) 22 = Ischaemum antephoroides.
arabica, Willd. ex Steud. Nom. ed. II. ii. 474 = Lepturus filiformis.
articulata, Thunb. ex Roem. & Schult. Syst. ii. 787 = compressa.
arundinacea, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 444 = exaltata.
aurita, *Steud. Syn. Pl. Gram.* 361.—Am. trop.
Balansae, *Hack. in Mart. Fl. Bras.* ii. III. 312.—Paraguay.
barbata, Spreng. Syst. i. 300 = Xerochloa barbata.

ROTTBOELLIA:—
barbata, Willd. ex Steud. Nom. ed. II. ii. 474 = Stenotaphrum subulatum.
Beyreichii, Trin. ex Steud. l. c. (Quid?).—Hab.?
biflora, Roth, Neue Beitr. i. 121 = Lepturus pannonicus.
biflora, Roxb. Hort. Beng. 82; Fl. Ind. i. 357 (Lepturi sp.).—Ind. or.
brevis, *Chauvin, ex Steud. Syn. Pl. Gram.* 361.—Ins. Wagiou.
campestris, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 151.—Am. bor.
ciliaris, Willd. ex Steud. Nom. ed. II. ii. 474 (= *Aegilops ciliaris*).—Ind. or.
ciliata, Nutt. Gen. Am. i. 83 = Elionurus tripsacoides.
Coelorachis, *Forst. f. Prod.* 9.—N. Caled.
complanata, Sw. in Ges. Naturf. Fr. Berl. Mag. iv. (1810) 89. t. 5 = Stenotaphrum glabrum.
compressa, Beauv. Agrost. 109. t. 21. f. 81 = Stenotaphrum glabrum.
compressa, Guss. ex Steud. Nom. ed. II. ii. 474 = Lepturus filiformis.
compressa, *Linn. f. Suppl.* 114.—Reg. trop.
corrugata, *Baldw. in Am. Journ. Sc.* Ser. I. i. (1819) 355.—Am. bor.
corymbosa, Linn. f. Suppl. 114 = Ophiurus corymbosus.
cylindrica, *Torr. Bot. Whippl. Exped.* 103; *Chapm. Fl. S. U. St.* 579.—Am. bor.
cylindrica, Willd. Sp. Pl. i. 464 = Lepturus cylindricus.
Cymbachne, *Willd. l. c.* 465.—Ind. or.
denudata, Steud. Syn. Pl. Gram. 362 = exaltata.
digitata, *Sibth. & Sm. Fl. Graec.* i. t. 92.—Oriens.
dimidiata, Thunb. Prod. Pl. Cap. 23; Sw. in Ges. Naturf. Fr. Berl. Mag. iv. (1810) 89 = Stenotaphrum glabrum.
diversiflora, Salzm. ex Steud. Syn. Pl. Gram. 361 = Andropogon imberbis.
elegantissima, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 474 = Elionurus Royleanus.
erecta, Savi, in Nuov. Giorn. Lett. Pisan. iv. (1806) 250 = Lepturus filiformis.
exaltata, *Linn. f. Suppl.* 114.—Reg. trop.
fasciculata, Brongn. in Duperr. Voy. Coq. Bot. t. 14 = R. Coelorachis.
fasciculata, Lam. Illustr. i. 204 = compressa.
filifolia, Wright, in Sauv. Fl. Cub. 200 = loricata.
filiformis, Roth, in Usteri, Ann. Bot. x. (1794) 38 = Lepturus filiformis.
foliata, Steud. Syn. Pl. Gram. 361 = latifolia.
foliosa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 291 = latifolia.
formosa, *R. Br. Prod.* 206.—Austral.
glabra, Roxb. Hort. Beng. 8; Fl. Ind. i. 353 = compressa.
glabrata, Benth. ex Hemsl. Biol. Centr. Am. Bot. iii. 521 = Andropogon imberbis.
glandulosa, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 250.—Burma; Malaya.
gracilis, Mill. ex Steud. Nom. ed. II. ii. 474 = Lepturus filiformis.
Hamiltoni, Trin. ex Steud. l. c. (Quid?).—Bengal.
hirsuta, *Vahl, Symb. Bot.* i. 11.—Afr. trop.; Arab.; Afghan.
hordeoides, *Munro, in Harv. Gen. S. Afr. Pl.* ed. II. 442.—Afr. austr.
impressa, *Griseb. Cat. Pl. Cub.* 235.—Cuba.
incurva, Roem. & Schult. Syst. ii. 799, in obs. = Lepturus incurvatus.
incurvata, Linn. f. Suppl. 114 = Lepturus incurvatus.
incurvata, Sibth. & Sm. Fl. Graec. i. 72 = Lepturus cylindricus.
laevis, Retz. Obs. iii. 11 = Ophiurus perforatus.
latifolia, *Steud. in Flora*, xxix. (1846) 21.—China; Japon.
loliacea, Bory & Chaub. Expéd. Mor. 46. t. 3. f. 2 = Lolium lepturoides.
loricata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 250.—Bras.; Cuba.
mollicoma, *Hance, in Journ. Bot.* ix. (1871) 134.—China.
monandra, Cav. Ic. i. 27. t. 39. f. 1 = Psilurus nardoides.
muricata, Benth. Fl. Austral. vii. 514 = ophiuroides.
muricata, *Retz. Obs.* iii. 12.—Ind. or.; Austral.
myurus, *Benth. in Journ. Linn. Soc.* xix. (1881) 68.—Ind. or.
nigrescens, *Thw. Enum. Pl. Zeyl.* 364.—Zeylan.

ROTTBOELLIA:—
ocreata, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 250 = Andropogon imberbis?
ophiuroides, *Benth. Fl. Austral.* vii. 514.—Austral.
paleacea, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 301 = Stenotaphrum glabrum.
paniculata, Salzm. ex Steud. Nom. ed. II. ii. 474 = Spartina bahiensis.
paniculata, Spreng. Syst. i. 300 = Schedonnardus texanus.
pannonica, Host, Gram. Austr. i. t. 24 = Lepturus pannonicus.
perforata, Roxb. Pl. Corom. ii. 43. t. 182 = Ophiurus perforatus.
pilosa, Roth, ex Roem. & Schult. Syst. ii. 785 (Enteropogonis sp.?).—Ind. or.
pilosa, Willd. Sp. Pl. i. 465 = Oropetium thomaeum.
pubescens, Bertol. Misc. Bot. i. 10 = Lepturus pubescens.
pulcherrima, Wall. ex Kunth, Enum. Pl. i. 468 = Ratzeburgia pulcherrima.
punctata, Retz. Obs. iii. 11 = Ophiurus corymbosus.
ramosa, Benth. ex Hemsl. Biol. Centr. Am. Bot. iii. 521 = aurita.
ramosa, Cav. in Anal. Cienc. Nat. iii. (1801) 11 = Lepturus cylindricus.
repens, Forst. f. Prod. 9 = Lepturus repens.
Rhytachne, *Hack. in Bol. Soc. Broter.* iii. (1885) 136.—Afr. trop.
Roxburghiana, Schult. Mant. ii. 430 = Lepturus pannonicus.
rugosa, *Nutt. Gen. Am.* i. 84.—Am. bor.
salina, Hort. ex Link, Enum. Hort. Berol. i. 101 = Lepturus pannonicus.
salsa, Fisch. ex Roem. & Schult. Syst. ii. 798 = Lepturus pannonicus.
Salzmanni, Trin. ex Steud. Syn. Pl. Gram. 361 = Andropogon imberbis?
Sandorii, Frivald. in Flora, xviii. (1835) 335 = digitata.
sanguinea, Retz. Obs. iii. 25 = Andropogon Pseudograya.
scorpioides, Poir. ex Steud. Nom. ed. II. ii. 474 = Ctenium carolinianum.
Selloana, *Hack. in Mart. Fl. Bras.* ii. III. 312.—Bras.
setacea, Roxb. Pl. Corom. ii. 17. t. 132 = Microchloa setacea.
setosa, *Meyen, Reise*, ii. 266, nomen.—Ins. Luzon.
setosa, J. & C. Presl, Rel. Haenk. i. 329 = exaltata.
spathacea, Tenore, Prod. Fl. Nap. p. xi = compressa.
stigmosa, *Trin. ex Rupr. in Bull. Acad. Brux.* ix. II. (1844) 243, nomen = aurita.
stipoides, Steud. Syn. Pl. Gram. 370 = Andropogon heteranthus.
stolonifera, Poir. Encyc. vi. 310 = Stenotaphrum glabrum.
striata, *Nees, ex Steud. Syn. Pl. Gram.* 361.—Ind. or.
stricta, Hort. ex Steud. Nom. ed. I. 705; ed. II. ii. 474 = Spartina stricta.
subulata, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 733 = Lepturus filiformis.
subulata, Savi, in Nuov. Giorn. Pisan. iv. (1806) 250 = Lepturus cylindricus.
tessellata, Steud. Syn. Pl. Gram. 362 = corrugata.
thomaea, Koen. in Naturf. xxiii. (1788) 210; Willd. Sp. Pl. i. 464 = Oropetium thomaeum.
Tranchelli, J. F. Gmel. Syst. 197 = compressa.
trichantha, Roth, Nov. Pl. Sp. 43 (Enteropogonis sp.?).—Ind. or.
tripsacoides, *Lam. Illustr.* i. 205.—Afr. trop.?
uncinata, Beauv. Agrost. 109; Spreng. Syst. i. 299 = compressa.
uniflora, A. Cunn. in Hook. Comp. Bot. Mag. ii. (1836) 371 = Zoysia pungens.
villosa, *Poir. Encyc.* vi. 313.—Ind. or.
Xerochloa, Spreng. Syst. i. 300 = Xerochloa imberbis.

ROTTBOLLA, Zumag. Fl. Pedem. i. 93 (1849), sphalm. = praec.

ROTTBOLLIA, Cav. Ic. i. 27 (1791) = praec.

ROTTLERA, Roxb. Pl. Corom. ii. 36. t. 168 (1798) = **Mallotus**, Lour. (Euphorb.).
acuminata, Baill. Etud. Gén. Euph. 426 = M. acuminatus.
acuminata, A. Juss. Tent. Euph. 33 = M. tiliaefolius.

ROTTLERA:—
acutifolia, Hassk. Cat. Hort. Bog. Alt. 238 = Mallotus acuminatus.
affinis, Baill. Etud. Gén. Euph. 424 = M. claoxyloides.
affinis, Hassk. in Flora, xxv. (1842) II. Beibl. 41 = M. philippinensis.
alba, Roxb. Hort. Beng. 73; Fl. Ind. iii. 829 = M. albus.
albicans, Hassk. Cat. Hort. Bog. Alt. 238 = Cephalocroton discolor.
albicans, Mor. ex Reichb. f. & Zoll. in Verh. Natuurk. Ver. Nederl. Indie, i. (1856) 21 = Cephalocroton albicans.
angulata, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. angulatus.
aurantiaca, Hook. & Arn. Bot. Beech. Voy. 270 = M. philippinensis.
aureopunctata, Dalz. in Hook. Kew Journ. iii. (1851) 122 = M. Lawii.
barbata, Wall. Cat. n. 7822 = M. barbatus.
bifaria, Wall. l. c. n. 7981 = Gelonium multiflorum.
Blumeana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. Blumeanus.
Blumei, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 486 = M. tiliaefolius.
borneensis, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. borneensis.
brasiliensis, Spreng. Syst. iii. 877 = Croton brasiliensis.
calcosa, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. calcosus.
cantoniensis, Spreng. Syst. iii. 878 = M. Apelta.
chinensis, A. Juss. Tent. Euph. 33 = M. Apelta.
congesta, Wall. Cat. n. 7981, B = Gelonium multiflorum.
cordifolia, Benth. Fl. Hongk. 307 = M. repandus.
dentata, Baill. Adansonia, i. (1860–61) 69 = M. oppositifolius.
diadena, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. diadenus.
dicocca, Roxb. Hort. Beng. 73; Fl. Ind. iii. 829 = M. repandus.
digyna, Thw. Enum. Pl. Zeyl. 273 = Macaranga digyna.
dioica, Baill. Etud. Gén. Euph. 423 = M. repandus.
dioica, Roxb. ex Wall. Cat. n. 7827, A = *dicocca*.
discolor, F. Muell. ex Benth. Fl. Austral. vi. 143 = M. discolor.
dispar, Blume, Bijdr. 608 = M. dispar.
dispar, Mor. Syst. Verz. Zoll. 17 = M. Moritzianus.
eglandulosa, Zipp. ex Span. in Linnaea, xv. (1841) 348, nomen (Quid?).—Ins. Timor.
eriocarpa, Thw. Enum. Pl. Zeyl. 273 = M. eriocarpus.
fasciculata, Wall. Cat. n. 7981, B = Gelonium multiflorum.
ferruginea, Roxb. Hort. Beng. 73; Fl. Ind. iii. 828 = M. ferrugineus.
flavigutta, Miq. Fl. Ind. Bat. Suppl. 453 = M. acuminatus.
floribunda, Hassk. Cat. Hort. Bog. Alt. 238 = M. floribundus.
Furetiana, Baill. Etud. Gén. Euph. 426, nomen = M. Furetianus.
fuscescens, Thw. Enum. Pl. Zeyl. 273 = M. fuscescens.
gigantea, Reichb. f. & Zoll. ex Kurz, in Tijdschr. Nederl. Indie, xxvii. (1864) 44 = Macaranga gigantea.
glaberrima, Hassk. in Flora, xxv. (1842) II. Beibl. 41 = M. glaberrimus.
glauca, Hassk. l. c. = Macaranga denticulata.
hexandra, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 829 = Macaranga hexandra.
hispida, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxiv. (1864) 43 = Macaranga hispida.
Hookeriana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. Hookerianus.
Inophyllum, Endl. in Ann. Wien. Mus. i. (1836) 183. t. 75 = Codiaeum Inophyllum.
integrifolia, Wall. Cat. n. 7810, F = Macaranga Tanarius.
japonica, Spreng. Syst. iii. 878 = M. japonicus (cf. Addend.).
javanica, Hassk. in Flora, xxv. (1842) II. Beibl. 41 = Macaranga javanica.
Korthalsii, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. Korthalsii.
laccifera, Voigt, Hort. Suburb. Calc. 157 = M. repandus.

ROTTLERA :—
Lambertiana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = Mallotus Lambertianus.
lanceolata, Wall. Cat. n. 7981 = Gelonium multiflorum.
lappacea, Wall. l. c. n. 7845 = Coelodiscus lappaceus.
longestipulata, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 43 = Macaranga longestipulata.
longifolia, Reichb. f. & Zoll. in Verh. Natuurk. Ver. Nederl. Indie, i. (1856) 31 = M. longifolius.
macrostachya, Miq. Fl. Ind. Bat. Suppl. 454 = M. macrostachyus.
mappoides, Dalz. in Hook. Kew Journ. iii. (1851) 122 = M. albus.
Miqueliana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124.—Borneo.
mollis, Wall. Cat. n. 7839 = M. philippinensis.
mollissima, Wall. l. c. n. 7825 = Alchornea mollis.
moluccana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 122 = M. moluccanus.
montana, Heyne, ex Wall. Cat. n. 7833 = Macaranga javanica.
montana, Wall. Cat. n. 7833 = M. philippinensis.
Moritziana, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. Moritzianus.
multiglandulosa, Blume, Bijdr. 609 = M. moluccanus.
muricata, Thw. Enum. Pl. Zeyl. 273 = M. muricatus.
oblongifolia, Miq. Fl. Ind. Bat. i. II. 396 = M. oblongifolius.
oppositifolia, Blume, Bijdr. 608 = M. Blumeanus.
oppositifolia, Thw. Enum. Pl. Zeyl. 273 = M. rhamnifolius.
paniculata, A. Juss. Tent. Euph. 33 = M. cochinchinensis.
paniculata, Wall. Cat. n. 7812, 7818 = M. albus, repandus.
peltata, Miq. Fl. Ind. Bat. i. II. 395 = M. ricinoides.
peltata, Roxb. Hort. Beng. 73; Fl. Ind. iii. 828 = M. Roxburghianus.
peltata, Wight, Ic. t. 1873 = M. albus.
philippinensis, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. philippinensis.
pycnostachys, F. Muell. Fragm. iv. 138 = M. ricinoides.
repanda, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 124 = M. repandus.
rhombifolia, Thw. Enum. Pl. Zeyl. 272 = M. repandus.
rhynchophylla, Miq. Fl. Ind. Bat. Suppl. 454 = M. subpeltatus.
ricinoides, A. Juss. Tent. Euph. 33 = M. ricinoides.
rufidula, Miq. Fl. Ind. Bat. Suppl. 453 = M. rufidulus.
scabrifolia, A. Juss. Tent. Euph. 111 = M. repandus.
scandens, Span. in Linnaea, xv. (1841) 348 = M. scandens.
Schimperi, Hochst. & Steud. in Flora, xxiv. (1841) I. Intell. 21 = Croton macrostachys.
spathulata, Panch. ex Baill. Adansonia, ii. (1861–62) 222 = Cleidion verticillatum.
sphaerocarpa, Miq. Fl. Ind. Bat. Suppl. 454 = M. sphaerocarpus.
stipulosa, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. stipulosus.
stylaris, Scheff. in Miq. l. c. = M. stylaris.
subfalcata, Reichb. f. & Zoll. in Verh. Natuurk. Ver. Nederl. Indie, i. (1856) 11 = Macaranga subfalcata.
subpeltata, Baill. Etud. Gén. Euph. 423 = M. subpeltatus.
suffruticosa, Buch.-Ham. ex Wall. Cat. n. 7843 = Baliospermum axillare.
Tanarius, Hassk. Cat. Hort. Bog. Alt. 238 = Macaranga Tanarius.
tetracocca, Roxb. Hort. Beng. 73; Fl. Ind. iii. 826 = M. albus.
Thwaitesii, Baill. Etud. Gén. Euph. 426 = Podadenia sapida.
tiliaefolia, Blume, Bijdr. 607 = M. tiliaefolius.
tinctoria, Roxb. Pl. Corom. ii. 36. t. 168 = M. philippinensis.
Tokbrai, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 122 = M. Tokbrai.
tomentosa, Hassk. Cat. Hort. Bog. Alt. 238 = Macaranga Tanarius.
trinervis, Zipp. ex Span. in Linnaea, xv. (1841) 348 = M. repandus.
urandra, Dalz. in Hook. Kew Journ. iii. (1851) 229 = Cleidion javanicum.

ROTTLERA :—
viscida, Blume, Bijdr. 608 = Mallotus repandus.
Zippelii, Hassk. Cat. Hort. Bog. Alt. 238 = M. ricinoides.
Zollingeri, Scheff. in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 125 = M. Zollingeri.
Zollingeriana, Miq. Fl. Ind. Bat. i. II. 394 (Xylosmatis sp. ?).—Java.

ROTTLERA, Vahl, Enum. i. 88 (1805) (Roettlera) = **Didymocarpus**, Wall. (Gesner.).
incana, Vahl, l. c. = D. Rottleriana.

ROTTLERA, Willd. in Goett. Journ. Wiss. i. (1797) 7. t. 1 = **Trewia**, Linn. (Euphorb.).
Hoperiana, Blume, ex Muell. Arg. in DC. Prod. xv. II. 953 = T. nudiflora.
indica, Willd. in Goett. Journ. Wiss. i. (1797) 8 = T. nudiflora.
operiana, Blume, ex Baill. Etud. Gen. Euph. 423 = T. nudiflora.
rusciflora, Spreng. Syst. iii. 878 = T. rusciflora.

ROTULA, Lour. Fl. Cochinch. 121 (1790) = **Rhabdia**, Mart. (Borag.).
aquatica, Lour. l. c. = Rhabdia lycioides.

ROUBIEVA, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 292. t. 10. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 52.
anthelmintica, Hook. & Arn. Bot. Beech. Voy. 387 = Chenopodium ambrosioides.
bonariensis, *Hook. f. in Benth. & Hook. f. Gen.* iii. 52.—Am. austr.
Botrys, Fuss, Fl. Trans. 552 = Chenopodium Botrys.
multifida, *Moq. in Ann. Sc. Nat.* Sér. II. i. (1834) 292. t. 10.—Am. austr.

ROUCELA, Dum. Comm. Bot. 14 (1822) = **Campanula**, Tourn.
Erinus, Dum. l. c. = C. Erinus.
hederacea, Dum. Fl. Belg. 58 = Wahlenbergia hederacea.

ROUCHERIA, Planch. in Hook. Lond. Journ. Bot. vi. (1847) 141. t. 2. *LINEAE*, Benth. & Hook. f. i. 243.
calophylla, *Planch. l. c.*—Guian.
Griffithiana, *Planch. l. c.* 143.—Ins. Molucc.
humiriifolia, Planch. l. c. = Hebepetalum humiriifolium.
latifolia, Spruce, ex Benth. & Hook. f. Gen. i. 245 = Hebepetalum latifolium.
macrophylla, *Miq. Fl. Ind. Bat.* i. II. 136.—Sumatra.
Schomburgkii, *Planch. in Hook. Lond. Journ. Bot.* vii. (1848) 526.—Guian.

ROUHAMON, Aubl. Pl. Gui. i. 93. t. 36 (1775) = **Strychnos**, Linn. (Logan.).
Bredemeyeri, DC. Prod. ix. 18.—Venezuela.
brevifolium, Spruce, ex Prog. in Mart. Fl. Bras. vi. I. 277 = S. brevifolia.
Curare, DC. Prod. ix. 17 = S. Curare.
divaricatum, DC. l. c. = S. Rouhamon.
glabrum, Sagot, ex Prog. in Mart. Fl. Bras. vi. I. 275 = S. glabra.
guianensis, Aubl. Pl. Gui. i. 93 = S. Rouhamon.
Medeola, Sagot, ex Prog. in Mart. Fl. Bras. vi. I. 282 = S. Medeola.
pedunculatum, A. DC. ex DC. Prod. ix. 561 = S. pedunculata.
stans, DC. Prod. ix. 18.—Bras.

ROULINIA, Brongn. in Ann. Sc. Nat. Sér. II. xiv. (1840) 320 = **Nolina**, Michx. (Liliac.).
acrotricha, Brongn. l. c = Dasylirion acrotrichum.
gracilis, Brongn. l. c. = Dasylirion acrotrichum.
Humboldtiana, Brongn. l. c. = N. parviflora.
Karwinskiana, Brongn. l. c. = N. longifolia.
longifolia, Brongn. l. c. = N. Hartwegiana.
pitcairniaefolia, Nuytens, ex Versch. Cat. n. 21 (1879–80), cum ic.—Hab.?
serratifolia, Brongn. in Ann. Sc. Nat. Sér. II. xiv. (1840) 320 = Dasylirion serratifolium.

ROULINIA, Decne. in DC. Prod. viii. 516 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 762.
acuta, *Decne. l. c.* 517.—Bras.
barbata, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) I. 253.—Bras.

ROULINIA:—
blanda, *Decne. in DC. Prod.* viii. 517.—Guian.
connivens, *Decne. l. c.* 518.—Bras.
convolvulacea, *Decne. l. c.* 517.—Bras.
corymbosa, *Decne. l. c.*—Bras.
fluminensis, *Decne. l. c.*—Reg. Argent.
guayanensis, *Decne. l. c.*—Guian.
Jacquinii, *Decne. l. c.*—Mexic.
jamaicensis, *Benth. & Hook. f. Gen.* ii. 757.—Ind. occ.
modesta, *Decne. in DC. Prod.* viii. 517.—Bras.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 115.—N. Mexic.
parviflora, *Decne. in DC. Prod.* viii. 518.—Bras.
Riedelii, *Fourn. in Mart. Fl. Bras.* vi. IV. 217.—Bras.
Selloana, *Fourn. l. c.* 216.—Bras.
Sprucei, *Fourn. l. c.*—Bras.
surinamensis, *Miq. Stirp. Surin. Sel.* 169.—Guian.
tamifolium, *Decne. in DC. Prod.* viii. 517.—Bras.
unifaria, *Engelm. ex Torr. Bot. Mex. Bound.* 160.—Texas.
versicolor, Herb. Gen. Amat. 4. t. 26; Decne. in DC. Prod. viii. 517 = tamifolium?

ROUMEA, DC. Prod. i. 256 (1824), sphalm = RUMEA, Poit. = **Xylosma**, Forst. (Bixin.).

ROUPALA, Aubl. Pl. Gui. i. 83. t. 32 (1775). *PROTEACEAE*, Benth. & Hook. f. iii. 178.
DICKNECKERIA, Vell. Fl. Flum. i. t. 105 (1827).
LEINKERIA, Scop. Introd. 345 (1777).
RHOPALA, Schreb. Gen. 62 (1789).
ROPALA, J. F. Gmel. Syst. ii. 225 (1791).
RUPALA, Vahl, Symb. Bot. iii. 20 (1794).
adiantifolia, *Klotzsch, in Linnaea*, xv. (1841) 56.—Bras.
affinis, *Pohl, Pl. Bras. Ic.* i. 112.—Bras.
attenuata, Jack, in Malay. Misc. i. (1820) n. v. 10 = Helicia attenuata.
aurea, *Linden, in Belg. Hortic.* xvi. (1866) 202.—Bras.
australis, *Linden, Cat.* n. 14 (1859) 8.—Bras.
Bleasdalii, *F. Muell. in Proc. Roy. Soc. N. S. Wales*, (1881) 28.—Austral.
Boissieriana, *Meissn. in DC. Prod.* xiv. 430.—N. Granat.
borealis, *Hemsl. Biol. Centr. Am. Bot.* iii. 78.—Mexic.
brasiliensis, *Klotzsch, in Linnaea*, xv. (1841) 55.—Bras.
chrysogenia, Mart. ex Meissn. in Mart. Fl. Bras. v. I. 93 (= *Adenostephanus incana*).—Bras.
cochinchinensis, R. Br. in Trans. Linn. Soc. x. (1810) 192 = Helicia cochinchinensis.
complicata, *H. B. & K. Nov. Gen. et Sp.* ii. 153.—N. Granat.
complicata, *Linden, Cat.* n. 8 (1853) 3; *et ex Schlecht. in Bot. Zeit.* xi. (1853) 717.—Am. trop.
corcovadensis, Hort. ex Meissn. in DC. Prod. xiv. 433 = Pohlii.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 152.—N. Granat.
dentata, *R. Br. in Trans. Linn. Soc.* x. (1810) 192.—Guian.
dentigera, *Knight, Prot.* 103.—Ins. Trinit.
Dickneckeria, Steud. Nom. ed. II. ii. 487 (= *Adenostephanus Sellowii*).—Bras.
diversifolia, *R. Br. in Trans. Linn. Soc.* x. (1810) 193.—Peruv.
diversifolia, Schott, in Wien. Med. Jahrb. vi. n. 11; ex Schult. Mant. iii. 287 = heterophylla.
diversifolia, Zucc. ex Meissn. in DC. Prod. xiv. 433 = Pohlii.
elegans, *Pohl, Pl. Bras. Ic.* i. 112.—Bras.
excelsa, Roxb. Hort. Beng. 83; Fl. Ind. i. 362 = Helicia excelsa.
ferruginea, *H. B. & K. Nov. et Sp.* ii. 152.—N. Granat.
frigida, Linden, Cat. n. 18 (1863), 10 (Quid?).—Hab.?
frondosa, Rich. ex Meissn. in DC. Prod. xiv. 427 = montana.
Gardneri, *Meissn. in Mart. Fl. Bras.* v. I. 83. t. 31. f. 1.—Bras.
glabrata, *Klotzsch, ex Meissn. l. c.* 84.—Bras.
glabrata, Wall. Cat. n. 1039 = Helicia robusta.
glaucophylla, *Linden, Cat.* n. 13 (1858); n. 14, (1859) 8.—Bras.

ROUPALA :—
gracilis, *Meissn. in Mart. Fl. Bras.* v. I. 90.—Bras.
granatensis, *Linden, Cat.* n. 24 (1869).—N. Granat.
grandifolia, *Klotzsch, in Linnaea,* xv. (1841) 56.—Bras.
hameliaefolia, Rudge, Pl. Guian. i. 22. t. 31 = Panopsis hameliaefolia.
heterophylla, Herb. Monac. ex Meissn. in DC. Prod. xiv. 433 = Pohlii.
heterophylla, *Pohl, Pl. Bras. Ic.* i. 111. t. 90.—Bras.
inaequalis, Pohl, l. c. 110. t. 89 (= *Adenostephanus inaequalis*).—Bras.
javanica, Steud. Nom. ed. II. ii. 487 = Helicia robusta.
laurifolia, Knight, Prot. 102 = nitida.
longepetiolata, Klotzsch, ex Meissn. in Mart. Fl. Bras. v. I. 85 = rhombifolia.
longepetiolata, *Pohl, Pl. Bras. Ic.* i. 109. t. 88.—Bras.
loranthoides, *Meissn. in DC. Prod.* xiv. 425.—Guatemala.
lucens, *Meissn. in Mart. Fl. Bras.* v. I. 84. t. 31. f. 2.—Bras.
macrophylla, *Pohl, Pl. Bras. Ic.* i. 112.—Bras.
macropoda, *Klotzsch, in Linnaea,* xx. (1847) 474.—N. Granat.
Martii, *Meissn. in Mart. Fl. Bras.* v. I. 87. t. 32. f. 3.—Bras.
media, *R. Br. in Trans. Linn. Soc.* x. (1810) 191.—Guian.
moluccana, R. Br. l. c. = Helicia moluccana.
moluccana, Jack, in Malay. Misc. i. (1820) n. v. 10 = Helicia petiolaris.
moluccana, Roxb. Hort. Beng. 83; Fl. Ind. i. 361 = Helicia attenuata.
montana, *Aubl. Pl. Gui.* i. 83.—Guian.
montana, Willd. Sp. Pl. i. 536 = media.
multiflora, Schott, in Spreng. Syst. iv. Add. 403 (= *Andriapetalum reticulatum*).—Bras.
myrtoidea, Poepp. & Endl. Nov. Gen. et Sp. ii. 35. t. 149 = Orites myrtoidea.
nervosa, *Klotzsch, in Linnaea,* xv. (1841) 56.—Bras.
nitida, *Rudge, Pl. Guian.* i. 26. t. 39.—Guian.
obovata, *H. B. & K. Nov. Gen. et Sp.* ii. 153.—N. Granat.
obtusata, *Klotzsch, in Linnaea,* xv. (1841) 54.—Bras.
ochrantha, Mart. ex Meissn. in Mart. Fl. Bras. v. I. 86 = brasiliensis.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 135 (= *Adenostephanus organensis*).—Bras.
ovalis, Klotzsch, in Linnaea, xv. (1841) 54 = tomentosa.
ovalis, *Pohl, Pl. Bras. Ic.* i. 107. t. 86.—Bras.
ovata, Jack, in Malay. Misc. ii. (1827) n. VII. 95 = Helicia ovata.
pamplonensis, Linden, Cat. n. 8 (1853), 3, in desc. = Boissieriana.
peruviana, *R. Br. in Trans. Linn. Soc.* x. (1810) 192.—Peruv.
pinnata, Lam. Illustr. i. 243 = Euplassa meridionalis.
Pohlii, *Meissn. in Mart. Fl. Bras.* v. I. 89. t. 33.—Bras.
polystachya, H. B. & K. Nov. Gen. et Sp. ii. 154 (= *Andriapetalum polystachyum*).—N. Granat.
Princeps, *Linden, in Belg. Hortic.* vii. (1857) 198.—N. Granat.
pyrifolia, *Knight, Prot.* 102.—Guian.
racemosa, Roxb. ex Meissn. in DC. Prod. xiv. 439 = Helicia attenuata.
rhombifolia, *Mart. ex Meissn. in Mart. Fl. Bras.* v. I. 85.—Bras.
robusta, Roxb. Hort. Beng. 83; Fl. Ind. i. 363 = Helicia robusta.
robusta, Spreng. Syst. i. 482, partim = Helicia cochinchinensis.
Rousselii, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 394.—N. Caled.
Sanctae-Catharinae, *Linden, Cat.* n. 17 (1862), 8.—Bras.
Schomburgkii, *Klotzsch, in Linnaea,* xx. (1847) 474.—Bras.
serrata, R. Br. in Trans. Linn. Soc. x. (1810) 192 = Helicia serrata.
serrata, Reinw. ex Blume, Cat. Gew. Buitenz. 42 = Helicia robusta.
serratifolia, *Linden, in Belg. Hortic.* xvi. (1866) 202.—Bras.

ROUPALA :—
sessilifolia, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Panopsis hameliaefolia.
silaifolia, Linden, Cat. n. 18 (1863), 10 (Quid?).—Hab. ?
suaveolens, *Klotzsch, in Linnaea,* xx. (1847) 473.—Guian.
Thomesiana, *Moric. Pl. Nouv. Am.* 172. t. 100.—Bras.
tomentosa, *Pohl, Pl. Bras. Ic.* i. 108. t. 87.—Bras.
velutina, Klotzsch, in Linnaea, xv. (1841) 54 = brasiliensis.
veraguensis, Klotzsch, ex Meissn. in Mart. Fl. Bras. v. I. 87 = Martii.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 229.—N. Caled.
Wallichii, Steud. Nom. ed. II. ii. 487 = Helicia robusta.
Warmingii, *Meissn. in Kjoeb. Vidensk. Meddel.* (1870) 148.—Bras.

ROUPALIA, T. Moore & Ayres, Gard. Mag. Bot. ii. (1850) t. ad p. 33, sphalm. = **Roupellia**, Wall. & Hook. (Apocyn.).
grata, T. Moore & Ayres, l. c. = Roupellia grata.

ROUPALLIA, Hassk. in Flora, xl. (1857) 103 = seq.

ROUPELLIA, Wall. & Hook. ex Benth. in Hook. Bot. Mag. t. 4466 (1849). *APOCYNACEAE*, Benth. & Hook. f. ii. 713.
grata, *Wall. & Hook. l. c.*; *et in Hook. Niger Fl.* 449.—Afr. trop.

ROUREA, Aubl. Pl. Gui. i. 467. t. 187 (1775). *CONNARACEAE*, Benth. & Hook. f. i. 432.
MALBRANCIA, Neck. Elem. ii. 366 (1790).
ROBERGIA, Schreb. Gen. 309 (1789).
acuminata, *Hook. f. Fl. Brit. Ind.* ii. 48.—Singapore.
acutipetala, *Miq. Fl. Ind. Bat. Suppl.* 528.—Java.
Afzelii, Planch. in Linnaea, xxiii. (1850) 418 = santaloides.
Balansaeana, *Baill. Adansonia,* xi. (1875) 309.—N. Caled.
Bowiniana, *Baill. Adansonia,* vii. (1866–67) 231.—Afr. trop.
brachyandra, *F. Muell. Fragm.* viii. 6.—Austral.
caudata, *Planch. in Linnaea,* xxiii. (1850) 419.—Reg. Himal.
coccinea, Hook. f. ex Hook. Niger Fl. 290 = Byrsocarpus coccineus.
commutata, *Planch. in Linnaea,* xxiii. (1850) 420.—Ind. or.; China.
concolor, *Blume, Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
cuspidata, *Benth. ex Baker, in Mart. Fl. Bras.* xiv. II. 181.—Bras.
dasyphylla, Miq. Fl. Ind. Bat. Suppl. 528 = Cnestis ramiflora.
discolor, *Baker, in Mart. Fl. Bras.* xiv. II. 180.—Bras.
diversifolia, *Miq. Fl. Ind. Bat. Suppl.* 528.—Sumatra.
Doniana, *Baker, in Mart. Fl. Bras.* xiv. II. 179.—Bras.
florida, *Miq. Fl. Ind. Bat. Suppl.* 206.—Sumatra.
fluminensis, Baill. Hist. Pl. ii. 5 = Bernardinia fluminensis.
fraterna, Planch. in Linnaea, xxiii. (1850) 416 = induta.
frutescens, *Aubl. Pl. Gui.* i. 467.—Guian.
fulgens, *Planch. in Linnaea,* xxiii. (1850) 423.—Singapore.
Gardneriana, *Planch. l. c.* 417.—Bras.
glabra, *H. B. & K. Nov. Gen. et Sp.* vii. 41.—Venezuela.
Grosourdyana, *Baill. Adansonia,* ix. (1868–70) 149.—Venezuela.
heterophylla, *Baker, in Oliver, Fl. Trop. Afr.* i. 456.—Afr. trop.
heterophylla, *Planch. in Linnaea,* xxiii. (1850) 419.—Ins. Philipp.
humilis, *Blume, Mus. Bot. Lugd. Bat.* i. 262.—Borneo.
induta, *Planch. in Linnaea,* xxiii. (1850) 417.—Bras.
javanica, *Blume, Mus. Bot. Lugd. Bat.* i. 262.—Java.
ligulata, *Baker, in Mart. Fl. Bras.* xiv. II. 181.—Bras.
lucida, *Planch. in Linnaea,* xxiii. (1850) 423.—Java.

ROUREA :—
macrophylla, *Baker, in Mart. Fl. Bras.* xiv. II. 183.—Bras.
Martiana, *Baker, l. c.* 178.—Bras.
microphylla, *Planch. in Linnaea,* xxiii. (1850) 421.—China.
Millettii, *Planch. l. c.* 420 (Milletii).—China.
mimosoides, *Planch. l. c.*—Ins. Nicobar.
multiflora, *Planch. l. c.* 418.—Ins. Philipp.
myriantha, *Baill. Adansonia,* viii. (1867–68) 198.—Afr. trop.
nitida, *Planch. in Linnaea,* xxiii. (1850) 423.—Java.
oblongifolia, Hook. & Arn. Bot. Beech. Voy. 283 = glabra.
orientalis, Baill. Adansonia, vii. (1866–67) 230 = Byrsocarpus orientalis.
Palisotii, *Baill. l. c.* 233, in textu.—Afr. trop.
parallela, Planch. in Linnaea, xxiii. (1850) 421 = similis.
parvifolia, Planch. l. c. 420 = concolor.
paucifoliata, Planch. l. c. 415 = glabra.
Pervilleana, *Baill. Adansonia,* vii. (1866–67) 232.—Ins. Mascar.
phyllanthoides, *Blume, Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
platysepala, *Baker, in Journ. Linn. Soc.* xxi. (1884) 336.—Madag.
polyphylla, *Blume, Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
puberula, *Baker, in Mart. Fl. Bras.* xiv. II. 179.—Bras.
pulchella, *Planch. in Linnaea,* xxiii. (1850) 419.—Malacca.
reticulata, Planch. l. c. 416 = induta.
revoluta, *Planch. l. c.* 415.—Guian.
rugosa, *Planch. l. c.* 422.—Malaya.
santaloides, *Wight & Arn. Prod.* 144.—Ind. or.
sclerocarpa, Wight & Arn. ex Hook. f. Fl. Brit. Ind. ii. 51 = Connarus Wightii.
similis, *Blume, Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
simplicifolia, *Blume, l. c.* 263.—Sumatra.
Solanderi, *Baker, in Oliver, Fl. Trop. Afr.* i. 456.—Afr. trop.
Sookurthoontee, Voigt, Hort. Suburb. Calc. 265 = commutata.
sororia, Planch. in Linnaea, xxiii. (1850) 421 = similis.
stenopetala, *Hook. f. Fl. Brit. Ind.* ii. 49.—Burma.
surinamensis, *Miq. in Linnaea,* xxvi. (1853) 221.—Guian.
villosa, *Planch. l. c.* xxiii. (1850) 422.—Burma; Sumatra.
Wallichiana, *Planch. l. c.* 421.—Burma.

ROUREOPSIS, Planch. in Linnaea, xxiii. (1850) 423. *CONNARACEAE* [Benth. & Hook. f. i. 432].
javanica, *Planch. l. c.*—Java.
pubinervis, *Planch. l. c.*—Malaya.

ROUSSAEA, DC. Prod. vii. 522 (1839) = seq.

ROUSSEA, Sm. Ic. Ined. i. 6. t. 6 (1789). *SAXIFRAGEAE*, Benth. & Hook. f. i. 645.
ROUSSAEA, DC. Prod. vii. 522 (1839).
ROUSSEAUVIA, Boj. Hort. Maurit. 246 (1837).
ROUSSOA, Roem. & Schult. Syst. iii. 3, 92 (1818).
salicifolia, Spreng. Syst. i. 419 = Lucuma Sellowii.
simplex, *Sm. Ic. Ined.* i. 6. t. 6.—Ins. Mascar.

ROUSSEAUVIA, Boj. Hort. Maurit. 246 (1837) = praec.

ROUSSEAUXIA, DC. Prod. iii. 152 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 753.
articulata, DC. l. c. = Dichaetanthera articulata.
chrysophylla, *DC. l. c.* 153.—Madag.

ROUSSELIA, Gaudich. in Freyc. Voy. Bot. 503 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 393.
LITHOCNIDE, Rafin. Fl. Tellur. iii. 48 (1836).
lappulacea, *Gaudich. in Freyc. Voy. Bot.* 503.—Ind. occ.

ROUSSINIA, Gaudich. Bot. Voy. Bonite, t. 21 (1844–52) = **Pandanus**, Linn.
indica, Gaudich. l. c. t. 21. ff. 1–4 = P. Leram.
indica, Gaudich. l. c. t. 21. ff. 5–9 = P. palustris.

ROUSSOA, Roem. & Schult. Syst. iii. 3, 92 (1818) = **Roussea**, Sm. (Saxifr.).

ROXBURGHIA, [Banks, in] Roxb. Pl. Corom. i. 29. t. 32 (1795) = **Stemona**, Lour. (Roxburgh.).
gloriosa, Pers. Syn. i. 412 = S. tuberosa.
gloriosoides, Roxb. Pl. Corom. i. 26. t. 32 = S. tuberosa.
gloriosoides, Zoll. ex Kunth, Enum. Pl. v. 288 = *javanica*.
japonica, Blume, Enum. Pl. Jav. i. 9 = S. japonica.
japonica, Miq. ex Franch. & Sav. Enum. Pl. Jap. ii. 92, in syn. = S. sessilifolia.
javanica, Kunth, Enum. Pl. v. 288.—Malaya; Austral.
moluccana, Blume, Enum. Pl. Jav. i. 9.—Ins. Molucc.
rostrata, Russ. Herb. ex Nees, in DC. Prod. xi. 57, 58 = Thunbergia fragrans.
ruscifolia, Zucc. ex Schnizl. Iconog. i. t. 56[b]. f. 17.—Hab. ?
sessilifolia, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 211 = S. sessilifolia.
Stemona, Steud. Nom. ed. II. ii. 475 = S. tuberosa.
viridiflora, Sm. Exot. Bot. i. 111 = S. tuberosa.

ROXBURGHIA, Koen. ex Roxb. Fl. Ind. ed. Carey, i. 168 (1820) = **Olax**, Schreb.
baccata, Koen. ex Roxb. l. c. = O. scandens.

ROYDSIA, Roxb. Hort. Beng. 42 (1814); Pl. Corom. iii. 86. t. 289 (1819); Fl. Ind. ii. 643 (1832). *CAPPARIDEAE*, Benth. & Hook. f. i. 110.
STIXIS, Lour. Fl. Cochinch. 295 (1790), nomen prius.
floribunda, Planch. ex Mast. in Hook. f. Fl. Brit. Ind. i. 409 = philippinensis.
obtusifolia, *Hook. f. & Thoms. in Hook. f. l. c.* 180.—Burma.
parviflora, *Griff. Notul.* iv. 578.—Burma.
philippinensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 329.—Ins. Philipp.
suaveolens, *Roxb. Hort. Beng.* 42; *Pl. Corom.* iii. 86. t. 289; *Fl. Ind.* ii. 643.—Reg. Himal.

ROYENA, Linn. Gen. ed. I. 114 (1737). *EBENACEAE*, Benth. & Hook. f. ii. 663.
ambigua, Salisb. Prod. 285 (Quid ?).—Afr. austr.
ambigua, *Vent. Jard. Malm.* 17.—Afr. austr.
angustifolia, *Salisb. Prod.* 285.—Hab. ?
angustifolia, Willd. Sp. Pl. ii. 633 = hirsuta.
brachiata, E. Mey. in Drège, Cat. Pl. Exs. Afr. Austr. 7, nomen; et ex A. DC. in DC. Prod. viii. 213 = pallens.
cistoides, *Welw. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 87.—Afr. trop.
cordata, *E. Mey. in Drège, Cat. Pl. Exs. Afr. Austr.* 7, nomen; *et ex A. DC. in DC. Prod.* viii. 211.—Afr. austr.
corylifolia, Salisb. Prod. 284 = villosa.
crassifolia, *Salisb. l. c.* 285.—Afr. austr.
cuneata, Poir. Encyc. vi. 322 = hirsuta ?
cuneata, Spreng. Syst. ii. 360 = pallens.
cuneifolia, E. Mey. in Drège, Cat. Pl. Exs. Afr. Austr. 7, nomen; et ex A. DC. in DC. Prod. viii. 214 = pallens.
decidua, Burch. Trav. i. 317 = pallens.
falcata, E. Mey. in Drège, Zwei Pfl. Docum. 217 = glabra.
glabra, *Linn. Sp. Pl.* 397.—Afr. austr.
glandulosa, *Harv. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 89.—Afr. austr.
hirsuta, Jacq. Coll. v. 110. t. 13. f. 1 = pallens.
hirsuta, *Linn. Sp. Pl.* 397.—Afr. austr.
hirsuta, Sieber, ex A. DC. in DC. Prod. viii. 214 = glabra.
hispidula, Harv. ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 85 = pallens.
latifolia, *Willd. Enum. Hort. Berol. Suppl.* 23, nomen. —Hab. ?
lucida, *Linn. Sp. Pl.* 397.—Afr. austr.
lycioides, Desf. Tabl. ed. I. 79, nomen; A. DC. in DC. Prod. viii. 214 = pallens.
macrophylla, E. Mey. ex A. DC. l. c. 218 = Euclea natalensis.
media, *Hort. ex Steud. Nom.* ed. II. ii. 475, nomen.—Afr. austr.
microphylla, Burch. Trav. i. 348 = hirsuta.
myrtifolia, Salisb. Prod. 285 = glabra.
nitens, *Harv. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 87.—Afr. austr.
oleifolia, Desf. ex Hiern, l. c. 85 = pallens.
opaca, E. Mey. in Drège, Zwei Pfl. Docum. 217 = cordata.

ROYENA:—
ovalifolia, Salisb. Prod. 285 = Euclea polyandra.
pallens, *Thunb. Prod. Pl. Cap.* 80.—Afr. austr.
parviflora, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 88.—Afr. austr.
polyandra, Linn. f. Suppl. 240 = Euclea polyandra.
pubescens, Willd. Enum. Hort. Berol. 457 = pallens.
ramulosa, E. Mey. ex A. DC. in DC. Prod. viii. 212 = pallens.
rufescens, E. Mey. in Drège, Zwei Pfl. Docum. 217 = Euclea ovata.
rugosa, E. Mey. in Drège, Cat. Pl. Exs. Afr. Austr. 7, nomen; et ex A. DC. in DC. Prod. viii. 212 = hirsuta.
scabra, Burm. f. Fl. Cap. Prod. 13 = villosa.
scabrida, *Harv. ex Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 82.—Afr. austr.
scandens, Burch. ex Hiern, l. c. = villosa.
sericea, Bernh. in Flora, xxvii. (1844) 824 = pallens.
sessilifolia, *Hiern, in Trans. Camb. Phil. Soc.* xii. (1873) 84.—Hab. ?
supra-cordata, Burch. ex Hiern, l. c. 81 = cordata.
villosa, *Linn. Syst.* ed. XII. 302.—Afr. austr.

ROYENIA, Houst. ex Linn. Gen. ed. I. 348 (1737) = **Loeselia**, Linn. (Polemon.).

ROYLEA, Nees, ex Steud. Nom. ed. II. ii. 475 (1841) = **Melanocenchris**, Nees (Gramin.).
Pennisetum, Nees, ex Steud. l. c. = M. Royleana.

ROYLEA, Wall. Pl. As. Rar. i. 57. t. 74 (1830). *LABIATAE*, Benth. & Hook. f. ii. 1212.
elegans, *Wall. l. c.*—Reg. Himal.

RUAGEA, Karst. Fl. Columb. ii. 51. t. 126 (1862 ?) = **Guarea**, Linn. (Meliac.).
glabra, Triana & Planch. in Ann. Sc. Nat. Sér. V. xv. (1872) 368 = G. Trianae.
pubescens, Karst. Fl. Columb. ii. 51. t. 126 = G. Ruagea.

RUBACHIA, Berg, in Linnaea, xxvii. (1854) 11 = **Marlierea**, Cambess. (Myrtac.).
Antonia, Berg, in Mart. Fl. Bras. xiv. I. 29.—Bras.
glomerata, Berg, l. c. 31, 536.—Bras.
involucrata, Berg, l. c. 29.—Bras.
lateriflora, Berg, in Linnaea, xxvii. (1854) 12.—Bras.
Neuwiedeana, Berg, in Mart. Fl. Bras. xiv. I. 29.—Bras.
obumbrans, Berg, l. c. 28.—Bras.
spathulata, Berg, l. c. 535.—Bras.
spiciflora, Berg, in Linnaea, xxvii. (1854) 12.—Bras.

RUBENTIA, Boj. ex Steud. Nom. ed. II. ii. 475 (1841) = **Toddalia**, Juss. (Rutac.).
angustifolia, Boj. ex Steud. l. c. = T. aculeata.

RUBENTIA, Comm. ex Juss. Gen. 378 (1789) = **Elaeodendron**, Jacq. f. (Tiliac.).
angustifolia, Colla, Hort. Ripul. App. ii. 356 = E. orientale.
longifolia, Desf. Tabl. ed. I. 201 = E. orientale.
mauritiana, Desf. l. c. = E. orientale.
olivina, J. F. Gmel. Syst. 408 = E. orientale.

RUBEOLA, Tourn. ex Adans. Fam. ii. 144 (1763) = **Crucianella**, Linn. (Rubiac.).
angustifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = C. angustifolia.
heterophylla, Moench, Meth. 526 = C. latifolia.
latifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = C. latifolia.
linearifolia, Moench, Meth. 525 = C. angustifolia.
maritima, Moench, l. c. 526 = C. maritima.

RUBIA, [Tourn.] Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 149.
acaliculata, Cav. Ic. ii. 75. t. 195 = tinctorum ?
aculeata, Royle, Illustr. Bot. Himal. 237, nomen = himalayensis.
acuminata, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 127 = Galium uropetalum.
affinis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 111 = *vilis*.
alata, Wall. in Roxb. Fl. Ind. ed. Carey, i. 384 = cordifolia.
albicaulis, *Boiss. Diagn.* Ser. I. iii. 53.—Persia.
anglica, Huds. Fl. Angl. ed. I. 54 = peregrina.

RUBIA:—
angustifolia, *Linn. Mant.* i. 39.—Lusit.; Ins. Balear.
angustissima, *Wall. Cat.* n. 6207.—Reg. Himal.; Burma.
Aparine, Baill. Hist. Pl. vii. 373 = Galium Aparine.
aspera, Pohl, ex DC. Prod. iv. 592 (Relbunii sp.).—Bras.
Aucheri, *Boiss. Diagn.* Ser. I. iii. 54.—Persia; Arab.; Afghan.
Aucheri, C. Koch, in Linnaea, xxiv. (1851) 463, partim = peregrina.
Bocconi, Petagn. Inst. Bot. ii. 255 = peregrina.
borealis, Baill. Hist. Pl. vii. 374 = Galium boreale.
brachypoda, Boiss. Diagn. Ser. I. x. 57 = Olivierii.
Braunii, Hochst. in Flora, xxiv. (1841) I. Intell. 27 = discolor.
brevicaulis, Boiss. Diagn. Ser. I. xiii. 100 = albicaulis.
Brownei, Michx. Fl. Bor. Am. i. 81 = Galium hispidulum.
Brownei, Spreng. Syst. i. 397 = Relbunium hypocarpium.
canariensis, Poir. Encyc. Suppl. ii. 706 = fruticosa.
charaefolia, Wall. Cat. n. 6210 = angustissima.
chilensis, Molina, Sagg. Chile, 141 = Galium chilense (in Addend.).
chilensis, Willd. Sp. Pl. i. 604 = Relbunium hypocarpium.
chinensis, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 76.—As. or.
ciliata, DC. Prod. iv. 591 = Relbunium ciliatum.
clematifolia, Reinw. ex Miq. Fl. Ind. Bat. ii. 338 = cordifolia.
cordata, Thunb. Fl. Jap. 60 = cordifolia.
cordifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 344 = discolor.
cordifolia, *Linn. Syst.* ed. XII. 229.—As. trop. et temp.; Afr. trop.
cordifolia, Thunb. Fl. Cap. ed. Schult. 151 = petiolaris.
corymbosa, DC. Prod. iv. 591 (Relbunii sp.).—Peruv.
cretica, Scheele, in Linnaea, xvii. (1843) 342 = Olivierii.
crocea, DC. Prod. iv. 591 (Relbunii sp.).—Peruv.
Cruciata, Baill. Hist. Pl. vii. 372 = Galium Cruciata.
dalmatica, Scheele, in Linnaea, xvii. (1843) 342 = peregrina.
debilis, *H. B. & K. Nov. Gen. et Sp.* iii. 340.—Ecuador.
diffusa, Pohl, ex DC. Prod. iv. 592 (Relbunii sp.).—Bras.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xviii. (1845) I. 310.—Abyss.
dolichophylla, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 115.—Soongar.
Doniettii, Griseb. Spicil. Fl. Rumel. ii. 156 = Olivierii.
Edgeworthii, *Hook. f. Fl. Brit. Ind.* iii. 203.—Reg. Himal.
ephedroides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 231.—Bras.
equisetoides, *Cham. & Schlecht. l. c.* 232.—Bras.
florida, *Boiss. Diagn.* Ser. I. vi. 70.—Persia.
fruticosa, *Ait. Hort. Kew.* ed. I. i. 147.—Ins. Canar.
galioides, Poir. Encyc. Suppl. ii. 705 = fruticosa.
glabra, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 112.—Bras.
glomerata, Pohl, ex DC. Prod. iv. 566 = Triodon glomeratus.
gracilis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 111 = Asperula Platygalium.
graeca, Baill. Hist. Pl. vii. 376 = Galium graecum.
guadaloupensis, *Spreng. Syst.* i. 397.—Ins. Guadal.
Haenkeana, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 363.—Chili.
himalayensis, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 86.—Reg. Himal.
hirsuta, Schlecht. ex Wedd. Chlor. And. ii. 37 = Galium hirsutum.
hirta, H. B. & K. Nov. Gen. et Sp. iii. 338 = Relbunium ciliatum.
hispida, Willd. ex Spreng. Syst. i. 397 = Relbunium hypocarpium.
hypocarpia, DC. Prod. iv. 591 = Relbunium hypocarpium.
iberica, C. Koch, in Linnaea, xxiv. (1851) 463 = tinctorum.
incana, H. B. & K. Nov. Gen. et Sp. iii. 338 = Relbunium hypocarpium.
indecora, Cham. & Schlecht. in Linnaea, iii. (1828) 229 = Relbunium hypocarpium.

RUBIA :—
intricata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 362 = Galium latoramosum.
javana, DC. Prod. iv. 588 = cordifolia.
Kotschyi, Boiss. Diagn. Ser. I. vi. 70 = albicaulis.
laevigata, DC. Prod. iv. 591 = Relbunium laevigatum.
laevis, Poir. Voy. Barb. ii. 111 = Galium Poiretianum.
laevis, Thunb. Fl. Cap. ed. Schult. 151 = Thunbergii.
Linnaeana, Baill. Hist. Pl. vii. 374 = Galium trifidum.
longifolia, Poir. Encyc. Suppl. ii. 705 = angustifolia.
lucida, Linn. Syst. ed. XII. 732 = peregrina.
lucida, Sibth. & Sm. Fl. Graec. ii. 32 = Olivierii.
Manjith, Roxb. ex Flem. in As. Res. xi. (1810) 177 = cordifolia.
micans, Fl. Corc. in Ionios Anthol. iv. 944 (Quid ?).—Ins. Corcyra.
mitis, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 112 = chinensis.
Mollugo, Baill. Hist. Pl. vii. 372 = Galium Mollugo.
monantha, *DC. Prod.* iv. 591.—Peruv.
montana, Steud. Nom. ed. II. i. 659, in syn. = praec. ?
mucronata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 363 = *diffusa* ?
Munjista, Roxb. Hort. Beng. [10] ; Fl. Ind. i. 374 = cordifolia.
nitida, H. B. & K. Nov. Gen. et Sp. iii. 339 (Relbunii sp.).—Ecuador.
noxia, A. St. Hil. Pl. Rem. Brés. 229 (Relbunii sp.).—Bras.
Olivierii, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 132.—Ins. Scio.
orinocensis, Cham. & Schlecht. in Linnaea, iii. (1828) 228 = Relbunium hypocarpium.
orinocensis, H. B. & K. Nov. Gen. et Sp. iii. 339 (Relbunii sp.).—Venezuela.
ovalis, DC. Prod. iv. 591 (Relbunii sp.).—Peruv.
palustris, Baill. Hist. Pl. vii. 373 = Galium palustre.
pauciflora, *Boiss. Diagn.* Ser. I. iii. 54.—Persia.
peregrina, *Linn. Sp. Pl.* 109.—Europ. ; Reg. Mediterr.
peregrina, Murith, Bot. Valais, 91 = tinctorum.
peregrina, Walt. Fl. Carol. 86 = Galium hispidulum.
peregrina, Willd. ex Ledeb. Fl. Ross. ii. 410 = Galium tataricum.
petiolaris, *DC. Prod.* iv. 588.—Afr. austr.
purpurea, Decne. in Jacquem. Voy. Bot. 84. t. 92 = cordifolia.
pusilla, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 363 = Galium pusillum, *Endl.*
ramosissima, Pohl, ex DC. Prod. iv. 591 = Relbunium hypocarpium.
Relbun, Cham. & Schlecht. in Linnaea, iii. (1828) 229 = Relbunium hypocarpium.
Requienii, Duby, Bot. Gall. ed. II. i. 247 = angustifolia.
Richardiana, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 362.—Chili.
rigida, Baill. Hist. Pl. vii. 372 = Galium rigidum.
rotundifolia, Poir. Encyc. Suppl. ii. 706 = peregrina.
rotundifolia, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 267.—Syria.
rupestris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 111 = Relbunium hypocarpium ?
scabra, *H. B. & K. Nov. Gen. et Sp.* iii. 340.—Ecuador.
scabra, Salisb. Prod. 59 = fruticosa.
scandens, Zoll. & Mor. Syst. Verz. Zoll. 66 = cordifolia.
secunda, Moon, Cat. Pl. Ceyl. 10 = cordifolia.
Sellowiana, Cham. in Linnaea, ix. (1834) 214 (Relbunii sp.).—Bras.
sikkimensis, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 188.—Reg. Himal.
splendens, Hoffmgg. & Link, Fl. Port. ii. 67. t. 85 = peregrina.
sylvestris, Brot. Fl. Lusit. i. 153 = peregrina.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 2 = tinctorum.
syrticola, Miq. in Nederl. Kruidk. Arch. iv. (1859) 111 = Asperula scoparia.
tatarica, Nym. Consp. 323, in syn. ; Baill. Hist. Pl. vii. 374 = Galium tataricum.
techensis, Rafin. Fl. Ludov. 76 = Galium hispidulum.
tenuifolia, Urv. in Mém. Soc. Linn. Par. i. (1822) 273 = Olivierii.
Thunbergii, *DC. Prod.* iv. 590.—Afr. austr.
tibetica, *Hook. f. Fl. Brit. Ind.* iii. 204.—Tibet occ.
tinctoria, Salisb. Prod. 59 = tinctorum.

RUBIA :—
tinctorum, Lapeyr. Hist. Abr. Pl. Pyr. 68 = Galium Aparine.
tinctorum, *Linn. Sp. Pl.* 109.—Europ. austr. ; Oriens.
tinctorum, With. Arr. Brit. Pl. ed. III. ii. 193 = peregrina.
trachyantha, Boiss. Diagn. Ser. I. vi. 70 = albicaulis.
tricornis, Baill. Hist. Pl. vii. 373 = Galium tricorne.
tuberculosa, Benth. Pl. Hartw. 195 = Relbunium tuberculosum.
valantioides, Cham. & Schlecht. in Linnaea, iii. (1828) 231 (Relbunii sp.).—Bras.
vera, Baill. Hist. Pl. vii. 372 = Galium verum.
vilis, Cham. & Schlecht. in Linnaea, iii. (1828) 230 (Relbunii sp.).— Bras.
Walteri, DC. Prod. iv. 590 = Galium hispidulum.

RUBINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Antidesma** ?, Burm. (Euphorb.).

RUBIOIDES, Soland. ex Gaertn. Fruct. i. 112 (1788) = **Opercularia**, Gaertn. (Rubiac.).
aspera, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 205 = O. aspera.
diphylla, Soland. ex A. Cunn. l. c. = O. diphylla.

RUBUS, [Tourn.] Linn. Syst. ed. I (1735). *ROSACEAE*, Benth. & Hook. f. i. 616.
AMETRON, AMPOMELE, CUMBATA, CYLASTIS, DYCTISPERMA, MANTEIA, SELNORITION, Rafin. Sylva Tellur. 159–161 (1838).
CYLACTIS, Rafin. in Am. Journ. Sc. Ser I. i. (1819) 377.
DALIBARDA, Linn. Sp. Pl. 491 (1753).
abortivus, *Kuntze, Method.* 69.—Archip. Luchu.
abruptus, Lindl. Syn. Brit. Fl. ed. I. 92 = ulmifolius.
absconditus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 167.—Europ.
acanthophyllos, *Focke, in Bremen, Abh.* iv. (1874) 161.—Venezuela.
acanthostachys, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 232 = pubescens, *Weihe.*
acaulis, Michx. Fl. Bor. Am. i. 298 = arcticus.
acerifolius, Wall. Cat. n. 744 = glomeratus.
aceroides, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 36 = trifidus.
acheruntinus, Tenore, Syll. Fl. Neap. 603 = caesius.
acicularis, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 203.—Europ.
aciculatissimus, *Hegetschw. Fl. Schw.* 490.—Helvet.
aciculatus, Weihe, ex Lej. & Court. Comp. Fl. Belg. ii. 172 = serpens.
acidacanthos, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 240 = tomentosus.
aciodontos, *Lefèv. ex P. J. Muell. in Pollichia*, xvi–xvii. (1859) 83.—Europ.
acridentulus, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 67 = hirtus.
acroleucophorus, Ripart, ex Genev. l. c. 290 = tomentosus, *Borck.*
aculeatissimus, Kaltenb. Fl. Aach. Beckens, 300 = rosaceus.
aculeatus, *Spreng. ex Steud. Nom.* ed. I. 706, nomen.—Hab. ?
aculeoidaeus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 224, nomen.—Gallia.
aculeolatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 228.—Europ.
acuminatissimus, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 146.—Java.
acuminatus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 92 = thyrsoideus.
acuminatus, *Sm. in Rees, Cycl.* xxx. n. 43.—Reg. Himal.
acutatus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 144.—Europ.
acutiflorus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 124.—Europ.
acutifolius, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 211.—Europ.
acutipetalus, Lefèv. & P. J. Muell. l. c. 174 = macrophyllus.
acutisepalus, *P. J. Muell. l. c.* 207.—Europ.
acutus, Pritz. Ic. Ind. ii. 247, sphalm. = Rumex acutus.
adenoleucus, Chabois. in Bull. Soc. Bot. Fr. vii. (1860) 267 = nemorosus.
adenophyllus, *G. Braun, ex Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 99.—Europ.

RUBUS :—
adenotrichos, *Schlecht. in Linnaea*, xiii. (1839) 267.—Mexic.
adornatus, *P. J. Muell. ex Wirtg. in Flora*, xlii. (1859) 234.—Europ.
adpressus, Ripart & Genev. in Genev. Monog. Rub. ed. II. 257 = arduennensis.
adscitus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 88 = hirtus.
adspersus, Weihe, ex Reichb. Fl. Germ. Excurs. 602 = carpinifolius.
aegopodioides, Ser. in DC. Prod. ii. 565 = triflorus.
affinis, Betcke, in Meklenburg. Archiv, iv. (1850) 93 = sulcatus.
affinis, Liebm. in Fl. Dan. t. 2539 = thyrsoideus.
affinis, Madden, ex Hook. f. Fl. Brit. Ind. ii. 337 = ellipticus.
affinis, *Weihe & Nees, Rub. Germ.* 18 (err. typ. 22).—Europ.
agastachys, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 204 = geniculatus.
aggregatus, Kaltenb. Fl. Aach. Beckens, 277 = obscurus.
agrestis, Waldst. & Kit. Pl. Rar. Hung. iii. 297. t. 268 = tomentosus, nemorosus.
agriacanthus, *Genev. Monog. Rub.* ed. II. 297.—Europ.
albatus, Bayer, Bot. Excurs. Oesterr. 289 = bifrons.
albescens, Roxb. Hort. Beng. 39 ; Fl. Ind. ii. 519 = lasiocarpus.
albicans, *Kit. ex Schult. Oesterr. Fl.* ed. II. ii. 80.—Hungar.
albicomus, Gremli, Beitr. Fl. Schw. 30 = saltuum.
albidus, E. Merc. in Reut. Cat. Pl. Vasc. Genèv. ed. II. 288 = villicaulis.
albomicans, *Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 277.—Europ.
alcaefolius, Poir. Encyc. vi. 247 = moluccanus.
alceaefolius, Vidal, Sinops. t. 46. f. C = Rolfei.
alpestris, *Blume, Bijdr.* 1108.—Java.
alpinus, *Macfad. Fl. Jamaic.* ii. 7.—Jamaic.
alterniflorus, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 160 = macrophyllus.
althaeifolius, Bab. Fl. Camb. 305 = tomentosus, *Borck.*
althaeoides, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 222.—China.
altheaefolius, Host, ex Tratt. Ros. iii. 37 = nemorosus.
amaeniflorus, Lefèv. in Lamotte, Prod. Fl. Centr. Fr. i. 256 = amoeniflorus.
ambifarius, P. J. Muell. ex Focke, Syn. Rub. Germ. 237 = tomentosus, *Borck.* ?
ambiguus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 258.—Europ.
amblycaulon, Boul. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 28 = nemorosus.
amblystachys, *P. J. Muell. in Bonplandia*, ix. (1861) 307.—Europ.
amiantinus, *Focke, Syn. Rub. Germ.* 192, 195.—Europ.
amictifolius, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 298 = tomentosus, *Borck.*
amictus, P. J. Muell. ex Genev. l. c. 156 = vestitus.
ammobius, *Focke, Syn. Rub. Germ.* 118.—Europ.
amoeniflorus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223.—Europ.
amoenus, Koehl. in Wimm. & Grab. Fl. Siles. (1829) i. II. 54 = hirtus.
amoenus, Portenschl. ex Lange, in Kjoeb. Vidensk. Meddel. (1865) 146 = ulmifolius.
amphibolus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 89.—Europ.
amphichloros, P. J. Muell. in Bonplandia, ix. (1861) 279 = Sprengelii.
amphitapos, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 240 = tomentosus, *Borck.*
amplificatus, E. Lees, in Steele, Handb. Field Bot. 58 = macrophyllus.
amplifolius, P. J. Muell. in Bonplandia, ix. (1861) 294 = hirtus.
amplithyrsus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
amygdalanthus, Focke, Syn. Rub. Germ. 174 = thyrsoideus.
anadenes, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 191 = macrophyllus.

RUBUS:—
analogus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 232 = hirtus.
anceps, Boul. & Letendre, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = umbelliformis.
anchostachys, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 32 = ulmifolius.
ancistrophorus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 268.—Europ.
Andersoni, *Hook. f. Fl. Brit. Ind.* ii. 333.—Reg. Himal.
Andersonii, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 218, 222.—Europ.
angulatus, Rafin. Fl. Ludov. 98 (Quid?).—Am. bor.
angulosus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 128.—Europ.
angustatus, *Chab. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 92.—Europ.
angustifolius, *Kaltenb. Fl. Aach. Beckens*, 274.—Europ.
angustifolius, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 119.—Europ.
anomalus, P. J. Muell. in Flora, xli. (1858) 136 = tomentosus, *Borck.*
anoplostachys, *P. J. Muell. in Bonplandia*, ix. (1861) 292.—Europ.
antarcticus, Kuntze, Method. 115 = geoides, radicans, *Cav.*, Gunnianus.
antennifer, *Hook. f. Fl. Brit. Ind.* ii. 337.—Reg. Himal.
apertionum, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 253.—Europ.
apetalus, *Poir. Encyc.* vi. 242.—Afr. trop.; Ins. Borbon.
apiculatus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 680.—Europ.
apiculiferus, P. J. Muell. & Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 126 = ulmifolius.
appendiculatus, *Tratt. Ros.* iii. 31.—Hungar.
approximatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 242.—Europ.
approximatus, Quest. ex Focke, Syn. Rub. Germ. 307 = Fuckeli.
apricus, Wimm. Fl. Schles. ed. III. (1857) 626 = Koehleri.
aralioides, *Hance, in Journ. Bot.* xxii. (1884) 41.—China.
archimonophyllus, *Kuntze, Method.* 124, in nota.—Hab.?
arcticus, *Linn. Sp. Pl.* 494.—Reg. bor. et arct.
arcuatus, Kuntze, Method. 69 = moluccanus.
arduennensis, *Libert, ex Lej. Fl. Spa*, ii. 317.—Europ.
arenarius, Koehl. ex Wimm. & Grab. Fl. Siles. (1829) i. ii. 26 = fruticosus.
arenarius, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 38 = tomentosus, *Borck.*
arenosus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223, nomen.—Europ.
Areschougii, Blytt, in Bot. Notiser (1875), 48 = caesius.
argentatus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 93 = pubescens, *Weihe.*
argenteus, C. C. Gmel. Fl. Bad. ii. 434 = tomentosus, *Borck.*
argenteus, Gremli, Beitr. Fl. Schw. 48 = thyrsoideus.
argenteus, Weihe & Nees, Rub. Germ. 45. t. 19 = pubescens, *Weihe.*
argenteus, Wirtg. ex Focke, Syn. Rub. Germ. 195 = amiantinus.
argutifolius, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 150 = Sprengelii.
argutus, Link, Enum. Hort. Berol. ii. 60 = villosus.
argyropsis, Focke, Syn. Rub. Germ. 170 = thyrsoideus.
armeniacus, *Focke, in Bremen, Abh.* iv. (1874) 183.—Armenia.
Arrhenii, Beurl. ex Nym. Consp. 221 = corylifolius.
Arrhenii, *Lange, Haandb. Dansk. Fl.* ed. I. 309.—Europ.
arvinus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 285.—Europ.
Aschoffii, Weihe, ex Steud. Nom. ed. II. ii. 476 = corylifolius.
asper, D. Don, Prod. 234 = rosaefolius.
asper, *J. & C. Presl, Delic. Prag.* 222.—Europ.
asper, Royle, ex Hook. f. Fl. Brit. Ind. ii. 336 = macilentus, *Jacq.*
asper, Weihe, ex Lej. & Court. Comp. Fl. Belg. ii. 174 = rudis.

RUBUS:—
aspericaulis, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 141.—Europ.
asperrimus, Ripart, ex Genev. Monog. Rub. ed. II. 104 = Sprengelii.
aspreticolus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 259.—Europ.
assamensis, *Focke, in Bremen, Abh.* iv. (1874) 197.—Reg. Himal.
asservatus, Ripart & Genev. in Genev. Monog. Rub. ed. II. 136 = hirtus.
assurgens, Boul. & Bouv. ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = nemorosus.
atlanticus, *Pomel, Nouv. Mat. Fl. Atl.* 314.—Alger.
atratus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 70, 82 = rudis.
atrocaulis, P. J. Muell. in Pollichia, xvi–xvii. (1859) 163 = villicaulis.
atrorubens, *P. J. Muell. & Lefèv. l. c.* 181.—Europ.
atrorubens, Wirtg. in Flora, xlii. (1859) 234 = adornatus.
atrovirens, P. J. Muell. in Flora, xli. (1858) 153 = rosaceus.
atrovirescens, P. J. Muell. ex Nym. Consp. 219 = Lejeunei?
attenuatus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 96 = Guentheri.
australis, *Forst. f. Prod.* 40.—N. Zel.
australis, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 155; Nov. Pl. Sp. Decas, iii. 32 = tomentosus, *Borck.*
austriacus, Focke, Syn. Rub. Germ. 174 = thyrsoideus.
avellanaefolius, Koehl. ex Wimm. & Grab. Fl. Siles. (1829) i. ii. 24 = plicatus.
avellanus, P. J. Muell. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 129 = ulmifolius.
aviicolus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 115.—Europ.
axillaris, *P. J. Muell. in Flora*, xli. (1858) 139.—Europ.
axilliflorus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 280.—Europ.
Babingtonii, Bell Salter, in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 307 = thyrsiflorus.
badius, *Focke, Syn. Rub. Germ.* 276.—Europ.
Bagnalli, Bloxam, ex Briggs, in Bot. Exch. Club Rep. 1876 (1878) 14 = Briggsii.
baldensis, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 144; Nov. Pl. Sp. Decas, iii. 21 = tomentosus, *Borck.*
Balfourianus, Bloxam, ex Bab. in Ann. & Mag. Nat. Hist. Ser. I. xix. (1847) 86 = corylifolius.
balticus, Focke, Syn. Rub. Germ. 350 = Koehleri.
Banningii, G. Braun, ex Focke, l. c. 148 = rhamnifolius.
Banningii, Focke, l. c. 261 = vulgaris.
Barbeyi, Favrat & Gremli, in Gremli, Excurs. Schw. ed. IV. 150 = arduennensis.
Bastardianus, Genev. in Mem. Soc. Acad. Maine-et-Loire, xiv. (1863) 87 = ulmifolius.
bavaricus, Focke, Syn. Rub. Germ. 351 = Koehleri.
Bayeri, *Focke, in Oestr. Bot. Zeitschr.* xviii. (1868) 99.—Europ.
begoniaefolius, *Holuby, l. c.* xxv. (1875) 315.—Europ.
Bellardi, Libert, ex Focke, Syn. Rub. Germ. 365 = serpens.
Bellardi, Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 688 = glandulosus.
bellidiflorus, Hort. ex C. Koch, Dendrol. i. 292 = ulmifolius.
bellulus, *P. J. Muell. & Timb. ex Martr. Fl. Tarn*, 215.—Europ.
belophorus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 111.—Europ.
Bergii, Eckl. & Zeyh. Enum. 262 = fruticosus.
berolinensis, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 16.—Europ.
Bertramii, *G. Braun, ex Focke, Syn. Rub. Germ.* 117.—Europ.
bertricensis ×, Wirtg. ex Focke, l. c. 240.—Europ.
Betckei, Marss. Fl. Neuvorpomm. 148 = Metschii.
betulinus, D. Don, Prod. Fl. Nap. 233 = acuminatus.
bhotanensis, Kuntze, Method. 67 = moluccanus.
bicolor, *Arrond. in Bull. Soc. Polym. Morb.* (1862) 113; *et ej. Cat Pl. Morb.* 34.—Gallia.
bifidus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 220 = stenopetalus.

RUBUS:—
biflorus, *Buch.-Ham. ex Sm. in Rees, Cycl.* xxx. n. 9.—Reg. Himal.
bifrons, *Vest, ex Tratt. Ros.* iii. 28.—Europ.
bijugus, Focke, in Bremen, Abh. iv. (1874) 200 = lasiocarpus.
Billotii, *P. J. Muell. in Bonplandia*, ix. (1861) 288.—Europ.
bipartitus, Boul. & Bouv. ex Genev. Monog. Rub. ed. II. 254 = arduennensis.
birmanicus, *Hook. f. Fl. Brit. Ind.* ii. 331.—Burma.
biturigensis, *Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 59.—Europ.
blanditus, P. J. Muell. ex Focke, Syn. Rub. Germ. 315 = Lejeunei.
Blondaei, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 105 = rudis.
Bloxami, Bor. Fl. Centr. Fr. ed. III. ii. 194 = thyrsiflorus.
Bloxamianus, Colem. ex Baker, in Bot. Exch. Club Rep. 1862 (1863) 8 = R. Radula.
Bloxamii, *Lees, in Steele, Handb. Field Bot.* 55.—Europ.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* vi. 220.—N. Granat.
boliviensis, *Focke, in Bremen, Abh.* iv. (1874) 158.—Bolivia.
bonariensis, *Steud. Nom.* ed. II. ii. 477, nomen.—Reg. Argent.
Boraeanus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 87.—Europ.
borbonicus, *Pers. Syn.* ii. 51.—Ins. Borbon.
borealis, Spach, ex Focke, in Bremen, Abh. iv. (1874) 147 = strigosus.
Borreri, Bell Salter, in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 306 = Sprengelii.
Boschianus, *Zoll. in Tijdschr. Nederl. Ind.* xiv. (1857) 176.—Java.
bosquetianus, *Timb. & P. J. Muell. ex Martr. Fl. Tarn*, 212.—Europ.
Bouveti, Genev. Monog. Rub. ed. II. 296 = ulmifolius.
brachyadenes, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 33 = foliosus.
brachyandrus, *Gremli, Beitr. Fl. Schw.* 29.—Europ.
brachyanthus, *Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 98.—Europ.
brachybotrys, *Focke, in Nuov. Giorn. Bot. Ital.* xvi. (1884) 170.—Europ.
brachyclados, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 128 = ulmifolius.
brachypetalus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 96.—Europ.
brachyphyllos, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 158 = arduennensis.
brachystachys, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 200.—Europ.
brachystemon, *Heimerl, in Oestr. Bot. Zeitschr.* xxxii. (1882) 109.—Austria.
bracteatus, Bor. Fl. Centr. Fr. ed. III. ii. 193 = thyrsiflorus.
bracteosus, A. Gray, ex Kuntze, Method. 54 = Sieboldii.
bracteosus, *Weihe, in Lej. & Court. Comp. Fl. Belg.* ii. 162.—Europ.
brasiliensis, *Mart. Cat. Hort. Monac.* (1829) 173.—Bras.
bregutiensis, Kern. ex Focke, Syn. Rub. Germ. 300 = vestitus.
brevicaulis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 94.—Europ.
breviflorus, *Gremli, l. c.* 97.—Europ.
brevipes, *Gremli, l. c.* 96.—Europ.
breviramus, Ripart & Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 47 = Sprengelii.
brevis, *Gremli, Beitr. Fl. Schw.* 33.—Europ.
Briggsii, *Bloxam, in Journ. Bot.* vii. (1869) 33.—Europ.
bucciburgensis, Banning & Focke, ex Focke, Syn. Rub. Germ. 148 = rhamnifolius.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 36.—Japon.
buhnensis, G. Braun, ex Focke, Syn. Rub. Germ. 311 = obscurus, *Kaltenb.*
bursonnensis, Lefèv. ex Genev. Monog. Rub. ed. II. 110 = thyrsiflorus.
caeruleus, Gilib. Fl. Lituan. ii. 242 = caesius.
caesio-idaeus, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 294.—Europ.

RUBUS :—
caesio-idaeus, P. J. Muell. in Flora, xli. (1858) 183 = caesius.
caesius, Boj. Hort. Maurit. 127 = fruticosus.
caesius, *Linn. Sp. Pl.* 493.—Europ.; Oriens; As. bor.
Caflischii, *Focke, Syn. Rub. Germ.* 278.—Europ.
calanthelos, Focke, l. c. 261 = vulgaris.
calcareus, P. J. Muell. in Flora, xli. (1858) 181 = virgultorum.
calcareus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 254 = pubescens, *Weihe*.
Caldesianus, *Focke, in Nuov. Giorn. Bot. Ital.* xvi. (1884) 169.—Europ.
caliculatus, Kaltenb. Fl. Aach. Beckens, 283 = serpens.
californicus, *Kuntze, Method.* 102.—Calif.
callianthus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 294 = nemorosus.
calliphyllus, P. J. Muell. in Flora, xli. (1858) 168 = hirtus.
calvatus, Lees, in Kirby, Fl. Leicest. 42 = silvaticus.
calycinoides, Kuntze, Method. 67, 78 = moluccanus.
calycinus, *Wall. in D. Don, Prod. Fl. Nep.* 235 —Reg. Himal.
campestris, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 284.—Europ.
canadensis, *Linn. Sp. Pl.* 494.—Am. bor.
canadensis, Torr. Fl. N. U. St. i. 488 = triflorus.
canaliculatus, *P. J. Muell. in Flora*, xli. (1858) 132.—Europ.
candefactus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 221, nomen.—Europ.
candicans, Lasch. in Linnaea, viii. (1833) 297 = thyrsoideus.
candicans, *Weihe, ex Reichb. Fl. Germ. Excurs.* 601.—Europ.
canescens, DC. Cat. Hort. Monsp. 139 = tomentosus, *Borck.*
cannabinus, Boul. & Letendre, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 222 = ulmifolius.
cannabinus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 127.—Europ.
canus, Kit. in Rochel, Pl. Banat. Rar. 26, nomen = tomentosus, *Borck.*
cardiophyllus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 86 = arduennensis.
carisiensis, Ripart & Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 55 = ulmifolius.
carneiflorus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 169.—Europ.
carneistylus, *P. J. Muell. in Flora*, xli. (1858) 179.—Europ.
carpinetorum, *Freyn, in Verh. Zool.-Bot. Ges. Wien*, xxxi. (1882) 373.—Europ.
carpinifolius, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 76 = macrophyllus.
carpinifolius, Lees, in Steele, Handb. Field Bot. 58 = rhamnifolius.
carpinifolius, J. & C. Presl, Delic. Prag. 220 = plicatus.
carpinifolius, Weihe & Nees, Rub. Germ. 36. t. 13.—Europ.
*Caspary*i, Wirtg. ex Focke, Syn. Rub. Germ. 324 = R. Radula.
castoreus, *Fries, Mant.* ii. 37.—Europ.
caucasicus, *Focke, in Bremen, Abh.* iv. (1874) 183.—Reg. Caucas.
cavatifolius, P. J. Muell. ex Bab. in Journ. Bot. xvi. (1878) 144 = Koehleri.
cedrorum, Kotschy, ex Boiss. Fl. Orient. ii. 694 = tomentosus, *Borck.*
celebicus, *Blume, Bijdr.* 1107.—Ins. Celebes.
celtidifolius, Focke, ex Gremli, Beitr. Fl. Schw. 33 = brachyandrus.
centronotus, *Kern. in Ber. Naturw. Ver. Innsbruck*, ii. (1871) 411; *Nov. Pl. Sp. Decas*, iii. 18.—Tirol.
cerasifolius, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 98.—Europ.
cerchoviensis, *Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 91.—Europ.
cerinostylus, *P. J. Muell. in Flora*, xli. (1858) 156.—Europ.
cernuus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 194 = hirtus.
Chaboissaei, P. J. Muell. l. c. 156 = macrophyllus.
chalaranthus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 35 = Sprengelii.
Chamaemorus, Fisch. ex DC. Prod. ii. 565 = stellatus.
Chamaemorus, *Linn. Sp. Pl.* 494.—Reg. bor. et arct.

RUBUS :—
chartaceus, *Kuntze, Method.* 61, 76.—Java.
chinensis, Ser. in DC. Prod. ii. 557 = rosaefolius.
chinensis, Thunb. Diss. Rub. 8 = parvifolius.
chloracanthus, Boul. & Gillot, ex Genev. Monog. Rub. ed. II. 224 = thyrsoideus.
chlorocarpus, Bor. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 239 = ulmifolius.
chlorophyllus, Gremli, Excurs. Schw. ed. II. 161 = orthocanthus.
chlorostachys, *P. J. Muell. in Bonplandia*, ix. (1861) 303.—Europ.
chlorothyrsos, Focke, in Bremen, Abh. ii. (1871) 462 = silvaticus.
chnoophyllus, P. J. Muell. ex Genev. Monog. Rub. ed. II. 327 = tomentosus, *Borck.*
chnoostachys, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 240 = tomentosus, *Borck.*
chrysacanthus, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 454.—Europ.
chrysocarpus, Mundt, ex Cham. & Schlecht. in Linnaea, ii. (1827) 17 = rigidus, *Sm.*
chrysophyllus, *Reinw. ex Miq. Fl. Ind. Bat.* i. 1. 380.—Java.
cicur, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 311.—Hungar.
cinerascens, Bor. Fl. Centr. Fr. ed. III. ii. 197 = foliosus.
cinerascens, Weihe, ex Lej. & Court. Comp. Fl. Belg. ii. 171 = obscurus, *Kaltenb.*
cinereus, Godr. ex Gren. & Godr. Fl. Fr. i. 544 = hirtus.
cinereus, Reichb. Fl. Germ. Excurs. 607 = tomentosus, *Borck.*
cinereus, Weihe, ex Reichb. l. c. 602 = pubescens, *Weihe.*
cissoides, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 245 = australis.
citriodorus, Lej. ex Lej. & Court. Comp. Fl. Belg. ii. 163 = plicatus.
Clarkei, *Hook. f. Fl. Brit. Ind.* ii. 337.—Reg. Himal.
clematitifolius, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 455.—Europ.
clethraphilus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 167 = carpinifolius.
coarctatus, P. J. Muell. in Flora, xli. (1858) 133 = candicans.
cochinchinensis, *Tratt. Ros.* iii. 97.—China.
Colemanni, *Bloxam, in Kirby, Fl. Leicest.* 38.—Europ.
collinus, *DC. Cat. Hort. Monsp.* 139.—Europ.; As. Min.
collinus, Delile, ex Nym. Consp. 218 = tomentosus, *Borck.*
collinus, Lej. & Court. Comp. Fl. Belg. ii. 167 = arduennensis.
collivagus, Ripart, ex Genev. Monog. Rub. ed. II. 292 = ulmifolius.
coloratus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 126 = hirtus.
comintanus, Blanco, Fl. Filip. ed. II. 298 = rosaefolius.
Commersonii, Poir. Encyc. vi. 240 = rosaefolius.
commiscibilis, *P. J. Muell. in Bonplandia*, ix. (1861) 301.—Europ.
commixtus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 112.—Europ.
communis, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 222 = truncifolius.
communis, Weihe, ex Reichb. Fl. Germ. Excurs. 602 = vulgaris.
compactiflorus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 221, nomen.—Europ.
compactus, *Benth. Pl. Hartw.* 129.—Peruv.
compar, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyss.
concinnus, Focke, Syn. Rub. Germ. 309 = obscurus, *Kaltenb.*
concolor, *Ley, in Rabenh. Bot. Centralbl.* i. (1846) 434.—Europ.
concolor, *Lowe, Man. Fl. Mad.* i. 249.—Ins. Madera.
concolor, Wall. Cat. n. 733 = niveus, *Wall.*
condensatus, *P. J. Muell. in Flora*, xli. (1858) 167.—Europ.
confinis, *Lindeb. in Bot. Notiser* (1883), 105.—Suecia.
confinis, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 262.—Europ.
confinis, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 309 = tomentosus, *Borck.*

RUBUS :—
confluentinus, Wirtg. in Flora, xlii. (1859) 232 = pubescens, *Weihe.*
confusus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 231 = ulmifolius.
congestus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 130 = ulmifolius.
conglomeratus, Boul. & Letendre, ex Lefèv. l. c. xxiv. (1877) 224 = nemorosus.
conothyrsos, *Focke, Syn. Rub. Germ.* 271.—Europ.
consimilis, *P. J. Muell. in Bonplandia*, ix. (1861) 278.—Europ.
consimilis, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 274 = villicaulis.
consociatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 173.—Europ.
conspectus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 19.—Europ.
conspersus, Pritz. Ic. Ind. ii. 247, sphalm. = Rumex conspersus.
conspicuus, P. J. Muell. in Flora, xli. (1859) 71; et Pollichia, xvi–xvii. (1859) 122 = vestitus.
constrictiflorus, Boul. & Timb. ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 219 = ulmifolius.
constrictifolius, *Boul. ex Malbranche, l. c.* xxvi. (1879) 126.—Europ.
constrictus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 79.—Europ.
contemptus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 245 = ulmifolius.
controversus, Ripart, ex Genev. l. c. 244 = ulmifolius.
convexifolius, *P. J. Muell. in Bonplandia*, ix. (1861) 302.—Europ.
coptophyllus, A. Gray, in Perry, Exped. Jap. ii. 311 = microphyllus.
corchorifolius, *Linn. f. Suppl.* 263.—China; Japon.
cordatus, *Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 671.—Europ.
cordifolius, D. Don, Prod. Fl. Nep. 233 = moluccanus.
cordifolius, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 26 = glabriusculus.
cordifolius, J. & C. Presl, Delic. Prag. 219 = cordatus.
cordifolius, Weihe & Nees, Rub. Germ. 21. t. 5 = rhamnifolius.
cordigerus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 370 = serpens.
cordylifolius, Walp. Rep. ii. 16, sphalm. = Wahlbergii.
coreanus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 34.—China; Corea.
coriaceus, *Holuby, in Oestr. Bot. Zeitschr.* xxxiv. (1884) 81.—Hungar.
coriaceus, *Poir. Encyc.* vi. 237.—Peruv.
coriifolius, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 157.—Mexic.
coronarius, Sweet, Hort. Brit. ed. I. 144 = rosaefolius.
coronatus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 222 = serratifolius.
corylifolius, Bess. Enum. Pl. Volh. 47 = nessensis, sulcatus.
corylifolius, DC. Fl. Fr. iv. 475 = nemorosus.
corylifolius, Hayne, Arzn. Gew. iii. t. 11 = plicatus.
corylifolius, *Sm. Fl. Brit.* 542.—Europ.
corylinus, *P. J. Muell. in Flora*, xli. (1858) 169.—Europ.
corymbiflorus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 283.—Europ.
corymbosus, P. J. Muell. in Flora, xli. (1858) 151 = orthocanthus.
corymbosus, Weihe, ex Reichb. Fl. Germ. Excurs. 600 = nitidus.
costaricanus, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 159.—Costa Rica.
costatus, Dum. ex Focke, Syn. Rub. Germ. 161 = thyrsoides.
Courtoisianus, Lej. in Lej. & Court. Comp. Fl. Belg. ii. 170 = vestitus.
crassifolius, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 236 = pubescens, *Weihe.*
crassus, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 381.—Hungar.
crataegifolius, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 98.—China.
Crippsii, *Clarke, Trav.* i. ed. II. *Add.* p. xxii.—Turcia.
crispus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 147.—Europ.
cruentatus, P. J. Muell. l. c. 294 = obscurus, *Kaltenb.*
Cumingii, *Kuntze, Method.* 76.—Ins. Philipp.

RUBUS:—
Cunctator, *Focke, Syn. Rub. Germ.* 281.—Europ.
cuneatus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 126 = ulmifolius.
cuneifolius, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 282.—Europ.
cuneifolius, *Pursh, Fl. Am. Sept.* i. 347.—Am. bor.
cuneifolius, Weihe, ex Reichb. Fl. Germ. Excurs. 602 = vulgaris.
Cupanianus, *Guss. Fl. Sic. Syn.* i. 570.—Sicil.
curtidens, Gremli, Excurs. Schw. ed. IV. 144 = foliosus.
curvistylus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 125 = vestitus.
cuspidatus, P. J. Muell. in Flora, xli. (1858) 180 = nemorosus.
cuspidiferus, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 89 = ulmifolius.
cuspidifolius, *P. J. Muell. l. c.* 172.—Europ.
cyclopetalus, Focke, Syn. Rub. Germ. 169 = thyrsoideus.
cynomorus, *Genev. in Mém. Soc. Acad. Maine-et Loire*, x. (1861) 21.—Europ.
Dalibarda, *Linn. Sp. Pl.* ed. II. 708.—Am. bor.
dalibardoides, Kuntze, Method. 110 = calycinus.
dalmaticus, Guss. Fl. Sic. Syn. i. 567 = ulmifolius.
dalmatinus, Tratt. ex Focke, Syn. Rub. Germ. 184 = ulmifolius.
darschilingensis, Kuntze, Method. 67 = moluccanus.
dasyclados, *Kern. in Ber. Naturw. Ver. Innsbruck*, ii. (1871) 155; *Nov. Pl. Sp. Decas*, iii. 38.—Europ.
Davidianus, *Kuntze, Method.* 58.—China.
debilis, *Ball, in Journ. Bot.* xi. (1873) 332.—Marocc.
debilis, *Boul. ex Genev. Monog. Rub.* ed. II. 115.—Europ.
debilitatus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 15 = caesius.
Decaisneanus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = papulosus.
Decheni, Wirtg. ex Focke, Syn. Rub. Germ. 220 = macrophyllus.
decipiens, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 158.—Europ.
Decloetii, Ortm. in Flora, xviii. (1835) 489 = nitidus.
decorus, *P. J. Muell. in Flora*, xli. (1858) 151.—Europ.
decumbens, *Thunb. Diss. Rub.* 5.—Am. bor.
decurtatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 210.—Europ.
degener, *Kuntze, Method.* 122.—Hab. ?
degener, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 22 = caesius.
delectus, P. J. Muell. ex Nym. Consp. 221 = nemorosus.
delicatulus, *Martr. Fl. Tarn*, 206.—Gallia.
delicatus, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 312.—Europ.
deliciosus, *Torr. in Ann. Lyc. N. York*, ii. (1828) 196.—Am. bor. occ.
deltaefolius, P. J. Muell. in Pollichia, xvi–xvii. (1859) 256 = caesius.
deltoideus, P. J. Muell. in Flora, xli. (1858) 181 = tomentosus, *Borck.*
demotus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xiv. (1863) 91 = ulmifolius.
densiflorus, Gremli, Beitr. Fl. Schw. 36 = silvaticus.
dentatus, Bab. in Ann. & Mag. Nat. Hist. Ser. I. xvii. (1846) 241 = Leightonianus.
dentatus, Bloxam, in Kirby, Fl. Leicest. 39 = glandulosus.
denticulatus, *Kern. ex Focke, Syn. Rub. Germ.* 282.—Europ.
dentosus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223, nomen.—Europ.
depauperatus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 274 = nemorosus.
derasus, *Lefèv. & P. J. Muell. l. c.* 239.—Europ.
Deseglisei, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 130.—Europ.
despectus, Genev. Monog. Rub. ed. II. 132 = hirtus.
Dethardingii, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1858) 18, nomen.—Europ.
dichromus, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 106 = Bayeri.
dicrophilus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 392 = orthocanthus.
diffusus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.

RUBUS:—
digitatus, *Spreng. Pugill.* i. 34.—Hab. ?
dimorphus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 124.—Europ.
discerptus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 146 = rudis.
discoideus, *P. J. Muell. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 58.—Europ.
discolor, Auct. Scand. ex Nym. Consp. 217 = villicaulis.
discolor, Gren. & Godr. Fl. Fr. i. 546 = bifrons.
discolor, Weihe & Nees, Rub. Germ. 46, partim = ulmifolius.
discolor, Weihe & Nees, l. c. t. 20 = pubescens, *Weihe.*
discriminatus, P. J. Muell. ex Nym. Consp. 221 = nemorosus.
disjectus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 132.—Europ.
disjunctus, *P. J. Muell. & Lefèv. l. c.* 216.—Europ.
disparatus, *P. J. Muell. l. c.* 107.—Europ.
dissectus, Tenore, Ind. Sem. Hort. Neap. (1830) 13 = tomentosus, *Borck.*
dissectus, *Thunb. Diss. Rub.* 5.—China.
dissociatus, Boul. & Malbranche, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = scaber.
distans, D. Don, Prod Fl. Nep. 256 = lasiocarpus.
distractus, P. J. Muell. in Bonplandia, ix. (1861) 296 = Koehleri.
divaricatus, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 382.—Hungar.
divaricatus, P. J. Muell. in Flora, xli. (1858) 130 = carpinifolius.
divergens, *P. J. Muell. l. c.* 182.—Europ.
divergens, *Neum. in Bot. Notiser* (1885), 90.—Europ.
diversifolius, Lindl. Syn. Brit. Fl. ed. I. 93 = nemorosus.
diversifolius, *Tineo, Pl. Rar. Sicil.* 41.—Sicilia.
divexiramus, P. J. Muell. ex Genev. Monog. Rub. ed. II. 88 = tomentosus, *Borck.*
domesticus, Pritz. Ic. Ind. ii. 247, sphalm. = Rumex domesticus.
Donianus, Spreng. Syst. v. 602 = foliolosus.
Douglasii, Steud. Nom. ed. II. ii. 477 = leucodermis.
Drejeri, G. Jens. in Fl. Dan. t. 3023 = mucronatus.
droseraceus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = exsolutus.
dryadeus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 28 = corylifolius.
drymophilus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 223.—Europ.
dubiosus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 237 = tomentosus, *Borck.* ?
dumetorum, Schlecht. in Linnaea, xiii. (1839) 267 = Schiedeanus.
dumetorum, Weihe, ex Boenn. Prod. Fl. Monast. 153 = nemorosus.
dumosus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 220, nomen.—Europ.
dumosus, Salisb. Prod. 364 = fruticosus.
duricorius, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 23 = tomentosus, *Borck.*
durus, *Wright, in Sauv. Fl. Cub.* 36.—Cuba.
dyscolus, Ripart & Genev. Monog. Rub. ed. II. 44 = nemorosus.
Ebneri, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 169; Nov. Pl. Sp. Decas, iii. 46 = orthocanthus.
echinaceus, *Kern. in Oestr. Bot. Zeitschr.* xix. (1869) 201.—Hungar.
echinatus, Lindl. Syn. Brit. Fl. ed. I. 94 = R. Radula.
echinatus, *P. J. Muell. in Flora*, xli. (1858) 171.—Europ.
echinophora, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 32 = echinatus.
Ecklonii, *Focke, in Bremen, Abh.* iv. (1874) 176.—Afr. austr.
effusus, *Lefèv. & P. J. Muell. in Pollicha*, xvi-xvii. (1859) 116.—Europ.
eglandulosus, *Lefèv. & P. J. Muell. l. c.* 164.—Europ.
Eglanteria, Tratt. Ros. iii. 9 = rosaefolius.
egregius, Focke, in Bremen, Abh. ii. (1871) 463 = thyrsoideus.
eifeliensis, Wirtg. in Flora, xlii. (1859) 235 = Menkei.
elatior, Focke, in Gremli, Beitr. Fl. Schw. 50 = thyrsoideus.
elegans, *P. J. Muell. in Flora*, xli. (1858) 170.—Europ.

RUBUS:—
ellipticus, *Sm. in Rees, Cycl.* xxx. n. 16.—Ind. or.
elongatus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 91 = macrophyllus.
elongatus, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 283.—Europ.
elongatus, *Sm. Pl. Ic. Ined.* t. 62.—Java.
emarginatus, *P. J. Muell. in Flora*, xli. (1858) 164.—Europ.
emersistylus, Bab. in Journ. Bot. xvi. (1878) 175 = Briggsii.
emersistylus, *P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 77.—Europ.
enoplostachys, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 130 = ulmifolius.
Enslenii, Tratt. Ros. iii. 63 = villosus.
entomodontos, P. J. Muell. ex Focke, Syn. Rub. Germ. 312 = obscurus.
epipsilos, Focke, l. c. 258 = villicaulis.
erectus, J. F. Gmel. Syst. 1616, sphalm. = *serratus.*
erectus, *Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles.* (1829) i. II. 52.—Europ.
ericetorum, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 218, 223 = linguifolius.
eriocarpa, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 162.—Mexic.
eriophyllus, Ripart, ex Genev. Monog. Rub. ed. II. 337 = tomentosus, *Borck.*
eriostachys, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 225.—Europ.
erubescens, Wirtg. in Flora, xlii. (1859) 234 = obscurus.
erythracanthus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 250.—Europ.
erythradenes, P. J. Muell. in Bonplandia, ix. (1861) 288 = hirtus.
erythrinus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 196 = Sprengelii.
erythroclados, *Mart. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 62.—Bras.
erythrostylos, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 157.—Europ.
eurypetalus, Focke, Syn. Rub. Germ. 209 = villicaulis.
eurypetalus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 14.—Europ.
euryphyllos, *P. J. Muell. in Bonplandia*, ix. (1861) 295.—Europ.
exactus, *P. J. Muell. in Flora*, xli. (1858) 167.—Europ.
exalbatus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 161.—Europ.
exasperatus, *Lefèv. & P. J. Muell. l. c.* 149.—Europ.
excavatus, Lefèv. & P. J. Muell. l. c. 135 = Sprengelii.
excelsior, P. J. Muell. ex Focke, Syn. Rub. Germ. 167 = candicans.
excultus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 221 = ulmifolius.
excurvatus, Kuntze, Method. 71 = moluccanus.
exilis, Lange, in Fl. Dan. t. 3027 = R. Radula.
expallescens, Timb. & Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 128 = ulmifolius.
expansus, Lefèv. & P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 44 = tomentosus, *Borck.*
exsecatus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 311 = obscurus.
exsolutus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 241.—Europ.
exsuccus, Steud. ex A. Rich. Tent. Fl. Abyss. i. 256 = apetalus.
extensifolius, Boul. & Ripart, ex Genev. Monog. Rub. ed. II. 334 = tomentosus, *Borck.*
fagicola, *Martr. Fl. Tarn*, 219.—Gallia.
fagifolius, *Cham. & Schlecht. in Linnaea*, v. (1830) 571.—Mexic.
Fairholmianus, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 5 = moluccanus.
falcarius, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 232 = pubescens, *Weihe.*
falcatus, Kaltenb. Fl. Aach. Beckens, 266 = candicans.
falciferus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 84.—Europ.
Falconeri, Kuntze, Method. 68 = moluccanus.
fallacinus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 263 = villicaulis.
fallax, T. M. Chab. ex P. J. Muell. in Pollichia, xvi–xvii. (1859) 155 = thyrsoideus.

RUBUS:—
fallax, Kuntze, Method. 55 = moluccanus.
fasciculatus, P. J. Muell. in Flora, xli. (1858) 182 = tomentosus, *Borck.*
fastigiatus, Bab. ex Nym. Consp. 221 = corylifolius.
fastigiatus, Weihe & Nees, Rub. Germ. 16. t. 2 = nessensis, sulcatus.
fastuosus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 220, nomen.—Europ.
ferox, Vest, ex Tratt. Ros. iii. 40 = horridus, nemorosus.
ferox, *Wall. Cat.* n. 724.—Reg. Himal.; Burma.
ferox, Weihe, ex Boenn. Prod. Fl. Monast. 153 = nemorosus.
ferriararum, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 40 = hirtus.
ferrugineus, *Wickst. in Vet. Acad. Handl. Stockh.* (1827) 68.—Ins. Guadaloup.
ferruginosus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 231.—Europ.
festivus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 314 = Lejeunei.
Fieberi, Ortm. in Opiz, Nom. 15 = nemorosus.
fimbrifolius, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 256 = thyrsoideus.
Finlaysonianus, *Wall. Cat.* n. 7109.—Siam.
firmulus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 94.—Helvet.
Fischii, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 5, 19.—Europ.
fissipetalus, P. J. Muell. in Flora, xli. (1858) 153 = rudis.
fissus, Lindl. Syn. Brit. Fl. ed. II. 92 = nessensis.
flaccidifolius, *P. J. Muell. in Bonplandia*, ix. (1861) 300.—Europ.
flaccidus, P. J. Muell. in Flora, xli. (1858) 134 = villicaulis.
flagellaris, Hook. Comp. Bot. Mag. i. (1835) 24 = trivialis.
flagellaris, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 261.—Europ.
flagellaris, Schrank, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 26 = triflorus.
flagellaris, Willd. Enum. Hort. Berol. 549 = canadensis.
flagelliformis, Koehl. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 689 = glandulosus.
flavescens, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 195.—Europ.
flavo-virens, Genev. in Lamotte, Prod. Fl. Centr. Fr. i. 255, nomen = vestitus.
flavus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 234 = ellipticus.
flexicaulis, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 71, 96 = Sprengelii.
flexipes, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 126 = ulmifolius.
flexuosus, *Lej. Rev. Fl. Spa*, 238, nomen.—Europ.
flexuosus, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 240 = saltuum.
floccosus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 128 = ulmifolius.
floribundus, *H. B. & K. Nov. Gen. et Sp.* vi. 219.—Ecuador.
floribundus, Ley, in Rabenh. Bot. Centralbl. i. (1846) 435 = Lejeunei.
floribundus, Weihe, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 684 = thyrsiflorus.
floridus, Tratt. Ros. iii. 73 = villosus.
Fockeanus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 206.—Reg. Himal.
foliolatus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 216.—Europ.
foliolosus, *D. Don, Prod. Fl. Nep.* 256.—Reg. Himal.
foliosus, Bab. Brit. Rub. 243 = silvaticus.
foliosus, Gremli, Beitr. Fl. Schw. 35 = vestitus.
foliosus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 682.—Europ.
fontinalis, *Kuntze, Method.* 60.—Java.
Fordii, Hance, in Journ. Bot. xxi. (1883) 298 = Hanceanus.
formidabilis, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 128.—Europ.
formosensis, *Kuntze, Method.* 73.—Ins. Formosa.
formosus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 127.—Europ.
fortis, *Focke, Syn. Rub. Germ.* 70.—Europ.
fossicola ×, Holuby, in Oestr. Bot. Zeitschr. xxiii. (1873) 381.—Europ.

RUBUS:—
Fragaria, *F. W. Schmidt, in Neue. Abh. Boehm. Ges.* i. (1791) 38.—Europ.
fragariaeflorus, *P. J. Muell. in Flora*, xli. (1858) 173.—Europ.
fragarioides, *Bertol. Misc. Bot.* xxii. 16.—Reg. Himal.
fragrans, Focke, Syn. Rub. Germ. 172 = thyrsoideus.
fragrans, Salisb. Prod. 364 = R. Idaeus.
frambaesianus, Lam. Fl. Fr. iii. 135 = R. Idaeus.
Frangipani, Tineo, ex Guss. Suppl. Fl. Sic. Prod. 157 = ulmifolius.
fraternus, Gremli, Beitr. Fl. Schw. 34 = vestitus.
fraxinifolius, *Poir. Encyc.* vi. 242.—Java.
frondosus, Bigel. Fl. Bost. ed. II. 199 = villosus.
fruticetorum, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 279.—Europ.
fruticosus, Gaertn. Mey. & Scherb. Fl. Wett. ii. 236 = plicatus.
fruticosus, *Linn. Sp. Pl.* 493.—Europ.
fruticosus, Lour. Fl. Cochinch. 325 = cochinchinensis.
fruticosus, Marsh. Arbust. 137 = villosus.
fruticosus, Pollich, Hist. Pl. Palat. ii. 58 = nessensis, sulcatus.
fruticosus, Roth, Tent. Fl. Germ. i. 220 = rhamnifolius.
fruticosus, Sm. Fl. Brit. ii. 543 = ulmifolius.
fruticosus, Weihe & Nees, Rub. Germ. 24 = candicans.
Fuckeli, *Wirtg. in Flora*, xlii. (1859) 235.—Europ.
fulcratus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 178.—Europ.
furfuraceus, Wall. Cat. n. 739 = lasiocarpus.
fuscoater, *Weihe, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 681.—Europ.
fuscoater, Wirtg. Fl. Preuss. Rheinprov. 156 = adornatus.
fuscus, Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 681 = pallidus.
gallicus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 219, 221.—Europ.
Gardenerianus, Kuntze, Method. 42 = moluccanus.
geifolius, Kuntze, l. c. 97 = incisus.
Genevierii, Bor. Fl. Centr. Fr. ed. III. ii. 193 = thyrsiflorus.
geniculatus, Focke, in Bremen, Abh. i. (1868) 289 = carpinifolius.
geniculatus, *Kaltenb. Fl. Aach. Beckens*, 267.—Europ.
geoides, *Sm. Pl. Ic. Ined.* t. 19.—Reg. Magellan.
Gerard-Martini, P. J. Muell. in Bonplandia, ix. (1861) 287 = Lejeunei.
geromensis, P. J. Muell. in Flora, xli. (1858) 185 = serpens.
giganteus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 72, 98.—Europ.
glaber, Mill. Gard. Dict. ed. VIII. n. 4 = R. Idaeus.
glaber, Weihe, ex Reichb. Fl. Germ. Excurs. 600 = nitidus.
glaberrimus, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 80 = leucanthus.
glabratus, *H. B. & K. Nov. Gen. et Sp.* vi. 221.—Am. austr.
glabrefactus, P. J. Muell. ex Focke, Syn. Rub. Germ. 227 = tomentosus, *Borck.*
glabriusculus, *Hassk. Cat. Hort. Bogor. Alt.* 267.—Malaya.
glandulosus, *Bell. in Mém. Acad. Turin*, 1790–91 (1793), 230.—Europ.; Oriens.
glandulosus, Guenth. Grab. & Wimm. Enum. Pl. Siles. (1824) 86 = Guentheri.
glandulosus, Menke, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 680 = Menkei.
glandulosus, Reichb. Fl. Germ. Excurs. 607 = hirtus.
glandulosus, Schleich. & Willd. ex Boenn. Prod. Fl. Monast. 152 = Schleicheri.
glandulosus, Spreng. ex Wallr. Sched. Crit. 231 = corylifolius.
glaphyrus, Ripart & Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 235 = pubescens, *Weihe.*
glaucicomans, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223, nomen.—Europ.
glaucifolius, Kellogg, in Proc. Calif. Acad. ed. II. i. (1873) 70 = leucodermis.
glaucocaulis, *Kuntze, Method.* 72.—Tibet.
glaucovirus, *Maass. in Verh. Bot. Ver. Brandenb.* xii. (1870) 162.—Europ.
glaucus, *Benth. Pl. Hartw.* 173.—N. Granat.
glomeratus, *Blume, Bijdr.* 1111.—Java.
glomeriflorus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 98.—Europ.

RUBUS:—
glomicellus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 221, nomen.—Europ.
Godroni, Gren. & Godr. Fl. Fr. i. 540 = pubescens, *Weihe.*
Godroni, Lec. & Lamotte, Cat. Pl. Centr. Fr. 151 = nemorosus.
Godroni, *P. J. Muell. in Flora*, xli. (1858) 184.—Europ.
goniophyllus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 80.—Europ.
gorizianus, *Kern. in Ber. Naturw. Ver. Innsbruck*, ii. (1871) 129; *Nov. Pl. Sp. Decas*, iii. 6.—Europ.
Gowreephul, Roxb. Hort. Beng. 39; Fl. Ind. ii. 517 = ellipticus.
Grabowskii, Bab. Brit. Rub. 123 = carpinifolius.
Grabowskii, Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 32 = thyrsoideus.
gracilentus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 245.—Europ.
gracilescens, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 104 = Bayeri.
gracilicaulis, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 125 = hirtus.
gracilis, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 380.—Europ.
gracilis, J. & C. Presl, Delic. Prag. 220 = nemorosus.
gracilis, Roxb. Hort. Beng. 39; Fl. Ind. ii. 519 = niveus, *Wall.*
grandiflorus, Kaltenb. Fl. Aach. Beckens, 291 = nemorosus.
grandiflorus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 108 = macrophyllus.
grandifolius, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 32.—Ins. Madera.
grandifolius, Salisb. Prod. 364 = odoratus.
granulatus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 154.—Europ.
gratiflorus, P. J. Muell. ex Genev. Monog. Rub. ed. II. 198 = macrophyllus.
gratiosus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 153.—Europ.
gratus, Focke, in Alpers, Verz. Gefaesspfl. Stade, 26 = villicaulis.
Grayanus, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 152.—Archip. Luchu.
Gremlii, *Focke, Syn. Rub. Germ.* 266.—Europ.
Grenierii, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 221 = goniophyllus.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 327.—Reg. Himal.
grypoacanthus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 133 = hirtus.
Guenteri, Bab. in Ann. & Mag. Nat. Hist. Ser. I. xix. (1847) 17 = seq.
Guentheri, Bloxam, in Kirby, Fl. Leicest. 41 = saltuum.
Guentheri, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 679.—Europ.
guestphalicus, Focke, Syn. Rub. Germ. 431 = obscurus.
Gunnianus, *Hook. Ic. Pl.* t. 291.—Tasmania.
guyanensis, *Focke, in Bremen, Abh.* iv. (1874) 160.—Guian.
gymnostachys, Genev. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 28 = thyrsiflorus.
gymnostylos, *P. J. Muell. in Bonplandia*, ix. (1861) 306.—Europ.
Haellstroemi, Kuntze, Method. 130 = arcticus.
hakonensis, Franch. & Sav. Enum. Pl. Jap. ii. 333 = Lambertianus.
Hamiltoni, *Hook. f. Fl. Brit. Ind.* ii. 328.—Reg. Himal.
Hamiltonianus, Ser. in DC. Prod. ii. 566 = moluccanus.
hamosus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 71, 94 = arduennensis.
hamulosus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 76 = nitidus.
Hanceanus, *Kuntze, Method.* 72.—China.
Harpago, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 100.—Europ.
Hasskarlii, Miq. Fl. Ind. Bat. i. I. 381 = glomeratus.
Hasskarlii, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 306 = Menkei.
hawaiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 504. t. 56.—Ins. Sandvic.

RUBUS:—
hebecarpos, *P. J. Muell. in Bonplandia*, ix. (1861) 282.—Europ.
hedycarpus, *Focke, Syn. Rub. Germ.* 190.—Europ.
helveticus, Gremli, Beitr. Fl. Schw. 36 = silvaticus.
hemistemon, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 314 = nessensis.
heptaphyllus, Opiz, in Flora, v. (1822) 269 = nessensis.
hercynicus, *G. Braun, ex Focke, Syn. Rub. Germ.* 370.—Europ.
heteracanthus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 102.—Europ.
heteracanthus, Weihe, ex Lej. Rev. Fl. Spa, 101 = rosaceus.
heterocaulon, Ortm. in Opiz, Nom. 14 = nessensis.
heteroclitus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 127 = carpinifolius.
heteromorphus, *Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 255.—Europ.
heterophyllus, *Host, Fl. Austr.* ii. 32.—Europ.
heterophyllus, *Willd. Enum. Hort. Berol. Suppl.* 38.—Am. bor.
hevellicus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 18.—Europ.
hexagynus, *Roxb. Hort. Beng.* 39; *Fl. Ind.* ii. 516.—Reg. Himal.; Burma.
hibiscifolius, *Focke, in Bremen, Abh.* iv. (1874) 197.—Reg. Himal.
hiemalis, Kuntze, Method. 69 = moluccanus.
Hillii, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 67 = moluccanus.
himalaicus, Kuntze, Method. 66 = moluccanus.
Hippolapathum, Pritz. Ic. Ind. ii. 247, sphalm. = Rumex Hippolapathum.
hirsutuosus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 72, 98.—Europ.
hirsutus, *J. & C. Presl, Delic. Prag.* 221.—Europ.
hirsutus, *Thunb. Diss. Rub.* 7.—Japon.
hirsutus, Wimm. Fl. Schles. ed. III. (1857) 627 = Koehleri.
hirsutus, Wirtg. Prod. Fl. Pr. Rheinl. 61 = vestitus.
hirtellus, Ripart, ex Genev. Monog. Rub. ed. II. 318 = villicaulis.
hirtifolius, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 290 = vestitus.
hirtocaulis, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 251.—Europ.
hirtus, Aspegren, Förs Blekingst. Fl. 37 = Koehleri.
hirtus, Reichb. Fl. Germ. Excurs. 607 = glandulosus.
hirtus, Roxb. Hort. Beng. 38; Fl. Ind. ii. 518 = ellipticus.
hirtus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 150. t. 141.—Europ.; Reg. Caucas.
hispanicus, Willk. in Bot. Zeit. ii. (1844) 819 = ulmifolius.
hispidulus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 51 = carpinifolius.
hispidulus, Weihe, ex Focke, Syn. Rub. Germ. 361 = Schleicheri.
hispidus, *Linn. Sp. Pl.* 493.—Am. bor.
hispidus, Marsh. Arbust. 137 = trivialis.
hispidus, E. Merc. in Reut. Cat. Pl. Vasc. Genèv. ed. II. 275 = obscurus, *Kaltenb.*
Histrix, Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 687 = *Hystrix*.
Histrix, Koehl. ex Bluff & Fingerh. l. c. 681 = Koehleri.
Hochstetterorum, *Seub. Fl. Azor.* 48.—Ins. Azor.
Hoffmeisterianus, Kunth & Bouché, Ind. Sem. Hort. Berol. (1847) 14 = niveus, *Wall.*?
Hollandrei, P. J. Muell. in Flora, xli. (1858) 185 = corylifolius.
hololeucos, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 72, 104 = villicaulis.
holorodos, Ripart, ex Genev. Monog. Rub. ed. II. 256 = arduennensis.
holosericeus, *Vest, ex Tratt. Ros.* iii. 48.—Europ.
holosericeus, Weihe, ex Boenn. Prod. Fl. Monast. 153 = nemorosus.
Hookeri, *Focke, in Bremen, Abh.* iv. (1874) 198.—Reg. Himal.
horrefactus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 252.—Europ.
horricaulis, C. Muell. in Walp. Ann. vii. 858 = hirtocaulis.
horridicaulis, *P. J. Muell. in Bonplandia*, ix. (1861) 284.—Europ.

RUBUS:—
horridulus, *Hook. f. Fl. Brit. Ind.* ii. 341.—Reg. Himal.
horridus, Betcke, in Meklenburg. Archiv, iv. (1850) 132 = Koehleri.
horridus, *Hartm. Handb. Skand. Fl.* ii. 139.—Suecia.
horridus, Koehl. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 680 = Schleicheri.
horridus, Schultz, Prod. Fl. Starg. Suppl. i. 30 = nemorosus.
horridus, *Weihe, ex Spreng. Syst.* ii. 529.—Europ.
horripilus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 126.—Europ.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 375 = lasiocarpus.
horstensis ×, Banning & Focke, ex Focke, Syn. Rub. Germ. 206.—Europ.
hostilis, P. J. Muell. & Wirtg. ex Focke, l. c. 324 = R. Radula.
humifusus, Bab. in Trans. Edinb. Bot. Soc. ii. (1846) 245 = saxicolus.
humifusus, Schlecht. in Linnaea, xiii. (1839) 270 = humistratus.
humifusus, Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 685 = Schleicheri.
humilis, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 246.—Europ.
humistratus, *Steud. Nom.* ed. II. ii. 478.—Mexic.
humorosus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 219.—Europ.
humulifolius, *C. A. Mey. Beitr. Pfl. Russ. Reich.* v. 57. t. 1.—Rossia.
hungaricus, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 314.—Hungaria.
hybridus, *Kaltenb. Fl. Aach. Beckens*, 278.—Europ.
hybridus, Vest, in Steyerm. Zeitschr. i. 162; ex Focke, Syn. Rub. Germ. 105 = nessensis.
hybridus, Vill. Prosp. 46; Hist. Pl. Dauph. iii. 559 = glandulosus.
hydrastifolius, A. Gray, in Perry, Jap. Exped. ii. 311 = trifidus.
hylonomus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 224.—Europ.
hylophilus, *Ripart, ex Genev. Monog. Rub.* ed. II. 238.—Europ.
hypargyrus, Edgew. in Trans. Linn. Soc. xx. (1846) 45 = niveus, *Wall.*
hypoleucos, Vest, in Syll. Ratisb. i. (1824) 235 = tomentosus, *Borck.*
hypoleucus, Focke, Syn. Rub. Germ. 218 = macrophyllus.
hypoleucus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 143.—Europ.
hypomalacus, *Focke, Syn. Rub. Germ.* 274.—Europ.
hypomallos, P. J. Muell. ex Focke, l. c. 238 = thyrsoideus, tomentosus, *Borck.*
Hystrix, Weihe & Nees, Rub. Germ. 92 = rosaceus.
idaeoides, Ruthe, ex Focke, Syn. Rub. Germ. 411 = caesius.
Idaeus, Blanco, Fl. Filip. ed. I. 428 = parvifolius.
Idaeus, Linn. Sp. Pl. 492.—Europ.; Oriens; As. bor.
Idaeus, Pursh, Fl. Am. Sept. i. 346 = strigosus.
Idaeus, Thunb. Fl. Jap. 216 = Thunbergii, *Sieb. & Zucc.*
Idaeus, Vell. Fl. Flum. 220; v. t. 81 = floribundus.
imbricatus, *Hort, in Trans. Bot. Soc. Edinb.* iv. (1853) 113.—Europ.
immitis, Bor. Fl. Centr. Fr. ed. III. ii. 188 = nemorosus.
imperialis, *Cham. & Schlecht. in Linnaea*, ii. (1827) 13.—Bras.
inaequiaculeatus, Kuntze, Method. 97 = incisus.
inaequidentatus, Boul. & Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
inamoenus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 126.—Europ.
inaspectus, *P. J. Muell. in Bonplandia*, ix. (1861) 297.—Europ.
incarnatus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 95 = villicaulis.
incisus, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 35 = trifidus.
incisus, *Thunb. Fl. Jap.* 217.—China; Japon.
inclinabilis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 124.—Europ.
inconspicuus, Lange, ex Willk. & Lange, Prod. Fl. Hisp. iii. 221 = minutiflorus.
incultus, Wirtg. ex Focke, Syn. Rub. Germ. 369 = serpens.
incurvatus, Bab. in Trans. Bot. Soc. Edinb. iii. (1850) 54 = corylifolius.

RUBUS:—
indicus, Heyne, ex. Wall. Cat. sub n. 742 = lasiocarpus.
indicus, Leschen. ex Ser. in DC. Prod. ii. 568 = hexagynus.
indicus, *Thunb. Diss. Rub.* 5.—Ind. or.?
indotatus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 128.—Europ.
indusiatus, *Focke, Syn. Rub. Germ.* 284.—Europ.
indutus, Boul. & Vendr. ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 220, nomen = fastuosus.
inermis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 326.—Europ.
inermis, Willd. Enum. Hort. Berol. 549 = villosus.
infestus, Bill. ex Focke, Syn. Rub. Germ. 307 = Fuckeli.
infestus, Hartm. Handb. Skand. Fl. ed. III. 120 = horridus, *Hartm.*
infestus, Kaltenb. Fl. Aach. Beckens, 293 = carpinifolius.
infestus, *Waldst. & Kit. ex Steud. Nom.* ed. II. ii. 478, nomen (an seq.?).—Europ.
infestus, *Weihe, ex Boenn. Prod. Fl. Monast.* 153.—Europ.
inflatus, C. Muell. in Walp. Ann. vii. 860 = irrufatus.
innominatus, *S. Moore, in Journ. Bot.* xiii. (1875) 226.—China.
inopacatus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 117.—Europ.
insectifolius, *Lefèv. & P. J. Muell. l. c.* 176.—Europ.
insericatus, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 233 = obscurus, *Kaltenb.*
insignis, *Hook. f. Fl. Brit. Ind.* ii. 329.—Reg. Himal.
insignitus, Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 130 = ulmifolius.
insolatus, P. J. Muell. in Flora, xli. (1858) 166 = hirtus.
insularis, *Aresch. Skånes Fl.* ed. II; *et in Bot. Notiser* (1881), 158.—Suecia.
intectus, P. J. Muell. in Pollichia, xvi–xvii. (1849) 247 = Sprengelii.
integribasis, P. J. Muell. ex Genev. Monog. Rub. ed. II. 345 = nessensis.
intermedius, *Ortm. in Flora*, xviii. (1835) 496.—Europ.
intricatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 277.—Europ.
inuncatus, *Lefèv. & P. J. Muell. l. c.* 105.—Europ.
irasuensis, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 160.—Costa Rica.
irrufatus, *P. J. Muell. in Bonplandia*, ix. (1861) 291.—Europ.
Ischelus, Buch. Ham. ex Hook. f. Fl. Brit. Ind. ii. 339 = lasiocarpus.
italicus, *Link, Handb.* ii. 110.—Ital.
jamaicensis, Blanco, Fl. Filip. ed. I. 427 = rosaefolius.
jamaicensis, *Linn. Mant.* i. 75.—Am. trop.
jambosoides, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 222.—China.
japonicus, *Focke, in Bremen, Abh.* iv. (1874) 198.—Japon.
japonicus, Linn. Mant. ii. 245 = Kerria japonica.
japonicus, *Veitch, ex Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 163.—Japon.
javanicus, *Blume, Bijdr.* 1108.—Java.
Jensenii, Lange, in Fl. Dan. t. 2833 = orthacanthus.
juglandifolius, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 455.—Europ.
Kaltenbachii, Metsch, in Linnaea, xxviii. (1856) 170 = Metschii, hirtus.
Kingii, Kuntze, Method. 55 = moluccanus.
Koehleri, Bor. Fl. Centr. Fr. ed. III. ii. 195 = Deseglisei.
Koehleri, Weihe, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 689 = glandulosus.
Koehleri, *Weihe & Nees, ex Bluff & Fingerh. l. c.* 681.—Europ.
Kurzianus, Kuntze, Method. 69 = moluccanus.
lacer, Kuntze, l. c. 103 = nutkanus.
laceratus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 229.—Europ.
laciniatus, *Willd. Hort. Berol.* t. 82.—Hab.?
lacteus, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 313.—Europ.
laetevirens, *Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 95.—Europ.
laetus, *Prog. l. c.* 92.—Europ.
laevefactus, P. J. Muell. ex Focke, Syn. Rub. Germ. 257 = thyrsoideus.

RUBUS:—
laevigatus, Wall. Cat. n. 1280 = lucens.
Lambertianus, *Ser. in DC. Prod.* ii. 567.—China.
Lamottei, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 21 = tomentosus, *Borck.*
lamprococcus, *Focke, in Bremen, Abh.* i. (1867) 307. —Europ.
lamprophyllus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 94 = Bayeri.
Lamyyi, Genev. Monog. Rub. ed. II. 92 = tomentosus, *Borck.*
lanatus, Focke, in Bremen, Abh. i. (1867) 294 = vestitus.
lanatus, *Wall. Cat.* n. 746.—Reg. Himal.
Langei, G. Jens. ex Focke, Syn. Rub. Germ. 221 = silvaticus.
lanuginosus, Schlecht. ex Ledeb. Fl. Ross. ii. 69 = glandulosus.
lanuginosus, *Stev. ex Ser. in DC. Prod.* ii. 564.—Reg. Caucas. ?
Laschii, *Focke, Syn. Rub. Germ.* 402.—Europ.
lasiocarpus, *Sm. in Rees, Cycl.* xxx. n. 6.—Ind. or.; Malaya.
lasioclados, Focke, Syn. Rub. Germ. 198 = pubescens, *Weihe.*
lasiococcus, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82). 201.—Am. bor. occ.
lasiostachys, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 106.—Europ.
latebrosus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 24 = caesius.
latifoliatus, Boul. & Bouv. ex Genev. Monog. Rub. ed. II. 66 = nemorosus.
latifolius, *Bab. Man. Brit. Bot.* ed. III. 94.—Europ.
latifolius, Kuntze, Method. 70 = moluccanus.
latifrons, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
laxatus, Ripart & Genev. in Genev. Monog. Rub. ed. II. 30 = corylifolius.
laxiflorus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 206.—Europ.
Lechleri, *Focke, in Bremen, Abh.* iv. (1874) 161.—Peruv.
Lecoqui, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 27 = caesius.
Leesii, Bab. in Steele, Handb. Field Bot. 60 = R. Idaeus.
Lefevrei, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 183.—Europ.
Leguei, Genev. Monog. Rub. ed. II. 31 = tomentosus, *Borck.*
Leightoni, Lees, ex Leight. Fl. Shrops. 233 = rudis.
Leightonianus, *Bab. in Ann. & Mag. Nat. Hist.* Ser. I. xvii. (1846) 240.—Europ.
Lejeunei, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 683.—Europ.
Lejeunei, Wirtg. ex Focke, Syn. Rub. Germ. 307 = Fuckeli.
Lemaitrei, *Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 257.—Europ.
lentiginosus, Lees, in Steele, Handb. Field Bot. 60 = affinis.
lephracanthus, Boul. & Timb. ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1817) 219, sphalm. = *tephracanthus.*
lepidus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 294 = amiantinus.
leptacanthos, P. J. Muell. ex Focke, Syn. Rub. Germ. 353 = Koehleri.
leptocalyx, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 105 = Bayeri.
leptocaulon, Boul. & Letendre, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = nemorosus.
leptoclados, *P. J. Muell. in Bonplandia*, ix. (1861) 299.—Europ.
leptopetalus, *Focke, Syn. Rub. Germ.* 386.—Europ.
leptopetalus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 95.—Europ.
leptostachys, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 165.—Europ.
leucadenes, *P. J. Muell. in Bonplandia*, ix. (1861) 304. —Europ.
leucandrus, Focke, in Alpers, Verz. Gefaesspfl. Stade, 27 = villicaulis.
leucanthemus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 122 = vestitus.
leucanthus, *Hance, in Walp. Ann.* ii. 468.—China.
leucocarpus, *Arv. in Nov. Act. Nat. Cur.* xviii. (1836) 334.—Zeylan.
leucodermis, *Dougl. ex Hook. Fl. Bor. Am.* 178; *et ex Torr. & Gray, Fl. N. Am.* i. 454.—Am. bor. occ.

RUBUS:—
leucophaeus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 264 = tomentosus, *Borck.*
leucophyllus, Vest, ex Focke, Syn. Rub. Germ. 226 = tomentosus, *Borck.*
leucostachys, Dougl. ex Steud. Nom. ed. II. ii. 478 = leucodermis.
leucostachys, Schleich. ex Sm. Engl. Fl. ii. 403 = vestitus.
levefactus, P. J. Muell. ex Nym. Consp. 216 = thyrsoideus.
Leyi, Focke, Syn. Rub. Germ. 268 = silvaticus.
Libertianus, *Weihe, in Lej. & Court. Comp. Fl. Belg.* ii. 163.—Europ.
Liebmannii, *Focke, in Bremen Abh.* iv. (1874) 158.—Mexic.
ligerinus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 68, 73 = caesius.
ligulatus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
lilacinus, Wirtg. ex Focke, Syn. Rub. Germ. 376 = hirtus.
Lindebergii, P. J. Muell. in Pollichia, xvi–xvii. (1859) 292 = bifrons.
Lindleianus, Lees, in Phytol. iii. (1848) 361 = vulgaris.
lineatus, *Reinw. ex Blume, Bijdr.* 1108.—Reg. Himal.; Java.
Lingua, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 686.—Europ.
linguifolius, *P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 83.—Europ.
lingulatus, *Lefèv. ex Genev. Monog. Rub.* ed. II. 148. —Europ.
Linkianus, *Ser. in DC. Prod.* ii. 560.—Hab. ?
lithuanicus, *Gilib. Hist. Pl. Europ.* ii. 321.—Europ.
lividus, G. Braun, ex Focke, Syn. Rub. Germ. 365 = serpens.
Lloydianus, Genev. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 26 = tomentosus, *Borck.*
lobatus, Wall. Cat. sub n. 737 = calycinus.
Lobbianus, *Hook. Ic. Pl.* t. 741, 742.—Java.
Loehrii, Wirtg. in Flora, xxxix. (1856) 236 = rudis.
longicaulis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 126.—Europ.
longicuspis, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 120 = rudis.
longiflorens, Ripart, in Genev. Monog. Rub. ed. II. 276 = pubescens, *Weihe.*
longifolius, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras..
longifolius, *Host, Fl. Austr.* ii. 80.—Europ.
longifolius, *Kunze, Method.* 96.—Hab. ?
longifolius, Wall. Cat. sub n. 727 = lasiocarpus.
longipetalus, *Dougl. ex Steud. Nom.* ed. II. ii. 478, nomen.—Am. bor. occ.
longisepalus, *P. J. Muell. in Bonplandia*, ix. (1861) 297.—Europ.
longithyrsiger, Lees, ex Bab. Brit. Rub. 231 = pyramidalis.
loxensis, *Benth. Pl. Hartw.* 128.—Peruv.
lucens, *Focke, in Bremen, Abh.* iv. (1874) 199.—Reg. Himal.
lucorum, *P. J. Muell. ex Lefèv. in Pollichia*, xvi–xvii. (1859) 260.—Europ.
Ludwigii, *Eckl. & Zeyh. Enum.* 263.—Afr. austr.
luteifolius, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 219 = carpinifolius.
lutescens, Lefèv. l. c. 220 = constrictus.
luxurians, Ripart, ex Genev. Monog. Rub. ed. II. 198 = macrophyllus.
Maassii, Focke, ex Bertram, Fl. Braunschw. 75 = rhamnifolius.
macilentus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 42.—Europ.
macilentus, *Jacquem. Voy. Bot.* 49 t. 60.—Reg. Himal.
Macraei, *A. Gray, Bot. U. St. Expl. Exped.* i. 505. t. 57. —Ins. Sandvic.
macranthelos, Marss. Fl. Neuvorpomm. 147 = vestitus.
macroacanthos, Weihe & Nees, Rub. Germ. 44 = vulgaris.
macroacanthos, Wirtg. ex Focke, Syn. Rub. Germ. 296 = vestitus.
macrobelos, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 126 = ulmifolius.
macrocardiophyllus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 337 = thyrsiflorus ?

RUBUS:—
macrocarpus, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 7 = moluccanus.
macrocarpus, *Benth. Pl. Hartw.* 129.—Ecuador.
macrocarpus, King, ex C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 141 = Hookeri.
macrodontus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 185.—Europ.
macropetalus, Dougl. ex Hook. Fl. Bor. Am. i. 178. t. 59 = ursinus.
macropetalus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 254.—Europ.
macrophylloides, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 172 = macrophyllus.
macrophyllus, *Weihe & Nees, Rub. Germ.* 35. t. 12.—Europ.
macropodus, Ser. in DC. Prod. ii. 557 = parvifolius.
macrostachys, *P. J. Muell. in Flora*, xli. (1858) 150. —Europ.
macrostemon, Focke, Syn. Rub. Germ. 193 = pubescens, *Weihe.*
macrothyrsus, Lange, in Fl. Dan. t. 2832 = vestitus.
magdeburgensis, Maass, ex Focke, Syn. Rub. Germ. 151 = rhamnifolius.
magnificus, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 158 = thyrsiflorus.
malacophyllus, P. J. Muell. ex Focke, Syn. Rub. Germ. 237 = tomentosus, *Borckh.*
malagassus, *Focke, in Bremen, Abh.* viii. (1884) 473.—Madag.
maleficus, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 314.—Europ.
Mandonii, *Focke, in Bremen, Abh.* iv. (1874) 162.—Bolivia.
marchicus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 15.—Europ.
marginum, P. J. Muell. in Pollichia, xvi–xvii. (1859) 271 = tomentosus, *Borck.*
mascarinensis, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 139.—Ins. Mascar.
massiliensis, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 126 = ulmifolius.
Maximowiczii, *Kuntze, Method.* 64.—Japon.
maximus, *Kuntze, l. c.* 62.—Java.
maximus, Linn. ex Nym. Consp. 221 = corylifolius.
maximus, *Marss. Fl. Neuvorpomm.* 151.—Europ.
maximus, *Neum. in Bot. Notiser* (1885), 90.—Suecia.
medius, *Kuntze, Method.* 95.—Japon.
megalacanthus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 78.—Europ.
megalococeus, *Focke, in Bremen, Abh.* iv. (1874) 157. —Bolivia.
megapolitanus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 14.—Europ.
megathamnus, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 150 = tomentosus, *Borck.*
melanochlamys, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 101 = hirtus.
melanoxylon, P. J. Muell. & Wirtg. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 133 = villicaulis.
Menkei, Weihe, ex Spreng. Syst. ii. 528 = vestitus.
Menkei, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 679.—Europ.
mentitus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 324 = R. Radula, thyrsiflorus.
menyhazensis, *Simk. in Termész. Fuzetek*, ix. (1885) 42. —Europ.
Menziesii, Hook. Fl. Bor. Am. i. 179 = spectabilis.
Mercierii, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 271 = arduennensis.
meridionalis, Kern. ex Focke, Syn. Rub. Germ. 226 = tomentosus, *Borck.*
Metschii, *Focke, l. c.* 359.—Europ.
mexicanus, *Kuntze, Method*, 102.—Mexic.
micans, *Gren. & Godr. Fl. Fr.* i. 546.—Europ.
micans, Weihe, ex Lej. & Court. Comp. Fl. Belg. ii. 168 = tomentosus, *Borck.*
micrandrus, Ripart, ex Genev. Monog. Rub. ed. II. 338 = tomentosus, *Borck.*
micranthus, D. Don, Prod. Fl. Nep. 235 = lasiocarpus.
microacanthus, Kaltenb. Fl. Aach. Beckens, 263 = nessensis.
microcarpus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 25.—Europ.
microdontus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 244.—Europ.

RUBUS:—
micropetalus, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 6 = moluccanus.
microphyllus, D. Don, Prod. Fl. Nep. 234 = foliolosus.
microphyllus, *Linn. f. Suppl.* 263.—China; Japon.
microphyllus, Reichb. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 688 = Koehleri.
Mikani, Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 56 = Koehleri.
minor, Kuntze, Method. 97 = crataegifolius.
minutiflorus, *Lange, Desc. Pl. Nov. Hisp.* fasc. i. 2. t. 3. —Europ.
minutiflorus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 235 = Schleicheri.
miser, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 156. —Costa Rica.
mitis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 98. —Europ.
mitissimus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 11 = caesius.
mixtus, Ripart, ex Genev. l. c. 260 = villicaulis.
modestus, Ripart, ex Genev. Monog. Rub. ed. II. 8 = caesius.
moestus, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 378.—Europ.
mollifolius, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 267.—Europ.
mollis, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 378.—Europ.
mollis, Jungh. ex Miq. Fl. Ind. Bat. i. 1. 379 = lineatus.
mollis, J. & C. Presl, Delic. Prag. 218 = nemorosus.
moluccanus, *Linn. Sp. Pl.* 1197.—Reg. Himal.; Malaya.
monanthus, Kuntze, Method. 142 = humulifolius.
mongolicus, *Kuntze, l. c.* 92.—Mongol.
montanus, Libert, ex Lej. Fl. Spa, ii. 317 = candicans.
montanus, *Ortm. ex Opiz, Seznam*, 85.—Europ.
montanus, Wimm. Fl. Schles. ed. III. (1857) 631 = nemorosus.
montanus, Wirtg. in Flora, xxxix. (1856) 236 = carpinifolius.
monticolus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 127.—Europ.
Moorei, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 67.—Austral.
morifolius, *P. J. Muell. in Flora*, xli. (1858) 164.—Europ.
morifolius, *Siebold, ex Franch. & Sav. Enum. Pl. Jap.* i. 125.—Japon.
mortuorum ×, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxxi. (1882) 376.—Europ.
Mougeoti, Bill. ex F. Schultz, Arch. Fl. Fr. et Allem. 166 = nemorosus.
mucronatus, Bloxam, in Kirby, Fl. Leicest. 43 = hirtus.
mucronatus, *Ser. in DC. Prod.* ii. 565.—Am. bor.
mucronipetalus, *P. J. Muell. in Bonplandia*, ix. (1861) 298.—Europ.
mucronulatus, Bor. Fl. Centr. Fr. ed. III. ii. 196 = hirtus.
Muelleri, *F. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 9.—Austral.
Muelleri, Wirtg. ex Nym. Consp. 220 = serpens.
Muenteri, Marss. Fl. Neuvorpomm. 144 = rhamnifolius.
Mullerii, *Lefèv. ex P. J. Muell. in Pollichia*, xvi–xvii. (1859) 180.—Europ.
multifidus, *Boulay & Malbranche, ex Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 222, nomen.—Europ.
multiflorus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 19.—Europ.
multiflorus, *Thunb. Diss. Rub.* 7.—Hab. ?
multisetosus, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 102 = hirtus.
Mundtii, Cham. & Schlecht. in Linnaea, ii. (1827) 18 = rigidus, *Sm.*
mutabilis, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 84 = rudis.
myriacanthus, Dougl. ex Hook. Fl. Bor. Am. i. 178 = ursinus.
myriacanthus, Focke, in Bremen, Abh. ii. (1871) 467 = nemorosus.
myriadenophorus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 39 = thyrsiflorus.
myrianthus, *Baker, in Journ. Linn. Soc.* xx. (1883) 136.—Madag.
myrianthus, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxxi. (1882) 372 = pubescens, *Weihe*.

RUBUS:—
Myricae, *Focke, in Alpers, Verz. Gefaesspfl. Stade*, 27. —Europ.
mysorensis, Heyne, ex Roth, Nov. Pl. Sp. 235 = lasiocarpus.
napophilus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 220.—Europ.
natkanus, G. Don, Gen. Syst. ii. 540, sphalm. = nutkanus.
Neesii, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 227 = ulmifolius.
neglectus, P. J. Muell. ex Focke, Syn. Rub. Germ. 237 = nemorosus.
neglectus, *Peck, Rep. Reg. N. York St. Univ.* xxii. 53. —Am. bor.
nemocharis, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 191 = Sprengelii.
nemophilus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 206 = arduennensis.
nemoralis, P. J. Muell. in Flora, xli. (1858) 139 = Sprengelii.
nemoralis, *Neum. in Bot. Notiser* (1885), 91.—Suecia.
nemorensis, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 198.—Europ.
nemorivagus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 27 = corylifolius.
nemorosus, Boenn. Prod. Fl. Monast. 153 = vestitus.
nemorosus, Guenth. ex Focke, Syn. Rub. Germ. 404 = oreogeton.
nemorosus, *Hayne, Arzn. Gew.* iii. t. 10.—Europ.; Reg. Caucas.
nemorum, Ripart, ex Genev. Monog. Rub. ed. II. 235 = carpinifolius.
neo-mexicanus, *A. Gray, Pl. Wright.* ii. 55.—N. Mexic.
nessensis, *W. Hall, in Trans. Roy. Soc. Edinb.* iii. (1794) *Hist.* 21.—Europ.
neurophanes, Boul. & Cornet, ex Genev. Monog. Rub. ed. II. 247 = arduennensis.
nexuosus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 309 = nitidus.
nigellus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = fruticetorum.
nigricatus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 204.—Europ.
nilagiricus, Focke, in Bremen, Abh. v. (1877) 406 = ramosus.
nitens, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = nemorosus.
nitidus, Bell Salt, in Phytol. ii. (1845) 101 = vulgaris.
nitidus, Kaltenb. Fl. Aach. Beckens, 265 = affinis.
nitidus, Rafin. Fl. Ludov. 98 (Quid?).—Am. bor.
nitidus, *Weihe & Nees, Rub. Germ.* 19. t. 4.—Europ.
nivalis, Dougl. ex Hook. Fl. Bor. Am. i. 181 = strigosus?
niveus, *Thunb. Diss. Rub.* 9.—Hab. ?
niveus, *Wall. Cat.* n. 734; *et ex G. Don, Gen. Syst.* ii. 530.—Reg. Himal.
nobilis, *Hort. Angl. ex Regel, Gartenfl.* vi. (1857) 86. —Hab. ?
nodiflorus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 95, nomen = orthocanthus.
novus, *Kuntze, Method.* 112.—Austral.
nubigenus, *H. B. & K. Gen. et Sp.* vi. 220.—Ecuador.
nubis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 585 = R. Chamaemorus.
numidicus, *Focke, in Bremen, Abh.* iv. (1874) 175.—Afr. bor.
nutans, Vest, in Syll. Ratisb. i. (1824) 238 = nessensis.
nutans, *Wall. Cat.* n. 738.—Reg. Himal.
nutkanus, *Moç. ex Ser. in DC. Prod.* ii. 566.—Am. bor. occ.
obcuneatus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 196.—Europ.
obliquus, Bloxam, in Journ. Bot. viii. (1870) 69 = rudis.
obliquus, Wirtg. in Flora, xlii. (1859) 235 = pallidus.
oblongatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 184.—Europ.
oblongifolius, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 304 = Menkei.
obotriticus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 15.—Europ.
obovalis, Michx. Fl. Bor. Am. i. 298 = hispidus.
obovatus, Ell. Sketch, i. 570 = hispidus.
obovatus, *Pers. Syn.* ii. 52.—Am. bor.
obrosus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 234.—Europ.
obscurifrons, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 315 = Lejeunei.
obscurus, *Kaltenb. Fl. Aach. Beckens*, 281.—Europ.

RUBUS:—
obscurus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 200.—Europ.
obsectifolius, P. J. Muell. ex Lamotte, Prod. Fl. Centr. Fr. i. 254, nomen = R. Radula.
obtruncatus, *P. J. Muell. in Flora*, xli. (1858) 152.—Europ.
obtusangulus, Gremli, Excurs. Schw. ed. II. 161 = arduennensis.
obtusatus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 95 = villicaulis.
obtusiflorus, Gremli, Excurs. Schw. ed. IV. 144 = foliosus.
obtusifolius, Pritz. Ic. Ind. ii. 247, sphalm. = Rumex obtusifolius.
obtusifolius, Weihe, ex Focke, Syn. Rub. Germ. 291 = vestitus.
obtusifolius, Willd. Berl. Baumz. ed. II. 409 = ulmifolius.
obvallatus, Boul. & Gillot, ex Genev. Monog. Rub. ed. II. 175 = vestitus.
occidentalis, Boland. Cat. 11, nomen = leucodermis.
occidentalis, Georgi, Beschr. Russ. Reich. III. iv. 1029 = R. Idaeus.
occidentalis, *Linn. Sp. Pl.* 493.—Am. bor.
occiduus, Boul. & Bouv. ex Genev. Monog. Rub. ed. II. 236 = carpinifolius.
ochlanthus, Hance, in Journ. Bot. xx. (1882) 260 = Lambertianus.
odoratus, *Linn. Sp. Pl.* 494.—Am. bor.
offensus, *P. J. Muell. in Bonplandia*, ix. (1861) 286. —Europ.
oigocladus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 134.—Europ.
Oldhami, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 34 = pungens.
oleifluus, *Steud. Nom.* ed. II. ii. 478, nomen.—Reg. Argent.
oligacanthus, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 178 = nemorosus.
oliganthos, *P. J. Muell. in Bonplandia*, ix. (1861) 287.—Europ.
Olivieri, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 35 = corchorifolius.
omalodontos, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 354 = Koehleri.
omissus, Ripart, ex Genev. Monog. Rub. ed. II. 252 = arduennensis.
opacus, Bertram, Fl. Braunschw. ed. II. 74 = Bertramii.
opacus ×, Focke, in Alpers, Verz. Gefaesspfl. Stade, 25 = plicatus.
opulentus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 136.—Europ.
opulifolius, *Bertol. Misc. Bot.* xxii. 16. t. 4.—Reg. Himal.
orbiculatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 209.—Europ.
orbifolius, Boul. & Letendre, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = bifrons.
oreades, P. J. Muell. & Wirtg. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 89 = orthocanthus.
oreogeton, *Focke, Syn. Rub. Germ.* 404.—Europ.
oreophilus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 227.—Europ.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 342.—Bras.
orgyalis, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 221, nomen.—Europ.
orthacanthus, Wimm. Fl. Schles. ed. III. (1857) 626 = seq.
orthoacanthus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 104 = seq.
orthocanthus, *Wimm. in Flora*, xl. (1857) 77.—Europ.
Ortmanni, Opiz, in André, Oekon. Neuigk. i. (1835); et in Lotos, iv. (1854) 70 = montanus, *Ortm.*
ovalifolius, Weihe, ex Focke, Syn. Rub. Germ. 263 = silvaticus.
oxyacanthus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 131.—Europ.
oxyphyllus, Wall. Cat. n. 7110 = acuminatus.
pacificus, *Hance, in Journ. Bot.* xii. (1874) 259.—China.
pallens, Weihe, ex Focke, Syn. Rub. Germ. 340 = scaber.
pallescens, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 248 = ulmifolius.
pallescens, Weihe, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 684 = scaber.

RUBUS:—
pallidiflorus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 187.—Europ.
pallidus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 103 = foliosus.
pallidus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 682.—Europ.
palmatoides, *Kuntze, Method.* 93, 96.—China?
palmatus, Thunb. Fl. Jap. 217 = microphyllus.
paniculatus, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 140 = rosaefolius.
paniculatus, Moon, Cat. Pl. Ceyl. 40 = ellipticus.
paniculatus, Roxb. Hort. Beng. 39; Fl. Ind. ii. 518 = parvifolius.
paniculatus, Schlecht. ex Link, Enum. Hort. Berol. ii. 61 = fruticosus.
paniculatus, *Sm. in Rees, Cyclop.* xxx. n. 41.—Reg. Himal.
pannosus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 304 = Menkei.
panormitanus, *Tineo, Pl. Rar. Sicil.* 24.—Sicilia.
Pappei, Eckl. & Zeyh. Enum. 263 = rosaefolius.
papulosus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 142.—Europ.
paradoxus, S. Moore, in Journ. Bot. xvi. (1878) 132 = leucanthus.
Parkeri, *Hance, l. c.* xx. (1882) 260.—China.
parviflorus, Nutt. Gen. Am. i. 308 = nutkanus.
parvifolius, *Linn. Sp. Pl.* 1197.—Malaya; China; Austral.
parvifolius, Moon, Cat. Pl. Ceyl. 40 = lasiocarpus?
parvifolius, Sm. in Rees, Cyclop. xxx. n. 21 = foliolosus.
parvifolius, Walt. Fl. Carol. 149 = cuneifolius, *Pursh.*
parvulus, Genev. in Mém. Soc. Acad. Maine-et Loire, viii. (1860) 74 = caesius.
parvus, *J. Buch. in Trans. N. Z. Inst.* vi. (1874) 243.—N. Zel.
patens, E. Merc. in Reut. Cat. Pl. Vasc. Genèv. ed. II. 265 = caesius.
patulus, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 97 = ulmifolius.
pauciflorus, *Baker, in Journ. Linn. Soc.* xx. (1883) 136.—Madag.
pauciflorus, Wall. ex Lindl. in Bot. Reg. t. 854 = lasiocarpus.
paucistamineus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 96.—Europ.
pectinelloides, *Kuntze, Method.* 79, 107.—Japon.
pectinellus, *Maxim. in Bull. Acad. Pétersb.* xvii. (1871) 147.—Japon.
pedatifolius, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, viii. (1860) 71, 93; x. (1861) 35.—Europ.
pedatus, Banks & Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 33 = grandifolius.
pedatus, *Sm. Pl. Ic. Ined.* t. 63.—Am. bor. occ.
pedunculatus, Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 131 = ulmifolius.
pedunculatus, Steud. Nom. ed. II. ii. 478 = seq.
pedunculosus, D. Don, Prod. Fl. Nep. 234 = niveus, *Wall.*
pellitus, Ripart, ex Genev. in Mem. Soc. Acad. Maine-et-Loire, xxiv. (1868) 297 = tomentosus, *Borck.*
peltaefolius, *Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 107.—Europ.
peltatus, *Maxim. in Bull. Acad. Pétersb.* xvii. (1871) 154.—Japon.
pendulinus, P. J. Muell. in Bonplandia, ix. (1861) 290 = hirtus.
peniculus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = nemorosus.
pensilvanicus, Poir. Encyc. vi. 246 = strigosus.
pentagonus, Wall. Cat. n. 731 = alpestris.
pentaphyllos, *P. J. Muell. in Bonplandia*, ix. (1861) 289.—Europ.
perarmatus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
permiscibilis, P. J. Muell. ex Focke, Syn. Rub. Germ. 236 = tomentosus, *Borck.*
perplexus, P. J. Muell. ex Focke, l. c. 375 = hirtus.
persicinus, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 137; Nov. Pl. Sp. Decas, iii. 14 = thyrsoides.
persicus, *Boiss. Fl. Orient.* ii. 693.—Persia.
petiolulatus, Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
Petitianus, *A. Rich. Tent. Fl. Abyss.* i. 259.—Abyss.
petraeus, Koehl. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 688 = Koehleri.
petraeus, Walp. Rep. ii. 19, sphalm. = Prunus Padus.

RUBUS:—
philyrophyllus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 190.—Europ.
phoenicolasius, *Maxim. in Bull. Acad. Pétersb.* xvii. (1871) 160.—Japon.
phyllophorus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 178 = glandulosus.
phyllostachys, P. J. Muell. in Flora, xli. (1858) 133 = thyrsoideus.
piletocaulon, P. J. Muell. in Bonplandia, ix. (1861) 281 = vestitus.
piletostachys, Gremli, Beitr. Fl. Schw. 43 = Gremlii.
piletostachys, Gren. & Godr. Fl. Fr. i. 548 = macrophyllus.
piletosus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 278 = villicaulis.
pilocalyx, Kuntze, Method. 77 = moluccanus.
pilocarpus, Gremli, Beitr. Fl. Schw. 42 = vestitus.
pilocephalus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 126.—Europ.
pinnatus, D. Don, Prod. Fl. Nep. 234 = lasiocarpus.
pinnatus, Willd. Sp. Pl. ii. 1081 = rosaefolius.
pistillatus, Sm. Exot. Bot. ii. 53. t. 86 = arcticus.
platyacanthus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 86.—Europ.
platycephalus, *Focke, Syn. Rub. Germ.* 329.—Europ.
platypetalus, *Timb. & P. J. Muell. ex Martr. Fl. Tarn*, 213.—Europ.
platyphyllos, Weihe, ex Focke, Syn. Rub. Germ. 210, 215 = villicaulis, macrophyllus.
platyphyllus, C. Koch, in Linnaea, xvi. (1842) 348 = glandulosus.
platyphyllus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 217.—Europ.
plicatus, C. A. Mey. Verz. Pfl. Cauc. 167 = persicus.
plicatus, *Weihe & Nees, Rub. Germ.* 15. t. 1.—Europ.
plinthostylus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 108 = Sprengelii.
podhradiensis ×, Holuby, in Oestr. Bot. Zeitschr. xxiii. (1873) 374.—Europ.
podophyllos, P. J. Muell. in Bonplandia, ix. (1861) 281 = hirtus.
poliophyllus, Kuntze, Method. 68 = moluccanus.
polyacanthus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 95 = Bayeri.
polyandrus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 26; ed. II. (1827) 84 (Quid?).—Java.
polyanthemus, *Lindeb. in Bot. Notiser* (1883), 105.—Suecia.
polyanthus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 96 = tomentosus, *Borck.*
polycarpus, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 313.—Europ.
polymorphus, *Host, Fl. Austr.* ii. 29.—Europ.
polymorphus, *Schimp. & Spenn. in Spenn. Fl. Friburg.* iii. 744.—Europ.
polytrichus, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 102 = hirtus.
populosus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 142.—Europ.
porphyracanthos, Focke, Syn. Rub. Germ. 148 = rhamnifolius.
praealtus, Lefèv. ex Genev. Monog. Rub. ed. II. 237 = vulgaris.
praecox, Kern. in Ber. Naturw. Ver. Innsbruck, ii. (1871) 124; Nov. Pl. Sp. Decas, iii. 1 = sulcatus.
praecox, Weihe, ex Reichb. Fl. Germ. Excurs. 600 = nitidus.
praeruptorum, *Boul. ex Bab. in Journ. Bot.* xvi. (1878) 143.—Europ.
praestabilis, P. J. Muell. ex Focke, Syn. Rub. Germ. 196 = amiantinus.
praetermissus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 208 = arduennensis.
praetervisus, Ripart, ex Genev. l. c. 247 = pubescens, *Weihe.*
prasinifolius, Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 131 = ulmifolius.
prasinus, Focke, in Bremen, Abh. i. (1867) 302 = orthocanthus.
prehensibilis, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 453.—Europ.
Preslii, Weitenweb. in Opiz, Natural. 346 = plicatus.
prionodontus, *Lefèv. ex P. J. Muell. in Pollichia*, xvi–xvii. (1859) 117.—Europ.
procerus, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 209 = thyrsoideus.

RUBUS:—
procumbens, Muhl. Cat. ed. I. 50; ed. II. 52 = canadensis.
prolectibilis, Ripart & Genev. in Genev. Monog. Rub. ed. II. 315 = villicaulis.
prolixatus, Ripart & Genev. l. c. 87 = tomentosus, *Borck.*
prolongatus, Boul. & Letendre, ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 221 = candefactus.
pronatiflorus, Timb. & P. J. Muell. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 128 = ulmifolius.
propinquus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 88 = ulmifolius.
propinquus, Richards. in Frankl. 1st Journ. ed. II. App. 19 = arcticus.
prostratus, DC. ex Steud. Nom. ed. I. 706 = tomentosus, *Borck.*
prostratus, Seidel, ex Steud. l. c. = hispidus.
Provostii, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 21 = caesius.
proximellus, Ripart, ex Genev. Monog. Rub. ed. II. 330 = tomentosus, *Borck.*
pruinosus, Arrhen. Rub. Suec. Monog. 15 = corylifolius.
pruinosus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
pruinosus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 587.—Java.
psammophilus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 47 = nemorosus.
Pseudo-americanus, Kuntze, Method. 90, 96 = crataegifolius.
Pseudo-arcticus, Kuntze, l. c. 151 = arcticus.
Pseudo-caesius, Weihe & Nees, Rub. Germ. 103. t. 46 B = caesius.
Pseudo-idaeus, P. J. Muell. in Flora, xli. (1858) 129 = nessensis.
Pseudo-idaeus, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 37 = R. Idaeus?
Pseudo-ideus, Weihe, ex Boenn. Prod. Fl. Monast. 151 = caesius.
Pseudo-incisus, Kuntze, Method. 93 = incisus.
pseudopsis, Gremli, ex Focke, Syn. Rub. Germ. 394 = orthocanthus.
Pseudo-radula, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 379.—Europ.
Pseudo-tiliaceus, Kuntze, Method. 60 = paniculatus?
Pseudo-triflorus, Kuntze, l. c. 129 = humulifolius.
pubescens, Rafin. in Med. Repos. N. York, Ser. III. ii. (1811) 333 = saxatilis.
pubescens, *Vest, ex Tratt. Ros.* iii. 34.—Europ.
pubescens, *Weihe, ex Boenn. Prod. Fl. Monast.* 152.—Europ.
pubicaulis, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 266 = nemorosus.
pubinervis, *Blume, Bijdr.* 1110.—Java.
Pugeti, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 131 = ulmifolius.
pulchellus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 97.—Europ.
pulcher, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 148.—Europ.
pulcherrimus, Hook. Ic. Pl. t. 729–730 = lineatus.
pulcherrimus, Neum. in Oefvers. Vet. Akad. Foerhandl. Stockh. (1883) n. viii. 65 = polyanthemus.
pumilus, *Focke, in Bremen, Abh.* iv. (1874) 155.—Mexic.
punctulatus, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 235 = thyrsoideus.
pungens, *Cambess. in Jacquem. Voy. Bot.* 48. t. 59.—Reg. Himal.; Japon.
Purchasii, *Bloxam, in Journ. Bot.* xvi. (1878) 208.—Europ.
purpureistylus, *P. J. Muell. in Flora*, xli. (1858) 171—Europ.
purpureus, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 98 = parvifolius.
purpureus, Holuby, in Oestr. Bot. Zeitschr. xxiii. (1873) 374 = thyrsoideus.
pusillus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 14 = caesius.
pycnanthus, Focke, in Bremen, Abh. iv. (1874) 196 = Lambertianus.
pycnanthus, Genev. Monog. Rub. ed. II. 210 = Sprengelii.
pycnostachys, P. J. Muell. in Pollichia, xvi–xvii. (1859) 101 = tomentosus, *Borck.*

RUBUS:—
pycnostylus, *P. J. Muell. in Bonplandia*, ix. (1861) 293.—Europ.
pygmaeopsis, Focke, Syn. Rub. Germ. 364 = Schleicheri.
pygmaeus, Bab. Man. Brit. Bot. ed. V. 104 = praeruptorum.
pygmaeus, Bor. Fl. Centr. Fr. ed. III. ii. 190 = rosaceus.
pygmaeus, Metsch, in Linnaea, xxviii. (1856) 178 = Bayeri.
pygmaeus, Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 687 = Koehleri.
pygmaeus, Wirtg. Fl. Preuss. Rheinprov. 161, partim = Schleicheri.
pyramidalis, *Bab. Man. Brit. Bot.* ed. III. 101.—Europ.
pyramidalis, Kaltenb. Fl. Aach. Beckens, 275 = vestitus.
pyramidalis, Quest. ex Bab. Brit. Rub. 235 = saltuum.
pyramidatus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 192 = Sprengelii.
pyramidatus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 291 = vulgaris.
pyrifolius, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. ii. 328 = Hamiltoni.
pyrifolius, Hook. f. & Thoms. ex Hook. f. l. c. = hexagynus.
pyrifolius, *Sm. Pl. Ic. Ined.* t. 61.—Java.
quadraticus, *P. J. Muell. in Pollichia*, xvi.–xvii. (1859) 182.—Europ.
Quartinianus, *A. Rich. Tent. Fl. Abyss.* i. 257.—Abyss.
Questierii, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 120.—Europ.
quinquelobus, Stokes, Bot. Mat. Med. iii. 142 = odoratus.
racemigerus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 128 = R. Radula.
racemosus, Genev. in Mém. Soc. Acad. Maine-et-Loire, x. (1861) 34 = macrophyllus.
racemosus, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii. 519.—Ind. or.
Raddeanus, *Focke, in Bremen, Abh.* iv. (1874) 182.—Reg. Caucas.
radicans, *Cav. Ic.* v. 7.—Chili.
radicans, Focke, in Bremen, Abh. v. (1877) 407 = Fockeanus?
Radula, *Weihe, in Boenn. Prod. Fl. Monast.* 152.—Europ.
ramosus, *Bloxam, in Journ. Bot.* ix. (1871) 330.—Europ.
ramosus, Neum. in Oefvers. Vet. Akad. Foerhandl. Stockh. xl. (1883) n. VIII. 82 = caesius.
rariflorus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 125.—Europ.
Ravi, Weihe, ex Reichb. Fl. Germ. Excurs. 602 = vulgaris.
reclinatus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 37.—Europ.
rectangulatus, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 14.—Europ.
rectangulatus, Maass, ex Focke, Syn. Rub. Germ. 209 = villicaulis.
rectangulifolius, *Kuntze, Method.* 60.—China.
rectispinus ×, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 239.—Europ.
reduncus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 211 = arduennensis.
reflexus, Ker-Gawl. in Bot. Reg. t. 461 = moluccanus.
Reichenbachii, Koehl. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 685 = villicaulis.
Reinwardtii, *Kuntze, Method.* 56.—Java.
remotus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 126 = brachyandrus.
repens, Gueldenst. Reisen, ii. 4, 197; ex Ledeb. Fl. Ross. ii. 72 (Quid?).—Ross.
reptans, *Herb. Par. ex Kuntze, Method.* 107.—Mexic.
reticulatus, *Kern. in Ber. Naturw. Ver. Innsbruck*, 165; *Nov. Pl. Sp. Decas*, iii. 42.—Europ.
reticulatus, *Wall. Cat.* n. 747.—Reg. Himal.
retrodentatus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 168.—Europ.
retrogressus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 20 = caesius.
retrospiciens, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
Reussii, Holuby, in Oestr. Bot. Zeitschr. xxv. (1875) 314 = Guentheri.
Reuteri, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 272.—Europ.
rhamnifolius, Bill. ex Genev. Monog. Rub. ed. II. 262 = ulmifolius.

RUBUS:—
rhamnifolius, *Weihe & Nees, Rub. Germ.* 22. t. 6.—Europ.
rhamnifolius, Wirtg. Fl. Preuss. Rheinprov. 150 = bifrons.
rhenanus, *P. J. Muell. in Flora*, xli. (1858) 184.—Europ.
rhodacantha, E. Mey. in Drège, Zwei Pfl. Docum. 217 = Ludwigii.
rhombifolius, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 233 = vulgaris.
rhombifolius, Weihe, ex Boenn. Prod. Fl. Monast. 151 = villicaulis.
rhombophyllus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 175.—Europ.
ribesifolius, Sieber, ex Hook. f. Fl. Tasm. i. 112 = parvifolius.
ribifolius, Sieb. & Zucc. Fl. Jap. i. 127 = incisus.
ribifolius, Willd. ex Ser. in DC. Prod. ii. 559 = spectabilis.
Ribis-folius, Gilib. Fl. Lituan. ii. 244 = R. Chamaemorus.
rigidatus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 127.—Europ.
rigidus, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 274.—Europ.
rigidus, *Sm. in Rees, Cycl.* xxx. n. 5.—Afr. austr.
Ripartii, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 300 = tomentosus, *Borck.*
rivalis, Genev. l. c. viii. (1860) 76 = caesius.
rivularis, P. J. Muell. in Flora, xlii. (1859) 237 = serpens.
robustus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 87 = pubescens, *Weihe.*
robustus, *Presl, Epim. Bot.* 196.—Peruv.
robustus, Wirtg. ex Nym. Consp. 218 = tomentosus, *Borck.*
Roesli, Regel, Gartenfl. (1875) 227; et in Act. Hort. Petrop. iii. II. (1875) 285 = deliciosus.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 171.—Ins. Philipp.
roridus, *Lindl. Bot. Reg.* t. 1607.—Madag.
rorulentus, *Haldscy, in Kern. Sched. Fl. Exsicc.* iii. 39.—Europ.
rosaceus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 685.—Europ.
rosaeflorus, Hook. Ic. Pl. t. 46 = roseus.
rosaeflorus, Roxb. Hort. Beng. 39; Fl. Ind. ii. 519 = lasiocarpus.
rosaefolius, *Sm. Pl. Ic. Ined.* iii. t. 60.—As. trop.
rosaefolius, Vidal, Sinops. t. 46. f. D = fraxinifolius.
roseiflorus, P. J. Muell. in Flora, xli. (1858) 179 = caesius.
roseus, *Poir. Encyc.* vi. 245.—Peruv.
rosifolius, Stokes, Bot. Mat. Med. iii. 147 = rosaefolius.
rostochiensis, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 18.—Europ.
rosulans, Kuntze, Method. 65, 78 = moluccanus.
rosulentus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 76 = plicatus.
Rothii, Focke, in Bremen, Abh. ii. (1871) 461 = vestitus.
rotundatus, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 177 = Sprengelii.
rotundifolius, *Bloxam, in Kirby, Fl. Leicest.* 39.—Europ.
rotundifolius, *P. J. Muell. in Flora*, xli. (1858) 178.—Europ.
rotundifolius, *Reinw. ex Miq. Fl. Ind. Bat.* i. 1. 384.—Java.
rotundifolius, Royle, Illustr. Bot. Himal. 203, nomen = lasiocarpus.
rotundifolius, Wall. Cat. n. 730 = ellipticus.
rotundifolius, Weihe, ex Focke, Syn. Rub. Germ. 291 = vestitus.
rotundipetalus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 82 = candicans.
Roxburghianus, Wall. Cat. n. 732 = parvifolius.
Roylei, Klotzsch, in Bot. Ergeb. Waldem. Reise, 154. t. 6 = foliolosus.
rubellus, *P. J. Muell. in Flora*, xli. (1858) 156.—Europ.
rubens, Weihe, ex Reichb. Fl. Germ. Excurs. 604 = Menkei.
rubeolus, Reichb. ex Nym. Consp. 216 = affinis.
rubeolus, Weihe, ex Reichb. Fl. Germ. Excurs. 600 = rhamnifolius.

RUBUS:—
ruber, *Gilib. Fl. Lituan.* ii. 243 = saxatilis.
rubescens, Lej. Fl. Spa, ii. 340 = rosaceus.
rubicaulis, Steud. Nom. ed. II. ii. 479, sphalm. = *rubricaulis.*
rubicolor, *Bloxam, in Syme, Engl. Bot.* ed. III. iii. 180.—Europ.
rubicundus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 310 = obscurus, *Kaltenb.*
rubifolius, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 19 = trifidus.
rubiginosus, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = rufescens.
rubiginosus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 207.—Europ.
rubricaulis, Weihe, ex Weihe & Nees, Rub. Germ. 69 = apiculatus.
rubriflorus, *Kuntze, Method.* 102.—Mexic.
ruderalis, Chab. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 272 = villicaulis.
rudicaulis, *Genev. l. c.* viii. (1860) 89.—Europ.
rudiformis, Genev. l. c. xxiv. (1868) 151 = rudis.
rudis, Bab. Brit. Rub. 189 = R. Radula.
rudis, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 687.—Europ.
rufescens, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 152.—Europ.
rugosus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 234 = moluccanus.
rugosus, G. Jens. ex Lange, Fl. Dan. t. 2905 = slesvicensis.
rugosus, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 149 = formosensis.
rugosus, Sm. in Rees, Cycl. xxx. n. 34 = moluccanus.
Ruizii, *Focke, in Bremen, Abh.* iv. (1874) 162.—Peruv.
rupestris, P. J. Muell. in Flora, xli. (1858) 180 = nemorosus.
russeicaulis, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 222, nomen.—Europ.
rusticanus, E. Merc. in Reut. Cat. Pl. Vasc. Genèv. ed. II. 279 = ulmifolius.
sabaudus, Focke, ex Gremli, Excurs. Schw. ed. II. 161 = ulmifolius.
sacer, Georgi, Beschr. Russ. Reich. III. iv. 1032 = ulmifolius.
saevus, Holuby, in Oestr. Bot. Zeitschr. xxiii. (1873) 383 = Koehleri.
salisburgensis, *Focke, Syn. Rub. Germ.* 280.—Europ.
Salteri, Bab. in Ann. & Mag. Nat. Hist. Ser. I. xvii. (1846) 172 = silvaticus.
saltuum, *Focke, in Gremli, Beitr. Fl. Schw.* 30.—Europ.
sanctus, Kuntze, Taschen-Fl. Leipz. 262 = caesius.
sanctus, Schreb. Ic. et Desc. Pl. Dec. 15. t. 8 = ulmifolius.
sanguineus, Frivald. in Flora, xviii. (1835) 334 = ulmifolius.
sanguinolentus, *Link, Enum. Hort. Berol.* ii. 61.—Ins. Borbon.
sapidus, *Schlecht. in Linnaea*, xiii. (1839) 269.—Mexic.
Sauli, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 33 = pubescens, *Weihe.*
saxatilis, Bigel. Fl. Bost. ed. II. 201 = triflorus.
saxatilis, *Linn. Sp. Pl.* 494.—Reg. bor. temp. et arct.
saxicolus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 202 = hirtus.
scaber, Kaltenb. Fl. Aach. Beckens, 289 = hirtus.
scaber, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 683.—Europ.
scabridus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 158.—Europ.
scabrifolius, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 96 = Bayeri.
scabripes, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 69, 81 = Sprengelii.
scabrosus, P. J. Muell. in Flora, xli. (1858) 185 = nemorosus.
scandens, Juss. ex Poir. Encyc. vi. 237 = saxatilis.
scandens, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 154.—Mexic.
scanicus, *Aresch. Skånes Fl.* ed. II; *et in Bot. Notiser* (1881), 158.—Europ.
Schefferi, *Focke, in Bremen, Abh.* viii. (1884) 472.—Java.
Schiedeanus, *Steud. Nom.* ed. II. ii. 479.—Mexic.
schistogenes, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 238 = nemorosus.

RUBUS:—
Schlechtendahlii, Weihe, ex Link, Enum. Hort. Berol. ii. 62 = macrophyllus.
Schleicheri, Holuby, ex Focke, Syn. Rub. Germ. 390 = orthocanthus.
Schleicheri, *Weihe, ex Tratt. Ros.* iii. 22.—Europ.
Schlickumi, Wirtg. in Flora, xlii. (1859) 235 = thyrsoideus.
Schmelleri, *Holuby, in Oestr. Bot. Zeitschr.* xxiii. (1873) 377.—Europ.
schmidelioides, A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 245 = australis.
Schomburgkii, *Klotzsch, in Schomb. Faun. et Fl. Brit. Gui.* 1102, nomen.—Guian.
Schottii, *Pohl, ex Focke, in Bremen, Abh.* iv. (1874) 157.—Bras.
Schultzii, Ripart, ex P. J. Muell. in Pollichia, xvi–xvii. (1859) 289 = tomentosus, *Borck.*
Schummelii, *Weihe, in Wimm. & Grab. Fl. Siles.* (1829) i. II. 47.—Europ.
Schwarzeri ×, Holuby, in Oestr. Bot. Zeitschr. xxiii. (1873) 376.—Europ.
sciaphilus, *Lange, in Fl. Dan.* t. 3026.—Europ.
sciophilus, *Lefèv. ex P. J. Muell. in Pollichia*, xvi–xvii. (1859) 205.—Europ.
scophilus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxiv. (1868) 178.—Europ.
Sellowii, *Cham. & Schlecht. in Linnaea*, ii. (1827) 15.—Arg. Argent.
semipellitus, Ripart, ex Genev. Monog. Rub. ed. II. 336 = tomentosus, *Borck.*
semivestitus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 197.—Europ.
semiviridis, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 131 = ulmifolius.
sempervirens, Bigel. Fl. Bost. ed. II. 201 = hispidus.
Sendtneri, *Prog. in Ber. Bot. Ver. Landshut*, viii. (1882) 109.—Europ.
senticetorum, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 255.—Europ.
senticosus, *Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles.* (1829) i. II. 51.—Europ.
separinus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 71, 90, 111 = vestitus.
sepicolus, Lefèv. & P. J. Muell. in Pollichia, xvi–xvii. (1859) 273 = nemorosus.
sepincolus, Boul. ex Lamotte, Prod. Fl. Centr. Fr. i. 255, nomen = R. Radula.
sepium, Ripart, ex Genev. Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 53 = nemorosus.
septicolus, Arrond. Cat. Pl. Morb. 33 = *sepicolus.*
septorum, P. J. Muell. in Pollichia, xvi–xvii. (1859) 271 = macrophyllus.
sericatus, *P. J. Muell. & Lefèv. l. c.* 125.—Europ.
sericatus, *P. J. Muell. in Flora*, xli. (1858) 184.—Europ.
sericeus, Gilib. Fl. Lituan. ii. 241 = R. Idaeus.
sericophyllus, P. J. Muell. & Wirtg. ex Focke, Syn. Rub. Germ. 240 = tomentosus, *Borck.*
serpens, Gren. & Godr. Fl. Fr. i. 538 = orthocanthus.
serpens, *Weihe, in Lej. & Court. Comp. Fl. Belg.* ii. 172.—Europ.
serratifolius, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 237.—Europ.
serratus, J. F. Gmel. Syst. 856 = villosus.
serriculatus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 250 = ulmifolius.
sertiflorus, P. J. Muell. ex Genev. l. c. 96 = rosaceus.
sessilifolius, Miq. ex Hook. f. Fl. Brit. Ind. ii. 336 = ellipticus.
setigerus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 222.—Europ.
setosolignosus, Kuntze, Method. 69 = moluccanus.
setosus, Bigel. Fl. Bost. ed. II. 198 = hispidus.
setulosus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 199.—Europ.
siculus, Presl, Fl. Sic. p. xxiv = tomentosus, *Borck.*
Sieboldii, *Blume, Bijdr.* 1110.—Java.
siekensis, Banning, in Focke, Syn. Rub. Germ. 271 = conothyrsos.
sikkimensis, *Hook. f. Fl. Brit. Ind.* ii. 336.—Reg. Himal.
sikkimensis, Kuntze, ex Hook. f. l. c. 341 = rosaefolius.
silesiacus, Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 53 = thyrsoideus.

RUBUS:—
silvaticus, *Weihe & Nees, Rub. Germ.* 41. t. 15.—Europ.
silvaticus, Wirtg. ex Focke, Syn. Rub. Germ. 209 = villicaulis.
silvestris, Focke, l. c. 330 = foliosus.
silvicolus, C. Muell. in Walp. Ann. vii. 855 = sylvicolus.
similatus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 114.—Europ.
similis, Kuntze, Method. 94 = microphyllus.
sinensis, Hort. ex Ser. in DC. Prod. ii. 556 = rosaefolius.
slesvicensis, *Lange, in Fl. Dan.* t. 2905.—Europ.
sorbifolius, Maxim. in Bull. Acad. Pétersb. xvii. (1871) 158 = rosaefolius.
sordidus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 98.—Europ.
sparsipilus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 73 = hirtus.
speciosus, P. J. Muell. in Flora, xli. (1858) 135 = bifrons.
spectabilis, James, Cat. 182 = deliciosus.
spectabilis, E. Merc. in Reut. Cat. Pl. Vasc. Genèv. ed. II. 291 = arduennensis.
spectabilis, *Pursh, Fl. Am. Sept.* i. 348. t. 16.—Am. bor.
speluncarum, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 453.—Europ.
sphenoides, Focke, ex Gremli, Excurs. Schw. ed. II. 161 = tomentosus, *Borck.*
sphenoiphyllos, G. Braun, ex Focke, Syn. Rub. Germ. 270 = glaucovirens.
sphenophyllus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 215.—Europ.
spicatus, Lefèv. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 57 = pubescens, *Weihe.*
spicatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 26 = elongatus? *Sm.*
spiculatus, Boul. & Bouv. ex Lamotte, Prod. Fl. Centr. Fr. i. 248 = nemorosus.
spidnostachys, Ripart & Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 275 = villicaulis.
Spina-curva, Boul. & Gillot, ex Genev. Monog. Rub. ed. II. 317 = villicaulis.
spinetorum, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 281.—Europ.
spinosissimus, P. J. Muell. in Flora, xli. (1858) 177 = nemorosus.
spinosissimus, Schur, Enum. Pl. Transs. 186 = heterophyllus, *Host.*
spinulicaulis, *P. J. Muell. in Bonplandia*, ix. (1861) 305.—Europ.
spinuliferus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 213.—Europ.
spinulifolius, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 127.—Europ.
spinulosus, *P. J. Muell. in Flora*, xli. (1858) 169.—Europ.
splendens, Chab. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 266 = villicaulis.
splendens, *Genev. l. c.* x. (1861) 36.—Europ.
splendidus, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 130.—Europ.
spoliatus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 48 = carpinifolius.
Sprengelii, *Weihe, in Flora*, ii. (1819) 18.—Europ.
squalidus, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, x. (1861) 20.—Europ.
stellatus, *Sm. Pl. Ic. Ined.* t. 64.—Am. bor.
stelliger, Turcz. ex Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 224 = nemorosus.
stellinus, Kuntze, ex Focke, Syn. Rub. Germ. 434 = tomentosus, *Borck.*
stenoacanthus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 110.—Europ.
stenopetalus, Cham. in Choris, Voy. Pittor. (Kamtchat). 10 = spectabilis.
stenopetalus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 94.—Europ.
stenophyllus, P. J. Muell. l. c. 98 = carpinifolius.
stenoplos, Focke, Syn. Rub. Germ. 434 = rhamnifolius.
stenothyrsos, *Holuby, in Oestr. Bot. Zeitschr.* xxv. (1875) 315.—Europ.
stereacanthos, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 189 = macrophyllus.
sterilis, Kuntze, Method. 69, 78 = moluccanus.

RUBUS:—
Steudneri, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 669.—Afr. trop.
stictocalyx, *P. J. Muell. in Bonplandia*, ix. (1861) 285.—Europ.
stipularis, *Benth. Pl. Hartw.* 173.—N. Granat.
striatus, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 128 = ulmifolius.
stricticaulis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 126.—Europ.
strigosus, *Michx. Fl. Bor. Am.* i. 297.—Am. bor.
suaveolens, Eichw. Skizze, 162 (Quid?).—Ross.
suaveolens, *Thunb. Diss. Rub.* 6.—Hab.?
suavifolius, Gremli, Beitr. Fl. Schw. 35 = vestitus.
subcanus, P. J. Muell. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 45 = Sprengelii.
subemarginatus, P. J. Muell. & Timb. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 127 = ulmifolius.
suberectus, Anders. ex Sm. Engl. Bot. t. 2572; et in Trans. Linn. Soc. xi. (1815) 218 = nessensis.
suberectus, Baer, in Flora, iv. (1821) 411 = Sprengelii.
suberectus, Hook. Fl. Bor. Am. i. 179 = villosus.
Suberti, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 281 = ulmifolius.
subfulgens, Prog. in Ber. Bot. Ver. Landshut, viii. (1882) 109 = Sendtneri.
subherbaceus, *Kuntze, Method.* 83.—Hab.?
subinermis, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 162.—Europ.
subinermis, Ruprecht, in Beitr. Pfl. Russ. Reich. iv. 66 = nessensis.
subintegrifolius, Kuntze, Method. 113 = humulifolius.
sublenis, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 131 = ulmifolius.
sublustris, Lees, in Steele, Handb. Field Bot. 54 = corylifolius.
subramiflorus, Kuntze, Method. 54, 79 = Sieboldii.
subvelutinus, Boul. & Ripart, ex Genev. Monog. Rub. ed. II. 319 = villicaulis.
sulcatus, Koehl. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 669 = fruticosus.
sulcatus, *Vest, ex Tratt. Ros.* iii. 42.—Europ.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* 307.—Sumatra.
sundaicus, *Blume, Bijdr.* 1111.—Java.
Swinhoii, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 211.—Ins. Formosa; China.
sylvarum, Ripart, ex Genev. Monog. Rub. ed. II. 196 = macrophyllus.
sylvaticus, Leight. ex Focke, Syn. Rub. Germ. 206 = villicaulis.
sylvestris, Kaltenb. Fl. Aach. Beckens, 288 = foliosus.
sylvicolus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 113.—Europ.
taeniarum, Lindeb. Nov. Fl. Scand. 5. t. 1 = infestus, *Weihe.*
tagallus, *Cham. & Schlecht. in Linnaea*, ii. (1827) 9.—Ins. Luzon; China.
Tanakae, *Kuntze, Method.* 94.—Japon.
tardiflorus, Focke, ex Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 96 = foliosus.
tauricus, Schlecht. ex Ledeb. Fl. Ross. ii. 68 = fruticosus.
tenacellus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 52 = ulmifolius.
tenellus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 248.—Europ.
tenuiarmatus, Lees, in Phytol. (1853) 818 = corylifolius.
tenuiflorus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 205 = thyrsoideus.
tenuiglandulosus, Gremli, in Oestr. Bot. Zeitschr. xxi. (1871) 97 = foliosus.
tenuis, Bell Salt, in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 305 = caesius.
tenuis, *Kuntze, Method.* 154.—Europ.
tephracanthos, Boul. ex Malbranche, in Bull. Soc. Bot. Fr. xxvi. (1879) 130 = ulmifolius.
tephrodes, *Hance, in Journ. Bot.* xii. (1874) 260.—China.
tereticaulis, P. J. Muell. in Flora, xli. (1858) 173 = tomentosus, *Borck.*
teretiusculus, Kaltenb. Fl. Aach. Beckens, 282 = vestitus.
ternatus, *Thunb. Diss. Rub.* 8.—Hab.?
tetraphyllus, *Willd. Berl. Baumz.* ed. II. 413.—Am. bor.
thamnocharis, P. J. Muell. in Pollichia, xvi–xvii. (1859) 263 = nemorosus.

RUBUS:—
thamnophilus, Ripart, ex Genev. Monog. Rub. ed. II. 247 = arduennensis.
thelybatos, *Focke, Syn. Rub. Germ.* 279.—Europ.
Thomsonii, *Focke, in Bremen, Abh.* iv. (1874) 198.—Reg. Himal.
Thuillieri, Poir. Encyc. Suppl. iv. 694 = rhamnifolius, tomentosus, *Borck.*
Thunbergii, Blume, Bijdr. 1109 = parvifolius.
Thunbergii, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* i. 18.—Japon.
thuringensis, Metsch, in Linnaea, xxviii. (1856) 199 = R. Radula.
thyrsanthoides, *Krause, in Verh. Bot. Ver. Brandenb.* xxvi. (1885) 15.—Europ.
thyrsanthus, Focke, Syn. Rub. Germ. 168 = thyrsoideus.
thyrsiflorus, Betcke, in Meklenburg. Archiv, iv. (1850) 125 = Metschii.
thyrsiflorus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 684.—Europ.
thyrsiger, Banning & Focke, ex Focke, Syn. Rub. Germ. 341 = scaber.
thyrsoideus, Bab. Brit. Rub. 108 = pubescens, *Weihe.*
thyrsoideus, Schur, Enum. Pl. Transs. 186 = candicans.
thyrsoideus, *Wimm. Fl. Schles.* (1832) 204.—Europ.
tiliaceus, Liebm. in Kjoeb. Vidensk. Meddel. (1852) 161 = tiliaefolius, *Focke.*
tiliaceus, Sm. in Rees, Cycl. xxx. n. 35 = paniculatus.
tiliaceus, Strach. & Winterb. ex Hook. f. Fl. Brit. Ind. ii. 331 = lanatus.
tiliaefolius, *Focke, in Bremen, Abh.* iv. (1874) 159.—Mexic.
tiliaefolius, *Weihe, ex Spreng. Syst.* ii. 529.—Europ.
titanophilus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 30 = nemorosus.
Tokkura, *Siebold, in Verh. Batav. Gen.* xii. (1830) 65.—Japon.
tomentellus, Ripart, ex Genèv. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 301 = tomentosus, *Borck.*
tomentosus, *Borck. in Roem. Neues Mag. Bot.* i. (1794) 2.—Europ.
tomentosus, Bory, Voy. ii. 375 = borbonicus.
tomentosus, *Thunb. Diss. Rub.* 5.—Europ.
tomentosus, *Willd. Sp. Pl.* ii. 1083.—Europ.; As. Min.
tonglooensis, Kuntze, Method. 66, 78 = moluccanus.
trachydermis, *Focke, Syn. Rub. Germ.* 324.—Europ.
trachypus, *Boul. & Gillot, in Ann. Soc. Bot. Lyon*, viii. (1881) 20.—Europ.
transalpinus ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 58.—Europ.
transiens, *Kuntze, Method.* 79, 107.—Japon.
tremulus, *Lefèv. & P. J. Muell. in Pollichia*, xvi–xvii. (1859) 138.—Europ.
Treutleri, *Hook. f. Fl. Brit. Ind.* ii. 331.—Reg. Himal.
Trevirani, Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 53 = thyrsoideus.
trichocarpus, Timer. ex Bab. Brit. Rub. 129 = scaber.
trichomallus, *Schlecht. in Linnaea*, xiii. (1839) 268.—Mexic.
tricordatus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223, nomen.—Europ.
trifidus, Steud. Nom. ed. II. ii. 479 = trilobus.
trifidus, *Thunb. Fl. Jap.* 217.—Japon.
triflorus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 233 = acuminatus.
triflorus, *Richards. in Frankl. 1st Journ. App.* ed. II. 19.—Am. arct.
trifoliatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = lineatus.
trifoliatus, *Weihe, ex Steud. Nom.* ed. II. ii. 479, nomen.—Germ.
trigonodonthos, Boul. ex Genev. Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 25 = nemorosus.
trilobus, [*Moç. & Sesse,*] *ex Ser. in DC. Prod.* ii. 566.—Mexic.
triphyllus, Bell. in Mém. Acad. Turin, 1790–91 (1793) 231 = tomentosus, *Borck.*
triphyllus, Thunb. Fl. Jap. 215 = parvifolius.
tristifrons, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 262 = villicaulis.
tristis, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 97.—Europ.
trivialis, *Michx. Fl. Bor. Am.* i. 296.—Am. bor. occ.
trivialis, Pursh, Fl. Am. Sept. i. 347 = canadensis.

RUBUS:—
truncifolius, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 139.—Europ.
tuberculatus, Bab. Fl. Camb. 306 = nemorosus.
tumidus, Gremli, Excurs. Schw. ed. II. 161 = arduennensis.
Uhdeanus, *Focke, in Bremen, Abh.* iv. (1874) 159.—Mexic.
ulmifolius, J. & C. Presl, Delic. Prag. i. 223 = nemorosus.
ulmifolius, *Schott, in Oken, Isis* (1818), fasc. v. 821.—Europ.
umbelliformis, *P. J. Muell. & Lefèv. in Pollichia*, xvi–xvii. (1859) 265.—Europ.
umbraticolus, *Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 97.—Europ.
umbraticus, P. J. Muell. in Pollichia, xvi–xvii. (1859) 114 = vestitus.
umbrosus, Lees, in Steele, Handb. Field Bot. 60 = nessensis.
umbrosus, Maass, ex Focke, Syn. Rub. Germ. 151 = rhamnifolius.
umbrosus, Weihe, ex Nym. Consp. 217 = villicaulis.
uncatus, Wall. Cat. n. 750 = macilentus, *Jacquem.*
uncinatus, Heut. ex Nym. Consp. 219 = rudis.
uncinatus, *P. J. Muell. in Flora*, xli. (1858) 154.—Europ.
uncinelliferus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 280 = villicaulis.
uncinellus, P. J. Muell. & Lefèv. in Pollichia, xvi–xvii. (1859) 286 = caesius.
undulatus, *E. Merc. in Reut. Cat. Pl. Vasc. Genèv.* ed. II. 283.—Europ.
uniflorus, *Kuntze, Method.* 91, 97.—Mexic.
universalis, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 218 = ulmifolius.
ursinus, *Cham. & Schlecht. in Linnaea*, ii. (1827) 11.—Am. bor. occ.
ursinus, Torr. & Gray, Fl. N. Am. i. 456, partim = spectabilis.
urticaefolius, *Poir. Encyc.* vi. 246.—Peruv.
ustulatus, Lefèv. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 58 = pubescens, *Weihe.*
valesiacus, Gremli, Excurs. Schw. ed. IV. 150 = arduennensis.
validulus, Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 20 = caesius.
validus, *Lefèv. in Bull. Soc. Bot. Fr.* xxiv. (1877) 223, nomen.—Europ.
vallicolus, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 188.—Europ.
variabilis, Lefèv. in Bull. Soc. Bot. Fr. xxiv. (1877) 223 = saltuum.
velatus, Lefèv. ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxviii. (1872) 34 = Sprengelii.
velicaulis, Koehl. ex Opiz, Boehm. Gew. (1823) 62 = villicaulis.
velutinus, Hook. & Arn. Bot. Beech. Voy. 140 = nutkanus.
velutinus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 221 = collinus.
velutinus, Vest, ex Tratt. Ros. iii. 47 = Menkei.
vendeanus, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 72, 98 = villicaulis.
vepretorum, *P. J. Muell. in Pollichia*, xvi–xvii. (1859) 276.—Europ.
vernus, *Focke, in Bremen, Abh.* v. (1877) 407.—Japon.
versistipulatus, *Kuntze, Method.* 84.—Hab.?
vespicum, P. J. Muell. ex Focke, Syn. Rub. Germ. 237 = tomentosus, *Borck.*
vestiferus, P. J. Muell. ex Lamotte, Prod. Fl. Centr. Fr. i. 256, nomen = thyrsiflorus.
Vestii, *Focke, Syn. Rub. Germ.* 155.—Europ.
vestitus, *Weihe & Nees, ex Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 684.—Europ.
vicinus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 283 = villicaulis.
Victoris, *Martr. Fl. Tarn*, 221.—Europ.
vigorosus, P. J. Muell. & Wirtg. ex Nym. Consp. 216 = affinis.
Villarsianus, Focke, in Gremli, Beitr. Fl. Schw. 28 = orthocanthus.
villicaulis, *Koehl. ex Weihe & Nees, Rub. Germ.* 43. t. 17.—Silesia.
villosulus, Ripart, ex Genev. Monog. Rub. ed. II. 291 = ulmifolius.
villosus, *Ait. Hort. Kew.* ed. I. ii. 210.—Am. bor.

RUBUS:—
villosus, Lasch. in Linnaea, viii. (1833) 297 = vestitus.
villosus, Thunb. Fl. Jap. 218 = corchorifolius.
villosus, Torr. Fl. N. York, i. 215 = heterophyllus, *Willd.*
vinetorum, Holandre, Fl. Moselle, ed. I. 267 = vestitus.
virescens, *G. Braun, ex Focke, Syn. Rub. Germ.* 224.—Europ.
virescens, P. J. Muell. ex Wirtg. in Flora, xlii. (1859) 235 = Fuckeli.
viretorum, P. J. Muell. in Pollichia, xvi–xvii. (1859) 275 = tomentosus, *Borck.*
virgultorum, *P. J. Muell. l. c.* 273.—Europ.
viridis, Kaltenb. Fl. Aach. Beckens, 284 = serpens.
viridis, Presl, ex Ortm. in Flora, xviii. (1835) 488 = nessensis.
viridulus, *Freyn, in Verh. Zool.-Bot. Ges. Wien*, xxxi. (1882) 376.—Europ.
viscosus, Weihe, ex Lej. & Court. Comp. Fl. Belg. ii. 170 = foliosus.
vitifolius, Cham. & Schlecht. in Linnaea, ii. (1827) 10 = ursinus.
vogesiacus, *P. J. Muell. in Flora*, xli. (1858) 170.—Europ.
Vrabelianus, *Kern. in Oestr. Bot. Zeitschr.* xix. (1869) 203.—Europ.
vulgaris, Bor. Fl. Centr. Fr. ed. III. ii. 200 = vestitus.
vulgaris, Focke, in Bremen, Abh. i. (1868) 287 = villicaulis.
vulgaris, *Weihe & Nees, Rub. Germ.* 38.—Europ.
vulgarius, *Genev. in Mém. Soc. Acad. Maine-et-Loire*, xxviii. (1872) 58.—Europ.
vulgatius, *Genev. l. c.* xx. (1866) 32.—Europ.
vulnificus, Lefèv. ex Genev. l. c. xxviii. (1872) 53 = ulmifolius.
vulpinus, Desf. Tabl. ed. I. 178 = Sprengelii.
Wahlbergii, *Arrhen. Rub. Suec. Monog.* 43.—Europ.
Wahlbergii, Focke, in Bremen, Abh. i. (1868) 305 = Laschii.
Wahlbergii, Godr. Monog. 17 = Godroni.
Wahlbergii, Neum. in Oefvers. Vet. Akad. Foerhandl. Stockh. xl. (1883) n. VIII. 78; et in Bot. Notiser (1885), 91 = corylifolius.
Wallichianus, Wight & Arn. Prod. 298 = ellipticus.
Warrenii, Bloxam, ex Bab. in Journ. Bot. xvi. (1878) 115 = villicaulis.
Wawrai, Kuntze, Method. 56 = moluccanus.
Weigelii, Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 55 = hirtus.
Weiheanus, *Focke, ex Gremli, in Oestr. Bot. Zeitschr.* xxi. (1871) 98.—Europ.
Weiheanus, Ripart, ex Genev. in Mém. Soc. Acad. Maine-et-Loire, xxiv. (1868) 253 = pubescens, *Weihe.*
Weihei, Koehl. ex Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 31 = nemorosus.
Weihei, Lej. in Lej. & Court. Comp. Fl. Belg. ii. 164 = sulcatus.
Weinteri, Genev. Monog. Rub. ed. II. 268 = pubescens.
Weitenweberi, Ortm. in Opiz, Nom. 16 = tomentosus, *Borck.*
Wiegmanni, Weihe, ex Spreng. Syst. ii. 528 = vestitus.
Wimmeri, Weihe, in Wimm. & Grab. Fl. Siles. (1829) i. II. 53 = macrophyllus.
Winteri, P. J. Muell. ex Focke, Syn. Rub. Germ. 196 = pubescens, *Weihe.*
Wirtgeni, *Auersw. ex Wirtg. Fl. Preuss. Rheinprov.* 155.—Europ.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 387.—Japon.
yessoicus, *Kuntze, Method.* 130.—Japon.
yunanicus, *Kuntze, l. c.* 71.—China.
Zahlbrucknerianus, Endl. Atakt. t. 35 = parvifolius.

RUCKERIA, DC. Prod. vi. 483 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 453.
euryopsidis, *DC. l. c.* 484.—Afr. austr.
othonnoides, *DC. l. c.*—Afr. austr.
tagetoides, *DC. l. c.*—Afr. austr.

RUCKIA, Regel, Ind. Sem. Hort. Petrop. (1867) 28; et Gartenfl. (1868) 65. t. 571 = **Rhodostachys**, Phil. (Bromel.).
Ellemeetiana, Hort.—Cf. Gard. Chron. (1874) i. 243 = seq.
Ellemeti, Regel, Ind. Sem. Hort. Petrop. (1867) 28; et Gartenfl. (1868) 65 = Rhod. andina.

RUDBECKIA, Adans. Fam. ii. 80 (1763) = **Conocarpus**, Gaertn. (Combret.).

RUDBECKIA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 365.
CENTROCARPHA, D. Don, in Sweet, Hort. Brit. ed. II. 310 (1830); et in Brit. Flow. Gard. Ser. II. t. 87 (1831).
DRACONTOPSIS, Lem. in Orb. Dict. v. 123 (1849).
DRACOPIS, Cass. in Dict. Sc. Nat. xxxv. 273 (1825).
DRACOPSIS, Lem. in Orb. Dict. v. 123 (1849).
HELIOPHTHALMUM, Rafin. Fl. Ludov. 72 (1817).
OBELISCOTHECA, Adans. Fam. ii. 128 (1763).
RATABIDA, Loud. Hort. Brit. Suppl. II. 665 (1839).
TITHONIA, Raeusch. Nom. ed. III. 251 (1797).
alata, Jacq. Ic. Pl. Rar. iii. 16. t. 593 = Helenium quadridentatum.
alata, *Neuenh. in Roem. Arch.* i. II. (1797) 30.—Am. bor.
alismaefolia, *Torr. & Gray, Fl. N. Am.* ii. 310.—Am. bor.
amplexicaulis, *Vahl, in Skriv. Nat. Selsk. Kjoeb.* ii. II. (1793) 29. t. 4.—Am. bor.; Mexic.
amplexifolia, Jacq. Ic. Pl. Rar. iii. 16. t. 592 = amplexicaulis.
angulata, Rafin. Atl. Journ. i. 48 (Quid?).—Am. bor.
angustifolia, Linn. Sp. Pl. ed. II. 1281 = Helianthus angustifolius.
apetala, Torr. ex Nutt. in Journ. Acad. Philad. vii. (1834) 77 = Helianthus Radula.
aristata, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 574 = triloba.
aspera, Desf. Tabl. ed. I. 107 = fulgida.
aspera, Pers. Syn. ii. 477 = speciosa?
asperrima, *Hornem. ex Maund, Bot. Gard.* vii. t. 647.—Am. bor.
atrorubens, *Nutt. in Journ. Acad. Philad.* vii. (1834) 80.—Am. bor.
bellidioides, Sm. in Rees, Cycl. xxx. n. 12 = Spilanthes arnicoides.
bicolor, *Nutt. in Journ. Acad. Philad.* vii. (1834) 81.—Am. bor.
biennis, Chapm. ex Torr. & Gray, Fl. N. Am. ii. 309 = triloba.
bupleuroides, *Shuttl. ex Chapm. Fl. S. U. St. Suppl.* 629.—Florida.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 357.—Calif.
chrysomela, Michx. Fl. Bor. Am. ii. 143 = fulgida.
cicutaefolia, *Spreng. Syst.* iii. 613.—Am. bor.
cinerea, Hort. ex DC. Prod. v. 559 = Lepachys pinnatifida.
citriodora, Hort. ex Steud. Nom. ed. I. 707 = Lepachys pinnatifida.
columnaris, Sims, Bot. Mag. t. 1601; Pursh, Fl. Am. Sept. ii. 575 = Lepachys columnaris.
columnifera, [Nutt.] in Fras. Cat. (1813) n. 75; Pursh, l. c. = Lepachys columnaris.
decumbens, Sm. in Rees, Cycl. xxx. n. 11 = Spilanthes arnicoides.
densifolia, Sm. l. c. n. 10 = Aspilia buphthalmiflora, Viguiera anchusaefolia.
dentata, Cav. Desc. 223 = Viguiera dentata.
digitata, Mill. Gard. Dict. ed. VIII. n. 6 = laciniata.
digitata, Willd. Sp. Pl. iii. 2247 = Lepachys pinnatifida.
discolor, Ell. Sketch, ii. 454 = hirta.
discolor, Pursh, Fl. Am. Sept. ii. 574 = fulgida.
Drummondii, *Paxt. Mag. Bot.* vi. (1839) 51.—Am. bor.
fulgida, *Ait. Hort. Kew.* ed. I. iii. 251.—Am. bor.
fulgida, Meehan, Nat. Flowers, Ser. II. i. t. 14 = speciosa.
glabra, DC. Prod. v. 556 = nitida.
globosa, Nutt. in Journ. Acad. Philad. vii. (1834) 79 = Lepachys Tagetes.
gracilis, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 574 = spathulata.
gracilis, Nutt. Gen. Am. ii. 178 = hirta.
grandiflora, *Gmel. ex DC. Prod.* 556.—Am. bor.
Heliopsidis, A. H. Curtiss, ex A. Gray, Syn. Fl. N. Am. i. II. 261 = spathulata.
Heliopsidis, *Torr. & Gray, Fl. N. Am.* ii. 310.—Am. bor.

RUDBECKIA:—
heterophylla, Schur, in Verh. Zool.-Bot. Ges. Wien, viii. (1858) Sitzb. 118 = amplexicaulis.
heterophylla, *Torr. & Gray, Fl. N. Am.* ii. 312.—Florida.
hirta, *Linn. Sp. Pl.* 907.—Am. bor.
hispida, Hoffmgg. Verz. Pfl. Nachtr. i. 223 = Echinacea purpurea.
laciniata, Linn. Sp. Pl. 906.—Am. bor.
laevigata, Nutt. Gen. Am. ii. 178 = nitida.
laevigata, *Pursh, Fl. Am. Sept.* ii. 574.—Am. bor.
laevis, Hoffmgg. Verz. Pfl. Nachtr. iii. 58 = Lepachys pinnatifida.
lanceolata, *Bertol. Misc. Bot.* vii. 40.—Am. bor.
lanceolata, *Bisch. Del. Sem. Hort. Heidelb.* (1848) 4; *ex Walp. Ann.* ii. 855.—Am. bor.
lancifolia, Bisch. Heidelb. Consp. Crit. (1852) 7 = praec.?
leucantha, *Lag. Gen. et Sp. Nov.* 32.—Mexic.
maxima, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 354.—Am. bor.
media, [*Spreng*]. *Bot. Gart. Halle*, 80 (Quid?).—Hab.?
Mohrii, A. Gray, in Proc. Am. Acad. xvii. (1881-82) 217 = bupleuroides.
mollis, *Ell. Sketch*, ii. 453.—Am. bor.
montana, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 217.—Am. bor. occ.
moschata, *Lodd. ex Steud. Nom.* ed. II. ii. 480.—Am. bor.
napifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 244.—Mexic.
Neumanni, Steud. Nom. ed. II. ii. 480 = Newmani.
Neumannii, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 990 = laciniata.
Newmani, Loud. Hort. Brit. 358 = fulgida.
Newmani, *Loud. l. c. Suppl.* ii. 584, in syn.—Am. bor.
nitida, *Nutt. in Journ. Acad. Philad.* vii. (1834) 78.—Am. bor.
nudicaulis, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 311 = grandiflora.
nudicaulis, Pers. Syn. ii. 477 = Viguiera nudicaulis.
nudiflora, *Bertol. Misc. Bot.* vii. 39.—Am. bor.
occidentalis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 355.—Am. bor. occ.
odorata, Hort. ex Pursh, Fl. Am. Sept. ii. 576 = Lepachys pinnatifida.
odorata, Nutt. in Journ. Acad. Philad. vii. (1834) 78 = subtomentosa.
oppositifolia, Hort. ex. Vilm. Fl. Pl. Terre, ed. I. 743 = amplexicaulis.
oppositifolia, Linn. Sp. Pl. 907 = Heliopsis laevis.
pallida, Nutt. in Journ. Acad. Philad. vii. (1834) 77 = Echinacea angustifolia.
pauciflora, [Zinn,] Steud. Nom. ed. I. 707 = Zinnia pauciflora.
perfoliata, Cav. Ic. iii. 27 = amplexicaulis.
pinnata, Vent. Jard. Cels, 71 = Lepachys pinnatifida.
Porteri, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 83 = Gymnolomia Porteri.
purpurea, Linn. Sp. Pl. 907 = Echinacea purpurea.
quinata, Mill. Gard. Dict. ed. VIII. n. 5 = laciniata.
Radula, Pursh, Fl. Am. Sept. ii. 575 = Helianthus Radula.
rupestris, Chick. in Coult. Bot. Gaz. vi. (1881) 188 = triloba.
scabra, Cav. Desc. 224 = triloba.
scabra, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 747 = fulgida.
serotina, Nutt. in Journ. Acad. Philad. vii. (1834) 80 = hirta.
serotina, Sweet, Brit. Flow. Gard. Ser. I. t. 4 = Echinacea purpurea.
spathulata, *Michx. Fl. Bor. Am.* ii. 144.—Am. bor.
spathulata, Nutt. Gen. Am. ii. 178 = amplexicaulis.
spathulata, Pursh, Fl. Am. Sept. ii. 574 = mollis.
speciosa, Hort. ex Link, Enum. Hort. Berol. ii. 353 = Echinacea purpurea.
speciosa, *Wender. in Flora*, xii. (1829) I. *Erg.* 30.—Am. bor.
strigosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 354 = hirta.
subtomentosa, *Pursh, Fl. Am. Sept.* ii. 575.—Am. bor.
Tagetes, James, in Long, Exped. ii. 256, 353 = Lepachys Tagetes.
tomentosa, Ell. Sketch, ii. 453 = subtomentosa.
triloba, *Linn. Sp. Pl.* 907.—Am. bor.

RUDBEKIA, Scop. Introd. 131 (1777) = praec.

RUDELIA, Oliver, in Hook. Ic. Pl. t. 1419 (1883). *SCITAMINEAE*.
curviflora, *Oliver, l. c.* — Archip. Ind.

RUDGEA, Salisb. in Trans. Linn. Soc. viii. (1807) 327. tt. 18, 19. *RUBIACEAE*, Benth. & Hook. f. ii. 125.
PACHYSANTHUS, Presl, Bot. Bemerk. 87 (1844); in Abh. Boehm. Ges. V. iii. (1845) 517.
amazonica, *Muell. Arg. in Flora*, lix. (1876) 449, 460.—Bras.
angustifolia, *Muell. Arg. l. c.* 453, 462.—Bras.
apoda, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 178.—Bras.
bacciflora, *Muell. Arg. l. c.* 164.—Bras.
Blanchetiana, *Muell. Arg. in Flora*, lix. (1876) 452, 461.—Bras.
brachyandra, *Muell. Arg. l. c.* 451, 461.—Bras.
breviflora, *Muell. Arg. l. c.* 456, 463.—Bras.
Burchelliana, *Muell. Arg. l. c.* 453, 462.—Bras.
calycina, *Benth. in Linnaea*, xxiii. (1850) 459.—Bras.
caribaea, *Benth. l. c.* 460.—Ind. occ.
Casarettoana, *Muell. Arg. in Flora*, lix. (1876) 452, 461.—Bras.
celastrinea, *Muell. Arg. l. c.* 450, 460.—Bras.
ciliata, Spreng. Syst. i. 755 = Coffea ciliata.
Clausseniana, *Benth. in Linnaea*, xxiii. (1850) 461.—Bras.
clavipes, *Benth. l. c.* 462.—Bras.
conocarpa, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
corniculata, *Benth. in Linnaea*, xxiii. (1850) 462.—Bras.
coronata, *Muell. Arg. in Flora*, lix. (1876) 449.—Bras.
corymbulosa, *Benth. in Linnaea*, xxiii. (1850) 461.—Bras.
cujabensis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 215.—Bras.
cyanosperma, *Muell. Arg. l. c.* 211.—Bras.
decipiens, *Muell. Arg. in Flora*, lix. (1876) 454, 462.—Bras.
discolor, *Benth. in Linnaea*, xxiii. (1850) 460.—Bras.
elliptica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 219.—Bras.
eriantha, *Benth. in Linnaea*, xxiii. (1850) 458.—Bras.
erioloba, *Benth. l. c.* 459.—Bras.
erythrocarpa, *Muell. Arg. in Flora*, lix. (1876) 451, 461.—Bras.
fissistipula, *Muell. Arg. l. c.* 449, 460.—Bras.
Francavillana, *Muell. Arg. l. c.* 454, 462.—Bras.
gardenioides, *Muell. Arg. l. c.* 455.—Bras.
Gaudichaudii, *Muell. Arg. l. c.* 453, 461.—Bras.
Glaziovii, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 217.—Bras.
Gomeziana, *Muell. Arg. l. c.* 177.—Bras.
goyazensis, *Muell. Arg. in Flora*, lix. (1876) 450, 461.—Bras.
Heurckii, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 209.—Bras.
Hostmanniana, *Benth. in Linnaea*, xxiii. (1850) 459.—Guian.
ilheotica, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 217.—Bras.
insignis, *Muell. Arg. l. c.* 203.—Bras.
intercedens, *Muell. Arg. l. c.* 205.—Bras.
interrupta, *Benth. in Linnaea*, xxiii. (1850) 457.—Bras.
involucrata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 219.—Bras.
irregularis, *Muell. Arg. in Flora*, lix. (1876) 451, 461.—Bras.
jacobinensis, *Muell. Arg. l. c.* 458, 461.—Bras.
japurensis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 177.—Bras.
jasminioides, *Muell. Arg. in Flora*, lix. (1876) 452.—Bras.
lacinulata, *Muell. Arg. l. c.* 454, 462.—Bras.
lanceaefolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 327. t. 18.—Guian.
lanceolata, *Benth. in Linnaea*, xxiii. (1850) 455.—Bras.
Langsdorffii, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 191.—Bras.
lasiostylis, *Muell. Arg. l. c.* 212.—Bras.

RUDGEA :—
leiocarpoides, *Muell. Arg. in Flora*, lix. (1876) 456.—Bras.
longiflora, *Benth. in Linnaea*, xxiii. (1850) 457.—Bras.
longistipula, *Muell. Arg. in Flora*, lix. (1876) 449, 460.—Bras.
loniceroides, *Muell. Arg. l. c.* 454, 462.—Bras.
Lundiana, *Muell. Arg. l. c.* 456, 463.—Bras.
macrophylla, *Benth. in Linnaea*, xxiii. (1850) 456.—Bras.
magnoliaefolia, *Muell. Arg. in Flora*, lix. (1876) 450.—Bras.
major, *Muell. Arg. l. c.* 452.—Bras.
Martiana, *Muell. Arg. l. c.* 453, 462.—Bras.
medians, *Muell. Arg. ll. cc.*—Bras.
mexicana, Bartl. ex DC. Prod. iv. 501 = Coffea mexicana.
micrantha, *Muell. Arg. in Flora*, lix. (1876) 454.—Bras.
microcephala, *Muell. Arg. l. c.* 456, 464.—Bras.
Miersiana, *Benth. in Linnaea*, xxiii. (1850) 461.—Bras.
mucronata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. (1850) 185.—Bras.
multicostata, *Muell. Arg. l. c.* 207.—Bras.
myrsinaefolia, *Benth. in Linnaea*, xxiii. (1850) 455.—Bras.
Nettoana, *Muell. Arg. in Flora*, lix. (1876) 450, 460.—Bras.
Neuwiedii, *Muell. Arg. l. c.* 455, 463.—Bras.
nitida, Spreng. Syst. i. 755 = Coffea nitida.
nivosa, *André, in Rev. Hortic.* (1866) 308; *et Berkeley, in Journ. Hort. Soc.* N. S. i. (1866) 156.—Bras.
nobilis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 172.—Bras.
nodosa, *Benth. in Linnaea*, xxiii. (1850) 456.—Bras.
ochroleuca, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
ovalifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 328. t. 19.—Guian.
ovalis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 462.—Bras.
pachyphylla, *Muell. Arg. l. c.* 183.—Bras.
pachysanthus, *Muell. Arg. l. c.* 208.—Bras.
palicoureoides, *Muell. Arg. in Flora*, lix. (1876) 455.—Bras.
paniculata, *Benth. in Linnaea*, xxiii. (1850) 457.—Bras.
panurensis, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
parquioides, *Muell. Arg. l. c.* 450.—Bras.
parvifolia, *Muell. Arg. l. c.* 454.—Bras.
psammophila, *Muell. Arg. l. c.* 453, 462.—Bras.
racemosa, Spreng. Syst. i. 755 = Coffea racemosa, *Ruiz & Pav.*
recurva, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
reticulata, *Benth. in Linnaea*, xxiii. (1850) 458.—Bras.
Riedeliana, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
Schomburgkiana, *Benth. in Linnaea*, xxiii. (1850) 459.—Guian.
Selloana, *Muell. Arg. in Flora*, lix. (1876) 453, 462.—Bras.
sessilis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 182.—Bras.
spicata, Spreng. Syst. i. 755 = Coffea spicata.
stenantha, *Muell. Arg. in Flora*, lix. (1876) 456, 463.—Bras.
subcordata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 178.—Bras.
subsessilis, *Benth. in Linnaea*, xxiii. (1850) 458.—Bras.
symplocoides, *Muell. Arg. in Flora*, lix. (1876) 455, 463.—Bras.
tenuifolia, *Muell. Arg. l. c.* 453, 462.—Bras.
tinguana, *Muell. Arg. l. c.* 455, 463.—Bras.
triflora, *Benth. in Linnaea*, xxiii. (1850) 460.—Bras.
umbellata, Muell. Arg. in Flora, lix. (1876) 456 = Psychotria Arrabidae.
umbrosa, *Muell. Arg. l. c.* 452, 461.—Bras.
vellerea, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 208.—Bras.
verticillata, Spreng. Syst. i. 755 = Coffea verticillata.

RUDGEA :—
viburnoides, *Benth. in Linnaea*, xxiii. (1850) 458.—Bras.
Weddelliana, *Muell. Arg. in Flora*, lix. (1876) 451, 461.—Bras.
ypanemensis, *Muell. Arg. l. c.* 452, 461.—Bras.

RUDOLPHIA, Medic. Malv. 111 (1787) = **Malpighia**, Plum.
edulis, Medic. l. c. 112 = M. urens.

RUDOLPHIA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 451. *LEGUMINOSAE*, Benth. & Hook. f. i. 532.
dubia, H. B. & K. Nov. Gen. et Sp. vi. 432. t. 591 = Centrosema hastatum.
elliptica, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 422 (= *Collaea rugosa*).—Bras.
frondosa, Poir. Encyc. vi. 333 = Butea frondosa.
peltata, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 453.—Ins. S. Doming.
portoricensis, Sweet, Hort. Brit. ed. II. 161 = volubilis.
rosea, *Tussac, Fl. Antill.* i. 149. t. 22 (rosaea).—Ins. S. Doming.
scandens, Poir. in Dict. Sc. Nat. xlvi. 419 = volubilis.
sericea, Steud. Nom. ed. II. i. 241 = volubilis.
superba, Poir. Encyc. vi. 332 = Butea superba.
volubilis, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 452.—Porto Rico; Mexic.

RUDOLPHO-ROEMERIA, Steud. ex Hochst. in Flora, xxvii. (1844) 30 = **Kniphofia**, Moench (Liliac.).
isoëtifolia, Steud. l. c. = K. isoëtifolia.

RUEHSSIA, Karst. in Verh. Ver. Gartenb. Preuss. xix. (1849) II. 304 = **Marsdenia**, R. Br. (Asclep.).
estebanensis, Karst. in Linnaea, xxviii. (1856) 424.—N. Granat.
glauca, Karst. l. c.—N. Granat.
macrophylla, Karst. l. c.—N. Granat.
pubescens, Karst. l. c.—N. Granat.
purpurea, Schlecht. l. c. xxvi. (1853–55) 669.—N. Granat.

RUELINGIA, Ehrh. Beitr. iii. 132 (1788) = **Anacampseros**, Linn. (Portul.).
Anacampseros, Ehrh. l. c. 133 = A. Telephiastrum.
angustifolia, Haw. Rev. Pl. Succ. 60 = A. angustifolia.
arachnoides, Haw. Syn. Pl. Succ. 125 = A. arachnoides.
filamentosa, Haw. l. c. = A. filamentosa.
lanceolata, Haw. l. c. 126 = A. lanceolata.
patens, Ehrh. Beitr. iii. 135 = Talinum patens.
polyphylla, Haw. Suppl. Pl. Succ. 65 = A. polyphylla.
reflexa, [Spreng.] Bot. Gart. Halle, 20.—Hab.?
rubens, Haw. Syn. Pl. Succ. 125 = A. arachnoides.
rufescens, Haw. Suppl. Pl. Succ. 64 = A. arachnoides.
triangularis, Ehrh. Beitr. iii. 134 = Talinum triangulare.
varians, Haw. Suppl. Pl. Succ. 64 = A. Telephiastrum.

RUELLIA, Plum. ex Linn. Syst. ed. I (1735). *ACANTHACEAE*, Benth. & Hook. f. ii. 1077.
ANTHEILEMA, Rafin. Fl. Tellur. iv. 66 (1836).
APHRAGMIA, Nees, in Lindl. Syst. Nat. ed. II. 444 (1836).
ARRHOSTOXYLUM, Mart. ex Nees, in DC. Prod. xi. 209 (1847).
AUBLETIA, Neck. Elem. i. 356 (1790).
COPIOGLOSSA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 294.
CRYPHIACANTHUS, Nees, in Del. Sem. Hort. Vrat. (1841); et in Linnaea, xvi. (1842) 298.
CYRTACANTHUS, Mart. ex Nees, in Mart. Fl. Bras. ix. 51 (1347).
DIPTERACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75 (1832).
DIZYGANDRA, Meissn. Gen. 294 (1840).
EURYCHANES, Nees, in Mart. Fl. Bras. ix. 52. t. 3 (1847).
FABRIA, E. Mey. in Drège, Zwei Pfl. Docum. 185, nomen (1843).
GYMNACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 126.
HEMONACANTHUS, Nees, in DC. Prod. xi. 725 (1847).

RUELLIA :—
HOLTZENDORFFIA, Klotzsch & Karst. ex Nees, in DC. Prod. xi. 727 (1847).
HOMOTROPIUM, Nees, in Mart. Fl. Bras. ix. 47. t. 2 (1847).
LARYSACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 123.
NEOWEDIA, Schrad. in Neuwied, Reise Bras. ii. 343 (1821).
OPHTHALMACANTHUS, Nees, in DC. Prod. xi. 219 (1847).
PATTERSONIA, J. F. Gmel. Syst. 925 (1791).
SCORODOXYLUM, Nees, in Benth. Pl. Hartw. 236 (1846).
SIPHONACANTHUS, Nees, in Mart. Fl. Bras. ix. 45. t. 1 (1847).
SOLENACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 122.
STEMONACANTHUS, Nees, in Mart. Fl. Bras. ix. 53. tt. 4, 205 (1847).
STEPHANOPHYSUM, Pohl, Pl. Bras. Ic. ii. 83. tt. 155, 156 (1831).
UPUDALIA, Rafin. Fl. Tellur. iv. 66 (1836)
acaulis, *R. Br. Prod.* 479.—Austral.
acaulis, Ruiz, ex Nees, in DC. Prod. xi. 282 = Stenandrium dulce.
achimeniflorum, *Hemsl. Biol. Centr. Am. Bot.* ii. 503.—Costa Rica.
achyranthifolia, *Desf. Cat. Hort Par.* ed. III. 391.—Hab.?
acicularis, Steud. Nom. ed. II. ii. 480, sphalm. = seq.
aciculata, Roth, Nov. Pl. Sp. 301 = Haplanthus tentaculatus.
acuminata, *Griseb. Fl. Brit. W. Ind.* 452.—Ind. occ.
acutangula, *Nees, ex Mart. in Flora*, xxi. (1838) II. *Beibl.* 61.—Bras.
adhaerens, *Forsk. Fl. Aegypt. Arab.* 114.—Arab.
affinis, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 269 = speciosa.
alata, *Blume, ex Steud. Nom.* ed. II. ii. 480, nomen.—Java.
alata, Wall. Pl. As. Rar. i. 26 = Strobilanthes Wallichii.
alba, *Nees, in Mart. Fl. Bras.* ix. 55.—Bras.
albicans, Bert. ex Nees, in DC. Prod. xi. 724 = albicaulis.
albicaulis, *Bert. ex Spreng. Syst.* ii. 822.—N. Granat.
alopecuroides, Vahl, Eclog. Am. ii. 49 = Lepidagathis alopecuroides.
alternata, Burm. f. Fl. Ind. 135 = Hemigraphis alternata.
amabilis, *S. Moore, in Journ. Bot.* xviii. (1880) 7.—Afr. trop.
amoena, *Nees, in DC. Prod.* xi. 203, in syn.—Am. austr.
amplectens, Wall. ex Nees, in Wall. Pl. As. Rar. iii. 86 = Strobilanthes auriculatus.
ampliata, Edgew. ex Nees, in DC. Prod. xi. 193 = Strobilanthes Wallichii.
amygdalaefolia, Hort. ex Steud. Nom. ed. II. ii. 480 = Strobilanthes anisophyllus.
Anagallis, Burm. f. Fl. Ind. 135 = Bonnaya veronicaefolia.
angusta, Boj. ex Nees, in DC. Prod. xi. 511 = Isoglossa angusta.
angustifolia, Poir. Encyc. vi. 338 = Hygrophila angustifolia.
angustifolia, Sw. Prod. Veg. Ind. Occ. 93 = Blechum angustifolium.
anisophylla, Wall. ex Hook. Exot. Fl. iii. t. 191 = Strobilanthes anisophyllus.
antipoda, Linn. Sp. Pl. 635 = Bonnaya veronicaefolia.
arborescens, Pav. ex Nees, in DC. Prod. xi. 217 = Bravaisia floribunda.
arcuata, Herb. Madr. ex Wall. Cat. n. 2334 = Stenosiphonium Russellianum.
argentea, Wall. Cat. n. 2339 = Strobilanthes Sabinianus.
aristata, Thunb. Prod. Pl. Cap. 104 = Calophanes Persoonii.
aristata, Vahl, Symb. Bot. ii. 73 = Lepidagathis hyalina.
aromatica, Roxb. ex Nees, in DC. Prod. xi. 127 = patula.
aruensis, *S. Moore, in Journ. Bot.* xvi. (1878) 134.—Ins. Aru.
aspera, Nees, in DC. Prod. xi. 147 = Strobilanthes scaber.
asperula, *Benth. & Hook. f. Gen.* ii. 1079.—Bras.
atropurpurea, Wall. Cat. n. 2398 = Strobilanthes atropurpureus.

RUELLIA:—
atroviridis, Wall. Cat. n. 2404 = Asystasia atroviridis.
attenuata, Wall. l c. n. 2345 = Echinacanthus attenuatus.
auriculata, Wall. l. c. n. 2341 = Strobilanthes auriculatus.
australis, *Cav. Ic.* vi. 62; *R. Br. Prod.* 479.—Austral.
axillaris, Steud. Nom. ed. II. ii. 480 = Ebermaiera axillaris.
axillaris, Wall. ex Nees, in DC. Prod. xi. 127 = patula.
balsamica, Linn. f. Suppl. 289 = Cardanthera balsamica.
barbata, Sieber, ex Steud. Nom. ed. II. i. 701, in syn. = Gonzalea Coccocypselum.
barbata, *Vahl, Symb. Bot.* iii. 83.—Ind. or.
barbata, Wall. Cat. ex Nees, in DC. Prod. xi. 155 = Strobilanthes barbatus.
barlerioides, Roth, Nov. Pl. Sp. 310 = Petalidium barlerioides.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 413.—Ind. or.
biceps, Steud. Nom. ed. II. ii. 480 = Strobilanthes capitatus.
bicolor, *Blume, Bijdr.* 795.—Java.
bicolor, Ruiz, ex Nees, in DC. Prod. xi. 136 = geminiflora.
biflora, Balb. ex Nees, l. c. 122 = strepens.
biflora, Boj. ex Nees, l. c. 726 = Phaylopsis longifolia.
biflora, Bosc, ex Nees, l. c. 108 = Calophanes humistratus.
biflora, Linn. Sp. Pl. 635 = Calophanes oblongifolius ?
biflora, *Pav. ex Nees, in DC. Prod.* xi. 133.—Peruv.
bignoniaeflora, *S. Moore, in Journ. Bot.* xviii. (1880) 198.—Afr. trop.
bilabiata, Reinw. ex Blume, Cat. Gew. Buitenz. 84, nomen = Sautiera tinctorum.
biloba, Hochst. in Flora, xxviii. (1845) 72 = Asystasia coromandeliana.
blechioides, Sw. Prod. Veg. Ind. Occ. 93 = Blechum laxiflorum.
Blechum, Linn. Syst. ed. X. 1120; Amoen. Acad. v. 400 = Blechum Brownei.
Blumeana, Nees, in DC. Prod. xi. 149 = Hemigraphis alternata.
Blumei, *Steud. Nom.* ed. II. ii. 480.—Java.
Bourgaei, *Hemsl. Diagn. Pl. Nov.* 35.—Mexic.
brachyopoda, Boj. ex Nees, in DC. Prod. xi. 726 = Phaylopsis longifolia, *T. Anders.*
brachysiphon, *Benth. & Hook. f. Gen.* ii. 1078; *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 73.—Bras.
bracteata, *R. Br. Prod.* 479.—Austral.
bracteata, Heyne, in Wall. Cat. n. 2357 = Strobilanthes Heyneanus.
bracteata, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 47 = Petalidium barlerioides.
bracteata, Vahl, ex Nees, in DC. Prod. xi. 466 = Blechum Brownei.
bracteata, Wall. Cat. n. 2357 B = Strobilanthes quadrangularis.
bracteata, Willd. ex Nees, in DC. Prod. xi. 155 = Hygrophila quadrivalvis.
brasiliensis, Mart. ex Nees, in Mart. Fl. Bras. ix. 29 = solitaria.
brasiliensis, Spreng. Syst. ii. 822 = Hygrophila portoricensis.
brevicaulis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 428.—Madag.
Brunoniana, Wall. Cat. n. 2368 = Strobilanthes Brunonianus.
callosa, Wall. l. c. n. 2359 = Strobilanthes callosus.
calycina, Wall. l. c. n. 2417 = Strobilanthes Helictus.
campestris, *Hemsl. Biol. Centr. Am. Bot.* ii. 504.—Costa Rica.
candida, *Ruiz & Pav. ex Nees, in DC. Prod.* xi. 134, in syn.—Peruv.
canescens, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 428 = Stenosiphonium Russellianum.
capitata, D. Don, Prod. Fl. Nep. 120 = Strobilanthes pentstemonoides.
capitata, Wall. Cat. n. 2351 = Strobilanthes capitatus, pentstemonoides.
carnea, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
carolinensis, Steud. Nom. ed. II. ii. 481 = strepens ?
cayenennsis, *Spreng. Syst.* ii. 823.—Guian.
Cephalotes, Wall. Cat. n. 2352 = Strobilanthes pentstemonoides.

RUELLIA:—
cernua, T. Anders. in Journ. Linn. Soc. ix. (1867) 260, partim = Beddomei.
cernua, Herb. Madr. ex Wall. Cat. n. 2380 A = Calophanes Nagchana.
cernua, Heyne, ex Wall. l. c. n. 2380, partim = Cardanthera balsamica.
cernua, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 45.—Ind. or.
chelonoides, Wall. Cat. n. 2335 = Asystasia crispata.
chenopodiifolia, Poir. Encyc. vi. 339 = Adenosma chenopodiifolia.
chinensis, Lam. ex Nees, in DC. Prod. xi. 475 = Dicliptera ciliaris.
chinensis, Lyall, ex Nees, l. c. 476 = Dicliptera madagascariensis.
chinensis, *Nees, l. c.* 147.—China.
ciliaris, Buch.-Ham. ex Wall. Cat. n. 2373 C = Hygrophila angustifolia.
ciliaris, Linn. Mant. i. 89 = Blepharis edulis ?
ciliaris, Pers. Syn. ii. 176 = pilosa, *Linn.*
ciliata, Herb. Madr. ex Wall. Cat. n. 2392 = Strobilanthes barbatus.
ciliata, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 53; *Hort. Hafn.* ii. 585.—Ind. or.; Burma.
ciliata, Ruiz, ex Nees, in DC. Prod. xi. 110 = Calophanes ciliatus.
ciliata, Wall. Cat. n. 2415 = Strobilanthes ciliatus.
ciliata, Wall. ex Nees, in DC. Prod. xi. 156 = Barleria cristata.
ciliatiflora, *Hook. Bot. Mag.* t. 3718.—Reg. Argent.
ciliosa, *Pursh, Fl. Am. Sept.* ii. 420.—Am. bor.
cirsioides, Nees, in Linnaea, xv. (1841) 354 = Crabbea nana.
clandestina, Hook. ex Nees, in DC. Prod. xi. 153 = obtusa.
clandestina, Linn. Sp. Pl. 634 = tuberosa.
coccinea, *Vahl, Symb. Bot.* iii. 83.—Ind. occ.
collina, Wall. Cat. n. 2343 = Strobilanthes collinus.
colorata, *Blume, Bijdr.* 795.—Java.
colorata, Vell. Fl. Flum. 267; vi. t. 99 (= *Lagochilium montanum*).—Bras.
colorata, Wall. Cat. n. 2388 = Strobilanthes coloratus.
commutata, Steud. Nom. ed. II. ii. 481 = Calophanes Nagchana.
comosa, Roxb. Hort. Beng. 46; Fl. Ind. iii. 43 = Strobilanthes rufescens.
comosa, Vell. Fl. Flum. 265; vi. t. 91 = Aphelandra squarrosa.
comosa, Wall. Cat. n. 2414 = Ebermaiera axillaris.
concavifolia, Orteg. Hort. Matr. Dec. 94 = patula.
confinis, Nees, in DC. Prod. xi. 148 = Hemigraphis confinis.
contorta, Blanco, Fl. Filip. ed. I. 496 = Peristrophe contorta.
cordata, Thunb. Prod. Pl. Cap. 104 = pilosa, *Linn.*
cordifolia, Vahl, Symb. Bot. iii. 84 = Stenosiphonium Russellianum.
coromandeliana, Herb. Madr. ex Wall. Cat. n. 2399 = Asystasia coromandeliana, violacea.
coromandelina, Herb. Monac. ex Nees, in DC. Prod. xi. 166 = Asystasia coromandeliana.
corymbosa, Wall. ex Nees, l. c. 167 = Strobilanthes Panichanga.
corynotheca, *F. Muell. ex Benth. Fl. Austral.* iv. 546.—Austral.
costata, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 76.—Bras.
crenata, Benth. ex Hohen. in Flora, xxxii. (1849) 558 = Hemigraphis latebrosa.
crispa, Linn. Herb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 446 = Strobilanthes scaber.
crispa, Linn. Sp. Pl. 635 = Hemigraphis crispa.
crispa, Nees, in Wall. Pl. As. Rar. iii. 83 = Hemigraphis venosa.
crispa, Roxb. ex T. Anders. in Journ. Linn. Soc. ix. (1867) 461 = Hemigraphis dura.
cristata, Andr. Bot. Rep. t. 506 = Aphelandra cristata.
Crossandra, Steud. Nom. ed. II. ii. 481 = Hemigraphis latebrosa.
Crucis, Steud. l. c. = tetragona, *Link.*
cucullata, Boj. ex Nees, in DC. Prod. xi. 726 = Phaylopsis parviflora.
Cumingiana, Nees, l. c. 148 = Hemigraphis Cumingiana.
Currori, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 24.—Afr. trop.
curviflora, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 50.—Bras.

RUELLIA:—
Cusia, Buch.-Ham. ex Wall. Cat. n. 2386 = Strobilanthes flaccidifolius.
cuspidata, Heyne, ex Wall. l. c. n. 2405 = Lepidagathis cuspidata.
cyanea, *Boj. ex Nees, in DC. Prod.* xi. 121, in syn.—Madag.
Decaisniana, Nees, l. c. 150 = Hemigraphis Decaisneana.
deccanensis, J. Grah. Cat. Pl. Bomb. 162 = prostrata.
decipiens, Nees, in Wall. Pl. As. Rar. iii. 83 = Hemigraphis decipiens.
densa, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 74.—Bras.
densiflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 504.—Panama.
denticulata, Wall. Cat. n. 2418 = Strobilanthes denticulatus.
dependens, Hook. ex Nees, in DC. Prod. xi. 470 = Rungia parviflora.
dependens, Roxb. Hort. Beng. 46; Fl. Ind. iii. 49 = Lepidagathis hyalina.
depressa, Linn. f. Suppl. 290 = Aptosimum depressum.
depressa, Wall. Cat. n. 2379 = Calophanes Nagchana.
Devosiana, *Hort. Makoy, ex E. Morr. in Belg. Hortic.* xxvii. (1877) 344. t. 19.—Bras.
dichotoma, Bert. ex Nees, in DC. Prod. xi. 142 = paniculata.
difformis, Linn. f. Suppl. 289 = Cardanthera triflora.
diffusa, *Griseb. Cat. Pl. Cub.* 195.—Ins. S. Doming.
diffusa, Royle, ex Nees, in DC. Prod. xi. 171 = Aechmanthera tomentosa.
diffusa, Vell. Fl. Flum. 266; vi. t. 94 = Nelsonia campestris.
diffusa, Wall. Cat. n. 2383 = Hemigraphis latebrosa.
Digitalis, Koen. ex Spreng. Syst. ii. 825 = Asystasia coromandeliana.
dimidiata, Nees, in DC. Prod. xi. 157 = Hygrophila angustifolia.
Dipteracanthus, *Hemsl. Biol. Centr. Am. Bot.* ii. 504.—Mexic.
discolor, Lecchi, ex Colla, Hort. Ripul. App. ii. 356 = *nova.*
discolor, *Nees, in DC. Prod.* xi. 149.—Java.
dissitifolia, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 73.—Bras.
disticha, Buch.-Ham. ex Wall. Cat. n. 2365 = Lepidagathis hyalina.
divaricata, Hoffmgg. Verz. Pfl. 193 = lactea.
divaricata, Wall. Cat. n. 2372 = Strobilanthes divaricatus.
diversifolia, *S. Moore, in Journ. Bot.* xviii. (1880) 198.—Afr. trop.
domingensis, *Spreng. ex Nees, in DC. Prod.* xi. 136, in syn.—Ins. S. Doming.
dorsiflora, Retz. Obs. vi. 31 = Phaylopsis parviflora.
Drummondiana, *A. Gray, Syn. Fl. N. Am.* ii. 1. 326.—Texas.
dubia, Vahl, ex Nees, in DC. Prod. xi. 79 = Ebermaiera elongata.
dulcis, Cav. Ic. vi. 62. t. 585. f. 2 = Stenandrium dulce.
dulcis, Wall. Cat. n. 7167 = Lepidagathis dulcis.
dulcissima, Steud. Nom. ed. II. ii. 481 = Lepidagathis dulcis.
dura, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 311 = Hemigraphis dura.
ebracteata, Dalz. in Hook. Kew Journ. ii. (1850) 342 = Hemigraphis ebracteolata.
ebracteolata, T. Anders. in Journ. Linn. Soc. ix. (1867) 462 = Hemigraphis ebracteolata.
echinata, Wall. Cat. n. 2356 = Strobilanthes pectinatus.
elegans, Hook. Bot. Mag. t. 3389 = Hemigraphis latebrosa.
elegans, Poir. Encyc. Suppl. iv. 727 = formosa.
elongata, Beauv. Fl. Owar. i. 46 = Whitfieldia longifolia.
erecta, Buch.-Ham. ex Wall. Cat. n. 2395 = Hemigraphis latebrosa.
erecta, Burm. f. Fl. Ind. 135 = Calophanes Nagchana.
erecta, Roth, Nov. Pl. Sp. 308 = patula.
eriostachya, Boj. ex Nees, in DC. Prod. xi. 67 = Nelsonia campestris.
eucoma, Steud. Nom. ed. II. ii. 481 = Ebermaiera axillaris.
extensa, Steud. l. c. = Strobilanthes extensus.
falcata, Steud. l. c. = Lepidagathis falcata.
fasciculata, Herb. Madr. ex Wall. Cat. n. 2394 = Hemigraphis latebrosa.

RUELLIA:—
fasciculata, Retz. Obs. iv. 28 = Lepidagathis fasciculata.
filicifolia, Salzm. ex Nees, in Mart. Fl. Bras. ix. 22 = Hygrophila costata.
filiformis, Heyne, ex Steud. Nom. ed. II. ii. 481 = Asystasia violacea.
filiformis, Klein, ex Nees, in DC. Prod. xi. 165 = Asystasia coromandeliana.
fimbriata, Wall. Cat. n. 2363 = Strobilanthes fimbriata.
flaccida, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 91 = Hemigraphis flaccida.
flagelliformis, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 471. —Ins. Molucc.
flava, *Pers. Syn.* ii. 177.—Ind. or.
flava, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 43 = Hemigraphis flava.
flava, Salzm. ex Nees, in DC. Prod. xi. 152 = ochroleuca.
flexuosa, Wall. Cat. n. 2412 = Strobilanthes pentstemonoides.
floribunda, *Hook. Bot. Misc.* ii. (1831) 236.—Peruv.
foetida, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 656 = rubicaulis.
foliosa, Schweinitz, ex Nees, in DC. Prod. xi. 122 = strepens.
formosa, *Andr. Bot. Rep.* t. 610.—Bras.
formosa, Humb. & Bonpl. Pl. Aequin. i. 169 = Aphelandra formosa.
formosa, Ledeb. ex Steud. Nom. ed. II. ii. 481, nomen (Quid?).—Hab.?
fragrans, Forst. f. Prod. 44 = Limnophila serrata.
fulgida, *Andr. Bot. Rep.* t. 527.—N. Granat.
Gairae, Rohr, ex Nees, in DC. Prod. xi. 466 = Blechum Brownei.
geminiflora, *H. B. & K. Nov. Gen. et Sp.* ii. 240.—Am. trop.
gemmipara, *Nees, in DC. Prod.* xi. 220, in syn.—Mexic.
geniculata, Boj. ex Nees, l. c. 107 (= *Dyschoriste geniculata*).—Madag.
geniculata, Vell. Fl. Flum. 265; vi. t. 93 = Stenandrum mandioccanum.
gigantea, Humb. & Bonpl. Pl. Aequin. ii. 75. t. 102 = Trichanthera gigantea.
glabra, Heyne, ex Roth, Nov. Pl. Sp. 312 = glandulosa.
glabrata, Herb. Petrop. ex Nees, in Mart. Fl. Bras. ix. 29 = solitaria.
glabrata, Wall. Cat. n. 2390 = Strobilanthes glabratus.
glandulosa, *Roth, Nov. Pl. Sp.* 312 (Gen.?).—Ind. or.
glaucescens, Wall. ex Nees, in Wall. Pl As. Rar. iii. 85, in syn. = Hemigraphis glaucescens.
glomerata, Herb. Madr. ex Nees, in Wall. l. c. 94 = Phaylopsis parviflora.
glomerata, Wall. Cat. n. 2361 = Strobilanthes glomeratus.
glutinosa, Roxb. ex Nees, in DC. Prod. xi. 261 = Phaylopsis parviflora.
glutinosa, Wall. Cat. n. 2350 = Strobilanthes glutinosus.
gossypina, Nees, in Wall. Pl. As. Rar. i. 38. t. 42; Wall. Cat. n. 2332 = Aechmanthera tomentosa.
grandiflora, Blanch. ex Nees, in Mart. Fl. Bras. ix. 30 = speciosa.
grandiflora, Pers. Syn. ii. 175 = longiflora.
grandiflora, *Poir. Encyc.* vi. 340.—Ins. Guadeloup.
grandiflora, Ruiz, ex Nees, in DC. Prod. xi. 134 = candida.
grandiflora, Salzm. ex Nees, in Mart. Fl. Bras. ix. 31 (= *Dipteracanthus subringens*).—Bras.
grandis, Hort. ex Hook. Bot. Mag. t. 4366 = Strobilanthes lactatus.
graveolens, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 202.—Nepal.
Griffithiana, Nees, in DC. Prod. xi. 149 = Hemigraphis Griffithiana.
guttata, *Forsk. Fl. Aegypt. Arab.* 114.—Arab.
Hamiltoniana, Steud. Nom. ed. II. ii. 481 = Strobilanthes coloratus.
Herbstii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 75.—Bras.
hirsuta, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 270, in syn. = Hemigraphis Decaisneana.
hirsuta, Ell. Sketch, ii. 109 = ciliosa.
hirsuta, Nees, in DC. Prod. xi. 148 = Hemigraphis hirsuta.
hirsuta, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 51 = Hygrophila phlomoides.

RUELLIA:—
hirsuta, Vell. Fl. Flum. 266; vi. t. 96 = geminiflora.
hirsuto-glandulosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 505.—Mexic.
hirta, D. Don, Prod. Fl. Nep. 119 = Strobilanthes nutans.
hirta, Vahl, Symb. Bot. iii. 84. t. 67 = Hemigraphis hirta.
hispida, DC. ex Nees, in DC. Prod. xi. 198 = lactea.
hispida, Rich. ex Nees, l. c. 87 = Hygrophila hispida.
hispida, Vahl, ex Nees, l. c. 137 = geminiflora.
hispidosa, Boj. ex Nees, l. c. 262 = Phaylopsis parviflora.
Hochstetteri, Steud. Nom. ed. II. ii. 481 = Calophanes radicans.
Hoffmannseggii, Steud. l. c. = lactea.
homonyma, Steud. l. c. = Asystasia Kunthiana.
Hookeriana, *Hemsl. Biol. Centr. Am. Bot.* ii. 505.—Mexic.
Humboldtiana, *Klotzsch, ex Nees, in DC. Prod.* xi. 207, in syn.—N. Granat.
humifusa, *Hemsl. Biol. Centr. Am. Bot.* ii. 505.—Mexic.
humifusa, *Pers. Syn.* ii. 177.—Ins. Seychell.
humilis, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 182 = ciliosa.
humilis, *Pohl, ex Nees, in Mart. Fl. Bras.* ix. 40.—Bras.
humistrata, Michx. Fl. Bor. Am. ii. 23 = Calophanes humistratus.
Huttonii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 25.—Afr. austr.
hybrida, Pursh, Fl. Am. Sept. ii. 420 = ciliosa.
hygrophila, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 65. —Bras.
hygrophiloides, *Hemsl. Biol. Centr. Am. Bot.* ii. 505. —Mexic.
hyssopifolia, Hochst. ex Nees, in DC. Prod. xi. 111 = Calophanes hyssopifolius.
imbricata, Forsk. Fl. Aegypt. Arab. 113 = Phaylopsis longifolia.
imbricata, Hook. ex Nees, in DC. Prod. xi. 181 = Strobilanthes verruculosus.
imbricata, Vahl, Symb. Bot. ii. 73 = Phaylopsis parviflora.
imbricata, Wall. Cat. n. 7156 = Strobilanthes imbricatus.
inaperta, *Spreng. in Schrad. Journ.* (1800) II. 198.—Hab.?
incana, Blume, Cat. Gew. Buitenz. 85 = Lepidagathis incana.
incana, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 408 = Hygrophila incana.
indigofera, Griff. Journ. 237 = Strobilanthes flaccidifolius.
indigotica, Fortune, Resid. Chin. 158 = Strobilanthes flaccidifolius.
inflata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 110.—Guian.
infundibuliformis, Andr. Bot. Rep. t. 542; Roxb. Fl. Ind. iii. 41 = Crossandra undulaefolia.
insignis, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotra.
intrusa, Forsk. Fl. Aegypt. Arab. 113 = Asystasia coromandeliana.
inundata, *H. B. & K. Nov. Gen. et Sp.* ii. 239.—N. Granat.
involucrata, Heyne, ex Wall. Cat. n. 2357 = Strobilanthes Heyneanus.
involucrata, *Vahl, Symb. Bot.* iii. 83 (Gen.?).—Ind. or.
isophylla, Steud. Nom. ed. II. ii. 482 = Strobilanthes isophyllus.
Jacquemontiana, Nees, in DC. Prod. xi. 145 = Strobilanthes glutinosus.
japonica, *Thunb. Fl. Jap.* 254.—Japon.
japurensis, Mart. ex Nees, in Mart. Fl. Bras. ix. 29 = solitaria.
Junghuhniana, *Miq. Fl. Ind. Bat.* ii. 790.—Java.
jussieuoides, *Schlecht. in Linnaea*, vi. (1831) 370.—Mexic.
justiciaeflora, Hook. Comp. Bot. Mag. i. (1835) 176 = Hygrophila lacustris.
justicioides, Wall. ex Nees, in Wall. Pl. As. Rar. iii. 94 = Phaylopsis parviflora.
Kunthiana, Wall. Cat. n. 2367 = Strobilanthes Kunthianus.
Kunthiana, Wall. l. c. n. 2419 = Asystasia Kunthiana.

RUELLIA:—
lactea, *Cav. Ic.* iii. 28. t. 255.—Mexic.
lactea, Willd. ex Nees, in DC. Prod. xi. 197 = tuberosa.
lacustris, Schlecht. & Cham. in Linnaea, v. (1830) 96 = Hygrophila lacustris.
lagopodes, Ryan, ex Nees, in DC. Prod. xi. 263 = Lepidagathis alopecuroides.
lamiifolia, Wall. Cat. n. 2347 = Strobilanthes lamiifolius.
lanato-viscosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 506.—Mexic.
latebrosa, Dalz. & Gibs. Bomb. Fl. 186 = Hemigraphis ebracteolata.
latebrosa, Roth, Nov. Pl. Sp. 307 = Hemigraphis latebrosa.
latebrosa, Roxb. Hort. Beng. 46; Fl. Ind. iii. 46 = Hemigraphis hirta.
latifolia, Roth, ex Nees, in DC. Prod. xi. 146 = Hemigraphis hirta.
leptostachya, Wall. Cat. n. 2411 = Dianthera Wallichii.
lilacina, Hook. Bot. Mag. t. 4147 = solitaria.
littoralis, Linn. f. Suppl. 289 = Calophanes littoralis.
liturata, Hort. Berol. ex Nees, in Mart. Fl. Bras. ix. 29 = solitaria.
longepetiolata, *Hemsl. Biol. Centr. Am. Bot.* ii. 506.—Mexic.
longiflora, DC. ex Nees, in DC. Prod. xi. 216 = curviflora.
longiflora, Drumm. ex Nees, l. c. 123 = ciliosa.
longiflora, Hort. ex Lindl. Bot. Reg. (1846) sub t. 13 = solitaria.
longiflora, Nees, in DC. Prod. xi. 140 = Whitfieldia longifolia.
longiflora, Pers. Syn. ii. 176 = longifolia, *Rich.*
longiflora, Salzm. ex Nees, in DC. Prod. xi. 210 (= *Arrhostoxylum glabrum*).—Bras.
longiflora, *Vahl, Symb. Bot.* i. 45.—Arab.
longifolia, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 460.—Beluchist.
longifolia, T. Anders. l. c. vii. (1864) 27 = Whitfieldia longifolia.
longifolia, Raeusch. Nom. ed. III. 180 = cayennensis.
longifolia, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 110.—Guian.
longifolia, Roth, Nov. Pl. Sp. 306 = Hygrophila angustifolia.
longifolia, Roxb. Hort. Beng. 46; Fl. Ind. iii. 50 = Hygrophila spinosa.
longifolia, Thunb. ex Nees, in DC. Prod. xi. 182 = Strobilanthes cernuus.
Lorentziana, *Griseb. in Goett. Abh.* xxiv. (1879) 259.—Reg. Argent.
lupulina, Wall. Cat. n. 2355 = Strobilanthes lupulinus, Heyneanus.
luzona, Nees, in DC. Prod. xi. 151 = Hemigraphis luzona.
Lysimachia, Hort. Monsp. ex Nees, l. c. 198 = tuberosa.
macrantha, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 37.—Bras.
macrocalyx, Ruiz, ex Nees, in DC. Prod. xi. 218 = Macrostegia Ruiziana.
macrocarpa, Wall. Cat. n. 2348 = Asystasia macrocarpa.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 241 = fulgida.
macrophylla, *Vahl, Symb. Bot.* ii. 72. t. 39.—Am. trop.
macrophylla, Willd. ex Nees, in DC. Prod. xi. 207 (= *Stemonacanthus Willdenoviana*).—Hab.?
macrosiphon, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 92.—Burma.
maculata, Tafalla, ex Nees, in DC. Prod. xi. 290 (= *Lagochilium maculatum*).—Hab.?
maculata, Wall. Pl. As. Rar. iii. 33. t. 250 = Strobilanthes maculatus.
maderensis, Hort. ex Nees, in DC. Prod. xi. 159 = solitaria.
madurensis, Poir. Encyc. vi. 341 = Calophanes littoralis.
magnifica, Mart. ex Nees, in Mart. Fl. Bras. ix. 37 = macrantha.
Maranhonis, Pohl, ex Nees, in DC. Prod. xi. 142 = viscidula.
matutina, Hochst. & Steud. ex Presl, Bot. Bemerk. 94 = patula.
menthoides, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 72.—Bras.

RUELLIA:—
Methia, Buch.-Ham. ex Wall. Cat. n. 2384 = prostrata.
microphylla, Cav. Ic. vi. 63. t. 586. f. 2 = Calophanes microphylla.
microphylla, Vahl, ex Nees, in DC. Prod. xi. 106 = Calophanes littoralis.
modesta, Mart. ex Nees, l. c. 465 = Pentstemonacanthus modestus.
mollissima, Klein, ex Nees, l. c. 91 = Hygrophila incana.
mollissima, Vahl, Symb. Bot. iii. 82 = Hypoëstes mollissima.
monadelpha, Wall. ex Steud. Nom. ed. II. ii. 482, sphalm. = Strobilanthes monadelphus.
monanthos, *Boj. ex Nees, in DC. Prod.* xi. 125, in syn. —Madag.
montana, Wall. Cat. n. 2391 = Barleria montana.
Moritziana, Klotzsch, ex Nees, in DC. Prod. xi. 137 = geminiflora.
mucronata, Wall. Cat. n. 2366 = Lepidagathis purpuricaulis, hyalina.
multicaulis, Hochst. in Flora, xxiv. (1841) i. Intell. 24 = Calophanes multicaulis.
mutabilis, *Zipp. in Alg. Konst- en Letter-Bode* (1829), i. 297, nomen.—Malaya.
mysurensis, Roth, Nov. Pl. Sp. 303 = Asystasia Lauriana.
Nagchana, Buch.-Ham. ex Wall. Cat. n. 2396 = Calophanes Nagchana.
nana, Nees, in Linnaea, xv. (1841) 355 = Crabbea nana.
napifera, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 573.—Java.
Neesiana, Klotzsch, ex Schlecht. in Linnaea, xiv. (1840) 289 = tetragona, *Link*.
Neesiana, Wall. Pl. As. Rar. i. 73. t. 83 = Asystasia Neesiana.
neglecta, Steud. Nom. ed. II. ii. 482 = Calophanes Nagchana.
nemoralis, Hilsenb. & Boj. ex Nees, in DC. Prod. xi. 159 (= *Rhytiglossa nemoralis*).—Afr. trop.?
nemoralis, Vahl, ex Nees, l. c. 165 = Asystasia coromandeliana.
nemorosa, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 574.—Java.
nemorum, Russ. ex Wall. Cat. n. 2399 C = Asystasia intrusa.
Nilbutia, Buch.-Ham. ex Wall. l. c. n. 2366 D = Lepidagathis purpuricaulis.
noctiflora, *A. Gray, Syn. Fl. N. Am.* ii. i. 326.—Am. bor.
nova, *Colla, Hort. Ripul. App.* ii. 356, nomen.—Hab.?
nubica, *Delile, Cent. Pl. Méroé*, 74.—Aegypt.
nutans, Steud. Nom. ed. II. ii. 482 = Strobilanthes glomeratus.
obliqua, *Pers. Syn.* ii. 177.—Java.
obliqua, Wight, ex Wall. Cat. n. 2399 D = Asystasia coromandeliana.
oblongifolia, Kinn, ex Nees, in DC. Prod. xi. 122 = strepens.
oblongifolia, Michx. Fl. Bor. Am. ii. 23 = Calophanes oblongifolius.
oblongifolia, Rafin. ex Nees, in DC. Prod. xi. 123 = ciliosa.
obovata, *Blume, ex Steud. Nom.* ed. II. ii. 482, nomen. —Java.
obovata, Buch.-Ham. in Wall. Cat. n. 2375 = Cardanthera triflora.
obovata, Roxb. Hort. Beng. 46; Fl. Ind. iii. 51 = Hygrophila phlomoides.
obovata, Vahl, ex Nees, in DC. Prod. xi. 75 = Ebermaiera glauca.
obscura, Russ. ex Wall. Cat. n. 2382 = Hemigraphis latebrosa.
obtusa, *Nees, in DC. Prod.* xi. 152.—N. Granat.
ochroleuca, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 56.—Bras.
ocymoides, Cav. Ic. v. 9 = patula.
ovalifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 506.—Mexic.
ovata, Cav. Ic. iii. 28. t. 254 = Calophanes linearis.
ovata, Delar. ex Nees, in DC. Prod. xi. 122 = strepens.
ovata, Gouan, ex Nees. l. c. 127 = patula.
ovata, *Thunb. Prod. Pl. Cap.* 104.—Afr. austr.
ovata, Willd. ex Nees, in Mart. Fl. Bras. ix. 49 = lactea.
pallida, *Vahl, Symb. Bot.* ii. 72.—Arab.

RUELLIA:—
pallida, Willd. ex Nees, in DC. Prod. xi. 126 = prostrata.
Panichanga, Buch.-Ham. ex Wall. Cat. n. 2387 = Strobilanthes Panichanga.
paniculata, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 409 = Nomaphila stricta.
paniculata, H. B. & K. Nov. Gen. et Sp. ii. 239 (= *Dipteracanthus turbacensis*).—N. Granat.
paniculata, Linn. Herb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 554 = Peristrophe bicalyculata.
paniculata, *Linn. Sp. Pl.* 635.—Am. trop.
paniculata, Scop. ex Nees, in DC. Prod. xi. 197 = tuberosa.
paniculata, Vahl, ex Nees, l. c. 141 = albicaulis.
paniculata, Wall. ex Nees, in Wall. Pl. As. Rar. iii. 92 = Barleria tomentosa.
parabolica, Nees, in DC. Prod. xi. 144 = Hemigraphis parabolica.
Parryi, *A. Gray, Syn. Fl. N. Am.* ii. i. 326.—Texas.
parva, *Hemsl. Biol. Centr. Am. Bot.* ii. 506.—Mexic.
parviflora, Baldw. ex Nees, in DC. Prod. xi. 107 = Calophanes oblongifolius.
parviflora, Hort. Tonell. ex Nees, l. c. 123 = Hemigraphis latebrosa.
patula, Hort. Par. ex Nees, in DC. Prod. xi. 126 (= *Dipteracanthus dubius*).—Hab.?
patula, *Jacq. Misc.* ii. 358.—Abyss.; Arab.; Ind. or.; Burma.
patula, Salzm. ex Nees, in DC. Prod. xi. 135 = trivialis.
pauciflora, Wall. Cat. n. 2369 = Codonacanthus pauciflorus.
Pavala, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 47 = Hemigraphis latebrosa.
pectinata, Wall. Cat. n. 2356 = Strobilanthes pectinatus.
pedunculata, *Torr. ex A. Gray, Syn. Fl. N. Am.* ii. i. 325.—Am. bor.
pedunculosa, *Nees, in DC. Prod.* xi. 211, in syn.—Peruv.
penstemonoides, Wall. Cat. n. 2340 = Strobilanthes penstemonoides.
persica, Burm. f. Fl. Ind. 135. t. 42. f. 1 = Blepharis edulis.
persicifolia, Lindl. in Bot. Reg. t. 955; Griff. Itin. Notes, 70 = Strobilanthes anisophyllus.
petiolaris, Boj. ex Nees, in DC. Prod. xi. 262 = Phaylopsis longifolia.
petiolata, Boj. ex Nees, l. c. 726 = praec.
phlomoides, Wall. Cat. n. 2376 = Hygrophila phlomoides.
picta, *Lodd. Bot. Cab.* t. 1448.—Ins. S. Doming.
pilosa, Herb. Vindob. ex Nees, in DC. Prod. xi. 200. (= *Siphonacanthus pubens*).—Bras.
pilosa, *Linn. f. Suppl.* 290.—Afr. austr.
pilosa, *Pav. ex Nees, in DC. Prod.* xi. 127, in syn.—Mexic.
pilosa, Vell. Fl. Flum. 268; vi. t. 100 (= *Dipteracanthus Neesianus*).—Bras.
polysperma, Roth, Nov. Pl. Sp. 305 = Hygrophila Serpyllum.
porrigens, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 41.—Bras.
Portellae, *Hook. f. Bot. Mag.* t. 6498.—Bras.
primulacea, *F. Muell. ex Benth. Fl. Austral.* iv. 546. —Austral.
primulifolia, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 382 = Hemigraphis primulaefolia.
prismatica, Vell. Fl. Flum, 267; vi. t. 98 = Aphelandra prismatica.
prostrata, Pav. ex Nees, in DC. Prod. xi. 198 = lactea.
prostrata, *Poir. Encyc.* vi. 349.—Ind. or; Afr. trop.
pubescens, Pers. Syn. ii. 176 = pilosa, *Linn.*
pubiflora, *Griseb. in Goett. Abh.* xxiv. (1879) 259.—Reg. Argent.
pulchella, Schott, ex Steud. Nom. ed. II ii. 482 = rubicaulis.
pulcherrima, *T. Anders. ex Hemsl. Biol. Centr.-Am. Bot.* ii. 507.—Mexic.
pumila, *Pav. ex Nees, in DC. Prod.* xi. 199, in syn.—Peruv.
Pumilio, R. Br. Prod. 479 = australis.
punctata, Nees, in DC. Prod. xi. 147 = Strobilanthes anceps.
Purdieana, *Hook. Bot. Mag.* t. 4298.—N. Granat.?
Puri, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 35.—Bras.

RUELLIA:—
purpuricaulis, Wall. ex Nees, in Wall. Pl. As. Rar. iii. 96 = Lepidagathis purpuricaulis.
quadrangularis, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 494 = Asystasia chelonoides.
quadrangularis, Vell. Fl. Flum. 267; vi. t. 97 = Aphelandra sciophila.
quadrangularis, Wall. Cat. n. 2358 = Strobilanthes quadrangularis.
quadrifaria, Wall. l. c. n. 7152 = Hemigraphis quadrifaria.
quadrivalvis, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 291 = Hygrophila quadrivalvis.
quaterna, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 284 = Asystasia scandens.
quitensis, H. B. & K. Nov. Gen. et Sp. ii. 240 = Calophanes quitensis.
quitensis, Schlecht. & Cham. in Linnaea, v. (1830) 96 = Calophanes Schiedeanus.
racemosa, Roxb. Hort. Beng. [95]; Fl. Ind. iii. 42 = Ebermaiera racemosa.
radicalis, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. iii. 96 = Lepidagathis Hamiltoniana.
radicans, Hochst. in Flora, xxiv. (1841) i. Intell. 24 = Calophanes radicans.
radicans, Wall. Cat. n. 2410 = Hygrophila angustifolia.
radiciflora, Vahl, ex Nees, in DC. Prod. xi. 160 = Lepidagathis cristata.
rasa, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 73. —Bras.
ravaccensis, *Nees, in DC. Prod.* xi. 144.—Ins. Rawak.
reflexa, Royle, ex Nees, in DC. l. c. 194 = Strobilanthes alatus.
reniformis, Wall. ex Nees, in Wall. Pl. As. Rar. iii. 94 = Phaylopsis parviflora.
repanda, Blume, Bijdr. 794 = Blumeana.
repanda, *Linn. Sp. Pl.* ed. II. 886.—Malaya.
repens, Blanco, Fl. Filip. ed. I. 493 = Hemigraphis repens.
repens, Burm. f. Fl. Ind. t. 41. f. 2 = Calophanes Nagchana.
repens, Hort. Madr. ex Nees, in DC. Prod. xi. 124 = prostrata.
repens, *Linn. Mant.* i. 89.—Burma; Malaya.
repens, Ruiz, ex Nees, in DC. Prod. xi. 109 = Calophanes repens.
repens, Wall. Cat. n. 2384 = prostrata, suffruticosa.
reptans, Forst. f. Prod. 44 = Hemigraphis reptans.
retrofacta, Wall. Cat. n. 7165 = Lepidagathis purpuricaulis.
rhamnifolia, Willd. ex Nees, in DC. Prod. xi. 142 = albicaulis.
rhytiphylla, Nees, l. c. 150 = Hemigraphis rhytiphylla.
ringens, Linn. Sp. Pl. 635 = Hygrophila angustifolia.
ringens, Roxb. Hort. Beng. 46; Fl. Ind. iii. 44 = prostrata.
rivinaefolia, Wall. Cat. n. 2413 = Justicia vasculosa.
rivularis, Schlecht. in Linnaea, vii. (1832) 396 = Hygrophila rivularis.
Rohrii, Vahl, ex Nees, in DC. Prod. xi. 466 = Blechum Brownei.
rosea, *Hemsl. Biol. Centr. Am. Bot.* ii. 507.—Mexic.
rosea, Mart. ex Nees, in Mart. Fl. Bras. ix. 61 (= *Arrhostoxylum roseum*).—Bras.
rosea, Wall. ex Nees, in DC. Prod. xi. 177 = Strobilanthes roseus.
rotundifolia, D. Don, Prod. Fl. Nep. 120 = Strobilanthes lamiifolius.
rubicaulis, *Cav. Ic.* v. 10.—Mexic.
rubicunda, Wight, ex Wall. Cat. n. 2337 = Strobilanthes rubicundus.
rubra, *Aubl. Pl. Gui.* ii. 666.—Guian.
rubra, Nees, in DC. Prod. xi. 211. (= *Arrhostoxylum Haenkeanum*).—Peruv.
rubra, Willd. ex Nees, l. c. 209. (= *Arrhostoxylum Willdenovianum*).—Ind. occ.
rufescens, Roth, Nov. Pl. Sp. 304 = Strobilanthes rufescens.
rupestris, T. Anders. in Journ. Linn. Soc. ix. (1867) 462 = Hemigraphis ebracteolata.
ruprestris, Sw. Prod. Veg. Ind. Occ. 93 = Stenandrium rupestre.
Russeliana, Wall. Cat. n. 2400 = Stenosiphonium Russellianum.
Sabiniana, Wall. l. c. n. 2338 = Strobilanthes Sabinianus.
salicifolia, Bert. ex Nees, in DC. Prod. xi. 92 = Hygrophila portoricensis.

RUELLIA:—
salicifolia, Boj. ex Nees, in DC. Prod. xi. 511 = Isoglossa angusta.
salicifolia, Herb. Madr. ex Wall. Cat. n. 2374 B = Hygrophila quadrivalvis.
salicifolia, Salzm. ex Nees, in DC. Prod. xi. 88 = Hygrophila costata.
salicifolia, Vahl, Symb. Bot. iii. 84 = Hygrophila angustifolia.
sanguinea, *Griseb. in Goett. Abh.* xxiv. (1879) 260.—Reg. Argent.
sarmentosa, Nees, in Wall. Pl. As. Rar. iii. 83 = Hemigraphis hirta.
satpoorensis, Wawra, in Oestr. Bot. Zeitschr. xxxi. (1881) 281 = Hemigraphis latebrosa.
saxicola, Boj. ex Nees, in DC. Prod. xi. 503 = Hypoëstes saxicola.
scabra, Gaudich. ex Nees, l. c. 150 = Hemigraphis Decaisneana.
scabra, Juss. ex Pers. Syn. ii. 178 = obliqua.
scabra, Wall. Cat. n. 2377 = Strobilanthes heteromallus.
scabra, Wall. l. c. n. 2393 = Strobilanthes scaber.
scabrida, Wall. l. c. n. 2393 C = Strobilanthus scaber.
scabrosa, Sw. Fl. Ind. Occ. ii. 1074 = Stenandrium scabrosum.
scariosa, Wall. Cat. n. 2354 = Lepidagathis scariosa.
scariosa, Wall. l. c. n. 7166 = Lepidagathis falcata.
Schlechtendaliana, *Hemsl. Biol. Centr. Am. Bot.* ii. 507.—Mexic.
sclerochiton, *S. Moore, in Journ. Bot.* xviii. (1880) 7.—Afr. trop.
seclusa, *S. Moore, l. c.* xiv. (1876) 208.—China.
secunda, Blanco, Fl. Filip. ed. I. 495 = Lepidagathis secunda.
secunda, Vahl, Symb. Bot. iii. 84 = Asystasia coromandeliana.
secunda, Wall. Cat. n. 2401 = Asystasia intrusa.
secunda, Wight, ex Nees, in Wall. Pl. As. Rar. iii. 99 = Strobilanthes exareolatus.
secundiflora, Boj. ex Nees, in DC. Prod. xi. 166 = Asystasia coromandeliana.
semitetrandra, *Jacq. ex G. Don, in Sweet, Hort. Brit.* ed. III. 555, nomen.—Hab. ?
serpens, *Nees, in DC. Prod.* xi. 145.—Java.
serrata, Thunb. in Nov. Act. Soc. Sc. Upsal. vii. (1815) 143 = Bonnaya penicillata.
sesamoides, Wall. Cat. n. 2408 = Strobilanthes consanguineus, Stenosiphonium Russellianum.
sessilis, Balb. ex Nees, in DC. Prod. xi. 198 = lactea.
setigera, Pers. Syn. ii. 176 = Calophanes Persoonii.
setigera, Wight, ex Nees, in Wall. Pl. As. Rar. iii. 81 = Calophanes Nagchana.
setigera, Zeyh. ex Nees, in DC. Prod. xi. 391 = Adhatoda divaricata.
Sibua, Buch.-Ham. ex Wall. Cat. n. 2384 F = suffruticosa.
silvaccola, Mart. ex Nees, in DC. Prod. xi. 213 = formosa.
simplex, *Wright, in Sauv. Fl. Cub.* 97.—Cuba.
siphonantha, Hilsenb. & Boj. ex Nees, in DC. Prod. xi. 112 = Calophanes siphonanthus.
solitaria, *Vell. Fl. Flum.* 266; vi. t. 95.—Bras.
speciosa, Mart. ex Nees, in DC. Prod. xi. 215 = formosa.
speciosa, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 30.—Bras.
speciosa, Schott, ex Nees, in DC. Prod. xi. 118 = macrantha.
spicata, Nees, in DC. Prod. xi. 161 (= *Lagochilium spicatum*).—Mexic.
spicata, Roth, Nov. Pl. Sp. 310 = Strobilanthes consanguineus.
spicata, Vell. Fl. Flum. 265; vi. t. 92 = Geissomeria distans.
spiciflora, *F. Muell. ex Benth. Fl. Austral.* iv. 547.—Austral.
spinescens, *Thunb. Prod. Pl. Cap.* 104.—Afr. austr.
stemonacanthoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 507.—Costa Rica.
strepens, Beck, Bot. N. & Mid. St. 285 = ciliosa.
strepens, Bert. ex Nees, in DC. Prod. xi. 136 = domingensis.
strepens, Forsk. Fl. Aegypt. Arab. 114 = pallida.
strepens, Hort. Madrit. ex Nees, in DC. Prod. xi. 198 = tuberosa.
strepens, *Linn. Sp. Pl.* 634.—Am. bor.; Ind. occ.
strepens, Pav. ex Nees, in DC. Prod. xi. 136 = geminiflora.
strepens, Sw. ex Nees, Mart. Fl. Bras. ix. 40 = humilis.
strepens, Wight, ex Nees, in DC. Prod. xi. 127 = patula.
strigosa, Nees, l. c. 148 = Hemigraphis strigosa.

RUELLIA:—
strobilina, Wall. Cat. n. 2362 = Strobilanthes nutans.
suaveolens, Span. in Linnaea, xv. (1841) 331 = Striga curviflora ?
subacaulis, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) *App.* 12.—Hab. ?
subcapitata, *Hemsl. Biol. Centr. Am. Bot.* ii. 507.—Nicaragua.
subdentata, *Klotzsch, ex Nees, in Mart. Fl. Bras.* ix. 33.—Bras.
subsericea, Wall. Cat. n. 2416 = Stenosiphonium Russellianum.
suffruticosa, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 53.—Ind. or.
sumatrensis, *Roth, Nov. Pl. Sp.* 311.—Sumatra.
sylvaccola, Mart. ex Nees, in Mart. Fl. Bras. ix. 60 = formosa.
tanacetifolia, Herb. Vahl, ex Nees, in DC. Prod. xi. 68 = Cardanthera triflora.
tentaculata, Linn. Cent. Pl. ii. 22; Amoen. Acad. iv. 320 = Haplanthus tentaculatus.
tenuis, Buch.-Ham. ex Wall. Cat. n. 2366 C = Lepidagathis purpuricaulis.
tetragona, Blume, Cat. Gew. Buitenz. 85 = Blumei.
tetragona, *Link, Enum. Hort. Berol.* ii. 133.—Bras.
tetragona, *Thunb. Fl. Javan.* 22.—Java; Zeylan.
tetragonostachya, Wall. Cat. n. 7168 = Neuracanthus tetragonostachyus.
tetrasperma, *Champ. ex Benth. in Kew Journ.* v. (1853) 132.—China.
thomana, *Spreng. in Schrad. Journ.* (1800) II. 199.—Ins. S. Thomas.
thunbergiaeflora, T. Anders. in Journ. Linn. Soc. vii. (1864) 24 = Distichocalyx thunbergiaeflora.
thymifolia, Herb. Vahl, ex Nees, in DC. Prod. xi. 462 = Calophanes Persoonii.
tomentosa, Wall. Cat. n. 2333 = Aechmanthera tomentosa.
trichotoma, *Nees, in DC. Prod.* xi. 149.—Java.
triflora, DC. ex Nees, l. c. 122 = ciliosa.
triflora, Pav. ex Nees, l. c. 198 = tuberosa.
triflora, Roxb. Hort. Beng. 46; Fl. Ind. iii. 52 = Cardanthera triflora.
trinervis, Buch.-Ham. ex Wall. Cat. n. 2500 A = Lepidagathis trinervis.
trivialis, *Blanch. ex Nees, in Mart. Fl. Bras.* ix. 39.—Bras.
tuberosa, *Linn. Sp. Pl.* 635.—Am. bor. et trop.
tubiflora, *H. B. & K. Nov. Gen. et Sp.* ii. 241.—N. Granat.
tubiflora, Le Conte, in Ann. Lyc. N. York, i. (1824) 142 = noctiflora.
tubulata, *Pav. ex Nees, in DC. Prod.* xi. 202.—Peruv.
Tweediana, *Griseb. in Goett. Abh.* xxiv. (1879) 259.—Reg. Argent.
uliginosa, Blanco, Fl. Filip. ed. I. 494 = Blechum Brownei.
uliginosa, Linn. f. Suppl. 290 = Cardanthera uliginosa.
uliginosa, Mart. ex Nees, in DC. Prod. xi. 68 = Cardanthera triflora.
uliginosa, Nees, l. c. 149 = discolor.
uliginosa, Wall. Cat. n. 2378, partim = Hygrophila polysperma.
ulmifolia, Wall. l. c. n. 2360 = Strobilanthes rufescens.
umbrosa, Herb. Madr. ex Wall. Cat. n. 2399 B = Asystasia coromandeliana, violacea.
undulata, Vahl, Symb. Bot. iii. 82 = Hygrophila quadrivalvis.
undulata, *Vis. Sem. Rar. Patav.* (1847) 4; *et ex Linnaea*, xxiv. (1851) 181.—Hab. ?
uniflora, Boj. ex Nees, in DC. Prod. xi. 125 = monanthos.
urens, Heyne, ex Roth, Nov. Pl. Sp. 302 = Hemigraphis dura.
urophylla, Wall. Cat. n. 2344 = Strobilanthes urophyllus.
urticifolia, Wall. l. c. n. 2346 = Strobilanthes alatus.
ustulata, Buch.-Ham. ex Wall. Cat. n. 2366 B = Lepidagathis hyalina.
variabilis, Macrae, ex Nees, in DC. Prod. xi. 165 = Asystasia coromandeliana.
variabilis, Zipp. in Hall, Bijdr. Nat. Wetensch. v. (1830) 180 = mutabilis.
varians, Heyne, ex Wall. Cat. n. 2492 B = Daedalacanthus montanus.
varians, Vent. Jard. Cels, 46 = Daedalacanthus nervosus.

RUELLIA:—
variegata, Lubbers, Cat. Pl. Rar. San-Donato (1880), 13 = Strobilanthes maculatus.
venosa, Blume, Bijdr. 795 = Hemigraphis confinis.
venosa, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 423 = Hemigraphis venosa.
ventricosa, *H. B. & K. Nov. Gen. et Sp.* ii. 241.—N. Granat.
ventricosa, Poir. Encyc. vi. 339 = cayennensis,
venusta, *Hance, in Journ. Bot.* vi. (1868) 92.—China.
verticillata, Pav. ex Nees, in DC. Prod. xi. 198 = lactea.
verticillata, Spreng. Syst. ii. 822 = Hygrophila longifolia.
verticillata, Wall. Cat. n. 2409 = Cardanthera verticillata.
verticilliflora, Steud. Nom. ed. II. ii. 483 = Cardanthera verticillata.
villosa, *Pohl, ex Nees, in DC. Prod.* xi. 200.—Bras.
vincaeflora, DC. ex Nees, l. c. 122 = strepens.
vincoides, *Willd. ex Nees, in Mart. Fl. Bras.* ix. 36.—Bras.
vindex, *Mart. ex Nees, l. c.* 42.—Bras.
violacea, *Aubl. Pl. Gui.* ii. 668.—Guian.
violacea, Ruiz, ex Nees, in Mart. Fl. Bras. ix. 29 = solitaria.
viscidula, *Mart. ex Schrank, Pl. Rar. Hort. Monac.* t. 93.—Bras.
viscosa, H. B. & K. Nov. Gen. et Sp. ii. 239 = paniculata.
viscosa, Pav. ex Nees, in Mart. Fl. Bras. ix. 25 = Calophanes Maranhonis.
viscosula, Wall. ex Nees, in DC. Prod. xi. 260 = Lepidagathis fasciculata.
Wightiana, Wall. Cat. n. 2397 = prostrata.
Zeyheri, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 25.—Afr. austr.
zeylanica, Koen. ex Roxb. Hort. Beng. 46; Fl. Ind. iii. 42 = Asystasia coromandeliana.

RUEPPELIA, A. Rich. Tent. Fl. Abyss. i. 203. t. 37 (1847) = **Aeschynomene**, Linn. (Legumin.).
abyssinica, A. Rich. l. c. = A. Ruppellii.

RUGELIA, Shuttlew. ex Chapm. Fl. S. U. St. 246 (1860) = **Senecio**, Tourn. (Compos.).
nudicaulis, Shuttlew. ex Chapm. l. c. = S. Rugelia.

RUGENDASIA, Schiede, ex Schlecht. Hort. Halens. 14 (1841) = **Weldenia**, Schult. f. (Commel.).
majalis, Schiede, ex Schlecht. l. c. = W. candida.

RUGENIA, Neck. Elem. ii. 78 (1790), sphalm. = **Eugenia**, Mich. (Myrtac.).

RUIZIA, Cav. Diss. iii. 117. tt. 36, 37 (1787). *STERCULIACEAE*, Benth. & Hook. f. i. 221.
aurea, Hort. ex Baker, Fl. Maurit. 59 = Cossignia pinnata.
cordata, *Cav. Diss.* iii. 117. t. 36. f. 2.—Ins. Borbon.
diversifolia, Jacq. Hort. Schoenb. iii. t. 295 = variabilis.
diversifolia, *Voigt, in Syll. Ratisb.* ii. (1828) 54.—Hab. ?
integrifolia, *Hort. ex Steud. Nom.* ed. II. ii. 483, nomen.—Ins. Borbon.
laciniata, Cav. Diss. iii. 119 = variabilis.
lobata, *Cav. l. c.* 118. t. 36. f. 1.—Ins. Borbon.
palmata, Cav. l. c. 119 = variabilis.
variabilis, *Jacq. Hort. Schoenb.* iii. 24.—Ins. Borbon.

RUIZIA, Ruiz & Pav. Prod. 135. t. 29 (1794) = **Peumus**, Molina (Laurin.).
fragrans, Ruiz & Pav. Syst. Veg. 267 = P. Boldus.

RULAC, Adans. Fam. ii. 383 (1763) = **Acer**, Tourn. (Sapind.).

RULINGIA, R. Br. in Bot. Mag. t. 2191 (1820). *STERCULIACEAE*, Benth. & Hook. f. i. 226.
ACHILLEOPSIS, Turcz. in Bull. Soc. Nat. Mosc. (1849) II. 9.
althaeifolia, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 151 = grandiflora.
cistifolia, A. Cunn. ex Steetz, in Lehm. Pl. Preiss. ii. 354, sphalm. = hermanniaefolia.
corylifolia, *R. Grah. in Edinb. N. Phil. Journ.* (Jul. 1832) 171.—Austral.

RULINGIA:—
corylifolia, Steud. in Lehm. Pl. Preiss. i. 237 = parviflora.
craurophylla, *F. Muell. Fragm.* ix. 59.—Austral.
crispa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 10 = Commersonia crispa.
cristifolia, A. Cunn. ex Benth. Fl. Austral. i. 240 = hermanniaefolia.
cuneata, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 151.—Austral.
dasyphylla, Sweet, Hort. Brit. ed. I. 55 = pannosa.
densiflora, *Benth. Fl. Austral.* i. 241.—Austral.
grandiflora, *Endl. in Enum. Pl. Hueg.* 12.—Austral.
hermanniaefolia, *Steetz, in Lehm. Pl. Preiss.* ii. 353.—Austral.
hexamera, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 151 = cuneata.
Kempeana, *F. Muell. Fragm.* xi. 113.—Austral.
loxophylla, *F. Muell. l. c.* i. 68.—Austral.
macrantha, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 503.—Madag.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 104.—Madag.
magniflora, *F. Muell. Fragm.* viii. 223.—Austral.
malvaefolia, *Steetz, in Lehm. Pl. Preiss.* ii. 356.—Austral.
nana, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 150 = parviflora.
oblongifolia, Steetz, in Lehm. Pl. Preiss. ii. 353 = hermanniaefolia.
pannosa, *R. Br. in Bot. Mag.* t. 2191.—Austral.
parviflora, *Endl. in Enum. Pl. Hueg.* 12.—Austral.
pauciflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 570.—Austral.
platycalyx, *Benth. Fl. Austral.* i. 240.—Austral.
pulchella, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 10 = Commersonia pulchella.
rotundifolia, *Turcz. l. c.* (1852) II. 152.—Austral.
rugosa, *Steetz, in Lehm. Pl. Preiss.* ii. 352.—Austral.
salvifolia, *Benth. Fl. Austral.* i. 238.—Austral.

RUMEA, Poit. in Mém. Mus. Par. i. (1815) 62. t. 4 = **Xylosma**, Forst. (Bixin.).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 34 = Aberia abyssinica.
chinensis, Wall. ex Steud. Nom. ed. II. ii. 475 (= *Phoberos chinensis*).—China.
cochinchinensis, Wall. ex Steud. l. c. = Scolopia cochinchinensis.
coriacea, Poit. in Mém. Mus. Par. i. (1815) 63. t. 4 = X. nitidum.
crocea, Steud. Nom. ed. II. i. 532, sphalm. = praec.
floribunda, Agardh, Theor. Syst. Pl. t. 13. f. 13–15.—Hab. ?
hebecarpa, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 9 = Aberia Gardnerii.
inermis, DC. Prod. i. 256 = Fluggea microcarpa.
Jangomas, Spreng. Syst. ii. 632.—Cochinch.

RUMEX, Linn. Syst. ed. I (1735). *POLYGONACEAE*, Benth. & Hook. f. iii. 100.
ACETOSA, Neck. Elem. ii. 214 (1790).
ACETOSELLA, Rafin. Fl. Tellur. iii. 46 (1836).
ANALITON, Rafin. l. c. (1836).
ATECOSA, Rafin. l. c. (1836).
EUTRALIA, Rafin. l. c. (1836).
LAPATHUM, [Tourn.] Adans. Fam. ii. 277 (1763).
MENOPHYLA, Rafin. Fl. Tellur. iii. 45 (1836).
OXYLAPATHON, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159.
PATIENTIA, Rafin. Fl. Tellur. iii. 46 (1836).
RHODOPTERA, Rafin. l. c. 45 (1836).
STEINMANNIA, Opiz, Seznam, 93 (1852).
TOMARIS, Rafin. Fl. Tellur. 46 (1836).
VIBONES, Rafin. l. c. 45 (1836).
abortivus ×, Ruhm. in Eichl. Jahrb. Berl. i. (1881) 253.—Europ.
abyssinicus, Desf. Tabl. ed. II. 48 = arifolius.
abyssinicus, *Jacq. Hort. Vindob.* iii. 48. t. 93.—Abyss.
Acetosa, *Linn. Sp. Pl.* 337.—Europ.; As. bor.
Acetosella, Holl, ex Meissn. in DC. Prod. xiv. 72 = suffruticosus.
Acetosella, *Linn. Sp. Pl.* 338.—Europ.; As. bor.
acetoselloides, *Bal. in Bull. Soc. Bot. Fr.* i. (1854) 282.—Oriens.
acetosus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 134 = R. Acetosa.
acidus, Salisb. Prod. 258 = R. Acetosa.

RUMEX:—
aculeatus, Linn. Syst. ed. X. 991 = bucephalophorus.
acuminatus, Campd. Monog. Rum. 116 = R. Acetosa.
acutus, All. Fl. Pedem. ii. 202 = obtusifolius.
acutus, Curt. Fl. Lond. fasc. iii. 21 = sanguineus.
acutus, Fries, ex Meissn. in DC. Prod. xiv. 52 = conspersus.
acutus, *Linn. Sp. Pl.* 335.—Europ.
acutus, Pollich, Hist. Pl. Palat. i. 359 = maritimus.
acutus, Schultz, Prod. Fl. Starg. Suppl. i. 22 = maximus.
acutus, Sm. Fl. Brit. i. 391 = conglomeratus.
acutus, Wahlenb. Fl. Suec. 233 = R. Hydrolapathum.
adulterinus, Wallr. Beitr. Fl. Hercyn. 197 = acutus.
aegyptiacus, Balb. ex Meissn. in DC. Prod. xiv. 58 = pulcher.
aegyptiacus, Bieb. Fl. Taur. Cauc. i. 290 = Marschallianus.
aegyptiacus, *Linn. Sp. Pl.* 335.—Aegypt.
aegyptius, Linn. Syst. ed. X. 990 = praec.
aetnensis, Presl, Delic. Prag. 68 = scutatus.
afer, *Savi, in Atti Stor. Nat. Bologn.* iii. (1830) 90.—Aegypt.
agrestis, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
agrestis, Schur, Enum. Pl. Transs. 580 = obtusifolius.
alexandrinus, *Hort. ex Schult. f. Syst.* vii. 1476, nomen.—Hab. ?
alismaefolius, *Fresen. in Flora*, xxi. (1838) 602.—Abyss.
Allioni, Link, Handb. i. 306 = montanus.
alpestris, Jacq. Enum. Stirp. Vindob. 62 = scutatus.
alpinus, Georgi, Reise, i. 207 = Gmelini.
alpinus, Hook. in Engl. Bot. Suppl. t. 2694 = longifolius.
alpinus, *Linn. Sp. Pl.* 334.—Europ.; Reg. Caucas.
altissimus, *Wood, in Proc. Am. Assoc.* (1853) 177.—Am. bor.
ambiguus, J. Gay, ex Gaud. Fl. Helv. ii. 589, in obs. = acutus.
ambiguus, Gren. in Gren. & Godr. Fl. Fr. iii. 44 = R. Acetosa.
americanus, Campd. Monog. Rum. 151 = vesicarius *vel* scutatus.
amplexicaulis, Lapeyr. Hist. Abr. Pl. Pyr. 200 = montanus.
amplexicaulis, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
amplexicaulis, Tenore, Fl. Nap. iii. 394 = tmoleus.
amurensis, *F. Schmidt, in Maxim. Prim. Fl. Amur.* 228.—Mandchuria.
angustatus, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
angustifolius, *Campd. Monog. Rum.* 63, 73. t. 1. f. 2.—As. Min.
angustifolius, Engelm. ex Meissn. in DC. Prod. xiv. 64 = Engelmanni.
angustissimus, Ledeb. in Mém. Acad. Pétersb. v. (1812) 536 = graminifolius.
Anthoxanthus, Murr. Prod. Gotting. 52 = maritimus.
Anthoxantum, Moench, Enum. Pl. Hass. 174 = praec.
antiscorbuticus, Salisb. Prod. 258 = R. Hydrolapathum.
aquaticus, Campd. Monog. Rum. 100 = maximus.
aquaticus, Linn. Sp. Pl. 336.—Europ.; As. bor.
aquaticus, Sm. Fl. Brit. i. 394 = R. Hydrolapathum.
arboreus, Salisb. Prod. 258 = R. Lunaria.
arcticus, *Trautv. in Middend. Sibir. Reise*, i. II. 29.—Sibir. arct.
arenarius, Poepp. ex Endl. Gen. Suppl. iv. II. 52 = cuneifolius.
Areschougii, *Eckstr. ex Aresch. in Bot. Notiser* (1866), 10, 54.—Europ.
arifolius, Ait. Hort. Kew. ed. I. i. 487; ed. II. ii. 324 = abyssinicus.
arifolius, All. Fl. Pedem. ii. 204 = montanus.
arifolius, *Linn. f. Suppl.* 212.—Afr. austr.
Aristidis, *Coss. in Bull. Soc. Bot. Fr.* v. (1858) 103.—Afr. bor.
armenus, *C. Koch, in Linnaea*, xxii. (1849) 209.—Armenia.
armoraciaefolius, *Neum. in Bot. Notiser* (1885), 155.—Europ.
arvensis, Dulac, Fl. Hautes-Pyr. 165 = R. Acetosella.
atriplicifolius, Reichb. ex Schult. f. Syst. vii. 1460 = pictus.
auratus, Sm. in Rees, Cycl. xxx. n. 30 = sanguineus.
aureus, Hoepfn. in Mag. Naturk. Helv. iv. (1789) 35 = sanguineus.
aureus, Mill. Gard. Dict. ed. VIII. n. 7 = maritimus.
auritus, *Campd. Monog. Rum.* 149.—Oriens.
Berlandieri, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 104 (an seq. ?).—Porto Rico.
Berlandieri, *Meissn. in DC. Prod.* xiv. 45.—Am. bor. occ.; Chili.
bidens, *R. Br. Prod.* 421.—Austral.

RUMEX:—
bidentula, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
biformis, *Lange, Ind. Sem. Hort. Haun.* (1857) 26.—*Cf. Linnaea*, xxix. (1857–58) 733.—Gallia.
biformis, Menyh. Kalocsa Vidék. Növényt. 161 = acutus.
bipinnatus, Linn. f. Suppl. 211 = pictus.
bonariensis, *Campd. Monog. Rum.* 144.—Reg. Argent.
brasiliensis, *Link, Enum. Hort. Berol.* i. 349.—Bras.
brevipes, *Meissn. in DC. Prod.* xiv. 55.—Mexic.
Britanica, Huds. Fl. Angl. ed. I. 135 = R. Hydrolapathum.
Britannica, *Linn. Sp. Pl.* 334.—Am. bor.
Britannicus, Ell. Sketch, i. 414 = verticillatus.
britannicus, Hoffmgg. Verz. Pfl. 239 = R. Britannica.
Brownianus, Raoul, Choix Pl. N. Zel. 42 = flexuosus.
Brownianus, Schult. f. Syst. vii. 1395 = seq.
Brownii, *Campd. Monog. Rum.* 64, 81.—Austral.
Brownii, Schlecht. in Linnaea, xx. (1837) 576 = pulcher.
bucephalophorus, *Linn. Sp. Pl.* 336.—Reg. Mediterr.
bulbosus, *Campd. Monog. Rum.* 151.—Hab. ?
Burchelli, Campd. l. c. 135 = luxurians.
caldeirarum, H. C. Wats. ex Meissn. in DC. Prod. xiv. 43 = aquaticus.
callosissimus, Meissn. in DC. Prod. xiv. 57 = dentatus.
calthaefolius, *Campd. Monog. Rum.* 150.—Aegypt.
campestris, Savi, Fl. Pis. i. 372 = conglomeratus.
Candollii, Mart. ex Meissn. in DC. Prod. xiv. 52 = alpinus.
cassius, *Boiss. Fl. Orient.* iv. 1013.—Syria.
chalepensis, *Mill. Gard. Dict.* ed. VIII. n. 11.—Syria.
chilensis, Herb. Berol. ex Meissn. in DC. Prod. xiv. 44 = longifolius.
chilensis, Hort. Par. ex Moris, in Mem. Acc. Torin. xxxviii. (1835) 46 = cuneifolius.
chinensis, Campd. Monog. Rum. 63, 75 = maritimus.
chrysocarpos, *Moris, Enum. Sem. Hort. Taurin.* (1813); *ex eod. in Mem. Acc. Torin.* xxxviii. (1835) 46. t. 2.—Mexic.
chrystallinus, Auct. = crystallinus.
Claytoni, Campd. Monog. Rum. 66, 99. = R. Britannica.
Commersonianus, D. Dietr. Syn. Pl. ii. 1221 = seq.
Commersonnii, Campd. Monog. Rum. 72, 148 = R. Acetosa.
comosus, Buch.-Ham. ex Meissn. in DC. Prod. xvi. 48 = maritimus.
comosus, Forsk. Fl. Aegypt. Arab. 76 = aegyptiacus.
condylodes, Bieb. Fl. Taur. Cauc. i. 288 = nemorosus.
confertus, Willd. Enum. Hort. Berol. 397 = alpinus.
confusus ×, Simk. in Termész. Füzetek, i. (1877) 238.—Hungar.
conglomeratus, *Murr. Prod. Gotting.* 52.—Europ. austr.; As. occ.
conspersus, *Hartm. Skand. Fl.* ed. I. 147.—Europ.
cordatus, *Desf. Tabl.* ed. I. 40, nomen; *Cat. Hort. Par.* ed. III. 389.—Afr. austr.
cordifolius, *Hornem. ex Campd. Monog. Rum.* 151, nomen; *et ex Reichb. Pl. Crit.* v. 52.—Europ.; As. bor.
creticus, Boiss. Diagn. Ser. I. xii. 102 = tuberosus.
creticus, Campd. Monog. Rum. 149 = bucephalophorus ?
crispatulus, Hort. ex Meissn. in DC. Prod. xiv. 59 = ucranicus.
crispatulus, Michx. Fl. Bor. Am. i. 217 = maritimus.
crispatus, Pers. Syn. i. 395 = maritimus.
crispus, Cham. & Schlecht. in Linnaea, iii. (1828) 59 = longifolius.
crispus, *Linn. Sp. Pl.* 335.—Europ.; As. bor.
crispus, Lour. Fl. Cochinch. 216 = maritimus.
cristatus, *DC. Cat. Hort. Monsp.* (1813) 139.—Hab. ?
cristatus, Fries, Nov. Fl. Suec. 100 = conspersus.
cristatus, *Hort. ex Schult. f. Syst.* vii. 1436 = cordifolius.
cristatus, Wall. Sched. Crit. 163 = acutus.
crystallinus, *Lange, Ind. Sem. Hort. Haun.* (1861) 28.—Austral.
cuneifolius, *Campd. Monog. Rum.* 66, 95.—Am. austr.
Cunninghamii, Meissn. in DC. Prod. xiv. 62 = flexuosus.
daghestanicus, C. Koch, in Linnaea, xxii. (1849) 209 = nemorosus.
dentatus, *Linn. Mant.* ii. 226.—Oriens; Ind. or.
dentatus, Wall. Cat. 1730 (sp. diversa).—Ind. or.
denticulatus, Campd. Monog. Rum. 143 = maritimus.
denticulatus, C. Koch, in Linnaea, xxii. (1849) 208 = pulcher.
dictyocarpus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 192 = obtusifolius.
digynus, Linn. Sp. Pl. 337 = Oxyria digyna.

RUMEX:—
dimorphus, Gren. in Gren. & Godr. Fl. Fr. iii. 43 = arifolius.
Dioscoridis, Hayne, Arzn. Gew. xiii. 5. t. 5 = orientalis.
distans, Dum. Fl. Belg. 17 = longifolius.
divaricatus, Fries, Mant. iii. 25 = obtusifolius.
divaricatus, Linn. Sp. Pl. ed. II. 477 = pulcher.
divaricatus, Thuill. Fl. Par. ed. II. i. 182 = conglomeratus.
domesticus, Hartm. Skand. Fl. ed. I. 148 = longifolius.
Dregeanus, *Meissn. in Linnaea*, xiv. (1840) 496—Afr. austr.
Dregei, Meissn. in DC. Prod. xiv. 68 = praec.
Drummondii, Meissn. in Lehm. Pl. Preiss. ii. 272 = pulcher.
Drummondii, F. Muell. ex Meissn. in DC. Prod. xiv. 61 = praec.?
dubius, Gunn. Fl. Noveg. ii. 26 = maritimus.
dubius, Retz. Fl. Scand. Prod. 64 = conglomeratus.
dumosus, *A. Cunn. ex Meissn. in DC. Prod.* xiv. 62.—Austral.
Ecklonianus, *Meissn. in Linnaea*, xiv. (1840) 493.—Afr. austr.
Ecklonii, Meissn. in DC. Prod. xiv. 50 = praec.
Ehrenbergii, Meissn. l. c. 56 = dentatus.
elbrusensis, *Boiss. Diagn.* Ser. I. xii. 100.—Persia.
elongatus, Guss. Pl. Rar. Neap. (1826) 150. t. 28 = crispus.
Engelmanni, *Ledeb. ex Meissn. in DC. Prod.* xiv. 64.—Am. bor. occ.
erosus, *Hort. ex Schult. f. Syst.* vii. 1476, nomen.—Hab.?
erubescens ×, Simk. in Termesz. Füzetek, i. (1877) 239.—Hungar.
exsanguis, Kit. ex Schult. Oesterr. Fl. ed. II. i. 576 = nemorosus.
falcarius, Willd. ex Ledeb. Fl. Ross. iii. 512 = R. Acetosella.
ferrugineus, Willd. ex Spreng. Syst. ii. 158 = conglomeratus.
fimbriatus, R. Br. Prod. 421 = Brownii.
fimbriatus, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 455 = flexuosus.
fimbriatus, *Poir. Encyc.* v. 65.—Afr. austr.
Fischeri, Reichb. Ic. Pl. Crit. iv. 58 = crispus.
flexuosus, *Soland. ex Forst. f. Prod.* 90.—N. Zel.
floridanus, *Meissn. in DC. Prod.* xiv. 46.—Florida.
foveolatus, Hochst. in Lorent, Wanderungen, 328 = pulcher.
Friesii, Aresch. in Oefvers. Vet. Akad. Handl. Stockh. 1862 (1863) 58, in syn. = aquaticus.
Friesii, Gren. & Godr. Fl. Fr. iii. 36 = obtusifolius.
frutescens, *Thou. Fl. Tristan d'Acugn.* 38.—Ins. Tristan d'Acugn.
garipensis, *Meissn. in Linnaea*, xiv. (1840) 491.—Afr. austr.
gentilis, Loennr. ex Aresch. in Bot. Notiser (1866), 7, 52 = acutus.
giganteus, *Ait. Hort. Kew.* ed. II. ii. 323.—Ins. Sandvic.
glaber, Forsk. Fl. Aegypt. Arab. 75 = Emex spinosa.
glaucus, Jacq. Coll. i. 63 = scutatus.
glomeratus, Schreb. Spicil. Fl. Lips. ed. II. 64 = conglomeratus.
glomeratus, Tenore, Fl. Nap. v. p. vi = sanguineus.
Gmelini, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 100; xxv. (1852) II. 444.—Sibir. baical.
gracilis, Eckstr. in Bot. Notiser (1866), 52 = acutus.
gracilis, Schur, Enum. Pl. Transs. 579 = sylvestris.
graecus, Boiss. & Heldr. Diagn. Ser. II. iv. 80 = orientalis.
gramineus, Pall. ex Ledeb. Fl. Ross. iii. 512 = graminifolius.
graminifolius, *Georgi, Beschr. Russ. Reich.* III. iv. 921.—Kamtschat.
granulosus, Wender. ex Steud. Nom. ed. II. ii. 484, nomen.—Hab.?
halophilus, *F. Muell. Fragm.* iv. 48.—Austral.
hamatus, Trevir. in Nov. Act. Nat. Cur. xiii. (1826) 174 = nepalensis.
hamulosus, Meissn. in DC. Prod. xiv. 693 = nepalensis.
haplorhizus, Czern. ex Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 445 = R. Acetosa.
hastaefolius, Campd. Monog. Rum. 148 = scutatus.
hastatus, *D. Don, Prod. Fl. Nep.* 74.—Reg. Himal.
hastatus, Link, ex Meissn. in DC. Prod. xiv. 70 = scutatus.
hastifolius, Bieb. Fl. Taur. Cauc. i. 290 = scutatus.
hastifolius, Guss. ex Nym. Consp. 635 = intermedius.

RUMEX:—
hastulatus, Baldw. ex Meissn. in DC. Prod. xiv. 64 = Engelmanni.
hastulatus, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
hastulatus, Sm. in Rees, Cycl. xxx. n. 29 = Muehlenbeckia chilensis.
Helolapathum, Drejer, ex Hornem. in Fl. Dan. t. 2288 = aquaticus
heteranthos ×, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 393.—Europ.
heterophylus, Rafin. New Fl. Am. iv. 52 (Quid?).—Am. bor.
heterophyllus, Schultz, Prod. Pl. Starg. Suppl. i. 21 = maximus.
hexagynus, Auct. ex Schult. f. Syst. vii. 1476, nomen (Quid?).—Hab.?
hippiatricus, *Remy, in C. Gay, Fl. Chil.* v. 281.—Chili.
Hippolapathum, Fries, Nov. Fl. Suec. ed. II. 106 = longifolius, aquaticus.
hirsutus, *Tenore, ex Steud. Nom.* ed. II. ii. 484, nomen.—Italia.
hispanicus, C. C. Gmel. Fl. Bad. ii. 112 = montanus.
hispanicus, Koch, Syn. Fl. Germ. ed. II. 709 = R. Acetosa.
horizontalis, *C. Koch, in Linnaea*, xxii. (1849) 211.—As. Min.
hortensis, Vis. ex Nym. Consp. 635 = R. Acetosa.
hostilis, *Lour. Fl. Cochinch.* 217.—Cochinch.
Hydrolapathum, Campd. Monog. Rum. 145 = conspersus.
Hydrolapathum, *Huds. Fl. Angl.* ed. II. 154 (Hydrolapatheum).—Europ.
hydrolapathus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
hymenosepalus, *Torr. Bot. Mex. Bound.* 177.—Am. bor. occ.
incomtus, Weinm. in Flora, iv. (1821) 28 = brasiliensis.
induratus, Boiss. & Reut. Pugill. Pl. Nov. 107 = scutatus.
infestus, Salisb. Prod. 258 = R. Acetosella.
integrifolia, *Rafin. New Fl. Am.* iv. 52.—Am. bor.
intermedius, *DC. Fl. Fr. Suppl.* 369.—Europ. austr.
intermedius, Guss. Fl. Sic. Prod. i. 449 = thyrsoides.
intermedius, Sturm, Deutschl. Fl. Heft 74; et ex Meissn. in DC. Prod. xiv. 65 = R. Acetosa.
italicus, Hort. Par. ex Campd. Monog. Rum. 112 = montanus.
japonicus, Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 56 = crispus.
Klotzschianus, Meissn. in DC. Prod. xiv. 57 = dentatus.
Knafii ×, Čelak. in Arch. Naturw. Landesd. Boehm. ii. (1873) 158.—Europ.
Kunthianus, Schult. f. Syst. vii. 1423 = seq.
Kunthii, Campd. Monog. Rum. 66, 97 = longifolius.
lacerus, Balb. in Mém. Acad. Turin (1802), 335 = pictus.
lacerus, Hort. ex Steud. Nom. ed. II. ii. 485 = bucephalophorus.
lachanus, *Forsk. Fl. Aegypt. Arab.* 209 (sp. dub.).—Anatol.
laevigatus, Willd. ex Spreng. Syst. ii. 159 = obtusifolius.
lanceolatus, *Thunb. Prod. Pl. Cap.* 67.—Afr. austr.
lanuginosus, *Campd. Monog. Rum.* 150.—Oriens.
Lappula, *Meissn. in DC. Prod.* xiv. 58.—Hab.?
latifolius, *Humb. ex Cham. & Schlecht. in Linnaea*, iii. (1828) 61.—Am austr.
latifolius, G. F. W. Mey. Chlor. Hannov. 478 = aquaticus.
lativalvis, *Meissn. in Linnaea*, xiv. (1848) 497.—Afr. austr.
lentatus, *Gouan, Herb. Monsp.* 268 (Quid?).—Hab.?
leucocephalus, Hort. ex Schult. f. Syst. vii. 1473 = bucephalophorus.
limoniastrum, *Jaub. & Spach, Illustr.* ii. 6. t. 106.—Arab.
limosus, Thuill. Fl. Par. ed. II. 182 = palustris.
linearis, *Campd. Monog. Rum.* 90.—Aft. austr.
lingulatus ×, Jungner, in Bot. Notiser (1885), 155.—Europ.
lingulatus, Schur, Enum. Pl. Transs. 580 = crispus.
litoralis, *H. B. & K. Nov. Gen. et Sp.* ii. 181.—Peruv.
longifolius, Auct. ex S. Wats. in Bot. Calif. ii. 8 = occidentalis.
longifolius, DC. Fl. Fr. Suppl. 368 = R. Hydrolapathum.
longifolius, *H. B. & K. Nov. Gen. et Sp.* ii. 180.—Am. austr.

RUMEX:—
longifolius, Krock, Fl. Siles. Suppl. ii. 40 (Quid?).—Europ.
Loureirianus, Schult. f. Syst. vii. 1474 = seq.
Loureirii, Camp. Monog. Rum. 142 = maritimus.
lucidus, Campd. l. c. 150 = pictus.
ludovicianus, Rafin. Fl. Ludov. 30 (Quid?).—Am. bor.
Luederi, Muent. in Act. Congr. Bot. Amst. 1877 (1879) 188 = crispus.
Lunaria, *Linn. Sp. Pl.* 336.—Ins. Canar.
luxurians, Hort. ex Meissn. in DC. Prod. xiv. 70 = scutatus.
luxurians, *Linn. Mant.* i. 64.—Afr. austr.
luxurians, Pourr. ex Nym. Consp. 635 = montanus.
macranthus, Boiss. Diagn. Ser. II. iv. 80 = angustifolius.
macrophyllus, Campd. Monog. Rum. 67 = arifolius.
macrophyllus, Wallr. ex Hayne, Arzn. Gew. xiii. sub t. 5 = orientalis.
macropterus, G. Ehrenb. ex Meissn. in DC. Prod. xiv. 53 = obtusifolius.
maderensis, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 12.—Ins. Canar. et Mader.
magellanicus, *Campd. Monog. Rum.* 144 (sp. dub.).—Reg. Magellan.
magellanicus, *Griseb. in Goett. Abh.* vi. (1854) 118.—Reg. Magellan.
maricola, *Remy, in C. Gay, Fl. Chil.* v. 281.—Chili.
maritimus, *Linn. Sp. Pl.* 335.—Reg. temp. et subtrop.
maritimus, E. Mey. ex Meissn. in Linnaea, xiv. (1840) 491 = garipensis.
maritimus, With. Arr. Brit. Pl. ed. III. ii. 356 = palustris.
Marschallianus, *Reichb. Ic. Pl. Crit.* iv. 58; vi. 13.—Sibir.; Taur.
maximus, Campd. Monog. Rum. 149 = alpinus?
maximus, C. C. Gmel. Fl. Bad. ii. 99 = R. Hydrolapathum.
maximus, *Schreb. in Schweigg. & Korte, Fl. Erlang.* i. 152.—Europ.
membranosus, Poir. Voy. Barb. ii. 155 = bucephalophorus.
menzalensis, Ehrenb. ex Boiss. Fl. Orient. iv. 1013 = dentatus.
mexicanus, *Meissn. in DC. Prod.* xiv. 45.—Mexic.
Meyeri, *Meissn. in Linnaea*, xiv. (1840) 494.—Afr. austr.
Meyerianus, Meissn. in DC. Prod. xiv. 50 = praec.
micranthus, Campd. ex Meissn. l. c. 65 = R. Acetosa.
microcarpus, *Campd. Monog. Rum.* 143.—China.
montanus, Campd. l. c. 149 = monticola.
montanus, *Desf. Tabl.* ed. II. 48.—Europ.
montevidensis, Spreng. Syst. iv. Cur. Post. 144 = cuneifolius.
monticola, *Schult. f. Syst.* vii. 1476.—Italia.
Muelleri, Meissn. in DC. Prod. xiv. 61 = pulcher?
multifidus, All. Fl. Pedem. ii. 205 = intermedius.
multifidus, Linn. Sp. Pl. ed. II. 482 = R. Acetosella, acetoselloides.
multiflorus, Pers. Syn. i. 395 = dentatus.
muricatus, Blanco, Fl. Filip. ed. II. 195 = maritimus.
muricatus, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 340.—Europ.
mutabilis, Campd. ex Meissn. in DC. Prod. xiv. 65 = R. Acetosa.
napalensis, Meissn. in Wall. Pl. As. Rar. iii. 64 = nepalensis.
nebroides, Campd. Monog. Rum. 72, 150 = intermedius.
neglectus, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 493.—N. Zel.
Nemolapathum, Ehrh. Beitr. i. 181 = conglomeratus.
Nemolapathum, Wallr. Sched. Crit. 158 = nemorosus.
nemorosus, *Schrad. ex Willd. Enum. Hort. Berol.* 397.—Europ.; Oriens; Am. bor. et austr.
nemorosus, Stev. ex Meissn. in DC. Prod. xiv. 52 = conspersus.
nepalensis, *Spreng. Syst.* ii. 159.—As. occ.; Ind. or.; Malaya; Afr. austr.
nervosus, *Vahl, Symb. Bot.* i. 27.—Arab.
nipponicus, Franch. & Sav. Enum. Pl. Jap. ii. 471 = pulcher.
nivalis, *Hegetschw. Fl. Schw.* 345.—Europ. austr.
oblongus, Campd. Monog. Rum. 115 = R. Acetosa.
obtusifolius, Auct. ex Meissn. in DC. Prod. xiv. 54 = acutus.
obtusifolius, Fries, ex Meissn. l. c. 50 = sylvestris.
obtusifolius, Herb. Buch.-Ham. ex Wall. Cat. n. 6289 = dentatus.

RUMEX:—
obtusifolius, *Linn. Sp. Pl.* 335.—Europ.; As. et Am. bor.
occidentalis, *S. Wats. in Proc. Am. Acad.* xii. (1877) 253.—Am. bor. occ.
ochotensis, *Hort. ex Schult. f. Syst.* vii. 1476, nomen (Quid?).—Sibir.?
olitoria, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
olympicus, Boiss. Diagn. Ser. I. v. 45 = R. Patientia.
oravicensis, Wierzb. ex Nym. Consp. 634 = sanguineus.
orbiculatus, *A. Gray, Man. Bot. N. U. St.* ed. V. 420.—Am. bor.
orientalis, *Bernh. ex Schult. f. Syst.* vii. 1433.—Graec.; As. Min.; Syria.
Oxylapathum, Gueldenst. ex Ledeb. Fl. Ross. iii. 503 = obtusifolius.
Oxylapathum, Wallr. ex Fries, Nov. Fl. Suec. ed. II. 101 = acutus.
oxyotus, Campd. Monog. Rum. 114 = R. Acetosa.
oxysepalus, Meissn. in Lehm. Pl. Preiss. i. 625 = pulcher.
pallidus, Bigel. Fl. Bost. ed. III. 153 = salicifolius.
paludosus, Huds. Fl. Angl. ed. II. i. 154 = aquaticus.
paludosus, With. Bot. Arr. Brit. Pl. ed. III. ii. 354 = conglomeratus.
paluster, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 353 = seq.
palustris, *Sm. Fl. Brit.* i. 394.—Europ.; As. bor.
palustroides ×, Simk. in Termész. Füzetek, i. (1877) 237.—Hungar.
Papilio, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 260.—Marocc.
papillaris, Boiss. & Reut. Pugill. Pl. Nov. 107 = R. Acetosa.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 160; *Contrib.* 52.—Paraguay.
parvifolius, Willd. ex Schult. f. Syst. vii. 1417 = litoralis.
patens, Ind. Sem. Hort. Berol. (1851); ex Meissn. in DC. Prod. xiv. 54 = obtusifolius.
Patientia, Gaud. ex Meissn. l. c. 46 = magellanicus.
Patientia, *Linn. Sp. Pl.* 333.—Europ. austr.; Oriens.
Patientia, Pall. ex Bieb. Fl. Taur. Cauc. i. 291 = alpinus.
pauciflorus, Campd. Monog. Rum. 104 = alpinus.
paucifolius, *Nutt. ex S. Wats. in Bot. King's Exped.* 314.—Am. bor. occ.
pectinatus, Ehrenb. ex Meissn. in DC. Prod. xiv. 56 = dentatus.
peregrinus, Boiss. Diagn. Ser. I. v. 46 = nepalensis.
peregrinus, *Schult. f. Syst.* vii. 1437.—Hab.?
persicarioides, Forsk. Fl. Aegypt. Arab. 76 = nervosus.
persicarioides, Hort. ex Meissn. in DC. Prod. xiv. 59 = ucranicus.
persicarioides, Linn. Sp. Pl. 335 = maritimus.
peruanus, Meyen, ex Endl. Gen. Suppl. iv. II. 52 = cuneifolius.
pictus, *Forsk. Fl. Aegypt. Arab.* 76.—Syria; Aegypt.
platyphyllus, Aresch. in Ofvers. Vet. Akad. Stockh. xix. (1862) 73 = aquaticus.
polygamus, Cav. Ic. i. 14 = R. Lunaria.
polygonifolius, Hort. Vindob. ex Meissn. in DC. Prod. xiv. 47 = R. Britannica.
pratensis, Dulac, Fl. Hautes-Pyr. 165 = R. Acetosa.
pratensis, Mert. & Koch, Deutschl. Fl. ii. 609 = acutus.
propinquus ×, Aresch. in Bot. Notiser (1840), 22.—Europ.
Pseudo-acetosa, Bertol. Fl. Ital. iv. 252 = R. Acetosa.
Pseudo-alpinus, Hoefft, Cat. Pl. Kursko, 26 = alpinus.
pubescens, C. Koch, in Linnaea, xxii. (1849) 211 = scutatus.
pulcher, Ehrenb. ex Meissn. in DC. Prod. xiv. 55 = brevipes.
pulcher, *Linn. Sp. Pl.* 336.—Europ.; Oriens.
pulcherrimus, Hort. ex Schult. f. Syst. vii. 1394 = ucranicus.
purpureus, *Link, Enum. Hort. Berol.* i. 350.—Hab.?
purpureus, Poir. Encyc. v. 63 = obtusifolius.
pusillus, Delarb. Fl. Auv. ed. II. 523 = conglomeratus.
pycnocarpus, Wallr. ex Steud. Nom. ed. II. ii. 486 = conglomeratus.
pyrenaicus, Pourr. ex Lapeyr. Suppl. Hist. Pl. Pyr. 49 = R. Acetosella.
quadridentatus, Ehrenb. ex Boiss. Fl. Orient. iv. 1013 = dentatus
ramulosus, Meissn. in DC. Prod. xiv. 55 = nepalensis.
Raulini, *Boiss. Diagn.* Ser. I. xii. 100.—Ins. Creta.
Regeli, F. Schmidt, Reisen Amurl. 167 = crispus.

RUMEX:—
reniformis, Radde, Reise in Sued-Ost Sibir. 131 = Oxyria digyna.
repandus, Bosc, ex Pers. Syn. i. 395 = fimbriatus.
reticulatus, Bess. Ind. Sem. Hort. Cremen. (1820); et ex Spreng. Neue Entdeck. iii. 18 (excl. syn.) = pulcher.
retroflexus, Lag. Elench. Pl. 13; ex Schult. f. Syst. vii. 1449 = alpinus.
rheifolius, Hort. ex Schult. f. l. c. 1436 = cordifolius.
Romassa, *Remy, in C. Gay, Fl. Chil.* v. 280.—Chili.
roseus, Aucher, ex Meissn. in DC. Prod. xiv. 71 = vesicarius.
roseus, *Linn. Sp. Pl.* 337.—Reg. Mediterr.; Oriens.
roseus, Poir. Encyc. v. 65 = pictus.
Roxburghianus, Schult. f. Syst. vii. 1402 = nepalensis.
Roxburghianus, Wall. Cat. n. 1731 = maritimus, dentatus.
rubellus, Steud. Nom. ed. I. 710; ed. II. ii. 486 = ucranicus.
Rugelii, Meissn. in DC. Prod. xiv. 54 = obtusifolius.
rugosus, Campd. Monog. Rum. 113 = R. Acetosa.
rupestris, *Le Gall, in Congr. Sc. Fr.* xvi. I. (1850) 143; *Fl. Morb.* 501.—Europ.
sagittatus, Thunb. Prod. Pl. Cap. 67 = luxurians.
sagittifolius, Campd. Monog. Rum. 150 = R. Acetosa.
sagittifolius, Plenk, ex Campd. l. c. 108 = abyssinicus.
salicifolius, *Weinm. in Flora*, iv. (1821) 28.—Am. bor.
sanguineus, *Linn. Sp. Pl.* 334.—Europ.
sanguineus, Poir. Encyc. v. 59 = obtusifolius.
sarcorhizus, Link, Enum. Hort. Berol. i. 351 = cordatus.
Saxei, Kellogg, in Pacif. Rural Press (June 1879); ex Curran, in Bull. Calif. Acad. i. n. III. (1885) 146 = hymenosepalus.
scandens, Burch. Trav. ii. 146 = luxurians.
Schimperi, *Meissn. in DC. Prod.* xiv. 67.—Abyss.
Schultzii, Meissn. l. c. 74 = Shultzii.
scutatus, Herb. Banks, ex Campd. Monog. Rum. 136 = cordatus.
scutatus, *Linn. Sp. Pl.* 337.—Europ.; Oriens.
sepium, Bieb. ex Schult. f. Syst. vii. 1728 = obtusifolius.
setaceus, Buch.-Ham. ex Meissn. in Wall. Pl. As. Rar. iii. 64 = maritimus.
Shultzii, *Rafin. in Med. Repos. N. York*, v. (1808) 353, nomen.—Am. bor.
silvester, Kern. in Oestr. Bot. Zeitschr. xxv. (1875) 353 = sylvestris.
sinuatus, Nathhort. ex Steud. Nom. ed. II. ii. 486 = tingitanus.
songaricus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ix. 85.—Sibir. altaic.
spathulatus, *Thunb. Prod. Pl. Cap.* 67.—Afr. austr.
spinosus, Linn. Sp. Pl. 337 = Emex spinosa.
spinosus, Thomas, ex Meissn. in DC. Prod. xiv. 58 = pulcher.
spinosus, Thunb. Prod. Pl. Cat. 67 = Emex australis.
spurius, *Rafin. in Med. Repos. N. York*, v. (1808) 353, nomen.—Am. bor.
Steini, Becker, Fl. Frankf. i. 165 = palustris.
stenophylloides ×, Simk. in Termész. Füzetek, i. (1877) 238.—Europ.
stenophyllus, *Ledeb. Fl. Alt.* ii. 58.—Reg. Cauc.; As. bor.
stenophyllus, Timb. & Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxii = R. Acetosa.
Steudelianus, Meissn. in DC. Prod. xiv. 56 = obtusifolius.
Steudelii, Hochst. ex A. Br. in Flora, xxiv. (1841) 278 = obtusifolius.
strepens, Hort. ex Schult. f. Syst. vii. 1424 = longifolius.
strictus, Link, Enum. Hort. Berol. i. 350 = dentatus.
strictus, Meissn. in DC. Prod. xiv. 57 = cassius.
subrotundus, *Campd. Monog. Rum.* 151.—Oriens.
subtrilobus, *Boiss. Diagn.* Sér. I. v. 46.—Lydia.
suffocatus, Moris, ex Bertol. Fl. Ital. iv. 242 = pulcher.
suffruticosus, *J. Gay, ex Meissn. in DC. Prod.* xiv. 72.—Hispan.
supinus, Campd. Monog. Rum. 147 = R. Acetosella.
sylvaticus, Herb. ex Steud. Nom. ed. II. ii. 486 = nemorosus.
sylvaticus, *Rafin. Ann. Nat.* (1820) 12.—Am. bor.
sylvaticus, Schrad. ex Link, Enum. Hort. Berol. i. 350 = acutus.
sylvestris, Campd. Monog. Rum. 146 = obtusifolius.
sylvestris, *Wallr. Sched. Crit.* 161.—Europ.
syriacus, Meissn. in DC. Prod. xiv. 53 = obtusifolius.
tauricus, Hort. ex Schult. f. Syst. vii. 1411 = conglomeratus.

RUMEX:—
thyrsiflorus, Fingerh. in Linnaea, iv. (1829) 380 = R. Acetosa.
thyrsoides, *Desf. Fl. Atlant.* i. 321.—Reg. Mediterr.
thyrsoides, Gren. & Godr. Fl. Fr. iii. 44 = intermedius.
tingitanus, *Linn. Syst.* ed. X. 991.—Reg. Mediterr.
tingitanus, Webb & Bourg. ex Meissn. in DC. Prod. xiv. 67 = maderensis.
tmoleus, *Boiss. Diagn.* Ser. I. xii. 101.—As. Min.
tolimensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 262.—N. Granat.
Tournefortii, Campd. Monog. Rum. 149 = intermedius.
triangularis, DC. Fl. Fr. Suppl. 368, in obs. = R. Acetosa.
triangularis, Guss. Fl. Sic. Syn. i. 434 = intermedius.
trifidus, *Campd. Monog. Rum.* 129.—Hab.?
tuberculatus, C. Koch, in Linnaea, xxii. (1849) 208 = pulcher.
tuberosus, Eckl. ex Steud. Nom. ed. II. ii. 486 = luxurians.
tuberosus, *Linn. Sp. Pl.* ed. II. 481.—Graec.; Afr. bor.; Oriens.
tuberosus, Poir. Encyc. v. 67 = R. Acetosa.
tuberosus, Roxb. ex Steud. Nom. ed. II. ii. 486 = nepalensis.
tuberosus, Sieber, ex Meissn. in DC. Prod. xiv. 67 = lativalvis.
tuberosus, Thunb. Prod. Pl. Cap. 67 = cordatus.
turcicus, Boiss. Diagn. Ser. II. iv. 79 = crispus
ucranicus, *Fisch. Cat. Hort. Gorenk.* (1812) 16; *et Bess. ex Spreng. Nov. Prov.* 36.—Rossia.
uliginosus, Guss. Pl. Rar. Neap. (1826) 151. t. 29 = palustris.
umbellatus, Houtt. Handleid. viii. 414 = Polygonum chinense.
uncatus, Hort. ex Schult. f. Syst. vii. 1401 = nepalensis.
uncinatus, Ehrenb. ex Meissn. in DC. Prod. xiv. 58 = pulcher.
uncinatus, Hort. ex Schult. f. Syst. vii. 1401 = nepalensis.
uncinulatus, Schrad. Ind. Sem. Hort. Gotting. (1823) 5 = nepalensis.
undulatus, Bess. ex Meissn. in DC. Prod. xiv. 43 = aquaticus.
undulatus, Desf. Cat. Hort. Par. ed. III. 389 = cristatus, alpinus.
undulatus, Schrank, Baier. Fl. i. 625 = conglomeratus.
undulatus, Willd. ex Spreng. Syst. ii. 160 = dentatus.
venosus, *Pursh, Fl. Am. Sept.* ii. 733.—Am. bor.
verticillatus, *Linn. Sp. Pl.* 334.—Am. bor.
verticillatus, Richards. in Frankl. 1st Journ. App. ed. II. 11 = salicifolius.
verticillatus, Ucria, Hort. Reg. Panorm. 164 = sanguineus.
vesicarius, *Linn. Sp. Pl.* 336.—Graecia; Afr. bor.; Oriens.
vinealis, Timb. & Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxii = R. Acetosa.
virgatus, Haenke, in Jirasek, Beob. Riesengeb. (1791) 39 = conglomeratus.
viridis, Steud. Nom. ed. I. 710 = sanguineus.
Wallichianus, Meissn. in Wall. Pl. As. Rar. iii. 64 = maritimus.
Wallichii, Meissn. in DC. Prod. xiv. 48 = praec.
Wallrothii, Nym. Syll. 327 = obtusifolius.
Warrenii, Trimen, ex Nym. Consp. 635 = *Knafii*.
Winterlii, Zuccagni, Syn. Pl. Hort. Florent. 45 = conglomeratus.
Woodsii, *De Not. Cat. Sem. Roma* (1875), 28; *et in Nuov. Giorn. Bot. Ital.* viii. (1876) 42.—Italia.
xanthorhizus, Pers. Syn. i. 394 = R. Britannica.

RUMFORDIA, DC. Prod. v. 549 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 359.
floribunda, *DC. l. c.* 550.—Mexic.

RUMIA, Hoffm. Gen. Umbell. ed. II. 171 (1816) = **Trinia**, Hoffm. (Umbell.).
athamantoides, DC. Prod. iv. 98 (= *Stenocoelium athamantoides*).—Sibir. altaic.
capensis, Link, Enum. Hort. Berol. i. 271 = Capnophyllum africanum.
depressa, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 126 = Trachydium depressum.
frigida, Boiss. & Heldr. Diagn. Ser. II. vi. 77.—Graecia.
Guicciardii, Boiss. & Heldr. l. c.—Graecia.
Kotschyi, Boiss. l. c. Ser. I. vi. 59 = Trachydium Kotschyi.

RUMIA :—
leiogona, C. A. Mey. Verz. Pfl. Cauc. 125 = Trinia Hoffmanni.
microcarpa, Hoffm. Gen. Umbell. ed. II. 175 = Hippomarathrum crispum.
multiflora, Ledeb. Fl. Ross. ii. 281.—Rossia.
seseloides, Hoffm. Gen. Umbell. ed. II. 174.— Sibir. altaic.
taurica, Hoffm. l. c. 173 = Sanicula crithmifolia.

RUMINIA, Parl. 'Due Nov. Gen. Monocot. 3'; ex ej. Fl. Ital. iii. 84 (1858) = **Leucojum**, Linn. (Amaryll.).
hyemalis, Parl. ll. cc. 4; 85 = L. hyemale.
nicaeënsis, Jord. & Four. Ic. Fl. Europ. 26. t. 65. f. 108 = L. hyemale.

RUMPFIA, Linn. Syst. ed. I (1735) = seq.

RUMPHIA, Linn. Sp. Pl. ed. I. 1193 (1753). *ANACARDIACEAE*, Benth. & Hook. f. i. 428.
Tsiemtani, Adans. Fam. ii. 303 (1763).
amboinensis, *Linn. Sp. Pl.* 1193.—Ind. or.
tiliaefolia, Poir. Encyc. vi. 352 = amboinensis.

RUMPUTRIS, Rafin. Fl. Tellur. iv. 92 (1836) = **Cassytha**, Linn. (Laurin.).
fasciculata, Rafin. l. c. = C. filiformis?

RUNCINA, Allem. in Nov. Act. Nat. Cur. iv. (1770) 94 (Quid?).

RUNGIA, Nees, in Wall. Pl. As. Rar. iii. 77 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1120.
apiculata, *Bedd. Ic. Pl. Ind. Or.* i. 60.—Ind. or.
Arnottiana, Wight, Ic. t. 1550 = Wightiana.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 549.—Burma.
Brandisii, *C. B. Clarke, l. c.*—Burma.
chinensis, *Benth. Fl. Hongk.* 266.—China.
ciliata, *Nees, in Wall. Pl. As. Rar.* iii. 111.—Am. austr.
crenata, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 518.—Ind. or.
diversiformis, *Nees, in DC. Prod.* xi. 472.—Ins. Philipp.
elegans, *Dals. & Gibs. Bomb. Fl.* 196.—Ind. or.
grandis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 46 —Afr. trop.
himalayensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 548.—Reg. Himal.
interrupta, Nees, in DC. Prod. xi. 732, in syn. = Dicliptera interrupta.
khasiana, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 518.—Reg. Himal.
laeta, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 546.—Ind. or.
latior, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 313. —Ind. or.
linifolia, *Nees, in Wall. Pl. As. Rar.* iii. 110.—Ind. or.
longifolia, T. Anders. in Thw. Enum. Pl. Zeyl. 235, partim = latior.
longifolia, Bedd. Ic. Pl. Ind. Or. t. 266 = parviflora.
longifolia, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 313.—Zeylan.
Mastersi, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 519.—Assam.
muralis, Royle, ex Nees, in DC. Prod. xi. 470 = parviflora.
origanoides, Nees, in Wall. Pl. As. Rar. iii. 110 = parviflora.
parviflora, *Nees, l. c.*—Ind. or.; Burma.
pectinata, Nees, in DC. Prod. xi. 470 = parviflora.
polygonoides, Nees, in Wall. Pl. As. Rar. iii. 110 = parviflora.
pubinervia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 46.—Afr. austr.
punduana, Nees, in Wall. Pl. As. Rar. iii. 110 = Justicia punduana.
repens, T. Anders. in Journ. Linn. Soc. ix. (1867) 518, partim = parviflora.
repens, *Nees, in Wall. Pl. As. Rar.* iii. 110.—Ind. or.
sisparensis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 518.—Ind. or.
stolonifera, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 547.—Reg. Himal.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 567.—Sumatra.
Wightiana, *Nees, in Wall. Pl. As. Rar.* iii. 110.—Ind. or.

RUPALA, Vahl, Symb. Bot. iii. 20 (1794) = **Roupala**, Aubl. (Proteac.).

RUPALLEYA, Morière, in Bull. Soc. Linn. Normand. viii. (1864) 317. t. 15 = **Stropholirion**, Torr. (Liliac.).
volubilis, Morière, l. c. = S. californicum.

RUPICAPNOS, Pomel, Mat. Fl. Atl. 16 (1860) = **Fumaria**, Tourn. (Papav.).
africana, Pomel, Nouv. Mat. Fl. Atl. 240 = F. africana.
Caput-plataleae, Pomel, l. c. 244 = F. numidica.
cerefolia, Pomel, l. c. 242 = F. africana.
corymbosa, Pomel, Mat. Fl. Atl. 16 = F. africana.
delicatula, Pomel, Nouv. Mat. Fl. Atl. 246 = F. numidica.
erosa, Pomel, l. c. 243 = F. numidica.
graciliflora, Pomel, l. c. 241 = F. africana.
longipes, Pomel, Mat. Fl. Atl. 16 = F. numidica.
muricaria, Pomel, Nouv. Mat. Fl. Atl. 245 = F. numidica.
numidica, Pomel, Mat. Fl. Atl. 16 = F. numidica.
ochracea, Pomel, Nouv. Mat. Fl. Atl. 242 = F. africana.
platycentra, Pomel, l. c. 242 = F. africana.
Reboudiana, Pomel, l. c. 379 = F. numidica.
sarcocapnoides, Pomel, Mat. Fl. Atl. 16 = F. numidica.
speciosa, Pomel, Nouv. Mat. Fl. Atl. 241 = F. africana.
tenuifolia, Pomel, l. c. 244 = F. numidica.

RUPIFRAGA, 'Linn.' ex Rafin. Fl. Tellur. ii. 67 (1836) = **Saxifraga**, Linn.
cuscutaeformis, Rafin. l. c. iv. 121 = S. sarmentosa.
sarmentosa, 'Linn.' ex Rafin. l. c. ii. 68 = S. sarmentosa.

RUPPELIA, Baker, in Oliver, Fl. Trop. Afr. ii. 149 (1871) = RUEPPELIA, A. Rich. = **Aeschynomene**, Linn. (Legumin.).

RUPPIA, Linn. Syst. ed. I (1735). *NAIADACEAE*, Benth. & Hook. f. iii. 1014.
Buccaferrea, Mich. ex Petagna, Inst. Bot. v. 1826 (1787).
antarctica, Labill. Nov. Holl. Pl. ii. 116 = Cymodocea antarctica.
aragonensis, Losc. Pard. Ser. Inconf. Pl. Arag. 105 = rostellata?
brachypus, J. Gay, in Coss. Notes Crit. 10 = maritima.
brevipes, Bertol. ex Griseb. Fl. Brit. W. Ind. 507 = maritima.
didyma, Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1825) 427 = maritima.
drepanensis, Tineo, ex Guss. Fl. Sic. Syn. ii. 878 = maritima.
marina, Fries, Summa Veg. Scand. 68 (err. typ. 69) = maritima.
maritima, Griff. Notul. iii. 196 = rostellata.
maritima, *Linn. Sp. Pl.* 127.—Reg. temp. et trop.
obliqua, Schur, ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 1. 355 = rostellata.
rostellata, *Koch, ex Reichb. Ic. Pl. Crit.* ii. 66.—Europ.; As. trop.
salina, Schur, Enum. Pl. Transs. 634 = rostellata.
spiralis, Dum. Fl. Belg. 164 = maritima.
subsessilis, Thw. Enum. Pl. Zeyl. 333 = Potamogeton pectinatus.
transsilvanica, Schur, in Oestr. Bot. Zeitschr. x. (1860) 356 = rostellata.
trichodes, Dur. in Duch. Rev. Bot. ii. (1846–47) 426 = maritima.

RUPRECHTIA, C. A. Mey. in Mém. Acad. Pétersb. Sér. VI. vi. (1840) 148. t. 4. *POLYGONACEAE*, Benth. & Hook. f. iii. 104.
Magonia, Vell. Fl. Flum. 165 (1825); iv. t. 60 (1827).
amentacea, *Meissn. in Mart. Fl. Bras.* v. 1. 56.—Bras.
apetala, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 268.—Bolivia.
brachysepala, *Meissn. in Mart. Fl. Bras.* v. 1. 58.—Bras.
brachystachya, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 630.—Guian.
carpinoides, *Meissn. in Mart. Fl. Bras.* v. 1. 58.—Bras.
corylifolia, *Griseb. in Goett. Abh.* xix. (1874) 113.—Reg. Argent.

RUPRECHTIA:—
costata, Meissn. in DC. Prod. xiv. 180 = Cumingii.
Cruegerii, Griseb. Fl. Brit. W. Ind. 710 = fagifolia.
Cumingii, *Meissn. in DC. Prod.* xiv. 179.—Panama.
excelsa, *Griseb. in Goett. Abh.* xix. (1874) 113.—Reg. Argent.
fagifolia, *Meissn. in Mart. Fl. Bras.* v. 1. 58.—Bras.
glauca, *Meissn. l. c.* 59.—Bras.
Jamesonii, *Meissn. in DC. Prod.* xiv. 179.—Peruv.
laurifolia, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. vi. (1840) 150.—Bras.
laxiflora, *Meissn. in Mart. Fl. Bras.* v. 1. 56.—Bras.
Lundii, *Meissn. l. c.* 54.—Bras.
Martii, *Meissn. l. c.* 58.—Bras.
mollis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 268.—Bolivia.
polystachya, *Griseb. in Goett. Abh.* xxiv. (1879) 89.—Reg. Argent.
ramiflora, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. vi. (1840) 150.—N. Granat.
salicifolia, *C. A. Mey. l. c.*—Bras.
tenuiflora, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 629.—Bras.
triflora, *Griseb. in Goett. Abh.* xxiv. (1879) 89.—Reg. Argent.
Viraru, *Griseb. l. c.* 90.—Reg. Argent.

RUPRECHTIA, Opiz, Seznam, 86 (1852) = **Thalictrum**, Tourn. (Ranunc.).
aquilegifolia, Opiz, l. c. = T. aquilegifolium.

RUSCUS, [Tourn.] Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 764.
aculeatus, *Linn. Sp. Pl.* 1041.—Europ.; Oriens.
androgynus, Linn. l. c. = Semele androgyna.
anthopus, Rafin. Fl. Tellur. iv. 26 = Semele androgyna.
dumosus, *Clarke, Trav.* i. ed. II. Add. p. xxii.—Rumelia.
flexuosus, Mill. Gard. Dict. ed. VIII. n. 6 = aculeatus.
frutescens, *Mill. l. c.* n. 8.—Mexic.
humilis, Salisb. Prod. 255 = R. Hypoglossum.
Hypoglossum, *Linn. Sp. Pl.* 1041.—Europ. austr.
Hypophyllum, *Linn. l. c.*—Reg. Mediterr. et Caucas.
latifolius, Rafin. Fl. Tellur. iv. 26 = Semele androgyna.
laurifolius, Hort. ex Steud. Nom. ed. II. ii. 487 = Danaë Laurus.
laxus, Sm. in Trans. Linn. Soc. iii. (1797) 334 = aculeatus.
lugubris, Salisb. Pord. 255 = R. Hypophyllum.
microglossus, Bertol. Fl. Ital. x. 401 = R. Hypoglossum.
parasiticus, Gueldenst. Reisen, i. 426 = aculeatus?
procerus, Salisb. Prod. 255 = Semele androgyna.
racemosus, Linn. Sp. Pl. 1041 = Danaë Laurus.
racemus, Crantz, Inst. i. 200, sphalm. = praec.
reticulatus, Thunb. Prod. Pl. Cap. 13 = Behnia reticulata.
terminalis, Salisb. Prod. 255 = Danaë Laurus.
trifoliatus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Ins. Graecia.
troadensis, *Clarke, Trav.* ii. 146.—Graecia.
volubilis, Thunb. Prod. Pl. Cap. 13 = Asparagus medeoloides.

RUSSEA, J. F. Gmel. Syst. 237 (1791) = **Roussea**, Sm. (Saxifrag.).

RUSSEGGERA, Endl. Nov. Stirp. Dec. 38 (1839) = **Lepidagathis**, Willd. (Acanth.).
collina, Endl. l. c. = L. radicalis.

RUSSELIA, Jacq. Enum. Pl. Carib. 6 (1760). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 940.
alata, *Cham. & Schlecht. in Linnaea*, iii. (1828) 3.—Bras.
alternifolia, *Pers. Syn.* ii. 164.—Peruv.
americana, Raeusch. Nom. ed. III. 180 = sarmentosa.
depressa, Walp. Rep. iii. 234 = Aptosimum depressum.
ephedroides, Benth. Pl. Hartw. 69 = verticillata.
equisetiformis, *Schlecht. & Cham. in Linnaea*, vi. (1831) 377.—Mexic.
flammea, Rafin. New Fl. Am. ii. 71 = sarmentosa.
floribunda, H. B. & K. Nov. Gen. et Sp. ii. 359 = rotundifolia.

RUSSELIA:—
juncea, *Zucc. in Flora*, xv. (1832) II. *Beibl.* 99.—Mexic.
multiflora, Sims, Bot. Mag. t. 1528 = sarmentosa.
paniculata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 19 = sarmentosa.
polyedra, Zucc. in Abh. Akad. Muench. ii. (1831–36) 328 = sarmentosa.
rotundifolia, *Cav. Ic.* v. 9. t. 415.—Mexic.
sarmentosa, *Jacq. Enum. Pl. Carib.* 25; *Select. Am.* 178.—Am. trop.
syringaefolia, Schlecht. & Cham. in Linnaea, vi. (1831) 376 = sarmentosa.
ternifolia, H. B. & K. Nov. Gen. et Sp. ii. 359 = sarmentosa.
verticillata, *H. B. & K. l. c.* 360.—Mexic.

RUSSELIA, Koen. ex Roxb. Fl. Ind. iii. 364 (1832) = **Ormocarpum**, Beauv. (Legumin.).

RUSSELIA, Linn. f. Suppl. 24 (1781) = **Vahlia**, Thunb. (Saxifrag.).
capensis, Linn. f. l. c. 175 = V. capensis.
oldenlandioides, Roxb. ex Wight & Arn. Prod. 449 = V. oldenlandioides.
viscosa, Roxb. l. c. = V. viscosa.

RUSSERIA, Buek, Index ad DC. Prod. iii. 389 (1858), sphalm. = **Bursera**, Jacq.

RUSTIA, Klotzsch, in Hayne, Arzn. Gew. xiv. sub t. 14; et sub t. 15 (1846). *RUBIACEAE*, Benth. & Hook. f. ii. 45.
HENLEA, Karst. Fl. Columb. 38 in textu (1858); 157. t. 78 (1861).
TRESANTHERA, Karst. l. c. 37. t. 19 (1858).
formosa, *Klotzsch, in Hayne, Arzn. Gew.* xiv. sub t. 15.—Bras.
occidentalis, *Hemsl. Biol. Centr. Am. Bot.* ii. 14.—Mexic.
Pohliana, Klotzsch, in Hayne, Arzn. Gew. xiv. sub t. 15 = formosa.
Sellowiana, Klotzsch, l. c. = formosa.

RUTA, [Tourn.] Linn. Syst. ed. I (1735). *RUTACEAE*, Benth. & Hook. f. i. 286.
APLOPHYLLUM, A. Juss. in Mém. Mus. Par. xii. (1825) 464.
DESMOPHYLLUM, Webb & Berth. Phyt. Canar. i. 14 (1836–50).
HAPLOPHYLLUM, Reichb. Fl. Germ. Excurs. 766 (1832).
RUTARIA, Webb, ex Benth. & Hook. f. Gen. i. 287 (1862).
acutifolia, *DC. Prod.* i. 711.—Persia.
albiflora, Hook. Exot. Fl. t. 79 = Boenninghausenia albiflora.
altera, Mill. Gard. Dict. ed. VIII. n. 2 = graveolens.
angustifolia, Pers. Syn. i. 464 = chalepensis.
Besseri, *Nym. Syll.* 219.—Rossia.
Biebersteinii, Neilr. Nachtr. 280 = suaveolens.
bracteosa, DC. Prod. i. 710 = chalepensis.
Buxbaumii, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 129 = patavina.
Buxbaumii, *Poir. Encyc.* vi. 336.—Oriens.
chalepensis, *Linn. Mant.* i. 69.—Reg. Mediterr.
chalepensis, Wall. Cat. n. 7113 = graveolens.
ciliata, Savi, ex Steud. Nom. ed. II. ii. 487 = glabra.
ciliata, Mill. Gard. Dict. ed. VIII. n. 5 = graveolens.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 (Quid?).—Java.
cordata, *D. Don, Prod. Fl. Nep.* 206.—Nepal.
coronata, *Nym. Syll.* 219.—Europ.
corsica, *DC. Prod.* i. 710.—Ins. Corsic.
crithmifolia, Moric. ex DC. l. c. = graveolens.
dahurica, *DC. Prod.* i. 712.—Dahuria.
Dampatis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 207 = Boenninghausenia albiflora.
dichotoma, *DC. Prod.* i. 711.—Persia.
divaricata, Salzm. in Flora, iv. (1821) 109 = corsica.
divaricata, Tenore, Prod. Fl. Nap. p. xxiv = graveolens.
diversifolia, Wender. ex Steud. Nom. ed. II. ii. 487, nomen (Quid?).—Hab.?
fernandezia, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 52.—Ins. J. Fernand.
frutescens, Siev. ex Fisch. Enum. Pl. Schrenk. i. 90 = acutifolia.

RUTA:—
fruticulosa, *Labill. Ic. Pl. Syr. Dec.* i. 13.—Syria.
fumariaefolia, Boiss. & Heldr. Diagn. Ser. I. viii. 125 = chalepensis.
glabra, *DC. Prod.* i. 711.—Assyria.
graveolens, *Linn. Sp. Pl.* 383.—Europ. austr.
holopetala, Kitt. Taschenb. ed. II. 794 = graveolens.
hortensis, Mill. Gard. Dict. ed. VIII. n. 1 = graveolens.
intermedia, Hort. ex Steud. Nom. ed. II. ii. 488 = graveolens.
japonica, Siebold, ex Hook. f. Fl. Brit. Ind. i. 486 = Boenninghausenia albiflora.
latifolia, Salisb. Prod. 320 = chalepensis.
legitima, All. Fl. Pedem. i. 280 = montana.
linearis, *DC. Prod.* i. 712.—Sibir.
linifolia, Bieb. Fl. Taur. Cauc. i. 309 = suaveolens?
linifolia, *Linn. Sp. Pl.* 384.—Hispan.; Afr. bor.
linifolia, Sibth. & Sm. Fl. Graec. Prod. i. 273 = Buxbaumii.
macrophylla, Moris, Stirp. Sard. Elench. 11 = chalepensis.
macrophylla, *Soland. in Bot. Mag.* t. 2018.—Afr. bor.
montana, *Mill. Gard. Dict.* ed. VIII. n. 7.—Reg. Mediterr. et Caucas.
Montbretii, Viv. Pl. Aegypt. Dec. iv. 13 = tuberculata.
obtusifolia, *Ledeb. ex Eichw. Pl. Casp. Cauc.* 37.—Reg. Casp.
officinalis, Pall. Reise Suedl. Statth. ii. 80 = graveolens.
Oreojasme, *Webb & Berth. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 130.—Ins. Canar.
parviflora, Desf. in Ann. Mus. Par. xi. (1808) 440 = villosa.
patavina, *Linn. Sp. Pl.* 384.—Europ. austr.
patavina, Poir. Encyc. vi. 335 = pubescens.
patavina, Portenschl. Enum. t. 7, ex Reichb. Fl. Germ. Excurs. 766 = linifolia.
perforata, *Bieb. Beschr. Casp.* 172.—Sibir. baical.
pinnata, *Linn. f. Suppl.* 232.—Ins. Canar.
pubescens, *Willd. Enum. Hort. Berol.* 446.—Hispan.
punctata, Presl, ex Mert. & Koch, Deutschl. Fl. iii. 88 = chalepensis.
rosmarinifolia, *Pers. Syn.* i. 465.—Hispan.
spatulata, Sibth. & Sm. Fl. Graec. iv. 63. t. 370 = Buxbaumii.
suaveolens, Bess. ex Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 179 = Besseri.
suaveolens, *DC. Prod.* i. 711.—Rossia; Thracia.
subtripinnata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 21.—Japon.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 3 = montana.
taurica, *Nym. Syll.* 220.—Taur.
tenuifolia, Vill. Prosp. 47; Gouan, Herbor. 26; Pourr. in Mém. Acad. Toul. iii. (1788) 326 = montana.
thesioides, *Fisch. ex DC. Prod.* i. 712.—Reg. Casp.
tuberculata, *Forsk. Fl. Aegypt. Arab.* 86.—Aegypt.; Arab.; Persia.
tuberculata, Viv. Pl. Aegypt. Dec. iv. 13 = glabra.
villosa, *Bieb. Fl. Taur. Cauc.* i. 310.—Reg. Caucas.
villosa, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = fruticulosa, Buxbaumii.

RUTAEA, M. Roem. Syn. Hesper. 93 (1846) = **Turraea**, Linn. (Meliac.).
floribunda, M. Roem. l. c. = T. floribunda.
heterophylla, M. Roem. l. c. 94 = T. heterophylla.
lobata, M. Roem. l. c. = T. heterophylla.
maculata, M. Roem. l. c. = T. maculata.
obtusifolia, M. Roem. l. c. 93 = T. obtusifolia.
quercifolia, M. Roem. l. c. 94 = T. heterophylla.
rigida, M. Roem. l. c. = Quivisia chilosantha.
sericea, M. Roem. l. c. 93 = T. sericea.

RUTARIA, Webb, ex Benth. & Hook. f. Gen. i. 287 (1862) = **Ruta**, Tourn.

RUTERIA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 380 = **Psoralea**, Linn. (Legumin.).
africana, Medic. l. c. = P. pinnata.
decidua, Moench, Meth. Suppl. 38 = P. aphylla.
pinnata, Medic. Phil. Bot. ii. 41 = P. pinnata.
tenuifolia, Moench, Meth. Suppl. 38 = P. tenuifolia.

RUTHEA, Bolle, in Verh. Bot. Ver. Brandenb. iii–iv. (1861–62) 174 = **Lichtensteinia**, Cham. & Schlecht. (Umbell.).
herbanica, Bolle, l. c.—Ins. Canar.

RUTICA, Neck. Elem. ii. 202 (1790), sphalm. = **Urtica**, Tourn.

RUTIDEA, DC. in Ann. Mus. Par. ix. (1807) 219. *RUBIACEAE*, Benth. & Hook. f. ii. 116.
RYTIDEA, Spreng. Syst. i. 515 (1825).
coriacea, *Baker, Fl. Maurit.* 149.—Ins. Mascar.
decorticata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 190.—Afr. trop.
ferruginea, *Hiern, l. c.* 189.—Afr. trop.
fuscescens, *Hiern, l. c.* 191.—Afr. trop.
glabra, *Hiern, l. c.* 190.—Afr. trop.
hispida, *Hiern, l. c.* 189.—Afr. trop.
membranacea, *Hiern, l. c.* 190.—Afr. trop.
mollis, *Blume, ex DC. Prod.* iv. 495.—Ins. Pinang.
olenotricha, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 189.—Afr. trop.
parviflora, Benth. in Hook. Niger Fl. 416, partim = olenotricha.
parviflora, *DC. in Ann. Mus. Par.* ix. (1807) 219.—Afr. trop.
rufipilis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 188.—Afr. trop.
Smithii, *Hiern, l. c.* 189.—Afr. trop.

RUTIDOCHLAMYS, Sond. in Linnaea, xxv. (1852) 497 = **Podolepis**, Labill. (Compos.).
Mitchellii, Sond. l. c. = P. Rhytidochlamys.

RUTIDOSIS, DC. Prod. vi. 158 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 316.
ACTINOPAPPUS, Hook. f. ex A. Gray, in Hook. Kew Journ. iv. (1852) 226.
PUMILO, Schlecht. in Linnaea, xxi. (1848) 448.
acoma, F. Muell. Fragm. ii. 89 = Acomis rutidosea.
arachnoidea, Hook. in Mitch. Journ. Trop. Austral. 341 = Podolepis Rhytidochlamys.
auricoma, F. Muell. in Linnaea, xxv. (1852) 408 = helichrysoides.
Brownii, *Benth. Fl. Austral.* iii. 594.—Austral.
helichrysoides, *DC. Prod.* vi. 159 (helychrysoides).—Austral.
leiolepis, *F. Muell. in Trans. Vict. Inst.* i. (1855) 131.—Austral.
leptorrhychoides, *F. Muell. Fragm.* v. 148.—Austral.
leucantha, *F. Muell. l. c.* i. 35.—Austral.
macra, F. Muell. l. c. iv. 145 = Acomis macra.
Murchisonii, *F. Muell. l. c.* i. 34.—Austral.
Pumilo, *Benth. Fl. Austral.* iii. 595.—Austral.

RUTILIA, Vell. Fl. Flum. 156 (1825); iv. t. 19 (1827) (Quid?).
ternata, Vell. ll. cc. 19.—Bras.

RUTOSMA, A. Gray, Gen. Illustr. ii. 143. t. 155 (1849) = **Thamnosma**, Torr. & Frém. (Rutac.).
texanum, A. Gray, l. c. 144. t. 155 = T. texanum.

RUTTYA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 27. *ACANTHACEAE*, Benth. & Hook. f. ii. 1105.
HABLANTHERA, Hochst. in Bot. Zeit. ii. (1844) 853 (sphalm.).
HAPLANTHERA, Hochst. in Flora, xxvi. (1843) 71.
ovata, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 27.—Afr. austr.

RUYSCHIA, Jacq. Enum. Pl. Carib. 2 (1760). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 181.
LOGHANIA, Scop. Introd. 236 (1777).
SOUROUBEA, Aubl. Pl. Gui. 244. t. 97 (1775).
amazonica, *Mart. Nov. Gen. et Sp.* iii. 176. t. 292.—Bras.
bahiensis, *Mart. l. c.* 178.—Bras.
bicolor, *Benth. Bot. Voy. Sulph.* 73. t. 29.—Panama.
clusiaefolia, *Jacq. Enum. Pl. Carib.* 17; *Select. Am.* 75.—Am. trop.
corallina, *Mart. Nov. Gen. et Sp.* iii. 177. t. 294.—Bras.
crassipes, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 378.—N. Granat.
cymbadenia, Delp. in Atti Soc. Ital. Sc. Nat. Milano, xii. (1869) 196 = clusiaefolia.
fragrans, Mor. ex Wittm. in Mart. Fl. Bras. xii. 1. 248 = clusiaefolia.
guianensis, Vitm. Summa Pl. ii. 250 = R. Souroubea.
laurifolia, Presl, Bot. Bemerk. 24 = clusiaefolia.
lepidota, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 83.—Guian.
mexicana, *Baill. Adansonia*, x. (1872) 241.—Mexic.

RUYSCHIA:—
Pavonii, *G. Don, Gen. Syst.* i. 625.—Peruv.
pilophora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 379.—N. Granat.
Souroubea, *Sw. Prod. Veg. Ind. Occ.* 50.—Guian.
sphaeradenia, *Delp. in Atti Soc. Ital. Sc. Nat. Milano*, xii. (1869) 180.—Peruv.
Spixiana, *Mart. Nov. Gen. et Sp.* iii. 177. t. 293.—Bras.
subsessilis, *Benth. Bot. Voy. Sulph.* 73. t. 29.—Panama.
Surubea, J. F. Gmel. Syst. 407 = R. Souroubea.

RUYSCHIA, Linn. Syst. ed. I (1735) = **Dracocephalum**, Tourn. (Labiat.).

RUYSCHIANA, Boerh. ex Mill. Gard. Dict. ed. VII. (1759) = **Dracocephalum**, Tourn. (Labiat.).
fasciculata, [Clairv.] Man. Herb. 190 = D. Ruyschiana.
laciniata, Mill. Gard. Dict. ed. VIII. n. 2 = D. austriacum.
spicata, Mill. l. c. n. 1 = D. Ruyschiana.
verticillata, Mill. l. c. n. 3. = D. peregrinum.

RYANAEA, DC. Prod. i. 255 (1824) = seq.

RYANIA, Vahl, Eclog. Am. i. 51. t. 9 (1796). *BIXINEAE*, Benth. & Hook. f. i. 126.
PATRISIA, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110.
TETRACOCYNE, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 555.
acuminata, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. i. 492.—Bras.
Candollei, *Miq. in Ann. & Mag. Nat. Hist.* Sér. I. xi. (1843) 16.—Guian.
canescens, *Eichl. in Mart. Fl. Bras.* xiii. i. 490.—Bras.
chocoënsis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 117.—N. Granat.
dentata, *Miq. in Ann. & Mag. Nat. Hist.* Ser. I. xi. (1843) 16; *et Eichl. in Mart. Fl. Bras.* xiii. i. 493.—Venezuel.
Kunthii, Miq. in Ann. & Mag. Nat. Hist. Ser. I. xi. (1843) 16 = dentata.
Mansoana, *Eichl. in Mart. Fl. Bras.* xiii. i. 490.—Bras.
parviflora, Eichl. l. c. 493 = Patrisii.
Patrisii, *Miq. in Ann. & Mag. Nat. Hist.* Ser. I. xi. (1843) 16.—Bras.; Guian.
Reideliana, *Eichl. in Mart. Fl. Bras.* xiii. i. 491.—Bras.
Sagotiana, *Eichl. l. c.*—Bras.; Guian.
speciosa, *Vahl, Eclog. Am.* i. 51. t. 9.—Am. trop.
tomentosa, Miq. in Ann. & Mag. Nat. Hist. Ser. I. xi. (1843) 15 = speciosa?

RYCKIA, Balf. f. in Journ. Linn. Soc. xvii. (1878) 35, 48, sphalm. = RYKIA, De Vriese = **Pandanus**, Rumph.

RYDITOPHYLLUM, Walp. Rep. ii. 721 (1843), sphalm. = **Rytidophyllum**, Mart. (Gesner.).

RYDITOSTYLIS, Walp. Rep. ii. 204 (1843), sphalm. = RYTIDOSTYLIS, Hook. & Arn. = **Elaterium**, Linn. (Cucurb.).

RYKIA, De Vriese, in Verh. Kon. Akad. Wetensch. ii. (1854) 203 = **Pandanus**, Rumph.
furcata, De Vriese, l. c. = P. furcatus.

RYMANDRA, Salisb. in Knight, Prot. 124 (1809) = **Knightia**, R. Br. (1810) (Proteac.).
excelsa, Knight, l. c. = K. excelsa.

RYMIA, Endl. Gen. 743 (1839) = **Euclea**, Murr. (Ebenac.).
polyandra, Endl. Cat. Hort. Acad. Vindob. ii. (1843) 123 = E. polyandra.

RYNCHOSIA, Macfad. Fl. Jamaic. i. 275 (1837), sphalm. = **Rhynchosia**, Lour. (Legumin.).

RYNCHOSPORA, Vahl, Enum. ii. 229 (1806). *CYPERACEAE*, Benth. & Hook. f. iii. 1058.
ASTEROSCHOENUS, Nees, in Mart. Fl. Bras. ii. i. 124 (1842).
CALYPTROLEPIS, Steud. Syn. Pl. Cyp. 151 (1855).
CALYPTROSTYLIS, Nees, in Linnaea, ix. (1834) 295.

RYNCHOSPORA:—
CEPHALOSCHOENUS, Nees, in Linnaea, ix. (1834) 295.
CERATOSCHOENUS, Nees, l. c. 296 (1834).
CHAETOSPORA, H. B. & K. Nov. Gen. et Sp. i. 229 (1815).
DIPLOCHAETE, Nees, in Linnaea, ix. (1834) 296.
ECHINOSCHOENUS, Nees, in Linnaea, ix. (1834) 297.
EPHIPPIORHYNCHIUM, Nees, in Mart. Fl. Bras. ii. i. 134. t. 12 (1842).
EXPHALOSCHOENUS, Nees, in Hook. Journ. Bot. ii. (1840) 398.
HALOSCHOENUS, Nees, in Linnaea, ix. (1834) 296.
HAPLOSTYLIS, Nees, l. c. 295 (1834).
HYGROCHARIS, Nees, in Mart. Fl. Bras. ii. i. 147, in nota (1842).
KLEISTOCALYX, Steud. in Flora, xxxiii. (1850) 229, nomen.
LEPTOSCHOENUS, Nees, in Hook. Journ. Bot. ii. (1840) 393.
LONCHOSTYLIS, Torr. in Ann. Lyc. N. York, iii. (1836) 370.
MICROCHAETA, Reichb. Nom. 192 (1841).
MITROSPORA, Nees, in Linnaea, ix. (1834) 295.
MORISIA, Nees, in Edinb. N. Phil. Journ. (Oct. 1834) 265.
NEMOCHLOA, Nees, in Mart. Fl. Bras. ii. i. 148. tt. 14, 15 (1842).
PACHYMITRA, Nees, l. c. 115 (1842).
PLEUROSTACHYS, Brongn. in Duperr. Voy. Coq. Bot. 172. t. 31 (1829).
PTEROCHAETE, Arn. ex Boeck. in Linnaea, xxxvii. (1871–73) 600.
PTEROTHECA, Presl, Symb. Bot. i. 55. t. 35 (1831).
PTILOCHAETA, Nees, in Mart. Fl. Bras. ii. i. 147. t. 8 (1842).
PTILOSCIADIUM, Steud. Syn. Pl. Cyp. 149 (1855).
RHYNCHOSPORA, Willd. Enum. Hort. Berol. 71 (1809).
SPERMODON, Beauv. in Lestib. Ess. Fam. Cyp. 27 (1819).
SPHAEROSCHOENUS, Hook. & Arn. in Hook. Bot. Beech. Voy. 227, in adnot. (1841).
TRICHOCHAETA, Steud. Syn. Pl. Cyp. 155 (1855).
TRIODON, Rich. ex Pers. Syn. i. 60, in nota (1805).
ZOSTEROSPERMUM, Beauv. in Lestib. Ess. Fam. Cyp. 28 (1819).
adulta, Schlecht. in Bot. Zeit. vii. (1849) 166 = polyphylla.
alba, *Vahl, Enum.* ii. 236, partim.—Europ.; Am. bor.
alba, Vahl, l. c. partim = fusca.
albiceps, *Kunth, Enum. Pl.* ii. 289.—Bras.
albida, *Boeck. in Linnaea*, xxxvii. (1871–73) 561.—Bras.
alpina, Spreng. Syst. i. 195 = Carpha alpina.
amazonica, *Poepp. ex Kunth, Enum. Pl.* ii. 292.—Am. austr.
anomala, Steud. Syn. Pl. Cyp. 149 (Fimbristylidis sp.).—Java.
aristata, *Boeck. in Flora*, xl. (1857) 36.—Mexic.
armerioides, *J. & C. Presl, Rel. Haenk.* i. 197. t. 31. f. 2.—Am. trop.
articulata, Schult. Mant. ii. 49 = aurea.
arundinacea, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 27.—Bras.
asperula, Steud. Syn. Pl. Cyp. 144 = aurea.
aurea, Sieber, ex Schult. Mant. ii. 43, in obs. = cyperoides.
aurea, *Vahl, Enum. Pl.* ii. 229.—Reg. trop.
auriculata, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 144 = albiceps.
avenacea, Spreng. Syst. i. 197 = Cyathochaeta avenacea.
bahiensis, Steud. Nom. ed. II. ii. 455 = pterocarpa.
Baldwinii, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 210. f. 19.—Am. bor.
barbata, Kunth, Enum. Pl. ii. 290 = pterocarpa.
Berteroana, *Boeck. in Flora*, xli. (1858) 645.—Ins. S. Doming.
Beyrichii, *Steud. Syn. Pl. Cyp.* 144.—Bras.
biceps, Torr. ex A. Gray, in Ann. Lyc. N. York, iii. (1835) 219, in obs. = cephalantha, *A. Gray*.
biflora, *Boeck. in Kjoeb. Vidensk. Meddel.* (1871) 156.—Bras.
binervis, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 269 = Hypolytrum mauritianum.
Boivini, *Boeck. in Linnaea*, xxxviii. (1874) 403.—Ins. Borbon.

RYNCHOSPORA:—
brachychaeta, Wright, in Sauv. Fl. Cub. 180 = pallida, *M. A. Curt.*
brasiliensis, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 26.—Bras.
brevirostris, *Griseb. Cat. Pl. Cub.* 246.—Cuba.
bromoides, Kunth, Enum. Pl. ii. 300 = comata.
Brownii, Roem. & Schult. Syst. ii. 86 = glauca.
brunnea, *Boeck. in Linnaea*, xxxvii. (1871–73) 593.—Peruv.
brunnea, Willd. ex Kunth, Enum. Pl. ii. 297 = glauca.
bulbocaulis, Boeck. in Flora, lxii. (1879) 567 (Actinoschoeni sp.).—Afr. trop.
caduca, *Ell. Sketch*, i. 62.—Am. trop.
cajennensis, *Boeck. in Linnaea*, xxxviii. (1874) 404.—Guian.
canaliculata, *Boeck. in Flora*, lxiii. (1880) 451.—Bras.
candida, *Boeck. in Linnaea*, xxxvii. (1871–73) 605.—Afr. trop.; Am.
capillacea, *Boj. Hort. Maurit.* 386.—Hab.?
capillacea, Steud. Syn. Pl. Cyp. 141 = fusca.
capillacea, *Torr. Fl. U. St.* i. 55.—Am. bor.
capitata, *Roem. & Schult. Syst.* ii. 88.—Venezuela.
capitellata, Ell. Sketch, i. 61 = glomerata.
capitellata, Vahl, Enum. Pl. ii. 235 = glauca.
caracasana, *Boeck. in Linnaea*, xxxvii. (1871–73) 607.—Venezuela.
cariciformis, *Nees, in Mart. Fl. Bras.* ii. i. 145.—Bras.
castanea, Nees & Meyen, Reise, ii. 119; et ex Kunth, Enum. Pl. ii. 298 = glauca.
cephalantha, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 218. f. 30.—Am. bor.
cephalantha, *A. Rich. Fl. Cub. Fanerog.* ii. 293.—Cuba.
cephalophora, *Steud. Syn. Pl. Cyp.* 146.—Bras.
Cephalotes, *Vahl, Enum.* ii. 237.—Am. austr.
cephalotoides, Griseb. Cat. Pl. Cub. 242 = comata.
cernua, *Griseb. l. c.* 248.—Cuba.
ceylonica, Kunth, Enum. Pl. ii. 294 = triflora.
Chapmanii, *M. A. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 409.—Florida.
chinensis, Nees & Meyen, ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 108 = glauca.
ciliata, *Vahl, Enum.* ii. 235.—Am. bor.
ciliatifolia, Steud. Syn. Pl. Cyp. 143 = pterocarpa.
ciliolata, *Boeck. in Linnaea*, xxxvii. (1871–73) 563.—Bras.
cinnamomea, *Boeck. in Flora*, xli. (1858) 645.—Am. bor.
clandestina, Spreng. Syst. i. 194 = Cyathochaeta clandestina.
comata, *Link, Hort. Berol.* i. 280.—Am. trop.
compressa, *J. Carey, ex Chapm. Fl. S. U. St.* 525.—Florida.
conferta, *Boeck. in Linnaea*, xxxvii. (1871–73) 615.—Bras.
conferta, Chapm. ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 409 = Chapmanii.
conica, Desv. ex Ham. Prod. Pl. Ind. Occ. 15 = R. Cephalotes.
consanguinea, *Boeck. in Linnaea*, xxxvii. (1871–73) 536.—Bras.
coriifolia, *Boeck. l. c.* xxxviii. (1874) 406.—Bras.
corniculata, A. Gray, in Ann. Lyc. N. York, iii. (1835) 205 = laxa.
corymbifera, Boeck. in Linnaea, xxxvii. (1871–73) 619 = longiflora.
corymbifera, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 107.—Austral.
crassipes, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 146.—Bras.
cubensis, Griseb. Cat. Pl. Cub. 246 = filiformis.
cubensis, Griseb. Fl. Brit. W. Ind. 576 = tenuis.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 294.—Cuba.
Curtisii, Steud. Syn. Pl. Cyp. 141 = pallida, *M. A. Curt.*
curvula, *Boeck. in Linnaea*, xxxvii. (1871–73) 596.—Bras.
curvula, *Griseb. Fl. Brit. W. Ind.* 574.—Ins. Trinit.
cymosa, Ell. Sketch, i. 58 = fascicularis, *Vahl.*
cymosa, *Nutt. Gen. Am.* i. 33.—Am. bor.
cyperoides, *Mart. in Denkschr. Akad. Muench.* vi. 1816–17 (1820) 149.—Afr. et Am. trop.
decurrens, Chapm. Fl. S. U. St. 525 = patula.
deflexa, *Griseb. Cat. Pl. Cub.* 243.—Cuba.
deusta, Spreng. Syst. i. 195 = Mesomplaena deusta.
diandra, Spreng. l. c. 197 = Cyathochaeta diandra.

RYNCHOSPORA:—
diodon, Griseb. Cat. Pl. Cub. 245 in obs.; Boeck. in Linnaea, xxxvii. (1871–73) 558 = lunata.
discolor, Hochst. ex Steud. Syn. Pl. Cyp. 148 = Wightiana.
dissitiflora, *Steud. in Lechl. Berb. Am. Austr.* 56, nomen; *et ex Boeck. l. c.* 598.—Peruv.
distans, Beyr. ex Kunth, Enum. Pl. ii. 297 = glauca.
distans, Ell. Sketch, i. 59 = Grayii.
distans, *Vahl, Enum.* ii. 235.—Am. bor.
distans, Willd. ex Kunth, Enum. Pl. ii. 296 = glomerata.
distichophylla, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 26.—Bras.
divergens, *Chapm. ex M. A. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 409.—Am. bor.
dodecandra, Baldw. ex A. Gray, in Ann. Lyc. N. York, iii. (1835) 207 = megalocarpa.
Drummondiana, Boeck. in Flora, xli. (1858) 644 = gracilenta.
echinata, Willd. ex Kunth, Enum. Pl. ii. 284, 290 = Actinoschoenus filiformis.
effusa, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 142 = glauca.
elatior, *Kunth, Enum. Pl.* ii. 289.—Bras.
Elliotti, *A. Dietr. Sp. Pl.* ii. 69.—Am. bor.
Elliottii, A. Gray, in Ann. Lyc. N. York, iii. (1835) 204. f. 12 = Grayii.
emaciata, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 149.—Bras.
etuberculata, Steud. Syn. Pl. Cyp. 142 = Scirpus leptolepis.
exaltata, *Kunth, Enum. Pl.* ii. 291.—Bras.
exilis, *Boeck. in Flora*, lxiii. (1880) 440.—Bras.
eximia, *Boeck. in Linnaea*, xxxvii. (1871–73) 601.—Mexic.
Fabri, Steud. Syn. Pl. Cyp. 147 = Panicum lanatum.
fascicularis, Beyr. ex Kunth, Enum. Pl. ii. 296 = distans.
fascicularis, *Nutt. Gen. Am.* i. 33.—Am. bor.
fascicularis, *Vahl, Enum.* ii. 234.—Am. bor.
ferruginea, Roem. & Schult. Syst. ii. 85 = glauca.
ferruginea, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 269 = polyphylla.
filifolia, Kunth, Enum. Pl. ii. 299 = Sellowiana, *Steud.*
filifolia, *Torr. in Ann. Lyc. N. York*, iii. (1836) 366.—Am. bor.
filiformis, Balb. ex Kunth, Enum. Pl. ii. 297 = distans.
filiformis, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 269 = cymosa.
filiformis, *Vahl, Enum.* ii. 232.—Am. centr.
filiformis, Willd. ex Kunth, Enum. Pl. ii. 281 = Dichromena caracasana.
flavida, *Boeck. in Flora*, lxiii. (1880) 439.—Bras.
florida, A. Dietr. Sp. Pl. ii. 71 = aurea.
florida, Griseb. Fl. Brit. West Ind. 575 = longiflora.
foliata, *Steud. Syn. Pl. Cyp.* 142.—Am. bor.
fusca, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 127—Europ.; Am. bor.
fusca, A. Gray, in Ann. Lyc. N. York, iii. (1835) 216 = gracilenta.
gigantea, *Link, Jahrb.* i. III. (1820) 76.—Bras.
glauca, Baldw. ex A. Gray, in Ann. Lyc. N. York, iii. (1835) 210 = Baldwinii.
glauca, *Vahl, Enum.* ii. 233.—Geront. trop.
Glaziovii, *Boeck. in Linnaea*, xxxviii. (1874) 405.—Bras.
globosa, Desv. ex Ham. Prod. Pl. Ind. Occ. 15 = pterocarpa.
globosa, *Roem. & Schult. Syst.* ii. 89.—Am. austr.
globulifera, Link, Jahrb. i. III. (1820) 76 = cyperoides.
glomerata, *Vahl, Enum.* ii. 234.—Am. bor.
glomerata, Vahl, ex Kunth, Enum. Pl. ii. 216 = Scirpus flaccifolius.
Gollmeri, *Boeck. in Linnaea*, xxxvii. (1871–73) 556.—Venezeula.
gracilenta, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 215. f. 27.—Am. bor.
gracilis, Nees, ex Kunth, Enum. Pl. ii. 298 = laxa.
gracilis, Vahl, Enum. ii. 234 = glauca.
gracilis, Willd. ex Kunth, Enum. Pl. ii. 300 = polyphylla.
gracillima, *Thw. Enum. Pl. Zeyl.* 435.—Zeylan.
gracillima, Wright, in Sauv. Fl. Cub. 181 = pallida, *M. A. Curt.*
Grayana, Chapm. ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 409 = Chapmanii.
Grayii, *Kunth, Enum. Pl.* ii. 539.—Am. bor.

RYNCHOSPORA:—
Griffithii, *Boeck. in Linnaea*, xxxviii. (1874) 404.—Ind. or.
Haenkei, J. & C. Presl, Rel. Haenk. i. 199 = Wallichiana.
Hamiltonii, Kunth, Enum. Pl. ii. 291 = pterocarpa.
Harveyi, W. Boott, in Coult. Bot. Gaz. ix. (1884) 85 = Torreyana.
heterophylla, Boeck. in Linnaea, xxxvii. (1871–73) 553 = Chapmanii.
Hildebrantii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 508.—Madag.
hirsuta, Griseb. Fl. Brit. W. Ind. 576 = Dichromena ciliata.
hirsuta, *Vahl, Enum.* ii. 231.—Am. trop.
hirta, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 146, *in obs.*; *et in Linnaea*, xxxvii. (1871–73) 543.—Bras.
hispidula, *Boeck. l. c.* (1871–73) 604.—Guian.
hispidula, *Griseb. Cat. Pl. Cub.* 245.—Cuba.
Hoffmanni, *Boeck. in Linnaea*, xxxvii. (1871–73) 637.—Costa Rica.
Hookeri, *Boeck. l. c.* 621.—Ind. or.
Humboldtiana, Griseb. Fl. Brit. W. Ind. 577 = Dichromena repens.
humilis, Beyr. ex Kunth, Enum. Pl. ii. 297 = cymosa.
ignorata, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 508.—Cuba.
inexpansa, Sieber, ex Steud. Nom. ed. II. ii. 456 = longispicata.
inexpansa, *Vahl, Enum.* ii. 232.—Am. bor.
intermedia, Beyr. ex Kunth, Enum. Pl. ii. 296 = cephalantha, *A. Gray.*
intermixta, Wright, in Sauv. Fl. Cub. 184 = pusilla, *Chapm.*
Jelskiana, *Boeck. in Linnaea*, xxxviii. (1874) 401.—Guian.
jubata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 254 = R. Marisculus.
juncea, Willd. ex Kunth, Enum. Pl. ii. 298 = glauca.
junciformis, *Boeck. in Flora*, xli. (1858) 646.—Guian.
Kamphoeveneri, Boeck. in Engl. Bot. Jahrb. v. (1884) 508 = gracillima.
Knieskernii, *J. Carey, in Am. Journ. Sc.* Ser. II. iv. (1847) 25.—Am. bor.
Kunthii, *Nees, in Linnaea*, ix. (1834) 297.—Mexic.
Kyssor, A. Dietr. Sp. Pl. ii. 74 = Scirpus grossus.
laevigata, *Steud. Syn. Pl. Cyp.* 145.—Guian.
laevis, Schlecht. in Linnaea, xxv. (1852) 166 = glauca.
lamprosperma, Wright, in Sauv. Fl. Cub. 183 = divergens.
lanceolata, Willd. ex Kunth, Enum. Pl. ii. 288 = capitata.
lavarum, *Gaudich. in Freyc. Voy. Bot.* 415.—Ins. Sandvic.
lavarum, Hook. & Arn. Bot. Beech. Voy. 98 = glauca.
laxa, R. Br. Prod. 230 = glauca.
laxa, Schult. Mant. ii. 46, partim = longiflora.
laxa, *Vahl, Enum.* ii. 231.—Am. bor.
Lechleri, *Steud. in Lechl. Berb. Am. Austr.* 56, nomen; *et ex Boeck. in Linnaea*, xxxvii. (1871–73) 633.—Peruv.
leptorhyncha, *Wright, in Sauv. Fl. Cub.* 180.—Cuba.
leptostachya, Boeck. in Linnaea, xxxvii. (1871–73) 591 = tenuis.
leucocarpa, *Ridl. in Journ. Linn. Soc.* xx. (1883) 335.—Madag.
leucocephala, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 144 = Dichromena leucocephala.
leucocephala, *Nees & Meyen, in Linnaea*, ix. (1834) 297, nomen.—Hab.?
Lindeniana, *Griseb. Cat. Pl. Cub.* 244.—Cuba.
longibracteata, *Boeck. in Linnaea*, xxxvii. (1871–73) 534.—Guian.
longiflora, *Presl, in Oken, Isis*, xxi. (1828) 269.—Ins. Martinic.
longirostris, Ell. Sketch, i. 59 = laxa.
longirostris, Steud. Syn. Pl. Cyp. 145 = amazonica.
longiseta, Steud. l. c. 140 = caduca.
longisetis, *R. Br. Prod.* 230.—Austral.
longisetis, F. Muell. Fragm. ix. 17 = tenuifolia.
longispicata, *Boeck. in Linnaea*, xxxvii. (1871–73) 600.—Bras.
lunata, *Griseb. Cat. Pl. Cub.* 244.—Am. trop.
Lundii, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 147.—Bras.
luzuliformis, *Boeck. in Linnaea*, xxxviii. (1871–73) 632.—Am. austr.

RYNCHOSPORA:—
macrocarpa, Boeck. in Linnaea, xxxvii. (1871–73) 629 = spectabilis.
macrochaeta, *Steud. in Lechl. Berb. Am. Austr.* 56, nomen; *et ex Boeck. l. c.* 632.—Peruv.
macrostachya, *Torr. ex A. Gray, in Ann. Lyc. N. York*, iii. (1835) 206. t. 6. f. 14.—Am. bor.
macrostylis, A. Dietr. Sp. Pl. ii. 74 = pterocarpa.
marginata, Steud. Syn. Pl. Cyp. 142 = globosa.
Marisculus, *Nees, in Linnaea*, ix. (1834) 297.—Am. trop.
Mauritii, Steud. Syn. Pl. Cyp. 149 = cyperoides.
megalocarpa, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 208. f. 16.—Am. bor.
mexicana, *Steud. Syn. Pl. Cyp.* 144.—Mexic.
mexicana, Wright, in Sauv. Fl. Cub. 182 = rufa.
micrantha, A. Gray, in Ann. Lyc. N. York, iii. (1835) 197 = Torreyana.
micrantha, Pursh, Fl. Am. Sept. i. 48 = rariflora.
micrantha, *Vahl, Enum.* ii. 231.—Mexic.
microcarpa, *Baldw. ex A. Gray, in Ann. Lyc. N. York* iii. (1835) 202. f. 9.—Am. bor.
microseta, Steud. Syn. Pl. Cyp. 142 = ciliata.
miliacea, A. Gray, in Ann. Lyc. N. York, iii. (1835) 198. f. 4 = sparsa.
millegrana, *Schrad. ex Steud. Syn. Pl. Cyp.* 139.—Am. trop.
Minarum, *Steud. l. c.* 143.—Bras.
minor, *Nees, in Mart. Fl. Bras.* ii. 1. 143.—Bras.
mollis, *Schult. Mant.* ii. 49.—Nepal.
monostachya, Steud. Syn. Pl. Cyp. 140 = Eleocharis tuberculosa.
Moritziana, *Boeck. in Linnaea*, xxxvii. (1871–73) 640.—Venezuela.
multiflora, A. Gray, in Ann. Lyc. N. York, iii. (1835) 200. f. 7 = Elliotti.
multiflora, Nees, in Mart. Fl. Bras. ii. 1. 143 = polyantha.
nardifolia, *Boeck. in Linnaea*, xxxvii. (1871–73) 560.—Bras.
Neesii, *Steud. Syn. Pl. Cyp.* 146.—Bras.
nervosa, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 143 = Dichromena ciliata, pulchella.
nitens, *A. Gray, Man. Bot. N. U. St.* ed. V. 568, in nota.—Am. bor.
nitida, Spreng. Neue Entdeck. iii. 10 = Tetraria compar.
nivea, Boeck. in Linnaea, xxxvii. (1871–73) 527 (Dichromenae sp.).—Am. bor.
ochrocephala, *Boeck. in Flora*, lxii. (1879) 568.—Afr. trop.
odorata, Griseb. Cat. Pl. Cub. 242 = R. Marisculus.
oligantha, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 212. f. 22.—Texas.
oligosantha, Endl. ex Kunth, Enum. Pl. ii. 216 = Scirpus flaccifolius.
oxycephala, Wright, in Sauv. Fl. Cub. 184 = eximia.
pallens, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 146 = Sellowiana, *Steud.*
pallida, *M. A. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 409.—Am. bor.
pallida, *Steud. Syn. Pl. Cyp.* 146.—Bras.
paniculata, *A. Gray. in Ann. Lyc. N. York*, iii. (1835) 211. f. 21.—Am. bor.; Ind. occ.
paniculata, Presl, in Oken, Isis, xxi. (1828) 269 = polyphylla.
paniculifera, Steud. Syn. Pl. Cyp. 146 = Schiedeana.
paraënsis, *Schrad. ex Kunth, Enum. Pl.* ii. 300.—Bras.
parva, Kunth, l. c. 302 = Wallichiana.
parviflora, Beyr. ex Kunth, l. c. 297 = microcarpa.
parviflora, Vahl, ex Kunth, l. c. 279 = micrantha.
patula, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 201. f. 8.—Am. bor.
pauciflora, Presl, in Oken, Isis, xxi. (1828) 269 = cymosa.
pauciflora, Willd. ex Kunth, Enum. Pl. ii. 284 = Actinoschoenus filiformis.
pendula, A. Gray, in Ann. Lyc. N. York, iii. (1835) 201 = Elliotti.
penniseta, Griseb. Cat. Pl. Cub, 244 = plumosa.
Persooniana, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 536 = Dichromena repens.
petraea, Nees, in Mart. Fl. Bras. ii. 1. 142 = glauca.
pilosa, *Boeck. in Linnaea*, xxxvii. (1871–73) 537.—Bras.
pilulifera, *Bertol. in Bologn. Opusc. Sc.* iii. (1819) 406.—Bras.
plumosa, *Ell. Sketch*, i. 58.—Am. bor.
podosperma, Wright, in Sauv. Fl. Cub. 183 = longispicata.

RYNCHOSPORA :—
Pohliana, Steud. Syn. Pl. Cyp. 142 = globosa.
polyantha, *Steud. l. c.* 147.—Bras.
polycephala, Presl, in Oken, Isis, xxi. (1828) 269 = polyphylla.
polycephala, Wydl. ex Kunth, Enum. Pl. ii. 291 = cyperoides.
polyphylla, Balb. ex Kunth, l. c. 279 = micrantha.
polyphylla, *Vahl, Enum.* ii. 230.—Ind. occ.
Prescottiana, Wall. Cat. n. 3423 = longisetis.
Presleana, Steud. Syn. Pl. Cyp. 142 = globosa.
pruinosa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 535.—Cuba.
Pseudo-mariscus, Kunth, Enum. Pl. ii. 303 = gigantea.
psilocaryoides, *Griseb. Cat. Pl. Cub.* 247.—Cuba.
pterocarpa, *Roem. & Schult. Syst.* ii. 89.—Am. austr.
pterochaeta, *F. Muell. Fragm.* ix. 17.—Austral.
pubera, Boeck. in Linnaea, xxxvii. (1871–73) 528 = Dichromena pubera.
punctata, *Boeck. l. c.* 580.—Mexic.; Venezuel.
punctata, Ell. Sketch, i. 60 = cymosa.
pungens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 253 = glauca.
pura, Griseb. Fl. Brit. W. Ind. 577 = Dichromena ciliata.
pusilla, *Chapm. ex M. A. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 409; *Fl. S. U. St.* 528.—Florida.
pusilla, *Griseb. in Goett. Abh.* vii. (1857) 271.—Ind. occ.
pycnocarpa, A. Gray, in Ann. Lyc. N. York, iii. (1835) 208. f. 17 = megalocarpa.
pyramidata, Desv. ex Ham. Prod. Pl. Ind. Occ. 15 = polyphylla.
racemosa, *Wright, in Sauv. Fl. Cub.* 182.—Cuba.
rariflora, *Ell. Sketch*, i. 58.—Am. bor.
Ravenelii, M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 410 = divergens.
recurvata, *Steud. Syn. Pl. Cyp.* 145.—Bras.
repens, Boeck. in Linnaea, xxxvii. (1871–73) 531 = Dichromena repens.
rigida, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 145.—Bras.
rigida, Schrad. ex Kunth, Enum. Pl. ii. 297 = glauca.
rigida, Spreng. Syst. i. 196 = polyphylla.
riparia, *Boeck. in Linnaea*, xxxvii. (1871–73) 561.—Bras.
robusta, *Boeck. l. c.* 616.—Bras.
Rudioi, *Boeck. in Flora*, lxv. (1882) 27.—Bras.
rufa, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 149.—Bras.
Ruiziana, *Boeck. in Linnaea*, xxxvii. (1871–73) 641.—Peruv.
ruppioides, Benth. in Hook. Ic. Pl. t. 1344 = Scirpus submersus.
Sartoriana, *Boeck. in Flora*, xl. (1857) 36.—Mexic.
scaberrima, *Boeck. l. c.* lxiii. (1880) 452.—Bras.
scabrata, Griseb. Cat. Pl. Cub. 243 = pruinosa.
scapigera, *Boeck. in Linnaea*, xxxviii. (1874) 402.—Guian.
Schaffneri, *Boeck. l. c.* xxxvii. (1871–73) 575.—Mexic.
Schiedeana, *Kunth, Enum. Pl.* ii. 300.—Mexic.
schoenoides, Wood, Class-Book (1861), 744 = Elliotti.
Schottmuelleri, Boeck. in Flora, lxv. (1882) 27 (Pleurostachys sp.).—Bras.
Schraderiana, Steud. Syn. Pl. Cyp. 146 = aurea.
scirpoides, *Griseb. Cat. Pl. Cub.* 247, in obs.; *A. Gray, Man. Bot. N. U. St.* ed. V. 568.—Am. bor.
sclerioides, Hook. & Arn. Bot. Beech. Voy. 99 = thyrsoidea.
scutellata, *Griseb. Cat. Pl. Cub.* 246.—Cuba.
Selloiana, *Boeck. in Linnaea*, xxxvii. (1871–73) 620.—Bras.
Sellowiana, *Steud. Nom.* ed. II. ii. 456; *et Syn. Pl. Cyp.* 145.—Bras.
semihirsuta, *Boeck. in Linnaea*, xxxvii. (1871–73) 614.—Bras.
semiinvolucrata, *J. & C. Presl, Rel. Haenk.* i. 198.—Peruv.
semiplumosa, *A. Gray, in Ann. Lyc. N. York*, iii. (1835) 213. t. 6. f. 23.—Am. bor.
senegalensis, *Steud. Syn. Pl. Cyp.* 149.—Afr. trop.
seslerioides, Griseb. Cat. Pl. Cub. 247 (Dichromenae sp.).—Cuba.
setacea, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 159 = tenerrima.
setacea, *Vahl, Enum.* ii. 233.—Am. austr.

RYNCHOSPORA :—
setacea, Willd. ex Kunth, Enum. Pl. ii. 297 = rariflora.
setigera, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 145.—Bras.
setigera, Griseb. Fl. Brit. W. Ind. 577 = longibracteata.
sinensis, Steud. Nom. ed. II. ii. 456 = *chinensis*.
sparganioides, Boj. Hort. Maurit. 386 = cyperoides.
sparganoides, Steud. Syn. Pl. Cyp. 144 = exaltata.
sparsa, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 268 = micrantha.
sparsa, *Vahl, Enum.* ii. 230.—Am. bor.
speciosa, *Boeck. in Linnaea*, xxxvii. (1871–73) 536.—Bras.
speciosa, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 138 = aurea.
spectabilis, *Hochst. ex Krauss, in Flora*, xxviii. (1845) 760.—Afr. austr.
spermodon, Griseb. Fl. Brit. W. Ind. 576 = tenerrima, *Nees*.
spicata, Spreng. Syst. i. 194 = Scleria pleostachya.
stellata, Griseb. in Goett. Abh. vii. (1857) 271 = Dichromena leucocephala.
stenocarpa, *Kunth, Enum. Pl.* ii. 294.—Bras.
stenophylla, *Chapm. Fl. S. U. St.* 525.—Florida.
stenorhyncha, Griseb. Fl. Brit. W. Ind. 575 = triflora.
stipitata, Chapm. Fl. S. U. St. Suppl. 660 = R. Marisculus.
stolonifera, *Boeck. in Linnaea*, xxxvii. (1871–73) 562.—Bras.
stricta, *Boeck. l. c.* 603.—Bras.; Guian.
subcapitata, Boeck. l. c. 547 = R. Minarum.
suberostris, Boeck. l. c. 579 = Grayii.
subimberbis, *Griseb. Cat. Pl. Cub.* 245.—Cuba.
sublateralis, Steud. Syn. Pl. Cyp. 148 (Schoeni sp.).—Austral.
subulirostris, Steud. l. c. 149 = aurea.
surinamensis, Nees, in Linnaea, ix. (1834) 297 = aurea.
sylvatica, Nees, l. c. = comata.
tenella, *Boeck. in Linnaea*, xxxvii. (1871–73) 595.—Bras.
tenerrima, *Boeck. l. c.* 554.—Venezuela.
tenerrima, *Nees, ex Spreng. Syst.* iv. *Cur. Post.* 26.—Am. trop.
tenuifolia, *Benth. Fl. Austral.* vii. 350.—Austral.
tenuifolia, Griseb. Cat. Pl. Cub. 244 = stenophylla.
tenuirostris, Kunth, Enum. Pl. ii. 303 = cyperoides.
tenuis, Bald. ex A. Gray, in Ann. Lyc. N. York, iii. (1835) 217 = distans.
tenuis, *Link, Jahrb.* i. III. (1820) 76.—Bras.
tenuiseta, Wright, in Sauv. Fl. Cub. 179 = R. Marisculus.
terminalis, *Steud. Syn. Pl. Cyp.* 143.—Bras.
testacea, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 148.—Bras.
texensis, Boeck. in Linnaea, xxxvii. (1871–73) 602 = nitens.
Thouarsii, Nees, in Linnaea, ix. (1834) 297 = glauca.
thyrsoidea, *Nees & Meyen, l. c.*—Ins. Sandvic.
Torreyana, *A. Gray, in Ann. Lyc. York*, iii. (1835) 197.—Am. bor.
Torreyana, Griseb. Cat. Pl. Cub. 243 = cymosa.
triceps, *Boeck. in Linnaea*, xxxvii. (1871–73) 642.—Peruv.
triceps, Hochst. in Flora, xxviii. (1845) 759 = cyperoides.
triceps, *Steud. Nom.* ed. II. ii. 456.—Am. austr.
trichophora, Nees, in Mart. Fl. Bras. ii. 1. 145 = caricinformis.
triflora, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 140 = oligantha.
triflora, *Vahl, Enum.* ii. 231.—As. et Am. trop.
trigyna, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 21.—Abyss.
trispicata, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 136 = triflora.
umbraticola, *Poepp. & Kunth, in Kunth, Enum. Pl.* ii. 300.—Peruv.
uniflora, *Boeck. in Flora*, lxiii. (1880) 439.—Bras.
Urvillei, Steud. Syn. Pl. Cyp. 149 = cyperoides.
Vahliana, Griseb. Fl. Brit. W. Ind. 577 = Dichromena ciliata.
velutina, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 149.—Bras.
vicozensis, Schult. Mant. ii. 46 = glauca.
vulcani, *Boeck. in Linnaea*, xxxvii. (1871–73) 638.—Costa Rica.

RYNCHOSPORA :—
Wallichiana, *Kunth, Enum. Pl.* ii. 289.—Ind. or.
Warmingii, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 143.—Bras.
Wightiana, *Steud. Syn. Pl. Cyp.* 148.—Ind. or.
zeylanica, Thw. Enum. Pl. Zeyl. 352 = triflora.

RYNCHOSTYLIS, Blume, Bijdr. 285. t. 49 (1825) = **Rhynchostylis**, Blume (Orchid.).

RYPARIA, Blume, Fl. Jav. Praef. 8 (1828) = **Ryparosa**, Blume (Euphorb.?).
caesia, Blume, ex Baill. Etud. Gén. Euph. 339 = Ryparosa caesia.
dubia, Blume, ex Baill. l. c.—Hab.?

RYPAROSA, Blume, Bijdr. 600 (1825). Gen. dub. *EUPHORBIACEARUM*.—Cf. Benth. & Hook. f. iii. 257.
RYPARIA, Blume, Fl. Jav. Praef. 8 (1828).
caesia, *Blume, Bijdr.* 601.—Java.

RYSODIUM, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 263 = **Astragalus**, Tourn. (Legumin.).
pterostylis, Stev. l. c. = A. pterostylis.

RYSSOPTERYS, Blume, ex A. Juss. in Deless. Ic. Sel. iii. 21. t. 35 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 257.
abutilifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 385.—N. Hibern.
chrysantha, *Hassk. Hort. Bog. Desc.* 133.—Java.
Cumingiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 386.—Ins. Philipp.
dealbata, *A. Juss. l. c.*—Ins. Philipp.
microstema, *A. Juss. l. c.* 384.—Ins. Philipp.; Java.
ovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 583.—Ins. Philipp.
rufescens, *Turcz. l. c.*—Java.
tiliaefolia, Blume, ex A. Juss. in Arch. Mus. Par. iii. (1843) 384 = microstema.
tiliaefolia, *A. Juss. l. c.*—Java.
timorensis, *Blume, ex A. Juss. in Deless. Ic. Sel.* iii. 21.—Ins. Timor.
tomentosa, Blume, ex A. Juss. in Arch. Mus. Par. iii. (1843) 514 = Aspidopterys tomentosa.

RYTACHNE, Endl. Gen. 109 (1837) = **Rhytachne**, Desv. (Gramin.).

RYTICARYUM, Becc. in Malesia, i. (1877) 120. *ICACINEAE*.
fasciculatum, *Becc. l. c.* 121.—N. Guin.
macrocarpum, *Becc. l. c.* (1878) 256.—N. Guin.
oleraceum, *Becc. l. c.* (1877) 121.—'Grande Kei.'
racemosum, *Becc. l. c.*—N. Guin.

RYTIDEA, Spreng. Syst. i. 515 (1825) = **Rutidea**, DC. (Compos.).

RYTIDOLOMA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 319 = **Dictyanthus**, Decne. (Asclep.).
reticulatum, Turcz. l. c. 320 = D. reticulatus.

RYTIDOPHYLLUM, Mart. Nov. Gen. et Sp. iii. 38 (1829) = **Rhytidophyllum**, Mart. (Gesnerac.).

RYTIDOSPERMA, Steud. Syn. Pl. Gram. 425 (1855) = **Deschampsia**, Beauv. (Gramin.).
Lechleri, Steud. l. c. = D. caespitosa?

RYTIDOSPERMUM, Benth. in Journ. Linn. Soc. xix. (1881) 96, sphalm. = praec.

RYTIDOSTYLIS, Hook. & Arn. Bot. Beech. Voy. 424. t. 97 (1841) = **Elaterium**, Jacq. (Cucurb.).
amazonica, Spruce, ex Cogn. Diagn. Cucurb. fasc. ii. 55 = E. amazonicum.
gracilis, Hook. & Arn. Bot. Beech. Voy. 425. t. 97 = E. gracile.

RYTIDOTUS, Hook. f. in Hook. Ic. Pl. t. 1071 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 101.
RHYTIDOTUS, Hook. f. ex Benth. & Hook. f. Gen. ii. 101 (1873).
sandvicensis, *Hook. f. in Hook. Ic. Pl.* t. 1071.—Ins. Sandvic.

RYTIGLOSSA, Steud. Nom. ed. II. ii. 488 (1841) = RHYTIGLOSSA, Nees = **Dianthera**, Linn. (Acanth.).

RYTIGYNIA, Blume, Mus. Bot. Lugd. Bat. i. 178 (1850) = **Vangueria**, Juss. (Rubiac.).
senegalensis, Blume, l. c. 179.—Afr. trop.

RYTILIX, Rafin. in Ser. Bull. Bot. i. (1830) 219 = **Manisuris**, Sw. (Gramin.).
glandulosa, Rafin. l. c. = M. granularis.

S

SABADILLA, Brandt & Ratzeb. in Hayne, Arzn. Gew. xiii. t. 27 (1837) = **Schoenocaulon**, A. Gray (Liliac.).
officinarum, Brandt & Ratzeb. l. c. = Sch. officinale.

SABAL, Adans. Fam. ii. 495 (1763). *PALMAE*, Benth. & Hook. f. iii. 922.
Adansoni, *Guerns. in Bull. Soc. Philom.* iii. (1803) 206. t. 25.—Am. bor.
adiantinum, Rafin. Fl. Ludov. 17 (Quid?).—Am. bor.
Blackburnianum, *Glazebrook, in Loud. Gard. Mag.* v. (1829) 52. ff. 10–13.—Ind. occ.
carolinianum, Hort. ex Poir. Encyc. vi. 356 = Adansoni.
coerulescens, *Hort.—Cf. Gard. Chron.* (1875) i. 589. —N. Granat.
columnare, *Hort. ex Mart. Hist. Nat. Palm.* iii. 320, 348, nomen.—Hab.?
elatum, Hort. ex Mart. l. c.—Hab.?
giganteum, Fulchir. ex Steud. Nom. ed. II. ii. 489 = mexicanum?
glaucescens, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 320, nomen.—Ins. Trinit.
graminifolium, *Lodd. ex Loud. Hort. Brit.* 141, nomen. —Am. austr.
havanensis, Lodd. ex Mart. Hist. Nat. Palm. iii. 320, 348 (Quid?).—Hab.?
Histrix, Pursh, ex Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 293 = Rhapidophyllum Hystrix.
Magdalenae, *Linden, in Illustr. Hortic.* xxviii. (1881) 32, nomen.—N. Granat.
mauritiiforme, *Griseb. & H. Wendl. in Griseb. Fl. Brit. W. Ind.* 514.—Ind. occ.
mexicanum, *Mart. Hist. Nat. Palm.* iii. 246. t. 8.—Mexic.
minimum, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 293 = Adansoni.
minus, Pers. Syn. i. 399 = Adansoni.
oleraceum, Hort. ex Mart. Hist. Nat. Palm. iii. 320, 348 (Quid?).—Hab.?
Palmetto, Hort. l. c. 246 = mexicanum.
Palmetto, *Lodd. ex Schult. f. Syst* vii. 1487.—Am. bor.
Palmetto, Rein, in Ber. Senckenb. Naturf. Ges. Frankf. (1873) 150 = Blackburnianum.
pumila, Ell. Sketch, i. 430 = Adansoni.
Sanfordi, *Linden, in Illustr. Hortic.* xxviii. (1881) 32, nomen.—Florida.
serrulatum, Schult. f. Syst. vii. 1486 = Brahea serrulata.
taurinum, Hort. ex Mart. Hist. Nat. Palm. iii. 320, 348 (Quid?).—Jamaic.
umbraculiferum, Hort. ex Mart. l. c. 246 = mexicanum.
umbraculiferum, Mart. l. c. 245. t. 130 = Blackburnianum.
Woodfordii, *Lodd. ex Mart. l. c.* 247, nomen.—Ins. Trinit.

SABATIA, Adans. Fam. ii. 503 (1763) = **Sabbatia**, Adans. (Gentian.).

SABAZIA, Cass. in Dict. Sc. Nat. xlvi. 480 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 362.
BAZIASA, Steud. Nom. ed. II. i. 192 (1840).
humilis, *Cass. in Dict. Sc. Nat.* xlvi. 481.—Mexic.
microglossa, *DC. Prod.* v. 497.—Mexic.
occidentalis, DC. l. c., sphalm. = *Siegesbeckia occidentalis.*
portoriccensis, DC. l. c., sphalm. = *Siegesbeckia portoriccensis.*
sarmentosa, *Less. in Linnaea*, v. (1830) 148.—Mexic.
urticaefolia, *DC. Prod.* v. 497.—Am. trop.

SABBATA, Vell. Fl. Flum. 352 (1825); viii. tt. 94–95 (1827). Gen. dub. *COMPOSITARUM.*—Cf. Benth. & Hook. f. ii. 223.
polyphylla, Vell. l. c. viii. t. 95.—Bras.
romana, Vell. l. c. 352; viii. t. 94.—Bras.

SABBATIA, Adans. Fam. ii. 503 (1763) (Sabatia). *GENTIANEAE*, Benth. & Hook. f. ii. 810.
LAPITHEA, Griseb. in DC. Prod. ix. 48 (1845).
NEUROLA, Rafin. New Fl. Am. iv. 92 (1836).
amena, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
amoena, G. Don, Gen. Syst. iv. 207 = stellaris.
anceps, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
angularis, *Pursh, Fl. Am. Sept.* i. 137.—Am. bor.
australis, Cham. & Schlecht. in Linnaea, i. (1826) 194 = Zygostigma australe.
Boykini, *A. Gray, ex Chapm. Fl. S. U. St.* 354.—Am. bor.
brachiata, *Ell. Sketch*, i. 285.—Am. bor.
brevifolia, Rafin. Atl. Journ. 147 (Quid?).—Am. bor.
calycosa, *Pursh, Fl. Am. Sept.* i. 138.—Am. bor.
campanulata, Torr. ex Griseb. Gen. et Sp. Gent. 120 = gracilis.
campestris, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 197.—Am. bor.
chloroides, *Pursh, Fl. Am. Sept.* i. 137.—Am. bor.
concinna, Wood, Class-Book (1861), 584 = brachiata.
corymbosa, *Baldw. ex Ell. Sketch*, i. 283.—Am. bor.
cymosa, G. Don, Gen. Syst. iv. 207 = corymbosa.
diffusa, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
Elliottii, *Steud. Nom.* ed. II. ii. 489 (Elliotii).—Am. bor.
formosa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 7 = campestris.
gentianoides, *Ell. Sketch*, i. 286.—Am. bor.
gracilis, Ell. l. c. 284 = stellaris.
gracilis, *Salisb. Parad. Lond.* t. 32.—Am. bor.
heterophyla, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
lanceolata, Torr. & Gray, ex A. Gray, Man. Bot. N. U. St. ed. I. 356 = corymbosa.
lingulata, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
macrophylla, *Hook. Comp. Bot. Mag.* i. (1835) 171.—Am. bor. occ.
maculata, *Benth. & Hook. f. Gen.* ii. 812, in obs.—Mexic.
nervosa, *Rafin. New Fl. Am.* iv. 92.—Am. bor.
nivea, *Rafin. Med. Fl.* ii. 77.—Am. bor.
obtusifolia, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
oligophylla, Featherman, in Univ. Mississip. Rep. (1871); ex A. Gray, Syn. Fl. N. Am. ii. 1. 115 = gentianoides.
paniculata, Ell. Sketch, i. 282 = Elliottii.
paniculata, *Pursh, Fl. Am. Sept.* i. 138.—Am. bor.
petiolata, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
pulchella, Rafin. ex Griseb. in DC. Prod. ix. 49 = gracilis.
pulchella, Spreng. Syst. i. 581 = Erythraea ramosissima?
pumila, Rafin. Fl. Tellur. iii. 30 (Quid?).—Hab.?
simplex, Bertol. Misc. Bot. x. 27 = Rhexia stricta.
stellaris, *Pursh, Fl. Am. Sept.* i. 137.—Am. bor.
stricta. Rafin. Fl. Tellur. iii. 30 = Curtia gentianoides?
tenuifolia, Rafin. l. c. (= *Schuebleria tenuifolia*?).—Am. trop.
umbellata, Rafin. l. c. (Quid?).—Hab.?
verticillaris, Spreng. Syst. i. 580 = Curtia gentianoides.

SABBATIA, Moench, Meth. 386 (1794) = **Micromeria**, Benth. (Labiat.).
corymbosa, Moench, l. c. = M. Juliana.

SABDARIFFA, Kostel. Allg. Med. Pharm. Fl. v. 1857 (1836) = **Hibiscus**, Linn. (Malv.).
digitata, Kostel. l. c. = H. digitatus.
rubra, Kostel. l. c. = H. Sabdariffa.

SABIA, Colebr. in Trans. Linn. Soc. xii. (1818) 355. t. 14. *SABIACEAE*, Benth. & Hook. f. i. 414.
ANDROGLOSSA, Benth. in Hook. Kew Journ. iii. (1851) 42.
ENANTIA, Falc. in Hook. Journ. Bot. iv. (1842) 75.
MENICOSTA, D. Dietr. Syn. Pl. ii. 923 (1839).
MENISCOSTA, Blume, Bijdr. 28 (1825).
Bullockii, Hance, in Journ. Bot. xvi. (1878) 9 = japonica.
campanulata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 311.—Reg. Himal.

SABIA:—
celastrinea, C. Muell. in Walp. Ann. vi. 1269 (Index) = limoniacea.
densiflora, *Miq. Fl. Ind. Bat. Suppl.* 520.—Sumatra.
elliptica, *Miq. l. c.*—Sumatra.
floribunda, Miq. l. c. = Meliosma elliptica.
japonica, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 430.—China; Japon.
lanceolata, *Colebr. in Trans. Linn. Soc.* xii. (1819) 355. t. 14.—Reg. Himal.
leptandra, *Hook. f. & Thoms. Fl. Ind.* i. 209.—Reg. Himal.
limoniacea, *Wall. Cat.* n. 1000.—Reg. Himal.
malabarica, *Bedd. Ic. Pl. Ind. Or.* i. 39.—Ind. or.
Meniscosta, *Blume, Mus. Bot. Lugd. Bat.* i. 369.—Malaya.
paniculata, *Edgew. ex Hook. f. & Thoms. Fl. Ind.* i. 211. —Reg. Himal.
paniculata, Seem. Bot. Voy. Herald, 362 = limoniacea.
parviflora, Wall. Cat. n. 1001 = purpurea.
parviflora, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 310. —Reg. Himal.
pauciflora, *Blume, Mus. Bot. Lugd. Bat.* i. 370.—Ins. Molucc.
purpurea, *Hook. f. & Thoms. Fl. Ind.* i. 209.—Reg. Himal.
sumatrana, *Blume, Mus. Bot. Lugd. Bat.* i. 370.—Sumatra.
tomentosa, *Hook. f. Fl. Brit. Ind.* ii. 3.—Assam?
viridissima, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 304.—Ins. Andam.

SABICEA, Aubl. Pl. Gui. i. 192. t. 75, 76 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 72.
PAIVA, Vell. Fl. Flum. 104 (1825); iii. t. 16 (1827).
SCHWENKFELDA, Schreb. Gen. 123 (1789).
aspera, *Aubl. Pl. Gui.* i. 194. t. 76.—Guian.
aurea, Steud. Nom. ed. I. 712 = Coccocypselum aureum.
calycina, *Benth. in Hook. Niger Fl.* 399.—Afr. trop.
cana, *Hook. Ic. Pl.* t. 247.—Peruv.
capitellata, *Benth. in Hook. Niger Fl.* 398.—Afr. trop.
cauliflora, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 77.—Afr. trop.
cinerea, *Aubl. Pl. Gui.* i. 192. t. 75.—Guian.
crinita, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 228.—Madag.
diversifolia, *Pers. Syn.* i. 203.—Ins. Mascar.
edulis, Seem. in Hook. Kew Journ. iii. (1851) 266 = aspera.
eriantha, DC. Prod. iv. 439 = cinerea.
ferruginea, *Benth. in Hook. Niger Fl.* 397.—Afr. trop.
geantha, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 78.—Afr. trop.
glabrescens, Benth. in Hook. Journ. Bot. iii. (1841) 219 = aspera.
grisea, Cham. & Schlecht. in Linnaea, iv. (1829) 192 = cinerea.
guianensis, Baill. Hist. Pl. vii. 320 = Patima guianensis.
hirsuta, H. B. & K. Nov. Gen. et Sp. iii. 417 = aspera.
hirta, Sw. Prod. Veg. Ind. Occ. 46 = aspera.
macrophylla, Steud. Nom. ed. II. ii. 489, sphalm. = *Nonatelia macrophylla.*
Moralesii, Griseb. Cat. Pl. Cub. 124 = Lasianthus Moralesii.
Perrottetii, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 228 = (Uncariae sp).—Afr. trop.
pilosa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 76.—Afr. trop.
pumila, *Bartl. ex DC. Prod.* iv. 440.—Peruv.
purpurea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 228.—Bras.
segregata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 77.—Afr. trop.
setosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 228.—Madag.
tomentosa, *A. Rich. l. c.*—Bras.
triflora, *DC. Prod.* iv. 439.—Madag.
umbellata, *Pers. Syn.* i. 203.—Peruv.
velutina, Benth. in Hook. Journ. Bot. iii. (1841) 219 = aspera.
venosa, *Benth. in Hook. Niger Fl.* 399.—Afr. trop.
villosa, Roem. & Schult. Syst. v. 265 = aspera.
Vogelii, *Benth. in Hook. Niger Fl.* 398.—Afr. trop.

SABINA, Hall. in Rupp. Fl. Jen. 336 (1745) = **Juniperus**, Tourn. (Conif.).
alba, Antoine, Cupress. Gatt. 70 = J. virginiana.
aquatica, Antoine, l. c. 70 = Taxodium heterophyllum.
bacciformis, Antoine, l. c. 69 = J. phoenicea.
bermudiana, Antoine, l. c. 65. t. 87 = J. bermudiana.
Cabiancae, Antoine, l. c. 41. t. 54 = J. chinensis.
caesia, Antoine, l. c. 68 = J. caesia.
californica, Antoine, l. c. 52. t. 72 = J. virginiana.
chinensis, Antoine, l. c. 54. t. 75 = J. chinensis.
Corneyana, Antoine, l. c. 54. t. 74 = Cupressus torulosa.
cupressifolia, Antoine, ex C. Koch, Dendrol. ii. II. 125 = J. Sabina.
davurica, Antoine, Cupress. Gatt. 56. t. 77 = J. davurica.
dealbata, Antoine, l. c. 68 = J. communis.
dimorpha, Antoine, l. c. 70 = J. chinensis.
excelsa, Antoine, l. c. 45. t. 60 = J. excelsa.
Fischeri, Antoine, l. c. 53. t. 73 = J. Pseudo-sabina.
flaccida, Antoine, l. c. 37. t. 49 = J. flaccida.
foetidissima, Antoine, l. c. 49. tt. 67–71 = J. foetidissima.
fragrans, Antoine, l. c. 70 = J. virginiana.
gigantea, Antoine, l. c. 36. t. 48 = J. mexicana.
glauca, Antoine, l. c. 71 = J. virginiana.
gossainthanea, Antoine, l. c. 69 = J. virginiana.
Grisebachii, Antoine, l. c. 61 = J. foetidissima.
isophyllos, Antoine, l. c. 48. tt. 64–66 = J. excelsa.
lycia, Antoine, l. c. 44. t. 59 = J. phoenicea.
mexicana, Antoine, l. c. 38. t. 51 = J. mexicana.
occidentalis, Antoine, l. c. 64. tt. 85–86 = J. occidentalis.
officinalis, Garcke, Fl. Deutschl. ed. IV. 387 = J. Sabina.
Olivierii, Antoine, Cupress. Gatt. 70 = J. excelsa.
osteosperma, Antoine, l. c. 51.—Hab. ?
pachyphlaea, Antoine, l. c. 39 = J. pachyphlaea.
phoenicea, Antoine, l. c. 42. t. 57 = J. phoenicea.
plochyderma, Antoine, l. c. 40. t. 52 = J. pachyphlaea.
polycarpos, Antoine, l. c. 47. t. 63 = J. excelsa.
procera, Antoine, l. c. 36. t. 47 = J. procera.
prostrata, Antoine, l. c. 57. t. 79 = J. Sabina.
Pseudothurifera, Antoine, l. c. 60. t. 82 = J. thurifera ?
racemosa, Antoine, l. c. 71 = J. racemosa.
recurva, Antoine, l. c. 67. t. 88 = J. recurva.
religiosa, Antoine, l. c. 47. t. 61 = J. excelsa.
sphaerica, Antoine, l. c. 52. t. 72 = J. sphaerica.
squamata, Antoine, l. c. 66. t. 89 = J. recurva.
struthiacea, Antoine, l. c. 69 = J. chinensis.
tamariscifolia, Hort. ex C. Koch, Dendrol. ii. II. 125 = J. Sabina.
tetragona, Antoine, Cupress. Gatt. 40. t. 53 = J. tetragona.
thurifera, Antoine, l. c. 51 = J. thurifera ?
turbinata, Antoine, l. c. 41. t. 56 = J. phoenicea.
virginiana, Antoine, l. c. 61. tt. 83, 84 = J. virginiana.
vulgaris, Antoine, l. c. 58. t. 80 = J. Sabina.

SABINEA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 92. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
carinalis, *Griseb. Fl. Brit. W. Ind.* 183.—Ind. occ.
dubia, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 92.—Ins. Martinic.
florida, *DC. l. c.*—Ind. occ.

SABSAB, Adans. Fam. ii. 31 (1763) = **Paspalum**, Linn. (Gramin.).

SABULINA, Reichb. Fl. Germ. Excurs. 785 (1832) = **Arenaria**, Rupp. (Caryoph.).
austriaca, Reichb. l. c. 787 = A. austriaca.
bannatica, Reichb. l. c. = A. banatica.
biflora, Reichb. l. c. 790 = Stellaria biflora.
caespitosa, Reichb. l. c. 788 = A. verna.
capillacea, Reichb. l. c. 786 = A. grandiflora.
conferta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = A. tenuifolia.
falcata, Schur, Enum. Pl. Transs. 111 = A. setacea.
fastigiata, Reichb. Fl. Germ. Excurs. 786 = A. rostrata.
frutescens, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 136 = A. setacea.
Gerardi, Reichb. Fl. Germ. Excurs. 788 = A. verna.
glareosa, Schur, Enum. Pl. Transs. 111 = A. recurva.
graminifolia, Reichb. Fl. Germ. Excurs. 789 = A. clandestina, graminifolia.
hybrida, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = A. tenuifolia.

SABULINA :—
lanceolata, Reichb. Fl. Germ. Excurs. 790 = Arenaria lanceolata.
laricifolia, Reichb. l. c. 789 = A. laricifolia.
laxa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = A. grandiflora.
macrocarpa, Reichb. Fl. Germ. Excurs. 788 = A. laricifolia.
marginata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 13 = A. setacea.
mediterranea, Reichb. Ic. Fl. Germ. v. 27 = A. mucronata.
mucronata, Reichb. Fl. Germ. Excurs. 786 = A. mucronata.
polygonoides, Reichb. l. c. 790 = A. polygonoides.
Ponae, Reichb. l. c. = A. bavarica.
procera, Reichb. l. c. 787 = A. graminifolia.
ramosissima, Schur, Enum. Pl. Transs. 113 = A. ramosissima.
recurva, Reichb. Fl. Germ. Excurs. 788 = A. recurva.
rostrata, Reichb. l. c. 787 = A. rostrata.
setacea, Reichb. l. c. 786 = A. setacea.
sphagnoides, Reichb. l. c. 790 = A. polygonoides.
striata, Reichb. l. c. 789 = A. laricifolia.
stricta, Reichb. l. c. = A. stricta.
tenella, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 73 = A. verna.
tenuifolia, Reichb. Fl. Germ. Excurs. 785 = A. tenuifolia.
verna, Reichb. l. c. 788 = A. verna.
Villarsii, Reichb. l. c. 787 = A. Villarsii.
viscosa, Reichb. l. c. 786 = A. tenuifolia.

SACAGLOTTIS, G. Don, Gen. Syst. i. 676 (1831), sphalm. = **Sacoglottis**, Mart. (Humir.).

SACCARUM, Sanguin. in Atti Acc. Pont. Lincei, Ser. I. v. (1852) 501, 555 = **Saccharum**, Linn. (Gramin.).

SACCELLIUM, Humb. & Bonpl. Pl. Aequin. i. 47. t. 13 (1808). *BORAGINEAE*, Benth. & Hook. f. ii. 839.
lanceolatum, *Humb. & Bonpl. l. c.*—Peruv.

SACCHARIFERA, Stokes, Bot. Mat. Med. i. 131 (1812) = **Saccharum**, Linn. (Gramin.).
officinalis, Stokes, l. c. 132 = Saccharum officinarum.
spontanea, Stokes, l. c. = Saccharum spontaneum.

SACCHAROPHORUM, Neck. Elem. iii. 209 (1790) = seq.

SACCHARUM, Linn. Gen. ed. I. 18 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1125.
ERIOCHRYSIS, Beauv. Agrost. 8. t. 4. f. 11 (1812).
PHRAGMITES, Adans. Fam. ii. 34 (1763).
PLAZERIUM, Willd. ex Kunth, Enum. Pl. i. 474 (1838).
SACCHARIFERA, Stokes, Bot. Mat. Med. i. 131 (1812).
SACCHAROPHORUM, Neck. Elem. iii. 209 (1790).
acinaciforme, Spreng. Syst. i. 282 = Dimeria acinaciformis.
aegyptiacum, Willd. Enum. Hort. Berol. 82 = spontaneum.
alopecuroideum, Nutt. Gen. Am. i. 60 = Erianthus saccharoides.
Alopecurus, Nees, in Hook. Kew Journ. ii. (1850) 100 = Imperata exaltata.
angustifolium, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 107 = stenophyllum.
angustifolium, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 92 = Erianthus saccharoides.
Antillarum, Roem. & Schult. Syst. ii. 877 = Setaria setosa.
appressum, Kit. ex Reichb. Fl. Germ. Excurs. 34 = Erianthus Hostii.
argenteum, Brouss. Elench. 50 = Andropogon saccharoides.
arundinaceum, *Retz. Obs.* iv. 14.—Ind. or.; China.
asperum, Steud. Syn. Pl. Gram. 407 = Erianthus asper.
aureum, Spreng. Pugill. ii. 17 = Pollinia aurea.
Baldwini, Spreng. Syst. i. 282 = Erianthus strictus.
benghalense, Retz. Obs. v. 16 = arundinaceum.
bicorne, Griseb. in Goett. Abh. vii. (1857) 266 = Andropogon bicornis.
biflorum, Forsk. Fl. Aegypt. Arab. 16 = spontaneum.

SACCHARUM :—
brasilianum, Trin. in Mém. Acad. Péters. Sér. VI. ii. (1833) 311 = Erianthus asper.
brevibarbe, Pers. Syn. i. 103 = Erianthus saccharoides.
brunneum, Desv. Opusc. 65 (Panici sp.).—Ind. or.
caducum, Tausch, in Flora, xix. (1836) 527 = spontaneum.
canaliculatum, Roxb. Hort. Beng. 6 ; Fl. Ind. i. 246 = spontaneum.
capense, Steud. Syn. Pl. Gram. 408 = Erianthus capensis.
caudatum, G. F. W. Mey. Prim. Fl. Esseq. 68 = Imperata exaltata.
cayennense, *Benth. in Journ. Linn. Soc.* xix. (1881) 66.—Am. trop.
chinense, Nees, in Hook. & Arn. Bot. Beech. Voy. 241 = spontaneum.
chinense, Roxb. Hort. Beng. 6, nomen = officinarum.
ciliare, Andersson. in Oevers. Vet. Akad. Handl. Stockh. (1855) 155 = S. Sara.
confertum, J. & S. Presl, Rel. Haenk. i. 346 = Imperata exaltata.
conjugatum, Spreng. Syst. i. 283 (= *Andropogon conjugatus*).—Ind. or.
contortum, Nutt. Gen. Am. i. 60 ; Spreng. Syst. i. 282 = Erianthus saccharoides.
contractum, H. B. & K. Nov. Gen. et Sp. i. 182 = Imperata exaltata.
cylindricum, Lam. Encyc. i. 594 = Imperata arundinacea.
densum, Nees, in Hook. Kew Journ. ii. (1850) 100 = Miscanthus japonicus.
diandrum, Koen. ex Retz. Obs. v. 16 = Imperata arundinacea.
dissitiflorum, Edgew. in Journ. As. Soc. Beng. xvi. (1847) II. 1219 = Panicum Teneriffae.
distichophyllum, Steud. ex Jardin, in Mém. Soc. Sc. Nat. Cherb. v. (1857) 299, 325 = Miscanthus japonicus.
dubium, H. B. & K. Nov. Gen. et Sp. i. 183 = Imperata exaltata.
Ecklonii, Steud. Syn. Pl. Gram. 408 = Erianthus capensis.
edule, Hassk. in Flora, xxv. (1842) II. Beibl. 3 = officinarum.
erianthoides, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 308 = Erianthus saccharoides.
europaeum, Pers. Syn. i. 103 = Imperata arundinacea.
exaltatum, Roxb. Hort. Beng. [81]; Fl. Ind. i. 245 = arundinaceum ?
fasciculatum, Willd. ex Steud. Nom. ed. II. ii. 490 = Andropogon leucostachyus.
fastigiatum, Nees, ex Steud. Syn. Pl. Gram. 409 = Erianthus fastigiatus.
fatuum, Bechey, ex Steud. Nom. ed. II. ii. 490, nomen (Quid ?).—Hab. ?
filifolium, Steud. Syn. Pl. Gram. 409 = Erianthus filifolius.
floridulum, Labill. Sert. Austr. Caled. 13. t. 18 = Miscanthus japonicus.
fulvum, R. Br. Prod. 203 = Pollinia Cumingii.
fuscum, *Roxb. Hort. Beng.* 6; *Fl. Ind.* i. 236.—Ind. or.
giganteum, Pers. Syn. i. 103 = Erianthus saccharoides.
giganteum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 311 (Erianthi sp.).—Am. trop.
glaucum, Steud. Syn. Pl. Gram. 408 (Polliniae sp.).—Java.
grandiflorum, Walp. Ann. iii. 792 = Panicum insigne.
Griffithii, Boiss. Fl. Orient. v. 453 = S. Sara.
Griffithii, Munro, ex Aitch. in Journ. Linn. Soc. xix. (1882) 191 = praec. ?
hirsutum, Forsk. Fl. Aegypt. Arab. 16 = Rottboellia hirsuta.
holcoides, *Hack. in Mart. Fl. Bras.* ii. III. 254.—Bras.; Paraguay.
indum, Pers. Syn. i. 103 = Imperata arundinacea.
infirmum, Steud. in Lechl. Berb. Am. Austr. 56 = officinarum.
insulare, Brongn. in Duperr. Voy. Coq. Bot. 99 = spontaneum.
irritans, R. Br. Prod. 203 = Pollinia irritans.
jaculatorium, Lour. Fl. Cochinch. 53 = Imperata exaltata.
jamaicense, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 312 = Erianthus Ravennae.
japonicum, Thunb. in Trans. Linn. Soc. ii. (1794) 328 = Miscanthus sinensis, japonicus.

SACCHARUM:—
Klagha, Jungh. ex Steud. Syn. Pl. Gram. 405 = spontaneum.
Koenigii, Retz. Obs. v. 16 = Imperata arundinacea.
laguroides, Pourr. in Mém. Acad. Toul. iii. (1788) 326 = Imperata arundinacea.
longifolium, *Munro, ex Benth. in Journ. Linn. Soc.* xix. (1881) 66.—Ind. or.
longisetum, Walp. Ann. iii. 793 (= *Tricholaena longiseta*).—Abyss.
macilentum, Chauv. ex Steud. Syn. Pl. Gram. 406 = Imperata exaltata.
maximum, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. (1838) 92 = Erianthus maximus.
mexicanum, Brouss. ex Hornem. Hort. Hafn. i. 75 = Imperata Hookeri.
monandrum, *Roland. in Act. Lit. Univ. Hafn.* (1777) 285.—Jamaic. ; Guian.
monandrum, Sw. ex Hack. in Mart. Fl. Bras. ii. III. 252, in obs. = Imperata exaltata.
Munja, Roxb. Hort. Beng. 6 ; Fl. Ind. i. 246 = S. Sara ?
Narenga, *Wall. Cat.* n. 8856.—Ind. or. ; China.
Nareya, Benth. in Journ. Linn. Soc. xix. (1881) 66, sphalm. = praec.
negrosense, Steud. Syn. Pl. Gram. 407 = Imperata exaltata.
obscurum, Steud. l. c. (Miscanthi sp.).—Japon.
obscurum, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 313 = Erianthus Hostii ?
occidentale, Sw. Adnot. Bot. 17 = monandrum.
officinale, Salisb. Prod. 17 = seq.
officinarum, *Linn. Sp. Pl.* 54.—Reg. Calid. cult.
Palisoti, Tausch, in Flora, xix. (1836) 527 = spontaneum.
pallidum, *Benth. in Journ. Linn. Soc.* xix. (1881) 66. —Afr. austr.
paniceum, Lam. Encyc. i. 595. t. 40. f. 1 = Pogonatherum saccharoideum.
panicosum, Griff. Notul. iii. 73 = Pogonatherum saccharoideum.
pappiferum, Lam. Illustr. i. 155 = Pappophorum alopecuroideum.
pedicellare, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 310 (Erianthi sp.).—Ins. Nukahiva.
polydactylum, Thunb. Fl. Jap. 42 = Miscanthus japonicus, sinensis.
polydactylum, Willd. ex Steud. Nom. ed. II. ii. 490 = Panicum saccharoides.
polystachyum, Siebold, ex Kunth, Enum. Pl. i. 124 = Panicum lanatum.
polystachyum, Sw. Prod. Veg. Ind. Occ. 21 = Panicum saccharoides.
praegrande, Steud. Syn. Pl. Gram. 408 = Miscanthus japonicus.
procerum, Roxb. Hort. Beng. 6 ; Fl. Ind. i. 243 = arundinaceum.
propinquum, Steud. Syn. Pl. Gram. 406 = spontaneum.
punctatum, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 46 = spontaneum.
rarum, Poir. Encyc. Suppl. ii. 70 = Perotis rara.
Ravennae, Bieb. Fl. Taur. Cauc. iii. 51 = Imperata arundinacea.
Ravennae, Murr. Syst. ed. XIII. 88 = Erianthus Ravennae.
repens, Willd. Sp. Pl. i. 322 (Tricholaenae sp.).—Afr. trop.
reptans, Lam. Illustr. i. 155 = Panicum repens ?
roseum, Reinw. ex Miq. Fl. Ind. Bat. iii. 518 = Miscanthus sinensis.
rufipilum, Steud. Syn. Pl. Gram. 409 = Erianthus fulvus.
rufum, Steud. l. c. = Pollinia argentea.
sagittatum, Aubl. Pl. Gui. i. 50 = Gynerium saccharoides.
Sapé, A. St. Hil. Voy. Distr. Diam. i. 368 = Imperata brasiliensis.
Sara, *Roxb. Hort. Beng.* 6 ; *Fl. Ind.* i. 244.—Ind. or. ; Afghan.
semidecumbens, Roxb. l. c. 236 = spontaneum.
sinense, Roxb. Fl. Ind. ed. Carey, i. 244 = officinarum.
Sisca, Cav. Ic. iii. 47 = Imperata arundinacea.
Sorghum, Steud. Syn. Pl. Gram. 408 = Erianthus Sorghum.
speciosissimum, Tausch, in Flora, xix. (1836) 527 = spontaneum.
sphacelatum, Walp. Ann. iii. 793 (= *Tricholaena sphacelata*).—Afr. trop.

SACCHARUM:—
spicatum, Burm. ex Kunth, Enum. Pl. i. 477 = Imperata arundinacea.
spicatum, Linn. Sp. Pl. 54 = Perotis latifolia.
spicatum, Thunb. Fl. Jap. 42 ; J. & C. Presl, Rel. Haenk. i. 346 = Imperata arundinacea.
spontaneum, *Linn. Mant.* ii. 183.—Geront. trop.
stenophyllum, *Buese, in De Vriese, Pl. Ind. Bat. Or.* 107.—Ins. Timor.
strictum, Nutt. Gen. Am. i. 60 = Erianthus strictus.
strictum, Spreng. Pugill. ii. 16 = Erianthus Hostii.
Teneriffae, Linn. f. Suppl. 106 = Panicum Teneriffae.
ternatum, Spreng. Syst. i, 283 = Andropogon ternatus.
Thunbergii, Retz. Obs. v. 17 = Imperata arundinacea.
tinctorium, Steud. Syn. Pl. Gram. 469 (Mischanthi sp.). —Japon.
tridentatum, Spreng. Syst. i. 283 = Apocopis tridentatus.
tripsacoides, Beauv. ex Roem. & Schultz. Syst. ii. 783 = Hemarthria compressa.
tristachyum, Steud. Syn. Pl. Gram. 408 = Pollinia argentea.
versicolor, Steud. l. c. 409 = Erianthus versicolor.
villosum, Steud. l. c. 408 (= Erianthi sp.).—Bras.
violaceum, Tussac, Fl. Antill. i. 160. t. 25 = officinarum.
Warmingianum, *Hack. in Mart. Fl. Bras.* ii. III. 254. —Bras.

SACCIDIUM, Lindl. Gen. et Sp. Orch. 301 (1835) = **Holothrix**, Rich. (Orchid.).
pilosum, Lindl. l. c.—H. pilosa.

SACCILABIUM, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 294 = **Nepeta**, Riv. (Labiat.).
basilicoides, Rottb. l. c. = N. basilicoides.
viscosa, Rottb. l. c. = N. viscosa.

SACCOCALYX, Coss. & Dur. in Ann. Sc. Nat. Sér. III. xx. (1853) 80. t. 5 = **Satureia**, Linn. (Labiat.).
satureioides, Coss. & Dur. l. c. 81.—Alger.

SACCOCALYX, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = **Astragalus**, Tourn. (Legumin.).
calycinus, Stev. l. c. = A. calycinus.
coluteoides, Stev. l. c. = A. coluteoides.
follicularis, Stev. l. c. = A. follicularis.
halicacabus, Stev. l. c. = A. halicacabus.
laguroides, Stev. l. c. = A. laguroides.
lineatus, Stev. l. c. = A. lineatus.
lupulinus, Stev. l. c. = A. lupulinus.
tumidus, Stev. l. c. = A. Clusii.
Vulnerariae, Stev. l. c. = A. Vulnerariae.

SACCOCHILUS, Blume, Fl. Jav. Praef. p. viii (1828) = **Saccolabium**, Blume (Orchid.).

SACCOGLOTTIS, Endl. Gen. 1040 (1840) = **Sacoglottis**, Mart. (Humir.).

SACCOLABIUM, Blume, Bijdr. 292 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 578.
CERATOCHILUS, Blume, Bijdr. 358. t. 25 (1825).
GASTROCHILUS, D. Don, Prod. Fl. Nep. 32 (1825).
MICROPERA, Dalz. in Hook. Kew Journ. iii. (1851) 282.
OECEOCLADES, Lindl. Bot. Reg. sub t. 1522 (1832) ; Gen. et Sp. Orch. 235 (1833).
OMOEA, Blume, Bijdr. 359 (1825).
ROBIQUETIA, Gaudich. in Freyc. Voy. Bot. 426. t. 34 (1826).
SACCOCHILUS, Blume, Fl. Jav. Praef. p. viii (1828).
acuminatum, Thw. Enum. Pl. Zeyl. 304 = Sarcanthus peninsularis.
acutifolium, *Lindl. Gen. et Sp. Orch.* 223.—Reg. Himal.
albo-lineatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 20.—Siam.
ampullaceum, *Lindl. in Wall. Cat.* n. 7307.—Reg. Himal. ; Burma.
aphyllum, Lindl. Gen. et Sp. Orch. 223 = Angraecum aphyllum.
ascendens, *Lindl. l. c.*—Ins. Molucc.
bellinum, *Reichb. f. in Gard. Chron.* (1884) I. 174.—Burma.
Berkeleyi, *Reichb. f. l. c.* (1883) I. 814.—Burma ?
Bertholdi, *Reichb. f. in Seem. Fl. Vit.* 297.—Ins. Fiji.
bifidum, *Lindl. Bot. Reg.* (1838) *Misc.* 3.—Ins. Philipp.

SACCOLABIUM:—
bigibbum, *Reichb. f. ex Hook. f. Bot. Mag.* t. 5767.—Burma ; Malaya.
bipunctatum, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 145 (Cleisostomatis sp.).—Burma.
Blumei, Lindl. Bot. Reg. (1841) Misc. 55 = Rhynchostylis retusa.
borneense, *Reichb. f. in Gard. Chron.* (1881) I. 563.—Borneo.
brevifolium, *Lindl. Gen. et Sp. Orch.* 225.—Zeylan.
buccosum, *Reichb. f. in Gard. Chron.* (1871) 938.—Reg. Himal. ; Burma.
calcaratum, F. Muell. Fragm. i. 192 = Cleiosostoma tridentatum.
calceolare, *Lindl. in Wall. Cat.* n. 7302.—Reg. Himal. ; Malaya.
calceolare, Paxt. Mag. Bot. vi. (1839) 97 = intermedium.
calopterum, *Reichb. f. in Gard. Chron.* (1882) II. 520. —N. Guin.
carinatum, Griff. Notul. iii. 354 = papillosum.
chionanthum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 35.—Java.
clavatum, *Lindl. Gen. et Sp. Orch.* 223.—Ind. or.
coeleste, *Reichb. f. in Gard. Chron.* (1885) I. 692.—Hab. ?
compressum, *Lindl. Bot. Reg.* (1840) *Misc.* 9.—Ins. Philipp.
constrictum, *Reichb. f. Otia Bot. Hamb.* 52.—Ins. Fiji.
coriaceum, Lindl. Gen. et Sp. Orch. 224 = Aërides coriaceum.
curvifolium, *Lindl. l. c.* 222.—Burma ; Java.
dasypogon, Lindl. l. c. = calceolare ?
densiflorum, Lindl. in Wall. Cat. n. 7311 = Cleisostoma spicatum.
dentatum, Reichb. f. in Walp. Am. vi. 883, sphalm. = seq.
denticulatum, Paxt. Mag. Bot. vii. (1840) 145 = acutifolium.
discolor, Reichb. f. Otia Bot. Hamb. 42 = S. Pumilio.
distichum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 36. —Reg. Himal.
dives, *Reichb. f. in Gard. Chron.* (1875) II. 180.—Hab. ?
fasciculatum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 238.—N. Guin.
filiforme, *Lindl. in Journ. Linn. Soc.* iii. (1859) 36.—Ind. or.
flexum, *Reichb. f. in Gard. Chron.* (1882) I. 520.—N. Guin.
flexuosum, Lindl. in Journ. Linn. Soc. iii. (1859) 36 (Cleisostomatis sp.).—Reg. Himal. ; Burma.
fragrans, *Par. & Reichb. f. in Journ. Bot.* xii. (1874) 197.—Burma.
galeatum, Gardn. ex Thw. Enum. Pl. Zeyl. 305 = Cleisostoma maculosum.
gemmatum, *Lindl. Bot. Reg.* (1838) *Misc.* 50.—Reg. Himal.
giganteum, Lindl. in Wall. Cat. n. 7306 = Vanda densiflora.
gracile, *Lindl. Gen. et Sp. Orch.* 225.—Zeylan.
Graeffei, *Reichb. f. in Gard. Chron.* (1881) II. 716.—Ins. Fiji.
Griffithii, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 145 = Microsaccus javensis.
gurwalicum, Lindl. in Journ. Linn. Soc. iii. (1859) 32 = Rhynchostylis retusa.
guttatum, Lindl. in Wall. Cat. n. 7308 = Rynchostylis retusa.
Hanissonii, Boxall, ex Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 241, sphalm. = seq. ?
Harrisonianum, *Hook. Bot. Mag.* t. 5433.—Malaya.
Heathi, Hort. ex Gard. Chron. (1885) II. 369 = Rhynchostylis retusa.
Hendersonianum, *Reichb. f. l. c.* (1875) II. 356.—Hab. ?
Hillii, *F. Muell. Fragm.* i. 192.—Austral.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 537.—Ins. Comor.
Huttoni, *Hook. f. Bot. Mag.* t. 5681.—Java.
intermedium, *Griff. ex Lindl. in Journ. Linn. Soc.* iii. (1859) 33.—Reg. Himal.
Jerdonianum, *Reichb. f. in Walp. Ann.* vi. 886.—Ind. or.
lineare, Lindl. in Wall. Cat. n. 7312 (Aëridis sp.).—Ind. or.
lineolatum, Thw. Enum. Pl. Zeyl. 304 = ochraceum.
littorale, *Reichb. f. in Gard. Chron.* (1881) II. 198.—Malacca.

SACCOLABIUM:—
Macphersonii, F. Muell. Fragm. vii. 96 = Cleisostoma Macphersoni.
macrostachyum, Lindl. Sert. Orch. t. 47 = Rhynchostylis retusa.
micranthum, *Lindl. in Wall. Cat.* n. 7300.—Reg. Himal.; Burma.
Mimus, *Reichb. f. in Gard. Chron.* (1878) i. 266.—Ins. Pacif.
miniatum, Hook. Bot. Mag. t. 5326 = curvifolium.
miniatum, *Lindl. Bot. Reg.* (1847) sub t. 26.—Java.
niveum, *Lindl. Gen. et Sp. Orch.* 224.—Zeylan.
obliquum, *Lindl. in Wall. Cat.* n. 7304.—Burma.
ochraceum, *Lindl. Bot. Reg.* (1842) *Misc.* 2.—Ind. or.
pachyglossum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 34.—Borneo.
pallens, *Cathcart, ex Lindl. l. c.* 35.—Reg. Himal.
paniculatum, Wight, Ic. t. 1676 = *lineare*.
papillosum, Dalz. & Gibs. Bomb. Fl. 264 (= *Acampe Wightiana*).—Ind. or.
papillosum, *Lindl. in Wall. Cat.* n. 7305; *et Bot. Reg.* t. 1552.—Ind. or.
papillosum, Wight, Ic. t. 1672 (= *Acampe congesta*).—Ind. or.
parvulum, Lindl. in Journ. Linn. Soc. iii. (1859) 36 = buccosum.
praemorsum, Lindl. Gen. et Sp. Orch. 22 = Rhynchostylis retusa.
Pumilio, *Reichb. f. in Walp. Ann.* vi. 886.—Reg. Himal.
pusillum, *Blume, Bijdr.* 292. t. 50.—Java.
pusillum, Lindl. in Journ. Linn. Soc. iii. (1859) 36 = S. Pumilio.
quinquefidum, *Lindl. in Hook. Journ. Bot.* ii. (1843) 238.—N. Hibern.
racemiferum, Lindl. Gen. et Sp. Orch. 224 = Sarcanthus pallidus.
radicosum, *A. Rich. Tent. Fl. Abyss.* ii. 285.—Abyss.
ramosum, Lindl. Gen. et Sp. Orch. 224 (Cleisostomatis sp.).—Reg. Himal.; Burma.
ramulosum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 34—Java.
reflexum, Lindl. in Wall. Cat. n. 7309 = Renanthera elongata.
repens, *Lindl. ex Steud. Nom.* ed. II. ii. 491, nomen.—Ind. or.
retusum, Voigt, Hort. Suburb. Calc. 630 = Rhynchostylis retusa.
Rheedii, Wight, Ic. v. 19 = Rhynchostylis retusa.
ringens, Lindl. in Wall. Cat. n. 7313 = Aërides radicosum.
roseum, *Lindl. Gen. et Sp. Orch.* 225.—Zeylan.
roseum, Lindl. in Journ. Linn. Soc. iii. (1859) 36 = filiforme.
rubrum, Lindl. in Wall. Cat. n. 7310 = curvifolium.
rubrum, Wight, Ic. t. 1673 = Aërides radicosum.
speciosum, Wight, l. c. t. 1674–75 = Aërides maculosum.
spicatum, Lindl. Gen. et Sp. Orch. 221 = Rhynchostylis retusa.
striatum, Lindl. l. c. 224 = Angræcum striatum.
tenerum, Lindl. in Journ. Linn. Soc. iii. (1859) 36 (Cleisostomatis sp.).—Ind. or.
trichromum, Reichb. f. in Hamb. Gartenz. xv. (1859) 51 = pallens.
undulatum, Lindl. in Wall. Cat. n. 7301 = Cleisostoma undulatum.
violaceum, Reichb. f. in Bonplandia, ii. (1854) 93 = Rhynchostylis violacea.
virescens, Gardn. ex Lindl. in Journ. Linn. Soc. iii. (1859) 35 = brevifolium.
viridiflorum, *Lindl. l. c.* 36.—Ind. or.
Walkerianum, Reichb. f. in Walp. Ann. vi. 887 = roseum.
Wightianum, Lindl. in Wall. Cat. n. 7303, partim = Aërides radicosum.
Witteanum, *Reichb. f. in Gard. Chron.* (1883) ii. 618.—Java.

SACCONIA, Endl. Gen. 541 (1838) = **Chione**, DC. (Rubiac.).
glabra, Endl. ex Walp. Rep. ii. 487 = C. glabra.

SACCOPETALUM, Benn. Pl. Jav. Rar. 165. t. 35 (1838). *ANONACEAE*, Benth. & Hook. f. i. 28.
Bidwilli, *Benth. Fl. Austral.* i. 53.—Austral.

SACCOPETALUM:—
Brahei, *F. Muell. Fragm.* viii. 159.—Austral.
carnosum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 418.—Java.
Horsfieldii, *Benn. Pl. Jav. Rar.* 165. t. 35.—Java.
longiflorum, *Hook. f. & Thoms. Fl. Ind.* 151.—Ind. or.
sclerocarpum, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 88.—Burma.
tomentosum, *Hook. f. & Thoms. Fl. Ind.* i. 152.—Ind. or.

SACCOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 117 = **Alloplectus**, Mart. (Gesner.).
strigosus, Oerst. l. c. 118 = A. strigosus.

SACCOSTOMA, Wall. ex Voigt, Hort. Suburb. Calc. 449 (1845). *LABIATAE* (Quid?).
urticaefolium, Wall. ex Voigt, l. c. nomen.—Ind. or.

SACCULARIA, Kellogg, in Proc. Calif. Acad (1863) 17 = **Antirrhinum**, Tourn. (Scrophul.). ii.
Veatchii, Kellogg, l. c. = A. junceum.

SACELLIUM, Spreng. Syst. iii. 851 (1826) = **Saccellium**, Humb. & Bonpl. (Borag.).

SACHARUM, Scop. Introd. 75 (1777) = **Saccharum**, Linn. (Gramin.).

SACHSIA, Griseb. Cat. Pl. Cub. 150 (1866). *COMPOSITAE*, Benth. & Hook. f. ii. 291.
divaricata, *Griseb, l. c.* 151.—Cuba.
polycephala, *Griseb. l. c.*—Cuba.
tricephala, *Griseb. l. c.*—Cuba.

SACODON, Rafin. Fl. Tellur. iv. 45 (1836) = **Cypripedium**, Linn. (Orchid.).
macranthon, Rafin. l. c. = C. macranthon.
ventricosum, Rafin. l. c. = C. macranthon.

SACOGLOTTIS, Mart. Nov. Gen. et Sp. ii. 146 (1827). *HUMIRIACEAE*, Benth. & Hook. f. i 247.
amazonica, Benth. in Hook. Kew Journ. v. (1853) 104 = guianensis.
amazonica, *Mart. Nov. Gen. et Sp.* ii. 146.—Bras.
cuspidata, *Urb. in Mart. Fl. Bras.* xii. ii. 444.—Bras.
densiflora, *Urb. l. c.* 445.—Bras.
dentata, *Urb. l. c.* 444.—Bras.
dichotoma, *Urb. l. c.* 446.—Bras.
gabonensis, Urb. l. c. 449 = Aubrya gabonensis.
guianensis, *Benth. in Hook. Kew Journ.* v. (1853) 103.—Guian.
macrophylla, *Urb. in Mart. Fl. Bras.* xii. ii. 446.—Bras.
oblongifolia, *Urb. l. c.* 447.—Bras.
obovata, *Urb. l. c.* 443.—Bras.
subcrenata, Urb. in Sitzb. Ges. Naturf. Berl. (1878) 5 = Humiria subcrenata.

SACOILA, Rafin. Fl. Tellur. ii. 86 (1836) = **Spiranthes**, Rich. (Orchid.).
lurida, Rafin. l. c. = Sp. orchioides.

SACRANTHUS, Endl. Ench. 331 (1841), sphalm. = SAIRANTHUS, G. Don = **Nicotiana**, Linn. (Solan.).

SADRUM, Soland. ex Baill. in DC. Prod. xvii. 18 (1873) = **Pyrenacantha**, Wight (Olacin.).
volubile, Soland. ex Baill. l. c. = P. scandens.

SADYMIA, Griseb. Fl. Brit. W. Ind. 25 (1859) = **Samyda**, Linn.
villosa, Griseb. l. c. = Samyda villosa.

SAELANTHUS, Forsk. Fl. Aegypt. Arab. 33 (1775) = **Vitis**, Linn. (Ampel.).
digitatus, Forsk. l. c. 35 (= *Cissus digitata*).—Arab.
glandulosus, Forsk. l. c. 34 (= *Cissus glandulosa*).—Arab.
quadragonus, Forsk. l. c. 33 = V. quadrangularis.
rotundifolius, Forsk. l. c. 35 (= *Cissus rotundifolia*).—Arab.
ternatus, Forsk. l. c. (= *Cissus ternata*).—Arab.

SAFRAN, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 473 = **Crocus**, Tourn. (Irid.).
officinarum, Medic. l. c. = C. sativus.

SAGERAEA, Dalz. in Hook. Kew Journ. iii. (1851) 207 = **Bocagea**, A. St. Hil. (Anonac.).
cauliflora, Scheff. in Ann. Jard. Buitenz. ii. (1885) 5.—Sumatra.
Dalzellii, Bedd. Ic. Pl. Ind. Or. i. 9. t. 42 = B. Dalzellii.
elliptica, Hook. f. & Thoms. Fl. Ind. i. 93 = B. elliptica.
Hookeri, Pierre, Fl. For. Cochin. t. 15 = B. elliptica.
lanceolata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 10.—Borneo.
laurina, Dalz. in Hook. Kew Journ. iii. (1851) 207 = B. Dalzellii.
Thwaitesii, Hook. f. & Thoms. Fl. Ind. i. 93 = B. Thwaitesii.

SAGERETIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 359. t. 13. f. 2. *RHAMNEAE*, Benth. & Hook. f. i. 379.
AFARCA, Rafin. Sylva Tellur. 30 (1838).
affinis, G. Don, Gen. Syst. ii. 29 = Rhamnus affinis.
Blumii, G. Don, l. c. 28 = oppositifolia.
Brandrethiana, *Aitch. in Journ. Linn. Soc.* viii. (1865) 62.—Oriens; Ind. bor. occ.
corymbosa, *G. Don, Gen. Syst.* ii. 29.—Ind. occ.
costata, *Miq. Fl. Ind. Bat.* i. 1. 645.—Java.
decussata, G. Don, Gen. Syst. ii. 29 = senticosa?
divergens, *Steud. Nom.* ed. II. ii. 491.—Chili.
elegans, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 359.—Mexic.; Peruv.
filiformis, G. Don, Gen. Syst. ii. 29 = oppositifolia.
guayaquilensis, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 360.—Ecuador.
hamosa, *Brongn. l. c.* 360.—Ind. or.; China.
lanceolata, G. Don, Gen. Syst. ii. 29 = Rhamnus lanceolata.
leprosa, G. Don, l. c. = Rhamnus leprosa.
Michauxii, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 360.—Am. bor.
oppositifolia, *Brongn. l. c.* 360.—Ind. or.; Malaya.
parviflora, G. Don, Gen. Syst. ii. 29 = oppositifolia.
ramosa, Royle, Illustr. Bot. Himal. 169 = hamosa?
riparia, *Steud. Nom.* ed. II. ii. 491.—Chili.
rugosa, *Hance, in Journ. Bot.* xvi. (1878) 9.—China.
senticosa, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 360.—Peruv.
spicata, *Brongn. l. c.*—Peruv.
splendens, G. Don. Gen. Syst. ii. 29 = Rhamnus splendens.
theezans, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 360.—Ind. bor. occ.; Burma; China.
trigyna, G. Don, Gen. Syst. ii. 29 = oppositifolia.
trinervis, *Gill. ex Hook. Bot. Misc.* iii. (1833) 172.—Chili.
Wrightii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 358.—Texas.

SAGINA, Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
PHALOË, Dum. Fl. Belg. 110 (1827).
SPERGELLA, Reichb. in Moessl. Handb. ed. II. i. p. lxv. (1827).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 47.—Abyss.
acicularis, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 756 = subulata, *Wimm.*
ambigua, Lloyd, Fl. Ouest Fr. ed. I. 74 = ciliata.
ammannioides, Heyne, ex Wall. Cat. n. 7504 = Bergia ammanioides.
apetala, *Linn. Mant.* ii. 559.—Europ.
breviflora, Gilib. Fl. Lituan. ii. 148 = procumbens.
bryoides, *Froel. ex Reichb. Fl. Germ. Excurs.* 793.—Europ.
caespitosa, Lange, Pl. Groenl. 133; Consp. Fl. Groenl. i. 22 = nivalis.
cerastioides, Sm. Engl. Bot. t. 166; et in Trans. Linn. Soc. ii. (1794) 343 = Cerastium tetrandrum.
chilensis, *C. Gay, Fl. Chil.* i. 282.—Chili.
ciliaris, Fenzl, Verbr. Alsin. tab. ad p. 57 = Spergula ciliaris.
ciliata, *Fries, in Liljeb. Sv. Fl.* ed. III. 713.—Europ.
corsica, *Jord. Obs. Pl. Crit.* vii. (1849) 15.—Corsica.

SAGINA:—
crassicaulis, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 191.—Calif.
crassifolia, Urv. in Mém. Soc. Linn. Par. iv. (1826) 617 = Colobanthus crassifolius.
debilis, Jord. Obs. Pl. Crit. iii. (1846) 50 = maritima.
decandra, Reichb. Ic. Pl. Crit. vii. 27. t. 875 = Arenaria verna, sedoides.
decumbens, *Torr. & Gray, Fl. N. Am.* i. 177.—Am. bor.
densa, Jord. Obs. Pl. Crit. iii. (1846) 49 = maritima.
depressa, Schultz, Prod. Fl. Starg. Suppl. i. 10 = ciliata
dichotoma, *Heuff. in Flora*, xxxvi. (1853) 626.—Hungar.
Elliottii, Fenzl, ex A. Gray, Man. Bot. N. U. St. ed. II. 61 = decumbens.
erecta, Lam. Encyc. vi. 390, partim = apetala.
erecta, Linn. Sp. Pl. 128 = Cerastium quaternellum.
erecta, O. F. Muell. in Fl. Dan. t. 845 = maritima.
fasciculata, *Boiss. Diagn.* Ser. II. i. 83.—Sibir.
fasciculata, *Poir. Encyc.* vi. 390.—Afr. bor.
filicaulis, Jord. Obs. Pl. Crit. vii. (1849) 16 = apetala.
filiformis, Pourr. ex Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 27 = maritima.
fontinalis, Short & Peter, in Transylv. Journ. Med. (1836) 600 = Stellaria crassifolia.
glabra, *Fenzl, Verbr. Alsin.* tab. ad p. 57.—Europ.
glandulifera, Nym. Consp. 1010 (Index) = nodosa.
glandulosa, Bess. ex Nym. l. c. 120 = nodosa.
graminifolia, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 292.—Bolivia.
Haenkeana, *C. Gay, Fl. Chil.* i. 284.—Chili.
Hochstetteri, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xxi. (1871) 34.—Ins. St. Paul (Mar. Ind. austr.).
humifusa, *Fenzl, ex Rohrb. in Mart. Fl. Bras.* xiv. II. 262.—Bras.
intermedia, Fenzl, in Ledeb. Fl. Ross. i. 339 = nivalis.
kevandensis, *Boiss. & Reut. Pugill. Pl. Nov.* 22.—Hispan.
Lamyi, *F. Schultz, Arch. Fl. Fr.* 174.—Gallia.
laricina, Fenzl, Verbr. Alsin. tab. ad p. 46 = Spergula laricina.
Linnaei, A. Gray, in Proc. Am. Acad. viii. (1873) 378 = occidentalis.
Linnaei, *Presl, Rel. Haenk.* ii. 14 (Linnei).—Reg. bor. et arct.
Loscosii, Boiss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 604 = sabuletorum.
macrocarpa, Losc. & Pard. Ser. Inconf. ed. II. 64 = sabuletorum.
macrocarpa, Maly, Enum. 293 = Linnaei.
magellanica, Willd. ex Phil. f. Cat. Pl. Vasc. Chil. 25 = Colobanthus quitensis.
maritima, *D. Don, in Gen. Engl. Bot.* t. 2195.—Europ.
maxima, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 382 = Linnaei.
media ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 71.—Europ.
melitensis, Gulia, ex Duthie, in Journ. Bot. xiii. (1875) 37 = apetala.
micrantha ×, Bor.—Cf. Bull. Soc. Bot. Fr. xxii. (1875) Bibl. 187.—Europ.
muscoides, Koch, Syn. Fl. Germ. ed. I. 108 = bryoides.
muscosa, Jord. Pugill. Pl. Nov. 32 = procumbens.
nevadensis, *Boiss. Pugill. Pl. Nov.* 22.—Hispan.
nivalis, *Fries, Nov. Mant.* iii. 31.—Reg. bor. et arct.
nodosa, *Fenzl, Verbr. Alsin* .tab. ad p. 18.—Reg. bor. temp.
occidentalis, *S. Wats. in Proc. Am. Acad.* x. (1875) 344.—Am. bor. occ.
octandra, Ziz, ex Mert. & Koch, Deutschl. Fl. i. 864 = Cerastium quaternellum.
oxysepala, *Boiss. Fl. Orient.* i. 663.—Armenia.
patula, Jord. Obs. Pl. Crit. i. (1846) 23 = ciliata.
pilifera, Fenzl, Verbr. Alsin. tab. ad p. 57 = glabra.
pilifera, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 756 = subulata, *Wimm.*
procumbens, Boland. Cat. 6 = occidentalis.
procumbens, *Hook. Bot. Misc.* iii. (1833) 147 an seq.?).—Chili.
procumbens, *Linn. Sp. Pl.* 128.—Reg. bor. temp.
procumbens, Pursh, Fl. Am. Sept. i. 219 = decumbens.
procumbens, Thunb. Fl. Jap. 80 = Linnaei.
quitensis, H. B. & K. Nov. Gen. et Sp. vi. 19 = Colobanthus quitensis.
repens, Burnat, ex Gremli, Excurs. Schw. ed. IV. 102 = glabra.
Reuteri, *Boiss. Diagn.* Ser. II. i. 82.—Hispan.
Revelieri, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 11 = subulata.

SAGINA:—
Rodriguezii, Willk. in Oestr. Bot. Zeitschr. xxv. (1875) 113 = maritima.
sabuletorum, *Lange, Desc. et Ic. Ill.* 3.—Hispan.
saxatilis, Wimm. Fl. Schles. (1841) 75 = Linnaei.
setigera, *Bisch. Del. Sem. Hort. Heidelb.* (1852) 8. —Hab.?
sinensis, Hance, in Journ. Bot. vi. (1868) 46 = Linnaei.
Spergella, Fenzl, Verbr. Alsin. tab. ad p. 18 = Linnaei.
stricta, Fries, Novit. Fl. Suec. iii. 122 = maritima.
subulata, *Presl, Fl. Sic.* 158.—Europ.
subulata, Torr. & Gray, Fl. N. Am. i. 178 = decumbens.
subulata, Urv. in Mém. Soc. Linn. Par. iv. (1826) 617 = Colobanthus subulatus.
subulata, *Wimm. Fl. Schles.* (1841) 76.—Hispan.
urbica, *Phil. in Linnaea*, xxvjii. (1856) 613.—Chili.
urceolata, Viv. Fl. Lib. Spec. 67; Fl. Cors. 3 = maritima.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864-65) 22.—Chili.
virginica, Linn. Sp. Pl. 128 = Bartonia tenella.
viscida, Steud. Nom. ed. II. ii. 491 = procumbens?

SAGITTA, Adans. Fam. ii. 459 (1763) = **Sagittaria**, Rupp. (Alismac.).
maior, Scop. Fl. Carn. ed II. ii. 239 = Sagittaria sagittifolia.

SAGITTARIA, Rupp. ex Linn. Syst. ed. I (1735).
ALISMACEAE, Benth. & Hook. f. iii. 1006.
DIPHOREA, Rafin. Neogenyt. 3 (1825).
DREPACHENIA, Rafin. l. c. 3 (1825).
LOPHIOCARPUS, Miq. Ill. Arch. Ind. Fl. 50 (1871).
SAGITTA, Adans. Fam. ii. 459 (1763).
acuminata, Sm. in Rees, Cycl. xxxi. n. 4 = sagittifolia.
acutifolia, Linn. ex Griseb. Cat. Pl. Cub. 218 = intermedia.
acutifolia, Linn. f. Suppl. 419 = pugioniformis.
acutifolia, Pursh, Fl. Am. Sept. ii. 397 = graminea.
affinis, *Seub. in Mart. Fl. Bras.* iii. 1. 111.—Bras.
alismaefolia, Phil. ex M. Mich. in DC. Monog. Phan. iii. 76 = montevidensis.
alpina, Willd. Sp. Pl. iv. 410 = sagittifolia.
andina, Phil. in Linnaea, xxix. (1857–58) 45 = montevidensis.
angustifolia, Lindl. Bot. Reg. t. 1141 = lancifolia.
aquatica, Lam. Fl. Fr. ii. 197 = sagittifolia.
Blumei, Kunth, Enum. Pl. iii. 158 = guayanensis.
bracteata, Willd. ex Seub. in Mart. Fl. Bras. iii. 1. 106 = guyannensis.
brasiliensis, Mart. ex M. Mich. in DC. Monog. Phan. iii. 57 = Echinodorus floribundus.
bulbosa, Donn, Hort. Cantab. ed. VI. 246; et ex Sims, Bot. Mag. sub t. 1632 = heterophylla.
calycina, *Engelm. ex Torr. in Bot. Mex. Bound.* 212. —Am. bor.
cerulea, Rafin. Fl. Ludov. 18 (Quid?).—Am. bor.
chilensis, Cham. & Schlecht. in Linnaea, ii. (1827) 155 = montevidensis.
chinensis, *Sims, Bot. Mag.* t. 1631.—China.
cordifolia, Lam. Encyc. ii. 504 = Echinodorus macrophyllus?
cordifolia, Roxb. Hort. Beng. 68; Fl. Ind. iii. 647 = guyanensis.
cristata, *Engelm. ex Arthur, Contrib. Fl. Iowa*, n. v; *ex Upham, Cat. Fl. Minnesota*, 138.—Am. bor.
Doniana, Sweet, Hort. Brit. ed. I. 375 = sagittifolia.
echinocarpa, Mart. Amoen. Bot. Monac. (Ausw. Merkw. Pfl.) 6. t. 3 = guayanensis.
edulis, Schlecht. in Linnaea, xviii. (1844) 432 = sagittifolia.
falcata, Pursh, Fl. Am. Sept. ii. 397 = lancifolia.
gigantea, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 759 = sagittifolia.
gracilis, Pursh, Fl. Am. Sept. ii. 396 = sagittifolia.
graminea, *Michx. Fl. Bor. Am.* ii. 190.—Am. bor.
guayanensis, *H. B. & K. Nov. Gen. et Sp.* i. 250.—Reg. trop.
guianensis, Schecht. in Linnaea, vi. (1831). 43 = praec.
guyanensis, Kunth, Syn. Pl. Aequin. I. 261 = praec.
hastata, Pursh, Fl. Am. Sept. ii. 396 = sagittifolia.
hermaphrodita, Buch.-Ham. in Wall. Cat. n. 4991 = sagittifolia.

SAGITTARIA:—
heterophylla, Bert. ex Steud. Nom. ed. II. ii. 491 = incrassata.
heterophylla, *Pursh, Fl. Am. Sept.* ii. 396.—Am. bor.
heterophylla, Schreb. in Schweig. & Koerte, Fl. Erlang. ii. 119 = sagittifolia.
hirundinacea, Blume, Enum. Pl. Jav. i. 34 = sagittifolia.
incrassata, *Steud. Nom.* ed. II. ii. 491.—Chili.
intermedia, *M. Mich. in DC. Monog. Phan.* iii. 80.—Ind. occ.
japonica, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1002 = sagittifolia.
lagoënsis, *Seub. ex Warm. in Kjoeb. Vidensk. Meddel.* (1872) 113.—Bras.
lancifolia, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 759 = sagittifolia.
lancifolia, *Linn. Syst.* ed. X. 1270.—Ind. occ.
lappula, D. Don, Prod. Fl. Nep. 22 = guayanensis.
latifolia, Willd. Sp. Pl. iv. 409 = sagittifolia.
longiloba, Engelm. in Torr. Mex. Bound. Surv. 212 = sagittifolia.
macrophylla, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1834) 137 = sagittifolia.
macrophylla, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 289.—Mexic.
major, Scop. Fl. Carn. ed. II. ii. 239; Schrank, Fl. Monac. iv. t. 336 = sagittifolia.
mexicana, Steud. Nom. ed. II. ii. 491 = macrophylla.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = sagittifolia.
monoeca, Gilib. Fl. Lituan. ii. 218 = sagittifolia.
montevidensis, *Cham. & Schlecht. in Linnaea*, ii. (1827) 156—Reg. Argent.
natans, *Michx. Fl. Am. Bor.* ii. 190.—Am. bor.
natans, Pall. Reise, iii. 757 = sagittifolia.
nymphaeifolia, Hochst. in Flora, xxiv. (1841) 1. Intell. 44; et ex Steud. Nom. ed. II. ii. 491 = Limnophyton obtusifolium.
obtusa, Thunb. Fl. Jap. 242 = sagittifolia.
obtusifolia, Linn. Sp. Pl. 993 = Limnophyton obtusifolium.
obtusissima, Hassk. Cat. Hort. Bog. Alt. 26 = guyanensis.
ovata, [Delile,] Red. Lil. t. 411 = lancifolia.
palaefolia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 21 = Echinodorus ellipticus.
papillosa, Buchen. in Bremen, Abh. ii. (1871) 27, 44 = lancifolia.
parviflora, Wall. Cat. ex Micheli, in DC. Monog. Phan. iii. 62 = guayanensis.
plantaginifolia, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 379 = lancifolia.
pubescens, *Muhl. Cat.* 86.—Am. bor.
pugioniformis, *Linn. Diss. Pl. Surin.* (1775) 15.—Guian.
Purshii, Rich. in Kunth, Enum. Pl. iii. 160 = graminea.
pusilla, Blume, Enum. Pl. Jav. i. 34 = guayanensis.
pusilla, Nutt. Gen. Am. ii. 213 = nutans.
pygmaea, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 138 = sagittifolia.
radicans, Nutt. in Trans. Am. Phil. Soc. v. (1837) 159 = Echinodorus radicans.
ranunculoides, Vell. Fl. Flum. t. 32 = Hydrocleys Commersonii.
rhombifolia, *Cham. in Linnaea*, x. (1836) 219.—Bras.
rhomboidalis, M. Mich. in DC. Monog. Phan. iii. 66 = rhombifolia.
rigida, *Pursh, Fl. Am. Sept.* ii. 397.—Am. bor.
sagittaefolia, Auct. = sagittifolia.
sagittata, Thunb. Fl. Jap. 242 = sagittifolia.
sagittifolia, *Linn. Sp. Pl.* 993.—Europ.; As.; Am. bor.
sagittifolia, Lour. Fl. Cochinch. 570 = chinensis.
sagittifolia, A. Rich. Fl. Cub. Fanerog. ii. 323 = pugioniformis.
sagittifolia, Vell. Fl. Flum. x. t. 31 = Echinodorus muricatus.
Sellowiana, Kunth, Enum. Pl. iii. 159 = lancifolia.
Seubertiana, *Mart. ex Seub. in Mart. Fl. Bras.* iii. 1. 110.—Bras.
simplex, Pursh, Fl. Am. Sept. ii. 397 = sagittifolia.
simplex, Torr. Consp. 356 = graminea.
sinensis, Sims, Bot. Mag. t. 1631 = sagittifolia.
Sprucei, *M. Mich. in DC. Monog. Phan.* iii. 80.—Bras.
stolonifera, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 234 = graminea.
subulata, Buchen. in Bremen, Abh. ii. (1871) 490, 500 = natans.

SAGITTARIA:—
subulata, *Schomb. Fauna & Fl. Brit. Gui.* 1117.—Guian.
teniaefolia, Phil. ex M. Mich. in DC. Monog. Phan. iii. 76 = montevidensis.
trachysepala, Engelm. ex M. Mich. l. c. 74 = lancifolia.
triandra, Dalz. in Hook. Kew Journ. ii. (1850) 144 = Wisneria triandra.
triflora, Hort. Bog. ex Miq. Fl. Ind. Bat. iii. 239 = Hydrocharis Morsus-ranae.
triflora, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = guayanensis.
trifolia, Linn. Sp. Pl. 993 = sagittifolia.
variabilis, Engelm. in A. Gray, Man. Bot. N. U. St. ed. V. 493 = sagittifolia.
vulgaris, Gueldenst. Reisen, ii. 45 = sagittifolia.

SAGMEN, Hill, Veg. Syst. iv. 35 (1762) = **Centaurea**, Linn. (Compos.).
jaceoides, Hill, Herb. Brit. i. 113 = C. Scabiosa.
Scabiosa, Hill, Hort. Kew. 66 = C. Scabiosa.
sempervirens, Hill, l. c. = C. sempervirens.

SAGONEA, Aubl. Pl. Gui. i. 285. t. 111 (1775) = **Hydrolea**, Linn. (Hydroph.).
aquatica, Poir. Encyc. vi. 392 = H. caroliniana.
palustris, Aubl. Pl. Gui. i. 285. t. 111 = H. caroliniana.

SAGOTIA, Baill. Adansonia, i. 53 (1860–61). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
racemosa, *Baill. l. c.* 54.—Guian.

SAGOTIA, Duchass. & Walp. in Linnaea, xxiii. (1850) 737 = **Desmodium**, Desv. (Legum.).
triflora, Duchass. & Walp. l. c. 738 = D. triflorum.

SAGRAEA, DC. Prod. iii. 170 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 766.
CAPITELLARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 103.
STAPHIDIASTRUM, Naud. l. c. xvii. (1852) 325.
acutiflora, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 9 = Henriettella fascicularis.
aphanantha, Triana, in Trans. Linn. Soc. xxviii. (1871) 137 = Clidemia aphanantha.
attenuata, *Triana, l. c.*—Ind. occ.
aureo-sericea, *Macfad. Fl. Jamaic.* ii. 100.—Jamaic.
Berterii, DC. Prod. iii. 171 = plumosa.
Bonplandii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 137.—Am. trop.
bracteolata, *Macfad. Fl. Jamaic.* ii. 55.—Jamaic.
capillaris, *DC. Prod.* iii. 170.—Jamaic.
capilliflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Bras.
capillipes, *Triana, l. c.*—Bras.
capitata, *Triana, l. c.* 139.—Guian.
cognata, Steud. in Flora, xxvii. (1844) 722 = Clidemia rubra.
columneaefolia, DC. Prod. iii. 170 = Clidemia rubra.
coriacea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Guian.
crossopetala, *Triana, l. c.* 139.—Jamaic.
discolor, *Triana, l. c.*—N. Granat.
domingensis, *DC. Prod.* iii. 171.—Ins. S. Doming.
dumosa, *Macfad. Fl. Jamaic.* ii. 53.—Jamaic.
epiphytica, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—N. Granat.
fascicularis, DC. Prod. iii. 170 = Henriettella fascicularis.
glomerata, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 96 = Ossaea glomerata.
Grisebachii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—Jamaic.
guadalupensis, *DC. Prod.* iii. 170.—Ins. Guadalup.
haeretica, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 100 = Oxymeris cornoides.
heteronervis, Naud. l. c. 98 = Ossaea heteronervis.
heteroneura, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 137.—Bras.
hirsuta, DC. Prod. iii. 171 = Ossaea hirsuta.
hirtella, DC. l. c. = Ossaea hirtella.
lanata, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 97 = Ossaea lanata.
latifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Ind. occ.
Lima, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 99 = Ossaea Lima.
Lindeniana, Naud. l. c. 96 = Ossaea scabrosa.

SAGRAEA:—
macrandra, *Wright, in Sauv. Fl. Cub.* 46.—Cuba.
melanophyllum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—Bras.
microphylla, DC. Prod. iii. 171 = Ossaea microphylla.
minutiflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 137.—Bras.
neurocarpa, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 94 = Octopleura micrantha.
obliqua, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—Peruv.
ossaeaeformis, *Triana, l. c.* 138.—Ind. occ.
pauciflora, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 100 = Calycogonium echinatum.
penninervis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—Cuba.
petiolata, Triana, l. c. = Clidemia Deppeana.
pilosa, *DC. Prod.* iii. 171.—Jamaic.
platyphylla, *Triana*, in Trans. Linn. Soc. xxviii. (1871) 137.—Bras.
plumosa, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 93.—Jamaic.; Venezuela.
polystachya, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Porto Rico.
racemifera, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 91, 102 = Miconia racemifera.
rariflora, *DC. Prod.* iii. 171.—Venezuela.
rubinervis, Sauv. Fl. Cub. 47 = domingensis.
rubra, Triana, in Trans. Linn. Soc. xxviii. (1871) 137 = Clidemia rubra.
salicifolia, *Triana, l. c.*—Ins. Trinit.
scabrida, DC. Prod. iii. 170 = Henriettella fascicularis.
scabrosa, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 97 = Ossaea glomerata.
scabrosa, Seem. Bot. Voy. Herald, 124 = Henriettella Seemannii.
scalpta, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 95 = Ossaea scalpta.
serpens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 139.—N. Granat.
sessiliflora, DC. Prod. iii. 170 = Clidemia rubra.
sessiliflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 137.—Am. trop.
sparsiflora, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 98 = Henriettella lateriflora.
sparsiflora, Wawra, Bot. Reis. Maximil. 24 = Henriettella umbelluliflora.
tetragona, *DC. Prod.* iii. 171.—Ins. Dominic.
trichotoma, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Cuba.
trinitensis, *Triana, l. c.* 137.—Ins. Trinit.
umbrosa, *DC. Prod.* iii. 171.—Ind. occ.
virgata, *Macfad. Fl. Jamaic.* ii. 52.—Jamaic.
Wrightii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 138.—Cuba.

SAGUERUS, Adans. Fam. ii. 24 (1763) = **Arenga**, Labill. (Palm.).
americanus, H. Wendl. in Kerch. Palm. 256 = Mauritia flexuosa.
australasicus, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 219.—Austral.
Gamuto, Houtt. Handleid. i. 410. t. 4. f. 2 = A. saccharifera.
Langbak, Blume, Rumphia, ii. 131. tt. 96, 125.—Malaya.
Rumphii, Roxb. Hort. Beng. 68; Fl. Ind. iii. 626 = A. saccharifera.
saccharifer, Blume, Rumphia, ii. 128. t. 123-124 = A. saccharifera.
Westerhoutii, H. Wendl. & Drude, in Kerch. Palm. 256 = A. obtusifolia.
Wightii, H. Wendl. & Drude, l. c. = A. Wightii.

SAGUS, Rumph. ex Gaertn. Fruct. i. 27 (1788) = **Metroxylon**, Rottb. (Palm.).
americana, Poir. Encyc. vi. 395 = Mauritia Sagus.
amicarum, H. Wendl. in Bot. Zeit. xxxvi. (1878) 115.—Ins. Pacif.
elata, Reinw. ex Blume, Rumphia, ii. 156. t. 128. f. 1 = Pigafettia elata.
farinifera, Gaertn. Fruct. ii. 186 = Raphia pedunculata.
filaris, Blume, Rumphia, ii. 154. t. 128 = Pigafettia filaris.
filifera, Gisek. Prael. 94.—Ind. or.
gemina, Steud. Nom. ed. II. ii. 492, sphalm. = *genuina*, Labill.

SAGUS:—
genuina, Gisek. Prael. 94 = Metroxylon Sagu.
genuina, Labill. ex Blume, Rumphia, ii. 153 = M. micracanthum.
hospita, Kerch. Palm. 162, in adnot.—Hab. ?
inermis, Roxb. Hort. Beng. 68; Fl. Ind. iii. 623 = M. Sagu.
Koenigii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 19 = M. Sagu.
laevis, Jack, in Malay. Misc. iii. [1823 ?] 9; Blume, Rumphia, ii. 147. tt. 86, 126, 127 = M. Sagu.
longispina, Blume, l. c. 154 = M. longispinum.
micracantha, Blume, l. c. 153 = M. micracanthum.
microcarpa, Zipp. ex Hall, Bijdr. Nat. Wet. v. (1830) 178, nomen = Pigafettia filaris.
microsperma, Zipp. ex Hall, l. c. 178 = Pigafettia filaris.
Palma-pinus, Gaertn. Fruct. i. 27. t. 10 = Raphia Gaertneri.
pedunculata, Poir. Encyc. Suppl. v. 13 = Raphia pedunculata.
Raphia, Poir. l. c. vi. 395 = Raphia vinifera.
Ruffia, Jacq. Fragm. 7 = Raphia pedunculata.
Rumphii, Willd. Sp. Pl. iv. 404 = M. Sagu.
spinosa, Roxb. Hort. Beng. 68, nomen = M. Sagu ?
sylvestris, Blume, Rumphia, ii. 153 = Pigafettia elata.
taedigera, Mart. Hist. Nat. Palm. ii. 54. t. 45 = Raphia vinifera.
vinifera, Pers. Syn. ii. 562 = Raphia vinifera.
vitiensis, H. Wendl. in Seem. Viti, App. 444; et in Fl. Vit. 279. t. 80 = M. vitiense.

SAHAGUNIA, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 316. *URTICACEAE*, Benth. & Hook. f. iii. 377.
ACANTHINOPHYLLUM, Fr. Allem. in Rev. Brazil. i. (1858) 368.
mexicana, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 316.—Mexic.

SAHLBERGIA, Reichb. Consp. 95 (1828) = SALHBERGIA, Neck. = **Gardenia**, Ellis (Rubiac.).

SAINTLEGERIA, Cordem. in Baill. Adansonia, iii. (1862–63) 300 = **Chloranthus**, Sw.
gracilis, Cordem. ex Baill. l. c.—China.

SAINTMORYSIA, Endl. Gen. 438 (1838) = **Athanasia**, Linn. (Compos.).
pachyphylla, Walp. Rep. vi. 211 = A. dentata.

SAIRANTHUS, G. Don, Gen. Syst. iv. 467 (1837) = **Nicotiana**, Linn. (Solan.).
glutinosus, G. Don, l. c. = N. glutinosa.

SAIROCARPUS, Nutt. ex A. DC. in DC. Prod. x. 592 (1846) = **Antirrhinum**, Tourn. (Scrophul.).
californicus, Nutt. l. c. = A. Nuttallianum.

SAIVALA, Jones, in As. Res. iv. (1799) 275 = **Blyxa**, Noronha (Hydroch.).
valisnerioides, Buch.-Ham. ex Wall. Cat. n. 5047 = B. Saivala.

SAJORIUM, Endl. Gen. Suppl. iii. 98 (1843) = **Plukenetia**, Linn. (Euphorb.).
africanum, Baill. Etud. Gén. Euph. 483.—Afr. austr.
corniculatum, Baill. l. c. 484 = P. corniculata.
corniculatum, D. Dietr. Syn. Pl. v. 331.—Java.
tamnoides, Baill. Etud. Gén. Euph. 484 = P. tamnoides.
volubile, Baill. l. c. 483 = P. volubilis.

SAKERSIA, Hook. f. in Benth. & Hook. f. Gen. i. 757 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
africana, *Hook. f. l. c.*—Afr. trop.
echinulata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 458.—Afr. trop.

SALABERTIA, Neck. Elem. ii. 238 (1790) = **Tapiria**, Juss. (Anacard.).

SALACCA, Reinw. in Syll. Ratisb. ii. (1828) 3 = **Zalacca**, Rumph. (Palm.).
edulis, Reinw. l. c. = Z. edulis.

SALACIA, Linn. Mant. ii. 159, 293 (1771). *CELASTRINEAE*, Benth. & Hook. f. i. 370.
ANTHODISCUS, Endl. Gen. 1091, in syn. (1841).
ANTHODON, Ruiz & Pav. Fl. Per. i. 45 (1798).
ANTHODUS, Mart. in Schult. Mant. i. 253, 348 (1822).
CALYPSO, Thou. Hist. Vég. Afr. i. 33. t. 6 (1805).
CLERCIA, Vell. Fl. Flum. i. tt. 73, 74 (1827).
CUSTINIA, Neck. Elem. ii. 223 (1790).
DIPLESTHES, Harv. in Hook. Lond. Journ. Bot. i. (1842) 19.
JOHNIA, Roxb. Fl. Ind. ed. Carey, i. 172 (1820).
KIPPISTIA, Miers, in Trans. Linn. Soc. xxviii. (1872) 414, 416.
MACAHANEA, Aubl. Pl. Gui. Suppl. 6. t. 371 (1775).
MACANEA, Juss. Gen. 257 (1789).
PERITASSA, Miers, in Trans. Linn. Soc. xxviii. (1872) 402.
PYRAMIDOSTYLIUM, Mart. ex Peyr. in Mart. Fl. Bras. xi. 1. 155 (1878).
RADDIA, Spreng. Syst. i. 177, in syn. (1825).
RADDISIA, Leandro, in Denkschr. Akad. Muench. vii. (1821) 237. t. 15.
SARCOCAMPSA, Miers, in Trans. Linn. Soc. xxviii. (1872) 408.
TONSELLA, Schreb. Gen. 34 (1789).
TONTELEA, Aubl. Pl. Gui. i. 31. t. 10 (1775).
affinis, Hook. f. in Hook. Niger Fl. 281 = senegalensis.
affinis, *Peyr. in Mart. Fl. Bras.* xi. 1. 159.—Bras.
africana, *DC. Prod.* i. 570.—Afr. trop.
alternifolia, *Hochst. in Flora*, xxvii. (1844) 306.—Afr. austr.
amygdalina, *Peyr. in Mart. Fl. Bras.* xi. 1. 159.—Bras.
anomala, *Peyr. l. c.* 144.—Bras.
arborea, Peyr. l. c. 156 = undulata.
attenuata, *Peyr. l. c.* 149.—Bras.
brachypoda, *Peyr. l. c.* 150.—Bras.
brasiliensis, *G. Don, Gen. Syst.* i. 628.—Bras.
Brunoniana, *Wight & Arn. Prod.* 105.—Ind. or.
Buddinghii, *Scheff. in Flora*, lii. (1869) 306; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 16.—Java.
Calypso, *DC. Prod.* i. 571.—Madag.
calypsoides, *A. St. Hil. Fl. Bras. Mer.* ii. 107.—Bras.
campestris, *Walp. Rep.* i. 402.—Bras.
capilliflora, Sagot, ex Peyr. in Mart. Fl. Bras. xi. 1. 134 = Hippocratea comosa.
celebica, *Blume, Rumphia*, iv. 19.—Ins. Celebes.
cerasifera, *Welw. ex Oliver, Fl. Trop. Afr.* i. 376.—Afr. trop.
cerasiformis, *Teijsm. & Binn. Cat. Hort. Bog.* 219.—Java.
chinensis, Linn. Mant. ii. 293 = Roxburghii?
chlorantha, *Oliver, Fl. Trop. Afr.* i. 375.—Afr. trop.
cochinchinensis, Lour. Fl. Cochinch. 526 = Roxburghii?
cognata, *Peyr. in Mart. Fl. Bras.* xi. 1. 144.—Bras.
conferta, *Peyr. l. c.* 159.—Bras.
cornifolia, *Hook. f. in Hook. Niger Fl.* 281.—Afr. trop.
coromandeliana, *Teijsm. & Binn. Cat. Hort. Bog.* 392, nomen.—Ind. or.
crassifolia, *G. Don, Gen. Syst.* i. 628.—Bras.
debilis, *Walp. Rep.* i. 402.—Afr. trop.
decussata, G. Don, Gen. Syst. i. 627 = Hippocratea decussata.
densiflora, *Wall. Cat.* n. 4224.—Ind. or.; Burma.
dentata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 334.—Madag.
diandra, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 151.—Ins. Celebes.
diandra, Thw. Enum. Pl. Zeyl. 53, 407 = reticulata.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 581.—Hab.?
diffusiflora, *Peyr. in Mart. Fl. Bras.* xi. 1. 143.—Bras.
distincta, *Peyr. l. c.* 151.—Bras.
Ducis-Wuertembergiae, *Hochst. in Flora*, xxvii. (1844) 307.—Afr. trop.
dulcis, *Benth. in Hook. Kew Journ.* iv. (1852) 9.—Bras.
elegans, *Welw. in Oliver, Fl. Trop. Afr.* i. 378.—Afr. trop.
elliptica, *G. Don, Gen. Syst.* i. 628.—Bras.
elongata, *Hook. f. in Hook. Niger Fl.* 282.—Afr. trop.
erecta, *Walp. Rep.* i. 403.—Afr. trop.

SALACIA:—
erythroxyloides, A. St. Hil. Fl. Bras. Mer. ii. 160 = elliptica.
evonymiflora, *Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 149.—Java.
exsculpta, *Korth. in Flora*, xxxi. (1848) 579.—Sumatra.
Finlaysonii, *Wall. Cat.* n. 4220.—Siam.
flavescens, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 300.—Burma; Penang.
floribunda, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 97.—Ins. Comor.
floribunda, Wight, Illustr. i. 134 = densiflora.
fluminensis, *Peyr. in Mart. Fl. Bras.* xi. 1. 149.—Bras.
fruticosa, *Wall. Cat.* n. 4223.—Ind. or.
glomerata, *G. Don, Gen. Syst.* i. 628.—Bras.
grandiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 300.—Burma; Penang.
grandiflora, *Peyr. in Mart. Fl. Bras.* xi. 1. 157.—Bras.
grandifolia, *G. Don, Gen. Syst.* i. 628.—Bras.
Griffithii, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 628.—Ind. or.
guianensis, Klotzsch, ex Peyr. in Mart. Fl. Bras. xi. 1. 158 = lacunosa.
hippocrateoides, *Peyr. l. c.* 142.—Bras.
javanensis, *Blume, Bijdr.* 219.—Java.
Jenkinsii, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 203.—Assam.
kalahiensis, *Korth. in Flora*, xxxi. (1848) 579.—Borneo.
Korthalsiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 152.—Java.
Kraussii, Harv. in Harv. & Sond. Fl. Cap. i. 230 = alternifolia.
lacunosa, *Peyr. in Mart. Fl. Bras.* xi. 1. 158.—Guian.
laevigata, *DC. Prod.* i. 570.—Hab.?
laevigata, Wight, in Hook. Bot. Misc. iii. (1833) 295. Suppl. t. 36 = Hippocratea obtusifolia.
lanceolata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 424.—Java.
latifolia, *Wall. Cat.* n. 4222.—Singapore.
laxiflora, *Peyr. in Mart. Fl. Bras.* xi. 1. 145.—Bras.
leptoclada, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 96.—Ins. Comor.
Lobbii, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 626.—Singapore.
longifolia, *Hook. f. ex M. Laws. l. c.*—Malaya.
longifolia, Wall. Pl. As. Rar. iii. 47. t. 278; Cat. n. 4225 = densiflora.
lucida, *Oliver, Fl. Trop. Afr.* i. 373.—Afr. trop.
macrocarpa, *Korth. in Flora*, xxxi. (1848) 579.—Borneo.
macrophylla, *Blume, Bijdr.* 221.—Cochinch.; Java.
macrophylla, *Miq. Stirp. Surin. Sel.* 85.—Guian.
macrosperma, *Wight, Spicil. Neilgherr.* i. 28. t. 32; *Ic.* t. 962.—Ind. or.
madagascariensis, *DC. Prod.* i. 570.—Madag.
Maingayi, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 626.—Penang.
Mannii, *Oliver, Fl. Trop. Afr.* i. 376.—Afr. trop.
Martiana, *Peyr. in Mart. Fl. Bras.* xi. 1. 146.—Bras.
melitocarpa, *Blume, Bijdr.* 220.—Java.
membranacea, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 627.—Reg. Himal.
memecyloides, Wall. Cat. n. 4218 = verrucosa.
micrantha, *G. Don, Gen. Syst.* i. 628.—Bras.
Miersii, *Peyr. in Mart. Fl. Bras.* xi. 1. 147.—Bras.
multiflora, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 109 = Hippocratea ovata.
multiflora, *DC. Prod.* i. 570.—Guian.
multiflora, Wight, Illustr. i. 134 = polyantha, *Steud.*
myrtifolia, *Griff. Notul.* iv. 470.—Burma.
oblonga, *Wall. Cat.* n. 4226.—Ind. or.
oblongifolia, *Blume, Bijdr.* 220.—Java.
oblongifolia, G. Don, Gen. Syst. i. 628 = elliptica.
oblongifolia, *Oliver, Fl. Trop. Afr.* i. 374.—Afr. trop.
oblongifolia, Rottl. ex Steud. Nom. ed. II. ii. 492 = oblonga.
obovata, Boiv. ex. Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 96 = multiflora.
obtusifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 105.—Bras.
oleoides, *Baker, in Journ. Linn. Soc.* xxi. (1884) 334.—Madag.

SALACIA:—
oppositifolia, Rottl. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 628 = oblonga.
ovalis, *Korth. in Flora*, xxxi. (1848) 579.—Java.
ovalis, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 627.—Malacca.
pachyphylla, *Peyr. in Mart. Fl. Bras.* xi. 1. 158.—Guian.
pallescens, *Oliver, Fl. Trop. Afr.* i. 376.—Afr. trop.
Pancheri, *Baill. Adansonia*, x. (1872) 184.—N. Caled.
paniculata, *G. Don, Gen. Syst.* i. 627.—Bras.
paniculata, Wall. Cat. n. 4214 B, partim = Hippocratea Grahami.
parviflora, *Sagot, ex Peyr. in Mart. Fl. Bras.* xi. 1. 143.—Guian.
patens, *Decne. ex Steud. Nom.* ed. II. ii. 492.—Ins. Timor.
patens, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 374.—N. Granat.
platyphylla, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 203.—Ins. Nicobar.
podopetala, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 581 = prinoides.
polyantha, *Korth. in Flora*, xxxi. (1848) 579.—Borneo.
polyantha, *Steud. Nom.* ed. II. ii. 492.—Burma.
pomifera, Wall. Cat. n. 4227 = oblonga.
pomifera, Wight, Arn. Prod. 105 = densiflora.
prinoides, *DC. Prod.* i. 571.—Ind. or.
pruinosa, Seem. Bot. Voy. Herald, 90 (Hippocrateae sp.).—Panama.
pyriformis, *Steud. Nom.* ed. II. ii. 492; *Walp. Rep.* i. 402.—Afr. trop.
Radula, *G. Don, Gen. Syst.* i. 628.—Bras.
reticulata, *Wight, Illustr.* i. 134.—Ind. or.
Richardii, *Peyr. in Mart. Fl. Bras.* xi. 1. 148.—Guian.
Riedeliana, *Peyr. l. c.* 147.—Bras.
Roxburghii, *Wall. Cat.* n. 4217.—Ind. or.
rubra, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 627.—Malacca.
rufescens, *Hook. f. in Hook. Niger Fl.* 283.—Afr. trop.
rugosa, *Wall. Cat.* n. 4228.—Ind. or.
rugulosa, *Sagot, in Ann. Sc. Nat.* Sér. VI. xi. (1881) 174.—Guian.
saigonensis, *Baill. Adansonia*, xi. (1873-76) 272.—Cochinch.
scabra, *DC. Prod.* i. 570.—Guian.
scandens, *Griseb. Fl. Brit. W. Ind.* 148.—Am. trop.
senegalensis, *DC. Prod.* i. 570.—Afr. trop.
serrata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 109.—Bras.
sinensis, Blanco, Fl. Filip. ed. I. 26 = prinoides.
sinensis, J. F. Gmel. Syst. 107 = *chinensis*.
sinensis, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 47 (Quid?).—Ins. Philipp.
sororia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 151.—N. Guin.
sylvestris, *Steud. Nom.* ed. II. ii. 92; *Walp. Rep.* i. 402.—Bras.
tennicula, *Peyr. in Mart. Fl. Bras.* xi. 1. 146.—Bras.
terminalis, Thw. Enum. Pl. Zeyl. 407 = Hippocratea Arnottiana.
tortuosa, *Griff. Notul.* iv. 471; *Ic. Pl. Asiat.* t. 581. f. 2.—Burma.
trigonocarpa, *Boiv. ex Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 96.—Madag.
trinervia, *G. Don, Gen. Syst.* i. 628.—Bras.
triplinervis, *Llanos, in Mem. Acad. Cienc. Mad.* ii. (1859) 500.—Hab.?
undulata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 107.—Bras.
Vahliana, *Cambess. l. c.* 105, in adnot.—Ins. Trinit.
velutina, *Cambess. l. c.* 108.—Bras.
verrucosa, *Wight, Illustr.* i. 134.—Burma; Ins. Philipp.
verrucosa, *Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 172, in syn.—Cuba.
viminea, *Wall. Cat.* n. 7267.—Burma.
viridiflora, *Payer, Organog.* 164. t. 35. f. 1-13.—Hab.?
Wightiana, Wall. Cat. n. 4221 = prinoides.
Zeyherii, *Spreng. ex Harv. & Sond. Fl. Cap.* i. 230.—Afr. austr.

SALACISTIS, Reichb. f. in Bonplandia, v. (1857) 36 = **Hetaeria**, Blume (Orchid.).
novembrilis, Reichb. f. l. c.—Java.

SALAKKA, Reinw. ex Blume, Cat. Gew. Buitenz. 112 (1823) = **Zalacca**, Rumph. (Palm.).
edulis, Reinw. l. c. = Z. edulis.

SALAXIS, Salisb. in Trans. Linn. Soc. vi. (1802) 317. *ERICACEAE*, Benth. & Hook. f. ii. 594.
COCCOSPERMA, Klotzsch, in Linnaea, xii. (1838) 215.
LAGENOCARPUS, Klotzsch, l. c (1838).
abietina, Willd. Enum. Hort. Berol. 415 = Philippia abietina.
arborescens, Willd. l. c. 415 = Philippia arborescens.
aristata, D. Dietr. Syn. Pl. ii. 1261 = Philippia aristata.
artemisioides, *Klotzsch, in Linnaea*, xii. (1838) 212.—Afr. austr.
axillaris, *Salisb. ex G. Don, Gen. Syst.* iii. 828.—Afr. austr.
brachyphylla, D. Dietr. Syn. Pl. ii. 1261 = Philippia brachyphylla.
brevifolia, D. Dietr. l. c. = ciliata.
Chamissonis, D. Dietr. l. c. 1260 = Philippia Chamissonis.
ciliata, *Benth. in DC. Prod.* vii. 711.—Afr. austr.
ciliata, D. Dietr. Syn. Pl. ii. 1260 = Philippia ciliata.
densa, D. Dietr. l. c. = Philippia densa.
flexuosa, *Klotzsch, in Linnaea*, xii. (1838) 213.—Afr. austr.
floribunda, D. Dietr. Syn. Pl. ii. 1260 = Philippia floribunda.
galioides, D. Dietr. l. c. = Philippia galioides.
Goudotiana, D. Dietr. l. c. = Philippia Goudotiana.
gracilis, D. Dietr. l. c. 1261 = Ericinella gracilis.
hexandra, *Klotzsch, in Linnaea*, ix. (1834) 352.—Afr. austr.
imbricata, Willd. ex Steud. Nom. ed. II. ii. 319, 493 = Philippia abietina.
Leeana, D. Dietr. Syn. Pl. ii. 1260 = Philippia Leeana.
mauritiana, Boj. ex Wall. Cat. n. 1525 = Philippia abietina.
micrantha, *Benth. in DC. Prod.* vii. 711.—Afr. austr.
montana, Boj. Hort. Maurit. 195 = Philippia brachyphylla.
montana, Willd. Enum. Hort. Berol. 415 = Philippia montana.
multiflora, D. Dietr. Syn. Pl. ii. 1261 = Ericinella multiflora.
octandra, *Klotzsch, in Linnaea*, ix. (1834) 353.—Afr. austr.
parviflora, D. Dietr. Syn. Pl. ii. 1260 = Philippia parviflora.
puberula, *Klotzsch, in Linnaea*, xii. (1838) 212.—Afr. austr.
ramosissima, D. Dietr. Syn. Pl. ii. 1260 = Philippia ramosissima.
rugosa, *Benth. in DC. Prod.* vii. 711.—Afr. austr.
Sieberi, *Benth. l. c.*—Afr. austr.
tenuifolia, D. Dietr. Syn. Pl. ii. 1260 = Philippia tenuifolia.
tenuissima, D. Dietr. l. c. 1261 = Philippia tenuissima.
viscosa, D. Dietr. l. c. = Philippia viscosa.

SALAZARIA, Torr. in Bot. Mex. Bound. 133. t. 39 (1858). *LABIATAE*, Benth. & Hook. f. ii. 1201.
mexicana, *Torr. l. c.*—Mexic.

SALCEDA, Blanco, Fl. Filip. ed. II. 374 (1854) = **Camellia**, Linn. (Ternstr.).
montana, Blanco, l. c. = C. lanceolata.

SALDANHA, Vell. Fl. Flum. 141 (1825); Ic. iii. tt. 157, 158 (1827) = **Hillia**, Jacq. (Rubiac.).
illustris, Vell. l. c. 142; iii. t. 157.—Bras.
nobilis, Vell. l. c. t. 158.—Bras.

SALDANHAEA, Bur. in Baill. Adansonia, viii. (1867–68) 354. tt. 7, 11, 12. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1035.
confertiflora, *Bur. l. c.* 356.—Bras.
lateriflora, *Bur. l. c.*—Bras.

SALDINIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 206. *RUBIACEAE*, Benth. & Hook. f. ii. 129.
Pseudomorinda, *A. Rich. l. c.* 206.—Madag.

SALGADA, Blanco, Fl. Filip. ed. II. 221 (1845) = **Eusideroxylon**, Teijsm. & Binn. (Laurin.).
lauriflora, Blanco, l. c. = E. Zwageri.

SALHBERGIA, Neck. Elem. i. 262 (1790) = **Gardenia**, Ellis (Rubiac.).

SALICA, Hill, Hort. Kew. 172/24 (1768) = **Lythrum**, Linn.
Lythastrum, Hill, l. c. = L. Thymifolia.

SALICARIA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Lythrum**, Linn.
Graefferi, Hort. Berol. ex Steud. Nom. ed. II. ii. 86, 493 L. Hyssopifolia.
Hyssopifolia, Lam. Fl. Fr. iii. 103 = L. Hyssopifolia.
Hyssopifolia, Moench, Meth. Suppl. 281 = L. Salicaria.
spicata, Lam. Fl. Fr. iii. 103 = L. Salicaria.
Thymifolia, Lam. l. c. 104 = L. Thymifolia.
virgata, Moench, Meth. 665 = L. virgatum.
vulgaris, Moench, l. c. = L. Salicaria.

SALICORNIA, [Tourn.] Linn. Sp. Pl. ed. I. 357 (1737). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 66.
SARCATHRIA, Rafin. Fl. Tellur. iii. 47 (1836).
acetaria, Pall. Illustr. tt. 1, 2. f. 1 = herbacea.
Alpini, Lag. Mem. Pl. Barrill. 48 = fruticosa.
ambigua, Michx. Fl. Bor. Am. i. 2 = fruticosa.
amplexicaulis, Vahl, Symb. Bot. ii. 1 = Halopeplis amplexicaulis.
anceps, Lag. Mem. Pl. Barill. 52 = fruticosa.
annua, Sm. Engl. Bot. t. 415 = herbacea.
appressa, *Dum. in Bull. Soc. Bot. Belg.* vii. (1866) 333.—Europ.
arabica, Linn. Sp. Pl. 3 = fruticosa.
arabica, Pall. Reise, i. 481 = Kalidium caspicum.
arabica, Raddi, ex Moq. in DC. Prod. xiii. II. 157 = Suaeda fruticosa.
Arbuscula, R. Br. Prod. 411 = Arthrocnemum arbusculum.
Arbuscula, DC. ex Moq. in DC. Prod. xiii. II. 151 = fruticosa.
Arbuscula, Moric. ex Steud. Nom. ed. II. ii. 493 = Arthrocnemum glaucum.
australis, *Soland. ex Forst. f. Prod.* 88.—Austral.
Bergii, *Lor. & Niederl. in Roca, Exped. Rio-Negro*, ii. 194.—Patag.
bidens, *Benth. Fl. Austral.* v. 204.—Austral.
biennis, Afzel. ex Sm. Fl. Brit. i. 2 = herbacea.
Bigelowii, *Torr. Bot. Mex. Bound.* 184.—Am. bor.
brachiata, Miq. Fl. Ind. Bat. i. 1. 1019 = Arthrocnemum ciliolatum.
brachiata, *Roxb. Hort. Beng.* [79]; *Fl. Ind.* i. 84.—Ind. or.
californica, Nutt. ex Moq. in DC. Prod. xiii. II. 151 = Arthrocnemum fruticosum.
caspica, Linn. Sp. Pl. 4 = Kalidium caspicum.
caspica, Pall. Reise, i. 480 = Halostachys caspica.
cinerea, *F. Muell. Fragm.* vi. 251.—Austral.
corticosa, Walp. ex Ung. Sternb. Vers. Syst. Salic. 60 = Gaudichaudiana.
cruciata, Coqueb. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876) 310 = herbacea.
cruciata, Forsk. Fl. Aegypt. Arab. 2 = Halocnemum strobilaceum.
Doeringii, *Lor. & Niederl. in Roca, Exped. Rio-Negro*, ii. 194.—Patag.
drepanensis, Tineo, ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 338 = Halocnemum strobilaceum.
Emerici, Duval-Jouve, in Bull. Soc. Bot. Fr. xv. (1868) 176 = herbacea.
equisetifolia, Willd. ex Moq. in DC. Prod. xiii. II. 145 = fruticosa.
europaea, Linn. Sp. Pl. 3 = fruticosa, herbacea.
falcata, J. F. Gmel. Syst. 1618, sphalm. = *foliata*.
fastigiata, Losc. & Pard. Ser. Inconf. Pl. Arag. ed. I. 90 = Microcnemum fastigiatum.
foliacea, Georgi, Beschr. Russ. Reich. iv. 610 = Kalidium foliatum.
foliata, Pall. Reise, i. 482 = Kalidium foliatum.
foliosa, Vahl, Enum. i. 12 = Kalidium foliatum.
frutescens, Friedrich. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 296 = fruticosa.
fruticosa, *Linn. Sp. Pl.* ed. II. 5.—Europ.; Afr. bor.
fruticosa, Moq. Chenop. Enum. 112 = Halocnemum strobilaceum.
fruticosa, Retz. Prod. Fl. Scand. ed. II. 1 = herbacea.
fruticosa, Sibth. & Sm. Fl. Graec. Prod. i. 1 = Arthrocnemum glaucum, macrostachyum.
fruticulosa, Tineo, Cat. Pl. Hort. Panorm. (1827) 280 = fruticosa.

SALICORNIA:—
Gaudichaudiana, *Moq. Chenop. Enum.* 115.—Bras.
glauca, Delile, Fl. Egypt. Illustr. 69 = Arthrocnemum glaucum.
glauca, Sieber, ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 338 = Halocnemum strobilaceum.
glauca, Stokes, Bot. Mat. Med. i. 8 = fruticosa?
herbacea, Fée, ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 296 = fruticosa.
herbacea, *Linn. Sp. Pl.* ed. II. 5.—Europ.
indica, R. Br. Prod. 411 = australis.
indica, Drège, Zwei Pfl. Docum. 159, 217 = natalensis.
indica, Ritt. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 333 = Heterostachys Ritteriana.
indica, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 111. t. 4. f. 4 = Arthrocnemum indicum.
intermedia, Woods, in Proc. Linn. Soc. ii. (1851) 110 = herbacea.
leiostachya, *Benth. Fl. Austral.* v. 203.—Austral.
lignosa, Woods, in Proc. Linn. Soc. ii. (1851) 111 = fruticosa?
macedonica, Bunge, ex Ung. Sternb. Vers. Syst. Salic. 48 = herbacea.
macrostachya, Moric. Fl. Venet. i. 2 = Arthrocnemum glaucum.
megastachya, *Woods, in Proc. Linn. Soc.* ii. (1851) 113.—Europ.
mucronata, Bigel. Fl. Bost. ed. III. 2 = Bigelowii.
mucronata, Lag. Mem. Pl. Barill. 58 = Arthrocnemum glaucum.
natalensis, *Bunge, ex Ung. Sternb. Vers. Syst. Salic.* 62.—Afr. austr.
Neei, Lag. Mem. Pl. Barill. 51 = fruticosa.
nodulosa, Delile, Fl. Egypte, 147 = Halopeplis amplexicaulis.
pachystachya, *Bunge, ex Ung. Sternb. Vers. Syst. Salic.* 51.—Madag.
Pallasiana, C. A. Mey. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 321 = Kalidium foliatum.
Pallasii, C. A. Mey. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) 361 = Kalidium caspicum.
patula, Duval-Jouve, in Bull. Soc. Bot. Fr. xv. (1868) 175 = herbacea.
peregrina, Weinm. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 309 = herbacea.
perennans, Willd. Sp. Pl. i. 24 = herbacea.
perennis, Mill. Gard. Dict. ed. VIII. n. 2 = herbacea.
perfoliata, Forsk. Fl. Aegypt. Arab. 3 = Halopeplis perfoliata.
peruviana, H. B. & K. Nov. Gen. et Sp. ii. 193 = fruticosa.
procumbens, Sm. Engl. Bot. t. 2475 = herbacea.
prostrata, Pall. Illustr. t. 3 = herbacea.
pusilla, Woods, in Proc. Linn. Soc. ii. (1851) 110 = herbacea.
pygmaea, Pall. Illustr. i. 8. t. 2. f. 2 = Halopeplis pygmaea.
quinqueflora, Bunge, ex Ung. Sternb. Vers. Syst. Salic. 59 = australis.
radicans, Mert. & Koch, in Rochl. Deutschl. Fl. ed. III. i. 292 = herbacea.
radicans, Sm. Engl. Bot. t. 1691 = fruticosa.
ramosissima, Woods, in Proc. Linn. Soc. ii. (1851) 110 = herbacea.
robusta, F. Muell. Fragm. vi. 251 = Pachycornia robusta.
Salsola, Coqueb. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876), 310 = herbacea.
sarmentosa, Duval-Jouve, in Bull. Soc. Bot. Fr. xv. (1868) 174 = fruticosa.
sempervirens, Sauvages, ex Steud. Nom. ed. I. 714 = fruticosa.
simplex, Stokes, Bot. Mat. Med. i. 7 = virginica.
stricta, Willd. ex Steud. Nom. ed. I. 714 = herbacea.
strobilacea, Delile, Fl. Egypte. t. 3. f. 2 = Halopeplis amplexicaulis.
strobilacea, Gaertn. Fruct. ii. 210. t. 127. f. 8 = Kalidium foliatum.
strobilacea, Pall. Reise, i. 481 = Halocnemum strobilaceum.
tenuis, *Benth. Fl. Austral.* v. 204.—Austral.
virginica, Forsk. Fl. Aegypt. Arab. 2 = Arthrocnemum glaucum.
virginica, *Linn. Sp. Pl.* 4 (virginia).—Am. bor.
virginica, Nutt. Gen. Am. i. 2 = Bigelowii.
virginica, Scholler, Barb. Suppl. 311 = fruticosa.

SALIMORI, Adans. Fam. ii. 177 (1763) = **Cordia**?, Linn. (Borag.).

SALIOLA, Schwag. ex Leplay, in Démid. Voy. Russ. Mér. ii. 236 (1842) = **Salsola**?, Linn. (Chenop.).
Loeselii, Schwag. l. c. (Quid?).—Ross.

SALISBURIA, Sm. in Trans. Linn. Soc. iii. (1797) 330 = **Ginkgo**, Linn. (Conif.).
adiantifolia, Sm. l. c. = G. biloba.
Billardierii, Rich. ex Carr. Conif. ed. I. 500 = Phyllocladus rhomboidalis.
biloba, Hoffmgg. Verz. Pfl. 109 = G. biloba.
Ginkgo, Rich. Conif. 133. t. 3bis = G. biloba.
macrophylla, Reyn. Cat. Sénécl. (1854) 40 = G. biloba.

SALISBURYA, Hoffmgg. Verz. Pfl. 109 (1824) = praec.

SALISIA, Lindl. Swan River App. 10 (1839) = **Kunzea**, Reichb. (Myrtac.).
pulchella, Lindl. l. c. = K. sericea.

SALISIA, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 373 = **Xanthostemon**, F. Muell. (Myrtac.).
aurantiaca, Panch. l. c. (= *Fremya aurantiaca*).—N. Caled.
flava, Panch. l. c. (= *Fremya flava*).—N. Caled.
rubra, Panch. l. c. = X. rubrum.
rugosa, Panch. l. c. 374 = X. Pancheri.

SALISIA, Regel, in Flora, xxxii. (1849) 179 = **Gloxinia**, L'Hérit. (Gesner.).
gloxiniaeflora, Regel, l. c. = G. maculata.
maculata, Regel, in Bot. Zeit. ix. (1851) 894 = G. maculata.
pallidiflora, Regel, l. c. = G. maculata.
suaveolens, Regel, l. c. = G. maculata.

SALIX, [Tourn.] Linn. Syst. ed I (1735); Gen. ed. I. 300 (1737). *SALICINEAE*, Benth. & Hook. f. iii. 411.
AMERINA, ARGORIPS, BIGGINA, DIAMARIPS, DIPLIMA, DIPLUSION, NECTOLIS, NECTOPIX, NECTUSION, NESTYLIX, OISODIX, PLEIARINA, PSATHERIPS, RIPSELAXIS, RIPSOCTIS, SOKOLOFIA, TELESMIA, URNECTIS, USIONIS, VETRIX, VIMEN, Rafin. Alsog. Am. 13–15 (1838).
CAPRAEA, Opiz, Seznam, 25 (1852).
GRUENERA, Opiz, l. c. 48 (1852).
KNAFIA, Opiz, l. c. 56 (1852).
LUSEKIA, Opiz, l. c. 61 (1852).
acmophylla, *Boiss. Diagn.* Ser. I. vii. 98.—Oriens; Ind. or.
acuminata, Lange, Dansk. Fl. ed. II. 658 = Smithiana.
acuminata, Mill. Gard. Dict. ed. VIII. n. 14; Pall. Fl. Ross. ii. 81 = cinerea.
acuminata, Schleich. ex Ser. Ess. Saul. 21 = grandifolia.
acuta, *Szub. Spis Drzew. Krzew. Krol. Polsk.* 136.—Polonia.
acutifolia, Borrer, ex Hook. Fl. Bor. Am. ii. 150 = boganidensis?
acutifolia, *Willd. Sp. Pl.* iv. 668 = daphnoides.
adenophylla, *Hook. Fl. Bor. Am.* ii. 146.—Labrador.
adscendens, Sm. Engl. Bot. t. 1962 = repens.
aegyptiaca, *Forsk. Fl. Aegypt. Arab.* p. lxxvi, 170 (an seq.?).—Aegypt.
aegyptiaca, Linn. Cent. Pl. i. 32; Amoen. Acad. iv. 295 = S. Safsaf.
aegyptiaca, Thunb. Prod. Pl. Cap. 6 = capensis.
affinis, Gren. & Godr. Fl. Fr. iii. 132 = Smithiana.
affinis, Schur, Enum. Pl. Transs. 619 = incana.
aglaja, Hort. ex C. Koch, Dendrol. ii. II. 501 = daphnoides.
ajanensis, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 107 = sitchensis.
alatavica, Kar. ex Stschégl. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 197 = subcordata.
alaternoides, Forbes, Salict. Woburn. 151. t. 76 = Lapponum.
alaternoides, Schleich. ex Ser. Ess. Saul. 68 = nigricans.
alba, Kern. in Oestr. Bot. Zeitschr. viii. (1858) 183 = hexandra.
alba, *Linn. Sp. Pl.* 1021.—Europ.; As. et Afr. bor.
alba, Thunb. Fl. Jap. 25 = Thunbergiana.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 463.—As. centr.

SALIX:—
albescens, Schleich. ex Ser. Ess. Saul. 68 = nigricans.
albicans, Bonjean, ex Ser. l. c. 21 = grandifolia.
albida, Schleich. ex Ser. l. c. 31 = glauca.
albiphila, Tausch, ex Wimm. Sal. Europ. 196 = Lapponum.
albovirens, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 127 = nigricans.
alnoides, Schang. & Siev. in Pall. N. Nord. Beitr. vii. 347, 349 = pyrolaefolia.
alopecuroides, Tausch, Ind. Hort. Canal. (1821); ex Opiz, Boehm. Gew. (1823) 111 = triandra.
alpestris, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 129 = ovata.
alpestris, Wulf. ex Willd. Sp. Pl. iv. 680 = S. Arbuscula.
alpigena, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 369 = retusa.
alpina, Scop. Fl. Carn. ed. II. ii. 255 = S. Arbuscula, Myrsinites.
altaica, *Lundstr. in Nov. Act. Soc. Sc. Upsal.* (1877) 36.—Sibir. altaic.
Amandae, Anderss. in DC. Prod. xvi. II. 283 = glauca.
Amaniana, Willd. Sp. Pl. iv. 663 = nigricans.
ambigua ×, Ehrh. Beitr. vi. 103.—Europ.
ambigua, Pollini, Fl. Veron. iii. 162 = cinerea.
ambigua, Pursh, Fl. Am. Sept. ii. 617 = nigra.
ambigua, Salisb. Prod. 394 = aurita.
ambigua, Schleich. ex Ser. Ess. Saul. 40 = repens.
americana, Walp. Ann. v. 747 = lanata.
amigdalina, Vill. Fl. Delphin. 109 = triandra.
amplexicaulis, Bory & Chaub. Fl. Pélop. 64. t. 36 = purpurea.
amygdalifolia, Gilib. Exercit. ii. 406 = triandra.
amygdalina, Boiss. ex Anderss. in DC. Prod. xvi. II. 211 = alba.
amygdalina, Linn. Sp. Pl. 1016 = triandra.
amygdaloides, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 114 = nigra.
Andersoniana, Sm. Engl. Bot. t. 2343 = nigricans.
androgyna, Hoppe, ex Willd. Sp. Pl. iv. 654 = triandra.
androgyna, *Roth, Tent. Fl. Germ.* ii. II. 523.—Europ.
Anglorum, Cham. in Linnaea, vi. (1831) 541 = phlebophylla.
angustata, Pursh, Fl. Am. Sept. ii. 613 = cordata.
angustifolia, Poir. in Duham. Arb. ed. Nov. iii. 125 = incana.
angustifolia, *Willd. Sp. Pl.* iv. 699.—Oriens; As. bor.
angustifolia, Wulf. in Jacq. Coll. iii. 48 = repens.
annularis, Forbes, Salict. Woburn. 41. t. 21 = babylonica.
Ansoniana, Forbes, l. c. 213. t. 107 = nigricans.
apiculata, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 470 = tetrasperma.
apoda, *Trautv. Ind. Sem. Hort. Petrop.* (1865) 37.—Reg. Caucas.
appendiculata, Vahl, in Fl. Dan. t. 1056 = glauca.
appendiculata, Vill. Hist. Pl. Dauph. iii. 775 = grandifolia.
aquatica, Lam. Fl. Fr. ii. 231 = S. Caprea?
aquatica, Ser. Ess. Saul. 20 = aurita.
aquatica, Sm. Fl. Brit. 1065; Engl. Bot. t. 1437 = cinerea.
arborescens, Hartig, Lehrb. Forstl. Pfl. 23; ex Wimm. Sal. Europ. 167, 169 = purpurea, oxycarpa.
Arbuscula, Anderss. in DC. Prod. xvi. II. 292 = fruticulosa.
Arbuscula, *Linn. Sp. Pl.* 1018.—Europ.; As. bor.
Arbuscula, Pall. Fl. Ross. ii. 83 = myrtilloides.
Arbuscula, Vill. Hist. Pl. Dauph. iii. 771 = hastata.
Arbuscula, Wahlenb. in Fl. Dan. t. 1055 = phylicifolia.
arbusculoides, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 147.—Am. arct.
arbutifolia, Pall. Fl. Ross. i. II. 79 = hastata.
arbutifolia, Ser. Ess. Saul. 44 = S. Myrsinites.
arbutifolia, *Willd. Sp. Pl.* iv. 682.—Europ.
arctica, *Pall. Fl. Ross.* i. II. 86.—Reg. arct.
arenaria, Linn. Sp. Pl. 1019 = repens.
arenaria, Willd. Sp. Pl. iv. 689 = Lapponum.
argentea, *Dum.-Cours. Bot. Cult.* ed. II. vi. 395.—Am. bor.
argentea, *Lindem. in Bull. Soc. Nat. Mosc.* xxiii. (1850) II. 528.—Rossia.
argentea, Sm. Fl. Brit. 1059 = repens.

SALIX:—
argophylla, Nutt. N. Am. Sylva, i. 71. t. 20 = longifolia.
arguta, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 32 = lasiandra.
argyrocarpa, *Anderss. l. c.* 107.—Am. bor.
asplenifolia, Hort. ex C. Koch, Dendrol. ii. I. 540 = incana.
atrocinerea, Brot. Fl. Lusit. i. 31 = cinerea.
atropurpurea, Forbes, Salict. Woburn. 284 = nigricans.
atrovirens, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
attenuata, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 246 = grandifolia.
aubonnensis, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
aurea, Salisb. Prod. 393 = alba.
auriculata, Mill. Gard. Dict. ed. VIII. n. 9 = triandra.
aurigerana, Lapeyr. Hist. Abr. Pl. Pyr. 598 = S. Caprea.
aurita, Host, Sal. 23–25 = nigricans.
aurita, *Linn. Sp. Pl.* 1019.—Europ.; As. bor.
auritoides, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 257 = purpurea.
Aurora, Laestad. ex Anderss. in Bot. Notiser (1866), 147 = fuscescens.
Austinae, *Bebb, in Bot. Calif.* ii. 88.—Calif.
australior, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 43 = fragilis.
australis, Forbes, Salict. Woburn. 205. t. 103 = cinerea.
australis, Hilsenb. & Boj. ex Fries, Mant. i. 77 = madagascariensis.
australis, Schleich. ex Spreng. Pugill. ii. 88 = nigricans.
austriaca, Host, Sal. 19. t. 64 = purpurea.
axillaris, Anderss. ex A. Rich. Tent. Fl. Abyss. ii. 275 = S. Safsaf.
Azaolana, Blanco, Fl. Filip. ed. II. 539 = tetrasperma.
babylonica, *Linn. Sp. Pl.* 1017.—Reg. Caucas.; As. bor.
badensis, Doell, Fl. Bad. 519 = nigricans.
baicalensis, Turcz. ex Anderss. in DC. Prod. xvi. II. 276 = speciosa.
balsamifera, Barratt, ex Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 125 = discolor.
Barclayi, *Anderss. l. c.*—Am. bor.
Barrattiana, *Hook. Fl. Bor. Am.* ii. 146. t. 187.—Am. bor.
Basfordiana, *Scaling, ex J. Salter, in Gard. Chron.* (1882) I. 298.—Gallia.
Baumgarteniana, Schur, Enum. Pl. Transs. 618 = purpurea.
bavarica ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxix. (1880) 117.—Europ.
berberifolia, *Pall. Reise*, iii. 759.—Sibir. altaic.
bicolor, Ehrh. Beitr. v. 162 = phylicifolia.
bicolor, Fries, Mant. i. 53 = livida.
bifida, Wulf. Fl. Nor. Phan. 780 = incana.
bifurcata, Chevall. Fl. Env. Par. ii. 357 = purpurea.
Bigelovii, Torr. in Pacif. Rail. Rep. iv. 139 = lasiolepis.
bigemmis, Hoffm. Deutschl. Fl. ed. II. 260 = daphnoides.
blanda, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 50 = alba.
Blumhardtiana, Schimp. ex Hochst. in Flora, xxiv. (1841) I. Intell. 21 = S. Safsaf.
Blyttii, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 210 = pyrenaica.
boganidensis, *Trautv. in Middend. Reise*, i. II. *Bot.* 154.—Sibir.
Bonplandiana, *H. B. & K. Nov. Gen. et Sp.* ii. 24. t. 101, 102.—Mexic.
borealis, Fries, in Bot. Notiser (1840), 193 = nigricans.
Borreri, S. F. Gray, Nat. Arr. Brit. Pl. ii. 732 = seq.
Borreriana, Sm. Engl. Fl. iv. 174 = phylicifolia.
brachycarpa, Nutt. N. Am. Sylv. i. 69 = longifolia.
brachylepis, *Franch. & Sav. Enum. Pl. Jap.* ii. 503.—Japon.
brachystachys, *Benth. Pl. Hartw.* 336.—Am. bor.
bracteata, Anderss. in DC. Prod. xvi. II. 213 = seq.
bracteosa, Turcz. ex Trautv. & Mey. in Middend. Reise (Fl. Ochot. 77) = macrolepis.
Brayi, Ledeb. Fl. Alt. iv. 289 = berberifolia.
Brayi, Nym. Consp. 670 = S. Myrsinites.
brevijulis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 387.—Sibir. baical.
brevistyla, *Richter, in Corresp. Ver. Naturk. Presb.* ii. (1863) 196.—Europ.

SALIX:—
Breweri, *Bebb, in Bot. Calif.* ii. 88.—Calif.
Brownei, *Lundstr. in Nov. Act. Soc. Sc. Upsal.* (1877) 6, 8, 12, 37.—Ins. N. Zembla.
Buergeriana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 28.—Japon.
buxifolia, Schleich. ex Ser. Ess. Saul. 54 = Lapponum.
buxifolia, Trev. ex Anderss. in DC. Prod. xvi. II. 290 = phlebophylla.
cærulea, Sm. Engl. Bot. t. 2431 = alba.
caesia, *Vill. Hist. Pl. Dauph.* iii. 768. t. 50.—Gall.; Helv.
californica, *Bebb, in Bot. Calif.* ii. 89.—Calif.
calliantha, J. Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 44 = purpurea.
callicarpaea, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 295 = arctica.
calodendron, Wimm. Sal. Europ. 187 = Smithiana.
calophylla, Wall. Cat. n. 9102 = tetrasperma?
caloptera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 26.—Japon.
calostachya, *Anderss. in Vet. Acad. Handl. Stockh.* 1850 (1851) 489.—Reg. Himal.
calyculata, *Hook. f. ex Anderss. in Journ. Linn. Soc.* iv. (1860) 55.—Reg. Himal.
campestris, Fries, Nov. Mant. ed. II. i. 68 = nigricans.
cana, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 344.—Mexic.
canaliculata, Bess. Enum. Pl. Volh. 77 = repens.
canariensis, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 159.—Marocc.; Ins. Canar.
candida, *Fluegge, ex Willd. Sp. Pl.* iv. 708.—Am. bor.
candidula, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
canescens, Fries, Mant. i. 58 = Lapponum.
canescens, Willd. Sp. Pl. iv. 687 = oleaefolia.
Canthiana ×, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 218.—Europ.
cantoniensis, *Hance, in Journ. Bot.* vi. (1868) 48.—China.
capensis, *Thunb. Fl. Cap.* i. 139.—Afr. austr.
Caprea, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 201 = cinerea.
Caprea, *Linn. Sp. Pl.* 1020.—Europ.; As. bor.
Caprea, Oed. Fl. Dan. t. 245 = lanata.
capraeformis, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 217 = viminalis.
capreoides, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 120 = brachystachys.
capreola, Kern. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 70 = S. Caprea.
caprina, Dulac, Fl. Hautes-Pyr. 148 = S. Caprea.
Capusii, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 251.—Turkestan.
cardiophylla, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 77).—As. bor. or.
carinata, Sm. Fl. Brit. 1055 = S. Arbuscula.
carinthiaca, Host, Sal. t. 113 = cinerea.
carniolica, Host, Fl. Austr. ii. 64 = purpurea.
caroliniana, Michx. Fl. Bor. Am. ii. 226 = nigra.
carpinifolia, Schleich. ex Spreng. Pugill. ii. 89 = nigricans.
caspica, Hort. ex Willd. Sp. Pl. iv. 668 = daphnoides.
caspica, Pall. Fl. Ross. i. II. 74 = volgensis.
caucasica, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 68 = silesiaca.
cerasifolia, Goldb. ex Ledeb. Fl. Ross. iii. 608 = nigricans.
cerasifolia, Schleich. ex Spreng. Pugill. ii. 88 = hastata.
Chalaf, *Forsk. Fl. Aegypt. Arab.* p. lxxvi.—Aegypt.
Chamissonis, *Anderss. in DC. Prod.* xvi. II. 290.—Am. bor.
chinensis, *Burm. f. Fl. Ind.* 211 (err. typ. 311).—China; Japon.
chinensis, Molina, Sagg. Chile, 137 = Humboldtiana.
chlorocarpa, Schur, Enum. Pl. Transs. 616 = fragilis.
chlorophana, Anderss. in DC. Prod. xvi. II. 260 = glabra.
chlorophylla, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 138.—Am. bor.
chlorostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 373.—Sibir.
chrysanthos, Vahl, in Fl. Dan. t. 1057 = lanata.
chrysophylla, Bertol. in Mem. Acad. Sc. Bologn. Ser. II. iv. (1864) 314 = Daltoniana.
ciliata, DC. Fl. Fr. 293 = pyrenaica.
cinerascens, Duby, Bot. Gall. 423 = grandifolia.
cinerascens, *Link, ex Willd. Sp. Pl.* iv. 706.—Europ. austr.

SALIX:—
cinerea, Kar. & Kir. ex Anderss. in DC. Prod. xvi. II. 269 = Smithiana.
cinerea, *Linn. Sp. Pl.* 1021.—Europ.; Oriens; As. bor.
cinerea, Schleich. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 69 = aurita.
cinerea, Sm. Fl. Brit. 1063 = daphnoides.
cinnamomea, Schleich. Cat. Pl. Helv. ed. III. 25 = aurita.
cistifolia, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
cladostema, Hayne, Dendr. Fl. Umg. Berl. 190 = aurita.
clavata, Wall. Cat. n. 3698 = Lindleyana.
clethraefolia, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
clorophana, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 178 (*chlorophana*) = glabra.
cochleata, *Dum. Fl. Belg.* 13.—Belgium.
coerulescens, Doell, Fl. Bad. 517 = livida.
coerulescens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 374 = acutifolia.
coluteoides, *Mirb. in Mém. Mus. Par.* xiv. (1827) 462.—Afr. trop.
commixta, Anderss. in DC. Prod. xvi. II. 225 = discolor.
comosa, Tausch, ex Opiz, Boehm. Gew. (1823) 111 nomen (Quid?).—Europ.
compacta ×, Anderss. in Bot. Notiser (1867), 112.—Europ.
composita, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
concolor, Host, Fl. Austr. ii. 639 = purpurea.
concolor, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
conformis, Forbes, Salict. Woburn. 47. t. 24 = Smithiana.
conformis, Schleich. ex Ser. Ess. Saul. 19 = aurita.
confusa, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 383 = lanata.
conifera, Wangh. Beytr. 123. t. 31 = phylicifolia.
connata, Bess. ex Link, Handb. ii. 453 = repens.
constricta, Guss. Enum. Pl. Inar. 403 = cinerea.
cordata, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 236.—Am. bor. et arct.
cordifolia, Banks, ex Pursh, Fl. Am. Sept. ii. 615 = cordata.
cordifolia, Hook. Fl. Bor. Am. ii. 152 = glauca.
cordifolia, Pursh, Fl. Am. Sept. ii. 611 = arctica.
coriacea, Schleich. ex Ser. Ess. Saul. 68; Forbes, Salict. Woburn. 223. t. 112 = nigricans.
coruscans, Jacq. Fl. Austr. t. 408 = S. Arbuscula.
coruscans, Willd. Sp. Pl. iv. 681 = glabra.
corylifolia, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 385 = pyrolaefolia.
cotinifolia, Sm. Engl. Bot. t. 1403 = nigricans.
Cotteti, *Kern. in Oestr. Bot. Zeitschr.* xiv. (1864) 368.—Europ.
Cotteti, Lagg. ex Anderss. in DC. Prod. xvi. II. 294 = retusa.
Coulteri, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 119 = sitchensis.
crassa, Barratt, Salic. Am. n. 7 = discolor.
crassa, *Peterm. in Flora*, xxvii. (1844) 374.—Europ.
crassifolia, Schleich. ex Ser. Ess. Saul. 68 = nigricans.
crassijulis, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. 308 = Pallasii.
crataegifolia, Bertol. in Desv. Journ. Bot. ii. (1813) 76 (1832) = glabra.
crispa, Forbes, Salict. Woburn. 83. t. 42 = nigricans.
crisposerrata, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
Croweana, Sm. Engl. Bot. t. 1115 = phylicifolia.
cuneata, Nutt. N. Am. Sylv. i. 66 = sitchensis.
cuneata, Turcz. ex Ledeb. Fl. Ross. iii. 623 = arctica.
curtiflora, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 130.—Am. bor.
curvirostris, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
cuspidata, D. Don, Prod. Fl. Nep. 59 = tetrasperma.
cuspidata, Schultz, Prod. Fl. Starg. Suppl. 47 = pentandra.
Cutleri, *Tuckerm. in Am. Journ.* Ser. I. xlv. (Oct. 1843) 36.—Am. bor.
cyanolimenaea, *Hance, in Journ. Bot.* xx. (1882) 294.—China.
cyathipoda, Anderss. ex A. Rich. Tent. Fl. Abyss. ii. 275 = S. Safsaf.

SALIX:—
cyclostigma, *Peterm. in Flora*, xxvii. (1844) 372.—Europ.
cydoniifolia, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
cyllenea, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 1188 = incana.
dahurica, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 377 = purpurea.
Daltoniana, *Anderss. in Journ. Linn. Soc.* iv. (1860) 49.—Reg. Himal.
damascena, Forbes, Salict. Woburn. 285. t. 157 = nigricans.
daphneola, Tausch, ex Anderss. in DC. Prod. xvi. II. 276 = Lapponum.
daphnoides, *Vill. Prosp.* 51; *Hist. Pl. Dauph.* iii. 765.—Europ.; As. bor.
dasycarpa, Turcz. ex Ledeb. Fl. Ross. iii. 618 = S. Myrsinites.
dasyclados, Wimm. in Flora, xxxii. (1849) 35 = Smithiana.
Davalliana, Sm. Engl. Fl. iv. 175 = phylicifolia.
Daviesii, *Boiss. Diagn.* Ser. I. vii. 98.—Persia.
dealbata, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851), 472 = acmophylla.
decipiens, Hoffm. Hist. Sal. ii. 9 = fragilis.
decumbens, Forbes, Salict. Woburn. 175. t. 88 = repens.
decumbens, *Schleich. Cat. Pl. Helv.* ed. III. 25.—Europ.
densa, Wall. Cat. n. 9103 = tetrasperma.
denticulata, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851), 481 = elegans.
denudata, Rafin. Fl. Ludov. 116 (Quid?).—Am. bor.
denudata, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
depressa, Fries, Mant. i. 57 = livida.
depressa, Hoffm. Hist. Sal. i. 39 = repens.
depressa, Linn. Fl. Suec. ed. II. 352 = lanata.
desertorum, *Richards. in Frankl. 1st Journ.* (1823) 753.—Am. bor.
dichroa, Doell, Fl Bad. 511 = purpurea.
Dicksoniana, Sm. Engl. Bot. t. 1390 = phylicifolia.
diffusa, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
digenea ×, J. Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 204.—Europ.
diplodictya, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 307 = Pallasii.
discolor, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 123 = chlorophylla.
discolor, Host, Sal. 18. tt. 60, 61 = purpurea.
discolor, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 234.—Am. bor.
discolor, Schrad. ex Willd. Sp. Pl. iv. 695 = phylicifolia.
disperma, Roxb. ex D. Don, Prod. Fl. Nep. 58 = tetrasperma.
divaricata, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 127 = nigricans.
divaricata, Pall. Fl. Ross. i. II. 80 = arctica.
divergens, *Anderss. in DC. Prod.* xvi. II. 316.—Reg. Himal.
diversifolia, Rochel, ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 116 = repens.
Doniana, *Sm. Engl. Fl.* iv. 213.—Europ.
dracunculifolia, Boiss. Diagn. Ser. I. vii. 99 = angustifolia.
dracunculoides, Anderss. ex Regel, in Act. Hort. Petrop. vi. (1879) 466 = Trautvetteriana.
Drummondiana, *Barratt, in Hook. Fl. Bor. Am.* ii. 144.—Am. bor.
dubia, Anderss. in DC. Prod. xvi. II. 314 = cinerea.
dubia, Suter, Fl. Helv. ii. 283 = S. Myrsinites.
dubia, Trautv. in. Bull. Acad. Sc. Pétersb. i. (1836) 132, nomen = nigra.
dumetorum, Suter, Fl. Helv. ii. 284 = cinerea.
dura, Schleich. Cat. Pl. Helv. ed. III. 25; Forbes, Salict. Woburn. 209. t. 105 = cardiophylla, nigricans.
Ehrhartiana, E. Mey. Chlor. Hannov. 486 = pentandra.
elaeagnifolia, Forbes, Salict. Woburn. 137. t. 69 = glauca.
elaeagnifolia, Tausch, ex Opiz, Boehm. Gew. (1823) 111 = purpurea.
elaeagnoides, Schleich. ex Ser. Ess. Saul. 91 = glauca
Elaeagnos, Scop. Fl. Carn. ed. II. ii. 257 = incana.
elbrusensis, Boiss. Diagn. Ser. I. xii. 117 = purpurea.
elegans, Bess. Enum. Pl. Volh. 77 = myrtilloides.
elegans, Host, Fl. Austr. ii. 652 = hastata.
elegans, *Wall. Cat.* n. 3699.—Reg. Himal.

SALIX:—
elegantissima, *C. Koch, Wochenschr.* xiv. (1871) 381.—Japon.
elliptica, Schleich. ex Ser. Ess. Saul. 44 = nigricans.
Erdingeri, Kern. in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 243 = daphnoides.
eriantha, Schleich. ex Link, Enum. Hort. Berol. ii. 415 = hastata.
eriocarpa, *Franch. & Sav. Enum. Pl. Jap.* ii. 502.—Japon.
eriocephala, Michx. Fl. Bor. Am. ii. 225 = discolor.
eriolepis, Less. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 171 = hastata.
eriophylla, *Anderss. in Journ. Linn. Soc.* iv. (1860) 48.—Reg. Himal.
eriostachya, *Wall. Cat.* n. 3704.—Reg. Himal.
erythrocoma, Barratt, ex Anderss. in DC. Prod. xvi. II. 206 = lasiandra.
erythrophleba, Tausch, ex Wimm. Sal. Europ. 196 = Lapponum.
estipulata, *Kit. in Linnaea*, xxxii. (1863) 356 (Quid?).—Europ.
euclasia, Goldb. ex Hoefft, Cat. Koursk, 63 = fragilis.
excelsa, J. F. Gmel. Syst. 74 = seq.?
excelsa, Tausch, ex Koch, Syn. Fl. Germ. ed. I. 643 = fragilis.
excelsior, Host, Sal. 8. t. 28 = fragilis.
exigua, Nutt. N. Am. Sylv. 75 = longifolia.
fagifolia, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
fagifolia, Willd. Sp. Pl. iv. 704 = silesiaca.
falcata, H. B. & K. Nov. Gen. et Sp. ii. 23 = Humboldtiana.
falcata, Pursh, Fl. Am. Sept. ii. 614 = nigra.
farinosa, Hartig, Nachtr. Naturg. Forstl. Cult. Pfl. Deutschl. 7 = Seringiana.
Fendleriana, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 115 = lasiandra.
Fenzliana, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 195 x. = retusa.
ferruginea, G. Anders. ex Forbes, Salict. Woburn. 255. t. 128 = Smithiana.
finmarchica, Willd. Berl. Baumz. ed. II. 441 = myrtilloides.
firma, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
fissa, Dum. Fl. Belg. 13 = pentandra.
fissa, Hoffm. Hist. Sal. i. 61 = purpurea.
flabellaris, *Anderss. in Vet. Acad. Handl. Stockh.* 1850 (1851) 497.—Reg. Himal.
flava, Schoepf. ex J. F. Gmel. Syst. 74 = humilis.
flavescens, Host, Sal. 31. t. 101 = S. Arbuscula.
flavescens, Nutt. N. Am. Sylv. i. 65 = brachystachys.
flavissima, Tausch, ex Opiz, Boehm. Gew. (1823) 111, nomen (Quid?).—Europ.
flavo-virens, Hornem. Cat. Pl. Hort. Hafn. Suppl. ii. 11 = nigra.
flexibilis, Gilib. Exercit ii. 406 = alba.
floribunda, Forbes, Salict. Woburn. 107. t. 54 = phylicifolia.
floridana, *Chapm. Fl. S. U. St.* 430.—Florida.
Flueggeana, Willd. Sp. Pl. iv. 709 = oleaefolia.
fluviatilis, Nutt. N. Am. Sylv. i. 73 = longifolia.
foetida, Schleich. ex DC. Fl. Fr. iii. 296 = S. Arbuscula.
foetida, Sm. Engl. Fl. iv. 208 = repens.
foliolosa, *Willd. Sp. Pl.* iv. 685.—Europ.
foliosa, Afzel. ex Koch, Syn. Fl. Germ. ed. I. 653 = livida.
Forbesiana, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 117 = repens.
Forbesiana, Loud. Arb. Brit. iii. 1581 = phylicifolia.
Forbyana, Sm. Fl. Brit. 1041 = purpurea.
formosa, Willd. Sp. Pl. iv. 680 = S. Arbuscula.
Forsteriana, Sm. Engl. Bot. t. 2344 = nigricans.
fragilior, Host, Sal. 6. t. 22 = fragilis.
fragilis, Forsk. Fl. Aegypt. Arab. p. lxxvi (Quid?).—Aegypt.
fragilis, Host, Sal. 5. tt. 18, 19 = pentandra.
fragilis, *Linn. Sp. Pl.* 1017.—Europ.; As. bor.
fragilissima, Host, Sal. 6. tt. 22, 23 = fragilis.
fragillima, Schur, Enum. Pl. Transs. 616 = fragilis.
fragrans, Salisb. Prod. 393 = pentandra.
Frangula, Willd. ex Schleichl. Cat. Pl. Helv. ed. III. 25 = nigricans.
Friesiana, Anderss. in Bot. Notiser (1866), 156 = repens.
Friesii, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 183 = pentandra.
fruticosa, Doell, Fl. Bad. 515 = Smithiana.
fruticulosa, *Anderss. in Journ. Linn. Soc.* iv. (1860) 53.—Reg. Himal.

SALIX:—
fruticulosa, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 368 = S. Arbuscula.
fulcrata, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 139.—Am. bor.
fumosa, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 384 = arctica.
furcata, Anderss. in DC. Prod. xvi. II. 291 = fruticulosa.
furcata, Tausch, ex Wimm. Sal. Europ. 173 = purpurea.
fusca, Fries, Mant. i. 69 = myrtilloides.
fusca, Hook. Fl. Bor. Am. ii. 151 = argyrocarpa.
fusca, Jacq. Fl. Austr. t. 409 = S. Myrsinites.
fusca, Linn. Sp. Pl. 1020 = repens.
fuscata, Pursh, Fl. Am. Sept. ii. 612 = petiolaris.
fuscata, Tausch, ex Anderss. in DC. Prod. xvi. II. 307 = purpurea.
fuscescens, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 97.—Kamtschat.
fuscior, *Anderss. l. c.* 163.—Am. bor.
gariepina, Burch. Trav. i. 317 = capensis.
geminata, Forbes, Salict. Woburn. 257. t. 129 = Smithiana.
gemmata, Wimm. ex Anderss. in DC. Prod. xvi. II. 304 = Seringeana.
Geyeriana, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 122.—Am. bor. occ.
glabra, Ledeb. Fl. Ross. iii. 623 = caesia.
glabra, *Scop. Fl. Carn.* ed. II. ii. 255.—Europ.
glabrata, Herb. Heyne, ex Wall. Cat. n. 7977 = Excoecaria crenulata.
glabrata, *Schleich. Cat Pl. Helv.* ed. IV. 25.—Europ.
glabrescens, Lindl. in Wall. Cat. n. 3706 = tetrasperma?
glabricarpa, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
glacialis, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 131.—Am. bor.
glareosa, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
glauca, *Linn. Sp. Pl.* 1019.—Europ.; As. et Am. bor.
glauca, Sm. Engl. Bot. t. 1810 = Lapponum.
glauca, Willd. Berl. Baumz. 388 = S. Arbuscula.
glaucescens, Host, Sal. 23 = nigricans.
glaucescens, Moench, Meth. Suppl. 116 = S. Arbuscula.
glaucoides, Anderss. in DC. Prod. xvi. II. 284 = S. Myrsinites.
glaucophylla, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 474 = acmophylla.
glaucophylla, Bess. Enum. Pl. Volh. 38 = Lapponum.
glaucophylla, Schleich. ex Ser. Ess. Saul. 68 = nigricans.
glaucops, Anderss. in DC. Prod. xvi. II. 281 = glauca.
Gmelini, Pall. Fl. Ross. i. II. 77 = viminalis.
Gmelini, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 390 = Lapponum.
Gmeliniana, Turcz. l. c. = praec.
gnaphaloides, Schleich. ex Anderss. in DC. Prod. xvi. II. 276 = Lapponum.
Goepperti, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 68 = hastata.
gongylocarpa, Shuttlew. ex Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 114 = nigra.
Govaniana, Wall. Cat. n. 3699 = elegans.
gracilenta, Tausch, in Flora, xvii. (1834) 447 = fragilis.
gracilis, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 127 = petiolaris.
gracilistyla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 26.—Japon.
Grahami, Borrer, ex Baker, in Journ. Bot. v. (1867) 157. t. 66 = retusa.
grandifolia, Fries, ex Nym. Consp. 667 = S. Caprea.
grandifolia, *Ser. Ess. Saul.* 20.—Europ.
grisea, Hook. Kew Journ. vii. (1855) 372 = seq.?
grisea, *Marsh. ex A. Gray, Man. Bot. N. U. St.* ed. II. 417.—Am. bor.
grisea, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid?).—Am. bor.?
grisea, Wall. Cat. n. 3700 G = Wallichiana.
grisea, Willd. Sp. Pl. iv. 699 = sericea.
grisescens ×, Anderss. in Bot. Notiser (1867), 112.—Europ.
grisonensis, Forbes, Salict. Woburn. 197. t. 99 = nigricans.
grisophylla, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
groenlandica, *Lundstr. in Nov. Act. Soc. Sc. Upsal.* (1877) n. xiv. 36.—Ins. N. Zembla.
gryonensis, Schleich. ex Ser. Ess. Saul. 67 = nigricans.

SALIX:—
guseniensis, Wimm. Sal. Europ. 231 = cinerea.
Halleri, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
Hartmanniana, Anderss. in DC. Prod. xvi. II. 275 = lanata.
Hartwegi, Benth. Pl. Hartw. 46 = lasiolepis.
hastata, Hoppe, ex Vill. Hist. Pl. Dauph. iii. 774. t. 50 = nigricans.
hastata, *Linn. Sp. Pl.* 1017.—Europ.; As. bor.
hastata, Thuill. Fl. Par. ed. II. 512 = triandra.
Heerii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 104.—Europ.
Hegetschweileri, Heer, in Hegetschw. Fl. Schw. 963 = phylicifolia.
Heimerli ×, H. Braun, in Oestr. Bot. Zeitschr. xxxi. (1881) 107.—Europ.
heliciflora, Tausch, ex Anderss. in DC. Prod. xvi. II. 308 = purpurea.
Helix, Linn. Sp. Pl. 1017 = purpurea.
helvetica, Forbes, Salict. Woburn. 287 = nigricans.
helvetica, Vill. Hist. Pl. Dauph. iii. 783 = Lapponum.
herbacea, *Linn. Sp. Pl.* 1018.—Reg. bor. et arct.
herbacea, Schrenk, Reise, ii. 523 = rotundifolia.
herbacea, Soland. in Phipps, Voy. 202; Hook. in Parry, Narr. 2nd Voy. (1825) 404 (1827) = polaris.
hermaphroditica, Linn. Sp. Pl. 1015 = pentandra.
heterophylla, Bray, in Denkschr. Bot. Ges. Regensb. i. (1815) 51 = alba.
heterophylla, Host, Sal. 26. t. 87 = aurita.
heterophylla, Schult. Fl. Starg. Suppl. 51 = repens.
heterophylla, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 141 = hastata.
hexandra, *Ehrh. Beitr.* vii. 138.—Austria.
himalensis, Klotzsch, ex Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 482 = elegans.
Hindsiana, Benth. Pl. Hartw. 335 = longifolia.
hippophaëfolia, Ledeb. Fl. Ross. iii. 601 = purpurea.
hippophaëfolia, Thuill. Fl. Par. ed. II. 514 = mollissima.
hippophaëfolia, Wimm. & Grab. Fl. Siles. ii. 363 = multiformis.
hircina ×, Kern. in Verh. Zool.-Bot. Ges. Wien, xiv. (1864) 99 = incana.
hirsuta, Thunb. Prod. Pl. Cap. 6 = capensis.
hirta, Sm. Engl. Bot. t. 1404 = nigricans.
hirtula, Anderss. in Bot. Notiser (1866), 147 = myrtilloides.
Hoffmanniana, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 568 = cinerea.
Hoffmanniana, Hook. & Arn. Bot. Beech. Voy. 159 = lasiandra.
Hoffmanniana, Sm. Engl. Bot. t. 2620 = triandra.
Hoffmanniana, Tausch, ex Wimm. Sal. Europ. 173 = purpurea.
holosericea, Ser. in Flora, xiii. (1830) 1. Erg. 41 = Seringeana.
holosericea, Willd. Sp. Pl. iv. 708 = Smithiana.
Hookeriana, *Barratt, in Hook. Fl. Bor. Am.* ii. 145. t. 180.—Am. bor.
Hoppeana, Willd. Sp. Pl. iv. 654; Sturm, Deutschl. Fl. Heft 25 = triandra.
Horsfieldiana, Miq. Fl. Ind. Bat. i. II. 461 = tetrasperma.
Hosteana, Willd. in Ges. Naturf. Fr. Berl. Mag. iv. (1810) 53 = S. Arbuscula?
Hostii ×, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 213 = viminalis.
Houstoniana, Pursh, Fl. Am. Sept. ii. 614 = nigra.
Huartina, Sm. Engl. Fl. iv. 203 = Lapponum.
Hugelii, Wimm. ex Anderss. in Journ. Linn. Soc. iv. (1860) 41 = tetrasperma.
Huguenini ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 117; xxv. (1882) 104.—Europ.
Humboldtiana, *Willd. Sp. Pl.* iv. 657.—Am. austr.
humilis, *Marsh. Arbust.* 140.—Am. bor.
humilis, Willd. Berl. Baumz. ed. II. 439 = phylicifolia.
humillima, *Anderss. in DC. Prod.* xvi. II. 248.—Am. arct.
hungarica, *Kern. in Oestr. Bot. Zeitschr.* xiv. (1864) 9.—Europ.
Huteri, Kern. in Oestr. Bot. Zeitschr. xvi. (1866) 370 = Lapponum.
hybrida, Hoffm. Deutschl. Fl. ed. II. ii. 263 = nigricans.
hybrida, *Opiz, in Flora*, v. (1822) 270.—Europ.

SALIX:—
hybrida, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid?).—Am. bor.?
hybrida, *Scop. Fl. Carn.* ed. II. ii. 253.—Europ.
hybrida, Vill. Hist. Pl. Dauph. iii. 778 = S. Caprea.
hyperborea, Anderss. Sal. Lapp. 51. f. 12 = hastata.
hypophaefolia, Forbes, Salict. Woburn. 13, sphalm. = *hippophaëfolia*.
ichnostachya, *Wall. Cat.* n. 3704; *Lindl. ex Wight, Ic.* t. 1953.—Ind. or.
ilicifolia, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 127 = nigricans.
ilicifolia, Willd. ex Cham. in Linnaea, vi. (1831) 542 = berberifolia.
iliensis, *Regel, in Act. Hort. Petrop.* vi. (1879) 464. —As. centr.
incana, Michx. Fl. Bor. Am. ii. 225 = candida.
incana, Schleich. ex Ser. Ess. Saul. 44 = nigricans.
incana, *Schrank, Baier. Fl.* i. 230.—Europ.
incanescens, Forbes, Salict. Woburn. 239. t. 120 = cinerea.
incerta, Lapeyr. Hist. Abr. Pl. Pyr. 594 = triandra.
incubacea, Linn. Sp. Pl. 1020 = repens.
indigena, Forsk. Fl. Aegypt. Arab. p. lxxvi = S. Safsaf.
insignis, *Anderss. in Journ. Linn. Soc.* iv. (1860) 47.—Reg. Himal.
integra, Thunb. Fl. Jap. 24 = repens.
intermedia, Host, Sal. 17. tt. 56, 57 = incana.
intermedia, Thomas, ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 145 = S. Arbuscula.
irrorata, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 117.—Mexic.
iserana, J. & C. Presl, Fl. Čech. 204 = aurita.
Jacquiniana, Willd. Sp. Pl. iv. 692 = S. Myrsinites.
Jacquinii, Host, Syn. 529 = praec.
japonica, Thunb. Fl. Jap. 24. t. 31 = babylonica.
jaspidea, Hort. ex C. Koch, Dendrol. ii. II. 501 = daphnoides.
javanica, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 3 = tetrasperma.
julacea, Anderss. l. c. 1850 (1851), 476 = S. Caprea.
Junghuhniana, *Anderss. ex Miq. Illustr. Fl. Ind. Arch.* 13.—Java.
juratensis, Schleich. ex Ser. Ess. Saul. 44 = nigricans.
kamaunensis, Lindl. in Wall. Cat. n. 3701 = elegans.
kanderiana, Ser. Ess. Saul. 37 = Seringeana.
Karelini, *Turcz. ex Stschégl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 196, 393.—As. centr.
kassanogluensis, Kotschy, ex Anderss. in DC. Prod. xvi. II. 212 = alba.
Kawajanagi, Siebold, in Verh. Batav. Gen. xii. (1830) 24 = Buergeriana, gracilistyla.
Kerneri, Erding, in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 78 = incana.
kilnensis, Anderss. in DC. Prod. xvi. II. 320 = myrtilloides.
Kirilowiana, *Stschégl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 198.—Sibir. altaic.
Kitaibeliana, Willd. Sp. Pl. iv. 683 = retusa.
Kochiana, Hartig, Naturg. Forstl. Cult. 569 = purpurea.
Kochiana, *Trautv. in Bull. Acad. Sc. Pétersb.* i. (1836) 132, nomen; *et in Mém. Sav. Etr. Pétersb.* iii. (1837) 632. t. 1.—Sibir. altaic.
Koernickei, Anderss. in DC. Prod. xvi. II. 263 = daphnoides.
koreensis, *Anderss. l. c.* 271.—Korea.
Kovatsii, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 190 = triandra.
kossoviensis, *Bess. in Mem. Pharm. Ac. Viln.* ii. 649. —Lithuan.
Kraettliana, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 117; xxv. (1882) 105.—Europ.
Krausei, Anderss. in Vet. Akad. Handl. Stockh. vi. (1867) n. 1. 29 = triandra.
kumaonensis, Auct. = *kamaunensis*.
lactea, Bray, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 43. t. 2 = Lapponum.
lacustris, Schleich. Cat. Pl. Helv. ed. III. 25; Forbes, Salict. Woburn. 231. t. 116 = nigricans.
Laestadiana, Hartm. Skand. Fl. ed. III. 237 = Lapponum.
laeta, Schultz, Prod. Fl. Starg. Suppl. i. 51 = repens.
laevigata, *Bebb, in Am. Natural.* viii. (1874) 202.—Calif.
laevis, Hook. Brit. Fl. ed. I. 482 = S. Myrsinites.
Laggeri, Wimm. in Flora, xxxvii. (1854) 162 = grandifolia.

SALIX:—
Lambertiana, Sm. Fl. Brit. 1041 = purpurea.
lanata, Lightf. Fl. Scot. ii. 602 = S. Caprea.
lanata, *Linn. Sp. Pl.* 1019.—Europ. et As. bor.
lanata, Vill. Hist. Pl. Dauph. iii. 777 = cinerea.
lanceolata, Fries, Novit. ed. II. 283 = Smithiana.
lanceolata, Ser. Ess. Saul. 37 = Seringeana.
lanceolata, Sm. Engl. Bot. t. 1436 = triandra.
lancifolia, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 34 = lasiandra.
lancifolia, Doell, Fl. Bad. 112 = Smithiana.
Lantana, Hartig, Naturg. Forstl. Cult. Pfl. 410. t. 41 = *ambigua*.
lanuginosa, Pall. Fl. Ross. ii. 83 = lanata.
lapponica, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.
Lapponum, *Linn. Sp. Pl.* 1019.—Europ.; As. bor.
Lapponum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 389 = speciosa.
lasiandra, *Benth. Pl. Hartw.* 335.—Am. bor. occ.
lasiocarpa, Wimm. ex Anderss. in DC. Prod. xvi. II. 218 = silesiaca.
lasiolepis, *Benth. Pl. Hartw.* 335.—Calif.
lasiostyla, *Peterm. in Flora*, xxvii. (1844) 373.—Europ.
latifolia, Forbes, Salict. Woburn. 235. t. 118 = nigricans.
latifolia, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 344.—Mexic.
latiuscula, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 165.—Am. bor.
laurifolia, Wesmael, in Bull. Congr. Bot. Brux. (1864) 280 = pentandra.
laurina, Sm. in Trans. Linn. Soc. vi. (1802) 122 = phylicifolia.
laurinea, Loisel. Fl. Gall. ed. II. ii. 343 = phylicifolia.
lavandulaefolia, Lapeyr. Hist. Abr. Pl. Pyr. 601 = incana.
laxa, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
laxiflora, Borrer, in Engl. Bot. Suppl. ii. t. 2749 = phylicifolia.
laxiflora, Kern. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 176 = glabra.
Ledebouriana, Trautv. in Bull. Sc. Acad. Pétersb. i. (1836) 132, nomen; et in Mém. Sav. Etr. Pétersb. iii. (1837) 631 = purpurea.
Lemana, Schleich. Cat. Pl. Helv. ed. III. 25 = nigricans.
Lemmoni, *Bebb, in Bot. Calif.* ii. 88.—Calif.
lenta, Fries, Fl. Suec. Mant. i. 78 = tetrasperma.
leptoclados, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 144.—Sibir.
leucantha, Tausch, ex Opiz, Boehm. Gew. (1823) 111, nomen (Quid?).—Europ.
leucophylla, Willd. Berl. Baumz. ed. II. 444 = Lapponum.
libanotica, Boiss. Fl. Orient. iv. 1185 = alba.
ligustrina, Host, Sal. t. 15–16 = triandra.
ligustrina, Michx. f. Hist. Arb. Am. iii. 226. t. 5 = nigra.
ligustroides, Schleich. ex Ser. Ess. Saul. 65 = hastata.
limnogena, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) 64 = aurita?
limnogena, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 187 = grandifolia.
limosa, Trautv. ex Anderss. in DC. Prod. xvi. II. 277 = speciosa.
limosa, Wahlenb. Fl. Lapp. 265 = Lapponum.
Lindleyana, *Wall. Cat.* n. 3697.—Reg. Himal.
linearis, Forbes, Salict. Woburn. 177. t. 89 = incana.
linearis, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 379 = viminalis.
lingulata, *Anderss. in DC. Prod.* xvi. II. 281.—Am. bor.
lithuania, Bess. ex Ledeb. Fl. Ross. iii. 608 = nigricans.
littoralis, Host, Sal. 15. t. 52 = repens.
livescens, Doell, Fl. Bad. 520 = aurita.
livescens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 386 = repens.
livida, *Wahlenb. Fl. Lapp.* 272. t. 16. f. 6.—Europ.
livida, Wimm. in Denkschr. Bresl. Ges. (1853) 147 = aurita.
longiflora, *Wall. ex Anderss. in Journ. Linn. Soc.* iv. (1860) 50.—Reg. Himal.
longifolia, Lam. Fl. Fr. ii. 232 = viminalis.

SALIX:—
longifolia, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 238.—Am. bor.
longifolia, Schleich. Cat. Pl. Helv. ed. III. 26 = Seringeana.
longipes, Hook. f. & Thoms. ex Anderss. in Journ. Linn. Soc. iv. (1860) 56 = S. Serpyllum.
longipes, Shuttlew. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 114 = nigra.
longirostris, Michx. Fl. Bor. Am. ii. 226 = tristis.
loniceraefolia, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 375 = Kochiana.
lucida, Jacquem. ex Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 497 = flabellaris.
lucida, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 239.—Am. bor.
ludoviciana, Rafin. Fl. Ludov. 115 (Quid?).—Am. bor.
Ludwigii, Bonj. ex Bertol. Fl. Ital. x. 313 = nigricans.
Ludwigii, Schkuhr, Handb. iv. 227. t. 317 = silesiaca.
lutea, Nutt. N. Am. Sylv. i. 63. t. 19 = cordata.
lutescens, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 253 = cinerea.
lutescens, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
Lyonii, *Forbes, Salict. Woburn.* 19. t. 10.—Helvet.
macilenta, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 141.—Kamtschat.
mackenziana, *Barratt, ex Hook. Fl. Bor. Am.* i. 149. —Am. arct.
macrocarpa, Anderss. in Vet. Acad. Handl. Sto kh. 1850 (1851) 479 = phylicifolia.
macrocarpa, *Nutt. N. Am. Sylv.* i. 83.—Am. bor. occ.
macrocarpa, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 292 = glauca.
macrolepis; *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 371.—As. bor.
macronectaria, *Peterm. in Flora*, xxvii. (1844) 373.—German.
macrophylla, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 64 = grandifolia.
macrophylla ×, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 247 = S. Caprea.
macropoda, *Stschégl. in Bull. Soc. Nat. Mosc.* (1854) 1. 197.—Sibir. altaic.
macrorhyncha, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 155 = phylicifolia.
macrostachya, Nutt. N. Am. Sylv. i. 72 = longifolia.
macrostipulacea, Forbes, Salict. Woburn. 259. t. 130 = nigricans.
macrostipularis, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
madagascariensis, *Boj. ex Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 15.—Madag.
magellanica, Poir. Encyc. Suppl. iv. 66 = Humboldtiana.
majalis, Wahlenb. Fl. Lapp. 270 = phylicifolia.
malifolia, Bess. Fl. Galic. ii. 313 = livida.
malifolia, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
malifolia, Sm. Fl. Brit. 1053 = hastata.
marginata, Wimm. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 21 = nigra.
marinifolia, Willd. Sp. Pl. iv. 697 = repens.
marrubiifolia, Tausch, ex Anderss. in DC. Prod. xvi. II. 276 = Lapponum.
Martiana, Leyb. in Mart. Fl. Bras. iv. 1. 228. t. 72 = Humboldtiana.
Mauckschii, Hartig, Naturg. Forstl. Cult. 396 = silesiaca.
mauternensis, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 261 = purpurea.
Medemii, Boiss. Diagn. Ser. I. vii. 100 = cinerea.
melanolepis, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 293 = glauca.
melanopsis, Nutt. N. Am. Sylv. i. 78 = amygdaloides.
membranacea, Thuill. Fl. Par. ed. II. 515 = purpurea.
menthaefolia, Host, Sal. 23–25 = nigricans.
Mesnyi, *Hance, in Journ. Bot.* xx. (1882) 38.—China.
mespilifolia, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
Meyeriana, Rostl. in Willd. Berl. Baumz. ed. II. 427; Forbes, Salict. Woburn. 65. t. 33; Borrer, in Hook. Brit. Fl. ed. I. 417; ed. III. 421 = pentandra.
micans, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 49 = alba.
Micheliana, Forbes. Salict. Woburn. t. 125 = Smithiana.
microdonta, Schleich. ex Ser. Ess. Saul. 67 = nigricans.

SALIX:—
microphylla, Cham. & Schlecht. in Linnaea, vi. (1831) 354 = taxifolia.
microphylla, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 37.—Hispan.
microstachya, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 377 = angustifolia.
Mielichhoferi, Saut. in Flora, xxxii. (1849) 662 = glabra.
Mielichoferi, Tausch, ex Anderss. in DC. Prod. xvi. II. 290 = S. Myrsinites.
minutiflora, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 394 = caesia.
Miquelii, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 166.—Japon.
mirabilis, Host, Sal. 13. t. 46 = purpurea.
mollis, Schleich. ex Ser. Ess. Saul. 44 = aurita.
mollissima, *Ehrh. Beitr.* vi. 101.—Europ.
mollissima, Sm. Fl. Brit. 1070 = Smithiana.
mollissima, Wahlenb. Fl. Carp. 317 = purpurea.
monadelpha, Koch, ex Schur, Enum. Pl. Transs. 618 = purpurea.
monandra, Hoffm. Hist. Sal. i. 18 = purpurea.
monandra, Host, Sal. 22. t. 71, 72 = grandifolia.
monica, *Bebb, in Bot. Calif.* ii. 90.—Calif.
monspeliensis, Forbes, Salict. Woburn. 59. t. 30 = fragilis.
montana, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
mucronata, Thunb. Prod. Pl. Cap. 6 = capensis.
Muehlenbergiana, Barratt, ex Torr. Fl. N. York, ii. 205 = humilis.
Muehlenbergiana, Willd. Sp. Pl. iv. 692 = tristis.
multiformis, *Doell, Fl. Bad.* 506.—Europ.
multiglandulosa, Wimm. Sal. Europ. 196 = Lapponum.
multinervis, Doell, Fl. Bad. 516 = cinerea.
murina, *Schleich. Cat. Pl. Helv.* ed. III. 26.—Europ.
mutabilis, Forbes, Salict. Woburn. 288. t. 160 = nigricans.
mutabilis, *Host, Sal.* tt. 42, 43.—Europ.
mutabilis, Schleich. Cat. Pl. Helv. ed. III. 26 = *ambigua*.
myricaefolia, Anderss. l. c. 1850 (1851), 483 = divergens.
myricoides, Muhl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 235 = cordata.
myrsinifolia, Salisb. Prod. 394 = nigricans.
Myrsinites, Cham. in Linnaea, vi. (1831) 540 = Chamissonis.
Myrsinites, Hoffm. Hist. Sal. 71 = nigricans.
Myrsinites, *Linn. Sp. Pl.* 1018.—Europ.
myrsinitoides, Fries, ex Anderss. in DC. Prod. xvi. II. 290 = nigricans.
myrtillacea, *Anderss. in Journ. Linn. Soc.* iv. (1860) 51.—Reg. Himal.
myrtillifolia, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 132 = S. Novae-Angliae.
myrtilloides, Cham. in Linnaea, vi. (1831) 539 = fuscescens.
myrtilloides, *Linn. Sp. Pl.* 1018.—Reg. bor.
myrtilloides, Vill. Hist. Pl. Dauph. iii. 770 = S. Arbuscula.
myrtilloides, Willd. Sp. Pl. iv. 686 = caesia.
myrtoides, Doell, Fl. Bad. 522 = livida.
myrtoides, Fries, Mant. i. 70; Hartm. Skand. Fl. ed. IV. 322 = myrtilloides.
Myurus, Wimm. ex Anderss. in Journ. Linn. Soc. iv. (1860) 41 = tetrasperma.
nana, Schleich. Cat. Pl. Helv. ed. III. 26 = repens?
Napoleona, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 616 = babylonica.
Napoleonis, F. Schultz, Arch. Fl. 239 = praec.
natalensis, Wimm. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 14 = capensis.
neglecta, Gorski, ex Anderss. l. c. 103 = myrtilloides.
Neilreichii, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 258 = purpurea.
Neisseana ×, Kern. l. c. 217.—Europ.
nepetaefolia, *F. & C. Presl, Fl. Čech.* 205.—Europ.
neriifolia, Schleich. Cat. Pl. Helv. ed. III. 26 (Quid).—Helvet.
nervosa, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
nevadensis, *S. Wats. in Am. Natural.* vii. (1873) 302.—Am. bor. occ.
nigra, *Marsh. Arbust.* 293.—Am. bor.
nigrescens, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
nigricans, *Sm. in Trans. Linn. Soc.* vi. (1802) 120.—Europ.; As. bor.

SALIX:—
nilagirica, Miq. Anal. Bot. Ind. iii. 22 = tetrasperma.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* ii. 502.—Japon.
nitens, G. Anders. ex Sm. Engl. Fl. iv. 174 = phylicifolia.
nitens, Gilib. Exercit. ii. 408 = Lapponum.
nitens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 380 = viminalis.
nitida, J. F. Gmel. Syst. 74 (Quid?).—Aegypt.?
nivalis, Hook. Fl. Bor. Am. ii. 152 = reticulata.
nivalis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 68 = herbacea.
nivea, Ser. Ess. Saul. 51 = Lapponum.
nivea, Sm. ex Anderss. in DC. Prod. xvi. II. 278 = candida.
nobilis, Fries, Novit. i. 78 = tetrasperma.
Normanni, Anderss. in DC. Prod. xvi. II. 288 = S. Myrsinites.
notha, Anderss. in Bot. Notiser (1867), 114 = S. Myrsinites.
Novae-Angliae, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 160.—Am. bor.
nummularia, *Anderss. in DC. Prod.* xvi. II. 298.—Sibir. arct.
nummulariaefolia, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 127 = nigricans.
obcordata, *Anderss. in DC. Prod.* xvi. II. 291.—Am. bor.
oblongifolia, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 81).—As. bor.
obovata, Wall. Cat. n. 3698 = flabellaris.
obscura, *Anderss. in DC. Prod.* xvi. II. 269.—Reg. Himal.
obtusa, *Richter, ex Corresp. Ver. Naturk. Presb.* ii. (1863) 195.—Europ.
obtusa, *Schleich. ex Link, Enum. Hort. Berol.* ii. 415.—Helvet.
obtusifolia, *Willd. Sp. Pl.* iv. 705.—Lappon.
occidentalis, *Bosc, ex Koch, Comment.* 16.—Ind. occ.
octandra, Delile, ex Aitch. Cat. Punjab Pl. 140 = acmophylla.
octandra, Sieber, ex A. Rich. Tent. Fl. Abyss. ii. 276 = S. Safsaf.
Oldhamiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 25.—Japon.
oleaefolia, *Vill. Hist. Pl. Dauph.* iii. 784.—Europ.
oleifolia, Host, ex Wimm. Sal. Europ. 161 = purpurea.
oleifolia, Ser. Ess. Saul. 11 = incana.
oleifolia, Sm. Fl. Brit. 1065 = cinerea.
oleifolia, Wimm. Herb. Sal. 66; ex Anderss. in DC. Prod. xvi. II. 305 = oleaefolia.
Olgae, *Regel, in Act. Hort. Petrop.* vi. (1879) 465, nomen; *Pl. Nov. Fedsch.* 79.—Turkestan.
olivacea, Thuill. Fl. Par. ed. II. 515 = purpurea.
onusta, Bess. Enum. Pl. Volh. 78 = myrtilloides.
onychiophylla, Anderss. ex Nym. Consp. 671 = herbacea.
oppositifolia, Host, Sal. 11. t. 38 = purpurea.
orbicularis, *Anderss. in DC. Prod.* xvi. II. 300.—Am. bor.
oreophila, *Hook. f. ex Anderss. in Journ. Linn. Soc.* iv. (1860) 57.—Reg. Himal.
ovalifolia, *Trautv. in Nouv. Mém. Soc. Nat. Mosc.* ii. (1832) 306.—Am. arct.
ovata, *Ser. Ess. Saul.* 92.—Europ.
ovata, Spreng. Pugill. i. 62 = nigricans.
oxycarpa, *Anderss. in Journ. Linn. Soc.* iv. (1860) 45.—Reg. Himal.
oxyphylla, H. B. & K. Nov. Gen. et Sp. ii. 24 = Humboldtiana.
padifolia, *Anderss. in Mem. Am. Acad.* N. S. vi. (1858–59) 451.—Japon.
Pallasii, *Anderss. in DC. Prod.* xvi. II. 285.—Kamtschat.
pallescens, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 33.—Am. bor.
pallescens, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
pallida, Forbes, Salict. Woburn. 191. t. 96 = oleaefolia.
pallida, H. B. & K. Nov. Gen. et Sp. ii. 25 = Bonplandiana.
pallida, Ledeb. Fl. Alt. iv. 261 = purpurea.
pallida, Salisb. Prod. 394 = alba.
pallida, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
paludosa, Hartm. Skand. Fl. ed. III. 236 = myrtilloides.

SALIX:—
paludosa, Link, Enum. Hort. Berol. ii. 419 = aurita.
paludosa, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
palustris, Host, Sal. 7. t. 24 = fragilis.
Pameachiana, Barratt, Salic. Am. n. 16 = alba.
pannosa, Schleich. Cat. Pl. Helv. ed. IV. 31; Forbes, Salict. Woburn. 245. t. 122 = nigricans.
paradoxa, H. B. & K. Nov. Gen. et Sp. ii. 25 = discolor.
parietariifolia, Host, Sal. t. 23–25 = nigricans.
parviflora, Host, l. c. 14. t. 49 = Doniana.
parvifolia, Anderss. in DC. Prod. xvi. II. 283 = glauca.
parvifolia, Sm. Engl. Bot. t. 1961 = repens.
patens, Forbes, Salict. Woburn. 77. t. 39 = phylicifolia.
patula, Kern. ex Anderss. in DC. Prod. xvi. II. 305 = oleaefolia.
patula, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
patula, Ser. l. c. 11 = incana.
Patzeana, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 117, 124 = repens.
Patzei, Wimm. ex Anderss. l. c., 93 = aurita.
pedicellata, *Desf. Fl. Atlant.* ii. 362.—Reg. Mediterr. occ.
pedicillaris, Pursh, Fl. Am. Sept. ii. 611 = myrtilloides.
pellita, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 139 = chlorophylla.
Peloritana, Prestandr. ex Tineo, Pl. Rar. Sicil. 31 = oxycarpa.
pendula, Moench, Meth. 336; Salisb. Prod. 393 = babylonica.
pendula, Ser. Ess. Saul. 79 = fragilis.
pendulina, Wender. in Schrift. Nat. Ges. Marb. ii. (1831) 235 = purpurea.
pensylvanica, Hort. ex Pursh, Fl. Am. Sept. ii. 611 = myrtilloides.
pensylvanica, Hort. l. c. 616 = petiolaris.
pentandra, Kern. in Oestr. Bot. Zeitschr. viii. (1858) 183 = hexandra.
pentandra, *Linn. Sp. Pl.* 1016.—Europ.; As. bor.
pentandra, Nutt. N. Am. Sylv. i. 61 = lucida.
pentandra, Thuill, Fl. Par. ed. II. 513 = triandra.
pentandra, Walt. Fl. Carol. 243 = nigra.
persica, *Boiss. Diagn.* Ser. I. vii. 99.—Persia.
persicaefolia, Tausch, ex Opiz, Boehm. Gew. (1823) 111 = fragilis?
persicifolia, Schleich. Cat. Pl. Helv. ed. II. 80 = fragilis.
petiolaris, *Sm. in Trans. Linn. Soc.* vi. (1802) 122.—Am. bor.
petraea, [G. Anders. in] Forbes, Salict. Woburn. 193. t. 97 = nigricans.
petrosa, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
phaeophylla, Anderss. in Bot. Notiser (1867), 116 = S. Myrsinites.
philicifolia, Crantz, Inst. i. 176 = phylicifolia.
phillyreifolia, Borrer, in Engl. Bot. Suppl. t. 2660 = phylicifolia.
phlebophylla, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 132.—Am. bor.
phlomoides, Bieb. Fl. Taur. Cauc. ii. 415; iii. 628 = cinerea.
phylicaefolia, Wulf. in Jacq. Coll. ii. 139 = phylicifolia.
phylicifolia, J. Carey, ex A. Gray, Man. Bot. N. U. St. ed. II. 416 = chlorophylla.
phylicifolia, Cham. in Linnaea, vi. (1831) 542 = macilenta.
phylicifolia, *Linn. Sp. Pl.* 1016.—Europ.
phylicifolia, Wahlenb. Fl. Lapp. 270 = nigricans.
phylicifolia, Willd. Enum. Hort. Berol. 1003 = glabra.
phylicoides, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 123.—Am. bor. occ.
Pierotii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 27.—Japon.
pilosa, Bray, ex Ledeb. Fl. Ross. iii. 608 = nigricans.
pilosa, Schleich. ex Ser. Ess. Saul. 47 = S. Myrsinites.
pinetorum, Czern. ex Trautv. in Act. Hort. Petrop. ix. (1884) 183, nomen (Quid?).—Rossia.
Pitcheriana, Baratt, Salic. Am. n. 22 = nigra?
planifolia, *Pursh, Fl. Am. Sept.* ii. 611.—Am. bor.; Labrador.
plicata, Fries, Novit. i. 66 = repens.
podophylla, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) 142.—Sibir.
Pokornyi, Kern. in Verh. Zool. Bot. Ges. Wien, x. (1860) 181 = pentandra.
polaris, *Wahlenb. Fl. Lapp.* 261.—Reg. arct.
poliphylla, Schleich. ex Ser. Ess. Saul. 67 = nigricans.

SALIX:—
polyandra, Gleditsch, Forstw. ii. 4. 43; ex Schrank, Baier. Fl. i. 229 = pentandra.
polygonifolia, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
polymorpha, Ehrh. Beitr. v. 160 = repens.
polymorpha, Host, Sal. t. 69 = cinerea.
pomeranica, Willd. Enum. Hort. Berol. Suppl. 66; Forbes, Salict. Woburn. 281. t. 153 = daphnoides.
pondicheriana, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 489 = ichnostachya.
Pontederae, Vill. Hist. Pl. Dauph. iii. 766 = hastata.
Pontederana, Schleich. Cat. Pl. Helv. ed. III. 25 = purpurea.
Pontederana, Willd. Sp. Pl. iv. 661 = hastata.
populifolia, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 494 = tetrasperma?
populifolia, *Schleich. ex Spreng. Pugill.* ii. 89.—Europ.
praecox, Dum. Fl. Belg. 12 = cinerea, acutifolia, daphnoides.
praecox, Hoppe, ex Willd. Sp. Pl. iv. 670 = daphnoides.
praecox, Salisb. Prod. 394 = S. Caprea.
pratensis, Host, Sal. 15. tt. 50, 51 = repens.
pratensis, Scop. Fl. Carn. ed. II. ii. 252 = purpurea.
prinoides, Pursh, Fl. Am. Sept. ii. 613 = brachystachys.
procumbens, Forbes, Salict. Woburn. 121. t. 61 = S. Myrsinites.
prolixa, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 94.—Am. bor. occ.
propendens, *Chaub. in Act. Linn. Soc. Bord.* xix. (1853) 234.—Gallia.
propendens, Ser. Ess. Saul. 73 = babylonica.
propinqua, Borrer, in Engl. Bot. Suppl. t. 2729 = phylicifolia.
prostrata, Ehrh. Beitr. vi. 103 = caesia.
prostrata, Muhl. Cat. 95 = Cutleri.
prostrata, Sm. Fl. Brit. 1060 = repens.
protea, Anderss. in Bot. Notiser (1866), 157 = nigricans.
proteaefolia, Forbes, Salict. Woburn. 149. t. 75 = *ambigua*.
pruinosa, Bess. Cat. Hort. Bot. Crem. 121 = acutifolia.
pruinosa, Schleich. ex Ser. Ess. Saul. 44 = nigricans.
pruinosa, Wendl. ex Reichb. Fl. Germ. Excurs. 172 = daphnoides.
prunifolia, Host, Sal. 23–25 = nigricans.
prunifolia, Sm. Fl. Brit. 1054 = S. Arbuscula.
Pseudo-aurita, Schur, Enum. Pl. Transs. 620 = lanata.
Pseudo-caprea, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 570 = acuminata.
Pseudo-cordata, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 161 = Novae-Angliae.
Pseudo-glauca, Anderss. ex Nym. Consp. 670 = glauca.
Pseudo-grisea, Steud. Nom. ed. II. ii. 498 = Wallichiana.
Pseudo-myrsinites, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 129.—Am. bor.
psilocarpa, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
psilostigma, Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 496 = eriophylla.
pubera, Koch, in Boenn. Prod. Fl. Monast. 300 = mollissima.
puberula, Doell, Fl. Bad. 518 = nigricans.
pubescens, Schleich. Cat. Pl. Helv. ed. III. 26 = grandifolia.
pulchella, Host, Sal. 30. t. 98 = S. Arbuscula.
pulchra, Wimm. Sal. Europ. 7 = daphnoides.
pumila, Salisb. Prod. 394 = herbacea.
punctata, Wahlenb. Fl. Lapp. 269 = nigricans.
purpurea, *Linn. Sp. Pl.* 1017.—Europ.; As. et Afr. bor.
Purshiana, Spreng. Syst. v. 608 (Index) = nigra.
pychnocarpa, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 139 = chlorophylla.
pycnostachya, *Anderss. in Journ Linn. Soc.* iv. (1860) 44.—Reg. Himal.
pyracanthoides, Turcz. ex Trautv. in Act. Hort. Petrop. ix. (1884) 184 = chlorostachya.
pyracanthos, Turcz. ex Anderss. in DC. Prod. xvi. II. 246 = chlorostachya.
pyramidalis, Budischtschew, ex Trautv. in Act. Hort. Petrop ix. (1884) 184, nomen (Quid?).—Rossia.
pyrenaica, *Gouan, Illustr.* 77.—Europ. austr.

SALIX:—
pyrifolia, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 162.—Am. bor.
pyrifolia, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
pyrina, Wall. Cat. n. 3705 = tetrasperma.
pyrocanthoides, Turcz. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 143 = chlorostachya.
pyrolaefolia, *Ledeb. Fl. Alt.* iv. 270.—Ross. arct.
pyrolaefolia, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
radicans, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 69 = triandra.
radicans, Sm. Engl. Bot. t. 2700 = phylicifolia.
ramifusca, Forbes, Salict. Woburn. 105. t. 53 = phylicifolia.
Rapini, Ayasse, in Bull. Soc. Bot. Fr. xxvi. (1879) 341 = hastata.
rectijulis, Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 313 = S. Myrsinites.
recurvata, Pursh, Fl. Am. Sept. ii. 609 = humilis.
recurvata, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
reflexa, Forbes, Salict. Woburn. 187. t. 94 = nigricans.
reflexa, *Rostk. in Willd. Baumg.* ed. II. 442; *ex Fl. Sedin.* 385 (Quid?).—Europ.
reflexiflora, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
regalis, Hort. ex Wesmael, in Bull. Congr. Bot. Brux. (1864) 280 = alba.
Regelii, *Anderss. in DC. Prod.* xvi. II. 309.—Soongaria.
Reichardtii, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 249 = S. Caprea.
Reinii, Franch. & Sav. Enum. Pl. Jap. i. 459 = glabra.
repanda, *Dum. Fl. Belg.* 12.—Europ.
repens, *Linn. Sp. Pl.* 1020.—Europ.; As. bor.
repens, Pursh, Fl. Am. Sept. ii. 610 = argyrocarpa.
reptans, *Rupr. Fl. Samojed. Cisural.* 54. t. 3.—Ross. arct.
reticulata, *Linn. Sp. Pl.* 1018.—Reg. arct. et bor.
reticuloides, Anderss. in DC. Prod. xvi. II. 301 = praec.
retractata, *F. Fellm. in Bull. Soc. Nat. Mosc.* viii. (1835) 287.—Europ. bor.
retusa, Hook, Fl. Bor. Am. ii. 153 = phlebophylla.
retusa, *Linn. Syst.* ed. X. 1287.—Europ.
retusa, Oakes, ex Tuckerm. in Am. Journ. Sc. Ser. I. xlv. (Oct. 1843) 36 = Cutleri.
retusa, Turcz. in Bull. Soc. Nat. Mosc. (1854) 1. 386 = nummularia.
retusa, With. Arr. Brit. Pl. t. 31 = S. Myrsinites.
retusoides, J. Kern. in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 1221 = retusa.
Reuteri, Mor. Fl. Schweiz, 459 (1844) = daphnoides.
reversifolia, Wahlenb. ex Schur, Enum. Pl. Transs. 621 = repens.
rhaetica, Kern. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 136 = phylicifolia.
rhamnifolia, Hook. & Arn. Bot. Beech. Voy. 117. t. 26 = fuscescens.
rhamnifolia, *Pall. Fl. Ross.* i. II. 84.—Sibir.
Rhodani, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
Richardsoni, *Hook. Fl. Bor. Am.* ii. 147. t. 182.—Am. arct.
rigida, Muhl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 237; Willd. Sp. Pl. iv. 667 = cordata.
riparia, Willd. l. c. 698 = incana.
Ritschelii, Anderss. in DC. Prod. xvi. II. 313 = purpurea.
rivalis, Host, Sal. 25. tt. 81, 82 = nigricans.
rivularis, Schleich. Cat. Pl. Helv. ed. III. 26; Forbes, Salict. Woburn. 203. t. 102 = nigricans.
rorismarinifolia, Gouan, Hort. Monsp. 501 = incana.
rosea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 231 = purpurea.
rosmarinifolia, Barratt & Hook. Fl. Bor. Am. ii. 148 = petiolaris.
rosmarinifolia, Laest. ex Anderss. in DC. Prod. xvi. II. 232 = fuscescens.
rosmarinifolia, Linn. Sp. Pl. 1020 = repens.
rosmarinifolia, Pursh, Fl. Am. Sept. ii. 612 = petiolaris.
Rossmaessleri, Willk. ex Anderss. in DC. Prod. xvi. II. 309 = purpurea.

SALIX:—
rostrata, Richards. in Frankl. Voy. 1st Journ. (1823) 753 = livida.
rostrata, Schleich. Cat. Pl. Helv. ed. IV. 31; ex Ser. Ess. Saul. 65 = nigricans.
rostrata, Thuill. Fl. Par. ed. II. 516 = repens.
Rothii, Hoppe, ex Reichb. Fl. Germ. Excurs. 169 = cinerea.
rotundata, Forbes, Salict. Woburn. 207. t. 104 = nigricans.
rotundifolia, *Nutt. N. Am. Sylv.* i. 75.—Am. bor. occ.
rotundifolia, Royle, ex Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 497 = flabellaris.
rotundifolia, *Trautv. in Nouv. Mém. Acad. Mosc.* ii. (1832) 304.—Europ. arct.
Roylei, Klotzsch, ex Anderss. in Vet. Acad. Handl. Stockh. 1850 (1851) 479 = hastata.
rubens, J. & C. Presl, Fl. Čech. 203 = silesiaca.
rubens, Schrank, Baier. Fl. i. 226 = nigra.
rubra, Huds. Fl. Angl. ed. I. 364 = purpurea.
rubra, Richards. in Frankl. Voy. 1st Journ. 752 = longifolia.
Rudbeckiana, Fries, ex Anderss. in DC. Prod. xvi. II. 241 = nigricans.
rufescens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 380 = viminalis.
rufinervis, DC. Fl. Fr. v. 371 = cinerea.
rugosa, Ser. Ess. Saul. 18 = aurita.
rugosa, Schleich. ex Ser. l. c. 67 = nigricans.
rugulosa, Anderss. in Bot. Notiser (1866), 145 = myrtilloides.
rugulosa, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
rupestris, Donn, ex Sm. Engl. Bot. t. 2342 = phylicifolia.
Russelliana, Sm. Fl. Brit. 1045 = fragilis.
sabulosa, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 385 = pyrolaefolia.
sachalinensis, *F. Schmidt, Reisen Amurl.* 173.—Ins. Sachalin.
Sadleri, Syme, in Journ. Bot. xiii. (1875) 33 = lanata.
Safsaf, *Forsk. Fl. Aegypt. Arab.* 76.—Afr. bor. et trop.
salviaefolia, Brot. Fl. Lusit. i. 29; Link, ex Willd. Sp. Pl. iv. 688 = oleaefolia.
salviaefolia, Koch, Syn. Fl. Germ. ed. I. 747 = incana.
sanguinea, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
Sansoniana, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) 1. 377 = purpurea.
sarawschanica, *Regel, in Act. Hort. Petrop.* vi. (1879) 466; *Pl. Nov. Fedsch.* 80.—As.
sarmentacea, *Fries, Novit.* i. 75.—Europ. bor.
saxatilis, *Turcz. in Ledeb. Fl. Ross.* iii. 621; *in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 391.—Sibir. baical.
Schleicheriana, Schleich. ex Ser. Ess. Saul. 65; Forbes, Salict. Woburn. 195. t. 98 = nigricans.
Schnelleriana, Richter, in Corresp. Ver. Naturk. Presb. ii. (1863) 194 (Quid?).—Europ.
Schraderiana, Willd. Sp. Pl. iv. 695 = phylicifolia.
Schrenkiana, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. 1. 179 = glabra.
Schultziana, Willd. ex Link, Handb. ii. 453 = repens.
sclerophylla, *Anderss. in Journ. Linn. Soc.* iv. (1860) 52.—Reg. Himal.
Scouleriana, Barratt, in Hook. Fl. Bor. Am. ii. 145 = brachystachys, sitchensis.
secta, Hook. f. ex Anderss. in Journ. Linn. Soc. iv. (1860) 57 = oreophila.
semicordata, Dulac, Fl. Hautes-Pyr. 147 = phylicifolia.
semihelix, Lasch, ex Wimm. Sal. Europ. 173 = purpurea.
semperflorens, Host, Sal. tt. 5–16 = triandra.
Sendtneri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 117.—Europ.
senegalensis, Mert. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 12 = coluteoides.
sensitiva, Barratt, Salic. Am. n. 8 = discolor.
sepiaria, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. 1. 127 = nigricans.
septentrionalis, Lodd. ex G. Don, in Loud. Hort. Brit. Suppl. ii. 671 = nigricans.
sericans, Tausch, ex Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 214 = Smithiana.
sericea, Marsh. Arbust. 140 = petiolaris.
sericea, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 239.—Am. bor.
sericea, Vill. Hist. Pl. Dauph. i. 382 = glauca.
sericocarpa, *Anderss. in Journ. Linn. Soc.* iv. (1860) 43.—Reg. Himal.

SALIX:—
Seringeana, Lecoq & Lamotte, Cat. 336 = oleaefolia.
Seringiana, *Gaud. Fl. Helv.* vi. 251.—Europ.
serotina, Pall. Reise, iii. 759 = viminalis.
serotina, Schur, Enum. Pl. Transs. 617 = triandra.
serpens, Tausch, ex Opiz, Boehm. Gew. (1823) 111 (Quid?).—Europ.
serpyllifolia, Scop. Fl. Carn. ed. II. ii. 255 = retusa.
Serpyllum, Anderss. in Journ. Linn. Soc. iv. (1860) 55. —Reg. Himal.
serrulata, Willd. Sp. Pl. iv. 664 = hastata.
sessilifolia, *Nutt. N. Am. Sylv.* i. 68.—Am. bor. occ.
sibirica, Pall. Fl. Ross. i. II. 78 = caesia.
Sieboldiana, *Blume, Bijdr.* 517.—Japon.
Sieboldii, Hort. ex C. Koch, Dendrol. ii. II. 505 = elegantissima.
Siegerti, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 68 = purpurea.
sikkimensis, *Anderss. in DC. Prod.* xvi. II. 269.—Reg. Himal.
silesiaca, Fries, ex Nym. Consp. 667 = S. Caprea.
silesiaca, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
silesiaca, *Willd. Sp. Pl.* iv. 660.—Europ.; Reg. Caucas.
sitchensis, *Sanson, ex Bong. in Mém. Acad. Pétersb.* ii. (1833) 162. t. 2.—Am. bor. occ.
skiaphila, Tausch, ex Opiz, Boehm. Gew. (1823) 111, nomen (Quid?).—Europ.
Smithiana, *Willd. Enum. Hort. Berol.* 1008.—Europ.
Sommerfeltii, Anderss. in DC. Prod. xvi. II. 291 = S. Myrsinites.
songarica, *Anderss. in Vet. Acad. Handl. Stockh.* vi. (1867) n. I. 53.—Soongar.
sordida, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 257 = purpurea.
sordida, Schleich. ex Ser. Ess. Saul. 65 = nigricans.
spadicea, Chaix, in Vill. Hist. Pl. Dauph. i. 373 = cinerea.
spathulata, Schultz, Prod. Fl. Starg. Suppl. i. 54 = repens.
spathulata, Willd. Sp. Pl. iv. 700 = repens.
speciosa, *Hook. & Arn. Bot. Beech. Voy.* 130.—Am. arct.; Sibir.
speciosa, Host, Sal. 5. t. 17 = triandra.
speciosa, Nutt. N. Am. Sylv. i. 58 = lasiandra.
spectabilis, Host, Sal. tt. 3, 4 = triandra.
sphacelata, Loisel. Fl. Gall. ii. 343 = grandifolia.
sphacelata, Schleich. ex Ser. Ess. Saul. 21 = grandifolia.
sphacelata, Sommerf. ex Reichb. Fl. Germ. Excurs. 845 = livida.
sphacelata, Willd. Sp. Pl. iv. 702 = S. Caprea.
sphaerocephala, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 240 = purpurea.
sphenocarpa, Tausch, ex Wimm. Sal. Europ. 196 = Lapponum.
sphenogyna, Wimm. l. c. = Lapponum.
spiraeaefolia, Schleich. ex Ser. Ess. Saul. 40 = aurita.
spiraeifolia, Willd. ex Link, Enum. Hort. Berol. ii. 416 = nigricans.
spissa, Anderss. in DC. Prod. xvi. II. 283 = subcordata.
splendens, Bray, ex Opiz, Boehm. Gew. (1823) 110 = alba.
splendens, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 380 = viminalis.
spuria, Schleich. ex Ser. Ess. Saul. 54; Willd. ex Anderss. in DC. Prod. xvi. II. 279 = Lapponum.
stagnalis, Nutt. N. Am. Sylv. i. 66 = brachystachys.
Starkeana, Willd. Sp. Pl. iv. 677 = livida.
stenoclados, Doell, Fl. Bad. 521 = livida.
stenostachya, Kern. in Oestr. Bot. Zeitschr. xiv. (1864) 188 = glabra.
stipulacea, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 343.—Mexic.
stipularis, Ser. Ess. Saul. 20 = grandifolia.
stipularis, Sm. Fl. Brit. 1079 = viminalis.
stolbergensis, Wallr. ex Anderss. in Vet. Akad. Handl. Stockh. vi. (1867) n. I. 171 = hastata.
strepida, Schleich. Cat. Pl. Helv. ed. III. 26; Forbes, Salict. Woburn. 199. t. 100 = nigricans.
Stuartiana, Sm. Engl. Bot. t. 2586; Hook. Fl. Bor. Am. ii. 151 = Lapponum.
Sturneana, Andrz. ex Trautv. in Act. Hort. Petrop. ix. (1884) 187 = livida?
stylaris, Ser. Ess. Saul. 62 = nigricans.
stylosa, DC. Fl. Fr. Suppl. 339 = nigricans.
suaveolens, Anderss. in Vet. Akad. Handl. Stockh. 1850 (1851) 491 = tetrasperma.

SALIX:—
subalpina, Forbes, Salict. Woburn. 185. t. 93 = repens.
subalpina, *Kern. in Oestr. Bot. Zeitschr.* xvi. (1866) 338.—Europ.
subaurita, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 67 = silesiaca.
subcaprea, Anderss. l. c. = silesiaca.
subcinerea, Anderss. l. c. = silesiaca.
ubconcolor, Ser. Ess. Saul. 54 = Lapponum.
subcordata, *Anderss. in Oefvers. Vet. Akad. Foerhandl.* xv. (1858) 128.—Am. bor.
subfragilis, *Anderss. in Mem. Am. Acad.* N. S. vi. (1858–59) 450.—Japon.
subglabra, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 237 = glabra.
subhastata, Laest. ex Anderss. Sal. Lapp. 21 = lanata.
sublanata, Anderss. ex Nym. Consp. 669 = lanata.
subnivalis, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 106.—Europ.
subopposita, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 28. —Japon.
subsericea, Doell, Fl. Bad. 517 = repens.
subserrata, Willd. Sp. Pl. iv. 671 = S. Safsaf.
subtriandra, *Neilr. in Verh. Zool.-Bot. Ges. Wien*, i. (1851) 119.—Europ.
sudetica, Host, Sal. 28 = Lapponum.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 474.—Sumatra.
taimyrensis, Trautv. in Middend. Sibir. Reise, 27. t. 5, 6 = arctica.
Tauschiana, Sieber, Cat. (1821); ex Wimm. Sal. Europ. 196; Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 68 = Lapponum.
taxifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 22.—N. Mexic.
tenera, *Anderss. in DC. Prod.* xvi. II. 288.—Am. bor. occ.
tenuiflora, Host, Sal. t. 5–16 = triandra.
tenuifolia, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
tenuifolia, Sm. Engl. Bot. t. 2186 = phylicifolia.
tenuifolia, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 376 = purpurea.
tenuijulis, Ledeb. Fl. Alt. iv. 262 = purpurea.
tenuior, Borrer, in Engl. Bot. Suppl. t. 2650 = phylicifolia.
tenuis, Host, Sal. 14. tt. 47, 48 = repens.
tenuis, Tausch, in Literat. Flora (1837), 174 (err. typ. 673) = purpurea.
tephrocarpa ×, Wimm. Sal. Europ. 205.—Europ.
Teplouchovi ×, Schroed. in Gard. Chron. (1882) II. 108.—Hab.?
testacea, Wright, ex Anderss. in DC. Prod. xvi. II. 201 = nigra.
tetrandra, *Willd. Enum. Hort. Berol. Suppl.* 66, nomen; *Reichb. Ic. Fl. Germ.* xi. 28. t. 612.—Europ.
tetrapla, Walk. Ess. 408; ex C. Koch, Dendrol. ii. I. 578 = phylicifolia.
tetrasperma, *Roxb. Pl. Corom.* i. 66. t. 97.—Ind. or.; Malaya.
thianschanica, *Regel, in Act. Hort. Petrop.* vi. (1879) 471.—Turkestan.
Thomasiana, Reichb. Ic. Fl. Germ. xi. 16. t. 560 = retusa.
Thomasii, *Anderss. in DC. Prod.* xvi. II. 297.—Helvet.
Thomsoniana, *Anderss. in Journ. Linn. Soc.* iv. (1860) 54.—Reg. Himal.
Thunbergiana, *Blume, ex Anderss. in DC. Prod.* xvi. II. 271.—Japon.
thymelaeoides, Schleich. ex Ser. Ess. Saul. 49 = S. Arbuscula.
tiliaefolia, Schleich. ex Spreng. Pugill. ii. 87 = nigricans.
Timmii, Schkuhr, ex Reichb. Fl. Germ. Excurs. 169 = cinerea.
tinctoria, Sm. in Rees, Cycl. xxxi. n. 13 = pentandra.
tomentella, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
tomentosa, Host, Sal. 28 = Lapponum.
tomentosa, Schrad. ex Anderss. in DC. Prod. xvi. II. 278 = candida.
tomentosa, Ser. Ess. Saul. 14 = S. Caprea.
torfacea, Schleich. ex Ser. l. c. 67 = nigricans.
Torreyana, *Barratt, Salic. Am.* n. 29.—Am. bor.
tortuosa, Host, Sal. t. 15 = cinerea.
Traunsteineri, Kern. ex Anderss. in DC. Prod. xvi. II. 308 = purpurea.
Trautvetteriana, *Regel, in Act. Hort. Petrop.* vi. (1879) 465.—As. centr.

SALIX:—
Trevirani, Wimm. ex Anderss. in DC. Prod. xvi. II. 204 = triandra.
Treviranii, Spreng. Pugill. i. 61 = mollissima.
triandra, *Linn. Sp. Pl.* 1016.—Europ.; Oriens; As. bor.
triandroides, Ehrh. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 28 = triandra.
trichocarpa, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
tridentata, Viv. Fl. Lib. Spec. 61. t. 7. f. 4 = Nitraria tridentata.
tristis, *Ait. Hort. Kew.* ed. I. iii. 393.—Am. bor.
tristis, Hort. ex Pursh, Fl. Am. Sept. ii. 614 = nigra.
tristis, *Muhl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 241.—Am. bor.
tuberculata, Moench, Meth. 336; Dum. Fl. Belg. 13 (Quid?).—Europ.
turfacea, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 127 = nigricans.
turgida, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
udensis, *Trautv. in Middend. Reise* (*Fl. Ochot.* 81).—Sibir.
uliginosa, Schleich. ex Ser. Ess. Saul. 21 = grandifolia.
uliginosa, Starke, ex Willd. Sp. Pl. iv. 677 = livida.
uliginosa, Willd. Enum. Hort. Berol. 1007 = aurita.
ulmifolia, Schleich. ex Ser. Ess. Saul. 21 = nigricans.
ulmifolia, Thuill. Fl. Par. ed. II. 518 = S. Caprea.
ulmifolia, Vill. Hist. Pl. Dauph. iii. 776 = aurita.
unalaschkensis, Cham. in Linnaea, vi. (1831) 539 = ovalifolia.
undulata, Ehrh. Beitr. vi. 101 = triandra.
undulata, Forbes, Salict. Woburn. 25. t. 13 = multiformis.
undulata, Schleich. ex Ser. Ess. Saul. 66 = nigricans.
uralensis, Hort. ex C. Koch, Dendrol. ii. I. 529 = purpurea.
urophylla, Lindl. in Wall. Cat. n. 3708 = tetrasperma.
Uva-ursi, Pursh, Fl. Am. Sept. ii. 610 = Cutleri.
Uva-ursi, Seem. Bot. Voy. Herald, 40 = ovalifolia.
vacciniifolia, Sm. Engl. Bot. t. 2341 = S. Arbuscula.
vaccinoides, Schleich. ex Ser. Ess. Saul. 67 = nigricans.
vagans, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1851) 121 = livida.
vagans, Hook. f. ex Anderss. in Journ. Linn. Soc. iv. (1860) 54 = Thomsoniana.
vallesiaca, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
varia, Host, Sal. t. 5–16 = triandra.
varians, Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 127 = nigricans.
vaudensis, Forbes, Salict. Woburn. 233. t. 117 = nigricans.
vaudensis, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 263 = cinerea.
velutina, Schleich. ex Ser. Ess. Saul. 53 = nivea.
velutina, Schrad. ex Koch, Syn. Fl. Germ. ed. I. 650 = Smithiana.
venulosa, Sm. Fl. Brit. 1055 = S. Arbuscula.
venusta, *Anderss. in DC. Prod.* xvi. II. 288.—Am. bor.
venusta, Host, Sal. t. 5–16 = triandra.
versicolor, Forbes, Salict. Woburn. 153. t. 77 = repens.
versifolia, Ser. Ess. Saul. 40 = repens.
versifolia, Wahlenb. Fl. Lapp. 271 = myrtilloides.
vestita, *Pursh, Fl. Am. Sept.* ii. 610.—Am. bor.
viadrina, Wimm. in Hartig, Naturg. Forstl. Cult. 6 = viminalis.
viburnoides, Schleich. ex Ser. Ess. Saul. 61 = hastata.
Villarsiana, Fluegge, ex Willd. Sp. Pl. iv. 655 = triandra.
villosa, D. Don, ex Hook. Fl. Bor. Am. ii. 144 = glauca.
villosa, Forbes, Salict. Woburn. 183. t. 92 = tristis.
villosa, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
villosula, Schleich. ex Ser. Ess. Saul. 43 = nigricans.
vilnensis, Gorski, ex Reichb. Ic. Fl. Germ. xi. 24 = myrtilloides.
viminalis, Balb. Fl. Taur. 165 = incana.
viminalis, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 451 = Kirilowiana.
viminalis, *Linn. Sp. Pl.* 1021.—Europ.; As. bor.
viminalis, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 377 = purpurea.
violacea, G. Anders. ex Sm. in Rees, Cycl. xxxi. n. 33 = daphnoides.
violacea, Bray, ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) n. I. 95 = myrtilloides.
violacea, Holl. ex Nym. Consp. 666 = mollissima.
violacea, Hort. ex Anderss. in DC. Prod. xvi. II. 250 = phylicifolia.

SALIX:—
violacea, Willd. Enum. Hort. Berol. Suppl. 66, nomen (Quid?).—Hab.?
virescens, Forbes, Salict. Woburn. 13 = mollissima.
virescens, Vill. Hist. Pl. Dauph. iii. 785 = viminalis.
virgata, Forbes, Salict. Woburn. 23. t. 12 = multiformis.
virgata, Schleich. Cat. Pl. Helv. ed. III. 26 = nigricans.
viridis, Fries, Novit. ed. II. 283 = fragilis.
viridula, Anderss. in Mem. Am. Acad. N. S. vi. (1858–59) 451 = padifolia.
vitellina, Linn. Sp. Pl. 1016 = alba.
volgensis, *Anderss. in DC. Prod.* xvi. II. 314.—Rossia.
vratislaviana ×, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 216.—Europ.
vulgaris, Clayton, ex Gronov. Fl. Virg. ed. II. (1762) 154 = nigra.
vulpina, Anderss. in Mem. Am. Acad. N. S. vi. (1858–59) 452 = Miquelii.
Wahlenbergii, Anderss. in Bot. Notiser (1867), 115 = nigricans.
Waldsteiniana, Willd. Sp. Pl. iv. 679 = S. Arbuscula.
Wallichiana, *Anderss. in Vet. Acad. Handl. Stockh.* 1850 (1851) 477.—Reg. Himal.
Wallichii, Wimm. ex Anderss. l. c. vi. (1867) n. I. 4 = tetrasperma.
wargiana, Lej. Fl. Spa, ii. 322 = fragilis.
washitana, Muhl. ex Rafin. Fl. Ludov. 138, nomen (Quid?).—Am. bor.
Weigeliana, Willd. Sp. Pl. iv. 678 = phylicifolia.
Weinmanniana, Spreng. ex Anderss. in Vet. Acad. Handl. Stockh. vi. (1867) I. 95 = mytilloides.
Wichurae, Anderss. l. c. vi. (1867) 135 = glauca.
Wichurae, Kern. ex Anderss. in DC. Prod. xvi. II. 305 = incana.
Wilhelmsiana, Bieb. Fl. Taur. Cauc. iii. 627 = angustifolia.
Willdenoviana, Forbes, Salict. Woburn. 81. t. 41 = nigricans.
Wimmeri, Kern. in Flora, xxxv. (1852) 511 = daphnoides.
Wimmeriana, Gren. & Godr. Fl. Fr. iii. 130 = purpurea.
Wolfii, *Bebb, in Wheeler, Rep. Bot. U. St. Geogr. Surv.* vi. 241.—Am. bor. occ.
Woolgariana, Borrer, in Engl. Bot. Suppl. t. 2651 = purpurea.
Wrightii, Anderss. in Oefvers. Vet. Akad. Foerhandl. xv. (1858) 115 = nigra.
Wulfeniana, Forbes, Salict. Woburn. 95 = phylicifolia.
Wulfeniana, Pollini, Fl. Veron. iii. 160 = nigricans.
Wulfeniana, Willd. Sp. Pl. iv. 660 = glabra.
Zedlitziana ×, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 217.—Europ.
Zeyheri, Sond. ex Fries, in Oefvers. Vet. Akad. Foerhandl. xiii. (1856) 121 = capensis.
Zollingeri, Miq. in Zoll. Syst. Verz. Ind. Archip. 108 = tetrasperma.
zygostemon, Boiss. & Hohen. Diagn. Ser. I. xii. 118 = oxycarpa.

SALIZARIA, A. Gray, Syn. Fl. N. Am. ii. I. 382 (1878), sphalm. = **Salazaria**, Torr. (Labiat.).

SALKEA, Steud. Nom. ed. II. ii. 500 (1841), sphalm. = seq.

SALKEN, Adans. Fam. ii. 322 (1763) = **Dalbergia**, Linn. (Legumin.).

SALLOA, Walp. Rep. v. 17 (1845), sphalm. = GALLOA, Hassk. = **Cocculus**, DC. (Menisp.).

SALMALIA, Schott & Endl. Melet. 35 (1832) = **Bombax**, Linn. (Malvac.).
insignis, Schott & Endl. l. c. 35 = B. insigne.
malabarica, Schott & Endl. l. c. = B. malabaricum.

SALMASIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317 = **Aira**, Linn. (Gramin.).
corymbosa, Bubani, l. c. = A. intermedia.

SALMASIA, Reichb. Handb. 295 (1837), sphalm. = SALMALIA, Schott & Endl. = **Bombax**, Linn. (Malvac.).

SALMASIA, Schreb. Gen. 201 (1789) = **Tachibota**, Aubl. (Inc. Sed.).
guianensis, J. F. Gmel. Syst. 500 = T. guianensis.
racemosa, Willd. Sp. Pl. i. 1502 = T. guianensis.
spinosa, Spreng. Neue Entdeck. ii. 166 = Zanthoxylum Ochroxylum.

SALMEA, DC. Cat. Hort. Monsp. 140 (1813). *COMPOSITAE*, Benth. & Hook. f. ii. 381.
FORNICARIA, Rafin. Sylva Tellur. 116 (1838).
HOPKIRKIA, Spreng. Nov. Prov. 23 (1818).
angustifolia, *Benth. Bot. Voy. Sulph.* 117.—Mexic.
caleoides, *Griseb. Cat. Pl. Cub.* 155.—Cuba.
curviflora, R. Br. in Trans. Linn. Soc. xii. (1818) 112 = Otopappus curviflorus.
Eupatoria, *DC. Cat. Hort. Monsp.* 140.—Ind. occ.
fruticulosa, Steud. Nom. ed. II. i. 775, in syn. (= *Armania fruticulosa*.).—Am. austr.
glaberrima, *Wright, ex Griseb. Cat. Pl. Cub.* 155.—Cuba.
grandiceps, Cass. in Dict. Sc. Nat. xlvii. 88 = S. Eupatoria.
hirsuta, *DC. Cat. Hort. Monsp.* 140.—Jamaic.
oligocephala, *Hemsl. Biol. Centr. Am. Bot.* ii. 194.—Mexic.
oppositiceps, Cass. in Dict. Sc. Nat. xlvii. 89 = hirsuta.
parviceps, Cass. l. c. 88 = S. Eupatoria.
pauciceps, *Griseb. Cat. Pl. Cub.* 155.—Ind. occ.
petrochioides, *Griseb. Fl. Brit. W. Ind.* 375.—Ind. occ.
salicifolia, *Brongn. in Rev. Hortic.* Sér. III. i. (1847) 87.—Hab.?
scandens, DC. in Cat. Hort. Monsp. 140 = S. Eupatoria.
sessilifolia, *Griseb. Fl. Brit. W. Ind.* 375.—Ind. occ.

SALMEOPSIS, Benth. in Benth. & Hook. f. Gen. ii. 381 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
Claussenii, *Benth. in Hook. Ic. Pl.* t. 1152.—Bras.

SALMIA, Cav. Ic. iii. 24. t. 246 (1794) = **Sansevieria**, Thunb. (Haemod.).
ebracteata, Cav. l. c. = S. lanuginosa.
guineensis, Cav. l. c. = S. guineensis.
spicata, Cav. l. c. = S. thyrsiflora.

SALMIA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399 = **Carludovica**, Ruiz & Pav. (Cyclanth.)
acuminata, Willd. l. c. 401 = C. acuminata.
angustifolia, Willd. l. c. = Cyclanthus Plumierii.
funifera, Spreng. Syst. iii. 772 = C. funifera.
humilis, Steud. Nom. ed. II. ii. 501 = C. humilis.
jamaicensis, Steud. l. c. = C. jamaicensis (in Addend.).
latifolia, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 401 = C. latifolia.
palmaefolia, Willd. l. c. = Cyclanthus Plumierii.
palmata, Willd. l. c. = C. palmata.
petrobioides, Griseb. Fl. Brit. W. Ind. 375.—Ind. occ.
subacaulis, Schult. Mant. iii. 341 = C. trigona.
tetragona, Roem. & Schult. Syst. iii. 496 = C. tetragona.
trigona, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 401 = C. trigona.

SALMONEA, Vahl, Enum. i. 8 (1805) = **Salomonia**, Lour. (Polygal.).
cantoniensis, Vahl, l. c. = Salomonia cantoniensis.

SALMONIA, Scop. Introd. 209 (1777) = **Vochysia**, Juss.

SALOMONIA, Heist. Syst. 5 (1748) (Quid?).

SALOMONIA, Lour. Fl. Cochinch. 14 (1790). *POLYGALEAE*, Benth. & Hook. f. i. 136.
EPIRIXANTHES, Blume, Cat. Gew. Buitenz. 25 (1823).
angulata, Griff. Notul. iv. 539 = ciliata?
aphylla, *Griff. in Proc. Linn. Soc.* i. (1844) 221; *et in Trans. Linn. Soc.* xiv. (1845) 112.—Malaya.
Arnottiana, Miq. Anal. Bot. Ind. iii. 3 = oblongifolia.
canarana, Hassk. in Miq. l. c. = oblongifolia.
cantoniensis, *Lour. Fl. Cochinch.* 14.—As. trop.
ciliata, *DC. Prod.* i. 334.—Ind. or.
cordata, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 322; et in Wight, Illustr. i. 49. t. 22 = ciliata.
cylindrica, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 79.—Burma.
edentula, *DC. Prod.* i. 334.—Reg. Himal.

SALMONIA:—
Horneri, Hassk. in Miq. Pl. Jungh. 123 = oblongifolia?
leptostachya, Wall. Cat. n. 4193 = oblongifolia.
longiciliata, *Kurz, in Journ. As Soc. Beng.* xli. (1872) II. 292.—Burma.
oblongifolia, *DC. Prod.* i. 334.—As. trop.; Austral.
obovata, Wight, Illustr. i. 49. t. 22 = oblongifolia.
parasitica, Griff. Notul. iv. 538 = aphylla.
petiolata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 200 = edentula.
ramosissima, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 352.—Ins. Philipp.
rigida, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 148 = oblongifolia.
sessiliflora, A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 207, sphalm. = seq.
sessilifolia, D. Don, Prod. Fl. Nep. 200 = oblongifolia.
setoso-ciliata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 149 = oblongifolia.
stricta, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 152 = oblongifolia.
subrotunda, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 146 = cantoniensis.
tenella, C. Muell. in Walp. Ann. vii. 243 = aphylla.
trinervata, Steud. Nom. ed. II. ii. 373 = cantoniensis.
uncinata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 148 = oblongifolia?

SALPIANTHUS, Humb. & Bonpl. Pl. Aequin. i. 154. t. 44 (1808) = **Boldoa**, Cav. (Nyctag.).
arenarius, Humb. & Bonpl. l. c. 155 = B. lanceolata.
purpurascens, Hook. & Arn. Bot. Beech. Voy. 309 = Cryptocarpus globosus.

SALPICHROA, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 321. *SOLANACEAE*, Benth. & Hook. f. ii. 899.
BUSBECKEA, Mart. Cat. Hort. Monac. (1829) 69.
PERIZOMA, Miers, ex Lindl. Veg. Kingd. 622 (1847).
PLANCHONIA, Dun. in DC. Prod. xiii. I. 471 (1852).
SALPICHROMA, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 333.
ciliata, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 329 = Lycium ciliatum.
dependens, *Miers, l. c.* 325.—Peruv.
diffusa, *Walp. Ann.* iii. 169.—Am. trop.
glandulosa, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 325.—Peruv.
hirsuta, *Miers, l. c.*—Peruv.
ramosissima, *Miers, l. c.* 326.—Peruv.
rhomboidea, *Miers, l. c.*—Reg. Argent.
tristis, *Walp. Rep.* iii. 170.—Peruv.
Wrightii, *Miers, ex A. Gray, Syn. Fl. N. Am.* ii. I. 231.—Am. bor. occ.

SALPICHROMA, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 333 = **Salpichroa**, Miers (Solan.).
breviflorum, Dun. in DC. Prod. xiii. I. 474.—Peruv.
ciliata, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 334 = Lycium ciliatum.
cuspidatum, Dun. in DC. Prod. xiii. I. 474.—Peruv.
dependens, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 334 = Salpichroa dependens.
Didieranum, Jaub. in Bull. Soc. Bot. Fr. viii. (1861) 117.—Peruv.
diffusum, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 335 = Salpichroa diffusa.
glandulosum, Miers, l. c. 334 = Salpichroa glandulosa.
hirsutum, Miers, l. c. = Salpichroa hirsuta.
Mandonianum, Wedd. Chlor. And. ii. 98.—Bolivia.
parviflorum, Dun. in DC. Prod. xiii. I. 476.—Bras.
ramosissimum, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 334 = Salpichroa ramosissima.
rhomboideum, Miers, l. c. = Salpichroa rhomboidea.
triste, Miers, l. c. 335 = Salpichroa tristis.
urceolatum, *Dun. in DC. Prod.* xiii. I. 475.—Mexic.

SALPIGLOSSIS, Ruiz & Pav. Prod. 94. t. 19 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 909.
atro-purpurea, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1827) 176 = sinuata.
aurea, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 762 = sinuata.
Barcklayana, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 599 = seq.

SALPIGLOSSIS :—
Barclayana, Sweet, Brit. Flow. Gard. Ser. II. t. 112 = sinuata.
coccinea, Lindl. & Paxt. Flow. Gard. iii. (1852–53) 135. t. 100 = sinuata.
fulva, *Court. in Mag. Hort.* i. (1833) 219. n. 411.—Chili.
hybrida, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 762 = sinuata.
integrifolia, Hook. Bot. Mag. t. 3113 = Petunia violacea.
integrifolia, *Lodd. Bot. Cab.* t. 1978.—Reg. Argent.
intermedia, Sweet, Hort. Brit. ed. II. 594; Brit. Flow. Gard. Ser. II. t. 112 = sinuata.
linearis, *Hook. Bot. Mag.* sub t. 3113; et t. 3256.—Reg. Argent.
parviflora, G. Don, ex Benth. in DC. Prod. x. 202 (Petuniae sp.).—Hab. ?
parviflora, Phil. Fl. Atac. 45 = Reyesia chilensis.
picta, Sweet, Brit. Flow. Gard. Ser. I. t. 258 = straminea.
prostrata, *Hook. & Arn. Bot. Beech. Voy.* 153.—Calif.
purpurea, *Miers, Trav.* ii. 531.—Chili.
sinuata, Hook. & Arn. ex Miers, in Hook. Lond. Journ. Bot. v. (1846) 190 = Petunia viscidula.
sinuata, *Ruiz & Pav. Syst. Veg.* 163.—Chili.
spinescens, *Clos, in C. Gay, Fl. Chil.* v. 127.—Chili.
straminea, *Hook. Exot. Fl.* iii. t. 229.—Chili.
variabilis, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 809 = sinuata.

SALPIGLOTTIS, Hort. ex C. Koch, in Neue Verh. Berl. Gartenb. i. (1853) 400 = praec.

SALPINGA, Mart. in DC. Prod. iii. 112 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 755.
cristata, DC. Prod. iii. 113 = Macrocentrum cristatum.
fasciculata, DC. l. c. = Macrocentrum fasciculatum.
longifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 80.—Bras.
margaritacea, *Triana, l. c.*—Bras.
parviflora, DC. Prod. iii. 113 = Macrocentrum cristatum.
secunda, *Schrank & Mart. ex DC. l. c.*—Bras.

SALPINGANTHA, Hort. ex Lem. in Fl. des Serres, iii. (1847) 240, sphalm. = SALPIXANTHA, Hook. = **Geissomeria**, Lindl. (Acanth.).

SALPINGOGLOTTIS, C. Koch, in Neue Verh. Berl. Gartenb. i. (1853) 401 = **Salpiglossis**, Ruiz & Pav. (Solanac.).

SALPIXANTHA, Hook. Bot. Mag. t. 4158 (1845) = **Geissomeria**, Lindl. (Acanth.).
coccinea, Hook. l. c. = G. coccinea.

SALSA, Feuillé, ex Ruiz & Pav. Fl. Per. iii. 70 (1802) = **Herreria**, Ruiz & Pav. (Liliac.).

SALSOLA, Linn. Syst. ed. I (1735). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 71.
CAROXYLON, Thunb. Nov. Gen. ii. 37 (1782).
HALOTHAMNUS, Jaub. & Spach, Illustr. ii. 50. t. 136 (1846).
ISGARUM, Rafin. Fl. Tellur. iii. 46 (1836).
KALI, Tourn. ex Adans. Fam. ii. 261 (1763).
SARCOMORPHIS, Boj. ex Moq. in DC. Prod. xiii. II. 179 (1849).
SODA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 145.
abrotanoides, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 366.—Mongol.
acicularis, Salisb. Prod. 153 = S. Kali.
acuminata, D. Dietr. Syn. Pl. ii. 998 = Suaeda acuminata.
affinis, *C. A. Mey. ex Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 360.—Sibir.
africana, Hort. ex Moq. in DC. Prod. xiii. II. 176 = aphylla.
agrigentina, Guss. Fl Sic. Prod. i. 303 = vermiculata, *Linn.*
alopecurioides, Delile, Fl. Egypte, 200. t. 21 = Halogeton alopecuroides.
altissima, Dryand. ex Moq. in DC. Prod. xiii. II. 170 = Helicilla altissima.
altissima, Linn. Sp. Pl. ed. II. 324 = Suaeda altissima.
annularis, Poir. Encyc. vii. 298 = Suaeda fruticosa.
anomala, C. A. Mey. ex Eichw. Pl. Casp.-Cauc. 14. t. 12 = Horaninovia juniperina.
anonyma, Poir. Encyc. vii. 299 = Suaeda anonyma.
aphylla, *Linn. f. Suppl.* 173.—Afr. austr.
aphylla, Spreng. Syst. i. 925 = Kochia aphylla.
arabica, Ehrenb. ex Boiss. Fl. Orient. iv. 970, in obs. = Anabasis Ehrenbergii.
arborescens, Linn. f. Suppl. 173 = S. Arbuscula.
Arbuscula, *Pall. Reise*, i. 487.—Rossia; Pers.; Sibir.
arenaria, Waldst. & Kit. Pl. Rar. Hung. i. 80 = Kochia arenaria.
articulata, Cav. Ic. iii. 43. t. 284 = Haloxylon articulatum.
articulata, Forsk. Fl. Aegypt. Arab. 55 = Anabasis articulata.
asparagoides, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 194.—Japon.
atriplicifolia, Spreng. Nachr. i. Bot. Gart. Halle, 35 = Cycloloma platyphyllum.
Atriplicis, Schult. Obs. Bot. 52 = Cycloloma platyphyllum.
aurantiaca, *Bunge, ex Boiss. Fl. Orient.* iv. 963.—Persia; Afghan.
Auricula, *Moq. Chenop. Enum.* 135.—Persia; Afghan.
australis, R. Br. Prod. 411 = S. Kali.
baccata, Poir. Encyc. vii. 298 = Suaeda baccata.
baccifera, Pall. Illustr. t. 23 = clavifolia.
baccifera, Schrad. Neues Journ. iii. (1809) 88 = Suaeda baccifera.
Bottae, *Boiss. Fl. Orient.* iv. 964, in obs.—Arab.
brachiata, *Pall. Illustr.* 30. t. 22.—As. med.
brachyphylla, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 959.—Persia.
brachyphylla, Spreng. Syst. i. 924 = Kochia brevifolia.
brachypteris, Moq. Chenop. Enum. 147 = S. Kali.
brevifolia, Desf. Fl. Atlant. i. 218 = vermiculata, *Linn.*
buxifolia, Hort. Par. ex Dum.-Cours. Bot. Cult. ed. I. i. 622 = praec. ?
caesia, Fisch. & Mey. in Bull. Soc. Nat. Mosc. xi. (1838) 357 = ericoides.
Calluna, *Fenzl, ex Moq. in DC. Prod.* xiii. II. 191, nomen.—Afr. austr.
camphorosmoides, Desf. Fl. Atlant. i. 218 = Noaea spinosissima.
cana, C. Koch, in Linnaea, xxii. (1849) 190 = crassa.
candida, *Fenzl, ex Moq. in DC. Prod.* xiii. II. 191, nomen.—Afr. austr.
canescens, *Boiss. Fl. Orient.* iv. 963.—As. Min.; Persia.
canescens, Desf. Tabl. ed. I. 41 = Chenolea diffusa.
carinata, *C. A. Mey. in Eichw. Pl. Casp.-Cauc.* 36. t. 28.—Reg. Casp.; Soongar.; Afghan.
carinata, Spreng. Syst. i. 923 = Chenopodium carinatum.
carnosa, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
caroliniana, Walt. Fl. Carol. 111 = S. Kali.
Caroxylon, Moq. Chenop. Enum. 148 = aphylla.
celtifolia, Schult. Syst. vi. 243 = ciliata.
chenopodiana, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
chenopodioides, Hort. ex Dum.-Cours. Bot. Cult. ed. II. ii. 463 = Cycloloma platyphyllum.
ciliata, *Willd. Enum. Hort. Berol. Suppl.* 14.—Hab. ?
cinerea, Waldst. & Kit. Pl. Rar. Hung. ii. 110. t. 106 = Kochia sedoides.
clavifolia, *Pall. Reise*, ii. 486.—Rossia; As. bor.
clavifolia, Unio, It. ex Moq. in DC. Prod. xiii. II. 157 = Suaeda setigera.
collina, *Pall. Illustr.* 34. t. 26.—Rossia; Reg. Himal.; Sibir.
conglomerata, *Gueldenst. Reisen*, ii. 98.—Ross. austr.
controversa, Todaro, ex Nym. Consp. 631 = S. Kali.
coquimbana, *Molina, Sagg. Chil.* 132, 330.—Chili.
corniculata, D. Dietr. Syn. Pl. ii. 998 = Suaeda corniculata.
corticosa, *Meyen, Reise*, i. 378.—Chili.
corymbosa, Hort. ex Moq. in DC. Prod. xiii. II. 60 = Cycloloma platyphyllum.
crassa, *Bieb. in Mém. Soc. Nat. Mosc.* i. (1811) 100.—Rossia; As. bor.
daghestanica, *Turcz. ex Bunge, Anab. Revis.* 26.—Reg. Caucas.
dasyantha, Pall. Illustr. 19. t. 11 = Kochia arenaria.
decumbens, Lam. Fl. Fr. iii. 241 = S. Kali.
decurrens, Jaub. & Spach, Illustr. ii. 53. t. 138 = S. Auricula.
dendroides, Pall. Illustr. 22. t. 14 = verrucosa.
depressa, Pursh, Fl. Am. Sept. i. 197 = Suaeda plattensis.
dichotoma, Pall. Reise, iii. 604 = Ofaiston paucifolium.
didyma, *Lour. Fl. Cochinch.* 17.—Afr. trop.
diffusa, Thunb. Prod. Pl. Cap. 48 = Kochia sericea.
dioeca, Spreng. Syst. i. 923 = Kochia dioica.
divaricata, Mass. ex Link, in Buch, Beschr. Canar. Ins. 141 = longifolia.
divergens, Poir. Encyc. vii. 299 = Suaeda hortensis.
Echinus, Labill. Ic. Pl. Syr. dec. ii. 10. t. 5 = Noaea spinosissima.
elata, Wight, in Wall. Cat. n. 6944 = Suaeda nudiflora.
ennaea, Jan, Elench. Pl. Parm. (1826) 4 = vermiculata *Linn.*
eriantha, Hort. ex Moq. in DC. Prod. xiii. II. 135 = Kochia eriophora.
ericaefolia, Adams, in Hoffm. Hort. Mosq. (1808) n. 2764 = ericoides.
ericifolia, Mass. ex Link, in Buch, Beschr. Canar. Ins. 141 = vermiculata, *Linn.*
ericoides, *Bieb. in Mem. Soc. Nat. Mosc.* i. (1811) 141.—Reg. Transcauc.
ericoides, Lag. ex Willk. & Lange, Prod. Fl. Hisp. i. 258 = Webbii.
ericoides, C. A. Mey. Verz. Pfl. Cauc. 160 = verrucosa.
eriophora, Steph. ex Bieb. in Mém. Soc. Nat. Mosc. iv. (1813) 17 = Kochia eriophora.
erubescens, Schrad. Ind. Sem. Hort. Gotting. (1834) 5 = collina.
exasperata, *C. A. Mey. in Eichw. Pl. Casp.-Cauc.* 7 = S. Arbuscula.
farinosa, Poir. Encyc. vii. 298 = Suaeda vera.
ferox, Delile, ex Moq. in DC. Prod. xiii. II. 218 = Cornulaca monacantha.
flavescens, Cav. Ic. iii. 45. t. 288 = vermiculata, *Linn.*
florida, Poir. Encyc. Suppl. v. 191 = Seidlitzia florida.
foetida, *Delile, Fl. Egypt. Illustr.* n. 310.—Afr. trop.; Arab.; Ind. or.
foetida, Vest, ex Schult. Syst. vi. 238 = Suaeda foetida.
foliosa, Schrad. ex Schult. l. c. 235 = clavifolia.
frutescens, Pall, Reise, i. 488 = crassa.
fruticosa, Bieb. Tabl. Prov. Casp. 149 = Suaeda dendroides.
fruticosa, Cav. Ic. iii. 44. t. 285 = longifolia.
fruticosa, Drège, ex Moq. in DC. Prod. xiii. II. 131 = Kochia villosa.
fruticosa, Linn. Sp. Pl. ed. II. 324 = Suaeda fruticosa.
fruticosa, Wall. Cat. n. 6944 = Suaeda nudiflora.
gemmascens, *Pall. Illustr.* 24.—Persia; As. bor.
genistoides, *Juss. ex Poir. Encyc.* vii. 294.—Hispan.
georgica, Bunge, ex Ledeb. Fl. Ross. iii. 814 = verrucosa.
glauca, *Bieb. Fl. Taur. Cauc.* i. 185.—Reg. Caucas.; Persia.
glauca, C. A. Mey. Verz. Pfl. Cauc. 160 = ericoides.
globulifolia, Poir. Encyc. vii. 298 = Suaeda vermiculata.
glomerata, Steph. ex Moq. in DC. Prod. xiii. II. 206 = Halogeton glomeratus.
glomerulata, Delile, Fl. Aegypt. 56 = Halogeton alopecuroides.
glomerulata, *Meyen, Reise*, i. 375.—Chili.
gossypina, *Bunge, ex Boiss. Fl. Orient.* iv. 956.—Persia.
grandiflora, *Link, ex Steud. Nom.* ed. II. ii. 502.—Lusitan.
Helenae, Bory, ex Schult. Syst. vi. 242 = Suaeda fruticosa.
heterocarpa, Bunge, ex Boiss. Fl. Orient. iv. 953 = sogdiana.
hirsuta, Linn. Sp. Pl. ed. II. 323 = Kochia hirsuta.
hispidula, *Boiss. Fl. Orient.* iv. 960.—Turkest.; Afghan.
hissopifolia, Nocca, ex Usteri, Ann. Bot. vi. (1793) 60 = Kochia hyssopifolia.
hortensis, Desf. Tabl. ed. II. 50; Cat. Hort. Par. ed. III. 73; et ex Moq. in DC. Prod. xiii. II. 158 = Suaeda altissima.
Hostii, Tratt. Arch. 93. t. 112 = Suaeda maritima.
hyssopifolia, Pall. Reise, i. 491 = Kochia hyssopifolia.
imbricata, *Forsk. Fl. Aegypt. Arab.* 57.—Arab.

SALSOLA :—
incanescens, C. A. Mey. in Eichw. Pl. Casp.-Cauc. 35. t. 26 = spissa.
indica, Herb. Madr. ex Wall. Cat. n. 6946 C = Suaeda fruticosa.
indica, Willd. Sp. Pl. i. 1317 = Suaeda maritima.
inermis, *Forsk. Fl. Aegypt. Arab.* 57.—Aegypt.; Syria.
Jacquemontii, Moq. in DC. Prod. xiii. II. 188 = S. Kali?
Kali, *Linn. Sp. Pl.* 222.—Reg. bor. et austr. temp.
Lana, Edgew. in Hook. Journ. Bot. ii. (1840) 286 = Suaeda fruticosa.
Lana, Stocks, ex Boiss. Fl. Orient. iv. 950 = Haloxylon recurvum.
lanata, Hort. ex Moq. in DC. Prod. xiii. II. 135 = Kochia eriophora.
lanata, Mass. ex Link, in Buch, Beschr. Canar. Ins. 141 = Chenolea canariensis.
lanata, *Pall. Reise*, ii. 736.—Ross.; As. bor.
lanata, Poll. ex Moq. in DC. Prod. xiii. II. 132 = Kochia prostrata.
lanata, Vahl, in Hornem. Hort. Hafn. 259 = Kochia hyssopifolia.
lancifolia, *Boiss. Fl. Orient.* iv. 958.—Palaestin.
laniflora, S. G. Gmel. Reise, i. 160. t. 37 = Kochia arenaria.
laniflora, Hort. ex Moq. in DC. Prod. xiii. II. 135 = Kochia eriophora.
laniflora, Linn. f. Suppl. 172 = lanata.
lanuginosa, Hort. ex Moq. in DC. Prod. xiii. II. 135 = Kochia eriophora.
lanuginosa, Steud. Nom. ed. II. ii. 502 = glomerulata.
laricina, *Pall. Illustr.* 21. t. 13.—Ross.; As. bor.
latifolia, Hort. Par. ex Poir. Encyc. vii. 298 = Cycloloma platyphyllum.
leiosperma, D. Dietr. Syn. Pl. ii. 997 = Suaeda altissima.
leucoxylum, Moq. in DC. Prod. xiii. II. 183 = ericoides.
linearis, Ell. Sketch, i. 332 = Suaeda maritima.
littoralis, *Moq. in DC. Prod.* xiii. II. 180.—Madag.
longiflora, J. F. Gmel. Syst. 1618 = seq.
longifolia, *Forsk. Fl. Aegypt. Arab.* 55.—Afr. bor.; Palaest.
longifolia, Lam. Fl. Fr. iii. 233 = S. Soda.
lutea, Delile, ex Moq. in DC. Prod. xiii. II. 215 = Anabasis setifera.
macrophylla, R. Br. Prod. 411 = S. Kali.
marina, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
maritima, Bieb. Tabl. Prov. Casp. 150; Poir. Encyc. vii. 291 = Suaeda maritima.
Marosteum, Moq. in DC. Prod. xiii. II. 191, sphalm. = *Moorcroftiana*.
mendocinensis, Gill. ex Moq. l. c. 156 = Suaeda divaricata.
microphylla, Cav. Ic. iii. 45. t. 287 = vermiculata, *Linn.*
mollis, A. Cunn. ex Moq. in DC. Prod. xiii. II. 138 = Sclerolaena bicornis.
mollis, Desf. Fl. Atlant. i. 218 = Suaeda vermiculata.
monandra, Pall. Reise, iii. 724 = Ofaiston paucifolium.
monobractea, Forsk. Fl. Aegypt. Arab. 55 = Kochia muricata.
monocrophylla, Steud. Nom. ed. I. 723, sphalm. = *macrophylla*.
monoica, Poir. Encyc. vii. 299 = Suaeda monoica.
monoptera, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 364.—Sibir.
Moorcroftiana, Wall. Cat. n. 6947 = foetida.
Moquiniana, Jaub. & Spach, Illustr. ii. 52. t. 137 = S. Auricula.
mucronata, Forsk. Fl. Aegypt. Arab. 56 = Noaea spinosissima.
muricata, Bieb. Tabl. Prov. Casp. 148 = Kochia sedoides.
muricata, Linn. Mant. i. 54 = Kochia muricata.
mutica, *C. A. Mey. ex Becker, in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 455.—Ross. austr.
nitida, *Clarke, Trav.* iii. 42.—Aegypt.
nitraria, Pall. Illustr. 23. t. 15 = spissa.
nitrosa, Gueldenst. ex Ledeb. Fl. Ross. iii. 809 = spissa.
nudiflora, Wall. Cat. n. 6945 = Suaeda maritima.
nudiflora, Willd. Sp. Pl. i. 1313 = Suaeda nudiflora.
obtusifolia, *Ledeb. ex Sweet, Hort. Brit.* ed. III. 372.—Sibir.

SALSOLA :—
oppositiflora, Pall. Reise, i. 735 = Girgensohnia oppositiflora.
oppositifolia, Desf. Fl. Atlant. 219 = longifolia.
oppositifolia, Pall. Reise, ii. 477 = Petrosimonia sibirica.
oppositifolia, Sieber, ex Moq. in DC. Prod. xiii. II. 190 = Sieberi.
orientalis, S. G. Gmel. Reise, iv. 47. t. 5.—Cf. J. F. Gmel. Syst. 453 = rigida.
Pallasiana, Vest. ex Schult. Syst. vi. 240 = brachiata.
paniculata, Hort. ex Moq. in DC. Prod. xiii. II. 60 = Cycloloma platyphyllum.
papillosa, *Willk. Iber. Balbins.* 146.—Hispan.
Passerina, Bunge, in Linnaea, xvii. (1843) 4, in adnot. = gemmascens.
persica, *Bunge, ex Boiss. Fl. Orient.* iv. 964.—Persia.
physophora, Schrad. Neues Journ. iii. (1809) 88 = Suaeda physophora.
pilosa, Pall. Reise, ii. 477 = Halimocnemis malacophylla.
pinnatifida, *Schult. Syst.* vi. 243.—Aegypt.
platyphylla, Michx. Fl. Bor. Am. i. 174 = Cycloloma platyphyllum.
polyclonos, Georgi, Beschr. Russ. Reich. iv. 823 (Halimocnemidis sp. ?).—Ross. austr.
polyclonos, *Linn. Mant.* i. 54.—Hispan.; Sicil.
prostrata, Linn. Sp. Pl. 222 = Kochia prostrata.
prostrata, Pall. Reise, i. 490 = Kochia arenaria.
pulla, C. Koch, in Linnaea, xxii. (1839) 190 = brachiata.
pygmaea, Schrad. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda microsperma?
radiata, Desf. in Ann. Mus. Par. ii. (1803) 28. t. 34 = Cycloloma platyphyllum.
recurva, Wall. Cat. n. 6943 = Haloxylon recurvum.
Richleri, Karel. ex Moq. in DC. Prod. xiii. II. 185 = S. Arbuscula.
rigida, *Pall. Illustr.* 20. t. 12.—Oriens; As. bor.
rosacea, Bieb. Tabl. Prov. Casp. App. 143 = crassa.
rosacea, Cav. Ic. iii. 44. t. 286 = S. Kali.
rosacea, *Linn. Sp. Pl.* 222.—As. med.; Sibir.
rosea, S. G. Gmel. ex Nym. Consp. 631 = praec.
rubescens, *Franch. in Révoil, Çomali*, 60.—Afr. trop.
salsa, Cav. Ic. iii. 46. t. 290 = Suaeda spicata.
salsa, Jacq. Hort. Vindob. iii. 44. t. 83 = Suaeda maritima.
salsa, Linn. Sp. Pl. ed. II. 324 = Suaeda salsa.
salsa, Pursh, Fl. Am. Sept. i. 197 = Suaeda plattensis.
salsa, Roxb. in Beatson, Tracts St. Helena, 322 = Suaeda fruticosa.
sativa, Cav. Ic. iii. 46. t. 291 = Suaeda spicata.
sativa, Herb. Madr. ex Moq. in DC. Prod. xiii. II. 191 = Suaeda fruticosa.
sativa, Linn. Sp. Pl. ed. II. 323 = Halogeton sativus.
sativa, Wight, in Wall. Cat. n. 6945 = Suaeda maritima.
sativa, Zeyh. ex Moq. in DC. Prod. xiii. II. 131 = Kochia villosa.
scabra, Hort. ex Moq. l. c. 161 = Suaeda maritima.
scariosa, Stokes, Bot. Mat. Med. ii. 31 = S. Kali.
sclerantha, C. A. Mey. in Eichw. Pl. Casp. Cauc. 35. t. 27 = carinata.
scoparia, Bieb. in Mém. Soc. Nat. Mosc. i. (1811) 144 = Kochia scoparia.
sedifolia, Salisb. Prod. 153 = Suaeda fruticosa.
sedoides, Pall. Reise, i. 492 = Kochia sedoides.
sericea, Ait. Hort. Kew. i. ed. I. 317 = Kochia sericea.
seriphioides, *Stev. ex Moq. in DC. Prod.* xiii. II. 190.—Ins. Canar.
setifera, Lag. Gen. et Sp. Nov. 12 = Suaeda setigera.
setigera, Steud. Nom. ed. II. ii. 502 = praec.
sibirica, Koel. ex Moq. in DC. Prod. xiii. II. 213 = Anabasis brachiata.
Sieberi, *Presl, Bot. Bemerk.* 108.—Aegypt.
Sieversiana, Pall. ex Steud. Nom. ed. II. i. 848 = Kochia scoparia.
Soda, Brot. Fl. Lusit. i. 404 = grandiflora.
Soda, *Linn. Sp. Pl.* 223.—Europ. austr.; As. et Afr. bor.
Soda, Pall. Reise, ii. 493 = collina.
Soda, Weinm. in Flora, v. (1822) 300 = S. Kali.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 473.—Persia; Turkest.
songarica, Siev. ex Pall. Illustr. 35. sub t. 38 = Kochia scoparia.

SALSOLA :—
Souda, Loefl. Iter, 132 = Halogeton sativus.
spicata, Pall. Illustr. 27. t. 19 = glauca.
spicata, Willd. Sp. Pl. i. 1311 = Suaeda spicata.
spinescens, *Moq. in DC. Prod.* xiii. II. 179.—Arab.
spinescens, Wight, Ic. t. 1795 = foetida.
Spinifex, Pall. Illustr. 32. t. 24 = Noaea spinosissima.
Spinifex, Poir. Encyc. Suppl. v. 190 = Kochia sedoides.
spinosa, Lam. Fl. Fr. iii. 240 = S. Kali.
spissa, *Bieb. in Mém. Soc. Nat. Mosc.* i. (1811) 103.—Rossia; As. or. et bor.
splendens, Pourr. in Mém. Acad. Toul. iii. (1788) 327 = Suaeda setigera.
squarrosa, *Stev. ex Moq. in DC. Prod.* xiii. II. 190.—Ins. Naxos.
stellata, Hort. ex Moq. l. c. 60 = Cycloloma platyphyllum.
Stocksii, Boiss. Diagn. Ser. II. iv. 75 = Haloxylon recurvum.
stricta, Spreng. Syst. i. 293 = Chenopodium strictum.
strobilifera, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
Suaeda, Poir. Encyc. vii. 299 = Suaeda vera.
suaedoides, Moq. Chenop. Enum. 148 = spissa.
subaphylla, *C. A. Mey. in Eichw. Pl. Casp. Cauc.* 34. t. 24.—Reg. Casp.; Afghan.
subdecandra, Lag. Elench. Pl. Hort. Matr. (1813) 13, nomen (Quid?).—Hab.?
succulenta, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
tamariscifolia, *Cav. Ic.* v. 69.—Europ.
tamariscifolia, Falk, Beitr. ii. 141 = tamariscina.
tamariscifolia, Lag. Gen. et Sp. Nov. 12 = genistoides.
tamariscina, *Pall. Reise*, iii. 604.—Rossia; Reg. Cauc.; As. bor.
tenuifolia, Bieb. Fl. Taur. Cauc. i. 188 = Kochia arenaria.
tenuifolia, Falk, Beitr. ii. 141 = tamariscina.
tetragona, *Delile, Fl. Egypt*, 203. t. 21. f. 4.—Aegypt.; Arab.
tetrandra, Forsk. Fl. Aegypt. Arab. 58 = tetragona.
tomentosa, Spach, ex Moq. in DC. Prod. xiii. II. 208 = canescens.
Tournefortii, Spach, ex Moq. l. c. 209 = Noaea Tournefortii.
Tragus, Linn. Cent. Pl. ii. 13; Amoen. Acad. iv. 310 = S. Kali.
triandra, Poir. Encyc. Suppl. v. 190 = Kochia hyssopifolia.
triflora, Pers. Syn. i. 296 = Suaeda altissima.
trigyna, Hort. ex Steud. Nom. ed. II. ii. 503 = Suaeda maritima.
trigyna, Willd. Sp. Pl. i. 1312 = Suaeda altissima.
tuberculata, *Fenzl, ex Moq. in DC. Prod.* xiii. II. 178.—Afr. austr.
turgida, Dum. Fl. Belg. 23 = S. Kali.
ulicina, Bunge, in Bull. Acad. Pétersb. xxv. (1879) 364 = Horaninovia ulicina.
ulicina, *Trautv. in Act. Hort. Petrop.* i. (1871) 29.—Turkestan.
vera, Schult. Syst. vi. 242 = Suaeda vera.
vermiculata, Bieb. Tabl. Prov. Casp. App. 145 = gemmascens.
vermiculata, *Brot. Fl. Lusit.* i. 404.—Lusitan.
vermiculata, Hort. ex Moq. in DC. Prod. xiii. II. 161 = Suaeda maritima.
vermiculata, *Linn. Sp. Pl.* 223.—Reg. Medit.; As. bor.
vermiculata, Pall. Reise, i. 488 = rigida.
verrucosa, *Bieb. in Mém. Soc. Nat. Mosc.* i. (1811) 103.—Oriens; Ind. bor. occ.
verrucosa, C. A. Mey. Verz. Pfl. Cauc. 160 = gemmascens.
verticillata, Schousb. Vextr. Marokko, 123 = longifolia.
villosa, Delile, Illustr. n. 309; D. Dietr. Syn. Pl. ii. 997 = rigida.
villosa, Sieber, ex Boiss. Fl. Orient. iv. 955 = inermis.
Webbii, *Moq. Chenop. Enum.* 139.—Hispan.

SALTIA, R. Br. in Salt, Abyss. App. 64 (1814), nomen = **Cometes**, Linn. (Illecebr.).
abyssinica, A. Br. l. c. = C. abyssinica.

SALTIA, R. Br. in Wall. Pl. As. Rar. i. 17 (1830). *AMARANTACEAE*, Benth. & Hook. f. iii. 29.
globosa, Hochst. & Steud. ex Moq. in DC. Prod. xiii. II. 325 = Cyathula globulifera.
papposa, *R. Br. in Wall. Pl. As. Rar.* i. 17.—Ind. or.
Seddera, Moq. in DC. Prod. xiii. II. 325 = papposa.

SALTZWEDELIA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 480 (1800) = **Genista**, Linn. (Legumin.).
sagittalis, Gaertn. Mey. & Scherb. l. c. 498 = G. sagittalis.
tetragona, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = G. sagittalis.

SALUTIAEA, Colla, in Mem. Acc. Torin. Ser. II. x. (1849) 208 = **Achimenes**, P. Br. (Gesner.).
grandiflora, Colla, l. c. = A. grandiflora.

SALUTIEA, Griseb. Jahresb. 1850 (1853), 367; ex Pfeiff. Nom. ii. 1028 = praec.

SALVADORA, Garcin, ex Linn. Diss. Chen. 36 (1751); in Amoen. Acad. iii. 21. *SALVADORACEAE*, Benth. & Hook. f. ii. 681.
PELLA, Gaertn. Fruct. i. 143. t. 28. f. 8 (1788).
biflora, *Lour. Fl. Cochinch.* 88.—Cochinch.
capitulata, *Lour. l. c.* 87.—Cochinch.
crassinervia, Hochst. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 30 = persica.
glabra, Baill. Adansonia, ix. (1870) 290 = Dobera Roxburghii.
indica, Royle, Illustr. Bot. Himal. 319; Wight, Illustr. ii. 229 = persica.
madurensis, Decne. in Jacquem. Voy. Bot. 140 = Azima sarmentosa.
oleoides, Decne. l. c. 140. t. 144 = persica.
paniculata, Zucc. ex Steud. Nom. ed. II. ii. 503 = persica.
persica, *Linn. Sp. Pl.* 122.—Oriens; Ind. or.; Afr. bor.
persica, Wall. Cat. n. 1042 = Cansjera Rheedii.
Stocksii, Wight, Illustr. 229 = persica.
surinamensis, Spreng. f. Tent. Suppl. Syst. 7 = Cybianthus myrianthos.
Wightiana, Planch. ex Thw. Enum. Pl. Zeyl. 190 = persica.

SALVERTIA, A. St. Hil. in Mém. Mus. Par. vi. (1820) 266. *VOCHYSIACEAE*, Benth. & Hook. f. i. 977.
convallariaeodora, *A. St. Hil. l. c.*—Bras.
thyrsiflora, Pohl, Pl. Bras. Ic. ii. 16. t. 110 = convallariaeodora.

SALVIA, [Tourn.] Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1194.
AETHIOPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134.
AITOPSIS, BELOSPIS, CODANTHERA, CROLOCOS, ELELIS, ENIPEA, EPIADENA, EURIPLES, FLIPANTA, GLUTINARIA, HEMATODES, HEMISTEGIA, KIOSMINA, LARNASTYRA, LESEMIA, MEGYATHUS, MELLIGO, OBOSKON, ORMIASTIS, ORMILIS, PIARADENA, PLEUDIA, RHODORMIS, SOBISO, TEREPIS, Rafin. Fl. Tellur. iii. 89–94 (1836).
COVOLA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 227.
DRYMOSPHACE, Opiz, Seznam, 38 (1852).
GALLITRICHUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134.
HEMISPHACE, Opiz, l. c. 50 (1852).
HORMINUM, Moench, Meth. 376 (1794).
JUNGIA, Moench, l. c. 378 (1794).
LEONIA, La Llave & Lex. Nov. Veg. Desc. fasc. ii. 6 (1825).
PLETHIOSPHACE, Opiz, Seznam, 75 (1852).
MELINUM, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 200.
RHODOCHLAMYS, Schau. in Linnaea, xx. (1847) 706.
SCHRADERIA, Medic. Phil. Bot. ii. 40 (1791).
SCLAREA, Moench, Meth. 374 (1794).
STENARRHENA, D. Don, Prod. Fl. Nep. 111 (1825).
STIEFIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 200; Phil. Bot. ii. 40 (1791).
abyssinica, R. Br. in Salt, Voy. Abyss. App. 62 = Meriandra benghalensis.
abyssinica, Jacq. Ic. Pl. Rar. i. 2 = nilotica.
abyssinica, Linn. f. Suppl. 88 = nudicaulis, spinosa.
acaulis, Vahl, Enum. i. 257 = lyrata?
acetabulosa, *Linn. Mant.* i. 25.—As. Min.; Syria.
acuminata, *Ruiz & Pav. Fl. Per.* i. 24. t. 37.—Peruv.
acuminata, Vent. Jard. Cels, t. 50 = azurea.
acuminatissima, Steud. Nom. ed. II. ii. 503 = azurea.
acutata, Brot. Phyt. Lusit. i. 203 = S. Verbenaca.
acutifolia, Lam. Illustr. i. 68 = nutans.
acutifolia, *Ruiz & Pav. Fl. Per.* i. 24. t. 38.—Peruv.
adglutinans, *Lag. Gen. et Sp. Nov.* 2.—Mexic.

SALVIA:—
aegyptiaca, Linn. Mant. i. 26 = spinosa.
aegyptiaca, *Linn. Sp. Pl.* 23.—Geront. trop. et subtrop.
Aethiopis, Brot. Fl. Lusit. i. 18 = argentea.
Aethiopis, *Linn. Sp. Pl.* 27.—Europ. austr.; Oriens.
affinis, *Schlecht. & Cham. in Linnaea*, v. (1830) 99.—Mexic.
affinis, Spreng. ex Steud. Nom. ed. I. 724 = virgata.
africana, *Linn. Sp. Pl.* ed. II. 38.—Afr. austr.
agglutinans, Roem. & Schult. Syst. i. 582 = adglutinans.
agrestis, Linn. Amoen. Acad. iii. 399 = pratensis.
agrestis, *Vell. Fl. Flum* 14; i. t. 35.—Bras.
agrestis, Vill. Hist. Pl. Dauph. ii. 402 = S. Verbenaca.
albicaulis, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 234.—Afr. austr.
albida, Jacq. Obs. 10 = candidissima.
albiflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 76.—Mexic.
albo-caerulea, *Linden, in Belg. Hortic.* vii. (1857) 199.—Mexic.
algeriensis, *Desf. Fl. Atlant.* i. 23. t. 3.—Afr. bor.
alpestris, Hort. ex Benth. Lab. Gen. et Sp. 237 = sylvestris.
altissima, *Pohl, Pl. Bras. Ic.* ii. 140. t. 195.—Bras.
Alvajaca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 38.—Costa Rica.
amabilis, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 13 = farinacea.
amara, Jacq. Hort. Schoenb. iii. 2 = amarissima.
amarissima, *Orteg. Hort. Matr. Dec.* 4.—Mexic.
ambigua ×, Čelak. in Arch. Naturw. Landesd. Boehm. ii. (1873) 353.—Europ.
ambigua, Rochebr. & Savat. Cat. Pl. La Charente, 167, in obs. = S. Verbenaca?
ambigua, Salisb. Prod. 74 = nubia.
amethystina, Salisb. l. c. = mexicana.
amethystina, *Sm. Ic. Ined.* ii. 27. t. 27.—N. Granat.
amoena, Sims, Bot. Mag. t. 1294 = lamiifolia.
amplexicaulis, Hort. ex Benth. Lab. Gen. et Sp. 236 = virgata.
amplexicaulis, Lam. Illustr. i. 68 = sylvestris.
angulata, *Benth. Lab. Gen. et Sp.* 721.—Venezuela.
angustifolia, *Cav. Ic.* iv. 9. t. 317.—Mexic.
angustifolia, Michx. Fl. Bor. Am. i. 15 = azurea.
angustifolia, Salisb. Prod. 73 = dentata.
antirrhiniflora, *Dehnh. Cat. Pl. Hort. Camald.* 27.—Hab.?
applanata, Willd. Enum. Hort. Berol. Suppl. 3 = nilotica.
aprica, Dupuy, ex Nym. Consp. 569 = pratensis.
aprica, Schur, Enum. Pl. Transs. 522 = nemorosa.
arachnoidea, Noé, ex Walp. Ann. iii. 256 = phlomoides.
arenaria, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 257.—Bras.
argaea, Boiss. & Bal. ex Boiss. Fl. Orient. iv. 601 = recognita.
argentea, Hort. ex Benth. Lab. Gen. et Sp. 223 = candidissima.
argentea, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 773 = officinalis.
argentea, *Linn. Sp. Pl.* ed. II. 38.—Reg. Mediterr.
arguta, Benth. Lab. Gen. et Sp. 283 = salicifolia.
arida, Salisb. Prod. 73 = aegyptiaca.
aristata, *Auch. ex Benth. in DC. Prod.* xii. 270.—Persia.
aristulata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 67.—Mexic.
arizonica, *A. Gray, Syn. Fl. N. Am.* ii. 1. 370.—Arizona.
armena, C. Koch, in Linnaea, xxi. (1848) 654 = candidissima.
Arrabidae, *Steud. Nom.* ed. II. ii. 503.—Bras.
asiodophylla, Steud. l. c., sphalm. = *aspidophylla*.
aspera, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 71.—Mexic.
asperata, *Falc. ex Benth. in DC. Prod.* xii. 282.—Reg. Himal.
asperifolia, *Benth. Lab. Gen. et Sp.* 281.—Reg. Himal.
asperula, Hort. ex Benth. l. c. 237 = sylvestris.
aspidophylla, Schult. Mant. i. 206 = lanceolata.
assurgens, *H. B. & K. Nov. Gen. et Sp.* ii. 293.—Mexic.
atlantica, Pers. Syn. i. 29 = argentea.
atomaria, Boiss. Diagn. Ser. II. iv. 18 = verbascifolia.

SALVIA:—
atropatana, *Bunge, Lab. Pers.* 47.—Persia.
Aucheri, *Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 38.—As. Min.
aurasiaca, Pomel, Nouv. Mat. Fl. Atl. 306 = argentea.
aurea, *Linn. Sp. Pl.* ed. II. 38.—Afr. austr.
auriculata, Mill. Gard. Dict. ed. VIII. n. 3 = triloba.
aurita, Schult. Obs. Bot. 10 = officinalis.
aurita, *Thunb. Prod. Pl. Cap.* 96.—Afr. austr.
austriaca, *Jacq. Fl. Austr.* ii. 8.—Europ. austr. or.
austriaca, C. Koch, in Linnaea, xxi. (1848) 656 = staminea.
axillaris, *Moç. & Sesse, ex Benth. Lab. Gen. et Sp.* 270.—Mexic.
ayavacensis, Schult. Mant. i. 195, sphalm. = seq.
ayavazensis, *H. B. & K. Nov. Gen. et Sp.* ii. 298.—Peruv.
azurea, Hort. ex Benth. Lab. Gen. et Sp. 299 = lamiifolia.
azurea, *Lam. Journ. Hist. Nat.* i. (1792) 409.—Am. bor.
baccifera, Etling. Salv. 18 = triloba.
bachtiarica, Bunge, Lab. Pers. 47 = atropatana.
baetica, Boiss. Elench. 73 = bullata.
Balansae, *Noé, ex Coss. Notes Crit.* 139.—Afr. bor.
balaustina, *Pohl, Pl. Bras. Ic.* ii. 133. t. 189.—Bras.
ballotaeflora, *Benth. Lab. Gen. et Sp.* 270.—Texas; Mexic.
barbata, Lam. Illustr. i. 72 = africana.
Barrelieri, Etling. Salv. 46 = inamoena.
Barrelieri, Hort. ex Benth. Lab. Gen. et Sp. 235 = virgata.
Barrelieri, Tenore, Fl. Nap. i. 9. t. 2 = pratensis.
Bauhini, Hort. ex Benth. Lab. Gen. et Sp. 235 = virgata.
Baumgartenii, Heuff. ex Griseb. & Schenk, in Wiegm. Arch. xviii. (1852) 328 = pratensis.
bavarica, Schrank, Baier. Fl. ii. 133 = austriaca?
Beckeri, *Trautv. in Act. Hort. Petrop.* iii. II. (1875) 276.—Reg. Caucas.
bengalensis, Roxb. Hort. Beng. 4; Fl. Ind. i. 145 = Meriandra benghalensis.
Benthamiana, Boiss. ex Benth. in DC. Prod. xii. 269 = Russeggeri.
Benthamiana, *Gardn. & Field. Sert. Pl.* t. 19.—Bras.
Benthamii, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 187 = brevipes.
Bernardina ×, Parish, ex Greene, in Bull. Calif. Acad. i. IV. (1885) 211.—Calif.
Bertolonii, *Vis. Fl. Dalm.* ii. 189.—Europ. austr.
Bethelli, Hort.—Cf. Gard. Chron. (1881) I. 51 = involucrata.
Betonica, Schult. Mant. i. 188 = elongata.
betonicaefolia, Etling. Salv. 48 = pendula.
betonicaefolia, Lam. Illustr. i. 70 = S. Verbenaca.
bicolor, *Lam. l. c.* 69.—Reg. Mediterr. occ.
bifida, Forsk. Fl. Aegypt. Arab. 202 = Forskohlei.
biflora, Ruiz & Pav. Fl. Per. i. 24 = tubiflora.
biserrata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 66.—Mexic.
Blancoana, Webb & Heldr. ex Walp. Ann. iii. 254 = S. Candelabrum.
Blodgettii, *Chapm. Fl. S. U. St.* 319.—Am. bor.
Bodeana, *Bunge, Lab. Pers.* 42.—Persia.
bogotensis, *Benth. in DC. Prod.* xii. 312.—N. Granat.
boliviana, *Van Houtte, in Fl. des Serres*, xi. (1856) 151.—Bolivia.
Bolleana, *Noé, ex Bolle, in Bonplandia*, viii. (1860) 284.—Ins. Canar.
Bonplandiana, F. G. Dietr. Vollst. Lexic. Gaertn. Nachtr. vii. 403 = discolor.
Boosiana, Jacq. f. Eclog. i. 69. t. 47 = lamiifolia.
Boosii, Tratt. Arch. Gew. i. 58. t. 92 = lamiifolia.
Boucheana, *Kunth, Ind. Sem. Hort. Berol.* (1845) 9.—Am. trop.
Bourgeana, *W. Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1885) 96.—Lycia.
brachiata, Roxb. Hort. Beng. 4; Fl. Ind. i. 145 = plebeia.
brachycalyx, *Boiss. Fl. Orient.* iv. 625.—As. Min.; Syria.
brachypoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = brevipes.
bracteata, Poir. Encyc. vi. 622 = hirsuta.
bracteata, Sims, Bot. Mag. t. 2320 = S. Sclarea.
bracteata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 242.—As. Min.; Syria.
brasiliensis, Spreng. Syst. i. 56 = splendens.

SALVIA :—
brevicalyx, *Benth. in DC. Prod.* xii. 309.—Mexic.
breviflora, Moç. & Sesse, ex Benth. Lab. Gen. et Sp. 274 = spicata.
brevipes, *Benth. in DC. Prod.* xii. 321.—Bras.
Broussonetii, *Benth. Lab. Gen. et Sp.* 227.—Ins. Canar.; Afr. bor.
Broussoneti, Bolle, in Bonplandia, viii. (1860) 284 = Bolleana.
bullata, Orteg. Hort. Matr. Dec. 109 = tenella.
bullata, *Vahl, Enum.* i. 265.—Hispan.; Lusitan.
bullulata, *Benth. in DC. Prod.* xii. 327.—Peruv.
bupleuroides, *Presl, ex Benth. Lab. Gen. et Sp.* 271.—Mexic.
byzantina, Hort. ex Benth. l. c. 240 = S. Verbenaca.
byzantina, *Juss. ex Steud. Nom.* ed. I. 725.—Turcia.
cabulica, *Benth. in DC. Prod.* xii. 268.—Afghan.
cacaliaefolia, *Benth. l. c.* 348.—Mexic.
cadmica, *Boiss. Diagn.* Ser. I. v. 6.—As. Min.
caduca, Vahl, ex Hornem. Hort. Hafn. i. 30 = virgata.
caerulea, *Linn. Sp. Pl.* 26.—Afr. austr.
caesia, Scheele, in Linnaea, xxii. (1849) 588 = farinacea.
caesia, Willd. Enum. Hort. Berol. 40 = polystachya.
caespitosa, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 39.—As. Min.
calaminthaefolia, *Vahl, Enum.* i. 233.—Ins. S. Doming.
Calverti, Boiss. Fl. Orient. iv. 626 = staminea.
calycina, *Sibth. & Sm. Fl. Graec.* i. 13. t. 16.—Graecia.
calycinflata, Moç. & Sesse, ex Benth. Lab. Gen. et Sp. 288 = Sessei.
camaraefolia, *Benth. in DC. Prod.* xii. 342.—N. Granat.
Camertoni, Regel, in Bot. Zeit. xi. (1853) 334 = elegans?
campanulata, *Wall. ex Benth. in Wall. Pl. As. Rar.* i. 67.—Reg. Himal.
campestris, Bieb. Fl. Taur. Cauc. i. 20 = virgata.
cana, Wall. Cat. n. 2145 = lanata.
canariensis, *Linn. Sp. Pl.* 26.—Ins. Canar.
candelabriformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Candelabrum.
Candelabrum, *Boiss. Elench.* 72.—Hispan.
candicans, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 61.—Mexic.
candidissima, Bieb. Fl. Taur. Cauc. i. 22 = verbascifolia.
candidissima, Guss. Fl. Sicil. Prod. i. 16 = argentea.
candidissima, *Vahl, Enum.* i. 278.—As. Min.
Candollei, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* i. (1872–73) 285.—Gallia.
canescens, Benth, Lab. Gen. et Sp. 250 = pallida.
canescens, *C. A. Mey. Verz. Pfl. Cauc.* 86.—Reg. Caucas.
capitata, *Schlecht. in Linnaea*, xxvi. (1853–55) 292.—Mexic.
Capusii, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 232.—Turkestan.
cardinalis, H. B. & K. Nov. Gen. et Sp. ii. 301. t. 152 = fulgens.
cardiophylla, *Benth. Lab. Gen. et Sp.* 721.—Am. austr.
carduacea, *Benth. l. c.* 302.—Calif.
carnea, *H. B. & K. Nov. Gen. et Sp.* ii. 300.—N. Granat.
carnosa, Dougl. ex Benth. in Bot. Reg. sub t. 1469 = Audibertia incana.
cataonica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 602.—As. Min.
catarioides, *Loisel. ex Drapiez, Herb. Amat.* vi. (1833). t. 390.—Hab.?
caucasica, Schrank, in Syll. Ratisb. ii. (1828) 58 = virgata.
Cedronella, *Boiss. Diagn.* Ser. I. v. 3.—Graecia.
cedrosensis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 212.—Calif.
ceratophylla, *Linn. Sp. Pl.* 27.—As. Min.; Persia.
ceratophylla, C. A. Mey. Verz. Pfl. Cauc. 68 = S. Verbenaca.
ceratophylla, Tenore, Syll. Fl. Neap. 21 = ceratophylloides.
ceratophylloides, Forsk. Fl. Aegypt. Arab. p. lix = lanigera.
ceratophylloides, *Linn. Mant.* i. 26.—Sicilia; Aegypt.
ceratophylloides, Vahl, Enum. i. 279 = S. Verbenaca.
cespitosa, Montb. & Auch. = caespitosa.

SALVIA :—
chamaedrifolia, Andr. Bot. Rep. t. 416 = chamaedryoides.
chamaedryoides, *Cav. Ic.* ii. 77. t. 197.—Texas; Mexic.
Chamaedrys, Willd. Hort. Berol. i. 29 = chamaedryoides.
Chamaeelaeagnea, Roem. & Schult. Syst. i. 634, sphalm. = seq.
Chamelaeagnus, Burm. f. Fl. Cap. Prod. 1 = paniculata.
Chapmani, *A. Gray, Syn. Fl. N. Am.* ii. I. 370.—Am. bor.
chilensis, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1857) 45.—Chili.
chinensis, *Benth. Lab. Gen. et Sp.* 725.—China.
chionantha, *Boiss. Diagn.* Ser. I. v. 8.—As. Min.
chnoodes, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 98.—Lycia; Mesopot.
chorassanica, *Bunge, Lab. Pers.* 45.—Persia.
chromatica, Hoffmgg. Verz. Pfl. 194 = officinalis.
chrysantha, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 72.—Mexic.
chrysophila, J. A. Schmidt, in Mart. Fl. Bras. viii. 187 = rigida.
chrysophylla, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 96.—Lycia.
ciliata, Benth. Lab. Gen. et Sp. 286 = coccinea.
ciliata, Hort. ex Poir. Encyc. vi. 588 = hirsuta.
ciliata, Rafin. Fl. Ludov. 41 (Quid?).—Am. bor.
cilicica, Boiss. Diagn. Ser. II. iv. 19 = crassifolia.
cinnabarina, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 63.—Mexic.
circinnata, Cav. Ic. iv. 9. t. 318 = amarissima.
clandestina, Linn. Sp. Pl. ed. II. 36 = S. Verbenaca.
clandestina, Sibth. & Sm. Fl. Graec. Prod. i. 16. t. 24 = controversa.
clandestinoides, Link, Jahrb. i. III. (1820) 169 = S. Verbenaca.
clausa, *Vell. Fl. Flum.* 13; i. t. 33.—Bras.
Claytoni, M. A. Curt. ex A. Gray, Syn. Fl. N. Am. ii. I. 370 = urticifolia.
Claytoni, Ell. Sketch, i. 32 = S. Verbenaca.
Claytoni, *Nutt. Gen. Am.* 17.—Am. bor.
cleistogama, *De Bary & Paul, Ind. Sem. Hort. Halens.* (1867) 6.—*Cf. Bot. Zeit.* xxix. (1871) 555.—Afr. austr.
clinopodioides, *H. B. & K. Nov. Gen. et Sp.* ii. 294. t. 145.—Mexic.
Clusii, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 773 = officinalis.
Clusii, Jacq. Hort. Schoenb. ii. 37 = triloba.
Clusii, Timb. in Mém. Acad. Toul. Sér. VII. ii. (1870) 237 = pratensis.
coarctata, Vahl, Enum. i. 253 = foetida.
coccinea, *Buc'hoz, ex Etling. Salv.* 23 (an seq.?).—Hab.?
coccinea, *Juss. ex Murr. in Comm. Gotting.* i. (1778) 86. t. 1.—Am. bor. et trop.
coelestina, *Carr. in Rev. Hortic.* (1878) 49.—Mexic.
coerulea, *Benth. Lab. Gen. et Sp.* 298.—Bras.
coerulea, Hort. ex Benth. l. c. 299 = lamiifolia.
collina, *H. B. & K. Nov. Gen. et Sp.* ii. 288.—Ecuador.
collina, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 18 = S. Verbenaca.
collina, Salisb. Prod. 73 = sylvestris.
collina, Schur, Enum. Pl. Transs. 522 = nemorosa.
colorans, Hort. ex Benth. Lab. Gen. et Sp. 287 = splendens.
colorata, Linn. Syst. ed. XII. 66 = aurea.
colorata, Thore, Chlor. Land. 17 = S. Horminum.
colorata, Vahl, Enum. i. 230 = africana.
Columbariae, *Benth. Lab. Gen. et Sp.* 302.—Am. bor. occ.
commutata, Benth. l. c. 222 = dominica.
comosa, *Peyr. in Linnaea*, xxx. (1859–60) 32.—Mexic.
comosa, Salisb. Prod. 73 = S. Horminum.
complanata, Wall. ex Steud. Nom. ed. II. ii. 504, sphalm. = campanulata.
compressa, *Vent. Jard. Cels*, 59.—Persia; Afghan.
concolor, *Lamb. ex Benth. Lab. Gen. et Sp.* 297.—Mexic.
confertiflora, *Pohl, Pl. Bras. Ic.* ii. 134. t. 190.—Bras.
confusa, *Benth. Lab. Gen. et Sp.* 210.—Europ. austr.
congesta, *A. Rich. Tent. Fl. Abyss.* ii. 194.—Abyss.
Conradi, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 97.—Caria.
controversa, *Tenore, Syll. Fl. Neap.* 18.—Reg. Mediterr.

SALVIA :—
cordata, *Benth. Lab. Gen. et Sp.* 268.—Bras.
coriifolia, Scheele, in Linnaea, xxii. (1839) 587 = azurea.
corrugata, *Vahl, Enum.* i. 252.—Peruv.
corymbosa, Heyne, ex Wall. Cat. n. 2149 = plebeia.
costaricensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 89.—Costa Rica.
crassifolia, *Desf. Arb.* i. 134.—Europ. austr.
crassifolia, Jacq. Fragm. 47. t. 60 = bicolor.
crassifolia, *Sibth. & Sm. Fl. Graec.* i. 19. t. 26.—As. Min.
cremenecensis, Bess. Enum. Pl. Volh. 40 = pendula.
crenata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 74.—Mexic.
cretica, *Linn. Sp. Pl.* 23.—Hispan.; Creta.
crispula, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 234.—Afr. austr.
Cruckshanksii, *Benth. Lab. Gen. et Sp.* 261 (Cruikshanksii).—Peruv.
cryptantha, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 40.—As. Min.
cryptanthos, Schult. Obs. Bot. 12 = hirsuta.
cryptoclada, *Baker, in Journ. Linn. Soc.* xviii. (1881) 275.—Madag.
cuneifolia, *Benth. Lab. Gen. et Sp.* 270.—Mexic.
cuphaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 288.—Peruv.
curviflora, Benth. Lab. Gen. et Sp. 284.—Mexic.
cuspidata, *Ruiz & Pav. Fl. Per.* i. 23. t. 40.—Peruv.
cyanea, *Benth. Lab. Gen. et Sp.* 296.—Mexic.
cyanea, Fisch. Cat. Hort Gorenk. 21 = lamiifolia.
cyanescens, *Boiss. & Bal. Diagn.* Ser. II. iv. 19.—Armenia.
cyanifera, Otto, ex Benth. in DC. Prod. xii. 336 = cyanea.
cyaniflora, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 301 = cyanea.
cyanocalyx, *Boiss. in Tchihat. Asie Min. Bot.* ii. 148, nomen.—As. Min.
cypria, *Ung. & Kotschy, Ins. Cyp.* 266.—Ins. Cyprus.
dasyantha, *Lem. in Rev. Hortic.* Sér. IV. viii. (1859) 438.—N. Granat.
decipiens, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 64 = patens.
deltoidea, Pers. Syn. i. 23 = S. Regla.
densiflora, *Benth. Lab. Gen. et Sp.* 721.—Ins. Dominic.?
dentata, *Ait. Hort. Kew.* ed. I. i. 37.—Afr. austr.
derasa, *Benth. Pl. Hartw.* 242.—N. Granat.
deserta, Schangin, in Ledeb. Ind. Sem. Hort. Dorpat. (1824) 6 = sylvestris.
deserti, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 248.—Arab.
desertorum, J. A. Schmidt, in Mart. Fl. Bras. viii. 189 = scabrida.
dianthera, Roth, ex Roem. & Schult. Syst. i. 263 = Meriandra benghalensis.
dichroa, Hook. f. Bot. Mag. t. 6004 = bicolor.
dichroantha, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 96.—Lycia.
dichroma, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 82.—Mexic.
dielytroides, Roezl, ex Van Houtte, in Fl. des Serres, xxiv. (1861) 31 = Roezli.
digyna, Stokes, Bot. Mat. Med. i. 58 = cretica.
discolor, *H. B. & K. Nov. Gen. et Sp.* ii. 294.—Peruv.
disermas, *Linn. Sp. Pl.* ed. II. 36.—Syria.
disermas, Sibth. & Sm. Fl. Graec. Prod. i. 16 = S. Verbenaca.
distans, Pohl, ex Steud. Nom. ed. I. 725 = austriaca.
divaricata, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 37.—Cappadoc.
diversifolia, Benth. in DC. Prod. xii. 274, in syn. = nivea.
diversifolia, Miq. Ann. Mus. Bot. Lugd. Bot. ii. 108 = chinensis?
doliostachys, Lag. ex Benth. Lab. Gen. et Sp. 724 = pulchella.
dominica, *Linn. Sp. Pl.* 25.—Syria.
dominica, Sw. Obs. Bot. 18. t. 1. f. 1 = serotina.
dracocephaloides, *Boiss. Diagn.* Ser. I. v. 4.—Persia.
Dregeana, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 234.—Afr. austr.
dubia, C. Koch, in Linnaea, xxi. (1848) 657 = pratensis.
dubia, Lowe, in Hook. Journ. Bot. i. (1834) 37 = S. Verbenaca.

SALVIA:—
dumetorum, *Andrz. in Bess. Enum. Pl. Volh.* 3, 40. —Europ. bor.; As. bor.;
Eckloniana, *Benth. in DC. Prod.* xii. 273.—Afr. austr.
elata, Host, Fl. Austr. i. 24 = sylvestris.
elata, Poir. Encyc. vi. 625 = azurea.
elegans, *Vahl, Enum.* i. 238.—Mexic.
elongata, *H. B. & K. Nov. Gen. et Sp.* ii. 287. t. 139. —Mexic.
elongata, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 75 = protracta.
elongata, Salisb. Prod. 74 = indica.
elongata, Spreng. Pugill. i. 3 = bullata.
elongata, Torr. in Ann. Lyc. N. York, ii. (1828) 231 = azurea.
elsholtzioides, Benth. Bot. Voy. Sulph. 152. t. 50 = hyptoides.
Engelmanni, A. Gray, in Proc. Am. Acad. viii. (1870) 368 (Salviastri sp.).—Texas.
Engelmanni, Hort. ex Schlecht. in Linnaea, xxvii. (1854) 504 = Roemeriana.
eremetica, Cerv. ex Lag. Gen. et Sp. Nov. 2 = polystachya.
eremophila, *Boiss. Diagn.* Ser. I. v. 12.—Persia.
eriocalyx, *Bert. ex Roem. & Schult. Syst.* i. *Addit.* II. 246.—Jamaic.
eriophora, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 611.—Cilicia.
erosa, Desf. Tabl. ed. II. 68; et ex Poir. Encyc. Suppl. v. 49 = S. Verbenaca.
euphratica, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 40.—Cappadoc.
exasperata, Cav. in Anal. Hist. Nat. ii. (1800) 113. t. 17 = ceratophylloides.
excelsa, *Benth. ex Lindl. Bot. Reg.* (1841) *Misc.* 90. —Mexic.
excisa, Ruiz & Pav. Fl. Per. i. 25 = tubiflora.
exserta, *Griseb. in Goett. Abh.* xxiv. (1879) 274.—Reg. Argent.
farinacea, *Benth. Lab. Gen. et Sp.* 274.—Texas; Mexic.
farinosa, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 78 = purpurea.
filamentosa, Tausch, in Flora, xxv. (1842) 282; Regel, Gartenfl. vii. (1858) t. 232 = coccinea.
filipes, *Benth. in DC. Prod.* xii. 309.—Mexic.
fimbriata, H. B. & K. Nov. Gen. et Sp. ii. 299. t. 149 = tiliaefolia.
flavescens, *Juss. ex Steud. Nom.* ed. I. 726.—Hab.?
flexuosa, *Presl, ex Benth. Lab. Gen. et Sp.* 248.— Mexic.
flexuosa, Schrank, in Syll. Ratisb. ii. (1828) 57 = limbata.
flocculosa, *Benth. Pl. Hartw.* 244.—Ecuador.
floribunda, *Brongn. in Lem. Hortic. Univ.* iii. (1841) 296.—Hab.?
florida, *Benth. in DC. Prod.* xii. 338.—Peruv.
foetida, *Lam. Illustr.* i. 69.—Marocc.
foliosa, Benth. in Bot. Reg. t. 1429 = rhombifolia.
fontinalis, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 177.—Ind. or.
formosa, L' Hérit. Stirp. Nov. 41 = leonuroides.
Forskählei, Linn. Mant. i. 26 = seq.
Forskohlei, *Linn. Syst.* ed. XII. 67.—As. Min.
Fortunii, *Benth. in DC. Prod.* xii. 354.—China.
fragifera, Etling, Salv. 19. n. 5 = pomifera.
frigida, *Boiss. Diagn.* Ser. I. v. 10.—As. Min.
frigida, Boiss. Fl. Orient. iv. 620 = xanthocheila.
frugifera, Benth. Lab. Gen. et Sp. 215 = pomifera.
fruticetorum, *Benth. l. c.* 284.—Bras.
fruticosa, Mill. Gard. Dict. ed. VIII. n. 5 = triloba.
fruticulosa, *Benth. Lab. Gen. et Sp.* 721.—Mexic.
fruticum, Vukot.—Cf. Oestr. Bot. Zeitschr. xxx. (1880) 163 = grandiflora.
fulgens, *Cav. Ic.* i. 15. t. 23.—Mexic.
Fuschii ×, Timb. in Mém. Acad. Toul. Sér. VII. ii. (1870) 213.—Gallia.
Gaillardoti, Boiss. Fl. Orient. iv. 628 = viscosa.
galatica, *C. Koch, in Linnaea*, xxi. (1848) 653.— Galatia.
galeata, *Ruiz & Pav. Fl. Per.* i. 27. t. 42.—Peruv.
Galeottii, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 75 = coccinea.
garganica, Tenore, Ind. Sem. Hort. Neap. (1829); ex ej. Syll. Fl. Neap. 18 = virgata.
gariepensis, *E. Mey. Comm. Pl. Afr. Austr.* 232.— Afr. austr.
Gerardiana, Benth. in Wall. Cat. n. 2150 = coccinea.

SALVIA:—
gesneraeflora, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 49. t. 47.—N. Granat.
gesneriaefolia, Lindl. ex Lem. Jard. Fleur. ii. (1852) t. 179, sphalm. = praec.
gigantea, Desf. Tabl. ed. II. 68; Cat. Hort. Par. ed. III. 394 = virgata.
Gilliesii, *Benth. Lab. Gen. et Sp.* 265.—Chili.
glabra, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 68.—Mexic.
glabrata, *H. B. & K. Nov. Gen. et Sp.* ii. 299.— Mexic.
glandulifera, *Cav. in Anal. Hist. Nat.* ii. (1800) 111. —Ecuador.
glaucescens, Pohl, Pl. Bras. Ic. ii. t. 192 = coccinea.
glechomaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 290. t. 141.—N. Mexic.
glumacea, *H. B. & K. l. c.* 298.—Mexic.
glutinosa, *Linn. Sp. Pl.* 26.—Europ. austr.; Oriens; Reg. Himal.
gossypina, Benth. Pl. Hartw. 330 = carduacea.
Goudotii, *Benth. in DC. Prod.* xii. 339.—N. Granat.
graciliflora, Avé-Lall. Ind. Sem. Hort. Petrop. x. 57 = aurita.
graciliflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 75.—Mexic.
gracilis, *Benth. Lab. Gen. et Sp.* 258.—Mexic.
Grahami, *Benth. in Bot. Reg.* t. 1370.—Mexic.
grandidentata, Tenore, Ind. Sem. Hort. Neap. (1829); ex Benth. Lab. Gen. et Sp. 519 = virgata.
grandiflora, *Etling. Salv.* 17.—As. Min.
grandiflora, Hornem. Hort. Hafn. i. 27 = pendula.
grandiflora, Née, ex Cav. Ic. v. 33 = patens.
grandiflora, Tenore, Fl. Nap. Prod. App. v. 3 = officinalis.
grandiflora, Vukot.—Cf. Oestr. Bot. Zeitschr. xxx. (1880) 163 (Quid?).—Croatia.
granitica, *Hochst. in Flora*, xxviii. (1845) 65.—Afr. austr.
grata, Vahl, Enum. i. 244 = oppositiflora.
graveolens, Vahl, l. c. 273 = dominica.
Greggii, *A. Gray, in Proc. Am. Acad.* viii. (1870) 369.—Texas; Mexic.
guaranitica, *A. St. Hil. ex. Benth. Lab. Gen. et Sp.* 298.—Bras.
Gundelia, *C. Koch, in Linnaea*, xxi. (1848) 653.— Oriens.
Gussonii, Boiss. ex Nym. Consp. 569 = argentea.
Hablitziana, *Pall. Tabl. Taur.* 44; *et ex Clarke, Trav.* i. 572.—Tauria.
Habliziana, Willd. Sp. Pl. i. 129 = scabiosaefolia.
haematodes, Linn. Sp. Pl. 24 = pratensis.
haematodes, Scop. Fl. Carn. ed. II. i. 29 = S. Sclarea.
Haenkei, *Benth. Lab. Gen. et Sp.* 283.—Peruv.
halaënsis, *Vicary, in Ann. & Mag. Nat. Hist.* Ser. II. i. (1848) 432.—Ind. or.
Hallieri, *Fisch. ex Steud. Nom.* ed. I. 726.—Hab.?
hastaefolia, *E. Mey. Comm. Pl. Afr. Austr.* 233.— Afr. austr.
hastata, Etling. Salv. 48 = nutans.
Hausknechtii, *Boiss. Fl. Orient.* iv. 605.—Cataonia.
Heerii, *Regel, Gartenfl.* iv. (1855) 77. t. 115.— Peruv.
Heldreichiana, Boiss. ex DC. Prod. xvii. 320 = Russeggeri.
helianthemifolia, *Benth. Lab. Gen. et Sp.* 254.— Mexic.
Henryi, *A. Gray, in Proc. Am. Acad.* viii. (1870) 368.—N. Mexic.
herbacea, *Benth. Lab. Gen. et Sp.* 720.—Mexic.
Hermani ×, Timb. ex Nym. Consp. 570.—Europ.
heterophylla, Hort. ex Benth. in DC. Prod. xii. 294 = S. Verbenaca.
hians, *Royle, ex Benth. in Hook. Bot. Misc.* iii. (1833) 373.—Reg. Himal.
hiemalis, Brot. Phyt. Lusit. ii. 3 = S. Verbenaca.
hierosolymitana, *Boiss. Diagn.* Ser. I. xii. 61.—Syria.
Hilarii, *Benth. Lab. Gen. et Sp.* 282.—Bras.
hirsuta, *Jacq. Hort. Schoenb.* iii. 1. t. 252.—Mexic.
hirta, *H. B. & K. Nov. Gen. et Sp.* ii. 296.—Peruv.
hirta, Schrank, in Syll. Ratisb. ii. (1828) 60 = scorodoniaefolia.
hirtella, *Vahl, Enum.* i. 249.—Peruv.
hispanica, Ettling. ex Willk. & Lange, Prod. Fl. Hisp. ii. 420 = officinalis.
hispanica, *Linn. Sp. Pl.* 25.—Am. trop.
hispanorum, Lag. Gen. et Sp. Nov. 1 = lavandulaefolia.

SALVIA:—
horminioides, Pourr. in Mem. Acad. Toul. iii. (1788) 327 = S. Verbenaca.
Horminum, *Linn. Sp. Pl.* 24.—Reg. Mediterr.
Hoveyi, Hort. ex Gard. Chron. (1880) II. 695 = splendens.
Humboldtiana, F. G. Dietr. Vollst. Lexic. Gaertn. vii. 418 = spicata.
Humboldtiana, Schult. Mant. i. 183 = lavanduloides.
Humboldtii, Spreng. Syst. i. 64 = spicata.
humilis, *Benth. Lab. Gen. et Sp.* 247.—Mexico.
hybrida ×, Schur, Enum. Pl. Transs. 522.—Europ.
Hydrangea, *DC. ex Benth. Lab. Gen. et Sp.* 717.— Persia.
hyemalis, Auct. = *hiemalis*.
hyemalis, Boose, ex A. Dietr. Sp. Pl. i. 276 = lamiifolia.
hypanica, Andrz. n Bess. Enum. Pl. Volh. 3 = virgata.
hypargeia, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 34 = Montbretii.
hypochionaea, *Boiss. Diagn.* Ser I. v. 8.—Oriens.
hypochionea, Buhse, ex Boiss. Fl. Orient. iv. 620 = atropatana.
hypoleuca, *Benth. in DC. Prod.* xii. 279.—Persia.
hypoleuca, Hochst. ex Benth. l. c. 282 = Schimperi.
hyptoides, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 74.—Mexic.
ianthina, *Otto & Dietr. Allg. Gartenz.* xv. (1847) 362. —Mexico.
illyrica, Schult. Obs. Bot. 8 = S. Verbenaca.
inamoena, *Vahl, Enum.* i. 269.—Hispan.
incana, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 68.—Mexic.
incarnata, Cav. Anal. Hist. Nat. ii. (1800) 112 = elegans.
incarnata, Etling. Salv. 25 = triloba, recognita.
incisa, *Benth. in DC. Prod.* xii. 352.—Afr. austr.
inconspicua, *Benth. Lab. Gen. et Sp.* 247.—Mexic.
incurvata, *Ruiz & Pav. Fl. Per.* i. 24. t. 42.—Peruv.
indica, Heyne, ex Roem. & Schult. Syst. i. 263 = Meriandra benghalensis.
indica, *Linn. Sp. Pl.* 26.—Ind. or.
indica, Linn. l. c., partim = brachycalyx.
integerrima, Mill. Gard. Dict. ed. VIII. n. 12 = africana.
integrifolia, Hardwicke, in As. Res. vi. (1799) 349 = lanata.
integrifolia, *Ruiz & Pav. Fl. Per.* i. 26. t. 36.— Peruv.
intermedia, Schult. Obs. Bot. 9 = nilotica.
interrupta, Hort. ex Benth. in DC. Prod. xii. 265 = confusa.
interrupta, *Schousb. Vextr. Marokko*, 18.—Marocc.
involucrata, *Cav. Ic.* ii. 3. t. 105.—Mexic.
Jaminiana, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 581. —Afr. bor.
japonica, *Thunb. Fl. Jap.* 22. t. 5.—Japon.
judaica, *Boiss. Diagn.* Ser. I. xii. 61.—Syria.
Karwinskii, *Benth. Lab. Gen. et Sp.* 725.—Mexic.
Keerlii, *Benth. l. c.* 263.—Mexic.
Kochiana, Kunze, Ind. Sem. Hort. Lips. (1847) 4 = S. Aethiopis.
Korolkowi, Regel & Schmalh. in Act. Hort. Petrop. vi. (1879) 356.—Turkestan.
Kotschyi, Benth. ex Boiss. Fl. Orient. iv. 600 = Russeggeri.
Kotschyi, Boiss. Diagn. Ser. I. vii. 46 = macrosiphon.
kurdica, *Boiss. & Hohen. ex Benth. in DC. Prod.* xii. 268.—Kurdistan.
lachnostachys, *Benth. Lab. Gen. et Sp.* 267.—Bras.
laciniata, Willd. Enum. Hort. Berol. Suppl. 2 = S. Verbenaca.
laciniosa, Jord. Gr. Rec. Grenoble (1849), 24 = pratensis.
laevigata, H. B. & K. Nov. Gen. & Sp. ii. t. 147 = involucrata.
laevis, *Benth. Lab. Gen. et Sp.* 251.—Mexic.
lamiifolia, *Jacq. Hort. Schoenb.* iii. 37. t. 318.—Ind. occ.
lampsanaefolia, Vahl, ex Steud. Nom. ed. I. 727 = verticillata.
lanata, *Roxb. Hort. Beng.* [80]; *Fl. Ind.* i. 146.—Reg. Himal.
lanata, Salisb. Prod. 74 = canariensis.
lanata, Stokes, Bot. Mat. Med. 52 = aethiopica.
lanceaefolia, Poir. Encyc. Suppl. v. 49 = lanceolata.
lanceolata, *Brouss. App. Elench. Pl. Hort. Monsp.* (1805) 15.—Am. bor. occ.; Mexico.
lanceolata, Lam. Illustr. i. 72 = nivea.

SALVIA:—
lanigera, *Desf. Cat. Hort. Par.* ed. III. 95, 394.—Afr. bor.
lanigera, Poir. Encyc. Suppl. v. 49 = controversa.
lantanaefolia, *Mart. & Gal. in Bull, Acad. Brux.* xi. II. (1844) 69.—Mexic.
lanuginosa, *Burm. f. Fl. Cap. Prod.* i.—Afr. austr.
lasiantha, *Benth. Lab. Gen. et Sp.* 276.—Mexic.
lasiocephala, *Hook. & Arn. Bot. Beech. Voy.* 306.—Mexico.
lasiostachys, *Benth. in DC. Prod.* xii. 350.—Afr. austr.
latens, *Benth. Pl. Hartw.* 242.—N. Granat.
latifolia, A. Anders. in Trans. Soc. Arts & Manuf. xxv. (1807) 210 = lamiifolia.
latifolia, *Vahl, Enum.* i. 271.—Afr. bor.
lavandulaefolia, Spreng. Syst. i. 58 = lavanduloides.
lavandulaefolia, *Vahl, Enum.* i. 222.—Europ. austr.
lavanduloides, *H. B. & K. Nov. Gen. et Sp.* ii. 287.—Mexic.
laxa, Benth. in DC. Prod. xii. 313 = ballotaeflora.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xx. (1885) 309.—Arizona.
Leonia, *Benth. Lab. Gen. et Sp.* 303.—Mexic.
leonuroides, *Gloxin, Obs. Bot.* 15. t. 2.—Peruv.
leptophylla, *Benth. Lab. Gen. et Sp.* 249.—Mexic.
leptostachys, *Benth. l. c.* 258.—Mexic.
leriaefolia, *Benth. in DC. Prod.* xii. 287.—Afghan.
leucantha, *Cav. Ic.* i. 16. t. 24.—Mexic.
leucanthera, Vis. in Linnaea, xiv. (1840) 137 = lamiifolia.
leucocephala, *H. B. & K. Nov. Gen. et Sp.* ii. 302.—Ecuador.
leucoclada, *Benth. in DC. Prod.* xii. 314.—Bras.
leucodermis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 276.—Madag.
leuconeura, Boiss. Diagn. Ser. II. iv. 20 = S. Aethiopis.
libanotica, *Boiss. & Gaill. Diagn.* Ser. II. iv. 16.—Syria.
limbata, *C. A. Mey. Verz. Pfl. Cauc.* 86.—As. Min.; Persia.
linarioides, *Hort. ex Henders. in Paxt. Flow. Gard.* vii. (1840) 77.—Am. austr.?
Lindenii, *Benth. in DC. Prod.* xii. 333.—Mexic.
linearifolia, Benth. l. c. 308, sphalm. = *lineatifolia*.
linearis, Ruiz & Pav. Fl. Peruv. i. t. 41 = revoluta.
lineata, *Benth. Lab. Gen. et Sp.* 724.—Mexic.
lineatifolia, Lag. Gen. et Sp. Nov. 2 = polystachya.
linifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 70.—Mexic.
Linkiana, Schult. Mant. i. 182 = pilantha.
Littae, *Vis. in Padov. Nuov. Saggi*, vi. (1847) 87.—Mexic.
longepetiolata, C. Koch, in Linnaea, xxi. (1848) 657 = Forskohlei.
longiflora, *Ruiz & Pav. Fl. Per.* i. 23. t. 10.—Peruv.
longiflora, Willd. Sp. Pl. i. 141 = tubifera.
longifolia, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 185 = azurea.
longifolia, Willd. Sp. Pl. i. 141 = stachyoides.
longispicata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 73.—Mexic.
longistyla, *Benth. Lab. Gen. et Sp.* 295.—Mexic.
lophantha, *Benth. in DC. Prod.* xii. 301.—Guatemala.
Lorentii, *Hochst. in Lorent, Wanderung.* 333.—Mesopot.
Lorentzii, *Griseb. in Goett. Abh.* xix. (1874) 239.—Reg. Argent.
Lowei, Steud. Nom. ed. II. ii. 506 = S. Verbenaca.
loxensis, *Benth. Pl. Hartw.* 145.—Peruv.
lupinoides, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1019 = pratensis.
lusitanica, *Jacq. f. Eclog.* i. 57.—Hispan.
lusitanica, Poir. Encyc. vi. 666 = sclaraeoides.
lutea, *H. B. & K. Nov. Gen. et Sp.* ii. 297. t. 148.—Ecuador.
lutea, *Linn. Sp. Pl.* 26.—Afr. austr.
lyraefolia, Salisb. Prod. 73 = lyrata.
lyrata, *Linn. Sp. Pl.* 23.—Am. bor.
macilenta, *Boiss. Diagn.* Ser. I. v. 13.—Arab.
macrantha, Schlecht. in Otto & Dietr. Allg. Gartenz. vi. (1838) 314 = patens.
macrantha, Schur, Enum. Pl. Transs. 520 = pratensis.
macrocalyx, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 133.—Bras.
macrochlamys, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 595.—Armenia.
macrophylla, *Benth. Lab. Gen. et Sp.* 725.—Peruv.
macrophylla, Tausch, in Flora, xxv. (1842) 282 = hians.

SALVIA:—
macrosiphon, *Boiss. Diagn.* Ser. I. v. 11.—Persia; Afghan.
macrosiphonia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
macrostachya, *H. B. & K. Nov. Gen. et Sp.* ii. 298.—Eucador.
madrensis, *Seem. Bot. Voy. Herald*, 327. t. 70.—Mexic.
magnifica, *Warsz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292, nomen.—Hab.?
malacophylla, *Benth. in DC. Prod.* xii. 312.—Peruv.
marginata, *E. Mey. Comm. Pl. Afr. Austr.* 236.—Afr. austr.
Mariana, *Mart. in Benth. Lab. Gen. et Sp.* 282.—Bras.
marrubioides, Vahl, Enum. i. 223 (*maruibioides*) = triloba.
Martensii, *Gal. in Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 77.—Mexic.
Mathewsii, *Benth. Lab. Gen. et Sp.* 722.—Peruv.
Matico, *Griseb. in Goett. Abh.* xix. (1874) 239.—Reg. Argent.
maurorum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 615.—Marocc.
maxima, Regel, Cat. Pl. Hort. Aksakov. 130 (Quid?).—Hab.?
melissaeflora, *Benth. in DC. Prod.* xii. 331.—Bras.
melissifolia, Desf. Cat. Hort. Par. ed. III. 94 = mexicana.
melissodora, Lag. Gen. et Sp. Nov. 2 = scorodoniaefolia.
membranacea, *Benth. Lab. Gen. et Sp.* 259.—Mexic.
menthaefolia, *Tenore, in Ann. Sc. Nat.* Sér. IV. ii. (1854) 379.—Hab.?
mentiens, *Pohl, Pl. Bras. Ic.* ii. 137. t. 193.—Bras.
Merjamie, Forsk. Fl. Aegypt. Arab. 10 = nudicaulis.
mexicana, Hemsl. in Gard. Chron. (1883) I. 341 = discolor.
mexicana, *Linn. Sp. Pl.* 25.—Mexic.
mexicana, Walt. Fl. Carol. 65 = azurea.
micrantha, Desf. Tabl. ed. I. 57, nomen = S. Verbenaca.
micrantha, Vahl, Enum. i. 235 = tenella.
microcalyx, *Scheele, in Linnaea*, xxii. (1849) 589.—Mexic.
microculis, Poir. Encyc. vi. 614 = elegans.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 294.—Mexic.
microphylla, Torr. Bot. Mex. Bound. 131 = Greggii.
microstegia, Boiss. & Bal. Diagn. Ser. II. iv. 17 = verbascifolia.
miltiorhiza, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 124 (*Enum. Pl. Chin. Bor.* 50).—China.
minor, *Gleichen, Ausl. Mikrosk. Entdeck.* 28.—Hab.?
minutiflora, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 124 (Enum. Pl. Chin. Bor. 50) = plebeia.
misella, *H. B. & K. Nov. Gen. et Sp.* ii. 290.—Mexic.
mitis, *Ruiz & Pav. Fl. Per.* i. 27.—Peruv.
Mocinoi, *Benth. Lab. Gen. et Sp.* 271.—Mexic.
modesta, *Boiss. Fl. Orient.* iv. 621.—As. Min.
mollis, Donn, Hort. Cantab. ed. III. 7 = verticillata.
mollis, H. B. & K. Nov. Gen. et Sp. ii. t. 153 = amethystina.
mollis, Jacq. f. Eclog. 56. t. 37 = virgata.
mollissima, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 71.—Mexic.
Moluccella, Benth. Lab. Gen. et Sp. 216 = acetabulosa.
monadelpha, *Stokes, Bot. Comment.* i. 133.—Hab.?
montana, *Gardn. & Field Sert. Pl. Sub.* t. 20.—Bras.
montana, Salisb. Prod. 74 = phlomoides.
Montbretii, *Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 42.—As. Min.; Syria.
monticola, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 238.—Afr. austr.
Moorcroftiana, *Wall. ex Benth. in Wall. Pl. As. Rar.* i. 67.—Reg. Himal.
moschata, *H. B. & K. Nov. Gen. et Sp.* ii. 293. t. 143.—N. Granat.
multicaulis, Vahl, Enum. i. 225 = acetabulosa.
multifida, Sibth. & Sm. Fl. Graec. Prod. i. 17 = S. Verbenaca.
mutilata, Medic. Bot. Beobacht. (1783) 115 (Quid?).—Hab.?
nana, *H. B. & K. Nov. Gen. et Sp.* ii. 289.—Peruv.
napaefolia, Auct. ex Benth. Lab. Gen. et Sp. 304 = nilotica.
napifolia, Auct. Croat. ex Nym. Consp. 571 = verticillata.
napifolia, *Jacq. Hort. Vindob.* ii. 71. t. 152.—As. Min.; Gallia; Ital.

SALVIA:—
neglecta, Tenore, Ind. Sem. Hort. Neap. (1829) 18 = S. Verbenaca.
nemorensis, Pall. Reise Prov. Russ. Reich. ii. 103 = nemorosa?
nemorosa, Baumg. ex Nym. Consp. 569 = pratensis.
nemorosa, Colm. Cat. Pl. Catalun. 122 = valentina.
nemorosa, *Crantz, Stirp. Austr.* ed. II. 242.—Europ.
nemorosa, Linn. Sp. Pl. ed. II. 35 = sylvestris.
neovidensis, *Benth. Lab. Gen. et Sp.* 284.—Bras.
nepetifolia, Desf. Tabl. ed. II. 68; Poir. Encyc. Suppl. v. 48 = hirsuta.
nepetoides, H. B. & K. Nov. Gen. et Sp. ii. 299. t. 150 = amarissima.
nervata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 77.—Mexic.
nervosa, *Benth. Lab. Gen. et Sp.* 268.—Bras.
nigricans, Hort.—Cf. Gard. Chron. (1883) I. 341; (1883) II. 588 = discolor.
nilotica, Juss. ex Jacq. Hort. Vindob. iii. 48. t. 92 = nubia.
nilotica, *Murr. in Comm. Gotting.* (1778) 98. t. 2.—Aegypt.
nipponica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 107.—Japon.
nitida, *Benth. in DC. Prod.* xii. 300.—Mexic.
nitidifolia, Orteg. Hort. Matr. Dec. 53 = mexicana.
nivea, Ehrenb. ex Boiss. Fl. Orient. iv. 619 = verbascifolia.
nivea, *Thunb. Prod. Pl. Cap.* 96.—Afr. austr.
nodosa, Ruiz & Pav. Fl. Per. i. 25 = leonuroides.
nubia, *Ait. Hort. Kew.* ed. I. i. 42.—Afr. bor. occ.
nubia, Murr. in Comm. Gotting. (1778) 90. t. 3 = nilotica.
nubicola, Wall. Pl. As. Rar. i. 68; Cat. n. 2146 = glutinosa.
nudicaulis, C. Koch, in Linnaea, xix. (1847) 24 = virgata.
nudicaulis, *Vahl, Enum.* i. 266.—Arab.
nutans, *Linn. Sp. Pl.* 27.—Europ. austr. or.
nutans, Willd. ex Ledeb. Fl. Ross. iii. 366 = pendula.
oblonga, C. Koch, in Linnaea, xix. (1847) 24 = virgata.
oblongata, Schur, Enum. Pl. Transs. 961 (Index) = pratensis.
oblongata, Vahl, Enum. i. 256 = S. Verbenaca.
oblongifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 79.—Mexic.
obovata, Ell. Sketch, i. 33 = lyrata.
obscura, Benth. Lab. Gen. et Sp. 245 = occidentalis.
obtusa, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 72.—Mexic.
obtusata, Brot. Phyt. Lusit. i. 203 = S. Verbenaca.
obtusata, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
occidentalis, *Sw. Prod. Veg. Ind. Occ.* 14.—Am. trop.
ochroleuca, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 254; xxii. (1875) 65, nomen.—Marocc.
ocimoides, Roxb. in Wall. Pl. As. Rar. i. 68; Cat. n. 2148 = lyrata?
odorata, Willd. Enum. Hort. Berol. 43 = candidissima.
officinalis, *Linn. Sp. Pl.* 23.—Reg. Mediterr.
officinalis, Pall. Tabl. Taur. 44 = grandiflora.
oligophylla, Auch. ex Benth. in DC. Prod. xii. 279 = hypoleuca.
oligostachya, *Gasparr. in Ann. Civ.* i. (1838) 117.—Europ.
oppositiflora, *Ruiz & Pav. Fl. Per.* i. 26. t. 43.—Ecuador.
orbicularis, *Benth. Bot. Voy. Sulph.* 151.—Panama.
Orbignaei, *Benth. in DC. Prod.* xii. 338.—Peruv.
oreades, Schott & Kotschy, ex Boiss. Fl. Orient. iv. 621 = frigida.
orgyalis, Fenzl, in Tchihat. Asie Min. Bot. ii. 136 = recognita.
orientalis, Hort. ex Poir. Encyc. vi. 629 = argentea.
orientalis, Mill. Gard. Dict. ed. VIII. n. 8 = rosaefolia.
Ottoniana, Lehm. Ind. Sem. Hort. Hamb. (1850) 12 = seq.?
Ottonis, *Lehm. in E. Otto, Hamb. Gartenz.* vi. (1850) 350; *Regel, in Gartenfl.* (1853) 129; *in Bot. Zeit.* xi. (1853) 335.—Mexic. *vel* Peruv.?
ovalifolia, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 267—Bras.
ovata, F. G. Dietr. Vollst. Lexik. Gaertn. vii. 465 = triloba.
Owerini, Trautv. in Act. Hort. Petrop. ii. (1873) 479 = aristata.
oxyodon, *Webb & Heldr. ex Walp. Ann.* iii. 254.—Hispan.

SALVIA:—
pachystachya, Trautv. in Bull. Soc. Nat. Mosc. xli. (1868) I. 462 = caespitosa.
palaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 303.—N. Granat.
palaestina, *Benth. Lab. Gen. et Sp.* 718.—Syria.
pallida, *Benth. l. c.* 250.—Bras.
pallidiflora, St. Amans, Fl. Agen. 10 = S. Verbenaca.
paniculata, *Linn. Mant.* i. 25.—Afr. austr.
papilionacea, Cav. Ic. iv. 9 = mexicana.
papillosa, Hoffmgg. Verz. Pfl. Nachtr. iii. 58 = officinalis.
paposana, *Phil. Fl. Atac.* 39.—Chili.
parquenensis, *Moç. & Sesse, ex Benth. Lab. Gen. et Sp.* 273.—Mexic.
Parryi, *A. Gray, in Proc. Am. Acad.* viii. (1870) 369.—Texas.
parviflora, Brot. Phyt. Lusit. i. 203 = S. Verbenaca.
parviflora, Roxb. Hort. Beng. 4 = plebeia.
parviflora, Salisb. Prod. 74 = nilotica.
parviflora, Vahl, Enum i. 268 = syriaca.
parvifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 232.—Madag.
patens, *Cav. Ic.* v. 33. t. 454.—Mexic.
patula, Desf. Fl. Atlant. i. 25 = argentea.
patula, Maund, Bot. Gard. v. 714 = praec.?
pauciflora, *H. B. & K. Nov. Gen. et Sp.* ii. 303.—Mexic.
pauciserrata, *Benth. Pl. Hartw.* 241.—N. Granat.
Pavonii, *Benth. Lab. Gen. et Sp.* 278.—Peruv.
pectinifolia, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 33 = caespitosa.
peloponnesiaca, *Boiss. & Heldr. Diagn.* Ser. I. vii. 47.—Graec.
pendula, Bess. Fl. Galic. i. 41 = nutans.
pendula, *Vahl, Enum.* i. 281.—Europ. or.
pentstemonoides, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 13.—Texas.
persepolitana, *Boiss. Diagn.* Ser. I. xii. 60.—Persia.
persicifolia, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 281.—Bras.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* ii. 304.—N. Granat.
phlomoides, *Asso, Oryctog. Arag.* 158. t. 4.—Hispan.
phlomoides, Cav. Ic. iv. 10. t. 320 = hirsuta.
phlyctidea, *C. Koch, in Linnaea*, xxi. (1848) 655.—Transcauc.
phoenicea, *H. B. & B. Nov. Gen. et Sp.* ii. 302. t. 155.—Ecuador.
Piasezkii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 528.—China.
pichinchensis, *Benth. Pl. Hartw.* 243.—Ecuador.
pilantha, *Link, Enum. Hort. Berol.* i. 10 (sp. dub.).—Oriens.
pilifera, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. (1836) 40.—As. Min.
pilosa, *Vahl, Enum.* i. 241.—Peruv.
Pinardi, *Boiss. Diagn.* Ser. I. xii. 59.—Syria.
pinifolia, Roem. & Schult. Syst. i. 220, sphalm. = *pinnifolia*.
pinnata, *Linn. Sp. Pl.* 27.—As. Min.; Syria.
pinnifolia; Pall. Tabl. Taur. 44 = scabiosaefolia.
pisidica, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 269.—As. Min.
Pitcheri, Torr. ex Benth. Lab. Gen. et Sp. 251 = azurea.
platycheila, *A. Gray, in Proc. Am. Acad.* viii. (1870) 292.—Calif.
plebeia, *R. Br. Prod.* 501.—As. or.; Austral.
plectranthoides, *Griff. Notul.* iv. 199.—Reg. Himal.
plumosa, *Ruiz & Pav. Fl. Per.* i. 26. t. 37.—Peruv.
pogonocalyx, *Hance, in Journ. Linn. Soc.* xiii. (1873) 85.—China.
polyadenia, Boiss. & Heldr. Diagn. Ser. I. v. 7 = limbata.
polyclonos, Boiss. Diagn. Ser. I. vii. 47 = compressa.
polymorpha, Hoffmgg. & Link, Fl. Port. i. 149. t. 19 = S. Verbenaca.
polystachya, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 77 = xalapensis.
polystachya, *Orteg. Hort. Matr. Dec.* 55.—Mexic.
pomifera, *Linn. Sp. Pl.* 24.—Graecia; Syria.
porphyrantha, Decne. in Rev. Hortic. Sér. IV. iii. (1854) 301. f. 16 = Roemeriana.
porphyrata, Hook. Bot. Mag. t. 4939, sphalm. = praec.
porphyrocalyx, *Baker, in Journ. Linn. Soc.* xviii. (1880) 277.—Madag.
potentillaefolia, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 270.—As. Min.; Syria.

SALVIA:—
praecox, Loisel. Not. 6 = virgata.
praecox, Savi, Fl. Pisan. i. 22 = S. Verbenaca.
praecox, Vahl, Enum. i. 274 = spinosa.
prasiifolia, *Benth. Bot. Voy. Sulph.* 151.—Mexic.
pratensis, Habl. Phys. Beschr. Taur. 143 = virgata.
pratensis, *Linn. Sp. Pl.* 25.—Europ.; Reg. Caucas.
Prionitis, *Hance, in Journ. Bot.* viii. (1870) 74.—China.
prismatica, Hemsl. Biol. Centr. Am. ii. 558 = hispanica.
privoides, *Benth. Bot. Voy. Sulph.* 150.—Am. trop.
procumbens, Ruiz & Pav. Fl. Per. i. 27 = occidentalis.
procurrens, *Benth. Lab. Gen. et Sp.* 266.—Am. austr.
propinqua, *Benth. l. c.* 267.—Bras.
prostratus, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 200.—Ins. Galapag.
protracta, *Benth. in DC. Prod.* xii. 309.—Mexic.
proxima, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 62 = vitifolia.
prunelloides, Benth. Pl. Hartw. 90 = nana.
prunelloides, *H. B. & K. Nov. Gen. et Sp.* ii. 289.—Mexic.
prysmatica, Cav. Desc. 14 = hispanica.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 526.—China.
Pseudo-aethiopis, Boiss. Fl. Orient. iv. 619 = verbascifolia.
Pseudo-coccinea, Jacq. Coll. ii. 302 = coccinea.
Pseudo-pendula, Schur, Enum. Pl. Transs. 961 (Index) = nutans.
Pseudo-silvestris, Stapf, in Denkschr. Akad. Wien, l. (1885) 42.—Persia.
pubescens, *Benth. Lab. Gen. et Sp.* 723.—Mexic.
pulchella, *DC. Cat. Hort. Monsp.* 142.—Am. austr.
pulchella, H. B. & K. Nov. Gen. et Sp. ii. 288. t. 140 = spicata.
pumila, Benth. Lab. Gen. et Sp. 726 = aegyptiaca.
punctata, *Ruiz & Pav. Fl. Per.* i. 27. t. 39.—Ecuador.
pungens, Hochst. ex Benth. in DC. Prod. xii. 350 = nilotica.
punicea, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 65 = elegans.
purpurascens, *Mart. & Gal. l. c* 69.—Mexic.
purpurea, *Cav. Ic.* ii. 52. t. 166.—Mexic.
purpurina, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 82.—Mexic.
pustulata, *Benth. in DC. Prod.* xii. 327.—Peruv.
pyramidalis, *Petagn. Inst. Bot.* ii. 54.—Calabr.
pyrenaica, *Linn. Sp. Pl.* 25.—Mont. Pyren.
pyrenaica, Vahl, Enum. i. 263 = bicolor.
pyrifolia, Domb. ex Cav. Desc. 13, in obs. = fulgens, leonuroides.
quercifolia, Hort. ex Benth. Lab. Gen. et Sp. 235 = virgata.
quitensis, *Benth. in DC. Prod.* xii. 339.—Ecuador.
racemosa, *Ruiz & Pav. Fl. Per.* i. 25.—Peruv.
radicans, Poir. Encyc. vi. 621 = occidentalis.
Radula, *Benth. in DC. Prod.* xii. 291.—Afr. austr.
raphanifolia, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 237.—Afr. austr.
Rascheyana, *Boiss. Diagn.* Ser. I. xii. 58.—Syria.
Rassami, Boiss. Fl. Orient. iv. 615 = palaestina.
recognita, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 33.—As. Min.
rectiflora, *Vis. Sem. Hort. Patav.* (1839) (*Ort. Bot. Pad.* 145).—Mexic.
recurva, *Benth. in DC. Prod.* xii. 336.—Mexic.
reflexa, Hornem. Enum. Pl. Hort. Hafn. (1807) 34 = lanceolata.
Regeliana, *Trautv. Ind. Sem. Hort. Petrop.* (1866) 93.—Reg. Caucas.
Regla, *Cav. Ic.* v. 33. t. 455.—Mexic.
remota, *Benth. in DC. Prod.* xii. 304.—Mexic.
repens, Benth. Lab. Gen. et Sp. 267 = ovalifolia.
repens, *Burch. ex Benth. l. c.* 306.—Afr. austr.
reptans, Jacq. Hort. Schoenb. iii. 38 = angustifolia.
reticulata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 64.—Mexic.
Reuteriana, *Boiss. Diagn.* Ser. I. v. 10.—Persia.
revoluta, *Ruiz & Pav. Fl. Per.* i. 28. t. 41.—Peruv.
rhinosina, *Griseb. in Goett. Abh.* xix. (1874) 239.—Reg. Argent.
rhombifolia, *Ruiz & Pav. Fl. Per.* i. 26. t. 36.—Peruv.

SALVIA:—
rhytidea, *Benth. in DC. Prod.* xii. 280.—Afghan.
rigida, *Benth. Lab. Gen. et Sp.* 269.—Bras.
rigida, Thunb. Prod. Pl. Cap. 96 = dentata.
ringens, *Sibth. & Sm. Fl. Graec.* i. 14. t. 18.—Graecia.
riparia, *H. B. & K. Nov. Gen. et Sp.* ii. 300.—Peruv.
rivularis, *Gardn. & Field. Sert. Pl.* t. 20.—Bras.
Roborowskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 527.—China.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 586.—Texas; Mexic.
Roemeriana, Torr. Bot. Mex. Bound. 132, partim = Henryi.
Roezli, *Scheidw. in Fl. des Serres*, xiv. (1861) 31.—Mexic.
rosaefolia, Hort. ex Schult. Mant. i. 183 = pilantha.
rosaefolia, *Sm. Ic. Ined.* i. 5. t. 5.—Armenia.
Rosani, Tenore, Fl. Nap. iii. 22 = viridis.
rosea, Vahl, Enum. i. 244 = coccinea.
rosmarinifolia, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 532 = lavandulaefolia.
rosmarinoides, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 269.—Bras.
Rosmarinus, Schleiden, Handb. Med. Pharm. Bot. i. 265 = Rosmarinus officinalis.
rostrata, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 8 = pratensis.
rotundifolia, Benth. in DC. Prod. xii. 273 = Eckloniana.
rotundifolia, Salisb. Prod. 74 = africana.
rotundifolia, Vis. in Mem. Ist. Venet. i. 1842 (1843), 42. t. 1 = grandiflora.
rubescens, *H. B. & K. Nov. Gen. et Sp.* ii. 301. t. 154.—Ecuador.
rubicunda, Wender. ex Benth. Lab. Gen. et Sp. 233 = pratensis.
rubifolia, *Boiss. Diagn.* Ser. I. xii. 56.—Syria.
rubiginosa, *Benth. in DC. Prod.* xii. 301.—Mexic.
rubra, Spreng. Syst. iv. Cur. Post. 17 = virgata.
rudis, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 235.—Afr. austr.
rufida, Willd. ex Schult. Mant. i. 189 = rufula.
rufula, *H. B. & K. Nov. Gen. et Sp.* iii. 291.—N. Granat.
rugosa, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
rugosissima, Zucc. in Abh. Akad. Muench. iii. (1837-43) 244. t. 7 = controversa.
rumicifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 305.—Ecuador.
runcinata, *Linn. f. Suppl.* 89.—Afr. austr.
Ruprechti, Boiss. Fl. Orient. iv. 624 = Beckeri.
Russeggeri, *Fenzl, Del. Sem. Hort. Vindob.* (1851) [4]; *et ex Tchihat. Asie Min. Bot.* ii. 138.—As. Min.
Russellii, *Benth. Lab. Gen. et Sp.* 312.—As. Min.; Mesopot.
ruthenica, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. VII. 55 = nutans.
rutilans, *Carr. in Rev. Hortic.* (1873) 251.—Hab.?
sabulicola, Pomel, Nouv. Mat. Fl. Atl. 121 = S. Verbenaca.
saccata, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 424 = argentea.
sagittata, *Ruiz & Pav. Fl. Per.* i. 23. t. 35.—Peruv.
sahendica, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 172.—Persia.
salicifolia, *Pohl, Pl. Bras. Ic.* ii. 140. t. 196.—Bras.
saligna, Willd. ex A. Dietr. Sp. Pl. i. 307 = spicata.
Salvatori, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 772 = pratensis.
Sanctae-luciae, *Seem. Bot. Voy. Herald*, 327.—Mexic.
santolinaefolia, *Boiss. Diagn.* Ser. I. v. 13.—Persia; Ind. bor. occ.
sarawschanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 62.—Turkestan.
saxicola, *Wall. ex Benth. in Wall. Pl. As. Rar.* i. 68; *Cat.* n. 2147.—Reg. Himal.
Scabiosae, Pers. Syn. i. 29 = seq.
scabiosaefolia, *Lam. Journ. Hist. Nat. Par.* ii. (1792) 44. t. 27.—Peruv.
scabra, Benth. Lab. Gen. et Sp. 305 = runcinata.
scabra, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
scabrida, Bertol. Fl. Ital. i. Sp. Add. 855 = Bertolonii.
scabrida, *Pohl, Pl. Bras. Ic.* ii. 140. t. 195.—Bras.
scapiformis, *Hance, in Journ. Bot.* xxiii. (1885) 368.—China.
Schielii, Walp. Ann. vi. 1273 (Index), sphalm. = Shielii.

SALVIA :—
Schimperi, *Benth. in DC. Prod.* xii. 282.—Abyss.
Schimperiana, Hochst. ex Benth. in DC. Prod. xii. 262 = Meriandra benghalensis.
Schmalhauseni, *Regel, in Act. Hort. Petrop.* vi. (1879) 356.—Bokhara.
Sclaraea, Crantz, Stirp. Austr. ed. II. 236 = austriaca.
sclaraeoides, *Brot. Fl. Lusit.* i. 17.—Lusitan.
Sclarea, *Linn. Sp. Pl.* 27.—Reg. Mediterr.; Oriens.
Scorodonia, Benth. Lab. Gen. et Sp. 264 = scorodoniaefolia.
scorodoniaefolia, *Desf. Tabl.* ed. II. 67; *Poir. Encyc. Suppl.* v. 46.—Mexic.
scrobiculata, *Meyen, Reise*, ii. 45.—Peruv.
scutellarioides, *H. B. & K. Nov. Gen. et Sp.* ii. 304. t. 158.—Ecuador.
secunda, *Benth. Lab. Gen. et Sp.* 285.—Bras.
Sellowiana, *Benth. in DC. Prod.* xii. 329.—Bras.
semiatrata, *Zucc. in Abh. Akad. Muench.* i. (1829–30) 298.—Mexic.
semperflorens, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 81.—Mexic.
serotina, *Linn. Mant.* i. 25.—Ind. occ.
serotina, Vahl, Enum. i. 232 = tenella.
serrata, *Benth. Lab. Gen. et Sp.* 257.—Bras.
Sessei, Benth. l. c. 288.—Mexic.
sessilifolia, *Baker, in Journ. Linn. Soc.* xviii. (1881) 276.—Madag.
Shielei, *Boiss. Diagn.* Ser. I. vii. 45.—Persia.
Sibthorpii, Bory & Chaub. Nouv. Fl. Pélop. 2 = S. Verbenaca.
Sibthorpii, Sibth. & Sm. Fl. Graec. i. 17. t. 22 = virgata.
sidaefolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 67.—Mexic.
Sideritidis, *Presl, ex Benth. Lab. Gen. et Sp.* 255.—Peruv.
Sideritidis, Vahl, Enum. i. 250 = hirsuta.
Sieberi, Presl, Bot. Bemerk. 100 = palaestina.
silvestris, Auct. = *sylvestris*.
simplex, Spreng. Syst. i. 58 = elongata.
Simsiana, Schult. Mant. i. 210 = S. Sclarea.
sinaica, Delile, ex Benth. Lab. Gen. et Sp. 718 = palaestina.
sinapifolia, C. Koch, in Linnaea, xxi. (1848) 659 = Russellii.
sipylea, Lam. Illustr. i. 68 = triloba.
smyrnaea, *Boiss. Diagn.* Ser. I. v. 5.—As. Min.
somalensis, *Vatke, in Linnaea*, xliii. (1880–82) 93.—Afr. trop.
Sonklari, *Pant. in Magyar, Növényt. Lapok*, v. (1881) 151.—Europ. austr.
sordida, *Benth. Pl. Hartw.* 241.—N. Granat.
Souchetii, Van Houtte, in Fl. des Serres, xiv. (1861) 32 = Roezli.
speciosa, *Presl, ex Benth. Lab. Gen. et Sp.* 272.—Peruv.
spectabilis, H. B. & K. Nov. Gen. et Sp. ii. 304 = patens.
speirematoides, *Wright, in Sauv. Fl. Cub.* 114.—Cuba.
sphacelioides, *Benth. in DC. Prod.* xii. 337.—N. Granat.
spicata, *Schult. Mant.* i. 202.—Mexic.
spicata, Willd. ex A. Dietr. Sp. Pl. i. 277 = collina.
Spielmanni, Scop. Del. Insub. iii. 31. t. 15 = viridis.
Spielmanni, Willd. Enum. Hort. Berol. i. 38 = S. Verbenaca.
Spielmanniana, Bieb. Fl. Taur. Cauc. i. 21 = S. Verbenaca.
spinosa, *Linn. Mant.* ii. 511.—Arab.; Mesopot.
spinosa, Moench, Meth. 373 = nilotica.
spinulosa, Montbr. & Auch. ex Benth. in DC. Prod. xii. 283 = frigida.
spiraeifolia, *Boiss. & Hohen. Diagn.* Ser. I. v. 5.—Syria.
splendens, *Ker-Gawl. in Bot. Reg.* t. 687; *Sellow, ex Nees, in Neuwied, Reise Bras.* ii. 335.—Bras.
squalens, *H. B. & K. Nov. Gen. et Sp.* ii. 297.—Peruv.
squarrosa, *Scheidw. in Hortic. Belge*, v. (1838) 187. t. 10.—Mexic.
stachydea, Klein, ex Schult. Mant. i. 216 = Meriandra benghalensis.
stachydifolia, *Benth. in L. C. Prod.* xii. 311.—Peruv.
stachyoides, *H. B. & K. Nov. Gen. et Sp.* ii. 287. t. 138.—Mexic.
staminea, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 65 = patens.

SALVIA :—
staminea, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 41.—As. Min.
stenophylla, *Burch. ex Benth. Lab. Gen. et Sp.* 306.—Afr. austr.
stolonifera, *Benth. Pl. Hartw.* 70.—Mexic.
striata, *Benth. in DC. Prod.* xii. 343.—Peruv.
strictiflora, *Hook. Bot. Misc.* ii. (1831) 234.—Peruv.
strobilanthoidea, *Wright, ex Griseb. Cat. Pl. Cub.* 214.—Cuba.
suaveolens, Pomel, Nouv. Mat. Fl. Atl. 306 = argentea.
subincisa, *Benth. Pl. Hartw.* 20.—N. Mexic.
sublobata, Schur, Enum. Pl. Transs. 520 = pratensis.
subrotunda, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 290.—Bras.
subsessilis, *E. Mey. Comm. Pl. Afr. Austr.* 237.—Afr. austr.
subspathulata, *Lehm. in E. Otto, Hamb. Gartenz.* vi. (1850) 457.—Afr. austr.
subtriflora, Roem. & Schult. Syst. i. 610 (Index) = tubiflora.
subtriloba, *Schrank, in Syll. Ratisb.* ii. (1828) 58 = triloba.
suffruticosa, *Montbr. & Auch. in Ann. Sc. Nat.* Sér. II. vi. (1836) 39.—As. Min.
superba, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 769 = coccinea.
sylvestris, Cav. Desc. 291 (*silvestris*) = valentina.
sylvestris, *Linn. Sp. Pl.* 24.—Europ. or.; As. bor.
sylvestris, Vell. Fl. Flum. 14; i. t. 34 = Arrabidae.
sylvestris, Vis. Stirp. Dalm. 46 = Bertolonii.
sylvicola, *Burch. ex Benth. Lab. Gen. et Sp.* 304.—Afr. austr.
sypilea, Lam. Illustr. i. 68 = triloba.
syriaca, Gouan, ex Benth. in DC. Prod. xii. 279 = dominica.
syriaca, *Linn. Syst.* ed. X. 854.—As. Min.; Syria; Persia.
Szovitsiana, *Bunge, Lab. Pers.* 43.—Persia.
Tafallae, *Benth. Lab. Gen. et Sp.* 260.—Peruv.
taraxacifolia, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 253; xxii. (1875) 65.—Marocc.
taurica, Habl. ex Willd. Sp. Pl. i. 129 = scabiosaefolia.
taurica, Hort. ex Benth. Lab. Gen. et Sp. 235 = virgata.
tauricola, Schott & Kotschy, ex Boiss. Fl. Orient. iv. 608 = acetabulosa.
taurina, Hort. ex Benth. Lab. Gen. et Sp. 237 = sylvestris.
Tchihatcheffii, *Boiss. Fl. Orient.* iv. 598.—Galatia.
tebesana, *Bunge, Lab. Pers.* 52.—Persia.
tenella, *Sw. Prod. Veg. Ind. Occ.* 14.—Am. trop.
Tenorii, *Spreng. Pugill.* i. 2 = pratensis.
tenuifolia, Burch. ex Benth. Lab. Gen. et Sp. 304 = triangularis.
tenuior, Desf. ex Roem. & Schult. Syst. i. 200 = lavandulaefolia.
tetragona, Moench, Meth. 373 = hispanica.
texana, *Torr. Bot. Mex. Bound.* 132.—Texas.
thymiflora, Willd. Enum. Hort. Berol. Suppl. 2 = lamiifolia.
thymoides, *Benth. Lab. Gen. et Sp.* 255.—Mexic.
thyrsiflora, *Benth. Bot. Voy. Sulph.* 151.—Mexic.
tiberina, Mauri, in Tenore, Viagg. Abruzz. 41 = pratensis.
tiliaefolia, *Vahl, Symb. Bot.* iii. 7.—Mexic.
tingitana, Etling. Salv. 35 = foetida.
tmolea, *Boiss. Diagn.* Ser. I. v. 9.—As. Min.
tolimensis, *H. B. & K. Nov. Gen. et Sp.* ii. 292.—N. Granat.
tomentella, *Pohl, Pl. Bras. Ic.* ii. 138. t. 194.—Bras.
tomentosa, Heyne, ex Wall. Cat. n. 2152 = Gomphostemma strobilinum.
tomentosa, Mill. Gard. Dict. ed. VIII. n. 2 = grandiflora.
tortuosa, *H. B. & K. Nov. Gen. et Sp.* ii. 292. t. 142.—N. Granat.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 57 = pratensis.
Trautvetteri, *Regel, in Act. Hort. Petrop.* vi. (1879) 355.—Turkestan.
triangularis, *Thunb. Prod. Pl. Cap.* 96.—Afr. austr.
trichocalycina, *Benth. in DC. Prod.* xii. 356.—Afghan.
trichoclada, *Benth. l. c.* 267.—Mesopot.
trichostemoides, Pursh, Fl. Am. Sept. i. 19 = lanceolata.
trichostyle, Bischoff, Del. Sem. Hort. Heidelb. (1847) in adnot.—Cf. Ann. Sc. Nat. Sér. III. ix. (1848) 319 = farinacea.

SALVIA :—
tricolor, Hort. ex Benth. in DC. Prod. xii. 290 = pratensis.
tricolor, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 773 = officinalis.
tricolor, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 71. t. 120.—Mexic.
tricuspidata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 78.—Mexic.
triloba, *Linn. f. Suppl.* 88.—Europ. or.; As. Min.
truncata, Willd. Enum. Hort. Berol. 34 = viridis.
tubaeformis, Auct. = *tubiformis*.
tubifera, *Cav. Ic.* i. 16. t. 25.—Mexic.
tubiflora, *Sm. Ic. Ined.* 26. t. 26.—Peruv.
tubiformis, Klotzsch, in Otto & Dietr. All. Gartenz. ix. (1841) 114 = rectiflora.
Turdi, *A. Rich. Tent. Fl. Abyss.* ii. 194.—Abyss.
uliginosa, *Benth. Lab. Gen. et Sp.* 251.—Bras.
umbratica, *Hance, in Journ. Bot.* viii. (1870) 75.—China.
undulata, *Benth. in DC. Prod.* xii. 275.—Afr. austr.
undulata, *Wender. ex Steud. Nom.* ed. II. ii. 508.—Hab.?
urmiensis, *Bunge, Lab. Pers.* 48.—Persia.
urticifolia, *Linn. Sp. Pl.* 24.—Am. bor.
uruguayensis, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 258.—Bras.
valentina, *Vahl, Enum.* i. 268.—Hispan.
vanensis, Boiss. & Noé, Diagn. Ser. II. iv. 17 = suffruticosa.
varia, Vahl, Enum. i. 273 = syriaca.
variabilis, *Loisel. ex Benth. Lab. Gen. et Sp.* 241 = S. Verbenaca.
variegata, *Host, Fl. Austr.* i. 27.—Hungar.
variegata, Waldst. & Kit. ex Willd. Enum. Hort. Berol. i. 36 = pratensis.
velutina, *Benth. Lab. Gen. et Sp.* 275.—Bras.
velutina, Vahl, Enum. i. 268 = sylvestris.
venusta, H. B. & K. Nov. Gen. et Sp. ii. 301. t. 153 = amethystina.
verbascifolia, *Bieb. Fl. Taur. Cauc.* iii. 24.—As. Min.; Persia.
Verbenaca, *Linn. Sp. Pl.* 25.—Europ.; Oriens.
verbenacoides, Brot. Fl. Lusit. i. 17 = S. Verbenaca.
verbenaefolia, Salisb. Prod. 73 = S. Verbenaca.
versicolor, *Lag. ex Steud. Nom.* ed. II. ii. 508.—Hispan.
verticillata, *Linn. Sp. Pl.* 26.—Europ. austr.; Oriens.
vestita, *Benth. in DC. Prod.* xii. 346.—N. Granat.
villosula, Benth. Lab. Gen. et Sp. 720 = Martensii.
violacea, Donn, Hort. Cantab. ed. II. 5; et ex A. Dietr. Sp. Pl. i. 276 = lamiifolia.
violacea, *Ruiz & Pav. Fl. Per.* i. 26. t. 43.—Ecuador.
virgata, *Ait. Hort. Kew.* ed. I. i. 39.—Europ. austr. or.; Oriens.
virgata, Orteg. Hort. Matr. Dec. 3 = angustifolia.
virgata, Savi, Bot. Etrusc. ii. 4 = pratensis.
viridis, *Linn. Sp. Pl.* 24.—Reg. Mediterr.; Oriens.
viscida, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 268.—Bras.
viscosa, *Jacq. Misc.* ii. 328.—Ital.
viscosa, Wight, ex Steud. Nom. ed. II. ii. 509 = Meriandra benghalensis.
vitifolia, *Benth. Lab. Gen. et Sp.* 724.—Mexic.
Viviani, Sieber, ex Reichb. Pl. Crit. vi. 18. t. 524 = S. Verbenaca.
vulnerariaefolia, Willd. Sp. Pl. i. 149 = scabiosaefolia.
Wagneriana, *Polak. in Linnaea*, xli. (1877) 591.—Costa Rica.
Warszewicziana, *Regel, in Flora*, xxxii. (1849) 184.—Guatemala.
Wiedemanni, *Boiss. Fl. Orient.* iv. 599.—Anatolia.
xalapensis, *Benth. in DC. Prod.* xii. 308.—Mexic.
xanthocheila, *Boiss. ex Benth. l. c.* 284.—As. Min.; Persia.

SALVIASTRUM, Heist. Syst. 16 (1748) = **Tarchonanthus**, Vaill. (Compos.).

SALVIASTRUM, Scheele, in Linnaea, xxii. (1849) 584. *LABIATAE*, Benth. & Hook. f. ii. 1196.
texanum, *Scheele, l. c.* 585.—Texas.

SALZMANNIA, DC. Prod. iv. 617 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 106.
nitida, *DC. l. c.*—Bras.

SALZWEDELIA, O. F. Lang, in Flora, xxvi. (1843) 770 = SALTZWEDELIA, Fl. Wett. = **Genista**, Linn. (Legumin.).

SAMADERA, Gaertn. Fruct. ii. 352. t. 159 (1791).
SIMARUBEAE, Benth. & Hook. f. i. 310.
BIPOREIA, Thou. Gen. Nov. Madag. 14 (1806).
LOCANDI, Adans. Fam. ii. 449 (1763).
MANUNGALA, Blanco, Fl. Filip. ed. I. 306 (1837).
MAUDUITA, Comm. ex DC. Prod. i. 592 (1824).
NIOTA, Lam. Illustr. t. 299 (1796 ?).
SAMANDURA, Linn. Fl. Zeyl. 202 (1747), nomen prius.
VITMANNIA, Vahl, Symb. Bot. iii. 51. t. 60 (1794).
brevipetala, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 410.—Java.
glandulifera, *Presl, Symb. Bot.* ii. 1. t. 51.—Ind. or.
indica, *Gaertn. Fruct.* ii. 352. t. 156.—Ind. or.
lucida, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 562.—Burma.
madagascariensis, *A. Juss. in Mém. Mus. Par.* xii. (1825) 516. t. 27.—Madag.
pentapetala, G. Don, Gen. Syst. i. 811 = indica.
tetrapetala, *G. Don, l. c.*—Madag.

SAMANDURA, Linn. Fl. Zeyl. 202 (1747) = praec.

SAMARA, Linn. Mant. ii. 144 (1771) (excl. syn. p. 199) = **Embelia**, Burm. (Myrsin.).
atropunctata, Arn. in Proc. Linn. Soc. i. (1847) 327 = E. viridiflora.
aurantiaca, Arn. l. c. = E. viridiflora.
australiana, F. Muell. Fragm. vi. 164.—Austral.
Blackburnia, Spreng. Syst. i. 441 = Zanthoxylum Blackburnia.
esculenta, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 147 = E. floribunda.
floribunda, Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 222 = E. Ribes, floribunda.
frondosa, King, ex Gamble, List Darjeeling Trees, 52 = E. Gamblei.
Helicia, D. Dietr. Syn. Pl. i. 497 = Helicia cochinchinensis.
Izquierda, Spreng. Syst. i. 442.—Peruv.
laeta, Linn. Mant. ii. 199 = Memecylon umbellatum ?
laeta, Sw. Prod. Veg. Ind. Occ. 151 = Myrsine laeta.
longifolia, Benth. Fl. Hongk. 205.—Ins. Hongkong.
microcalyx, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 221 = E. microcalyx.
Myrtillus, Kurz, l. c. 223 = E. Myrtillus.
Nagushia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 147 = E. Nagushia.
obovata, Benth. in Hook. Kew Journ. iv. (1852) 301.—Ins. Hongkong.
paniculata, Roxb. Hort. Beng. 83; Fl. Ind. i. 414 = Ardisia Samara.
parviflora, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 223 = E. parviflora.
philippensis, Vidal, Sinops. t. 61. f. D = E. philippinensis.
picta, Buch.-Ham. ex Wall. Cat. n. 2302 = E. robusta.
polygama, Roxb. Hort. Beng. 83; Fl. Ind. i. 414 = Ardisia polygama.
racemosa, Hassk. Cat. Hort. Bog. Alt. 232 = E. racemosa.
Rheedii, Wight, Ic. t. 1591 = E. viridiflora.
Ribes, Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 222 = E. Ribes.
robusta, Benth. & Hook. f. ex Kurz, l. c. = E. robusta.
scandens, Hassk. Cat. Hort. Bog. Alt. 232 = E. javanica.
sessiliflora, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 222 = E. sessiliflora.
undulata, Arn. in Proc. Linn. Soc. i. (1847) 327 = E. Nagushia.
vestita, Benth. & Hook. f. ex Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 223 = E. vestita, nutans.
viridiflora, Arn. in Proc. Linn. Soc. i. (1847) 327 = E. viridiflora.
viridiflora, Thw. Enum. Pl. Zeyl. 173 = E. viridiflora.

SAMARA, Sw. Prod. Veg. Ind. Occ. 32 (1788); Fl. Ind. Occ. i. 261 (1797) = **Myrsine**, Linn.
coriacea, Sw. Prod. Veg. Ind. Occ. 32 = M. coriacea.
floribunda, Willd. Sp. Pl. i. 665 = M. floribunda.
guianensis, Sw. Obs. Bot. 51 = M. floribunda.
laeta, Sw. Prod. Veg. Ind. Occ. 151 = M. laeta.
myricoides, Willd. ex Schult. Mant. iii. 220 = M. popayanensis.

SAMARA:—
myrtifolia, Willd. ex Schult. Mant. iii. 220 = Myrsine myrtifolia.
pentandra, Ait. Hort. Kew. ed. I. i. 160 = M. pentandra.
pentandra, A. DC. in DC. Prod. viii. 98 = M. floribunda.
Potama, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 146 = M. africana.
saligna, Willd. ex Schult. Mant. iii. 220 = M. saligna.
sessilis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 147 = M. semiserrata.
subspinosa, Buch.-Ham. l. c. = M. semiserrata.

SAMAROPYXIS, Miq. Fl. Ind. Bat. Suppl. 464 (1860) = **Hymenocardia**, Wall. (Euphorb.).
elliptica, Miq. l. c. 465 = H. punctata.

SAMARPSES, Rafin. New Fl. Am. iii. 93 (1836) = **Fraxinus**, Tourn. (Oleac.).
triptera, Rafin. l. c. = F. platycarpa.

SAMBUCUS, Tourn. ex Linn. Syst. ed. I (1735).
CAPRIFOLIACEAE, Benth. & Hook. f. ii. 3.
EBULUM, Garcke, Fl. Deutschl. ed. VII. 184 (1865).
PHYTEUMA, Lour. Fl. Cochinch. 138 (1790).
TRIPETELUS, Lindl. in Mitch. Three Exped. ii. 14 (1839).
adnata, *Wall. Cat.* n. 482; *et ex DC. Prod.* iv. 322.—Reg. Himal.
angustifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = javanica.
alba, Rafin. Alsog. Am. 48 (Quid ?).—Am. bor.
arborescens, Gilib. Exercit. i. 5 = nigra.
arborescens, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 13 = racemosa.
australis, *Cham. & Schlecht. in Linnaea*, iii. (1828) 140.—Bras.
bipinnata, *Cham. & Schlecht. l. c.* v. (1830) 171.—Mexic.
bipinnata, Moench, Meth. 506 = canadensis.
caerulea, *Rafin. Alsog. Am.* 48 (cerulea).—Am. bor. occ.
californica, Hort. ex C. Koch, Dendrol. ii. I. 72 = caerulea ?
canadensis, Burm. f. Fl. Ind. 75 = Turpinia pomifera.
canadensis, *Linn. Sp. Pl.* 269.—Am. bor.
canadensis, Thunb. Fl. Jap. 126 = javanica.
chinensis, Lindl. in Trans. Hort. Soc. vi. (1826) 297 = javanica.
cochinchinensis, Spreng. Syst. i. 935 = Loureiriana.
columnaris, Hort. ex Lebas, in Rev. Hortic. (1884) 398 = pyramidata.
dimidiata, Rafin. Alsog. Am. 49 (Quid ?).—Am. bor.
dissecta, Hort. ex C. Koch, Dendrol. ii. I. 71 = nigra.
ebuloides, *Desv. ex DC. Prod.* iv. 323.—China.
Ebulus, *Linn. Sp. Pl.* 269.—Europ.; Oriens; Reg. Himal.
elegans, Hort. ex C. Koch, Dendrol. ii. I. 71 = nigra.
floribunda, Hort. l. c. = nigra.
florida, Salisb. Prod. 173 = nigra.
Fontenaysii, Carr. in Rev. Hortic. (1868) 99 = caerulea ?
Gaudichaudiana, *DC. Prod.* iv. 322.—Austral.
glauca, Benth. Pl. Hartw. 313 = mexicana.
glauca, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 13 = caerulea.
graveolens, Willd. ex Schult. Syst. vi. 641 = peruviana.
herbacea, Gilib. Fl. Lituan. i. 250 = S. Ebulus.
hirta, *Tausch, in Flora*, xxi. (1838) 737.—Hab. ?
humilis, Mill. Gard. Dict. ed. VIII. n. 5 = S. Ebulus.
humilis, Rafin. Ann. Nat. 13 = canadensis.
integerrima, Stokes, Bot. Mat. Med. ii. 173 = Loureiriana.
japonica, *Thunb. Fl. Jap.* 125.—Japon.
javanica, *Reinw. ex Blume, Bijdr.* 657.—As. trop.
laciniata, Mill. Gard. Dict. ed. VIII. n. 2 = nigra.
lanceolata, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 31 = nigra.
leucocarpa, Hort. ex C. Koch, Dendrol. ii. I. 71 = nigra.
linearis, Hort. l. c. = nigra.
Loureiriana, *DC. Prod.* iv. 323.—China.
lucida, *Tausch, in Flora*, xxi. (1838) 737.—Hab. ?

SAMBUCUS:—
maderensis, *Lowe, Man. Fl. Mad.* i. 381.—Ins. Mader.
medullosa, Gilib. Fl. Lituan. i. 2 = nigra.
melanocarpa, *A. Gray, in Proc. Am. Acad.* xix. (1883) 76.—Am. bor. occ.
mexicana, *Presl, ex DC. Prod.* iv. 322.—Am. bor. occ.
monstrosa, Hort. ex Loud. Hort. Brit. Suppl. ii. 671 = nigra.
nigra, *Linn. Sp. Pl.* 269.—Europ.; As. bor.
nigra, Lour. Fl. Cochinch. 181 = Loureiriana.
nigra, Marsh. Arbust. 141 = canadensis.
nigra, Thunb. Fl. Jap. 126 = racemosa.
palmensis, *Link, in Buch, Beschr. Canar. Ins.* 151.—Ins. Canar.
paucijuga, *Stev. in Bull. Soc. Nat. Mosc.* xxi. (1848) II. 277.—Transcauc.
peruviana, *H. B. & K. Nov. Gen. et Sp.* iii. 429.—Peruv.
phyteumoides, *DC. Prod.* iv. 323.—Cochinch.
pubens, Michx. Fl. Bor. Am. i. 181 = racemosa.
pubescens, Pers. Syn. i. 328 = racemosa.
pulverulenta, Hort. ex C. Koch, Dendrol. ii. I. 71 = nigra.
pyramidata, *Hort. ex Lebas, in Rev. Hortic.* (1884) 398.—Hab. ?
racemosa, *Linn. Sp. Pl.* 270.—Reg. bor. temp.
repens, Rafin. Alsog. Am. 47 (Quid ?).—Am. bor.
rotundifolia, Lodd. ex Sweet, Hort. Brit. ed. II. 257 = nigra.
rubra, Buch.-Ham. ex Wall. Cat. n. 482, p. 235 = javanica.
rupestris, Rafin. Alsog. Am. 50 (Quid ?).—Am. bor.
Sieboldiana, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 265 = racemosa.
suaveolens, DC. Prod. iv. 323, sphalm. = peruviana.
Thunbergiana, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 265 = javanica.
Thunbergii, G. Don, Gen. Syst. iii. 438 = javanica.
tiliaefolia, Wall. Cat. n. 483; et ex DC. Prod. iv. 257 = Toricellia tiliaefolia.
velutina, Durand & Hilg. in Pacif. Rail. Rep. v. 8; et in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 39 = mexicana.
verrucosa, Rafin. Alsog. Am. 49 = racemosa.
virescens, Desf. Hist. Arbr. i. 348 = nigra.
vulgaris, Neck. Delic. Gallo-Belg. i. 156 = nigra.
Wightiana, Wall. Cat. n. 6303 = javanica.
Williamsii, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 217 = racemosa.
xanthocarpa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 42.—Austral.

SAMBULUS, Mohl & Schlecht. in Bot. Zeit. v. (1847) 424, err. typ. = SUMBULUS, H. Reinsch = **Ferula**, Tourn. (Umbell.).

SAMERARIA, Desv. Journ. Bot. iii. (1813) 161 = **Isatis**, Linn. (Crucif.).
armena, Desv. l. c. = I. armena.
cardiocarpa, Trautv. in Act. Hort. Petrop. i. (1871) 31.—Reg. Casp.
elegans, Boiss. Fl. Orient. i. 375.—Persia.
glastifolia, Boiss. l. c.—Armenia.
leiocarpa, Boiss. l. c.—Persia.
stylophora, Boiss. l. c.—Persia.

SAMODIA, Baudo, in Ann. Sc. Nat. Sér. II. xx. (1843) 350 = **Samolus**, Tourn. (Primul.).
ebracteata, Baudo, l. c. = Samolus ebracteatus.

SAMOLUS, [Tourn.] Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 638.
SAMODIA, Baudo, in Ann. Sc. Nat. Sér. II. xx. (1843) 350.
SHEFFIELDIA, Forst. Char. Gen. t. 9 (1776).
africanus, Burm. f. Fl. Cap. Prod. 5 = porosus ?
ambiguus, R. Br. Prod. 429 = repens ?
americanus, Spreng. Syst. i. 702 = S. Valerandi.
aquaticus, Lam. Fl. Fr. iii. 329 = S. Valerandi.
Beccabungae-facie, Gilib. Fl. Lituan. i. 60 = S. Valerandi.
bracteatus, Stokes, Bot. Mat. Med. i. 344 = S. Valerandi.
campanuloides, R. Br. Prod. 429 = porosus.
caulescens, Willd. ex Roem. & Schult. Syst. v. 4, in obs. = S. Valerandi.
ebracteatus, *H. B. & K. Nov. Gen. et Sp.* ii. 223. t. 129.—Am. bor.; Ind. occ.

SAMOLUS:—
floribundus, H. B. & K. Nov. Gen. et Sp. ii. 181 = S. Valerandi.
geniculatus, Dulac, Fl. Hautes-Pyr. 422 = S. Valerandi.
junceus, *R. Br. Prod.* 428.—Austral.
latifolius, *Duby, in DC. Prod.* viii. 74.—Chili.
littoralis, R. Br. Prod. 428 = repens.
littoralis, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 34.—Hab. ?
longipes, Hook. ex Shuttl. in Bot. Zeit. iii. (1845) 222 = ebracteatus.
parviflorus, Nees, in Lehm. Pl. Preiss. i. 337 = repens.
parviflorus, Rafin. in Am. Monthly Mag. (1818) 176 = S. Valerandi ?
platyphyllus, *F. Muell. Census*, 91.—Austral.
porosus, *Thunb. Fl. Cap.* ii. 32.—Afr. austr.
repens, *Pers. Syn.* i. 171.—Austral.; N. Zel.
spathulatus, *Duby, in DC. Prod.* viii. 74.—Am. austr.
subnudicaulis, *A. St. Hil. ex Compt. Rend. Acad. Sc. Par.* (1838) II. 98.—Bras.
superbus, *Brickell, in Med. Repos. N. York*, vi. (1803) 327 (Quid ?).—Am. bor.
Valerandi, *Linn. Sp. Pl.* 443.—Reg. temp.

SAMUDRA, Rafin. Fl. Tellur. iv. 72 (1836) = **Argyreia**, Lour. (Convolv.).
speciosa, Rafin. l. c. = A. speciosa.

SAMYDA, Linn. Coroll. Gen. 21 (1737). *SAMYDACEAE*, Benth. & Hook. f. i. 797.
GEUNZIA, Neck. Elem. ii. 440 (1790).
SADYMIA, Griseb. Fl. Brit. W. Ind. 25 (1859).
affinis, *Spreng. Syst.* ii. 354.—Ins. S. Doming.
arborea, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Casearia stipularis.
bazanica, Buch.-Ham. ex Wall. Cat. n. 7192 C ? = Casearia tomentosa.
Canziala, Buch.-Ham. in Trans. Linn. Soc. xvii. (1836) 225 = Casearia tomentosa.
comocladifolia, Poir. Encyc. Suppl. v. 32 = Casearia comocladifolia.
coriacea, Poir. l. c. = Casearia coriacea.
crenata, Poir. Encyc. vi. 491 = Casearia nitida.
decandra, Jacq. Enum. Pl. Carib. 21 = Casearia parvifolia.
decurrens, *Spreng. Syst.* ii. 354.—Bras.
denticulata, Poir. in Dict. Sc. Nat. xlvii. 159 = serrulata.
dodecandra, Jacq. Enum. Pl. Carib. 21 = serrulata.
fragilis, Poir. Encyc. Suppl. v. 32 = Casearia fragilis.
glabra, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 227; et in Wall. Cat. n. 7195 A = Casearia graveolens.
glabrata, *Sw. Prod. Veg. Ind. Occ.* 68.—Ind. occ.
grandiflora, *Griseb. Cat. Pl. Cub.* 11.—Cuba.
grandiflora, *Spreng. Syst.* ii. 354.—Bras.
greviaefolia, Poir. Encyc. Suppl. v. 32 = Casearia greviaefolia.
Guidonia, Linn. Sp. Pl. 443 = Casearia brevipes.
hirsuta, Poir. Encyc. vi. 493 = Casearia hirsuta.
icosandra, Sw. Fl. Ind. Occ. iii. 1962 = Casearia icosandra.
ilicifolia, Poir. Encyc. Suppl. v. 31 = Casearia ilicifolia.
Iroucana, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Casearia ramiflora.
lucida, *Hort. ex Steud. Nom.* ed. II. ii. 509.—Ind. occ.
macrocarpa, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 48.—Mexic.
macrophylla, Willd. Sp. Pl. ii. 625 = Casearia viridiflora.
multiflora, *Cav. Ic.* i. 48. t. 67.—Ind. occ.
nitida, Linn. Syst. ed. X. 1025 = Casearia nitida.
Niviana, Poir. Encyc. vi. 493 = Casearia serrulata.
parviflora, Linn. Syst. ed. X. 1025 = Casearia parvifolia.
parviflora, Poir. Encyc. vi. 491 = Casearia sylvestris.
parvifolia, Poir. l. c. 492 = Casearia parvifolia.
petiolaris, Spruce, ex Eichl. in Mart. Fl. Bras. xiii. I. 454 = Laetia suaveolens.
Piscidia, Buch.-Ham. ex Wall. Cat. n. 7193 D = Casearia tomentosa.
Pitumba, Poir. Encyc. vi. 492 = Casearia macrophylla.
polyandra, Willd. Sp. Pl. ii. 626 = Casearia Melistaurum.
procera, Poepp. & Endl. Nov. Gen. et Sp. iii. 67 = Laetia procera, casearioides.

SAMYDA:—
pubescens, Blanco, Fl. Filip. ed. II. 263 = Casearia tomentosa.
pubescens, *Linn. Sp. Pl.* ed. II. 758.—Ind. occ.
pubescens, Linn. l. c., partim = serrulata.
rosea, Sims, Bot. Mag. t. 550 = serrulata.
rubra, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 48.—Mexic.
Selloi, Spreng. Syst. ii. 354 = Casearia inaequilatera.
serrulata, Blanco, Fl. Filip. ed. I. 374 = Casearia tomentosa.
serrulata, *Linn. Sp. Pl.* ed. II. 558.—Ind. occ.
silvestris, Poir. Encyc. vi. 492 = *sylvestris*.
spinescens, Sw. Prod. Veg. Ind. Occ. 68 = Casearia brevipes.
spinosa, Linn. Sp. Pl. ed. II. 557 = Casearia ramiflora.
spinulosa, Vent. Choix, 43. t. 43 = glabrata.
stipularis, Poir. Encyc. Suppl. v. 32 = Casearia stipularis.
suaveolens, Poepp. & Endl. Nov. Gen. et Sp. iii. 66 = Laetia suaveolens.
sylvestris, Poir. Encyc. vi. 492 (*silvestris*) = Casearia sylvestris.
tinifolia, Poir. Encyc. Suppl. v. 32 = Casearia tinifolia ?
tomentosa, Sw. Prod. Veg. Ind. Occ. 68 = Casearia hirta.
velutina, *DC. Prod.* ii. 48.—Ins. S. Doming.
villosa, *Sw. Prod. Veg. Ind. Occ.* 68.—Jamaic.
viridiflora, Poir. Encyc. vi. 493 = Casearia viridiflora.

SANAMUNDA, Adans. Fam. ii. 285 (1763) = **Passerina**, Linn. (Thymel.).
Myrsinites, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 147 = Thymelaea Sanamunda.
tomentosa, Zumag. Fl. Pedem. ii. 156 = P. hirsuta.

SANAMUNDA, Neck. Elem. ii. 179 (1790) = **Daphne**, Linn. (Thymel.).

SANCHEZIA, Ruiz & Pav. Prod. 5. t. 32 (1794). *ACANTHACEAE*, Benth. & Hook. f. ii. 1083.
ANCYLOGYNE, Nees, in Mart. Fl. Bras. ix. 63. t. 7 (1847).
glabra, Pers. Syn. i. 24 = oblonga.
glaucophylla, Hort.—Cf. Gard. Chron. (1869) 587 = nobilis.
hirsuta, Pers. Syn. i. 24 = ovata.
longiflora, *Hook. f. ex Planch. in Fl. des Serres*, xxiii. (1880–83) 257. t. 2460.—Ecuador.
munita, *Planch. l. c.*—Am. trop.
nobilis, *Hook. f. Bot. Mag.* t. 5594.—Ecuador.
oblonga, *Ruiz & Pav. Fl. Per.* i. 7. t. 8. f. 6.—Peruv.
ovata, *Ruiz & Pav. l. c.*—Peruv.

SANDERSONIA, Hook. Bot. Mag. t. 4716 (1853). *LILIACEAE*, Benth. & Hook. f. iii. 830.
aurantiaca, *Hook. Bot. Mag.* t. 4716.—Natal.
littonioides, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 262 = Littonia Welwitschii.

SANDORICUM, Cav. Diss. vii. 359. tt. 202, 203 (1789). *MELIACEAE*, Benth. & Hook. f. i. 333.
Beccarianum, *Baill. Adansonia*, xi. (1873–76) 264.—Borneo.
borneense, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 33.—Malaya.
dasyneuron, *Baill. Adansonia*, xi. (1873–76) 265.—Borneo.
emarginatum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 553.—Molucca.
glaberrimum, Hassk. Retzia, i. 145 = indicum.
indicum, *Cav. Diss.* vii. 359. tt. 202, 203.—As. trop.
Maingayi, Hiern, in Hook. f. Fl. Brit. Ind. i. 554 = borneense.
nervosum, Blume, Bijdr. 163 = indicum.
serratum, *G. Don, Gen. Syst.* i. 680.—Ind. or.
ternatum, Blanco, Fl. Filip. ed. I. 346 = indicum.
venosum, M. Roem. Syn. Hesper. 109 = indicum.

SANFORDIA, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 53 = **Geleznovia**, Turcz. (Rutac.).
calycina, J. Drumm. ex Harv. l. c. 54 = G. calycina.

SANGUINARIA, Dill. ex Linn. Syst. ed. I (1735). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
acaulis, Moench, Meth. 227 = canadensis.
canadensis, *Linn. Sp. Pl.* 505.—Am. bor.

SANGUINARIA:—
grandiflora, Rosc. Fl. Illustr. Seasons, t. 8; ex Sweet, Brit. Flow. Gard. Ser. II. v. t. 147 = canadensis.
stenopetala, Steud. Nom. ed. II. ii. 510 = canadensis.
vernalis, Salisb. Prod. 376 = canadensis.
virginiana, Gaertn. Fruct. ii. 479 = canadensis.

SANGUINELLA, Gleichen, Mikrosk. Untersuch. 4 [bis] t. 8 (1764) = **Panicum**, Linn.
Thunbergii, Gleichen, ex Beauv. Agrost. 177 = Hemarthria compressa.
tripsacoides, Gleichen, ex Beauv. l. c. = Hemarthria compressa.

SANGUISORBA, Rupp. ex Linn. Syst. ed. I (1735) = **Poterium**, Linn. (Rosac.).
alpina, Bunge, in Ledeb. Fl. Alt. i. 142.—Sibir. altaic.
altissima, Moench, Meth. 519 = P. officinale.
Andersonii, G. Don, Gen. Syst. ii. 594.—Sibir.
angustifolia, Radde, Ber. Reis. Sued. Ost-Sibir. 466 = P. tenuifolium.
annua, Nutt. ex Torr. & Gray, Fl. N. Am. i. 429 = P. annuum.
armena, Boiss. Diagn. Ser. II. v. 95.—Kurdistan.
auriculata, Scop. Fl. Carn. ed. II. i. 110 = P. officinale.
bracteosa, Bess. ex Eichw. Skizze, 163 = P. officinale.
canadensis, Linn. Sp. Pl. 117 = P. canadense.
carnea, Fisch. ex Link, Enum. Hort. Berol. i. 144 = P. officinale.
cernua, Bess. in Flora, xvii. (1834) I. Beibl. 11 = P. officinale.
cordifolia, Opiz, Boehm. Gew. (1823) 22 = P. officinale.
decandra, Benth. & Hook. f. Gen. i. 624, sphalm. = seq.
diandra, Wall. Cat. n. 710 = P. diandrum.
dictyocarpa, Franch. Fl. Cher-et-Loire, 180 = P. Sanguisorba.
dodecandra, Moretti, in Bibl. Ital. lxx. (1833) 436; et in Flora, xvi. (1833) 655 = P. dodecandrum (in Addend.).
garganica, Bertol. Fl. Ital. ii. 191 = P. garganicum.
hispanica, Mill. Gard. Dict. ed. VIII. n. 3 = P. officinale.
ingrica, Sweet, Hort. Brit. ed. II. 193 (Quid ?).—Rossia.
longifolia, Bertol. in Mem. Acad. Sc. Bologn. xii. (1861) 234 = P. longifolium.
macrostachya, Comolli & Jan, in Cat. Pl. Hort. Parm. (1833); et in Fl. Comen. i. 200 = P. dodecandrum (in Addend.).
major, Gilib. Fl. Lituan. i. 18 = P. officinale.
mauritanica, Desf. Fl. Atlant. i. 142 = P. verrucosum.
media, Hook. Fl. Bor. Am. i. 197 = P. sitchense.
media, Linn. Sp. Pl. ed. II. 169 = P. canadense.
microcephala, Presl, Epim. Bot. 202.—Am. bor. occ.
minor, Bertol. Fl. Ital. ii. 189 = P. polygamum.
minor, Scop. Fl. Carn. ed. II. i. 110 = P. Sanguisorba.
montana, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 212 = P. officinale.
muricata, Franch. Fl. Cher-et-Loire, 181 = P. polygamum.
myriophylla, Reichb. ex A. Br. & Bouché, Ind. Sem. Hort. Berol. (1867) App. i. 10.—Hab. ?
neglecta, G. Don, Gen. Syst. ii. 594 = P. officinale.
obtusa, Maxim. in Bull. Acad. Pétersb. xix. (1874) 160.—Japon.
occidentalis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 429 = P. annuum.
officinalis, Linn. Sp. Pl. 116 = P. officinale.
palustris, Rafin. Atl. Journ. 153 (Quid ?).—Am. bor.
Pimpinella, Spenn. Fl. Friburg. iii. 1085, in textu = P. Sanguisorba.
polygama, Nyland. Spicil. Fl. Fenn. Cent. i. 10 = P. polygamum ?
Poterium, [Weber, in] Wigg. Prim. Fl. Holsat. 14 = P. Sanguisorba.
praecox, Bess. ex Reichb. Fl. Germ. Excurs. 611 = P. officinale.
rubra, Schrank, Pl. Rar. Hort. Monac. t. 69 = P. officinale.
sabauda, Mill. Gard. Dict. ed. VIII. n. 2 = P. officinale.
serotina, Jord. Pugill. Pl. Nov. (1852) 71 = P. officinale.
sitchensis, C. A. Mey. in Middend. Reise (Fl. Ochot. 35) = P. sitchense.
spinosa, Bertol. Fl. Ital. ii. 191 = P. spinosum.

SANGUISORBA:—
tenuifolia, Fisch. ex Link, Enum. Hort. Berol. i. 144 = Poterium tenuifolium.
Vallistellinae, Bertol. Fl. Ital. ii. 772 = P. dodecandrum (in Addend.).

SANHILARIA, Leandr. ex DC. Prod. vii. 26 (1838) = **Stifftia**, Mikan (Compos.).

SANICULA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 70 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 880.
TRICLINIUM, Rafin. Fl. Ludov. 79 (1817).
arctopoides, *Hook. & Arn. Bot. Beech. Voy.* 141.—Am. bor. occ.
azorica, *Guthnick, ex Seub. Fl. Azor.* 41.—Ins. Azor.
bipinnata, *Hook. & Arn. Bot. Beech. Voy.* 347.—Calif.
bipinnatifida, *Dougl. in Hook. Fl. Bor. Am.* i. 258.—Am. bor. occ.
brasiliensis, D. Dietr. Syn. Pl. ii. 935 = Klotzschia brasiliensis.
canadensis, *Linn. Sp. Pl.* 235.—Am. bor.
canadensis, Thunb. Fl. Jap. 116 = elata.
canadensis, Thunb. Prod. Pl. Cap. 49 = europaea.
capensis, Eckl. & Zeyh. Enum. 339 = europaea.
chilensis, Bert. ex Colla, in Mem. Acc. Torin. xxxvii. (1834) 84 = graveolens.
chinensis, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 106 = europaea.
ciliaris, Sm. ex H. C. Wats. in Godman, Nat. Hist. Azor. 165 = azorica.
ciliata, Soland. ex H. C. Wats. l. c. = azorica.
crassicaulis, Poepp. ex DC. Prod. iv. 84 = S. Liberta.
crithmifolia, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 419.—Tauria.
elata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 183 = europaea.
europaea, *Linn. Sp. Pl.* 235.—Europ.; As. bor.; Afr. austr.
graveolens, *Poepp. ex DC. Prod.* iv. 85.—Chili.
hermaphrodita, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 183 = europaea.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 46 = Cryptotaenia canadensis.
javanica, Blume, Bijdr. 882 = europaea.
laciniata, *Hook. & Arn. Bot. Beech. Voy.* 347.—Am. bor. occ.
lamelligera, *Hance, in Journ. Bot.* xvi. (1878) 11.—China.
Liberta, *Cham. & Schlecht. in Linnaea*, i. (1826) 353 (err. typ. 253).—Chili.
macrorhiza, Colla, in Mem. Acc. Torin. xxxvii. (1834) 84 = graveolens.
marilandica, *Linn. Sp. Pl.* 235.—Am. bor.
marilandica, Torr. & Gray, Fl. N. Am. i. 602, partim = canadensis.
maritima, *Kellogg, ex S. Wats. Bibliog. Ind. N. Am. Bot.* 431, nomen; *et in Bot. Calif.* ii. 451.—Calif.
marylandica, DC. Prod. iv. 84 = marilandica.
Menziesii, *Hook. & Arn. Bot. Beech. Voy.* 142.—Am. bor. occ.
mexicana, *DC. Prod.* iv. 84.—Mexic.
montana, Reinw. ex Blume, Bijdr. 882 = europaea.
nevadensis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 139.—Calif.
nudicaulis, Hook. & Arn. Bot. Beech. Voy. 347 = Menziesii.
officinalis, Gouan, Hort. Monsp. 131; [Clairv.] Man. Herb. 78 = europaea.
officinarum, Neck. Delic. Gallo-Belg. i. 137 = europaea.
orthacantha, *S. Moore, in Journ. Bot.* xiii. (1875) 227.—China.
pinnatifida, Torr. in Bot. U. St. Expl. Exped. ii. 314 = bipinnata.
rubriflora, *F. Schmidt, in Maxim. Prim. Fl. Amur.* 123.—China.
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 705. t. 88.—Ins. Sandvic.
sylvatica, Salisb. Prod. 159 = europaea.
triciinaria, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.
Triclinium, DC. Prod. iv. 85 = canadensis.
trilobata, Gilib. Fl. Lituan. ii. 6 = europaea.
tuberculata, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 431.—Korea.
tuberosa, *Torr. in Pacif. Rail. Rep.* iv. 91.—Am. bor. occ.
vulgaris, Fries, Summ. Veg. Scand. 24 = europaea.

SANILUM, Rafin. Fl. Tellur. iv. 70 (1836), nomen prius = **Hewittia**, Wight & Arn. (1837) (Convolv.).
bicolor, Rafin. l. c. = H. bicolor.
bracteatum, Rafin. l. c. = H. bicolor.
copticum, Rafin. l. c. = Ipomoea dissecta.
humile, Rafin. l. c. = Ipomoea humilis, *Rafin.*
parviflorum, Rafin. l. c. = Ipomoea parviflora.

SANSEVERA, Stokes, Bot. Mat. Med. ii. 260 (1812) = seq.

SANSEVERINA, Thunb. Fl. Cap. ii. 322 (1817) = **Sansevieria**, Thunb. (Haemod.).

SANSEVIELLA, Reichb. Consp. 44 (1828), nomen prius = **Reineckea**, Kunth (Liliac.).
carnea, Reichb. ex Steud. Nom. ed. II. ii. 510 = R. carnea.

SANSEVIERA, Willd. Sp. Pl. ii. 159 (1799) = seq.

SANSEVIERIA, Thunb. Prod. Pl. Cap. 65 (1794). *HAEMODORACEAE*, Benth. & Hook. f. iii. 679.
ACYNTHA, Medic. Theodora, 76 (1786).
SALMIA, Cav. Ic. iii. 24. t. 246 (1794).
aethiopica, Thunb. Prod. Pl. Cap. 65 = zeylanica.
angolensis, Welw. Apont. 543, nomen; et ex Carr. in Rev. Hortic. (1861) 447 = cylindrica.
angustiflora, Lindb. in Act. Soc. Sc. Fenn. x. (1871) 130 = thyrsiflora.
angustifolia, Baker, in Journ. Linn. Soc. xiv. (1875) 547 = thyrsiflora.
bracteata, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 253.—Afr. trop.
Aubrytiana, *Carr. in Rev. Hortic.* (1861) 449.—Afr. trop.
canaliculata, *Carr. l. c.*—Afr. trop.
cannaefolia, Spreng. Syst. ii. 93 = Cordyline terminalis.
carnea, Andr. Bot. Rep. t. 361 = Reineckia carnea.
cylindrica, *Boj. Hort. Maurit.* 349; *Hook. Bot. Mag.* t. 5093.—Afr. trop.
Ehrenbergii, *Schweinf. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 549.—Abyss.
ensifolia, Haw. Syn. Pl. Suec. 66 = zeylanica.
flexuosa, Blume, Enum. Pl. Jav. i. 11 = Dracaena angustifolia.
fragrans, Jacq. Fragm. 29 = Dracaena fragrans.
fruticosa, Blume, Enum. Pl. Jav. i. 11 = Dracaena angustifolia.
fulvocincta, Haw. Succ. Pl. Suppl. 30 = thyrsiflora.
glauca, Haw. Syn. Pl. Succ. 65 = guineensis.
grandicuspis, Haw. l. c. 67 = zeylanica.
guineensis, *Willd. Sp. Pl.* ii. 159.—Afr. trop.
hyacinthoides, Hort. ex Steud. Nom. ed. II. ii. 511 = guineensis.
javanica, Blume, Enum. Pl. Jav. i. 11 = Dracaena elliptica.
laetevirens, Haw. Syn. Pl. Succ. 66 = guineensis.
lanuginosa, *Willd. Sp. Pl.* ii. 160.—Ind. or.
latifolia, Boj. Hort. Maurit. 348 = guineensis.
longiflora, *Sims, Bot. Mag.* t. 2634.—Afr. trop.
nilotica, *Baker, in Journ. Linn. Soc.* xiv. (1875) 548.—Afr. trop.
polyphylla, Haw. Syn. Pl. Succ. 65 = guineensis.
pumila, Haw. l. c. 67 = zeylanica.
rosea, F. G. Dietr. Vollst. Lexik. Gaertn. viii. 504 = Reineckia carnea.
Roxburghiana, *Schult. f. Syst.* vii. 357.—Ind. or.
rufocincta, Hort. Angl. ex Baker, in Journ. Linn. Soc. xiv. (1875) 548 = thyrsiflora.
sarmentosa, Jacq. Fragm. 83. t. 132. f. 3 = Reineckia carnea.
senegambica, *Baker, in Journ. Linn. Soc.* xiv. (1875) 548.—Afr. trop.
sessiliflora, Ker-Gawl. in Bot. Mag. t. 739 = Reineckia carnea.
spicata, Haw. Syn. Pl. Succ. 66 = thyrsiflora.
stenophylla, Link, Enum. Hort. Berol. i. 342 = zeylanica.
striata, Don, ex Steud. Nom. ed. II. ii. 511 (Quid?).—Hab.?
sulcata, Boj. ex Baker, in Journ. Linn. Soc. xiv. (1875) 549 = cylindrica.
thyrsiflora, *Thunb. Prod. Pl. Cap.* 65.—Afr. austr.
venosa, Don, ex Steud. Nom. ed. II. ii. 511 (Quid?).—Hab.?
zeylanica, Roxb. Pl. Corom. ii. 43. t. 184 = Roxburghiana.
zeylanica, *Willd. Sp. Pl.* ii. 159.—Zeylan.

SANSOVINIA, Scop. Introd. 228 (1777) = **Leea**, Linn. (Ampel.).

SANTALUM, Linn. Gen. ed. II. 165 (1742). *SANTALACEAE*, Benth. & Hook. f. iii. 224.
SIRIUM, Schreb. Gen. 82 (1789).
acuminatum, A. DC. in DC. Prod. xiv. 684 = Fusanus acuminatus.
album, *Linn. Sp. Pl.* 349.—Ind. or.
angustifolium, A. DC. in DC. Prod. xiv. 685 = Fusanus acuminatus.
austro-caledonicum, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 61.—N. Caled.
capense, Spreng. ex A. DC. in DC. Prod. xiv. 635, 686 = Rhoiacarpus capensis.
cognatum, Miq. in Lehm. Pl. Preiss. i. 616 = Fusanus acuminatus.
crassifolium, A. DC. in DC. Prod. xiv. 685 = Fusanus crassifolius.
Cunninghamii, Hook. f. Fl. N. Zel. i. 223 (Fusani sp.).—N. Zel.
Cygnorum, Miq. in Lehm. Pl. Preiss. i. 615 = Fusanus spicatus.
diversifolium, A. DC. in DC. Prod. xiv. 684 = Fusanus persicarius.
ellipticum, Gaudich. in Freyc. Voy. Bot. 442 = Freycinetianum.
ellipticum, Zipp. ex Span. in Linnaea, xv. (1841) 335 = album.
Freycinetianum, *Gaudich. in Freyc. Voy. Bot.* 442. t. 45.—Ins. Sandvic.
Hornei, *Seem. Fl. Vit.* 210.—Ins. Fiji.
insulare, Bert. ex A. DC. in DC. Prod. xiv. 685 = Freycinetianum.
lanceolatum, *R. Br. Prod.* 356.—Austral.
lanceolatum, Schlecht. in Linnaea, xx. (1847) 579 = Fusanus acuminatus.
Mida, Hook. Ic. Pl. t. 563, 575 = *Cunninghamii*.
myrtifolium, Roxb. Hort. Beng. 83; Fl. Ind. i. 444 = album.
oblongatum, R. Br. Prod. 355 = lanceolatum.
obtusifolium, *R. Br. l. c.* 356.—Austral.
ovatum, *R. Br. l. c.* 355.—Austral.
paniculatum, Hook. & Arn. Bot. Beech. Voy. 94 = Freycinetianum.
persicarium, F. Muell. in Trans. Vict. Inst. i. (1855) 41 = Fusanus persicarius.
Preissianum, Miq. in Lehm. Pl. Preiss. i. 615 = Fusanus acuminatus.
Preissii, F. Muell. Fragm. ii. 179 = praec.?
pyrularium, *A. Gray, in Proc. Am. Acad.* iv. (1859) 327.—Ins. Sandvic.
spicatum, A. DC. in DC. Prod. xiv. 685 = Fusanus spicatus.
venosum, R. Br. Prod. 355 = lanceolatum.
Yasi, *Seem. in Bonplandia*, ix. (1861) 258; *Fl. Vit.* 210.—Ins. Fiji.

SANTIA, Savi, in Mem. Soc. Ital. Mod. viii. (1798) II. 479 = **Polypogon**, Desf. (1800) (Gramin.).
elongata, Parl. ex Webb & Berth. Phyt. Canar. iii. 393 = P. elongatus.
maritima, Fior. in Nuov. Giorn. Pis. Lett. xvii. Sc. (1828) 111 = P. maritimus.
monspeliensis, Parl. Fl. Palerm. i. 73 = P. monspeliensis.
plumosa, Savi, in Mem. Soc. Ital. Mod. viii. (1798) II. 479 = P. monspeliensis.
unioloides, Rafin. Chlor. Aetn. 7 = Demazeria sicula.

SANTIA, Wight & Arn. Prod. 422 (1834) = **Lasianthus**, Jack (Rubiac.).
venulosa, Wight & Arn. l. c. = L. venulosus.

SANTIERA, Span. in Linnaea, xv. (1841) 328, sphalm. = **Sautiera**, Decne. (Acanth.).

SANTIRIA, Blume, Mus. Bot. Lugd. Bat. i. 209. f. 40 (1850). *BURSERACEAE*, Benth. & Hook. f. i. 325.
apiculata, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 537.—Malacca.
Beccarii, *Engl. in DC. Monog. Phan.* iv. 162.—Borneo.
bornensis, *Engl. Bot. Jahrb.* i. (1881) 43.—Borneo.
caesia, *Engl. in DC. Monog. Phan.* iv. 166.—Borneo.

SANTIRIA:—
conferta, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 537.—Malacca.
costata, *A. W. Benn. l. c.*—Malacca.
fasciculata, *A. W. Benn. l. c.* 539.—Malacca.
glabrifolia, *Engl. in DC. Monog. Phan.* iv. 164.—Borneo.
Griffithii, *Engl. Bot. Jahrb.* i. (1881) 43.—Malacca.
laevigata, *Blume, Mus. Bot. Lugd. Bat.* i. 211.—Sumatra.
Maingayi, A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 538 = oblongifolia.
mollis, *Engl. in DC. Monog. Phan.* iv. 157.—Borneo.
montana, *Blume, Mus. Bot. Lugd. Bat.* i. 212.—Borneo.
multiflora, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 538.—Malacca.
oblongifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 211.—Malaya.
parviflora, *Engl. in DC. Monog. Phan.* iv. 155.—Borneo.
pilosa, *Engl. l. c.* 159.—Borneo.
Planchoni, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 536.—Malacca.
puberula, *A. W. Benn. l. c.* 537.—Malacca.
rostrata, *Blume, Mus. Bot. Lugd. Bat.* i. 213.—Borneo.
rubiginosa, *Blume, l. c.*—Sumatra.
rufescens, *Blume, l. c.*—Sumatra.
rugosa, *Blume, l. c.* 212.—Sumatra.
serrulata, *Engl. in DC. Monog. Phan.* iv. 160.—Borneo.
Teysmanni, *Engl. l. c.* 162.—Ins. Ceram.
tomentosa, *Blume, Mus. Bot. Lugd. Bat.* i. 211.—Sumatra.
virgata, *Blume, l. c.* 213.—Borneo.

SANTOLINA, Tourn. ex Linn. Gen. ed. I. 248 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 420.
africana, Jord. & Fourr. Ic. Fl. Eur. ii. 8 = S. Chamaecyparissus.
Ageratum, Baill. Hist. Pl. viii. 311 = Achillea Ageratum.
alpina, Bertol. Amoen. Ital. 43 = Lasiospermum pedunculare.
alpina, Linn. Sp. Pl. ed. II. 1180 = Anthemis montana.
alpina, Sieber, ex Steud. Nom. ed. II. ii. 511 = Achillea cretica.
Amellus, Linn. Syst. ed. X. 1207 = Salmea Eupatoria.
annua, Linn. Sp. Pl. 842 = Lonas inodora.
anthemoides, Linn. l. c. ed. II. 1180 = Achillea cretica.
atrata, Baill. Hist. Pl. viii. 311 = Achillea atrata.
Benthamiana, Jord. & Fourr. Ic. Fl. Eur. ii. 10 = S. Chamaecyparissus.
brevicaulis, Jord. l. c. = S. Chamaecyparissus.
brevidentata, Stokes, Bot. Mat. Med. iv. 172 = S. Chamaecyparissus.
brevifolia, Jord. & Fourr. Ic. Fl. Eur. ii. 11 = S. Chamaecyparissus.
canescens, Lag. Gen. et Sp. Nov. 25 = rosmarinifolia.
capitata, *Linn. Pl. Rar. Afr.* 18 = Athanasia capitata.
Chamaecyparissus, *Linn. Sp. Pl.* 842.—Europ. austr.
chamaemelifolia, *Mill. Gard. Dict.* ed. VIII. n. 7.—Hab.?
coronopifolia, Crantz, Inst. i. 287 = Athanasia coronopifolia.
corsica, Jord. & Fourr. Ic. Fl. Eur. ii. 9 = S. Chamaecyparissus.
crenata, *Linn. Sp. Pl.* 842.—Afr. trop.
crithmifolia, *Hort. Berol. ex Hornem. Hort. Hafn. Add.* 971.—Hab.?
crithmifolia, Linn. Sp. Pl. 843 = Athanasia crithmifolia.
cuneifolia, Baill. Hist. Pl. viii. 311 = Achillea Herbarota.
cupressiformis, Lam. Fl. Fr. ii. 42 = viridis.
decumbens, *Mill. Gard. Dict.* ed. VIII. n. 3.—Hab.?
dentata, *Linn. Sp. Pl.* 843.—Afr. trop.
dentata, Moench, Meth. 580 = S. Chamaecyparissus.
diversifolia, Jord. & Fourr. Ic. Fl. Eur. ii. 9 = S. Chamaecyparissus.
elegans, *Boiss. ex DC. Prod.* vii. 296.—Hispan.
erecta, Pers. Syn. ii. 407 = Anthemis montana.
erecta, Poir. Encyc. vi. 508 = Lasiospermum pedunculare.

SANTOLINA:—
ericoides, Hort. ex Poir. Encyc. vi. 504 = S. Chamaecyparissus.
eriosperma, Desf. Tabl. ed. I. 99; Pers. Syn. ii. 407 = Lasiospermum pedunculare.
falcata, Baill. Hist. Pl. viii. 311 = Achillea falcata.
flava, Forsk. Fl. Aegypt. Arab. p. xxxi = Anthemis tinctoria.
fragrantissima, Forsk. l. c. 147 = Achillea fragrantissima.
glabrescens, Jord. & Fourr. Ic. Fl. Eur. ii. 11 = S. Chamaecyparissus.
grandiflora, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 80 = pectinata.
heterophylla, Willk. & Cutand. in Linnaea, xxx. (1859) 106 = oblongifolia.
homophylla, Jord. & Fourr. Ic. Fl. Eur. ii. 8 = S. Chamaecyparissus.
impressa, Hoffmgg. & Link, Fl. Port. ii. 363 = rosmarinifolia.
incana, Lam. Fl. Fr. ii. 43 = S. Chamaecyparissus.
intricata, Jord. & Fourr. Ic. Fl. Eur. ii. 11 = S. Chamaecyparissus.
jamaicensis, Linn. Syst. ed. X. 1207 = Calea jamaicensis.
laevigata, Linn. Pl. Rar. Afr. 18 = Athanasia aspera.
leucantha, Bertol. Amoen. Ital. 43 = pinnata.
linearifolia, Jord. & Fourr. Ic. Fl. Eur. ii. 12 = S. Chamaecyparissus.
lobata, Jord. l. c. 12 = S. Chamaecyparissus.
macrophylla, Baill. Hist. Pl. viii. 311 = Achillea macrophylla.
magna, Baill. l. c. = Achillea magna.
maritima, Crantz, Inst. i. 287 = Diotis candidissima.
maschalantha, Spreng. in Schrad. Journ. (1800) II. 199 = Lasiospermum pedunculare.
microcephala, Jord. & Fourr. Ic. Fl. Eur. ii. 11 = S. Chamaecyparissus.
Millefolium, Baill. Hist. Pl, viii. 61, 311 = Achillea Millefolium.
minor, Mill. Gard. ed. VIII. n. 6 = rosmarinifolia.
montana, Sibth. & Sm. Fl. Graec. ix. 39. t. 852 = Anthemis Sibthorpii.
moschata, Baill. Hist. Pl. viii. 311 = Achillea moschata.
muricata, Guss. ex DC. Prod. vi. 15 = Achillea muricata.
neapolitana, Jord. & Fourr. Ic. Fl. Eur. ii. 10 = S. Chamaecyparissus.
nobilis, Baill. Hist. Pl. viii. 311 = Achillea nobilis.
oblongifolia, *Boiss. Diagn.* Ser. II. iii. 18.—Hispan.
oppositifolia, Linn. Syst ed. X. 1207 = Isocarpha oppositifolia.
pallida, Salisb. Prod. 189 = S. Chamaecyparissus.
pectinata, *Benth. Cat. Pl. Pyr.* 117.—Mont. Pyren.
pectinata, Lag. Gen. et Sp. Nov. 25 = rosmarinifolia.
pinnata, Donn, Hort. Cantab. ed. II. 105 = Lasiospermum pedunculare.
pinnata, *Viv. Elench. Hort. Dinegro*, 31.—Ital.
provincialis, Jord. & Fourr. Ic. Fl. Eur. ii. 9 = S. Chamaecyparissus.
Ptarmica, Baill. Hist. Pl. viii. 311 = Achillea Ptarmica.
ptarmicoides, Lam. Illustr. Gen. t. 671. f. 5 = Achillea fragrantissima.
pubescens, *Linn. Cent. Pl.* ii. 31.—Afr. trop.
pygmaea, Sieber, ex Steud. Nom. ed. II. ii. 511 (Quid?).—Hab.?
rigida, Sibth. & Sm. Fl. Graec. Prod. ii. 166; Fl. Graec. t. 853 = Anthemis cretica.
rigidula, Jord. & Fourr. Ic. Fl. Eur. ii. 10 = S. Chamaecyparissus.
rosmarinifolia, *Linn. Sp. Pl.* 842.—Europ. austr.
scariosa, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocc.
semidentata, *Hoffmgg. & Link, Fl. Port.* ii. 362.—Hispan.
sericea, Jord. & Fourr. Ic. Fl. Eur. ii. 13 = S. Chamaecyparissus.
simplicifolia, L'Hérit. ex DC. Prod. vi. 37 = Achillea fragrantissima.
sinaica, Fresen. in Mus. Senckenb. i. (1833) 83 = Tanacetum sinaicum.
sinaitica, Boiss. Fl. Orient. iii. 353, sphalm. = praec.
squarrosa, Linn. Pl. Rar. Afr. 18; Cent. Pl. ii. 30 = Relhania squarrosa.
squarrosa, Willd. Sp. Pl. iii. 1798 = S. Chamaecyparissus.
suaveolens, *Pursh, Fl. Am. Sept.* ii. 520.—Am. bor.

SANTOLINA:—
terrestris, Forsk. Fl. Aegypt. Arab. 147 = Anacyclus alexandrinus.
tinctoria, Molina, Sagg. Chile, 142 = Cephalophora glauca.
tomentosa, Lam. Fl. Fr. ii. 41 = Diotis candidissima.
tomentosa, Pers. Syn. ii. 406 = S. Chamaecyparissus.
trifurca, Hill, Veg. Syst. iii. 126, sphalm. = seq.
trifurcata, *Linn. Sp. Pl.* 843.—Afr. trop.
tuberculosa, Lam. Fl. Fr. ii. 42 = rosmarinifolia.
valida, Jord. & Fourr. Ic. Fl. Eur. ii. 9 = S. Chamaecyparissus.
variifolia, Salisb. Prod. 189 = rosmarinifolia.
villosa, Mill. Gard. Dict. ed. VIII. n. 2 = S. Chamaecyparissus.
villosissima, Poir. Encyc. vi. 505 = S. Chamaecyparissus.
virens, Mill. Gard. Dict. ed VIII. n. 4 = viridis.
viridis, Hort. ex Poir. Encyc. vi. 504 = seq.?
viridis, *Willd. Sp. Pl.* iii. 1798.—Europ. austr.
viscosa, *Lag. Gen. et Sp. Nov.* 25.—Hispan.

SANVITALIA, Gualt. in Lam. Journ. Nat. Hist. ii. (1792) 176. t. 33. *COMPOSITAE*, Benth. & Hook. f. ii. 357.
ANAITIS, DC. Prod. v. 628 (1836).
LAURENTIA, Steud. Nom. ed. I. 466 (1821).
LORENTEA, Orteg. Hort. Matr. Dec. 41. t. 5 (1797).
Aberti, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 87.—N. Mexic.
acapulcensis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 154.—Mexic.
acinifolia, DC. Prod. v. 628 = procumbens.
angustifolia, *Engelm. ex A. Gray, Pl. Wright.* i. 12.—Mexic.
fruticosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 155.—Mexic.
helianthoides, *Rich. ex Willd. Sp. Pl.* iii. 2190.—Peruv.
mexicana, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1238 = Zinnia Haageana.
ocymoides, *DC. Prod.* v. 628.—Mexic.
procumbens, *Lam. Journ. Hist. Nat. Par.* ii. (1792) 176. t. 33.—Mexic.
tragiaefolia, DC. Prod. v. 628 = ocymoides.
versicolor, *Griseb. in Goett. Abh.* xxiv. (1879) 189.—Reg. Argent.
villosa, Cav. Ic. iv. 31. t. 35 = procumbens.

SANVITALIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 357 (1873) = **Zinnia**, Linn. (Compos.).

SAOUARI, Aubl. Pl. Gui. i. 599. t. 240 (1775) = **Caryocar**, Linn. (Ternstr.).
glabra, Aubl. l. c. = C. glabrum.
villosa, Aubl. l. c. 601 = C. villosum.

SAPINDUS, Tourn. ex Linn. Gen. ed. I. 359 (1737). *SAPINDACEAE*, Benth. & Hook. f. i. 404.
APHANIA, Blume, Bijdr. 236 (1825).
DIDYMOCOCCUS, Blume, Rumphia, iii. 103, in obs. (1847).
ELECTRA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2.
PSEUDIMA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 300, 309; ix. (1879) 495, 515.
TRIGONACHRAS, Radlk. l. c. (1878) 299, 304; ix. (1879) 500.
abruptus, Lour. Fl. Cochinch. 238 = S. Mukorossi.
abstergens, Roxb. ex Wight & Arn. Prod. 111 = trifoliatus, *Linn.*
abyssinicus, *Fresen. in Mus. Senckenb.* ii. (1837) 277.—Abyss.
acuminatus, Rafin. New Fl. Am. iii. 22 = marginatus.
acuminatus, Wall. Cat. n. 8035 = S. Mukorossi.
acutus, Roxb. ex Wight & Arn. Prod. 111 = trifoliatus, *Linn.*
acutus, Wall. Cat. n. 8096 (Engelhardtiae sp.).—Ind. or.
adenophyllus, Wall. l. c. n. 8044 = Cupania adenophylla.
alatus, Salisb. Prod. 280 = S. Saponaria.
alternifolius, Buch.-Ham. ex Wight & Arn. Prod. 112 = Erioglossum edule.
angulatus, *Poir. Encyc.* vi. 665.—Hab.?
angustifolius, *Blume, Rumphia*, iii. 99.—Hab.?

SAPINDUS:—
angustifolius, *Wall. ex Voigt, Hort. Suburb. Calc.* 94 (sp. diversa).—Mont. Khasia.
arborescens, *Aubl. Pl. Gui.* i. 357. t. 139.—Guian.
attenuatus, *Wall. Cat.* n. 8037.—Reg. Himal.
australis, *Benth. Fl. Austral.* i. 464.—Austral.
baccatus, *Blanco, Fl. Filip.* ed. I. 290.—Ins. Philipp.
balicus, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 396.—Ins. Bali.
benghalensis, Roxb. ex Wight & Arn. Prod. 113 = Nephelium Longana.
bifoliolatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 684.—Ind. or.
bijugus, Wall. Cat. n. 8045 = Hemigyrosa canescens.
capensis, Hochst. in Flora, xxvi. (1843) 80 = oblongifolius.
capensis, *Sond. in Harv. & Sond. Fl. Cap.* i. 240.—Afr. austr.
cerasinus, *Benth. in Hook. Kew Journ.* iii. (1851) 197.—Bras.
chinensis, Murr. Syst. XIII. 315 = Koelreuteria paniculata.
cinereus, A. Cunn. ex A. Gray, Bot. U. St. Expl. Exped. i. 258 = Cupania subcinerea.
cinereus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 402 = Nephelium Longana.
cultratus, *Turcz. l. c.* 403.—Ins. Philipp.
cuspidatus, *Blume, Rumphia*, iii. 98.—N. Guin.
Danura, *Voigt, Hort. Suburb. Calc.* 94.—Ind. or.; Burma.
deficiens, Wight & Arn. Prod. 111 = Hemigyrosa deficiens.
detergens, Roxb. Hort. Beng. 29; Fl. Ind. ii. 280 = S. Mukorossi.
detergens, Wall. Cat. n. 8042 = Dittelasma Rarak.
divaricatus, Hoffmgg. ex Schlecht. & Cham. in Linnaea, vi. (1831) 419 = marginatus.
divaricatus, *Willd. ex A. St. Hil. Fl. Bras. Mer.* i. 390.—Bras.
Drummondii, Hook. & Arn. Bot. Beech. Voy. 281 (excl. var.) = marginatus.
edulis, Ait. Hort. Kew. ed. I. ii. 36 = Nephelium Lit-chi.
edulis, Blume, Cat. Gew. Buitenz. 64 = Erioglossum edule.
emarginatus, Vahl, Symb. Bot. iii. 54 = trifoliatus, *Linn.*
erectus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 683.—Zeylan.
esculentus, *A. St. Hil. Pl. Us. Brés.* t. 68.—Bras.
falcatus, Rafin. Med. Bot. ii. 261 = marginatus.
Forsythii, DC. Prod. i. 607 = marginatus.
fraxinifolius, DC. l. c. 608 = Erioglossum edule.
fraxinifolius, Herb. Par. ex Blume, Rumphia, iii. 149 = Scorodendron pallens.
frutescens, *Aubl. Pl. Gui.* i. 355. t. 138.—Guian.
fruticosus, *Roxb. Hort. Beng.* 29; *Fl. Ind.* ii. 283.—Ins Molucc.
fuscatus, Buch.-Ham. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 685 = Dittelasma Rarak.
glabratus, Wall. Cat. n. 8095 = Xerospermum Noronhianum.
glabrescens, Hook. & Arn. Bot. Beech. Voy. 281 = Guarea fulva.
Glenieii, Thw. Enum. Pl. Zeyl. 408 = Glenniea unijuga.
guineensis, *G. Don, Gen. Syst.* i. 666.—Afr. trop.
inaequalis, DC. Prod. i. 608 = marginatus.
indicus, *Poir. Encyc.* vi. 667.—Ind. or.
juglandifolius, *A. St. Hil. Fl. Bras. Mer.* i. 391.—Bras.
Koelreuteria, Blanco, Fl. Filip. ed. I. 289 (= *Hemigyrosa Perrottetii*).—Ins. Philipp.
laurifolius, Balb. ex DC. Prod. i. 608 = marginatus.
laurifolius, Vahl, Symb. Bot. iii. 54 = trifoliatus, *Linn.*
lepidotus, Wall. Cat. n. 8036 = Aglaia Wallichii.
longifolius, Buch.-Ham. ex Wight & Arn. Prod. 112 = Erioglossum edule.
longifolius, *Vahl, Symb. Bot.* iii. 53.—Ins. Molucc.
lucidus, *Desv. in Ham. Prod. Pl. Ind. Occ.* 36.—Ins. S. Doming.
maduriensis, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 144.—Java.
marginatus, *Willd. Enum. Hort. Berol.* 432.—Am. bor.
mauritianus, *Hort. Par. ex Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 361.—Ins. Maurit.

SAPINDUS:—
microcarpus, D. Dietr. Syn. Pl. ii. 1318 = arborescens.
microcarpus, Jardin, Hist. Nat. Iles Marquises, 25 = S. Saponaria.
microcarpus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 205.—Siam.
microcarpus, *Ruiz & Pav. Fl. Per.* iv. t. 341.—Peruv.
microcarpus, Wight & Arn. Prod. 112 = Meliosma Arnottiana.
mollis, Blume, Rumphia, iii. 98 = trifoliatus, *Linn.*?
monogynus, Heyne, ex Wall. Cat. n. 8049 = Nephelium Longana.
montanus, *Blume, Rumphia*, iii. 97.—Java.
Mukorossi, *Gaertn. Fruct.* i. 342. t. 70.—As. trop.
multijugus, Wall. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 685 = Chiocheton paniculatus?
oahuensis, *Hillebr. ex Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 401.—Ins. Sandvic.
oblongifolius, *Sond. in Harv. & Sond. Fl. Cap.* i. 240.—Afr. austr.
oblongus, *Benth. in Hook. Kew Journ.* iii. (1851) 198.—Bras.
obovatus, Wight & Arn. Prod. 111 = Blighia sapida.
Pappea, *Sond. in Harv. & Sond. Fl. Cap.* i. 241.—Afr. austr.
peruvianus, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 312.—Peruv.
pinnatus, Mill. Gard. Dict. ed. VIII. n. 3 = Dittelasma Rarak.
pinnatus, Roxb. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 672 = Erioglossum edule.
polyphyllus, Roxb. Hort. Beng. 29, nomen = S. Mukorossi?
pubescens, *Zoll. & Mor. Syst. Verz. Zoll.* 22.—Java.
Rarak, DC. Prod. i. 608 = Dittelasma Rarak.
rigidus, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. austr.
ruber, *Kurz, For. Fl. Brit. Burma*, i. 298.—Burma.
rubiginosus, Roxb. Pl. Corom. i. 44. t. 62 = Erioglossum edule.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 404.—Bras.
salicifolius, DC. Prod. i. 608 = Atalaya salicifolia.
Saponaria, Lam. Illustr. ii. 441. t. 307 = marginatus.
Saponaria, *Linn. Sp. Pl.* 367 (*excl. syn. Pluk.*).—Am. bor. et austr.
Saponaria, Lour. Fl. Cochinch. 238 = Dittelasma Rarak.
senegalensis, *Poir. Encyc.* vi. 666.—Afr. trop.
serratus, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 284 = Jagera Roxburghii.
simplicifolius, G. Don, Gen. Syst. i. 666 = Sorindeia juglandifolia.
sinensis, J. F. Gmel. Syst. 642 = *chinensis*.
spinosus, Linn. Sp. Pl. ed. II. 526 = Zanthoxylum emarginatum.
squamosus, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 282 = Cupania glabrata.
stellulatus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 403 = Euphoria stellulata.
stenopterus, *DC. Prod.* i. 607.—Ins. S. Doming.
surinamensis, *Poir. Encyc.* vi. 665.—Guian.
tetraphyllus, Vahl, Symb. Bot. iii. 54 = Hemigyrosa canescens.
Thwaitesii, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 683.—Zeylan.
tomentosus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 204.—Burma.
travancorensis, Wall. Cat. n. 8047 = Canarium commune.
trifoliatus, *Linn. Sp. Pl.* 367.—As. trop.
trifoliatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 586.—N. Granat.
Turczaninowii, *Vidal, Phan. Cuming. Philipp.* 8, 105.—Ins. Philipp.
undulatus, Wall. ex Voigt, Hort. Suburb. Calc. 94 = Nephelium Longana?
unijugus, Thw. Enum. Pl. Zeyl. 56 = Glenniea unijuga.
vitiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 251.—Ins. Fiji.
xanthocarpus, Klotzsch, in Peters, Reise Mossamb. Bot. 119 = Deinbollia xanthocarpa.

SAPIOPSIS, Muell. Arg. in DC. Prod. xv. ii. 1205, in syn. (1866) = **Sapium**, P. Br. (Euphorb.).
cremostachys, Muell. Arg. l. c. = Sapium Aucuparium.

SAPIUM, P. Br. Hist. Jamaic. 338 (1756). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 334.
CARUMBIUM, Kurz, For. Fl. Brit. Burma, ii. 411 (1877).
CONOSAPIUM, Muell. Arg. in Linnaea, xxxii. (1863) 87.
FALCONERIA, Royle, Illustr. Bot. Himal. 354. t. 84 a (1839).
GYMNOBOTHRYS, Wall. ex Baill. Etud. Gén. Euph. 526 (1858).
SAPIOPSIS, Muell. Arg. in DC. Prod. xv. ii. 1205, in syn. (1866).
STILLINGFLEETIA, Boj. Hort. Maurit. 284 (1837).
TRIADICA, Lour. Fl. Cochinch. 610 (1790).
acutifolium, Benth. Pl. Hartw. 90 = Stillingia acutifolia.
adenodon, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 160 = Bonania adenodon.
aereum, *Klotzsch, ex Muell. Arg. in Linnaea*, xxxii. (1863) 119.—Peruv.
annuum, Torr. Bot. Mex. Bound. 201 = Stillingia annua.
aquifolium, J. Sm. in Proc. Linn. Soc. i. (1839) 41 = Alchornea ilicifolia.
Aucuparium, *Jacq. Enum. Pl. Carib.* 21; *Select. Am.* 249.—Am. trop.
Aucuparium, Willd. Sp. Pl. iv. 572 = S. Laurocerasum.
aureum, Buek, Index ad DC. Prod. iv. 364, sphalm. = aereum.
baccatum, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 694.—As. trop.
Balansae, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 53.—Reg. Argent.
berberifolium, Hort. ex Meissn. in DC. Prod. xv. ii. 907 = Alchornea ilicifolia.
biglandulosum, Muell. Arg. in Linnaea, xxxii. (1863) 116 - S. Aucuparium.
bingerium, Roxb. ex Willd. Sp. Pl. iv. 572 = indicum.
bingyricum, Roxb. ex Muell. Arg. in DC. Prod. xv. ii. 1216 = praec.
Brownei, Banks, ex Griseb. Fl. Brit. W. Ind. 49 = S. Laurocerasum.
cordifolium, Roxb. Hort. Beng. 104; Fl. Ind. iii. 693 = Alchornea mollis.
cuneatum, *Griseb. Fl. Brit. W. Ind.* 49.—Jamaic.
Dacdece, Buch.-Ham. ex Wall. Cat. n. 7965 = baccatum.
daphnoides, Griseb. in Goett. Nachr. (1865) 176 = S. Aucuparium.
discolor, *Muell. Arg. in Linnaea*, xxxii. (1863) 121.—China; Malaya.
Drummondii, Jacques, in Lem. Hort. Univ. (1844) 289 (= *Eriudaphus Clossianus*).—Hab.?
erythrospermum, Muell. Arg. in Linnaea, xxxii. (1863) 119 = Bonania erythrosperma.
eugeniaefolium, *Buch.-Ham. ex Wall. Cat.* n. 7971.—Reg. Himal.
haematospermum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 217.—Bras.
herbertiaefolium, *Jacques, in Lem. Hort. Univ.* (1844) 289.—Austral.
hexandrum, Wall. Cat. n. 7965 = baccatum.
Hippomane, G. F. W. Mey. Prim. Fl. Esseq. 275 = S. Aucuparium.
Hurmais, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 229 = indicum.
ilicifolium, Willd. Sp. Pl. iv. 573 = Hippomane spinosa.
indicum, *Willd. Sp. Pl.* iv. 572.—Ind. or.
insigne, *Trimen, Cat. Pl. Zeyl.* 83.—Ind. or.
jamaicense, Sw. Adnot. Bot. 62 = S. Aucuparium.
laevifolium, Thou. ex Baill. Etud. Gén. Euph. 513 = Stillingia obtusifolia.
laevigatum, Lam. Encyc. ii. 735 = Stillingia obtusifolia.
laurifolium, Griseb. Fl. Brit. W. Ind. 49 = S. Laurocerasum.
Laurocerasum, *Desf. Cat. Hort. Par.* ed. III. 411.—Ind. occ.
leucogynum, *Wright, ex Griseb. in Goett. Nachr.* (1865) 176.—Cuba.
leucospermum, *Griseb. l. c.* 177.—Cuba.
lineatum, Lam. Encyc. ii. 734 = Stillingia obtusifolia.
longifolium, Boj. Hort. Maurit. 284, nomen (Quid?).—Hab.?
macrocarpum, *Muell. Arg. in Linnaea*, xxxii. (1863) 119.—Peruv.
marginatum, *Muell. Arg. l. c.* 120.—Bras.

SAPIUM:—
montevidense, Klotzsch, ex Baill. Adansonia, v. (1864–65) 320 = S. Aucuparium.
Moritzianum, Klotzsch, in Seem. Bot. Voy. Herald, 100 = S. Aucuparium.
myricifolium, Wright, ex Griseb. in Goett. Nachr. (1865) 178 = Bonania myricifolia.
obovatum, *Klotzsch, ex Muell. Arg. in Linnaea*, xxxii. (1863) 120.—Bras.
obtusifolium, H. B. & K. Nov. Gen. et Sp. ii. 65 = Excoecaria obtusifolia.
obtusifolium, Lam. Encyc. ii. 735 = Stillingia obtusifolia.
obtusilobum, *Muell. Arg. in Linnaea*, xxxii. (1863) 116.—Venezuel.
oppositifolium, Klotzsch, ex Baill. Etud. Gén. Euph. 513 = Stillingia oppositifolia.
peruvianum, Steud. Nom. ed. II. ii. 512 = Excoecaria obtusifolia.
populifolium, Wall. Cat. n. 7966 = baccatum.
prunifolium, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 45 = S. Aucuparium.
salicifolium, H. B. & K. Nov. Gen. et Sp. ii. 65 = S. Aucuparium.
salpingadenium, *Muell. Arg. in Linnaea*, xxxii. (1863) 121.—Bolivia.
sebiferum, *Roxb. Hort. Beng.* 69; *Fl. Ind.* iii. 693.—Reg. trop.
Sellowianum, Klotzsch, ex Baill. Adansonia, v. (1864–65) 320 = S. Aucuparium.
serratum, Klotzsch, ex Baill. l. c. = S. Aucuparium.
sterculiaceum, Wall. Cat. n. 7974 = Baccaurea bracteata.
stylare, *Muell. Arg. in Linnaea*, xxxii. (1863) 119.—Venezuel.
suberosum, *Muell. Arg. l. c.* xxxiv. (1865–66) 217.—Ins. Barbados.
sylvaticum, Torr. Bot. Mex. Bound. 201 = Stillingia sylvatica.
tanguinum, Muell. Arg. in Linnaea, xxxii. (1863) 116 = Stillingia obtusifolia.
terminale, Muell. Arg. l. c. = Stillingia terminalis.
zelayense, H. B. & K. Nov. Gen. et Sp. ii. 65 = Stillingia zelayensis.

SAPONARIA, Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 146.
BOOTIA, Neck. ex DC. Prod. i. 365 (1824).
PROTEINIA, Reichb. Nom. 205 (1841).
SMEGMATHAMNIUM, Fenzl, ex Reichb. Ic. Fl. Germ. vi. 42. t. 244 (1841).
VACCARIA, Medic. Phil. Bot. i. 96 (1789).
acerosa, Boiss. Diagn. Ser. II. i. 69 = Gypsophila acerosa.
aenesia, *Heldr. in Atti Congr. Bot. Firenze*, 1874 (1876), 238.—Graec.
aetnensis, *Rafin. Chl. Aetn.* 9.—Ital.
alpina, Pall. Reise, iii. 34 = Silene paucifolia.
alsinoides, Viv. Fl. Cors. Prod. App. ii. 7, in obs. = ocymoides.
amplissima, Mill. Gard. Dict. ed. VIII. n. 4 (*amplicimus*) = S. Vaccaria.
andicola, *Kunze, ex Rohrb. Monog. Silene*, 231.—Chili.
artegioides, Boiss. in Tchihat. Asie Min. Bot. t. 11, sphalm. = *ortegioides*.
astroides, Fenzl, in Flora, xxvi. (1843) 402 = tridentata.
atocioides, Boiss. Diagn. Ser. I. i. 16 = prostrata.
barradensis, Boiss. l. c. viii. 72 = Gypsophila hirsuta.
Baumgartenii, Janka, in Linnaea, xxx. (1859–60) 559 = Silene dinarica.
bellidifolia, Lapeyr. Hist. Abr. Pl. Pyr. 239 = Valeriana apula.
bellidifolia, *Sm. Spicil. Bot.* 5.—Europ. or.
Bodeana, *Boiss. Fl. Orient.* i. 527.—Persia.
caespitosa, Bivona, in Rafin. Statis. Gen. Sicil. 27 = depressa.
caespitosa, *DC. Rapp. Voy.* ii. 78; *ex ej. Fl. Fr. Suppl.* 601.—Mont. Pyren.
caespitosa, Sibth. & Sm. Fl. Graec. iv. 79. t. 339 = Silene Saxifraga.
calabrica, *Guss. Pl. Rar.* 164. t. 31.—Ital.; Graec.
calabrica, Heldr. ex Nym. Consp. 98 = graeca.
Calverti, Boiss. Diagn. Ser. II. v. 51 = prostrata.
cerastoides, *Fisch. ex C. A. Mey. Verz. Pfl. Cauc.* 212.—Reg. Casp.
chloraefolia, *Kunze, Ind. Hort. Lips.* (1846).—*Cf. Linnaea*, xxiv. (1851) 163.—As. Min.

SAPONARIA:—
corrugata, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 237.—Turkestan.
cretica, Linn. Sp. Pl. ed. II. 584 = Tunica cretica.
depressa, *Bivona, Stirp. Nov. Kar. Sicil.* ii. [15].—Sicil.
diffusa, Lam. Fl. Fr. ii. 540 = Gypsophila repens.
dioica, Cham. & Schlecht. in Linnaea, i. (1826) 38 = Silene antirrhina.
dioica, Moench, Meth. 76 = Lychnis dioica.
elegans, Lapeyr. Hist. Abr. Pl. Pyr. 238 = caespitosa.
Elisabethae, Fenzl, ex Nym. Consp. 93 = Silene Elisabethae.
fasciculata, Boiss. & Reut. Diagn. Ser. I. viii. 73 = Gypsophila fasciculata.
fastigiata, Lam. Fl. Fr. ii. 541 = Gypsophila fastigiata.
filicaulis, Boiss. Diagn. Ser. I. viii. 71 = Gypsophila hirsuta.
filipes, Boiss. l. c. Ser. II. i. 69 = Gypsophila floribunda.
floribunda, Boiss. l. c. 70 = Gypsophila floribunda.
fruticulosa, Bory & Chaub. Nouv. Fl. Pélop. 26. t. 12 = Gypsophila fruticulosa.
glutinosa, *Bieb. Fl. Taur. Cauc.* i. 322.—Reg. Medit. et Cauc.
graeca, *Boiss. Fl. Orient.* i. 528.—Graecia.
Griffithiana, *Boiss. Diagn.* Ser. II. i. 70.—Afghan.
hirsuta, Labill. Ic. Pl. Syr. Dec. iv. 9 = Gypsophila hirsuta.
hirta, Steud. Nom. ed. I. 732, sphalm. = praec.
hispanica, Mill. Gard. Dict. ed. VIII (in Errat.) = S. Vaccaria.
holopetala, Ledeb. Fl. Ross. i. 301 = prostrata.
hybrida, Mill. Gard. Dict. ed. VIII. n. 2 = officinalis.
illyrica, Linn. Mant. i. 70 = Tunica illyrica.
inclusa, C. Koch, in Linnaea, xv. (1841) 711 = S. Vaccaria?
Kotschyi, *Boiss. Diagn.* Ser. I. i. 16.—As. Min.
liniflora, *Boiss. & Haussk. in Boiss. Fl. Orient.* i. 525.—Mesopot.
lutea, *Linn. Sp. Pl.* ed. II. 585 (*excl. syn. Bocc.*).—Europ.
lutea, Pall. ex Ledeb. Fl. Ross. i. 306 = Silene paucifolia.
mesogitana, *Boiss. Diagn.* Ser. I. i. 16.—As. Min.
mollis, Boiss. l. c. viii. 72 = Gypsophila hirsuta.
multiflora, Hort. ex Regel, Gartenfl. i. (1852) 101 = calabrica.
muralis, Lam. Fl. Fr. ii. 540 = Gypsophila muralis.
nervosa, Gilib. Fl. Lituan. ii. 163 = officinalis.
noctiflora, Fenzl, ex Nym. Consp. 93 = Silene noctiflora.
nodiflora, *Boiss. Diagn.* Ser. I. i. 15.—Syria.
ocymifolia, Salisb. Prod. 302 = seq.
ocymoides, *Linn. Sp. Pl.* 409.—Europ. austr.
officinalis, *Linn. l. c.* 408.—Europ.; As. occ.
officinarum, Rupr. Fl. Cauc. 176 = praec.
orientalis, *Linn. Sp. Pl.* 409.—Oriens.
ortegioides, Boiss. & Bal. Diagn. Ser. II. vi. 25 = Gypsophila ortegioides.
oxyodonta, Boiss. Fl. Orient. i. 525 = S. Vaccaria.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 74.—As. Min.
parvula, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 181.—Turkestan.
pentagona, Hort. ex Steud. Nom. ed. II. ii. 512 = S. Vaccaria.
perfoliata, Gilib. Fl. Lituan. ii. 163; Roxb. ex Willd. Enum. Hort. Berol. 464 = S. Vaccaria.
persica, *Boiss. Diagn.* Ser. I. i. 18.—Persia.
picta, Boiss. in Tchihat. Asie Min. Bot. i. 202 = Gypsophila picta.
plumbaginea, *Stschégl. in Bull. Soc. Nat. Mosc.* xxvi. 1. (1853) 323. t. 4.—Reg. Cauc.
polygonoides, Jaub. & Spach, Illustr. v. 2. t. 402 = Gypsophila hirsuta.
porrigens, Linn. Mant. ii. 239 = Gypsophila porrigens.
prostrata, *Willd. Enum. Hort. Berol.* 465.—As. Min.
pulvinaris, *Boiss. Diagn.* Ser. I. viii. 74.—As. Min.; Syria.
Pumilio, Boiss. l. c. i. 18 = pulvinaris.
pungens, Bunge, in Ledeb. Fl. Alt. ii. 133 = Acanthophyllum spinosum.
repens, Lam. Fl. Fr. ii. 542 = ocymoides.
rubra, Lam. l. c. 541 = S. Vaccaria.
saxatilis, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) 13.—Hispan.

SAPONARIA:—
segetalis, Neck. Delic. Gallo-Belg. i. 194 = S. Vaccaria.
Sewerzowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 246.—Turkestan.
sicula, Rafin. Specch. ii. 7 = depressa.
Smithii, Ser. in DC. Prod. i. 367 = Silene Saxifraga.
Struthium, Loefl. Iter Hisp. 73 = Gypsophila Struthium.
syriaca, *Boiss. Diagn.* Ser. I. viii. 73.—Syria.
thessala, Jaub. & Spach, Illustr. v. 2 = Gypsophila hirsuta.
thymifolia, Boiss. Diagn. Ser. I. i. 17 = Gypsophila hirsuta.
tridentata, *Boiss. l. c.*—Syria.
tubulosa, F. Muell. Native Pl. Vict. i. 136 = Gypsophila tubulosa.
Vaccaria, *Linn. Sp. Pl.* 409.—Europ.; As. Min.; Sibir.
villosa, Rafin. Fl. Ludov. 93 (Quid?).—Am. bor.
viscosa, Dulac, Fl. Hautes-Pyr. 264 = ocymoides.
viscosa, *C. A. Mey. Verz. Pfl. Cauc.* 212.—Armen.; Persia.
vulgaris, Pall. Reise, iii. 655 = officinalis.

SAPOTA, Plum. ex Mill. Gard. Dict. ed. VI (1752) = **Achras**, Linn. (Sapotac.).
Achras, Mill. Gard. Dict. ed. VIII. n. 1 = A. Sapota.
armata, A. DC. in DC. Prod. viii. 175 = Sideroxylon tomentosum.
australis, A. DC. l. c. = Sideroxylon australe.
Baueri, Montr. in Mém. Acad. Lyon, x. (1860) 229.—N. Caled.
cerasifera, Welw. Apont. 585. n. 17.—Afr. trop.
costata, A. DC. in DC. Prod. viii. 175 = Sideroxylon costatum.
curvifolia, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 68 = Lucuma curvifolia.
elengoides, A. DC. in DC. Prod. viii. 176 = Sideroxylon tomentosum.
elongata, Gaertn. f. Fruct. iii. 136.—Hab.?
Endlicheri, Montr. in Mém. Acad. Lyon, x. (1860) 228.—N. Caled.
ferruginea, Walp. Rep. vi. 455 = A. Sapota.
gonocarpa, Mart. & Eichl. ex Miq. in Mart. Fl. Bras. vii. 60.—Bras.
Lessertii, A. DC. in DC. Prod. viii. 174 = Sideroxylon Lessertii.
mammosa, Mill. Gard. Dict. ed. VIII. n. 2 = Lucuma mammosa.
marginata, Decne. ex Webb, in Hook. Niger Fl. 169. t. 13.—Ins. Cap. Virid.
Mulleri, Blume, ex Bleekrod. in Ann. Sc. Nat. Sér. IV. vii. (1857) 225 = Mimusops Kauki.
nigra, J. F. Gmel. Syst. 750 = Diospyros Ebenum.
parvifolia, A. DC. in DC. Prod. viii. 175 = Sideroxylon parvifolium.
Pohlmaniana, F. Muell. Fragm. v. 184 = Sideroxylon Pohlmanianum.
polita, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 517.—Cuba.
pyrulifera, A. Gray, in Proc. Am. Acad. v. (1862) 328.—Ins. Fiji.
rugosa, Griseb. Fl. Brit. W. Ind. 399.—Ind. occ.
sandwicensis, A. Gray, in Proc. Am. Acad. v. (1861–62) 328.—Ins. Sandvic.
Sellowiana, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 67 = Lucuma marginata.
sericea, A. DC. in DC. Prod. viii. 176 = A. sericea.
Sideroxylon, Griseb. Fl. Brit. W. Ind. 399 = A. Sapota.
spectabilis, Miq. Fl. Ind. Bat. Suppl. 580.—Sumatra.
tomentosa, A. DC. in DC. Prod. viii. 175 = Sideroxylon tomentosum.
venosa, A. DC. l. c. 174 = Lucuma venosa.
vitiensis, A. Gray, in Proc. Am. Acad. v. (1862) 328.—Ins. Fiji.

SAPRANTHUS, Seem. Journ. Bot. iv. (1866) 369. t. 54 = **Porcelia**, Ruiz & Pav. (Anonac.).
nicaraguensis, Seem. l. c.—Am. centr.

SAPRIA, Griff. in Proc. Linn. Soc. i. (1844) 216. *CYTINACEAE*, Benth. & Hook. f. iii. 119.
Griffithii, Hook. f. in DC. Prod. xvii. 112, in syn. sphalm. = seq.
himalayana, *Griff. in Proc. Linn. Soc.* i. (1844) 217.—Assam.

SAPROSMA, Blume, Bijdr. 956 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 131.
DYSODIDENDRON, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 1.
DYSOSMIA, Miq. Fl. Ind. Bat. ii. 325 (1859).
arboreum, *Blume, Bijdr.* 956.—Java.
ceylanicum, *Bedd. Forest. Man. Bot.* 134/12.—Ind. or.
consimile, *Kurz, For. Fl. Brit. Burma*, ii. 29.—Reg. Himal.; Burma.
corymbosum, *Bedd. Forest. Man. Bot.* 134/12.—Ind. or.
dispar, *Hassk. in Flora*, xxviii. (1845) 226.—Java.
fragrans, *Bedd. Forest. Man. Bot.* 134/11.—Ind. or.
fruticosum, *Blume, Bijdr.* 956.—Java.
glomeratum, Bedd. Forest. Man. Bot. 134/11 = indicum.
hirsutum, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 251.—Sumatra.
indicum, *Dalz. in Hook. Kew Journ.* iii. (1851) 37.—Ind. or.
scabridum, *Bedd. Forest. Man. Bot.* 134/12.—Zeylan.
ternatum, *Benth. & Hook. f. Gen.* ii. 131.—Reg. Himal.; Burma.
Wightii, Bedd. Forest. Man. Bot. 134/11 = ceylanicum.

SARACA, Linn. Mant. i. 13 (1767). *LEGUMINOSAE*, Benth. & Hook. f. i. 583.
JONESIA, Roxb. in As. Res. iv. (1795) 355.
arborescens, Burm. f. Fl. Ind. 85 = indica.
cauliflora, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 272.—Malacca.
declinata, *Miq. Fl. Ind. Bat.* i. 1. 84, 1080.—Sumatra.
indica, *Linn. Mant.* i. 98.—Ind. or.; Malaya.
Lobbiana, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 272.—Burma.
macroptera, *Miq. Fl. Ind. Bat.* i. 1. 1080.—Sumatra.
minor, Miq. l. c. 84 = indica.
obtusifolia, *Miq. l. c.* 85.—Java.
palembanica, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 272.—Sumatra.
triandra, *Baker, l. c.*—Burma; Malaya.
Zollingeriana, Miq. Fl. Ind. Bat. i. 1. 84 = indica.

SARACENA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Sarracenia**, Linn.

SARACENIA, Spreng. Gen. i. 433 (1830) = praec.

SARACHA, Ruiz & Pav. Prod. 31. t. 34 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 891.
BELLINIA, Roem. & Schult. Syst. iv. 56 (1819).
DISKION, Rafin. Sylva Tellur. 55 (1838).
JALTOMATA, Schlecht. Ind. Sem. Hort. Hal. (1838).
JALTONIA, Steud. Nom. ed II. i. 797 (1840).
acutifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 449 (Physalidis sp. ?).—Calif.
alata, *Dun. in DC. Prod.* xiii. 1. 431.—Peruv.
allogona, *Schlecht. in Linnaea*, xix. (1847) 308.—Mexic.
angulata, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 133, nomen.—Mexic.
auriculata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 448.—N. Granat.
biflora, Ruiz & Pav. Fl. Per. ii. 42. t. 179 = Hebecladus biflorus.
Boissieri, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 93 (= *Lycioplesium Boissieri*).—Venezuel.
brasiliensis, Klotzsch, in Linnaea, xiv. (1840) 290 = Athenaea pogogena.
ciliata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 446.—Peruv.
conoides, Benth. & Hook. f. Gen. ii. 891 = Chamaesaracha sordida ?
conspersa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 449.—Mexic.
contorta, *Ruiz & Pav. Fl. Per.* ii. 43. t. 180.—Peruv.
Coronopus, A. Gray, in Proc. Am. Acad. x. (1874) 62 = Chamaesaracha Coronopus.
dentata, *Ruiz & Pav. Fl. Per.* ii. 43. t. 179.—Peruv.
diffusa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 447 = laxa.
frondosa, *Miers, l. c.* xi. (1853) 93 = Poecilochroma frondosum.
Funkiana, Miers, l. c. = Poecilochroma Funckianum.
geniculata, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 133 = Physalis geniculata.
glabrata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 448.—Mexic.
glandulosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 450.—N. Granat.
guttata, Miers, l. c. xi. (1853) 93 = Poecilochroma guttatum.
Jaltomata, *Schlecht. Ind. Sem. Hort. Hal.* (1839) 10; *ex Dun. in DC. Prod.* xiii. 1. 432.—Mexic.
Jaltonia, Steud. Nom. ed. II. i. 796, in syn., sphalm. = praec.
laxa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 447.—Mexic.
Lindeniana, Miers, l. c. xi. (1853) 93 = Poecilochroma Lindenianum.
Lobbiana, Miers, l. c. = Poecilochroma Lobbianum.
maculata, Miers, l. c. = Poecilochroma maculatum.
micrantha, DC. Cat. Hort. Monsp. (1813) 58, nomen = Zuccagniana.
Miersii, Dun. & A. DC. in DC. Prod. xiii. 1. 684 = laxa.
nana, A. Gray, in Proc. Am. Acad. x. (1874) 62 = Chamaesaracha nana.
peruviana, F. G. Dietr. Vollst. Lexik. Gaertn. viii. 522 (sp. confusa).
procumbens, Math. ex Dun. & A. DC. in DC. Prod. xiii. 1. 683 = propinqua.
procumbens, *Ruiz & Pav. Fl. Per.* ii. 43. t. 180.—Peruv.
propinqua, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 446.—Peruv.
pubescens, *Humb. ex Roem. & Schult. Syst.* iv. 690.—Peruv.
punctata, *Ruiz & Pav. Fl. Per.* ii. 42. t. 178.—Peruv.
quitoënsis, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 93 = Poecilochroma quitensis.
Sellowiana, *Miers. l. c.*—Bras.
solanacea, Miers, l. c. iii. (1849) 450 = Bassovia solanacea.
sordida, A. Gray, in Proc. Am. Acad. x. (1874) 62 = Chamaesaracha sordida.
stapelioides, *Decne. ex Regel, Ind. Sem. Hort. Petrop.* (1864) *Suppl.* 18.—Hab. ?
umbellata, DC. Cat. Hort. Monsp. (1813) 142 = procumbens.
umbellata, *G. Don, Gen. Syst.* iv. 451.—Peruv.
vestita, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iii. (1849) 449.—N. Granat.
villosa, *G. Don, Gen. Syst.* iv. 452.—Peruv.
viscosa, *Link, ex D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 323.—Peruv.; Chili.
viscosa, Schrad. Ind. Sem. Hort. Gotting. (1832) 5; ex Dun. in DC. Prod. xiii. 1. 433 (Physalidis sp.).—Mexic.
Zuccagniana, *G. Don, Gen. Syst.* iv. 452.—Am. austr.

SARAGODRA, Hort. ex Steud. Nom. ed. II. ii. 513 (1841) = **Gelonium**, Roxb. (Euphorb.).
glabra, Hort. ex Steud. l. c. = G. multiflorum.

SARANA, Fisch. ex Baker, in Journ. Linn. Soc. xiv. (1874) 251 = **Fritillaria**, Tourn. (Liliac.).
edulis, Fisch. ex Baker, l. c. 274 = F. camschatcensis.

SARANTHE, Eichl. in Berl. Phys. Abh. 1883 (1884), 85 = **Myrosma**, Linn. f. (Scitam.).
cuiabensis, Eichl. l. c. 86 = Maranta cuyabensis.
Klotzschiana, Eichl. l. c. = Maranta Klotzschiana.
leptostachya, Eichl. l. c. = Maranta leptostachya.
Moritziana, Eichl. l. c. = Maranta Moritziana.
pygmaea, Eichl. l. c. = Maranta pygmaea.
Riedeliana, Eichl. l. c. = Maranta Riedeliana.
unilateralis, Eichl. l. c. = Myrosma unilaterale.

SARCANDRA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 348 = **Chloranthus**, Sw.
chloranthoides, Gardn. l. c. = C. brachystachys.

SARCANTHEMUM, Cass. in Bull. Soc. Philom. (1818) 74 = **Psiadia**, Jacq. (Compos.).
Coronopus, Cass. l. c. = P. Coronopus.

SARCANTHERA, Rafin. Fl. Tellur. iv. 64 (1836) = **Gymnostachyum**, Nees (Acanth.).
venusta, Rafin. l. c. = G. venustum.

SARCANTHIDION, Baill. Adansonia, xi. (1874) 199. *OLACINEAE*.
sarmentosum, *Baill. l. c.*—N. Caled.

SARCANTHUS, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 209 = **Heliotropium**, Linn. (Borag.).
asperrimus, Anderss. l. c.—Ins. Galapag.

SARCANTHUS, Lindl. Coll. Bot. t. 39 B (1821). *ORCHIDEAE*. Benth. & Hook. f. iii. 580.
arietinus, *Reichb. f. in Gard. Chron.* (1869) 416.—Assam.
armeniacus, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 219.—Afr. trop.
aspersus, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 297.—Burma.
belophorus, *Reichb. f. in Gard. Chron.* (1883) II. 262.—Hab. ?
callosus, Reichb. f. in Bonplandia, v. (1857) 41 = Cleisostoma callosum.
chrysomelas, *Reichb. f. in Gard. Chron.* (1869) 662.—Burma.
croceus, *Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 634, nomen.—Ins. Philipp.
dealbatus, *Reichb. f. in Walp. Ann.* vi. 892.—Ins. Philipp.
densiflorus, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 136 = Cleisostoma spicatum.
erinaceus, *Reichb. in Bot. Zeit.* xxii. (1864) 298.—Burma.
filiformis, *Lindl. Bot. Reg.* (1842) *Misc.* 61.—Reg. Himal.
filiformis, Wight, Ic. t. 1684 = Saccolabium filiforme.
flexus, *Reichb. f. in Gard. Chron.* (1881) II. 492.—Borneo.
geminatus, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 243.—Ins. Molucc.
guttatus, Lindl. Bot. Reg. t. 1443 = Rhynchostylis retusa.
Hincksiana, *Reichb. f. in Gard. Chron.* (1878) I. 73.—Hab. ?
insectifer, *Reichb. f. in Bot. Zeit.* xv. (1857) 159.—Ind. or.; Burma.
laxus, *Reichb. f. l. c.* xxiv. (1866) 378.—Burma.
Lendyanus, *Reichb. f. in Gard. Chron.* (1884) I. 44.—Annam.
macrodon, *Reichb. f. l. c.* (1872) 1555.—Ind. or.
mirabilis, *Reichb. f. l. c.* (1878) II. 300.—Burma ?
nagarensis, *Reichb. f. in Seem. Fl. Vit.* 298.—Ins. Fiji.
ornithorrhynchus, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 219.—Hab. ?
oxyphyllus, *Wall. ex Lindl. Bot. Reg.* (1840) *Misc.* 58.—Burma.
pallidus, *Lindl. Bot. Reg.* (1840) *Misc.* 78.—Reg. Himal.; Burma.
paniculatus, *Lindl. in Bauer, Illustr. Orch. Pl. Gen.* t. 9.—China.
Parishii, *Hook. Bot. Mag.* t. 5217.—Burma.
pauciflorus, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 21.—Sumatra.
pauciflorus, Wight, Ic. t. 1747 = peninsularis.
peninsularis, *Dalz. in Hook. Kew Journ.* iii. (1851) 343.—Ind. or.
praemorsus, Lindl. ex Spreng. Syst. iii. 721 = Saccolabium papillosum.
pugioniformis, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 219.—Venezuel.
racemifer, Reichb. f. in Walp. Ann. vi. 891 = pallidus.
roseus, Wight, Ic. t. 1685 = Saccolabium filiforme.
rostratus, *Lindl. Coll. Bot.* t. 39 B.—China.
secundus, *Griff. Notul.* iii. 362.—Reg. Himal.
Stowellianus, Batem. in Hook. Bot. Mag. sub t. 5630 = erinaceus.
striolatus, *Reichb. f. in Gard. Chron.* (1882) II. 168.—Ins. Philipp.
suaveolens, *Reichb. f. in Bonplandia*, v. (1857) 40.—Java.
subulatus, *Reichb. f. l. c.* 41.—Java.
succisus, *Lindl. in Bot. Reg.* t. 1014.—China.
teretifolius, *Lindl. Gen. et Sp. Orch.* 234.—China.
teretifolius, *Reichb. f. in. Trans. Linn. Soc.* xxx. (1874) 136.—Burma.
terminus, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 15.—Java.
timorensis, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 365.—Ins. Timor.
tricolor, Reichb. f. in Bonplandia, iii. (1855) 219 = pallidus.
Walkerianus, Wight, Ic. t. 1686 = Saccolabium roseum.
Williamsoni, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 353.—Burma.

SARCATHRIA, Rafin. Fl. Tellur. iii. 47 (1836) = **Salicornia**, Tourn. (Chenop.).
ambigua, Rafin. l. c. = Arthrocnemum ambiguum.
indica, Rafin. l. c. = Arthrocnemum indicum.
procumbens, Rafin. l. c. = Sal. herbacea.
radicans, Rafin. l. c. = Sal. fruticosa.
strobilacea, Rafin. l. c. = Halocnemum strobilaceum.
virginica, Rafin. l. c. = Sal. fruticosa.

SARCAULIS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 310. *SAPOTACEAE.*
macrophyllus, *Radlk. l. c.* 311.—Bras.

SARCHOCHILUS, Vidal, Phan. Cuming. Philipp. 150 (1885) = **Sarcochilus**, R. Br. (Orchid.).

SARCINANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. 196 (1857) = **Carludovica**, Ruiz & Pav. (Cyclanth.).
utilis, Oerst. l. c. 197 = C. utilis.

SARCOBATUS, Nees, in Neuwied, Reise Nord-Am. ii. 446 (1841). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 76.
FREMONTIA, Torr. in Emory, Notes Mil. Reconnois. 95. 317. t. 3 (1845).
Maximiliani, *Nees, in Neuwied, Reise Nord-Am.* ii. 447.—Am. bor.
vermiculatus, Torr. in Emory, Notes Mil. Reconnois. 149 = Maximiliani.

SARCOBODIUM, Beer, Prakt. Orch. 306 (1854) = **Bulbophyllum**, Thou. (Orchid.).
Lobbii, Beer, l. c. = B. Lobbii.

SARCOCA, Rafin. Fl. Tellur. iii. 55 (1836) = **Phytolacca**, Tourn.
abyssinica, Rafin. l. c. = P. abyssinica.
dioica, Rafin. l. c. = P. dioica.
icosandra, Rafin. l. c. = P. icosandra.

SARCOCALYX, Walp. in Linnaea, xiii. (1839) 479 = **Aspalathus**, Linn. (Legumin.).
capensis, Walp. l. c. 480 = A. sarcoides.

SARCOCALYX, Zipp. in Alg. Konst- en Letter-Bode, i. (1829) 298 = **Exocarpus**, Labill. (Santal.)
miniatus, Zipp. l. c. = E. latifolius.

SARCOCAMPSA, Miers, in Trans. Linn. Soc. xxviii. (1872) 408 = **Salacia**, Linn. (Celastrin.).
campestris, Miers, l. c. = Sal. campestris.
silvestris, Miers, l. c. 409 = Sal. sylvestris.

SARCOCAPNOS, DC. Syst. ii. 129 (1821). *PAPAVERACEAE*, Benth. & Hook. f. i. 56.
APLECTROCAPNOS, Boiss. & Reut. Diagn. Ser. I. v. 79 (1844).
baetica, *Nym. Consp.* 26.—Hispan.
crassifolia, *DC. Syst.* ii. 130.—Mauritania.
enneaphylla, *DC. l. c.* 129.—Lusitan.
speciosa, Boiss. Diagn. Ser. II. i. 14 = crassifolia.

SARCOCARPON, Blume, Bijdr. 21 (1825) = **Kadsura**, Kaempf. (Magnol.).
scandens, Blume, l. c. = K. scandens.

SARCOCAULON, Sweet, Hort. Brit. ed. I. 73 (1826). *GERANIACEAE*, Benth. & Hook. f. i. 272.
Burmanni, *Sweet, l. c.*—Afr. austr.
L'Heritieri, *Sweet, l. c.*—Afr. austr.
Patersonii, *Eckl. & Zeyh. Enum.* 57.—Afr. austr.

SARCOCEPHALUS, Afzel. ex R. Br. in Tuckey, Congo, App. v. 467 (1818); et ex Sabine, in Trans. Hort Soc. v. (1824) 442. t. 18. *RUBIACEAE*, Benth. & Hook. f. ii. 29.
CEPHALINA, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 105 (1828).
PLATANOCARPUM, Korth. Obs. Naucl. Ind. 18 (1839).
Bartlingii, *Miq. Fl. Ind. Bat.* ii. 133.—Ins. Philipp.
buruensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 179.—Malaya.
Cadamba, Kurz, For. Fl. Brit. Burma, ii. 63; et in Journ. As. Soc. Beng. xlvi. (1877) II. 125 = Anthocephalus morindaefolius.

SARCOCEPHALUS:—
cordatus, *Miq. Fl. Ind. Bat.* ii. 133.—As. et Austral. trop.
dasyphyllus, *Miq. l. c. Suppl.* 538.—Sumatra.
esculentus, *Afzel. ex Sabine, in Trans. Hort. Soc.* v. (1824) 442. t. 18.—Afr. trop.
glaberrimus, *Miq. Fl. Ind. Bat.* ii. 134.—Ins. Philipp.
Horsfieldii, Miq. l. c. = cordatus.
Junghuhnii, *Miq. l. c.*—Malaya.
mitragynus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 180.—Ins. Amboin.
Russeggeri, *Kotschy, ex Schweinf. Reliq. Kotsch.* 49. t. 33.—Afr. trop.
subditus, *Miq. Fl. Ind. Bat.* ii. 134.—Malaya.
undulatus, *Miq. l. c.* 133.—Ins. Celebes.

SARCOCHILUS, R. Br. Prod. 332 (1810). *ORCHIDEAE*, Benth and Hook. f. iii. 575.
ADENONCOS, Blume, Bijdr. 381 (1825).
CAMAROTIS, Lindl. in Wall. Cat. n. 7329; ex ej. Gen. et Sp. Orch. 219 (1833).
CHILOSCHISTA, Lindl. Bot. Reg. (1832) sub t. 1522.
CYLINDROCHILUS, Thw. Enum. Pl. Zeyl. 307 (1861).
DENDROCOLLA, Blume, Bijdr. 286. t. 67 (1825).
GROSOURDYA, Reichb. f. in Bot. Zeit. xxii. (1864) 297.
GUNNIA, Lindl. Bot. Reg. (1834) sub t. 1699.
MICROPERA, Lindl. Bot. Reg. sub t. 1522 (1832).
ORNITHARIUM, Lindl. & Paxt. Flow. Gard. i. (1850–51) 188. f. 117.
ORSIDICE, Reichb. f. in Bonplandia, ii. (1854) 93.
PTEROCERAS, Hassk. in Flora, xxv. (1842) II. Beibl. 6.
STEREOCHILUS, Lindl. in Journ. Linn. Soc. iii. (1859) 38.
THRIXSPERMUM, Lour. Fl. Cochinch. 519 (1790), nomen prius.
acuminatissimus, *Reichb. f. in Walp. Ann.* vi. 498.—Java.
adversus, *Hook. f. Fl. N. Zel.* i. 241.—N. Zel.
amplexicaulis, *Reichb. f. in Walp. Ann.* vi. 499.—Java.
anceps, *Reichb. f. l. c.* 500.—Java.
angustifolius, *Reichb. f. l. c.*—Java.
Arachnites, *Reichb. f. l. c.* 498.—Reg. Himal.; Malaya.
Armitii, *F. Muell. Fragm.* ix. 49.—Austral.
aurifer, *Reichb. f. in Walp. Ann.* vi. 498.—China.
australis, Reichb. f. l. c. 501 = parviflorus.
Baileyi, F. Muell. ex Benth. Fl. Austral. vi. 291 = Taeniophyllum Muelleri.
Barklyanus, F. Muell. Fragm. i. 89 = parviflorus.
Beckleri, F. Muell. Census, 111 = Cleisostoma Beckleri.
Berkeleyi, *Reichb. f. in Gard. Chron.* (1882) I. 557, in syn.—Ins. Nicobar.
brevilabris, F. Muell. Census, 111 = Cleisostoma brevilabre.
breviscapa, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 332.—N. Zel.
calcaratus, F. Muell. Fragm. ii. 181 = Cleisostoma tridentatum.
Calceolus, *Lindl. Bot. Reg.* (1846) t. 19.—Ins. Philipp.
Ceciliae, *F. Muell. Fragm.* v. 42. t. 42.—Austral.
compressus, *Reichb. f. in Walp. Ann.* vi. 498.—Java.
croceus, *Lindl. Bot. Reg.* (1846) sub t. 19.—Ins. Philipp.
dilatatus, F. Muell. Fragm. i. 191 = olivaceus.
divitiflorus, *F. Muell. ex Benth Fl. Austral.* vi. 292.—Austral.
emarginatus, *Reichb. f. in Walp. Ann.* vi. 500.—Java.
erectus, F. Muell. Census, 111 = Cleisostoma erectum.
falcatus, *R. Br. Prod.* 332.—Austral.
Fitzgeraldi, *F. Muell. Fragm.* vii. 115.—Austral.
fusco-luteus, *Lindl. Bot. Reg.* (1847) sub t. 18.—Borneo.
gracilentus, *Reichb. f. in Walp. Ann.* vi. 499.—Malaya.
Gunnii, F. Muell. Fragm. i. 90 = parviflorus.
Hartmanni, *F. Muell. l. c.* viii. 248.—Austral.
Hillii, *F. Muell. l. c.* ii. 94.—Austral.
Hystrix, *Reichb. f. in Walp. Ann.* vi. 500.—Burma.
japonicus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 206.—Japon.
Keffordii, F. Muell. Census, Suppl. ii. 3 = Cleisostoma Keffordii.
lilacinus, *Griff. Notul.* iii. 334.—Malaya.
Macphersonii, *F. Muell. Fragm.* viii. 248.—Austral.

SARCOCHILUS:—
maculatus, Pfitzer, Vergl. Morph. Orch. 15; Benth. & Hook. f. Gen. iii. 575 (Saccolabii sp.).—Ind. or.
minimus, *Reichb. f. in Walp. Ann.* vi. 500.—Java.
nepalensis, Spreng. Syst. iii. 721 = Saccolabium calceolare.
obtusus, *Reichb. f. in Walp. Ann.* vi. 500.—Java.
olivaceus, *Lindl. Bot. Reg.* (1839) *Misc.* 32.—Austral.
pallidus, *Reichb. f. in Walp. Ann.* vi. 500.—Malaya.
parviflorus, *Lindl. Bot. Reg.* (1838) *Misc.* 34.—Austral.
philippinensis, *Vidal, Phan. Cuming. Philipp.* 82, 150.—Ins. Philipp.
phyllorrhizus, *F. Muell. Fragm.* v. 201.—Austral.
pictus, Reichb. f. in Walp. Ann. vi. 501 = olivaceus.
praemorsus, Spreng. Syst. iii. 721 = Saccolabium papillosum.
pulchellus, *Trimen, Syst. Cat. Pl. Ceyl.* 89.—Zeylan.
purpurascens, *Reichb. f. in Walp. Ann.* vi. 498.—Java.
purpureus, *Benth. & Hook. f. (in Herb. Kew.)*.—Ind. or.
pusillus, *Reichb. f. in Walp. Ann.* vi. 500.—Java.
rhopalorrachis, *Reichb. f. l. c.*—Java.
rubricentrum, *Fitzg. in Gard. Chron.* (1880) II. 38.—Austral.
serraeformis, Reichb. f. in Walp. Ann. vi. 498 = S. Arachnites.
Sillemianus, *Reichb. f. in Gard. Chron.* (1882) I. 524, in syn.—Burma?
spurius, *Reichb. f. in Ann. Walp.* vi. 500.—Java.
subulatus, *Reichb. f. in Walp. Ann.* vi. 500.—Java.
teres, *Reichb. f. l. c.* 499.—Java.
tridentatus, Reichb. f. l. c. 500 = Cleisostoma tridentatum.
unguiculatus, *Lindl. in Bot. Reg.* (1840) *Misc.* 67.—Ins. Philipp.
usneoides, *Reichb. f. in Walp. Ann.* vi. 497.—Reg. Himal.
Zollingeri, *Reichb. f. l. c.* 500.—Java.

SARCOCHLAENA, Spreng. Syst. ii. 594 (1825) = **Sarcolaena**, Thou. (Chlaenac.).

SARCOCHLAMYS, Gaudich. Bot. Voy. Bonite, t. 89 (1844–52). *URTICACEAE*, Benth. & Hook. f. iii. 389.
pulcherrima, *Gaudich. l. c.*—Ind. or.; Malay.

SARCOCLINIUM, Wight, Ic. v. II. 24. t. 1887–88 (1852) = **Agrostistachys**, Dalz. (Euphorb.).
Gaudichaudi, Baill. Etud. Gén. Euph. 310, nomen = A. Gaudichaudi.
Hookeri, Thw. Enum. Pl. Zeyl. 279 = A. Hookeri.
longifolium, Wight, Ic. t. 1887–88 = A. longifolia.
sessilifolium, Kurz, in Flora, lviii. (1875) 31.—Singapore.

SARCOCOCCA, Lindl. Bot. Reg. t. 1012 (1826). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 266.
AUSTROBUXUS, Miq. Fl. Ind. Bat. Suppl. 444 (1860).
LEPIDOPELMA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 118. t. 22 (1862).
coriacea, Sweet, Hort. Brit. ed. I. 491 = pruniformis.
Hookeriana, Baill. Monog. Bux. 53 = pruniformis.
laurifolius, Kunth, Enum. Pl. v. 190 = pruniformis.
nepalensis, Royle, Illustr. Bot. Himal. 327 = pruniformis.
pruniformis, *Lindl. Bot. Reg.* t. 1012.—Ind. or.; Malaya.
salicifolia, Baill. Monog. Bux. 49 = pruniformis.
saligna, Muell. Arg. in DC. Prod. xvi. I. 11 = pruniformis.
sumatrana, Blume, Mus. Bot. Lugd. Bat. ii. 191 = pruniformis.
trinervia, Wight, Ic. t. 1877 = pruniformis.
zeylanica, Baill. Monog. Bux. 52 = pruniformis.
Zollingeri, Baill. l. c. 54 = pruniformis.

SARCOCODON, N. E. Br. in Journ. Linn. Soc. xvii. (1878) 169. *ASCLEPIADEAE.*
speciosus, *N. E. Br. l. c.* 170.—Afr. trop.

SARCOCOLLA, Linn. Syst. ed. I (1735). *PENAEACEAE*, Benth. & Hook. f. iii. 202.
BRACHYSIPHON, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 24. t. 2.
acuta, *Kunth, in Linnaea*, v. (1830) 678.—Afr. austr.
formosa, *Kunth, l. c.*; *et A. Juss. in Ann. Sc. Nat.* Sér. III. vi. (1846) 25.—Afr. austr.

SARCOCOLLA:—
fucata, *Endl. Gen. Suppl.* iv. II. 74; *A. DC. in DC. Prod.* xiv. 489.—Afr. austr.
fucata, Zeyh. ex A. DC. l. c. = squamosa.
imbricata, *Benth. & Hook. f. Gen.* iii. 202.—Afr. austr.
Lessertiana, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 26 (= *Glischrocolla Lessertiana*).—Afr. austr.?
Linnaei, *A. Juss. l. c.* 25.—Afr. austr.
longiflora, *Kunth, in Linnaea*, v. (1830) 678; *A. DC. in DC. Prod.* xiv. 489.—Afr. austr.
minor, *A. DC. l. c.*—Afr. austr.
retzioides, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 104 = Endonema retzioides.
rupestris, *Eckl. & Zeyh. ex Sond. l. c.* 102.—Afr. austr.
speciosa, *Eckl. & Zeyh. ex Sond. l. c.* 103.—Afr. austr.
squamosa, *Kunth, in Linnaea*, v. (1830) 678.—Afr. austr.

SARCOCORDYLIS, Wall. Cat. n. 7249 (1832?) = **Balanophora**, Forst.

SARCOCYPHULA, Harv. Thes. Cap. ii. 58. t. 191 (1863) = **Cynanchum**, Linn. (Asclep.).
Gerrardi, Harv. l. c.—Afr. austr.

SARCODACTILIS, Gaertn. f. Fruct. iii. 39. t. 185 (1805) = **Citrus**, Linn. (Rutac.).
helicteroides, Gaertn. f. l. c. = C. decumana.

SARCODACTYLIS, Steud. Nom. ed. II. ii. 513 (1841) = praec.

SARCODES, Torr. in Smithson. Contrib. vi. (1854) (Pl. Frémont.) 17. t. 10. *MONOTROPEAE*, Benth. & Hook. f. ii. 606.
PTEROSPOROPSIS, Kellogg, fide A. Gray, ex Benth. & Hook. f. Gen. ii. 606 (1876).
sanguinea, *Torr. in Smithson. Contrib.* vi. (1854) (*Pl. Frémont.*) 18.—Calif.

SARCODISCUS, Griff. Notul. iv. 380 (1854) = **Kibara**, Endl. (Monim.).
chloranthiformis, Griff. l. c. = K. Blumei.

SARCODIUM, Pers. Syn. ii. 352 (1807) = seq.

SARCODUM, Lour. Fl. Cochinch. 462 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 498.
scandens, *Lour. l. c.*—Cochinch.

SARCOGLOSSUM, Beer, Prakt. Orch. 306 (1854) = **Cirrhaea**, Lindl. (Orchid.).
suaveolens, Beer, l. c. = C. tristis.

SARCOGLOTTIS, Presl, Rel. Haenk. i. 95. t. 15 (1827) = **Spiranthes**, Rich. (Orchid.).
cerina, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 634 = Spiranthes cerina.
diaphana, W. Baxt. l. c. = Spiranthes diaphana.
diuretica, W. Baxt. l. c. = Spiranthes diuretica.
grandiflora, Klotzsch, in Otto & Dietr. Allg. Gartenz. x. (1842) 107 = Spiranthes picta.
lobata, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 634 = Spiranthes lobata.
picta, Klotzsch, in Otto & Dietr. Allg. Gartenz. x. (1842) 106 = Spiranthes picta.
rosulata, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 634 = Spiranthes rosulata.
rufescens, Klotzsch, in Otto & Dietr. Allg. Gartenz. x. (1842) 107 = Spiranthes rufescens.
speciosa, Presl, Rel. Haenk. i. 96. t. 15 = Spiranthes speciosa, *Lindl.*

SARCOGONUM, G. Don, in Sweet, Hort. Brit. ed. III. 577 (1839) = **Muehlenbeckia**, Meissn. (Polygon.).
adpressum, G. Don, l. c. = M. adpressa.
complexum, Kunze, in Linnaea, xvii. (1843) 570 = M. complexa.
depressum, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1354 = M. australis.

SARCOLAENA, Thou. Hist. Vég. Afr. Austr. 37. tt. 9, 10 (1806). *CHLAENACEAE*, Benth. & Hook. f. i. 195.
SARCOCHLAENA, Spreng. Syst. ii. 594 (1825).
TANTALUS, Noronha, ex Thou. Hist. Vég. Afr. Austr. 39 (1806).

SARCOLAENA:—
Bojeriana, *Baill. Adansonia*, x. (1872) 177.—Madag.
eriophora, *Thou. Hist. Vég. Afr. Austr.* 40.—Madag.
grandiflora, *Thou. l. c.* t. 9.—Madag.
multiflora, *Thou. l. c.* t. 10.—Madag.

SARCOLEMMA, Griseb. ex Lorentz, Veget. Nord. Entre Rios, 80 (1878) = **Sarcostemma**, R. Br. (Asclep.).

SARCOLIPES, Eckl. & Zeyh. Enum. 290 (1837) = **Crassula**, Dill.
pubescens, Eckl. & Zeyh. l. c. = C. Sarcolipes.

SARCOLOBUS, R. Br. in Mem. Wern. Soc. i. (1809) 34. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 769.
Banksii, *Schult. Syst.* vi. 58.—Java.
carinatus, Griff. Notul. iv. 53 = globosus.
carinatus, *Wall. in As. Res.* xii. (1816) 570. t. 4.—Ind. or.; Burma.
globosus, *Wall. l. c.* 568. t. 5.—Ind. or.; Malaya.
narcoticus, *Miq. Fl. Ind. Bat.* ii. 502.—Java.
Spanoghei, *Miq. l. c.*—Java.
virulentus, Griff. Notul. iv. 54 = carinatus?

SARCOMERIS, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 340 = **Pachyanthus**, A. Rich. (Melastom.).
coriacea, Naud. l. c. = P. cubensis.

SARCOMORPHIS, Boj. ex Moq. in DC. Prod. xiii. II. 179 (1849) = **Salsola**, Linn. (Chenop.).
littoralis, Boj. ex Moq. l. c. 180 = Salsola littoralis.

SARCOMPHALIUM, Dulac, Fl. Hautes-Pyr. 113 (1867) = **Hyacinthus**, Tourn. (Liliac.).
amethystinum, Dulac, l. c. = H. amethystinus.

SARCOMPHALUS, P. Br. Hist. Jamaic. 179 (1756). *RHAMNEAE*, Benth. & Hook. f. i. 376.
acutifolius, *Griseb. Cat. Pl. Cub.* 32.—Cuba.
carolinianus, Rafin. Sylva Tellur. 29 = Rhamnus caroliniana.
divaricatus, *Griseb. Cat. Pl. Cub.* 31.—Cuba.
graecus, Rafin. Sylva Tellur. 29 = Rhamnus Sibthorpiana.
havanensis, *Griseb. Cat. Pl. Cub.* 31.—Ind. occ.
laurinus, Griseb. Fl. Brit. W. Ind. 100 = retusus.
levigatus, Rafin. Sylva Tellur. 29 = Ceanothus laevigatus.
mauritianus, Rafin. l. c. = Zizyphus Jujuba.
prunifolius, Rafin. l. c. = Rhamnus prunifolia.
retusus, *Rafin. l. c.*—Ind. occ.
Shortianus, Rafin. l. c. = Rhamnus lanceolata.

SARCOPERIS, Rafin. Fl. Tellur. ii. 16 (1836) = **Campelia**, Rich. (Commel.).
bibracteata, Rafin. l. c. = C. Zanonia.

SARCOPETALUM, F. Muell. Pl. Vict. i. 26, Suppl. t. 3 (1860). *MENISPERMACEAE*, Benth. & Hook. f. i. 37.
Harveyanum, *F. Muell. l. c.* 27; *Suppl.* t. 3.—Austral.

SARCOPHYLLUM, E. Mey. Comm. Pl. Afr. Austr. 32 (1835) = **Lebeckia**, Thunb. (Legumin.).
carnosum, E. Mey. l. c. = L. contaminata.
carnosum, Sims, Bot. Mag. t. 2502 = L. Simsiana.
carnosum, Willd. ex Walp. in Linnaea, xiii. (1839) 472 = Coelidium ciliare.
gnidioides, Willd. ex Walp. l. c. = Coelidium Vogelii.
grandiflorum, E. Mey. Comm. Pl. Afr. Austr. 32 = L. grandiflora?

SARCOPHYLLUM, Willd. Sp. Pl. iii. 968 (1800) = seq.

SARCOPHYLLUS, Thunb. in Schrad. Journ. i. (1799) 323 = **Aspalathus**, Linn. (Legumin.).
carnosus, Thunb. l. c. 324 = A. sarcodes.

SARCOPHYSA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 190 = **Juanulloa**, Ruiz & Pav. (Solan.).
speciosa, Miers, l. c. 191 = J. speciosa.

SARCOPHYTE, Sparrm. in Vet. Akad. Handl. Stockh. xxvii. (1776) 300. t. 7. *BALANOPHOREAE*, Benth. & Hook. f. iii. 234.
ICHTHYOSMA, Schlecht. in Linnaea, ii. (1827) 671. t. 8.
sanguinea, *Sparrm. in Vet. Akad. Handl. Stockh.* xxvii. (1776) 300. t. 7.—Afr. austr.

SARCOPODIUM, Lindl. in Lindl. & Paxt. Flow. Gard. i. (1850) 155 = **Bulbophyllum**, Thou. (Orchid.).
affine, Lindl. & Paxt. l. c., in obs. = B. affine.
amplum, Lindl. & Paxt. l. c. = Dendrobium amplum.
Cheiri, Lindl. & Paxt. l. c. = B. Cheiri.
Dearei, Hort.—Cf. Gard. Chron. (1883) II. 108. f. 17.—Hab.?
fuscescens, Lindl. Fol. Orchid. Sarc. 2 = Dendrobium fuscescens.
grandiflorum, Lindl. l. c. 3 = B. grandiflorum.
Griffithii, Lindl. l. c. 6 = B. Griffithii.
leopardinum, Lindl. & Paxt. Flow. Gard. i. (1850–51) 155, in obs. = B. leopardinum.
Lobbii, Lindl. & Paxt. l. c. = B. Lobbii.
macranthum, Lindl. & Paxt. l. c., in obs. = B. macranthum.
megalanthum, Lindl. Fol. Orchid. Sarc. 4 = B. megalanthum.
pileatum, Lindl. l. c. = B. pileatum.
psittacoglossum, Reichb. f. ex Hook. Bot. Mag. t. 5408 = B. psittacoglossum.
purpureum, Reichb. f. in Bonplandia, iv. (1856) 330.—Java.
Reinwardtii, Lindl. Fol. Orchid. Sarc. 4 = B. Reinwardtii.
rotundatum, Lindl. l. c. 2 (Dendrobii sp.).—Reg. Himal.
striatum, Lindl. l. c. 5 = B. striatum.
triadenium, Lindl. l. c. 2 = B. triadenium.
uniflorum, Lindl. l. c. 6 = B. uniflorum.

SARCOPOTERIUM, Spach, in Ann. Sc. Nat. Sér. III. v. (1846) 43 = **Poterium**, Linn. (Rosac.).
spinosum, Spach, l. c. = P. spinosum.

SARCOPTERYX, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 500, 658. *SAPINDACEAE*.
coriacea, *Radlk. l. c.* 659.—Ins. Waigiou.
Martyana, *Radlk. l. c.*—Austral.
melanophloea, *Radlk. l. c.*—N. Guin.
squamosa, Radlk. l. c. 659 = Cupania glabrata.
stipitata, Radlk. l. c. = Ratonia stipitata.

SARCOPYRAMIS, Wall. Tent. Fl. Nap. 32. t. 23 (1824). *MELASTOMACEAE*, Benth. & Hook. f. i. 754.
grandiflora, Griff. Notul. iv. 678 = napalensis.
javanensis, Triana, in Trans. Linn. Soc. xxviii. (1871) 77, sphalm. = seq.
javanica, Zoll. & Mor. Syst. Verz. Zoll. 12 = napalensis.
lanceolata, Wall. Cat. n. 6290 = napalensis.
napalensis, *Wall. Tent. Fl. Nep.* 32. t. 23.—Reg. Himal.; Malay.

SARCORHYNA, Presl, Bot. Bemerk. 103 (1844) = **Rostellaria**, Gaertn. f. (Acanth.).
Lessertiana, Presl, l. c. = R. Lessertiana.

SARCOSIPHON, Blume, Mus. Bot. Lugd. Bat. i. 65. f. 18 (1849) = **Bagnisia**, Becc. (Burmann.).
clandestinum, Blume. l. c.—Java.

SARCOSPERMA, Hook. f. in Benth. & Hook. f. Gen. ii. 655 (1876). *SAPOTACEAE*, Benth. & Hook. f. l. c.
PERONIA, R. Br. in Wall. Cat. n. 7543 (1832).
arboreum, *Hook. f. in Benth. & Hook. f. Gen.* ii. 655.—Reg. Himal.
Griffithii, *Hook. f. l. c.*—Reg. Himal.
laurinum, *Hook. f. l. c.* 649, in obs., 655.—China.

SARCOSTEMMA, R. Br. in Mem. Wern. Soc. i. (1809) 50. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 763.
MONOSTEMMA, Turcz. in Bull. Soc. Nat. Mosc. (1848) i. 255.
SARCOLEMMA, Griseb. ex Lorentz, Veget. Nord. Entre Rios, 80 (1878).
acidum, Voigt, Hort. Suburb. Calc. 542 = brevistigma.

SARCOSTEMMA:—
annulare, Roth, Nov. Pl. Sp. 178 = Holostemma Rheedianum.
aphyllum, Hochst. ex Decne. in DC. Prod. viii. 538 = viminale.
aphyllum, *Schult. Syst.* vi. 116.—Afr. austr.
apiculatum, *Decne. in DC. Prod.* viii. 539.—Bras.
arenarium, *Benth. Bot. Voy. Sulph.* 34.—Calif.
australe, *R. Br. Prod.* 463.—Austral.
barbatum, *Mart. ex Fourn. in Mart. Fl. Bras.* vi. IV. 235.—Bras.
bicolor, Decne. in DC. Prod. viii. 541 = Philibertia elegans.
bifidum, *Fourn. in Mart. Fl. Bras.* vi. IV. 235.—Bras.
bilobum, Hook. & Arn. Bot. Beech. Voy. 438 = Philibertia cynanchoides.
bonariense, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 296.—Reg. Argent.
brachystigma, Hook. Ic. Pl. t. 861, sphalm. = seq.
brevistigma, *Wight & Arn. in Wight, Contrib.* 59.—Ind. or.; Burma.
brevistylum, Royle, Illustr. Bot. Himal. 274, sphalm. = praec.
Brownii, G. F. W. Mey. Prim. Fl. Esseq. 139 = Philibertia viminalis.
Brunonianum, *Wight & Arn. in Wight, Contrib.* 59.—Ind. or.
campanulatum, *Lindl. Bot. Reg.* (1846) t. 36.—Peruv.?
clausum, Schult. Syst. vi. 114 = Philibertia viminalis.
crassifolium, Chapm. Fl. S. U. St. 368 = Philibertia viminalis.
crassifolium, Decne. in DC. Prod. viii. 540 = Philibertia crassifolia.
crispum, Benth. Pl. Hartw. 291 = Philibertia crispa.
cumanense, H. B. & K. Nov. Gen. et Sp. iii. 195 = Philibertia cumanensis.
cuspidatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 233.—Bras.
cynanchoides, Decne. in DC. Prod. viii. 540 = Philibertia cynanchoides.
Daltoni, *Decne. in Hook. Niger Fl.* 149. t. 14.—Ins. Cap. Virid.
Dombeyanum, *Decne. in DC. Prod.* viii. 539.—Peruv.
Donianum, Decne. l. c. 542 = Philibertia gracilis.
elegans, Decne. l. c. 541 = Philibertia elegans.
flavum, Decne. l. c. 540 = Philibertia flava?
Forskålianum, *Schult. Syst.* vi. 117.—Arabia.
Gardneri, Fourn. in Mart. Fl. Bras. vi. IV. 233.—Bras.
Gilliesii, Decne. in DC. Prod. viii. 542 = Philibertia solanoides.
glaucum, *H. B. & K. Nov. Gen. et Sp.* iii. 194. t. 229.—Am. austr.; Venezuela.
gracile, Decne. in DC. Prod. viii. 539 = Philibertia gracilis.
grandiflorum, Decne. l. c. 541 = Philibertia gracilis.
hastatum, *Decne. l. c.*—Peruv.
heterophyllum, Engelm. ex Torr. in Pacif. Rail. Rep. v. 362 = Philibertia linearis.
incanum, Decne. in DC. Prod. viii. 542 = Philibertia solanoides.
intermedium, *Decne. l. c.* 538.—Ind. or.
Jacquini, *Decne. l. c.* 542.—N. Granat.
Lindenianum, Decne. l. c. 541 = Philibertia Lindeniana.
lineare, Decne. in Benth. Pl. Hartw. 25 = Philibertia linearis.
lineare, *Spreng. Neue Entdeck.* iii. 35.—Venezuela.
luridum, Kunze, in Linnaea, xx. (1847) 26 = Philibertia lurida.
marsupiflorum, *Decne. in DC. Prod.* viii. 541.—Peruv.
mauritianum, Boj. Hort. Maurit. 214 = Decanema Bojerianum.
nudum, C. Sm. ex Decne. in Hook. Niger Fl. 149 = Daltoni.
Odontolepis, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 17.—Ins. Rodrig.
ovatum, E. Mey. Comm. Pl. Afr. Austr. 221 = Vincetoxicum Meyeri.
pallidum, *Fourn. in Mart. Fl. Bras.* vi. IV. 235.—Bras.
pannosum, Decne. in DC. Prod. viii. 540 = Philibertia pannosa.
pedunculatum, *Fourn. in Mart. Fl. Bras.* vi. IV. 235.—Bras.
pubescens, *H. B. & K. Nov. Gen. et Sp.* iii. 194.—Am. austr.

SARCOSTEMMA:—
pyrotechnicum, Schult. Syst. vi. 116 = Leptadenia Spartium.
quadriflorum, *Decne. in DC. Prod.* viii. 542.—Peruv.
riparium, Decne. l. c. 540.—Bras.
rotundifolium, Decne. l. c. = Philibertia rotundifolia.
Schottii, *Fourn. in Mart. Fl. Bras.* vi. IV. 234.—Bras.
solanoides, Decne. in DC. Prod. viii. 541 = Philibertia solanoides.
stellare, *Griseb. in Lorentz, Veget. Nord. Entre Rios*, 80, nomen.—Reg. Argent.
stipitaceum, *Schult. Syst.* vi. 116.—Arabia.
Stocksii, *Hook. f. Fl. Brit. Ind.* iv. 27.—Sind.
Swartzianum, Schult. Syst. vi. 115 = Philibertia viminalis.
tetrapterum, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) I. 255.—Afr. austr.
texanum, Engelm. ex Torr. Bot. Mex. Bound. 160 = Philibertia cynanchoides.
Thunbergii, G. Don, Gen. Syst. iv. 156 = aphyllum.
undulatum, Torr. Bot. Mex. Bound. 161 = Philibertia undulatum.
variifolium, *Decne. in DC. Prod.* viii. 541.—Peruv.
viminale, *R. Br. in Mem. Wern. Soc.* i. (1809) 51.—Afr. trop.
viminale, Wall. ex Decne. in DC. Prod. viii. 538 = brevistigma.
viminale, Wight & Arn. Contrib. 59 = intermedium.

SARCOSTIGMA, Wight & Arn. in Edinb. N. Phil. Journ. xiv. (1853) 299. *OLACINEAE*, Benth. & Hook. f. i. 354.
edule, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 298.—Ins. Andaman.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 241. t. 47.—Java.
Kleinii, *Wight & Arn. in Edinb. N. Phil. Journ.* xiv. (1833) 299.—Ind. or.; Malaya.
Roxburghii, *Wall. ex Griff. in Proc. Linn. Soc.* i. (1846) 282.—Ind. or.
Vogelii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. x. (1852) 117.—Afr. trop.
Wallichii, *Baill. Adansonia*, x. (1872) 282.—Burma.

SARCOSTOMA, Blume, Bijdr. 339. t. 45 (1825) = **Dendrobium**, Sw. (Orchid.).
javanica, Blume, Bijdr. 340. t. 45 = D. Sarcostoma.
juncea, Steud. Nom. ed. II. ii. 514 (err. typ.) = praec.

SARCOSTYLES, Presl, ex DC. Prod. iv. 15 (1830) = **Hydrangea**, Gronov. (Saxifr.).
peruviana, Presl, l. c. 16 = H. peruviana.

SARCOTHECA, Blume, Mus. Bot. Lugd. Bat. i. 241 (1850). *LINEAE*, Benth. & Hook. f. i. 245.
macrophylla, *Blume, l. c.* 242.—Malaya.

SARCOTHECA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 474 = **Schinus**, Linn. (Anacard.).
bahiensis, Turcz. l. c. = Sch. terebinthifolius.

SARCOTOECHIA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 501, 659 = **Ratonia**, DC. (Sapind.).
cuneata, Radlk. l. c. 659 = R. tenax.
protracta, Radlk. l. c. 660 = R. tenax.

SARCOZYGIUM, Bunge, in Linnaea, xvii. (1843) 7. t. 1. *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 266.
xanthoxylum, *Bunge, l. c.* 8.—Mongolia.

SARDINIA, Vell. Fl. Flum. 144 (1825); iii. t. 163 (1827) = **Guettarda**, Vent. (Rubiac.).
bella, Vell. l. c.—Bras.

SARIAVA, Reinw. in Syll. Ratisb. ii. (1828) 12 = **Symplocos**, Linn. (Styrac.).

SARIBUS, Blume, Rumphia, ii. 48. t. 95, 96 (1836) = **Livistona**, R. Br. (Palm.).
chinensis, Blume, l. c. 49, in obs. = L. chinensis.
cochinchinensis, Blume, l. c., in obs. = L. cochinchinensis
olivaeformis, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 176 = L. olivaeformis.
rotundifolius, Blume, Rumphia, ii. 49. t. 95, 96 = L. rotundifolia.
subglobosus, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 177 = L. subglobosa.

SARISSUS, Gaertn. Fruct. i. 118. t. 25 (1788) = **Hydrophylax**, Linn. f. (Rubiac.).
anceps, Gaertn. l. c. = H. maritima.

SARMENTARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 140 = **Adelobotrys**, DC. (Melastom.).
decora, Naud. l. c. 141 = A. adscendens.

SARMIENTA, Ruiz & Pav. Prod. 4 (1794). *GESNERACEAE*, Benth. & Hook f. ii. 1012.
URCEOLARIA, Molina, Sagg. Chile, ed. II. 136, 277 (1810).
repens, *Ruiz & Pav. Fl. Per.* i. 8. t. 7.—Chili.
scandens, Pers. Syn. i. 13 = repens.

SARNA, Karst. in Linnaea, xxviii. (1856) 414 = **Apodanthes**, Poit. (Cytinac.).
Caulotreti, Karst. l. c. 415 = A. Blanchetii?
Ingae, Karst. l. c. = A. Calliandrae?

SAROSANTHERA, Korth. Verh. Nat. Gesch. Bot. 103. t. 16 (1840?) = **Adinandra**, Jack (Ternstr.).
excelsa, Korth. l. c. 104.—Borneo.
lasiopetala, Thw. Enum. Pl. Zeyl. 41 = A. lasiopetala.

SAROTES, Lindl. Swan River App. 19 (1839) = **Guichenotia**, J. Gay (Stercul.).
latifolia, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 150 = Lasiopetalum rosmarinifolium.
ledifolia, Lindl. Swan River App. 19 = G. Sarotes.
micrantha, Steetz, in Lehm. Pl. Preiss. ii. 346 = G. micrantha.
pogonanthera, F. Muell. Fragm. ii. 7 = G. micrantha.
rosmarinifolia, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 149 = Lasiopetalum rosmarinifolium.
semihastata, F. Muell. Fragm. ii. 4 = G. semihastata.

SAROTHAMNOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.

SAROTHAMNUS, Wimm. Fl. Schles. (1832) 278 = **Cytisus**, Linn. (Legumin.).
affinis, Boiss. Voy. Bot. Espagne, ii. 134. t. 40 = C. affinis, *Boiss.*
arboreus, Boiss. l. c. 137 = C. baeticus.
arboreus, Webb, Iter Hisp. 52, in obs. = C. arboreus.
baeticus, Webb, l. c. = C. baeticus.
Balansae, Boiss. Diagn. Ser. II. ii. 7 = C. Balansae.
Bourgaei, Boiss. l. c. 6.—Hispan.
cantabricus, Bourg. ex Nym. Consp. 150 = C. eriocarpus.
cantabricus, Willk. in Flora, xxxiv. (1851) 616.—Hispan.
Carlieri, Comp. ex Nym. Consp. 150 = *catalaunicus*.
catalaunicus, Webb, in Ann. Sc. Nat. Sér. III. ix. (1848) 63.—Ins. Canar.
commutatus, Willk. in Oestr. Bot. Zeitschr. xxxi. (1881) 105.—Hispan.
eriocarpus, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 10 = C. eriocarpus.
gaditanus, Boiss. & Reut. l. c. = C. baeticus.
grandiflorus, Webb, Otia Hisp. 45. t. 39 = C. grandiflorus.
malacitanus, Boiss. Voy. Bot. Espagne, ii. 137 = C. malacitanus.
oxyphyllus, Boiss. Diagn. Ser. II. ii. 7.—Hispan.
parviflorus, Willk. & Cutand. in Linnaea, xxx. (1859–60) 95 = C. albus.
patens, Plan. Fl. Gallega, 164 = *Welwitschii*.
patens, Webb, Iter Hisp. 51 = C. patens.
purgans, Gren. & Godr. Fl. Fr. i. 349 = C. purgans.
scoparius, Koch, Syn. Fl. Germ. ed. 1. 152 = C. scoparius.
virgatus, Webb, Iter Hisp. 52 = C. grandiflorus.
vulgaris, Wimm. Fl. Schles. (1832) 278 = C. scoparius.
Welwitschii, Boiss. & Reut. Pugill. Pl. Nov. (1852) 28.—Hispan.

SAROTHECA, Nees, in Mart. Fl. Bras. ix. 113. t. 18 (1847) = **Justicia**, Linn. (Acanth.).
elegans, Nees, l. c. = J. elegans, *Pohl* (cf. Addend.).
rostrata, Nees, in DC. Prod. xi. 383 = J. rostrata.
salviaeflora, Nees, l. c. = J. salviaeflora.
scabra, Nees, in Mart. Fl. Bras. ix. 114 = Aphelandra pectinata?

SAROTHRA, Linn. Diss. Chen. 14 (1751); Amoen. Acad. iii. 11 (1787) = **Hypericum**, Tourn.
blentinensis, Savi f. in Nuov. Giorn. Pisan. xxxix. (1839) 225 = H. mutilum.
canadensis, Rafin. in Am. Monthly Mag. (1818) 267 = H. canadense.
cuneifolia, Rafin. Atl. Journ. 17 = H. Sarothra?
Drummondii, Grev. & Hook. in Hook. Bot. Misc. iii. (1833) 236. t. 107 = H. Drummondii.
fastigiata, Rafin. Fl. Tellur. iii. 79 = H. adpressum.
gentianoides, Linn. Sp. Pl. 272 = H. Sarothra.
gentianoides, Lour. Fl. Cochinch. 182.—Cochinchina.
hypericoides, Nutt. Gen. Am. i. 204 = H. Sarothra.
Loureiriana, Schult. Syst. vi. 679 = *gentianoides*, Lour.
parviflora, Rafin. in Am. Monthly Mag. (1819) 192 = H. mutilum.
pauciflora, Rafin. Fl. Tellur. iii. 79 = H. canadense.
quinquenervia, Rafin. in Am. Monthly Mag. (1818) 267 = H. mutilum.
trinervia, Rafin. Fl. Tellur. iii. 79 = H. triplinerve?

SAROTHROSTACHYS, Klotzsch, in Wiegm. Archiv, vii. (1841) 185 = **Sebastiania**, Spreng. (Euphorb.).
daphniphylla, Buek, Index ad DC. Prod. iv. 366 = Seb. daphniphylla.
Laureola, Buek, l. c. = Seb. Laureola.
Luschnathiana, Klotzsch, in Wiegm. Archiv, vii. (1841) 185 = Seb. multiramea.
multiramea, Klotzsch, l. c. = Seb. multiramea.

SARRACENA, Tourn. ex Linn. Hort. Cliff. 497 (1737) = seq.

SARRACENIA. Linn. Sp. Pl. ed. I. 510 (1753). *SARRACENIACEAE*, Benth. & Hook. f. i. 48.
SARRAZINIA, Hoffmgg. Verz. Pfl. Nachtr. i. 223 (1824).
adunca, Sm. Exot. Bot. i. 103 = variolaris.
atropurpurea, Hort. Bull. in Gard. Chron. (1880) i. 759 = flava.
atrosanguinea, Hort. Bull, ex André, in Illustr. Hortic. xxvii. (1880) 87. t. 386 = praec.
calceolata, Nutt. in Trans. Am. Phil. Soc. N. S. iv. (1834) 49. t. 1 = psittacina.
Catesbaei, Ell. Sketch, ii. 11 = flava.
Chelsoni ×, Hort. Veitch. in Gard. Chron. (1880) i. 722. f. 125.
Courtii ×, Hort.—Cf. Gard. Chron. (1881) ii. 381.
crispata ×, Hort. Bull, ex André, in Illustr. Hortic. xxvii. (1880) 87. t. 387.
Drummondii, *Croom, in Ann. Lyc. N. York*, iv. (1848) 100.—Am. bor.
Drummondii, Lindl. & Paxt. Flow. Gard. i. (1850–51) 1. t. 1 = undulata.
erythropus ×, Hort. ?—Cf. Gard. Chron. (1882) i. 713.—Hab.?
flava, *Linn. Sp. Pl.* 510.—Am. bor.
formosa ×, Hort. Veitch.—Cf. Gard. Chron. (1881) ii. 281.
Gronovii, Wood, Class-Book (1861), 222; Am. Bot. & Flor. 30 = flava, rubra.
heterophylla, Eaton, Man. Bot. ed. IV. 447 = purpurea.
heterophylla, Ell. ex Steud. Nom. ed. II. ii. 514 = flava.
lacunosa, Bartr. Trav. ed. II. (1794) p. xii. 415 = variolaris?
leucophylla, Rafin. Fl. Ludov. 14 = Drummondii?
melanorhoda ×, Hort. Veitch, in Gard. Chron. (1882) ii. 281.—Hab.?
minor, Sweet, Brit. Flow. Gard. Ser. II. t. 138 = Sweetii.
minor, Wall. Fl. Carol. 153 = variolaris.
Mooreana ×.—Cf. Gard. Chron. (1877) i. 425.
Moorei ×, Mast. in Gard. Chron. (1881) ii. 40.
nutans, D. Dietr. Syn. Pl. iii. 260 = Heliamphora nutans.
Popei ×, Hort.—Cf. Gard. Chron. (1881) i. 691.
psittacina, *Michx. Fl. Bor. Am.* i. 311 (psyttacina).—Am. bor.
pulchella, Croom, in Am. Journ. Sc. Ser. I. xxv. (1834) 75 = psittacina.
purpurea, *Linn. Sp. Pl.* 510.—Am. bor.
rubra, Planch. in Fl. des Serres, Sér. I. x. (1854–55) 243. t. 1074 = Sweetii.
rubra, *Walt. Fl. Carol.* 152.—Am. bor.
Rugelii, Shuttl. ex A. DC. in DC. Prod. xvii. 6 = flava.
Stevensi ×, Hort. ex Mast. in Gard. Chron. (1881) ii. 40. f. 7.

SARRACENIA:—
Sweetii, *A. DC. in DC. Prod.* xvii. 5.—Am. bor.
undulata, *Decne. in Rev. Hortic.* Sér. IV. i. (1852) 126.—Am. bor.
variolaris, *Michx. Fl. Bor. Am.* i. 310.—Am. bor.
Williamsii ×, Hort.—Cf. Gard. Chron. (1874) ii. 49; (1881) i. 629.

SARRACHA, Reichb. in Moessl, Handb. ed. II. 1. p. xxxix (1827) = **Saracha**, Ruiz & Pav. (Solan.).

SARRATIA, Moq. in DC. Prod. xiii. ii. 233, 268 (1849) = **Amaranthus**, Linn.
Berlandieri, Moq. l. c. 268 = A. polygonoides.
polygonoides, Moq. l. c. 270 = A. polygonoides.
scariosa, Moq. l. c. 269 = A. scariosus.
urceolata, Moq. l. c. = A. urceolatus.

SARRAZINIA, Hoffmgg. Verz. Pfl. Nachtr. i. 223 (1824) = **Sarracenia**, Linn.

SARRIZINIA, Steud. Nom. ed. II. ii. 514 (1841), sphalm. = praec.

SARTORIA, Boiss. Diagn. Ser. I. ix. 109 (1849) = **Onobrychis**, Linn. (Legumin.).
hedysaroides, Boiss. & Heldr. l. c.—Taurus.

SARTWELLIA, A. Gray, Pl. Wright. i. (in Smithson. Contrib. iii.) 122. t. 6 (1852). *COMPOSITAE*, Benth. & Hook. f. ii. 407.
Flaveriae, *A. Gray, in Pl. Wright.* i. 122.—Texas.
mexicana, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1882) 107, nomen; xix. (1884) 34.—Mexic.

SASALI, Adans. Fam. ii. 305 (1763) = **Grewia**, Linn. (Tiliac.).

SASANQUA, Nees, in Liter. Flora, iv. (1834) 144 = **Camellia**, Linn. (Ternstr.).
malliflora, Rafin. Sylva Tellur. 140 = C. Sasanqua.
ochroleuca, Rafin. l. c. = Gordonia anomala.
odorata, Rafin. l. c.—Hab.?
oleifera, Rafin. l. c. = C. Sasanqua.
vulgaris, Nees, in Liter. Flora, iv. (1834) 144 = C. Sasanqua.

SASSA, Bruce, ex J. F. Gmel. Syst. 1038 (1791) = **Acacia**, Willd. (Legumin.).
gummifera, J. F. Gmel. l. c. = A. gummifera.

SASSAFRAS, Linn. Syst. ed. I (1735); Nees, Laur. Expos. 17 (1833). *LAURINEAE*, Benth. & Hook. f. iii. 160.
albidum, Nees, Syst. Laurin. 490 = officinale.
Goesianum, Teijsm. & Binn. Cat. Hort. Bog. 94 = Massoia aromatica.
Loureiri, Kostel. Allg. Med.-Pharm. Fl. ii. 481 = Cinnamomum Parthenoxylon.
mauritianum, Boj. Hort. Maurit. 273 = Hernandia peltata.
officinale, *Nees & Eberm. Handb. Med.-Pharm. Bot.* ii. 418; *Nees, Syst. Laurin.* 488.—Am. bor.
officinale, Siebold, in Verh. Batav. Gen. xii. (1830) 23 = Lindera triloba.
officinarum, J. S. Presl, Rostl. ii. 68 = officinale.
Parthenoxylon, Nees, Syst. Laurin. 491 = Cinnamomum Parthenoxylon.
rubrum, Rafin. Sylva Tellur. 134 = officinale?
Thunbergii, Siebold, in Verh. Batav. Gen. xii. (1830) 23 = Lindera umbellata.

SASSAFRIDIUM, Meissn. in DC. Prod. xv. i. 171 (1864). *LAURINEAE*, Benth. & Hook. f. iii. 160.
veraguense, *Meissn. l. c.*—Mexic.

SASSEA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 128.—Cf. in Abh. Akad. Berl. 1854 (1855), 132. t. 12 C = **Begonia**, Linn.
columnaris, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 128 = B. columnaris.
glabra, Klotzsch, l. c. = B. columnaris.
Hoffmanniana, Klotzsch, ex A. DC. in DC. Prod. xv. i. 275 = B. Urticae.
Urticae, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 128 = B. Urticae.

SASSIA, Molina, Sagg. Chile, 145 (1782) = **Oxalis**, Linn. (Geran.).
perdicaria, Molina, l. c. 146 = O. lobata.
tinctoria, Molina, l. c. = O. violacea.

SATANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 = **Flacourtia**, Comm. (Bixin.).

SATANOCRATER, Schweinf. in Verh. Zool.-Bot. Ges. Wien, xviii. (1868) 676. *ACANTHACEAE*, Benth. & Hook. f. ii. 1085.
fellatensis, *Schweinf. l. c.*—Afr. trop.

SATARIA, Rafin. New Fl. Am. iv. 20 (1836) = **Peucedanum**, Tourn. (Umbell.).
linearis, Rafin. l. c. 21 = P. triternatum, *Nutt.*

SATIRIUM, Neck. Delic. Gallo-Belg. ii. 372 (1768) = **Satyrium**, Linn. (Orchid.).

SATTADIA, Fourn. in Mart. Fl. Bras. vi. iv. 231 (1885). *ASCLEPIADEAE*.
Burchellii, *Fourn. l. c.* 232.—Bras.

SATUREGIA, Leers, Fl. Herborn. 135 (1775) = seq.

SATUREIA, Linn. Coroll. Gen. 9. n. 961 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1187.
FURERA, Adans. Fam. ii. 193 (1763).
NOSTELIS, Rafin. Sylva Tellur. 76 (1838).
PILOBLEPHIS, Rafin. New Fl. Am. iii. 52 (1836).
SACCOCALYX, Coss. & Dur. in Ann. Sc. Nat. Sér. III. xx. (1853) 80. t. 5.
SATURIASTRUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137.
Acinos, Scheele, in Flora, xxvi. (1843) 577 = Calamintha Acinos.
alpina, Scheele, l. c. = Calamintha alpina.
alternipilosa, *C. Koch, in Linnaea*, xxi. (1848) 668.—Reg. Caucas.
americana, Poir. Encyc. vi. 571 = Hyptis scoparia.
angustifolia, Presl, Fl. Sic. p. xxxvi = Micromeria graeca.
approximata, Bivona, Stirp. Rar. Sicil. Manip. iv. 13 = Micromeria approximata.
approximata, *Frivald. in Flora*, xiv. (1836) 438.—Graec.
argaea, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 33 = Thymus Serpyllum.
atropatana, *Bunge, Lab. Pers.* 36.—Persia.
bachtiarica, *Bunge, l. c.* 37.—Persia.
Boissieri, *Haussk. ex Boiss. Fl. Orient.* iv. 565.—As. Min.
brachiata, Stokes, Bot. Mat. Med. iii. 300 = hortensis.
brevis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 44 = montana.
Calamintha, Scheele, in Flora, xxvi. (1843) 577 = Calamintha officinalis.
Campanella, *Ehrenb. Naturg. Reise*, 13.—Afr. bor.
canescens, Guss. Pl. Rar. 228. t. 42 = Micromeria canescens.
capitata, Desf. Fl. Atlant. ii. 9 = Micromeria nervosa.
capitata, Linn. Sp. Pl. 568 = Thymus capitatus.
ciliata, Avé-Lall. in Ind. Sem. Hort. Petrop. xi. Suppl. 66 = montana.
Clinopodium, Caruel, in Parl. Fl. Ital. vi. 135 = Calamintha Clinopodium.
coccinea, Bertol. Misc. Bot. viii. 23 = Calamintha coccinea.
coerulea, *Janka, in Kanitz, Pl. Roman.* 94.—Banat.
collina, Salisb. Prod. 77 = S. Thymbra.
condaea, Juss. ex Poir. Encyc. vi. 571 = Hyptis scoparia.
congesta, Hornem. Hort. Hafn. 541 = Micromeria graeca.
consentina, Tenore, Prod. Fl. Nap. App. v. 17 = Micromeria graeca.
cordata, Moris, ex Bertol. Fl. Ital. x. 519 = Micromeria cordata.
corsica, Caruel, in Parl. Fl. Ital. vi. 137 = Calamintha corsica.
cristata, Nym. Syll. 102 = Micromeria cristata.
cuneata, Reichb. Ic. Fl. Germ. xviii. t. 1272, sphalm. = seq.
cuneifolia, *Tenore, Prod. Fl. Nap.* 33.—Reg. Mediterr.
dalmatica, Nym. Syll. 102 = Micromeria dalmatica.

SATUREIA:—
diffusa, Benth. ex Boiss. Fl. Orient. iv. 566 = alternipilosa.
ericoides, Willd. ex Benth. in Linnaea, xi. (1837) 328 = Gardoquia microphylla.
fasciculata, Rafin. Prec. Découv. 39 = Micromeria approximata.
fasciculata, Tausch, in Syll. Ratisb. ii. (1828) 248 = cuneifolia.
Fenzlii, Nym. Syll. Suppl. 20 = Micromeria Pulegium.
filicaulis, Schott, ex Boiss. Fl. Orient. iv. 562 = hortensis.
filiformis, Desf. Fl. Atlant. ii. 8 = Micromeria microphylla.
filiformis, Nym. Syll. 102 = Micromeria filiformis.
filiformis, Tenore, Marin. Viagg. i. 299 = Micromeria nervosa.
flexuosa, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 45 = montana.
glandulosa, Caruel, in Parl. Fl. Ital. vi. 125 = Calamintha glandulosa.
globifera, Steud. Nom. ed. I. 733, sphalm. = seq.
globulifera, *Desf. Tabl.* ed. I. 58, nomen.—Hab. ?
glomerata, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 613 = inodora.
graeca, Linn. Sp. Pl. 568 = Micromeria graeca.
grandiflora, Scheele, in Flora, xxvi. (1843) 577 = Calamintha grandiflora.
graveolens, Caruel, in Parl. Fl. Ital. vi. 143 = Calamintha graveolens.
hirsuta, J. & C. Presl, Delic. Prag. 79 = Micromeria Juliana.
hirta, Host, Fl. Austr. ii. 134 = cuneifolia.
hispida, *Ehrh. Beitr.* vii. 147.—Graecia ?
hispida, Nym. Syll. 102 = Micromeria hispida.
hortensis, *Linn. Sp. Pl.* 568.—Reg. Medit.; Oriens.
Hostii, Caruel, in Parl. Fl. Ital. vi. 125 = Calamintha thymifolia.
hyssopifolia, Bertol. in Ann. Stor. Nat. (1829) iii. 406 = montana.
hyssopifolia, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 428 = obovata.
illyrica, Host, Fl. Austr. iii. 133 = montana.
incana, Spreng. Syst. ii. 719 = Micromeria obovata.
inodora, Host, Fl. Austr. ii. 135 = Micromeria parviflora.
inodora, *Salzm. ex Benth. Lab. Gen. et Sp.* 354.—Reg. Medit. occ.
intermedia, Benth. in DC. Prod. xii. 210 = mutica.
intermedia, *C. A. Mey. Verz. Pfl. Cauc.* 91.—Reg. Caucas.
intricata, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 96.—Hispan.
Juliana, Linn. Sp. Pl. 567 = Micromeria Juliana.
Juliana, Pall. Tabl. Taur. 53 = montana.
Kitaibelii, Wierzb. ex Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 177 = montana.
lanata, C. Sm. ex Link, in Buch, Beschr. Canar. Ins. 143 = Micromeria lanata.
laxiflora, C. Koch, in Linnaea, xxi. (1848) 668 = hortensis.
Libanotis, [*Spreng.*] *Bot. Gart. Halle*, 84 (Quid ?).—Syria.
longiflora, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 566.—Persia.
longiflora, Presl, Fl. Sicul. p. xxxvi = Micromeria graeca.
macrantha, *C. A. Mey. in Ind. Sem. Hort. Petrop.* xi. *Suppl.* 67.—Reg. Caucas.
marifolia, Caruel, in Parl. Fl. Ital. vi. 125 = Micromeria marifolia.
Mastichina, Linn. Sp. Pl. 567 = Thymus Mastichina.
micrantha, Hoffmgg. & Link, Fl. Port. i. 142 = Micromeria graeca.
microphylla, Guss. Fl. Sic. Prod. ii. 120 = Micromeria microphylla.
mollis, *Webb, ex Steud. Nom.* ed. II. ii. 515, nomen.—Ins. Teneriff.
montana, *Linn. Sp. Pl.* 568.—Europ. austr.; Reg. Caucas.
mucronifolia, Stokes, Bot. Mat. Med. iii. 300 = montana.
mutica, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 49.—Reg. Caucas.
nemorosa, *Tardent, Ess. Hist. Nat. Bessarab.* 54, nomen (Quid ?).—Bessarab.
Nepeta, Scheele, in Flora, xxvi. (1843) 577 = Calamintha Nepeta.

SATUREIA:—
nervosa, Desf. Fl. Atlant. ii. 9. t. 121. f. 2 = Micromeria nervosa.
obovata, *Lag. Gen. et Sp. Nov.* 18.—Hispan.
officinarum, Crantz, Inst. i. 526 = hortensis.
Olla, *Ehrenb. Naturg. Reise*, 13.—Afr. bor.
origanoides, Linn. Sp. Pl. 568 = Cunila Mariana.
ovata, R. Br. in Salt, Abyss. App. 64 = Micromeria ovata.
pachyphylla, C. Koch, in Linnaea, xvii. (1843) 295 = hortensis.
parnassica, Heldr. & Sart. ex Boiss. Fl. Orient. iv. 563 = approximata.
parviflora, Presl, Fl. Sic. p. xxxvi = Micromeria Juliana.
parviflora, Vis. in Flora, xii. (1829) i. Erg. 13 = Micromeria parviflora.
petraea, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 45 = montana.
Piperella, Bertol. Fl. Ital. vi. 50 = Micromeria croatica, M. Piperella.
plumosa, Hampe, in Flora, xxv. (1842) i. Beibl. 127 = Micromeria nervosa.
provincialis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 45 = montana.
punctata, R. Br. in Salt, Abyss. App. 64 = Micromeria punctata.
pygmaea, Sieber, ex Vis. Stirp. Dalm. Spec. 11 = montana.
pyrenaica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 89 = montana.
rigida, *Bartr. ex Benth. Lab. Gen. et Sp.* 354.—Florida.
rigidula, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 46 = montana.
rupestris, Wulf. in Jacq. Coll. ii. 130 = Micromeria rupestris.
scabra, *Presl, Fl. Sic.* p. xxxvi.—Sicilia.
sessiliflora, Presl, l. c. = Micromeria graeca.
sicula, Guss. Fl. Sic. Syn. ii. 89 = Micromeria graeca.
Sideritis, Benth. in DC. Prod. xii. 212, sphalm. = Micromeria obovata.
Sophia, Hort. ex Steud. Nom. ed. I. 733 = Micromeria Juliana.
spicata, Crantz, Inst. i. 525 = Micromeria Juliana.
spicata, Vis. Fl. Dalm. ii. 195 = cuneifolia.
spicigera, Boiss. Fl. Orient. iv. 566 = alternipilosa.
spinosa, Heldr. ex Nym. Consp. 591 = approximata.
spinosa, *Linn. Cent. Pl.* ii. 19; *Amoen. Acad.* iv. 317.—Ins. Creta.
stricta, [Soland. in] Russ. Aleppo, ed. II. ii. 255 = Nepeta heliotropifolia.
subdentata, *Boiss. Fl. Orient.* iv. 565.—Reg. Caucas.
subspicata, Bartl. ex Vis. Stirp. Dalm. Rar. ii. t. 4 = montana.
Teneriffae, Willd. Enum. Hort. Berol. Suppl. 40 = Micromeria microphylla.
tenuifolia, Botteri, ex Scheele, in Linnaea, xxii. (1849) 593 = Micromeria longifolia.
tenuifolia, Tenore, Prod. Fl. Nap. 33 = Micromeria graeca.
tenuis, Link, in Buch, Beschr. Canar. Ins. 143 = Micromeria tenuis.
Thymbra, *Linn. Sp. Pl.* 567.—Reg. Mediterr. or.
thymifolia, Scop. Fl. Carn. ed. II. i. 428 = Micromeria rupestris.
thymoides, Nym. Syll. 102 = Micromeria thymoides.
thymoides, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 19 = Micromeria varia.
Tragoriganum, Tausch, in Syll. Ratisb. ii. (1828) 248 = S. Thymbra.
trifida, Moench, Meth. 386 = montana.
varia, Webb & Berth. Phyt. Canar. iii. 71 = Micromeria varia.
variegata, Host, Fl. Austr. iii. 134 = montana.
viminea, Burm. f. Fl. Ind. 126 = hortensis.
viminea, Linn. Syst. ed. X. 1096 = Micromeria obovata.
virgata, Stokes, Bot. Mat. Med. iii. 299 = Micromeria Juliana.
virgata, Vis. ex Benth. Lab. Gen. et Sp. 353 = cuneifolia.
virginiana, Linn. Sp. Pl. 567 = Pycnanthemum lanceolatum, linifolium.
virginica, Hill, Veg. Syst. xviii. 51 = praec.

SATUREJA, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = praec.

SATURIASTRUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 133 = **Satureia**, Linn. (Labiat.).
montanum, Fourr. l. c. = Satureia montana.
petraeum, Fourr. l. c. = Satureia montana.

SATURNA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 (Euphorb. ?).

SATURNIA, Maratt. Pl. Romul. et Saturn. 18. t. 2 (1772) = **Allium**, Tourn. (Liliac.).
cernua, Maratt. Fl. Rom. i. 258 = A. Chamaemoly.
Chamaemoly, Salisb. Gen. Pl. Fragm. 92 = A. Chamaemoly.
etrusca, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 59 = A. Chamaemoly.
littoralis, Jord. & Fourr. Ic. Fl. Eur. ii. 21 = A. Chamaemoly.
rubrinervis, Jord. & Fourr. l. c. 22 = A. Chamaemoly.
viridula, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 60 = A. Chamaemoly.

SATYRIA, Klotzsch, in Linnaea, xxiv. (1851) 21. *VACCINIACEAE*, Benth. & Hook. f. ii. 567.
Riedelia, Meissn. in Mart. Fl. Bras. vii. 171 (1863).
clonantha, *Klotzsch, in Linnaea*, xxiv. (1851) 22.—Am. centr.
panurensis, *Benth. & Hook. f. Gen.* ii. 568.—Bras.
Warszewiczii, *Klotzsch, in Linnaea*, xxiv. (1851) 22.—Am. centr.

SATYRIDIUM, Lindl. Gen. et Sp. Orch. 345 (1838) = **Satyrium**, Linn. (Orchid.).
rostratum, Lindl. l. c. = Satyrium rhynchanthum.

SATYRIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 270 (1737), partim; Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 214. *ORCHIDEAE*, Benth. & Hook. f. iii. 629.
Aviceps, Lindl. Gen. et Sp. Orch. 345 (1838).
Diplecthrum, Pers. Syn. ii. 508 (1807).
Satyridium, Lindl. Gen. et Sp. Orch. 345 (1838).
aculeatum, Linn. f. Suppl. 402 = Eulophia aculeata.
acuminatum, *Lindl. Gen. et Sp. Orch.* 339.—Afr. austr.
adnatum, Sw. Prod. Veg. Ind. Occ. 118 = Pelexia adnata.
albidum, Linn. Sp. Pl. 944 = Habenaria albida.
albiflorum, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 76 = nepalense.
alpinum, Pers. Syn. ii. 507 = Herminium alpinum.
alpinum, F. W. Schmidt, Fl. Boëm. i. 63 = Habenaria viridis.
amoenum, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 29.—Ins. Borbon.
anthropomorpha, Pers. Syn. ii. 507 = Aceras anthropophora.
anthropophora, Pers. l. c. = Aceras anthropophora.
Atherstonei, *Reichb. f. in Flora*, lxiv. (1881) 328.—Afr. austr.
atlanticum, Lindl. Gen. et Sp. Orch. 300 = Habenaria intacta.
aureum, Paxt. Mag. Bot. xv. (1849) 31 = coriifolium.
barbatum, Thunb. Prod. Pl. Cap. 5 = Disa barbata.
bicallosum, *Thunb. l. c.* 6.—Afr. austr.
bicorne, *Thunb. l. c.*—Afr. austr.
bifidum, Thunb. l. c. 5 = Disa bifida.
bifolium, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 273.—Abyss.
bifolium, Wahlenb. Fl. Suec. 579 = Habenaria bifolia.
brachypetalum, *A. Rich. Tent. Fl. Abyss.* ii. 299.—Abyss.
bracteale, Salisb. in Trans. Hort. Soc. i. (1812) 290 = Habenaria viridis.
bracteatum, Lindl. Gen. et Sp. Orch. 342 = Lindleyanum.
bracteatum, Pers. Syn. ii. 507 = Habenaria viridis.
bracteatum, *Thunb. Prod. Pl. Cap.* 6.—Afr. austr.
calceatum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 520.—Madag.
candidum, *Lindl. Bot. Reg.* (1838) *Misc.* 82.—Afr. austr.
capense, Houtt. Handleid. xii. 502 = Eulophia tristis.
capense, Linn. Pl. Rar. Afr. 27; Amoen. Acad. vi. 110 = Eulophia aculeata.
carneum, *R. Br. in Ait. Hort. Kew.* ed. II. v. 196.—Afr. austr.
cassideum, Lindl. Gen. et Sp. Orch. 341 = parviflorum.
cernuum, Thunb. Prod. Pl. Cap. 5 = Disa physodes.

SATYRIUM:—
chrysostachyum, Herschel, in Bot. Reg. (1838) Misc. 84 = coriifolium.
ciliatum, Lindl. Gen. et Sp. Orch. 341 = nepalense.
conopseum, Wahlenb. Fl. Suec. 579 = Habenaria conopsea, *Benth.*
cordifolium, *Lindl. Gen. et Sp. Orch.* 344.—Afr. austr.
coriifolium, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 216.—Afr. austr.
coriophoroides, *A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 274.—Abyss.
cornutum, *Burm. f. Fl. Cap. Prod.* 30 (err. typ. 26).—Afr. austr.
cornutum, Thunb. Prod. Pl. Cap. 5 = Disa cornuta.
cristatum, *Sond. in Linnaea*, xix. (1847) 84.—Afr. austr.
cucullatum, Lodd. Bot. Cab. t. 104 = coriifolium.
cucullatum, Sw. in Vet. Acad. Handl. Stockh. xxi. (1800) 216 = bicorne.
cylindricum, Thunb. Prod. Pl. Cap. 5 = Disa cylindrica.
debile, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 67.—Afr. austr.
densiflorum, Brot. Fl. Lusit. i. 22 = Habenaria intacta.
densiflorum, Lindl. Gen. et Sp. Orch. 340 = parviflorum.
diphyllum, Link, in Schrad. Journ. ii. (1799) 323 = Habenaria cordata.
draconis, Thunb. Prod. Pl. Cap. 5 = Disa draconis.
elatum, Sw. Prod. Veg. Ind. Occ. 119 = Spiranthes elata.
emarcidum, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 67.—Afr. austr.
Epipogium, Linn. Sp. Pl. 945 = Epipogum aphyllum.
erectum, Lindl. Gen. et Sp. Orch. 340 = coriifolium.
erectum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 216.—Afr. austr.
eriostomum, *Lindl. Gen. et Sp. Orch.* 342.—Afr. austr.
excelsum, Thunb. Prod. Pl. Cap. 5 = Disa excelsa.
ferrugineum, F. W. Schmidt, Fl. Boëm. i. 62 = Habenaria viridis.
ferrugineum, Thunb. Prod. Pl. Cap. 5 = Disa ferruginea.
flexuosum, Thou. Orch. Isles Austr. Afr. t. 7 (= *Peristylus flexuosus*).—Ins. Mascar.
flexuosum, Thunb. Prod. Pl. Cap. 5 = Disa flexuosa.
foliosum, Heyne, ex Wall. Cat. n. 7054 (= *Coeloglossum secundum*).—Ind. or.
foliosum, Lindl. Gen. et Sp. Orch. 336 = Hallackii.
foliosum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 216.—Afr. austr.
fuscum, Huds. Fl. Angl. ed. I. 337 = Habenaria viridis.
giganteum, Linn. f. Suppl. 402 (= *Cyrtopera gigantea*).—Afr. austr.
gracile, *Lindl. Gen. et Sp. Orch.* 338.—Madag.
gramineum, Thou. Orch. Isles Austr. Afr. t. 6 = Habenaria graminea.
grandiflorum, Thunb. Prod. Pl. Cap. 4 = Disa uniflora.
Hallackii, *Bolus, in Journ. Linn. Soc.* xx. (1884) 476.—Afr. austr.
Herscheliae, *Harv. Gen. S. Afr. Pl.* 326, nomen.—Afr. austr.
hians, Linn. f. Suppl. 401 = Disa lacera ?
hircinum, Linn. Sp. Pl. 944 = Orchis hircina.
hirsutum, Gilib. Exercit. ii. 484 = Goodyera repens.
hirtellum, *Sw. Prod. Veg. Ind. Occ.* 118.—Jamaic.
humile, *Lindl. Gen. et Sp. Orch.* 339.—Afr. austr.
Ivantalae, *Reichb. f. in Flora*, xlviii. (1865) 183.—Afr. trop.
lanceum, Pers. Syn. ii. 507 = Ophrys lancea.
latifolium, Thou. Orch. Isles Austr. Afr. t. 10 = Habenaria chlorantha.
leucocomos, *Reichb. f. in Flora*, xlviii. (1865) 182.—Afr. trop.
ligulatum, *Lindl. Gen. et Sp. Orch.* 342.—Afr. austr.
Lindleyanum, *Bolus, in Journ. Linn. Soc.* xx. (1884) 474.—Afr. austr.
lineatum, Lindl. Gen. et Sp. Orch. 343 = bracteatum.
lingulatum, Vill. Hist. Pl. Dauph. ii. 42 (err. typ. 24) = Habenaria viridis.
longicauda, *Lindl. Gen. et Sp. Orch.* 337.—Afr. austr.
longicolle, Lindl. l. c. 335 = maculatum.
lupulinum, *Lindl. l. c.* 338.—Afr. austr.
lydenburgense, Reichb. f. in Flora, lxiv. (1881) 328 = eriostomum.

SATYRIUM:—
macrophyllum, *Lindl. Gen. et Sp. Orch.* 338.—Afr. austr.
macrostachyum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 298 = coriophoroides.
maculatum, *Burch. ex Lindl. Gen. et Sp. Orch.* 337.—Afr. austr.
maculatum, Desf. Fl. Atlant. ii. 319 = Habenaria intacta.
marginatum, *Bolus, in Journ. Linn. Soc.* xx. (1884) 476.—Afr. austr.
membranaceum, *Sw in Vet. Acad. Handl. Stockh.* xxi. (1800) 216.—Afr. austr.
militare, Lindl. Gen. et Sp. Orch. 342 = sphaerocarpum.
Monorchis, Pers. Syn. ii. 507 = Herminium Monorchis.
muticum, *Lindl. Gen. et Sp. Orch.* 344.—Afr. austr.
nepalense, *D. Don, Prod. Fl. Nep.* 26.—Ind. or.
nigrum, Linn. Sp. Pl. 944 = Habenaria nigra.
obsoletum, Pers. Syn. ii. 507 = Habenaria obsoleta.
ochroleucum, *Bolus, in Journ. Linn. Soc.* xxii. (1885) 66.—Afr. austr.
odoratissimum, Wahlenb. Fl. Suec. 578 = Habenaria odoratissima.
odorum, *Sond. in Linnaea*, xix. (1847) 86.—Afr. austr.
orchioides, Sw. Prod. Veg. Ind. Occ. 118 = Spiranthes orchioides.
orientale, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 27 (Quid ?).—Java.
orobanchoides, Linn. f. Suppl. 402 = Corycium orobanchoides.
pallidum, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 77 = nepalense.
paludosum, *Reichb. f. in Flora*, xlviii. (1865) 182.—Afr. trop.
papillosum, Lindl. Gen. et Sp. Orch. 341 = erectum.
parviflorum, Lindl. l. c. 336 = marginatum.
parviflorum, Pers. Syn. ii. 506 = Habenaria intacta.
parviflorum, *Sw. in Vet. Acad. Handl. Stockh.* xxi. (1800) 216.—Afr. austr.
pedicellatum, Linn. f. Suppl. 402 = Eulophia aculeata.
Perrottetianum, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 76 = nepalense.
pictum, Lindl. Gen. et Sp. Orch. 344 = bracteatum.
plantagineum, Linn. Syst. ed. X. 1244 = Spiranthes orchioides.
praealtum, Thou. Orch. Isles Austr. Afr. t. 11 = Habenaria praealta.
pumilum, *Thunb. Prod. Pl. Cap.* 6.—Afr. austr.
pustulatum, Lindl. Bot. Reg. (1840) t. 18 = erectum.
pygmaeum, *Sond. in Linnaea*, xix. (1847) 86.—Afr. austr.
repens, Linn. Sp. Pl. 945 = Goodyera repens.
repens, Michx. Fl. Bor. Am. ii. 157 = Goodyera pubescens.
retusum, *Lindl. Gen. et Sp. Orch.* 343.—Afr. austr.
rhynchanthum, *Bolus, in Journ. Linn. Soc.* xix. (1882) 342.—Afr. austr.
riparium, *Reichb. f. in Flora*, xlviii. (1865) 183.—Afr. trop.
rosellatum, Thou. Orch. Isles Austr. Afr. t. 8 = Habenaria mascarenensis.
rostratum, *Lindl. Gen. et Sp. Orch.* 338.—Madag.
rufescens, Thunb. Prod. Pl. Cap. 5 = Disa rufescens.
sagittale, Thunb. l. c. = Disa sagittalis.
saxicolum, *Bolus, in Journ. Linn. Soc.* xx. (1884) 474.—Afr. austr.
scanense, Linn. ex Steud. Nom. ed. I. 734 = Habenaria albida.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 300.—Abyss.
secundum, Thunb. Prod. Pl. Cap. 4 = Disa racemosa.
spathulatum, Thunb. l. c. 5 = Disa spathulata.
sphaerocarpum, *Lindl. Gen. et Sp. Orch.* 337.—Afr. austr.
spirale, Sw. Prod. Veg. Ind. Occ. 118 = Spiranthes tortilis.
spirale, Thou. Orch. Isles Austr. Afr. t. 9 = Habenaria spiralis, *A. Rich.*
stenopetalum, *Lindl. Gen. et Sp. Orch.* 336.—Afr. austr.
striatum, Ker-Gawl. in Brand. Journ. Sc. viii. (1820) 221 = bracteatum.
striatum, *Thunb. Prod. Pl. Cap.* 6.—Afr. austr.
tabulare, Linn. f. Suppl. 402 = Cymbidium tabulare.
tenellum, Thunb. Prod. Pl. Cap. 5 = Disa tenella.
tortum, Thunb. l. c. = Disa torta.

SATYRIUM:—
trifidum, Vill. Hist. Pl. Dauph. ii. 42 = Habenaria albida.
trinerve, *Lindl. Gen. et Sp. Orch.* 344.—Madag.
triste, Linn. f. Suppl. 402 = Eulophia tristis.
utriculatum, Sond. in Linnaea, xix. (1847) 84 = candidum.
vallisoletanum, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 7.—Mexic.
venosum, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Hab. ?
virescens, Pers. Syn. ii. 507 = Habenaria herbiola.
viride, Linn. Sp. Pl. 944 = Habenaria viridis.
Welwitschii, *Reichb. f. in Flora*, xlviii. (1865) 182.—Afr. trop.
Wightianum, Lindl. Gen. et Sp. Orch. 340 = nepalense.

SAUBINETIA, Remy, in C. Gay, Fl. Chil. iv. 282. t. 49 (1848) = **Verbesina**, Linn. (Compos.).
helianthoides, Remy, l. c. 284.—Chili.

SAUERIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 122; et in Abh. Akad. Berl. 1854 (1855), 40. t. 2 A = **Begonia**, Linn.
crenata, Hassk. in Versl. Kon. Akad. Wetensch. iv. (1855) 139 = B. crenata.
sulcata, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 122; et in Abh. Akad. Berl. 1854 (1855), 41 = B. sulcata.

SAUL, Roxb. ex Wight & Arn. Prod. 84 (1834) = **Shorea**, Linn. (Dipteroc.).
Iallarea, Roxb. ex Wight & Arn. l. c. = Shorea Talura.
Tumbagia, Steud. Nom. ed. II. ii. 516 = seq.
Tumbugaia, Roxb. ex Wight & Arn. l. c. = Shorea Tumbugaia.

SAULCYA, Michon, Voy. Relig. Or. ii. 383 (1854) = **Odontospermum**, Neck (Compos.).
hierochuntica, Michon, l. c. (= *Asteriscus pygmaeus*).—Afr. bor.; Palaest.

SAUNDERSIA, Reichb. f. in Bot. Congr. Lond. 120 (1866); Beitr. Orch. 17. t. 6 (1869). *ORCHIDEAE*, Benth. & Hook. f. iii. 561.
mirabilis, *Reichb. f. ll. cc.* 120; 19.—Bras.

SAURAVIA, Spreng. Anleit. ii. II. 818 (1818) = **Saurauja**, Willd. (Ternstr.).

SAUROBROMA, Rafin. Sylva Tellur. 32 (1838) = **Celtis**, Tourn. (Urtic.).
iguanense, Rafin. l. c. = C. aculeata.

SAURAUJA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 407. t. 4. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 184.
APATELIA, DC. in Mém. Soc. Phys. Genèv. i. (1821) 426.
BLUMIA, Spreng. Syst. iii. 126 (1826).
DRAYTONIA, A. Gray, Bot. U. St. Expl. Exped. i. 206. t. 15 (1854).
LEUCOTHEA, Moç. & Sesse, ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 419.
MARUMIA, Reinw. ex Blume, Cat. Gew. Buitenz. 79 (1823).
OBELANTHERA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 148.
OVERSTRATIA, Deschamps, ex R. Br. in Benn. Pl. Jav. Rar. 171 (1840).
PALAUA, Ruiz & Pav. Prod. 100. t. 22 (1794), nomen prius.
REINWARDTIA, Blume, ex Nees, in Syll. Ratisb. i. (1824) 96.
SAURAVIA, Spreng. Anleit. ii. II. 818 (1818).
SCAPHA, Noronha, in Verh. Batav. Gen. v. (1770) ed. I. Art. IV. 3.
TONSHIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 225 (1825).
VANALPHIMIA, Lechen. ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 419.
aesculifolia, *De Vriese, Pl. Ind. Bat. Or.* 42.—Java.
altissima, *Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 108.—N. Guin.
Andreana, *Oliver, ex F. Muell. Census, Suppl.* ii. 3, nomen.—Austral.
angustifolia, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 50 = lanceolata.

SAURAUJA :—
angustifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 242.—Mexic.
anisopoda, *Turcz. l. c.*—Mexic.
armata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 59.—Burma.
aspera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 242.—Mexic.
barbigera, Hook. Ic. Pl. t. 331 = leucocarpa.
biserrata, *Spreng. Syst.* iv. *Cur. Post.* 211.—Peruv.
Blumeana, Spreng. l. c. 210 = bracteosa.
Blumia, *Steud. Nom.* ed. II. ii. 516.—Java.
Blumiana, *Benn. Pl. Jav. Rar.* 174.—Java.
brachybotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 245.—N. Granat.
bracteolata, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 422.—Java.
bracteosa, *DC. l. c.* 423. t. 6.—Java.
brevirostris, *Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 106.—N. Guin.
callithrix, *Miq. l. c.* 111.—Ins. Celebes.
camptodonta, Miq. Fl. Ind. Bat. Suppl. 482 = tristyla.
cauliflora, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 425.—Java.
ceramica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 111.—Ins. Ceram.
cerea, *Griff. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 288.—Reg. Himal.
costata, *Reinw. ex De Vriese, Pl. Ind. Bat. Or.* 56.—Ins. Celebes.
crenulata, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 424.—Java.
crenulata, Wight, ex Wall. Cat. n. 1459 = Gordonia obtusa.
Cumingiana, De Vriese, Pl. Ind. Bat. Or. 38 = latebractea.
cuneata, Blume, Bijdr. 127 = bracteolata.
cuspidella, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 108.—Sumatra.
dasyantha, *De Vriese, Pl. Ind. Bat. Or.* 49.—Sumatra.
dicalyx, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 112.—Ins. Ceram.
distasosa, *Korth. Verh. Nat. Gesch. Bot.* 131.—Java.
euryolepis, *De Vriese, Pl. Ind. Bat. Or.* 48.—Ins. Celebes.
exasperata, *De Vriese, l. c.* 56.—Ins. Philipp.
excavata, *Korth. Verh. Nat. Gesch. Bot.* 130.—Borneo.
excelsa, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 407.—N. Granat.
fasciculata, *Wall. Pl. As. Rar.* ii. 40.—Reg. Himal.
ferox, *Korth. Verh. Nat. Gesch. Bot.* 132.—Borneo.
ferruginea, *Griff. Itin. Notes*, 120 (sp. dub.).—Reg. Himal.
floccifera, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 267.—N. Granat.
gigantea, Blume, Bijdr. 129 = bracteosa.
gigantea, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 424.—Java.
glabrata, *Steud. Nom.* ed. II. ii. 516; *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 119.—Peruv.
grandifolia, *Zoll. & Mor. Syst. Verz. Zoll.* 26.—Java.
Griffithii, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 286.—Assam.
Hasskarliana, *Miq. Fl. Ind. Bat.* i. II. 484.—Java.
hirsuta, *Blume, Bijdr.* 128.—Java.
horrida, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 162.—Borneo.
inflexidens, Miq. Fl. Ind. Bat. Suppl. 481 = tristyla.
Jackiana, Korth. Verh. Nat. Gesch. Bot. 127 = cauliflora.
Junghuhnii, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 148.—Java.
Kegeliana, *Schlecht. in Bot. Zeit.* xi. (1853) 694.—Guatemala.
Korthalsii, Zoll. & Mor. Syst. Verz. Zoll. 26 = leprosa.
laevigata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 267.—N. Granat.
lanceolata, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 421. t. 4.—Java.
latibractea, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 148.—Ins. Philipp.
latipetala, *Hemsl. Diagn. Pl. Nov.* 4.—Mexic.
lepidicalyx, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 109.—Ins. Celebes.

SAURAUJA :—
leprosa, *Korth. Verh. Nat. Gesch. Bot.* 131.—Java.
leucocarpa, *Schlecht. in Linnaea*, x. (1836) 249.—Mexic.
leucophloia, Korth. Verh. Nat. Gesch. Bot. 125 = tristyla.
Lindeni, *Regel, Ind. Sem. Hort. Petrop.* (1868) 90.—Hab.?
longifolia, *Oliver, in Journ. Linn. Soc.* xv. (1876) 98.—Ins. Celebes.
macrophylla, *Hort. ex Regel, Ind. Sem. Hort. Petrop.* (1866) 102.—Hab.?
macrophylla, Linden, in Lindl. & Paxt. Flow. Gard. ii. (1851–52) 27 = villosa?
macrotricha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 60.—Burma; China.
media, Korth. Verh. Nat. Gesch. Bot. 125 = tristyla.
micrantha, *Blume, Bijdr.* 127.—Java.
micrantha, Griff. Itin. Notes, 71 = Roxburghii?
microphylla, *De Vriese, Pl. Ind. Bat. Or.* 49.—Java.
mollis, *Hassk. Cat. Hort. Bog. Alt.* 210.—Java.
monadelpha, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 8.—N. Guin.
montana, Seem. Bot. Voy. Herald, 87. t. 16 = veraguensis.
Moritziana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 243.—Venezuela.
muricata, *Reinw. ex De Vriese, Pl. Ind. Bat. Or.* 46.—Ins. Celebes.
muricata, *Steud. Nom.* ed. II. ii. 516.—Java.
napaulensis, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 421.—Reg. Himal.
nigrescens, *Korth. Verh. Nat. Gesch. Bot.* 128.—Borneo.
Noronhiana, Blume, Bijdr. 127 = nudiflora.
novo guineensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 7.—N. Guin.
nudiflora, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 422. t. 5.—Java.
Obelanthera, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 245 = villosa.
oligolepis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 109.—Ins. Celebes.
oreophila, *Hemsl. Diagn. Pl. Nov.* 3.—Mexic.
oxyphylla, Miq. Fl. Ind. Bat. Suppl. 480 = tristyla.
paniculata, Wall. ex G. Don, Gen. Syst. i. 567 = napaulensis.
parviflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 268.—N. Granat.
pauciserrata, *Hemsl. Diagn. Pl. Nov.* 3.—Mexic.
pedunculares, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 267.—N. Granat.
pedunculata, Hook. Ic. Pl. t. 341–342.—Mexic.
pendula, *Blume, Bijdr.* 127.—Java.
Planchoni, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 161.—Malaya.
polyantha, De Vriese, Pl. Ind. Bat. Or. 36 = vulcani.
polyodon, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 107.—Molucc. *vel* Celebes?
polytricha, Miq. Fl. Ind. Bat. Suppl. 480 = setigera.
punduana, *Wall. Pl. As. Rar.* ii. 40.—Reg. Himal.
pycnotricha, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 244.—Venezuela.
Reinwardtiana, *Blume, Bijdr.* 128.—Java.
rosea, *Jungh. in Hoev. & De Vriese, Tijdschr.* viii. (1841) 384.—Java.
Roxburghii, *Wall. Pl. As. Rar.* ii. 40.—Reg. Himal.; Burma.
rubicunda, *Seem. Fl. Vit.* 14.—Ins. Fiji.
rubiformis, *Vatke, in Linnaea*, xl. (1876) 221.—Costa Rica.
rugosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 245.—Ins. Philipp.
Ruiziana, *Steud. Nom.* ed. II. ii. 516.—Peruv.
sakoembangensis, *Korth. Verh. Nat. Gesch. Bot.* 134.—Borneo.
sarapiquensis, *Carr. in Rev. Hortic.* (1877) 60.—Costa Rica.
scabra, *Poepp. ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 118.—Peruv.
scabrida, *Hemsl. Diagn. Pl. Nov.* 3.—Mexic.
serrata, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 420. t. 3; *Prod.* i. 526.—Mexic.
setidens, Miq. Fl. Ind. Bat. Suppl. 481 = vulcani.
setigera, *Korth. Verh. Nat. Gesch. Bot.* 133.—Sumatra.
singalangensis, *Korth. l. c.* 134.—Sumatra.
spadicea, Blume, Bijdr. 128 = lanceolata.

SAURAUJA :—
spectabilis, *Hook. Bot. Mag.* t. 3982.—Am. austr.
squamellicaula, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 112.—Ins. Molucc.
sterculifolia, Griff. Itin. Notes, 198 = napaulensis?
strigillosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 266.—N. Granat.
subcordata, *Korth. Verh. Nat. Gesch. Bot.* 124.—Borneo.
tewensis, *Korth. l. c.* 133.—Borneo.
Teysmanniana, Miq. Fl. Ind. Bat. Suppl. 481 = tristyla.
tomentosa, *Spreng. Syst.* iv. *Cur. Post.* 211.—N. Granat.; Ecuador.
tristyla, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 423. t. 7.—As. trop. or.
ursina, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 265.—N. Granat.
veraguensis, *Seem. Bot. Voy. Herald*, 249.—Panama.
villosa, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 420. t. 2; *Prod.* i. 525.—Mexic.
vulcani, *Korth. Verh. Nat. Gesch. Bot.* 128.—Sumatra.
xanthotricha, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 243.—Venezuela.

SAUROGLOSSUM, Lindl. Bot. Reg. t. 1618 (1833) = **Spiranthes**, Rich. (Orchid.).
elatum, Lindl. l. c.—Bras.
monophyllum, Griseb. Cat. Pl. Cub. 269 = Cranichis monophylla.
tenue, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 334.—Cuba.

SAUROMATUM, Schott, in Schott & Endl. Melet. i. 17 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 966.
abyssinicum, *Schott, Syn. Aroid.* 25.—Abyss.
asperum, C. Koch, in Belg. Hortic. xv. (1865) 101 = Dracontium asperum.
ferox, Linden, ex Engl. in DC. Monog. Phan. ii. 305 = Anchomanes dubius.
guttatum, *Schott, in Schott & Endl. Melet.* i. 17.—Ind. bor. occ.
Horsfieldii, *Miq. Fl. Ind. Bat.* iii. 196.—Java.
nubicum, *Schott, Syn. Aroid.* 25.—Afr. trop.
pedatum, Schott, in Schott & Endl. Melet. i. 17 = guttatum.
pulchrum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 221.—Sumatra.
punctatum, C. Koch, in C. Koch & Fint. Wochenschr. i. (1858) 263 = guttatum.
sessiliflorum, Kunth, Enum. Pl. iii. 28 = guttatum.
simlense, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 349 = guttatum.
venosum, Kunth, Enum. Pl. iii. 28; Schott, Prod. Aroid. 71 = guttatum.

SAUROPUS, Blume, Bijdr. 595 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 271.
CERATOGYNUM, Wight, Ic. v. II. 26. t. 1900 (1852).
albicans, *Blume, Bijdr.* 596.—Ind. or.; Malaya.
assimilis, *Thw. Enum. Pl. Zeyl.* 284.—Zeylan.
brevipes, *Muell. Arg. in Linnaea*, xxxii. (1863) 73.—Burma.
Ceratogynum, Baill. Etud. Gén. Euph. 635 = quadrangularis.
compressus, *Muell. Arg. in DC. Prod.* xv. II. 243.—Reg. Himal.; Burma.
Gardnerianus, Wight, Ic. vi. 6. t. 1951 = albicans.
indicus, Wight, Ic. t. 1952 = albicans.
macranthus, *Hassk. Retzia*, i. 166.—Java.
macranthus, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 187 = albicans.
quadrangularis, *Muell. Arg. in Linnaea*, xxxii. (1863) 73.—Ind. or.
repandus, *Muell. Arg. in Flora*, lv. (1872) 2.—Reg. Himal.
retroversus, *Wight, Ic.* vi. 6. t. 1951.—Zeylan.
rhamnoides, *Blume, Bijdr.* 596.—Java.
rigidus, *Thw. Enum. Pl. Zeyl.* 284.—Zeylan.
rostratus, *Miq. Fl. Ind. Bat. Suppl.* 447.—Sumatra.
spectabilis, *Miq. l. c.* 446.—Sumatra.
sumatranus, *Miq. l. c.*—Sumatra.
trinervius, *Hook. f. & Thoms. ex Muell. Arg. in Linnaea*, xxxii. (1863) 72.—Reg. Himal.
zeylanicus, Wight, Ic. t. 1952 = albicans.

SAURUROPSIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 589 = **Saururus**, Plum. (Piperac.).
chinensis, Turcz. l. c. 590 = Saururus Loureiri.
Cumingii, C. DC. in DC. Prod. xvi. I. 239 = Saururus Loureiri.

SAURURUS, Plum. ex Linn. Gen. ed. I. 108 (1737). *PIPERACEAE*, Benth. & Hook. f. iii. 127.
Mattuschkia, J. F. Gmel. Syst. ii. 589 (1791).
Saururopsis, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 589.
Spathium, Lour. Fl. Cochinch. 217 (1790).
cernuus, *Linn. Sp. Pl.* 341.—Am. bor.
cernuus, Thunb. Fl. Jap. 154 = Loureiri.
chinensis, *Hort. ex Loud. Hort. Brit.* 144, nomen; *et Encyc. Pl.* 298.—China.
Loureiri, *Decne. in Ann. Sc. Nat.* Sér. III. iii. (1845) 102.—China; Japon.
lucidus, Donn, Hort. Cantab. ed. III. 66 = cernuus.
natans, Linn. Mant. ii. 227 = Aponogeton monostachyum.
sinensis, Teijsm. & Binn. Cat. Hort. Bog. 77 = chinensis.

SAUSSUREA, DC. in Ann. Mus. Par. xvi. (1810) 156, 198. tt. 10, 13. *COMPOSITAE*, Benth. & Hook. f. ii. 471.
Aplotaxis, DC. in Guill. Archiv. Bot. ii. (1833) 330.
Aucklandia, Falc. in Trans. Linn. Soc. xix. (1842) 23.
Bennettia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 440 (1821).
Cyathidium, Lindl. in Royle, Illustr. Bot. Himal. 251. t. 56 (1835).
Eriocoryne, Wall. ex DC. Prod. vi. 514 (1837).
Eriostemon, Less. Syn. Comp. 12 (1832).
Frolovia, Ledeb. ex DC. Prod. vi. 538 (1837).
Hemistepta, Bunge, in Dorp. Jahrb. Litt. i. (1833) 221.
Heterotrichum, Bieb. Fl. Taur. Cauc. iii. 551 (1819).
Lagurostemon, Cass. in Dict. Sc. Nat. liii. 466 (1828).
Poecilotriche, Dulac, Fl. Hautes-Pyr. 528 (1867).
Theodorea, Cass. in Dict. Sc. Nat. xlvii. (1827) 500, 513; liii. 463 (1828).
acanthocarpa, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 378 = Jurinea ceratocarpa.
acaulis, Klatt, in Sitzb. Akad. Muench. (1878) 91 = Thomsoni.
acuminata, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 37 = alpina.
affinis, *Spreng. ex DC. Prod.* vi. 540.—As. trop.
alaschanica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 492.—As. centr.
alata, DC. in Ann. Mus. Par. xvi. (1810) 202 = japonica.
Alberti, *Regel & Winkler, in Act. Hort. Petrop.* vi. (1879) 298.—Turkestan.
albescens, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 233.—Reg. Himal.
alpina, *DC. in Ann. Mus. Par.* xvi. (1810) 198.—Reg. bor. et arct.
amara, *DC. l. c.* 200.—Sibir.
amara, Less. in Linnaea, ix. (1834) 169 = glomerata.
americana, *Eaton, in Coult. Bot. Gaz.* vi. (1881) 283.—Am. bor. occ.
amoena, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 447 (*amaena*) = salicifolia.
amurensis, Turcz. ex DC. Prod. vi. 534 = eriolepis.
Andersoni, *C. B. Clarke, Comp. Ind.* 226.—Reg. Himal.
andryaloides, *Hook. f. & Thoms. ex C. B. Clarke, l. c.* 227.—Reg. Himal.
angustifolia, DC. in Ann. Mus. Par. xvi. (1810) 200 = alpina.
Apus, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 490.—As. centr.
arenaria, *Maxim. l. c.*—China.
asperifolia, Sch. Bip. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 12 = elongata.
Atkinsoni, *C. B. Clarke, Comp. Ind.* 226.—Reg. Himal.
atriplicifolia, Bieb. ex Bess. in Flora, xvii. (1834) i. Beibl. 15 = serrata.
blanda, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 354.—Sibir. altaic.
borealis, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 38 = alpina.
bracteata, *Decne. in Jacquem. Voy. Bot.* 94. t. 102.—Reg. Himal.
brahuica, Boiss. Fl. Orient. iii. 566 = candicans.

SAUSSUREA:—
Bungei, Benth. & Hook. f. ex Franch. & Sav. Enum. Pl. Jap. i. 255 = affinis.
caespitosa, Wall. Cat. n. 2913 = taraxacifolia.
cana, *Ledeb. Ic. Pl. Fl. Ross.* i. 18. t. 75.—Sibir. altaic.
candicans, *C. B. Clarke, Comp. Ind.* 232.—Reg. Himal.
Candolleana, C. B. Clarke, l. c. 230 = Clarkei.
Candolleana, *Wall. Cat.* n. 3289.—Reg. Himal.; Soongar.
carthamoides, Buch.-Ham. ex DC. Prod. vi. 540 = affinis.
centauroides, *Tausch, in Flora*, xi. (1828) ii. 484.—Sibir. or.
ceratocarpa, Decne. in Jacquem. Voy. Bot. 93. t. 101 = Jurinea ceratocarpa.
chenopodifolia, Klatt, in Sitzb. Akad. Muench. (1878) 92 = albescens.
chinensis, Sch. Bip. in Flora, xxxv. (1852) 48 = japonica.
Clarkei, *Hook. f. Fl. Brit. Ind.* iii. 372.—Reg. Himal.
compacta, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 23 = alpina.
congesta, *Turcz. ex DC. Prod.* vi. 535.—Dahuria.
conica, C. B. Clarke, Comp. Ind. 224 = uniflora.
controversa, DC. in Ann. Mus. Par. xvi. (1810) 199 = discolor.
coronata, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 107.—Soongaria.
crassifolia, *DC. in Ann. Mus. Par.* xvi. (1810) 201.—Sibir.
crepidifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) ii. 47.—As. bor.
daurica, *Adams, in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 251.—Dahuria.
Davidi, *Franch. Pl. Davidian.* i. (1884) 180.—China.
deltoidea, *C. B. Clarke, Comp. Ind.* 235.—Reg. Himal.
denticulata, *Ledeb. Ic. Pl. Fl. Ross.* i. 18. t. 74.—Sibir.
denticulata, *Wall. Cat.* n. 2917.—Reg. Himal.
depressa, Gren. in Mém. Acad. Besanç. (1849); ex Gren. & Godr. Fl. Fr. ii. 272 = alpina.
depressa, Heuff. ex Nym. Consp. 415 = discolor.
discolor, *DC. in Ann. Mus. Par.* xvi. (1810) 199.—Europ.; As. bor.
dissecta, Ledeb. Fl. Alt. iv. 19, in nota = japonica.
duiensis, *F. Schmidt, Reisen Amurl.* 152.—Ins. Sachalin.
elata, *Ledeb. Ic. Pl. Fl. Ross.* i. 20. t. 80.—Sibir. altaic.
elegans, *Ledeb. l. c.* 19. t. 77.—Sibir. altaic.
elliptica, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 372.—Reg. Himal.
elongata, *DC. in Ann. Mus. Par.* xvi. (1810) 201.—Sibir.
epilobioides, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 495.—China.
eriolepis, *Bunge, ex DC. Prod.* vi. 535.—China; Manchuria.
eriostemon, *Wall. Cat.* n. 2912.—Reg. Himal.
esthonica, *Baer, in Rupr. Stirp. Fl. Petrop.* 21 (*cf. Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* vii. (1885) 5).—Ross.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 369.—Reg. Himal.
filifolia, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 312.—As. centr.
flavescens, Spreng. Syst. iii. 380 = Serratula flavescens.
foliosa, *Hook. f. Fl. Brit. Ind.* iii. 373.—Reg. Himal.
foliosa, *Ledeb. Ic. Pl. Fl. Ross.* i. 17. t. 69.—Sibir. altaic.
Frolowii, *Ledeb. l. c.* iv. 16. t. 352.—Sibir. altaic.
fruticulosa, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 448 = cana.
Gebleriana, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* vii. (1855) 6.—Sibir. altaic.
glacialis, *Herd. in Bull. Soc. Nat. Mosc.* xi. (1867) ii. 144.—Turkestan.
glanduligera, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 371.—Reg. Himal.
glauca, Ledeb. ex Bess. in Flora, xvii. (1834) i. Beibl. 15 = Laxmanni.
glomerata, *Poir. Encyl. Suppl.* v. 71.—Sibir.
Gmelini, G. Don, in Sweet, Hort. Brit. ed. III. 392 = Serratula Gmelini.
Gmelini, Hort. Dorp. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 48 = amara.

SAUSSUREA:—
gossipiphora, *D. Don, in Mem. Wern. Soc.* iii. (1821) 414.—Reg. Himal.
gossypifera, Spreng. Syst. iii. 381 = praec.
gossypina, Wall. Pl. As. Rar. ii. 32. t. 138 = praec.
gossypiphora, Wall. Cat. n. 2910 B = sacra.
gracilis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 518.—Japon.
graminifolia, *Wall. Cat.* n. 2911.—Reg. Himal.
grandifolia, *Maxim. Prim. Fl. Amur.* 169.—Reg. Amur.
Griffithii, *Boiss. Diagn.* Ser. II. iii. 47.—Afghan.
hieracioides, *Hook. f. Fl. Brit. Ind.* iii. 371.—Reg. Himal.
Hookeri, *C. B. Clarke, Comp. Ind.* 230.—Reg. Himal.
hypoleuca, *Spreng. ex DC. Prod.* vi. 541.—Reg. Himal.
integerrima, Spreng. l. c. 536 = graminifolia.
intermedia, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 193 = japonica.
iodostegia, *Hance, in Journ. Bot.* xvi. (1878) 109.—China.
Jacea, *C. B. Clake, Comp. Ind.* 233.—Reg. Himal.
japonica, *DC. in Ann. Mus. Par.* xvi. (1810) 203.—Japon.
Karelinii, *Stschégl. in Bull. Soc. Nat. Mosc.* xxi. (1848) ii. 244.—As. centr.
kaschgarica, *Rupr. Sert. Tiansch.* 54.—Turkestan.
Katschaete, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 491.—China.
Kunthiana, *C. B. Clarke, Comp. Ind.* 225.—Reg. Himal.
laciniata, *Ledeb. Ic. Pl. Fl. Ross.* i. 16. t. 64.—Sibir. altaic.
laevigata, Tausch, ex Steud. Nom. ed. II. ii. 517 = glomerata.
Lappa, *C. B. Clarke, Comp. Ind.* 233.—Reg. Himal.
latifolia, *Ledeb. Ic. Pl. Fl. Ross.* i. 17. t. 70.—Sibir. altaic.
Laxmanni, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 15.—Sibir. baical.
Ledebouri, Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 41 = alpina.
leucantha, Jord. in F. Schultz, Arch. Fl. 2 = alpina.
leucophylla, Schrenk, in Bull. Soc. Acad. Pétersb. x. (1842) 354 = pygmaea.
liatroides, Fisch. ex DC. Prod. vi. 532 = pycnocephala.
linearis, Champ. & Benth. in Hook. Kew Journ. iv. (1852) 235 = japonica.
lyrata, *Fisch. ex Loud. Hort. Brit.* 329, nomen.—Sibir.
lyrata, Franch. in Mém. Soc. Sc. Nat. Cherb. xxiv. (1884) 229 = affinis.
macrocephala, Less. in Linnaea, vi. (1831) 87 = glomerata.
macrophylla, Bunge, ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) ii. 17 = latifolia.
macrophylla, Sauter, in Flora, xxiii. (1840) 413 = alpina.
malitiosa, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 493.—China.
marginata, *Borszcz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 482, nomen.—As. centr.
Maximowiczii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1869) ii. 14.—Japon.
Medusa, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 488.—China; Mongolia.
mollis, Spreng. Syst. iii. 381 = Jurinea humilis.
monocephala, Cass. in Dict. Sc. Nat. l. 444 = pygmaea.
monticola, Richards. in Frankl. Narr. 1st Journ. App. ed. II. 29 = alpina.
mucronata, Spreng. ex DC. Prod. v. 376 = Pluchea lanceolata.
multiflora, DC. in Ann. Mus. Par. xvi. (1810) 199 = salicifolia.
multiflora, Richards. in Frankl. Narr. 1st Journ. App. 747 = alpina.
nepalensis, Spreng. Syst. iii. 380 = eriostemon.
nigrescens, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 491.—China.
nikoënsis, *Franch. & Sav. Enum. Pl. Jap.* ii. 407.—Japon.
nipponica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 183.—Japon.
nivea, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. vii. 153 = eriolepis.

SAUSSUREA :—
nuda, Ledeb. Ic. Pl. Fl. Ross. i. 15. t. 61 = alpina.
obvallata, *Wall. Cat.* sub n. 2906.—Reg. Himal.
odontolepis, *Sch. Bip. ex Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) II. 13.—China; Manchuria.
ovata, *Benth. in Henders. & Hume, Lahore to Yarkand*, 325.—Turkestan.
paleata, *Maxim. Prim. Fl. Amur.* 168.—Reg. Amur.
papposa, Turcz. ex DC. Prod. vi. 534 = crassifolia.
parviflora, DC. in Ann. Mus. Par. xvi. (1810) 200 = serrata.
pectinata, *Bunge, ex DC. Prod.* vi. 538.—China.
peguensis, *C. B. Clarke, Comp. Ind.* 235.—Burma.
phaeantha, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 489.—China.
piptathera, *Edgew. in Trans. Linn. Soc.* xx. (1846) 76, in obs.—Reg. Himal.
Poljakowi, *Glehn, in Act. Hort. Petrop.* iv. (1876) 60.—Sibir.
polyphylla, Schrenk, ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 5 = pygmaea.
polypodiifolia, DC. in Ann. Mus. Par. xvi. (1810) 201 = elongata.
polypodiifolia, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) II. 46 = runcinata.
polystichoides, *Hook. f. Fl. Brit. Ind.* iii. 369.—Reg. Himal.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 494.—China.
pterocaulon, Decne. in Jacquem. Voy. Bot. 95. t. 103 = Candolleana.
pujolica, *Costa, Suplem. Fl. Catalon.* 47.—Hispan.
pulchella, Fisch. ex DC. Prod. vi. 537 = japonica.
pulvinata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 493.—China.
Pumilio, Fisch. ex DC. Prod. vi. 535 = alpina.
pycnocephala, *Ledeb. Ic. Pl. Fl. Ross.* i. 15. t. 59.—Sibir. altaic.
pygmaea, *Spreng. Syst.* iii. 381.—Europ.; Sibir.
religiosa, Royle, Illustr. Bot. Himal. 26, 251 = Jurinea macrocephala, *Benth.*
Riederi, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) II. 35.—Kamtchat.
rigida, *Ledeb. Ic. Pl. Fl. Ross.* i. 19. t. 79.—Sibir. altaic.
robusta, *Ledeb. l. c.* i. 16. t. 65.—Sibir. altaic.
Roylei, *C. B. Clarke, Comp. Ind.* 229.—Reg. Himal.
runcinata, *DC. in Ann. Mus. Par.* xvi. (1810) 202.—Sibir.
sachalinensis, *F. Schmidt, Reisen Amurl.* 152.—Reg. Amur.
sacra, *Edgew. in Trans. Linn. Soc.* xx. (1846) 76.—Reg. Himal.
salicifolia, *DC. in Ann. Mus. Par.* xvi. (1810) 200.—Sibir.
salsa, *Spreng. Syst.* iii. 381.—Reg. Cauc.; As. bor.
scabiosaefolia, Turcz. ex Ledeb. Fl. Ross. iii. 666 = elongata.
scabra, Less. in Linnaea, ix. (1834) 180 = glomerata.
scaposa, *Franch. & Sav. Enum. Pl. Jap.* ii. 480.—Japon.
Schanginiana, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 5 = pygmaea.
Schlagintweitii, Klatt, in Sitzb. Akad. Muench. (1878) 94 = bracteata?
Schultzii, *Hook. f. Fl. Brit. Ind.* iii. 366.—Reg. Himal.
Semenovii, *Herd. in Bull. Soc. Nat. Mosc.* xl. (1867) II. 142.—Turkestan.
serrata, *DC. in Ann. Mus. Par.* xvi. (1810) 199.—Sibir. altaic.
serratuloides, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) II. 35.—Sibir. altaic.
setifolia, Klatt, in Sitzb. Akad. Muench. (1878) 95 = subulata.
sordida, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 389.—Sibir. altaic.
sorocephala, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 226.—Reg. Himal.
sorocephala, *Schrenk, Enum. Pl. Nov.* ii. 38.—Sibir. altaic.
squarrosa, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) II. 44.—Dahuria.
Stella, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 490.—China.
stemmaphora, Klatt, in Sitzb. Akad. Muench. (1878) 93 = Jurinea ceratocarpa.
Stoliczkae, *C.B. Clarke, Comp. Ind.* 225.—Reg. Himal.

SAUSSUREA :—
stricta, Spreng. ex DC. Prod. vi. 540 = affinis.
Stubendorffii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) II. 26.—Sibir.
subsinuata, Ledeb. Ic. Pl. Fl. Ross. i. 15. t. 62 = alpina.
subulata, *C. B. Clarke, Comp. Ind.* 226, partim.—Reg Himal.
subulata, C. B. Clarke, l. c., partim = wernerioides.
Sughoo, *C. B. Clarke, l. c.* 225.—Reg. Himal.
sylvatica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 495.—China.
Tanakae, *Franch. & Sav. ex Maxim. l. c.* xix. (1874) 516.—Japon.
tangutica, *Maxim. l. c.* xxvii. (1881) 489.—China.
taraxacifolia, *Wall. Cat.* n. 2914.—Reg. Himal.
tenuis, *Ledeb. Ic. Pl. Fl. Ross.* i. 19. t. 78.—Sibir. altaic.
Thomsoni, *C. B. Clarke, Comp. Ind.* 227.—Reg. Himal.
Tilesii, Ledeb. Ic. Pl. Fl. Ross. i. 15. t. 60 = alpina.
triangulata, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 58).—Sibir. or.
tridactyla, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 377.—Reg. Himal.
triptera, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 515.—Japon.
uniflora, C. B. Clarke, Comp. Ind. 229 = Roylei.
uniflora, *Wall. Cat.* n. 2916.—Reg. Himal.
ussuriensis, *Maxim. Prim. Fl. Amur.* 167.—Manchuria.
wernerioides, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 367.—Reg. Himal.
Yakla, *C. B. Clarke, Comp. Ind.* 227.—Reg. Himal.

SAUSSUREA, Salisb. in Trans. Linn. Soc. viii. (1807) 11 = **Funkia**, Spreng. (Liliac.).
caerulea, Salisb. l. c. = F. ovata.
cordifolia, Salisb. l. c. = Lilium cordifolium.

SAUSSURIA, Moench, Meth. 388 (1794) = **Nepeta**, Linn. (Labiat.).
pinnatifida, Moench, l. c. = N. lavandulacea.

SAUSSURIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = **Saussurea**, DC. (Compos.).

SAUTIERA, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 383. *ACANTHACEAE*, Benth. & Hook. f. ii. 1088.
SANTIERA, Span. in Linnaea, xv. (1841) 328, sphalm.
Decaisnii, Nees, in DC. Prod. xi. 98 = tinctorum.
tinctorum, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 383.—Timor.

SAUVAGEA, Linn. Gen. ed. II. 241 (1742) = seq.

SAUVAGESIA, Linn. Sp. Pl. ed. I. 203 (1753). *VIOLARIEAE*, Benth. & Hook. f. i. 120.
IRON, P. Br. Hist. Jamaic. 179. t. 12. f. 3 (1756).
SAUVAGEA, Linn. Gen. ed. II. 241 (1742).
Adima, Aubl. Pl. Gui. i. 251 = erecta.
deflexifolia, *Gardn. in Hook. Ic. Pl.* t. 484.—Bras.
deflexifolia, Miq. ex Eichl. in Mart. Fl. Bras. xiii. 1. 409 = erecta.
elata, *Benth. in Hook. Journ. Bot.* iv. (1842) 107.—Ind. occ.; Guiap.
elegantissima, A. St. Hil. in Mém. Mus. Par. ix. (1822) 325 = Lavradia elegantissima.
erecta, Aubl. Pl. Gui. i. 254; Spreng. Neue Entdeck. i. 296 = Sprengelii.
erecta, *Linn. Sp. Pl.* 203.—Am. et Afr. trop.
ericoides, Ging. in DC. Prod. i. 316 = Lavradia alpestris.
fruticosa, *Mart. Nov. Gen. et Sp.* i. 38.—Bras.
geminiflora, [Moç. & Sesse,] ex Ging. in DC. Prod. i. 315 = erecta.
gracilis, Steud. Nom. ed. II. ii. 517, in syn. = erecta.
Kappleri, Miq. Stirp. Surin. Sel. 105 = Sprengelii.
laxa, Mart. Nov. Gen. et Sp. i. 38 = erecta.
linearifolia, A. St. Hil. in Mém. Mus. Par. xi. (1824) 106 = pusilla.
longifolia, *Eichl. in Mart. Fl. Bras.* xiii. 1. 407.—Bras.
nutans, Pers. Syn. i. 253 = erecta.
ovata, Mart. Nov. Gen. et Sp. i. 36 = racemosa.
pendula, Mart. ex Ging. in DC. Prod. i. 316 = Lavradia alpestris.
peruviana, Roem. & Schult. Syst. v. 457 = erecta.

SAUVAGESIA :—
pulchella, *Planch. in Seem. Bot. Voy. Herald*, 80.—Am. trop.
pusilla, *Mart. Nov. Gen. et Sp.* i. 35.—Bras.
racemosa, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 98.—Bras.
ramosissima, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 409.—Bras.
rubiginosa, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 101.—Bras.
salicifolia, *Ging. in DC. Prod.* i. 316.—Bras.
Salzmannii, Benth. ex Eichl. in Mart. Fl. Bras. xiii. 1. 412 = tenella.
serpyllifolia, Mart. Nov. Gen. et Sp. i. 37. t. 25 = Sprengelii.
Sprengelii, *A. St. Hil. in Mém. Mus. Par.* xi. (1824) 99.—Am. trop.
tenella, *Lam. Illustr.* ii. 119.—Am. trop.

SAUVAGIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = praec.

SAUVALLEA, Wright, in Sauv. Fl. Cub. 156 (1873). *COMMELINACEAE*, Benth. & Hook. f. iii. 855.
Blainii, *Wright, l. c.* 157.—Cuba.

SAVA, Adans. Fam. ii. p. (21) (1763) = **Onosma**, Linn. (Borag.).

SAVASTANA, Schrank, Baier. Fl. i. 100, 337 (1789) = **Hierochloë**, S. G. Gmel. (Gramin.).
hirta, Schrank, l. c. 337 = H. borealis.

SAVASTANIA, Scop. Introd. 213 (1777) = **Tiboucchina**, Aubl. (Melastom.).

SAVASTONIA, Neck. ex Steud. Nom. ed. II. ii. 518, err. typ. (1841) = praec.

SAVIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Amphicarpaea**, Ell. (Legumin.).
volubilis, Rafin. l. c. = A. monoica.

SAVIA, Willd. Sp. Pl. iv. 771 (1805). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 270.
CHARIDIA, Baill. Etud. Gén. Euph. 572 (1858).
GONATOGYNE, Klotzsch, ex Muell. Arg. in Mart. Fl. Bras. xi. II. 13 (1873).
PETALODISCUS, Baill. Etud. Gén. Euph. 571. t. 22 (1858).
Actephila, Hassk. Cat. Hort. Bogor. Alt. 243 = Actephila excelsa.
Bojeriana, *Baill. Adansonia*, viii. (1868) 345.—Madag.
clusiifolia, *Griseb. in Goett. Nachr.* (1865) 164.—Cuba.
dictyocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 704.—Bras.
elegans, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Ins. Mascar.
erythroxyloides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 157.—Cuba.
Laureola, *Baill. Adansonia*, ii. (1861–62) 33.—Madag.
laurifolia, *Griseb. in Goett. Nachr.* (1865) 163.—Cuba.
mimosoides, *Baill. Adansonia*, ii. (1861–62) 34.—Madag.
oblongifolia, *Baill. l. c.* 35.—Madag.
platyrachis, *Baill. l. c.* 33.—Madag.
pulchella, *Baill. l. c.* 35.—Madag.
sessiliflora, *Willd. Sp. Pl.* iv. 771.—Ins. S. Doming.
zeylanica, Baill. Etud. Gén. Euph. 571, nomen = Actephila excelsa.

SAVIGNYA, DC. Syst. ii. 283 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 85.
aegyptiaca, Coss. in Bull. Soc. Bot. Fr. iv. (1857) 395 = longistyla.
aegyptiaca, *DC. Syst.* ii. 283.—Aegypt.
longistyla, *Boiss. Diagn.* Ser. II. v. 27.—Algeria.
parviflora Webb, Fragm. Fl. Aethiop. 14 = aegyptiaca.

SAVINIONA, Webb & Berth. Phyt. Canar. i. 30 (1836) = **Lavatera**, Linn. (Malvac.).
acerifolia, Webb & Berth. l. c. 31 = L. acerifolia.
brachyfolia, Walp. Rep. i. 291, sphalm. = seq.
brachyloba, Webb & Berth. Phyt. Canar. i. 32 = L. brachyfolia.

SAVIONA, Pritz. Ic. Ind. i. 989 (1855) = praec.

SAXEGOTHAEA, Lindl. in Journ. Hort. Soc. vi. (1851) 258. *CONIFERAE*, Benth. & Hook. f. iii. 434.
SAXEGOTHEA, Benth. & Hook. f. Gen. iii. 434 (1880), err. typ.
SQUAMATAXUS, [Nelson] Senilis, Pinaceae, 168 (1866).
conspicua, *Lindl. in Journ. Hort. Soc.* vi. (1851) 258; *et in Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 111.—Chili.
gracilis, Hort. ex Gord. Pinet. 280 = Podocarpus nubigena.

SAXEGOTHEA, Benth. & Hook. f. Gen. iii. 434 (1880) err. typ. = praec.

SAXIFRAGA, Tourn. ex Linn. Gen. ed. I. 131 (1737). *SAXIFRAGEAE*, Benth. & Hook. f. i. 635.
ADENOGYNE, APHOMONIX, EVAIEZOA, HEMIEVA, HETERISIA, HEXAPHOMA, PIAROPHYLA, PONISTA, STEIRANISIA, TELESONIX, TULORIMA, Rafin. Fl. Tellur. ii. 65–70 (1836).
ANTIPHYLLA, AULAXIS, CHRONDOSEA, CILIARIA, DERMASEA, HIRCULUS, LEPTASEA, LOBARIA, MEGASEA, MUSCARIA, SPATULARIA, Haw. Saxifr. Enum. 6–47 (1821).
BERGENIA, Moench, Meth. 664 (1794).
DIPTERA, Borck. in Roem. N. Mag. i. (1794) 29.
GERYONIA, Schrank, ex Hoppe, in Flora, i. (1818) 229.
HYDATICA, Neck. Elem. ii. 387 (1790).
KINGSTONIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 531 (1821).
LEPTARRHENA, Reichb. Handb. 230 (1837).
LIGULARIA, Duval, Pl. Succ. 11 (1809).
MICRANTHES, Haw. Syn. Pl. Succ. 320 (1812).
MISCOPETALUM, Haw. l. c. 323 (1812).
ROBERTSONIA, Haw. Syn. Pl. Succ. 321 (1812).
RUPIFRAGA, 'Linn.' ex Rafin. Fl. Tellur. ii. 67 (1836).
SAXIFRAGIA, Tourn. ex Linn. Syst. ed. I (1735).
SEKIKA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 209; Phil. Bot. ii. 49 (1791).
abyssinica, Rich. ex Engl. Monog. Saxifr. 76 = hederaefolia.
acaulis, Gaud. in Nat. Anz. Schw. Ges. i. (1818) 68; Fl. Helvet. iii. 127 = muscoides.
aciculata, Royle, Illustr. Bot. Himal. 226 = flagellaris?
aconitifolia, Field. & Gardn. Sert. Pl. t. 57 = Boykinia aconitifolia.
adenodes, *Poepp. ex Sternb. Rev. Saxifr. Suppl.* ii. 75. t. 26.—Am. austr.
adenophora, *C. Koch, in Linnaea*, xix. (1847) 40.—Macedon.; Cappadoc.
adoxoidea, Griff. Notul. iv. 428; Ic. Pl. Asiat. t. 570 = Chrysosplenium lanuginosum.
adscendens, Haw. Misc. Nat. 162 = altifida.
adscendens, Jacq. Coll. i. 197. t. 11 = controversa.
adscendens, *Linn. Sp. Pl.* 405.—Reg. bor. et arct.
adscendens, Vahl, in Skrivt. Nat. Selsk. Kjoeb. ii. 1. (1792) 12 = aquatica.
aestivalis, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 37 = punctata.
affinis, D. Don, in Trans. Linn. Soc. xiii. (1822) 418 = hypnoides.
afghanica, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 56.—Afghan.
aggregata, Lej. Rev. Fl. Spa, 81 = hypnoides.
aizoidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.
aizoides, *Linn. Sp. Pl.* 403.—Reg. bor. et arct.
aizoidoides, *Miégev. in Bull. Soc. Bot. Fr.* xii. (1865) 22.—Mont. Pyren.
Aizoon, Guss. ex DC. Prod. iv. 20 = lingulata.
Aizoon, *Jacq. Fl. Austr.* v. 18. t. 438.—Reg. bor. et arct.
Aizoon, Tenore, Prod. Fl. Nap. p. xxv = marginata.
aizoonia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Aizoon.
ajugaefolia, Wahlenb. Fl. Carpat. 122 = perdurans.
ajugifolia, *Linn. Cent. Pl.* 11; *Amoen. Acad.* iv. 271.—Europ. austr.
Alberti, *Regel & Schmalh. in Act. Hort. Petrop* v. (1877) 584.—Turkestan.
alchemilloides, *Griseb. in Goett. Abh.* xxiv. (1879) 142.—Reg. Argent.
Allioni, Baumg. Enum. Stirp. Transs. i. 378 = pedemontana.

SAXIFRAGA:—
Allionii, Gaud. in Nat. Anz. Schw. Ges. i. (1818) 68 = muscoides.
almeriensis, Willk. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 525 = Camposii.
alpina, Pav. ex D. Don, in Trans. Linn. Soc. xiii. (1822) 421 = peruviana.
alpina, Salisb. Prod. 306 = aizoides.
altaica, Steph. ex Ledeb. Fl. Ross. ii. 212 = melaleuca.
altifida, *Haw. Saxifr. Enum.* 24.—Hab.?
altissima, *Kern. in Zeitschr. Ferdinand. Tirol.* III. xv. (1870) 273.—Europ.
ambigua ×, DC. Fl. Fr. vi. 517.—Europ.
amplexifolia, Sternb. Rev. Saxifr. Suppl. i. 2. t. 2 = Leptarrhena amplexifolia.
ampullacea, Tenore, Fl. Nap. Prod. App. iv. 21; ex ej. Fl. Nap. 59, 60 = muscoides.
andicola, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 427.—Ecuador.
andicola, H. B. & K. Nov. Gen. et Sp. vi. 48 = cordillerarum.
Andrewsii ×, Harv. in Hook. Lond. Journ. Bot. vii. (1848) 570.
androsacea, Comolli, Prod. Fl. Comens. 78 = sedoides.
androsacea, *Linn. Sp. Pl.* 399.—Europ.
androsacea, Pursh, Fl. Am. Sept. i. 310 = caespitosa?
androsacea, Sibth. & Sm. Fl. Graec. Prod. i. 276 = exarata.
angulosa, Schott, Nym. & Kotschy, Analect. Bot. 29 = rotundifolia.
angustifida, Hort. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 526 = hypnoides.
angustifolia, Hall. f. ex Gaud. in Nat. Anz. Schw. Ges. i. (1818) 67 = Seguieri.
angustifolia, Haw. Misc. 166; Hort. ex DC. Prod. iv. 31; Schleich, ex Sternb. Rev. Saxifr. Suppl. ii. 70 = hypnoides.
annua, Lapeyr. Fl. Pyren. 59 = tridactylites.
aphylla, *Sternb. Rev. Saxifr.* 40. t. 11.—Europ.
aphylla, Tenore, Fl. Nap. Prod. App. iv. 21 = sedoides.
apennina, Bertol. f. in Bull. Sc. Med. Ser. II. ix. 65; ex Bertol. Fl. Ital. iv. 479 = cuneifolia.
aquatica, Bieb. Fl. Taur. Cauc. i. 317 = irrigua.
aquatica, *Lapeyr. Fl. Pyren.* t. 28.—Mont. Pyren.
arachnoidea, *Sternb. Rev. Saxifr.* 23. t. 15.—Helvet.
arachnoidea, Zanted. in Comm. Ateneo Brescia, 1815 (1818), 90 = praec.?
arctica, Willd. ex Sternb. Saxifr. Suppl. ii. 66 = muscoides.
arenarioides, Brign. Fasc. Pl. Forojul. 24 = tenella.
aretioides ×, Bluff & Fingerh. Comp. Fl. Germ. ed. II. i. 11. 60.—Europ.
aretioides, *Lapeyr. Fl. Pyren.* 28. t. 13.—Mont. Pyren.
arguta, D. Don, in Trans. Linn. Soc. xiii. (1822) 356 = punctata.
aristulata, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 68.—Reg. Himal.
arundana, Boiss. Voy. Bot. Espagne, ii. 693. t. 64 B = dichotoma.
asarifolia, Sternb. Rev. Saxifr. Suppl. t. 24 = odontophylla.
ascendens, Auct. = adscendens.
aspera, Bieb. Fl. Taur. Cauc. i. 314 = flagellaris.
aspera, *Linn. Sp. Pl.* 402.—Europ. austr.
atlantica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 48.—Hispan.
atrata, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 117.—China.
atropurpurea, Wulf. ex Sternb. Rev. Saxifr. 41. t. 11. b. f. 2 = muscoides.
atrorubens, Bertol. in Desv. Journ. Bot. ii. 11. (1813) 76; et Amoen. Ital. 99 = aizoides.
atrorubens, Sternb. Rev. Saxifr. Suppl. i. 10 = muscoides.
aurea, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 397 = flagellaris.
aurea, Steud. Nom. ed. I. 735 = Chrysosplenium alternifolium, oppositifolium.
australis, Moric. Fl. Venet. i. 431 = lingulata.
autumnalis, Jacq. Obs. i. 31 = S. Aizoon.
autumnalis, Pall. ex Sternb. Rev. Saxifr. Suppl. ii. 19 = serpyllifolia.
autumnalis, Linn. Sp. Pl. 402 = aizoides.
axillaris, Dulac, Fl. Hautes-Pyr. 268 = ajugifolia.
axillaris, Kit. in Linnaea, xxxii. (1863) 477 = perdurans.
Baumgarteni, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 126 = retusa.

SAXIFRAGA:—
Baumgarteni, Schur, Enum. Pl. Transs. 239 = adscendens.
Bellardi, All. Fl. Pedem. ii. 72 = controversa.
Besleri, Sternb. Rev. Saxifr. Suppl. ii. 52 = Hostii.
bicolor, Sternb. l. c. 49 = serpyllifolia.
bicuspidata, *Hook. f. Fl. Antarct.* 281. t. 97.—Reg. Magell.
bifida, Hook. ex Engl. Monog. Saxifr. 202 = bicuspidata.
biflora, *All. Fl. Pedem.* ii. 71. t. 21. f. 1.—Europ. austr.
biflora, Ledeb. Fl. Ross. ii. 205 = oppositifolia.
bimaculata, Turcz. in Bull. Soc. Nat. Mosc. xiii. (1840) 71 = Sieversiana.
biternata, *Boiss. Voy. Bot. Espagne*, ii. 231. t. 64.—Hispan.
Blanca, *Willk. Fl. Hisp. Balear.* 8.—Hispan.
bohemica, Panz. ex Koch, Syn. Fl. Germ. ed. I. 274 = hypnoides.
Boissieri, *Engl. in Verh. Zool.-Bot. Ges. Wien*, xix. (1869) 528.—Hispan.
Bongardi, Presl, ex Engl. l. c. = leucanthemifolia.
Bonplandii, D. Don, in Trans. Linn. Soc. xiii. (1822) 430 = peruviana.
Boryi, *Boiss. & Heldr. Diagn.* Ser. II. ii. 65.—Graecia.
Bourgaeana, *Boiss. & Reut. l. c.* 71.—Hispan.
Boussingaultii, *Brongn. in Ann. Sc. Nat.* Sér. II. iii. (1835) 49. t. 1. f. 1.—Ecuador.
brachypoda, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 378.—Reg. Himal.
bracteata, *D. Don, l. c.* 367.—Sibir.
bronchialis, *Linn. Sp. Pl.* 400.—Reg. bor. et arct.
Brunoniana, *Wall. Cat.* n. 444.—Reg. Himal.
Brunonis, Ser. in DC. Prod. iv. 45 = praec.
bryoides, Linn. Sp. Pl. 400 = aspera.
bryoides, Pall. ex D. Don, in Trans. Linn. Soc. xiii. (1822) 382 = nitida.
bryophora, *A. Gray, in Proc. Am. Acad.* vi. (1865) 533.—Calif.
bulbifera, *Linn. Sp. Pl.* 403.—Europ. austr.
bulbifera, Oed. Fl. Dan. t. 390 = cernua.
bulbosa, Hochst. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 529 = granulata.
Burseriana, Lapeyr. Hist. Abr. Pl. Pyr. i. 226 = aretioides.
Burseriana, *Linn. Sp. Pl.* 400.—Europ. austr.
caerulea, Pers. Syn. i. 488 = oppositifolia.
caesia, *Linn. Sp. Pl.* 399.—Europ. austr.
caesia, Scop. Fl. Carn. ed. II. i. 294 = squarrosa.
caespitosa, Asso, Syn. Stirp. Arag. 52 = pentadactylis.
caespitosa, A. Gray, in Proc. Am. Acad. viii. (1873) 383 = Spiraea pectinata.
caespitosa, Lag. & Rodr. in Anal. Cienc. Nat. v. (1802) 278; D. Don, in Trans. Linn. Soc. xiii. (1822) 428 = mixta.
caespitosa, *Linn. Sp. Pl.* 404.—Reg. bor. et arct.
caespitosa, Scop. Fl. Carn. ed. II. i. 293 = muscoides.
caespitosa, Sm. Engl. Fl. ii. 273 = hypnoides.
caespitosa, Vill. Hist. Pl. Dauph. iii. 673 = exarata.
californica, Nutt. ex Torr. & Gray, Fl. N. Am. i. 585 = Tellima Cymbalaria.
callosa, Sm. in Rees, Cycl. xxxi. [n. 2] = longifolia.
calyciflora, Lapeyr. Fl. Pyren. 28. t. 12 = media.
calycina, Sternb. Rev. Saxifr. Suppl. ii. 10 = davurica.
Camposii, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 47.—Hispan.
canaliculata, *Boiss. & Reut. ex Engl. Monog. Saxifr.* 169.—Europ.
Candollei, Tausch, in Syll. Ratisb. ii. (1828) 242 = geranioides.
Candollii, Salzm. ex Pouz. in Mém. Soc. Linn. Par. iv. (1826) 561 = pedemontana.
canescens, *Haw. Syn. Pl. Succ.* 325.—Hab.?
cantabrica, Boiss. & Reut. ex Engl. Monog. Saxifr. 191 = hypnoides.
capillipes, Reichb. ex Engl. l. c. 229 = cuneifolia.
capitata, Lapeyr. Fl. Pyren. 55. t. 30 = aquatica.
Careyana, *A. Gray, in Hook. Lond. Journ. Bot.* ii. (1843) 116.—Am. bor.
carinthiaca, Schott, Nym. & Kotschy, Analect. Bot. 24 = S. Aizoon.
carnosa, Lucé, Topog. Nachr. Oesel, 137 = granulata.
caroliniana, *A. Gray, in Mem. Am. Acad.* N. S. iii. (1848) 39.—Am. bor.
caroliniana, Hort. ex Ser. in DC. Prod. iv. 39 = pensylvanica.

SAXIFRAGA :—
caroliniana, Schleich. ex Ser. in DC. Prod. iv. 41 = erosa.
carpathica, *Reichb. Fl. Germ. Excurs.* 552.—Transsylv.
carpetana, Boiss. & Reut. Diagn. Pl. Nov. 12 = atlantica.
cartilaginea, *Willd. ex Sternb. Rev. Saxifr.* 5. t. 3.—Reg. Caucas. ; Persia.
castellana, Boiss. & Reut. ex Nym. Consp. 273 = granulata.
catalaunica, Boiss. & Reut. Diagn. Ser. II. ii. 64 = lingulata.
ceratophylla, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 70 = trifurcata.
ceratophylla, Willd. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 530 = pentadactylis.
cernua, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 409, partim = debilis.
cernua, Lapeyr. Hist. Abr. Pl. Pyr. i. 232 = granulata.
cernua, *Linn. Sp. Pl.* 403.—Reg. bor. et arct.
cervicornis, Viv. Fl. Cors. Prod. App. 2 = pedemontana.
cespitosa, Scop. Ann. II. Hist. Nat. (1769) 51 = *caespitosa*, Scop., etc.
Chamissoi, Sternb. Rev. Saxifr. Suppl. i. 12. t. 10. f. 1 = tricuspidata.
Chamissonis, Sternb. ex Ser. in DC. Prod. iv. 35 = rivularis.
cherlerioides, D. Don, in Trans. Linn. Soc. xiii. (1822) 382 = bronchialis.
cherlerioides, Presl, ex Ledeb. Fl. Ross. ii. 208 = nitida.
chilensis, Poepp. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 530 = adenodes.
chinensis, Lour. Fl. Cochinch. 281 = sarmentosa.
chlorantha, Dulac, Fl. Hautes-Pyr. 269 = S. Aizoon.
chlorantha, Schur, Enum. Pl. Transs. 233 = luteoviridis.
chrysantha, *A. Gray, in Proc. Am. Acad.* xii. (1877) 83.—Am. bor. occ.
chrysosplenifolia, *Boiss. Diagn.* Ser. I. iii. 20.—Graecia ; Creta.
ciliaris, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 55 = mixta.
ciliata, *Lindl. Bot. Reg.* (1843) t. 65.—Reg. Himal.
ciliata, Royle, Illustr. Bot. Himal. 226. t. 49. f. 2 = ligulata.
ciliata, Walp. Rep. ii. 365 ; v. 827 = imbricata.
ciliolata, Schur, Enum. Pl. Transs. 231 = S. Aizoon.
circinnata, Hort. Monac. ex Reichb. Fl. Germ. Excurs. 552 = controversa.
citrina, *Hegetschw. Fl. Schw.* 398.—Helvet.
Clusii, Auct. Germ. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 132 = stellaris.
Clusii, *Gouan, Illustr.* 28.—Mont. Pyren.
Clusii, Kunze, in Linnaea, xx. (1847) 43 = leucanthemifolia.
cochlearіaefolia ×, Schrad. ex Kunze, l. c. xvi. (1842) 318.—Europ.
cochlearis, Reichb. Fl. Germ. Excurs. 559 = lingulata.
coerulea, Auct. = *caerulea*.
columnaris, Steud. Nom. ed. I. 735 = pedemontana.
comosa, Fellm. in Bull. Soc. Bot. Fr. x. (1863) 500 = stellaris.
compacta, *Adams, in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 239.—Sibir. or.
compacta, Sternb. Rev. Saxifr. Suppl. i. 9. t. 1. f. 3 = valdensis.
condensata, D. Dietr. Syn. Pl. ii. 1537, partim = sileniflora.
condensata, C. C. Gmel. Fl. Bad. ii. 226. t. 3 = hypnoides.
confusa, Lej. Rev. Fl. Spa, 80 = hypnoides.
congesta, Schleich. Cat. Pl. Helvet. ed. IV. 63 = hypnoides.
congesta, Willd. ex Ledeb. Fl. Ross. ii. 207 = bronchialis.
conifera, *Coss. & Dur. in Bull. Soc. Bot. Fr.* xi. (1864) 332.—Hispan.
controversa, Hort. Lond. ex Haw. Saxifr. Enum. 32 = hypnoides.
controversa, *Sternb. Rev. Saxifr.* 43. t. 16.—Europ. austr. ; Am. bor.
corbariensis, Timb. in Mém. Acad. Toul. Sér. VII. vii. (1875) 469 = geranioides.
cordifolia, Haw. Misc. Nat. 157 = crassifolia.
cordigera, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 68.—Reg. Himal.
cordillerarum, *Presl, Rel. Haenk.* ii. 55.—Peruv.
coriacea, *Adams, in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 240.—Sibir. arct.

SAXIFRAGA :—
coriacea, Fisch. ex Ser. in DC. Prod. iv. 48 = Leptarrhena pyrolaefolia.
coriophylla, Griseb. Spicil. Fl. Rumel. i. 333 = Rochеliana.
coronensis, Schur, Enum. Pl. Transs. 239 = adscendens.
corsica, Gren. & Godr. Fl. Fr. i. 642 = granulata.
cortusaefolia, *Sieb. & Zucc. in Akad. Muench.* iv. II. (1843) 190.—Japon.
corymbosa, *Boiss. Diagn.* Ser. I. iii. 17.—Caria.
corymbosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 70.—Reg. Himal.
corymbosa, Lucé, Topog. Nachr. Oesel, 137 = granulata.
Cossoniana, Boiss. & Reut. Diagn. Ser. II. ii. 70 = granulata.
Cotyledon, Bieb. Fl. Taur. Cauc. i. 313 = cartilaginea.
Cotyledon, *Linn. Sp. Pl.* 398.—Europ.
Cotyledon, Tenore, Prod. Fl. Nap. p. xxv = marginata.
Cotyledon, Wulf. ex Jacq. Coll. iv. 291 = S. Aizoon.
crassifolia, *Linn. Sp. Pl.* 401.—Sibir.
crateriformis, Hort. Oxon. ex Ser. in DC. Prod. iv. 32 = hypnoides.
crenata, Steud. Nom. ed. I. 735 ; Sweet, Hort. Brit. ed. I. 181 = umbrosa.
crocea, Gaud. in Nat. Anz. Schw. Ges. i. (1818) 69 = muscoides.
crustacea, Hoppe, Taschenb. (1805) 237 = seq.
crustata, *Vest, in Bot. Zeit. Regensb.* iii. (1804) 95 ; iv. (1805) 42 ; *Man. Bot.* 656.—Europ. austr.
cultrata, Schott, Nym. & Kotschy, Analect. Bot. 23 = S. Aizoon.
cuneata, *Willd. Sp. Pl.* ii. 658.—Hispan.
cuneifolia, Cav. Ic. iii. 25 = cuneata.
cuneifolia, *Linn. Syst.* ed. X. 1026.—Europ. austr.
curvata, Schleich. ex Ser. in DC. Prod. iv. 31 = hypnoides.
cuscutaeformis, *Lodd. Bot. Cab.* t. 186.—Japon. ?
cuspidata, Schleich. Cat. Pl. Helvet. ed. IV. 63 = pedatifida.
Cymbalaria, *Linn. Sp. Pl.* 405.—As. Min. ; Persia.
Cymbalaria, Sibth. & Sm. Fl. Graec. iii. 72. t. 378 = Sibthorpii.
Cymbalaria, Sternb. in Linnaea, vi. (1831) 555 = rivularis.
Cymbalaria, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 261 = sibirica.
cymbalariaefolia, Ehrenb. ex Engl. Monog. Saxifr. 77 = hederacea.
cymosa, Waldst. & Kit. Pl. Rar. Hung. i. 91. t. 88 = pedemontana.
dahurica, Lyall, ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 542 = Lyallii.
dahurica, Schleich. ex Ser. in DC. Prod. iv. 42 = cuneifolia.
davurica, Torr. & Gray, Fl. N. Am. i. 569, partim = Lyallii.
davurica, *Willd. Sp. Pl.* ii. 645.—As. bor. ; Am. bor. occ.
Debeauxii, Pomel, Nouv. Mat. Fl. Atl. 152 = dichotoma.
debilis, *Engelm. ex A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 62.—Am. bor. occ.
decipiens, Ehrh. Beitr. v. 47, 175 = hypnoides.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 6.—China.
demissa, Schott & Kotschy, in Oestr. Bot. Zeitschr. ix. (1859) 8 = mutata.
densa, Haw. Misc. Nat. 166 = hypnoides.
densa, Willd. ex Sternb. Rev. Saxifr. Suppl. ii. 26 = nitida.
densifolia, Schleich. ex Ser. in DC. Prod. iv. 31 = hypnoides.
dentata, Link, Enum. Hort. Berol. i. 413 = umbrosa.
denticulata, Dulac, Fl. Hautes-Pyr. 270 = aretioides.
denudata, D. Don, in Trans. Linn. Soc. xiii. (1822) 421 = hypnoides.
denudata, Nutt. ex Torr. & Gray, Fl. Bor. Am. i. 567 = punctata.
depressa, Sternb. Rev. Saxifr. 42. t. 11 = androsacea.
diapensioides, *Bell. in Mém. Acad. Turin*, v. 1790–1791 (1793), 227. t. 5.—Europ. austr.
diapensioides, Fisch. ex Ser. in DC. Prod. iv. 22 = serpyllifolia.
diapensioides, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 54 = media.

SAXIFRAGA :—
diapensordes, Kurz, in Flora, lv. (1872) 286, sphalm. = diapensioides ?
dichotoma, *Willd. ex Sternb. Rev. Saxifr.* 51.—Hispan.
digitata, Lapeyr. ex Steud. Nom. ed. I. 736 = pentadactylis.
digitata, Pourr. ex Steud. l. c. ed. II. ii. 520 = muscoides.
dilatata, Schott, Nym. & Kotschy, Analect. Bot. 24 = S. Aizoon.
dissimilis, G. Don, in Loud. Hort. Brit. 176 = rivularis ?
divaricata, Ram. ex DC. Fl. Fr. iv. 374 = exarata.
diversifolia, *Wall. Cat.* n. 452.—Reg. Himal.
diversifolia, *Schleich. Cat. Pl. Helvet.* ed. IV. 63.—Europ. centr.
dubia, Schleich. ex Ser. in DC. Prod. iv. 32 = hypnoides.
egregia, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 113.—China.
elata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 575 = Boykinia occidentalis.
elatior, Mert. & Koch, Deutschl. Fl. iii. 115 = Hostii.
elatior, Wimm. Fl. Schles. (1857) 470 = S. Aizoon.
elegans, Mackay, Fl. Hibern. i. 65 = S. Geum.
elegans, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 573.—Am. bor. occ.
elegans, Sternb. Rev. Saxifr. Suppl. ii. 34 = exilis.
elegans, Zeyh. ex Schrank, in Syll. Ratisb. i. (1824) 208 = seq. ?
elegantula, Boiss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 449 = granulata.
elongata, Panz. ex Ser. in DC. Prod. iv. 32 = hypnoides.
elongata, Sternb. Rev. Saxifr. 9. t. 4 = virginiensis.
elongella, Haw. Misc. Nat. 167 = muscoides.
elongella, Sm. in Trans. Linn. Soc. x. (1811) 340 = hypnoides.
Engleri, *Dalla Torre, Atl. Alpenfl.* 102.—Europ.
erecta, Tenore, Prod. Fl. Nap. p. xxv = S. Aizoon.
erioblasta, *Boiss. & Reut. Diagn.* Ser. II. ii. 67.—Hispan.
eriophora, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 372.—Am. bor.
erosa, *Pursh, Fl. Am. Sept.* i. 311.—Am. bor.
Eschholzii, *Sternb. Rev. Saxifr. Suppl.* i. 9. t. 10. f. 3.—Am. bor. et arct.
Eschscholtzii, Cham. ex DC. Prod. iv. 22 = serpyllifolia.
evolvuloidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.
evolvuloides, Wall. Cat. n. 447 = hispidula.
exarata, All. Fl. Pedem. ii. 73. t. 88. f. 2 = muscoides.
exarata, Griseb. Spicil. Fl. Rumel. i. 335 = adenophora.
exarata, Hook. f. Fl. Antarct. ii. 280 = cordillerarum.
exarata, Hook. Fl. Bor. Am. i. 244 (excl. syn.) = caespitosa.
exarata, *Vill. Prosp.* 47 ; *Hist. Pl. Dauph.* iii. 674. t. 44.—Europ. austr.
excavata, Kunze, in Linnaea, xvi. (1842) 319 = cuneifolia.
exilis, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 29 = tridactylites.
exilis, *Steph. ex Sternb. Rev. Saxifr. Suppl.* i. 8. t. 3. f. 1.—Sibir. ; Am. arct.
Facchini, Koch, in Flora, xxv. (1841) 624 ; Syn. Fl. Germ. ed. II. 445 = muscoides.
Federici-Augusti, Biasoletti, Rel. Viagg. Dalm. 199, 213 = porophylla.
ferruginea, R. Grah. in Edinb. N. Phil. Journ. (Jul.-Oct. 1829) 349 = leucanthemifolia.
filamentosa, Cham. ex Ser. in DC. Prod. iv. 45 = flagellaris.
filicaulis, *Wall. Cat.* n. 445.—Reg. Himal.
fimbriata, D. Don, in Trans. Linn. Soc. xiii. (1822) 404 = Eschholtzii.
fimbriata, *Wall. Cat.* n. 443.—Reg. Himal.
Fischeri, Ser. in DC. Prod. iv. 22 = serpyllifolia.
flabellifolia, R. Br. ex Torr. & Gray, Fl. N. Am. i. 569 = davurica.
flagellaris, *Willd. ex Sternb. Rev. Saxifr.* 25. t. 6.—Reg. bor. et arct.
flava, Lam. Fl. Fr. iii. 529 = S. Hirculus.
flavescens, Sternb. Rev. Saxifr. Suppl. i. 13. t. 9. f. 2 = hypnoides.
Flittneri, Heuff. in Flora, xxxvii. (1854) 289 = perdurans.
floribunda, D. Dietr. Syn. Pl. ii. 1532 = florulenta.

SAXIFRAGA:—
florulenta, *Moretti, in Brugnat. Giorn. Fis.* II. vii. (1824) 105.—Europ. austr.
foliolosa, R. Br. in Parry, 1st Voy. App. 275 = stellaris.
foliosa, D. Dietr. Syn. Pl. ii. 1527 = praec.
foliosa, Orsini, ex Tenore, Syll. Fl. Neap. 203 = androsacea.
fonticola, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 90 = rotundifolia.
Forbesii, *Vasey, in Am. Entomologist & Botanist*, ii. (1870) 288.—Am. bor.
Forsteri ×, Stein, in Oestr. Bot. Zeitschr. xxvii. (1877) 291.—Europ.
Fortunei, *Hook. Bot. Mag.* t. 5377.—China.
fragarioides, *Greene, in Bull. Torrey Bot. Club.* vii. (1881) 121.—Calif.
fragilis, Hort. Schweig. ex Schrank, Pl. Rar. Hort. Monac. t. 92 = trifurcata.
fragilis, Sieber, ex Steud. Nom. ed. I. 736 = rotundifolia.
Frederici-Augusti, Biasoletto, Rel. Viagg. Dalm. t. 1 = porophylla.
furcata, *Lapeyr. ex Steud. Nom.* ed. I. 736.—Mont. Pyren.
fusca, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 38. —Japon.
Gaudini ×, Bruegg. Cat. Hort. Turic. (1868) 5.—Europ.
geifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Geum.
gemmifera, Pers. Syn. i. 490 = hypnoides.
gemmulosa, *Boiss. in Bibl. Univ. Genèv.* N. S. xiii. (1838) 409.—Hispan.
geranioides, Geners. Elench. Scepus. 31 = perdurans.
geranioides, Georgi, Beschr. Russ. Reich. iv. 965 = neglecta.
geranioides, Host, Syn. Fl. Austr. 231 = petraea, *Linn.*
geranioides, *Linn. Cent. Pl.* i. 10; *Amoen. Acad.* iv. 271.—Mont. Pyren.
geranioides, Prey. & Zirk. ex Bab. in Journ. Linn. Soc. xi. (1870) 308 = petraea, *Prey. & Zirk.*
Geum, *Linn. Sp. Pl.* 401.—Europ. occ.
Geum, Pall. Reise, ii. 35; Pursh, Fl. Am. Sept. i. 311 = punctata.
Geum, Sibth. & Sm. Fl. Graec. Prod. i. 276 = taygetea.
gibraltarica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 47 = globulifera.
Girtanneri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 84.—Europ.
glabella, *Bertol. in Giorn. Arcad.* xxi. (1824) 192 [bis]. —Ital.; Graec.
glandulifera, Jacquem. Journ. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 395 = ramulosa.
glandulosa, Wall. Cat. n. 442 = brachypoda.
glandulosa, Willd. ex Ser. in DC. Prod. iv. 46 = bronchialis.
glauca, [Clairv.] Man. Herb. 140 = diapensoides.
glaucescens, Reut. Pugill. Pl. Nov. (1852) 131 = granulata.
globulifera, *Desf. Fl. Atlant.* i. 342. t. 96.—Reg. Mediterr. occ.
globulifera, Schur, Enum. Pl. Transs. 232 = S. Aizoon.
globulifera, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 448 = Maweana.
Gmelini, Host, Fl. Austr. i. 512 = hypnoides.
gracilis, Schleich. ex Ser. in DC. Prod. iv. 31 = erosa.
gracilis, Steph. ex Sternb. Rev. Saxifr. Suppl. i. 7. t. 5. f. 1 = punctata.
gracilis, Sternb. ex Ser. in Nat. Anz. Schw. Ges. i. (1818) 74 = hypnoides.
graeca, Boiss. & Heldr. ex Boiss. Fl. Orient. ii. 807 = granulata.
granatensis, Boiss. & Reut. Pugill. Pl. Nov. (1852) 46 = globulifera.
grandiflora, Sternb. Rev. Saxifr. 20. t. 12 = sibirica.
granigera, C. Koch, in Linnaea, xix. (1847) 40 = sibirica.
granulata, Bieb. Fl. Taur. Cauc. i. 314 = sibirica.
granulata, Desf. Fl. Atlant. i. 341 = atlantica.
granulata, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1857) 63 = sibirica.
granulata, *Linn. Sp. Pl.* 403.—Europ.
granulata, Pollini, Fl. Veron. ii. 29 = bulbifera.
groenlandica, Colm. ex Costa, Fl. Catal. 92 = exarata.

SAXIFRAGA:—
groenlandica, Lapeyr. Fl. Pyr. 39. t. 19 = mixta.
groenlandica, Linn. Sp. Pl. 404 = caespitosa.
Grzegorzekii, Janka, in Oestr. Bot. Zeitschr. vi. (1856) 241 = perdurans.
Guthriana ×, Hort. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 537.
Haenseleri, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 13. —Hispan.
Hausmanni ×, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 105.—Europ.
hederacea, Bieb. Fl. Taur. Cauc. i. 317 = S. Cymbalaria.
hederacea, *Linn. Sp. Pl.* 405.—Europ. austr.; As. Min.
hederaefolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 317.—Abyss.
Hegetschweileri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 91.—Europ.
helianthemifolia, Willd. ex Sternb. Saxif. Suppl. ii. 19 = Merkii.
hemisphaerica, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 62.—Reg. Himal.
heterantha, Hook. Fl. Bor. Am. i. 252. t. 78 = Mertensiana.
heteromalla, *Georgi, Beschr. Russ. Reich.* iv. 966.—Sibir. or.
heterophylla, Sternb. Rev. Saxifr. 50 = pedemontana.
heucherifolia, Griseb. & Schenk, in Wiegm. Archiv, xxviii. (1852) 1. 317 = rotundifolia.
Heuffelii, Schott, Nym. & Kotschy, Analect. Bot. 28 = rotundifolia.
hibernica, Haw. Saxifr. 29 = hypnoides.
hibernica, Sternb. Rev. Saxifr. Suppl. ii. t. 11 = umbrosa.
hieracifolia, Porter, in Hayd. Rep. (1871) 782 = integrifolia.
hieracifolia, *Waldst. & Kit. Pl. Rar. Hung.* i. 17. t. 18.—Reg. bor. et arct.
hirculoides, Decne. in Jacquem. Voy. Bot. 67 = S. Hirculus.
hirculoides, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 112.—China.
Hirculus, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 409 = chrysantha?
Hirculus, *Linn. Sp. Pl.* 402.—Reg. bor. et arct.
hirsuta, Linn. Syst. ed. X. 1026 = umbrosa.
hirta, Donn, Hort. Cantab. ed. V. 107; Sm. Engl. Bot. t. 2291 = hypnoides.
hirta, Hook. f. in Journ. Linn. Soc. i. (1857) 117 = caespitosa.
hispanica, Coss. ex Nym. Consp. 273 = dichotoma.
hispidula, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 380.—Reg. Himal.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 146 = flagellaris.
Hohenwartii, Vest, ex Sternb. Rev. Saxifr. 26. t. 7 = sedoides.
Hornungii, *Shuttlew. in Mag. Zool. & Bot.* ii. (1838) 512.—Helvet.
Hostii, *Tausch, in Syll. Ratisb.* ii. (1828) 240.—Europ. austr.
Huetiana, *Boiss. Diagn.* Ser. II. ii. 72.—Kurdistan.
Huteri ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 147.—Europ.
hybrida, Haw. Misc. Nat. 164 = hypnoides.
hybrida, *Sternb. Rev. Saxifr.* 17. t. 8. f. 3.—Mont. Pyren.
hybrida, Vill. ex Lapeyr. Fl. Pyren. 51 = umbrosa.
hyperborea, R. Br. in Parry, 1st Voy. App. 274 = rivularis.
hypnoides, All. Fl. Pedem. ii. 72 = exarata.
hypnoides, *Linn. Sp. Pl.* 405.—Europ.
hypnoides, Scop. Fl. Carn. ed. II. i. 296 = adscendens.
hypnoides, Vitm. Sagg. Ist. Erb. Alp. 29 = caespitosa.
Idsuroei, *Franch. & Sav. Enum. Pl. Jap.* ii. 353.—Japon.
imbricata, Bertol. in Ann. Stor. Nat. iv. (1830) 260 = squarrosa.
imbricata, Lam. Fl. Fr. iii. 531 = retusa.
imbricata, *Royle, Illustr. Bot. Himal.* 226. t. 49.—Reg. Himal.
immaculata, Fl. Nicro. Danub. ex Steud. Nom. ed. II. ii. 521 = intacta.
incisata, Schur, Enum. Pl. Transs. 238 = muscoides.

SAXIFRAGA:—
inclinata ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 146.—Europ.
incrustata, Vest, in Bot. Zeit. Regensb. iii. (1804) 95, 96 = crustata.
incurva, Mack. ex Ser. in DC. Prod. iv. 32 = caespitosa.
incurvifolia, D. Don, in Trans. Linn. Soc. xiii. (1822) 423 = caespitosa.
Infundibulum, *Lange, Ind. Sem. Hort. Haun.* (1863) 5.—Hab.?
intacta, *Willd. Enum. Hort. Berol.* 459.—Europ.
integrifolia, *Hook. Fl. Bor. Am.* i. 249. t. 86.—Am. bor. occ.
integrifolia, Schleich. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 539 = muscoides.
intermedia, Gaud. Fl. Helvet. iii. 288 = exarata.
intermedia, Hegetschw. Fl. Helv. i. 288 = aspera.
intermedia, *Sweet, Hort. Brit.* ed. I. 181.—Hab.?
intermedia, Tausch, ex Link, Enum. Hort. Berol. i. 415 = hypnoides.
intricata, Lapeyr. in Bull. Soc. Bot. Fr. xii. (1865) 20 = exarata.
iratiana, F. Schultz, Arch. Fl. Fr. et Allem. (1850) 176 = mixta.
irrigua, *Bieb. Fl. Taur. Cauc.* ii. 460.—Tauria.
irrigua, Fisch. ex Spreng. Nov. Prov. 36 (Quid?).—Sibir.
Jacggiana ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxii. (1879) p. xxix.—Europ.
Jacquemontiana, *Decne. in Jacquem. Voy. Bot.* 68. t. 78. f. 2.—Reg. Himal.
Jamesiana, Walp. Rep. ii. 367 = seq.
Jamesii, *Torr. in Ann. Lyc. N. York*, ii. (1828) 204.—Am. bor. occ.
juniperifolia, *Adams, in Weber & Mohr, Beitr.* i. 53.—Reg. Caucas.
juniperina, Boiss. Fl. Orient. ii. 804, sphalm. = praec.
Kochii, Hornung, in Flora, xviii. (1835) 465 = biflora.
Kolenatiana, Regel, Ind. Sem. Hort. Petrop. (1865) 39 = cartilaginea.
Kotschyi, *Boiss. Diagn.* Ser. II. ii. 65.—As. Min.
Kruhsiana, Fisch. ex Ser. in DC. Prod. iv. 46 = nitida.
Kunzeana, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 121 = dichotoma.
lactea, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 71.—Sibir. or.
ladanifera, Balb. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 540 = pedemontana.
ladanifera, Lapeyr. Fl. Pyren. 65. t. 42 = geranioides.
laeta, Schott, Nym. & Kotschy, Analect. Bot. 24 = S. Aizoon.
laetevirens, D. Don, in Trans. Linn. Soc. xiii. (1822) 451 = hypnoides.
laevis, *Bieb. Fl. Taur. Cauc.* i. 314.—Reg. Caucas.
laevis, Donn, Hort. Cantab. ed. V. 107 = hypnoides?
laevis, Haw. Misc. 165 = hypnoides.
lanceolata, *Haw. Syn. Pl. Succ.* 324.—Europ. austr.
lantoscana, Boiss. & Reut. Diagn. Ser. II. ii. 63 = lingulata.
lanuginosa, Jacquem. ex Decne. in Jacquem. Voy. Bot. 67 = S. Hirculus.
lanuginosa, *Royle, Illustr. Bot. Himal.* 226.—Ind. or.
Lapeyrousii, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 397.—Mont. Pyren.
Lapeyrousii, Herbich, Stirp. Rar. Bucov. 35 = luteoviridis.
Lapeyrousii, Sternb. Rev. Saxifr. 51 = hypnoides.
lasiophylla, Schott, Nym. & Kotschy, Analect. Bot. 29 = rotundifolia.
lasiostigma, Pers. Syn. i. 490 = caespitosa.
latepetiolata, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 120.—Hispan.
latifida, Haw. Syn. Pl. Succ. 325; D. Don, in Trans. Linn. Soc. xiii. (1822) 420 = hypnoides.
latiflora, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 71.—Reg. Himal.
latifolia, DC. Prod. iv. 32 = hypnoides.
laurentiana, Ser. in DC. l. c. 35 = rivularis.
laxa, Schleich. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 540 = hypnoides.
lepismigena, Plan. Fl. Gallega, 224 = Clusii.
leptophylla, Froel. ex Ser. in DC. Prod. iv. 24 = aphylla.
leptophylla, Pers. Syn. i. 490 = hypnoides.

SAXIFRAGA:—
leucantha, Thomas, ex Sternb. Rev. Saxifr. Suppl. ii. 70 = exarata.
leucanthemifolia, Lapeyr. Fl. Pyr. 49. t. 25 = Clusii.
leucanthemifolia, *Michx. Fl. Bor. Am.* i. 260.—Am. bor. occ.
ligulata, Murr. in Comm. Gotting. (1781) 26. t. 1 = sarmentosa.
ligulata, *Wall. in As. Res.* xiii. (1820) 398.—Reg. Himal.
linearis, Hort. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 541 = hypnoides.
lineata, Sternb. Rev. Saxifr. Suppl. t. 7 = muscoides.
lingulata, *Bell. in Mém. Acad. Turin*, v. 1790–91 (1793) 226.—Europ. austr.
lingulata, Moretti, Tentat. 5 = intacta.
Linnaei, Boiss. Diagn. Ser. II. ii. 69 = adscendens.
longifolia, Auct. ex Nym. Consp. 268 = crustata.
longifolia, Host, Fl. Austr. i. 409 = Hostii.
longifolia, *Lapeyr. Fl. Pyren.* 26. t. 11.—Europ. austr.
longifolia, Moretti, Notiz. ed. I. 11 = intacta.
longifolia, Tenore, Prod. Fl. Nap. p. xxv = marginata.
longiscapa, D. Don, in Trans. Linn. Soc. xiii. (1822) 388 = nivalis.
lutea, Gilib. Fl. Lituan. ii. 179 = S. Hirculus.
luteopurpurea, Lapeyr. Fl. Pyren. 29. t. 14 = Lapeyrousii.
luteopurpurea, *Sternb. Rev. Saxifr.* 7; *Suppl.* ii. 48.—Transsylv.
luteopurpurea, Waldst. & Kit. ex Schult. Oestr. Fl. ed. II. i. 637 = luteoviridis.
luteoviridis, *Schott & Kotschy, ex Schott, in Oestr. Bot. Zeitschr.* ix. (1851) 65.—Transsylv.
Lyallii, *Engl. in Verh. Zool.-Bot. Ges. Wien*, xix. (1869) 542.—Am. bor. occ.
Lychnitis, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 68.—Reg. Himal.
lysimachoides, Klotzsch, in Bot. Ergeb. Waldem. Reise, 145 = diversifolia, *Wall.*
Macnabiana, Hort.—Cf. Gard. Chron. (1855) i. 705, 770 = S. Cotyledon.
macrantha, Boiss. Fl. Orient. ii. 807 = sibirica.
macropetala, *Kern. ex Engl. Monog. Saxifr.* 280.—Europ.
maculata, *Schrank, Baier. Fl.* i. 690.—Bavar.
maderensis, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 414.—Ins. Mader.
magellanica, *Poir. Encyc.* vi. 686.—Reg. Magellan.
malvaefolia, *Greene, in Bull. Torrey Bot. Club.* ix. (1882) 121.—Calif.
Malyi, Schott, Nym. & Kotschy, Analect. Bot. 23 = S. Aizoon.
marginata, Bory & Chaub. Nouv. Fl. Pélop. 26 = Boryi.
marginata, *Sternb. Rev. Saxifr. Suppl.* i. 1. t. 1. f. 1.—Italia.
marylandica, Hort. ex Sternb. l. c. i. 4 = pensylvanica.
Maweana, *Baker, in Gard. Chron.* (1871) 1355.—Marocc.
media, *Gouan, Illustr.* 27.—Mont. Pyren.
media, Sibth. & Sm. Fl. Graec. iv. 70. t. 376 = porophylla.
media, Tausch, in Flora, xii. (1829) 10 = hypnoides.
melaena, Boiss. Diagn. Ser. II. ii. 66 = mixta.
melaleuca, [*Fisch.*] *Hort. Gorenk.* ed. I. (1808) 99; *et in Spreng. Pugill.* ii. 61.—Mont. Altai.
Merkii, *Fisch. in Sternb. Rev. Saxifr. Suppl.* i. 1. t. 1. f. 3.—Sibir.
Mertensiana, *Bongard, in Mém. Acad. Pétersb.* Sér. VI. ii. (1835) 141.—Am. bor. occ.
Meyeri, *Sternb. Rev. Saxifr. Suppl.* ii. 11. t. 21.—Sibir.
micrantha, *Edgew. in Trans. Linn. Soc.* xx. (1846) 50.—Reg. Himal.
micrantha, Fisch. ex DC. Prod. iv. 48 = Leptarrhena pyrolaefolia.
micranthifolia, Steud. Nom. ed. I. 736 = erosa.
microphylla, *Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 72.—Reg. Himal.
microphylla, *Schleich. Cat. Pl. Helv.* ed. IV. 63.—Europ. centr.
Milesii, Hort. Leichtl. ex Baker, in Gard. Chron. (1882) ii. 102 = ligulata.
minor, *Sieber, ex Steud. Nom.* ed. I. 736; *et ex Engl. in Verh. Zool.-Bot. Ges. Wien*, xix. (1869) 543, nomen.—Hab.?
minuta, Koch, Syn. Fl. Germ. ed. I. 276 = tridactylites.

SAXIFRAGA:—
mixta, Bieb. Fl. Taur. Cauc. ii. 460 = exarata.
mixta, *Lapeyr. Fl. Pyren.* 41. t. 20.—Mont. Pyren.
modesta, Reichb. Ic. Fl. Germ. vii. 12. t. 626 = S. Geum.
mollis, Sm. ex Spreng. Syst. ii. 364 = sibirica.
monodactylites, Bory, ex Thore, Promen. Gascogne, 24 = tridactylites?
Moorcroftiana, Wall. Cat. n. 453 = diversifolia, *Wall.*
moschata, D. Don, in Trans. Linn. Soc. xiii. (1822) 436 = hypnoides.
moschata, Lapeyr. ex DC. Prod. iv. 28, partim = pubescens.
moschata, Wulf. in Jacq. Misc. ii. (1781) 128. t. 21. f. 2 = muscoides.
mucronulata, Royle, Illustr. Bot. Himal. 27. t. 50 f. 2 = flagellaris.
multicaulis, Lange, Ind. Sem. Hort. Haun. (1864); ex Engl. Monog. Saxifr. 228 = cuneifolia.
multifida, *Rosbach, in Bull. Soc. Bot. Belg.* xiv. (1875) 111.—Hab.?
multiflora, Ehrh. ex Ser. in DC. Prod. iv. 19 = S. Cotyledon.
multiflora, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 532.—Sibir.
multinervis, Dulac, Fl. Hautes-Pyr. 267 = androsacea.
muralis, Salisb. Prod. 306 = tridactylites.
Mureti ×, Ramb. in Verh. Schw. Nat. Ges. Chur, (1863) 191.—Europ.
Murithiana, *Tissière, in Bull. Soc. Murith.* i. (1868) 28.—Helvet.
muscoides, All. Fl. Pedem. ii. 70 = planifolia.
muscoides, Bieb. Fl. Taur. Cauc. ii. 460 = exarata.
muscoides, *Wulf. in Jacq. Coll.* ii. 123.—Europ.; As. Min.; Persia.
muscosa, Schrank, Baier. Fl. i. 700 = muscoides.
muscosa, Suter, Fl. Helvet. i. 251 = pubescens.
mutata, *Linn. Sp. Pl.* ed. II. 570.—Europ. austr.
myosotifolia, D. Don, in Trans. Linn. Soc. xiii. (1822) 373 = Merkii.
myosotifolia, Pall. ex Spreng. Syst. ii. 364 = flagellaris.
nana, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 118.—China.
neglecta, *Bray, ex Sternb. Rev. Saxifr. Suppl.* i. 9. t. 6.—Sibir.; Kamtsch.
neglecta, Tenore, Fl. Nap. Prod. App. 19 = S. Aizoon.
Nelsoniana, D. Don, in Trans. Linn. Soc. xiii. (1822) 355 = punctata.
Nelsoniana, Hook. & Arn. Bot. Beech. Voy. 124. t. 29 = Boykinia Richardsoni.
nervosa, Lapeyr. Fl. Pyren. 63. t. 39 = exarata.
nervosa, *Miégev. in Bull. Soc. Bot. Fr.* xii. (1865) 19.—Mont. Pyren.
nevadensis, Boiss. Diagn. Ser. II. ii. 67 = mixta.
nigricans, Willd. ex Sternb. Rev. Saxifr. Suppl. ii. 46 = nivalis.
nitida, Ledeb. Fl. Ross. ii. 207 = bronchialis.
nitida, *Schreb. ex Sternb. Rev. Saxifr.* 32. t. 10. f. 4.—Hab.?
nivalis, *Linn. Sp. Pl.* 401.—Reg. bor. et arct.
nivalis, Muhl. Cat. 44 = virginiensis.
nivalis, Pall. Reise, iii. 444 = davurica.
norica ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 147.—Europ.
notata, Schott, Nym. & Kotschy, Analect. Bot. 22 = S. Aizoon.
notkana, Moç. ex Ser. in DC. Prod. iv. 40 = stellaris?
nuda, Ser. l. c. 41 = erosa.
nudicaulis, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 366.—Am. bor. occ.
nutans, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 242 = S. Hirculus.
nutans, D. Don, in Trans. Linn. Soc. xiii. (1822) 368 = Romanzoffia unalaschcensis *vel* sitchensis.
nutans, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 69.—Reg. Himal.
obscura, *Gren. & Godr. Fl. Fr.* i. 646.—Mont. Pyren.
obtusifida, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 416.—Hispan.
obtusifolia, *Schleich. Cat. Pl. Helvet.* ed. IV. 61.—Helvet.
odontophylla, *Wall. Cat.* n. 454.—Reg. Himal.
odontophylla, Wall. l. c., partim = sibirica.
officinarum, Crantz, Inst. ii. 426 = granulata.
olympica, Boiss. Diagn. Ser. I. iii. 19 = rotundifolia.
oppositifolia, *Linn. Sp. Pl.* 402.—Reg. bor. et arct.

SAXIFRAGA:—
oranensis, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Alger.
orientalis, Jacq. Obs. ii. 9. t. 34 = S. Cymbalaria.
Pacumbis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 209 = ligulata.
Padellae ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 85.—Europ.
Pallasiana, Sternb. Rev. Saxifr. Suppl. ii. 19 = serpyllifolia.
pallida, *Wall. Cat.* n. 450.—Reg. Himal.
palmata, Lapeyr. Fl. Pyren. 64. t. 41 = geranioides.
palmata, Sm. Engl. Bot. t. 455 = hypnoides.
palmata, Steph. ex Sternb. Rev. Saxifr. Suppl. i. 8. t. 6. f. 2 = sibirica.
palpebrata, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 67.—Reg. Himal.
palustris, Link, Enum. Hort. Berol. i. 412 = pensylvanica.
palustris, Salisb. Prod. 306 = S. Hirculus.
paniculata, *Cav. Desc.* 473; *Willk. & Lange, Prod. Fl. Hisp.* iii. 112.—Hispan.
paniculata, Hornem. Hort. Hafn. i. 402 = trifurcata.
paniculata, Mill. Gard. Dict. ed. VIII. n. 3 = S. Cotyledon.
paniculata, Schleich. ex Ser. in DC. Prod. iv. 30 = geranioides.
paniculata, Willk. in Bot. Zeit. v. (1847) 431 = granulata.
paradoxa, *Sternb. Rev. Saxifr.* 22. n. 35. t. 14.—Carinth.
Parisii, Pomel, Nouv. Mat. Fl. Atl. 154 = atlantica?
parnassica, Boiss. & Heldr. Diagn. Ser. II. ii. 19 = adscendens.
parnassifolia, D. Don, in Trans. Linn. Soc. xiii. (1822) 405 = diversifolia, *Wall.*
parnassifolia, Wall. Cat. n. 451, partim = corymbosa, *Hook. f. & Thoms.*
parnassiodes, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 27.—Turkestan.
Parryi, *Torr. Bot. Mex. Bound.* 69.—Am. bor. occ.
parvicalyx, Jacquem. ex Decne. in Jacquem. Voy. Bot. 68 = Jacquemontiana.
parviflora, Bivona, Stirp. Rar. Sicil. Manip. iv. 11. t. 3 = hederacea.
parviflora, Fisch. ex Cham. in Linnaea, vi. (1831) 556 = Leptarrhena pyrolaefolia.
patens, Gaud. in Nat. Anz. Schw. Ges. i. (1818) 65 = caesia.
patens ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 146.—Europ.
patens, Sternb. Rev. Saxifr. Suppl. ii. t. 12 = valdensis.
pauciflora, Sternb. l. c. Suppl. i. 6. t. 4 = nivalis.
Pavonii, *D. Don, in Trans. Linn. Soc.* xiii. (1822) 434.—Peruv.
pectinata, Pursh, Fl. Am. Sept. i. 312 = Spiraea pectinata.
pectinata, Schott, Nym. & Kotschy, Analect. Bot. 21 = S. Aizoon.
pedatifida, *Ehrh. ex Sm. in Trans. Linn. Soc.* x. (1811) 340.—Europ. austr.
pedemontana, *All. Fl. Pedem.* ii. 73.—Europ. austr.
pedemontana, Willd. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 547 = aquatica.
pekinensis, *Maxim. Ind. Pekin. in Prim. Fl. Amur. Suppl.* 120.—China.
peltata, *Torr. ex Benth. Pl. Hartw.* 311.—Calif.
penduliflora, Bast. in Desv. Journ. Bot. iii. (1814) 17 = granulata.
pensylvanica, *Linn. Sp. Pl.* 399.—Am. bor.
pentadactylis, *Lapeyr. Fl. Pyren.* 64. t. 40.—Mont. Pyren.
perdurans, *Kit. ex Wahlenb. Fl. Carpat.* 123.—Transsylv.
perporosa, Schur, Enum. Pl. Transs. 234 = retusa.
perpusilla, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 72.—Reg. Himal.
peruviana, *Sternb. Rev. Saxifr.* 55. t. 22.—Peruv.
petiolaris, R. Br. in Ross, Voy. ed. II. ii. 192 = rivularis.
petraea, Gouan, Illustr. 29. t. 18. f. 3 = aquatica.
petraea, Gunn. Fl. Norveg. ii. 30. n. 427 = controversa.
petraea, *Linn. Syst.* ed. X. 1027.—Europ. austr.
petraea, Pall. Ind. Taur. 147 = irrigua.
petraea, *Prey. & Zirk. ex Bab. in Journ. Linn. Soc.* xi. (1870) 308.—Islandia.

SAXIFRAGA:—
petraea, Roth, Tent. Fl. Germ. i. 184 = hypnoides.
petraea, Willd. Sp. Pl. ii. 654 = adscendens.
petrosa, Vahl, ex Steud. Nom. ed. II. ii. 522 = petraea, *Linn.*
pilifera, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 66.—Reg. Himal.
pilosa, Haw. Misc. Nat. 157 = virginiensis.
planifolia, Lapeyr. Fl. Pyren. 31 = muscoides.
planifolia, Sternb. ex Cham. in Linnaea, vi. (1831) 555 = serpyllifolia.
planipetala, D. Dietr. Syn. Pl. ii. 1535 = hypnoides.
plantaginifolia, Hook. in Parry's 2nd Voy. Suppl. 394 = hieracifolia.
platypetala, Sm. in Trans. Linn. Soc. x. (1811) 391 = hypnoides.
polita, Hort. ex Ser. in DC. Prod. iv. 42 = umbrosa.
polita, Link, Enum. Hort. Berol. i. 414 = hypnoides.
polydactyla, Pall. Reise, iii. 320 = davurica.
Ponae, Sternb. Rev. Saxifr. 47. t. 11. f. 6 = petraea, *Linn.*
porophylla, Bertol. in Desv. Journ. Bot. ii. (1813) 76. —Europ. austr.
porphyrantha, Hort. Gotting. ex Engl. Monog. Saxifr. 176 = muscoides.
portosanctana, *Boiss. Diagn.* Ser. II. ii. 68.—Ins. Mader.
praecox, Bernh. ex Engl. Monog. Saxifr. 84 = tridactilites.
primulaefolia, Willd. ex Sternb. Rev. Saxifr. Suppl. ii. 46 = nivalis.
procumbens, Juss. ex Steud. Nom. ed. II. ii. 522 = hypnoides.
propaginea, Pourr. ex Engl. Monog. Saxifr. 134 = Clusii.
propinqua, R. Br. in Ross, Voy. ed. II. ii. 192 = S. Hirculus.
prorepens, Fisch. ex Sternb. Rev. Saxifr. 59 = flagellaris.
Prostiana, Nym. Consp. 270 = mixta.
Prostii, Sternb. Rev. Saxifr. Suppl. 34 = pedatifida.
protensa, *Schleich. Cat. Pl. Helv.* ed. IV. 63.—Helvet.
Pseudo-adscendens, Schur, Enum. Pl. Transs. 238 = tridactylites.
Pseudo-bronchialis, *Fisch. Hort. Gorenk.* ed. II (1812) 61.—Hab.?
Pseudo-Burseriana, Fisch. ex Cham. in Linnaea, vi. (1831) 555 = bronchialis.
Pseudo-caesia, Rochel, Pl. Banat. Rar. 35. t. 3 = Rocheliana.
Pseudo-hypnoides, Murith, Bot. Val. 94 = exarata.
Pseudo sancta, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 176.—Balkan.
pubescens, DC. Fl. Fr. iv. 375 = mixta.
pubescens, *Pourr. in Mém. Acad. Toul.* iii. (1788) 327. —Europ.
pubescens, Sternb. in Linnaea, vi. (1831) 556 = caespitosa.
pulchella, D. Don, in Trans. Linn. Soc. xiii. (1822) 426 = exarata.
punctata, Gunn. Fl. Norveg. ii. 140 = cuneifolia.
punctata, Haw. ex Nym. Consp. 294 = umbrosa.
punctata, Hook. Lond. Journ. Bot. vi. (1847) 231 = Mertensiana.
punctata, *Linn. Sp. Pl.* 401.—Sibir.; Am. bor. occ.
punctata, Pall. Reise, iii. 731 = davurica.
punctata, Rottb. in Skrift. Kjoeb. Selsk. x. (1770) 445. t. 3. f. 10 = nivalis.
pungens, [Clairv.] Man. Herb. 140 = Vandellii.
purpurascens, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 61.—Reg. Himal.
purpurea, All. Fl. Pedem. ii. 71 = retusa.
pusilla, *Sternb. Rev. Saxifr. Suppl.* ii. 35.—Ins. Aleut.
pygmaea, Haw. Misc. Nat. 168 = muscoides.
pyramidalis, Lapeyr. Fl. Pyren. 32 = S. Cotyledon.
pyramidalis, Salisb. Prod. 305 = S. Aizoon.
pyramidalis, Tenore, Prod. Fl. Nap. p. xxv = lingulata.
pyramidata, Mill. Gard. Dict. ed. VIII. n. 4 = S. Cotyledon.
pyrenaica, Scop. Fl. Carn. ed. II. i. 296 = androsacea.
pyrenaica, Vill. Hist. Pl. Dauph. iii. 371 = muscoides.
pyrolifolia, D. Don, in Trans. Linn. Soc. xiii. (1822) 389 = Leptarrhena pyrolaefolia.
quinquedens, Haw Syn. Pl. Succ. 325 = hypnoides.
quinquefida, Haw. Misc. Nat. 163 = hypnoides.
quinquefida, Lam. Fl. Fr. iii. 533 = geranioides.
radicans, Jacquem. Voy. Bot. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 397 = flagellaris.

SAXIFRAGA:—
ramosissima, Schur, Enum. Pl. Transs. 239 = adscendens.
ramulosa, *Wall. Cat.* n. 446.—Reg. Himal.
ranunculifolia, *Hook. Fl. Bor. Am.* i. 246. t. 83.—Am. bor.
ranunculoides, Haw. Enum. Saxifr. 25 = irrigua.
recta, Lapeyr. Fl. Pyren. 33. t. 15 = S. Aizoon.
recta, Tenore, Prod. Fl. Nap. p. xxv = marginata.
recurva, Bree, ex Schleich. Cat. Pl. Helv. ed. IV. 63 = hypnoides.
recurvifolia, Lapeyr. Fl. Pyren. 30 = caesia.
Redofskyi, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 241 = stellaris.
Redowskyana, Presl, ex Ledeb. Fl. Ross. ii. 212 = davurica.
reflexa, *Hook. Fl. Bor. Am.* i. 249. t. 85.—Am. bor.
reflexa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Hirculus.
Regelii, *Kern. in Oestr. Bot. Zeitschr.* xiii. (1863) 107. —Europ.
repanda, Willd. ex Sternb. Rev. Saxifr. 17. t. 5 = rotundifolia.
reticulata, Cham. in Linnaea, vi. (1831) 555 = Sieversiana.
reticulata, Willd. ex Sternb. Rev. Saxifr. 21. t. 13 = S. Cymbalaria.
retroflexa, Hort. ex Ser. in DC. Prod. iv. 31 = hypnoides.
retusa, Cham. in Linnaea, vi. (1831) 556 = oppositifolia.
retusa, *Gouan, Illustr.* 28. t. 18. f. 1.—Europ. austr.
Reuteriana, *Boiss. Voy. Bot. Espagne*, ii. 730.—Hispan.
Rhei, Schott, Nym. & Kotschy, Analect. Bot. 30 = muscoides.
Richardsonii, Hook. Fl. Bor. Am. i. 247 = Boykinia Richardsoni.
rigens, Kit. in Linnaea, xxxii. (1863) 474 = Rocheliana.
rigida, C. Sm. ex Ser. in DC. Prod. iv. 39 = hieracifolia.
rivalis, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 372 = seq.
rivularis, *Linn. Sp. Pl.* 404.—Reg. bor. et arct.
rivularis, Schult. Oestr. Fl. ed. II. i. 640 = carpathica.
rivularis, Thomas, ex DC. Prod. iv. 35 = granulata.
rivularis, S. Wats. in Bot. King's Exped. v. 93 = debilis.
robusta, *Schott, Nym. & Kotschy, Analect. Bot.* 22 = S. Aizoon.
Rocheliana, *Sternb. Rev. Saxifr. Suppl.* ii. 55.—Banatus.
rorida, Bieb. ex Ledeb. Fl. Ross. ii. 218 = irrigua.
rosacea, Moench, Meth. 106 = hypnoides.
rosea, Lapeyr. ex Steud. Nom. ed. II. ii. 523 = biflora.
Rossii, *Oliver, in Hook. Ic. Pl.* t. 1258.—Manchuria.
rosularis, Haw. Misc. Nat. 159 = S. Aizoon.
rosulenta, Ehrh. ex Ser. in DC. Prod. iv. 19 = S. Aizoon.
rotundifolia, Geners. Elench. Scepus. n. 372; ex Wahlenb. Fl. Carpat. 119 = hieracifolia.
rotundifolia, *Linn. Sp. Pl.* 403.—Graec.; Reg. Caucas.
rotundifolia, Pall. ex Georgi, Beschr. Russ. Reich. iv. 964 = punctata.
rotundifolia, Sibth. & Sm. Fl. Graec. iv. 71. t. 377 = chrysosplenifolia.
Rudolphiana, Hornsch. ex Koch, Syn. Fl. Germ. ed. I. 269 = oppositifolia.
rupestris, Lapeyr. Hist. Abr. Pl. Pyr. i. 233 = adscendens.
rupestris, Salisb. Prod. 306; Schleich. ex Ser. in DC. Prod. iv. 31 = hypnoides.
rupestris, Thomas, ex Ser. l. c. 22 = valdensis.
rupestris, Willd. Sp. Pl. ii. 653 = petraea, *Linn.*
Russi, Presl, Delic. Prag. 140 = bulbifera.
sabulicola, Pomel, Nouv. Mat. Fl. Atl. 153 = atlantica.
sachalinensis, *F. Schmidt, Reisen Amurl.* 133.—Ins. Sachalin.
saginoides, Froel. ex Steud. Nom. ed. II. ii. 523 = sedoides.
saginoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 68.—Reg. Himal.
sancta, *Griseb. Spicil. Fl. Rumel.* i. 332.—Graecia; As. Min.
sarmentosa, *Linn. f. Suppl.* 240.—China; Japon.
scapigera, Willd. ex Sternb. Rev. Saxif. 59; Suppl. ii. 19 = serpyllifolia.

SAXIFRAGA:—
scardica, *Griseb. Spicil. Fl. Rumel.* i. 332.—Graecia.
scariosa, Schleich. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 551 = hypnoides.
Schleicheri, D. Don, ex G. Don, in Loud. Hort. Brit. 176 = stellaris.
Schmidti, Regel, Gartenfl. (1878) 225 = ligulata?
Schraderi, Sternb. Rev. Saxifr. Suppl. 11. t. 8 = hypnoides.
Scopolii, Vill. Prosp. 47; Hist. Pl. Dauph. iii. 670 = adscendens.
scotophila, Boiss. Diagn. Ser. I. iii. 23 = hederacea.
scrobiculata, Schur, ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 551 = retusa.
secunda, Moretti, in Brugnat. Giorn. Fis. II. viii. (1825) 285 = muscoides.
secunda, Willd. ex Cham. in Linnaea, ii. (1827) 608 = Romanzoffia unalaschcensis, sitchensis.
sednensis, Clarion, ex Sternb. Rev. Saxifr. 53 = lingulata.
sedoides, *Linn. Sp. Pl.* 404.—Europ. austr.
sedoides, Wahlenb. Veg. Helvet. 79, partim; Murith, Bot. Val. 94 = Seguieri.
Seguieri, *Spreng. Fl. Hal. Mant.* 40.—Helvet.
semidecandra, Fisch. ex DC. Prod. iv. 42 = umbrosa.
semidodecandra, Walp. Rep. ii. 364 = punctata.
semipubescens, Sweet, Hort. Suburb. Lond. 97 = pensylvanica.
Sempervivum, C. Koch, in Linnaea, xix. (1847) 40 = porophylla.
sendaica, *Maxim. in Bull, Acad. Pétersb.* xviii. (1873) 38.—Japon.
septifida, *Haw. ex Steud. Nom.* ed. II. ii. 523 (Quid?). —Hibernia.
serpyllifolia, A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864), 62 = chrysantha.
serpyllifola, *Pursh, Fl. Am. Sept.* i. 311.—Reg. bor. et arct.
serrata, Sternb. Rev. Saxifr. Suppl. ii. 5 = umbrosa.
serratifolia, Mackay, ex Reichb. in Moessl. Handb. ed. II. i. 709 = umbrosa.
setigera, Pursh, Fl. Am. Sept. i. 312 = flagellaris.
sibirica, *Linn. Syst.* ed. X. 1027.—As. bor. et arct.
sibirica, Pall. ex Ledeb. Fl. Ross. ii. 212 = davurica.
sibirica, Pursh, Fl. Am. Sept. i. 312 = cernua?
sibirica, Sternb. Rev. Saxif. 23 = Sieversiana.
sibirica, Wahlenb. Fl. Carpat. 121 = carpathica.
Sibthorpii, *Boiss. Diagn.* Ser. I. iii. 22.—Graecia.
Sieversiana, *Sternb. Rev. Saxifr. Suppl.* ii. 16.—Sibir.
sileniflora, *Hook. Fl. Bor. Am.* i. 245.—Am. bor. occ.
silenifolia, Engl. Monog. Saxif. 172, sphalm. = praec.
sobolifera, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 243 = flagellaris.
spathularis, Brot. Fl. Lusit. 172 = umbrosa.
spathulata, Boiss. Voy. Bot. Espagne, ii. 231 = erioblasta.
spathulata, *Desf. Fl. Atlant.* i. 342. t. 96. f. 2.—Afr. bor.
spatulata, Haw. Misc. Nat. 165 = hypnoides.
sphaeroidea, Sweet, Hort. Brit. ed. 1. 181 = umbrosa.
spicata, D. Don, in Trans. Linn. Soc. xiii. (1822) 354 = punctata.
spinulosa, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 96 = bronchialis.
spinulosa, Royle, Illustr. Bot. Himal. 227. t. 50. f. 2 = flagellaris.
sponhemica, C. C. Gmel. Fl. Bad. ii. 224 = hypnoides.
Spruneri, *Boiss. Diagn.* Ser. I. iii. 18.—Thessal.
spuria ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 146. —Europ.
squarrosa, *Sieber, in Flora*, iv. (1821) 99.—Tyrol.
Stabiana, Tenore, Syll. Fl. Neap. 201 = S. Aizoon.
Steinmanni, Tausch, Hort. Canal. dec. i. 76; et in Flora, vii. (1824) 85 = hypnoides.
Stella-aurea, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 72.—Reg. Himal.
stellaris, *Linn. Sp. Pl.* 400.—Reg. bor. et arct.
stellata, *Pav. ex D. Don, in Trans. Linn. Soc.* xiii. (1822) 430.—Peruv.
Stelleriana, *Merk. ex Ser. in DC. Prod.* iv. 46.—Sibir.
stenoglossa, *Tausch, ex Schott, Nym. & Kotschy, Analect. Bot.* 22.—Europ.
stenopetala, Gaud. in Nat. Anz. Schw. Ges. i. (1818) 70 = aphylla.
stenophylla, Royle, Illustr. Bot. Himal. 227. t. 50 f. 1 = flagellaris.

SAXIFRAGA:—
Stephaniana, Sternb. Rev. Saxifr. Suppl. i. 8. t. 6 = sibirica.
Sternbergii, Willd. Enum. Hort. Berol. 462 = caespitosa.
stolonifera, Meerb. Afbeeld. t. 23. (1775); Jacq. Misc. ii. 327 = sarmentosa.
Stracheyi, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 61.—Reg. Himal.
striata, Hall. f. ex Gaud. Fl. Helv. iii. 129 = muscoides.
stricta, C. Sm. ex Nym. Consp. 274 = hieracifolia.
strigosa, *Wall. Cat.* n. 448.—Reg. Himal.
Sturmiana, Schott, Nym. & Kotschy, Analect. Bot. 25 = S. Aizoon.
stylosa, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 236.—Bolivia.
subacaulis, Hegetsch. Fl. Schw. 397 = muscoides.
submutica, Hort. ex Steud. Nom. ed. II. ii. 523 = S. Aizoon.
subverticillata, *Boiss. Fl. Orient.* ii. 803.—Reg. Caucas.
Sullivantii, Torr. & Gray, Fl. N. Am. i. 575 = Sullivantia Ohionis.
tangutica, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 114.—China.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. x. 19.—Graecia.
tecta, Kit. in Linnaea, xxii. (1863) 473 = luteoviridis.
tellimoides, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 216.—Japon.
tenella, *Wulf. in Jacq. Coll.* iii. 144. t. 17.—Europ. austr.
tenera, Sternb. Rev. Saxifr. t. 9 B = androsacea.
tenera, Suter, Fl. Helvet. i. 245 = planifolia.
tenerrima, Willk. in Oestr. Bot. Zeitschr. xxv. (1875) 111 = tridactylites.
Tenorii, *Sternb. Rev. Saxifr. Suppl.* ii. 69. t. 14.—Europ. austr.
terektensis, *Bunge, in Ledeb. Fl. Alt.* ii. 123.—Sibir. altaic.
texana, *Buckl. in Proc. Acad. Sc. Philad.* 1861 (1862) 455.—Am. bor. occ.
thessalica, Schott, Nym. & Kotschy, Analect. 26, in obs. = porophylla.
thyrsoidea, *Tausch, in Syll. Ratisb.* ii. (1828) 240.—Italia.
thysanodes, Lindl. Bot. Reg. (1846) t. 33 = ligulata.
Tilingiana, *Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 94.—As. or.
tiroliensis ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 146.—Europ.
Tolmiei, *Torr. & Gray, Fl. N. Am.* i. 567 (Tolmaei).—Am. bor. occ.
tombeanensis, Boiss. ex Engl. in Verh. Bot.-Zool. Ges. Wien, xix. (1869) 554 = diapensioides.
transsilvanica, *Fuss, Fl. Transs.* 236.—Transsylv.
trichodes, Scop. Fl. Carn. ed II. i. 295 = sedoides.
tricuspidata, *Rottb. in Skrift. Kioeb. Selsk.* x. (1770) 446.—Reg. bor. et arct.
tridactylites, *Linn. Sp. Pl.* 404.—Europ.; Afr. bor.; Oriens.
tridens, Haw. Syn. Pl. Succ. 324 = hypnoides?
tridens, Jan, ex DC. Prod. iv. 24 = androsacea.
tridentata, D. Don, in Trans. Linn. Soc. xiii. (1822) 426 = hypnoides.
trifida, Gilib. Fl. Lituan. ii. 181 = tridactylites.
trifida, Haw. Misc. Nat. 165 = hypnoides.
trifida, *Vand. Fasc. Pl.* 13 (Quid?).—Ital.
trifurcata, *Schrad. Hort. Gotting.* fasc. i. 13. t. 7.—Hispan.
trigyna, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 235.—Chili.
tuberosa, Sternb. Rev. Saxifr. Suppl. ii. 9 = punctata.
uliginosa, Fisch. ex Ser. in DC. Prod. iv. 39 = hieracifolia.
umbellulata, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 71.—Reg. Himal.
umbrosa, *Linn. Sp. Pl.* ed. II. 574.—Europ. occ.
unalaschcensis, Fisch. ex Sternb. Rev. Saxifr. Suppl. i. 2 = Leptarrhena pyrolaefolia.
unalaschcensis, Presl, ex Ledeb. Fl. Ross. ii. 212 = davurica.
unguiculata, *Engl. in Bull. Acad. Pétersb.* xxix. (1883) 115.—China.
uniflora, R. Br. in Parry, 1st Voy. App. 275 = caespitosa.
uniflora, Froel. ex DC. Prod. iv. 24 = aphylla.
uniflora, Sternb. Rev. Saxifr. Suppl. 13. t. 9. f. 1 = hypnoides.

SAXIFRAGA:—
vaginalis, Turcz. ex Ledeb. Fl. Ross. ii. 220 = neglecta.
vaginata, Presl, ex Sternb. Rev. Saxifr. Suppl. ii. 39 = rivularis.
Vahlii, Ram. ex DC. Fl. Fr. iv. 370 = adscendens.
valdensis, *DC. l. c. Suppl.* 517.—Europ. austr.
Vandellii, Lapeyr. Hist. Abr. Pl. Pyr. 636 = arctioides.
Vandellii, *Sternb. Rev. Saxifr.* 34. t. 10.—Europ. austr.
varians, Sieber, ex Ser. in DC. Prod. iv. 25 = muscoides.
venosa, Haw. Enum Saxifr. 28 = caespitosa.
verna, Hort. ex Steud. Nom. ed. II. ii. 524 = virginiensis.
vernalis, Willd. Hort. Berol. i. 43 = virginiensis.
veronicaefolia, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 325 = atlantica.
veronicaefolia, Pers. Syn. i. 489 = bulbifera.
villosa, Willd. Enum. Hort. Berol. 462 = hypnoides.
virginiana, Poit. in Rev. Hortic. (1879) 260 = virginiensis.
virginica, Nutt. Gen. Am. i. 285 = seq.
virginiensis, *Michx. Fl. Bor. Am.* i. 269.—Am. bor.
viscidula, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1857) 69.—Reg. Himal.
viscosa, Hort. ex Haw. Misc. Nat. 164 = hypnoides?
vivipara, Vest, in Flora, iii. (1820) 405 = bulbifera.
Wahlenbergii, Ball, in Bot. Zeit. iv. (1846) 401 = perdurans.
Wallacei, McNab.—Cf. Gard. Chron. (1883) i. 670 = Camposii.
Wallichiana, Sternb. Rev. Saxifr. Suppl. i. t. 22 = fimbriata.
Wettsteinii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 87.—Europ.
Wildiana, *Kunze, in Linnaea*, xvii. (1843) 575.—Hab.?
Willkommiana, Boiss. ex Engl. in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 556 = Camposii.
Wolleana, Torr. & Gray, Fl. N. Am. i. 569 = erosa.
Wormskioldii, Fisch. ex DC. Prod. iv. 42 = umbrosa.
Wulfeniana, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 126 = retusa.
Zeleborii, *Schott, l. c.* 246.—Europ. austr.
Zimmeteri ×, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 146.—Europ.

SAXIFRAGIA, Tourn. ex Linn. Syst. ed. I (1735) = **Saxifraga**, Linn.

SAXO-FRIDERICIA, Rob. Schomb. Rapat. Frid.-Aug. Braunschw. (1845) 13. t. 2. *RAPATEACEAE*, Benth. & Hook. f. iii. 858.
aculeata, *Koern. in Linnaea*, xxxvii. (1871–73) 457.—Guian.
regalis, *Rob. Schomb. Rapat. Frid.-Aug. Braunschw.* (1845) 14. t. 2.—Guian.
subcordata, *Koern. in Linnaea*, xxxvii. (1871–73) 459.—Bras.

SCABIOSA, [Tourn.] Linn. Syst. ed. I (1735). *DIPSACEAE*, Benth. & Hook. f. ii. 159.
Anisodens, Dulac, Fl. Hautes-Pyr. 465 (1867).
Asterocephalus, Lag. Gen. et Sp. Nov. 8 (1816).
Astrocephalus, Euptilia, Gonokeros, Lomelosia, Pentena, Sixalix, Tereiphas, Thlasidia, Tremastelma, Xetola, Rafin. Fl. Tellur. iv. 95, 96 (1836).
Callistemma, Boiss. Fl. Orient. iii. 146 (1875).
Chetaria, Steud. Nom. ed. II. ii. 521 (1841).
Chetastrum, Neck. Elem. i. 712 (1790).
Columbaria, J. & C. Presl, Fl. Čech. 32 (1819).
Cyrtostemma, Spach, Hist. Vég. Phan. x. 321 (1841).
Diototheca, Rafin. Fl. Ludov. 74 (1817).
Knautia, Linn. Gen. ed. I. 23 (1737).
Pterocephalus, Vaill. ex Adans. Fam. ii. 152 (1763).
Pycnocomon, Hoffmgg. & Link, Fl. Port. ii. 93. t. 88 (1820).
Sclerostemma, Schott, ex Roem. & Schult. Syst. iii. 64 (1818).
Spongostemma, Reichb. Nom. 72 (1841).
Succisa, Neck. Elem. i. 109 (1790).
Trichera, Schrad. Cat. Sem. Hort. Gotting. (1814); ex Roem. & Schult. Syst. iii. 54 (1819).
Trochocephalus, Opiz, ex Bercht. Oek. Fl. Poehm. ii. 222 (1838).
acaulis, Thunb. Prod. Pl. Cap. 29 = S. Columbaria.
achaeta, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1863) 465.—Serbia.
acutiflora, Reichb. Pl. Crit. iv. 24. t. 326 = maritima.
affinis, Gren. & Godr. Fl. Fr. ii. 78 = S. Columbaria.
afghanica, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 67.—Afghan.
africana, *Linn. Sp. Pl.* ed. II. 145.—Afr. austr.
africana, Sibth. & Sm. Fl. Graec. Prod. i. 83 = cretica.
agrestis, Láng, ex Nym. Consp. 344 = Scopolii.
agrestis, F. W. Schmidt, Fl. Boëm. iii. 77 = arvensis.
agrestis, Waldst. & Kit. Pl. Rar. Hung. ii. 226 = gramuntia.
alba, Scop. Delic. Insub. iii. 33 = ucranica.
albescens, Willd. Enum. Hort. Berol. 144 = Cephalaria leucantha.
aleppica, Coult. Mem. Dipsac. 35. t. 2. f. 8 = palaestina.
alpestris, Jord. Pugill. Pl. Nov. 84 = lucida.
alpestris, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 536.—Sibir.
alpigena, Schur, Enum. Pl. Transs. 962 (Index) = arvensis.
alpina, Georgi, Beschr. Russ. Reich. Nachtr. 259 = Cephalaria tatarica.
alpina, Linn. Sp. Pl. 98 = Cephalaria alpina.
alpina, Sibth. & Sm. Fl. Graec. Prod. i. 79 = Cephalaria ambrosioides.
altissima, Jacq. Hort. Vindob. ii. 87 = africana.
altissima, Mill. Gard. Dict. ed. VIII. n. 6 = Cephalaria tatarica.
ambigua, Avé-Lall. Diss. Pl. Quib. 8 = pyrenaica.
ambigua, *Frivald. in Flora*, xix. (1836) 438.—Macedon.
ambigua, Tenore, Fl. Neap. App. v. 7 = maritima.
ambrosioides, Sibth. & Sm. Fl. Graec. Prod. i. 80; Fl. Graec. ii. 5 t. 103 = Cephalaria ambrosioides.
amoena, *Jacq. f. Eclog.* i. 86. t. 59.—As. Min.; Persia.
amplexicaulis, *Linn. Mant.* ii. 195.—Hab.?
anatolica, Boiss. Diagn. Ser. I. x. 82 = rotata.
anethifolia, Steud. Nom. ed. II. ii. 524, sphalm. = S. Columbaria.
angulata, Rafin. Caratteri, 81 = maritima.
angustifolia, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 206.—Hab.?
angustifolia, Salisb. Prod. 54 = Cephalaria syriaca.
anthemifolia, Eckl. & Zeyh. Enum. 371 = S. Columbaria.
arenaria, *Forsk. Fl. Aegypt. Arab.* p. lxi.—Aegypt.; Arab.
argentea, Boiss. Voy. Bot. Espagne, ii. 296 = sicula.
argentea, Linn. Sp. Pl. 100 = ucranica.
arvensis, *Linn. l. c.* 99.—Europ.; Reg. Caucas.
arvensis, Lenore, Syll. Fl. Neap. 59 = integrifolia.
arvensis, Ucria, ex DC. Prod. iv. 657 = maritima.
asterocephala, Thuill. Fl. Par. ed. II. 72 = suaveolens.
atrata, Lag. Elench. Hort. Madrit. (1805); ex ej. Gen. et Sp. Nov. 7. n. 103 = Cephalaria tatarica.
atropurpurea, Linn. Sp. Pl. 100 = maritima.
attenuata, Linn. f. Suppl. 118 = Cephalaria attenuata.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 111.—Syria; Assyria.
australis, *Wulf. in Roem. Arch.* iii. iii. (1803) 316.—Europ. austr.
baetica, *Boiss. Elench. Pl. Nov. Hisp.* 58.—Hispan.
Bailleti, Timb.—Cf. Bull. Soc. Bot. Fr. xv. (1868) p. xviii = maritima.
Balansae, Reut. Cat. Hort. Genev.; ex Boiss. Fl. Orient. iii. 145 = palaestina.
banatica, *Waldst. & Kit. Pl. Rar. Hung.* i. 10.—Bannat.
bellidifolia, Lam. Fl. Fr. iii. 347 = arvensis.
bellidis-folia, Gilib. Fl. Lituan. i. 169 = S. Columbaria.
bicolor, *Kotschy, ex Boiss. Fl. Orient.* iii. 143.—Kurdistan.
bidens, *Sibth. & Sm. Fl. Graec. Prod.* i. 80; *Fl. Graec.* ii. 6. t. 104.—As. Min.
Biebersteinii, Roem. & Schult. Syst. iii. 75 = micrantha.
bipinnata, C. Koch, in Linnaea, xvii. (1843) 37 = ochroleuca.

SCABIOSA :—

bipinnata, Nym. Consp. 343 = stellata.
bohemica, F. W. Schmidt, Fl. Boëm. iii. 79 = arvensis.
borealis, Salisb. Prod. 54 = S. Succisa.
brachiata, *Sibth. & Sm. Fl. Graec. Prod.* i. 83 ; *Fl. Graec.* ii. 9. t. 109.—Europ. austr. ; As. Min.
brachycarpa, *Boiss. & Hohen. Diagn.* Ser. I. ii. 113. —Kurdistan.
brevicoma, Nym. Consp. 343 = stellata.
brevis, D. Dietr. Syn. Pl. i. 448 = papposa.
breviscapa, Boiss. & Heldr. Diagn. Ser. I. vi. 73 = crenata.
breviseta, Jord. Pugill. Pl. Nov. 101 = gramuntia.
brigantiaca, Jord. l. c. 97. = S. Columbaria.
Broussonetii, *D. Dietr. Syn. Pl.* i. 448.—Hispan.
Brunoniana, Wall. Cat. n. 429 = Dipsacus Leschenaultii.
Buekiana, *Eckl. & Zeyh. Enum.* 371.—Afr. austr.
caerulea, Kielm. Diss. Pl. Hort. Tuebing. (1814) 16 ; Schult. Mant. iii. 43 = caucasica.
caerulea, *Gueldenst. Reisen*, ii. 10 (Quid ?).—Rossia.
caerulea, Hort. ex Steud. Nom. ed. II. ii. 525 = S. Columbaria.
caesaraugustana, *Echeand. Fl. Caesaraug.* 25.—Hispan.
calocephala, Boiss. Diagn. Ser. I. x. 80 = palaestina.
calyptocarpa, St. Amans, Fl. Agen. 60 = maritima.
camelorum, *Coss. & Dur. Bull. Soc. Bot. Fr.* ii. (1855) 251.—Algeria.
campestris, Andrz. ex Bess. Enum. Pl. Volh. 7 = arvensis.
cana, *D. Dietr. Syn. Pl.* i. 448.—Oriens.
candicans, Jord. Pugill. Pl. Nov. 99 = pyrenaica.
Candolliana, *Wall. Cat.* n. 430.—Reg. Himal.
Candollii, DC. Prod. iv. 654 = praec.
canescens, Hort. Taur. ex DC. Prod. iv. 651 = arvensis.
canescens, Waldst. & Kit. Pl. Rar. Hung. i. 50 = suaveolens.
capillata, Roem. & Schult. Syst. iii. 64 = gramuntia.
carpathica, Fisch. ex Spreng. Syst. i. 377 = arvensis.
caucasea, Sims, Bot. Mag. t. 886 = seq.
caucasica, *Bieb. Fl. Taur. Cauc.* i. 98.—Reg. Caucas. ; Soongaria.
centauroides, Lam. Illustr. i. 250 = Cephalaria corniculata.
centauroides, *Mill. Gard. Dict.* ed. VIII. n. 4 (Quid ?). —Hispan.
ceratophylla, *Roem. & Schult. Syst.* iii. 521.—Sibir.
ceratophylla, Tenore, Prod. Fl. Nap. p. xii = S. Columbaria.
ciliata, *Spreng. in Schrad. Journ.* (1880) II. 199.— Graec.
cilicica, *Boiss. ex Tchihat. Asie Min. Bot.* ii. 229, nomen.—As. Min.
cinerascens, Jord. Pugill. Pl. Nov. 89 = holosericea.
cinerea, Lam. Illustr. i. 251 = pyrenaica.
cochinchinensis, *Lour. Fl. Cochinch.* 68 (sp. dub.).— Cochinch. ; China.
coerulea, Auct. = caerulea.
colchica, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 371, in textu.—Reg. Caucas.
collina, *Req. in Guerin, Desc. Vaucl.* ed. II. 248.— Hispan.
collina, Salisb. Prod. 54 = stellata.
collina, *F. W. Schmidt, Fl. Boëm.* iii. 78.—Bohem.
Columbaria, *Linn. Sp. Pl.* 99.—Europ. ; As. bor. ; Afr. bor. et austr.
Columbaria, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 17 = maritima.
Columnae, Tenore, Fl. Nap. i. 29. t. 7 = gramuntia.
commutata, Roem. & Schult. Syst. iii. 65 = S. Columbaria.
comosa, Fisch. ex Roem. & Schult. l. c. 84 = Fischeri.
connata, Hornem. Hort. Hafn. i. 128 = caucasica.
coriacea, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 416 = Cephalaria corniculata.
corniculata, Waldst. & Kit. Pl. Rar. Hung. i. 11 = Cephalaria corniculata.
Cornucopia, *Viv. Fl. Lib. Spec.* 67.—Italia.
coronopifolia, Sibth. & Sm. Fl. Graec. Prod. i. 85.; Fl. Graec. ii. 13. t. 114 = crenata.
coronopifolia, Willd. ex Roem. & Schult. Syst. iii. 521 = ceratophylla.
cosmoides, *Boiss. Diagn.* Ser. I. ii. 113.—Graec.
crassicaulis, E. Mey. in Drège, Zwei Pfl. Docum. 218 = S. Columbaria.
crenata, *Cyrill. Pl. Rar. Neap.* i. 11. t. 3.—Reg. Mediterr.

SCABIOSA :—

crenata, Lucé, Topog. Nachr. Oesel, 28 (Quid ?).— Rossia.
cretacea, Pall. ex Bieb. Fl. Taur. Cauc. i. 93 = Cephalaria corniculata.
cretica, *Linn. Sp. Pl.* 100.—Ins. Creta ; Sicil.
cretica, Willich, ex Roem. & Schult. Syst. iii. 53 = plumosa.
crinita, *Kotschy & Boiss. in Boiss. Fl. Orient.* iii. 133. —Kurdistan.
Cupani, Guss. Fl. Sic. Prod. i. 160 = maritima.
Dahuriae, Fisch. ex DC. Prod. iv. 658 = Fischeri.
Dallaportae, *Heldr. ex Boiss. Fl. Orient.* iii. 138.—Ins. Cephalon.
daucoides, *Desf. Fl. Atlant.* i. 23. t. 28.—Alger.
davurica, Willd. ex Roem. & Schult. Syst. iii. 521 = isetensis.
decemaristata, Portenschl. ex Schult. Mant. iii. 36 = brachiata.
decurrens, Sibth. & Sm. Fl. Graec. Prod. i. 80 = Cephalaria graeca.
decurrens, Thunb. Prod. Pl. Cap. 28 = Cephalaria attenuata.
diandra, Lag. in Varied. Cienc. ii. IV. (1805) 39 = papposa.
dichotoma, Lam. Illustr. i. 249 = Cephalaria syriaca.
dichotoma, *Ucria, in Roem. Arch.* i. I. (1796) 68. —Mauritan. ; Sicil.
diffusa, *Reichb. Hort. Bot.* i. 13. t. 16.—Ins. Canar.
dipsacifolia, Host, Fl. Austr. i. 191 = sylvatica.
dipsacoides, *Boiss. ex Tchihat. Asie Min. Bot.* 229, nomen.—As. Min.
dissecta, Dulac, Fl. Hautes-Pyr. 465 = gramuntia.
divaricata, Clarke, Trav. ii. 644 = papposa.
divaricata, Jacq. Hort. Vindob. i. 5 = ucranica.
divaricata, Lam. Illustr. i. 250 = rutaefolia.
diversifolia, Baumg. Enum. Stirp. Transs. i. 75 = banatica.
dubia, Moench, Enum. Pl. Indig. Hass. 62 = arvensis.
dumetorum, *Brouss. ex Willd. Enum. Hort. Berol.* 146.—Ins. Canar.
eburnea, Sibth. & Sm. Fl. Graec. Prod. i. 82 ; Fl. Graec. ii. 7. t. 106 = ucranica.
elata, Hornem. Hort. Hafn. i. 126 = Cephalaria tatarica.
elegans, Spreng. Pugill. ii. 24 = caucasica.
eremophila, *Boiss. Diagn.* Ser. I. x. 79.—Arab.
fenestrata, Pomel, Nouv. Mat. Fl. Atl. 63 = arenaria.
fimbriata, Mill. Gard. Dict. ed. VIII. n. 19 = palaestina.
Fischeri, *DC. Prod.* iv. 658.—Dahuria.
flava, Sibth. & Sm. Fl. Graec. Prod. ii. 356 = Cephalaria graeca.
flavescens, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 350 = ochroleuca.
flavida, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 141. —Persia.
Fleischmanni, Hladn. ex Nym. Consp. 347 = arvensis.
Frivaldskyi, Steud. Nom. ed. II. ii. 525 = ambigua.
frutescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 252.— Afr. trop.
frutescens, Mill. Gard. Dict. ed. VIII. n. 12 = cretica.
fruticulosa, *C. Sm. ex DC. Prod.* iv. 654.—Ins. Canar.
fumarioides, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 466.—Banat.
Fussiana, Schur, Enum. Pl. Transs. 962 (Index) = Cephalaria radiata.
Gemonae, D. Dietr. Syn. Pl. i. 452 = humilis.
gigantea, Ledeb. Ind. Sem. Hort. Dorpat. (1811) 4 = Cephalaria tatarica.
glabrata, Schott, in Roem. & Schult. Syst. iii. 61 = S. Succisa.
glabrescens, Jord. Pugill. Pl. Nov. 82 = lucida.
Gmelini, [Honck.] Verz. Aller Gew. Teutschl. 397 (Quid ?).—Europ.
Gmelini, A. St. Hil. in Bull. Soc. Philom. iii. (1812) 149 ; et in Journ. Phys. lxxv. (1812) 433 = ucranica.
gracilis, Boiss. Fl. Orient. iii. 135 = semipapposa.
gracilis, Roem. & Schult. Syst. iii. 61 = gramuntia.
graeca, D. Dietr. Syn. Pl. i. 447 = Cephalaria graeca.
graminifolia, *Linn. Cent. Pl.* i. 6 ; *Amoen. Acad.* iv. 267.—Europ. austr.
gramontia, Linn. Sp. Pl. ed. II. 143 = gramuntia.
gramuntia, Brot. Fl. Lusit. i. 145 = lusitanica.
gramuntia, *Linn. Syst.* ed. X. 889.—Reg. Mediterr.
gramuntiana, Gouan, Hort. Monsp. 62 = praec.

SCABIOSA :—

grandiflora, Guss. Fl. Sic. Prod. i. 160 = S. Cornucopia.
grandiflora, Scop. Delic. Insub. iii. 29. t. 14 = maritima.
grandiflora, Steud. Nom. ed. II. ii. 525 = caucasica.
Guitardi, Timb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxviii, in textu = lucida.
gumbetica, *Boiss. Fl. Orient.* iii. 137.—Reg. Caucas.
Gussonii, Tineo, ex Nym. Consp. 342 = crenata.
Hacquetii, Lam. Illustr. i. 250 = Cephalaria transylvanica.
Heraclea, *Willd. ex Roem. & Schult. Syst.* iii. 521 ; (sp. dub.).—Fret. Gadit.
heterophyllos, S. G. Gmel. It. i. 150. t. 28 ; ex J. F. Gmel. Syst. 229 = ochroleuca.
heterotricha, Opiz, in Lotos, ii. (1852) 229 (Quid ?).— Europ.
hirsuta, Lapeyr. Hist. Abr. Pl. Pyr. 59 = arvensis.
hirsuta, [Mazziari,] in Ionios, Anthol. iii. (1834) 691 = S. Succisa.
hirta, Willd. in Roem. & Schult. Syst. iii. 521 = ucranica.
hispanica, *Willd. l. c.*—Hispan.
hispida, Portenschl. ex Roem. & Schult. l. c. 58 = ciliata.
hispidula, *Boiss. Diagn.* Ser. I. ii. 112.—Graecia ; As. Min.
Hladnikiana, Host, Fl. Austr. i. 195 = lucida.
holosericea, *Bertol. Rar. Ital. Pl. Dec.* iii. 49.—Europ. austr.
holosericea, DC. Fl. Fr. Suppl. 489 = lucida.
holosericea, Panč. ex Nym. Consp. 343 = ochroleuca.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 218.—Reg. Himal.
Hornemanni, *Spreng. Syst.* iv. *Cur. Post.* 38.—Hab. ?
humilis, *Host, Fl. Austr.* i. 195.—Carniol.
humilis, Thunb. Prod. Pl. Cap. 28 = Cephalaria attenuata.
hybrida, All. Auct. Fl. Pedem. 9 = integrifolia.
hybrida, Bouch. Fl. Abb. 12 ; ex Roem. & Schult. Syst. iii. 54 = arvensis.
hymettia, *Boiss. & Sprun. Diagn.* Ser. I. ii. 111. —Graecia.
hyrcanica, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 370, in textu.—Reg. Caucas.
incana, Roem. & Schult. Syst. iii. 521 = suaveolens.
incana, Willd. ex Roem. & Schult. l. c. = ucranica.
incisa, Mill. Gard. Dict. ed. VIII. n. 18 = africana.
indurata, Linn. Mant. ii. 196 = africana.
inflexa, Kluk, in Bess. Prim. Fl. Galic. ii. 336 = australis.
integrata, Hoffmgg. & Link, Fl. Port. ii. 88 = maritima.
integrifolia, [Honck.] Verz. Aller Gew. Teutschl. 383 (Quid ?).—Germania.
integrifolia, *Linn. Sp. Pl.* 99.—Graec. ; As. Min.
integrifolia, Roth, Tent. Fl. Germ. ii. 164 = arvensis.
integrifolia, Savi, Fl. Pisan. i. 162 = sylvatica.
intermedia ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 114.—Europ.
intermedia, Roem. & Schult. Syst. iii. 76 = lusitanica ?
involucrata, Sibth. & Sm. Fl. Graec. Prod. i. 84 ; Fl. Graec. ii. 11. t. 112 = papposa.
isetensis, Colm, Cat. Pl. Catal. 76 = pulsatilloides.
isetensis, *Linn. Mant.* i. 37.—Reg. Caucas. ; As. bor.
Jacquini, Roem. & Schult. Syst. iii. 77, in obs. = stellata.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 113.— Japon.
joppensis, Reichb. Hort. Bot. i. 13. t. 17 = Cephalaria joppensis.
Jordani, Timb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxviii, in textu – pyrenaica.
Kitaibeli, Schult. Obs. Bot. 18 – ciliata.
laciniata, Licht. ex Roem. & Schult. Syst. iii. 87 = S. Columbaria.
laciniata, Lucé, Topog. Nachr. Oesel, 28 = praec. ?
laevigata, Waldst. & Kit. Pl. Rar. Hung. ii. 255 = Cephalaria corniculata.
lanata, *Hill, Veg. Syst.* v. 46.—Afr. austr.
lasiosperma, D. Dietr. Syn. Pl. i. 448 = fruticulosa.
lasiosperma, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 193.—Ins. Canar.
laxiflora, Willd. ex Link, Enum. Hort. Berol. i. 128 = pyrenaica.
legionensis, *Lag. Gen. et Sp. Nov.* 9.—Hispan.
leiocephala, Hoppe, ex Mert. & Koch, Deutschl. Fl. i. 754 = gramuntia.

SCABIOSA :—
leptopoda, Boiss. Diagn. Ser. I. x. 78 = arenaria.
leucantha, Linn. Sp. Pl. 98 = Cephalaria leucantha.
leucanthema, Frivald. ex Nym. Consp. 346 = Cephalaria graeca.
leucanthema, *Hill, Veg. Syst.* v. 39. t. 20.—Gallia.
leucocephala, Link, Handb. i. 856 (Index) = Cephalaria leucantha.
limonifolia, *Vahl, Symb. Bot.* ii. 27.—Sicilia.
linifolia, C. Koch, in Linnaea, xvii. (1843) 36 = rotata.
lobata, Nym. Consp. 343 = stellata.
longepedunculata, Fisch. Hort. Gorenk. ed. II (1812), 43, nomen = amoena.
longifolia, *Waldst. & Kit. Pl. Rar. Hung.* i. 4.—Europ.
Loretiana, Timb. ex Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxxii = gramuntia.
lucida, Ait. Hort. Kew. ed. II. i. 229 = nitens.
lucida, *Vill. Prosp.* 18; *Hist. Pl. Dauph.* ii. 293.—Europ.
lusitanica, *D. Dietr. Syn. Pl.* i. 448.—Lusitan.
lutea, *Hort. ex Loud. Hort. Brit.* i. 44.—Rossia.
lutescens, Gilib. Fl. Lituan. i. 169 = ochroleuca.
lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 109.—As. Min.
lyrata, Forsk. Fl. Aegypt. Arab. 203 = sicula.
lyrata, Lam. Illustr. i. 250 = integrifolia.
lyrophylla, Vis. in Mem. Ist. Venet. xv. (1870) t. 19 (Pl. Serb.) = seq.
macedonica, *Vis. l. c.* 12.—Graecia.
macrochaete, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 143.—Persia.
macropoda, Costa, ex Willk. in Linnaea, xxx. (1859–60) 103 = pulsatilloides.
major, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 786 = atropurpurea.
maritima, Lag. Elench. Hort. Madrit. (1808); ex ej. Gen. et Sp. Nov. 8. n. 111 = lusitanica?
maritima, *Linn. Cent. Pl.* ii. 8; *Amoen. Acad.* iv. 304.—Reg. Mediterr.; Ins. Canar.
maritima, Thunb. Prod. Pl. Cap. 29 = S. Columbaria.
maritima, Vill. Hist. Pl. Dauph. ii. 295 = pyrenaica.
maritima, Wulf. in Roem. Arch. iii. III. (1805) 317 = ucranica.
mediterranea, Viv. App. Fl. Cors. Prod. 1 = Cephalaria leucantha.
micrantha, *Desf. in Ann. Mus. Par.* xi. (1808) 168.—Serbia; Reg. Caucas.; Armen.
microcephala, *Nym. Syll.* 59.—Hispan.
mixta, *De Not. Prosp. Fl. Ligust.* 52.—Liguria.
moldavica, Hort. ex Roem. & Schult. Syst. iii. 58 = integrifolia.
mollis, Schleich. Cat. Pl. Helv. ed. III. 27 = pyrenaica.
mollis, Willd. Enum. Hort. Berol. Suppl. 7 = gramuntia.
mollissima, DC. Fl. Fr. Suppl. 490 = pyrenaica.
mollissima, Viv. Ann. Bot. i. II. (1804) 161 = gramuntia.
monspeliaca, *Georgi, Beschr. Russ. Reich.* III. iv. 727.—Tauria.
monspeliensis, DC. Prod. iv. 655 = stellata.
monspeliensis, Jacq. Misc. ii. 320 = praec.?
montana, *Bieb. Fl. Taur. Cauc.* i. 95.—As. Min.; Reg. Ural.
montana, Mill. Gard. Dict. ed. VIII. n. 5 = arvensis.
montana, Schur, ex Nym. Consp. 343 = ochroleuca.
multiseta, Vis. Stirp. Dalm. Rar. 1. t. 1 = brachiata.
neglecta, *Hornem. Hort. Hafn.* i. 127.—Italia.
nitens, *Roem. & Schult. Syst.* iii. 82.—Ins. Azores.
nitida, Bernh. ex Roem. & Schult. l. c. 68 = amoena.
nivea, *Agardh, ex Roem. & Schult. l. c.* 81.—Hispan.
norica, Vest, in Bot. Zeit. Regensb. iv. (1805) 40 = lucida.
nudicaulis, Lam. Illustr. i. 252 = S. Columbaria.
nudicaulis, Sieber, Reise Kreta, ii. 321 = sphaciotica.
ochroleuca, *Linn. Sp. Pl.* 101.—Europ.; As. bor.
ochroleuca, Thunb. Prod. Pl. Cap. 29 = S. Columbaria.
odontostemma, Kotschy, ex Boiss. Fl. Orient. iii. 145 = palaestina.
odora, Salisb. Prod. 55 = maritima.
officinarum-arvensis, Crantz, Inst. i. 352 = arvensis.
Olivieri, *Coult. Mém. Dipsac.* 36. t. 2. f. 10.—Arab.; Persia; Afghan.; Soongar.
orientalis, *Lag. Gen. et Sp. Nov.* 9.—Oriens.
orophila, Timb. & Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxxi = S. Columbaria.
ovatifolia, Lag. Gen. et Sp. Nov. 9 = sylvatica.
Owerini, *Boiss. Fl. Orient.* iii. 133.—Reg. Caucas.
palaestina, D. Dietr. Syn. Pl. i. 448 = brachiata.

SCABIOSA :—
palaestina, *Linn. Mant.* i. 87.—As. Min.; Syria; Persia.
pallida, Láng, ex Nym. Consp. 343, 344 = ochroleuca, banatica.
pallida, E. Mey. in Drège, Zwei Pfl. Docum. 218 = S. Columbaria.
pallida, Salisb. Prod. 55 = cretica.
pannonica, Jacq. Enum. Vindob. 22 = sylvatica.
papposa, Gaertn. Fruct. ii. 38. t. 86. f. 4 = plumosa.
papposa, *Linn. Sp. Pl.* 101.—Ins. Cyprus; Syria; Persia.
parviflora, Desf. Fl. Atlant. i. 119 = dichotoma.
patens, Jord. Pugill. Pl. Nov. 94 = S. Columbaria.
pauciseta, DC. Prod. iv. 658 = gramuntia.
pectinata, *Lag. Elench. Fl. Madrit.* (1803) 33; *ex ej. Gen. et Sp. Nov.* 8. n. 108.—Arab.
permixta, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 319 = S. Columbaria.
persica, Boiss. Diagn. Ser. I. x. 81 = palaestina.
phrygia, Boiss. l. c. Ser. II. vi. 96 = palaestina.
pilosa, Roem. & Schult. Syst. iii. 80 = ucranica.
pinnata, *Gueldenst. Reisen*, ii. 10 (Quid?).—Rossia.
plumosa, Link, in Schrad. Neues Journ. ii. (1807) II. 300 = lusitanica.
plumosa, *Sibth. & Sm. Fl. Graec. Prod.* i. 84; *Fl. Graec.* ii. 11. t. 111.—Graecia; Syria; Persia.
polymorpha, F. W. Schmidt, Fl. Boem. iii. 76 = arvensis.
polymorpha, *Weigel, Obs.* 23 = S. Columbaria, ochroleuca.
porphyrostephana, Boiss. ex Boiss. Fl. Orient. iii. 146 = rotata.
praemorsa, Gilib. Fl. Lituan. i. 168 = S. Succisa.
pratensis, Jord. Pugill. Pl. Nov. 91 = S. Columbaria.
pratensis, F. W. Schmidt, Fl. Boëm. iii. 78 = arvensis.
procera, Salisb. Prod. 55 = africana.
prolifera, *Linn. Syst.* ed. X. 889.—Syria; Ins. Cyprus.
prolifera, [Mazziari,] in Ionios, Anthol. iii. (1834) 691 = S. Succisa.
propontica, *Lag. Gen. et Sp. Nov.* 9.—As. Min.
Pseudaustralis, Roem. & Schult. Syst. iii. 61 = australis.
Pseudo-banatica, Schur, Enum. Pl. Transs. 300 = lucida.
Pterocephala, *Linn. Sp. Pl.* 100.—Graecia.
pubescens, Jord. Pugill. Pl. Nov. 90 = lucida.
pubescens, Wahlenb. Fl. Carpat. 38 = ciliata.
pubescens, Waldst. & Kit. ex Roem. & Schult. Syst. iii. 57 = collina.
pulsatilloides, *Boiss. Elench.* 58.—Hispan.
pumila, Burm. f. Fl. Cap. Prod. 4 = S. Columbaria.
pyrenaica, *All. Fl. Pedem.* i. 140. t. 25.—Europ. austr.
pyrenaica, Hornem. ex Spreng. Syst. iv. Cur. Post. 38 = Hornemanni.
pyrenaica, C. A. Mey. Verz. Pfl. Cauc. 48 = amoena.
radiata, F. W. Schmidt, Fl. Boëm. iii. 78 = arvensis.
repens, Brign. Fasc. Pl. Forojul. 19 = australis.
repens, [Honck.] Verz. Aller Gew. Teutschl. 397 (Quid?).—Germania.
Reuteriana, *Boiss. Diagn.* Ser. I. ii. 112.—Caria.
reversa, *Ledeb. Ind. Hort. Dorpat. Suppl.* (1824) 6.—Sibir.
rhizantha, Viv. Fl. Lib. Spec. 6. t. 3 = arenaria.
rhodantha, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 435 = Olivieri.
rigida, Linn. Pl. Rar. Afr. 8; in Amoen. Acad. vi. 84 = Cephalaria rigida.
rigida, Mill. Gard. Dict. ed. VIII. n. 7 = Cephalaria leucantha.
rosea, Hort. ex Steud. Nom. ed. II. ii. 526 = maritima.
rotata, *Bieb. Fl. Taur. Cauc.* iii. 102.—As. Min.; Persia.
rubella, Opiz, ex Mert. & Koch, Deutschl. Fl. i. 751; et Natural. viii. (1824) 76 = S. Columbaria.
rupestris, Bieb. Fl. Taur. Cauc. i. 95 = isetensis.
rutaefolia, *Vahl, Symb. Bot.* ii. 26.—Reg. Mediterr. occ.
ruthenica, *Tchertkoff, in Bull. Soc. Nat. Mosc.* vi. (1833) 16, nomen.—Ross. centr.
Salcedi, *Lag. ex Roem. & Schult. Syst.* iii. 55.—Hispan.
Saviana, Reichb. Hort. Bot. i. 38. t. 53 = maritima.
saxatilis, *Cav. Ic.* ii. 68. t. 184.—Hispan.
scabra, Thunb. Prod. Pl. Cap. 29 = Cephalaria scabra.
Schimperiana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 112 = palaestina.

SCABIOSA :—
schizopetala, Fenzl, ex Boiss. Fl. Orient. iii. 145 = palaestina.
Scopolii, *Link, Enum. Hort. Berol.* i. 128.—Europ. austr.
scrophulariaefolia, Boiss. Fl. Orient. iii. 133 = crinita.
semipapposa, *Salzm. ex DC. Prod.* iv. 658.—Reg. Mediterr. occ.
sericea, Jord. Pugill. Pl. Nov. 98 = graminifolia.
serrata, Lam. Illustr. i. 249 = integrifolia.
setifera, *Lam. l. c.* 251.—Europ. austr.
setulosa, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 31 = palaestina.
sibirica, Lam. Illustr. i. 249 = Cephalaria syriaca.
Sibthorpiana, Sibth. & Sm. Prod. Fl. Graec. i. 84; Fl. Graec. ii. 10. t. 110 = brachiata.
sicula, Bieb. Fl. Taur. Cauc. i. 98 = micrantha.
sicula, *Linn. Mant.* ii. 196.—Europ. austr.; As. Min.
silenifolia, *Waldst. & Kit. Pl. Rar. Hung.* ii. 170.—Europ.
simplex, DC. Fl. Fr. iv. 231 = stellata.
simplex, *Desf. Fl. Atlant.* i. 125.—Afr. bor.
solitaria, [*Mazziari,*] *in Ionios, Anthol.* iii. (1834) 698.—Ins. Corcyra.
songarica, *Schrenk, Enum. Pl. Nov.* i. 55.—Sibir.
spathulata, *D. Dietr. Syn. Pl.* i. 448.—Hispan.
spathulata, *Kit. in Linnaea*, xxxii. (1863) 369.—Europ.
speciosa, *Royle, Illustr. Bot. Himal.* 245. t. 55. f. 2.—Reg. Himal.
sphaciotica, Frivald. ex Nym. Consp. 341 = S. Pterocephala.
sphaciotica, *Roem. & Schult. Syst.* iii. 86.—Ins. Creta.
sphakiottica, Sieber, Herb. Cret. 2, nomen = praec.?
spreta, Jord. Pugill. Pl. Nov. 92 = S. Columbaria.
squamaeflora, Sieber, ex Schult. Mant. iii. 45 (Quid?).—Hab.?
stellaris, Salisb. Prod. 54 = stellata.
stellata, Bieb. Fl. Taur. Cauc. i. 98 = rotata.
stellata, *Linn. Sp. Pl.* 100.—Reg. Mediterr. occ.
stellata, Pall. ex Bieb. Fl. Taur. Cauc. i. 98, in syn. = micrantha.
stellata, Schangin, in Pall. N. Nord. Beitr. vi. 76 = ochroleuca.
stricta, Seidel, ex Roem. & Schult. Syst. iii. 55, in syn. = arvensis.
stricta, Waldst. & Kit. Pl. Rar. Hung. ii. 147 = lucida.
styriaca, *Vest, in Flora*, iv. (1821) 146.—Styria.
suaveolens, *Desf. Tabl.* ed. I. 110.—Europ.; As. Min.
subescens, Reichb. Ic. Fl. Germ. xxii. 18 (err. typ.) = *pubescens*.
subpubescens, Hort. ex Schult. Mant. iii. 28 = arvensis.
Succisa, *Linn. Sp. Pl.* 98.—Europ.; As. bor. trop. & austr.
sulphurea, *Boiss. & Huet, Diagn.* Ser. II. ii. 123.—Armenia.
sylvatica, *Linn. Sp. Pl.* ed. II. 142.—Europ.; Reg. Caucas.
sympaganthera, Ruiz & Pav. Fl. Per. i. 49 = Calycera balsamitaefolia.
syriaca, D. Dietr. Syn. Pl. i. 451, sphalm. = styriaca.
syriaca, Linn. Sp. Pl. 98 = Cephalaria syriaca.
syriaca, Urv. ex Coult. Mém. Dips. 25 = Cephalaria transylvanica.
tatarica, Habl. Phys. Beschr. Taur. 154 = Cephalaria corniculata.
tatarica, Hornem. Hort. Hafn. i. 126 = montana.
tatarica, Linn. Sp. Pl. 99 = Cephalaria tatarica.
tatarica, Lumnitz, Fl. Poson. 55 = ciliata.
tatarica, Viv. ex Coult. Mém. Dips. 25 = Cephalaria transylvanica.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. vi. 73.—Graecia.
tenuifolia, Frivald. ex Steud. Nom. ed. II. ii. 527 = triniaefolia.
tenuifolia, Lag. Elench. Hort. Madrit. (1803); ex ej. Gen. et Sp. Nov. 8. n. 111 = lusitanica?
tenuifolia, *Presl, Symb. Bot.* i. 20. t. 12.—Hab.?
tenuifolia, Roth, Tent. Fl. Germ. i. 59 = ochroleuca.
tenuifolia, Savi, Bot. Etrusc. i. 122 = gramuntia.
tenuifolia, Willd. ex Roem. & Schult. Syst. iii. 521 = ucranica.
tenuis, *Sprun. in Boiss. Diagn.* Ser. I. ii. 114.—Graecia.
tenuisecta, Jord. Pugill. Pl. Nov. 96 = gramuntia.
Thunbergii, Roem. & Schult. Syst. iii. 76, in nota = S. Columbaria.

SCABIOSA :—

Timeroyi, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = sylvatica.
tmolea, Boiss. Fl. Orient. iii. 139 = ucranica.
tolosana, Timb. & Jeanb. in Bull. Soc. Bot. Fr. xi. (1864) p. lxxxi = S. Columbaria.
tomentosa, *Cav. Ic.* ii. 66. t. 183.—Hispan.
tomentosa, J. F. Gmel. Syst. 229 = pyrenaica.
tomentosa, E. Mey. in Drège, Zwei Pfl. Docum. 218 = Buekiana.
tomentosa, Sibth. & Sm. Fl. Graec. Prod. i. 85 = sphaciotica.
transsilvanica, Sprun. ex Nym. Consp. 343 = maritima.
transylvanica, All. Fl. Pedem. i. 138 = Cephalaria corniculata.
transylvanica, Linn. Sp. Pl. 98 = Cephalaria transylvanica.
transylvanica, Sieber, ex DC. Prod. iv. 648 = Cephalaria joppensis.
trenta, Hacq. Pl. Alp. 14. t. 4 = Cephalaria transylvanica.
triandra, Linn. Sp. Pl. 99 = gramuntia.
trichotoma, Moench, Meth. 490 = integrifolia.
trifida, Thunb. Prod. Pl. Cap. 28 = Cephalaria attenuata.
triniaefolia, *Frivald. in Flora*, xviii. (1835) 333.—Graecia ; Rumelia.
Triumfetti, Moretti, ex Roem. & Schult. Syst. iii. 61 = australis.
trivialis, F. W. Schmidt, Fl. Boëm. iii. 78 = arvensis.
turicensis ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 114.—Europ.
ucranica, *Linn. Syst.* ed. X. 889.—Europ. ; As. Min. ; Persia.
uniflora, [*Mazziari,*] *in Ionios, Anthol.* iii. (1834) 698.—Ins. Corcyra.
uniseta, Savi, Fl. Pisan. i. 167 = gramuntia.
uralensis, Murr. in Comm. Gotting. (1782) 13. t. 4 = Cephalaria corniculata.
uralensis, Willd. Sp. Pl. i. 551 ; et ex Ledeb. Fl. Ross. ii. 447 = Cephalaria tatarica.
urceolata, Desf. Fl. Atlant. i. 122 = rutaefolia.
Urvillei, D. Dietr. Syn. Pl. i. 447 = integrifolia.
ustulata, Thunb. Prod. Pl. Cap. 29 = Cephalaria corniculata.
Vahlii, Coult. Mém. Dips. 35. t. 2. f. 6 = palaestina.
Vaillantii, Bieb. Fl. Taur. Cauc. iii. 99 = Cephalaria syriaca.
varia, Gilib. Fl. Lituan. i. 167 = arvensis.
variifolia, *Boiss. Fl. Orient.* iii. 137.—Ins. Rhodus.
velutina, Jord. Pugill. Pl. Nov. 87 = pyrenaica.
venusta, *Hort. Bonon. ex Pasq. Cat. Ort. Bot. Nap.* (1867) 93, nomen.—Hab. ?
verbascifolia, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 459.—Gallia.
verbenacea, Lam. Illustr. i. 250 = Cephalaria attenuata.
vestina, *Facc. ex Koch, Syn. Fl. Germ.* ed. II. 447 ; *et in Zeitschr. Ferdinand. Tirol.* III. iv. (1854) 13.—Europ.
vestita, Jord. Pugill. Pl. Nov. 86 = pyrenaica.
villosa, [Honck.] Verz. Aller Gew. Teutschl. 383 (Quid ?).—Germania.
villosa, *Schousb. ex Steud. Nom.* ed. II. ii. 527, nomen.—Marocc.
virens, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 194.—Ins. Canar.
virescens, Nym. Consp. 342 = graminifolia.
Virga-pastoris, Mill. Gard. Dict. ed. VIII. n. 9 = suaveolens.
vivipara, Nees, ex Steud. Nom. ed. II. ii. 527 (Quid ?).—Hab. ?
Webbiana, D. Don, in Bot. Reg. t. 717 = ochroleuca.
Willichii, Link, in Schrad. Neues Journ. ii. (1807) II. 299 = plumosa.
Wulfeniana, Hort. ex Reichb. Ic. Fl. Germ. iv. 24 = maritima.
Wulfenii, Roem. & Schult. Syst. iii. 76, in nota = ucranica.

SCABRITA, Linn. Mant. i. 3 (1767) = **Nyctanthes**, Linn. (Oleac.).
scabra, Linn. Syst. ed. XII. 115 = N. Arbor-tristis.
triflora, Linn. Mant. i. 37 = N. Arbor-tristis.

SCADIANUS, Rafin. Atl. Journ. 164 (1833) = **Crinum**, Linn. (Amaryll.).
multiflorus, Rafin. Fl. Tellur. iv. 23 = C. americanum.

SCADOXUS, Rafin. Fl. Tellur. iv. 19 (1836) = **Haemanthus**, Tourn. (Amaryll.).
multiflorus, Rafin. l. c. = H. multiflorus.

SCADUAKINTOS, Rafin. Fl. Tellur. iv. 23 (1836) = **Brodiaea**, Sm. (Liliac.).
umbellaris, Rafin. l. c. = B. lactea.

SCAEVOLA, Linn. Mant. ii. 145 (1771). *GOODENOVIEAE*, Benth. & Hook. f. ii. 539.
BAUDINIA, Lesch. ex DC. Prod. vii. 511 (1839).
CAMPHUSIA, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 148, partim.
CERBERA, Lour. Fl. Cochinch. 136 (1790).
CROSSOTOMA, Spach, Hist. Vég. Phan. ix. 583 (1840).
GLYPHA, Lour. ex Endl. Gen. 506, in syn. (1838).
HEMICHARIS, Salisb. ex DC. Prod. vii. II. 510 (1839).
LOBELIA, Adans. Fam. ii. 157 (1763).
MERKUSIA, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 150.
MOLKENBOERIA, De Vriese, in Nat. Verh. Haarl. x. (1854) 38. t. 7.
POGONANTHERA, Spach, Hist. Vég. Phan. ix. 583 (1840).
POGONETES, Lindl. Introd. Nat. Syst. ed. II. 443. (1836).
ROEMERIA, Dennst. Schluess. Hort. Malab. 24 (1818).
TEMMINCKIA, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 141, partim.
XEROCARPA, Spach, Hist. Vég. Phan. ix. 583 (1840).
aemula, R. Br. Prod. 584 = saligna.
amblyanthera, *F. Muell. Fragm.* i. 121.—Austral.
amboinensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 210.—Ins. Amboin.
anchusaefolia, *Benth. in Enum. Pl. Hueg.* 68.—Austral.
angulata, *R. Br. Prod.* 586.—Austral.
apterantha, *F. Muell. Fragm.* i. 121.—Austral.
atriplicina, *F. Muell. l. c.* ii. 18.—Austral.
attenuata, *R. Br. Prod.* 583.—Austral.
auriculata, *Benth. Fl. Austral.* iv. 99.—Austral.
Bela-Modagani, Roem. & Schult. Syst. v. 163 = Koenigii.
Benthamea, De Vriese, in Lehm. Pl. Preiss. i. 411 = striata.
Billardieri, D. Dietr. Syn. Pl. i. 786 = Koenigii.
Brookeana, *F. Muell. in Vict. Natural.* i. (Dec. 1884) 122.—Austral.
caespitosa, R. Br. Prod. 585 = globulifera.
calliptera, Benth. in Enum. Pl. Hueg. 70 = striata.
Candollei, De Vriese, in Lehm. Pl. Preiss. i. 405 = platyphylla.
canescens, *Benth. in Enum. Pl. Hueg.* 69.—Austral.
Chamissoniana, *Gaudich. in Freyc. Voy. Bot.* 461. t. 82.—Ins. Sandvic.
chlorantha, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 137 = Koenigii.
ciliata, G. Don, Gen. Syst. iii. 728 = Chamissoniana.
clandestina, F. Muell. Fragm. i. 206 = Diaspasis filifolia.
collaris, *F. Muell. Rep. Bab. Exped.* 15.—Austral.
coriacea, *Nutt. in Trans. Am. Phil. Soc.* N. S. viii. (1843) 253.—Polynes.
Corsoniana, *G. Don, in Loud. Gard. Mag.* xviii. (1842) 370.—Ins. Molucc.
crassifolia, *Labill. Nov. Holl. Pl.* i. 56. t. 78.—Austral.
cumana, *De Vriese, in Verh. Haarl.* x. (1854) 35.—Am. centr.
cuneiformis, *Labill. Nov. Holl. Pl.* i. 56. t. 80.—Austral.
Cunninghamii, *DC. Prod.* vii. 508.—Austral.
Cunninghamii, Steud. Nom. ed. II. ii. 527, partim = Hymenanthera crassifolia.
depauperata, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 83.—Austral.
depressa, De Vriese, in Lehm. Pl. Preiss. i. 410 = longifolia ?
Drummondi, DC. Prod. vii. 508 = nitida.
enantophylla, *F. Muell. Fragm.* viii. 58.—Austral.
fasciculata, *Benth. in Enum. Pl. Hueg.* 68.—Austral.
fastigiata, De Vriese, in Lehm. Pl. Preiss. i. 406 = nitida.
flaccida, De Vriese, l. c. 407 = thesioides.
floribunda, *A. Gray, in Proc. Am. Acad.* v. (1861) 152.—Ins. Fiji.
Gaudichaudi, *Hook. & Arn. Bot. Beech. Voy.* 89.—Ins. Sandvic.

SCAEVOLA :—

Gaudichaudiana, Cham. in Linnaea, viii. (1833) 226 = Chamissoniana.
geniculata, De Vriese, in Lehm. Pl. Preiss. i. 404 = Goodenia geniculata.
glabra, *Hook. & Arn. Bot. Beech. Voy.* 89.—Ins. Sandvic.
glandulifera, *DC. Prod.* vii. 510.—Austral.
glaucescens, De Vriese, in Lehm. Pl. Preiss. i. 410 = canescens.
globulifera, *Labill. Nov. Holl. Pl.* i. 55. t. 79.—Austral.
goodeniacea, F. Muell. Fragm. i. 121 = Catosperma Muelleri.
gracilis, *Hook. f. in Journ. Linn. Soc.* i. (1857) 129.—Ins. Kermadec.
grandiflora, Benth. in Enum. Pl. Hueg. 70 = Leschenaultia linarioides.
Groeneri, *F. Muell. Fragm.* vi. 15.—Austral.
hainanensis, *Hance, in Journ. Bot.* xvi. (1878) 229.—China.
hispida, *Cav. Ic.* vi. 7. t. 510.—Austral.
holosericea, *De Vriese, in Lehm. Pl. Preiss.* i. 408.—Austral.
Hookeri, *F. Muell. ex Hook. f. Fl. Tasm.* i. 231. t. 67.—Australia.
humifusa, *De Vriese, in Lehm. Pl. Preiss.* i. 410.—Austral.
humilis, *R. Br. Prod.* 585.—Austral.
ivaefolia, *L'Hérit. ex Sweet, Hort. Brit.* ed. II. 325.—Ins. Trinit.
Koenigii, *Vahl, Symb. Bot.* iii. 36.—Reg. trop.
laevigata, Pers. Syn. i. 195 = microcarpa.
Lambertiana, De Vriese, in Nederl. Kruidk. Arch. ii. (1851) 139 = Koenigii.
lanceolata, *Benth. in Enum. Pl. Hueg.* 69.—Austral.
lasiantha, F. Muell. Fragm. i. 207 = lanceolata.
lativaga, Hance, in Walp. Ann. ii. 1054 = Koenigii.
Leschenaultii, DC. Prod. vii. 506 = Koenigii.
ligustrifolia, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 253 = Chamissoniana.
linearis, *R. Br. Prod.* 586.—Austral.
Lobelia, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 250 = Koenigii.
Lobelia, *Murr. Syst.* ed XIII. 178; *Thunb. Prod. Pl. Cap.* 38.—Reg. trop.
longifolia, *De Vriese, in Lehm. Pl. Preiss.* i. 410.—Austral.
lycioides, DC. Prod. vii. 512 = spinescens.
lyratifolia, *De Vriese, in Lehm. Pl. Preiss.* i. 405 (al. gen. ?).—Austral.
Macraei, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 30 = S. Lobelia.
macrocalyx, De Vriese, l. c. 138 = Koenigii.
macrodonta, DC. Prod. vii. 511 = striata ?
macrophylla, *Benth. Fl. Austral.* iv. 98.—Austral.
macropoda, DC. Prod. vii. 509 = striata.
macrostachya, *Benth. Fl. Austral.* iv. 97.—Austral.
Maitlandi, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 497 ; et in Edinb. N. Phil. Journ. N. S. xvii. (Jan.–Apr. 1863) 231 = Cunninghamii.
membranacea, Benth. in Enum. Pl. Hueg. 69 = pilosa ?
Menziesiana, Cham. in Linnaea, viii. (1833) 227 = Chamissoniana.
micrantha, *Presl, Rel. Haenk.* ii. 58.—Ins Philipp.
microcarpa, *Cav. Ic.* vi. 6. t. 509.—Austral.
microphylla, *Benth. Fl. Austral.* iv. 100.—Austral.
mollis, *Hook. & Arn. Bot. Beech. Voy.* 89.—Ins. Sandvic.
montana, Gaudich. in Freyc. Voy. Bot. 460 = Gaudichaudi.
montana, Labill. Sert. Austro-Caled. 41. t. 42 = Koenigii.
multiflora, Lindl. Swan River App. 26 = nitida.
nitida, *R. Br. Prod.* 584.—Austral.
Novae-Zelandiae, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 52 = Hymenanthera crassifolia.
Oldfieldii, *F. Muell. Fragm.* ii. 19.—Austral.
oleoides, DC. Prod. vii. 512 = spinescens.
oppositifolia, *Roxb. Hort. Beng.* 85 ; *Fl. Ind.* i. 528.—Ins. Molucc.
ovalifolia, *R. Br. Prod.* 584.—Austral.
oxyclona, *F. Muell. Fragm.* x. 58.—Austral.
pallida, R. Br. Prod. 585 = microcarpa.
paludosa, *R. Br. l. c.* 586.—Austral.
paniculata, De Vriese, in Lehm. Pl. Preiss. i. 407 = thesioides.

SCAEVOLA :—
parvifolia, *F. Muell. ex Benth. Fl. Austral.* iv. 91.—Austral.
patens, F. Muell. Fragm. iii. 33 = depauperata.
phlebopetala, *F. Muell. l. c.* ii. 18.—Austral.
piliplena, *Miq. Fl. Ind. Bat.* ii. 581.—Java.
pilosa, *Benth. in Enum. Pl. Hueg.* 69; *DC. Prod.* vii. 511.—Austral.
platyphylla, *Lindl. Swan River App.* 26.—Austral.
plumerioides, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 252 = Koenigii.
Plumieri, Blume, Bijdr. 730 = Koenigii.
Plumieri, Vahl, Symb. Bot. ii. 36 = S. Lobelia.
polystachya, DC. Prod. ii. 508 = thesioides.
porocarya, *F. Muell. Fragm.* ii. 19.—Austral.
prostrata, De Vriese, in Lehm Pl. Preiss. i. 406 = Dampiera diversifolia.
pterosperma, De Vriese, l. c. i. 408 = Goodenia incana.
pubescens, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 253 = Chamissoniana.
pusilla, De Vriese, in Lehm. Pl. Preiss. i. 412 = Goodenia tenella.
Reinwardtii, De Vriese, l. c. 409 = Verreauxia Reinwardtii.
repens, De Vriese, l. c. 406 = paludosa.
restiacea, *Benth. Fl. Austral.* iv. 91.—Austral.
revoluta, *R. Br. Prod.* 586.—Austral.
rufa, De Vriese, in Lehm. Pl. Preiss. i. 405 = glandulifera.
saligna, *Forst. f. Prod.* 89.—N. Caled.
semiamplexicaulis, DC. Prod. vii. 509 = platyphylla.
semiamplexicaulis, De Vriese, in Nat. Verh. Haarl. x. (1854) 41 = auriculata.
senegalensis, Presl, Rel. Haenk. ii. 59 = S. Lobelia.
sericea, Forst. f. Prod. 89; Vahl, Symb. Bot. ii. 37 = Koenigii,
sericophylla, *F. Muell. ex Benth. Fl. Austral.* iv. 102.—Austral.
Sieberi, De Vriese, in Nat. Verh. Haarl. x. (1854) 33 = S. Lobelia.
sinuata, R. Br. Prod. 584 = saligna.
sphaerocarpa, De Vriese, in Lehm. Pl. Preiss. i. 409 = holosericea.
spinescens, *R. Br. Prod.* 586.—Austral.
squarrosa, Lindl. Swan River App. 26 = thesioides.
stenophylla, *Benth. Fl. Austral.* iv. 104.—Austral.
striata, *R. Br. Prod.* 586.—Austral.
stricta, De Vriese, in Lehm. Pl. Preiss. i. 408 = Goodenia scapigera.
stricta, Roem. & Schult. Syst. v. 167, sphalm. = striata.
suaveolens, *R. Br. Prod.* 585.—Austral.
Taccada, Roxb. Hort. Beng. 15; Fl. Ind. i. 527 = Koenigii.
tenera, De Vriese, in Lehm. Pl. Preiss. i. 409 = Goodenia caerulea.
thesioides, *Benth. in Enum. Pl. Hueg.* 68.—Austral.
Thunbergii, Eckl. & Zeyh. Enum. 387 = S. Lobelia.
tomentosa, *Gaudich. in Freyc. Voy. Bot.* 460. t. 81.—Austral.
tortuosa, *Benth. Fl. Austral.* iv. 91.—Austral.
Tournoa, Willd. Enum. Hort. Berol. Suppl. 10 (Quid?).—Hab.?
trinervia, De Vriese, in Lehm. Pl. Preiss. i. 407 = canescens.
umbellata, De Vriese, l. c. 411 = Goodenia filiformis.
uvifera, Stocks, in Wight, Ic. t. 1613 = S. Lobelia.
velutina, Presl, Rel. Haenk. ii. 58 = Koenigii.
Verreauxii, *F. Muell. Bot. Teachings* (1877), 65.—Austral.

SCALESIA, Arn. in Lindl. Introd. Nat. Syst. ed. II. 443 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 367.
affinis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 212.—Ins. Galapag.
aspera, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 180.—Ins. Galapag.
atractyloides, *Arn. in Lindl. Introd. Nat. Syst.* ed. II. 443.—Ins. Galapag.
Darwinii, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 211.—Ins. Galapag.
decurrens, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 182.—Ins. Galapag.
devisa, *Anderss. l. c.* 179.—Ins. Galapag.
gummifera, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 212.—Ins. Galapag.
incisa, *Hook. f. l. c.* 210.—Ins. Galapag.

SCALESIA :—
ovata, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855), 181.—Ins. Galapag.
pedunculata, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 211.—Ins. Galapag.

SCALIA, Sims, Bot. Mag. t. 956 (1806) = **Podolepis**, Labill. (Compos.).
jaceioides, Sims, l. c. = P. acuminata.

SCALIGERA, Adans. Fam. ii. 323 (1763) = **Aspalathus**, Linn. (Legumin.).
orientalis, Rafin. Sylva Tellur. 69 = A. orientalis.

SCALIGERIA, DC. Coll. Mém. v. 70 (1829); Prod. iv. 248 (1830) = **Conopodium**, Koch (Umbell.).
cretica, Boiss. Diagn. Ser. I. x. 52 (= *Bunium creticum*, Urv.).—Europ. austr.
elata, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 877 (= *Elaeosticta meifolia*).—Syria; Mesopot.
glaucescens, Boiss. l. c. (= *Elaeosticta glaucescens*).—Persia.
lazica, Boiss. l. c. 876.—As. Min.
meifolia, Boiss. l. c. (= *Elaeosticta meifolia*).—As. Min.; Syria.
microcarpa, DC. Prod. iv. 249 (= *Bunium creticum*, Urv.).—Oriens.
nodosa, Boiss. Fl. Orient. ii. 878 (= *Elaeosticta nodosa*).—Persia.
podagrarioides, Boiss. l. c. 876.—As. Min.
rotundifolia, Boiss. l. c. = Pimpinella rotundifolia.
Tournefortii, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 70 (= *Bunium creticum*, Urv.).—Graecia.

SCALIOPSIS, Walp. in Linnaea, xiv. (1840) 318 = **Podolepis**, Labill. (Compos.).
Lucaeana, Walp. l. c. = P. longipedata.

SCANDALIDA, Adans. Fam. ii. 326 (1763) = **Lotus**, Tourn. (Legumin.).
flava, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384 = L. siliquosus.
rubra, Medic. l. c. = L. Tetragonolobus.
siliquosa, Medic. Phil. Bot. i. 205 = L. siliquosus.
tetragonoloba, Medic. l. c. = L. Tetragonolobus.

SCANDEDERIS, Thou. Orch. Isles Afr. 3me Tabl. (1822) = SCAREDEDERIS, Thou. = **Dendrobium**, Sw. (Orchid.).
scandens, Thou. l. c.—Ins. Maurit.

SCANDIX, Tourn. ex Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 899.
CYCLOTAXIS, Boiss. Diagn. Ser. I. x. 48 (1849).
DASYSPERMUM, Neck. Elem. i. 276 (1790), partim.
PECTEN, Lam. Fl. Fr. iii. 437 (1778).
WYLIA, Hoffm. Gen. Umb. 3. t. 2 (1814).
aegyptiaca, Hort. ex Steud. Nom. ed. II. ii. 528 (Quid?).—Hab.?
affinis, Hort. ex Boiss. l. c. ii. 915 = macrorrhyncha.
affinis, C. Koch, ex Boiss. Fl. Orient. ii. 917 = grandiflora.
Anthriscus, Linn. Sp. Pl. 257 = Anthriscus vulgaris.
apiculata, DC. Prod. iv. 221 = iberica.
aromatica, Wahlenb. in Oken, Isis, xxi. (1826) 984 = Chaerophyllum aromaticum.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 58.—Oriens; Afghan.
aurea, Roth, Tent. Fl. Germ. i. 123 = Chaerophyllum aureum.
australis, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 80 = microcarpa.
australis, *Linn. Sp. Pl.* 257.—Reg. Mediterr.
Balansae, *Reut. Cat. Hort. Genev.* (1867) 2.—Cilicia.
brachycarpa, Boiss. Fl. Orient. ii. 916 = Balansae.
brachycarpa, *Guss. Ind. Sem. Hort. Boccad.* (1825) 10; *Fl. Sic. Prod.* i. 350.—Sicil.; Syria.
brevirostris, *Boiss. & Reut. Cat. Hort. Genev.* (1868) 4.—Hispan.
Bulbocastanum, Moench, Meth. 101 = Carum Bulbocastanum.
bulbosa, Roth, Tent. Fl. Germ. i. 123 = Chaerophyllum bulbosum.
Cerefolium, Linn. Sp. Pl. 257 = Anthriscus Cerefolium.
chilensis, *Molina, Sagg. Chil.* 125.—Chili.
clavata, Banks & Soland. ex Hook. Fl. Antarct. ii. 288 = Osmorhiza chilensis.

SCANDIX :—
clavata, DC. Prod. iv. 283, in syn., sphalm. = Chaerophyllum macrospermum.
cornuta, Gilib. Fl. Lituan. ii. 27 = S. Pecten-Veneris.
cretica, Mill. Gard. Dict. ed. VIII. n. 4 = S. Pecten-Veneris.
dulcis, Muhl. Cat. 41; DC. Prod. iv. 232 = Osmorhiza brevistylis.
falcata, Bieb. Fl. Taur. Cauc. i. 230 = iberica.
falcata, Londes, in Mém. Soc. Nat. Mosc. i. (1811) 31 = grandiflora.
fumarioides, Chaub. & Bory, Nouv. Fl. Pélop. 19 = Anthriscus tenerrimus.
fumarioides, Waldst. & Kit. Pl. Rar. Hung. iii. 249 = Anthriscus fumarioides.
georgica, C. Koch, ex Boiss. Fl. Orient. ii. 915 = iberica.
gilanica, S. G. Gmel. Reise, iii. 303. t. 31. f. 2 = Chaerophyllum macrospermum.
glaberrima, Desf. Fl. Atlant. i. 260 (= *Bunium glaberrimum*).—Alger.
glochidiata, Labill. Nov. Holl. Pl. i. 75. t. 102 = Daucus brachiatus.
grandiflora, *Linn. Sp. Pl.* 257.—Europ. or.; As. Min.
hirsuta, Biasol. ex Steud. Nom. ed. II. ii. 528 = grandiflora.
hirsuta, Scop. Fl. Carn. ed. II. i. 211 = Chaerophyllum hirsutum.
hispanica, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 57 = macrorhyncha.
hispanica, Cutand. Fl. Matr. 340 = S. Pecten-Veneris.
hispida, Gilib. Fl. Lituan. ii. 27 = Anthriscus vulgaris.
hispidula, Bertol. in Nov. Comm. Acad. Bologn. v. (1862) 427 = pinnatifida.
iberica, *Bieb. Fl. Taur. Cauc.* i. 425.—As. Min.; Persia.
iberica, Heldr. ex Boiss. Diagn. Ser. I. x. 50 = pisidica.
inferta, Jacq. ex Steud. Nom. ed. II. ii. 528, sphalm. = seq.
infesta, Linn. Syst. ed. XII. 732 = Caucalis infesta.
laeta, Salisb. Prod. 167 = Anthriscus vulgaris.
latifolia, Sibth. & Sm. Fl. Graec. iii. 77. t. 284 = Lecokia cretica.
macrorhyncha, *C. A. Mey. in Ind. Sem. Hort. Petrop.* ix. 86.—As. Min.
macrosperma, Willd. ex Schult. Syst. vi. 507 = Chaerophyllum macrospermum.
microcarpa, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 42.—Hispan.
millefolia, Willd. ex Spreng. Syst. i. 903 = Chaerophyllum millefolium.
nodosa, Linn. Sp. Pl. 257 = Chaerophyllum nodosum.
nutans, Moench, Meth. 101 = Chaerophyllum temulum.
odorata, Linn. Sp. Pl. 257 = Myrrhis odorata.
orientalis, Griseb. Spicil. Fl. Rumel. i. 369 = grandiflora.
palaestina, *Boiss. Fl. Orient.* ii. 918.—Palaestin.
parviflora, Retz. f. Pugill. 12 = pinnatifida?
Pecten, Scop. Fl. Carn. ed. II. i. 211 = seq.
Pecten-Veneris, *Linn. Sp. Pl.* 256 (Pecten ♀).—Europ.; Oriens.
pectinifera, Stokes, Bot. Mat. Med. ii. 122 = seq.
pectiniformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70, 134 = praec.
persica, Mart. in Linnaea, xii. (1838) Litt. 87 = S. Pecten-Veneris.
petraea, Auct. ex Steud. Nom. ed. II. ii. 528 = australis.
pinnatifida, *Vent. Jard. Cels*, t. 14.—Reg. Mediterr.; Oriens.
pinnatifida, Willk. in Flora, xxxiv. (1851) 728 = S. Pecten-Veneris.
pisidica, *Boiss. & Heldr. Diagn.* Ser. I. x. 49.—As. Min.
procumbens, Linn. Sp. Pl. 257 = Chaerophyllum procumbens.
radiata, Salisb. Prod. 167 = australis.
rostrata, Salisb. l. c. 166 = S. Pecten-Veneris.
Russeliana, Griseb. Spicil. Fl. Rumel. i. 369 = pinnatifida.
Stella, Steud. Nom. ed. II. ii. 528, sphalm. = seq.
stellata, [Soland. in] Russ. Aleppo, ed. II. ii. 249 = pinnatifida.
taurica, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1846) II. 355.—Tauria.
temula, Roth, Tent. Fl. Germ. i. 122 = Chaerophyllum temulum.

SCANDIX :—
tenuifolia, Salisb. Prod. 166 = Anthriscus Cerefolium.
ternata, Moench, Meth. 101 = Cryptotaenia canadensis.
tinctoria, Scop. Fl. Carn. ed. II. i. 212 = Chaerophyllum aromaticum.
trichosperma, Linn. Mant. i. 57 = Caucalis trichosperma.
trichosperma, Schur, Enum. Pl. Transs. 963 (Index) = Anthriscus Cerefolium.
turgida, *Boiss. Fl. Orient.* ii. 918.—Cilicia.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 503 = S. Pecten-Veneris.
Wilhelmsii, C. Koch, ex Boiss. Fl. Orient. ii. 915 = S. Pecten-Veneris.

SCAPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Saurauja**, Willd. (Ternstr.).
acuminata, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 121 = Ternstroemia acuminata.
Candollii, Choisy, l. c. = Saur. tristyla.
cauliflora, Noronha, ex DC. Prod. i. (1821) 425 = Saur. cauliflora.
elegans, Choisy, l. c. xiv. (1855) 121.—Ins. Philipp.
ferox, Choisy, l. c. = Saur. ferox.
gigantea, Noronha, ex DC. l. c. i. (1821) 424 = Saur. gigantea.
leucophloia, Choisy, l. c. xiv. (1855) 121 = Saur. leucophloia.
nigrescens, Choisy, l. c. = Saur. nigrescens.
pinangiana, Choisy, l. c. = Saur. tristyla.
rubiginosa, Choisy, l. c. = Saur. Jackiana.
serrata, Choisy, l. c. = Ternstroemia serrata.
spadicea, Choisy, l. c. = Saur. spadicea.
subcordata, Choisy, l. c. = Saur. subcordata.

SCAPHESPERMUM, Edgew. in Trans. Linn. Soc. xx. (1846) 58 = **Seseli**, Linn. (Umbell.).
trilobum, Edgew. l. c. = Seseli trilobum.

SCAPHISPATHA, Brongn. ex Schott, Prod. Aroid. 214 (1860). *AROIDEAE*, Benth. & Hook. f. iii. 978.
gracilis, *Brongn. ex Schott, l. c.*—Bolivia.

SCAPHIUM, Schott & Endl. Melet. 33 (1832) = **Sterculia**, Linn.
Wallichii, Schott & Endl. l. c. = St. scaphigera.

SCAPHOPETALUM, Mast. in Journ. Linn. Soc. x. (1869) 28. *STERCULIACEAE*, Benth. & Hook. f. i. 983.
Blackii, *Mast. l. c.* 29. t. 2.—Afr. trop.
longepedunculatum, *Mast. l. c.* 28. t. 3.—Afr. trop.
Mannii, *Mast. l. c.* 29. t. 3.—Afr. trop.

SCAPHYGLOTTIS, Poepp. & Endl. Nov. Gen. et Sp. i. 58 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 525.
CLADOBIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
affinis, *Poepp. & Endl. Nov. Gen. et Sp.* i. 59.—Peruv.
Behrii, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 219.—Panama.
conferta, *Poepp. & Endl. Nov. Gen. et Sp.* i. 59.—Peruv.
dubia, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 219.—Mexic.
fasciculata, *Hook. Ic. Pl.* t. 317.—Guian.
graminifolia, *Poepp. & Endl. Nov. Gen. et Sp.* i. 59.—Peruv.
Kienastii, *Hemsl. Biol. Centr. Am. Bot.* iii. 219.—Mexic.
leucantha, *Reichb. f. in Linnaea*, xxii. (1849) 856.—Bras.
mesocopis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 220.—Costa Rica.
pallidiflora, *Lindl. in Sweet, Hort. Brit.* ed. III. 646.—Ins. Trinit.
parviflora, *Poepp. & Endl. Nov. Gen. et Sp.* i. 58.—Peruv.
pendula, *Poepp. & Endl. l. c.*—Peruv.
reflexa, *Lindl. Bot. Reg.* (1839) *Misc.* 20.—Guian.
rosea, Hook. Ic. Pl. t. 313 = violacea.
ruberrima, Lindl. Orch. Linden. 22 = Ornithidium ruberrimum.

SCAPHYGLOTTIS:—
stellata, *Lodd. ex Lindl. Bot. Reg.* (1889) *Misc.* 44.—Guian.
Tafallae, *Reichb. f. in Linnaea*, xxii. (1849) 855.—Peruv.
violacea, *Lindl. Bot. Reg.* t. 1901.—Guian.

SCAREDEDERIS, Thou. Orch. Iles Afr. t. 91 (1822) = **Dendrobium**, Sw. (Orchid.).
scandens, Thou. l. c. (Quid?).—Ins. Maurit.

SCARIOLA, F. W. Schmidt, Samml. Phys. Aufs. i. 270 (1795) = **Lactuca**, Linn. (Compos.).
viminea, F. W. Schmidt, l. c. = L. viminea.

SCELOCHILUS, Klotzsch, in Otto & Dietr. Allg. Gartenz. ix. (1841) 261. *ORCHIDEAE*, Benth. & Hook. f. iii. 558.
heterophyllus, *Reichb. f. in Linnaea*, xli. (1877) 105.—Ecuador.
Jamiesoni, *Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 88.—Ecuador.
Lindenii, *Lindl. & Paxt. l. c.*—Venezuela.
Ottonis, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* ix. (1841) 261.—Venezuela.
saccatus, *Reichb. f. in Walp. Ann.* vi. 689.—Peruv.
stenochilus, Reichb. f. l. c. = Lindenii.

SCEPA, Lindl. Introd. Nat. Syst. ed. II. 441 (1836) = **Aporosa**, Blume (Euphorb.).
aurita, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 254 = A. aurita.
chinensis, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 72 = A. microcalyx.
Lindleyana, Wight, Ic. t. 361 = A. Lindleyana.
microstachya, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 256 = A. microstachya.
stipulacea, Lindl. Introd. Nat. Syst. ed. II. 441 = A. microstachya.
villosa, Lindl. l. c. = A. villosa.

SCEPASMA, Blume, Bijdr. 582 (1825) = **Phyllanthus**, Linn. (Euphorb.).
buxifolia, Blume, l. c. = P. buxifolius.
longifolia, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1848) 143 = P. acutissimus.
parvifolia, Reichb. f. & Zoll. ex Teijsm. & Binn. Cat. Hort. Bog. 229.—Malaya.

SCEPINIA, Neck. Elem. i. 78 (1790) = **Pteronia**, Linn. (Compos.).
dichotoma, Cass. in Dict. Sc. Nat. xlviii. 45 = P. oppositifolia.
lepidophylla, Cass. l. c. = P. glomerata.

SCEPOCARPUS, Wedd. in DC. Prod. xvi. i. 98 (1869) = **Urera**, Gaudich. (Urtic.).
Mannii, Wedd. l. c.—Afr. trop.

SCEPSEOTHAMNUS, Cham. in Linnaea, ix. (1834) 248 = **Alibertia**, A. Rich. (Rubiac.).
gardenioloides, Cham. l. c. 249.—Bras.
rotunda, Cham. l. c. 250.—Bras.
vinosa, Cham. l. c. 249 = *rotunda*.

SCEPSOTHAMNUS, Steud. Nom. ed. II. ii. 528 (1841), sphalm. = praec.

SCEPTRANTHES, R. Grah. in Edinb. N. Phil. Journ. xx. (1836) 413 = **Cooperia**, Herb. (Amaryll.).
Drummondii, R. Grah. l. c. = C. paniculata.

SCEPTRANTHUS, Benth. & Hook. f. Gen. iii. 723 (1883), err. typ. = praec.

SCEPTROCNIDE, Maxim. in Bull. Acad. Pétersb. xxii. (1877) 238. *URTICACEAE*, Benth. & Hook. f. ii. 382.
macrostachya, *Maxim. l. c.* 240.—Japon.

SCEPTRUM-CAROLINUM, Linn. Fl. Lapp. 197 (1737) = **Pedicularis**, Linn. (Scrophul.).

SCEURA, Forsk. Fl. Aegypt. Arab. 37 (1775) = **Avicennia**, Linn. (Verben.).
marina, Forsk. l. c. = A. officinalis.

SCHACHTIA, Karst. in Linnaea, xxx. (1859–60) 156. *RUBIACEAE*, Benth. & Hook. f. ii. 80.
dioica, *Karst. l. c.* 157.—N. Granat.

SCHAEFFERA, Schreb. Gen. ii. 681 (1791), sphalm. = seq.

SCHAEFFERIA, Jacq. Enum. Pl. Carib. 10 (1760). *CELASTRINEAE*, Benth. & Hook. f. i. 367.
Berterii, *Griseb. Cat. Pl. Cub.* 56.—Ins. S. Doming.
buxifolia, Nutt. Sylv. N. Am. ii. 42. t. 56 = frutescens.
completa, Sw. Prod. Veg. Ind. Occ. 38 = frutescens.
cuneifolia, *A. Gray, Pl. Wright.* i. 35.—Texas.
frutescens, *Jacq. Enum. Pl. Carib.* 33; *Select. Am.* 259; *Karst. Fl. Columb.* i. 183.—Am. bor.; Ind. occ.
lateriflora, *Sw. Prod. Veg. Ind. Occ.* 38.—Ins. S. Doming.
Marchii, *Griseb. Fl. Brit. W. Ind.* 710.—Jamaic.
paniculata, *Spreng. Neue Entdeck.* iii. 49.—Bras.
racemosa, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 41.—Mexic.
viridescens, [Moç. & Sesse,] ex DC. l. c. = frutescens.

SCHAEFFNERA, Benth. & Hook. f. Gen. ii. 492 (1873), sphalm. = SCHAFFNERA, Sch. Bip. = **Dicoma**, Cass. (Compos.).

SCHAENOLAENA, Lindl. Veg. Kingd. 778 (1847), sphalm. = SCHOENOLAENA, Bunge = **Xanthosia**, Rudge (Umbell.).

SCHAENOPRASUM, Franch. & Sav. Enum. Pl. Jap. i. 76 (1879), sphalm. = SCHOENOPRASUM, H. B. & K. = **Allium**, Tourn. (Liliac.).

SCHAENUS, Gouan, Fl. Monsp. 388 (1765), err. typ. = **Schoenus**, Linn. (Cyperac.).

SCHAETZELIA, Sch. Bip. in Walp. Ann. ii. 949 (1852) = **Schaetzellia**, Sch. Bip. (Compos.).

SCHAETZELLIA, Klotzsch, in Otto & Dietr. Allg. Gartenz. xvii. (1849) 82 = **Onoseris**, DC. (Compos.).
Deckeri, Klotzsch, l. c. = O. Isotypus.

SCHAETZELLIA, Sch. Bip. in Flora, xxxiii. (1850) 419. *COMPOSITAE*, Benth. & Hook. f. ii. 242.
DICHAETA, Sch. Bip. l. c. xxxiii. (1850).
SCHAETZELIA, Sch. Bp. in Walp. Am. ii. 949 (1852).
mexicana, *Sch. Bip. in Flora*, xxxiii. (1850) 419.—Mexic.

SCHAETZELLIA, Sch. Bip. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 192, nomen = **Hinterhubera**, Sch. Bip. (Compos.).
columbica, Sch. Bip. l. c. = H. columbica.

SCHAFFNERA, Benth. in Journ. Linn. Soc. xix. (1881) 63. *GRAMINEAE*, Benth. & Hook. f. iii. 1124.
gracilis, *Benth. in Hook. Ic. Pl.* xiv. 59. t. 1378.—Mexic.

SCHAFFNERA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 42 = **Dicoma**, Cass. (Compos.).
carduoides, Sch. Bip. l. c. = D. tomentosa.

SCHANGINIA, C. A. Mey. in Ledeb. Fl. Alt. i. 394 (1829) = **Suaeda**, Forsk. (Chenop.).
altissima, Hohen. ex Ledeb. Fl. Ross. iii. 781 = Suaeda heterocarpa.
altissima, C. A. Mey. Verz. Pfl. Cauc. 159 = Suaeda altissima.
Arbuscula, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 733.—Sibir. altaic.
baccata, Moq. Chenop. Enum. 119 = Suaeda baccata.
baccifera, Ledeb. Fl. Ross. iii. 776 = Suaeda baccifera.
Borszczowii, Bunge, ex Trautv. in Act. Hort. Petrop. ix. (1884) 131, nomen.—Reg. Ural.
hortensis, Moq. Chenop. Enum. 119 = Suaeda hortensis.
inderiensis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 460.—Reg. Ural.

SCHANGINIA :—
linifolia, C. A. Mey. in Ledeb. Fl. Alt. i. 395 = Suaeda linifolia.
linifolia, Moq. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 320. t. 23 = Suaeda hortensis.
tribracteata, C. Koch, in Linnaea, xxii. (1869) 188.—Rossia.
trigyna, Heynh. Nom. ii. 640 = Suaeda trigyna.

SCHAPHESPERMUM, Edgew. in Proc. Linn. Soc. i. (1845) 253, sphalm. = SCAPHESPERMUM, Edgew. = **Seseli**, Linn. (Umbell.).

SCHAUERA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 202, in adnot. (1836) = **Aydendron**, Nees (Laurin.).

SCHAUERIA, Hassk. in Flora, xxv. (1842) II. Beibl. 25 = **Hyptis**, Jacq. (Labiat.).
graveolens, Hassk. l. c. = H. suaveolens.

SCHAUERIA, Nees, Ind. Sem. Hort. Vratisl. (1838). —Cf. Linnaea, xiii. (1839) Litt. 119. *ACANTHACEAE*, Benth. & Hook. f. ii. 1116.
FLAVICOMA, Rafin. Fl. Tellur. iv. 63 (1836).
caduciflora, *Griseb. in Goett. Abh.* xxiv. (1879) 261.—Reg. Argent.
calicotricha, *Nees, Ind. Sem. Hort. Vratisl.* (1838).—*Cf. Linnaea*, xiii. (1839) *Litt.* 119.—Bras.
capitata, *Nees, in DC. Prod.* xi. 316.—Bras.
decipiens, *Nees, l. c.* 318.—Mexic.
flavicoma, *N. E. Br. in Gard. Chron.* (1883) I. 14.—Bras.
gonyostachya, *Nees, in Mart. Fl. Bras.* ix. 106.—Bras.
hirsuta, *Nees, l. c.* 103.—Bras.
humuliflora, *Nees, l. c.* 102.—Bras.
lachnostachya, *Nees, l. c.* 105.—Bras.
linearifolia, Torr. Bot. Mex. Bound. 123 = Dianthera linifolia.
linifolia, Benth. & Hook. f. Gen. ii. 1114 = praec.
lophura, *Nees, in Mart. Fl. Bras.* ix. 104.—Bras.
macrophylla, *Nees, l. c.* 105.—Bras.
malifolia, *Nees, l. c.* 103.—Bras.
marginata, *Nees, l. c.* 102.—Bras.
Maximiliani, *Nees, l. c.* 106.—Bras.
paniculata, *Nees, l. c.*—Bras.
parvifolia, Torr. Bot. Mex. Bound. 122 = Dianthera parvifolia.
Schottii, *Nees, in Mart. Fl. Bras.* ix. 107.—Bras.
sulphurea, *Nees, l. c.* 104.—Bras.
virginea, *Nees, l. c.* 105.—Bras.

SCHEADENDRON, Bertol. f. in Mem. Acad. Sc. Bologn. ii. (1850) 572, sphalm. = SHEADENDRON, Auct. = **Combretum**, Linn.

SCHEDONNARDUS, Steud. in Flora, xxxiii. (1850) 228, 229, nomen; Syn. Pl. Gram. 146 (1855). *GRAMINEAE*, Benth. & Hook. f. iii. 1167.
texanus, *Steud. ll. cc.*—Texas.

SCHEDONORUS, Beauv. Agrost. 99. t. 19. f. 2 (1812) = **Festuca**, Linn. et **Bromus**, Linn. (Gramin.).
alpinus, Hoppe, Agrost. Dec. 4; ex Schult. Mant. ii. 389 = F. montana.
altissimus, Beauv. Agrost. 99 = F. sylvatica.
americanus, Roem. & Schult. Syst. ii. 706 = F. poaeoides.
arundinaceus, Dum. Obs. Gram. Belg. 106 = F. elatior.
arundinaceus, Roem. & Schult. Syst. ii. 700 = Graphephorum arundinaceum.
asper, Gaud. ex Roem. & Schult. Syst. ii. 643 = B. asper.
auratus, Beauv. Agrost. 99 = F. violacea.
aureus, Beauv. l. c. = F. spadicea.
Benekeni, Lange, in Fl. Dan. t. 2826 = B. asper.
Beneki, Lange, in Bull. Soc. Bot. Belg. xiii. (1874) 310, sphalm. = praec.
Billardierianus, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 419 = F. littoralis.
bonariensis, Roem. & Schult. Syst. ii. 709 = Poa pallens.
bromoides, Wight, ex Steud. Syn. Pl. Gram. 316 = F. scaberrima, Wightiana.
caerulescens, Roem. & Schult. Syst. ii. 699 = F. caerulescens.
calamarius, Beauv. Agrost. 99 = F. sylvatica.
Caldasii, Roem. & Schult. Syst. ii. 709 = F. Caldasii.
ciliatus, Kjellm. in Nordensk. Vega-Exped. Vetensk. Iakтt. i. 557 = B. ciliatus.
compressus, Roem. & Schult. Syst. ii. 700 = F. spadicea.

SCHEDONORUS:—
consobrinus, Timb. in Bull. Soc. Hist. Nat. Toul. iii. (1869) 130 = Festuca spadicea.
curvatus, Beauv. Agrost. 99 = F. elatior.
deciduus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 115 = F. sylvatica.
divaricatus, Roem. & Schult. Syst. ii. 703 = Cutandia divaricata.
dumetorum, Beauv. Agrost. 99 = F. rubra.
elatior, Beauv. l. c. = F. elatior.
erectus, Gaud. ex Roem. & Schult. Syst. ii. 637 = B. erectus.
Eskia, Beauv. Agrost. 99 = F. varia.
exaltatus, Schult. Mant. iii. 644 = F. montana.
ferrugineus, Vest, in Syll. Ratisb. ii. (1828) 221 = F. spadicea.
Gerardi, Beauv. Agrost. 99 (Quid ?).—Europ.
giganteus, Gaud. ex Roem. & Schult. Syst. ii. 644 = Bromus giganteus.
glaucus, Beauv. Agrost. 99 = F. glauca.
Hookerianus, Benth. Fl. Austral. vii. 656 = F. Hookeriana.
inermis, Beauv. Agrost. 99 = B. inermis.
lanatus, Roem. & Schult. Syst. ii. 708 = F. mollis.
latifolius, Roem. & Schult. l. c. 710 = F. latifolia.
littoralis, Beauv. Agrost. 99: Roem. & Schult. l. c. 707 = F. littoralis.
lividus, Roem. & Schult. l. c. = F. livida.
loliaceus, Beauv. Agrost. 99 = *F. loliacea.*
longifolius, Trin. ex Steud. Nom. ed. II. ii. 529 = B. inermis.
modesta, Nees, ex Steud. Syn. Pl. Gram. 316 = F. modesta.
montana, Opiz, Seznam, 89 = F. montana.
nigrescens, Beauv. Agrost. 99 = F. rubra.
nitidus, Roem. & Schult. Syst. ii. 706 = F. rubra.
nutans, Roem. & Schult. l. c. 704 = F. pulchella.
obtusus, Roem. & Schult. l. c. 710 = F. nutans.
patulus, Roem. & Schult. l. c. 710 = F. patula.
phoenicoides, Roem. & Schult. l. c. 699 = Brachypodium ramosum.
pilosus, Beauv. Agrost. 99 = Poa violacea.
poaeformis, Beauv. l. c. = Poa violacea.
pratensis, Beauv. l. c. = F. elatior.
procerus, Roem. & Schult. Syst. ii. 708 = F. eminens.
pulchellus, Beauv. Agrost. 99 = F. pulchella.
pumilus, Roem. & Schult. Syst. ii. 706 = F. varia.
quadridens, Roem. & Schult. l. c. 710 = Triodia cuprea.
radicans, Dum. Obs. Gram. Belg. 106 = F. elatior.
remotiflorus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 283 = F. pauciflora ?
rhaeticus, Beauv. Agrost. 99 = Poa violacea.
scaber, Nees, ex Steud. Nom. ed. II. ii. 529 = F. Neesiana.
Scheuchzeri, Beauv. Agrost. 99 = F. pulchella.
scirpoideus, Benth. Fl. Austral. vii. 655 = F. scirpoidea.
serotinus, Beauv. Agrost. 99 = Diplachne serotina.
serotinus, Rostrup, in Kjoeb. Vidensk. Meddel. 1864 (1864), 119 = B. asper.
spadiceus, Roem. & Schult. Syst. ii. 700 = F. spadicea.
spicatus, Roem. & Schult. l. c. 707 = F. spicata.
sterilis, Fries, Summ. Veg. 76 = B. sterilis.
sylvaticus, Beauv. Agrost. 99 = F. sylvatica.
tectorum, Fries, Summ. Veg. 76 = B. tectorum.
tenellus, Beauv. Agrost. 99 = F. tenella, *Willd.*
tenuifolius, Andrz. ex Trautv. in Act. Hort. Petrop. ix. (1884) 327, nomen (Quid ?).—Rossia.
trichodes, Schult. Mant. iii. 645 = Eragrostis mayurensis.
triflorus, Roem. & Schult. Syst. ii. 709 = F. triflora.
unioloides, Roem. & Schult. l. c. 708 = B. unioloides.
varius, Beauv. Agrost. 99 = F. varia.
versicolor, Opiz, Seznam, 89 = F. varia.
violaceus, Beauv. Agrost. 99 = F. rubra.
violaceus, Link, Hort. Berol. i. 171 = Poa violacea.

SCHEELEA, Karst. in Linnaea, xxviii. (1856) 264. *PALMAE*, Benth. & Hook. f. iii. 947.
attaleoides, *Karst. l. c.* 265; *Fl. Columb.* i. 135. t. 67.—N. Granat.
butyracea, *Karst. ex H. Wendl. in Kerch. Palm.* 256.—Hab. ?
Cephalotes, *Karst. in Linnaea*, xxviii. (1856) 269.—Bras.
excelsa, *Karst. l. c.* 267.—N. Granat.
imperialis, *Hort.—Cf. Gard. Chron.* (1875) I. 589.—N. Granat.

SCHEELEA:—
insignis, *Karst. in Linnaea*, xxviii. (1856) 269.—Bras.
macrocarpa, *Karst. l. c.* 268.—Venezuela.
Maripa, *Hort. ex H. Wendl. in Kerch. Palm.* 256, nomen.—Hab. ?
Princeps, *Karst. in Linnaea*, xxviii. (1856) 269.—Bras.
regia, *Karst. l. c.* 266; *Fl. Columb.* ii. 145. t. 167.—N. Granat.

SCHEERIA, Seem. in Bot. Mag. t. 4743 (1853) = **Achimenes**, P. Br. (Gesner.).
ichthyostoma, Seem. Bot. Voy. Herald, 185 = A. ichthyostoma.
lanata, Hanst. in Linnaea, xxvii. (1854) 714 = A. lanata.
mexicana, Seem. in Bot. Mag. t. 4743 = A. Scheerii.
panamensis, Seem. Bot. Voy. Herald, 185 = A. panamensis.

SCHEFFIELDIA, Scop. Introd. 183 (1777), sphalm. = SHEFFIELDIA, Forst. = **Samolus**, Linn. (Primul.).

SCHEFFLERA, Forst. Char. Gen. 45. t. 23 (1775). *ARALIACEAE*, Benth. & Hook. f. i. 940.
affinis, *Baill. Adansonia*, xii. (1876–79) 141.—N. Caled.
Andraeana, *Baill. l. c.*—N. Caled.
apioidea, *Baill. l. c.* 145.—N. Caled.
Balansaeana, *Baill. l. c.* 142.—N. Caled.
Candelabrum, *Baill. l. c.* 146.—N. Caled.
crassipes, *Baill. l. c.* 144.—N. Caled.
Cunninghami, *Miq. in Linnaea*, xviii. (1844) 89.—N. Zel.
Cussoniae, *Baill. Adansonia*, xii. (1876–79) 143.—N. Caled.
digitata, *Forst. Char. Gen.* 46.—N. Zel.
dipyrena, Baill. Hist. Pl. vii. 162 = Heptapleurum dipyrenum.
elongatum, *Baill. Adansonia*, xii. (1876–79) 144.—N. Caled.
Emiliana, *Baill. l. c.* 143.—N. Caled.
Fagueti, *Baill. l. c.* 142.—N. Caled.
Gabriellae, *Baill. l. c.* 144.—N. Caled.
Golip, *Baill. l. c.* 142.—Ins. Lifu.
indivisa, *Baill. l. c.* 140.—N. Caled.
Marcellana, *Baill. l. c.*—N. Caled.
Nono, *Baill. l. c.* 141.—N. Caled.
Pancheri, *Baill. l. c.* 143.—N. Caled.
parvifolia, *Baill. l. c.* 144.—N. Caled.
stellata, Baill. Hist. Pl. vii. 161 = Heptapleurum stellatum.
Toto, *Baill. Adansonia*, xii. (1876–79) 140.—N. Caled.
umbellifera, Baill. Hist. Pl. vii. 162 = Cussonia umbellifera.
Vieillardi, *Baill. Adansonia*, xii. (1876–79) 146.—N. Caled.
vitiensis, *Seem. in Journ. Bot.* iii. (1865) 176.—Ins. Fiji.

SCHEIDWEILERIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 123; et in Abh. Akad. Berl. 1854 (1855), 58. t. 46 = **Begonia**, Linn.
digitata, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 123 = B. digitata.
ferrata, Klotzsch, l. c. 123, nomen (an seq. ?).—Bras.
inciso-serrata, Klotzsch, in Abh. Akad. Berl. 1854 (1855), 61 = B. inciso-serrata.
luxurians, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 123 = B. luxurians.
muricata, Klotzsch, l. c. = B. muricata.
parviflora, Klotzsch, in Abh. Akad. Berl. 1854 (1855), 59 = B. parviflora.
repens, Hassk. Hort. Bog. Desc. 325 = B. mollis.
Sellowiana, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 123, nomen.—Bras.

SCHELHAMERIA, Heist. Syst. 9 (1748) = **Cheiranthus**, Linn. (Crucif.).

SCHELHAMMERA, R. Br. Prod. 273 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 829.
PARDUYNA, Salisb. Gen. Pl. Fragm. 58 (1866).
glandulifera, Loisel. ex Steud. Nom. ed. II. ii. 529 (Quid ?).—Hab. ?
multiflora, *R. Br. Prod.* 274.—Austral.
multiflora, Lodd. Bot. Cab. t. 1511 = Kreysigia multiflora.
undulata, *R. Br. Prod.* 274.—Austral.

SCHELHAMMERIA, Moench, Meth. Suppl. 119 (1802) = **Carex**, Dill. (Cyperac.).
capitata, Moench, l. c. = C. cyperoides.
cyperoides, Dum. Fl. Belg. 145; Reichb. Fl. Germ. Excurs. 72 = C. cyperoides.

SCHELLINGIA, Steud. in Flora, xxxiii. (1850) 231. t. 1 = **Aegopogon**, Beauv. (Gramin.).
tenera, Steud. l. c. 232. t. 1 = A. geminiflorus.

SCHELVERIA, Nees & Mart. in Flora, iv. (1821) 299, 328 = **Angelonia**, Humb. & Bonpl. (Scrophul.).
arguta, Nees & Mart. l. c. = A. procumbens.

SCHENKIA, Griseb. in Bonplandia, i. (1853) 226 = **Erythraea**, Rich. (Gentian.).
sebaeoides, Griseb. l. c. = E. sebaeoides.

SCHENODORUS, Beauv. Agrost. 177 (Index) (1812); Roem. & Schult. Syst. ii. 698 (1817).—Cf. Poir. Encyc. Suppl. v. 84 (1817) = SCHEDONORUS, Beauv. = **Festuca**, Linn. et **Bromus**, Linn. (Gramin.).

SCHEPERIA, Rafin. Sylva Tellur. 109 (1838), sphalm. = seq.

SCHEPPERIA, Neck. Elem. ii. 67 (1793) = **Cadaba**, Forsk. (Cappar.).
aphylla, Eckl. & Zeyh. Enum. 14 = C. juncea.
juncea, DC. Prod. i. 245 = C. juncea.

SCHERARDIA, Neck. Delic. Gallo-Belg. i. 82 (1768), sphalm. = **Sherardia**, Linn. (Rubiac.).

SCHETTI, Adans. Fam. ii. 146 (1763) = **Ixora**, Linn. (Rubiac.).

SCHEUCHLERIA, Heynh. Nom. i. 729 (1840), nomen. *COMPOSITAE* (Quid?).
patula, Heynh. l. c.—China.

SCHEUCHZERA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.

SCHEUCHZERIA, Linn. Syst. ed. I (1735). *NAIADACEAE*, Benth. & Hook. f. iii. 1012.
PAPILLARIA, Dulac, Fl. Hautes-Pyr. 45 (1867).
TELMATOPHILA, Ehrh. Beitr. iv. 147 (1789).
asiatica, Miq. Fl. Ind. Bat. iii. 243 = palustris.
palustris, *Linn. Sp. Pl.* 338.—Reg. bor. temp.
paniculata, Gilib. Exercit. ii. 502 = palustris.
Pseudo-asphodelus, Scop. Fl. Carn. ed. II. i. 263 = Tofieldia calyculata.
unicapsularis, Comm. ex Lam. Encyc. iii. 266 = Rostkovia grandiflora, *Hook. f.*

SCHEUKZERIA, Hill, Veg. Syst. xxvi. 13 (1774), sphalm. = praec.

SCHIDIOMYRTUS, Schau. in Linnaea, xvii. (1843) 237 = **Baeckea**, Linn. (Myrtac.).
crenulata, Schau. l. c. 237 = B. crenulata.
diosmifolia, Schau. l. c. = B. decemfida.
ericacea, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 148 = Micromyrtus microphylla.
micrantha, Schau. in Linnaea, xvii. (1843) 237 = Thryptomene baeckeacea.
Sieberi, Schau. l. c. = B. crenulata.
tenella, Schau. l. c. = Micromyrtus microphylla.

SCHIDOSPERMUM, Griseb. in Lechl. Berb. Am. Austr. 56 (1857) = **Chlorophytum**, Ker-Gawl. (Liliac.).
Sanseviera, Griseb. l. c. = C. Schidospermum.

SCHIECKEA, Karst. in Bot. Zeit. vi. (1848) 898. Genus dubium *SAPINDACEARUM*.—Cf. Benth. & Hook. f. Gen. i. 392.

SCHIECKIA, Benth. & Hook. f. Gen. iii. 674 (1883), sphalm. = **Schiekia**, Meissn. (Haemod.).

SCHIEDEA, Bartl. Ord. 210 (1830) = **Richardsonia**, Kunth (Rubiac.).
mexicana, Bartl. ex DC. Prod. iv. 569 = R. Haenkeana.

SCHIEDEA, Cham. & Schlecht. in Linnaea, i. (1826) 46. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
EUCLADUS, Nutt. ex Hook. Ic. Pl. t. 650 (1844).
amplexicaulis, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 310.—Ins. Sandvic.

SCHIEDEA :—
diffusa, *A. Gray, Bot. U. St. Expl. Exped.* i. 138.—Ins. Sandvic.
globosa, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 311.—Ins. Sandvic.
Hookeri, *A. Gray, Bot. U. St. Expl. Exped.* i. 133.—Ins. Sandvic.
Kaalae, *Wawra, in Flora*, lvi. (1873) 175.—Ins. Sandvic.
ligustrina, *Cham. & Schlecht. in Linnaea*, i. (1826) 46.—Ins. Sandvic.
Menziesii, *Hook. Ic. Pl.* sub t. 649–650.—Ins. Sandvic.
Nuttallii, *Hook. l. c.* t. 649–650.—Ins. Sandvic.
oahuensis, Wawra, in Flora, lvi. (1873) 175 = Nuttallii.
Remyi, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 310.—Ins. Sandvic.
spergulina, *A. Gray, Bot. U. St. Expl. Exped.* i. 135.—Ins. Sandvic.
stellarioides, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 310.—Ins. Sandvic.
viscosa, *H. Mann, l. c.* 311.—Ins. Sandvic.

SCHIEDEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 186 = **Tertrea**, DC. (Rubiac.).
martinicensis, A. Rich. l. c. 187 = T. martinicensis.

SCHIEKIA, Meissn. Gen. 397 (1842). *HAEMODORACEAE*, Benth. & Hook. f. iii. 674.
TROSCHELIA, Klotzsch, in Rich. Schomb. Reise Guian. 1066 (1848).
orinocensis, *Meissn. Gen. Comm.* 300.—Guian.

SCHIEVERECKIA, Nym. Consp. 54 (1878) = SCHIVERECKIA, Andrz. = **Alyssum**, Tourn. (Crucif.).

SCHILLERA, Reichb. Consp. 204 (1828) = **Eriolaena**, DC. (Stercul.).

SCHILLERIA, Kunth, in Linnaea, xiii. (1839) 609 = **Piper**, Linn.
abutiloides, Kunth, l. c. 721 = P. abutiloides.
aequalis, Kunth, l. c. 687 = P. aequale.
aurita, Kunth, l. c. 713 = P. auritum.
barbata, Kunth, l. c. 712 = P. barbatum.
callosa, Kunth, l. c. 689 = P. callosum.
Carpunya, Kunth, l. c. 693 = P. Carpunya.
catalpaefolia, Kunth, l. c. 718 = P. marginatum.
caudata, Kunth, l. c. 716 = P. marginatum.
Churumayu, Kunth, l. c. 706 = P. Churumayu.
colubrina, Kunth, l. c. 684 = P. colubrinum.
cornifolia, Kunth, l. c. 697 = P. cornifolium.
coruscans, Kunth, l. c. 708 = P. coruscans.
corylifolia, Kunth, l. c. 699 = P. dilatatum.
cumanensis, Kunth, l. c. 632 = P. cumanense.
grandifolia, Kunth, l. c. 709 = P. grandifolium.
halesiaefolia, Kunth, l. c. 681 = P. halesiaefolium.
heterophylla, Kunth, l. c. 688 = P. heterophyllum.
Hilariana, Kunth, l. c. 711 = P. Hilarianum.
inaequalifolia, Kunth, l. c. 631, in syn. = Peperomia inaequalifolium.
indecora, Kunth, l. c. 722 = P. indecorum.
Jussiaeana, Kunth, l. c. 680 = P. Jussiaeanum.
Karwinskiana, Kunth, l. c. 700 = P. Karwinskianum.
lanceaefolia, Kunth, l. c. 702 = P. lanceaefolium.
lapathifolia, Kunth, l. c. 714 = P. lapathifolium.
laurifolia, Kunth, l. c. 691 = P. laurinum.
lentaginoides, Kunth, l. c. 683 = P. lentaginoides.
leptura, Kunth, l. c. 679 = P. lepturum.
lineata, Kunth, l. c. 704 = P. lineatum.
longifolia, Kunth, l. c. 725 = P. longifolium.
Lucaeana, Kunth, l. c. 678 = P. Lucaeanum.
macrophylla, Kunth, l. c. 715 = P. macrophyllum.
marginata, Kunth, l. c. 718 = P. marginatum.
Martiana, Kunth, l. c. 691 = P. Martianum.
melastomoides, Kunth, l. c. 724 = P. melastomoides.
Mikaniana, Kunth, l. c. 719 = P. Mikanianum.
oblonga, Kunth, l. c. 696 = P. oblongum.
obovata, Kunth, l. c. 686 = P. obovatum.
persicariaefolia, Kunth, l. c. 684 = P. persicariaefolium.
pilulifera, Kunth, l. c. 722 = P. piluliferum.
populifolia, Kunth, l. c. 709 = P. Kunthianum.
riparia, Kunth, l. c. 707 = P. grandifolium.
rivinoides, Kunth, l. c. 682 = P. rivinoides.
rudis, Kunth, l. c. 703 = P. rude.
ulmifolia, Kunth, l. c. 698 = P. dilatatum.
umbrosa, Kunth, l. c. 726 = P. umbrosum.
variegata, Kunth, l. c. 695 = P. variegatum.
velutina, Kunth, l. c. 701 = P. velutinum.
xylosteoides, Kunth, l. c. 677 = P. xylosteoides.

SCHIMA, Auct. ex Steud. Nom. ed. II. ii. 530 (1841) = SEHIMA, Forsk. = **Ischaemum**, Linn. (Gramin.).
inscalpta, Hochst. ex Walp. Ann. iii. 803, 1135 = I. laxum.
macrostachyum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 472 = I. laxum.

SCHIMA, Reinw. ex Blume, Cat. Gew. Buitenz. 80 (1823). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 185.
antherisosa, *Korth. Verh. Nat. Gesch. Bot.* 145.—Sumatra.
bancana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 113.—Malaya.
crenata, Korth. Verh. Nat. Gesch. Bot. 143 = Noronhae.
excelsa, Blume, Cat. Gew. Buitenz. 80 = Gordonia excelsa.
hypoglauca, Miq. Fl. Ind. Bat. Suppl. 484 = Wallichii.
khasiana, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 289.—Reg. Himal.
mollis, *Dyer, l. c.* 288.—Burma.
monticola, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 93, nomen.—Burma.
Noronhae, *Reinw. ex Blume, Cat. Gew. Buitens.* 80; *Bijdr.* 130.—Malaya.
oblata, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65 = Noronhae.
rigida, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 113.—Java.
sulcinervia, *Miq. l. c.*—Sumatra.
superba, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 246 = Noronhae.
Wallichii, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 144 = Adinandra villosa.
Wallichii, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 144.—Reg. Himal.; Malaya.

SCHIMPERA, Hochst. ex Steud. in Endl. Gen. 889 (1839). *CRUCIFERAE*, Benth. & Hook. f. i. 96.
TRAILLIA, Lindl. ex Endl. Gen. Suppl. 1419 (1841).
arabica, *Hochst. & Steud. ex Steud. Nom.* ed. II. ii. 530.—Arab.
persica, *Boiss. Diagn.* Ser. I. vi. 18.—Persia.

SCHINNONGIA, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 223. *IRIDEAE* (Quid?).
ciliata, Schrank, l. c.—Afr. austr.

SCHINOPSIS, Engl. in Mart. Fl. Bras. xii. II. 403 (1876) = **Quebrachia**, Griseb. (Anacard.).
Balansae, Engl. Bot. Jahrb. vi. (1885) 286.—Paraguay.
brasiliensis, Engl. in Mart. Fl. Bras. xii. II. 404. t. 87.—Bras.
Haenkeana, Engl. l. c. 406.—Peruv.
Lorentzii, Engl. Bot. Jahrb. i. (1881) 46 = Q. Lorentzii.
marginata, Engl. in DC. Monog. Phan. iv. 464.—Reg. Argent.
peruviana, Engl. in Mart. Fl. Bras. xii. II. 405. t. 86. f. 2.—Peruv.

SCHINOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = seq.

SCHINUS, Linn. Gen. ed. I. 358 (1737). *ANACARDIACEAE*, Benth. & Hook. f. i. 422.
DUVAUA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 340.
MOLLE, [Tourn.] Adans. Fam. ii. 341 (1763).
SARCOTHECA, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 474.
Areira, Linn. Sp. Pl. 389 = S. Molle.
Aroeira, Vell. Fl. Flum. x. t. 135 = terebinthifolius.
aurantiodorus, Ruiz, ex Engl. in DC. Monog. Phan. iv. 326 = Mauria aurantiodora.
benghalensis, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 194 = Protium serratum.
bituminosus, Salisb. Prod. 171 = S. Molle.
Bonplandianus, March. Rev. Anacard. 164 = dependens.
Chichita, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1883) 96.—Reg. Argent.
chilensis, March. Rev. Anacard. 164 = latifolius.
crenatus, *Engl. Bot. Jahrb.* i. (1881) 423.—Chili.
dentatus, *Andr. Bot. Rep.* t. 620.—Ins. Sandvic.
dependens, *Orteg. Hort. Matr. Dec.* 102.—Bras.
discolor, *Benth. Bot. Voy. Sulph.* 11.—Calif.
Fagara, *Linn. Sp. Pl.* 389.—Jamaic.
Huygan, Molina, Sagg. Chil. ed. I. 169, 355 = dependens.

SCHINUS:—
indicus, Burm. f. Fl. Ind. 215 (err. typ. 315) (Quid?).—Ind. or.
latifolius, *Engl. in Mart. Fl. Bras.* xii. II. 389.—Chili.
lentiscifolius, *March. Rev. Anacard.* 164.—Bras.
leucocarpus, Mart. ex Engl. in Mart. Fl. Bras. xii. II. 394 = Lithraea Aroeirinha.
Limonia, Linn. Sp. Pl. 389 = Limonia acidissima.
Lithi, Domb. ex March. Rev. Anacard. 183 = Lithraea venenosa.
Mellisii, *Engl. Bot. Jahrb.* i. (1881) 423.—Ins. S. Helena.
Molle, Hort. ex Engl. in DC. Monog. Phan. iv. 346 = Lithraea Aroeirinha.
Molle, *Linn. Sp. Pl.* 388.—Am. trop.
molleoides, Vell. Fl. Flum. x. t. 134 (*moleoides*) = Lithraea Aroeirinha.
montanus, *Engl. Bot. Jahrb.* i. (1881) 422.—Chili.
mucronulata, Mart. Syst. Mat. Med. Bras. 52 = terebinthifolius.
myricoides, Linn. Sp. Pl. 388 = Cuscuta africana.
Niara, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 194 = Protium serratum.
Pearcei, *Engl. Bot. Jahrb.* i. (1881) 423.—Peruv.
procerus, March. Rev. Anacard. 164 = Cyrtocarpa procera.
pubescens, Spreng. ex Engl. in Mart. Fl. Bras. xii. II. 176 = Zanthoxylum rhoifolium.
rhoifolius, Mart. in Flora, xx. (1847) II. Beibl. 101 = terebinthifolius.
Saheria, Buch.-Ham. in Trans. Linn. Soc. xvii. (1835) 193 = Protium serratum.
sinuatus, *Engl. Bot. Jahrb.* i. (1881) 423.—Reg. Argent.
spinosus, *Engl. in Mart. Fl. Bras.* xii. II. 388.—Bras.
tenuifolius, Steud. Nom. ed. II. ii. 530, sphalm. = *ternifolius*.
terebinthifolius, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 399.—Bras.
ternifolius, Gill. ex Hook. Bot. Misc. iii. (1833) 177 = Lithraea Aroeirinha.
tragodes, Linn. Sp. Pl. 389 = Zanthoxylum tragodes.
virgatus, *Sweet, Hort. Brit.* ed. I. 98, nomen.—Peruv.
weinmanniaefolius, *Engl. in Mart. Fl. Bras.* xii. II. 385.—Bras.

SCHINZIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Alsodeia** ?, Thou. (Violar.).
inconspicua, Dennst. l. c. (Quid?).—Ind. or.

SCHISACHYRIUM, Munro, in Journ. Linn. Soc. vi. (1862) 53, sphalm. = SCHIZACHYRIUM, Nees = **Andropogon**, Linn. (Gramin.).

SCHISANDRA, Michx. Fl. Bor. Am. ii. 218. t. 47 (1803) = **Schizandra**, Michx. (Magnol.).

SCHISANTHES, Haw. Suppl. Pl. Succ. 128 (1819) = **Narcissus**, Tourn. (Amaryll.).
orientalis, Haw. l. c. 129 = N. Tazetta.

SCHISMATOCLAEA, Baker, in Journ. Linn. Soc. xx. (1883) 159. *RUBIACEAE*.
concinna, *Baker, l. c.* xxi. (1885) 407.—Madag.
psychotrioides, *Baker, l. c.* xx. (1883) 160.—Madag.
viburnoides, *Baker, l. c.* xxi. (1885) 407.—Madag.

SCHISMATOGLOTTIS, Zoll. & Mor. Syst. Verz. Zoll. Ind. Archip. 83 (1854). *AROIDEAE*, Benth. & Hook. f. iii. 984.
APOBALLIS, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 318.
COLOBOGYNIUM, Schott, l. c. xv. (1865) 34.
ZANTEDESCHIA, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 9.
acuminatissima, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 281.—Borneo.
asperata, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 297.—Borneo.
barbata, *Engl. l. c.* 298.—Borneo.
Beccariana, *Engl. l. c.* 297.—Borneo.
calyptrata, *Zoll. & Mor. Syst. Verz. Ind. Archip.* 83.—Malaya.
concinna, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 281 = acuminatissima.
conoidea, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 298.—Borneo.

SCHISMATOGLOTTIS:—
crispata, *Hook. f. Bot. Mag.* t. 6576.—Borneo.
decora, *Bull, Cat.* (1884) 16; *et in Gard. Chron.* (1884) I. 711.—Borneo.
elongata, Engl. in Bull. Soc. Tosc. Ortic. iv. (1879) 298 = Rhynchopyle elongata.
Lansbergiana, Linden, Cat. n. 109 (1883) 5; et in Gard. Chron. (1883) I. 510 = Lavalleei.
latifolia, Miq. Fl. Ind. Bat. iii. 214 = rupestris.
Lavalleei, *Linden, in Illustr. Hortic.* xxviii. (1881) t. 418.—Malaya.
longipes, Miq. Fl. Ind. Bat. iii. 214 = calyptrata.
longispatha, *Bull, Cat.* (1881) 20, 6, cum ic.; *et in Illustr. Hortic.* xxix. (1882) 159. t. 466.—Borneo.
marginata, Engl. in Bull. Soc. Tosc. Ortic. iv. (1879) 298 = Rhynchopyle marginata.
modesta, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 125.—Borneo.
neoguineensis, *N. E. Br. in Gard. Chron.* (1885) II. 776.—N. Guin.
ovata, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 125.—Borneo.
picta, *Schott, in Oestr. Bot. Zeitschr.* xiii. (1863) 317.—Java.
picta, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 25 (an praec. ?).—Java.
pulchra, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) t. 520.—Borneo.
pusilla, *Engl. Bot. Jahrb.* i. (1881) 184.—Ins. Philipp.
riparia, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 281 = calyptrata.
rupestris, *Zoll. & Mor. ex Zoll. Syst. Verz. Ind. Archip.* 77.—Java.
siamensis, Bull, Cat. (1885) 15 (Aglaonematis sp.?).—Siam.
sumatrana, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 80.—Sumatra.
variegata, *Hook. ex Engl. in DC. Monog. Phan.* ii. 353.—Borneo.
variegata, Hort. ex N. E. Br. in Gard. Chron. (1885) II. 776 = neoguineense.

SCHISMATOPERA, Klotzsch, in Wiegm. Archiv, vii. (1841) 178 = **Pera**, Mutis (Euphorb.).
distichophylla, Hochst. in Hook. Lond. Journ. Bot. ii. (1843) 44 = P. distichophylla.
laurina, Benth. in Hook. Kew Journ. vi. (1854) 324 = P. distichophylla.
Martiana, Klotzsch, in Wiegm. Arch. vii. (1841) 178 = P. distichophylla.

SCHISMAXON, Steud. in Bot. Zeit. xvi. (1856) 391, nomen = **Xyris**, Gronov.
distichioides, Steud. l. c. = X. subulata.

SCHISMOCERAS, Presl, Rel. Haenk. i. 96. t. 13. f. 2 (1827) = **Dendrobium**, Sw. (Orchid.).
disticha, Presl, l. c. = D. distichum.

SCHISMUS, Beauv. Agrost. 73. t. 15. f. 4 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1195.
ELECTRA, Panz. in Denkschr. Akad. Muench. (1813) 299.
HEMISACRIS, Steud. in Flora, xii. (1829) 490.
arabicus, Nees, Fl. Afr. Austr. 422 = minutus.
brevifolius, Nees, l. c.—Afr. austr.
calycinus, *Coss. in Expl. Sc. Alg.* 138 (an seq. ?).—Marocc.
calycinus, Coss. & Dur. ex Coss. in Bull. Soc. Bot. Fr. iv. (1857) 399 = marginatus.
fasciculatus, Beauv. Agrost. 74 = marginatus, Panicum sanguinale.
Gouani, Trin. Fund. Agrost. 148 = Deschampsia juncea.
hirsutus, Trin. l. c. 148 = minutus.
marginatus, *Beauv. Agrost.* 74. t. 15.—Europ. austr.; As. med.; Afr. austr.
marginatus, Willk. in Flora, xxxv. (1852) 530 = Koeleria cantabrica.
minutus, *Roem. & Schult. Syst.* ii. 584.—Reg. Caucas.
ovalis, *Nees, Fl. Afr. Austr.* 421.—Afr. austr.
patens, J. & C. Presl, Rel. Haenk. i. 371 = Leptochloa patens.
scaberrimus, *Nees, Fl. Afr. Austr.* 423.—Afr. austr.
spectabilis, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 255 = minutus.
tenuis, *Steud. Syn. Pl. Gram.* 295.—Afr. austr.
Villarsii, Trin. Fund. Agrost. 148 = Deschampsia juncea.

SCHISTACHNE, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 252 = **Aristida**, Linn. (Gramin.).
ciliata, Fig. & De Not. l. c. = A. ciliata.

SCHISTANTHE, Kunze, Ind. Sem. Hort. Lips. (1841).—Cf. Linnaea, xvi. (1842) Litt. 109 = **Alonsoa**, Ruiz & Pav. (Scrophul.).
pedunculares, Kunze, ll. cc.—Afr. austr.

SCHISTOCARPHA, Less. in Linnaea, vi. (1831) 409. *COMPOSITAE*, Benth. & Hook. f. ii. 437.
NEILREICHIA, Fenzl, in Denkschr. Akad. Wien. i. (1850) 258. t. 30.
bicolor, *Less. in Linnaea*, vi. (1831) 409.—Mexic.

SCHISTOCARPIA, Pritz. Ic. Ind. i. 999 (1855) = praec.

SCHISTOCODON, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 362 = **Toxocarpus**, Wight & Arn. (Asclep.).
Meyenii, Schau. l. c. 363 = T. Wightianus.

SCHISTOGYNE, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 292. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 758.
Decaisneana, *Karst. Fl. Columb.* ii. 121.—Bras.
sylvestris, Decne. in DC. Prod. viii. 588 = Decaisneana.
sylvestris, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 292.—Uruguay.

SCHISTOPHRAGMA, Benth. ex Endl. Gen. 679 (1839) = **Conobea**, Aubl. (Scrophul.).
pusilla, Benth. in DC. Prod. x. 392 = C. pusilla.

SCHISTOSTEPHIUM, Less. Syn. Comp. 251 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 432.
argyreum, Fenzl, ex Harv. & Sond. Fl. Cap. iii. 168 = flabelliforme.
crataegifolium, *Fenzl, ex Harv. & Sond. l. c.* 169.—Afr. austr.
flabelliforme, *Less. Syn. Comp.* 251.—Afr. austr.
rotundifolium, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 169.—Afr. austr.

SCHIVERECKIA, Andrz. ex DC. Syst. ii. 300 (1821) = **Alyssum**, Tourn. (Crucif.).
contorta, Andrz. ex Ledeb. Fl. Ross. i. 136, in nota.—Kamtschat.
iberidea, Boiss. Diagn. Ser. II. v. 29 (= *Ptilotrichum cappadocicum*).—As. Min.
imeretica, Rupr. Fl. Cauc. 106, 291 = Draba imeretica.
podolica, Andrz. & Bess. ex DC. Syst. ii. 300 = A. podolicum.
spathulata, Andrz. ex DC. Syst. ii. 304 = A. spathulatum.

SCHIWERECKIA, Andrz. ex DC. in Ann. Mus. Par. vii. (1821) 232 = praec.

SCHIWERETZKIA, Rupr. Fl. Cauc. 106 (1869) = praec.

SCHIZACHYRIUM, Nees, Agrost. Bras. 331 (1829) = **Andropogon**, Linn. (Gramin.).
brevifolium, Nees, l. c. 332 = A. brevifolius.
condensatum, Nees, l. c. 333 = A. condensatus.
filiforme, Nees, l. c. 338 = A. tener.
hirtiflorum, Nees, l. c. 334 = A. hirtiflorus.
intermedium, Nees, l. c. = A. consanguineus.
Myosurus, Nees, in Linnaea, xix. (1847) 695 = A. hirtiflorus.
paradoxum, Buese, in Miq. Pl. Jungh. 359 = A. brevifolius.
semiberbe, Nees, Agrost. Bras. 336 = A. semiberbis.
striatum, Nees, l. c. 332 = Ischaemum laxum.
tenerum, Nees, l. c. 336 = A. tener.
zeylanicum, Nees, ex Steud. Nom. ed. II. ii. 530; Wight & Arn. ex Steud. Syn. Pl. Gram. 365 = A. Pseudo-graya.

SCHIZANDRA, Michx. Fl. Bor. Am. ii. 218. t. 47 (1803) (Schisandra); DC. Syst. i. 544 (1818) *MAGNOLIACEAE*, Benth. & Hook. f. i. 19.
COSBAEA, Lem. Illustr. Hortic. ii. (1855) 71.
MAXIMOVITZIA, Benth. & Hook. f. Gen. i. 19 (1862), sphalm.

SCHIZANDRA:—
MAXIMOWICZIA, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 142, 124.
SPHAEROSTEMA, Blume, Bijdr. 22 (1825).
STELLANDRIA, Brickell, in Med. Repos. N. York, vi. (1803) 327.
axillaris, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 45.—Reg. Himal.; Java.
chinensis, *C. Koch, Dendrol.* i. 386; *Baill. Hist. Pl.* i. 148.—China; Japon.
coccinea, *Michx. Fl. Bor. Am.* ii. 219. t. 47; *Sims, Bot. Mag.* t. 1413.—Am. bor.
elongata, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 44.—Reg. Himal.; Java.
grandiflora, *Hook. f. & Thoms. l. c.*—Reg. Himal.
Hanceana, *Baill. Hist. Pl.* i. 150.—China.
japonica, Baill. l. c. = Kadsura japonica.
japonica, Hance, in Journ. Bot. xviii. (1880) 258 = chinensis.
nigra, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 144.—Japon.
propinqua, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 45.—Reg. Himal.

SCHIZANGIUM, Bartl. ex DC. Prod. iv. 571 (1830) = **Mitracarpum**, Zucc. (Rubiac.).
durum, Bartl. ex DC. l. c. 572 = M. Schizangium.

SCHIZANTHERA, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 322 = **Miconia**, Ruiz & Pav. (Melastom.).
bullata, Turcz. l. c. = M. bullata.

SCHIZANTHES, Endl. Gen. 179 (1837) = SCHISANTHES, Haw. = **Narcissus**, Tourn. (Amaryll.).

SCHIZANTHUS, Ruiz & Pav. Prod. 6. t. 1 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 909.
SCIZANTHUS, Pers. Syn. ii. 161 (1807).
alpestris, *Poepp. ex Benth. in DC. Prod.* x. 202.—Chili.
calycosus, *Phil. in Anal. Univ. Chil.* (1873) 529.—Chili.
candidus, *Lindl. Bot. Reg.* (1843) t. 45.—Chili.
duodecemfidus, Stokes, Bot. Comment. i. 117 = pinnatus.
Evansianus, *Paxt. Mag. Bot.* viii. (1841) 171 (sp. dub.).—Hab.?
Gayanus, *Phil. in Linnaea*, xxx. (1859-60) 198.—Chili.
Gilliesii, *Phil. l. c.* xxix. (1857-58) 28.—Chili.
gracilis, *Clos, in C. Gay, Fl. Chil.* v. 153.—Chili.
Grahami, *Gill. in Hook. Bot. Mag.* t. 3044.—Chili.
Hookeri, *Gill. ex R. Grah. in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 176; *et in Bot. Mag.* t. 3070.—Chili.
humilis, Hort. ex Steud. Nom. ed II. ii. 530 = pinnatus.
incanus, Hort. ex C. Morr. in Belg. Hortic. ii. (1852) 42. t. 6. f. 2 = retusus.
integrifolius, *Phil. in Anal. Univ. Chil.* (1873) 530.—Chili.
lacteus, *Phil. Fl. Atac.* 45.—Chili.
laetus, *Phil. l. c.*—Chili.
latifolius, *Phil. in Linnaea*, xxxiii. (1864-65) 214.—Chili.
lilacinus, *Kunze, in Linnaea*, xxiv. (1851) 229.—Chili.
major, Hoffmgg. ex Steud. Nom. ed. II. ii. 530 = pinnatus.
pinnatifidus, Lindl. Bot. Reg. (1843) sub. t. 45 = pinnatus.
pinnatus, *Ruiz & Pav. Fl. Per.* i. 13. t. 17.—Chili.
porrigens, R. Grah. in Edinb. Phil. Journ. xi. (1824) 401; Hook. Exot. Fl. t. 86 = pinnatus.
Priestii, Paxt. Mag. Bot. i. (1835) t. 31 = pinnatus.
retusus, *Hook. Bot. Mag.* t. 3045.—Chili; Peruv.
violaceus, Hort.—Cf. Gard. Chron. (1853) 758 = pinnatus.

SCHIZOBASIS, Baker, in Journ. Bot. xi. (1873) 105. *LILIACEAE*, Benth. & Hook. f. iii. 786.
ADENOTHECA, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 255.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 255.—Afr. trop.
cuscutoides, *Benth. & Hook. f. Gen.* iii. 786.—Afr. austr.

SCHIZOBASIS:—
flagelliformis, *Baker, in Journ. Linn. Soc.* xv. (1876) 261.—Afr. austr.
intricata, *Baker, in Journ. Bot.* xii. (1874) 368.—Afr. austr.
Macowani, *Baker, l. c.* xi. (1873) 105.—Afr. austr.

SCHIZOCALYX, Berg, in Linnaea, xxvii. (1854) 319 = **Calycorectes**, Berg (Myrtac.).
Pohlianus, Berg, l. c. 320.—Bras.

SCHIZOCALYX, Hochst. in Flora, xxvii. (1844) Beil. 1 = **Dobera**, Juss. (Salvador.).
coriaceus, Hochst. l. c. 2 = D. Roxburghii.

SCHIZOCALYX, Scheele, in Flora, xxvi. (1843) 575 = **Origanum**, Tourn. (Labiat.).
smyrnaeus, Scheele, l. c. = O. Onites.
syriacus, Scheele, l. c. = O. Maru?

SCHIZOCALYX, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 73. *RUBIACEAE*, Benth. & Hook. f. ii. 39.
bracteosa, *Wedd. l. c.* 74.—N. Granat.
neo-caledonica, *Brongn. & Gris, l. c.* Sér. V. xiii. (1870-71) 382.—N. Caled.
rubiginosa, *Brongn. & Gris, l. c.* 380.—N. Caled.

SCHIZOCAPSA, Hance, in Journ. Bot. xix. (1881) 292. *TACCACEAE*, Benth. & Hook. f. iii. 741.
plantaginea, *Hance, l. c.*—China.

SCHIZOCARPUM, Schrad. Ind. Sem. Hort. Gotting. (1830).—Cf. Linnaea, vi. (1831) Litt. 73 = *CUCURBITACEAE*, [Benth. & Hook. f. i. 825].
filiforme, *Schrad. ll. cc.*—Mexic.
Liebmannii, *Cogn. in DC. Monog. Phan.* iii. 553.—Mexic.

SCHIZOCARYA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 325 = **Gaura**, Linn. (Onagrar.).
crispa, Spach, l. c. 384 = G. crispa.
Drummondii, Spach, l. c. 382 = G. Drummondii.
Kunthii, Spach, l. c. 383 = G. coccinea.
micrantha, Spach, l. c. 382 = G. parviflora.

SCHIZOCASIA, Schott, in Bonplandia, x. (1862) 148. *AROIDEAE*, Benth. & Hook. f. iii. 978.
acuta, *Engl. Bot. Jahrb.* i. (1881) 186.—N. Guin.
Portei, *Schott, in Bonplandia*, x. (1862) 148.—Ins. Philipp.

SCHIZOCENTRON, Meissn. Gen. Comm. 355 (1843) = **Heeria**, Schlecht. (Melastom.).
elegans, Meissn. l. c. = H. elegans.

SCHIZOCHILUS, Sond. in Linnaea, xix. (1847) 78. *ORCHIDEAE*, Benth. & Hook. f. iii. 632.
Zeyheri, *Sond. l. c.*—Afr. austr.

SCHIZOCHITON, Spreng. Syst. iv. Cur. Post. 251 (1827) = **Chisocheton**, Blume (Meliac.).

SCHIZOCHLAENA, Spreng. Syst. ii. 595 (1825) = **Schizolaena**, Thou. (Chlaenac.).

SCHIZOCODON, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1843) 723. t. 2. *DIAPENSIACEAE*, Benth. & Hook. f. ii. 620.
ilicifolius, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 71.—Japon.
soldanelloides, *Sieb. & Zucc. in Abh. Akad. Muench.* iii. (1843) 725. t. 2. f. 1.—Japon.
uniflorus, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 71.—Japon.

SCHIZODIUM, Lindl. Gen. et Sp. Orch. 358 (1838) = **Disa**, Berg. (Orchid.).
arcuatum, Lindl. Gen. et Sp. Orch. 359 = D. torta.
bifidum, Reichb. f. in Flora, lxvi. (1883) 460 = D. bifida.
clavigerum, Lindl. Gen. et Sp. Orch. 360.—Afr. austr.
flexuosum, Lindl. l. c. 359 = D. flexuosa.
Gueinzii, Reichb. f. in Linnaea, xx. (1847) 694.—Afr. austr.

SCHIZODIUM:—
inflexum, Lindl. Gen. et Sp. Orch. 360 = Disa inflexa.
longipetalum, Lindl. l. c. 359 = D. torta?
maculatum, Lindl. l. c. 360 = D. maculata.
obliquum, Lindl. l. c. 359 = D. bifida?
obtusatum, Lindl. l. c. = D. bifida?
rigidum, Lindl. l. c. 360 = D. bifida.
torta, Steud. Nom. ed. II. ii. 530 = D. torta.

SCHIZOGLOSSUM, E. Mey. Comm. Pl. Afr. Austr. 218 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 753.
ASPIDOGLOSSUM, E. Mey. Comm. Pl. Afr. Austr. 200 (1837).
LAGARINTHUS, E. Mey. l. c. 202 (1837).
MACKENIA, Harv. Gen. S. Afr. Pl. ed. II. 233 (1868).
RHINOLOBIUM, Arn. in Mag. Zool. & Bot. ii. (1838) 420.
atropurpureum, *E. Mey. Comm. Pl. Afr. Austr.* 219.—Afr. austr.
bidens, *E. Mey. l. c.* 220.—Afr. austr.
cordifolium, E. Mey. l. c. 219 = Cynanchum cordifolium.
euphorbioides, *E. Mey. l. c.*—Afr. austr.
Gerrardi, Benth. & Hook. f. Gen. ii. 753 (Gerardi) = Gomphocarpus Gerrardi.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 109.—Afr. trop.
hamatum, *E. Mey. Comm. Pl. Afr. Austr.* 220.—Afr. austr.
hirsutum, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 256.—Afr. austr.
interruptum, *Benth. & Hook. f. Gen.* ii. 753.—Afr. austr.
Petherickianum, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 109.—Afr. trop.
virens, *E. Mey. Comm. Pl. Afr. Austr.* 219.—Afr. austr.

SCHIZOGYNE, Cass. in Dict. Sc. Nat. lvi. 23 (1828) = **Inula**, Linn. (Compos.).
glaberrima, DC. Prod. v. 473.—Ins. Canar.
obtusifolia, Cass. in Dict. Sc. Nat. lvi. 23.—Ins. Canar.
sericea, DC. Prod. v. 473 = *obtusifolia*.

SCHIZOICA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 249 = **Napaea**, Linn. (Malvac.).
laevis, Alef. l. c. = N. dioica.
palmata, Alef. l. c. = N. dioica.

SCHIZOLAENA, Thou. Hist. Vég. Afr. 43. t. 12 (1807). *CHLAENACEAE*, Benth. & Hook. f. i. 195.
SCHIZOCHLAENA, Spreng. Syst. ii. 595 (1825).
cauliflora, *Thou. Hist. Vég. Afr.* 43. t. 12.—Madag.
elongata, *Thou. l. c.* 44.—Madag.
exinvolucrata, *Baker, in Journ. Linn. Soc.* xx. (1883) 97.—Madag.
rosea, *Thou. Hist. Vég. Afr.* 43. t. 12.—Madag.

SCHIZOLEPIS, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 186. t. 26 (1842) = **Scleria**, Berg. (Cyperac.).
abortiva, Nees, l. c. 187 = Scleria abortiva.
arguta, Nees, l. c. 189 = Scleria arguta.
foliosa, Nees, l. c. 188 = Scleria plusiophylla.
latifolia, Nees, l. c. 186 = Scleria arundinacea.
myricocarpa, Nees, l. c. 189 = Scleria myricocarpa.
olyroides, Nees, l. c. 187 = Scleria olyroides.
panicoides, Nees, l. c. 188 = Scleria panicoides.
rufescens, Schrad. ex Nees, l. c. 187 = Scleria olyroides.
Sieberi, Nees, in Mart. Fl. Bras. ii. 1. 188 = Scleria Sieberi.
silvestris, Nees, l. c. 223–224 = Scleria sylvestris.
trigonocarpa, Nees, l. c. 186 = Scleria arundinacea.

SCHIZOLOBIUM, Vog. in Linnaea, xi. (1837) 399. *LEGUMINOSAE*, Benth. & Hook. f. i. 569.
excelsum, *Vog. l. c.*—Bras.
glutinosum, *Tul. in Arch. Mus. Par.* iv. (1846) 157.—Bras.?

SCHIZOMERIA, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 94. *SAXIFRAGEAE*, Benth. & Hook. f. i. 651.
ovata, *D. Don, l. c.* 95.—Austral.

SCHIZONEPHOS, Griff. Notul. iv. 383 (1854) = **Piper**, Linn.
glaucescens, Griff. l. c. = P. Schizonephos.

SCHIZONOTUS, A. Gray, in Proc. Am. Acad. xii. (1876) 66. *ASCLEPIADEAE.*
purpurascens, *A. Gray, l. c.*—Calif.

SCHIZONOTUS, Lindl. ex Wall. Cat. n. 703 (1830) = **Spiraea**, Linn. (Rosac.).
discolor, Rafin. New Fl. Am. iii. 75 = Sp. discolor.
Lindleyanus, Wall. ex Steud. Nom. ed. II. ii. 531 = Sp. sorbifolia.
sorbifolius, Lindl. ex Wall. Cat. n. 703 = Sp. sorbifolia.
tomentosus, Lindl. Bot. Reg. (1840) Misc. 71 = Sp. sorbifolia.

SCHIZOPEPON, Maxim. Prim. Fl. Amur. 110 (1859). *CUCURBITACEAE*, Benth. & Hook. f. i. 839.
bryoniaefolius, *Maxim. l. c.* 111. t. 6.—Manchuria.

SCHIZOPETALON, Sims, Bot. Mag. t. 2379 (1823). *CRUCIFERAE*, Benth. & Hook. f. i. 76.
PERREYMONDIA, Barn. in Ann. Sc. Nat. Sér. III. iii. (1845) 168.
Gayanum, *Barn. in C. Gay, Fl. Chil.* i. 145.—Chili.
maritimum, *Barn. in Ann. Sc. Nat.* Sér. III. iii. (1845) 167.—Chili.
Walcheri, Barn. l. c. 168, sphalm. = seq.
Walkeri, *Sims, Bot. Mag.* t. 2379; *Hook. Exot. Fl.* i. t. 74.—Chili.

SCHIZOPETALUM, DC. Prod. i. 236 (1824) = praec.

SCHIZOPHRAGMA, Sieb. & Zucc. Fl. Jap. 58. t. 26 (1835). *SAXIFRAGEAE*, Benth. & Hook. f. i. 641.
hydrangioides, *Sieb. & Zucc. l. c.* 60.—Japon.

SCHIZOPHYLLUM, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 452 = **Lipochaeta**, DC. (Compos.).
micranthum, Nutt. l. c. 453 = L. micrantha.

SCHIZOPLEURA, Endl. Gen. 1228 (1840) = **Beaufortia**, R. Br. (Myrtac.).
Dampieri, Walp. Rep. ii. 162 = B. Dampieri.
macrostemon, Walp. l. c. = B. macrostemon.
purpurea, Walp. l. c. = B. purpurea.

SCHIZOPOGON, Reichb. Consp. 52 (1828) = **Andropogon**, Linn. (Gramin.).
decumbens, Reichb. ex Steud. Nom. ed. II. ii. 531 = A. brevifolius.

SCHIZOPSERA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 181 = **Schizoptera**, Turcz. (emend. a C. Muell.).

SCHIZOPSIS, Bur. ex Baill. Adansonia, v. (1864–65) 371. t. 10, 11 = **Tynanthus**, Miers (Bignon.).
chimonantha, Bur. ex Baill. l. c. 375.—Bras.
fasciculata, Bur. ex Baill. l. c. 379.—Bras.
Goudotiana, Bur. ex Baill. l. c. 374.—N. Granat.
labiata, Bur. ex Baill. l. c. 373.—Bras.
panurensis, Bur. ex Baill. l. c.—Bras.
polyantha, Bur. ex Baill. l. c. 378.—Peruv.
Regnelliana, Bur. ex Baill. l. c. 379.—Bras.

SCHIZOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 181 (Schizopsera). *COMPOSITAE*, Benth. & Hook. f. ii. 349.
trichotoma, *Turcz. l. c.* 182.—Ecuador.

SCHIZOSPERMUM, Boiv. ex Baill. Adansonia, xii. (1879) 235 = **Cremaspora**, Benth. (Rubiac.).
tubulosum, Boiv. ex Baill. l. c. = C. tubulosa.

SCHIZOSTACHYUM, Nees, Agrost. Bras. 535 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1214.
CHLOOTHAMNUS, Buese, in Miq. Pl. Jungh. 386 (1854).
acutiflorum, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 137.—Ins. Philipp.
Apus, Steud. Syn. Pl. Gram. 332 = Bambusa Apus.
Bitung, Steud. l. c. = Bambusa Bitung.

SCHIZOSTACHYUM :—
Blumei, Munro, in Trans. Linn. Soc. xxvi. (1868) 136 = hindostanicum.
Blumii, *Nees, Agrost. Bras.* 535.—Java.
brachycladum, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 89.—Ins. Molucc.
capitatum, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 136 = Guadua capitata.
chilianthum, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 88 = Melocanna gracilis.
dumetorum, *Munro, in Seem. Bot. Voy. Herald*, 424. —China.
Durie, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 136 = Bambusa Blumeana.
elegantissimum, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 90 (= *Beesha elegantissima*).—Java.
glaucifolium, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 137.—Ins. Fiji.
Hasskarlianum, *Kurz, Bamboo*, 68.—Java.
hindostanicum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 252.—Ind. or.
Iraten, *Steud. Syn. Pl. Gram.* 332.—Java.
Irratun, Kurz, Bamboo, 66, sphalm. = praec.
longispiculatum, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 89.—Java.
loriforme, Munro, in Trans. Linn. Soc. xxvi. (1868) 150 = Dendrocalamus flagellifer.
parviflorum, Munro, l. c. 153 = Dinochloa Tjankorreh.
parvifolium, *Munro, l. c.* 136.—Madag.
serpentinum, *Kurz, Bamboo*, 67.—Java.
Zollingeri, Steud. Syn. Pl. Gram. 332 = Melocanna Zollingeri.

SCHIZOSTEMMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1834) 344 = **Oxypetalum**, R. Br. (Asclep.).
Kingii, Phil. in Anal. Univ. Chil. (1873) 509 = Vincetoxicum boerhaavifolium.
longifolium, Decne. in Ann. Sc. Nat. Sér. II. ix. (1834) 334 = O. solanoides.
microphyllum, Decne. l. c. 345 = O. microphyllum.
parviflorum, Decne. l. c. = O. parviflorum.
solanoides, Decne. l. c. = O. solanoides.

SCHIZOSTEPHANUS, Hochst. ex Benth. & Hook. f. Gen. ii. 762 (1876) = **Vincetoxicum**, Moench (Asclep.).

SCHIZOSTIGMA, Arn. in Ann. Nat. Hist. Ser. I. iii. (1842) *RUBIACEAE*, Benth. & Hook. f. ii. 72.
hirsutum, *Arn. l. c.*—Ind. or.

SCHIZOSTIGMA, Arn. in Hook. Journ. Bot. iii. (1841) 275 = **Cucurbitella**, Walp. (Cucurb.).
asperata, Arn. l. c. = Ç. asperata.

SCHIZOSTYLIS, Backh. & Harv. in Bot. Mag. t. 5422 (1864). *IRIDEAE*, Benth. & Hook. f. iii. 702.
coccinea, *Backh. & Harv. l. c.*—Afr. austr.
ixioides, Harv. ex Baker, in Journ. Linn. Soc. xvi. (1877) 108 = pauciflora.
pauciflora, *Klatt, in Linnaea*, xxxv. (1867–68) 380.—Afr. austr.

SCHIZOTECHIUM, Reichb. Nom. 205 (1841) = **Stellaria**, Linn. (Caryoph.).

SCHIZOTHECA, Ehrenb. in Abh. Akad. Berl. i. (1832) 429 = **Thalassia**, Soland. (Hydroch.).
Hemprichii, Ehrenb. l. c.—Afr. bor.

SCHIZOTHECA, Lindl. Veg. Kingd. 513 (1847) = **Atriplex**, Tourn. (Chenop.).
Halimus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 143 = A. Halimus.
hastata, Fourr. l. c.—A. hastata.
laciniata, Fourr. l. c. = A. laciniata.
littoralis, Fourr. l. c. = A. littoralis.
patula, Fourr. l. c. = A. patula.
rosea, Fourr. l. c. = A. rosea.
tatarica, Čelak. in Arch. Naturw. Landesd. Boehm. ii. (1873) 149 = A. tatarica.

SCHIZOTRICHIA, Benth. in Benth. & Hook. f. Gen. ii. 410 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
eupatorioides, *Benth. l. c.*—Peruv.

SCHKUHRIA, Moench, Meth. 566 (1794) (Sckuhria); Cass. in Dict. Sc. Nat. xlix. 115 (1827) = **Siegesbeckia**, Linn. (Compos.).
dichotoma, Moench, l. c.; Cass. l. c. = Sieg. flosculosa.

SCHKUHRIA, Roth, Catalect. fasc. i. 116 (1797). *COMPOSITAE*, Benth. & Hook. f. ii. 403.
ACHYROPAPPUS, H. B. & K. Nov. Gen. et Sp. iv. 257. t. 390 (1820).
CHAMAESTEPHANUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140.
HOPKIRKIA, DC. Prod. v. 660 (1836).
MIERIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 9 (1825).
TETRACARPUM, Moench, Meth. Suppl. 241 (1802).
abrotanifolia, Steud. Nom. ed. II. ii. 280, in syn., sphalm. = abrotanoides.
abrotanoides, [D.] Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 322 = bonariensis.
abrotanoides, *Roth, Catalect.* fasc. i. 117.—Peruv.
anthemoides, Wedd. Chlor. And. i. 74, in obs. = Bahia anthemoides.
Bigelovii, A. Gray, in Proc. Am. Acad. ix. (1874) 199 = Bahia Bigelovii.
biternata, A. Gray, l. c.; Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot. ii. 212 = Bahia biternata.
bonariensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 321.—Chili.
Hopkirkia, *A. Gray, Pl. Wright.* ii. 94.—Mexic.
integrifolia, A. Gray, in Am. Natural. viii. (1874) 213 = Bahia nudicaulis.
isopappa, *Benth. Pl. Hartw.* 205.—N. Granat.
multiflora, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 322.—Reg. Argent.
neo-mexicana, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 96 = Bahia neo-mexicana.
octoaristata, *DC. Prod.* v. 654.—Peruv.
oolepis, Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 80; et in Linnaea, xxxiv. (1865–66) 529 = pusilla.
pedata, A. Gray, in Proc. Am. Acad. ix. (1874) 199 = Bahia pedata?
pusilla, *Wedd. Chlor. And.* i. 73.—Peruv.; Bolivia.
pusilla, Wedd. ex Benth. & Hook. f. Gen. ii. 403 = Bahia anthemoides?
senecioides, *Nees, Del. Sem. Hort. Bonn.* (1831); *ex DC. Prod.* v. 654.—Mexic.
virgata, *DC. l. c.*—Mexic.
Wislizeni, *A. Gray, in Mem. Am. Acad.* iv. (1849) 96.—Mexic.
Woodhousei, A. Gray, in Proc. Am. Acad. ix. (1874) 199 = Bahia Woodhousei.
Wrightii, *A. Gray, Pl. Wright.* ii. 95.—N. Mexic.

SCHLAGINTWEITIA, Griseb. in Goett. Abh. v. (1855) 156 = **Hieracium**, Tourn. (Compos.).
intybacea, Griseb. l. c. = H. intybaceum.

SCHLECHTENDAHLIA, Benth. & Hook. f. Gen. ii. 408 (1873), sphalm. = **Adenophyllum**, Pers. (Compos.).

SCHLECHTENDALIA, Less. in Linnaea, v. (1830) 242. *COMPOSITAE*, Benth. & Hook. f. ii. 484.
luzulaefolia, *Less. l. c.* 243.—Bras.

SCHLECHTENDALIA, Spreng. Syst. iv. Cur. Post. 296 (1827) = **Mollia**, Mart. (Tiliac.).
speciosa, Spreng. l. c. = M. speciosa.

SCHLECHTENDALIA, Willd. Sp. Pl. iii. 2125 (1803) = **Adenophyllum**, Pers. (Compos.).
capilliacea, [Moç. & Sesse,] ex DC. Prod. v. 638 = A. coccineum.
glandulosa, Willd. Sp. Pl. iii. 2125 = A. coccineum.

SCHLEGELIA, Miq. in Bot. Zeit. ii. (1844) 785. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1052.
axillaris, *Griseb. Fl. Brit. W. Ind.* 445.—Ins. Dominic.
brachyantha, *Griseb. Cat. Pl. Cub.* 191.—Cuba.
elongata, *Miq. in Linnaea*, xxii. (1849) 73.—Guian.
lilacina, *Miq. in Bot. Zeit.* ii. (1844) 786.—Guian.
parasitica, *Griseb. Fl. Brit. W. Ind.* 445.—Ind. occ.
violacea, Griseb. l. c. (Tabebuiae sp.).—Ind. occ.

SCHLEICHERA, Willd. Sp. Pl. iv. 1096 (1805) *SAPINDACEAE*, Benth. & Hook. f. i. 404.
CUSSAMBIUM, [Rumph.] ex Buch.-Ham. in Mem. Wern. Soc. v. ii. (1826) 356.

SCHLEICHERA:—
aculeata, Kostel. Allg. Med.-Pharm. Flora, v. 1829 = trijuga.
amoena, Walp. Rep. v. 366 = Melicocca amoena.
pentapetala, Roxb. Hort. Beng. 29 ; Fl. Ind. ii. 275 = Cupania pentapetala.
ptychocarpa, F. Muell. Fragm. ix. 97 = Dysoxylum ptychocarpum.
pubescens, Roth, Nov. Pl. Sp. 385 = trijuga.
revoluta, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. 575, in obs.—Ins. Philipp.
subundulata, *Turcz. l. c.* 574.—Ins. Philipp.
trijuga, Mor. Syst. Verz. Zoll. 22 = Cupania Lessertiana.
trijuga, *Willd. Sp. Pl.* iv. 1396.—Ind. or. ; Malaya.

SCHLEIDENIA, Endl. Gen. 646 (1839) = **Heliotropium**, Tourn. (Borag.).
bahiensis, Fresen. in Mart. Fl. Bras. viii. 37 = H. polyphyllum.
barbata, Fresen. l. c. 43 = H. barbatum.
Clausseni, Fresen. l. c. 38 = H. Clausseni.
dasycarpa, Fresen. l. c. 37.—Bras.
elliptica, Fresen. l. c. 42.—Bras.
filiformis, Fresen. l. c. 40 = A. filiforme.
fumana, Fresen. l. c.—Bras.
Gardneri, Fresen. l. c. 35.—Bras.
glomerata, Fresen. l. c. 39 = H. polyphyllum.
hispida, Fresen. l. c. 37 = H. fruticosum.
humistrata, Fresen. l. c. 34 = H. humistratum.
incana, Fresen. l. c. 38.—Bras.
inundata, Fresen. l. c. 43 = H. inundatum.
lagoënsis, Warm. in Kjoeb. Vidensk. Meddel. (1867) 15.—Bras.
leptostachya, Fresen. in Mart. Fl. Bras. viii. 43.—Bras.
linifolia, DC. Prod. ix. 558.—Bras.
longepetiolata, Fresen. in Mart. Fl. Bras. viii. 42.—Bras.
macrantha, Fresen. l. c. 40.—Bras.
macrodon, Fresen. l. c. 35.—Bras.
ocellata, Fresen. l. c. 41 = H. ocellatum.
ovalifolia, Fresen. l. c.—Bras.
paradoxa, DC. Prod. ix. 557.—Bras.
parciflora, DC. l. c.—Bras.
polyphylla, Fresen. in Mart. Fl. Bras. viii. 37 = H. polyphyllum.
pubescens, Fresen. l. c. 35.—Bras.
pullulans, Fresen. l. c. 34.—Bras.
salicoides, Fresen. l. c. 38 = H. salicoides.
stenostachya, DC. Prod. ix. 558.—Bras.
strictissima, Fresen. in. Mart. Fl. Bras. viii. 39 = H. strictissimum.
subracemosa, Warm. in Kjoeb. Vidensk. Meddel. (1867) 16.—Bras.

SCHLERANTHUS, Bertol. in Mem. Acad. Sc. Bologn. ii. (1850) 295, sphalm. = **Scleranthus**, Linn. (Illecebr.).

SCHLIMIA, Regel, in Act. Hort. Petrop. iii. ii. (1875) 285 = **Lisianthus**, Aubl. (Gentian.).
Princeps, Regel, l. c. 286 = L. Princeps.

SCHLIMMIA, Planch. & Linden, in Linden, Cat. (1852) ; ex Lindl. & Paxt. Flow. Gard. iii. (1852–53) 115. *ORCHIDEAE*, Benth. & Hook. f. iii. 553.
alpina, *Reichb. & Warsz. in Bonplandia*, ii. (1854) 98.—N. Granat.
jasminodora, *Planch. & Linden, in Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 115.—N. Granat.
trifida, *Reichb. f. in Gard. Chron.* (1876) ii. 708.—N. Granat.

SCHLOSSERIA, Mill. ex Steud. Nom. ed. II. ii. 531 (1841) = **Coccoloba**, Linn. (Polygon.).

SCHLOSSERIA, Vukot. in Oestr. Bot. Wochenbl. vii. (1857) 350 = **Peucedanum**, Tourn. (Umbell.).
heterophilla, Vukot. l. c. = P. carvifolium, *Vill.*

SCHLUMBERGERA, Lem. in Rev. Hortic. Sér. IV. vii. (1858) 253 = **Epiphyllum**, Haw. (Cact.).
epiphylloides, Lem. l. c. = E. Russelianum.

SCHLUMBERGERIA, E. Morr. Belg. Hortic. xxviii. (1878) 311. *BROMELIACEAE*, Benth. & Hook. f. iii. 668.
Lindeni, *E. Morr. Belg. Hortic.* xxxiii. (1883) 121. t. 10–12.—Peruv.

SCHLUMBERGERIA:—
Morreniana, *E. Morr. Belg. Hortic.* xxxiii. (1883) 46, t. 4–6.—Ecuador *vel* Peruv.
Roezli, *E. Morr. l. c.* xxviii. (1878) 311 ; xxix. (1879) 360. t. 19.—Peruv.
virescens, E. Morr. l. c. xxix. (1879) 225, 360 = Roezli.

SCHMALTZIA, ' Desv.' ex Steud. Nom. ed. II. ii. 531 (1841) = seq.

SCHMALZIA, Desv. ex DC. Prod. ii. 72 (1825) = **Rhus**, Tourn. (Anacard.).
aromatica, Desv. ex Steud. Nom. ed. II. ii. 531 = Rhus aromatica.

SCHMARDAEA, Karst. Fl. Columb. i. 187. t. 93. (1861 ?) = **Elutheria**, M. Roem. (Meliac.).
microphylla, Karst. ex C. Muell. in Walp. Ann. vii. 560 = E. microphylla.
nobilis, Karst. Fl. Columb. i. 187. t. 93 = E. microphylla.

SCHMIDELIA, Linn. Mant. i. 10 (1767). *SAPINDACEAE*, Benth. & Hook. f. i. 396.
Aporetica, Forst. Char. Gen. 131. t. 66 (1776).
Azamara, Hochst. ex Reichb. Nom. 145 (1841).
Gemella, Lour. Fl. Cochinch. 648 (1790).
Nassavia, Vell. Fl. Flum. iii. tt. 155, 156 (1827).
Ornitrophe, Juss. Gen. 247 (1789).
Toxicodendron, Gaertn. Fruct. i. 207. t. 44 (1788).
abyssinica, Hochst. in Flora, xxvi. (1843) 80 = Allophylus zeylanicus.
acuminata, Thw. Enum. Pl. Zeyl. 55 = Allophylus zeylanicus.
adenophylla, Wall. Cat. n. 8063 = Allophylus Cobbe.
affinis, *Guill. & Perr. Fl. Seneg. Tent.* 121.—Afr. trop.
africana, DC. Prod. i. 611 = Allophylus zeylanicus.
allophylla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.
Allophyllus, DC. Prod. i. 611 = Allophylus zeylanicus.
alnifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 422.—Afr. trop.
amazonica, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 6.—Bras.
angustata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 371.—N. Granat.
anodonta, F. Muell. Fragm. i. 2 = Ratonia anodonta.
aporetica, Wall. Cat. n. 8058 ; Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 74 = Allophylus Cobbe.
bahiensis, Turcz. in Bull. Sc. Nat. Mosc. xxxi. (1858) i. 398 = Connarus Blanchetii.
bantamensis, Blume, Bijdr. 231 = Allophylus fulvinervis.
bidentata, Wall. Cat. n. 8062 = Allophylus zeylanicus.
Bojeriana, Cambess. in Mém. Mus. Par. xviii. (1829) 38 = Allophylus Cobbe.
caracasana, *Karst. Fl. Columb.* ii. 83. t. 142.—Venezuela.
chartacea, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 183.—Reg. Himal.
Cobbe, DC. Prod. i. 610 = Allophylus Cobbe.
cochinchinensis, DC. l. c. 611 = Allophylus Cobbe.
cochinchinensis, *Spreng. Syst.* ii. 222.—Cochinch.
Cominia, *Sw. Fl. Ind. Occ.* ii. 697.—Ind. occ.
conduplicata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1180, nomen.—Guian.
conferta, Blanco, Fl. Filip. ed. II. 217 = Allophylus zeylanicus.
decipiens, *Arn. in Hook. Journ. Bot.* iii. (1841) 152, 153.—Afr. austr.
dentata, Wall. ex Voigt, Hort. Suburb. Calc. 93 = Allophylus Cobbe.
dioica, Mart. in Flora, xxii. (1839) i. Beibl. 5 = heterophylla.
distachya, DC. Prod. i. 610 = Allophylus Cobbe.
Dregeana, Sond. in Harv. & Sond. Fl. Cap. i. 239 = monophylla.
edulis, *A. St. Hil. Pl. Us. Brés.* t. 67.—Bras.
erosa, *Arn. in Hook. Journ. Bot.* iii. (1841) 152, 153 ; *Presl, Bot. Bemerk.* 41.—Afr. austr.
excelsa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 371.—N. Granat.
floribunda, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Peruv.

SCHMIDELIA:—
foetidissima, Wall. Cat. n. 8064 (an Gynocardiae sp. ?).—Ind. or.
fulvinervis, Blume, Bijdr. 231 = Allophylus fulvinervis.
Gemella, *Cambess. in Mém. Mus. Par.* xviii. (1829) 24.—Cochinch.
glabra, Roxb. ex Wall. Cat. n. 8057 = Allophylus Cobbe.
glabrata, *H. B. & K. Nov. Gen. et Sp.* v. 122.—Am. trop.
Goudotii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 372.—N. Granat.
grandifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 421.—Afr. trop.
grossedentata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 401 = Allophylus Cobbe.
guaranitica, *A. St. Hil. in Bull. Soc. Philom.* (1823) 133.—Bras.
guianensis, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1180, nomen.—Guian.
heterophylla, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 383. t. 82.—Bras.
hirtella, *Hook. f. in Hook. Niger Fl.* 248.—Afr. trop.
hispida, Thw. Enum. Pl. Zeyl. 55 = Allophylus zeylanicus.
inaequilatera, Mart. in Flora, xxii. (1839) i. Beibl. 6. = occidentalis.
integerrima, Wall. Cat. n. 8065 = Evodia Roxburghiana.
integrifolia, DC. Prod. i. 610 = Allophylus Cobbe.
javensis, *Blume, Bijdr.* 232.—Java.
Kobbe, Lam. Illustr. ii. 443. t. 312. f. 2 = Allophylus Cobbe.
laevigata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 399.—Bras.
leptostachya, *Blume, Rumphia*, iii. 141.—Java.
leucocarpa, *Arn. in Hook. Journ. Bot.* iii. (1841) 152, 153.—Afr. austr.
levis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 382.—Bras.
littoralis, *Blume, Bijdr.* 232.—Java.
macrocarpa, A. Rich. Ess. Fl. Cub. 283 = Picrodendron arboreum.
macrophylla, *DC. Prod.* i. 610.—Guian.
macrophylla, Zipp. ex Span. in Linnaea, xv. (1841) 180, nomen = Allophylus Cobbe.
magica, *Baker, in Oliver, Fl. Trop. Afr.* i. 423.—Afr. trop.
melanocarpa, *Arn. in Hook. Journ. Bot.* iii. (1841) 152, 153.—Afr. austr.
Meyeri, Planch. ex Baker, in Oliver, Fl. Trop. Afr. i. 424 = monophylla.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 123.—N. Granat.
monophylla, Hook. f. in Hook. Ic. Pl. t. 775 = hirtella.
monophylla, *Presl, Bot. Bemerk.* 40.—Afr. trop. et austr.
mutabilis, *Blume, Rumphia*, iii. 140.—Java.
natalensis, Sond. in Harv. & Sond. Fl. Cap. i. 239 = erosa.
nervosa, A. Rich. Ess. Fl. Cub. 281 = Thouinia nervosa.
nitidula, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 370.—N. Granat.
oblongifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 424.—Afr. trop.
obovata, A. Gray, in Bot. U. St. Expl. Exped. i. 249 = Allophylus Cobbe.
occidentalis, *Sw. Fl. Ind. Occ.* ii. 665.—Am. trop.
orientalis, Sw. l. c. 666 = Allophylus Cobbe.
ornitrophioides, Wall. Cat. n. 8055 = Allophylus Cobbe.
paniculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Peruv.
panigera, *G. Don, Gen. Syst.* i. 663.—N. Caled.
parviflora, Zipp. ex Span. in Linnaea, xv. (1841) 180, nomen = Allophylus Cobbe.
pinnata, DC. Prod. i. 611 = Pometia pinnata.
puberula, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 382.—Bras.
punctata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38. t. 244.—Peruv.
pyriformis, F. Muell. Fragm. i. 2 = Ratonia pyriformis.
quercifolia, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 7.—Bras.
racemosa, Linn. Mant. i. 67 = Allophylus Cobbe.

SCHMIDELIA:—
reflexa, *Baker, in Oliver, Fl. Trop. Afr.* i. 425.—Afr. trop.
repanda, *Baker, l. c.* 422.—Afr. trop.
Rheedei, Wight, Ic. t. 964; Spicil. Neilgherr. i. 30. t. 33 = Allophylus Cobbe.
rhomboidalis, Nadeaud, Enum. Pl. Tahiti, 70 = Allophylus Cobbe.
rigida, *Sw. Fl. Ind. Occ.* ii. 663.—Ins. S. Doming.
rubrifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 103.—Abyss.
scrobiculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 37. t. 244.—Peruv.
semidentata, Miq. in Linnaea, xxii. (1849) 798 = occidentalis.
senegalensis, Reichb. ex Steud. Nom. ed. II. ii. 531 = africana.
sericea, Cambess. in A. St. Hil. Fl. Bras. Mer. i. 380 = occidentalis.
serrata, DC. Prod. i. 610 = Allophylus Cobbe.
spicata, DC. l. c. 611 = Allophylus Cobbe.
ternata, Cambess. in Mém. Mus. Par. xviii. (1829) 24 = Allophylus Cobbe.
thyrsoides, *Baker, in Oliver, Fl. Trop. Afr.* i. 423.—Afr. trop.
timoriensis, DC. Prod. i. 611 = Allophylus Cobbe.
tomentosa, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 164.—Borneo.
triandra, Blanco, Fl. Filip. ed. II. 218 = Schleichera trijuga.
undulata, Arn. in Hook. Journ. Bot. iii. (1841) 152, 153 = decipiens.
unguicularis, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 399 = occidentalis.
varians, Thw. Enum. Pl. Zeyl. 408 = Allophylus zeylanicus.
velutina, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1838) i. 400.—Guian.
vestita, Wall. Cat. n. 8066 = Allophylus Cobbe.
villosa, Wight, Ic. t. 401 = Allophylus Cobbe.
viticifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 215.—Cuba.

SCHMIDIA, Wight, Ic. t. 1848 (1852) = **Thunbergia**, Linn. (Acanth.).
bicolor, Wight, l. c. = T. Wightiana.

SCHMIDTIA, Moench, Meth. Suppl. 217 (1802) = **Tolpis**, Adans. (Compos.).
abyssinica, Sch. Bip. ex Hochst. in Flora, xxix. (1841) i. Intell. 27 = T. abyssinica.
ambigua, Cass. in Dict. Sc. Nat. xlviii. 433 = T. virgata.
anethifolia, Reichb. in Flora, xiii. (1830) 131 = T. pectinata.
capensis, Reichb. ex Froel. in DC. Prod. vii. 218 = Hieracium capense.
coronopifolia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 404 = T. coronopifolia.
farinulosa, Webb, in Hook. Niger Fl. 143.—Ins. Cap. Virid.
filiformis, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 399 = T. filiformis.
fruticosa, Moench, Meth. Suppl. 218 = T. fruticosa.
laciniata, Sch. Bip. ex Webb & Berth. Phyt. Canar. ii. 406 = T. laciniata.
lagopoda, Sch. Bip. l. c. 408 = T. lagopoda.
nobilis, Sch. Bip. l. c. 399 = T. nobilis.
parviflora, Sch. Bip. l. c. 404 = T. coronopifolia.
quercifolia, Reichb. in Flora, xiii. (1830) 131 = T. umbellata.
Webbii, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 409 = T. Webbii.

SCHMIDTIA, Steud. in J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 144 (1852). *GRAMINEAE*, Benth. & Hook. f. iii. 1175.
pappophoroides, *Steud. l. c.* 145.—Ins. Cap. Virid.
quinqueseta, *Benth. ex Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 31.—Afr. trop.

SCHMIDTIA, Tratt. Fl. Austr. i. 12. t. 451 (1811) = **Coleanthus**, Seid. (Gramin.).
subtilis, Tratt. l. c. = C. subtilis.
utriculata, J. & C. Presl, Fl. Čech. 12 = C. subtilis.
utriculosa, Sternb. in Flora, ii. (1819) 6 = C. subtilis.

SCHMIEDELIA, Murr. Syst. ed. XIII. 310 (1774) = **Schmidelia**, Linn. (Sapind.).

SCHNELLA, Raddi, in Mem. Mod. xviii. Fis. (1820) 411 = **Bauhinia**, Linn. (Legumin.).
brachystachya, Benth. in Hook. Journ. Bot. ii. (1840) 98 = cumanensis.
columbiensis, Benth. Bot. Voy. Sulph. 89 = B. cumanensis.
excisa, Griseb. Fl. Brit. W. Ind. 214 = B. excisa.
emarginata, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1210, nomen.—Guian.
flexuosa, Walp. Rep. v. 572 = B. flexuosa.
longipetala, Benth. in Hook. Journ. Bot. ii. (1840) 98 = B. longipetala.
macrophylla, Griseb. in Goett. Abh. vii. (1857) 210 = B. macrophylla.
macrostachya, Raddi, in Mem. Mod. xviii. Fis. (1820) 411 = B. Raddiana.
microstachya, Raddi, l. c. = B. Langsdorffiana.
rosea, Walp. Rep. v. 572 = B. rosea.
rubiginosa, Benth. in Hook. Journ. Bot. ii. (1840) 97 = B. rubiginosa.
smilacina, G. Don, Gen. Syst. ii. 459 = B. smilacina.
splendens, Benth. in Hook. Journ. Bot. ii. (1840) 97 = B. splendens.
vestita, Benth. Pl. Hartw. 171.—N. Granat.

SCHNITTSPAHNIA, Sch. Bip. in Flora, xxv. (1842) 436 = **Landtia**, Less. (Compos.).
Rueppellii, Sch. Bip. l. c. 439 = L. Rueppellii.
Schimperi, Sch. Bip. l. c. = L. Schimperi.

SCHNITZLEINIA, 'Steud.' ex Walp. Ann. iii. 625 (1852-53) = SCHNIZLEINIA, Steud. = **Vellozia**, Vand. (Amaryll.).

SCHNIZLEINIA, Mart. ex Engl. in Mart. Fl. Bras. xii. ii. 46 (1872) = **Emmotum**, Desv. (Olac.).
nitida, Mart. ex Engl. l. c. = E. nitens.

SCHNIZLEINIA, Steud. ex Hochst. in Flora, xxvii. (1844) 31 = **Vellozia**, Vand. (Amaryll.).
amica, Steud. l. c. (= *Hypoxis Schnizleinia*).—Abyss.

SCHNIZLEINIA, Steud. Nom. ed. II. i. 213, in syn. (1840) = **Boissiera**, Hochst. (Gramin.).

SCHNIZLEINIA, Steud. Nom. ed. II. ii. 532, 720 (1841) = **Trochiscanthes**, Koch (Umbell.).
nodiflora, Steud. l. c. = T. nodiflora.

SCHOBERA, Scop. Introd. 158 (1777) = **Heliotropium**, Tourn. (Borag.).
hirsuta, Rafin. Sylva Tellur. 158 = H. parviflorum?

SCHOBERIA, C. A. Mey. in Ledeb. Fl. Alt. i. 395 (1829) = **Suaeda**, Forsk. (Chenop.).
acuminata, C. A. Mey. l. c. 398 = Suaeda acuminata.
altissima, C. A. Mey. l. c. 402, in obs. = Suaeda altissima.
americana, C. A. Mey. l. c. = Suaeda plattensis.
australis, C. A. Mey. l. c.—Austral.
baccifera, Hohen. ex Moq. in DC. Prod. xiii. ii. 168 = Suaeda setigera.
baccifera, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Suaeda baccifera.
Biebersteiniana, C. A. Mey. l. c. = Suaeda microphylla.
calceoliformis, Moq. in DC. Prod. xiii. ii. 166 = Suaeda calceoliformis.
Clemente, Steud. Nom. ed. II. ii. 532 = Suaeda setigera.
corniculata, C. A. Mey. in Ledeb. Fl. Alt. i. 399 = Suaeda corniculata.
crassifolia, Steud. Nom. ed. II. ii. 532.—Hispan.
cycloptera, Bunge, ex Trautv. in Bull. Soc. Nat. Mosc. xl. (1867) ii. 64.—Soongar.
dendroides, C. A. Mey. Verz. Pfl. Cauc. 159 = Suaeda dendroides.
Dumortieri, Steud. Nom. ed. II. ii. 532 = Suaeda maritima.
fruticosa, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Suaeda fruticosa.
fruticulosa, Less. in Linnaea, ix. (1834) 200.—Sibir.
glauca, Bunge, Enum. Pl. Chin. Bor. 56 = Suaeda glauca.
heterophylla, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 734 = Suaeda heterophylla.

SCHOBERIA:—
hortensis, Steud. Nom. ed. II. ii. 532 = Suaeda hortensis.
indica, Kostel. Allg. Med.-Pharm. Fl. iv. 1438 = Suaeda maritima.
leiosperma, C. A. Mey. in Ledeb. Fl. Alt. i. 397 = Suaeda altissima.
linearis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 465 = Suaeda linearis.
linifolia, Nutt. ex Moq. in DC. Prod. xiii. ii. 161 = Suaeda maritima.
macrocarpa, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Suaeda maritima.
maritima, C. A. Mey. l. c. 400 = Suaeda maritima.
microphylla, Eichw. Pl. Casp.-Cauc. 3, 6 = Suaeda dendroides.
microphylla, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Suaeda microphylla.
microsperma, C. A. Mey. in Eichw. Pl. Casp.-Cauc. 14. t. 13 = Suaeda microsperma.
monoica, Steud. Nom. ed. II. ii. 532 = Suaeda monoica.
montana, Nutt. ex Moq. in DC. Prod. xiii. ii. 161 = Suaeda maritima.
nudiflora, Kostel. Allg. Med.-Pharm. Fl. iv. 1438 = Suaeda nudiflora.
obtusifolia, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 466 = Suaeda obtusifolia.
occidentalis, S. Wats. in Bot. King's Exped. 295.—Am. bor. occ.
Pallasiana, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Kochia hirsuta.
parviflora, Less. in Linnaea, ix. (1834) 201 = Suaeda microsperma.
physophora, C. A. Mey. in Ledeb. Fl. Alt. i. 396 = Suaeda physophora.
prostrata, C. A. Mey. l. c. 402, in obs. = Suaeda microsperma.
pterantha, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 734 = Suaeda pterantha.
pygmaea, Kar. & Kir. l. c. (1842) 436 = Suaeda minima.
salinaria, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 62.—Europ.
salsa, C. A. Mey. in Ledeb. Fl. Alt. i. 401 = Suaeda salsa.
setigera, C. A. Mey. l. c. 402, in obs. = Suaeda setigera.
spicata, C. A. Mey. l. c. = Suaeda spicata.
suffruticulosa, Less. ex Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 465 = Suaeda physophora.
tortuosa, Moq. ex Bunge, l. c. 464.—Hab.?
transoxana, Bunge, l. c. 467 = Suaeda pterantha.
trigyna, C. A. Mey. in Ledeb. Fl. Alt. i. 402, in obs. = Suaeda altissima.
vera, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 465 = Suaeda vera.

SCHOEBERA, Neck. Elem. i. 278 (1790), sphalm. = SCHOBERA, Scop. = **Heliotropium**, Tourn. (Borag.).

SCHOENANTHUM, Mich. ex Linn. Syst. ed. I (1735) = **Andropogon**, Linn. (Gramin.).

SCHOENANTHUS, Adans. Fam. ii. 38 (1763) = praec.

SCHOENEFELDIA, Kunth, Rév. Gram. 283. t. 53 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1164.
gracilis, *Kunth, l. c.*—Afr. trop.
nutans, *Steud. Syn. Pl. Gram.* 202.—Afr. trop.
pallida, *Edgew. in Journ. As. Soc. Beng.* xxi. 1852 (1853), 183.—Ind. or.
ramosa, *Trin. Sp. Gram. Ic.* iii. t. 359.—Ins. Cap. Virid.
stricta, Steud. Syn. Pl. Gram. 202 = gracilis.

SCHOENFELDIA, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 183 (Schaenfeldia), sphalm. = praec.

SCHOENIA, Steetz, in Lehm. Pl. Preiss. i. 480 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 314.
Cassiniana, *Steetz, in Lehm. Pl. Preiss.* i. 481.—Austral.
chlorocephala, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 193 = Helipterum chlorocephalum.

SCHOENIA:—
Humboldtiana, Steetz, ex Walp. Rep. vi. 244 = Helipterum Humboldtianum.
oppositifolia, Steetz, in Lehm. Pl. Preiss. i. 480 = Cassiniana.

SCHOENIDIUM, Nees, in Linnaea, ix. (1834) 291 = **Ficinia**, Schrad. (Cyperac.).
laterale, Nees, l. c. = F. lateralis.

SCHOENISSA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
caucasea, Salisb. l. c. = A. globosum.
rosea, Salisb. l. c. = A. Schoenoprasum.
schoenoprasa, Salisb. l. c. = A. Schoenoprasum.

SCHOENLEINIA, Klotzsch, in Hayne, Arzn. Gew. xiv. sub t. 15 (1843) = **Bathysa**, Presl (Rubiac.).
barbinervis, Klotzsch, l. c. = B. stipulata.
cuspidata, Klotzsch, l. c.—Bras.
thyrsoidea, Miers, in Proc. Roy. Hort. Soc. iv. (1864) 186.—N. Granat.

SCHOENLEINIA, Klotzsch, ex Lindl. Veg. Kingd. 182 (1847) = **Ponthieva**, R. Br. (Orchid.).

SCHOENOBIBLOS, Endl. Gen. 330 (1838) = seq.

SCHOENOBIBLUS, Mart. Nov. Gen. et Sp. i. 65 (1824). *THYMELAEACEAE*, Benth. & Hook. f. iii. 189.
daphnoides, *Mart. l. c.*—Bras.

SCHOENOCAULON, A. Gray, in Ann. Lyc. N. York, iv. (1837) 127. *LILIACEAE*, Benth. & Hook. f. iii. 836.
ASAGRAEA, Lindl. Bot. Reg. t. 33 (1839).
SABADILLA, Brandt & Ratzeb. in Heyne, Arzn. Gew. xiii. t. 27 (1837).
SKOINOLON, Rafin. Fl. Tellur. iv. 27 (1836), nomen prius.
Coulteri, *Baker, in Journ. Linn. Soc.* xvii. (1879) 477.—Mexic.
Drummondii, *A. Gray, in Hook. & Arn. Bot. Beech. Voy.* 388, in nota.—Mexic.
gracile, *A. Gray, in Ann. Lyc. N. York*, iv. (1837) 127.—Am. bor.
intermedium, *Baker, in Journ. Linn. Soc.* xvii. (1879) 477.—Mexic.
officinale, *A. Gray, in Benth. Pl. Hartw.* 29.—Mexic.
texanum, Scheele, in Linnaea, xxv. (1852) 262 = Drummondii.

SCHOENOCEPHALIUM, Seub. in Mart. Fl. Bras. iii. 1. 130 (1847). *RAPATEACEAE*, Benth. & Hook. f. iii. 859.
arthrophyllum, *Seub. l. c.* 131.—Bras.
Martianum, *Seub. l. c.* 130.—Bras.

SCHOENODORUS, Roem. & Schult. Syst. ii. 42 (1817) = **Schedonorus**, Beauv. (Gramin.).

SCHOENODUM, Labill. Nov. Holl. Pl. ii. 79. t. 29 (1806) = **Lyginia**, R. Br. et **Leptocarpus**, R. Br. (Restiac.).
chilense, C. Gay, Fl. Chil. vi. 152 = Leptocarpus chilensis.
gracile, Poir. ex Steud. Nom. ed. II. ii. 532 = Anarthria gracilis.
laevis, Poir. ex Steud. l. c. = Anarthria laevis.
pauciflorum, Poir. ex Steud. l. c. = Anarthria prolifera.
proliferum, Poir. ex Steud. l. c. = Anarthria prolifera.
scabra, Poir. ex Steud. l. c. = Anarthria scabra.
scariosum, Kunth, Enum. Pl. iii. 447 = Leptocarpus scariosus.
simplex, Kunth, l. c. 446 = Leptocarpus Brownii.
tenax, Labill. Nov. Holl. Pl. ii. 80. t. 229 = Leptocarpus tenax, Lyginia barbata.

SCHOENOLAENA, Bunge, in Lehm. Pl. Preiss. i. 289 (1845) = **Xanthosia**, Rudge (Umbell.).
juncea, Bunge, l. c. = X. juncea.
tenuior, Bunge, l. c. 290 = X. tenuior.

SCHOENOLIRION, Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 103. *LILIACEAE*, Benth. & Hook. f. iii. 785.
AMBLOSTIMA, Rafin. Fl. Tellur. ii. 26 (1836).
OXYTRIA, Rafin. l. c. (1836).
album, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 103.—Calif.
croceum, *A. Gray, in Am. Natural.* x. (1876) 427.—Am. bor.
Elliottii, *Feay, ex A. Gray, l. c.*—Florida.
Michauxii, *Torr. Bot. Mex. Bound.* 220.—Am. bor.
Michauxii, Torr. l. c., partim = texanum.
texanum, *A. Gray, in Am. Natural.* x. (1876) 427.—Am. bor.

SCHOENOPRASUM, H. B. & K. Nov. Gen. et Sp. i. 277 (1815) = **Allium**, Tourn. (Liliac.).
lineare, H. B. & K. l. c. = A. Kunthii.
longifolium, H. B. & K. l. c. = A. longifolium.
vulgare, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. Schoenoprasum.

SCHOENOPSIS, Beauv. in Lestib. Ess. Fam. Cyp. 341 (1819) = **Tetraria**, Beauv. (Cyperac.).
Burmanni, Nees, in Linnaea, vii. (1832) 528.—Afr. austr.
flexuosa, Nees, l. c.—Afr. austr.

SCHOENORCHIS, Reinw. Cat. Gew. Buitenz. 100 (1823), nomen; Blume, Bijdr. 361 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 581.
juncifolia, *Reinw. l. c.*; *Blume, l. c.*—Java.
micrantha, Reinw. l. c.; Blume, l. c. 362 = Saccolabium chionanthum.
paniculata, Blume, l. c. = Saccolabium ramulosum.

SCHOENOXIPHIUM, Nees, in Linnaea, vii. (1832) 581. *CYPERACEAE*, Benth. & Hook. f. iii. 1072.
HEMICAREX, Benth. in Benth. & Hook. f. Gen. iii. 1072 (species africanae).
Burkei, C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 386 = Dregeanum.
capense, *Nees, in Linnaea*, vii. (1832) 533.—Afr. austr.
Dregeanum, *Kunth, Enum. Pl.* ii. 529.—Afr. austr.
Ecklonii, *Nees, in Linnaea*, x. (1836) 200.—Afr. austr.
Lehmanni, Kunth, ex Steud. Syn. Pl. Cyp. 245 = Carex Esenbeckiana.
Ludwigii, Hochst. in Flora, xxviii. (1845) 764 = rufum.
Meyerianum, Kunth, Enum. Pl. ii. 530 = capense.
rufum, *Nees, in Linnaea*, x. (1836) 201.—Afr. austr.
Sickmannianum, *Kunth, Enum. Pl.* ii. 530.—Afr. austr.
Thunbergii, Nees, in Linnaea, ix. (1834) 305 = rufum.
Thunbergii, *Nees, l. c.* x. (1836) 201.—Afr. austr.

SCHOENUS, Linn. Coroll. Gen. 2 (1737), pro parva parte; Benth. Fl. Austral. vii. 352 (1878). *CYPERACEAE*, Benth. & Hook. f. iii. 1062.
CHAETOSPORA, R. Br. Prod. 232 (1810).
CYCLOCAMPE, Steud. Syn. Pl. Cyp. 156 (1855).
CYCLOCARPA, Miq. Fl. Ind. Bat. iii. 339 (1855).
GYMNOCHAETA, Steud. Syn. Pl. Cyp. 156 (1855).
HELOTHRIX, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 45.
ISOSCHOENUS, Nees, l. c. 49 (1841).
STREBLIDIA, Link, Hort. Berol. i. 276 (1827).
aculeatus, Linn. Sp. Pl. 42 = Crypsis aculeata.
acuminatus, *R. Br. Prod.* 231.—Austral.
acutus, Labill. Nov. Holl. Pl. i. 18. t. 18 = Cladium schoenoides.
aggregatus, Spreng. ex Kunth, Enum. Pl. ii. 202 = Scirpus flaccifolius.
aggregatus, Steud. ex Kunth, l. c. 202, in obs. = Ficinia tenuifolia.
aggregatus, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 5; Fl. Cap. i. 349 = nigricans.
albus, Linn. Sp. Pl. 44 = Rynchospora alba.
altissimus, Salzm. ex Steud. Syn. Pl. Cyp. 152 = Cladium germanicum.
anceps, Poir. Encyc. Suppl. ii. 252 = Mesomelaena anceps.
angularis, Spreng. ex Kunth, Enum. Pl. ii. 128 = Kyllinga peruviana.

SCHOENUS:—
angustifolius, Vahl, Enum. ii. 225 = Dulichium spathaceum.
aphyllus, *Boeck. in Linnaea*, xxxviii. (1874) 280.—Austral.
aphyllus, Vahl, Enum. ii. 209 = Actinoschoenus Thouarsii.
apogon, *Roem. & Schult. Syst.* ii. 77.—Austral.; N. Zel.
arenarius, Schrad. Anal. Cyp. 27. t. 4. f. 2 = Tetraria compar.
aristatus, Nees, ex Kunth, Enum. Pl. ii. 309 (= *Elynanthus cuspidatus*).—Afr. austr.
aristatus, *Noronha, ex Thou. Prod. Phyt.* 2, *in Mélang.* (1811), nomen.—Madag.
aristatus, Steud. in Flora, xii. (1829) 135 = Hypodiscus duplicatus.
aristatus, Thunb. Prod. Pl. Cap. 16 (= *Elynanthus aristatus*).—Afr. austr.
Armeria, *Boeck. in Linnaea*, xxxviii. (1874) 279.—Austral.
articulatus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 184 = Rynchospora aurea.
arundinaceus, *Soland. ex Forst. f. Prod.* 89, nomen; *Vahl, Enum.* ii. 220.—N. Zel.; Ins. Molucc.; N. Caled.
asper, Schrad. in Schult. Mant. ii. 41 = Fimbristylis aspera.
asperocarpus, *F. Muell. Fragm.* ix. 29.—Austral.
atratus, Schrad. in Goett. Gel. Anz. (1821) III. 2069 = Ficinia setiformis.
axillaris, Lam. Illustr. i. 137 = Rynchospora cephalantha.
axillaris, *Poir. Encyc. Suppl.* ii. 251.—N. Zel.
barbatus, *Boeck. in Linnaea*, xxxviii. (1874) 277.—Austral.
barbatus, Vahl, Eclog. Am. ii. 4 = Rynchospora pterocarpa.
Benthami, F. Muell. Census, 127 = compressus.
bifidus, *Boeck. in Linnaea*, xxxviii. (1874) 282.—Austral.
Bobartiae, *Retz. Obs.* iv. 9.—Zeylan.
Bonanni, Presl, Fl. Sic. p. xliii = nigricans.
breviculmis, *Benth. Fl. Austral.* vii. 364.—Austral.
brevifolius, *R. Br. Prod.* 231.—Austral.
brevisetis, *Poir. Encyc. Suppl.* ii. 251.—Austral.
bromoides, Lam. Illustr. i. 137 (= *Lepidosperma thermale*).—Afr. austr.
Brownii, Hook. f. Handb. N. Zeal. Fl. 298 = apogon.
brunneus, Soland. ex Roem. & Schult. Syst. ii. 65 = Scirpus rufus.
bulbosus, Linn. Mant. ii. 178 = Ficinia bulbosa.
bulbosus, Thunb. Prod. Pl. Cap. 16 = Ficinia secunda.
Burmanni, Vahl, Enum. ii. 227 (= Tetrariae sp.).—Afr. austr.
calostachyus, *Poir. Encyc. Suppl.* ii. 251.—Austral.; Malaya.
capensis, Linn. Cent. Pl. i. 4; Amoen. Acad. iv. 264 = Ficinia ramosissima.
capillaceus, Thunb. Prod. Pl. Cap. 16 (= Elynanthi sp.).—Afr. austr.
capillaris, F. Muell. Census, 128 (*Elynanthus capillaceus*).—Austral.
capillaris, Sw. Prod. Veg. Ind. Occ. 20 = Mariscus capillaris.
capitatus, Crantz, Inst. i. 408 = Kyllinga brevifolia, Mariscus Dregeanus.
capitatus, Dubois, Méth. ed. II. 247 = Scirpus Michelianus.
capitatus, *F. Muell. Fragm.* ix. 58.—Austral.
capitatus, Pers. Syn. i. 60 = Rynchospora glomerata.
capitatus, Poir. Encyc. Suppl. v. 617 = Rynchospora capitata.
capitellatus, Michx. Fl. Bor. Am. i. 36 = Rynchospora glomerata.
Capitellum, Thunb. Prod. Pl. Cap. 16 = Ficinia setiformis.
caricoides, Steud. in Flora, xii. (1829) 137 = Hemichlaena angustifolia.
castaneus, Willd. ex Kunth, Enum. Pl. ii. 303 = Cladium germanicum.
Cephalotes, Rottb. Desc. Nov. Pl. 61; Progr. 14 = Rynchospora Cephalotes.
cerasiformis, Poir. Encyc. Suppl. v. 617 = Rynchospora globosa.
ciliaris, Michx. Fl. Bor. Am. i. 36 = Rynchospora ciliata.

SCHOENUS:—
ciliaris, Muhl. Desc. Gram. 11 = Rynchospora plumosa.
ciliatus, G. F. W. Mey. Prim. Fl. Esseq. 23 = Dichromena ciliata.
circinalis, Schrad. ex Schult. Mant. ii. 43 (Tetrariae sp.).—Afr. austr.
Cladium, Sw. Prod. Veg. Ind. Occ. 19 = Cladium germanicum.
coarctatus, Seenus, Reise Istrien, 74 = Juncus maritimus.
coloratus, Linn. Sp. Pl. 43 = Dichromena leucocephala.
comatus, Link, Enum. Hort. Berol. i. 41 = Rynchospora comata.
compar, Linn. Mant. ii. 177 = Tetraria compar.
compressus, *Benth. Fl. Austral.* vii. 357.—Austral.
compressus, Linn. Sp. Pl. 43 = Scirpus Caricis.
compressus, Presl, Cyp. et Gram. Sic. 12 = nigricans.
concinnus, *Hook. f. Handb. N. Zeal. Fl.* 299.—N. Zel.
confervoides, Willd. ex Kunth, Enum. Pl. ii. 173 = Scirpus confervoides.
congestus, Willd. ex Link, Jahrb. i. III. (1820) 76 = Cladium germanicum.
corniculatus, Lam. Illustr. i. 137 = Rynchospora laxa.
coronatus, Steud. in Flora, xii. (1829) 136 = Ficinia laxa.
corymbosus, Pers. Syn. i. 59 = Rynchospora aurea.
cruentus, *Benth. Fl. Austral.* vii. 357.—Austral.
curvifolius, *Poir. Encyc. Suppl.* ii. 251.—Austral.
cuspidatus, Rottb. Desc. Nov. Pl. 66; Progr. 13 (= *Elynanthus cuspidatus*).—Afr. austr.
cygneus, *Nees, in Lehm. Pl. Preiss.* ii. 81.—Austral.
cymosus, Willd. Sp. Pl. i. 265 = Rynchospora cymosa.
cyperinus, J. F. Gmel. Syst. 1621, sphalm. = seq.
cyperoides, Retz. Obs. iv. 8 = Fimbristylis tristachya.
cyperoides, Sw. Prod. Veg. Ind. Occ. 19 = Rynchospora cyperoides.
dactyloides, Nees, ex Kunth, Enum. Pl. ii. 321 (= *Lepidosperma thermale*).—Afr. austr.
dactyloides, Vahl, Enum. ii. 224 = Asterochaete glomerata.
deformis, *Poir. Encyc. Suppl.* ii. 251.—Austral.
deustus, Berg. Desc. Pl. Cap. 10 = Ficinia scariosa.
deustus, F. Muell. Census, 128 = Mesomelaena deusta.
dispar, Spreng. Neue Entdeck. iii. 8 = Ficinia secunda.
distans, Michx. Fl. Bor. Am. i. 36 = Rynchospora distans.
distans, F. Muell. Census, 127 = calostachyus.
distans, Muhl. Desc. Gram. 11 = Rynchospora Grayii.
distans, Poir. Encyc. Suppl. ii. 249 = Rynchospora longispicata.
Drummondii, *Benth. Fl. Austral.* vii. 359.—Austral.
Drummondii, Steud. Syn. Pl. Cyp. 165 = Gahnia polyphylla.
effusus, Sw. Prod. Veg. Ind. Occ. 19 = Cladium germanicum.
efoliatus, *F. Muell. Fragm.* ix. 32.—Austral.
elatus, Boeck. in Flora, lviii. (1875) 117 = falcatus.
elevatus, Soland. ex Forst. f. Prod. 89 = Cladium germanicum.
elongatus, Willd. ex Kunth, Enum. Pl. ii. 312 = Asterochaete elongata.
ericetorum, *R. Br. Prod.* 231.—Austral.
erinaceus, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 165.—Afr. trop.
erraticus, Hook. f. in Journ. Linn. Soc. vi. (1862) 22 (Fimbristylidis sp.).—Afr. trop.
Fabri, Rottb. Desc. Nov. Pl. 62; Progr. 28 = Panicum lanatum.
falcatus, *R. Br. Prod.* 232.—Austral.
falcatus, Nees, in Linnaea, ix. (1834) 298 = Cladium schoenoides.
fasciatus, Rottb. Desc. Nov. Pl. 67. t. 16. f. 5 = (Tetrariae sp.).—Afr. austr.
fascicularis, Michx. Fl. Bor. Am. i. 37 = Rynchospora fascicularis, *Nutt.*
fascicularis, *Nees, in Ann. & Mag. Nat. Hist.* Ser. I. vi. (1841) 48.—Austral.
ferrugineus, Huds. Fl. Angl. ed. I. 14 = Scirpus rufus.
ferrugineus, Krock. Fl. Siles. i. 57. t. 8 = Juncus bufonius.
ferrugineus, *Linn. Sp. Pl.* 43.—Europ.
festucoides, Willd. ex Kunth, Enum. Pl. ii. 20 (= *Isolepis festucoides*).—Madag.

SCHOENUS:—
filiformis, Lam. Illustr. i. 135 = Ficinia filiformis.
filiformis, Thunb. Prod. Pl. Cap. 16 = Ficinia fastigiata.
filiformis, Willd. ex Kunth, Enum. Pl. ii. 265 = Ficinia bracteata.
Filum, Labill. Nov. Holl. Pl. i. 18. t. 18 = Gahnia trifida.
fimbristyloides, F. Muell. Census, 128 = Cladium undulatum.
flaviculmis, Nees, in Lehm. Pl. Preiss. ii. 81 = brevisetis.
flavus, *Boeck. in Linnaea*, xxxviii. (1874) 278.—Austral.
flavus, Humb. ex Link, Jahrb. i. III. (1820) 75 (= *Monotrema flavum*).—Venezuela.
flexuosus, Steud. ex Kunth, Enum. Pl. ii. 309 (= *Elynanthus cuspidatus*).—Afr. austr.
flexuosus, Thunb. Prod. Pl. Cap. 16 (Tetrariae sp.).—Afr. austr.
floribundus, Willd. ex Presl, in Oken, Isis, xxi. (1828) 270 = Cladium germanicum.
floridus, Rudge, Pl. Gui. 15. t. 18 = Rynchospora aurea.
fluitans, *Hook. f. Fl. Tasm.* ii. 81. t. 141.—Tasman.
fragiferus, Rudge, Pl. Gui. 15. t. 17 = Rynchospora cyperoides.
Fribri, J. F. Gmel. Syst. 136, sphalm. = *Fabri*.
fuscus, Linn. Sp. Pl. ed. II. 1664 = Rynchospora fusca.
fuscus, Muhl. Desc. Gram. 6 = Rynchospora Grayii.
glaucus, Poir. Encyc. Suppl. ii. 249 = Rynchospora rariflora.
globifer, Nees, in Lehm. Pl. Preiss. ii. 81 = brevifolius.
globosus, H. B. & K. Nov. Gen. et Sp. i. 229 = Dichromena ciliata.
globosus, Rudge, Pl. Gui. 14. t. 15 = Rynchospora pterocarpa.
globosus, Willd. ex Kunth, Enum. Pl. ii. 288 = Rynchospora globosa.
globuliferus, Willd. ex Kunth, l. c. 291 = Rynchospora cyperoides.
glomeratus, Linn. Sp. Pl. 44 = Rynchospora glomerata.
glomeratus, Thunb. Prod. Pl. Cap. 17 = Asterochaete glomerata.
gracilis, Sw. Prod. Veg. Ind. Occ. 19 = Rynchospora racemosa.
grammatophyllus, *F. Muell. Fragm.* ix. 31.—Austral.
grandiflorus, *F. Muell. l. c.* 30.—Austral.
Halleri, [Honck.] Verz. Aller Gew. Teutschl. 127 (Quid ?).—Europ.
Hasskarli, Steud. Syn. Pl. Cyp. 166 = Gahnia javanica.
hemisphaericus, Humb. ex Link, Jahrb. i. III. (1820) 75 = Rynchospora pterocarpa.
hirsutus, Vahl, Eclog. Am. i. 6 = Rynchospora hirsuta.
hirtus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 114 = Rynchospora pilosa.
hispidulus, Vahl, Enum. ii. 219 = Rynchospora consanguinea.
holoschaenoides, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Rynchospora cyperoides.
humilis, *Benth. Fl. Austral.* vii. 374.—Austral.
hypomelas, Spreng. Neue Entdeck. iii. 8 = nigricans.
Hystrix, Vahl, Enum. ii. 226 = Hypodiscus aristatus.
imberbis, *R. Br. Prod.* 231.—Austral.
imberbis, Poir. l. c. 251 = apogon.
inanis, Thunb. Prod. Pl. Cap. 16 = Scirpus spathaceus.
incanus, *Forsk. Fl. Aegypt. Arab.* 12.—Aegypt.
indicus, Lam. Encyc. i. 740 = Ficinia setiformis.
indutus, *F. Muell. ex Benth. Fl. Austral.* vii. 372.—Austral.
inermis, Willd. ex Kunth, Enum. Pl. ii. 304 = Scleria triglomerata.
inexpansus, Michx. Fl. Bor. Am. i. 35 = Rynchospora inexpansa.
intermedius ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxv. (1882) 108.—Europ.
involucratus, Rottb. Desc. Nov. Pl. 64. t. 19. f. 1 (Tetrariae ? sp.).—Afr. austr.
junceus, Willd. Phyt. 2 = Scirpus supinus.
juncoides, Vahl, Enum. ii. 211 (= *Isolepis consanguinea*).—Bras.
lacustris, Bernh. Syst. Verz. Erf. 26 = Scirpus lacustris.
laevis, Thunb. Prod. Pl. Cap. 17 (Tetrariae sp.).—Afr. austr.
lanatus, *Labill. Nov. Holl. Pl.* i. 19. t. 20.—Austral.

SCHOENUS:—
lanatus, Willd. ex Kunth, Enum. Pl. ii. 206 (= *Isolepis lanata*).—Am. austr.
lanceus, Thunb. Prod. Pl. Cap. 17 = Schoenoxiphium Sickmannianum.
lateralis, Vahl, Enum. ii. 211 = Ficinia lateralis.
latifolius, Vahl, l. c. 226 = Scleria communis.
laxiflorus, Steud. Syn. Pl. Cyp. 166 = apogon.
laxus, Poir. Encyc. Suppl. ii. 252 = Rynchospora glauca.
lithospermus, Linn. Sp. Pl. ed. II. 65 = Scleria Flagellum-nigrorum.
longifolius, Rudge, Pl. Gui. 14. t. 16 = Hoppia microcephala.
longirostris, Michx. Fl. Bor. Am. i. 35 = Rynchospora laxa.
longirostris, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 269 = Rynchospora longiflora.
longisetis, Poir. Encyc. Suppl. ii. 252 = Rynchospora longisetis.
macrocephalus, Boeck. in Flora, lxi. (1878) 144 (= Tetrariae sp.).—Afr. austr.
macrostachyus, Noé, ex Reichb. Ic. Fl. Germ. viii. 34 = Scirpus maritimus.
marginatus, Kunze, ex Kunth, Enum. Pl. ii. 421 = Carex aphylla.
mariscoides, Muhl. Desc. Gram. 4 = Scleria triglomerata.
Marisculus, Salzm. ex Steud. Syn. Pl. Cyp. 147 = Rynchospora Marisculus.
Mariscus, Linn. Sp. Pl. 42 = Cladium germanicum.
maritimus, Lam. Fl. Fr. iii. 543 = Cyperus aegyptiacus.
maritimus, Salzm. ex Steud. Syn. Pl. Cyp. 137 = Rynchospora tenuis.
maschalinus, Roem. & Schult. Syst. ii. 77 = axillaris.
melanostachys, *R. Br. Prod.* 231.—Austral.
microstachys, Vahl, Enum. ii. 220 (Tetrariae sp.).—Afr. austr.
miliaceus, Lam. Illustr. i. 137 = Rynchospora sparsa.
minimus, T. F. Forst. in Symons, Syn. Pl. Brit. 197 = Juncus capitatus.
minutulus, *F. Muell. Fragm.* ix. 32.—Austral.
moniliferus, Sparrm. ex Spreng. Syst. iii. 830 = Scleria margaritifera.
monocephalus, Salzm. ex Steud. Syn. Pl. Cyp. 143 = Rynchospora pterocarpa.
monoicus, Sm. Engl. Bot. t. 1410 = Kobresia caricina.
Moorei, *Benth. Fl. Austral.* vii. 367.—Austral.
Moorei, T. Kirk, in Trans. N. Z. Inst. xiii. (1881) 384 = pauciflorus.
mucronatus, Linn. Sp. Pl. 42 = Cyperus aegyptiacus.
multiglumis, *Benth. Fl. Austral.* vii. 368.—Austral.
nanus, *Benth. l. c.* 364.—Austral.
natans, *Benth. l. c.* 375.—Austral.
Neesii, F. Muell. Census, 128 = Tricostularia Neesii.
nemorum, Vahl, Symb. Bot. iii. 8 = Hypolytrum latifolium.
nigricans, Hoppe, Taschenb. (1800) 135; (1802) 55 = ferrugineus.
nigricans, *Linn. Sp. Pl.* 43.—Cosmop.
nitens, *Poir. Encyc. Suppl.* ii. 252.—Austral.; N. Zel.; Chili.
nitens, Willd. ex Kunth, Enum. Pl. ii. 313 = Asterochaete nitens.
niveus, Murr. Syst. ed. XIII. 81 = Kyllinga triceps.
nudus, Steud. Syn. Pl. Cyp. 165 = Cladium Gunnii.
obtusifolius, *Boeck. in Linnaea*, xxxviii. (1874) 281.—Austral.
octandrus, F. Muell. Fragm. ix. 31 (= *Elynanthus octandrus*).—Austral.
odontocarpus, *F. Muell. l. c.* 32.—Austral.
odoratus, Aubl. Pl. Gui. i. 44 = Kyllinga odorata.
oliganthos, Kunth, Enum. Pl. ii. 253, in syn., sphalm. = Ficinia filiformis.
oligostachyus, F. Muell. Fragm. ix. 38 = bifidus.
pallens, Schrad. Anal. Cyp. 25 = Ficinia setiformis.
paludosus, Poir. Encyc. Suppl. ii. 251 = Tricostularia paludosa.
palustris, Bernh. Syst. Verz. Erf. 26 = Eleocharis palustris.
paniculatus, Burm. f. Fl. Ind. 19 = Scleria elata, sumatrensis.
paniculatus, Hassk. Cat. Hort. Bog. Alt. 296 = Gahnia javanica.
paniculatus, *Noronha, ex Thou. Prod. Phyt.* 2; *in Mélang.* (1811), nomen.—Madag.

SCHOENUS:—
paradoxus, Spreng. Syst. i. 190 (= *Isolepis paradoxa*). —Venezuela.
pauciflorus, *Hook. f. Handb. N. Zeal. Fl.* 298.—N. Zel.
pauciflorus, F. Muell. Census, 128 = Tricostularia pauciflora.
pauciflorus, Willd. ex Kunth, Enum. Pl. ii. 298 = Rynchospora glauca.
pedicellatus, *Poir. Encyc. Suppl.* ii. 251.—Austral.
pilosus, Willd. Phyt. 3 (= *Abildgaardia pilosa*).—Afr. bor.
pleiostemoneus, *F. Muell. Fragm.* ix. 52.—Austral.
polycephalus, Pers. Syn. i. 59 = Rynchospora cyperoides.
polymorphus, Rottb. Desc. Nov. Pl. 67; Progr. 13 = Fimbristylis ferruginea.
polyphyllus, Vahl, Eclog. Am. ii. 5 = Rynchospora polyphylla.
pterocarpus, Poir. Encyc. Suppl. v. 617 = Rynchospora pterocarpa.
puberulus, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 196 (Fimbristylidis sp.).—Reg. Himal.
pubescens, H. B. & K. Nov. Gen. et Sp. i. 228 = Dichromena pubera.
pubigerus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 116 = Rynchospora robusta.
punctatus, *R. Br. Prod.* 232.—Austral.
punctorius, Vahl, Enum. ii. 217 (Tetrariae sp.).—Afr. austr.
pungens, Willd. ex Kunth, Enum. Pl. ii. 308 = Tetraria compar.
pusillus, Sw. Prod. Veg. Ind. Occ. 20 (Rynchosporae sp.).—Ind. occ.
quadrangularis, *Boeck. in Linnaea*, xxxviii. (1874) 274.—Afr. austr.
radiatus, Forst. ex Vahl, Enum. ii. 220 = arundinaceus.
radiatus, Linn. f. Suppl. 101 = Ficinia radiata.
rariflorus, Michx. Fl. Bor. Am. i. 35 = Rynchospora rariflora.
recurvatus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 132 = Rynchospora recurvata.
restioides, Sw. Prod. Veg. Ind. Occ. 19 (Cladii sp.). —Ind. occ.
ruber, Lour. Fl. Cochinch. 52 = Rynchospora Wallichiana.
rubiginosus, Soland. ex Forst. f. Prod. 89 = Cladium glomeratum.
rufus, Huds. Fl. Angl. ed. II. 15 = Scirpus rufus.
rugosus, Vahl, Eclog. Am. ii. 5 = Rynchospora glauca.
scabripes, *Benth. Fl. Austral.* vii. 368.—Austral.
scariosus, Thunb. Prod. Pl. Cap. 16 = Ficinia bracteata.
scariosus, Vahl, Enum. ii. 210 = Ficinia scariosa.
Scheuchzeri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 119.—Europ.
sculptus, *Boeck. in Linnaea*, xxxviii. (1874) 286.—Austral.
secans, Linn. Syst. ed. X. 865 = Scleria Flagellum-nigrorum.
secundus, Vahl, Enum. ii. 215 = Ficinia secunda.
setaceus, Berg. in Act. Helv. vii. (1772) 130. t. 9 = Rynchospora tenerrima.
setaceus, Mnhl. Desc. Gram. 6 = Rynchospora capillacea.
setaceus, Thuill. Fl. Par. ed. II. 19 = Rynchospora fusca.
setaceus, Vahl, ex Kunth, Enum. Pl. ii. 297 = Rynchospora distans.
setifolius, Benth. Fl. Austral. vii. 359 = Ficinia tenuifolia.
Sodalium, *Hariot, in Bull. Soc. Bot. Fr.* xxxi. (1884) 154.—Fuegia.
spadiceus, H. B. & K. Nov. Gen. et Sp. i. 227 (= *Isolepis paradoxa*).—Venezuela.
spadiceus, Vahl, Enum. ii. 210 = Scirpus spadiceus.
sparganoides, Lindl. ex Kunth, Enum. Pl. ii. 291 = Rynchospora cyperoides.
sparsus, Michx. Fl. Bor. Am. i. 35 = Rynchospora sparsa.
sparteus, *R. Br. Prod.* 231.—Austral.
spathaceus, Linn. Sp. Pl. ed. II. 63 = Dulichium spathaceum.
Spermodon, Schrad. ex Kunth, Enum. Pl. ii. 280 = Rynchospora tenerrima, *Nees*.
sphaerocephalus, Poir. Encyc. Suppl. ii. 252 = Mesomelaena sphaerocephala.
spicatus, Burm. f. Fl. Cap. Prod. 3 (Tetrariae sp.). —Afr. austr.
spicatus, Thunb. Prod. Pl. Cap. 16 = Ficinia bulbosa.

SCHOENUS:—
spilocarpus, Soland. ex Seem. Fl. Vit. 316 = Gahnia schoenoides.
stellatus, Lam. Encyc. i. 741 = Dichromena leucocephala.
striatus, Thunb. Prod. Pl. Cap. 16 = Ficinia setiformis.
stygius, Poir. Encyc. Suppl. ii. 252 = Mesomelaena stygia.
subbulbosus, Benth. Fl. Austral. vii. 358 = capitatus.
subserratus, Steud. in Flora, xii. (1829) 136 = Ficinia setiformis, brevifolia.
surinamensis, Rottb. Desc. Nov. Pl. 68; Progr. 14 = Rynchospora aurea.
surinamensis, *Soland. ex Forst. f. Prod.* 89 (an praec.?). —Ins. Tahiti.
sylvaticus, Bernh. Syst. Verz. Erf. 27 = Scirpus sylvaticus.
sylvaticus, Salzm. ex Schlecht. in Bot. Zeit. vii. (1849) 167 = Rynchospora comata.
tenax, Hook. f. Handb. N. Zeal. Fl. 298 = breviflorus.
Tendo, *Banks & Soland. ex Hook. f. l. c.*—N. Zel.
tenellus, Benth. Fl. Austral. vii. 375 = fluitans.
tener, *Spreng. Neue Entdeck.* iii. 9.—Afr. austr.
tenuiflorus, *Willd. ex Link, Jahrb.* i. III. (1820) 75.—Am. austr.?
tenuifolius, H. B. & K. Nov. Gen. et Sp. i. 228 = Dichromena tenuifolia.
tenuis, *T. Kirk, in Trans. N. Z. Inst.* i. ed. II. (1871) 94.—N. Zel.
tenuissimus, Benth. Fl. Austral. vii. 365 = Lepidospora tenuissima.
Tepperi, F. Muell. Fragm. xi. 106 = trachycarpus.
tetragonus, Poir. Encyc. Suppl. ii. 252 = Mesomelaena tetragona.
thermalis, Linn. Mant. ii. 172 (= *Lepidosperma thermale*).—Afr. austr.
trachycarpus, *F. Muell. Fragm.* ix. 33.—Austral.
triceps, Vahl, Eclog. Am. ii. 4 = Rynchospora cyperoides.
Tricostularia, F. Muell. Census, 128 = Tricostularia compressa.
tristachyus, Thunb. Prod. Pl. Cap. 16 (Tetrariae sp.). —Afr. austr.
tuberosus, Burm. f. Fl. Ind. 190 = Cyperus rotundus.
turbinatus, *Poir. Encyc. Suppl.* ii. 251.—Austral.
umbellatus, Jacq. Coll. i. 113 = Mariscus ovularis.
umbellatus, Walt. Fl. Carol. 70 = Rynchospora laxa.
unispiculatus, *F. Muell. ex Benth. Fl. Austral.* vii. 365.—Austral.
ustulatus, Linn. Mant. ii. 178 (= *Elynanthus ustulatus*).—Afr. austr.
vacillans, T. Kirk, in Trans. N. Z. Inst. x. (1878) 421 = axillaris.
vaginatus, *F. Muell. ex Benth. Fl. Austral.* vii. 371.—Austral.
velutinus, Schrad. ex Kunth, Enum. Pl. ii. 282 = Rynchospora velutina.
villosus, *R. Br. Prod.* 231.—Austral.
viscosus, Schrad. Anal. Cyp. 26 = Tetraria compar.

SCHOEPFIA, Schreb. Gen. 129 (1789). *OLACINEAE*, Benth. & Hook. f. i. 348.
CODONIUM, Rohr, ex Vahl, in Skrivt. Naturh. Selsk. Kjoeb. ii. (1792) 206.
DIPLOCALYX, A. Rich. Fl. Cub. Fanerog. ii. 81. t. 54 (1853).
HAENKEA, Ruiz & Pav. Fl. Per. iii. 8. t. 231 (1802).
SCHOEPFIOPSIS, Miers, in Journ. Linn. Soc. xvii. (1878) 75.
acuminata. *Wall. Cat.* n. 486; *et ex DC. Prod.* iv. 320. —Reg. Himal.
americana, Willd. Sp. Pl. i. 996 = Schreberi.
angulata, *Planch. ex Hemsl. Diagn. Pl. Nov.* 5.—Mexic.
arborescens, Roem. & Schult. Syst. v. 160 = Schreberi.
brasiliensis, *A. DC. in DC. Prod.* xiv. 622.—Bras.
chinensis, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 308.—China.
chrysophylloides, *Planch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 261, in obs.—Ind. occ.
confertiflora, Planch. ex Miers, in Journ. Linn. Soc. xvii. (1878) 74 = quintuplinervis.
didyma, *Wright, ex Griseb. Cat. Pl. Cub.* 119.—Cuba.
flexuosa, *Roem. & Schult. Syst.* v. 160.—Peruv.
fragrans, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 188.—Reg. Himal.
grandifolia, *Baill. Adansonia*, iii. (1862–63) 117.—Bras.

SCHOEPFIA:—
jasminodora, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 135.—Ins. Luchu; Japon.
Marchii, Griseb. Fl. Brit. W. Ind. 310 = chrysophylloides.
mexicana, *A. DC. in DC. Prod.* xiv. 622.—Mexic.
nigricans, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 249 = brasiliensis.
obliquifolia, *Turcz. l. c.* (sp. dub.).—Bras.
obovata, *Wright, Sauv. Fl. Cub.* 21.—Cuba.
odorata, Wall. Cat. n. 485 = fragrans.
parvifolia, *Planch. ex Hemsl. Diagn. Pl. Nov.* 5 (an praec.?).—Mexic.
quintuplinervis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 248.—Bras.
Schreberi, *J. F. Gmel. Syst.* 376.—Peruv.
vacciniflora, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 382, nomen.—N. Granat.
vacciniiflora, *Planch. ex Hemsl. Diagn. Pl. Nov.* 5 (an praec.?).—Guatemala.

SCHOEPFIOPSIS, Miers, in Journ. Linn. Soc. xvii. (1878) 75 = **Schoepfia**, Schreb. (Olacin.).
acuminata, Miers, l. c. = Schoepfia acuminata.
chinensis, Miers, l. c. = Schoepfia chinensis.
fragrans, Miers, l. c. 76 = Schoepfia fragrans.
jasminodora, Miers, l. c. 77 = Schoepfia jasminodora.

SCHOLERA, Hook. f. Fl. Brit. Ind. iv. 62, in syn. (1883), sphalm. = SCHOLLIA, Jacq. f. = **Hoya**, R. Br. (Asclep.).

SCHOLLERA, Rohr, in Skrivt. Naturh. Selsk. Kjoeb. ii. (1792) 210 = **Microtea**, Sw. (Phytol.).
debilis, Rohr, l. c. = M. debilis.

SCHOLLERA, Roth, Tent. Fl. Germ. i. 165, 170 (1788) = **Oxycoccus**, Tourn. (Vaccin.).
erecta, Steud. Nom. ed. I. 746 = Vaccinium erythrocarpum.
erythrocarpa, Steud. l. c. = Vaccinium erythrocarpum.
europaea, Steud. l. c. = O. palustris.
hispidula, Steud. l. c. = Chiogenes serpyllifolia.
macrocarpa, Steud. l. c. = O. macrocarpus.
Oxycoccos, Roth, Tent. Fl. Germ. i. 170 = O. palustris.
paludosa, Baumg. Enum. Stirp. Transs. i. 331 = O. palustris.
palustris, Steud. Nom. ed. I. 746 = O. palustris.

SCHOLLERA, Schreb. Gen. 785 (1791) = **Heteranthera**, Ruiz & Pav. (Ponted.).
graminea, Rafin. in Am. Monthly Mag. (1818) 175; A. Gray, Man. Bot. N. U. St. ed. II. 485 = H. graminea.
graminifolia, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 438, in obs., nomen = H. graminea.
limosa, Rafin. in Am. Monthly Mag. (1818) 175 = H. limosa.

SCHOLLIA, Jacq. f. Eclog. Pl. i. 5. t. 2 (1811) = **Hoya**, R. Br. (Asclep.).
carnosa, Schrank, ex Steud. Nom. ed. I. 414, 746 = H. carnosa.
chinensis, Jacq. f. Eclog. Pl. i. 7 = H. carnosa.
cochinchinensis, Jacq. f. l. c. = H. cochinchinensis.
crassifolia, Jacq. f. l. c. 5. t. 2 = H. carnosa.
viminalis, Jacq. f. l. c. 8 = Philibertia viminalis.
volubilis, Jacq. ex Steud. Nom. ed. I. 414, 746 = Dregea volubilis.

SCHOLTZIA, Schau. in Linnaea, xvii. (1843) 241. *MYRTACEAE*, Benth. & Hook. f. i. 700.
PIPTANDRA, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 323.
PRITZELIA, Schau. in Flora, xxvi. (1843) 405, 407, nomen.
capitata, *F. Muell. ex Benth. Fl. Austral.* iii. 69.—Austral.
ciliata, *F. Muell. Fragm.* iv. 76.—Austral.
decandra, F. Muell. l. c. 75 = Thryptomene saxicola.
denticulata, F. Muell. l. c. = Thryptomene denticulata.
Drummondii, *Benth. Fl. Austral.* iii. 70.—Austral.
laxiflora, *Benth. l. c.* 69.—Austral.
leptantha, *Benth. l. c.*—Austral.
obovata, *Schau. in Linnaea*, xvii. (1843) 241.—Austral.
oligandra, *F. Muell. ex Benth. Fl. Austral.* iii. 70.—Austral.
parviflora, *F. Muell. Fragm.* iv. 76.—Austral.

SCHOLTZIA:—
spathulata, *Benth. Fl. Austral.* iii. 68.—Austral.
teretifolia, *Benth. l. c.* 70.—Austral.
uberiflora, *F. Muell. Fragm.* iv. 74.—Austral.
umbelliflora, *F. Muell. l. c.* 75.—Austral.

SCHOMBURGHIA, DC. Prod. vii. 293 (1838) = **Geissopappus**, Benth. (Compos.).
caleoides, DC. l. c. 294 = G. caleoides.

SCHOMBURGKIA, Benth. & Hook. f. Gen. ii. 396 (1873) = praec.

SCHOMBURGKIA, Lindl. Sert. Orch. tt. 10, 13 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 534.
Brysiana, Lem. Jard. Fleur. i. (1851) Misc. 34 = tibicinis.
carinata, Griseb. Fl. Brit. W. Ind. 621 = Lyonsii.
crispa, *Lindl. Sert. Orch.* t. 10.—Guian.
Galeottiana, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = tibicinis.
gloriosa, Reichb. f. in Hamb. Gartenz. xvi. (1860) 178 = crispa.
Humboldtii, *Reichb. f. in Bonplandia*, iv. (1856) 328.—Venezuela.
Luddemani, Prill. in Journ. Soc. Imp. Hortic. Par. viii. (1862) 123 = undulata.
Lyonsii, *Lindl. in Gard. Chron.* (1853) 615.—Jamaic.
marginata, Hook. Bot. Mag. t. 3729, partim = crispa.
marginata, *Lindl. Sert. Orch.* t. 13.—Guian.
rosea, *Linden, ex Lindl. Bot. Reg.* (1845) sub t. 53.—N. Granat.
tibicinis, *Batem. Orch. Mex. et Guat.* t. 30; *et ex Lindl. Bot. Reg.* (1841) *Misc.* 57.—Am. centr.
undulata, *Lindl. l. c.* (1844) sub t. 23; *Misc.* 13.—Venezuela.
violacea, Paxt. Mag. Bot. xvi. (1849) 123 = undulata.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 116.—N. Granat.

SCHORIGERAM, Adans. Fam. ii. 355 (1763) = **Tragia**, Plum. (Euphorb.).

SCHORTIA, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 794 (1863) = **Actinolepis**, DC. (Compos.).
californica, Hort. ex Vilm. l. c. = A. coronaria.

SCHOTIA, Jacq. Coll. i. 93 (1786). *LEGUMINOSAE*, Benth. & Hook. f. i. 581.
OMPHALOBIUM, Jacq. ex DC. Prod. ii. 508 (1825); et Eclog. Pl. ii. 6. t. 126 (1844).
THEODORA, Medic. Theod. 16 (1786).
afra, Thunb. Prod. Pl. Cap. 79 = speciosa.
alata, *Thunb. Fl. Cap. ed. Schult.* 389.—Afr. austr.
angustifolia, E. Mey. Comm. Pl. Afr. Austr. 161 = speciosa.
brachypetala, *Sond. in Linnaea*, xxiii. (1850) 39.—Afr. austr.
capitata, *Bolle, in Peters, Reise Mossamb. Bot.* 18.—Afr. trop.
diversifolia, Walp. in Linnaea, xiii. (1839) 541 = latifolia.
humboldtioides, *Oliver, Fl. Trop. Afr.* ii. 310.—Afr. trop.
latifolia, *Jacq. Fragm.* 23. t. 15. f. 4.—Afr. austr.
parvifolia, *Dehnh. Rivist. Napol.* i. III. 179.—Afr. austr.
parvifolia, *Jacq. Fragm.* 85. t. 136. f. 4.—Afr. austr.
simplicifolia, Vahl, ex DC. Prod. ii. 508 = Bandeiraea simplicifolia.
speciosa, Blanco, Fl. Filip. ed. I. 356 = Cynometra ramiflora.
speciosa, *Jacq. Coll.* i. 93; *Ic. Pl. Rar.* i. 8. t. 75.—Afr. austr.
stipulata, Ait. Hort. Kew. ed. II. iii. 33 = speciosa.
tamarindifolia, Afzel. ex Sims, Bot. Mag. t. 1153 = speciosa.

SCHOUSBEA, Rafin. Princ. Somiol. 29 (1814) = SCHOUSBOEA, Willd.

SCHOUSBOEA, Schum. & Thonn. Beskr. Guin. Pl. 449 (1827) = **Alchornea**, Sw. (Euphorb.).
cordifolia, Schum. & Thonn. l. c. = A. cordifolia.

SCHOUSBOEA, Willd. Sp. Pl. ii. 578 (1799) = **Cacoucia**, Aubl. (Combret.).
coccinea, Willd. l. c. = C. coccinea.
commutata, Spreng. Syst. ii. 332 = Laguncularia racemosa.

SCHOUTENIA, Korth. in Nederl. Kruidk. Arch. i. (1848) 312. *TILIACEAE*, Benth. & Hook. f. i. 237.
ACTINOPHORA, Wall. Cat. n. 1163 (1830).
Godefroyana, *Baill. Adansonia*, xi. (1876) 370.—Siam.
ovata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 313.—Java.

SCHOUTENSIA, Endl. Prod. Fl. Norf. 79, in obs. (1833) = **Pittosporum**, Banks.
monophylla, Endl. l. c. = P. tenuifolium.

SCHOUWIA, DC. Syst. ii. 643 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 89.
CYCLOPTERYGIUM, Hochst. in Flora, xxxi. (1848) 175.
SUBULARIA, Forsk. Fl. Aegypt. Arab. 117 (1775).
arabica, *DC. Syst.* ii. 644.—Arab.
arabica, Steud. & Hochst. ex Jaub. & Spach, Illustr. iii. 145 = Schimperi.
brassicaefolia, Jaub. & Spach, l. c. = arabica.
brevisepta, Hochst. in Flora, xxxi. (1848) 175 = Schimperi.
glastifolia, Jaub. & Spach, Illustr. iii. 146 = arabica.
Schimperi, *Jaub. & Spach, l. c.* 145.—Aegypt.; Arab.
thebaica, Webb, in Parl. Giorn. Bot. Ital. ii. (1847) 219 = Schimperi.

SCHOUWIA, Schrad. in Goett. Gel. Anz. i. (1821) 717 = **Goethea**, Nees & Mart. (Malvac.).
semiserrata, Schrad. l. c. = G. sempervirens.

SCHOWIA, Sweet, Hort. Brit. ed. III. 43 (1839), sphalm. = praec.

SCHRADERA, Vahl, Eclog. Am. i. 35. t. 5 (1796). *RUBIACEAE*, Benth. & Hook. f. ii. 66.
FUCHSIA, Sw. Prod. Veg. Ind. Occ. 62 (1788).
URCEOLARIA, Willd. in Cothen. Dispos. Veg. 10 (1790).
brasiliensis, *Mart. ex Schult. f. Syst.* vii. 164.—Bras.
capitata, *Vahl, Eclog. Am.* i. 35.—Ind. occ.
cephalophora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1868) 504.—Cuba.
Cephalotes, *Willd. Sp. Pl.* ii. 238.—Jamaic.
cephalotis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
jamaicensis, Pers. Syn. i. 201 = S. Cephalotes.
ligularis, Rudge, Pl. Gui. i. 29. t. 45 = Cephaëlis ligularis.
longifolia, Spruce, ex Hook. f. in Hook. Ic. Pl. sub t. 1068 = Stachyarrhena longifolia.
polycephala, *DC. Prod.* iv. 444.—Guian.
spicata, Spruce, ex Hook. f. in Hook. Ic. Pl. sub t. 1068 = Stachyarrhena spicata.
stellata, *Benth. Bot. Voy. Sulph.* 106. t. 40.—N. Granat.
umbellata, *Presl, Bot. Bemerk.* 81.—Peruv.

SCHRADERA, Willd. in Goett. Journ. Wiss. i. (1797) 1 = **Croton**, Linn. (Euphorb.).
scandens, Willd. l. c. = C. trilobatus.

SCHRADERIA, Medic. Phil. Bot. ii. 40 (1791) = **Salvia**, Tourn. (Labiat.).
hastata, Moench, Meth. 378 = Salvia canariensis.

SCHRANCKIA, Scop. ex J. F. Gmel. Syst. 515 (1790) = **Goupia**, Aubl. (Celastr.).
quinquefaria, J. F. Gmel. l. c. = G. glabra.

SCHRANKIA, Medic. Pflanzeng. 42 (1792) = **Rapistrum**, Tourn. (Crucif.).
divaricata, Moench, Meth. Suppl. 86 = R. perenne.
rugosa, Medic. Pflanzeng. 44 = R. rugosum.
sulcata, Moench, Meth. Suppl. 87 = R. orientale

SCHRANKIA Willd. Sp. Pl. iv. 1041 (1806). *LEGUMINOSAE*, Benth. & Hook. f. i. 593.
LEPTOGLOTTIS, DC. Mém. Lég. 451 (1825).
aculeata, *Willd. Sp. Pl.* iv 1041.—Mexic.
angustata, *Torr. & Gray, Fl. N. Am.* i. 400.—Am. bor.

SCHRANKIA:—
brachycarpa, Benth. in Hook. Journ. Bot. ii. (1840) 130 = Mimosa invisa.
distachya, [*Moç. & Sesse,*] *ex DC. Prod.* ii. 443.—Mexic.
elata, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 309 = Mimosa Ervendbergii.
glabra, Benth. in Hook. Lond. Journ. Bot. v. (1846) 94 = Mimosa spiciflora.
hamata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1042.—Am. trop.
leptocarpa, *DC. Prod.* ii. 443.—Am. trop.
macrostachya, Benth. in Hook. Journ. Bot. iv. (1842) 414 = Mimosa millefoliata.
mexicana, Rafin. New Fl. Am. i. 43 = aculeata.
nova, Steud. Nom. ed. II. ii. 535 (Quid?).—Mexic.
platycarpa, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 183.—Texas.
rhodostachya, Benth. in Hook. Journ. Bot. iv. (1842) 414 = Mimosa rhodostachya.
Sellowii, Benth. l. c. 415 = Mimosa Selloi.
subinermis, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 350.—Mexic.
uncinata, Pritz. Ic. Ind. ii. 254, sphalm. = Smilax pubera.
uncinata, *Willd. Sp. Pl.* iv. 1043.—Am. bor.

SCHREBERA, Linn. Sp. Pl. ed. II. 1662 (1763) = **Cuscuta**, Tourn. (Convolv.).
schinoides, Linn. l. c. = C. africana.

SCHREBERA, Retz. Obs. vi. 25 (1791) = **Elaeodendron**, Jacq. f. (Tiliac.).
albens, Retz. l. c. = E. glaucum.
Pongati, DC. Prod. vii. 548, sphalm. = Sphenoclea zeylanica.

SCHREBERA, Roxb. Pl. Corom. ii. 1. t. 101 (1798). *OLEACEAE*, Benth. & Hook. f. ii. 675.
NATHUSIA, Hochst. in Flora, xxiv. (1841) I. Intell. 25; II. 671.
alata, Welw. in Trans. Linn. Soc. xxvii. (1869) 41 = Saundersiae.
golungensis, *Welw. l. c.* 40.—Afr. trop.
pubescens, Kurz, in Flora, lv. (1872) 399 = swietenioides.
Saundersiae, *Harv. Thes. Cap.* ii. 40.—Afr. austr. et trop.
swietenioides, *Roxb. Pl. Corom.* ii. 1. t. 101.—Ind. or.; Burma.
trichoclada, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 41.—Afr. trop.

SCHREBERA, Schreb. in Nov. Act. Soc. Sc. Upsal. i. (1773) t. 5. f. 1 (cf. p. 91) = **Hartogia**, Linn. f. (Celastr.).
schinoides, Thunb. Prod. Pl. Cap. 28 = H. capensis.

SCHREIBERIA, Steud. Nom. ed. II. ii. 535 (1841), sphalm. = seq.

SCHREIBERSIA, Pohl, in Flora, viii. (1825) 183 = **Augusta**, Pohl (Rubiac.).

SCHRENCKIA, Benth. & Hook. f. Gen. i. 883 (1867), sphalm. = seq.

SCHRENKIA, Fisch. & Mey. in Schrenk, Enum. Pl. Nov. 63 (1841) = **Hippomarathrum**, Link (Umbell.).
vaginata, Fisch. & Mey. l. c. 64 = H. vaginatum.

SCHRENKIA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 606. *UMBELLIFERAE*.
involucrata, *Regel & Schmalh. l. c.*—Turkestan.
papillaris, *Regel & Schmalh. l. c.* 608.—Turkestan.
pungens, *Regel & Schmalh. l. c.* 607.—Turkestan.

SCHROPHULARIA, Medic. Beobacht. (1783) 132, err. typ. = **Scrophularia**, Linn.

SCHTSCHUROWSKIA, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 40 (1882). *UMBELLIFERAE*.
meifolia, *Regel & Schmalh. l. c.*—Kokan.

SCHUBERTA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = SCHUBERTIA, Auct.

SCHUBERTIA, Blume, Bijdr. 884 (1826) = **Horsfieldia**, Blume (Umbell.).
aculeata, Blume, l. c. 885 = H. aculeata.

SCHUBERTIA, Mart. Nov. Gen. et Sp. i. 55. t. 33 (1824) = **Araujia**, Brot. (Asclep.).
auricoma, Lem. in Fl. des Serres, ii. (1846) t. 169 = *graveolens*.
Grahami, Decne. in DC. Prod. viii. 535 = *graveolens*.
grandiflora, Mart. Nov. Gen. et Sp. i. 57.—Bras.
graveolens, Lindl. Bot. Reg. (1838) Misc. 2.—Bras.
hamata, Fourn. in Mart. Fl. Bras. vi. IV. 297.—Bras.
Kefersteinii, Schlecht. in Bot. Zeit. x. (1852) 553.—N. Granat.
longiflora, Mart. Nov. Gen. et Sp. i. 57.—N. Granat.
multiflora, Mart. l. c. i. 56. t. 33.—Bras.
muricata, Manso, ex Fourn. in Mart. Fl. Bras. vi. IV. 296.—Bras.
tristis, Seem. Bot. Voy. Herald, 160 = Macroscepis tristis.

SCHUBERTIA, Mirb. in Nouv. Bull. Soc. Philom. iii. (1812) 123 = **Taxodium**, Rich. (Conif.).
capensis, Spreng. Syst. iii. 890 = Callitris arborea.
disticha, Mirb. in Mém. Mus. Par. xiii. (1825) 75 = T. distichum.
japonica, Spach, Hist. Vég. Phan. xi. 352 = T. heterophyllum.
nucifera, Denh. ex Endl. Syn. Conif. 70 = T. heterophyllum.
sempervirens, Spach, Hist. Vég. Phan. xi. 353 = Sequoia sempervirens.

SCHUDIA, Molina, ex C. Gay, Fl. Chil. iii. 143 (1847) = **Osmorhiza**, Rafin. (Umbell.).
chilensis, Molina, ex C. Gay, l. c. = O. Berterii.

SCHUEBLERIA, Mart. Nov. Gen. et Sp. ii. 113. tt. 186–188 (1827) = **Curtia**, Cham. & Schlecht. (Gentian.).
coarctata, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 442 = *tenuifolia*.
conferta, Mart. Nov. Gen. et Sp. ii. 115. t. 186. f. 2.—Bras.
diffusa, Mart. l. c. 114 = C. diffusa.
intermedia, Prog. in Mart. Fl. Bras. vi. I. 218.—Bras.
obtusifolia, Benth. in Hook. Kew Journ. vi. (1854) 194.—Bras.
patula, Mart. Nov. Gen. et Sp. ii. 117.—Bras.
pusilla, Griseb. Gen. et Sp. Gent. 163.—Bras.
stricta, Mart. Nov. Gen. et Sp. ii. 116 = C. gentianoides.
tenella, Benth. in Ann. Nat. Hist. Ser. I. ii. (1839) 442 = *tenuifolia*.
tenella, Mart. Nov. Gen. et Sp. ii. 117 = C. tenella.
tenuifolia, G. Don, Gen. Syst. iv. 202.—Am. trop.

SCHUECHIA, Endl. Gen. 1178 (1840) = **Qualea**, Aubl. (Vochys.).
brasiliensis, Endl. ex Walp. Rep. ii. 68 = Q. ecalcarata.

SCHUENKIA, Rafin. Princ. Somiol. 30 (1814) = **Schwenkia**, Linn. (Solanac.).

SCHUERMANNIA, F. Muell. in Linnaea, xxv. (1852) 386 = **Darwinia**, Rudge (Myrtac.).
homoranthoides, F. Muell. l. c. 387 = D. Schuermanni.

SCHUFIA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 177 = **Fuchsia**, Plum. (Onagr.).
arborescens, Spach, l. c. = F. arborescens.

SCHULTESIA, Mart. Nov. Gen. et Sp. ii. 103. tt. 180–182 (1827). *GENTIANEAE*, Benth. & Hook. f. ii. 811.
REICHERTIA, Karst. in Bot. Zeit. vi. (1848) 397.
XESTAEA, Griseb. in Linnaea, xxii. (1849) 35.
angustifolia, *Griseb. in DC. Prod.* ix. 68.—Bras.
aptera, *Cham. in Linnaea*, viii. (1833) 9.—Bras.
Aubletii, G. Don, Gen. Syst. iv. 196 = stenophylla.
australis, *Griseb. Gen. et Sp. Gent.* 127.—Bras.
Benthamiana, *Klotzsch, ex Griseb. in Linnaea*, xxii. (1849) 34.—Guian.
brachyptera, *Cham. in Linnaea*, viii. (1833) 8.—Bras.
crenuliflora, *Mart. Nov. Gen. et Sp.* ii. 104.—Bras.
Doniana, *Prog. in Mart. Fl. Bras.* vi. I. 209.—Bras.
gracilis, *Mart. Nov. Gen. et Sp.* ii. 105. t. 181.—Bras.
heterophylla, *Miq. in Linnaea*, xix. (1847) 136.—Am. trop.

SCHULTESIA:—
lisianthoides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 348.—Am. centr.
neuroptera, Griseb. ex Prog. in Mart. Fl. Bras. vi. I. 208 = heterophylla.
pachyphylla, *Griseb. in DC. Prod.* ix. 67.—Bras.
pallens, Bunbury, in Proc. Linn. Soc. i. (1841) 110 = stenophylla.
Pohliana, *Prog. in Mart. Fl. Bras.* vi. I. 205.—Bras.
Schomburgkiana, *Prog. l. c.* 206.—Guian.
stenophylla, *Mart. Nov. Gen. et Sp.* ii. 106. t. 182.—Am. trop.
subcrenata, *Klotzsch, ex Griseb. in Linnaea*, xxii. (1849) 34.—Guian.

SCHULTESIA, Roth, Enum. Pl. Phaen. Germ. i. I. 690 (1827) = **Wahlenbergia**, Schrad. (Campan.).
hederacea, Roth, l. c. = W. hederacea.

SCHULTESIA, Schrad. in Goett. Gel. Anz. i. (1821) 708 = **Gomphrena**, Linn. (Amarant.).
capitata, Schrad. l. c. = G. rupestris.

SCHULTESIA, Spreng. Pugill. ii. 17 (1815) = **Chloris**, Sw. (Gramin.).
petraea, Spreng. l. c. = C. petraea.

SCHULTZIA, Nees, in Nov. Act. Nat. Cur. xi. (1823) 63 = **Herpetacanthus**, Nees (Acanth.).
brasiliensis, Nees, in DC. Prod. xi. 366, in syn. = H. Schultzia.

SCHULTZIA, Rafin. in Med. Repos. N. York, v. (1808) 350 = **Obolaria**, Linn. (Gentian.).
obolarioides, Rafin. l. c. 356 = O. virginica.

SCHULTZIA, Spreng. Pl. Umb. Prod. 30 (1813). *UMBELLIFERAE*, Benth. & Hook. f. i. 909.
Benthami, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 697.—Ind. or.
compacta, *Ledeb. Fl. Ross.* ii. 258.—Sibir. altaic.
crinita, *Spreng. Pl. Umb. Prod.* 30.—Sibir. altaic.

SCHULTZIA, Wall. Cat. n. 589 (1830) = **Cortia**, DC. (Umbell.).
Lindlei, Wall. Cat. l. c.; et ex DC. Prod. iv. 187 = C. Lindleyi.

SCHUMACHERIA, Spreng. Gen. 232 (1830) = **Wormskioldia**, Thonn. (Turner.).
raphanoides, Spreng. ex D. Dietr. Syn. Pl. ii. 1008 = W. heterophylla.

SCHUMACHERIA, Vahl, in Skrivt. Naturh. Selsk. Kjoeb. vi. (1810) 122. *DILLENIACEAE*, Benth. & Hook. f. i. 13.
PLEURODESMIA, Arn. in Edinb. N. Phil. Journ. xvi. (1834) 316.
alnifolia. *Hook. f. & Thoms. Fl. Ind.* i. 66.—Zeylan.
angustifolia, *Hook. f. & Thoms. l. c.*—Zeylan.
castaneifolia, *Vahl, in Skrift. Naturh. Selsk. Kjoeb.* vi. (1810) 122.—Zeylan.

SCHURMANSIA, Blume, Mus. Bot. Lugd. Bat. i. 177. t. 32 (1850). *VIOLARIEAE*, Benth. & Hook. f. i. 120.
angustifolia, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 157.—Borneo.
elegans, *Blume, Mus. Bot. Lugd. Bat.* i. 178.—Ins. Amboin.

SCHWABEA, Endl. Nov. Stirp. Dec. 81 (1839). *ACANTHACEAE*, Benth. & Hook. f. ii. 1108.
POGONOSPERMUM, Hochst. in Flora, xxvii. (1844) Beil. 5.
ciliaris, *Nees, in DC. Prod.* xi. 384.—Afr. trop.
ciliata, Nees, l. c. 431, sphalm. = praec.
modesta, *Endl. Nov. Stirp. Dec.* 81.—Afr. trop.
spicigera, *Nees, in DC. Prod.* xi. 384.—Afr. trop.

SCHWAEGRICHENIA, Reichb. Consp. 147 (1828) = **Hedwigia**, Sw. (Burser.).

SCHWAEGRICHENIA, Spreng. Pugill. ii. 58 (1815) = **Anigozanthos**, Labill. (Haemod.).
flavida, Spreng. l. c. = A. flavida.
rufa, Spreng. l. c. = A. rufa.

SCHWALBEA, Linn. Gen. ed. I. 188 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 975.
americana, *Linn. Sp. Pl.* ed. I. 606.—Am. bor.

SCHWANNIA, Endl. Gen. 1058 (1840). *MALPIGHIACEAE*, Benth. & Hook. f. i. 262.
FIMBRIARIA, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 63. t. 173 (1833).
anisandra, A. Juss. in Arch. Mus. Par. iii. (1843) 608 = Janusia anisandra.
caudata, A. Juss. l. c. 607 = Janusia caudata.
crassistipula, A. Juss. l. c. 610 = Janusia muricata.
cujabensis, Presl, Pot. Bemerk. 23 = elegans.
elegans, *A. Juss. in Arch. Mus. Par.* iii. (1843) 611.—Bras.
intermedia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 395.—Bras.
janusioides, A. Juss. in Arch. Mus. Par. iii. (1843) 606 = Janusia amazonica.
muricata, A. Juss. l. c. 609 = Janusia muricata.

SCHWARTZIA, Vell. Fl. Flum. 221 (1825); v. t. 84 (1827) = **Norantea**, Aubl. (Ternstr.).
glabra, Vell. ll. cc.—Bras.

SCHWEIGGERA, Mart. Nov. Gen. et Sp. iii. 166. t. 297 (1829) = **Renggeria**, Meissn. (Guttif.).
comans, Mart. l. c. t. 297. f. 2 = R. comata.
comata, Steud. Nom. ed. II. ii. 535, sphalm. = praec.

SCHWEIGGERA, E. Mey. ex Baker, in Journ. Linn. Soc. xvi. (1877) 170, 178 = **Gladiolus**, Tourn. (Irid.).
montana, E. Mey. ex Baker, l. c. 178 = G. arenarius.
nemorosa, E. Mey. ex Baker, l. c. = G. montanus.

SCHWEIGGERIA, Spreng. Neue Entdeck. ii. 167 (1821). *VIOLARIEAE*, Benth. & Hook. f. i. 117.
GLOSSARRHEN, Mart. in DC. Prod. i. 290 (1824).
floribunda, A. St. Hil. Pl. Rem. Brés. 282 = fruticosa.
fruticosa, *Spreng. Neue Entdeck.* ii. 167.—Bras.
mexicana, *Schlecht. in Linnaea*, xii. (1838) 204.—Mexic.
pauciflora, Lindl. Bot. Reg. (1841) t. 40 = fruticosa.

SCHWEINFURTHIA, A. Br. in Sitzb. Ges. Naturf. Fr. xx. (1866) 24. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 933.
papilionacea, Boiss. Fl. Orient. iv. 387 = sphaerocarpa.
pedicellata, *Benth. & Hook. f. Gen.* ii. 934 (lapsu, pedicellaris).—Arab.
pterosperma, *A. Br. in Sitzb. Ges. Naturf. Fr.* xx. (1866) 24.—Aegypt.; Arab.
sphaerocarpa, *A. Br. l. c.*—Afghan.

SCHWEINITZIA, Ell. Sketch, i. 478, in obs. (1817); et ex Nutt. Gen. Am. ii. Add. (1818). *MONOTROPEAE*, Benth. & Hook. f. ii. 606.
MONOTROPSIS, Schwein. ex Ell. Sketch, i. 478 (1817).
caroliniana, G. Don, Gen. Syst. iii. 867 = odorata.
odorata, *Ell. Sketch*, i. 478, 479.—*Cf. DC. Prod.* vii. II. 780.—Am. bor.
Reynoldsiae, *A. Gray, in Proc. Am. Acad.* xx. (1885) 301.—Florida.

SCHWENKFELDA, Schreb. Gen. 123 (1789) = **Sabicea**, Aubl. (Rubiac.).
aspera, Spreng. Neue Entdeck. i. 281 = Coccocypselum canescens.
aspera, Willd. Sp. Pl. i. 982 = Sabicea aspera.
aurea, Spreng. Neue Entdeck. i. 280 = Coccocypselum aureum.
cinerea, Sw. Fl. Ind. Occ. i. 452 = Sabicea cinerea.
diversifolia, Spreng. Syst. i. 765 = Sabicea diversifolia.
eriantha, D. Dietr. Syn. Pl. i. 793 = Sabicea cinerea.
glabra, Spreng. Syst. iv. Cur. Post. 85 = Urophyllum glabrum.
hirsuta, Spreng. l. c. i. 765 = Sabicea aspera.
hirta, Sw. Fl. Ind. Occ. i. 450 = Sabicea aspera.
macrophylla, Spreng. Syst. i. 765 (= *Nonatelia macrophylla*).—Venezuela.
malaccensis, Spreng. l. c. iv. Cur. Post. 85 = Urophyllum villosum.
pumila, D. Dietr. Syn. Pl. i. 794 = Sabicea pumila.
triflora, D. Dietr. l. c. 793 = Sabicea triflora.
umbellata, Ruiz & Pav. Fl. Per. ii. 55 = Sabicea umbellata.
villosa, D. Dietr. Syn. Pl. i. 793 = Urophyllum villosum.
villosa, Willd. ex Spreng. Syst. i. 765 = Sabicea aspera.

SCHWENKFELDIA, Willd. Sp. Pl. i. 982 (1797) = praec.

SCHWENKIA, Linn. Gen. ed. VI. 567 (1764). *SOLANACEAE*, Benth. & Hook. f. ii. 911.
CHAETOCHILUS, Vahl, Enum. i. 101 (1805).
MATHAEA, Vell. Fl. Flum. 22 (1825).
MATTHISSONIA, Raddi, in Mem. Mod. xviii. Fis. (1820) 390.
SCHUENKIA, Rafin. Princ. Somiol. 30 (1814).
adscendens, [Willd.] Link, Jahrb. i. III. (1820) 51 = hirta.
americana, H. B. & K. Nov. Gen. et Sp. ii. 375. t. 180 = hirta.
americana, *Linn. Gen.* ed. VI. 567.—Bras.
angustifolia, *Benth. in DC. Prod.* x. 195.—Bras.
brasiliensis, *Poir. Encyc. Suppl.* v. 88.—Bras.
breviseta, *Casar. Nov. Stirp. Bras. Dec.* 29.—Bras.
browallioides, *H. B. & K. Nov. Gen. et Sp.* ii. 375. t. 181.—Venezuela.
chenopodiacea, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1155, nomen.—Guian.
curviflora, *Benth. in DC. Prod.* x. 196.—Bras.
deflorata, *Vatke, in Ind. Sem. Hort. Berol. App.* (1875) [2].—Bras.
discolor, *Kunze, in Linnaea*, xvi. (1842) 312.—Hab.?
divaricata, *Benth. in DC. Prod.* x. 193.—Bras.
fasciculata, Benth. l. c. 195 = Melananthus dipyrenoides.
glabrata, *H. B. & K. Nov. Gen. et Sp.* ii. 374. t. 178.—N. Granat.
grandiflora, *Benth. in DC. Prod.* x. 193.—Guian.
guianensis, *Benth. l. c.* 195.—Guian.
guineënsis, *Schum. & Thonn. Beskr. Guin. Pl.* 8.—Afr. trop.
Hilariana, *DC. Pl. Rar. Hort. Genèv.* 35. t. 10.—Bras.
hirta, *Klotzsch, in Linnaea*, xiv. (1840) 289.—Bras.
hyssopifolia, *Benth. in DC. Prod.* x. 195.—Bras.
Karstenii, *Vatke, in Ind. Sem. Hort. Berol. App.* (1875) [2].—N. Granat.
longiseta, *Casar. Nov. Stirp. Bras. Dec.* 30.—Bras.
micrantha, *Benth. in DC. Prod.* x. 195.—Bras.
mollissima, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 541 = divaricata.
mollissima, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 47.—Bras.
ovalifolia, *F. A. Schmidt, in Mart. Fl. Bras.* viii. 253.—Bras.
pallens, D. Dietr. Syn. Pl. i. 41 = patens.
patens, *H. B. & K. Nov. Gen. et Sp.* ii. 374. t. 179.—Venezuela.
pubescens, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 48.—Bras.
tenuis, *Griseb. in Goett. Abh.* xix. (1874) 214.—Reg. Argent.
Tweediana, Benth. in DC. Prod. x. 196 = curviflora.
ventricosa, *Vatke, in Ind. Sem. Hort. Berol. App.* (1875) [2].—Bras.
volubilis, *Benth. in DC. Prod.* x. 193.—Bras.

SCHWERINIA, Karst. Ausw. Gew. Venez. 12 (1848) = **Meriania**, Sw. (Melastom.).
barbinervis, Klotzsch, in Karst. l. c. 13 = M. barbinervis.
cordata, Karst. Fl. Columb. i. 35. t. 18. f. b = M. cordata.
superba, Karst. Ausw. Gew. Venez. 13 = M. Karstenii.
Trianae, Karst. Fl. Columb. i. 35. t. 18. f. a = M. speciosa.

SCHWEYCKERTA, C. C. Gmel. Fl. Bad. i. 447 (1805) = **Limnanthemum**, S. T. Gmel. (Gentian.).
nymphoides, C. C. Gmel. l. c. = L. nymphoides.

SCHWEYKERTA, Griseb. in DC. Prod. ix. 138 (1845) = praec.

SCHYCHOWSKIA, Wedd. in DC. Prod. xvi. I. 70 (1869) = seq.

SCHYCHOWSKYA, Endl. in Ann. Wien. Mus. i. (1836) 187. t. 13 = **Fleurya**, Gaudich. (Urtic.).
ruderalis, Endl. l. c. = F. ruderalis.

SCIADIARA, Rafin. Fl. Tellur. iv. 70 (1836) = **Convolvulus**, Tourn.
Cneorum, Rafin. l. c. = C. Cneorum.
ruber, Rafin. l. c. = Calystegia setifera?

SCIADICARPUS, Hassk. in Flora, xxv. (1842) II. Beibl. 20 = **Kibara**, Endl. (Monim.).
Brongniartii, Hassk. l. c. = K. Blumei.

SCIADIOSERIS, Kunze, in Bot. Zeit. ix. (1851) 349 = **Senecio**, Tourn. (Compos.).
vaginata, Kunze, l. c. = Sen. cervariaefolius.

SCIADIPHYLLUM, Hassk. Cat. Hort. Bog. Alt. 165 (1844) = **Sciadophyllum** P. Br. (Araliac.).

SCIADOCALYX, Regel, Gartenfl. ii. (1853) 257. t. 61 = **Isoloma**, Decne. (Gesner.).
digitalaeflora, Linden & Andr. in Illustr. Hortic. xvii. (1870) 95. t. 17.—N. Granat.
Luciani ×, Andr. l. c. xxi. (1874) 151. t. 182.
Warszewiczii, Regel, Gartenfl. ii. (1853) 257, cum ic.—N. Granat.

SCIADODENDRON, Griseb. in Bonplandia, vi. (1858) 7. Genus dubium *ARALIACEARUM*.—Cf. Benth. & Hook. f. i. 935.
excelsum, *Griseb. l. c.*—Panama.

SCIADONARDUS, Steud. in Flora, xxxiii. (1850) 229, nomen. *GRAMINEAE*.
distichophyllus, Steud. l. c. (Quid?).—Am. bor.

SCIADOPANAX, Seem. in Journ. Bot. iii. (1865) 73. t. 27 = **Panax**, Linn. (Araliac.).
Boivini, Seem. l. c. 74 = P. Boivini.

SCIADOPHILA, Phil. in Linnaea, xxviii. (1856) 618 = **Rhamnus**, Linn.
maytenoides, Phil. l. c.—Chili.

SCIADOPHYLLUM, P. Br. Hist. Jamaic. 190. t. 19 (1756) (Sciodaphyllum); Reichb. Consp. 145 (1828). *ARALIACEAE*, Benth. & Hook. f. i. 940.
ACTINOPHYLLUM, Ruiz & Pav. Prod. 51. t. 8 (1794).
abyssinicum, Steud. Nom. ed. II. ii. 537 = Heptapleurum abyssinicum.
acuminatum, *Poir. Encyc.* vi. 746.—Peruv.
angulatum, *Poir. l. c.* 745.—Peruv.
anomalum, *G. Don, Gen. Syst.* iii. 391.—Ins. Trinit.
assamicum, *C. Koch, in Wochenschr.* ii. (1859) 364, 365 (sp. dub.).—Ind. or.
avene, Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 19 = Heptapleurum avene.
Baikiei, Seem. in Journ. Bot. iii. (1865) 267 = Heptapleurum Baikiei.
Barteri, Seem. l. c. = Heptapleurum Barteri.
Belangeri, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 92.—Ins. Martinic.
Brownii, *Spreng. Syst.* i. 953.—Jamaic.
Burchellianum, *Baill. Adansonia*, xii. (1876-79) 147.—Bras.?
capitatum, Griseb. Fl. Brit. W. Ind. 306 = Oreopanax capitatum.
caudatum, *Decne. & Planch. in Rev. Hort.* Sér. IV. iii. (1854) 107, nomen.—Hab.?
confusum, *March. in Mart. Fl. Bras.* xi. I. 244.—Bras.
conicum, *Poir. Encyc.* vi. 746.—Peruv.
decaphyllum, *Seem. in Journ. Bot.* iii. (1865) 266.—Guian.
digitatum, G. Don, ex Steud. Nom. ed. II. ii. 537 = Pachira insignis.
divaricatum, Blume, Bijdr. 876 = Heptapleurum divaricatum.
elatum, Seem. in Journ. Bot. iii. (1865) 267 = Heptapleurum elatum.
ellipticum, Blume, Bijdr. 878 = Heptapleurum venulosum.
farinosum, Blume, l. c. 875 = Heptapleurum tomentosum.
ferrugineum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107.—Peruv.
Gayanum, *Decne. & Planch. l. c.*, nomen.—Hab.?
Goudotii, *Planch. & Linden, ex Seem. Journ. Bot.* iii. (1865) 266.—N. Granat.
heterotrichum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen.—N. Granat.
Humboldtianum, *Decne. & Planch. ex Seem. in Journ. Bot.* iii. (1865) 268.—N. Granat.
humile, Blume, Bijdr. 875 = Heptapleurum parasiticum.

SCIADOPHYLLUM:—
Jacquini, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 190 = Hedera arborea.
japurense, *Mart. & Zucc. ex March. in Mart. Fl. Bras.* xi. I. 244.—Bras.
Karstenianum, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 93.—Venezuela.
longifolium, Blume, Bijdr. 876 = Heptapleurum longifolium.
lucidum, Blume, l. c. 877 = Heptapleurum longifolium.
macrostachyum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 222 = Brassaia macrostachya.
Manni, Seem. in Journ. Bot. iii. (1865) 267 = Heptapleurum Mannii.
Mathewsii, *Seem. l. c.* 268.—Peruv.
micranthum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen; et (1884) 321.—N. Granat.
ochroleucum, *Decne. & Planch. l. c.* 107, nomen.—Hab?
oxyanthum, *Planch. & Linden, ex Seem. in Journ. Bot.* iii. (1865) 266, nomen.—N. Granat.
palmatum, Blume, Bijdr. 875 = Trevesia sundaica.
parasticum, Blume, l. c. 877 = Heptapleurum parasiticum.
pedicillatum, *Poir. Encyc.* vi. 746.—Peruv.
pentandrum, *Poir. l. c.* 747.—Peruv.
Planchonianum, *March. in Bull. Soc. Bot. Belg.* xix. (1880) 92.—Ecuador.
pulchellum, Hort. ex Seem. in Journ. Bot. iii. (1865) 269 = Tupidanthus calyptratus.
pulchrum, *Hort. ex Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 106.—Hab.?
quinduense, *DC. Prod.* iv. 261.—N. Granat.
racemiferum, Miq. ex Seem. in Journ. Bot. iii. (1865) 269, nomen (Quid?).—Hab.?
racemiflorum, *Miq. in Linnaea*, xxv. (1852) 651.—Chili.
rotundifolium, Hort. ex Seem. in Journ. Bot. iii. (1865) 269. (= Heptapleuri sp.).—Hab.?
rubiginosum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen.—Venezuela.
samydifolium, *Wright, ex Griseb. Cat. Pl. Cub.* 117.—Cuba.
scandens, Blume, Bijdr. 878 = Heptapleurum scandens, *Seem.*
sphaerocoma, *Benth. Bot. Voy. Sulph.* 102.—N. Granat.
Sprucei, *Seem. in Journ. Bot.* iii. (1865) 268.—Peruv.
subavene, Blume, Bijdr. 876 = Heptapleurum Coronaesylvae.
tomentosum, Blume, l. c. 877 = Heptapleurum tomentosum.
Trianae, *Planch. & Linden, ex Seem. in Journ Bot.* iii. (1865) 268, nomen.—N. Granat.
verticillatum, Span. in Linnaea, xv. (1841) 208 = Heptapleurum verticillatum.
villosum, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 107, nomen.—N. Granat.

SCIADOPITYS, Sieb. & Zucc. Fl. Jap. ii. I. t. 101, 102 (1841?). *CONIFERAE*, Benth. & Hook. f. iii. 437.
verticillata, *Sieb. et Zucc. l. c.*—Japon.

SCIADOSERIS, C. Muell. in Walp. Ann. v. 143 (1858), err. typ. = SCIADIOSERIS, Kunze = **Senecio**, Tourn. (Compos.).

SCIADOTAENIA, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 51 = seq.

SCIADOTENIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43. *MENISPERMACEAE*, Benth. & Hook. f. i. 39.
SCIADOTAENIA, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 51.
amazonica, *Eichl. in Flora*, xlvii. (1864) 395.—Bras.
cayennensis, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 51.—Guian.
ramiflora, *Eichl. in Flora*, xlvii. (1864) 395.—Peruv.

SCIAPHILA, Blume, Bijdr. 514 (1825). *TRIURIDEAE*, Benth. & Hook. f. iii. 1002.
APHYLLEIA, Champ. in Calc. Journ. Nat. Hist. vii. (1847) 468.

SCIAPHILA:—
Hyalisma, Champ. in Calc. Journ. Nat. Hist. vii. (1847) 468.
Soridium, Miers, in Trans. Linn. Soc. xxi. (1852) 47, 49. t. 6.
albescens, *Benth. in Hook. Kew Journ.* vii. (1855) 11.—Bras.
consimilis, *Blume, Mus. Bot. Lugd. Bat.* i. 322.—Ins. Philipp.
corymbosa, *Benth. in Hook. Kew Journ.* vii. (1855) 12.—Bras.
erubescens, *Miers, in Proc. Linn. Soc.* ii. (1850) 74; *et in Trans. Linn. Soc.* xxi. (1852) 48.—Zeylan.
janthina, *Thw. Enum. Pl. Zeyl.* 294.—Ind. or.; Zeylan.
khasiana, *Benth. & Hook. f. Gen.* iii. 1003.—Reg. Himal.
maculata, *Miers, in Proc. Linn. Soc.* ii. (1850) 73; *et in Trans. Linn. Soc.* xxi. (1852) 48.—Ins. Philipp.
nana, *Blume, Mus. Bot. Lugd. Bat.* i. 322.—Java.
picta, *Miers, in Proc. Linn. Soc.* ii. (1850) 73; *et in Trans. Linn. Soc.* xxi. (1852) 48.—Venezuela.
purpurea, *Benth. in Hook. Kew Journ.* vii. (1855) 11.—Bras.
secundiflora, *Thw. ex Benth. l. c.* 10.—Zeylan.
tenella, *Blume, Bijdr.* 515.—Java.

SCILLA, Linn. Syst. ed. I (1735); Gen. ed. I. 95 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 814.
Adenoscilla, Gren. & Godr. Fl. Fr. iii. 187 (1856).
Agraphis, Link, Handb. i. 166 (1829).
Barnardia, Lindl. Bot. Reg. t. 1029 (1826).
Basaltogeton, Salisb. Gen. Pl. Fragm. 27 (1866).
Caloscilla, Jord. et Fourr. Ic. Fl. Europ. ii. 14. t. 245 (1869).
Endymion, Dum. Fl. Belg. 140 (1840).
Epimenidion, Henonix, Kozola, Lagocodes, Oncostema, Skilla, Syncodium, Tractema, Rafin. Fl. Tellur. ii. 13, 22, 25, 57.
Eratobotrys, Fenzl, in Endl. Gen. Pl. Suppl. ii. 13 (1842).
Hyacinthoides, Medic. in Usteri, Ann. Bot. ii. (1791) 9.
Hylomenes, Salisb. Gen. Pl. Fragm. 26 (1866).
Ioncomelos, Rafin. Fl. Tellur. iii. 62 (1836).
Ledebouria, Roth, Nov. Pl. Sp. 194 (1821).
Monocallis, Salisb. Gen. Pl. Fragm. 27 (1866).
Nectaroscilla, Parl. Nov. Gen. Monocot. 27 (1854).
Othocallis, Salisb. Gen. Pl. Fragm. 28 (1866).
Petranthe, Salisb. l. c. 27 (1866).
Prospero, Salisb. l. c. 28 (1866).
Rinopodium, Salisb. l. c. 28 (1866).
Somera, Salisb. l. c. 26 (1866).
Stellaris, Dill. ex Moench, Meth. 303 (1794).
Sugillaria, Salisb. Gen. Pl. Fragm. 18 (1866).
Simira, Rafin. Fl. Tellur. iv. 28 (1836).
Usteria, Medic. in Act. Theod. Palat. vi. Phys. (1790) 480.
Xeodolon, Salisb. Gen. Pl. Fragm. 18 (1866).
acaulis, *Baker, in Journ. Bot.* xii. (1874) 364.—Peruv.
alliifolia, Lapeyr. Hist. Abr. Pl. Pyr. i. 188 = verna.
alpina, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 90; iv. (1853) 75 = bifolia.
alvesiana, Welw. ex Nym. Consp. 729 = vincentina.
amara, Crantz, Inst. i. 495 = amoena.
amethystina, Salisb. Prod. 243 = Hyacinthus amethystinus.
amethystina, Vis. in Flora, xii. (1829) i. Erg. 11 = pratensis.
amoena, Bieb. Fl. Taur. Cauc. i. 278; [DC. in] Red. Lil. iii. t. 130 = sibirica.
amoena, Bory & Chaub. Nouv. Fl. Pélop. 23 = messeniaca.
amoena, *Linn. Sp. Pl.* 309.—Europ. austr.
amoenula, Hornem. Hort. Hafn. i. 331 = sibirica.
angusta, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 237 = Camassia Fraseri.
anthericoides, Poir. Voy. Barb. ii. 149 = Urginea anthericoides.
anthericoides, Willd. ex Kunth, Enum. Pl. iv. 332 = Urginea Scilla.
arenaria, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 249.—Afr. trop.
Aristides, *Coss. in Bull. Soc. Bot. Fr.* v. (1858) 104.—Alger.
autumnalis, Curt. Bot. Mag. t. 919 = obtusifolia.
autumnalis, *Linn. Sp. Pl.* 309.—Reg. Mediterr. & Caucas.

SCILLA:—
azurea, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 125 = sibirica.
Barberi, *Baker, in Journ. Linn. Soc.* xiii. (1873) 247.—Afr. austr.
Barnardia, *C. Koch, in C. Koch & Fint. Wochenschr.* i. (1858) 316.—China.
Bellii, *Baker, in Gard. Chron.* (1884) ii. 488.—Persia.
benguellensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 249.—Afr. trop.
Berthelotii, *Webb & Berth. Phyt. Canar.* iii. 337.—Ins. Canar.
Bertolonii, Duby, Bot. Gall. ed. II. i. 465 = italica.
biflora, Ruiz & Pav. Fl. Per. iii. 69. t. 302 = Ornithogalum biflorum.
bifolia, Hort. Berol. ex Vilm. Fl. Pl. Terre, ed. III. 1046 = sibirica.
bifolia, *Linn. Sp. Pl.* 309.—Europ.; As. Min.
bifolia, Oed. Fl. Dan. t. 568 = verna.
bifolia, Petter, ex Baker, in Journ. Linn. Soc. xi. (1871) 428 = Hyacinthus dalmaticus.
bifolia, Retz. in Vet. Akad. Handl. Stockh. (1769) 246 = italica.
bifolia, Sieber, ex Schult. f. Syst. vii. 581 = Chionodoxa nana.
bifolia, Thunb. Fl. Jap. 138 = orientalis.
bifolia, Ucria, Hort. Reg. Panorm. 156 = obtusifolia.
bipartita, Salzm. ex Baker, in Journ. Linn. Soc. xiii. (1873) 244 = lingulata.
bithynica, Boiss. Diagn. Ser. I. vii. 111 = amoena.
blanda, Salisb. Prod. 244 = Lachenalia orchioides.
brachyphylla, Schult. f. Syst. vii. 573 = Hyacinthus brevifolius.
brevifolia, Ker-Gawl. in Bot. Mag. t. 1468; Schult. f. Syst. vii. 574 = Hyacinthus brevifolius.
brumalis, Haw. ex G. Don, in Sweet, Hort. Brit. ed. III. 697, nomen (Quid?).—Hab.?
byzantina, Hort. Par. ex Poir. Encyc. vi. 739 = italica.
camerooniana, *Baker, in Saund. Refug. Bot.* iii. *App.* 9.—Afr. trop.
campanulata, Ait. Hort. Kew. ed. I. i. 444 = hispanica.
campestris, Savi, Fl. Pis. i. 350 = Ornithogalum umbellatum.
candida, Guss. Fl. Sic. Syn. ii. 814, in obs. = peruviana.
capensis, Burm. f. Fl. Cap. Prod. 10 = Urginea altissima.
carnea, Sweet, Hort. Brit. ed. II. 528 = bifolia.
cernua, [Delar. in] Red. Lil. sub t. 298 = sibirica.
cernua, Janka, in Oestr. Bot. Wochenbl. v. (1855) 230 = bifolia.
cernua, Salisb. Prod. 242 = festalis.
chinensis, *Benth. Fl. Hongk.* 373.—China.
Chladnii, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 39 = bifolia.
chloroleuca, Kunth, Enum. Pl. iv. 325 = Ornithogalum biflorum.
ciliaris, Hort.—Cf. Gard. Chron. (1875) i. 306 = peruviana.
ciliolata, *Pomel, Nouv. Mat. Fl. Atl.* 385.—Alger.
Clusiana, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 795 = peruviana.
Clusii, Parl. Fl. Ital. ii. 462 = peruviana.
comata, Hoffmgg. Verz. Pfl. 111 = peruviana.
comosa, Salisb. Prod. 243 = Muscari comosum.
concinna, *Baker, in Saund. Refug. Bot.* t. 235.—Afr. austr.
concinna, Salisb. Prod. 242 = lusitanica.
concolor, *Baker, in Saund. Refug. Bot.* iii. *App.* 13.—Afr. austr.
congesta, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 250.—Afr. trop.
Cooperi, *Hook. f. Bot. Mag.* t. 5580.—Afr. austr.
coromandeliana, Roxb. Hort. Beng. 24; Fl. Ind. ii. 147 = Urginea indica.
coronaria, Salisb. Prod. 243 = Hyacinthus orientalis.
corsica, Boullu, in Ann. Soc. Bot. Lyon, v. (1878) 88 = autumnalis.
corymbosa, Ker-Gawl. in Bot. Mag. sub t. 1468 = Hyacinthus corymbosus.
Cundra, Buch.-Ham. in Wall. Cat. n. 5062 = Urginea indica.
Cupani, *Guss. Ind. Sem. Hort. Boccad.* (1825) 10; *Fl. Sic. Prod.* i. 416.—Sicilia.
Cupaniana, Schult. f. Syst. vii. 559 = praec.
Currori, *Baker, in Journ. Linn. Soc.* xiii. (1873) 249.—Afr. trop.

SCILLA:—
dalmatica, Presl, ex Avé-Lall. in Ind. Sem. Hort. Petrop. xi. 71 = Muscari pallens.
dasyantha, Webb & Berth. Phyt. Canar. iii. 337. t. 231 = haemorrhoidalis.
decora, Salisb. Prod. 243 = Lachenalia pendula.
denudata, Buch.-Ham. in Wall. Cat. n. 5062 = Urginea indica.
dubia, C. Koch, in Linnaea, xix. (1847) 315 = bifolia.
dumetorum, Balansa, ex Baker, in Journ. Linn. Soc. xiii. (1873) 234 = autumnalis.
elongata, Parl. Nuov. Gen. e Sp. 24 = peruviana.
eriophora, Mill. Gard. Dict. ed. VIII. n. 10 = hyacinthoides.
esculenta, Ker-Gawl. in Bot. Mag. t. 1574 = Camassia Fraseri.
exigua, *Baker, in Journ. Linn. Soc.* xiii. (1873) 247.—Afr. austr.
fallax, Steinheil, in Ann. Sc. Nat. Sér. II. i. (1834) 103 = obtusifolia.
fastigiata, Viv. App. Fl. Cors. Prod. 1 = Hyacinthus fastigiatus.
festalis, *Salisb. Prod.* 242.—Europ.
firmifolia, *Baker, in Saund. Refug. Bot.* iii. *App.* 7.—Afr. austr.
fistulosa, Rafin. Préc. Découv. 45 = Cupani.
flaccidula, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 249.—Afr. trop.
flexuosa, *Baker, in Journ. Linn. Soc.* xiii. (1873) 245.—Afr. austr.
floribunda, *Baker, in Saund. Refug. Bot.* t. 188.—Afr. austr.
fragrans, Salisb. Prod. 243 = Muscari moschatum.
Fraseri, A. Gray, Man. Bot. N. U. St. ed. V. 533 = Camassia Fraseri.
fugax, Munby, Fl. Alger. 36 = Urginea fugax.
gallica, Todaro, Ind. Sem. Hort. Panorm. (1872) 39; et in Nuov. Giorn. Bot. Ital. v. (1873) 157 = autumnalis.
geminiflora, Kunth, Enum. Pl. iv. 325 = Ornithogalum biflorum.
Gerrardi, *Baker, in Journ. Linn. Soc.* xiii. (1873) 237.—Afr. austr.
haemorrhoidalis, *Webb & Berth. Phyt. Canar.* iii. 336. t. 230.—Ins. Canar.
Hanburii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 235.—Oriens.
hemisphaerica, Boiss. Voy. Bot. Espagne, ii. 613 = peruviana.
hirsuta, Willd. ex Kunth, Enum. Pl. iv. 319 = villosa.
hispanica, *Mill. Gard. Dict.* ed. VIII. n. 8.—Hispan.
hispida, J. Gay, ex Baker, in Journ. Linn. Soc. xi. (1871) 430 = Hyacinthus hispidus.
hispidula, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 248.—Afr. trop.
Hohenackeri, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 79.—Reg. Caucas.
Hohenackeri, Janka, in Oestr. Bot. Wochenbl. v. (1855) 230 = bifolia.
Hughii, Bertol. Fl. Ital. iv. 108, in syn. = peruviana.
humifusa, *Baker, in Gard. Chron.* (1881) i. 626.—Afr. austr.
hyacinthoides, Desf. ex Link, in Buch, Beschr. Canar. Ins. 140 = haemorrhoidalis.
hyacinthoides, Jacq. Coll. i. 61 = hispanica.
hyacinthoides, *Linn. Syst.* ed. XII. 243.—Europ. austr.
indica, *Baker, in Saund. Refug. Bot.* iii. *App.* 12.—Ind. or.; Abyss.
indica, Roxb. Hort. Beng. 24; Fl. Ind. ii. 147 = Urginea indica.
inflata, Salisb. Prod. 243 = Muscari botryoides.
intermedia, Guss. Ind. Sem. Hort. Boccad. (1825); Fl. Sic. Prod. i. 417 = obtusifolia.
iridifolia, Webb & Berth. Phyt. Canar. iii. 338 = latifolia.
italica, DC. Fl. Fr. iii. 213 = hyacinthoides.
italica, Host, Fl. Austr. i. 434 = pratensis.
italica, Linn. Herb. ex Baker, in Journ. Linn. Soc. xiii. (1873) 242 = verna.
italica, *Linn. Sp. Pl.* 308.—Europ. austr.
Jacquini, J. F. Gmel. Syst. 552 = hispanica.
japonica, *Baker, in Journ. Linn. Soc.* xiii. (1873) 233.—Japon.
japonica, Thunb. Fl. Jap. 137 = Heloniopsis japonica.
Kirkii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 254.—Afr. trop.

SCILLA :—
Kladnii, Schur, Enum. Pl. Transs. 668 = bifolia.
Kraussii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 248.—Afr. austr.
lanceaefolia, *Baker, in Saund. Refug. Bot.* t. 182.—Afr. austr.
lanceolata, *Baker, l. c.* iii. *App.* 14.—Afr. austr.
lanceolata, Viv. App. Fl. Cors. Prod. ii. 3 = Urginea Scilla.
latifolia, *Willd. ex Schult. f. Syst.* vii. 566.—Ins. Canar.
laxa, Schur, Enum. Pl. Transs. 669 = bifolia.
laxiflora, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 250.—Afr. trop.
ligulata, Steud. Nom. ed. I. 747 = lingulata.
lilacina, *Baker, in Saund. Refug. Bot.* iii. *App.* 13.—Nubia.
Lilio-hyacinthus, *Linn. Sp. Pl.* 308.—Europ. austr.
linearifolia, *Baker, in Saund. Refug. Bot.* t. 184.—Afr. austr.
lingulata, *Poir. Voy. Barb.* ii. 151.—Afr. bor.
livida, *Baker, in Gard. Chron.* (1883) ii. 166.—Afr. austr.
lorata, *Baker, in Saund. Refug. Bot.* iii. *App.* 14.—Afr. austr.
Ludwigii, *Baker, l. c.* 9.—Afr. austr.
lusitanica, *Linn. Syst.* ed. XII. 243.—Hispan.
lusitanica, Sims, Bot. Mag. t. 1999 = latifolia.
lusitanica, Willd. ex Kunth, Enum. Pl. iv. 321 = monophyllos.
Macowani, *Baker, in Gard. Chron.* (1875) i. 748.—Afr. austr.
macrogona, *Link, in Schrad. Journ.* ii. (1799) 319.—Lusitan.
maculata, *Baker, in Journ. Linn. Soc.* xiii. (1873) 250.—Ind. or.
maculata, Schrank, Pl. Rar. Hort. Monac. t. 100 = lanceaefolia.
maesta, *Baker, in Saund. Refug. Bot.* iii. *App.* 10.—Afr. trop.
maritima, Linn. Sp. Pl. 308 = Urginea Scilla.
mauritanica, Schousb. Vextr. Marokko, 168 = vincentina.
mauritiana, Steud. Nom. ed. I. 747 = praec.
messeniaca, *Boiss. Diagn.* Ser. I. vii. 110.—Graecia.
micrantha, A. Rich. Tent. Fl. Abyss. ii. 328 = Urginea micrantha.
microscypha, *Baker, in Gard. Chron.* (1881) ii. 102.—Afr. austr.
minima, *Baker, in Saund. Refug. Bot.* iii. *App.* 6.—Afr. austr.
minor, C. Koch, in Linnaea, xix. (1847) 9 = bifolia.
monanthos, C. Koch, l. c. xxii. (1849) 251 = sibirica.
monophylla, Plan., Ens. Fl. Gallega, 380 = seq.
monophyllos, *Link, in Schrad. Journ.* ii. (1799) 319.—Lusitan.
monstrosa, Salisb. Prod. 243 = Muscari comosum.
montana, Savi, Due Cent. 89 = Ornithogalum narbonense.
natalensis, *Planch. in Fl. des Serres*, x. (1854–55) 185. t. 1043.—Afr. austr.
nivalis, Boiss. Diagn. Ser. I. v. 63 = bifolia.
nonscripta, Hoffmgg. & Link, in Ges. Naturf. Fr. Neue Schrift. iv. (1803) 19 = festalis.
numidica, *Poir. Voy. Barb.* ii. 150.—Alger.
nutans, Sm. Engl. Bot. t. 377 = festalis.
obtusifolia, *Poir. Voy. Barb.* ii. 149.—Sardin.; Afr. bor.
odorata, *Link, in Schrad. Journ.* ii. (1799) 319; *Brot. Fl. Lusit.* i. 527.—Hispan.
orientalis, *Thunb. in Trans. Linn. Soc.* ii. (1794) 334.—Japon.
ovalifolia, Hort.—Cf. Gard. Chron. (1871) 6 (Quid ?).—Hab. ?
ovatifolia, Baker, in Saund. Refug. Bot. t. 183 = lanceaefolia.
pallida, Salisb. Prod. 241 = italica.
pallidiflora, *Baker, in Saund. Refug. Bot.* t. 179.—Afr. austr.
Pancration, Nym. Syll. 369 = Urginea Scilla.
paradoxa, Bieb. Fl. Taur. Cauc. iii. 267 = Allium paradoxum.
parviflora, Desf. Fl. Atlant. i. 300. t. 87 = numidica.
parviflora, Guss. Cat. Rar. Hort. Boccad. 57; ex ej. Fl. Sic. Prod. i. 418 = obtusifolia.
parviflora, Salisb. Prod. 242 = hyacinthoides.
patula, DC. Fl. Fr. iii. 211 = hispanica.
pauciflora, Link, ex Schult. f. Syst. vii. 561 = italica.

SCILLA :—
paucifolia, *Baker, in Saund. Refug. Bot.* t. 181.—Afr. austr.
pendula, *Baker, l. c.* iii. *App.* 14.—Afr. austr.
peruviana, *Linn. Sp. Pl.* 309.—Reg. Mediterr.
Petitiana, A. Rich. Tent. Fl. Abyss. ii. 328 = Urginea Petitiana.
platyphylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 250.—Afr. trop.
plumbea, *Lindl. Bot. Reg.* t. 1355.—Afr. austr.
polyantha, *Baker, in Gard. Chron.* (1878) i. 104.—Afr. austr.
polyphylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 249.—Afr. trop.
polyphylla, Royle, Illustr. Bot. Himal. 391 (= *Ornithogalum polyphyllum*).—Ind. or.
pomeridiana, DC. Cat. Hort. Monsp. (1813) 143 = Chlorogalum pomeridianum.
praebracteata, Haw. ex G. Don, in Sweet, Hort. Brit. ed. III. 697 = peruviana.
praecox, Donn, Hort. Cantab. ed. II. 41 = sibirica.
praecox, Willd. Sp. Pl. ii. 128 = bifolia.
prasina, *Baker, in Saund. Refug. Bot* iii. *App.* 10.—Afr. austr.
pratensis, Berger. ex Schult. f. Syst. vii. 563 = verna.
pratensis, C. Koch, in Linnaea, xix. (1847) 9 = sibirica.
pratensis, *Waldst. & Kit. Pl. Rar. Hung.* ii. 207. t. 189.—Croatia.
Princeps, *Baker, in Saund. Refug. Bot.* t. 189.—Afr. austr.
pubens, Welw. ex Lindl. Bot. Reg. (1845) Misc. 40 = peruviana.
pulchella, Munby, in Bull. Soc. Bot. Fr. ii. (1855) 286 = autumnalis.
pumila, Brot. Fl. Lusit. i. 527 = monophyllos.
purpurea, Mill. Gard. Dict. ed. VIII. n. 9 = italica.
puschkinioides, *Regel, in Act. Hort. Petrop.* iii. ii. (1875) 295.—Turkestan.
pusilla, *Baker, in Journ. Bot.* xiv. (1876) 183.—Afr. austr.
Quartiniana, A. Rich. Tent. Fl. Abyss. ii. 329 = Urginea Quartiniana.
racemosa, Balansa, ex Baker, in Journ. Linn. Soc. xiii. (1873) 234 = autumnalis.
Raewskiana, *Regel, in Act. Hort. Petrop.* viii. (1884) 655.—Turkestan.
Ramburei, Boiss. Elench. 86 = verna.
revoluta, *Baker, in Saund. Refug. Bot.* iii. *App.* 6.—Afr. austr.
Richardiana, *Buching. ex Baker, l. c.* 11.—Abyss.
rigidifolia, *Kunth, Enum. Pl.* iv. 330.—Afr. austr.
romana, Ker-Gawl. in Bot. Mag. t. 939 = Hyacinthus romanus.
rosea, Lehm. Ind. Sem. Hort. Hamb. (1828) 17 = bifolia.
Roseni, C. Koch, in Linnaea, xxii. (1849) 250 = sibirica.
Sandersoni, *Baker, in Saund. Refug. Bot.* iii. *App.* 5.—Afr. austr.
saturata, *Baker, in Journ. Bot.* xii. (1874) 365.—Afr. austr.
secunda, Janka, in Oestr. Bot. Wochenbl. vi. (1856) 194 = bifolia.
serotina, Ker-Gawl. in Bot. Mag. t. 859 = Dipcadi serotinum.
serotina, Schousb. Vextr. Marokko, 165 = Urginea anthericoides.
sibirica, *Andr. Bot. Rep.* t. 365.—Ross.; As. Min.
sicula, Tineo, in Guss. Fl. Sic. Syn. ii. 813 = peruviana.
simensis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 327 = Urginea simensis.
simiarum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 249.—Afr. trop.
socialis, *Baker, in Saund. Refug. Bot.* t. 180.—Afr. austr.
sordida, Salisb. Prod. 243 = Dipcadi serotinum.
spathulata, *Baker, in Saund. Refug. Bot.* t. 187.—Afr. austr.
spicata, *Baker, in Journ. Bot.* xvi. (1878) 328.—Afr. trop.
squamosa, Dulac, Fl. Hautes-Pyr. 109 = S. Lilio-hyacinthus.
stellaris, Salisb. Prod. 241 = peruviana.
Strangwaysii, *Tenore, Ind. Sem. Hort. Neap.* (1839) 12. n. 11.—Byzantium.
suaveolens, Salisb. Prod. 243 = Muscari racemosum.

SCILLA :—
subglauca, *Baker, in Saund. Refug. Bot.* t. 186.—Afr. austr.
subnivalis, Nym. Consp. 730 = bifolia.
subsecunda, *Baker, in Gard. Chron.* (1881) ii. 38.—Afr. austr.
subtriphylla, Schur, Enum. Pl. Transs. 668 = bifolia.
sylvatica, Czetz, 'Eod. Muz. (1872) 20' = bifolia.
sylvestris, Savi, Due Cent. 88 = Ornithogalum pyrenaicum.
tetraphylla, Linn. f. Suppl. 200 = Chlorophytum tetraphyllum.
tingitana, Schousb. Vextr. Marokko, 169 = monophyllos.
tricolor, *Baker, in Gard. Chron.* (1880) ii. 230.—Afr. austr.
triflora, *Phil. Fl. Atac.* 51.—Chili.
trifolia, Schur, Enum. Pl. Transs. 668 = bifolia.
Ughii, Tineo, ex Guss. Fl. Sic. Prod. 162 = peruviana.
umbellata, Ram. in Bull. Soc. Philom. ii. (1800) 130. t. 8. f. 6 = verna.
undulata, *Baker, in Saund. Refug. Bot.* iii. *App.* 11.—Afr. austr.
undulata, Desf. Fl. Atlant. 300. t. 88 = Urginea undulata.
uniflora, Willd. ex Kunth, Enum. Pl. iv. 318 = sibirica.
unifolia, Linn. Sp. Pl. 309 = Ornithogalum unifolium.
verna, Bourg. ex Nym. Consp. 729 = lusitanica.
verna, *Huds. Fl. Angl.* ed. II. 142.—Europ. occ.
verna, Koch, Syn. Fl. Germ. ed. I. 714 = italica.
verna, Moris, Stirp. Sard. Elench. 47 = Hyacinthus fastigiatus.
vernalis, Salisb. Prod. 242 = verna.
versicolor, *Baker, in Saund. Refug. Bot.* t. 305.—Afr. austr.
versicolor, Salisb. Prod. 214 = Lachenalia tricolor.
villosa, *Desf. Fl. Atlant.* i. 299. t. 85. f. 2.—Afr. bor.
vincentina, *Link & Hoffmgg. in Ges. Naturf. Fr. Neue Schrift.* iv. (1803) 17.—Lusitan.
viridiflora, *Baker, in Journ. Linn. Soc.* xiii. (1873) 255.—Afr. austr.
viridis, Salisb. Prod. 243 = Dipcadi filamentosum.
Vivianii, Bertol. Fl. Ital. x. 517 = peruviana.
xanthandra, C. Koch, in Linnaea, xix. (1847) 316, in obs. = sibirica.
zambesiaca, *Baker, in Saund. Refug. Bot.* iii. *App.* 8.—Afr. trop.
zebrina, *Baker, l. c.* t. 185.—Afr. austr.

SCILLOPSIS, Lem. in Illustr. Hortic. ii. (1855) Misc. 96; iii. (1856) Misc. 33 = **Lachenalia**, Jacq. (Liliac.).
anguinea, Lem. l. c. iii. (1856) 34 = L. anguinea.
angustifolia, Lem. l. c. 33 = L. contaminata.
bifolia, Lem. l. c. 35 = L. isopetala.
contaminata, Lem. l. c. 34 = L. contaminata.
fragrans, Lem. l. c. 35 = L. versicolor.
hyacinthoides, Lem. l. c. = L. contaminata.
isopetala, Lem. l. c. = L. isopetala.
liliiflora, Lem. l. c. = L. liliiflora.
lucida, Lem. l. c. 34 = L. lucida.
mediana, Lem. l. c. 35 = L. mediana.
nervosa, Lem. l. c. 34 = L. nervosa.
orthopetala, Lem. l. c. = L. orthopetala.
patula, Lem. l. c. 35 = L. patula.
purpurea, Lem. l. c. = L. versicolor.
purpureo-coerulea, Lem. l. c. = L. purpureo-coerulea.
pustulata, Lem. l. c. 34 = L. pustulata.
racemosa, Lem. l. c. = L. racemosa.
rosea, Lem. l. c. 35 = L. isopetala.
unicolor, Lem. l. c. 34 = L. versicolor.
unifolia, Lem. l. c. = L. unifolia.
violacea, Lem. l. c. 35 = L. violacea.

SCINDAPSUS, Schott, Melet. 21 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 992.
CUSCUARIA, Schott, in Bonplandia, v. (1857) 45.
angulatus, Miq. Fl. Ind. Bat. iii. 219 = Raphidophora montana.
angustifolius, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 164 = Rhaphidophora angustifolia.
anomalus, Carr. in Rev. Hortic. (1884) 536 = Monstera acuminata.
arborum, Presl, Epim. Bot. 242 = Pothos macrostachyus.
argyraeus, Engl. in DC. Monog. Phan. ii. 255 = pictus.
aruensis, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 270.—Ins. Aru.

SCINDAPSUS:—
bipinnatifidus, Teijsm. & Binn. Cat. Hort. Bog. 65 = Epipremnum mirabile.
Beccarii, *Engl. Bot. Jahrb.* i. (1881) 182.—Sumatra.
cannaefolius, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 271.—Borneo.
caudatus, Schott, Melet. 21 = Epipremnum mirabile.
coriaceus, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 271.—Borneo.
crassipes, *Engl. Bot. Jahrb.* i. (1881) 182.—Borneo.
Cuscuaria, *Presl, Epim. Bot.* 242.—Malaya.
decursivus, Mor. Syst. Verz. Zoll. 82 = Epipremnum mirabile.
decursivus, Schott, Melet. 21 = Raphidophora decursiva.
dilaceratus, C. Koch & H. Sello, Ind. Sem. Hort. Berol. (1853) App. 5 = Monstera dilacerata.
erectus, Presl, Epim. Bot. 242 = Aglaonema marantifolium.
Forsteri, Endl. in Ann. Wien. Mus. i. (1836) 161 = Raphidophora pertusa.
geniculatus, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 271.—Borneo.
giganteus, Schott, Melet. 21 = Epipremnum giganteum.
glaucus, Schott, l. c. = Raphidophora glauca.
Haenkii, Presl, Epim. Bot. 240 = Aglaonema Haenkei.
hederaceus, *Schott, in Bonplandia*, v. (1857) 45.—Malaya.
inaequilaterus, Presl, Epim. Bot. 240 = Pothos Cumingianus.
inquinatus, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 283 = hederaceus.
lingulatus, Hassk. Cat. Hort. Bog. Alt. 58 = Raphidophora lingulata.
longipes, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 270.—Borneo.
marantaefolius, Miq. Fl. Ind. Bat. iii. 187 = S. Cuscuaria.
medius, Zoll. & Mor. Syst. Verz. Zoll. 82 = Epipremnum medium.
microstachyus, De Vriese & Miq. in Miq. Fl. Ind. Bat. iii. 188 = Anadendrum affine.
montanus, Kunth, Enum. Pl. iii. 64 = Anadendrum montanum.
occidentalis, Poepp. & Endl. Nov. Gen. et Sp. iii. 88 = Monstera occidentalis.
officinalis, *Schott, Melet.* 21.—Ind. or.; Burma.
Peepla, Schott, l. c. = Raphidophora Peepla.
Peepla, Thw. Enum. Pl. Zeyl. 336 = Raphidophora pertusa.
pertusus, Schott, Melet. 21 = Raphidophora pertusa.
pictus, *Hassk. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 164.—Malaya.
pinnatifidus, Schott, Melet. 21 = Epipremnum mirabile.
pinnatus, Schott, l. c. = Epipremnum mirabile.
pothoides, Miq. Fl. Ind. Bat. iii. 184 = hederaceus.
pothoides, Schott, Prod. Aroid. 394 = pictus.
pteropodus, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* xxvii. (1864) 23.—Sumatra.
Rumphii, Presl, Epim. Bot. 242 = Pothos Rumphii.
sylvestris, Kunth, Enum. Pl. iii. 64 = Raphidophora angustifolia.
tener, Presl, Epim. Bot. 242 = Pothos gracilis, tener.

SCIOBIA, Reichb. Nom. 66 (1841) = SCIOPHILA, Gaudich. = **Procris**, Comm. (Urtic.).

SCIODAPHYLLUM, Blume, Bijdr. 875 (1826) = **Heptapleurum**, Gaertn. (Araliac.).

SCIODAPHYLLUM, P. Br. Hist. Jamaic. 190. t. 19 (1756) = **Sciadophyllum**, P. Br. (Araliac.).

SCIOPHILA, Gaudich. in Freyc. Voy. Bot. 493. t. 120 (1826) = **Procris**, Comm. (Urtic.).
cephalida, Gaudich. l. c. 494 = P. cephalida.
frutescens, Gaudich. l. c. 493, in nota absque diagn. = P. frutescens?
salicifolia, Gaudich. l. c. 494 = P. cephalida?
sessile, Gaudich. l. c. = Elatostema sessile.
sessilifolia, Gaudich. l. c. = Elatostema sessile?
torresiana, Gaudich. l. c. 493. t. 120 = P. cephalida.

SCIOPHILA, Wibel, Prim. Fl. Werth. 147 (1799) = **Maianthemum**, Weber (Liliac.).
convallarioides, Wibel, l. c. = M. Convallaria.

SCIOPHYLLA, Heller, Fl. Wirceb. 158 (1810) = praec.

SCIOTHAMNUS, Endl. Gen. 780 (1839) = **Peucedanum**, Tourn. (Umbell.).
capensis, Endl. ex Walp. Rep. ii. 412 = P. capense.
collinus, Endl. l. c. 413 = P. collinum.
montanus, Endl. l. c. = P. Ecklonianum.
virgatus, Endl. l. c. = P. capense.

SCIRPIDIUM, Nees, in Linnaea, ix. (1834) 293 = **Eleocharis**, R. Br. (Cyperac.).
aciculare, Nees, l. c. 293 = E. acicularis.
grande, Nees, in Mart. Fl. Bras. ii. 1. 97 (Fimbristylidis sp.).—Bras.
nigrescens, Nees, l. c. = E. nigrescens.
piauhiense, Nees, l. c. = E. sulcata.
sulcatum, Nees, l. c. 98 = E. sulcata.
tenue, Nees, in Linnaea, ix. (1834) 293 = E. tenuis.

SCIRPODENDRON, Zipp. ex Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) II. 84. *CYPERACEAE*, Benth. & Hook. f. iii. 1056.
PTYCHOCARYA, R. Br. in Wall. Cat. n. 3538 (1830).
costatum, *Kurz, in Journ. As. Soc. Beng.* xxxviii. (1869) II. 85.—As. trop.
pandaniforme, Zipp. ex Kurz, l. c. = costatum.
sulcatum, Miq. Illustr. Fl. Arch. Ind. 65. t. 28 = costatum.

SCIRPUS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 12 (1737). *CYPERACEAE*, Benth. & Hook. f. iii. 1049.
ANDROCOMA, Nees, in Hook. Journ. Bot. ii. (1840) 396.
ANTHOPHYLLUM, Steud. Syn. Pl. Cyp. 160 (1855).
APLOSTEMON, Rafin. in Journ. Phys. lxxxix. (1819) 105.
BAEOTHRYON, Ehrh. Beitr. iv. 147 (1789).
BLEPHAROLEPIS, Nees, in Mart. Fl. Bras. ii. 1. 91 (1847).
BLYSMUS, Panz. ex Schult. Mant. ii. 41 (1824).
CLAVULA, Dum. Fl. Belg. 143 (1827), pro parte.
CREPIDOCARPUS, Klotzsch, ex Boeck. in Linnaea, xxxvi. (1869-70) 414.
DESMOSCHOENUS, Hook. f. Fl. N. Zel. i. 271 (1853).
DICHELOSTYLIS, Endl. Gen. 118, in syn. (1836).
DICHISMUS, Rafin. in Journ. Phys. lxxxix. (1819) 106.
DICHOSTYLIS, Beauv. in Lestib. Ess. Fam. Cyp. 39 (1819).
DIPLARINUS, Rafin. in Journ. Phys. lxxxix. (1819) 106.
DISTICHMUS, Endl. Gen. 119 (1837).
ELEOGITON, Link, Hort. Berol. i. 293 (1827).
ELYTROSPERMUM, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 200. t. 2.
HELEOPHILA, Schult. Mant. ii. 2 (1824).
HELEOPHYA, Flora, iv. (1821) 25.
HELEOPHYLAX, Beauv. ex Lestib. Ess. Fam. Cyp. 41 (1819).
HELIOPHYLAX, Steud. Nom. ed. II. i. 743 (1840).
HEMICARPHA, Nees, in Edinb. N. Phil. Journ. xvii. (1834) 263.
HOLOSCHOENUS, Link, Hort. Berol. i. 293 (1827).
HYMENOCHAETA, Beauv. in Lestib. Ess. Fam. Cyp. 43 (1819).
ISOLEPIS, R. Br. Prod. 221 (1810).
LEIOPHYLLUM, Ehrh. Beitr. iv. 146 (1789).
LIMNOCHLOA, Reichb. Fl. Germ. Excurs. 78, pro majore parte.
MALACOCHAETE, Nees, in Linnaea, ix. (1834) 292.
MEGADENUS, Rafin. Neogenyt. 4 (1825).
NEMOCHARIS, Beurl. in Bot. Notiser (1853), 52.
NEMUM, Desv. in Ham. Prod. Pl. Ind. Occ. 13 (1825).
NOMOCHLOA, Beauv. in Lestib. Ess. Fam. Cyp. 37 (1819).
OXYCARYUM, Nees, in Mart. Fl. Bras. ii. 1. 90 (1847).
PTEROLEPIS, Schrad. in Goett. Gel. Anz. (1821) III. 2071.
PTERYGOLEPIS, Reichb. Nom. 236 (1841).
REIGERA, Opiz, Seznam, 83 (1852).
SEIDLIA, Opiz, Natural. xi. (1826) 349.
STENOPHYLLUS, Rafin. Neogenyt. 4 (1825).
ablepharus, Griseb. Cat. Pl. Cub. 240 = cubensis.

SCIRPUS:—
acaulis, *Phil. in Linnaea*, xxix. (1857-58) 78.—Chili.
acicularis, Linn. Sp. Pl. 48 = Eleocharis acicularis.
acicularis, Oed. in Fl. Dan. t. 167 = Eleocharis palustris.
acicularis, Ucria, Hort. Reg. Panorm. 52 = cernuus.
acicularis, Willd. ex Kunth, l. c. 142 = Eleocharis exigua.
acicularis, Zanard. ex Parl. Fl. Ital. ii. 67 = Eleocharis atropurpurea.
acrostachys, Steud. in Flora, xii. (1829) 148 = Ficinia composita.
acuminatus, Muhl. Desc. Gram. 27 = Eleocharis acuminata.
acuminatus, Steud. in. Flora, xii. (1829) 146 = Ficinia acuminata.
acuta, Poir. Encyc. Suppl. v. 102 = Eleocharis acuta.
acutangulus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 213 = Eleocharis fistulosa.
acutus, Muhl. Desc. Gram. 33 = lacustris.
acutus, J. & C. Presl, Rel. Haenk. i. 192 = mucronatus.
acutus, Spreng. Syst. i. 203 = Eleocharis acuta.
acutus, Willd. ex Kunth, Enum. Pl. ii. 141 = Eleocharis acicularis.
adscendens, *Willd. ex Kunth, l. c.* 198.—Aegypt.
aegyptiacus, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 196 = littoralis.
aegyptiacus, *Poir. Encyc.* vi. 770.—Aegypt.; Ital.
aemulans, Steud. Syn. Pl. Cyp. 87 = grossus.
aestivalis, Retz. Obs. iv. 12 = Fimbristylis aestivalis.
aestivalis, Wall. in Roxb. Fl. Ind. ed. Carey, i. 230 = Fimbristylis squarrosa.
affinis, Roth, Nov. Pl. Sp. 30 = maritimus.
afflatus, Benth. Fl. Hongk. 394 = Eleocharis afflata.
aggregatus, Ruiz, ex Boeck. in Linnaea, xxxvi. (1869-70) 709 = pungens.
albescens, Phil. f. Cat. Pl. Vass. Chil. 310 = inundatus.
albo-ater, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 102 = Eleocharis ocreata.
albovittatus, Schur, Enum. Pl. Transs. 694, in syn. = S. Holoschoenus.
albus, Salisb. Prod. 31 = Rynchospora alba.
alpinus, Schleich. Cat. Pl. Helv. ed. IV. 32 = pumilus.
altissimus, Gilib. Exercit. ii. 514 = lacustris.
ambiguus, Schult. Mant. ii. 85 = divaricatus.
amentaceus, Griseb. in Goett. Abh. vii. (1857) 270 = Fimbristylis complanata.
americanus, Pers. Syn. i. 68 = pungens.
anceps, Poir. Encyc. vi. 782 = Cladium iridifolium.
anceps, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 287 = Fimbristylis complanata.
andinus, *Phil. in Anal. Univ. Chil.* (1873) 554.—Chili.
androgyna, Poir. Encyc. Suppl. v. 98 = Fimbristylis androgyna.
Andrzejowscii, Bess. ex Schult. Mant. ii. 536 = lacustris.
angachillensis, Boeck. in Linnaea, xxxvi. (1869-70) 496 = inundatus.
angulatus, Willd. ex Kunth, Enum. Pl. ii. 155 = Eleocharis fistulosa.
anisochaetus, *Wright, in Sauv. Fl. Cub.* 174.—Cuba.
annuus, All. Fl. Pedem. ii. 277 = Fimbristylis diphylla.
annuus, Host, Gram. Austr. iii. 42. t. 63 = Fimbristylis dichotoma.
annuus, Thuill. Fl. Par. ed. II. 22 = Eleocharis ovata.
anomalus, Retz. Obs. v. 15 = Hypolytrum latifolium.
antarcticus, *Linn. Mant.* ii. 181.—Afr. austr.
antarcticus, Thunb. Prod. Pl. Cap. 17 = diabolicus?
antarcticus, Vahl, Enum. ii. 261, partim (= *Isolepis barbata*).—Geront. trop.
apus, A. Gray, in Proc. Am. Acad. x. (1874) 78 = Frimbristylis Vahlii.
aquatilis, *Rafin. in Med. Repos. N. York*, v. (1808) 352, nomen.—Am. bor.
arabicus, Ehrenb. ex Boeck. in Linnaea, xxxvi. (1869-70) 716 = littoralis.
arenarius, *Benth. Fl. Austral.* vii. 325.—Austral.
arenarius, Boeck. in Linnaea, xxxvi. (1869-70) 741 = Fimbristylis arenaria.
argenteus, Rottb. Desc. Nov. Pl. 51; Progr. 27 = Fimbristylis argentea.
aristatus, Willd. Sp. Pl. i. 300; Vahl, Enum. ii. 271 = Fuirena glomerata.

SCIRPUS:—
articulatus, *Linn. Sp. Pl.* 47.—Afric.; Ind. or.; Austral.
arvensis, Retz. Obs. iv. 11 = Fimbristylis ferruginea.
arvensis, Roxb. Fl. Ind. i. 224 = Fimbristylis diphylla.
asper, *J. & C. Presl, Rel. Haenk.* i. 194.—Chili.
asperulus, Poir. Encyc. Suppl. v. 104; Spreng. Syst. i. 213 (Bulbostylidis sp.).—Am. trop.
atacamensis, Boeck. in Linnaea, xxxvi. (1869–70) 482 = rigidus?
atricha, Poir. Encyc. Suppl. v. 103 = Eleocharis atricha, *R. Br.*
atrofactus, Steud. Syn Pl. Cyp. 317 = Eleocharis capitata.
atropurpureo-vaginatus, Boeck. in Flora, lxv. (1882) 14 = nodosus.
atropurpureus, Retz. Obs. v. 14 = Eleocharis atropurpurea.
atropurpureus, Willd. ex Kunth, Enum. Pl. ii. 150 = Eleocharis capitata.
atrovirens, Willd. Enum. Hort. Berol. 79; Hort. Berol. t. 99 = sylvaticus.
attenuatus, *Franch. & Sav. Enum. Pl. Jap.* ii. 543.—Japon.
aucklandicus, *Boeck. in Linnaea*, xxxvi. (1869–70) 491.—N. Zel.
auro-nitens, Nees & Ehrenb. ex Boeck. l. c. 723 = maritimus.
australis, Murr. Syst. ed. XIII. 85 = S. Holoschoenus.
autumnalis, Linn. Mant. ii. 180 = Fimbristylis autumnalis.
autumnalis, Rottb. Desc. Nov. Pl. 58. t. 17 = Cyperus Haspan.
badius, J. & C. Presl, Rel. Haenk. i. 193 = pungens.
Baeothryon, Linn. f. Suppl. 103 = pauciflorus.
Baeothryon, Timm. ex Kunth, Enum. Pl. ii. 159 = caespitosus.
Baeothryon, Wohlleb. Suppl. 3; ex Schrad. Fl. Germ. i. 127 = Eleocharis palustris.
bahiensis, Steud. Syn. Pl. Cyp. 83 = Eleocharis Sellowiana.
Bailii, Kohts, in Oestr. Bot. Zeitschr. xix. (1869) 334 = caespitosus.
Balduinianus, Schult. Mant. ii. 85 = Fimbristylis diphylla.
balearicus, Willd. ex Kunth, Enum. Pl. ii. 166 = littoralis.
bangalorensis, Heyne, ex Wall. Cat. n. 3464 = corymbosus.
barbata, Poir. Encyc. Suppl. v. 104. (Bulbostylidis sp.).—Geront. trop.
barbatulus, Kunth, Enum. Pl. ii. 583 (Index) = lacustris.
barbatus, Rottb. Desc. Nov. Pl. 52; Progr. 27 = Fimbristylis barbata.
barbatus, Vahl, ex Kunth, Enum. Pl. ii. 211 = Fimbristylis vestita.
barbulatus, Kunth, l. c. 165 = lacustris.
Bellardi, Wahlenb. Fl. Lapp. 15 = Kobresia scirpina.
bengalensis, Pers. Syn. i. 68 = Fimbristylis miliacea.
Bergianus, Spreng. Syst. i. 212 = antarcticus.
Berlandieri, Steud. Syn. Pl. Cyp. 103. (Fimbristylidis sp.).—Mexic.
bicapitatus, Poir. Encyc. vi. 761 = Ficinia bulbosa.
biceps, Roem. & Schult. Syst. ii. 134 = Ficinia bulbosa.
bicolor, Spreng. Syst. iv. Cur. Post. 28 = Thouarsianus.
bifolius, Wallr. Sched. Crit. 27 = rufus.
bispicatus, Koen. ex Roxb. Hort. Beng. 6; Fl. Ind. i. 220 = Fimbristylis schoenoides.
bisumbellatus, Forsk. Fl. Aegypt. Arab. 15 = Fimbristylis dichotoma.
bivalvis, *Lam. Illustr.* i. 141.—Madag.
bonariensis, Poir. Encyc. vi. 763 = Fimbristylis ferruginea.
brachyceras, Hochst. in Flora, xxiv. (1841) i. Intell. 21, nomen; et ex A. Rich. Tent. Fl. Abyss. ii. 496 = corymbosus.
brachyphyllus, Willd. ex Kunth, Enum. Pl. ii. 212 = Fimbristylis barbata.
bracteatus, *Bigel. in New Engl. Med. Journ.* v. (1816) 335 (Quid?).—Am. bor.
Brayii, Hoppe, ex Roem. & Schult. Syst. ii. 137 = lacustris.
breviculmis, Boeck. in Linnaea, xxxvi. (1869–70) 742 (Fimbristylidis sp.).—Afr. austr.
brevifolius, Bernh. ex Kunth, Enum. Pl. ii. 212 = Fimbristylis barbata.

SCIRPUS:—
brevifolius, Poir. Encyc. Suppl. v. 99 = Fimbristylis ferruginea.
brevifolius, Rottb. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 190 (Quid.?).—Ind. or.
brevifolius, Roxb. Hort. Beng. [81]; Fl. Ind. i. 229 = Fimbristylis diphylla.
brevis, Urv. in Mém. Soc. Linn. Par. iv. (1826) 600 = cernuus.
brevivaginatus, Boeck. in Flora, xli. (1858) 413 = pungens.
Bridgesii, Boeck. in Linnaea, xxxvi. (1869–70) 508 = inundatus.
brizoides, *Benth. Fl. Austral.* vii. 326.—Austral.
brizoides, Willd. ex Link, Jahrb. i. III. (1820) 80 = lineatus.
bromeliaefolius, Rudge, Pl. Gui. 19. t. 24 = Diplasia karataefolia.
Brownii, Spreng. Syst. i. 204 = Eleocharis capitata.
brunneo-vaginatus, Boeck. in Kjoeb. Vidensk. Meddel. (1879–80) 25. (Fimbristylidis sp.).—Bras.
brunneus, Muhl. Desc. Gram. 42 = polyphyllus.
bufonius, Poir. Encyc. Suppl. v. 105; Spreng. Syst. i. 213 = Fimbristylis capillaris.
bulbiferus, *Boeck. in Linnaea*, xxxvi. (1869–70) 482.—Afr. austr.
bulbosus, Rottb. Desc. Nov. Pl. 46; Progr. 25 = Ficinia scariosa.
caducus, Delile, Fl. Egypte, 153 = Eleocharis caduca.
caespitosus, Boeck. in Linnaea, xxxvi. (1869–70) 488, partim = pumilus.
caespitosus, *Linn. Sp. Pl.* 48.—Europ.; Am. bor.
caespitosus, Willd. ex Kunth, Enum. Pl. ii. 148 = Eleocharis palustris.
californicus, Steud. Nom. ed. II. ii. 538 = riparius.
campestris, Roth, in Usteri, Ann. Bot. xiv. (1795) 18 = pauciflorus.
campestris, Willd. ex Kunth, Enum. Pl. ii. 148 = Eleocharis palustris.
camptotrichus, *Wright, in Sauv. Fl. Cub.* 173.—Cuba.
camtschaticus, Kunth, Enum. Pl. ii. 158 = Eleocharis palustris.
canaliculato-triqueter, Steud. Syn. Pl. Cyp. 318 = grossus.
Canbyi, A. Gray, in Proc. Acad. Sc. Philad. (1864) 18 = leptolepis.
Candelabrum, Willd. ex Kunth, Enum. Pl. ii. 232 = Fimbristylis diphylla.
capensis, Burm. f. Fl. Cap. Prod. 3 = maritimus.
capensis, Rottb. Desc. Nov. Pl. 53; Progr. 24 = Ficinia bulbosa.
capillaceus, Griseb. Cat. Pl. Cub. 239 = Eleocharis capillacea.
capillaceus, Michx. Fl. Bor. Am. i. 30 = nanus, Eleocharis acicularis.
capillaris, Linn. Sp. Pl. 49 = Fimbristylis capillaris.
capillifolius, *Parl. Fl. Ital.* ii. 83.—Afr. austr.
capitatus, Burm. f. Fl. Ind. 21 = Fuirena uncinata.
capitatus, Linn. Sp. Pl. 48 = Eleocharis capitata.
capitatus, Muhl. Desc. Gram. 30 = Eleocharis albida.
capitatus, Schreb. Spicil. 60 = Eleocharis ovata.
capitatus, Willd. ex Kunth, Enum. Pl. ii. 145 = Eleocharis tenuis.
capsularis, *Lour. Fl. Cochinch.* 44.—China.
caribaeus, Griseb. in Goett. Abh. vii. (1857) 269 = Eleocharis ochreatus.
caribaeus, Rottb. Desc. Nov. Pl. 46; Progr. 24 = Eleocharis capitata.
caricinus, Schrad. Fl. Germ. i. 132 = S. Caricis.
Caricis, *Retz. Fl. Scand. Prod.* 11.—Europ.; As. media.
carinatus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 392.—Am. bor.
carinatus, Sm. Engl. Fl. i. 60 = lacustris.
carolinianus, Lam. Illustr. i. 142 = Fimbristylis castanea.
carolinianus, Panz. ex Kunth, Enum. Pl. ii. 282 = Rynchospora nitens.
cartilagineus, Benth. Fl. Austral. vii. 328 = Gunnii.
cartilagineus, Poir. Encyc. Suppl. v. 103; Spreng. Syst. i. 208 = antarcticus.
castaneus, Michx. Fl. Bor. Am. i. 31 = Fimbristylis castanea.
castaneus, Muhl. Desc. Gram. 38 = Fimbristylis coarctata.
castaneus, Willd. ex Kunth, Enum. Pl. ii. 227, 282 = Fimbristylis autumnalis, Rynchospora nitens.

SCIRPUS:—
Cephalotes, Heyne, ex Roth, Nov. Pl. Sp. 30 = maritimus.
Cephalotes, Jacq. Hort. Vindob. i. 42. t. 97 = Kyllinga monocephala.
Cephalotes, Linn. Sp. Pl. ed. II. 76 = Rynchospora Cephalotes.
Cephalotes, Walt. Fl. Carol. 71 = Lipocarpha Humboldtiana.
cernuus, *Vahl, Enum.* ii. 245.—Cosmop.
cespitosus, Ehrh. Beitr. vi. 84, nomen = caespitosus.
cespitosus, Poir. Encyc. Suppl. v. 99 = Fimbristylis caespitosa.
cespitosus, Pollich, Hist. Pl. Palat. i. 39 = pauciflorus.
Chaetà, Schult. Mant. ii. 72 = Eleocharis acicularis.
chaetarius, Spreng. Syst. i. 203; Thw. Enum. Pl. Zeyl. 435 = Eleocharis Chaetaria.
chaetodes, Link, in Buch, Beschr. Canar. Ins. 138 = cernuus.
Chamissoi, Schrad. ex Nees, in Mart. Fl. Bras. ii. i. 107 = riparius.
chilensis, Meyen, Reise, i. 380, nomen; et Nees & Meyen, in Linnaea, ix. (1834) 293 = pungens.
chinensis, *Munro, in Seem. Bot. Voy. Herald*, 423.—Ind. or.; China.
chinensis, Osb. It. 220 = squarrosus.
Chirigota, Wright, in Sauv. Fl. Cub. 176 (Fimbristylidis sp.).—Cuba.
ciliaris, Linn. Mant. ii. 182 = Fuirena glomerata.
ciliaris, Pers. Syn. i. 77 = Fuirena pubescens.
ciliatifolius, Ell. Sketch, i. 82; Darlingt. Florul. Cestr. 7 (Fimbristylidis sp.).—Am. bor.
ciliatus, Steud. Syn. Pl. Cyp. 86 = lacustris.
ciliolatus, Boeck. in Kjoeb. Vidensk. Meddel. (1871) 152 = Fimbristylidis sp.).—Guian.
cinereofuscus, Willd. ex Kunth, Enum. Pl. ii. 236, 237 = Fimbristylis ferruginea.
cinnamometorum, Vahl, Enum. ii. 278 = Fimbristylis cyperoides.
cinnamomeus, Boeck. in Engl. Bot. Jahrb. v. (1884) 505 (Fimbristylidis sp.).—Madag.
circinatus, Boeck. in Linnaea, xxxvi. (1869–70) 762 (Fimbristylidis sp.).—Am. trop.
clathratus, Reichb. Ic. Fl. Germ. viii. 39, in obs.—setaceus?
Clintonii, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxviii. (1864) 290.—Am. bor.
coarctatus, Ell. Sketch, i. 83 = Fimbristylis coarctata.
cognatus, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 228 = mucronatus.
coleotrichus, Boeck. in Linnaea, xxxvi. (1869–70) 763 (Fimbristylidis sp.).—Afr. trop.
collinus, Boeck. l. c. 746 = Fimbristylis contexta.
comosus, *Dum. Fl. Belg.* 143.—Belgium.
comosus, Wall. in Roxb. Fl. Ind. ed. Carey, i. 234 = Eriophorum comosum.
compactus, Hoffm. Deutschl. Fl. ed. II. i. 25 = maritimus.
compactus, Poir. Encyc. Suppl. v. 102; Spreng. Syst. i. 203 = Eleocharis compacta.
complanatus, Retz. Obs. v. 14 = Fimbristylis complanata.
compressus, Moench, Meth. 349 = Eleocharis ovata.
compressus, Pers. Syn. i. 66 = S. Caricis.
confervoides, Poir. Encyc. vi. 755.—Madag.
congestus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 41 = Eleocharis congesta.
congestus, Reichb. ex Kunth, Enum. Pl. ii. 211 (= *Isolepis junciformis*).—Am. austr.
conglomeratus, H. B. & K. Nov. Gen. et Sp. i. 227 = pungens.
conicus, Presl, ex Mert. & Koch, Deutschl. Fl. i. 427, in syn. = Eleocharis palustris.
coniferus, Poir. Encyc. vi. 756 = Lepironia mucronata.
conostachys, Boeck. in Linnaea, xxxvi. (1869–70) 756 (Fimbristylidis sp.).—Bras.
consanguineus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 188; et in Linnaea, xxxvi. (1869–70) 754 (Fimbristylidis sp.).—Bras.; Reg. Argent.
conspersus, Boeck. in Linnaea, xxxvi. (1869–70) 505 = inundatus.
constrictus, Schrad. ex Kunth, Enum. Pl. ii. 153 = Eleocharis geniculata.
cornatus, Wright, in Sauv. Fl. Cub. 177 = Fimbristylis squarrosa.
coronarius, Vahl, Enum. ii. 261 = Cyperus coronarius.

SCIRPUS:—
corymbiferus, Wright, in Sauv. Fl. Cub. 176 = Rynchospora ignorata.
corymbosus, Forsk. Fl. Aegypt. Arab. 14 = maritimus.
corymbosus, *Heyne, in Roth, Nov. Pl. Sp.* 28.—Afr.; Ind. or.
corymbosus, Linn. Cent. Pl. ii. 7; Amoen. Acad. iv. 303 = Rynchospora aurea.
costatus, *Boeck. in Linnaea*, xxxvi. (1869–70) 511.—Tasman.; Abyss.
crassiusculus, *Benth. Fl. Austral.* vii. 326.—Austral.
crinalis, Griseb. in Goett. Abh. xix. (1874) 265 (Eleocharidis sp.).—Reg. Argent.
criniger, *A. Gray, in Proc. Am. Acad.* vii. (1868) 392.—Calif.
crinitus, Poir. Encyc. vi. 752 = Ficinia filiformis.
cubensis, *Poepp. & Kunth, in Enum. Pl.* ii. 172.—Am. trop.; Afr. trop.
Cumingii, Boeck. in Linnaea, xxxvi. (1869–70) 491 = Carex setifolia.
cuspidatus, Roem. & Schult. Syst. ii. 67, sphalm., in citat. (Tetrariae sp.).—Afr. austr.
cuspidatus, Roth, Nov. Pl. Sp. 31 = Fuirena Wallichiana.
cuspidatus, Willd. ex Kunth, Enum. Pl. ii. 213 (= *Isolepis puberula*).—Madag.
Custoris, Hegetschw. Fl. Schw. 49 = lacustris.
cylindraceus, Willd. ex Kunth, Enum. Pl. ii. 238 = Fimbristylis spadicea.
cymosus, Lam. Illustr. i. 141 = Fimbristylis rigida.
cymosus, Poir. Encyc. Suppl. v. 99 = Fimbristylis cymosa.
cymosus, Spreng. Syst. i. 209 = Rynchospora distans.
cyperiformis, Muhl. Desc. Gram. 41 = Mariscus glomeratus.
cyperinus, Kunth, Enum. Pl. ii. 170 = S. Eriophorum.
cyperoides, *Hemsl. Biol. Centr. Am. Bot.* iii. 461.—Mexic.
cyperoides, Lam. Fl. Fr. iii. 553 = maritimus.
cyperoides, Linn. Mant. ii. 181 = Mariscus Sieberianus.
cyperoides, *Spreng. Syst.* i. 208.—Austral.
debilis, Lam. Illustr. i. 141 = Fimbristylis ferruginea.
debilis, Pursh, Fl. Am. Sept. i. 55 = erectus.
decipiens, A. St. Hil. ex Nees, in Mart. Fl. Bras. ii. 1. 208 = riparius.
decumanus, Willd. ex Kunth, Enum. Pl. ii. 168 = maritimus.
densus, Wall. in Roxb. Fl. Ind. ed. Carey, 231 (Fimbristylidis sp.).—Ind. or.
denudata, Poir. Encyc. Suppl. v. 99 = Fimbristylis denudata.
depauperata, Poir. l. c. 98; Muhl. ex Kunth, Enum. Pl. ii. 233 = Fimbristylis diphylla.
depressus, Vell. Fl. Flum. 35; i. t. 83 = Eleocharis geniculata.
deserticola, *Phil. Fl. Atac.* 53.—Chili.
Desprauxii, Steud. Syn. Pl. Cyp. 86 = littoralis.
diabolicus, *Steud. in Flora*, xii. (1829) 147.—Afr. austr.
dichotomus, Bieb. Fl. Taur. Cauc. i. 39 = Fimbristylis diphylla.
dichotomus, Linn. Sp. Pl. 50 = Fimbristylis dichotoma.
dichotomus, G. F. W. Mey. Prim. Fl. Esseq. 43 = Fimbristylis spadicea.
dichotomus, Willd. ex Kunth, Enum. Pl. ii. 235, 239 = Fimbristylis diphylla, junciformis.
dictyospermus, Wright, in Sauv. Fl. Cub. 175 = Eleocharis cellulosa.
Dietrichiae, *Boeck. in Flora*, lviii. (1875) 109.—Austral.
digitatus, Boeck. in Linnaea, xxxvi. (1869–70) 691 = flaccifolius.
dioëcus, *Boeck. l. c.* 719.—Afr. austr.
diphyllus, Retz. Obs. v. 15 = Fimbristylis diphylla.
dipsaceus, Rottb. Desc. Nov. Pl. 56 = Fimbristylis dipsacea.
dipsaceus, Vahl, ex Kunth, Enum. Pl. ii. 204 = S. Hystrix.
dissitus, Duthie, in Saharunpore Report (1885) = S. Caricis.
distichus, Peterm. in Flora, xxvii. (1844) 343 = S. Caricis.
divaricatus, *Ell. Sketch*, i. 88. t. 2. f. 4.—Am. bor.
domingensis, *Ham. Prod. Pl. Ind. Occ.* 14.—Ins. S. Doming.
domingensis, Pers. Syn. i. 67 = Fimbristylis spadicea.
Donianus, Spreng. Syst. iv. Cur. Post. 28 = quadrangulus.

SCIRPUS:—
Drummondii, Steud. Syn. Pl. Cyp. 318 = lineatus.
dubius, Roxb. Hort. Beng. [81]; Fl. Ind. i. 215 = Eleocharis plantaginea?
Duvalii, Hoppe, in Sturm, Deutschl. Fl. Heft 36 = lacustris.
ebenocarpus, T. Kirk, in Trans. N. Z. Inst. xvii. (1885) 224 = antarcticus.
echinatus, Linn. Sp. Pl. 50 = Mariscus paniceus.
echinatus, Muhl. Desc. Gram. 39 = Mariscus ovularis.
Eckloneus, Steud. in Flora, xii. (1829) 148 = Ficinia Ecklonea.
effusus, *Gueldenst. Reisen*, ii. 179; *ex Ledeb. Fl. Ross.* iv. 252, nomen (Quid?).—Ross. austr.
Ehrenbergii, Boeck. in Linnaea, xxxvi. (1869–70) 712 = maritimus.
elatus, Boeck. l. c. 694 = S. Holoschoenus?
elatus, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 252 (Quid?).—Ross. austr.
elatus, Poir. Encyc. Suppl. v. 99 = Fimbristylis denudata.
elegans, H. B. & K. Nov. Gen. et Sp. i. 226 = Eleocharis geniculata.
elegans, Salisb. Prod. 31 = Cyperus Luzulae.
elegantulus, Steud. Syn. Pl. Cyp. 317 = Eleocharis ovata.
Elliottii, Spreng. Syst. iv. Cur. Post. 28 = Fimbristylis diphylla.
ellipticus, Willd. ex Kunth, Enum. Pl. ii. 146 = Eleocharis tenuis, montana.
ellychniarius, *Molina, Sagg. Chil.* 153 (Quid?).—Chili.
elongatus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 40 = Eriophorum comosum.
elongatus, Poir. Encyc. Suppl. v. 99 = Fimbristylis diphylla.
equisetoides, Ell. Sketch, i. 79 = Eleocharis interstincta.
erectus, *Poir. Encyc.* vi. 761.—As. temp.; Am. bor.; Austral.
Eriophorum, *Michx. Fl. Bor. Am.* i. 33.—As. or.; Am. bor.
eriophorus, Poepp. ex Presl, in Oken, Isis, xxi. (1828) 269 = stramineus.
erraticus, Rota, ex De Not. in Ann. Sc. Nat. Sér. III. v. (1846) 366 = Eleocharis atropurpurea.
exaltatus, Pursh, Fl. Am. Sept. i. 56 = polyphyllus, *Vahl.*
excelsus, Salisb. Prod. 31 = Cladium germanicum.
exiguus, H. B. & K. Nov. Gen. et Sp. i. 225 = Eleocharis acicularis.
exilis, Poir. Encyc. Suppl. v. 105; Willd. ex Kunth, Enum. Pl. ii. 228 = Fimbristylis exilis.
expallescens, *Boeck. in Linnaea*, xxxvi. (1869–70) 509.—Afr. austr.
falcatus, Vahl, Enum. ii. 275 = Fimbristylis junciformis.
fasciculatus, Willd. ex Kunth, Enum. Pl. ii. 188 = Ludwigii.
fastigiatus, Thunb. Prod. Pl. Cap. 18 = Ficinia fastigiata.
ferrugineus, Linn. Sp. Pl. 50 = Fimbristylis ferruginea.
festucoides, Poir. Encyc. vi. 752 (= *Isolepis festucoides*).—Madag.
ficinioides, *Kunth, Enum. Pl.* ii. 172.—Afr. austr.
filamentosus, Vahl, Enum. ii. 262 (= *Isolepis filamentosa*).—Afr. trop.
filiculmis, Schrad. ex Griseb. in Goett. Abh. xxiv. (1879) 311 = Eleocharis sulcata
filiformis, Burm. f. Fl. Cap. Prod. 3 (Ficiniae sp.).—Afr. austr.
filiformis, Lam. Illustr. i. 138 = Eleocharis tenuis.
filiformis, Savi, Fl. Pisan. i. 46 = cernuus.
fimbriatus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 138 (Fimbristylidis sp.).—Bras.
fimbriatus, Poir. Encyc. vi. 779 = Fimbristylis diphylla.
fimbriatus, Willd. ex Kunth, Enum. Pl. ii. 235 = Fimbristylis pilosa.
fimbrisetus, Delile, Fl. Egypte, 155 = littoralis.
fistulosus, Forsk. Fl. Aegypt. Arab. 14 = articulatus.
fistulosus, Poir. Encyc. vi. 749 = Eleocharis fistulosa.
flaccidus, Reichb. ex Spreng. f. Tent. Suppl. Syst. 3 = Eleocharis capitata.
flaccifolius, *Steud. Syn. Pl. Cyp.* 83.—Afr. austr.
flavescens, Gromow, ex Trantv. in Act. Hort. Petrop. ix. (1884) 293, nomen (Quid?).—Hab.?
flavescens, Poir. Encyc. vi. 756 = Eleocharis capitata.

SCIRPUS:—
floccosus, Griseb. Cat. Pl. Cub. 241. (Fimbristylidis sp.).—Cuba.
floridanus, Michx. ex Kunth, Enum. Pl. ii. 158 = nanus.
fluitans, *Linn. Sp. Pl.* 48.—Geront.
fluviatilis, A. Gray, Man. Bot. N. U. St. ed. II. 500 = maritimus.
foliatus, Hook. f. in Hook. Lond. Journ. Bot. iii (1844) 414 = Schoenus axillaris.
fragrans, *Ruiz & Pav. Fl. Per.* i. 47.—Peruv.
frondosus, *Banks & Soland. ex Boeck. in Flora*, lxi. (1878) 141.—N. Zel.
furva, Poir. Encyc. Suppl. v. 99 = Fimbristylis furva.
fuscescens, Link, Jahrb. i. III. (1820) 81 = Fimbristylis dichotoma.
fuscus, Willd. ex Kunth, Enum. Pl. ii. 212 (= *Isolepis asperula*).—Am. trop.
Gaudichaudianus, Kunth, Enum. Pl. ii. 157 = Eleocharis ocreata.
Gaudichaudii, Boeck. in Linnaea, xxxvi. (1869–70) 511 = inundatus.
geniculatus, Linn. Sp. Pl. 48 = Eleocharis geniculata.
giganteus, *Kunth, Enum. Pl.* ii. 172.—Bras.
giganteus, *Noronha, ex Thou. Prod. Phyt.* 2, *in Mélang.* (1811), nomen.—Madag.
Giraudyi, Boeck. in Linnaea, xxxvi. (1869–70) 483 = fluitans.
glaucescens, Willd. Enum. Hort. Berol. 76 = Eleocharis glaucescens.
glaucophyllus, Boeck. in Kjoeb. Vidensk. Meddel. (1871) 153. (Fimbristylidis sp.).—Bras.
glaucus, Lam. Illustr. i. 142 = maritimus.
glaucus, Nees & Meyen, in Linnaea, ix. (1834) 293 = asper.
glaucus, Smith, Engl. Bot. t. 2321 = lacustris.
glaucus, Torr. Fl. U. St. 44 = Eleocharis glaucescens.
Glaziovii, Boeck. in Kjoeb. Vidensk. Meddel. (1871) 152 (Fimbristylidis sp.).—Bras.
globifer, Welw. ex Steud. Syn. Pl. Cyp. 87 = lacustris.
globiferus, Linn. f. Suppl. 104 = S. Holoschoenus.
globosus, Spreng. Syst. iv. Cur. Post. 27 = nodosus.
globulosus, Retz. Obs. vi. 19 = Fimbristylis globulosa.
globulosus, Roxb. Hort. Beng. [81]; Fl. Ind. 217 = Fimbristylis ferruginea.
glomeratus, Heyne, ex Wall. Cat. n. 3460 = Cyperus arenarius.
glomeratus, Linn. Sp. Pl. 52 = Mariscus Dregeanus.
glomeratus, Retz. Obs. iv. II. = Fimbristylis spathacea.
glomeratus, Roxb. Fl. Ind. i. 224 = Fimbristylis diphylla.
glomeratus, Scop. Fl. Carn. ed. II. i. 47 = mucronatus.
glutinosus, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 270 = nodosus.
Gmelini, [Honck.] Verz. Aller Gew. Teutschl. 138 (Quid?).—Europ.
gracilis, Koch, in Salzm. Enum. Pl. Rar. Gall. 9 = cernuus.
gracilis, Kunth, Enum. Pl. ii. 585 = Eleocharis maculosa.
gracilis, Poir. Encyc. vi. 763 = Ficinia Poiretii.
gracilis, Rudge, in Trans. Linn. Soc. x. (1811) 289. t. 15. f. 1 = nodosus.
gracilis, Willd. ex Kunth, Enum. Pl. ii. 227 = Fimbristylis tenuis.
gracilis, Zeyh. ex Kunth, l. c. 256 = Ficinia gracilis.
gracillimus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 140; et in Linnaea, xxxvi. (1869–70) 761 (Fimbristylidis sp.).—Abyss.
gracillimus, Kohts, in Oestr. Bot. Zeitschr. xix. (1869) 333 = setaceus.
gramineus, Neck. Delic. Gallo-Belg. 27 = sylvaticus.
Griffithii, Boeck. in Flora, xli. (1858) 595 = grossus.
grossus, *Linn. f. Suppl.* 104.—Ind. or.; Malaya.
Gunnii, *Boeck. in Linnaea*, xxxvi. (1869–70) 493.—Tasman.
hakonensis, Franch. & Sav. Enum. Pl. Jap. ii. 110 = Eleocharis tetraquetra.
Halleri, Vill. Hist. Pl. Dauph. ii. 188 = pauciflorus.
Halleri, Vitm. Summa Pl. i. 150 = supinus.
Hallii, A. Gray, Man. Bot. N. U. St. ed. V. 563 = supinus.
hamulosus, *Stev. in Mém. Soc. Nat. Mosc.* v. (1814) 356.—Reg. Casp.
hemisphaericus, Roth, Nov. Pl. Sp. 29 = Lipocarpha sphacelata.
heterocarpus, S. Wats. in Proc. Am. Acad. xviii. (1883) 171 = supinus.

SCIRPUS:—
heteromorphus, Steud. ex Phil. f. Cat. Pl. Vasc. Chil. 311 = Eleocharis nodulosa.
hexatrichus, Ehrnb. ex Boiss. Fl. Orient. v. 387 = Eleocharis macrantha.
Hildebrandtii, Boeck. in Flora, lviii. (1875) 263 (Fimbristylidis sp.).—Afr. trop.
hirsutus, Willd. ex Kunth, Enum. Pl. ii. 210 = Fimbristylis vestita.
hirtellus, Vahl, Eclog. Am. iii. 1 = Fimbristylis hirtella.
hirtus, Griseb. Cat. Pl. Cub. 241 (Fimbristylidis sp.).—Cuba.
hirtus, Poir. Encyc. Suppl. v. 106; Willd. ex Kunth, Enum. Pl. ii. 224 = Fimbristylis squarrosa.
hispanicus, Gueldenst. Reisen, i. 109; ex Ledeb. Fl. Ross. iv. 252 (Quid?).—Ross. austr.
hispidulus, *Vahl, Enum.* ii. 276.—Afr. trop.
Hochstetteri, Boeck. in Linnaea, xxxvi. (1869–70) 739 (Fimbristylidis sp.).—Abyss.
Holoschoenus, *Linn. Sp. Pl.* 49.—Geront.
Holoschoenus, Oed. Fl. Dan. t. 454 = lacustris.
Holoschoenus, Thunb. Fl. Cap. i. 368 = Thunbergii.
Hookeri, *Kunth, Enum. Pl.* ii. 173.—Chili.
Hoppii, Weihe, in Flora, xi. (1828) 118 = triqueter.
hottentottus, Linn. Mant. ii. 82 = Fuirena hirta.
Humboldtii, A. Dietr. Sp. Pl. ii. 112 = Hemicarpha subsquarrosa.
Humboldtii, Spreng. Syst. i. 213 (= *Isolepis junciformis*).—Am. austr.
humilis, *Vell. Fl. Flum.* 35; i. t. 84.—Bras.
humilis, Wallr. Sched. Crit. 27 = nanus.
humilis, Willd. ex Kunth, Enum. Pl. ii. 212 (= *Isolepis scabra*).—Peruv.
humillimus, *Benth. Fl. Austral.* vii. 324.—Austral.
hyalinolepis, Steud. ex Jardin, in Bull. Soc. Linn. Normand. Sér. II. ix. (1875) 278, 280 = maritimus.
Hystrix, *Thunb. Prod. Pl. Cap.* 17.—Afr. austr.
inanis, Steud. Syn. Pl. Cyp. 86 = spathaceus.
inclinatus, Aschers. & Schweinf. ex Boiss. Fl. Orient. v. 381 = corymbosus.
incomtulus, *Boeck. in Linnaea*, xxxvi. (1869–70) 693.—Afr. austr.
incurvatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 214 = articulatus.
intermedius, Čelak. in Arch. Naturw. Landesd. Boehm. iv. (1881) 742 = sylvaticus.
intermedius, A. Gray, ex Torr. in Ann. Lyc. N. York, iii. (1836) 300 = Eleocharis olivacea.
intermedius, Muhl. Desc. Gram. 31 = Eleocharis intermedia.
intermedius, Poir. Encyc. vi. 767 = S. Holoschoenus.
intermedius, Thuill. Fl. Par. ed. II. 21 = Eleocharis palustris.
interstinctus, Poepp. ex Presl, in Oken, Isis, xxi. (1828) 269 = Eleocharis geniculata.
interstinctus, Vahl, Enum. ii. 149 = Eleocharis interstincta.
intricatus, Linn. Mant. ii. 182 = Cyperus aristatus.
inundatus, *Poir. Encyc. Suppl.* v. 103; *Spreng. Syst.* i. 207.—Austral.; Am.; Austr.
iridifolius, Bory, Voy. ii. 94; Poir. Encyc. vi. 783 = Cladium iridifolium.
Isolepis, *Boeck. in Linnae*, xxxvi. (1869–70) 498 — Ind. or.; Afr. austr.
Janii, Bess. ex Schult. Mant. ii. 536 = lacustris.
japonicus, Franch. & Sav. Enum. Pl. Jap. ii. 109 = Eleocharis affata.
Jardinei, Steud. & Jardin, in Bull. Soc. Linn. Normand. Sér. II. ix. (1875) 278, 280 = lacustris.
javanus, Nees, in Wight, Contrib. 112 = mucronatus.
junceus, Forst. f. Prod. 6 = Fimbristylis juncea.
junciformis, Nees, in Wight, Contrib. 112 = erectus.
junciformis, Poir. Encyc. Suppl. v. 105 (= *Isolepis junciformis*.).—Am. austr.
junciformis, *Retz. Obs.* vi. 19.—China.
juncoides, Roxb. Hort. Beng. [81]; Fl. Ind. i. 216 = erectus.
juncoides, Willd. ex Kunth, Enum. Pl. ii. 211 (= *Isolepis junciformis*.).—Am. austr.
Junghuhnii, *Miq. Fl. Ind. Bat.* iii. 307.—Java.
Kalli, Forsk. Fl. Aegypt. Arab. 15 = Cyperus aegypticus.
kamtschaticus, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 198 = Eleocharis palustris.
Kielmeyeri, Schimp. ex Steud. Nom. ed. II. ii. 539 = Fimbristylis complanata.

SCIRPUS:—
Koshewnikowii, *Litwinow, ex Kotsom, in Bull. Soc. Nat. Mosc.* lvii. (1882) 220, nomen.—Ross.
Kunthii, Boeck. in Linnaea, xxxvi. (1869–70) 508 = ficinioides.
kyllingioides, *Boeck. l. c.* 733.—Ind. or.; Afr. trop.
Kysoor, Roxb. Hort. Beng. 6; Fl. Ind. i. 230 = grossus.
laciniatus, Nees & Ehrenb. ex Boeck. in Linnaea, xxxvi. (1869–70) 723 = maritimus.
laciniatus, Thunb. Prod. Pl. Cap. 17 = Ficinia laciniata.
lacustris, *Linn. Sp. Pl.* 48.—Cosmop.
laetevirens, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 103 = Eleocharis atropurpurea.
lanatus, Poir. Encyc. Suppl. v. 104 (= *Isolepis lanata*).—Am. austr.
laniferus, Boeck. in Linnaea, xxxvi. (1869–70) 768 (Bulbostylidis sp).—Afr. trop.
lappaceus, Lam. Illustr. i. 139 = Cyperus aristatus.
lateralis, Forsk. Fl. Aegypt. Arab. 15 = supinus.
lateralis, Retz. Obs. iv. 11 = erectus?
lateriflorus, J. F. Gmel. Syst. 127 = praec.?
latifolius, Bertol. in Bologn. Opusc. Sc. iii. (1819) 406 = Pleurostachys tenuiflora.
latifolius, Gilib. Exercit. ii. 515 = sylvaticus.
lavarum, Poir. Encyc. vi. 783 = Cladium iridifolium.
Lawianus, Boeck. in Linnaea, xxxvi. (1869–70) 497 = Fimbristylis digitata.
laxifolius, Thw. Enum. Pl. Zeyl. 435 = Eleocharis variegata.
laxus, Poir. Encyc. Suppl. v. 101 = Fimbristylis diphylla.
Lejeunei, Weihe, in Flora, xi. (1828) 116 = triqueter.
lemburgensis, Schult. Mant. ii. 85, sphalm. = *limburgensis*.
lenticularis, *Poir. Encyc. Suppl.* v. 103; *Spreng. Syst.* i. 208.—Austral.; N. Zel.
lenticularis, Torr. in Ann. Lyc. N. York, iii. (1836) 328 = sylvaticus.
leptaleus, Koch, in Syll. Ratisb. i. (1824) 8 = cernuus.
leptocarpus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 109 = Lipocarpha microcephala.
leptolepis, *Chapm. Fl. S. U. St.* 520.—Am. bor.
leptopus, Boeck. in Flora, xli. (1858) 414 = asper.
leptos, Wright, in Sauv. Fl. Cuba, 176 = nanus.
leptostachyus, *Boeck. in Linnaea*, xxxvi. (1869–70) 485.—Afr. austr.
Lereschii, Thom. ex Koch, Syn. Fl. Germ. ed. II. 853 = Eleocharis atropurpurea.
leucostachyus, Poir. Encyc. Suppl. v. 104 (= *Isolepis leucostachya*).—Am. trop.
Lichtensteinianus, Boeck. in Linnaea, xxxvi. (1869–70) 745 = Fimbristylis Lichtensteiniana.
limburgensis, Lej. Fl. Spa, ii. 289 = Eleocharis palustris.
limosus, Schrad. in Goett. Gel. Anz. (1821) iii. 2069 = Eleocharis limosa.
lineatus, *Michx. Fl. Bor. Am.* i. 32.—Am. bor.
lineatus, Muhl. Desc. Gram. 45 = divaricatus.
lineolatus, Franch. & Sav. Enum. Pl. Jap. ii. 545 = erectus.
lithospermus, Linn. Sp. Pl. 51 = Scleria lithosperma.
lithospermus, *Noronha, ex Thou. Prod. Phyt.* 2, *in Mélang.* (1811), nomen.—Madag.
lithuanicus, Bess. ex Schult. Mant. ii. 535 = lacustris.
littoralis, Fluegge, ex Reichb. Fl. Germ. Excurs. 74 = rufus.
littoralis, *Schrad. Fl. Germ.* i. 142. t. 5. f. 7.—Europ.; Asia; Aegypt.
longebracteatus, Salzm. ex Reichb. in Flora, xiii. (1830) 497 = S. Holoschoenus.
longifolius, Hook. & Arn. Bot. Beech. Voy. 49 = pungens.
longifolius, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Hypolytrum longifolium.
Lorentzii, Boeck. in Linnaea, xxxviii. (1874) 378 (Bulbostylidis sp.).—Reg. Argent.
lucidus, Less. ex Kunth, Enum. Pl. ii. 168 = maritimus.
Ludwigii, *Boeck. in Linnaea*, xxxvi. (1869–70) 486.—Afr. austr.
lupulinus, Spreng. Fl. Hal. Mant. 30 = Mariscus glomeratus.
luteus, Gilib. Exercit. ii. 513 = caespitosus.
luzonensis, J. & C. Presl, Rel. Haenk. i. 193 = erectus.
Luzulae, Linn. Syst. ed. X. 868 = Cyperus Luzulae.

SCIRPUS:—
luzulinus, Reichb. ex Kunth, Enum. Pl. ii. 211 (= *Isolepis juncifformis*).—Am. trop.
Lyallii, Baker, in Journ. Linn. Soc. xx. (1883) 297 = corymbosus.
macer, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 503.—Madag.
macranthus, Boeck. in Flora, xli. (1858) 415 = leptolepis.
macrophyllus, Bess. ex Schult. Mant. ii. 535, in obs. = lacustris.
macrostachyos, Lam. Illustr. i. 142 = maritimus.
macrostachys, Willd. Enum. Hort. Berol. 78 = praec.
macrostylis, Spreng. Syst. i. 208 = Rynchospora pterocarpa.
maculosus, Vahl, Enum. ii. 247 = Eleocharis maculosa.
madagascariensis, Boeck. in Bremen, Abh. vii. (1880) 37 = supinus.
margaritiferus, Boeck. in Linnaea, xxxvi. (1869–70) 697 = Schoenus apogon.
marginatus, Muhl. Desc. Gram. 28 = Eleocharis mutata.
marginatus, Thunb. Prod. Pl. Cap. 17 = Ficinia filiformis.
maritimus, *Linn. Sp. Pl.* 51.—Cosmop.
Martii, Dufour, ex Roem. & Schult. Syst. ii. 117 = Cladium germanicum.
mauritanicus, Steud. Syn. Pl. Cyp. 84 = mucronatus.
maximus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 231 = grossus.
medius, S. F. Gray, Nat. Arr. Brit. Pl. ii. 75 = lacustris.
medius, Roxb. Fl. Ind. ed. Carey, i. 216 = Eleocharis fistulosa.
megapotamicus, Spreng. f. Tent. Suppl. Syst. 4 = Rynchospora luzuliformis.
megastachyus, Steud. Syn. Pl. Cyp. 87 = maritimus.
melanocarpus, Baldw. ex Torr. in Ann. Lyc. N. York, iii. (1836) 311 = Eleocharis melanocarpa.
melanocarpus, Griseb. Cat. Pl. Cub. 239 = Eleocharis pachystyla.
melanocephalus, Griseb. in Goett. Abh. xxiv. (1879) 311 = Eleocharis albibracteata.
melanorrhizus, Boeck. in Engl. Bot. Jahrb. v. (1884) 504 = pungens.
melanospermus, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 199 = supinus.
melanospermus, Salzm. ex Steud. Syn. Pl. Cyp. 78 = Eleocharis ocreata.
melanostachys, Urv. in Mém. Soc. Linn. Par. iv. (1826) 600 = Eleocharis macrorhiza.
melas, Kunze, ex Kunth, Enum. Pl. ii. 169 = pungens.
membranaceus, Thunb. Prod. Pl. Cap. 17 = Ficinia membranacea.
Meyenii, Nees, in Lehm. Pl. Preiss. ii. 75 = lacustris.
Meyeri, Trautv. in Act. Hort. Petrop. v. (1877) 121, in textu = pumilus.
Michauxii, Pers. Syn. i. 68 = Fimbristylis autumnalis.
Michelianus, *Linn. Sp. Pl.* 52.—Europ.; Afr. bor.; Asia.
Michelianus, Sieber, ex Kunth, Enum. Pl. ii. 19 = Cyperus pygmaeus.
micranthus, Vahl, Enum. ii. 254.—Am. bor. et austr.
microcarpus, J. & C. Presl, Rel. Haenk. i. 195 = sylvaticus.
microlepis, Griseb. Cat. Pl. Cub. 239 = Eleocharis minima.
miliaceus, Linn. Syst. ed. X. 868 = Fimbristylis quadrangularis?
miliaceus, Roxb. Fl. Ind. ed. Carey, i. 231 = Fimbristylis diphylla.
miliaceus, Thunb. Fl. Jap. 37 = Fimbristylis ferruginea.
Minaae, Todaro, Rar. Pl. Sic. Decad. i; in Atti Acc. Sc. Palermo, N. S. i. (1845) 15 = cernuus.
minimus, Hook. & Arn. Bot. Beech. Voy. 49 = cernuus.
minimus, Pursh, Fl. Am. Sept. i. 55 = micranthus.
minimus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 219 = Fimbristylis dipsacea.
minimus, Sieber, ex Kunth, Enum. Pl. ii. 195 = antarcticus.
minimus, Vahl, Enum. ii. 253 = Fimbristylis barbata.
minimus, Willd. ex Kunth, Enum. Pl. ii. 268 = S. Isolepis.
mitratus, Franch. & Sav. Enum. Pl. Jap. ii. 544 = Eleocharis palustris?
mitratus, Griseb. Fl. Brit. W. Ind. 570 (Eleocharidis sp.).—Am. trop.

SCIRPUS:—
modestus, Phil. f. Cat. Pl. Vasc. Chil. 311 = cernuus.
mollis, Wall. in Roxb. Fl. Ind. ed. Carey, i. 227 = Rynchospora mollis.
monander, Rottb. Desc. Nov. Pl. 50 = Fimbristylis argentea.
monander, Roxb. Hort. Beng. [81]; Fl. Ind. i. 222 (= *Isolepis barbata*).—Geront. trop.
monander, Willd. Sp. Pl. i. 311 = Lipocarpha sphacelata.
monophyllus, J. & C. Presl, Rel. Haenk. i. 193 = pungens.
monostachyos, Koen. ex Roxb. Hort. Beng. 6; Fl. Ind. i. 219 = Fimbristylis schoenoides.
montanus, H. B. & K. Nov. Gen. et Sp. i. 226 = Eleocharis montana.
Moseleyanus, *Boeck. in Flora*, lviii. (1875) 262.—Ins. Nightingale.
mucronatus, Host, Gram. Austr. iii. 44 = triqueter.
mucronatus, *Linn. Sp. Pl.* 50 = Geront.
mucronatus, Pollich, Hist. Pl. Palat. i. 44 = maritimus.
mucronatus, Roth, Tent. Fl. Germ. i. 23 = pungens.
mucronatus, Roxb. Fl. Ind. ed. Carey, i. 219 = supinus.
mucronulatus, Michx. Fl. Bor. Am. i. 31 = Fimbristylis autumnalis.
Muehlenbergianus, Schult. Mant. ii. 74 = Eleocharis Muehlenbergianus.
Muehlenbergii, Spreng. Syst. i. 207 = Fimbristylis barbata.
multicaulis, C. C. Gmel. Fl. Bad. i. 96 = Eleocharis ovata.
multicaulis, Sm. Fl. Brit. i. 48 = Eleocharis palustris.
multicaulis, Vahl, ex Kunth, Enum. Pl. ii. 146 = Eleocharis tenuis, montana.
multicostatus, Baker, in Journ. Linn. Soc. xx. (1883) 298 = costatus.
muricatus, Poir. Encyc. vi. 765 = Rynchospora cyperoides.
muscosus, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 224, nomen.—N. Zel.
mutatus, Linn. Syst. ed. X. 867 = Eleocharis mutata.
muticus, D. Don, Prod. Fl. Nep. 41 = mucronatus.
mysurensis, Heyne, ex Wall. Cat. n. 3467 = mucronatus.
nanus, Phil. f. Cat. Pl. Vasc. Chil. 311 = inundatus.
nanus, Poir. Encyc. vi. 759 = Fimbristylis argentea.
nanus, *Spreng. Pugill.* i. 4.—Europ.; Aegypt.; Am. bor. et centr.
natans, Boj. Hort. Maurit. 383 = supinus.
natans, *Griseb. Cat. Pl. Cub.* 238.—Cuba.
natans, Spreng. ex Kunth, Enum. Pl. ii. 194 = rivularis.
natans, Thunb. Prod. Pl. Cap. 17 = tenuissimus.
Neesii, *Boeck. in Linnaea*, xxxvi. (1869–70) 504.—Afr. austr.
nervosus, *Boeck. l. c.* 484.—Abyss.
nevadensis, *S. Wats. in Bot. King's Exp.* 360.—Am. bor.
nigrescens, Salzm. ex Steud. Syn. Pl. Cyp. 91 = Eleocharis nigrescens.
nigricans, Poir. Encyc. Suppl. v. 104; Spreng. Syst. i. 212 = inundatus.
nigricans, Steud. Nom. ed. II. ii. 540 = Schoenus nigricans.
niloticus, Blanco, Fl. Filip. ed. I. 33 = Fimbristylis miliacea.
niloticus, J. F. Gmel. Syst. 126 = Fimbristylis dichotoma.
nindensis, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 27.—Afr. trop.
nitens, Boeck. in Linnaea, xxxvi. (1869–70) 696 = Schoenus nitens.
nitens, Vahl, Enum. ii. 272 = Rynchospora nitens.
nitidulus, Willd. ex Kunth, Enum. Pl. ii. 232 = Fimbristylis diphylla.
nobilis, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 159 = maritimus.
nodosus, *Rottb. Desc. Nov. Pl.* 52; *Progr.* 24.—Reg. austr. temp.
nodulosus, Roth, Nov. Pl. Sp. 29 = Eleocharis nodulosa.
nubigenus, Phil. in Anal. Univ. Chil. (1873) 554 = acaulis.
nudipes, Griseb. in Goett. Abh. xxiv. (1879) 312 = cernuus.
nudissimus, Steud. & Jardin, in Bull. Soc. Linn. Normand. Sér. II. ix. (1875) 278, 280 = Eleocharis palustris.

SCIRPUS:—
nudus, Rafin. Atl. Journ. 18 (Quid?).—Am. bor.
numidianus, Vahl, Enum. ii. 254 = cernuus.
nutans, Retz. Obs. iv. 12 = Fimbristylis nutans.
obtuse-trigonus, Salzm. ex Steud. Syn. Pl. Cyp. 80 = Eleocharis mutata.
obtusifolius, Heyne, ex Kunth, Enum. Pl. ii. 239 = Fimbristylis junciformis.
obtusifolius, Lam. Illustr. i. 141 = Fimbristylis obtusifolia.
obtusus, Willd. Enum. Hort. Berol. 76 = Eleocharis ovata.
ochreatus, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 535 = Eleocharis ochreata.
oliganthus, Steud. in Flora, xii. (1829) 146 = Ficinia filiformis.
oligostachyus, *Boeck. in Linnaea*, xxxvi. (1869–70) 743.—Abyss.
Oliveri, Boeck. in Flora, lviii. (1875) 260 = Thouarsianus.
Olneyi, *A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 238.—Am. bor.
Onaei, Franch. & Sav. Enum. Pl. Jap. ii. 544 = Eleocharis tetraquetra.
orgyalis, Rafin. Ann. Nat. 16 = lacustris.
ovatus, Gilib. Exercit. ii. 512 = Eleocharis palustris.
ovatus, Roth, Tent. Fl. Germ. ii. II. 562 = Eleocharis ovata.
pachystylus, Wright, in Sauv. Fl. Cub. 174 (Eleocharidis sp.).—Cuba.
pallescens, Boeck. ex Hemsl. Bot. Challenger, i. II. 158 = Thouarsianus.
pallescens, Roxb. Hort. Beng. [81]; Fl. Ind. i. 229 = Fimbristylis dichotoma.
pallescens, Willd. ex Kunth, Enum. Pl. ii. 237 = Fimbristylis ferruginea.
pallescens, Willd. ex Link, Hort. Berol. i. 292 = Fimbristylis diphylla.
palmaris, Willd. ex Kunth, Enum. Pl. ii. 150 = Eleocharis ochreata.
paludicola, *Kunth, l. c.* 163.—Afr. austr.
palustris, Linn. Sp. Pl. 47 = Eleocharis palustris.
panormitanus, Parl. Fl. Ital. ii. 99 = S. Holoschoenus.
paradoxus, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 136; *et in Linnaea*, xxxvi. (1869–70) 739.—Bras.
Parlatoris, Bivona f. Piante Ined. 2; et in Parl. Fl. Ital. ii. 96 = S. Holoschoenus.
parviflorus, Poir. Encyc. Suppl. v. 99 = Fimbristylis diphylla dichotoma.
parviflorus, Willd. ex Kunth, Enum. Pl. ii. 230 = Fimbristylis miliacea.
parvulus, Roem. & Schult. Syst. ii. 124 = nanus.
pauciflorus, *Lightf. Fl. Scot.* ii. 1078.—Europ.; Asia.
pauciflorus, Poir. Encyc. Suppl. v. 98 = Fimbristylis pauciflora.
pauciflorus, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid?).—Am. bor.
pectinatus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 218 = littoralis.
pendulus, Muhl. Desc. Gram. 44 = lineatus.
pentagonus, Roxb. Hort. Beng. 6; Fl. Ind. i. 218 = Fimbristylis quinquangularis.
perpusillus, *Boeck. in Linnaea*, xxxvi. (1869–70) 496.—Chili.
petasatus, Maxim. in Bull. Soc. Nat. Mosc. liv. (1879) I. 64 = Eleocharis tetraquetra.
phaeocarpus, *Phil. f. Cat. Pl. Vasc. Chil.* 312.—Chili.
Philippi, Tineo, Pl. Rar. Sicil. 36 = littoralis.
pictus, Boeck. in Linnaea, xxxvi. (1869–70) 503 = cernuus?
pileatus, Franch. & Sav. Enum. Pl. Jap. ii. 110 = Eleocharis palustris.
pilosus, Retz. Obs. vi. 19 = Fuirena glomerata.
pilosus, *Thunb. Prod. Pl. Cap.* 18.—Afr. austr.
planiculmis, F. Schmidt, Reisen Amurl. 190 = erectus.
planifolius, *Muhl. Desc. Gram.* 32.—Am. bor.
plantagineus, Retz. Obs. v. 14 = Eleocharis plantaginea.
plantagineus, Roxb. Hort. Beng. 6; Fl. Ind. i. 212 = Fimbristylis quinquangularis.
plantagineus, Sw. Fl. Ind. Occ. i. 123 = Eleocharis interstincta.
plantaginoides, Rottb. Desc. Nov. Pl. 45 = Eleocharis plantaginea.
plicarhachis, Griseb. Cat. Pl. Cub. 239 = Eleocharis variegata.

SCIRPUS:—
plumosus, R. Br. Prod. 223; Regel, in Act. Hort. Petrop. vii. (1880) 559 = littoralis.
pollicaris, Delile, Fl. Aeg. Illustr. 50 = nanus.
Pollichii, Gren. & Godr. Fl. Fr. iii. 374 = triqueter.
polycoleus, De Not. in Ann. Sc. Nat. Sér. III. ix. (1848) 326 = supinus.
polymorphus, Dulac, Fl. Hautes-Pyr. 60 = S. Holoschoenus.
polymorphus, Salzm. ex Steud. Syn. Pl. Cyp. 74 (Eleocharidis sp.).—Bras.
polyphyllus, *Vahl, Enum.* ii. 274.—Am. bor.
polystachyus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 108.—Austral.
polytrichoides, Retz. Obs. iv. 11 = Fimbristylis polytrichoides.
praelongatus, Poir. Encyc. vi. 764 = articulatus.
Preslii. A. Dietr. Sp. Pl. ii. 175 = mucronatus.
prolifer, *Rottb. Desc. Nov. Pl.* 55; *Progr.* 25.—Afr. austr.; Austral.
prolifer, Thou. Fl. Tristan d'Acugn. 36. t. 7 = Thouarsianus.
prolifero-ramosus, Boeck. in Flora, lviii. (1875) 261 = Thouarsianus.
propinqua, Poir. Encyc. Suppl. v. 99 = Fimbristylis squarrosa.
Pseudo-schoenus, Steud. in Flora, xii. (1829) 147 = Ficinia tristachya.
Pseudo-triqueter, Steud. Syn. Pl. Cyp. 86 = riparius.
pterigosperma, Poir. Encyc. Suppl. v. 98 = Fimbristylis pterigosperma.
Pterolepis, Kunth, Enum. Pl. ii. 166 = littoralis.
puberulus, Michx. Fl. Bor. Am. i. 31 = Fimbristylis diphylla.
puberulus, Poir. Encyc. vi. 767 (= *Isolepis puberula*).—Madag.
pubescens, Lam. Illustr. i. 139 = Fuirena pubescens.
pubescens, Pers. Syn. i. 68; Link, Jahrb. i. III. (1820) 80 = Fimbristylis diphylla.
pubigerus, Bernh. ex Link, Hort. Berol. i. 289 = Fimbristylis diphylla.
pulchellus, *Boeck. in Linnaea*, xxxvi. (1869–79) 698.—Afr. austr.
pulcher, Rudge, Pl. Gui. 18. t. 23 = Hypolytrum pungens.
Pumilio, Spreng. Syst. i. 204 = Eleocharis pusilla.
pumilus, *Vahl, Enum.* ii. 243.—Reg. bor.
pumilus, Willd. ex Kunth, Enum. Pl. ii. 191 = cernuus.
punctatus, Poir. Encyc. Suppl. v. 98 = Fimbristylis punctata.
punctatus, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 93 = Eleocharis nana.
pungens, Poir. Encyc. Suppl. v. 102 = Hypolytrum pungens.
pungens, *Vahl, Enum.* ii. 255.—Europ.; Am.
pungens, Willd. ex Kunth, Enum. Pl. ii. 164 = lacustris.
purpureo-ater, Boeck. in Linnaea, xxxviii. (1874) 379 (Fimbristylidis sp.).—Abyss.
pusillus, Herb. Lucae, ex Kunth, Enum. Pl. ii. 141 = Eleocharis acicularis.
pusillus, Poir. Encyc. Suppl. v. 103 = Eleocharis pusilla.
pusillus, Vahl, Enum. ii. 246 = nanus?
pusillus, Willd. ex Kunth, Enum. Pl. ii. 143 = Eleocharis intermedia.
pycnostachys, Salzm. ex Steud. Syn. Pl. Cyp. 129 = Lipocarpha Humboldtiana.
pygmaeus, A. Gray, in Proc. Am. Acad. vii. (1868) 392 = cernuus.
pygmaeus, Lam. Illustr. i. 139 = Eleocharis Chaetaria.
quadrangularis, Thou. ex Cat. Hort. Maurit. (1822) 4 = Fimbristylis miliacea?
quadrangulatus, Michx. Fl. Bor. Am. i. 30 = Eleocharis mutata.
quadrangulus, *D. Don, Prod. Fl. Nep.* 40 (sp. dub.).—Nepal.
quadriflorus, Boeck. in Linnaea, xxxvi. (1869–70) 761 (Fimbristylidis sp.).—Bras.
quinquangularis, Vahl, Enum. ii. 279 = Fimbristylis quinquangularis.
quinquefarius, *Buch.-Ham. ex Boeck. in Linnaea*, xxxvi. (1869–70) 701.—Ind. or.; Afghan.
quinqueflorus, Vitm. Summa Pl. i. 150 = pauciflorus.
racemosus, *Spreng. Syst.* i. 213.—Bras.
radiatus, Steud. Nom. ed. II. ii. 541, sphalm. = *radicans*, Poir.
radiatus, Thunb. Prod. Pl. Cap. 18 = Ficinia radiata.
radicans, Poir. Encyc. vi. 751 = Eleocharis acicularis.

SCIRPUS:—
radicans, *Schkuhr, in Usteri, Ann. Bot.* iv. (1793) 49. t. 1.—Europ.
ramosus, Boeck. in Linnaea, xxxviii. (1874) 377 = fluitans.
rarus, Poir. Encyc. Suppl. v. 99 = Fimbristylis rara.
reclinatus, Beyr. ex Kunth, Enum. Pl. ii. 143 = Eleocharis intermedia.
Rehmanni, Ridl. in Trans. Linn. Soc. Ser. II. ii. (1884) 159 = S. Hystrix.
Reichenbachii, Boeck. in Linnaea, xxxvi. (1869–70) 737 (Bulbostylidis sp.).—Am. trop.
remireoides, *Griseb. in Goett. Abh.* xix. (1874) 266.—Reg. Argent.
Renschii, Boeck. in Engl. Bot. Jahrb. v. (1884) 504 (Bulbostylidis sp.).—Madag.
repens, Willd. ex Schult. Mant. ii. 84, in obs. = Eleocharis capitata.
reptans, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Dichromena repens.
reptans, Thuill. Fl. Par. ed. II. 22 = Eleocharis palustris.
reticulatus, Lam. Illustr. i. 142 = polyphyllus?
retroflexus, Poir. Encyc. vi. 753 = Eleocharis Chaetaria.
retrofractus, Linn. Sp. Pl. 50 = Mariscus retrofractus.
retusus, Koen. ex Wall. Cat. n. 3428 = Rynchospora Wallichiana.
rigescens, Willd. ex Schult. Mant. ii. 84, in obs. = Fimbristylis spathacea.
rigidus, *Boeck. in Linnaea*, xxxvi. (1869–70) 492.—Peruv.
rigidus, Griseb. in Goett. Abh. xxiv. (1879) 312 (Fimbristylidis sp.).—Reg. Argent.
rigidus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 107 = riparius.
riparius, Pers. Syn. i. 68 = maritimus.
riparius, Poir. Encyc. Suppl. v. 103; Spreng. Syst. i. 208 = cernuus.
riparius, *J. & C. Presl, Rel. Haenk.* i. 193.—Am. austr.; Calif.
rivularis, *Boeck. in Linnaea*, xxxvi. (1869–70) 504.—Afr. austr.
rivularis, Don, ex Nees, l. c. ix. (1834) 288 = Fuirena umbellata.
robustus, Pursh, Fl. Am. Sept. i. 56 = maritimus.
romanus, Linn. Sp. Pl. 49 = S. Holoschoenus.
Rosellinii, Ces. Pass. & Gibelli, Compend. Fl. Ital. 99 = lacustris.
rostellatus, Torr. in Ann. Lyc. N. York, iii. (1836) 318 = Eleocharis rostellata.
Rothii, Hoppe, in Sturm, Deutschl. Fl. Heft 36 = pungens.
rufescens, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 138; et in Linnaea, xxxvi. (1869–70) 755 (Bulbostylidis sp.).—Bras.
rufus, *Schrad. Fl. Germ.* i. 133. t. 1. f. 3.—Europ.; Asia.
ruppioides, Thw. ex Sauv. Fl. Cub. 176, in obs. = submersus.
salbundius, Buch.-Ham. ex Wall. Cat. n. 3499 = Fimbristylis quinquangularis.
salinus, Schmidt, ex Steud. Nom. ed. II. ii. 541 = maritimus.
Savii, Sebast. & Mauri, Fl. Rom. Prod. 22 = cernuus.
scaber, Roxb. Hort. Beng. [81]; Fl. Ind. i. 220 = Fimbristylis acuminata.
scaberrimus, Boeck. in Flora, xli. (1858) 595 = grossus.
scariosus, Roem. & Schult. Syst. ii. 61, sphalm. = Ficinia bracteata.
Scheuchzeri, Vitm. Summa Pl. i. 150 = Eriophorum vaginatum.
Schimperianus, Boeck. in Linnaea, xxxvi. (1869–70) 750 (Bulbostylidis sp.).—Abyss.
schoenoides, Boeck. l. c. 745 = Fimbristylis Kunthiana.
schoenoides, Ell. Sketch, i. 89 = Rynchospora Elliotii.
schoenoides, Retz. Obs. v. 14 = Fimbristylis schoenoides.
schoenoides, Roxb. Hort. Beng. 6; Fl. Ind. i. 221 = Fimbristylis monostachya.
Schomburgkianus, Boeck. in Linnaea, xxxvi. (1869–70) 757 (Rynchosporae sp.).—Guian.
Schweinfurthianus, Boeck. l. c. 758 (Bulbostylidis sp.).—Afr. trop.
Sellovianus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 138; et l. c. 748 (Fimbristylidis sp.).—Bras.
Sellowianus, Griseb. in Goett. Abh. xxiv. (1879) 312 = Eleocharis Sellowiana.
semisubterraneus, *Boeck. in Linnaea*, xxxvi. (1869–70) 495.—Peruv.

SCIRPUS:—
senegalensis, Lam. Illustr. i. 140 = Lipocarpha argentea.
sepium, [Honck.] Verz. Aller Gew. Teutschl. i. 78 = pauciflorus.
sericeus, Poir. Encyc. Suppl. v. 99 = Fimbristylis sericea.
serrulatus, Vahl, Eclog. Am. iii. 1 = Fimbristylis diphylla.
sesquipollicaris, Willd. ex Kunth, Enum. Pl. ii. 203 = micranthus.
setaceus, Linn. Mant. ii. 321 = cernuus.
setaceus, *Linn. Sp. Pl.* 49, partim.—Austral.; Asia; Europ.
setaceus, Thunb. Prod. Pl Cap. 17 = antarcticus.
setifolius, Boeck. in Flora, lxii. (1879) 160 (Bulbostylidis sp.).—Paraguay.
simplex, Ell. Sketch, i. 76 = Eleocharis tortilis.
sinensis, Kunth, Enum. Pl. ii. 202, in syn. = squarrosus.
Smithii, A. Gray, Man. Bot. N. U. St. ed. V. 563 = erectus.
soloniensis, Dubois, Meth. ed. II. 249 = Eleocharis ovata.
spadiceus, *Boeck. in Linnaea*, xxxvi. (1869–70) 493.—Ind. occ. et Afr. trop.
spadiceus, Linn. Sp. Pl. 51 = Fimbristylis spadicea.
spadiceus, Phil. in Linnaea, xxix. (1857–58) 80 = Schoenus nitens.
Sparmanni, Lam. Illustr. i. 140 = Ficinia sylvatica.
spathaceus, *Hochst. in Flora*, xxviii. (1845) 759.—Afr. austr.
spathaceus, Michx. Fl. Bor. Am. i. 32 = Dulichium spathaceum.
speciosus, Salisb. Prod. 31 = Eriophorum angustifolium.
sphacelatus, Poir. Encyc. Suppl. v. 102; Spreng. Syst. i. 204 = Eleocharis sphacelata.
sphaerocarpus, Boeck. in Linnaea, xxxvi. (1869–70) 741 (Bulbostylidis sp.).—Afr. trop.
sphaerocephalus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 137; et in Linnaea, xxxvi. (1869–70) 748 (Bulbostylidis sp.).—Bras.
sphaerolepis, Boeck. ll. cc. 140; 766 (Bulbostylidis sp.).—Bras.
spiralis, Bosc. ex Kunth, Enum. Pl. ii. 145 = Eleocharis tortilis.
spiralis, Rottb. Desc. Nov. Pl. 42 = Eleocharis spiralis.
spiralis, Willd. ex Kunth, Enum. Pl. ii. 153 = Eleocharis plantaginea.
spiraloides, Roxb. Hort. Beng. [81], nomen = Eleocharis fistulosa.
squamosus, Link, Jahrb. i. III. (1820) 79 = squarrosus.
squarrosulus, Steud. Syn. Pl. Cyp. 88 = maritimus.
squarrosus, *Linn. Mant.* ii. 181.—Afr.; Asia.
squarrosus, Poir. Encyc. Suppl. v. 100 = Fimbristylis squarrosa.
squarrosus, Spreng. Syst. iv. Cur. Post. 28 = Thouarsianus.
stans, *Ham. Prod. Pl. Ind. Occ.* 14.—Ind. occ.
stenophyllus, Ell. Sketch, i. 83 (Bulbostylidis sp.).—Am. bor.
sterilis, Salzm. ex Steud. Syn. Pl. Cyp. 118 = Fimbristylis spadicea.
Steudneri, *Boeck. in Linnaea*, xxxvi. (1869–70) 733.—Abyss.
stolonifer, Roth, in Usteri, Ann. Bot. x. (1794) 36 = fluitans.
stramineus, *Presl, in Oken, Isis*, xxi. (1828) 269.—Am. bor.
striatulus, Griseb. in Goett. Abh. xix. (1874) 266 = Eleocharis bonariensis.
strictus, Poir. Encyc. Suppl. v. 99 = Fimbristylis diphylla.
strictus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 226 = Fimbristylis spathacea.
strobiliferus, Steud. Syn. Pl. Cyp. 317 = maritimus.
strobilinus, Roxb. Hort. Beng. 6; Fl. Ind. i. 219 = maritimus.
stygius, Ferber, ex Fleischer, Syst. Verz. Ostsee-Prov. Bek. Gew. Phan. (1830) (Lithogr.) 120 = Juncus stygius.
subaphyllus, Boeck. in Linnaea, xxxvi. (1869–70) 705 = subcapitatus.
subappendiculatus, Steud. Syn. Pl. Cyp. 84 = pungens.
subarticulatus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 215 = articulatus.
subcapitatus, *Thw. Enum. Pl. Zeyl.* 351.—Ind. or.; Sumatra; China.
subcucullatus, Berggren, Ny-Zeel. Fanerog. 12. t. 5. f. 16–20 = aucklandicus.
subdistichus, Boeck. in Linnaea, xxxvi. (1869–70) 490 = natans.

SCIRPUS:—
submersus, *Wright, in Sauv. Fl. Cub.* 175.—Am. trop.; Zeylan.
subprolifer, Boeck. in Linnaea, xxxvi. (1869–70) 492 = cernuus.
subquadriflorus, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 139 = *quadriflorus*.
subsquarrosus, Muhl. Desc. Gram. 39 = micranthus.
subterminalis, *Torr. Fl. U. St.* 47.—Am. bor.
subulatus, Vahl, Enum. ii. 268 = littoralis.
sulcatus, Ell. Sketch, i. 56 = Fimbristylis diphylla.
sulcatus, Roth, Nov. Pl. Sp. 30 = Eleocharis Rothiana.
sulcatus, *Thou. Fl. Tristan d'Acugn.* 36. t. 7.—Ins. Trist. da Cunha.
sulciculmis, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 103 = Eleocharis atropurpurea.
sundanus, Miq. Fl. Ind. Bat. iii. 304 = mucronatus.
supinus, Boeck. in Linnaea, xxxvi. (1869–70) 700, partim = erectus.
supinus, *Linn. Sp. Pl.* 49.—Reg. temp.
swampianus, Bosc, ex Kunth, Enum. Pl. ii. 168 = maritimus.
sylvaticus, *Linn. Sp. Pl.* 49.—Europ.; Am. bor.
sylvaticus, Vahl, Enum. ii. 271, partim = radicans.
Tabernaemontani, C. C. Gmel. Fl. Bad. i. 101 = lacustris.
Tabernaemontani, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 702 = littoralis.
Tatora, Kunth, Enum. Pl. ii. 166 = riparius.
tegetalis, Burch. Trav. i. 260, in nota = spathaceus.
tenellus, Link, Jahrb. i. III. (1820) 80 = Fimbristylis tenella.
tenellus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 224 = Fimbristylis tenera.
tener, Spreng. Syst. i. 204 = Eleocharis gracilis.
tenerrimus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 94 = Eleocharis minima.
tenuiculus, Schrad. ex Schult. Mant. ii. 74 = Eleocharis sulcata.
tenuifolius, DC. Fl. Fr. Suppl. 300 = pungens.
tenuifolius, Rudge, Pl. Gui. 18. t. 22 = Fimbristylis capillaris.
tenuis, Schreb. ex Schweigg. & Koerte, Fl. Erlang. 12 = Eleocharis palustris.
tenuis, *Spreng. in Flora*, xii. (1829) 1. *Beil.* 12.—Afr. austr.
tenuis, Willd. Enum. Hort. Berol. 76 = Eleocharis tenuis.
tenuispicatus, Boeck. in Linnaea, xxxvi. (1859–60) 740 (Bulbostylidis sp.).—Mexic.
tenuissimus, *Boeck. l. c.* 481.—Afr. austr.
tereticulmis, Steud. Syn. Pl. Cyp. 85 = riparius.
ternatanus, Reinw. ex Miq. Fl. Ind. Bat. iii. 307 = chinensis.
tetragonus, Poir. Encyc. Suppl. v. 98 = Fimbristylis tetragona.
tetragonus, Poir. Encyc. vi. 767 = Fimbristylis miliacea.
tetraqueter, Thw. Enum. Pl. Zeyl. 434 = Eleocharis tetraquetra.
thermalis, Spruce, ex Benth. & Hook. f. Gen. iii. 1051 = semisubterraneus.
Thouarsi, Roem. & Schult. Syst. ii. 134 (Bulbostylidis sp.).—Madag.?
Thouarsianus, *Schult. Mant.* ii. 84.—Ins. Trist. da Cunha.
Thuarsii, Spreng. Syst. iv. Cur. Post. 27 = sulcatus.
Thunbergii, Boeck. in Linnaea, xxxvi. (1869–70) 719 = S. Holoschoenus.
Thunbergii, *Spreng. f. Tent. Suppl. Syst.* 4.—Afr. austr.
Thwaitesii, Boeck. in Linnaea, xxxviii. (1874) 380 (Fimbristylidis sp.).—Ind. or.
thyrsiflorus, Willd. Enum. Hort. Berol. 78 = S. Eriophorum.
timorensis, Kunth, Enum. Pl. ii. 162 = erectus.
tomentosus, Poir. Encyc. Suppl. v. 101 = Fimbristylis diphylla.
Torreyi, *Olney, in Proc. Franklin Soc. Providence*, i. (1847) 32; *et ex A. Gray, Man. Bot. N. U. St.* ed. II. 499.—Am. bor.
tortilis, Willd. ex Kunth, Enum. Pl. ii. 144 = Eleocharis tortilis.
tranquebariensis, Roth, Catalect. iii. 6 = Fimbristylis ferruginea.
translucens, Le Gall, ex Lloyd, Fl. Loire-Inf. 275; et in Congr. Sc. Fr. xvi. 1. (1850) 146 = nanus.
trialatus, Boeck. in Flora, xli. (1859) 445 = Cyperus turgidulus.

SCIRPUS:—
triangularis, Steud. Nom. ed. I. 752 = pungens.
triangulatus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 217 = mucronatus.
tricarinatus, Pers. Syn. i. 68 = mucronatus.
trichobasis, *Baker, in Journ. Linn. Soc.* xx. (1883) 298.—Madag.
trichodes, Muhl. Desc. Gram. 30 = Eleocharis acicularis.
trichoides, H. B. & K. Nov. Gen. et Sp. i. 225 = Eleocharis trichoides.
trichophyllus, Willd. ex Kunth, Enum. Pl. ii. 224 = Fimbristylis Vahlii.
tridentatus, Roxb. Hort. Beng. [81]; Fl. Ind. i. 225 = maritimus.
triflorus, Poir. Encyc. ii. 249 = Rynchospora triflora.
trigonus, Nolte, Novit. Fl. Holsat. 9 = lacustris.
trigonus, Roth, Neue Beytr. i. 90 = triqueter.
trigynus, Linn. Mant. ii. 180 = Ficinia scariosa.
triqueter, R. Br. Prod. 223 = pungens.
triqueter, Gren. & Godr. Fl. Fr. iii. 373 = littoralis.
triqueter, *Linn. Mant.* i. 29.—Europ.; Austral.
triqueter, Torr. in Pacif. Rail. Rep. iv. 153 = Olneyi.
trispicatus, Linn. f. Suppl. 103 = Ficinia sylvatica.
trispicatus, Thunb. Prod. Fl. Cap. 17 = Ficinia membranacea.
tristachyos, Boj. Hort. Maurit. 383 (= *Isolepis barbata*).—Geront. trop.
tristachyos, Rottb. Desc. Nov. Pl. 48; Progr. 25 = Ficinia tristachya.
tristachyos, Roxb. Hort. Beng. 6; Fl. Ind. i. 221 = Fimbristylis ferruginea.
tristachyos, Vahl, ex Kunth, Enum. Pl. ii. 254 = Ficinia Bergiana.
tristachyos, Zoll. ex Steud. Syn. Pl. Cyp. 96 = supinus.
truncatus, Sieber, ex Kunth, Enum. Pl. ii. 254 = Ficinia lateralis.
truncatus, Thunb. Prod. Pl. Cap. 17 = Ficinia truncata.
tuberculatus, Beyr. ex Kunth, Enum. Pl. ii. 145 = Eleocharis tortilis.
tuberculatus, Ell. Sketch, i. 78 = Eleocharis tuberculata.
tuberculosus, Michx. Fl. Bor. Am. i. 30 = Eleocharis tuberculosa.
tuberosus, Desf. Fl. Atlant. i. 50 = maritimus.
tuberosus, Roxb. Hort. Beng. 6; Pl. Corom. iii. 25. t. 231 = Eleocharis tuberosa.
tumidus, Roxb. Hort. Beng. 6; Fl. Ind. i. 212 = Eleocharis plantaginea.
turgidus, Pers. Syn. i. 66 = Eleocharis ovata.
turgidus, Thuill. Fl. Par. ed. II. 22 = lacustris.
typhinus, Rafin. Atl. Journ. 18 (Quid ?).—Am. bor.
uliginosus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 520 = lacustris.
uliginosus, C. Koch, in Linnaea, xxi. (1848) 620 = Eleocharis acicularis.
umbellaris, Lam. Illustr. i. 141 = Fimbristylis globulosa.
umbellatus, Noronha, ex Thou. Prod. Phyt. 2, in Mélang. (1811), nomen (Quid ?).—Madag.
umbellatus, Roxb. ex Kunth, Enum. Pl. ii. 294 = Rynchospora aurea.
uncinatus, Willd. Sp. Pl. i. 300 = Fuirena uncinata.
uniflorus, Trautv. in Act. Hort. Petrop. v. (1877) 120 = pumilus.
uniglumis, Link, Jahrb. i. III. (1820) 77 = Eleocharis palustris.
uninodis, Boiss. Fl. Orient. v. 380, in syn. = supinus.
Urvillei, Boeck. in Linnaea, xxxvi. (1869–70) 510 = inundatus.
vaginatus, Salisb. Prod. 31 = Eriophorum vaginatum.
vaginatus, Thunb. Prod. Pl. Cap. 17 = Ficinia bulbosa.
Vahlii, Lam. Illustr. i. 139 = Fimbristylis Vahlii.
Vahlii, Willd. ex Kunth, Enum. Pl. ii. 85 = Mariscus filiformis.
validus, Benth. Pl. Hartw. 27 = riparius.
validus, Vahl, Enum. ii. 268 = lacustris.
variabilis, Poir. Encyc. Suppl. v. 99 = Fimbristylis diphylla.
varians, Boeck. in Linnaea, xxxvi. (1869–70) 698 = inundatus.
variegatus, Poir. Encyc. vi. 749 = Eleocharis variegata.
varius, Schreb. ex Schweigg. & Koerte, Fl. Erlang. 11 = Eleocharis palustris.
velatus, Poir. Encyc. Suppl. v. 99 = Fimbristylis squarrosa.

SCIRPUS:—
venustulus, *Boeck. in Linnaea*, xxxvi. (1869–70) 479.—Afr. austr.
verrucosulus, *Steud. in Flora*, xii. (1829) 145.—Afr. austr.
vestitus, Reichb. ex Boeck. in Linnaea, xxxvi. (1869–70) 753 = Fimbristylis vestita.
villosus, Poir. Encyc. vi. 780 = Fimbristylis diphylla.
virens, Boeck. in Flora, lviii. (1875) 261 = Thouarsianus.
viscosus, Lam. Illustr. i. 142 = Cyperus elegans.
viviparus, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 93 = Eleocharis tenuissima.
vulgaris, [Mazziari,] in Ionios Anthol. ii. (1834) 460 = maritimus.
Wallichianus, Spreng. Syst. iv. Cur. Post. 27 = Fimbristylis Wallichiana.
Wallichii, Nees, in Wight, Contrib. 112 = erectus.
Wallrothii, Bernh. in Flora, viii. (1825) 607 = rufus.
Warmingii, Boeck. in Kjoeb. Vidensk. Meddel. (1869) 136; et in Linnaea, xxxvi. (1869–70) 744 (Fimbristylidis sp.).—Bras.
Wichurai, Boeck. in Linnaea, xxxvi. (1869–70) 729 = Eriophorum cyperinum.
Wightianus, Boeck. l. c. 765 (Fimbristylidis sp.).—As. et Afr. trop.
Willemetii, *Steud. Nom.* ed. I. 752.—Ins. Maurit.
Wolfgangii, Bess. ex Schult. Mant. ii. 536 = lacustris.
Wolfii, A. Gray, in Proc. Am. Acad. x. (1874) 77 (Eleocharidis sp.).—Am. bor.
Xyridis, Poir. Encyc. Suppl. v. 98 = Fimbristylis Xyridis.
yokoscencis, Franch. & Sav. Enum. Pl. Jap. ii. 543 = nanus.
Zeyheri, Boeck. in Linnaea, xxxvi. (1869–70) 752 (Bulbostylidis sp.).—Afr. austr.

SCIRRHOPHORUS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 188 = SKIRRHOPHORUS, DC. = **Angianthus**, Wendl. (Compos.).

SCIURIS, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 150, 153 = **Galipea**, Aubl. (Rutac.).
aromatica, Vahl, Enum. i. 101 = G. aromatica.
bracteata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 156 = G. bracteata.
corymbosa, Spreng. Syst. i. 38 = G. trifoliata.
multiflora, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 155 = G. multiflora.
odoratissima, D. Dietr. Syn. Pl. i. 100 = Cusparia odoratissima.
simplicifolia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 157 = G. simplicifolia.
simplicifolia, Spreng. Syst. i. 39 = Cusparia macrophylla.
trifoliata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 155 = G. trifoliata.

SCIURIS, Schreb. Gen. i. 24 (1789) = **Raputia**, Aubl. (Rutac.).
aromatica, J. F. Gmel. Syst. 49 = R. aromatica.
Ossana, Nees & Mart. ex Steud. Nom. ed. II. ii. 542 = R. Ossana.

SCIURUS, D. Dietr. Syn. Pl. i. 32, 100 (1839), sphalm. = praec.

SCIZANTHUS, Pers. Syn. ii. 161 (1807), sphalm. = **Schizanthus**, Ruiz & Pav. (Solan.).

SCKUHRIA, Moench, Meth. 566 (1794) = **Siegesbeckia**, Linn. (Compos.).
dichotoma, Moench, l. c. = Siegesbeckia flosculosa.

SCLAERANTHUS, Thunb. Prod. Pl. Cap. 80 (1794) err. tpy. = **Scleranthus**, Linn. (Illecebr.).

SCLARAEA, Steud. Nom. ed. I. 752 (1821) = seq.

SCLAREA, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Salvia**, Tourn. (Labiat.).
Aethiopis, Mill. l. c. ed. VIII. n. 2 = Salvia Aethiopis.
argentea, Mill. l. c. n. 15 = Salvia argentea.
ceratophylla, Mill. l. c. n. 8 = Salvia ceratophylla.
decidua, Moench, Meth. 375 = Salvia Verbenaca.
distans, Moench, l. c. = Salvia austriaca.
glutinosa, Mill. Gard. Dict. ed. VIII. n. 11 = Salvia glutinosa.
indica, Mill. l. c. n. 9 = Salvia indica.

SCLAREA:—
lanata, Moench, Meth. 374 = Salvia Aethiopis.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = Salvia lusitanica.
mexicana, Mill. l. c. n. 14 = Salvia mexicana.
nemorosa, Mill. l. c. n. 6 = Salvia sylvestris.
orientalis, Mill. l. c. n. 10 = Salvia rosaefolia.
pratensis, Mill. l. c. n. 4 = Salvia pratensis.
punctata, Moench, Meth. 374 = Salvia sylvestris.
spinosa, Rafin. Fl. Tellur. iii. 94 = Salvia spinosa.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 7 = Salvia sylvestris.
syriaca, Mill. l. c. n. 5 = Salvia sylvestris.
tingitana, Rafin. Fl. Tellur. iii. 94 = Salvia foetida.
tomentosa, Mill. Gard. Dict. ed. VIII. n. 13 = Salvia canariensis.
triloba, Rafin. Fl. Tellur. iii. 94 = Salvia triloba.
tuberosa, Mill. Gard. Dict. ed. VIII. n. 12 = Salvia pratensis.
viscosissima, Moench, Meth. 376 = Salvia pyrenaica.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = Salvia Sclarea.

SCLEPSION, Rafin. ex Wedd. in Ann. Sc. Nat. Sér. IV. vii. (1857) 355 = **Laportea**, Gaudich. (Urtic.).
divaricatum, Rafin. ex Wedd. in DC. Prod. xvi. 1. 78 = L. canadensis.

SCLERACHNE, R. Br. in Benn. Pl. Jav. Rar. 15. t. 4 (1838). *GRAMINEAE*, Benth. & Hook. f. iii. 1113.
punctata, *R. Br. l. c.* 15.—Java.

SCLERACHNE, Torr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) 273 = **Thurberia**, Benth. (Gramin.).
arkansana, Torr. ex Trin. l. c. 274 = T. arkansana.
cyathopoda, F. Muell. Fragm. viii. 116 = Chionachne cyathopoda.
pilosa, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) 275 = T. arkansana.

SCLERANTHUS, Linn. Syst. ed. I (1735). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 19.
DITOCA, Banks & Soland. ex Gaertn. Fruct. ii. 196. t. 126 (1791).
KNAVEL, Adans. Fam. ii. 506 (1763).
MNIARUM, Forst. Char. Gen. 1. t. 1 (1776).
abyssinicus, adsurgens, aequidens, anisodus, argillophilus, arrectifolius, Aschersoni, banaticus, bigeminus, Bilimeckii, brachyphyllus, byzantinus, cancellatus, canescens, capitulatus, caudiger, comatus, condensatus, conocephalus, corymbulosus, crinitus, crispathus, Decandollii, densifolius, densiramaeus, diander, dicranifolius, diffusus, dinsteinensis, dissitiflorus, Drogeanus, echinophorus, Falkii, fasciculifolius, Fenzlii, filicaulis, filiferricaulis, floribundus, foliosus, Friedrichsthalii, Fritzscheanus, fuscus, gemmulosus, glomerifer, griseus, Haepi, hamulatus, Haslingeri, Hasskarlii, Hausmanni, Heldreichii, Herbichii, heterogeneus, Heufleri, Hoemeanus, Hohenackeri, holanthus, horridus, humilis, Janii, Kessneri, Kittelii, Kohlianus, Kotschyanus, laricifolius, laxus, lepidus, leptodus, leucoloma, longidens, longisetus, macilentus, macrocephalus, macronemus, Medusellus, micranthus, microcephalus, microdus, mollis, myrianthus, myriocarpus, nivemontanus, orthocarpus, orthoxys, oxystephanus, paradoxus, picturatus, pictus, pityophilus, pleiotomus, polyclados, porphyrocaulis, Poscharskyi, procerus, Pseudopolycarpus, pycnocarpus, Reichardti, remotiflorus, repens, retropuberulus, riparius, Rochelii, Schimperi, Schottii, Schroederi, scoparius, Sekerae, semiglaber, Serpentini, setifer, spergulifolius, squarrosus, stenoloma, stipatus, striatipes, sulcifer, Tabernaemontani, tenuis, tortuosus, valgus, variflexus, verticillifer, Wilkommii, Wollmari, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 114–115 (Quid ?).—Europ.
aetnensis, Strobl, l. c. xxiv. (1874) 148 = perennis.
anisochaitos, bjelagoranus, curvibracteatus, erlmagellus, jastrebicanus, micromeroides, montenegrinus, Pantocsekii, polychaitos, seticeps, subuncinatus, uncinellus, unculatus, unguiculatus, venustus, Reichb. ex Pant. in Oestr. Bot. Zeitschr. xxiv. (1874) 25–26 (Quid ?).—Europ.
annuus, *Linn. Sp. Pl.* 406.—Europ.; As. temp.

SCLERANTHUS:—
arenarius, Schur, Enum. Pl. Transs. 224 = perennis.
ascriviensis, Reichb. ex Vis. in Mem. Ist. Venet. xx. (1876–79) 218, nomen.—Europ.
attenuatus, Reichb. ex Lindem. in Bull. Soc. Nat. Mosc. xlix. (1875) II. 79, nomen (Quid?).—Rossia.
Augustae, Carthusianorum, chersonensis, cinereus, crispatulus, debilis, decipiens, dumosus, gypsophilanthus, hyssopifolius, inflexus, Johannis, leiphaemus, leucocarpus, leucococcus, madonius, marginellus, moenchiaeformis, myosoteidos, obscurus, oligoconus, pelviger, polybotrys, profusus, sanctus, Seladonius, setibracteatus, sordidus, Sprengelii, stenodus, subspathaceus, suprafastigiatus, tenellus, tenuicaulis, tenuifolius, Theodori, vagans, Reichb. ex Hoeme, in Oestr. Bot. Zeitschr. xxiv. (1874) 143–148. (Quid?).—Europ.
biennis, Reut. in Compt. Rend. Soc. Hallér. (1853–54) 20 = annuus.
biflorus, *Hook. f. Fl. N. Zel.* i. 74.—N. Zel.; Austral.
borborygius, Reichb. ex Holuby, in Oestr. Bot. Zeitschr. xxiv. (1874) 55 = annuus.
brachycarpus, brachyodon, Follioti, implicatus, leucoperos, longirameus, longiramosus, obsoletus, Petronellae, serpyllaceus, stenoleucus, Reichb. ex Wiesbaur, in Oestr. Bot. Zeitschr. xxvi. (1876) 152–154 (Quid?).—Europ.
brumalis, clevemontanus, compactus, concinnus, Durandoi, expansus, Havranus, Holubyi, Knäzicanus, laevigatus, leptochaetos, leucocarpus, paganus, rusticus, serotinus, soroanus, Tauscheri, Reichb. ex Holuby, in Oestr. Bot. Zeitschr. xxiv. (1784) 55–57 (Quid?).—Europ.
campestris, Schur, Enum. Pl. Transs. 224 = annuus.
caniceps, Reichb. ex Lindem. in Bull. Soc. Nat. Mosc. xlix. (1875) II. 79, nomen (Quid?).—Rossia.
cattaroanus, Reichb. ex Vis. in Mem. Ist. Venet. xx. (1876–79) 218, nomen = *ascriviensis*.
celtiniensis, Reichb. l. c.—Europ.
chaitophyllus, Cumani, cymulosus, engadineus, graniti, insignis, intricatus, juvencus, modestus, ptyocarpus, sparsiflorus, spectabilis, subcrinitus, verecundus, Warnstorfi, Reichb. ex Hausskn. in Oestr. Bot. Zeitschr. xxiii. (1873) 127–129 (Quid?).—Europ.
cinereus, Reichb. ex Holuby, in Oestr. Bot. Zeitschr. xxiv. (1874) 55 = annuus.
collinus, Hornung, ex Opiz, Natural. x. (1825) 232 = verticillatus.
collinus, Schur, Enum. Pl. Transs. 224 = annuus.
comosus, Dum. Fl. Belg. 23 = annuus.
corrigioleflorus, Reichb. ex Nym. Consp. 257 = perennis.
Csetzii, Reichb. in Oestr. Bot. Zeit. xxii. (1872) 115 = uncinatus.
curvibracteatus, Reichb. ex Vis. in Mem. Ist. Venet. xx. (1876–79) 218, nomen (Quid?).—Europ.
Delorti, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 205, 206 = verticillatus.
Delorti, Sauzé & Maill. Cat. Deux-Sèvres, 30 = S. Pseudopolycarpos.
diander, *R. Br. Prod.* 412.—Austral.
dichotomus, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 10 = perennis.
echinulifer, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 114 = annuus.
divaricatus, Dum. Fl. Belg. Prod. 23 = verticillatus.
fallax, Boenn. Prod. Fl. Monast. 123 = perennis.
fasciculatus, Hook. f. Fl. Tasman. i. 42 = biflorus.
fastigiatus, Hochst. in Flora, viii. (1825) 536 = perennis.
ferrugineus, Reichb. ex Nym. Consp. 257 = annuus.
Fussii, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 115 = uncinatus.
gracilidens, Reichb. ex Lindem. in Bull. Soc. Nat. Mosc. xlix. (1875) II. 79, nomen (Quid?).—Rossia.
glomeratus, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 115 = verticillatus *vel* annuus.
gypsophilanthus, Reichb. l. c. = perennis.
hamosus, Pouz. Fl. Gard, i. 371. t. 3 = uncinatus.
Haynaldi, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 115 = uncinatus.
hirsutus, J. & C. Presl, Delic. Prag. 66 = perennis.
intermedius, Boenn. ex Reichb. Fl. Germ. Excurs. 565 = perennis.
intermedius, Reichb. ex Nym. Consp. 257 = annuus.
Jankei, Reichb. in Oestr. Bot. Zeitschr. xxii. (1872) 115 = uncinatus.

SCLERANTHUS:—
Lindemanni, Reichb. ex Lindem. in Bull. Soc. Nat. Mosc. xlix. (1875) II. 79, nomen (Quid?).—Rossia.
marginatus, Guss. Fl. Sic. Prod. 486 = perennis.
marginellus, Reichb. ex Hausskn. in Oestr. Bot. Zeitschr. xxiii. (1873) 129 = annuus.
Martini, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 206 = uncinatus.
microdichodon, Reichb. ex Tauscher, in Oestr. Bot. Zeitschr. xxiii. (1873) 323 = dichotomus.
mniaroides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 13.—Austral.
neglectus, Rochel, ex Baumg. Enum. Stirp. Transs. iii. 345; et in Pl. Banat. Rar. 35. t. 3. f. 10 = perennis.
Nočanus, Reichb. ex Nym. Consp. 257 = perennis.
Orzinii, Reichb. ex Vis. in Mem. Ist. Venet. xx, (1876–79) 218, nomen (Quid?).—Europ.
perennis, *Linn. Sp. Pl.* 406.—Europ.
polycarpos, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 206 = uncinatus.
polycarpus, DC. ex Nym. Consp. 257 = verticillatus.
polycarpus, Linn. Cent. Pl. 216; Amoen. Acad. iv. 313 = verticillatus.
polycnemoides, *Willk. & Costa, in Linnaea*, xxx. (1859–60) 98.—Hispan.
praecox, Wallr. ex Hoeme, in Oestr. Bot. Zeitschr. xxiv. (1874) 144 = verticillatus.
Pseudopolycarpos, *Lacroix, in Bull. Soc. Bot. Fr.* vi. (1859) 558.—Europ.
Pseudoverticillatus, Reichb. ex Pant. in Oestr. Bot. Zeitschr. xxiv. (1874) 26 = verticillatus.
pungens, *R. Br. Prod.* 412.—Austral.
Reichenbachii, Tauscher, ex Hoeme, in Oestr. Bot. Zeitschr. xxiv, (1874) 145, nomen (Quid?).—Hungar.
rodnanus, Reichb. ex Nym. Consp. 257 = uncinatus.
tauricus, Presl, ex Nym. l. c. = annuus.
tenuissimus, Reichb. ex Nym. l. c. = verticillatus.
umagellus, Reichb. ex Vis. in Mem. Ist. Venet. xx. (1876–79) 218, nomen (Quid?).—Europ.
uncinatus, *Schur, in Verh. Siebenb. Ver. Naturw.* i. (1850) 107; ii. (1851) 10.—Europ. austr.
verticillatus, Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 205, 206 = S. Pseudopolycarpus.
verticillatus, *Tausch, in Flora*, xii. (1829) I. *Erg.* 50.—Europ.; As. Min.
Visianii, Reichb. ex Vis. in Mem. Ist. Venet. xx. (1876–79) 218, nomen.—Europ.

SCLERIA, Berg. in Vet. Acad. Handl. Stockh. xxvi. (1765) 142. tt. 4, 5. *CYPERACEAE*, Benth. & Hook. f. ii. 1070.
ANERMA, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 170 (1847).
CHONDROLOMIA, Nees, l. c. 173 (1847).
CRYPTOPODIUM, Schrad. ex Nees, l. c. 180 (1847).
CYLINDROPUS, Nees, in Linnaea, ix. (1834) 303; et in Edinb. N. Phil. Journ. xv. (Oct. 1834) 266.
DIAPHORA, Lour. Fl. Cochinch. 578 (1790).
DIPLACRUM, R. Br. Prod. 241 (1810).
DIPLOSCYPHUS, Liebm. in Vidensk. Selsk. Skr. V. ii. (1850) 262.
HYMENOLYTRUM, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 174. t. 22 (1842).
HYPOPORUM, Nees, in Linnaea, ix. (1834) 303.
MACROLOMIA, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 181. t. 24 (1847).
MASTIGOSCLERIA, Nees, l. c. 177 (1847).
OMOSCLERIA, Nees, l. c. 180 (1847).
OPHRYDIUM, Schrad. ex Nees, l. c. 183 (1847).
OPHRYOSCLERIA, Nees, l. c. 182. t. 25 (1847).
SCHIZOLEPIS, Schrad. ex Nees, l. c. 186. t. 26 (1847).
SPHAEROPUS, Boeck. in Flora, lvi. (1873) 89.
TRACHYLOMIA, Nees, in Mart. Fl. Bras. ii. I. 174 (1847.)
abortiva, *Nees, in Linnaea*, ix. (1834) 302.—Madag.
acanthocarpa, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 154.—Bras.
adamantina, Steud. Syn. Pl. Cyp. 177 = Lagenocarpus adamantinus.
affinis, Presl, ex Steud. Nom. ed. II. ii. 542 = pratensis.
africana, *Benth. in Benth. & Hook. f. Gen.* iii. 1071.—Afr. trop.
alata, Moon, Cat. Pl. Ceyl. 62 = hebecarpa.
albonigra, *A. St. Hil. Voy. Distr. Diam.* i. 371.—Bras.
alta, *Boeck. in Linnaea*, xxxviii. (1874) 485.—Ind. or.
anceps, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 257.—Mexic.
androgyna, *Nees, in Wight, Contrib.* 117.—Ind. or.
angusta, *Nees, in Linnaea*, ix. (1834) 303.—Afr. austr.

SCLERIA:—
angustata, Steud. Nom. ed. II. ii. 542, sphalm. = praec.
annularis, *Steud. Syn. Pl. Cyp.* 176.—Ind. or.; China.
approximata, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 118 = sumatrensis.
arguta, *Steud. Syn. Pl. Cyp.* 172.—Bras.
arundinacea, *Kunth, Enum. Pl.* ii. 347.—Am. trop.
aspera, Boeck. in Linnaea, xxxviii. (1874) 482 = S. Radula.
asperata, Presl, in Oken, Isis, xxi. (1828) 268 = pratensis.
asperrima, *Steud. Syn. Pl. Cyp.* 170.—Nicaragua.
atropurpurea, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 29.—Bras.
atrosanguinea, Hochst. ex Steud. Syn. Pl. Cyp. 175 = bulbifer.
auricoma, Otto, ex Sweet, Hort. Brit. ed. II. 549 (Quid?).—Am. bor.
Baldwinii, *Steud. Syn. Pl. Cyp.* 175.—Am. bor.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 602.—Malaya.
Barteri, *Boeck. in Linnaea*, xxxviii. (1874) 504.—Afr. trop.
Bertolonii, Martens, in Flora, xl. (1857) 570 = oryzoides.
biflora, Roxb. Hort. Beng. 66; Fl. Ind. iii. 573 = tesselata.
Boivini, *Steud. Syn. Pl. Cyp.* 173.—Madag.
Bourgeaui, *Boeck. in Flora*, lxv. (1882) 29.—Mexic.
bracteata, *Cav. Ic.* v. 34. t. 457.—Am. austr.
Brekiana, *Schlecht. in Bot. Zeit.* iii. (1845) 491.—Bras.
Brownei, *Kunth, Enum. Pl.* ii. 349.—Austral.
bulbifera, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 510.—Abyss.
caespitosa, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 167.—Afr. trop.
campestris, Steud. Syn. Pl. Cyp. 178 = Cryptangium campestre.
canaliculato-triquetra, *Boeck. in Flora*, lxii. (1879) 573.—Afr. trop.
capillaris, R. Br. Prod. 240 = lithosperma.
capitata, Willd. Sp. Pl. iv. 319 (Pteroscleriae sp.).—Guian.
caricina, *Benth. Fl. Austral.* vii. 426.—As. et Austral. trop.
caroliniana, Willd. Sp. Pl. iv. 318 = pauciflora.
cenchroides, Kunth, Enum. Pl. ii. 352 = hirtella.
cervina, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 171.—Afr. trop.
ceylanica, Kunth, Enum. Pl. ii. 358 = zeylanica.
chinensis, *Kunth, Enum. Pl.* ii. 357.—As. et Austral. trop.
chlorantha, *Boeck. in Linnaea*, xxxviii. (1874) 506.—Ins. S. Doming.
ciliaris, Nees, in Wight, Contrib. 117 = chinensis.
ciliata, *Michx. Fl. Bor. Am.* ii. 167.—Am. bor.; Ind. occ.
ciliolata, Boeck. in Flora, lxv. (1882) 31 = racemosa.
cincta, Steud. Syn. Pl. Cyp. 177 = Rhynchospora tenuis.
clathrata, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 510.—Abyss.
communis, Kunth, Enum. Pl. ii. 340 = pratensis.
comosa, *Steud. Syn. Pl. Cyp.* 171.—Bras.
complanata, *Boeck. in Flora*, lxii. (1879) 571.—Afr. trop.
composita, *Boeck. in Linnaea*, xxxviii. (1874) 444.—Bras.
conspersa, Sellow, ex Nees, in Mart. Fl. Bras. ii. I. 179 = pterota?
coriacea, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 259.—Costa Rica.
corymbifera, Boeck. in Linnaea, xxxviii. (1874) 537 = androgyna.
corymbosa, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 574 = androgyna.
cyanocarpa, Kunth, Enum. Pl. ii. 347 = sylvestris.
cymosa, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 155, 158.—Bras.
cyperina, *Kunth, Enum. Pl.* ii. 345.—Guian.
debilis, Wright, in Sauv. Fl. Cub. 186 = hemitaphra.
densifolia, Steud. Syn. Pl. Cyp. 178 = Cryptangium Minarum.
depauperata, *Boeck. in Flora*, lviii. (1875) 118.—Polynes.
dictyocarpa, Wright, ex Griseb. Cat. Pl. Cub. 249 = Torreyana.
Dietrichiae, Boeck. in Flora, lviii. (1875) 121 = hebecarpa.

SCLERIA:—
diffusa, Michx. ex Kunth, Enum. Pl. ii. 359 = verticillata.
Dillonia, Boeck. in Flora, lxi. (1878) 38 = foliosa.
dioica, Leconte, ex Britton, in Ann. N. York Acad. Sc. iii. (1885) 229 = Baldwinii.
discolor, *Boeck. in Linnaea*, xxxviii. (1874) 541.—Bras.
distans, R. Br. Prod. 240 = Brownei.
distans, *Poir. Encyc.* vii. 4.—Ind. occ.
diurensis, *Boeck. in Flora*, lxii. (1879) 573.—Afr. trop.
divaricata, *Nees, in Linnaea*, ix. (1834) 303.—Am. trop.
Doederleiniana, Boeck. in Engl. Bot. Jahrb. v. (1884) 512 = scrobiculata.
Dregeana, *Kunth, Enum. Pl.* ii. 354.—Afr. austr.
dumicola, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 169.—Afr. austr.
elata, *Thw. Enum. Pl. Zeyl.* 353.—As. trop.
elata, *Wright, in Sauv. Fl. Cub.* 186.—Cuba.
Elliottii, Chapm. Fl. S. U. St. 531 = ciliata.
elongata, *J. & C. Presl, Rel. Haenk.* i. 202.—Mexic.
erythrorrhiza, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 167.—Afr. trop.
exaltata, Boeck. in Engl. Bot. Jahrb. v. (1884) 511 = hebecarpa.
fasciculata, Willd. ex Kunth, Enum. Pl. ii. 301 = Rynchospora Cephalotes.
fenestrata, Franch. & Sav. Enum. Pl. Jap. ii. 549 = tessellata.
filiformis, Sw. Prod. Veg. Ind. Occ. 19 = lithosperma.
Flabellum, Poir. Encyc. vii. 1, sphalm. = S. Flagellum-nigrorum.
flaccida, Steud. Syn. Pl. Cyp. 174 = triglomerata.
flagellata, Sw. ex Boeck. in Linnaea, xxxviii. (1874) 506 = pterota.
Flagellum, Benth. in Hook. Niger Fl. 555 = ovuligera.
Flagellum, Griseb. Fl. Brit. W. Ind. 578 = reflexa.
Flagellum, Sw. Fl. Ind. Occ. i. 88 = seq.
Flagellum-nigrorum, *Berg. in Vet. Akad. Handl. Stockh.* xxvi. (1765) 144. t. 4.—Am. trop.
floribunda, H. B. & K. Nov. Gen. et Sp. i. 233 = bracteata.
foliosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 509.—Abyss.
foliosa, Wright, in Sauv. Fl. Cub. 187 = microcarpa.
foveolata, *Cav. Ic.* v. 35.—Panama.
foveolata, Llanos, Fragm. Fl. Filip. 103 = scrobiculata.
glandiformis, *Boeck. in Flora*, xli. (1858) 647.—Afr. trop.
glaucescens, J. & C. Presl, Rel. Haenk. i. 202 = lithosperma.
Glazioviana, *Boeck. in Flora*, lxv. (1882) 352.—Bras.
glomerulata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 170.—Afr. trop.
gracilis, *Ell. Sketch*, ii. 557.—Am. bor.; Ind. occ.
gracilis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = lithosperma.
gracillima, *Boeck. in Flora*, xlii. (1879) 570.—Afr. trop.
Graeffeana, Boeck. l. c. lviii. (1875) 121 = margaritifera.
grandifolia, Miq. in Linnaea, xix. (1847) 230 = arundinacea.
guahensis, Schrad. ex Boeck. l. c. xxxviii. (1874) 512 = margaritifera.
guianensis, Steud. Syn. Pl. Cyp. 177 = Lagenocarpus guianensis.
haematostachys, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 512.—Java.
Harlandi, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 248.—Ins. Hongk.
Hasskarliana, Boeck. in Engl. Bot. Jahrb. v. (1884) 511 = elata, *Thw.*
hebecarpa, *Nees, in Wight, Contrib.* 117.—As. et Austral. trop.
hemitaphra, *Steud. Syn. Pl. Cyp.* 169.—Am. bor.; Ind. occ.
Hildebrandtii, *Boeck. in Flora*, lxiii. (1880) 454.—Afr. trop.
Hilsenbergii, *Ridl. in Journ. Bot.* xxii. (1884) 16.—Madag.
hirsuta, *Boeck. in Linnaea*, xxxviii. (1874) 489.—Java.
hirsuta, Moon, Cat. Pl. Ceyl. 103 = junciformis, *Thw.*
hirta, *Boeck. in Flora*, lxv. (1882) 29.—Mexic.
hirta, Willd. ex Kunth, Enum. Pl. ii. 352 = hirtella.
hirtella, Boeck. in Linnaea, xxxviii. (1874) 439 = melanotricha, distans.

SCLERIA:—
hirtella, Michx. Fl. Bor. Am. ii. 168 = ciliata.
hirtella, *Sw. Prod. Veg. Ind. Occ.* 19.—Afr. et Am. trop.
hispidula, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 511.—Abyss.
holcoides, *Kunth, Enum. Pl.* ii. 354.—Afr. austr.
homonyma, Steud. Syn. Pl. Cyp. 179 = scabriuscula.
Hookeriana, *Boeck. in Linnaea*, xxxviii. (1874) 498.—Ind. or.
Hostmanniana, Steud. Syn. Pl. Cyp. 178 = Cryptangium leptocladum.
humilis, Britton, in Ann. N. York Acad. Sc. iii. (1885) 235 = hirtella.
humilis, Steud. Syn. Pl. Cyp. 178 = Cryptangium humile.
hypoxis, Schweinf. ex Boeck. in Linnaea, xxxviii. (1874) 465 = Schimperiana.
interrupta, Michx. Fl. Bor. Am. ii. 168 = hirtella.
interrupta, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 112.—Guian.
interrupta, Schlecht. in Linnaea, xx. (1847) 544 = hispidula.
japonica, Steud. Syn. Pl. Cyp. 169 = hebecarpa.
Jelskiana, Boeck. in Linnaea, xxxviii. (1874) 469 = Torreyana.
junciformis, Kunth, Enum. Pl. ii. 356 = Cryptangium junciforme.
junciformis, *Thw. Enum. Pl. Zeyl.* 354.—Zeylan.
junciformis, *Welw. ex Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 168.—Afr. trop.
Junghuhniana, *Boeck. in Linnaea*, xxxviii. (1874) 499.—Java.
Kappleriana, Hochst. ex Steud. Syn. Pl. Cyp. 172 = arundinacea.
Kunthiana, Steud. l. c. 176 = verticillata, *Muhl.*
Kunthii, Miq. Sert. Exot. 4. t. 2 = Cryptangium leptocladum.
lacustris, *Wright, in Sauv. Fl. Cub.* 185.—Cuba.
laevis, Willd. Sp. Pl. iv. 13 = levis.
lagoënsis, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 151.—Am. trop.
lateriflora, Boeck. in Linnaea, xxxviii. (1874) 455 = ceylanica.
latifolia, Moon, Cat. Pl. Ceyl. 103 = oryzoides.
latifolia, Nees, in Flora, xi. (1828) 303 = Sieberi.
latifolia, Reichb. ex Nees, in Mart. Fl. Bras. ii. 1. 183 = mitis.
latifolia, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 268 = pratensis, pterota.
latifolia, *Sw. Prod. Veg. Ind. Occ.* 18.—Am. trop.
laxa, *R. Br. Prod.* 240.—Austral.
laxa, Torr. in Ann. Lyc. N. York, iii. (1836) 376 = Torreyana.
leptocladia, Poepp. ex Kunth, Enum. Pl. ii. 355 = Cryptangium leptocladum.
leptoculmis, Wood, Class Book (1861), 746 = oligantha.
leptostachya, *Kunth, Enum. Pl.* ii. 354.—Bras.
levis, *Retz. Obs.* iv. 13.—As. trop.
Liebmanni, *Steud. Syn. Pl. Cyp.* 179.—Am. trop.
lithosperma, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 574 = elata, *Thw.*
lithosperma, *Sw. Prod. Veg. Ind. Occ.* 18.—Amphig. trop.
longifolia, *Boeck. in Flora*, lxv. (1882) 30.—Madag.
lucida, Steud. Syn. Pl. Cyp. 168 = mitis.
luzulaeformis, Wright, in Sauv. Fl. Cub. 188 = Liebmanni.
mackaviensis, *Boeck. in Flora*, lviii. (1875) 119.—Austral.
macrantha, Boeck. l. c. xli. (1858) 647 = ciliata.
macrantha, Boeck. l. c. lxii. (1879) 572 = melanomphala.
macrocarpa, Nees, in Mart. Fl. Bras. ii. 1. 183 = racemosa.
macrocarpa, Salzm. ex Boeck. in Linnaea, xxxviii. (1874) 521 = paludosa.
macrocarpa, Wall. Cat. n. 3538 = Scirpodendron costatum.
macrophylla, J. & C. Presl, Rel. Haenk. i. 200 = bancana.
macrosperma, Nees, ex Kunth, Enum. Pl. ii. 344 = racemosa.
madagascariensis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 514.—Madag.
major, Moon, Cat. Pl. Ceyl. 62 = androgyna.

SCLERIA:—
malaccensis, Boeck. in Linnaea, xxxviii. (1874) 507 = bancana.
margaritifera, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 268 = pratensis.
margaritifera, *Willd. Sp. Pl.* iv. 312.—Austral.; Polynes.
Martii, *Steud. Syn. Pl. Cyp.* 171.—Am. trop.
Mechowiana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 510.—Afr. trop.
melaleuca, *Reichb. ex Schlecht. & Cham. in Linnaea*, vi. (1831) 29.—Am. trop.
melanocephala, Drège, Zwei Pfl. Docum. 147, sphalm. = seq.
melanomphala, *Kunth, Enum. Pl.* ii. 345.—Afr. austr. et trop.
melanosperma, Nees & Arn. ex Boeck. in Linnaea, xxxviii. (1874) 515 = hebecarpa.
melanostoma, Boeck. l. c. 514 = hebecarpa.
melanotricha, *Hochst. & A. Rich. Tent. Fl. Abyss.* ii. 511.—Abyss.
melicoides, *Schlecht. in Linnaea*, xxvi. (1853) 127.—N. Granat.
mexicana, *Boeck. l. c.* xxxviii. (1874) 500.—Mexic.
Meyeriana, *Kunth, Enum. Pl.* ii. 354.—Afr. austr.
Michauxii, Chapm. Fl. S. U. St. 522 = hirtella.
micrantha, Poir. Encyc. Suppl. v. 108 = hemitaphra.
microcarpa, *Nees, in Linnaea*, ix. (1834) 302.—Am. trop.
micrococca, *Steud. Syn. Pl. Cyp.* 179.—Nicaragua.
microdiscus, Steud. l. c. 175 = Rynchospora Berteroana.
microstachya, *Boeck. in Flora*, lxiii. (1880) 454.—Bras.
Minarum, Steud. Syn. Pl. Cyp. 178 = Cryptangium Minarum.
minutiflora, Rich. ex Spreng. Syst. iii. 831 = Rynchospora micrantha.
mitis, *Berg. in Vet. Akad. Handl. Stockh.* xxvii. (1765) 145. t. 5.—Am. austral.
mollis, Kunth, Enum. Pl. ii. 352 = hirtella.
Moritziana, *Boeck. in Linnaea*, xxxviii. (1874) 460.—Venezuela.
mucronata, *Poir. Encyc.* vii. 3.—Ind. or.
Muehlenbergii, Steud. Nom. ed. II. ii. 543; Schlecht. in Bot. Zeit. iii. (1845) 474 = Torreyana.
Muhlenbergiana, Liebm. in Kjoeb. Vidensk. Selsk. Skr. ii. (1851) 258 = hemitaphra.
multifoliata, *Boeck. in Linnaea*, xxxviii. (1874) 510.—Malaya.
muricata, *Boeck. l. c.* 540.—Am. trop.
myricocarpa, *Kunth, Enum. Pl.* ii. 347.—Bras.
myriocarpa, Steud. Syn. Pl. Cyp. 172, sphalm. = praec.
Naumanniana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 94.—Afr. trop.
Neesiana, Hook. & Arn. Bot. Beech. Voy. 229 = scrobiculata.
Neesii, *Kunth, Enum. Pl.* ii. 358.—Zeylan.; Borneo.
nervosa, *Wikstr. in Vet. Akad. Handl. Stockh.* (1827) 75.—Ins. Guadal.
nigricans, *J. & C. Presl, Rel. Haenk.* i. 201.—N. Granat.
nitida, Willd. ex Kunth, Enum. Pl. ii. 350 = triglomerata.
notoptera, Nees, in Linnaea, ix. (1834) 303 = Pteroscleria guianensis.
Novae-Hollandiae, Boeck. in Flora, lviii. (1875) 120 = laxa.
nutans, Kunth, Enum. Pl. ii. 351 = hirtella.
Oakesiana, Robbins, ex Britton, in Ann. N. York Acad. Sc. iii. (1885) 234 = pauciflora.
oligantha, *Michx. Fl. Am. Bor.* ii. 167.—Am. bor.
olyroides, *Kunth, Enum. Pl.* ii. 348.—Bras.
Onoei, *Franch. & Sav. Enum. Pl. Jap.* ii. 549.—Japon.
orizoides, Boeck. in Linnaea, xxxviii. (1874) 492 = seq.
oryzoides, *J. & C. Presl, Rel. Haenk.* i. 201.—Geront. trop.
Ottonis, *Boeck. in Linnaea*, xxxviii. (1874) 490.—Cuba.
ovuligera, *Nees, l. c.* ix. (1834) 303.—Afr. trop.
ovuligera, Reichb. ex Nees, l. c. = microcarpa.
pallidiflora, Boeck. in Flora, lviii. (1875) 119 = Brownei.
palmifolia, Hoffmgg. & Schlecht. in Bot. Zeit. iii. (1845) 492 = paludosa.

SCLERIA:—
paludosa, *Kunth, Enum. Pl.* ii. 344.—Am. trop.
panicoides, *Kunth, l. c.* 348.—Am. trop.
papillata, Willd. ex Kunth, l. c. 345 = bracteata.
parvula, Steud. Syn. Pl. Cyp. 174 = tessellata.
pauciflora, *Muhl. ex Willd. Sp. Pl.* iv. 318.—Am. bor.
pauciflora, Sauv. Fl. Cub. 183 = gracilis.
pergracilis, *Kunth, Enum. Pl.* ii. 354.—Geront. trop.
phyllomacra, *Steud. Syn. Pl. Cyp.* 173.—Hab. ?
phylloptera, *Wright, ex Griseb. Cat. Pl. Cub.* 248.—Cuba.
pilosa, Boeck. in Linnaea, xxxviii. (1874) 473 = junciformis, *Thw.*
pleostachya, *Kunth, Enum. Pl.* ii. 355.—Bras.
Ploemii, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 513.—Java.
plusiophylla, *Steud. Syn. Pl. Cyp.* 172.—Bras.
poaeformis, *Retz. Obs.* iv. 13.—Ind. or.
poeoides, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 170.—Afr. trop.
Poeppigii, *Steud. Syn. Pl. Cyp.* 171.—Am. trop.
polycarpa, Boeck. in Linnaea, xxviii. (1874) 509 = margaritifera.
polyphylla, Steud. Syn. Pl. Cyp. 178 = Cryptangium polyphyllum.
porphyrorhiza, *Wright, in Sauv. Fl. Cub.* 188.—Cuba.
praealta, Salzm. ex Schlecht. in Bot. Zeit. iii. (1845) 461 = mitis.
pratensis, *Lindl. ex Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 121.—Am. trop.
propinqua, Steud. Syn. Pl. Cyp. 169 = tessellata.
pterota, *Presl, in Oken, Isis*, xxi. (1828) 268.—Ind. occ.
pubescens, *Steud. Syn. Pl. Cyp.* 168.—Ins. Hongkong.
pulchella, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 168.—Afr. trop.
purpurascens, Benth. Fl. Hongk. 400 = Harlandi.
purpurascens, Steud. Syn. Pl. Cyp. 169 = sumatrensis.
purpurea, Poir. Encyc. vii. 4 = lithosperma.
purpureo-vaginata, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 513.—Ins. Philipp.
purpureo-zonata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 258 = oligantha.
pygmaea, *R. Br. Prod.* 240.—Austral.
pygmaea, Nees, ex Spreng. Syst. iv. Cur. Post. 313 = Rynchospora micrantha.
pyramidalis, Hochst. ex Steud. Syn. Pl. Cyp. 171 = Martii.
racemosa, Boj. Hort. Maurit. 385 = Sieberi.
racemosa, *Poir. Encyc.* vii. 6.—Madag.
Radula, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 232.—China.
reflexa, Benth. in Hook. Niger Fl. 555 = Barteri.
reflexa, *H. B. & K. Nov. Gen. et Sp.* i. 232.—Am. trop.
remota, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 169.—Afr. trop.
Renggeriana, Steud. Syn. Pl. Cyp. 173 = reflexa.
reticularis, *Michx. Fl. Bor. Am.* ii. 167.—Am. bor.
reticularis, Muhl. Desc. Gram. 266 = Torreyana.
reticularis, J. & C. Presl, Rel. Haenk. i. 202 = hemitaphra.
rigida, Kunth, Enum. Pl. ii. 355 = Lagenocarpus rigidus.
riparia, *Poepp. & Kunth, ex Kunth, l. c.* 341.—Peruv.
rugosa, *R. Br. Prod.* 240.—Austral.
Rutenbergiana, *Boeck. in Bremen, Abh.* vii. (1880) 40.—Madag.
salebrosa, Spreng. ex Boeck. in Linnaea, xxxviii. (1874) 438 = pergracilis.
scabra, *Willd. Sp. Pl.* iv. 315.—Venezuela.
scabriuscula, *Schlecht. in Linnaea*, xx. (1847) 544.—Mexic.
Schiedeana, *Schlecht. in Bot. Zeit.* iii. (1845) 489.—Mexic.
Schiedeana, Schlecht. l. c. 494 = scabriuscula.
Schimperiana, *Boeck. in Linnaea*, xxxviii. (1874) 466.—Abyss.
Schweinfurthiana, *Boeck. in Flora*, lxii. (1879) 570.—Afr. trop.
scindens, Nees, in Linnaea, ix. (1834) 303 = pterota.
scrobiculata, Mor. Syst. Verz. Zoll. 98 = chinensis.
scrobiculata, *Nees & Meyen, in Wight, Contrib.* 117.—China; Malaya.

SCLERIA:—
scrobiculata, Zoll. Syst. Verz. Ind. Archip. ii. 61 = hebecarpa, multifoliata.
Selloana, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 179 = pratensis.
Sellowiana, *Kunth, Enum. Pl.* ii. 350.—Bras.
setacea, Poir. Encyc. vii. 4 = hemitaphra.
setifera, Boeck. in Flora, xli. (1858) 648 = Schoenus axillaris.
setigera, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 575 = sumatrensis.
setoso-asperula, Boeck. in Flora, lviii. (1875) 120 = Brownei.
setuloso-ciliata, *Boeck. l. c.* lxv. (1882) 30.—Am. centr.
Sieberi, *Nees, in Linnaea*, ix. (1834) 303.—Ins. Maurit.
silvestris, Nees, in Mart. Bras. ii. 1. 188 = sylvestris.
simplicior, Steud. Syn. Pl. Cyp. 169 = pratensis.
Sororia, *Kunth, Enum. Pl.* ii. 343.—Bras.
sphacelata, *F. Muell. Fragm.* ix. 20.—Austral.
spicaeformis, *Benth. in Hook. Niger Fl.* 556.—Afr. trop.
Steudeliana, *Miq. Fl. Ind. Bat.* iii. 344.—Austral.
stipularis, *Nees, in Hook. Journ. Bot.* ii. (1840) 394.—Am. trop.
Stocksiana, *Boeck. in Linnaea*, xxxviii. (1874) 474.—Ind. or.
stricta, Kunth, Enum. Pl. ii. 355 = Cryptangium uliginosum.
stricta, Moon, Cat. Pl. Ceyl. 62 = Neesii.
sumatrensis, *Retz. Obs.* v. 19. t. 2.—Ind. or.; Malaya.
sylvestris, *Poepp. & Kunth, in Enum. Pl.* ii. 346.—Am. trop.
tenacissima, *Steud. Syn. Pl. Cyp.* 175.—Am. trop.
tenella, Griseb. Cat. Pl. Cub. 249 = distans, Liebmanni.
tenella, *Kunth, Enum. Pl.* ii. 353.—Guian.
tenuiflora, Willd. ex Kunth, l. c. = verticillata.
tenuis, Retz. Obs. iv. 13 = lithosperma.
tessellata, Benth. Fl. Hongk. 399 = Steudeliana.
tessellata, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 362 = scrobiculata.
tessellata, *Willd. Sp. Pl.* iv. 315.—As. et Austral. trop.
testacea, *Nees, in Linnaea*, ix. (1834) 303.—Ins. Sandvic.
Thomsoniana, *Boeck. in Linnaea*, xxxviii. (1874) 479.—Reg. Himal.
Thwaitesiana, Boeck. l. c. 454 = zeylanica.
timorensis, Nees, in Linnaea, ix. (1834) 303 = scrobiculata.
topazina, Steud. Syn. Pl. Cyp. 177 = Lagenocarpus topazinus.
Torreyana, *Walp. Ann.* iii. 696.—Am. bor. et austr.
tremula, Steud. Syn. Cyp. 177 = Lagenocarpus tremulus.
trialata, Bert. ex Boeck. in Linnaea, xxxviii. (1874) 521 = mitis.
trialata, *Poir. Encyc.* vii. 6.—Madag.
trichopoda, Wright, ex Britton, in Ann. N. York Acad. Sc. iii. (1885) 232 = Torreyana.
triglomerata, *Michx. Fl. Bor. Am.* ii. 168.—Am. bor.
trigonocarpa, Steud. Syn. Pl. Cyp. 171 = arundinacea.
tristis, *A. St. Hil. Voy. Distr. Diam.* i. 370.—Bras.
tuberculata, *Boeck. in Linnaea*, xxxviii. (1874) 540.—Am. trop.
uliginosa, Hochst. ex Boeck. l. c. 471 = tessellata.
ustulata, *Ridl. in Trans. Linn. Soc.* Ser. II. ii. (1884) 168.—Afr. trop.
vaginata, *Steud. Syn. Pl. Cyp.* 179.—Am. bor.
variegata, *Steud. l. c.* 176.—Bras.
velutina, Steud. l. c. 177 = Lagenocarpus velutinus.
verrucosa, *Willd. Sp. Pl.* iv. 313.—Afr. trop.
verticillata, *Muhl. ex Willd. l. c.* 317.—Am. bor.
verticillata, Nees, ex Kunth, Enum. Pl. ii. 354 = tenella.
vestita, Boeck. in Linnaea, xxxviii. (1874) 482 = hebecarpa.
virgata, *Steud. Syn. Pl. Cyp.* 176.—Bras.
viridifolia, Steud. l. c. 177 = Lagenocarpus Martii.
waigiouensis, Steud. l. c. 173 = scrobiculata.
Warmingiana, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 152.—Bras.
Weigeltiana, Schrad. ex Boeck. in Linnaea, xxxviii. (1874) 504 = reflexa.
Wichurai, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 510.—Ins. Philipp.
Wightiana, Steud. Syn. Pl. Cyp. 176 = lithosperma.
zelanica, Moon, Cat. Pl. Ceyl. 62 = levis.
zeylanica, *Poir. Encyc.* vii. 3.—Ind. or.; Malaya.

SCLEROBASIS, Cass. in Bull. Soc. Philom. (1818) 73 = **Senecio**, Tourn. (Compos.).
rigida, Cass. in Dict. Sc. Nat. xlviii. 146 = Sen. rigidus.
Sonneratii, Cass. in Bull. Soc. Philom. (1818) 74 = Sen. Sonneratii.

SCLEROCALYX, Nees, in Benth. Bot. Voy. Sulph. 145 (1844). *ACANTHACEAE*, Benth. & Hook. f. ii. 1083.
GYMNACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
mexicanum, *Nees, in Benth. Bot. Voy. Sulph.* 145.—Mexic.

SCLEROCARPA, Sond. in Linnaea, xxiii. (1850) 26, sphalm. = **Sclerocarya**, Hochst. (Anacard.).

SCLEROCARPUS, Jacq. Ic. Pl. Rar. t. 176 (1782); et in Act. Helv. ix. (1787) 34. t. 2. *COMPOSITAE*, Benth. & Hook. f. ii. 364.
ALDAMA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 14 (1824).
DICHOTOMA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 364 (1873).
africanus, *Jacq. Ic. Pl. Rar.* i. 17. t. 176.—Afr. trop.
dentatus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 164.—Mexic.
discoideus, *Vatke, in Linnaea*, xxxix. (1875) 495.—Afr. trop.
divaricatus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 164.—Mexic.
exiguus, *Sm. in Rees, Cycl.* xxxi. n. 3.—Am. bor.
gracilis, *Sm. l. c.* n. 2.—Am. bor.
Kerberi, *Fourn. in Bull. Soc. Bot. Fr.* xxx. (1883) 183.—Mexic.
oracilis, Steud. Nom. ed. II. ii. 543, err. typ. = gracilis.
Schiedeanus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 164.—Mexic.
uniserialis, *Benth. & Hook. f. Gen.* ii. 364.—Mexic.

SCLEROCARYA, Hochst. in Flora, xxvii. (1844) Bes. Beil. 1. *ANACARDIACEAE*, Benth. & Hook. f. i. 427.
SCLEROCARPA, Sond. in Linnaea, xxiii. (1850) 26, sphalm.
Birroea, *Hochst. in Flora*, xxvii. (1844) *Bes. Beil.* 1.—Abyss.
caffra, *Sond. in Linnaea*, xxiii. (1850) 26.—Afr. austr.
castanea, *Baker, Fl. Maurit.* 63.—Ins. Mascar.
Shakua, *Baker, l. c.*—Ins. Mascar.

SCLEROCEPHALUS, Boiss. Diagn. Ser. I. iii. 12 (1843). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 17.
arabicus, *Boiss. l. c.*—Afr. bor.; Arab.; Pers.
Aucheri, Walp. Rep. v. 75 = arabicus.

SCLEROCHAETIUM, Nees, in Linnaea, vii. (1832) 511 = **Tetraria**, Beauv. (Cyperac.).
angustifolium, Hochst. in Flora, xxviii. (1845) 762.—Afr. austr.
giganteum, Steud. Syn. Pl. Cyp. 159 = *thermale*.
involucratum, Nees, in Linnaea, vii. (1832) 511.—Afr. austr.
Koenigii, Nees, ex Hochst. in Flora, xxviii. (1845) 763 = *Rottboellii*.
Rottboellii, Nees, in Linnaea, viii. (1833) 89; ix. (1835) 302; x. (1835-36) 198.—Afr. austr.
spirale, Hochst. in Flora, xxviii. (1845) 761.—Afr. austr.
thermale, Nees, in Linnaea, vii. (1832) 512.—Afr. austr.

SCLEROCHITON, Harv. in Hook. Lond. Journ. Bot. i. (1842) 27. *ACANTHACEAE*, Benth. & Hook. f. ii. 1090.
ISACANTHUS, Nees, in DC. Prod. xi. 278 (1845).
Harveyanus, *Nees, in DC. Prod.* xi. 279.—Afr. austr.
Vogelii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 37.—Afr. trop.

SCLEROCHLAMYS, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 76 = **Kochia**, Roth (Chenop.).
brachyptera, F. Muell. l. c. = K. brachyptera.

SCLEROCHLOA, Beauv. Agrost. 97 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1194.
AMBLYCHLOA, Link, in Linnaea, xvii. (1843) 399.
angusta, *Nees, Fl. Afr. Austr.* 381.—Afr. austr.
arenaria, Nees, l. c. 380 = Glyceria distans.
articulata, Link, Enum. Hort. Berol. i. 90 = Festuca divaricata.
Borreri, Bab. Man. Brit. Bot. ed. I. 370 = Glyceria Borreri.
dichotoma, Link, Enum. Hort. Berol i. 90 = Cutandia maritima.
distans, Bab. Man. Brit. Bot. ed. I. 370 = Glyceria distans.
divaricata, Beauv. Agrost. 98 = Sphenopus Gouani.
divaricata, Link, Hort. Berol. ii. 273 = Festuca divaricata.
dura, *Beauv. Agrost.* 98.—Europ.; Afr. bor.; Oriens.
expansa, Link, Hort. Berol. ii. 274 = Sphenopus Gouani.
hemipoa, Guss. Fl. Sic. Syn. i. 93 = Festuca hemipoa.
loliacea, Syme, Engl. Bot. ed. III. xi. 110. t. 1759 = *Festuca loliacea*, Huds.
loliacea, J. Woods, Tourist Fl. 411 = Desmazeria loliacea.
maritima, Lindl. in Bab. Man. Brit. Bot. ed. I. 370 = Glyceria maritima.
maritima, Sweet, Hort. Brit. ed. I. 453 = Cutandia maritima.
memphitica, Nym. Syll. Suppl. 74 = Cutandia scleropoides.
multiculmis, Syme, Engl. Bot. ed. III. xi. 103 = Glyceria distans.
patens, Presl, Cyp. & Gram. Sic. 45 = Festuca rigida.
philistaea, Steud. Syn. Pl. Gram. 428, sphalm. = Cutandia philistaea.
poaeformis, Link, Handb. i. 69, in syn. = Poa violacea.
procumbens, Beauv. Agrost. 98 = Glyceria procumbens.
pumila, Steud. Syn. Pl. Gram. 428, sphalm. (= *Scleropoa pumila*).—Arab.
rigida, Beauv. Agrost. 97; et ex Panz. in Denkschr. Bot. Regensb. i. II. (1818) 180 = Festuca rigida.
rigida, Parl. Fl. Palerm. i. 213 = Festuca hemipoa.
vestita, De Not. Ind. Sem. Hort. Genuen (1847).—Cf. Ann. Sc. Nat. Sér. III. ix. (1848) 326 = Festuca dichotoma.
Zwierleinii, *Lojac. in Natural. Sicil.* iv. (1885) 138.—Sicil.

SCLEROCHLOA, Reichb. Ic. Fl. Germ. xi. 23. t. 58 (1834) = **Festuca**, Tourn. (Gramin.).

SCLEROCHORTON, Boiss. Fl. Orient. ii. 968 (1872). *UMBELLIFERAE.*
Haussknechtii, *Boiss. l. c.*—Persia.
junceum, *Boiss. l. c.* 969.—Graecia.

SCLEROCLADUS, Rafin. Sylva Tellur. 35 (1838) = **Bumelia**, Sw. (Sapot.).
tenax, Rafin. l. c. = B. tenax.

SCLEROCOCCUS, Bartl. Ord. 210 (1830) = **Hedyotis**, Linn. (Rubiac.).
angustifolius, Bartl. ex DC. Prod. iv. 436 = H. angustifolia.
ferrugineus, Bartl. ex DC. l. c. (= *Metabolos ferrugineus*).—Ins. Philipp.
laevigatus, Bartl. ex DC. l. c. = H. laevigata.
lineatus, Bartl. ex DC. l. c. 435 = H. vestita.
radicans, Bartl. ex DC. l. c. = H. radicans.

SCLEROCROTON, Hochst. in Flora, xxviii. (1845) 85 = **Excoecaria**, Linn. (Euphorb.).
ellipticus, Hochst. l. c.—Afr. austr.
integerrimus, Hochst. l c. = E. reticulata.
reticulatus, Hochst. l. c. = E. reticulata.

SCLEROLAENA, R. Br. Prod. 410 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 61.
DISSOCARPUS, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 75.
KENTROPSIS, Moq. Chenop. Enum. 83 (1840).
bicornis, *Lindl. in Mitch. Three Exped. Austral.* ii. 47.—Austral.
biflora, *R. Br. Prod.* 410.—Austral.
coriacea, Moq. in DC. Prod. xiii. II. 123 (sp. dub.).—Austral.

SCLEROLAENA:—
diacantha, *Benth. Fl. Austral.* v. 194.—Austral.
lanicuspis, *F. Muell. ex Benth. l. c.* 195.—Austral.
paradoxa, *R. Br. Prod.* 410.—Austral.
uniflora, *R. Br. l. c.*—Austral.
uniflora, Lindl. in Mitch. Journ. Trop. Austral. 72 = Kochia ciliata.

SCLEROLEIMA, Hook. f. in Hook. Lond. Journ. Bot. v. (1847) 444. t. 14 = **Abrotanella**, Cass. (Compos.).
forsterioides, Hook. f. l. c. = A. forsterioides.

SCLEROLEPIS, Cass. in Bull. Soc. Philom. (1816) 198. *COMPOSITAE*, Benth. & Hook. f. ii. 240.
verticillata, *Cass. in Dict. Sc. Nat.* xxv. 365; xlviii. 155.—Am. bor.

SCLEROLEPIS, Monn. Ess. Hierac. 81 (1829) = **Rodigia**, Spreng. (Compos.).
Kalmii, Monn. l. c. (= *Pachylepis Kalmii*).—Am. bor.

SCLEROLOBIUM, Vog. in Linnaea, xi. (1837) 395. *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
COSYMBE, Tul. in Arch. Mus. Par. iv. (1844) 168.
aureum, *Baill. Hist. Pl.* ii. 90; *Benth. in Mart. Fl. Bras.* xv. II. 50.—Bras.
chrysophyllum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 60.—Bras.
densiflorum, *Benth. in Mart. Fl. Bras.* xv. II. 51.—Bras.
denudatum, *Vog. in Linnaea*, xi. (1837) 396.—Bras.
guianense, *Benth. in Hook. Kew Journ.* ii. (1850) 237.—Guian.; Bras.
hippoboscum, Walp. Ann. ii. 441, sphalm. = seq.
hypoleucum, *Benth. in Hook. Kew Journ.* ii. (1850) 236.—Bras.
macrophyllum, Vog. in Linnaea, xi. (1837) 397 = paniculatum.
odoratissimum, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 48.—Bras.
paniculatum, *Vog. in Linnaea*, xi. (1837) 397.—Bras.
Poeppigianum, *Baill. Hist. Pl.* ii. 91.—Bras.
rubiginosum, Mart. Herb. Fl. Bras. n. 1147; ex Tul. in Arch. Mus. Par. iv. (1844) 123 = paniculatum.
rugosum, *Mart. l. c.* n. 1155; *ex Benth. in Hook. Kew Journ.* ii. (1850) 237.—Bras.
sericeum, Tul. in Arch. Mus. Par. iv. (1844) 125 = chrysophyllum.
tinctorium, *Benth. in Hook. Kew Journ.* ii. (1850) 236.—Bras.

SCLEROMITRION, Wight & Arn. Prod. 412 (1834). **Hedyotis**, Linn. (Rubiac.).
angustifolium, Benth. in Hook. Kew Journ. iv. (1852) 172 = H. tenelliflora.
approximatum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 137 = H. tenelliflora.
caespitosum, Miq. Fl. Ind. Bat. Suppl. 540 = H. pinifolia.
capitatum, Miq. Fl. Ind. Bat. ii. 186 = H. monocephala, *Miq.*
coronarium, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 136 = H. coronaria.
crassifolium, Miq. Fl. Ind. Bat. ii. 185 = H. crassifolia.
hispidum, Korth. in Nederl. Kruidk. Arch. ii. (1851) 155 = H. hispida.
nitidum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 137 = H. nitida.
paradoxum, Kurz, l. c. = H. paradoxa.
pauciflorum, Miq. Fl. Ind. Bat. ii. 187 = H. pauciflora.
rigidum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 136 = H. pinifolia.
rugosum, Kurz, l. c. 137 = H. rugosa.
sinense, Miq. in Journ. Bot. Néerl. i. (1861) 108.—China.
tenelliflorum, Korth. in Nederl. Kruidk. Arch. ii. (1851) 155 = H. tenelliflora.
tetrandrum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 136 = H. macrophylla.
tetraquetrum, Miq. Fl. Ind. Bat. ii. 186 = H. tenelliflora.

SCLEROMPHALOS, Griff. Notul. iv. 96; Ic. Asiat. Pl. t. 414, f. 1 (1854) = **Withania**, Pauq. (Solan.).

SCLERONEMA, Benth. in Journ. Linn. Soc. vi. (1862) 109 = **Catostemma**, Benth. (Malvac.).
Spruceana, Benth. l. c.—Bras.

SCLERONEMA, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. ii. (1864) 166 = **Xeronema**, Brongn. (Liliac.).
Moorii, Brongn. & Gris, l. c. 167 = X. Moorei.

SCLEROOLAENA, Baill. in Adansonia, x. (1872) 236. *CHLAENACEAE.*
XYLOOLAENA, Baill. Dict. ii. 2, in obs. (1879).
Richardi, *Baill. Hist. Pl.* iv. 224, in nota.—Madag.

SCLEROON, Benth. in Lindl. Bot. Reg. (1843) Misc. 65 = **Petitia**, Jacq. (Verben.).
oleinum, Lindl. l. c. = P. oleina.

SCLEROPHYLAX, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 18. *SOLANACEAE*, Benth. & Hook. f. ii. 912.
STERRHYMENIA, Griseb. in Goett. Abh. xix. (1874) 230.
Arnottii, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 20.—Reg. Argent.
Cynocrambe, *Griseb. in Goett. Abh.* xxiv. (1879) 268.—Reg. Argent.
Gilliesii, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 21.—Reg. Argent.
Lorentzianus, *O. Hoffm. in Linnaea*, xliii. (1880–82) 136.—Reg. Argent.
spinescens, *Miers, in Hook. Lond. Journ. Bot.* vii. (1848) 19.—Reg. Argent.

SCLEROPHYLLUM, Gaud. Fl. Helv. v. 47 (1829) = **Phaecasium**, Cass. (Compos.).
pulchrum, Gaud. l. c. 48 = P. lampsanoides.

SCLEROPOA, Griseb. Spicil. Fl. Rumel. ii. 431 (1844) = **Festuca**, Tourn. (Gramin.).
caspica, C. Koch, in Linnaea, xxi. (1848) 409 = Cutandia scleropoides.
dichotoma, Parl. Fl. Ital. i. 471 = F. dichotoma.
divaricata, Parl. l. c. 470; C. Koch, in Linnaea, xxi. (1848) 409 = F. divaricata.
hemipoa, Parl. l. c. 472 = F. hemipoa.
loliacea, Gren. & Godr. Fl. Fr. iii. 557 = Desmazeria loliacea.
maritima, Parl. Fl. Ital. i. 468 = Cutandia maritima.
memphitica, Boiss. Diagn. Ser. I. xiii. 62 = Cutandia scleropoides.
philistaea, Boiss. l. c. 60 = Cutandia philistaea.
procumbens, Parl. Fl. Ital. i. 474 = Glyceria procumbens.
pumila, Boiss. Diagn. Ser. I. xiii. 61.—Arab.
rigida, Griseb. Spicil. Fl. Rumel. ii. 431 = F. rigida.
stenostachya, Boiss. Fl. Orient. v. 638.—As. Min.

SCLEROPOGON, Phil. Sert. Mendoc. ii. 47 (1871). *GRAMINEAE*, Benth. & Hook. f. iii. 1177.
LESOURDIA, Fourn. in Bull. Soc. Bot. Fr. xxvii. (1880) 102. t. 3, 4.
brevifolius, *Phil. Sert. Mendoc.* ii. 48.—Chili.
Karwinskyanus, Benth. ex S. Wats. in Proc. Am. Acad. xviii. (1883) 181 = brevifolius.

SCLEROPTERIS, Scheidw. in Otto & Dietr. Allg. Gartenz. vii. (1839) 407 = **Cirrhaea**, Lindl. (Orchid.).
flava, Scheidw. l. c.—Bras.

SCLEROPUS, Schrad. Ind. Sem. Hort. Gotting. (1835).—Cf. Linnaea, xi. (1837) Litt. 89 = **Amarantus**, Linn.
amaranthoides, Schrad. ll. cc. = A. polygonoides.
crassipes, Moq. in DC. Prod. xiii. II. 271 = A. polygonoides.
squamulatus, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 162.—Ins. Galapag.
urceolatus, Anderss. l. c. = A. urceolatus?

SCLEROPYRON, Endl. Gen. Suppl. ii. 105 (1842) = seq.

SCLEROPYRUM, Arn. in Mag. Zool. & Bot. ii. (1838) 549. *SANTALACEAE*, Benth. & Hook. f. iii. 228.
HEYDIA, Dennst. Schluess. Hort. Malab. 30 (1818).
Wallichianum, *Arn. in Mag. Zool. & Bot.* ii. (1838) 550.—Ind. or.

SCLEROSCIADIUM, Koch, in DC. Coll. Mém. v. 43 (1829); Prod. iv. 140 (1830) = **Capnophyllum**, Gaertn. (Umbell.).
humile, Koch, ex DC. Coll. Mém. v. 43. t. 2.—Marocc.
nodiflorum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 59.—Afr. bor.

SCLEROSIA, Klotzsch, in Schomb. Faun. Fl. Gui. 1098 (1848), nomen. *OCHNACEAE.*
apiculata, Klotzsch, l. c.—Guian.

SCLEROSPERMA, G. Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 427. *PALMAE*, Benth. & Hook. f. iii. 919.
Manni, *H. Wendl. l. c.*—Afr. trop.

SCLEROSTYLIS, Blume, Bijdr. 133 (1825) = **Atalantia**, Correa (Rutac.).
amyridoides, M. Roem. Syn. Hesper. 43 = A. Roxburghiana.
Arnottiana, Wight, Illustr. Ind. Bot. i. 109 = A. ceylanica.
atalantioides, Wight, Ic. t. 71 = A. racemosa.
atalantioides, Wight & Arn. Prod. 93 = A. buxifolia.
buxifolia, Benth. in Hook. Kew Journ. iii. (1851) 326 = A. buxifolia.
ceylanica, Wight, Illustr. Ind. Bot. i. 109 = A. ceylanica.
Hindsii, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 327 = A. Hindsii.
lanceolata, Blume, Bijdr. 134 = Glycosmis simplicifolia.
macrophylla, Blume, l. c. 135 = Glycosmis pentaphylla?
nitida, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 249 = A. nitida.
ovalifolia, Wight, Illustr. Ind. Bot. i. 109 = A. racemosa.
parvifolia, Wight, Ic., in textu ad t. 71 = A. racemosa.
pentaphylla, Blume, Bijdr. 135.—Java.
racemosa, Wight, Illustr. Ind. Bot. 109 = A. racemosa.
rotundifolia, Thw. Enum. Pl. Zeyl. 46 = A. ceylanica.
Roxburghiana, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 513 = A. caudata.
Roxburghii, Wight, Ic. i. t. 72 = A. caudata.
spinosa, Blume, Bijdr. 134 (Atalantiae sp.).—Java.
timoriensis, M. Roem. Syn. Hesper. 43 = *pentaphylla*.
trifoliata, Blume, Bijdr. 134 = Glycosmis trifoliata.
venosa, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 327 = A. Hindsii.
zeylanica, Hook. f. Fl. Brit. Ind. i. 513 = A. ceylanica.

SCLEROTHAMNUS, R. Br. in Ait. Hort. Kew. ed. II. iii. 16 (1811) = **Eutaxia**, R. Br. (Legumin.).
microphyllus, R. Br. l. c. = E. empetrifolia.

SCLEROTHECA, A. DC. in DC. Prod. vii. 356 (1839). *CAMPANULACEAE*, Benth. & Hook. f. ii. 548.
arborea, *DC. l. c.* 357.—Ins. Societ.

SCLEROTHRIX, Presl, Symb. Bot. ii. 3. t. 53 (1833). *LOASEAE*, Benth. & Hook. f. i. 803.
ANCYROSTEMMA, Poepp. & Endl. Nov. Gen. et Sp. iii. 65. t. 272 (1845).
fasciculata, *Presl, Symb. Bot.* ii. 3. t. 53.—Mexic.

SCLEROXYLON, Bertol. in Mem. Acc. Sc. Bologn. viii. (1857) 237 (Misc. Bot. xviii. 15) (1858) = **Chrysophyllum**?, Linn. (Sapot.).
edule, Bertol. ll. cc.—Afr. trop.

SCLEROXYLON, Steud. Nom. ed. II. ii. 544 (1841) = seq.

SCLEROXYLUM, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 57; et in Enum. Hort. Berol. 249 (1809) = **Myrsine**, Linn.
canariense, Willd. l. c. 59 = M. canariensis.
Manglillo, Willd. l. c. = M. Manglilla.
melanophleum, Willd. l. c. 58 = M. melanophleos.
mite, Willd. l. c. = M. mitis.
racemosum, Willd. ex Roem. & Schult. Syst. iv. 803 = M. racemosa.
venulosum, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 58 = M. melanophleos.

SCOBEDIA, Labill. ex Steud. Nom. ed. I. 753 (1821). *LABIATAE*?
asperifolia, Labill. ex Steud. l. c. (Quid?).—Hab.?

SCOBIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Premna**?, Linn. (Verben.).

SCOLIOCHILUS, Reichb. f. Xenia Orch. ii. 118, in obs. (1872) = **Appendicula**, Blume (Orchid.).

SCOLIOPUS, Torr. in Pacif. Rail. Rep. iv. 145. t. 22 (1856). *LILIACEAE*, Benth. & Hook. f. iii. 832.
Bigelovii, *Torr. l. c.*—Calif.
Hallii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 272.—Am. bor. occ.

SCOLOBUS, Rafin. in Journ. Phys. lxxxix. (1819) 259 = **Thermopsis**, R. Br. (Legumin.).

SCOLOCHLOA, Link, Hort. Berol. i. 136 (1827) = **Graphephorum**, Desv. (Gramin.).
festucacea, Link, l. c. = G. festucaceum.

SCOLOCHLOA, Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 374, 528 = **Arundo**, Tourn. (Gramin.).
arundinacea, Mert. & Koch, l. c. 530 = A. Donax.
Donax, Gaud. Fl. Helvet. i. 202 = A. Donax.
spiculosa, F. Schmidt, Reise Amurl. 201.—Reg. Amur.

SCLEROSTEMMA, Schott, ex Roem. & Schult. Syst. iii. 64 (1818) = **Scabiosa**, Tourn. (Dipsac.).
agreste, Schott, l. c. 67 = Scab. gramuntia.
altissimum, Schott, l. c. 82 = Scab. africana.
argenteum, Schott, l. c. 79 = Scab. ucranica.
atropurpureum, Schott, l. c. 78 = Scab. maritima.
Biebersteinianum, Schott, l. c. 75 = Scab. micrantha.
caucasicum, Schott, l. c. 84 = Scab. caucasica.
Columbarium, Schott, l. c. 64 = Scab. Columbaria.
connatum, Schott, l. c. 84 = Scab. caucasica.
isetense, Schott, l. c. 83 = Scab. isetensis.
lucidum, Schott, l. c. 66 = Scab. lucida.
pyrenaicum, Schott, l. c. 73 = Scab. pyrenaica.
rupestre, Schott, l. c. 64 = Scab. isetensis.
Scopolii, Schott, l. c. 76 = Scab. ucranica.
siculum, Schott, l. c. 74 = Scab. ucranica.

SCOLODIA, Rafin. Sylva Tellur. 128 (1838) = **Cassia**, Tourn. (Legumin.).
viminea, Rafin. l. c. = C. viminea.

SCOLODRYS, Rafin. Alsog. Am. 29 (1838) = **Quercus**, Tourn. (Cupul.).
lyrata, Rafin. l. c. = Q. lyrata.
rigida, Rafin. l. c. (Quid?).—Am. bor.

SCOLOPACIUM, Eckl. & Zeyh. Enum. 59 (1835) = **Erodium**, L'Hérit. (Geran.).
incarnatum, Eckl. & Zeyh. l. c. = E. incarnatum.

SCOLOPIA, Schreb. Gen. 335 (1789). *BIXINEAE*, Benth. & Hook. f. i. 127.
ADENOGYRUS, Klotzsch, in Walp. Ann. iv. (1857) 226.
DASIANTHERA, Presl, Rel. Haenk. ii. 90. t. 66 (1831).
ERIUDAPHUS, Nees, in Eckl. & Zeyh. Enum. 271 (1835).
LIMONIA, Gaertn. Fruct. i. 278. t. 58. f. 4 (1789).
PHOBEROS, Lour. Fl. Cochinch. 317 (1790).
RHINANTHERA, Blume, Bijdr. 1121 (1826).
SCOPOLIA, Lam. Illustr. t. 860 (1798).
acuminata, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 251 = crenata.
Brownii, *F. Muell. Fragm.* iii. 11.—Austral.
chinensis, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 249 = crenata.
cochinchinensis, *Clos, l. c.* 253.—Cochinch.
coriacea, *Tul. l. c.* Sér. V. ix. (1868) 336.—Madag.
crassipes, *Clos, l. c.* Sér. IV. viii. (1857) 251 = crenata.
crenata, *Clos, l. c.* 250.—Ind. or.; China.
Dasyanthera, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 12.—Ins. Philipp.
elliptica, *Regel, Ind. Sem. Hort. Petrop. Suppl.* 1866 (1867) 35.—As. temp. or.
foetida, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 253 = Flacourtia sumatrana.
Gaertneri, Thw. Enum. Pl. Zeyl. 400 = Schreberi.

SCOLOPIA:—
Gerrardi, *Harv. in Harv. & Sond. Fl. Cap.* ii. 584.—Afr. austr.
lanceolata, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 252 = crenata.
macrophylla, Clos, l. c. 253 = rhinanthera.
Oldhamii, Hance, l. c. Sér. V. v. (1866) 206 = crenata.
Pseudo-crenata, Clos, l. c. Sér. IV. viii. (1857) 250 = crenata.
pusilla, Willd. Sp. Pl. ii. 981 = Schreberi.
rhinanthera, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 252.—Malaya.
Roxburghii, *Clos, l. c.* 250.—Burma; Malaya.
saeva, Hance, l. c. xviii. (1862) 217 = crenata.
Schreberi, *J. F. Gmel. Syst.* 793.—Zeylan.
Wightiana, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 252 = crenata.

SCOLOPOSPERMUM, Hemsl. Biol. Centr. Am. Bot. ii. 144 (1881), sphalm. = SCOLOSPERMUM, Less. = **Baltimora**, Linn. (Compos.).

SCOLOSANTHUS, Vahl, Eclog. Am. i. 11. t. 10 (1796). *RUBIACEAE*, Benth. & Hook. f. ii. 107.
ANTACANTHUS, Rich. ex DC. Prod. iv. 484 (1830).
ECHINODENDRUM, A. Rich. Fl. Cub. Fanerog. ii. 18 (1855).
crucifer, *Wright, in Sauv. Fl. Cub.* 67.—Cuba.
parviflorus, *Wright, l. c.*—Cuba.
triacanthus, DC. Prod. iv. 484 = Chomelia triacantha.
versicolor, *Vahl, Eclog. Am.* i. 11. t. 10.—Ind. occ.
Wrightii, *Wright, in Sauv. Fl. Cub.* 68.—Cuba.

SCOLOSPERMA, Rafin. Sylva Tellur. 111 (1838) = **Cleome**, Linn. (Cappar.).
dendroides, Rafin. l. c. = C. dendroidea.

SCOLOSPERMUM, Less. in Linnaea, v. (1830) 152 = **Baltimora**, Linn. (Compos.).
baltimoroides, Less. l. c. t. 2. f. 19–31 = B. Scolospermum.
Fougerouxiae, DC. Prod. v. 509 = B. recta, B. Scolospermum.

SCOLYMANTHUS, Willd. ex DC. Prod. vii. 63 (1838) = **Perezia**, Lag. (Compos.).
cnicoides, Willd. l. c. 64 = P. multiflora.

SCOLYMUS, Tourn. ex Linn. Gen. ed. I. 237 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 504.
MYSCOLUS, Cass. in Dict. Sc. Nat. xxv. 60 (1822).
angiospermus, Gaertn. Fruct. ii. 355. t. 157 = hispanicus.
annuus, Gerard, ex Steud. Nom. ed. I. 753 = maculatus.
congestus, Lam. Fl. Fr. ii. 116 = hispanicus.
giganteus, Roem. Hort. Turic. ex Steud. Nom. ed. II. ii. 544 (Quid?).—Hab.?
grandiflorus, *Desf. Fl. Atlant.* ii. 240.—Afr. bor.
gymnospermus, Gaertn. Fruct. ii. 355 = maculatus.
hispanicus, *Linn. Sp. Pl.* 813.—Reg. Mediterr.
maculatus, *Linn. l. c.*—Reg. Mediterr.
paniculatus, *Ucria, in Roem. Arch.* i. 1. (1796) 70.—Sicil.
pectinatus, Cass. in Dict. Sc. Nat. xxxiv. 87 = maculatus.
perennis, Gerard, ex Steud. Nom. ed. I. 753 = hispanicus.
vivax, Sauvages, ex DC. Prod. vii. 76 = hispanicus.

SCOPARIA, Linn. Syst. ed. VI. 87 (1748). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 959.
KREIDEK, Adans. Fam. ii. 225 (1763).
annua, *Cham. & Schlecht. in Linnaea*, vi. (1831) 375.—Mexic.
arborea, Linn. f. Suppl. 125 = Chilianthus oleaceus.
australis, Sieber, ex Cham. in Linnaea, viii. (1833) 22 = Teucrium corymbosum.
dulcis, *Linn. Sp. Pl.* 116.—Reg. trop.
elliptica, *Cham. in Linnaea*, viii. (1833) 21.—Bras.
ericacea, *Cham. & Schlecht. l. c.* ii. (1827) 604.—Bras.
flava, *Cham. & Schlecht. l. c.* 603.—Reg. Argent.
fruticosa, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 286.—Hab.?
gypsophyloides, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 394.—China.

SCOPARIA:—
multifida, G. Don, Gen. Syst. iv. 648, sphalm. = seq.
pinnatifida, Cham. in Linnaea, viii. (1833) 22 = flava.
plebeia, Cham. & Schlecht. l. c. ii. (1827) 605 = flava.
procumbens, Jacq. Enum. Pl. Carib. 12 = dulcis.
ternata, Forsk. Fl. Aegypt. Arab. 31 = dulcis.

SCOPOLA, Jacq. Obs. i. 32. t. 20 (1764) = **Scopolia**, Jacq. (Solan.).

SCOPOLIA, Adans. Fam. ii. 419 (1763) = **Ricotia**, Linn. (Crucif.).

SCOPOLIA, Forst. Char. Gen. 139. t. 70 (1776) = **Griselinia**, Forst. f. (Celastr.).
lucida, Forst. l. c. 140 = G. lucida.

SCOPOLIA, Jacq. Obs. i. 32. t. 20 (1764) (Scopola); Link, Enum. Hort. Berol. i. 178 (1821). *SOLANACEAE*, Benth. & Hook. f. ii. 902.
ANISODUS, Link & Otto, Ic. Pl. Sel. 77. t. 35 (1824).
SCOPOLINA, Schult. Oestr. Fl. ed. II. i. 335 (1814).
WHITLEYA, D. Don, in News Lit. & Fashion, iii. 108; ex Sweet, Brit. Flow. Gard. Ser. I. t. 125 (1825).
atropoides, Bercht. & Presl, Rostl. i. Solanac. 12 = carniolica.
Boreana, Dun. in DC. Prod. xiii. 553 = Hyoscyamus muticus.
carniolica, *Jacq. Obs.* i. 32. t 20.—Europ.
Datora, Dun. in DC. Prod. xiii. 553 = Hyoscyamus muticus.
Hladnikiana, Fleischm. ex Nym. Consp. 524 = carniolica.
japonica, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 57.—Japon.
lurida, *Dun. in DC. Prod.* xiii. 555.—Reg. Himal.
mutica, Dun. l. c. 552 = Hyoscyamus muticus.
orientalis, Dun. l. c. 554 = Physochlaina orientalis.
physaloides, *Dun. in DC. Prod.* xiii. 554.—Sibir. altaic.
praealta, Dun. l. c. = Physochlaina praealta.
Pseudo-physaloides, Dun. l. c. 556 = Hyoscyamus Pseudo-physaloides.
tangutica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 508.—China.
trichotoma, Moench, Meth. 462 = carniolica.

SCOPOLIA, Lam. Illustr. t. 860 (1798), sphalm. = **Scolopia**, Schreb. (Bixin.).

SCOPOLIA, Linn. f. Suppl. 60 (1781) = **Daphne**, Tourn. (Thymel.).
composita, Linn. f. l. c. = D. pendula.
involucrata, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 357 = D. involucrata.
pendula, Rafin. Fl. Tellur. iv. 106 = D. pendula.

SCOPOLIA, Sm. Ic. Ined. ii. t. 34 (1790) = **Toddalia**, Juss. (Rutac.).
aculeata, Sm. l. c. = T. aculeata.
angustifolia, Spreng. Syst. i. 779 = T. aculeata.
inermis, Willd. Sp. Pl. i. 1116 = T. paniculata.
lanceolata, Spreng. Syst. i. 779 = T. lanceolata.
megapotamica, Spreng. Syst. iv. Cur. Post. 91 = T. megapotamica.
micracantha, Blume, Bijdr. 1167 = T. micrantha.
micrantha, Steud. Nom. ed. II. ii. 545 = praec.
nitida, Willd. ex Spreng. Syst. i. 779 = T. Willdenowii.
paniculata, Spreng. l. c. = T. paniculata.
venosa, Spreng. l. c. iv. Cur. Post. 91 = T. venosa.

SCOPOLINA, Schult. Oestr. Fl. ed. II. i. 335 (1814) = **Scopolia**, Jacq. (Solan.).
atropoides, Schult. l. c. 383 = Scopolia carniolica.
carniolica, Schur, Enum. Pl. Transs. 479 = Scopolia carniolica.
Hladnikiana, Frey. ex Koch, Syn. Fl. Germ. ed. II. 585 = Scopolia carniolica.
viridiflora, Frey. ex Koch, l. c. = Scopolia carniolica.

SCOPULARIA, Lindl. Bot. Reg. sub t. 1701 (1834) = **Holothrix**, Rich. (Orchid.).
Burchelli, Lindl. Gen. et Sp. Orch. 304 = H. Burchellii.
grandiflora, Sond. in Linnaea, xix. (1847) 79 = H grandiflora.
secunda, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 207 = H. Scopularia.

SCORBION, Rafin. Fl. Tellur. iii. 85 (1836) = **Teucrium**, Tourn. (Labiat.).
spinosum, Rafin. l. c. = T. spinosum.

SCORDIUM, Gilib. Fl. Lituan. i. 95 (1781); Cav. Ic. i. 19. t. 31 (1791) = **Teucrium**, Tourn. (Labiat.).
altaicum, Gilib. l. c. = T. Scordium.
officinale, Gueldenst. ex Ledeb. Fl. Ross. iii. 443 = T. Scordium.
palustre, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138 = T. Scordium.
scordioides, Fourr. l. c. = T. scordioides.
spinosum, Cav. Ic. i. 19. t. 31 = T. spinosum.

SCORIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Carya**, Nutt. (Jugland.).
alba, Rafin. l. c. = C. alba.
globosa, Rafin. l. c.—Am. bor.
mucronata, Rafin. l. c. = C. sulcata.
pyriformis, Rafin. l. c.—Am. bor.
tomentosa, Rafin. l. c. = C. tomentosa.

SCORIAS, Rafin. ex Endl. Gen. 1126 (1841) = praec.

SCORODENDRON, Blume, Rumphia, iii. 149 (1847). *SAPINDACEAE*, Benth. & Hook. f. i. 403.
pallens, *Blume, l. c.*—Ins. Timor.

SCORODOCARPUS, Becc. in Nuov. Giorn. Bot. Ital. x. (1877) 274. *OLACINEAE*.
borneënsis, *Becc. l. c.*—Borneo.

SCORODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160 = **Allium**, Tourn. (Liliac.).
moschatum, Fourr. l. c. = A. moschatum.

SCORODONIA, Adans. Fam. ii. 188 (1763) = **Teucrium**, Linn. (Labiat.).
Arduini, Reichb. Fl. Germ. Excurs. 315 = T. Arduini.
betonicaefolia, Moench, Meth. Suppl. 133 = T. betonicum.
Botrys, Ser. Bull. Bot. (1830) 317 = T. Botrys.
cordata, Moench, Meth. 385 = T. massiliense.
Fontanesiana, Ser. Bull. Bot. (1830) 315 = T. Pseudo-scorodonia.
heteromalla, Moench, Meth. 384 = T. Scorodonia.
lancifolia, Moench, l. c. = T. lancifolium.
macrophylla, Moench, Meth. Suppl. 133 = T. canadense.
massiliensis, Link, Handb. i. 458 = T. massiliense.
Salviastrum, Link, l. c. = T. Pseudo-scorodonia.
sicula, Rafin. Fl. Tellur. iii. 85 = T. Scorodonia.
solitaria, Stokes, Bot. Mat. Med. iii. 374 = T. Scorodonia.
spicata, Moench, Meth. 385 = T. hyrcanicum.
spinosa, Link, Handb. i. 458 = T. spinosum.
sylvestris, Link, l. c. = T. Scorodonia.
trivialis, Rafin. Fl. Tellur. iii. 85 = T. Scorodonia?

SCORODOSMA, Bunge, Ind. Sem. Hort. Dorpat. (1846) 3; et in Mém. Sav. Etr. Pétersb. vii. (1854) 309 = **Ferula**, Tourn. (Umbell.).
foetidum, Bunge, ll. cc. = F. Assa-foetida.

SCORODOXYLUM, Nees, in Benth. Pl. Hartw. 236. (1846) = **Ruellia**, Linn. (Acanth.).
Hartwegianum, Nees, l. c.—N. Granat.
terminale, Nees, ex Benth. Bot. Voy. Sulph. 145.—Ecuador.

SCORPAENA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 (Quid?).

SCORPIANTHES, Rafin. Sylva Tellur. 89 (1838) = **Heliotropium**, Tourn. (Borag.).
lineatum, Rafin. l. c. = H. lineatum.

SCORPIOIDES, Gilib. Fl. Lituan. i. 20 (1781) = **Myosotis**, Linn. (Borag.).
glaber, Gilib. l. c. = M. palustris.
hirsutus, Gilib. l. c. = M. intermedia.

SCORPIOIDES, Tourn. ex Adans. Fam. ii. 328 (1763) = **Scorpiurus**, Linn. (Legumin.).
muricata, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 370 = Scorpiurus muricata.
sulcata, Medic. l. c. = Scorpiurus sulcata.
vermiculata, Medic. l. c = Scorpiurus vermiculata.

SCORPIURUS, Linn. Syst. ed. I (1735); Gen. ed. I. 226 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 508.
SCORPIUS, Loisel. Fl. Gall. 468 (1806).
SCORPIOIDES, Tourn. ex Adans. Fam. ii. 328 (1763).
acutifolia, Viv. Fl. Lib. Spec. 43. t. 19. f. 4 = subvillosa.
echinata, Lam. Encyc. i. 726 = muricata, subvillosa, sulcata.
echinata, A. Rich. Tent. Fl. Abyss. i. 200 = sulcata.
laevigata, Sibth. & Sm. Fl. Graec. viii. 13. t. 718 = muricata.
muricata, *Linn. Sp. Pl.* 745.—Reg. Mediterr.
pinnata, *Mill. Gard. Dict.* ed. VIII. n. 5.—Europ. austr.
purpurea, *Desf. Fl. Atlant.* ii. 174.—Algeria.
subvillosa, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. i. 200 = sulcata.
subvillosa *Linn. Sp. Pl.* 745.—Reg. Mediterr.
sulcata, *Linn. l. c.*—Reg. Mediterr.
sulcata, Sibth. & Sm. Fl. Graec. viii. 14. t. 719 = subvillosa.
vermiculata, *Linn. Sp. Pl.* 744.—Reg. Mediterr.

SCORPIUS, Loisel. Fl. Gall. 468 (1806) = praec.

SCORPIUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 369 = **Coronilla**, Tourn. (Legumin.).
scorpioides, Medic. l. c. = C. scorpioides.

SCORPIUS, Moench, Meth. 134 (1794) = **Genista**, Linn. (Legumin.).
anglicus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 428 = G. anglica.
Clusii, Moench, Meth. 134 = G. germanica.
falcatus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 429 = G. falcata.
hispanicus, Pourr. ex Willk. & Lange, l. c. 427 = G. hispanica.
juniperiformis, Pourr. ex Willk. & Lange, l. c. = G. hispanica.
parviflorus, Pourr. ex Willk. & Lange, l. c. 425 = G. triacanthos.
sericeus, Pourr. ex Willk. & Lange, l. c. 426 = G. hirsuta.
serpyllifolius, Pourr. ex Willk. & Lange, l. c. 429 = G. falcata.
spinosus, Moench, Meth. 134 = G. germanica.
vulgaris, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 430 = G. Scorpius.

SCORZONELLA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 426 = **Microseris**, D. Don (Compos.).
glauca, Nutt. l. c. = M. leptosepala.
laciniata, Nutt. l. c. = M. laciniata.
leptosepala, Nutt. l. c. = M. leptosepala.
nutans, Hook. Lond. Journ. Bot. vi. (1847) 253 = M. nutans.
sylvatica, Benth. Pl. Hartw. 320 = M. sylvatica.

SCORZONERA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 241 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 531.
ACHYROSERIS, Sch. Bip. in Nov. Act. Nat. Cur. xxi. (1845) 165.
FLEISCHERIA, Steud. & Hochst. ex Endl. Gen. 497 (1838).
GELASIA, Cass. in Bull. Soc. Philom. (1818) 33.
LASIOSPERMUM, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 34.
LASIOSPORA, Cass. in Dict. Sc. Nat. xxv. 306 (1822).
PODOSPERMUM, DC. Fl. Fr. iv. 61 (1805).
acanthoclada, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 333.—Turkestan.
acaulis, Willd. Sp. Pl. iii. 1508 = Taraxacum officinale.
acrolasia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 378.—Turkest.; Beluchist.
acuminata, *Boiss. & Bal. Diagn.* Ser. II. vi. 119.—As. Min.
africana, *Poir. Encyc. Suppl.* v. 114.—Afr. trop.
albicans, *Coss. Notes Crit.* 119.—Hispan.
albicaulis, *Bunge, Enum. Pl. Chin. Bor.* 40.—China.
alexandrina, *Boiss. Fl. Orient.* iii. 760.—Afr. bor.
alpina, Hoppe, in Sturm, Deutschl. Fl. Heft 51 = aristata.
alpina, Pollini, Viagg. Lag. Gard. 100 = Candollei.

SCORZONERA :—
ambigua, DC. Prod. vii. 126 = mollis.
ammophila, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 378.—Turkest.
ancyritana, DC. Prod. vii. 124 = tomentosa.
angustifolia, Asso, Syn. Stirp. Arag. 108 = graminifolia.
angustifolia, DC. Prod. vii. 120 = austrica, Candollei.
angustifolia, *Linn. Sp. Pl.* 791.—Europ. austr.
angustifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 128. t. 122 = villosa.
angustifolia, [Weber, in] Wigg. Prim. Fl. Holsat. 55 = humilis.
angustifolia, Wohlleb. ex Wallr. Ann. Bot. 96 = hispanica.
araneosa, *Sibth. & Sm. Fl. Graec.* viii. 62. t. 785.—Graecia.
araneosa, Urv. in Mém. Soc. Linn. Par. i. (1822) 354 = mollis.
argyrea, *Boiss. Diagn.* Ser. I. iv. 26.—Caria.
aristata, *Ram. ex DC. Fl. Fr.* iv. 922.—Europ. austr.
armeniaca, *Boiss. Fl. Orient.* iii. 760.—Armen.
aspera, Desf. in Ann. Mus. Par. i. (1802) 133. t. 9 = Hymenonema Tournefortii.
asperrima, Willd. Sp. Pl. iii. 1507 = Leontodon asperrimus.
asphodeloides, Wallr. Ann. Bot. 95 = hispanica.
astrachanica, DC. Prod. vii. 118 = mollis.
Aucheri, Boiss. & Huet, Diagn. Ser. II. iii. 92 = suberosa.
Aucheriana, *DC. Prod.* 125.—Mesopot.
austriaca, Balb. Misc. Alt. 24 = Candollei.
austriaca, *Willd. Sp. Pl.* iii. 1498.—Europ.; As. bor.
autumnalis, Lam. Fl. Fr. ii. 82 = Leontodon autumnalis.
boetica, *Boiss. Voy. Bot. Espagne*, ii. 382.—Hispan.
bohemica, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 55 = humilis.
brevicaulis, Vahl, Symb. Bot. ii. 88 = coronopifolia.
bulbipes, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 759.—Persia.
buphthalmoides, DC. Prod. vii. 121 = Tragopogon persicus.
bupleuroides, *D. Don, Prod. Fl. Nep.* 162 (Quid ?).—Nepal.
caerulea, Saut. ex Reichb. Fl. Germ. Excurs. 855 = Lactuca perennis.
caespitosa, Pomel, Nouv. Mat. Fl. Atl. 266 = pygmaea.
calcitrapifolia, *Vahl, Symb. Bot.* ii. 87.—Europ.
callosa, Moris, Elench. Stirp. Sard. i. 29 = villosa.
calvescens, DC. Prod. vii. 123 = laciniata.
calyculata, *Boiss. Diagn.* Ser. I. xi. 42.—Persia.
Candollei, *Vis. Fl. Dalm.* ii. 106.—Dalmatia.
capensis, Thunb. Prod. Pl. Cap. 141 = Lactuca capensis.
caricifolia, Pall. Reise, iii. 756 = parviflora.
cariensis, *Boiss. Diagn.* Ser. I. iv. 24.—Caria.
cenopleura, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 378.—Turkestan.
chondrilloides, *Pourr. ex Willd. Sp. Pl.* iii. 1505.—Hispan.
chondrilloides, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 54.—Turkestan.
ciliata, Forsk. Fl. Aegypt. Arab. 143 = Picridium tingitanum.
cilicica, *Boiss. Diagn.* Ser. II. iii. 94.—Cilicia.
cinerea, *Boiss. l. c.* Ser. I. xi. 44.—As. Min.; Persia.
circinnata, Pall. Reise, iii. 677 = pusilla.
Clusii, Asso, Syn. Stirp. Arag. 108 = humilis.
Columnae, Guss. Ind. Sem. Hort. Boccad. (1825); Pl. Rar. 321. t. 54 = villosa.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 (Quid ?).—Java.
coronopifolia, *Desf. Fl. Atlant.* ii. 220. t. 44.—Afr. bor.
crepioides, Poir. Encyc. vii. 25 = Sonchus crepioides.
cretica, *Willd. Sp. Pl.* iii. 1504.—Ins. Creta.
crispa, Bieb. Fl. Taur. Cauc. ii. 234 = austriaca.
crispatula, Boiss. Voy. Bot. Espagne, ii. 741 = hispanica.
crocifolia, *Sibth. & Sm. Fl. Graec.* viii. 63. t. 785.—Graecia.
Cypria, Kotschy, in Unger & Kotschy, Ins. Cyp. 253 = parviflora.
dalmatica, Host, Fl. Austr. ii. 391 = villosa.
decumbens, Guss. Fl. Sic. Syn. ii. 386 = calcitrapifolia.
deliciosa, *Guss. ex DC. Prod.* vii. 118.—Sicilia.
dentata, Kit. in Linnaea, xxxii. (1863) 409 = hispanica.
denticulata, Lam. Fl. Fr. ii. 82 = hispanica.

SCORZONERA :—
dichotoma, Vahl, Symb. Bot. ii. 89 = Launaea nudicaulis.
divaricata, Auch. ex DC. Prod. vii. 125 = Aucheriana.
divaricata, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 200.—Mongol.
edulis, Moench, Meth. 548 = hispanica.
elata, *Boiss. Diagn.* Ser. I. iv. 25.—As. Min.
elongata, Willd. Sp. Pl. iii. 1508 = Hymenonema Tournefortii.
ensifolia, *Bieb. Fl. Taur. Cauc.* ii. 235.—Rossia; As. bor.
eriocarpa, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 225 = hirsuta.
eriophora, *DC. Prod.* vii. 125.—As. Min.; Mesopot.
eriosperma, Auch. ex DC. Prod. vii. 125 = ramosissima.
eriosperma, *Bieb. Fl. Taur. Cauc.* ii. 236; *Beschr. Casp.* 117. n. 28.—Reg. Caucas.
eriosperma, Gouan, Illustr. 52 = hirsuta.
eriosperma, Willd. ex Ledeb. Fl. Ross. ii. 783 = mollis.
fasciata, Pomel, Nouv. Mat. Fl. Atl. 9 = coronopifolia.
filifolia, *Boiss. Fl. Orient.* iii. 774.—Reg. Caucas.
fistulosa, *Brot. Fl. Lusit.* i. 329.—Lusitan.
floccosa, *Boiss. Fl. Orient.* iii. 758.—As. Min.
gageoides, *Boiss. Diagn.* Ser. II. iii. 95.—Beluchist.
Gebleri, Bess. ex DC. Prod. vii. 120 = austriaca.
glabra, *Rupr. in Beitr. Pfl. Russ. Reich.* ii. 11.—Reg. Ural.
glandulosa, Pourr. ex DC. Prod. vii. 183 (= *Zollikofera pumila*).—Hispan.; Arab.
glastifolia, Hegetschw. Fl. Schw. ii. 157 = humilis.
glastifolia, Willd. Sp. Pl. iii. 1499 = hispanica.
glauca, Salisb. Prod. 179 = Picridium tingitanum.
graeca, Hort. ex Steud. Nom. ed. II. ii. 546 = Hymenonema graecum.
graminea, Krock. Fl. Siles. ii. 11. 263 = tomentosa.
graminifolia, All. Fl. Pedem. i. 228 = aristata.
graminifolia, Auch. ex DC. Prod. vii. 121 = Tragopogon persicus.
graminifolia, Bory & Chaub. Nouv. Fl. Pélop. 53 = crocifolia.
graminifolia, Griseb. Spicil. Fl. Rumel. ii. 263 = mollis.
graminifolia, Hoffm. Deutschl. Fl. ed. I. 272 = hispanica.
graminifolia, Lag. ex DC. Prod. vii. 122 = macrocephala.
graminifolia, *Linn. Sp. Pl.* 791 (sp. confusa).—Sibir.; Lusitan.
graminifolia, Portenschl. ex Vis. Fl. Dalm. ii. 107 = villosa.
graminifolia, Schur, Enum. Pl. Transs. 362 = austriaca.
graminifolia, Sibth. & Sm. Fl. Graec. viii. 62. t. 784 = elata.
graminifolia, Tausch, ex Nym. Consp. 464 = parviflora.
grandiflora, Lapeyr. Hist. Abr. Pl. Pyr. 457 = aristata.
halophila, Fisch. & Mey. in DC. Prod. vii. 122 = parviflora.
hemilasia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 377.—Turkestan.
heterophylla, Pomel, Nouv. Mat. Fl. Atl. 265 = mollis.
hirsuta, *Linn. Mant.* ii. 278.—Europ. austr.
hirsuta, Pett. ex Nym. Consp. 465 = villosa.
hispanica, *Linn. Sp. Pl.* 791.—Europ. austr.
hispida, Bieb. Casp. App. 210 = Leontodon asperrimus.
hispida, *Forsk. Fl. Aegypt. Arab.* 215.—Galatia.
hispidula, Spreng. Syst. iii. 667 = Leontodon hispidulus.
Hoppeana, Sieber, ex Reichb. Fl. Germ. Excurs. 276 = aristata.
humifusa, *Darracq, ex Woods, in Journ. Linn. Soc.* ii. (1858) 120.—Hispan.
humilis, Jacq. Fl. Austr. t. 36 = austriaca.
humilis, *Linn. Sp. Pl.* 790.—Europ.
inaequiscapa, *Boiss. ex Boiss. Fl. Orient.* iii. 762.—Armenia.
incisa, *DC. Prod.* vii. 119.—Cappadoc.
intermedia, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 377 = hemilasia.
intermedia, Guss. Fl. Sic. Syn. ii. 387 = laciniata ?
intricata, *Boiss. Diagn.* Ser. I. vii. 9.—Persia.
ispahanica, *Boiss. l. c.* 7.—Persia.
Jacquiniana, *Boiss. Fl. Orient.* iii. 757.—Europ. or.; As. Min.
julia, Vest, in Flora, iii. (1820) 6 = villosa.
juncea, *Hochst. ex Steud. Nom.* ed. II. ii. 366, in syn.—Arab.

SCORZONERA :—
Kotschyi, *Boiss. Fl. Orient.* iii. 780.—Syria; Mesopot.
Kotschyi, Boiss. l. c. 762 = mollis.
kurdica, Boiss. & Noé, Diagn. Ser. II. iii. 93 = papposa.
lacera, *Boiss. & Bal. l. c.* v. 116.—Cilicia.
laciniata, Jacq. Fl. Austr. t. 356 = Jacquiniana.
laciniata, *Linn. Sp. Pl.* 791.—Reg. Mediterr. et Caucas.
laciniata, Vahl, ex DC. Prod. vii. 123 = coronopifolia.
lanata, *Bieb. Fl. Taur. Cauc.* ii. 237.—Graecia; Oriens.
lanata, Schrank, Baier. Fl. ii. 306 = humilis.
lanigera, *Fisch. & Mey. in Tchihat. Asie Min. Bot.* ii. 366.—As. Min.
lanuginosa, Baumg. Enum. Stirp. Transs. iii. 9 = humilis.
lasiospora, Boiss. & Bal. Diagn. Ser. II. vi. 118 = nervosa.
latifolia, DC. Prod. vii. 124 = nervosa.
latifolia, Vis. Fl. Dalm. ii. 106. t. 5. f. 2 = austriaca.
Lawrencii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 124 = Microseris Forsteri.
libanotica, *Boiss. Diagn.* Ser. I. xi. 43.—Syria.
limnophila, *Boiss. l. c.* vii. 7.—Persia.
linarifolia, *Wender. Fl. Hass.* 265.—Europ.
lorea, Griseb. Spicil. Fl. Rumel. ii. 263 = Jacquiniana.
mackmeliana, *Boiss. Diagn.* Ser. I. xi. 44.—Syria.
macrocephala, *DC. Prod.* vii. 122.—Hispan.
macrophylla, *Fisch. & Mey. in Schrenk, Enum. Pl. Nov.* 39, in adnot.—Sibir.
macrorhiza, Schleich. Cat. Pl. Helv. ed. IV. 32 = humilis.
macrosperma, *Turcz. in DC. Prod*, vii. 121.—Dahur.
Marschalliana, C. A. Mey. ex Bong. & Mey. Verz. Pfl. Saisang-Nor, 44 = stricta.
messeniaca, Bory & Chaub. Nouv. Fl. Pélop. 53. t. 30 = laciniata.
mollis, *Bieb. Fl. Taur. Cauc.* iii. 522.—Graec.; Oriens.
mollis, Fisch. & Mey. ex DC. Prod. vii. 117 = mollis.
montana, Mutel, Hist. Pl. Dauph. ii. 268 = hispanica.
multifida, *Link, ex Steud. Nom.* ed. II. ii. 546.—Hab. ?
multiscapa, *Boiss. Diagn.* Ser. I. xi. 41.—Oriens.
muricata, Balb. Misc. 25 = laciniata.
nana, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 138.—As. Min.; Persia.
nervosa, Lam. Fl. Fr. ii. 81 = humilis.
nervosa, *Trevir. Del. Sem. Hort. Bonn.* (1833).—*Cf. Linnaea*, x. (1836) *Litt.* 73.—Armen.; Persia.
nivalis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 765.—Kurdist.
nova, Desf. ex Walp. Rep. vi. 340 = Hypochoeris glabra.
octangularis, Hort. ex Willd. Sp. Pl. iii. 1506 = laciniata.
octangularis, Roth, ex Steud. Nom. ed. II. ii. 546 = Jacquiniana.
odontopleura, Boiss. Fl. Orient. iii. 766 = incisa.
orientalis, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 249 = Crepis albida.
orientalis, Linn. Syst. ed. X. 1191 = Picridium tingitanum.
ovata, *Trautv. in Act. Hort. Petrop.* i. (1871–72) 275.—Turkest.
paniculata, Boiss. Diagn. Ser. I. vii. 8 = nervosa.
papposa, *DC. Prod.* vii. 119.—Syria; Persia.
paradoxa, Fisch. & Mey. ex DC. l. c. = picridioides.
parviflora, *Jacq. Fl. Austr.* iv. 3. t. 305.—Europ.; As. bor. et occ.
paucifida, Lam. Fl. Fr. ii. 83 = laciniata.
persepolitana, *Boiss. Fl. Orient.* iii. 760.—Persia.
persica, Boiss. l. c. 777 = psychrophila.
persica, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 139.—Persia.
petiolaris, Lapeyr. Hist. Abr. Pl. Pyr. 120 = laciniata.
phaeopappa, Boiss. Fl. Orient. iii. 764 = multiscapa.
picridioides, *Boiss. Diagn.* Ser. I. vii. 6.—Persia; Beluchist.
picroides, Linn. Sp. Pl. 792 = Picridium vulgare.
pinifolia, Gouan, Illustr. 53 = macrocephala.
pinnatifida, Lour. Fl. Cochinch. 479 = africana.
pinnatifida, Michx. Fl. Bor. Am. ii. 89 = Pyrrhopappus carolinianus.
plantaginea, Gaud. Fl. Helv. v. 20 = seq.
plantaginifolia, Schleich. Cat. Pl. Helv. ed. III. 27 = humilis.
plurifida, Lam. Fl. Fr. ii. 83 = calcitrapifolia.

SCORZONERA :—
Prescottii, Comp. ex Boiss. Fl. Orient. iii. 770 = austriaca.
psychrophila, *Boiss. & Haussk. ex Boiss. l. c.* 777.—Persia.
pubescens, *DC. Prod.* vii. 122.—Sibir. altaic.
pumila, Cav. Ic. ii. 19 (= *Zollikoferia pumila*).—Hispan.; Arab.
purpurea, *Linn. Sp. Pl.* 791.—Europ.; As. bor.
pusilla, *Pall. Reise*, ii. 744.—Rossia; As. occ.
pygmaea, *Sibth. & Sm. Fl. Graec.* viii. 61. t. 783.—Bithyn.
quitensis, Humb. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 120 (= *Achyrophora quitensis*).—Ecuador.
racemosa, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 332.—Turkest.
radians, *Fisch. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 18.—As. bor.
radiata, Fisch. ex DC. Prod. vii. 122 = praec.
radicosa, *Boiss. Diagn.* Ser. I. xi. 43.—Persia.
ramosissima, *DC. Prod.* vii. 125.—Persia.
resedifolia, Linn. Sp. Pl. 1198 = Launaea chondrilloides.
resedifolia, Retz. Obs. iii. 42 = laciniata.
rigida, *Auch. ex DC. Prod.* vii. 123.—As. Min.
rosea, Waldst. & Kit. Pl. Rar. Hung. ii. 127. t. 121 = purpurea.
Roylei, DC. Prod. vii. 125 = divaricata.
rumicifolia, *Hort. Vindob. ex Link, Enum. Hort. Berol.* ii. 278.—Hab.?
runcinata, *Pall. Reise Suedl. Statth. Russ. Reich.* i. 798.—Rossia.
sardoa, Moris, ex Spreng. Syst. iv. Cur. Post. 347 = villosa.
sativa, Gater. Desc. Pl. Montaub. 136 = hispanica.
scapigera, Forst. f. Prod. 91 = Microseris Forsteri.
scoparia, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 774 = mackmeliana.
Scowitzii, Steud. Nom. ed. II. ii. 546, in syn. = *Szowitzii*.
Seidlitzii, *Boiss. Fl. Orient.* iii. 775.—Armenia.
semicana, *DC. Prod.* vii. 119.—Assyria.
sericea, *Auch. ex DC. l. c.* 123.—Cappadoc.
serrulata, Viv. Fl. Lib. Spec. 49. t. 17. f. 4 = undulata, trachysperma.
sessilis, Humb. ex Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 120 (= *Achyrophora quitensis*.).—Ecuador.
stenocephala, Boiss. Diagn. Ser. I. vii. 6 = mollis.
Stewartii, *Hook. f. Fl. Brit. Ind.* iii. 419.—Ind. bor. occ.
stricta, DC. Prod. vii. 122 = ensifolia.
stricta, *Hornem. Hort. Hafn.* ii. 750.—Reg. Cauc.; Sibir.
stylosa, Pers. Syn. ii. 361 = Hymenonema Tournefortii.
subaphylla, *Boiss. Diagn.* Ser. I. vii. 8.—Persia.
suberosa, *C. Koch, in Linnaea*, xvii. (1843) 275.—Armenia.
subulata, Lam. Fl. Fr. ii. 81 = laciniata.
sylvatica, Benth. Pl. Hartw. 388 (Index) = Microseris sylvatica.
syriaca, Boiss. & Blanch. Diagn. Ser. II. iii. 93 = mollis.
Szowitzii, DC. Prod. vii. 117 = mollis.
Taraxaci, Roth, Beitr. Bot. i. (1782) 120 = Taraxacum palustre.
taraxacifolium, Jacq. Ic. Pl. Rar. t. 160 = Hypochoeris glabra.
taurica, Bieb. Fl. Taur. Cauc. ii. 234 = hispanica.
taurica, Janka, ex Nym. Consp. 463 = stricta.
Tenorii, Presl, Delic. Prag. 116 = laciniata.
tetuanensis, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 551 = deliciosa.
tenuifolia, Bertol. Fl. Ital. viii. 360 = aristata.
tenuifolia, Schrad. in DC. Prod. vii. 120 = Candollei.
Thirkei, Sch. Bip. in Linnaea, xix. (1847) 314 = eriophora.
tingitana, Linn. Sp. Pl. 791 = Picridium tingitanum.
tomentosa, *Linn. l. c.* ed. II. 1112.—As. Min.
tomentosa, Pall. Reise, iii. 513 = mollis.
tomentosa, Siev. ex Ledeb. Fl. Alt. iv. 163 = stricta.
tortuosissima, *Boiss. Fl. Orient.* iii. 775.—Persia; Beluchist.
trachysperma, *Guenth. ex Spreng. Syst.* iii. 666.—Ital.
tragopogonoides, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 635.—Turkestan.
tuberosa, *Pall. Reise*, iii. 757.—Turkestan.
tuberosa, Walp. Rep. vi. 351, sphalm. = suberosa.

SCORZONERA :—
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 333.—Turkestan.
undulata, Vahl, Symb. Bot. ii. 86 = mollis.
vaginata, Bertol. Misc. Bot. ii. 235 = lanata.
variifolia, Salisb. Prod. 179 = Picridium vulgare.
velutina, DC. Prod. vii. 123 = mollis.
veratrifolia, *Fenzl, in Flora*, xxvi. (1843) 399.—Kurdist.
verrucosa, *Boiss. Fl. Orient.* iii. 764.—Cilicia.
villosa, Bieb. Fl. Taur. Cauc. ii. 235 = stricta.
villosa, *Scop. Fl. Carn.* ed. II. ii. 97. t. 46.—Europ. austr.
virgata, DC. Prod. vii. 125 = divaricata.

SCORZONEROIDES, Moench, Meth. 549 (1794) = **Leontodon**, Linn. (Compos.).
autumnalis, Moench, l. c. = L. autumnalis.

SCOTANTHUS, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 172 = **Gymnopetalum**, Arn. (Cucurb.).
Porteanus, Naud. l. c. Sér. V. v. (1866) 25 = G. quinquelobum.
tubiflorus, Naud. l. c. Sér. IV. xvi. (1861) 173. t. 3 = G. cochinchinense.
Weberi, Naud. l. c. Sér. V. v. (1866) 26 = G. Weberi.

SCOTANUM, Adans. Fam. ii. 459 (1763) = **Ranunculus**, Tourn.

SCOTIA, Thunb. Nov. Gen. ix. 129 (1798), sphalm. = **Schotia**, Jacq. (Legumin.).

SCOTTEA, DC. Prod. ii. 118 (1825), sphalm. = seq.

SCOTTIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 268 (1812) = **Bossiaea**, Vent. (Legumin.).
angustifolia, Lindl. Bot. Reg. t. 1266 = B. dentata.
dentata, R. Br. in Ait. Hort. Kew. ed. II. iv. 269 = B. dentata.
laevis, Lindl. Bot. Reg. t. 1652 = B. dentata.
trapeziformis, Mackay, ex Sweet, Hort. Brit. ed. II. 126 (Quid?).—Austral.

SCOVITZIA, Walp. Rep. ii. 420 (1843), sphalm. = **Szowitsia**, Fisch. & Mey. (Umbell.).

SCRIBAEA, Borkh. in Rhein. Mag. i. (1793) 591 = **Cucubalus**, Tourn. (Caryoph.).
baccifera, Schur, Enum. Pl. Transs. 100 = C. baccifer.
Cucubalus, Borkh. in Rhein. Mag. i. (1793) 591 = C. baccifer.
divaricata, Gaertn. Mey. & Scherb. Fl. Wett. ii. 96 = C. baccifer.

SCROBICARIA, Cass. in Dict. Sc. Nat. xlviii. 456 (1827) = **Gynoxys**, Cass. (Compos.).
ilicifolia, Cass. l. c. = G. ilicifolia.

SCROFULARIA, Spreng. Fl. Hal. 178 (1806) = seq.

SCROPHULARIA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 178 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 937.
CERAMANTHE, Dum. Not. sur Scroph. 7 (1834).
TOMIEPHYLLUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125.
VENILIA, Fourr. l. c. (1869).
adscendens, *Willd. Enum. Hort. Berol.* 644.—Hab.?
aestivalis, *Griseb. Spicil. Fl. Rumel.* ii. 36.—Macedon.
affinis, Opiz, in Lotos, iv. (1854) 71 = alata?
aintabensis, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 413 = xanthoglossa.
alata, *Gilib. Fl. Lituan.* i. 127.—Europ.; Oriens; As. bor.
alata, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 401.—Japon.
alpestris, *J. Gay, ex Benth. in DC. Prod.* x. 307.—Mont. Pyren.
altaica, *Murr. in Comm. Gotting.* (1781) 35. t. 2.—Sibir.
amgunensis, *F. Schmidt, Reisen Amurl.* 57.—Manchur.
amplexicaulis, *Benth. in DC. Prod.* x. 310.—Persia.
Anagae, *Bolle, in Verh. Zool.-Bot. Ges. Wien*, xi. (1861) 199.—Ins. Canar.
Ani, C. Koch, in Linnaea, xvii. (1843) 285 = variegata.
anomala, *Vest, in Flora*, xii. (1829) 1. *Erg.* 62.—Hab.?
anthemifolia, *Parrot, in Engelh. & Parrot, Reise Krym*, ii. 125.—Reg. Caucas.

SCROPHULARIA :—
appendiculata, Balb. ex Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 157 = auriculata.
appendiculata, Jacq. Hort. Schoenb. iii. 19 = trifoliata.
aquatica, Koch, Syn. Fl. Germ. ed. I. 515 = alata, *Gilib.*
aquatica, *Linn. Sp. Pl.* 620.—Europ.; Caucas.
arguta, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 342.—Afr. bor.; Ins. Canar.; Arab.
armena, *C. Koch, in Linnaea*, xxii. (1849) 711.—Armenia.
ascendens, Walp. Rep. iii. 176 = adscendens.
atropurpurea, Moretti, in Bibl. Ital. xii. (1818) 372 = canina.
auriculata, Asso, Syn. Stirp. Arag. 81 = oblongifolia.
auriculata, Brot. Fl. Lusit. i. 201 = trifoliata.
auriculata, Heldr. ex Boiss. Fl. Orient. iv. 401 = cretica.
auriculata, *Linn. Sp. Pl.* 620.—Hispan.
auriculata, Scop. Fl. Carn. ed. II. i. 446 = Scopolii.
Balbisii, Hornem. Hort. Hafn. ii. 577 = auriculata.
Balbisii, Koch, Syn. Fl. Germ. ed. I. 515 = oblongifolia.
Benthamiana, *Boiss. Diagn.* Ser. I. xii. 39.—Persia.
Berthelotii, *Bolle, in Verh. Zool.-Bot. Ges. Wien*, xi. (1861) 202.—Ins. Canar.
betonicaefolia, Lapeyr. Hist. Abr. Pl. Pyr. 356 = alpestris.
betonicaefolia, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 151 = Scopolii.
betonicefolia, Viv. Fl. Cors. 10 = oblongifolia.
betonicifolia, Linn. Mant. i. 87 = S. Scorodonia.
bicolor, Gueldenst. ex Ledeb. Fl. Ross. iii. 221 = variegata.
bicolor, Sibth. & Sm. Fl. Graec. Prod. i. 437 = canina.
biserrata, *Willd. Enum. Hort. Berol.* 644.—Hab.?
Boissieriana, *Jaub. & Spach, Illustr.* iii. 30.—Persia.
Bourgaeana, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* ii. 550.—Hispan.
Buergeriana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 116.—Japon.
byzantina, Benth. in DC. Prod. x. 303 = Kotschyana.
cabulica, *Benth. l. c.* 316.—Afghan.
caesia, Griseb. Spicil. Fl. Rumel. ii. 38 = laciniata.
caesia, Sibth. & Sm. Fl. Graec. Prod. i. 438; Fl. Graec. vii. 3. t. 604 = heterophylla.
californica, *Cham. & Schlecht. in Linnaea*, ii. (1827) 585.—Calif.
calliantha, *Webb & Berth. Phyt. Canar.* iii. 126.—Ins. Canar.
Calverti, Boiss. Diagn. Ser. II. iii. 152 = ilwensis.
calycina, *Benth. Scroph. Ind.* 18.—Reg. Himal.
calycina, Boiss. Fl. Orient. iv. 390 = chrysantha.
canescens, *Bong. in Bull. Sc. Acad. Pétersb.* viii. (1841) 340.—Sibir. altaic.
canina, Bourg. ex Reut. in Bull. Soc. Bot. Fr. xi. (1864) 46 = frutescens.
canina, Hoppe, ex Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 165 = Hoppii.
canina, *Linn. Sp. Pl.* 621.—Europ.; As. Min.; Afr. bor.
canina, Sibth. & Sm. Fl. Graec. vi. 78. t. 598 = sphaerocarpa.
capillaris, *Boiss. & Bal. ex Boiss. Fl. Orient.* iv. 397.—As. Min.
Castagneana, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 149 = Scopolii.
catariaefolia, *Boiss. & Heldr. Diagn.* Ser. I. xii. 36.—Lycaonia.
čechica, Opiz, Seznam, 90 = nodosa?
chamaedrifolia, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 418 = variegata.
chinensis, Linn. Mant. ii. 250; Sm. in Rees, Cycl. xxxii. sub n. 94 = Celsia coromandeliana *cum* Salvia plebeia.
chlorantha, *Kotschy & Boiss. in Boiss. Fl. Orient.* iv. 399.—Kurdistan.
chrysantha, *Jaub. & Spach, Illustr.* iii. 26. t. 220.—Reg. Caucas.
chrysanthemifolia, Balb. ex Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 164 = canina.
chrysanthemifolia, Bory & Chaub. Nouv. Fl. Pélop. 40 = heterophylla.
chrysanthemifolia, Reichb. ex Nym. Consp. 534 = Hoppii.
chrysanthemifolia, Willd. Hort. Berol. i. 59 = canina.
cinerascens, Boiss. in Tchihat. Asie Min. Bot. ii. 16 = variegata.

SCROPHULARIA:—
cinerea, Dum. Not. sur Scroph. 11 = aquatica.
clandestina, Rupr. ex Boiss. Fl. Orient. iv. 392 = lateriflora.
Clausii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 163.—Persia.
coccinea, *A. Gray, in Torr. Bot. Mex. Bound.* 111.—N. Mexic.
coccinea, Linn. Sp. Pl. 621 = Russelia sarmentosa.
congesta, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) i. 348, in textu = chrysantha.
cordata, Mill. Gard. Dict. ed. VIII. n. 4 = S. Scorodonia.
cordata, Waldst. & Kit. Pl. Rar. Hung. i. 75. t. 73 = vernalis.
crassicaulis, *Boiss. Diagn.* Ser. I. xii. 37.—Persia.
crenophila, *Boiss. l. c.* vii. 41.—Persia.
cretacea, *Fisch. Hort. Gorenk.* ed. II. (1812) 24, nomen; *et ex Spreng. Syst.* ii. 788.—Sibir.
cretica, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 400.—Ins. Creta.
crithmifolia, *Boiss. Voy. Bot. Espagne*, ii. 447.—Hispan.
cryptophilla, *Boiss. & Heldr. Diagn.* Ser. I. xii. 31.—Persia.
cymigera, C. Koch, in Linnaea, xix. (1847) 22 = alata.
Decaisniana, Presl, Bot. Bemerk. 91 = variegata.
decipiens, Boiss. & Kotschy, Diagn. Ser. II. iii. 156 = xanthoglossa.
decomposita, Royle, ex Benth. Scroph. Ind. 18 = lucida.
decora, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 70 = sambucifolia.
decumbens, Fisch. Mey. & Avé-Lall. l. c. x. 58 = Scopolii.
dentata, *Royle, ex Benth. Scroph. Ind.* 19.—Reg. Himal.
depauperata, Boiss. Diagn. Ser. I. iv. 68 = variegata.
deserti, *Delile, Fl. Egypte*, 240. t. 33. f. 1.—Aegypt.; Arab.; Persia.
digitalifolia, *Richt. ex Stapf, in Denkschr. Akad. Wien*, l. (1885) 23.—Persia.
divaricata, *Ledeb. Ind. Sem. Hort. Dorpat.* (1822) 17.—Reg. Caucas.
Durandii, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 91.—Marocc.
ebulifolia, Bieb. Fl. Taur.-Cauc. ii. 77 = orientalis.
ebulifolia, Hoffmgg. & Link, Fl. Port. i. 270 = sublyrata.
Edgeworthii, *Benth. in DC. Prod.* x. 304.—Reg. Himal.
Ehrharti, Stevens, in Ann. Nat. Hist. Ser. I. v. (1840) 3 = alata, *Gilib.*
elatior, *Benth. in Wall. Cat.* n. 3921.—Reg. Himal.
elongata, *Webb, ex A. DC. in DC. Prod.* x. 593.—Ins. Canar.
expansa, Reut. ex Boiss. Fl. Orient. iv. 413 = xanthoglossa.
farinosa, *Boiss. Diagn.* Ser. I. iii. 40.—Persia.
ficifolia, Pritz. Ic. Ind. i. 1010, sphalm. = seq.
filicifolia, *Mill. Gard. Dict.* ed. VIII. n. 10.—Graecia.
filicifolia, Sibth. & Sm. Fl. Graec. vi. 79. t. 600 = lucida.
floribunda, Boiss. & Bal. Diagn. Ser. II. iii. 158 = canina.
fluminensis, Vell. Fl. Flum. 263; vi. t. 87 = Alectra brasiliensis.
foetida, *Wydl. in Mém. Soc. Phys. Nat. Genèv.* vi. (1828) 154.—Hab. ?
foliosa, Pomel, Nouv. Mat. Fl. Atl. 101 = laevigata.
fontana, Kotschy, ex Boiss. Fl. Orient. iv. 396 = Scopolii.
frigida, *Boiss. Diagn.* Ser. I. vii. 42.—Persia.
frutescens, DC. Fl. Fr. iii. 729 = ramosissima.
frutescens, *Linn. Sp. Pl.* 621.—Reg. Mediterr.
frutescens, Sibth. & Sm. Fl. Graec. Prod. i. 437 = laciniata.
geminiflora, Lam. Fl. Fr. ii. 336 = peregrina.
georgica, Benth. in DC. Prod. x. 305 = divaricata.
glabrata, *Ait. Hort. Kew.* ed. I. ii. 341.—Ins. Canar.
glabrata, Schimp. ex Benth. in DC. Prod. x. 309 = auriculata.
glabrata, Spreng. Syst. ii. 784 = nepetaefolia.
glandulifera, Clarke, Trav. i. ed. II. App. p. xxi = Scopolii.

SCROPHULARIA:—
glandulosa, Waldst. & Kit. Pl. Rar. Hung. ii. 238. t. 214 = Scopolii.
glauca, *Decne. ex Benth. in DC. Prod.* x. 310.—Persia.
glauca, Sibth. & Sm. Fl. Graec. vi. 78. t. 599 = lucida.
Gmelini, Turcz. ex Benth. in DC. Prod. x. 311 = incisa.
grandidentata, Tenore, Fl. Neap. Suppl. ii. 69 = Scopolii.
grandiflora, DC. Cat. Hort. Monsp. 143 = sambucifolia.
grandifolia, *C. Koch, in Linnaea*, xxii. (1849) 707.—Banat.
Grenieri, *Reut. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 554.—Hispan.
Griffithii, Benth. in DC. Prod. x. 312 = lucida.
Gussonei, Nym. Consp. 534 = frutescens.
haematantha, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 415.—As. Min.
Halleri, Gueldenst. ex Ledeb. Fl. Ross. iii. 219 = nodosa.
hastata, *Rafin. Fl. Ludov.* 44.—Am. bor.
Heldreichii, *Boiss. Diagn.* Ser. II. iii. 158.—As. Min.
hemschinica, C. Koch, in Linnaea, xxii. (1849) 708 = nodosa.
Herminii, *Hoffmgg. & Link, Fl. Port.* i. 266. t. 53.—Lusitan.
heterophylla, Sibth. & Sm. Fl. Graec. vii. 2. t. 603 = laciniata.
heterophylla, *Willd. Sp. Pl.* iii. 274.—Graecia.
heucheriaeflora, *Schrenk, Enum. Pl. Nov.* i. 25.—Soongar.
hierochuntina, Boiss. Diagn. Ser. I. xii. 35 = Michoniana.
himalensis, *Royle, ex Benth. Scroph. Ind.* 18.—Reg. Himal.
hirsuta, Hornem. Hort. Hafn. Suppl. 68 = Scopolii.
hirta, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 21.—Ins. Mader.
hispanica, Coss. Notes Crit. 124 = sciaphila.
hispida, *Desf. Fl. Atlant.* ii. 55.—Afr. bor.
hispidula, Boiss. & Bal. Diagn. Ser. II. vi. 157 = xanthoglossa.
Hoppii, *Koch, in Mert. & Koch, Deutschl. Fl.* ii. 410.—Carniol.; Ital.
humifusa, Timb. ex Nym. Consp. 534 = canina.
humilis, *Desf. Cat. Hort. Par.* ed. III. 396.—Hab. ?
hypericifolia, *Wydl. in Mém. Soc. Phys. Genèv.* iv. (1828) 166. t. 5.—Mesopot.
hyssopifolia, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 418.—As. Min.
ilwensis, *C. Koch, in Linnaea*, xvii. (1843) 284.—As. Min.
incisa, C. Koch, ex Boiss. Fl. Orient. iv. 418 = variegata.
incisa, *Weinm. Ind. Sem. Hort. Dorpat.* (1810) 136.—Sibir.
italica, Mill. Gard. Dict. ed. VIII. n. 6 = nodosa.
juncea, *Richt. ex Stapf, in Denkschr. Akad. Wien*, l. (1885) 24.—Persia.
juratensis, Schleich. ex Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 164 = Hoppii.
kakudensis, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 87.—Japon.
Kotschyana, *Benth. in DC. Prod.* x. 303.—As. Min.
Kotschyi, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 256 = dentata.
lacera, *Fisch. Hort. Gorenk.* ed. II. (1812) 24, nomen.—Hab. ?
laciniata, C. Koch, in Linnaea, xxiii. (1850) 710 = variegata.
laciniata, *Waldst. & Kit. Pl. Rar. Hung.* ii. 185. t. 170.—Europ. austr.
laevigata, *Vahl, Symb. Bot.* ii. 67.—Afr. bor.
lanata, *Fisch. ex A. DC. in DC. Prod.* x. 593, nomen.—Sibir.
lanceolata, Pursh, Fl. Am. Sept. ii. 419 = nodosa.
lancifolia, *Presl, Fl. Sic.* i. p. xxxv.—Sicil.
Langeana, *Bolle, in Verh. Zool.-Bot. Ges. Wien*, xi. (1861) 196.—Ins. Canar.
lasiocaulis, Schur, Enum. Pl. Transs. 486 = rupestris.
lateriflora, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 396.—Reg. Caucas.
latifolia, Host, Fl. Austr. ii. 206 = vernalis.
laxa, *Boiss. & Heldr. Diagn.* Ser. II. iii. 154.—Graecia.

SCROPHULARIA:—
laxiflora, Lange, in Kjoeb. Vidensk. Meddel. (1877-78) 231 = nodosa.
lazica, Boiss. & Bal. ex Boiss. Fl. Orient. iv. 409 = olympica.
lepidota, *Boiss. Diagn.* Ser. I. iv. 71.—As. Min.
leucoclada, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 424.—Turkestan.
libanotica, Boiss. Diagn. Ser. I. xii. 36 = variegata.
livida, Heldr. ex Nym. Consp. 534 = lucida.
livida, Sibth. & Sm. Fl. Graec. Prod. i. 437 = sphaerocarpa.
longiflora, Benth. in DC. Prod. x. 313 = farinosa.
longifolia, *Benth. l. c.* 309.—Ins. Mader.
lucida, Bieb. Fl. Taur. Cauc. ii. 78 = rutaefolia.
lucida, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 554 = crithmifolia.
lucida, *Linn. Syst.* ed. X. 1114.—Europ. austr.; As. occ.; Reg. Himal.
lucida, Pall. ex Bieb. Fl. Taur. Cauc. ii. 78, in syn. = canina.
lunariaefolia, *Boiss. & Bal. ex Boiss. Fl. Orient.* iv. 390.—As. Min.
luridiflora, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 40.—Anatolia.
lutea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 320 = vernalis.
lyrata, Willd. Enum. Hort. Berol. 645 = auriculata.
macrobotrys, Ledeb. Fl. Ross. iii. 217 = nodosa.
macrophylla, *Boiss. Diagn.* Ser. I. xii. 32.—Syria.
mandshurica, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 35.—China.
marginata, *Boiss. Diagn.* Ser. I. iv. 72.—Persia.
marilandica, Linn. Sp. Pl. 619 = nodosa.
marylandica, Georgi, Beschr. Russ. Reich. III. iv. 1108 = altaica.
melissaefolia, Salisb. Prod. 100 = S. Scorodonia.
melissaefolia, Urv. in Mém. Soc. Linn. Par. i. (1822) 331 = Scopolii.
mellifera, *Ait. Hort. Kew.* ed. I. ii. 343 = sambucifolia.
mellifera, Loisel. ex Nym. Consp. 532 = trifoliata.
meridionalis, Linn. f. Suppl. 280 = Alonsoa caulialata.
meridionalis, Presl, Fl. Sic. i. p. xxxv = peregrina.
mesogitana, *Boiss. Fl. Orient.* iv. 407.—Lydia.
mesopotamica, *Boiss. Diagn.* Ser. I. xii. 34.—Mesopot.
Michoniana, *Coss. & Kral. Cat. Pl. Syr. et Palest.* 13.—Syria.
micrantha, Desv. ex Ham. Prod. Fl. Ind. Occ. 47 (Quid ?).—Ins. S. Doming.
micrantha, Urv. in Mém. Soc. Linn. Par. i. (1822) 331 = heterophylla.
minima, Benth. in DC. Prod. x. 303 = chrysantha.
minima, *Bieb. Fl. Taur. Cauc.* ii. 79.—Reg. Caucas.
minor, Savi, Fl. Pisan. ii. 81 = peregrina.
Moellendorffi, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 501.—China.
multicaulis, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 76.—Sibir.
multifida, Lam. Fl. Fr. ii. 336 = canina.
multifida, Willd. Enum. Hort. Berol. 646 = laciniata.
murcica, Guir. ex Willk. in Linnaea, xxx. (1859-60) 136 = sciaphila.
myriophylla, *Boiss. & Heldr. Diagn.* Ser. I. xii. 39.—As. Min.
Neesii, Wirtg. in Verh. Nat. Ver. Rheinl. i. (1844) 29 = aquatica.
nepetaefolia, Boiss. & Heldr. ex Boiss. Fl. Orient. iv. 407 = catariaefolia.
nepetaefolia, *Presl, Delic. Prag.* 73 (nepethaefolia).—Hispan.
nepetaefolia, Sm. in Rees, Cyclop. xxxii. n. 7 = Scopolii.
nervosa, *Benth. in DC. Prod.* x. 303.—Persia.
nitida, *Richt. ex Stapf, in Denkschr. Akad. Wien.* l. (1885) 24.—Persia.
nodosa, *Linn. Sp. Pl.* 619.—Reg. bor. temp.
nudicaulis, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 168 = Celsia persica.
obliqua, *Megg. Kanitz & Knapp, in Verh. Zool.-Bot. Ges. Wien*, xvi. (1866) 125.—Austria.
oblongifolia, *Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 418.—Ins. Corsic.
obtusa, *Edgew. ex Hook. f. Fl. Brit. Ind.* iv. 254.—Reg. Himal.
obtusifolia, *Kit. ex A. DC. in DC. Prod.* x. 593, nomen.—Hungar.
Oldhami, *Oliver, in Journ. Linn. Soc.* ix. (1867) 167.—Japon.
oligantha, Boiss. & Heldr. Diagn. Ser. I. xii. 33 = Scopolii.

SCROPHULARIA:—
Oliveriana, C. Koch, in Linnaea, xxii. (1849) 709 = olympica.
Olivieri, Jaub. & Spach, Illustr. iii. 29. t. 222 = nervosa.
Olivieriana, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 159 = heterophylla.
olympica, *Boiss. Diagn.* Ser. I. iv. 69.—As. Min.
orientalis, Boiss. Fl. Orient. iv. 393 = Boissieriana.
orientalis, Ehrenb. ex Boiss. l. c. 413 = xanthoglossa.
orientalis, Habl. Phys. Desc. Taur. 164 = rupestris.
orientalis, *Linn. Sp. Pl.* 620.—As. Min. et bor.
oxysepala, *Boiss. Fl. Orient.* iv. 398.—Persia.
pallida, *Fisch. ex Sweet*, *Hort. Brit.* ed. II. 392; *A. DC. in DC. Prod.* x. 593, nomen.—Reg. Caucas.
paniculata, Seenus, Reise Istrien, 67 = peregrina.
Pantocsekii, Griseb. ex Pantoč. in Oest. Bot. Zeitschr. xxiii. (1873) 267 = Scopolii.
papillaris, Boiss. & Reut. Pugill. Pl. Nov. (1852) 90 = S. Scorodonia.
Patriniana, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 159 = incisa.
pauciflora, *Benth. Scroph. Ind.* 17.—Reg. Himal.
pellucida, Pomel, Nouv. Mat. Fl. Atl. 101 = laevigata.
peregrina, *Linn. Sp. Pl.* 621.—Reg. Mediterr.
persica, Benth. in DC. Prod. x. 314 = subaphylla, xanthoglossa, variegata.
petraea, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 180.—Afghan.
physalodes, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xv. (1827) 353 = Melasma hispidum.
Pinardi, *Boiss. Diagn.* Ser. I. iv. 70.—As. Min.
pinnata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 719.—Sibir.
pinnata, Mill. Gard. Dict. ed. VIII. n. 15 = canina.
pinnatifida, Brot. Fl. Lusit. i. 202 = canina.
pinnatifida, Guss. Fl. Sic. Syn. ii. 1. 129 = frutescens.
pisidica, Boiss. & Heldr. Diagn. Ser. I. xii. 33 = alata, *Gilib.*
polyantha, *Royle, ex Benth. Scroph. Ind.* 18.—Reg. Himal.
polybotrys, *Boiss. Diagn.* Ser. I. xii. 32.—Persia.
prasiifolia, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 416.—Persia.
procumbens, Vell. Fl. Flum. 264; vi. t. 89 = Herpestis lanigera.
pruinosa, *Boiss. Diagn.* Ser. I. xii. 38.—Persia.
puberula, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 396 = Scopolii.
puberula, *Boiss. l. c.* 418.—Persia.
pubescens, Hort. ex Sweet, Hort. Brit. ed. II. 392 = canina.
pulverulenta, *Boiss. & Noé*, *Diagn.* Ser. II. iii. 155.—As. Min.
pulverulenta, *Janka, in Termész. Füzetek*, iv. (1880) 309.—Europ. austr. occ.
pumila, Adams, ex Ledeb. Fl. Ross. iii. 215 = minima.
pyramidalis, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 165 = canina.
pyrenaica, *Benth. in DC. Prod.* x. 306.—Mont. Pyren.
pyrrolopha, Boiss. Fl. Orient. iv. 409 = olympica.
racemosa, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 20.—Ins. Mader.
ramosissima, C. Koch, in Linnaea, xvii. (1843) 285 = variegata.
ramosissima, *Loisel. Fl. Gall.* ed. II. ii. 381.—Ins. Corsic.
ramosissima, Urv. in Mém. Soc. Linn. Par. i. (1822) 331 = canina.
rivularis, Moris, Stirp. Sard. Elench. App. 8 = aquatica.
rostrata, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 163.—Persia.
rostrata, Hochst. ex Boiss. Fl. Orient. iv. 395 = arguta.
rotundifolia, Host, ex Schur, Enum. Pl. Transs. 486 = vernalis.
rubricaulis, Boiss. Diagn. Ser. I. xii. 34 = Michoniana.
rugosa, Hort. Lugd. ex Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 151 = betonicifolia.
rugosa, Willd. Enum. Hort. Berol. Suppl. 42 = Scopolii.
rupestris, *Bieb. ex Willd. Sp. Pl.* iii. 274.—Reg. Caucas.
Ruprechti, *Boiss. Fl. Orient.* iv. 410.—Reg. Caucas.
rutaefolia, *Boiss. l. c.* 404.—Reg. Caucas.
Samaritanii, Boiss. & Heldr. ex Boiss. l. c. 400 = aquatica.

SCROPHULARIA:—
sambucifolia, Fisch. Ind. Sem. Hort. Petrop. viii. 70 = trifoliata.
sambucifolia, *Linn. Sp. Pl.* 620.—Reg. Medit. occ.
saxatilis, Boeb. ex Ledeb. Fl. Ross. iii. 221 = rupestris.
scabiosaefolia, *Benth. in DC. Prod.* x. 311.—Reg. Himal.; Afghan.
scariosa, *Boiss. Diagn.* Ser. I. iv. 67.—Syria.
Schousboei, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* ii. 553.—Hispan.
sciaphila, *Willk. in Bot. Zeit.* viii. (1850) 77.—Hispan.
Sckellii, Spreng. Syst. ii. 783 = nodosa.
Scopolii, Loisel. Not. Pl. Fr. 95; DC. Fl. Fr. Suppl. 406 = alpestris.
Scopolii, *Hoppe, ex Pers. Syn.* ii. 160.—Europ. or.; As. Min.; Reg. Himal.
Scorodonia, Host, Fl. Austr. ii. 214 = Scopolii.
Scorodonia, *Linn. Sp. Pl.* 620.—Europ.; As. et Afr. bor.
scorodonifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 320 = S. Scorodonia.
sexangularis, Moench, Meth. 445 = peregrina.
Sibthorpiana, Spreng. Syst. ii. 786 = laciniata.
silaifolia, Clarke, Trav. ii. 261 = laciniata.
sinaica, Benth. in DC. Prod. x. 314 = deserti.
sinensis, J. F. Gmel. Syst. 938 = *chinensis*.
Skellii, Walp. Rep. iii. 177, lapsu = *Sckellii*.
Smithii, *Hornem. Hort. Hafn. Suppl.* 68.—Ins. Canar.
smyrnaea, Boiss. Diagn. Ser. I. iv. 66 = Scopolii.
sphaerantha, *Trautv. ex Becker, in Bull. Soc. Nat. Mosc.* xlviii. (1874) 1. 213, nomen.—Reg. Caucas.
sphaerocarpa, *Boiss. & Reut. Diagn.* Ser. II. iii. 158, in textu.—As. Min.; Syria.
Stelleri, Ledeb. in Denkschr. Bot. Ges. Regensb. iii. (1841) 98 = multicaulis.
Stelleriana, Steud. Nom. ed. II. ii. 548 = multicaulis.
striata, *Boiss. Fl. Orient.* iv. 413.—Persia.
subaphylla, *Boiss. Diagn.* Ser. I. vii. 41.—Persia.
subcrispa, Pomel, Nouv. Mat. Fl. Atl. 102 = hispida.
subhastata, Vell. Fl. Flum. 264; vi. t. 88 = Stemodia subhastata.
sublyrata, *Brot. Phyt. Lusit.* ii. 156.—Lusitan.
subverticillata, Moris, Stirp. Sard. Elench. App. 8 = auriculata.
suffruticans, Pall. ex Steud. Nom. ed. I. 756 = tanacetifolia.
sulphurea, Mill. Gard. Dict. ed. VIII. n. 13 = auriculata.
sylvatica, *Boiss. & Heldr. Diagn.* Ser. II. iii. 153.—Graecia.
syriaca, *Benth. in DC. Prod.* x. 316.—Palaestin.
tagetifolia, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 401.—Syria.
tanacetifolia, *Willd. Hort. Berol.* t. 56.—Am. bor.
taurica, Hort. ex Schur, Enum. Pl. Transs. 485 = Scopolii.
taygetea, *Boiss. Diagn.* Ser. I. iv. 68.—Graecia.
tenuipes, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 175.—Algeria.
tenuisecta, Jord. ex Nym. Consp. 534 = canina.
ternata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 53 = nodosa.
thesioides, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 164.—Armenia.
Tinantii, Dum. in Bull. Soc. Bot. Belg. xiv. (1875) 110 = S. Scorodonia.
tmolea, Boiss. Diagn. Ser. I. iv. 66 = Scopolii.
trichopoda, *Boiss. & Bal. l. c.* Ser. II. iii. 156.—Cilicia.
trifoliata, Desf. Fl. Atlant. ii. 54 = laevigata.
trifoliata, Hoffmgg. & Link, Fl. Port. i. 267 = auriculata.
trifoliata, *Linn. Syst.* ed. X. 1114.—Reg. Mediterr.
umbrosa, Dum. Prod. Fl. Belg. 37 = aquatica.
umbrosa, Salzm. ex Benth. in DC. Prod. x. 309 = auriculata.
uniflora, *Richt. in Denkschr. Akad. Wien*, l. (1885) 90.—Lycia.
urticifolia, *Wall. Cat.* n. 3922.—Reg. Himal.
Urvillaei, Walp. Ann. v. 624 = seq.
Urvilleana, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 252 = variegata.
Urvilleana, Wydl. in Mém. Soc. Phys. Genèv. iv. (1828) 160 = heterophylla.
valentina, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 124.—Hispan.
varia, *Bieb. Tabl. Casp.* 58 (an seq. ?).—Reg. Casp.

SCROPHULARIA:—
variegata, *Bieb. Beschr. Casp.* 177, *App.* n. 48; *et Fl. Taur. Cauc.* ii. 78.—As. Min.; Reg. Himal.
variegata, Reichb. ex Nym. Consp. 534 = laciniata.
verbenaefolia, Desf. Tabl. ed. II. 77 = caesia.
verna, Mill. Gard. Dict. ed. VIII. n. 13 = vernalis.
vernalis, Bieb Fl. Taur. Cauc. ii. 76 = chrysantha.
vernalis, Lapeyr. Hist. Abr. Pl. Pyr. 356 = pyrenaica.
vernalis, *Linn. Sp. Pl.* 620.—Europ.
versicolor, *Boiss. Diagn.* Ser. II. iii. 155.—Armenia.
verticillata, *Moris, Stirp. Sard. Elench. App.* 7.—Ins. Sardoa.
viridiflora, Poir. Voy. Barb. ii. 194 = sambucifolia.
viscosa, Boiss. Fl. Orient. iv. 391 = Kotschyana.
Wirtgenii, Koch, ex Opiz, Seznam, 90 = nodosa ?
xanthoglossa, *Boiss. Diagn.* Ser. I. xii. 38.—As. Min.; Syria.
xylorrhiza, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 406.—Syria; Mesopot.

SCROPHULARIOIDES, Forst. f. Prod. 91 (1786) = **Premna**, Linn. (Verben.).
arborea, Forst. f. l. c. = P. taitensis.

SCUBALIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3 (Quid ?).

SCULERIA, Rafin. Fl. Tellur. ii. 51 (1836) = **Vancouveria**, Decne. (Berber.).
geminata, Rafin. l. c. = V. hexandra.

SCURIA, Rafin. in Journ. Phys. lxxxix. (1819) 106 = **Carex**, Dill. (Cyperac.).

SCURRULA, Linn. Sp. Pl. ed. I. 110 (1753) = **Loranthus**, Linn.
amplexifolia, G. Don, Gen. Syst. iii. 424 = L. longiflorus.
Belvisii, G. Don, l. c. 423 = L. lanceolatus.
bicolor, G. Don, l. c. 424 = L. longiflorus.
biflora, G. Don, l. c. 422 = L. Scurrula ?
bracteata, G. Don, l. c. 425 = L. bracteatus.
budleioides, G. Don, l. c. 422 = L. Scurrula.
canescens, G. Don, l. c. 423 = L. canescens.
chinensis, G. Don, l. c. 421 = L. chinensis.
cinerea, Rafin. Sylva Tellur. 159 = L. canescens ?
clavata, G. Don, Gen. Syst. iii. 423 = L. clavatus.
cordifolia, G. Don, l. c. 421 = L. cordifolius.
Corynitis, G. Don, l. c. 422 = L. Corynitis.
curvata, G. Don, l. c. 425 = L. curvatus.
cylindrica, G. Don, l. c. 424 = L. cylindricus.
dodoneaefolia, G. Don, l. c. 423 = L. dodoneaefolius.
elastica, G. Don, l. c. 425 = L. elasticus.
elegans, G. Don, l. c. 424 = L. elegans, *Wall.*
elliptica, Rafin. Sylva Tellur. 125 = L. tetrapetalus.
falcata, G. Don, Gen. Syst. iii. 424 = L. longiflorus.
fasciculata, G. Don, l. c. 421 = L. fasciculatus.
fusca, G. Don, l. c. = L. fuscus.
glauca, G. Don, l. c. 423 = L. glaucus.
goodeniaefolia, G. Don, l. c. 425 = L. cuneatus.
Haenkeana, G. Don, l. c. 423 = L. Haenkeanus.
heterantha, G. Don, l. c. 424 = L. heteranthus.
incarnata, G. Don, l. c. 423 = L. incarnatus.
indica, G. Don, l. c. 424 = L. longiflorus ?
Koenigiana, G. Don, l. c. 425 = L. longiflorus.
laevigata, G. Don, l. c. 422 = L. Scurrula.
lepidota, G. Don, l. c. = L. lepidotus.
lobeliaeflora, G. Don, l. c. 425 = L. cuneatus.
longiflora, G. Don, l. c. 423 = L. longiflorus.
malifolia, G. Don, l. c. = L. malifolius.
obovata, G. Don, l. c. 422 = L. Scurrula.
oleaefolia, G. Don, l. c. 423 = L. oleaefolius.
parasitica, Linn. Sp. Pl. 110 = L. Scurrula.
Pentagonia, G. Don, Gen. Syst. iii. 423 = L. Pentagonia.
pentandra, G. Don, l. c. 424 = L. pentandrus.
philippensis, G. Don, l. c. 422 = L. philippensis.
praelonga, G. Don, l. c. 425 = L. praelongus.
pulverulenta, G. Don, l. c. 421 = L. pulverulentus.
Reinwardtiana, G. Don, l. c. 425 = L. Reinwardtianus.
Roxburghii, G. Don, l. c. 421 = L. Scurrula.
rufescens, G. Don, l. c. 422 = L. rufescens.
rufidula, G. Don, l. c. = L. Scurrula.
rugulosa, G. Don, l. c. = L. rugulosus.
sessilifolia, G. Don, l. c. = L. sessilifolius.
Thonningii, G. Don, l. c. = L. Thonningii.
turbinata, G. Don, l. c. 424 = L. turbinatus.

SCURRULA :—
umbellata, Rafin. Sylva Tellur. 125 = Loranthus tetrandrus.
umbellifera, G. Don, Gen. Syst. iii. 421 = L. umbellifer.
venosa, G. Don, l. c. 424 = L. venosus.
vestita, G. Don, l. c. 422 = L. vestitus, *Wall.*

SCUTELLARIA, Riv. ex Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1201.
Cassida, Moench, Meth. 413 (1794).
Hastifolia, Ehrh. Beitr. iv. 148 (1789).
Adamsii, A. Ham. Monog. Scutell. 34. t. 2. f. 3 = baicalensis.
Adamsii, Spreng. Syst. ii. 701 = scordiifolia.
adenophylla, *Miq. in Journ. Bot. Néerl.* i. (1861) 117. —China.
adenotricha, Boiss. & Heldr. ex Boiss. Fl. Orient. iv. 688 = peregrina.
africana, Hochst. ex Benth. in DC. Prod. xii. 420 = peregrina.
agrestis, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 427.—Bras.
albida, Benth. in DC. Prod. xii. 420 = alpina.
albida, *Linn. Mant.* ii. 248.—Europ. or.; Oriens.
albo-rosea, *Lem. Illustr. Hortic.* xvi. (1869) t. 584.—Bras.
alpina, *Linn. Sp. Pl.* 599.—Europ.; As. bor.
altaica, Ledeb. ex Spreng. f. Tent. Suppl. Syst. 16 = alpina.
altissima, A. Ham. Monog. Scutell. 48 = Columnae.
altissima, *Linn. Sp. Pl.* 600.—Europ. austr.; Reg. Caucas.
amaranthina, *Linden, ex Galeotti, in Journ. Hort. Prat.* Sér. II. ii. 217. t. 19. f. 2.—N. Granat.
ambigua, Nutt. Gen. Am. ii. 37 = parvula.
angulosa, *Benth. in Wall. Pl. As. Rar.* i. 67.—Reg. Himal.
angustifolia, Adams, ex Benth. in DC. Prod. xii. 424, in nota = cordifolia.
angustifolia, *Pursh, Fl. Am. Sept.* ii. 412.—Am. bor. occ.
antirrhinoides, *Benth. in Bot. Reg.* sub t. 1493. n. 35. —Am. bor. occ.
arabica, *Jaub. & Spach, Illustr.* iv. 114. t. 376.—Arab.
arguta, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 175 = saxatilis.
arguta, *Soland. ex Coss. Notes Pl. Nouv.* 172.—Hispan.
atriplicifolia, *Benth. in Bot. Reg.* sub t. 1493. n. 8.—Peruv.
aurata, *Lem. Illustr. Hortic.* x. (1863) t. 368.—Bras.
baicalensis, *Georgi, Reise*, i. 228.—Sibir.; As. or.
balearica, *Barc. in Anal. Soc. Esp. Hist. Nat.* vi. (1877) 399.—Ins. Balear.
barbata, D. Don, Prod. Fl. Nep. 109 = rivularis.
bicolor, Hochst. in Lorent, Wanderungen, 333 = orientalis.
Bolanderi, *A. Gray, in Proc. Am. Akad.* vii. (1868) 387.—Calif.
bonariensis, Willd. ex Benth. in Linnaea, xi. (1837) 345 = rumicifolia.
brevibracteata, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 99.—Lycia.
breviflora, *Benth. in Bot. Reg.* sub t. 1493. n. 7.—Peruv.
breviflora, Turcz. ex Benth. Lab. Gen. et Sp. 439 = minor.
brevifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 380.—Texas.
californica, *A. Gray, l. c.* 381.—Calif.
cana, Wall. Pl. As. Rar. i. 67 = repens.
canescens, *Nutt. Gen. Am.* ii. 38.—Am. bor.
cardiophylla, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 227.—Am. bor. occ.
caroliniana, Lam. Encyc. vii. 706 = integrifolia.
caroliniana, Walt. Fl. Carol. 163 = pilosa, *Michx.*
caucasica, A. Ham. in Ser. Bull. Bot. (1832) 282 = orientalis.
celtidifolia, A. Ham. Monog. Scutell. 27 = angulosa.
Chamaedrys, Shuttl. ex Benth. in DC. Prod. xii. 422 = saxatilis.
chorassanica, *Bunge, Lab. Pers.* 65.—Persia.
coccinea, *H. B. & K. Nov. Gen. et Sp.* ii. 325.—N. Granat.
coerulea, *Moç. & Sesse, ex Benth. in Bot. Reg.* sub t. 1493. n. 49.—Mexic.
Colebrookiana, Wall. Pl. As. Rar. i. 67 = violacea.

SCUTELLARIA :—
Colebrookiana, Zoll. & Mor. Syst. Verz. Zoll. 54 = discolor.
Columnae, *All. Fl. Pedem.* i. 40. t. 84. f. 2.—Europ. austr.
Columnae, Host, Fl. Austr. ii. 178 = altissima.
commutata, Guss. Fl. Sic. Prod. ii. 136 = altissima.
compressa, A. Ham. Monog. Scutell. 29 = alpina
cordifolia, Benth. in Hook. Bot. Mag. t. 4290 = splendens.
cordifolia, Muhl. Cat. ex Benth. in DC. Prod. xii. 422 = versicolor.
cordifolia, Walp. Rep. iii. 757, sphalm. = scordiifolia.
costaricana, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 29.—Costa Rica.
cretacea, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 685.—Mesopot.
cretica, Linn. Sp. Pl. 600 = Teucrium lamiifolium.
cretica, Mill. Gard. Dict. ed. VIII. n. 2 = albida.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 158.—Cuba.
cumanensis, *H. B. & K. Nov. Gen. et Sp.* ii. 325.—Venezuela.
cuneata, Willd. ex Benth. in Linnaea, xi. (1837) 345 = pilosa, *Michx.*
cyrtopoda, *Miq. Fl. Ind. Bot.* ii. 973.—Java.
davurica, Pall. ex Ledeb. Fl. Ross. iii. 397 = baicalensis.
decumbens, Sieber, ex Reichb. Fl. Germ. Excurs. 332 = albida.
decumbens, Sieber, ex Spreng. Syst. ii. 702 = hirta.
deltoidea, Rafin. Fl. Ludov. 43 (Quid?).—Am. bor.
dependens, *Maxim. Prim. Fl. Amur.* 219.—Japon.
depressa, Fisch. ex Ledeb. Fl. Ross. iii. 398 = scordiifolia.
diffusa, *Benth. in DC. Prod.* xii. 421.—As. Min.; Syria.
discolor, *Colebr. in Wall. Pl. As. Rar.* i. 66.—Ind. or.; Burma; Java.
Drummondii, *Benth. Lab. Gen. et Sp.* 441.—Mexic.; Texas.
dumetorum, *Schlecht. in Linnaea*, vii. (1832) 400.—Mexic.
elliptica, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 173, nomen = pilosa, *Michx.*
epilobiifolia, A. Ham. Monog. Scutell. 32 = galericulata.
Felisberti, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 77 = purpurascens.
filicaulis, *Regel, Pl. Nov. Fedsch.* 68.—Turkest.
floribunda, Benth. in DC. Prod. xii. 418 = violacea.
floridana, *Chapm. Fl. S. U. St.* 324.—Florida.
fruticosa, *Desf. Tabl.* ed. I. 63.—Syria; Persia.
fruticosa, *Soyer- Willem. in Usteri, Ann. Bot.* xviii. (1796) 42.—Ins. Maurit.
galericulata, *Linn. Sp. Pl.* 599.—Reg. bor. temp.
gardoquioides, Benth. in DC. Prod. xii. 430 = Perilomia gardoquioides.
glandulosa, *Hook. f. Fl. Brit. Ind.* iv. 669.—Burma.
glechomoides, *Boiss. Diagn.* Ser. I. xii. 67.—Persia.
glutinosa, *Benth. in DC. Prod.* xii. 419.—Afghan.
gracilis, Nutt. Gen. Am. ii. 37 = nervosa.
grandiflora, Adam, ex Benth. in DC. Prod. xii. 424 = baicalensis.
grandiflora, Sims, Bot. Mag. t. 635 = orientalis.
grossa, *Wall. Pl. As. Rar.* i. 67.—Reg. Himal.
Guilielmi, *A. Gray, in Am. Ass. Adv. Sc.* xxi. (1872) 25.—Am. bor.?
Gussonii, Tenore, Fl. Nap. v. 29 = Columnae.
Hartwegii, *Benth. Pl. Hartw.* 244.—N. Granat.
hastaefolia, Tenore, ex Steud. Nom. ed. II. ii. 548 = minor.
hastata, Rafin. Fl. Ludov. 44 (Quid?).—Am. bor.
hastifolia, *Linn. Sp. Pl.* 599.—Europ.; Reg. Caucas.
Haussknechtii, *Boiss. Fl. Orient.* iv. 684.—Armenia.
havanensis, *Jacq. Enum. Pl. Carib.* 25.—Cuba.
hederacea, Kunth & Bouché, Ind. Sem. Hort. Berol. (1845) 10 = dependens?
heterophylla, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 45.—Cilicia; Syria.
heterophylla, Willd. ex Benth. in Linnaea, xi. (1837) 345 = rumicifolia.
heteropoda, Miq. Fl. Ind. Bat. ii. 972 = discolor.
Heydei, *Hook. f. Fl. Brit. Ind.* iv. 667.—Reg. Himal.
hirsuta, Kit. ex Sadl. Fl. Pest. ii. 83 = Columnae.
hirta, *Sibth. & Sm. Fl. Graec.* vi. 66. t. 583.—Graecia; Ins. Creta; Cyprus.
hirta, Sieber, ex Boiss. Fl. Orient. iv. 689 = Sieberi.

SCUTELLARIA :—
hirta, Sieber, ex Reichb. Fl. Germ. Excurs. 332 = albida.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 974.—Java.
humilis, *R. Br. Prod.* 507.—Austral.
hyssopifolia, Linn. Sp. Pl. 599 = integrifolia.
incana, Spreng. Fl. Hal. Mant. 44 = canescens.
incarnata, *Vent. Choix Pl.* 39. t. 39.—N. Granat.?
incurva, *Wall. Pl. As. Rar.* i. 67.—Burma.
indica, Blume, Bijdr. 839; D. Don, Prod. Fl. Nep. 109 = discolor.
indica, *Linn. Sp. Pl.* 600.—Japon; China; Java.
indica, Roxb. Hort. Beng. 45; Fl. Ind. iii. 24 = violacea.
integrifolia, *Linn. Sp. Pl.* 599.—Am. bor.
integrifolia, Pursh, Fl. Am. Sept. ii. 413 = serrata.
japonica, Burm. f. Fl. Ind. 130 = Plectranthus menthoides.
japonica, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 315 = dependens.
javanensis, *Jungh. Java*, i. 662, in adnot.—Java.
khasiana, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 669.—Reg. Himal.
lanceolaria, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 110 = baicalensis.
lateriflora, *Linn. Sp. Pl.* 598.—Am. bor.
Lehmanni, *Hort.—Cf. Gard. Chron.* (1883) II. 726 (seq.?).—Ecuador.
Lehmanni, *Regel, Gartenfl.* (1884) 129. t. 1152.—N. Granat.
leucodasys, *Miq. in Journ. Bot. Néerl.* i. (1861) 116. —China.
Lindeniana, *Benth. in DC. Prod.* xii. 421.—Venezuela.
linearis, *Benth. in Wall. Pl. As. Rar.* i. 66.—Reg. Himal.; Afghan.
Linnaeana, *Caruel, in Parl. Fl. Ital.* vi. 322.—Europ.; As. temp.; Abyss.
longifolia, *Benth. in Bot. Reg.* sub t. 1493. n. 5.—Mexic.
lupulina, Linn. Sp. Pl. 835 = alpina.
luridiflora, Fisch. & Mey. ex Benth. in DC. Prod. xii. 431, err. typ. = Scrophularia luridiflora.
luzonica, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 315. —Ins. Philipp.
macrantha, Fisch. ex Reichb. Ic. Bot. v. 52. t. 488 = baicalensis.
malvaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 324.—N. Granat.
microphylla, *Moç. & Sesse, ex Benth. in Bot. Reg.* sub t. 1493. n. 43.—Mexic.
minor, *Huds. Fl. Angl.* ed. I. 232.—Europ.; As. bor.
mississipiana, Martens, in Bull. Acad. Brux. viii. 1. (1841) 66 = versicolor.
Mociniana, *Benth. Lab. Gen. et Sp.* 442.—Mexic.
mollis, *R. Br. Prod.* 507.—Austral.
montana, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 11.—Am. bor.
Moriciniana, Carr. in Rev. Hortic. (1872) 350, sphalm. = Mociniana.
mucida, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 48. —Persia.
multicaulis, *Boiss. Diagn.* Ser. I. vii. 61.—Persia; Afghan.
multiflora, *Benth. in Bot. Reg.* sub t. 1493. n. 6.—Austral.
mussooriensis, Wawra, in Oestr. Bot. Zeitschr. xxiii. (1873) 266 = grossa.
nana, *A. Gray, in Proc. Am. Acad.* xi. (1876) 100.—Am. bor. occ.
nepetaefolia, Benth. in DC. Prod. xii. 414 = multicaulis.
nervosa, *Pursh, Fl. Am. Sept.* ii. 412.—Am. bor.
nigrescens, Spreng. Syst. ii. 702 = albida.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* i. 377.—Japon.
nodulosa, A. Ham. Monog. Scutell. 26 = violacea.
Novae-Zelandiae, *Hook. f. Fl. N. Zel.* ii. 335.—N. Zel.
nummularia, Fenzl, ex Boiss. Fl. Orient, iv. 687 = diffusa.
nummulariaefolia, *Hook. f. Fl. Antarct.* ii. 336.—Reg. Magellan.
oblonga, *Benth. in Bot. Reg.* sub t. 1493. n. 20.—Zeylan.
Oldhami, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 197. —Korea.
orbicularis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 436.—As. centr.
orientalis, *Linn. Sp. Pl.* 598.—Europ. austr.; Oriens.
ovalifolia, Pers. Syn. ii. 136 = pilosa, *Michx.*

SCUTELLARIA:—
ovata, Hill, Hort. Kew. 242 = parvula.
pallida, Bieb. Fl. Taur. Cauc. ii. 65 = albida.
pallida, Guss. Fl. Sic. Prod. ii. 137 = Columnae.
parnassica, Heldr. & Sart. ex Boiss. Fl. Orient. iv. 688 = peregrina.
parviflora, Rafin. ex A. Ham. Monog. Scutell. 37 = nervosa.
parvula, *Michx. Fl. Am. Bor.* i. 12.—Am. bor.
pauciflora, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 266 = galericulata.
pectinata, Hohen. ex Steud. Nom. ed. II. ii. 549 = orientalis.
pectinata, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Ser. II. vi. (1836) 44.—Cappadoc.
pekinensis, *Maxim. Prim. Fl. Amur.* 476.—China.
peregrina, *Linn. Sp. Pl.* 599.—Europ. austr.; Oriens.
peregrina, Roxb. Hort. Beng. 45; Fl. Ind. iii. 24 = rivularis.
peregrina, Waldst. & Kit. Pl. Rar. Hung. 132. ii. t. 125 = altissima.
physocalyx, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 68.—Kokania.
Pichleri, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 47.—Persia.
pilosa, *Hill, Veg. Syst.* xiii. 64.—Am. bor.
pilosa, *Michx. Fl. Bor. Am.* ii. 11.—Am. bor.
pilosiuscula, Nutt. ex Benth. in DC. Prod. xii. 429 = tuberosa.
pinnatifida, *A. Ham. Monog. Scutell.* 16. t. 2. f. 2.—Persia.
polymorpha, A. Ham. l. c. 38 = integrifolia, pilosa, *Michx.*
pontica, *C. Koch, in Linnaea*, xxi. (1848) 701.—Pontus.
prostrata, *Jacquem. ex Benth. Lab. Gen. et Sp.* 733.—Reg. Himal.
pubescens, Martr. Fl. Tarn, 572 = galericulata.
pubescens, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 173 = canescens.
pulchella, Bunge, Verz. Suppl. Fl. Alt. 57 = orientalis.
pulchella, *Hort. ex Heer, in Gartenfl.* (1857) 297. t. 205.—Hab.?
pumila, Salisb. Prod. 89 = minor.
purpurascens, *Sw. Prod. Veg. Ind. Occ.* 89.—Am. trop.
pygmaea, Willd. ex Benth. in Linnaea, xi. (1837) 345 = parvula.
racemosa, Pers. Syn. ii. 136 = rumicifolia.
radicata, Rafin. Atl. Journ. 16 (Quid?).—Am. bor.
repens, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 110.—Reg. Himal.; Burma.
reptans, Pall. ex Ledeb. Fl. Ross. iii. 398 = scordiifolia.
resinosa, *Torr. in Ann. Lyc. N. York*, ii. (1828) 232.—Am. bor. occ.
resinosa, S. Wats. in Bot. King's Exped. 237 = antirrhinoides.
rivularis, *Wall. Pl. As. Rar.* ii. 66.—Ind. or.; Burma; China.
robusta, Benth. in DC. Prod. xii. 418 = violacea.
rubicunda, Willd. Enum. Hort. Berol. Suppl. 42, nomen; et Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 56 = peregrina.
rugosa, Wood, in Proc. Am. Assoc. (1853) 176 = versicolor.
rumicifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 324.—Mexic.; Chili.
rupestris, Boiss. & Heldr. Diagn. Ser. I. vii. 60 = hirta.
russeliaefolia, *Vatke, in Bot. Zeit.* xxx. (1872) 715.—Ins. Philipp.
salviaefolia, *Benth. Lab. Gen. et Sp.* 433.—Graecia.
sanguinea, Steud. Nom. ed. II. ii. 549 = alpina.
saxatilis, *Riddell, Cat. Ohio Pl. Suppl.* (1836) 14.—Am. bor.
scandens, D. Don, Prod. Fl. Nep. 110 = angulosa.
scarlatina, *Planch. & Linden, Cat.* n. 10 (1855) 6; ex *Lem. Illustr. Hortic.* iii. (1856) t. 104.—Am. centr.
sciaphila, *S. Moore, in Journ. Bot.* xiii. (1875) 228.—China.
scordiifolia, *Fisch. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 55.—Sibir.
serrata, *Andr. Bot. Rep.* t. 494.—Am. bor.
serrata, Spreng. Syst. ii. 703 = canescens.
Sibthorpii, Boiss. & Reut. ex Boiss. Fl. Orient. iv. 688 = peregrina.
Sieberi, *Benth. in DC. Prod.* xii. 420.—Ins. Creta et Rodos.

SCUTELLARIA:—
Sieversii, Bunge, in Ledeb. Fl. Alt. ii. 394 = orientalis.
siphocampyloides, A. Gray, Syn. Fl. Am. ii. 1. 381, sphalm. = seq.
Siphocampylus, Vatke, in Bot. Zeit. xxx. (1872) 717 = angustifolia.
speciosa, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 389 = baicalensis.
spicata, *Trimen, in Journ. Bot.* xxiii. (1885) 241.—Zeylan.
splendens, *Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol.* i. 31. t. 13.—Mexic.
squamulosa, A. Ham. Monog. Scutell. 35 = rivularis.
Stocksii, *Boiss. Diagn.* Ser. II. iv. 28.—Beluchist.
sumatrana, *Miq. Fl. Ind. Bat.* ii. 974.—Sumatra.
supina, Linn. Sp. Pl. 598 = alpina.
Szovitziana, *Bunge, Lab. Pers.* 65.—Persia.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 464.—Japon.
tenerifolia, Sm. in Rees, Cycl. xxxii. n. 15 = nervosa.
tomentosa, Bertol. Misc. Bot. ii. 11 = fruticosa, *Desf.*
Tournefortii, *Benth. in Bot. Reg.* sub t. 1493. n. 15.—Persia; Reg. Caucas.
Trianaei, *Planch. & Linden, in Linden, Cat.* n. 10 (1855) 6; *et in Belg. Hortic.* vi. (1856) 232.—N. Granat.
tuberosa, *Benth. Lab. Gen. et Sp.* 441.—Calif.
tubiflora, *Benth. in Bot. Reg.* sub t. 1493. n. 10.—Bras.
uliginosa, *A. St. Hil. ex Benth. Lab. Gen. et Sp.* 427.—Bras.
utriculata, *Labill. Ic. Pl. Syr. Dec.* iv. 11. t. 6.—Syria.
variegata, Spreng. f. Tent. Suppl. Syst. 16 = alpina.
Ventenatii, *Hook. Bot. Mag.* t. 4271.—N. Granat.
verna, Bess. Prim. Fl. Galic. ii. 43 = alpina.
versicolor, *Nutt. Gen. Am.* ii. 38.—Am. bor.
Vigineixii, *Marès & Vigin. Cat. Pl. Balear.* 220.—Ins. Majorc.
villosa, *Ell. Sketch*, ii. 90.—Am. bor.
villosa, *Hook. Bot. Mag.* t. 4789.—Bras.
villosa, Rafin. Atl. Journ. 17 (Quid?).—Am. bor.
violacea, *Heyne, ex Wall. Pl. As. Rar.* i. 66.—Ind. or.; Burma; Java.
virens, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 682.—Kurdistan.
viscida, Spreng. f. Tent. Syst. Suppl. 15 = alpina.
viscidula, *Bunge, Enum. Pl. Chin. Bor.* 52.—China.
volubilis, *H. B. & K. Nov. Gen. et Sp.* ii. 325.—N. Granat.
Wallichiana, A. Ham. Monog. Scutell. 28. t. 2. f. 1 = repens.
Wightiana, Benth. in Wall. Pl. As. Rar. i. 67 = violacea.
Wrightii, *A. Gray, in Proc. Am. Acad.* viii. (1870) 370.—Am. bor.

SCUTIA, Comm. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 362. *RHAMNEAE*, Benth. & Hook. f. i. 379.
BLEPETALON, Rafin. Sylva Tellur. 30 (1838).
arenicola, *Reiss. in Mart. Fl. Bras.* xi. 1. 93.—Bras.
buxifolia, *Reiss. l. c.*—Reg. Argent.
capensis, G. Don, Gen. Syst. ii. 33 = Commersonii.
Commersonii, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 363.—As. trop.; Ins. Mascar.
discolor, Klotzsch, in Peters, Reise Mossamb. Bot. 110 = Berchemia discolor.
ferrea, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 363.—Ind. occ.
ferrea, Chapm. Fl. S. U. St. 72 = Reynosia latifolia.
indica, Brongn. in Ann. Sc Nat. Sér. I. x. (1827) 363 = Commersonii.
laevigata, G. Don, Gen. Syst. ii. 34 = Reynosia latifolia.
lucida, G. Don, l. c. 33 = Commersonii.
myrtina, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 168 = Commersonii.
natalensis, Hochst. in Flora, xxvii. (1844) 346 = Commersonii.
obcordata, *Bivona, ex Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 116.—Madag.
paniculata, G. Don, Gen. Syst. ii. 34 = Celastrus paniculatus.
pauciflora, G. Don, l. c. = Ceanothus pauciflorus.
Rheediana, Wight, Ic. t. 1071 = Commersonii.
Sarcomphalus, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 364 = Sarcomphalus retusus.
sphaerosperma, G. Don, Gen. Syst. ii. 34 = Rhamnus sphaerosperma.

SCUTICARIA, Lindl. Bot. Reg. (1843). Misc. 14 *ORCHIDEAE*, Benth. & Hook. f. iii. 554.
Dodgsoni, Hort. ex Reichb. f. in Gard. Chron. (1881) I. 9 = Hadwenii?
Hadwenii, *Hort. ex Planch. in Fl. des Serres*, vii. (1851–52) 239.—Bras.
Steelii, *Lindl. Bot. Reg.* (1843) *Misc.* 14.—Guian.

SCUTINANTHE, Thw. in Hook. Kew Journ. viii. (1856) 266. t. 8 = **Canarium**, Linn. (Burser.).
brunnea, Thw. l. c. 267 = C. brunneum.

SCUTULA, Lour. Fl. Cochinch. 235 (1790) = **Memecylon**, Linn. (Melastom.).
scutellata, Lour. l. c. = M. edule.
umbellata, Lour. l. c. 236 = M. umbellatum.

SCYBALIUM, Schott & Endl. Melet. 3. t. 2 (1832). *BALANOPHOREAE*, Benth. & Hook. f. iii. 237.
PHYLLOCORYNE, Hook. f. in Trans. Linn. Soc. xxii. (1856) 51. t. 11.
SPHAERORHIZON, Hook. f. l. c. 50. t. 10 (1856).
depressum, *Eichl. in DC. Prod.* xvii. 133.—N. Granat.
fungiforme, *Schott & Endl. Melet.* 3. t. 2.—Bras.
Glaziovii, *Eichl. in DC. Prod.* xvii. 133.—Bras.
jamaicense, *Schott & Endl. Melet.* 12.—Ind. occ.

SCYNOPSOLE, Reichb. Handb. 164 (1837) = CYNOPSOLE, Endl. = **Balanophora**, Forst.

SCYPHAEA, Presl, Scyph. Nov. Pl. Gen. (1829); et Symb. Bot. i. 7. t. 4 (1830) = **Marila**, Sw. (Ternstr.).
racemosa, Presl, ll. cc. = M. racemosa.

SCYPHANTHUS, [D. Don, in] Sweet, Brit. Flow. Gard. Ser. I. t. 238 (1828) = **Grammatocarpus**, Presl (Loaseae).
Cumingii, Phil. f. Cat. Pl. Vasc. Chil. 84 = G. Cumingii.
elegans, [D. Don, in] Sweet, Brit. Flow. Gard. Ser. I. t. 238 = G. volubilis.
grandiflorus, Sweet, ex Steud. Nom. ed. II. ii. 569 = G. volubilis.

SCYPHARIA, Miers, in Ann. & Mag. Nat. Hist. Sér. III. vi. (1860) 8 = **Colletia**, Comm. (Rhamn.).
guayaquilensis, Miers, l. c. 9 = Rhamnus guayaquilensis.
parviflora, Miers, l. c. 10 = Discaria parviflora.
senticosa, Miers, l. c. 11 = Rhamnus senticosa.
tetragona, Miers, l. c. 12 = C. tetragona.

SCYPHELLANDRA, Thw. Enum. Pl. Zeyl. 21 (1864). *VIOLARIEAE*, Benth. & Hook. f. i. 120.
virgata, *Thw. l. c.*—Zeylan.

SCYPHIPHORA, Gaertn. f. Fruct. iii. 91. t. 196 (1805). *RUBIACEAE*, Benth. & Hook. f. ii. 99.
EPITHINIA, Jack, in Malay. Misc. i. (1820) v. 12.
hydrophyllacea, *Gaertn. f. Fruct.* iii. 91. t. 196.—Ins. Molucc.
malayana, *Bedd. Fl. Sylv. Anal. Gen.* t. 29. f. 5.—Malay.

SCYPHOCHLAMYS, Balf. f. in Journ. Linn. Soc. xvi. (1877) 14; et in Baker, Fl. Maurit. 149 (1877). *RUBIACEAE.*
revoluta, *Balf. f. ll. cc.* 15; 149.—Ins. Rodrig.

SCYPHOCORONIS, A. Gray, in Hook. Kew Journ. iv. (1852) 225. *COMPOSITAE*, Benth. & Hook. f. ii. 316.
viscosa, *A. Gray, l. c.*—Austral.

SCYPHOGLOTTIS, Pritz. Ic. Ind. i. 1012 (1858), sphalm. = **Scaphyglottis**, Poepp. & Endl. (Orchid.).

SCYPHOGYNE, Brongn. Voy. Coq. Bot. t. 54 (1828?). *ERICACEAE*, Benth. & Hook. f. ii. 594.
BLEPHARISPERMUM, Klotzsch, ex Benth. in DC. Prod. vii. 709 (1839).
BLEPHAROPHYLLUM, Klotzsch, in Linnaea, xii. (1838) 216.
CODONOSTIGMA, Klotzsch, ex Benth. in DC. Prod. vii. 709 (1839).
COILOSTIGMA, Klotzsch, in Linnaea, xii. (1838) 234.

SCYPHOGYNE:—
OMPHALOCARYON, Klotzsch, in Linnaea, xii. (1838) 243.
THAMNIUM, Klotzsch, l. c. 223 (1838).
TRISTEMON, Klotzsch, l. c. 245 (1838).
brevifolia, *Benth. in DC. Prod.* vii. 710.—Afr. austr.
capitata, *Benth. l. c.*—Afr. austr.
divaricata, *Benth. l. c.*—Afr. austr.
eglandulosa, *Benth. l. c.*—Afr. austr.
fasciculata, *Benth. l. c.* 709.—Afr. austr.
inconspicua, *Brongn. Voy. Coq. Bot.* t. 54.—Afr. austr.
muscosa, Steud. Nom. ed. II. i. 568, in syn. = praec.
puberula, *Benth. in DC. Prod.* vii. 709.—Afr. austr.
urceolata, *Benth. l. c.*—Afr. austr.

SCYPHONYCHIUM, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 495. *SAPINDACEAE.*
multiflorum, *Radlk. l. c.* 519, 545.—Bras.

SCYPHOPETALUM, Hiern, in Hook. f. Fl. Brit. Ind. i. 675 (1875). *SAPINDACEAE.*
ramiflorum, *Hiern, l. c.* 676.—Burma.

SCYPHOSTACHYS, Thw. Enum. Pl. Zeyl. 157 (1859). *RUBIACEAE*, Benth. & Hook. f. ii. 94.
coffeoides, *Thw. l. c.*—Zeylan.
pedunculatus, *Thw. l. c.*—Zeylan.

SCYPHOSTIGMA, M. Roem. Syn. Hesper. 80, 94 (1846) = **Turraea**, Linn. (Meliac.).
Bennetii, M. Roem. l. c. 94 = Quivisia tetramera.
philippinense, M. Roem. l. c. = T. octandra.

SCYPHOSYCE, Baill. Adansonia, xi. (1875) 293. *URTICACEAE*, Benth. & Hook. f. iii. 371.
Manniana, *Baill. l. c.*—Afr. trop.

SCYRTOCARPA, Miers, in Journ. Linn. Soc. xvii. (1879) 292 = seq.

SCYRTOCARPUS, Miers, in Lindl. Veg. Kingd. 593 a (1847) = **Symplocos**, Linn. (Styrac.).

SCYTALA, E. Mey. ex DC. Prod. vii. 12 (1838) = **Oldenburgia**, Less. (Compos.).
Papionum, E. Mey. l. c. = O. Papionum.

SCYTALANTHUS, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 361 = **Skytanthus**, Meyen (Apocyn.).
acutus, Schau. l. c. 362 = Skytanthus acutus.

SCYTALIA, Gaertn. Fruct. i. 197. t. 42 (1788) = **Nephelium**, Linn. (Sapindac.).
benghalensis, Roxb. ex Wight & Arn. Prod. 113 = N. Longana.
chinensis, Gaertn. Fruct. i. 197. t. 42. f. 3 = N. Lit-chi.
crinita, Raeusch. Nom. ed. III. 113; Stokes, Bot. Mat. Med. ii. 386 = N. lappaceum.
Danura, Roxb. Hort. Beng. 29; Fl. Ind. ii. 274 = Sapindus Danura.
glabra, Buch.-Ham. ex Wall. Cat. n. 8048 E = Walsura robusta.
informis, Raeusch. Nom. ed. III. 113 = Euphoria informis.
Litschi, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 269 = N. Lit-chi.
Longan, Roxb. ll. cc. 29; 270 = N. Longana.
oppositifolia, Roxb. ll. cc. [88]; 273 = N. oppositifolium.
parviflora, Roxb. ll. cc. = N. parviflorum.
Ramboutan, Roxb. ll. cc. 29; 271 = N. lappaceum.
rimosa, Roxb. ll. cc. 29; 272 = N. rimosum.
rubia, Roxb. ll. cc. = Sapindus attenuatus.
sinensis, J. F. Gmel. Syst. 612 = *chinensis*.
squamosa, Stokes, Bot. Mat. Med. ii. 385 = N. Lit-chi.
trijuga, Roxb. ex DC. Prod. i. 615 = Schleichera trijuga.
verruculosa, Stokes, Bot. Mat. Med. ii. 386 = N. Longana.
verticillata, Roxb. Hort. Beng. 29; Fl. Ind. ii. 273 = Sapindus Danura.

SCYTALIS, E. Mey. Comm. Pl. Afr. Austr. 144 (1835) = **Vigna**, Savi (Legumin.).
anomala, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 32 = V. retusa.
Catjang, E. Mey. Comm. Pl. Afr. Austr. 145 = V. Catjang.
Helicopus, E. Mey. l. c. 146 = V. glabra.
hispida, E. Mey. l. c. = V. hispida.
protracta, E. Mey. l. c. = V. triloba.
retusa, E. Mey. l. c. 147 = V. retusa.
tenuis, E. Mey. l. c. 145 = V. tenuis, *D. Dietr.*

SCYTANTHUS, T. Anders. ex Benth. & Hook. f. Gen. ii. 1093 (1876). *ACANTHACEAE*, Benth. & Hook. f. ii. 1093.
laurifolius, *T. Anders. l. c.*—Afr. trop.

SCYTANTHUS, Hook. Ic. Pl. t. 605–625 (184) = **Hoodia**, Sweet (Asclep.).
Burkei, Hook. l. c. sub t. 625 = H. Currori.
Currori, Hook. l. c. t. 605–606 = H. Currori.
Gordoni, Hook. l. c. t. 625 = H. Gordoni.

SCYTANTHUS, Liebm. in Foerhandl. Skand. Naturf. 4de Mode (1844), 177 = **Cytinus**, Tourn.
Bambusarum, Liebm. l. c. 183 = C. americanus.

SCYTOPHYLLUM, Eckl. & Zeyh. Enum. 124 (1835) = **Elaeodendron**, Jacq. f. (Celastrin.).
angustifolium, Sond. in Harv. & Sond. Fl. Cap. i. 472 = E. angustifolium.
laurinum, Eckl. & Zeyh. Enum. 124 = Celastrus laurinus.
obtusum, Eckl. & Zeyh. l. c. = Celastrus laurinus.
oleoides, Eckl. & Zeyh. l. c. = Celastrus oleoides.

SCZEGLEEWIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 233 = **Pterospermum**, Schreb. (Stercul.).
involucrata, Turcz. l. c. 234 = P. obliquum.

SCZEGLEEWIA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 212 = **Symphorema**, Roxb. (Verben.).
luçonensis, Turcz. l. c. 213 = Sym. luzonicum.

SCZUKINIA, Turcz. in Bull. Soc. Nat. Mosc. (1840) 165 = **Swertia**, Linn. (Gentian.).
diluta, Turcz. l. c. 166 = Sw. diluta.

SEAFORTHIA, R. Br. Prod. 267 (1810) = **Ptychosperma**, Labill., etc. (Palm.).
angustifolia, Mart. Hist. Nat. Palm. iii. 314.—N. Guin.
appendiculata, Juss. ex Kunth, Enum. Pl. iii. 192 = Drymophloeus appendiculatus.
Blumei, Juss. ex Kunth, l. c. = Drymophloeus olivaeformis, leprosus.
caesia, Mart. Hist. Nat. Palm. iii. 313 = Pinanga caesia.
Calapparia, Mart. l. c. = Actinorhytis Calapparia.
cochinchinensis, Mart. l. c. = Pinanga cochinchinensis.
communis, Blume, Rumphia, ii. 78 = Drymophloeus communis.
coronata, Mart. Hist. Nat. Palm. iii. 185 = Pinanga coronata.
costata, Mart. l. c. 313 = Pinanga Kuhlii.
Dicksoni, Mart. l. c. 184 = Pinanga Dicksonii.
disticha, Mart. l. c. = Pinanga disticha.
elegans, R. Br. Prod. 267 = Ptychosperma elegans.
elegans, Hook. Bot. Mag. t. 4961 = Archontophoenix Cunninghami.
furfuracea, Mart. Hist. Nat. Palm. iii. 313 = Pinanga furfuracea.
gracilis, Mart. l. c. 184 = Pinanga gracilis.
inaequalis, Mart. l. c. 313 = Pinanga inaequalis.
jaculatoria, Mart. l. c. 314 = Drymophloeus appendiculatus.
Kuhlii, Mart. l. c. 185. t. 158. f. 4 = Pinanga Kuhlii.
latisecta, Mart. l. c. 313 = Pinanga latisecta.
malaiana, Mart. l. c. 183. t. 158. f. 3 = Pinanga malaiana.
minor, Mart. l. c. 313 = Pinanga minor.
montana, Blume, in Mart. l. c. 185 = Pinanga coronata.
olivaeformis, Mart. l. c. 314 = Drymophloeus olivaeformis.
oryzaeformis, Mart. l. c. 185 = Pinanga Kuhlii.
patula, Mart. l. c. 313 = Pinanga patula.

SEAFORTHIA:—
Ptychosperma, Mart. Hist. Nat. Palm. iii. 182. tt. 128, 129 = Ptychosperma gracilis.
Reinwardtiana, Mart. l. c. 183. t. 158. f. 2 = Pinanga noxa.
robusta, Hort. ex H. Wendl. in Kerch. Palm. 257 = Rhopalostylis Baueri.
Rumphiana, Mart. Hist. Nat. Palm. iii. 134 = Pinanga ternatensis.
salicifolia, Mart. l. c. 313 = Pinanga salicifolia.
saxatilis, Blume, ex Mart. l. c. 134. t. 158. f. 6 = Drymophloeus saxatilis.
sylvestris, Blume, l. c. 185 = Pinanga javana.
Veitchii, Hort.—Cf. Gard. Chron. (1870) 37.—Austral.
vestiaria, Mart. Hist. Nat. Palm. iii. 313 = Mischophloeus paniculatus.

SEALA, Adans. Fam. ii. 131 (1763) = **Pectis**, Linn. (Compos.).

SEBAEA, Soland. ex R. Br. Prod. 451 (1810). *GENTIANEAE*, Benth. & Hook. f. ii. 804.
LAGENIAS, E. Mey. Comm. Pl. Afr. Austr. 186 (1837).
albens, *R. Br. Prod.* 452.—Afr. austr.
albidiflora, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 46.—Austral.
ambigua, *Cham. in Linnaea*, vi. (1831) 346.—Afr. austr.
aurea, *R. Br. Prod.* 452.—Afr. austr.
Boyeri, *Griseb. Gent. Gen. et Sp.* 169.—Madag.
brachyphylla, *Griseb. l. c.* 170.—Madag.
brasiliensis, Spreng. Syst. iv. Cur. Post. 338 = Schultesia crenuliflora.
capitata, *Schlecht. in Linnaea*, i. (1826) 193.—Afr. austr.
capitata, Schrad. ex Griseb. in DC. Prod. ix. 52 = congesta.
carinata, Spreng. Syst. i. 426 = Exacum petiolare.
congesta, *Schrad. ex Griseb. in DC. Prod.* ix. 52.—Afr. austr.
cordata, *R. Br. Prod.* 452.—Afr. austr.
crassulaefolia, *Schlecht. in Linnaea*, i. (1826) 193.—Afr. austr.
cymosa, Jarosz, Pl. Nov. Cap. 10 = aurea.
elongata, *E. Mey. Comm. Pl. Afr. Austr.* 184.—Afr. austr.
emirnensis, Boj. ex Griseb. in DC. Prod. ix. 53 = brachyphylla.
gracilis, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 45 = ovata.
gracilis, Spreng. Syst. iv. Cur. Post. 338 = Schultesia gracilis.
grandis, *Steud. Nom.* ed. II. ii. 550.—Afr. austr.
guianensis, Spreng. Syst. i. 426 = Schultesia stenophylla.
involucrata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 271.—Afr. trop.
jorullensis, Spreng. Syst. i. 426 = Schultesia stenophylla.
khasiana, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 428 = aurea.
minima, Jarosz, Pl. Nov. Cap. 11 = aurea.
Ohlendorffiana, Eckl. ex Griseb. Gen. et Sp. Gent. 166 = Belmontia Ohlendorffii.
oldenlandioides, S. Moore, in Journ. Bot. xv. (1877) 68 = Exacum quinquenervium.
ovata, *R. Br. Prod.* 452.—Austral.; N. Zel.
pallida, *E. Mey. Comm. Pl. Afr. Austr.* 185.—Afr. austr.
pentandra, *E. Mey. l. c.* 184.—Afr. austr.
pusilla, *Cham. in Linnaea*, vi. (1831) 346.—Afr. austr.
Rutenbergiana, *Vatke, in Bremen, Abh.* ix. (1885) 126.—Madag.
spathulata, *Steud. Nom.* ed. II. ii. 550.—Afr. austr.
sulcata, Spreng. Syst. i. 426 = Exacum pedunculatum.
sulphurea, *Schlecht. in Linnaea*, i. (1826) 192.—Afr. austr.
tabularis, Eckl. ex Steud. Nom. ed. II. ii. 550 = sulphurea.

SEBASTIANA, Benth. & Hook. f. Gen. ii. 389 (1873), err. typ. = SEBASTIANIA, Bertol. = **Chrysanthellum**, Rich. (Compos.).

SEBASTIANA, Spreng. Syst. iii. 24 (1826) = **Sebastiania**, Spreng. (Euphorb.).

SEBASTIANIA, Bertol. Lucubr. 37 (1822) = **Chrysanthellum**, Rich. (Compos.).
heterophylla, Bertol. l. c. = C. procumbens.

SEBASTIANIA, Spreng. Neue Entdeck. ii. 118. t. 3 (1821). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 336.
ADENOGYNE, Klotzsch, in Wiegm. Archiv, vii. (1841) 183.
CLONOSTACHYS, Klotzsch, l. c. 185 (1841).
CNEMIDOSTACHYS, Mart. Nov. Gen. et Sp. i. 66. t. 40 (1824).
DITRYSINIA, Rafin. Neogenyt. 2 (1825).
ELACHOCROTON, F. Muell. in Hook. Kew Journ. ix. (1857) 17.
GUSSONIA, Spreng. Neue Entdeck. ii. 119. t. 2 (1821).
MICROSTACHYS, A. Juss. Euph. Tent. 48. t. 15 (1824).
SAROTHROSTACHYS, Klotzsch, in Wiegm. Archiv, vii. (1841) 185.
TRAGIOPSIS, Karst. in C. Koch, Wochenschr. (1859) 5.
actinostemoides, Muell. Arg. in DC. Prod. xv. II. 1184 = Gymnanthes actinostemoides.
albicans, *Wright, in Sauv. Fl. Cub.* 132.—Cuba.
angustifolia, *Muell. Arg. in DC. Prod.* xv. II. 1179.—Bras.
anisodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 556.—Bras.
bahiensis, *Muell. Arg. in DC. Prod.* xv. II. 1183.—Bras.
bilocularis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 374.—N. Mexic.
brachyclada, *Muell. Arg. in DC. Prod.* xv. II. 1178.—Bras.
brachypoda, *Wright, in Sauv. Fl. Cub.* 132.—Cuba.
brasiliensis, *Spreng. Neue Entdeck.* ii. 118.—Bras.
brevifolia, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183.—Bras.
chaetodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 577.—Bras.
Chamaelea, *Muell. Arg. in DC. Prod.* xv. II. 1175.—Geront. trop.
corniculata, *Muell. Arg. l. c.* 1168.—Bras.
daphniphylla, *Muell. Arg. l. c.* 1180.—Bras.
daphnoides, *Muell. Arg. l. c.* 1167.—Bras.
desertorum, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183, nomen.—Bras.
dimorphocalyx, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 574.—Bras.
discolor, *Muell. Arg. in DC. Prod.* xv. II. 1185.—Bras.
ditassoides, *Muell. Arg. l. c.* 1174.—Bras.
divaricata, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183, nomen.—Bras.
echinocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 584.—Bras.
elliptica, *Muell. Arg. in DC. Prod.* xv. II. 1181.—Jamaic.
foveata, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183, nomen.—Bras.
Gaudichaudii, *Muell. Arg. in DC. Prod.* xv. II. 1177.—Bras.
glandulosa, Muell. Arg. l. c. 1186 = Gymnanthes glandulosa.
glaucophylla, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 571.—Bras.
granatensis, *Muell. Arg. in DC. Prod.* xv. II. 1189.—N. Granat.
Grisebachiana, *Muell. Arg. l. c.* 1183.—Ind. occ.
guyanensis, *Muell. Arg. l. c.*—Guian.
heteroica, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 708.—Bras.
hypoleuca, Muell. Arg. in DC. Prod. xv. II. 1184 = Gymnanthes hypoleuca.
jacobinensis, *Muell. Arg. l. c.* 1188.—Bras.
Klotzschiana, Muell. Arg. l. c. 1178 = Gymnanthes marginata.
Laureola, *Muell. Arg. l. c.* 1180.—Bras.
ligustrina, *Muell. Arg. l. c.* 1165.—Am. bor.
longipes, Muell. Arg. l. c. 1184 = Gymnanthes longipes.
lucida, Muell. Arg. l. c. 1181 = Gymnanthes lucida.
macrocarpa, *Muell. Arg. l. c.* 1188.—Bras.
macrophylla, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183, nomen.—Bras.
marginata, *Muell. Arg. in DC. Prod.* xv. II. 1166.—Bras.

SEBASTIANIA:—
membranifolia, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 579.—Bras.
mucronata, Muell. Arg. in DC. Prod. xv. II. 1180 = Schottiana.
multiramea, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 53; *Muell. Arg. in DC. Prod.* xv. II. 1177.—Bras.; Afr. occ.
myricifolia, *Wright, in Sauv. Fl. Cub.* 132.—Cuba.
nervosa, *Muell. Arg. in DC. Prod.* xv. II. 1183.—Bras.
oleoides, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 548.—Bras.
pachystachys, *Muell. Arg. in DC. Prod.* xv. II. 1182.—Bras.
pallens, Muell. Arg. l. c. 1189 = Gymnanthes pallens.
Pavoniana, *Muell. Arg. l. c.*—Mexic.
phyllanthiformis, Muell. Arg. l. c. 1176 = Schottiana.
pteroclada, *Muell. Arg. l. c.* 1190.—Bras.
reticulata, *Klotzsch, in Wiegm. Archiv*, vii. (1841) 183.—Bras.
rhombifolia, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 590.—Bras.
Riedelii, *Muell. Arg. l. c.* 578.—Bras.
rigida, *Muell. Arg. in DC. Prod.* xv. II. 1180.—Bras.
riparia, Muell. Arg. l. c. 1185 = Excoecaria potamophila.
riparia, Schrad. in Goett. Gel. Anz. i. (1821) 713 = Gymnanthes riparia.
Schlechtendaliana, Muell. Arg. in DC. Prod. xv. II. 1181 = Gymnanthes riparia.
Schottiana, *Muell. Arg. l. c.* 1176.—Bras.
serrata, *Muell. Arg. l. c.* 1179.—Bras.
serrulata, *Muell. Arg. l. c.* 1167.—Bras.
stipulacea, *Muell. Arg. l. c.* 1176.—Bras.
Treculiana, *Muell. Arg. l. c.* 1165.—Am. trop.
trinervia, *Muell. Arg. l. c.* 1182.—Bras.
vestita, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 575.—Bras.
viminea, Nees, in Flora, iv. (1821) 328 = Gymnanthes riparia.
virgata, *Muell. Arg. in DC. Prod.* xv. II. 1173.—Bras.
Weddelliana, *Muell. Arg. l. c.* 1188.—Bras.
Widgreni, *Muell. Arg. l. c.* 1178.—Bras.
ypanemensis, *Muell. Arg. l. c.* 1179.—Bras.

SEBASTIANO-SCHAUERIA, Nees, in Mart. Fl. Bras. ix. 158 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1098.
oblongata, *Nees, l. c.* 159.—Bras.

SEBEEKIA, Steud. Nom. ed. II. ii. 550 (1841), sphalm. = seq.

SEBEOKIA, Neck. Elem. ii. 13 (1790) = **Gentiana**, Linn.

SEBESTEN, Adans. Fam. ii. 177 (1763) = seq.

SEBESTENA, [Dill.] Gaertn. Fruct. i. 364. t. 76. f. 1 (1788) = **Cordia**, Linn. (Borag.).
glomerata, Lem. Jard. Fleur. iv. (1854) sub t. 326 = C. glomerata.
indica, Rafin. Sylva Tellur. 38 = C. Myxa.
officinalis, Gaertn. Fruct. i. 364. t. 76 = C. Myxa.
repanda, Rafin. Sylva Tellur. 38 (Quid?).—Hab.?
scabra, Rafin. l. c. = C. scabra.
senegalensis, Rafin. l. c. 39 = C. senegalensis.

SEBIFERA, Lour. Fl. Cochinch. 637 (1790) = **Litsea**, Lam. (Laurin.).
Balongay, Blanco, Fl. Filip. ed. I. 820 = L. ligustrina.
glutinosa, Blanco, l. c. 819 = L. Sebifera?
glutinosa, Lour. Fl. Cochinch. 638 = L. Sebifera.

SEBIPIRA, Mart. Reise Bras. ii. 787 (1828) = **Bowdichia**, H. B. & K. (Legumin.).
ferruginea, Vog. in Linnaea, xi. (1837) 387 = B. virgilioides.
major, Mart.? ex Vog. l. c. 386 = B. virgilioides.

SEBIZIA, Mart. ex Meissn. Gen. Comm. 376 (1843). Gen. *AMPELIDEARUM*, nondum descr.

SEBOPHORA, Neck. Elem. ii. 188 (1790) = **Myristica**, Linn.

SEBORIUM, Rafin. Sylva Tellur. 63 (1838) = **Stillingia**, Linn. (Euphorb.).
chinense, Rafin. l. c. = Sapium sebiferum.
nutans, Rafin. l. c. = St. nutans.

SECALE, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 17 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1203.
aestivum, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 368 = cereale?
anatolicum, Boiss. Diagn. Ser. I. v. 76 = cereale.
arundinaceum, *Trautv. Schilfroggen* (1840), t. 1, 2; *ex Steud. Syn. Pl. Gram.* 347.—Hab.?
barbatum, Moench, Meth. Suppl. 59 = cereale.
bromoides, Linn. Herb. ex Munro, in Journ. Linn. Soc. vi. (1862) 47 = Brachypodium distachyum.
campestre, Kit. ex Schult. Oestr. Fl. ed. II. i. 197 = fragile.
cereale, Bieb. Fl. Taur. Cauc. i. 84 = fragile.
cereale, *Linn. Sp. Pl.* 84.—Oriens.
compositum, Poir. Encyc. vii. 54 = cereale.
cornutum, Offic. ex Nees, Duesseld. Samml. Suppl. t. 74; ex Pritz. Ic. Ind. i. 1013 = Claviceps purpurea (Crypt.).
creticum, Linn. Sp. Pl. 84 = Triticum creticum.
creticum, Sieber, ex Kunth, Enum. Pl. i. 449 = cereale.
dalmaticum, *Vis. Fl. Dalm.* i. 97.—Dalmat.
fragile, *Bieb. Fl. Taur. Cauc. Suppl.* 93.—Europ.; As. temp.
glaucum, Urv. in Mém. Soc. Linn. Par. i. (1822) 268 = fragile.
hirtum, Poir. Encyc. vii. 55 = Agropyron orientale.
hybernum, Poir. l. c. 54 = cereale.
montanum, Guss. Ind. Sem. Hort. Boccad. (1825); ex ej. Fl. Sic. Prod. i. 145 = cereale.
orientale, Linn. Sp. Pl. 84 = Agropyron orientale.
orientale, Schreb. ex Roth, Nov. Pl. Sp. 59 = Hordeum bifarium.
prostratum, Pall. Reise, i. 485 = Agropyron triticeum.
pumilum, Pers. Syn. i. 109 = Agropyron triticeum.
pungens, Pers. l. c. 108 = Agropyron patulum.
reptans, Pall. Reise, i. 168 = Agropyron triticeum.
sibiricum, *Georgi, Beschr. Russ. Reich.* III. iv. 714.—Sibir.
spontaneum, Fisch. ex Steud. Nom. ed. I. 757 = fragile.
strictum, Presl, Fl. Sic. p. xlvi = cereale.
sylvestre, Host, Gram. iv. 7. t. 11 = fragile.
triflorum, Beauv. Agrost. 105, 178 (Index) = cereale.
vernum, Poir. Encyc. vii. 54 = cereale.
villosum, Huds. Fl. Angl. ed. I. 46 = Elymus europaeus.
villosum, Linn. Sp. Pl. 84 = Agropyron villosum.

SECALIDIUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91 = **Agropyron**, J. Gaertn. (Gramin.).
pectinatum, Schur, Enum. Pl. Transs. 807 = A. cristatum.
pilosum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91 = A. cristatum.

SECAMONE, R. Br. in Mem. Wern. Soc. i. (1809) 55. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 746.
acutifolia, *Sond. in Linnaea*, xxiii. (1850) 76.—Afr. austr.
aegyptiaca, Ait. Hort. Kew. ed. II. ii. 75 = Oxystelma esculentum.
Alpini, Schult. Syst. vi. 125 = Oxystelma esculentum.
amygdalina, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 313.—Java; Sumatra.
angustifolia, *Decne. in DC. Prod.* viii. 502.—Madag.
attenuata, Decne. l. c. 501 = elliptica.
auriculata, *Blume, Bijdr.* 1051.—Java.
bicolor, *Decne. in DC. Prod.* viii. 503.—Madag.
Blumei, *Decne. l. c.* 502.—Java.
buxifolia, *Decne. l. c.*—Madag.
canescens, Sm. in Rees, Cycl. xxxii. n. 3 = Leptadenia reticulata.
crassifolia, Wall. ex Decne. in DC. Prod. viii. 506 = Genianthus crassifolius.
elliptica, *R. Br. Prod.* 464.—Austral.; Ins. Philipp.
emetica, *R. Br. in Mem. Wern. Soc.* i. (1809) 56.—Ind. or.
emetica, F. Muell. Fragm. v. 161 = elliptica.
Finlaysoni, *Wight, Contrib.* 61.—Siam.
frutescens, Decne. in DC. Prod. viii. 501 = Astephanus frutescens.
fulva, Blume, ex Decne. l. c. 506 = Toxocarpus Blumei.
insularis, *Miq. Fl. Ind. Bat. Suppl.* 557.—Ins. Sunda.

SECAMONE:—
lanceolata, *Blume, Bijdr.* 1051.—Java.
ligustrifolia, *Decne. in DC. Prod.* viii. 502.—Madag.
lineata, *Blume, Bijdr.* 1051.—Java.
macrophylla, *Blume, l. c.* 1050.—Java.
macrostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) I. 252.—Ins. Philipp.
maritima, *Blume, Bijdr.* 1050.—Java.
mauritiana, G. Don, in Sweet, Hort. Brit. ed. III. 469 = Tylophora laevigata.
micrantha, *Decne. in DC. Prod.* viii. 501.—Ins. Timor.
multiflora, *Decne. l. c.*—Ins. Philipp.
myrtifolia, *Benth. in Hook. Niger Fl.* 453.—Afr. trop.
obovata, *Decne. in DC. Prod.* viii. 503.—Madag.
occidentalis, *Spreng. Neue Entdeck.* iii. 34.—Ind. occ.
oleaefolia, *Decne. in DC. Prod.* viii. 503.—Madag.
ovata, *R. Br. Prod.* 464.—Austral.
punctulosa, *Decne. in DC. Prod.* viii. 502.—Afr. trop.
saligna, *Decne. l. c.* 503.—Ins. Mascar.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 79.—Ins. Socotra.
tenuifolia, *Decne. in DC. Prod.* viii. 503.—Madag.
Thouarsii, *Decne. l. c.* 502.—Madag.
Thunbergii, *E. Mey. Comm. Pl. Afr. Austr.* 220.—Afr. austr.
timorensis, *Decne. in DC. Prod.* viii. 502.—Ins. Timor.
uniflora, *Decne. l. c.* 503.—Madag.
vestita, *Zipp. ex Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
villosa, Blume, Bijdr. 1050 = Toxocarpus villosus.

SECHIOPSIS, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 23. *CUCURBITACEAE*, Benth. & Hook. f. i. 838.
triqueter, *Naud. l. c.* 24.—Mexic.

SECHIUM, P. Br. Hist. Jamaic. 355 (1756). *CUCURBITACEAE*, Benth. & Hook. f. i. 837.
CHAYOTA, Jacq. Select. Am. ed. Pict. t. 245 (1780).
CHOCHO, Adans. Fam. ii. 500 (1763).
amazonicum, Poepp. & Endl. Nov. Gen. et Sp. ii. 57 = Cayaponia amazonica.
americanum, Lam. Encyc. vii. 50 = edule.
Chayota, Hemsl. Biol. Centr. Am. Bot. i. 491, in syn. = edule.
edule, *Sw. Fl. Ind. Occ.* ii. 1150.—Ind. occ.
palmatum, Ser. in DC. Prod. iii. 313 = Microsechium palmatum.
peruvianum, Poepp. & Endl. Nov. Gen. et Sp. ii. 56. t. 176 = Cayaponia peruvianum.

SECONDATIA, A. DC. in DC. Prod. viii. 445 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 723.
arborea, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 110.—Bras.
densiflora, *A. DC. in DC. Prod.* viii. 445.—Bras.
difformis, Benth. & Hook. f. Gen. ii. 710, 723 = Trachelospermum difforme.
ferruginea, Miers, Apocyn. S. Am. 227 = Echites ferruginea.
floribunda, *A. DC. in DC. Prod.* viii. 446.—Bras.
foliosa, *A. DC. l. c.*—Bras.
peruviana, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 71. t. 281.—Peruv.
Schlimiana, *Muell. Arg. in Linnaea*, xxx. (1859–60) 416.—N. Granat.

SECRETANIA, Muell. Arg. in DC. Prod. xv. II. 227 (1866) = **Hyeronyma**, Allem. (Euphorb.).
loranthacea, Muell. Arg. l. c. 228.—Guian.

SECURIDACA, Linn. Gen. ed. II. 347 (1742). *POLYGALEAE*, Benth. & Hook. f. i. 138.
CORYTHOLOBIUM, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 93.
ELSOTA, Adans. Fam. ii. 358 (1763).
LOPHOSTYLIS, Hochst. in Flora, xxv. (1842) 229.
RODSCHIEDIA, Miq. in Linnaea, xviii. (1844) 585.
acuminata, *A. St. Hil. Fl. Bras. Mer.* ii. 70.—Bras.
acuminata, Schlecht. in Linnaea, xiv. (1840) 382 = Schlechtendaliana.
affinis, *Bernh. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 360.—Guian.
angustifolia, Miq. in Tijdschr. Wetensch. i. (1848) 152 = longepedunculata.

SECURIDACA:—
atropurpurea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 359.—N. Granat.
bialata, *Benth. in Hook. Kew Journ.* iii. (1851) 162.—Bras.
bracteata, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 208.—Malacca.
brasiliensis, *Spreng. Syst.* iii. 175.—Bras.
Brownii, *Griseb. Fl. Brit. W. Ind.* 30.—Jamaica.
capparidifolia, Mart. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 66 = longifolia.
complicata, *H. B. & K. Nov. Gen. et Sp.* v. 423.—Venezuela.
coriacea, Bonpl. ex Steud. Nom. ed. II. ii. 551 = volubilis.
coriacea, Walp. Rep. i. 247 = Seguieria coriacea.
corymbosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 135.—N. Granat.
corymbosa, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 360 = Cumingii.
Corytholobium, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 68.—Bras.
Cumingii, *Hassk. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 190.—Ins. Philipp.
dasycarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 359.—N. Granat.
declinata, Vell. Fl. Flum. 294; vii. t. 73 = tomentosa.
decora, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 66.—Peruv.
divaricata, Nees & Mart. in Nov. Act. Nat. Cur. xii. I. (1824) 25 = rivinaefolia.
elliptica, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 358.—Cuba.
erecta, *Jacq. Enum. Pl. Carib.* 27.—Ind. occ.
erecta, Linn. Sp. Pl. ed. II. 992 = volubilis
erecta, Vell. Fl. Flum. 294; vii. t. 72 = lanceolata.
flexuosa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 560 = lanceolata.
floribunda, Oliver, Fl. Trop. Afr. i. 135 = Welwitschii.
floribunda, Walp. Rep. i. 247 = Seguieria floribunda.
Goudotiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 135.—N. Granat.
hebeclada, *DC. Prod.* i. 341.—Bras.
Hostmanni, *Miq. in Tijdschr. Wetensch.* i. (1848) 153; *Stirp. Surin. Sel.* 30.—Guian.
inappendiculata, Hassk. Pl. Jav. Rar. 295; Cat. Hort. Bog. Alt. 227 = tavoyana.
incrassata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1097, nomen.—Guian.
Lamarckii, *Griseb. Fl. Brit. W. Ind.* 30.—Ind. occ.
lanceolata, *A. St. Hil. Fl. Bras. Mer.* ii. 69.—Bras.
lateralis, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 68.—Bras.
latifolia, Benth. in Hook. Journ. Bot. iv. (1842) 104 = longifolia.
longepedunculata, *Fresen. in Mus. Senckenb.* ii. (1837) 275.—Abyss.
longifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 66.—Bras.
longifolia, Walp. Rep. i. 247 = Seguieria longifolia.
Luschnathiana, Presl, Bot. Bemerk. 17 = tomentosa.
macrocarpa, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 63.—Bras.
macrophylla, Walp. Rep. i. 247 = Seguieria macrophylla.
macrostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 560.—Ins. S. Vincent.
major, *Sagot, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 64.—Guian.
marginata, *Benth. in Hook. Journ. Bot.* iv. (1842) 103.—Guian.
Martiana, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. I. (1863) 559 = volubilis.
mollis, H. B. & K. Nov. Gen. et Sp. v. 421 = volubilis.
obcordata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 358.—Cuba.
oblongifolia, *Benth. & Hook. f. ex Harv. & Sond. Fl. Cap.* ii. 586.—Afr. austr.
ovalifolia, *A. St. Hil. Fl. Bras. Mer.* ii. 68.—Bras.
paniculata, Poir. Encyc. vii. 52 = volubilis.
paniculata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
paniculata, Roxb. ex Wall. Cat. sub n. 4195 = tavoyana.
paniculata, Roxb. Hort. Beng. 53; Fl. Ind. iii. 219 = scandens.

SECURIDACA:—
parvifolia, Spruce, ex A. W. Benn. in Mart Fl. Bras. xiii. III. 65 = rivinaefolia.
parvifolia, Walp. Rep. i. 247 = Seguieria parvifolia.
pendula, Bonpl. ex Steud. Nom. ed. II. ii. 551 = complicata.
polylopha, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 360.—Bras.
pubescens, *DC. Prod.* i. 341.—Guian.
pubescens, Seem. Bot. Voy. Herald, 80 = volubilis.
pubiflora, Benth. in Hook. Journ. Bot. iv. (1842) 104 = longifolia.
reticulata, Presl, Bot. Bemerk. 17 = rivinaefolia.
retusa, *Benth. in Hook. Kew Journ.* ii. (1850) 210.—Bras.
retusa, *Spruce, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 67.—Bras.
rivinaefolia, *A. St. Hil. Fl. Bras. Mer.* ii. 69.—Bras.
rufescens, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 134.—N. Granat.
rufescens, Sond. in Linnaea, xxii. (1849) 546 = tomentosa.
scandens, Buch.-Ham. ex Wall. Cat. n. 4195 = tavoyana.
scandens, *Jacq. Enum. Pl. Carib.* 27.—Ind. occ.
scandens, Lam. Encyc. vii. 51 = Lamarckii.
scandens, Tussac, Fl. Antill. iv. t. 20 = virgata.
scandens, Vell. Fl. Flum. vii. t. 60 = rivinaefolia.
Schlechtendaliana, *Walp. Rep.* i. 246.—Mexic.
Schlimii, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 135.—N. Granat.
Sellowiana, *Klotzsch, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 63.—Bras.
sulcata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1006, nomen.—Guian.
surinamensis, *Miq. in Tijdschr. Wetensch.* i. (1848) 153.—Guian.
sylvestris, *Schlecht. in Linnaea*, xiv. (1840) 381.—Mexic.
tavoyana, *Wall. Cat.* n. 4196.—Ind. or.; Malaya.
tomentosa, *A. St. Hil. Fl. Bras. Mer.* ii. 71. t. 96.—Bras.
virgata, Macfad. Fl. Jamaic. i. 49 = Brownii.
virgata, Sieber, ex Presl, Bot. Bemerk. 17 = volubilis.
virgata, *Sw. Prod. Veg. Ind. Occ.* 104.—Ind. occ.
volubilis, Blanco, Fl. Filip. ed. I. 556 = tavoyana.
volubilis, *Linn. Sp. Pl.* 707.—Am. austr.
Welwitschii, *Oliver, Fl. Trop. Afr.* i. 135.—Afr. trop.

SECURIDACA, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Securigera**, DC. (Legumin.).
legitima, Gaertn. Fruct. ii. 337. t. 153. f. 3 = Securigera Coronilla.
lutea, Mill. Gard. Dict. ed. VIII = Securigera Coronilla.
vera, Hort. ex Steud. Nom. ed. I. 758 = Securigera Coronilla.

SECURIDAEA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 559 (err. typ.) = **Securidaca**, Linn. (Polygal.).

SECURIGERA, DC. Fl. Fr. iv. 609 (1805). *LEGUMINOSAE*, Benth. & Hook. f. i. 489.
BONAVERIA, Scop. Introd. 310 (1777), nomen prius.
SECURIDACA, Tourn. ex Mill. Gard. Dict. ed. VI (1752).
SECURILLA, Gaertn. ex Steud. Nom. ed. I. 228 (1821).
SECURINA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 368.
atlantica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 41 = Coronilla atlantica.
Brownii, C. Muell. in Walp. Ann. vii. 702, sphalm. = Securidaca Brownii.
Coronilla, *DC. Fl. Fr.* iv. 609.—Europ. austr.
grandiflora, Boiss. Diagn. Ser. I. v. 88 = Coronilla grandiflora.
Lamarkii, C. Muell. in Walp. Ann. vii. 702, sphalm. = Securidaca Lamarckii.
legitima, Desv. in Mém. Soc. Linn. Par. iv. (1826) 301 = S. Coronilla.
Securidaca, supra, i. p. 620, lapsu = S. Coronilla.

SECURILLA, Gaertn. ex Steud. Nom. ed. I. 228 (1821) = **Securigera**, DC. (Legumin.).
Securidaca, Gaertn. ex Steud. l. c. = Securigera Coronilla.

SECURINA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 368 = **Securigera**, DC. (Legumin.).
lutea, Medic. l. c. 369 = Securigera Coronilla.

SECURINEGA, Comm. ex Juss. Gen. 388 (1789). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 275.
BESSERA, Spreng. Pugill. ii. 90 (1815).
COLMEIROA, Reut. in Mém. Soc. Phys. Genèv. x. (1843) 240. t. 6.
GEBLERA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 28 (1835).
MEINECKIA, Baill. Etud. Gén. Euph. 586 (1858).
PLEIOSTEMON, Sond. in Linnaea, xxiii. (1850) 135.
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 256 = Fluggea microcarpa.
Acidothamnus, *Muell. Arg. in DC. Prod.* xv. II. 451. —Ind. occ.
Bailloniana, *Muell. Arg. l. c.*—Afr. trop.
buxifolia, *Muell. Arg. l. c.* 452.—Europ. austr.
Commersonii, G. Don, in Sweet, Hort. Brit. ed. III. 768, nomen = durissima.
congesta, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 76.—Bras.
durissima, *J. F. Gmel. Syst.* 1008.—Ins. Borbon.
elliptica, *Muell. Arg. in DC. Prod.* xv. II. 448.—Am. centr.
flexuosa, *Muell. Arg. l. c.* 450.—Ins. Philipp.
fluggeoides, *Muell. Arg. l. c.*—Japon.
grisea, *Muell. Arg. l. c.* 451.—Reg. Himal.
Hilariana, *Muell. Arg. l. c.* 1273.—Bras.
hysterantha, Boj. Hort. Maurit. 278 = Phyllanthus indicus.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 128 = fluggeoides.
Leucopyrus, Brand. Forest Fl. 456 = Fluggea microcarpa.
Leucopyrus, Muell. Arg. in DC. Prod. xv. II. 451 = Fluggea Leucopyrus.
microcarpa, Muell. Arg. l. c. 434–6, passim = *obovata*.
Muelleriana, Baill. Adansonia, vi. (1865–66) 333 = Neoroepera buxifolia.
nitida, Lindl. Coll. Bot. t. 9 (= *Lithoxylon Lindleyi*).—Ins. Tahiti.
nitida, Willd. Sp. Pl. iv. 761 = durissima.
obovata, Muell. Arg. in DC. Prod. xv. II. 449 = Fluggea microcarpa.
phyllanthoides, *Muell. Arg. l. c.* 448.—Arab.
ramiflora, *Muell. Arg. l. c.* 449.—As. or.
Schuechiana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 78.—Bras.
Schweinfurthii, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 411.—Ins. Socotr.
trichogynis, *Muell. Arg. in DC. Prod.* xv. II. 447.—Madag.
virosa, Baill. Adansonia, vi. (1865–66) 334 = Fluggea Leucopyrus.

EDDERA, Hochst. in Flora, xxvii. (1844) Bes. Beil. 7. t. 5 = **Breweria**, R. Br. (Convolv.).
arabica, Choisy, in DC. Prod. ix. 441.—Arab.
Bottae, Jaub. & Spach, Illustr. iv. 108. t. 371.—Arab.
evolvuloides, Wight, Ic. t. 1369 = B. evolvuloides.
intermedia, Hochst. & Steud. ex Hochst. in Flora, xxvii. (1844) Bes. Beil. 8 = B. intermedia (in Addend.).
latifolia, Hochst. & Steud. l. c. = B. latifolia.
secundiflora, Jaub. & Spach, Illustr. iv. 110. t. 372.—Arab.
virgata, Hochst. & Steud. ex Hochst. in Flora, xxvii. (1844) Bes. Beil. 8 = B. virgata.

SEDGWICKIA, Griff. in As. Res. xix. (1836) 98. t. 15 = **Altingia**, Noronha (Hamamel.).
cerasifolia, Griff. l. c. 99. t. 15 = A. excelsa.

SEDUM, Adans. Fam. ii. 248 (1763 = **Sempervivum**, Rupp. (Crassul.).

SEDUM, Tourn. ex Linn. Syst. ed. I (1735). *CRASSULACEAE*, Benth. & Hook. f. i. 659.
AITHALES, Webb & Berth. Phyt. Canar. i. 178 (1836 ?).
ANACAMPSEROS, Haw. Syn. Pl. Succ. 111 (1812).
CEPAEA, Caesalp. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 384.
ENCHYLUS, Ehrh. Beitr. iv. 147 (1789).
LEUCOSEDUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 384.
RHODIA, Linn. Syst. ed. I (1735).
RHODIOLA, Linn. Gen. ed. I. 318 (1737).
PROCRASSULA, Griseb. Spicil. Fl. Rumel. i. 323 (1843).
TELMISSA, Fenzl, Pugill. Pl. Syr. 14 (1842).
TETRADIUM, Dulac, Fl. Hautes-Pyr. 321 (1867).
acre, *Linn. Sp. Pl.* 432.—Europ.; As. bor.
acutifolium, *Ledeb. Fl. Ross.* ii. 187.—As. Min.
adenotrichum, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 101, partim = Griffithii.
adenotrichum, *Wall. Cat.* n. 7231.—Reg. Himal.
aestivum, All. Fl. Pedem. ii. 121 = annuum.
aetnense, *Tineo, in Guss. Fl. Sic. Syn.* ii. 826.—Sicil.
affine, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 122 = S. Telephium.
aizoides, DC. Pl. Grass. t. 4 = Sempervivum aizoides.
aizoides, *Salm-Dyck, Hort. Dyck.* 371.—Hab. ?
Aizoon, *Linn. Sp. Pl.* 430.—Sibir.
Aizoon, Sibth. Herb. ex Boiss. Fl. Orient. iii. 961 = Trachelium Jacquini.
albellum, Willk. in Willk. et Lange, Prod. Fl. Hisp. iii. 141 = album.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 299.—Turkestan.
albescens, Haw. Rev. Pl. Succ. 28 = rupestre.
albicans, Steud. Nom. ed. I. 758 = S. Telephium.
alboroseum, *Baker, in Saund. Refug. Bot.* t. 33.—Japon.
album, Eichw. Beschr. Casp.-Cauc. 31 = acutifolium.
album, *Linn. Sp. Pl.* 432.—Europ.; As. bor.
Alfredi, *Hance, in Journ. Bot.* viii. (1870) 7.—China.
algidum, *Ledeb. Fl. Alt.* ii. 194.—Turkestan.; Sibir.
alpestre, *Vill. Prosp.* 49; *Hist. Pl. Dauph.* iii. 684.—Mont. Pyren.
alsinefolium, All. Fl. Pedem. ii. 119 = S. Cepaea.
altaicum, *G. Don, Gen. Syst.* iii. 114.—Sibir.
altissimum, Poir. Encyc. iv. 634 = nicaeënse.
altum, Clarke, Trav. ii. 420 = nicaeënse.
americanum, Banks, ex Pursh, Fl. Am. Sept. i. 324 = ternatum.
amplexicaule, *DC. Rapp. Voy.* ii. 80; *ex ej. Fl. Fr. Suppl.* 526.—Reg. Mediterr.
Anacampseros, *Linn. Sp. Pl.* 430.—Europ.
anatolicum, C. Koch, in Linnaea, xix. (1847) 41 = S. Listoniae.
andegavense, *DC. Prod.* iii. 406.—Europ. austr.
Andersonii, G. Don, in Loud. Hort. Brit. 184 = hispanicum.
andinum, *Ball, in Journ. Linn. Soc.* xxii. (1885) 38. —Peruv.
anglicum, *Huds. Fl. Angl.* ed. II. 196.—Europ.
angulatum, Hort. ex Mast. in Gard. Chron. (1878) II. 717 = album.
angustum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 138.—China.
annuum, All. Fl. Pedem. ii. 122 = alpestre.
annuum, Gunn. ex DC. Prod. iii. 405 = anglicum.
annuum, *Linn. Sp.* 432.—Europ.; As. Min.; Groenl.
anopetalum, *DC. Rapp. Voy.* ii. 80; *ex ej. Fl. Fr. Suppl.* 526.—Europ. austr.; As. Min.
arboreum, *Mast. in Gard. Chron.* (1878) II. 717.—Hab. ?
arboreum, Orteg. ex Steud. Nom. ed. II. ii. 552 = Crassula portulacea.
arduennense, Callay, ex Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 126 = S. Telephium.
arenarium, Brot. Fl. Lusit. ii. 212 = anglicum.
argutum, *Sweet, Hort. Brit.* ed. I. 179.—Mont. Carpath.
aristatum, Tenore, Fl. Nap. i. 250 = hispanicum.
aristatum, Vill. Hist. Pl. Dauph. iv. 680. t. 45 = anopetalum.
armenum, Boiss. & Huet, Diagn. Ser. II. ii. 61 = hispanicum.
arrigens, Gren. ex Nym. Consp. 261 = rupestre.
asiaticum, Aitch. in Journ. Linn. Soc. xix. (1882) 162 = heterodontum.

SEDUM:—
asiaticum, *Spreng. Syst.* iv. *Cur. Post.* 188; *DC. Prod.* iii. 401.—Reg. Himal.
assyriacum, *Boiss. Diagn.* Ser. I. vi. 55.—Syria; Assyria.
athoma, D. Dietr. Syn. Pl. ii. 1621, sphalm. = seq.
athoum, DC. Prod. iii. 407 = album.
atlanticum, Pers. Syn. i. 513 = hispidum.
atratum, Aubry, ex DC. Prod. iii. 405 = anglicum.
atratum, Bast. Fl. Maine-et-Loire, 167 = andegavense.
atratum, *Linn. Sp. Pl.* ed. II. 1673.—Europ. med.
atratum, Lucé, Fl. Osil. 143 = annuum.
atratum, Willem. ex Steud. Nom. ed. I. 758 = villosum.
atropurpureum, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 70.—Sibir.
aureum, Wirtg. ex F. Schultz, Arch. Fl. 200; et Fl. Preuss. Rheinprov. 184 = pruinatum.
azureum, Desf. Fl. Atlant. i. 362 = coeruleum, *Vahl.*
azureum, Royle, Illustr. Bot. Himal. 222. t. 48. f. 2 = Ewersii.
balticum, Hartm. f. Skand. Fl. ed. IX. 128 = album.
Baselicis, *Tenore, ex Steud. Nom.* ed. II. ii. 552, nomen.—Neapol.
Batesii, *Hemsl. Diagn. Pl. Nov.* 12.—Mexic.
Beyrichianum, *Mast. in Gard. Chron.* (1878) II. 376. —Hab. ?
bicolor, *H. B. & K. Nov. Gen. et Sp.* vi. 45.—Venezuela.
bithynicum, Boiss. Diagn. Ser. I. x. 17 = hispanicum.
Bohuslawii, *Rupr. in Beitr. Pfl. Russ. Reich.* vii. 18. —Ross.
boloniense, Loisel. in Desv. Journ. Bot. ii. (1809) 327 = sexangulare.
Boryanum, *DC. Prod.* iii. 410.—Hispan.
Botterii, *Hemsl. Diagn. Pl. Nov.* 10.—Mexic.
Bourgaei, *Hemsl. l. c.* 11.—Mexic.
bracteatum, *Viv. Fl. Lib. Spec.* 24. t. 8. f. 3.—Afr. bor.
brevifolium, *DC. Rapp. Voy.* ii. (1808) 79; *ex ej. Fl. Fr. Suppl.* 524.—Reg. Mediterr. occ.
Brunfelsii, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 121 = S. Telephium.
Bulliardi, Bor. l. c. 123 = S. Telephium.
bupleuroides, *Wall. Cat.* n. 7229.—Reg. Himal.
caeruleum, *Linn. Mant.* ii. 241 (Quid ?).—Afr. austr.
caesium, *Bor. ex Pérard, in Bull. Soc. Bot. Fr.* xviii. (1871) 325.—Gallia.
caespitosum, *DC. Prod.* iii. 405.—Reg. Mediterr. et Cauc.
calabricum, *Tenore, ex Steud. Nom.* ed. II. ii. 552, nomen.—Ital.
Callayanum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 127 = S. Telephium.
callichroum, *Boiss. Diagn.* Ser. I. vi. 56.—Persia.
Calverti, Boiss. Diagn. Ser. II. ii. 62 = acutifolium.
capense, *Burm. f. Fl. Cap. Prod.* 13.—Afr. austr.
capitatum, *Royle, Illustr. Bot. Himal.* 222, nomen.—Reg. Himal.
cariense, Jaub. & Spach, Illustr. i. 16. t. 7 = Listoniae.
carinatum, Link, ex Spreng. Syst. ii. 435 = amplexicaule.
Carioni, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 126 = S. Telephium.
carpaticum, Reuss, Květ. Slov. (1853) 162 = S. Telephium.
Cepaea, *Linn. Sp. Pl.* 431.—Europ.
cepaeum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
cespitosum, Boiss. Fl. Orient. ii. 794 = caespitosum.
chrisianum, Hort. Flor. ex Mast. in Gard. Chron. (1878) II. 717 (an *Clusianum* ?) = album.
chrysastrum, *Hance, in Journ. Bot.* viii. (1870) 6.—China.
ciliare, Sweet, Hort. Brit. ed. I. 179 = dentatum.
Clusianum, Guss. Fl. Sic. Syn. i. 516 = album.
coccineum, Royle, Illustr. Bot. Himal. 223. t. 48. f. 3 = quadrifidum.
coerulescens, Haw. in Phil. Mag. (1825) 174 = stenopetalum.
coeruleum, *Vahl, Symb. Bot.* ii. 51.—Reg. Mediterr.
coeruleum, Webb, ex Willk. & Lange, Prod. Fl. Hisp. iii. 140 = dasyphyllum.
cognense, Muell. Arg. in Compt.-Rend. Soc. Hallér. (1854–56) 130 = villosum.
collinum, Willd. Enum. Hort. Berol. Suppl. 24 = rupestre.
commutatum, Spreng. ex Steud. Nom. ed. II. ii. 552 = andegavense.

SEDUM:—
complanatum, Gilib. Fl. Lituan. ii. 181 = S. Telephium.
confertiflorum, *Boiss. Diagn.* Ser. I. iii. 15.—As. Min.
confertum, Bor. Fl. Centr. Fr. ed. III. ii. 253 = S. Telephium.
confertum, *Delile, Fl. Aeg. Illustr.* 62.—Aegypt.
confusum, *Hemsl. Diagn. Pl. Nov.* 10.—Mexic.
congestum, C. Koch, ex Boiss. Fl. Orient. ii. 779 = crenatum.
controversum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 125 = S. Telephium.
Cooperi, *Clémenc. in Rev. Hortic.* (1868) 474.—Hab.?
cordifolium, *Baker, in Saund. Refug. Bot.* t. 34.—Hab.?
coriaceum, *Wall. Cat.* n. 7238.—Reg. Himal.
corsicum, Duby, Bot. Gall. ed. II. i. 202 = dasyphyllum.
corymbiferum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 122 = S. Telephium.
Cotyledon, Jacq. f. Eclog. Pl. i. 27. t. 17 = Cotyledon caespitosa.
Cotyledon, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = creticum.
crassicaule, Hort. ex Link, Enum. Hort. Berol. i. 438 = rupestre.
crassipes, Wall. Cat. n. 7234 = asiaticum.
crenatum, *Boiss. Fl. Orient.* ii. 779.—Armenia.
crenulatum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 96.—Reg. Himal.
creticum, *Boiss. & Heldr. Diagn.* Ser. I. x. 16.—Ins. Creta.
creticum, *Presl, in Oken, Isis*, xxi. (1828) 273.—Ins. Creta.
cristatum, Schrad. Hort. Gotting. 16 = rupestre.
cruciatum, Desf. Tabl. ed. I. 162 = monregalense.
cupressoides, *Hemsl. Diagn. Pl. Nov.* 11.—Mexic.
cyaneum, *Rudolph, in Mém. Acad. Pétersb.* iv. (1811) 341.—Sibir.
dasyphyllum, *Linn. Sp. Pl.* 431.—Europ.; Afr. bor.
debile, *S. Wats. in Bot. King's Exp.* 102.—Am. bor. occ.
decumbens, Lucé, Topog. Nachr. Oesel, 142 = S. Telephium.
deficiens, Donn, Hort. Cantab. ed. VI. 126 = ternatum.
deltoideum, Tenore, Prod. Fl. Nap. p. xxvi = stellatum.
dendroideum, *Moç. & Sessé, ex DC. Prod.* iii. 409.—Mexic.
dentatum, *DC. Prod.* iii. 403.—Reg. Cauc.
dentatum, Steph. ex Cham. in Linnaea, vi. (1831) 549 = Stephani.
denticulatum, Donn, Hort. Cantab. ed. VIII. ex DC. Prod. iii. 403 = oppositifolium.
dioicum, Donn, l. c. ed. III. 84 = nicaeënse.
dioicum, Stokes, Bot. Mat. Med. ii. 549 = roseum, *Scop.*
divaricatum, Ait. Hort. Kew. ed. I. ii. 108 = Sempervivum aizoides.
divaricatum, Lapeyr. Hist. Abr. Pl. Pyr. 260 = annuum.
divaricatum, Schlecht. ex Ledeb. Fl. Ross. ii. 185 = sempervivoides.
divaricatum, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 372.—Am. bor. occ.
divergens, *S. Wats. l. c.*—Am. bor. occ.
dodecandrum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 138 = pruinatum.
dolomiticum, Hort. ex Mast. in Gard. Chron. (1878) II. 626 = nicaeënse.
Douglasii, *Hook. Fl. Bor. Am.* i. 228.—Am. bor. occ.
Douglasii, S. Wats. in Bot. Calif. i. 210 = radiatum.
drymarioides, *Hance, in Seem. Journ. Bot.* iii. (1865) 379.—China.
dumulosum, *Franch. Pl. Davidian.* i. (1884) 129.—China.
ebracteatum, [*Moç. & Sessé,*] *ex DC. Prod.* iii. 409.—Mexic.
edule, Nutt. ex Torr. & Gray, Fl. N. Am. i. 560 = Cotyledon edulis.
elatinoides, *Franch. Pl. Davidian.* i. (1884) 131.—China.
elegans, Lej. Fl. Spa, i. 205 = pruinatum.
ellipticum, *Miégev. Bull. Soc. Bot. Fr.* xiv. (1867) 145.—Mont. Pyren.
elongatum, Ledeb. Fl. Alt. ii. 193 = roseum, *Scop.*
elongatum, *Wall. Cat.* n. 7233; *Hook. f. in Journ. Linn. Soc.* ii. (1858) 98.—Reg. Himal.

SEDUM:—
engadinense, *Bruegg. in Jahresb. Nat. Ges. Graub.* II. xxv. (1882) 96.—Europ.
epidendrum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 314.—Abyss.
eriocarpum, Sibth. & Sm. Fl. Graec. Prod. i. 310; et Fl. Graec. v. 36. t. 449 = hispanicum.
erraticum ×, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxv. (1882) 58, 96.—Europ.
erythraeum, *Griseb. Spicil. Fl. Rumel.* i. 326.—Macedon.
erythromelanum, Fr. Braun, ex Nym. Consp. 260 = maximum.
erythrostictum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 155 = S. Telephium.
euphorbioides, *Schlecht. ex Ledeb. Fl. Ross.* ii. 177.—Sibir.
Ewersii, *Ledeb. Fl. Alt.* ii. 191.—Reg. Himal.; Sibir. altaic.
Fabaria, Koch, Syn. Fl. Germ. ed. I. 258 = S. Telephium.
Fabaria, Vilm. ex Vilm. Fl. Pl. Terre, ed. I. 811 = spectabile.
fabarium, Hort. Angl. ex Lem. Illustr. Hortic. viii. (1861) t. 271 = spectabile.
farinosum, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 31.—Ins. Mader.
fastigiatum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 98.—Reg. Himal.
filiforme, Poir. Encyc. Suppl. iv. 210 = hispidum.
fimbriatum, Franch. Pl. Davidian. i. (1884) 128 = Cotyledon fimbriata.
formosanum, *N. E. Br. in Gard. Chron.* (1885) II. 134.—Ins. Formosa.
Forsteri, Haw. Syn. Pl. Succ. 117 = pruinatum.
Forsterianum, Reichb. Pl. Crit. ix. 15 = sexangulare.
Forsterianum, Sm. Engl. Bot. t. 1802 = pruinatum.
fragile, *Dum. Fl. Belg. Prod.* 85.—Belgium.
fruticulosum, Brot. Fl. Lusit. ii. 206 = nicaeënse.
Funkii, Reichb. Fl. Germ. Excurs. Index 117, sphalm. = Sempervivum Funckii.
fuscum, *Hemsl. Biol. Centr. Am. Bot.* i. 395.—Mexic.
fusiforme, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 31. t. 3. f. 1.—Ins. Mader.
galioides, All. Fl. Pedem. ii. 120. t. 65. f. 3 = S. Cepaea.
gelidum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 355.—Sibir. altaic.
Gerardianum, Wall. Cat. n. 7235; Royle, Illustr. Bot. Himal. 222 = Ewersii.
glaciale, Clarion, ex DC. Fl. Fr. iv. 393 = acre.
glanduliferum, Guss. Fl. Sic. Prod. i. 519 = dasyphyllum.
glandulosum, *Moris, Stirp. Sard. Elench.* 20.—Ins. Sardoa.
glandulosum, Walp. Rep. v. 795 = Telephium glandulosum.
glaucum, Lam. Fl. Fr. iii. 84 = dasyphyllum.
glaucum, Mayer, ex Tenore, Syll. Fl. Neap. 226 = pallidum.
glaucum, Sm. Engl. Fl. ii. 321 = rupestre.
glaucum, Waldst. & Kit. Pl. Rar. Hung. ii. 198. t. 181 = hispanicum.
globiferum, Pall. Reise, i. 256 = Sempervivum globiferum.
globiferum, Pourr. in Mém. Acad. Toul. iii. (1788) 327 = brevifolium.
globulifolium, *Moris, ex G. Don, in Sweet, Hort. Brit.* ed. III. 269, nomen.—Europ.
gracile, DC. ex Steud. Nom. ed. I. 759 = dasyphyllum.
gracile, *Haensel. ex Willk. in Bot. Zeit.* iv. (1846) 312.—Hispan.
gracile, *C. A. Mey. Verz. Pfl. Cauc.* 150.—Reg. Cauc.
grandidentatum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 124 = S. Telephium.
graniticum, Pérard, in Bull. Soc. Bot. Fr. xvi. (1869) 186 = rupestre.
granulatum, Salisb. Prod. 307 = dasyphyllum.
Greggii, *Hemsl. Diagn. Pl. Nov.* 12.—Mexic.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 421.—Reg. Himal.
Grisebachii, *Boiss. & Heldr. Diagn.* Ser. II. ii. 61.—Graecia.
guatemalense, *Hemsl. Diagn. Pl. Nov.* 11.—Guatemala.
guestphalicum, Boenn. ex Reichb. Fl. Germ. Excurs. 550 = S. Telephium.

SEDUM:—
Guetardi, Vill. Hist. Pl. Dauph. iii. 678. t. 45 = anglicum, alpestre.
gypsicolum, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 13.—Hispan.
haematodes, Mill. Gard. Dict. ed. VIII. n. 15 = telephioides?
haematodes, Scop. Fl. Carn. ed. II. i. 323 = atratum.
haematodeum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
heptapetalum, *Fisch. ex Hornem. Suppl. Hort. Hafn.* 138.—Rossia.
heptapetalum, Poir. Voy. Barb. ii. 169 = coeruleum, *Vahl.*
heterodontum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 95.—Reg. Himal.
hexangulare, J. B. Fisch. Naturg. von Livl. 498 = sexangulare.
hexangulare, Frivald. ex Nym. Consp. 264 = annuum.
hexapetalum, *Haw. Rev. Pl. Succ.* 26.—Hab.?
Hillebrandii, Fenzl, in Verh. Zool.-Bot. Ges. Wien, vi. (1856) 449 = sexangulare.
himalayanum, Wall. Cat. n. 7236 = seq.
himalense, *D. Don, Prod. Fl. Nep.* 212.—Reg. Himal.
hirsutum, *All. Fl. Pedem.* ii. 122. t. 65. f. 5.—Europ. austr.
hirsutum, Brot. Fl. Lusit. ii. 212 = anglicum.
hispanicum, DC. Fl. Fr. iv. 395 = anopetalum.
hispanicum, *Linn. Cent. Pl.* i. 12; *Amoen. Acad.* iv. 273.—Europ. austr.; As. Min.
hispidum, *Desf. Fl. Atlant.* i. 361.—Afr. bor. occ.
hispidum, Poir. Encyc. iv. 633 = hirsutum.
Hudsonianum, Lange.—Cf. Linnaea, xxix. (1857–58) 736 = anglicum.
humile, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 99.—Reg. Himal.
hungaricum, Poir. Encyc. Suppl. iv. 208 = hispanicum.
hybridum, *Linn. Sp. Pl.* 431.—Sibir.
hybridum, Trautv. & C. A. Mey. in Middend. Reise, (Fl. Ochot. 40) = Middendorffianum.
hybridum, Urv. in Mém. Soc. Linn. Par. i. (1822) 307 = stoloniferum.
hypericifolium, Wall. Cat. n. 7237 = himalense.
ibericum, Stev. ex Bieb. Fl. Taur. Cauc. iii. 312 = stoloniferum.
imbricatum, Walp. Ann. i. 325; Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 101 = roseum, *Scop.*
incertum, *Hemsl. Diagn. Pl. Nov.* 11.—Mexic.
insulare, *Moris, Enum. Sem. Hort. Taurin.* (1861) 33.—Ins. Sardoa.
intermedium, Déségl. ex Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 123 = S. Telephium.
intermedium, *Rupr. Fl. Ingr.* 404.—Reg. Cauc.
involucratum, *Bieb. Fl. Taur. Cauc.* i. 352.—Reg. Cauc.
Jacquini, Haw. in Phil. Mag. (1825) 174 = nicaeënse.
Jaeschkei, *Kurz, in Journ. Bot.* v. (1867) 240.—Reg. Himal.
japonicum, *Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 156.—China; Japon.
Jullianum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 117 = S. Telephium.
kagamontanum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 137.—Japon.
kamtschaticum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vii. 54.—As. or.
kamtschaticum, Hort. ex Mast. in Gard. Chron. (1878) II. 268 = Selskianum.
Kirilowii, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 92.—Soongar.; Turkestan.
kokanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 26.—Kokan.
Kotschyanum, *Boiss. Diagn.* Ser. I. vi. 56.—Persia.
laconicum, *Boiss. & Heldr. Diagn.* Ser. I. vi. 55.—Graecia; As. Min.
laconicum, Heldr. & Sart. ex Nym. Consp. 262 = acre.
lactifolium, Salisb. Prod. 307 = S. Telephium.
Lampusae, *Boiss. Fl. Orient.* ii. 787.—Ins. Cyprus.
lanceolatum, Torr. in Ann. Lyc. N. York, ii. (1828) 205 = stenopetalum.
latifolium, Bertol. Amoen. Ital. 366 = maximum.
laxiflorum, *DC. Prod.* iii. 409.—Ins. Canar.
lazicum, Boiss. Diagn. Ser. II. ii. 63 = crenatum.
Ledebourii, Hort. ex Steud. Nom. ed. II. ii. 552 = roseum, *Scop.*
libanoticum, Linn. Syst. ed. X. 1037 = Cotyledon serrata.
Liebmannianum, *Hemsl. Diagn. Pl. Nov.* 12.—Mexic.
lilacinum, Ledeb. in Mém. Acad. Pétersb. v. (1812) 535 = cyaneum.

SEDUM:—
lineare, *Thunb. Fl. Jap.* 187.—Japon.
linearifolium, *Royle, Illustr. Bot. Himal.* 222. t. 48. f. 1.—Reg. Himal.
linifolium, Nutt. ex Torr. & Gray, Fl. N. Am. i. 559 = pulchellum.
Listoniae, *Vis. in Mem. Ist. Venet.* i. 1842 (1843), 57. t. 6.—As. Min.
litoreum, *Guss. Pl. Rar.* 185. t. 37. f. 2.—Ital.; Graec.; As. Min.
lividum, Drejer, in Fl. Dan. t. 686 = S. Telephium.
lividum, Hort.—Cf. Gard. Chron. (1877) II. 521 = lydium.
lividum, *Willd. Enum. Hort. Berol. Suppl.* 24.—Hab.?
Lobelii, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 125 = S. Telephium.
lugdunense, Nym. Consp. 260 = S. Telephium.
lusitanicum, Brot. Phyt. Lusit. ii. t. 178 = nicaeënse.
lusitanum, Brot. l. c. 220, sphalm. = praec.
luteolum, *Chabois. in Bull. Soc. Bot. Fr.* x. (1863) 296.—Europ.
lydium, *Boiss. Diagn.* Ser. I. iii. 17.—As. Min.
macrophyllum, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 811 maximum.
magellense, *Tenore, Prod. Fl. Nap.* p. xxvi.—Ital.; Graec.; As. Min.
majus, Gueldenst. ex Ledeb. Fl. Ross. ii. 180 = maximum.
malacophyllum, Steud. Nom. ed. I. 759; Franch. Pl. Davidian. i. (1884) 129 = Cotyledon malacophylla.
Marichalii, Lloyd, ex Nym. Consp. 264 = litoreum.
maritimum, *Bogusl. in Erman, Archiv*, vi. (1848) 66.—Ross.
maritimum, Sass. Phaner. Fl. Oesels, 55, sphalm. = maximum.
matrense, *Kit. in Linnaea*, xxxii. (1863) 470.—Hungar.
Maximowiczii, Regel, Gartenfl. (1866) 355. t. 528; et Ind. Sem. Hort. Petrop. (1866) 98 = S. Aizoon.
maximum, *Suter, Fl. Helvet.* i. 270.—Europ.; Reg. Cauc.
Meehani, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 105.—Am. bor.
melanantherum, *DC. Prod.* iii. 410.—Hispan.
melanoleucum, Schlecht. ex Ledeb. Fl. Ross. ii. 172 = caespitosum.
micranthum, Bast. ex DC. Fl. Fr. Suppl. 523 = album.
microstachyum, *Boiss. Fl. Orient.* ii. 787.—Ins. Cyprus.
Middendorffianum, *Maxim. Prim. Fl. Amur.* 116.—Reg. Amur.
Millii, *Baker, in Journ. Bot.* xiii. (1875) 236.—Anatolia.
minus, Haw. in Phil. Mag. (1825) 175 = rupestre.
miserum, *Lindl. in Bot. Reg.* (1838) *Misc.* 65.—Mexic.
mite, Gilib. Fl. Lituan. ii. 182 = sexangulare.
mixtum, *Linn. Syst.* ed. X. 1037.—Hab.?
modestum, *Ball, in Journ. Bot.* xi. (1873) 333.—Marocc.
modestum, Boiss. Fl. Orient. ii. 789 = hispanicum.
monregalense, *Balb. Misc. Bot.* 23. t. 6.—Europ. austr.
montanum, Song. & Perrier, in Billotia, i. (1864) 77 = anopetalum.
Moorcroftianum, Wall. Cat. n. 7228 = Sempervivum acuminatum.
moranense, *H. B. & K. Nov. Gen. et Sp.* vi. 44.—Mexic.
mucronatum, Edgew. in Trans. Linn. Soc. xx. (1846) 48 = linearifolium.
multicaule, *Wall. Cat.* n. 7232; *et ex Lindl. Bot. Reg.* (1840) *Misc.* 58.—Reg. Himal.; China.
multiceps, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 171.—Alger.
multiflorum, Boenn. ex Salm-Dyck, Hort. Dyck. 250, nomen (Quid?).—Hab.?
mutabile, Wender. ex Steud. Nom. ed. II. ii. 553, nomen (Quid?).—Hab.?
nanum, *Boiss. Diagn.* Ser. I. vi. 57.—Persia.
napiferum, *Peyr. in Linnaea*, xxx. (1859-60) 50.—Mexic.
neapolitanum, Tenore, Syll. Fl. Neap. 536 = dasyphyllum.
nebrodense, Gaspar. in Rendic. Acc. Nap. i. (1842) 49 = dasyphyllum.
neglectum, Tenore, Ind. Sem. Hort. Neap. (1830) 12; et Syll. Fl. Neap. 227 = acre.
nevadense, Coss. Notes Crit. 163 = villosum.

SEDUM:—
Nevii, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858) 373.—Am. bor.
nicaeënse, *All. Fl. Pedem.* ii. 122.—Reg. Mediterr.
Notarjanni, *Tenore, Prod. Fl. Nap.* p. xxvi.—Ital.
nudum, *Ait. Hort. Kew.* ed. I. ii. 112.—Ins. Mader.
nudum, Schousb. ex Coss. in Bull. Soc. Bot. Fr. xx. (1873) 248 = brevifolium.
Nuttalianum, Rafin. Atl. Journ. 146 (Quid?).—Am. bor.
oblongum, Haw. Rev. Pl. Succ. 29 = anglicum.
obtusatum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 342.—Calif.
obtusifolium, *C. A. Mey. Verz. Pfl. Cauc.* 150.—Reg. Cauc.
occidentale, Bor. ex Mast. in Gard. Chron. (1878) II. 303 = S. Telephium.
ochroleucum, Chaix, in Vill. Hist. Pl. Dauph. i. 325 = anopetalum.
ochroleucum, Vill. l. c. iv. 680 = nicaeënse.
octogonum, Hort. ex DC. Prod. iii. 403 = ternatum.
Oederi, Retz. Fl. Scand. 85 = saxatile.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 26.—Turkestan.
olympicum, Boiss. Diagn. Ser. I. iii. 16 = magellense.
oppositifolium, *Sims, Bot. Mag.* t. 1807.—Reg. Cauc.; Persia.
oreganum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 559.—Am. bor. occ.
orientale, Boiss. Diagn. Ser. I. x. 17 = hispanicum.
oxypetalum, *H. B. & K. Nov. Gen. et Sp.* vi. 45.—Mexic.
pachyclados, *Aitch. ex Hemsl. in Journ. Linn. Soc.* xviii. (1880) 58.—Afghan.
palaestinum, *Boiss. Diagn.* Ser. I. x. 18.—Syria.
pallidum, *Bieb. Fl. Taur. Cauc.* i. 353.—As. Min.; Persia; Ind. bor. occ.
pallidum, Tenore, Syll. Fl. Neap. 226 = hispanicum.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 355.—Mexic.
paludum, Link, Handb. ii. 27, sphalm. = pallidum.
palustre, Oed. Fl. Dan. i. 6. t. 24 = villosum.
paniculatum, Lam. Fl. Fr. iii. 83 = S. Cepaea.
paniculatum, Wall. Cat. n. 7227 = Crassula indica.
parnassicum, Boiss. & Heldr. Diagn. Ser. II. ii. 61 = annuum.
parvum, *Hemsl. Diagn. Pl. Nov.* 50.—Mexic.
paucidens, *Sweet, Hort. Brit.* ed. I. 179.—Europ.
pauciflorum, Edgew. in Trans. Linn. Soc. xx. (1846) 49 = linearifolium.
pectinatum, *Sweet, Hort. Brit.* ed. I. 179; *DC. Prod.* iii. 403.—Hab.?
pedicellatum, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 13.—Hispan.
pentandrum, Bor. Fl. Centr. Fr. ed. II. n. 779; et ed. III. ii. 256 = villosum.
Penthorum, Crantz, Inst. ii. 443 = Penthorum sedoides.
perpusillum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 103.—Reg. Himal.
peruvianum, *A. Gray, Bot. U. St. Expl. Exped.* i. 691.—Peruv.
pilosum, *Bieb. Fl. Taur. Cauc.* i. 352.—Reg. Cauc.; As. Min.
populifolium, *Pall. Reise*, iii. 730.—Sibir.
porphyreum, *Kotschy, in Unger & Kotschy, Ins. Cypern*, 313.—Ins. Cyprus.
portulacoides, Willd. Enum. Hort. Berol. 484 = ternatum.
praealtum, *A. DC.* 10me *Not. Pl. Rar. Jard. Genèv.* 21.—Mexic.
procumbens, *Schrank, Baier. Fl.* i. 726.—Bavaria.
pruinatum, *Brot. Fl. Lusit.* ii. 209.—Reg. Mediterr. occ.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 156.—China.
Pseudo-aizoon, Debeaux, in Act. Linn. Soc. Bord. xxxi. (1877) 159 (Fl. du Tché-fou, 64) = S. Aizoon.
Pseudo-fabaria, *Lange, Ind. Sem. Hort. Haun.* (1868) 23.—Hab.?
puberulum, DC. Mém. Crass. 33 = hispanicum.
pubescens, *Vahl, Symb. Bot.* ii. 52.—Reg. Tunet.
pulchellum, *Michx. Fl. Bor. Am.* i. 277.—Am. bor.
pulchrum, DC. Prod. iii. 403 = pulchellum.
pulligerum, Pomel, Nouv. Mat. Fl. Atl. 326 = dasyphyllum.
pumilum, *Benth. Pl. Hartw.* 310.—Calif.
punctatum, *Kit. in Linnaea*, xxxii. (1863) 469.—Hungar.

SEDUM:—
purpurascens, Koch, Syn. Fl. Germ. ed. II. 284 = S. Telephium.
purpureum, Link, Enum. Hort. Berol. i. 437 = S. Telephium.
pusillum, *Michx. Fl. Bor. Am.* i. 276.—Am. bor.
pusillum, Pursh, Fl. Am. Sept. i. 323 = pulchellum.
pyrenaicum, Lange.—Cf. Linnaea, xxix. (1857-58) 735 = anglicum.
pyriforme, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 101 = rosulatum.
quadrifidum, *Pall. Reise*, iii. 730.—Reg. Himal.; As. arct.
quinquefidum, *Haw. Rev. Pl. Succ.* 26.—Hab.?
quitense, *H. B. & K. Nov. Gen. et Sp.* vi. 46.—Ecuador.
racemiferum, Griseb. ex Nym. Consp. 264 = Grisebachii.
radiatum, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 193.—Calif.
radicosum, Boiss. & Hohen. Diagn. Ser. I. x. 15 (= *Umbilicus platyphyllus*).—Persia; Soongar.
Raji, Lange.—Cf. Linnaea, xxix. (1857-58) 735 = anglicum.
ramosissimum, Franch. Pl. Davidian. i. (1884) 128 = Cotyledon fimbriata.
recurvatum, Willd. Enum. Hort. Berol. Suppl. 24 = rupestre.
reflexum, Cutanda, Fl. Madrit. 312 = pruinatum.
reflexum, Frivald. ex Nym. Consp. 261 = amplexicaule.
reflexum, Linn. Fl. Suec. App. 463 = rupestre.
repens, Nym. Consp. 260 = S. Telephium.
repens, Schleich. ex DC. Fl. Fr. Suppl. 525 = alpestre.
reticulatum, Schrank, in Denkschr. Bot. Ges. Regensb. i. (1815) 107 = dasyphyllum.
retorridum, Hort. ex Mast. in Gard. Chron. (1878) II. 626 = amplexicaule.
retusum, *Hemsl. Diagn. Pl. Nov.* 51.—Mexic.
rhodanense, Nym. Consp. 260 = S. Telephium.
rhodanthum, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 405.—Am. bor. occ.
Rhodiola, DC. Fl. Fr. iv. 386 = roseum, *Scop.*
Rhodiola, Ledeb. Fl. Alt. ii. 194 = altaicum.
Rhodiola, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 244 = elongatum.
rhodioloides, Rafin. Atl. Journ. 146 = roseum, *Scop.*
rigidum, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xx. (1866) 117.—Europ.
rivulare, Boiss. Elench. 43 = melanantherum.
rivulare, Griseb. Spicil. Fl. Rumel. i. 327 = erythraeum.
Roborowskii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 154.—China.
roseum, *Scop. Fl. Carn.* ed. II. i. 326.—Europ.
roseum, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 263.—Reg. Cauc.
rostratum, Tenore, Prod. Fl. Nap. p. xxvi = amplexicaule.
rosulatum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 48.—Reg. Himal.
rotundifolium, Lam. Fl. Fr. iii. 82 = S. Anacampseros.
rubellum, Wender. Ind. Sem. Hort. Marb. (1824); ex Steud. Nom. ed. II. ii. 553 (Quid?).—Hab.?
rubens, DC. Prod. iii. 405 = pallidum.
rubens, Haenke, in Jirasek, Beob. Riesengeb. (1791) 114 = alpestre.
rubens, Jacq. ex Nym. Consp. 264 = atratum.
rubens, *Linn. Sp. Pl.* 432.—Reg. Mediterr.
rubens, Oed. Fl. Dan. t. 82 = anglicum.
rubens, Pollini, Fl. Veron. ii. 71 = hispanicum.
rubrum, Royle, Illustr. Bot. Himal. 222 = Ewersii.
rufescens, Tenore, Prod. Fl. Nap. p. xxvii = nicaeënse.
rupestre, All. ex Reut. in Bull. Soc. Bot. Fr. ii. (1855) 643 = anglicum.
rupestre, Hacq. ex Nym. Consp. 262 = alpestre.
rupestre, *Linn. Sp. Pl.* 431.—Europ.
rupestre, Oed. Fl. Dan. t. 59 = saxatile.
rupestre, Plan. Ens. Fl. Gallega, 221 = pruinatum.
rupestre, Vill. Hist. Pl. Dauph. iii. 679 = anopetalum.
sajanense, Pall. ex Ledeb. Fl. Ross. ii. 183 = S. Aizoon.
Salsovitzianum, Hort. ex Mast. in Gard. Chron. (1878) II. 626 = nicaeënse.
sanguineum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 790.—Mesopot.
sanguineum, *Ortega, Hort. Matr. Dec.* 125.—Hispan.
sarmentosum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 104.—China.
Sartorianum, Boiss. Diagn. Ser. II. ii. 62 = acre.

SEDUM:—
saxatile, All. Fl. Pedem. ii. 121. t. 65. f. 6 = alpestre.
saxatile, DC. Fl. Fr. iv. 394 = annuum.
saxatile, Pall. Tabl. Taur. 51 = pallidum.
saxatile, [*Weber, in*] *Wigg. Prim. Fl. Holsat.* 35.—Europ.
Scheuchzeri, Schleich. ex Mert. & Koch, Deutschl. Fl. iii. 314 = alpestre.
Schimperi, *Britten, in Oliver, Fl. Trop. Afr.* ii. 400.—Abyss.
schistosum, Lej. Fl. Spa, i. 206 = sexangulare.
Selskianum, *Regel & Maack, in Tent. Fl. Ussur.* 66.—Reg. Amur.
Semenovii, *Mast. in Gard. Chron.* (1878) II. 267.—Turkestan.
semiglobosum, *Boiss. & Huet, in Tchihat. Asie Min. Bot.* i. 397, nomen.—Armenia.
sempervivoides, *Fisch. ex Bieb. Fl. Taur. Cauc.* iii. 313.—As. Min.; Reg. Cauc.
Sempervivum, Ledeb. ex Spreng. Syst. ii. 434 = sempervivoides.
septangulare, Haw. Syn. Pl. Succ. 116 = rupestre.
serotinum, *Bellynck, Fl. Namur* (1855), 322.—Belgium.
serratum, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 417 = heterodontum.
sexangulare, *Linn. Sp. Pl.* 432.—Europ.
sexangulare, Loisel. ex Boiss. Fl. Orient. ii. 784 = acre.
sexfidum, Bieb. Fl. Taur. Cauc. i. 354 = hispanicum.
Sheareii, *S. Moore, in Journ. Bot.* xiii. (1875) 227.—China.
Sieboldii, *Hort. ex G. Don, in Sweet, Hort. Brit.* ed. III. 270, nomen; *et ex Hook. Bot. Mag.* t. 5358.—Japon.
simplex, Steud. ex Boiss. Fl. Orient. ii. 790 = Steudelii.
sinuatum, Royle, Illustr. Bot. Himal. 222 = trifidum.
soluntinum, Tineo, in Guss. Fl. Sic. Syn. ii. 827 = nicaeënse.
sordidum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 142.—Japon.
sparsiflorum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 559 = Torreyi.
sparsiflorum, Torr. in Bot. U. St. Expl. Exped. ii. 307 = oreganum ?
spathulatum, D. Dietr. Syn. Pl. ii. 1618, sphalm. = spathulifolium.
spathulatum, Waldst. & Kit. Pl. Rar. Hung. ii. 108 = S. Cepaea.
spathulifolium, *Hook. Fl. Bor. Am.* i. 227.—Am. bor. occ.
spectabile, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xx. (1866) 116.—Japon.
sphaericum, Lapeyr. Hist. Abr. Pl. Pyr. 259 = brevifolium.
spicatum, [Moç. & Sessé,] ex DC. Prod. iii. 401 = Cotyledon coccinea.
spinosum, Thunb. Fl. Jap. 186; Willd. Enum. Hort. Berol. 485 = Cotyledon spinosa.
spirale, Haw. in Phil. Mag. (1825) 176 = sexangulare.
spurium, Bieb. Fl. Taur. Cauc. i. 352 = stoloniferum.
spurium, Sibth. ex Nym. Consp. 485 = Trachelium rumelianum.
stellariaefolium, Franch. Pl. Davidian. i. (1884) 130 = drymarioides.
stellatum, Frivald. ex Nym. Consp. 265 = caespitosum.
stellatum, Habl. Desc. Phys. Taur. 161 = pallidum.
stellatum, *Linn. Sp. Pl.* 431.—Europ. austr.
stellatum, Lour. Fl. Cochinch. 287 = Alfredi.
stellatum, Pollini, Fl. Veron. ii. 73 = Sempervivum Seguieri.
stellatum, Urv. in Mém. Soc. Linn. Par. i. (1822) 307 = litoreum.
stelliforme, *S. Wats. in Proc. Am. Acad.* xx. (1885) 365.—Arizona.
stenopetalum, *Pursh, Fl. Am. Sept.* i. 324.—Am. bor. occ.
Stephani, *Cham. in Linnaea*, vi. (1831) 549.—Sibir.
Steudelii, *Boiss. Fl. Orient.* ii. 790.—Cilicia.
stoloniferum, *S. T. Gmel. Reise*, iii. 311. t. 35. f. 2.—As. Min.; Persia.
Stracheyi, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 96 = tibeticum.
strictum, C. Koch, in Linnaea, xix. (1847) 42 = S. Cepaea.
strictum, Schlecht. ex Ledeb. Fl. Ross. ii. 177 = algidum.

SEDUM:—
subclavatum, Haw. in Phil. Mag. (1831) 414 = stenopetalum ?
subopposítum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 132.—China.
subtile, Debeaux, Fl. Shangh. 32 = Alfredi.
subtile, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 156.—Japon.
subulatum, *Boiss. Fl. Orient.* ii. 783.—Reg. Cauc.
surculosum, Coss. in Bull. Soc. Bot. Fr. xx. (1873) 248 = Monanthes atlantica.
Taeschkei, Kurz, in Journ. Bot. v. (1867) 240, sphalm. = Jaeschkei.
Tatarinowii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 134.—China.
tectorum, Schur, Enum. Pl. Transs. 965 (Index) = annuum.
tectorum, Scop. Fl. Carn. ed. II. i. 325 = Sempervivum tectorum.
telephioides, *Michx. Fl. Bor. Am.* i. 277.—Am. bor.
Telephium, *Linn. Sp. Pl.* 430.—Europ.; As. bor.
Telephium, Willd. ex Ledeb. Fl. Ross. ii. 180, 181 = maximum.
tenellum, *Bieb. Fl. Taur. Cauc.* iii. 315.—Reg. Cauc.; As. Min.
tenuifolium, DC. Prod. iii. 407, in syn. = amplexicaule.
teretifolium, Lam. Fl. Fr. iii. 84 = album.
teretifolium, [Moç. & Sessé,] ex DC. Prod. iii. 401 = Cotyledon teretifolia.
ternatum, *Michx. Fl. Bor. Am.* i. 277.—Am. bor.
tetramerum, *Trautv. in Act. Hort. Petrop.* vii. (1880) 454.—Reg. Cauc.
tetraphyllum, Sibth. & Sm. Fl. Graec. Prod. i. 309; Fl. Graec. v. 36. t. 448 = S. Cepaea.
Thirkeanum, C. Koch, in Linnaea, xix. (1847) 41 = S. Telephium.
thyrsoideum, Bor. in Mém. Soc. Acad. Maine-et-Loire, xx. (1866) 120 = S. Telephium.
tibeticum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 96.—Reg. Himal.
Tilingiana, C. Muell. in Walp. Ann. vii. 921, sphalm. = Saxifraga Tilingiana.
Torreyi, *G. Don, Gen. Syst.* iii. 121.—Am. bor. occ.
tortuosum, *Hemsl. Diagn. Pl. Nov.* 10.—Mexic.
transbaikalense, Schlecht. ex Ledeb. Fl. Ross. ii. 187 = album.
trevericum, Rosbach, ex Nym. Consp. 261 = pruinatum.
trifidum, *Wall. Cat.* n. 7230.—Reg. Himal.
triphyllum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 540 = S. Telephium.
tristriatum, *Boiss. & Heldr. Diagn.* Ser. I. x. 16.—Lycaonia; Ins. Creta.
trullipetalum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 102.—Reg. Himal.
tuberosum, *Coss. & Letourn. in Bull. Soc. Bot. Fr.* xxii. (1875) 9.—Afr. bor.
turgidum, Ram. ex DC. Fl. Fr. iv. 390 = album.
turgidum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 452 = brevifolium.
umbilicoides, *Regel, Gartenfl.* (1877) 290; *et in Act. Hort. Petrop.* v. (1877) 263.—Turkestan.
uniflorum, *Hook. & Arn. Bot. Beech. Voy.* 263.—Ins. Luchu.
uniflorum, Rafin. Caratteri, 73 = S. Cepaea.
uralense, *Rupr. in Hofm. Noerdl. Ural*, ii. *App.* 11, 34.—Mont. Ural.
Urvillei, DC. Prod. iii. 408 = pallidum.
variegatum, *S. Wats. in Proc. Am. Acad.* xi. (1876) 137.—Calif.
Verloti, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 606 = anopetalum.
vermiculare, *Pall. Reise Suedl. Statth. Russ. Reich.* i. 337, 345.—Reg. Cauc.
verticillatum, Latour, ex DC. Prod. iii. 404 = S. Cepaea.
verticillatum, *Linn. Sp. Pl.* 430.—Kamtschat.
villosum, *Linn. l. c.* 432.—Europ.; Greenl.
villosum, Webb, Iter Hisp. 46 = dasyphyllum.
virens, Ait. Hort. Kew. ed. I. ii. 110 = rupestre.
virescens, Willd. Enum. Hort. Berol. Suppl. 25 = rupestre.
viridulum, *Haw. in Phil. Mag.* (1827) 183.—Europ.
viviparum, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 137.—Manchur.
vogesiacum, Nym. Consp. 260 = S. Telephium.

SEDUM:—
vulgare, Link, Enum. Hort. Berol. i. 437; S. F. Gray, Nat. Arr. Brit. Pl. ii. 539 = S. Telephium.
Wallichianum, Hook. Ic. Pl. t. 604 = asiaticum.
Whitmanni, Hort. ex Mast. in Gard. Chron. (1878) II. 751 = seq. ?
Wightmannianum, Hort. ex Mast. l. c. = seq. ?
Witmanni, Hort. l. c. 685 = hispanicum.
Wrightii, *A. Gray, Pl. Wright.* i. 76.—Texas.
yantaiense, *Debeaux, in Act. Linn. Soc. Bord.* xxxi. (1877) 160 (*Fl. du Tché-fou*, 65).—China.

SEEMANNANTHA, Alef. in Bonplandia, x. (1862) 264 = **Tephrosia**, Pers. (Legumin.).
strigosa, Alef. l. c. = T. tenuis.
tenuis, Alef. l. c. = T. tenuis.

SEEMANNIA, Hook. Lond. Journ. Bot. vii. (1848) 567 = **Pentagonia**, Benth. (Rubiac.).

SEEMANNIA, Regel, Gartenfl. iv. (1855) 133. t. 126. *GESNERACEAE*, Benth. & Hook. f. ii. 1000.
Benaryi, *Regel, l. c.* (1874) 353.—Am. trop.
silvatica, Hanst. in Linnaea, xxix. (1857-58) 540 = ternifolia.
ternifolia, *Regel, Gartenfl.* iv. (1855) 133. t. 126.—Peruv.

SEETZENIA, R. Br. in Denh. & Clapp. Trav. App. 231 (1826). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 266.
africana, *R. Br. l. c.*—Afr. austr.
orientalis, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 281.—Sind.; Afr. bor. et austr.; Arab.
prostrata, Eckl. & Zeyh. Enum. 98 = africana.

SEEZENIA, Nees, in R. Br. Verm. Bot. Schr. iv. 46 (1830), sphalm. = praec.

SEGERETIA, G. Don, Gen. Syst. ii. 28 (1832), sphalm. = **Sageretia**, Brongn. (Rhamn.).

SEGUIERA, Adans. Fam. ii. 443 (1763), sphalm. = **Seguieria**, Loefl. (Phytol.).

SEGUIERA, Reichb. ex Oliver, Fl. Trop. Afr. ii. 424 (1871) = **Combretum**, Loefl.
nudiflora, Reichb. l. c. = C. aculeatum.

SEGUIERIA, Loefl. Iter Hisp. 191 (1758). *PHYLOLACCACEAE*, Benth. & Hook. f. iii. 83.
aculeata, Jacq. Select. Am. 170 = americana.
americana, *Linn. Syst.* ed. X. 1074.—Am. austr.
asiatica, *Lour. Fl. Cochinch.* 341.—Cochinch.
coriacea, *Benth. in Trans. Linn. Soc.* xviii. (1839) 235.—Bras.
floribunda, *Benth. l. c.*—Bras.
foliosa, *Benth. l. c.* 236.—Guian.
guaranitica, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1883) 88.—Reg. Argent.
Langsdorffii, *Moq. in DC. Prod.* xiii. II. 6.—Bras.
longifolia, *Benth. in Trans. Linn. Soc.* xviii. (1839) 235.—Bras.
macrophylla, *Benth. l. c.*—Guian.
parvifolia, *Benth. l. c.*—Guian.
Vauthieri, *Moq. in DC. Prod.* xiii. II. 7.—Bras.

SEGUINUM, Rafin. Fl. Tellur. iii. 66 (1836) = **Dieffenbachia**, Schott (Aroid.).
maculatum, Rafin. l. c. 67 = D. Seguine.

SEHIMA, Forsk. Fl. Aegypt. Arab. 178 = **Ischaemum**, Linn. (Gramin.).
inscalptum, Hochst. in Flora, xxvii. (1844) 247 = I. laxum.
ischaemoides, Forsk. Fl. Aegypt. Arab. 178 = I. laxum.
Kotschyi, Hochst. in Flora, xxvii. (1844) 247 = I. laxum.

SEIDELIA, Baill. Etud. Gén. Euph. 465. t. 9 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 310.
Mercurialis, *Baill. l. c.* 466.—Afr. austr.
pumila, Baill. l. c. = S. Mercurialis.

SEIDELLA, Benth. & Hook. f. Gen. iii. 310 (1880), sphalm. = praec.

SEIDLIA, Kostel. Allg. Med.-Pharm. Fl. v. 1945 (1836) = **Vatica**, Linn. (Dipteroc.).
lanceaefolia, Kostel. l. c. = V. lanceaefolia.

SEIDLIA, Opiz, Natural. xi. (1826) 349 = **Scirpus**, Tourn. (Cyperac.).
atrovirens, Opiz, l. c. = Scirpus Holoschoenus.
Jechlii, Opiz, ex Nym. Consp. 763 = Scirpus sylvaticus.
radicans, Opiz, Natural. xi. (1826) 349 = Scirpus sylvaticus.
sylvatica, Opiz, l. c. = Scirpus sylvaticus.

SEIDLITZIA, Bunge, ex Boiss. Fl. Orient. iv. 950 (1879). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 69.
flordia, *Bunge, l. c.*—As. Min.; Persia.
Rosmarinus, *Bunge, l. c.* 951.—Arab.

SEKIKA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 209; Phil. Bot. ii. 49 (1791) = **Saxifraga**, Tourn.
cyclaminea, Medic. ll. cc. = Sax. sarmentosa.
sarmentosa, Moench, Meth. Suppl. 77 = Sax. sarmentosa.

SELAGO, Adans. Fam. ii. 268 (1763) = **Polycnemum**, Linn. (Chenop.).

SELAGO, Linn. Syst. ed. I (1735). *SELAGINEAE*, Benth. & Hook. f. ii. 1128.
KURITIS, Rafin. Fl. Tellur. ii. 63 (1836).
MACRIA, Spach, Hist. Vég. Phan. ix. 245 (1840).
MANETTIA, Crantz, Inst. i. 580 (1766).
NOLTIA, Eckl. ex Steud. Nom. ed. II. i. 197, 554 (1841).
VORMIA, Adans. Fam. ii. 284 (1763).
WALAFRIDA, E. Mey. Comm. Pl. Afr. Austr. 372 (1837).
abietina, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
acaulis, Herb. Russ. ex Nees, in DC. Prod. xi. 256 = Lepidagathis cristata.
adpressa, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 103. t. 4.—Afr. austr.
albida, *Choisy, in DC. Prod.* xii. 18.—Afr. austr.
angustifolia, Sieber, ex Choisy, l. c. 19 = decipiens.
angustifolia, Thunb. Prod. Pl. Cap. 99 = Agathelpis angustifolia.
apiculata, *E. Mey. Comm. Pl. Afr. Austr.* 256.—Afr. austr.
appressa, Eckl. & Zeyh. ex Choisy, in DC. Prod. xii. 9 = glabrata.
arguta, *E. Mey. Comm. Pl. Afr. Austr.* 260.—Afr. austr.
articulata, *Thunb. Prod. Pl. Cap.* 99.—Afr. austr.
ascendens, *E. Mey. Comm. Pl. Afr. Austr.* 259.—Afr. austr.
aspera, *Choisy, in DC. Prod.* xii. 12.—Afr. austr.
bracteata, Thunb. Prod. Pl. Cap. 100 = Microdon lucidus.
Burmanni, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 108.—Afr. austr.
canescens, *Linn. f. Suppl.* 284.—Afr. austr.
canescens, E. Mey. Comm. Pl. Afr. Austr. 265 = albida.
capitata, Berg. Desc. Pl. Cap. 157 = Microdon ovatus.
capitata, *Linn. Mant.* ii. 568.—Afr. austr.
capituliflora, *Rolfe, in Journ. Linn. Soc.* xx. (1883) 355.—Afr. austr.
cephalophora, E. Mey. Comm. Pl. Afr. Austr. 256 = nutans.
cephalophora, *Thunb. Prod. Pl. Cap.* 100; *Fl. Cap. ed. Schult.* 464.—Afr. austr.
Choisiana, E. Mey. Comm. Pl. Afr. Austr. 268 = paniculata.
ciliata, *Linn. f. Suppl.* 285.—Afr. austr.
cinerascens, Choisy, in DC. Prod. xii. 9 = albida.
cinerascens, E. Mey. Comm. Pl. Afr. Austr. 270 = cinerea.
cinerea, Eckl. & Zeyh. ex Choisy, in DC. Prod. xii. 18 = witbergensis.
cinerea, Krauss, ex Choisy, l. c. 9 = glabrata.
cinerea, *Linn. f. Suppl.* 285.—Afr. austr.
coccinea, Linn. Pl. Afr. Rar. 12; Amoen. Acad. vi. 89 = spuria.
coerulea, Bernh. ex Steud. Nom. ed. I. 760, nomen.—Hab.?
coerulea, *Burch. ex Krauss, in Flora*, xxviii. (1845) 70, nomen.—Afr. austr.
comosa, *E. Mey. Comm. Pl. Afr. Austr.* 255.—Afr. austr.

SELAGO:—
congesta, *Rolfe, in Journ. Linn. Soc.* xx. (1883) 356.—Afr. austr.
cordata, E. Mey. Comm. Pl. Afr. Austr. 257 = decumbens.
cordata, Thunb. Prod. Pl. Cap. 100 = Phyllopodium heterophyllum.
corymbosa, Linn. Herb. ex Rolfe, in Journ. Linn. Soc. xx. (1883) 342 = stricta.
corymbosa, *Linn. Sp. Pl.* 629.—Afr. austr.
decipiens, *E. Mey. Comm. Pl. Afr. Austr.* 270.—Afr. austr.
decumbens, Choisy, in DC. Prod. xii. 18 = quadrangularis.
decumbens, *Thunb. Prod. Pl. Cap.* 100.—Afr. austr.
densifolia, Hochst. in Flora, xxviii. (1845) 69 = spuria.
dentata, Poir. Encyc. vii. 57 = spuria.
diffusa, Sieber, ex Choisy, in DC. Prod. xii. 11 = triquetra.
diffusa, Thunb. Herb., partim, ex Rolfe, in Journ. Linn. Soc. xx. (1883) 352 = nigrescens.
diffusa, *Thunb. Prod. Pl. Cap.* 99.—Afr. austr.
diffusa, Zeyh. ex Choisy, in DC. Prod. xii. 11 = tenuis.
distans, *E. Mey. Comm. Pl. Afr. Austr.* 266.—Afr. austr.
divaricata, Linn. f. Suppl. 284 = triquetra.
Dregei, *Rolfe, in Journ. Linn. Soc.* xx. (1883) 353.—Afr. austr.
dubia, Linn. Sp. Pl. 629 = Agathelpis angustifolia.
Eckloniana, *Choisy, in DC. Prod.* xii. 13.—Afr. austr.
elata, *Choisy, l. c.* 12.—Afr. austr.
elegans, *Choisy, l. c.* 14.—Afr. austr.
ericina, *E. Mey. Comm. Pl. Afr. Austr.* 254.—Afr. austr.
ericoides, Linn. Mant. i. 87 = Stilbe ericoides.
fasciculata, Linn. l. c. 250 = serrata.
fruticosa, *Linn. l. c.* i. 87.—Afr. austr.
fruticosa, Linn. f. ex Rolfe, in Journ. Linn. Soc. xx. (1883) 351 = Dischisma fruticosum.
fruticosa, Thunb. ex Choisy, in DC. Prod. xii. 9 = Thunbergii.
fruticosa, Thunb. Prod. Pl. Cap. 98, partim = congesta.
fulvomaculata, Link, Enum. Hort. Berol. ii. 123 = spuria.
geniculata, Choisy, in DC. Prod. xii. 9 = articulata.
geniculata, *Linn. f. Suppl.* 284.—Afr. austr.
Gillii, Hook. Bot. Mag. t. 3028 = myrtifolia.
glabrata, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 104.—Afr. austr.
glandulosa, Choisy, in DC. Prod. xii. 20 = scabrida.
glomerata, E. Mey. Comm. Pl. Afr. Austr. 264, partim = Dregei.
glomerata, *Thunb. Prod. Pl. Cap.* 99.—Afr. austr.
glutinosa, *E. Mey. Comm. Pl. Afr. Austr.* 255.—Afr. austr.
guttata, *E. Mey. l. c.* 259.—Afr. austr.
hamulosa, *E. Mey. l. c.* 257.—Afr. austr.
herbacea, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 109.—Afr. austr.
hermannioides, *E. Mey. Comm. Pl. Afr. Austr.* 267.—Afr. austr.
heterophylla, *E. Mey. l. c.* 256.—Afr. austr.
heterophylla, Thunb. Prod. Pl. Cap. 99 = spuria.
hirta, Choisy, in Mém. Soc. Phys. Genèv. ii. II. (1824) 107 = quadrangularis.
hirta, *Linn. f. Suppl.* 285.—Afr. austr.
hispida, Choisy, in DC. Prod. xii. 10 = stricta.
hispida, *Linn. f. Suppl.* 284.—Afr. austr.
hispida, Sieber, ex Choisy, in DC. Prod. xii. 7 = Dischisma capitatum.
hyssopifolia, *E. Mey. Comm. Pl. Afr. Austr.* 262.—Afr. austr.
incisa, *Hochst. in Flora*, xxviii. (1845) 69.—Afr. austr.
lacunosa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 255.—Afr. trop.
lanceolata, *Choisy, in DC. Prod.* xii. 15.—Afr. austr.
laxiflora, *Choisy, l. c.* 8.—Afr. austr.
leptostachya, E. Mey. Comm. Pl. Afr. Austr. 266 = geniculata.
lobeliacea, Hochst. in Flora, xxviii. (1845) 69 = herbacea.
lucida, Vent. Jard. Malm. 26 = Microdon lucidus.
luxurians, *Choisy, in DC. Prod.* xii. 9.—Afr. austr.
lychnidea, Linn. Pl. Rar. Afr. 12; Amoen. Acad. vi. 89 = Lyperia fragrans.
Meyeri, *Choisy, in DC. Prod.* xii. 8.—Afr. austr.

SELAGO:—
micrantha, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 98.—Afr. austr.
micrantha, Zeyh. ex Choisy, in DC. Prod. xii. 9 = articulata.
minutissima, *Choisy, in Mém. Soc. Phys. Genèv.* ii. II. (1824) 100. t. 3.—Afr. austr.
muralis, *Benth. & Hook. f. Gen.* ii. 1129.—Madag.
myrtifolia, *Reichb. Ic. Bot. Exot.* iii. 10. t. 223.—Afr. austr.
nigrescens, *Rolfe, in Journ. Linn. Soc.* xx. (1883) 352.—Afr. austr.
nutans, *Rolfe, l. c.* 354.—Afr. austr.
Ohlendorffiana, *Lehm. Del. Sem. Hort. Hamb.* (1831) 6, 7.—*Cf. Linnaea*, vi. (1831) *Litt.* 74.—Afr. austr.
ovata, L'Hérit. ex Ait. Hort. Kew. ed. I. ii. 355 = Microdon ovatus.
pallida, Salisb. Prod. 93 = spuria.
paniculata, *Thunb. Prod. Pl. Cap.* 99.—Afr. austr.
peduncularis, *E. Mey. Comm. Pl. Afr. Austr.* 262.—Afr. austr.
philicifolia, Herb. Berol. ex Choisy, in DC. Prod. xii. 10 = cinerea.
Pinastra, Linn. Sp. Pl. ed. II. 876 = Stilbe vestita.
pinea, Walp. Rep. iv. 151, sphalm. = spinea.
Pinguicula, *E. Mey. Comm. Pl. Afr. Austr.* 255.—Afr. austr.
polycephala, *Otto, ex Walp. Rep.* iv. 164.—Afr. austr.
polygaloides, Choisy, in DC. Prod. xii. 11 = geniculata.
polygaloides, Linn. f. Suppl. 284 = Microdon cylindricus.
polystachya, Licht. ex Choisy, in DC. Prod. xii. 9 = Thunbergii.
polystachya, *Linn. Mant.* ii. 250.—Afr. austr.
Prunastri, Linn. Syst. ed. X. 1117 = Stilbe Pinastra.
pterophylla, *Otto, ex Sweet, Hort. Brit.* ed. II. 415, nomen.—Afr. austr.
pulchella, Salisb. Prod. 93 = spuria.
purpurea, Cels, Cat. Arb. Pl. (1817) 34, nomen (Quid?).—Hab.?
pusilla, Thunb. Prod. Pl. Cap. 99 = Vandellia scabra.
quadrangularis, *Choisy, in DC. Prod.* xii. 15.—Afr. austr.
racemosa, *Bernh. in Flora*, xxviii. (1845) 69.—Afr. austr.
ramulosa, *Link, Enum. Hort. Berol.* ii. 124.—Afr. austr.
ramulosa, E. Mey. Comm. Pl. Afr. Austr. 265 = Meyeri.
rapunculoides, Linn. Cent. Pl. ii. 22; Amoen. Acad. vi. 89 = spuria.
recurva, Choisy, in DC. Prod. xii. 19 = fruticosa.
recurva, *E. Mey. Comm. Pl. Afr. Austr.* 271.—Afr. austr.
rotundifolia, *Linn. f. Suppl.* 285.—Afr. austr.
rotundifolia, Thunb. Herb. ex Rolfe, in Journ. Linn. Soc. xx. (1883) 354 = Microdon lucidus.
saxatilis, *E. Mey. Comm. Pl. Afr. Austr.* 269.—Afr. austr.
scabrida, *Thunb. Prod. Pl. Cap.* 99.—Afr. austr.
serrata, *Berg. Desc. Pl. Cap.* 159.—Afr. austr.
spicata, Hochst. ex Choisy, in DC. Prod. xii. 14 = spuria.
spicata, Link, Enum. Hort. Berol. ii. 124 = Microdon ovatus.
spinea, *Link, l. c.* 123.—Afr. austr.
spinosa, Burm. f. Fl. Cap. Prod. 17 = Stoebe aethiopica.
spuria, *Linn. Sp. Pl.* 629.—Afr. austr.
spuria, Lodd. Bot. Cab. t. 391 = cinerea.
spuria, Sieber, ex Choisy, in DC. Prod. xii. 15 = serrata.
squarrosa, *Choisy, l. c.* 16.—Afr. austr.
stricta, *Berg. Desc. Pl. Cap.* 155.—Afr. austr.
stricta, Choisy, in Mém. Soc. Phys. Genèv. ii. II. (1824) 103 = tephrodes.
tenuis, *E. Mey. Comm. Pl. Afr. Austr.* 266.—Afr. austr.
tephrodes, *E. Mey. l. c.* 264.—Afr. austr.
tephrodes, E. Mey. l. c., partim = stricta, adpressa.
teretifolia, Link, Enum. Hort. Berol. ii. 124 = spuria.
teretifolia, Walp. Rep. iv. 154 = Walpersii.
teucriifolia, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
Thomsoni, *Rolfe, in Journ. Linn. Soc.* xxi. (1885) 402—Afr. trop.
Thunbergii, *Choisy, in DC. Prod.* xii. 9.—Afr. austr.
tomentosa, Linn. Pl. Rar. Afr. 13; Amoen. Acad. vi. 90 = Manulea tomentosa.

SELAGO:—
trinervia, *E. Mey. Comm. Pl. Afr. Austr.* 261.—Afr. austr.
triquetra, *Linn. f. Suppl.* 284.—Afr. austr.
verbenacea, *Linn. f. l. c.* 285.—Afr. austr.
Walpersii, *Choisy, in DC. Prod.* xii. 20.—Afr. austr.
witbergensis, *E. Mey. Comm. Pl. Afr. Austr.* 270.—Afr. austr.
Zeyheri, *Choisy, in DC. Prod.* xii. 19.—Afr. austr.

SELAS, Spreng. Syst. ii. 216 (1825) = **Acronychia**, Forst. (Rutac.).
lanceolatum, Spreng. l. c. = A. laurifolia.

SELATIUM, D. Don, ex G. Don, Gen. Syst. iv. 196 (1837) = **Gentiana**, Linn.
assurgens, D. Don, l. c. = G. guayaquilensis.
attenuatum, D. Don, l. c. = G. attenuata.
coccineum, D. Don, l. c. = G. coccinea.
diffusum, D. Don, l. c. = G. Selatium.
foliosum, D. Don, l. c. = G. Donii.
incurvum, G. Don, l. c. = G. incurva.
multicaulis, D. Don, l. c. = G. Pavonii.
thyrsoidea, G. Don, l. c. = G. thyrsoidea.

SELBYA, M. Roem. Syn. Hesper. 89, 126 (1846) = **Aglaia**, Lour. (Meliac.).
montana, M. Roem. l. c. 126 (= *Milnea montana*, Jack).—Sumatra.

SELENIA, Nutt. in Journ. Acad. Philad. v. (1825) 132. t. 6. *CRUCIFERAE*, Benth. & Hook. f. i. 72.
aurea, *Nutt. l. c.*—Am. bor.
dissecta, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 160. t. 1.—Calif.

SELENIPEDIUM, Reichb. f. in Bonplandia, ii. (1854) 116; Xenia Orch. i. 3. tt. 2, 27, 44, 62, 181 (1854). *ORCHIDEAE*, Benth. & Hook. f. iii. 635.
SOLENIPEDIUM, Beer, Prakt. Orch. 310 (1854), sphalm.
UROPEDIUM, Lindl. Orch. Linden. 28 (1846), nomen prius.
Ainsworthii ×, Reichb. f. in Gard. Chron. (1879) i. 748.
albo-pictum, Linden, Cat. n. 21 (1867) 16 (Quid?).—Hab.?
albo-purpureum ×, Reichb. f. in Gard. Chron. (1877) ii. 38.
Boissierianum, *Reichb. f. in Bonplandia*, ii. (1854) 116; iv. (1856) 217.—Peruv.
cardinale ×, Reichb. f. in Gard. Chron. (1882) ii. 488.
caricinum, *Reichb. f. in Bonplandia*, ii. (1854) 116.—Bolivia.
caudatum, *Reichb. f. l. c.* ii. (1854) 116; iv. (1856) 217.—Am. trop.
Chica, *Reichb. f. l. c.* ii. (1854) 116; *Xenia Orch.* i. 3. t. 2.—Panama.
Czerwiakowianum, *Reichb. f. in Bonplandia*, ii. (1854) 116.—Peruv.
dariense, Reichb. f. in Nov. Act. Nat. Cur. xxxv. (1869) ii. 8 = longifolium.
Dominianum ×, Hort. ex Reichb. f. in Gard. Chron. (1870) 1181.
grande ×, Reichb. f. in Gard. Chron. (1882) ii. 488.
Hartwegii, Reichb. f. in Bonplandia, ii. (1854) 116 = longifolium.
Hincksianum, Reichb. f. in Gard. Chron. (1878) i. 202, in nota = longifolium.
Isabelianum, *Rodrig. Orch. Nov.* i. 201.—Bras.
kaieteurium, N. E. Br. in Gard. Chron. (1885) ii. 262 = Lindleyanum.
Klotzschianum, *Reichb. f. in Bonplandia*, ii. (1854) 116.—Guian.
laevigatum, May, in Rev. Hortic. (1885) 301 = Cypripedium philippense.
Lindleyanum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 116; *Xenia Orch.* i. 3.—Am. trop.
longifolium, *Reichb. f. ll. cc.*—Guian.
palmifolium, *Reichb. f. in Bonplandia*, ii. (1854) 116.—Guian.
Parishii, Jolibois, in Rev. Hortic. (1885) 132 = Cypripedium Parishii.
Pearcei, Reichb. f. in Hamb. Gartenz. xxi. (1865) 298 = caricinum.
porphyreum ×, Reichb. f. in Gard. Chron. (1878) i. 366.
Reichenbachii, Endres, ex Reichb. f. l. c. (1869) 1206 = longifolium.
reticulatum, Reichb. f. Xenia Orch. ii. 223 = Boissierianum.

SELENIPEDIUM:—
Roezli, *Reichb. f. in Regel, Gartenfl.* (1871) 164 = longifolium.
Schlimii, *Linden, ex Reichb. f. in Bonplandia*, ii. (1854) 277.—N. Granat.
Sedeni ×, Reichb. f. in Gard. Chron. (1873) 1431.
stenophyllum ×, Reichb. f. l. c. (1876) i. 461.
vittatum, *Reichb. f. in Illustr. Hortic.* xxiii. (1876) 57. t. 238.—Bras.
Wallisii, Reichb. f. Otia Bot. Hamb. 30 = caudatum.
Warszewiczianum, Reichb. f. in Bonplandia, ii. (1854) 116; Xenia Orch. i. 3 = caudatum.

SELENOCARPAEA, Eckl. & Zeyh. Enum. 10 (1835) = **Heliophila**, Burm. f. (Crucif.).
diffusa, Eckl. & Zeyh. l. c. = H. diffusa.
Peltaria, Eckl. & Zeyh. l. c. = H. flacca.
pinnata, Eckl. & Zeyh. l. c. = H. Peltaria.

SELENOCERA, Zipp. ex Span. in Linnaea, xv. (1841) 316. *RUBIACEAE*.
secundiflora, *Zipp. l. c.*—Ins. Timor.

SELENOGYNE, DC. Prod. vi. 147 (1837), sphalm. = SOLENOGYNE, Cass. = **Lagenophora**, Cass. (Compos.).

SELEPSION, Rafin. Fl. Tellur. iii. 48 (1836) = **Urtica**?, Tourn.

SELERANTHUS, Hill, Hort. Kew. 372 (1768), sphalm. = **Scleranthus**, Linn. (Illecebr.).

SELERIA, Boeck. in Linnaea, xxxviii. (1874) 436, sphalm. = **Scleria**, Berg. (Cyperac.).

SELINOCARPUS, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 262. *NYCTAGINEAE*, Benth. & Hook. f. iii. 6.
angustifolius, *Torr. Bot. Mex. Bound.* 170.—Texas.
chenopodioides, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 262.—N. Mexic.
diffusus, *A. Gray*, l. c.—N. Mexic.
Palmeri, *Hemsl. Biol. Centr. Am. Bot.* iii. 6.—Mexic.

SELINON, Adans. Fam. ii. (23) (1763) = **Apium**, Tourn. (Umbell.).

SELINON, Rafin. in Journ. Phys. lxxxix. (1819) 259 = **Selinum**, Linn.
carvifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 134 = Selinum Carvifolia.

SELINOPSIS, Coss. & Dur. ex Munby, Cat. Pl. Alg. ed. II. 13 (1859), nomen = **Carum**, Rupp. (Umbell.).
foetida, Coss. & Dur. l. c.—Alger.
montana, Coss. & Dur. l. c.—Alger.

SELINUM, Linn. Gen. ed. I. 73 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 914.
ALLINUM, Neck. Elem. i. 179 (1790).
ANTHOSCIADIUM, Fenzl, in Endl. Gen. Suppl. iv. iii. 9 (1850).
CENOLOPHIUM, Koch, in Nov. Act. Nat. Cur. xii. i. (1824) (Gen. Umbell.) 103.
CNIDIUM, Cusson, ex DC. Prod. iv. 152 (1830).
GNIDIUM, G. Don, in Loud. Hort. Brit. 107 (1830).
HETEROPTILIS, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 534.
HYALAENA, C. Muell. in Walp. Ann. v. 70 (1858).
HYALOLAENA, Bunge, in Mém. Sav. Étr. Pétersb. vii. (1851) 304.
LITHOSCIADIUM, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 730.
LOMATOCARUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 59 (1839).
MICRANGELIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 388.
MYLINUM, Gaud. Fl. Helv. ii. 344 (1828).
OREOCOMA, Edgew. in Trans. Linn. Soc. xx. (1846) 54.
SPHENOSCIADIUM, A. Gray, in Proc. Am. Acad. vi. (1866) 536.
THYSSELINUM, Adans. Fam. ii. 100 (1763).
acaule, Cav. Ic. v. 59 = Laretia acaulis.
acaule, Pursh, Fl. Am. Sept. ii. 732 = Cymopterus glomeratus.
acaule, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 13.—Sibir. baical.

SELINUM:—
acutangulum, Gilib. Fl. Lituan. ii. 22 = S. Carvifolia.
alatum, Poir. Encyc. Suppl. v. 126 = Ligusticum alatum.
alpinum, Bieb. Fl. Taur. Cauc. i. 236 = Seseli annuum?
alsaticum, Crantz, Stirp. Austr. ed. I. fasc. iii. 17; ed. II. 155 = Peucedanum alsaticum.
anesorrhizum, F. Muell. Sel. Pl. Indust. Cult. [ed. II.] 225 = Annesorhiza capensis.
Anethum, Roth, Tent. Fl. Germ. i. 134 = Peucedanum graveolens.
Angelica, Roth, l. c. 133 = Angelica sylvestris.
Angelicastrum, Link, ex Spreng. Syst. i. 916 = Angelica lancifolia.
angulatum, Lam. Fl. Fr. iii. 419 = S. Carvifolia.
annuum, Steud. Nom. ed. II. ii. 572, in syn., sphalm. = Seseli annuum.
appianum, Pers. Syn. i. 309 = Carum rigidulum.
appuanum, Viv. Cat. Hort. Dinegro, 33; ex ej. Fl. Ital. Fragm. 14 = Carum rigidulum.
Archangelica, Link, Enum. Hort. Berol. i. 269 = Archangelica officinalis.
arenosum, Crantz, ex DC. Prod. iv. 165 = Peucedanum austriacum.
argenteum, All. Fl. Pedem. ii. 9 = Peucedanum involucratum.
argenteum, Crantz, Stirp. Austr. ed. I. fasc. iii. 43; ed. II. 174 = Peucedanum austriacum.
aristatum, Ait. ex Link, Handb. i. 330 = Ligusticum pyrenaicum.
atropurpureum, Link, Enum. Hort. Berol. i. 269 = Archangelica atropurpurea.
austriacum, Bertol. Pl. Gen. 44; et Amoen. Ital. 137 = Peucedanum alsaticum.
austriacum, Jacq. Enum. Stirp. Vindob. 49, 220 = Peucedanum austriacum.
austriacum, Tenore, Fl. Nap. iii. 287 = Peucedanum sulcatum.
baicalense, Redowsk. ex Willd. Enum. Hort. Berol. 306 = Peucedanum baicalense.
banaticum, *Kit. in Rochel, Pl. Banat. Index*, nomen.—Banat.
Bellardi, Balb. Cat. Hort. Taur. (1821) 51 = Peucedanum austriacum.
Benthami, S. Wats. Bibliog. Ind. N. Am. Bot. 432 = Gmelini.
canadense, *Michx. Fl. Bor. Am.* i. 165.—Am. bor.
Candollei, Edgew. in Trans. Linn. Soc. xx. (1846) 55 = tenuifolium.
Candollii, *DC. Prod.* iv. 165.—Reg. Himal.
cantabrigense, Fisch. ex DC. l. c. 179 = Peucedanum palustre.
capitellatum, *Benth. & Hook. f. ex S. Wats. in Bot. King's Exp.* 126.—Am. bor.
carniolicum, Hoppe, ex Nym. Consp. 285 = Peucedanum austriacum.
Carvifolia, Gilib. ex DC. Prod. iv. 152 (= *Cenolophium Fischeri*).—Rossia; Reg. Cauc.
Carvifolia, *Linn. Sp. Pl.* ed. II. 350.—Europ.
carvifolium, Crantz, Inst. ii. 126. n. 5 = praec.
carvifolium, Crantz, Stirp. Austr. ed. I. fasc. iii. 22; ed. II. 162 = Peucedanum carvifolium.
caucasicum, Bieb. Fl. Taur. Cauc. i. 213 = Peucedanum caucasicum.
cenolophioides, Turcz. ex Ledeb. Fl. Ross. ii. 284 = sylvestre.
Cervaria, Linn. Sp. Pl. 1194 = Peucedanum Cervaria.
cervicaria, Roth, Tent. Fl. Germ. i. 132 = praec.
Chabraei, DC. Prod. iv. 176, in syn. = Peucedanum carvifolium.
Chabraei, Rebent. & Kunth, Fl. Berol. 78 = Monnieri.
cicutarium, Benth. & Hook. f. Gen. i. 914 = tenuifolium.
cnidiifolium, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 72.—Sibir. or.
collinum, Kit. ex DC. Prod. iv. 179 = Peucedanum baicalense.
coniifolium, Benth. & Hook. f. Gen. i. 914 = tenuifolium.
decipiens, Schrad. & Wendl. Sert. Hannov. i. iii. 23. t. 13 (= *Melanoselinum decipiens*).—Ins. Canar.
decussatum, [Clairv.] Man. Herb. 80 = Peucedanum carvifolium?
densiflorum, Salisb. Prod. 162 = Monnieri.
dimidiatum, DC. Fl. Fr. iv. 323 = Seseli annuum.
dissectum, *Wall. Cat.* n. 581; *et ex DC. Prod.* iv. 178. (sp. dub.).—Nepal.

SELINUM:—
divaricatum, *Buch, Beschr. Canar. Ins.* 195.—Ins. Canar.
dubium, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 735 = Gmelini.
elegans, Balb. in Mém. Acad. Turin (1810), 364; Hort. Taur. Fasc. 28. t. 4 = Peucedanum austriacum.
frutescens, Pers. Syn. i. 309 = *decipiens*.
Galbanum, Spreng. in Schult. Syst. vi. 563 = Peucedanum Galbanum.
glaucum, Lam. Fl. Fr. iii. 419 = Peucedanum Cervaria.
glaucum, Wall. Cat. n. 580; et ex DC. Prod. iv. 179 = Peucedanum glaucum.
Gmelini, *Bray, in Denkschr. Bot. Ges. Regensb.* i. II. (1818) 36.—Reg. bor.
gummiferum, Spreng. in Schult. Syst. vi. 564 = Seseli gummiferum.
humile, Turcz. in Bull. Soc. Nat. Mosc. (1838) 93 = Peucedanum salinum.
Imperatoria, Crantz, Stirp. Austr. ed. I. fasc. iii. 42; ed. II. 174 = Peucedanum Ostruthium.
imperatorioides, Link, Enum. Hort. Berol. i. 269 = Peucedanum angustifolium.
intermedium, Bess. Prim. Fl. Galic. i. 205 = Peucedanum palustre.
Kingii, *S. Wats. in Bot. King's Exp.* 126.—Am. bor. occ.
Kraussianum, *Benth. & Hook. f. Gen.* i. 914.—Afr. austr.
Lachenalii, C. C. Gmel. Fl. Bad. i. 640 = pyrenaicum.
lactescens, Gilib. Fl. Lituan. ii. 22 = Peucedanum palustre.
lactescens, Lam. Fl. Fr. iii. 418 = Peucedanum carvifolium.
laevicaule, Stokes, Bot. Mat. Med. ii. 75 = sylvestre.
latifolium, Bieb. Fl. Taur. Cauc. i. 213 = Peucedanum latifolium.
lineare, Schumach. Enum. Pl. Saelland. i. 95 = sylvestre.
lineare, Spreng. in Schult. Syst. vi. 560 = Peucedanum austriacum.
livonicum, Ledeb. ex DC. Prod. iv. 169 = Angelica chaerophyllea.
membranaceum, Vill. Cat. Jard. Strasb. t. 6 = S. Carvifolia.
microphyllum, Cav. Ic. v. 59 = Mulinum microphyllum.
Monnieri, *Linn. Cent. Pl.* i. 9; *Amoen. Acad.* iv. 269. —Europ.
montanum, Schleich. ex Willd. Enum. Hort. Berol. 306 = Peucedanum austriacum.
neglectum, Láng, ex Reichb. Fl. Germ. Excurs. 461 (Quid?).—Banat.
nigricans, Gaud. Fl. Helvet. ii. 333 = Peucedanum austriacum.
nigrum, Lam. Fl. Fr. iii. 420 = Peucedanum austriacum.
obscurum, Fisch. ex Trev. Ind. Sem. Hort. Wratisl. App. iii. (1821) [3] = Peucedanum baicalense.
odoratum, Bernh. ex Spreng. Sp. Umbellif. 60 = Angelica palustris.
Opopanax, Crantz, Class. Umbell. 63 = Opopanax Chironium.
Oreoselinum, Crantz, Stirp. Austr. ed. I. fasc. iii. 33; ed. II. 169 = Peucedanum Oreoselinum.
orientale, *Benth. & Hook. f. Gen.* i. 914.—As. Min.; Syria.
Ostruthium, Wallr. Ann. Bot. 41 = Peucedanum Ostruthium.
pacificum, *S. Wats. in Proc. Am. Acad.* xi. (1876) 140.—Am. bor. occ.
palustre, Crantz, Stirp. Austr. ed. I. fasc. iii. 39; ed. II. 172 = S. Carvifolia.
palustre, Linn. Sp. Pl. 244 = Peucedanum palustre.
palustre, Thuill. Fl. Par. ed. II. 139 = Peucedanum carvifolium.
paniculatum, Spreng. Pugill. ii. 50 = Peucedanum Sprengelii.
papyraceum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 701.—Reg. Himal.
Pastinaca, Crantz, Stirp. Austr. ed. I. fasc. iii. 21; ed. II. 161 = Peucedanum sativum.
peregrinum, Willd. Enum. Hort. Berol. 306 = Carum rigidulum.
peucedanoides, Brot. Phyt. Lusit. ii. 196. t. 167 = Peucedanum lancifolium.

SELINUM:—
peucedanoides, Desf. Tabl. ed. II. 142 = Peucedanum involucratum.
Peucedanum, Crantz, Stirp. Austr. ed. I. fasc. iii. 30; ed. II. 167; Roth, Tent. Fl. Germ. i. 133 = Peucedanum officinale.
podolicum, Bess. Prim. Fl. Galic. ii. 392 = Peucedanum carvifolium.
polymorphum, Spreng. Pugill. ii. 50 = Peucedanum austriacum.
pratense, Spreng. Fl. Hal. 92. t. 2 = sylvestre.
proliferum, Cav. Ic. v. 58 = Mulinum proliferum.
Pseudo-carvifolia, All. Fl. Pedem. ii. 9 = S. Carvifolia.
pubescens, Moench, Meth. 80 = Angelica sylvestris.
pyrenaeum, *Gouan, Illustr.* ii. t. 5.—Mont. Pyren.
rablense, Spreng. Pugill. ii. 51 = Peucedanum austriacum.
ramosissimum, Wall. Cat. n. 578 = Peucedanum ramosissimum.
Razulii, Link, Enum. Hort. Berol. i. 269 = Angelica Razulii.
rigidulum, Viv. Fl. Ital. Fragm. i. 14. t. 15 = Carum rigidulum.
Rochelii, Heuff. ex Rochel, Bot. Reise Banat, 78 = Silaus carvifolius.
scabrum, Lapeyr. Hist. Abr. Pl. Pyr. 147 = Xatardia scabra.
Schiwerekii, Bess. Prim. Fl. Galic. i. 205 = Peucedanum palustre.
Seguirei, Baumg. Enum. Stirp. Transs. i. 224 = S. Carvifolia.
Seguieri, Jacq. Hort. Vindob. i. 24. t. 61 = Ligusticum Seguieri.
sibiricum, Retz. Obs. ii. 16 (= *Peucedanum selinoides*). —Sibir.
silvestre, Pourr. ex Nym. Consp. 283 = Angelica sylvestris.
spinosum, Cav. Ic. v. 59 = Mulinum spinosum.
spinulifolia, Stokes, Bot. Mat. Med. ii. 73 = Peucedanum Cervaria.
stellatum, D. Don, Prod. Fl. Nep. 185 = Pleurospermum stellatum.
striatum, *Benth. & Hook. f. Gen.* i. 914.—Reg. Himal.
sublactescens, Gilib. Fl. Lituan. ii. 21 = Peucedanum palustre.
suffruticosum, *Benth. & Hook. f. Gen.* i. 914.—Afr. austr.
sulcatum, Bertol. Fl. Ital. iii. 367 = Peucedanum sulcatum.
sylvestre, Crantz, Stirp. Austr. ed. I. fasc. iii. 47; ed. II. 177 = Angelica sylvestris.
sylvestre, *Linn. Sp. Pl.* 244.—Europ.
sylvestre, Oed. Fl. Dan. t. 412 = Peucedanum palustre.
tenuifolium, Salisb. Prod. 162 = S. Carvifolia.
tenuifolium, *Wall. Cat.* n. 579; *et ex DC. Prod.* iv. 181.—Reg. Himal.
terebinthinaceum, Fisch. ex Trev. Ind. Sem. Hort. Wratisl. App. iii. (1821) [3] = Peucedanum terebinthaceum.
terebinthinum, Hook. Fl. Bor. Am. i. 266. t. 95 = Cymopterus terebinthinus.
turfosum, Baumg. Enum. Stirp. Transs. i. 223 = sylvestre.
Tysselinum, Crantz, Stirp. Austr. ed. I. fasc. iii. 35; ed. II. 170 = Peucedanum palustre.
uliginosum, Link, ex Steud. Nom. ed. I. 762 = Laserpitium peucedanoides.
univittatum, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 735 = Gmelini.
vaginatum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 700.—Reg. Himal.
venetum, Spreng. Sp. Umbellif. 73 = Peucedanum venetum.
verticillare, Link, Enum. Hort. Berol. i. 269 = Peucedanum verticillare.

SELKIRKIA, Hemsl. Bot. Challenger, i. III. 47 (1884). *BORAGINEAE.*
Berteroi, *Hemsl. l. c.* 48.—Ins. J. Fernand.

SELLIERA, Cav. in Anal. Hist. Nat. i. (1799) 41. t. 5. f. 2; et Ic. v. 49. t. 474 (1799). *GOODENOVIEAE*, Benth. & Hook. f. ii. 539.
exigua, *F. Muell. Fragm.* iii. 142.—Austral.
fasciculata, *F. Buch. in Trans. N. Z. Inst.* iii. (1871) 211.—N. Zel.

SELLIERA:—
herpystica, Schlecht. in Linnaea, xx. (1847) 598 = radicans.
radicans, *Cav. in Anal. Hist. Nat.* i. (1799) 41. t. 5. f. 2; Ic. v. 49. t. 474.—Austral.
repens, De Vriese, in Verh. Haarl. x. (1854) 162 = radicans.

SELLIERIA, J. Buch. in Trans. N. Z. Inst. iii. (1871) 211 = praec.

SELLOA, H. B. & K. Nov. Gen. et Sp. iv. 265. t. 395 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 361.
FEAEA, Spreng. Syst. iii. 362 (1826).
linearis, *DC. Prod.* v. 612 (an Spilanthes sp.?).—Reg. Argent.
plantaginea, *H. B. & K. Nov. Gen. et Sp.* iv. 266. t. 395.—Mexic.

SELLOA, Spreng. Nov. Prov. Hort. Hal. 36 (1818) = **Gymnosperma**, Less. (Compos.).
capensis, Spreng. Syst. iii. 496 = Denekia capensis.
glutinosa, Spreng. Nov. Prov. Hort. Hal. 36 = Baccharis glutinosa.
nudata, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 300 = Flaveria linearis.

SELLOWIA, Roth, ex Roem. & Schult. Syst. v. p. xxxi. 407 (1819) = **Ammannia**, Linn. (Lythrar.).
uliginosa, Roth, l. c = A. pentandra.

SELLUNIA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 358 = **Vicia**, Tourn. (Legumin.).
lunata, Alef. l. c. 359 = V. lunata.

SELNORITION, Rafin. Sylva Tellur. 160 (1838) = **Rubus**, Tourn. (Rosac.).
canadensis, Rafin. l. c. = R. canadensis.
cesius, Rafin. l. c. = R. caesius.
obovalis, Rafin. l. c. = R. hispidus.
saxatilis, Rafin. l. c. = R. saxatilis.

SELONIA, Regel, in Bull. Soc. Nat. Mosc. xli. (1868) I. 457 = **Eremurus**, Bieb. (Liliac.).
soogdiana, Regel, l. c. 458 = E. soogdiana (in Addend.).

SELWYNIA, F. Muell. Fragm. iv. 153 (1864). *MENISPERMACEAE.*
laurina, *F. Muell. l. c.*—Austral.

SELYSIA, Cogn. in DC. Monog. Phan. iii. 735 (1881). *CUCURBITACEAE.*
cordata, *Cogn. l. c.* 736.—N. Granat.
prunifera, *Cogn. l. c.*—Peruv.

SEMARILLA, Rafin. Sylva Tellur. 146 (1838) = **Celastrus**, Linn.
bicolor, Rafin. l. c.—Hab.?

SEMARILLARIA, Ruiz & Pav. Prod. 54. t. 9 (1794) = **Paullinia**, Linn. (Sapind.).
acutangula, Ruiz & Pav. Syst. Veg. 93 = P. acutangula.
alata, Ruiz & Pav. Fl. Per. iv. t. 340 = P. alata.
Cururu, Ruiz & Pav. ex Steud. Nom. ed. II. ii. 556 = Serjania nodosa.
enneaphylla, Ruiz & Pav. Fl. Per. iv. t. 341 = P. enneaphylla.
nitida, Ruiz & Pav. l. c. 339 = P. neglecta.
obovata, Ruiz & Pav. Syst. Veg. 93 = P. obovata.
subrotunda, Ruiz & Pav. l. c. 92 = P. subrotunda.

SEMECARPOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = seq.

SEMECARPUS, Linn. f. Suppl. 285 (1781). *ANACARDIACEAE*, Benth. & Hook. f. i. 424.
acuminata, *Thw. Enum. Pl. Zeyl.* 76.—Zeylan.
acuminata, Wall. ex Voigt, Hort. Suburb. Calc. 271; Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 75 = S. Cassuvium.
albescens, *Kurz, l. c.* xl. (1871) II. 51.—Burma.
Anacardium, Blanco, Fl. Filip. ed. I. 217 = pubescens.
Anacardium, Blume, Bijdr. 1156 = heterophylla.
Anacardium, *Linn. f. Suppl.* 182.—As. et Austral. trop.

SEMECARPUS:—
aruensis, *Engl. in DC. Monog. Phan.* iv. 484.—Ins. Aru.
atra, Vieill. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 71 = Oncocarpus vitiensis.
auriculata, *Bedd. Fl. Sylv.* ii. t. 232.—Ind. or.
australiensis, *Engl. in DC. Monog. Phan.* iv. 482.—Austral.
Balansae, *Engl. l. c.* 484.—N. Caled.
caesia, Blume, Mus. Bot. Lugd. Bat. i. 189 = heterophylla.
Cassuvium, *Roxb. Hort. Beng.* 22; *Fl. Ind.* ii. 85; *Spreng. Syst.* i. 936.—Burma; Malaya.
cochinchinensis, *Engl. in DC. Monog. Phan.* iv. 489.—Cochinch.
coriacea, *Thw. Enum. Pl. Zeyl.* 76.—Zeylan.
cuneata, *Engl. in DC. Monog. Phan.* iv. 493 (sp. dub.).—Zeylan.
cuneifolia, Kurz, in Pegu, Rep. A. 42 = pandurata.
cuneifolia, Roxb. Hort. Beng. 22; Fl. Ind. ii. 86 = S. Anacardium.
cuneiformis, Blanco, Fl. Filip. ed. I. 220 = microcarpa.
Forstenii, *Blume, Mus. Bot. Lugd. Bat.* i. 188.—Ins. Molucc.
fulvinervis, *Blume, l. c.* 189.—Borneo.
Gardneri, *Thw. Enum. Pl. Zeyl.* 76, 410.—Zeylan.
glauca, *Engl. in DC. Monog. Phan.* iv. 478.—Borneo.
Grahamii, Wight, Ic. t. 235 = Holigarna Grahamii.
grandifolia, Wall. Cat. n. 985 = Campnosperma Griffithii.
heterophylla, *Blume, Mus. Bot. Lugd. Bat.* i. 187.—Java.
heterophylla, Hook. f. Fl. Brit. Ind. ii. 35 = albescens.
insularum, *Seem. Fl. Vit.* 50.—N. Guin.
Kurzii, *Engl. in DC. Monog. Phan.* iv. 489.—Ins. Nicobar.
laevigata, *Thw. ex Hook. f. Fl. Brit. Ind.* ii. 35.—Zeylan.
latifolia, Pers. Syn. i. 324 = S. Anacardium.
longifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 188.—Ins. Timor.
lurida, *Hook. f. Fl. Brit. Ind.* ii. 34.—Malacca.
mangifera, Wight & Arn. in Edinb. N. Phil. Journ. xv. (1833) 177, in obs., sphalm. = S. Anacardium.
marginata, *Thw. Enum. Pl. Zeyl.* 77.—Zeylan.
microcarpa, *Wall. Cat.* n. 989.—Burma.
monogyna, Soyer-Willem. in Usteri, Ann. Bot. xviii. (1796) 25 (Quid?).—Ins. Maurit.
Moonii, *Thw. Enum. Pl. Zeyl.* 77.—Zeylan.
neocaledonica, *Engl. in DC. Monog. Phan.* iv. 482.—N. Caled.
nicobarensis, *Engl. l. c.* 480.—Ins. Nicobar.
nigroviridis, *Thw. Enum. Pl. Zeyl.* 76, 410.—Zeylan.
oblongifolia, Thw. l. c. 76 = obscura.
obovata, *Moon, Cat. Pl. Ceyl.* 22, nomen; *Thw. l. c.* 77.—Zeylan.
obscura, *Thw. l. c.* 76.—Zeylan.
obscura, Thw. l. c. 410, partim = Walkeri.
pandurata, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 209.—Burma.
parvifolia, *Thw. Enum. Pl. Zeyl.* 77.—Zeylan.
Perrottetii, *March. Rév. Anacard.* 169.—Ins. Celebes.
philippinensis, *Engl. in DC. Monog. Phan.* iv. 481.—Ins. Philipp.
pubescens, *Thw. Enum. Pl. Zeyl.* 77.—Zeylan.
Roxburghii, *Blume, Mus. Bot. Lugd. Bat.* i. 188.—Borneo.
scabrida, *Blume, l. c.* 189.—Borneo.
subpanduraeformis, *Wall. Cat.* n. 987.—Ind. or.
subpeltata, *Thw. Enum. Pl. Zeyl.* 75.—Zeylan.
subracemosa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 304.—Burma.
Thwaitesii, Hook. f. Fl. Brit. Ind. ii. 31 = pubescens.
travancorica, *Bedd. Fl. Sylv.* ii. t. 232.—Ind. or.
vitiensis, Engl. in DC. Monog. Phan. iv. 483 = Oncocarpus vitiensis.
Walkeri, *Hook. f. Fl. Brit. Ind.* ii. 33.—Zeylan.
zeylanica, Blume, Mus. Bot. Lugd. Bat. i. 189 = obscura?

SEMEIANDRA, Hook. & Arn. Bot. Beech. Voy. 291. t. 59 (1839). *ONAGRARIEAE*, Benth. & Hook. f. i. 791.
grandiflora, *Hook. & Arn. l. c.*—Mexic.

SEMEIOCARDIUM, Zoll. in Tijdschr. Nederl. Ind. xvii. (1858) 245 = **Polygala**, Linn.
Arriensii, Zoll. l. c.—Java.
glaucescens, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 151 = P. triphylla.
Hamiltonii, Hassk. l. c. = P. triphylla.
hyalinum, Hassk. l. c. = P. triphylla.

SEMEIONOTIS, Schott, in Wien. Zeitschr. iii. (1829) 804; ex Endl. in Linnaea, vi. (1831) Litt. 55 = **Dalbergia**, Linn. (Legumin.).
floribunda, Schott, ll. cc. (Quid?).—Bras.

SEMELE, Kunth, in Abh. Akad. Berl. (1842) 49. *LILIACEAE*, Benth. & Hook. f. iii. 765.
AMPHION, Salisb. Gen. Pl. Fragm. 66 (1866).
androgyna, *Kunth, Enum. Pl.* v. 277.—Ins. Canar.

SEMENOVIA, Regel & Herd. in Bull. Soc. Nat. Mosc. xxxix. (1866) ii. 78. *UMBELLIFERAE*.
transiliensis, *Regel & Herd. l. c.* 79.—Turkestan.

SEMETOR, Rafin. Sylva Tellur. 69 (1838) = **Aspalathus**, Linn. (Legumin.).
arborea, Rafin. l. c. = A. arborea.

SEMIDOPSIS, Zumag. Fl. Pedem. i. 249 (1849) = **Alnus**, Tourn. (Cupul.).
viridis, Zumag. l. c. 250 = A. viridis.

SEMILTA, Rafin. Sylva Tellur. 63 (1838) = **Croton**, Linn. (Euphorb.).
altheaefolia, Rafin. l. c. = C. altheaifolius.

SEMIRAMISIA, Klotzsch, in Linnaea, xxiv. (1851) 25. *VACCINIACEAE*, Benth. & Hook. f. ii. 569.
Karsteniana, *Klotzsch, l. c.* 26.—Venezuela.
speciosa, *Klotzsch, l. c.* 25.—Ecuador.

SEMNOS, Rafin. Sylva Tellur. 161 (1838) = **Chilianthus**, Burch. (Logan.).
paniculata, Rafin. l. c. = C. oleaceus.

SEMONVILLEA, J. Gay, in Bull. Féruss. xviii. (1829) 412. *FICOIDEAE*, Benth. & Hook. f. i. 859.
DITROCHE, E. Mey. ex DC. Prod. xiii. ii. 19 (1849).
chenopodioides, J. Gay, ex Moq. in DC. Prod. xiii. ii. 19 = pterocarpa.
fenestrata, *Fenzl, in Nov. Stirp. Dec.* 42.—Afr. austr.
portulacoides, Mus. Par. ex Moq. in DC. Prod. xiii. ii. 20 = Limeum viscosum.
pterocarpa, *J. Gay, in Bull. Féruss.* xviii. (1829) 412.—Afr. trop.
punctata, Steud. Nom. ed. II. ii. 556 = pterocarpa.

SEMPERVIVUM, Rupp. ex Linn. Syst. ed. I (1735). *CRASSULACEAE*, Benth. & Hook. f. i. 660.
AEONIUM, Webb & Berth. Phyt. Canar. i. 184. t. 28 (1836?).
AICHRYSON, Webb & Berth. l. c. 181. t. 27 (1836?).
GREENOVIA, Webb & Berth. l. c. 198. t. 36 (1836?).
JOVIBARBA, Opiz, Seznam, 54 (1852).
SEDUM, Adans. Fam. ii. 248 (1763).
abyssinicum, *Hochst. in Flora*, xxiv. (1841) i. Intell. 30; *et ex A. Rich. Tent. Fl. Abyss.* i. 315.—Abyss.
acuminatum, *Jacquem. Voy. Bot.* 63. t. 74. f. 1.—Reg. Himal.
acuminatum, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 28 = glaucum.
adoxum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 40 = tectorum.
affine, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 269.—Europ.
africanum, *Mill. Gard. Dict.* ed. VIII. n. 7.—Afr. austr.
agriostaphis, *Webb & Berth. Phyt. Canar.* i. 216 (Index), nomen.—Ins. Canar.
aizoides, *Lam. Encyc.* iii. 290.—Ins. Madera.
alatum, *Scheele, in Flora*, xxvi. (1843) 453.—Europ.
albidum, *Schnittsp. & C. B. Lehm. l. c.* xxxviii. (1855) 4.—Hab.?
album, Edgw. in Trans. Linn. Soc. xx. (1846) 49 = sedoides.
Allionii, Nym. Consp. 260 = hirtum.

SEMPERVIVUM:—
alpestre, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 308 = montanum.
alpinum, Griseb. & Schenk, in Linnaea, xxv. (1852) 600 = arvernense.
ambiguum, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 271.—Europ.
angustifolium, *A. Kern. in Zeitschr. Ferdinand. Tirol.* III. xv. (1870) 267.—Tyrol.
annuum, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 154.—Ins. Canar.
anomalum, Hort. ex Baker, in Gard. Chron. (1879) ii. 107 = pumilum?
anomalum, Lag. Gen. et Sp. Nov. 17 = Sedum amplexicaule.
arachnoideum, *Linn. Sp. Pl.* 465.—Europ. austr.
arboreum, *Linn. l. c.* 464.—Reg. Mediterr.
arenarium, *Koch, Syn. Fl. Germ.* ed. I. 833.—Europ. austr.
arenarium, Stev. ex Baker, in Gard. Chron. (1874) ii. 104 = ruthenicum.
armenum, Boiss. & Huet, Diagn. Ser. II. ii. 60 = globiferum.
arvernense, *Lecoq & Lamotte, Cat. Pl. Vasc. Fr.* 179.—Gallia.
assimile, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 19.—Transsylv.
atlanticum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 454.—Marocc.
aureum, *C. Sm. ex Hornem. Suppl. Hort. Hafn.* 61.—Ins. Canar.
austriacum, Nym. Consp. 260 = hirtum.
balsamiferum, *Webb & Berth. Phyt. Canar.* i. 192.—Ins. Canar.
barbatum, Hornem. Suppl. Hort. Hafn. 61 = caespitosum.
barbatum, *C. Sm. Hort. Berol.* 37; *ex Link, Enum. Hort. Berol.* ii. 20.—Ins. Canar.
barbulatum, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 91 = arachnoideum.
beugesiacum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 30 = tectorum.
bifurcum, *W. Baxt. ex Loud. Hort. Brit. Suppl.* ii. 673, nomen.—Ins. Canar.
blandum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 29.—Transsylv.
Boissieri, *Hort. ex Baker, in Gard. Chron.* (1879) ii. 39.—Hab.?
Boutignyanum, Billot & Gren. in F. Schultz, Arch. Fl. Fr. et Allem. 263, 302 = arvernense.
brachiatum, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 267.—Europ.
brachycaulon, *Webb & Berth. Phyt. Canar.* i. 216 (Index) (Quid?).—Ins. Canar.
bracteatum, *Viv. ex Steud. Nom.* ed. II. ii. 556, nomen.—Afr. bor.
Brassaii, Schur, Enum. Pl. Transs. 229 = patens.
Braunii, *Funk, ex Koch, in Sturm, Deutschl. Fl.* Heft 67.—Europ. austr.
Braunii, Ledeb. Fl. Ross. ii. 190 = globiferum.
breviramum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 36 = tectorum.
brevistylum, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 264.—Europ.
bulbosum, Soland. ex Webb & Berth. Phyt. Canar. i. 202 = Monanthes brachycaulon.
caespitosum, *C. Sm. Hort. Berol.* 38; *ex Link, Enum. Hort. Berol.* ii. 20.—Ins. Canar.
calcaratum, *Hort. ex Baker, in Gard. Chron.* (1874) ii. 386.—Hab.?
calcareum, *Jord. Obs. Pl. Crit.* vii. (1849) 26.—Europ. austr.
californicum, Hort. ex Baker, in Gard. Chron. (1874) ii. 103 = calcareum.
calyciforme, Haw. Suppl. Pl. Succ. 69 = aureum.
campaniforme, Schur, Enum. Pl. Transs. 229 = globiferum.
canariense, *Linn. Sp. Pl.* 464.—Ins. Canar.
cantalicum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 41 = tectorum.
cappadocicum, Boiss. Fl. Orient. ii. 797 = globiferum.
caucasicum, *Rupr. ex Boiss. l. c.* 796.—Reg. Cauc.
cebennense, Lamotte, Prod. Fl. Centr. Fr. i. 313 = arvernense.
celsicaule, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 33 = tectorum.
Chavini, *Lagg. in Bull. Soc. Murith.* ii. (1873) 32.—Helvet.

SEMPERVIVUM:—
chrysanthum, *Hochst. ex Britten, in Oliver, Fl. Trop. Afr.* ii. 400.—Abyss.
ciliare, Haw. Rev. Pl. Succ. 64 = caespitosum.
ciliatum, Gilib. Fl. Lituan. ii. 183 = tectorum.
ciliatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 27 = globiferum.
ciliatum, Sims, Bot. Mag. t. 1978 = caespitosum.
ciliatum, *Willd. Enum. Hort. Berol.* 508.—Ins. Canar.
claviculatum, *Sieber, ex DC. Prod.* iii. 414, nomen.—Hab.?
Clusianum, Tenore, Fl. Nap. iv. 268 = montanum.
collinum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 39 = tectorum.
columnare, Jord. & Fourr. l. c. 45 = tectorum.
Comollii, Rota, Fl. Berg. 100 = glaucum.
compactum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 289 = tectorum.
complanatum, *A. DC. in Lem. Jard. Fleur.* i. (1851) *Misc.* 35.—Ins. Canar.
constrictum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 43 = tectorum.
corymbosum, Jord. & Fourr. l. c. 44 = tectorum.
crassifolium, Salisb. Prod. 308 = tectorum.
cruentum, *Webb & Berth. Phyt. Canar.* i. 186.—Ins. Canar.
cuneatum, *Webb & Berth. l. c.* 197.—Ins. Canar.
cuspidatum, Haw. Misc. Nat. 186 = Umbilicus spinosus.
debile, *Schott, in Oestr. Bot. Wochenbl.* ii. (1852) 18.—Europ. occ.
decoloratum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 30 = tectorum.
Delasoieii, *C. B. Lehm. & Schnittsp. in Ber. Offenb. Ver. Naturk.* i. (1860) 33.—Hab.?
densum, *C. B. Lehm. & Schnittsp. l. c.* 56.—Hab.?
dichotomum, DC. Pl. Rar. Genève, 78 t. 21 = annuum.
dicranocladon, Jord. & Fourr. Pl. Brev. Nov. fasc. ii. 42 = tectorum.
divaricatum, Lowe, Man. Fl. Mad. i. 326 = punctatum.
dodrantale, *Willd. Enum. Hort. Berol.* 508.—Ins. Canar.
Doellianum, C. B. Lehm. in Flora, xxxiii. (1850) 449 = arachnoideum.
dolomiticum, Facc. in Zeitschr. Ferdinand. Tirol. III. iv. (1854) 56 = Funckii.
dumosum, *Lowe, Man. Fl. Mad.* i. 328.—Ins. Mader.
elegans, *Lagg. in Bull. Soc. Murith.* ii. (1873) 50.—Helvet.
erubescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 42 = arvernense.
Fauconnetti, Reut. Cat. Pl. Vasc. Genèv. ed. II. 298 = arachnoideum.
fimbriatum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 144. t. 43 = sedoides.
fimbriatum, *Schnittsp. & C. B. Lehm. in Flora*, xxxviii. (1855) 17.—Tyrol.
flagelliforme, *Fisch. ex Link, Enum. Hort. Berol.* ii. 20.—Sibir.
foliosum, C. Sm. Hort. Berol. 38; ex Link, Enum. Hort. Berol. ii. 19 = Smithii.
Fontanae ×, Bruegg. in Cat. Hort. Turic. xii. (1864).—Europ.
frigidum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 310 = montanum.
frutescens, *Haw. in Phil. Mag.* (1827) 125.—Ins. Canar.
Funckii, *F. Braun, ex Koch, in Flora*, xv. (1832) 4. t. 1.—Europ. austr.
Funkii, Lej. ex Nym. Consp. 258 = Schnittspahni.
glandulosum, *Ait. Hort. Kew.* ed. I. ii. 148.—Ins. Mader.
glaucum, *Tenore, Ind. Sem. Hort. Neap.* (1830) 13.—Tyrol.
globiferum, *Linn. Sp. Pl.* 464.—Europ. or.; As. Min.
globiferum, Wulf. in Jacq. Fl. Austr. v. 50. t. 40 = Wulfeni.
glutinosum, *Ait. Hort. Kew.* ed. I. ii. 147.—Ins. Mader.
Goochiae, *Webb & Berth. Phyt. Canar.* i. 190.—Ins. Canar.
grandiflorum, Haw. Rev. Pl. Succ. 66 = globiferum.
Greenii, *Baker, in Gard. Chron.* (1877) II. 230.—Hab.?
Guillemotii, *Lamotte, in Bull. Soc. Bot. Fr.* iii. (1856) 457.—Gallia.
Hausmanni, Auersd. ex Nym. Consp. 259 = arachnoideum.

SEMPERVIVUM:—
Haworthii, *Hort. Angl. ex Salm-Dyck, Hort. Dyck.* 253, nomen; *et ex Webb & Berth. Phyt. Canar.* i. 193.—Ins. Canar.
Heerianum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 97.—Europ.
heterotrichum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 83.—Europ. or.
Heuffelii, Schott, l. c. ii. (1852) 18 = patens.
Hillebrandtii, Schott, l. c. = patens.
himalayense, Klotzsch, in Bot. Ergeb. Waldem. Reise, 145. t. 43 = acuminatum.
hirtellum, Schott, in Verh. Siebenb. Ver. Naturw. (1857) 171 = arenarium.
hirtum, Jacq. Fl. Austr. i. 10. t. 12 = globiferum.
hirtum, *Linn. Cent. Pl.* i. 12; *Amoen. Acad.* iv. 273.—Europ. austr.
hirtum, Sibth. & Sm. Fl. Graec. Prod. i. 334 = patens.
hispanicum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 136 = Sedum amplexicaule.
hispanicum, *Willd. Enum. Hort. Berol.* 508.—Hispan.
holochrysum, *Webb & Berth. Phyt. Canar.* i. 194.—Ins. Canar.
hybridum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 93.—Europ.
hybridum, Sweet, Hort. Brit. ed. I. 180 = caespitosum.
juratense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 28 = montanum.
Kochii, Facc. in Zeitschr. Ferdinand. Tirol. III. iv. (1854) 57 = arenarium.
kopaonikense, *Panč. Fl. Princ. Serbiae*, (1874) 314; *ex ej. Elem. Fl. Bulg.* 30.—Serbia.
laetevirens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 37 = tectorum.
Laggeri, Schnittsp. ex Nym. Consp. 259 = arachnoideum.
Lamottei, Bor. in Mém. Soc. Acad. Maine-et-Loire, (1859) 86 = tectorum.
latifolium, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 208.—Ins. Canar.
latifolium, Salisb. Prod. 308 = canariense.
lautareticum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 296 = arachnoideum.
laxum, *Haw. Rev. Pl. Succ.* 65.—Ins. Canar.
Legrandi, F. Schultz, in Flora, l. (1867) 469 = arvernense.
leptopetalum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 43 = tectorum.
lesurinum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 282 = arvernense.
leucanthum, *Panč. Elem. Fl. Bulg.* 30.—Europ.
Lindleyi, *Webb & Berth. Phyt. Canar.* i. 189.—Ins. Canar.
lineolare, Haw. Suppl. Pl. Succ. 69 = barbatum.
longifolium, *Schnittsp. ex Lagg. in Flora*, xli. (1858) 659, nomen.—Hab.?
Lowei, Paiva, in Journ. Bot. v. (1867) 21 = Monanthes brachycaulon.
luxurians, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 34 = tectorum.
macranthum, Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 559 = arachnoideum.
Maitrea, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 273 = tectorum.
majus, Neck. Delic. Gallo-Belg. i. 213 = tectorum.
marmoreum, Griseb. Spicil. Fl. Rumel. i. 329 = tectorum.
Masferreri, *Hillebr. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 137.—Ins. Canar.
Mettenianum, Schnittsp. & C. B. Lehm. in Flora, xxxviii. (1855) 4 = tectorum.
micranthes, *Webb, ex Steud. Nom.* ed. II. ii. 556 (sp. dub.).—Ins. Canar.
minimum, Timb. in Bull. Soc. Bot. Fr. xi. (1864) 136, 138 = arvernense.
minutum, Bourg. ex Nym. Consp. 258 = tectorum.
modestum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 38 = tectorum.
Moggridgei, *De Smet, ex Hook. f. Bot. Mag.* t. 6610.—Europ. austr.
molle, *Vis. Sem. Hort. Patav.* (1841) 3; *ex Walp. Rep.* ii. 264.—Afr. trop.
Monanthes, Ait. Hort. Kew. ed. I. ii. 149 = Monanthes polyphylla.
montanum, *Linn. Sp. Pl.* 465.—Europ. med.
montanum, C. A. Mey. Verz. Pfl. Cauc. 152 = caucasicum.
montanum, Sibth. & Sm. Fl. Graec. Prod. i. 335 = tectorum.

SEMPERVIVUM:—
monticolum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 37 = tectorum.
monticolum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 306, partim = montanum.
mucronatum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 49.—Reg. Himal.
multiflorum, Jacquem. Voy. Bot. 61 = Crassula indica.
Murithii, *Lagg. in Bull. Soc. Murith.* ii. (1873) 49.—Helvet.
mutabile, *W. Schlecht. ex Breit, Hort. Breit.* 472.—Ins. Creta.
Neilreichii, Schott, Nym. & Kotschy, Analect. Bot. 19 = arenarium.
obovatum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 32 = tectorum.
oligotrichium, *Baker, in Gard. Chron.* (1879) II. 166.—Tyrol.
Paivae, *Lowe, in Hook. Bot. Mag.* t. 5593.—Ins. Canar.
pallescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 40 = tectorum.
pallidum, Jord. & Fourr. l. c. 33 = tectorum.
paniculatum, Soland. ex Lowe, Man. Fl. Mad. i. 337 = arboreum.
parviflorum, *C. B. Lehm. & Schnittsp. in Ber. Offenb. Ver. Naturk.* v. (1864) 55.—Hab.?
parvulum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 38 = tectorum.
patens, *Griseb. & Schenk, in Wiegm. Archiv*, xviii. (1852) I. 315.—Banat.
patina, Lowe, Man. Fl. Mad. i. 322 = glandulosum.
pauciflorum, Hort. ex Baker, in Gard. Chron. (1879) II. 107 = pumilum?
penninum, *Lagg. ex La Soie, in Bull. Soc. Murith.* iii. (1875) 15.—Helvet.
piliferum, Jord. Obs. Pl. Crit. vii. (1849) 27 = arachnoideum.
Pittonii, *Schott, Nym. & Kotschy, Analect. Bot.* 19.—Styria.
Pomelii, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 303.—Gallia.
praestabile, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 28 = tectorum.
pruinatum, Spreng. Syst. ii. 169 = Sedum pruinatum.
Pseudo-arachnoideum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 294 = arachnoideum.
pulchellum, *Walp. Rep.* ii. 935.—Hab.?
pulverulentum, Dulac, Fl. Hautes-Pyr. 325 = arvernense.
pumilum, *Bieb. Fl. Taur. Cauc.* i. 381.—Reg. Caucas.
punctatum, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 155.—Ins. Canar.
purpurascens, Schott, ex Boiss. Fl. Orient. ii. 796 = tectorum.
pygmaeum, Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 559 = arachnoideum.
pygmaeum, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 155.—Ins. Canar.
pyrenaicum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 44 = tectorum.
pyrenaicum, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 284.—Europ.
racemosum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 44 = tectorum.
Reginae-Amaliae, Heldr. & Sart. ex Boiss. Fl. Orient. ii. 796 = tectorum.
Requieni, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 572 = tectorum.
retusum, *Haw. in Phil. Mag.* (1827) 125.—Ins. Canar.
rhaeticum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 95.—Europ.
rhodanicum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 31 = tectorum.
rigidum, Jord. & Fourr. l. c. 35 = tectorum.
robustum, Jord. & Fourr. l. c. 32 = tectorum.
roseum, Hut. ex Nym. Consp. 259 = fimbriatum.
rubellum, *Timb. in Bull. Soc. Bot. Fr.* v. (1858) 14.—Europ. austr.
rubicundum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 67 = blandum.
rupicolum, *Kern. in Zeitschr. Ferdinand. Tirol.* III. xv. (1870) 270.—Europ.
ruthenicum, *Koch, Syn. Fl. Germ.* ed. II. 289; *Schnittsp. & C. B. Lehm. in Flora*, xxxviii. (1855) 5.—Europ. or.

SEMPERVIVUM :—
sabaudum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 29 = tectorum.
sanguineum, *Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* iii. (1875–76) 558 = arachnoideum.
saxosum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 36 = tectorum.
Schlehani, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 12 = glaucum.
Schnittspahni, *Lagg. in Flora*, xli. (1858) 659.—Europ.
Schottii, Baker, in Gard. Chron. (1874) II. 103 = glaucum.
Schottii, *C. B. Lehm. & Schnittsp. in Ber. Offenb. Ver. Naturk.* i. (1860) 35.—Hab.?
sediforme, Jacq. Hort. Vindob. i. 35. t. 81 = Sedum nicaeënse.
sedoides, J. F. Gmel. Syst. 769 = praec.
sedoides, *Jacquem. Voy. Bot.* 63. t. 74. f. 2.—Reg. Himal.
Seguieri, *DC. Prod.* iii. 414.—Ital.
seusanum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 35 = tectorum.
simense, Hochst. ex Britten, in Oliver, Fl. Trop. Afr. ii. 399 = Cotyledon simensis.
Simsii, Sweet, Hort. Suburb. Lond. 230 = caespitosum.
Smithii, *Sims, Bot. Mag.* t. 1980.—Ins. Canar.
soboliferum, Fleisch. & Lindem. Fl. Ostseeprov. 171 = hirtum.
soboliferum, Sims, Bot. Mag. t. 1457 = globiferum.
spathulatum, Hornem. Suppl. Hort. Hafn. 60 = barbatum.
speciosum, *Lamotte, in Mém. Acad. Sc. Clermont*, vi. (1864) 265.—Europ.
sphaeroideum, *Hort.—Cf. Gard. Chron.* (1876) I. 603.—Ins. Canar.
spinulosum, *Lagg. ex La Soie, in Bull. Soc. Murith.* iii. (1875) 17, nomen.—Helvet.
stellatum, Pollini, Fl. Veron. ii. 114 = Seguieri.
stellatum, *Sm. in Trans. Linn. Soc.* i. (1791) 251.—Ins. Mader.
stenopetalum, *Schnittsp. & C. B. Lehm. in Flora*, xxxviii. (1855) 18.—Hab.?
stramineum, *Baker, in Gard. Chron.* (1874) II. 104.—Hungar.
stramineum, Nym. Consp. 260 = patens.
strepsicladum, *Webb & Berth. Phyt. Canar.* i. 187.—Ins. Canar.
subvillosum, *Lowe, Man. Fl. Mad.* i. 331.—Ins. Mader.
tabulaeforme, *Haw. Suppl. Pl. Succ.* 69.—Ins. Mader.
tectorum, Eichw. Beschr. Casp.-Cauc. 31 = Braunii.
tectorum, *Linn. Sp. Pl.* 464.—Europ.; Oriens.
tenuifolium, Sibth. & Sm. Fl. Graec. Prod. i. 335 = Sedum amplexicaule.
Theobaldi ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 97.—Europ.
Thomasii, *Lagg. ex La Soie, in Bull. Soc. Murith.* iii. (1875) 17.—Helvet.
Tissieri, *Lagg. l. c.* ii. (1873) 32.—Helvet.
tomentosum, C. B. Lehm. & Schnittsp. in Flora, xxxix. (1856) 57 = arachnoideum.
tortuosum, *Ait. Hort. Kew.* ed. I. ii. 148.—Ins. Canar.
transylvanicum, Baker, in Gard. Chron. (1874) II. 104 = globiferum.
trifurcum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 41 = tectorum.
triste, *Hort. ex Baker, in Gard. Chron.* (1879) II. 39.—Hab.?
undulatum, *Webb & Berth. Phyt. Canar.* i. 197.—Ins. Canar.
urbicum, Lindl. Bot. Reg. t. 1741 = holochrysum.
urbicum, *C. Sm. ex Hornem. Suppl. Hort. Hafn.* 60.—Ins. Canar.
uviferum, *Webb, ex Steud. Nom.* ed. II. ii. 556, nomen.—Ins. Canar.
valesiaceum, *Lagg. ex La Soie, in Bull. Soc. Murith.* iii. (1875) 15.—Helvet.
validum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 31 = tectorum.
vellavum, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 280 = arvernense.
venustum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 29 = tectorum.
Verlotı, Lamotte, in Mém. Acad. Sc. Clermont, vi. (1864) 275 = Pomelii.

SEMPERVIVUM :—
vermiculare, Gueldenst. Reisen, i. 102; ex Ledeb. Fl. Ross. ii. 190 (Quid?).—Transcauc.
villosum, *Ait. Hort. Kew.* ed. I. ii. 148.—Ins. Canar.
villosum, Lindl. Bot. Reg. t. 1553 = Lindleyi.
violascens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 34 = tectorum.
Webbianum, Hort. ex C. B. Lehm. & Schnittsp. in Flora, xxxix. (1856) 57 = arachnoideum.
Widderi, *C. B. Lehm. & Schnittsp. in Ber. Offenb. Ver. Naturk.* i. (1860) 36.—Hab.?
Wulfeni, *Hoppe, ex Mert. & Koch, Deutschl. Fl.* iii. 386.—Austria.
Youngianum, *Webb & Berth. Phyt. Canar.* i. 197.—Ins. Canar.
Zelebori, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 245.—Serbia.

SENACIA, Comm. ex DC. Prod. i. 347 (1824) = **Pittosporum**, Banks.
nepalensis, DC. l. c. = P. floribundum.

SENACIA, Lam. Illustr. ii. 95 (1793). 'Genus delendum.'—Cf. Benth. & Hook. f. i. 360.
elliptica, Lam. l. c. 96 = Celastrus dubius.
glauca, Lam. l. c. 97 = Elaeodendrum glaucum.
lanceolata, Lam. l. c. 96 = Celastrus lanceolatus.
Maytenus, Lam. l. c. = Maytenus Boaria.
octogona, Lam. l. c. = Maytenus octogona.
undulata, Lam. l. c. = Pittosporum Senacia.

SENAPEA, Aubl. Pl. Gui. Suppl. 22. f. 381 (1775). Genus ignotum.
guianensis, Aubl. l. c. 22.—Guian.

SENCKENBERGIA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 413 (1800) = **Lepidium**, Linn. (Crucif.).
ruderalis, Gaertn. Mey. & Scherb. l. c. 433 = L. ruderale.

SENEBIERA, DC. in Mém. Soc. Hist. Nat. Par. (1799) 140. tt. 8, 9. *CRUCIFERAE*, Benth. & Hook. f. i. 967.
CARARA, Medic. Pflanzengatt. i. 34 (1792).
CORONOPUS, Rupp. ex Linn. Syst. ed. I (1735), nomen prius; Gaertn. Fruct. ii. 293 (1791).
COTYLISCUS, Desv. Journ. Bot. iii. (1813) 164, 175.
EUDISTEMON, Rafin. Med. Fl. ii. 219 (1830).
NASTURTIOLUM, Medic., Pflanzengatt. i. 82 (1792).
australis, *Hook. f. Fl. Antarct.* 241.—Patagon.
coronopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Coronopus.
Coronopus, *Poir. Encyc.* vii. 76.—Cosmop.
didyma, Pers. Syn. ii. 185 = pinnatifida.
diffusa, *Bert. ex Steud. Nom.* ed. II. ii. 557, nomen.—Chili.
dubia, H. B. & K. Nov. Gen. et Sp. v. 76 = Lepidium Humboldtii.
Figariana, *Parl. Giorn. Bot. Ital.* i. 1. (1844) 308.—Aegypt.?
Heleniana, DC. Syst. ii. 523 = pinnatifida?
incisa, Willd. Enum. Hort. Berol. 668 = pinnatifida.
integrifolia, *DC. in Mém. Soc. Hist. Nat. Par.* (1799) 144. t. 8.—China; Afric.; Austral.
lepidioides, *Coss. in Bull. Soc. Bot. Fr.* ii. (1855) 245.—Alger.
linoides, DC. Syst. ii. 522 = integrifolia.
mexicana, Hook. & Arn. Bot. Beech. Voy. 276 = Lepidium virginicum.
nilotica, *DC. Syst.* ii. 527.—Aegypt.
pectinata, DC. l. c. 523 = pinnatifida.
pinnatifida, *DC. in Mém. Soc. Hist. Nat. Par.* (1799) 144 t. 9.—Cosmop.
Raddii, *Walp. Rep.* ii. 764.—Aegypt.
rhytidocarpa, *Hook. Lond. Journ. Bot.* ii. (1843) 506.—Patagon.
serrata, *Poir. Encyc.* vii. 76.—Reg. Argent.
squamata, C. Muell. in Walp. Ann. vii. 156 = S. Coronopus.
supina, *Thore, Ess. Chl. Land.* 275.—Gallia.
violacea, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 282.—Alger.

SENECIASTRUM, Heist. Syst. 7 (1748) (Quid?).

SENECILLIS, Gaertn. Fruct. ii. 453, t. 173 (1791) = **Senecio**, Tourn. (Compos.).
carpathica, Schott, Nym. & Kotschy, Analect. Bot. 5 = *glauca*, Schott.
filiformis, Spreng. ex Sch. Bip. in Flora, xxvii. (1844) 702 = Euryops trifurcatus.
glauca, Gaertn. Fruct. ii. 453.—Sibir.
glauca, Ledeb. Fl. Alt. iv. 113 = Senecio altaicus.
glauca, Schott, Nym. & Kotschy, Analect. Bot. 5 = Transsylv.
heterophylla, Schur, Enum. Pl. Transs. 342 = *glauca*, Schott.
Jacquemontiana, Decne. in Jacquem. Voy. Bot. 90. t. 98 = Senecio Jacquemontianus.
purpurata, Gaertn. Fruct. ii. 453 = Cineraria purpurata.
Schmidtii, Maxim. in Bull. Acad. Pétersb. xvi. (1871) 222 = Senecio Schmidtii.

SENECIO, [Tourn.] Linn. Syst. ed. I. 251 (1735); Gen. ed. I. 251 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 446.
ACLEIA, DC. Prod. vi. 340 (1837).
ADENOTRICHIA, Lindl. Bot. Reg. t. 1190 (1828).
ADONIGERON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404.
AETHEOLAENA, Cass. in Dict. Sc. Nat. xlviii. 450 (1827).
ANECIO, Neck. Elem. i. 28 (1790).
ASPELINA, Cass. in Dict. Sc. Nat. xli. 166 (1826).
BETHENCOURTIA, Choisy, in Buch, Beschr. Canar. Ins. 148 (1825).
BRACHYPAPPUS, Sch. Bip. in Flora, xxxviii. (1855) 119.
BRACHYRHYNCHOS, Less. Syn. Comp. 392 (1832).
BRANICIA, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 476.
CACALIA, Linn. Sp. Pl. ed. I. 834 (1753).
CARDERINA, Cass. in Dict. Sc. Nat. xxxv. 272 (1825).
CENTROPAPPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 124.
CINERARIA, Linn. Sp. Pl. ed. II. 1242 (1762), partim.
CISSAMPELOPSIS, Miq. Fl. Ind. Bat. ii. 102 (1856).
CLADOPOGON, Sch. Bip. in Ind. Sem. Hort. Hamb. (1852) 8; ex Lehm. in Ann. Sc. Nat. Sér. III. xix. (1853) 359.
DANAA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 27. t. 28.
DELAIRIA, Lem. in Ann. Sc. Nat. Sér. III. i. (1844) 379.
DORIA, Thunb. Nov. Gen. xii. 162 (1800).
DOROBAEA, Cass. in Dict. Sc. Nat. xlviii. 453 (1827).
ERYTHROCHAETE, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 188 (Fam. Nat. Fl. Jap. ii. 64).
EUDORUS, Cass. in Bull. Soc. Philom. (1818) 165; et in Dict. Sc. Nat. xv. 525 (1819).
FARFUGIUM, Lindl. in Gard. Chron. (1857) 4.
FAROBAEA, Schrank, ex Colla, Hort. Ripul. App. iii. 36, in nota (1829); et iv. 19. t. 9 (1831).
HAPLOSTICHA, Phil. in Linnaea, xxx. (1859–60) 193.
HERBICHIA, Zawadski, in Guill. Arch. Bot. i. (1832) 450; et Enum. Pl. Galic. 198 (1835).
HOPPEA, Reichb. in Flora, vii. (1824) 245.
HUBERTIA, Bory, Voy. i. 334. t. 14 (1804).
JACOBAEA, Thunb. Fl. Cap. Prod. Praef. (1794).
JACOBANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 403.
KLEINIA, Haw. Syn. Pl. Succ. 312 (1812).
LACHANODES, DC. in Guill. Arch. Bot. ii. (1833) 332.
LEUCOSERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404.
LIGULARIA, Cass. in Bull. Soc. Philom. (1816) 198.
MADACARPUS, Wight, Ic. t. 1152 (1846).
MADARACTIS, DC. Prod. vi. 322 (1837).
MESADENIA, Rafin. in Loud. Gard. Mag. viii. (1832) 247.
MESOGRAMMA, DC. Prod. vi. 304 (1837).
MESONEURIS, A. Gray, in Proc. Am. Acad. viii. (1873) 661.
METACANTHUS, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 280.
METAZANTHUS, Meyen, Reise, i. 356 (1834).
MICROCHAETE, Benth. Pl. Hartw. 209 (1845).

SENECIO :—
Moerkensteinia, Opiz, Seznam, 66 (1852).
Obaejaca, Cass. in Dict. Sc. Nat. xxxv. 270 (1825).
Odontotrichum, Zucc. in Abh. Akad. Muench. i. (1832) 311.
Pentacalia, Cass. in Dict. Sc. Nat. xlviii. 461 (1827).
Pentanthus, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 32.
Pericalia, Cass. in Dict. Sc. Nat. xlviii. 459 (1827).
Pericallis, Webb & Berth. Phyt. Canar. ii. tt. 103, 106 (1835–60).
Pithosillum, Cass. in Dict. Sc. Nat. xli. 164 (1826).
Pladaroxylon, Hook. f. in Hook. Ic. Pl. t. 1055 (1870).
Psacalium, Cass. in Dict. Sc. Nat. xliii. 461 (1820).
Roldana, La Llave & Lex. Nov. Veg. Desc. fasc. ii. 10 (1825).
Rugelia, Shuttlew. in Chapm. Fl. S. U. St. 246 (1860).
Sciadioseris, Kunze, in Bot. Zeit. ix. (1851) 349.
Sclerobasis, Cass. in Bull. Soc. Philom. (1818) 73.
Senecillis, Gaertn. Fruct. ii. 453. t. 173 (1791).
Senecіuncuius, Opiz, Seznam, 90 (1852).
Synarthrum, Cass. in Dict. Sc. Nat. li. 457 (1827).
Syneilesis, Maxim. Prim. Fl. Amur. 165 (1859).
Synosma, Rafin. in Loud. Gard. Mag. viii. (1832) 247.
Tephroseris, Schur, Enum. Pl. Transs. 343 (1866).
Traversia, Hook. f. Handb. N. Zeal. Fl. 163 (1864).
Willkommia, Sch. Bip. ex Nym. Consp. 357 (1879).
abadianus, *DC. Prod.* vi. 423.—Peruv.
abietinus, *Willd. ex Wedd. Chlor. And.* i. 100, 101.—N. Granat.
abrotanifolius, Georgi, Beschr. Russ. Reich. III. iv. 1242 = S. Jacobaea.
abrotanifolius, Gouan, Hort. Monsp. 440 = adonidifolius.
abrotanifolius, *Linn. Sp. Pl.* 869.—Europ.
abrotanoides, E. Mey. ex DC. Prod. vi. 397 = pinnatifidus.
abruptus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
abyssinicus, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) I. *Intell.* 27, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 438.—Abyss.
acanthifolius, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 46.—Patagon.
acanthifolius, Kostel. Allg. Med.-Pharm. Fl. ii. 714 = S. Cineraria.
acaulis, *Phil. in Anal. Univ. Chil.* (1861) 52.—Chili.
acaulis, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
acerifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 235.—Mexic.
achilleaefolius, *DC. Prod.* vi. 396.—Afr. austr.
aconitifolius, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. vii. 155.—China.
Actinella, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 87.—Arizona.
actinoleucus, *F. Muell. in Trans. Roy. Soc. Victoria*, viii. (1867) 40.—Afr. austr.
acuminatus, *Wall. Cat.* n. 3107.—Reg. Himal.
acutangulus, *Hemsl. Biol. Centr. Am. Bot.* ii. 235.—Guatemala.
acutidentatus, A. Rich. Tent. Fl. Abyss. i. 436 = picridifolius.
acutifolius, *DC. Prod.* vi. 402.—Afr. austr.
adamantinus, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 97.—Bras.
adenodontus, *DC. Prod.* vi. 378.—Madag.
adenophylloides, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
adenophyllus, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 282.—Peruv.
adenostyloides, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 486.—Japon.
adenotrichius, *DC. Prod.* vi. 416.—Chili.
adfinis, Less. ex Harv. & Sond. Fl. Cap. iii. 363 = erubescens, *Ait.*
adnatus, *DC. Prod.* vi. 388.—Afr. austr.

SENECIO :—
adonidifolius, *Loisel. Fl. Gall.* ed. I. 566; ed. II. ii. 239.—Europ.
adscendens, *Boj. ex DC. Prod.* vi. 378.—Madag.
adscendens, DC. l. c. 423 = andinus.
aegyptius, *Linn. Sp. Pl.* 867.—Aegypt.
aethnensis, *Jan, ex DC. Prod.* vi. 345.—Ital.
affinis, DC. l. c. 385 = albanensis.
africanus, Hort. ex Poir. Encyc. Suppl. v. 129 = Gynura haematophylla.
agapatensis, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
Aitonianus, G. Don, in Sweet, Hort. Brit. ed. III. 386 = Cineraria canescens.
aizoides, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
akrabatensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 312.—Turkestan.
alatus, *Wall. Cat.* n. 3114.—Reg. Himal.
alatus, Willd. ex Ledeb. Fl. Ross. ii. 633 = lampsanoides.
albanensis, *DC. Prod.* vi. 385.—Afr. austr.
albens, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
albicans, Pers. Syn. ii. 438 = lanatus.
albicaulis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 344.—Patagon.
albicaulis, Sch. Bip. in Flora, xxvii. (1844) 702 = polymorphus.
albidus, E. Mey. ex DC. Prod. vi. 398 = grandiflorus.
albiflorus, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Chili.
albifolius, *DC. Prod.* vi. 389.—Afr. austr.
albo-dentatus, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 144 = S. Cacaliaster.
albolanatus, *Phil. Fl. Atac.* 32.—Chili.
albo-lutescens, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
alcicornis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 334.—Chili.
Alciope, Sch. Bip. in Flora, xxviii. (1845) 500 = Alciope lanata.
algens, *Wedd. Chlor. And.* i. 104.—Bolivia.
Almeidae, *Phil. Fl. Atac.* 33.—Chili.
aloides, *DC. Prod.* vi. 402.—Afr. austr.
alpester, Kern. in Oestr. Bot. Zeitschr. xxi. (1871) 202 = seq.
alpestris, *DC. Prod.* vi. 360.—Europ.
alpestris, Gaud. Fl. Helv. v. 296 = Fuchsii.
alpinus, C. B. Clarke, Comp. Ind. 190 = bracteolatus, *Hook. f.*
alpinus, Linn. f. Suppl. 371 = lyratifolius.
alpinus, *Scop. Fl. Carn.* ed. II. ii. 164.—Europ.
alsophilus, *Phil. in Linnaea*, xxix. (1857–58) 2.—Chili.
altaicus, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Sibir. altaic.
altissimus, Mill. Gard. Dict. ed. VIII. n. 9 = S. Doria.
amabilis, *DC. Prod.* vi. 406.—Afr. austr.
amabilis, Vell. Fl. Flum. viii. t. 107 = brasiliensis.
Ambavilla, *Pers. Syn.* ii. 436.—Ins. Borbon.
ambiguus, *DC. Prod.* vi. 356.—Ital.
ambraceus, *Turcz. ex DC. l. c.* 348.—China.
ambrosioides, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 322 = brasiliensis.
americanus, *DC. Prod.* vi. 310.—N. Granat.
ammophilus, Greene, in Bull. Calif. Acad. i. iv. (1885) 193 = californicus.
ammophilus, *Phil. in Linnaea*, xxviii. (1856) 748.—Chili.
amoenus, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 443 = Werneria africana.
amphibolus, *C. Koch, in Linnaea*, xxiv. (1851) 361.—Reg. Caucas.
amphibolus, *Wedd. Chlor. And.* i. 109.—Bolivia.
amplectens, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 240.—Am. bor.
amplexicaulis, *H. B. & K. Nov. Gen. et Sp.* iv. 181.—Ecuador.
amplexicaulis, *Wall. Cat.* n. 3139.—Reg. Himal.
amplifolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
amplus, *Remy, in C. Gay, Fl. Chil.* iv. 195.—Chili.
ampullaceus, *Hook. Bot. Mag.* t. 3487.—Texas.
amygdalifolius, *F. Muell. Fragm.* i. 232.—Austral.
Anacampserotis, *DC. Prod.* vi. 374.—Austral.
anacephalus, *Griseb. in Goett. Abh.* xxiv. (1879) 208.—Reg. Argent.

SENECIO :—
analogus, DC. Prod. vi. 366 = chrysanthemoides.
Anampoza, *Baker, in Journ. Linn. Soc.* xx. (1883) 191.—Madag.
Andersoni, *C. B. Clarke, Comp. Ind.* 204.—Burma.
Andersoni, *Hook. f. Fl. Antarct.* 312.—Patagon.
andicola, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 91.—Ecuador.
andinus, *Buek, Ind. ad DC. Prod.* ii. p. vi.—Peruv.
andinus, Nutt. in Trans. Am. Phil. Soc. vii. (1841) 409 = S. Serra, *Hook.*
Andrieuxii, *DC. Prod.* vi. 430.—Mexic.
andryaloides, DC. l. c. 420 = crassifolius.
anethifolius, *A. Cunn. in DC. l. c.* 371.—Austral.
angavonensis, *Boj. ex DC. l. c.* 377.—Madag.
angulatus, *Linn. f. Suppl.* 369.—Afr. austr.
angulifolius, *DC. Prod.* vi. 431.—Mexic.
angulosus, Wall. Cat. n. 3117 = densiflorus.
angustifolius, Forst. f. Prod. 91 = lautus.
angustifolius, Sond. in Linnaea, xxv. (1852) 526 = anethifolius.
angustifolius, Wall. ex DC. Prod. vi. 367 = saxatilis.
angustifolius, *Willd. Sp. Pl.* iii. 1973.—Afr. austr.
angustilobus, F. Muell. in Linnaea, xxv. (1852) 418 = anethifolius.
angustissimus, *Phil. l. c.* xxviii. (1856) 746.—Chili.
anomalus, *DC. Prod.* vi. 310.—Graecia.
anonymus, *Wood, Am. Bot. Flor.* 187.—Am. bor.
Antennaria, *Wedd. Chlor. And.* i. 106.—Peruv.; Bolivia.
Anteuphorbium, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
anthemidiphyllus, *Remy, in C. Gay, Fl. Chil.* iv. 177.—Chili.
anthemifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 376.—Afr. austr.
anthemifolius, Sieber, ex Steud. Nom. ed. II. ii. 558 = delphinifolius.
anthemoides, *Steud. l. c.*—Ind. or.
anthoraefolius, Presl, ex Reichb. Fl. Germ. Excurs. 243 = praealtus.
Antisanae, *Benth. Pl. Hartw.* 210.—N. Granat.
antucensis, *Phil. in Linnaea*, xxviii. (1856) 747.—Chili.
apargiaefolius, Walp. l. c. xiv. (1840) 309 = Erechtites arguta.
apenninus, Tausch, in Syll. Ratisb. ii. (1828) 252 = S. Doronicum.
apiculatus, *Sch. Bip. ex Wedd. Chlor. And.* i. 128.—Venezuela.
appendiculatus, *DC. Prod.* vi. 376.—Ins. Maurit.
appendiculatus, Poir. Encyc. vii. 102 = auriculatus, *Vahl.*
appendiculatus, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 327.—Ins. Canar.
apuanus, *Tausch, ex Reichb. Fl. Germ. Excurs.* 851.—Europ.
apulus, Tenore, ex Nym. Consp. 357 = crassifolius.
aquaticus, Boiss. Voy. Bot. Espagne, ii. 332 = praealtus.
aquaticus, *Hill, Veg. Syst.* ii. 120; *Huds. Fl. Angl.* ed. I. 317.—Europ.
aquaticus, Loisel. Fl. Gall. ii. 566; et ed. II. ii. 240; Sibth. & Sm. Fl. Graec. Prod. ii. 178 = erraticus.
aquifoliaceus, *DC. Prod.* vi. 394.—Afr. austr.
arabicus, *Linn. Mant.* i. 114.—Aegypt.; Arab.
arachnoideus, *Phil. in Linnaea*, xxviii. (1856) 739.—Chili.
arachnoideus, Sieber, ex DC. Prod. vi. 357 = lanatus, *Scop.*
arachnoideus, Wall. Cat. n. 3163 = araneosus.
arachnolomus, *Wedd. Chlor. And.* i. 104.—Peruv.
araneosus, *DC. Prod.* vi. 364.—Ind. or.; Java.
araneosus, E. Mey, ex DC. l. c. 391 = arnicaeflorus.
araucanus, *Phil. in Linnaea*, xxxiii. (1864–65) 158.—Chili.
arborescens, *Steetz, in Seem. Bot. Voy. Herald*, 162. t. 31.—Panama.
arbutifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 182.—Peruv.
arcticus, *Rupr. Beitr. Pfl. Russ. Reich.* ii. 44.—As. bor. or.
Arechavaletae, *Baker, in Mart. Fl. Bras.* vi. iii. 310.—Uruguay.
arenarius, Bieb. in Bess. Cat. Hort. Crem. (1816) 129 = praealtus.
arenarius, Lange, ex Nym. Consp. 356 = erucifolius.

SENECIO:—
arenarius, Salzm. ex DC. Prod. vi. 345 = crassifolius.
arenarius, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
argaeus, *Boiss. & Bal. Diagn.* Ser. II. vi. 100.—As. Min.
argenteus, *Kunze, ex DC. Prod.* vi. 415.—Chili.
argentinus, *Baker, in Mart. Fl. Bras.* vi. III. 307, in obs.—Reg. Argent.
argillosus, *Baker, l. c.* 310.—Uruguay.
argophylloides, *Griseb. in Goett. Abh.* xix. (1874) 189.—Reg. Argent.
argophyllus, *Phil. in Linnaea*, xxviii. (1856) 747.—Chili.
argunensis, Debeaux, Fl. Tché-Fou, 85 = ambraceus.
argunensis, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) II. 18.—China.
argutus, *H. B. & K. Nov. Gen. et Sp.* iv. 188.—Mexic.
argutus, A. Rich. Fl. N. Zél. 258 = Erechtites arguta.
Aristianus, *Remy, in C. Gay, Fl. Chil.* iv. 181.—Chili.
arizonicus, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 87.—Arizona.
Arleguianus, *Remy, in C. Gay, Fl. Chil.* iv. 190.—Chili.
armenius, *Jaub. & Spach, Illustr.* iv. 143. t. 398.—Armenia.
armenius, Ledeb. Fl. Ross. ii. 634, sphalm. = praealtus.
armoracifolius, Burch. ex Baker, in Mart. Fl. Bras. vi. III. 316 = icoglossus.
arnicaeflorus, *DC. Prod.* vi. 390.—Afr. austr.
arnicoides, Hook. & Arn. Bot. Beech. Voy. 32 = plantagineus.
arnicoides, *Wall. Cat.* n. 3138.—Reg. Himal.
Arnottianus, DC. Prod. vi. 367 = neelgherryanus.
Arnottii, *Hook. f. Fl. Antarct.* 314.—Patagon.
aromaticus, *Gilib. Exerc. Bot. in Schola Princip. Univ. Viln.* 14.—Lithuan.
aronicoides, *DC. Prod.* vi. 426.—Calif.
Arrabidae, Steud. Nom. ed. II. ii. 558 = brasiliensis.
artemisiaefolius, *Gilib. Exerc. Bot. in Schola Princip. Univ. Viln.* 14.—Lithuan.
artemisiaefolius, Pers. Syn. ii. 435 = adonidifolius.
articulatus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
asarifolius, Boj. ex DC. Prod. vi. 330 = Faujasia reticulata.
Aschenbornianus, *Schau. in Linnaea*, xx. (1847) 698.—Mexic.
asclepiadeus, *DC. Prod.* vi. 422.—Am. austr.
Aspelina, *DC. l. c.* 436.—Amer.
asper, Ait. Hort. Kew. ed. I. iii. 196 = rosmarinifolius.
aspericaulis, *Remy, in C. Gay, Fl. Chil.* iv. 174.—Chili.
asperulus, *DC. Prod.* vi. 386.—Afr. austr.
asplenifolius, *Griseb. in Goett. Abh.* xxiv. (1879) 207.—Reg. Argent.
assuayensis, *DC. Prod.* vi. 422.—Am. austr.
assuayensis, Sch. Bip. ex Wedd. Chlor. And. i. 97 = nevadensis.
Atkinsoni, C. B. Clarke, Comp. Ind. 207 = S. Ligularia.
Atkinsoniae, F. Muell. Fragm. v. 88 = Erechtites Atkinsoniae.
atlanticus, Boiss. & Reut. Pugill. Pl. Nov. (1852) 58 = vernalis.
atlanticus, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 706.—Alger.
atriplicifolius, *Hook. Fl. Bor. Am.* i. 332.—Am. bor.
attenuatus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
Aucheri, *DC. Prod.* vi. 361.—Graec.; Bithyn.
aurantiacus, Less. in Linnaea, ix. (1834) 191; DC. l. c. = campestris.
aurantiacus, Reichb. Ic. Fl. Germ. xvi. 44. t. 982 = palustris.
auratus, *DC. l. c.* 348.—Sibiria.
aureus, Georgi, Beschr. Russ. Reich. III. iv. 1243 = paludosus.
aureus, *Linn. Sp. Pl.* 870.—Am. bor.
aureus, Wall. Cat. n. 3113 = densiflorus.
auricomus, Link, ex DC. Prod. vi. 354 = caespitosus, *Brot.*
Auricula, *Bourg. ex Coss. Notes Crit.* 169.—Hispan.
auriculatus, *Burm. f. Fl. Ind.* 181.—Java.
auriculatus, Desf. Fl. Atlant. ii. 272 = auritus.
auriculatus, Poepp. ex DC. Prod. vi. 417 = glabratus.

SENECIO:—
auriculatus, Reichb. ex Steud. Nom. ed. II. ii. 558 = alpinus.
auriculatus, *Vahl, Symb. Bot.* i. 72. t. 18.—Arab.
auriculatus, Wall. Cat. n. 3115 = rufinervis.
aurinitens, Schur, Enum. Pl. Transs. 350 = S. Jacobaea.
auritus, *Willd. Sp. Pl.* iii. 1982.—Afr. bor.
Austinae, *Greene, in Bull. Calif. Acad.* i. n. III. (1885) 93.—Calif.
australis, Willd. Sp. Pl. iii. 1981 = lautus.
axillaris, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 333.—Mexic.
axillaris, *Vell. Fl. Flum.* viii. t. 112.—Bras.
ayapatensis, Baker, in Mart. Fl. Bras. vi. II. 314, in obs. sphalm. = agapatensis.
ayapatensis, Griseb. in Goett. Abh. xxiv. (1879) 206 = trichocodon.
baccharidifolius, *DC. Prod.* vi. 414.—Chili.
baccharoides, H. B. & K. Nov. Gen. et Sp. iv. 186 = Gynoxys baccharoides.
Badaroi, Moris, ex Badaro, in Brugnat. Giorn. Fis. Dec. II. vii. (1824) 367 = delphinifolius.
bahioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 336.—Chili.
Balansae, *Baker, in Mart. Fl. Bras.* vi. III. 317 = Paraguay.
Balansae, Boiss. & Reut. Diagn. Ser. II. iii. 32 = nebrodensis.
Balbisianus, *DC. Prod.* vi. 360.—Europ.
Balbisianus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 445 = campestris.
baldensis, Poir. Encyc. vii. 109 = alpinus, S. Doronicum.
balsamicus, *Phil. in Anal. Univ. Chil.* (1873) 499.—Chili.
Balsamitae, Muhl. ex Willd. Sp. Pl. iii. 1998 = aureus.
Banksii, *Hook. f. Fl. N. Zel.* i. 146.—N. Zel.
Barba-Johannis, *DC. Prod.* vi. 430.—Mexic.
barbareaefolius, Krock. Fl. Siles. ii. II. 421 = erraticus.
barbareaefolius, Reichb. Fl. Germ. Excurs. 244 = aquaticus.
barbareaefolius, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 90.—Afr. austr.
barbatus, *DC. Prod.* vi. 406.—Afr. austr.
barbatus, [*D.*] *Don, ex Hook. et Arn. in Hook. Journ. Bot.* iii. (1841) 334.—Reg. Argent.
barbellatus, *DC. Prod.* vi. 388.—Afr. austr.
barckhausiaefolius, *Boiss. & Heldr. Diagn.* Ser. II. vi. 101.—Graecia.
barkhausioides, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 86 = georgianus.
Barrelieri, Gouan, Illustr. 68 = S. Doronicum.
Barrelieri, Hoppe, ex DC. Prod. vi. 354 = nemorensis.
bathurstianus, Sch. Bip. in Flora, xxviii. (1845) 498 = Erechtites arguta.
bayonnensis, Boiss. Diagn. Ser. II. iii. 34 = nemorensis.
Bedfordii, F. Muell. Cat. Hort. Melb. (1858) 26 = Bedfordia salicina.
Beecheyanus, Sch. Bip. in Flora, xxviii. (1845) 499 = sessilifolius.
Behrianus, *Sond. & F. Muell. in Linnaea*, xxv. (1852) 527.—Austral.
belbeysius, *Delile, Fl. Egypte*, 270.—Aegypt.
belbeyticus, Poir. Encyc. Suppl. v. 229 = praec.
belenensis, *Griseb. in Goett. Abh.* xix. (1874) 192 = Reg. Argent.
belgaumensis, *C. B. Clarke, Comp. Ind.* 200.—Ind. or.
bellidifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 175.—Mexic.
bellidifolius, A. Rich. Tent. Fl. Abyss. i. 439 = abyssinicus.
bellidioides, *Hook. f. Fl. N. Zel.* i. 144.—N. Zel.
Bellis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 369.—Afr. austr.
belophyllus, *Remy, in C. Gay, Fl. Chil.* iv. 164.—Chili.
benaventianus, *Remy, l. c.* 141.—Chili.
Benthamii, *Griseb. in Goett. Abh.* xxiv. (1879) 206.—Reg. Argent.
Benthamii, Sch. Bip. ex Wedd. Chlor. And. i. 96 = Humboldtianus.
Bergianus, E. Mey, ex DC. Prod. vi. 400 = rosmarinifolius.
Bergii, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 357.—Reg. Argent.

SENECIO:—
Berlandieri, *Sch. Bip. in Flora*, xxviii. (1845) 498; *Hemsl. Biol. Centr. Am. Bot.* ii. 236.—Mexic.
Berterianus, *Colla, in Mem. Acc. Torin.* xxxviii. (1835) 32.—Chili.
betonicaefolius, *DC. Prod.* vi. 422—Am. austr.
betsiliensis, *Baker, in Journ. Bot.* xx. (1882) 170.—Madag.
Bhot, *C. B. Clarke, Comp. Ind.* 187.—Reg. Himal.
Biafrae, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 420.—Afr. trop.
bicolor, Balb. Cat. Hort. Taur. (1804) 42 = Gynura haematophylla.
bicolor, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura bicolor.
bicolor, *Todaro, Ind. Sem. Hort. Panorm.* (1859) 22, 30.—*Cf. Linnaea*, xxx. (1859–60) 759.—Ital.; Graec.
bidentatus, *Willd. Sp. Pl.* iii. 1976.—Afr. austr.
Bidwillii, *Hook. f. Fl. N. Zel.* i. 150.—N. Zel.
bifistulosus, *Hook. f. l. c.* 144.—N. Zel.
biflorus, *Burm. f. Fl. Ind.* 181.—Java.
biflorus, *Vahl, Symb. Bot.* i. 72.—Arab.
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 111.—Am. bor. occ.
Billardierii, F. Muell. Cat. Hort. Melb. (1858) 26 = Bedfordia linearis.
bipinnatifidus, Hook. & Arn. Bot. Beech. Voy. 32 = Berterianus, sinuatilobus.
bipinnatus, DC. Prod. vi. 397 = foeniculoides.
bipinnatus, *Less. Syn. Comp.* 391.—Afr. austr.
blattarioides, *DC. Prod.* vi. 393.—Afr. austr.
Blumeanus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura aurantiaca.
Blumei, DC. Prod. vi. 334, in syn. = araneosus.
bogotensis, *Spreng. Syst.* iii. 556.—N. Granat.
bohemicus, Tausch, in Syll. Ratisb. ii. (1828) 253 = paludosus.
Boissieri, *DC. Prod.* vii. 300.—Hispan.
Boivini, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 375.—Ins. Maurit.
Bojeri, *DC. Prod.* vi. 37.—Madag.
Bojeri, Hook. f. in Journ. Linn. Soc. vii. (1864) 202 = Biafrae.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 362.—Calif.
boliviensis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
Bolusii, *Oliver, in Hook. Ic. Pl.* t. 1456.—Afr. austr.
bonariensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 340.—Reg. Argent.
Bonplandianus, *DC. Prod.* vi. 421.—Ecuador.
Boryanus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura auriculata.
borysthenicus, Andrz. ex DC. Prod. vi. 351 = praealtus.
Boscianus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Carolina.
Boutoni, *Baker, Fl. Maurit.* 178.—Ins. Rodriguez.
Bowenkampi, *Phil. in Anal. Univ. Chil.* (1873) 500.—Chili.
brachyatus, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 344, in obs. = erucifolius.
brachychaetus, *DC. Prod.* vi. 362.—Europ.; As. bor.
brachycodon, *Baker, in Mart. Fl. Bras.* vi. III. 319.—Bras.
brachyglossus, *F. Muell. ex Sond. in Linnaea*, xxv. (1852) 525.—Austral.
brachyglossus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 87 = glutinosus, *Thunb.*
brachylaenus, DC. Prod. vi. 370 = Cunninghamii.
brachylepis, *Phil. in Linnaea*, xxxiii. (1864–65) 162.—Chili.
brachylobus, *Phil. l. c.* 156.—Chili.
brachypodus, *DC. Prod.* vi. 403.—Afr. austr.
brachyrhynchoides, DC. l. c. 383 = hastulatus.
bracteatus, Boiss. & Orph. Diagn. Ser. II. iii. 36 = macedonicus.
bracteolatus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 347.—Reg. Argent.
bracteolatus, *Hook. f. Fl. Brit. Ind.* iii. 339.—Reg. Himal.
bracteosus, *Remy, in C. Gay, Fl. Chil.* iv. 189.—Chili.
brasiliensis, *Less. in Linnaea*, vi. (1831) 249.—Bras.
brasiliensis, Spreng. ex Baker, in Mart. Fl. Bras. vi. III. 303 = Pohlii.

SENECIO :—

breviculus, *Phil. in Anal. Univ. Chil.* (1862) II. 398.—Chili.
breviscapus, *DC. Prod.* vi. 418.—Chili.
Bridgesii, *Hook. & Arn. Bot. Beech. Voy.* 57.—Chili.
Brunonianus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 337.—Chili.
bryoniaefolius, *Harv. in Harv. in Sond. Fl. Cap.* iii. 404.—Afr. austr.
Buchanani, Armstr. in Trans. N. Z. Inst. xiii. (1881) 339 = elaeagnifolius.
Buglossa, *Phil. in Linnaea*, xxxiii. (1864–65) 157.—Chili.
Buimalia, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 178.—Reg. Himal.
bulbiferus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 485.—China; Japon.
bulbinefolius, *DC. Prod.* vi. 402.—Afr. austr.
bulbosus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura auriculata.
bullatus, *Benth. Pl. Hartw.* 208.—Ecuador.
Bungei, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 313.—Turkestan.
bupleuriformis, Sch. Bip. in Flora, xxvii. (1844) 696 = bupleuroides.
bupleuroides, *DC. Prod.* vi. 387.—Afr. trop. et austr.
Burchellii, *DC. l. c.* 401.—Afr. austr.
Burtoni, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 202.—Afr. trop.
bustillosianus, *Remy, in C. Gay, Fl. Chil.* iv. 155.—Chili.
buxifolius, H. B. & K. Nov. Gen. et Sp. iv. 188 = Gynoxys buxifolia.
byzantinus, *Linn. Sp. Pl.* 871.—Turc.
byzantinus, Willd. Sp. Pl. iii. 2009 = paludosus.
cacaliaefolius, Sch. Bip. in Flora, xxviii. (1845) 50 = S. Ligularia.
cacaliaeformis, Reichb. f. Ic. Fl. Germ. xvi. 43. t. 977. f. 1–9 = L. Ligularia.
Cacaliaster, Blanco, Fl. Filip. ed. II. 441 = Gynura angulosa.
Cacaliaster, Guss. ex DC. Prod. vi. 354 = sarracenicus.
Cacaliaster, *Lam. Fl. Fr.* ii. 132.—Gallia.
cacalioides, Fisch. ex Spreng. Nov. Prov. 37 = Erechtites praealta.
cachinalensis, *Phil. Fl. Atac.* 33.—Chili.
caducifolius, Sch. Bip. ex Wedd. Chlor. And. i. 94 = ledifolius.
caespitosus, *Brot. Fl. Lusit.* i. 390.—Lusitan.
caespitosus, *Phil. in Linnaea*, xxviii. (1856) 741.—Chili.
cakilefolius, *DC. Prod.* vi. 408.—Afr. austr.
calamifolius, Hook. Bot. Mag. t. 4011 = scaposus.
calcarius, *H. B. & K. Nov. Gen. et Sp.* iv. 184.—Mexic.
calcicola, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 283.—Peruv.
Caledoniae, *Schlecht. ex DC. Prod.* vi. 373.—N. Caled.
californicus, *DC. l. c.* 426.—Calif.
callosus, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
calocephalus, *Hemsl. Biol. Centr. Am. Bot.* ii. 237.—Guatemala.
calocephalus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 58.—Chili.
calophyllus, Hemsl. Biol. Centr. Am. Bot. ii. 237 = albo-lutescens.
calthaefolius, *Hook. f. Fl. Brit. Ind.* iii. 350.—Reg. Himal.
calthaefolius, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 220, 221.—Manchur.
Calverti, Boiss. Diagn. Ser. II. iii. 32 = viscosus.
Calverti, Boiss. Fl. Orient. iii. 393 = mollis.
calvescens, *Moris & De Not. Fl. Ins. Caprar.* 74. t. 1.—Ital.
campanulatus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
campester, Sleńdz.—Cf. Oestr. Bot. Zeitschr. xxxiv. (1884) 32 = seq.
campestris, *DC. Prod.* vi. 361.—Europ.; As. bor.
campestris, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 295, 332 = lugens, Hookeri.
camporum, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 423 = adamantinus.
campylodes, DC. Prod. vi. 370 = scandens.
canabinaefolius, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 341 = brasiliensis.

SENECIO :—

canadensis, Linn. Sp. Pl. 869 = artemisiaefolius.
canaliculatus, *Boj. ex DC. Prod.* vi. 378.—Madag.
canalipes, *DC. l. c.* 405.—Afr. austr.
canalipes, E. Mey. in Drège, Zwei Pfl. Docum. 220 = angulatus.
cancellatus, *DC. Prod.* 435.—Ins. S. Doming.
Candelariae, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 108.—Costa Rica.
candicans, *Wall. Cat.* n. 3123.—Ind. or.
candidans, *DC. Prod.* vi. 412.—Ins. Falkland.
candidus, *DC. l. c.* 355.—Sicilia.
Candolleanus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 345.—Patagon.
Candolleanus, *Wall. Cat.* n. 3119.—Reg. Himal.
canescens, Bernh. ex DC. Prod. vi. 414 = erucifolius.
cannabifolius, Less. in Linnaea, vi. (1831) 242 = palmatus.
cantabricus, Willk. in Flora, xxxiv. (1851) 745 = pyrenaicus.
canus, *Hook. Fl. Bor. Am.* i. 333. t. 116.—Am. arct.
canus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 118 = leucophyllus.
capillaris, *Gaudich. in Freyc. Voy. Bot.* 468.—Ins. Sandvic.
capillifolius, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 123.—Tasman.
capitatus, *Steud. Nom.* ed. II. ii. 559; *Andrae, in Bot. Zeit.* xiii. (1855) 310.—Reg. Carpath. et Caucas.
Cappa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 179 = densiflorus.
cappadocicus, *Boiss. Fl. Orient.* iii. 398.—Cappadoc.
caracasanus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 331.—Venezuela.
Cardamine, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 98.—Am. bor.
cardaminefolius, *DC. Prod.* vi. 382.—Afr. austr.
cardiophyllus, Hemsl. Biol. Centr. Am. Bot. ii. 237 = ovatifolius, *Sch. Bip.*
carduifolius, Desf. Cat. Hort. Par. ed. III. 177 = Erechtites praealta.
caricifolius, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 345.—Patagon.
cariensis, *Boiss. Fl. Orient.* iii. 398.—As. Min.
carniolicus, *Willd. Sp. Pl.* iii. 1993.—Europ.
carnosus, Lam. Fl. Fr. ii. 131 = S. Doria.
carnosus, *Phil. in Anal. Univ. Chil.* (1862) II. 382.—Chili.
carnosus, Presl, Fl. Sic. p. xxviii = aethnensis.
carnosus, Thunb. Prod. Pl. Cap. 158 = elegans.
carnulentus, DC. Prod. vi. 372 = lautus.
carolinianus, Spreng. Syst. iii. 559 = lobatus.
carpathicus, Herbich, Addit. Fl. Galic. 43 = abrotanifolius.
carpetanus, *Boiss. & Reut. Pugill. Pl. Nov.* 59.—Hispan.
carroënsis, *DC. Prod.* vi. 396.—Afr. austr.
carthagenensis, *DC. l. c.* 420.—Am. austr.
casapaltensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 47.—Peruv.
caspicus, Less. in Linnaea, ix. (1834) 170 = racemosus.
cassinioides, *Hook. f. Handb. N. Zeal. Fl.* 163.—N. Zel.
Castagneanus, *DC. Prod.* vi. 354.—Turc.
castaneaefolius, *DC. l. c.* 425.—Venezuela.
caucasicus, *DC. l. c.* 348.—Reg. Caucas.
caudatus, *DC. l. c.* 384.—Afr. austr.
caulopterus, *DC. l. c.* 388.—Afr. austr.
cedrosensis, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 194.—Calif.
Centropappus, *F. Muell. Cat. Hort. Melb.* (1858) 26.—Austral.
ceratophylloides, Griseb. in Goett. Abh. xxiv. (1879) 206 = chilensis.
ceratophyllus, [D.] Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 332 = chilensis.
ceratophyllus, Nees, in Neuwied, Reise Nord-Am. ii. 48 = aureus.
cerberoanus, *Remy, in C. Gay, Fl. Chil.* iv. 185.—Chili.
cernuus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 239.—Am. bor.
cernuus, Linn. f. Suppl. 370 = Gynura cernua.
cervariaefolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
Cervi-cornis, *Phil. in Anal. Univ. Chil.* (1873) 497.—Chili.

SENECIO :—

cervilingua, Sch. Bip. ex Wedd. Chlor. And. i. 125 = formosus.
chamaecephalus, *Wedd. l. c.* 132.—Chili?
chamaedrifolius, Less. in Linnaea, vi. (1831) 247 = nigrescens.
chamaemelifolius, DC. Prod. vi. 383 = diffusus.
chamomillaefolius, *Phil. in Anal. Univ. Chil.* (1865) II. 349.—Chili.
Cheesemani, Hook. f. in Hook. Ic. Pl. t. 1201 = myrianthos.
cheiranthifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 176.—Mexic.
chenopodifolius, *DC. Prod.* vi. 364.—Reg. Himal.
chenopodioides, *H. B. & K. Nov. Gen. et Sp.* iv. 179.—Mexic.
chiapensis, *Hemsl. in Biol. Centr. Am. Bot.* ii. 238.—Mexic.
chilensis, *Less. in Linnaea*, vi. (1831) 248.—Chili.
chiloensis, *Phil. l. c.* xxix. (1857–58) 2.—Chili.
chinensis, DC. Prod. vi. 363 = scandens.
Chionogeton, *Wedd. Chlor. And.* i. 91.—N. Granat.
chionopappus, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Ind. or.
chordifolius, *Hook. f. Bot. Mag.* t. 6216.—Afr. austr.
Chrysactis, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 237.—Mexic.
chrysanthemifolius, Bivona, Pl. Sic. Cent. ii. 52 = aethnensis.
chrysanthemifolius, DC. Prod. vi. 345 = coronopifolius.
chrysanthemifolius, Moretti, in Bibl. Ital. xii. (1818) 374 = praealtus.
chrysanthemifolius, Poir. Encyc. vii. 96 = squalidus.
chrysanthemoides, *DC. Prod.* vi. 365.—Reg. Himal.
chrysanthemoides, Schrank, Baier. Fl. ii. 386 = squalidus.
Chrysocoma, Meerb. Afbeeld. t. 39 = paniculatus.
chrysocomoides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 346.—Patagon.
chrysolepis, *Phil. Fl. Atac.* 32.—Chili.
cicatricosus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
cicutarius, Stev. ex DC. Prod. vi. 349 = palmatus.
ciliatus, Walt. Fl. Carol. 208 = Erigeron canadensis.
cilicius, *Boiss. Diagn.* Ser. II. iii. 37.—As. Min.
ciliolatus, *DC. Prod.* vi. 374.—Austral.
Cineraria, *DC. l. c.* 355.—Reg. Mediterr.
Cinerariae, Spreng. Syst. iv. Cur. Post. 346 = praealtus.
cinerarioides, *H. B. & K. Nov. Gen. et Sp.* iv. 183.—Mexic.
cinerarioides, A. Rich. Sert. Astrol. 128 = australis.
cinerarioides, Viv. Fl. Lib. Spec. 68 = erucifolius.
cinerarius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Cineraria.
cinerascens, *Ait. Hort. Kew.* ed. I. iii. 191.—Afr. austr.
cinereus, *Colla, in Mem. Acc. Torin.* xxxviii. (1835) 33.—Chili.
cinereus, Moretti, Elench. Sem. Hort. Tic. (1829); ex DC. Prod. vi. 436 = Morettii.
cirsiifolius, *Sch. Bip. in Flora*, xxvii. (1845) 499.—Mexic.
cirsioides, Hemsl. Biol. Centr. Am. Bot. ii. 238 = roseus.
cissampelinus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
cladobotrys, *Ledeb. Fl. Ross.* ii. 644.—Reg. Caucas.
clarenceanus, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 14.—Afr. trop.
clarioneaefolius, *Remy, in C. Gay, Fl. Chil.* iv. 156.—Chili.
Clarkianus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 362.—Calif.
Claussenii, *Decne. in Rev. Hortic.* Sér. IV. iv. (1855) 181.—Bras.
claviseta, Pomel, Nouv. Mat. Fl. Atl. 60 = Decaisnei.
clematoides, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 442.—Abyss.
Clevelandi, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 87.—Calif.
clivicolus, *Wedd. Chlor. And.* i. 130.—Bolivia.
clivorum, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 220, 221.—China; Japon.
Clusianus, Reichb. f. Ic. Fl. Germ. xvi. 44. t. 980 = alpestris.
Clusii, Sch. Bip. in Flora, xxviii. (1845) 498 = Doronicum hirsutum.
cochlearifolius, *Boj. ex DC. Prod.* vi. 379.—Madag.
Colensoi, *Hook. f. Fl. N. Zel.* i. 147.—N. Zel.

SENECIO:—
coleophyllus, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 90.—Afr. austr.
collinus, *DC. Prod.* vi. 420.—Peruv.
colpodes, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 98.—Bras.
columbaria, Remy, in C. Gay, Fl. Chil. iv. 208 = calocephalus, *Poepp. & Endl.*
Commersonii, Spreng. Syst. iii. 554 = Faujasia pinifolia.
commutatus, Heller, Fl. Wirceb. ii. 293; Spenn. Fl. Friburg. ii. 525 = Fuchsii.
comosus, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
compactus, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 395.—N. Zel.
concolor, *DC. Prod.* vi. 407.—Afr. austr.
concolor, Harv. in Harv. & Sond. Fl. Cap. iii. 362 = speciosus.
confertus, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 441.—Abyss.
congestus, DC. Prod. vi. 363 = palustris.
consanguineus, *DC. l. c.* 381.—Afr. austr.
controversus, F. Schultz, ex Nym. Consp. 353 = nemorensis.
conyzaefolius, *Baker, in Mart. Fl. Bras.* vi. III. 303.—Bras.
conyzoides, *DC. Prod.* vi. 412.—Ins. S. Doming.
coquimbensis, *Phil. in Linnaea*, xxviii. (1856) 749.—Chili.
cordato-rotundus, *Steud. Nom.* ed. II. ii. 559.—Chili.
cordatus, Koch, in Flora, xvii. (1834) 614 = alpinus.
cordatus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 411.—Am. bor.
cordatus, Wall. Cat. n. 3109 = tetranthus.
cordifolius, [*Clairv.*] *Man. Herb.* 241.—Europ.
cordifolius, Link, Handb. i. 748 = alpinus.
cordifolius, *Linn. f. Suppl.* 372.—Afr. austr.
cordovensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 238.—Mexic.
coriaceus, *Ait. Hort. Kew.* ed. I. iii. 195.—Oriens.
Cornu-cervi, *MacOwan, in Journ. Linn. Soc.* x. (1869) 482.—Afr. austr.
coronatus, Harv. in Harv. & Sond. Fl. Cap. iii. 369 = lasiorhizus.
coronopifolius, Burm. f. Fl. Ind. 181 = javanicus.
coronopifolius, *Desf. Fl. Atlant.* ii. 273.—Europ.; Oriens; As. bor.; Afr. bor.
coronopifolius, Willd. ex Ball, in Journ. Linn. Soc. xvi. (1878) 514 = gallicus.
coronopodiphyllus, *Remy, in C. Gay, Fl. Chil.* iv. 156.—Chili.
Coronopus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 413 = californicus.
corsicus, *Sch. Bip. in Flora*, xxviii. (1845) 498 = Doronicum corsicum.
corymbiferus, *DC. Prod.* vi. 402.—Afr. austr.
corymbosus, C. B. Clarke, Comp. Ind. 205 = pachycarpus.
corymbosus, Dalla Torre, Atl. Alpenfl. 135 = incanus.
corymbosus, *Wall. Cat.* n. 3121.—Ind. or.
Cotyledonis, *DC. Prod.* vi. 402.—Afr. austr.
Coxi, *Phil. in Linnaea*, xxxiii. (1864–65) 150.—Chili.
crambefolius, Boiss. & Kotschy, in Boiss. Fl. Orient. iii. 396 = eriospermus.
crassicaulis, *Brongn. in Rev. Hortic.* Sér. III. i. (1847) 87.—Mexic.
crassiflorus, *DC. Prod.* vi. 412.—Am. austr.
crassifolius, Sibth. & Sm. Fl. Graec. ix. 53. t. 868 = vernalis.
crassifolius, *Willd. Sp. Pl.* iii. 1982.—Europ.
crassiusculus, *DC. Prod.* vi. 401.—Afr. austr.
crassulaefolius, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
crassulus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 54.—Am. bor. occ.
crassus, Vell. Fl. Flum. viii. t. 111 = Erechtites valerianaefolia.
crenatus, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
crenulatus, *DC. Prod.* vi. 385.—Afr. austr.
crepidifolius, *DC. l. c.* 425.—Peruv.
crepidioides, Aschers. in Schweinf. Beitr. Fl. Aethiop. 155 = Gynura crepidioides.
crinitus, Bertol. in Opusc. Sc. Bologn. (1817) i. 234 = delphinifolius.
crispatus, *DC. Prod.* vi. 359.—Europ.
crispus, Kitt. Taschenb. ed. II. 588; Kern. in Oestr. Bot. Zeitschr. xxi. (1871) 202 = crispatus.

SENECIO:—
crispus, *Phil. Fl. Atac.* 33.—Chili.
crispus, *Thunb. Prod. Pl. Cap.* 157.—Afr. austr.
cristatus, Thunb. l. c. = rosmarinifolius.
crithmifolius, A. Rich. Sert. Astrol. 116 = lautus.
crithmifolius, Scop. Fl. Carn. ed. II. ii. 163 = Inula crithmoides.
crithmoides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 347.—Chili.
croaticus, Waldst. & Kit. Pl. Rar. Hung. ii. 153 = S. Cacaliaster.
croceus, DC. Prod. vi. 359 = crispatus.
crucaefolius, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 120, in syn., sphalm. = erucifolius.
cruciatus, Linn. f. Suppl. 371 = juniperinus.
cruentus, *DC. Prod.* vi. 410.—Ins. Canar.
cruentus, *Roth, in Roem. Arch.* i. III. (1798) 44.—Hab.?
crymophilus, *Wedd. Chlor. And.* i. 95.—N. Granat.
cryptocephalus, Sch. Bip. ex Wedd. l. c. 90 = repens.
cucullatus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 331.—Venezuel.
culcitioides, *Sch. Bip. in Bonplandia*, iv. (1856) 55; *et in Lechl. Berb. Am. Austr.* 57, nomen.—Peruv.
Culcitium, *Remy, in C. Gay, Fl. Chil.* iv. 200.—Chili.
Cumingii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 338.—Chili.
Cumingii, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 330 = Arnica cordifolia *vel* latifolia.
cuneatus, *Hook. f. Fl. Antarct.* 318.—Patagon.
cuneifolius, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 126.—Bras.
cuneifolius, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
Cunninghami, *DC. Prod.* vi. 371.—Austral.
curvatus, *Baker, in Journ. Linn. Soc.* xx. (1883) 190.—Madag.
curvidens, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
cuspidatus, DC. Prod. vi. 419 = chilensis.
cydoniaefolius, *Steud. Nom.* ed. II. ii. 560.—Chili.
cygnorum, Steetz, in Lehm. Pl. Preiss. i. 483 = ramosissimus.
Cymbalaria, Pursh, Fl. Am. Sept. ii. 530 = aureus.
cymbalariaefolius, *Less. Syn. Comp.* 391.—Afr. austr.
cymbalarioides, Buek, Ind. Cand. ii. p. vi; Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 412 = aureus.
cymosus, *Remy, in C. Gay, Fl. Chil.* iv. 170.—Chili.
cypriaefolius, Steud. Nom. ed. II. ii. 560, sphalm. = oxyriaefolius.
Cyri, C. Koch, in Linnaea, xxiv. (1851) 362 = racemosus.
dahuricus, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) II. 19 = palmatus.
dahuricus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—As. or.; Japon.
dalmaticus, Vis. Fl. Dalm. ii. 69, in syn. = S. Jacobaea.
Daltoni, *F. Muell. Fragm.* vi. 27.—Austral.
Dalzellii, *C. B. Clarke, Comp. Ind.* 201.—Ind. or.
Danai, *A. Gray, in Proc. Am. Acad.* v. (1861) 142.—Peruv.
Danyausii, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 48 = Laseguei.
daphnoides, Poir. Encyc. Suppl. v. 133 = S. Hubertia.
Darwinii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 333.—Patagon.
davuricus, Franch. & Sav. Enum. Pl. Jap. i. 250 = dahuricus.
dealbatus, *Phil. in Linnaea*, xxviii. (1856) 747.—Chili.
debilis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 400.—Afr. austr.
debilis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 408 = aureus.
debilis, *Phil. in Linnaea*, xxviii. (1856) 752.—Chili.
Decaisnei, *DC. Prod.* vi. 342.—Afr. bor.; Oriens.
decurrens, *DC. l. c.* 388.—Afr. austr.
deferens, *Griseb. in Goett. Abh.* xix. (1874) 193.—Reg. Argent.
Delessertii, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura bicolor?
delphinifolius, Reichb. Pl. Crit. iv. 29 = erucifolius.
delphinifolius, *Vahl, Symb. Bot.* ii. 91. t. 45.—Reg. Mediterr.

SENECIO:—
deltoideus, *Less. Syn. Comp.* 392.—Afr. austr.
deltophyllus, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 487.—China.
demissus, *Phil. in Anal. Univ. Chil.* (1870) II. 181.—Chili.
densiflorus, Martens, in Bull. Acad. Brux. viii. (1841) 67 = lobatus.
densiflorus, *Wall. Cat.* n. 3116.—Reg. Himal.
dentatus, *Jacq. Coll.* v. 150. t. 6. f. 2.—Afr. austr.
dentatus, Wall. Cat. n. 3112 = Wallichii.
denticulatus, *DC. Prod.* vi. 416.—Am. austr.
denticulatus, O. F. Muell. in Fl. Dan. t. 791 = vulgaris.
denudatus, D. Don, Prod. Fl. Nep. 179 = nudicaulis.
Deppeanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 239.—Mexic.
depressus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 342.—Chili.
desertorum, *Hemsl. Biol. Centr. Am. Bot.* ii. 239.—Mexic.
desiderabilis, Vell. Fl. Flum. viii. t. 108 = ellipticus.
desideratus, Baker, in Mart. Fl. Bras. vi. III. 318 = ellipticus.
desideratus, *DC. Prod.* vi. 413.—Patagon.
desmatus, *Klatt, in Linnaea*, xxxvii. (1871–73) 511.—Madag.
desquamatus, Willd. Enum. Hort. Berol. Suppl. 58; ex DC. Prod. vi. 346 = gallicus.
Diazi, *Phil. in Anal. Univ. Chil.* (1862) II. 399.—Chili.
dichrous, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Bras.
diclinus, *Wedd. Chlor. And.* i. 107.—Peruv.
dictyophyllus, *Benth. Pl. Hartw.* 43.—Mexic.
difficilis, Dufour, in Ann. Sc. Nat. Sér. I. v. (1825) 428. t. 11 = gallicus.
diffusus, *Linn. f. Suppl.* 371.—Afr. austr.
digitalifolius, *DC. Prod.* vi. 397.—Afr. austr.
Diodon, *DC. l. c.* 399.—Afr. austr.
diosmoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 210.—Peruv.
discifolius, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 100.—Afr. trop.
discolor, Boj. ex DC. Prod. vi. 379 = hypargyraeus.
discolor, *DC. Prod.* vi. 412.—Jamaic.
discolor, Desf. Tabl. ed. I. 104 = Gynura haematophylla.
disjectus, *Wedd. Chlor. And.* i. 115.—Peruv.
divaricatus, Linn. Sp. Pl. 866 = Gynura divaricata.
divergens, F. Schultz, ex Nym. Consp. 355 = erraticus.
diversifolius, Dum. Fl. Belg. Prod. 66 = Kickxii.
diversifolius, Harv. in Harv. & Sond. Fl. Cap. iii. 375 = polymorphus.
diversifolius, *Phil. in Anal. Univ. Chil.* (1873) 495.—Chili.
diversifolius, A. Rich. Tent. Fl. Abyss. i. 437 = Gynura crepidioides.
diversifolius, *Wall. Cat.* n. 3128.—Reg. Himal.
Doeringii, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 358.—Reg. Argent.
Dombeyanus, *DC. Prod.* vi. 418.—Am. austr.
Donianus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 332.—Chili.
doratophyllus, *Benth. Pl. Hartw.* 87.—Guatemala.
Doria, Kickx, Fl. Brux. 283 = Kickxii.
Doria, C. Koch, in Linnaea, xvii. (1843) 48 = macrophyllus.
Doria, *Linn. Syst.* ed. X. 1215.—Europ. austr.; Oriens.
doriaeformis, *DC. Prod.* vi. 352.—As. Min.; Syria.
doroniciflorus, DC. l. c. 385 = albanensis.
doroniciflorus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 338 = Selloi.
Doronicum, *Linn. Syst.* ed. X. 1215.—Europ.
Doronicum, Willk. in Flora, xxxiv. (1851) 745 = Lagascanus.
Douglasii, *DC. Prod.* vi. 429.—Calif.
dracunculoides, *DC. l. c.* 400.—Afr. austr.
Dregeanus, *DC. l. c.* 407.—Afr. austr.
dryadeus, Sieber, ex DC. l. c. 374 = australis.
drymophilus, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 69 = Gynura Pseudo-china.
dryophyllus, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 283.—Peruv.
dubius, Beck, Bot. N. & M. St. 200 = tomentosus.
dubius, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 342 = pedunculatus.

SENECIO :—
dubius, *Ledeb. Fl. Alt.* iv. 112.—As. bor. et centr.
dubius, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura dubia.
dumetorum, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 422.—Bras.
dunensis, *Dum. Fl. Belg. Prod.* 66.—Belgium.
Duriaei, *J. Gay, Ann. Sc. Nat.* Sér. II. vi. (1836) 346.—Hispan.
ecalyculatus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Emilia sonchifolia.
echinatus, *DC. Prod.* vi. 411.—Ins. Canar.
Ecklonianus, DC. Prod. vi. 384 = spiraeifolius.
Edgeworthii, *Hook. f. Fl. Brit. Ind.* iii. 346.—Ind. or.
Eightsii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 332.—Patagon.
elaeagnifolius, *Hook. f. Fl. N. Zel.* i. 150. t. 41.—N. Zel.
elatior, Steud. Nom. ed. II. ii. 560 = multiflorus, *DC.*
elatoides, *Wedd. Chlor. And.* i. 101.—N. Granat.
elatus, *H. B. & K. Nov. Gen. et Sp.* iv. 180.—Peruv.
elbrusensis, *Boiss. Fl. Orient.* iii. 410.—Persia.
elegans, *Linn. Sp. Pl.* 869.—Afr. austr.
elegans, Thunb. Fl. Cap. ed. Schult. 685 = arenarius.
Elliottii, Torr. & Gray, Fl. N. Am. ii. 443 = aureus.
ellipticus, *DC. Prod.* vi. 420.—Bras.
elodes, *Boiss. ex DC. l. c.* vii. 301.—Hispan.
elongatus, Pursh, Fl. Am. Sept. ii. 529 = aureus.
emilioides, *Baker, in Mart. Fl. Bras.* vi. III. 304.—Bras.
emilioides, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 158.—Afr. trop.
emirnensis, *DC. Prod.* vi. 377.—Madag.
emodensis, Sch. Bip. in Flora, xxviii. (1845) 50 = S. Ligularia.
Encelia, *Remy, in C. Gay, Fl. Chil.* iv. 204.—Chili.
Endlicheri, *DC. Prod.* vi. 373.—Austral.
erechthitoides, *Baker, in Journ. Bot.* xx. (1882) 171.—Madag.
eremophilus, *Phil. Fl. Atac.* 32.—Chili.
eremophilus, *Richards. in Frankl. 1st Journ. App.* ed. II. 31.—Am. bor.
ericaefolius, *Benth. Pl. Hartw.* 208.—Ecuador.
eriobasis, *DC. Prod.* vi. 388.—Afr. austr.
eriocladus, *Wedd. Chlor. And.* i. 131.—Peruv.
eriophyton, *Remy, in C. Gay, Fl. Chil.* iv. 159.—Chili.
eriopus, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 116.—Hispan.
eriospermus, *DC. Prod.* vi. 358.—As. Min.
erisithalifolius, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 321.—Bras.
erodiifolius, DC. Prod. vi. 381 = repandus.
erosus, *Linn. f. Suppl.* 370.—Afr. austr.
erosus, *Wedd. Chlor. And.* i. 228.—Bolivia.
erraticus, *Bertol. Rar. Ital. Pl. Dec.* iii. 62.—Europ.; Oriens.
erraticus, Willk. in Flora, xxxiv. (1851) 744 = aquaticus.
erubescens, *Ait. Hort. Kew.* ed. I. iii. 190.—Afr. austr.
erubescens, *Panč. Elem. Fl. Bulg.* 41.—Europ.
erucaefolius, Kunze, in Flora, xxix. (1846) 655 = praealtus.
erucaefolius, Ledeb. Fl. Alt. iv. 110 = S. Jacobaea.
erucaeformis, *Remy, in C. Gay, Fl. Chil.* iv. 180.—Chili.
erucifolius, *Linn. Fl. Suec.* ed. II. 291.—Europ.; As. bor.
erysimoides, *DC. Prod.* vi. 382.—Afr. austr.
eubaeus, *Boiss. & Heldr. Diagn.* Ser. II. iii. 36.—Graec.
eudorus, *DC. Prod.* vi. 352.—Hab.?
eupatorioides, *Sch. Bip. in Flora*, xxvii. (1844) 702.—Afr. austr.
euriopoides, *DC. Prod.* vi. 396.—Afr. austr.
eurycephalus, *Torr. & Gray, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 109.—Calif.
evacoides, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
exaltatus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 410 = lugens.
exilis, Homb. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 50 = patagonicus, *Hook. & Arn.*
eximius, Hemsl. Biol. Centr. Am. Bot. ii. 239 = callosus.
expansus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 394.—Afr. austr.

SENECIO :—
expansus, *Wedd. Chlor. And.* i. 107.—Bolivia.
exsertus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Madag.
exsquameus, Brot. Fl. Lusit. i. 388 = gallicus.
Exul, *Hance, in Journ. Bot.* vi. (1868) 174.—China.
Fadyenii, *Griseb. Fl. Brit. W. Ind.* 382.—Jamaic.
falklandicus, *Hook. f. Fl. Antarct.* 316.—Ins. Falkland.
farfaraefolius, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 33.—As. Min.
farfaraefolius, *Sch. Bip. ex C. Koch & Fint. Wochenschr.* i. (1858) 212; *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 483.—Japon.
farfarus, Hemsl. Biol. Centr. Am. Bot. ii. 239 = tussilaginoides.
Farfugium, C. Koch, in C. Koch & Fint. Wochenschr. i. (1858) 209 = Kaempferi.
farinaceus, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 443.—Abyss.
farinifer, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 331.—Chili.
Farrisii, *Baker, in Saund. Refug. Bot.* t. 250.—Peruv.
fasciculatus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 344.—Chili.
fasciculatus, Wall. ex DC. Prod. v. 439 = Laggera flava.
fastigiatus, A. Gray, Pl. Wright. ii. 99 = Douglasii.
fastigiatus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 410.—Am. bor. occ.
fastigiatus, Schwein. ex Ell. Sketch, ii. 331 = aureus.
faugasioides, *Baker, in Journ. Bot.* xx. (1882) 170.—Madag.
Fendleri, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 108.—N. Mexic.
ferulifolius, Stokes, Bot. Mat. Med. iv. 216 = adonidifolius.
ficoides, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
filaginoides, *DC. Prod.* vi. 413.—Chili.
filifolius, Berg. Desc. Pl. Cap. 278 = umbellatus.
filifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 399.—Afr. austr.
filifolius, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 414 = Douglasii.
filipes, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 87 = matricariaefolius.
Finlaysonianus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura sarmentosa.
Fischeri, Sch. Bip. in Flora, xxviii. (1845) 498 = Erechtites praealta.
Fischerianus, *Steud. Nom.* ed. II. ii. 560.—Armenia.
fistulosus, *Poepp. ex DC. Prod.* vi. 418.—Chili.
flabellatus, Viv. App. Fl. Cors. Prod. 6 = delphinifolius.
flaccidifolius, *Wedd. Chlor. And.* i. 113.—Peruv.
flaccidus, *Bolle, in Bonplandia*, viii. (1860) 134.—Ins. Canar.
flaccidus, *Less. in Linnaea*, v. (1830) 161.—Mexic.
flaccidus, A. Rich. Sert. Astrolab. 110 = Erechtites prenanthoides.
flagellisectus, Griseb. in Goett. Abh. xix. (1874) 191 = brasiliensis.
flammeus, *DC. Prod.* vi. 362.—Dahuria.
flavus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 319. t. 107 = Decaisnei.
flexicaulis, Edgew. in Trans. Linn. Soc. xx. (1846) 74 = scandens.
flexuosus, *Clarke, Trav.* ed. II. i. *Add.* p. xxi.—Thracia.
flexuosus, *Less. Syn. Comp.* 392.—Ins. Maurit.
flexuosus, Wall. Cat. n. 3110 = scandens.
floccidus, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 50 = patagonicus.
flocciferus, DC. Prod. vi. 426 = Malacothrix obtusa.
floccosus, Schur, Enum. Pl. Transs. 350 = S. Jacobaea.
flocculidens, *Sch. Bip. ex Wedd. Chlor. And.* i. 127.—Venezuela.
flocoso-araneosus, Steud. Nom. ed. II. ii. 560 = andinus.
floribundus, Wall. Cat. n. 3111 = triligulatus.
floridanus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Am. bor.
flosculosus, Gilib. Fl. Lituan. i. 198 = vulgaris.
flosculosus, Jord. Cat. Jard. Dijon (1848), 30 = S. Jacobaea.
fluminensis, *Steud. Nom.* ed. II. ii. 560.—Bras.

SENECIO :—
fluviatilis, Wallr. in Linnaea, xiv. (1840) 64 = sarracenicus.
foeniculaceus, Tenore, Prod. Fl. Nap. p. xlix = lividus.
foeniculoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 395.—Afr. austr.
foetens, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura nepalensis.
foliosus, Colm. Catal. Pl. Cataluña, 87 = jacobaeoides.
foliosus, *Salzm. ex DC. Prod.* vi. 351.—Afr. bor.
fontanus, Wallr. in Linnaea, xiv. (1840) 353 = nemorensis.
Forbesii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 420.—Afr. trop.
formosus, *H. B. & K. Nov. Gen. et Sp.* iv. 177.—Mexic.
Forskâhlii, Baill. Hist. Pl. viii. 308 = odorus.
Forsteri, *Phil. in Anal. Univ. Chil.* (1873) 494.—Chili.
Forsteri, Schlecht. in DC. Prod. vi. 373 = Brachyglottis Forsteri.
Fradini, Pomel, Nouv. Mat. Fl. Atl. 61 = leucanthemifolius, *Poir.*
fragrans, *Fisch. ex Spreng. Syst.* iv. Cur. Post. 302.—Sibir.
fragrans, Lehm. ex Spreng. l. c. iii. 518 = Erigeron fragrans.
fragrans, Riddell, ex Torr. & Gray, Fl. N. Am. ii. 444 = Douglasii.
Fremontii, *Torr. & Gray, l. c.* 445.—Am. bor.
Fresenii, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 160.—Afr. bor.
frigidus, *Less. in Linnaea*, vi. (1831) 239.—Sibir.; Am. arct.
frigidus, Rupr. in Beitr. Pfl. Russ. Reich. vii. 28 (an praec.?).—Sibir.
frigidus, Steud. Nom. ed. II. ii. 561 = linifolius, *Linn.*
frondosus, Tausch, ex Reichb. Fl. Germ. Excurs. 245 = nemorensis.
fruticosus, *Vell. Fl. Flum.* viii. t. 109.—Bras.
fruticulosus, *Sibth. & Sm. Fl. Graec. Prod.* ii. 178.—Ins. Creta.
Fuchsii, *C. C. Gmel. Fl. Bad.* iii. 444.—Europ.
fuliginosus, H. B. & K. Nov. Gen. et Sp. iv. 186 = Gynoxis fuliginosa.
Funckii, *Sch. Bip. ex Wedd. Chlor. And.* i. 125.—Venezuela.
furfuraceus, Kotschy, ex Boiss. Fl. Orient. iii. 400 = farfaraefolius, *Boiss. & Kotschy.*
Fussii, *Andrae, in Bot. Zeit.* xiii. (1855) 310.—Hungar.
gabonicus, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 421.—Afr. trop.
gaffatensis, Vatke, in Linnaea, xxxix. (1875) 505 = vulgaris.
Galeottii, *Hemsl. Biol. Centr. Am. Bot.* ii. 239.—Mexic.
Gallerandianus, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 365.—Alger.
gallicus, *Chaix, in Vill. Hist. Pl. Dauph.* i. 371; iii. 230.—Europ.
gamolepoides, *Sch. Bip. in Journ. Bot.* i. (1863) 236.—Guian.
Gardneri, *C. B. Clarke, Comp. Ind.* 202.—Zeylan.
Gaudichaudianus, *A. Rich. Sert. Astrolab.* 98.—Austral.
Gayanus, *DC. Prod.* vi. 414.—Chili.
geifolius, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Reg. Caucas.
Geissei, *Phil. in Anal. Univ. Chil.* (1865) II. 348.—Chili.
gelidus, *Wedd. Chlor. And.* i. 95.—N. Granat.
geniculatus, *Steud. Nom.* ed. II. ii. 561.—Afr. austr.
georgianus, *DC. Prod.* vi. 371.—Austral.
Gerardi, Gren. & Godr. Fl. Fr. ii. 122 = S. Doronicum.
Gerardi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 380.—Afr. austr.
gerberaefolius, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 240.—Mexic.
Germaini, *Phil. in Linnaea*, xxix. (1857-58) 1.—Chili.
germanicus, Wallr. Sched. Crit. 476 = nemorensis.
Ghiesbreghtii, Hort. Hal. ex Regel, Ind. Sem. Hort. Petrop. (1860) 36 = grandifolius.
gibbosus, *DC. Prod.* vi. 355.—Ital.
Gibsoni, *Hook. f. Fl. Brit. Ind.* iii. 347.—Ind. or.
giganteus, *Desf. Fl. Atlant.* ii. 273. t. 234.—Alger.
Gigas, *Vatke, in Linnaea*, xxxix. (1875) 506.—Afr. trop.

SENECIO:—
Gilberti, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 208.—Anstral.
Gilliesianus, *Hieron. in Bol. Acad. Nat. Córdova*, iii. (1879) 356.—Chili.
Gilliesii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 343.—Chili.
Gilliesii, Phil. in Anal. Univ. Chil. (1870) ii. 182 = Gilliesianus.
glabellus, DC. Prod. vi. 360 = campestris.
glabellus, Poir. Encyc. vii. 102 = lobatus.
glaber, [Clairv.] Man. Herb. 240 = Adenostyles viridis.
glaber, *Less. in Linnaea*, vi. (1831) 248.—Chili.
glaber, Ucria, in Roem. Arch. i. i. (1796) 70 = squalidus.
glaberrimus, *DC. Prod.* vi. 403.—Afr. austr.
glaberrimus, Schur, Enum. Pl. Transs. 351 = S. Doronicum.
glabratus, *Hook. & Arn. Bot. Beech. Voy.* 32.—Chili.
glabrescens, Sch. Bip. in Flora, xxviii. (1845) 498 = Erechtites quadridentata.
glabrifolius, *DC. Prod.* vi. 406.—Afr. austr.
glacialis, Sch. Bip. in Flora, xxviii. (1845) 498 = Doronicum glaciale.
glacialis, *Wedd. Chlor. And.* i. 113.—Bolivia.
glandulosus, A. Cunn. ex DC. Prod. vi. 295 = Erechtites quadridentata?
glandulosus, [*D.*] *Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 334.—Chili.
glareosus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
glastifolius, *Hook. f. Fl. N. Zel.* i. 147. t. 39.—N. Zel.
glastifolius, *Linn. f. Suppl.* 372.—Afr. austr.
glaucescens, DC. Prod. vi. 395 = achilleaefolius.
glaucescens, Spreng. Syst. iii. 561 = squalidus.
glaucus, DC. Prod. vi. 344 = leucanthemifolius, *Poir.*
glaucus, *Linn. Sp. Pl.* 868.—Afr. bor.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. iii. 305.—Bras.
glomeratus, Desf. Tabl. ed. II. 124 = Erechtites arguta?
glutinarius, *DC. Prod.* vi. 381.—Afr. austr.
glutinosus, E. Mey. ex DC. l. c. 381 = praec.
glutinosus, *Roth, Catalecta*, fasc. ii. 109.—Hab.?
glutinosus, Schur, Enum. Pl. Transs. 347 = viscosus.
glutinosus, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
Gmelini, *Ledeb. Fl. Ross.* ii. 645.—Sibir. altaic.
gnaphalodes, *Sieber, Reise Kreta*, i. 352.—Ins. Creta.
gnidioides, *Phil. in Linnaea*, xxviii. (1856) 742.—Chili.
Godmanii, *Hemsl. Biol. Centr. Am. Bot.* ii. 240.—Guatemala.
Goeringii, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 126, nomen.—Java.
gonocaulos, *DC. Prod.* vi. 425.—Am. austr.
gonocladus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
goyazensis, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 421.—Bras.
graciliflorus, *DC. Prod.* vi. 365.—Reg. Himal.
gracilipes, *A. Gray, in Proc. Am. Acad.* v. (1861) 142.—Peruv.
gracilis, Arn. in Nov. Act. Nat. Cur. xviii. i. (1836) 349 = zeylanicus.
gracilis, *Pursh, Fl. Am. Sept.* ii. 529.—Am. bor.
Grahami, Benth. Pl. Hartw. 18 = S. Barba-Johannis.
Grahami, *Hook. f. Fl. Brit. Ind.* iii. 347.—Ind. or.
gramineus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 372.—Afr. austr.
graminicolus, Sch. Bip. in Flora, xxvii. (1844) 698 = crenatus.
graminifolius, Jacq. Misc. ii. 322 = paniculatus.
graminifolius, *Phil. in Anal. Univ. Chil.* (1865) ii. 330.—Chili.
granatensis, Boiss. Voy. Bot. Espagne, ii. 334 = Boissieri.
grandidentatus, Ledeb. Fl. Ross. ii. 636 = erucifolius.
grandiflorus, *Berg. Desc. Pl. Cap.* 280.—Afr. austr.
grandiflorus, Hoffmgg. & Link, Fl. Port. ii. 307. t. 100 = S. Doria.
grandifolius, *Less. in Linnaea*, v. (1830) 162.—Mexic.
grandis, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 422.—Bras.
grandissimus, Schur, Enum. Pl. Transs. 354 = sarracenicus.
graveolens, *Wedd. Chlor. And.* i. 111.—Bolivia.

SENECIO:—
Grayanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 241.—Mexic.
Grayi, Parry, ex A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864) 67, in nota = S. Soldanella.
Greenei, *A. Gray, in Proc. Am. Acad.* x. (1874) 75.—Calif.
Gregorii, *F. Muell. Rep. Babb. Exped.* 14.—Austral.
Greyii, *Hook. f. Fl. N. Zel.* i. 148. t. 38.—N. Zel.
Griffithii, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 193.—Reg. Himal.
grindeliaefolius, *DC. Prod.* vi. 416.—Chili.
Grisebachii, *Baker, in Mart. Fl. Bras.* vi. iii. 313.—Reg. Argent.
guatimalensis, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Guatemala.
guineensis, Sch. Bip. l. c. = Gynura cernua.
gummifer, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 331 = phagnaloides.
Gundelius, C. Koch, in Linnaea, xxiv. (1851) 364 = armenius.
gynoxoides, *Baker, in Mart. Fl. Bras.* vi. iii. 306.—Bras.
gynoxoides, Sch. Bip. ex Wedd. Chlor. And. i. 78 = Gynoxys albiflora.
gypsophilus, *Phil. in Anal. Univ. Chil.* (1873) 498.—Chili.
Haastii, *Hook. f. Handb. N. Zeal. Fl.* 159.—N. Zel.
hadiensis, *Forsk. Fl. Aegypt. Arab.* 149.—Arab.
haematophyllus, Willd. Enum. Hort. Berol. 875 = Gynura haematophylla.
Haenkei, *DC. Prod.* vi. 413.—Chili.
hakeaefolius, *Bert. ex DC. l. c.* 416.—Chili.
halimifolius, *Linn. Sp. Pl.* 871.—Afr. austr.
Haloragis, *Remy, in C. Gay, Fl. Chil.* iv. 172.—Chili.
hamamelidifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 179 (hamamelifolius).—Ecuador.
Hamiltonianus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura Pseudo-china.
Hartwegi, *Benth. Pl. Hartw.* 18.—Mexic.
hastatus, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 99.—Bras.
hastatus, Hook. f. Fl. Brit. Ind. iii. 353, in nota = sagittatus.
hastatus, *Linn. Sp. Pl.* 868.—Afr.
hastatus, Willd. Sp. Pl. iii. 1987 = hastulatus.
hastifolius, Less. Syn. Comp. 392 = cymbalarifolius.
hastulatus, *Linn. Sp. Pl.* ed. II. 1218.—Afr. austr.
Haworthii, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
Haworthii, *Steud. Nom.* ed. II. ii. 561.—Afr. austr.
hebespermus, *DC. Prod.* vii. 301.—Mesopot.
hebetatus, *Wedd. Chlor. And.* i. 114.—Peruv.
Hectori, *F. Buch. in Trans. N. Z. Inst.* v. (1873) 348.—N. Zel.
hederaefolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 241.—Mexic.
Heldreichii, *Boiss. Diagn.* Ser. I. vi. 95.—Graec.
helianthemoides, *Wedd. Chlor. And.* i. 115.—Boliv.
helichrysoides, F. Muell. in Trans. Vict. Inst. i. (1855) 39 = georgianus.
Hellerii, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 333.—Mexic.
helodes, *Benth. Pl. Hartw.* 42.—Mexic.
heracleifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 241.—Mexic.
Herbichia, Steud. Nom. ed. II. ii. 561 = abrotanifolius.
Heritieri, *DC. Prod.* vi. 409.—Ins. Canar.
Hervaloira, Link, ex Nym. Consp. 354 = caespitosus, *Brot.*
heteroclinius, DC. Prod. vi. 380 = abruptus.
heterogamus, *Hemsl. Biol. Centr. Am. Bot.* ii. 242.—Costa Rica.
heterophyllus, DC. Prod. vi. 414 = pilquensis.
heterophyllus, Drège, ex Harv. & Sond. Fl. Cap. iii. 355 = littoreus.
heterophyllus, Thunb. Prod. Pl. Cap. 157 = abruptus.
heteroschizus, *Baker, in Mart. Fl. Bras.* vi. iii. 325.—Bras.
heterotrichius, *DC. Prod.* vi. 419.—Bras.
hewrensis, *Hook. f. Fl. Brit. Ind.* iii. 346.—Ind. or.
Heyneanus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura nitida.
hieracifolius, Forsk. Fl. Aegypt. Arab. p. lxxiii = arabicus.
hieracifolius, Linn. Sp. Pl. 866 = Erechtites praealta.

SENECIO:—
hieracioides, *DC. Prod.* vi. 384.—Afr. austr.
Hieracium, *Remy, in C. Gay, Fl. Chil.* iv. 202.—Chili.
Hieronymii, *Griseb. in Goett. Abh.* xxiv. (1879) 205.—Reg. Argent.
himalensis, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 78.—Reg. Himal.
Hindsii, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = scandens.
hirsutulus, *Phil. in Linnaea*, xxxiii. (1864–65) 163.—Chili.
hirsutus, [Clairv.] Man. Herb. 240 = Adenostyles albida.
hirtellus, *DC. Prod.* vi. 401.—Afr. austr.
hirtifolius, *DC. l. c.* 394.—Afr. austr.
hispidulus, A. Rich. Sert. Astrolab. 92 = Erechtites hispidula.
hispidulus, A. Rich. l. c. t. 34 = odoratus.
Hochstetteri, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) i. *Intell.* 27, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 435.—Abyss.
Hohenackeri, *Hook. f. Fl. Brit. Ind.* iii. 345.—Ind. or.
Hohenackeri, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen; *Wedd. Chlor. And.* i. 131.—Peruv.
holophyllus, *Remy, in C. Gay, Fl. Chil.* iv. 147.—Chili.
homogynefolius, Kotschy, ex Boiss. Fl. Orient. iii. 400 = farfaraefolius, *Boiss.*
Hookeri, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrolab. et Zél. 50 = patagonicus, *Hook. & Arn.*
Hookeri, *Torr. & Gray, Fl. N. Am.* ii. 438.—Am. bor.
Howellii, Greene, in Bull. Torrey Bot. Club, viii. (1881) 98 = canus.
huachucanus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 54.—Arizona.
Hualtata, *Bert. ex DC. Prod.* vi. 417.—Chili.
Hubertia, *Pers. Syn.* ii. 436.—Ins. Borbon.
Humboldtianus, *DC. Prod.* vi. 421.—Ecuador.
humilis, *Desf. Fl. Atlant.* ii. 271. t. 233.—Sicil.; Afr. bor.
humilis, Duby, ex Nym. Consp. 357 = crassifolius.
humillimus, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen; *Wedd. Chlor. And.* i. 104. t. 19 B.—Peruv.
Huntii, *F. Muell. Enum. Pl. Chath. Isl.* 23. t. 3.—N. Zel.
hybridus, Hort. ex Regel, Gartenfl. (1863) t. 394 = cruentus.
hydrophilus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 411.—Am. bor. occ.
hydrophilus, *Hoffmgg. & Link, ex Nym. Consp.* 356.—Lusitan.
hyemalis, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 514 = leucanthemifolius, *Poir.*
hyoseridifolius, *Wedd. Chlor. And.* i. 108.—Peruv.
hypargyraeus, *DC. Prod.* vi. 379.—Madag.
hypochionaeus, *Boiss. Diagn.* Ser. I. iv. 14.—As. Min.
hypochoerideus, *DC. Prod.* vi. 386.—Afr. austr.
hypoleucus, *F. Muell. ex Benth. Fl. Austral.* iii. 672.—Austral.
hypsobates, *Wedd. Chlor. And.* i. 91.—N. Granat.
hypsophilus, *Phil. in Anal. Univ. Chil.* (1862) ii. 382.—Chili.
icoglossus, *DC. Prod.* vi. 420.—Bras.
ignobilis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 299 = Erechtites ignobilis.
ilicifolius, Berg. Desc. Pl. Cap. 281 = lyratus.
ilicifolius, Less. Syn. Comp. 392 = Gynoxys ilicifolia.
ilicifolius, *Linn. Sp. Pl.* 871 (Quid?).—Afr. austr.
ilicifolius, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
illapelinus, *Phil. in Linnaea*, xxxiii. (1864–65) 153.—Chili.
illinitus, *Phil. l. c.* xxviii. (1856) 750.—Chili.
imbricatifolius, *Sch. Bip. ex Wedd. Chlor. And.* i. 96.—N. Granat.
imbricatus, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 421.—Bras.
immunis, Wallr. Beitr. Fl. Hercyn. 278 = paludosus.
imoleus, Boiss. & Heldr. ex Orph. in Bull. Congr. Bot. Pétersb. 1869 (1870) 128, sphalm. = *tmoleus*.
imparipinnatus, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 333 = lobatus.
inaequidens, DC. Prod. vi. 401 = pellucidus.
inaequilobus, Sch. Bip. ex Boiss. Fl. Orient. iii. 410 = paucilobus.
inamoenus, DC. Prod. vi. 382 = glutinosus, *Thunb.*
incanus, Boeber, in Pall. N. Nord. Beitr. vi. 262 = erucifolius.

SENECIO:—
incanus, Lapeyr. Hist. Abr. Pl. Pyr. 515 = leucophyllus.
incanus, *Linn. Sp. Pl.* 869.—Europ.
incanus, Scop. Fl. Carn. ed. II. ii. 166 = carniolicus.
incarnatus, Lowe, ex DC. Prod. vi. 344 = crassifolius.
incertus, *DC. l. c.* 433.—Afr. austr.
incisus, Presl, Fl. Sic. p. xxviii = squalidus.
incisus, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
incomptus, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 143 = Erechtites quadridentatus.
incomptus, *DC. Prod.* vi. 386.—Afr. austr.
inconstans, DC. l. c. 382 = repandus.
incrassatus, Guss. Fl. Sic. Syn. ii. 1. 473 = leucanthemifolius. *Poir.*
incrassatus, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 16.—Ins. Mader.
infestus, Salisb. Prod. 197 = S. Jacobaea.
inops, *Boiss. & Bal. Diagn.* Ser. II. vi. 101.—Cappadoc.
inornatus, *DC. Prod.* vi. 385.—Afr. austr.
insignis, *Hemsl. Biol. Centr. Am. Bot.* ii. 242.—Mexic.
insularis, *Benth. Fl. Austral.* iii. 666.—Austral.
intactus, *Moench, Meth.* 568 = arabicus.
integerrimus, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 238 = crassulus.
integerrimus, *Nutt. Gen. Am.* ii. 165.—Am. bor.
integrifolius, [Clairv.] Man. Herb. 241 = spathulaefolius.
integrifolius, Hook. Fl. Bor. Am. i. 334 = Hookeri.
integrifolius, Kern. in Oestr. Bot. Zeitschr. xxi. (1871) 202 = campestris?
integrifolius, Nutt. Gen. Am. ii. 165 = canus.
intermedius ×, Wiesb. in Oestr. Bot. Zeitschr. xxiv. (1874) 109.—Europ.
intermedius, Wight, Ic. t. 1135 = scandens.
involucratus, *DC. Prod.* vi. 422.—Peruv.
iodopappus, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen (jodopappus); *Wedd. Chlor. And.* i. 116. t. 20 B.—Peruv.
isatideus, *DC. l. c.* 387.—Afr. austr.
Iserni, *Phil. in Anal. Univ. Chil.* (1873) 499.—Chili.
Jacobaea, Brot. ex Willk. & Lange, Prod. Fl. Hisp. ii. 120 = foliosus.
Jacobaea, D. Don, Prod. Fl. Nep. 179 = chrysanthemoides.
Jacobaea, Kunze, in Flora, xxix. (1846) 676 = praealtus.
Jacobaea, *Linn. Sp. Pl.* 870.—Europ.; As. bor.
Jacobaea, Losc. & Pard. Ser. Inconf. 52 = jacobaeoides.
jacobaeoides, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 119.—Hispan.
jacobaeaformis, *Remy, in C. Gay, Fl. Chil.* iv. 182.—Chili.
Jacquemontianus, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 350.—Reg. Himal.
Jacquinianus, Reichb. Pl. Crit. iii. 80 = nemorensis.
jagensis, *Steud. Nom.* ed. II. ii. 561.—Chili.
jamaicensis, DC. Prod. vi. 412 = Gynoxys incana.
japonicus, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Japon.
japonicus, Thunb. Fl. Jap. 315 = Gynura pinnatifida.
jathrophoides, *Sch. Bip. l. c.* 498.—Mexic.
javanicus, *Willd. Sp. Pl.* iii. 1984.—Java.
Julieti, *Phil. in Anal. Univ. Chil.* (1873) 496.—Chili.
junceus, J. F. Gmel. Syst. 1624, sphalm. = lanceus.
junceus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 407.—Afr. austr.
Jungei, *Phil. in Linnaea*, xxxiii. (1864–65) 149.—Chili.
juniperinus, *Linn. f. Suppl.* 371.—Afr. austr.
Jurgensenii, *Hemsl. Biol. Centr. Am. Bot.* ii. 242.—Mexic.
jurineaefolius, *Boiss. & Bal. Diagn.* Ser. II. vi. 99.—As. Min.
Kaempferi, *DC. Prod.* vi. 363.—Japon.
Kalmii, Hook. & Arn. Bot. Beech. Voy. 126 = lugens.
Kalmii, Less. in Linnaea, vi. (1831) 244 = palustris.
Kalmii, Nutt. Gen. Am. ii. 165 = artemisiaefolius.
kamtschaticus, Ledeb. Fl. Ross. ii. 636 = palmatus.
kermesinus, *Hemsl. Biol. Centr. Am. Bot.* ii. 242.—Mexic.
Kickxii, *Steud. Nom.* ed. II. ii. 562.—Belgium.
Kingii, *Hook. f. Fl. Antarct.* 314.—Patagon.
kirghisicus, DC. Prod. vi. 362 = racemosus.

SENECIO:—
Kirilowii, Turcz. ex DC. Prod. vi. 361 = campestris.
Kleinia, *Less. in Linnaea*, vi. (1831) 252; *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 321.—Ins. Canar.
kleinioides, *Oliver & Hiern, in Oliver Fl. Trop. Afr.* iii. 421.—Afr. trop.
Kochii, Ledeb. Fl. Ross. ii. 643 = crispatus.
Kolenatianus, *C. A. Mey. in Beitr. Pfl. Russ. Reich.* vi. 34.—Reg. Caucas.
korintjiana, Boerl. in Veth, Midden-Sumatra, iv. II. Flora, 20 = sumatranus.
Kotschyanus, *Boiss. Diagn.* Ser. I. vi. 93.—Persia.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 406.—Japon.
Kraussii, *Sch. Bip. in Flora*, xxvii. (1844) 698.—Afr. austr.
Kunthianus, *Wall. Cat.* n. 3118.—Reg. Himal.
Kurzii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 341.—Burma.
lacerus, Boiss. Diagn. Ser. I. xi. 32 = coronopifolius.
laciniatus, Bertol. in Desv. Journ. Bot. ii. (1813) 76 = nebrodensis.
laciniatus, DC. Prod. vi. 411 = Swartzianus.
laciniatus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 470 = S. Jacobaea.
laciniatus, Hook. f. Fl. Brit. Ind. iii. 339, sphalm. = *laciniosus*, Wall.
laciniatus, *H. B. & K. Nov. Gen. et Sp.* iv. 175.—Ecuador.
laciniatus, Schousb. ex Willk. & Lange, Prod. Fl. Hisp. ii. 120 = foliosus.
lacinifolius, Steud. Nom. ed. II. ii. 562 = laciniatus.
laciniosus, Arn. in Nov. Act. Nat. Cur. xviii. 1. (1836) 350 = tenuifolius.
laciniosus, *Phil. in Linnaea*, xxviii. (1856) 744.—Chili.
laciniosus, Wall. Cat. n. 3124 = chrysanthemoides.
lactescens, Brot. Fl. Lusit. i. 391 = Crepis foetida.
laetus, Edgew. in Trans. Linn. Soc. xx. (1846) 74 = chrysanthemoides.
laevicaulis, *DC. Prod.* vi. 414.—Chili.
laevigatus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
Lagascanus, *DC. Prod.* vi. 357.—Hispan.
Laggeri, Nym. Consp. 356 = incanus.
lagopus, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 119. t. 18.—N. Zel.
lampsanoides, *DC. Prod.* vi. 348.—Reg. Caucas.
lanatus, Burm. ex DC. l. c. 389 = verbascifolius.
lanatus, *DC. l. c.* 422.—N. Granat.
lanatus, Lec. & Lamotte, Cat. Auv. 232 = Gerardi.
lanatus, Linn. Syst. ed. X. 1216 = populifolius.
lanatus, Linn. f. Suppl. 370 = Cineraria tomentosa.
lanatus, *Scop. Fl. Carn.* ed. II. ii. 165.—Europ.
lanceolatus, *Burm. f. Fl. Cap. Prod.* 26.—Afr. austr.
lanceolatus, Torr. & Gray, Fl. N. Am. ii. 440 = S. Serra.
lanceus, *Ait. Hort. Kew.* ed. I. iii. 194.—Afr. austr.
lancifolius, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 89.—Afr. austr.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864-65) 151.—Chili.
lanifer, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 312 = vernonioides.
laniger, Boj. ex DC. Prod. vi. 379 = adenodontus.
lanigerus, *Burm. f. Fl. Cap. Prod.* 26.—Afr. austr.
lanuginosus, [D.] Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 332 = Donianus.
lanuginosus, *Spreng. Syst.* iii. 565.—Hab.?
laricifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 185.—Peruv.
Laseguei, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 48.—Reg. Magellan.
lasiorhizoides, Sch. Bip. ex Harv. & Sond. Fl. Cap. iii. 370 = seq.
lasiorhizus, *DC. Prod.* vi. 387.—Afr. austr.
Lastarrianus, *Remy, in C. Gay, Fl. Chil.* iv. 146.—Chili.
latiflorus, *Wedd. Chlor. And.* i. 125.—N. Granat.
latifolius, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 145.—N. Zel.
latifolius, DC. Prod. vi. 387 = barbellatus.
latronum, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 402 = racemosus.
laurifolius, H. B. & K. Nov. Gen. et Sp. iv. 185 = Gynoxis laurifolia.
lautus, *Soland. ex Forst. f. Prod.* 91, nomen; *et ex Hook. f. Fl. N. Zel.* i. 145.—Austral.

SENECIO:—
lautus, Willd. Sp. Pl. iii. 1981 = oligocephalus?
lavandulaefolius, *Wall. Cat.* n. 3130.—Ind. or.
Lawii, *C. B. Clarke, Comp. Ind.* 201.—Ind. or.
laxiflorus, Viv. Fl. Lib. Spec. 55. t. 11. f. 3 = gallicus.
laxifolius, *J. Buch. in Trans. N. Z. Inst.* ii. (1870) 89.—N. Zel.
laxus, *DC. Prod.* vi. 380.—Afr. austr.
Layneae, Greene, in Bull. Torrey Bot. Club, x. (1883) 87 = fastigiatus.
lazicus, *Boiss. & Bal. in Boiss. Fl. Orient.* iii. 400.—As. Min.
Lechleri, *Phil. in Anal. Univ. Chil.* (1873) 494.—Reg. Magellan.
Lechleri, Sch. Bip. in Bonplandia, iv. (1856) 55, nomen = cymosus.
Ledebourii, Sch. Bip. in Flora, xxviii. (1845) 50.—Sibir.
ledifolius, *DC. Prod.* vi. 421.—N. Granat.
legionensis, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 81.—Hispan.
leiocarpus, DC. Prod. vi. 390 = albanensis.
leiophyllus, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 209.—Chili.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 220.—Arizona.
leontodontifolius, Boj. ex DC. Prod. vi. 376 = Bojeri.
Leontodontis, *DC. l. c.* 386.—Afr. austr.
leptocarpus, DC. l. c. 372 = pectinatus.
leptolobus, DC. l. c. 419 = pinnatus.
leptophyllus, *DC. l. c.* 400.—Afr. austr.
leptophyllus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 345 = valparadisiensis.
leptoschizus, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 99.—Bras.
Lessingianus, *C. B. Clarke, Comp. Ind.* 198.—Ind. or.
Lessingii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 56.—Afr. austr.
Lessoni, F. Muell. Cat. Hort. Melb. (1858) 26 = seq.
Lessonianus, Sch. Bip. in Flora, xxviii. (1845) 498 = Erechtites arguta.
Lessonianus, Steud. Nom. ed. II. ii. 562 = australis.
Leucadendron, *Benth. & Hook. f. Gen.* ii. 448.—Ins. St. Helena.
leucanthemifolius, *Phil. in Anal. Univ. Chil.* (1873) 495.—Chili.
leucanthemifolius, *Poir. Voy. Barb.* ii. 238.—Reg. Mediterr.
leuciscus, *Phil. in Anal. Univ. Chil.* (1870) II. 181.—Chili.
leucoglossus, *F. Muell. Fragm.* ii. 15.—Austral.
leucoglossus, Sond. in Harv. & Sond. Fl. Cap. iii. 393 = actinoleucus.
leucomallus, *A. Gray, in Proc. Am. Acad.* v. (1861) 141.—Terr. Magellan.
leucomallus, *Phil. in Linnaea*, xxxiii. (1864–65) 148.—Chili.
leucopappus, *Boj. ex DC. Prod.* vi. 331, in syn.—Madag.
leucophyllus, *DC. Cat. Hort. Monsp.* (1813) 144.—Mont. Pyren.
leucophyton, *Phil. in Linnaea*, xxviii. (1856) 738.—Chili.
leucostachys, *Baker, in Mart. Fl. Bras.* vi. III. 307, in obs.—Patagon.
Levingii, *C. B. Clarke, Comp. Ind.* 301.—Reg. Himal.
Leyboldi, *Phil. in Anal. Univ. Chil.* (1873) 493.—Chili.
libanoticus, Ehrenb. ex Boiss. Fl. Orient. iii. 407 = doriaeformis.
Ligularia, *Hook. f. Fl. Brit. Ind.* iii. 349.—Reg. Himal.
lilacinus, *Jacq. f. ex Spreng. f. Tent. Suppl. Syst.* 24.—Hab.?
lilacinus, *Schrad. Blumenbachia*, 39. t. 4. f. 1.—Afr. austr.
limbardioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 347.—Patagon.
limbardioides, Hook. & Arn. l. c. 333 = Arnottii.
linariaefolius, Drège, ex Harv. & Sond. Fl. Cap. iii. 400 = debilis, *Harv.*
linariaefolius, *Poepp. ex DC. Prod.* vi. 414; *Wedd. Chlor. And.* i. 133.—Chili.
Lindenii, *Sch. Bip. ex Wedd. l. c.* 101.—N. Granat.
linearifolius, Colla, in Mem. Acc. Torin. xxxviii. (1835) 31. t. 29 = argenteus.
linearifolius, A. Rich. Sert. Astrolab. 129 = australis.

SENECIO:—
linearilobus, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 99.—Bras.
linearilobus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 346 = platycodon.
linearis, *DC. Prod.* vi. 375.—Ins. Maurit.
lineatus, *DC. l. c.* 395.—Afr. austr.
lineolatus, DC. l. c. 427 = sinuatus.
Lingua, *Poir. Encyc. Suppl.* v. 130.—Ins. Maurit.
linifolius, *C. B. Clarke, Comp. Ind.* 202.—Reg. Himal.
linifolius, DC. Prod. vi. 400, in syn. = rosmarinifolius.
linifolius, Forsk. Fl. Aegypt. Arab. 150 = biflorus, *Vahl.*
linifolius, *Linn. Syst.* ed. X. 1215.—Hispan.
linifolius, Thunb. Fl. Cap. ed. Schult. 678, excl. syn. = filifolius.
linoides, *DC. Prod.* vi. 397.—Afr. austr.
littoralis, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 104.—Ins. Falkland.
littoralis, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 51 = Andersoni.
littoralis, Poepp. ex DC. Prod. vi. 417 = sinuatilobus.
littoralis, Scop. Fl. Carn. ed. II. ii. 162 = Inula viscosa.
littoreus, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
lividus, *Linn. Sp. Pl.* 867.—Reg. Mediterr.
lividus, Nolte, Novit. Fl. Holsat. 72 = sylvaticus.
lividus, Sm. Engl. Bot. t. 2515 = viscosus.
lobatus, *Pers. Syn.* ii. 436.—Am. bor.
lobelioides, *DC. Prod.* vi. 382.—Afr. austr.
longepenicillatus, Sch. Bip. ex Wedd. Chlor. And. i. 97 = nevadensis.
longidentatus, DC. Prod. vi. 428 = triangularis.
longiflorus, *Sch. Bip. in Flora*, xxviii. (1845) 499; *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 421.—Afr. trop. et austr.
longifolius, Berg. Desc. Pl. Cap. 279 = rosmarinifolius.
longifolius, Labill. ex DC. Prod. vi. 374 = australis.
longifolius, *Linn. Sp. Pl.* ed. II. 1222.—Afr. austr.
longilobus, Benth. Pl. Hartw. 18 = Douglasii.
longipes, *Hook. f. Fl. Antarct.* 314.—Patagon.
longiradiatus, Trautv. in Ind. Sem. Hort. Petrop. (1865) 37 = amphibolus, *C. Koch.*
longiscapus, *Boj. in DC. Prod.* vi. 376.—Madag.
Lopezii, *Boiss. Elench.* 60.—Hispan.
Lorentii, Hochst. in Lorent, Wanderungen, 330 = armenius.
Lorentzii, *Griseb. in Goett. Abh.* xix. (1874) 193.—Reg. Argent.
Loureiroi, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 160 = Gynura pinnatifida.
lucidus, *DC. Prod.* vi. 411.—Ind. occ.
ludens, *C. B. Clarke, Comp. Ind.* 199.—Ind. or.
lugens, *Richards. in App. Frankl. Journ.* ed. II. 31.—Am. arct.
luridus, *Phil. Fl. Atac.* 33.—Chili.
luridus, Salisb. Prod. 196 = japonicus.
Lyallii, *Hook. f. Fl. N. Zel.* i. 146.—N. Zel.
lycopersicifolius, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura lycopersicifolia.
lycopifolius, Desf. Tabl. ed. II. 124 = erucifolius.
lyratifolius, *Reichb. Pl. Crit.* ii. 24.—Europ.
lyratipartitus, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 439 = Cineraria Schimperi.
lyratus, DC. Prod. vi. 380, excl. syn. = expansus, *Harv.*
lyratus, Forsk. Fl. Aegypt. Arab. 148 = auriculatus, *Vahl.*
lyratus, *Linn. f. Suppl.* 369.—Afr. austr.
lyratus, Michx. Fl. Bor. Am. ii. 120 = lobatus.
lyratus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) II. 16 = resedifolius.
macedonicus, *Griseb. Spicil. Fl. Rumel.* ii. 221.—Graecia.
macquariensis, DC. Prod. vi. 372 = lautus.
macranthus, C. B. Clarke, Comp. Ind. 205 = japonicus.
macranthus, *A. Rich. Sert. Astrolab.* 126.—Austral.
macrobotrys, *Hemsl. Biol. Centr. Am. Bot.* ii. 243.—Mexic.
macrocephalus, *DC. Prod.* vi. 407.—Afr. austr.
macrodontus, DC. l. c. 373 = australis.
macroglossus, *DC. l. c.* 404.—Afr. austr.
macropappus, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 436.—Abyss.
macrophyllus, *Bieb. Fl. Taur. Cauc.* ii. 308.—Europ. or.; Reg. Caucas.

SENECIO:—
macrophyllus, H. B. & K. Nov. Gen. et Sp. iv. 178 = bogotensis.
macropodus, DC. Prod. vi. 403 = angulatus.
macrorrhizus, *Wedd. Chlor. And.* i. 108.—Peruv.
macrospermus, *DC. Prod.* vi. 385.—Afr. austr.
macrotis, *Baker, in Mart. Fl. Bras.* vi. III. 308.—Bras.
macrotus, *Kunze, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 281.—Chili.
madagascariensis, *Poir. Encyc. Suppl.* v. 130.—Madag.
maderensis, *DC. Prod.* vi. 409.—Ins. Madera.
madrensis, *A. Gray in Proc. Am. Acad.* xix. (1883) 55.—Mexic.
magellanicus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 343 = Culcitium magellanicum.
magellensis, *Tenore, Ind. Sem. Hort. Neap.* (1857).—*Cf. Ann. Sc. Nat.* Sér. IV. xii. (1859) 78.—Ital.
magnificus, *F. Muell. in Linnaea*, xxv. (1852) 418.—Austral.
Mairetianus, *DC. Prod.* vi. 430.—Mexic.
maldonadensis, *Baker, in Mart. Fl. Bras.* vi. III. 312.—Bras.; Uruguay.
malvaefolius, *DC. Prod.* vi. 410.—Ins. Azor.
Mandonianus, *Wedd. Chlor. And.* i. 228.—Bolivia.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 14.—Afr. trop.
marginatus, *Linn. f. Suppl.* 370.—Afr. austr.
marginatus, Thunb. Fl. Cap. ed. Schult. 680, partim = lanceus.
maritimus, *Linn. f. Suppl.* 369.—Afr. austr.
maritimus, Reichb. in Moessl. Handb. ed. II. ii. 1479 = S. Cineraria.
Marmorae, Moris, Stirp. Sard. Elench. ii. 6 = crassifolius.
Mathewsii, *Wedd. Chlor. And.* i. 114.—Peruv.
matricariaefolius, *DC. Prod.* vi. 383.—Afr. austr.
mauritanicus, Pomel, Nouv. Mat. Fl. Atl. 62 = leucanthemifolius, *Poir.*
mauritianus, *Soyer-Willem. in Usteri, Ann. Bot.* xviii. (1796) 52.—Ins. Maurit.
medicinalis, *Phil. in Linnaea*, xxviii. (1856) 745.—Chili.
medius, *G. Don, in Sweet, Hort. Brit.* ed. III. 386.—Hungar.
medullosus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
megacephalus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 410.—Am. bor. occ.
megaglossus, *F. Muell. in Linnaea*, xxv. (1852) 419.—Austral.
megalophron, Fenzl, ex Boiss. Fl. Orient. iii. 407 = doriaeformis.
megapotamicus, Buek, Ind. ad DC. Prod. ii. p. vi = brasiliensis.
megapotamicus, Spreng. Syst. iii. 547 = pinnatus.
melanolepis, *DC. Prod.* vi. 424.—Peruv.
membranifolius, DC. l. c. 381 = repandus.
mendocinensis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 362.—Calif.
mendocinus, *Phil. Sert. Mendoc. Alt.* 24.—Chili.
mesembryanthemoides, *Boj. ex DC. Prod.* vi. 378.—Madag.
Meyeri, *Regel, Ind. Sem. Hort. Petrop.* (1856) 25.—Austral.
micaniaeformis, Steud. Nom. ed. II. ii. 562, sphalm. = *mikaniaeformis.*
micranthus, *Hornem. Hort. Hafn. Suppl.* 97.—Hab. ?
Microchaeta, *Wedd. Chlor. And.* i. 100.—N. Granat.
microdon, *Wedd. Chlor. And.* i. 102.—Ecuador.
microdontus, *Baker, in Journ. Linn. Soc.* xviii. (1881) 271.—Madag.
microglossus, *DC. Prod.* vi. 395.—Afr. austr.
microphyllus, *Phil. in Linnaea*, xxix. (1857–58) 1.—Chili.
microphyllus, Spreng. Syst. iii. 556, sphalm. = macrophyllus.
micropifolius, *DC. Prod.* vi. 413.—Chili.
microspermus, *DC. l. c.* 391.—Afr. austr.
Mikaniae, DC. l. c. 405 = deltoideus.
mikaniaefolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Peruv.
mikaniaeformis, DC. Prod. vi. 405 = deltoideus.
mikanioides, *Otto, ex Walp. in Otto & Dietr. Allg. Gartens.* xiii. (1845) 42.—Afr. austr.
Millefolium, *Torr. & Gray, Fl. N. Am.* ii. 444.—Am. bor.
minimus, Poir. Encyc. Suppl. v. 130 = Erechtites minima.

SENECIO:—
minutus, *DC. Prod.* vi. 346.—Hispan.
miser, *Hook. f. Fl. Antarct.* 314.—Reg. Magell.
Mishmi, *C. B. Clarke, Comp. Ind.* 203.—Reg. Himal.
mississipianus, DC. Prod. vi. 427 = lobatus.
mixtus, A. Rich. Sert. Astrolab. 112. t. 36 = Erechtites mixta.
modestus, *Phil. in Linnaea*, xxviii. (1856) 745.—Chili.
modestus, *Wedd. Chlor. And.* i. 105.—Peruv.
moharensis, *A. Gray, Syn. Fl. N. Am.* i. II. 446.—Calif.
Molinae, *Phil. in Anal. Univ. Chil.* (1873) 497.—Chili.
mollis, *DC. Prod.* vi. 358 = cappadocicus.
mollis, Poepp. ex DC. l. c. 417 = sinuatilobus.
mollis, *Willd. Sp. Pl.* iii. 2011.—As. Min.; Persia.
moluccanus, Roxb. Hort. Beng. [101]; Fl. Ind. iii. 433 = Gynura bicolor ?
mongolicus, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Mongolia.
monocephalus, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 107; iv. (1853) 41 = abrotanifolius.
Monroi, *Hook. f. Fl. N. Zel.* ii. 333.—N. Zel.
montanus, Hoppe, ex Nym. Consp. 356 = nebrodensis.
montanus, Lam. Fl. Fr. ii. 130 = S. Doronicum.
montanus, Willd. Sp. Pl. iii. 1989 = nebrodensis.
montevidensis, *Baker, in Mart. Fl. Bras.* vi. III. 307.—Uruguay.
monticola, *DC. Prod.* vi. 384.—Afr. austr.
Monttianus, *Remy, in C. Gay, Fl. Chil.* iv. 158.—Chili.
Moreliae, *Hemsl. Biol. Centr. Am. Bot.* ii. 243.—Mexic.
Morettii, *Buek, Ind. ad DC. Prod.* ii. p. vi.—Hab. ?
Mortoni, *C. B. Clarke, Comp. Ind.* 208.—Reg. Himal.
mucronatus, *Willd. Sp. Pl.* iii. 1974.—Afr. austr.
mucronulatus, Sch. Bip. in Flora, xxvii. (1844) 701 = purpureus.
Muehlenbergii, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Am. bor.
Muelleri, *T. Kirk, in Trans. N. Z. Inst.* xv. (1883) 359.—N. Zel.
Muelleri, *Regel, Ind. Sem. Hort. Petrop.* (1863) 31.—Austral.
mulgediifolius, *Schau. in Linnaea*, xix. (1847) 733.—Mexic.
multangularis, Tausch, in Syll. Ratisb. ii. (1828) 253 = nebrodensis.
multibracteatus, *Baker, in Journ. Linn. Soc.* xx. (1883) 192.—Madag.
multibracteatus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 361.—Afr. austr.
multicaulis, *DC. Prod.* vi. 398.—Afr. austr.
multicaulis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 58.—Chili.
multicaulis, A. Rich. Sert. Astrolab. 105 = Erechtites arguta.
multidentatus, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 243.—Mexic.
multifidus, Domb. ex DC. Prod. vi. 416 = glaber.
multifidus, Willd. Sp. Pl. iii. 1986 = tenuifolius.
multiflorus, *DC. Prod.* vi. 410.—Ins. Canar.
multiflorus, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 325 (an praec. ?).—Ins. Canar.
multifoliosus, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 375.—Ins. Borbon.
multilobatus, *Torr. & Gray, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 109.—Am. bor. occ.
multinervis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
multivenius, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 109.—Costa Rica.
munitus, Wallr. Beitr. Fl. Hercyn. 278 = paludosus.
muricatus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
murorum, *Remy, in C. Gay, Fl. Chil.* iv. 149.—Chili.
myosotidifolius, Benth. in Enum. Pl. Hueg. 66 = Millotia tenuifolia.
myrianthos, *Cheesm. in Trans. N. Z. Inst.* vii. (1875) 348.—N. Zel.
myriocephalus, *Baker, in Mart. Fl. Bras.* vi. III. 319.—Bras.
myriocephalus, *Sch. Bip. in A. Rich. Tent. Fl. Abyss.* i. 441.—Abyss.
myriophyllus, *Phil. Fl. Atac.* 33.—Chili.
myrrhifolius, Thunb. Prod. Pl. Cap. 159 = arenarius.

SENECIO:—
Myrsinites, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 211.—Ecuador.
nanus, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 443 = Werneria africana.
napeaefolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
napellifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 244.—Mexic.
napifolius, Schrad. ex Reichb. Fl. Germ. Excurs. 243 = erucifolius.
natalensis, *Sch. Bip. in Flora*, xxvii. (1844) 700.—Afr. austr.
Neaei, *DC. Prod.* vi. 413.—Chili.
nebrodensis, DC. Fl. Fr. iv. 162 = lividus.
nebrodensis, Gilib. ex Georgi, Beschr. Russ. Reich. III. iv. 1242 = vernalis.
nebrodensis, *Linn. Sp. Pl.* ed. II. 1217.—Europ. austr.; Afr. bor.
neelgherryanus, *DC. in Wight, Contrib.* 23.—Ind. or.
neglectus, Desv. Obs. Pl. Andeg. 129 = S. Jacobaea.
neglectus, A. Rich. Fl. N. Zél. 258 = lautus.
nemophilus, Dum. Fl. Belg. Prod. 66 = Fuchsii.
nemorensis, *Linn. Sp. Pl.* 870.—Europ. austr.; As. bor.
nemorensis, Lorey & Duret, Fl. Côte-d'Or, i. 474 = Fuchsii.
nemorosus, Jord. Cat. Jard. Dijon (1848), 30 = S. Jacobaea.
neo-mexicanus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 55.—Am. bor. occ.
nepalensis, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura nepalensis.
neriifolius, Baill. Hist. Pl. viii. 308 = S. Kleinia.
nervatus, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 302.—Afr. austr.
nevadensis, Boiss. & Reut. Pugill. Pl. Nov. 60 = linifolius, *Linn.*
nevadensis, *Sch. Bip. ex Wedd. Chlor. And.* i. 97.—Venezuela.
nigrescens, *Hook. & Arn. Bot. Beech. Voy.* 32.—Chili.
nikoënsis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 182.—Japon.
nilagerensis, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 337 = Notonia Walkeri.
nilgheryanus, Hook. f. Fl. Brit. Ind iii. 341 = neelgherryanus.
nitidus, *DC. Prod.* vi. 421.—Ecuador.
nitidus, Less. ex Harv. & Sond. Fl. Cap. iii. 379 = isatidens.
nitidus, Mérat, Fl. Par. Addit. (1845) 495 = S. Jacobaea.
nitidus, *Phil. in Linnaea*, xxviii. (1856) 742.—Chili.
niveus, Wall. Cat. n. 3143 = Gerbera nivea.
niveus, *Willd. Sp. Pl.* iii. 1974.—Afr. austr.
Noëanus, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xiv. (1856) 231.—Mesopot.?
nubigenus, *H. B. & K. Nov. Gen. et Sp.* iv. 174.—Ecuador.
nudicaulis, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 179.—Reg. Himal.
nudicaulis, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura Pseudo-china.
nudiusculus, *DC. Prod.* vi. 384.—Afr. austr.
nutans, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
Nuttallii, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Am. bor.
oaxacanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 244.—Mexic.
obliquus, Dulac, Fl. Hautes-Pyr. 507 = S. Cacaliaster.
oblongifolius, *Steud. Nom.* ed. II. ii. 563.—Chili.
obovatus, Muhl. ex Willd. Sp. Pl. iii. 1999 = aureus.
obscurus, *Fisch. ex Ledeb. Fl. Ross.* ii. 638.—Sibir.
obscurus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) ii. 20 = nemorensis.
obtusatus, Banks, ex Pursh, Fl. Am. Sept. ii. 530 = aureus.
obtusatus, *Wall. Cat.* n. 3133.—Reg. Himal.; Burma.
ochrocarpus, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 416.—Afr. trop.
ochroleucus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 340 = rhodaster, fistulosus.
octoglossus, DC. Prod. vi. 354 = nemorensis.
octolepis, *Griseb. in Goett. Abh.* xxiv. (1879) 204.—Reg. Argent.
octophyllus, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 581, nomen.—Bolivia.

SENECIO:—
odontopterus, *DC. Prod.* vi. 406.—Afr. austr.
odoratus, *Hornem. Hort. Hafn.* ii. 809.—Austral.
odorus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Arab.
oederiaefolius, *DC. Prod.* vi. 391.—Afr. austr.
Oerstedianus, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 109.—Costa Rica.
Oldhamianus, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 219.—China.
oleaefolius, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 305 = trixoides.
oleosus, Vell. Fl. Flum. viii. t. 104 = Selloi.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 45.—Turkestan.
oliganthus, *DC. Prod.* vi. 390.—Afr. austr.
oliganthus, Spreng. Syst. v. 636 (Index) = S. Vulneraria.
oligocephalus, *DC. Prod.* vi. 373.—Austral.
oligolepis, *Boiss. Diagn.* Ser. I. vi. 94.—Persia.
oligoleucus, *Baker, in Mart. Fl. Bras.* vi. iii. 311.—Reg. Argent.
oligophyllus, *Baker, l. c.* 317.—Bras.
olympicus, *Boiss. Diagn.* Ser. I. iv. 13.—As. Min.
oporinus, Willd. Sp. Pl. iii. 2008 = lanceus.
opuntiaefolius, *DC. Prod.* vi. 435 (sp. dub.).—Am. bor.
orbicularis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 376.—Afr. austr.
oreophyton, *Remy, in C. Gay, Fl. Chil.* iv. 158.—Chili.
organensis, *Casar. Nov. Stirp. Bras. Dec.* 77; *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 127.—Bras.
orientalis, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 221 = macedonicus.
orientalis, Mill. Gard. Dict. ed. VIII. n. 10 = coriaceus.
orientalis, Pacho, Voy. Marm. t. 97 = macedonicus?
orientalis, Tenore, Fl. Nap. ii. 222 = S. Doronicum.
orientalis, *Willd. Sp. Pl.* iii. 2006.—As. Min.; Persia.
orizabensis, *Sch Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 244.—Mexic.
oronocensis, *DC. Prod.* vi. 423.—Venezuel.
Othonnae, *Bieb. Fl. Taur. Cauc.* ii. 308.—Serbia; Reg. Caucas.; Persia.
othonnaeflorus, *DC. Prod.* vi. 386.—Afr. austr.
othonnaeoides, A. Cunn. ex DC. l. c. v. 398 = Minuria Cunninghamii.
otiphyllus, A. Cunn. ex DC. l. c. vi. 371 = odoratus.
Otites, *Kunze, ex DC. l. c.* 417.—Chili.
otophorus, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 219.—Japon.
otophorus, *Wedd. Chlor. And.* i. 98.—N. Granat.
otopterus, *Griseb. in Goett. Abh.* xix. (1874) 192.—Reg. Argent.
ovalifolius, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 208.—Chili.
ovalis, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura auriculata.
ovatifolius, *Boiss. Fl. Orient.* iii. 406.—Armenia.
ovatifolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
ovatus, Willd. Sp. Pl. iii. 2004 = Fuchsii.
ovirensis, DC. Prod. vi. 360 = alpestris.
oxyodontus, *DC. l. c.* 392.—Afr. austr.
oxyphyllus, *DC. l. c.* 419.—Bras.
oxyriaefolius, *DC. l. c.* 405.—Afr. austr.
paarlensis, *DC. l. c.* 383.—Afr. austr.
pachycarpus, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 349.—Reg. Himal.
pachyphyllos, *Remy, in C. Gay, Fl. Chil.* iv. 147.—Chili.
pachyphyllus, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 410.—N. Zel.
Pallasii, Sch. Bip. in Flora, xxviii. (1845) 498 = Doronicum altaicum.
pallens, Boj. ex DC. Prod. vi. 377 = emirnensis.
pallens, Wall. Cat. n. 3132 = nudicaulis.
pallidus, *Vell. Fl. Flum.* viii. t. 105.—Bras.
palmatus, Lapeyr. Suppl. Hist. Pl. Pyr. 134 = leucophyllus.
palmatus, Less. Syn. Comp. 392 = S. Syneilesis.
palmatus, *Pall. Reise*, iii. 321.—Sibir.; Ins. Sachalin.
palmensis, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 148.—Ins. Canar.
Palmeri, *A. Gray, in Proc. Am. Acad.* xi. (1876) 80.—Calif.
paludicola, *Steud. Nom.* ed. II. ii. 563.—Bras.
paludosus, *Linn. Sp. Pl.* 870.—Europ.; As. bor.
palustris, *Hook. Fl. Bor. Am.* i. 334.—Reg. bor. temp.
palustris, Vell. Fl. Flum. viii. t. 110 = paludicola.

SENECIO:—
pamplonensis, Sch. Bip. ex Wedd. Chlor. And. i. 94 = ledifolius.
panduratus, DC. Prod. vi. 383 = hastulatus.
panduratus, Less. Syn. Comp. 392 = erosus, *Linn. f.*
pandurifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 380.—Afr. austr.
pandurifolius, *C. Koch, in Linnaea*, xvii. (1843) 48.—Mingrelia.
paniculatus, *Berg. Desc. Pl. Cap.* 227.—Afr. austr.
paniculatus, Drège, ex Harv. & Sond. Fl. Cap. iii. 400 = debilis, *Harv.*
papaverifolius, A. Rich. Tent. Fl. Abyss. i. 437 = picridifolius.
papillaris, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
papillosus, *F. Mueil. in Hook. Kew Journ.* ix. (1857) 301.—Tasman.
paposanus, *Phil. Fl. Atac.* 32.—Chili.
papposus, Less. in Linnaea, vi. (1831) 244 = lugens.
papyraceus, *DC. Prod.* vi. 410.—Ins. Canar.
paradoxus, Hoppe, ex Willd. Sp. Pl. iii. 1989 = nebrodensis.
Paralias, *Phil. in Linnaea*, xxviii. (1856) 750.—Chili.
parasiticus, *Hemsl. Biol. Centr. Am. Bot.* ii. 244.—Mexic.
Parkeri, *Baker, in Journ. Linn. Soc.* xx. (1883) 191.—Madag.
Parnassi, *Boiss. & Heldr. in Boiss. Fl. Orient.* iii. 412.—Graec.
Parryi, *A. Gray, in Torr. Bot. Mex. Bound.* 103.—Am. bor. occ.
parviflorus, All. Fl. Pedem. i. 200. t. 38. f. 3 = incanus.
parvifolius, *DC. Prod.* vi. 396.—Afr. austr.
patagonicus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 344.—Patagon.
patagonicus, *Phil. in Linnaea*, xxxiii. (1864–65) 159.—Patagon.
patens, *DC. Prod.* vi. 423.—Ecuador.
patulus, Burm. ex DC. l. c. 404 = deltoideus.
paucidentatus, *DC. l. c.* 414.—Chili.
pauciflorus, *Baker, in Journ. Bot.* xx. (1882) 171.—Madag.
pauciflorus, H. B. & K. Nov. Gen. et Sp. iv. 176. t. 365 = S. Vulneraria.
pauciflorus, Pursh, Fl. Am. Sept. ii. 529 = aureus.
pauciflorus, Thunb. ex Harv. & Sond. Fl. Cap. iii. 392 = grandiflorus.
paucifolius, *DC. Prod.* vi. 403.—Afr. austr.
paucifolius, S. G. Gmel. It. i. 171 = S. Doria.
paucijugus, *Baker, in Mart. Fl. Bras.* vi. iii. 323.—Bras.
pauciligulatus, A. Rich. Sert. Astrolab. 121. t. 38 = australis.
paucilobus, *DC. Prod.* vi. 352.—Persia.
paulensis, *Bong. in Bull. Sc. Acad. Pétersb.* v. (1838) 98.—Bras.
pauperculus, Michx. Fl. Bor. Am. ii. 120 = aureus.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864–65) 157.—Chili.
pectinatus, *DC. Prod.* vi. 372.—Austral.
pectinatus, D. Don, ex Baker, in Mart. Fl. Bras. vi. iii. 323 = pinnatus.
pedicularifolius, DC. ex Steud. Nom. ed. II. ii. 563 = spiraeifolius.
pedicularifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 173.—Ecuador.
pedunculatus, Griseb. Spicil. Fl. Rumel. ii. 224 = vernalis.
pedunculatus, *Edgew. in Trans. Linn. Soc.* xx. (1846) 74.—Reg. Himal.
pedunculatus, Sch. Bip. in Flora, xxvii. (1844) 702 = polymorphus.
pedunculosus, *Trautv. in Act. Hort. Petrop.* iii. ii. (1875) 277.—Armen.
peguanus, *DC. Prod.* vi. 365.—Burma.
pellitus, *A. Gray, in Proc. Am. Acad.* v. (1861) 143.—Peruv.
pellucidinervis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. iii. 319.—Bras.
pellucidus, *DC. Prod.* vi. 380.—Afr. austr.
peltatus, DC. l. c. 405 = oxyriaefolius.
peltiferus, *Hemsl. Biol. Centr. Am. Bot.* ii. 245.—Mexic.
peltiformis, DC. Prod. vi. 405 = oxyriaefolius.
pendulus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Arab.

SENECIO:—
penicillatus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Ins. Borbon.
penninervius, *DC. Prod.* vi. 395.—Afr. austr.
pentaphyllus, *Phil. in Linnaea*, xxviii. (1856) 744.—Chili.
Pentlandianus, *DC. Prod.* vi. 421.—Bolivia.
perdicioides, *Hook. f. Fl. N. Zel.* i. 149.—N. Zel.
peregrinus, *Griseb. in Goett. Abh.* xxiv. (1879) 205.—Reg. Argent.
Perralderianus, Coss. in Bull. Soc. Bot. Fr. ix. (1862) 173 = atlanticus.
Perrottetianus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 329.—Ind. or.
Perrottetii, *DC. Prod.* vi. 343.—Afr. trop.
persicaefolius, Ram. in Bull. Soc. Philom. ii. (1799–1801) 146. t. 11. f. 3 = Tournefortii.
persicaefolius, A. Rich. Sert. Astrolab. 123 = australis.
persicifolius, Burm. ex DC. Prod. vi. 395 = lineatus.
persicifolius, *Linn. Pl. Afr. Rar.* 20; *Amoen. Acad.* vi. 100.—Afr. austr.
persicus, *Boiss. Diagn.* Ser. I. vi. 94.—Persia.
persicus, Sch. Bip. in Flora, xxviii. (1845) 50 = taraxacifolius.
Persoonii, De Not. Repert. Fl. Ligust. 229 = incanus.
peruvianus, *Pers. Syn.* ii. 436.—Peruv.
Petasitis, *DC. Prod.* vi. 431.—Am. austr.
petiolaris, *DC. l. c.* 434.—Afr. austr.
petiolaris, Less. ex Harv. & Sond. Fl. Cap. iii. 357 = Lessingii.
Petitianus, *A. Rich. Tent. Fl. Abyss.* i. 442.—Afr. trop.
petraeus, *Boiss. & Reut. Pugill. Pl. Nov.* 59.—Hispan.
petraeus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 330.—Am. bor. occ.
peucedanifolius, Linn. f. Suppl. 372 = paniculatus.
phagnaloides, *DC. Prod.* vi. 415.—Chili.
phalacrocarpus, *Hance, in Journ. Bot.* xix. (1881) 151.—China.
phalacrolaenus, DC. Prod. vi. 393 = incisus.
Philippicus, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1857) 43.—Chili.
Philippii, *Sch. Bip. ex Wedd. Chlor. And.* i. 134.—Chili.
phylicaefolius, *Poepp. ex DC. Prod.* vi. 415.—Chili.
phyllomorphus, Steud. Nom. ed. II. ii. 564 = pilquensis.
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 142.—Peruv.
picridifolius, *DC. Prod.* vi. 386.—Afr. austr.
Picridis, *S. Schau. in Linnaea*, xix. (1847) 733.—Mexic.
Pierotii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 182 = campestris.
pilosiusculus, C. B. Clarke, Comp. Ind. 203 = saxatilis.
pilquensis, *Buek, Ind. ad DC. Prod.* ii. p. vi.—Chili.
pimpinellaefolius, *H. B. & K. Nov. Gen. et Sp.* iv. 174.—Am. austr.
pinceus, *Sch. Bip. in Flora*, xxvii. (1844) 702.—Afr. austr.
pinetorum, *Hemsl. Biol. Centr. Am. Bot.* ii. 245.—Mexic.
pinguiculus, Pomel, Nouv. Mat. Fl. Atl. 62 = crassifolius.
pinguifolius, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
pinifolius, *Lam. Encyc.* iii. 263; *Spreng. Syst.* iii. 554.—Afr. austr.
pinifolius, Pers. Syn. ii. 436 = Faujasia pinifolia.
pinifolius, Willd. ex Wedd. Chlor. And. i. 101 = abietinus.
pinnatifidus, C. B. Clarke, Comp. Ind. 198 = Hohenackeri, *Hook. f.*
pinnatifidus, DC. Prod. vi. 397, excl. syn. = foeniculoides.
pinnatifidus, *Less. Syn. Comp.* 391.—Afr. austr.
pinnatifolius, A. Rich. Sert. Astrolab. 117 = lautus.
pinnatilobatus, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
pinnatipartitus, *Sch. Bip. ex Oliver, Fl. Trop. Afr.* iii. 412.—Abyss.
pinnatisectus, *DC. Prod.* vi. 427.—Mexic.
pinnatulus, Steud. Nom. ed. II. ii. 564 = Burchelii.
pinnatulus, Steud. l. c. = pinnulatus.
pinnatus, *Poir. Encyc. Suppl.* v. 131.—Bras.
pinnatus, Sch. Bip. in Flora, xxviii. (1845) 499 = Othonnae.

SENECIO:—
pinnulatus, Sieber, ex Harv. & Sond. Pl. Cap. iii. 405 = crassiusculus.
pinnulatus, *Thunb. Prod. Pl. Cap.* 157.—Afr. austr.
pisidicus, Boiss. Diagn. Ser. II. iii. 35 = Castagneanus.
Pissisi, *Phil. in Linnaea*, xxxiii. (1864-65) 151.—Chili.
Pithosillum, *DC. Prod.* vi. 375.—Ins. Maurit.
Placillae, *Steud. Nom.* ed. II. ii. 564.—Chili.
plantagineus, *Bert. ex Colla, in Mem. Acc. Torin.* xxxviii. (1835) 32.—Chili.
platanifolius, *Benth. Pl. Hartw.* 43.—Mexic.
plattensis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 413 = aureus.
platycodon, *Baker, in Mart. Fl. Bras.* vi. III. 307.—Reg. Argent.; Chili.
platylepis, *DC. Prod.* vi. 371.—Austral.
platyphyllus, *DC. l. c.* 352.—Reg. Caucas.
plebeius, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 142 = Erechtites arguta.
plebeius, DC. Prod. vi. 379 = laevigatus.
plumbeus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 515.—Cuba.
Poeppigii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 342.—Chili.
Pohlii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 303.—Bras.
polifolius, Linn. Syst. ed. XIII. 632 = populifolius.
pollicaris, DC. ex Boj. Hort. Maurit. 188 = Faujasia flexuosa.
polyanthemoides, Sch. Bip. in Flora, xxvii. (1844) 697 = pterophorus.
polyanthemus, DC. Prod. vi. 391 = juniperinus.
polycephalus, *C. B. Clarke, Comp. Ind.* 197.—Ind. or.
polycephalus, Ledeb. Fl. Ross. ii. 630 = vernalis.
polylepis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 536, nomen.—Bolivia.
polymerus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 332.—Venezuel.
polymorphus, *Sch. Bip. in Flora*, xxvii. (1844) 702.—Afr. austr.
polyodon, *DC. Prod.* iv. 386.—Afr. austr.
polyphlebius, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 515.—Cuba.
polyphyllus, *Kunze, ex DC. Prod.* vi. 415.—Chili.
polyphyllus, Sch. Bip. ex Wedd. Chlor. And. i. 119 = Schultzii, *Wedd.*
polyrhizus, *Baker, in Journ. Linn. Soc.* xx. (1883) 192.—Madag.
populifolius, Burm. ex DC. Prod. vi. 389 = verbascifolius.
populifolius, *DC. l. c.* 409.—Ins. Canar.
populifolius, Linn. Sp. Pl. ed. II. 1224 = halimifolius.
porophylloides, *Remy, in C. Gay, Fl. Chil.* iv. 154.—Chili.
Portalesianus, *Remy, l. c.* 175.—Chili.
portulacoides, *Remy, l. c.* 163.—Chili.
potamogetonifolius, *Sch. Bip. ex Phil. f. Cat. Pl. Vasc. Chil.* 183, nomen.—Chili.
potosianus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 331.—Bolivia.
Pottsii, *Armstr. in Trans. N. Z. Inst.* iv. (1872) 290.—N. Zel.
praealtus, *Bertol. in Opusc. Sc. Bologn.* iii. (1819) 183. t. 7.—Europ.
praecox, *DC. Prod.* vi. 431.—Mexic.
praeruptorum, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 530.—Bolivia.
pratensis, DC. Prod. vi. 360 = campestris.
pratensis, *Phil. in Linnaea*, xxx. (1859–60) 194.—Chili.
pratensis, Richt. in Klett & Richt. Fl. Leipz. 692 = aquaticus.
pratensis, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 61 = S. Jacobaea.
prenanthiflorus, *Benth. & Hook. f. Gen.* ii. 448.—Ins. S. Helena.
prenanthifolius, *Phil. in Linnaea*, xxxiii. (1864–65) 159.—Chili.
prenanthoides, A. Rich. Sert. Astrolab. 96 = Erechtites prenanthoides.
primulifolius, *F. Muell. in Hook. Kew Journ.* ix. (1857) 300.—Tasmania.
prionites, *MacOwan, in Journ. Linn. Soc.* xviii. (1881 392.—Afr. austr.
procerus, *Griseb. Spicil. Fl. Rumel.* ii. 219.—Serbia: Graec.

SENECIO:—
procerus, Salisb. Prod. 197 = sarracenicus.
procumbens, *H. B. & K. Nov. Gen. et Sp.* iv. 17.—Mexic.
proteus, *Remy, in C. Gay, Fl. Chil.* iv. 140.—Chili.
prunifolius, *Wedd. Chlor. And.* i. 102.—N. Granat.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 493.—Mongolia.
psammophilus, *Griseb. in Goett. Abh.* xix. (1874) 189.—Reg. Argent.
Pseudo-arnica, *Less. in Linnaea*, vi. (1831) 240.—Am. arct.; Sibir.
Pseudo-china, Andr. Bot. Rep. t. 291 = speciosus.
Pseudo-china, Linn. Sp. Pl. 867 = Gynura Pseudo-china.
Pseudo-discoideus, *Sch. Bip. ex Phil. f. Cat. Pl. Vasc. Chil.* 183, nomen.—Chili.
Pseudo-doronicum, Schur, Enum. Pl. Transs. 351 = S. Doronicum.
Pseudo-elegans, Less. Syn. Comp. 391 = elegans.
Pseudo-longifolius, Sch. Bip. in Flora, xxvii. (1844) 699 = longifolius.
Pseudo-sceptrum, Steud. Nom. ed. II. ii. 564 = spiraeifolius.
Pseud-otites, *Griseb. in Goett. Abh.* xix. (1874) 191.—Reg. Argent.
Pseudo-vernalis, Zabel, ex Nym. Consp. 357 = vernalis.
psychrophilus, *Wedd. Chlor. And.* i. 112.—Boliv.
ptarmicaefolius, *Bory, Voy.* iii. 110.—Ins. Borbon.
pterocarpus, Steud. Nom. ed. II. ii. 564, sphalm. = pterophorus.
pteroneurus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Marocc.
pterophorus, *DC. Prod.* vi. 389.—Afr. austr.
pteropogon, *Griseb. in Goett. Abh.* xxiv. (1879) 207.—Reg. Argent.
puberulus, *DC. Prod.* vi. 380.—Afr. austr.
pubigerus, *Linn. Pl. Afr. Rar.* 21; *Amoen. Acad.* vi. 101.—Afr. austr.
pugioniformis, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Hab.?
pulchellus, *DC. Prod.* vi. 421.—N. Granat.
pulchellus, H. B. & K. Nov. Gen. et Sp. iv. 187 = Gynoxys pulchella.
pulcher, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 337.—Bras.; Uruguay.
pulicarioides, *Baker, in Mart. Fl. Bras.* vi. III. 312.—Bras.
pullus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 333.—Mexic.
pumilus, Poir. Encyc. Suppl. v. 130 = Erechtites arguta?
punduahensis, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura angulosa.
pungens, *DC. Prod.* vi. 421.—Peruv.
purpuratus, *Phil. in Linnaea*, xxviii. (1856) 741.—Chili.
purpureo-viridis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 419.—Madag.
purpureus, *Linn. Syst.* ed. X. 1214.—Afr. austr.
purpureus, Soyer-Willem. in Usteri, Ann. Bot. xviii. (1796) 51 = Gynura cernua.
Purshianus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 412 = canus.
pusillus, Guss. ex DC. Prod. vi. 341 = pygmaeus.
pusillus, A. Rich. Sert. Astrolab. 99 = Erechtites arguta.
pycnanthus, *Phil. in Linnaea*, xxviii. (1856) 740.—Chili.
pygmaeus, *DC. Prod.* vi. 341.—Sicil.
pyramidatus, *DC. l. c.* 402.—Afr. austr.
pyrenaicus, *Gren. & Godr. Fl. Fr.* ii. 124.—Mont. Pyren.
pyrifolius, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 318 = ellipticus.
pyroglossus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 386 = campestris.
pyrophilus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 266.—Java.
pyrrhopappus, Sch. Bip. in Flora, xxviii. (1845) 50 = amplexicaulis, *Wall.*
quadridentatus, Labill. Nov. Holl. Pl. ii. 48. t. 194 = Erechtites quadridentata.
Quartinianus, Aschers. in Schweinf. Beitr. Fl. Aethiop. 157 = abyssinicus.
quercifolius, Pourr. ex Steud. Nom. ed. I. 765 = aquaticus.
quercifolius, Thunb. Prod. Pl. Cap. 158 = ilicifolius, *Thunb.*

SENECIO:—
quillotensis, *Phil. in Linnaea*, xxix. (1857–58) 3.—Chili.
quinquangulatus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
quinqueflorus, DC. Prod. vi. 396 = bipinnatus.
quinqueligulatus, *Winkl. in Del. Sem. Hort. Petrop.* (1881) 15.—Turkestan.
quinquelobus, *DC. Prod.* vi. 404.—Afr. austr.
quinquelobus, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 209.—Reg. Himal.
quinquenervius, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 391.—Afr. austr.
quinqueradiatus, *Boiss. ex DC. Prod.* vii. 300.—Hispan.
Rabani, C. B. Clarke, Comp. Ind. 202 = saxatilis.
racemosus, *DC. Prod.* vi. 358.—Rossia; Reg. Caucas.; As. bor.
racemosus, Wall. Cat. n. 3140 = S. Ligularia.
radicans, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
radiolatus, F. Muell. Enum. Pl. Chath. Isl. 24. t. 4 = lautus.
radulaefolius, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
ramentaceus, *Baker, in Mart. Fl. Bras.* vi. III. 305.—Bras.
ramosissimus, *DC. Prod.* vi. 371.—Austral.
ramosus, *Wall. Cat.* n. 3129.—Reg. Himal.
rancaguensis, *Remy, in C. Gay, Fl. Chil.* iv. 169.—Chili.
rancoënsis, *Sch. Bip. ex Phil. f. Cat. Pl. Vasc. Chil.* 183, nomen.—Chili.
raphanifolius, Wall. Cat. n. 3125 = diversifolius, *Wall.*
rapifolius, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 409.—Am. bor.
rapistroides, DC. Prod. vi. 346 = vernalis.
rariflorus, C. Koch, in Linnaea, xvii. (1843) 47 = Othonnae.
reclinatus, Linn. f. Suppl. 369 = paniculatus.
recurvatus, *H. B. & K. Nov. Gen. et Sp.* iv. 184.—Peruv.
Reedi, *Phil. in Anal. Univ.* (1870) II. 182.—Chili.
reflexus, *H. B. & K. Nov. Gen. et Sp.* iv. 182.—Ecuador.
regiomontanus, DC. Prod. vi. 429 = Douglasii.
regularis, Sch. Bip. in Flora, xxvii. (1844) 695 = littoreus.
Reinoldi, Endl. ex Hook. f. Handb. N. Zeal. Fl. 734 = rotundifolius.
Renardi, *Winkl. in Trautv. Regel, Maxim. & Winkl. Decas. Pl. Nov.* 6.—Turkestan.
renifolius, *Porter & Coulter, Fl. Colorado*, 83.—Colorado.
renifolius, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Reg. Caucas.
reniformis, Wall. Cat. n. 3141 = Cremanthodium reniforme.
repandus, Hook. f. ex C. Muell. in Walp. Ann. v. 341 = Brachyglottis repanda.
repandus, Thunb. ex Harv. & Sond. Fl. Cap. iii. 357, partim = Lessingii.
repandus, *Thunb. Prod. Pl. Cap.* 158.—Afr. austr.
repens, Baill. Hist. Pl. viii. 308 = succulentus, *Sch. Bip.*
repens, *DC. Prod.* vi. 423.—Ecuador.
repens, Dulac, Fl. Hautes-Pyr. 507 = erucifolius.
repens, Stokes, Bot. Mat. Med. iv. 212 = sarracenicus.
reptans, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 207.—Afr. austr.
resectus, *Boj. ex DC. Prod.* vi. 378.—Madag.
resedifolius, *Less. in Linnaea*, vi. (1831) 243.—Reg. arct.
reticulatus, C. B. Clarke, Comp. Ind. 199 = Grahami.
reticulatus, *DC. Prod.* vi. 431.—Mexic.
retortus, *Benth. in Hook. Ic. Pl.* t. 1114.—Afr. austr.
retrorsus, DC. Prod. vi. 387 = barbellatus.
retusus, *Wall. Cat.* n. 3142.—Reg. Himal.
rhizocephalus, *A. Gray, ex Wedd. Chlor. And.* i. 129.—Peruv.
rhizocephalus, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 210.—Ecuador; N. Granat.
rhodaster, *Baker, in Mart. Fl. Bras.* vi. III. 316.—Bras.
rhombifolius, *Bolle, in Bonplandia*, vii. (1859) 296.—Ins. Canar.
rhombifolius, Sch. Bip. in Flora, xxviii. (1845) 499 = platyphyllus.

SENECIO:—
rhomboideus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 377.—Afr. austr.
rhyncholaenus, *DC. Prod.* vi. 397.—Afr. austr.
Richardianus, DC. l. c. 374 = australis.
Richii, *A. Gray, in Proc. Am. Acad.* v. (1861) 142.—Peruv.
Riddellii, Torr. & Gray, Fl. N. Am. ii. 444 = Douglasii.
rigens, Linn. Sp. Pl. ed. II. 1224 = Othonna parviflora.
rigescens, Jacq. Coll. v. 150. t. 6. f. 1 = rosmarinifolius.
rigidus, *Linn. Sp. Pl.* 872.—Afr. austr.
ringens, Steud. Nom. ed. II. ii. 564 = Othonna parviflora.
riparius, *DC. Prod.* vi. 377.—Madag.
riparius, Salisb. Prod. 197 = S. Doria.
riparius, Wallr. Sched. Crit. 482 = paludosus.
rivularis, DC. Prod. vi. 359 = crispatus.
rivularis, *Remy, in C. Gay, Fl. Chil.* iv. 191.—Chili.
robertiaefolius, *DC. Prod.* vi. 384.—Afr. austr.
Roborowskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 487.—China.
robustus, *F. Buch. in Trans. N. Z. Inst.* vi. (1874) 243.—N. Zel.
robustus, *Sch. Bip. in Flora*, xxviii. (1845) 50.—Sibir.
Rochelianus, Fuss, Fl. Transs. 353 = S. Doronicum.
Rodriguezii, *Willk. ex Rodrig. in Anal. Soc. Esp. Hist. Nat.* iii. (1874) 36.—Ins. Balear.
Roldana, *DC. Prod.* vi. 431.—Mexic.
rosellatus, *Boj. ex DC. l. c.* 377.—Madag.
roseus, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Mexic.
rosmarinifolius, *Linn. f. Suppl.* 369.—Afr. austr.
rostellatus, Steud. Nom. ed. II. ii. 564, sphalm. = rosellatus.
rotundifolius, *Hook. f. Fl. N. Zel.* i. 149.—N. Zel.
rotundifolius, Lapeyr. Hist. Abr. Pl. Pyr. 517 = S. Doronicum.
rotundifolius, Stokes, Bot. Mat. Med. iv. 215 = aureus.
Royleanus, DC. Prod. vi. 366 = graciliflorus.
rubens, Jacq. Hort. Vindob. iii. 50 = Gynura cernua.
rubescens, *S. Moore, in Journ. Bot.* xiii. (1875) 228.—China.
rudbeckiaefolius, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 283.—Peruv.
ruderalis, Harv. in Harv. & Sond. Fl. Cap. iii. 355 = pellucidus.
rudis, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 88 = leontodontis.
Rueppellii, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 157 = coronopifolius.
rufescens, *DC. Prod.* vi. 413.—Chili.
rufinervis, *DC. l. c.* 369.—Reg. Himal.
Rugelia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 54.—Am. bor.
rumicifolius, MacOwan & Bolus, in Journ. Linn. Soc. xviii. (1881) 392 = trullaefolius.
runcinatus, Less. in Linnaea, vi. (1831) 410 = Erechtites runcinata.
rupestris, Waldst. & Kit. Pl. Rar. Hung. ii. 136 = nebrodensis.
rupicola, A. Rich. Sert. Astrolab. 119 = lautus.
Rusbyi, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 64.—N. Mexic.
rutaceus, *Phil. in Linnaea*, xxviii. (1856) 751.—Chili.
ruthenensis, Maz. & Timb. Note Nouv. Senecio, 3 = S. Doronicum.
sabulosus, *DC. Prod.* vi. 422.—Ecuador.
Sadleri, Láng, ex Reichb. Fl. Germ. Excurs. 245 = paludosus.
Sagitta, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 483.—China.
sagittarius, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura Sagittaria.
sagittatus, *Sch. Bip. in Flora*, xxviii. (1845) 498.—Ross.; As. or.
sagittifolius, *Baker, in Mart. Fl. Bras.* vi. III. 315.—Reg. Argent.
salicetorum, Godr. Fl. Lorr. ed. I. ii. 11 = sarracenicus.
salicifolius, *Pers. Syn.* ii. 436.—Ins. Borbon.
salicifolius, Wallr. Sched. Crit. 478 = Fuchsii.
salignus, *DC. Prod.* vi. 430.—Mexic.
salsus, *Griseb. in Goett. Abh.* xix. (1874) 190.—Reg. Argent.

SENECIO:—
saltensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 338.—Reg. Argent.
salviifolius, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 160, nomen.—Afr. trop.
Sandersoni, *Harv. in Harv. & Sond. Fl. Cap.* iii. 365.—Afr. austr.
sandwicensis, *Less. in Linnaea*, vi. (1831) 526.—Ins. Owahu.
Sanguisorbae, *DC. Prod.* vi. 427.—Mexic.
saracenicus, Auct. ex Nym. Consp. 353 = nemorensis.
saracenicus, Koch, Syn. Fl. Germ. ed. I. 390 = sarracenicus.
sarcobasis, Boj. ex DC. Prod. vi. 300 = Gynura sarcobasis.
sarcophyllus, *Phil. in Linnaea*, xxviii. (1856) 740.—Chili.
sarcophyllus, *Steud. Nom.* ed. II. ii. 565.—Chili.
sarmentosus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 159 = Gynura sarmentosa.
sarracenicus, C. C. Gmel. Fl. Bad. iii. 443 = Fuchsii.
sarracenicus, *Linn. Sp. Pl.* 871.—Europ.; Sibir.
Sartorii, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 247.—Mexic.
Savatieri, Franch. Pl. Davidian. i. (1884) 175 = Oldhamianus.
saxatilis, *Wall. Cat.* n. 3131.—Ind. or.
saxicolus, *Wedd. Chlor. And.* i. 131.—Peruv.
saxifragoides, *Hook. f. Fl. N. Zel.* i. 144.—N. Zel.
scaber, Willd. Sp. Pl. iii. 1977 = rigidus.
scabriusculus, DC. Prod. vi. 396 = pinnulatus.
scabrosus, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 147 = Banksii.
scandens, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 178.—Reg. Himal.
scandens, DC. Prod. vi. 404 = mikanioides.
scandens, Juss. ex DC. l. c. 326 = Gynoxys cordifolia.
scandens, Wall. Cat. n. 3120 = S. Buimalia.
scanorensis, Linn. ex Steud. Nom. ed. I. 765 = erucifolius.
scapiger, Sieber, in Spreng. Syst. iii. 559 = Brachycome scapigera.
scaposus, *DC. Prod.* vi. 403.—Afr. austr.
Scheuchzeri, Gaud. Fl. Helv. v. 294 = gallicus.
Schimperi, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) I. *Intell.* 27, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 435.—Abyss.
Schkuhrii, DC. Prod. vi. 359 = crispatus.
Schlechtendahlii, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 322 = brasiliensis.
Schmidtii, *Franch. & Sav. Enum. Pl. Jap.* i. 246.—Japon.
Schultesii, Hochst. ex Field. & Gardn. Sert. Pl. t. 59, sphalm. = seq.
Schultzii, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 444.—Abyss.
Schultzii, *Wedd. Chlor. And.* i. 119.—Chili.
Schumannianus, *S. Schau. in Linnaea*, xx. (1847) 698.—Mexic.
Schweinitzianus, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 413 = lobatus.
sciadophilus, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 119.—N. Zel.
sclerophyllus, *Hemsl. Biol. Centr. Am. Bot.* ii. 247.—Mexic.
scoparius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 389.—Afr. austr.
Scopolii, Hoppe & Hornsch. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 380 = lanatus, *Scop.*
scopulorum, *Poepp. & Endl. Nov. Gen. et Sp. Pl.* iii. 58.—Chili.
scorpioides, Sch. Bip. in Flora, xxviii. (1845) 498 = Doronicum scorpioides.
scorzoneraefolius, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 282.—Peruv.
scorzoneroides, Hook. f. Fl. N. Zel. i. 146 = Lyallii.
Scotti, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 841.—Ins. Socotra.
Scrobicaria, DC. Prod. vi. 422 = Gynoxys ilicifolia.
scrobicarioides, *DC. l. c.* 430.—Mexic.
scytophyllus, *H. B. & K. Nov. Gen. et Sp.* iv. 181.—Ecuador.
sechellensis, *Baker, Fl. Maurit.* 178.—Ins. Seychelles.
sectilis, *Griseb. in Goett. Abh.* xix. (1874) 190.—Reg. Argent.
Seemannii, Sch. Bip. in Seem. Bot. Voy. Herald, 311 = Hartwegi.

SENECIO:—
Segethi, *Phil. in Linnaea*, xxxiii. (1864–65) 156.—Chili.
segmentatus, *Oliver, in Hook. Ic. Pl.* t. 1488.—Afr. austr.
Selloi, *DC. Prod.* vi. 419.—Bras.
semidentatus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 331.—N. Granat.
seminudus, Bory, in Ann. Gén. Sc. Phys. i. (1819) 303. t. 12 = Erechtites praealta.
semperflorens, Stokes, Bot. Mat. Med. iv. 210 = vulgaris.
sempervivus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Arab.
sepium, *Sch. Bip. in Linnaea*, xxxiv. (1865-66) 531, nomen.—Bolivia.
serenensis, *Remy, in C. Gay, Fl. Chil.* iv. 171.—Chili.
Serra, *Hook. Fl. Bor. Am.* i. 333.—Am. bor. occ.
Serra, *Sond. in Linnaea*, xxiii. (1850) 68.—Natal.
serratulaefolius, Weinm. in Syll. Ratisb. ii. (1828) 176 = glastifolius, *Linn. f.*
serratuloides, *DC. Prod.* vi. 395.—Afr. austr.
serratus, MacOwan & Bolus, in Journ. Linn. Soc. xviii. (1881) 392 = incomptus.
serratus, Sond. ex Harv. & Sond. Fl. Cap. iii. 366 = Leontodontis.
serrulatus, *DC. Prod.* vi. 401.—Afr. austr.
serrurioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 89.—Afr. austr.
sessilifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 247.—Mexic.
sessilifolius, Sch. Bip. in Flora, xxviii. (1845) 50 = Werneria nana.
sessilis, Thunb. Prod. Pl. Cap. 158 = halimifolius.
sessilis, Vell. Fl. Flum. viii. t. 106 = adamantinus.
setulosus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 330.—N. Granat.
sibiricus, C. B. Clarke, Comp. Ind. 206 = S. Ligularia.
sibiricus, *Linn. f. Suppl.* 370.—Sibir. altaic.
siculus, Jan, ex DC. Prod. vi. 351 = erucifolius.
Sieberianus, *DC. Prod.* vi. 373.—Austral.
Sieboldii, Sch. Bip. in Flora, xxviii. (1845) 50 = Kaempferi.
Siegfriedi ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 106.—Europ.
Simonsii, *C. B. Clarke, Comp. Ind.* 188.—Reg. Himal.
simplicissimus, *Boj. ex DC. Prod.* vi. 377.—Madag.
sinuatidens, Peterm. in Flora, xxvii. (1844) 474 = aquaticus.
sinuatilobus, *DC. Prod.* vi. 417.—Chili.
sinuatus, Gilib. Hist. Pl. Europ. ii. 245 = vernalis.
sinuatus, *H. B. & K. Nov. Gen. et Sp.* iv. 180.—Mexic.
sisymbrifolius, *DC. Prod.* vi. 382.—Afr. austr.
sisymbriiformis, DC. l. c. 366 = chrysanthemoides.
Skinneri, *Hemsl. Biol. Centr. Am. Bot.* ii. 247.—Guatemala.
skirrhodon, *DC. Prod.* vi. 401.—Afr. austr.
Smithii, *DC. l. c.* 412.—Reg. Magellan.
sneeuwbergensis, *Bolus, in Hook. Ic. Pl.* t. 1067.—Afr. austr.
socialis, *Wedd. Chlor. And.* i. 109.—Peruv.
solanoides, Sch. Bip. in Flora, xxvii. (1844) 702 = incertus.
solanoides, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 159.—Abyss.
Soldanella, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 67.—Am. bor. occ.
solidagineus, Spreng. Syst. iii. 552 = seq.
solidaginoides, Berg. Desc. Pl. Cap. 284 = halimifolius.
sonchifolius, Moench, Meth. Suppl. 231 = Emilia sonchifolia.
sonchoides, *H. B. & K. Nov. Gen. et Sp.* iv. 178.—Peruv.
sonchoides, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 145 = Vukotinovici.
Sonneratii, *Steud. Nom.* ed. II. ii. 565.—Afr. austr.
soongaricus, *Schrenk, Enum. Pl. Nov.* 62.—Soongar.
sophioides, *DC. Prod.* vi. 382.—Afr. austr.
Soratae, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen.—Bolivia.
sparsiflorus, Steud. Nom. ed. II. ii. 565, sphalm. = seq.
sparsifolius, *Boj. ex DC. Prod.* vi. 378.—Madag.
spartioides, Torr. & Gray, Fl. N. Am. ii. 438 = Douglasii.
spathulaefolius, *DC. Prod.* vi. 362.—Europ.
spathulaefolius, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 154 = campestris.
spathulatus, *A. Rich. Sert. Astrolab.* 125.—Austral.

SENECIO:—
spatulifolius, Kellogg, in Proc. Calif. Acad. i. ed. II. (1873) 56 = fastigiatus.
speciosus, *Willd. Sp. Pl.* iii. 1991.—Afr. austr.
spectabilis, Wall. Cat. n. 3127 = chrysanthemoides.
spicatus, *Sweet, Hort. Brit.* ed. II. 304, nomen.—Hab.?
spinescens, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
spinosus, *DC. Prod.* vi. 420.—Peruv.
spiraeifolius, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
squalidus, Bieb. Fl. Taur. Cauc. ii. 306 = vernalis.
squalidus, *Linn. Sp. Pl.* 869.—Europ.
squalidus, Urv. in Mém. Soc. Linn. Par. i. (1822) 366 = coronopifolius.
squalidus, Willd. Sp. Pl. iii. 1991 = gallicus.
squamosus, *DC. Prod.* vi. 376.—Ins. Borbon.
squamosus, Thunb. Prod. Pl. Cap. 158 = pubigerus.
squarrosus, A. Rich. Sert. Astrolab. 107 = Erechtites hispidula.
Stauntonii, *DC. Prod.* vi. 363.—China.
Steetzii, *Bolle, in Bonplandia*, vii. (1859) 296.—Ins. Canar.
Stenactis, Sch. Bip. in Flora, xxviii. (1845) 50 = amplexicaulis, *Wall.*
stenocephalus, *Boiss. Fl. Orient.* iii. 408.—Reg. Caucas.
stenocephalus, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 218.—China; Japon.
stenophyllus, Bert. ex DC. Prod. vi. 448 = Robinsonia gracilis.
stenophyllus, *Phil. in Linnaea*, xxviii. (1856) 743.—Chili.
stenopteris, *Griseb. in Goett. Abh.* xix. (1874) 193.—Reg. Argent.
Sternbergianus, *DC. Prod.* vi. 425.—Peruv.
Steudelii, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 440.—Abyss.
steudelioides, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 157.—Afr. bor.
Stewartiae, *Armstr. in Trans. N. Z. Inst.* xiii. (1881) 339.—N. Zel.
stigophlebius, *Baker, in Mart. Fl. Bras.* vi. III. 321.—Bras.
stipulatus, Vell. Fl. Flum. viii. t. 113 = fluminensis.
stipulatus, Wall. Cat. n. 3122 = scandens.
stipulatus, Wall. Cat. n. 3126 = chrysanthemoides.
stoechadiformis, DC. Prod. vi. 429 = Douglasii.
striatifolius, *DC. l. c.* 387.—Afr. austr.
striatus, Thunb. Prod. Pl. Cap. 157 = Osteospermum Bidens.
strictus, Banks, ex Pursh, Fl. Am. Sept. ii. 529 = gracilis.
strictus, DC. Prod. vi. 343 = Perrottetii.
suaveolens, *Ell. Sketch*, ii. 328.—Am. bor.
suaveolens, Gilib. Hist. Pl. Europ. ii. 245 = erucifolius.
subalpinus, *Koch, in Flora*, xvii. (1834) 614.—Europ. austr.
subarachnoideus, *Sch. Bip. ex Wedd. Chlor. And.* i. 127.—N. Granat.
subaraneosus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Java.
subcandidus, *A. Gray, in Proc. Am. Acad.* v. (1861) 141.—Peruv.
subcordatus, Boj. ex DC. Prod. vi. 330 = Faujasia flexuosa.
subcorymbosus, Rupr. ex Boiss. Fl. Orient. iii. 405 = amphibolus, *C. Koch.*
subdecurrens, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Peruv.
subdentatus, Ledeb. Fl. Alt. iv. 110 = coronopifolius.
subdentatus, *Phil. in Linnaea*, xxviii. (1856) 748.—Chili.
subensiformis, DC. l. c. 363 = campestris.
sublobatus, DC. Prod. vi. 310 = S. Roldana.
subnebrodensis, *Simk. in Termész. Füzetek*, v. (1881) 51.—Europ.
subnudus, DC. Prod. vi. 405 = oxyriaefolius.
subnudus, DC. l. c. 428 = aureus.
subpeltatus, *Sch. Bip. in Seem. Bot. Voy. Herald*, 311.—Mexic.
subpeltatus, Steud. Nom. ed. II. ii. 565 = oxyriaefolius.
subradiatus, *Sch. Bip. in Flora*, xxviii. (1845) 500.—Afr. austr.
subscandens, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 434.—Abyss.
subsessilis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 415.—Afr. trop.

SENECIO:—
subsinuatus, *DC. Prod.* vi. 401.—Afr. austr.
subulatus, [*D.*] *Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 330.—Chili; Patagon.
subumbellatus, Phil. in Anal. Univ. Chil. (1861) 64 = thermarum.
subverticillatus, DC. Prod. vi. 430 = Andrieuxii.
succulentus, *DC. l. c.* 402.—Afr. austr.
succulentus, Forsk. Fl. Aegypt. Arab. 149 = Inula crithmoides.
succulentus, *Sch. Bip. in Flora*, xxviii. (1845) 499.—Afr. austr.
sudeticus, DC. Prod. vi. 359 = crispatus.
sulcatus, DC. l. c. 384 = crispus, *Thunb.*
sumatranus, *Martelli, in Nouv. Giorn. Bot. Ital.* xv. (1883) 301.—Sumatr.
surculosus, *MacOwan, in Journ. Linn. Soc.* x. (1869) 480.—Afr. austr.
Swartzianus, *Buek, Ind. ad DC. Prod.* ii. p. vi.—Jamaic.
Swartzii, *DC. Prod.* vi. 411.—Jamaic.
Sweetii, Heynh. Nom. ii. 656 = Cineraria pulchella.
sylvarum, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 531, nomen = Bolivia.
sylvaticus, Gueldenst. Reisen, ii. 56; ex Ledeb. Fl. Ross. ii. 630 = vernalis.
sylvaticus, *Linn. Sp. Pl.* 868.—Europ.; As. bor.
sylvaticus, Nestl. ex DC. Prod. vi. 353 = Fuchsii.
sylvestris, Grindel. Bot. Taschenb. 254 = sylvaticus.
Syneilesis, *Franch. & Sav. Enum. Pl. Jap.* i. 249.—Japon.
syrmiensis, Kit. ex Nym. Consp. 357 = vernalis.
Tabacon, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 91.—Venez.; Ecuador.
tabidus, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 141 = Erechtites prenanthoides.
tabularis, *Hemsl. Biol. Centr. Am. Bot.* ii. 248.—Mexic.
tabularis, Sch. Bip. in Flora, xxviii. (1845) 500 = Alciope tabularis.
talinoides, *Sch. Bip. l. c.* 499.—Afr. austr.
tamoides, *DC. Prod.* vi. 403.—Afr. austr.
tampicanus, *DC. l. c.* 427.—Mexic.
tampicanus, A. Gray, Pl. Wright. ii. 89 = multilobatus.
tanacetoides, Kunth & Bouché, Ind. Sem. Hort. Berol. (1845) 12 = graciliflorus.
tanacetoides, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 394.—Afr. austr.
tanguticus, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 486.—China.
taraxacifolius, *DC. Prod.* vi. 348.—Reg. Caucas.; Persia.
tasconensis, Martr. in Soc. Sc. Tarn-et-Garonne (1853); (Pl. Crit. Tarn.) 27 = erucifolius.
Tasna, Welw. ex Willk. & Lange, Prod. Fl. Hisp. ii. 120 = foliosus.
tataricus, Less. in Linnaea, ix. (1834) 192 = auratus.
taxifolius, Poir. Encyc. Suppl. v. 133 = S. Ambavilla.
taygeteus, *Boiss. & Heldr. Diagn.* Ser. I. vi. 95.—Graec.
Tchihatchewi, Boiss. in Tchihat. Asie Min. Bot. ii. 293 = mollis.
Tedliei, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 420.—Afr. trop.
telephifolius, Jacq. Fragm. i. 1. t. 1. f. 3 = maritimus.
tenellus, *DC. Prod.* vi. 379.—Afr. austr.
Teneriffae, Sch. Bip. ex Bolle, in Bonplandia, viii. (1860) 133 = vulgaris.
tenuicaulis, *Sch. Bip. in Linnae*, xxxiv. (1865–66) 531, nomen.—Bolivia.
tenuiculus, *Steud. Nom.* ed. II. ii. 565.—Chili.
tenuiflorus, Sieber, ex DC. Prod. vi. 296 = Erechtites quadridentata.
tenuifolius, *Burm. f. Fl. Ind.* 181.—Ind. or.; Java.
tenuifolius, DC. Prod. vi. 398 = adonidifolius.
tenuifolius, Jacq. Fl. Austr. iii. 42 = erucifolius.
tenuilobus, DC. Prod. vi. 398 = umbellatus.
tenuiscapus, *Boj. ex DC. l. c.* 378.—Madag.
tephrosioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 92.—Ecuador.
teretifolius, *DC. Prod.* vi. 420.—Ecuador.
tetrandrus, *Buch.-Ham. ex Wall. Cat.* n. 3129.—Reg. Himal.
tetranthus, *DC. Prod.* vi. 370.—Reg. Himal.
thapsoides, *DC. l. c.* vii. 301.—Graec.
theaefolius, *Benth. Pl. Hartw.* 210.—N. Granat.
thermarum, *Phil. in Linnaea*, xxxiii. (1864–65) 153.—Chili.

SENECIO:—
thianschanicus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 311.—As. centr.
thinophilus, *Phil. in Anal. Univ. Chil.* (1873) 500.—Chili.
Thomasii, *Klatt, in Abh. Naturf. Ges. Halle,* xv. (1882) 332.—Mexic.
Thomsoni, *C. B. Clarke, Comp. Ind.* 205.—Reg. Himal.
Thunbergianus, *Schrad. Blumenbachia,* 41. t. 4. f. 2.—Afr. austr.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 379.—Afr. austr.
Thurberi, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 68.—N. Mexic.
thurifer, Bert. ex DC. Prod. vi. 448 = Robinsonia thurifera.
thyrsoideus, *DC. l. c.* 406.—Afr. austr.
thyrsoideus, G. Don, in Sweet, Hort. Brit. ed. III. 386 = Fischerianus.
thyrsophorus, C. Koch, in Linnaea, xxiv. (1851) 362 = racemosus.
tibeticus, *Hook. f. Fl. Brit. Ind.* iii. 340.—Tibet. occ.
tillandsiaefolius, Poepp. ex DC. Prod. vi. 415 = argenteus.
tmoleus, Boiss. Diagn. Ser. I. iv. 13 = Castagneanus.
Tocornali, *Phil. in Anal. Univ. Chil.* (1862) II. 383.—Chili.
tolimensis, *Sch. Bip. ex Wedd. Chlor. And.* i. 98, 99.—N. Granat.
toluccanus, *DC. Prod.* vi. 428.—Mexic.
tomentosus, [Clairv.] Man. Herb. 240 = Adenostyles candidissima.
tomentosus, C. B. Clarke, Comp. Ind. 200 = Edgeworthii.
tomentosus, Dulac, Fl. Hautes-Pyr. 507 = S. Doronicum.
tomentosus, Host, Fl. Austr. ii. 476 = paludosus.
tomentosus, *Michx. Fl. Bor. Am.* ii. 119.—Am. bor.
tomentosus, Rohde, ex DC. Prod. vi. 356 = leucophyllus.
tomentosus, Salisb. Prod. 196 = cinerascens.
tortuosus, *DC. Prod.* vi. 392.—Afr. austr.
Tournefortii, *Lapeyr. Hist. Abr. Pl. Pyr.* 516.—Mont. Pyren.
trachycarpus, Sch. Bip. in Flora, xxvii. (1844) 702 = incertus.
trachylaenus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 358.—Afr. austr.
transilvanicus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 208 = sarracenicus.
transylvanicus, Boiss. Diagn. Ser. II. iii. 34 = S. Doronicum.
trapezuntinus, *Boiss. Fl. Orient.* iii. 393.—Armenia.
Traversii, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1861) 154 = bellidioides.
tredecimsquamosus, Sch. Bip. in Flora, xxvii. (1844) 700 = angulatus.
triangularis, *Hook. Fl. Bor. Am.* i. 332. t. 115.—Am. arct.
Trianii, *Klatt, in Abh. Naturf. Ges. Halle,* xv. (1882) 332.—N. Granat.
trichocaulon, *Baker, in Mart. Fl. Bras.* vi. III. 315.—Bras.
trichocodon, *Baker, l. c.* 314.—Paraguay.
tricuspidatus, Banks & Soland. ex Hook. f. Fl. Antarct. 317 = Darwinii.
tricuspidatus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 346.—Chili.
tridentatus, *DC. Prod.* vi. 424.—Peruv.
trifidus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 346.—Chili.
triflorus, Linn. Sp. Pl. 867 = aegyptius.
trifurcatus, *Less. Syn. Comp.* 392.—Reg. Magellan.
triligulatus, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 178.—Reg. Himal.
trilobus, Linn. Sp. Pl. 868 = lividus.
trineurus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 514.—Cuba.
triodon, *Phil. in Linnaea,* xxviii. (1856) 749.—Chili.
tripartitus, DC. Prod. vi. 418 = brasiliensis.
tripartitus, A. Rich. Sert. Astrolab. 114 = lautus.
triplinervius, *DC. Prod.* vi. 395.—Afr. austr.
triplipartitus, Steud. Nom. ed. II. ii. 566 = lautus.
triqueter, *Less. in Linnaea,* vi. (1831) 251.—Afr. austr.
tristis, *Phil. in Linnaea,* xxviii. (1856) 739.—Chili.
trixoides, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 421.—Bras.

SENECIO:—
Troncosi, *Phil. Fl. Atac.* 32.—Chili.
tropaeolifolius, *MacOwan, ex F. Muell. in Journ. Bot.* v. (1867) 242.—Afr. austr.
trullaefolius, *MacOwan, in Journ. Linn. Soc.* xviii. (1881) 392.—Afr. austr.
truncatus, Buch.-Ham. ex Wall. Cat. n. 3137 = nudicaulis.
truncatus, Dulac, Fl. Hautes-Pyr. 507 = S. Jacobaea.
tschabanicus, *C. Koch, in Linnaea,* xxiv. (1851) 370.—Oriens.
tuberosus, Harv. in Harv. & Sond. Fl. Cap. iii. 375 = incertus.
tuberosus, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 434.—Abyss.
Turczaninovii, *DC. Prod.* vi. 360.—Dahuria.
Turczaninowii, Maxim. in Bull. Acad. Pétersb. xix. (1874) 488 (cum cit. falsa) = aconitifolius.
Tussilaginis, *Less. Syn. Comp.* 392.—Ins. Canar.
tussilaginoides, Heynh. Nom. ii. 656 = S. Tussilaginis.
tussilaginoides, *Sch. Bip. in Flora,* xxviii. (1845) 498.—Mexic.
tussilaginoides, Walt. Fl. Carol. 208 = aureus.
Tweediei, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 339.—Reg. Argent.
tymphresteus, Boiss. & Heldr. Diagn. Ser. II. vi. 99 = viscosus.
Ualtata, Bert. ex Steud. Nom. ed. II. ii. 566 = S. Hualtata.
uliginosus, DC. Prod. vi. 348 = gallicus.
uliginosus, *Phil. in Anal. Univ. Chil.* (1873) 501.—Chili.
umbellatus, *Linn. Pl. Afr. Rar.* 21; *Amoen. Acad.* vi. 101.—Afr. austr.
umbrosus, *Waldst. & Kit. Pl. Rar. Hung.* iii. 232. t. 210.—Hungar.
uncinellus, DC. Prod. vi. 368 = densiflorus.
undulatus, *Thunb. Prod. Pl. Cap.* 157.—Afr. austr.
uniflorus, *All. Fl. Pedem.* i. 200. t. 17.—Europ.
uniflorus, Retz. Obs. iii. 42 = Gynura cernua.
Unionis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 440.—Abyss.
uniserialis, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 126, nomen.—Malaya.
uspallatensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 335.—Chili.
ustulatus, DC. Prod. vi. 380 = littoreus.
vaccinioides, *Wedd. Chlor. And.* i. 99.—N. Granat.
vagans, *Wall. Cat.* n. 3108.—Reg. Himal.
vaginaefolius, *Sch. Bip. in Flora,* xxxviii. (1855) 117.—Reg. Magellan.
vaginatus, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 331 = littoralis.
vagus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 46.—Austral.
valerianaefolius, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 127.—Bras.
valerianifolius, Wolf, Ind. Sem. Hort. Berol. (1825); ex Reichb. Ic. Bot. Exot. i. 59 = Erechtites valerianaefolia.
vallestris, *DC. Prod.* vi. 421.—Peruv.
valparadisiacus, Colla, in Mem. Acc. Torin. xxxviii. (1835) 32 = glabratus.
valparadisiensis, *Remy, in C. Gay, Fl. Chil.* iv. 157.—Chili.
variabilis, Sch. Bip. in Flora, xxvii. (1844) 700 = erubescens, *Ait.*
varicosus, *Linn. f. Dec.* i. 9. t. 5.—Aegypt.
variegatus, *Sch. Bip. in Linnaea,* xxxiv. (1865–66) 536, nomen.—Bolivia.
variifolius, DC. Prod. vi. 393 = lyratus.
velleioides, *A. Cunn. ex DC. Prod.* vi. 374.—Austral.
venosus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 379.—Afr. austr.
venustus, Ait. Hort. Kew. ed. I. iii. 192 = grandiflorus.
verbascifolius, *Burm. Fl. Cap. Prod.* 26.—Afr. austr.
verbascifolius, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrolab. et Zél. 47 = Smithii.
verbenaefolius, Jacq. Hort. Vindob. i. 2. t. 3 = aegyptius.
vernalis, Hoppe, ex DC. Prod. vi. 351 = nebrodensis.
vernalis, *Waldst. & Kit. Pl. Rar. Hung.* i. 23. t. 24.—Europ. austr.; Oriens.
vernicosus, *Sch. Bip. ex Wedd. Chlor. And.* i. 94.—Venezuel.; N. Granat.
vernonioides, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 312.—Bras.
vernonioides, Sch. Bip. in Flora, xxvii. (1844) 701 = erubescens, *Ait.*

SENECIO:—
vernus, Bivona, Sicul. Pl. Cent. i. 73. t. 5 = leucanthemifolius, *Poir.*
vernus, DC. Prod. vi. 430 = salignus.
vernus, Urv. in Mém. Soc. Linn. Par. i. (1822) 366 = vernalis.
verrucosus, *Klatt, in Abh. Naturf. Ges. Halle,* xv. (1882) 332.—N. Granat.
verrucosus, Sch. Bip. ex Wedd. Chlor. And. i. 77 = Gynoxys verrucosa.
verticillatus, *Klatt, in Abh. Naturf. Ges. Halle,* xv. (1882) 331.—Peruv.
vestitus, *Berg. Desc. Pl. Cap.* 282.—Afr. austr.
vestitus, Sweet, ex Steud. Nom. ed. II. ii. 566 = Haworthii, *Steud.*
villosus, Kostel. Allg. Med.-Pharm. Fl. ii. 715 = palustris?
vimineus, *DC. Prod.* vi. 400.—Afr. austr.
vimineus, *Harv. in Harv. et Sond. Fl. Cap.* iii. 401.—Afr. austr.
violaceus, Wall. Cat. n. 3134 = nudicaulis.
virgatus, *Linn. Pl. Afr. Rar.* 20; *Amoen. Acad.* vi. 101.—Afr. austr.
Virgaurea, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 484.—China.
viridulus, Martr. Pl. Crit. Tarn, 27 = erucifolius.
viscidulus, Scheele, in Linnaea, xviii. (1844) 480 = sylvaticus.
viscosissimus, Colla, in Mem. Acc. Torin. xxxviii. (1835) 33 = glaber.
viscosus, *Linn. Sp. Pl.* 868.—Europ.; As. Min.
viscosus, Webb, Iter Hisp. 40 = Durieui.
Visianus, *Papaf. ex Vis. Fl. Dalm.* ii. 72. t. 8.—Dalmat.
vitellinus, Aschers. in Schweinf. Beitr. Fl. Aethiop. 286 = Gynura vitellina.
vittarifolius, *Boj. ex DC. Prod.* vi. 377.—Madag.
Volckameri, Sch. Bip. in Doell, Rhein. Fl. 470 = arenarius.
Volckmanni, *Phil. in Linnaea,* xxxiii. (1864–65) 152.—Chili.
volubilis, *Hook. Bot. Misc.* ii. (1831) 226.—Peruv.
Vukotinovici, *Schloss. in Oestr. Bot. Zeitschr.* xxxi. (1881) 5.—Europ. austr.
vulcanicus, *Boiss. Diagn.* Ser. I. xi. 31.—Persia.
vulcanicus, *Phil. in Linnaea,* xxviii. (1856) 746.—Chili.
vulgaris, *Linn. Sp. Pl.* 867.—Europ. (Cosmop. inquil.).
Vulneraria, *DC. Prod.* vi. 428.—Mexic.
Walkeri, Arn. in Nov. Act. Nat. Cur. xviii. I. (1836) 349 = araneosus.
Walkeri, Thw. Enum. Pl. Zeyl. 167 = Notonia Walkeri.
Wallichii, *DC. Prod.* vi. 364.—Reg. Himal.
Walteri, *Sch. Bip. in Flora,* xxviii. (1845) 499.—Am. bor.
Warszewiczii, *A. Br. & Bouché, Ind. Sem. Hort. Berol. App.* (1851) 13.—*Cf. Linnaea,* xxv. (1852) 298.—Guatemala.
Wartmanni ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 106.—Europ.
Websteri, *Hook. f. Fl. Antarct.* 317.—Reg. Magellan.
werneriaefolius, *A. Gray, in Proc. Am. Acad.* xix. (1883) 54.—Am. bor. occ.
wernerioides, *Wedd. Chlor. And.* i. 128.—Peruv.
Weylii ×, Vatke, in Verh. Bot. Ver. Brandenb. xiv. (1872) 48.—Hab.?
Whippleanus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 54.—Calif.
Wightianus, DC. Prod. vi. 370 = scandens.
Wightii, Benth. ex C. B. Clarke, Comp. Ind. 197 = saxatilis.
xerophilus, *Phil. Fl. Atac.* 32.—Chili.
Yakla, C. B. Clarke, Comp. Ind. 204 = amplexicaulis, *Wall.*
Zahlbruckneri, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxviii–xxiv. (1880) 106 (Quid?).—Europ.
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 89.—Afr. austr.
zeylanicus, *DC. Prod.* vi. 365.—Zeylan.
zimapanicus, *Hemsl. Biol. Centr. Am. Bot.* ii. 248.—Mexic.
Zollingerianus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 160 = Gynura mollis.
zosteraefolius, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 342.—Chili.
Zuccarinii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 486.—Japon.

SENECIUNCULUS, Opiz, Seznam, 90 (1852) = **Senecio**, Tourn. (Compos.).
viscosus, Opiz, l. c. = Senecio viscosus.

SENEFELDERA, Mart. in Flora, xxiv. (1841) II. Beibl. 29. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 332.
angustifolia, Klotzsch, in Wiegm. Archiv, vii. (1841) 184 = multiflora.
dodecandra, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 529.—Bras.
grandifolia, Baill. Etud. Gén. Euph. 536. t. 9 = multiflora.
inclinata, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 530.—Bras.
latifolia, Klotzsch, in Wiegm. Archiv, vii. (1841) 184 = multiflora.
multiflora, *Mart. in Flora*, xxiv. (1841) II. Beibl. 29.—Bras.

SENEGA, Spach, Hist. Vég. Phan. vii. 128 (1839) = **Polygala**, Linn.
officinalis, Spach, l. c. 129 = P. Senega.

SENEGALIA, Rafin. Sylva Tellur. 119 (1838) = **Acacia**, Willd. (Legumin.).
retusa, Rafin. l. c. = A. Senegal.
triacantha, Rafin. l. c. = A. glaucophylla.

SENEICO, Hill, Hort. Kew. 25 (1768), sphalm. = **Senecio**, Tourn. (Compos.).

SENFTENBERGIA, Klotzsch & Karst. ex Klotzsch, in Linnaea, xx. (1847) 460 = **Langsdorffia**, Mart. (Balanoph.).
Moritziana, Klotzsch & Karst. ex Eichl. in Mart. Fl. Bras. iv. II. 10 = L. hypogaea.

SENITES, Adans. Fam. ii. 39. (1763) = **Zeugites**, Schreb. (Gramin.).

SENKEBERGIA, Neck. Elem. i. 366 (1790) = **Justicia**, Houst. (Acanth.).
debilis, Rafin. Sylva Tellur. 70 = J. debilis.

SENKENBERGIA, Schau. in Linnaea, xix. (1847) 711. *NYCTAGINEAE*, Benth. & Hook. f. iii. 5.
LINDENIA, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 357.
TINANTIA, Mart. & Gal. l. c. xi. I. (1844) 240.
annulata, *Schau. in Linnaea*, xix. (1847) 711.—Mexic.
Coulteri, *Hook. f. in Benth. et Hook. f. Gen.* iii. 6.—Mexic.
erecta, *Benth. & Hook. f. l. c.*—Am. bor. occ.
gypsophiloides, Benth. & Hook. f. l. c. = annulata.
stricta, Benth. & Hook. f. l. c. 1217, sphalm. = erecta.

SENNA, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768) = **Cassia**, Tourn. (Legumin.).
Absus, Roxb. Fl. Ind. ii. 340 = C. Absus.
acutifolia, Link, Handb. ii. 140; Batka, in Proc. Linn. Soc. ii. (1854) 282 = C. acutifolia.
alata, Roxb. Fl. Ind. ii. 349 = C. alata.
alexandrina, Mill. Gard. Dict. ed. VIII. n. 1 = C. Sophera.
angustifolia, Batka, in Proc. Linn. Soc. ii. (1854) 283 = C. angustifolia.
arborescens, Roxb. Fl. Ind. ii. 345 = C. glauca.
aurata, Roxb. l. c. 342 = C. laevigata.
auriculata, Roxb. l. c. 349 = C. auriculata.
bicapsularis, Roxb. l. c. 342 = C. bicapsularis.
dimidiata, Roxb. l. c. 352 = C. Kleinii.
esculenta, Roxb. l. c. 346 = C. Sophera.
exigua, Roxb. l. c. 339 = C. Absus.
glauca, Roxb. l. c. 351 = C. montana.
Hookeriana, Batke, in Bot. Zeit. xxi. (1863) 264 = C. adenensis.
italica, Mill. Gard. Dict. ed. VIII. n. 2 = C. obovata, acutifolia.
lanceolata, Link, Handb. ii. 140 = C. Sophera.
marylandica, Link, l. c. = C. marylandica?
obovata, Link, l. c.; Batka, in Proc. Linn. Soc. ii. (1854) 281 = C. obovata.
obtusa, Roxb. Fl. Ind. ii. 344 = C. obovata.
obtusata, Link, Handb. ii. 140 = C. obovata.
occidentalis, Link, l. c.; Roxb. Fl. Ind. ii. 343 = C. occidentalis.
officinalis, Gaertn. Fruct. ii. 312 = C. Sophera.

SENNA:—
officinalis, Roxb. Fl. Ind. ii. 346 = Cassia angustifolia.
orientalis, Walp. Rep. i. 816 = C. occidentalis.
ovalifolia, Batka, Monog. Senn. 35, 50. t. 4 = C. holosericea.
prostrata, Roxb. Fl. Ind. ii. 352 = C. pumila.
pubescens, Mart. Vers. Monog. Sennesbl. (1857) 61 = *tomentosa*.
purpurea, Roxb. Fl. Ind. ii. 342 = C. Sophera.
riparia, Rafin. Med. Fl. i. 94 = C. marylandica.
sensitiva, Roxb. Fl. Ind. ii. 353 = C. mimosoides.
Sophora, Roxb. l. c. 347 = C. Sophera.
speciosa, Roxb. l. c. = C. glauca.
sumatrana, Roxb. l. c. = C. siamea.
tenella, Roxb. l. c. 354 = C. mimosoides.
tomentosa, Batka, in Proc. Linn. Soc. ii. (1854) 283; et Bot. Zeit. xii. (1854) 115 = C. holosericea.
Tora, Roxb. Fl. Ind. ii. 340 = C. Tora.
toroides, Roxb. l. c. 341 = C. Tora.

SENNEBIERA, Willd. Enum. Hort. Berol. 668 = **Senebiera**, DC. (Crucif.).

SENNEFELDERA, Endl. Gen. Suppl. ii. 88 (1842) = SENEFELDERA, Mart. (Euphorb.).

SENRA, Cav. Diss. ii. 83, 104. t. 35. f. 3 (1786). *MALVACEAE*, Benth. & Hook. f. i. 207.
DUMREICHERA, Hochst. in Flora, xxi. (1838) I. Intell. 26.
SERRAEA, Spreng. Syst. iii. 78 (1826).
arabica, Webb, Fragm. Fl. Aethiop. 48 = incana.
incana, *Cav. Diss.* ii. 83.—As. et Afr. trop.
nubica, Webb, Fragm. Fl. Aethiop. 49 = incana.

SENRAEA, Willd. Sp. Pl. iii. 695 (1800) = **Senra**, Cav. (Malvac.).

SENSITIVA, Rafin. Sylva Tellur. 119 (1838) = **Mimosa**, Linn. (Legumin.).
pudica, Rafin. l. c. = M. pudica.

SENTIS, F. Muell. Fragm. iv. 47 (1863) = **Pholidia**, R. Br. (Myopor.).
rhynchocarpa, F. Muell. l. c. 48 = P. divaricata.

SEPTACANTHUS, Wight, Ic. t. 1527 (1850), sphalm. = LEPTACANTHUS, Nees = **Strobilanthes**, Blume (Acanth.).

SEPTAS, Linn. Pl. Afr. Rar. 10 (1760); Amoen. Acad. vi. 87 (1764) = **Crassula**, Dill.
capensis, Linn. ll. cc. = C. Septas.
globifera, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. ii. 357 = C. Saxifraga.
globifera, Sims, Bot. Mag. t. 1472 = C. Septas.
trichotoma, Lam. Illustr. ii. 409.—Afr. austr.
Umbella, Haw. Syn. Pl. Succ. 62 = C. Umbella.

SEPTAS, Lour. Fl. Cochinch. 392 (1790) = **Herpestis**, Gaertn. f. (Scrophul.).
repens, Lour. l. c. = H. Monnieria.

SEPTILIA, Rafin. Fl. Tellur. iv. 68 (1836) = **Herpestis**, Gaertn. f. (Scrophul.).
repens, Rafin. l. c. = H. Monnieria.

SEPTINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Cinnamomum**?, Tourn. (Laurin.).

SEPTIS, Hook. f. Fl. Brit. Ind. iv. 272 (1884), err. typ. = SEPTAS, Lour. = **Herpestis**, Gaertn. f. (Scrophul.).

SEQUOIA, Endl. Syn. Conif. 197 (1847). *CONIFERAE*, Benth. & Hook. f. iii. 429.
CONDYLOCARPUS, Salisb. ex Lamb. Gen. Pin. ed. 8vo. 120, sub t. 64 (1832).
GIGANTABIES, [Nelson] Senilis, Pinaceae, 77 (1866).
WASHINGTONIA, Winsl. in Calif. Farm. (Sept. 1854); ex Hook. Kew Journ. vii. (1855) 29.
WELLINGTONIA, Lindl. in Gard. Chron. (1853) 823.
gigantea, Endl. Syn. Conif. 198 = sempervirens.
gigantea, *Lindl. & Gord. in Journ. Hort. Soc.* v. (1850) 222; *Decne. in Bull. Soc. Bot. Fr.* i. (1854) 70, 71; *Torr. ex A. Gray, in Am. Journ. Sc.* Ser. II. xviii. (1854) 287.—Calif.

SEQUOIA:—
pyramidata, Hort. ex Carr. Conif. ed. II. 211 = sempervirens.
Rafinesquei, Carr. l. c. 213 = Picea sitchensis?
religiosa, Presl, Epim. Bot. 237 = sempervirens.
sempervirens, *Endl. Syn. Conif.* 198.—Calif.
taxifolia, Hort. ex C. Koch, Dendrol. ii. II. 194 = sempervirens.
Wellingtonia, Seem. in Bonplandia, iii. (1855) 27; vi. (1858) 343 = gigantea.

SERANGIUM, W. Wood, ex Salisb. Gen. Pl. Fragm. 5 (1866) = **Monstera**, Adans. (Aroid.).

SERAPHYTA, Fisch. & Mey. in Bull. Sc. Acad. Pétersb. viii. (1840) 24. *ORCHIDEAE*, Benth. & Hook. f. iii. 523.
multiflora, *Fisch. & Mey. l. c.* 25.—Mexic.

SERAPIAS, Linn. Syst. ed. I (1735); Gen. ed. I. 271 (1737). *ORCHIDEAE*, Benth. & Hook. f. iii. 620.
HELLEBORINE, Tourn. ex Hall. Enum. Stirp. Helv. i. 274 (1742), partim; Pers. Syn. ii. 512 (1807).
ISIAS, De Not. in Mem. Acc. Torin. Ser. II. vi. (1844) 413.
abortiva, Scop. Fl. Carn. ed. II. ii. 205 = Limodorum abortivum.
aculeata, Thunb. Prod. Pl. Cap. 3 = Eulophia aculeata.
alba, Salisb. Prod. 8 = Cephalanthera pallens.
anthropophora, Jundz. Fl. Lithuan. 267 = Aceras anthropophora.
aphylla, Vell. Fl. Flum. ix. t. 57 (Spirantheos sp.?).—Bras.
athensis, Hocquart, Fl. Jemmape, 236 = Epipactis microphylla.
athensis, Lej. Fl. Spa, ii. 196 = Orchis Morio.
atrorubens, Hoffm. Deutschl. Fl. ed. II. i. II. 182 = Epipactis atrorubens.
autumnalis, Poir. ex Steud. Nom. ed. I. 766 = Pterostylis cucullata.
biflora, Vell. Fl. Flum. ix. t. 50 (Spirantheos sp.).—Bras.
brevibracteata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Lingua.
camschatea, Steud. Nom. ed. I. 766 = Neottia camtschatica.
capensis, Linn. Mant. ii. 293 = Eulophia barbata.
Caravata, Aubl. Pl. Gui. ii. 816 = Elleanthus Caravata.
coccinea, Vell. Fl. Flum. ix. t. 48 = Spiranthes colorata, *N. E. Br.*
congesta, Vell. l. c. t. 54 (Spirantheos sp.).—Bras.
convallarioides, Steud. Nom. ed. I. 766 = Listera cordata.
cordata, Steud. l. c. = Listera cordata.
cordigera, Bertol. Fl. Ital. ix. 603, partim = neglecta.
cordigera, Bieb. Fl. Taur. Cauc. ii 370 = longipetala.
cordigera, Griseb. Spicil. Fl. Rumel. ii. 367 = S. Pseudo-cordigera.
cordigera, *Linn. Sp. Pl.* ed. II. 1345—Europ. austr.; Afr. bor.
cucullata, Pers. Syn. ii. 513 = Eriochylus autumnalis.
Damasonium, Mill. Gard. Dict. ed. VIII. n. 2 = Cephalanthera pallens.
ensifolia, Murr. Syst. ed. XIV. 815 = Cephalanthera ensifolia.
epidendraea, Retz. Obs. vi. 65 = Eulophia virens.
Epigogium, Steud. Nom. ed. I. 766, sphalm. = Epipogum aphyllum.
erecta, Thunb. Fl. Jap. 27 = Cephalanthera pallens.
falcata, Thunb. l. c. 28 = Cephalanthera falcata.
fasciculata, Vell. Fl. Flum. ix. t. 53 (Spirantheos sp.).—Bras.
flava, Sw. Prod. Veg. Ind. Occ. 119 = Corymbis flava.
Fontanae ×, Rigo & Goiran, in Nouv. Giorn. Bot. Ital. xv. (1883) 32 = purpurea.
Gavila, Pav. ex Steud. Nom. ed. II. ii. 567 = Chloraea Pavonii.
glabra, Lapeyr. Hist. Abr. Pl. Pyr. 552 = S. Lingua.
grandiflora, Linn. Syst. ed. XII. 594 = Cephalanthera pallens.
grandiflora, Oed. Fl. Dan. t. 506 = Cephalanthera ensifolia.
Helleborine, Linn. Sp. Pl. 949 = Cephalanthera ensifolia, Epipactis latifolia.
hirsuta, Lapeyr. Hist. Abr. Pl. Pyr. 551 = S. Pseudo-cordigera.
intermedia ×, Forest. ex F. Schultz, Arch. Fl. Fr. et Allem. 265.—Europ.
lancifera, St. Amans, Fl. Agen. 378 = S. Pseudo-cordigera.

SERAPIAS:—
lancifolia, Roth, Tent. Fl. Germ. ii. II. 409 = Cephalanthera pallens.
lancifolia, Steud. Nom. ed. II. ii. 567, sphalm. = *lancifera*.
latifolia, Huds. Fl. Angl. ed. I. 341 = Epipactis latifolia.
latifolia, O. F. Muell. in Fl. Dan. t. 811 = Epipactis atrorubens.
laxiflora, Chaub. in Bory & Chaub. Nouv. Fl. Pélop. 62 = S. Pseudo-cordigera.
laxiflora, Reichb. f. Ic. Fl. Germ. xiii-xiv. 13 = occultata.
Lingua, Bertol. Amoen. Ital. 202 = S. Pseudo-cordigera.
Lingua, *Linn. Sp. Pl.* 950.—Europ. austr.
Lingua, Merl. ex Steud. Nom. ed. II. ii. 567 = cordigera.
Lingua, Savi, Fl. Pisan. ii. 304 = neglecta.
Loeselii, Hoffm. Deutschl. Fl. ed. II. i. II. 181 = Liparis Loeselii.
Lonchophyllum, Linn. f. Suppl. 405 = Cephalanthera pallens.
longiflora, Asso, Syn. Stirp. Arag. 131 = Epipactis palustris.
longifolia, Huds. Fl. Angl. ed. I. 341 = Epipactis palustris.
longifolia, Scop. Fl. Carn. ed. II. ii. 202 = Cephalanthera ensifolia.
longipetala, Pollini, Fl. Veron. iii. 30 = S. Pseudo-cordigera.
lutea, Comm. ex Pers. Syn. ii. 513 = Chloraea Gaudichaudii.
melaleuca, Thunb. Prod. Pl. Cap. 3 = Disa melaleuca.
memoralis, Salisb. Prod. 8 = Epipactis latifolia.
microphylla, Ehrh. Beitr. iv. 42 = Epipactis microphylla.
neglecta, *De Not. Prosp. Fl. Ligust.* 55.—Ital.; Corsic.
neglecta ×, Forest. ex Nym. Consp. 690.—Europ.
Neotria, Raeusch. Nom. ed. III. 261, sphalm. = seq.
Neottia, J. F. Gmel. Syst. 59 = Spiranthes orchidioides.
Nidus-avis, Steud. Nom. ed. I. 766 = Neottia Nidus-avis.
nigra, *Gueldenst. Reisen*, ii. 56; *ex Ledeb. Fl. Ross.* 88 iv. (Quid ?).—Ross. austr.
nitida, Vell. Fl. Flum. ix. t. 52. (Spirantheos sp.).—Bras.
nivea, Vill. Hist. Pl. Dauph. ii. 52 = Cephalanthera ensifolia.
occultata, *F. Gay, in Ann. Sc. Nat.* Sér. II. vi. (1836) 119.—Europ. austr.
ochroleuca, Steud. Nom. ed. I. 766 = Cephalanthera pallens.
ovalis, Rich. in Mém. Mus. Par. iv. (1818) 54 = cordigera.
ovata, Steud. Nom. ed. I. 766 = Listera ovata.
oxyglottis, Reichb. Fl. Germ. Excurs. 130 = S. Pseudo-cordigera.
oxyglottis, Willd. Sp. Pl. iv. 71 = S. Lingua.
pallens, Jundz. Fl. Lithuan. 268 = Cephalanthera pallens.
pallida, Wahlenb. Fl. Carpat. 291 = Cephalanthera pallens.
palustris, Mill. Gard. Dict. ed. VIII. n. 3; Scop. Fl. Carn. ed. II. ii. 204 = Epipactis palustris.
parviflora, Parl. in Giorn. Sc. Lett. Sicil. lix. (1837) 66 = occulta.
patens, Thunb. Prod. Pl. Cap. 3 = Disa tenuifolia.
pedicellata, Thunb. l. c. = Eulophia aculeata.
polyaden, Vell. Fl. Flum. ix. t. 56. (Spirantheos sp.).—Bras.
polystachya, Sw. Prod. Veg. Ind. Occ. 119 (= *Chloidia vernalis*).—Jamaic.
porrifolia, Steud. Nom. ed. I. 767 = Microtis porrifolia.
Pseudo-cordigera, *Moric. Fl. Venet.* 374.—Europ. austr.
pubescens, Steud. Nom. ed. I. 767 = Ponthieva glandulosa.
pumila, Vell. Fl. Flum. ix. t. 55 (Spirantheos sp.).—Bras.
purpurea ×, Doumenj. Suppl. Herb. (1851) 54.—Gallia.
purpurea, Poir. Illustr. iii. 306 = Bletia verecunda.
reflexa, Pers. Syn. ii. 513 = Chiloglottis diphylla.
regularis, Forst. f. Prod. 59 = Thelymitra longifolia.
repens, Vill. Hist. Pl. Dauph. ii. 53 = Goodyera repens.
Roselliniana ×, Goiran, in Nouv. Giorn. Bot. Ital. xv. (1883) 33 = *Tommasinii*.
rubra, Linn. Syst. ed. XII. 594 = Cephalanthera rubra.
Salassia, Steud. Nom. ed. I. 767 = Liparis purpurascens.
speciosa, J. F. Gmel. Syst. 59 = Spiranthes colorata.

SERAPIAS:—
spiralis, Scop. Fl. Carn. ed. II. ii. 201 = Spiranthes autumnalis.
stricta, Welw. ex Woods, in Journ. Linn. Soc. ii. (1858) 116 = occulta.
sylvestris, Murr. ex Steud. Nom. ed. I. 766, 767 = Epipactis viridiflora.
tabularis, Thunb. Prod. Pl. Cap. 3 = Cymbidium tabulare.
Todari, *Tineo, Pl. Rar. Sicil.* 12.—Sicil.
tomentosa, Vell. Fl. Flum. ix. t. 51 (Spirantheos sp.).—Bras.
Tommasinii ×, Kern. in Verh. Zool.-Bot. Ges. Wien, xv. (1865) 231.—Ital.; Istria.
tota-alba, Gilib. Exercit. ii. 493 = Cephalanthera ensifolia.
triloba ×, Dupuy, ex Noul. Fl. Pyr. Suppl. 33.—Gallia.
triloba ×, Lloyd, Fl. Loire-Inf. ed. I. 255.—Gallia.
triloba, Viv. Fl. Ital. Fragm. 11, t. 12. f. 1.—Istria; Ital.
ventricosa, Vell. Fl. Flum. ix. t. 47 (Spirantheos sp.).—Bras.
viridiflora, Hoffm. Deutschl. Fl. ed. II. i. II. 182 = Epipactis viridiflora.
viridis, Vell. Fl. Flum. ix. t. 49 (Spirantheos sp.).—Bras.
Xiphophyllum, Linn. f. Suppl. 404 = Cephalanthera ensifolia.

SERENA, Rafin. Fl. Tellur. iv. 20 (1836) = **Haemanthus**, Tourn. (Amaryll.).
carnea, Rafin. l. c. = H. carneus.
lancifolia, Rafin. l. c. = H. lanceaefolius.

SERENAEA, Hook. f. in Benth. & Hook. f. Gen. iii. 926 (1883), sphalm. = seq.

SERENOA, Hook. f. in Benth. & Hook. f. Gen. iii. 1228, 926 (1883). *PALMAE*, Benth. & Hook. f. ll. cc.
serrulata, *Hook. f. ll. cc.*—Am. bor.

SEREREA, Rafin. Sylva Tellur. 107 (1838) = **Bignonia**, Tourn.
heterophyla, Rafin. l. c. = B. Kerere.

SERGILUS, Gaertn. Fruct. ii. 409. t. 174 (1791) = **Baccharis**, Linn. (Compos.).
scoparius, Gaertn. l. c. = B. scoparia.

SERIANA, Linn. Syst. ed. I (1735); Willd. Sp. Pl. ii. 464 (1799), sphalm. = seq.

SERIANIA, Plum. [Gen. t. 35] ex Schum. in Skrivt. Naturh. Selsk. Kjoeb. iii. II. (1794) 125 = **Serjania**, Plum. (Sapind.).

SERIANTHES, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 84, in nota = **Albizzia**, Durazz. (Legumin.).

SERIANTHES, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 225. *LEGUMINOSAE*, Benth. & Hook. f. i. 599.
calycina, *Benth. in Trans. Linn. Soc.* xxx. (1875) 600.—N. Caled.
grandiflora, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 225, partim.—Malacc.; Ins. Philip.
myriadenia, *Planch. ex Benth. l. c.* v. (1846) 108.—Ins. Pacif.
tenuiflora, *Benth. in Trans. Linn. Soc.* xxx. (1875) 599.—Hab. ?
vitiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 485.—Ins. Fiji.

SERICANDRA, Rafin. Sylva Tellur. 119 (1838) = **Albizzia**, Durazz. (Legumin.).
Julibrissin, Rafin. l. c. = A. Julibrissin.
lophantha, Rafin. l. c. = A. lophantha.
pennata, Rafin. l. c. = Acacia pennata.

SERICOBONIA, Linden & André, Illustr. Hortic. xxii. (1875) 39. t. 198 = **Jacobinia**?, Moric. (Acanth.).
ignea ×, Linden & André, l. c.

SERICOCARPUS, Nees, Gen. et Sp. Aster. 148 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 270. OLIGACTIS, Rafin. Fl. Tellur. ii. 44 (1836).
californicus, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 90.—Calif.
Collinsii, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 302 = tortifolius.

SERICOCARPUS:—
conyzoides, *Nees, Gen. et Sp. Aster.* 150.—Am. bor.
oregonensis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 302 = rigidus.
rigidus, *Lindl. in DC. Prod.* v. 261.—Am. bor. occ.
solidagineus, *Nees, Gen. et Sp. Aster.* 149.—Am. bor.
tortifolius, *Nees, l. c.* 151.—Am. bor.
Woodhousii, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457 = Bigelowia Wrightii.

SERICOCOMA, Fenzl, in Endl. Gen. Suppl. iii. 33 (1842). *AMARANTACEAE*, Benth. & Hook. f. iii. 30.
angustifolia, *Hook. f. in Benth. & Hook. f. Gen.* iii. 30.—Afr. austr.
avolans, *Fenzl, in Linnaea*, xvii. (1843) 328.—Afr. austr.
Bainesii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 31.—Afr. trop.
capensis, *Moq. in DC. Prod.* xiii. II. 307.—Afr. austr.
capitata, Moq. l. c. 308 = Calicorema capitata.
chrysurus, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 547 (err. typ, 447).—Afr. austr.
denudata, *Hook. f. in Benth. & Hook. f. Gen.* iii. 30.—Afr. trop.
pallida, *S. Moore, in Journ. Bot.* xv. (1877) 70.—Afric. trop.
pungens, *Fenzl, in Linnaea*, xvii. (1843) 326.—Afr. austr.
remotiflora, *Benth. & Hook. f. Gen.* iii. 30.—Afr. austr.
somalensis, S. Moore, in Journ. Bot. xv. (1877) 70 = Chionothrix somalensis.
trichinioides, *Fenzl, in Linnaea*, xvii. (1843) 324.—Afr. austr.
Welwitschii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 30.—Afr. trop.

SERICODES, A. Gray, Pl. Wright. i. 28. (in Smithson. Contrib. iii.) (1852). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 265.
Greggii, *A. Gray, Pl. Wright.* i. 28.—Mexic.

SERICOGRAPHIS, Nees, in Mart. Fl. Bras. ix. 107 ((1847) = **Jacobinia**, Moric. (Acanth.).
acuminata, Nees, l. c. 109.—Bras.
Aschenborniana, Nees, in DC. Prod. xi. 362 = J. Aschenborniana.
aurantiaca, Linden, Cat. n. 17 (1862) 6.—Mexic.
californica, A. Gray, in Bot. Mex. Bound. 125 = J. californica.
caripensis, Nees, in Hook. Lond. Journ. Bot. iv. (1845) 637 = Dianthera secunda.
Clausseniana, Nees, in Mart. Fl. Bras. ix. 111.—Bras.
colorata, Nees, in Benth. Pl. Hartw. 148.—Peruv.
cordata, Nees, in Mart. Fl. Bras. ix. 108.—Bras.
Ghiesbregtiana, Nees, in DC. Prod. xi. 730 = J. Ghiesbrechtiana.
haplostachya, Nees, l. c. 362 = Aschenborniana.
heterophylla, Nees, l. c. 365 = J. heterophylla.
hirsuta, Nees, in Mart. Fl. Bras. ix. 112.—Bras.
incana, Nees, in DC. Prod. xi. 361 = J. incana.
macrophylla, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 149 = J. macrophylla.
Mohintli, Nees, in DC. Prod. xi. 361 = J. Mohintli.
mollis, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 149.—Mexic.
monticola, Nees, in Mart. Fl. Bras. ix. 111 = J. monticola.
neglecta, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 151.—Mexic.
palustris, Nees, in Mart. Fl. Bras. ix. 110.—Bras.
parabolica, Nees, l. c. 111.—Bras.
pauciflora, Nees, l. c. 110.—Bras.
polita, Nees, l. c. 109 = J. polita.
rigida, Nees, l. c. 108 = Dianthera rigida.
scandens, Nees, l. c. 109.—Bras.
Selloviana, Nees, l. c. 111 = J. Selloviana.
squarrosa, Nees, in DC. Prod. xi. 364 = Dianthera nodosa.
tinctoria, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 150. t. 3. f. 15, 16 = J. tinctoria.
verrucosa, Oerst. l. c. 150.—Mexic.
vestita, Nees, in DC. Prod. xi. 364.—Peruv.

SERICOLA, Rafin. Sylva Tellur. 93 (1838) = **Miconia**, Ruiz & Pav. (Melastom.).
brachiata, Rafin. l. c. 94 = M. holosericea.

SERICOMA, Hochst. in Flora, xxviii. (1845) 73, sphalm. = **Sericocoma**, Fenzl (Amarant.).

SERICOSPORA, Nees, in DC. Prod. xi. 444 (1847). Genus dubium *ACANTHACEARUM*.—Cf. Benth. & Hook. f. Gen. ii. 1071.
crinita, *Nees, l. c.*—Ind. occ.

SERICOSTOMA, Stocks, in Wight, Ic. iv. II. 14. t. 1377 (1848). *BORAGINEAE*, Benth. & Hook. f. ii. 861.
albidum, *Franch. in Revoil, Çomal.* 46.—Afr. trop.
Kotschyi, *Benth. & Hook. f. Gen.* ii. 861.—Persia.
parviflorum, Walp. Ann. iii. 135, sphalm. = seq.
pauciflorum, *Stocks, in Wight, Ic.* iv. II. 15. t. 1377.—Sind.

SERICOTHECA, Rafin. Sylva Tellur. 152 (1838) = **Spiraea**, Linn. (Rosac.).
argentea, Rafin. l. c. = Sp. argentea.

SERICROSTIS, Rafin. Neogenyt. 4 (1825) = **Muehlenbergia**, Schreb. (Gramin.).

SERICURA, Hassk. in Flora, xxv. (1842) II. Beibl. 2 = **Pennisetum**, Rich. (Gramin.).
elegans, Hassk. l. c. = P. macrostachyum
japonica, Steud. in Flora, xxix. (1846) 20 = Phragmites communis.

SERIDIA, Juss. Gen. 173 (1789) = **Centaurea**, Linn. (Compos.).
Adami, Sweet, Hort. Brit. ed. I. 218 = C. solstitialis.
aspera, Sweet, l. c. = C. aspera.
auriculata, Sweet, l. c. = C. Seridis.
cruenta, Sweet, l. c. = C. cruenta?
ferox, Sweet, l. c. = C. ferox.
glomerata, Cass. in Dict. Sc. Nat. xlviii. 499 = C. glomerata.
Isnardi, Sweet, Hort. Brit. ed. I. 218 = C. aspera?
latifolia, Sweet, l. c. = C. Seridis.
longifolia, Willk. & Lange, Prod. Fl. Hisp. ii. 141, sphalm. = C. sonchifolia.
megacephala, Cass. in Dict. Sc. Nat. xlviii. 499 = C. Seridis.
melitensis, Sweet, Hort. Brit. ed. I. 218 = C. melitensis.
microcephala, Cass. in Dict. Sc. Nat. xlviii. 499 = aspera.
napifolia, Sweet, Hort. Brit. ed. I. 218 = C. napifolia.
polyacantha, Sweet, l. c. = C. polyacantha.
praetermissa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. aspera.
prolifera, Sweet, Hort. Brit. ed. I. 218 = C. glomerata.
pullata, Sweet, l. c. = C. pullata.
romana, Sweet, l. c. = C. romana.
sicula, Sweet, l. c. = C. sicula?
solstitialis, Sweet, l. c. = C. solstitialis.
sonchifolia, Sweet, Hort. Brit. ed. I. 218; Cass. in Dict. Sc. Nat. xlviii. 499 = C. sonchifolia.
sphaerocephala, Webb, Iter Hisp. 33 = C. sphaerocephala.
straminea, Sweet, Hort. Brit. ed. I. 218 = C. glomerata.
sulphurea, Sweet, l. c. = C. sulphurea?

SERIGROSTIS, Steud. Nom. ed. II. ii. 568 (1841), sphalm. = SERICROSTIS, Rafin. = **Muehlenbergia**, Schreb. (Gramin.).

SERINGEA, F. Muell. Fragm. x. 96 (1877) = seq.

SERINGIA, J. Gay, in Mém. Mus. Par. vii. (1821) 442. tt. 16, 17. *STERCULIACEAE*, Benth. & Hook. f. i. 226.
ACTINOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. (1859) I. 259.
GAYA, Spreng. Syst. i. 535, 971 (1825).
adenolasia, *F. Muell. Fragm.* x. 96.—Austral.
corollata, Steetz, in Lehm. Pl. Preiss. ii. 350 = Keraudrenia Hookeriana.
grandiflora, F. Muell. Fragm. i. 142 = Keraudrenia integrifolia.
Hillii, F. Muell. Census, 16 = Keraudrenia Hillii.
Hookeri, F. Muell. Fragm. ii. 5 = Keraudrenia Hookeriana.
integrifolia, F. Muell. l. c. = Keraudrenia integrifolia.
lanceolata, Steetz, in Lehm. Pl. Preiss. ii. 349 = Keraudrenia lanceolata.

SERINGIA:—
microphylla, F. Muell. Fragm. ii. 5 = Keraudrenia hermanniaefolia.
nephrosperma, F. Muell. in Hook. Kew Journ. ix. (1857) 15 = Keraudrenia nephrosperma.
platyphylla, *J. Gay, in Mém. Mus. Par.* vii. (1821) 443.—Austral.
velutina, F. Muell. Fragm. ii. 5 = Keraudrenia integrifolia.

SERINGIA, Spreng. Anleit. ii. 694 (1818); Syst. i. 441 (1825) = **Ptelidium**, Thou. (Celastr.).
lanceolata, Blanco, Fl. Filip. ed. II. 45 = Harpullia cupanioides.
ovata, Spreng. Syst. i. 441 = P. ovatum.

SERINIA, Rafin. Fl. Ludov. 149 (1817) = **Krigia**, Schreb. (Compos.).
cespitosa, Rafin. l. c. = K. virginica.

SERIOLA, Linn. Sp. Pl. ed. II. 1139 (1763) = **Hypochoeris**, Linn. (Compos.).
aethnensis, Linn. l. c. = H. aetnensis.
albicans, Tineo, Pl. Rar. Sicil. i. (1817) 20 = H. Achyrophorus.
Alliatae, Bivona, Sic. Pl. Cent. ii. 57 = H. Achyrophorus.
andina, Poepp. ex DC. Prod. vii. 92 = H. andina.
apargioides, Less. Syn. Comp. 131 (= *Achyrophorus Lessingii*).—Chili.
brasiliensis, Less. Syn. Comp. 131 = H. brasiliensis.
commutata, Less. l. c. = Rodigia commutata.
cretensis, Linn. Sp. Pl. ed. II. 1139 = H. cretensis.
depressa, Viv. Fl. Cors. Prod. App. ii. 4 = H. aetnensis.
glauca, Tineo, Pl. Rar. Sicil. i. (1817) 19 = H. Achyrophorus.
hieracioides, Raeusch. Nom. ed. III. 228, nomen.—Afr. bor.
hispida, Moench, Meth. 543 = H. aetnensis.
hyoseroides, Nym. Syll. 55 = Leontodon creticus.
incana, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 42.—Am. austr.
laevigata, Linn. Sp. Pl. ed. II. 1139 = H. Achyrophorus.
petiolaris, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 31 = H. petiolaris.
pubescens, Link, Handb. i. 794, sphalm. = seq.
rubescens, Tineo, Pl. Rar. Sicil. i. (1817) 21 = H. Achyrophorus.
scabra, Hort. ex Steud. Nom. ed. II. ii. 568, nomen.—Hab.?
taraxacifolia, Salzm. in Flora, iv. (1821) 111 = H. radicata.
taraxacoides, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 30 = H. Scorzonerae.
taraxacoides, Loisel. Fl. Gall. ed. I. i. 530. t. 18 = H. taraxacoides.
tenuiflora, Nym. Syll. 55 = H. tenuiflora.
tenuifolia, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 31 = H. tenuifolia.
Tweedii, Hook. & Arn. in Hook. l. c. = H. brasiliensis.
uniflora, Bivona, Sic. Pl. Cent. ii. 55 = H. taraxacoides.
urens, All. Fl. Pedem. i. 230. t. 29. f. 1 = H. aetnensis.
urens, Lam. Illustr. iii. 235 = Crepis aspera.
urens, Linn. Sp. Pl. ed. II. 1139 = H. cretensis.
urens, Viv. Prod. Fl. Cors. 14 = H. aetnensis.

SERIPHIUM, Linn. Sp. Pl. ed. I. 928 (1753) = **Stoebe**, Linn. (Compos.).
adpressum, DC. Prod. vi. 263 = St. adpressa.
aethiopicum, Steud. Nom. ed. I. 767 = St. aethiopica.
alopecuroides, Lam. Encyc. i. 271 = St. alopecuroides.
ambiguum, Murr. Syst. ed. XIII. 665 = St. artemisioides.
asperum, Pers. Syn. ii. 501 = Helichrysum ericaefolium.
candicans, Mundt, ex Harv. in Harv. & Sond. Fl. Cap. iii. 282 = St. phlaeoides.
capitatum, Less. Syn. Comp. 352 = St. capitata.
cernuum, Pers. Syn. ii. 501 = Elytropappus Rhinocerotis.
cinereum, Berg. Desc. Pl. Cap. 340 = St. artemisioides.
cinereum, Linn. Sp. Pl. 982 = St. cinerea.
corymbiferum, Linn. Mant. i. 119 = St. gomphrenoides.

SERIPHIUM:—
distichum, Lam. Encyc. i. 271 (= *Trichogyne seriphioides*).—Afr. austr.
ericoides, Pers. Syn. ii. 500 = Disparago ericoides.
fasciculatum, Pers. l. c. 501 (= *Trichogyne seriphioides*).—Afr. austr.
filagineum, DC. Prod. vi. 263 = Stoebe filaginea.
flavescens, DC. l. c. = St. spiralis.
fuscum, Linn. Sp. Pl. 928 = St. fusca.
gnaphaloides, Linn. Mant. i. 481 = St. gomphrenoides.
gomphrenoides, Lam. Encyc. i. 272 = St. gomphrenoides.
incanum, Lam. l. c. 271 = St. incana.
indicum, Pers. Syn. ii. 501 = St. incana.
juniperifolium, Lam. Encyc. i. 272 = St. aethiopica.
laricifolium, Lam. l. c. 271 (= *Trichogyne laricifolia*).—Afr. austr.
nervigerum, DC. Prod. vi. 263 = St. nervigera.
niveum, Pers. Syn. ii. 501 = Helichrysum niveum.
passerinoides, Lam. Encyc. i. 271 = St. passerinoides.
perotrichoides, Less. Syn. Comp. 352 = St. capitata.
phleoides, DC. Prod. vi. 262 = St. phlaeoides.
phylicoides, Pers. Syn. ii. 500 = St. phylicoides.
phyllostachyum, DC. Prod. vi. 262 = St. phlaeoides.
plumosum, Linn. Sp. Pl. 928 = St. cinerea.
prostratum, Lam. Encyc. i. 273 = St. prostrata.
reflexum, Pers. Syn. ii. 501 (= *Trichogyne reflexa*).—Afr. austr.
Rhinocerotis, Pers. l. c. = Elytropappus Rhinocerotis.
scabrum, Pers. l. c. 500 = Elytropappus glandulosus.
squarrosum, Forst. ex DC. Prod. vi. 262 = St. cinerea.
tenuifolium, Salisb. Prod. 212 = St. cinerea.
tomentosum, Moç. ex DC. Prod. vi. 264 = Olearia Forsteri?
uniflorum, Moç. ex DC. l. c. v. 77 = Vernonia salicifolia.
vermiculatum, DC. Prod. vi. 263 = St. artemisioides.
virgatum, Pers. Syn. ii. 501 = St. cinerea.

SERIS, Less. in Linnaea, v. (1830) 253. *COMPOSITAE*, Benth. & Hook. f. ii. 492.
amplexifolia, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 456.—Bras.
angustifolia, Gardn. l. c. 457 = Trichocline angustifolia.
denticulata, DC. Prod. vii. 20 = Trichocline polymorpha.
discoidea, *Less. in Linnaea*, v. (1830) 255.—Bras.
polymorpha, Gardn. in Hook. Ic. Pl. t. 501 = Trichocline arenaria.
polymorpha, Less. in Linnaea, v. (1830) 254 = Trichocline polymorpha.
polyphylla, *Baker, in Mart. Fl. Bras.* vi. III. 354.—Bras.
vaginata, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 456 = amplexifolia.

SERIS, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139 = **Onoseris**, DC. (Compos.).
onoseroides, Willd. l. c. = O. Isotypus.

SERISSA, Comm. ex Juss. Gen. 209 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 138.
DEMOCRITEA, DC. Prod. iv. 540 (1830).
DYSODA, Lour. Fl. Cochinch. 145 (1790).
capensis, Thunb. Nov. Gen. 131 = Plectronia ventosa.
ceylanica, Thw. Enum. Pl. Zeyl. 150 = Saprosma ceylanicum.
corymbosa, Bedd. Ic. Pl. Ind. Or. i. 3 = Saprosma corymbosum.
Democritea, *Baill. Hist. Pl.* vii. 386.—China.
fetida, *Lam. Illustr.* ii. 211.—China; Japon.
fragrans, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 50 = Saprosma fragrans.
Gardneri, Thw. Enum. Pl. Zeyl. 150 = Saprosma indicum.
glomerata, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 50 = Saprosma indicum.
japonica, Thunb. Nov. Gen. 132 = fetida.
myrtifolia, Blanco, Fl. Filip. ed. I. 164 = Randia densiflora, angatensis.
pinnata, Blanco, Fl. Filip. ed. I. 163 = Gardenia obscura.
scabrida, Thw. Enum. Pl. Zeyl. 151 = Saprosma scabridum.
ternata, Kurz, Andam. Rep. App. A. 40 = Saprosma ternatum.
Wightii, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 50 = Saprosma ceylonicum.

SERJANIA, Plum. [Gen. 34. t. 35 (SERIANIA)] ex Schum. in Skrivt. Naturh. Selsk. Kjoeb. iii. 11. (1794) 125. *SAPINDACEAE*, Benth. & Hook. f. i. 393.
acapulcensis, H. B. & K. Nov. Gen. et Sp. v. 108 = emarginata.
acoma, *Radlk. Monog. Serj.* 127.—Bras.
aculeata, *Radlk. l. c.* 146.—Bras.
acuminata, *Radlk. l. c.* 347.—Bras.
acuta, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 349.—Panama.
acutidentata, *Radlk. Monog. Serj.* 232.—Bras.
acutifolia, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 215 = oxyphylla.
adusta, *Radlk. Monog. Serj.* 156.—Venezuela.
albopunctata, Wright, ex Griseb. Cat. Pl. Cub. 44 = subdentata.
altissima, *Radlk. Monog. Serj.* 125.—Peruv.
Ampelopsis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 350.—N. Granat.
amplifolia, *Radlk. Mong. Serj.* 223.—N. Granat.
angustifolia, Hartw. Hort. Carlsruhan. (1825) 260 = mexicana.
angustifolia, *Willd. Sp. Pl.* ii. 466.—Ind. occ.
apiculata, Wright?, ex Radlk. Monog. Serj. 52, 110 = sinuata.
aquatica, *Vell. Fl. Flum.* 24; i. t. 53.—Bras.
Arnottiana, Walp. Rep. i. 412, sphalm. = Laruotteana.
atrolineata, *Wright, in Sauv. Fl. Cub.* 24.—Cuba.
australis, Spreng. Syst. ii. 248 = Dodonaea pinnata.
baramensis, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1180 = membranacea.
bignonioides, Klotzsch, l. c. = caracasana.
brachycarpa, *A. Gray, ex Radlk. Monog. Serj.* 259.—Texas.
brachyphylla, *Radlk. l. c.* 345.—N. Granat.
brachystachya, *Radlk. l. c.* 310.—Mexic.
bracteosa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 585 = caracasana.
brevipes, *Benth. Bot. Voy. Sulph.* 76.—Ecuador.
californica, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 222.—Calif.
Cambessedeana, *Schlecht. & Cham. in Linnaea*, v. (1830) 214.—Mexic.
caracasana, Seem. Bot. Voy. Herald, 93 = mucronulata.
caracasana, *Willd. Sp. Pl.* ii. 465.—Am. centr.
cardiospermoides, *Schlecht. & Cham. in Linnaea*, vi. (1831) 418.—Mexic.
chartacea, *Radlk. Monog. Serj.* 114.—Guian.
circumvallata, *Radlk. l. c.* 345.—N. Granat.
cirrhiflora, Sieber, ex Radlk. l. c. 352 = Urvillea ulmacea.
cissoides, *Radlk. l. c.* 294.—Bras.
clematidea, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 349.—N. Granat.
clematidifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 361.—Bras.
comata, *Radlk. Monog. Serj.* 131.—Bras.
communis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 362.—Bras.
confertiflora, *Radlk. Monog. Serj.* 117.—Bras.
cornigera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) 1. 267.—Panama.
corrugata, *Radlk. Monog. Serj.* 131.—Bras.
crassifolia, *Radlk. l. c.* 225.—Bolivia.
crassinervis, *Radlk. l. c.* 200.—Cuba.
crenata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 168.—Cuba.
crenata, Juss. ex H. B. & K. Nov. Gen. et Sp. v. 112, in obs. = subdentata.
cubensis, La Sagra, Hist. Econ. Polit. Cuba (1831), 353 = diversifolia.
cuneolata, *Radlk. Monog. Serj.* 128.—Ind. occ.
curassavica. *Radlk. l. c.* 311.—Ind. occ.
cuspidata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 356.—Bras.
cystocarpa, *Radlk. in Fest. Meth. Bot. Syst.* (1883) 32.—Mexic.
dasyclados, *Radlk. Monog. Serj.* 300.—N. Granat.
decemstriata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 223.—Reg. Argent.
deflexa, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 337.—Bras.
deltoidea, *Radlk. Monog. Serj.* 332.—Peruv.
dentata, *Radlk. l. c.* 144.—Bras.
dibotrya, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 35. t. 242.—Peruv.

SERJANIA:—
diffusa, *Radlk. Monog. Serj.* 302.—Bolivia.
divaricata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 168 = scatens.
divaricata, Schum. in Skrivt. Naturf. Selsk. Kjoeb. iii. 11. (1794) 126 = mexicana.
diversifolia, Radlk. Monog. Serj. 136 = lupulina.
Dombeyana, Juss. ex Cambess. in A. St. Hil. Fl. Bras. Mer. i. 364 = caracasana.
dubia, Spreng. Syst. ii. 247 = lucida.
dumicola, *Radlk. Monog. Serj.* 111.—Bolivia.
dura, *Radlk. l. c.* 134.—Bras.
elegans, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 358.—Bras.
elegans, Mart. in Flora, xx. (1837) 11. Beibl. 92 = corrugata.
emarginata, *H. B. & K. Nov. Gen. et Sp.* v. 109.—Mexic.
equestris, *Macfad. Fl. Jamaic.* i. 156.—Jamaic.
erecta, *Radlk. Monog. Serj.* 160.—Bras.
eucardia, *Radlk. l. c.* 121.—Bras.
exarata, *Radlk. l. c.* 166.—Bras.
faveolata, *Radlk. l. c.* 145.—Bras.
filicifolia, *Radlk. l. c.* 287.—Ind. occ.
floribunda, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 348 = mexicana.
foveata, *Griseb. in Goett. Abh.* xix. (1874) 108.—Tucuman.
fulta, Griseb. l. c. = glabrata.
fuscifolia, *Radlk. Monog. Serj.* 221.—Bras.
glabrata, Benth. Pl. Hartw. 115 = mucronulata.
glabrata, Griseb. ex Radlk. Monog. Serj. 165 = membranacea.
glabrata, *H. B. & K. Nov. Gen. et Sp.* v. 110.—Peruv.
glutinosa, *Radlk. Monog. Serj.* 126.—Bras.
goniocarpa, *Radlk. l. c.* 309.—Mexic.
gracilis, *Radlk. l. c.* 134.—Bras.
grammatophora, *Radlk. l. c.* 258.—Hab.?
grandiflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 360.—Bras.
grandiflora, Griseb. in Goett. Abh. xxiv. (1879) 79, partim = caracasana.
grandifolia, *Sagot, ex Radlk. Monog. Serj.* 166.—Guian.
grandis, *Seem. Bot. Voy. Herald*, 92.—Panama.
Grayii, Hemsl. Biol. Centr. Am. Bot. i. 206, sphalm. = seq.
Grosii, *Schlecht. in Linnaea*, xviii. (1844) 42 (sphalm. 58).—Mexic.
Guarumina, Mart. in Flora, xx. (1837) 11. Beibl. 91 = cuspidata.
Guaruminea, Walp. Rep. i. 413, sphalm. = praec.
hamuligera, *Radlk. Monog. Serj.* 339.—Bras.
hebecarpa, *Benth. in Hook. Kew Journ.* iii. (1851) 192.—Bras.
heterophylla, DC. Prod. i. 604 = mexicana.
hirsuta, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 367.—Bras.
ichthyctona, *Radlk. Monog. Serj.* 230.—Bras.
ilicifolia, Tausch, ex Opiz, in Oestr. Bot. Wochenbl. i. (1851) 339 = sinuata.
impressa, *Radlk. Monog. Serj.* 323.—Mexic.
incisa, *Torr. Bot. Mex. Bound.* 47.—Mexic.; Texas.
inebrians, *Radlk. Monog. Serj.* 346.—Costa Rica.
inflata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 36.—Peruv.
inflata, S. Wats. in Proc. Am. Acad. xvii. (1882) 337 = cystocarpa.
insignis, *Radlk. Monog. Serj.* 331.—Mexic.
lamprophylla, *Radlk. l. c.* 233.—Bras.
lanceolata, Cambess. in A. St. Hil. Fl. Bras. Mer. i. 356 = Urvillea ulmacea.
Laroutteana, D. Dietr. Syn. Pl. ii. 1317, sphalm. = seq.
Laruotteana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 368.—Bras.
laxiflora, *Radlk. Monog. Serj.* 135.—Bras.
leptocarpa, *Radlk. l. c.* 112.—Bras.
lethalis, *A. St. Hil. Pl. Rem. Brés.* i. 206, 235.—Bras.
longipes, *Radlk. Monog. Serj.* 256.—Ecuador.
lucida, Griseb. Fl. Brit. W. Ind. 123 = diversifolia.
lucida, *Schum. in Skrivt. Naturh. Selsk. Kjoeb.* iii. 11. (1794) 128.—Ins. S. Doming.
lupulina, Benth. Bot. Voy. Sulph. 76 = Paullinia velutina.
lupulina, Griseb. Cat. Pl. Cub. 44 = crassinervis.
lupulina, *Schum. in Skrivt. Nat. Selsk.* iii. 11. (1794) 127.—Ind. occ.

SERJANIA:—
macrococca, *Radlk. Monog. Serj.* 270.—Mexic.
macrostachya, *Radlk. l. c.* 135.—Bras.
Mansiana, *Mart. in Flora*, xxii. (1839) 1. *Beibl.* 9.—Bras.
marginata, *Casar. Nov. Stirp. Bras. Dec.* v. 44.—Bras.
mariquitensis, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 347 = caracasana.
megalocarpa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1858) 1. 396 = Toulicia megalocarpa.
membranacea, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 105.—Guian.
meridionalis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 366.—Bras.; Uruguay.
meridionalis, Griseb. in Goett. Abh. xxiv. (1879) 79 = hebecarpa.
mexicana, Seem. Bot. Voy. Herald, 92 = Paullinia velutina.
mexicana, *Willd. Sp. Pl.* ii. 465.—Am. centr.
micrantha, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1180 = oblongifolia.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 108.—Peruv.
mollis, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 345 = grandis.
mollis, Willd. ex Schlecht. in Linnaea, xviii. (1844) 45 (sphalm. 61) = mexicana.
monogyna, Hoffmgg. ex Schlecht. l. c. 30 (sphalm. 55) = Paullinia monogyna.
Moritziana, Schlecht. l. c. 44 (sphalm. 60) = Urvillea ulmacea.
mucronulata, *Radlk. Monog. Serj.* 282.—Ecuador.
multiflora, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 365.—Bras.
nigricans, *Radlk. Monog. Serj.* 336.—Bras.
nitidula, Benth. in Hook. Kew Journ. iii. (1851) 191 = caracasana.
nodosa, *Radlk. Monog. Serj.* 341.—Ind. occ.
noxia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 363.—Bras.
nutans, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 36. t. 242.—Peruv.
oblongifolia, *Radlk. Monog. Serj.* 219.—Guian.
obtusidentata, *Radlk. l. c.* 233.—Bras.
orbicularis, *Radlk. l. c.* 292.—Bras.
Ossana, DC. Prod. i. 603 = diversifolia.
Ossana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 396 = subdentata.
ovalifolia, *Radlk. Monog. Serj.* 218.—Bras.
oxypetala, G. Don, Gen. Syst. i. 659, sphalm. = oxyphylla.
oxyphylla, *H. B. & K. Nov. Gen. et Sp.* v. 112.—Peruv.
oxytoma, *Radlk. Monog. Serj.* 234.—Bras.
paleata, *Radlk. l. c.* 130.—Bras.
paludosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 368.—Bras.
paniculata, Griseb. Fl. Brit. W. Ind. 123 = equestris.
paniculata, *H. B. & K. Nov. Gen. et Sp.* v. 111. t. 441.—Mexic.
paniculata, Poepp. ex Presl, Bot. Bemerk. 25 = subdentata.
paniculata, Seem. Bot. Voy. Herald, 92 = mexicana.
paniculata, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 348 = insiguis.
paradoxa, *Radlk. Monog. Serj.* 133.—Bras.
parviflora, DC. Prod. i. 604 (sphalm.) = parvifolia.
parvifolia, *H. B. & K. Nov. Gen. et Sp.* v. 113.—Peruv.
paucidentata, *DC. Prod.* i. 603.—Bras.
paucidentata, Seem. Bot. Voy. Herald, 92 = rhombea.
pedicellaris, *Radlk. Monog. Serj.* 124.—Guian.
perulacea, *Radlk. l. c.* 227.—Bras.
pinnatifolia, *Radlk. l. c.* 226.—Bras.
piscatoria, *Radlk. l. c.* 340.—Bras.
platycarpa, *Benth. in Hook. Kew Journ.* iii. (1851) 192.—Bras.
plicata, *Radlk. Monog. Serj.* 167.—Mexic.
Plumeriana, Spreng. Syst. ii. 247 = sinuata.
polyphylla, Bert. ex Colla, Hort. Ripul. 101 = filicifolia.
polyphylla, *Poir. ex Steud. Nom.* ed. II. ii. 568.—Guian.
polyphylla, Willd. ex Schlecht. & Cham. in Linnaea, v. (1830) 215; Radlk. Monog. Serj. 179 = lucida.
polystachya, *Radlk. l. c.* 276.—Mexic.
pubescens, Griseb. in Bonplandia, vi. (1858) 3 = rhombea.

SERJANIA:—
pubescens, H. B. & K. Nov. Gen. et Sp. v. 110 = mexicana.
pubescens, Seem. Bot. Voy. Herald, 92 = Paullinia velutina.
purpurascens, *Radlk. Monog. Serj.* 335.—Bras.
pyramidata, *Radlk. l. c.* 155.—Peruv.
racemosa, Hook. & Arn. Bot. Beech. Voy. 413 = triquetra.
racemosa, *Schum. in Skrivt. Naturh. Selsk. Kjoeb.* iii. II. (1794) 127.—Am. centr.
racemosa, Seem. in Bonplandia, v. (1857) 74 = goniocarpa.
Regnellii, *Schlecht. in Linnaea*, xviii. (1844) 41 (sphalm. 57).—Bras.
reticulata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 359.—Bras.
rhombea, *Radlk. Monog. Serj.* 324.—Venezuela.
rigida, *Radlk. l. c.* 283.—Bolivia.
rubens, Schlecht. in Linnaea, xviii. (1844) 63 = reticulata.
rubicaulis, Benth. ex Radlk. Monog. Serj. 254 = seq.
rubifolia, *Klotzsch, ex Trev. in Bot. Zeit.* v. (1847) 393.—Peruv.
rufa, *Radlk. Monog. Serj.* 324.—Peruv.
Salzmanniana, *Schlecht. in Linnaea*, xviii. (1844) 46 (sphalm. 62).—Bras.
Salzmanniana, Seem. Bot. Voy. Herald, 92 = acuta.
samydea, *Griseb. in Bonplandia*, vi. (1858) 3.—N. Granat.
samydea, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 346 = rhombea.
scatens, *Radlk. Monog. Serj.* 213.—Mexic.
Schiedeana, *Schlecht. in Linnaea*, xviii. (1844) 44 (sphalm. 60).—Mexic.
scopulifera, *Radlk. Monog. Serj.* 229.—Bras.
Seemanni, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 346.—Panama.
Selloviana, Klotzsch, ex Trev. in Bot. Zeit. v. (1847) 393 = Laruotteana.
serrata, *Radlk. Monog. Serj.* 293.—Bras.
setulosa, *Radlk. l. c.* 337.—Mexic.
sinuata, Hook. Bot. Misc. iii. (1833) 159 = Urvillea uniloba.
sinuata, Presl, Bot. Bemerk. 25 = Urvillea ulmacea.
sinuata, *Schum. in Skrivt. Naturh. Selsk. Kjoeb.* iii. II. (1794) 126.—Ins. S. Doming.
sordida, *Radlk. Monog. Serj.* 272.—Mexic.
spectabilis, Schum. in Skrivt. Naturh. Selsk. Kjoeb. iii. II. (1794) 127 = mexicana.
sphaerococca, *Radlk. Monog. Serj.* 153.—Peruv.
sphenocarpa, *Radlk. l. c.* 269.—Mexic.
squarrosa, *Radlk. l. c.* 115.—Peruv.
stans, Schott, in Spreng. Syst. iv. App. 405 = Toulicia stans.
stenopterygia, *Radlk. Monog. Serj.* 110.—Bras.
striata, *Radlk. l. c.* 280.—Peruv.
subdentata, *Juss. ex Poir. Encyc.* v. 102.—Cuba.
subimpunctata, *Radlk. Monog. Serj.* 123.—Bras.
subtriplinervis, *Radlk. l. c.* 273.—Mexic.
sufferruginea, *Radlk. l. c.* 299.—Peruv.
tenuifolia, *Radlk. l. c.* 220.—Bras.
tenuis, *Radlk. l. c.* 98.—Bras.
thoracoides, *Radlk. l. c.* 170.—Bras.
tinctoria, *Sandifort, Elench. Lugd. Bat.* (1822) 31.—Java.
trachygona, *Radlk. Monog. Serj.* 327.—Mexic.
trichomisca, *Radlk. l. c.* 114.—Bras.
triquetra, *Radlk. l. c.* 305.—Mexic.
tristis, *Radlk. l. c.* 293.—Bras.
triternata, Seem. Bot. Voy. Herald, 92 = trachygona.
triternata, Willd. Sp. Pl. ii. 466 = lucida.
velutina, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 357. t. 75.—Bras.
velutina, Herb. ex Radlk. Monog. Serj. 305 = triquetra.
vesicosa, *Radlk. l. c.* 277.—Mexic.

SEROPHYTON, Benth. Bot. Voy. Sulph. 52 (1844) = **Argithamnia**, Sw. (Euphorb.).
Drummondi, Benth. l. c. 53 = A. mercurialina.
lanceolatum, Benth. l. c. 52 = A. lanceolata.
pilosissimum, Benth. l. c. 53 = A. pilosissima.

SERPAEA, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 296 = **Oyedaea**, DC. (Compos.).
oblonga, Gardn. l. c. 395 = Aspilia oblonga.
ovata, Gardn. l. c. 296 = O. ovata.
rotundifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 208 = O. rotundifolia.

SERPICULA, Linn. Mant. i. 16 (1767). *HALORAGEAE*, Benth. & Hook. f. i. 675.
EPILITHES, Blume, Bijdr. 734 (1826).
LAUREMBERGIA, Berg. Desc. Pl. Cap. 350 (1767).
brasiliensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 250.—Bras.
brevipes, Wight & Arn. Prod. 338 = indica.
capensis, Raeusch. Nom. ed. III. 271 = repens.
Epilithes, *Blume, Mus. Bot. Lugd. Bat.* i. 110.—Java.
hirsuta, Wight & Arn. Prod. 338 = indica.
indica, *Thw. Enum. Pl. Zeyl.* 123, partim.—Ind. or.
indica, Thw. l. c., partim = zeylanica.
javanica, *Mig. Fl. Ind. Bat.* i. 1. 1090.—Java.
repens, *Linn. Mant.* i. 124.—Afr. austr.
rubicunda, Burch. Trav. i. 37 = repens.
serpyllifolia, Burm. f. Fl. Cap. Prod. 30 (err. typ. 26) = repens.
thymifolia, *Burm. l. c.*—Afr. austr.
veronicaefolia, *Bory, Voy.* iii. 174.—Ins. Borbon.
zeylanica, *Arn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 431.—Zeylan.

SERPICULA, Linn. f. Suppl. 416 (1781), partim = **Hydrilla**, Rich. (Hydrochar.).
verticillata, Linn. f. l. c. = H. ovalifolia.

SERPICULA, Pursh, Fl. Am. Sept. i. 33 (1817) = **Elodea**, Michx. (Hydrochar.).
canadensis, Eaton, Man. (1833) 336 = E. canadensis.
occidentalis, Pursh, Fl. Am. Sept. i. 33 = E. canadensis.
verticillata, Rostk. & Schmidt, Fl. Sedin. 370 = E. canadensis.

SERPILLARIA, Heist. Syst. 6 (1748) (Quid?).

SERPYLLUM, Tourn. ex Rupp. Jen. ed. Hall. 234 (1745) = **Thymus**, Linn. (Labiat.).
angustifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 133 = T. Serpyllum.
Chamaedrys, Fourr. l. c. = T. Serpyllum.
citri-odora, Pall. Reise Suedl. Statth. Russ. Reich. ii. 80 = T. Serpyllum.
lanuginosum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 133 = T. Serpyllum.
vulgare, Fourr. l. c. = T. Serpyllum.

SERRA, Cav. Diss. ii. 83 (1786), sphalm. = **Senra**, Cav. (Malvac.).

SERRAEA, Spreng. Syst. iii. 78 (1826) = **Senra**, Cav. (Malvac.).
incana, Spreng. l. c. = Senra incana.
rupestris, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 168 = Senra incana.

SERRAFALCUS, Parl. Rar. Pl. Sic. ii. 14 (1840); Pl. Nov. 75 (1842) = **Bromus**, Linn. (Gramin.).
alopecuroides, Parl. Fl. Ital. i. 399 = B. Alopecuros.
arduennensis, Crép. Man. Bot. Belg. ed. I. 211 = B. arduennensis.
arvensis, Godr. Fl. Lorr. ed. I. iii. 185 = B. arvensis.
Cavanillesii, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 101 = B. molliformis.
Chiapporianus, De Not. ex Parl. Fl. Ital. i. 394 = B. patulus.
commutatus, Bab. Man. Brit. Bot. ed. I. 374; Godr. Fl. Lorr. ed. I. iii. 184 = B. racemosus.
contortus, Parl. in Giorn. Bot. i. 1. (1844) 326 = B. Alopecuros.
divaricatus, Dulac, Fl. Hautes-Pyr. 97 = B. secalinus.
hordeaceus, Gren. & Godr. Fl. Fr. iii. 590 = B. racemosus?
Hughi, Todaro, Ind. Sem. Hort. Panorm. (1866) 33 = B. intermedius.
intermedius, Parl. Rar. Pl. Sic. ii. 17 = B. intermedius.
lanceolatus, Parl. l. c. 18 = B. macrostachys.
Lloydianus, Gren. & Godr. Fl. Fr. iii. 591 = B. molliformis.
macrostachys, Parl. Fl. Ital. i. 397 = B. macrostachys.
massiliensis, Gren. in Mém. Soc. Emul. Doubs, Sér. III. iv. (1860) 391 = P. massiliensis.
molliformis, F. Schultz, Arch. Fl. 320 = B. molliformis.
mollis, Parl. Rar. Pl. Sic. ii. 16 = B. mollis.
neglectus, Parl. Fl. Ital. i. 391 = B. racemosus.

SERRAFALCUS:—
patulus, Parl. Fl. Ital. i. 394 = Bromus patulus.
racemosus, Parl. Rar. Pl. Sic. ii. 14 = B. racemosus.
scoparius, Parl. l. c. 19 = B. scoparius.
secalinus, Bab. Man. Brit. Bot. ed. I. 374; Godr. Fl. Lorr. ed. I. iii. 182 = B. secalinus.
squarrosus, Bab. l. c. 375 = B. squarrosus.
velutinus, Parl. Rar. Pl. Sic. ii. 16 = B. racemosus.

SERRARIA, Burm. Afr. 204. t. 99 (1737) = **Serruria**, Salisb. (Proteac.).

SERRATULA, Dill. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 475.
CRUPINASTRUM, Schur, in Ber. Bot. Rundsch. (1853) 37; ex ej. Enum. Pl. Transs. 411.
KLASEA, Cass. in Dict. Sc. Nat. xxxv. 17 (1825); et xli. 321.
MASTRUCIUM, Cass. l. c. (1825).
PEREUPHORA, Hoffmgg. Verz. Pfl. Nachtr. ii. 173 (1826).
PSILOSANTHUS, Neck. Elem. i. 69 (1790).
STEMMACANTHA, Cass. in Bull. Soc. Philom. (1817) 12.
acaulis, DC. Diss. Comp. ii. 31 = Centaurea chamaerhaponticum.
acutifolia, Poir. Encyc. vi. 554 = Lucilia acutifolia.
alata, Desf. Tabl. ed. II. 108 = Jurinea alata.
alata, S. T. Gmel. Reise, i. 155. t. 35. f. 2; ex Ledeb. Fl. Ross. ii. 733 (= *Cirsium elodes*?).—Reg. Caucas.
alata, Poir. Encyc. vi. 553 = Jurinea cyanoides.
alata, Sweet, Hort. Brit. ed. I. 213 = Saussurea japonica.
alatavica, *C. A. Mey. ex Rupr. Sert. Tiansch.* 56.—Turkest.
albescens, Wall. Cat. n. 2918 = Saussurea albescens.
albida, DC. ex Pers. Syn. ii. 391 = Heterocoma albida.
Alcalae, *Coss. Notes Crit.* 40.—Hispan.
alpina, Linn. Sp. Pl. 816 = Saussureae sp. variae.
altissima, DC. Prod. vi. 671 = Vernonia altissima.
altissima, Steph. ex Ledeb. Fl. Ross. ii. 761, in nota = coronata.
amara, Linn. Sp. Pl. 819 = Saussurea amara.
amara, Pall. Reise, i. 500 = Centaurea Picris.
ambigua, DC. Diss. Comp. ii. 30 = Jurinea ambigua.
angulata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 453.—Soongar.
angustifolia, Willd. Sp. Pl. iii. 1642 = Saussurea alpina.
anthelmintica, Roxb. Hort. Beng. 60; Fl. Ind. iii. 405 = Vernonia anthelmintica.
apiculata, Poir. Encyc. Suppl. v. 43 = Staehelina apiculata.
arborescens, Lam. Fl. Fr. ii. 40 = Staehelina arborescens.
arguta, Fisch. ex Hornem. Suppl. Hort. Hafn. 156 = tinctoria.
arvensis, Linn. Sp. Pl. 820 = Cnicus arvensis.
aspera, *Link, Enum. Hort. Berol.* ii. 297.—Nepal.
atriplicifolia, *Benth. & Hook. f. Gen.* ii. 475.—Dahuria.
autaretica, Poir. Encyc. vi. 563 = Cnicus heterophyllus.
babylonica, Linn. Syst. ed. X. 1199 = Centaurea babylonica.
bachtiarica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 587.—Persia.
Barrelieri, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 156 = pinnatifida.
Behen, Lam. Illustr. iii. 242. t. 666 = Centaurea cerinthifolia.
bifrons, Pers. Syn. ii. 391 = Chronopappus bifrons.
blanda, Bieb. Fl. Taur. Cauc. iii. 549 = Jurinea arachnoidea.
Bocconi, Guss. Ind. Sem. Hort. Boccad. (1825); Tenore, Succ. Rel. Viagg. Abruzzo, 84 = Jurinea humilis.
boetica, *Boiss. ex DC. Prod.* vii. 306.—Hispan.
caespitosa, Steud. Nom. ed. II. ii. 569, sphalm. = Saussurea taraxacifolia.
campestris, Schweigg. ex DC. Prod. vi. 671 = Cnicus arvensis.
canariensis, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 370.—Ins. Canar.
carniolica, Poir. Encyc. vi. 563 = Cnicus rufescens.
caroliniana, Mill. Gard. Dict. ed. VIII. n. 7 = Vernonia oligophylla.
carthamoides, Buch.-Ham. ex Roxb. Hort. Beng. 60; Fl. Ind. iii. 407 = Saussurea affinis.

SERRATULA:—
carthamoides, Poir. Encyc. vi. 561 = Centaurea carthamoides.
caspia, Pall. Reise, ii. 743 (= *Karelinia caspia*).—Reg. Casp.
caspica, Willd. Sp. Pl. iii. 1643 = praec.
caucasica, *Boiss. Fl. Orient.* iii. 590.—Reg. Caucas.
caulescens, Coss. in Bull. Soc. Bot. Fr. xx. (1873) 251 = Centaurea Cossoniana.
centauroides, Bieb. Fl. Taur. Cauc. ii. 264 = radiata.
centauroides, *Linn. Sp. Pl.* 820.—Sibir.
cerinthefolia, *Sibth. & Sm. Fl. Graec. Prod.* ii. 567.—Syria; Persia.
cernua, Poir. Encyc. vi. 560 = Carduus cernuus.
chaetocarpa, Ledeb. Fl Alt. iv. 42 = Jurinea chaetocarpa.
Chamaepeuce, Linn. Sp. Pl. 819 = Cnicus Chamaepeuce.
chinensis, *S. Moore, in Journ. Bot.* xiii. (1875) 228.—China.
cichoracea, *DC. Prod.* vi. 670.—Ital.
ciliata, Vahl, Symb. Bot. i. 67; Bieb. Fl. Taur. Cauc. ii. 267 = Cnicus arvensis.
cinerea, Roxb. Hort. Beng. 60; Fl. Ind. iii. 406 = Vernonia cinerea.
cirsioides, *Tenore, Succ. Rel. Viagg. Abruzzo*, 85.—Europ. austr.
complanata, Schweigg. in Hagen, Chlor. Bor. 303 = Cnicus arvensis.
complicata, Koch, Syn. Fl. Germ. ed. I. 400, sphalm. = praec.
compta, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 508 = Liatris spicata.
conica, Lam. Fl. Fr. ii. 40 = Staehelina dubia.
conifera, Brot. Phyt. Lusit. i. 67. t. 31 = Leuzea conifera.
cordata, Cass. in Dict. Sc. Nat. l. 468 = cerinthefolia.
coriacea, *Fisch. & Mey. ex DC. Prod.* vi. 668.—Armen.; Persia.
coronaria, Siev. in Pall. N. Nord. Beitr. vii. 345 = coronata.
coronata, DC. Fl. Fr. iv. 85 (excl. desc.); Suppl. 458 = tinctoria.
coronata, *Linn. Sp. Pl.* ed. II. 1144.—Sibir.
corymbosa, Poir. Encyc. vi. 559 = Vernonia corymbosa.
crupina, Vill. Hist. Pl. Dauph. iii. 38 = Crupina vulgaris.
cyanoides, DC. Diss. Comp. ii. 30 = Jurinea cyanoides.
cyanoides, Gaertn. Fruct. ii. t. 162 = Jurinea mollis.
cynarifolia, Poir. Encyc. vi. 561 = Centaurea carthamoides.
cynaroides, DC. Fl. Fr. iv. 87 = Cnicus cynaroides.
cynoglossifolia, Willd. ex Spreng. Syst. iii. 381 = Saussurea serrata.
daurica, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 251 = Saussurea crassifolia.
decurrens, Ledeb. Ind. Sem. Hort. Dorpat. (1811) 4 = Saussurea japonica.
dentata, Poir. Encyc. vi. 562 = Carthamus dentatus.
depressa, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 52.—Kokan.
depressa, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 61 = Jurinea depressa.
discolor, Poir. Encyc. vi. 565 = Cnicus altisimus.
discolor, Willd. Sp. Pl. iii. 1641 = Saussurea discolor.
dissecta, *Ledeb. Fl. Alt.* iv. 40.—Sibir. altaic.
dissecta, Steph. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 46 = Saussurea japonica.
divaricata, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 37 = Volutarella divaricata.
dubia, Brot. Fl. Lusit. i. 350 = Staehelina dubia.
dubia, Scop. ex Steud. Nom. ed. II. ii. 569 (Quid?).—Am. bor.
echinocephala, Poir. Encyc. vi. 561 = Cnicus echinocephalus.
elegans, Ait. Hort. Kew. ed. I. iii. 138 = Liatris elegans.
elegans, Stev. in Trans. Linn. Soc. xi. (1815) 414. t. 37 = Jurinea elegans.
fasciculata, Poir. Encyc. vi. 560 = Lachnospermum ericifolium.
flavescens, *Poir. l. c.* 562.—Hispan.
flexicaulis, *Rupr. Sert. Tiansch.* 57.—Turkest.
foliosa, Hill, ex Steud. Nom. ed. II. ii. 570 (Quid?).—Am. bor.
frigida, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 37 = Saussurea alpina.
fruticosa, Lam. Illustr. iii. 242 = Staehelina fruticosa.

SERRATULA:—
furcata, Spreng. ex DC. Prod. vi. 563 = Tricholepis furcata.
glandulifera, *Turcz. ex DC. l. c.* 670.—Rossia.
glauca, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 560.—Reg. Caucas.; Sibir.
glauca, Linn. Sp. Pl. 818 = Vernonia noveboracensis.
Gmelini, *Ledeb. ex DC. Prod.* vi. 668.—Sibir.
Gmelini, Tausch, in Flora, xi. (1828) II. 485 = radiata.
gnaphaloides, Poir. Encyc. vi. 559 = Helipterum gnaphaloides.
gossypiphora, *D. Don, Prod. Fl. Nep.* 168.—Reg. Himal.
gracilis, Hill, ex Steud. Nom. ed. II. ii. 570 (Quid?).—Am. bor.
graeca, Spreng. Syst. iii. 388 = Jurinea glycacantha.
Griffithii, Boiss. Diagn. Ser. II. iii. 49 = Volutarella divaricata.
hastata, Poir. Encyc. vi. 557 = Vernonia atriplicifolia.
Hausknechtii, *Boiss. Fl. Orient.* iii. 589.—Persia.
heleniifolia, *Sch. Bip. ex Reichb. Ic. Fl. Germ.* xv. 47.—Europ. centr.
heterophylla, *Desf. Tabl.* ed. I. 93.—Europ.; Reg. Caucas.
heterophylla, Vill. Cat. Strasb. 198 = radiata.
humilis, Desf. Fl. Atlant. ii. 244 = Jurinea humilis.
ilicifolia, Poir. Encyc. vi. 558 = Gynoxys ilicifolia.
imbricata, Poir. l. c. 559 = Metalasia uniflora.
incana, Gmel. It. i. 155 = Cnicus arvensis.
indica, Klein, ex Willd. Sp. Pl. iii. 1642 = Goniocaulon glabrum; Tricholepis glaberrima.
indica, Russ. ex Wall. Cat. n. 2985 F = Echinops echinatus.
indivisa, Poir. ex DC. Prod. vi. 667 = tinctoria.
inermis, Gilib. Fl. Lituan. i. 183 = tinctoria.
inermis, Poir. Encyc. vi. 564 = Cnicus centauroides.
integrifolia, *Krock. Fl. Siles. Suppl.* ii. 277.—Siles.
isophylla, *Claus, in Beitr. Pfl. Russ. Reich.* viii. 118, 301.—Ross. med.
japonica, Thunb. Fl. Jap. 305 = Saussurea japonica.
Kitaibelii, G. Don, ex Loud. Hort. Brit. 329 = heterophylla.
Kotschyi, *Boiss. Fl. Orient.* iii. 590.—Kurdist.
lanata, Poir. Encyc. vi. 565 = Cnicus arvensis.
latifolia, *Boiss. Diagn.* Ser. I. x. 96.—Persia.
Laxmanni, Fisch. ex DC. Prod. vi. 669 = glauca.
leucantha, DC. l. c. 670 = flavescens.
leucocephala, Poir. Encyc. vi. 565 = Cnicus leucocephalus.
liatroides, Adams, ex DC. Prod. vi. 532 = Saussurea pycnocephala.
linearifolia, DC. Diss. Comp. ii. 29 = Jurinea linearifolia.
lucida, Poir. Encyc. vi. 555 (= *Cirsium lucidum*).—Hab.?
lyrata, Bess. Cat. Hort. Crem. (1816) 129 = Jurinea alata.
lyratifolia, *Scheele, in Linnaea*, xviii. (1844) 481.—Reg. Caucas.
lyratifolia, *Schrenk, Enum. Pl. Nov.* i. 45.—Sibir. altaic.
macrocephala, *Bertol. Fl. Ital.* viii. 604.—Ital.
macrocephala, Wall. Cat. n. 2904 = Jurinea macrocephala.
macrophylla, Wender. ex Steud. Nom. ed. II. ii. 570 (Quid?).—Hab.?
marginata, Tausch, in Flora, xi. (1828) 484 = Gmelini.
marilandica, Crantz, Inst. i. 259 = glauca.
media, Poir. Encyc. vi. 564 = Cnicus medius.
melanocheila, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 589.—Persia.
mollis, Cav. Ic. i. 62. t. 90. f. 1 = Jurinea humilis.
mollis, Koch, Syn. Fl. Germ. ed. I. 408 = Jurinea mollis.
mollis, Wallr. Sched. Crit. 452 = praec.?
Monardi, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 155 = Jurinea Monardi.
montana, Poir. Encyc. vi. 564 = Cnicus montanus.
monticola, Bor. Fl. Centr. Fr. ed. III. ii. 364 = macrocephala.
mucronata, Desf. Fl. Atlant. ii. 243 = flavescens.
multicaulis, Wall. Cat. n. 2897 = Saussurea affinis.
multiflora, Blanco, Fl. Filip. ed. I. 617 = Vernonia chinensis.
multiflora, Linn. Sp. Pl. 817 = Jurinea linearifolia.
multiflora, *Lour. Fl. Cochinch.* 483.—China.

SERRATULA:—
multiflora, Pall. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) 20 = Saussurea serrata.
nemorum, Lavy, Orig. 13; ex DC. Prod. vi. 671 (Quid?).—Hab.?
nitens, Spreng. Syst. iii. 389 = Centaurea nitens.
nitida, Bess. Enum. Pl. Volh. 32 = heterophylla.
nitida, *Fisch. ex Spreng. Syst.* iii. 390.—Rossia; Sibir.
nivea, Wall. Cat. sub n. 2898 = Saussurea deltoidea.
noveboracensis, Linn. Sp. Pl. 818 = Vernonia noveboracensis.
nudicaulis, *DC. Fl. Fr.* iv. 86.—Gallia; Hispan.
numidica, Poir. Encyc. vi. 555 (= *Cirsium numidicum*).—Afr. bor.
oleracea, Neck. in Act. Acad. Theod. Palat. ii. (1770) 465 = Cnicus oleraceus.
oligocephala, *DC. Prod.* vi. 669.—As. Min.; Syria.
orientalis, Poir. Encyc. vi. 565 = Galactites tomentosa.
pallida, *DC. Prod.* vi. 670.—Reg. Himal.
palustris, Neck. in Act. Acad. Theod. Palat. ii. (1770) 465 = Cnicus palustris.
pannonica, Spreng. Syst. iii. 389 = Cnicus pannonicus.
parviflora, Poir. Encyc. vi. 554 = Saussurea serrata.
pauciflora, Poir. l. c. 563 = Cnicus heterophyllus.
pectinata, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 94 = centauroides.
pedunculata, Pers. Syn. ii. 391 = Vernonia pedunculata.
phrygia, *Karamysch. in Linn. Amoen. Acad.* vii. 456.—Ross.
Picris, Bieb. Fl. Taur. Cauc. iii. 546 = Centaurea Picris.
pilosa, Ait. Hort. Kew. ed. I. iii. 138 = Liatris spicata.
pilosa, Bosc, ex DC. Prod. v. 63, sphalm. = Vernonia angustifolia.
pinnata, Kit. in Schult. Oestr. Fl. ed. II. ii. 454 = tinctoria.
pinnata, Poir. Encyc. Suppl. v. 43 = Jurinea pinnata.
pinnatifida, Desf. Tabl. ed. I. 93 = radiata.
pinnatifida, *Poir. Encyc.* vi. 561.—Hispan.
Pollichii, DC. Cat. Hort. Monsp. (1813) 145 = Jurinea cyanoides.
polyclonos, DC. Diss. Comp. 30 = Jurinea polyclonos.
polygyna, A. Rich. Tent. Fl. Abyss. i. 457 = Laggera purpurascens.
praealta, Linn. Sp. Pl. 818 = Vernonia noveboracensis.
procumbens, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 165.—As. centr.
propinqua, Pomel, Nouv. Mat. Fl. Atl. 25 = flavescens.
pulchella, Fisch. & Mey. ex DC. Prod. vi. 676 = Jurinea pulchella.
pulchella, Sims, Bot. Mag. t. 2589 = Saussurea japonica.
pumila, Thore, ex DC. Prod. vi. 667 = tinctoria.
pungens, Franch. & Sav. Enum. Pl. Jap. ii. 416 = atriplicifolia.
pungens, Poir. Encyc. vi. 566 = Cnicus pungens.
purpurata, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 20 = Saussurea serrata.
pygmaea, Jacq. Fl. Austr. v. 20 = Saussurea pygmaea.
quinquefolia, *Bieb. Fl. Taur. Cauc.* ii. 264.—Reg. Caucas.
radiata, *Bieb. l. c.* iii. 545.—Europ.; Reg. Caucas.
rhapontica, DC. Fl. Fr. iv. 87 = Centaurea Rhapponticum.
rosmarinifolia, Nym. Consp. 401 = Staehelina dubia.
runcinata, Demid. ex DC. Prod. vi. 537 = Saussurea runcinata.
salicifolia, Linn. Sp. Pl. 817 = Saussurea salicifolia.
salicina, Pall. Reise, iii. 682 = Jurinea linearifolia.
salina, Pall. Reise, iii. 314 = Saussurea crassifolia.
salisburgensis, Poir. Encyc. vi. 564 = Cnicus rivularis.
salsa, Pall. Reise, ii. 444; Bieb. Fl. Taur. Cauc. ii. 266 = Saussurea crassifolia.
salsa, Spreng. Fl. Hal. Mant. 49 = praec.?
scariosa, Linn. Sp. Pl. 818 = Liatris scariosa.
scordioidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
Scordium, *Lour. Fl. Cochinch.* 483.—China.
semiserrata, Willd. ex DC. Prod. v. 33 = Vernonia serratuloides.
serrata, Sweet, Hort. Brit. ed. I. 213 = Saussurea serrata.
setosa, Willd. Sp. Pl. iii. 1645 = Cnicus arvensis.
simplex, DC. Diss. Comp. ii. 30 = Jurinea mollis.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 367.—Turkest.
sonchifolia, *Bess. ex Rochel, Pl. Banat. Rar.* (*Index*) (Quid?).—Banat.

SERRATULA:—
speciosa, Ait. Hort. Kew. ed. I. iii. 138 = Liatris elegans.
spicata, Linn. Sp. Pl. 819 = Liatris spicata.
spinosa, *Forsk. Fl. Aegypt. Arab.* 217.—Reg. Byzant.
spinosa, Gilib. Fl. Lituan. i. 184 = Cnicus arvensis.
spinosa, Poir. ex DC. Prod. v. 475 = Iphiona juniperifolia.
spinulosa, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 329.—Turkestan.
squalida, Steph. ex DC. Prod. vi. 537 = Saussurea alata.
squarrosa, Linn. Sp. Pl. 818 = Liatris squarrosa.
Stelleria, Steph. ex DC. Prod. vi. 534 = Saussurea elongata.
stoechadifolia, Bieb. Fl. Taur. Cauc. ii. 266 = Jurinea staechadifolia.
subacaulis, Poir. Encyc. vi. 550 = Jurinea humilis.
subulata, Poir. l. c. 558 = Iphiona juniperifolia.
suffruticosa, Trautv. in Act. Hort. Petrop. viii. (1883) 510, sphalm. = seq.
suffruticulosa, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 110.—Soongaria.
tenuifolia, *Bong. in Bull. Sc. Acad. Pétersb.* viii. (1841) 340.—Sibir. altaic.
thracica, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 178 = Centaurea monacantha.
Tilesii, Ledeb. in Mém. Acad. Pétersb. v. (1812) 562 = Saussurea alpina.
tincta, Cass. in Dict. Sc. Nat. xlvii. 476 = Jurinea linearifolia.
tinctoria, *Linn. Sp. Pl.* 816.—Europ.
tinctoria, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 183 = Saussurea affinis.
tomentosa, Hoffm. ex Ledeb. Fl. Ross. ii. 762 = Jurinea staechadifolia.
transylvanica, Spreng. Syst. iii. 388 = Jurinea mollis.
Trautvetteriana, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 321.—Turkestan.
tuberosa, Neck. in Act. Acad. Theod. Palat. ii. (1770) 465 = Cnicus tuberosus.
uniflora, Poir. Encyc. vi. 560; Spreng. Syst. iii. 388 = Centaurea monantha.
uniflosculosa, Poir. Encyc. Suppl. v. 43 = Staehelina uniflosculosa.
varia, Hill, ex Steud. Nom. ed. II. ii. 570 (Quid?).—Am. bor.
viciaefolia, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 586.—Persia.
Vulpii, Fisch.-Ooster, in Flora, xxxvii. (1854) 98 = macrocephala.
Wolffii, Andrae, in Bot. Zeit. xiii. (1855) 321 = coronata.
xeranthemoides, *Bieb. Fl. Taur. Cauc.* ii. 265.—Rossia; Reg. Caucas.

SERRESIA, Montr. in Mém. Acad. Lyon, x. (1860) 179. *INCERT. SEDIS.*—Cf. Benth. & Hook. f. Gen. i. 970.
asarifolia, *Montr. l. c.* 180.—N. Caled.

SERRONIA, Gaudich. in Deless. Ic. Sel. iii. 54. t. 90 (1837) = **Piper**, Linn.
acuminata, Moric. Pl. Nouv. Am. 86. t. 57 = P. corcovadense.
Jaborandi, Gaudich. & Guill. in Deless. Ic. Sel. iii. 54 = P. Jaborandi.

SERRULATA, DC. Prod. v. 63 (1836), sphalm. = **Serratula**, Dill. (Compos.).

SERRURIA, Salisb. Parad. Lond. sub t. 67 (1807). *PROTEACEAE*, Benth. & Hook. f. iii. 170.
HOLDERLINIA, Neck. Elem. i. 106 (1790).
SERRARIA, Burm. Afr. 264. t. 99 (1737).
abrotanifolia, *Knight, Prot.* 79.—Afr. austr.
acrocarpa, *R. Br. in Trans. Linn. Soc.* x. (1810) 113.—Afr. austr.
adscendens, *R. Br. l. c.* 127.—Afr. austr.
adscendens, Drège, ex Meissn. in DC. Prod. xiv. 287 = saxicola.
aemula, *Knight, Prot.* 89.—Afr. austr.
Aitoni, *R. Br. in Trans. Linn. Soc.* x. (1810) 114.—Afr. austr.
albicans, *Roem. & Schult. Syst.* iii. 378.—Afr. austr.
Andrewsii, Endl. Gen. Suppl. iv. II. 79 = phylicoides, odorata.
anemonefolia, *Knight, Prot.* 83.—Afr. austr.
anethifolia, *Knight, l. c.* 84.—Afr. austr.

SERRURIA:—
arenaria, *R. Br. in Trans. Linn. Soc.* x. (1810) 117.—Afr. austr.
arenaria, Knight, Prot. 87 = emarginata.
argentiflora, Buek, ex Meissn. in DC. Prod. xiv. 297 = triternata.
artemisiaefolia, *Knight, Prot.* 80.—Afr. austr.
barbigera, *Knight, l. c.* 90.—Afr. austr.
Bergii, *R. Br. in Trans. Linn. Soc.* x. (1810) 220.—Afr. austr.
Brownii, *Meissn. in DC. Prod.* xiv. 290.—Afr. austr.
Burmanni, R. Br. in Trans. Linn. Soc. x. (1810) 310 = fasciflora
callosa, *Knight, Prot.* 80.—Afr. austr.
candicans, *R. Br. in Trans. Linn. Soc.* x. (1810) 130.—Afr. austr.
candicans, Roem. & Schult. Syst. iii. 378 = Nivenia concava.
chlamydiflora, *Knight, Prot.* 91.—Afr. austr.
ciliata, *R. Br. in Trans. Linn. Soc.* x. (1810) 123.—Afr. austr.
collina, *Knight, Prot.* 86.—Afr. austr.
colorata, *Buek, ex Drège, Zwei Pfl. Docum.* 109, 221.—Afr. austr.
colorata, Zeyh. ex Meissn. in DC. Prod. xiv. 286 = cygnea.
commutata, Endl. Gen. Suppl. iv. II. 79 = triplicato-ternata.
compar, *R. Br. in Trans. Linn. Soc.* x. (1810) 129.—Afr. austr.
concinna, *Knight, Prot.* 88.—Afr. austr.
congesta, R. Br. in Trans. Linn. Soc. x. (1810) 123 = millefolia.
crithmifolia, *Knight, Prot.* 83; *R. Br. l. c.* 132.—Afr. austr.
cyanea, Drège, ex Meissn. in DC. Prod. xiv. 286 = cygnea.
cyanoides, R. Br. in Trans. Linn. Soc. x. (1810) 117 = pulchella.
cygnea, *R. Br. l. c.* 113.—Afr. austr.
decipiens, *R. Br. l. c.* 129.—Afr. austr.
decumbens, *R. Br. l. c.* 126.—Afr. austr.
delphiniifolia, *Knight, Prot.* 82.—Afr. austr.
diffusa, *R. Br. in Trans. Linn. Soc.* x. (1810) 115.—Afr. austr.
Dregei, *Meissn. in DC. Prod.* xiv. 289.—Afr. austr.
elevata, *R. Br. in Trans. Linn. Soc.* x. (1810) 114.—Afr. austr.
elongata, R. Br. l. c. 132 = crithmifolia.
elumbis, *Knight, Prot.* 83.—Afr. austr.
emarginata, *Sweet, Hort. Brit.* ed. I. 348.—Afr. austr.
erecta, *Meissn. in DC. Prod.* xiv. 298.—Afr. austr.
eriocephala, Steud. Nom. ed. II. ii. 571 = squarrosa.
fallax, *Knight, Prot.* 90.—Afr. austr.
fasciflora, *Knight, l. c.* 85.—Afr. austr.
flagellaris, *R. Br. in Trans. Linn. Soc.* x. (1810) 127.—Afr. austr.
flagellifolia, *Knight, Prot.* 84.—Afr. austr.
flava, *E. Mey. ex Meissn. in DC. Prod.* xiv. 287.—Afr. austr.
florida, *R. Br. in Trans. Linn. Soc.* x. (1810) 126.—Afr. austr.
florida, *Knight, Prot.* 92.—Afr. austr.
foeniculacea, *R. Br. in Trans. Linn. Soc.* x. (1810) 122.—Afr. austr.
foeniculacea, Sieber, ex Meissn. in DC. Prod. xiv. 294 = glomerata.
foliosa, *Knight, Prot.* 89.—Afr. austr.
frondosa, *Knight, l. c.* 85.—Afr. austr.
fucifolia, *Knight, l. c.* 81.—Afr. austr.
furcellata, *R. Br. in Trans. Linn. Soc.* x. (1810) 118.—Afr. austr.
glaberrima, *R. Br. l. c.* 112.—Afr. austr.
glomerata, *R. Br. l. c.* 128.—Afr. austr.
gracilis, *Knight, Prot.* 81.—Afr. austr.
gremiiflora, *Knight, l. c.* 88.—Afr. austr.
helvola, Steud. Nom. ed. II. ii. 571 = cygnea.
heterophylla, *Meissn. in DC. Prod.* xiv. 284.—Afr. austr.
hirsuta, Berg. ex Meissn. l. c. 290 = Brownii.
hirsuta, *R. Br. in Trans. Linn. Soc.* x. (1810) 120.—Afr. austr.
hirsuta, Buek, ex Meissn. in DC. Prod. xiv. 289 = scoparia.
hyemalis, *Knight, Prot.* 84.—Afr. austr.
incrassata, *Buek, ex E. Mey. in Drège, Zwei Pfl. Docum.* 98.—Afr. austr.
Kraussii, *Meissn. in DC. Prod.* xiv. 296.—Afr. austr.

SERRURIA:—
linearis, *Knight, Prot.* 82.—Afr. austr.
millefolia, *Knight, l. c.* 79.—Afr. austr.
montana, *Knight, l. c.* 80.—Afr. austr.
multifida, E. Mey. in Drège, Zwei Pfl. Docum. 74 = Aitoni.
nervosa, *Meissn. in DC. Prod.* xiv. 290.—Afr. austr.
nitida, *R. Br. in Trans. Linn. Soc.* x. (1810) 124.—Afr. austr.
Niveni, *R. Br. l. c.* 121.—Afr. austr.
odorata, *Sweet, Hort. Brit.* ed. I. 348.—Afr. austr.
parilis, *Knight, Prot.* 91.—Afr. austr.
pedunculares, *Knight, l. c.* 81.—Afr. austr.
pedunculata, R. Br. in Trans. Linn. Soc. x. (1810) = artemisiaefolia.
phylicoides, R. Br. l. c. 126 = barbigera.
pilosa, *Roem. & Schult. Syst.* iii. 378.—Afr. austr.
pinnata, *R. Br. in Trans. Linn. Soc.* x. (1810) 116.—Afr. austr.
plumosa, *Meissn. in DC. Prod.* xiv. 291.—Afr. austr.
pulchella, *Knight, Prot.* 89.—Afr. austr.
quinquemestris, *Knight, l. c.* 87.—Afr. austr.
rangiferina, *Knight, l. c.* 86.—Afr. austr.
rostellaris, *Knight, l. c.* 88.—Afr. austr.
Roxburghii, *R. Br. in Trans. Linn. Soc.* x. (1810) 130.—Afr. austr.
rubricaulis, *R. Br. l. c.* 128.—Afr. austr.
saxicola, *Buek, ex E. Mey. in Drège, Zwei Pfl. Docum.* 76.—Afr. austr.
scariosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 118.—Afr. austr.
scariosa, Drège, ex Meissn. in DC. Prod. xiv. 285 = furcellata.
scoparia, *R. Br. in Trans. Linn. Soc.* x. (1810) 119.—Afr. austr.
simplicifolia, *R. Br. l. c.* 115.—Afr. austr.
sphaerocephala, Steud. Nom. ed. II. ii. 571 = hirsuta, pedunculata, scariosa.
squarrosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 124.—Afr. austr.
Stilbe, *R. Br. l. c.* 120.—Afr. austr.
subcorymbosa, *Meissn. in DC. Prod.* xiv. 285.—Afr. austr.
Thunbergi, Endl. Gen. Suppl. iv. II. 79 = pilosa.
tomentosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 76.—Afr. austr.
tridentata, D. Dietr. Syn. Pl. i. 520 = triternata.
trilopha, *Knight, Prot.* 90.—Afr. austr.
triplicato-ternata, *Roem. & Schult. Syst.* iii. 378.—Afr. austr.
triternata, R. Br. in Trans. Linn. Soc. x. (1810) 131 = antheifolia.
vallaris, *Knight, Prot.* 78.—Afr. austr.
vestita, *Buek, ex E. Mey. in Drège, Zwei Pfl. Docum.* 104.—Afr. austr.
villosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 122.—Afr. austr.
zanthophylla, *Knight, Prot.* 86.—Afr. austr.
Zeyheri, *Meissn. in DC. Prod.* xiv. 297.—Afr. austr.

SERSALISIA, R. Br. Prod. 529 (1810) = **Lucuma**, Molina (Sapot.).
cotinifolia, F. Muell. Fragm. v. 161 = Hormogyne cotinifolia.
galactoxylon, F. Muell. ex Benth. Fl. Austral. iv. 279 = L. galactoxylon.
glabra, A. Gray, in Proc. Am. Acad. v. (1862) 327 = Sideroxylon Richardi.
laurifolia, A. Rich. Sert. Astrolab. 84. t. 31 = Sideroxylon Richardi.
obovata, R. Br. Prod. 529 = Sideroxylon Brownii.
sericea, R. Br. l. c. 530 = L. sericeum.

SERTIFERA, Lindl. ex Reichb. f. in Linnaea, xli. (1877) 63. *ORCHIDEAE*, Benth. & Hook. f. iii. 591.
purpurea, *Reichb. f. l. c.* 64.—Ecuador.
virgata, *Reichb. f. l. c.*—Peruv.

SERTULA, Linn. Syst. ed. I (1735) = **Trifolium**, Linn. (Legumin.).

SERTUERNERA, Mart. Nov. Gen. et Sp. ii. 36. t. 136 (1826) = **Pfaffia**, Mart. (Amarant.).
glauca, Mart. l. c. 37. t. 136 = P. glauca.
guianensis, Klotzsch, in Rich. Schomb. Faun. Fl. Brit. Gui. 933, nomen.—Guian.

SERTUERNERA :—
iresinoides, Mart. Nov. Gen. et Sp. ii. 48. t. 138 = Gomphrena iresinoides.
luzulaeflora, Mart. l. c. 39 = Gomphrena luzulaeflora.
paraguayensis, Parodi, in Anal. Soc. Cient. Argent. vi. (1878) 40 ; Contrib. 92.—Paraguay.
Schomburgkii, Klotzsch, in Rich. Schomb. Faun. Fl. Brit. Gui. 933, nomen.—Guian.

SESAMELLA, Reichb. Consp. 186. (1828) = **Astrocarpus**, Neck. (Resed.).
spathulata, Reichb. in Flora, xiii. (1830) 130 = A. Clusii.
Tournefortii, Reichb. l. c. = A. Sesamoides.

SESAMOIDES, Tourn. ex Reichb. Consp. 186 (1828) = **Astrocarpus**, Neck. (Resed.).

SESAMOPTERIS, DC. ex Meissn. Gen. 298 ; Comm. 206 (1840) = **Sesamum**, Linn. (Pedal.).
alata, DC. ex Meissn. Gen. Comm. 206 = Sesamum capense.
pentaphylla, DC. Prod. ix. 251 = Sesamum capense.
radiata, DC. ex Meissn. Gen. Comm. 206 = Sesamum radiatum.

SESAMOTHAMNUS, Welw. in Trans. Linn. Soc. xxvii. (1869) 49. t. 18. *PEDALINEAE*, Benth. & Hook. f. ii. 1058.
benguellensis, *Welw. l. c.* 50.—Afr. trop.

SESAMUM, Adans. Fam. ii. 213 (1763) = **Martynia**, Linn. (Pedal.).

SESAMUM, Linn. Coroll. Gen. 11 (1737). *PEDALINEAE*, Benth. & Hook. f. ii. 1058.
ANTHADENIA, Van Houtte, Fl. des Serres, ii. (Apr. 1846) 10. t. 6.
DYSOSMON, Rafin. Fl. Ludov. 48 (1817).
GANGILA, Bernh. in Linnaea, xvi. (1842) 42.
SESAMOPTERIS, DC. ex Meissn. Gen. 298 ; Comm. 206 (1840).
SIMSIMUM, Bernh. in Linnaea, xvi. (1842) 42.
africanum, Todaro, Ind. Sem. Hort. Panorm. (1866) 34 = indicum ?
alatum, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 284 = capense.
angolense, *Welw. Apont.* 588 ; *et in Trans Linn. Soc.* xxvii. (1871) 51.—Afr. trop.
auriculatum, *Presl, in Oken, Isis*, xxi. (1828) 275.—Ins. Creta.
brasiliense, *Vell. Fl. Flum.* 264 ; vi. t. 90.—Bras.
calycinum, *Welw. in Trans. Linn. Soc.* xxvii. (1871) 52.—Afr. trop.
capense, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
edule, Hort. ex Steud. Nom. ed. I. 769 = indicum.
gracile, Endl. in Linnaea, vii. (1832) 10. t. 3 = capense.
indicum, *Linn. Sp. Pl.* 634.—Reg. trop. (cult.).
javanicum, Burm. f. Fl. Ind. 133 = Artanema sesamoides.
laciniatum, *Klein, ex Willd. Sp. Pl.* iii. 359.—Ind. or.
laciniatum, Willd. l. c., partim = trifoliatum ?
luteum, Retz, Obs. vi. 31 = indicum.
macranthum, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 131.—Afr. trop.
occidentalis, Heer & Regel, Ind. Sem. Hort. Turic. (1842) ; ex DC. Prod. ix. 250 = indicum.
oleiferum, Moench, Meth. Suppl. 174 = indicum.
orientale, Linn. Sp. Pl. 634 = indicum.
orientale, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 275 = auriculatum.
pentaphyllum, E. Mey. in Drège, Zwei Pfl. Docum. 50, 54 ; et ex DC. Prod. ix. 251 = capense.
prostratum, *Retz. Obs.* iv. 28.—Ind. or.
pterospermum, R. Br. in Salt, Abyss. App. 64 = capense.
radiatum, *Schum. & Thonn. Beskr. Guin. Pl.* 282.—Afr. trop.
rigidum, *Peyr. in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 572.—Afr. trop.
rostratum, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = capense.
trifoliatum, *Mill. Gard. Dict.* ed. VIII. n. 3.—Ind. or.

SESBAN, Adans. Fam. ii. 327 (1763) = seq.

SESBANA, R. Br. in Ait. Hort. Kew. ed. II. iv. 330 (1812) = seq.

SESBANIA, Scop. Introd. 308 (1777). *LEGUMINOSAE*, Benth. & Hook. f. i. 502.
AGATI, Adans. Fam. ii. 376 (1763) ; Desv. Journ. Bot. i. (1813) 119. t. 4.
DARWINIA, Rafin. Fl. Ludov. 106 (1817).
DAUBENTONIA, DC. Mém. Lég. 285 (1825).
GLOTTIDIUM, Desv. Journ. Bot. i. (1813) 119. t. 1.
MONOPLECTRA, Rafin. Fl. Ludov. 106 (1817).
RESUPINARIA, Rafin. Sylva Tellur. 115 (1838).
aculeata, *Poir. Encyc.* vii. 128 ; Pers. Syn. ii. 316.—Cosmop. trop.
aegyptiaca, Buch.-Ham. ex Wall. Cat. n. 5655 B = aculeata.
aegyptiaca, *Poir. Encyc.* vii. 128 ; *Pers. Syn.* ii. 316.—Geront. trop.
affinis, Schrad. Ind. Sem. Hort. Gotting. (1819) ; ex DC. Prod. ii. 265 = aculeata.
arabica, *Hochst. ex Steud. Nom.* ed. II. ii. 572.—Arab.
australis, F. Muell. in Trans. Vict. Inst. i. (1855) 36 = aculeata.
bispinosa, Steud. Nom. ed. II. ii. 572 = aculeata.
brachycarpa, *F. Muell. Fragm.* xi. 32.—Austral.
Candira, Buch.-Ham. ex Wall. Cat. n. 5655 C = aculeata.
cannabina, Poir. Encyc. vii. 130 ; Pers. Syn. ii. 316 = aculeata.
cassioides, *G. Don, Gen. Syst.* ii. 240.—Ecuador.
Cavanillesii, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 342.—Am. bor. ; Mexic.
cinerascens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 134.—Afr. trop.
coccinea, Poir. Encyc. vii. 127 = grandiflora.
cochinchinensis, *DC. Prod.* ii. 266.—Cochinch.
disperma, Pursh, Fl. Am. Sept. ii. 485 = platycarpa.
dubia, H. B. & K. Nov. Gen. et Sp. vii. 268. t. 660 = Coursetia dubia.
dubia, Steud. Nom. ed. II. ii. 572 = exasperata.
exasperata, *H. B. & K. Nov. Gen. et Sp.* vi. 534.—Am. trop.
ferruginea, Hochst. in Flora, xxiv. (1841) i. Intell. 31 = Millettia ferruginea.
filiformis, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 133 = aegyptiaca.
filiformis, Hochst. ex Baker, l. c. 135 = leptocarpa.
fusca, Desv. in Mém. Soc. Linn. Par. iv. (1826) 300 = Aeschynomene fusca.
geminiflora, Schrad. ex Steud. Nom. ed. II. ii. 572 (Quid ?).—Hab. ?
gracilis, *Schrad. Cat. Sem. Gotting.* 1819 ; *ex DC. Prod.* ii. 265.—Hab. ?
grandiflora, *Poir. Encyc.* vii. 127 ; *Pers. Syn.* ii. 316.—Ind. or. ; Malaya ; Austral.
guianensis, *Klotzsch, in Rich. Schomb. Faun. Fl. Brit. Gui.* 1199, nomen.—Guian.
javanica, *Miq. Fl. Ind. Bat.* i. i. 288.—Java.
leptocarpa, *DC. Prod.* ii. 265.—Afr. trop.
longifolia, *DC. l. c.*—Mexic.
macrocarpa, *Muhl. ex Rafin. Fl. Ludov.* 137 ; *et Ell. Sketch*, ii. 221.—Am. bor.
macroptera, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) vii. n. 14.—Paraguay.
marginata, *Benth. in Mart. Fl. Bras.* xv. i. 43.—Am. trop.
mossambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. 45 = aculeata.
multijuga, Schweinf. ex Baker, in Oliver, Fl. Trop. Afr. ii. 133 = aegyptiaca.
muricata, Macfad. Fl. Jamaic. i. 257 = aculeata.
occidentalis, Griseb. in Goett. Abh. vii. (1857) 200 = aculeata.
occidentalis, *Poir. Encyc.* vii. 129 ; *Pers. Syn.* ii. 316.—Ind. occ.
pachycarpa, DC. Prod. ii. 265 = aegyptiaca.
paludosa, Jacq. Hort. Vindob. ex DC. Prod. ii. 265 = aculeata.
peruviana, *G. Don, Gen. Syst.* ii. 240.—Peruv.
picta, Lindl. ex Griseb. Fl. Brit. W. Ind. 184 = occidentalis.
picta, Poir. Encyc. vii. 129 ; Pers. Syn. ii. 316 = aegyptiaca.
platycarpa, *Pers. l. c.*—Am. bor.
polyphylla, *Miq. Fl. Ind. Bat.* i. i. 283.—Java.
procumbens, *Wight & Arn. Prod.* 215.—Ind. or.
pubescens, *DC. Prod.* ii. 265.—Guian.
punctata, DC. l. c. = aegyptiaca.
punicea, *Benth. in Mart. Fl. Bras.* xv. i. 43.—Reg. Argent.
sericea, *DC. Prod.* ii. 266.—Ind. occ.

SESBANIA :—
sericea, Link, Enum. Hort. Berol. ii. 244 ; et ex Desv. in Mém. Soc. Linn. Par. iv. (1826) 300 = praec. ?
sericea, Welw. Apont. 585 = pubescens.
simpliciuscula, *F. Muell. ex Benth. Fl. Austral.* ii. 213.—Austral.
sphaerocarpa, *Welw. Apont.* 590 ; *et ex Baker, in Oliver, Fl. Trop. Afr.* ii. 135.—Afr. trop.
stricta, *Schrad. Ind. Sem. Hort. Gotting.* (1834).—*Cf. Linnaea*, x. (1836) *Litt.* 111.—Ind. or.
tetraptera, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 136.—Afr. trop.
Thurberi, *S. Wats. Bibliog. Ind. N. Am. Bot.* i. 258.—Am. bor. occ.
tomentosa, *Hook. & Arn. Bot. Beech. Voy.* 286.—Mexic.
triflora, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 346, nomen.—Ins. Timor.
uliginosa, Sweet, Hort. Brit. ed. I. 129 = aculeata.
vesicaria, Ell. Sketch, ii. 222 = platycarpa.
virgata, Poir. Encyc. vii. 129 ; Pers. Syn. ii. 316 = Coursetia virgata.

SESELI, Linn. Gen. ed. I. 80 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 901.
BUBON, Linn. Gen. ed. I. 339 (1737).
DELA, Adans. Fam. ii. 499 (1763).
DETHAWIA, Endl. Gen. 775 (1839).
GALBANOPHORA, Neck. Elem. i. 175 (1790).
HIPPOMARATHRUM, Hall. in Rupp. Fl. Jen. ed. II. 280 (1745).
LIBANOTIS, Riv. ex Hall. Enum. Helv. ii. 450 (1742).
LOMATOPODIUM, Fisch. et Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 305.
MARATHRUM, Link, Handb. i. 348 (1829).
SCAPHESPERMUM, Edgew. in Trans. Linn. Soc. xx. (1846) 58.
STENOCOELIUM, Ledeb. Fl. Alt. i. 297 (1829).
TELEOPHUS, Dulac, Fl. Hautes-Pyr. 353 (1867).
WALLROTHIA, Spreng. Pugill. ii. 52 (1815).
Aegopodium, Scop. Fl. Carn. ed. II. i. 215 = Aegopodium Podagraria.
algens, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 104.—Austral.
alpinum, Bieb. Fl. Taur. Cauc. i. 236 = Carum Lomatocarum.
Ammi, Savi, Due Cent. Fl. Etrusc. 71 = Apium Ammi.
Ammi, Steud. Nom. ed. II. ii. 572 = Carum ammoides.
ammoides, Hort. ex Link, Enum. Hort. Berol. i. 285 = Apium Ammi.
ammoides, Jacq. Hort. Vindob. t. 52 = Carum copticum.
ammoides, Linn. Sp. Pl. 260 = Carum ammoides.
Amomum, Scop. Fl. Carn. ed. II. i. 213 = Sison Amomum.
annuum, *Linn. Sp. Pl.* 260.—Europ.
annuum, Pall. Tabl. Taur. 49 ; Bieb. Fl. Taur. Cauc. i. 235 = tortuosum.
arenarium, Bieb. l. c. iii. 242 = campestre.
aristatum, Ait. Hort. Kew. ed. I. i. 359 = Ligusticum pyrenaicum.
articulatum, Crantz, Stirp. Austr. ed. I. fasc. iii. 91 ; ed. II. 205 = S. Hippomarathrum.
aspergillifolium, *Bogusl. in Erman, Archiv*, vi. (1848) 63.—Rossia.
asperum, *Sond. in Harv. & Sond. Fl. Cap.* ii. 550.—Afr. austr.
athamantoides, *C. A. Mey. ex Ledeb. Fl. Alt.* i. 342.—Sibir. altaic.
atlanticum, Boiss. Diagn. Ser. II. ii. 87 = glaucum.
bengalense, Roxb. Hort. Beng. 22 ; Fl. Ind. ii. 94 = Oenanthe benghalensis.
bienne, Crantz, Stirp. Austr. ed. I. fasc. iii. 88 ; ed. II. 204 = annuum.
biternatum, Torr. & Gray, Fl. N. Am. i. 627, in syn., sphalm. = triternatum.
Bocconi, *Guss. Adnot. Cat. Pl. Boccad.* (1821) 12.—Sicil.
brevicaule, Jord. in F. Schultz, Arch. Fl. 1 = annuum.
buchtormense, *Koch, in Nov. Act. Nat. Cur.* xii. i. (1824) 111.—Sibir.
Bunius, Vill. Prosp. 24 ; Hist. Pl. Dauph. ii. 586 = Carum Bunius.
caespitosum, *Sibth. & Sm. Fl. Graec. Prod.* i. 200.—Bithynia.
caffrum, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 533 = Annesorhiza caffra (in Addend.).
campestre, *Bess. Enum. Pl. Volh.* 44.—Europ. or.

SESELI :—

cantabricum, *Lange, Ind. Sem. Hort. Haun.* (1855) 27. —*Cf. Linnaea*, xxviii. (1856) 363.—Hispan.
carinatum, *Kit. in Linnaea*, xxxii. (1863) 461.—Europ. austr.
Carum, Scop. Fl. Carn. ed. II. i. 215 = Carum Carvi.
Carvi, Spreng. in Schult. Syst. vi. 414 = Carum Carvi.
Carvifolia, Linn. Sp. Pl. 260 = Selinum Carvifolia.
carvifolium, Vill. Prosp. 24 ; Hist. Pl. Dauph. ii. 586 = praec.
cervariaefolium, DC. Cat. Hort. Monsp. (1813) 145 = Athamanta cervariaefolia.
cespitosum, Spreng. in Schult. Syst. vi. 843 = caespitosum.
chaerophylloides, Hort. ex DC. Prod. iv. 146 = varium.
chaerophylloides, *Thunb. Prod. Pl. Cap.* 51.—Afr. austr.
coloratum, Ehrh. Beitr. v. 179 = annuum.
condensatum, *Reichb. f. Ic. Fl. Germ.* xxi. 37.—Sibir.
connatum, Walp. Rep. ii. 402, sphalm. = coronatum.
coronatum, *Ledeb. Fl. Alt.* i. 336.—Sibir. altaic.
corsicum, Bess. Cat. Hort. Crem. (1816) 130 ; Link, Enum. Hort. Berol. i. 283 = Carum ammoides.
corymbosum, *Boiss. & Heldr. Diagn.* Ser. I. x. 29.—As. Min.
crassifolium, Hort. ex DC. Prod. iv. 147 = Pallasii.
creticum, *Presl, Bot. Bemerk.* 74 (Quid ?).—Ins. Creta.
crithmifolium, *Boiss. Fl. Orient.* ii. 962.—Archip. Graec.
cuneifolium, *Bieb. Beschr. Casp.* 159 ; *Fl. Taur. Cauc.* i. 236.—Reg. Caucas.
curvifolium, J. F. Gmel. Syst. 489, sphalm. = Selinum Carvifolia.
daucifolium, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 693 = Selinum Monnieri.
decipiens, Ledeb. ex Reichb. Fl. Germ. Excurs. 467 = varium.
defoliatum, *Ledeb. Fl. Alt.* i. 343.—Sibir. altaic.
denudatum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 298.—Persia.
dichotomum, *Pall. Tabl. Taur.* 49 ; *Bieb. Fl. Taur. Cauc.* i. 255.—Tauria.
dioicum, Dufour, ex DC. Prod. iv. 104 = Trinia Dufourii.
dioicum, Vill. Hist. Pl. Dauph. ii. 579 = Trinia vulgaris.
divaricatum, Pursh, Fl. Am. Sept. ii. 732 = Musenium divaricatum.
dubium, Schk. Handb. i. 217 = Selinum sylvestre.
elatum, *Linn. Sp. Pl.* ed. II. 375 ; *Gouan, Illustr.* 16. t. 8.—Europ. austr.
elatum, Reichb. Fl. Germ. Excurs. 467 = osseum.
elatum, Thuill. Fl. Par. 118 = varium.
Falcaria, Crantz, Stirp. Austr. ed. II. 208 = Falcaria vulgaris.
Fedtschenkoanum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 30.—Turkestan.
ferulaceum, Stev. ex DC. Prod. iv. 147 = annuum.
ferulaefolium, Poir. Encyc. vii. 135 = Ligusticum pyrenaicum.
filifolium, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 203.—Balkan.
filifolium, Thunb. Prod. Pl. Cap. 51 = Oenanthe filiformis.
filiforme, Gilib. Fl. Lituan. ii. 36 = Carum Bunius.
foeniculifolium, Poir. Encyc. vii. 137 = Carum copticum.
fragile, *Gouan, Illustr.* 15.—Europ.
Gilliesii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 354 = Carum copticum ?
glabratum, Schlecht. ex Ledeb. Fl. Ross. ii. 274 = tenuifolium.
glabratum, Willd. ex Schult. Syst. vi. 406 (= *Rumia seseloides*).—Sibir. altaic.
glaucescens, Jord. ex Nym. Consp. 297 = montanum.
glaucum, *Bieb. Fl. Taur. Cauc.* i. 234.—Europ. austr. ; Reg. Cauc.
glaucum, Jacq. Fl. Austr. ii. 27 = osseum.
glaucum, Lam. Fl. Fr. iii. 436 = Trinia vulgaris.
glaucum, Linn. Sp. Pl. 260 = montanum.
glaucum, Pett. ex Nym. Consp. 297 = Tommasinii.
globiferum, *Vis. in Flora*, xiii. (1830) 50.—Dalmatia.
Gouani, Koch, Syn. Fl. Germ. ed. I. 294 = Pimpinella Gouani.
gracile, *Waldst. & Kit. Pl. Rar. Hung.* ii. 122. t. 117. —Transsylv.
granatense, *Willk. in Bot. Zeit.* v. (1847) 431.—Hispan.
graveolens, *Ledeb. Fl. Alt.* i. 340.—Rossia.

SESELI :—

graveolens, Scop. Fl. Carn. ed. II. i. 215 = Apium graveolens.
gummiferum, Boiss. Diagn. Ser. I. x. 29 = corymbosum.
gummiferum, Less. ex Ledeb. Fl. Ross. ii. 273 = Lessingianum.
gummiferum, *Pall. ex Sm. Exot. Bot.* ii. 121.—Tauria.
Hallii, A. Gray, in Proc. Am. Acad. viii. (1870) 288 = Carum Hallii.
Harveyanum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 104.—Austral.
heterophyllum, *Janka, in Linnaea*, xxx. (1859-60) 572.—Europ. or.
Hippomarathrum, *Jacq. Enum. Stirp. Vindob.* 52, 224.—Europ.
Hippomarathrum, Pall. Tabl. Taur. 49 ; et ex Bieb. Fl. Taur. Cauc. i. 235 = petraeum.
hispidum, Bess. Cat. Hort. Crem. (1812) ; ex Schult. Syst. vi. 407 (Quid ?).—Hab. ?
hypoleucum, *Meissn. ex Krauss, in Flora*, xxvii. (1844) 431.—Afr. austr.
indicum, *Wight & Arn. Prod.* 371.—Ind. or.
intricatum, *Boiss. Elench.* 48.—Hispan.
inundatum, Crantz, Cl. Umb. Emend. 91 = Apium inundatum.
junceum, Sibth. & Sm. Fl. Graec. Prod. i. 200 = Sclerochorton junceum.
Ledebourii, G. Don, Gen. Syst. iii. 308 = S. Hippomarathrum.
Lehmannianum, *Boiss. Fl. Orient.* ii. 967.—Turkestan.
leiocarpum, Hook. Fl. Bor. Am. i. 263. t. 93 = Peucedanum leiocarpum.
Lessingianum, *Turcz. ex Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 429.—Sibir. altaic.
leucospermum, *Waldst. & Kit. Pl. Rar. Hung.* i. 92. t. 89.—Hungar.
Libanotis, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 111.—Europ. ; Oriens.
libanotum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
littorale, Willk. in Flora, xxxiv. (1851) 724 = tortuosum.
Lobelianum, Hornem. ex Reichb. Fl. Germ. Excurs. 467 = elatum.
Lobelianum, *Hort. Dorpat.* (1819) *ex Schult. Syst.* vi. 407 ; *Bess. ex Sweet, Hort. Brit.* ed. II. 248 (Quid ?). —Europ. austr.
lucanum, *Barbasita, in Atti Ist. Incorr. Nap.* vii. (1847) 34.—Italia.
lucidum, [Nutt.] Fras. Cat. (1813) n. 78, nomen ; ex Nutt. Gen. Am. i. 195 = Musenium divaricatum.
macedonicum, *Benth. & Hook. f. ex Baill. Hist. Pl.* vii. 192.—Reg. Mediterr.
macrophyllum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 31.—Turkestan.
Malyi, *Kern. in Oestr. Bot. Zeitschr.* xxxi. (1881) 37.—Europ. austr.
Meum, Scop. Fl. Carn. ed. II. i. 212 = Meum athamanticum.
Meyeri, D. Dietr. Syn. Pl. ii. 956 = Ferula Meyeri.
montanum, *Linn. Sp. Pl.* 259.—Europ.
montanum, Pall. ex Ledeb. Fl. Ross. ii. 274 = tenuifolium.
montanum, Vis. Fl. Dalm. iii. 42 = Tommasinii.
multicaule, Jacq. Hort. Vindob. ii. 59 = montanum.
Mutellina, Steud. Nom. ed. I. 530, 771 = Ligusticum Mutellina.
nanum, *Dufour, Lettres Mont. Maudit.* 363.—Mont. Pyren.
natalense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 550.—Afr. austr.
nitidum, Reichb. f. Ic. Fl. Germ. xxi. 36 = athamantoides.
nodiflorum, Scop. Fl. Carn. ed. II. i. 213 = Apium nodiflorum.
Nuttalli, *A. Gray, in Proc. Am. Acad.* viii. (1870) 287.—Mexic.
oligophyllum, *Griseb. Spicil. Fl. Rumel.* i. 359.—Macedon.
Olivieri, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 298.—Persia.
osseum, *Crantz, Stirp. Austr.* ed. I. fasc. iii. 92 ; ed. II. 207.—Europ.
Pallasii, *Bess. Cat. Hort. Crem.* (1816) 130.—Rossia.
parnassicum, *Boiss. & Heldr. Diagn.* Ser. II. vi. 80.—Graecia.
patens, Gueldenst. Reisen, i. 108, 190.—Cf. Schult. Syst vi. 401 = tortuosum.

SESELI :—

petraeum, *Bieb. Fl. Taur. Cauc.* i. 235.—Reg. Caucas.
peucedanifolium, Bess. Enum. Pl. Volh. 44 = rigidum.
peucedanifolium, Mérat, Fl. Par. 115 = montanum.
pimpinelloides, Linn. Sp. Pl. 259 = Carum rigidulum.
pimpinelloides, Spreng. Syst. i. 885 = Peucedanum carvifolium.
polyphyllum, *Tenore, Ind. Sem. Hort. Neap.* (1825) 12. n. 25.—Italia.
pratense, Crantz, Stirp. Austr. ed. I. fasc. iii. 96 ; ed. II. 209 = Silaus flavescens
proliferum, Brocchi, in Bibl. Ital. xxviii. (1822) 218 = Trinia vulgaris.
proliferum, Spreng. Pugill. ii. 57 = rigidum.
promonense, Vis. Fl. Dalmat. iii. t. 29 = varium.
puberulum, DC. Prod. iv. 147 = tortuosum.
pumilum, All. ex DC. l. c. 104 = Trinia vulgaris.
pumilum, Linn. Syst. ed. X. 962 = Trinia Kitaibelii.
purpurascens, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 177.—Balkan.
purpureum, Gilib. Fl. Lituan. ii. 35 = annuum.
pusillum, Brot. Fl. Lusit. i. 457 = Carum ammoides.
pyrenaeum, Linn. Sp. Pl. 261 = Selinum pyrenaeum.
pyrenaicum, *Siev. in Pall. N. Nord. Beitr.* vii. 324 (sp. dub.).—Soongaria.
quinquefolium, Bess. Cat. Hort. Crem. (1811) ; ex Schult. Syst. vi. 407 (Quid ?).—Hab. ?
radicosum, Boiss. & Hohen. Diagn. Ser. I. x. 30 = Olivieri.
rigidum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 156. t. 146.—Europ. or.
saxifragum, Linn. Sp. Pl. 261 = Carum Bunius.
saxifragum, Scholler, Fl. Barb. 307 = Selinum sylvestre.
scabriusculum, Kit. in Linnaea, xxxii. (1863) 461 = carinatum.
scariosum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 361.—Sibir. altaic.
segetum, Crantz, Cl. Umb. Emend. 91 = Carum segetum.
selinoides, Bess. Cat. Hort. Crem. (1816) 130 = Selinum sylvestre.
selinoides, Jacq. Enum. Stirp. Vindob. 51, 227 = Silaus flavescens.
sessiliflorum, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 307.—Soongaria.
sibiricum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 693.—Reg. Himal.
sibiricum, Garcke, Fl. Deutschl. ed. I. 139 ; Reichb. f. Ic. Fl. Germ. xxi. 36. t. 1916 ; Boiss. Fl. Orient. ii. 967 = S. Libanotis.
Sibthorpii, Gren. & Godr. Fl. Fr. i. 711 (= *Libanotis Candollei*).—Europ.
simplex, Poir. Encyc. vii. 137 = annuum.
Sisarum, Crantz, Cl. Umb. Emend. 92 = Sium Sisarum.
striatum, Thunb. Prod. Pl. Cap. 51 = Peucedanum collinum.
strictum, *Ledeb. Fl. Alt.* i. 338.—Sibir. altaic.
stylare, Boiss. & Heldr. Diagn. Ser. I. vi. 60 = Sclerochorton junceum.
Tachiroei, *Franch. & Sav. Enum. Pl. Jap.* ii. 373.—Japon.
tauricum, Link, ex Spreng. Syst. i. 884 = tortuosum.
tenuifolium, *Ledeb. Ic. Pl. Alt.* i. t. 97.—Sibir. altaic.
tenuisectum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 31.—Turkestan.
teres, Moench, Meth. 88 = elatum.
tomentosum, *Vis. Stirp. Dalm. Rar.* 6. t. 3.—Dalmatia.
Tommasinii, *Reichb. f. Ic. Fl. Germ.* xxi. 34. t. 2045. —Dalmat. ; Graec.
tortuosum, Boeber, in Pall. N. Nord. Beitr. vi. 275 = campestre.
tortuosum, *Linn. Sp. Pl.* 260.—Europ. austr. ; As. Min.
tortuosum, Pollich, Hist. Pl. Palat. i. 300 = annuum.
tortuosum, Sibth. & Sm. Fl. Graec. Prod. i. 200 = Johrenia graeca.
transsilvanicum, Schur, Enum. Pl. Transs. 257 = varium.
trilobum, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 694.—Ind. or.
triternatum, Pursh, Fl. Am. Sept. i. 197 = Peucedanum triternatum.
Turbith, Linn. Cent. Pl. ii. 14 ; Amoen. Acad. iv. 310 = Athamanta Matthioli.
Turbith, Tenore, Prod. Fl. Nap. p. xx = Carum ammoides.

SESELI:—
vaginatum, *Ledeb. Fl. Alt.* i. 336.—Dahur.
varium, Trev. Ind. Sem. Hort. Vratisl. (1818) 6 = glaucum.
venosum, Hoffm. Deutschl. Fl. 144 = Selinum sylvestre.
verticillatum, Crantz, Cl. Umb. Emend. 92 = Carum verticillatum.
verticillatum, Desf. Fl. Atlant. i. 260 = Carum ammoides.
verticillatum, Reichb. f. Ic. Fl. Germ. xxi. 37 = Carum graecum.
vulgatum, Bor. Fl. Centr. Fr. ed. III. ii. 279 = Carum Lomatocarum.
Webbii, Coss. in Bull. Soc. Bot. Fr. iii. (1856) 57 = Foeniculum tortuosum.
zeylanicum, Herb. Rottl. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 684 = Pimpinella Heyneanum.

SESLERA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.

SESLERIA, Nutt. Gen. Am. i. 64 (1818) = **Buchloë**, Engelm. (Gramin.).
dactyloides, Nutt. l. c. 65 = B. dactyloides.

SESLERIA, Scop. Fl. Carn. ed. I. 63 (1760). *GRAMINEAE*, Benth. & Hook. f. iii. 1181.
DIPTYCHUM, Dulac, Fl. Hautes-Pyr. 81 (1867).
PSILATHERA, Link, Hort. Berol. i. 121 (1827).
alba, Sibth. & Sm. Fl. Graec. i. 56. t. 72 = argentea.
albicans, *Kit. in Schult. Oestr. Fl.* ed. II. i. 216.—Mont. Carpat.
americana, *Nees, ex Steud. Syn. Pl. Gram.* 296.—Am. bor.
Arduini, Desf. ex Steud. Nom. ed. II. ii. 573 = coerulea.
argentea, *Savi, Bot. Etrusc.* i. 68.—Europ. austr.; As. Min.
autumnalis, F. Schultz, Arch. Fl. 318 = argentea.
Bielzii, Griseb. in Wiegm. Archiv, xxv. (1852) 361, in syn. = rigida.
Bielzii, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 109; iv. (1853) 84 = coerulans.
brevifolia, Panz. in Denkschr. Akad. Muench. 1813 (1814) 269 = coerulea.
calcaria, Pers. Syn. i. 72 = coerulea.
capitata, Schur, Enum. Pl. Transs. 743 = coerulans.
caucasica, Scheele, in Flora, xxvii. (1844) 55 = phleoides.
coerulans, *Frivald. in Flora*, xix. (1836) 438.—Transsylv.; Banat.
coerulea, *Arduin. Animad.* ii. 18. t. 6. f. 3–5.—Europ.
cristata, [Clairv.] Man. Herb. 23 = coerulea?
cylindrica, DC. Syn. Gall. 134; Fl. Fr. Suppl. 279 = argentea.
disticha, Pers. Syn. i. 72 = Oreochloa disticha.
dura, Kunth, Rév. Gram. i. 110 = Sclerochloa dura.
echinata, Lam. Illustr. i. 198; Host, Gram. Austr. iii. 6. t. 8 = Echinaria capitata.
elongata, Host, l. c. ii. 69. t. 97 = argentea.
filifolia, Heuff. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 361 = coerulans.
filifolia, Hoppe, in Flora, xvii. (1834) 384 = tenuifolia.
flexuosa, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 91 = coerulans.
gracilis, Schur, l. c. iv. (1853) 84 = tenuifolia.
Haynaldiana, Fuss, Fl. Transs. 717 (1866) = coerulans.
Haynaldiana, Schur, in Verh. Zool.-Bot. Ges. Wien, vi. (1856) 207 = rigida.
Heufleriana, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 84.—Europ.
Heufleriana, Schur, in Verh. Zool.-Bot. Ges. Wien, vi. (1856) 207 = coerulans.
interrupta, Vis. Fl. Dalmat. i. 87 = tenuifolia.
juncifolia, Suffren, Cat. Pl. Frioul, 113 = tenuifolia.
lagopodioides, Spreng. Pugill. ii. 22 = Aeluropus pubescens.
leucocephala, DC. Fl. Fr. iii. 76 = sphaerocephala.
macrocephala, Link, Handb. i. 81, sphalm. = *microcephala*.
marginata, Griseb. Spicil. Fl. Rumel. ii. 442 = coerulans.
microcephala, DC. Fl. Fr. iii. 76 = tenella.
mollis, Spreng. Syst. i. 329 = Triraphis mollis.
mutica, Badaro, in Brugnat. Giorn. Fis. Dec. II. vii. (1824) 363 = coerulea.
nitida, Heldr. ex Nym. Consp. 796 = Heufleriana.
nitida, C. A. Mey. Verz. Pfl. Cauc. 21 = phleoides.

SESLERIA:—
nitida, Tenore, Prod. Fl. Nap. p. x. = argentea.
ovata, Kern. Sched. Fl. Exsicc. i. 109 = tenella.
pallescens, Hartm. ex Kunth, Enum. Pl. i. 322 = argentea.
pedemontana, Reut. in Boiss. & Reut. Pugill. 127 = Oreochloa pedemontana.
permixta, Schur, Enum. Pl. Transs. 744 = coerulans.
phleoides, *Stev. ex Roem. & Schult. Syst.* ii. 604.—As. Min.; Persia.
polyathera, C. Koch, in Linnaea, xxi. (1848) 399 = phleoides.
Pseudo-rigida, Schur, Enum. Pl. Transs. 745 = rigida.
Pumilio, Spreng. Syst. iv. Cur. Post. 34 = Triraphis Pumilio.
pungens, Spreng. l. c. i. 329 = Triraphis pungens.
quitensis, Spreng. l. c. = Festuca quadridentata.
rigida, *Heuff. ex Reichb. Fl. Germ. Excurs.* 140*.—Banat.
rigida, Schur, Enum. Pl. Transs. 743 = coerulans.
robusta, Pavai, in Oestr. Bot. Zeitschr. xii. (1862) 214 = transilvanica.
robusta, Schott, Nym. & Kotschy, Analect. Bot. 1 = argentea.
Sadleriana, Janka, in Oestr. Bot. Zeitschr. xxxii. (1882) 309; et in Termesz. Füzetek, viii. (1884) 28 = transilvanica.
sphaerocephala, *Arduin. Animad.* ii. 20. t. 7.—Europ. alp.
spicata, Spreng. Pugill. ii. 21 = Elytrophorus articulatus.
subacaulis, Bal. ex Boiss. Fl. Orient. v. 566 = Ammochloa subacaulis.
tenella, *Host, Gram. Austr.* ii. 71. t. 100.—Europ. alp.
tenella, Schur, Enum. Pl. Transs. 746 = rigida.
tenuifolia, *Schrad. Fl. Germ.* i. 272.—Europ. austr.
tenuifolia, Schur, Enum. Pl. Transs. 746 = rigida.
transilvanica, *Schur, in Verh. Zool.-Bot. Ges. Wien*, vi. (1856) 205.—Transsylv.
transsilvanica, Schur, Enum. Pl. Transs. 745 = Heufleriana.
vaginalis, Boiss. & Orph. Diagn. Ser. II. iv. 130 = coerulans.
vallesiaca, Beauv. ex Steud. Nom. ed. I. 339, 772 = Festuca ovina.
variegata, [Clairv.] Man. Herb. 23 = Koeleria setacea.

SESQUICELLA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 255 = **Callirhoë**, Nutt. (Malvac.).
alceoides, Alef. l. c. 256 = C. alceoides.
digitata, Alef. l. c. = C. digitata.
involucrata, Alef. l. c. = C. involucrata.
macrorhiza, Alef. l. c. = C. alcaeoides.
Papaver, Alef. l. c. = C. Papaver.
pedata, Alef. l. c. = C. pedata.
triangulata, Alef. l. c. = C. triangulata.

SESSEA, Ruiz & Pav. Prod. 21. t. 33 (1794). *SOLANACEAE*, Benth. & Hook. f. ii. 905.
atrovirens, *Benth. & Hook. f. Gen.* ii. 904.—Peruv.
corymbosa, *Goudot, in Phil. Mag.* iii. (1828) 132.—N. Granat.
dependens, *Ruiz & Pav. Fl. Per.* ii. 9. t. 116.—Peruv.
Dombeyi, *Dun. in DC. Prod.* xiii. 1. 596.—Peruv.
petiolaris, *Spreng. Syst.* i. 584.—Peruv.
stipulata, *Ruiz & Pav. Fl. Per.* ii. 9. t. 115.—Peruv.
vestita, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 155.—N. Granat.

SESSLERIA, Spreng. Pugill. ii. 21 (1815), sphalm. = **Sesleria**, Scop. (Gramin.).

SESTINIA, Boiss. Diagn. Ser. I. v. 40 (1844) = **Hymenocrater**, Fisch. & Mey. (Labiat.).
calycina, Boiss. l. c. 41 = H. calycinus.

SESTINIA, Boiss. & Hohen. Diagn. Ser. I. iii. 55 (1843) = **Wendlandia**, Bartl. (Rubiac.).
ligustroides, Boiss. & Hohen. l. c. = W. Kotschyi.

SESTOCHILOS, Breda, Orch. Kuhl & Hass. [t. 3] (1827) = **Bulbophyllum**, Thou. (Orchid.).
uniflorum, Breda, l. c. = B. Lobbii.

SESUVIUM, Linn. Syst. ed. X. 1058 (1759). *FICOIDEAE*, Benth. & Hook. f. i. 855.
PSAMMANTHE, Hance, in Walp. Ann. ii. 659 (1852).
PYXIPOMA, Fenzl, in Ann. Wien. Mus. ii. (1838) 293.
acutifolium, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 75.—Guian.
brevifolium, Schum. & Thonn. Beskr. Guin. Pl. 233 = S. Portulacastrum.
congense, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 586.—Afr. trop.
crithmoides, *Welw. Apont.* 586.—Afr. trop.
crystallinum, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 586.—Afr. trop.
digynum, *Welw. ex Oliver, l. c.*—Afr. trop.
Edmonstonei, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 221.—Ins. Galapag.
edule, Wight, ex Wall. Cat. n. 6836 D = S. Portulacastrum.
longifolium, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 521 = S. Portulacastrum.
mesembrianthemoides, *Wawra, in Sitzb. Akad. Wien, Math.-Nat.* xxxviii. (1860) 564.—Afr. trop.
mesembryanthemoides, Welw. Apont. 557 = S. Portulacastrum.
microphyllum, *Willd. Enum. Hort. Berol.* 521.—Cuba.
Ortegae, Spreng. Nachr. i. Bot. Gart. Halle, 36 = S. Portulacastrum.
parviflorum, DC. Prod. iii. 453 = S. Portulacastrum.
pedunculatum, Pers. Syn. ii. 39 = S. Portulacastrum.
pentandrum, Ell. Sketch, i. 556 = S. Portulacastrum?
Portulaca, Crantz, Inst. i. 136 = seq.
Portulacastrum, *Linn. Syst.* ed. X. 1058.—Reg. trop.
quadrifidum, *F. Muell. Rep. Babb. Exp.* 9.—Austral.
repens, Willd. Enum. Hort. Berol. 521 = S. Portulacastrum.
revolutifolium, *Colla, ex Ces. in Atti Soc. Acc. Sc. Nap.* v. (1873) n. VII. 10.—Reg. Argent.
revolutifolium, Ortega, Hort. Matr. Dec. 19 = S. Portulacastrum.
revolutum, Pers. Syn. ii. 39 = S. Portulacastrum.
sessile, Pers. l. c. = S. Portulacastrum.
sessiliflorum, Domb. ex Rohrb. in Mart. Fl. Bras. xiv. II. 310 = S. Portulacastrum.
spathulatum, H. B. & K. Nov. Gen. et Sp. vi. 87 = microphyllum.
verrucosum, Rafin. New Fl. Am. iv. 16 = S. Portulacastrum?

SETACHNA, Dulac, Fl. Hautes-Pyr. 518 (1867) = **Centaurea**, Linn. (Compos.).
amara, Dulac, l. c. = C. amara.
Calcitrapa, Dulac, l. c. 519 = C. Calcitrapa.
collina, Dulac, l. c. 520 = C. collina.
Cyanus, Dulac, l. c. = C. Cyanus.
fimbriata, Dulac, l. c. 519 = C. Jacea.
microptilon, Dulac, l. c. = C. nigra.
montana, Dulac, l. c. = C. montana.
nigra, Dulac, l. c. = C. nigra.
nigrescens, Dulac, l. c. = C. nigrescens.
solstitialis, Dulac, l. c. 520 = C. solstitialis.
sylvatica, Dulac, l. c. = C. Scabiosa.

SETARIA, Beauv. Fl. Owar. ii. 80 (1807). *GRAMINEAE*, Benth. & Hook. f. iii. 1105.
IXOPHORUS, Schlecht. in Linnaea, xxxi. (1861–62) 420.
affinis, Schult. Mant. ii. 275 = viridis.
Alopecurus, Fisch. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) 227 = macrostachya.
ambigua, Guss. Fl. Sic. Syn. i. 114 = viridis.
ambigua, Schrad. in Linnaea, xii. (1838) 430 = Panicum Pseudoholcus.
Antillarum, Kunth, Rév. Gram. i. 46 = setosa.
asiatica, Hort. ex Reichb. Fl. Germ. Excurs. 29 = italica.
aspera, Link, ex Loud. Hort. Brit. 25 (Quid?).—Afr. austr.
aspera, *Miq. Fl. Ind. Bat.* iii. 469.—Malaya.
atroseta, Steud. Syn. Pl. Gram. 102 = Pennisetum compressum.
aurea, Hochst. ex A. Br. in Flora, xxiv. (1841) 276 = glauca.
auricoma, Link, ex Loud. Hort. Brit. 25; et ex A. Br. l. c. 277 = glauca *vel* helvola.
auriculata, *Fourn. ex Hemsl. Biol. Centr. Am.* iii. 503. —Mexic.

SETARIA :—
barbata, Kunth, Rév. Gram. i. 47 = Panicum prostratum.
Berteroniana, Schult. Mant. ii. 276, in obs. = corrugata ?
biconvexa, *Griseb. Fl. Brit. W. Ind.* 555.—Ins. Trinit.
brachiata, *Kunth, Rév. Gram.* i. 47.—Bras.
Brownii, Desv. Opusc. 81, in obs. = Panicum gracile.
californica, *Kellogg, in Proc. Calif. Acad.* i. ed. II. (1873) 26 (sp. dub.).—Calif.
canescens, *Kunth, Rév. Gram.* i. 47.—Ind. or.
caudata, Roem. & Schult. Syst. ii. 495 = setosa.
cenchroides, Roem. & Schult. l. c. = Pennisetum setosum.
cernua, *H. B. & K. Nov. Gen. et Sp.* i. 111.—Ecuador.
chlorantha, Schur, Enum. Pl. Transs. 723 = viridis.
chrysantha, Heynh. Nom. ii. 661 = glauca.
cirrosa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 504.—Mexic.
comosa, *Miq. Fl. Ind. Bat.* iii. 468.—Ins. Philipp.
compacta, Schur, Enum. Pl. Transs. 967 (Index) = italica.
composita, H. B. & K. Nov. Gen. et Sp. i. 111 = macrostachya.
compressa, Kunth, Rév. Gram. i. 46; Enum. Pl. i. 150 = Pennisetum compressum.
conglomerata, *Fries, ex Schlecht. in Linnaea*, xxxi. (1861–62) 491.—Hab. ?
corrugata, *Schult. Mant.* ii. 276.—Cuba.
Crus-ardeae, Kunth, Enum. Pl. i. 155 = Panicum Crus-ardeae.
cylindrica, Schult. Mant. ii. 277 = setosa.
dasyura, *Hort. Hal. ex Schlecht. in Linnaea*, xxvi. (1853) 463.—Ind. or.
decipiens, C. Schimp. ex Nym. Consp. 787 = viridis.
dioica, Hochst. in Flora, xxiv. (1841) I. Intell. 19 = Pennisetum adoënse.
disticha, H. B. & K. Nov. Gen. et Sp. i. 112 = Panicum pilosum.
effusa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 504.—Mexic.
Elliotiana, Schult. Mant. ii. 279 = Cenchrus Elliotii.
elongata, *Spreng. in Schult. l. c.* 280.—Am. bor.
erubescens, Beauv. Agrost. 51; Roem. & Schult. Syst. ii. 491 = Pennisetum erubescens.
erythrosperma, Hornem. ex Reichb. Fl. Germ. Excurs. 29 = italica.
erythrosperma, Spreng. Syst. i. 304 = italica, viridis ?
excurrens, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 275.—Japon.
falcifolia, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 504.—Mexic.
Fieldingii, *C. Muell. in Bot. Zeit.* xix. (1861) 323.—Ind. or.
flava, Kunth, Rév. Gram. i. 46 = glauca.
flavida, Hornem. ex Reichb. Fl. Germ. Excurs. 29 = italica.
floriana, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 138.—Ins. Galapag.
floribunda, Spreng. Syst. i. 305 = verticillata.
geniculata, *Beauv. Agrost.* 51; *Roem. & Schult. l. c.* ii. 491.—Mexic.
geniculata, Sieber, ex Kunth, Rév. Gram. i. 50 = Pennisetum Sieberi.
germanica, Beauv. Agrost. 51. t. 13 = italica.
glauca, *Beauv. Agrost.* 51.—Europ.; As. temp.
glauca, H. B. & K. Nov. Gen. et Sp. i. 109 = penicillata, *J. Presl.*
globularis, J. Presl, in Rel. Haenk. i. 314 = italica.
globulifera, Griseb. in Goett. Abh. xxiv. (1879) 307 = Panicum globuliferum.
gracilis, *H. B. & K. Nov. Gen. et Sp.* i. 109.—N. Granat.
gracilis, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) 226 = P. Onurus.
granosa, *Desv. Opusc.* 79.—Bras.
Grisebachii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 504.—Mexic.
helvola, Roem. & Schult. Syst. ii. 491 = glauca.
hirsuta, *Kunth, Rév. Gram.* i. 47.—Ind. or.
hirta, *Kunth, l. c.*—Ind. or.
hirtella, Schult. Mant. ii. 276 = Oplismenus undulatifolius.
Hochstetteri, *Kunth, in Linnaea*, xix. (1847) 380.—Nubia.
humifusa, Dum. Obs. Gram. Belg. 139 = glauca.
imberbis, Roem. & Schult. Syst. ii. 891 = viridis.
impressa, *Kunth, Enum. Pl.* i. 154.—Bras.

SETARIA :—
intermedia, Roem. & Schult. Syst. ii. 489 = verticillata.
interrupta, *Desv. Opusc.* 80.—Bras.
italica, *Beauv. Agrost.* 51.—Reg. trop. et subtrop. (cult.).
italica, Miq. Fl. Ind. Bat. iii. 467 = Panicum sanguinale.
Itieri, *Delile, Ind. Sem. Hort. Monsp.* (1849); *ex Steud. Syn. Pl. Gram.* 51.—Aegypt.
japonica, Pynaert, in Fl. des Serres, xiv. (1861) 7 = italica ?
Jurgensenii, *Fourn. ex Hemsl. Biol. Centr. Am.* iii. 505.—Mexic.
lachnea, Kunth, Enum. Pl. i. 154 = setosa.
laevigata, *Schult. Mant.* ii. 275.—Am. bor.
laevis, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
leiophylla, *Nees, ex Kunth, Enum. Pl.* i. 154.—Bras.
lenis, *Miq. Fl. Ind. Bat.* iii. 468.—Java.
leucophylla, D. Dietr. Syn. Pl. i. 293 = Panicum leiophyllum.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
longicauda, *Desv. Opusc.* 77.—Bras.
longiflora, Desv. l. c. 81, in obs. = Chamaeraphis paradoxa.
longifolia, *Link, Hort. Berol.* i. 220.—Hab. ?
longipila, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
longiseta, *Beauv. Fl. Owar.* ii. 81. t. 110.—Afr. trop.
luxurians, *Spreng. Syst.* i. 305.—Ind. or.
macrochaeta, Hochst. ex Steud. Nom. ed. II. ii. 574 = Pennisetum adoënse.
macrochaeta, Schult. Mant. ii. 274 = italica.
macrochaeta, *Spreng. Syst.* i. 305.—Ind. or.
macrophylla, *Anderss. in Peters, Reise Mossamb. Bot.* 550.—Afr. trop.
macrostachya, Hochst. ex Steud. Syn. Pl. Gram. 53 = composita.
macrostachya, *H. B. & K. Nov. Gen. et Sp.* i. 110.—As. et Austral. trop.
magna, *Griseb. Fl. Brit. W. Ind.* 554.—Ind. occ.
maritima, Roem. & Schult. Syst. ii. 492 = italica.
mauritiana, Spreng. l. c. i. 305 = Panicum plicatum.
maxima, *Desv. Opusc.* 78.—Bras.
megapotamica, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) 226 = setosa.
Melinis, Link, ex Steud. Nom. ed. II. ii. 574 = italica.
Meyeri, Kunth, Rév. Gram. i. 47 = Panicum pilosum.
multiseta, Dum. Obs. Gram. Belg. 138 = setosa.
muricata, Beauv. Agrost. 51; Roem. & Schult. Syst. ii. 495 = Panicum Crus-galli.
nana, Dum. Obs. Gram. Belg. 139 = viride.
nubica, Link, Hort. Berol. i. 220 = verticillata.
ocreata, *Spreng. Syst.* i. 305.—Ind. occ.
Onurus, *Griseb. Fl. Brit. W. Ind.* 555.—Am. trop.
paniculifera, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
panis, Jessen, Deutschl. Graes. 248 = italica.
penicillata, *E. Desv. in C. Gay, Fl. Chil.* vi. 247.—Chili.
penicillata, Nees, in Wall. Cat. n. 8640 = glauca, viridis.
penicillata, *J. Presl, in Rel. Haenk.* i. 314.—Mexic.
Pennisetum, Roem. & Schult. Syst. ii. 494 = Panicum pilosum.
persica, Hort. ex Reichb. Fl. Germ. Excurs. 29 = italica.
pilifera, *Desv. Opusc.* 81.—Am.
pilifera, Spreng. Syst. iv. Cur. Post. 33 = Panicum javanicum.
pilosa, Kunth, Rév. Gram. i. 47 = Panicum pilosum.
Poiretiana, *Kunth, l. c.*—Bras.
polygonata, Kunth, l. c. = Panicum pilosum.
polystachya, *Scheele, in Linnaea*, xxii. (1849) 339.—Texas.
polystachya, Schrad. ex Schult. Mant. ii. 277 = macrostachya.
Pseudo-verticillata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
Pseudo-verticillata, Hort. Vindob. ex Schlecht. in Linnaea, xxxi. (1861–62) 491 = conglomerata.
pumila, Roem. & Schult. Syst. ii. 891 = glauca.
pumila, *Schult. Mant.* ii. 274.—Hab. ?
purpurascens, H. B. & K. Nov. Gen. et Sp. i. 110; Opiz, Boehm. Gew. (1823) 12 = imberbis.
purpurea, *Beauv. Agrost.* 51; *Roem. & Schult. Syst.* ii. 488.—Peruv.
pyramidata, *Desv. Opusc.* 78.—Am. trop.
rariflora, Mikan, ex Spreng. Neue Entdeck. ii. 78 = setosa.

SETARIA :—
rariflora, *J. Presl, in Rel. Haenk.* i. 313.—Mexic.
refracta, F. Muell. Fragm. iii. 147 = Plagiosetum refractum.
respiciens, Walp. Ann. iii. 721 = verticillata.
Rottleri, Spreng. Syst. i. 304 = verticillata.
rubicunda, Dum. Obs. Gram. Belg. 139 = italica.
rubiginosa, *Miq. Fl. Ind. Bat.* iii. 467.—Ins. Philipp.
sagittifolia, *Walp. Ann.* iii. 721.—Afr. trop.
scabrifolia, *Kunth, Rév. Gram.* ii. t. 212.—Bras.
scandens, *Schrad. ex Schult. Mant.* ii. 279.—Bras.
Schiedeana, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 505.—Mexic.
Schraderi, Kunth, Rév. Gram. i. 47 = Panicum pilosum.
sciuroidea, *C. Muell. in Bot. Zeit.* xix. (1861) 316.—Ind. or.
semirugosa, *Kunth, Enum. Pl.* i. 152.—Mexic.
sericea, *Beauv. Agrost.* 51; *Roem. & Schult. Syst.* ii. 488.—Ind. or.
setosa, *Beauv. l. c.*; *Roem. & Schult. l. c.* 494.—Reg. trop.
stipaeculmis, *C. Muell. in Bot. Zeit.* xix. (1861) 323.—Bras.
streptobotrys, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 506.—Mexic.
stricta, Kunth, Rév. Gram. i. 47 = Panicum Kunthii.
subcordata, Kunth, l. c. = Panicum prostratum.
sulcata, *Raddi, Agrost. Bras.* 50.—Mexic.
tejucensis, Kunth, Enum. Pl. i. 150 = glauca.
tenacissima, Schrad. ex Schult. Mant. ii. 279 = scandens.
tenax, *Desv. Opusc.* 78.—Guian.
tenella, *Desv. l. c.* 77.—Peruv.
Teysmanni, *Miq. Fl. Ind. Bat.* iii. 468.—Java.
tomentosa, *Kunth, Rév. Gram.* i. 47.—Ind. or.
Trinii, Kunth, Enum. Pl. i. 151 = scandens.
umbrosa, Beauv. Agrost. 51 = Panicum prostratum.
uniseta, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 506.—Mexic.
vaginata, *Spreng. Syst.* iv. *Cur. Post.* 33.—Bras.
Ventenatii, Kunth, Rév. Gram. i. 251. t. 37 = glauca.
verticillata, *Beauv. Agrost.* 51.—Cosmop.
verticilliformis, Dum. Fl. Belg. 150 = verticillatum.
villosa, Beauv. Agrost. 51 = viride.
violacea, Hornem. ex Reichb. Fl. Germ. Excurs. 29 = italica.
viridis, *Beauv. Agrost.* 51.—Cosmop.
viscidula, *Desv. Opusc.* 80.—Ins. Maurit.
vulpina, Beauv. Agrost. 51 = Pennisetum cenchroides.
vulpiseta, Raddi, Agrost. Bras. 49 = macrostachya.
vulpiseta, *Roem. & Schult. Syst.* ii. 495.—Am. trop.
Wienmanni, Roem. & Schult. l. c. 490 = viridis.

SETHIA, H. B. & K. Nov. Gen. et Sp. v. 175, in nota (1821) = **Erythroxylum**, P. Br. (Lineae).
acuminata, Arn. in Nov. Act. Nat. Cur. xviii. I. (1836) 324 = E. lucidum.
erythroxyloides, Wight, Illustr. i. 136 = E. lanceolatum.
indica, DC. Prod. i. 576 = E. monogynum.
Kunthiana, Wall. Cat. n. 6849 = E. Kunthianum, *Kurz.*
lanceolata, Wight, Illustr. i. 136 = E. lanceolatum.
magnoliaefolia, G. Don, Gen. Syst. i. 633 = E. magnoliaefolium.
microphylla, G. Don, l. c. = E. microphyllum.
oblonga, Wight, Ic. t. 97, sphalm. = Salacia oblonga.
obtusifolia, Thw. Enum. Pl. Zeyl. 54 = E. obtusifolium.

SEUBERTIA, Kunth, Enum. Pl. iv. 475 (1843) = **Brodiaea**, Sm. (Liliac.).
crocea, Wood, in Proc. Acad. Nat. Sc. Philad. (1868) 172 = B. crocea.
laxa, Kunth, Enum. Pl. iv. 475 = B. laxa.

SEUBERTIA, H. C. Wats. in Hook. Lond. Journ. Bot. iii. (1844) 602 = **Bellis**, Tourn. (Compos.).
azorica, H. C. Wats. l. c. = B. azorica.

SEUTERA, Reichb. Consp. 131 (1828) = **Vincetoxicum**, Moench (Asclep.).
maritima, Decne. in DC. Prod. viii. 590 = V. palustre.

SEVADA, Moq. in DC. Prod. xiii. II. 154 (1849) = **Suaeda**, Forsk. (Chenop.).
Schimperi, Moq. l. c. = Suaeda baccata.

SEVERINIA, Tenore, Ind. Sem. Hort. Neap. (1840) 3 = **Atalantia**, Correa (Rutac.).
buxifolia, Tenore, l. c. = A. buxifolia.

SEWERZOWIA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 580 = **Astragalus**, Tourn. (Legumin.).
turkestanica, Regel & Schmalh. l. c. 581 = A. Schmalhauseni.

SEXILIA, Rafin. New Fl. Am. iv. 87 (1836) = **Polygala**, Linn.
missurica, Rafin. l. c. 89 = P. missurica.
verticillata, Rafin. l. c. 87 = P. verticillata.

SEYMERIA, Pursh, Fl. Am. Sept. ii. 736 (1814). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 971.
AFZELIA, J. F. Gmel. Syst. 927 (1791).
auriculata, Spreng. Syst. ii. 810 = Gerardia auriculata.
bipinnatisecta, *Seem. Bot. Voy. Herald*, 323. t. 59.—Am. bor. occ.
decurva, *Benth. in DC. Prod.* x. 512.—Mexic.
heterophyla, Rafin. New Fl. Am. ii. 68 = pectinata.
Jacksoni, Ell. Sketch, ii. 123, in obs. = pectinata.
macrophylla, *Nutt. Gen. Am.* ii. 49 —Am. bor.
pectinata, *Pursh, Fl. Am. Sept.* ii. 737.—Am. bor.
pinnatifida, *Hemsl. Biol. Centr. Am. Bot.* ii. 458.—Mexic.
scabra, *A. Gray, in Torr. Bot. Mex. Bound.* 118.—Texas.
tenuifolia, *Pursh, Fl. Am. Sept.* ii. 737.—Am. bor.
virgata, *Benth. in DC. Prod.* x. 511.—Mexic.

SEYMOURIA, Sweet, Geran. iii. t. 206 (1830) = **Pelargonium**, L'Hérit. (Geran.).
asarifolia, Sweet, l. c. = P. asarifolium.
Heritieri, Sweet, l. c. sub t. 206 = P. dipetalum.

SGNGONIUM, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 178, err. typ. = **Syngonium**, Schott (Aroid.).

SHAKUA, Boj. Hort. Maurit. 82 (1837) = **Spondias**, Linn. (Anacard.).
excelsa, Boj. l. c. 82, nomen.—Madag.
minor, Boj. l. c.—Madag.
pubescens, Boj. l. c. = Spondias pubescens, *Baker*.

SHALLONIUM, Rafin. in Am. Monthly Mag. (1818) 266 = **Gaultheria**, Kalm (Eric.).
serrulatum, Rafin. l. c. = G. Shallon.

SHAWIA, Forst. Char. Gen. 95. t. 48 (1776) = **Olearia**, Moench (1802) (Compos.).
aculeata, Sch. Bip. in Pollichia, xviii–xix. (1861) 174 = O. ramulosa.
affinis, Sch. Bip. l. c. = O. heleophila.
albida, Sch. Bip. l. c. 175 = O. albida.
alpina, Sch. Bip. l. c. 174 = O. persoonioides.
arborescens, Sch. Bip. l. c. 172 = O. nitida.
argophylla, Baill. Hist. Pl. viii. 299 = O. argophylla.
argyrophylla, Sch. Bip. in Pollichia, xviii–xix. (1861) 173 = O. argophylla.
aspera, Sch. Bip. l. c. 174 = O. strigosa.
avicenniaefolia, Raoul, Choix Pl. Nouv. Zél. 19. t. 13 = O. avicenniaefolia.
axillaris, Sch. Bip. in Pollichia, xviii–xix. (1861) 172 O. axillaris.
brachyglossa, Sch. Bip. l. c. = O. axillaris.
candidissima, Sch. Bip. l. c. 174 = O. axillaris.
capitellata, Sch. Bip. l. c. 173 = O. axillaris.
chrysophylla, Sch. Bip. l. c. = O. chrysophylla.
chrysotricha, Sch. Bip. l. c. 174 (= *Eurybia chrysotricha*).—Austral.
ciliata, Sch. Bip. l. c. = O ciliata.
corymbosa, DC. Prod. vi. 264 = O. Forsteri.
cuneifolia, Sch. Bip. in Pollichia, xviii–xix. (1861) 174 = O. pimelioides.
Cunninghamii, Sch. Bip. l. c. 175 = O. Cunninghamii.
cydoniaefolia, Sch. Bip. l. c. 173 = O. cydoniaefolia.
Dampieri, Sch. Bip. l. c. = O. axillaris.
decurrens, Sch. Bip. l. c. = O. decurrens.
dentata, Sch. Bip. l. c. 175 = O. ilicifolia.
elaeophila, Sch. Bip. l. c. 174 = O. heleophila.
epileia, Sch. Bip. l. c. = O. ramulosa.
ericoides, Sch. Bip. l. c. = O. ramulosa.
erubescens, Sch. Bip. l. c. 173 = O. myrsinoides.

SHAWIA:—
floribunda, Sch. Bip. in Pollichia, xviii–xix. (1861) 175 = Olearia floribunda.
furfuracea, Raoul, ex Sch. Bip. l. c. 173 = O. furfuracea.
glandulosa, Sch. Bip. l. c. = O. glandulosa.
glutinosa, Sch. Bip. l. c. 174 = O. glutinosa.
gracilis, Sch. Bip. l. c. (= *Eurybia gracilis*).—Austral.
Gunniana, Sch. Bip. l. c. 173 = O. stellulata.
hygrophila, Sch. Bip. l. c. 174 = O. hygrophila.
imbricata, Sch. Bip. l. c. 175 = O. imbricata.
ledifolia, Sch. Bip. l. c. 173 = O. ledifolia.
Lehmanniana, Sch. Bip. l. c. 174 = O. heleophila.
lepidophylla, Sch. Bip. l. c. = O. lepidophylla.
leptophylla, Sch. Bip. l. c. 175 (= *Eurybia leptophylla*).—Austral.
linearifolia, Sch. Bip. l. c. 173 = O. axillaris.
linifolia, Sch. Bip. l. c. 175 = O. glutinosa.
lyrata, Sch. Bip. l. c. 173 = O. stellulata.
microphylla, Sch. Bip. l. c. 174 = O. ramulosa.
muricata, Sch. Bip. l. c. = O. muricata.
myrsinoides, Sch. Bip. l. c. 173 = O. myrsinoides.
nummularifolia, Sch. Bip. l. c. 175 = O. nummularifolia.
obcordata, Sch. Bip. l. c. = O. obcordata.
oligantha, Sch. Bip. l. c. 173 = O. axillaris.
paniculata, Forst. Char. Gen. 95. t. 48 = O. Forsteri.
paucidentata, Sch. Bip. in Pollichia, xviii–xix. (1861) 174 = O. paucidentata.
persoonioides, Sch. Bip. l. c. 173 = O. persoonioides.
pimeleoides, Sch. Bip. l. c. = O. pimeleoides.
pinifolia, Sch. Bip. l. c. 175 = O. pinifolia.
propinqua, Sch. Bip. l. c. 174 = O. ramulosa.
ramosissima, Sch. Bip. l. c. = O. ramosissima.
ramulosa, Sch. Bip. l. c. = O. ramulosa.
rosmarinifolia, Sch. Bip. l. c. 173 = O. rosmarinifolia.
rudis, Sch. Bip. l. c. 174 = O. rudis.
salicinia, Sch. Bip. l. c. 173 = O. stellulata.
scabra, Sch. Bip. l. c. 174 = O. rudis.
Solandri, Sch. Bip. l. c. 175 = O. Solandri.
Steeaii, Sch. Bip. l. c. 174 = O. heleophila.
strigosa, Sch. Bip. l. c. = O. strigosa.
subrepanda, Sch. Bip. l. c. 173 = O. stellulata.
tenuifolia, Sch. Bip. l. c. 174 = O. tenuifolia.
virgata, Sch. Bip. l. c. 175 = O. virgata.
viscosa, Sch. Bip. l. c. 173 = O. viscosa.

SHEADENDRON, Bertol. f. in Mem. Acc. Bologn. ii. (1850) 574. t. 40 = **Combretum**, Linn.
butyrosum, Bertol. f. l. c. 572 = C. butyrosum.
molle, Klotzsch, in Peters, Reise Mossamb. Bot. 76 = C. holosericum.
pisoniaeflorum, Klotzsch, l. c. = C. pisoniaeflorum.

SHEAREREIA, S. Moore, in Journ. Bot. xiii. (1875) 227. t. 165. *COMPOSITAE*, Benth. & Hook. f. ii. 1234.
nana, *S. Moore, l. c.*—China.
Polii, Franch. l. c. xvi. (1878) 257. t. 198 = nana.

SHEFFIELDIA, Forst. Char. Gen. 17. t. 9 (1776) = **Samolus**, Linn. (Primul.).
incana, Labill. Nov. Holl. Pl. i. 40. t. 54 = Samolus repens.
repens, Forst. Char. Gen. 18 = Samolus repens.

SHEPHERDIA, Nutt. Gen. Am. ii. 240 (1818). *ELAEAGNACEAE*, Benth. & Hook. f. iii. 204.
LEPTARGYREIA, Schlecht. in DC. Prod. xiv. 607 (1857).
argentea, Hort. ex Schlecht. in DC. Prod. xiv. 609 = Elaeagnus argentea.
argentea, *Nutt. Gen. Am.* ii. 240.—Am. bor.
canadensis, *Nutt. l. c.* 241.—Am. bor.
rotundifolia, *Parry, in Am. Natural.* ix. (1875) 350.—Am. bor. occ.

SHERARDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 176 = **Sherardia**, Dill. (Rubiac.).

SHERARDIA, Adans. Fam. ii. 198 (1763) = **Stachytarpheta**, Vahl (Verben.).

SHERARDIA, Dill. ex Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 151.
HEXODONTOCARPUS, Dulac, Fl. Hautes-Pyr. 467 (1867).

SHERARDIA:—
arvensis, *Linn. Sp. Pl.* 102.—Europ.; As. temp.
erecta, Sibth. & Sm. Fl. Graec. ii. 14. t. 116 = Galium verticillatum.
foetida, Lam. Encyc. iv. 326 = Putoria calabrica.
foetidissima, Cyrill. Char. Comm. (1784) 54 = Putoria calabrica.
fruticosa, Linn. Sp. Pl. 103 = Hedyotis Adscensionis.
minuta, Pall. ex Georgi, Beschr. Russ. Reich. III. iv. 727 = Galium verticillatum.
muralis, Linn. Sp. Pl. 103 = Galium murale.
muralis, Pall. ex Georgi, Beschr. Russ. Reich. iv. 727 = Galium verticillatum.
muralis, Stev. in Nouv. Mém. Soc. Nat. Mosc. i. (1829) 276 = Galium recurvatum.
neglecta, Guep. ex Nym. Consp. 335 = arvensis.
pusilla, Bory & Chaub. Nouv. Fl. Pélop. 10. t. 16. f. 1 = Galium Boryanum.
umbellata, Gilib. Exercit. i. 26 = arvensis.

SHERBOURNIA, G. Don, in Loud. Encyc. Pl. Suppl. ii. 1323 (1855). *RUBIACEAE*.
AMARALIA, Welw. ex Benth. & Hook. f. Gen. ii. 90 (1873).
foliosa, *G. Don, in Loud. Encyc. Pl. Suppl.* ii. 1322.—Afr. trop.

SHOREA, Roxb. in Gaertn. f. Fruct. iii. 48. t. 186 (1805). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 193.
PARASHOREA, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65.
PENTACME, A. DC. in DC. Prod. xvi. II. 626 (1868).
SAUL, Roxb. ex Wight & Arn. Prod. 84 (1834).
acuminata, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 305.—Malacca.
assamica, *Dyer, l. c.* 307.—Assam.
bracteolata, *Dyer, l. c.* 305.—Malacca.
brevipetiolaris, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 205 (Balanocarpi sp.).—Zeylan.
camphorifera, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 616.—Sumatra.
costata, J. S. Presl, Rostl. ii. 66 = Dryobalanops aromatica.
Dyerii, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 204.—Zeylan.
eximia, *Scheff. in Flora*, liii. (1870) 247; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 349.—Sumatra.
floribunda, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 60.—Burma.
furfuracea, *Miq. Fl. Ind. Bat. Suppl.* 488.—Sumatra.
gratissima, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 307.—Singapore.
Guiso, *Blume, Mus. Bot. Lugd. Bat.* ii. 34.—Ins. Philipp.
Helferi, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 100.—Burma.
hypochra, *Hance, in Journ. Bot.* xiv. (1876) 242.—Cambodia.
laccifera, Heyne, ex Wall. Cat. n. 967 = S. Talura.
laurifolia, Wall. ex Steud. Nom. ed. II. ii. 575 = S. Talura.
lepida, *Blume, Mus. Bot. Lugd. Bat.* ii. 32.—Sumatra.
leprosula, *Miq. Fl. Ind. Bat. Suppl.* 487.—Sumatra.
leucobotrya, Miq. Ann. Mus. Bot. Lugd. Bat. i. 215 = obtusa.
lissophylla, *Thw. Enum. Pl. Zeylan.* 402.—Zeylan.
longisperma, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 618.—Penang.
lucida, *Miq. Fl. Ind. Bat. Suppl.* 487.—Sumatra.
macroptera, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 308.—Malaya.
Malaanonan, *Blume, Mus. Bot. Lugd. Bat.* ii. 34.—Ins. Philipp.
Mangachapoi, *Blume, l. c.*—Ins. Philipp.
Martiniana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 408.—Borneo.
Mayapis, *Blume, Mus. Bot. Lugd. Bat.* ii. 33.—Ins. Philipp.
nervosa, *Kurz, For. Fl. Brit. Burma*, i. 119.—Burma.
nitens, *Miq. Fl. Ind. Bat. Suppl.* 488.—Sumatra.
oblongifolia, *Thw. Enum. Pl. Zeyl.* 36.—Zeylan.
obtusa, *Wall. Cat.* n. 966.—Burma.
ovalis, *Blume, Mus. Bot. Lugd. Bat.* ii. 33.—Borneo.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 487.—Sumatra.
parvifolia, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 305.—Malacca.

SHOREA:—
penicillata, A. DC. in DC. Prod. xvi. II. 630 = S. Tumbuggaia.
Pierrei, *Hance, in Journ. Bot.* xvi. (1878) 302.—Cambodia.
Pinanga, *Scheff. in Flora*, liii. (1870) 247; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 350.—Malaya.
purpurea, A. DC. in DC. Prod. xvi. II. 632, sphalm. = furfuracea.
reticulata, *Thw. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 307.—Zeylan.
robusta, A. DC. in DC. Prod. xvi. II. 628 = floribunda.
robusta, *Gaertn. f. Fruct.* iii. 48. t. 186.—Ind. or.
robusta, Roth, Nov. Pl. Sp. 221 = S. Talura.
Roxburghii, G. Don, Gen. Syst. i. 813 = S. Talura.
Schefferiana, *Hance, in Journ. Bot.* xvi. (1878) 303.—Borneo.
selanica, *Blume, Mus. Bot. Lugd. Bat.* ii. 33.—Molucca.
sericea, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 306.—Malacca.
siamensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 214.—Burma; Siam.
squamata, *Benth. & Hook. f. Gen.* i. 193.—Malaya.
stellata, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 304.—Burma.
stipularis, *Thw. Enum. Pl. Zeyl.* 36.—Zeylan.
suavis, Wall.—Cf. Dyer, in Hook. f. Fl. Brit. Ind. i. 304 = siamensis.
sublacunosa, Scheff. in Flora, liii. (1870) 247; et in Tijschr. Nederl. Ind. xxxi. (1870) 350 = eximia.
subpeltata, *Miq. Fl. Ind. Bat. Suppl.* 487.—Sumatra.
Talura, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 618.—Ind. or.
trinervia, Banks, ex Gaertn. f. Fruct. iii. 51 = Dipterocarpus turbinatus.
Tumbuggaia, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 617.—Ind. or.

SHORTIA, Torr. & Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 48; et Ser. II. xlv. (1868) 402. *DIAPENSIACEAE*, Benth. & Hook. f. ii. 620.
californica, Sisley, ex Carr. Rev. Hortic. (1883) 41, cum. fig. (Compos.).—Am. bor.
galacifolia, *Torr. & Gray, in Am. Journ. Sc.* xlii. (1842) 48.—Am. bor.
uniflora, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 225.—Japon.

SHULTZIA, Rafin. in Med. Repos. N. York, v. (1808) 356 = SCHULTZIA, Rafin. = **Obolaria**, Linn. (Gentian.).

SHURIA, Hort. ex Hérincq, L'Hortic. Franç. Sér. II. ii. (1861) 159 = **Achimenes**, P. Br. (Gesner.).
mexicana, Hort. ex Hérincq, l. c.—Mexic.

SHUTEREIA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 486 (Conv. Or. 103) = **Hewittia**, Wight (Convolv.).
bicolor, Choisy, ll. cc. = H. bicolor.

SHUTERIA, Wight & Arn. Prod. 207 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 529.
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 190.—Afr. trop.
densiflora, Benth. in Miq. Pl. Jungh. 232 = vestita.
ferruginea, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 182.—Reg. Himal.
glabrata, Wight & Arn. Prod. 207 = vestita.
hirsuta, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 182.—Reg. Himal.
involucrata, Wight & Arn. Prod. 207 = vestita.
rotundifolia, *Miq. Fl. Ind. Bat.* i. 1. 221.—Java.
suffulta, *Benth. in Miq. Pl. Jungh.* 232.—Burma.
trisperma, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 51.—Japon.
vestita, Benth. in Journ. Linn. Soc. viii. (1865) 267 = Glycine javanica.
vestita, *Wight & Arn. Prod.* 207.—Ind. or.

SHUTTELWORTHIA, Steud. Nom. ed. II. ii. 575 (1841), sphalm. = seq.

SHUTTLEWORTHIA, Meissn. Gen. 290 (1846) = **Verbena**, Linn.
Berterii, Meissn. Gen. Comm. 198 = V. Berterii.
Diceras, Meissn. l. c. = V. sulphurea.
dissecta, Walp. Rep. iv. 12 = V. dissecta.

SHUTTLEWORTHIA:—
pulchella, Meissn. Gen. Comm. 198 = Verbena tenera.
Ruiziana, Walp. Rep. iv. 12 = V. clavata.
Selloi, Walp. l. c. 13 = V. erinoides.
sulphurea, Meissn. Gen. Comm. 198 = V. sulphurea.
tenera, Walp. Rep. iv. 14 = V. tenera.

SIAGONANTHUS, Poepp. & Endl. Nov. Gen. et Sp. i. 40. t. 69 (1835) = **Ornithidium**, Salisb. (Orchid.).
multicaulis, Poepp. & Endl. l. c. = O. multicaule.

SIAGONARRHEN, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 146 (1858) = **Hyptis**, Jacq. (Labill.).
subicanus, Mart. ex J. A. Schmidt, l. c. = H. leucophylla.

SIALODES, Eckl. & Zeyh. Enum. 329 (1835) = **Galenia**, Linn. (Ficoid.).
glauca, Eckl. & Zeyh. l. c. = G. glauca.

SIBALDIA, Linn. Fl. Lapp. 77 (1737), sphalm. = SIBBALDIA, Linn. = **Potentilla**, Linn. (Rosac.).

SIBANGEA, Oliver, in Hook. Ic. Pl. xv. t. 1411 (1883). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 1223.
arborescens, *Oliver, l. c.*—Afr. trop.

SIBBALDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 177 = seq.

SIBBALDIA, Linn. Sp. Pl. ed. I. 284 (1753) = **Potentilla**, Linn. (Rosac.).
adpressa, Bunge, in Ledeb. Fl. Alt. i. 428.—Sibir. altaic.
altaica, Linn. f. Suppl. 191 = Chamaerhodos altaica.
californica, Spreng. Syst. iv. Cur. Post. 341 = P. multijuga.
congesta, D. Dietr. Syn. Pl. ii. 1020 = P. multijuga.
cuneata, Hornem. Ind. Sem. Hort. Hafn. (1842) et (1845).—Cf. Kunze, in Linnaea, xx. (1847) 59; Edgew. in Trans. Linn. Soc. xx. (1846) 44 = P. Sibbaldi.
erecta, Linn. Sp. Pl. 284 = Chamaerhodos erecta.
Fragariastrum, Turcz. ex Ledeb. Fl. Ross. ii. 33 = *adpressa*.
grandiflora, Pall. ex Schult. Syst. vi. 770 = Chamaerhodos grandiflora.
octopetala, Mill. Gard. Dict. ed. VIII. n. 2 = P. Sibbaldi.
parviflora, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 125. t. 5. f. 2 = P. Sibbaldi.
polygyna, Willd. ex Schult. Syst. vi. 770 = Chamaerhodos erecta.
potentilloides, Cambess. in Jacquem. Voy. Bot. 54 = P. micropetala.
procumbens, Linn. Sp. Pl. 284 = P. Sibbaldi.
purpurea, Royle, Illustr. Bot. Himal. 208. t. 40. f. 3 = P. purpurea.
sabulosa, Steud. Nom. ed. II. ii. 575 = Chamaerhodos sabulosa.
semiglabra, C. A. Mey. in Beitr. Pfl. Russ. Reich. vi. 44.—Reg. Caucas.
tetrandra, Bunge, Verz. Suppl. Fl. Alt. 25 = P. tetrandra.

SIBERTIA, Steud. Syn. Pl. Gram. in obs. (1855), sphalm. = LIBERTIA, Lej. = **Bromus**, Linn. (Gramin.).

SIBIRAEA, Maxim. in Act. Hort. Petrop. vi. (1879) 116, in obs. 213 = **Spiraea**, Linn. (Rosac.).
laevigata, Maxim. l. c. 215 = Spiraea laevigata.

SIBTHORPIA, Linn. Diss. Chen. 38 (1751); Amoen. Acad. iii. 21 (1756). *SCROPHULARINEAE*, Benth & Hook. f. ii. 959.
DISANDRA, Linn. f. Suppl. 32 (1781).
HORNEMANNIA, Benth. in DC. Prod. x. 428 (1846).
WILLICHIA, Mutis, ex Linn. Mant. ii. 553 (1771).
africana, *Linn. Sp. Pl.* 631.—Oriens.
europaea, *Linn. l. c.*—Europ. occ.
evolvulacea, Linn. f. Suppl. 288 = Dichondra repens.
nectarifera, *Wedd. Chlor. And.* ii. 111.—Bolivia.
parvifolia, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 25 = pichinchensis.
peregrina, *Linn. Sp. Pl.* 631.—Ins. Maurit.
pichinchensis, *H. B. & K. Nov. Gen. et Sp.* ii. 390. t. 175.—Mexic.; Andes.
pichinensis, Steud. Nom. ed. II. ii. 575, sphalm. = praec.

SIBTHORPIA:—
pinnata, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 288.—Nepal.
prostrata, Salisb. Prod. 94 = europaea.
retusa, H. B. & K. Nov. Gen. et Sp. ii. 391. t. 177 = pichinchensis.

SIBTORPIA, Scop. Introd. 178 (1777), err. typ. = praec.

SIBURATIA, Thou. Gen. Nov. Madag. 12 (1806) = **Maesa**, Forsk. (Myrsin.).

SICANA, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 180. t. 8. *CUCURBITACEAE*, Benth. & Hook. f. i. 829.
excisa, Griseb. Cat. Pl. Cub. 112 = Cayaponia excisa.
odorifera, *Naud. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 181.—Peruv.

SICELIUM, P. Br. Hist. Jamaic. 144 (1756) = **Coccocypselum**, P. Br. (Rubiac.).

SICKINGIA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 445. *RUBIACEAE*, Benth. & Hook. f. ii. 34.
ARARIBA, Mart. in Sitzb. Akad. Muench. (1860) 323.
cordifolia, *Hook. f. in Benth. & Hook. f. Gen.* ii. 34.—N. Granat.
Erythroxylon, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 446.—N. Granat.
longifolia, *Willd. l. c.* 447.—Venezuela.
orinocensis, Spreng Syst. i. 622 = Platycarpum orinocense.
pisoniaeformis, *Baill. Adansonia*, xii. (1879) 307.—Bras.

SICKLERA, M. Roem. Syn. Hesper. 34, 49 (1846) = **Murraya**, Linn. (Rutac.).
longifolia, M. Roem. l. c. 49 = M. longifolia.

SICKLERA, Sendtn. in Flora, xxix. (1846) 194 (err. typ. 178) = **Brachistus**, Miers (Solan.).
cuspidata, Dun. in DC. Prod. xiii. I. 502.—N. Granat.
solanacea, Sendtn. in Flora, xxix. (1846) 195 (err. typ. 179) = B. solanaceus.
tetrandra, A. Br. & Bouché, in Ind. Sem. Hort. Berol. (1853) App. 10.—Venezuela.

SICKMANNIA, Nees, in Linnaea, ix. (1834) 292 = **Ficinia**, Schrad. (Cyperac.).
radiata, Nees, l. c. = F. radiata.

SICYDIUM, Schlecht. in Linnaea, vii. (1832) 388; A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 194. *CUCURBITACEAE*, Benth. & Hook. f. i. 833.
TRICERATIA, A. Rich. Ess. Fl. Cub. 614. t. 44 (1845).
coriaceum, *Cogn. in DC. Monog. Phan.* iii. 906.—N. Granat.
diffusum, *Cogn. in Mart. Fl. Bras.* vi. IV. 112.—Bras.
gracile, *Cogn. l. c.* 113. t. 36.—Bras.
Lindheimeri, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 194 = Maximowiczia Lindheimeri.
monospermum, *Cogn. in Mart. Fl. Bras.* vi. IV. 114. t. 36. f. 1.—Bras.
Schiedeanum, *Schlecht. in Linnaea*, vii. (1832) 388.—Bras.
tamnifolium, Cogn. in DC. Monog. Phan. iii. 905 = Schiedeanum.
tenellum, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 167. t. 1 A = Maximowiczia tripartita.
tripartitum, Naud. l. c. 166 = Maximowiczia tripartita.

SICYOCARPUS, Boj. Hort. Maurit. 214 (1837) = **Marsdenia**, R. Br. (Asclep.).
verrucosus, Boj. l. c. = M. verrucosa.

SICYOIDES, Tourn. ex Medic. Phil. Bot. i. 36 (1789) = **Sicyos**, Linn. (Cucurb.).
angulata, Medic l. c. = Sicyos angulata.

SICYOMORPHA, Miers, in Trans. Linn. Soc. xxviii. (1872) 410. *CELASTRINEAE*.
nervosa, *Miers, l. c.* 412.—Peruv.
peruviana, *Miers, l. c.* 411.—Peruv.
pruinosa, Miers, l. c. (Hippocrateae sp.).—Panama.

SICYOS, Linn. Syst. ed. I (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 837.
BADEROA, Bert. ex Hook. Bot. Misc. iii. (1833) 324.
BRYONIASTRUM, Heist. Syst. 10 (1748).
SICYOIDES, Tourn. ex Medic. Phil. Bot. i. 36 (1789).
acuta, Rafin. Fl. Ludov. 113 = angulata.
Andreana, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 200.—Ecuador.
angulata, Berg. Desc. Pl. Cap. 352 = Kedrostis nana.
angulata, Forst. f. Prod. 368 = australis.
angulata, *Linn. Sp. Pl.* 1013.—Am. bor.
australis, *Endl. Prod. Fl. Norf.* 67.—Ins. Norfolk.
Baderoa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 324 = bryoniaefolia.
bogotensis, *Cogn. in DC. Monog. Phan.* iii. 874.—N. Granat.
Bourgaeana, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 22 = microphylla.
bryoniaefolia, *Moris, Enum. Sem. Hort. Taur.* (1831); *et in Mem. Acad. Torin.* xxxvii. (1834) 106. t. 6.—Chili.
cucumerina, *A. Gray, Bot. U. St. Expl. Exped.* i. 652. t. 82.—Ins. Sandvic.
depauperata, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 23 = parviflora.
Deppei, G. Don, Gen. Syst. iii. 34 = microphylla.
edulis, Jacq. Enum. Pl. Carib. 32 = Sechium edule.
eremocarpa, Peyr. in Linnaea, xxx. (1859–60) 56 = Echinocystis gemella.
eremocarpa, Schau. l. c. xx. (1847) 722 = Cyclanthera eremocarpa.
fretensis, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 473 = australis.
fusiformis, *Cogn. in Mart. Fl. Bras.* vi. IV. 108. t. 34.—Bras.
Galeottii, *Cogn. in DC. Monog. Phan.* iii. 883.—Mexic.
Garcini, Burm. f. Fl. Ind. 211 (err. typ. 311) = seq.?
Garcini, Linn. Mant. ii. 297 = Blastania Garcini.
glandulosa, Poir. Encyc. vii. 155 = Bowlesia oppositifolia.
glandulosa, Roem. Syn. Pepon. 105 = microphylla.
gracillima, *Cogn. in DC. Monog. Phan.* iii. 886.—Peruv.
Gunnii, Hook. f. ex Cogn. l. c. 875 = australis.
gymnacantha, Griseb. ex Cogn. l. c. 803 = Echinocystis torquata.
hederaefolia, Walp. Rep. ii. 204 = seq.
hederifolia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 450 = Gynostemma hederaefolium.
Helleri, Peyr. in Linnaea, xxx. (1859–60) 56 = Microsechium ruderale.
Kunthii, *Cogn. in DC. Monog. Phan.* iii. 888.—Peruv.
laciniata, Descourt. Fl. Antill. v. 103 = Sechium dule.
laciniata, *Linn. Sp. Pl.* i. 1013.—Ind. occ.
laevis, *A. Gray, Bot. U. St. Expl. Exped.* i. 648.—Peruv.
lobata, Michx. Fl. Bor. Am. ii. 217 = Echinocystis lobata.
longisetosa, *Cogn. in DC. Monog. Phan.* iii. 882.—Ecuador?
macrocarpa, *Cogn. l. c.* 893.—Ecuador.
macrophylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 651. t. 81.—Ins. Sandvic.
malvifolia, *Griseb. in Goett. Abh.* xix. (1874) 147.—Reg. Argent.
Martii, *Cogn. in Mart Fl. Bras.* vi. IV. 109.—Bras.
Maximowiczii, *Cogn. in DC. Monog. Phan.* iii. 895.—Ins. Sandvic.
microcarpa, *H. Mann, in Proc. Am. Acad.* vii. (1867) 167.—Ins. Sandvic.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 119.—Mexic.
montana, Griseb. in Goett. Abh. xix. (1874) 147 = Warmingii.
montana, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 53. t. 172.—Peruv.
muricata, Galeotti, ex Cogn. in DC. Monog. Phan. iii. 839 = Cyclanthera ribiflora.
oregana, Torr. & Gray, Fl. N. Am. i. 542 = Echinocystis oregana.
oxyantha, Wall. Cat. n. 6683 = Actinostemma tenerum.
pachycarpa, *Hook. & Arn. Bot. Beech. Voy.* 83.—Ins. Sandvic.
palmata, Moç. & Sesse, ex Cogn. in Mart. Fl. Bras. vi. IV. 111 = Microsechium palmatum.

SICYOS:—
palmatiloba, *Cogn. in DC. Monog. Phan.* iii. 891.—Ecuador.
parviflora, A. Gray, Pl. Wright. ii. 62 = laciniata.
parviflora, H. B. & K. Nov. Gen. et Sp. ii. 119 = Kunthii.
parviflora, *Willd. Sp. Pl.* iv. 626.—Ecuador.
parvifolia, A. Gray, ex Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 22 = laciniata.
peltata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 27 = Cocculus glaucescens.
pentandra, Wall. Cat. n. 6682 = Natsiatum herpeticum.
polyacanthos, *Cogn. in Mart. Fl. Bras.* vi. IV. 107.—Bras.
quinquelobata, *Cogn. l. c.* 109.—Bras.
Remyana, *Cogn. in DC. Monog. Phan.* iii. 898.—Ins. Sandvic.
scaberrima, Mart. & Gal. ex Peyr. in Linnaea, xxx. (1859–60) 57 = Microsechium ruderale.
Schimperi, Naud. in Schweinf. & Aschers. App. Beitr. Fl. Aethiop. 268 = australis.
subcorymbosa, *Cogn. in DC. Monog. Phan.* iii. 887.—Bolivia.
trifoliata, Linn. Sp. Pl. 1013 = Vitis acida.
triquetra, Moç. & Sesse, ex DC. Prod. iii. 309 = Sechiopsis triqueter.
villosa, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 223.—Ins. Galapag.
vitifolia, Hook. & Arn. Bot. Beech. Voy. 292 = microphylla.
vitifolia, *Willd. Sp. Pl.* iv. 626.—Mexic.
Warmingii, *Cogn. in Mart. Fl. Bras.* vi. IV. 108.—Bras.

SICYOSPERMA, A. Gray, Pl. Wright. ii. 62 (in Smithson. Contrib. v.) (1853). *CUCURBITACEAE*, Benth. & Hook. f. i. 837.
gracile, *A. Gray, l. c.*—N. Mexic.

SIDA, Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 203.
DIADESMA, Rafin. New Fl. Am. i. 41 (1836).
DICTYOCARPUS, Wight, in Madr. Journ. Sc. v. (1837) 309; et in Ann. Sc. Nat. Sér. II. xi. (1839) 169.
FLEISCHERIA, Steud. in Lehm. Pl. Preiss. i. 236 (1844).
LAMARKIA, Medic. Phil. Bot. i. 28 (1789).
MALVELLA, Jaub. & Spach, Illustr. v. 47. t. 444 (1855?).
MALVINDA, Dill. ex Medic. Malv. 23 (1787).
abscissa, Willd. ex Spreng. Syst. iii. 117 = serrata.
abutilifolia, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. austr.
abutilifolia, Moench, ex Steud. Nom. ed. I. 773 = Abutilon Avicennae.
abutiloides, Jacq. Obs. i. 17 t. 7 = Abutilon Jacquini.
abutiloides, Macfad. Fl. Jamaic. i. 86 = Abutilon indicum.
Abutilon, Linn. Sp. Pl. 685 = Abutilon Avicennae.
abyssinica, D. Dietr. Syn. Pl. iv. 859 (= Abutilonis sp.).—Abyss.
acaulis, Cav. Ic. v. 13. t. 422. f. 2 = Malvastrum Cavanillesii.
acerifolia, *Lag. Gen. et Sp. Nov.* 21.—Mexic.
acerifolia, Medic. Malv. 21 = Hibiscus Solandra.
acerifolia, Zuccagni, in Roem. Collect. 148 = Anoda hastata.
acrantha, *Link, Enum. Hort. Berol.* ii. 203.—Bras.
acuminata, R. Br. in Salt, Abyss. App. 65 = Abutilon longicuspe.
acuminata, *DC. Prod.* i. 462.—Ins. S. Doming.
acuta, *Burm. f. Fl. Ind.* 147.—Am. centr.
acutangula, Steud. Nom. ed. II. ii. 576 = Abutilon intermedium.
acutifolia, *Steud. l. c.*—Abyss.
adscendens, *A. St. Hil. Fl. Bras. Mer.* i. 182.—Bras.
affinis, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 285 = spinosa.
affinis, Spreng. Syst. iii. 121 = Abutilon rivulare.
africana, Beauv. Fl. Owar. ii. 87 = cordifolia.
aggregata, *Presl, Rel. Haenk.* ii. 106.—Reg. trop.
alata, *S. Wats, in Proc. Am. Acad.* xx. (1885) 356.—Mexic.
alba, Cav. Diss. i. 22 = rhombifolia.
alba, Linn. Sp. Pl. ed. II. 960 = spinosa.
albida, Willd. Enum. Hort. Berol. 722 = Abutilon indicum.
alcaeoides, Michx. Fl. Bor. Am. ii. 44 = Callirhoë alcaeoides.
alnifolia, Linn. Sp. Pl. 684 = spinosa.

SIDA:—
alpestris, A. St. Hil. Fl. Bras. Mer. i. 186 = paniculata.
altheaefolia, Sw. Prod. Veg. Ind. Occ. 101 = cordifolia.
ambigua, *Tenore, ex Steud. Nom.* ed. II. ii. 576.—Hab.?
americana, Linn. Sp. Pl. ed. II. 963 = Abutilon indicum.
amoena, *Desf. Cat. Hort. Par.* ed. III. 247.—Bras.
amoena, Wall. Cat. n. 1848 = Abutilon fruticosum.
amoena, *Weinm. in Syll. Ratisb.* i. (1824) 12.—China.
amplexicaulis, Lam. Encyc. i. 7 = Abutilon crispum.
amplexicaulis, Vell. Fl. Flum. 279; vii. t. 21 = Wissadula rostrata.
amplexifolia, [Moç. & Sesse,] ex DC. Prod. i. 469 = Abutilon crispum *vel* amplexifolium.
amplissima, Linn. Sp. Pl. 685 (Wissadulae sp.).—Am. trop.
angulata, *Vell. Fl. Flum.* 277; vii. t. 11.—Bras.
angulosa, Boj. Hort. Maurit. 34 = Abutilon intermedium.
angustifolia, Lam. Encyc. i. 4 = spinosa.
angustifolia, Medic. Beobacht. (1783) 193 (Quid?).—Hab.?
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3 = rhombifolia.
angustissima, Miq. Stirp. Surin. Sel. 102 = linifolia.
angustissima, *A. St. Hil. Fl. Bras. Mer.* i. 179.—Bras.
angusturensis, Willd. ex Spreng. Syst. iii. 104 = Hibiscus affinis.
anomala, *A. St. Hil. Fl. Bras. Mer.* i. 177. t. 33.—Am. trop.
anthemidifolia, Remy, in Ann. Sc. Nat. Sér. III. vi. (1846) 356 = Malvastrum anthemidifolium.
arabica, J. F. Gmel. Syst. 1049 = supina.
arborea, Linn. f. Suppl. 397; L'Hérit. Stirp. Nov. 131. t. 63 = Abutilon arboreum.
arborescens, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 60.—Bras.
Arbuscula, Zipp. ex Span. in Linnaea, xv. (1841) 171 = rhombifolia.
arguta, Fisch. ex Link, Enum. Hort. Berol. ii. 206 = verruculata.
arguta, *Presl, Rel. Haenk.* ii. 106.—Mexic.
arguta, Sw. Prod. Veg. Ind. Occ. 101 = ulmifolia.
aristata, Willd. ex Spreng. Syst. iii. 116 = cordifolia.
Arnottiana, *D. Dietr. Syn. Pl.* iv. 857.—Chili.
Arnottiana, Gill. ex Hook. Bot. Misc. iii. (1833) 154 = Abutilon crispum.
Arrabidae, Steud. Nom. ed. II. ii. 576 = truncata.
ascendens, Auct. = adscendens.
asiatica, Linn. Cent. Pl. ii. 26; Amoen. Acad. iv. 324 = Abutilon indicum.
asiatica, Thunb. Fl. Cap. ed. Schult. 548 = Abutilon Sonneratianum.
asiatica, Wall. Cat. n. 1852 D = Abutilon muticum.
atropurpurea, Blume, Bijdr. 77 = Abutilon auritum.
atropurpurea, Moris, in Mem. Acc. Torin. xxxvi. (1833) 182 (an praec.?).—Java.
astrosanguinea, Jacq. Ic. Pl. Rar. 1. t. 136 = paniculata.
aurantiaca, *A. St. Hil. Fl. Bras. Mer.* i. 185. t. 35.—Bras.
aurea, G. Don, Gen. Syst. i. 497 = Gaya aurea.
aurea, Lodd. Bot. Cab. t. 1842 (Abutilonis sp.).—Ind. or.
aurita, Hornem. ex Schrank, in Syll. Ratisb. ii. (1828) 73; Wall. Cat. n. 1860 = Abutilon auritum.
australis, Hort. Berol. ex Presl, Bot. Bemerk. 20 = Abutilon polyandrum.
Balbisiana, DC. Prod. i. 460 = acuta.
balica, Miq. Fl. Ind. Bat. i. II. 141 = veronicaefolia.
Bastardia, DC. Prod. i. 467 = Bastardia viscosa.
Bedfordiana, Hook. Bot. Mag. t. 3892 = Abutilon Bedfordianum.
begonioides, Griseb. in Bonplandia, vi. (1858) 3 = veronicaefolia.
bella, Steud. Nom. ed. II. ii. 576 = Abutilon elegans.
Beloëre, L'Hérit. Stirp. Nov. 130 = Abutilon indicum.
Berlandieri, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 197 = acuta?
Berteriana, Balb. ex DC. Prod. i. 460 = glomerata.
Berteriana, Spreng. Syst. iii. 110 = acuta.
Berthae, F. Muell. Fragm. v. 103 = Plagianthus Berthae.
betonicaefolia, Balb. ex DC. Prod. i. 463 = supina.
betonicaefolia, Pav. ex Hemsl. Biol. Centr. Am. Bot. i. 106 = spinosa.

SIDA :—

betulaefolia, Hort. ex Steud. Nom. ed. II. ii. 576 = Malvastrum tricuspidatum.
betulaefolia, *Schrank, in Syll. Ratisb.* ii. (1828) 74.—Bras.
betulina, *Hornem. Hort. Hafn.* ii. 646.—Am. austr.
betulina, Lag. ex Spreng. Syst. iii. 111 = acuta.
bibracteolata, D. Dietr. Syn. Pl. iv. 856 = Abutilon mexicanum.
bicallosa, Rafin. Fl. Ludov. 91 (Quid ?).—Am. bor.
bicolor, *Cav. Ic.* iv. 6. t. 311.—Mexic.
bicuspidata, J. F. Gmel. Syst. 1050 = spinosa.
bidentata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 68 = Abutilon bidentatum.
biflora, Cav. Diss. i. 37. t. 9. f. 1 = Abutilon ibarrense ?
bihamata, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 54.—Bras.
biloba, *Herb. Hook. ex Seem. Fl. Vit.* 15.—Polynes.
bivalvis, Cav. Diss. i. 13. t. 11. f. 2 = Bastardia bivalvis.
bonariensis, *Willd. ex Spreng. Syst.* iii. 120.—Reg. Argent.
borbonica, Cav. Diss. i. 9. t. 10. f. 2 = cordifolia.
Boriara, Wall. Cat. n. 1872 = spinosa.
borussica, Meyen, Reise, ii. 31 = Malvastrum borussicum.
brachyantha, D. Dietr. Syn. Pl. iv. 858 (Anodae sp.).—Am. austr.
brachypetala, DC. Prod. i. 460 = acuta.
brachystemon, [*Moç. & Sesse,*] *ex DC. Prod.* i. 459.—Mexic.
bracteolata, DC. Prod. i. 460 = glomerata *vel* Malvastrum tricuspidatum.
brasila, Schrank, ex Link, Enum. Hort. Berol. ii. 203 = acuta.
brasiliensis, *Cav. Diss.* i. 37. t. 34. f. 1.—Bras.
breviflora, Steud. ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 177 = urens.
brevipes, DC. Prod. i. 467 = Bastardia viscosa.
buettneriacea, Klotzsch, in Bot. Zeit. iv. (1846) 102, nomen = micrantha.
byssina, Schrank, in Syll. Ratisb. ii. (1828) 70 = cordifolia.
californica, Nutt. ex Torr. & Gray, Fl. N. Am. i. 233 = Sidalcea californica.
callifera, *Griseb. Cat. Pl. Cub.* 24.—Cuba.
calycina, Cav. Diss. i. 9. t. 8. f. 2 = veronicaefolia ?
calyptrata, Cav. l. c. 55 = Gaya calyptrata (in Addend.).
calyxhymenia, *J. Gay, in DC. Prod.* i. 462.—Austral.
campestris, *Benth. Pl. Hartw.* 113.—Ecuador.
campi, Vell. Fl. Flum. 278; vii. t. 19 = linifolia.
canariensis, Willd. Sp. Pl. iii. 755 = rhombifolia.
candicans, DC. Prod. i. 466 = Gaya calyptrata (in Addend.).
canescens, Cav. Diss. i. 23 = rhombifolia.
capensis, Cav. l. c. t. 12. f. 3.; ii. 49 = acuta ?
capensis, DC. Prod. i. 461 = rhombifolia.
capensis, *Eckl. & Zeyh. Enum.* 40; *Harv. & Sond. Fl. Cap.* i. 167.—Afr. austr.
capillaris, *Cav. Diss.* i. 10 = paniculata.
capitata, Linn. Sp. Pl. 685 = Malachra capitata.
cardanisea, Rafin. Fl. Ludov. 90 (Quid ?).—Am. bor.
cardiophylla, *F. Muell. Fragm.* viii. 242.—Austral.
carnea, D. Dietr. Syn. Pl. iv. 853 = Abutilon carneum.
carnea, [*Moç. & Sesse,*] *ex DC. Prod.* i. 473 (Kosteletzkyae sp. ?).—Mexic.
carpinifolia, Bourg. ex Griseb. Fl. Brit. W. Ind. 74 = rhombifolia.
carpinifolia, Linn. f. Suppl. 307 = acuta.
carpinoides, DC. Prod. i. 461 = Malvastrum tricuspidatum.
Castelneana, Griseb. in Goett. Abh. xxiv. (1879) 44 (Malvastri sp.).—Reg. Argent.
caudata, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 52.—Bras.
centrota, Spreng. Syst. iv. Cur. Post. 259 = Anoda hastata.
ceratocarpa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 154 = Abutilon ceratocarpum.
chaetodonta, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 199 = veronicaefolia.
chilensis, Spreng. Syst. iii. 109 = Cristaria betonicaefolia.
chinensis, Retz. Obs. iv. 29 = rhombifolia.
chlorophis, Wall. ex Steud. Nom. ed. II. ii. 577 (Quid ?).—Ind. or.
ciliaris, *Linn. Syst.* ed. X. 1145.—Am. bor.; Ind. occ.
ciliosa, Boj. ex Baker, Fl. Maurit. 19 = cordifolia.
circinata, *Willd. ex Spreng. Syst.* iii. 119.—Bras.
cistiflora, Blume, Bijdr. 77 = Abutilon Sonneratianum ?

SIDA :—

cistiflora, L'Hérit. Stirp. Nov. 127. t. 61 = Abutilon mollissimum.
cistoides, Hort. ex Steud. Nom. ed. II. ii. 577 = Abutilon mollissimum.
clandestina, Phil. Fl. Atac. 12 (Malvastri sp.).—Chili.
cleisocalyx, *F. Muell. Fragm.* x. 73.—Austral.
coccinea, DC. Prod. i. 465 = Malvastrum coccineum.
Collaei, D. Dietr. Syn. Pl. iv. 853 = Abutilon elegans.
collina, *Schlecht. in Linnaea*, xi. (1834) 364.—Mexic.
compacta, C. Gay, Fl. Chil. i. 329 = Malvastrum compactum.
compacta, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 54.—Reg. Argent.
compressa, DC. Prod. i. 462; Wall. Cat. n. 1866 = rhombifolia.
concinna, *Phil. Fl. Atac.* 12.—Chili.
conferta, Link, Enum. Hort. Berol. ii. 207 = cordifolia.
conferta, Salzm. ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 177 = urens.
confertiflora, D. Dietr. Syn. Pl. iv. 856 = Abutilon confertiflorum.
congensis, D. Dietr. l. c. 859 = urens.
contracta, Link, Enum. Hort. Berol. ii. 204 = Wissadula rostrata.
corchorifolia, *Tenore, Ind. Sem. Hort. Neap.* (1857).—Hab. ?
cordifolia, Forsk. Fl. Aegypt. Arab. 124 = Abutilon indicum.
cordifolia, Griseb. in Goett. Abh. xxiv. (1879) 44 = potentilloides.
cordifolia, *Linn. Sp. Pl.* 684.—Reg. trop.
cordilleriana, D. Dietr. Syn. Pl. iv. 857 = Cristaria Haenkeana.
cornuta, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 724.—Am. austr.
coromandelina, Hort. ex Steud. Nom. ed. II. ii. 577 = Malvastrum tricuspidatum.
coronata, Scop. Delic. Insubr. iii. 1 = Abutilon indicum.
corrugata, *Lindl. in Mitch. Three Exped.* ii. 13.—Austral.
corylifolia, *Wall. Cat.* n. 1865.—Burma.
corynocarpa, Wall. Cat. n. 1870 = rhombifolia.
costata, *Schlecht. in Linnaea*, xi. (1834) 365.—Mexic.
crassifolia, L'Hérit. Stirp. Nov. 125. t. 60 = Abutilon Jacquini.
crenata, *G. Don, Gen. Syst.* i. 492.—Afr. trop.
crenatiflora, Pers. Syn. ii. 247 = Anoda parviflora ?
crenatiflora, Willd. ex Schlecht. in Linnaea, xi. (1837) 217 = Anoda crenatiflora.
crispa, Linn. Sp. Pl. 685 = Abutilon crispum.
crispifolia, Cav. Ic. v. 11. t. 419 (Sphaeralceae sp.).—Am. austr.
cristata, Linn. Sp. Pl. 685 = Anoda lavateroides.
cryphiopetala, *F. Muell. Fragm.* iii. 4.—Austral.
cryptopetala, F. Muell. l. c. ii. 11 = Abutilon cryptopetalum.
cubensis, D. Dietr. Syn. Pl. iv. 851 = Gaya occidentalis (in Addend.).
cuneifolia, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 165.—Texas.
cuneifolia, Roxb. Fl. Ind. iii. 170 = Schimperiana.
cuneifolia, Roxb. Hort. Beng. 50 = Melochia corchorifolia.
debilis, G. Don, Gen. Syst. i. 499 = urens.
decagyna, Schum. & Thonn. Beskr. Guin. Pl. 307 = cordifolia.
decipiens, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 54 = Malvastrum campanulatum.
decumbens, *A. St. Hil. & Naud. l. c.* 51.—Bras.
deflexa, *Cav. in Anal. Cienc. Nat.* vi. (1803) 337 (sp. dub.).—Cuba.
delphinifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 235 = Sidalcea malvaeflora.
deltoidea, Hornem. Enum. Pl. Hort. Hafn. (1807) 36 = Anoda triangularis *vel* hastata.
densiflora, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 155.—Am. austr.
densiflora, A. Rich. Tent. Fl. Abyss. i. 66 = urens.
denticulata, Fresen. in Mus. Senckenb. i. (1833) 182 = Abutilon fruticosum.
denudata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 100 (Anodae sp.).—Bras.
depauperata, Hook. f. in Trans. Linn. Soc. xx. (1847) 232 = Abutilon depauperatum.
depressa, Benth. Bot. Voy. Sulph. 69 = Malvastrum scoparium.
diantha, D. Dietr. Syn. Pl. iv. 856 = Abutilon dianthum.

SIDA :—

dictyocarpa, F. Muell. ex Benth. in Journ. Linn. Soc. vi. (1862) 103 (Plagianthi sp.).—Austral.
Diellii, A. Gray, Bot. U. St. Expl. Exped. i. 162 = rotundifolia.
diffusa, *H. B. & K. Nov. Gen. et Sp.* v. 257.—Reg. trop.
digitata, Spreng. Syst. iv. Cur. Post. 259 = Callirhoë digitata.
Dilleniana, Willd. Sp. Pl. iii. 764 = Anoda lavateroides.
dioica, Cav. Diss. v. 278. t. 132 = Napaea dioica.
diploscypha, Torr. & Gray, Fl. N. Am. i. 234 = Sidalcea diploscypha.
discissa, Bertol. in Mem. Soc. Ital. Moden. xxiii. (1844) 305. t. 5 = Abutilon discissum.
discolor, Hook. Journ. Bot. i. (1834) 250 = Plagianthus sidoides.
dissecta, Nutt. ex Torr. & Gray, Fl. N. Am. i. 235 = Malvastrum coccineum.
dissecta, Steud. Nom. ed. II. ii. 577 = Cristaria dissecta.
disticha, Cav. Ic. v. 12. t. 432 = Gaya calyptrata (in Addend.).
divergens, Benth. Bot. Voy. Sulph. 69 = Wissadula divergens.
diversifolia, Spreng. Syst. iii. 116 = Hibiscus Solandra.
Dombeyana, DC. Prod. i. 463 = veronicaefolia.
Doniana, D. Dietr. Syn. Pl. iv. 857 = Abutilon indicum.
dubia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 50.—Bras.
dumosa, Backhouse, ex Walp. Ann. ii. 158 = Abutilon Fraseri.
dumosa, Sw. Prod. Veg. Ind. Occ. 101 = pyramidata.
echinata, *Willd. ex Spreng. Syst.* iii. 120.—Ecuador.
echinocarpa, *F. Muell. Fragm.* xi. 62.—Austral.
ecornis, Vell. Fl. Flum. 278; vii. t. 16 = Abutilon pauciflorum.
elata, Macfad. Fl. Jamaic. i. 87 = Abutilon giganteum.
elegans, Colla, in Mem. Acc. Torin. xxxv. (1831) 155 = Abutilon elegans.
elegans, D. Dietr. Syn. Pl. iv. 852 = bella.
Elliotti, *Torr. & Gray, Fl. N. Am.* i. 231.—Am. bor.
elliptica, Steud. Nom. ed. II. ii. 577 = Abutilon ellipticum.
elongata, Blume, Bijdr. 76 = veronicaefolia.
emarginata, Willd. Sp. Pl. iii. 757 = ulmifolia.
Endlicheriana, Presl, Rel. Haenk. ii. 111 = glutinosa.
erecta, Macfad. Fl. Jamaic. i. 80 = spinosa.
eriantha, Steud. Nom. ed. II. ii. 577 = Cristaria eriantha.
erosa, Link, Enum. Hort. Berol. ii. 203 = ciliaris *vel* rhombifolia.
erubescens, *Schrank, ex Steud. Nom.* ed. II. ii. 577 (Quid ?).—Am. trop. ?
esculenta, Steud. l. c. = Abutilon esculentum.
Eteromischos, Cav. Diss. ii. 55 = Abutilon indicum.
excelsior, Cav. l. c. i. 27. t. 5. f. 3 = Wissadula rostrata.
exstipularis, Cav. l. c. ii. 56. t. 14. f. 2 = Abutilon stipulare.
fallax, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 306 = rotundifolia.
fasciculata, Torr. & Gray, Fl. N. Am. i. 231 = anomala.
fasciculata, Willd. ex Spreng. Syst. iii. 113 = glutinosa.
fasciculiflora, Miq. Fl. Ind. Bat. i. II. 140 = glutinosa.
ferruginea, DC. Prod. i. 468 (Wissadulae sp.).—Peruv.
fibulifera, Lindl. in Mitch. Three Exped. ii. 45 = corrugata.
filicaulis, Torr. & Gray, Fl. N. Am. i. 232 = diffusa.
filiformis, A. Cunn. in Mitch. Journ. Trop. Austral. 361 = corrugata.
filiformis, Jacq. Obs. Bot. ii. 23 = Abutilon crispum.
filiformis, Moric. Pl. Am. Rar. i. 10. t. 8; ex Bull. Férussac, xxiii. (1830) 79 = diffusa.
filipes, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 164.—Texas.
flavescens, *Cav. Diss.* i. 14.—Bras.
floribunda, H. B. & K. Nov. Gen. et Sp. v. 258 = paniculata.
fluminensis, Steud. Nom. ed. II. ii. 577 = *pilosa*, Vell.
fluviatilis, Vell. Fl. Flum. 278; vii. t. 18 (Abutilonis sp.).—Bras.
foetida, Cav. Diss. vi. 349 = Bastardia viscosa.
foliosa, Splitg. ex De Vriese, in Nederl. Kruidk. Arch. i. (1848) 340 = acuta ?
Forsteri, Montr. in Mém. Acad. Lyon, x. (1860) 182 = rhombifolia.

SIDA:—
fragrans, L'Hérit. Stirp. Nov. 111. t. 53 (Bastardiae sp.).—Ins. S. Doming.
Fraseri, Hook. in Mitch. Journ. Trop. Austr. 368 = Abutilon Fraseri.
frutescens, Cav. Diss. i. 12. t. 10. f. 1 = acuta.
frutescens, DC. Prod. i. 461 = rhombifolia.
fruticosa, Mill. Gard. Dict. ed. VIII. n. 18 = Pavonia Typhalaea.
fulva, A. St. Hil. Fl. Bras. Mer. i. 176 = ciliaris.
Garckeana, Polak. in Linnaea, xli. (1877) 551 = acuta.
Gaudichaudiana, G. Don, Gen. Syst. i. 497 = Gaya Gaudichaudiana.
Gaya, DC. Prod. i. 466 = Gaya hermannioides.
geminiflora, DC. l. c. 470 = Abutilon ibarrense.
geniculata, *Roxb. Hort. Beng.* 50, nomen.—' Am.'
geraniifolia, D. Dietr. Syn. Pl. iv. 857 = Cristaria geraniifolia.
geranioides, DC. Prod. i. 474 = Abutilon geranioides.
Ghisbreghtiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 200 = Lindeniana.
gigantea, Jacq. Hort. Schoenb. ii. 8. t. 141 = Abutilon giganteum.
glabra, Mill. Gard. Dict. ed. VIII. n. 14 = ulmifolia.
glabra, Nutt. in Journ. Acad. Philad. vii. (1834) 90 = acuta.
glandulifera, Benth. Bot. Voy. Sulph. 69 = pyramidata.
glandulosa, Roxb. ex Wight & Arn. Prod. 58 = spinosa.
glauca, Cav. Ic. i. 8. t. 11 = Abutilon muticum.
glaucophylla, Spreng. Syst. iii. 109 = Cristaria glaucophylla.
glechomaefolia, D. Dietr. Syn. Pl. iv. 852 = Abutilon glechomaefolium.
globiflora, Hook. Bot. Mag. t. 2821 (Abutilonis sp.).—Ins. Maurit.
glochidiata, *Regel, in Act. Hort. Petrop.* iv. (1876) 338.—As. temp.
glomerata, *Cav. Diss.* i. 18. t. 2. f. 6.—Am. trop.
glutinosa, *Cav. l. c.* 16. t. 2. f. 8; *Roxb. Hort. Beng.* [97]; *Fl. Ind.* iii. 172.—Am. et As. trop.; Ins. Mascar.
gracilis, R. Br. in Salt, Abyss. App. 65 = Abutilon fruticosum.
gracilis, Ell. Sketch, ii. 159 = Elliottii.
gracilis, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Ind. occ.
gracilis, Salisb. Prod. 382 = Abutilon crispum.
graminifolia, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
grandiflora, Poir. Encyc. Suppl. i. 31 = Abutilon arboreum.
grandiflora, Walp. Ann. i. 103 = Malva grandiflora.
grandifolia, Willd. Enum. Hort. Berol. 724 = Abutilon mollissimum.
graveolens, Roxb. Hort. Beng. 50; Fl. Ind. iii. 179 = Abutilon graveolens.
Greveana, Baill. in Bull. Soc. Linn. Par. i. (1885) 504 = Abutilon intermedium.
Grevilliana, Gill. ex Hook. Bot. Misc. iii. (1833) 154 = Abutilon Grevillianum.
grewioides, *Guill. & Perr. Fl. Seneg. Tent.* i. 71.—Ind. bor. occ.; Afr. trop.
grossulariaefolia, Hook. & Arn. Bot. Beech. Voy. 326 = Malvastrum coccineum.
Guichenotiana, Steud. Nom. ed. II. ii. 577 = Abutilon Guichenotianum.
Guilleminiana, Steud. l. c. = Abutilon indicum?
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 307 = Abutilon asiaticum?
Haenkeana, Presl, Rel. Haenk. ii. 104 = rhombifolia.
hamulosa, Salzm. ex Griseb. Fl. Brit. W. Ind. 76 = cordifolia.
hastata, *A. St. Hil. Fl. Bras. Mer.* i. 190. t. 34. f. 2.—Bras.
hastata, Sims, Bot. Mag. t. 1541 = acerifolia.
hastata, Willd. Sp. Pl. iii. 763 = Anoda hastata.
hederacea, Torr. ex A. Gray, in Mem. Am. Acad. iv. (1849) 23 = obliqua.
hederaefolia, Cav. Diss. i. 8. t. 9. f. 3 = veronicaefolia.
herbacea, Cav. l. c. 19. t. 13. f. 1 = cordifolia.
hermanniaefolia, *Willd. ex Spreng. Syst.* iii. 117.—Mexic.
hermannioides, H. B. & K. Nov. Gen. et Sp. v. 258 = jamaicensis.
hernandioides, L'Hérit. Stirp. Nov. 121. t. 58 = Wissadula rostrata.

SIDA:—
heterocarpa, Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 163 = spinosa.
heterophylla, Cav. Ic. v. 12. t. 421 = Cristaria heterophylla.
heterophylla, Klein, ex Spreng. Syst. iii. 116 = Hibiscus Solandra.
heterosperma, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 182 = Wissadula rostrata.
heterosperma, *Vis. in Linnaea*, xvi. (1842) *Litt.* 113.—Afr. bor.
heterotricha, Zipp. ex Span. in Linnaea, xv. (1841) 171 = Abutilon timoriense.
hibisciformis, *Bertol. Fl. Guatim.* 28.—Guatemala.
Hilaireana, Steud. Nom. ed. II. ii. 577 = Abutilon carneum.
Hilariana, Presl, Rel. Haenk. ii. 107 = pyramidata.
Hilariana, Walp. Rep. i. 317 = prostrata *vel* flavescens.
hirsuta, *Mill. Gard. Dict.* ed. VIII. n. 9.—Hab.?
hirsuta, Vell. Fl. Flum. 279; vii. t. 20 = *megalantha*.
hirsutissima, D. Dietr. Syn. Pl. iv. 850 (Bastardiae sp.).—Mexic.
hirsutissima, Mill. Gard. Dict. ed. VIII. n. 11 = glomerata.
hirta, Lam. Encyc. i. 7 = Abutilon hirtum.
hirta, Wall. Cat. n. 1852 B, 1858 G = Abutilon graveolens.
hirta, Wall. Cat. n. 1855 = glutinosa.
hispida, Bert. ex DC. Prod. i. 463 = supina.
hispida, Pursh, Fl. Am. Sept. ii. 452 = Malvastrum angustum.
Hoheri, Hook. Ic. Pl. sub t. 565-566 = Hoheria populnea.
holosericea, Willd. ex Spreng. Syst. iii. 112 = cordifolia.
hondensis, H. B. & K. Nov. Gen. et Sp. v. 260 = rhombifolia.
Hookeri, D. Dietr. Syn. Pl. iv. 856 = Abutilon albidum.
Hookeri, *Steud. Nom.* ed. II. ii. 578.—Chili.
Hookeriana, *Miq. in Lehm. Pl. Preiss.* i. 242.—Austral.
Hulseana, Torr. & Gray, Fl. N. Am. i. 233 = Abutilon Hulseanum.
Humboldtii, D. Dietr. Syn. Pl. iv. 848 = paniculata.
humilis, Cav. Diss. v. 277 = veronicaefolia.
humillima, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 12 = corrugata.
hyssopifolia, Presl, Rel. Haenk. ii. 109 = spinosa.
ibarrensis, DC. Prod. i. 470 = Abutilon ibarrense.
imberbis, DC. l. c. 469 = Abutilon crispum.
inaequalis, Link & Otto, Ic. Pl. Sel. 75. t. 34 (Abutilonis sp.).—Bras.
incana, Link, Enum. Hort. Berol. ii. 204 = Abutilon incanum.
incarnata, Spreng. Syst. iii. 115 = Anoda punicea.
incerta, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 51 = rhombifolia?
inclusa, *Benth. Fl. Austral.* i. 197.—Austral.
indica, Cav. Diss. i. 33. t. 7. f. 10 = Abutilon crispum.
indica, Linn. Cent. Pl. ii. 26; Amoen. Acad. iv. 323 = Abutilon indicum.
integerrima, Hook. Bot. Mag. t. 4360 = Abutilon integerrimum.
integrifolia, Montr. in Mém. Acad. Lyon, x. (1860) 183 (Abutilonis sp.).—N. Caled.
intermedia, A. St. Hil. Fl. Bras. Mer. i. 188. t. 34. f. 1 = macrodon.
interrupta, *Balb. ex DC. Prod.* i. 464.—Am. trop.
interstans, F. Muell. in Linnaea, xxv. (1852) 383 = corrugata.
intricata, *F. Muell. ex Trans. Phil. Soc. Vict.* i. (1855) 12.—Austral.
involucrata, A. Rich. Ess. Fl. Cub. 162 = anomala.
Jacquini, D. Dietr. Syn. Pl. iv. 855 = Abutilon Jacquini.
jamaicensis, *Linn. Syst.* ed. X. 1145.—Am. trop.
jamaicensis, Mill. Gard. Dict. ed. VIII. n. 5 = Malvastrum tricuspidatum.
jamaicensis, Vell. Fl. Flum. vii. t. 10 = acuta.
jatrophoides, *L'Hérit. Stirp. Nov.* 117. t. 56.—Peruv.
javensis, Cav. Diss. i. 10 = veronicaefolia.
Julianae, Steud. Nom. ed. II. ii. 578 = Abutilon Julianae.
juncea, *Banks & Soland. ex Steud. Nom.* ed. I. 774; ed. II. ii. 578.—Hab.?
Jussiaeana, DC. Prod. i. 463 = veronicaefolia.

SIDA:—
Keerlena, Steud. Nom. ed. II. ii. 578 = Abutilon floribundum.
Kohautiana, Walp. Rep. i. 320 = rhombifolia.
Kotschyi, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 188 = Abutilon fruticosum.
Kunthiana, *Presl, Rel. Haenk.* ii. 106.—Mexic.
lanceolata, Retz. Obs. iv. 28 = acuta.
lasiostega, Link, Enum. Hort. Berol. ii. 20 = Abutilon crispum.
Lawrencia, F. Muell. Vict. Pl. i. 162 = Plagianthus spicatus.
laxiflora, Steud. Nom. ed. II. ii. 578 = Wissadula rostrata.
leiophloia, Miq. in Lehm. Pl. Preiss. i. 241 = Hookeriana.
leiophylla, *Spreng. Syst.* iii. 121.—Porto Rico.
lepida, *F. Muell. Fragm.* vi. 168.—Austral.
lepidota, *A. Gray, Pl. Wright.* i. 18; ii. 21.—N. Mexic.
Leschenaultiana, DC. Prod. i. 468 = Wissadula rostrata.
leucanthema, D. Dietr. Syn. Pl. iv. 852 = Abutilon leucanthemum.
leucopetala, F. Muell. Fragm. ii. 12 = Abutilon leucopetalum.
lignosa, Cav. Diss. i. 28. t. 6. f. 2 = Abutilon Jacquini.
Lindeniana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 200.—Mexic.
Lindheimeri, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 213.—Texas.
linearifolia, *A. St. Hil. Fl. Bras. Mer.* i. 180.—Bras.
linearifolia, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 303 = linifolia.
linearis, Cav. Ic. iv. 6. t. 312. f. 1 = spinosa.
lineata, Vell. Fl. Flum. 280; vii. t. 25 (Abutilonis sp.).—Bras.
linifolia, *Cav. Diss.* i. 14. t. 2. f. 1.—Am. et Afr. trop.
Lonchitis, A. St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 50 = rhombifolia.
longicuspis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 69 = Abutilon longicuspe.
longipes, *A. Gray, Pl. Wright.* i. 19; ii. 21.—N. Mexic.
longipes, Harv. & Sond. Fl. Cap. i. 167 = rhombifolia.
longipes, E. Mey. in Drège, Zwei Pfl. Docum. 222 = capensis.
luciana, DC. Prod. i. 468 = Wissadula rostrata.
Luschnathiana, *Steud. Nom.* ed. II. ii. 578.—Bras.
macrantha, Steud. l. c. = Abutilon macranthum.
macrodon, *DC. Prod.* i. 464.—Bras.
macrophylla, Hilsenb. & Boj. ex Baill. in Bull. Soc. Linn. Par. i. (1885) 504 = Abutilon intermedium.
macrophylla, Wall. Cat. n. 1851 D = Abutilon polyandrum.
macropoda, *F. Muell. ex Benth. Fl. Austral.* i. 193.—Austral.
macropodon, Steud. Nom. ed. II. ii. 578 = Abutilon macropodum.
macrorhiza, James, in Long's Exped. ii. 121 = Callirhoë alcaeoides.
macrostipula, Zipp. ex Span. in Linnaea, xv. (1841) 171 = Abutilon Guichenotianum.
maculata, Bert. ex DC. Prod. i. 463 = acuminata.
maculata, Cav. Diss. i. 20. t. 3. f. 7 = cordifolia.
maderensis, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 35 = rhombifolia.
Magdalenae, DC. Prod. i. 467 = Bastardia viscosa.
malacophylla, Link & Otto, Ic. Pl. Sel. 67. t. 30 = Pavonia velutina.
malvacea, Vell. Fl. Flum. 277; vii. t. 13 = Pavonia sepium.
malvaeflora, Hook. & Arn. Bot. Beech. Voy. 326 = Sidalcea malvaeflora.
malvaeflora, Lindl. in Bot. Reg. t. 1036 = Sidalcea campestris.
malvaeflora, [Moç. & Sesse,] ex DC. Prod. i. 474 = Sidalcea malvaeflora.
Malvaviscus, [Moç. & Sesse,] ex DC. l. c. 459 = Anoda punicea.
maritiana, Miq. in Hoev. & De Vriese, Tijdschr. x. (1843) 79 = Plagianthus spicatus.
Martiana, A. St. Hil. Fl. Bras. Mer. i. 187 = aurantiaca.
Mathewsii, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 565 = veronicaefolia *vel* Melochia pyramidata.
maura, *Endl. Fl. Suedseeinseln*, 182.—Ins. Marian.
mauritiana, Jacq. Misc. ii. 352 = Abutilon Avicennae.
megalantha, Steud. Nom. ed. II. ii. 578 (Abutilonis sp.).—Bras.
megalorrhiza, Phil. Fl. Atac. 11 (Malvastri sp.).—Chili.

SIDA :—

megapotamica, Spreng. f. Tent. Suppl. Syst. 19 = Abutilon megapotamicum.
Mendanha, Vell. Fl. Flum. 279; vii. t. 23 (Abutilonis sp. ?).—Bras.
meridionalis, Salisb. Prod. 382 = Abutilon indicum.
mexicana, *Scop. Delic. Insubr.* i. 22.—Mexic.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 307.—Ins. Sandvic.
micans, Cav. Diss. i. 19. t. 3. f. 1 = cordifolia.
micrantha, *A. St. Hil. Fl. Bras. Mer.* i. 190.—Bras.
micrantha, *Schrank, in Syll. Ratisb.* ii. (1828) 72 (sp. diversa).—Bras.
microphylla, Cav. Diss. i. 22. t. 12. f. 2 = rhombifolia.
microphylla, *DC. Prod.* i. 461.—Ind. or.; Polynes.
microsperma, Cav. Diss. i. 29. t. 13. f. 4 = Abutilon microspermum.
Milleri, DC. Prod. i. 472 = spinosa *vel* rhombifolia.
minor, Macfad. Fl. Jamaic. i. 79 = spinosa.
Miqueliana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 259.—Bras.
Mociniana, D. Dietr. Syn. Pl. iv. 858 (Kosteletzkyae sp.).—Mexic.
modesta, Phil. Fl. Atac. 12 (Malvastri sp.).—Chili.
mollicoma, Willd. Enum. Hort. Berol. 725 = Abutilon mollicomum.
mollis, Herb. Banks, ex Griseb. Fl. Brit. W. Ind. 76 = cordifolia.
mollis, Orteg. Hort. Matr. Dec. 65 = Abutilon molle.
mollis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 111 = glomerata.
mollissima, Cav. Diss. ii. 49. t. 14. f. 1 = Abutilon mollissimum.
montana, D. Dietr. Syn. Pl. iv. 854 = Abutilon montanum.
montana, Roxb. ex Wight & Arn. Prod. 56 = Abutilon crispum.
morifolia, Cav. Diss. i. 9. t. 1. f. 1 = veronicaefolia.
mucronulata, DC. Prod. i. 461 = Malvastrum tricuspidatum.
multicaulis, Cav. Diss. i. 10. t. 1. f. 6 = veronicaefolia.
multifida, Cav. l. c. i. 25 = Cristaria multifida.
multiflora, Cav. l. c. 18. t. 3. f. 3 = cordifolia.
multiflora, Jacq. Obs. Bot. ii. 23 = paniculata.
multiflora, Pers. Syn. ii. 243 = Wissadula nudiflora.
muricata, Cav. Ic. vi. 78. t. 597. f. 2 = ciliaris.
mutica, Delile, Illustr. Fl. Egypt. 60. n. 45 = Abutilon muticum.
myriantha, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 179.—N. Granat.
mysorensis, Herb. Madr. ex Wight & Arn. Prod. 59 = glutinosa.
Napaea, Cav. Diss. v. 277. t. 132. f. 1 = Napaea dioica.
neelgerrensis, Steud. Nom. ed. II. ii. 578 = Abutilon neilgherrense.
neglecta, *Passerini, Select. Sem. Hort. Parmen.* (1847). —*Cf. Linnaea*, xxiv. (1851) 180.—Hab. ?
nematopoda, F. Muell. in Linnaea, xxv. (1852) 382 = corrugata.
nemoralis, G. Don, Gen. Syst. i. 497 = Abutilon crispum.
nervosa, DC. Prod. i. 465 = glutinosa.
nervosa, Wall. Cat. n. 1853 = veronicaefolia.
nubigena, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 307 (Malvastri sp.).—Peruv.; Bolivia.
nudiflora, L'Hérit. Stirp. Nov. 123. t. 59 = Wissadula nudiflora.
nutans, L'Hérit. l. c. 119bis, t. 57 = Gaya calyptrata (in Addend.).
obliqua, [Moç. & Sesse,] ex DC. Prod. i. 462 = bicolor.
obliqua, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 233.—Am. bor.
obovata, Wall. Cat. n. 1864 = rhombifolia.
obtusa, Cav. Diss. i. 34. t. 9. f. 2 = Abutilon umbellatum.
obtusa, A. Rich. Ess. Fl. Cub. 161 = acuta.
occidentalis, Linn. Cent. Pl. ii. 26; Amoen. Acad. iv. 324 = Gaya occidentalis (in Addend.).
oculata, *Lowe, Fl. Mader.* 592.—Ins. Canar.
olens, Buch.-Ham. ex Wall. Cat. n. 1874 = glutinosa.
oligophylla, D. Dietr. Syn. Pl. iv. 858 (Hibisci sp.).—Ins. Timor.
olygantha, D. Dietr. l. c. 854 (Abutilonis sp. ?).—Bras.
opaca, Sieber, ex Presl, Bot. Bemerk. 19 = Pavonia hastata.

SIDA :—

orbiculata, DC. Prod. i. 471 = Abutilon indicum.
oregana, Nutt. ex Torr. & Gray, Fl. N. Am. i. 234 = Sidalcea malvaeflora.
orientalis, Cav. Diss. i. 21. t. 12. f. 1 = rhombifolia.
orientalis, DC. Prod. i. 461 = acuta.
Ortegae, Steud. Nom. ed. II. ii. 578 = Anoda crenatiflora.
ostryaefolia, Webb, Fragm. Fl. Aethiop. 49 = rhombifolia.
ovalis, *Kostel. Allg. Med.-Pharm. Fl.* v. 1867.—Peruv.
ovata, Cav. Diss. vi. 350 = supina.
ovata, G. Don, Gen. Syst. i. 492 = acuta.
ovata, *Forsk. Fl. Aegypt. Arab.* 124.—Arab.
oxycarpa, F. Muell. Fragm. ii. 12 = Abutilon oxycarpum.
oxyphylla, [Moç. & Sesse,] ex DC. Prod. i. 465 (Abutilonis sp.).—Mexic.
oxyphylla, Wall. Cat. n. 1850 = Abutilon polyandrum.
paeoniaeflora, Hook. Bot. Mag. t. 4170 = Abutilon paeoniaeflorum.
palmata, *Cav. Diss.* i. 20.—Peruv.
palmata, Jacq. Ic. Pl. Rar. iii. t. 547 = jatrophoides.
palmata, [Moç. & Sesse,] ex DC. Prod. i. 472 = acerifolia.
paniculata, *Linn. Syst.* ed. X. 1145.—Am. trop.
paniculata, Salzm. ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 187 = Wissadula hirsuta.
pannosa, R. Br. in Salt, Abyss. App. 65 = Abutilon glaucum.
pannosa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 565 = glutinosa.
parnassiaefolia, Hook. Ic. Pl. t. 385 = Malvastrum parnassiaefolium.
parviflora, Willd. Enum. Hort. Berol. 726 = Anoda parviflora.
parvifolia, DC. Prod. i. 461 = microphylla.
patens, Andr. Bot. Rep. t. 571 (Abutilonis sp. ?).—Abyss.
patens, D. Dietr. Syn. Pl. iv. 851 (Wissadulae sp.).—Bras.
patula, Pers. Syn. ii. 243 = glomerata.
pauciflora, D. Dietr. Syn. Pl. iv. 851 = Wissadula hirsuta.
paucifolia, DC. Prod. i. 472 (Hibisci sp.).—Ins. Timor.
pedata, Spreng. Syst. iv. Cur. Post. 259 = Callirhoë digitata.
pedicularifolia, Meyen, Reise, i. 460 = Malvastrum pedicilariaefolium.
pedunculoris, DC. Prod. i. 469 = Abutilon pedunculare.
pedunculata, A. Cunn. ex Benth. Fl. Austral. i. 193 = corrugata.
pellita, H. B. & K. Nov. Gen. et Sp. v. 263 = cordifolia.
pellita, Willd. ex Spreng. Syst. iii. 115 = paniculata.
pentacarpos, Roxb. ex Hornem. Hort. Hafn. Suppl. 78 (Abutilonis sp.).—Ind. or.
periplocifolia, Linn. Sp. Pl. 684 = Wissadula rostrata.
periplocifolia, Macfad. Fl. Jamaic. i. 85 = Wissadula divergens.
periptera, Sims, Bot. Mag. t. 1644 = Anoda punicea.
permollis, Willd. Enum. Hort. Berol. 728 = Abutilon permolle.
permutata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 67. —Abyss.
Perrottetiana, D. Dietr. Syn. Pl. iv. 855 = Abutilon fruticosum.
persica, Burm. ex Cav. Diss. i. 35 = Abutilon polyandrum.
peruviana, Cav. l. c. 36 = Abutilon arboreum.
petiolaris, DC. Prod. i. 470 = Abutilon petiolare.
petrophila, *F. Muell. in Linnaea*, xxv. (1852) 381.—Austral.
phaeotricha, F. Muell. l. c. 382 = virgata.
philippica, DC. Prod. i. 463 = rhombifolia.
phlebococca, Griseb. Cat. Pl. Cub. 25 = micrantha.
phyllanthos, Cav. Diss. v. 276. t. 127. f. 4 = Malvastrum phyllanthos.
physaloides, Presl, Rel. Haenk. ii. 105 = macrodon.
physocalyx, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 163 = hastata.
physocalyx, *F. Muell. Fragm.* iii. 3.—Austral.
pichinchensis, Humb. & Bonpl. Pl. Aequin. ii. 115 = Malvastrum pichinchense.
picta, Gill. ex Hook. Bot. Misc. iii. (1833) 154 = Abutilon striatum.
pilosa, Cav. Diss. i. 9. t. 8 = supina *vel* veronicaefolia.
pilosa, L'Hérit. Stirp. Nov. 130 = Abutilon graveolens.

SIDA :—

pilosa, Mill. Gard. Dict. ed. VIII. n. 17 = Melochia venosa.
pilosa, Retz. Obs. ii. 23 = veronicaefolia.
pilosa, Vell. Fl. Flum. 280; vii. t. 26 (Abutilonis sp.).—Bras.
pimpinellifolia, Mill. Gard. Dict. ed. VIII. n. 4 = spinosa.
pinnata, Cav. Ic. v. 13. t. 422. f. 1 = Malvastrum pinnatum.
planicaulis, Cav. Diss. i. 24. t. 3 = Malvastrum tricuspidatum.
planicaulis, DC. Prod. i. 461 = acuta.
planiflora, Cav. Diss. i. 32. t. 7. f. 4 = Abutilon Avicennae.
platycalyx, *F. Muell. ex Benth. Fl. Austral.* i. 197.—Austral.
pleiantha, *F. Muell. ex Benth. l. c.* 195.—Austral.
plumosa, Cav. Diss. i. 7 = ciliaris.
Pohliana, *Presl, Rel. Haenk.* ii. 108.—Peruv.
polyandra, Roxb. Hort. Beng. 50; Fl. Ind. iii. 173 = Abutilon polyandrum.
polyantha, Schlecht. ex Link, Enum. Hort. Berol. ii. 204 = Wissadula rostrata.
polycarpa, D. Dietr. Syn. Pl. iv. 855 = Abutilon indicum.
polycarpa, C. Sm. in Tuckey, Congo, 250 = Abutilon muticum.
polystachia, Vell. Fl. Flum. 279; vii. t. 22 = Abutilon ramiflorum, Wissadula rostrata.
populifolia, Lam. Encyc. i. 7 = Abutilon indicum.
portoricensis, *Spreng. Syst.* iii. 114.—Porto Rico.
potentilloides, *A. St. Hil. Fl. Bras. Mer.* i. 178.—Bras.
Presliana, D. Dietr. Syn. Pl. iv. 856 = Abutilon Haenkeanum.
procumbens, Sw. Prod. Veg. Ind. Occ. 101 = supina.
prostrata, Cav. Diss. i. 13 = flavescens.
prostrata, G. Don, Gen. Syst. i. 490 = acuta.
Pseudo-abutilon, *Desf. ex Steud. Nom.* ed. I. 774; ed. II. ii. 579.—Hab. ?
pterosperma, L'Herit. Stirp. Nov. 119. t. 57 = Cristaria multifida.
pubescens, Cav. Diss. i. 33. t. 7. f. 6 = Abutilon indicum.
pulchella, Willd. Enum. Hort. Berol. 724; Bonpl. Desc. Pl. Rar. Malm. 5. t. 2 = Plagianthus pulchellus.
pulcherrima, Hort. Turic. ex Steud. Nom. ed. II. ii. 579 (Quid ?).—Hab. ?
pulchra, Bert. ex Colla, Hort. Rip. 129. t. 34 (Abutilonis sp.).—Ecuador.
pungens, H. B. & K. Nov. Gen. et Sp. v. 263 = cordifolia.
purpurascens, Link, Enum. Hort. Berol. ii. 206 = Abutilon esculentum.
purpurea, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 239 (Kosteletzkyae sp. ?).—Porto Rico.
pusilla, Cav. Diss. i. 6. t. 1. f. 4; v. 127. f. 1 = spinosa.
pygmaea, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 238 = Malvastrum pygmaeum.
pyramidata, *Cav. Diss.* ii. 11. t. 1. f. 10; vi. t. 194. f. 1. —Ind. occ.
quaternata, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 60; *et in Syll. Ratisb.* ii. (1828) 73.—Bras.
quinqueangulata, D. Dietr. Syn. Pl. iv. 857 = seq.
quinqueloba, [Moç. & Sesse,] ex DC. Prod. i. 459 = Anoda hastata.
quinquenervia, *Duchass. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 176.—Panama.
racemiflora, Steud. Nom. ed. II. ii. 578 = Abutilon racemosum.
racemosa, Burm. f. Fl. Ind. 148 = glutinosa.
racemosa, Vell. Fl. Flum. 278; vii. t. 15 = Wissadula rostrata.
radiata, Linn. Sp. Pl. ed. II. 965 = Malachra radiata.
radicans, Cav. Diss. i. 8 = veronicaefolia.
radiciflora, Presl, Rel. Haenk. ii. 104 (Malvastri sp.).—Peruv.
ramiflora, D. Dietr. Syn. Pl. iv. 852 = Abutilon ramiflorum.
ramosa, Cav. Diss. i. 28. t. 6. f. 1 = Abutilon ramosum.
ramosa, Willd. ex Spreng. Syst. iii. 120 = diffusa.
ramosissima, D. Dietr. Syn. Pl. iv. 853 = Abutilon ramosissimum.
recisa, Link, Enum. Hort. Berol. ii. 203 = rhombifolia.
reclinans, Kunze, in Linnaea, xvii. (1843) 579 = veronicaefolia.

SIDA:—
reflexa, Cav. Diss. i. 36. t. 7; vi. t. 195. f. 1 = Abutilon reflexum.
Regelii, *Hort. ex Pasq. Cat. Ort. Bot. Nap.* 95, nomen. —Hab. ?
repanda, Roth, Nov. Pl. Sp. 328 = acuta.
repens, Cav. Diss. i. 7 = veronicaefolia.
reticulata, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 60; et in Syll. Ratisb. ii. (1828) 72 (Wissadulae sp.).—Bras.
retrofracta, DC. Prod. i. 467 = Bastardia viscosa.
retrorsa, L'Hérit. Stirp. Nov. 133. t. 64 = Abutilon reflexum.
retusa, Linn. Sp. Pl. ed. II. 961 = rhombifolia.
retusa, Wight, ex Mast. in Hook. f. Fl. Brit. Ind. i. 323 = spinosa.
retusifolia, Stokes, Bot. Mat. Med. iii. 527 = rhombifolia.
Retzii, J. F. Gmel. Syst. 1042 = veronicaefolia.
rhombifolia, *Linn. Sp. Pl.* 684.—Amphig. trop.
rhomboidea, Roxb. ex Flem. in As. Res. xi. (1810) 178 = rhombifolia.
ribifolia, *A. St. Hil. Fl. Bras. Mer.* i. 183. t. 34.—Bras.
Ricini, Spreng. Syst. iii. 116 = palmata.
ricinoides, L'Hérit. Stirp. Nov. 115. t. 55 = palmata.
rigida, D. Dietr. Syn. Pl. iv. 856 = Abutilon rigidum.
riparia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 65 = rhombifolia.
rivularis, D. Dietr. Syn. Pl. iv. 854 = Abutilon rivulare.
rosea, Link & Otto, Ic. Pl. Sel. 71. t. 32 = Abutilon esculentum.
rostrata, Schum. & Thonn. Beskr. Guin. Pl. 306 = Wissadula rostrata.
rosula, Meyen, ex Wedd. Chlor. And. ii. 278 = Malvastrum borussicum.
rotundifolia, *Hook. & Arn. Bot. Beech. Voy.* 79.—Ins. Sandvic.
rotundifolia, Lam. Encyc. i. 5 = cordifolia.
rotundifolia, *Perr. in Mém. Soc. Linn. Par.* iii. (1825) 145.—Java.
rubifolia, *A. St. Hil. Fl. Bras. Mer.* i. 183.—Bras.
rubra, D. Dietr. Syn. Pl. iv. 848 = micrantha.
rubra, Tenore, Cat. Ort. Nap. (1813) 96 = Anoda punicea.
ruderata, Macfad. Fl. Jamaic. i. 80 = rhombifolia.
rufescens, A. St. Hil. Fl. Bras. Mer. i. 185 = urens ?
rufinervis, Steud. Nom. ed. II. ii. 578 = Abutilon rufinerve.
rugosa, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 304 = acuta.
rupestris, Miq. in Lehm. Pl. Preiss. i. 241 = Hookeriana.
sabeana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 449 = Melochia pyramidata.
salicifolia, Forst. ex A. Gray, Bot. U. St. Expl. Exped. i. 158 = microphylla.
Saltii, *Steud. Nom.* ed. II. ii. 579.—Abyss.
salviaefolia, Presl, Rel. Haenk. ii. 110 = rhombifolia *vel* spinosa.
Saxifraga, Humb. & Bonpl. Pl. Aequin. ii. 116 = Malvastrum phyllanthos.
scabra, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 305 = spinosa.
scabrida, Wight & Arn. Prod. 57 = rhombifolia.
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 66.—Ind. or.; Afr. trop.
Schlechtendahlii, Steud. Nom. ed. II. ii. 579 = Abutilon erosum.
Schmitzii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 565.—Mexic.
Schrankii, DC. Prod. i. 472 = acuta.
scoparia, Lour. Fl. Cochinch. 414 = acuta.
scoparia, Vell. Fl. Flum. 277; vii. t. 12 = rhombifolia.
Sellowiana, Klotzsch, in Otto & Dietr. Allg. Gartenz. iv. (1836) 9 = Abutilon Sellovianum.
semicrenata, Link, Enum. Hort. Berol. ii. 202 = rhombifolia.
semidentata, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 51.—Bras.
sericea, Cav. Desc. 165 = Abutilon mollicomum.
sericea, Mill. Gard. Dict. ed. VIII. n. 15 = jamaicensis.
serrata, *Willd. ex Spreng. Syst.* iii. 111.—Venezuela.
sertum, Nutt. ex A. Gray, Bot. U. St. Expl. Exped. i. 163 = rotundifolia.
Sessei, Lag. Gen. et Sp. Nov. 21 (Abutilonis sp.).—Mexic.

SIDA:—
sessiliflora, G. Don, Gen. Syst. i. 499 = urens.
sessiliflora, Hook. Bot. Mag. t. 2857 = Abutilon crispum.
sessilis, Vell. Fl. Flum. 280; vii. t. 27 = Abutilon crispum.
setifera, *Presl, Rel. Haenk.* ii. 105.—Mexic.
Sherardiana, *Benth. in Journ. Linn. Soc.* vi. (1862) 101, in obs.—Europ. austr.; Oriens.
sinensis, J. F. Gmel. Syst. 1048 = *chinensis*.
Sonneratiana, Cav. Diss. i. 29. t. 6. f. 4 = Abutilon Sonneratianum.
sordida, Willd. ex Spreng. Syst. iii. 120 = Bastardia viscosa.
sparmannioides, Steud. Nom. ed. II. ii. 579 = Abutilon ramosum.
speciosa, Willd. ex Spreng. Syst. iii. 119 = Abutilon esculentum.
Spenceriana, *F. Muell. in Wing's South. Sc. Record* (April 1885).—Austral.
spicata, Cav. Diss. i. 24. t. 8. f. 1 = Gaya occidentalis (in Addend.).
spiciflora, DC. Prod. i. 468 = Wissadula spicata.
spinifex, [Moç. & Sesse,] ex DC. Prod. i. 472 = acerifolia.
spinosa, *Linn. Sp. Pl.* 683.—Amphig. trop.
spinosa, E. Mey. ex C. Muell. in Walp. Ann. vii. 394 = capensis.
spireaefolia, Willd. Enum. Hort. Berol. Suppl. 49 = acuta.
spodochroma, F. Muell. in Linnaea, xxv. (1852) 383 = corrugata.
Stauntoniana, DC. Prod. i. 460 = acuta.
stellata, Cav. Diss. i. 27 = Wissadula rostrata.
stellata, Torr. in Ann. Lyc. N. York, ii. (1828) 171 = Sphaeralcea angustifolia.
Steudelii, D. Dietr. Syn. Pl. iv. 857 (Cristariae sp.).—Chili.
stipularis, D. Dietr. l. c. 855 = Abutilon auritum.
stipulata, Cav. Diss. i. 22. t. 3. f. 10 = acuta.
stipulata, Chapm. Fl. S. U. St. 55 = rhombifolia.
stolonifera, Salzm. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 199 = decumbens.
striata, D. Dietr. Syn. Pl. iv. 852 = Abutilon striatum.
stylosa, *Baxt. ex Loud. Hort. Brit. Suppl.* ii. 673, nomen (Quid?).—Sibir. altaic.
subcordata, *Spun. in Linnaea*, xv. (1841) 172.—Ins. Timor.
subcuneata, *A. St. Hil. Fl. Bras. Mer.* i. 184.—Bras.
subdistans, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 50.—Bras.
suberosa, D. Dietr. Syn. Pl. iv. 853 = Bastardia bivalvis.
suberosa, Hort. ex Steud. Nom. ed. I. 775 = Anoda triangularis.
suberosa, L'Hérit. Stirp. Nov. 113. t. 54 = cordifolia.
suborbicularis, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 53.—Bras.
subsessilis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1863) 1. 199.—Bras.
subspicata, *F. Muell. ex Benth. Fl. Austral.* i. 195.—Austral.
subtriloba, DC. Prod. i. 466 = Gaya subtriloba.
sulphurea, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 23 = obliqua.
sundaica, Blume, Bijdr. 78 = Wissadula rostrata.
sundaica, Zipp. ex Miq. Fl. Ind. Bat. i. II. 141 = veronicaefolia.
sundensis, Spreng. Syst. iv. Cur. Post. 259 = Wissadula rostrata.
supina, *L'Hérit. Stirp. Nov.* 109bis, t. 52.—Am. trop.
surinamensis, Miq. in Linnaea, xxii. (1849) 469 = rhombifolia.
Swartzii, D. Dietr. Syn. Pl. iv. 847 = ulmifolia.
sylvatica, Cav. Diss. ii. 56; v. 276. t. 133. f. 2 (Abutilonis sp.).—Peruv.
tasmanica, Hook. f. in Hook. Journ. Bot. ii. (1840) 412 = Plagianthus pulchellus.
tenax, Buch.-Ham. ex Wight & Arn. Prod. 59 = glutinosa.
tenax, Salisb. Prod. 382 = Abutilon Avicennae.
tenuicaulis, Hook. f. in Trans. Linn. Soc. xx. (1847) 232 = spinosa.
terminalis, Cav. Diss. i. 29. t. 6. f. 7 = Abutilon terminale.

SIDA:—
ternata, *Linn. f. Suppl.* 307.—Afr. austr.
tetracocca, D. Dietr. Syn. Pl. iv. 859 = Wissadula rostrata.
thyrsiflora, *Miq. Pl. Jungh.* 287.—Java.
tiliaefolia, Fisch. Cat. Hort. Gorenk. (1808); ex DC. Prod. i. 470 = Abutilon Avicennae.
timoriensis, DC. l. c. 468 = Abutilon timoriense.
tomentella, *Miq. in Linnaea*, xxii. (1849) 553.—Bras.
tomentosa, Cav. Desc. 164 = crassifolia.
tomentosa, Roxb. Hort. Beng. 50; Fl. Ind. iii. 178 = Abutilon muticum.
tomentosa, Vell. Fl. Flum. 277; vii. t. 14 = cordifolia.
tomentosa, Wall. Cat. n. 1852 = Abutilon graveolens.
tragiaefolia, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 164.—Texas.
triangularis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 725 = hastata, Anoda triangularis.
trichoda, D. Dietr. Syn. Pl. iv. 856 = Abutilon crispum.
trichopoda, F. Muell. in Linnaea, xxv. (1852) 384 = corrugata.
trichosperma, D. Dietr. Syn. Pl. iv. 857 = Abutilon calycinum.
tricuspidata, Cav. Diss. i. 30 = Abutilon Jacquini.
tricuspis, *Desf. Tabl.* ed. II. 171.—Hab. ?
tridentata, Cav. Ic. iv. 6 = ciliaris.
triflora, Vell. Fl. Flum. 280; vii. t. 24 = Abutilon esculentum.
triloba, *Cav. Diss.* i. 11. t. 1. f. 11; v. t. 131. f. 1; *Thunb. Prod. Pl. Cap.* 117.—Afr. austr.
triloba, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 67 = permutata.
trinervia, Splitg. ex De Vriese, in Nederl. Kruidk. Arch. i. (1848) 339 = rhombifolia.
triquetra, Linn. Sp. Pl. ed. II. 963 = Abutilon triquetrum.
trisecta, D. Dietr. Syn. Pl. iv. 857 (Cristariae sp.).—Chili.
tristis, Schlecht. in Linnaea, iii. (1828) 271 = jamaicensis.
trisulcata, Jacq. Enum. Pl. Carib. 26 = Abutilon triquetrum.
trivialis, Macfad. Fl. Jamaic. i. 78 = acuta.
truncata, Cav. Diss. i. 35. t. 6. f. 7 = cordifolia.
truncata, L'Hérit. Stirp. Nov. 107 = ulmifolia ?
truncata, *Vell. Fl. Flum.* 278; vii. t. 17.—Bras.
truncatula, Blanco, Fl. Filip. ed. I. 548 = rhombifolia.
truncatula, J. F. Gmel. Syst. 1049 = *truncata*.
tubulosa, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austr. 390 = Abutilon tubulosum.
ulmifolia, *Cav. Diss.* i. 15. t. 2. f. 4.—Am. trop.
ulmifolia, Hook. & Arn. Bot. Beech. Voy. 79 = Meyeniana.
ulmifolia, Mill. Gard. Dict. ed. VIII. n. 1 = acuta.
ulmifolia, Retz. Obs. iii. 37 = spinosa.
ulmifolia, Spreng. Syst. iii. 113 = Gaya occidentalis (in Addend.).
ulmifolia, Willd. Enum. Hort. Berol. Suppl. 49 = Malvastrum tricuspidatum.
umbellata, Linn. Syst. ed. X. 1145 = Abutilon umbellatum.
umbellifera, D. Dietr. Syn. Pl. iv. 853 = praec.
unidentata, D. Dietr. l. c. 858 = Anoda lanceolata.
unilocularis, L'Hérit. Stirp. Nov. 117. bis, t. 56 = veronicaefolia.
urens, *Linn. Syst.* ed. X. 1145.—Afr. et Am. trop.
urticaefolia, A. St. Hil. Fl. Bras. Mer. i. 189 = flavescens.
urticaefolia, Wight & Arn. Prod. 59 = glutinosa.
Velloziana, Steud. Nom. ed. II. ii. 579 = cordifolia.
velutina, Willd. ex Spreng. Syst. iii. 115 = cordifolia.
venosa, A. Dietr. in Otto & Dietr. Allg. Gartenz. iii. (1835) 33 = Abutilon venosum.
venusta, Schlecht. in Linnaea, xii. (1838) 364 = filipes ?
veronicaefolia, *Lam. Encyc.* i. 5.—Am. trop.
verruculata, *DC. Prod.* i. 473.—Bras.
verticillata, Cav. Diss. i. 13. t. 1. f. 12 = urens.
Vescoana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 504.—Madag.
vesicaria, Cav. Diss. ii. 55. t. 14. f. 3 = Abutilon indicum.

SIDA:—
vestita, Steud. Nom. ed. II. ii. 579 = cordifolia.
viarum, *A. St. Hil. Fl. Bras. Mer.* i. 182.—Bras.
villosa, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. austr.
villosa, Wall. Cat. n. 1856 C = Abutilon muticum.
viminea, Fisch. ex Link, Enum. Hort. Berol. ii. 202 = linifolia.
virgata, Cav. Ic. i. 53. t. 73 = Abutilon virgatum.
virgata, *Hook. in Mitch. Journ. Trop. Austral.* 361.—Austral.
viridis, A. St. Hil. & Naud. ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 175 = glomerata.
viscidula, Blume, Bijdr. 76 = glutinosa.
viscidula, Klotzsch, ex Steud. Nom. ed. II. ii. 580 = Luschnathiana.
viscosa, Linn. Syst. ed. X. 1145 = Bastardia viscosa.
viscosa, Macfad. Fl. Jamaic. i. 84 = Bastardia bivalvis.
vitellina, *Hoffmgg. Preisv. Pfl.* viii. *Nachtr.* 33; *ex Walp. Rep.* i. 315.—Java.
vitifolia, Cav. Ic. v. 12. t. 428 = Abutilon vitifolium.
Vogelii, Hook. f. in Hook. Niger Fl. 231 = acuta.
Wallichii, Steud. Nom. ed. II. ii. 580 = Abutilon polyandrum.
waltheriaefolia, Boj. ex Baker, Fl. Maurit. 19 = cordifolia.
Weinmanni, *Heynh. Nom.* ii. 663 (Quid?).—China.
Wightiana, D. Dietr. Syn. Pl. iv. 845 = glutinosa.
Willdenowii, D. Dietr. l. c. 847 = glutinosa.
Zollingeriana, *Miq. Fl. Ind. Bat.* i. II. 141.—Java.
Zuccagnii, Spreng. Syst. iii. 121 = Anoda hastata.

SIDALCEA, A. Gray, ex Benth. Pl. Hartw. 300 (1848); et in Mem. Am. Acad. N. S. iv. (1849) 18.
MALVACEAE, Benth. & Hook. f. i. 201.
asprella, *Greene, in Bull. Calif. Acad.* i. III. (1885) 78.—Am. bor. occ.
atacosa, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 449 = Sphaeralcea Fendleri.
californica, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 18.—Calif.
calycosa, *M. Jones, in Am. Natural.* xvii. II. (1883) 875.—Calif.
campestris, *Greene, in Bull. Calif. Acad.* i. III. (1885) 76.—Calif.
candida, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 20, 24.—N. Mexic.
delphinifolia, A. Gray, ex Benth. Pl. Hartw. 300 = malvaeflora.
diploscypha, *A. Gray, ex Benth. l. c.*—Calif.
glaucescens, *Greene, in Bull. Calif. Acad.* i. III. (1885) 77.—Am. bor. occ.
Hartwegii, *A. Gray, ex Benth. Pl. Hartw.* 300.—Calif.
hirsuta, *A. Gray, Pl. Wright.* i. 16.—Calif.
humilis, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 20 = malvaeflora.
malachroides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 332.—Calif.
malvaeflora, *A. Gray, ex Benth. Pl. Hartw.* 300.—Am. bor. occ.
malvaeflora, A. Gray, Pl. Wright. i. 16, partim = neo-mexicana.
malvaeflora, S. Wats. in Bot. King's Exped. 46, partim = glaucescens.
neo-mexicana, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 23.—N. Mexic.
nodosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 566.—Peruv.
oregana, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 20 = malvaeflora.
peruviana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 566.—Peruv.
spicata, *Greene, in Bull. Calif. Acad.* i. III. (1885) 76.—Calif.
sulcata, Curran, ex Greene, l. c. 79 = calycosa.
tenella, Greene, l. c. I. (1884) 7 = Hartwegii.
triloba, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1863) I. 566.—Chili.
vitifolia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 332.—Calif.

SIDE, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = **Sida**, Linn. (Malvac.).
abutila, St. Lag. l. c. = Abutilon Avicennae.
phyllantha, St. Lag. l. c. = Malvastrum phyllanthos.

SIDERANTHUS, [Nutt.] in Fras. Cat. (1813) nn. 80, 81; et ex Sweet, Hort. Brit. ed. I. 227 (1826) = **Aplopappus**, Cass. (Compos.).
integrifolius, [Nutt.] l. c. n. 80 = Chrysopsis villosa.
pinnatifidus, [Nutt.] l. c. n. 81 = A. spinulosus.
spinulosus, Sweet, Hort. Brit. ed. I. 227 = A. spinulosus.
villosus, Sweet, l. c. = Chrysopsis villosa.

SIDERASIS, Rafin. Fl. Tellur. iii. 67 (1836) = **Tradescantia**, Linn. (Commel.).
acaulis, Rafin. l. c. = T. fuscata.

SIDERITIS, Tourn. ex Linn. Gen. ed. I. 171 (1737).
LABIATAE, Benth. & Hook. f. ii. 1205.
BURGSDORFIA, Moench, Meth. 392 (1794).
EMPEDOCLIA, Rafin. Caratteri, 78 (1810).
FRACASTORA, Adans. Fam. ii. 505 (1763).
FRANCASTORA, Steud. Nom. ed. II. ii. 580, in syn. (1841), sphalm.
HESIODIA, Moench, Meth. 391 (1794).
LEUCOPHAR, Webb & Berth. Phyt. Canar. iii. 99, 168, 172 (1835–60).
MARRUBIASTRUM, Tourn. ex Moench, Meth. 391 (1794).
NAVICULARIA, Heist. ex Adans. Fam. ii. 188 (1763).
aegyptiaca, Pers. Syn. ii. 117 = Stachys rosea.
alopecuros, Scop. Fl. Carn. ed. II. i. 413 = Stachys alopecuros.
alpina, Pourr. in Mém. Acad. Toul. iii. (1788) 328 = hyssopifolia.
ambigua, Fenzl, in Tchihat. Asie Min. Bot. ii. 163 = libanotica.
angustifolia, Lag. Gen. et Sp. Nov. 18. n. 231 = Lagascana.
angustifolia, *Lam. Encyc.* ii. 168.—Hispan.
approximata, Gasp. ex Guss. Fl. Sic. Prod. Suppl. 191 = romana.
arborescens, *Salzm. ex Benth. Lab. Gen. et Sp.* 579.—Hispan.
argosphacelus, *Clos, in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 80.—Ins. Canar.
arguta, Boiss. Diagn. Ser. II. iv. 32 = libanotica.
atlantica, Pomel, Nouv. Mat. Fl. Atl. 120 = hyssopifolia.
baetica, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 18.—Hispan.
Balansae, *Boiss. Diagn.* Ser. II. iv. 35.—As. Min.
Balansae, Coss. in Bull. Soc. Bot. Fr. xx. (1873) 256; xxii. (1875) 65 = Cossoniana.
Bourgaeana, Boiss. & Reut. Diagn. Ser. II. iv. 34 = leucantha.
Bourgaei, Boiss. Fl. Orient. iv. 712 = libanotica.
brutia, Tenore, Fl. Nap. ii. 16 = sicula.
bullata, H. B. & K. Nov. Gen. et Sp. ii. 306 = Sphacele bullata.
calycantha, Bieb. Fl. Taur. Cauc. iii. 393 = Stachys lavandulaefolia.
canariensis, *Linn. Sp. Pl.* 574.—Ins. Canar.
candicans, *Ait. Hort. Kew.* ed. I. ii. 289.—Ins. Mader.
candida, Salisb. Prod. 79 = candicans.
cassia, Boiss. Diagn. Ser. I. xii. 76, sphalm. = Stachys cassia.
Cavanillesii, Lag. Gen. et Sp. Nov. 18 = scordioides.
chamaedrifolia, Cav. Ic. iv. 1. t. 301 = hirsuta.
chamaedrifolia, Hoffmgg. & Link, Fl. Port. i. 99 = hyssopifolia.
chamaedryoides, Duf. ex Willk. & Lange, Prod. Fl. Hisp. ii. 455 = scordioides.
ciliata, *Thunb. Fl. Jap.* 245.—Japon.
cilicica, *Boiss. & Bal. Diagn.* Ser. II. iv. 29.—Cilicia.
condensata, *Boiss. & Heldr. Diagn.* Ser. I. xii. 71.—As. Min.
Cossoniana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 622.—Maroce.
crenata, Lapeyr. Hist. Abr. Pl. Pyr. 331 = hyssopifolia.
cretica, Boiss. Fl. Orient. iv. 708 = syriaca.
cretica, *Linn. Sp. Pl.* 574.—Ins. Creta.
cretica, Sibth. & Sm. Fl. Graec. Prod. i. 400 = theezans.
crispata, *Willd. Enum. Hort. Berol.* 606.—Hispan.
curvidens, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 100.—Lycia.
dasygnaphala, *Clos, in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 80.—Ins. Canar.
decumbens, Moench, Meth. 390 = sicula.

SIDERITIS:—
decumbens, Sieber, ex Benth. in DC. Prod. xii. 440 = pullulans.
decumbens, Thunb. Prod. Pl. Cap. 95 = Stachys Lamarckii.
Dendro-chahorra, *Bolle, in Bonplandia*, viii. (1860) 285.—Ins. Canar.
deserti, Noé, ex Coss. in Bull. Soc. Bot. Fr. iv. (1857) 472 = Marrubium deserti.
discolor, Willd. ex Benth. in Linnaea, xi. (1837) 330; Bolle, in Bonplandia, viii. (1860) 285 = macrostachyos.
distans, Willd. Sp. Pl. iii. 66 = taurica.
dura, Benth. Lab. Gen. et Sp. 576 = taurica.
ebracteata, Asso, Mant. Stirp. Arag. 171 = montana.
elegans, Murr. in Comm. Gotting. (1778) 92. t. 4 = lanata.
Endressi, Willk. in Bot. Zeit. xvii. (1859) 276 = hirsuta.
errata, Walp. Rep. iii. 851, sphalm. = *serrata*.
erythrantha, *Boiss. & Heldr. Diagn.* Ser. I. xii. 70.—As. Min.
fetida, Poir. Encyc. Suppl. ii. 382 = scordioides.
florida, Boiss. & Heldr. Diagn. Ser. II. iv. 31 = scardica.
foetens, Benth. Lab. Gen. et Sp. 581 = arborescens.
foetens, Clemente, ex Lag. Gen. et Sp. Nov. 18 = lasiantha.
foetida, Desf. Tabl. ed. I. 59 = scordioides.
fragrans, *Costa, ex Willk. in Bot. Zeit.* xvii. (1859) 289.—Hispan.
fruticulosa, Pourr. in Mém. Acad. Toul. iii. (1788) 328 = scordioides.
Funkiana, Willk. in Bot. Zeit. xvii. (1859) 282 = angustifolia.
georgica, C. Koch, in Linnaea, xxi. (1848) 694 = taurica.
glacialis, *Boiss. ex Willk. in Bot. Zeit.* xvii. (1859) 276.—Hispan.
glauca, *Cav. Ic.* ii. 68. t. 185.—Hispan.
gomeraea, *Bolle, in Bonplandia*, viii. (1860) 286.—Ins. Canar.
gracilis, *Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1885) 98.—Lycia.
grandiflora, *Salzm. ex Benth. Lab. Gen. et Sp.* 577.—Hispan.; Marocc.
Guillonii, Timb. in Mém. Acad. Toul. Sér. VII. iv. (1872) 382 = hyssopifolia.
Guyoniana, *Boiss. & Reut. Pugill.* 98.—Afr. bor.
heraclea, Col. ex Benth. in DC. Prod. xii. 447 = Stachys heraclea.
hirsuta, Eichw. Beschr. Casp.-Cauc. 24 = montana.
hirsuta, Gouan, Fl. Monsp. 85 = Stachys recta.
hirsuta, *Linn. Sp. Pl.* 575.—Reg. Mediterr.
hirsuta, Sieber, ex Benth. Lab. Gen. et Sp. 536 = Stachys italica.
hirta, Roth, in Usteri, Ann. Bot. x. (1794) 48 = scordioides.
hirtula, Brot. Fl. Lusit. i. 161 = hyssopifolia.
hispanica, Mill. Gard. Dict. ed. VIII. n. 7 = hirsuta.
hololeuca, *Boiss. & Heldr. Diagn.* Ser. I. xii. 69.—As. Min.
hyssopifolia, Georgi, Beschr. Russ. Reich. v. 1081 = Stachys recta.
hyssopifolia, *Linn. Sp. Pl.* 575.—Europ. austr.
hyssopifolia, Willd. ex Benth. in Linnaea, xi. (1837) 331 = scordioides.
ilicifolia, Willd. Enum. Hort. Berol. 606 = fragrans.
incana, Benth. in DC. Prod. xii. 444 = scordioides.
incana, Habl. Beschr. Taur. 148 = taurica.
incana, *Linn. Sp. Pl.* ed. II. 802.—Hispan.
infernalis, *Bolle, in Bonplandia*, viii. (1860) 286.—Ins. Canar.
integrifolia, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137, nomen = hyssopifolia.
juncea, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 459 = lasiantha.
jurana, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137, nomen = hyssopifolia.
Lagascana, *Willk. in Bot. Zeit.* xvii. (1859) 282.—Hispan.
lanata, *Linn. Amoen. Acad.* iv. 459.—Graec.; As. Min.
lanata, Steud. & Hochst. ex Benth. in DC. Prod. xii. 446 = remota.
lasiantha, *Juss. ex Pers. Syn.* ii. 117.—As. Min.
lavandulacea, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 458 = incana.
leucantha, *Cav. Ic.* iv. 2. t. 304.—Hispan.

SIDERITIS:—
libanotica, *Labill. Ic. Pl. Syr. Dec.* iv. 13. t. 8.—As. Min.; Syria.
linearifolia, Boiss. Voy. Bot. Espagne, ii. 507 = angustifolia.
linearifolia, Bourg. ex Nym. Consp. 584 = arborescens.
linearifolia, Lag. Gen. et Sp. Nov. 18 = pungens.
linearifolia, Lam. Encyc. ii. 168 = Stachys recta?
linearifolia, Willd. ex Benth. in Linnaea, xi. (1837) 331 = scordioides.
linearifolia, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 458 = Lagascana.
litoralis, Timb. ex Nym. Consp. 584 = hirsuta.
longicaulis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 136, nomen = hyssopifolia.
lucida, J. Gay, ex Nym. Consp. 584 = hyssopifolia.
lurida, *J. Gay, ex Benth. in DC. Prod.* xii. 443.—Hispan.
lycaonica, *Boiss. & Heldr. Diagn.* Ser. II. iv. 35.—Lycaonia.
Lyciae, *Boiss. & Heldr. l. c.* Ser. I. xii. 70.—Lycia.
macrostachyos, *Poir. Encyc. Suppl.* ii. 381.—Ins. Canar.
maculata, Gilib. Fl. Lituan. i. 81 = Stachys recta.
marmorea, *Bolle, in Bonplandia*, viii. (1860) 285.—Ins. Canar.
Massoniana, *Benth. Lab. Gen. et Sp.* 573.—Ins. Mader.
microstegia, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 712 = libanotica.
montana, *Linn. Sp. Pl.* 575.—Reg. Mediterr.; Oriens.
montana, Sibth. & Sm. Fl. Graec. vi. 42. t. 551 = remota.
mucronata, Sieber, ex Benth. in DC. Prod. xii. 440 = pullulans.
mutica, Boiss. Fl. Orient. iv. 706 = romana.
nemausensis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137 = hirsuta.
nepetaefolia, Willd. ex Benth. in Linnaea, xi. (1837) 329 = Stachys nepetaefolia.
nervosa, Desf. ex Steud. Nom. ed. I. 775, 836 = Micromeria graeca.
nigricans, Lam. Encyc. ii. 168 = lanata.
ochroleuca, *Noé, ex Willk. in Bot. Zeit.* xvii. (1859) 282.—Algeria.
Ocymastrum, Linn. Syst. ed. X. 1098; Gouan, Hort. Monsp. 278 = Stachys hirta.
oleaefolia, *Mill. Gard. Dict.* ed. VIII. n. 4.—Ins. Creta.
orophila, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137, nomen = hyssopifolia.
ovata, *Cav. Ic.* i. 36. t. 48.—Peruv.
pallida, Thunb. Prod. Pl Cap. 95 = Stachys rosmarinifolia.
paniculata, H. B. & K. Nov. Gen. et Sp. ii. 307 = Sphacele paniculata.
parvifolia, *Lam. Illustr.* iii. 63. t. 505. f. 2.—Hab.?
patula, Dulac, Fl. Hautes-Pyr. 411 = scordioides.
peloponnesiaca, Boiss. & Heldr. Diagn. Ser. II. iv. 32 = theezans.
perfoliata, *Linn. Sp. Pl.* 575.—As. Min.; Syria.
peruviana, Willd. ex Benth. in Linnaea, xi. (1837) 330 (Salviae sp.).—Hab.?
Peyrei, Timb. in Mém. Acad. Toul. Sér. VII. iv. (1872) 383 = scordioides.
phleoides, Lange, ex Willk. & Lange, Prod. Fl. Hisp. ii. 459 = lasiantha.
phlomifolia, Salisb. Prod. 79 = canariensis.
phlomoides, *Boiss. & Bal. Diagn.* Ser. II. iv. 33.—Cappadoc.
pisidica, *Boiss. & Heldr. l. c.* Ser. I. xii. 72.—As. Min.
plumosa, Thunb. Prod. Pl. Cap. 95 = Stachys integrifolia.
prostrata, *Zuccagni, in Roem. Collect.* 144.—Hab.?
provincialis, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137 = hirsuta.
pullulans, *Vent. Jard. Cels*, t. 98.—Syria.
punctata, *C. Sm. in Tuckey, Congo*, 251, nomen.—Ins. Canar.
pungens, *Benth. Lab. Gen. et Sp.* 579.—Hispan.
purpurea, *Talbot, in Benth. l. c.* 742.—Ins. Ion.
pyrenaica, Endr. ex Nym. Consp. 584 = hirsuta.
pyrenaica, Poir. Encyc. Suppl. ii. 383 = hyssopifolia.
Raeseri, *Boiss. & Heldr. Diagn.* Ser. II. iv. 30.—Graecia.

SIDERITIS:—
remota, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 322.—Ins. Samos.
remotiflora, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 136, nomen = hyssopifolia.
rhytidea, Spreng. Syst. v. 644 (Index) = hololeuca.
rigida, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 (Quid?).—Java.
romana, Georgi, Beschr. Russ. Reich. v. 1081 = montana.
romana, *Linn. Sp. Pl.* 575.—Reg. Mediterr.
rosea, Desf. in Ann. Mus. Par. x. (1807) 302. t. 24 = Stachys rosea.
rotundifolia, Willd. ex Benth. in Linnaea, xi. (1837) 331 = hirsuta.
rugosa, Poir. Encyc. Suppl. ii. 383 = hololeuca.
rugosa, Thunb. Prod. Pl. Cap. 95 = Stachys Burchellii.
rugosa, Willd. ex Benth. in Linnaea, xi. (1837) 330 = Sphacele bullata.
ruscinonensis, Timb. in Mém. Acad. Toul. Sér. VII. iv. (1872) 380 = hyssopifolia.
saetabensis, Rouy, in Bull. Soc. Bot. Fr. xxix. (1882) 125 = angustifolia.
salviaefolia, H. B. & K. Nov. Gen. et Sp. ii. 307 = Sphacele salviaefolia.
scardica, *Griseb. Spicil. Fl. Rumel.* ii. 144.—Macedon.
scordioides, Koch, Syn. Fl. Germ. ed. I. 570 = hyssopifolia.
scordioides, *Linn. Syst.* ed. X. 1098.—Europ. austr.
sericea, Pers. Syn. ii. 118 = incana.
serrata, Lag. Gen. et Sp. Nov. 18 = spinosa.
sicula, *Ucria, in Roem. Archiv*, i. 1. (1796) 69.—Europ. austr.
sipylea, *Boiss. Diagn.* Ser. I. v. 32.—As. Min.
soluta, *Clos, in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 81.—Ins. Canar.
spathulata, Lam. Fl. Fr. ii. 377 = romana.
spinosa, Benth. Cat. Pl. Pyr. 121 = scordioides.
spinosa, *Lam. Encyc.* ii. 169.—Hispan.; Afr. bor.
stachydioides, *Willk. in Bot. Zeit.* viii. (1850) 78.—Hispan.
stricta, *Boiss. & Heldr. Diagn.* Ser. I. xii. 72.—As. Min.
stricta, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 137, nomen = hyssopifolia.
subspinosa, Cav. Ic. iii. 5 = spinosa.
suffruticosa, Mill. ex Benth. in DC. Prod. xii. 444 = incana.
syriaca, Bory & Chaub. ex Boiss. Fl. Orient. iv. 711 = theezans.
syriaca, *Linn. Sp. Pl.* 574.—Oriens.
syriaca, Pall. ex Bieb. Fl. Taur. Cauc. ii. 43 = taurica.
taurica, *Willd. Sp. Pl.* iii. 66.—As. Min.; Reg. Caucas.
teucriifolia, *Delile, Fl. Aeg. Illustr.* 65.—Aegypt.
theezans, *Boiss. & Heldr. Diagn.* Ser. I. vii. 58.—Graecia.
tomentosa, Pourr. in Mém. Acad. Toul. iii. (1788) 328 = hirsuta.
tomentosa, *Ucria, in Roem. Archiv*, i. 1. (1796) 69.—Sicil.
Tragoriganum, Lag. Gen. et Sp. Nov. 18 = incana.
trinervia, Gilib. Fl. Lituan. i. 81 = Stachys annua.
vestita, Boiss. Diagn. Ser. II. iv. 33 = glacialis.
villosa, *Coss. & Ball, in Bull. Soc. Bot. Fr.* xx. (1873) 255; xxii. (1875) 65.—Marocc.
virgata, Desf. Fl. Atlant. ii. 15. t. 125 = incana.
viridifolia, Dulac, Fl. Hautes-Pyr. 411 = hyssopifolia.

SIDERODENDRON, Roem. & Schult. Syst. iii. 10, 185 (1818) = seq.

SIDERODENDRUM, Schreb. Gen. 71 (1789) = **Ixora**, Linn. (Rubiac.).
ferreum, J. F. Gmel. Syst. 242 = I. ferrea.
floribundum, A. Rich. Fl. Cub. Fanerog. ii. 24 = I. floribunda.
laxiflorum, Benth. in Hook. Journ. Bot. iii. (1841) 235 = I. laxiflora.
macrophyllum, Benth. l. c. 234 = I. macrophylla.
multiflorum, Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 184 = Myonima multiflora.
paniculatum, Willd. ex Cham. & Schlecht. in Linnaea, iv. (1829) 26 = Psychotria cordifolia.
triflorum, Vahl, Eclog. Am. i. 10 = I. ferrea.

SIDEROXYLOIDES, Jacq. Stirp. Am. 19. t. 175. f. 9 (1763) = **Ixora**, Linn. (Rubiac.).
ferreum, Jacq. l. c. = I. ferrea.

SIDEROXYLON, [Dill.] Linn. Syst. ed. I (1735); Hort. Cliff. 69 (1737). *SAPOTACEAE*, Benth. & Hook. f. ii. 655.
DECATELES, Rafin. Sylva Tellur. 35 (1838).
MASTICHODENDRON, R. Hedw. Gen. 116 (1806).
OLIGOTHECA, A. DC. in DC. Prod. viii. 174 (1844).
ROBERTSIA, Endl. Gen. 740 (1839).
ROEMERIA, Thunb. Nov. Gen. ix. 130 (1798), pro parte.
XANTOLIS, Rafin. Sylva Tellur. 35 (1838).
Acouma, A. DC. in DC. Prod. viii. 184 = obovatum, *Gaertn. f.*
arboreum, Buch.-Ham. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 536 = Sarcosperma arboreum.
argenteum, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 16.—Java.
argenteum, Spreng. Syst. i. 666 = Brownii.
argenteum, *Thunb. Prod. Pl. Cap.* 36.—Afr. austr.
armatum, Roth, Nov. Pl. Sp. 138 = tomentosum.
arnhemicum, *Benth. & Hook. f. ex F. Muell. Census*, 91.—Austral.
assamicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 537.—Assam.
atrovirens, Lam. Encyc. i. 245 = inerme.
atrovirens, Willd. Enum. Hort. Berol. Suppl. 11 = inerme, Myrsine mitis.
attenuatum, A. DC. in DC. Prod. viii. 178 = ferrugineum.
australe, *Benth. & Hook. f. ex F. Muell. Census*, 92.—Austral.
avenium, *Burck, in Ann. Jard. Buitenz.* v. (1885) 16.—Ins. Banca.
Balitbitan, *Blanco, Fl. Filip.* ed. I. 130.—Ins. Philipp.
bancanum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 10.—Ins. Banca.
Bilitbitan, A. DC. in DC. Prod. viii. 186, sphalm. = S. Balitbitan.
Bojeranum, *A. DC. l. c.* 179.—Ins. Mascar.
borbonicum, *A. DC. l. c.*—Ins. Borbon.
borneense, *Burck, in Ann. Jard. Buitenz.* v. (1885) 15.—Borneo.
Boutonianum, *A. DC. in DC. Prod.* viii. 179.—Ins. Mascar.
brevipes, *Baker, in Oliver, Fl. Trop. Afr.* iii. 502.—Afr. trop.
Brownii, *F. Muell. Census*, 92.—Austral.
Brownlessianum, *F. Muell. Fragm.* viii. 111.—Austral.
cantoniense, *Lour. Fl. Cochinch.* 122.—China.
carolinense, Sargent, Rep. For. N. Am. (U. St. 10th Census, ix.) 101 = Bumelia tenax.
celebicum, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 8.—Ins. Celebes.
chartaceum, *Benth. & Hook. f. ex F. Muell. Census*, 91.—Austral.
chrysophylloides, Michx. Fl. Bor. Am. i. 123 = Bumelia tenax.
chrysophyllum, *De Vriese, Pl. Ind. Bat. Or.* 59.—Java.
cinereum, Boj. Hort. Maurit. 197; Wall. Cat. n. 4157 = Bojerianum.
cinereum, *Lam. Encyc.* i. 244.—Ins. Borbon.
confertum, *Wright, in Sauv. Fl. Cub.* 86.—Cuba.
Corsanii, Boj. ex Baker, Fl. Maurit. 193 = Boutonianum.
costatum, *F. Muell. Census*, 92.—Austral.
crassipedicellatum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 57.—Bras.
cuneatum, A. DC. in DC. Prod. viii. 181 = obovatum, *Gaertn. f.*
cuspidatum, *A. DC. l. c.* 183.—Guian.
cylindrocarpon, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 72. t. 282.—Bras.; Peruv.
cymosum, Linn. f. Suppl. 152 = Olinia cymosa.
cyrtobotryum, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 57.—Bras.
decandrum, Linn. Mant. i. 48 = Bumelia lycioides.
decumbens, J. F. Gmel. Syst. 1626, sphalm. = praec.
densiflorum, *Baker, in Oliver, Fl. Trop. Afr.* iii. 503.—Afr. trop.
dentatum, *Burm. f. Fl. Cap. Prod.* 6.—Afr. austr.
dictyoneurum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 517.—Ind. occ.

SIDEROXYLON :—
diospyroides, *Baker, in Oliver, Fl. Trop. Afr.* iii. 502. —Afr. trop.
Duclitan, *Blanco, Fl. Filip.* ed. I. 129.—Ins. Philipp.
dulcificum, *A. DC. in DC. Prod.* viii. 183.—Afr. trop.
Dulitan, A. DC. l. c. 185, sphalm. = S. Duclitan.
durum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 56 = cuspidatum.
egense, A. DC. in DC. Prod. viii. 182 = rugosum.
elegans, *A. DC. l. c.* 183.—Guian.
ellipticum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 56 = cuspidatum.
euphlebium, *F. Muell. Census*, 92.—Austral.
ferrugineum, *Hook. & Arn. Bot. Beech. Voy.* 266. t. 55.—As. trop.; Ins. Seychelles.
fimbriatum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 76.—Ins. Socotr.
firmum, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 17.—Ins. Banca.
floribundum, *Griseb. Fl. Brit. W. Ind.* 399.—Ind. occ.
foetidissimum, *Jacq. Enum. Pl. Carib.* 15.—Ind. occ.
Gamblei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 538.—Reg. Himal.
Gardnerianum, *A. DC. in DC. Prod.* viii. 182.—Bras.
glabrescens, *Miq. Fl. Ind. Bat. Suppl.* 580.—Ins. Banca.
grandiflorum, *A. DC. in DC. Prod.* viii. 180.—Ins. Mascar.
grandifolium, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 348. —Ind. or.; Burma.
guadalupense, *Spreng. Syst.* i. 666.—Ins. Guadalup.
guyanense, *A. DC. in DC. Prod.* viii. 182.—Guian.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 537.—Reg. Himal.
Howeanum, *F. Muell. Census*, 92.—Austral.
imbricarioides, *A. DC. in DC. Prod.* viii. 180.—Ins. Maurit.
indicum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 13. —Malaya.
inerme, *Forsk. Fl. Aegypt. Arab.* 204.—Arab.
inerme, Krauss, ex A. DC. in DC. Prod. xiv. 634 = Colpoon compressum.
inerme, *Linn. Sp. Pl.* 192.—Afr. austr.
javense, *Burck, in Ann. Jard. Buitenz.* v. (1885) 14. —Java.
laeve, Walt. Fl. Carol. 100 = Bumelia lycioides.
lanceolatum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 14.—Ins. Banca.
lanuginosum, Michx. Fl. Bor. Am. i. 122 = Bumelia lanuginosa.
laurifolium, Comm. ex A. DC. in DC. Prod. viii. 180 = imbricarioides.
laurifolium, Lam. Encyc. i. 244 = Myrsine melanophleos.
Lessertii, *Baker, Fl. Maurit.* 194.—Ins. Maurit.
linggense, *Burck, in Ann. Jard. Buitenz.* v. (1885) 15.—Malaya.
longifolium, *Boj. ex A. DC. in DC. Prod.* viii. 180.—Ins. Maurit.
longistylum, *Baker, in Oliver, Fl. Trop. Afr.* iii. 502. —Afr. trop.
lucidum, *Soland. ex Lam. Encyc.* i. 246.
lyciifolium, Salisb. Prod. 139 = Bumelia lycioides.
lycioides, Linn. Sp. Pl. ed. II. 279 = Bumelia lycioides.
lycioides, Wangenh. ex Steud. Nom. ed. I. 776 = 'Ilicis *aut* Prinos species.'—Am. bor.
macrophyllum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 20.—Borneo.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 536.—Malacca.
malaccense, *C. B. Clarke, l. c.* 537.—Malacca.
Manglilla, Lam. Encyc. i. 244 = Myrsine Manglilla.
Marmulana, C. Sm. in Tuckey, Congo, 252 = seq.
Marmulano, *Banks, ex Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 22.—Ins. Mader.
Mastichodendron, *Jacq. Coll.* 2. t. 17. f. 5.—Ind. occ.
Mastichodendrum, Balb. ex A. DC. in DC. Prod. viii. 99 = Myrsine laeta.
melanophleos, Linn. Mant. i. 48 = Myrsine melanophleos.
mexicanum, *Hemsl., Biol. Centr. Am. Bot.* ii. 296.—Mexic.
micranthum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 54 = elegans.
microcarpum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 17.—Ins. Amboin.

SIDEROXYLON :—
Mirmulana, Buch, Beschr. Canar. Ins. 193 = S. Marmulano.
mite, Linn. Syst. ed. XII. 178 = Myrsine mitis.
mite, Sims, Bot. Mag. t. 1858 (Ilicis sp.).—Afr. austr.
mite, Willd. ex A. DC. in DC. Prod. viii. 97 = Myrsine melanophleos.
moluccanum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 19.—Ins. Gebeh.
multiflorum, Lam. Illustr. ii. 43 (non Sapotacea).—Afr. austr.
myrsinoides, *Benth. & Hook. f. ex F. Muell. Census*, 92.—Austral.
nervosum, *Wall. Cat.* n. 4159.—Burma.
nigricans, Dum.-Cours. ex A. DC. in DC. Prod. viii. 186 = Myrsine mitis.
nigrum, Gaertn. f. Fruct. iii. 123 = Dipholis salicifolia.
nigrum, Soland. ex A. DC. in DC. Prod. viii. 186 = Myrsine salicifolia.
nitidum, *Blume, Bijdr.* 675.—Java.
nitidum, Lam. Illustr. ii. 42 = lucidum.
nodosum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 12.—Java.
obovatum, *Burck, l. c.* 11.—Java.
obovatum, *Gaertn. f. Fruct.* iii. 125.—Hab.?
obovatum, Griff. Notul. iv. 290 = ferrugineum.
obovatum, *Lam. Illustr.* ii. 42.—Ind. occ.
oppositifolium, Mill. Gard. Dict. ed. VIII. n. 2 = foetidissimum.
orichalcium, *Zipp. in Alg. Konst- en Letter-bode* (1829) I. 297, nomen; *et ex Mackl. Bijdr. Nat. Wetensch.* v. (1830) 178.—N. Guin.
oxyedrum, *Miq. Fl. Ind. Bat. Suppl.* 580.—Sumatra.
Pacurero, *Loefl. It.* 204.—Venezuela.
pallidum, Spreng. Syst. i. 666 = S. Mastichodendron.
parviflorum, *Boj. ex A. DC. in DC. Prod.* viii. 184.—Ins. Maurit.
parvifolium, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 124.—Ins. Philipp.
pauciflorum, *Jacq. Enum. Pl. Carib.* 15.—Ins. S. Doming.
pauciflorum, Lam. Illustr. ii. 42 = Dipholis salicifolia.
pentagonum, *A. DC. in DC. Prod.* viii. 185.—Ins. Dominic.
Pohlmanianum, *Benth. & Hook. f. ex F. Muell. Census*, 91.—Austral.
pomiforme, *A. DC. in DC. Prod.* viii. 182.—Jamaic.
puberulum, *A. DC. l. c.* 179.—Ins. Mascar.
punctatum, Lam. Illustr. ii. 42 = Myrsine floribunda.
reclinatum, Michx. Fl. Bor. Am. i. 122 = Bumelia tenax.
regium, Wall. Cat. n. 4156 = grandifolium, Dichopsis polyantha.
revolutum, *Baker, in Oliver, Fl. Trop. Afr.* iii. 503.—Afr. trop.
Richardi, *F. Muell. Census*, 92.—Austral.
rigidum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 18. —Ins. Celebes.
robustum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 56.—Bras.
rufum, *Mart. & Eichl. l. c.* 52.—Bras.
rugosum, *Roem. & Schult. Syst.* iv. 501.—Bras.
rugosum, Wall. Cat. n. 4158 = Wallichianum.
salicifolium, Gaertn. f. Fruct. iii. 124. t. 202 = Dipholis salicifolia.
Sapota, Jacq. Enum. Pl. Carib. 15 = Achras Sapota.
sericeum, Ait. Hort. Kew. ed. I. i. 262 = Lucuma sericea.
sericeum, Walt. Fl. Carol. 100 = Bumelia tenax.
spectabile, *Burck, in Ann. Jard. Buitenz.* v. (1885) 10.—Sumatra.
spinosum, Linn. Sp. Pl. 193 = Argania Sideroxylon.
spinosum, Willd. Sp. Pl. i. 1091 = Flacourtia sepiaria.
Spruceanum, *Mart. & Miq. in Mart. Fl. Bras.* vii. 53. —Bras.
strigosum, Willd. Enum. Hort. Berol. Suppl. 11 = Bumelia strigosa.
sundaicum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 9.—Ins. Sundaic.
tenax, Linn. Mant. i. 48 = Bumelia tenax.
tenax, Walt. Fl. Carol. 100 = Bumelia lanuginosa.
Teysmannianum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 19.—Malaya.
tomentosum, *Roxb. Pl. Corom.* i. 28. t. 28.—Ind. or.
tomentosum, Wall. Cat. n. 4153 = assamicum, Isonandra Wightiana.
undulatum, *Burck, in Ann. Jard. Buitenz.* v. (1885) 12.—Ins. Banca.

SIDEROXYLON :—
venulosum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 52.—Bras.
Vrieseanum, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 18.—Ins. Batjan.
Wallichianum, *G. Don, Gen. Syst.* iv. 28.—Ins. Penang.
Wightianum, Hook. & Arn. Bot. Beech. Voy. 196. t. 41 = Isonandra Wightiana.
Wightianum, Wall. Cat. n. 4154 = Isonandra lanceolata.
xerocarpum, *Benth. & Hook. f. ex F. Muell. Census*, 91.—Austral.

SIDEROXYLUM, Salisb. Prod. 138 (1796) = praec.

SIEBERA, J. Gay, in Mém. Soc. Hist. Nat. Par. iii. (1827) 343. *COMPOSITAE*, Benth. & Hook. f. ii. 464.
FLEUROTIA, Reichb. Nom. 90 (1841).
Gayana, Cass. in Dict. Sc. Nat. lix. 126 = pungens.
pungens, *J. Gay, in Mém. Soc. Hist. Nat. Par.* iii. (1827) 343.—As. Min.; Syria; Persia.

SIEBERA, Hoppe, in Flora, ii. (1819) 24 = **Arenaria**, Rupp. (Caryoph.).
argentea, Steud. Nom. ed. I. 777; ed. II. ii. 582 = A. aretioides.
cherlerioides, Hoppe, in Flora, ii. (1819) 24 = A. aretioides.

SIEBERA, Presl, in Oken, Isis, xxi. (1828) 275 = **Anredera**, Juss. (Chenop.).
baselloides, Presl, l. c. = A. spicata.

SIEBERA, Reichb. Consp. 145 (1828) = **Platysace**, Bunge (Umbell.).
Billardieri, Benth. Fl. Austral. iii. 356.—Austral.
cirrosa, Benth. l. c. 354 = P. cirrosa.
commutata, Benth. l. c. 355.—Austral.
compressa, Benth. l. c. 352.—Austral.
deflexa, Benth. l. c. 355.—Austral.
dissecta, Benth. l. c. 354.—Austral.
effusa, Benth. l. c. 355 = P. trachymenioides.
ericoides, Benth. l. c. 356.—Austral.
haplosciadia, Benth. l. c. 353.—Austral.
heterophylla, Benth. l. c. 354.—Austral.
juncea, Benth. l. c. 353.—Austral.
linearifolia, Benth. l. c. 356.—Austral.
Stephensonii, Benth. l. c. 357.—Austral.
tenuissima, Benth. l. c. 354.—Austral.
valida, Benth. l. c. 355 = P. valida.

SIEBERIA, Spreng. Anleit. ii. 282 (1817) = **Habenaria**, Willd. (Orchid.).
albida, Spreng. l. c. = H. albida.
bifolia, Spreng. l. c. = H. bifolia.
nigra, Spreng. l. c. = H. nigra.
viridis, Spreng. l. c. = H. viridis.

SIEBOLDIA, Heynh. Nom. ii. 644 (1846) = **Simethis**, Kunth (Liliac.).
bicolor, Heynh. l. c. = Sim. bicolor.

SIEBOLDIA, Hoffmgg. Preisv. (1842) Nachtr. 28.—Cf. Linnaea, xvi. (1842) Litt. 281 = **Clematis**, Dill. (Ranunc.).
bicolor, Hoffmgg. ex Heynh. Nom. ii. 665 = C. florida.
florida, Hoffmgg. ex Heynh. l. c. = C. florida.
japonica, Hoffmgg. Preisv. (1842) Nachtr. 28.—Cf. Linnaea, xvi. (1842) Litt. 282 = C. japonica.

SIEGESBECKIA, Linn. Hort. Cliff. 412 (1737); Gen. ed. I. 352 (1737) (Sigesbeckia); Gled. in Mém. Acad. Berl. v. (1749) 131. *COMPOSITAE*, Benth. & Hook. f. ii. 359.
LIMNOGENNETON, Sch. Bip. in Walp. Rep. vi. 146 (1846).
MINYRANTHES, Turcz. in Bull. Soc. Nat. Mosc. (1851) I. 180.
SCHKUHRIA, Moench, Meth. 566 (1794) (Sckuhria).
TRIMERANTHES, Cass. in Dict. Sc. Nat. lix. 237 (1829).
abyssinica, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 372.—Afr. trop.
agrestis, Poepp. & Endl. Nov. Gen. et Sp. iii. 45. t. 256 = flosculosa.
brachiata, Roxb. Hort. Beng. 62; Fl. Ind. iii. 439 = orientalis.
caspia, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) 285 = orientalis.
cordifolia, H. B. & K. Nov. Gen. et Sp. iv. 283 = orientalis.

SIEGESBECKIA:—
dichotoma, Schkr. Handb. iii. 144 = flosculosa.
droseroides, Sweet, Hort. Brit. ed. I. 235; et in Brit. Flow. Gard. Ser. I. t. 203 = orientalis.
emirnensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 188.—Madag.
flosculosa, *L'Hérit. Stirp. Nov.* 37. t. 19.—Peruv.
glutinosa, Wall. Cat. n. 3182 = orientalis.
gracilis, DC. Prod. v. 496 = orientalis.
iberica, Willd. Sp. Pl. iii. 2220 = orientalis.
jorullensis, H. B. & K. Nov. Gen. et Sp. iv. 284 = orientalis.
laciniata, Poir. Encyc. vii. 158 = Verbesina virginica.
Mandonii, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81; *et in Linnaea*, xxxiv. (1865–66) 534, nomen.—Bolivia.
melanolepsis, *Sch. Bip. ll. cc.*, nomen.—Bolivia.
microcephala, DC. Prod. v. 496 = orientalis.
occidentalis, Linn. Sp. Pl. 900 = Verbesina occidentalis.
orientalis, *Linn. l. c.*—Cosmop. trop.
portoriccensis, Bert. ex DC. Prod. v. 546 = Eleutheranthera ovata.
serrata, DC. l. c. 496 = orientalis.
triangularis, Cav. Ic. iii. 27 = orientalis.
trinervata, Moench, Meth. 582 = Verbesina occidentalis.

SIEGLINGIA, Bernh. Syst. Verz. Erf. 20, 44 (1800) = **Triodia**, R. Br. (1810) (Gramin.).
decumbens, Bernh. l. c. = T. decumbens.

SIEMSSENIA, Steetz, in Lehm. Pl. Preiss. i. 467 (1845) = **Podolepis**, Labill. (Compos.).
capillaris, Steetz, l. c. = P. Siemssenia.

SIERUELA, Rafin. Sylva Tellur. 112 (1838) = **Cleome**, Linn. (Cappar.).
viscosa, Rafin. l. c. = C. monophylla.

SIEVEKINGIA, Reichb. f. Beitr. Syst. Pfl. 3 (1871). Genus dubium *ORCHIDEARUM*.—Cf. Benth. & Hook. f. iii. 477.
suavis, *Reichb. f. l. c.*—Costa Rica.

SIEVERSIA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 397 = **Geum**, Linn. (Rosac.).
adnata, G. Don, Gen. Syst. ii. 527 = G. elatum.
albiflora, Hook. f. Fl. Antarct. 9. t. 7 = G. parviflorum.
andicola, Phil. in Linnaea, xxviii. (1856) 636 = G. Lechlerianum.
anemonoides, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 398 = G. anemonoides.
atlantica, G. Don, Gen. Syst. ii. 527 = G. sylvaticum.
calthifolia, D. Don, ex Torr. & Gray, Fl. N. Am. i. 425 = G. calthifolium.
caroliniana, G. Don, Gen. Syst. ii. 528 = G. album.
ciliata, G. Don, l. c. = G. triflorum.
congesta, R. Br. in Parry's 1st Voy. App. 276 = G. calthifolium.
dilatata, R. Br. l. c. = G. calthifolium.
dryadoides, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 125 = G. anemonoides.
elata, Royle, Illustr. Bot. Himal. 207. t. 39. t. 1 = G. elatum.
geoides, Spreng. Syst. ii. 543 = Coluria potentilloides.
glacialis, R. Br. in Parry's 1st Voy. App. 276 = G. glaciale.
humilis, R. Br. l. c. = G. Rossii.
humilis, Royle, Illustr. Bot. Himal. 207 = G. elatum.
karatavica, Trautv. in Act. Hort. Petrop. viii. (1883) 309 = G. karatavicum.
micropetala, Nym. Syll. 273 = G. micropetalum.
montana, R. Br. in Parry's 1st Voy. App. 276 = G. montanum.
paradoxa, D. Don, in Trans. Linn. Soc. xiv. (1825) 576. t. 22. f. 7–10 = Fallugia paradoxa.
Peckii, R. Br. in Parry's 1st Voy. App. 276 = G. radiatum.
radiata, R. Br. l. c. = G. radiatum.
reptans, R. Br. l. c. = G. reptans.
rosea, R. Grah. in Edinb. N. Phil. Journ. (Apr.–June, 1831) 193 = G. triflorum.
Rossii, R. Br. in Parry's 1st Voy. App. 276 = G. Rossii.

SIEVERSIA:—
rotundifolia, Cham. & Schlecht. in Linnaea, ii. (1827) 4 = Geum calthifolium.
silenifolia, Fisch. ex F. Schmidt, Reisen Amurl. 39 = G. anemonoides.
triflora, R. Br. in Parry's 1st Voy. App. 276 = G. triflorum.

SIGESBECKIA, Linn. Gen. ed. I. 352 (1737), etc. = **Siegesbeckia**, Linn. (Compos.).

SIGILLARIA, Rafin. in Journ. Phys. lxxxix. (1819) 261; et in Am. Monthly Mag. (1819) 357 = **Smilacina**, Desf. (Liliac.).
ciliata, Rafin. Atl. Journ. 151 = Sm. racemosa.
multiflora, Rafin. Med. Fl. ii. 83 = Sm. racemosa?

SIGMATOPHYLLUM, D. Dietr. Syn. Pl. ii. 1587 (1840) = **Stigmaphyllum**, A. Juss. (Malpig.).

SIGMATOSTALIX, Reichb. f. in Bot. Zeit. x. (1852) 769. *ORCHIDEAE*, Benth. & Hook. f. iii. 565.
aurosanguinea, *Reichb. f. in Linnaea*, xli. (1877) 68.—Peruv.
brachycion, *Griseb. in Goett. Abh.* xxiv. (1879) 336.—Reg. Argent.
dilatata, *Reichb. f. in Linnaea*, xli. (1877) 16.—N. Granat.
graminea, *Reichb. f. in Bot. Zeit.* x. (1852) 769.—Peruv.
malleifera, *Reichb. f. in Gard. Chron.* (1883) II. 360.—N. Granat.
picta, *Reichb. f. in Walp. Ann.* vi. 859.—Ecuador.
radicans, *Reichb. f. l. c.*—Bras.
reversa, *Reichb. f. in Linnaea*, xli. (1877) 103.—N. Granat.
Wallisii, *Reichb. f. l. c.*—N. Granat.

SIGMODOSTYLES, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 93 = **Rhynchosia**, Lour. (Legumin.).
villosa, Meissn. l. c. 94 = R. sigmodes.

SILAMNUS, Rafin. Sylva Tellur. 60 (1838) = **Cephalanthus**, Linn. (Rubiac.).
procumbens, Rafin. l. c. 61 = C. procumbens.

SILAUS, Bernh. Syst. Verz. Erf. 116 (1800). *UMBELLIFERAE*, Benth. & Hook. f. i. 910.
GASPARINIA, Bertol. Fl. Ital. iii. 614 (1837).
alatus, Link, Handb. i. 329 = Ligusticum alatum.
alpestris, *Bess. in Schult. Syst.* vi. p. xxxvi.—Ross. austr.; Sibiria.
Besseri, DC. Prod. iv. 161 = alpestris.
carvifolius, *C. A. Mey. Verz. Pfl. Cauc.* 125.—Banat.; As. Min.
flavescens, *Bernh. Syst. Verz. Erf.* 174.—Europ.
Gasparinii, Nym. Syll. 154 = carvifolius.
Gilibertii, *Bess. ex Eichw. Skizze*, 156.—Rossia.
gracilis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 301.—Turkestan.
longifolius, *Ledeb. Fl. Alt.* i. 323.—Sibir.
Matthioli, Koch, in Nov. Act. Nat. Cur. xii. I. (1824) = tenuifolius.
meoides, Griseb. Spicil. Fl. Rumel. i. 362 = Carum graecum.
peucedanoides, Boiss. Fl. Orient. ii. 974 = carvifolius.
peucedanoides, DC. Prod. iv. 161 = Eulophus peucedanoides.
pratensis, Bess. in Schult. Syst. vi. p. xxxvi = flavescens.
serotinus, Reichb. Fl. Germ. Excurs. 471 = tenuifolius.
tenuifolius, *DC. 3me Not. Pl. Rar. Jard. Genèv.* 23.—Hungar.
virescens, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 301 = carvifolius.

SILENANTHE, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 300 = **Silene**, Linn. (Caryoph.).
Zawadzkii, Griseb. & Schenk, l. c. = Silene Zawadzkii.

SILENE, Linn. Syst. ed. I (1735); Gen. ed. I. 132 (1737). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 147.
ATOCION, Adans. Fam. ii. 254 (1763).
BEHENANTHA, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 130.

SILENE:—
CARPOPHORA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 139. t. 32 (1862).
CHEIROPETALUM, E. Fries, Ind. Sem. Hort. Upsal. (1857); ex Urban, Addit. Ind. Sem. Hort. Berol. (1881) 11.
DIPLOGAMA, Opiz, Seznam, 38 (1852).
ELISANTHE, Reichb. Nom. 206 (1841).
HELIOSPERMA, Reichb. Nom. 206 (1841); et Ic. Fl. Germ. vi. t. 277 (1844).
KALERIA, Adans. Fam. ii. 506 (1763).
LEPTOSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344.
MUSCIPULA, Rupp. Fl. Jen. ed. Hall. 125 (1745).
OBERNA, Adans. Fam. ii. 255 (1763).
ONCERUM, Dulac, Fl. Hautes-Pyr. 255 (1867).
OTITES, Adans. Fam. ii. 255 (1763).
PETROCOMA, Rupr. Fl. Cauc. 200 (1869).
PETROSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344.
SILENANTHE, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 300.
VISCAGO, Hall. Enum. Stirp. Helv. i. 373 (1742).
acarpophora, Griseb. Spicil. Fl. Rumel. i. 172 = coniflora.
acaulis, *Linn. Sp. Pl.* ed. II. 603.—Reg. bor. et arct.
acuta, E. Mey. in Drège, Zwei Pfl. Docum. 222 = Burchellii.
acutifolia, *Link, ex Rohrb. Monog. Silene*, 141.—Lusitan.
adhaerens, Ehrenb. ex Rohrb. l. c. 162 = linearis.
adscendens, *Lag. Gen. et Sp. Nov.* 15.—Hispan.
adusta, Ball, in Journ. Linn. Soc. xvi. (1877) 360 = corrugata.
aegyptiaca, *Linn. f. Suppl.* 241 = Afr. bor.; Syria.
aethiopica, *Burm. f. Fl. Cap. Prod.* 13.—Afr. austr.
aetolica, *Heldr. in Atti Congr. Bot. Firenze*, 1874 (1876), 239.—Graecia.
affghanica, *Rohrb. in Ind. Sem. Hort. Berol.* (1867) *App.* ii. 5.—Afghan.
affinis, Boiss. Diagn. Ser. II. i. 72.—Sinai.
affinis, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 417 = micropetala.
agrestina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 6 = gallica.
agrigentina, Lojac. in Natural. Sicil. ii. (1883) 295 = fruticosa.
Agrostemma, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 8 = S. Psammitis.
alba, Muhl. ex Rohrb. Monog. Silene, 87 = nivea.
albescens, Boiss. Diagn. Ser. I. i. 28 = eriocalycina.
Allamanni, Otth, in DC. Prod. i. 379 = laciniata.
allophila, Clemente, ex Lag. Gen. et Sp. Nov. 15 = adscendens.
Almolae, *F. Gay, ex Coss. Notes Crit.* ii. 31.—Hispan.
alpestris, *Jacq. Fl. Austr.* i. 60. t. 96.—Europ. or.
alpestris, Willd. ex Nym. Consp. 92 = rupestris.
alpina, S. F. Gray, Nat. Arr. Brit. Pl. ii. 648 = acaulis.
alpina, Pall. Reise, iii. 34; Tabl. Taur. 51 = alpestris.
alpina, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 177; iii. (1852) 85; iv. (1853) 12 = seq.?
alpina, Thomas, Cat. Pl. Suiss. (1837) 45 = S. Cucubalus.
alsinoides, *Pourr. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 114.—Hispan.
alsinoides, Viv. Fl. Cors. Sp. Nov. 6 = Saponaria ocymoides.
altaica, Otth, in DC. Prod. i. 374 = turgida.
altaica, *Pers. Syn.* i. 497.—Sibir.
amana, *Boiss. Fl. Orient.* i. 634.—Syria.
ambigua, Cambess. ex Soyer-Willem. in Mém. Acad. Nancy, 1850 (1851), 158; Willk. Ic. et Desc. Pl. Hisp. i. 42. t. 28 = glauca.
ambigua, Salisb. Prod. 302 = S. Behen.
ambigua, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 576 = tenuis.
amblevana, Lej. Fl. Spa, i. 199 = nutans.
ammophila, *Boiss. & Heldr. Diagn.* Ser. I. viii. 82.—Ins. Creta.
amoena, Hill, Veg. Syst. xiii. 55 = maritima.
amoena, Hort. ex Fenzl, Adv. Bot. [2] in Del. Sem. Hort. Vindob. (1851) = cretica.
amoena, Hort. ex Fenzl, l. c. [9] = nocturna, antirrhina, gallica.
amoena, Huds. Fl. Angl. ed. I. 164 = S. Cucubalus.
amoena, Linn. Sp. Pl. 417 = tenuis.
amoena, Linn. l. c. ed. II. 596 = maritima.
amphorina, Pomel, Nouv. Mat. Fl. Atl. 330 = colorata.

SILENE :—
ampullata, *Boiss. Diagn.* Ser. I. i. 26.—As. Min.
amurensis, Pomel, Nouv. Mat. Fl. Atl. 209 = italica.
anastomosans, Lag. Gen. et Sp. Nov. 15 = picta.
andicola, Gill. in Hook. Bot. Misc. iii. (1833) 147 = Saponaria andicola.
andryalaefolia, *Pomel, Nouv. Mat. Fl. Atl.* 331.—Alger.
angistoma, Fenzl, ex Boiss. Fl. Orient. i. 622 = persica.
anglica, Linn. Sp. Pl. 416 = gallica.
angustifolia, Bieb. Fl. Taur. Cauc. i. 337 = petraea.
angustifolia, D. Dietr. Syn. Pl. ii. 1568 = clandestina.
angustifolia, Guss. Fl. Sic. Prod. i. 500 = S. Cucubalus.
angustifolia, Poir. Voy. Barb. ii. 164 = Tunica compressa.
angustissima, Bor. Fl. Centr. Fr. ed. III. ii. 95, in obs. = S. Cucubalus.
anisoloba, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 198.—As. centr.
annulata, *Hort. ex Fenzl, Adv. Bot.* [2, 9]; *in Del. Sem. Hort. Vindob.* (1851).—Hab.?
annulata, Thore, Chlor. Land. 173 = cretica.
anomala, Duf. ex Rohrb. Monog. Silene, 118 = apetala.
antelopum, Steud. Nom. ed. II. i. 450, in syn. = S. Cucubalus
antirrhina, Hort. Madr. ex Otth, in DC. Prod. i. 378 = rubella.
antirrhina, *Linn. Sp. Pl.* 419.—Am. bor.
apetala, Host, Fl. Austr. i. 532 = nocturna.
apetala, *Willd. Sp. Pl.* ii. 307.—Europ. austr.; Persia.
aprica. *Turcz. in Linnaea*, x. (1836) *Litt.* 102.—Sibir. or.; China.
arabica, Boiss. Fl. Orient. i. 593 = affinis.
araxina, *Trautv. in Act. Hort. Petrop.* iii. II. (1875) 278.—Armenia.
Arbuscula, Fenzl, ex Boiss. Fl. Orient. i. 612 = spergulifolia.
arenaria, Desf. Fl. Atlant. i. 354 = nicaeënsis.
arenaria, Ehrenb. ex Rohrb. Monog. Silene, 231 = Gypsophila hirsuta.
arenarioides, Desf. Fl. Atlant. i. 355 = clandestina.
arenicola, Presl, Fl. Sic. 153 = nicaeënsis.
arenosa, *C. Koch, in Linnaea*, xv. (1841) 712.—As. Min.; Oriens.
argaea, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 36.—Cappadoc.
argentea, *Ledeb. Fl. Ross.* i. 311.—As. Min.
argillosa, Munby, in Bull. Soc. Bot. Fr. xi. (1864) 44 = rubella.
arguta, Boiss. & Buhse, in Boiss. Fl. Orient. i. 642 = peduncularis.
arguta, *Fenzl, Pugill. Pl. Nov. Syr.* 8.—As. Min.
Aristidis, *Pomel, Nouv. Mat. Fl. Atl.* 330.—Alger.
armena, *Boiss. Diagn.* Ser. I. i. 29.—As. Min.
armeniaca, *Rohrb. in Ind. Sem. Hort. Berol.* (1867) *App.* ii. 5; *Monog. Silene*, 209.—Armenia.
Armeria, Asso, Syn. Stirp. Arag. 54 = rubella.
Armeria, Bieb. Fl. Taur. Cauc. i. 339 = compacta.
Armeria, *Linn. Sp. Pl.* 420.—Europ. austr.
Armeria, Pall. Ind. Taur. 51 = Saponaria glutinosa.
articulata, Kunze, in Flora, xxix. (1846) 696 = fuscata.
articulata, Viv. Pl. Lib. Spec. 23. t. 12. f. 1 = gallica.
arvatica, Lag. in Varied. Cienc. ii. IV. (1805) 212 = ciliata.
arvensis, Losc. Trat. Pl. Arag. 31 = S. Muscipula.
arvensis, Salisb. Prod. 301 = gallica.
aspera, A. Br. ex Aschers. Fl. Brandenb. i. 85 = Lychnis Coeli-rosa.
aspera, Hort. ex Rohrb. Monog. Silene, 118 = apetala.
asphaltica, Kotschy, ex Rohrb. l. c. 105 = setacea.
Asterias, *Griseb. Spicil. Fl. Rumel.* i. 168.—Banatus.
atlantica, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 307.—Alger.
atocia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = S. Atocion.
atocioides, *Boiss. Diagn.* Ser. I. v. 83.—As. Min.
Atocion, Murr. Syst. ed. XIII. 421 = aegyptiaca.
Aucheriana, *Boiss. Diagn.* Ser. I. i. 27.—Peruv.
auriculaefolia, Pomel, Nouv. Mat. Fl. Atl. 332 = mollissima.
auriculata, *Sibth. & Sm. Fl. Graec. Prod.* i. 292.—Ital.; Graec.
austriaca, Hort. ex Steud. Nom. ed. II. ii. 583 = alpestris.

SILENE :—
axillaris, Leavenw. in Am. Journ. Sc. Ser. I. vii. (1824) 62 = Cuphea viscosissima.
azurea, *Hort. Piln. ex Pasq. Cat. Orto Bot. Nap.* 95, nomen.—Hab.?
babylonica, Boiss. & Noé, Diagn. Ser. II. i. 71 = rubella.
baccifera, Roth, Tent. Fl. Germ. i. 192 = Cucubalus baccifer.
baicalensis, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 575 = tenuis.
Balansae, *Boiss. Diagn.* Ser. II. vi. 31.—Cappadoc.
Baldwinii, *Nutt. Gen. Am.* i. 288.—Am. bor.
barbata, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 820 = pendula.
bassanensis, Sternb. Reise Rhet. Alp. 40 = italica.
Bastardi, Bor. ex Lloyd, Fl. Ouest Herbor. (1876-77), 4; et in Bull. Soc. Dauph. (1880) 254 = maritima.
Baumgarteni, Schott & Kotschy, ex Rohrb. Monog. Silene, 145 = dinarica.
Behen, *Linn. Sp. Pl.* 418.—Graecia; As. Min.
Behen, Wirzén, Enum. Pl. Offic. Fenn. 36 = S. Cucubalus.
bella, Clark, Trav. i. 746 = compacta.
bellidifolia, *Eckl. & Zeyh. Enum.* 33.—Afr. austr.
bellidifolia, Jacq. Hort. Vindob. iii. 44. t. 81 = vespertina.
bellidifolia, Thunb. Prod. Pl. Cap. 81 = ornata.
bellidioides, Sond. in Harv. & Sond. Fl. Cap. i. 125 = bellidifolia.
Bergeri, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
Bergeri, Schott, ex Spreng. Syst. ii. 411 = echinata.
Biafrae, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 183.—Afr. trop.
biappendiculata, *Ehrh. ex Rohrb. in Bot. Zeit.* xxv. (1867) 82.—Aegypt.
bicolor, Frivald. ex Rohrb. Monog. Silene, 158 = rigidula.
bicolor, Moench, Meth. 706 = nocturna.
bicolor, Reichb. Pl. Cult. t. 72 = picta.
bicolor, Thore, Ess. Chlor. Land. 174 = portensis.
bifida, Retz. in Hoffm. Phytog. Blaetter, i. (1803) 37 = colorata.
biflora, Willd. ex Rohrb. Monog. Silene, 124 = ciliata.
bipartita, Desf. Fl. Atlant. i. 352 = colorata.
bipartita, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = vespertina.
bohemica, Jord. & Fourr. Brev. Pl. Nov. i. fasc. 6 = gallica.
Boissieri, *J. Gay, ex Coss. Notes Crit.* ii. 32.—Hispan.
Boissieri, Tineo, ex Rohrb. Monog. Silene, 115 = colorata.
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 330 = Hookeri.
Borderi, Jord. in Ann. Soc. Linn. Lyon, N. S. xii. (1866) 445 = elegans.
Boryi, *Boiss. Elench.* 19.—Hispan.
brachiata, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 94 = S. Cucubalus.
brachyantha, Schott, ex Rohrb. Monog. Silene, 213 = saxatilis.
brachyantha, Schur, ex Nym. Consp. 86 = S. Cucubalus.
brachycarpa, *Boiss. & Bal. Diagn.* Ser. II. vi. 29.—As. Min.
brachypetala, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = apetala.
brachypetala, Rob. & Cast. ex DC. Fl. Fr. Suppl. 607 = nocturna.
brachystachys, *Webb, Fragm. Fl. Aethiop.* 34.—Nubia.
bracteata, Boiss. in Tchihat. Asie Min. Bot. i. 200 = olympica.
bracteosa, Bertol. Fl. Ital. vi. 627 = S. Muscipula.
brahuica, *Boiss. Fl. Orient.* i. 615.—Afghan.
brevicaulis, *Boiss. Diagn.* Ser. I. i. 34.—Cataonia.
Bridgesii, *Rohrb. Ind. Sem. Hort. Berol.* (1867) *App.* ii. 5.—Calif.
Broussonetiana, Schott, ex Webb & Berth. Phyt. Canar. i. 141 = nutans.
bryoides, Jord. Pugill. Pl. Nov. 30 = acaulis.
buglossifolia, Sm. in Rees, Cyclop. xxxii. [n. 44] = italica.
bupleuroides, Heldr. ex Boiss. Diagn. Ser. I. viii. 90 = caramanica.
bupleuroides, Ledeb. ex Boiss. Fl. Orient. i. 639 = longiflora.
bupleuroides, *Linn. Sp. Pl.* 421.—As. Min.
bupleuroides, Schang. ex Bunge, Verz. Alt. 31 = tenuis.
Burchellii, *Otth, in DC. Prod.* i. 374.—Afr. austr.

SILENE :—
byzantina, Castagne, ex Boiss. Fl. Orient. i. 627 = S. Fabaria.
caesarea, *Boiss. & Bal. Diagn.* Ser. II. vi. 31.—As. Min.
caesia, Jan, Elench. 7; Sibth. & Sm. Fl. Graec. Prod. i. 294; Fl. Graec. v. t. 417 = italica.
caesia, Sieber, ex Boiss. Fl. Orient. i. 632 = Sieberi.
caespitosa, Salisb. Prod. 302 = acaulis.
caespitosa, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 262.—Reg. Caucas.
caffra, *Fenzl, in Drège, Zwei Pfl. Docum.* 222, nomen; *et ex C. Muell. in Walp. Ann.* vii. 276.—Afr. austr.
californica, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 83.—Calif.
calycina, *Presl, Fl. Sic.* 155.—Sicil.
calycina, Salzm. ex Rohrb. Monog. Silene, 99 = tridentata.
calyculata, C. Koch, in Linnaea, xix. (1847) 56 = dianthoides.
Cambessedesii, Boiss. & Reut. Pugill. Nov. Pl. (1852) 18 = littorea.
Campanula, Lapeyr. Hist. Abr. Pl. Pyr. 248 = elegans.
Campanula, *Pers. Syn.* i. 500.—Italia.
campanulata, Saut. in Flora, xiii. (1830) 485 = S. Cucubalus.
campanulata, *S. Wats. in Proc. Am. Acad.* x. (1875) 341.—Calif.
cana, Otth, in DC. Prod. i. 382 = italica.
canariensis, Willd. in Spreng. Neue Entdeck. iii. 60 (nom. fals.?) = corrugata?
candicans, *Čelak. in Oestr. Bot. Zeitschr.* xxvi. (1876) 321.—Syria.
Candollii, Jord. ex Guss. Enum. Pl. Inar. 37 = gallica.
canescens, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica, Lychnis sylvestris.
canescens, Tenore, Prod. Fl. Nap. p. xxv; Fl. Nap. i. 236. t. 39 = colorata.
canopica, Boiss. Fl. Orient. i. 596 = biappendiculata.
canopica, Delile, Fl. Aeg. Ill. 88; ex Spreng. Syst. ii. 406 = villosa.
cantabrica, Jord. & Fourn. Brev. Pl. Nov. fasc. i. 5 = gallica.
capensis, *Otth, in DC. Prod.* i. 379.—Afr. austr.
capillipes, *Boiss. & Heldr. Diagn.* Ser. II. v. 55.—Cilicia.
capitellata, *Boiss. Diagn.* Ser. I. i. 25.—As. Min.
cappadocica, *Boiss. & Heldr. l. c.* viii. 86.—As. Min.
caramanica, *Boiss. & Heldr. l. c.* 90.—As. Min.
cariensis, *Boiss. l. c.* i. 40.—Caria.
carnea, Salisb. Prod. 301 = fruticosa.
carnosa, Moench, Meth. Suppl. 306 = nocturna.
caroliniana, Walt. Fl. Carol. 142 (err. typ. 241) = pensylvanica.
caryophylloides, *Otth, in DC. Prod.* i. 369.—As. Min.
caspica, *Pers. Syn.* i. 497.—Reg. Caucas.; Turkest.
cassia, *Boiss. Diagn.* Ser. I. viii. 78.—Syria.
Catesbaei, Walt. Fl. Carol. 141 = virginica.
catholica, *Ait. Hort. Kew.* ed. II. iii. 85.—Europ. austr.
catholica, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 665 = italica.
caucasica, *Boiss. Fl. Orient.* i. 622.—Reg. Caucas.
Ceccariana, Boiss. & Heldr. Diagn. Ser. II. i. 78 = linifolia.
cephalantha, *Boiss. Fl. Orient.* i. 613.—Armenia.
cephallenia, *Heldr. Fl. Céph.* 26.—Ins. Cephal.
cephalonica, Schimp. & Wiest, ex Rohrb. Monog. Silene, 81 = S. Fabaria.
cerastiifolia, Boiss. Fl. Orient. i. 626 = odontopetala.
cerastoides, All. Fl. Pedem. ii. 79 = gallica.
cerastoides, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = disticha.
cerastoides, *Linn. Sp. Pl.* 417.—Reg. Mediterr.
cernua, Bartl. in Linnaea, vii. (1832) 623 = Burchellii.
cernua, Thunb. Prod. Pl. Cap. 81 = clandestina.
cespitosa, Stev. = caespitosa.
chaetodonta, *Boiss. Diagn.* Ser. I. i. 39.—Syria; Persia.
chamarensis, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 578 = tenuis.
cheiranthifolia, Salzm. ex Rohrb. Monog. Silene, 102 = obtusifolia.
cheiranthifolia, Schott, in Oken, Isis (1818), 821 = micropetala.

SILENE:—
cheiranthoides, Poir. Encyc. vii. 176 = pensylvanica.
chia, Spreng. Syst. ii. 409 = gallica.
chilensis, Cham. & Schlecht. ex Rohrb. Monog. Silene, 97 = gallica.
chirensis, *A. Rich. Tent. Fl. Abyss.* i. 44.—Abyss.
chloraefolia, *Sm. Pl. Ic. Ined.* i. 13. t. 13.—As. Min.; Persia.
chlorantha, *Ehrh. Beitr.* vii. 144.—Europ.; Tauria.
chlorantha, Hort. ex Fenzl, Adv. Bot. [7, 9]; in Del. Sem. Hort. Vindob. (1851) = disticha.
chloropetala, *Rupr. Fl. Caucas.* 195.—Reg. Caucas.
Choulettii, *Coss. in Bull. Soc. Bot. Fr.* ix. (1862) 169.—Alger.
chromodonta, *Boiss. & Reut. Diagn.* Ser. II. i. 71.—Thessalia.
ciliata, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = obtusiloba, vespertina.
ciliata, *Pourr. in Mém. Acad. Toul.* iii. (1788) 329.—Gallia.
ciliata, Urv. in Mém. Soc. Linn. Par. i. (1822) 304 = discolor.
ciliata, Willd. Sp. Pl. ii. 692 = colorata.
cinerea, *Desf. Fl. Atlant.* i. 335.—Alger.
cirtensis, Pomel, Nouv. Mat. Fl. Atl. 328 = cinerea.
cisplatensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 163.—Bras.
clandestina, Duby, Bot. Gall. 77 = cretica.
clandestina, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vendob. (1851) = antirrhina, apetala.
clandestina, *Jacq. Coll. Suppl.* 111.—Afr. austr.
clavaecarpa, Rochel, ex Rohrb. Monog. Silene, 220 = italica.
clavata, Moench, Meth. 706 = vallesia.
clavicarpa, Rochel, ex Reichb. Fl. Germ. Excurs. 821 = italica.
Clusii, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica.
Clusii, Schott, ex Steud. Nom. ed. II. ii. 583 = cerastoides.
coarctata, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = disticha.
coarctata, Lag. Gen. et Sp. Nov. 15 = cerastoides.
coccinea, Moench, Meth. Suppl. 306 = virginica.
Coeli-rosa, A. Br. in Flora, xxvi. (1843) 373 = Lychnis Coeli-rosa.
colorata, Frivald. in Flora, xviii. (1835) 334 = Friwaldskyana.
colorata, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = gallica.
colorata, *Poir. Voy. Barb.* ii. 163.—Reg. Mediterr.
colorata, Schousb. ex Hornem. Hort. Hafn. i. 412 = obtusifolia.
commelinaefolia, *Boiss. Diagn.* Ser. I. i. 35.—As. Min.; Persia.
commuta, 'Bert.' ex Nym. Consp. 88 = maritima.
commutata, Guss. Fl. Sic. Prod. i. 499 = S. Cucubalus.
commutata, Schur, in Verh. Siebenb. Ver. Naturw. (1859) 66 = nutans.
comollina, *Ser. Fl. Jard.* iii. (1849) 337.—Austria.
compacta, *Fisch. Hort. Gorenk.* ed. II. (1812) 60; *et ex Hornem. Hort. Hafn.* i. 417.—Europ. or.; As. Min.
congesta, Griseb. Spicil. Fl. Rumel. i. 175 = Roemeri.
congesta, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = disticha.
congesta, Panč. ex Rohrb. Monog. Silene, 199 = Sendtneri.
congesta, Schott, ex Steud. Nom. ed. II. ii. 583 = gallica, disticha.
congesta, *Sibth. & Sm. Fl. Graec. Prod.* i. 300.—Graecia.
conica, Geners. ex Rohrb. Monog. Silene, 97 = gallica.
conica, Hampe, in Flora, xx. (1837) 226 = subconica.
conica, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = conoidea.
conica, *Linn. Sp. Pl.* 418.—Europ.; Oriens.
conifera, Steud. Nom. ed. II. ii. 583, sphalm. = seq.
coniflora, *Nees, ex Otth, in DC. Prod.* i. 371.—Syria; Persia.
conoidea, Huds. Fl. Angl. ed. II. 189 = conica.
conoidea, *Linn. Sp. Pl.* 418.—Europ.; Oriens.
conringiaefolia, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 136, nomen.—Podolia.
Constantia, *Eckl. & Zeyh. Enum.* 32.—Afr. austr.

SILENE:—
Constantia, Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = clandestina.
cordata, Link, Handb. ii. 244, sphalm. = seq.
cordifolia, *All. Fl. Pedem.* ii. 82. t. 23. f. 3.—Pedemont.
coronaria, Clairv. ex Reichb. Fl. Germ. Excurs. 826 = Lychnis coronaria.
corrugata, *Ball, in Journ. Bot.* xi. (1873) 301.—Marocc.
corsica, DC. Fl. Fr. iv. 756 = succulenta.
corsica, Hort. Erf. ex Link, Enum. Hort. Berol. i. 428 = fruticosa.
corsica, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = nocturna.
corymbifera, Bertol. Fl. Ital. iv. 591 = S. Muscipula.
costata, Otth, in DC. Prod. i. 376 = clandestina.
Coulteriana, Otth, l. c. 368 = S. Cucubalus.
crassicaulis, *Willk. & Costa, in Linnaea*, xxx. (1859–60) 91.—Hispan.
crassifolia, *Linn. Sp. Pl.* 597.—Afr. austr.
crassifolia, Thore, Promen. Gascogne, 53, 54 = Thorei.
crassifolia, Thunb. Prod. Pl. Cap. 81 = Thunbergiana.
crassipes, *Fenzl, Pugill. Pl. Nov. Syr.* 8.—Syria.
cretacea, *Fisch. in Spreng. Syst.* ii. 405.—Sibir.
cretica, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = disticha, Cuphea procumbens.
cretica, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1070 = pendula.
cretica, *Linn. Sp. Pl.* 420.—Europ. austr.
crispa, Moench, Meth. 706 = pendula.
crispa, Poir. Encyc. vii. 162 = rubella.
crispata, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 315, in textu = S. Cucubalus.
croatica, Waldst. & Kit. ex Rohrb. Monog. Silene, 140 = Waldsteinii.
cruentata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 8 = gallica.
cryptantha, *Viv. Fl. Lib. Spec.* 23.—Lybia.
Cserei, Baumg. Enum. Stirp. Transs. iii. 345 = S. Fabaria.
Cucubalus, *Wibel, Prim. Fl. Werth.* 241.—Europ.; Afr. bor.; Reg. Himal.
cuneifolia, J. F. Gmel. Syst. 714 = gallica.
cuprea, Hort. ex Reichb. Ic. Fl. Germ. iii. 52 = nutans.
cuspidata, Leveillé, in Demid. Voy. Russ. Mér. ii. 234, sphalm. = Erysimum cuspidatum.
cylindriflora, Griseb. Spicil. Fl. Rumel. i. 171 = subconica.
cylindriflora, Kotschy, ex Rohrb. Monog. Silene, 90 = juvenalis.
cylindriflora, Otth, in DC. Prod. i. 371 = conoidea.
Czerei, Boiss. Fl. Orient. i. 629 = S. Fabaria.
dagestanica, *Rupr. Fl. Cauc.* 194.—Reg. Caucas.
dahurica, *Fisch. ex Sweet, Hort. Brit.* ed. II. 53.—Dahuria.
dalmatica, Scheele, in Flora, xxvi. (1843) 430 = Waldsteinii.
damascena, Boiss. & Gaill. Diagn. Ser. II. vi. 34 = palaestina.
dasyphylla, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 577 = tenuis.
decipiens, *Ball, in Journ. Linn. Soc.* xvi. (1877) 357.—Marocc.
decipiens, *Barc. in Anal. Soc. Esp. Hist. Nat.* viii. (1879) 340.—Ins. Majorc.
decumbens, Bivona, Cent. Sic. Pl. i. 75 = colorata.
decumbens, Salzm. ex Rohrb. Monog. Silene, 117 = glauca.
decumbens, Schreb. ex DC. Prod. i. 369 = procumbens.
delicatula, Bertol. Fl. Ital. iv. 624 = sedoides.
delicatula, *Boiss. Diagn.* Ser. I. i. 41.—Cilicia.
delphica, Boiss. & Heldr. Diagn. Ser. II. i. 73 = congesta.
densiflora, Urv. in Mém. Soc. Linn. Par. i. (1822) 303 = S. Otites.
depressa, Baumg. Enum. Stirp. Transs. iii. 826 = dinarica.
depressa, *Bieb. Fl. Taur. Cauc.* i. 336.—Reg. Caucas.
dianthifolia, *J. Gay, in Tchihat. Asie Min. Bot.* i. 193.—Cilicia.
dianthifolia, Otth, in DC. Prod. i. 373 = altaica.
dianthoides, *Pers. Syn.* i. 500.—As. Min.
dianthoides, Schimp. ex Rohrb. Monog. Silene, 178 = chloraefolia.
dichotoma, *Ehrh. Beitr.* vii. 144.—Europ.
dichotoma, Gilib. Fl. Lituan. ii. 169 = noctiflora.
diffusa, Forsk. ex Steud. Nom. ed. II. ii. 584 = Forskohlei.
diffusa, Otth, in DC. Prod. i. 373 = colorata.
Dilleniana, *Schott, ex Desf. Cat. Hort. Par.* ed. III. 264.—Gallia.

SILENE:—
Dillenii, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = cretica.
dinarica, *Spreng. Syst.* ii. 405.—Transsylv.
dioica, Fisch. & Mey. Ind. Sem. Hort. Petrop. vii. 6 = conoidea.
discolor, Retz. in Hoffm. Phytog. Blaetter, i. (1803) 38 = nutans.
discolor, *Sibth. & Sm. Fl. Graec. Prod.* i. 292; *Fl. Graec.* v. 8. t. 410.—As. Min.
distachya, Brot. Fl. Lusit. ii. 189 = colorata.
distans, *Otth, in DC. Prod.* i. 370.—Hab. ?
disticha, *Willd. Enum. Hort. Berol.* 476.—Reg. Mediterr. occ.
distycha, Webb, Iter Hisp. 63 = gallica.
diurna, Gren. & Godr. Fl. Fr. i. 217 = Lychnis dioica.
diurniflora, Kunze, in Linnaea, xvii. (1843) 578 = undulata, *Ait.*
divaricata, *Clemente, ex Lag. Gen. et Sp. Nov.* 15.—Hispan.
divaricata, Ehrenb. ex Rohrb. Monog. Silene, 178 = chloraefolia.
divaricata, Otth, in DC. Prod. i. 153 = fuscata.
divaricata, Sibth. & Sm. Fl. Graec. Prod. i. 293; Fl. Graec. v. 10. t. 414 = dichotoma.
diversifolia, Otth, in DC. Prod. i. 378 = rubella.
Dorrii, Kellogg, in Proc. Calif. Acad. iii. (1863) 44 = Menziesii.
Douglasii, *Hook. Fl. Bor. Am.* i. 88.—Am. bor. occ.
Douglasii, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = glauca.
Drummondii, A. Gray, in Proc. Am. Acad. viii. (1873) 377 = Scouleri.
Drummondii, *Hook. Fl. Bor. Am.* i. 89.—Am. bor. occ.
Drummondii, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = glauca.
dschabanica, C. Koch, ex Rohrb. Monog. Silene, 180 = armena.
dubia, Herb. ex Rohrb. l. c. 217 = nutans.
Duriaei, Spach, ex Rohrb. l. c. 116 = colorata.
Durieui, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = vespertina.
echinata, Jaub. & Spach, Illustr. i. 8 = squamigera.
echinata, Kotschy, ex Rohrb. Monog. Silene, 172 = crassipes.
echinata, *Otth, in DC. Prod.* i. 380.—Graecia.
echinosperma, *Boiss. & Heldr. Diagn.* Ser. I. viii. 78.—Graecia.
Echinus, *Boiss. & Heldr. l. c.* Ser. II. v. 56.—Pisidia.
Eckloniana, *Sond. ex Harv. & Sond. Fl. Cap.* i. 126.—Afr. austr.
effusa, *Fisch. Hort. Gorenk.* ed. II. (1812) 59, nomen.—Hab. ?
effusa, Otth, in DC. Prod. i. 370 = S. Otites.
Ehrenbergiana, *Rohrb. in Bot. Zeit.* xxv. (1867) 83.—Syria.
elata, Otth, in DC. Prod. i. 370 = chlorantha.
elegans, *Brot. Fl. Lusit.* ii. 185.—Lusitan.
Elisabethae, *Jan, ex Reichb. Fl. Germ. Excurs.* 822.—Ital.
elongata, Bell. Osserv. Bot. 60 = acaulis.
elongata, *Forsk. ex Steud. Nom.* ed. II. ii. 584 (Quid ?).—Aegypt.
Engelmanni, Rohrb. in Ind. Sem. Hort. Berol. (1867) App. ii. 5 = Bridgesii.
eranthema, Wibel, ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. 556 = stricta.
eremitica, *Boiss. Fl. Orient.* i. 644.—Persia.
eremophila, Bien. ex Boiss. Fl. Orient. i. 594 = colorata.
erigens, Willd. ex Rohrb. Monog. Silene, 155 = rubella.
eriocalycina, *Boiss. Diagn.* Ser. I. i. 28.—Mesopot.
eriophora, Willd. ex Spreng. Syst. ii. 408 = colorata.
eriophylla, *Jur. ex Nym. Consp.* 87.—Syria.
euxina, Rupr. Fl. Cauc. 184 = dichotoma.
exaltata, Frivald. in Flora, xviii. (1835) 333 = S. Otites.
exaltata, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
exaltata, Nees, ex Reichb. Fl. Germ. Excurs. 814, in obs. = colorata.
expansa, Hort. ex Rohrb. Monog. Silene, 211 = longipetala.
exscapa, All. Fl. Pedem. ii. 83 = acaulis.
exsudans, Boiss. & Heldr. Diagn. Ser. I. viii. 76 = microsperma.
fabaefolia, Schott, ex Rohrb. Monog. Silene, 81 = S. Fabaria.

SILENE :—

Fabaria, Bertol. Fl. Ital. iv. 627 = S. Cucubalus.
Fabaria, *Sibth. & Sm. Fl. Graec. Prod.* i. 293; *Fl. Graec.* v. 11. t. 415.—Sicil.; Creta.
falcata, *Sibth. & Sm. ll. cc.* i. 301; 25. t. 436.—Bithyn.
Falconeriana, *Royle, Illustr. Bot. Himal.* 79. t. 20. f. a.—Reg. Himal.
fallax, Willk. Pl. Haloph. 107 = ramosissima.
fasciculata, Boiss. & Heldr. in Boiss. Fl. Orient. i. 613 = brachycarpa.
Fenzlii, *Boiss. & Bal. Diagn.* Ser. II. vi. 30.—Cilicia.
filiformis, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) 549 = inaperta.
filiformis, Otth, in DC. Prod. i. 376 = picta.
fimbriata, Baldw. ex Ell. Sketch, i. 515 = Baldwinii.
fimbriata, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = S. Cucubalus.
fimbriata, *Sims, Bot. Mag.* t. 908.—Reg. Caucas.
firma, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 166 = aprica.
Fischeri, Steud. Nom. ed. II. ii. 584 = S. Cucubalus.
fissa, Salisb. Prod. 302 = Cucubalus baccifer.
fissipetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 371.—China.
fistulosa, G. Don, Gen. Syst. i. 405 = italica.
flammulaefolia, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 43.—Abyss.
flavescens, *Waldst. & Kit. Pl. Rar. Hung.* ii. 131. t. 175.—Hungar.; Banat.
foetida, *Link, ex Spreng. Syst.* ii. 406.—Lusitan.
foetida, *Pourr. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 114, nomen.—Hispan.
foliosa, *Maxim. Prim. Fl. Amur.* 53.—Manchuria.
foliosa, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) II. 559 = macrostyla.
fontana, Tenore, Fl. Nap. App. i. 26 = quadrifida.
Forskohlei, *Steud. Nom.* ed. II. ii. 584.—Aegypt.
Fortunei, *Vis. Ind. Sem. Hort. Patav.* (1847).—*Cf. Linnaea*, xxiv. (1851) 181.—China.
Fortunii, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
Friwaldskyana, *Hampe, in Flora*, xx. (1837) 227.—Thracia.
frutescens, Bieb. Beschr. Casp. 175 = altaica ?
fruticosa, Georgi, Besch. Russ. Reich. iv. 974 = altaica.
fruticosa, *Linn. Sp. Pl.* 417.—Graecia; Sicilia.
fruticosa, Salzm. ex Rohrb. Monog. Silene, 226 = velutina.
fruticulosa, Dietr. ex Otth, in DC. Prod. i. 381 = caspica.
fruticulosa, Fenzl, ex Rohrb. Monog. Silene, 128 = dianthifolia.
fruticulosa, *Sieber, ex Otth, in DC. Prod.* i. 376.—Ins. Creta.
fulgida, Kotschy, ex Rohrb. Monog. Silene, 103 = palaestina.
fuscata, *Link, ex Brot. Fl. Lusit.* ii. 187.—Reg. Mediterr.
gallica, Hort. Par. ex Tenore, Syll. Fl. Neap. 212 = reflexa.
gallica, *Linn. Sp. Pl.* 417.—Oriens.
Gallinyi, Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = squamigera.
Gallinyi, Heuff. ex Reichb. Fl. Germ. Excurs. 815 = trinervia.
Gebleriana, *Schrenk, Enum. Pl. Nov.* 91.—Soongar.
geminiflora, Willd. Enum. Hort. Berol. 472 = obtusifolia.
geniculata, *Lag. Gen. et Sp. Nov.* 14.—Gallia.
geniculata, Pourr. in Mém. Acad. Toul. iii. (1788) 328 = ciliata.
germana, J. Gay, ex Coss. Notes Crit. ii. 31 = Boissieri.
getula, *Pomel, Nouv. Mat. Fl. Atl.* 329.—Alger.
gibraltarica, Boiss. Elench. 20 = mollissima.
gigantea, *Linn. Sp. Pl.* 418.—Reg. Mediterr.
Giraldii, Guss. Enum. Pl. Inar. 36. t. 1. f. 2 = gallica.
glabra. Koch, Syn. Fl. Germ. ed. II. 1147, sphalm. = Sagina glabra.
glandulosa, Bertol. Fl. Ital. vi. 626 = gallica.
glandulosa, Hampe, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 302 = plutonica.
glareosa, Jord. Pugill. Pl. Nov. 31 = S. Cucubalus.
glauca, Hort. Par. ex Rohrb. Monog. Silene, 118 = apetala.
glauca, *Lag. Elench. Hort. Matr.* (1803); *ex ej. Gen. et Sp. Nov.* 15.—Reg. Mediterr. occ.
glauca, Salisb. Prod. 302 = S. Armeria.
glauca, Sm. ex Steud. Nom. ed. II. ii. 584 = S. Cucubalus.

SILENE :—

glauca, Zea, ex Poir. Encyc. Suppl. v. 153 = crassipes.
glaucifolia, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = apetala.
glaucifolia, Lag. in Varied. Cienc. ii. IV. (1805) 213 = Lychnis pyrenaica.
glaucophylla, *Lag. ex Willk. & Lange, Prod. Fl. Hisp.* iii. 641.—Hispan.
glomerata, C. Gay, Fl. Chil. i. 259 = gallica.
glutinosa, Duf. ex Cambess. in Mém. Mus. Par. xiv. (1827) 219 = rubella.
glutinosa, Pers. Syn. i. 498 = catholica.
glutinosa, Rohrb. Monog. Silene, 80, sphalm. = plutonica.
glutinosa, Zois, in Sitzb. Zool.-Bot. Ges. Wien, viii. (1858) 80 = eriophylla.
goniocaula, *Boiss. Diagn.* Ser. I. i. 31.—Persia.
gonocalyx, Boiss. l. c. 38 = crassipes.
gracilis, DC. Cat. Hort. Monsp. (1813) 145 = longicaulis.
gracilis, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = apetala.
gracillima, *Rohrb. in Linnaea*, xxxvi. (1869-70) 679.—Japon.
graeca, *Boiss. & Sprun. Diagn.* Ser. I. i. 36.—Graecia.
graeca, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = conoidea.
graeca, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 865 = pendula.
Graefferi, Guss. Pl. Rar. Sic. 170. t. 34 = ciliata.
graminea, Vis. ex Reichb. Ic. Fl. Germ. vi. 52 = vallesia.
graminifolia, Heldr. ex Boiss. Diagn. Ser. II. vi. 32 = Marschallii.
graminifolia, Ledeb. Fl. Alt. ii. 143 = turgida.
graminifolia, Otth, in DC. Prod. i. 368 = tenuis.
grammatocalyx, Fenzl, ex Boiss. Diagn. Ser. I. viii. 82 = macrodonta.
grata, Donnersm. Adumbr. Pl. 28 = reticulata.
graveolens, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 242 = ramosissima.
Grayi, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 291.—Calif.
Greggii, A. Gray, Pl. Wright. ii. 17 = laciniata.
Griffithii, *Boiss. Diagn.* Ser. II. i. 76.—Reg. Himal.
grisea, *Boiss. l. c.* Ser. I. viii. 88.—Syria.
Guicciardii, Boiss. & Heldr. Diagn. Ser. II. vi. 32 = Marschallii.
Guilelmi-Waldemari, Klotzsch, in Bot. Ergeb. Waldem. Reise, 140 = tenuis.
gymnotheca, *Panč. ex Nym. Consp.* 92 = fruticulosa.
Gysophila, Desf. Cat. Hort. Par. ed. III. 184 = repens.
gypsophiloides, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 45 = repens.
Haenkeana, Presl, Rel. Haenk. ii. 19 = gallica.
halopetala, Hort. ex Fenzl, Adv. Bot. [9]; Del. Sem. Hort. Vindob. (1851) = S. Saxifraga.
halophila, Link, ex Willk. & Lange, Prod. Fl. Hisp. iii. 650 = littorea.
halophila, *Willd. ex Spreng. Syst.* ii. 406.—Hispan.
Heldreichii, *Boiss. Diagn.* Ser. I. viii. 81.—As. Min.
Hellmanni, *Claus, in Beitr. Pfl. Russ. Reich.* viii. 289.—Rossia.
Heufleri, Hausm. ex Nym. Consp. 87 = eriophylla.
hirsuta, *Lag. in Varied. Cienc.* ii. IV. (1805) 212.—Reg. Mediterr. occ.
hirsuta, Poir. Voy. Barb. ii. 163 = vespertina.
hirsuta, Presl, ex Mert. & Koch, Deutschl. Fl. iii. 23 = gallica.
hirsuta, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1877) 356 = disticha.
hirsutissima, Otth, ex DC. Prod. i. 372 = hirsuta.
hirta, Willd. Hort. Berol. t. 23 = sericea.
hispanica, Jacq. Fragm. 47 = colorata.
hispanica, Otth, in DC. Prod. i. 369 = littorea.
hispida, Desf. Fl. Atlant. i. 348 = vespertina.
hispida, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica.
hispida, Hort. ex Willd. Enum. Hort. Berol. 476; et Poir. Encyc. Suppl. v. 151 = disticha.
hispida, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 358 = hirsuta ?
Hochstetteri, *Rohrb. in Bot. Zeit.* xxv. (1867) 81.—Abyss.
Hoefftiana, *Fisch. ex C. A. Mey. Verz. Pfl. Cauc.* 215.—Reg. Caucas.
Hohenackeri, Boiss. Diagn. Ser. II. i. 75 = Aucheriana.
holopetala, *Bunge, in Ledeb. Fl. Alt.* ii. 142.—Sibir. altaic.
Honigbergeri, *Fenzl, in Endl. & Fenzl, Sert. Cabul.* 3.—Afghan.

SILENE :—

Hookeri, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = glauca.
Hookeri, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 192.—Am. bor. occ.
Hornemanni, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = Lychnis dioica.
Hornemanni, *Steud. Nom.* ed. II. ii. 584.—Europ. austr.
humilis, *C. A. Mey. Verz. Pfl. Cauc.* 215.—Reg. Caucas.
Hussoni, *Boiss. Diagn.* Ser. I. viii. 76.—Arabia; Aegypt.
iberica, Bieb. Fl. Taur. Cauc. i. 335 = dichotoma.
iconia, Boiss. & Heldr. Diagn. Ser. I. viii. 85 = Marschallii.
ignobilis, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) 549 = S. Behen.
imbricata, *Desf. Fl. Atlant.* i. 349. t. 98.—Afr. bor.
inamoena, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = apetala.
inaperta, DC. Fl. Fr. iv. 750 = rubella.
inaperta, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = annulata, linicola.
inaperta, *Linn. Sp. Pl.* 419.—Reg. Mediterr. occ.
inaperta, Sibth. & Sm. Fl. Graec. v. 14 = multicaulis.
incarnata, Lodd. Bot. Cab. t. 41 = pensylvanica.
inclusa, Hornem. Hort. Hafn. i. 413 = nocturna.
incompta, A. Gray, in Proc. Am. Acad. vii. (1868) 330 = Bridgesii.
inconspicua, Roem. ex Steud. Nom. ed. II. ii. 585.—Cf. Rohrb. Monog. Silene, 228 (Quid ?).—Hab. ?
incurvifolia, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 391.—Sibir.
indica, Roxb. Hort. Beng. 34; Fl. Ind. ii. 446 = Lychnis indica.
inflata, Sm. Fl. Brit. 467 = S. Cucubalus.
infracta, Waldst. & Kit. Pl. Rar. Hung. iii. 237 = nutans.
insubrica, Gaud. Fl. Helvet. iii. 173 = italica.
insularis, *W. Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1885) 94.—Ins. Karpathos (prope Rhodon).
integrifolia, Hort. ex Steud. Nom. ed. II. ii. 585, sphalm. ? = seq.
integripetala, *Bory & Chaub. Nouv. Fl. Pélop.* 27.—Graecia.
intrusa, Wight & Arn. Prod. 42 = colorata.
involuta, *Forsk. Fl. Aegypt. Arab. Suppl.* 210. n. 47.—Turcia.
ispirensis, Boiss. & Huet, Diagn. Ser. II. v. 55 = lazica.
italica, *Pers. Syn.* i. 498.—Reg. Mediterr.; Oriens.
italica, Pollini, Fl. Veron. ii. 61 = nutans.
italica, Viv. Fl. Cors. 6 = Salzmanni.
japonica, *Rohrb. in Linnaea*, xxxvi. (1869-70) 689.—Japon.
jenisea, Poir. Encyc. Suppl. v. 154 = tenuis.
jenisea, *Steph. ex Bunge, Verz. Alt.* 30.—Sibir. altaic.
jenisensis, Willd. Enum. Hort. Berol. 473 = jenisea.
jucunda, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 8 = gallica.
juncea, Roth, Catalect. i. 54 = longiflora.
juncea, Sibth. & Sm. Fl. Graec. Prod. i. 295 = picta.
juniperina, Kunth, ex Rohrb. Monog. Silene, 193 = Urvillei.
juvenalis, *Delile, Ind. Sem. Hort. Monsp.* (1836) 28.—As. Min.
Kaulfussii, Spreng. Pugill. ii. 64 = rupestris.
Keiskei, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 77 = repens.
khasyana, *Rohrb. in Linnaea*, xxxvi. (1869-70) 258.—Reg. Himal.
Kitaibelii, Vis. Fl. Dalm. iii. 169 = Waldsteinii.
Kotschyi, Boiss. Diagn. Ser. I. i. 40 = microsperma.
Kowalenskyi, Stschégl. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 322 = arenosa.
Kremeri, *Soyer-Willem. & Godr. in Mém. Soc. Sc. Acad. Nancy*, 1850 (1851) 165.—Alger.
kunawarensis, *Royle, Illustr. Bot. Himal.* 79.—Reg. Himal.
Kuschakewiczii, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 246.—Turkestan.
lacera, Sims, Bot. Mag. t. 2255 = fimbriata.
laciniata, *Cav. Ic.* vi. 44. t. 564.—Am. bor. occ.
laconica, *Boiss. & Orph. Diagn.* Ser. II. vi. 34.—Graecia.
laeta, *A. Br. in Flora*, xxvi. (1843) 373; *Gren. & Godr. Fl. Fr.* i. 220.—Reg. Mediterr. occ.
laeta, Ehrenb. ex Rohrb. Monog. Silene, 150 = Reuteriana.

SILENE:—
laevigata, *Sibth. & Sm. Fl. Graec. Prod.* i. 295; *Fl. Graec.* v. 13. t. 418.—Ins. Cyprus.
Lagascae, Boiss. in Welw. It. Lusit. 490 = apetala.
lagenocalyx, *Fenzl, ex Boiss. Fl. Orient.* i. 587.—Persia.
lagunensis, C. Sm. ex Link, in Buch, Beschr. Canar. Ins. 154 = nutans.
lanceolata, *A. Gray, Bot. U. St. Expl. Exped.* i. 111.—Ins. Sandvic.
lanuginosa, Bertol. in Desv. Journ. Bot. ii. (1813) 76 = auriculata.
lasiantha, C. Koch, in Linnaea, xv. (1841) 712 = Marschallii.
lasiopetala, Fenzl, Pugill. Pl. Nov. Syr. 8 = Marschallii.
lasiostyla, Boiss. Diagn. Ser. I. viii. 79 = S. Psammitis.
latiflora, Link, in Schrad. Neue Journ. ii. 1. (1807) 99, sphalm. = *laxiflora*.
latifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 647 = S. Armeria.
latifolia, Hornem. Suppl. Hort. Hafn. 49 = Hornemanni.
latifolia, *Poir. Voy. Barb.* ii. 165.—Afr. bor.
laxa, *Boiss. & Kotschy, in Boiss Fl. Orient.* i. 638.—Armenia.
laxiflora, Brot. Fl. Lusit. ii. 188 = hirsuta.
lazica, *Boiss. Diagn.* Ser. I. i. 35.—Reg. Caucas.
Ledebourii, Heynh. Nom. ii. 667 = pygmaea.
legionensis, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = vespertina, nocturna.
legionensis, *Lag. Gen. et Sp. Nov.* 14.—Hispan.
Lemmonii, *S. Wats. in Proc. Am. Acad.* x. (1875) 342.—Calif.
lepta, J. Gay, in Boiss. Fl. Orient. i. 602 = microsperma.
leptoclada, *Boiss. Fl. Orient.* i. 647.—Lycia.
leptopetala, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 198.—Soongar.
Lerchenfeldiana, *Baumg. Enum. Stirp. Trans.* i. 398.—Transsylv.; Banat.
leucojifolia, Orsini, ex Rohrb. Monog. Silene, 229 = multicaulis.
leucophaea, Sibth. & Sm. Fl. Graec. Prod. i. 297; Fl. Graec. v. 17. t. 431 = nicaeënsis.
leucophylla, *Boiss. Diagn.* Ser. I. i. 29.—Arab.
leyseroides, Boiss. l. c. 41 = arenosa.
libanotica, *Boiss. l. c.* viii. 89.—Syria.
ligulata, *Viv. Fl. Lib. Spec.* 24. t. 12. f. 3.—Tripoli.
linaccola, Wibel, ex Spreng. Syst. ii. 412, sphalm. = seq.
linaeola, Wibel, ex Otth, in DC. Prod. i. 379, sphalm. = stricta.
linearifolia, Otth, l. c. 374 = petraea.
linearis, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 276.—Arab.
linearis, Sweet, Hort. Brit. ed. II. 51 = S. Cucubalus.
lineata, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 37.—Persia.
linicola, *C. C. Gmel. Fl. Bad.* iv. 304.—Europ.
linifolia, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = gallica.
linifolia, *Sibth. & Sm. Fl. Graec.* v. 23. t. 433.—Graecia.
linifolia, Willd. Enum. Hort. Berol. 473 = clandestina.
linoides, Otth, in DC. Prod. i. 384 = linifolia.
linoides, Soleir. ex Rohrb. Monog. Silene, 165 = inaperta.
lithophila, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 167.—Soongar.
litigiosa, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 198 = supina.
littoralis, Jord. Cat. Jard. Dijon (1848), 31 = gallica.
littoralis, Plan. Ens. Fl. Gallega, 123 = hirsuta.
littoralis, Pourr. in Mém. Acad. Toul. iii. (1788) 329 = nicaeënsis.
littorea, *Brot. Fl. Lusit.* ii. 186.—Hispan.
littorea, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = gallica.
livida, Wierzb. ex Rohrb. Monog. Silene, 219 = italica.
livida, Willd. Enum. Hort. Berol. 474 = nutans.
Loeflingiana, Bernh. ex Godr. Inflor. Gen. Silene, 34 = longicaulis.
Loiseleurii, *Gren. & Godr. Fl. Fr.* i. 220.—Gallia.
longicaulis, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = annulata.
longicaulis, *Pourr. ex Lag. Gen. et Sp. Nov.* 15.—Hispan.

SILENE:—
longicilia, Otth, in DC. Prod. i. 377 = nutans.
longiflora, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 14 = Boryi.
longiflora, *Ehrh. Beitr.* vii. 144.—Europ. or.; Oriens.
longifolia, D. Dietr. Syn. Pl. i. 1576 = nutans.
longipetala, *Vent. Jard. Cels*, 33. t. 33.—As. Min.; Syria.
lurida, Hort. ex Steud. Nom. ed. II. ii. 585 = nutans.
lusitanica, Linn. Sp. Pl. 416 = gallica.
lusitanica, Sibth. & Sm. Fl. Graec. Prod. i. 290 = vespertina.
Lyallii, *S. Wats. in Proc. Am. Acad.* x. (1875) 342.—Am. bor. occ.
lychnidea, *C. A. Mey. Verz. Pfl. Cauc.* 213.—Reg. Caucas.
lychnidiflora, Otth, in DC. Prod. i. 375 = cretica.
lycia, Boiss. Diagn. Ser. I. viii. 90 = rhynchocarpa.
lydia, *Boiss. l. c.* i. 37.—As. Min.
macroclada, Boiss. l. c. viii. 84 = S. Otites.
macrodonta, *Boiss. l. c.* i. 37.—Graecia.
macropetala, Spreng. Syst. ii. 414 = longipetala.
macrophylla, *Lag. Gen. et Sp. Nov.* 15.—Reg. Caucas.
macrophylla, Spreng. ex Rohrb. Monog. Silene, 81 = S. Fabaria.
macrorhiza, J. Gay, ex Rohrb. l. c. 142 = foetida.
macrosolen, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 44.—Abyss.
macrostemma, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 46.—Galatia.
macrostyla, *Maxim. Prim. Fl. Amur.* 54.—Reg. Amur.
makmeliana, Boiss. Diagn. Ser. I. viii. 89 = chloraefolia.
makmeliana, Boiss. & Buhse, ex Boiss. Fl. Orient. i. 639 = longiflora.
Mandralisci, Parl. in L'Occhio (1839), n. VIII. 61; ex Bertol. Fl. Ital. iv. 609 = sericea.
marginata, Kit. in Linnaea, xxxii. (1863) 536 = S. Cucubalus.
marginata, Schott, in Syll. Ratisb. i. (1824) 210 = gallica.
maritima, Host, Fl. Austr. i. 528 = S. Cucubalus.
maritima, *With. Bot. Arr. Brit. Pl.* ed. III. ii. 414.—Europ.
Marschallii, *C. A. Mey. Verz. Pfl. Cauc.* 214.—Reg. Caucas.; Oriens.
masmenaea, *Boiss. Diagn.* Ser. II. v. 57.—As. Min.
matutina, Presl, Fl. Sic. 149 = nocturna.
mauritânica, *Pomel, Nouv. Mat. Fl. Atl.* 328.—Algér.
mauritanica, Pott, ex Rohrb. Monog. Silene, 84 = maritima.
Maximowicziana, *Rohrb. in Linnaea*, xxxvi. (1869–70) 680.—Japon. (cult.).
megaspilaea, Boiss. & Heldr. ex Rohrb. Monog. Silene, 140 = multicaulis.
melandroides, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 233.—Hispan.
melandryiformis, Maxim. Prim. Fl. Amur. 54 = aprica.
melanophylla, Boiss. Diagn. Ser. I. i. 24 = dianthoides.
mellifera, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 9.—Hispan.
membrancea, Poir. Encyc. vii. 165 = noctiflora.
Menziesii, *Hook. Fl. Bor. Am.* i. 90. t. 30.—Am. bor. occ.
mexicana, Moç. & Sesse, ex Otth, in DC. Prod. i. 379 = laciniata.
Meyeri, Fenzl, ex Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 38 = depressa.
micrantha, Link, ex Lag. Gen. et Sp. Nov. 15 = micropetala.
microloba, Schott, Nym. & Kotschy, Analect. Bot. 59 = S. Cucubalus.
micropetala, DC. Cat. Hort. Monsp. (1813) 146 = cerastoides.
micropetala, *Lag. in Varied. Cienc.* ii. IV. (1805) 213.—Hispan.
micropetala, Roem. ex Schrank, Pl. Rar. Hort. Monac. i. t. 39; et in Syll. Ratisb. i. (1824) 212 = disticha.
micropetala, Tenore, Fl. Nap. v. 344 = gallica.
microphylla, *Boiss. Diagn.* Ser. I. i. 33.—Persia.
microphylla, DC. Prod. i. 372, sphalm. = *micropetala*, Roem.
microsperma, Fenzl, in Russeg. Reise, ii. 916 = linearis.

SILENE:—
microsperma, *Fenzl, Pugill. Pl. Nov. Syr.* 9.—Syria.
miniata, Rafin. Ann. Nat. 12 (Quid ?).—Am. bor.
minutiflora, *Jord. & Fourr. Brev. Pl. Nov.* fasc. i. 4.—Hispan.
mixta, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = S. Armeria, gallica.
Mociniana, DC. Prod. i. 382 = laciniata.
modesta, Boiss. & Blanche, Diagn. Ser. II. vi. 33 = chaetodonta.
modesta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 5 = gallica.
mogadorensis, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 55 = corrugata ?
mollis, Hornem. Hort. Hafn. i. 418 = flavescens.
mollissima, Frivald. ex Rohrb. Monog. Silene, 204 = gigantea.
mollissima, Hort. ex Fenzl, Adv. Bot. [9]; in Del. Sem. Hort. Vindob. (1851) = annulata, cretica.
mollissima, Otth, in DC. Prod. i. 382 = italica.
mollissima, Pers. Syn. i. 498 = velutina.
mollissima, *Sibth. & Sm. Fl. Graec. Prod.* i. 298.—Reg. Mediterr. occ.
molopica, Fenzl, ex Boiss. Fl. Orient. i. 578 = coniflora.
monachorum, Vis. ex Nym. Consp. 87 = quadridentata.
monantha, S. Wats. in Proc. Am. Acad. x. (1875) 340 = Douglasii.
montana, Arrond. in Bull. Soc. Polym. Morb. (1863) 58; et in Cat. Pl. Morb. 15 = maritima.
montana, *S. Wats. in Proc. Am. Acad.* x. (1875) 343.—Am. bor. occ.
Montbretiana, *Boiss. Diagn.* Ser. I. i. 26.—Armenia.
Moorcroftiana, *Wall. Cat.* n. 626.—Reg. Himal.
Morisii, *Todaro, Ind. Sem. Hort. Panorm.* (1872) 39; *et in Nuov. Giorn. Bot. Ital.* v. (1873) 158.—Italia.
multicaulis, Boiss. Fl. Orient. i. 651 = clavata.
multicaulis, Durand. in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855); ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 108 = Bridgesii.
multicaulis, *Guss. Pl. Rar.* 172. t. 35.—Europ. austr.
multicaulis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 192, 675 = Douglasii.
multifida, Edgew. in Trans. Linn. Soc. xx. (1846) 34 = Griffithii.
multifida, Rohrb. Monog. Silene, 88 = fimbriata.
multiflora, *Pers. Syn.* i. 497.—Hungar.; Sibir.
Mundiana, *Eckl. & Zeyh. Enum.* 32.—Afr. austr.
muricata, Host, ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = squamigera.
Muscipula, *Linn. Sp. Pl.* 420.—Reg. Mediterr.
Muscipula, Otth, in DC. Prod. i. 379 = stricta.
Mussini, Hornem. Hort. Hafn. i. 415 = catholica.
mutabilis, Linn. Cent. Pl. ii. 16; Amoen. Acad. iv. 314 = nocturna.
mutabilis, *Siev. in Pall. N. Nord. Beitr.* vii. 229.—Rossia.
mutica, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = disticha.
myloptera, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 8 = gallica.
nana, Cambess. ex Nym. Consp. 96 = littorea.
nana, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 169.—Persia; Turkest.
nardifolia, Boiss. & Huet, Diagn. Ser. II. v. 56 = caryophylloides.
natalensis, Buching. ex Krauss, in Flora, xxvii. (1844) 292 = crassifolia.
nebrodensis, Jan, ex Rohrb. Monog. Silene, 219 = italica.
neglecta, Tenore, Prod. Fl. Neap. App. v. 13; et in Linnaea, iii. (1828) Litt. 99 = reflexa.
nemoralis, Waldst. & Kit. Pl. Rar. Hung. iii. 271. t. 249 = italica.
Nentvichii, Frivald. ex Rohrb. Monog. Silene, 198 = Roemeri.
nevadensis, *Boiss. Voy. Bot. Espagne*, ii. 721.—Hispan.
nicaeënsis, *All. Misc. Taur.* v. (1770–73) 88 (niceensis); *Fl. Pedem.* ii. 81.—Reg. Mediterr.
nicaeënsis, Cham. & Schlecht. in Linnaea, i. (1826) 41 = verecunda.
nicaeënsis, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica.
Niederi, *Heldr. in Boiss. Diagn.* Ser. II. vi. 32.—Graecia.

SILENE:—
nitida, Lag. Gen. et Sp. Nov. 15 = fruticosa.
nivalis, *Rohrb. Monog. Silene*, 142.—Transsylv.
nivea, *Muhl. Cat. ex DC. Prod.* i. 377.—Am. bor.
nocteolens, *Webb & Berth. Phyt. Canar.* i. 141.—Ins. Canar.
noctiflora, *Linn. Sp. Pl.* 419.—Europ.; Oriens.
nocturna, *Linn. l. c.* 416.—Reg. Mediterr.
nocturna, Pall. Tabl. Taur. 50; et ex Georgi, Beschr. Russ. Reich. iv. 974 = dichotoma.
nocturna, Schimp. ex Rohrb. Monog. Silene, 119 = colorata.
nodulosa, *Boiss. Diagn.* Ser. I. i. 31.—Persia.
nodulosa, Viv. Fl. Cors. Sp. Nov. 6 = pauciflora.
norwegica, Pers. Syn. i. 500 = acaulis.
Notarisii, Cesati, in Bibl. Ital. xci. (1838) 346 = multicaulis.
novissima, Hort. ex Steud. Nom. ed. II. ii. 586 (Quid ?).—Hab. ?
nubigena, *Phil. in Anal. Univ. Chil.* (1862) II. 378.—Chili.
nudicaulis, Ehrenb. ex Rohrb. Monog. Silene, 105 = biappendiculata.
nutans, *Linn. Sp. Pl.* 417.—Europ.; As. bor.
nyctantha, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = clandestina.
nyctantha, Willd. Enum. Hort. Berol. 472 = nocturna.
oblongifolia, Otth, in DC. Prod. i. 369 = rubella.
obtusifolia, *Willd. Enum. Hort. Berol.* 473.—Reg. Mediterr. occ.
occidentalis, *S. Wats. in Proc. Am. Acad.* x. (1875) 343.—Calif.
occitanica, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 6 = gallica.
ocimoides, Desf. Cat. Hort. Par. ed. III. 184 = italica.
odontopetala, *Fenzl, Pugill. Pl. Nov. Syr.* 9.—As. Min.; Syria.
odoratissima, *Bunge, in Ledeb. Fl. Alt.* ii. 148.—Turkestan.; Sibir.
Oldhamiana, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 187 = aprica.
oleracea, Bor. Fl. Centr. Fr. ed. III. ii. 95 = S. Cucubalus.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 14.—Turkestan.
Olgae, *Rohrb. in Linnaea*, xxxvi. (1869–70) 687.—Manchuria.
oligantha, Bess. ex Ledeb. Fl. Ross. i. 321 = depressa.
oligantha, Boiss. & Heldr. Diagn. Ser. II. i. 75 = radicosa.
Oliveriana, Boiss. ex Rohrb. Monog. Silene, 105 = biappendiculata.
Oliveriana, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica.
Oliveriana, Otth, in DC. Prod. i. 373 = colorata.
olympica, *Boiss. Diagn.* Ser. I. i. 24.—As. Min.
olympica, Panč. ex Rohrb. Monog. Silene, 139 = fruticulosa.
oppositifolia, S. G. Gmel. ex J. F. Gmel. Syst. 717 = conica.
orchidea, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = nocturna.
orchidea, Linn. f. Suppl. 241 = aegyptiaca.
Oreades, Boiss. & Heldr. Diagn. Ser. I. viii. 92 = S. Saxifraga.
oregana, *S. Wats. in Proc. Am. Acad.* x. (1875) 343.—Am. bor. occ.
oreophila, *Boiss. Fl. Orient.* i. 617.—Cappadoc.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 10.—Oriens.
ornata, *Ait. Hort. Kew.* ed. I. ii. 96.—Afr. austr.
Orphanidis, *Boiss. Fl. Orient.* i. 651.—Graecia.
Orsiniana, Reichb. ex Cesati, Stirp. Ital. i. 8 = multicaulis.
Otites, *Sm. Fl. Brit.* ii. 469; *Pers. Syn.* i. 497.—Europ.; Oriens.
otitis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 134 = praec.
Ouensae, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 55, nomen.—Afr. bor.
ovata, *Pursh, Fl. Am. Sept.* i. 316.—Am. bor.
oxyodonta, *C. & W. Barbey, Herbor. au Levant*, 121. t. 11.—Palaest.
pachyphylla, Hort. ex Fenzl, Adv. Bot. [2, 7]; in Del. Sem. Hort. Vindob. (1851) = gallica, obtusifolia.
palaestina, *Boiss. Diagn.* Ser. I. viii. 80.—Palaestin.
palinotricha, *Fenzl, ex Boiss. Fl. Orient.* i. 621.—Persia.
pallida, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 12 = longiflora.

SILENE:—
pallidiflora, Link, ex Rohrb. Monog. Silene, 118 = apetala.
Palmeri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 124.—Calif.
pamphylica, Boiss. & Heldr. Diagn. Ser. I. viii. 81 = macrodonta.
paniculata, Ehrenb. ex Rohrb. Monog. Silene, 195 = stenobotrys.
paniculata, Otth, in DC. Prod. i. 382 = italica.
panormitana, Parl. ex Bertol. Fl. Ital. iv. 610 = cretica.
papillosa, *Boiss. Diagn.* Ser. I. i. 39.—As. Min.
paradoxa, Lapeyr. Hist. Abr. Pl. Pyr. 246 = nutans.
paradoxa, *Linn. Sp. Pl.* ed. II. 1673.—Europ. austr.
Parishii, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 366.—Calif.
parnassica, Boiss. & Sprun. Diagn. Ser. I. viii. 91 = fruticulosa.
parviflora, *Boiss. Fl. Orient.* i. 607.—Reg. Caucas.
parviflora, Hampe, in Flora, xx. (1837) 233 = Roemeri.
parviflora, Moench, Meth. 708 = cretica.
parviflora, Pers. Syn. i. 497 = S. Otites.
parviflora, Turcz. ex Bess. in Flora, xvii. (1831) I. Beibl. 8 = jenisea.
parviflora, Zea, ex Poir. Encyc. Suppl. v. 150 = littorea.
parvifolia, Otth, in DC. Prod. i. 376 = disticha.
parvula, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 7 = gallica.
patens, Peete, in Engl. Bot. Suppl. t. 2748 = italica.
patula, Desf. Fl. Atlant. i. 356 = italica.
patula, Lag. in Anal. Cienc. Nat. (1802); ex Duf. in Bull. Soc. Bot. Fr. vii. (1860) 242 = rubella.
pauciflora, *Kit. in Linnaea*, xxxii. (1863) 537 (Quid ?).—Europ.
pauciflora, *Ledeb. Fl. Ross.* i. 306.—Ross. arct.; Sibir.
pauciflora, *Salzm. ex Otth, in DC. Prod.* i. 382.—Ins. Corsic.; Sardoa.
pauciflora, Ucria, in Roem. Arch. i. I. (1796) 68 = viridiflora.
pectinata, *S. Wats. in Proc. Am. Acad.* x. (1875) 344.—Calif.
pedicellata, Boiss. Fl. Orient. i. 608 = S. Otites.
pedicellata, Poir. Encyc. Suppl. v. 150 = italica.
peduncularis, *Boiss. Diagn.* Ser. I. i. 30.—Persia.
pelidna, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = nocturna, Lychnis dioica.
pelidna, Reichb. Fl. Germ. Excurs. 821; Ic. Fl. Germ. iii. 53 = nutans.
pendula, *Linn. Sp. Pl.* 418.—Reg. Mediterr.
pendula, Salzm. ex Willk. Ic. et Desc. Pl. Hisp. i. 49 = littorea.
pensylvanica, *Michx. Fl. Bor. Am.* i. 272.—Am. bor.
pentelica, Boiss. Diagn. Ser. I. viii. 74 = sedoides.
perfoliata, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = S. Schafta.
perfoliata, Otth, in DC. Prod. i. 384 = chloraefolia.
permixta, Jord. Pugill. Pl. Nov. 32 = nocturna.
pernoctans, Hort. ex Fenzl, Adv. Nov. [2]; in Del. Sem. Hort. Vindob. (1851) = noctiflora, S. Schafta.
pernoctans, Link, ex Spreng. Syst. ii. 408 = colorata.
persica, *Boiss. Diagn.* Ser. I. i. 27.—Persia.
Persoonii, Schott, ex Reichb. Fl. Germ. Excur. 823 = S. Cucubalus.
Pestalozzae, Boiss. Diagn. Ser. I. viii. 87 = cappadocica.
petraea, *Adam, in Weber & Mohr, Beitr.* i. (1805) 58.—Reg. Caucas.
petraea, Ledeb. & Schrank, in Syll. Ratisb. i. (1824) 211 = pygmaea.
petraea, Waldst. & Kit. Pl. Rar. Hung. ii. 178. t. 164 = S. Saxifraga.
pharnaceifolia, *Fenzl, Pugill. Pl. Nov. Syr.* 9.—Cilicia.
phrygia, *Boiss. Fl. Orient.* i. 644.—As. Min.
physalodes, *Boiss. Diagn.* Ser. I. viii. 83.—Palestin.
physocalyx, Ledeb. Fl. Ross. i. 321 = odontopetala.
picta, Desf. Tabl. ed. I. 159 = portensis.
picta, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = nocturna, vespertina.
picta, *Pers. Syn.* i. 498.—Reg. Mediterr.
picta, Reichb. Fl. Germ. Excurs. 816 = Reichenbachii.
pilosa, Hort. ex Willd. Enum. Hort. Berol. 476 = disticha.
pilosa, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 652 = colorata.
pilosa, Spreng. Syst. ii. 411 = italica.
pilosellaefolia, Cham. & Schlecht. in Linnaea, i. (1826) 41 = Burchellii.

SILENE:—
piloselloides, G. Don, Gen. Syst. i. 404 = Burchellii.
pinetorum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 75.—Ins. Creta.
pinguis, Vahl, ex Hornem. Suppl. Hort. Hafn. 49 = nocturna.
pittodes, Bien. ex Boiss. Fl. Orient. i. 606 = chaetodonta.
platyota, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 366.—Calif.
platypetala, Otth, in DC. Prod. i. 383 = pensylvanica.
plicata, S. Wats. in Proc. Am. Acad. xvii. (1882) 366 = Thurberi.
plutonica, *Naud. ex C. Gay, Fl. Chil.* i. 258.—Chili.
Pollinii, Nym. Syll. 232 = nutans.
polygama, Stokes, Bot. Mat. Med. ii. 536 = S. Saxifraga.
polygonoides, *Pers. Syn.* i. 500.—Ins. Naxos.
polyphylla, Baumg. Enum. Stirp. Transs. i. 397 = nutans.
polyphylla, Bieb. Fl. Taur. Cauc. i. 338 = spergulaefolia.
polyphylla, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
polyphylla, Linn. Sp. Pl. 420 = portensis.
polyphylla, Vill. Hist. Pl. Dauph. iii. 609 = inaperta.
polytricoides, *Zumag. Fl. Pedem.* ii. 269; *et Cur. Post.* p. iv.—Pedem.
pompeiopolitana, *J. Gay, ex Boiss. Fl. Orient.* i. 595.—Cilicia.
porrigens, Gouan, ex Linn. Syst. ed. XII. 230 = Gypsophila porrigens.
portensis, Frivald. ex Rohrb. Monog. Silene, 167 = cretica.
portensis, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = gallica.
portensis, *Linn. Sp. Pl.* 420.—Hispan.
Pourretii, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
Pourretii, Poir. Encyc. Suppl. v. 151 = ciliata.
praecox, Hort. Par. ex Rohrb. Monog. Silene, 118 = apetala.
pratensis, Gren. & Godr. Fl. Fr. i. 216 = Lychnis dioica.
primulaeflora, *Eckl. & Zeyh. Enum.* 32.—Afr. austr.
procumbens, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = vespertina.
procumbens, *Murr. in Comm. Gotting.* 1784, 1785 (1786), 83. t. 2.—Sibir.
prostrata, Willd. ex Spreng. Syst. ii. 406 = halophila.
pruinosa, Boiss. Diagn. Ser. I. i. 23 = supina.
psammica, Boiss. ex Rohrb. Monog. Silene, 110 = discolor.
Psammitis, *Link, in Spreng. Nov. Prov.* 39.—Afr. bor.
Pseudo-atocion, *Desf. Fl. Atlant.* i. 353.—Afr. bor.
Pseudo-atocion, Guss. Pl. Rar. 180 = fuscata.
Pseudo-behen, Boiss. Diagn. Ser. I. i. 36 = S. Behen.
Pseudo-behen, Schur, ex Nym. Consp. 88 = S. Cucubalus.
Pseudo-linum, Ram. ex DC. Fl. Fr. iv. 750 = rubella.
Pseudo-otites, Bess. ex Reichb. Fl. Germ. Excurs. 819 = S. Otites.
pteropleura, Boiss. & Reut. Pugill. Pl. Nov. 18 = stricta.
ptolemaica, Ehrenb. ex Rohrb. Monog. Silene, 105 = biappendiculata.
puberula, Bertol. in Nov. Comm. Acad. Bonon. vi. (1842) 221 = colorata.
puberula, *Boiss. Fl. Orient.* i. 636.—Armenia.
puberula, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 94 = S. Cucubalus.
pubescens, Loisel. Fl. Gall. 727 = sericea.
pudibunda, Hoffmgg. ex Reichb. Fl. Germ. Excurs. 817 = quadridentata.
pulchra, Torr. & Gray, Fl. N. Am. i. 675 = laciniata.
pumila, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
Pumilio, *Wulf. in Jacq. Fl. Austr.* v. 26; *Sturm, Deutschl. Fl.* Heft 22.—Europ. alp.
punctata, Bubani, ex Nym. Consp. 92 = elegans.
pungens, *Boiss. Diagn.* Ser. I. i. 32.—As. Min.
purpurea, Knowl. & Westc. Flor. Cab. i. (1837) 87 = compacta.
pusilla, Waldst. & Kit. Pl. Rar. Hung. iii. 235 = quadridentata.
pygmaea, *Adam, in Weber & Mohr, Beitr.* i. (1805) 59.—Europ.
pygmaea, Herd. Ind. Sem. Hort. Petrop. (1858) 44 = gallica.

SILENE:—
pygmaea, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = noctiflora.
pygmaea, Less. ex Ledeb. Fl. Ross. i. 323 = caspica.
pyrenaica, Pourr. ex Rohrb. Monog. Silene, 115 = colorata.
pyriformis, Dur. ex Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 221 = glauca.
quadridentata, DC. Fl. Fr. iv. 748 = quadrifida.
quadridentata, *Pers. Syn.* i. 500.—Europ. austr.
quadrifida, Georgi, Reise, i. 215.—Sibir. baical.
quadrifida, *Linn. Syst.* ed. X. 1032.—Europ.
quadrifida, Otth, in DC. Prod. i. 378 = nutans.
quadriloba, Turcz. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 167 = viscosa.
quinquevulnera, Linn. Sp. Pl. 416 = gallica.
racemosa, Otth, in DC. Prod. i. 384 = dichotoma.
Raddeana, *Trautv. in Act. Hort. Petrop.* ii. (1873) 472.—Armenia.
radians, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 168 = odoratissima.
radicosa, *Boiss. & Heldr. Diagn.* Ser. I. vi. 24.—Graecia.
Ramburiana, Webb, Iter Hisp. 64 = Boryi.
ramosa, G. Don, Gen. Syst. i. 409 = ramosissima.
ramosa, Otth, in DC. Prod. i. 380 = sedoides.
ramosissima, Boiss. Voy. Bot. Espagne, ii. 93. t. 24 = Boissieri.
ramosissima, *Desf. Fl. Atlant.* i. 354.—Afr. bor.
ramosissima, Sibth. & Sm. Fl. Graec. Prod. i. 599 = sedoides.
ramosissima, Willk. Pl. Haloph. 108 = divaricata.
recta, Bartl. in Linnaea, vii. (1832) 623 = clandestina.
reflexa, *Ait. Hort. Kew.* ed. II. iii. 86.—Europ. austr.
reflexa, Hort. Taur. ex Rohrb. Monog. Silene, 94 = dichotoma.
reflexa, Moench, Meth. 705 = gallica.
regia, *Sims, Bot. Mag.* t. 1724.—Am. bor.
Reichenbachii, *Vis. Fl. Dalm.* iii. 169.—Dalmat.
Reinholdii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 238 = S. Behen.
Reinwardtii, Roth, Catalect. fasc. iii. 42 = picta.
remotiflora, *Vis. Fl. Dalm.* iii. 166.—Dalmat.
repens, Boiss. Fl. Orient. i. 614 = tenuis.
repens, *Patrin, in Pers. Syn. Pl.* i. 500.—Reg. Caucas.; Sibir. or.
Requienii, *Otth, in DC. Prod.* i. 381.—Graecia.
reticulata, *Desf. Fl. Atlant.* i. 350. t. 99.—Alger.
reticulata, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = cretica.
reticulata, Hort. ex Link, Enum. Hort. Berol. i. 426 = picta.
retroflexa, Steud. Nom. ed. I. 780; ed. II. ii. 587 = aegyptiaca.
Reuteriana, *Boiss. & Blanche, Diagn.* Ser. II. v. 54.—Syria.
rhodopea, *Janka, in Termész. Füzetek*, ii. (1878) 28.—Thracia.
rhynchocarpa, *Boiss. Diagn.* Ser. I. i. 33.—As. Min.
rigida, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 252.—Syria.
rigidula, Heldr. ex Boiss. Diagn. Ser. I. viii. 79 = echinosperma.
rigidula, Linn. Cent. Pl. ii. 16; Amoen. Acad. iv. 313 = cerastoides.
rigidula, *Sibth. & Sm. Fl. Graec. Prod.* i. 299.—Graecia.
rimarum, *J. Gay, ex Tchihat. Asie Min. Bot.* i. 200.—Cilicia.
Roemeri, *Frivald. in Flora*, xix. (1836) 439.—Banat.
Roeseri, Boiss. & Heldr. Diagn. Ser. II. i. 77 = ciliata.
rosea, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = vespertina.
rosea, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 865 = pendula.
Rosella, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 7 = gallica.
rostrata, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 244 = cerastoides.
rosulata, Soyer-Willem. & Godr. Expl. Sc. Algér. t. 82; et in Mém. Soc. Sc. Nancy, 1850 (1851), 184 = mollissima.
rotundifolia, *Nutt. Gen. Am.* i. 288.—Am. bor.
Royeni, Pers. Syn. i. 497 = viscosa.
rubella, Bory, ex Rohrb. Monog. Silene, 94 = graeca.
rubella, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 821 = colorata.

SILENE:—
rubella, *Linn. Sp. Pl.* 419.—Reg. Mediterr.; Mesopot.
rubella, Suffren, Cat. Pl. Frioul, 141 = cretica.
rubens, Vest, in Flora, iv. (1821) 150 = nutans.
rubescens, Mert. & Koch, Deutschl. Fl. iii. 572 (Index), sphalm. = praec.
rubicunda, A. Dietr. in Otto & Dietr. Allg. Gartenz. iii. (1835) 195 = pensylvanica?
rudbarica, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 36 = Aucheriana.
rugosa, Pers. Syn. i. 498 = Lychnis Coeli-rosa.
rupestris, Jacq. Enum. Stirp. Vindob. 77 = alpestris.
rupestris, *Linn. Sp. Pl.* 421.—Europ.
rupestris, Sibth. & Sm. Fl. Graec. Prod. i. 295 = chromodonta.
rupicola, Bor. Fl. Centr. Fr. ed. III. ii. 95 = S. Cucubalus.
rupicola, Huet, ex Rohrb. Monog. Silene, 138 = S. Saxifraga.
ruthenica, Otth, in DC. Prod. i. 370 = tatarica.
sabuletorum, Duby, Bot. Gall. 1025 = vespertina.
sabuletorum, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1850) = gallica.
sabuletorum, Link, in Spreng. Nov. Prov. 39 = hirsuta.
sachalinensis, *F. Schmidt, Reisen Amurl.* 116.—Ins. Sachalin.
sahendica, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 38 = olympica.
sajanensis, Less. ex Ledeb. Fl. Ross. i. 307 = tenuis.
salsa, Boiss. Diagn. Ser. I. viii. 77 = arenosa.
Salzmanni, *Badaro, ex Moretti, in Brugnat. Giorn.* ix. (1826) 78.—Europ. austr.
Salzmanni, Otth, in DC. Prod. i. 381 = velutina.
Saponaria, Cav. ex Duf. in Bull. Soc. Bot. Fr. vii. (1860) 244 = glauca.
Saponaria, Fries, ex Willk. & Lange, Prod. Fl. Hisp. iii. 671 = Saponaria officinalis.
saponariaefolia, Láng & Szovits, ex Rohrb. Monog. Silene, 81 = S. Fabaria.
saponariaefolia, Schott, ex Bess. Enum. Pl. Volh. 46 = S. Cucubalus.
sarawschanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 14.—Turkestan.
sardoa, Moris, ex Reichb. Fl. Germ. Excurs. 813, in obs. = gallica.
Sargentii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 290.—Am. bor. occ.
Sartorii, *Boiss. & Heldr. Diagn.* Ser. II. v. 53.—Ins. Cyclad.
Sassiana, Bertol. Fl. Ital. iv. 609 = fuscata.
saxatilis, *Sims, Bot. Mag.* t. 689; *Bieb. Fl. Taur. Cauc.* i. 338.—As. Min.; Persia.
saxicola, Lag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 641 = Lychnis Lagascae.
saxicola, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 43.—Hispan.
Saxifraga, Auct. ex Nym. Consp. 92 = fruticulosa.
Saxifraga, *Linn. Sp. Pl.* 421.—Europ.
Saxifraga, [Orsini], in Cappello, Opusc. Sc. 287, nomen = multicaulis.
scabra, Bertol. Fl. Ital. iv. 614 = inaperta.
scabra, Fisch. ex Sweet, Hort. Brit. ed. III. 66 = S. Cucubalus.
scabra, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 683.—Hungar.
scabrida, *Soyer-Willem. & Godr. Expl. Sc. Algér.* t. 81; *et in Mém. Soc. Sc. Nancy*, 1850 (1851), 167.—Alger.
scabridula, Boiss. Fl. Orient. i. 643 = armena.
scabriflora, Brot. Fl. Lusit. ii. 184 = pendula.
scepusiensis, Waldst. & Kit. ex Steud. Nom. ed. I. 780 (*stepusiensis*) (Quid?).—Hungar.
Schafta, *Gmel. ex Hohen. in Bull. Soc. Nat. Mosc.* xii. (1838) 397.—Reg. Caucas.
Schimperiana, Boiss. Diagn. Ser. I. i. 31 = chloraefolia.
Schimperiana, Hochst. ex Rohrb. Monog. Silene, 120 = Hochstetteri.
schirensis, Rohrb. l. c. 106, sphalm. = chirensis.
schizolepis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 370.—Mexic.
Schlosseri, *Vukot. in Oestr. Bot. Zeitschr.* xxx. (1880) 382.—Austria.
Schottiana, Schur, Enum. Pl. Transs. 103 = S. Cucubalus.

SILENE:—
Schweinfurthii, *Rohrb. in Bot. Zeit.* xxv. (1867) 82.—Nubia.
sciotica, Otth, in DC. Prod. i. 372 = gallica.
sclerocarpa, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 245 = cerastoides.
Sconleri, *Hook. Fl. Bor. Am.* i. 88.—Am. bor. occ.
secunda, Steud. Nom. ed. II. ii. 587 = tatarica.
secundiflora, Otth, in DC. Prod. i. 375 = glauca.
sedoides, *Jacq. Coll. Suppl.* 112. t. 14. f. 1.—Reg. Mediterr.
segetalis, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 241 = rubella.
Semenovii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) I. 536.—As. centr.
semiglabra, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 7 = gallica.
Sendtneri, *Boiss. Fl. Orient.* i. 608.—Banat.
sericea, *All. Fl. Pedem.* ii. 81. t. 79. f. 3.—Europ. austr.
sericea, Bertol. Amoen. Ital. 27 = colorata.
sericea, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = disticha, vespertina, S. Cucubalus.
serpyllifolia, Fisch. ex Steud. Nom. ed. I. 780 (Quid?).—Hab.?
serrulata, *Boiss. Fl. Orient.* i. 643.—Lycia.
sersuensis, *Pomel, Nouv. Mat. Fl. Atl.* 209.—Alger.
sessiliflora, Poir. Encyc. Suppl. v. 154 = dichotoma.
setabensis, Duf. ex Rohrb. Monog. Silene, 117 = glauca.
setacea, Otth, in DC. Prod. i. 372 = discolor.
setacea, *Viv. Fl. Lib. Spec.* 23.—Afr. bor.; Mesopot.
setifolia, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 8 = jenisea.
sibirica, *Pers. Syn.* i. 497.—Sibir.
sibirica, Schrank, in Syll. Ratisb. i. (1824) 212 = noctiflora?
Sibthorpiana, Reichb. Fl. Germ. Excurs. 815, in textu = dichotoma.
sicula, Bertol. ex Rohrb. Monog. Silene, 153 = fuscata.
sicula, Cyrilli, ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 46 = sedoides?
sicula, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = laeta.
sicula, Ker-Gawl. ex Rohrb. Monog. Silene, 115 = colorata.
sicula, Presl, Fl. Sic. i. 155 = italica, cretica.
sicula, Ucria, in Roem. Arch. i. I. (1796) 68 = italica.
siderophila, Boiss. & Gaill. in Boiss. Fl. Orient. i. 594 = palaestina.
Sieberi, Fenzl, in Russeg. Reise, ii. 912 = Fenzlii.
Sieberi, *Fenzl, Pugill. Pl. Nov. Syr.* 8.—Graecia; Ins. Creta.
Siegeri, Baumg. Enum. Stirp. Transs. i. 400 = nivalis.
silvestris, Schott, ex Reichb. Fl. Germ. Excurs. 812 = gallica.
simplicicorona, Mutel, Fl. Fr. i. 148, 169 = fuscata.
sinaica, Boiss. Diagn. Ser. I. i. 25 = odontopetala.
sisianica, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 36.—Reg. Cauc.
Smithii, Boiss. & Heldr. Diagn. Ser. II. i. 77 = S. Saxifraga.
Smithii, J. F. Gmel. Syst. 714 = chloraefolia.
sobolifera, Schott, ex Reichb. Fl. Germ. Excurs. 820 = repens.
solenantha, *Trautv. in Act. Hort. Petrop.* vii. (1880) 421.—Reg. Caucas.
Spaldingii, *S. Wats. in Proc. Am. Acad.* x. (1875) 344.—Am. bor. occ.
spathulaefolia, Jord. Cat. Jard. Dijon (1848), 31 = nutans.
spathulata, Bieb. Fl. Taur. Cauc. i. 341 = pygmaea.
spathulata, Schur, ex Nym. Consp. 90 = nutans.
spathulata, Willd. ex Rohrb. Monog. Silene, 156 = aegyptiaca.
speciosa, Paxt. Mag. Bot. x. (1843) 219 = laciniata.
spergulifolia, *Bieb. Fl. Taur. Cauc.* iii. 305.—Armen.
spergulifolia, Griseb. Spicil. Fl. Rumel. i. 174 = supina.
spergulifolia, Reichb. ex Nym. Consp. 90 = nutans.
spergulifolia, Schur, Enum. Pl. Transs. 101 = saxatilis.
spicata, DC. Fl. Fr. iv. 759 = nocturna.
spicata, *Ehrh. ex Rohrb. in Bot. Zeit.* xxv. (1867) 82.—Afr. or.

SILENE:—
spinescens, *Sibth. & Sm. Fl. Graec. Prod.* i. 299.—Graecia.
splendens, *Boiss. Fl. Orient.* i. 631.—Phrygia.
squamigera, *Boiss. Diagn.* Ser. I. i. 38.—As. Min.; Syria.
ssahendica, Rohrb. Monog. Silene, 198, sphalm. = *sahendica*.
ssojutica, Less. ex Ledeb. Fl. Ross. i. 308 = repens.
staminea, Bertol. Fl. Ital. iv. 585 = Roemeri.
staticefolia, Fenzl, ex Rohrb. Monog. Silene, 180 = armena.
staticefolia, C. Koch, ex Boiss. Fl. Orient. i. 639 = bupleuroides.
staticefolia, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1865) 114 = legionensis.
staticifolia, Sibth. & Sm. Fl. Graec. Prod. i. 301 = longiflora.
stellata, *Ait. Hort. Kew.* ed. II. iii. 84.—Am. bor.
stellata, Lapeyr. Hist. Abr. Pl. Pyr. 245; Fl. Pyr. 250. t. 91 = ciliata.
stellerioides, Nutt. ex Torr. & Gray, Fl. N. Am. i. 193 = Menziesii.
stenobotrys, *Boiss. & Haussk. in Boiss. Fl. Orient.* i. 611.—Syria.
stenopetala, *Ledeb. ex Sweet, Hort. Brit.* ed. III. 66, nomen.—Sibir.
stenophylla, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 240 = inaperta.
stenophylla, Ledeb. Fl. Ross. i. 306 = tenuis.
stenophylla, Plan. Ens Fl. Gallega, 121 = maritima.
stentoria, Fenzl, Pugill. Pl. Nov. Syr. 9 = caryophylloides.
stepusiensis, Steud. Nom. ed. I. 780; ed. II. ii. 587, sphalm. = *scepusiensis*.
Stevenii, Spreng. Cat. Sem. Hal. ex Steud. l. c. ed. II. ii. 587 (Quid?).—Rossia?
Stracheyi, *Edgew. in Hook. Fl. Brit. Ind.* i. 221.—Reg. Himal.
striata, *Ehrenb. ex Rohrb. in Bot. Zeit.* xxv. (1867) 83.—Syria.
stricta, Lapeyr. Hist. Abr. Pl. Pyr. 246 = S. Muscipula.
stricta, *Linn. Cent. Pl.* ii. 17; *Amoen. Acad.* iv. 314.—Europ. austr.
stricta, Spreng. Syst. ii. 412 = linicola.
struthioloides, *A. Gray, Bot. U. St. Expl. Exped.* i. 109.—Ins. Sandvic.
stylosa, *Bunge, in Ledeb. Fl. Alt.* ii. 144.—Sibir. altaic.
stylosa, Hort. ex Fenzl, Adv. Bot. [7]; Del. Sem. Hort. Vindob. (1851) = S. Cucubalus.
suaveolens, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 168 = Griffithii.
suavis, Schrad. ex Sweet, Hort. Brit. ed. III. 66 = vespertina?
subacaulis, Vill. ex Steud. Nom. ed. II. ii. 587 = acaulis.
subconica, Frivald. in Flora, xviii. (1835) 334.—Thracia.
subnutans, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 78 = nutans.
subocculta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 5 = gallica.
subspathulata, Schur, Enum. Pl. Transs. 102 = chlorantha.
subulata, Boiss. Diagn. Ser. I. i. 33 = stentoria.
subulata, Heldr. ex Boiss. Diagn. Ser. II. v. 56 = S. Echinus.
subvinosa, Delile, Ind. Sem. Hort. Monsp. (1838) 12 = nicaeënsis.
succulenta, *Forsk. Fl. Aegypt. Arab.* p. lxvi; *Delile, Fl. Aegypt. Descr.* 89.—Aegypt.; Syria.
suffrutescens, Bieb. Fl. Taur. Cauc. i. 340 = caspica.
suffruticosa, Desf. ex Webb, Iter Hisp. 64 = fruticosa, velutina.
suffruticosa, Pall. Reise, ii. 522 = altaica.
supina, *Bieb. Fl. Taur. Cauc.* i. 338.—Reg. Caucas.
swertiaefolia, Boiss. Diagn. Ser. I. i. 32 = chloraefolia.
sylvestris, Reichb. Fl. Germ. Excurs. Index, 121 = *silvestris*.
syriaca, Reut. Cat. Gr. Jard. Genèv. (1857) 4.—Cf. Linnaea, xxxix. (1857–58) 724 = picta.
tachtensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 239.—Turkestan.
tatarica, *Pers. Syn.* i. 497.—Europ. or.; As. occ.
Tatarinowii, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) II. 562.—China.
taurica, *Fisch. ex Sweet, Hort. Brit.* ed. III. 66 (Quid?).—Tauria.

SILENE:—
tegedensis, Boiss. Elench. 20 = Boryi.
tenella, Huet, ex Schenk, Ind. Sem. Hort. Wirceb. (1861) = quadrifida.
tenella, C. Koch, ex Rohrb. Monog. Silene, 191 = petraea.
tenella, *C. A. Mey. Verz. Pfl. Cauc.* 216.—Reg. Caucas.
tenerrima, J. & C. Presl, Delic. Prag. 217 = hirsuta.
Tenoreana, Colla, Herb. Pedem. i. 328; ex Soyer-Willem. & Godr. in Mém. Soc. Sc. Nancy, 1850 (1851) 141 = S. Cucubalus.
tenuiflora, Guss. Pl. Rar. Sic. 177 = cretica.
tenuiflora, Hort. ex Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = cretica.
tenuifolia, Otth, in DC. Prod. i. 379 = tenuis.
tenuis, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 575 = jenisea.
tenuis, *Willd. Enum. Hort. Berol.* 474.—Reg. Himal.; Sibir.
tergestina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 6 = gallica.
thebana, *Orph. ex Boiss. Fl. Orient.* i. 627.—Graecia.
thessalonica, Boiss. & Heldr. Diagn. Ser. II. i. 74 = flavescens.
Thirkeana, C. Koch, in Linnaea, xix. (1847) 56 = dichotoma.
Thorei, *Duf. in Ann. Sc. Nat.* Sér. I. v. (1825) 84.—Gallia.
Thunbergiana, *Eckl. & Zeyh. Enum.* 32.—Afr. austr.
Thunbergiana, Krauss, ex Rohrb. Monog. Silene, 122 = primulaeflora.
Thunbergii, E. Mey. in Drège, Zwei Pfl. Docum. 87 = undulata.
Thurberi, *S. Wats. in Proc. Am. Acad.* x. (1875) 343.—N. Mexic.
thymifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 292.—Ins. Cyprus.
thysanodes, *Fenzl, in Endl. Nov. Stirp. Dec.* 39.—Am. austr.
Tilingi, Regel, in Act. Hort. Petrop. i. (1871) 99 = californica.
tincta, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 174 = Friwaldskyana.
tmolea, *J. Gay, in Tchihat. Asie Min. Bot.* i. 200.—Lydia.
tomentosa, Otth, in DC. Prod. i. 383 = mollissima.
Tommasinii, Vis. in Flora, xii. (1829) I. Erg. 12 = sedoides.
tragacantha, *Fenzl, ex Boiss. Fl. Orient.* i. 621.—Persia.
transsilvanica, Schur, in Oestr. Bot. Zeitschr. viii. (1858) 22; x. (1860) 181 = nutans.
tricuspidata, Desf. Tabl. ed. II. 184, nomen = gallica, disticha.
tridentata, *Desf. Fl. Atlant.* i. 349.—Reg. Mediterr. occ.
tridentata, Ram. ex DC. Fl. Fr. iv. 758 = gallica.
trifida, *Tenore, ex Steud. Nom.* ed. II. ii. 588.—Hab.
trinervia, Fenzl, Adv. Bot. [2]; in Del. Sem. Hort. Vindob. (1851) = squamigera.
trinervia, *Sebast. & Mauri, Fl. Rom.* 152.—Europ. austr.
trinervis, [Soland. in] Russ. Aleppo, ed. II. ii. 252 = dichotoma.
tristis, Salisb. Prod. 301 = undulata, *Ait.*
Tuba, Ehrenb. ex Rohrb. Monog. Silene, 130 = Boryi.
tubiflora, Duf. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 370 = colorata.
tunicoides, *Boiss. Diagn.* Ser. I. i. 34.—Lycia.
turbinata, Guss. Fl. Sic. Prod. i. 506 = rubella.
turgida, *Bieb. ex Bunge, Verz. Alt.* 23.—Sibir. altaic.
turgida, Schrenk, ex Rohrb. Monog. Silene, 187 = tenuis.
turkestanica, *Regel, in Act. Hort. Petrop.* ii. (1873) 436.—Turkestan.
Uhdeana, *Rohrb. Ind. Sem. Hort. Berol.* (1867) *App.* ii. 2.—Mexic.
umbellata, Gilib. Fl. Lituan. ii. 170 = S. Armeria.
undulaefolia, Sweet, Hort. Brit. ed. III. 65 = fuscata.
undulata, *Ait. Hort. Kew.* ed. I. ii. 96.—Afr. austr.
undulata, *Moench, Meth.* 707.—Hab.?
undulata, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 660 = rubella.
undulatifolia, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = gallica.
undulatifolia, Moris, Stirp. Sard. Elench. 8 = fuscata.
Ungeri, *Fenzl, in Unger, Reise Griechenl.* 136; *Boiss. Fl. Orient.* i. 601.—Ins. Ithaca.

SILENE:—
uniflora, Bertol. Fl. Ital. iv. 632 = S. Cucubalus?
uniflora, Roth, in Usteri, Ann. Bot. x. (1794) 46; et in Catalect. fasc. i. 52 = maritima.
unilocularis, Schrank, Baier. Fl. i. 712 = Arenaria spathulata?
Urvillii, *Schott, ex Urv. in Mém. Soc. Linn. Par.* i. (1822) 304.—Ins. Cos; Lydia.
vallesia, Bory & Chaub. Nouv. Fl. Pélop. 27 = paradoxa.
vallesia, *Linn. Syst.* ed. X. 1032.—Europ.
vallesia, C. A. Mey. Verz. Pfl. Cauc. 215 = caucasica.
vallesia, Pollini, Fl. Veron. ii. 58 = Elisabethae.
vallesiaca, Link, Handb. ii. 244 = nicaeënsis.
variegata, *Boiss. & Heldr. Diagn.* Ser. I. viii. 82.—Graecia.
variegata, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = S. Armeria.
velutina, Hort. ex Fenzl, l. c. = annulata, gallica.
velutina, *Pourr. ex Loisel. in Desv. Journ. Bot.* ii. (1809) 324.—Ins. Corsic.
velutinoides, *Pomel, Nouv. Mat. Fl. Atl.* 208.—Alger.
venosa, Aschers. Fl. Brandenb. i. 86 = S. Cucubalus.
verecunda, *S. Wats. in Proc. Am. Acad.* x. (1875) 344.—Calif.
vernalis, Poepp. ex Rohrb. Monog. Silene, 97 = gallica.
verticillata, Otth, in DC. Prod. i. 370 = spergulifolia.
vesicaria, Schrad. ex Reichb. Fl. Germ. Excurs. 822 = S. Cucubalus.
vesiculifera, *J. Gay, ex Boiss. Fl. Orient.* i. 586.—Cilicia.
vespertina, Curt. Bot. Mag. t. 677 = colorata.
vespertina, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = S. Cucubalus.
vespertina, *Retz. Obs.* iii. 31.—Reg. Mediterr.; Afr. bor.
vestita, Soyer-Willem. & Godr. in Expl. Sc. Algér. t. 81; et in Mém. Soc. Sc. Nancy, 1850 (1851), 154 = micropetala.
vilipensa, Kunze, in Ind. Sem. Hort. Lips. (1845); et in Flora, xxii. (1846) 641 = portensis, inaperta.
vilis, Bartl. Ind. Sem. Hort. Gotting. (1845) = portensis.
vilis, Fenzl, ex Boiss. Fl. Orient. i. 596 = apetala.
vilis, Kunze, ex Nym. Consp. 95 = inaperta.
villosa, Boiss. Voy. Bot. Espagne, ii. 90 = S. Psammitis.
villosa, *Cambess. in Mém. Mus. Par.* xiv. (1827) 221. t. 41.—Hispan.
villosa, *Forsk. Fl. Aegypt. Arab.* 88.—Aegypt.
villosa, Forsk. l. c. 210 = gallica.
villosa, Holl. ex Rohrb. Monog. Silene, 112 = littorea.
villosa, Hort. ex Fenzl, Adv. Bot. [7]; in Del. Sem. Hort. Vindob. (1851) = obtusifolia, vespertina.
villosa, Moench, Meth. 708 = nicaeënsis.
virginica, Benth. Pl. Hartw. 299 = californica.
virginica, Linn. Sp. Pl. 419.—Am. bor.
virginica, Rafin. ex DC. Prod. i. 383 = pensylvanica.
virginica, Willd. Sp. Pl. ii. 702 = pensylvanica?
viridella, Link, Handb. ii. 242 = nutans.
viridiflora, Gueldenst. ex Ledeb. Fl. Ross. i. 319 = chlorantha.
viridiflora, Kotschy, ex Rohrb. Monog. Silene, 214 = amana.
viridiflora, *Linn. Sp. Pl.* ed. II. 597.—Europ. austr.
viscaginoides, Hornem. Suppl. Hort. Hafn. 49 = tenuis.
viscaginoides, Otth, in DC. Prod. i. 368 = jenisea.
viscariaefolia, Boiss. Diagn. Ser. I. i. 30 = bupleuroides.
viscariaefolia, Bourg. ex Rohrb. Monog. Silene, 179 = longiflora.
viscida, Moench, Meth. 708 = noctiflora.
viscida, Spreng. Pugill. ii. 65 = alpestris.
viscosa, *Pers. Syn.* i. 497.—Europ.; As. bor.
viscosa, Schleich. Cat. Pl. Helv. 33 = italica.
viscosissima, Rochel, ex Rohrb. Monog. Silene, 220 = italica.
viscosissima, Tenore, Prod. Fl. Nap. p. xxvi = nicaeënsis.
Vivianii, Steud. Nom. ed. II. ii. 588 = setacea.
volgensis, Spreng. Syst. ii. 413 = wolgensis.
vulgaris, Garcke, Fl. Deutschl. ed. IX. 64 = S. Cucubalus.
Waldsteinii, *Griseb. Spicil. Fl. Rumel.* i. 179.—Europ. austr.
Wallichiana, Klotzsch, in Bot. Ergeb. Waldem. Reise, 139 = S. Cucubalus.
Webbiana, *Wall. Cat.* n. 627.—Reg. Himal.
Willdenowii, Sweet, Hort. Brit. ed. II. 51 = Thorei.
Willkommiana, J. Gay, ex Coss. Notes Crit. ii. 32 = divaricata.

SILENE:—
wolgensis, *Bess. ex Spreng. Ind. Sem. Hort. Hal.* (1818) 6; *Ledeb. Fl. Alt.* ii. 140 (an seq.?).—Europ. or.
wolgensis, Otth, in DC. Prod. i. 370 = S. Otites.
Wrightii, *A. Gray, Pl. Wright.* ii. 17.—N. Mexic.
xeranthema, *Viv. Fl. Lib. Spec.* 67; *Fl. Cors. Sp. Nov.* 7.—Ins. Corsic.
Zawadskii, Hort. ex Fenzl, Adv. Bot.[7]; in Del. Sem. Hort. Vindob. (1851) = S. Cucubalus.
Zawadzkii, *Herbich, in Zawad. Fl. Galic.* 191.—Bucovina.

SILENOPSIS, Willk. in Bot. Zeit. v. (1847) 237 = **Lychnis**, Tourn. (Caryoph.).
Lagascae, Willk. l. c. = L. Lagascae.

SILER, Crantz, Stirp. Austr. ed. I. fasc. iii. 60 (1767); ed. II. 185 (1769). *UMBELLIFERAE*, Benth. & Hook. f. i. 908.
AGASYLLIS, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259.
BRADLAEIA, Neck. Elem. i. 169 (1790).
alpinum, Baumg. Enum. Stirp. Transs. i. 229 = trilobum.
aquilegifolium, Gaertn. Fruct. i. 92. t. 22 = trilobum.
caucasicum, *Spreng. Sp. Umbellif.* 90. — Reg. Caucas.
Chironium, Crantz, Cl. Umb. Emend. 69 = Opoponax Chironium.
divaricatum, *Benth. & Hook. f. Gen.* i. 909.—Sibir. baical.
gallicum, Crantz, Stirp. Austr. ed. I. fasc. iii. 65; ed. II. 188 = Laserpitium gallicum.
lancifolium, Hoffmgg. & Link, Fl. Port. ii. t. 109 = Peucedanum lancifolium.
lancifolium, Moench, Meth. 85 = Laserpitium Siler.
montanum, Crantz, Stirp. Austr. ed. I. fasc. iii. 60; ed. II. 185 = Laserpitium Siler.
salsum, Spreng. Sp. Umbellif. 90 = Peucedanum redivivum.
trilobum, *Crantz, Stirp. Austr.* ed. I. fasc. iii. 62; ed. II. 186.—Oriens.

SILIQUARIA, Forsk. Fl. Aegypt. Arab. 78 (1775) = **Cleome**, Linn. (Cappar.).
arabica, Rafin. Sylva Tellur. 111 = C. arabica.
glandulosa, Forsk. Fl. Aegypt. Arab. 78 = C. arabica.

SILIQUASTRUM, Tourn. ex Adans. Fam. ii. 317 (1763) = **Cercis**, Linn. (Legumin.).
Arbor-Judae, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 339 = C. Siliquastrum.
canadense, Medic. l. c. = C. canadensis.
cordatum, Moench, Meth. 54 = C. canadensis.
orbiculatum, Moench, l. c. = C. Siliquastrum.

SILLYBUM, Hassk. Cat. Hort. Bog. Alt. 104 (1844), sphalm. = **Silybum**, Vaill. (Compos.).

SILOXERUS, Labill. Nov. Holl. Pl. ii. 57. t. 209 (1806) = **Angianthus**, Wendl. (Compos.).
humifusus, Labill. l. c. 58 = A. humifusus.

SILPHION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = **Silphium**, Linn. (Compos.).
asteriscum, St. Lag. l. c. = Silphium Astericus.

SILPHIOSPERMA, Steetz, in Lehm. Pl. Preiss. i. 433 (1845) = **Brachycome**, Cass. (Compos.).
collinum, Sond. in Linnaea, xxv. (1852) 483 = B. collina.
glandulosum, Steetz, in Lehm. Pl. Preiss. i. 433 = B. glandulosa.
perpusillum, Steetz, l. c. = B. collina.

SILPHIUM, Linn. Coroll. Gen. 16 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 350.
albiflorum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 4.—Texas.
arborescens, *Mill. Gard. Dict.* ed. VIII. n. 4.—Mexic.
asperrimum, *Hook. Comp. Bot. Mag.* i. (1835) 99.—Am. bor. occ.
Asteriscus, Hornem. Hort. Hafn. 854 = Berlandiera tomentosa.
Asteriscus, *Linn. Sp. Pl.* 920.—Am. bor.
atropurpureum, Ketz. ex Willd. Sp. Pl. iii. 2334 = trifoliatum.

SILPHIUM:—
betonicaefolium, Hook. Comp. Bot. Mag. i. (1835) 99 = Berlandiera texana.
brachiatum, *Gattinger, in Coult. Bot. Gaz.* ix. (1884) 192.—Am. bor.
compositum, *Michx. Fl. Bor. Am.* ii. 145.—Am. bor.
conjunctum, Willd. Enum. Hort. Berol. 933 = perfoliatum.
connatum, Linn. Mant. ii. 574 = perfoliatum.
dentatum, Ell. Sketch, ii. 468 = S. Asteriscus.
dissectum, *Poir. Encyc. Suppl.* v. 157.—Am. bor.?
doronicifolium, *Kunze, Del. Sem. Hort. Lips.* (1846) 4; *et in Linnaea*, xx. (1847) 20.—Afr. austr.
edule, Baill. Hist. Pl. viii. 306 = Polymnia edulis.
elatum, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 579 = compositum.
equatoriale, Spruce, ex Benth. & Hook. f. Gen. ii. 350 = Wedelia equatorialis.
erythrocaulon, Bernh. ex Spreng. Syst. iii. 630 = perfoliatum.
glaucum, *Weinm. Hort. Dorpat.* 143; *ex DC. Prod.* v. 571, in obs.—Am. bor.
gracile, *A. Gray, in Proc. Am. Acad.* viii. (1873) 653.—Texas.
gummiferum, Ell. Sketch, ii. 460 = laciniatum.
helianthoides, Linn. Sp. Pl. 920 = Heliopsis laevis.
Hornemanni, Schrad. ex DC. Prod. v. 514 = perfoliatum.
integrifolium, *Michx. Fl. Bor. Am.* ii. 146.—Am. bor.
laciniatum, *Linn. Sp. Pl.* 919.—Am. bor.
laciniatum, Walt. Fl. Carol. 217 = compositum.
laeve, *Hook. Lond. Journ. Bot.* vi. (1847) 244.—Am. bor.
laevigatum, Ell. Sketch, ii. 465 = trifoliatum.
laevigatum, Pursh, Fl. Am. Sept. ii. 578 = integrifolium.
lanceolatum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 341 = S. Asteriscus.
latifolium, *Hill, Hort. Kew.* 11.—Am. bor.
nudicaule, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 127 = compositum.
Nuttallianum, Torr. in Ann. Lyc. N. York, ii. (1828) 216 = Berlandiera lyrata.
patulum, Banks, ex Pursh, Fl. Am. Sept. ii. 578 = Berlandiera tomentosa.
perfoliatum, *Linn. Syst.* ed. X. 1232.—Am. bor.
peristenium, Rafin. Atl. Journ. 146 (Quid?).—Am. bor.
petiolatum, Poir. Encyc. Suppl. v. 157 = perfoliatum.
pinnatifidum, Ell. Sketch, ii. 462 = terebinthinaceum.
platypterum, Spruce, ex Benth. & Hook. f. Gen. ii. 350 = Verbesina platyptera.
Pseudo-laciniatum, Fras. ex DC. Prod. v. 512 = compositum.
pumilum, Michx. Fl. Bor. Am. ii. 183 = Berlandiera tomentosa.
quadrangulum, Hedw. in Roem. Arch. iii. (1803) 123 = perfoliatum.
quercifolium, *DC. Prod.* v. 513.—Hab.?
Radula, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 341 = asperrimum.
reniforme, Rafin. ex Nutt. l. c. 342 = compositum.
reticulatum, Moench, Meth. 606 = S. Asteriscus.
reticulatum, Pursh, Fl. Am. Sept. ii. 579 = Berlandiera tomentosa.
reticulatum, Rafin. Atl. Journ. 148 = praec.?
rigidum, *Moench, Meth. Suppl.* 258.—Hab.?
scaberrimum, *Ell. Sketch*, ii. 466.—Am. bor.
scabrum, Moench, Meth. 606 = perfoliatum.
scabrum, Walt. Fl. Carol. 217 = S. Asteriscus.
sinuatum, Banks, ex Pursh, Fl. Am. Sept. ii. 577 = compositum.
solidaginoides, Linn. Sp. Pl. 920 = Heliopsis laevis.
speciosum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 341 = integrifolium.
spicatum, Poir. Encyc. Suppl. v. 157 = laciniatum.
subacaule, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 301 = Berlandiera subacaulis.
terebinthinaceum, Ell. Sketch, ii. 463 = compositum.
terebinthinaceum, *Jacq. Hort. Vindob.* i. 16.—Am. bor.
ternatum, Retz. ex Willd. Sp. Pl. iii. 2333 = trifoliatum.
ternifolium, Michx. Fl. Bor. Am. ii. 146 = trifoliatum.
Tetragonotheca, Gaertn. Fruct. ii. 444. t. 171. f. 2 = Tetragonotheca helianthoides.
tetragonum, Moench, Meth. 606 = perfoliatum.
tomentosum, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 579 = Berlandiera tomentosa.

SILPHIUM:—
trachopus, *Rafin. Herb.* 24; *Atl. Journ.* 178.—Am. bor.
trifoliatum, *Linn. Sp. Pl.* 920.—Am. bor.
trilobatum, Linn. Syst. ed. X. 1232 = Wedelia carnosa.

SILVAEA, Hook. & Arn. Bot. Beech. Voy. 211 (1836) = **Trigostemon**, Blume (Euphorb.).
Hookeriana, Baill. Etud. Gén. Euph. 342 = T. semperflorens.
semperflorens, Hook. & Arn. Bot. Beech. Voy. 211 = T. semperflorens.

SILVAEA, Meissn. in DC. Prod. xv. 1. 84 (1864) = **Endiandra**, R. Br. (Laurin.).
Navalium, Meissn. l. c. = E. Navalium (in Addend.).

SILVAEA, Phil. Fl. Atac. 21. t. 1 C (1860). *PORTULACEAE*, Benth. & Hook. f. i. 159.
amarantoides, *Phil. l. c.* 22.—Chili.
celosioides, *Phil. l. c.*—Chili.
corrigioloides, *Phil. l. c.*—Chili.
pschyphylla, *Phil. l. c.* 21.—Chili.

SILVIA, Allem. Diss. Rio Janeiro (1845).—Cf. Bot. Zeit. xii. (1854) 454 = **Endiandra**, R. Br. (Laurin.).
Navalium, Allem. l. c. = E. Navalium (in Addend.).

SILVIA, Benth. in DC. Prod. x. 513 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 972.
prostrata, *Benth. in DC. Prod.* x. 513.—Mexic.
serpyllifolia, *Benth. l. c.*—Mexic.

SILVIA, Vell. Fl. Flum. 55 (1825); i. t. 149 (1827) = **Escobedia**, Ruiz & Pav. (Scrophul.).
curialis, Vell. ll. cc. = E. scabrifolia.

SILVIANTHUS, Hook. f. in Hook. Ic. Pl. t. 1048 (1868). *RUBIACEAE*, Benth. & Hook. f. ii. 64.
bracteatus, *Hook. f. in Hook. Ic. Pl.* t. 1048.—Reg. Himal.

SILYBON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = seq.

SILYBUM, Vaill. ex Adans. Fam. ii. 116 (1763); Gaertn. Fruct. ii. 378. t. 162 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 470.
MARIANA, Hill, Veg. Syst. iv. 19 (1762), nomen prius.
atriplicifolium, Fisch. Ind. Hort. Petrop. (1824); ex DC. Prod. vi. 663 = Serratula atriplicifolia.
cernuum, Gaertn. Fruct. ii. 378. t. 162. f. 2 = Carduus cernuus, *Steud.*
cernuum, Ledeb. Fl. Alt. iv. 43 = Carduus stenolepis.
eburneum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 366.—Afr. bor.
hispanicum, Loscos & Pardo, Ser. Inconf. Arag. ed. Willk. 56 = eburneum.
intermedium, *Willk. Ind. Sem. Hort. Dorpat.* (1869) 6.—Hab.?
leucanthum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 83 = S. Marianum.
maculatum, Moench, Meth. 555 = S. Marianum.
Mariae, S. F. Gray, Nat. Arr. Brit. Pl. ii. 436 = S. Marianum.
Marianum, *Gaertn. Fruct.* ii. 378. t. 168.—Europ.
pygmaeum, Cass. in Dict. Sc. Nat. l. 469 = S. Marianum.
squarrosum, Tausch, in Flora, xix. (1836) 396 = Carduus cernuus, *Steud.*

SILYMBRIUM, Neck. in Act. Acad. Theod. Palat. ii. (1770) 480, sphalm. = **Sisymbrium**, Linn. (Crucif.).

SIMABA, Aubl. Pl. Gui. i. 400. t. 153 (1775). *SIMARUBEAE*, Benth. & Hook. f. i. 308.
ARUBA, Aubl. Pl. Gui. i. 293. t. 115 (1775).
HOMALOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 575.
PHYLLOSTEMA, Neck. Elem. ii. 301 (1790).
ZWINGERA, Schreb. Gen. ii. 802 (1791).
africana, Baill. Adansonia, vii. (1866–67) 381 = Quassia africana.
angustifolia, Benth. in Hook. Kew Journ. iii. (1851) 371 = guianensis?
Aruba, A. St. Hil. ex DC. Prod. i. 734 = guianensis.

SIMABA:—
bahiensis, Moric. in Mém. Soc. Phys. Genèv. vii. (1836) 251 = ferruginea?
bicolor, *Zucc. in Abh. Akad. Muench*, i. (1829–30) 353.—Mexic.
Blanchetii, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 444 = ferruginea.
Cedron, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 566.—Am. trop.
crustacea, *Engl. in Mart. Fl. Bras.* xii. II. 211. —Bras.
cuneata, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 138.—Bras.
cuspidata, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. II. 212.—Bras.
ferruginea, *A. St. Hil. in Bull. Soc. Philom.* (1823) 129; *Pl. Rem. Brés.* i. 127.—Bras.
floribunda, *A. St. Hil. ll. cc.* 129; 126.—Bras.
foetida, Benth. in Hook. Kew Journ. iii. (1851) 370 = guianensis?
glabra, *Engl. in Mart. Fl. Bras.* xii. II. 218.—Bras.
glandulifera, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 169.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 400. t. 153.—Am. trop.
indica, Baill. Hist. Pl. iv. 440 = Samadera indica.
insignis, A. St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 137 = glandulifera.
laevis, *Casar. Nov. Stirp. Bras. Dec.* 10.—Bras.
longifolia, Casar. l. c. 9 = glandulifera.
maiana, *Casar. l. c.* 10.—Bras.
multiflora, A. Juss. in Mém. Mus. Par. xii. (1825) t. 27 = guianensis?
nigrescens, *Engl. in Mart. Fl. Bras.* xii. II. 213. —Bras.
obovata, *Spruce, ex Engl. l. c.* 210.—Bras.
orinocensis, *H. B. & K. Nov. Gen. et Sp.* vi. 18. t. 514. —Venezuela.
quassioides, D. Don, Prod. Fl. Nep. 248 = Picrasma quassioides.
salubris, *Engl. in Mart. Fl. Bras.* xii. II. 219.—Bras.
suaveolens, *A. St. Hil. in Bull. Soc. Philom.* (1823) 129; *Pl. Rem. Brés.* 128.—Bras.
subcymosa, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 137.—Bras.
suffruticosa, *Engl. in Mart. Fl. Bras.* xii. II. 213. —Bras
trichilioides, Planch. in Hook. Lond. Journ. Bot. v. (1846) 565, partim = S. Cedron.
trichilioides, *A. St. Hil. in Bull. Soc. Philom.* (1823) 129; *Pl. Rem. Brés.* 129.—Bras.
undulata, *Guill. & Perr. Fl. Seneg. Tent.* i. 136. —Afr. trop.
Warmingiana, *Engl. in Mart. Fl. Bras.* xii. II. 217.— Bras.

SIMARUBA, Aubl. Pl. Gui. ii. 859. t. 331, 332 (1775) (Simarouba); DC. in Ann. Mus. Par. xvii. (1811) 423. *SIMARUBEAE*, Benth. & Hook. f. i. 309.
amara, *Aubl. l. c.* 860.—Am. trop.
amara, Hayne, Arzn. Gew. ix. 15 = glauca.
antisyphilitica, Mart. ex Engl. in Mart. Fl. Bras. xii. II. 225 = versicolor.
excelsa, DC. in Ann. Mus. Par. xvii. (1811) 424 = Picraena excelsa.
glauca, *DC. l. c.*—Am. trop.
laevis, *Griseb. Cat. Pl. Cub.* 49.—Cuba.
medicinalis, Endl. Mediz. 528 = glauca.
monophylla, *Oliver, in Hook. Ic. Pl.* t. 1387.—Guian.
officinalis, DC. in Ann. Mus. Par. xvii. (1811) 423 = amara.
officinalis, Macfad. Fl. Jamaic. i. 198 = glauca.
versicolor, *A. St. Hil. Pl. Us. Bras.* t. 5.—Bras.

SIMBLOCLINE, DC. Prod. v. 297 (1836) = **Diplostephium**, H. B. & K. (Compos.).
Haenkei, DC. l. c. = D. Haenkei.

SIMBULETA, Forsk. Fl. Aegypt. Arab. 115 (1775) = **Anarrhinum**, Desf. (1800) (Scrophul.).
arabica, Poir. in Dict. Sc. Nat. xlix. 244 = A. arabicum.
Forskâhlii, J. F. Gmel. Syst. 242 = A. arabicum.

SIMETHIS, Kunth, Enum. Pl. iv. 618 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 784.
MORGAGNIA, Bubani, in Nouv. Ann. Sc. Nat. Bologn. ix. (1843) 92.

SIMETHIS:—
POGONELLA, Salisb. Gen. Pl. Fragm. 70 (1866).
PUBILARIA, Rafin. Fl. Tellur. ii. 27 (1836), nomen prius.
SIEBOLDIA, Heynh. Nom. ii. 664 (1846).
bicolor, *Kunth, Enum. Pl.* iv. 618.—Europ.
planifolia, Gren. & Godr. Fl. Fr. iii. 222 = bicolor.

SIMIRA, Aubl. Pl. Gui. i. 170. t. 65 (1775) = **Psychotria**, Linn. (Rubiac.).
nitida, Poir. Encyc. vii. 196 = P. nitida.
Palicourea, Poir. l. c. = Palicourea guianensis.
tinctoria, Aubl. Pl. Gui. i. 170 = P. parviflora, *Willd.*

SIMIRA, Rafin. Fl. Tellur. iv. 28 (1836) = **Scilla**, Linn. (Liliac.).
japonica, Rafin. l. c. = Sc. japonica.

SIMMONDSIA, Nutt. in Hook. Lond. Journ. Bot. iii. (1844) 400. t. 16. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 265.
BROCCHIA, Mauri, in Tenore, Cat. Ort. Napoli (1845), 80.
californica, *Nutt. in Hook. Lond. Journ. Bot.* iii. (1844) 400. t. 16.—Calif.
pabulosa, Kellogg, in Proc. Calif. Acad. ii. (1859) 21 = californica.

SIMOCHEILUS, Klotzsch, in Linnaea, xii. (1838) 236. *ERICACEAE*, Benth. & Hook. f. ii. 593.
ANOMALANTHUS, CODONANTHEMUM, MACROLINUM, OCTOGONIA, PACHYCALYX, PLAGIOSTEMON, SYNDESMANTHUS, THAMNUS, THORACOSPERMA, Klotzsch, l. c. 230–242.
barbiger, *Klotzsch, l. c.* 237.—Afr. austr.
bicolor, *Benth. in DC. Prod.* vii. 703.—Afr. austr.
carneus, *Klotzsch, in Linnaea*, xii. (1838) 236.—Afr. austr.
depressus, *Benth. in DC. Prod.* vii. 704.—Afr. austr.
Ecklonianus, *Benth. l. c.* 705.—Afr. austr.
glabellus, *Benth. l. c.* 704.—Afr. austr.
glaber, *Benth. l. c.* 705.—Afr. austr.
hirsutus, *Benth. l. c.* 704.—Afr. austr.
hispidus, *Benth. l. c.* 705.—Afr. austr.
Klotzschianus, *Benth. l. c.* 703.—Afr. austr.
multiflorus, *Klotzsch, in Linnaea*, xii. (1838) 236.— Afr. austr.
oblongus, *Benth. in DC. Prod.* vii. 705.—Afr. austr.
obovatus, *Benth. l. c.* 703.—Afr. austr.
patens, *Benth. l. c.* 704.—Afr. austr.
pubescens, *Klotzsch, in Linnaea*, xii. (1838) 236.— Afr. austr.
quadrifidus, *Benth. in DC. Prod.* vii. 703.—Afr. austr.
quadriflorus, Steud. Nom. ed. II. ii. 589, sphalm. = praec.
submuticus, *Benth. in DC. Prod.* vii. 704.—Afr. austr.

SIMONISIA, Nees, in Mart. Fl. Bras. ix. 144. t. 23 (1847) = **Beloperone**, Nees (Acanth.).
asclepiadea, Nees, l. c. 145.—Bras.
Riedeliana, Nees, l. c. = B. Riedeliana.
rosea, Nees, l. c. 144 = B. rosea.

SIMPHITUM, Neck. Delic. Gallo-Belg. i. 101 (1768) = **Symphytum**, Tourn. (Borag.).

SIMPLOCARPUS, F. Schmidt, Reisen Amurl. 178 (1868), sphalm. = **Symplocarpus**, Salisb. (Aroid.).

SIMPLOCOS, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 22 (1824), sphalm. = **Symplocos**, Linn. (Styrac.).

SIMSIA, R. Br. in Trans. Linn. Soc. x. (1810) 152 = **Stirlingia**, Endl. (Proteac.).
abrotanoides, F. Muell. Census, 66 = St. abrotanoides.
anethifolia, R. Br. in Trans. Linn. Soc. x. (1810) 153 = St. tenuifolia.
latifolia, R. Br. Prot. Nov. 9 = St. latifolia.
simplex, F. Muell. Census, 66 = St. simplex.
tenuifolia, R. Br. in Trans. Linn. Soc. x. (1810) 152 = St. tenuifolia.
teretifolia, F. Muell. Census, 66 = St. teretifolia.

SIMSIA, Pers. Syn. ii. 478 (1807) = **Encelia**, Adans. (Compos.).
amplexicaulis, Pers. l. c. = E. amplexicaulis.
auriculata, DC. Prod. v. 577 = E. mexicana.
calva, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 228 = E. calva.
canescens, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 85 = E. eriocephala.
cordata, Cass. in Dict. Sc. Nat. lix. 137 = E. cordata.
Dombeyana, DC. Prod. v. 578.—Peruv.?
exaristata, A. Gray, Pl. Wright. ii. 87 = E. exaristata.
ficifolia, Pers. Syn. ii. 478 = E. foetida.
frutescens, A. Gray, in Torr. Bot. Mex. Bound. 89 = E. frutescens.
grandiflora, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 92 = E. grandiflora.
heterophylla, DC. Prod. v. 577 = E. heterophylla.
heterophylla, Pers. Syn. ii. 478 = Echinacea heterophylla.
hispida, Cass. in Dict. Sc. Nat. lix. 137 = E. hispida.
Kunthiana, Cass. l. c. = E. heterophylla.
lagascaeformis, DC. Prod. v. 577 = E. lagascaeformis.
lagascaeformis, A. Gray, Pl. Wright. i. 107 = E. exaristata.
pastoënsis, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 40.—N. Granat.
polycephala, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 93 = E. polycephala.
pubescens, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 40.—N. Granat.
sanguinea, A. Gray, Pl. Wright. i. 107 = E. sanguinea.
scaposa, A. Gray, l. c. ii. 88 = E. scaposa.
subaristata, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 84 = E. subaristata.

SIMSIMUM, Bernh. in Linnaea, xvi. (1842) 42 = **Sesamum**, Linn. (Pedal.).
rostratum, Bernh. l. c. = Sesamum capense.

SINAPI, Dulac, Fl. Hautes-Pyr. 194 (1867) = **Brassica**, Tourn. (Crucif.).
pinnipartitum, Dulac, l. c. = B. Cheiranthos.

SINAPIDENDRON, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 36 = **Brassica**, Tourn. (Crucif.).
angustifolium, Lowe, Man. Fl. Mad. i. 30 = B. frutescens.
frutescens, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 37 = B. frutescens.
glaucum, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 267.—Ins. Cap. Virid.
gracile, Webb, in Hook. Niger Fl. 99. t. 1.—Ins. Cap. Virid.
rupestre, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 37 (= *Sinapis rupestris*).—Ins. Mader.
salicifolium, Lowe, l. c. = B. frutescens.
Vogelii, Webb, in Hook. Niger Fl. 100. t. 2.—Ins. Cap. Virid.

SINAPIS, Linn. Syst. ed. I (1735) = **Brassica**, Tourn. (Crucif.).
abyssinica, A. Br. ex Regel, Ind. Sem. Hort. Petrop. (1856) 34.—Hab.?
adpressa, Schloss. & Vukot. Fl. Croat. 237 = B. adpressa.
alba, Linn. Sp. Pl. 668 = B. alba.
Allionii, Jacq. Hort. Vindob. ii. 79 = B. oleracea.
alpina, Mart. ex Steud. Nom. ed. II. ii. 590 = Conringia alpina.
amplexicaulis, DC. Syst. ii. 610.—Alger.
angustifolia, DC. l. c. 624 = B. frutescens.
apula, Tenore, Prod. Fl. Nap. p. xxxix = Diplotaxis apula.
aristidis, Pomel, Nouv. Mat. Fl. Atl. 358.—Alger.
arvensis, Linn. Sp. Pl. 668 = B. Sinapistrum.
arvensis, O. F. Muell. in Fl. Dan. t. 678 = Raphanus Raphanistrum.
assurgens, Delile, Ind. Sem. Hort. Monsp. (1847) 7 = Diplotaxis pachypoda.
auriculata, DC. Syst. ii. 611 (pl. cult.).—Hab.?
baetica, Boiss. Elench. 11 = B. boetica.
benghalensis, Roxb. ex DC. Syst. ii. 198 = Nasturtium indicum.
bipinnata, Desf. Fl. Atlant. ii. 97 = Rapistrum bipinnatum.
bracteata, Steud. Nom. ed. II. ii. 590, sphalm. = seq.
bracteolata, G. Don, in Sweet, Hort. Brit. ed. III. 42 = B. bracteolata.
brassicata, Linn. Syst. ed. XII. 231 = B. juncea.
Burmanni, Eckl. ex Zeyh. Enum. 7 = *retrorsa*.
campestris, Bess. Prim. Fl. Galic. ii. 88.—Europ.

SINAPIS :—
campestris, Jacq. ex Steud. Nom. ed. II. ii. 590 = Brassica juncea.
caspica, Willd. ex Ledeb. Fl. Ross. i. 217 = B. Tournefortii.
cernua, Poir. Encyc. iv. 342 = B. juncea.
cernua, Thunb. Fl. Jap. 261 = B. nigra.
Cheiranthus, Mert. & Koch, Deutschl. Fl. iv. 717 = B. Cheiranthos.
chelidonifolia, Hort. ex Steud. Nom. ed. II. ii. 590, nomen (Quid ?).—Hab. ?
chinensis, Linn. Mant. i. 95 = B. juncea.
Choulettiana, Coss. & Dur. ex Coss. in Bull. Soc. Bot. Fr. ix. (1862) 295 = *procumbens*.
circinnata, Desf. Fl. Atlant. ii. 96 = *pubescens*.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = B. campestris.
crassifolia, Rafin. Caratteri, 78 = Diplotaxis pendula.
cuneifolia, Roxb. Hort. Beng. 48 ; Fl. Ind. iii. 121 = B. juncea.
dichotoma, Roxb. ex Flem. in As. Res. xi. (1810) 179 = B. campestris.
dissecta, Lag. Gen. et Sp. Nov. 20 = B. dissecta.
divaricata, Roxb. Hort. Beng. 48 ; Fl. Ind. iii. 123 = Nasturtium indicum.
elongata, Spach, ex Steud. Nom. ed. II. ii. 590 ; Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 308 = B. elongata.
Eruca, [Clairv.] Man. Herb. 220 = Eruca sativa.
erucoides, Linn. Cent. Pl. ii. 24 ; Amoen. Acad. iv. 322 = Diplotaxis erucoides.
erysimoides, Roxb. Hort. Beng. 48; Fl. Ind. iii. 123 = B. nigra.
exotica, Hort. ex DC. Syst. ii. 638 = Eruca sativa.
flexuosa, Poir. in Lam. Encyc. iv. 341 = B. hispida.
foliosa, Willd. Enum. Hort. Berol. 688 = B. alba.
frutescens, Ait. Hort. Kew. ed. I. ii. 404 = B. frutescens.
geniculata, Desf. Fl. Atlant. ii. 98 = B. adpressa.
glauca, Roxb. Hort. Beng. 48; Fl. Ind. iii. 118 = B. campestris.
Gorraea, Buch.-Ham. ex Wall. Cat. sub n. 4790 = B. sinapoides (in Addend.).
Harra, Forsk. Fl. Aegypt. Arab. 118 = Diplotaxis pendula.
hastata, Desf. Tabl. ed. II. 151 = Diplotaxis virgata.
heterophylla, Lag. Gen. et Sp. Nov. 20 = B. adpressa.
hispanica, Banks, ex DC. Prod. i. 230 = B. adpressa.
hispanica, Lag. ex DC. Syst. ii. 632 = Diplotaxis catholica.
hispanica, Linn. Sp. Pl. 669 = Erucaria tenuifolia.
hispanica, Thuill. Fl. Par. ed. II. 343 = Sisymbrium hispanicum.
hispida, Balb. Misc. Bot. 33 = B. Sinapistrum.
hispida, Schousb. Vextr. Marokko, 196 = B. hispida.
incana, Linn. Cent. Pl. i. 19 ; Amoen. Acad. iv. 280 = B. adpressa.
incana, Thuill. Fl. Par. ed. II. 343 = B. nigra.
indurata, Coss. in Bull. Soc. Bot. Fr. ix. (1862) 168. —Alger.
inflexa, Walp. Rep. i. 186 ; D. Dietr. Syn. Pl. iii. 725 = B. incana.
integrifolia, [Vahl, in] West, Ste-Croix, 296; Willd. Hort. Berol. 14. t. 14 = B. juncea.
integrifolia, Walp. Rep. i. 187 = B. incana ?
japonica, Thunb. Fl. Jap. 262 = B. juncea.
juncea, Linn. Sp. Pl. 668 = B. juncea.
Kaber, DC. Syst. ii. 617 = B. campestris.
laevigata, Burm. f. Fl. Cap. Prod. 18 = *retrorsa*.
laevigata, Linn. Cent. Pl. i. 20 ; Amoen. Acad. iv. 281 = B. juncea.
laevigata, Pall. Tabl. Taur. 54 = B. elongata.
laevigata, Poir. Encyc. iv. 345 = *auriculata*.
lanceolata, DC. Syst. ii. 611 = B. juncea.
leptopetala, DC. l. c. 610 = B. leptopetala.
longirostris, Boiss. ex Amo, Fl. Iber. vi. 472 = B. longirostra.
maritima, All. Fl. Pedem. i. 264 = Sisymbrium austriacum.
mesopotamica, Spreng. in Schrad. Journ. ii. 1800 (1801), 199 = Moricandia arvensis.
millefolia, Jacq. Coll. i. 41 = Sisymbrium millefolium.
monensis, Bab. Man. Brit. Bot. ed. II. 25 = B. monensis.
montana, Bourg. ex Nym. Consp. 47 = B. Cheiranthos.
muralis, R. Br. in Ait. Hort. Kew. ed. II. iv. 128 = Diplotaxis muralis.

SINAPIS :—
Napus, Brot. Fl. Lusit. i. 586 = Brassica campestris.
nasturtiifolia, Poir. in Lam. Encyc. iv. 346 = B. Erucastrum.
nigra, Forsk. ex DC. Syst. ii. 612 = B. juncea.
nigra, Linn. Sp. Pl. 668 = B. sinapoides (in Addend.).
nudicaulis, Lag. Gen. et Sp. Nov. 20 = B. oxyrrhina.
Nuttalliana, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 551.—Hab. ?
oleracea, Presl, Delic. Prag. 18 = B. juncea.
Oliveriana, DC. Syst. ii. 609 = Sisymbrium pannonicum.
orgyalis, Willd. ex Ledeb. Fl. Ross. i. 217 = B. sinapoides (in Addend.).
orientalis, Linn. Cent. Pl. i. 19 ; Amoen. Acad. v. 280 = B. Sinapistrum.
oxyrrhina, Coss. Notes Crit. 26 = B. oxyrrhina.
paniculata, Boiss. Fl. Orient. i. 390 = B. adpressa.
panormitana, Spreng. Syst. iv. Cur. Post. 244.—Sicilia.
parviflora, Lippi, ex Delile, Fl. Aeg. Illustr. 67, nomen. —Aegypt.
patens, Roxb. ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 139 = Nasturtium indicum.
patens, Roxb. Hort. Beng. 48 ; Fl. Ind. iii. 124 = B. juncea.
pekinensis, Lour. Fl. Cochinch. 400 = B. campestris.
pendula, E. Mey. in Drège, Zwei Pfl. Docum. 222, nomen.—Afr. austr.
philaeana, Delile, Fl. Egypte, 243 = Moretția canescens.
polymorpha, Geners. ex Schult. Obs. Bot. 133.—Hab. ?
procumbens, Poir. Encyc. Suppl. iv. 12.—Afr. bor.
prostrata, Roxb. Hort. Beng. 48.—China.
pubescens, Linn. Mant. i. 95.—Europ. austr.
pusilla, Roxb. Hort. Beng. 48 ; Fl. Ind. iii. 125 = Nasturtium montanum.
pyrenaica, Linn. Sp. Pl. ed. II. 934 = Sisymbrium austriacum.
pyrenaica, Spielm. ex Pourr. in Mém. Acad. Toul. iii. (1788) 329 (an praec. ?).—Mont. Pyren.
radicata, Desf. Fl. Atlant. ii. 98 = B. fruticulosa.
ramosa, Roxb. ex Flem. in As. Res. xi. (1810) 179 = B. juncea.
Rapa, Brot. Fl. Lusit. i. 586 = B. campestris.
Raphanistrum, Gueldenst. ex Ledeb. Fl. Ross. i. 225 = Raphanus Raphanistrum.
recurvata, All. Fl. Pedem. i. 265 = B. Cheiranthos.
retrohirsuta, Bess. ex Steud. Nom. ed. I. 782 = B. Sinapistrum.
retrorsa, Burch. ex DC. Syst. ii. 609.—Afr. austr.
rubella, Pall. Reise Suedl. Statth. Russ. Reich. i. 86, 125, 141, 221 = Chorispora tenella.
rugosa, Roxb. Hort. Beng. 48; Fl. Ind. iii. 122 = B. juncea.
rupestris, Steud. Nom. ed. II. ii. 591.—Ins. Mader.
Schkuhriana, Reichb. Ic. Fl. Germ. ii. 20 = B. Sinapistrum.
Schlosseri, Heuff. ex Nym. Consp. 47 = B. Sinapistrum.
setigera, J. Gay, ex Lange, in Kjoeb. Vidensk. Meddel. (1865) 76 = B. setigera.
sicula, Bertol. Fl. Ital. vii. 169 = B. boetica.
sinensis, J. F. Gmel. Syst. 989 = B. juncea ?
subbipinnatifida, Lag. Gen. et Sp. Nov. 20.—Hispan.
subularia, Sweet, Hort. Brit. ed. II. 34, nomen.—Hab. ?
taurica, Bieb. Fl. Taur. Cauc. iii. 450 = B. adpressa.
taurica, Fisch. Cat. Hort. Gorenk. (1812) 51 = B. Sinapistrum.
tenella, Moench, Meth. 253 = B. juncea.
tenuifolia, R. Br. in Ait. Hort. Kew. ed. II. iv. 128 = Diplotaxis tenuifolia.
tetraëdra, J. & C. Presl, Delic. Prag. 19 = Erysimum cuspidatum.
timoriana, DC. Syst. ii. 616 = B. juncea.
torosa, Gilib. Fl. Lituan. ii. 69 = B. Sinapistrum.
torulosa, Pers. Syn. ii. 207 = B. nigra.
Tournefortii, All. Fl. Pedem. i. 265 = B. Cheiranthos.
trilocularis, Roxb. Hort. Beng. 48 ; Fl. Ind. iii. 121 = B. trilocularis.
tripinnata, Burch. Trav. i. 318 = Sisymbrium tripinnatum.
tuberosa, Poir. in Lam. Encyc. iv. 346 = B. campestris.
turgida, Delile, Fl. Aeg. Illustr. 67.—Aegypt.
turgida, Pers. Syn. ii. 207 = B. sinapoides (in Addend.).
ucranica, Czern. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 312.—Rossia.
villosa, Mérat, Fl. Par. 215 ; ed. II. ii. 311 = B. sinapoides (in Addend.).
virgata, Cav. Desc. 428 = Diplotaxis virgata.
virgata, J. & C. Presl, Delic. Prag. 19 = B. juncea.
zeylanica, Hort. ex Steud. Nom. ed. II. ii. 591, nomen (Quid ?).—Hab. ?

SINAPISTRUM, Medic. Phil. Bot. i. 108 (1789) = **Gynandropsis**, DC. (Cappar.).
pentaphyllum, Medic. l. c. = G. pentaphylla.

SINAPISTRUM, Spach, Hist. Vég. Phan. vi. 343 (1838) = **Brassica**, Tourn. (Crucif.).
arvense, Spach, l. c. 345 = B. oleracea, arvensis, Conringia orientalis, Eruca sativa.

SINAPISTRUM, Tourn. ex Hall. in Rupp. Fl. Jen. 269 (1745) = **Cleome**, Linn. (Cappar.).
decandrum, Moench, Meth. 250 = C. graveolens.
giganteum, Moench, l. c. = C. gigantea.
lanatum, Moench, l. c. = C. arabica.
violaceum, Moench, Meth. Suppl. 83 = C. graveolens.
viscosum, Moench, l. c. = C. viscosa.

SINAPODENDRON, Ball, in Journ. Linn. Soc. xvi. (1877) 325, sphalm. = SINAPIDENDRON, Lowe = **Brassica**, Tourn. (Crucif.).

SINCARPIA, Tenore, in Mem. Soc. Ital. Moden. xxii. (1841) 230 = **Syncarpia**, Tenore (Myrtac.).

SINCLAIREA, Sch.-Bip. in Flora, xxxvi. (1853) 34 = seq.

SINCLAIRIA, Hook. & Arn. Bot. Beech. Voy. 433 (1841) = **Liabum**, Adans. (Compos.).
andromachioides, Sch.-Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 231 = L. andromachioides.
discolor, Hook. & Arn. Bot. Beech. Voy. 433 = L. discolor.

SINDORA, Miq. Fl. Ind. Bat. Suppl. 287 (1860).
LEGUMINOSAE, Benth. & Hook. f. i. 584.
ECHINOCALYX, Benth. in Benth. & Hook. f. Gen. i. 584 (1865).
cochinchinensis, *Baill. Adansonia*, x. (1871) 104.—Cochinch.
siamensis, Teysm. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 86 = Wallichii.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 288.—Sumatra.
velutina, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 269.—Malacca.
Wallichii, *Benth. in Hook. Ic. Pl.* t. 1017–18.—Singapore ; Malacca.

SINGANA, Aubl. Pl. Gui. i. 574. t. 230 (1775). Genus dubium *LEGUMINOSARUM*.—Cf. Benth. & Hook. f. i. 465.
MYLOSPHORA, Neck. Elem. ii. 280 (1790).
STERBECKIA, Schreb. Gen. i. 360 (1789).
guianensis, *Aubl. Pl. Gui.* i. 574.—Guian.

SINGLINGIA, Benth. in Journ. Linn. Soc. xix. (1881) 19, sphalm. = SIEGLINGIA, Bernh. = **Triodia**, R. Br. (Gramin.).

SINISTROPHORUM, Schrank, ex Endl. Gen. 882 (1840) = **Camelina**, Crantz (Crucif.).

SINNINGIA, Nees, in Ann. Sc. Nat. Sér. I. vi. (1825) 297. *GESNERACEAE*, Benth. & Hook. f. ii. 1004.
BIGLANDULARIA, Seem. in Gard. Chron. (1868) 738.
LIGERIA, Decne. in Rev. Hortic. (1848) 463.
ROSANOWIA, Regel, Gartenfl. xxi. (1872) 33. t. 712.
STENOGASTRA, Hanst. in Linnaea, xxvi. (1853–56) 205.
TAPEINOTES, DC. Prod. vii. 544 (1839).
TAPINA, Mart. Nov. Gen. et Sp. iii. 59. t. 225. f. 1 (1829).
calycina, Hort. ex G. Don, in Loud. Hort. Brit. 241 = Pentarhaphia calycina.
Carolinae, *Benth. & Hook. f. Gen.* ii. 1004.—Bras.
floribunda, *Hort. Par. ex A. Dietr. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 242.—Hab. ?
gloxiniflora, *Scheidw. l. c.* xv. (1847) 225.—Mexic.
guttata, *Lindl. Bot. Reg.* t. 1112.—Bras.
Helleri, Lindl. l. c. t. 997 = Pentarhaphia calycina.
Helleri, *Nees, in Ann. Sc. Nat.* Sér. I. vi. (1825) 297. —Bras.
Lindleyi, Schau. in Flora, xvii. (1834) 642 = Pentarhaphia calycina.
punctata, *Ysabeau, in Journ. Hort. Prat.* vii. (1850) 331.—Bras.

SINNINGIA:—
sanguinea, Regel, Cat. Pl. Hort. Aksakov. 132, nomen (Quid ?).—Hab. ?
speciosa, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 91.—Bras.
velutina, Hook. Bot. Mag. t. 4212 = Helleri.
velutina, *Lindl. Bot. Reg.* sub t. 1112.—Bras.
villosa, *Lindl. l. c.* tt. 1112, 1134.—Bras.
Youngeana ×, Marnock, ex Paxt. Mag. Bot. vii. (1840) 51.

SIOJA, Buch.-Ham. ex Lindl. Introd. Nat. Syst. ed. II. 82 (1836) = **Cardiopteris**, Wall. (Olacin.).
sanguinaria, Buch.-Ham. ex Wall. Cat. sub n. 8033 = C. lobata.

SIOLMATRA, Baill. in Bull. Soc. Linn. Par. i. (1885) 458. *CUCURBITACEAE.*
brasiliensis, *Baill. l. c.*—Bras.

SION, Adans. Fam. ii. 498 (1763) = **Sium**, Tourn. (Umbell.).

SIONA, Salisb. Gen. Pl. Fragm. 67 (1866) = **Dichopogon**, Kunth (Liliac.).
fimbriata, Salisb. l. c. = D. Sieberianus ?

SIPANEA, Aubl. Pl Gui. i. 147. t. 56 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 52.
PTYCHODEA, Willd. ex Cham. & Schlecht. in Linnaea, iv. (1829) 168.
VIRECTA, Linn. f. Suppl. 17 (1781).
angustifolia, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 276 = Virecta multiflora ?
biflora, *Cham. & Schlecht. in Linnaea*, iv. (1829) 168.—Am. trop.
carnea, *Neumann, in Rev. Hortic.* Sér. II. ii. (1843–44) 445.—Am. austr.
dichotoma, H. B. & K. Nov. Gen. et Sp. iii. 397 = pratensis.
elatior, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 276 = Virecta procumbens ?
erythraeoides, *Cham. in Linnaea*, ix. (1834) 242.—Bras.
glomerata, *H. B. & K. Nov. Gen. et Sp.* iii. 398.—Venezuela.
limnophila, Spruce, ex Hook. f. in Hook. Ic. Pl. sub t. 1050 = Limnosipanea Spruceana.
palustris, Seem. Bot. Voy. Herald, 136 = Limnosipanea palustris.
pratensis, *Aubl. Pl. Gui.* i. 147. t. 56.—Guian.
radicans, Endl. Atakt. Bot. 7. t. 7 = biflora.
trichantha, Miq. in Linnaea, xviii. (1844) 293 = pratensis.

SIPANIA, Seem. Bot. Voy. Herald, 136 (1853) = **Limnosipanea**, Hook. f. (Rubiac.).

SIPARUNA, Aubl. Pl. Gui. ii. 864. t. 333 (1775). *MONIMIACEAE*, Benth. & Hook. f. iii. 144.
CITROSMA, Ruiz & Pav. Prod. 134. t. 29 (1794).
LEONIA, Mutis, ex Kunth, Syn. Pl. i. 462 (1822).
alternifolia, *A. DC. in DC. Prod.* xvi. II. 656.—Bras.
amazonica, *Mart. ex A. DC. l. c.* 653.—Bras.
amplifolia, *A. DC. l. c.* 649.—N. Granat.
andina, *A. DC. l. c.* 648.—Mexic.
apicifera, *A. DC. l. c.* 652.—Ecuador.
Apiosyce, *A. DC. l. c.* 645.—Bras.
aspera, *A. DC. l. c.*—Peruv.
asperula, *A. DC. l. c.* 647.—Bras.
auriculata, *A. DC. in Journ. Bot.* iii. (1865) 219.—Peruv.
bifida, *A. DC. in DC. Prod.* xvi. II. 652.—Bras.
brasiliensis, *A. DC. l. c.* 656.—Bras.
budleiaefolia, *A. DC. l. c.* 652.—Ecuador.
camporum, *A. DC. l. c.* 654.—Bras.
cauliflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 69.—Panama.
chiridota, *A. DC. in DC. Prod.* xvi. II. 649.—N. Granat.
cristata, *A. DC. l. c.* 655.—Bras.
cujabana, *A. DC. l. c.* 644.—Bras.
cuspidata, *A. DC. l. c.* 655.—Guian.
decipiens, *A. DC. l. c.* 643.—Guian.
discolor, *A. DC. l. c.* 656.—Am. trop.
echinata, *A. DC. l. c.* 652.—N. Granat.
eriocalyx, *A. DC. l. c.* 646.—Peruv.
erythrocarpa, *A. DC. l. c.* 643.—Bras.
estrellensis, *A. DC. l. c.* 644.—Bras.
foliosa, *A. DC. l. c.* 650.—Bras.

SIPARUNA:—
fulva, *A. DC. in Journ. Bot.* iii. (1865) 219.—Peruv.
gesnerioides, *A. DC. in DC. Prod.* xvi. II. 646.—N. Granat.
glabra, *A. DC. l. c.* 656.—Bras.
glabrescens, *A. DC. l. c.* 648.—Ind. occ.
Goudotiana, *A. DC. l. c.* 651.—N. Granat.
guianensis, *Aubl. Pl. Gui.* ii. 865. t. 333.—Guian.
hispida, *A. DC. in Journ. Bot.* iii. (1865) 219.—Peruv.; Boliv.
Holtonii, *A. DC. in DC. Prod.* xvi. II. 652.—N. Granat.
Kunthii, *A. DC. l. c.* 649.—N. Granat.
Lagopus, *A. DC. l. c.* 648.—N. Granat.
lanceolata, *A. DC. l. c.* 644.—Bras.
laurifolia, *A. DC. l. c.* 651.—N. Granat.
lepidota, *A. DC. l. c.*—N. Granat.
limoniodora, *A. DC. l. c.* 646.—Peruv.
Lindeni, *A. DC. l. c.* 656.—Bras.
macrophylla, *A. DC. l. c.* 646.—N. Granat.
micrantha, *A. DC. in Journ. Bot.* iii. (1865) 219.—Bras.
mollicoma, *A. DC. in DC. Prod.* xvi. II. 653.—Bras.
mollis, *A. DC. l. c.*—N. Granat.
muricata, *A. DC. l. c.* 651.—Peruv.
Mutisii, *A. DC. l. c.* 650.—N. Granat.
neglecta, *A. DC. l. c.* 647.—Peruv.
nicaraguensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 69.—Nicaragua.
oblongifolia, *A. DC. in DC. Prod.* xvi. II. 656.—Peruv.
obovata, *A. DC. l. c.* 644.—Bras.
ovalis, *A. DC. l. c.* 650.—Peruv.
panamensis, *A. DC. in Journ. Bot.* iii. (1865) 219.—Panama.
paniculata, *A. DC. in DC. Prod.* xvi. II. 656.—Bras.
pauciflora, *A. DC. l. c.*—Panama.
pellita, *A. DC. l. c.* 645.—Bolivia.
petiolaris, *A. DC. l. c.* 651.—N. Granat.
Poeppigii, *A. DC. l. c.* 653.—Bras.
polyantha, *A. DC. l. c.* 646.—Bolivia.
radiata, *A. DC. l. c.* 656.—Peruv.
Reginae, *A. DC. l. c.* 654.—Bras.
riparia, *A. DC. l. c.* 647.—Mexic.
ruficeps, *A. DC. l. c.* 645.—Bras.
sessiliflora, *A. DC. l. c.* 647.—N. Granat.
Sprucei, *A. DC. in Journ. Bot.* iii. (1865) 219.—N. Granat.
suaveolens, *A. DC. in DC. Prod.* xvi. II. 647.—Peruv.
subinodora, *A. DC. l. c.* 649.—Peruv.
Thea, *A. DC. l. c.* 657.—Bras.
thecaphora, *A. DC. l. c.* 657.—Peruv.
tomentosa, *A. DC. l. c.*—Peruv.
Trianaei, *A. DC. in Journ. Bot.* iii. (1865) 219.—N. Granat.

SIPHANTHERA, Pohl, Pl. Bras. Ic. i. 102. t. 84, 85 (1827). *MELASTOMACEAE*, [Benth. & Hook. f. i. 739].
MEISNERIA, DC. Prod. iii. 114 (1828).
arenaria, *Cogn. in Mart. Fl. Bras.* xiv. III. 193.—Bras.
cordata, *Pohl, Pl. Bras. Ic.* i. 103. t. 84.—Bras.
discolor, *Cogn. in Mart. Fl. Bras.* xiv. III. 194.—Bras.
gracilis, Miq. in Linnaea, xxii. (1849) 537 = cordata, Miqueliana.
Hostmannii, *Cogn. in Mart. Fl. Bras.* xiv. III. 199.—Guian.
microlicioides, *Cogn. l. c.* 194.—Guian.
Miqueliana, *Cogn. l. c.* 197.—Bras.
paludosa, *Cogn. l. c.* 191.—Bras.
pusilla, *Cogn. l. c.* 197.—Bras.
robusta, *Cogn. l. c.* 195.—Bras.
subtilis, *Pohl, Pl. Bras. Ic.* i. 105. t. 85. f. 2.—Bras.
tenera, *Pohl, l. c.* i. 104. t. 85. f. 1.—Bras.
villosa, *Cogn. in Mart. Fl. Bras.* xiv. III. 192.—Bras.

SIPHAULAX, Rafin. Fl. Tellur. iii. 74 (1836) = **Nicotiana**, Linn. (Solan.).
glabra, Rafin. l. c.—Hab. ?

SIPHIDIA, Reichb. Handb. 173 (1837), sphalm. = seq.

SIPHISIA, Rafin. Med. Fl. i. 62 (1828) = **Aristolochia**, Tourn.
angustifolia, Klotzsch, in Monatsb. Akad. Berl. (1859) 603 = A. saccata.
glabra, Rafin. Med. Fl. i. 65 = A. macrophylla.

SIPHISIA:—
Griffithii, Klotzsch, ex Duch. in DC. Prod. xv. I. 437 = Aristolochia Griffithii.
macrophylla, Aschers. Fl. Brandenb. i. 598 = A. macrophylla.
platanifolia, Klotzsch, in Monatsb. Akad. Berl. (1859) 603 = A. platanifolia.
reticulata, Klotzsch, l. c. 604 = A. reticulata.
saccata, Klotzsch, l. c. 603 = A. saccata.
sericea, Klotzsch, l. c. = A. sericea, *Benth.*
Sipho, Klotzsch, l. c. 602 = A. macrophylla.
Sipho, Rafin. Med. Fl. i. 62 = A. Sipho.
Thwaitesii, Klotzsch, in Monatsb. Akad. Berl. (1859) 604 = A. Thwaitesii.
tomentosa, Rafin. Med. Fl. i. 62, 65 = A. tomentosa.
tripteris, Rafin. l. c. 62 = A. tripteris.

SIPHOCAMPYLUS, Pohl, Pl. Bras. Ic. ii. 104. tt. 169–177 (nec t. 168) (1831). *CAMPANULACEAE*, Benth. & Hook. f. ii. 547.
BYRSANTHES, Presl, Prod. Monog. Lobel. 41 (1836).
CANONANTHUS, G. Don, Gen. Syst. iii. 718 (1834).
CREMOCHILUS, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 174.
LOBELIA, Presl, Prod. Monog. Lobel. 33 (1836).
acuminatus, G. Don, in Sweet, Hort. Brit. ed. III. 424 = Lobelia acuminata.
amoenus, Planch. in Fl. des Serres, vi. (1850–51) 273. t. 619 = villosulus.
Andropogon, G. Don, Gen. Syst. iii. 703 = Centropogon Andropogon.
angustiflorus, *Schlecht. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
asper, *Benth. Pl. Hartw.* 213.—N. Granat.
attenuatus, *A. DC. in DC. Prod.* vii. 398.—Peruv.
ayavacensis, Steud. Nom. ed. II. ii. 63 = umbellatus.
barbatus, G. Don, Gen. Syst. iii. 701 = Centropogon barbatus (in Addend.).
Benthamianus, *Walp. Rep.* vi. 379.—N. Granat.
Berterianus, G. Don, Gen. Syst. iii. 703 = Centropogon Berterianus.
beslerioides, G. Don, l. c. 702 = Centropogon beslerioides.
betulaefolius, *G. Don, l. c.* 703.—Bras.
bicolor, G. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 389 = Lobelia laxiflora.
biserratus, *A. DC. in DC. Prod.* vii. 397.—Peruv.
bracteatus, *Decne. ex Linden, Cat.* n. 5. (1850) 7.—N. Granat.
brevidens, *Regel, Ind. Sem. Hort. Petrop.* (1868) 84.—Am. subtrop.
campanulatus, *A. DC. in DC. Prod.* vii. 401.—Ecuador.
canescens, *A. DC. l. c.* 402.—Mexic.
canus, Pohl, Pl. Bras. Ic. ii. 106. t. 169 = macropodus.
Caoutchouc, *G. Don, Gen. Syst.* iii. 701.—N. Granat.
cardiophyllus, Pohl, Pl. Bras. Ic. ii. 110. t. 173 = corymbiferus.
Cavanillesianus, G. Don, Gen. Syst. iii. 702 = biserratus.
cernuus, *Griseb. Cat. Pl. Cub.* 159.—Cuba.
Chamissonianus, A. DC. in DC. Prod. vii. 407 = Westinianus.
chimapilophyllus, *Schlecht. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
cirsiifolius, G. Don, Gen. Syst. iii. 703 = Lobelia cirsiifolia.
coccineus, Hook. Bot. Mag. t. 4178 = Centropogon coccineus.
coleoides, *Vatke, in Linnaea*, xxxviii. (1874) 733.—Costa Rica.
Columnae, *G. Don, Gen. Syst.* iii. 701.—N. Granat.
comosus, *G. Don, l. c.* 702.—Mexic.
convolvulaceus, *G. Don, l. c.* 703.—Bras.
cordifolius, Benth. Pl. Hartw. 213 = Benthamianus.
cordifolius, *Otto & Dietr. Allg. Gartenz.* xii. (1844) 371.—Hab. ?
corymbiferus, *Pohl, Pl. Bras. Ic.* ii. 112. t. 175.—Bras.
corymbiflorus, Warsz. in C. Koch, Wochenschr. vii. (1864) 60 = Centropogon Warscewiczii.
corymbosus, Steud. Nom. ed. II. ii. 59, sphalm., in syn. = praec.
Costaricae, *Vatke, in Linnaea*, xxxviii. (1874) 731.—Costa Rica.
crenatifolius, *Pohl, Pl. Bras. Ic.* ii. 107. t. 170.—Bras.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 68.—Cuba.
decumbens, *Juss. ex A. DC. in DC. Prod.* vii. 397.—Ins. S. Doming.

SIPHOCAMPYLUS:—
densiflorus, *Planch. in Fl. des Serres*, vi. (1850–51) 18.—N. Granat.
denticulosus, *Planch. l. c.* 19.—N. Granat.
dependens, *G. Don, Gen. Syst.* iii. 704.—Peruv.
divaricatus, *Benth. Pl. Hartw.* 138.—Peruv.
Dombeyanus, *A. DC. in DC. Prod.* vii. 406.—Peruv.
domingensis, *A. DC. l. c.* 397.—Ins. S. Doming.
duploserratus, Cham. in Linnaea, viii. (1833) 201 = Westinianus.
duploserratus, *Pohl, Pl. Bras. Ic.* ii. 114. t. 177.—Bras.
Eichleri, *Kanitz, in Mart. Fl. Bras.* vi. IV. 148.—Bras.
elegans, *Planch. in Fl. des Serres*, vi. (1850–51) 19.—N. Granat.
ellipticus, *Vatke, in Linnaea*, xxxviii. (1874) 724.—Am. austr.
erianthus, Benth. Pl. Hartw. 139 = Centropogon erianthus (in Addend.).
eximius, *Planch. in Fl. des Serres*, vi. (1850–51) 16.—N. Granat.
ferrugineus, *G. Don, Gen. Syst.* iii. 701.—N. Granat.
fimbriatus, Regel, Gartenfl. xvii. (1868) 355. t. 600; et in Ind. Sem. Hort. Petrop. (1868) 84 = longepedunculatus.
floccosus, *Planch. ex Linden, Cat.* n. 5. (1850) 7.—N. Granat.
foetidus, *G. Don, Gen. Syst.* iii. 702.—N. Granat.
foliosus, *Griseb. in Goett. Abh.* xix. (1874) 201.—Reg. Argent.
fulgens, *Lebas, in Rev. Hortic.* (1872) 118.—Hab. ?
Funckeanus, *Planch. in Fl. des Serres*, vi. (1850–51) 19.—N. Granat.
fuscus, *G. Don, Gen. Syst.* iii. 704.—Peruv.
giganteus, *G. Don, l. c.* 702.—Ecuador.
glabratus, *G. Don, l. c.* 702.—N. Granat.
glandulosus, *Hook. Bot. Mag.* t. 4331.—N. Granat.
grandidentatus, *A. DC. in DC. Prod.* vii. 407.—Mexic.
grandis, G. Don, Gen. Syst. iii. 702 = Centropogon grandis.
guatemalensis, Vatke, in Linnaea, xxxviii. (1874) 730 = Centropogon cordifolius.
Gutierrezii, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 155.—Am. centr.
Hallianus, *Vatke, in Linnaea*, xxxviii. (1874) 732.—Ecuador.
hamatus, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 138.—Bras.
Hartwegi, Benth. Pl Hartw. 139 = Centropogon Hartwegi (in Addend.).
hirtus, G. Don, Gen. Syst. iii. 702 = Centropogon hirtus.
hispidus, *Benth. Pl. Hartw.* 214.—N. Granat.
Humboldtianus, *A. DC. in DC. Prod.* vii. 398.—Peruv.
imbricatus, *G. Don, Gen. Syst.* iii. 703.—Bras.
Jamesonianus, *A. DC. in DC. Prod.* vii. 402.—Ecuador.
Kraussii, G. Don, in Sweet, Hort. Brit. ed. III. 424 = Lobelia Kraussii.
laciniatus, *G. Don, Gen. Syst.* iii. 704.—Peruv.
laevigatus, *Planch. in Fl. des Serres*, vi. (1850–51) 34.—N. Granat.
Lamarckii, *A. DC. in DC. Prod.* vii. 397.—Ins. S. Doming.
lanatus, *Benth. Pl. Hartw.* 214.—N. Granat.
lanceolatus, Regel, Cat. Pl. Hort. Aksakov. 132, nomen (Quid ?).—Hab. ?
lantanifolius, *A. DC. in DC. Prod.* vii. 399.—Venezuela.
lasiandrus, *Planch. in Fl. des Serres*, vi. (1850–51) 36.—N. Granat.
Lindleyi, *Lem. Jard. Fleur.* ii. (1852) t. 142.—N. Granat.
longepedunculatus, *Pohl, Pl. Bras. Ic.* ii. 109. t. 172.—Bras.
longipes, Van Houtte, ex Regel, Gartenfl. (1854) 3; Vatke, in Linnaea, xxxviii. (1874) 733 = Centropogon longipes.
lycioides, *G. Don, Gen. Syst.* iii. 703.—Bras.
macranthus, Pohl, Pl. Bras. Ic. ii. 105. t. 168 = Centropogon surinamensis.
macrophyllus, *G. Don, Gen. Syst.* iii. 704.—Peruv.
macropodus, *G. Don, l. c.* 702.—Bras.
macrostemon, *A. DC. in DC. Prod.* vii. 403.—Bras.
manettiaeflorus, *Hook. Bot. Mag.* t. 4403.—N. Granat.?
Matthiaei, *A. DC. in DC. Prod.* vii. 405.—Peruv.

SIPHOCAMPYLUS:—
microstoma, *Hook. Bot. Mag.* t. 4286.—N. Granat.
microstoma, Lindl. & Paxt. Flow. Gard. ii. (1851) t. 44 = Lindleyi.
mollis, *Planch. in Fl. des Serres*, vi. (1850–51) 36.—Venezuela.
mollis, *Regel, in Flora*, xxxiii. (1850) 353.—Guatemala.
Mutisianus, *G. Don, Gen. Syst.* iii. 703.—N. Granat.
nemoralis, *Griseb. in Goett. Abh.* xix. (1874) 201.—Reg. Argent.
nitidus, *De Jonghe, ex C. Morr. in Ann. Soc. Hort. Gand*, ii. (1846) 319. t. 78.—Cuba.
nitidus, Hort. ex Hook. Bot. Mag. sub t. 4403 = manettiaeflorus.
nitidus, *Pohl, Pl. Bras. Ic.* ii. 110. t. 174.—Bras.
niveus, *Vatke, in Linnaea*, xxxviii. (1874) 732.—N. Granat.
odontosepalus, *Vatke, l. c.* 729.—N. Granat.
Orbignianus, *A. DC. in DC. Prod.* vii. 405.—Bolivia.
patens, *Griseb. Cat. Pl. Cub.* 159.—Cuba.
penduliflorus, *Decne. ex Planch. in Fl. des Serres*, viii. (1852–53) 35. t. 763.—Venezuela.
pendulifolius, *A. DC. in DC. Prod.* vii. 398.—Peruv.
persicifolius, G. Don, in Sweet, Hort. Brit. ed. III. 424 = Lobelia persicifolia.
peruvianus, *A. DC. in DC. Prod.* vii. 401.—Peruv.
polyphyllus, *Planch. in Fl. des Serres*, vi. (1850–51) 36.—N. Granat.
popayanensis, Benth. Pl. Hartw. 213 = Centropogon popayanensis (in Addend.).
prunifolius, *A. DC. in DC. Prod.* vii. 401.—Am. austr.
psilophyllus, *Pohl, Pl. Bras. Ic.* ii. 113. t. 176.—Bras.
pubescens, *Benth. Pl. Hartw.* 139.—Peruv.
pulchellus, *Planch. & Linden, Cat.* n. 10 (1855) 7.—N. Granat.
Purdiaeanus, *Planch. in Fl. des Serres*, vi. (1850–51) 18.—N. Granat.
Regelii, *Vatke, in Linnaea*, xxxviii. (1874) 732.—Costa Rica.
resinosus, *Schlecht. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
reticulatus, *Klotzsch & Karst. ex Vatke, in Linnaea*, xxxviii. (1874) 728.—Venezuela.
retrorsus, *Vatke, l. c.* 724.—Am. austr.
revolutus, *R. Grah. in Edinb. N. Phil. Journ.* xxx. (Apr. 1841) 429.—Hab. ?
rosmarinifolius, *G. Don, Gen. Syst.* iii. 704.—Peruv.
rugosus, *A. DC. in DC. Prod.* vii. 399.—Peruv.
rugosus, *Poit. in Rev. Hortic.* Sér. II. i. (1842) 186.—Hab. ?
scaber, A. DC. in DC. Prod. vii. 399 = longepedunculatus.
scabrifolius, *Schlecht. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
scandens, *G. Don, Gen. Syst.* iii. 703.—Peruv.
Sceptrum, *Decne. ex Linden, Cat.* n. 5. (1850) 7.—N. Granat.
Schlimmianus, *Planch. in Fl. des Serres*, vi. (1850–51) 34.—N. Granat.
Sellowianus, A. DC. in DC. Prod. vii. 407 = Westinianus.
sessilifolius, *Vatke, in Linnaea*, xxxviii. (1874) 731.—N. Granat.
spectabilis, G. Don, Gen. Syst. iii. 702 = Centropogon surinamensis.
subcarnosus, *Benth. Pl. Hartw.* 138.—Ecuador.
surinamensis, G. Don, Gen. Syst. iii. 702 = Centropogon surinamensis.
Thomesianus, Moric. Pl. Nouv. Am. 142. t. 85 = imbricatus.
thysanopetalus, *Vatke, in Linnaea*, xxxviii. (1874) 731.—Costa Rica.
umbellatus, *G. Don, Gen. Syst.* iii. 702.—Peruv.
verbascifolius, *A. DC. in DC. Prod.* vii. 402.—Peruv.
verticillatus, *G. Don, Gen. Syst.* iii. 703.—Bras.
villosulus, *Pohl, Pl. Bras. Ic.* ii. 108. t. 171.—Bras.
virgatus, *A. DC. in DC. Prod.* vii. 398.—Peruv.
volubilis, *G. Don, Gen. Syst.* iii. 703.—Venezuela.
Warmingii, *Kanitz, in Mart. Fl. Bras.* vi. IV. 148.—Bras.
Warszewiczii, *Regel, in Schweiz. Zeitschr. Gartenb.* (1850) 131; *ex Walp. Ann.* v. 390.—Guatemala.
Westianus, G. Don, Gen. Syst. iii. 702 = seq.
Westinianus, *Pohl, Pl. Bras. Ic.* ii. 115.—Bras.

SIPHOCODON, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 175. *CAMPANULACEAE*, Benth. & Hook. f. ii. 560.
spartioides, *Turcz. l. c.* 176.—Afr. austr.

SIPHOMERIS, Boj. Hort. Maurit. 170 (1837) = **Lecontea**, A. Rich. (Rubiac.).
argentifolia, Boj. l. c. 171 = L. argentea.
foetens, Hiern, in Oliver, Fl. Trop. Afr. iii. 229.—Afr. trop.
Lingun, Boj. Hort. Maurit. 170 = L. Bojeriana.

SIPHONACANTHUS, Nees, in Mart. Fl. Bras. ix. 45. t. 1 (1847) = **Ruellia**, Linn. (Acanth.).
densus, Nees, l. c. 47 = R. densa.
diffusus, Nees, l. c. 46.—Bras.
erythropus, Nees, l. c. 47.—Bras.
pubens, Nees, l. c. 46.—Bras.
repens, Nees, l. c. 47 (= *Stephanophysum Baikiei*).—Bras.
rubiginosus, Nees, l. c. 46.—Bras.
villosus, Nees, l. c. = R. villosa.

SIPHONANDRA, Klotzsch, in Linnaea, xxiv. (1851) 24 = **Ceratostema**, Juss. (Vaccin.).
elliptica, Klotzsch, l. c. = C. ellipticum.

SIPHONANDRA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 581 = **Chiococca**, Linn. (Rubiac.).
mexicana, Turcz. l. c. 582.—Mexic.

SIPHONANTHUS, Linn. Gen. ed. II. 526 (1742) = **Clerodendron**, Linn. (Verben.).
angustifolia, Willd. Sp. Pl. i. 606 = C. Siphonanthus.
hastata, Roxb. Hort. Beng. 46; Fl. Ind. iii. 67 = C. hastatum.
indica, Linn. Sp. Pl. 109 = C. Siphonanthus.

SIPHONANTHUS, Schreb. ex Baill. Etud. Gén. Euph. 324 (1858) = **Hevea**, Aubl. (Euphorb.).
elasticus, Schreb. ex Baill. l. c. 326 = H. guianensis.

SIPHONEUGENA, Berg, in Linnaea, xxvii. (1854) 344 = **Eugenia**, Mich. (Myrtac.).
aromatica, Berg, l. c. 345; et in Mart. Fl. Bras. xiv. I. 379 = E. aromatica, *Sond.* (in Addend.).
densiflora, Berg, ll. cc.—Bras.
Widgreniana, Berg, ll. cc.—Bras.

SIPHONIA, Benth. Pl. Hartw. 84 (1841) = **Lindenia**, Benth. (Rubiac.).
rivalis, Benth. l. c. = L. rivalis.

SIPHONIA, Rich. in Schreb. Gen. 656 (1791) = **Hevea**, Aubl. (Euphorb.).
apiculata, Spruce, ex Baill. Adansonia, iv. (1863–64) 285 = H. lutea.
brasiliensis, *H. B. & K. Nov. Gen. et Sp.* vii. 171 = H. brasiliensis.
Cahuchu, Rich. ex Willd. Sp. Pl. iv. 567 = H. guianensis.
discolor, Spruce, ex Benth. in Hook. Kew Journ. vi. (1854) 369 = H. discolor.
elastica, Pers. Syn. ii. 588 = H. guianensis.
guyanensis, Juss. ex Baill. Etud. Gén. Euph. 326 = H. guianensis.
Kunthiana, Baill. l. c. = H. brasiliensis.
lutea, Spruce, ex Benth. in Hook. Kew Journ. vi. (1854) 370 = H. lutea.
pauciflora, Spruce, ex Benth. l. c. = H. pauciflora.
rigidifolia, Spruce, ex Benth. l. c. 371 = H. rigidifolia.
Spruceana, Benth. l. c. 370 = H. Spruceana.

SIPHONIDIUM, Armstr. in Trans. N. Z. Inst. xiii. (1881) 340. *SCROPHULARINEAE.*
longiflorum, *Armstr. l. c.* 341.—N. Zel.

SIPHONIOPSIS, Karst. Fl. Columb. i. 139. t. 69 (1860 ?) = **Cola**, Schott (Stercul.).
monoica, Karst. l. c. = C. acuminata ?

SIPHONODON, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 246. t. 14. *CELASTRINEAE*, Benth. & Hook. f. i. 370.
ASTROGYNE, Wall. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 629 (1875).
australis, *Benth. Fl. Austral.* i. 403.—Austral.
celastrineus, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 247.—Burma; Malaya.

SIPHONOGLOSSA, Oerst. in Kjoeb. Vidensk. Meddel. 159 (1854). *ACANTHACEAE*, Benth. & Hook. f. ii. 1110.
longiflora, *A. Gray, Syn. Fl. N. Am.* ii. 1. 328.—Am. bor. occ.
nummularia, *S. Moore, in Journ. Bot.* xviii. (1880) 40. —Afr. austr.
pilosella, *Torr. Bot. Mex. Bound.* 124.—Arizona.
ramosa, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 160. t. 5.—Mexic.

SIPHONOGYNE, Cass. in Dict. Sc. Nat. l. 493 (1827) = **Eriocephalus**, Linn. (Compos.).

SIPHONOSTEGIA, Benth. in Hook. & Arn. Bot. Beech. Voy. 203. t. 44 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 975.
LESQUEREUXIA, Boiss. & Reut. in Boiss. Diagn. Ser. II. xii. 43 (1853).
PRISMATANTHUS, Hook. & Arn. Bot. Beech. Voy. 203 (1835).
chinensis, *Benth. in Hook. & Arn. Bot. Beech. Voy.* 203. t. 44.—China.
laeta, *S. Moore, in Journ. Bot.* xviii. (1880) 5.—China.
syriaca, *Boiss. Fl. Orient.* iv. 471.—Cilicia.

SIPHONOSTEMA, Griseb. in Lechl. Berb. Am. austr. 58 (1857) = **Ceratostema**, Juss. (Vaccin.).
costatum, Griseb. l. c. = C. costatum.
myrtifolium, Griseb. l. c. = C. myrtifolium.

SIPHONOSTOMA, Benth. & Hook. f. Gen. ii. 570 (1876), sphalm. = praec.

SIPHONYCHIA, Torr. & Gray, Fl. N. Am. i. 173 (1838). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 16.
americana, *Torr. & Gray, l. c.*— Am. bor.
diffusa, *Chapm. Fl. S. U. St.* 47.—Am. bor.
erecta, *Chapm. l. c.*—Am. bor.
Rugelii, *Chapm. l. c.*—Am. bor.

SIPHOSTIGMA, Rafin. Fl. Tellur. ii. 16 (1836) = **Cyanotis**, D. Don (Commel.).
cristata, Rafin. l. c. = C. cristata.

SIPHOTOMA, Rafin. Fl. Tellur. iv. 22 (1836) = **Hymenocallis**, Salisb. (Amaryll.).
calathina, Rafin. l. c. = H. calathina?

SIPHOTOXIS, Boj. ex Benth. in DC. Prod. xii. 457 (1848) = **Achyrospermum**, Blume (Labiat.).
rosea, Boj. ex Benth. l. c. 458 = A. fruticosum.

SIPHOTRIA, Rafin. Fl. Tellur. iv. 50 (1836) = **Renealmia**, Linn. f. (Scitam.).
squamosa, Rafin. l. c. = R. exaltata.

SIPHVALIS, Rafin. Fl. Tellur. iv. 17 (1836) = **Polygonatum**, Tourn. (Liliac.).
nitida, Rafin. l. c. = P. oppositifolium.

SIPONIMA, A. DC. in DC. Prod. viii. 246 (1844), sphalm. = CIPONIMA, Aubl. = **Symplocos**, Linn. (Styrac.).

SIRAITOS, Rafin. Fl. Tellur. iv. 26 (1836) = **Chamaelirium**, Willd. (Liliac.).
aquaticus, Rafin. l. c. 27 = C. carolinianum.

SIRIUM, Linn. Mant. ii. 145 (1771) = **Santalum**, Linn.
myrtifolium, Linn. l. c. 200; Roxb. Pl. Corom. i. t. 2 = Sant. album.

SISARUM, [Tourn.] Adans. Fam. ii. 97 (1763) = **Pimpinella**, Linn. (Umbell.).
Germanorum, Schur, Enum. Pl. Transs. 248 = Sium latifolium.
macrophyllum, Schur, l. c. = Sium latifolium.

SISIMBRYUM, [Clairv.] Man. Herb. 218 (1811), sphalm. = **Sisymbrium**, Linn. (Crucif.).

SISMONDAEA, Delponte, in Mém. Acc. Torin. Ser. II. xiv. (1854) 394 = **Dioscorea**, Plum.
dioscoreoides, Delponte, l. c. = D. piperifolia?

SISON, Linn. Coroll. Gen. 5 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 892.
ALACOSPERMUM, Neck. Elem. i. 167 (1790).
acaule, Sieber, ex Steud. Nom. ed. II. ii. 592 = Peucedanum creticum.
aegopodioides, Spreng. Syst. i. 886 = Smyrnium aegopodioides.
alpinum, Sieber, Reise Kreta, i. 472; ii. 317 = Peucedanum creticum.
Ammi, Herb. Madr. ex Wall. Cat. sub n. 7208 = Ammi indicum.
Ammi, Jacq. Hort. Vindob. t. 200 = Carum copticum.
Ammi, Linn. Sp. Pl. 252 = Apium leptophyllum?
Ammi, Ucria, Hort. Reg. Panorm. 132 = Carum ammoides.
Amomum, *Linn. Sp. Pl.* 252.—Europ.; As. Min.
amomus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
Anisum, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Pimpinella Anisum.
arvense, Brot. Fl. Lusit. i. 424 = Carum segetum.
aureum, Spreng. in Schult. Syst. vi. 410 = Thaspium aureum.
bulbosum, Michx. Fl. Bor. Am. i. 169 = Erigenia bulbosa.
canadense, Linn. Sp. Pl. 252 = Cryptotaenia canadensis.
capense, Spreng. Sp. Umbellif. 111 = Pimpinella capensis.
capillaceum, Spreng. Syst. i. 887 = Apium leptophyllum, Discopleura capillacea.
carvifolium, Bert. ex DC. Prod. iv. 116 = Carum flexuosum.
carvifolium, Wall. Cat. n. 591 = Vicatia coniifolia.
cernuum, Salisb. Prod. 165 = Carum segetum.
coniifolium, Wall. ex DC. Prod. iv. 243 = Vicatia coniifolia.
crinitum, Pall. in Act. Acad. Petrop. ii. (1779) 250. t. 7 = Schultzia crinita.
cuneifolium, Steud. Nom. ed. II. ii. 592, sphalm. = *carvifolium*.
divaricatum, Spreng. Sp. Umbellif. 113 = Apium divaricatum, Carum Bunius.
diversifolium, Wall. Cat. n. 592; et ex DC. Prod. iv. 110 = Carum diversifolium.
erectum, Salisb. Prod. 165 = S. Amomum.
exaltatum, *Boiss. Diagn.* Ser. I. x. 21.—Syria.
fasciculatum, Pohl, ex DC. Prod. iv. 105 = Apium Ammi.
flexuosum, Dum. Fl. Belg. 78 = Conopodium denudatum.
flexuosum, Tenore, Prod. Fl. Nap. p. lxvii = Carum flexuosum.
Gussonianum, Balb. ex DC. Prod. iv. 83 = Petagnia saniculaefolia.
Gussonii, Spreng. Syst. iv. Cur. Post. 118 = Petagnia saniculaefolia.
Haenkei, Presl, ex DC. Prod. iv. 105 = Apium Ammi.
heterophyllum, Moench, Meth. 97 = S. Amomum.
integerrimum, Spreng. Syst. i. 887 = Pimpinella integerrima.
inundatum, Linn. Sp. Pl. 253 = Apium inundatum.
laciniatum, L'Hérit. ex DC. Prod. iv. 105 (= *Helosciadium laciniatum*).—Peruv.
lateriflorum, Bertol. Fl. Ital. iii. 283 = Apium Ammi.
majus, Eaton & Wright, N. Am. Bot. ed. VIII. 429 = Ammi majus.
marginatum, Michx. Fl. Bor. Am. i. 168 = Peucedanum ternatum.
nodiflorum, Brot. Fl. Lusit. i. 423 = Apium nodiflorum.
peregrinum, Spreng. in Schult. Syst. vi. 409 = Carum peregrinum.
Podagraria, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Aegopodium Podagraria.
proliferum, Brochi, ex DC. Prod. iv. 116 = Carum flexuosum.
pumilum, Brot. Fl. Lusit. i. 425 = Ammi pumilum.
pusillum, Michx. Fl. Bor. Am. i. 168 = Apium divaricatum.
ranunculaefolium, Presl, ex DC. Prod. iv. 111 (= *Helosciadium ranunculaefolium*).—Chili.
rotundifolium, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Pimpinella rotundifolia.
rubricaule, Eaton & Wright, N. Am. Bot. ed. VIII. 429 = Discopleura capillacea.
Ruta, Burm. f. Fl. Cap. Prod. 7 = Apium graveolens.
salsum, Linn. f. Suppl. 181 = Peucedanum redivivum.
segetum, Linn. Sp. Pl. 252 = Carum segetum.
Sieberianum, DC. Prod. iv. 111 = Peucedanum creticum

SISON:—
sylvaticum, Brot. Fl. Lusit. i. 423 = Physospermum commutatum.
tenerum, Wall. Cat. n. 593 = Pimpinella tenera.
Thomasii, Tenore, Prod. Fl. Nap. p. lxvii = Pimpinella Thomasii.
trifidum, Burm. ex DC. Prod. iv. 106 = Apium graveolens.
trifoliatum, Michx. Fl. Bor. Am. i. 168 = Thaspium aureum.
trinervium, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 184. —Nepal.
verticillato-inundatum, Thore, Ess. Chl. Land. 101 = Carum inundatum.
verticillatum, Linn. Sp. Pl. 253 = Carum verticillatum.
verticillatum, Pall. Reise, iii. 513 = Peucedanum redivivum.

SISYMBRELLA, Spach, Hist. Vég. Phan. vi. 422 (1838) = **Sisymbrium**, Tourn. (Crucif.).
aspera, Spach, l. c. 426 = Sisymbrium asperum.
sylvestris, Spach, l. c. 425 = Nasturtium sylvestre.

SISYMBRION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113, 135 = **Sisymbrium**, Tourn. (Crucif.).
alliarium, St. Lag. l. c. = S. Alliaria.

SISYMBRIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 199 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 77.
ALLIARIA, Adans. Fam. ii. 418 (1763).
ARABIDOPSIS, Schur, Enum. Pl. Transs. 55. (1866).
CHAMAEPLIUM, Wallr. Sched. Crit. i. 376 (1822).
DESCURAINIA, Webb & Berth. Phyt. Canar. i. 72 (1835–60).
DISCUREA, Schur, Enum. Pl. Transs. 54 (1866).
DRABOPSIS, C. Koch, in Linnaea, xv. (1841) 253.
ERYSIMUM, Tourn. ex Adans. Fam. ii. 418 (1763).
HUGUENINIA, Reichb. Fl. Germ. Excurs. 691 (1832).
IRIO, Linn. Syst. ed. I (1735).
KIBERA, Adans. Fam. ii. 417 (1763).
LEPTOBASIS, Dulac, Fl. Hautes-Pyr. 198 (1867).
LEPTOCARPAEA, DC. Syst. ii. 201 (1821).
MARESIA, Pomel, Nouv. Mat. Fl. Atl. 228 (1874).
NORTA, Adans. Fam. ii. 417 (1763).
PACHYPODIUM, Webb & Berth. Phyt. Canar. i. 75 (1835–60).
PILOSELLA, Kostel. Hort. Prag. 104 (1844).
ROBESCHIA, Hochst. ex Fourn. Thèse Crucif. 146 (1865).
SISYMBRELLA, Spach, Hist. Vég. Phan. vi. 422 (1838).
SOPHIA, Linn. Syst. ed. I (1735); Adans. Fam. ii. 417 (1763).
TONGUEA, Endl. Gen. 1419 (1841).
TRICHOLOBOS, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 290.
VALARUM, Schur, Enum. Pl. Transs. 54 (1866).
VANDALEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 331.
absinthoides, S. F. Gray, Nat. Arr. Brit. Pl. ii. 680 = S. Sophia.
abyssinicum, *Fourn. Thèse Crucif.* 71.—Abyss.
acaule, *Sieber, ex Steud. Nom.* ed. II. ii. 593, nomen. —Ins. Corsic.
acre, Lam. Fl. Fr. ii. 520 = Diplotaxis tenuifolia.
aculeolatum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 75 = Malcomia aculeolata.
acutangulum, DC. Fl. Fr. iv. 670 = austriacum.
acuticarpum, *M. E. Jones, in Am. Natural.* xvii. ii. (1883) 875.—Mexic.
adpressum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 367.—Turkestan.
adpressum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 303.—Ecuador.
aegyptium, Juss. ex DC. Syst. ii. 630 = Diplotaxis pendula.
affine, Willd. Enum. Hort. Berol. Suppl. 44 = austriacum, orientale.
album, Pall. Reise, iii. 739 = Smelowskia cinerea.
alliaceum, Fourn. Thèse Crucif. 114, in syn. = seq.
Alliaria, *Scop. Fl. Carn.* ed. II. ii. 26.—Europ.; Oriens; Reg. Himal.
alpestre, F. Muell. Fragm. vii. 20 = Blennodia alpestris.
alpinum, Fourn. Thèse Crucif. 131 = Braya alpina.
alpinum, Neck. in Act. Acad. Theod. Palat. ii. (1770) 481 = strictissimum.
altissimum, Asso, Syn. Stirp. Arag. 86 = Assoanum.
altissimum, *Linn. Sp. Pl.* 659.—Europ.

SISYMBRIUM:—
amarum, Salisb. Prod. 270 = Nasturtium officinale.
amphibium, Huds. Fl. Angl. ed. II. 296, partim = Nasturtium palustre.
amphibium, Linn. Sp. Pl. 657 = Nasturtium amphibium.
amplexicaule, Desf. Fl. Atlant. ii. 81 (= *Sinapis amplexicaulis*).—Alger.
amplexicaule, *A. Gray, Bot. U. St. Expl. Exped.* i. 61.—Peruv.
amplexicaule, Jan, ex Fourn. Thèse Crucif. 142 = Moricandia arvensis.
amplexicaule. *Phil. Fl. Atac.* 8.—Chili.
anceps, Wahlenb. Fl. Upsal. 223 = Nasturtium anceps.
andinum, *Phil. in Linnaea*, xxviii. (1856) 665.—Chili.
angustifolium, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 9.—Turkestan.
anomalum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 289.—Bolivia.
antarcticum, *Fourn. Thèse Crucif.* 65.—Reg. Magellan.
apenninum, Bubani, ex Fourn. Thèse Crucif. 99 = Zanonii.
apetalum, Lour. Fl. Cochinch. 400 = Nasturtium indicum.
aquaticum, Mill. Gard. Dict. ed. VIII. n. 4 = Nasturtium amphibium.
arabidoides, Hook. Fl. Bor. Am. i. 63 = Arabis lyrata.
arcticum, *Fourn. Thèse Crucif.* 58.—Sibir. arct.
arenosum, Linn. Fl. Suec. ed. II. 233 = Arabis petraea.
arenosum, Linn. Sp. Pl. 658 = Arabis arenosa.
argutum, E. Mey. in Drège, Zwei Pfl. Docum. 222 = capense.
Arnottianum, *Gill. ex Hook. Bot. Misc.* iii. (1833) 138.—Chili.
arundanum, *Boiss. Voy. Bot. Espagne*, ii. 30.—Hispan.
arundanum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 802 = granatense.
asperum, *Linn. Sp. Pl.* 659.—Europ. austr.
asperum, Pall. Reise, iii. 740 = Dontostemon pectinatus.
asperum, Schimp. ex Boiss. Fl. Orient. i. 226 = Malcomia aculeolata.
Assoanum, *Losc. & Pard. Ser. Inconf.* ed. I. 6.—Hispan.
athrocarpum, *A. Gray, Bot. U. St. Expl. Exped.* i. 59.—Peruv.
atomarium, Adans. ex DC. Syst. ii. 484 = Dontostemon integrifolius.
atriplicifolium, *Fisch. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 7, nomen.—Reg. baical.
atrovirens, Hornem. Suppl. Hort. Hafn. 72 = Nasturtium indicum.
auriculatum, *A. Gray, Pl. Wright.* i. 8.—Texas.
austriacum, *Jacq. Fl. Austr.* iii. 35.—Europ. med.
axillare, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 162.—Reg. Himal.
Barbarea, Crantz, Stirp. Austr. ed. I. fasc. i. 54; ed. II. 52 = Barbarea vulgaris.
Barbarea, Seguier, ex Pourr. in Mém. Acad. Toul. iii. (1788) 329 = praec.?
Barbareae, All. Auct. Pedem. 19 = Barbarea praecox.
Barbareae, Linn. Sp. Pl. ed. II. 921 = Barbarea plantaginea.
barbareaefolium, Delile, Fl. Aeg. Illustr. 67 = Nasturtium palustre.
Barrelieri, Linn. Sp. Pl. ed. II. 919 = Diplotaxis Barrelieri.
Barrelieri, Thuill. Fl. Par. ed. II. 334 = Diplotaxis muralis.
Belangerii, *Fourn. Thèse Crucif.* 141, 149.—Persia.
bellidifolium, Poir. Encyc. vii. 220 = Arabis Commersonii.
Berlandieri, *Fourn. Thèse Crucif.* 105.—Mexic.
Berteroanum, *Phil. in Linnaea*, xxviii. (1856) 666.—Chili.
binerve, C. A. Mey. Verz. Pfl. Cauc. 189 = Malcomia nana.
Blennodia, F. Muell. Fragm. vii. 20 = Blennodia canescens.
Boissieri, *Nym. Consp.* 44.—Lusit.; Hisp
bonariense, Poir. Encyc. vii. 205 = Nasturtium bonariense.
Boryi, Nym. Consp. 44 = pinnatifidum.
Bourgaeanum, *Fourn. Thèse Crucif.* 55.—Ins. Teneriff.

SISYMBRIUM:—
brachycarpon, Richards. in Frankl. Narr. 1st Journ. App. 744 = canescens.
brachycarpum, Hook. & Arn. Bot. Beech. Voy. 323 = incisum?
brachylobum, Otto, ex Steud. Nom. ed. II. ii. 593, nomen; Herb. Berol. ex Fourn. Thèse Crucif. 143 = Boissieri.
brachypetalum, C. A. Mey. in Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 88 = altissimum.
brachypodum, F. Muell. Fragm. vii. 20 = Blennodia brevipes.
brassicaeforme, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 129.—Sibir.
brevicaule, Wibel, Prim. Fl. Werth. 348 = Diplotaxis viminea.
brevipes, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 154 = nudum.
Bungei, Steud. Nom. ed. II. ii. 593 = Dontostemon dentatus.
Burchellii, *DC. Syst.* ii. 472.—Afr. austr.
Burchellii, Eckl. & Zeyh. Enum. 5 = capense.
burgundiacum, Hort. Taur. ex DC. Syst. ii. 601 = Brassica Cheiranthos.
bursifolium, Gouan, Illustr. 42 = pinnatifidum.
bursifolium, *Linn. Cent. Pl.* ii. 24; *Amoen. Acad.* iv. 322.—Europ. austr.
bursifolium, Patrin, ex DC. Syst. ii. 194 = Nasturtium bursifolium.
bursifolium, Pourr. ex Nym. Consp. 44 = Boissieri.
cabulicum, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 161 = pumilum.
caeruleum, Gilib. Fl. Lituan. ii. 73 = Arabis arenosa.
californicum, S. Wats. in Bot. King's Exped. 23 = incisum.
calycinum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 289.—Bolivia.
canescens, Benth. Pl. Hartw. 9 = incisum.
canescens, Griseb. ex Fourn. Thèse Crucif. 65 = antarcticum.
canescens, *Nutt. Gen. Am.* ii. 68.—Am. bor. et austr.
capense, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
capillaceum, Gilib. Fl. Lituan. ii. 72 = Nasturtium palustre.
cardaminefolium, Gilib. l. c. 70 = Nasturtium officinale.
cardaminoides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 34.—Austral.
carnosulum, Phil. Fl. Atac. 7 = stenophyllum.
cartilagineum, Pall. ex DC. Syst. ii. 471 = junceum.
catholicum, Boeber, in Pall. N. Nord. Beitr. vi. 258 = Diplotaxis tenuifolia.
catholicum, Linn. Mant. i. 93 = Diplotaxis catholica.
ceratophyllum, Desf. Fl. Atlant. ii. 82 = Nasturtium ceratophyllum.
cheiranthoides, Eaton & Wright, N. Am. Bot. ed. VIII. 429; Trautv. in Act. Hort. Petrop. v. (1877) 27 = Erysimum cheiranthoides.
Cheiranthus, Trautv. l. c. iv. (1876) 108 = Erysimum lanceolatum, altaicum.
chrysanthum, Jord. Diagn. i. 148 = austriacum.
ciliolatum, *Fourn. Thèse Crucif.* 129.—Peruv.
cinereum, Desf. Fl. Atlant. ii. 83. t. 157 = Ammosperma cinerea.
cinereum, Nutt. ex Fourn. Thèse Crucif. 63 = Cumingianum.
clandestinum, Hort. Berol. ex Spreng. Nov. Prov. 29 = Nasturtium clandestinum.
coeruleum, Auct. = *caeruleum*.
colchaguense, *Wedd. ex Fourn. Thèse Crucif.* 134.—Chili.
Columnae, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 157 = heteromallum.
Columnae, Jacq. Fl. Austr. iv. 12 = orientale.
compressum, Moench, Meth. Suppl. 83 = austriacum.
confertum, Stev. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 304 = polyceratium.
confusum, *Fourn. Thèse Crucif.* 95.—Afr. austr.
confusum, *Janka, in Termèsz. Füzetek*, viii. (1884) 31.—Hungar.
contortum, *Cav. Desc.* 436.—Hispan.
contortuplicatum, DC. Syst. ii. 483 = Malcomia contortuplicata.
contortuplicatum, C. A. Mey. Verz. Pfl. Cauc. 190 = Malcomia Ledebourii.
corniculatum, *Cav. Desc.* 437.—Hispan.
corniculatum, Lam. Fl. Fr. ii. 520 = polyceratium.
coronopifolium, Desf. Fl. Atlant. ii. 82 = Nasturtium coronopifolium.

SISYMBRIUM:—
corymbosum, *Phil. in Linnaea*, xxviii. (1856) 666.—Chili.
Coulteri, *Hemsl. Diagn. Pl. Nov.* 18.—Mexic.
crassifolium, *Cav. Desc.* 437.—Hispan.
crassipes, Trautv. in Act. Hort. Petrop. ii. (1873) 500 = Erysimum crassipes.
crithmifolium, Roth, Nov. Pl. Sp. 325 (err. typ. 225) = Heliophila crithmifolia.
Cumingianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 38.—Chili.
Cumingii, Presl, Bot. Bemerk. 9 = praec.
curvipes, F. Muell. Fragm. vii. 20 = Blennodia curvipes.
curvisiliquum, Bertol. Misc. Bot. ii. 13. t. 10 = Malcomia torulosa.
curvisiliquum, Hook. Fl. Bor. Am. i. 61 = Nasturtium curvisiliqua.
dahuricum, *Turcz. ex Fourn. Thèse Crucif.* 97.—Dahuria.
damascenum, *Boiss. & Gaill. Diagn.* Ser. II. vi. 11.—Syria.
dareaefolium, Poepp. ex Fourn. Thèse Crucif. 63, in syn. = Cumingianum.
decipiens, *Bunge, in Linnaea*, xviii. (1844) 505.—Turkestan.
deflexum, Harv. ex Torr. in Pacif. Rail. Rep. iv. 66, sphalm. = reflexum.
Delilei, Coss. in Bull. Soc. Bot. Fr. xxvii. (1880) 71 = Malcomia pygmaea.
deltoideum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 163.—Reg. Himal.
dentatum, All. Fl. Pedem. i. 275. t. 57. f. 3 = pinnatifidum.
dentatum, Torr. in Short, Pl. Kentucky, Suppl. iii. 338; ex S. Wats. Bibliog. Ind. N. Am. Bot. 47 = Arabis dentata.
derelictum, Jord. Diagn. i. 150 = austriacum.
diffusum, *A. Gray, Pl. Wright.* i. 8.—Texas.
dissectum, Nutt. ex Fourn. Thèse Crucif. 143 = Nasturtium palustre.
diversifolium, Stokes, Bot. Mat. Med. iii. 450 = Nasturtium amphibium.
domingense, Poir. Encyc. Suppl. v. 161 = Nasturtium pyrenaicum.
Doumetianum, *Coss. in Bull. Soc. Bot. Fr.* xxvii. (1880) 70.—Afr. bor.
dubium, Pers. Syn. ii. 199 = Nasturtium indicum.
eckartsbergense, Willd. Sp. Pl. iii. 502 = austriacum.
Edwardsii, Trautv. in Act. Hort. Petrop. i. (1871) 59 = Eutrema Edwardsii.
eglandulosum, DC. Syst. ii. 485 = Dontostemon dentatus.
elatum, *C. Koch, in Ind. Sem. Hort. Berol.* (1854) *App.* 14.—Armenia.
eremigenum, F. Muell. Fragm. ii. 143 = Blennodia eremigera.
erosum, E. Mey. in Drège, Zwei Pfl. Docum. 222 = Brassica strigosa.
erucaefolium, Gilib. Fl. Lituan. ii. 74 = S. Irio.
erucastrifolium, Trautv. in Act. Hort. Petrop. iv. (1876) 107 = Erysimum erucastrifolium.
Erucastrum, Gouan, Illustr. 42 = Diplotaxis muralis.
Erucastrum, Pollich, Hist. Pl. Palat. ii. 234 = Brassica Erucastrum.
erucoides, Desf. Fl. Atlant. ii. 83 = Diplotaxis erucoides.
Erysimastrum, Lam. Fl. Fr. ii. 521 = S. Irio, orientale.
erysimifolium, Pourr. in Mém. Acad. Toul. iii. (1788) 329 = austriacum.
erysimoides, *Desf. Fl. Atlant.* ii. 84. t. 158.—Reg. Mediterr.; Afr. trop.; Arab.
exacoides, DC. Syst. ii. 463 = Malcomia exacoides.
exasperatum, *Sond. in Linnaea*, xxiii. (1850) 3.—Afr. austr.
falcatum, *Fourn. Thèse Crucif.* 135.—Abyssin.
fastigiatum, *Phil. in Anal. Univ. Chil.* (1870) II. 670.—Chili.
filicifolium, Georgi, Beschr. Russ. Reich. Nachtr. 287 = Leptaleum filifolium.
filifolium, F. Muell. in Trans. Vict. Inst. i. (1855) 155, in obs. = Blennodia filifolia.
filifolium, Willd. Sp. Pl. iii. 495 = Leptaleum filifolium.
filiforme, Dulac, Fl. Hautes-Pyr. 197 = S. Irio.
filiformium, Poir. Encyc. vii. 208 = Leptaleum filifolium.
flavissimum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 382 = Smelowskia flavissima.

SISYMBRIUM:—
flexuosum, Dulac, Fl. Hautes-Pyr. 197 = orientale.
fluviatile, Vell. Fl. Flum. 271; vi. t. 109 = Nasturtium officinale.
foliosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 160.—Reg. Himal.
foliosum, *Phil. in Linnaea*, xxxiii. (1864–65) 10.—Chili.
fragile, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 288.—Peruv.
frutescens, *Gill. ex Hook. Bot. Misc.* iii. (1833) 139.—Chili.
fugax, *Lag. Gen. et Sp. Nov.* 20.—Hispan.
Gaillardoti, Boiss. Fl. Orient. i. 214, in syn. = pumilum.
Galeottianum, *Fourn. Thèse Crucif.* 59.—Mexic.
gallicum, Schleich. ex Soyer-Willem. in Ann. Sc. Nat. Sér. II. ii. (1834) 116 = Brassica Cheiranthos.
gallicum, Willd. Enum. Hort. Berol. 678 = S. Irio.
gangeticum, Buch.-Ham. ex Wall. Cat. n. 4777 = Nasturtium indicum.
gariepinum, *Burch. ex DC. Syst.* ii. 472.—Afr. austr.
Gayanum, Barn. in C. Gay, Fl. Chil. i. 122 = leptocarpum.
gelidum, Trautv. in Act. Hort. Petrop. ii. (1873) 501 = Erysimum gelidum.
geraniifolium, Poir. Encyc. vii. 218 = Cardamine geraniifolia.
glabrum, Willd. Enum. Hort. Berol. Suppl. 44 = S. Irio.
glaciale, Forst. f. in Comm. Gotting. ix. (1789) 32 = Cardamine glacialis.
glaucescens, Jord. Diagn. i. 144 = austriacum.
glaucum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 93.—Am. bor.
glaucum, Sieber, ex Steud. Nom. ed. I. 784, nomen (Quid?).—Hab.?
gracile, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 288.—Bolivia.
granatense, *Boiss. Diagn.* Ser. II. i. 24.—Hispan.
grandiflorum, Molina, Sagg. Chil. ed. II. 292 = Cardamine glacialis.
grandiflorum, Wedd. in Ann. Sc. Nat. Sér. V. i. (1864) 290 (= *Eudema grandiflora*).—Am. austr.
Griffithianum, *Boiss. Diagn.* Ser. II. i. 23.—Afghan.
halophilum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 143.—Sibir. altaic.
Harra, Steud. Nom. ed. I. 784 = Diplotaxis pendula.
Hartwegianum, Fourn. Thèse Crucif. 66 = incisum.
hastatum, *Phil. Sert. Mendoc. Alt.* 2.—Chili.
hesperidiflorum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 22 = Hesperis leucoclada.
heteromallum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 132.—Sibir. altaic.
heterophyllum, Forst. f. Prod. 46 = Cardamine debilis.
hieracifolium, Trautv. in Act. Hort. Petrop. v. (1877) 27 = Erysimum hieracifolium.
himalaicum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 160.—Reg. Himal.
hirsutum, Gilib. Fl. Lituan. ii. 74 = Loeselii.
hirsutum, Host, ex Fourn. Thèse Crucif. 76 = supinum.
hirsutum, Lag. ex DC. Syst. ii. 478 = runcinatum.
hirtulum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 240.—Turkestan.
hirtum, Host, Fl. Austr. ii. 261 = Brassica Erucastrum.
hispanicum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 799 = contortum.
hispanicum, *Jacq. Coll.* i. 69; *Ic. Pl. Rar.* i. 12. t. 124.—Hispan.
hispidulum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 63.—N. Granat.
hispidum, Moench, Meth. Suppl. 83 = Loeselii.
hispidum, *Phil. in Anal. Univ. Chil.* (1870) II. 670.—Chili.
hispidum, Poir. Encyc. Suppl. v. 161 = Nasturtium palustre.
hispidum, Vahl, Symb. Bot. ii. 77 = Diplotaxis pendula.
hispidum, Willd. ex DC. Syst. ii. 480 = Malcomia torulosa?
Hookeri, *Fourn. Thèse Crucif.* 120.—Reg. Himal.
Hueti, *Boiss. Fl. Orient.* i. 957.—Armenia.
humifusum, J. Vahl, in Fl. Dan. t. 2297 = Arabis lyrata.
humile, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 137.—Reg. Himal.; As. et Am. bor.
hungaricum, Lumn. Fl. Poson. 282 = altissimum.
hybridum, Thuill. Fl. Par. ed. II. 331 = Nasturtium palustre.

SISYMBRIUM:—
ibericum, Trautv. in Act. Hort. Petrop. iv. (1876) 108 = Erysimum ibericum.
imbricatifolium, *Wedd. ex Fourn. Thèse Crucif.* 132.—Chili.
incanum, Bernh. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 38 = canescens.
incisum, *Engelm. ex A. Gray, in Mem. Acad.* N. S. iv. (1849) 8.—Am. bor. occ.
indicum, Linn. Mant. i. 93 = Nasturtium indicum.
indicum, Walt. Fl. Carol. 174 = Nasturtium amphibium.
integrifolium, Boeber, in Pall. N. Nord. Beitr. vi. 260 = junceum.
integrifolium, Gilib. Fl. Lituan. ii. 72 = Cochlearia amphibia.
integrifolium, Linn. Sp. Pl. 660 = Dontostemon integrifolius.
intermedium, *Hose, in Usteri, Ann. Bot.* xxi. (1797) 37.—Europ.
Irio, Crantz, ex Steud. Nom. ed. I. 784 = orientale.
Irio, C. Koch, in Linnaea, xv. (1841) 253 = heteromallum.
Irio, *Linn. Sp. Pl.* 659.—Europ.; As. et Afr. bor.
Irio, Wibel, ex Steud. Nom. ed. II. ii. 694 = Diplotaxis viminea.
irioides, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 76 = S. Irio.
Iris, Retz. Fl. Scand. 126, sphalm. = S. Irio.
islandicum, Oed. Fl. Dan. t. 409 = Nasturtium palustre.
jacobeaefolium, Berger. ex Vill. Cat. Strasb. 259 = Brassica Erucastrum.
Jacquemontii, *Fourn. Thèse Crucif.* 111.—Ind. or.
junceum, *Bieb. Fl. Taur. Cauc.* ii. 114.—Europ. or.; As. et Am. bor.
junceum, Hook. Fl. Bor. Am. i. 61 = linifolium.
Kahikir, *Mart. ex Spreng. Syst.* ii. 901.—Hab.?
Kochii, *Petri, in Ind. Sem. Hort. Berol.* (1868) 24.—Hab.?
kokanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 9.—Turkestan.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 240.—Turkestan.
Kralikii, Fourn. Thèse Crucif. 74 = S. Irio.
kunawurense?—Cf. *runawurense*.
laevigatum, Willd. Sp. Pl. iii. 500 = Diplotaxis tenuifolia.
Lagascae, *Amo, Fl. Iber.* vi. 529.—Hispan.
lasiocarpum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 162.—Reg. Himal.
lasiocarpum, F. Muell. Fragm. vii. 20 = Blennodia lasiocarpa.
lasiolobum, *Boiss. Diagn.* Ser. II. i. 24.—America?
latifolium, S. F. Gray, Nat. Arr. Brit. Pl. ii. 679 = S. Irio.
laxiflorum, *Boiss. Elench.* 9.—Hispan.
Lechleri, *Fourn. Thèse Crucif.* 129.—Chili.
leiocarpum, Jord. Diagn. i. 139 = officinale.
leptocarpum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 139.—Chili.
leptopetalon, Rafin. Fl. Ludov. 84 (Quid?).—Am. bor.
leptophyllum, Trautv. in Act. Hort. Petrop. ii. (1873) 500 = Erysimum leptophyllum.
Limosella, *Fourn. Thèse Crucif.* 132.—Mont. Altai.
lineare, DC. Syst. ii. 464 = junceum.
linifolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 91.—Am. bor.
lippizense, Wulf. in Jacq. Coll. ii. 161 = Nasturtium lippizense.
litorale, *Phil. in Linnaea*, xxviii. (1856) 667.—Chili.
Loeselii, *Linn. Cent. Pl.* i. 18; *Amoen. Acad.* iv. 279.—Europ.; Oriens; Reg. Himal.
Loeselii, Nym. Syll. 194, partim = Assoanum.
Loeselii, Thuill. Fl. Par. ed. II. 335 = orientale.
longepedicellatum, Fourn. Thèse Crucif. 59 (excl. syn.) = incisum.
Lucae, *F. Muell. Census*, 5.—Austral.
ludovicianum, Nutt. ex Hook. Journ. Bot. i. (1834) 191 = Cardamine ludoviciana.
lyratum, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
lyratum, Meissn. ex Sond. in Harv. & Sond. Fl. Cap. i. 32 (= *Sinapis retrorsa*).—Afr. austr.
macroloma, *Pomel, Nouv. Mat. Fl. Atl.* 368.—Alger.
macrophyllum, Barn. in C. Gay, Fl. Chil. i. 129 = Cumingianum.
macrostachyum, *Phil. in Linnaea*, xxxiii. (1864–65) 9 (sp. dub.).—Chili.

SISYMBRIUM:—
magellanicum, *Hook. f. Fl. Antarct.* ii. 243.—Reg. Magell.
malcolmioides, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 431.—Alger.
Mandonii, *Fourn. Thèse Crucif.* 109.—Boliv.
matthioloides, Wedd. ex Fourn. l. c. 144 = Mathewsia matthioloides.
maximum, Hochst. ex Fourn. l. c. 72 = S. Irio.
mexicanum, Sesse, ex Spreng. Syst. ii. 882 = Nasturtium palustre.
Meyeri, Ledeb. Fl. Ross. i. 184 = Malcomia Ledebourii.
Meyeri, Steud. Nom. ed. II. ii. 594 = Dontostemon micranthus.
micranthum, Roth, Nov. Pl. Sp. 324 = Nasturtium micranthum.
micranthum, Willd. Enum. Hort. Berol. Suppl. 44, nomen (Quid?).—Hab.?
millefolium, *Ait. Hort. Kew.* ed. I. ii. 391.—Ins. Teneriff.
minus, Lam. Fl. Fr. ii. 517 = Diplotaxis muralis.
minutiflorum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 158.—Reg. Himal.
molle, Jacq. Coll. i. 68 = pumilum.
mollissimum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 140.—Reg. Himal.; Sibir.
monanthos, Viv. Fl. Lib. Spec. 68 = Morisia hypogaea.
monense, Gilib. Elem. ii. 184; ex DC. Syst. ii. 601 = Brassica Cheiranthos.
monense, Linn. Sp. Pl. ed. I. 658 = Brassica monensis.
monense, Linn. l. c. ed. II. 918 = Diplotaxis saxatilis.
monense, Thuill. Fl. Par. ed. II. 333 = Diplotaxis muralis.
monense, Vill. Hist. Pl. Dauph. iii. 350. t. 39 = Brassica repanda.
montanum, E. Mey. ex Presl, Bot. Bemerk. 9 = capense.
montivagum, Jord. Diagn. i. 146 = austriacum.
multifidum, Gilib. Fl. Lituan. ii. 75 = S. Sophia.
multifidum, Willd. ex Spreng. Syst. ii. 904 = junceum.
multisiliquosum, Hoffm. Deutschl. Fl. ed. II. i. II. 50 = austriacum.
murale, Bieb. Fl. Taur. Cauc. ii. 112 = Diplotaxis saxatilis.
murale, Bory & Chaub. Nouv. Fl. Pélop. 43 = Diplotaxis viminea.
murale, Linn. Sp. Pl. 658 = Diplotaxis muralis.
muricatum, Willd. ex DC. Syst. ii. 485 = Dontostemon pectinatus.
myriophyllum, *H. B. & K. ex DC. l. c.* 477.—Ecuador.
nanum, DC. Syst. ii. 486 = Malcomia nana.
nanum, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 281 = Braya versicolor.
nasturtiifolium, Gilib. Fl. Lituan. ii. 71 = Nasturtium sylvestre.
nasturtioides, F. Muell. in Trans. Vict. Inst. i. (1855) 115, in obs. = Blennodia nasturtioides.
Nasturtium, Muhl. Cat. 60 = Thelypodium pinnatifidum.
Nasturtium, Thunb. Fl. Jap. 260 = seq.
Nasturtium-aquaticum, Linn. Sp. Pl. 657 = Nasturtium officinale.
nebrodense, Poir. Encyc. Suppl. v. 161 = Lepidium nebrodense.
niagarense, Fourn. Thèse Crucif. 85 = officinale.
nitidulum, Lag. ex Spreng. Syst. ii. 901 = erysimoides.
nitidum, Zea, ex Desf. Tabl. ed. II. 153 = erysimoides.
niveum, *Phil. Fl. Atac.* 8.—Chili.
Novae-Zelandiae, *Hook. f. Handb. N. Zeal. Fl.* 11.—N. Zel.
nubigenum, Wedd. in Ann. Sc. Nat. Sér. V. i. (1864) 290 (= *Eudema nubigena*).—Ecuador.
nudum, *Boiss. Fl. Orient.* i. 214.—Persia.
Nuttallii, Colla, Herb. Pedem. i. 206. n. 29 = Cumingianum.
obtusangulum, Auch. ex Fourn. Thèse Crucif. 91 = altissimum.
obtusangulum, Schleich. Cat. 48; ex Willd. Sp. Pl. iii. 504 = Brassica Erucastrum.
officinale, *Scop. Fl. Carn.* ed. II. ii. 26.—Europ.; Afr. bor.
officinarum, Crantz, Stirp. Austr. ed. I. fasc. i. 54; ed. II. 52 = praec.
oliganthum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 289.—Bolivia.

SISYMBRIUM:—
Orbignyanum, *Fourn. Thèse Crucif.* 107.—Bolivia.
orientale, *Linn. Cent. Pl.* ii. 24; *Amoen. Acad.* iv. 322.—Europ.; Oriens; Reg. Himal.
orophilum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 288.—Bolivia.
Pallasii, Spreng. Syst. ii. 901 = junceum.
pallescens, Jord. Diagn. i. 145 = austriacum.
Palmeri, *Hemsl. Diagn. Pl. Nov.* 19.—Mexic.
palustre, Garsault, Fig. Pl. t. 550; ex DC. Syst. ii. 197 = Nasturtium amphibium.
palustre, Leyss. Fl. Hal. 166 = Nasturtium palustre.
paniculatum, Steph. ex Fourn. Thèse Crucif. 145 = Arabis incarnata.
pannonicum, Badaro, in Moretti, in Brugnat. Giorn. ix. (1826) 74 = orientale.
pannonicum, Jacq. Coll. i. 70 = altissimum.
Parra, Linn. Mant. ii. 255 = Brassica sabularia.
Parryi, *Hemsl. Diagn. Pl. Nov.* 19.—Mexic.
parviflorum, Lam. Fl. Fr. ii. 519 = S. Sophia.
parvifolium, *Phil. in Linnaea*, xxviii. (1856) 667.—Chili.
patens, *Moench, Meth.* 251 (sp. dub.).—Hab.?
patulum, *Fourn. Thèse Crucif.* 104.—Mexic.
pauciflorum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 93.—Am. bor.
pectinatum, Fisch. ex DC. Syst. ii. 485 = Dontostemon pectinatus.
pendulum, Desf. Fl. Atlant. ii. 82. t. 156 = Diplotaxis pendula.
pendulum, Lag. ex DC. Syst. ii. 629 = Diplotaxis Lagascana.
perenne, Steud. Nom. ed. II. ii. 594 = Dontostemon perennis.
perfoliatum, C. A. Mey. Verz. Pfl. Cauc. 188 = Conringia clavata.
persicum, Schrad. ex Spreng. Nov. Prov. 39 = S. Sophia.
peruvianum, *DC. Syst.* ii. 477.—Peruv.
petraeum, Delarb. Fl. Auv. ed. II. 359 = Arabis petraea.
petraeum, *Phil. in Linnaea*, xxviii. (1856) 668.—Chili.
Peyrousianum, Nym. Consp. 44 = pinnatifidum.
Piasezkii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 421.—China.
pilosissimum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 369.—Turkestan.
pimpinellifolium, Barn. in C. Gay, Fl. Chil. i. 129 = Cumingianum.
pinnatifidum, *DC. Fl. Fr.* iv. 667.—Europ.
pinnatifidum, Forsk. Fl. Aegypt. Arab. 118 = S. Irio.
pinnatum, *Barn. in C. Gay, Fl. Chil.* i. 125.—Chili.
planisiliquum, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 159 = Conringia planisiliqua.
polyceratium, *Linn. Sp. Pl.* 658.—Europ. austr.; As. Min
polyspermum, *Fourn. Thèse Crucif.* 103.—Mexic.
polystachyum, *Colla, Herb. Pedem.* i. 206 (sp. dub.).—Italia.
polystachyum, Hort. ex Tausch, in Flora, xix. (1836) 410 = Halimolobos patula.
praecox, *Colla, Herb. Pedem.* i. 205 (sp. dub.).—Italia.
Preauxianum, *Webb, in Ann. Sc. Nat.* Sér. II. xiii. (1840) 137.—Ins. Canar.
primulaefolium, Thoms. in Hook. Kew Journ. v. (1853) 18 = Eutrema primulaefolium.
procumbens, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1885) 67.—Austral.
propinquum, Jord. Diagn. i. 148 = austriacum.
Pseudo-Sophia, Boiss. & Huet, Diagn. Ser. II. v. 24 = S. Sophia.
pulchellum, Trautv. in Act. Hort. Petrop. ii. (1873) 501 = Erysimum rupestre.
Pumilio, *Oliver, Fl. Trop. Afr.* i. 64.—Afr. trop.
pumilum, Lam. Fl. Fr. ii. 516 = Diplotaxis viminea.
pumilum, Spach, ex Fourn. Thèse Crucif. 96 = scholare.
pumilum, *Steph. ex Willd. Sp. Pl.* iii. 507.—Oriens; As. med.
pusillum, Vill. Fl. Delph. 69; Hist. Pl. Dauph. iii. 341 = Nasturtium palustre.
pusillum, *Wedd. ex Fourn. Thèse Crucif.* 131.—Chili.
pygmaeum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 91 = linifolium?
pygmaeum, Trautv. in Act. Hort. Petrop. i (1871) 60 = Cheiranthus pygmaeus.

SISYMBRIUM:—
pyrenaicum, Linn. Syst. ed. X. 1132 = Nasturtium pyrenaicum.
pyrenaicum, Vill. Hist. Pl. Dauph. iii. 341 = austriacum.
ramosum, *Fourn. Thèse Crucif.* 140.—Persia.
ramulosum, *Delile, Fl. Aeg. Illustr.* 67.—Aegypt.
ramulosum, Poir. Encyc. Suppl v. 161 = S. Irio.
Reboudianum, Verlot, Cat. Pl. Jard. Grenoble (1857); et in Bull. Soc. Bot. Fr. iv. (1857) 726 = S. Irio.
Redowskii, Weinm. ex Steud. Nom. ed. I. 785 = Erysimum Redowskii.
reflexum, Kellogg, in Proc. Calif. Acad. ii. (1863) 101. t. 29 = Arabis Holboellii.
reflexum, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 183.—Am. bor. occ.
Relangerii, Fourn. Thèse Crucif. 141, sphalm. = Belangerii.
Remyanum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 290.—Ecuador.
repandum, Poir. Encyc. vii. 209 = Diplotaxis saxatilis.
repandum, Willd. Sp. Pl. iii. 497 = Brassica repanda.
reticulatum, Tratt. Arch. t. 350; ex Pritz. Ic. Ind. i. 1039 (sp. dub.).—Hab.?
rhedonense, Degland, ex Gren. & Godr. Fl. Fr. i. 95 = austriacum.
Richardsii, *F. Muell. Census*, 5.—Austral.
Richardsoni, Sweet, Hort. Brit. ed. II. 30 = canescens.
rigidulum, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 272 = altissimum.
rigidulum, Lag. Gen. et Sp. Nov. 20 = erysimoides.
rigidum, Bieb. Fl. Taur. Cauc. iii. 439 = Malcomia torulosa.
Robesetti, *Steud. Nom.* ed. II. ii. 594.—Arab.
Roripa, Scop. Fl. Carn. ed. II. ii. 25 = Nasturtium amphibium.
ruderale, Jord. Diagn. i. 138 = officinale.
runawurense, Royle, ex Fourn. Thèse Crucif. 122 (recte *kunawurense*?) = rupestre.
runcinatum, *Lag. ex DC. Syst.* ii. 478.—Reg. Mediterr. et Caucas.; Oriens.
rupestre, *Edgew. in Trans. Linn. Soc.* xx. (1846) 33.—Reg. Himal.
rupestre, Wedd. in Ann. Sc. Nat. Sér. V. i. (1864) 290 = Eutrema Humboldtii.
rupestricolum, Jord. Diagn. i. 142 = austriacum.
sagittatum, Ait. Hort. Kew. ed. I. ii. 390 = Alyssopsis deflexa.
sagittatum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 139.—Chili.
salsugineum, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 159 = Conringia persica.
salsugineum, *Pall. Reise*, ii. 740.—Sibir.
salsugineum, Schlecht. ex DC. Syst. ii. 220 = toxophyllum.
salsuginosum, Willd. Sp. Pl. iii. 507 = salsugineum.
salsum, Georgi, Reise, i. 224; ex Ledeb. Fl. Ross. i. 185 = salsugineum.
saxatile, Lam. Fl. Fr. ii. 517 = Diplotaxis saxatilis.
Schaffneri, *Hemsl. Diagn. Pl. Nov.* 19.—Mexic.
Schimperi, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 76; Presl, Bot. Bemerk. 9 = S. Sophia.
Schimperi, J. Gay, ex Fourn. Thèse Crucif. 124 = pumilum.
scholare, *Fourn. l. c.* 96.—Hab.?
scorpiuroides, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 74 = Malcomia torulosa.
Scorpiurus, *Pomel, Nouv. Mat. Fl. Atl.* 227.—Alger.
septaceum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 289.—Bolivia.
septulatum, DC. Syst. ii. 471 = altissimum.
serratum, Thunb. Prod. Pl. Cap. 109 = Lepidium ruderale.
setosum, Ledeb. ex Spreng. Syst. ii. 902 = Brassica Tournefortii.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 274.—Turkestan.
Sieversianum, Redowsk. ex DC. Syst. ii. 485 = Dontostemon dentatus.
siliquosum, *Fourn. Thèse Crucif.* 133.—As. bor.
simplex, Viv. Fl. Lib. Spec. 36. t. 16. f. 2 (Diplotaxidis sp.).—Afr. bor.
simplicissimum, S. T. Gmel. Reise, iii. 303. t. 31. f. 1 = Thalianum.
simplicissimum, Lapeyr. Hist. Abr. Pl. Pyr. 382 = Arabis perfoliata.
sinapifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = Nasturtium indicum.
Sinapios, Retz. Obs. iii. 37 = altissimum.

SISYMBRIUM:—
Sinapis, Burm. f. Fl. Ind. 140, excl. syn. = Nasturtium indicum.
Sinapistrum, Crantz, Stirp. Austr. ed. II. 52 = altissimum.
sinapoides, R. Br. in Ait. Hort. Kew. ed. II. iv. 112 = austriacum.
sinuatum, Nutt. ex Fourn. Thèse Crucif. 146 (Nasturtii sp.?).—Am. bor.
solidagineum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 64.—N. Granat.
Sophia, Barn. in C. Gay, Fl. Chil. i. 127 = Cumingianum.
Sophia, A. Gray, in Proc. Acad. Philad. (1863) 57, partim = incisum.
Sophia, *Linn. Sp. Pl.* 659.—Reg. temp.
Sophia, Pursh, Fl. Am. Sept. ii. 440 = canescens.
sophioides, Fisch. ex Hook. Fl. Bor. Am. i. 61. t. 20 = S. Sophia.
spathulaefolium, *A. Gray, Bot. U. St. Expl. Exped.* i. 60.—Peruv.
spathulatum, Poir. Encyc. vii. 221 = Arabis spathulata.
spectabile, *Hook. & Thoms. ex Fourn. Thèse Crucif.* 121.—Reg. Himal.
spurium, Hort. ex Steud. Nom. ed. I. 785, nomen.—(Quid?).
stenophyllum, *Gill. ex Hook. & Arn. in Bot. Misc.* iii. (1833) 139.—Chili.
stenophyllum, Schlecht. in Linnaea, xxviii. (1856) 475 = Lechleri.
stigmatosum, *Steud. Nom.* ed. II. ii. 595.—Arab.
stoloniferum, J. & C. Presl, Fl. Čech. 137 = Nasturtium amphibium.
streptocarpum, *Fourn. Thèse Crucif.* 58.—Mexic.
strictissimum, *Linn. Sp. Pl.* 660.—Europ. med.
strictissimum, Pall. ex Bieb. Fl. Taur. Cauc. ii. 115 = junceum.
strictum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 161.—Reg. Himal.
strictum, Phil. Fl. Atac. 7 = Cumingianum.
strigosum, Thunb. Herb. ex Sond. in Harv. & Sond. Fl. Cap. i. 24, 32 = capense, Brassica leptopetala.
strigosum, Thunb. Prod. Pl. Cap. 109 = Brassica strigosa.
subbipinnatifidum, Hornem. Suppl. Hort. Hafn. 72 (= *Sinapis subbipinnatifida*).—Hispan.
subhastatum, Edgew. ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 157 – S. Irio.
subhastatum, Hornem. Enum. Pl. Hort. Hafn. (1807) 37; Willd. Enum. Hort. Berol. 679 = orientale.
subspinescens, *Bunge, Arb. Nat. Ver. Riga*, i. 11. (1847) 151.—Reg. Casp.; Soongar.
subulatum, *Fourn. Thèse Crucif.* 72.—Abyss.
suecicum, *Nym. Consp.* 44.—Europ. bor.
suffruticosum, *Fourn. Thèse Crucif.* 132.—Chili.
supinum, Asso, Mant. 173 = runcinatum.
supinum, *Linn. Sp. Pl.* 657.—Europ. med.
sylvestre, Burm. ex DC. Syst. ii. 472 = lyratum.
sylvestre, Burm. f. Fl. Cap. Prod. 17 = Diplotaxis tenuifolia.
sylvestre, Linn. Sp. Pl. 657 = Nasturtium sylvestre.
sylvestre, Mohr, ex Bab. in Journ. Linn. Soc. xi. (1870) 293 = Nasturtium palustre.
tanacetifolium, *Linn. Sp. Pl.* 659.—Europ.
tanacetifolium, Walt. Fl. Carol. 174 = Nasturtium tanacetifolium.
taraxacifolium, DC. Fl. Fr. iv. 670 = austriacum.
tenuifolium, Geners. Elench. Scepus. n. 613; ex Wahlenb. Fl. Carp. 201 = altissimum.
tenuifolium, Linn. Cent. Pl. i. 18; Amoen. Acad. iv. 279 = Diplotaxis tenuifolia.
tenuifolium, Pall. ex Spreng. Syst. ii. 901 = junceum.
tenuifolium, Salisb. Prod. 270 = S. Sophia.
tenuissimum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 382 = S. Sophia.
tenuissimum, Phil. in Linnaea, xxxiii. (1864-65) 9 = Cumingianum.
teres, Torr. & Gray, Fl. N. Am. i. 93 = Nasturtium tanacetifolium.
terrestre, Wahlenb. Fl. Lappon. 179 = Nasturtium amphibium.
terrestre, With. Arr. Brit. Pl. ed. III. iii. 582 = Nasturtium palustre.
tetragonum, Trautv. in Act. Hort. Petrop. iv. (1876) 350 = Conringia orientalis.
tetromallum, C. Muell. in Walp. Ann. vii. 135, sphalm. = heteromallum.

SISYMBRIUM:—
Thalianum, *J. Gay, in Ann. Sc. Nat.* Sér. I. vii. (1826) 399.—Reg. temp.
Thomsoni, *Hook. f. in Journ. Linn. Soc.* v. (1861) 161.—Reg. Himal.
tibeticum, *Journ. Thèse Crucif.* 128.—Reg. Himal.
Tilesii, Ledeb. in Mém. Acad. Pétersb. v. (1815) 548 = Arabis petraea.
Tilingii, *Journ. Thèse Crucif.* 133.—Sibir.
Tillieri, Bell. ex Willd. Sp. Pl. iii. 497 = austriacum.
titicacense, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 249.—Peruv.
tomentosum, Stev. ex DC. Syst. ii. 210 = Arabis incarnata.
torulosum, Desf. Fl. Atlant. ii. 84. t. 159 = Malcomia torulosa.
Tournefortii, Schleich. ex Fourn. Thèse Crucif. 146 = Brassica Erucastrum.
toxophyllum, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 142.—As. bor.
trinerve, Steud. Nom. ed. II. ii. 595, sphalm. = binerve.
tripinnatum, *DC. Syst.* ii. 475.—Afr. austr.
trisectum, F. Muell. in Trans. Vict. Inst. i. (1855) 114 = Blennodia trisecta.
truncatum, Dulac, Fl. Hautes-Pyr. 197 = S. Alliaria.
tuberosum, Lag. ex DC. Syst. ii. 254 = Cardamine tuberosa.
Turczaninowii, *Sond. in Harv. & Sond. Fl. Cap.* i. 26.—Afr. austr.
turritopsis, C. Muell. in Walp. Ann. vii. 135, sphalm. = sagittatum.
uniflorum, *Fourn. Thèse Crucif.* 133.—Reg. Himal.
valentinum, Juss. ex DC. Syst. ii. 636 = Diplotaxis saxatilis.
valentinum, Linn. Sp. Pl. ed. II. 920 = Brassica valentina.
vernum, *Fourn. Thèse Crucif.* 127.—Reg. Caucas.
versicolor, Fourn. l. c. 141 = Braya versicolor.
Villarsii, Jord. Diagn. i. 143 = austriacum.
villosum, Lag. ex Spreng. Syst. ii. 902 = runcinatum.
villosum, Moench, Meth. 251 = orientale.
vimineum, Bieb. Fl. Taur. Cauc. ii. 112 = Diplotaxis saxatilis.
vimineum, Linn. Sp. Pl. ed. II. 919 = Diplotaxis viminea.
vineale, Gater. Desc. Pl. Montaub. 120 = Diplotaxis viminea.
virgatum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 93.—Am. bor.
Virletii, *Fourn. Thèse Crucif.* 62.—Mexic.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1870) II. 669.—Chili.
volgense, *Bieb. ex Fourn. Thèse Crucif.* 97.—Rossia.
vulgare, Pers. Syn. ii. 196 = Nasturtium sylvestre.
Wallichii, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 158.—Reg. Himal.
Walteri, Ell. Sketch, ii. 146 = Nasturtium tanacetifolium.
Waltheri, Crantz, Stirp. Austr. ed. II. 51 = orientale.
Weddellii, *Fourn. Thèse Crucif.* 104.—Boliv.
Zanonii, Caruel, in Atti Soc. Sc. Nat. Milan. v. (1863) 149 (= *Erucastrum Zanonii*).—Italia.
Zanonii, J. Gay, in Bull. Soc. Bot. Fr. vii. (1860) 881 = pinnatifidum.
Zeae, Spreng. Syst. ii. 904 = erysimoides.

SISYNDITE, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 354 (1860). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 265.
spartea, *E. Mey. ex Sond. l. c.*—Afr. austr.

SISYRANTHUS, E. Mey. Comm. Pl. Afr. Austr. 197 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 779.
imberbis, *Harv. Thes. Cap.* ii. 11. t. 116.—Afr. austr.
virgatus, *E. Mey. Comm. Pl. Afr. Austr.* 197.—Afr. austr.

SISYRINCHIUM, Eckl. Top. Verz. 16 (1827) = **Aristea**, Soland. (Irid.).
inundatum, Eckl. l. c. = A. melaleuca.
melaleucum, Eckl. l. c. = A. melaleuca.
pusillum, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 111 = A. pusilla.
spirale, Eckl. Top. Verz. 16 = A. spiralis.

SISYRINCHIUM, Linn. Gen. ed. I. 273 (1737). *IRIDEAE*, Benth. & Hook. f. iii. 698.
BERMUDIANA, Adans. Fam. ii. 60 (1763).
ECHTHRONEMA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85.
ERIPHILEMA, Herb. l. c. (1843).
GLUMOSIA, Herb. l. c. (1843).
HYDASTYLUS, Dryand. ex Salisb. in Trans. Hort. Soc. i. (1812) 310.
OLSYNIUM, Rafin. New Fl. Am. i. 72 (1836).
PANEGUIA, Rafin. Fl. Tellur. iv. 34 (1836).
POGADELPHA, Rafin. l. c. 29 (1836).
SOUZA, Vell. Fl. Flum. 273 (1825).
acaule, Klatt, in Linnaea, xxxiv. (1865–66) 736 = Solenomelus acaulis.
acre, *H. Mann, in Proc. Am. Acad.* vii. (1867) 206.—Ins. Sandvic.
acuminatum, Herb. in Lindl. Bot. Reg. (1843) Misc. 84 = angustifolium.
adenocarpum, *Steud. Nom.* ed. II. ii. 595.—Chili.
adenostemon, Phil. in Linnaea, xxix. (1857–58) 62 = cuspidatum.
adscendens, Phil. f. Cat. Pl. Vasc. Chil. 281 = graminifolium.
affine, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 111.—Mexic.
alatum, *Hook. Ic. Pl.* t. 219.—Am. trop.
albidum, Rafin. Atl. Journ. 17 = S. Bermudiana.
altissimum, Tenore, in Atti 3ª Riun. Sc. Ital. (1841) 504 = palmifolium.
anceps, Cav. Diss. vi. 345. t. 190. f. 2 = angustifolium.
andinum, Phil. in Linnaea, xxix. (1857–58) 62 = Chamelum andinum.
angustifolium, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.; Hibern.
aphyllum, Spreng. Syst. i. 166 = Bobartia aphylla.
arenarium, Poepp. Fragm. Syn. Pl. Chile, 3 = cuspidatum.
ascendens, Poepp. l. c. 4 = graminifolium.
asperulum, Phil. in Linnaea, xxxiii. (1864–65) 252 = junceum.
aurantiacum, Griseb. in Goett. Abh. xxiv. (1879) 327 = monostachyum.
aurantiacum, Zuccagni, in Roem. Collect. 145 = Homeria collina.
aureum, *Vell. Fl. Flum.* ix. t. 69.—Bras.
avenaceum, *Klatt, in Linnaea*, xxxi. (1861–62) 373.—Bras.
azureum, *Phil. Fl. Atac.* 50.—Chili.
Bakeri, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 378 = trinerve.
bellum, S. Wats. in Proc. Am. Acad. xii. (1877) 277 = angustifolium.
Bermudiana, *Linn. Sp. Pl.* 954, partim.—Ins. Bermud.
Bermudiana, Linn. l. c., partim = angustifolium.
Berteroanum, Phil. in Linnaea, xxix. (1857–58) 58 = leucanthum, *Colla*.
Berteroi, Steud. ex Baker, in Journ. Linn. Soc. xvi. (1877) 117 = chilense.
Berteroi, Steud. ex Klatt, in Mart. Fl. Bras. iii. 1. 537 = iridifolium.
biflorum, *Griseb. in Goett. Abh.* xxiv. (1879) 328.—Reg. Argent.
bogotense, *H. B. & K. Nov. Gen. et Sp.* i. 323.—N. Granat.
brevifolium, Klotzsch, ex Klatt, in Mart. Fl. Bras. iii. 1. 538 = incurvatum.
bulbosum, Mill. Gard. Dict. ed. VIII. n. 3 = Eleutherine plicata.
californicum, [*Dryand. in*] *Hort. Kew.* ed. II. iv. 135.—Am. bor. occ.
capitatum, Pers. Syn. i. 49 = Eleutherine plicata.
Cervantesii, *E. Mey. ex Presl, Rel. Haenk.* i. 117.—Mexic.
chilense, *Hook. Bot. Mag.* t. 2786.—Am. trop.
Claraxii, Baker, in Journ. Linn. Soc. xxi. (1884) 235 = monostachyum.
coerulescens, F. Muell. Fragm. vii. 91 = Libertia caerulescens.
coeruleum, *Vell. Fl. Flum.* ix. t. 66.—Bras.
colchaguense, Phil. in Linnaea, xxxiii. (1864–65) 253 = junceum.
collinum, Cav. Diss. vi. 346 = Homeria collina.
Comes, *Steud. Nom.* ed. II. ii. 595.—Bras.
commutatum, *Klatt, in Hamb. Gartenz.* xvi. (1860) 164.—Bras.
congestum, Klatt, in Linnaea, xxxi. (1861–62) 98 = palmifolium?

SISYRINCHIUM:—
convolutum, Klatt, in Linnaea, xxxiv. (1865–66) 735 = californicum.
convolutum, *Nocca, Ticin. Hort. Pl. Select.* i. t. 1.—Am. trop.
crenatum, Vell. Fl. Flum. ix. t. 67 = Trimezia lurida.
cultrifolium, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 (Quid?).—Java.
Cummingii, *Penny, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 673, nomen.—Chili.
cuspidatum, *Poepp. Fragm. Syn. Pl. Chile*, 3.—Chili.
cyaneum, Lindl. Bot. Reg. t. 1090 = Orthrosanthus multiflorus.
dasycarpum, Phil. in Linnaea, xxix. (1857–58) 61 = graminifolium.
depauperatum, Phil. l. c. = leucanthum, *Colla*.
dichroum, Poepp. ex Klatt, in Mart. Fl. Bras. iii. 1. 537 = iridifolium.
Douglasii, A. Dietr. Sp. Pl. ii. 504 = grandiflorum.
elatum, Seub. ex Klatt, in Mart. Fl. Bras. iii. 1. 544 = palmifolium.
elegans, [DC. in] Red. Lil. t. 171 = Homeria collina.
elegans, Willd. Sp. Pl. iii. 577 = Homeria elegans.
excisum, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 447 = S. Bermudiana.
fasciculatum, *Klatt, in Linnaea*, xxxi. (1861–62) 97.—Uruguay; Paraguay.
fernandezianum, *Steud. & Hochst. ex Steud. Nom.* ed. II. ii. 595.—Ins. J. Fernand.
filifolium, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 101.—Ins. Falkland.
filiforme, Gaudich. in Freyc. Voy. Bot. 421 = filifolium.
filiforme, Rafin. Atl. Journ. 178 (Quid?).—Am. bor.
filiforme, Spreng. Syst. i. 166 = Bobartia filiformis.
filiforme, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 595 = junceum.
fimbriatum, Domb. ex Klatt, in Mart. Fl. Bras. iii. 1. 537 = iridifolium.
flavidum, Kellogg, in Proc. Calif. Acad. ii. (1863) 50. t. 3 = californicum.
flavum, Hoffmgg. ex Steud. Nom. ed. II. ii. 595 = convolutum.
flavum, Miers, ex Steud. l. c. = Solenomelus chilensis.
flexuosum, Lindl. in Bot. Reg. sub t. 1067 = cuspidatum.
flexuosum, *Phil. in Linnaea*, xxxiii. (1864–65) 251.—Chili.
flexuosum, Spreng. Syst. i. 167 = Hexaglottis longifolia.
floribundum, Phil. in Linnaea, xxix. (1857–58) 59 = junceum.
fluminense, Vell. Fl. Flum. ix. t. 68 = Trimezia lurida.
formosum, F. Muell. Fragm. vii. 91 = Libertia formosa.
frigidum, Poepp. Fragm. Syn. Pl. Chile, 1 = Chamelum andinum.
galaxioides, Gomes, in Mem. Acad. Lisboa, iii. (1812) Mem. dos Corresp. 99 = Trimezia lurida.
Gaudichaudii, A. Dietr. Sp. Pl. ii. 505 = filifolium.
geniculatum, Herb. in Lindl. Bot. Reg. (1843) Misc. 84 = angustifolium.
giganteum, Tenore, Cat. Ort. Nap. (1845) 96 = palmifolium.
gladiatum, Spreng. Syst. i. 166 = Bobartia gladiata.
glaucophyllum, *Penny, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 601, nomen.—Am. bor.
Glaziovii, *Baker, in Journ. Bot.* xiv. (1876) 268.—Bras.
gracile, *Klotzsch, ex Baker, l. c.*—Bras.
gracile, Phil. in Linnaea, xxix. (1857–58) 60 = junceum.
gramineum, Lam. Encyc. i. 408; Curt. Bot. Mag. t. 464 = angustifolium.
graminifolium, Bert. ex Steud. Nom. ed. II. ii. 596 = chilense.
graminifolium, *Lindl. in Bot. Reg.* t. 1067.—Chili.
grandiflorum, Cav. Diss. vi. 345 = Tigridia lutea.
grandiflorum, *Dougl. ex Lindl. Bot. Reg.* t. 1364.—Am. bor. occ.
grandiflorum, Poepp. Fragm. Syn. Pl. Chile, 4 = Calydorea speciosa.
guttatum, *Moris, Enum. Hort. Taur.* (1833) 30.—Chili.
Hartwegii, Baker, in Journ. Linn. Soc. xvi. (1877) 119, nomen = tenuifolium.
hirtellum, *Penny, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 673, nomen.—Am. bor.

SISYRINCHIUM:—
homomallum, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 378 = angustifolium ?
Huismo, Domb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 120 = sessiliflorum.
Illmu, Molina, Sagg. Chile, ed. II. 110 = Conanthera bifolia.
incurvatum, *Gardn. in Hook. Ic. Pl.* t. 513.—Bras.
iridifolium, *H. B. & K. Nov. Gen. et Sp.* i. 324.—Am. trop.
iridioides, Curt. Bot. Mag. t. 94 = S. Bermudiana.
ixioides, Forst. f. Prod. 61 = Libertia ixioides.
Jamesoni, *Baker, in Journ. Bot.* xiv. (1876) 269.—Ecuador.
junceum, *E. Mey. in Presl, Rel. Haenk.* i. 118.—Reg. Andin.
juncifolium, Herb. in Lindl. Bot. Reg. (1843) Misc. 85 = junceum ?
junciforme, Poepp. Fragm. Syn. Pl. Chile, 2 = junceum.
laterale, *Baker, in Journ. Bot.* xiv. (1876) 269.—Bolivia.
latifolium, Sw. Prod. Veg. Ind. Occ. 17 = Eleutherine plicata.
laxum, Otto, ex Sims, Bot. Mag. t. 2312 = iridifolium.
Lechleri, Steud. ex Klatt, in Mart. Fl. Bras. iii. 1. 535 = chilense.
leucanthum, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 18.—Chili; Reg. Argent.
leucanthum, *Steud. Nom.* ed. II. ii. 596.—Chili.
libertioides, *Steud. & Hochst. ex Steud. l. c.*—Chili.
lineatum, Torr. in Pacif. Rail. Rep. iv. 143 = californicum.
longistylum, Lem. in Fl. des Serres, iii. (1847) t. 255 = Solenomelus chilensis.
lutescens, Lodd. Bot. Cab. t. 1870 = striatum.
luteum, Bert. ex Steud. Nom. ed. II. ii. 596 = tinctorium.
luteum, Fisch. ex Link, Enum. Hort. Berol. ii. 177 = convolutum.
Luzula, *Klotzsch, ex Klatt, in Linnaea*, xxxi. (1861–62) 89, 376.—Bras.
macranthum, *Griseb. in Goett. Abh.* xxiv. (1879) 326.—Reg. Argent.
macrocarpum, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 71.—Reg. Argent.
macrocephalum, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1833) 176 = palmifolium.
maculatum, Hook. Bot. Mag. t. 3197 = graminifolium.
majale, Link, Klotzsch & Otto, Ic. Pl. Hort. Berol. 24. t. 10 = graminifolium.
majus, Redhead, in Journ. Linn. Soc. ix. (1867) 209 = Iris Sisyrinchium.
Mandoni, *Baker, in Journ. Bot.* xiv. (1876) 269.—Bolivia.
Marchio, Steud. Nom. ed. II. ii. 596 – alatum.
marginatum, Klatt, in Linnaea, xxxi. (1861–62) 83 = palmifolium.
micranthemum, Pers. Syn. i. 50 = seq.
micranthum, *Cav. Diss.* vi. 345. t. 191. f. 2.—Peruv.
minus, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 263 = angustifolium.
minutiflorum, *Klatt, in Linnaea*, xxxi. (1861–62) 71.—Reg. Argent.
monostachyum, *Baker, in Journ. Bot.* xiv. (1876) 268.—Reg. Argent.
Moritzianum, Klatt, in Linnaea, xxxi. (1861–62) 378 = Orthrosanthus chimboracensis.
mucronatum, Michx. Fl. Bor. Am. ii. 33 = angustifolium.
multicaule, Schau. in Linnaea, xix. (1847) 704 = chilense.
multiflorum, *Lem. Illustr. Hortic.* vi. (1859) t. 221.—Am. bor.
multiflorum, Steud. Nom. ed. II. ii. 596 = Orthrosanthus multiflorus.
narcissoides, Cav. Diss. vi. 347. t. 191. f. 3 = Symphyostemon narcissoides.
nervosum, *Phil. in Linnaea*, xxxiii. (1864–65) 251.—Chili.
nigricans, C. Gay, Fl. Chil. vi. 20 = striatum.
Nunna, *Steud. & Hochst. ex Steud. Nom.* ed. II. ii. 596.—Chili.
Nuno, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 17. t. 54.—Chili.
Nuttallii, Sweet, Hort. Brit. ed. I. 395 = S. Bermudiana.

SISYRINCHIUM:—
Occissapungum, Ruiz, ex Klatt, in Linnaea, xxxi. (1861–62) 379 = Orthrosanthus chimboracensis.
odoratissimum, Lindl. Bot. Reg. t. 1283 = Symphyostemon odoratissimus.
pachyrhizum, *Baker, in Journ. Bot.* xiv. (1876) 269.—Bras.
palmifolium, Cav. Diss. vi. 348 = Eleutherine plicata.
palmifolium, *Linn. Mant.* i. 122, excl. syn.—Am. austr.
paniculatum, R. Br. Prod. 305 = Libertia paniculata.
patagonicum, *Phil. ex Baker, in Journ. Linn. Soc.* xvi. (1877) 117, nomen.—Patagon.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864–65) 251.—Chili.
pedunculatum, Gill. ex Hook. Bot. Mag. t. 2965 = Solenomelus chilensis.
Philippi, Klatt, in Linnaea, xxxi. (1861–62) 89 = junceum.
piliferum, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 379.—Paraguay.
plicatum, Spreng. Syst. i. 167 = Eleutherine plicata.
polystachyum, F. Muell. Census, 114 = Orthrosanthus polystachyus.
Proregia, *Steud. Nom.* ed. II. ii. 596.—Bras.
pubescens, Cham. ex Steud. Nom. ed. II. ii. 596 = iridifolium.
pulchellum, R. Br. Prod. 305 = Libertia pulchella.
pumilum, Hook. f. Fl. Antarct. t. 129 = Tapeinia magellanica.
pusillum, *H. B. & K. Nov. Gen. et Sp.* i. 323.—Ecuador.
pygmaeum, Klotzsch, ex Phil. f. Cat. Pl. Vasc. Chil. 282 = tenuifolium.
quinquevulnerum, Domb. ex Klatt, in Mart. Fl. Bras. iii. 1. 536 = chilense.
racemosum, Pers. Syn. i. 49 = palmifolium.
ramosum, Herb. in Lindl. Bot. Reg. (1843) Misc. 84 = angustifolium, chilense.
restioides, *Spreng. Syst.* i. 166.—Am. trop.
reticulatum, Auct. ex Pers. Syn. i. 50 = striatum.
rigidum, Lehm. in Hamb. Gartenz. vi. (1850) 415 = tinctorium.
roseum, Herb. in Lindl. Bot. Reg. (1843) Misc. 85; Phil. in Linnaea, xxix. (1857–58) 60 = junceum
scabriusculum, *Steud. Nom.* ed. II. ii. 596.—Chili.
scabrum, Phil. in Linnaea, xxix. (1857–58) 58 = junceum.
scabrum, Schlecht. & Cham. l. c. vi. (1831) 57 = chilense.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 160.—Mexic.
scirpeum, Phil. in Linnaea, xxix. (1857–58) 59 = junceum.
scirpiforme, Poepp. Fragm. Syn. Pl. Chile, 2 = junceum.
secundiflorum, *Klatt, in Linnaea*, xxxi. (1861–62) 91.—Bras.
Segethi, Phil. l. c. xxix. (1857–58) 61 = Solenomelus Lechleri.
Sellowianum, *Klatt, l. c.* xxxi. (1861–62) 375.—Uruguay.
Sellowii, Klotzsch, ex Baker, in Journ. Linn. Soc. xvi. (1877) 119 = pachyrhizum.
sertiflorum, Salisb. in Trans. Hort. Soc. i. (1812) 309 = striatum.
sessiliflorum, Hook. & Arn. Bot. Beech. Voy. 47 = Libertia caerulescens.
sessiliflorum, *Poepp. Fragm. Syn. Pl. Chile*, 2.—Chili.
setaceum, *Klatt, in Linnaea*, xxxi. (1861–62) 85.—Reg. Argent.
spathaceum, Pers. Syn. i. 50 = Bobartia indica.
speciosum, Hook. Bot. Mag. t. 3544 = Calydorea speciosa.
spicatum, Cav. Ic. ii. 2 = striatum.
spicatum, Seub. ex Klatt, in Linnaea, xxxi. (1861–62) 377 = Orthrosanthus spicatus.
striatum, *Sm. Ic. Pict.* t. 9.—Chili; Reg. Argent.
sulcatum, Gill. ex Hook. Ic. Pl. t. 218 = vaginatum.
tenuifolium, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 691.—Mexic.
teres, Rafin. Atl. Journ. 148 (Quid ?).—Am. bor.
tinctorium, *H. B. & K. Nov. Gen. et Sp.* i. 324.—Reg. Andin.; Mexic.
tingens, Steud. Nom. ed. II. ii. 596 = tinctorium.
tolucense, *Peyr. in Linnaea*, xxx. (1859–60) 18.—Mexic.
triflorum, *Schrank, Pl. Rar. Hort. Monac.* t. 61.—Hab.?
trinerve, *Baker, in Journ. Bot.* xiv. (1876) 267.—Bolivia.
triquetrum, Hort. ex Steud. Nom. ed. II. ii. 596 = convolutum.

SISYRINCHIUM:—
unguiculatum, *Griseb. in Goett. Abh.* xxiv. (1879) 326.—Reg. Argent.
uniflorum, C. Gay, ex Phil. in Linnaea, xxxix. (1857–58) 63 = chilense.
unispathaceum, *Klatt, in Linnaea*, xxxiv. (1865–66) 737.—Bolivia.
vaginatum, *Spreng. Syst.* i. 166.—Am. trop.
Weirii, *Baker, in Journ. Bot.* xiv. (1876) 268.—Bras.
xiphioides, Poepp. Fragm. Syn. Pl. Chile, 4 = Calydorea speciosa.

SISYRRINCHIUM, Hook. & Arn. Bot. Beech. Voy. 46 (1830), sphalm. = praec.

SITANION, Rafin. in Journ. Phys. lxxxix. (1819) 103 = **Elymus**, Linn. (Gramin.).
elymoides, Rafin. l. c. = E. Sitanion.

SITILIAS, Rafin. New Fl. Am. iv. 85 (1836) = **Pyrrhopappus**, DC. (Compos.).
caroliniana, Rafin. l. c. = P. carolinianus.

SITOCODIUM, Salisb. Gen. Pl. Fragm. 27 (1866) = **Camassia**, Lindl. (Liliac.).
esculentum, Salisb. l. c. = C. esculenta.

SITODIUM, Banks, ex Gaertn. Fruct. i. 344. tt. 71, 72 (1788) = **Artocarpus**, Forst. (Urtic.).
cauliflorum, Gaertn. l. c. 345. tt. 71, 72 = A. integrifolia.

SITOSPELOS, Adans. Fam. ii. 36 (1763) = **Elymus**, Linn. (Gramin.).

SIUM, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. iii. 893.
BERULA, Hoffm. ex Bess. Enum. Pl. Volh. 44 (1822).
MAUCHARTIA, Neck. Elem. i. 172 (1790).
Amomum, Roth, Tent. Fl. Germ. ii. 1. 336 = Sison Amomum.
angustifolium, Linn. Sp. Pl. ed. II. 1672 = erectum.
angustifolium, Thunb. Prod. Pl. Cap. 50 = Thunbergii.
annuum, Roth, Tent. Fl. Germ. i. 128 = Seseli annuum.
Apium, Roth, l. c. = Apium graveolens.
aquaticum, Gueldenst. ex Ledeb. Fl. Ross. ii. 259 = latifolium.
aromaticum, Lam. Fl. Fr. iii. 458 = Sison Amomum.
asperum, Thunb. Prod. Pl. Cap. 51 = Seseli asperum.
Berula, J. F. Gmel. Syst. 482 = latifolium.
Berula, Gouan, Fl. Monsp. 218 = erectum.
brevifolium, Hort. ex DC. Prod. iv. 124 = S. Sisarum.
Bulbocastanum, Spreng. in Schult. Syst. vi. 538 = Carum Bulbocastanum.
bulbosum, Poir. Encyc. Suppl. i. 622 = Erigenia bulbosa.
bulbosum, Thore, in Desv. Journ. Bot. i. (1808) 193 = Carum inundatum.
Burchellii, *Hemsl. Bot. Voy. Challenger*, i. 11. 68.—Ins. S. Helena.
californicum, *Baill. Hist. Pl.* vii. 181, 182.—Calif.
canadense, Lam. Encyc. i. 407 = Cryptotaenia canadensis.
Carsonii, *Durand, ex A. Gray, Man. Bot. N. U. St.* ed. V. 196.—Am. bor. occ.
Carum, [Weber, in] Wigg. Prim. Fl. Holsat. 24 = Carum Carvi.
Carvi, Bernh. Syst. Verz. Erf. 173 = Carum Carvi.
Cicuta, [Weber, in] Wigg. Prim. Fl. Holsat. 24 = Cicuta virosa.
cicutaefolium, *Schrank, Baier. Fl.* i. 558.—Reg. bor. temp.
cicutarium, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 639, nomen = cicutaefolium.
Cordiennii, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 404 = Silaus carvifolius.
crassipes, Spreng. Syst. iv. Cur. Post. 120 = Apium crassipes.
crinitum, Poir. Encyc. Suppl. i. 622 = Schultzia crinita.
crispum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 27 = Ceratopteris thalictroides (Crypt.).
cuneifolium, Boiss. Fl. Orient. ii. 1150, sphalm. = lancifolium.
Cyminosma, *Basin. in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 204.—Reg. Casp.
decumbens, *Thunb. Fl. Jap.* 118.—Japon.

SIUM:—
denticulatum, Baldw. ex Ell. Sketch, i. 354 = Peucedanum ternatum.
Douglasii, DC. Prod. iv. 125 = Conium maculatum ?
erectum, *Huds. Fl. Angl.* ed. I. 103.—Europ.; Caucas.
Falcaria, Linn. Sp. Pl. 252 = Falcaria vulgaris.
falcatum, Dubois, Méthod. ed. I. 442; ed. II. 407 = Falcaria vulgaris.
Ferula, Raeusch. Nom. ed. III. 81 = erectum.
ferulaefolium, Spreng. Sp. Umbellif. 95 = Carum ferulaefolium.
filifolium, Steud. Nom. ed. II. ii. 596 = Peucedanum teretifolium.
filifolium, Thunb. Prod. Pl. Cap. 50 = Oenanthe filiformis.
gallabatense, Schweinf. Beitr. Fl. Aethiop. 273 = Thunbergii.
graecum, Guss. ex Spreng. Syst. i. 914 = Ferula nudicaulis.
graecum, Linn. Sp. Pl. 252 = Bonannia resinifera.
graecum, Lour. Fl. Cochinch. 179 = decumbens.
grandiflorum, Thunb. Prod. Pl. Cap. 50 (= *Levisticum grandiflorum.*).—Afr. austr.
graveolens, Vest, Man. Bot. 517 = Apium graveolens.
helenianum, *Hook. f. in Hook. Ic. Pl.* t. 1032.—Ins. S. Helena.
Hippomarathrum, Roth, Tent. Fl. Germ. i. 128 = Seseli Hippomarathrum.
hispidum, Thunb. Prod. Pl. Cap. 51 = Carum hispidum.
incisum, Pers. Syn. i. 316 = erectum.
intermedium, DC. Fl. Fr. iv. 302 = Carum inundatum.
intermedium, Tenore, Fl. Nap. iii. 310 = Apium nodiflorum.
inundatum, Lam. Fl. Fr. iii. 460 = Apium inundatum.
japonicum, *Thunb. Fl. Jap.* 118.—Japon.
javanicum, Blume, Bijdr. 881 = Falcaria javanica.
laciniatum, Blume, l. c. = Falcaria laciniata.
lancifolium, *Bieb. Fl. Taur. Cauc.* iii. 230.—Reg. Caucas.
lancifolium, Schrank, Baier. Fl. i. 556 = latifolium.
latifolium, Bieb. Fl. Taur. Cauc. i. 225 = lancifolium.
latifolium, Bigel. Fl. Bost. 69 = Apium lineare.
latifolium, *Linn. Sp. Pl.* 251.—Europ.; Am. bor.
latijugum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 683.—Reg. Himal.
latissimum, Bor. Fl. Centr. Fr. ed. III. ii. 274, in obs. = latifolium.
limosum, Moris, ex DC. Prod. iv. 104 = Apium crassipes.
lineare, Michx. Fl. Bor. Am. i. 167 = Apium lineare.
lobatum, Moon, Cat. Pl. Ceyl. 22 = Heracleum nepalense.
longifolium, J. & C. Presl, Fl. Čech. 65 = latifolium.
longifolium, Pursh, Fl. Am. Sept. i. 194 = Peucedanum ternatum.
luteum, Spreng. Sp. Umbellif. 92 = Muretia tanaicensis.
marginatum, Poir. Encyc. Suppl. i. 622 = Peucedanum ternatum.
medium, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 19.—Sibir. alt.; Soongar.
napiforme, Schult. Syst. vi. 539 = Carum ferulaefolium.
Ninsi, Linn. Sp. Pl. 25 (Quid ?).—China.
Ninsi, Thunb. Fl. Jap. 118 = Aralia quinquefolia.
nipponicum, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 286.—Japon.
nodiflorum, Linn. Sp. Pl. 251 = Apium nodiflorum.
nodiflorum, Oed. Fl. Dan. t. 247 = erectum.
nudicaule, Lam. Encyc. i. 407 = Peucedanum redivivum.
oppositifolium, *Kit. ex Schult. Oester. Fl.* ed. II. i. 495.—Hungar.
paniculatum, Thunb. Prod. Pl. Cap. 51 = Rhyticarpus rugosus.
patulum, Thunb. l. c. = Selinum suffruticosum.
peucedanoides, Spreng. Sp. Umbellif. 92 = Muretia tanaicensis.
Podagraria, [Weber, ex] Wigg. Prim. Fl. Holsat. 24 = Aegopodium Podagraria.
podolicum, Bess. ex Reichb. Fl. Germ. Excurs. 479 = lancifolium.
podolicum, Hort. ex G. Don, in Loud. Hort. Brit. 473 = S. Sisarum.
pusillum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 611 = erectum.
pusillum, Poir. Encyc. Suppl. i. 622 = Apium divaricatum.
radiatum, Viv. Fl. Lib. Spec. 16. t. 9. f. 2 = Apium nodiflorum.
repens, Jacq. Fl. Austr. iii. 34 = Apium nodiflorum.

SIUM:—
repens, Linn. f. Suppl. 181 = Apium repens.
rigidius, Linn. Sp. Pl. 251; Walt. Fl. Carol. 114 = Peucedanum ternatum.
rigidum, Hill, Veg. Syst. vi. 34 = praec. ?
rugosum, Rafin. Med. Fl. ii. 264; Atl. Journ. 151 = Apium lineare.
saxifragum, Roth, Tent. Fl. Germ. i. 128 = Carum Bunius.
segetum, Lam. Fl. Fr. iii. 458 = Carum segetum.
siculum, Linn. Sp. Pl. 252 = Kundmannia sicula.
Silaus, Roth, Tent. Fl. Germ. i. 129 = Silaus flavescens.
simense, J. Gay, ex A. Rich. Tent. Fl. Abyss. i. 324 = Pimpinella simensis.
sisaroideum, DC. Prod. iv. 124 = lancifolium.
Sisarum, *Linn. Sp. Pl.* 251.—As. or.
stoloniferum, Guss. Suppl. Fl. Sic. Prod. 77 = Apium nodiflorum.
suave, Walt. Fl. Carol. 115 = Apium lineare.
sulcatum, Pers. Syn. i. 316 = latifolium.
tenuifolium, Muhl. Cat. 30 = Apium lineare.
teretifolium, Ell. Sketch, i. 354 = Peucedanum teretifolium.
Thunbergii, *DC. Prod.* iv. 125.—Afr. austr.
tortuosum, Roth, Tent. Fl. Germ. i. 128 = Seseli annuum.
tricuspidatum, Ell. Sketch, i. 354 = Peucedanum ternatum.
trifoliatum, Poir. Encyc. Suppl. i. 622 = Thaspium aureum.
triternatum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 57 = Ligusticum acutilobum.
triternatum, Moon, Cat. Pl. Ceyl. 22 = Pimpinella Heyneana.
verrucosum, J. Gay, ex A. Rich. Tent. Fl. Abyss. i. 324 = Pimpinella simensis.
verticillatum, Lam. Fl. Fr. iii. 460 = Carum verticillatum.
verticillatum, Pett. ex Reichb. Fl. Germ. Excurs. 474 = Carum ammoides.
villosum, Thunb. Prod. Pl. Cap. 51 = Annesorhiza villosa.
virens, Hort. ex Steud. Nom. ed. II. ii. 597 = Selinum sylvestre.
virescens, Spreng. Sp. Umbellif. 94 = Silaus carvifolius.
Visnaga, Stokes, Bot. Mat. Med. ii. 106 = Ammi Visnaga.
vulgare, Bernh. Syst. Verz. Erf. 173 = Aegopodium Podagraria.
vulgare, Gueldenst. ex Ledeb. Fl. Ross. ii. 259 = latifolium.

SIUMIS, Rafin. New Fl. Am. iv. 31 (1836) = **Apium**, Tourn. (Umbell.).
heterophyla, Rafin. l. c. = A. lineare.

SIXALIX, Rafin. Fl. Tellur. iv. 95 (1836) = **Scabiosa**, Tourn. (Dipsac.).
daucoides, Rafin. l. c. 96 = Sc. daucoides.

SIZYGIUM, Duch. in Orb. Dict. viii. 554 (1849), sphalm. = SYZYGIUM, Gaertn. = **Eugenia**, Michx. (Myrtic.).

SKAPHIUM, Miq. Fl. Ind. Bat. Suppl. 357 (1860) = **Xanthophyllum**, Roxb. (Polygal.).
lanceatum, Miq. l. c.—Sumatra.

SKIDANTHERA, Rafin. Sylva Tellur. 60 (1838) = **Elaeocarpus**, Burm. (Tiliac.).
dentata, Rafin. l. c. = E. dentatus.

SKILLA, Rafin. Fl. Tellur. ii. 13 (1836) = **Scilla**, Linn. (Liliac.).

SKIMMI, Adans. Fam. ii. 364 (1763) = seq.

SKIMMIA, Thunb. Nov. Gen. 57 (1783). *RUTACEAE*, Benth. & Hook. f. i. 302.
ANQUETILIA, Decne. in Jacquem. Voy. Bot. 161. t. 161 (1848 ?).
LAUREOLA, M. Roem. Syn. Hesper. 74 (1846).
SKIMMI, Adans. Fam. ii. 364 (1763).
Foremanni, Hort. ex H. Knight, in Flor. & Pomol. (1881) 70 = japonica.
fragrans, Carr. in Rev. Hortic. (1869) 258 = japonica ?

SKIMMIA:—
fragrantissima, Hort. ex T. Moore, in Proc. Linn. Soc. (1866–67) 7 = japonica.
intermedia, Carr. in Rev. Hortic. (1870) 200 = japonica.
japonica, Lindl. in Gard. Chron. (1852) 183; et Fortune, l. c. 739 (non seq.).—China.
japonica, *Thunb. Nov. Gen.* 58.—Japon.
Laureola, *Sieb. & Zucc. ex Walp. Rep.* v. 405.—Reg. Himal.
oblata, T. Moore, in Gard. Chron. (1864) 962 = japonica.
rubella, *Carr. in Rev. Hortic.* (1874) 311.—China.
Veitchii, *Carr. l. c.* (1869) 259.—Japon.

SKINNERA, Forst. Char. Gen. 57. t. 29 (1776) = **Fuchsia**, Plum. (Onagrar.).
antarctica, Forst. ex A. DC. in DC. Prod. xvii. 295 (= *Blackwellia antarctica*, Gaertn).—Hab. ?
coccinea, Moench, Meth. Suppl. 270 = F. coccinea.
excorticata, Forst. Char. Gen. 58 = F. excorticata.

SKINNERIA, Choisy in Mém. Soc. Phys. Genèv. vi. (1833) 487 (Conv. Or. 105. t. 6) = **Ipomoea**, Linn. (Convolv.).
caespitosa, Choisy, ll. cc. = I. linifolia.

SKIOPHILA, Hanst. in Linnaea, xxvi. (1853-55) 207 = **Episcia**, Mart. (Gesner.).
melittifolia, Hanst. l. c. = E. melittifolia.
pulchella, Hanst. l. c. = Tussaca pulchella.

SKIRRHOPHORUS, DC. ex Lindl. Introd. Nat. Syst. ed. II. 260 (1836) = **Angianthus**, Wendl. (Compos.).
Cunninghami, DC. Prod. vi. 150 = A. Cunninghamii.
demissus, A. Gray, in Hook. Kew Journ. iii. (1851) 149 = A. demissus.
Drummondii, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 188 = A. Drummondii.
eriocarpus, F. Muell. Fragm. iii. 156 = Gnephosis eriocarpa ?
eriocephalus, Hook. f. ex A. Gray, in Hook. Kew Journ. iii. (1851) 148 = A. eriocephalus.
mucronulatus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 72 = A. pygmaeus.
Muellerianus, Sond. in Linnaea, xxv. (1852) 486 = A. strictus.
phyllocephalus, F. Muell. Fragm. iii. 159 = A. phyllocephalus.
Preissianus, Steetz, in Lehm. Pl. Preiss. i. 439 = A. Preissianus.
pygmaeus, A. Gray, in Hook. Kew Journ. iii. (1851) 148 = A. pygmaeus.
strictus, A. Gray, l. c. 149 = A. strictus.

SKIRROPHORUS, C. Muell. in Walp. Ann. v. 263 (1859), sphalm. = praec.

SKIZIMA, Rafin. Fl. Tellur. ii. 32 (1836) = **Wurmbea**, Thunb. (Liliac.).
pumila, Rafin. l. c. = W. capensis.

SKOFITZIA, Hassk. & Kanitz, in Oestr. Bot. Zeitschr. xxii. (1872) 147 = **Tradescantia**, Linn. (Commel.).
boliviana, Hassk. & Kanitz, l. c. = T. ambigua.

SKOINOLON, Rafin. Fl. Tellur. iv. 27 (1836) = **Schoenocaulon**, A. Gray (Liliac.).

SKOLEMORA, Arruda, in Koster, Trav. Braz. 498 (1816). (Quid ?).
pernambucensis, Arruda, l. c.—Bras.

SKYTALANTHUS, Endl. Gen. Suppl. iii. 75 (1843) = seq.

SKYTANTHUS, Meyen, Reise, i. 376 (1834). *APOCYNACEAE*, Benth. & Hook. f. ii. 704.
HABSBURGIA, Mart. Syst. Mat. Med. Bras. 36 (1843).
NERIANDRA, A. DC. in DC. Prod. viii. 422 (1844).
SCYTALANTHUS, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 361.
acutus, *Meyen, Reise*, i. 376.—Chili.
hancorniaefolius, *Miers, Apocyn. S. Am.* 109.—Bras.
havanensis, Miers, l. c. 110 = Cameraria havanensis (in Addend.).
Martianus, *Miers, l. c.*—Bras.

SLACKIA, Griff. Itin. Notes, 187 (1848) = **Decaisnea**, Hook. f. & Thoms. (Berber.).
insignis, Griff. l. c. = D. insignis.

SLACKIA, Griff. Notul. iv. 158 (1854). *GESNERACEAE*, Benth. & Hook. f. ii. 1017.
Griffithii, *C. B. Clarke, in DC. Monog. Phan.* v. 188. —Burma.

SLACKIA, Griff. Notul. iii. 162 (1854) = **Iguanura**, Blume (Palmae).
geonomiformis, Griff. l. c. = I. geonomaeformis.

SLADENIA, Kurz, in Journ. Bot. xi. (1873) 194. *TERNSTROEMIACEAE*.
celastrifolia, *Kurz, l. c.*—China.

SLATERIA, Desv. Journ. Bot. i. (1808) 243 = **Ophiopogon**, Ker-Gawl. (Haemod.).
coerulea, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 143 = O. japonicus.
Jaburan, Siebold, in Verh. Bat. Gen. xii. (1830) 15 = O. Jaburan.
japonica, Desv. Journ. Bot. i. (1808) 244 = O. japonicus.
repens, Siebold, in Verh. Batav. Gen. xii. (1830) 15 = Liriope spicata.

SLEVOGTIA, Reichb. Consp. 133 (1828) = **Enicostema**, Blume (Gentian.).
maritima, Dalz. in Hook. Kew Journ. ii. (1850) 137 = E. littorale.
occidentalis, Griseb. in DC. Prod. ix. 66 = E. littorale.
orientalis, Griseb. l. c. = E. littorale.
verticillata, D. Don, in Trans. Linn. Soc. xvii. (1837) 532 = E. littorale.

SLOANA, Linn. Gen. ed. I. 153 (1737) = seq.

SLOANEA, Linn. Hort. Cliff. 210 (1737). *TILIACEAE*, Benth. & Hook. f. i. 238.
ABLANIA, Aubl. Pl. Gui. 585. t. 234 (1775).
ADENOBASIUM, Presl, Symb. Bot. i. 39. t. 27 (1830).
BLONDEA, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 100.
DASYCARPUS, Oerst. in Kjoeb. Vidensk. Meddel. (1857) 27.
DASYNEMA, Schott, in Spreng. Syst. iv. App. 408 (1827).
FORGETINA, Bocquill. ex Baill. Adansonia, vii. (1866) 49.
FOVEOLARIA, Meissn. Gen. Comm. 28 (1836).
OXYANDRA, Reichb. Handb. 303 (1837).
TRICHOCARPUS, Schreb. Gen. 366 (1789).
alnifolia, *Mart. in Flora*, xx. (1837) II. *Beibl.* 94.—Bras.
amygdalina, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 165.—Cuba.
Aubletii, Sw. Fl. Ind. Occ. ii. 940 = sinemariensis.
australis, F. Muell. Fragm. iv. 91 = Echinocarpus australis.
Berteriana, *Choisy, ex DC. Prod.* i. 516.—Ins. S. Doming.
brevipes, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 68.—Bras.
castanocarpa, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 345 = multiflora.
corymbiflora, *DC. Prod.* i. 516.—Afr. trop.
cuneifolia, *Mart. in Flora*, xx. (1837) II. *Beibl.* 94.—Bras.
curatellifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 164.—Cuba.
dentata, *Linn. Sp. Pl.* 512.—Bras.; Guian.
durissima, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 65.—Bras.
egensis, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1881) 329 = pubescens, *Benth.*
emarginata, Linn. Sp. Pl. 512 = Achras Sapota?
Fendleriana, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 70.—Venezuela.
floribunda, *Spruce, ex Benth. l. c.* 66.—Bras.
grandiflora, Sm. in Rees, Cycl. xxxiii. n. 2 = dentata.
guianensis, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 69 = cuneifolia.
hirsuta, *Planch. ex Benth. l. c.* 70 = monosperma.
jamaicensis, *Hook. Ic. Pl.* tt. 693–96.—Jamaic.
Langii, *A. Muell. Fragm.* v. 28.—Austral.

SLOANEA:—
laurifolia, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 70.—Bras.
laxiflora, *Spruce, ex Benth. l. c.* 65.—Venezuela.
longiseta, *Wright, ex Griseb. Cat. Pl. Cub.* 30.—Cuba.
Macbrydei, *F. Muell. Fragm.* vi. 170.—Austral.
macrophylla, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 67.—Bras.
Massoni, *Sw. Prod. Veg. Ind. Occ.* 82.—Ind. occ.
microcarpa, Planch. ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 69 = cuneifolia.
monosperma, Benth. l. c. 70 = ochrocarpa.
monosperma, *Vell. Fl. Flum.* 225; v. t. 100.—Bras.
multiflora, *Karst. Fl. Columb.* ii. 31. t. 116.—N. Granat.
nitida, *G. Don, Gen. Syst.* i. 555.—Guian.
obtusa, *Planch. ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 68.—Bras.
ochrocarpa, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* xii. (1882) 329.—Bras.
oppositifolia, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 70.—Bras.
paniculata, *Spruce, ex Benth. l. c.* 66.—Bras.
paradisearum, *F. Muell. Papuan Pl.* i. 84.—N. Guin.
parviflora, *Planch. ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 68.—Guian.
Plumerii, Aubl. Pl. Gui. i. 536 = dentata.
Plumieri, G. Don, Gen. Syst. i. 555 = praec.
pubescens, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 69.—Bras.
pubescens, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* xii. (1882) 329.—Bras.
pubiflora, *Planch. & Linden, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 67.—N. Granat.
pulverulenta, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 329 = monosperma.
Purdiaei, *Griseb. ex Benth. & Hook. f. Gen.* ii. 653, in obs.—Ind. occ.
quadrivalvis, *Seem. Bot. Voy. Herald*, 85. t. 15.—Am. centr.; Bras.
riparia, *Planch. ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 69.—Bras.
rufa, *Planch. ex Benth. l. c.* 68.—Guian.
Schomburgkii, *Spruce, ex Benth. l. c.* 66.—Guian.
sinemariensis, *Aubl. Pl. Gui.* i. 534. t. 212.—Guian.; Ind. occ.
stipitata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 68.—Venezuel.
synandra, *Spruce, ex Benth. l. c.* 66.—Bras.
Woollsii, *F. Muell. Fragm.* vi. 171.—Austral.

SLOANIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 175 = praec.

SLOETIA, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 291, nomen; Kurz, in Journ. Linn. Soc. viii. (1864) 167. t. 13. *URTICACEAE*, Benth. & Hook. f. iii. 365.
eusideroxylon, Kurz, in Journ. Linn. Soc. viii. (1864) 167 = S. Sideroxylon.
Sideroxylon, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 291, nomen; *Kurz, l. c.*—Malaya.

SLOTERIA, Auct. ex Steud. Nom. ed. II. ii. 597 (1841). sphalm. = **Slateria**, Desv. (Haemod.).

SMEATHMANNIA, [Soland.] R. Br. in Tuckey, Congo, 439 (1818); et in Trans. Linn. Soc. xiii. (1821) 220. *PASSIFLOREAE*, Benth. & Hook. f. i. 812.
BUELOWIA, Schum. & Thonn. Beskr. Guin. Pl. 246 (1827).
decandra, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 304.—Afr. trop.
emarginata, *Lem. Jard. Fleur.* i. (1851) *Misc.* 71.—Afr. trop.
illustris, *Endl. ex Walp. Rep.* ii. 218.—Afr. trop.
laevigata, *Soland. ex R. Br. in Trans. Linn. Soc.* xiii. (1821) 221.—Afr. trop.
media, R. Br. l. c. = pubescens.
pubescens, *Soland. ex R. Br. l. c.*—Afr. trop.
rosea, *Lem. Jard. Fleur.* i. (1851) *Misc.* 71.—Afr. trop.

SMEGMADERMOS, Ruiz & Pav. Prod. 144. t. 31 (1794) = **Quillaja**, Molina (Rosac.).
emarginata, Ruiz & Pav. Syst. Veg. 288 = Q. Saponaria.

SMEGMARIA, Willd. Sp. Pl. iv. 1123 (1806) = SMEGMADERMOS, Ruiz & Pav. = **Quillaja**, Molina (Rosac.).
enarginata, Willd. l. c. = Q. Saponaria.

SMEGMATHAMNIUM, Fenzl, ex Reichb. Ic. Fl. Germ. vi. t. 244 (1844) = **Saponaria**, Linn. (Caryoph.).
bellidifolium, Reichb. l. c. = Sap. bellidifolia.
luteum, Reichb. l. c. = Sap. lutea.

SMELOPHYLLUM, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 330. *SAPINDACEAE*.
capense, *Radlk. l. c.* 331.—Afr. austr.

SMELOWSKIA, C. A. Mey. in Ledeb. Fl. Alt. iii. 165 (1831). *CRUCIFERAE*, Benth. & Hook. f. i. 79.
alpina, C. A. Mey. l. c. 170, in obs. = Hutchinsia alpina.
annua, *Rupr. Sert. Tiansch.* 39.—Turkestan.
asplenifolia, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 285 = cinerea.
bifurcata, Turcz. l. c. (1838) 88; xv. (1842) 286 = cinerea.
brevicaulis, Steud. Nom. ed. II. ii. 597 = Hutchinsia alpina.
californica, A. Gray, in Proc. Am. Acad. vi. (1865) 520 = Sisymbrium incisum.
calycina, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 170.—As. et Am. arct.
cinerea, *G. A. Mey. l. c.* 171.—Sibir.
cordifolia, Walp. Rep. ii. 761 = Eutrema cordifolium.
flavissima, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 156.—Soongar.
Fremontii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 123. —Calif.
integrifolia, C. A. Mey. in Ledeb. Fl. Alt. iii. 168 = Eutrema alpestre.
parviflora, Walp. Rep. ii. 761 = Eutrema Edwardsii.

SMILACINA, Desf. in Ann. Mus. Par. ix. (1807) 51. t. 9. *LILIACEAE*, Benth. & Hook. f. iii. 770.
ASTERANTHEMUM, Kunth, Enum. Pl. v. 151 (1850).
JOCASTE, Kunth, l. c. 154 (1850).
MEDORA, Kunth, l. c. 155 (1850).
NEOLEXIS, Salisb. Gen. Pl. Fragm. 64 (1866).
POLYGONASTRUM, Moench, Meth. 637 (1794).
SIGILLARIA, Rafin. in Journ. Phys. lxxxix. (1819) 261.
STYRANDRA, Rafin. l. c. (1819) 102.
TOVARIA, Neck. Elem. ii. 190 (1790), nomen prius.
alpina, Royle, Illustr. Bot. Himal. 380 = Clintonia alpina.
amoena, H. Wendl. in Otto & Dietr. Allg. Gartenz. xviii. (1850) 137 = paniculata.
amplexicaulis, Nutt. in Journ. Acad. Philad. vii. (1834) 58 = racemosa.
anomala, C. Sm. in Tuckey, Congo, 249.—Cf. Steud. Nom. ed. II. ii. 598, nomen (Quid?).—Ins. Cap. Virid.
Bertolonii, Kunth, Enum. Pl. v. 151 = flexuosa.
bifolia, Schult. f. Syst. vii. 307 = Maianthemum Convallaria.
bootanensis, Griff. Ic. Pl. Asiat. t. 279 = fusca.
borealis, Ker-Gawl. in Bot. Mag. tt. 1155, 1403 = Clintonia umbellata, borealis.
canadensis, Pursh, Fl. Am. Sept. i. 233 = Maianthemum Convallaria.
ciliata, Desf. in Ann. Mus. Par. ix. (1807) 53. t. 9 = racemosa.
cordifolia, Becker, Fl. Frankf. i. 136 = Maianthemum Convallaria.
dahurica, *Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 38; *et in Bull. Soc. Nat. Mosc.* (1838) 102; xxvii. (1854) II. 110.—Dahur.; Japon.
dilatata, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1875) 563 = Maianthemum Convallaria.
divaricata, Wall. Cat. n. 4202 = fusca.
Eschscholtziana, Andrz. & Bess. ex Baker, in Journ. Linn. Soc. xiv. (1875) 564 = Maianthemum Convallaria.
flexicaulis, Wender. in Schrift. Marb. ii. (1831) 249 = ciliata?
flexuosa, *Bertol. in Nov. Comm. Acad. Bonon.* iv. (1840) 411. t. 39.—Guatemala.
flexuosa, Hook. Ic. Pl. t. 529 = praec.?
Forskäliana, *Schult. f. Syst.* vii. 304.—Arab.
fusca, *Wall. Pl. As. Rar.* iii. 37. t. 257.—Reg. Himal.

SMILACINA:—
hirta, Maxim. Prim. Fl. Amur. 276 = japonica.
japonica, *A. Gray, in Perry, Jap. Exped.* ii. 321.—Japon.
latifolia, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1875) 570 = racemosa.
laxiflora, *Hemsl. Biol. Centr. Am. Bot.* iii. 368.—Guatemala.
macrophylla, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 387 = scilloidea.
nervulosa, *Hemsl. Biol. Centr. Am. Bot.* iii. 368.—Mexic.
oleracea, *Hook. f. & Thoms. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 569.—Reg. Himal.
pallida, Royle, Illustr. Bot. Himal. 380 = purpurea.
paniculata, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 388.—Mexic.
purpurea, *Wall. Pl. As. Rar.* ii. 38. t. 144.—Reg. Himal.
racemosa, *Desf. in Ann. Mus. Par.* ix. (1807) 51.—Am. bor.
ramosa, *G. Don, ex Loud. Hort. Brit.* 131, nomen.—Sibir.
Salvini, *Hemsl. Biol. Centr. Am. Bot.* iii. 368.—Guatemala.
scilloidea, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 388.—Mexic.
sessilifolia, *Nutt. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 566.—Am. bor. occ.
stellata, *Desf. in Ann. Mus. Par.* ix. (1807) 52.—Am. bor.
streptopoides, Ledeb. Fl. Ross. iv. 128 = Streptopus roseus.
thyrsoidea, *Hemsl. Biol. Centr. Am. Bot.* iii. 368.—Mexic.
trifolia, *Desf. in Ann. Mus. Par.* ix. (1807) 52.—Sibir.; Am. arct.
umbellata, Desf. l. c. 53 = Clintonia umbellata.
uniflora, Menz. ex Hook. Fl. Bor. Am. ii. 175. t. 190 = Clintonia uniflora.
yesoënsis, Franch. & Sav. Enum. Pl. Jap. ii. 523 = dahurica.

SMILAX, [Tourn.] Linn. Syst. ed. I (1735).
LILIACEAE, Benth. & Hook. f. iii. 763.
COPROSMANTHUS, Kunth, Enum. Pl. v. 263 (1850).
NEMEXIA, Rafin. Neogenyt. 3 (1825).
PARILLAX, Rafin. Med. Fl. U. St. ii. 264 (1828).
PLEIOSMILAX, Seem. Journ. Bot. vi. (1868) 193. t. 8.
acanthophylla, Vell. Fl. Flum. x. t. 114 = lappacea.
aculeata, *Vell. l. c.* t. 110.—Bras.
acuminata, *Willd. Sp. Pl.* iv. 779.—Ind. or.
acutifolia, *Schlecht. in Linnaea*, xviii. (1844) 449.—Mexic.
aequatorialis, *A. DC. in DC. Monog. Phan.* i. 141.—Bras.
alba, *Bertol. Misc. Bot.* xvi. 24. t. 2.—Am.
alba, *Pursh, Fl. Am. Sept.* i. 250.—Am. bor.
Alpini, Willd. Enum. Hort. Berol. Suppl. 67 = S. Bona-nox.
anceps, *Willd. Sp. Pl.* iv. 782.—Ins. Mascar.
angustifolia, *A. DC. in DC. Monog. Phan.* i. 67.—Costa Rica.
aristolochiaefolia, *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
aristolochiaefolia, Willd. Sp. Pl. iv. 786 = hastata.
aspera, DC. Organogr. ii. 262 = rotundifolia, *Linn.*
aspera, *Linn. Sp. Pl.* 1028.—Europ.; Oriens; Ind. or.
aspericaulis, *Wall. Cat.* n. 5129.—Reg. Himal.; Malaya.
aspero-variabilis, Pers. Syn. ii. 618 = S. Bona-nox.
Assumptionis, *A. DC. in DC. Monog. Phan.* i. 132.—Am. trop.
auriculata, Chapm. Fl. S. U. St. 245 = Beyrichii.
auriculata, *Walt. Fl. Carol.* 245.—Am. bor.
australis, *R. Br. Prod.* 293.—Austral.
australis, A. Cunn. ex A. DC. in DC. Monog. Phan. i. 215 = Rhipogonum discolor.
Balbisiana, Griseb. Fl. Brit. W. Ind. 586 = celastroides.
Balbisiana, *Kunth, Enum. Pl.* v. 183.—Ind. occ.
balearica, Burnat & Barbey, Not. Voy. Bot. Baléar. 12, 61 = aspera.
barbata, *Wall. Cat.* n. 5125.—Malaya.
bauhinioides, *Kunth, Enum. Pl.* v. 243.—Cochinchina.
Benthamiana, *A. DC. in DC. Monog. Phan.* i. 114.—Am. trop.

SMILAX:—
bermudensis, Duham. Arb. ed. Nov. i. 241 = S. Bona-nox.
Berterii, *Spreng. Syst.* ii. 102.—Ins. S. Doming.
Beyrichii, *Kunth, Enum. Pl.* v. 207.—Am. bor.
biflora, *Siebold, ex Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 87.—Japon.
Blancoi, *Kunth, Enum. Pl.* v. 262.—Ins. Philipp.
Blumei, *A. DC. in DC. Monog. Phan.* i. 202.—Malaya.
Bona-nox, *Linn. Sp. Pl.* 1030.—Carolina; Georg.
borbonica, Hort. Petrop. ex A. DC. in DC. Monog. Phan. i. 169 = anceps.
borneensis, *A. DC. l. c.* 202.—Borneo.
Botterii, *A. DC. l. c.* 89.—Mexic.
bracteata, *Presl, Rel. Haenk.* i. 131.—Ins. Philipp.
brasiliensis, *Spreng. Syst.* ii. 100.—Bras.
caduca, Ell. Sketch, ii. 701 = Walteri.
caduca, Linn. Sp. Pl. 1030 = rotundifolia.
caduca, Lour. Fl. Cochinch. 622 = incerta.
calaris, A. DC. in DC. Monog. Phan. i. 132, sphalm. = scalaris.
californica, *A. Gray, in Bot. Calif.* ii. 186.—Calif.
calophylla, *Wall. Cat.* n. 5131.—Malaya.
calophylla, Wall. l. c., partim = singaporensis.
campestris, *Griseb. in Mart. Fl. Bras.* iii. I. 15.—Bras.
canariensis, H. C. Wats. in Godman, Azor. 227, partim = excelsa.
canariensis, *Willd. Sp. Pl.* iv. 784.—Ins. Canar.
Candelariae, *A. DC. in DC. Monog. Phan.* i. 70.—Costa Rica.
canellaefolia, *Mill. Gard. Dict.* ed. VIII. n. 10.—Jamaic.
capitata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 49 = aspera.
catalonica, Poir. Encyc. vi. 467 = aspera.
celastroides, *Kunth, Enum. Pl.* v. 184.—Jamaic.
celebica, *Blume, Enum. Pl. Jav.* i. 18.—Malaya.
cercidifolia, *Pursh, Fl. Am. Sept.* ii. 733.—Am. bor.
China, Duham. Arbr. ed. Nov. i. 239 = anceps.
China, S. T. Gmel. Reise, iii. 52. t. 6 = excelsa.
China, *Linn. Sp. Pl.* 1029.—China; Cochinch.; Japon.
China, Vell. Fl. Flum. x. tt. 105, 106 = fluminensis.
China, Walt. Fl. Carol. 245 = Walteri.
ciliata, Steud. ex A. DC. in DC. Monog. Phan. i. 74 = rotundifolia.
cinnamomea, *Desf. ex A. DC. l. c.* 146.—Guian.
cissoides, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. I. 8.—Bras.
cognata, *Kunth, Enum. Pl.* v. 175.—Mexic.
collina, *Kunth, l. c.* 261.—Ind. or.
columnifera, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 49 = macrophylla.
corbularia, *Kunth, Enum. Pl.* v. 262.—Cochinch.
corcovadensis, Kunth, l. c. 226 = nitida.
cordato-ovata, Balb. ex A. DC. in DC. Monog. Phan. i. 152 = populnea.
cordato-ovata, Mus. Par. ex A. DC. l. c. 122 = havanensis.
cordato-ovata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113.—Guian.
cordifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 778.—Mexic.; Ind. occ.
coriacea, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 120.—Porto Rico.
coriacea, *Spreng. Syst.* ii. 103.—Ins. S. Doming.
coriifolia, *A. DC. in DC. Monog. Phan.* i. 112.—Bras.
Costaricae, Vatke, in Linnaea, xl. (1876) 223 = mexicana.
cumanensis, Bert. ex A. DC. in DC. Monog. Phan. i. 209 = coriacea, *Spreng.*
cumanensis, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 783.—Am. trop.
cumanensis, Schlecht. & Cham. in Linnaea, vi. (1831) 48 = mexicana.
cumanensis, Spreng. Syst. ii. 102 = celastroides.
cuspidata, *Duham. Arb.* ed. Nov. i. 242.—Guian.
Davidiana, *A. DC. in DC. Monog. Phan.* i. 104.—China.
decipiens, *Spreng. Pugill.* ii. 91.—Ind. or.
densiflora, *A. DC. in DC. Monog. Phan.* i. 88.—Mexic.
dentata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 774 = havanensis.
discolor, *Schlecht. in Linnaea*, xviii. (1844) 454.—Mexic.
divaricata, Blanco, Fl. Filip. ed. I. 795 = Dioscorea myriantha.

SMILAX:—
divaricata, *Soland. ex H. C. Wats. in Hook. Lond. Journ. Bot.* iii. (1844) 608.—Ins. Azor.
domingensis, Hohen. ex A. DC. in DC. Monog. Phan. i. 89 = Botterii.
domingensis, Schlecht. & Cham. in Linnaea, vi. (1831) 48 = Schlechtendalii.
domingensis, Spreng. ex A. DC. in DC. Monog. Phan. i. 152 = populnea.
domingensis, *Willd. Sp. Pl.* iv. 783.—Ins. S. Doming.
dulcis, Desf. Tabl. ed. I. 218 = glycyphylla.
Ehrenbergiana, *Kunth, Enum. Pl.* v. 174.—Ins. S. Doming.
elastica, *Griseb. in Mart. Fl. Bras.* iii. I. 22.—Bras.
elegans, A. DC. in DC. Monog. Phan. i. 107 = parvifolia.
elegans, *Wall. Cat.* n. 5117.—Reg. Himal.
elliptica, R. Br. Prod. 293 = australis.
elliptica, Desv. in Ham. Prod. Pl. Ind. Occ. 58 = zeylanica.
Engelmanniana, Kunth, Enum. Pl. v. 221 = rotundifolia.
erythrocarpa, *Kunth, l. c.* 234.—Mexic.
eucalyptifolia, *Kunth, l. c.* 230.—Peruv.
excelsa, Duham. Arb. ed. Nov. i. 236 = mauritanica.
excelsa, *Linn. Sp. Pl.* 1029.—Reg. Caucas.; Oriens.
extensa, Wall. Cat. n. 5126 A = myosotiflora.
extensa, *Wall. l. c.* B.—Penang.
febrifuga, *Kunth, Enum. Pl.* v. 201.—Peruv.
ferox, Wall. Cat. n. 5119 = S. China.
ficifolia, *Griseb. in Mart. Fl. Bras.* iii. I. 21.—Bras.
fistulosa, *Blanco, Fl. Filip.* ed. I. 796.—Ins. Philipp
flavescens, Desv. in Ham. Prod. Pl. Ind. Occ. 58 = salutaris.
flexuosa, Hemsl. Biol. Centr. Am. Bot. iii. 367, sphalm. = Smilacina flexuosa.
floribunda, Desv. in Ham. Prod. Pl. Ind. Occ. 58 = cuspidata.
floribunda, *Kunth, Enum. Pl.* v. 229.—Peruv.
fluminensis, *Steud. Nom.* ed. II. ii. 598.—Bras.
fulgens, Wall. Cat. n. 5122 = aspera.
Gaudichaudiana, Kunth, Enum. Pl. v. 252 = Heterosmilax Gaudichaudiana.
gemina, *Vell. Fl. Flum.* x. t. 113.—Bras.
glabra, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 792.—Reg. Himal.; China.
glauca, Mart. Reise Bras. i. 283 = brasiliensis.
glauca, *Walt. Fl. Carol.* 245.—Am. bor.
glaucocarpos, *Schlecht. in Linnaea*, xviii. (1844) 450.—Mexic.
glaucophylla, Klotzsch, in Bot. Ergeb. Waldem. Reise, 45 = parvifolia.
globifera, *G. F. W. Mey. Prim. Fl. Esseq.* 281.—Guian.
globulifera, Splitg. ex A. DC. in DC. Monog. Phan. i. 119 = Hostmanniana.
glyciphylla, *Sm. in White's Voy.* 230. t. 24.—Austral.
glycyphylla, Hassk. in Flora, xxv. (1842) II. Beibl. 4 = leucophylla.
goyazana, *A. DC. in DC. Monog. Phan.* i. 112.—Bras.
Goudotiana, *A. DC. l. c.* 172.—Madag.
grandifolia, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 171 = hispida.
grandifolia, Poepp. ex A. DC. in DC. Monog. Phan. i. 142 = Poeppigii.
grandifolia, *Regel, Ind. Sem. Hort. Petrop.* (1856) 16; *et cf. Ann. Sc. Nat.* Sér. IV. vi. (1856) 73.—Bras.
grandifolia, Voigt, Hort. Suburb. Calc. 648 = macrophylla.
grandis, Wall. ex Voigt, l. c. 649 = macrophylla.
Griffithii, *A. DC. in DC. Monog. Phan.* i. 198.—Burma?
guianensis, Vitm. Summ. Veg. v. 422 = megalophylla.
hastata, *Jacq. Enum. Pl. Carib.* 33; *Select. Am.* 262.—Ins. S. Doming.
hastata, Willd. Sp. Pl. iv. 782 = S. Bona-nox.
havanensis, Griseb. Fl. Brit. W. Ind. 586 = ilicifolia.
havanensis, *Jacq. Enum. Pl. Carib.* 33; *Select. Am.* 262.—Cuba.
hawaiensis, Seem. Fl. Vit. 309 = sandwicensis.
hederaefolia, Beyr. ex Kunth, Enum. Pl. v. 209 = S. Bona-nox.
hederaefolia, Mill. Gard. Dict. ed. VIII. n. 12 = S. Pseudo-china.
Helferi, *A. DC. in DC. Monog. Phan.* i. 176.—Burma; Malaya.

SMILAX:—
herbacea, *Linn. Sp. Pl.* 1030.—Am. bor.; Japon.
herbacea, Thunb. Fl. Cap. ed. Schult. 341 = Telfaireana.
heterophylla, *Rafin. in Med. Repos. N. York*, v. (1808) 359; *et in Desv. Journ. Bot.* i. (1808) 226.—Am. bor.
higoënsis, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 88.—Japon.
Hilariana, *A. DC. in DC. Monog. Phan.* i. 131.—Bras.
hispida, *Muhl. Cat.* 92; ed. II. 97.—Am. bor.
Hohenackeri, Kunth, Enum. Pl. v. 240 = zeylanica.
hongkongensis, Seem. Bot. Voy. Herald, 420 = Heterosmilax Gaudichaudiana.
Hookeri, Kunth, Enum. Pl. v. 162 = glabra.
horrida, Desf. Tabl. ed. I. 21 = S. Bona-nox.
Hostmanniana, *Kunth, Enum. Pl.* v. 170.—Guian.
Houstoniana, Steud. Nom. ed. II. ii. 598 = spinosa.
humilis, Mill. Gard. Dict. ed. VIII. n. 11 = herbacea.
hypoglauca, *Benth. Fl. Hongk.* 369.—Ins. Hongkong.
ilicifolia, *Kunth, Enum. Pl.* v. 171.—Cuba.
incerta, *Kunth, l. c.* 263.—Cochinch.
indica, Burm. f. Fl. Ind. 213 (err. typ. 313) = zeylanica.
inermis, Walt. Fl. Carol. 244 = herbacea.
insignis, *Kunth, Enum. Pl.* v. 200.—Peruv.
invenusta, *Kunth, l. c.* 234.—Mexic.
irrorata, Benth. ex A. DC. in DC. Monog. Phan. i. 114 = Benthamiana.
irrorata, Hohen. ex A. DC. l. c. 156 = Schomburgkiana.
irrorata, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. 1. 10.—Bras.
Jacquini, Kunth, Enum. Pl. v. 254 = hastata.
jalapensis, *Schlecht. in Linnaea*, xviii. (1844) 451.—Mexic.
Japicanga, *Griseb. in Mart. Fl. Bras.* iii. 1. 6.—Bras.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1857) 412 = S. China.
javensis, *A. DC. in DC. Monog. Phan.* i. 175.—Java.
Klotzschii, *Kunth, Enum. Pl.* v. 245.—Java.
Korthalsii, *A. DC. in DC. Monog. Phan.* i. 48.—Borneo.
Kraussiana, *Meissn. in Flora*, xxviii. (1845) 312.—Afr. austr.
laevis, Gueldenst. ex Ledeb. Fl. Ross. iv. 129 = excelsa.
laevis, *Wall. Cat.* n. 5116.—Malaya; China.
lanceaefolia, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 792.—Reg. Himal.; China.
lanceolata, Burm. f. Fl. Ind. 213 (err. typ. 313) = glabra.
lanceolata, Engelm. ex A. DC. in DC. Monog. Phan. i 76 = rotundifolia, *Linn.*
lanceolata, *Linn. Sp. Pl.* 1031.—Am. bor.
lanceolata, Ruiz, ex A. DC. in DC. Monog. Phan. i. 147 = insignis.
lanceolata, Walt. Fl. Carol. 245 = hastata.
lappacea, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 777.—Venezuela.
larvata, *Griseb. in Mart. Fl. Bras.* iii. 1. 19.—Bras.
lasioneuron, Hook. Fl. Bor. Am. ii. 173. t. 187 = herbacea.
latifolia, Blanco, Fl. Filip. ed. II. 548 = vicaria.
latifolia, R. Br. Prod. 293 = australis.
latifolia, Hassk. in Flora, xxv. (1842) II. Beibl. 4 (nomen delendum).—Java.
latifolia, *Soland. ex Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 12.—Ins. Mader.
laurifolia, Drège, ex A. DC. in DC. Monog. Phan. i. 212 = Asparagus crispus.
laurifolia, Hohen. ex A. DC. l. c. 188 = zeylanica.
laurifolia, *Linn. Sp. Pl.* 1030.—Am. bor.
laurifolia, Poepp. ex A. DC. in DC. Monog. Phan. i. 99 = lanceolata.
laurifolia, Roxb. Hort. Beng. 72; Fl. Ind. iii. 793 = prolifera.
laurina, Kunth, Enum. Pl. v. 248 = prolifera.
Lessertiana, *A. DC. in DC. Monog. Phan.* i. 173.—Madag.
leucophylla, *Blume, Enum. Pl. Jav.* i. 18.—Malaya.
ligustrifolia, *A. DC. in DC. Monog. Phan.* i. 63.—N. Caled.
longifolia, Lem. Illustr. Hortic. xiv. (1867) t. 521 = salicifolia.
longifolia, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = papyracea.

SMILAX:—
longifolia, Wats. Dendrol. ii. t. 110 = herbacea.
lutescens, Vell. Fl. Flum. x. t. 109 = ramiflora.
luzonensis, Presl, Rel. Haenk. i. 131 = Helferi?
Macabucha, Duch. ex Vidal & Garcia, Cat. Sem. Manila, 14 = seq.?
Macalucha, *Royle, Illustr. Bot. Himal.* 383, nomen.—Ins. Philipp.
macrocarpa, *Blume, Enum. Pl. Jav.* i. 19.—Java.
macrophylla, Griseb. Fl. Brit. W. Ind. 586, partim = populnea.
macrophylla, Poepp. ex A. DC. in DC. Monog. Phan. i. 142 = Poeppigii.
macrophylla, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 793.—Ind. or.
macrophylla, Willd. Sp. Pl. iv. 786 = megalophylla.
macropoda, *A. DC. in DC. Monog. Phan.* i. 115.—Bras.
maculata, Roxb. Hort. Beng. 72; Fl. Ind. iii. 796 = aspera.
marginulata, Mart. ex Griseb. in Mart. Fl. Bras. iii. 1. 16 = campestris.
mauritanica, Desf. Fl. Atlant. ii. 367 = aspera.
mauritanica, *Poir. Voy. Barb.* ii. 263.—Mauritan.
mauritanica, Webb & Berth. Phyt. Canar. iii. 322 = pendulina.
maypurensis, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 776.—Am. trop.
medica, Bott. ex A. DC. in DC. Monog. Phan. i. 89 = Botterii.
medica, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 390 = S. Bona-nox.
medica, Petz. Arbor. Muscav. 740 = tamnoides.
medica, *Schlecht. & Cham. in Linnaea*, vi. (1831) 47.—Mexic.
megacarpa, *A. DC. in DC. Monog. Phan.* i. 186.—Malaya.
megalophylla, *Duham. Arb.* ed. Nov. i. 244.—Ind. occ.
melastomifolia, *Sm. in Rees, Cycl.* xxxiii. [n. 5].—Ins. Sandvic.
menispermoidea, A. DC. in DC. Monog. Phan. i. 108 = elegans.
mexicana, *Griseb. ex Kunth, Enum. Pl.* v. 167.—Mexic.
micrantha, Blume, Enum. Pl. Jav. i. 18 = Heterosmilax japonica.
micropoda, A. DC. in DC. Monog. Phan. i. 58 = lanceaefolia.
Milleri, Steud. Nom. ed. II. ii. 599 = aristolochiaefolia.
Minarum, *A. DC. in DC. Monog. Phan.* i. 113.—Bras.
minutiflora, *A. DC. l. c.* 109.—Reg. Himal.
modesta, *A. DC. l. c.* 185.—Java.
mollis, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 785.—Mexic.
montana, Griseb. in Mart. Fl. Bras. iii. 1. 15 = campestris.
montevidensis, Hort. ex Steud. Nom. ed. II. ii. 599, nomen (Quid?).—Reg. Argent.
moranensis, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 389.—Mexic.
Morsaniana, Kunth, Enum. Pl. v. 241 = Kraussiana.
mossambicensis, Garcke, in Peters, Reise Mossamb. Bot. 518 = Kraussiana.
multiflora, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 390.—Mexic.
myosotiflora, *A. DC. in DC. Monog. Phan.* i. 65.—Java.
Myrtillus, *A. DC. l. c.* 106.—Reg. Himal.
Nageliana, *A. DC. l. c.* 184.—Java.
narcotica, Buch.-Ham. ex Wall. Cat. n. 5099 = Dioscorea daemona.
nigra, Willd. Sp. Pl. iv. 773 = aspera.
nilagirensis, Steud. ex A. DC. in DC. Monog. Phan. i. 167 = aspera.
nipponica, Miq. in Versl. en Med. Kon. Akad. Wetensch. II. ii. (1868) 87 = herbacea.
nitida, *Griseb. in Mart. Fl. Bras.* iii. 1. 9.—Bras.
obliqua, *Spreng. Pugill.* ii. 91.—Ind. or.
obliquata, *Duham. Arb.* ed. Nov. i. 242.—Peruv.
oblongata, Hort. Petrop. ex A. DC. in DC. Monog. Phan. i. 160 = grandifolia.
oblongata, *Sw. Prod. Veg. Ind. Occ.* 59.—Ind. occ.
oblongifolia, *Pohl, ex Griseb. in Mart. Fl. Bras.* iii. 1. 13.—Bras.
obtusa, *Benth. Bot. Voy. Sulph.* 175.—Am. trop.
ocreata, A. DC. in DC. Monog. Phan. i. 191 = Roxburghiana.
odontoloma, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. 1. 18.—Bras.

SMILAX:—
odoratissima, *Blume, Enum. Pl. Jav.* i. 19.—Java.
officinalis, Griseb. in Mart. Fl. Bras. iii. 1. 6 = procera.
officinalis, Herb. Lucaean. ex Kunth, Enum. Pl. v. 227 = salutaris.
officinalis, H. B. & K. Nov. Gen. et Sp. i. 271.—N. Granat.
officinalis, Poepp. ex A. DC. in DC. Monog. Phan. i. 150 = papyracea.
Oldhami, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 86.—Japon.
orbiculata, *Labill. Sert. Nov. Caled.* 18. t. 23.—N. Caled.
orbiculata, Vieill. ex A. DC. in DC. Monog. Phan. i. 65, partim = purpurata.
ornata, *Lem. Illlustr. Hortic.* xii. (1865) t. 439.—Mexic.
orthoptera, A. DC. in DC. Monog. Phan. i. 192 = Roxburghiana.
ovalifolia, A. DC. l. c. 199 = prolifera.
ovalifolia, Roxb. Hort. Beng. 72; Fl. Ind. iii. 794 = macrophylla.
ovata, Duham. Arb. ed. Nov. i. 242 = havanensis.
ovata, Ell. Sketch, ii. 698 = Beyrichii.
ovata, *Pursh, Fl. Am. Sept.* i. 249.—Am. bor.
oxyphylla, Kunth, Enum. Pl. v. 239 = laevis.
oxyphylla, *Wall. Cat.* n. 5128.—Reg. Himal.
pallescens, *A. DC. in DC. Monog. Phan.* i. 198.—Reg. Himal.
pandurata, Pursh, Fl. Am. Sept. i. 251 = S. Bona-nox.
paniculata, Herb. Desv. ex A. DC. in DC. Monog. Phan. i. 156 = Schomburgkiana.
papyracea, *Duham. Arb.* ed. Nov. i. 242.—Guian.
papyracea, Spruce, ex A. DC. in DC. Monog. Phan. i. 153 = Spruceana.
parviflora, *Wall. Cat.* n. 5118.—Reg. Himal.
pedunclaris, Muhl. ex Willd. Sp. Pl. iv. 786 = herbacea.
peguana, *A. DC. in DC. Monog. Phan.* i. 62.—Burma.
pekingensis, *A. DC. l. c.* 108.—China.
pendulina, Lowe, in Trans. Cambr. Phil. Soc. iv. 1. (1833) 12 = latifolia.
perfoliata, Blume, Enum. Pl. Jav. i. 18 = Blumei.
perfoliata, *Lour. Fl. Cochinch.* 622.—Cochinch.
phyllobola, *Mart. ex Griseb. in Mart. Fl. Bras.* iii. 1. 21, partim.—Bras.
phyllobola, Mart. ex Griseb. l. c., partim = Riedeliana, Selloana.
picta, Hort. ex C. Koch, Dendrol. ii. II. 337 = aspera.
pirarensis, *Kunth & Rich. Schomb. ex Kunth, Enum. Pl.* v. 165.—Guian.
platycentron, Schlecht. in Linnaea, xviiii. (1844) 452 = S. Bona-nox.
plurifurcata, *A. DC. in DC. Monog. Phan.* i. 65.—N. Caled.
Poeppigii, *Kunth, Enum. Pl.* v. 192.—Am. trop.
Pohliana, Kunth, l. c. 228 = procera.
Poiretii, Kunth, l. c. 261 = havanensis.
polyacantha, *Wall. Cat.* n. 5127.—Ins. Penang.
polyantha, *Griseb. in Mart. Fl. Bras.* iii. 1. 8.—Bras.
populnea, *Kunth, Enum. Pl.* v. 192.—Ins. S. Doming.
procera, *Griseb. in Mart. Fl. Bras.* iii. 1. 6.—Bras.
prolifera, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 795.—Ind. or.
prolifera, Wall. Cat. n. 5124 = Roxburghiana.
pruinosa, *Griseb. in Mart. Fl. Bras.* iii. 1. 14.—Bras.
Pseudo-china, Bert. ex Kunth, Enum. Pl. v. 183 = Balbisiana.
Pseudo-china, Blanco, Fl. Filip. ed. I. 795 = Blancoi.
Pseudo-china, Descourt, Fl. Antill. vii. 47. t. 462 (Quid?).—Ind. occ.
Pseudo-china, Herb. Madr. ex Wall. Cat. n. 5130 = zeylanica.
Pseudo-china, *Linn. Sp. Pl.* 1031.—Am. bor.; Ind. occ.
Pseudo-china, Lour. Fl. Cochinch. 623 = corbularia.
Pseudo-china, Mart. & Gal. ex A. DC. in DC. Monog. Phan. i. 87 = moranensis.
Pseudo-china, Roxb. Hort. Beng. 72; Fl. Ind. iii. 792 = collina.
Pseudo-china, Schiede, ex A. DC. in DC. Monog. Phan. i. 84 = cordifolia.
Pseudo-china, Sieber, ex A. DC. l. c. 170 = Telfaireana.
Pseudo-sarsa, Vis. Sem. Hort. Pat. (1839).—Cf. Linnaea, xiv. (1840) Litt. 138 = S. Bona-nox.
Pseudo-syphilitica, *Kunth, Enum. Pl.* v. 188.—Bras.
pteropus, Miq. Journ. Bot. Néerl. i. (1861) 89 = S. China.

SMILAX:—
pubens, Willd. Enum. Hort. Berol. Suppl. 67 = S. Bona-nox.
pubera, Michx. Fl. Bor. Am. ii. 238 = pumila.
puberula, Kunth, Enum. Pl. v. 193 = pumila.
pulverulenta, Michx. Fl. Bor. Am. ii. 238 = herbacea.
pumila, *Walt. Fl. Carol.* 244.—Am. bor.
Purampui, Ruiz, ex A. DC. in DC. Monog. Phan. i. 159 = febrifuga.
purpurata, *Forst. f. Prod.* 70.—N. Caled.
Purtrampui, Ruiz, ex A. DC. in DC. Monog. Phan. i. 97 = eucalyptifolia.
quadrangularis, Muhl. ex Willd. Sp. Pl. iv. 775 = rotundifolia, *Linn.*
quadrata, *A. DC. in DC. Monog. Phan.* i. 183.—Reg. Himal.; Burma.
quinquenervia, *Vell. Fl. Flum.* x. t. 108.—Bras.
ramiflora, *Griseb. in Mart. Fl. Bras.* iii. 1. 7.—Bras.
reticulata, Desv. in Ham. Prod. Pl. Ind. Occ. 58 = lanceolata.
retusa, Roxb. Hort. Beng. 72; Fl. Ind. iii. 793 = macrophylla.
Riedeliana, *A. DC. in DC. Monog. Phan.* i. 141.—Bras.
rigida, Russ. ex Steud. Nom. ed. I. 788 = aspera.
rigida, *Wall. Cat.* n. 5120.—Reg. Himal.
riparia, *A. DC. in DC. Monog. Phan.* i. 55.—China.
Ripogonum, Forst. f. Prod. 70 = Rhipogonum scandens.
robusta, *Griseb. in Mart. Fl. Bras.* iii. 1. 12.—Bras.
rotundifolia, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 120.—Porto Rico.
rotundifolia, *Linn. Sp. Pl.* 1030.—Am. bor.
Roxburghiana, *Wall. Cat.* n. 5115.—Reg. Himal.
Roxburghiana, Wall. l. c., partim = aspericaulis.
Roxburghii, Kunth, Enum. Pl. v. 852 = macrophylla.
rubens, *Wats. Dendrol.* t. 108.—Am. bor.
rubens, Willd. ex A. DC. in DC. Monog. Phan. i. 78 = S. Bona-nox.
rubiginosa, Griseb. in Mart. Fl. Bras. iii. 1. 14 = campestris.
rubra, Link, in Buch, Beschr. Canar. Ins. 140 = canariensis.
rubra, Willd. Enum. Hort. Berol. 1015 = Tamus communis.
rufescens, *Griseb. in Mart. Fl. Bras.* iii. 1. 9.—Bras.
Ruiziana, *Kunth, Enum. Pl.* v. 200.—Peruv.
sagittaefolia, Lodd. Bot. Cab. t. 1799 = aspera.
sagittata, Desv. in Ham. Prod. Pl. Ind. Occ. 58 = aspera.
salicifolia, *Griseb. in Mart. Fl. Bras.* iii. 1. 19.—Bras.
salutaris, *Kunth, Enum. Pl.* v. 227.—Bras.
sandwicensis, *Kunth, l. c.* 253.—Ins. Sandvic.
santaremensis, *A. DC. in DC. Monog. Phan.* i. 115.—Bras.
Sarsaparilla, *Linn. Sp. Pl.* 1029, partim (sp. dub.).—Am. bor.
Sarsaparilla, Linn. l. c., partim = glauca.
scabriuscula, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 783.—Am. austr.
scalaris, Griseb. in Mart. Fl. Bras. iii. 1. 16 = campestris.
Schiedeana, Kunth, Enum. Pl. v. 236 = cordifolia.
Schlechtendalii, *Kunth, l. c.* 224.—Mexic.
Schomburgkiana, *Kunth, l. c.* 187.—Guian.
Sebeana, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 149 = S. China.
Selloana, *A. DC. in DC. Monog. Phan.* i. 154.—Bras.
semiamplexicaulis, Boj. Hort. Maurit. 353 = Telfaireana.
senticosa, Kunth, Enum. Pl. v. 209 = S. Bona-nox.
setosa, Miq. Fl. Ind. Bat. Suppl. 610 = barbata.
Shuttleworthii, *Hort.—Cf. Gard. Chron.* (1876) 1. 35.—Hab.?
Sieboldi, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 87.—Japon.
Sieboldii, Hort. Bog. ex Hassk. Cat. Hort. Bog. Alt. 32, nomen (Quid?).—Japon.
singaporensis, *A. DC. in DC. Monog. Phan.* i. 177.—Singapore.
siphilitica, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 780.—Am. austr.
solanifolia, *A. DC. in DC. Monog. Phan.* i. 161.—Am. trop.
spicata, *Vell. Fl. Flum.* x. t. 111.—Bras.
spicata, Vell. l. c. t. 112 = Dioscorea marginata.

SMILAX:—
spinescens, Miq. in Linnaea, xviii. (1844) 83 = australis.
spinosa, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
spinosa, Poir. Encyc. vi. 467 = havanensis.
spinulosa, Sm. in Rees, Cycl. xxxiii. [n. 9] = glauca.
Sprengelii, Kunth, Enum. Pl. v. 207 = rotundifolia.
Spruceana, *A. DC. in DC. Monog. Phan.* i. 153.—Bras.
staminea, *Griseb. in Mart. Fl. Bras.* iii. 1. 11.—Bras.
stans, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 170.—Japon.
stenopetala, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–59) 412.—Japon.
stenophylla, *A. DC. in DC. Monog. Phan.* i. 130.—Bras.
stipulacea, *Boj. Hort. Maurit.* 352.—Hab.?
subaculeata, *Spreng. Syst.* ii. 102.—Am. trop.
subinermis, Presl, Rel. Haenk. i. 132 = eucalyptifolia.
subpubescens, *A. DC. in DC. Monog. Phan.* i. 69.—Mexic.
subsessiliflora, Duham. Arb. ed. Nov. i. 243 = salutaris.
surinamensis, Miq. in Linnaea, xvii. (1843) 63 = cumanensis.
sylvatica, Kunth, Enum. Pl. v. 234 = cordifolia.
sylvatica, Kunth, l. c., partim = moranensis.
syphilitica, Auct. vide siphilitica.
syphilitica, Griseb. in Mart. Fl. Bras. iii. 1. 20 = S. Pseudo-syphilitica.
syphilitica, Hort. Berol. ex Kunth, Enum. Pl. v. 190 = Sarcococca pruniformis?
syphilitica, Mart. Herb. ex A. DC. in DC. Monog. Phan. i. 134 = irrorata.
syphilitica, Mart. Reise Bras. iii. 1280 = papyracea.
syringoides, Griseb. in Mart. Fl. Bras. iii. 1. 11 = fluminensis.
Talbotiana, *A. DC. in DC. Monog Phan.* i. 152.—Guian.
tamnifolia, Michx. Fl. Bor. Am. ii. 238 = tamnoides.
tamnoides, A. Gray, Man. Bot. N. U. St. ed. I. 485 = S. Bona-nox.
tamnoides, *Linn. Sp. Pl.* 1030.—Am. bor.
Telfaireana, *Wall. Cat.* n. 5121 A.—Ins. Maurit.
Telfaireana, Wall. l. c. B = Heterosmilax Gaudichaudiana.
tetragona, Linn. f. Suppl. 427 = aspera.
tetragona, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 390 = rotundifolia.
tetragona, Seub. Fl. Azor. 24 = excelsa.
Thomsoniana, A. DC. in DC. Monog. Phan. i. 104 = S. China.
tijucensis, *A. DC. l. c.* 94.—Bras.
timorensis, *A. DC. l. c.* 189.—Ins. Timor.
timorensis, Zipp. ex Span. in Linnaea, xv. (1841) 479, nomen (Quid?).—Ins. Timor.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* i. 272.—N. Granat.
trifurcata, *Seem. Fl. Vit.* 309.—Ins. Fiji.
trinervula, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 87.—Japon.
triplinervia, Griseb. Cat. Pl. Cub. 251 = mollis.
triplinervia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 784.—Venezuela.
tuberculata, *Presl, Rel. Haenk.* i. 132.—Ecuador.
umbellata, Herb. Madr. ex Wall. Cat. n. 5130 = zeylanica.
umbellata, *Vell. Fl. Flum.* x. t. 107.—Bras.
umbellifera, Herb. Heyne, ex Wall. Cat. n. 5124 = prolifera.
undulata, *Pohl, ex A. DC. in DC. Monog. Phan.* i. 135.—Bras.
vaginata, *Decne. in Jacquem. Voy. Bot.* 169. t. 169.—Reg. Himal.
variabilis, Pers. Syn. ii. 618 = aspera.
variegata, Walt. Fl. Carol. 244 = S. Bona-nox.
verrucosa, *Griseb. in Mart. Fl. Bras.* iii. 1. 17.—Bras.
vicaria, *Kunth, Enum. Pl.* v. 262.—Ins. Philipp.
Villandia, *Buch.-Ham. ex Wall. Cat.* n. 5123 (sp. dub.).—Ind. or.
viminea, *Griseb. in Mart. Fl. Bras.* iii. 1. 13.—Bras.
virginiana, *Mill. Gard. Dict.* ed. VIII. n. 9.—Jamaic.
virosa, Buch.-Ham. ex Wall. Cat. n. 5099 = Dioscorea daemona.
viscifolia, *Duham. Arb.* ed. Nov. i. 243.—Ins. S. Doming.

SMILAX:—
vitiensis, *A. DC. in DC. Monog. Phan.* i. 204.—Ins Fiji.
Wagneriana, *A. DC. l. c.* 143.—Panama.
Wallichii, Kunth, Enum. Pl. v. 246 = Roxburghiana?
Wallichii, Steud. Nom. ed. II. ii. 599 = rigida.
Walteri, *Pursh, Fl. Am. Sept.* i. 249.—Am. bor.
Watsoni, Sweet, Hort. Brit. ed. I. 414 = herbacea.
Wightii, *A. DC. in DC. Monog. Phan.* i. 174.—Ind. or.
xalapensis, Kunth, Enum. Pl. v. 258 = jalapensis.
zeylanica, Blume, Enum. Pl. Jav. i. 17 = Klotzschii.
zeylanica, *Linn. Sp. Pl.* 1029.—Ind. or.; Malaya.
zeylanica, Wight, Ic. t. 2057–58 = Wightii.
Zollingeri, Kunth, Enum. Pl. v. 241 = zeylanica.

SMIRNOWIA, Bunge, in Act. Hort. Petrop. iv. (1876) 339. *LEGUMINOSAE.*
turkestana, *Bunge, l. c.*—Turkestan.

SMITHIA, Ait. Hort. Kew. ed. I. iii. 496. t. 13 (1789). *LEGUMINOSAE*, Benth. & Hook. f. i. 516.
DAMAPANA, Adans. Fam. ii. 323 (1763).
KOTSCHYA, Endl. Stirp. Nov. Dec. 4 (1839).
PETAGNANA, J. F. Gmel. Syst. 1078 (1791).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 203 = sensitiva.
aeschynomenoides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 153.—Afr. trop.
aspera, Roxb. Hort. Beng. 56; Fl. Ind. iii. 343 = Aeschynomene indica.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
bigemina, *Dalz. in Hook. Kew Journ.* iii. (1851) 208.—Ind. or.
bigeminata, Blanco, Fl. Filip. ed. II. 395 = Zornia diphylla.
blanda, Hohen. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 150 = salsuginea.
blanda, *Wall. Cat.* n. 5669.—Ind. or.
capitata, *Dals. in Hook. Kew Journ.* iii. (1851) 208.—Ind. or.
capitata, Desv. Journ. Bot. i. (1813) 121 = geminiflora.
capitulifera, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 152.—Afr. trop.
Chamaecrista, *Benth. in Miq. Pl. Jungh.* 211.—Madag.
ciliata, *Royle, Illustr. Bot. Himal.* 201. t. 35. f. 2.—Reg. Himal.
coerulescens, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, iii. (1846) 76.—Java.
conferta, Sm. in Rees, Cycl. xxxiii. n. 2 = geminiflora.
cristata, Wight, ex Wight & Arn. Prod. 218 = Geissaspis cristata.
dichotoma, Dalz. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 150 = salsuginea.
disperma, Wight, ex Steud. Nom. ed. II. ii. 599 = Geissaspis cristata.
domingensis, Balb. ex DC. Prod. ii. 321 = Aeschynomene pleuronervia.
elaphroxylon, Baill. in Bull. Soc. Linn. Par. i. (1883) 404 = Herminiera elaphroxylon.
geminiflora, *Roth, Nov. Pl. Sp.* 352.—Ind. or.; Malaya; Austral.
gracilis, *Benth. in Linnaea*, xxiv. (1851) 642.—Ind. or.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 404.—Madag.
grandis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 151.—Reg. Himal.
hirsuta, Dalz. in Hook. Kew Journ. iii. (1851) 135 = blanda.
hispidissima, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 76 = geminiflora.
japonica, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 389.—Japon.
javanica, *Benth. in Miq. Pl. Jungh.* 211.—Java.
Kotschyi, *Benth. l. c.*—Afr trop.
laxiflora, Benth. l. c. = purpurea.
martinicensis, *Spreng. Syst.* iii. 323.—Ins. Martinic.
nodulosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 153.—Afr. trop.
paniculata, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 330 = blanda.
pumila, Royle, ex Wight & Arn. Prod. 220 = ciliata.
purpurea, Hook. Bot. Mag. t. 4283.—Ind. or.

SMITHIA :—
pycnantha, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 150.— Ind. or.
racemosa, Heyne, in Wall. Cat. n. 5670 = blanda.
salsuginea, *Hance, in Journ. Bot.* vii. (1869) 164.— China.
sensitiva, *Ait. Hort. Kew.* ed. I. iii. 496.—As. et Afr. trop.
sensitiva, Wight & Arn. Prod. 220 = geminiflora.
sensitiva, Zoll. & Mor. Syst. Verz. Zoll. 6 = javanica.
setulosa, *Dalz. in Hook. Kew Journ.* iii. (1851) 208.— Ind. or.
spicata, Spreng. Neue Entdeck. ii. 160 (nom. delend.). —Afr. trop.
strigosa, *Benth. in Miq. Pl. Jungh.* 211.—Madag.; Afr. trop.
strobilantha, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 154.—Afr. trop.

SMITHIA, J. F. Gmel. Syst. 388 (1791) = **Humbertia**, Lam. (Convolv.).
Thouinia, J. F. Gmel. l. c. 388 = H. madagascariensis.

SMITHIA, Scop. Introd. 322 (1777) = **Clusia**, Linn. (Guttif.).

SMODINGIUM, E. Mey. in Drège, Zwei Pfl. Docum. 222 (1843), nomen; et ex Harv. & Sond. Fl. Cap. i. 523 (1859–60). *ANACARDIACEAE*, Benth. & Hook. f. i. 422.
Andrieuxii, Baill. Adansonia, xi. (1874) 182 = Pseudosmodingium Andrieuxii.
argutum, *E. Mey. in Drège, Zwei Pfl. Docum.* 222, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 523.— Afr. austr.
Virletii, Baill. Adansonia, xi. (1874) 182 = Pseudosmodingium Virletii.

SMYRNIOPSIS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 72 = **Smyrnium**, Tourn. (Umbell.).
Aucheri, Boiss. l. c.—As. Min.; Persia.
cachroides, Boiss. Fl. Orient. ii. 928.—Syria; Mesopot.
syriaca, Boiss. Diagn. Ser. I. x. 51.—Syria.

SMYRNIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 81 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 885.
ANOSMIA, Bernh. in Linnaea, vii. (1832) 608.
SMYRNIOPSIS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 72.
acuminatum, Sm. in Rees, Cycl. xxxiii. n. 8 = Thaspium aureum.
aegopodioides, *H. B. & K. Nov. Gen. et Sp.* v. 16.— Mexic.
aegyptiacum, Linn. Cent. Pl. i. 10; Amoen. Acad. iv. 270 = connatum?
americanum, D. Dietr. Syn. Pl. ii. 989 = Eulophus americanus.
apiifolium, *Willd. Sp. Pl.* i. 1468.—Ins. Creta.
atropurpureum, Lam. Encyc. iii. 667 = Thaspium trifoliatum.
aureum, Linn. Sp. Pl. 262 = Thaspium aureum.
barbinode, Muhl. Cat. 32; et ex Pursh, Fl. Am. Sept. i. 196 = Thaspium barbinode.
cicutarium, Bieb. Fl. Taur. Cauc. i. 239 = Eleutherospermum grandifolium.
connatum, *Boiss. & Kotschy, in Unger & Kotschy, Ins. Cypern*, 309.—As. Min.; Syria.
cordatum, Walt. Fl. Carol. 114 = Thaspium trifoliatum.
cordifolium, *Boiss. Diagn.* Ser. I. vi. 64.—Persia; Assyr.
creticum, Mill. Gard. Dict. ed. VIII. n. 4 = apiifolium.
Dioscoridis, Spreng. Sp. Umbellif. 25 = perfoliatum.
Dodonaei, Spreng. l. c. 24 = rotundifolium.
heterophyllum, DC. Prod. iv. 110, sphalm. = Sison Amomum.
integerrimum, Linn. Sp. Pl. 263 = Pimpinella integerrima.
laserpitioides, Crantz, Cl. Umb. Emend. 73 = Prangos ferulacea.
laterale, Thunb. Prod. Pl. Cap. 51 = Apium graveolens.
lineare, Benth. Pl. Hartw. 83 = Eulophus peucedanoides.
luteum, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 166, nomen = Thaspium aureum.
maritimum, Salisb. Prod. 168 = S. Olusatrum.
Matthioli, Presl, Delic. Prag. 127 = rotundifolium.

SMYRNIUM:—
nodiflorum, All. Fl. Pedem. ii. 23. t. 72 = Trochiscanthes nodiflora.
nudicaule, Bieb. Fl. Taur. Cauc. i. 238 = Physospermum commutatum.
nudicaule, Pursh, Fl. Am. Sept. i. 196 = Peucedanum nudicaule.
Olusatrum, *Linn. Sp. Pl.* 262.—Europ.; Reg. Mediterr.
Orphanidis, *Boiss. Fl. Orient.* ii. 925.—Graecia.
perfoliatum, *Linn. Sp. Pl.* 262.—Europ. austr.; Reg. Cauc.
ramosum, Urv. in Mém. Soc. Linn. Par. i. (1822) 289 = rotundifolium.
rotundifolium, *Mill. Gard. Dict.* ed. VIII. n. 2.—Reg. Mediterr.
trifoliatum, Muhl. Cat. 31; Nutt. Gen. Am. i. 195 = Thaspium trifoliatum.
vulgare, S. F. Gray, Nat. Arr. Brit. Pl. ii. 525 = S. Olusatrum.

SMYTHEA, Seem. in Bonplandia, ix. (1861) 255; x. (1862) 35. *RHAMNEAE*, Benth. & Hook. f. i. 375.
calpicarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 301.—Burma.
novo-guineënsis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 14.—N. Guin.
pacifica, *Seem. in Bonplandia*, ix. (1861) 255; x. (1862) 35. t. 9.—Ins. Fiji.

SOALA, Blanco, Fl. Filip. ed. I. 437 (1837) = **Cyathocalyx**, Champ. (Anonac.).
litoralis, Blanco, l. c. = C. zeylanicus.

SOARESIA, Allem. in Rev. Braz. i. (1857) 210; et in Arch. Palestr. Sc. Rio Jan. (1858) 142 = **Clarisia**, Ruiz & Pav. (Urtic.).
nitida, Allem. ll. cc.—Bras.

SOARESIA, Sch. Bip. in Pollichia, xx–xxi. (1863) 376. *COMPOSITAE*, Benth. & Hook. f. ii. 236.
velutina, *Sch. Bip. l. c.* 377.—Bras.

SOBERBAEA, D. Dietr. Syn. Pl. ii. 1129 (1840), sphalm. = **Sowerbaea**, Sm. (Liliac.).

SOBISO, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
japonica, Rafin. l. c. = Salvia japonica.

SOBOLEWSKIA, Bieb. Fl. Taur. Cauc. iii. 421 (1819); et Cent. Pl. Rar. ii. t. 59 (1832). *CRUCIFERAE*, Benth. & Hook. f. i. 95.
MACROSPORUM, DC. Syst. ii. 575, in obs. (1821).
clavata, *Fenzl, in Tchihat. Asie Min. Bot.* i. 348.— As. Min.
lithophila, *Bieb. Fl. Taur. Cauc.* iii. 421; *et Cent. Pl. Rar.* ii. t. 59.—Tauria; Reg. Caucas.
lithophila, Fenzl, in Tchihat. Asie Min. Bot. i. 347 = clavata.
stylosa, Boiss. & Hohen. Diagn. Ser. I. viii. 41 = Physalidium stylosum.

SOBRALIA, Ruiz & Pav. Prod. 120. t. 26 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 590.
CYATHOGLOTTIS, Poepp. & Endl. Nov. Gen. et Sp. i. 55. t. 94 (1835).
amplexicaulis, Ruiz & Pav. Syst. Veg. 233 (al. gen.). —Peruv.
aurantiaca, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.?
biflora, Ruiz & Pav. Syst. Veg. 232 (al. gen.).— Peruv.
Bletiae, *Reichb. f. in Bot. Zeit.* x. (1852) 713.— Panama.
bletioides, Brongn. ex Decne. in Rev. Hortic. Sér. III. i. (1847) 421, cum tab. = Bletia hyacinthina.
candida, *Reichb. f. in Fl. des Serres*, viii. (1852–53) 247.—Peruv.
Caravata, Lindl. Gen. et Sp. Orch. 177 = Elleanthus Caravata.
Cattleya, *Reichb. f. in Gard. Chron.* (1877) i. 72.— N. Granat.
chlorantha, Hook. Bot. Mag. t. 4682 = macrophylla.
chrysantha, *Lindl. Fol. Orch. Sobr.* 3.—N. Granat.
chrysoleuca, *Reichb. f. Xenia Orch.* ii. 179.—Bolivia.
citrina, La Llave & Lex. Nov. Veg. Desc. fasc ii. (Orch. Opusc.) 21 = Cattleya citrina.

SOBRALIA:—
crocea, *Reichb. f. in Fl. des Serres*, viii. (1852–53) 247.—Peruv.
decora, *Batem. Orch. Mex. et Guat.* t. 26.—Guatemala.
dichotoma, *Ruiz & Pav. Syst. Veg.* 232.—Peruv.
D'Orbignyana, *Reichb. f. Xenia Orch.* ii. 179.— Bolivia.
Elisabethae, *Rob. Schomb. in Verh. Befoerd. Gartenb. Preuss.* xv. (1841) 137. tt. 1, 2.—Guian.
Fenzliana, *Reichb. f. in Bot. Zeit.* x. (1852) 714.— Nicaragua.
fimbriata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 54.— Peruv.
fragrans, *Lindl. in Gard. Chron.* (1853) 598.— N. Granat.
Galeottiana, *A. Rich. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic.
gloriosa, *Reichb. f. Xenia Orch.* ii. 178.—Ecuador.
Klotzscheana, *Reichb. f. in Linnaea*, xxii. (1849) 815. —Peruv.
labiata, *Warsz. & Reichb. f. in Bot. Zeit.* x. (1852) 714.—Panama.
lepida, *Reichb. f. Beitr. Orch. Centr. Am.* 68.—Costa Rica.
leucoxantha, *Reichb. f. l. c.*—Costa Rica.
Liliastrum, *Lindl. Gen. et Sp. Orch.* 177, 432.—Guian.; Bras.
Lindleyana, *Reichb. f. in Bot. Zeit.* x. (1852) 713. —Costa Rica.
macrantha, *Lindl. Sert. Orch.* sub t. 29.—Mexic.
macrophylla, *Reichb. f. in Bot. Zeit.* x. (1852) 713.— Panama.
Mandonii, *Reichb. f. Xenia Orch.* ii. 175.—Bolivia.
paludosa, *Linden, Cat.* n. 6 (1851) 8.—N. Granat.
paradisiaca, *Reichb. f. in Linnaea*, xxii. (1849) 816.— N. Granat.
porphyrocharis, *Reichb. f. Xenia Orch.* ii. 177.— Ecuador.
rigidissima, *Linden, ex Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.
Roezlii, *Reichb. f. Xenia Orch.* ii. 178.—N. Granat.
rosea, Lindl. Fol. Orch. Sobr. 2 = glauca.
rosea, *Poepp. & Endl. Nov. Gen. et Sp.* i. 54. t. 93.— Peruv.; N. Granat.
roseo-alba, *Reichb. f. Beitr. Orch. Centr. Am.* 7.—Am. centr.
Ruckeri, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 278.—N. Granat.
Ruckeri, Linden, ex Lindl. Fol. Orch. Sobr. 3 = rosea.
scopulorum, *Reichb. f. Xenia Orch.* ii. 176.—Bolivia.
sessilis, Hook. Bot. Mag. t. 4570 = decora.
sessilis, *Lindl. Bot. Reg.* (1841) *Misc.* 3. t. 17.— Guian.
setigera, *Poepp. & Endl. Nov. Gen. et Sp.* i. 54.— Peruv.
stenophylla, *Lindl. Fol. Orch. Sobr.* 2.—Guian.
suaveolens, *Reichb. f. in Gard. Chron.* (1878) i. 622. —Panama.
violacea, *Linden, ex Lindl. Orchid. Linden.* 26.— Venezuela.
Warszewiczi, *Reichb. f. in Bot. Zeit.* x. (1852) 714.— Panama.

SOBREYRA, Ruiz & Pav. Prod. 109. t. 23 (1794) = **Enhydra**, Lour. (Compos.).
oblonga, Ruiz & Pav. Syst. Veg. 197 = E. oblonga.
sessiliflora, Ruiz & Pav. l. c. = E. Sobreyra.

SOBRYA, Pers. Syn. ii. 473 (1807) = praec.

SOCOTORA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 77. *APOCYNACEAE*.
aphylla, *Balf. f. l. c.*—Ins. Socotr.

SOCRATEA, Karst. in Linnaea, xxviii. (1856) 263. *PALMAE*, Benth. & Hook. f. iii. 900.
durissima, *H. Wendl. in Bonplandia*, viii. (1860) 103. —Nicaragua.
elegans, *Karst. in Linnaea*, xxviii. (1856) 264.— N. Granat.
exorrhiza, *H. Wendl. in Bonplandia*, viii. (1860) 103. —Mexic.
fusca, Karst. Fl. Columb. i. 109. t. 54 = Dictyocaryum fuscum.
Orbignyana, *Karst. in Linnaea*, xxviii. (1856) 264.— N. Granat.
philonotia, Benth. & Hook. f. Gen. iii. 900 = exorrhiza.

SOCRATESIA, Klotzsch, in Linnaea, xxiv. (1851) 22 = **Cavendishia**, Lindl. (Vaccin.).
melastomoides, Klotzsch, l. c. = C. melastomoides.

SODA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 145 = **Salsola**, Linn. (Chenop.).
inermis, Fourr. l. c. = Salsola Soda.

SODADA, Forsk. Fl. Aegypt. Arab. 81 (1775) = **Capparis**, Tourn.
decidua, Forsk. l. c.; Delile, Fl. Egypt. 74. t. 26 = C. aphylla.

SODIROA, André, in Bull. Soc. Bot. Fr. xxiv. (1877) 167. *BROMELIACEAE*, Benth. & Hook. f. iii. 668.
caricifolia, *André, l. c.*—N. Granat.
graminifolia, *André, l. c.*—N. Granat.

SOEMMERINGIA, Mart. Diss. Soemmer. (1828) 27. *LEGUMINOSAE*, Benth. & Hook. f. i. 516.
psittacorhyncha, Webb, in Hook. Niger Fl. 123 = Geissaspis lupulina.
semperflorens, *Mart. Diss. Soemm.* (1828) 28.—Bras.

SOGALGINA, Cass. in Bull. Soc. Philom. (1818) 31 = **Tridax**, Linn. (Compos.).
balbisioides, Cass. in Dict. Sc. Nat. xlix. 398 = T. balbisioides.
dentata, Lilja, in Linnaea, xvii. (1843) 110 = T. trilobata?
trilobata, Cass. in Dict. Sc. Nat. xlix. 397 = T. trilobata.

SOGALIGNA, Steud. Nom. ed. I. 788 (1821), sphalm. = praec.

SOHREA, Steud. Nom. ed. I. 788 (1821), sphalm. = **Shorea**, Roxb. (Dipteroc.).

SOJA, Moench, Meth. 153 (1794) (Soia) = **Glycine**, Linn. (Legumin.).
angustifolia, Miq. Fl. Ind. Bat. i. 1. 223 = G. Soja.
hamata, Miq. l. c. 224 = G. pendula (in Addend.).
hispida, Moench, Meth. 153 = G. Soja.
japonica, Savi, in Pisa, Nuov. Giorn Litt. viii. (1824) 113; (Mem. Phas. ii. 16) = G. Soja.
javanica, R. Grah. in Wall. Cat. n. 5528 = G. javanica.
viridis, Savi, Cose Bot. 9.—Hab.?
Wightii, R. Grah. in Wall. Cat. n. 5530 = G. javanica.

SOKOLOFIA, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
rupestris, Rafin. l. c. 15 = Salix nigricans.

SOLAENACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 122 = **Ruellia**, Linn. (Acanth.).
longiflorus, Oerst. l. c. = R. longiflora.

SOLANANDRA, Pers. Syn. ii. 215 (1807), sphalm. = SOLENANDRIA, Beauv. = **Galax**, Linn. (Diapens.).

SOLANDRA, Linn. Syst. ed. X. 1269 (1759) = **Hydrocotyle**, Tourn. (Umbell.).
capensis, Linn. l. c. = H. Solandra.

SOLANDRA, Murr. in Comm. Gotting. vi. 1783–84 (1785), 20. t. 1 = **Hibiscus**, Linn. (Malvac.).
aculeata, Poir. Encyc. vii. 225 = H. ficulneus.
lobata, Murr. in Comm. Gotting. vi. 1783–84 (1785), 20 = H. Solandra.
spinosa, Poir. in Dict. Sc. Nat. xlix. 406 = H. ficulneus.
squamea, Poir. Encyc. vi. 225 = Lagunaria Patersonii.
ternata, Cav. Diss. v. 279. t. 136. f. 2 = H. ternatus.

SOLANDRA, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 300. t. 11. *SOLANACEAE*, Benth. & Hook. f. ii. 901.
SWARTZIA, J. F. Gmel. Syst. 360 (1791).
grandiflora, Salisb. in Trans. Linn. Soc. vi. (1802) 100 = hirsuta.
grandiflora, *Sw. in Vet. Akad. Handl. Stockh.* viii. (1787) 300. t. 11.—Am. trop.
guttata, *D. Don, in Bot. Reg.* t. 1551.—Mexic.
herbacea, Hort. ex Dun. in DC. Prod. xiii. 1. 544 = Datura ceratocaula.
hirsuta, *Dun. l. c.* 535.—Bras.

SOLANDRA:—
laevis, Hook. Bot. Mag. t. 4345 = longiflora.
longiflora, Tussac, Fl. Antill. ii. 49. t. 12.—Ind. occ.
longipes, Sendt. in Mart. Fl. Bras. x. 159 = Dyssochroma longipes.
macrantha, Dun. in DC. Prod. xiii. 1. 533 = grandiflora.
minor, *Griseb. Fl. Brit. W. Ind.* 433.—Ins. Dominic.
nitida, Zuccagni, in Roem. Collect. 128 = grandiflora.
oppositifolia, Moon, Cat. Pl. Ceyl. 15 = Fagraea ceilanica.
platantha, Walp. Ann. iii. 152 = Portlandia platantha.
scandens, Willd. ex Sendt. in Mart. Fl. Bras. x. 160 = grandiflora.
viridiflora, Sims, Bot. Mag. t. 1948 = Dyssochroma viridiflora.

SOLANOIDES, Tourn. ex Moench, Meth. 307 (1794) = **Rivina**, Plum. (Phytolac.).
laevis, Moench, l. c. = R. humilis.
pubescens, Moench, l. c. = R. humilis.
undulata, Moench, Meth. Suppl. 106 = R. humilis.

SOLANUM, [Tourn.] Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 888.
ANDROCERA, Nutt. Gen. Am. i. 129 (1818).
AQUARTIA, Jacq. Enum. Pl. Carib. 12 (1760).
CERANTHERA, Rafin. in Am. Monthly Mag. (1818) 176.
CLIOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 141.
DIAMONON, Rafin. Fl. Tellur. ii. 76 (1836).
DULCAMARA, Moench, Meth. 514 (1794).
NORMANIA, Lowe, Man. Fl. Mad. ii. 70 (1868).
NYCTERIUM, Vent. Jard. Malm. 85. t. 85 (1803).
PSEUDOCAPSICUM, Medic. Phil. Bot. i. 122 (1789).
PSOLANUM, Neck. Elem. ii. 60 (1790).
abyssinicum, Jacq. ex Vitm. Summa Pl. i. 492 = marginatum.
acanthifolium, Aubl. ex Dun. in DC. Prod. xiii. 1. 255 = stramonifolium.
acanthifolium, Hort. ex Dun. l. c. 215 = S. Juripeba.
acanthifolium, Hort. Par. ex Dun. l. c. 260 = torvum.
acanthifolium, Leman, ex Dun. l. c. 200 = jamaicense.
acanthifolium, Mill. Gard. Dict. ed. VIII. n. 14 = fuscatum, *Linn.*
acanthocalyx, *Klotzsch, in Peters, Reise Mossamb. Bot.* 232.—Afr. trop.
acanthocarpum, *Poit. in Rev. Hortic.* (1867) 290.—Cochinchina.
acanthodes, *Hook. f. Bot. Mag.* t. 6283.—Bras.
acanthoideum, *Drège, ex Dun. in DC. Prod.* xiii. 1. 364.—Afr. austr.
acerifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 41.—N. Granat.
acerosum, *Sendt. in Mart. Fl. Bras.* x. 61.—Bras.
acetosaefolium, Lam. Illustr. ii. 24; Encyc. iv. 306 = trilobatum.
aciculare, *Sw. in Roem. & Schult. Syst.* iv. 647.—Bras.
acropterum, *Griseb. Fl. Brit. W. Ind.* 437.—Ind. occ.
aculeastrum, *Dun. in DC. Prod.* xiii. 1. 366.—Afr. austr.
aculeatissimum, *Jacq. Coll.* i. 100.—As. et Am. trop.
aculeatissimum, *Moench, Meth.* 475.—Hab.?
aculeatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
aculeiger, *Moric. ex Dun. in DC. Prod.* xiii. 1. 301.—Bras.
aculeolatum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 143.—Mexic.
acuminatum, Dun. in DC. Prod. xiii. 1. 139 = apodum.
acuminatum, *Ruiz & Pav. Fl. Per.* ii. 34. t. 159.—Peruv.
acutangulum, *Griseb. in Lechl. Berb. Am. Austr.* 58.—Peruv.
acutifolium, Kit. in Linnaea, xxxii. (1863) 440 = nigrum.
acutifolium, *Ruiz & Pav. Fl. Per.* ii. 33. t. 162. f. b.—Peruv.
acutilobum, *Dun. ex Poir. Encyc. Suppl.* iii. 772.—Bras.
adelense, *Delile, in Rochet, Sec. Voy. Choa*, 345.—Abyss.
adenophorum, *F. Muell. Fragm.* ii. 162.—Austral.

SOLANUM:—
adenotrichum, *Dun. Hist. Solan.* 236.—Bras.
adhaerens, Willd. ex Roem. & Schult. Syst. iv. 669 = scabrum.
adoënse, *Hochst. ex A. Br. in Flora*, xxiv. (1841) 1. 278.—Abyss.
adscendens, *Sendt. in Mart. Fl. Bras.* x. 17.—Bras.
adulterinum, Buch.-Ham. ex Wall. Cat. sub n. 2616 = verbascifolium.
aegyptiacum, Forsk. Fl. Aegypt. Arab. 45 = nigrum.
aethiopicum, *Jacq. Hort. Vindob.* i. 4. t. 12.—Hab.?
aethiopicum, *Linn. Cent. Pl.* ii. 10; *Amoen. Acad.* iv. 307.—As. et Afr. trop.
aethiopoides, Cav. ex Roem. & Schult. Syst. iv. 669 (Quid?).—Hab.?
affine, *Dun. in DC. Prod.* xiii. 1. 168.—Cuba.
affine, Sendtn. in Mart. Fl. Bras. x. 63 = jatrophaefolium.
africanum, *Dun. Hist. Solan.* 238.—Africa.
africanum, Mill. Gard. Dict. ed. VIII. n. 26 = crassifolium.
Afzelii, *Dun. in DC. Prod.* xiii. 1. 363.—Afr. trop.
aggerum, *Dun. l. c.* 103.—Afr. austr.
aggregatum, *Jacq. Coll.* iv. 124.—Ind. occ.; Afr. trop.
agglutinatum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 48.—Bras.
agrarium, *Sendt. in Mart. Fl. Bras.* x. 69.—Bras.
agreste, Brouss. ex Roem. & Schult. Syst. iv. 669 (Quid?).—Hab.?
agreste, Roth, Nov. Pl. Sp. 130 = indicum.
agrimoniaefolium, Pav. ex Dun. in DC. Prod. xiii. 1. 25 = Lycopersicum agrimoniaefolium.
Ahlbergi, *Dun. l. c.* 372.—Bras.
alatum, Dun. l. c. 234 = decurrens.
alatum, Moench, Meth. 474 = nigrum.
alatum, Seem. & Schmidt, in Flora, xxvii. (1844) 497 = robustum.
albicaule, *Kotschy, ex Dun. in DC. Prod.* xiii. 1. 204.—Afr. trop.
albidum, *Dun. Hist. Solan.* 206.—Peruv.
albiflorum, *Phil. in Linnaea*, xxxiii. (1864–65) 202.—Chili.
album, *Lour. Fl. Cochinch.* 129.—Ins. Molucc.; Cochinch.
album, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 26 = S. Melongena.
aligerum, *Schlecht. in Linnaea*, xix. (1847) 301.—Mexic.
allogonum, Bernh. l. c. viii. (1833) 252 = Saracha allogona.
aloysiaefolium, *Dun. in DC. Prod.* xiii. 1. 73.—Bolivia.
Alphonsi, *Dun. l. c.* 69.—Chili.
alpinum, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 571.—Java.
alterato-pinnatum, *Steud. Nom.* ed. II. ii. 600.—Bras.
amarantoides, *Dun. in DC. Prod.* xiii. 1. 55.—Bras.
amazonium, *Ker-Gawl. in Bot. Reg.* t. 71.—Mexic.
ambiguum, *Dun. Solan. Syn.* 32.—Ins. Martinic.
amblophyllum, *Hook. Bot. Misc.* ii. (1831) 231.—Peruv.
amblycalycinum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
amblycalyx, Dun. in DC. Prod. xiii. 1. 371 = eriocalyx.
amblymerum, *Dun. l. c.* 294.—Austral.
ambrosiacum, *Vell. Fl. Flum.* 90; ii. t. 131.—Bras.
americanum, Mill. Gard. Dict. ed. VIII. n. 5 = nigrum.
amicorum, *Benth. in Hook. Lond. Journ Bot.* ii. (1843) 227.—Ins. Pacif.
amictum, *Moric. ex Dun. in DC. Prod.* xiii. 1. 263.—Mexic.
amoenum, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 300 = torvum.
amplexicaule, *Sendt. in Mart. Fl. Bras.* x. 14.—Bras.
amygdalifolium, *Steud. Nom.* ed. II. ii. 600.—Bras.
anacamptocarpum, *Dun. in DC. Prod.* xiii. 1. 59.—Java.
anacanthum, *Dun. l. c.* 183.—Mexic.
anceps, *Ruiz & Pav. Fl. Per.* ii. 36. t. 169.—Peruv.
Andreanum, *Baker, in Journ. Linn. Soc.* xx. (1884) 498.—N. Granat.
Andrieuxi, *Dun. in DC. Prod.* xiii. 1. 165.—Mexic.
Anghivi, Boj. ex Dun. l. c. 31 = seq.
Anguivi, Herb. Lamb. ex Dun. in DC. Prod. xiii. 1. 308 = occidentale.

SOLANUM:—
Anguivi, Hook. Exot. Fl. t. 199 = indicum.
Anguivi, *Lam. Illustr.* ii. 23 (an praec. ?).—Madag.
angulatum, Ruiz & Pav. Fl. Per. ii. 36 = quitoënse.
angulosum, Heyne, ex Nees, in Trans. Linn. Soc. xvii. (1834) 42 = bigeminatum.
Angurium, *Dun. Hist. Solan.* 243.—Mexic.
angustiflorum, *Mart. ex Sendt. in Mart. Fl. Bras.* x. 107.—Bras.
angustifolium, *Lam. Illustr.* ii. 18.—Reg. Argent.
angustifolium, Mill. Gard. Dict. ed. VIII. n. 15 = S. Angurium.
angustifolium, Ruiz & Pav. Fl. Per. ii. 33 = pulverulentum.
anisophyllum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 52.—Peruv.
anoacanthum, *Sendt. in Mart. Fl. Bras.* x. 66.—Bras.
anomalum, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 126.—Afr. trop.
anonaefolium, *Dun. Solan. Syn.* 15.—N. Granat.
anoplocladum, *Dun. in DC. Prod.* xiii. 1. 346.—Mexic.
anthropophagorum, Seem. in Bonplandia, ix. (1861) 258; x. (1862) 274. t. 14 = S. Uporo.
apalophyllum, *Dun. in DC. Prod.* xiii. 1. 42.—Peruv.
apiculatum, Dun. l. c. 99 = madagascariense.
apiculatum, *Sendt. in Mart. Fl. Bras.* x. 84.—Bras.
apocynifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 213.—Madag.
apodum, *Dun. in DC. Prod.* xiii. 1. 139.—Bras.
appendiculatum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 5.—Mexic.
appendiculatum, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 135 = Galeotti.
Aquartia, *Dun. Hist. Solan.* 187.—Ins. S. Doming.
aquaticum, *Dum.-Cours. Bot. Cult.* ed. I. *Suppl.* 152.—Hab. ?
arabicum, *Dun. Hist. Solan.* 240.—Arab.
Aracatscha, Bess. Cat. Pl. Hort. Crem. (1816) 135; et ex Roem. & Schult. Syst. iv. 669 = tuberosum.
arborescens, Moench, Meth. 474 = bonariense.
arboreum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 20.—Venezuela.
arcuatum, *Sendt. in Mart. Fl. Bras.* x. 60.—Bras.
ardisioides, *Blume, Bijdr.* 696.—Java.
arenarium, Schur, Enum. Pl. Transs. 477 = nigrum.
arenarium, *Sendt. in Mart. Fl. Bras.* x. 26.—Bras.
argenteum, Blanch. ex Dun. in DC. Prod. xiii. 1. 135 = Swartzianum.
argenteum, *Dun. ex Poir. Encyc. Suppl.* iii. 755.—Bras.
argenteum, Heyne, ex Wall. Cat. n. 2610 = giganteum.
argillicolum, *Dun. in DC. Prod.* xiii. 1. 236.—Bras.
argyacantha, *Dum.-Cours. Bot. Cult.* ed. II. iii. 161.—Hab. ?
armatum, *R. Br. Prod.* 446.—Austral.
armatum, Forsk. Fl. Aegypt. Arab. 47 = arabicum.
Arrabidae, Steud. Nom. ed. II. ii. 600 = Cyphomandra Velloziana.
Arrebenta, Vell. Fl. Flum. 89; ii. t. 127 = aculeatissimum, *Jacq.*
artense, *Montr in Mém. Acad. Lyon*, x. (1860) 245.—N. Caled.
asarifolium, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 10.—Venezuela.
aspero-lanatum, Ruiz & Pav. Fl. Per. ii. 39 = lanatum.
asperum, Hornem. ex Walp. Rep. iii. 49 = nigrum.
asperum, Pers. Syn. i. 229 = lanatum.
asperum, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = S. Radula.
asperum, Sieber, ex Dun. in DC. Prod. xiii. 1. 107 = callicarpaefolium.
asperum, *Vahl, Eclog. Am.* ii. 17.—Am. trop.
asphananthum, *Baker, in Journ. Bot.* xx. (1882) 220.—Madag.
assimile, Frivald. in Flora, xix. (1836) 439 = S. Dulcamara.
asterigerum, *Bertol. in Mém. Acad. Sc. Bologn.* ix. (1858) 167 (*Misc. Bot.* xix. 3).—China.
asterocormum, *Dun. in DC. Prod.* xiii. 1. 339.—Am. trop.
asteroideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = astroites.
asterophorum, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 79.—Bras.

SOLANUM:—
astroites, Forst. f. Prod. 89 = nigrum.
astroites, *Jacq. Eclog.* 97. t. 65.—Am. austr.
astrophorum, Jan, Elench. 4 = sodomeum.
athroanthum, *Dun. in DC. Prod.* xiii. 1. 208.—Java.
atriplicifolium, Desp. ex Dun. Solan. Syn. 12 = nigrum.
atriplicifolium, *Gill. ex Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 386.—Am. austr.
Atropo, Schum. & Thonn. Beskr. Guin. Pl. 124 = Thonningianum.
atropurpureum, *Schrank, in Syll. Ratisb.* i. (1824) 200.—Bras.
atrosanguineum, Hort. ex Dun. in DC. Prod. xiii. 1. 243 = atropurpureum.
atroviolaceum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 410.—Java.
atrovirens, *Dun. Hist. Solan.* 243.—Ind. or.
aturense, *Humb. & Bonpl. ex Dun. Solan. Syn.* 32.—Venezuela.
aurantiacum, *Sendt. in Mart. Fl. Bras.* x. 55.—Bras.
aureum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 16.—Ecuador.
aureum, Sw. ex Roem. & Schult. Syst. iv. 602 = Swartzianum.
aureum, Thunb. Pl. Bras. Dec. ii. 20 = Ahlbergi.
auriculatum, *Ait. Hort. Kew.* ed. I. i. 246.—As. trop.
auriculatum, Mart. ex Dun. in DC. Prod. xiii. 1. 115 = granuloso-leprosum.
austro-caledonicum, *Seem. in Journ. Bot.* i. (1863) 209.—Ins. Pacif.
aviculare, *Forst. f. Prod.* 18.—Austral.; N. Zel.
axilliflorum, *Dun. Hist. Solan.* 238.—Mexic.
baccatum, Hort. ex Steud. Nom. ed. II. ii. 601 = bonariense.
bahamense, Forsk. Fl. Aegypt. Arab. 46 = glabratum.
bahamense, *Linn. Sp. Pl.* 188.—Ind. occ.
bahamense, Mill. Gard. Dict. ed. VIII. n. 24 = verbascifolium.
Balbisii, Boj. ex Dun. in DC. Prod. xiii. 1. 326 = Richardi.
Balbisii, Dun. Hist. Solan. 232. t. 3 = sisymbrifolium.
Baencks, Vand. Fasc. Pl. 10, sphalm. = seq.
Banksianum, *Dun. Hist. Solan.* 242 (Bancksianum).—Bras.
barbadense, Crantz, Inst. ii. 374 = Rivina humilis.
barbisetum, *Nees, in Trans. Linn. Soc.* xvii. (1834) 51.—Ind. or.
Bassovia, *Dun. in Poir. Encyc. Suppl.* iii. 754.—Bras.
Bassovia, Mart. ex Dun. in DC. Prod. xiii. 1. 406 = Bassovia Richardi.
Bauerianum, *Endl. Prod. Fl. Norf.* 54.—Ins. Norfolk.
bejarense, Moric. ex Dun. in DC. Prod. xiii. 1. 330 = rostratum.
Belfort, Vand. Fasc. Pl. 11 = paniculatum.
Belfortianum, Dun. Hist. Solan. 242 = paniculatum.
benguelense, *Peyr. in Sitzb. Akad. Wien, Math.-Nat.* xxxviii. (1860) 575.—Afr. trop.
Benthami, *Dun. in DC. Prod.* xiii. 1. 101.—Mexic.
Berterianum, *Dun. l. c.* 167.—Ins. S. Doming.
Berterii, *Hort. Par. ex Steud. Nom.* ed. II. ii. 601.—Chili.
Besseri, Weinm. ex Roem. & Schult. Syst. iv. 593 = nigrum.
Besserianum, Weinm. Elench. Hort. Pawlowsk. (1824) 100 = praec.
betaceum, Cav. in Anal. Hist. Nat. i. (1799) 44; Ic. t. 524 = Cyphomandra betacea.
biceps, *Dun. in DC. Prod.* xiii. 1. 341.—Bras.
bicolor, Mart. Herb. Fl. Bras. n. 1254 = Swartzianum, argenteum.
bicolor, Willd. ex Roem. & Schult. Syst. iv. 661 = verbascifolium.
bicorne, *Dun. in DC. Prod.* xiii. 1. 232.—Am. trop.
bifidum, *Dun. l. c.* 122.—N. Granat.
bifidum, Dun. l. c. 341 = biceps.
bifissum, Vell. Fl. Flum. 86; ii. t. 111 = biceps.
bifissum, *Vell. l. c.* 90; ii. t. 129.—Bras.
biflorum, R. Br. Prod. 445 = dianthophorum.
biflorum, D. Don, Prod. Fl. Nep. 96 = pachypetalon.
biflorum, *Lour. Fl. Cochinch.* 129.—Malaya; China.
bifolium, Aubl. ex Dun. in DC. Prod. xiii. 1. 149 = leucocarpon.
biformifolium, *Ruiz & Pav. Fl. Per.* ii. 32. t. 162.—Peruv.

SOLANUM:—
bifurcum, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 24, nomen.—Abyss.
bigeminatum, *Nees, in Trans. Linn. Soc.* xvii. (1834) 42.—Ind. or.
bimense, *Miq. Fl. Ind. Bat.* ii. 642.—Ins. Sunda.
bipatens, *Dun. in DC. Prod.* xiii. 1. 209.—Bras.
blattarioides, Koen. ex Dun. l. c. 127 = pubescens.
Blodgettii, *Chapm. Fl. S. U. St.* 349.—Florida.
Blumei, *Nees, ex Blume, Bijdr.* 696.—Java.
boerhaviaefolium, *Sendt. in Mart. Fl. Bras.* x. 48.—Bras.
bogotense, *Dun. in DC. Prod.* xiii. 1. 121.—N. Granat.
Boisseri, *Dun. l. c.* 188.—Guian.
Bojeri, *Dun. l. c.* 344.—Afr. trop.
boldoënse, *Dun. & A. DC. l. c.* 679.—Cuba.
boliviense, *Dun. l. c.* 43.—Bolivia.
bombense, Banks, ex Dun. l. c. 271. = stenorchis.
bombense, Jacq. Enum. Pl. Carib. 15 = glabrum.
bonariense, *Linn. Sp. Pl.* 185.—Reg. Argent.
Botelhianum, *Dun. Hist. Solan.* 239.—Bras.
Botelho, Vand. Fasc. Pl. 10 = praec.
brachyacanthum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 32.—N. Granat.
brachyantherum, *Phil. in Anal. Univ. Chil.* (1873) 522.—Chili.
brachybotryon, *Dun. in DC. Prod.* xiii. 1. 212.—Guian.
brachylobum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 71.—Peruv.
brachypodon, Dun. in DC. Prod. xiii. 1. 411 = Bassovia brachypoda.
brachystachys, *Dun. l. c.* 128.—Mexic.
bracteatum, *Thunb. in Act. Gorenk.* (1812); *ex ej. Fl. Cap.* ii. 57.—Afr. austr.
brancaefolium, Jacq. Eclog. 14. t. 7 = sisymbrifolium.
brasilianum, *Dun. Solan. Syn.* 48.—Bras.
brevidens, *Phil. in Anal. Univ. Chil.* (1873) 521.—Chili.
brevifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 22.—Ecuador.
brevipes, *Dun. in DC. Prod.* xiii. 1. 203.—Bras.
brevipilosum, *Dun. l. c.* 199.—Guian.
brevipilum, Dun. Hist. Solan. 191. t. 21 = jamaicense.
Bridgesii, *Phil. in Linnaea*, xxxiii. (1864–65) 203.—Chili.
Brownii, Dun. Hist Solan. 201 = violaceum.
Brucei, Delile, ex Dun. in DC. Prod. xiii. 1. 282 = adoënse.
buddleiaefolium, *Sendt. in Mart. Fl. Bras.* x. 83.—Bras.
budleiaefolium, Dun. in DC. Prod. xiii. 1. 115 = granuloso-leprosum.
Bulbocastanum, *Dun. in Poir. Encyc. Suppl.* iii. 749.—Mexic.
bullato-rugosum, Dun. in DC. Prod. xiii. 1. 236 = cyanocarphium.
bullatum, *Vell. Fl. Flum.* 84; ii. t. 104.—Bras.
bumeliaefolium, *Dun. in DC. Prod.* xiii. 1. 292.—Madag.
Burchellii, *Dun. l. c.* 291.—Afr. austr.
Bustillosi, *Phil. in Linnaea*, xxix. (1857–58) 24.—Chili.
byrsinum, Voigt, Hort. Suburb. Calc. 511 = havanense, *Jacq.*
Caavurana, *Vell. Fl. Flum.* 86; ii. t. 112.—Bras.
cabiliense, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 713 = marginatum.
caeruleum, Vell. Fl. Flum. 86; ii. t. 110 = indigoferum.
caesium, *Griseb. in Goett. Abh.* xxiv. (1879) 254.—Reg. Argent.
cajanumense, H. B. & K. Nov. Gen. et Sp. iii. 47 = Cyphomandra cajanumensis.
Caldasii, *Humb. & Bonpl. ex Dun. Solan. Syn.* 6.—Ecuador.
calicinum, *Dun. in Poir. Encyc. Suppl.* iii. 747.—Mexic.
californicum, Dun. in DC. Prod. xiii. 1. 86 = umbelliferum.
Calleryanum, Dun. l. c. 178 = biflorum.
callicarpaefolium, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1853) *App.* 10.—Am. trop.
callicarpum, Hort. Berol. ex A. Br. l. c. = Karstenii, *A. Br. & Bouché.*

SOLANUM:—
calophyllum, *Phil. in Anal. Univ. Chil.* (1862) II. 403.—Chili.
calvefactum, Dun. in DC. Prod. xiii. I. 226 = lanceaefolium.
calycinum, Hort. ex Dun. l. c. 431 = Saracha procumbens.
calycinum, Moç. & Sesse, ex Dun. Solan. Syn. 10 = calicinum.
calycinum, Nees, in Trans. Linn. Soc. xvii. (1834) 60 = pubescens.
calygnaphalum, Ruiz & Pav. Fl. Per. ii. 31 = gnaphaloides.
campaniforme, *Roem. & Schult. Syst.* iv. 662.—Bras.
campanulatum, *R. Br. Prod.* 446.—Austral.
campanulatum, Willd. ex Sendt. in Mart. Fl. Bras. x. 22 = campaniforme.
campechiense, Banks, ex Ait. Hort. Kew. ed. I. i. 251 = aculeatissimum, *Jacq.*
campechiense, Hort. Par. ex Dun. in DC. Prod. xiii. I. 260 = torvum.
campechiense, *Linn. Sp. Pl.* 187.—Am. trop.
campechiense, Linn. l. c., partim = fuscatum, *Linn.*
campylacanthum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 24, nomen; *Dun. in DC. Prod.* xiii. I. 336.—Abyss.
campylocladum, *Dun. l. c.* 173.—Mexic.
canaranum, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 236 = trilobatum.
candicans, *Dun. Solan. Syn.* 23.—Peruv.
candidum, *Lindl. Bot. Reg.* (1839) *Misc.* 73.—Mexic.
canescens, Blume, Bijdr. 701 = indicum.
canescens, Kit. in Linnaea, xxxii. (1863) 440 = nigrum.
canescens, Pav. ex Dun. in DC. Prod. xiii. I. 343 = mexicanum.
canescens, Rich. ex Dun. l. c. 107 = S. Radula.
cano-cinereum, *Dun. l. c.* 213.—Guian.
capense, *Linn. Syst.* ed. X. 935; *Linn. f. Suppl.* 147.—Afr. austr.
capoerum, *Dun. in DC. Prod.* xiii. I. 112.—Bras.
Capsicastrum, *Link, ex Schau. in Otto & Dietr. Allg. Gartenz.* i (1833) 228.—Bras.
capsicoides, Hort. Par. ex Lam. Illustr. ii. 21 = aculeatissimum, *Jacq.*
capsicoides, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 78.—Bras.
caracasanum, Roem. & Schult. Syst. iv. 664 = Cyphomandra caracasana.
cardiophyllum, Dun. in DC. Prod. xiii. I. 89 = boldoënse.
cardiophyllum, *Lindl. in Journ. Hort. Soc.* iii. (1848) 70.—Mexic.
cardiospermum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 (Quid?).—Java.
carduiforme, *F. Muell. Fragm.* ii. 163.—Austral.
carense, *Dun. in DC. Prod.* xiii. I. 105.—Arab.
Cari, *Molina, Sagg. Chile*, 132.—Chili.
caribaeum, *Dun. in DC. Prod.* xiii. I. 48.—Ind occ.
caripense, *Humb. & Bonpl. ex Dun. Solan. Syn.* 8.—Venezuela.
carolinense, *Linn. Sp. Pl.* 187.—Am. bor.
caroliniense, Mill. Gard. Dict. ed. VIII. n. 21 = amazonium.
cataphractum, *A. Cunn. ex Benth. Fl. Austral.* iv. 459.—Austral.
catombelense, *Peyr. in Sitzb. Akad. Wien, Math.-Nat.* xxxviii. (1860) 576.—Afr. trop.
caudatum, *Dun. in DC. Prod.* xiii. I. 140.—Peruv.
caulorhizum, Dun. l. c. 181 = biflorum.
Cavanillesii, Dun. l. c. 61 = pinnatum.
cearense, *Dun. l. c.* 90.—Bras.
čechicum, *Opiz, in Lotos*, iv. (1854) 94 (Quid?).—Bohem.
celastroides, Sendt. ex Mart. in Flora, xxiv. (1841) II. Beibl. 86 = Bassovia lucida.
cerasiferum, *Dun. in DC. Prod.* xiii. I. 365.—Afr. bor.
cerinum, Hort. Dresd. ex Otto, in Ind. Sem. Hort. Berol. (1853) 8 = villosum, *Willd.*
cernuum, *Vell. Fl. Flum.* 84; ii. t. 103.—Bras.
Cervantesii, *Lag. Gen. et Sp. Nov.* 10.—Mexic.
cestrifolium, Jacq. ex Weinw. Elench. Hort. Pawlowsk. (1824) 100; et in Spreng. Syst. i. 680 = nigrum.
cestrifolium, Willd. ex Dun. in DC. Prod. xiii. I. 186 = seq.
cestrophyllum, *Dun. l. c.*—Hab. ?
chaeacanthum, *Griseb. Cat. Pl. Cub.* 190.—Cuba.

SOLANUM:—
chaetacanthum, *Dun. in DC. Prod.* xiii. I. 186.—Bras.
chenopodifolium, *Dun. l. c.* 44.—Reg. Argent.
chenopodinum, *F. Muell. Fragm.* ii. 165.—Austral.
chenopodioides, Hort. ex Dun. in DC. Prod. xiii. I. 55 = atriplicifolium.
chenopodioides, Lam. Illustr. ii. 18 = nigrum.
chinense, Dun. Hist. Solan. 240 = indicum.
chloranthum, *DC. Pl. Rar. Jard. Genèv.* t. 13.—Reg. Argent.
chloranthum, Poepp. ex Sendt. in Mart. Fl. Bras. x. 103 = Poeppigianum.
chloranthum, Salzm. ex Dun. in DC. Prod. xiii. I. 236 = argillicolum.
chloranthum, Spreng. Syst. i. 682 = Sprengelii.
chlorocarpum, Schur, Enum. Pl. Transs. 478 = nigrum.
chloroleucum, *Dun. in DC. Prod.* xiii. I. 279.—Bras.
chloropetalon, *Schlecht. in Linnaea*, xix. (1847) 291.—Mexic.
chlorophloium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = umbellatum, *Mill.*
chrysacanthum, *Dun. in DC. Prod.* xiii. I. 330.—Mexic.
chrysobotrys, Hort. Berol. ex Walp. Rep. iii. 100 = Lycopersicum peruvianum.
chrysophyllum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 25.—Ecuador.
chrysotrichum, *Schlecht. in Linnaea*, xix. (1847) 304.—Mexic.
ciliare, Willd. Enum. Hort. Berol. 237 = aculeatissimum, *Jacq.*
ciliatum, Blume, ex Miq. Fl. Ind. Bat. ii. 643 = nematosepalum.
ciliatum, Lam. Illustr. ii. 21 = aculeatissimum, *Jacq.*
ciliolatum, *Mart. & Gal. in Bull. Acad. Brux.* xii. I. (1845) 140.—Mexic.
cinerescens, Pav. ex Dun. in DC. Prod. xiii. I. 343 = mexicanum.
cinereum, R. Br. in Salt, Abyss. App. 62 = Saltianum.
cinereum, *R. Br. Prod.* 446.—Austral.
cinnamomeum, *Sendt. in Mart. Fl. Bras.* x. 44.—Bras.
cirrhosum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 9. (cyrrhosum).—Venezuela.
citrifolium, *Willd. ex Roem. & Schult. Syst.* iv. 662.—N. Granat.
citrullifolium, A. Br. in Ind. Sem. Hort. Friburg. (1849); ex Ann. Sc. Nat. Sér. III. xii. (1849) 356 = heterodoxum.
cladotrichodes, Mart. ex Sendt. in Mart. Fl. Bras. x. 28 = megalochiton.
cladotrichum, *Dun. Hist. Solan.* 236.—Bras.
cladotrichum, Mart. ex Dun. in DC. Prod. xiii. I. 223 = paratyense.
clathratum, *Sendt. in Mart. Fl. Bras.* x. 35.—Am. trop.
Clausseni, *Dun. in DC. Prod.* xiii. I. 271.—Bras.
claviceps, *Griseb. in Goett. Abh.* xix. (1874) 221.—Reg. Argent.
coagulans, Forsk. Fl. Aegypt. Arab. 47 = incanum, *Linn.*
coagulans, Jacq. Hort. Schoenb. iv. 35. t. 469 = schoenbrunnense.
coagulans, Schimp. ex Dun. in DC. Prod. xiii. I. 370 = marginatum.
coarense, *Sendt. in Mart. Fl. Bras.* x. 19.—Bras.
coccineum, Hort. Par. ex Dun. in DC. Prod. xiii. I. 310 = indicum.
coccineum, Jacq. Misc. ii. 329 = tomentosum, *Linn.*
coeruleum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = cyanocarphium.
collinum, Dun. in DC. Prod. xiii. I. 36 = Commersonii.
columbianum, Dun. l. c. 33 = tuberosum.
Comitis, *Dun. l. c.* 345.—Java.
comitum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
Commersonii, *Dun. ex Poir. Encyc. Suppl.* iii. 746.—Reg. Argent.
commutatum, Spreng. Pugill. i. 18 = Lycopersicum peruvianum.
compactum, Hort.—Cf. Gard. Chron. (1872) 641 = S. Pseudo-capsicum.
conanthum, Dun. in DC. Prod. xiii. I. 127 = pubescens.
concavum, *Lindl. in Bot. Reg.* (1842) *Misc.* 57.—Chili.
concinnum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 36.—Bras.

SOLANUM:—
concisum, *Dun. in DC. Prod.* xiii. I. 327.—Peruv.
confertum, Dun. l. c. 142 = indigoferum.
confine, *Dun. l. c.* 137.—Peruv.
congense, *Link, Enum. Hort. Berol.* i. 187.—Afr. trop.
congestiflorum, *Dun. in DC. Prod.* xiii. I. 92.—Chili.
congestum, Vell. Fl. Flum. 82; ii. t. 90 (Lycii sp).—Bras.
conglobatum, *Dun. in DC. Prod.* xiii. I. 112.—Bras.
conicum, *Ruiz & Pav. Fl. Per.* ii. 38. t. 172. f. b.—Peruv.
conicum, Vell. Fl. Flum. 83; ii. t. 96 = Cyphomandra sciadostylis.
coniodes, Mor. ex Dun. in DC. Prod. xiii. I. 64 = Chamaesaracha sordida.
conocarpum, *Dun. in Poir. Encyc. Suppl.* iii. 748.—Ind. occ.
convolvuliforme, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
Convolvulus, *Sendt. in Mart. Fl. Bras.* 48.—Bras.
cordatum, *Forsk. Fl. Aegypt. Arab.* 47.—Arab.
cordatum, Pav. ex Dun. in DC. Prod. xiii. I. 396 = Cyphomandra splendens.
cordifolium, *Dun. in Poir. Encyc. Suppl.* iii. 764.—Bras.
coriaceum, *Dun. Hist. Solan.* 197.—Guian.
coriaceum, Hook. Bot. Mag. t. 2703 = havanense, *Jacq.*
corifolium, F. Muell. Fragm. ii. 166 = discolor.
cormanthum, *Vell. Fl. Flum.* 86; ii. t. 113.—Bras.
corniculatum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 45.—Bras.
cornifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 21.—N. Granat.
cornigerum, *André, in Rev. Hortic.* (1868) 33.—Afr. trop.
cornigerum, *Dun. in DC. Prod.* xiii. I. 75.—Bras.
cornutum, Hort. Monsp. ex Dun. in DC. Prod. xiii. I. 329 = rostratum.
cornutum, *Lam. Illustr.* ii. 25.—Mexic.
coronatum, Vell. Fl. Flum. 82; ii. t. 92 = sambuciflorum.
coronopoideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
Coronopus, Dun. in DC. Prod. xiii. I. 64 = Chamaesaracha Coronopus.
cortex-virens, Dun. l. c. 372 = umbellatum, *Mill.*
corymbiferum, J. F. Gmel. Syst. 384 = corymbosum.
corymbosum, Herb. Banks, ex Dun. in DC. Prod. xiii. I. 347 = macrophyllum.
corymbosum, *Jacq. Coll.* i. 78.—Peruv.
corymbosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = verbascifolium.
corymbosum, Wight, ex Nees, in Trans. Linn. Soc. xvii. (1834) 48 = vagum.
cosmeticum, *Delile, in Rochet, Sec. Voy. Choa*, 340, nomen.—Abyss.
Cotoneaster, Dun. in DC. Prod. xiii. I. 158 = Orbignianum.
cotopaxense, *Dun. l. c.* 139.—Ecuador.
Coxi, *Phil. in Linnaea*, xxxiii. (1864–65) 200.—Chili.
crassiflorum, Schott, ex Sendt. in Mart. Fl. Bras. x. 20 = S. Caavurana.
crassifolium, *Lam. Illustr.* ii. 16.—Afr. austr.
crassifolium, Ortega, Hort. Matr. Dec. 117 = Cyphomandra betacea.
crassifolium, Salisb. Prod. 133 = macrocarpon.
crassipetalum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 256.—Reg. Himal.
crassipetalum, Wight, ex C. B. Clarke in Hook. f. Fl. Brit. Ind. iv. 231 = denticulatum.
crenato-dentatum, Dun. in DC. Prod. xiii. I. 54 = nigrum.
Crepini, *Heurck, Obs. Bot.* 89.—Afr. trop.
crinitipes, *Dun. in DC. Prod.* xiii. I. 317.—N. Granat.
crinitum, *Lam. Illustr.* ii. 20.—Guian.
crispum, Bert. ex Dun. in DC. Prod. xiii. I. 92 = congestiflorum.
crispum, *Ruiz & Pav. Fl. Per.* ii. 31. t. 158.—Chili.
croceum, Dun. Hist. Solan. 241 = jamaicense.
crotonifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 18.—Am. austr.
crotonoides, *Lam. Illustr.* ii. 24.—Ind. occ.
crotonoides, Michx. ex Dun. in DC. Prod. xiii. I. 260 = torvum.

SOLANUM:—
crotonoides, Sieber, ex Presl, Bot. Bemerk. 101 = lanceaefolium.
cryptolobum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 63.—Hab.?
cryptopodium, Phil. f. Cat. Pl. Vasc. Chil. 227, sphalm. = cyrtopodium.
cubense, *Dun. in DC. Prod.* xiii. 1. 379, nomen.—Cuba.
cumanense, Schlecht. ex Roem. & Schult. Syst. iv. 662 = sylvaticum.
Cumingii, Dun. in DC. Prod. xiii. 1. 363 = S. Melongena.
cuneatum, Moench, Meth. 476 = indicum.
cuneifolium, Dun. Hist. Solan. 193 = jamaicense.
cuneifolium, Hort. Avens. ex Dun. in DC. Prod. xiii. 1. 53 = nigrum.
cuneifolium, Willd. ex Roem. & Schult. Syst. iv. 623 = Willdenovianum.
Cunninghamii, *Benth. Fl. Austral.* iv. 465.—Austral.
cupulatum, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 231 = bigeminatum.
curvispinum, *Dun. in DC. Prod.* xiii. 1. 265.—Bras.
cuspidatum, Dun. l. c. 111 = cinnamomeum.
cuspistorta, Dun. l. c. 231 = Poeppigianum.
cutacense, *H. B. & K. Nov. Gen. et Sp.* iii. 38.—Peruv.
cyananthum, *Dun. in DC. Prod.* xiii. 1. 316.—Bras.
cyanocarpium, *Blume, Bijdr.* 700.—Java.
cyathocalyx, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 65.—Peruv.
cydoniaefolium, *Dun. in DC. Prod.* xiii. 1. 209.—Hab.?
cylindricum, Vell. Fl. Flum. 87; ii. t. 119 = Cyphomandra cylindrica.
cymosum, Banks, ex Dun. in DC. Prod. xiii. 1. 73 = viride.
cymosum, Ortega, Hort. Matr. Dec. 12 = macrophyllum.
cymosum, Ruiz & Pav. Fl. Per. ii. 31. t. 160 = corymbosum.
cynanchoides, Dun. in DC. Prod. xiii. 1. 82 = jasminoides.
cyrtopodium, *Dun. l. c.* 195.—Chili.
Dallachii, *Benth. Fl. Austral.* iv. 456.—Austral.
darfurense, Delile, ex Dun. in DC. Prod. xiii. 1. 204 = albicaule.
dasyphyllum, *Schum. & Thonn. Beskr. Guin. Pl.* 126.—Afr. trop.
dasypodum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
dasypus, *Drège, ex Dun. in DC. Prod.* xiii. 1. 161.—Afr. austr.
daturaefolium, *Dun. l. c.* 261.—Bras.
dealbatum, Lindl. in Trans. Hortic. Soc. vii. (1830) 52 = elaeagnifolium.
debile, *Dun. in DC. Prod.* xiii. 1. 74.—Bras.
decemdentatum, Roxb. Hort. Beng. 16; Fl. Ind. i. 565 = biflorum.
decemfidum, Pav. ex Dun. in DC. Prod. xiii. 1. 173 = lentum.
decipiens, *Opiz, in Bercht. & Opiz, Oekon. Fl. Boehm.* iii. 24 (Quid?).—Bohemia.
decompositifolium, *Sendt. in Mart. Fl. Bras.* x. 88.—Bras.
decorticans, *Sendt. l. c.* 47.—Bras.
decorticatum, Dun. in DC. Prod. xiii. 1. 135 = praec.
decorum, *Sendt. in Mart. Fl. Bras.* x. 83.—Bras.
decurrens, Balb. in Mém. Acad. Turin (1810), 360 = sisymbrifolium.
decurrens, Banks, ex Dun. in DC. Prod. xiii. 1. 200 = jamaicense.
decurrens, *Vell. Fl. Flum.* 89; ii. t. 126.—Bras.
decurrens, Wall. ex Dun. in DC. Prod. xiii. 1. 50 = nigrum.
defensum, *F. Muell. Fragm.* v. 193.—Austral.
delagoënse, *Dun. in DC. Prod.* xiii. 1. 349.—Afr. trop.
Delilei, *Dun. l. c.* 81.—Mexic.
demerarense, Dun. l. c. 257 = stramonifolium.
demissum, Lindl. in Journ. Hortic. Soc. iii. (1848) 70 = tuberosum.
densevestitum, *F. Muell. ex Benth. Fl. Austral.* iv. 456.—Austral.
densiflorum, Dun. in DC. Prod. xiii. 1. 208 = athroanthum.
densiflorum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 141.—Mexic.

SOLANUM:—
densiflorum, Sendt. in Mart. Fl. Bras. x. 93 = piluliferum.
denticulatum, *Blume, Bijdr.* 697.—Ind. or.; Malaya.
denticulatum, Nees, in Trans. Linn. Soc. xvii. (1834) 41 = subtruncatum.
depauperatum, *Dun. in DC. Prod.* xiii. 1. 227.—Bras.
Deppei, Hort. ex Dun. l. c. 52 = nigrum.
Desvauxii, *Ham. Prod. Pl. Ind. Occ.* 26.—Ind. occ.
diantherum, Vell. Fl. Flum. 83; ii. t. 99 = concinnum.
dianthophorum, *Dun. Hist. Solan.* 183.—Austral.
dibrachiatum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 59.—Peruv.
dichotomum, *Lour. Fl. Cochinch.* 129.—China.
dichotomum, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 138 = Martensii.
dichotomum, Pav. ex Dun. in DC. Prod. xiii. 1. 233 = bicorne.
dichotomum, Rich. ex Poir. Encyc. Suppl. iii. 757 = uniflorum.
dichotomum, Ruiz & Pav. Fl. Per. ii. 34 = riparium.
dichotomum, Vand. Fasc. Pl. 9 = Vandellianum.
dichroandrum, *Dun. in DC. Prod.* xiii. 1. 86.—Venezuela.
dichrophyllum, Dun. l. c. 48 = alpinum.
dictyoticum, *Roem. & Schult. Syst.* iv. 661.—Hab.?
didymanthum, *Dun. in DC. Prod.* xiii. 1. 289.—Afr. austr.
didymum, *Dun. Hist. Solan.* 236.—Bras.
diffusum, Link, ex Roem. & Schult. Syst. iv. 601 = Linkianum.
diffusum, Roxb. Hort. Beng. 17; Fl. Ind. i. 568 = xanthocarpum.
diffusum, *Ruiz & Pav. Fl. Per.* ii. 37. t. 171. f. b.—Peruv.
diffusum, Vell. Fl. Flum. 83; ii. t. 98 = nigrum.
diflorum, Vell. l. c. 84; ii. t. 102 = S. Capsicastrum.
Dilleni, Schult. Oestr. Fl. ed. II. i. 393 = nigrum.
dimidiatum, *Sendt. in Mart. Fl. Bras.* x. 102.—Bras.
diphyllum, Banks, ex Dun. in DC. Prod. xiii. 1. 170 = retrofractum.
diphyllum, Forsk. Fl. Aegypt. Arab. 63 = microcarpum.
diphyllum, *Linn. Sp. Pl.* 184.—Am. trop.
diphyllum, Osbeck, Voy. i. 328; et ex D. Don, Prod. Fl. Nep. 96 = biflorum.
discolor, *R. Br. Prod.* 445.—Austral.
discolor, Dun. in DC. Prod. xiii. 1. 111 = leucodendron.
dispar, Loisel. ex Dun. l. c. 69 = aviculare.
distichophyllum, *Sendt. in Mart. Fl. Bras.* x. 85.—Bras.
distichum, *Schum. & Thonn. Beskr. Guin. Pl.* 122.—Afr. trop.
divaricatum, Dun. in DC. Prod. xiii. 1. 125 = didymum.
divaricatum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 136.—Mexic.
divergens, Dun. in DC. Prod. xiii. 1. 411 = Bassovia leptopoda.
diversiflorum, *F. Muell. Fragm.* vi. 146.—Austral.
diversifolium, Humb. & Bonpl. ex Dun. Solan. Syn. 8 = Cyphomandra caracasana.
diversifolium, *Schlecht. in Linnaea*, xix. (1846) 297.—Mexic.
Dombeyi, *Dun. in DC. Prod.* xiii. 1. 162.—Peruv.
domingense, *Dun. l. c.* 80.—Ins. S. Doming.
domingense, Willd. ex Roem. & Schult. Syst. iv. 665 = insulanum.
Donianum, Walp. Rep. iii. 54 = bahamense.
Douglasii, Dun. in DC. Prod. xiii. 1. 48 = nigrum.
Douvillei, *Moric. ex Dun. l. c.* 271.—Bras.
Dregei, *Dun. l. c.* 288.—Afr. austr.
Dregei, Presl, Bot. Bermerk. 101 = aculeastrum.
dryadeum, *Dun. in DC. Prod.* xiii. 1. 196.—Bras.
dubium, Dun. Hist. Solan. 240 = jamaicense.
dubium, *Fresen. in Mus. Senckenb.* i. (1833) 166.—Arab.
Dulcamara, *Linn. Sp. Pl.* 185.—Europ.; As. et Afr. bor.
dulcamaroides, Dun. in Poir. Encyc. Suppl. iii. 751 = macrantherum.
dulcamaroides, Rich. ex Dun. in DC. Prod. xiii. 1. 87 = pyrifolium.
dulcamarum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Dulcamara.

SOLANUM:—
dumetorum, Dun. in DC. Prod. xiii. 1. 112 = rufescens.
dumetorum, *Willd. ex Roem. & Schult. Syst.* iv. 662.—Hab.?
Dunalianum, *Gaudich. in Freyc. Voy. Bot.* 448. t. 58.—Ins. Molucc.
duplo-sinuatum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 233.—Afr. trop.
eccremocarpum, *Dun. in DC. Prod.* xiii. 1. 225.—Bras.
Echegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, ix. (1881) 58.—Reg. Argent.
echidnaeforme, *Dun. in DC. Prod.* xiii. 1. 324.—Bras.
echinatum, *R. Br. Prod.* 447.—Austral.
echinatum, Pav. ex Dun. in DC. Prod. xiii. 1. 173 = lentum.
echinatum, Rich. ex Dun. l. c. 317 = crinitum.
echinocarpum, *Bredem. ex Roem. & Schult. Syst.* iv. 659.—Venezuela.
echinocaulon, Moric. ex Dun. in DC. Prod. xiii. 1. 316 = cyananthum.
Edmonstonei, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 201.—Ins. Galapag.
edule, Schum. & Thonn. Beskr. Guin. Pl. 125 = S. Melongena.
edule, Vell. Fl. Flum. 88; ii. t. 121 = sisymbrifolium.
Edwardi, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1846); *ex Dun. in DC. Prod.* xiii. 1. 107.—Cuba.
elachophyllum, *F. Muell. Fragm.* ii. 164.—Austral.
elaeagnifolium, *Cav. Ic.* iii. 22. t. 243.—Am. trop.
elaeagnifolium, Willd. ex Steud. Nom. ed. II. ii. 602 = obtusifolium.
elatum, *Link, Enum. Hort. Berol.* i. 186.—Hab.?
elegans, Dun. ex Poir. Encyc. Suppl. iii. 769 = cinereum.
elegans, Vell. Fl. Flum. 83; ii. t. 95 = Cyphomandra Velloziana.
ellipticum, *R. Br. Prod.* 446.—Austral.
ellipticum, Vell. Fl. Flum. 84; ii. t. 100 = Cyphomandra elliptica.
elongatum, Hort. ex Roem. & Schult. Syst. iv. 611 = virgatum.
Endlicheri, *Dun. in DC. Prod.* xiii. 1. 67.—Peruv.
enneodonton, Delile, ex Dun. l. c. 257 = robustum.
enoplocalyx, *Dun. l. c.* 222.—Peruv.
ensifolium, *Dun. l. c.* 186.—Mexic.
eremanthum, *Dun. l. c.* 129.—Bras.
eremophilum, *F. Muell. in Linnaea*, xxv. (1852) 432.—Austral.
erianthum, D. Don, Prod. Fl. Nep. 96 = verbascifolium.
eriocalyx, *Dun. Hist. Solan.* 237.—N. Granat.
eriocalyx, Dun. in DC. Prod. xiii. 1. 124 = trichocalyx.
eriocarpum, Dun. l. c. 131 = gnaphalocarpum.
eriophorum, Dun. l. c. 249 = barbisetum.
eriophyllum, Dun. Solan. Syn. 30 = lasiophyllum, *Dun.*
erosum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 83 = indicum.
erysimifolium, *Delile, in Rochet, Sec. Voy. Choa*, 342.—Abyss.
erythracanthum, *Boj. ex Dun. in DC. Prod.* xiii. 1. 201.—Madag.
erythraeum, Dun. Hist. Solan. 238 = nigrum.
erythrocarpon, G. F. W. Mey. Prim. Fl. Esseq. 109 = nigrum.
Eschweilerianum, Zipp. ex Miq. Fl. Ind. Bat. ii. 642 = Blumei.
esculentum, Drège, ex Dun. in DC. Prod. xiii. 1. 367 = subexarmatum.
esculentum, Dun. Hist. Solan. 208. t. 3 = S. Melongena.
esculentum, Neck. Delic. Gallo-Belg. i. 119 = tuberosum.
Esenbeckii, Steud. Nom. ed. II. ii. 602 = pubescens.
esuriale, *Lindl. in Mitch. Three Exped.* ii. 43.—Austral.
etuberosum, *Lindl. Bot. Reg.* t. 1712.—Chili.
euacanthum, *Phil. Sert. Mendoc. Alt.* 37.—Chili.
euracanthum, Dun. in DC. Prod. xiii. 1. 257 = robustum.
evonymoides, *Remy, in C. Gay, Fl. Chil.* v. 81.—Chili.
evonymoides, *Sendt. in Flora*, xxiv. (1841) II. *Beibl.* 87.—Bras.
exasperatum, *Drège, ex Dun. in DC. Prod.* xiii. 1. 104.—Afr. austr.

SOLANUM:—
excelsum, *A. St. Hil. Voy. Distr. Diam.* i. 104.—Bras.
excelsum, Salisb. Prod. 133 = bonariense.
extraaxillare, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
farinosum, Wall. Cat. n. 2610 = giganteum.
fasciatum, Roem. & Schult. Syst. iv. 664; Steud. Nom. ed. II. ii. 602, sphalm. = fuscatum, *Roem. & Schult.*?
fasciculatum, Mart. in Flora, xxi. (1838) Beibl. II. 81 = Bassovia lucida.
fasciculatum, F. Muell. Fragm. i. 123 = simile.
fasciculatum, Vell. Fl. Flum. 85; ii. t. 106 = Bassovia fasciculata.
fastigiatum, *Willd. Enum. Hort. Berol.* 235.—Mexic.
Fendleri, A. Gray, in Am. Journ. Sc. Ser. II. xxii. (1856) 285 = tuberosum.
Fendleri, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 130.—Panama.
fernandezianum, *Phil. in Linnaea*, xxix. (1857–58) 23.—Chili.
ferocissimum, *Lindl. in Mitch. Three Exped.* ii. 58.—Austral.
ferox, Burm. f. Fl. Ind. 56 = xanthocarpum?
ferox, Jungh. ex Miq. Fl. Ind. Bat. ii. 650 = indicum.
ferox, *Linn. Sp. Pl.* ed. II. 267.—As. trop.
ferox, Mill. ex Dun. in DC. Prod. xiii. 1. 299 = tomentosum.
ferrugineum, Jacq. Hort. Schoenb. iii. 46. t. 334 = torvum.
Feuillei, *Dun. in DC. Prod.* xiii. 1. 70.—Chili.
ficifolium, Ortega, Hort. Matr. Dec. 116 = torvum.
ficifolium, Pav. ex Dun. in DC. Prod. xiii. 1. 350 = Zuccagnianum.
Fieldingi, *Moric. ex Dun. l. c.* 188.—Bras.
filiforme, Pav. ex Dun. l. c. 431 = Saracha alata.
filiforme, *Ruiz & Pav. Fl. Per.* ii. 31. t. 159.—Peruv.
filipes, Dun. in DC. Prod. xiii. 1. 138 = rivulare.
fistulosum, Dun. ex Poir. Encyc. Suppl. iii. 749 = nigrum.
flaccidum, *Vell. Fl. Flum.* 87; ii. t. 115.—Bras.
flagellare, *Sendt. in Mart. Fl. Bras.* x. 68.—Bras.
flagelliferum, *Baker, in Journ. Linn. Soc.* xx. (1883) 213.—Madag.
flagrans, Tenore, Sem. Ort. Bot. Nap. (1839) 12; et in Ann. Sc. Nat. Sér. II. xiii. (1840) 381 = inaequale.
flavescens, Dun. Solan. Syn. 42 = hirtum.
flavidum, Torr. in Ann. Lyc. N. York, ii. (1828) 227 = elaeagnifolium.
flavum, Kit. in Schult. Oestr. Fl. ed. II. ii. 394 = nigrum.
flexicaule, *Benth. Bot. Voy. Sulph.* 141.—Am. austr.
flexuosum, Banks, ex Dun. in DC. Prod. xiii. 1. 170 = retrofractum.
flexuosum, Heyne, ex Nees, in Trans. Linn. Soc. xvii. (1834) 42 = bigeminatum.
flexuosum, Pav. ex Dun. in DC. Prod. xiii. 1. 178 = campylocladum.
flexuosum, Vahl, Eclog. Am. i. 22 = lanceaefolium.
flexuosum, *Willd. ex Roem. & Schult. Syst.* iv. 662.—Ins. S. Doming.
floccosum, Dun. in DC. Prod. xiii. 1. 133 = pachimatium.
floccosum, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 141 = Hartwegi.
floccosum, *Zipp. ex Span. in Linnaea*, xv. (1841) 337, nomen.—Ins. Timor.
floribundum, Dun. in DC. Prod. xiii. 1. 92 = congestiflorum.
floribundum, *Sendt. in Mart. Fl. Bras.* x. 63.—Bras.
floridanum, *Shuttl. ex Dun. in DC. Prod.* xiii. 1. 306 (sp. dub.).—Florida.
fluminense, Steud. Nom. ed. II. ii. 602 = Cyphomandra sciadostylis.
foetidum, H. B. & K. Nov. Gen. et Sp. iii. 33 = S. Caavurana.
foetidum, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 287 = Rottboellianum.
foetidum, *Ruiz & Pav. Fl. Per.* ii. 39. t. 165. f. b.—Peruv.
foliosum, Link, in Buch, Beschr. Canar. Ins. 144 = Lycopersicum Humboldtii.
Fontanesianum, *Dun. in Poir. Encyc. Suppl.* iii. 777.—Bras.
Fontanesianum, Schrank, Pl. Rar. Hort. Monac. t. 18 = nigrum, tectum.
formosum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 35.—Venezuela.
formosum, Weinm. Cat. Hort. Dorpat. (1810) 145 = sisymbrifolium.

SOLANUM:—
Forskålii, *Dun. Hist. Solan.* 237.—Arab.
Forsteri, *Seem. in Journ. Bot.* i. (1863) 207.—Ins. Pacif.
fossarum, *Dun. in DC. Prod.* xiii. 1. 145.—Bras.
fragile, Wedd. Chlor. And. ii. 105 = atriplicifolium.
fragile, *Wright, ex Griseb. Cat. Pl. Cub.* 189.—Cuba.
fragrans, Hook. Bot. Mag. t. 3684 = Cyphomandra fragrans.
fraxinifolium, *Dun. in DC. Prod.* xiii. 1. 39.—N. Granat.
fructu-tecto, Cav. Ic. iv. 5 = tectum.
frutescens, *A. Br. & Bouché, in Ind. Sem. Hort. Berol.* (1853) *App.* 9.—Venezuela.
fruticosum, Mill. Gard. Dict. ed. VIII. n. 18 = bahamense.
fugax, Bert. ex Dun. in DC. Prod. xiii. 1. 170 = sylvaticum.
fugax, *Jacq. Coll.* iv. 128.—Am. austr.
fultum, *Schrank, ex Sendt. in Mart. Fl. Bras.* x. 49.—Bras.
fulvum, *Hiern, ex Warm. in Kjoeb. Vidensk. Meddel.* (1877–78) 49.—Bras.
furcatum, *Dun. in Poir. Encyc. Suppl.* iii. 750.—Peruv.
furfuraceum, *R. Br. Prod.* 446.—Austral.
fuscatum, *Linn. Sp. Pl.* ed. II. 268.—Am. centr.; Ind. occ.
fuscatum, *Roem. & Schult. Syst.* iv. 872 (Index).—Am. austr.
fuscum, Heyne, ex Wall. Cat. n. 2622 B = trilobatum.
galeatum, *André, in Rev. Hortic.* (1862) 331.—Austral.
Galeotti, *Dun. in DC. Prod.* xiii. 1. 82.—Mexic.
Gardneri, *Sendt. in Mart. Fl. Bras.* x. 69.—Bras.
Gaudichaudi, *Dun. in DC. Prod.* xiii. 1. 60.—Chili.
Gayanum, *Phil. f. Cat. Pl. Vasc. Chil.* 228.—Chili.
gemellum, Mart. ex Sendt. in Fl. Bras. x. 28 = didymum.
geminatum, *Vahl, Eclog. Am.* i. 21.—Guian.
geminiflorum, Cham. & Schlecht. in Linnaea, v. (1830) 112 = Schlechtendalianum.
geminiflorum, Drège, ex Dun. in DC. Prod. xiii. 1. 289 = didymanthum.
geminiflorum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 142.—Mexic.
geminifolium, Schlecht. in Linnaea, viii. (1833) 255 = Schlechtendalianum.
geminifolium, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 121.—Afr. trop.
geniculatum, *Drège, ex Dun. in DC. Prod.* xiii. 1. 105.—Afr. austr.
genistoides, Dun. l. c. 85 = umbelliferum.
Germaini, *Phil. in Linnaea*, xxix. (1857–58) 23.—Chili.
giftbergense, *Dun. in DC. Prod.* xiii. 1. 288.—Afr. austr.
giganteum, *Jacq. Coll.* iv. 125.—Ind. or.
Gilo, *Raddi, in Mem. Soc. Ital. Moden.* xviii. *Fis.* (1820) 410.—Bras.
Gilo, Requien, ex Dun. in DC. Prod. xiii. 1. 360 = integrifolium.
glabellum, Dun. l. c. 142 = indigoferum.
glaberrimum, Dun. in Poir. Encyc. Suppl. iii. 747 = aviculare.
glabratum, *Dun. Solan. Syn.* 49.—Arabia.
glabrescens, Schrank, ex Sendt. in Mart. Fl. Bras. x. 39 = rufescens.
glabrum, *Dun. in DC. Prod.* xiii. 1. 102.—Mexic.
glandulosum, Banks, ex Dun. in DC. Prod. xiii. 1. 166 = guianense, *Dun.*
glandulosum, *Ruiz & Pav. Fl. Per.* ii. 35. t. 167.—Peruv.
glandulosum, Sendt. in Mart. Fl. Bras. x. 52 = japurense.
glaucescens, Bacle, ex Dun. in DC. Prod. xiii. 1. 100 = glaucum.
glaucescens, *Zuccar. in Abh. Akad. Muench.* ii. (1831–36) 325.—Mexic.
glaucophyllum, Desf. Cat. Pl. Hort. Par. ed. III. 396 = glaucum.
glaucum, *Dun. in DC. Prod.* xiii. 1. 100; *Bertol. Misc. Bot.* xii. 46.—Bras.
Glaziovii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 47.—Bras.
globiferum, *Dun. in DC. Prod.* xiii. 1. 251.—Ins. Martinic.
glomuliflorum, *Sendt. in Mart. Fl. Bras.* x. 24.—Bras.

SOLANUM:—
glutinosum, *Dun. in Poir. Encyc. Suppl.* iii. 79.—Mexic.
gnaphalocarpum, *Vell. Fl. Flum.* 82; ii. t. 91.—Bras.
gnaphaloides, *Pers. Syn.* i. 223.—Peruv.
gomphodes, *Dun. in DC. Prod.* xiii. 1. 339.—Bras.
gomphoidellum, *Moric. ex Dun. l. c.* 215.—Bras.
gonocladum, *Dun. l. c.* 93.—Bolivia.
gossypinum, *Dun. l. c.* 208.—Bras.
Gouakai, Dun. l. c. 177 = denticulatum.
Goudoti, *Dun. l. c.* 158.—Peruv.
gracile, Moric. ex Dun. l. c. 238 = litocladum.
gracile, *Otto, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 673; *Dun. l. c.* 54.—Am. trop.
gracile, Sendt. in Mart. Fl. Bras. x. 13 = Endlicheri.
graciliflorum, *Dun. ex Poir. Encyc. Suppl.* iii. 763.—Java.
gracilipes, *Decne. in Jacquem. Voy. Bot.* 113. t. 119.—Ind. bor. occ.
gracillimum, *Sendt. in Mart. Fl. Bras.* x. 36.—Bras.
grandiflorum, Desf. Hist. Arb. i. 169 = macrophyllum.
grandiflorum, Hort. ex Steud. Nom. ed. I. 791 = bonariense.
grandiflorum, *Ruiz & Pav. Fl. Per.* ii. 35. t. 168.—Peruv.
grandiflorum, Sieber, ex Dun. in DC. Prod. xiii. 1. 325 = Richardi.
grandiflorum, Vahl, ex Dun. l. c. 313 = dasyphyllum.
granuloso-leprosum, *Dun. l. c.* 115.—Mexic.
graveolens, *Bunbury, in Proc. Linn. Soc.* i. (1841) 110.—Bras.
greviaefolium, *Dun. in DC. Prod.* xiii. 1. 119.—Bras.
grossedentatum, *A. Rich. Tent. Fl. Abyss.* ii. 101.—Abyss.
guaraniticum, *A. St. Hil. in Mém. Mus. Par.* xii. (1825) 321.—Bras.
guianense, *Dun. in DC. Prod.* xiii. 1. 166.—Guian.
guianense, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
guiencense, Hill, Veg. Syst. ix. 34, sphalm. = *guineense*, Linn.
guineense, Lam. Illustr. ii. 18 = nigrum.
guineense, Linn. Sp. Pl. 184 = aggregatum.
Gula, Buch.-Ham. in Trans. Linn. Soc. xiv. (1825) 267 = xanthocarpum.
Gussonei, Dun. in DC. Prod. xiii. 1. 380 = cestrophyllum?
haematocladum, *Dun. l. c.* 230.—Bolivia.
hainanense, *Hance, in Journ. Bot.* vi. (1868) 331.—China.
haitense, *Dun. in DC. Prod.* xiii. 1. 167.—Ins. S. Doming.
Hartwegi, *Benth. Pl. Hartw.* 68.—Mexic.
hastatum, *Mart. ex Sendt. in Mart. Fl. Bras.* x. 61.—Bras.
hastifolium, *Hochst. ex Dun. in DC. Prod.* xiii. 1. 284.—Afr. trop.
havanense, *Jacq. Enum. Pl. Carib.* 15; *Linn. Mant.* i. 47.—Am. trop.
havanense, *Vell. Fl. Flum.* 82; ii. t. 94.—Bras.
hebecarpum, *Salzm. ex Sendt. in Fl. Bras.* x. 41.—Bras.
hebephorum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 7.—Peruv.
hedjazense, *Dun. in DC. Prod.* xiii. 1. 292.—Arab.
heliotropiifolium, Dun. l. c. 117 = Sellovianum.
hendecandraefolium, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
Herbertianum, *Paxt. Mag. Bot.* v. (1838) 269.—Hab.?
Hermanni, Dun. Hist. Solan. 212 = sodomeum.
Hernandesii, *Dun. in Poir. Encyc. Suppl.* iii. 771.—Mexic.
heteracanthum, Dun. l. c. 773 = S. Melongena.
heterandrum, Juss. in Ann. Mus. Par. iii. (1804) 120, in textu = cornutum.
heterandrum, *Pursh, Fl. Am. Sept.* i. 156; ii. 731. t. 7.—Am. bor.
Heteranthera, Willd. ex Roem. & Schult. Syst. iv. 663 = cotopaxense.
heteroclitum, *Sendt. in Flora*, xxix. (1846) 193.—Guatemala.
heterodoxum, Andrieux, ex Dun. in DC. Prod. xiii. 1. 330 = chrysacanthum.
heterodoxum, *Dun. Hist. Solan.* 235. t. 25.—Mexic.
heterogonum, Hort. ex Dun. in DC. Prod. xiii. 1. 52 = nigrum.
heteromorphum, *Dun. l. c.* 80.—Bras.
heterophyllum, Balb. ex Dun. l. c. 260 = torvum.

SOLANUM:—
heterophyllum, Lam. Illustr. ii. 22 = S. Juripeba.
heterophyllum, Willd. ex Roem. & Schult. Syst. iv. 664 = Cyphomandra caracasana.
heterotricum, Dun. Hist. Solan. 192. t. 20 = jamaicense.
Heudeloti, *Dun. in DC. Prod.* xiii. i. 205.—Afr. trop
hexandrum, Steud. Nom. ed. II. i. 87, sphalm. = rostratum.
hexandrum, Vell. Fl. Flum. 88; ii. t. 122 = maroniense.
Heynii, Roem. & Schult. Syst. iv. 669 = indicum.
hierochunticum, Dun. in DC. Prod. xiii. i. 369 = incanum.
Hildebrandtii, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1874) 8.—Afr. trop.
himalense, Dun. in DC. Prod. xiii. i. 300 = indicum.
Hindsianum, Benth. Bot. Voy. Sulph. 39 = elaeagnifolium.
hirsutum, Balb. ex Dun. in DC. Prod. xiii. i. 198 = cordifolium.
hirsutum, Dun. Hist. Solan. 158 = nigrum.
hirsutum, Herb. Peurari, ex Dun. in DC. Prod. xiii. i. 252 = stramonifolium.
hirsutum, Nutt. in Journ. Acad. Philad. vii. (1834) 109 = carolinense.
hirsutum, Roxb. Hort. Beng. 17; Fl. Ind. i. 571 = ferox.
hirtulum, *Steud. ex Dun. in DC. Prod.* xiii. i. 44.—Abyss.
hirtum, *Vahl, Symb. Bot.* ii. 40.—Ins. Trinitat.
hispidum, *Pers. Syn.* i. 228.—Peruv.
Hoffmanseggii, *Sendt. in Mart. Fl. Bras.* x. 112.—Bras.
Hohenackeri, Heurck & Muell. Arg. in Heurck, Obs. Bot. 87 = Wightii.
Hookerianum, Spreng. f. Tent. Suppl. Syst. 9 = havanense, *Jacq.*
horridissimum, Hort. Par. ex Sendt. in Mart. Fl. Bras. x. 59 = aculeatissimum, *Jacq.*
horridum, *Dun. in Poir. Encyc. Suppl.* iii. 763.—Austral.; Timor.
horridum, Salisb. Prod. 134 = aculeatissimum, *Jacq.*
Hostmanni, *Dun. in DC. Prod.* xiii. i. 184.—Guian.
Houstonii, *Dun. Hist. Solan.* 243.—Mexic.
Hovei, *Dun. in DC. Prod.* xiii. i. 311.—Ind. or.
Humboldti, *Dun. l. c.* 216.—Mexic.
Humboldtii, Willd. Hort. Berol. i. 27 = Lycopersicum esculentum.
humifusum, *Dun. in DC. Prod.* xiii. i. 237.—Bras.
humile, Bernh. ex Willd. Enum. Hort. Berol. 236 = nigrum.
humile, Lam. Illustr. ii. 23 = rigescens.
humile, Salisb. Prod. 134 = nigrum.
hybridum, Dun. ex Griseb. Fl. Brit. W. Ind. 441 = inclusum.
hybridum, *Jacq. Hort. Vindob.* ii. 51. t. 113.—Afr. trop.
hyemale, Salisb. Prod. 133 = S. Pseudo-capsicum.
hygrophilum, *Schlecht. in Linnaea*, viii. (1833) 254.—Mexic.
hyporhodium, *A. Br. & Bouché, in Ind. Sem. Hort. Berol.* (1853) *App.* 10.—Austral.
hystricinum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
Hystrix, *R. Br. Prod.* 446.—Austral.
Hystrix, Dun. in DC. Prod. xiii. i. 295 = armatum.
igneum, *Linn. Sp. Pl.* ed. II. 270.—Ind. occ.
ilicifolium, *Dun. in DC. Prod.* xiii. i. 190.—Bras.
imamense, *Dun. l. c.* 85.—Madag.
immane, Hance, ex Walp. Ann. iii. 165 = ferox.
immite, Dun. in DC. Prod. xiii. i. 32 = tuberosum.
impediens, *Weinm. in Syll. Ratisb.* ii. (1828) 162.—Bras.
inaequale, *Hornem. Suppl. Hort. Hafn.* 26.—Hab.?
inaequale, Presl, Bot. Bemerk. 101 = lanceaefolium.
inaequale, *Vell. Fl. Flum.* 87; ii. t. 116.—Bras.
inamoenum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 228 = tetrandrum.
incanum, Kit. ex Schult. Oestr. Fl. ed. II. i. 395 = nigrum.
incanum, *Linn. Sp. Pl.* 188; *Forsk. Fl. Aegypt. Arab.* 45.—Afr. occ.; As. or.
incanum, Pav. ex Dun. in DC. Prod. xiii. i. 369 = pyriforme.
incanum, Ruiz & Pav. Fl. Per. ii. 40 = albidum.
incanum, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 39.—N. Guin.

SOLANUM:—
incarceratum, *Ruiz & Pav. Fl. Per.* ii. 40. t. 176.—Am. trop.
incertum, Dun. Hist. Solan. 155 = nigrum.
incisum, *Griseb. in Goett. Abh.* xxiv. (1879) 251.—Reg. Argent.
inclusum, *Griseb. Fl. Brit. W. Ind.* 441.—Ind. occ.
incompletum, *Dun. in DC. Prod.* xiii. i. 311.—Ins. Pacif.
incurvum, *Ruiz & Pav. Fl. Per.* ii. 34. t. 164.—Peruv.
indecorum, *A. Rich. Fl. Cub. Fanerog.* ii. 121.—Cuba.
indicum, *Linn. Sp. Pl.* 187.—Reg. trop.
indicum, Roxb. Hort. Beng. 17; Fl. Ind. i. 570 = S. Melongena.
indigoferum, *A. St. Hil. in Mer. & De Lens, Dict. Mat. Méd.* vi. 416.—Bras.
inflatum, Hornem. Hort. Hafn. i. 221 = sisymbrifolium.
inodorum, *Vell. Fl. Flum.* 85; ii. t. 107.—Bras.
inops, *Dun. in DC. Prod.* xiii. i. 55.—Mexic.
insanum, J. B. Fisch. Livl. 202; ex Ledeb. Fl. Ross. iii. 189 (Quid?).—Ross.
insanum, Linn. Mant. i. 46 = S. Melongena.
insidiosum, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 79.—Bras.
insigne, *Lowe, in Journ. Hort. Soc.* N. S. i. (1866) 178; *Man. Fl. Mad.* ii. 84.—Ins. Mader.
insulanum, *Dun. in DC. Prod.* xiii. i. 373.—Ins. S. Doming.
integerrimum, *Dun. Hist. Solan.* 236.—Bras.
integrifolium, *Poir. in Lam. Encyc.* iv. 301.—As. et Afr.
intermedium, Dun. in DC. Prod. xiii. i. 311 = Hovei.
intermedium, *Sendt. in Mart. Fl. Bras.* x. 22.—Bras.
involucratum, *Blume, Bijdr.* 701.—Malaya
involucratum, Kurz, For. Fl. Brit. Burm. ii. 224 = barbisetum.
iodotrichum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 81.—Bolivia.
Ipomoea, *Sendt. in Mart. Fl. Bras.* x. 50.—Bras.
ipomoeum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
Isabelli, *Dun. in DC. Prod.* xiii. i. 153.—Reg. Argent.
isodynamum, *Sendt. in Mart. Fl. Bras.* x. 33.—Bras.
Izquierdi, *Phil. in Anal. Univ. Chil.* (1873) 522.—Chili.
Jacquemonti, Dun. in DC. Prod. xiii i. 287 = gracilipes.
Jacquini, Miq. Fl. Ind. Bat. ii. 651 = sarmentosum.
Jacquini, Willd. Sp. Pl. i. 1041 = xanthocarpum.
jamaicense, *Mill. Gard. Dict.* ed. VIII. 17; *Sw. Fl. Ind. Occ.* i. 454.—Am. trop.
Jamesii, *Torr. in Ann. Lyc. N. York*, ii. (1828) 227.—Am. bor. occ.
japonicum, Hort. ex Carr. in Rev. Hortic. (1868) 420 = Rantonnetii.
japurense, *Dun. in DC. Prod.* xiii. i. 174.—Bras.
jasminifolium, *Sendt. in Mart. Fl. Bras.* x. 13.—Bras.
jasminoides, *Paxt. Mag. Bot.* viii. (1841) t. 5.—Bras.
jatrophaefolium, *Dun. in DC. Prod.* xiii. i. 246.—Bras.
javanicum, Dun. l. c. 176 = biflorum.
javitense, H. B. & K. Nov. Gen. et Sp. iii. 42 = monachophyllum.
jubatum, Dun. in DC. Prod. xiii. i. 132 = cernuum.
jubatum, Willd. ex Roem. & Schult. Syst. iv. 669 = cyananthum.
Jubeba, Vell. Fl. Flum 89; ii. t. 124 = paniculatum, insidiosum.
Juciri, Mart. ex Sendt. in Mart. Fl. Bras. x. 110 = alternato-pinnatum.
judaicum, Bess. Fl. Galic. i. 183 = nigrum.
judaicum, Hackel, ex Dun. in DC. Prod. xiii. i. 53 = Tauschii
juglandifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 6.—N. Granat.
julocrotonoides, *Klotzsch, ex Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
Junghuhnii, Miq. Fl. Ind. Bat. ii. 649 = indicum.
Juripeba, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107.—Am. trop.
Juripeba, Vell. ex Steud. Nom. ed. II. ii. 603 = pycnanthemum.
Jussiaei, *Dun. in Poir. Encyc. Suppl.* iii. 767.—Bras.
kaitisis, *Dun. in DC. Prod.* xiii. i. 157.—Ind. or.

SOLANUM:—
Karstenii, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1853) *App.* 10.—Venezuela.
Karstenii, *Dun. in DC. Prod.* xiii. i. 151.—N. Granat.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 234.—Reg. Himal.
Kieseritzkii, *C. A. Mey. Verz. Pfl. Cauc.* 113.—Reg. Cauc.
Kitaibeli, Schult. Oestr. Fl. ed. II. i. 395 = nigrum.
Krauseanum, *Phil. in Linnaea*, xxxiii. (1864–65) 204.—Chili.
Kraussianum, Buching. ex Krauss, in Flora, xxvii. (1844) 831 = nigrum.
laciniatum, Ait. Hort. Kew. ed. I. i. 247 = aviculare.
laciniatum, Ruiz & Pav. Fl. Per. ii. 40 = nemorense.
lacteum, *Vell. Fl. Flum.* 82; ii. t. 93.—Bras.
lacunarium, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 18.—Austral.
laetum, *Kunze, in Linnaea*, xvi. (1842) 315.—Hab.?
laetum, *Miq. Stirp. Surin. Sel.* 135.—Guian.
laeve, *Dun. ex Poir. Encyc. Suppl.* iii. 751.—Ind. or.
laevigatum, H. B. & K. Nov. Gen. et Sp. iii. 32 = levigatum.
lagenarium, *Dun. in DC. Prod.* xiii. i. 368.—Hab.?
Lalandi, *Dun. l. c.* 210.—Bras.
Lamberti, *Dun. l. c.* 268.—Mexic.
lanaevestitum, *Dun. l. c.* 273.—Bras.
lanatum, *Dun. Hist. Solan.* 205.—Peruv.
lanceaefolium, *Jacq. Coll.* ii. 286.—Am. trop.
lanceaefolium, Salisb. Prod. 134 = bahamense.
lanceolatum, *Cav. Ic.* iii. 23. t. 245.—Mexic.
lanceolatum, Ruiz & Pav. Fl. Per. ii. 33 = patulum.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864–65) 204.—Chili.
laniflorum, *Sendt. in Mart. Fl. Bras.* x. 69.—Bras.
lanigerum, *Dun. in DC. Prod.* xiii. i. 273.—Bras.
Lantana, *Sendt. in Mart. Fl. Bras.* x. 36.—Bras.
lantanum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
lanuginosum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 25.—Ecuador.
lasiantherum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 48.—Peruv.
lasianthum, *Dun. in DC. Prod.* xiii. i. 204.—N. Granat.
lasiocarpum, Dun. Hist. Solan. 222; Blume, Bijdr. 701 = ferox.
lasiophyllum, *Dun. in Poir. Encyc. Suppl.* iii. 764.—Austral.
lasiophyllum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 25.—N. Granat.
lasiopodium, *Dun. in DC. Prod.* xiii. i. 679.—N. Granat.
latiflorum, Dun. l. c. 388 = Cyphomandra calycina.
latifolium, Dun. l. c. 353 = repandum.
latifolium, *Poir. in Lam. Encyc.* iv. 303.—Am. austr.
laurifolium, Deless. ex Dun. in DC. Prod. xiii. i. 80 = domingense.
laurifolium, Dun. Hist. Solan. 165. t. 8 = Riedlei.
laurifolium, Hort. ex Dun. in DC. Prod. xiii. i. 373 = Linkii.
laurifolium, *Linn. f. Suppl.* 148.—Am. austr.
laurifolium, Mill. Gard. Dict. ed. VIII. n. 20 = subinerme.
laurifolium, Sieber, ex Dun. in DC. Prod. xiii. i. 115 = verbascifolium.
laurinum, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
laurinum, *Dun. in DC. Prod.* xiii. i. 154.—Bras.
laxiflorum, Dun. l. c. 234 = viridipes.
laxiflorum, *Sendt. in Mart. Fl. Bras.* x. 21.—Bras.
laxum, Dun. in DC. Prod. xiii. i. 124 = megalochiton.
laxum, Royle, Illustr. Bot. Himal. 279 = S. Dulcamara.
laxum, *Spreng. Syst.* i. 684.—Reg. Argent.
Leiboldianum, Dun. in DC. Prod. xiii. i. 314 = vellereum.
leiophyllum, *Benth. Pl. Hartw.* 146.—Peruv.
leiophyllum, Dun. in DC. Prod. xiii. i. 141 = inaequale.
lentum, *Cav. Ic.* iv. 4. t. 308.—Am. trop.
lentum, Kunth, ex Dun. in DC. Prod. xiii. i. 173 = campylocladum.
leontopodium, *Sendt. in Mart. Fl. Bras.* x. 38.—Bras.
lepidotum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 17.—N. Granat.
leprosum, Ortega, Hort. Matr. Dec. 115 = elaeagnifolium.
leptanthum, *Dun. ex Poir. Encyc. Suppl.* iii. 747.—Am. trop.
leptocaulon, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 40.—Bolivia.

SOLANUM:—
leptophyllum, F. Muell. Fragm. ii. 164 = parvifolium.
leptopodum, Dun. in DC. Prod. xiii. 1. 411 = Bassovia leptopoda.
leptopodum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 57.—Peruv.
leptostachys, *Dun. in DC. Prod.* xiii. 1. 306.—Bras.
leucocarpon, *Dun. in Poir. Encyc. Suppl.* iii. 756.—Am. trop.
leucodendron, *Sendt. in Mart. Fl. Bras.* x. 43.—Bras.
leucophaeum, *Dun. in DC. Prod.* xiii. 1. 290.—Afr. austr.
leucophyllum, Dun. l. c. 126 = leontopodium.
leucosporum, *Dun. l. c.* 99.—Guian.
levigatum, Dun. Solan. Syn. 20 = triste.
Leycesterianum, *Savi, ex Delile, Hort. Monsp. Ic. Pict.* vii. t. 682.—Mexic.
Lhotskyanum, *Dun. in DC. Prod.* xiii. 1. 248.—Bras.
Lichtensteinii, *Willd. Enum. Hort. Berol.* 238.—Afr. austr.
ligustrinum, Lodd. Bot. Cab. t. 1963 = crispum.
Lima, Banks, ex Dun. in DC. Prod. xiii. 1. 107 = S. Radula.
Lindheimerianum, Scheele, in Linnaea, xxi. (1848) 766 = triquetrum.
lineatum, *Ruiz & Pav. Fl. Per.* ii. 31.—Peruv.
linifolium, Dun. in DC. Prod. xiii. 1. 152 = spissifolium.
Linkianum, *Roem. & Schult. Syst.* iv. 601.—Bras.
Linkii, *G. Don, Gen. Syst.* iv. 408.—Hab. ?
Linsecumii, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 6 = Chamaesaracha sordida.
lithophilum, F. Muell. in Linnaea, xxv. (1852) 434 = ellipticum.
litocladum, *Dun. in DC. Prod.* xiii. 1. 238.—Bras.
littorale, Raab, in Flora, ii. (1819) 414 = S. Dulcamara.
lividum, *Link, Enum. Hort. Berol.* i. 188.—Madag.
lividum, Willd. ex Dun. in DC. Prod. xiii. 1. 310 = indicum.
Lobelii, Tenore, Ind. Sem. Hort. Neap. (1851); et in Atti R. Ist. Incorr. Nap. viii. (1855) 328 = integrifolium.
longepedunculatum, Bert. ex Dun. in DC. Prod. xiii. 1. 171 = retrofractum.
longiflorum, *Vahl, Eclog. Am.* i. 20.—Guian.
longifolium, *Dun. Hist. Solan.* 163.—Hab. ?
longifolium, Pav. ex Dun. in DC. Prod. xiii. 1. 186 = ensifolium.
longipes, *Dun. l. c.* 85.—Afr. austr.
longum, Roxb. Hort. Beng. 16; Fl. Ind. i. 567 = S. Melongena.
loxense, *Humb. & Bonpl. ex Dun. Solan. Syn.* 16.—Ecuador.
lucidum, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 137 = brachystachys.
lucidum, Moric. Pl. Nouv. Am. 26. t. 18 = Bassovia lucida.
ludovicianum, *Rafin. Fl. Ludov.* 45 (Quid ?).—Am. bor.
Lundianum, *Dun. in DC. Prod.* xiii. 1. 270.—Bras.
luridum, Dun. l. c. 88 = flaccidum.
luridum, *Pav. ex Dun. l. c.* 113.—Mexic.
luridum, Salisb. Prod. 134 = Lycopersicum esculentum.
lusitanicum, *Dun. Hist. Solan.* 238.—Lusitan.
luteiflorum, Dun. in DC. Prod. xiii. 1. 64 = Physalis lobata.
luteoalbum, *Pers. Syn.* i. 221; *Dun. Hist. Solan.* 168.—Peruv.
luteoliflorum, A. Gray, Syn. Fl. N. Am. ii. 1. 233, sphalm. = *luteiflorum*.
luteo-virescens, Dun. in DC. Prod. xiii. 1. 238 = arcuatum.
luteo-virescens, C. C. Gmel. Fl. Bad. iv. 177 = nigrum.
luteum, Mill. Gard. Dict. ed. VIII. n. 3 = nigrum.
lycioides, Domb. ex Dun. in DC. Prod. xiii. 1. 162 = Dombeyi.
lycioides, *Linn. Mant.* i. 46.—Peruv.
lycioides, Ruiz & Pav. Fl. Per. ii. 41. t. 177 = candicans.
lycocarpum, *A. St. Hil. Voy. Distr. Diam.* i. 11. 333.—Bras.
lycoctonum, Hort. Panorm. ex Dun. in DC. Prod. xiii. 1. 103 = Cervantesii.
lycopersicoides, *Dun. l. c.* 38.—Bolivia.
Lycopersicum, Blanco, Fl. Filip. ed. I. 134 = Lycopersicum philippinarum.
Lycopersicum, Linn. Sp. Pl. 185 = Lycopersicum esculentum.

SOLANUM:—
lyratum, Thunb. Fl. Jap. 92 = S. Dulcamara.
lysimachioides, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 257.—Ind. or.
macaonense, Dun. in DC. Prod. xiii. 1. 264 = torvum.
Maccai, Bert. ex Dun. l. c. 260 = torvum.
Maccai, *Dun. in Poir. Encyc. Suppl.* iii. 771.—Guian.
macilentum, *A. Rich. Tent. Fl. Abyss.* ii. 105.—Abyss.
macracanthum, *A. Rich. l. c.* 106.—Abyss.
macranthemum, Roem. & Schult. Syst. iv. 596, sphalm. = seq.
macrantherum, *Dun. Solan. Syn.* 16.—Am. trop.
macranthum, Bert. ex Dun. in DC. Prod. xiii. 1. 167 = haitense.
macranthum, *Dun. Solan. Syn.* 43.—Bras.
macranthum, Hook. Bot. Mag. t. 4138 = maroniense.
macranthum, Hort. ex Carr. in Rev. Hortic. (1867) 132, cum ic. = Wrightii.
macranthum, Mart. & Gal. in Bull. Acad. Brux. xii. 1. (1845) 143 = porphyranthum.
macrobotryon, Dun. in DC. Prod. xiii. 1. 277 = pelliceum.
macrocalyx, *Dun. l. c.* 337.—Bras.
macrocarpon, *Linn. Mant.* ii. 205.—Ins. Mascar.
macrocarpon, *Molina, Sagg. Chile*, ed. II. 282.—Chili.
macrocarpon, Pav. ex Dun. in DC. Prod. xiii. 1. 352 = quitoënse.
macrodon, Wall. Cat. n. 2621 = pachypetalon.
macronema, *Sendt. in Mart. Fl. Bras.* x. 88.—Bras.
macrophyllum, *Hort. ex Dun. Hist. Solan.* 199. t. 17.—Mexic.
macropus, Dun. in DC. Prod. xiii. 1. 134 = bullatum.
madagascariense, *Dun. l. c.* 99.—Madag.
Maglia, *Schlecht. Hort. Hal.* 6.—Chili.
magnifolium, *F. Muell. Fragm.* vi. 27.—Austral.
malacoxylon, *Sendt. in Mart. Fl. Bras.* x. 52.—Reg. Argent.
mammiforme, Robin, Voy. Louis, iii. 398 (Quid ?).—Am. bor.
mammosum, Herb. ex Dun. in DC. Prod. xiii. 1. 260 = torvum.
mammosum, *Linn. Sp. Pl.* 187.—Am. trop.
mammosum, Lour. Fl. Cochinch. 162 = stramonifolium.
mammosum, Pav. ex Dun. in DC. Prod. xiii. 1. 140 = caudatum.
Mandonis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 78.—Bolivia.
Manoelii, Moric. Pl. Nouv. Am. 27 = paniculatum.
marginatum, *Linn. f. Suppl.* 147.—Afr. trop.
Mariannae, *Dun. in DC. Prod.* xiii. 1. 307.—Bras.
maritimum, *Meyen, ex Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 384.—Chili.
maroniense, *Poit. in Ann. Soc. Hort. Par.* vii. (1830) 155.—Bras.
Marsilianum, Tenore, Cat. Orto Bot. Nap. (1845) 96 = inaequale.
Martensii, *Dun. in DC. Prod.* xiii. 1. 140.—Mexic.
Martii, Dun. l. c. 399 = Cyphomandra Fraxinella.
Martii, *Sendt. in Mart. Fl. Bras.* x. 41.—Bras.
Martinisi, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 73.—Peruv.
Martiusianum, Steud. Nom. ed. II. ii. 604 = aculeatissimum, *Jacq.*
mauritianum, Blanco, Fl. Filip. ed. I. 134 = verbascifolium.
mauritianum, Scop. Delic. Insub. iii. 16 = auriculatum.
mauritianum, *Willd. ex Roth, Nov. Sp. Pl.* 131 = sisymbrifolium.
megalochiton, *Mart. in Flora*, xxi. (1838) 11. *Beibl.* 63.—Bras.
megalochitonium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
megalonychinum, St. Lag. l. c. 135 = seq.
megalonychum, St. Lag. l. c. viii. (1881) 161, 203 = seq.
megalonyx, *Sendt. in Mart. Fl. Bras.* x. 92.—Bras.
melanandrum, *Moric. ex Dun. in DC. Prod.* xiii. 1. 241.—Bras.
melancholicum, *Salzm. ex Dun. l. c.* 200.—Bras.
melaniferum, Moric. ex Dun. l. c. 43 = muricatum.
melanocarpum, Dun. l. c. 355 = S. Melongena.
melanocarpum, Hort. ex Rev. Hortic. (1877) 291 = Wallisii.
melanocerasum, Willd. Enum. Hort. Berol. 237 = nigrum.
melanospermum, *F. Muell. Fragm.* ii. 163.—Austral.

SOLANUM:—
melanoxylon, *Link, Enum. Hort. Berol.* i. 188.—Hab. ?
Melongena, *Linn. Sp. Pl.* 186.—Geront. trop.
Melongena, Wall. Cat. n. 2628 = sarmentosum.
melongenum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Melongena.
membranaceum, Thw. Enum. Pl. Zeyl. 216 = bigeminatum.
membranaceum, Wall. Cat. n. 2625 = laeve.
membranaceum, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 231 = denticulatum.
membranifolium, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 77.—Hab. ?
memphiticum, J. F. Gmel. Syst. 385; Mart. Hort. Erlang. (1814) 63 = nigrum.
mendax, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 61.—Peruv.
mendocianum, Gill. ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 387 = atriplicifolium.
mendocinum, *Phil. in Anal. Univ. Chil.* (1862) 11. 403.—Chili.
Menziesii, *Dun. in DC. Prod.* xiii. 1. 159.—Calif.
metrobotryon, *Dun. l. c.* 276.—Bras.
mexicanum, *Dun. ex Poir. Encyc. Suppl.* iii. 770.—Mexic.
mexicanum, Hill, Veg. Syst. ix. 39 = modestum ?
mexicanum, Willd. ex Roem. & Schult. Syst. iv. 663 = modestum.
micracanthos, *Lam. Illustr.* ii. 24.—Am. trop.
micranthum, D. Dietr. Syn. Pl. i. 710 = micracanthos.
micranthum, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 430 = Saracha Zuccagniana.
micranthum, Willd. ex Roem. & Schult. Syst. iv. 663 = triste.
microcalyx, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
microcarpum, Cerv. ex Lag. Gen. et Sp. Nov. 10 = Cervantesii.
microcarpum, Pav. ex Dun. in DC. Prod. xiii. 1. 72 = leptanthum.
microcarpum, *Vahl, Symb. Bot.* ii. 40.—Aegypt.
microphyllum, *Dun. Hist. Solan.* 187.—Ins. S. Doming.
microphyllum, Pav. ex Dun. in DC. Prod. xiii. 1. 86 = umbelliferum.
microspermum, *Dun. Solan. Syn.* 12.—Hab. ?
Milleri, Jacq. Coll. iv. 209 = capense.
Milnei, *Seem. in Journ. Bot.* i. (1863) 210.—N. Hebrid.
minax, Mart. ex Sendt. in Mart. Fl. Bras. x. 71 = maroniense.
miniatum, Bernh. ex Willd. Enum. Hort. Berol. 236 = nigrum.
minutiflorum, Dun. in DC. Prod. xiii. 1. 58 = nigrum.
mite, *Ruiz & Pav. Fl. Per.* ii. 38. t. 173. f. a.—Peruv.
mitlense, *Dun. in DC. Prod.* xiii. 1. 314.—Mexic.
Mocinianum, Dun. l. c. 164 = Mozinianum.
modestum, *Roem. & Schult. Syst.* iv. 663.—Mexic.
Moenchii, *Dun. Hist. Solan.* 242.—Hab. ?
moestum, *Dun. in DC. Prod.* xiii. 1. 284.—Afr. austr.
molle, Bert. ex Dun. l. c. 185 = persicaefolium.
molle, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
molle, *Humb. & Bonpl. ex Dun. Solan. Syn.* 28.—N. Granat.
molle, Schrank, Pl. Rar. Hort. Monac. t. 81 = aculeatissimum, *Jacq.*
mollissimum, Blume, Bijdr. 698 = biflorum.
monachophyllum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 33.—Venezuela.
monadelphum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 42.—Peruv.
monanthos, Dun. in DC. Prod. xiii. 1. 406 = Bassovia lucida.
monanthum, Roem. & Schult. Syst. iv. 608 = Mozinianum.
monocarpum, Dun. in DC. Prod. xiii. 1. 146 = glomuliflorum.
monodynamum, Dun. l. c. 131 = gnaphalocarpum.
monodynamum, Vis. Sem. Hort. Patav. (1841) 3 = tricolor.
montanum, *Linn. Sp. Pl.* 186.—Peruv.
montanum, Ruiz & Pav. Fl. Per. ii. 32 = tuberiferum.
montevidense, *Spreng. Syst.* i. 684.—Reg. Argent.
monticolum, *Dun. in DC. Prod.* xiii. 1. 161.—Afr. austr.
Moricandi, Dun. l. c. 319 = stagnale.
moschatum, J. & C. Presl, Delic. Prag. 77 = nigrum.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 235.—Afr. trop.

SOLANUM :—
Mozinianum, *Dun. ex Poir. Encyc. Suppl.* iii. 757.—Mexic.
Mullus, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 331 = heterodoxum.
multiangulatum, *Vell. Fl. Flum.* 91; ii. t. 133.—Bras.
multifidum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 96 = biflorum.
multifidum, Lam. Illustr. ii. 17 = senecioides.
multifidum, *Ruiz & Pav. Fl. Per.* ii. 37. t. 171.—Peruv.
multiflorum, Roth, Nov. Pl. Sp. 130 = indicum.
multinervium, *Dun. in DC. Prod.* xiii. 1. 127.—Mexic.
muricatum, *Ait. Hort. Kew.* ed. I. i. 250.—Peruv.; Chili.
muricatum, Bert. ex Dun. in DC. Prod. xiii. 1. 150 = rancaguense.
murinum, *Sendt. in Mart. Fl. Bras.* x. 29.—Bras.
Myosotis, *Dun. in DC. Prod.* xiii. 1. 102.—Bras.
myosotoideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
myoxotrichum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 426.—Madag.
myriacanthum, Dun. Hist. Solan. 218. t. 19 = aculeatissimum, *Jacq.*
myrtifolium, *Lodd. Bot. Cab.* t. 1431.—Am. austr.
Naratida, Buch.-Ham. ex Wall. Cat. n. 2619 B = spirale.
narense, *Humb. & Bonpl. ex Dun. Solan. Syn.* 18.—Am. austr.
Nava, *Webb & Berth. Phyt. Canar.* iii. 123.—Ins. Canar.
Neesianum, Dalz. & Gibs. Bomb. Fl. 175 = bigeminatum.
Neesianum, D. Dietr. Syn. Pl. i. 697 = pubescens.
Neesianum, Wall. ex Nees, in Trans. Linn. Soc. xvii. (1834) 42 = subtruncatum.
neglectum, Bert. ex Dun. in DC. Prod. xiii. 1. 170 = sylvaticum.
neglectum, *Dun. Hist. Solan.* 177.—Ind. occ.
Nelsoni, *Dun. in DC. Prod.* xiii. 1. 123.—Ins. Sandvic.
Nelsonii, Zipp. ex Span. in Linnaea, xv. (1841) 337 = indicum.
nematopus, *Sendt. in Mart. Fl. Bras.* x. 93.—Bras.
nematosepalum, *Miq. Fl. Ind. Bat.* ii. 643.—Java.
nemophilum, *F. Muell. Fragm.* ii. 161.—Austral.
nemorense, *Dun. Hist. Solan.* 194.—Peruv.
nicotianaefolium, Lag. Gen. et Sp. Nov. 10 = glutinosum.
nigrescens, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 140.—Mexic.
nigricans, *Mart. & Gal.* l. c. 134.—Mexic.
nigrum, Acerb. ex Dun. in DC. Prod. xiii. 1. 59 = pruinosum.
nigrum, Leschen. ex Dun. l. c. 177 = denticulatum.
nigrum, *Linn. Sp. Pl.* 186.—Amphigaea.
nigrum, Tausch, ex Dun. in DC. Prod. xiii. 1. 53 = Tauschii.
nilagiricum, *Schlecht. in Linnaea*, xxvi. (1853–55) 484.—Ind. or.
Nilghiry, Noton, ex Dun. in DC. Prod. xiii. 1. 259 = giganteum.
nitens, *Baker, in Journ. Bot.* xx. (1882) 220.—Madag.
nitens, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* ii. 21.—Bohemia.
nitidum, Pav. ex Dun. in DC. Prod. xiii. 1. 127 = multinervium.
nitidum, *Ruiz & Pav. Fl. Per.* ii. 33. t. 163.—Peruv.
niveum, All. ex Vitm. Summa Pl. i. 492 = marginatum.
niveum, Thunb. Prod. Pl. Cap. 36; Vahl, Symb. Bot. ii. 41 = giganteum.
noctiflorum, Hort. Aven. ex Dun. in DC. Prod. xiii. 1. 34 = stoloniferum.
nodiflorum, Desv. ex Dun. l. c. 46 = Desvauxii.
nodiflorum, Jacq. Ic. Pl. Rar. ii. 288. t. 326 = nigrum.
Nolitangere, *Salzm. ex Dun. in DC. Prod.* xiii. 1. 320.—Bras.
nossibeënse, *Vatke, in Linnaea*, xliii. (1880–82) 329.—Madag.
novemlobum, *Dun. in DC. Prod.* xiii. 1. 70.—Chili.
novo-granatense, *Dun. l. c.* 680.—N. Granat.
nudum, Humb. & Bonpl. ex Dun. Solan. Syn. 20 = triste.

SOLANUM :—
nutans, Poepp. ex Dun. in DC. Prod. xiii. 1. 137 = urceolatum.
nutans, *Ruiz & Pav. Fl. Per.* ii. 34. t. 166. f. a.—Am. trop.
nyctaginoides, *Dun. in DC. Prod.* xiii. 1. 172.—Mexic.
oaxacanum, *Dun. l. c.* 204.—Mexic.
obliquum, Bert. ex Dun. l. c. 393 = Cyphomandra betacea.
obliquum, Ruiz & Pav. Fl. Per. ii. 35. t. 165. f. a = Cyphomandra obliqua.
oblitum, *Dun. in DC. Prod.* xiii. 1. 126.—Peruv.
oblongifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 14.—N. Granat.
oblongum, *Ruiz & Pav. Fl. Per.* ii. 34.—Peruv.
obovatum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 21.—Ecuador.
obscurum, Sieber, ex Dun. in DC. Prod. xiii. 1. 224 = micracanthos.
obscurum, Vahl, Symb. Bot. ii. 41 = S. Juripeba.
obtusifolium, Hartw. ex Sendt. in Mart. Fl. Bras. x. 95 = torvum.
obtusifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 26.—Mexic.
obtusifolium, Schlecht. & Cham. in Linnaea, v. (1830) 113 = amazonium.
obtusifolium, Willd. Enum. Hort. Berol. Suppl. 11 = Zuccagnianum.
obtusilobum, Dun. in DC. Prod. xiii. 1. 294 = amblymerum.
obtusilobum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 142.—Mexic.
occidentale, *Dun. in DC. Prod.* xiii. 1. 308.—Amer.
ochrandrum, *Dun. l. c.* 110.—Bras.
ochranthum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 6.—Ecuador.
ochroleucum, Bast. in Desv. Journ. Bot. iii. (1814) 20 = nigrum.
ochroneurum, *Link, Enum. Hort. Berol.* i. 186.—Bras.
ochrophyllum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 50.—Peruv.
odoriferum, *Vell. Fl. Flum.* 85; ii. t. 108.—Bras.
Ohrondii, Carr. in Rev. Hortic. (1883) 497 = Commersoni.
Oldfieldii, *F. Muell. Fragm.* ii. 161.—Austral.
oleraceum, Dun. ex Poir. Encyc. Suppl. iii. 750 = nigrum.
oleraceum, Vell. Fl. Flum. 89; ii. t. 125 = alternato-pinnatum.
oligacanthum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 19.—Austral.
oliganthum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 8.—Am. austr.
oliganthum, Link, Enum. Hort. Berol. i. 182 = Linkii.
oocarpum, *Sendt. in Mart. Fl. Bras.* x. 106.—Bras.
opacum, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1853) *App.* 8.—Austral.
oporinum, Willd. Enum. Hort. Berol. 238 = macrophyllum.
oppositifolium, Ruiz & Pav Fl. Per. ii. 35 = urceolatum.
opuliflorum, Portenschl. ex Dun. in DC. Prod. xiii. 1. 326 = sisymbrifolium.
orbiculatum, Dun. in Poir. Encyc. Suppl. iii. 762 = parvifolium.
Orbignianum, *Sendt. in Mart. Fl. Bras.* x. 34.—Bras.
organorum, Dun. in DC. Prod. xiii. 1. 229 = schizandrum.
organum, Lapeyr. ex Dun. l. c. 171 = virgatum.
orinocense, *Humb. & Bonpl. ex Dun. Solan. Syn.* 42.—Venezuela.
Osbeckii, Dun. in DC. Prod. xiii. 1. 179 = biflorum.
Otites, *Dun. l. c.* 39.—Venezuela.
ovalifolium, *Humb. & Bonpl. ex Dun. Solan. Syn.* 37.—N. Granat.
oviferum, Salisb. Prod. 134 = S. Melongena.
ovigerum, Dun. Hist. Solan. 210 = S. Melongena.
ovoideum, Zipp. ex Miq. Fl. Ind. Bat. ii. 647 = ferox.
oxycarpum, *Schiede, in Schlecht. Hort. Hal.* i. 5. t. 3.—Mexic.
pachimatium, *Dun. in DC. Prod.* xiii. 1. 133.—Bras.
pachycalyx, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 92.—Afr. trop.
pachypetalon, *Spreng. Syst.* iv. *Cur. Post.* 72.—Nepal.

SOLANUM :—
paleatum, Schott, ex Sendt. in Mart. Fl. Bras. x. 42 = cernuum.
palinacanthum, *Dun. in DC. Prod.* xiii. 1. 245.—Bras.
palmetorum, *Dun. l. c.* 282.—Arab.
Palqui, *Molina, Sagg. Chile*, ed. II. 282.—Chili.
paludosum, Dun. in DC. Prod. xiii. 1. 57 = nigrum.
paludosum, *Moric. Pl. Nouv. Am.* 29. t. 20.—Bras.
palustre, Poepp. ex Walp. Rep. iii. 39 = Caldasii.
panamense, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* ii. 128.—Panama.
panduraeforme, *Drège, ex Dun. in DC. Prod.* xiii. 1. 370.—Afr. austr.
panduriforme, Dun. l. c. 346 = anoplocladum.
paniculatum, *Linn. Sp. Pl.* ed. II. 267.—Bras.
Papa, Valenzuela, ex Palacio, in Mém. Mus. Par. ii. (1815) 340 = Valenzuelae.
papilionaceum, Dum.-Cours. Bot. Cult. ed. I. Suppl. 153; ex Walp. Rep. iii. 78 = giganteum.
papillosum, Sendt. in Mart. Fl. Bras. x. 37 = cladotrichum.
paposanum, Phil. Fl. Atac. 42 = phyllanthum.
paradoxum, Schott, ex Dun. in DC. Prod. xiii. 1. 462 = Athenaea Martiana.
paraguariense, *Dun. l. c.* 376.—Paraguay.
parasiticum, *Blume, Bijdr.* 697.—Java.
paratyense, *Vell. Fl. Flum.* 90; ii. t. 130.—Bras.
Parmentieri, Molina, ex Walp. Rep. iii. 38 = tuberosum.
parviflorum, Cav. Ic. iii. 19 = polyacanthos.
parviflorum, Moris, ex Badaro, in Brugnat. Giorn. Fis. Dec. II. vii. (1824) 364 = nigrum.
parviflorum, Usteri, Ann. Bot. vi. (1793) 61 = corymbosum.
parvifolium, *R. Br. Prod.* 446.—Austral.
parvifolium, Steud. Nom. ed. I. 794 = igneum.
patellare, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 53.—Peruv.
patens, *Lowe, Man. Fl. Mad.* ii. 74.—Ins. Mader.
patulum, *Pers. Syn.* i. 223.—Peruv.
patulum, Roth, Catalect. fasc. ii. 23 = nigrum.
pauciflorum, Sendt. in Mart. Fl. Bras. x. 104 = dryadeum.
pauciflorum, *Vahl, Eclog. Am.* i. 20.—Ins. Martinic.
Pavoni, *Dun. in DC. Prod.* xiii. 1. 226.—Mexic.
pectinatum, A. Cunn. ex Benth. Pl. Austral. iv. 459 = cataphractum.
pectinatum, Dun. in DC. Prod. xiii. 1. 250 = mammosum.
pectinatum, Gill. ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. (1843) 385 = pinnatum.
pedunculare, *Schlecht. in Linnaea*, xix. (1847) 305.—Mexic.
pedunculatum, *Willd. ex Roem. & Schult. Syst.* iv. 662.—Hab. ?
pelliceum, *Sendt. in Mart. Fl. Bras.* x. 82.—Bras.
pendulipes, *Dun. in DC. Prod.* xiii. 1. 271.—Bras.
pendulum, Link, ex Steud. Nom. ed. I. 794 = pensile.
pendulum, *Ruiz & Pav. Fl. Per.* ii. 39. t. 174. f. a.—Peruv.
pendulum, Schott, ex Dun. in DC. Prod. xiii. 1. 76 = odoriferum.
pensile, *Sendt. in Mart. Fl. Bras.* x. 50.—Bras.
pentadactylon, *G. Don, in Loud. Hort. Brit.* 72.—Ins. Trinit.
pentapetaloides, Boj. ex Dun. in DC. Prod. xiii. 1. 85 = imamense.
pentapetaloides, *Roxb. Hort. Beng.* 17, nomen; *et ex Hornem. Suppl. Hort. Hafn.* 27.—Bras.
pentapetalum, Hort. Calc. ex Schlecht. in Linnaea, xxv. (1852) 197 = praec. ?
Pentlandi, *Dun. in DC. Prod.* xiii. 1. 51.—Peruv.
perianthomega, Vell. Fl. Flum. 87; ii. t. 118 = megalochiton.
persicaefolium, *Dun. Hist. Solan.* 185.—Ind. occ.
persicaefolium, Mart. in Flora, xxi. (1838) II. Beibl. 78 = amygdalifolium.
persicum, Willd. ex Roem. & Schult. Syst. iv. 662 = S. Dulcamara.
peruvianum, Jacq. Ic. Pl. Rar. ii. t. 337 = Lycopersicum peruvianum.
peruvianum, Linn. Sp. Pl. 186 = Lycopersicum peruvianum ?
petrophilum, *F. Muell. in Linnaea*, xxv. (1852) 433.—Austral.
pharmacum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 234.—Afr. trop.
phaseoloides, *Polak. in Linnaea*, xli. (1877) 585.—Costa Rica.

SOLANUM:—
phillyreoides, *Humb. & Bonpl. ex Dun. Solan. Syn.* 24.—N. Granat.
phlomidifolium, Moric. ex Dun. in DC. Prod. xiii. 1. 192 = Gardneri.
phlomoides, *A. Cunn. ex Benth. Fl. Austral.* iv. 464. —Austral.
phoricum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 234.—Afr. trop.
phyllanthum, *Cav. Ic.* iv. 35. t. 359.—Peruv.
physaloides, Dun. in DC. Prod. xiii. 1. 388 = Cyphomandra physaloides.
piliferum, *Benth. Pl. Hartw.* 68.—Mexic.
piligerum, Dun. in DC. Prod. xiii. 1. 152 = spissifolium.
pilosissimum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 139.—Mexic.
pilosiusculum, *Mart. & Gal. l. c.* 136.—Mexic.
pituliferum, *Dun. in DC. Prod.* xiii. 1. 265.—Bras.
pimpinellifolium, Linn. Cent. Pl. i. 8; Amoen. Acad. iv. 268 = Lycopersicum pimpinellifolium.
pimpinellifolium, Mill. ex Dun. in DC. Prod. xiii. 1. 25 = Lycopersicum Humboldtii.
pimpinelloides, *Hill, Veg. Syst.* ix. 36.—Peruv.
pinnatifidum, Lam. Illustr. ii. 17 = aviculare.
pinnatifidum, Roth, Nov. Pl. Sp. 129 = indicum.
pinnatifidum, *Ruiz & Pav. Fl. Per.* ii. 37.—Peruv.
pinnatifolium, Salisb. Prod. 133 = aviculare.
pinnatisectum, *Dun. in DC. Prod.* xiii. 1. 40.—Mexic.
pinnatum, Bert. ex Dun. l. c. 70 = Alphonsi.
pinnatum, *Cav. Ic.* v. 23. t. 439.—Chili.
pinnatum, Willd. ex Roem. & Schult. Syst. iv. 664 = fuscatum, *Roem. & Schult.*
piperiferum, *A. Rich. Tent. Fl. Abyss.* ii. 106.—Abyss.
platacanthum, *Dun. in DC. Prod.* xiii. 1. 285.—Arab.
platanifolium, *Hook. Bot. Mag.* t. 2618.—N. Granat.
platanifolium, Sendt. in Mart. Fl. Bras. x. 58 = palinacanthum.
platyphyllum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 38. —Am. austr.
platyphyllum, Torr. in Ann. Lyc. N. York, ii. (1828) 227 = Torreyi.
plebejum, *A. Rich. Tent. Fl. Abyss.* ii. 100.—Abyss.
Pleei, Dun. in DC. Prod. xiii. 1. 305 = carolinense.
Plukenetii, *Dun. Hist. Solan.* 237.—Am.?
Plukenetii, Dun. in DC. Prod. xiii. 1. 368 = pyriforme.
Plumieri, Dun. l. c. 352 = S. Melongena.
Poeppigianum, Dun. l. c. 195 = cyrtopodium.
Poeppigianum, *Sendt. in Mart. Fl. Bras.* x. 103.—Peruv.
pogogenum, Moric. Pl. Nouv. Am. 25 = Athenaea pogogena.
Pohlii, *Sendt. in Mart. Fl. Bras.* x. 54.—Bras.
Poka, *Dun. in Poir. Encyc. Suppl.* iii. 768.—Java.
polyacanthos, *Lam. Illustr.* ii. 23; *Vahl, Eclog. Am.* i. 24.—Ind. occ.
polyacanthos, L'Hérit. ex Dun. in DC. Prod. xiii. 1. 244 = aculeatissimum, *Jacq.*
polyanthemum, *Hochst. ex Dun. l. c.* 94.—Abyss.
polygamum, Boiss. ex Dun. l. c. 334 = tridynamum.
polygamum, *Vahl, Symb. Bot.* iii. 39. t. 55.—Ind. occ.
polytrichum, *Moric. Pl. Nouv. Am.* 32.—Bras.
pomiferum, Cav. Desc. 112 = Lycopersicum pyriforme.
porphyranthum, *Dun. in DC. Prod.* xiii. 1. 244.—Mexic.
portoricense, Dun. l. c. 374 = Willdenovianum.
praealtum, Sendt in Mart. Fl. Bras. x. 44 = excelsum.
premnaefolium, Dun. in DC. Prod. xiii. 1. 392 = Cyphomandra premnaefolia.
Preslei, Dun. l. c. 681 = lanceaefolium.
pressum, Dun. Hist. Solan. 217 = S. Melongena.
prinophyllum, Dun. in DC. Prod. xiii. 1. 296 = armatum.
procumbens, DC. Fl. Fr. iii. 611 = Saracha procumbens.
procumbens, *Lour. Fl. Cochinch.* 132.—Cochinch.
procumbens, Poir. Encyc. iv. 308 = Lycopersicum procumbens.
propinquum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 143.—Mexic.
prostratum, Raeusch. Nom. ed. III. 60 = procumbens.
pruinosum, *Dun. in DC. Prod.* xiii. 1. 58.—Mexic.
prunifolium, *Willd. ex Roem. & Schult. Syst.* iv. 662. —Venezuela.
pruriens, *Dun. in DC. Prod.* xiii. 1. 120.—Bras.
Pseudocapsicum, *Linn. Sp. Pl.* 184.—Amphig.
Pseudo-china, Spreng. Syst. iv. Cur. Post. 71, sphalm. = S. Pseudo-quina.

SOLANUM:—
Pseudo-glabrescens, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1154, nomen.—Guian.
Pseudo-lycopersicum, Jacq. Hort. Vindob. i. 4 = Lycopersicum esculentum.
Pseudo-melongena, *Tenore, Cat. Sem. Hort. Neap.* (1851); *et in Atti R. Ist. Incorr. Nap.* viii. (1855) 323.—Hab.?
Pseudo-quina, *A. St. Hil. Pl. Us. Brés.* t. 21.—Bras.
Pseudo-saponaceum, Blume, Bijdr. 702 = torvum.
Pseudo-undatum, Blume, l. c. 699 = S. Melongena.
Pseudo-villosum, Schur, Enum. Pl. Transs. 477 = nigrum.
psychotrioides, *Humb. & Bonpl. ex Dun. Solan. Syn.* 21.—N. Granat.
pteleaefolium, *Sendt. in Mart. Fl. Bras.* x. 15.—Bras.
pterocaulon, Reichb. ex Nym. Consp. 526 = seq.
pterocaulum, Dun. Hist. Solan. 153 = nigrum.
pterocladum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 44.—Peruv.
pteropodum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 14. —Ecuador.
ptychanthum, Dun. in DC. Prod. xiii. 1. 54 = nigrum.
puberulum, *Nutt. ex Seem. in Journ. Bot.* i. (1863) 207.—Ins. Pacif.
puberulum, *Phil. in Linnaea*, xxix. (1857–58) 22.—Chili.
pubescens, Heyne, ex Walp. Rep. iii. 83 = indicum.
pubescens, Roxb. Hort. Beng. 16; Fl. Ind. i. 564 = verbascifolium.
pubescens, Ruiz & Pav. Fl. Per. ii. 36 = luteoalbum.
pubescens, *Willd. Phyt.* i. 5.—Ind. or.
pubigerum, Dun. Hist. Solan. 160. t. 6 = Cervantesii.
pulchellum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 18 = esuriale.
pulchellum, *Phil. in Anal. Univ. Chil.* (1873) 523.—Chili.
pulchrum, *Dun. in Poir. Encyc. Suppl.* iii. 750.—Bras.
pulverulentum, Linn. Syst. ed. X. 935 = tomentosum.
pulverulentum, Nutt. ex Seem. in Journ. Bot. i. (1863) 208 = puberulum, *Nutt.*
pulverulentum, *Pers. Syn.* i. 2[illegible].—Peruv.
pulverulentum, Salisb. Prod. 133 = auriculatum.
pulvinare, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 39.—N. Guin.
pumilum, Dun. in DC. Prod. xiii. 1. 287 = carolinense.
punctulatum, *Dun. in DC. Prod.* xiii. 1. 122.—Jamaic.
pungetium, *R. Br. Prod.* 446.—Austral.
pungetium, Sieber, ex Dun. in DC. Prod. xiii. 1. 296 = armatum.
puniceum, C. C. Gmel. Fl. Bad. iv. 176 = nigrum.
purpureum, *Dun. in DC. Prod.* xiii. 1. 372.—Hab.?
pycnanthemum, *Mart. in Flora*, xx. (1837) II. *Beibl.* 120.—Bras.
pygmaeum, *Cav. Ic.* v. 23. t. 439. f. 2.—Reg. Argent.
Pyracantha, Sm. Exot. Bot. ii. 9 = Richardi?
pyracanthos, Boj. ex Dun. in DC. Prod. xiii. 1. 325 = Richardi.
pyracanthum, *Jacq. Hort. Schoenb.* iv. 36; *Lam. Illustr.* n. 2364.—Afr. trop.
pyrethrifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 250. —Reg. Argent.
pyrifolium, *Lam. Illustr.* ii. 19.—Ind. occ.
pyriforme, *Dun. in DC. Prod.* xiii. 1. 368 = Ins. Barbad.
pyriforme, Poir. Encyc. iv. 291 = Lycopersicum pyriforme.
pyrrhocarpum, *Phil. in Anal. Univ. Chil.* (1862) II. 383.—Chili.
quadrangulare, *Linn. f. Suppl.* 147.—Afr. austr.
quadriflorum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 139.—Mexic.
quadriloculare, Spreng. Syst. iv. Cur. Post. 72 = ferox.
quadriloculatum, *F. Muell. Fragm.* ii. 161.—Austral.
quadripartitum, *Dun. in DC. Prod.* xiii. 1. 72.—Bolivia.
quercifolium, Banks, ex Dun. l. c. 345 = S. Poka.
quercifolium, *Linn. Sp. Pl.* 185.—Peruv.
quercifolium, Mill. Gard. Dict. ed. VIII. n. 16 = Houstonii.
quercifolium, Poepp. ex Dun. in DC. Prod. xiii. 1. 70 = runcinatum.
quercifolium, Ruiz & Pav. Fl. Per. ii. 36 = radicans, *Linn. f.*

SOLANUM:—
quercifolium, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 121.—Afr. trop.
quinquangulare, *Willd. ex Roem. & Schult. Syst.* iv. 669.—Venezuela.
quitense, Hook. & Arn. Bot. Beech. Voy. 67 = repandum.
quitense, Hort. ex Dun. in DC. Prod. xiii. 1. 318 = maroniense.
quitense, H. B. & K. Nov. Gen. et Sp. ii. 35 = seq.
quitoënse, *Lam. Illustr.* ii. 16.—Peruv.
racemiflorum, *Dun. Hist. Solan.* 147.—Am. austr.
racemosum, *Jacq. Enum. Pl. Carib.* 15.—Ind. occ.
racemosum, Mill. Gard. Dict. ed. VIII. n. 28 = eriocalyx.
racemosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = indicum.
radiatum, *Sendt. in Mart. Fl. Bras.* x. 53.—N. Granat.
radicans, *Hill, Veg. Syst.* ix. 40 (Quid?).—Peruv.
radicans, *Linn. f. Dec.* 1. t. 10.—N. Granat.
Radula, *Vahl, Eclog. Am.* ii. 16.—Am. trop.
radulanum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
ramosissimum, *Dun. in DC. Prod.* xiii. 1. 144.—Bras.
ramosum, Lam. Encyc. iv. 281, sphalm. = racemosum.
ramulosum, *Sendt. in Mart. Fl. Bras.* x. 45.—Bras.
rancaguense, *Dun. in DC. Prod.* xiii. 1. 150.—Chili.
rancaguinum, *Phil. in Anal. Univ. Chil.* (1873) 523. —Chili.
Rantonetii, Carr. in Rev. Hortic. (1868) 420, err. typ. = seq.
Rantonnetii, *Carr. ex Lescuy. in Hérincq, L'Horticult. Franç.* Sér. II. i. (1859) 197.—Reg. Argent.
rariflorum, Steud. Nom. ed. II. ii. 605 = Linkii.
receptum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* i. 46.—Mexic.
reclinatum, L'Hérit. ex Pers. Syn. i. 225 = aviculare.
recurvum, Hoffmgg. Verz. Pfl. 113 = lividum.
recurvum, Poir. Encyc. Suppl. iii. 754 = incurvum.
reflexiflorum, *Moric. ex Dun. in DC. Prod.* xiii. 1. 213. —Bras.
reflexum, Schrank, Pl. Rar. Hort. Monac. t. 81 = aculeatissimum, *Jacq.*
refractifolium, *Sendt. in Mart. Fl. Bras.* x. 31.—Bras.
refractum, *Hook. & Arn. Bot. Beech. Voy.* 304.—Mexic.
Regnelii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 54.—Bras.
Reichenbachii, *Vatke, in Linnaea*, xliii. (1880–82) 330.—Abyss.
Reineggeri, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* iii. II. p. xix.—Bohemia.
Renggeri, *Schlecht. Hort. Hal.* fasc. i. 6.—Paraguay.
Renschii, *Vatke, in Linnaea*, xliii. (1880–82) 328.—Afr. trop.
repandum, *Forst. f. Prod.* 18.—Ins. Pacif.
repandum, F. Muell. Fragm. vi. 145 = Dallachii.
repandum, Vell. Fl. Flum. 88; ii. t. 123 = variabile.
repens, *Dun. Hist. Solan.* 239.—Bras.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = nigrum.
repens, Pav. ex Dun. in DC. Prod. xiii. 1. 39 = appendiculatum.
reptans, *Bunb. in Proc. Linn. Soc.* i. (1841) 110.—Bras.
Requieni, *Dun. in DC. Prod.* xiii. 1. 363.—Hab.?
reticulatum, Dun. in Poir. Encyc. Suppl. iii. 764 = crotonoides.
reticulatum, Juss. ex Dun. in DC. Prod. xiii. 1. 132 = Vellozianum.
reticulatum, Willd. ex Roem. & Schult. Syst. iv. 661 = dictyoticum.
retroflexum, Dun. in DC. Prod. xiii. 1. 50 = nigrum.
retrofractum, *Vahl, Eclog. Am.* i. 21.—Am. trop.
revolutum, *Hornem. Suppl. Hort. Hafn.* 26.—Bras.
rhamnoides, *Dun. in DC. Prod.* xiii. 1. 100.—Bolivia.
rhinozerothis, Blume, Bijdr. 695 = nigrum.
rhytidoandrum, *Sendt. in Mart. Fl. Bras.* x. 85.—Bras.
Richardi, *Dun. Solan. Syn.* 45.—Ins. Martinic.
Richardi, Sieber, ex Dun. in DC. Prod. xiii. 1. 254 = hirtum.

SOLANUM:—
Riedlei, *Dun. in DC. Prod.* xiii. 1. 109.—Ind. occ.
rigescens, *Jacq. Hort. Schoenb.* i. 19. t. 42.—Afr. austr.
rigidum, *Lam. Illustr.* ii. 23.—Hab.?
riparium, *Pers. Syn.* i. 221.—Am. trop.
ripense, *Dun. in DC. Prod.* xiii. 1. 680.—Venezuela.
rivulare, Dun. l. c. 304 = rubetorum.
rivulare, *Mart. ex Sendt. in Mart. Fl. Bras.* x. 25.—Bras.
robustum, *H. Wendl. in Flora*, xxvii. (1844) 784.—Bras.
Rodschiedii, G. F. W. Mey. Prim. Fl. Esseq. 111 = S. Juripeba.
Roemerianum, Scheele, in Linnaea, xxi. (1848) 767 = elaeagnifolium.
Rossii, *Knowl. & Westc. Flor. Cab.* ii. (1838) 141.—Mexic.
rostratum, *Dun. Hist. Solan.* 234.—Mexic.
rostratum, L'Hérit. ex Dun. in DC. Prod. xiii. 1. 328 = cornutum.
Rothelianum, Steud. Nom. ed. II. ii. 605, sphalm. = S. Botelhianum.
Rothii, *Dun. in DC. Prod.* xiii. 1. 383, nomen.—Hab.?
Rottboellianum, *Roem. & Schult. Syst.* iv. 668.—Guian.
rotundifolium, Moric. ex Dun. in DC. Prod. xiii. 1. 301 = Nelsoni.
Roxburghii, Dun. ex D. Don, Prod. Fl. Nep. 96 = biflorum.
Roxburghii, Dun. in DC. Prod. xiii. 1. 57 = nigrum.
rubetorum, *Dun. l. c.* 304.—Afr. austr.
rubiginosum, *Vahl, Eclog. Am.* ii. 17. t. 13.—Am. trop.
rubrum, Drège, ex Dun. in DC. Prod. xiii. 1. 77 = quadrangulare.
rubrum, *Linn. Syst.* ed. XII. 173.—Hab.?
rubrum, Mill. Gard. Dict. ed. VIII. n. 4 = nigrum.
rubrum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 = Blumei.
rude-pannum, *Dun. in DC. Prod.* xiii. 1. 264.—Mexic.
ruderale, *Phil. f. Cat. Pl. Vasc. Chil.* 229.—Chili.
ruderale, Salisb. Prod. 133 = S. Dulcamara.
rufescens, Dun. in DC. Prod. xiii. 1. 228 = sideroxanthum.
rufescens, *Sendt. in Mart. Fl. Bras.* x. 39.—Bras.
rufum, Dun. in DC. Prod. xiii. 1. 213 = brachybotryon.
rufum, Rich. ex Dun. l. c. 229 = rubiginosum.
rufum, *Sendt. in Mart. Fl. Bras.* x. 91.—Bras.
rugosum, *Dun. in DC. Prod.* xiii. 1. 108.—Bras.
rugosum, Rich. ex Poir. Encyc. Suppl. iii. 755 = Bassovia Richardi.
Rumphii, Dun. Hist. Solan. 157 = nigrum.
runcinatum, *Ruiz & Pav. Fl. Per.* ii. 36.—Peruv.; Chili.
runcinatum, Wendl. Bot. Beobacht. 43 = pyracanthum.
rupestre, F. W. Schmidt, Fl. Boëm. ii. 96 = S. Dulcamara.
rupicolum, Boj. ex Dun. in DC. Prod. xiii. 1. 85 = imamense.
rupincola, *Sendt. in Mart. Fl. Bras.* x. 100.—Bras.
sabeanum, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 6 = sisymbrifolium.
Sadae, *Phil. in Linnaea*, xxxiii. (1864-65) 203.—Chili.
Sagraeanum, *A. Rich. Fl. Cub. Fanerog.* ii. 124.—Cuba.
salicifolium, *Phil. Sert. Mendoc. Alt.* 37.—Chili.
salignum, Willd. ex Roem. & Schult. Syst. iv. 663 = cirrhosum.
Saltianum, *Roem. & Schult. l. c.* 669.—Abyss.
salviaefolium, *Dun. in DC. Prod.* xiii. 1. 187.—Bras.
salvifolium, Lam. Illustr. ii. 14 = S. Radula.
Salzmanni, Dun. in DC. Prod. xiii. 1. 206 = paludosum.
sambuciflorum, *Sendt. in Mart. Fl. Bras.* x. 38.—Bras.
Sanctae-Catharinae, *Dun. in DC. Prod.* xiii. 1. 109.—Nicaragua.
Sanctae-Crucis, *Dun. l. c.* 274.—Bolivia.
Sanctae-Nevadae, *Dun. & A. DC. l. c.* 678.—Venezuela.
sanctum, Jan, ex Dun. l. c. 260 = torvum.
sanctum, Linn. Sp. Pl. ed. II. 269 = incanum, *Linn.*
sanctum, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 605 = dubium.
sandwicense, *Hook. & Arn. Bot. Beech. Voy.* 92.—Ins. Sandvic.
sandwichense, Dun. in DC. Prod. xiii. 1. 268 = praec.

SOLANUM:—
sandwichianum, Dun. l. c. 311 = incompletum.
sanguineum, Hort. ex Dun. in DC. Prod. xiii. 1. 243 = atropurpureum.
saponaceum, *Dun. Hist. Solan.* 206.—Peruv.
saponaceum, Hook. Bot. Mag. t. 2697 = elaeagnifolium.
saponaceum, *Welw. Apont.* 588.—Afr. trop.
sarmentosum, Lam. Illustr. ii. 24 = lanceaefolium.
sarmentosum, *Nees, in Trans. Linn. Soc.* xvii. (1837) 58.—Malaya.
sarmentosum, Pav. ex Dun. in DC. Prod. xiii. 1. 87 = macrantherum.
sarrachoides, *Sendt. in Mart. Fl. Bras.* x. 18.—Bras.
sativum, Dun. in DC. Prod. xiii. 1. 360 = S. Melongena.
scaberrimum, Dun. l. c. 349 = asterophorum.
scabridum, *Dun. l. c.* 253.—N. Granat.
scabrum, H. B. & K. Nov. Gen. et Sp. iii. 41 = Humboldti.
scabrum, Jacq. Hort. Schoenb. iii. 45 = racemiflorum.
scabrum, Lam. Illustr. ii. 18 = muricatum.
scabrum, Mill. Gard. Dict. ed. VIII. n. 6 = nigrum.
scabrum, Ruiz & Pav. Fl. Per. ii. 39 = saponaceum.
scabrum, *Vahl, Eclog. Am.* i. 22.—Am. austr.
scabrum, Zuccagni, in Roem. Collect. 133 = Zuccagnianum.
scandens, Comm. ex Dun. in DC. Prod. xiii. 1. 88 = pyrifolium.
scandens, Linn. Pl. Surin. 5; Amoen. Acad. viii. 253 = S. Convolvulus?
scandens, *Linn. f. Suppl.* 147.—Guian.
scandens, Mill. Gard. Dict. ed. VIII. n. 19 = axilliflorum.
scandens, Moç. & Sesse, ex Dun. in DC. Prod. xiii. 1. 87 = macrantherum.
scandens, Neck. Delic. Gallo-Belg. i. 119; Lam. Fl. Fr. ii. 257 = S. Dulcamara.
scandens, Schomb. ex Dun. in DC. Prod. xiii. 1. 84 = pensile.
scandens, Sw. Prod. Veg. Ind. Occ. 47 = scabrum.
Schiedeanum, Schlecht. in Linnaea, xix. (1847) 295 = Hartwegi.
Schimperianum, *Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 24; *Dun. in DC. Prod.* xiii. 1. 94.—Abyss.
Schiru-schuna, *Mill. Gard. Dict.* ed. VIII. n. 32 (Quid?).—Ind. or.
schizandrum, *Sendt. in Mart. Fl. Bras.* x. 85.—Bras.
schizopodium, *Sendt. l. c.* 46.—Bras.
Schlechtendalianum, *Walp. Rep.* iii. 61.—Mexic.
Schneideri, Phil. f. Cat. Pl. Vasc. Chil. 229 = Alphonsi.
schoenbrunnense, *Dun. in DC. Prod.* xiii. 1. 365.—Hab.?
Schomburgkii, *Sendt. in Mart. Fl. Bras.* x. 55.—Am. trop.
Schrankii, *Steud. Nom.* ed. II. ii. 606.—Bras.
Schreberianum, *Sendt. in Mart. Fl. Bras.* x. 85.—Bras.
Schuchii, *Sendt. l. c.* 101.—Bras.
Schultesii, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* iii. 24.—Bohemia.
sclareaefolium, Dun. in DC. Prod. xiii. 1. 118 = cladotrichum.
Seaforthianum, *Andr. Bot. Rep.* t. 504.—Am. trop.
Seedii, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 369.—Ins. Fiji.
Sellovianum, *Sendt. in Mart. Fl. Bras.* x. 38.—Bras.
Sellowii, *Dun. in DC. Prod.* xiii. 1. 134.—Bras.
semiarmatum, *F. Muell. Fragm.* ii. 163.—Austral.
sempervirens, *Dun. in DC. Prod.* xiii. 1. 88.—Guian.
sempervirens, Mill. Gard. Dict. ed. VIII. n. 25 = aggregatum.
Sendtnerianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 75.—Peruv.
senecioides, *Domb. ex Dun. in DC. Prod.* xiii. 1. 65.—Peruv.
senegambicum, *Dun. l. c.* 194.—Afr. trop.
sepicula, *Dun. l. c.* 283.—Arab.
septemlobum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 122.—China.
sericeum, *Ruiz & Pav. Fl. Per.* ii. 33. t. 161.—Peruv.
sericeum, Vell. Fl. Flum. 83; ii. t. 97 = cinnamomeum.
serpentinum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26; Desf. Cat. Hort. Par. ed. III. 115, 397 = S. Melongena?
serratifolium, *A. DC. in DC. Prod.* xiii. 1. 680.—Bras.
serratum, Blanco, Fl. Filip. ed. I. 136 = Physalis minima.
serratum, Dun. in DC. Prod. xiii. 1. 146 = serratifolium.
sessile, *Ruiz & Pav. Fl. Per.* ii. 35. t. 167.—Peruv.

SOLANUM:—
sessiliflorum, *Dun. Solan. Syn.* 43.—Bras.
Shanesii, *F. Muell. Fragm.* vi. 144.—Austral.
sideroxanthum, *Dun. in DC. Prod.* xiii. 1. 228.—Bras.
sideroxyloides, *Schlecht. in Linnaea*, viii. (1833) 253.—Mexic.
Sieberi, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 56.—Ins. Trinit.
Sieberianum, Dun. in DC. Prod. xiii. 1. 107 = callicarpaefolium.
Silarium, Buch.-Ham. ex Wall. Cat. n. 2627 = torvum.
simile, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 19.—Austral.
simplex, *Dun. in DC. Prod.* xiii. 1. 272.—Bras.
sinaicum, *Boiss. Diagn.* Ser. I. xi. 135.—Arab.
sinense, Blanco, Fl. Filip. ed. I. 137 = tuberosum.
singuliflorum, Steud. Nom. ed. II. ii. 606 = S. Pseudocapsicum.
sinuatifolium, Vell. Fl. Flum. 91; ii. t. 132 = aculeatissimum, *Jacq.*
sinuatum, *Ait. ex Loud. Hort. Brit. Suppl.* ii. 673.—Hab.?
sinuatum, Salisb. Prod. 134 = sodomeum.
sinuatum, Willd. Sp. Pl. i. 1043 = tectum.
Sisleyanum, *Lambertye, in Rev. Hortic.* (1868) 470.—Reg. Argent.
sisymbrifolium, *Lam. Illustr.* ii. 25.—Am. bor.; Mexic.
sodomaeum, Drège, Zwei Pfl. Docum. 147; et ex Dun. in DC. Prod. xiii. 1. 367 = aculeastrum.
sodomaeum, Hort. ex Dun. l. c. 365 = schoenbrunnense.
sodomeum, *Linn. Sp. Pl.* 187.—Geront. calid.
sodomeum, Russ. ex Nees, in Trans. Linn. Soc. xvii. (1834) 55 = indicum.
solitudinum, *Delile, ex Dun. in DC. Prod.* xiii. 1. 285.—Aegypt.
somalense, *Franch. in Révoil, Çomali*, 47.—Afr. trop.
somniculentum, *Kunze, ex Schlecht. in Linnaea*, xix. (1847) 306.—Mexic.
sordidum, *Sendt. in Mart. Fl. Bras.* x. 53. t. 4.—Bras.
speciosum, *Dun. Hist. Solan.* 179.—Ind. occ.
speciosum, Hort. ex Dun. in DC. Prod. xiii. 1. 52 = nigrum.
spectabile, Steud. Nom. ed. II. ii. 606 = bifissum.
sphaerocarpum, Moric. Pl. Nouv. Am. 31 = aculeatissimum, *Jacq.*
spinatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 26 (Quid?).—Java.
spinosissimum, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 74, nomen.—Hab.?
spiraeoides, *Dun. in DC. Prod.* xiii. 1. 157.—Chili.
spirale, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 564.—Reg. Himal.; Burma.
spissifolium, *Sendt. in Mart. Fl. Bras.* x. 33.—Bras.
sporadotrichum, *F. Muell. in Melb. Chemist* (Oct. 1882).—Austral.
Sprengelii, *Dun. in DC. Prod.* xiii. 1. 372.—Bras.
Sprucei, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 67.—Peruv.
spurium, Balb. Cat. Hort. Taur. (1813) 73 = Lycopersicum pyriforme.
spurium, J. F. Gmel. Syst. 384 = Lycopersicum esculentum.
squamulosum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 140.—Mexic.
stagnale, *Moric. Pl. Nouv. Am.* 34. t. 23.—Bras.
stamineum, Steud. in Flora, xxvi. (1843) 764 = S. Juripeba.
Stauntonii, Dun. ex D. Don, Prod. Fl. Nep. 96 = biflorum.
stellatum, *Jacq. Coll.* iii. 254.—Am. austr.
stellatum, Moench, Meth. 475 = Moenchii.
stellatum, Ruiz & Pav. Fl. Per. ii. 40 = hispidum.
stelligerum, *Sm. Exot. Bot.* ii. 57.—Austral.
stenandrum, Dun. in DC. Prod. xiii. 1. 270 = stenorchis.
stenandrum, *Sendt. in Mart. Fl. Bras.* x. 68.—Bras.
stenolobum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 69.—Peruv.
stenopetalum, A. Br. in Ind. Sem. Hort. Berol. (1853) App. 8 = nigrum.
stenophyllum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 15.—Peruv.

SOLANUM:—
stenorchideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
stenorchis, *Dun. in DC. Prod.* xiii. 1. 270.—Am. austr.
stephanodes, *Schlecht. in Linnaea*, xix. (1847) 290.—Mexic.
stipulaceum, Brouss. ex Roem. & Schult. Syst. iv. 669 (Quid ?).—Hab. ?
stipulaceum, *Willd. ex Roem. & Schult. Syst.* iv. 662.—Bras.
stipulatum, Vell. Fl. Flum. 87; ii. t. 117 = rivulare.
stoloniferum, *Schlecht. & Bouché, in Verh. Ver. Preuss. Gartenb.* ix. (1833) 319. t. 2.—Mexic.
stoloniferum, Schlecht. in Linnaea, viii. (1833) 255 = oxycarpum.
stramonifolium, Dun. Hist. Solan. 222 = ferox.
stramonifolium, *Jacq. Misc.* ii. 298.—Cosmop. trop.
stramonifolium, Lam. Illustr. ii. 22 = torvum.
stramonifolium, Pav. ex Dun. in DC. Prod. xiii. 1. 346 = anoplocladum.
stramoniifolium, Hort. ex Benth. in Maund, Botanist, iv. [1839] sub t. 192 = vestitum.
strictum, Zuccagni, in Roem. Collect. 132 = nigrum.
strigosum, *Vand. Fasc. Pl.* 10.—Bras.
Sturtianum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 19.—Austral.
Styleanum, *Dun. in DC. Prod.* xiii. 1. 44.—Chili.
styracifolium, Willd. ex Roem. & Schult. Syst. iv. 663 = S. Juripeba.
suaveolens, Boj. Hort. Maurit. 239 = Bojeri.
suaveolens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 14.—Mexic.
subarmatum, *Willd. Enum. Hort. Berol.* 240.—Ind. occ.
subbiflorum, Ortega, Hort. Matr. Dec. 118 = capense.
subcordatum, *Sendt. in Mart. Fl. Bras.* x. 102.—Bras.
subenervium, *Dun. in DC. Prod.* xiii. 1. 104.—Chili.
subexarmatum, *Dun. l. c.* 367.—Afr. austr.
subinerme, *Jacq. Enum. Pl. Carib.* 15.—Ind. occ.
sublentum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 53.—Bras.
sublobatum, *Willd. ex Roem. & Schult. Syst.* iv. 664.—Reg. Argent.
subrepandum, *Desf. Tabl.* ed. I. 222.—Am. trop.
subscandens, *Vell. Fl. Flum.* 90; t. 128.—Bras.
subserratum, *Dun. in DC. Prod.* xiii. 1. 212.—N. Granat.
subsessile, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 964, nomen.—Guian.
subspatulatum, *Sendt. in Mart. Fl. Bras.* x. 45.—Bras.
subtruncatum, *Wall. Cat* n. 2620.—Reg. Himal.
subumbellatum, *Vell. Fl. Flum.* 85; ii. t. 105.—Bras.
subviscidum, *Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 35.—Bras.
suffruticosum, Schousb. ex Willd. Enum. Hort. Berol. 1236 = nigrum.
sulcatum, *Roeusch. Nom.* ed. III. 60.—Am. trop.?
supinum, *Dun. in DC. Prod.* xiii. 1. 289.—Afr. austr.
surattense, Burm. f. Fl. Ind. 57 = virginianum.
surinamense, Sendt. in Mart. Fl. Bras. x. 103 = canocinereum.
surinamense, *Steud. in Flora*, xxvi. (1843) 764.—Guian.
Swartzianum, *Roem. & Schult. Syst.* iv. 602.—Bras.
sycocarpum, Mart. & Sendt. in Flora, xxiv. (1841) II. Beibl. 85 = Cyphomandra sycocarpa.
sycophanta, *Dun. in DC. Prod.* xiii. 1. 357.—Venezuela.
sylvaticum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 24.—Am. austr.
sylvestre, Audib. ex Dun. in DC. Prod. xiii. 1. 32 = tuberosum.
symphysicaulis, *Pav. ex Dun. in DC. Prod.* xiii. 1. 106.—Mexic.
synantherum, *Sendt. in Flora*, xxix. (1846) 178.—Guatemala.
syringaefolium, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 10.—Chili.
tabaccifolium, Vell. Fl. Flum. 81; ii. t. 89 = auriculatum.
tabacifolium, *Salzm. ex Dun. in DC. Prod.* xiii. 1. 261.—Bras.
Tagore, J. F. Gmel. Syst. 383, sphalm. = *S. Tegore.*
taitense, *Vatke, in Linnaea*, xliii. (1880–82) 327.—Afr. trop.

SOLANUM:—
tampicense, *Dun. in DC. Prod.* xiii. 1. 284.—Mexic.
tarapotense, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 38.—Peruv.
Tauschii, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* iii. 20.—Bohemia.
tectum, *Pers. Syn.* i. 227.—Mexic.
Tegore, Aubl. Pl. Gui. i. 212. t. 84 = Cyphomandra Tejore.
tejucense, *Dun. in DC. Prod.* xiii. 1. 248.—Bras.
tenue, *Sendt. in Mart. Fl. Bras.* x. 13.—Bras.
tenuiflorum, Steud. Nom. ed. II. ii. 606 = nigrum.
tenuifolium, *Dun. in DC. Prod.* xiii. 1. 87.—N. Granat.
tenuissimum, *Sendt. in Mart. Fl. Bras.* x. 67.—Bras.
terminale, *Forsk. Fl. Aegypt. Arab.* 45.—Arab.
terminale, Vell. Fl. Flum. 84; ii. t. 101 = isodynamum.
ternatum, *Ruiz & Pav. Fl. Per.* ii. 38. t. 172.—Peruv.
tetramerum, *Dun. in DC. Prod.* xiii. 1. 681.—Cuba.
tetrandrum, Blume, Cat. Gew. Buitenz. 55 = ardisioides.
tetrandrum, *R. Br. Prod.* 445.—Austral.
tetrathecum, *F. Muell. Fragm.* ii. 165.—Austral.
tetricum, *Dun. in DC. Prod.* xiii. 1. 205.—Bras.
tettense, *Klotzsch, in Peters, Reise Mossamb. Bot.* 237.—Afr. trop.
texanum, Dun. in DC. Prod. xiii. 1. 359 = integrifolium ?
texanum, Hort. ex Tenore, in Atti R. Ist. Incorr. Nap. viii. (1855) 328 = integrifolium.
texense, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 227 = elaeagnifolium.
thelopodium, *Sendt. in Mart. Fl. Bras.* x. 46.—Bras.
thomasiaefolium, *Sendt. l. c.* 74.—Bras.
Thonningianum, *Jacq. f. Eclog.* i. 123.—Afr. trop.
Thouinii, C. C. Gmel. Cat. Hort. Carlsr. 254 = sisymbrifolium.
thymifolium, *Willd. ex Roem. & Schult. Syst.* iv. 664.—Reg. Argent.
tinctorium, Welw. Apont. 590 = nodiflorum.
tolimense, *Wedd. Chlor. And.* ii. 106.—N. Granat.
Tomatillo, *Phil. f. Cat. Pl. Vasc. Chil.* 229.—Chili.
tomentellum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 236.—Afr. trop.
tomentosum, Herb. Hasselt, ex Miq. Fl. Ind. Bat. ii. 654 = S. Melongena.
tomentosum, *Linn. Sp. Pl.* 188.—Am. bor.; Afr. austr.
tomentosum, *Phil. f. Cat. Pl. Vasc. Chil.* 229.—Chili.
Topiro, *Humb. & Bonpl. ex Dun. Solan. Syn.* 10.—Am. austr.
Torreyi, *A. Gray, in Proc. Am. Acad.* vi. (1862) 44.—Am. bor. occ.
torvum, Buch.-Ham. ex Wall. Cat. n. 2627 = stramonifolium.
torvum, Schlecht. in Linnaea, v. (1830) 113 = Hartwegi.
torvum, *Sw. Prod. Veg. Ind. Occ.* 47.—Cosmop. trop.
toxicarium, Rich. in Act. Soc. Hist. Nat. Par. ii. (1792) 107 = stramonifolium.
trachyphyllum, *Humb. in Bonpl. ex Dun. Solan. Syn.* 18.—N. Granat.
triangulare, Lam. Illustr. ii. 18 = nigrum.
tribulosum, *Schau. in Linnaea*, xx. (1847) 711.—Mexic.
trichocalyx, *Dun. in DC. Prod.* xiii. 1. 679.—Bras.
trichocarpum, *Miq. Stirp. Surin. Sel.* 134.—Guian.
tricolor, *Dun. in Poir. Encyc. Suppl.* iii. 756.—Mexic.
tricuspidatum, *Rich. ex Dun. in Poir. l. c.* iii. 771.—Guian.
tridynamum, *Dun. in Poir. l. c.* 776.—Mexic.
triferum, *Poepp. ex Schlecht. Hort. Hal.* i. 6.—Peruv.
triflorum, *Nutt. Gen. Am.* i. 128.—Am. bor. occ.
trifoliatum, Burm. f. Fl. Ind. 57. f. 3 = Triphasia Aurantiola.
trifolium, *Dun. in DC. Prod.* xiii. 1. 68.—Ecuador.
trifurcum, *Dun. l. c.* 272.—Bolivia.
trilobatum, *Linn. Sp. Pl.* 188; ed. II. 270.—As. trop.
trilobatum, Linn. ll. cc., partim = capense.
trilobatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 26 = S. Melongena.
tripartitum, *Dun. in DC. Prod.* xiii. 1. 72.—Bolivia.
triphyllum, Vell. Fl. Flum. 88; ii. t. 120 = pruniifolium.
triquetrum, *Cav. Ic.* iii. 30. t. 259.—Mexic.
trisectum, *Dun. in DC. Prod.* xiii. 1. 36.—Ins. Mader.
triste, Glocker, ex Dun. l. c. 147 = acuminatum.

SOLANUM:—
triste, *Jacq. Enum. Pl. Carib.* 15.—Am. trop.
Trongum, Poir. in Lam. Encyc. iv. 308 = S. Melongena.
Trutel, Jacq. ex Steud. Nom. ed. I. 796 = triste.
tuberculatum, Pav. ex Dun. in DC. Prod. xiii. 1. 66 = conicum.
tuberiferum, *Dun. l. c.* 63.—Mexic.
tuberosum, Bert. ex Walp. Rep. iii. 39 = etuberosum.
tuberosum, *Linn. Sp. Pl.* 185.—Am. bor. et austr.
tuberosum, Poepp. ex Walp. Rep. iii. 39 = S. Maglia.
tubiflorum, Dun. in DC. Prod. xiii. 1. 318 = maroniense.
tucumanense, *Griseb. in Goett. Abh.* xxiv. (1879) 254.—Reg. Argent.
Tweedianum, *Hook. Bot. Mag.* t. 3385.—Reg. Argent.
uliginosum, Blume, Bijdr. 695 = nigrum.
ulmoides, *Dun. in DC. Prod.* xiii. 1. 130.—Mexic.
umbellatum, Banks, ex Dun. l. c. 200 = jamaicense.
umbellatum, *Dun. l. c.* 62.—Ins. S. Doming.
umbellatum, *Mill. Gard. Dict.* ed. VIII. n. 27.—Mexic.
umbellatum, Willd. ex Roem. & Schult. iv. 662 = sylvaticum.
umbelliferum, *Eschsch. in Mem. Acad. Pétersb.* (1821–22) 283.—Calif.
umbelliferum, Torr. in Pacif. Rail. Rep. vii. 12 = nigrum.
umbrosum, Balb. ex Dun. in DC. Prod. xiii. 1. 185 = persicaefolium.
umbrosum, Bert. ex Dun. l. c. 185 = bahamense.
umbrosum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 33.—Am. austr.
umbrosum, Vahl, ex Dun. in DC. Prod. xiii. 1. 259 = anomalum.
uncinatum, *R. Br. in Salt, Abyss. App.* 64, nomen.—Abyss.
uncinatum, Schrank, in Syll. Ratisb. i. (1824) 200 = Schránkii.
uncinellum, *Lindl. Bot. Reg.* (1840) t. 15.—Bras.
undatifolium, *Dun. in DC. Prod.* xiii. 1. 104.—Bras.
undatum, Bouton, ex Dun. l. c. 366 = sodomeum.
undatum, Lam. Illustr. ii. 22 = S. Melongena.
undatum, Walsh, Not. xi. 310; ex Dun. in DC. Prod. xiii. 1. 338 = lycocarpum.
undecimangulare, Willd. ex Roem. & Schult. Syst. iv. 669 = platyphyllum.
undulatum, *Dun. in DC. Prod.* xiii. 1. 138.—Bras.
undulatum, Poir. Encyc. iv. 301 = S. Melongena.
ungens, *Weinm. in Flora*, iv. (1821) 27.—Bras.
unguiculatum, *A. Rich. Tent. Fl. Abyss.* ii. 102.—Abyss.
uniflorum, *Dun. in Poir. Encyc. Suppl.* iii. 757.—Ins. S. Doming.
uniflorum, Meyen, ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 388 = elaeagnifolium.
uniflorum, Sesse, ex Lag. Gen. et Sp. Nov. 10 = Mozinianum.
uniflorum, Vell. Fl. Flum. 86; ii. t. 114 = S. Pseudocapsicum.
Uporo, *Dun. in DC. Prod.* xiii. 1. 138.—Ins. Tahiti.
urceolatum, *Pers. Syn.* i. 223.—Peruv.
urens, *Bonhomme, ex Dun. in DC. Prod.* xiii. 1. 311.—Ins. S. Vincent.
urticans, *Dun. l. c.* 324.—Bolivia.
utile, Klotzsch, in Allg. Gartenz. xvii. (1849) 315 = tuberosum.
uvariaefolium, Dun. in DC. Prod. xiii. 1. 88 = flaccidum.
vacciniifolium, Dun. l. c. 160 = havanense, *Jacq.*
vagum, *Heyne, ex Wall. Cat.* n. 2624.—Ind. or.
Vaillantii, *Dun. Solan. Syn.* 38.—Hab. ?
Valadares, Vand. Fasc. Pl. 10 = spectabile.
valdiviense, *Dun. in DC. Prod.* xiii. 1. 195.—Chili.
Valenzuelae, *Palacio, in Mém. Mus. Par.* ii. (1815) 340.—N. Granat.
Vandellianum, *Dun. Hist. Solan.* 241.—Bras.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 85.—Peruv.
variabile, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 80.—Bras.
variegatum, Ruiz & Pav. Fl. Per. ii. 32 = muricatum.
vellereum, *Schlecht. in Linnaea*, xix. (1847) 302.—Mexic.
velleum, *Sw. ex Roem. & Schult. Syst.* iv. 643 (an seq.?).—Bras.

SOLANUM:—
velleum, *Thunb. Pl. Bras. Dec.* ii. 21.—Bras.
Vellozianum, *Dun. Hist. Solan.* 236.—Bras.
velutinum, *Dun. in Poir. Encyc. Suppl.* iii. 766.—Guian.
venosum, *Humb. & Bonpl. ex Dun. Solan. Syn.* 19.—N. Granat.
venosum, Sendt. in Mart. Fl. Bras. x. 27 = brachystachys.
venustum, Kunth, Ind. Sem. Hort. Berol. (1845) 10 = Seaforthianum.
Verae-crucis, Steud. Nom. ed. II. ii. 607 = amazonium.
verbascifolium, Banks, ex Dun. in DC. Prod. xiii. 1. 271 = stenorchis.
verbascifolium, H. B. & K. Nov. Gen. et Sp. iii. 80 = asperum.
verbascifolium, *Linn. Sp. Pl.* 184.—Amphig. trop.
vernicatum, *Lindl. Bot. Reg.* (1838), *Misc.* 74.—Reg. Argent.
verrucosum, Schlecht. Hort. Hal. i. 3. t. 2 = tuberosum.
vescum, F. Muell. in Trans. Vict. Inst. i. (1855) 69 = aviculare.
Vespertilio, *Ait. Hort. Kew.* ed. I. i. 252.—Ins. Canar.
vespertilioneum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
vestissimum, *Dun. in DC. Prod.* xiii. 1. 322.—Venezuela.
vestitum, *Benth. in Maund, Botanist*, iv. [1839] t. 192.—Mexic.
vestitum, Nutt. ex Seem. Fl. Vit. 174 = Nelsoni.
vestitum, Salzm. ex Dun. in DC. Prod. xiii. 1. 324 = polytrichum.
viarum, *Dun. l. c.* 240.—Bras.
viliflorum, *Sendt. in Mart. Fl. Bras.* x. 46.—Bras.
villosissimum, Zuccagni, in Roem. Collect. 132 = mammosum.
villosum, F. G. Dietr. Vollst. Lexic. Gaertn. ix. 341 = tectum.
villosum, Forsk. Fl. Aegypt. Arab. 47 = Forskålii.
villosum, Mill. Gard. Dict. ed. VIII. n. 2; Lam. Illustr. ii. 18 = nigrum.
villosum, *Moench, Meth.* 474.—Hab.?
villosum, Opiz, ex Dun. in DC. Prod. xiii. 1. 51 = decipiens.
villosum, *Willd. Prod. Fl. Berl.* 87.—Europ.; As.; Afr.
Vincentii, Delile, ex Dun. in DC. Prod. xiii. 1. 310 = indicum.
violaceum, Blume, Cat. Gew. Buitenz. 55 = Blumei.
violaceum, *R. Br. Prod.* 445.—Austral.
violaceum, DC. ex Dun. in DC. Prod. xiii. 1. 359 = S. Melongena.
violaceum, Hort. ex Dun. l. c. 334 = tridynamum.
violaceum, Ortega, Hort. Matr. Dec. 56; Jacq. Fragm. 82 = indicum.
violaefolium, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 403.—Bras.
virescens, Hiern, in Kjoeb. Vidensk. Meddel. (1877–78) 39 = rufescens.
virgatum, Endl. ex Sendt. in Mart. Fl. Bras. x. 13 = Endlicheri.
virgatum, *Lam. Illustr.* ii. 14.—Ins. Canar.; Ind. occ.
virginianum, Jacq. Ic. Pl. Rar. ii. 11. t. 332 = xanthocarpum.
virginianum, *Linn. Sp. Pl.* 187.—Am. bor.
virginianum, Pav. ex Dun. in DC. Prod. xiii. 1. 331 = heterodoxum.
virginianum, Pers. Syn. i. 227 = nigrum.
virginianum, Russ. ex Wall. Cat. n. 2613 D = indicum.
virginicum, Linn. Mant. ii. 340 = virginianum?
viride, *R. Br. Prod.* 445.—Austral.
viride, Schur, Enum. Pl. Transs. 478 = nigrum.
viride, Soland. ex Forst. f. Pl. Escul. 72 = S. Uporo.
viridescens, Hort. ex Reichb. Fl. Germ. Excurs. 391 = nigrum.
viridiflorum, Ruiz & Pav. Fl. Per. ii. 38. t. 173. f. b = Cyphomandra viridiflora.
viridiflorum, Schlecht. in Linnaea, xix. (1847) 292 = aculeatissimum, *Jacq.*
viridifolium, Dun. in DC. Prod. xiii. 1. 73 = viride.
viridipes, *Dun. l. c.* 234.—Bras.
viscidissimum, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 571.—Java.
viscidum, Schweigg. Enum. Pl. Hort. Regiom. 67 = sisymbrifolium.
viscosissimum, *Sendt. in Mart. Fl. Bras.* x. 14.—Bras.
viscosum, Lag. Gen. et Sp. Nov. 10 = sisymbrifolium.
viscosum, *Moench, Meth. Suppl.* 179.—Hab.?

SOLANUM:—
viscosum, Salzm. ex Dun. in DC. Prod. xiii. 1. 206 = paludosum.
vitiense, *Seem. in Journ. Bot.* i. (1863) 206; *Fl. Vit.* 176.—Ins. Fiji.
vitocense, *Dun. in DC. Prod.* xiii. 1. 176.—Peruv.
volubile, Sw. Fl. Ind. Occ. i. 458 = scabrum.
volubile, Thib. ex Dun. in DC. Prod. xiii. 1. 220 = ambiguum.
vulgare, Hegetschw. Fl. Schw. 219 = nigrum.
vulgatum, Willd. ex Steud. Nom. ed. II. ii. 607 = nigrum.
Wallisii, *Carr. in Rev. Hortic.* (1877) 291.—Peruv.
Warmingii, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 44.—Bras.
Warszewiczii, *Hort. ex Lambertye, in Rev. Hortic.* (1865) 429.—Hab.?
Wightii, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 234 = torvum.
Wightii, *Nees, in Trans. Linn. Soc.* xvii. (1834) 51.—Ind. or.
Wildianum, *Wender. ex Steud. Nom.* ed. II. ii. 607.—Hab.?
Willdenovianum, *G. Don, Gen. Syst.* iv. 427.—Ind. occ.
Willdenowii, Roem. & Schult. Syst. iv. 661 = Zuccagnianum.
woahense, Dun. in DC. Prod. xiii. 1. 268 = sandwicense.
Wrightii, *Benth. Fl. Hongk.* 243.—Bolivia.
xanthacanthum, Willd. ex Walp. Rep. vi. 601 = sisymbrifolium.
xanthocarpum, *Schrad. & Wendl. Sert. Hanov.* i. 8. t. 2.—Geront. trop.
Xanti, *A. Gray, in Proc. Am. Acad.* xi. (1876) 90.—Calif.
xylopiaefolium, *Dun. in DC. Prod.* xiii. 1. 179.—N. Granat.
xylorhagodes, Schrank, in Denkschr. Akad. Muench. vii. (1821) 241 (Quid?).—Europ.
xylostoideum, Moric. ex Dun. in DC. Prod. xiii. 1. 158 = Orbignianum.
Zanoni, *Gouan, Illustr.* 7.—Europ. austr.
zanzibarense, *Vatke, in Linnaea*, xliii. (1880–82) 326.—Afr. trop.
zeylanicum, Blanco, Fl. Filip. ed. I. 136 = ferox.
zeylanicum, Scop. Delic. Insub. i. 1 = S. Melongena.
Zollingeri, Dun. in DC. Prod. xiii. 1. 176 = biflorum.
Zuccagnianum, *Dun. Hist. Solan.* 149. t. 11.—Hab.?

SOLARIA, Phil. in Linnaea, xxix. (1857–58) 72.
LILIACEAE, Benth. & Hook. f. iii. 805.
SYMEA, Baker, in Saund. Refug. Bot. t. 260 (1871).
miersioides, *Phil. in Linnaea*, xxix. (1857–58) 72.—Chili.

SOLDANELLA, Linn. Syst. ed. I (1735). *PRIMULACEAE*, Benth. & Hook. f. ii. 633.
GOLIA, Adans. Fam. ii. 231 (1763).
affinis, Cree, ex Loud. Hort. Brit. Suppl. ii. 673 = montana.
alpina, *Linn. Sp. Pl.* 144.—Europ. alp.
alpina, F. W. Schmidt, Fl. Boëm. ii. 49 = montana.
Clusii, Gaud. Fl. Helv. ii. 76 = pusilla.
Clusii, F. W. Schmidt, Fl. Boëm. ii. 50 = alpina.
crenata, *Penny, ex G. Don, in Loud. Hort. Brit. Suppl.* i. 600.—Hab.?
crenata, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 94 = Schizocodon soldanelloides.
Ganderi, Hut. in Oestr. Bot. Zeitschr. xxiii. (1873) 122 = minima.
hybrida ×, Kern. l. c. xxv. (1875) 160.—Europ.
media ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 100.—Europ.
minima, *Hoppe, ex Sturm, Deutschl. Fl.* Heft 20.—Europ. med.
minima, Hoppe, in Flora, i. (1818) 14 = pusilla.
montana, *Willd. Enum. Hort. Berol.* 192.—Bohem.; Transsylv.
pusilla, *Baumg. Enum. Stirp. Transs.* i. 138.—Transsylv.
pyrolaefolia, Schott, Nym. & Kotschy, Analect. Bot. 16 = alpina.
sinuata, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 94 = Schizocodon soldanelloides.
villosa, Darracq, in Ann. Soc. Linn. Bord. vi. Mélang. 2; ex Gren. & Godr. Fl. Fr. ii. 461 = montana.

SOLDEVILLA, Lag. in Varied. Cienc. ii. iv. (1805) 37 = **Hispidella**, Barnad. (Compos.).
setosa, Lag. l. c. 38 = H. hispanica.

SOLEA, Spreng. in Schrad. Journ. Bot. iv. (1800) 192 = **Ionidium**, Vent. (Violar.).
alba, Spreng. Syst. iv. Cur. Post. 97 = I. album.
anomala, Spreng. Syst. i. 804 = I. anomalum.
atropurpurea, Spreng. Syst. iv. Cur. Post. 98 = I. atropurpureum.
bicolor, Spreng. l. c. 97 = I. bicolor.
bigibbosa, Spreng. l. c. 98 = I. bigibbosum.
brevicaulis, Spreng. l. c. = I. brevicaule.
buxifolia, Spreng. Syst. i. 804 = I. buxifolium.
Calceolaria, Spreng. l. c. 805 = I. Ipecacuanha.
calycina, Spreng. l. c. 804 = I. calycinum.
capensis, Spreng. l. c. = I. capense.
circaeoides, Spreng. l. c. = I. circaeoides.
communis, Spreng. Syst. iv. Cur. Post. 97 = I. commune.
concolor, Ging. in DC. Prod. i. 306 = I. concolor.
enneasperma, Spreng. Syst. i. 804 = I. heterophyllum.
erecta, Spreng. l. c. = I. heterophyllum.
glutinosa, Spreng. l. c. = I. commune.
guaranitica, Spreng. Syst. iv. Cur. Post. 98 = I. commune.
heterophylla, Spreng. Syst. i. 804 = I. heterophyllum.
Ipecacuanha, Spreng. l. c. 805 = I. Ipecacuanha.
Itoubou, Spreng. l. c. = I. Ipecacuanha.
lanata, Spreng. Syst. iv. Cur. Post. 97 = I. lanatum.
linifolia, Spreng. Syst. i. 803 = I. linifolium.
longifolia, Spreng. l. c. = I. strictum.
microphylla, Spreng. l. c. = I. glutinosum.
monopetala, Spreng. l. c. 400 = I. filiforme.
montana, Zipp. ex Span. in Linnaea, xv. (1841) 167 = I. heterophyllum.
nana, Spreng. Syst. iv. Cur. Post. 98 = I. album.
oppositifolia, Spreng. Syst. i. 803 = I. oppositifolium.
parietarifolia, Spreng. l. c. 804 = I. oppositifolium.
parviflora, Spreng. l. c. = I. glutinosum.
parvifolia, Eichl. in Mart. Fl. Bras. xiii. 1. 374 = praec.
Poaya, Spreng. Syst. iv. Cur. Post. 97 = I. Poaya.
riparia, Spreng. Syst. i. 803 = I. oppositifolium.
scariosa, Spreng. Syst. iv. Cur. Post. 97 = I. brevicaule.
setigera, Spreng. l. c. = I. setigerum.
stricta, Spreng. Pugill. i. 22 = I. concolor.
stricta, Spreng. Syst. i. 803 = I. strictum.
sylvatica, Spreng. Syst. iv. Cur. Post. 97 = I. commune.
urticaefolia, Spreng. l. c. 98 = I. circaeoides.
verbenacea, Spreng. Syst. i. 803 = I. verbenaceum.
verticillata, Spreng. in Schrad. Journ. Bot. iv. (1800) 192 = I. polygalaefolium.
viscidula, Spreng. Syst. i. 803 = I. oppositifolium.

SOLEIROLIA, Gaudich. in Freyc. Voy. Bot. 504 (1826) = **Helxine**, Req. (Urtic.).
corsica, Gaudich. l. c. = H. Soleirolii.

SOLENA, Lour. Fl. Cochinch. 514 (1790) = **Melothria**, Linn. (Cucurb.).
heterophylla, Lour. l. c. = M. heterophylla.

SOLENA, Willd. Sp. Pl. i. 961 (1797) = **Posoqueria**, Aubl. (Rubiac.).
armata, D. Dietr. Syn. Pl. i. 800 = Basanacantha armata.
Bowieana, D. Dietr. l. c. 801 = Randia macrantha.
calycina, D. Dietr. l. c. 799 (Basanacanthae sp.).—Bras.
capitata, D. Dietr. l. c. 800 = Randia capitata.
decora, D. Dietr. l. c. 799 = P. latifolia.
dumetorum, D. Dietr. l. c. = Randia dumetorum.
echinocarpa, D. Dietr. l. c. = Randia echinocarpa.
fasciculata, D. Dietr. l. c. 800 = Randia fasciculata.
ferox, D. Dietr. l. c. = Randia ferox.
floribunda, D. Dietr. l. c. = Randia dumetorum.
genipaeflora, D. Dietr. l. c. 801 = Randia genipaeflora.
gracilis, Rudge, Pl. Gui. i. 27. t. 41 = P. gracilis.
havanensis, D. Dietr. Syn. Pl. i. 799 = P. havanensis.
horrida, D. Dietr. l. c. 800 = Randia horrida.
Humboldtiana, D. Dietr. l. c. = Basanacantha Humboldtiana.
latifolia, Rudge, Pl. Gui. i. 26. t. 40 = P. latifolia.
longiflora, Willd. Sp. Pl. i. 961 = P. longiflora.
longispina, D. Dietr. Syn. Pl. i. 799 = Randia dumetorum.

longistyla, D. Dietr. Syn. Pl. i. 801 = Macrosphyra longistyla.
macrantha, D. Dietr. l. c. = Randia macrantha.
maculata, D. Dietr. l. c. = Randia maculata, *DC.*
madagascariensis, D. Dietr. l. c. = Randia madagascariensis.
multiflora, D. Dietr. l. c. 800 = Randia longiflora.
Mussaendae, D. Dietr. l. c. 801 = Randia Mussaenda.
nitida, D. Dietr. l. c. 800 = Randia nitida.
nutans, D. Dietr. l. c. 799 = Randia dumetorum.
obovata, D. Dietr. l. c. = Randia obovata.
parviflora, D. Dietr. l. c. 800 = Randia parviflora.
pubescens, D. Dietr. l. c. 799 = Randia pubescens.
Randia, D. Dietr. l. c. = Randia aculeata.
revoluta, D. Dietr. l. c. = Posoqueria latifolia.
rigida, D. Dietr. l. c. 800 = Randia fasciculata.
rotundifolia, D. Dietr. l. c. 799 = Randia rotundifolia.
Ruiziana, D. Dietr. l. c. 801 = Randia Ruiziana.
scandens, D. Dietr. l. c. 800 = Randia longiflora.
sinensis, D. Dietr. l. c. = Randia sinensis.
speciosa, D. Dietr. l. c. 801 = Randia speciosa, *DC.*
stricta, D. Dietr. l. c. 800 = Hyptianthera stricta.
Talangiana, D. Dietr. l. c. 801.—Madag.
tetracantha, D. Dietr. l. c. 800 = Basanacantha heteracantha.
tetrandra, D. Dietr. l. c. = Randia tetrandra.
triflora, D. Dietr. l. c. = Randia triflora.
Trinitatis, D. Dietr. l. c. 799 = P. Trinitatis.
uliginosa, D. Dietr. l. c. = Randia uliginosa.

SOLENACANTHUS, C. Muell. in Walp. Ann. v. 638 (1858), sphalm. = SOLAENACANTHUS, Oerst. = **Ruellia**, Linn. (Acanth.).

SOLENACHNE, Steud. Syn. Pl. Gram. 12 (1854) = **Spartina**, Schreb. (Gramin.).
phalaroides, Steud. l. c. = Spartina ciliata.

SOLENANDRA, Benth. & Hook. f. Gen. ii. 620 (1876), sphalm. = SOLENANDRIA, Beauv. = **Galax**, Linn. (Diapens.).

SOLENANDRA, Hook. f. in Benth. & Hook. f. Gen. ii. 43 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
ixoroides, *Hook. f. in Hook. Ic. Pl.* t. 1150.—Cuba.

SOLENANDRIA, Beauv. ex Vent. Jard. Malm. t. 69 (1803) = **Galax**, Linn. (Diapens.).
cordifolia, Beauv. ex Vent. l. c. = G. aphylla.

SOLENANTHA, G. Don, Gen. Syst. ii. 39 (1832) = **Hymenanthera**, R. Br. (Violar.).
spinosa, G. Don, l. c. = H. dentata?

SOLENANTHUS, Ledeb. Ic. Fl. Ross. t. 26 (1828); Fl. Alt. i. 193 (1829). *BORAGINEAE*, Benth. & Hook. f. ii. 848.
CERINTHOPSIS, Kotschy, ex Benth. & Hook. f. Gen. ii. 849 (1876).
TRACHELANTHUS, Kunze, in Bot. Zeit. viii. (1850) 665.
amplifolius, *Boiss. Diagn.* Ser. I. xi. 126.—Syria.
angustifolius, *Schrenk, Enum. Pl. Nov.* ii. 28.—Sibir.
apenninus, *Hohen. in Bull. Soc. Nat. Mosc.* xi. (1838) 306.—Europ.
baicalensis, DC. & A. DC. in Prod. x. 166 = Craniospermum subvillosum.
Biebersteinii, *DC. l. c.* 165.—Tauria.
brachystemon, *Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc.* xi. (1838) 306.—Reg. Casp.
cerinthoides, *Boiss. Diagn.* Ser. I. xi. 127.—Persia.
circinatus, *Ledeb. Fl. Alt.* i. 194.—Soongar.; Sibir.
conglobatus, A. DC. in DC. Prod. x. 166 = Tournefortii.
coronatus, *Regel, Pl. Nov. Fedsch.* 59.—Turkestan.
dubius, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. xi. (1838) 306 = Biebersteinii.
eriocalycinus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 156.—Reg. Caucas.
hirsutus, *Regel, Pl. Nov. Fedsch.* 60.—Turkestan.
kokanicus, *Regel, l. c.* 89.—Turkestan.
lanatus, *A. DC. in DC. Prod.* x. 165.—Armenia.
mollissimus, *DC. & A. DC. in Prod.* x. 166.—Persia.
nigricans, *Schrenk, Enum. Pl. Nov.* ii. 29.—Sibir.
Olgae, *Regel & Smirnow, ex Regel, Pl. Nov. Fedsch.* 59, 60.—Turkestan.
petiolaris, *DC. Prod.* x. 164.—Mesopot.; Persia.

SOLENANTHUS:—
rumicifolius, Boiss. Pl. Or. Nov. Dec. ii. 9 = petiolaris.
tenuiflorus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 194.—Soongar.
Tournefortii, *DC. Prod.* x. 164.—As. Min.; Persia.

SOLENANTHUS, Steud. teste Klatt, ex Baker, in Journ. Linn. Soc. xvi. (1877) 159 = **Acidanthera**, Hochst. (Irid.).

SOLENARIUM, Dulac, Fl. Hautes-Pyr. 112 (1867) = **Gagea**, Salisb. (Liliac.).
fistulosum, Dulac, l. c. = G. Liotardi.
luteum, Dulac, l. c. = G. fascicularis.

SOLENIDIUM, Lindl. Orch. Linden. 15 (1846). *ORCHIDEAE*, Benth. & Hook. f. iii. 564.
racemosum, *Lindl. l. c.*—N. Granat.

SOLENIPEDIUM, Beer, Prakt. Orch. 310 (1854), sphalm. = **Selenipedium**, Reichb. f. (Orchid.).

SOLENISCIA, DC. Prod. vii. 737 (1839) = **Styphelia**, Sm. (Epacrid.).
elegans, DC. l. c. 738 = Styph. tenuiflora.
pulchella, Stschégl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 3 = Styph. Leucopogon.

SOLENISIA, Steud. Nom. ed. II. ii. 607 (1841), sphalm. = praec.

SOLENIXORA, Baill. in Bull. Soc. Linn. Par. i. (1880) 242. *RUBIACEAE*.
Pervilleana, *Baill. l. c.* 243.—Madag.

SOLENOCARPUS, Wight & Arn. Prod. 171 (1834). *ANACARDIACEAE*, Benth. & Hook. f. i. 422.
indica, *Wight & Arn. l. c.*—Ind. or.

SOLENOGYNE, Cass. in Dict. Sc. Nat. l. 493 (1827) = **Eriocephalus**, Linn. (Compos.).

SOLENOGYNE, Cass. in Dict. Sc. Nat. lvi. 174 (1828) = **Lagenophora**, Cass. (Compos.).
bellidioides, Cass. l. c. 175 = L. Solenogyne.
bellioides, Sond. in Linnaea, xxv. (1852) 480 = L. Emphysopus.
brachycomoides, F. Muell. Fragm. v. 63 = L. Solenogyne.

SOLENOMELUS, Miers, in Proc. Linn. Soc. i. (1841) 122; et Trans. Linn. Soc. xix. (1842) 95. t. 8. *IRIDEAE*, Benth. & Hook. f. iii. 700.
CRUCKSHANKSIA, Miers, Trav. ii. 529 (1826), nomen.
LECHLERA, Griseb. in Lechl. Berb. Am. Austr. 53 (1857).
PHAIOPHLEPS, Rafin. Fl. Tellur. iv. 29 (1836).
acaulis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 121.—Reg. Andin.
andinus, Baker, l. c. = Chamelum andinum.
biflorus, Baker, l. c. = Symphyostemon odoratissimus.
Chamaelum, Baker, l. c. = Chamelum luteum.
chilensis, *Miers, in Proc. Linn. Soc.* i. (1841) 122; *et in Trans. Linn. Soc.* xix. (1842) 96.—Chili.
chimboracensis, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 380 = Orthrosanthus chimboracensis.
frigidus, Klatt, l. c. = Chamelum andinum.
gladioloides, Klatt, l. c. = Orthrosanthus chimboracensis.
Lechleri, *Baker, in Journ. Linn. Soc.* xvi. (1877) 121.—Reg. Andin.
nigricans, *Baker, l. c.*—Chili.
punctatus, Miers, in Proc. Linn. Soc. i. (1841) 122; et in Trans. Linn. Soc. xix. (1842) 96 = chilensis.
spicatus, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 380 = Orthrosanthus spicatus.

SOLENOPHORA, Benth. Pl. Hartw. 68 (1840). *GESNERACEAE*, Benth. & Hook. f. ii. 1005.
ARCTOCALYX, Fenzl, in Otto & Dietr. Allg. Gartenz. xvi. (1848) 305.
HIPPODAMIA, Decne. in Rev. Hortic. (1848) 464.
coccinea, *Benth. Pl. Hartw.* 68.—Mexic.
Endlicheriana, *Hanst. in Linnaea*, xxxiv. (1865-66) 314.—Mexic.
guttata, Decne. ex Benth. & Hook. f. Gen. ii. 1005 = Endlicheriana.
insignis, *Hanst. in Linnaea*, xxxiv. (1865-66) 314.—Mexic.
obscura, *Hanst. l. c.* 315.—Mexic.

SOLENOPSIS, Presl, Prod. Monog. Lobel. 32 (1836) = **Laurentia**, Michx. (Campan.).
Bivonaeana, Presl, l. c. = L. tenella.
canariensis, Presl, l. c. = L. canariensis.
Dregeana, Presl, in E. Mey. Comm. Pl. Afr. Austr. 290 = L. Dregeana.
Laurentia, Presl, Prod. Monog. Lobel. 32 = L. Michelii.
minuta, Presl, l. c. = L. minuta.
Salzmanniana, Presl, l. c. = L. Michelii.
tenella, Steud. Nom. ed. II. ii. 61, in syn. = L. tenella.

SOLENOSPERMUM, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 168. *CELASTRINEAE*.
javanicum, *Zoll. l. c.* 169.—Java.

SOLENOSTEMMA, Hayne, Arzn. Gew. ix. t. 38 (1825). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 749.
ARGELIA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 331. t. 11 G.
Argel, *Hayne, Arzn. Gew.* ix. t. 38.—Nubia.

SOLENOSTEMON, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 271 (1827). *LABIATAE*, Benth. & Hook. f. ii. 1175.
ocymoides, *Schum. & Thonn. l. c.*—Afr. trop.

SOLENOSTIGMA, Endl. Prod. Fl. Norf. 41 (1833) = **Celtis**, Tourn. (Urtic.).
angustifolium, Miq. Fl. Ind. Bat. Suppl. 412 = C. angustifolia.
brevinerve, Blume, Mus. Bot. Lugd. Bat. ii. 67 = C. paniculata.
consimile, Blume, l. c. 68 = C. Wightii.
Djungiel, Blume, l. c. 69 = C. Djungiel.
Hasseltii, Blume, l. c. 68 = C. Hasseltii.
latifolium, Blume, l. c. 67 = C. latifolia.
laurifolium, Blume, l. c. 68 = C. laurifolia.
mauritianum, Blume, l. c. 67 = C. mauritiana.
pacificum, Blume, l. c. = C. pacifica.
paniculatum, Endl. Prod. Fl. Norf. 42 = C. paniculata.
philippinensis, Blume, Mus. Bot. Lugd. Bat. ii. 67 = C. philippinensis.
rigescens, Miq. Fl. Ind. Bat. Suppl. 411 = C. rigescens.
strychnoides, Blume, Mus. Bot. Lugd. Bat. ii. 67 = C. philippinensis.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 411 = C. sumatrana.
Wightii, Blume, Mus. Bot. Lugd. Bat. ii. 68 = C. Wightii.
Zippelii, Blume, l. c. 67 = C. Zippelii.

SOLENOSTIGMA, Klotzsch, ex Walp. Rep. iii. 125 (1844-45) = **Retzia**, Thunb. (Solan.).
Kraussii, Klotzsch, ex Walp. l. c. = R. capensis.

SOLENOSTYLES, Hort. ex Pasq. Cat. Orto Bot. Nap. 97 (1867). *ACANTHACEAE*.
aurantiaca, Hort. ex Pasq. l. c. (Quid?).—Hab.?

SOLENOTHECA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 371 = **Tagetes**, Linn. (Compos.).
tenella, Nutt. l. c. 372 = T. tenuifolia?

SOLENOTINUS, Spach, Hist. Vég. Phan. viii. 305 (1839) = **Viburnum**, Linn. (Caprifol.).
erubescens, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 295 = V. erubescens.
foetens, Oerst. l. c. 294 = V. foetens.
nervosus, Oerst. l. c. 295 = V. nervosum.

SOLENOTUS, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = **Astragalus**, Tourn. (Legumin.).
adsurgens, Stev. l. c. 267 = A. adsurgens.
albicaulis, Stev. l. c. xxix. (1856) ii. 144 = A. albicaulis.
angarensis, Stev. l. c. iv. (1832) 266 = A. angarensis.
asper, Stev. l. c. 267 = A. asper.
bayonnensis, Stev. l. c. = A. bayonnensis.
brachypus, Stev. l. c. xxix. (1856) ii. 144 = A. brachypus.
canadensis, Stev. l. c. iv. (1832) 267 = A. canadensis.
candidissimus, Stev. l. c. xxix. (1856) ii. 144 = A. candidissimus.
cognatus, Stev. l. c. = A. cognatus.
dealbatus, Stev. l. c. iv. (1832) 267 = A. dealbatus.

SOLENOTUS :—
elongatus, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 267 = Astragalus elongatus.
fruticosus, Stev. l. c. 266 = A. fruticosus.
Gebleri, Stev. l. c. xxix. (1856) II. 144 = A. Gebleri.
glaber, Stev. l. c. iv. (1832) 267 = A. glaber.
gyzensis, Stev. l. c. 266 = A. gyzensis.
humifusus, Stev. l. c. 267 = A. humifusus.
hypoglottis, Stev. l. c. = A. hypoglottis.
hyrcanus, Stev. l. c. xxix. (1856) II. 144 = A. hyrcanus.
lasioglottis, Stev. l. c. iv. (1832) 267 = A. lasioglottis.
leontinus, Stev. l. c. = A. leontinus.
macrourus, Stev. l. c. xxix. (1856) II. 144 = A. macrourus.
odoratus, Stev. l. c. iv. (1832) 267 = A. odoratus.
pallescens, Stev. l. c. = A. pallescens.
podocarpus, Stev. l. c. xxix. (1856) II. 144 = A. podocarpus.
purpureus, Stev. l. c. iv. (1832) 267 = A. purpureus.
rariflorus, Stev. l. c. xxix. (1856) II. 144 = A. rariflorus.
resupinatus, Stev. l. c. = A. fragrans.
Schelichovii, Stev. l. c. = A. Schelichowii.
scoparius, Stev. l. c. = A. scoparius.
semibilocularis, Stev. l. c. iv. (1832) 267 = A. semibilocularis.
stenophyllus, Stev. l. c. 266 = A. stenophyllus.
uliginosus, Stev. l. c. 267 = A. uliginosus.
unifultus, Stev. l. c. = A. unifultus.
vesicarius, Stev. l. c. = A. vesicarius.
virgatus, Stev. l. c. 266 = A. virgatus.
xerophilus, Stev. l. c. xxix. (1856) II. 144 = A. xerophilus.

SOLIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3 = **Premna**, Linn. (Verbenac.).

SOLIDAGO, [Vaill.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 256.
AMPHIRAPHIS, DC. Prod. v. 343 (1836).
APLACTIS, Rafin. Fl. Tellur. ii. 42 (1836).
CHRYSOMA, Nutt. in Journ. Acad. Philad. vii. (1834) 67.
DASIORIMA, Rafin. Fl. Tellur. ii. 43 (1836).
DECTIS, Rafin. l. c. (1836).
DORIA, Adans. Fam. ii. 124 (1763).
EUTHAMIA, Ell. Sketch, ii. 391, in syn. (1824).
LEIOLIGO, Rafin. Fl. Tellur. ii. 42 (1836).
LEPIACTIS, Rafin. l. c. 43 (1836).
acaulis, Spreng. Syst. iii. 543 = Liabum acaule.
alata, Wall. ex DC. Prod. vi. 368 = Senecio alatus.
alba, Mill. Gard. Dict. ed. VIII. n. 26 = bicolor.
aliena, Spreng. ex Baker, in Mart. Fl. Bras. vi. III. 9 = Leucopsis Tweedici.
alpestris, Waldst. & Kit. in Willd. Sp. Pl. iii. 218 = S. Virgaurea.
alpina, Linn. Sp. Pl. 880 = Senecio alpinus.
alpina, Salisb. Prod. 199 = minuta.
altissima, Ait. Hort. Kew. ed. I. iii. 212 = rugosa.
altissima, Linn. Sp. Pl. 878 = canadensis.
altissima, Michx. Fl. Am. Bor. ii. 118 = pilosa.
ambigua, Ait. Hort. Kew. ed. I. iii. 217 = S. Virgaurea ?
amplexicaulis, Martens, in Bull. Acad. Brux. viii. (1841) 8 = Riddellii.
amplexicaulis, *Torr. & Gray*, *Fl. N. Am.* ii. 218.—Am. bor.
angulata, Mart. Hort. Erlang. (1814) 180 = patula.
angulosa, Wall. ex DC. Prod. vi. 369 = Senecio densiflorus.
angusta, Torr. & Gray, Fl. N. Am. ii. 204 = petiolaris.
angustifolia, Ell. Sketch, ii. 388 = stricta.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3 = Senecio sarracenicus.
anisata, *Lodd. Cat. ex Loud. Hort. Brit.* 348, nomen. —Am. bor.
arborescens, A. Cunn. ex Hook. f. Fl. N. Zel. i. 148 = Senecio glastifolius, *Hook. f.*
arborescens, Forst. f. Prod. 56 = Olearia nitida.
arcuata, Tausch, in Flora, xi. (1828) 498 = canadensis.
arenaria, Hort. Par. ex Poir. Encyc. Suppl. v. 461 = S. Virgaurea.
argentea, Hornem. Suppl. Hort. Hafn. 99 = caesia.
argentea, Lodd. Cat. ex Loud. Hort. Brit. 348 = puberula.

SOLIDAGO :—
arguta, *Ait. Hort. Kew.* ed. I. iii. 213.—Am. bor.
arguta, Spreng. Syst. iii. 540 = caesia.
arguta, Torr. & Gray, Fl. N. Am. ii. 214 = juncea.
aspera, Ait. Hort. Kew. ed. I. iii. 212 = rugosa.
asperata, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 538 = patula.
asperata, Soland. ex A. Gray, Syn. Fl. N. Am. i. II. 153 = rugosa.
asperula, Desf. Cat. Hort. Par. ed. III. 403 = rugosa ?
asperula, Juss. ex DC. Prod. v. 337 = rigida.
aurea, *Spreng. Syst.* iii. 542 (sp. dub.).—Nepal.
axillaris, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 542 = caesia.
axillaris, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 581 = S. Virgaurea.
azorica, Hochst. in Seubert, Fl. Azor. 31. t. 10 = sempervirens.
bicolor, *Linn. Mant.* i. 114.—Am. bor.
Bigelovii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 80.—Am. bor.
bonariensis, D. Don, ex Baker, in Mart. Fl. Bras. vi. III. 11 = microglossa.
Boottii, *Hook. Comp. Bot. Mag.* i. (1835) 97.—Am. bor.
Bosciana, Moretti, ex DC. Prod. v. 333 = rugosa.
brachyphylla, Chapm. ex Torr. & Gray, Fl. N. Am. ii. 218 = Boottii.
Buckleyi, *Torr. & Gray*, *l. c.* 198.—Am. bor.
caesia, *Linn. Sp. Pl.* 879.—Am. bor.
californica, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 328.—Am. bor. occ.
cambrica, Huds. Fl. Angl. ed. I. 319 = S. Virgaurea.
canadensis, *Linn. Sp. Pl.* 878.—Am. bor.
canescens, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 117 = Olearia Cunninghami.
cantoniensis, Lour. Fl. Cochinch. 501 = S. Virgaurea.
capitata, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) I. 395 = S. Virgaurea.
carinata, Schrad. in DC. Prod. v. 337 = sempervirens.
carnosa, Mill. Gard. Dict. ed. VIII. n. 20 = sempervirens.
carpathica, Schrad. ex Reichb. Fl. Germ. Excurs. 852 = S. Virgaurea.
ceylanica, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 347 = Microglossa zeylanica.
Chapmani, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 80.—Florida.
chilensis, Meyen, Reise, i. 311 = microglossa.
chinensis, Spreng. ex DC. Prod. v. 342 = Senecio vagans.
chrysantha, *Wallr. in Linnaea*, xiv. (1840) 643.—Europ.
ciliaris, Muhl. ex Willd. Sp. Pl. iii. 2056 = juncea.
cinerascens, Schwein. ex Ell. Sketch, ii. 375 = nemoralis.
Cleliae, DC. Prod. v. 331 = elliptica.
compacta, Turcz. in Bull. Soc. Nat. Mosc. (1840) 73 = multiradiata.
conferta, Hort. Par. ex Poir. Encyc. viii. 469 = nemoralis.
conferta, *Mill. Gard. Dict.* ed. VIII. n. 27.—Am. bor.
confertiflora, *DC. Prod.* v. 339.—Am. bor.
confertiflora, Fisch. & Mey. Ind. Sem. Hort. Petrop. vii. 57 = humilis.
confertiflora, Nutt. in Journ. Acad. Philad. vii. (1834) 102 = squarrosa.
confinis, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 191.—Calif.
coquimbana, Phil. in Linnaea, xxxiii. (1864–65) 138 = microglossa.
cordata, Short & Peter, in Trans. Journ. Med. xxviii. (1836) 599 = Brachychaeta cordata.
cordifolia, Moench, Meth. 599 = Aster cordifolius.
corymbosa, *Ell. Sketch*, ii. 378.—Am. bor.
corymbosa, Hort. ex Poir. Encyc. Suppl. v. 461 = S Virgaurea.
corymbosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 328 = multiradiata.
cuneifolia, Roxb. in Beats. Tracts St. Helena, App. 324 = Commidendron spurium.
Curtisii, *Torr. & Gray*, *Fl. N. Am.* ii. 200.—Am. bor.
cuspidata, Wall. Cat. n. 3244 = Inula cuspidata.
cylindrica, *Scheele, in Linnaea*, xxi. (1848) 600.—Am. bor.
Cymbalaria, Moench, Meth. Suppl. 249 = Aster Cymbalariae.
decemflora, DC. Prod. v. 332 = nemoralis.
decemflora, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 223 = S. Radula.

SOLIDAGO :—
decurrens, Lour. Fl. Cochinch. 501 = S. Virgaurea.
deflexa, Moench, Meth. 599 = serotina.
densiflora, Wall. Cat. n. 1821 = Senecio densiflorus.
diffusa, A. Gray, in Proc. Am. Acad. v. (1861) 159 = Bigelowia diffusa.
discoidea, *Torr. & Gray*, *Fl. N. Am.* ii. 195.—Am. bor.
domingensis, *Spreng. Syst.* iii. 639.—Ins. S. Doming.
Doronicum, Linn. Sp. Pl. 880 = Senecio Doronicum.
Drummondii, *Torr. & Gray*, *Fl. N. Am.* ii. 217.—Am. bor.
dubia, *Schult. Obs. Bot.* 178.—Hab. ?
dubia, Scop. Delic. Insub. ii. 19 = elliptica ?
elata, Ell. Sketch, ii. 389 = petiolaris.
elata, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 543 = sempervirens.
elata, Hook. Fl. Bor. Am. ii. 5 = elongata.
Elliottii, *Torr. & Gray*, *Fl. N. Am.* ii. 218.—Am. bor.
elliptica, *Ait. Hort. Kew.* ed. I. iii. 214.—Am. bor.
elliptica, Ell. Sketch, ii. 376 = Elliottii.
elongata, Hort. Par. ex A. Gray, Syn. Fl. N. Am. i. II. 153 = Elliottii.
elongata, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 327.—Am. bor. occ.
eminens, Bischoff, ex A. Gray, Syn. Fl. N. Am. i. II. 157 = canadensis.
erecta, DC. Prod. v. 340 = bicolor.
erecta, Ell. Sketch, ii. 385 = speciosa.
erecta, *Herb. Banks, ex Pursh, Fl. Am. Sept.* ii. 542. —Am. bor.
erecta, Nutt. Gen. Am. ii. 161 = petiolaris.
fistulosa, Mill. Gard. Dict. ed. VIII. n. 19 = pilosa.
flabellata, Schrad. ex Spreng. Syst. iii. 540 = caesia.
flabelliformis, *Wendl. ex DC. Prod.* v. 336.—Hab. ?
flavovirens, *Chapm. Fl. S. U. St.* 211.—Florida.
flexicaulis, Linn. Sp. Pl. 879 = latifolia.
flexuosa, Gilib. Fl. Lituan. i. 203 = S. Virgaurea.
floribundus, Wall. Cat. n. 3111 = Senecio triligulatus.
foliosa, Boj. ex DC. Prod. vi. 377 = Senecio riparius.
foliosa, *Hill*, *Hort. Kew.* 22.—Am. bor.
fragrans, Hort. Par. ex A. Gray, Syn. Fl. N. Am. i. II. 156 = serotina.
fragrans, Willd. Enum. Hist. Berol. Suppl. 59 = elliptica.
Frankii, Hochst. & Steud. ex A. Gray, Syn. Fl. N. Am. i. II. 152 = patula.
fruticosa, *Mill. Gard. Dict.* ed. VIII. n. 32.—Mexic.
fuscata, *Desf. Cat. Hort. Par.* ed. III. 402.—Am. bor.
Gattingeri, *Chapm. ex A. Gray*, *Syn. Fl. N. Am.* i. II. 156.—Am. bor.
genistoides, Bertol. Misc. Bot. vii. 37 = stricta.
gigantea, Ait. Hort. Kew. ed. I. iii. 211 = serotina.
gigantea, Hook. Fl. Bor. Am. ii. 2, partim = lepida.
glaberrima, Martens, in Bull. Acad. Brux. viii. (1841) 68 = missouriensis.
glabra, Desf. Cat. Hort. Par. ed. III. 402 = serotina.
glabra, Mill. Gard. Dict. ed. VIII. n. 22 = sempervirens.
glauca, S. G. Gmel. ex DC. Prod. v. 342 (= *Senecillis glauca*).—Sibir.
glomerata, *Michx. Fl. Bor. Am.* ii. 117.—Am. bor.
glutinosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 328 = confertiflora.
gonoclada, DC. Prod. v. 334 = odora.
gracilis, *Hill*, *Hort. Kew.* 22.—Am. bor.
gracilis, Hort. Par. ex Poir. Encyc. viii. 476 ; Schrad. Hort. Gotting. ex DC. Prod. v. 336 = caesia.
gracillima, *Torr. & Gray*, *Fl. N. Am.* ii. 215.—Florida.
graminifolia, Salisb. Prod. 199 = lanceolata, *Linn.*
grandiflora, Desf. Cat. Hort. Par. ed. III. 403 = littoralis ?
grandiflora, Rafin. in Med. Repos. N. York, v. (1808) 359 = rigida.
graveolens, Lam. Fl. Fr. ii. 145 = Inula graveolens.
Guiradonis, A. Gray, in Bot. Calif. i. 319, partim = spectabilis.
Guiradonis, *A. Gray, in Proc. Am. Acad.* vi. (1865) 543.—Calif.
heterophylla, Nutt. ex A. Gray, Syn. Fl. N. Am. i. II. 148 = multiradiata.
heterotricha, Wall. Cat. n. 3225 = Inula eupatorioides.
hieracioides, Spreng. Syst. iii. 543 = Liabum hieracioides.
hirsuta, L'Hérit. ex DC. Prod. v. 330 = canadensis.
hirsuta, Nutt. in Journ. Acad. Philad. vii. (1834) 103 = bicolor.

SOLIDAGO:—
hirsutissima, Mill. Gard. Dict. ed. VIII. n. 15 = canadensis.
hirta, Willd. Enum. Hort. Berol. 891 = rugosa.
hispida, Muhl. ex Willd. Sp. Pl. iii. 2063 = bicolor *vel* nemoralis.
Houghtoni, *Torr. & Gray, ex A. Gray, Man. Bot. N. U. St.* ed. I. 211.—Am. bor.
humilis, Desf. Cat. Hort. Par. ed. III. 402 = rugosa.
humilis, *Herb. Banks, ex Pursh, Fl. Am. Sept.* ii. 543. —Am. bor.
humilis, Mill. Gard. Dict. ed. VIII. n. 16 = canadensis.
incana, *Torr. & Gray, Fl. N. Am.* ii. 221.—Am. bor.
integerrima, Mill. Gard. Dict. ed. VIII. n. 24 = sempervirens.
integrifolia, Desf. Tabl. ed. I. 103 = sempervirens.
integrifolia, Roxb. in Beats. Tracts St. Helena App. 323 = Melanodendrum integrifolium.
juncea, *Ait. Hort. Kew.* ed. I. iii. 213.—Am. bor.
juncea, DC. Prod. v. 334 = Boottii.
laevigata, Ait. Hort. Kew. ed. I. iii. 215 = sempervirens.
lanata, Hook. Fl. Bor. Am. ii. 4 = bicolor.
lancastriensis, Spreng. Nachr. I. Bot. Gart. Halle, 38 = S. Virgaurea?
lanceolata, Bosc, ex DC. Prod. v. 334 = odora.
lanceolata, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
lanceolata, Cham. & Schlecht. in Linnaea, vi. (1831) 502 = occidentalis.
lanceolata, *Linn. Mant.* i. 114.—Am. bor.
lanceolata, Michx. Fl. Bor. Am. ii. 116 = tenuifolia.
lancifolia, *Chapm. Fl. S. U. St.* 209.—Am. bor.
lapponica, With. Arr. Brit. Pl. ed. III. iii. 728 = S. Virgaurea.
lateriflora, Ait. Hort. Kew. ed. I. iii. 211 = ulmifolia.
lateriflora, Linn. Sp. Pl. 879 = Aster diffusus.
lateriflora, Rafin. ex DC. Prod. v. 335 = caesia.
lateriflora, Willd. Sp. Pl. iii. 2057 = elliptica.
latifolia, *Linn. Sp. Pl.* 879.—Am. bor.
latifolia, Mill. Gard. Dict. ed. VIII. n. 1 = S. Virgaurea.
latissimifolia, Mill. l. c. n. 14 = elliptica.
laxa, Tausch, ex Schrank, in Syll. Ratisb. ii. (1828) 254 = S. Virgaurea.
Leavenworthii, *Torr. & Gray, Fl. N. Am.* ii. 223.—Am. bor.
leiocarpa, DC. Prod. v. 339 = macrophylla.
lepida, *DC. l. c.*—Am. bor.
leptocephala, *Torr. & Gray, Fl. N. Am.* ii. 226.—Am. bor.
Leucadendron, Forst. ex Spreng. Syst. iii. 539 = Senecio Leucadendron.
Leucadendron, Willd. Sp. Pl. iii. 2054 = Senecio prenanthiflorus.
leucanthemifolia, *Baxt. in Loud. Hort. Brit. Suppl.* ii. 673.—Hab.?
limonifolia, Pers. Syn. ii. 449 = sempervirens.
Lindeniana, *A. Rich. Fl. Cub. Fanerog.* ii. 46.—Cuba.
Lindheimeriana, *Scheele, in Linnaea*, xxi. (1848) 599. —Texas.
linearifolia, DC. Prod. v. 341 = microglossa.
linifolia, Linn. Sp. Pl. 881 = Senecio linifolius, *Linn.*
linoides, Soland. ex A. Gray, Syn. Fl. N. Am. i. II. 150 = stricta.
linoides, Torr. & Gray, Fl. N. Am. ii. 216 = neglecta.
lithospermifolia, Willd. Enum. Hort. Berol. 891 = sempervirens.
littoralis, *Savi, ex Pers. Syn.* ii. 450.—Europ. austr.
livida, Willd. Enum. Hort. Berol. 891 = caesia?
longifolia, Schrad. Hort. Gotting. ex DC. Prod. v. 330 = canadensis.
macrophylla, Bigel. Fl. Bost. ed. II. 305 = latifolia.
macrophylla, *Herb. Banks, ex Pursh, Fl. Am. Sept.* ii. 542.—Am. bor.
macrophylla, Schloss. & Vukot. Syll. Fl. Croat. 54 = S. Virgaurea.
macrorrhiza, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 39 = S. Virgaurea.
marginella, DC. Prod. v. 332 = microglossa.
Marshalli, *Rothr. in Wheeler, Rep. Bot.* 146.—Mexic.
marylandica, *Mill. Gard. Dict.* ed. VIII. n. 10.—Am. bor.
mexicana, Berland. ex DC. Prod. v. 332 = velutina.
mexicana, H. B. & K. Nov. Gen. et Sp. iv. 104 = paniculata.
mexicana, Linn. Sp. Pl. 879 = sempervirens.
microglossa, *DC. Prod.* v. 332.—Reg. Argent.

SOLIDAGO:—
minor, Mill. Gard. Dict. ed. VIII. n. 4; Brot. Fl. Lusit. i. 382 = S. Virgaurea.
minuta, *Linn. Sp. Pl.* ed. II. 1235.—Europ. austr.
minuta, Thore, ex Willk. & Lange, Prod. Fl. Hisp. ii. 39 = S. Virgaurea.
missouriensis, *Nutt. in Journ. Acad. Philad.* vii. (1834) 32.—Am. bor.
mollis, Bartl. Ind. Sem. Hort. Gotting. (1836) 5 = nemoralis.
mollis, Rothr. in Wheeler, Rep. Bot. 146 = canadensis.
montana, *Hill, Veg. Syst.* ii. 107 (Quid?).—Gallia.
montana, Linn. Sp. Pl. 881 = Senecio Cacaliaster.
montana, Poir. Encyc. viii. 474 = S. Virgaurea.
monticola, *Chapm. Fl. S. U. St.* 209.—Am. bor.
monticola, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 324 = S. Virgaurea.
Muhlenbergii, Torr. & Gray, Fl. N. Am. ii. 214 = arguta.
multiflora, Desf. Tabl. ed. I. 103 = ulmifolia.
multiradiata, *Ait. Hort. Kew.* ed. I. iii. 218.—Am. bor.
multiradiata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 328 = macrophylla.
nana, *Nutt. l. c.* 327.—Am. bor. occ.
narbonensis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 329.—Gallia.
neglecta, *Torr. & Gray, Fl. N. Am.* ii. 213.—Am. bor.
nemoralis, *Ait. Hort. Kew.* ed. I. iii. 213.—Am. bor.
nepalensis, D. Don, Prod. Fl. Nep. 177 = S. Virgaurea.
nitida, *Torr. & Gray, Fl. N. Am.* ii. 210.—Am. bor.
nitidula, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 10 = microglossa.
noveboracensis, Linn. Sp. Pl. 880 = Aster tataricus?
noveboracensis, Mill. Gard. Dict. ed. VIII. n. 23 = sempervirens.
nudiflora, Viv. Fl. Lib. Spec. 68; Fl. Cors. Sp. Nov. 15 = S. Virgaurea.
nutans, Desf. Cat. Hort. Par. ed. III. 402 = canadensis.
obliqua, Moench, Meth. Suppl. 248 = sempervirens.
obovata, Bertol. Misc. Bot. vii. 36 = puberula.
occidentalis, *Torr. & Gray, Fl. N. Am.* ii. 226.—Am. bor. occ.
occitanea, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 581 = S. Virgaurea.
odora, *Ait. Hort. Kew.* ed. I. iii. 214.—Am. bor.
odora, Chapm. Fl. S. U. St. 213, partim = Chapmani.
odora, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 45, partim = microglossa.
odora, Michx. Fl. Bor. Am. ii. 118 = tortifolia.
ohioënsis, *Riddell, Syn. Fl. West. U. St.* 57.—Am. bor. occ.
palmata, Pall. Reise, iii. 439 = Senecio palmatus.
paniculata, *DC. Prod.* v. 340.—Mexic.
parviflora, Rafin. Ann. Nat. 14 = rupestris.
patagonica, *Phil. in Linnaea*, xxxiii. (1864–65) 137.—Patagon.
patula, *Muhl. in Willd. Sp. Pl.* iii. 2059.—Am. bor.
pauciflora, *Rafin. in Med. Repos. N. York*, v. (1808) 359.—Am. bor.
paucifloculosa, *Michx. Fl. Bor. Am.* ii. 116.—Am. bor.
petiolaris, *Ait. Hort. Kew.* ed. I. iii. 216.—Am. bor.
petiolaris, A. Gray, in Bot. Mex. Bound. 79 = Bigelovii.
petiolaris, Hook. & Arn. Bot. Beech. Voy. 145, partim = californica.
petiolaris, Less. in Linnaea, vi. (1831) 502 = spathulata.
petiolaris, Muhl. Cat. 79 = speciosa.
petiolata, Mill. Gard. Dict. ed. VIII. n. 29 (Quid?).—Am. bor.
pilosa, Mill. l. c. n. 9 = rugosa.
pilosa, *Walt. Fl. Carol.* 207.—Am. bor.
Pitcheri, Nutt. in Journ. Acad. Philad. vii. (1834) 101 = serotina.
plantaginea, Desf. Cat. Hort. Par. ed. III. 402 = elliptica.
polifolia, *Lodd. Cat. ex Loud. Hort. Brit.* 348.—Am. bor.
polyglossa, DC. Prod. v. 332 = microglossa.
praecox, Moench, Meth. 599 = canadensis.
procera, Ait. Hort. Kew. ed. I. iii. 211 = canadensis.
pubens, M. A. Curt. ex Torr. & Gray, Fl. N. Am. ii. 198 = Curtisii.
puberula, Cham. & Schlecht. in Linnaea, vi. (1831) 502 = californica.

SOLIDAGO:—
puberula, DC. Prod. v. 333 = nemoralis.
puberula, *Nutt. Gen. Am.* ii. 162.—Am. bor.
pubescens, Ell. Sketch, ii. 381 = puberula.
pubescens, Wall. Cat. n. 3226 = S. Virgaurea.
pubescens, Walp. Rep. ii. 960 = Curtisii.
pulverulenta, Nutt. Gen. Am. ii. 161 = puberula.
pumila, Crantz, Inst. i. 326 = minuta.
pumila, Pers. Syn. ii. 450 = S. Virgaurea.
pumila, *Torr. & Gray, Fl. N. Am.* ii. 210.—Am. bor.
puncticulata, DC. Prod. v. 332 = odora.
pygmaea, Bertol. in Nov. Comm. Bologn. v. (1842) 215. t. 17. f. 1 = S. Virgaurea.
pyramidalis, *Moench, Meth. Suppl.* 248.—Hab.?
pyramidata, Pursh, Fl. Am. Sept. ii. 537 = pilosa.
Radula, *Nutt. in Journ. Acad. Philad.* vii. (1834) 102. —Am. bor.
recurvata, Mill. Gard. Dict. ed. VIII. n. 28 = rugosa.
recurvata, *Willd. Enum. Hort. Berol.* 889.—Am. bor.
reflexa, Ait. Hort. Kew. ed. I. iii. 211 = canadensis.
repens, D. Don, ex Baker, in Mart. Fl. Bras. vi. III. 11 = microglossa.
reticulata, Lapeyr. Hist. Abr. Pl. Pyr. 520 = S. Virgaurea.
retrorsa, Michx. Fl. Bor. Am. ii. 117 = odora.
retrorsa, Pursh, Fl. Am. Sept. ii. 539 = tortifolia.
retusa, Burm. f. Fl. Cap. Prod. 27 = Nidorella foliosa.
Riddellii, *Frank, in Riddell, Syn. Fl. West U. St.* 57.—Am. bor. occ.
rigida, *Linn. Sp. Pl.* 880.—Am. bor.
rigidula, Bosc, ex DC. Prod. v. 333 = rugosa.
rosmarinifolia, Cass. in Dict. Sc. Nat. lvi. 167 = Olearia rosmarinifolia?
rotundifolia, DC. Prod. v. 339 = S. Radula.
rotundifolia, Roxb. in Beats. Tracts St. Helena App. 324 = Psiadia rotundifolia.
rubricaulis, Wall. Cat. n. 3223 = Inula rubricaulis.
rugosa, *Mill. Gard. Dict.* ed. VIII. n. 25.—Am. bor.
rupestris, *Rafin. Ann. Nat.* 14.—Am. bor.
salicifolia, Salisb. Prod. 199 = stricta.
salicifolia, Wall. ex DC. Prod. v. 343 = Microglossa albescens.
salicina, Ell. Sketch, ii. 389 = patula.
Sarothrae, Pursh, Fl. Am. Sept. ii. 540 = Gutierrezia Euthamiae.
Saulii, Bor. Fl. Centr. Fr. ed. III. ii. 324, in obs. = S. Virgaurea.
scaberrima, Torr. & Gray, Fl. N. Am. ii. 221 = S. Radula.
scabra, Hook. Comp. Bot. Mag. i. (1835–36) 97 = patula.
scabra, Muhl. ex Torr. & Gray, Fl. N. Am. ii. 224 = canadensis.
scabra, Muhl. ex Willd. Sp. Pl. iii. 2059 = rugosa.
scabrida, *DC. Prod.* v. 331.—Mexic.
Schraderi, DC. l. c. 336 = caesia.
scrophulariifolia, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. bor.
semifloculosa, Walp. Rep. ii. 589, in syn. = paucifloculosa.
sempervirens, A. Gray, in Bot. Calif. i. 319 = confinis.
sempervirens, *Linn. Sp. Pl.* 878.—Am. bor.
sempervirens, Michx. Fl. Bor. Am. ii. 119, partim = speciosa.
Sera, J. F. Gmel. Syst. 1236 = serotina.
serotina, *Ait. Hort. Kew.* ed. I. iii. 211.—Am. bor.
serotina, Hook. Comp. Bot. Mag. i. (1835) 97 = missouriensis.
serotina, Retz. Obs. ii. 26 (Quid?).—Am. bor.
serratifolia, Bor. Fl. Centr. Fr. ed. III. ii. 324, in obs. = S. Virgaurea.
serratifolia, Gilib. Fl. Lituan. i. 202 = Senecio paludosus.
Shortii, *Torr. & Gray, Fl. N. Am.* ii. 222.—Am. bor.
sibirica, *Fisch. Hort. Gorenk.* ed. II. (1812) 39, nomen. —Sibir.
simplex, *H. B. & K. Nov. Gen. et Sp.* iv. 163.—Mexic.
sparsiflora, *A. Gray, in Proc. Am. Acad.* xii. (1877) 58. —Am. bor. occ.
spathulata, *DC. Prod.* v. 339.—Mexic.
speciosa, *Nutt. Gen. Am.* ii. 160.—Am. bor.
spectabilis, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 193.—Am. bor.
sphacelata, Rafin. Ann. Nat. (1820) 14 = Brachychaeta cordata.
spiciformis, Torr. & Gray, Fl. N. Am. ii. 202 = spathulata.

SOLIDAGO:—
spithamaea, *M. A. Curt. ex A. Gray, in Am. Journ. Sc.* Ser. I. xlii. (1842) 42.—Am. bor.
spuria, Forst. in Comm. Gotting. ix. (1789) 68 = Commidendrum spurium.
squarrosa, *Muhl. Cat.* 79; *Nutt. Gen. Am.* ii. 161.—Am. bor.
squarrosa, Nutt. in Journ. Acad. Philad. vii. (1834) 102 = petiolaris.
stricta, *Ait. Hort. Kew.* ed. I. iii. 216.—Am. bor.
stricta, Hook. Fl. Bor. Am. ii. 4 = humilis, uliginosa.
stricta, Less. in Linnaea, vi. (1831) 502 = elongata.
stricta, *Moench, Meth. Suppl.* 249.—Hab. ?
suaveolens, *J. F. Gmel. Syst.* 1235.—Hab. ?
tardiflora, Moench, Meth. 599 = sempervirens.
tenuifolia, *Pursh, Fl. Am. Sept.* ii. 540.—Am. bor.
Terrae-Novae, *Torr. & Gray, Fl. N. Am.* ii. 206.—Am. bor.
thyrsoidea, E. Mey. Pl. Labrad. 63 = macrophylla.
Tolmieana, *A. Gray, Syn. Fl. N. Am.* i. ii. 151.—Am. bor. occ.
tortifolia, *Ell. Sketch*, ii. 377.—Am. bor.
Tournefortii, Tausch, in Flora, xi. (1828) 500 = canadensis.
uliginosa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 101.—Am. bor.
ulmifolia, Hook. Comp. Bot. Mag. i. (1835) 97 = Drummondii.
ulmifolia, *Muhl. ex Willd. Sp. Pl.* iii. 2060.—Am. bor.
urticaefolia, Mill. Gard. Dict. ed. VIII. n. 30 = Calea urticaefolia.
valesiaca, Bor. ex Déségl. in Feuilles Jeunes Nat. viii. (1877) 18 = S. Virgaurea.
velutina, *DC. Prod.* v. 332.—Mexic.
varia, Hill, Hort. Kew. 21 (Quid ?).—Am. bor.
verna, *M. A. Curt. in Torr. & Gray, Fl. N. Am.* ii. 205.—Am. bor.
verrucosa, Schrad. Hort. Gotting. 12. t. 6 = elliptica *vel* arguta.
villosa, Ell. Sketch, ii. 372 = pilosa.
villosa, Pursh, Fl. Am. Sept. ii. 537 = rugosa.
viminea, Ait. Hort. Kew. ed. I. iii. 215 = sempervirens.
viminea, Bosc, ex DC. Prod. v. 341 = bicolor.
Virga, Dulac, Fl. Hautes-Pyr. 570 = S. Virgaurea.
Virga-aurea, Auct. = S. Virgaurea.
virgata, Michx. Fl. Bor. Am. ii. 117 = stricta.
virgata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = S. Virgaurea.
Virgaurea, Bigel. Fl. Bost. ed. II. 306 (excl. var.) = macrophylla.
Virgaurea, *Linn. Sp. Pl.* 880.—Reg. bor. temp.
virginiana, Mill. Gard. Dict. ed. VIII. n. 11 = rugosa.
viscosa, Lam. Fl. Fr. ii. 144 = Inula viscosa.
viscosa, Schrad. Sert. Hannov. 12 = Psiadia glutinosa.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 2; Lam. Fl. Fr. ii. 145 = S. Virgaurea.
vulneraria, Mart. Syst. Mat. Med. Bras. 9, nomen = microglossa.
Wrightii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 80.—Texas.
zeylanica, Hook. f. Fl. Brit. Ind. iii. 257 = *ceylanica*.

SOLIERA, Clos, in C. Gay, Fl. Chil. iv. 489. t. 53 (1849). *LABIATAE*, Benth. & Hook. f. ii. 1187.
pulchella, *Clos, l. c.*—Chili.

SOLIVA, Ruiz & Pav. Prod. 113. t. 24 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 430.
Gymnostyles, Juss. in Ann. Mus. Par. iv. (1804) 258. t. 61.
acaulis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 326 (araulis).—Reg. Argent.
aculeata, Ruiz & Pav. ex Phil. f. Cat. Pl. Vasc. Chil. 185 = Plagiocheilus solivaeformis.
alata, DC. Prod. vi. 143 = sessilis.
anthemifolia, *R. Br. in Trans. Linn. Soc.* xii. (1817) 102.—Am. austr.; Austral.
aphanes, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
Barcklayana, DC. Prod. vi. 143 = sessilis.
daucifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 403 = sessilis.
dioica, *Sch. Bip. in Lechl. Berb. Am. Austr.* 53.—Chili.
lusitanica, Less. Syn. Comp. 268 = nasturtiifolia.
mexicana, DC. Prod. vi. 143 = Cotula pygmaea.
microloma, Phil. in Anal. Univ. Chil. (1865) ii. 331.—Chili.
minuta, Sweet, Hort. Brit. ed. II. 317 = Cotula pygmaea.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iv. 302.—N. Granat.
nasturtiifolia, *DC. Prod.* vi. 142.—Reg. Argent.
pedicellata, Ruiz & Pav. Syst. Veg. 215 = Cotula pygmaea ?
pedunculata, Ruiz & Pav. ex Steud. Nom. ed. II. ii. 609, sphalm. = praec.
pterosperma, Less. Syn. Comp. 268 = sessilis.
pygmaea, H. B. & K. Nov. Gen. et Sp. iv. 303 = Cotula pygmaea.
radiata, Poepp. ex DC. Prod. vii. 288 = Blennosperma chilense.
sessilis, *Ruiz & Pav. Prod.* 113. t. 24; *Syst. Veg.* 215.—Am. bor. et austr.
stolonifera, Loud. Hort. Brit. 364 = nasturtiifolia.
tenella, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 128 = Cotula australis.
triniifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 202.—Reg. Argent.
urbica, Phil. in Linnaea, xxix. (1857–58) 7 = nasturtiifolia.
valdiviana, *Phil. l. c.* xxxiii. (1864–65) 168.—Chili.

SOLIVAEA, Cass. in Dict Sc. Nat. xxix. 177 (1823) = praec.

SOLLYA, Lindl. Bot. Reg. (1830) t. 1466. *PITTOSPOREAE*, Benth. & Hook. f. i. 133.
Xerosollya, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 362.
angustifolia, Lindl. Bot. Reg. sub t. 1466 = Billardiera scandens.
Drummondi, C. Morr. in Belg. Hortic. iv. (1854) 196. t. 33. f. 3–8 = heterophylla ?
fusiformis, Hort. ex Payer, Organog. t. 34 = heterophylla.
heterophylla, *Lindl. Bot. Reg.* t. 1466.—Austral.
linearis, Lindl. l. c. (1839) Misc. 83 = heterophylla.
parviflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 361.—Austral.
salicifolia, Marnoek, Flor. Mag. iv. (1839–40) 62, 136. t. 44. f. 2 = heterophylla ?

SOLMSIA, Baill. Adansonia, x. (1871) 37. *TILIACEAE*.
calophylla, *Baill. l. c.* 38.—N. Caled.
chrysophylla, *Baill. l. c.*—N. Caled.

SOLORI, Adans. Fam. ii. 327 (1763) = **Dalbergia**, Linn. (Legumin.).

SOLSTITIARIA, Hill, Veg. Syst. iv. 21 (1762) = **Centaurea**, Linn. (Compos.).
flava, Hill, Herb. Brit. i. 77 = C. solstitialis.
melitensis, Hill, Hort. Kew. 62 = C. melitensis.
solstitialis, Hill, l. c. = C. solstitialis.

SOLTMANNIA, [Klotzsch], ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 132 = **Miconia**, Ruiz & Pav. (Melastom.).
splendens, Klotzsch, ex Naud. l. c. = M. mexicana.

SOMERA, Salisb. Gen. Pl. Fragm. 26 (1866) = **Scilla**, Linn. (Liliac.).
italica, Salisb. l. c. 27 = Scilla italica.

SOMERAUERA, Hoppe, in Flora, ii. (1819) 26 = **Arenaria**, Rupp. (Caryoph.).
quadrifaria, Hoppe, l. c. = A. aretioides.
quadrifida, Reichb. Ic. Fl. Germ. v. t. 204 = praec.

SOMMEA, Bory, in Ann. Gén. Sc. Phys. vi. (1820) 94. t. 87 = **Acicarpha**, Juss. (Calycer.).
Calcitrapa, Bory, l. c. = A. spathulata.

SOMMERA, Schlecht. in Linnaea, ix. (1834) 602. *RUBIACEAE*, Benth. & Hook. f. ii. 79.
arborescens, *Schlecht. l. c.*—Mexic.
grandis, Griseb. Fl. Brit. W. Ind. 326 = arborescens.

SOMMERAUERA, Endl. Ench. 505 (1841), sphalm. = SOMERAUERA, Hoppe = **Arenaria**, Rupp. (Caryoph.).

SOMMERFELDTIA, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 331 (1827) = **Drepanocarpus**, G. F. W. Mey. (Legumin.).
obovata, Schum. l. c. = D. lunatus.

SOMMERFELTIA, Less. Syn. Comp. 189 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 275.
spinulosa, *Less. l. c.* 190.—Bras.

SOMMERINGIA, Lindl. Veg. Kingd. 554 (1847), sphalm. = **Soemmeringia**, Mart. (Legumin.).

SOMMIERA, Benth. & Hook. f. Gen. iii. 909 (1883) err. typ. = seq.

SOMMIERIA, Becc. Malesia, i. (1877) 66. *PALMAE*, Benth. & Hook. f. iii. 909.
elegans, *Becc. l. c.* 68.—N. Guin.
leucophylla, *Becc. l. c.* 67.—N. Guin.

SOMPHOXYLON, Eichl. in Flora, xlvii. (1864) 396; et in Mart. Fl. Bras. xiii. 1. 205. t. 37 (1864), Genus dubium *MENISPERMACEARUM*.—Cf. Benth. & Hook. f. Gen. i. 959.
Wullschlagelii, *Eichl. ll. cc.*—Guian.

SONCHIDIUM, Pomel, Nouv. Mat. Fl. Atl. 6 (1874) = **Sonchus**, Tourn. (Compos.).
maritimum, Pomel, l. c. 7 = Sonchus maritimus.
palustre, Pomel, l. c. = Sonchus palustris.

SONCHOS, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = **Sonchus**, Tourn. (Compos.).

SONCHOSERIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 102 = **Sonchus**, Tourn. (Compos.).
arvensis, Fourr. l. c. 103 = Sonchus arvensis.
decora, Fourr. l. c. 102 = Sonchus decorus.
maritima, Fourr. l. c. = Sonchus maritimus.
palustris, Fourr. l. c. 103 = Sonchus palustris.

SONCHUS, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 528.
Atalanthus, Don, in Edinb. N. Phil. Journ. (1828–29) 311.
Sonchidium, Pomel, Nouv. Mat. Fl. Atl. 6 (1874).
Sonchoseris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 102.
Trachodes, D. Don, in Trans. Linn. Soc. xvi. (1830) 182.
abbreviatus, Link, in Buch, Beschr. Canar. Ins. 146 = congestus.
acanthifolius, [Clairv.] Man. Herb. 231 = Lactuca Plumieri.
acanthoides, Steud. Nom. ed. II. ii. 610 = Lactuca Plumieri.
acaulis, Dum.-Cours. Bot. Cult. ed. II. iv. 12 = congestus.
acidus, *Schousb. ex Willd. Sp. Pl.* iii. 1511.—Marocc.
acuminatus, Bigel. Fl. Bost. ed. II. 290 = Lactuca alpina.
acuminatus, Willd. Sp. Pl. iii. 1521 = Lactuca canadensis.
adscendens, Boj. ex DC. Prod. vi. 303 = Emilia adscendens.
agrestis, Sw. Prod. Veg. Ind. Occ. 110 = Erechtites praealta.
albanus, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 59 = Lactuca racemosa.
alpestris, [Clairv.] Man. Herb. 231 = Lactuca alpina.
alpinus, Linn. Sp. Pl. 794 = Lactuca alpina.
altissimus, Vahl, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 431 = congestus.
amboinensis, Hort. ex Steud. Nom. ed. II. ii. 610 = Emilia flammea.
amplexicaulis, *Gueldenst. Reisen*, ii. 209; *ex Ledeb. Fl. Ross.* ii. 836.—Rossia.
angustifolius, Desf. Fl. Atlant. ii. 225 = fruticosus.
angustifolius, Neck. Delic. Gallo-Belg. ii. 326 = maritimus.
angustissimus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 203.—Afr. trop.
aquatilis, Pourr. in Mém. Acad. Toul. iii. (1788) 330 = maritimus.
arborescens, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 549 = tenerrimus.
arboreus, *DC. Prod.* vi. 189.—Ins. Canar.
arvensis, *Linn. Sp. Pl.* 793.—Europ.; As. bor.
asper, Gaertn. Mey. & Scherb. Fl. Wett. iii. 125 = oleraceus.
asper, *Hill, Herb. Brit.* i. 47; *Vill. Hist. Fl. Dauph.* iii. 158.—Cosmop.

SONCHUS:—
asper, Wulf. ex DC. Prod. vii. 190 = Picris hieracioides.
australis, Hort. ex Trev. Ind. Sem. Hort. Vratisl. (1818) 6 = longifolius.
azureus, Ledeb. Fl. Alt. iv. 138 (= *Mulgedium azureum*).—Sibir.
balthicus, Fries, ex Link, Handb. i. 784, in syn. = maritimus.
biennis, Moench, Meth. 545 = Lactuca floridana.
Bipontini, Aschers. in Schweinf. Beitr. Fl. Aethiop. 160 = lactucoides.
Bocconi, Jan, Cat. 658; ex DC. Prod. vii. 183 = chondrilloides.
Bollei, Sch. Bip. ex Bolle, in Bonplandia, vii. (1859) 298 = gorgadensis.
Bourgeauii, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 446.—Ins. Canar.
brachylobus, *Webb & Berth. l. c.* 438.—Ins. Canar.
brachyotus, DC. Prod. vii. 186 = arvensis.
brasiliensis, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 293 = Erechtites praealta.
Broussonetii, Desf. Tabl. ed. II. 101 = congestus.
Brunneri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 459.—Afr. trop.
bupleurifolius, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 797.—Armenia.
cacaliaefolius, Bieb. Fl. Taur. Cauc. ii. 241 (= *Mulgedium cacaliaefolium*).—Reg. Cauc.
caeruleus, Sm. Fl. Brit. 815 = Lactuca alpina.
californicus, Hook. & Arn. Bot. Beech. Voy. 361 = Malacothrix saxatilis.
calyculatus, Roxb. ex DC. Prod. vii. 140 = Lactuca Heyneana.
canadensis, Froel. in Usteri, Ann. Bot. i. (1791) 29 = Lactuca macrophylla.
canadensis, Linn. Sp. Pl. 793 = Lactuca alpina.
Candolleanus, Jaub. & Spach, Illustr. iii. 110 = Launaea chondrilloides.
capensis, *Schlecht. Hort. Hal.* (1850) 7.—*Cf. Linnaea*, xxiv. (1851) 211.—Afr. austr.
capitatus, Spreng. Syst. iii. 650 = Launaea glomerata.
carolinianus, Walt. Fl. Carol. 192 = asper.
Cassinianus, Jaub. & Spach, Illustr. iii. 112 (= *Zollikofera Cassiniana*).—Aegypt.
caucasicus, *Biehler, Cent. Pl. Rar.* 12. n. 31; *et ex Hornem. Suppl. Hort. Hafn.* 88.—Reg. Cauc.
caucasicus, *Spreng. Fl. Hal. Mant.* i. 49.—Reg. Cauc.
cervicornis, Nym. Syll. 38 = Launaea spinosa.
cervicornis, *Willk. Ind. Sem. Hort. Prag.* (1875).—Hab.?
chinensis, *Fisch. Hort. Gorenk.* ed. II. (1812) 33, nomen; *et ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 65.—China.
chondrilloides, *Desf. Fl. Atlant.* ii. 226.—Alger.
chondrilloides, Sibth. & Sm. Fl. Graec. viii. 67 = Picridium vulgare.
chuquitensis, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 294 = congestus.
cichoraceus, Moench, Meth. Suppl. 222 = Lactuca Plumieri.
ciliatus, Lam. Fl. Fr. ii. 87 = oleraceus.
ciliatus, Perr. ex DC. Prod. vii. 177 = Lactuca goraeënsis.
coeruleus, Auct. = caeruleus.
congestus, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 136.—Ins. Canar.
cordifolius, Desf. Tabl. ed. I. 87 = Lactuca macrophylla.
cornutus, *Hochst. ex Steud. Nom.* ed. II. ii. 610, nomen; *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 459.—Afr. trop.
crassifolius, *Pourr. ex Willd. Sp. Pl.* iii. 1509.—Hispan.
crepioides, *Sm. in Rees, Cycl.* xxxiii. n. 24.—Afr. bor.
crispus, *Poir. Encyc. Suppl.* iii. 289.—Ins. Canar.
cristulatus, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 318 (= *Dubyaea cristulata*).—Austral.
crocifolius, Hort. ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 449 = asper.
Cumbulus, Buch.-Ham. ex Wall. Cat. n. 3251 = arvensis.
cuspidatus, *Blume, Bijdr.* 888.—Java.
cyaneus, D. Don, Prod. Fl. Nep. 164 = Lactuca hastata.
Daltonii, *Webb, in Hook. Niger Fl.* 144. t. 15.—Ins. Cap. Virid.
decorus, *Cast. Cat. Pl. Env. Marseille*, 91.—Hispan.

SONCHUS:—
dentatus, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 23 = ustulatus.
dentatus, *Ledeb. Fl. Alt.* iv. 141.—Sibir. altaic.
dichotomus, Bieb. Fl. Taur. Cauc. ii. 240 = Picridium dichotomum.
dichotomus, Savi, Cose Bot. 10 = Launaea nudicaulis.
divaricatus, Buch, Beschr. Canar. Ins. 147 = nudicaulis.
divaricatus, Desf. in Ann. Mus. Par. ii. (1803) 212 = Launaea nudicaulis.
divaricatus, Sieber, ex Steud. Nom. ed. II. ii. 610 = Picridium dichotomum.
Dregeanus, *DC. Prod.* vii. 184.—Afr. austr.
dulosus, *Fisch. & Mey. ex Sweet, Hort. Brit.* ed. III. 414.—Hab.?
Ecklonianus, *DC. Prod.* vii. 184.—Afr. austr.
elongatus, *Kunze, ex Schlecht. Ind. Sem. Hort. Hal.* (1849) 7.—Hab.?
eriopus, *Royle, Illustr. Bot. Himal.* 247.—Ind. or.
eryngioides, DC. Prod. vii. 185 = asper.
erythropappus, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 293 = Erechtites valerianaefolia.
exaltatus, Wallr. in Linnaea, xiv. (1840) 659 (Beitr. Fl. Hercyn. 289) = arvensis?
fallax, Wallr. Ann. Bot. 98; Sched. Crit. 432 = asper.
ferox, Wall. Cat. n. 3248 = asper.
flexuosus, *Ledeb. Hort. Dorpat. Suppl.* i. 5.—Reg. Cauc.
floridanus, Ait. Hort. Kew. ed. I. iii. 116 = Lactuca alpina.
floridanus, Linn. Sp. Pl. 794 = Lactuca floridana.
floridanus, Lour. Fl. Cochinch. 480 = Lactuca brevirostris.
floridanus, Michx. Fl. Bor. Am. ii. 85, partim = Lactuca acuminata.
fragilis, *Ball, in Journ. Bot.* xi. (1873) 372.—Afr. bor. occ.
frondosus, Salisb. Prod. 180 = congestus.
fruticosus, Jacq. Coll. i. 13 = congestus.
fruticosus, *Linn. f. Suppl.* 346.—Ins. Mader.
fruticosus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 27 = Centratherum frutescens (in Addend.).
gibbosus, *Savi, Fl. Ital.* i. 47.—Italia.
glaber, Gilib. Fl. Lituan. i. 242 = oleraceus.
glaber, Schult. Obs. Bot. 162 = arvensis.
glaber, Thunb. Prod. Pl. Cap. 139 = asper.
glandulosus, Schur, Enum. Pl. Transs. 371 = arvensis.
glaucescens, *Ford. Obs. Pl. Crit.* v. (1847) 75. t. 5.—Reg. Mediterr.
glaucus, *Thunb. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 135.—Hab.?
glutinosus, Lam. Fl. Fr. ii. 88 = Hieracium glutinosum.
goraeënsis, Lam. Encyc. iii. 397 = Lactuca goraeënsis.
goraeënsis, Less. ex Steud. Nom. ed. II. ii. 610 = Prenanthes alata.
gorgadensis, *Bolle, in Bonplandia*, vii. (1859) 298.—Ins. Canar.
graecus, Reut. ex E. Weiss, in Verh. Zool.-Bot. Ges. Wien, xix. (1869) 45 = glaucescens.
gramineus, Boj. ex DC. Prod. vi. 303 = Emilia graminea.
gummifer, *Link, in Buch, Beschr. Canar. Ins.* 146.—Ins. Canar.
hantoniensis, Sweet, Hort. Brit. ed. II. 278 = arvensis?
hastatus, Less. in Linnaea, vi. (1831) 99 = Prenanthes alata.
hastatus, Wall. Cat. n. 3250 = Lactuca hastata.
hieracioides, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 240.—Hispan.
hispanicus, Jacq. Hort. Schoenb. ii. 9 = Picridium tingitanum.
hispidulus, Bové, ex DC. Prod. vii. 182 = Picridium tingitanum.
hispidus, Gilib. Fl. Lituan. i. 241 = arvensis.
hispidus, Ledeb. Fl. Alt. iv. 140 (= *Mulgedium hispidum*).—Sibir.
Hochstetteri, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 160 = Lactuca goraeënsis.
Hotha, *C. B. Clarke, Comp. Ind.* 276.—Burma.
hyoseridifolius, Hornem. Hort. Hafn. ii. 752 = ustulatus.
hypochaeroides, *Schlecht. Ind. Sem. Hort. Hal.* (1850) 7.—*Cf. Linnaea*, xxiv. (1851) 212.—Hab.?
infestus, Poepp. ex DC. Prod. vii. 185 = asper.
integrifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 528.—Afr. austr.
intermedius, Jan, ex DC. Prod. vii. 182 = Picridium intermedium.

SONCHUS:—
intermedius, Koch, Syn. Fl. Germ. ed. I. 434, cum. cit. falsa = arvensis.
intermedius, Krock. Fl. Siles. Suppl. ii. 234 (Quid?).—Europ.
italicus, Spreng. Syst. iii. 651 = tenerrimus.
Jacquini, DC. Cat. Hort. Monsp. (1813) 147 = congestus.
javanicus, Spreng. Syst. iii. 648 = Prenanthes javanica.
lacerus, Willd. Sp. Pl. iii. 1513 = oleraceus.
laciniatus, *T. Moore, in Flor. & Pom.* (1871) 138.—Hab.?
lactucoides, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 114 (Enum. Pl. Chin. Bor. 40) = Lactuca tatarica.
lactucoides, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 467.—Abyss.
laevigatus, *Willd. Enum. Hort. Berol. Suppl.* 54.—Ins. Mader.
laevis, Vill. Hist. Pl. Dauph. iii. 158 = oleraceus.
laevissimus, Schur, Enum. Pl. Transs. 371 = arvensis.
lampsanoideus, Lam. Fl. Fr. ii. 90 = Crepis lampsanoideus.
lapponicus, Froel. in Usteri, Ann. Bot. i. (1791) 26, in textu = Lactuca floridana?
leptocephalus, *Cass. in Dict. Sc. Nat.* xliii. 281.—Ins. Canar.
leucodon, Fisch. & Mey. ex Ledeb. Fl. Ross. ii. 832 = Launaea nudicaulis.
leucophaeus, Willd. Sp. Pl. iii. 1520 = Lactuca alpina.
libycus, Spach, ex Boiss. Fl. Orient. iii. 823 = fruticosus.
littoralis, Reichb. Fl. Germ. Excurs. 274 = maritimus.
longifolius, *Trev. Ind. Sem. Hort. Vratisl.* (1818) 6; *et in Nov. Act. Nat. Cur.* xiii. (1826) 197.—Hab.?
longifolius, Wall. Cat. n. 3251 = arvensis.
ludovicianus, Nutt. Gen. Am. ii. 125 = Lactuca ludoviciana.
luxurians, Schrad. ex Steud. Nom. ed. I. 799 = Launaea macrophylla.
lyratus, Willd. Enum. Hort. Berol. Suppl. 53 = laevigatus.
macranthus, Poir. Encyc. Suppl. iii. 289 = congestus.
macrophyllus, Willd. Sp. Pl. iii. 1519 = Lactuca macrophylla.
macrotus, Fenzl, in Flora, xxvii. (1844) 312 = oleraceus.
malayanus, *Miq. Fl. Ind. Bat.* ii. 113.—Java.
maritimus, Echeand. ex Losc. & Pard. Ser. Inconf. ed. I. 64 = palustris.
maritimus, *Linn. Syst.* ed. X. 1192.—Europ.; Reg. Mediterr.
maritimus, Pall. ex Ledeb. Fl. Ross. ii. 843 = Lactuca sibirica.
maritimus, Turcz. in Bull. Soc. Nat. Mosc. (1838) 96 = arvensis.
massaviensis, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 160 = Heterachaena massauiensis.
mauritanicus, *Boiss. & Reut. Pugill. Pl. Nov.* 70.—Alger.
melanolepis, *Fresen. in Mus. Senckenb.* iii. (1839) 76.—Abyss.
Metzianus, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 396 = Crepis acaulis.
montanus, Lam. Encyc. iii. 401 = Lactuca alpina.
multiflorus, Desf. Cat. Hort. Par. ed. III. 145 = Lactuca alpina.
nanus, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 528.—Afr. austr.
nigricans, *Tausch, in Flora*, xii. (1829) 35.—Ital.
nitidus, Vill. Prosp. 33; Hist. Pl. Dauph. iii. 160 = arvensis.
Novae-Zelandiae, *Benth. & Hook. f. Gen.* ii. 528.—N. Zel.
nudicaulis, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 427.—Aegypt.
Nymani, Tineo & Guss. in Guss. Fl. Sic. Syn. ii. 860 = arvensis.
occidentalis, Spreng. Neue Entdeck. ii. 143 = Erechtites praealta.
oleraceus, *Linn. Sp. Pl.* 794.—Cosmop.
oleraceus, Wall. Cat. n. 3252 F = asper.
oreophilus, *Miq. Fl. Ind. Bat.* ii. 114.—Java.
orixensis, Roxb. Hort. Beng. 60; Fl. Ind. iii. 402 = arvensis.
oxyspermus, Wallr. in Linnaea, xiv. (1840) 658 (Beitr. Fl. Hercyn. 288) = tenerrimus?
pallescens, *Pančić. Elench. Crna Gora*, 55.—Montenegr.

SONCHUS:—
pallidus, Torr. Compend. Fl. N. St. 279 = Lactuca alpina.
pallidus, Willd. Sp. Pl. iii. 1521 = Lactuca canadensis.
paludosus, Gueldenst. ex Ledeb. Fl. Ross. ii. 836 = palustris.
palustris, *Linn. Sp. Pl.* 793.—Europ.; As. Min.
paniculatus, *Moç. & Sesse, ex DC. Prod.* vii. 191.—Mexic.
parviflorus, Lej. ex Reichb. Fl. Germ. Excurs. 274 = oleraceus.
pauciflorus, *Baker, in Journ. Bot.* xx. (1882) 171.—Madag.
pauciflorus, Echeand. ex Willk. & Lange, Prod. Fl. Hisp. ii. 240 = maritimus.
pectinatus, DC. Rapp. Voy. ii. 78 = tenerrimus.
pendulus, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 421 = Prenanthes pendula.
pensylvanicus, Spreng. ex Steud. Nom. ed. II. ii. 611 (Quid?).—Am. bor.?
pentaphyllus, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 160 = Lactuca taraxacifolia.
perfoliatus, Gueldenst. ex Ledeb. Fl. Ross. ii. 836 (Quid?).—Rossia.
picroides, Lam. Encyc. iii. 398 = Picridium vulgare.
picroides, Sibth. & Sm. Fl. Graec. viii. 68 = Picridium intermedium.
pinnatifidus, Cav. in Anal. Cienc. Nat. iv. (1801) 78 = acidus.
pinnatifidus, Lag. ex DC. Prod. vii. 182 = Picridium tingitanum.
pinnatus, *Ait. Hort. Kew.* ed. I. iii. 116.—Ins. Mader.
platylepis, *Webb & Berth. Phyt. Canar.* ii. 433.—Ins. Canar.
Plumerianus, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 171 = seq.
Plumerii, St. Lag. l. c. = seq.
Plumieri, Linn. Syst. ed. X. 1192 = Lactuca Plumieri.
Plumieri, Panč. ex Vis. in Mem. Ist. Venet. ix. (1860) 173 (= *Mulgedium Pančičii*).—Europ. austr.
polyodon, Webb & Berth. Phyt. Canar. ii. 431 = congestus.
pratensis, Schur, Enum. Pl. Transs. 968 (Index), sphalm. = arvensis.
prenanthoides, Bieb. Fl. Taur. Cauc. ii. 242 (= *Mulgedium prenanthoides*).—Reg. Cauc.
prenanthoides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 459.—Afr. trop.
pulchellus, Pursh, Fl. Am. Sept. ii. 502 = Lactuca pulchella.
pustulatus, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 242.—Hispan.
quercifolius, *Desf. Fl. Atlant.* ii. 225. t. 213.—Afr. bor.
racemosus, Lam. Encyc. iii. 400 = Lactuca alpina.
radiatus, Link, Enum. Hort. Berol. ii. 279, sphalm. = seq.
radicatus, *Ait. Hort. Kew.* ed. I. iii. 116.—Ins. Mader.; Teneriff.
rarifolius, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 460.—Afr. trop.
reflexus, Lag. Elench. 15; ex ej. Gen. et Sp. Nov. 24 = Launaea nudicaulis.
reversus, E. Mey. ex DC. Prod. vii. 186 = oleraceus.
rivularis, *Phil. in Linnaea*, xxx. (1859–60) 194.—Chili.
robustus, Wall. Cat. n. 3249 = Lactuca hastata.
roseus, Bess. in Spreng. Syst. iii. 651 = oleraceus.
Royleanus, DC. Prod. vii. 184 = oleraceus.
runcinatus, *Vent. ex Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 434.—Ins. Canar.
sagittatus, Moench, Meth. 544 = palustris.
Schimperi, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1857) App. 1 = oleraceus.
Schmidianus, C. Koch, l. c. (1853) App. 12 = oleraceus.
Schweinfurthii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 458.—Afr. trop.
scorzoneraeformis, Lag. Gen. et Sp. Nov. 24 (= *Zollikoferia pumila*).—Hispan.; Arab.
serratus, Perr. ex DC. Prod. vii. 138 = Lactuca taraxacifolia.
sespedalis, Gilib. Fl. Lituan. i. 241 = palustris.
setosus, Ehrenb. ex Sweet, Hort. Brit. ed. III. 414 (Quod?).—Hab.?
shzucinianus, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 189 = arvensis.
sibiricus, Bieb. Fl. Taur. Cauc. ii. 240 = Lactuca racemosa.

SONCHUS:—
sibiricus, Linn. Sp. Pl. 795 = Lactuca sibirica.
sibiricus, Richards. App. Frankl. 1st Voy. ed. II. 30 = Lactuca pulchella.
simplicissimus, Lag. Gen. et Sp. Nov. 24 = crassifolius.
Spachii, *Schweinf. Beitr. Fl. Aethiop.* 307.—Nubia.
spicatus, Lam. Encyc. iii. 401 = Lactuca alpina.
spinosus, DC. Prod. vii. 189 = Launaea spinosa.
spinosus, Lam. Fl. Fr. ii. 86 = asper.
spinosus, *Marès, Cat. Pl. Ins. Baléar.* 172.—Ins. Balear.
spinulosus, Bigel. Fl. Bost. ed. II. 292 = asper.
squamosus, Lam. Fl. Fr. ii. 87 = Picridium vulgare.
squarrosus, DC. Cat. Hort. Monsp. (1813) 147 = fruticosus.
subacaulis, Dum.-Cours. ex Steud. Nom. ed. II. ii. 611 (Quid?).—Hab.?
sulphureus, Boiss. Fl. Orient. iii. 796 = asper.
sundaicus, *Blume, Bijdr.* 888.—Java.
taraxacifolius, Spreng. Syst. iii. 649 = Lactuca intybacea.
taraxacifolius, Willd. Sp. Pl. iii. 1511 = Lactuca taraxacifolia.
tataricus, Linn. Mant. ii. 572 = Lactuca tatarica.
tenax, Fisch. ex Sweet, Hort. Brit. ed. II. 278 (Quid?).—Sibir.
tener, Salisb. Prod. 179 = seq.
tenerrimus, *Linn. Sp. Pl.* 794.—Reg. Mediterr.
tenerrimus, Schur, Enum. Pl. Transs. 371 = oleraceus.
tenuifolius, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 438 = tenerrimus.
tigridus, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xiv. (1856) 231.—Mesopot.
tingitanus, Lam. Encyc. iii. 397 = Picridium tingitanum.
tomentosus, Palau, ex Willk. & Lange, Prod. Fl. Hisp. ii. 249 = Crepis albida.
torulosus, Savi, Fl. Ital. i. t. 15 = gibbosus.
tuberosus, Murr. Syst. ed. XIII. 594; Linn. f. Suppl. 346 = Lactuca tuberosa.
uliginosus, Bieb. Fl. Taur. Cauc. ii. 238 = arvensis.
umbellatus, E. Mey. ex DC. Prod. vii. 185 = asper.
umbellifer, Thunb. Prod. Pl. Cap. 139 = oleraceus.
ustulatus, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 22.—Ins. Mader.
vaginatus, *Hort. Erf. ex Steud. Nom.* ed. I. 799.—Hab.?
virgatus, Desf. ex Steud. l. c. 655, 799 = Prenanthes virgata.
volhynicus, Bess. ex Nym. Consp. 433 = Lactuca tatarica.
Wallichianus, DC. Prod. vii. 185; et in Deless. Ic. Sel. iv. t. 97 = arvensis.
Webbii, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 444.—Ins. Canar.
Wightianus, DC. Prod. vii. 187 = arvensis.
zacinthoides, DC. l. c. 184 = oleraceus.

SONDARIA, Dennst. Schluess. Hort. Malab. 31 (1818). *RHAMNEAE*?
cranganoorensis, Dennst. l. c. (Quid?).—Ind. or.

SONDERA, Lehm. Pugill. viii. 44 (1844) = **Drosera**, Linn.
macrantha, Lehm. l. c. = D. macrantha.
Preissii, Lehm. l. c. 45 = D. Preissii.

SONERILA, Roxb. Hort. Beng. [5] (1814); Fl. Ind. i. 176 (1832). *MELASTOMACEAE*, Benth. & Hook. f. i. 753.
acaulis, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 46 = Wallichii.
affinis, Arn. in Hook. Comp. Bot. Mag. ii. (1836) 307 = zeylanica.
affinis, Thw. Enum. Pl. Zeyl. 109 = Brunonis.
amabilis, *Bedd. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 76, nomen.—Ind. or.
amabilis, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 53.—Reg. Himal.
angustata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 76.—Zeylan.
angustifolia, Roxb. Hort. Beng. [5]; Fl. Ind. i. 178 = maculata.
arguta, *R. Br. in Wall. Cat.* n. 4095.—Reg. Himal.
Arnottiana, Thw. Enum. Pl. Zeyl. 108 = Wightiana.
axillaris, Wight, Ic. t. 1058 = versicolor.
begoniaefolia, Blume, in Flora, xiv. (1831) 490 = moluccana.
Bensoni, *Hook. f. Bot. Mag.* t. 6049.—Ind. or.
biflora, *Zoll. & Mor. Syst. Verz. Zoll.* ii.—Java.

SONERILA:—
brachyandra, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 320 = tenera.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 53.—Burma.
Brunonis, *Wight & Arn. Prod.* 321.—Ind. or.
bullata, Griff. Notul. iv. 675 = Driessenia bullosa.
bullosa, Triana, in Trans. Linn. Soc. xxviii. (1871) 73, sphalm. = praec.
ceylanica, Arn. in Hook. Comp. Bot. Mag. ii. (1836) 307 = zeylanica.
elegans, Hook. Bot. Mag. t. 4978 = speciosa.
elegans, *Wight, Spicil. Neilgh.* i. 57. t. 67.—Ind. or.
emaculata, Roxb. Hort. Beng. [5]; Fl. Ind. i. 178 = maculata.
erecta, *Jack, in Malay. Misc.* i. (1820) n. v. 7.—Burma; Malaya.
firma, Thw. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 75, nomen = Gardneri.
Fordii, *Oliver, in Hook. Ic. Pl.* t. 1457.—China.
Gardneri, *Thw. Enum. Pl. Zeyl.* 107.—Zeylan.
glaberrima, Arn. in Hook. Comp. Bot. Mag. ii. (1836) 307 = zeylanica.
grandiflora, *R. Br. in Wall. Cat.* n. 4099.—Ind. or.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 539.—Malacca.
Guneratnei, Trimen, in Journ. Bot. xxiii. (1885) 208 = linearis.
Hartwegi, Triana, in Trans. Linn. Soc. xxviii. (1871) 76, sphalm. = seq.
Harveyi, *Thw. Enum. Pl. Zeyl.* 107.—Zeylan.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 540.—Burma.
Hendersoni, Hort.—Cf. Gard. Chron. (1874) ii. 83; et ex Fourn. Illustr. Hortic. xxiii. (1876) 11. t. 230 = margaritacea.
heterophylla, *Jack, in Malay. Misc.* ii. (1822) n. vii. 16.—Malaya.
heterostema, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 326 = obliqua.
hirsutula, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 307.—Zeylan.
Hookeriana, Arn. l. c. 308 = zeylanica.
insignis, *Blume, Mus. Bot. Lugd. Bat.* i. 11.—Sumatra.
Junghuhniana, *Miq. Fl. Ind. Bat.* i. i. 566.—Sumatra.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 539.—Reg. Himal.
Kurzii, C. B. Clarke, l. c. = amabilis.
laeviuscula, *Zoll. & Mor. ex Miq. Fl. Ind. Bat.* i. i. 564.—Malaya.
lanceolata, *Thw. Enum. Pl. Zeyl.* 107.—Zeylan.
linearis, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 76.—Burma.
maculata, *Roxb. Hort. Beng.* [5]; *Fl. Ind.* i. 177.—Reg. Himal.
magnifica, *Miq. Fl. Ind. Bat. Suppl.* 320.—Sumatra.
Mamei, Linden, in Illustr. Hortic. xxiii. (1876) 146. t. 254 = margaritacea.
margaritacea, *Lindl. in Gard. Chron.* (1854) 727.—Burma.
moluccana, *Roxb. Fl. Ind. ed. Carey*, i. 182.—Malaya.
Naudiniana, Miq. Fl. Ind. Bat. i. i. 565 = Sarcopyramis nepalensis.
nudiscapa, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 78, 77.—Burma.
obliqua, *Korth. Verh. Nat. Gesch. Bot.* 250.—Malaya.
orbiculata, Lindl. in Journ. Hort. Soc. viii. (1853) 56 = speciosa.
paradoxa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 321 = moluccana.
pauciflora, Blume, in Flora, xiv. (1831) 491 = heterophylla.
pedunculosa, *Thw. Enum. Pl. Zeyl.* 109.—Zeylan.
picta, Griff. Notul. iv. 676 = maculata.
picta, *Korth. Verh. Nat. Gesch. Bot.* 249. t. 52.—Burma; Sumatra.
pilosula, *Thw. Enum. Pl. Zeyl.* 108.—Zeylan.
pumila, Thw. l. c. 109 = zeylanica.
Rheedii, *Wall. Cat.* n. 4096; *Wight & Arn. Prod.* 321.—Ind. or.
rhombifolia, Thw. Enum. Pl. Zeyl. 108 = zeylanica.
robusta, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 308.—Zeylan.
rostrata, Thw. Enum. Pl. Zeyl. 108 = zeylanica.
Rottleri, Wall. Cat. n. 4097 = stricta, pedunculosa.
rotundifolia, *Bedd. in Madr. Journ. Sc.* Ser. II. xxii. (1861) 74; *et in Trans. Linn. Soc.* xxv. (1865) 216.—Ind. or.

SONERILA:—
scapigera, *Dalz. in Hook. Lond. Journ. Bot.* vii. (1848) 672.—Ind. or.
secunda, *R. Br. in Wall. Cat.* n. 4094.—Burma.
solanoides, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 324 = speciosa.
speciosa, *Zenker, in Ann. Sc. Nat.* Sér. II. vi. (1830) 151.—Ind. or.
squarrosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 182.—Reg. Himal.
stricta, *Hook. Bot. Mag.* t. 4394.—Burma.
tenella, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 46 = Arnottiana.
tenera, R. Br. in Wall. Cat. n. 4098, partim = stricta.
tenera, *Royle, Illustr. Bot. Himal.* 215. t. 45. f. 2.—Ind. or.; China; Ins. Philipp.
tenuifolia, *Blume, in Flora*, xiv. (1831) 491.—Malaya.
Teysmanniana, *Miq. Fl. Ind. Bat. Suppl.* 320.—Sumatra.
tomentella, *Thw. Enum. Pl. Zeyl.* 109.—Zeylan.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 33.—Ind. or.
versicolor, *Wight, Ic.* t. 1057.—Ind. or.
violaefolia, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 77.—Ind. or.
Wallichii, *Benn. Pl. Jav. Rar.* 215.—Ind. or.
Wightiana, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 307.—Ind. or.
zeylanica, *Wight & Arn. Prod.* 322 (err. typ. 332).—Zeylan.; Borneo.

SONINNIA, Kostel. Allg. Med.-Pharm. Fl. iii. 1091 (1834), sphalm. = SONNINIA, Reichb. = **Diplolepis**, R. Br. (Asclep.).

SONNERATIA, Linn. f. Suppl. 38 (1781). *LYTHRARIEAE*, Benth. & Hook. f. i. 784.
Aubletia, Gaertn. Fruct. i. 379. t. 78 (1788).
Blatti, Adans. Fam. ii. 88 (1763).
Chiratia, Montr. in Mém. Acad. Lyon, x. (1860) 202.
Kambala, Rafin. Sylva Tellur. 19 (1838).
Mycostylis, Rafin. l. c. (1838).
acida, Benth. Fl. Austral. iii. 301 = alba.
acida, *Linn. f. Suppl.* 252.—Ind. or.; Malaya.
alba, Griff. Notul. iv. 652 = Griffithii.
alba, *Sm. in Rees, Cycl.* xxxiii. n. 2.—Geront. trop.
apetala, [*Buch.-Ham.*] *in Syme, Embassy*, ed. II. iii. 313.—Ind. or.; Burma.
evenia, *Blume, Mus. Bot. Lugd. Bat.* i. 337.—Ins. Timor.
Griffithii, *Kurz, Pegu For. Rep. App.* B. 54; *For. Fl. Brit. Burma*, i. 527.—Burma.
lanceolata, *Blume, Mus. Bot. Lugd. Bat.* i. 337.—Borneo.
mossambicensis, Klotzsch in Peters, Reise Mossamb. Bot. 66. t. 12 = alba.
neglecta, Blume, Mus. Bot. Lugd. Bat. i. 338 = Griffithii.
obovata, *Blume, l. c.* 337.—Java.
ovalis, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 198.—Java.
Pagatpat, *Blanco, Fl. Filip.* ed. I. 424; *Blume, Mus. Bot. Lugd. Bat.* i. 336.—Malaya.
punctata, J. F. Gmel. Syst. 675 = Clausena Wampi.

SONNINIA, Reichb. Consp. 131 (1828) = **Diplolepis**, R. Br. (Asclep.).
Menziezii, Decne. in DC. Prod. viii. 589 = D. Menziesi.
vomitoria, Kostel. Allg. Med.-Pharm. Fl. iii. 1091 = D. Menziesi.

SONZAYA, Marchand, in Baill. Adansonia, viii. (1867–68) 64. *BURSERACEAE*.
australiana, *Marchand, l. c.* 71.—Austral.

SOPHANDRA, Meissn. Gen. Comm. 152; et in Ind. (1840), sphalm. = LOPHANDRA, D. Don = **Erica**, Tourn.

SOPHIA, Linn. Pl. Surin. n. 86 (1775); Amoen. Acad. viii. 259 = **Pachira**, Aubl. (Malvac.).
carolina, Linn. l. c. = P. aquatica.

SOPHIA, Linn. Syst. ed. I (1735); Adans. Fam. ii. 417 (1763) = **Sisymbrium**, Tourn. (Crucif.).
Lobelii, Rupr. Fl. Cauc. 88 = Sis. Sophia.
multifida, Gilib. Exercit. i. 243 = Sis. Sophia.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = Sis. Sophia.

SOPHOCLESIA, Klotzsch, in Linnaea, xxiv. (1851) 29. *VACCINIACEAE*, Benth. & Hook. f. ii. 576.
apophysata, Griseb. Fl. Brit. W. Ind. 143 = Findlaya apophysata.
cordifolia, *Klotzsch, in Linnaea*, xxiv. (1851) 29.—N. Granat.
majus, *Benth. & Hook. f. Gen.* ii. 576.—Ind. occ.; Peruv.
nummulariaefolia, *Klotzsch, in Linnaea*, xxiv. (1851) 30.—N. Granat.
ovata, *Klotzsch, l. c.*—N. Granat.
subscandens, *Klotzsch, l. c.*—N. Granat.

SOPHONODON, Miq. Fl. Ind. Bat. i. ii. 592 (1859), sphalm. = **Siphonodon**, Grif. (Celastr.).

SOPHORA, Linn. Gen. ed. I. 125 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 555.
Agastianis, Rafin. New Fl. Am. iii. 85 (1836).
Broussonetia, Orteg. Hort. Matr. Dec. 61. t. 7 (1798).
Dermatophyllum, Scheele, in Linnaea, xxi. (1848) 458.
Edwardsia, Salisb. in Trans. Linn. Soc. ix. (1808) 298. t. 26.
Goebelia, Bunge, ex Boiss. Fl. Orient. ii. 628 (1872).
Patrinia, Rafin. in Journ. Phys. lxxxix. (1819) 97.
Pseudosophora, Sweet, Hort. Brit. ed. II. 122 (1830).
Radiusa, Reichb. Consp. 148 (1828).
Styphnolobium, Schott, in Wien. Zeitschr. iii. (1830) 844; ex Endl. in Linnaea, vi. (1831) Litt. 54.
Vibexia, Rafin. Atl. Journ. 144 (1832).
Zanthyrsis, Rafin. New Fl. Am. iii. 84 (1836).
acuminata, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 250.—Reg. Himal.
acuminata, *Desv. Journ. Bot.* iii. (1814) 75.—'Am.'
affinis, *Torr. & Gray, Fl. N. Am.* i. 390.—Am. bor. occ.
alba, Linn. Syst. ed. XII. 287 = Baptisia alba.
albicans, [Jaume St. Hil.] in Duham. Arb. ed. Nov. iii. 86 = alopecuroides.
alopecuroides, *Linn. Sp. Pl.* 373.—As. Min.; Reg. Himal.
alpina, Pall. Astragal. 121. t. 90. f. 1 = Thermopsis corgonensis.
angustifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1843) 118 = flavescens.
arenicola, Nees, in Flora, iv. (1821) 297 = tomentosa.
argentea, Pall. in Nov. Act. Petrop. x. (1797) 373. t. 8 = Ammodendron Sieversii.
argentea, Salisb. Prod. 328 = Podalyria myrtillifolia.
arizonica, S. Wats. in Proc. Am. Acad. xi. (1876) 135.—Am. bor. occ.
aurea, Ait. Hort. Kew. ed. I. ii. 44 = Calpurnia lasiogyne.
australis, Linn. Syst. ed. XII. 287 = Baptisia australis.
biflora, Houtt. Handleid. v. 8. t. 24. f. 1 = Podalyria calyptrata.
biflora, Linn. Syst. ed. X. 1015 = Podalyria myrtillifolia.
biflora, Retz. Obs. i. 36 = Podalyria argentea.
bifolia, Pall. Astragal. 124. t. 91 = Ammodendron Sieversii.
buxifolia, Retz. Obs. i. 35 = Podalyria buxifolia.
calyptrata, Retz. l. c. 36 = Podalyria calyptrata.
capensis, Linn. Mant. i. 67 = Virgilia capensis.
chinensis, *G. Don, in Loud. Hort. Brit.* 162, nomen.—China.
chrysophylla, *Seem. Fl. Vit.* 66.—Polynes.
coerulea, Moench, Meth. 69 = Baptisia australis.
cordata, Thunb. Prod. Pl. Cap. 79 = Podalyria Burchelli, cordata.
crassifolia, *Hassk. Cat. Hort. Bog. Alt.* 285.—Java.
crassifolia, [Jaume St. Hil.] in Duham. Arb. ed. Nov. iii. 87 = tomentosa.
cuneifolia, Steud. Nom. ed. I. 799 = Podalyria cuneifolia.
denudata, *Bory, Voy.* ii. 399.—Ins. Borbon.

SOPHORA:—
fabacea, Pall. Astragal. 122. t. 90. f. 2 = Thermopsis fabacea.
flavescens, *Ait. Hort. Kew.* ed. I. ii. 43.—Sibir.
Fraseri, *Benth. Fl. Austral.* ii. 274.—Austral.
galegoides, Pall. Astragal. 118. t. 88 = flavescens.
galioides, Berg. Desc. Pl. Cap. 141 = Cyclopia galioides, genistoides.
genistaefolia, Salisb. Prod. 328 = seq.
genistoides, Linn. Sp. Pl. 373; Burm. f. Prod. Fl. Cap. 12 = Cyclopia genistoides.
glabra, Hassk. in Flora, xxv. (1842) ii. Beibl. 47 = tomentosa.
glabra, Moench, Meth. Suppl. 22 = flavescens.
glauca, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 98.—Ind. or.
glauca, Salisb. Prod. 328 = Baptisia alba.
Griffithii, Stocks, in Hook. Kew Journ. Bot. iv. (1852) 147 = Keyserlingia Griffithii.
havanensis, Jacq. Enum. Pl. Carib. 20 = tomentosa.
heptaphylla, *Linn. Sp. Pl.* 373.—Zeylan; China.
heptaphylla, Wight, Ic. t. 1155 = Wightii.
hirsuta, Ait. Hort. Kew. ed. I. ii. 46 = Podalyria cordata.
Houghiana, Wall. Cat. n. 5336 B = mollis.
inhambanensis, Kotzsch, in Peters, Reise Mossamb. Bot. 26 = tomentosa.
intermedia, Hort. ex Steud. Nom. ed. II. ii. 612 (Quid?).—Hab. ?
interrupta, *Bedd. Forest. Man. Bot.* 90.—Ind. or.
Jabandas, *Montr. in Mém. Acad. Lyon*, x. (1860) 196.—N. Caled.
japonica, *Linn. Mant.* i. 68.—Japon; China.
Jaubertii, Spach, ex Jaub. et Spach, Illustr. iv. 45 = alopecuroides.
juncea, Schrad. Sert. Hannov. 9. t. 3 = Viminaria denudata.
kentukea, Dum.-Cours. Bot. Cult. ed. II. vi. 56 (Quid?).—Am. bor.
Kronei, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 219 = flavescens.
lanceolata, Walt. Fl. Carol. 135 = Baptisia lanceolata.
linearifolia, *Griseb. in Gott. Abh.* xxiv. (1879) 110.—Reg. Argent.
littoralis, Schrad. in Goett. Gel. Anz. i. (1821) 709 = tomentosa.
lupinoides, Linn. Sp. Pl. 374 = Thermopsis lanceolata.
macrocarpa, Sm. in Rees, Cycl. xxxiii. n. 6 = tetraptera.
macrosperma, DC. Prod. ii. 96 = seq.
mecosperma, [Jaume St. Hil.] in Duham. Arb. ed. Nov. iii. 86, 87 = flavescens.
microphylla, Ait. Hort. Kew. ed. I. ii. 43 = tetraptera.
microphylla, Meyen, Reise, i. 407 = Zuccagnia microphylla.
mollis, *R. Grah. in Wall. Cat.* n. 5335.—Reg. Himal.; China.
mollis, Span. in Linnaea, xv. (1841) 189 = glauca.
monosperma, Sw. Prod. Veg. Ind. Occ. 66 = Ormosia dasycarpa.
Moorcroftiana, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 249.—Reg. Himal.; China.
mutabilis, Salisb. Prod. 328 = Podalyria cordata.
myrtillifolia, Retz. Obs. i. 36 = Podalyria myrtillifolia.
nitens, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 266.—Afr. austr.
nitens, Schum. & Thonn. Beskr. Guin. Pl. 201 = tomentosa.
nitida, *Sm. in Rees, Cycl.* xxxiii. n. 7.—Ins. Borbon.
obliqua, Pers. Syn. i. 452 = Piptadenia moniliformis.
occidentalis, Linn. Syst. ed. X. 1015 = tomentosa.
oligophylla, *Baker, in Oliver, Fl. Trop. Afr.* ii. 254.—Afr. trop.
orientalis, Pall. Reise, ii. 505 = alopecuroides.
oroboides, Berg. Desc. Pl. Cap. 142 = Virgilia capensis.
pachycarpa, *Schrenk, ex C. A. Mey. Ind. Sem. Hort. Petrop.* ix. 89.—Soongar.
pallida, Salisb. Prod. 328 = alopecuroides.
pedunculata, Thunb. Prod. Pl. Cap. 79 = Podalyria argentea.
pentaphylla, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 407.—Peruv.
perfoliata, Walt. Fl. Carol. 135 = Baptisia perfoliata.
platycarpa, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 398.—Japon.

SOPHORA:—
prostrata, *F. Buch. in Trans. N. Z. Inst.* xvi. (1884) 395.—N. Zel.
pubescens, *Tausch, in Flora,* xvii. (1834) 489.—Hab. ?
retusa, Pers. Syn. i. 452 = denudata.
Retzii, J. F. Gmel. Syst. 665 = Podalyria argentea.
rhynchocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 110.—Reg. Argent.
robinoides, *Walp. in Otto & Dietr. Allg. Gartenz.* vii. (1839) 337.—Ind. or.
robusta, Roxb. Hort. Beng. [31], nomen = Ormosia robusta.
rotundifolia, Berg. Desc. Pl. Cap. 138; Thunb. Prod. Pl. Cap. 79 = Podalyria myrtillifolia.
secundiflora, *Lag. ex DC. Cat. Hort. Monsp.* (1813) 148.—Mexic.
sempervirens, Engelm. in Bost. Journ. Nat. Hist vi. (1850) 178 = speciosa.
senegalensis, Deless. ex DC. Prod. ii. 168 = Tephrosia obcordata.
sericea, Andr. Bot. Rep. t. 440 = Podalyria sericea.
sericea, [Jaume St. Hil.] in Duham. Arb. ed. Nov. iii. 86 = nitida.
sericea, *Nutt. Gen. Am.* i. 280.—Am. bor.
sinica, Rosier, in Journ. Phys. xiv. (1779) 248 = japonica.
songarica, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 307.—Soongar.
sororia, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 210 = flavescens.
speciosa, Benth. in Bost. Journ. Nat. Hist. vi. (1850) 178 = secundiflora.
speciosa, Torr. in Pacif. Rail. Rep. iv. 82 = arizonica.
stenophylla, *A. Gray, in Ives, Rep. Colorado River, Bot.* 10.—Mexic.
sylvatica, Burch. Trav. ii. 146; DC. Prod. ii. 98 = Calpurnia sylvatica.
ternata, Thunb. Prod. Pl. Cap. 79 = Aspalathus ferruginea.
tetraptera, *F. Mill. Ic. Pl.* t. 1.—N. Zel.; Chili.
tiloënsis, Blume, ex Miq. Fl. Ind. Bat. i. 1. 125 = heptaphylla.
tinctoria, Linn. Sp. Pl. 373 = Baptisia tinctoria.
tomentosa, *Linn. l. c.*—Reg. calid.
trifolia, Steud. Nom. ed. II. ii. 612, sphalm. = seq.
trifoliata, *Thunb. Prod. Pl. Cap.* 79 = Argyrolobium sericeum.
triphylla, Sweet, Hort. Brit. ed. I. 107, in syn., sphalm. = praec.
velutina, Lindl. Bot. Reg. t. 1185 = glauca.
viciaefolia, Salisb. Prod. 328 = Virgilia capensis.
viciifolia, *Hance, in Journ. Bot.* xix. (1881) 209.—China.
villosa, Walt. Fl. Carol. 134 = Baptisia villosa.
violacea, *Thw. Enum. Pl. Zeyl.* 94.—Zeylan.
violacea, Zipp. ex Span. in Linnaea, xv. (1841) 192 = Aeschynomene atropurpurea.
Wightii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 250.—Ind. or.
zambesiaca, *Baker, in Oliver, Fl. Trop. Afr.* ii. 253.—Afr. trop.

SOPHOROCAPNOS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 570 = **Corydalis**, Vent. (Papav.).
pallida, Turcz. l. c. 571 = C. pallida.

SOPHRONANTHE, Benth. in Lindl. Introd. Nat. Syst. ed. II. 445 (1836) = **Gratiola**, Rupp. (Scrophul.).
floridana, Walp. Rep. iii. 298 = G. floridana.
hispida, Benth. in Lindl. Introd. Nat. Syst. ed. II. 445 = G. subulata.

SOPHRONIA, Licht. ex Roem. & Schult. Syst. i. 482 (1817) = **Lapeyrousia**, Pourr. (Irid.).
caespitosa, Licht. l. c.—Afr. austr.
caespitosa, Spreng. Syst. iii. 147, in syn. = Klattia partita.

SOPHRONIA, Lindl. Bot. Reg. t. 1129 (1828) = **Sophronitis**, Lindl. (Orchid.).
cernua, Lindl. l. c. = Sophronitis cernua.
modesta, Lindl. l. c. sub t. 1147 = Sophronitis cernua.

SOPHRONITIS, Lindl. Bot. Reg. sub t. 1147 (1828).
ORCHIDEAE, Benth. & Hook. f. iii. 535.
LOPHOGLOTIS, Rafin. Fl. Tellur. iv. 49 (1836).
SOPHRONIA, Lindl. Bot. Reg. t. 1129 (1828).

SOPHRONITIS:—
cernua, *Lindl. Bot. Reg.* sub t. 1147.—Cf. t. 1129.—Bras.
coccinea, Reichb. f. in Walp. Ann. vi. 465 = grandiflora.
grandiflora, *Lindl. Sert. Orch.* (1838) t. 5. f. 2.—Bras.
Hoffmannseggii, Reichb. ex Hoffmgg. in Linnaea, xvi. (1842) Litt. 286 = cernua.
isopetala, Hoffmgg. Verz. Orch.—Cf. Bot. Zeit. i. (1843) 834 = cernua.
militaris, Reichb. f. in Walp. Ann. vi. 465 = grandiflora.
modesta, Lindl. Bot. Reg. sub t. 1147 = cernua.
nutans, Hoffmgg. in Linnaea, xvi. (1842) Litt. 286 = cernua.
pterocarpa, *Lindl. & Paxt. Flow. Gard.* iii. (1853) 11.—Bras.
purpurea, Reichb. f. in Gard. Chron. (1878) i. 462 = grandiflora.
rosea, Hort. ex Gostling, in Gard. Chron. (1884) i. 243 = grandiflora.
Rossiteriana, *Rodrig. Orch. Nov.* i. 77.—Bras.
violacea, *Lindl. Bot. Reg.* (1840) *Misc.* 18.—Bras.
Wittigiana, *Rodrig. in Rev. Hortic.* (1878) 24; *ex ej. Orch. Nov.* ii. 159.—Bras.

SOPUBIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 88 (1825). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 970.
GERDARIA, Presl, Bot. Bemerk. 91 (1844).
LOPHANTHERA, Rafin. New Fl. Am. ii. 58 (1836).
RAPHIDOPHYLLUM, Hochst. in Flora, xxiv. (1841) 666.
RHAPHIDOPHYLLUM, Benth. in DC. Prod. x. 521 (1846).
cana, *Harv. Thes. Cap.* ii. 29.—Afr. austr.
delphinifolia, *G. Don, Gen. Syst.* iv. 560.—Ind. or.
Dregeana, *Benth. in DC. Prod.* x. 522.—Afr. austr.
filiformis, *G. Don, Gen. Syst.* iv. 560.—Afr. trop.
Hildebrandtii, *Vatke, in Linnaea,* xliii. (1880-82) 314.—Afr. trop.
karaguensis, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 123.—Afr. trop.
kituiensis, *Vatke, in Linnaea,* xliii. (1880-82) 313.—Afr. trop.
obtusifolia, *G. Don, Gen. Syst.* iv. 560.—Madag.
ramosa, *Hochst. in Flora,* xxvii. (1844) 27.—Afr. trop.
scabra, *G. Don, Gen. Syst.* iv. 560.—Afr. austr.
scopiformis, *Vatke, in Linnaea,* xliii. (1880-82) 313.—Afr. trop.
simplex, *Hochst. in Flora,* xxvii. (1844) 27.—Afr. austr.
stricta, *Baker, in Journ. Linn. Soc.* xxi. (1885) 427.—Madag.
stricta, *G. Don, Gen. Syst.* iv. 559.—Ind. or.; Malaya.
trifida, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 88.—Ind. or.; Nepal.
triphylla, *Baker, in Journ. Bot.* xx. (1882) 220.—Madag.

SORAMIA, Aubl. Pl. Gui. i. 552. t. 219 (1775) = **Dolicarpus**, Roland. (Dillen.).
guianensis, Aubl. l. c. = D. Soramia.

SORANTHE, Salisb. in Knight, Prot. 71 (1809) = **Sorocephalus**, R. Br. (1810) (Proteac.).
ciliciiflora, Knight, Prot. 72 = Soroc. lanatus.
clavigera, Knight, l. c. 73.—Afr. austr.
glanduligera, Knight, l. c. 71 = Soroc. imbricatus.
montana, Knight, l. c. 73.—Afr. austr.
pinifolia, Knight, l. c. 72.—Afr. austr.
rupestris, Knight, l. c.—Afr. austr.
tenuifolia, Knight, l. c. = Soroc. tenuifolius.

SORANTHUS, Ledeb. Ic. Pl. i. 20. t. 82 (1829); Fl. Alt. i. 344 (1829) = **Ferula**, Tourn. (Umbell.).
Meyeri, Ledeb. ll. cc. = F. Meyeri.

SORBARIA, A. Br. in Aschers. Fl. Brandenb. 177 (1864) = **Spiraea**, Linn. (Rosac.).
grandiflora, Maxim. in Act. Hort. Petrop. vi. (1879) 223 = Sp. grandiflora.
Kirilowi, Maxim. l. c. 225 = Sp. sorbifolia.
Lindleyana, Maxim. l. c. 224 = Sp. sorbifolia.
sorbifolia, A. Br. in Aschers. Fl. Brandenb. 177 = Sp. sorbifolia.

SORBUS, [Tourn.] Linn. Syst. ed. I. (1735) = **Pyrus**, Tourn. (Rosac.).
alandica, Bergstr. Naturalh. Anteckn. om Åland, 19.—Suecia.
alnifolia, C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 249 = Crataegus alnifolia.
alpina, Heynh. Nom. ii. 684 = P. Chamaemespilus ?
ambigua, Michalet, Bot. Jura, 156 = P. Chamaemespilus.
Amelanchier, Crantz, Stirp. Austr. ed. I. fasc. ii. 53; ed. II. 90 = Amelanchier vulgaris.
americana, Marsh. Arbust. 145; Willd. Enum. Hort. Berol. 520 = P. americana.
Arbuscula, Bosc, ex Poir. Encyc. Suppl. v. 164 = P. Aucuparia.
arbutifolia, Heynh. Nom. i. 772 = P. arbutifolia.
Aria, Crantz, Stirp. Austr. ed. I. fasc. ii. 46; ed. II. 86* = P. Aria.
Aria, Wimm. ex Nym. Consp. 242 = P. Chamaemespilus.
arioides, Michalet, Bot. Jura, 156 = P. intermedia.
atrovirens, Hort. ex C. Koch, Dendrol. i. 194 = P. rotundifolia.
Aucuparia, Linn. Sp. Pl. 477 = P. Aucuparia.
Aucuparia, Poir. Encyc. vii. 234, partim = P. americana.
Aucuparia, Stell. in Pall. Fl. Nord. Beitr. ii. 300 = P. sambucifolia.
auriculata, Pers. Syn. ii. 38 = P. pinnatifida.
betulaefolia, Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. betulaefolia.
Chamaemespilus, Crantz, Stirp. Austr. ed. I. fasc. ii. 40; ed. II. 83* = P. Chamaemespilus.
corymbosa, C. Koch, Hort. Dendrol. 176 = P. communis.
crataegifolia, Wenzig, in Eichl. Jahrb. Berl. ii. (1883) 295 = P. crataegifolia.
crenata, Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. vestita ?
Cydonia, Crantz, Stirp. Austr. ed. I. fasc. ii. 57; ed. II. 93 = P. Cydonia.
densiflora, Heynh. Nom. ii. 684 = P. densiflora.
depressa, Heynh. l. c. i. 772 = P. arbutifolia.
domestica, Linn. Sp. Pl. 477 = P. Sorbus.
edulis, C. Koch, ex Kirch. in C. Koch & Fint. Wochenschr. ii. (1859) 382 = P. Aria.
fennica, Kalm, ex Fries, Summa Veg. Scand. 42 = P. pinnatifida.
fennica, C. Koch, Dendrol. i. 194 = P. intermedia.
flabellifolia, Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. Aria.
florentina, Nym. Syll. 266 = P. crataegifolia.
floribunda, Heynh. Nom. i. 773 = P. arbutifolia.
foliolosa, Spach, Hist. Vég. Phan. ii. 96 = P. foliolosa.
foliosa, Wenzig, in Linnaea, xxxviii. (1874) 75; Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 159, sphalm. = P. foliolosa.
fruticosa, Steud. Nom. ed. I. 800 = P. Chamaemespilus.
glabra, Gilib. Fl. Lituan. ii. 233 = P. glabra.
glabrata, Hort. Vrat. ex Kirch. in C. Koch & Fint. Wochenschr. ii. (1859) 382 = P. Aria.
gracilis, Wenzig, in Linnaea, xxxviii. (1874) 74; Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 159 = P. sambucifolia.
graeca, Lodd. ex Steud. Nom. ed. II. ii. 612 = P. Aria.
grandifolia, Heynh. Nom. i. 773 = P. arbutifolia.
heterophylla, Reichb. Fl. Germ. Excurs. 628 = P. spuria.
Hostii, Heynh. Nom. ii. 685 = P. Aria.
humifusa, S. Wats. Bibliog. Ind. N. Am. Bot. 290 = P. americana.
hybrida, Linn. Sp. Pl. ed. II. 684 = P. pinnatifida.
intermedia, Pers. Syn. Pl. ii. 38 = P. intermedia.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 67 = P. japonica.
kamunensis, Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. lanata.
lanata, Schau. l. c. = P. lanata.
lanuginosa, Kit. in Schult. Oester. Fl. ed. II. ii. 50 = P. Aucuparia.
latifolia, Pers. Syn. ii. 38 = P. rotundifolia.
Malus, Crantz, Stirp. Austr. ed. I. fasc. ii. 57; ed. II. 93 = P. Malus.
melanocarpa, Heynh. Nom. i. 773 = P. arbutifolia.
micrantha, Hort. Angl. ex Dum.-Cours. Bot. Cult. v. 464 = P. americana.

SORBUS:—
microcarpa, Pursh, Fl. Am. Sept. i. 341 = Pyrus americana.
microphylla, Wenzig, in Linnaea, xxxviii. (1874) 76; Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 159 = P. microphylla.
Mougeoti, Godr. & Soyer-Willem. in Bull. Soc. Bot. Fr. v. (1858) 447 = P. intermedia.
nivea, Steud. Nom. ed. II. ii. 612 = P. Aria.
oblongifolia, Reichb. ex Nym. Consp. 242 = P. Aria.
obtusata, Heynh. Nom. ii. 685 = P. obtusata.
pubens, Heynh. l. c. i. 773 = P. arbutifolia.
praemorsa, Nym. Syll. 265 = P. Aucuparia.
pumilus, Rafin. Med. Fl. ii. 265 = P. americana.
purpurea, Dulac, Fl. Hautes-Pyr. 318 = P. Chamaemespilus.
Pyrus, Crantz, Stirp. Austr. ed. I. fasc. ii. 56; ed. II. 93 = P. communis.
riparia, Rafin. New Fl. Am. iii. 15 = P. americana.
rivularis, Heynh. Nom. i. 773 = P. rivularis.
rupicola, Nym. Consp. 242 = P. Aria.
sabauda, Nym. l. c. = P. Chamaemespilus.
sambucifolia, M. Roem. Syn. Rosifl. 139 = P. sambucifolia.
scandica, Fries, Fl. Halland. 83 = P. intermedia.
sikkimensis, Wenzig, in Linnaea, xxxviii. (1874) 58 = P. sikkimensis.
sitchensis, M. Roem. Syn. Rosifl. 139 = P. sambucifolia.
speciosa, Hort. ex C. Koch, Dendrol. i. 190 = P. Aucuparia.
spuria, Pers. Syn. ii. 38 = P. spuria.
subfusca, Boiss. Fl. Orient. ii. 658 = P. subfusca.
subserrata, Opiz, in Flora, vii. (1824) i. Beil. 83 = P. Aucuparia.
syrmiensis, Kit. in Linnaea, xxxii. (1863) 585 = P. syrmiensis.
tianschanica, Rupr. Sert. Tiansch. 46 = P. tianschanica.
Tommasinii, Fleischm. Fl. Krain, 206 = P. torminalis.
Toringo, De Vriese, in Tuinbouw-Fl. iii. (1856) t. ad p. 368 = P. Sieboldi.
torminalis, Crantz, Stirp. Austr. ed. I. fasc. ii. 45; ed. II. 85* = P. torminalis.
trilobata, Heynh. Nom. i. 773 = P. trilobata.
ursina, Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. foliosa.
variolosa, Schau. l. c. = P. Pashia.
vestita, Lodd. Cat. (1836); Schau. in Otto & Dietr. Allg. Gartenz. xvii. (1849) 84 = P. vestita.

SOREMA, Lindl. Bot. Reg. (1844) sub t. 46 = **Nolana**, Linn. (Convolv.).
acuminata, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 370 = N. acuminata.
atriplicifolia, Lindl. Bot. Reg. (1844) sub t. 46 = N. atriplicifolia.
bracteosa, Phil. Fl. Atac. 43.—Chili.
cordata, Remy, in Ann. Sc. Nat. Sér. III. vi. (1846) 351 = N. cordata.
elegans, Phil. Fl. Atac. 43.—Chili.
glutinosa, Phil. l. c. 44.—Chili.
lanceolata, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 498 = N. lanceolata.
linearis, Miers, l. c. 499 = N. linearis.
littoralis, Miers, l. c. 370 = N. littoralis.
longifolia, Miers, l. c. 498 = N. longifolia.
paradoxa, Lindl. Bot. Reg. (1844) sub t. 46 = N. paradoxa.
parviflora, Phil. Fl. Atac. 44.—Chili.
petiolata, Phil. in Linnaea, xxxiii. (1864–65) 207.—Chili.

SORGHUM, Linn. Syst. ed. I (1735) (Sorgum); Moench, Meth. 207 (1794). *GRAMINEAE*, Benth. & Hook. f. iii. 1135.
BLUMENBACHIA, Koel. Gram. Gall. et Germ. 28 (1802).
album, Hort. ex Roem. & Schult. Syst. ii. 836 = vulgare.
anomalum, *Desv. Opusc.* 72.—Am. calid.?
Arduini, Jacq. Eclog. Gram. 25. t. 18 = vulgare?
asperum, Beauv. Agrost. 131 = Danthonia papillosa.
arundinaceum, Roem. & Schult. Syst. ii. 813 = halepense.
avenaceum, Beauv. Agrost. 131 = Danthonia curvifolia.
avenaceum, Chapm. Fl. S. U. St. 583 = Chrysopogon avenaceus.
Balansae, *Hack. in Mart. Fl. Bras.* ii. III. 277, in obs.—Paraguay.
barbatum, Steud. Syn. Pl. Gram. 238 = Tristachya barbata.

SORGHUM:—
bicolor, Moench, Meth. 207 = vulgare.
caffrorum, Beauv. Agrost. 131 = vulgare.
campanum, Tenore & Guss. in Atti Acc. Sc. Nap. v. (1843) 303 = vulgare.
canescens, *Hack. in Mart. Fl. Bras.* ii. III. 277.—Bras.
capillare, Beauv. Agrost. 131; Roem. & Schult. Syst. ii. 840 = Andropogon capillaris.
caucasicum, Griseb. in Ledeb. Fl. Ross. iv. 476 = Andropogon intermedius.
cernuum, Willd. Enum. Hort. Berol. 1036 = vulgare.
commune, Beauv. Agrost. 131 = vulgare.
Crupina, Link, Hort. Berol. i. 246 = halepense.
cubanicus, Schult. Mant. ii. 519 (Index) = Holcus cubanicus.
decolor, Beauv. Agrost. 131 = halepense.
decolorans, Roem. & Schult. Syst. ii. 838 = halepense.
Dora, *Griseb. Fl. Brit. W. Ind.* 560.—Ind. occ.
Drummondii, Nees, ex Steud. Syn. Pl. Gram. 393 = vulgare.
dubium, C. Koch, in Linnaea, xxi. (1848) 443 = halepense.
elongatum, Beauv. Agrost. 131 = Chrysopogon elongatus.
flavum, Beauv. l. c. = seq.
fulvum, Beauv. l. c. 164 = Andropogon serratus.
fuscum, Miq. Fl. Ind. Bat. iii. 503 = Andropogon serratus.
giganteum, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 181 = halepense.
glycychylum, Passer. in Hamb. Gartenz. xiii. (1857) 33 = halepense.
halepense, *Pers. Syn.* i. 101.—Reg. calid.
intrans, F. Muell. ex Benth. Fl. Austral. vii. 541 = Andropogon intrans.
Junghuhnii, Miq. Fl. Ind. Bat. iii. 753 = Andropogon serratus.
Minarum, Hack. in Mart. Fl. Bras. ii. III. 276 = Chrysopogon stipoides.
muticum, Wight, ex Steud. Nom. ed. II. ii. 612 = Andropogon serratus.
nervosum, Bess. ex Schult. Mant. ii. Add. II. 669 = vulgare.
nigricans, Hort. ex Roem. & Schult. Syst. ii. 837 = vulgare.
nigrum, Roem. & Schult. l. c. = vulgare.
nitidum, Pers. Syn. i. 101 = Andropogon serratus.
nutans, A. Gray, Man. Bot. N. U. St. ed. II. 584 = Chrysopogon nutans.
pallidum, Beauv. Agrost. 131 = Chrysopogon pallidus.
parviflorum, Beauv. l. c. 132 = Chrysopogon violascens.
parviflorum, Desv. in Ham. Prod. Pl. Ind. Occ. 12 = Chrysopogon nutans.
pauciflorum, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 20.—Florida.
plumosum, Beauv. Agrost. 132 = Andropogon australis.
purpureo-sericeum, Schweinf. & Aschers. Beitr. Fl. Aethiop. 310 = Andropogon purpureo-sericeus.
pyramidale, Hort. ex Roem. & Schult. Syst. ii. 836 = vulgare.
Quartinianum, Schweinf. & Aschers. Beitr. Fl. Aethiop. 310 = Andropogon Quartinianus.
rubens, Willd. Enum. Hort. Berol. 1036 = vulgare.
saccharatum, Host, Gram. Austr. iv. t. 48 = Oplismenus undulatifolius.
saccharatum, Moench, Meth. 207 = vulgare.
Schreberi, Tenore, Fl. Neap. Syll. App. IV. p. iv. = halepense.
secundum, *Chapm. Fl. S. U. St.* 583.—Florida.
serratum, Roem. & Schult. Syst. ii. 839 = Panicum serratum.
setifolium, Roem. & Schult. l. c. 840 = Andropogon setifolius.
striatum, *Beauv. Agrost.* 132.—Hab.?
subglabrescens, Schweinf. & Aschers. Beitr. Fl. Aethiop. 310 = vulgare.
Thonizzi, *Hort. Berol. ex Pasq. Cat. Orto Bot. Nap.* 98, nomen.—Hab.?
timorense, Buese, ex De Vriese, Pl. Ind. Bat. Reinw. 103 = Andropogon australis.
tropicum, Buese, in Miq. Pl. Jungh. 359 = Andropogon serratus.
truchmenorum, C. Koch, in Linnaea, xxi. (1848) 442 = vulgare?
usorum, Nees, Fl. Afr. Austr. 87 = vulgare.
versicolor, *Anderss. in Peters, Reise Mossamb. Bot.* 563.—Afr. trop.
vulgare, *Pers. Syn.* i. 101.—Reg. trop. et sub-trop.

SORGUM, Linn. Syst. ed. I (1735) = praec.

SORIA, Adans. Fam. ii. 421 (1763) = **Euclidium**, R. Br. (Crucif.).
syriaca, Desv. ex Steud. Nom. ed. I. 126, 800 = E. syriacum.

SORIDIUM, Miers, in Proc. Linn. Soc. ii. (1850) 74; et in Trans. Linn. Soc. xxi. (1852) 49. t. 6 = **Sciaphila**, Blume (Triurid.).
Spruceanum, Miers, ll. cc. 75; 50.—Bras.

SORINDEIA, Thou. Gen. Nov. Madag. 23 (1806). *ANACARDIACEAE*, Benth. & Hook. f. i. 419.
DUPUISIA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 148. t. 38 (1832).
africana, *C. Sm. in Tuckey, Congo*, 12.—Afr. trop.
aurantiodora, Engl. in Mart. Fl. Bras. xii. II. 391 = Mauria aurantiodora.
Biringo, March. Rev. Anacard. 167 = Mauria Biringo.
calabarensis, March. l. c. = Trichoscypha Mannii.
elongata, *Blume, Mus. Bot. Lugd. Bat.* i. 205.—Madag.
falcata, March. Rev. Anacard. 167 = Euroschinus falcatus.
ferruginea, March. l. c. = Mauria ferruginea.
glaberrima, Engl. in Mart. Fl. Bras. xii. II. 391 = Mauria Biringo.
glaberrima, Hassk. Cat. Hort. Bog. Alt. 245 = madagascariensis.
heterandra, March. Rev. Anacard. 165 = madagascariensis.
heterophylla, *Hook. f. in Hook. Niger Fl.* 286. - Afr. trop.
heterophylla, March. Rev. Anacard. 167 = Mauria heterophylla.
juglandifolia, *Planch. ex Oliver, Fl. Trop. Afr.* i. 440.—Afr. trop.
longifolia, Oliver, l. c. 442 = Trichoscypha longifolia.
lucens, March. Rev. Anacard. 167 = Trichoscypha lucens.
macrophylla, *Planch. ex Oliver, Fl. Trop. Afr.* i. 442.—Afr. trop.
madagascariensis, *DC. Prod.* ii. 80.—Madag.
Mannii, Oliver, Fl. Trop. Afr. i. 441 = Trichoscypha acuminata.
ovalifolia, March. Rev. Anacard. 167 = Mauria ovalifolia.
patens, Oliver, Fl. Trop. Afr. i. 441 = Trichoscypha patens.
pinnata, Desf. Cat. Hort. Par. ed. III. 331 = madagascariensis.
puberula, March. Rev. Anacard. 167 = Mauria heterophylla.
simplicifolia, March. l. c. = Mauria simplicifolia.
suaveolens, March. l. c. = Mauria suaveolens.
trimera, *Oliver, Fl. Trop. Afr.* i. 441.—Afr. trop.
venulosa, Engl. in Mart. Fl. Bras. xii. II. 391 = Mauria suaveolens.

SOROCEA, A. St. Hil. in Mém. Mus. Par. vii. (1821) 473. *URTICACEAE*, Benth. & Hook. f. iii. 378.
PSEUDOSOROCEA, Baill. Adansonia, xi. (1875) 296.
affinis, *Hemsl. Biol. Centr. Am. Bot.* iii. 150.—Panama.
amazonica, *Miq. in Mart. Fl. Bras.* iv. I. 111.—Am. trop.
Celsiana, *Heynh. Nom.* ii. 685, nomen.—Bras.
Guilleminiana, *Gaudich. Bot. Voy. Bonite*, t. 74.—Am. trop.
Hilariana, Bur. in DC. Prod. xvii. 254 = seq.
Hilarii, *Gaudich. Bot. Voy. Bonite*, t. 71.—Bras.
Houllettiana, *Gaudich. l. c.* t. 74.—Am. trop.
ilicifolia, *Miq. in Mart. Fl. Bras.* iv. I. 114.—Am. trop.
Klotzschiana, *Baill. Adansonia*, i. (1860-61) 212.—Bras.
macrophylla, *Gaudich. Bot. Voy. Bonite*, t. 73.—Am. trop.
mexicana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 335 = Trophis mexicana.
muriculata, *Miq. in Mart. Fl. Bras.* iv. I. 113.—Am. trop.
pubivena, *Hemsl. Biol. Centr. Am. Bot.* iii. 150.—Guatemala.
racemosa, *Gaudich. Bot. Voy. Bonite*, t. 72.—Am. trop.
Uriamem, *Mart. ex Miq. in Mart. Fl. Bras.* iv. I. 113.—Am. trop.

SOROCEPHALUS, R. Br. in Trans. Linn. Soc. x. (1810) 139. *PROTEACEAE*, Benth. & Hook. f. iii. 171.
SORANTHE, Salisb. in Knight, Prot. 71 (1809), nomen prius.

SOROCEPHALUS:—
diversifolius, *R. Br. in Trans. Linn. Soc.* x. (1810) 143.—Afr. austr.
diversifolius, Drège, ex Meissn. in DC. Prod. xiv. 301 = Nivenia Dregei.
Dregei, *Buek, in Drège, Zwei Pfl. Docum.* 117, 222; *et ex Meissn. in DC. Prod.* xiv. 305.—Afr. austr.
imberbis, *R. Br. in Trans. Linn. Soc.* x. (1810) 140.—Afr. austr.
imberbis, Klotzsch, ex Meissn. in DC. Prod. xiv. 304 = scabridus.
imbricatus, *R. Br. in Trans. Linn. Soc.* x. (1810) 142.—Afr. austr.
intermedius, Buek, ex Meissn. in DC. Prod. xiv. 305 = Dregei.
lanatus, *R. Br. in Trans. Linn. Soc.* x. (1810) 142.—Afr. austr.
nivalis, Mund, ex Meissn. in DC. Prod. xiv. 304 = scabridus.
phylicoides, *Meissn. l. c.*—Afr. austr.
salsoloides, *R. Br. in Trans. Linn. Soc.* x. (1810) 140.—Afr. austr.
scabridus, *Meissn. in DC. Prod.* xiv. 303.—Afr. austr.
setaceus, *R. Br. in Trans. Linn. Soc.* x. (1810) 140.—Afr. austr.
spatalloides, *R. Br. l. c.* 141.—Afr. austr.
spatalloides, Sieber, ex Meissn. in DC. Prod. xiv. 305 = lanatus.
tenuifolius, *R. Br. in Trans. Linn. Soc.* x. (1810) 141.—Afr. austr.
verticillatus, Roem. & Schult. Syst. iii. 391 = Leucadendron verticillatum.

SOROSTACHYS, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Cyp. 71 (1856) = **Cyperus**, Michx.
kyllingioides, Steud. ll. cc. = C. leucocephalus.

SOTOR, Fenzl, in Ber. 21ste Versamml. Deutsch. Naturf. 1843 (1844) 168 = **Kigelia**, DC. (1845) (Bignon.).
aethiopum, Fenzl, l. c. = K. pinnata.

SOTULARIA, Rafin. Sylva Tellur. 98 (1838) = **Lagerstroemia**, Linn. (Lythrar.).
malabarica, Rafin. l. c. = L. hirsuta.

SOUARI, Endl. Gen. 1076 (1840), sphalm. = SAOUARI, Aubl. = **Caryocar**, Linn. (Ternstr.).

SOUBEYRANIA, Neck. Elem. i. 365 (1790) = **Barleria**, Linn. (Acanth.).

SOULAMEA, Lam. Encyc. i. 449 (1783). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
CARDIOCARPUS, Reinw. in Syll. Ratisb. ii. (1828) 14.
CARDIOPHORA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 216.
amara, *Lam. Encyc.* i. 449.—Malaya.
cardioptera, *Baill. Adansonia*, x. (1871-72) 344.—N. Caled.
fraxinifolia, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 244.—N. Caled.
Muelleri, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. iii. (1865) 231.—N. Caled.
Pancheri, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 243.—N. Caled.
terminalioides, *Baker, Fl. Maurit.* 42.—Ins. Seychelles.
tomentosa, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 243.—N. Caled.
trifoliata, *Baill. Adansonia*, x. (1871-72) 344.—N. Caled.

SOULANGIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 377. t. 10. f. 3 = **Phylica**, Linn. (Rhamn.).
arborea, G. Don, Gen. Syst. ii. 43 = P. nitida.
arborescens, Eckl. & Zeyh. Enum. 136 = P. paniculata.
axillaris, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 378 = P. axillaris.
buxifolia, Brongn. l. c. 378 = P. buxifolia.
cordata, Brongn. l. c. = P. buxifolia.
dioica, G. Don, Gen. Syst. ii. 43 = P. dioica.
epacridifolia, A. Dietr. in Otto & Dietr. Allg. Gartenz. vii. (1839) 387.—Afr. austr.
glauca, Hort. ex A. Dietr. l. c. 388.—Hab. ?
hirsuta, Eckl. & Zeyh. Enum. 136 = P. axillaris.
ledifolia, Eckl. & Zeyh. l. c. = P. ledifolia.
lutescens, Eckl. & Zeyh. l. c. = P. lutescens.

SOULANGIA:—
marifolia, Bernh. ex Krauss, in Flora, xxvii. (1844) 348 = Phylica paniculata.
microphylla, Eckl. & Zeyh. Enum. 135 = P. microphylla.
myrtifolia, A. Dietr. in Otto & Dietr. Allg. Gartenz. vii. (1839) 381 = P. paniculata.
oleaefolia, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 378 = P. oleaefolia.
oppositifolia, Reiss. in Endl. Nov. Stirp. Dec. 79 = Nesiota elliptica.
orientalis, Sweet, Hort. Brit. ed. II. 115 = P. orientalis.
paniculata, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 378 = P. paniculata.
parviflora, Presl, Bot. Bemerk. 38 = P. rosmarinifolia.
pedicellata, G. Don, Gen. Syst. ii. 42 = P. villosa.
pinea, Eckl. & Zeyh. Enum. 136 = P. villosa, axillaris.
plumosa, Presl, Bot. Bemerk. 38 = P. ambigua.
reclinata, G. Don, Gen. Syst. ii. 42 = P. reclinata, *Wendl.*
rosmarinifolia, Harv. Gen. S. Afr. Pl. 62 = P. rosmarinifolia.
rubra, Hort. ex Otto & Dietr. Allg. Gartenz. vii. (1839) 381 = P. paniculata.
rubra, Lindl. Bot. Reg. t. 1498 = P. rubra.
subcanescens, Presl, Bot. Bemerk. 38 = P. cryptandroides.
thymifolia, Brongn. in Ann. Sc. Sér. I. x. (1827) 378 = P. thymifolia.
thymifolia, Presl, Bot. Bemerk. 38.—Afr. austr.
Willdenowiana, A. Dietr. in Otto & Dietr. Allg. Gartenz. vii. (1839) 386 = P. Willdenowiana.

SOULEYETIA, Gaudich. Bot. Voy. Bonite, t. 19 (1844-49) = **Pandanus**, Rumph.
freycinetioides, Gaudich. l. c. = P. freycinetioides.

SOUROUBEA, Aubl. Pl. Gui. i. 244. t. 97 (1775) = **Ruyschia**, Jacq. (Ternstr.).
amazonica, Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 275 = R. amazonica.
Aubletii, G. F. W. Mey. Prim. Fl. Esseq. 119 = R. Souroubea.
auriculata, Delp. in Att. Soc. Ital. Sc. Nat. Milan. xii. (1869) 180 = R. Souroubea.
bahiensis, Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 276 = R. bahiensis.
bicolor, Delp. l. c. = R. bicolor.
corallina, Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 276 = R. corallina.
crassipes, Wittm. in Mart. Fl. Bras. xii. 1. 254 = R. crassipes.
exauriculata, Delp. in Att. Soc. Ital. Sc. Nat. Milan. xii. (1869) 180 = R. clusiaefolia (in Addend.).
guianensis, Aubl. Pl. Gui. i. 244. t. 97 = R. Souroubea.
lepidota, Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 276 = R. lepidota.
Meyeri, Delp. l. c. 275 = R. Souroubea.
pileata, Delp. l. c. 258, 275 = R. pilophora.
pilophora, Wittm. in Mart. Fl. Bras. xii. 1. 254 = R. pilophora.
Spitziana, Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 275 = R. Spixiana.

SOUTHWELLIA, Salisb. Parad. Lond. t. 69 (1807) = **Sterculia**, Linn.
angustifolia, Wight, Illustr. Ind. Bot. i. 77 = St. Balanghas.
Balanghas, Spach, Hist. Vég. Phan. iii. 508 = St. Balanghas.
Blumii, Wight, Illustr. Ind. Bot. i. 77 = St. Blumei.
coccinea, Voigt, Hort. Suburb. Calc. 104 = St. coccinea.
cordifolia, Spach, Hist. Vég. Phan. xiv. 402 = St. cordifolia.
longifolia, Rafin. Sylva Tellur. 73 = Cola acuminata ?
nobilis, Salisb. Parad. Lond. t. 69 = St. nobilis.
parviflora, Voigt, Hort. Suburb. Calc. 104 = St. parviflora.
Roxburghiana, Spach, Hist. Vég. Phan. xiv. 402 = St. Roxburghii.
Tragacanthae, Wight, Illustr. Ind. Bot. i. 77 = St. Tragacanthae.
urceolata, Spach, Hist. Vég. Phan. xiv. 402 = St. urceolata.
versicolor, Wight, Illustr. Ind. Bot. i. 77 = St. versicolor.

SOUZA, Vell. Fl. Flum. 273 (1825); vii. t. 2 (1827) = **Sisyrinchium**, Linn. (Irid.).
Comes, Vell. l. c. = Sis. Comes.
Marchio, Vell. l. c. t. 1 = Sis. alatum.
Proregia, Vell. l. c. 274. t. 3 = Sis. Proregia.

SOWERBAEA, Sm. in Trans. Linn. Soc. iv. (1800) 218. *LILIACEAE*, Benth. & Hook. f. iii. 797.
alliacea, *F. Muell. Fragm.* vi. 180.—Austral.
americana, Spreng. ex Baker, in Journ. Linn. Soc. xi. (1871) 381 = Milla macrostemon.
juncea, *Sm. in Trans. Linn. Soc.* v. (1800) 159.—Austral.
laxiflora, *Lindl. Bot. Reg.* (1841) t. 10; *Swan River App.* 58.—Austral.

SOYA, Benth. in Ann. Wien. Mus. ii. (1838) 113 = SOJA, Moench = **Glycine**, Linn. (Legumin.).

SOYAUXIA, Oliver, in Hook. Ic. Pl. t. 1393 (1882). *PASSIFLOREAE.*
gabonensis, *Oliver, l. c.*—Afr. trop.

SOYERA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881). 171 = seq.

SOYERIA, Monn. Ess. Hierac. 75 (1829) = **Crepis**, Vaill. (Compos.).
alpestris, Mutel, Fl. Fr. ii. 223 = C. alpestris.
alpina, Gren. & Godr. Fl. Fr. ii. 327 = Lactuca alpina.
aurea, Mutel, Fl. Fr. ii. 223, 418 = C. aurea.
blattarioides, Monn. Ess. Hierac. 76 = C. blattarioides.
chrysantha, D. Dietr. Syn. Pl. iv. 1331 = C. chrysantha.
crocifolia, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 322 = C. crocifolia.
grandiflora, Monn. Ess. Hierac. 76 = C. grandiflora.
hieracioides, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 321 = C. hieracioides.
hyoseridifolia, Koch, Syn. Fl. Germ. ed. I. 442 = C. hyoseridifolia.
lampsanoides, Monn. Ess. Hierac. 77 = C. lampsanoides.
lapsanoides, Mutel, Fl. Fr. ii. 223 = praec.
lyrata, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 322 = C. lyrata.
montana, Monn. Ess. Hierac. 75 = C. montana.
paludosa, Godr. Fl. Lorr. ii. 72 = C. paludosa.
praemorsa, Mutel, Fl. Fr. ii. 224 = C. praemorsa.
pyrenaica, Sch. Bip. in Pollichia, xxii-xxiv. (1866) 322 = C. blattarioides.
serbica, Sch. Bip. l. c. = C. viscidula.
sibirica, Monn. Ess. Hierac. 77 = C. sibirica.
viscidula, D. Dietr. Syn. Pl. iv. 1332 = Hieracium praealtum.

SOYMIDA, A. Juss. in Mém. Mus. Par. xix. (1830) 250. t. 22. f. 26. *MELIACEAE*, Benth. & Hook. f. i. 338.
febrifuga, *Juss. l. c.* 251. t. 22. f. 26.—Ind. or.
roupalifolia, *Schweinf. Reliq. Kotschy.* 37.—Afr. trop.

SPACHEA, A. Juss. in Deless. Ic. Sel. iii. 19. t. 31 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 255.
MECKELIA, Mart. ex A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 326, in syn.
elegans, *A. Juss. in Deless. Ic. Sel.* iii. 19.—Guian.
littoralis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 328.—Ind. occ.
ossana, *A. Juss. l. c.* 327.—Ind. occ.
parviflora, *A. Juss. l. c.* 326.—Ind. occ.
perforata, *A. Juss. l. c.* 329.—Ind. occ.
tenuifolia, *Griseb. in Linnaea*, xxii. (1849) 9.—Guian.
tricarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 329.—Bras.

SPACHIA, Lilja, in Traedgardtidning, (1840) 62; ex Linnaea, xv. (1841) 262 = **Fuchsia**, Linn. (Onagr.).

SPADACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827) = **Atractylis**, Linn. (Compos.).
flava, Cass. l. c. = A. flava.
radiciflora, Cass. l. c. = A. radiciflora.

SPADONIA, Less. Syn. Comp. 99 (1832) = **Moquinia**, DC. (Compos.).
cinerea, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 109 = M. polymorpha.
paniculata, Less. Syn. Comp. 100 = M. paniculata.
polymorpha, Less. l. c. 99 = M. polymorpha.
racemosa, Less. l. c. 100 = M. racemosa.

SPADOSTYLES, Benth. in Ann. Wien. Mus. ii. (1838) 81 = **Pultenaea**, Sm. (Legumin.).
Benthamii, Endl. Nov. Stirp. Dec. 3 = P. humilis.
concolor, Endl. l. c. 24 = P. villosa.
Cunninghami, Benth. in Ann. Wien. Mus. ii. (1838) 81 = P. ternata.
Huegelii, Endl. Nov. Stirp. Dec. 3 = P. humilis.
ramulosa, Endl. l. c. 25 = P. villosa.
Sieberi, Benth. in Ann. Wien. Mus. ii. (1838) 81 = P. euchila.
ternata, F. Muell. First Gen. Rep. 12 = P. ternata.

SPAENDONCEA, Desf. in Dec. Phil. vii. 259 (1796 ?); ex Usteri, Neue Ann. Bot. xiii. (1796) 15 = **Cadia**, Forsk. (Legumin.).
tamarindifolia, Desf. l. c. 18 = C. purpurea.

SPALANTHUS, Walp. Rep. ii. 68 (1843), sphalm. = SPHALANTHUS, Jack = **Quisqualis**, Rumph. (Combret.).

SPALLANZANIA, DC. Prod. iv. 406. (1830) = **Mussaenda**, DC. (Rubiac.).
corymbosa, DC. l. c. = Madag.

SPALLANZANIA, Neck. Elem. ii. 79 (1790) = **Gustavia**, Linn. (Myrtac.).

SPALLANZANIA, Pollini (Hort. Veron. Pl. Nov. 10), in Brugnat. Giorn. Fis. ix. (1816) 28 = **Agrimonia**, Linn. (Rosac.).
agrimonoides, Pollini, ll. cc. = A. Eupatoria.

SPANANTHE, Jacq. Coll. iii. 247 (1789). *UMBELLIFERAE*, Benth. & Hook. f. i. 876.
angulosa, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 171 = paniculata.
Huanaca, Lag. Amen. Nat. ii. 93 = Huanaca Cavanillesii.
paniculata, *Jacq. Coll.* iii. 247.—Reg. Andin.
saniculaefolia, Spreng. in Schult. Syst. vi. 357 = Diposis saniculaefolia.
setosa, Moench, Meth. Suppl. 34 = paniculata.
sinuata, Spreng. Syst. i. 878 = Hydrocotyle acutifolia.

SPANIOPTILON, Less. Syn. Comp. 10 (1832) = **Cnicus**, Linn. (Compos.).
lineare, Less. l. c. = C. linearis.

SPANIZIUM, Griseb. Spicil. Fl. Rumel. i. 180 (1843) = **Saponaria**, Linn. (Caryoph.).
ocymoides, Griseb. l. c. = Sap. prostrata.

SPANOGHEA, Blume, Rumphia, iii. 173 (1850) *SAPINDACEAE*, Benth. & Hook. f. i. 406.
connata, F. Muell. in Tarns. Phil. Inst. Vict. iii. (1859) 26 = Nephelium connatum.
ferruginea, *Blume, Rumphia*, iii. 173.—N. Guin.
glabra, *Blume, l. c.* 174.—Ins. Timor.
nepheleoides, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 25; Vict. Pl. i. 227 = Nephelium leiocarpum.

SPANOTRICHUM, E. Mey. ex DC. Prod. vi. 291 (1837) = **Osmites**, Linn. (Compos.).
pinnatifidum, E. Mey. ex DC. l. c. = O. pinnatifida.

SPARATTANTHELIUM, Mart. in Flora, xxiv. (1841) ii. Beibl. 40. *COMBRETACEAE*, Benth. & Hook. f. i. 690.
amazonum, *Mart. l. c.* 42.—Bras.
bororum, *Mart. in Denkschr. Bot. Ges. Regensb.* iii. (1841) 302.—Bras.
botocudorum, *Mart. l. c.* 301. t. 10. f. 2; t. 11. f. 1.—Bras.
cordatum, *Meissn. in Mart. Fl. Bras.* v. ii. 294.—Bras.
tarapotanum, *Meissn. l. c.*—Bras.
tupinambazum, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 41.—Bras.
tupiniquinorum, *Mart. l. c.* 40.—Bras.

SPARATTOSPERMA, Mart. ex DC. in Meissn. Gen. 300; Comm. 208 (1840). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1041.
lithontripticum, *Mart. in DC. Prod.* ix. 203.—Bras.
psammophilum, *Mart. l. c.*—Bras.

SPARATTOSYCE, Bur. in Ann. Sc. Nat. Sér. V. xi. (1869) 379. t. 6. *URTICACEAE*, Benth. & Hook. f. iii. 369.
dioica, *Bur. l. c.* 380.—N. Caled.

SPARAXIS, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 225. *IRIDEAE*, Benth. & Hook. f. iii. 708.
ANACTORION, Rafin. Fl. Tellur. iv. 34 (1836).
albiflora, Eckl. Top. Verz. 28 = bulbifera.
anemoneflora, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 225 = Ixia anemonaeflora.
atropurpurea, Hort. ex Baker, in Journ. Linn. Soc. xvi. (1877) 99 = Dierama pulcherrimum.
atropurpurea, Klatt, in Abh. Naturf. Ges. Halle. xv. (1882) 389 = grandiflora.
bicolor, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 225 = Synnottia bicolor.
blanda, Sweet, Hort. Brit. ed. I. 398 = tricolor.
bulbifera, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 226.—Afr. austr.
cana, *Eckl. Top. Verz.* 28.—Afr. austr.
cuprea, Klatt, in Linnaea, xxxv. (1867–68) 378 = Streptanthera cuprea.
fimbriata, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 226 = grandiflora.
fragrans, Ker-Gawl. l. c. 225 = Ixia fragrans.
galeata, Ker-Gawl. l. c. = Synnottia galeata.
grandiflora, *Ker-Gawl. l. c.*—Afr. austr.
Griffini, Sweet, Hort. Brit. ed. I. 398 = tricolor.
Jauberti, Lodd. ex Fenzl, Del. Sem. Hort. Vindob. (1855), sphalm. = seq.
Joubertii, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 164 = Freesia refracta.
lacera, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 226 = grandiflora.
Liliago, Sweet, Hort. Brit. ed. I. 398 = grandiflora.
lineata, Sweet, Brit. Flow. Gard. Ser. II. t. 131 = grandiflora.
lutea, *Eckl. Top. Verz.* 28.—Afr. austr.
luteo-violacea, Eckl. l. c. 27 = Synnottia variegata.
Meleagris, *Eckl. l. c.*—Afr. austr.
miniata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 389.—Afr. austr.
orchidiflora, *Lodd. Bot. Cab.* t. 1099.—Afr. austr.
pendula, Ker-Gawl. in Bot. Mag. sub t. 1482; Lindl. Bot. Reg. t. 1360 = Dierama ensifolium.
pulcherrima, Hook. f. Bot. Mag. t. 5555 = Dierama pulcherrimum.
reflexa, *Villet, ex Voigt, Hort. Suburb. Calc.* 610, nomen.—Afr. austr.
stellaris, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 383 = grandiflora.
Thouberti, Hort. ex Klatt, in Gartenfl. (1874) 289 = Freesia Leichtlini.
tricolor, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 225.—Afr. austr.
versicolor, Sweet, Brit. Flow. Gard. Ser. I. t. 160 = tricolor.
violacea, *Eckl. Top. Verz.* 27.—Afr. austr.
Wattii, Harv. Gen. S. Afr. Pl. 330 = Synnottia variegata.

SPARGANION, Adans. Fam. ii. 471 (1763) = seq.

SPARGANIUM, [Tourn.] Linn. Syst. ed. I (1735). *TYPHACEAE*, Benth. & Hook. f. iii. 955.
PLATANARIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 39 (1821).
affine, Schnitz. Typh. 27 = simplex.
alpinum, D. Don, ex G. Don, in Loud. Hort. Brit. 375, nomen = simplex.
americanum, Nutt. Gen. Am. ii. 203 = angustifolium.
angustifolium, *Michx. Fl. Bor. Am.* ii. 189.—Am. bor.; Austral.
Borderi, *Focke, in Bremen. Abh.* v. (1877) 409.—Europ.
boreale, Laest. ex Beurl. in Oefvers. Vet. Akad. Foerhand. ix. (1852) 192 = simplex.
californicum, *Greene, in Bull. Calif. Acad.* i. iii. (1884) 11.—Calif.
carinatum, *Falc. in Proc. Linn. Soc.* i. (1839) 18, nomen (Quid ?).—Cashmir.
emersum, Rehm. in Verh. Nat. Ver. Bruenn, x. 1871 (1872) 80 = simplex.
erectum, Linn. Sp. Pl. 971 = ramosum.

SPARGANIUM:—
erectum, Wahlenb. Fl. Suec. ii. 604 = simplex.
eurycarpum, Engelm. in A. Gray, Man. Bot. N. U. St. ed. V. 481, partim = angustifolium.
fluitans, *Fries, in Bot. Notiser* (1849), 14, in obs.—Europ.
Friesii, Beurl. l. c. (1854), 136 = natans.
glomeratum, Laest. ex Beurl. in Oefvers. Vet. Akad. Foerhand. ix. (1852) 192 = fluitans.
gramineum, Georgi, Reise, i. 232 (Quid ?).—Rossia.
gramineum, Wallr. Beitr. Fl. Hercyn. 297 = minimum.
hyperboreum, *Laest. ex Beurl. in Oefvers. Vet. Akad. Foerhand.* ix. (1852) 192.—Island.; Scand. bor.
lanceolatum, Georgi, Reise, i. 233 = natans.
longifolium, Turcz. in Bull. Soc. Nat. Mosc. (1838) 103 = natans.
minimum, *Fries, Summa Veg. Scand.* 68, 560.—Europ.
natans, Fries, in Bot. Notiser (1849), 13; (1852), 92 = erectum.
natans, *Linn. Sp. Pl.* 971.—Europ.
natans, Oed. Fl. Dan. t. 260 = minimum.
neglectum, *Beeby, in Journ. Bot.* xxiii. (1885) 26, 193.—Europ.
oligocarpon, *Ångstr. in Bot. Notiser* (1853), 149.—Scand.
pubescens, *Poir. Voy. Barb.* ii. 254.—Afr. bor.
ramosum, *Curt. Fl. Loud.* fasc. v. t. 66.—Europ.; Sibir.; Am. bor.; Ind. or.
rectum, Neck. Delic. Gallo-Belg. ii. 377 = ramosum.
rostratum, Larss. Fl. Werml. (1859) 260 = minimum.
simplex, *Huds. Fl. Angl.* ed. II. 401.—Europ.; Sibir.; Am. bor.
simplex, Muhl. Cat. ed. I. 82; ed. II. 87 = angustifolium.
stoloniferum, Buch.-Ham. in Wall. Cat. n. 4990 = ramosum.
tenuifolium, *Poir. Voy. Barb.* ii. 254.—Afr. bor.
trifidum, Poir. Encyc. Suppl. iv. 723 = ramosum.
vaginatum, Larss. Fl. Werml. (1859) 259 = simplex.

SPARGANOPHORUS, Vaill. ex Crantz, Inst. i. 261 (1766); Gaertn. Fruct. ii. 395. t. 165 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 223.
ATHENAEA, Adans. Fam. ii. 121 (1763).
STRUCHIUM, P. Br. Hist. Jamaic. 312. t. 34 (1756).
africanus, Steud. Nom. ed. I. 801 = Vaillantii.
ageratoides, Lag. Gen. et Sp. Nov. 25 = Ageratum corymbosum.
Ethulia, Crantz, Inst. i. 261 = Ethulia conyzoides.
fasciatus, Lam. Illustr. i. 670; Encyc. vii. 302 = Vaillantii.
fasciculatus, Steud. Nom. ed. I. 801, sphalm. = praec.
fuscatus, Steud. l. c. ed. II. ii. 614, sphalm. = praec.
obtusifolius, Lag. Gen. et Sp. Nov. 25 = Ageratum conyzoides.
sonchifolius, Spreng. Syst. iii. 458 = Cephalopappus sonchifolius.
Struchium, Poir. Encyc. vii. 302 = Vaillantii.
Vaillantii, *Crantz, Inst.* i. 261.—Ind. occ.
verticillata, Michx. Fl. Bor. Am. ii. 95 = Sclerolepis verticillata.

SPARMANNIA, Buc'hoz, Pl. Nouv. Découv. t. 1 (1779) = **Rehmannia**, Libosch. (1835) (Scrophul.).
sinensis, Buc'hoz, l. c. = R. chinensis.

SPARMANNIA, Linn. f. Suppl. 41 (1781). *TILIACEAE*, Benth. & Hook. f. i. 235.
abyssinica, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 29, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 79.—Abyss.
acerifolia, Hort. ex Steud. Nom. ed. II. ii. 614 = africana.
africana, *Linn. f. Suppl.* 265.—Afr. austr.
discolor, *Baker, in Journ. Linn. Soc.* xx. (1883) 102.—Madag.
palmata, Hort. ex Lindl. Bot. Reg. (1838) Misc. 66 = Entelea palmata.
palmata, *E. Mey. ex Presl, Bot. Bemerk.* 19.—Afr. austr.
rugosa, Bivona ?, ex Sweet, Hort. Brit. ed. II. 70 (Quid ?).—Afr. austr.
subpalmata, *Baker, in Journ. Linn. Soc.* xx. (1883) 101.—Madag.

SPARTHOTHAMNUS, Buek, Ind. ad DC. Prod. iii. 442 (1858), sphalm. = **Spartothamnus**, A. Cunn. (Verben.).

SPARTIANTHUS, Link, Enum. Hort. Berol. ii. 223 (1822) = **Spartium**, Linn. (Legumin.).
americanus, Steud. Nom. ed. II. ii. 614 = Spartium junceum.
junceus, Link, Enum. Hort. Berol. ii. 223 = Spartium junceum.

SPARTIDIUM, Pomel, Nouv. Mat. Fl. Atl. 173 (1874) = **Genista**, Linn. (Legumin.).
Saharae, Pomel, l. c. = G. Saharae.

SPARTINA, Schreb. Gen. 43 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
CHAUVINIA, Steud. Syn. Pl. Gram. 362 (1855).
LIMNETIS, Pers. Syn. i. 72 (1805).
PONCELETIA, Thou. Fl. Trist. Acugn. 36 (1811).
SOLENACHNE, Steud. Syn. Pl. Gram. 12 (1854).
TRACHYNOTIA, Michx. Fl. Bor. Am. i. 63 (1803).
TRISTANIA, Poir. Encyc. Suppl. iv. 526, in obs. (1816).
alterniflora, Loisel. Fl. Gall. 719; ed. II. i. 41 = stricta.
arundinacea, *Carmich. in Trans. Linn. Soc.* xii. (1819) 504.—Ins. Trist. Acuna; Amst.
bahiensis, *Steud. Nom.* ed. II. ii. 474, in syn.—Bras.
brasiliensis, *Raddi, Agrost. Bras.* 211.—Bras.
capensis, *Nees, Fl. Afr. Austr.* 260.—Afr. austr.
ciliata, *Kunth, Rév. Gram.* i. 94.—Am. austr.
coarctata, Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 110 = ciliata.
cynosuroides, Roth, Catalect. fasc. iii. 10; Willd. Enum. Hort. Berol. 80 = Schreberi.
densiflora, *Brongn. in Duperr. Voy. Coq. Bot.* 14.—Chili.
dissitiflora, *Steud. Syn. Pl. Gram.* 215.—Bras.
Duriaei, Parl. Fl. Ital. i. 230 = versicolor.
fasciculata, *Beauv. Agrost.* 25.—Am. austr.
foliosa, Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 114 = stricta.
geniculata, *Beauv. Agrost.* 25.—Java.
glabra, Muhl. Desc. Gram. i. 54.—Am. bor.
glabriuscula, *Hassk. Cat. Hort. Bog. Alt.* 18, nomen.—Java.
Gouini, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 509.—Mexic.
gracilis, *Hook. ex Chapm. Fl. S. U. St.* 556.—Florida.
gracilis, *Trin. in Mém Acad. Pétersb.* Sér. VI. v. II. (1840) 110.—Am. bor.
humilis, [Willd. ex] Link, Jahrb. i. III. (1820) 93 = Bouteloua ovata.
juncea, *Willd. Enum. Hort. Berol.* 81.—Am. bor.
junciformis, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 238.—Am. bor.
laevigata, *Bosc, ex Link, Jahrb.* i. III. (1820) 92.—Am. bor.
leiantha, *Benth. Bot. Voy. Sulph.* 56.—Mexic.
patens, *Muhl. Desc. Gram.* 55.—Am. bor.
pectinata, *Bosc, ex Link, Jahrb.* i. III. (1820) 92.—Am. bor.
phleoides, Roth, Neue Beytr. i. 101 = Heleochloa sehoenoides.
polystachya, *Willd. Enum. Hort. Berol.* 81, in obs.—Am. bor.
pubera, *Hassk. Cat. Hort. Bog. Alt.* 18, nomen.—Java.
pumila, *Roth, Catalect.* fasc. iii. 10.—Am. bor.
pungens, Beauv. Agrost. 25 = stricta.
Schreberi, *J. F. Gmel. Syst.* 123.—Am. bor.
stricta, *Roth, Neue Beytr.* i. 101.—Europ.; Am. bor. et austr.
Townsendi, H. & J. Groves, in Bot. Exch. Club Rep. 1880 (1881), 37 = stricta.
versicolor, *Fabre, in Ann. Sc. Nat.* Sér. III. xiii. (1849) 123. t. 3.—Gallia.

SPARTIUM, Linn. Syst. ed. I (1735); Gen. ed. I. 218 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 483.
SPARTIANTHUS, Link, Enum. Hort. Berol. ii. 223 (1822).
acanthocladon, Spreng. Syst. iv. Cur. Post. 266 = Genista acanthoclada.
acutifolium, Lindl. Bot. Reg. t. 1974 = junceum.
aetnense, Bivona, Stirp. Rar. Sicil. ii. [13] = Genista aethnensis.
affine, *Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 296.—Ins. Canar.
albicans, Cav. in Anal. Cienc. Nat. iv. (1801) 64 (= *Teline canariensis*).—Ins. Canar.
album, Desf. Fl. Atlant. ii. 132 = Cytisus albus.

SPARTIUM:—
alpinum, Host, ex Steud. Nom. ed. II. ii. 615, nomen = Laburnum alpinum?
ambiguum, *Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 295.—Ins. Canar.
americanum, Meyen, Reise, i. 445 = junceum.
amoenum, Salisb. Prod. 330 = Cytisus linifolius.
angulatum, Linn. Sp. Pl. 709 = Genista parviflora.
angulatum, Rafin. Préc. Découv. 37 = Cytisus scoparius.
angulosum, Gilib. Fl. Lituan. ii. 79 = Cytisus scoparius.
apetalum, Labill. ex Steud. Nom. ed. I. 801 = Genista apetala.
aphyllum, Pall. Reise, iii. 742 = Eremosparton aphyllum.
aragonense, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 434 = Genista cinerea.
arborescens, Mill. Gard. Dict. ed. VIII. n. 10 = Brya Ebenus.
arboreum, Desf. Fl. Atlant. ii. 131 = Cytisus arboreus.
argenteum, Hort. Angl. ex Steud. Nom. ed. II. ii. 615, nomen (Quid?).—Ins. Canar.
aspalatoides, Desf. Fl. Atlant. ii. 136 = Genista aspalathoides.
biflorum, Desf. l. c. 133. t. 179 = Cytisus Fontanesii.
Bovei, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 297 = Genista monosperma.
capense, Linn. Pl. Afr. Rar. 14; Amoen. Acad. vi. 91 = Rafnia opposita.
capitatum, Barnado, ex Cav. in Anal. Cienc. Nat. iv. (1801) 61 = Genista umbellata.
capitatum, Cav. l. c. 63 = Genista clavata.
cinereum, Vill. Prosp. 40 = Genista cinerea.
Clusii, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 290. t. 16. f. 3 = Genista monosperma.
complicatum, Gouan, Hort. Monsp. 356 = Adenocarpus grandiflorus?
complicatum, Linn. Sp. Pl. 709 = Adenocarpus intermedius.
congestum, Willd. Enum. Hort. Berol. 744 = Cytisus congestus.
contaminatum, Ait. Hort. Kew. ed. II. iv. 261 = Indigofera filifolia.
contaminatum, Linn. Mant. ii. 268 = Lebeckia contaminata.
corsicum, Loisel. Fl. Gall. ed. I. 440 = Genista corsica.
creticum, Desf. Tabl. ed. I. 185 = Anthyllis Hermanniae.
Cupani, Spreng. Syst. iii. 177 = Genista Cupani.
cuspidatum, Cav. in Anal. Cienc. Nat. iv. (1801) 56 = Genista hirsuta.
cuspidosum, Burch. Trav. i. 348 = Lebeckia psiloloba.
cytisoides, Linn. f. Suppl. 320 = Lebeckia cytisoides.
dalmaticum, Spreng. Syst. iv. Cur. Post. 266 = Genista aristata.
decumbens, Ait. Hort. Kew. ed. I. iii. 11 = Cytisus decumbens.
dispermum, Willd. in Roem. & Usteri, Mag. iv. 11tes St. (1790) 35 = Cytisus albus.
dubium, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 295 = Genista monosperma.
Duriaei, *Spach, l. c.* 289. t. 16. f. 2.—Afr. bor.
ellipticum, Willd. ex Spreng. Syst. iii. 177 = Genista elliptica.
ephedroides, Spreng. l. c. iv. Cur. Post. 266 = Genista ephedroides.
erinaceoides, Loisel. Fl. Gall. ed. I. 441 = Genista aspalathoides.
ferox, Poir. Voy. Barb. ii. 206 = Genista ferox.
filifolium, Willd. ex Walp. in Linnaea, xiii. (1839) 461 = Viminaria denudata.
fragrans, Lam. ex Steud. Nom. ed. I. 801, 802 = Cytisus fragrans.
Gasparinii, Guss. Ind. Sem. Hort. Boccad. (1825) 11 = Genista ephedroides.
glabrum, Mill. Gard. Dict. ed. VIII. n. 7 = Cytisus scoparius.
gracile, Salisb. Prod. 329 = Genista monosperma.
grandiflorum, Brot. Fl. Lusit. ii. 80 = Cytisus grandiflorus.
grandiflorum, Salisb. Prod. 329 = Genista sphaerocarpa.
gymnopterum, Viv. App. Fl. Cors. Prod. 6 = Genista ephedroides.
Halleri, Auct. ex Lam. Encyc. ii. 618 = Cytisus decumbens.
heterophyllum, L'Hérit. Stirp. Nov. 183. t. 87 = Genista ferox.

SPARTIUM:—
hirsutum, Mill. Gard. Dict. ed. VIII. n. 6 = Cytisus Milleri.
hispanicum, Spreng. Syst. iii. 177 = Genista hispanica.
horridum, Sibth. & Sm. Fl. Graec. vii. 69. t. 674 = Genista acanthoclada.
horridum, Vahl, Symb. Bot. i. 51 = Genista horrida.
humile, Salisb. Prod. 330 = Genista germanica.
infestum, Presl, Delic. Prag. 33 = Genista infesta.
interruptum, Cav. in Anal. Cienc. Nat. iv. (1801) 58 = Genista triacanthos.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 43.—Japon.
junceum, *Linn. Sp. Pl.* 708.—Geront. temp.
Laburnum, Cav. ex Steud. Nom. ed. II. ii. 615 = Laburnum vulgare.
lanigerum, Desf. Fl. Atlant. ii. 135 = Calycotome villosa.
lanigerum, [Soland. in] Russ. Aleppo, ed. II. ii. 269 (Quid?).—Syria.
lasiocarpum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 460 = Cytisus patens.
linifolium, Desf. Fl. Atlant. ii. 134. t. 181 = Cytisus linifolius.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 5 = Cytisus grandiflorus.
microcarpum, *Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 292. t. 16. f. 5.—Ins. Canar.
microphyllum, Cav. in Anal. Cienc. Nat. iv. (1801) 63 = Adenocarpus foliolosus.
molle, Cav. l. c. 67 = Genista mollis.
monospermum, Linn. Sp. Pl. 708 = Genista monosperma.
monospermum, *Viv. Fl. Lib. Spec.* 39 = Genista Raetam.
multicaule, Hort. ex Lam. Encyc. ii. 617 = Genista multicaulis.
multiflorum, Ait. Hort. Kew. ed. I. iii. 21 = Cytisus albus.
nitens, Willd. ex Spreng. Syst. iii. 177 = Genista nitens.
nubigenum, Ait. Hort. Kew. ed. I. iii. 13 = Cytisus fragrans.
nudum, Willd. ex Spreng. Syst. iii. 176 = Genista nuda.
odoratissimum, D. Don, ex Steud. Nom. ed. II. ii. 615 = junceum.
odoratum, Dulac, Fl. Hautes-Pyr. 275 = junceum.
Osyris, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 433 = Genista ramosissima.
ovatum, Berg. Desc. Pl. Cap. 197 = Rafnia cuneifolia.
parviflorum, Vent. Jard. Cels, 87 = Genista parviflora.
patens, Murr. Syst. ed. XIII. 535 = Cytisus patens.
persicum, Willd. Sp. Pl. iii. 931 = Crotalaria furfuracea.
pilosum, Roth, Tent. Fl. Germ. i. 303 = Genista pilosa.
pinastrifolium, Burm. f. Fl. Cap. Prod. 20 = Lebeckia sepiaria.
procerum, Willd. Enum. Hort. Berol. 743 = Cytisus procerus.
pungens, Willd. ex Spreng. Syst. iii. 225 = Cytisus pungens.
purgans, Linn. Syst. ed. XII. 474 = Cytisus purgans.
pusillum, Salisb. Prod. 330 = Genista anglica.
radiatum, Barnado, ex Willk. & Lange, Prod. Fl. Hisp. iii. 423 = Genista Barnadesii.
radiatum, Linn. Sp. Pl. 708 = Genista radiata.
radiatum, Vill. Hist. Pl. Dauph. iii. 419 = Genista horrida.
Raetam, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 288. t. 16. f. 1 = Genista Raetam.
ramosissimum, Desf. Fl. Atlant. ii. 132. t. 178 = Genista ramosissima.
rhodorhizoides, Walp. Rep. ii. 840 = semperflorens.
rigidum, Viv. Fl. Lib. Spec. 40. t. 17. f. 1 = Calycotome villosa.
rostratum, *Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 293.—Ins. Canar.
rostratum, Spreng. Syst. iii. 177 = Genista triacanthos.
sagittale, Roth, Tent. Fl. Germ. i. 302 = Genista sagittalis.
scoparium, Linn. Sp. Pl. 709 = Cytisus scoparius.
Scorpius, Linn. l. c. 708 = Genista Scorpius.
semperflorens, *Spach, in Ann. Sc. Nat.* Sér. II. xix. (1843) 294. t. 16. f. 10.—Ins. Canar.
sepiarium, Linn. Amoen. Acad. vi. 91 = Lebeckia sepiaria.
sericeum, Ait. Hort. Kew. ed. I. iii. 12 = Genista sericea.
sericeum, Presl, ex Steud. Nom. ed. I. 802 = Genista infesta.

SPARTIUM:—
sericeum, Vent. Choix, 17 = Genista clavata.
sessilifolium, Spreng. Syst. iv. Cur. Post. 266 = Genista sessilifolia.
sophoroides, Berg. Desc. Pl. Cap. 198 = Hypocalyptus obcordatus.
sphaerocarpum, Desf. ex Spach, in Ann. Sc. Nat. Sér. III. ii. (1844) 244 = Genista numidica.
sphaerocarpum, Lapeyr. Hist. Abr. Pl. Pyr. 402 = Genista cinerea.
sphaerocarpum, Linn. Mant. ii. 571 = Genista sphaerocarpa.
spinosum, Asso, ex Willk. & Lange, Prod. Fl. Hisp. iii. 430 = Genista Scorpius.
spinosum, Brot. Fl. Lusit. ii. 85 = Calycotome villosa.
spinosum, Linn. Sp. Pl. 709 = Calycotome spinosa.
stylosum, Spreng. Syst. iii. 176 = Genista stylosa.
supranulium, Linn. f. Suppl. 319 = Cytisus fragrans.
terminale, Salisb. Prod. 330 = Cytisus canariensis.
thebaicum, Delile, Fl. Egypt. 250. t. 37. f. 1 = Crotalaria thebaica.
tinctorium, Roth, Tent. Fl. Germ. i. 302 = Genista tinctoria.
tricuspidatum, Cav. in Anal. Cienc. Nat. iv. (1801) 57 = Genista tricuspidata.
tridens, Cav. l. c. 59 = Genista tridens.
triquetrum, Steud. Nom. ed. I. 802 = Genista triquetra, *Lam.*
trispermum, Sm. in Rees, Cycl. xxxii. n. 5 = Genista aethnensis.
umbellatum, L'Hérit. Stirp. Nov. 183 = Genista umbellata.
valentinum, Willd. ex Spreng. Syst. iii. 176 = Genista valentina.
villosum, Poir. Voy. Barb. ii. 207 = Calycotome villosa.
villosum, Sieber, ex Presl, Bot. Bemerk. 51 = Cytisus creticus.
virgatum, Ait. Hort. Kew. ed. I. iii. 11 = Genista virgata.
Webbii, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 291. t. 16. f. 2 = Genista monosperma.

SPARTOCYSUS, Willk. & Lange, Prod. Fl. Hisp. iii. 457 (1880), sphalm. = seq.

SPARTOCYTISUS, Webb & Berth. Phyt. Canar. ii. 49 (1840) = **Cytisus**, Linn. (Legumin.).
albus, Webb & Berth. l. c. 45 = C. albus.
filipes, Webb & Berth. l. c. 52 = C. filipes.
nubigenus, Webb & Berth. l. c. 50 = C. fragrans.
purgans, Webb & Berth. l. c. 45 = C. purgans.
sessilifolius, Webb & Berth. l. c. = C. sessilifolius.
triflorus, Webb & Berth. l. c. = C. triflorus.

SPARTOTAMNUS, Webb & Berth. ex Presl, Bot. Bemerk. 138 (1844) = **Cytisus**, Linn. (Legumin.).
albus, Presl, l. c. = C. albus.
sessilifolius, Presl, l. c. = C. sessilifolius.

SPARTOTHAMNUS, A. Cunn. in Loud. Hort. Brit. Suppl. 600 (1830), nomen; et ex Walp. Rep. vi. 694 (1846–47). *VERBENACEAE*, Benth. & Hook. f. ii. 1141.
junceus, *A. Cunn. ll. cc.*—Austral.
puberulus, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 55, in obs.—Austral.
teucriiflorus, *F. Muell. l. c.* 55.—Austral.

SPARTOTHAMNUS, Walp. Ann. i. 221 (1848), sphalm. = SPARTOTAMNUS, Webb & Berth.

SPARTUM *vel* SPARTIUM, Heist. Syst. 12 (1748) = **Nardus**, Linn. (Gramin.).

SPATALANTHUS, Sweet, Brit. Flow. Gard. Ser. I. t. 300 (1829) = **Romulea**, Maratt. (Irid.).
speciosus, Sweet, l. c. (= *Trichonema monadelphum*).—Afr. austr.

SPATALLA, Salisb. Parad. Lond. sub t. 67 (1807) *PROTEACEAE*, Benth. & Hook. f. iii. 172.
abietina, Roem. & Schult. Syst. iii. 397 = curvifolia.
barbigera, *Knight, Prot.* 76.—Afr. austr.
bombycina, *Knight, l. c.*—Afr. austr.
bracteata, *R. Br. in Trans. Linn. Soc.* x. (1810) 146. Afr. austr.
brevifolia, *R. Br. l. c.* 151.—Afr. austr.
caudaeflora, *Knight, Prot.* 75.—Afr. austr.

SPATALLA:—
caudata, R. Br. in Trans. Linn. Soc. x. (1810) 150 = caudaeflora.
colorata, *Meissn. in DC. Prod.* xiv. 308.—Afr. austr.
curvifolia, *Knight, Prot.* 77.—Afr. austr.
ericaefolia, *Knight, l. c.* 74.—Afr. austr.
gracilis, *Knight, l. c.* 77.—Afr. austr.
incurva, R. Br. in Trans. Linn. Soc. x. (1810) 149 = procera.
laxa, R. Br. l. c. 146 = gracilis.
longifolia, *Knight, Prot.* 77.—Afr. austr.
mollis, *R. Br. in Trans. Linn. Soc.* x. (1810) 144.—Afr. austr.
nana, *Knight, Prot.* 76.—Afr. austr.
nivea, *R. Br. in Trans. Linn. Soc.* x. (1810) 145.—Afr. austr.
parilis, *Knight, Prot.* 75.—Afr. austr.
pedunculata, R. Br. in Trans. Linn. Soc. x. (1810) 144 = curvifolia.
polystachya, *R. Br. l. c.* 148.—Afr. austr.
polystachya, Zeyh. ex Meissn. in DC. Prod. xiv. 308 = sericea.
procera, *Knight, Prot.* 76.—Afr. austr.
prolifera, *Knight, l. c.* 75; *R. Br. in Trans. Linn. Soc.* x. (1810) 147.—Afr. austr.
propinqua, *R. Br. l. c.* 150.—Afr. austr.
pyramidalis, R. Br. l. c. 148 = parilis.
ramulosa, *R. Br. l. c.* 145.—Afr. austr.
ramulosa, Sieber, ex Meissn. in DC. Prod. xiv. 308 = sericea.
sericea, *R. Br. in Trans. Linn. Soc.* x. (1810) 147.—Afr. austr.
squamata, *Meissn. in DC. Prod.* xiv. 310.—Afr. austr.
Thunbergii, *R. Br. in Trans. Linn. Soc.* x. (1810) 150.—Afr. austr.
thyrsiflora, *Knight, Prot.* 74.—Afr. austr.

SPATANTHUS, Juss. in Dict. Sc. Nat. xxiv. 239 (1822) = **Spathanthus**, Desv. (Rapat.).

SPATELLARIA, Reichb. Consp. 189 (1828) = SPATHULARIA, A. St. Hil. = **Amphirrox**, Spreng. (Violar.).

SPATHANDRA, Guill. & Perr. Fl. Seneg. Tent. i. 313. t. 71 (1832) = **Memecylon**, Linn. (Melastom.).
caerulea, Guill. & Perr. l. c. = M. Spathandra.
fasciculatis, Planch. ex Benth. in Hook. Niger Fl. 357 = M. fasciculare.
memecyloides, Benth. l. c. = M. Vogelii.

SPATHANDUS, Steud. Nom. ed. II. ii. 615 (1841), sphalm. = **Spathanthus**, Desv. (Rapat.).

SPATHANTHEUM, Schott, in Bonplandia, vii. (1859) 164. *AROIDEAE*, Benth. & Hook. f. iii. 989.
GAMOCHLAMYS, Baker, in Gard. Chron. (1876) 164.
heterandrum, *Benth. & Hook. f. Gen.* iii. 989; *N. E. Br. in Gard. Chron.* (1883) II. 70.—Hab.?
Orbignyanum, *Schott, in Bonplandia*, vii. (1859) 165.—Bolivia.

SPATHANTHUS, Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 45. t. 4. *RAPATEACEAE*, Benth. & Hook. f. iii. 859.
unilateralis, *Desv. l. c.*—Guian.

SPATHE, P. Br. Hist. Jamaic. 187 (1756) = **Spathelia**, Linn. (Simarub.).
simplex, Crantz, Inst. ii. 464 = Spathelia simplex.

SPATHELIA, Linn. Sp. Pl. ed. II. 386 (1763). *SIMARUBEAE*, Benth. & Hook. f. i. 315.
SPATHE, P. Br. Hist. Jamaic. 187 (1756).
glabrescens, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 581.—Jamaic.
ramosa, Mart. ex Engl. in Mart. Fl. Bras. xii. II. 52 = Discophora guianensis.
rhoifolia, *DC. Prod.* ii. 84.—Mexic.
simplex, *Linn. Sp. Pl.* ed. II. 386.—Jamaic.
vernicosa, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 581.—Cuba.

SPATHESTIGMA, Hook. & Arn. Bot. Beech. Voy. 204 (1836) = **Adenosma**, R. Br. (Scrophul.).
glutinosum, Hook. & Arn. l. c. = A. grandiflorum.

SPATHICARPA, Hook. Bot. Misc. ii. (1831) 146. t. 77. *AROIDEAE*, Benth. & Hook. f. iii. 989.
Bridgesii, Schott, in Bonplandia, vi. (1858) 124, nomen = Spathantheum Orbignyanum.
cornuta, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 34.—Bras.
Gardneri, Schott, in Bonplandia, vi. (1858) 124 = sagittifolia.
hastifolia, *Hook. Bot. Misc.* ii. (1831) 147. t. 77.—Am. trop.
lanceolata, *Engl. in DC. Monog. Phan.* ii. 531.—Paraguay.
longicuspis, Schott, in Bonplandia, x. (1862) 87 = sagittifolia.
platyspatha, Schott, l. c. = sagittifolia.
sagittifolia, *Schott, l. c.* vi. (1858) 124.—Bras.
Tweedieana, *Schott, l. c.*—Bras.

SPATHIOSTEMON, Blume, Bijdr. 621 (1825) = **Homonoia**, Lour. (Compos.).
javense, Blume, l. c. 622 = H. riparia.
salicinum, Hassk. Hort. Bog. Desc. 41 = H. riparia.

SPATHIPHYLLOPSIS, Teysm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 400 = **Spathiphyllum**, Schott (Aroid.).
Minahassae, Teysm. & Binn. l. c. = Spathiphyllum commutatum.

SPATHIPHYLLUM, Schott, Melet. i. 22 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 997.
AMOMOPHYLLUM, Engl. in Gard. Chron. (1877) I. 139.
HYDNOSTACHYON, Liebm. in Kjoeb. Vidensk. Meddel. (1849) 24.
LEUCOCHLAMYS, Poepp. ex Engl. in DC. Monog. Phan. ii. 230 (1879).
MASSOWIA, C. Koch, in Bot. Zeit. x. (1852) 277.
SPATHIPHYLLOPSIS, Teysm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 400.
amazonicum, Spruce, ex Engl. in DC. Monog. Phan. ii. 278 = Urospatha sagittaefolia.
atrovirens, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 179.—Costa Rica.
Beccarii, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 268.—N. Guin.
blandum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 159.—Guian.
Bonplandii, Schott, l. c. 158 = candicans.
brevirostre, *Schott, Aroid.* i. 2.—Mexic.
candicans, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 85. t. 295.—Am. trop.
candidum, *N. E. Br. in Gard. Chron.* (1878) II. 783.—N. Granat.
Candolleanum, *Schott, Prod. Aroid.* 429.—Guian.
cannaefolium, Schott, Aroid. i. 1. t. 1 = candicans.
cannaeforme, Engl. in Mart. Fl. Bras. iii. II. 103. t. 16. f. 3 = candicans.
caudatum, Poepp. & Endl. Nov. Gen. et Sp. iii. 85. t. 296 = Urospatha caudata.
cochlearispathum, *Engl. in DC. Monog. Phan.* ii. 221.—Mexic.
commutatum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 158.—Ins. Philipp.
cuspidatum, *Schott, l. c.*—Guian.
Fendleri, Schott, l. c. 159 = Friedrichsthalii.
floribundum, *N. E. Br. in Gard. Chron.* (1878) II. 783.—N. Granat.
Friedrichsthalii, *Schott, Aroid.* i. 2. t. 4.—Am. centr.
fulvovirens, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 179; *Prod. Aroid.* 427.—Costa Rica.
Gardneri, *Schott, Aroid.* i. 2. t. 3.—Bras.
Glaziovii, *Engl. in Kjoeb. Vidensk. Meddel.* (1880) 358.—Bras.
heliconiaefolium, Schott, Aroid. i. 2. tt. 5, 6 = cochlearispathum.
Humboldtii, Schott, l. c. t. 2 = Candolleanum.
hybridum ×, N. E. Br. in Gard. Chron. (1883) I. 500.
lancaefolium, *Schott, Melet.* i. 22.—Venezuel.
lanceolatum, C. Koch, Berl. Allg. Gartenz. (1857) 174 = Friedrichsthalii.
Lechlerianum, *Schott, Prod. Aroid.* 425.—Peruv.
Liebmanni, Schott, Aroid. i. 2 = cochlearispathum.
longirostre, Schott, l. c. = cochlearispathum.
Minahassae, Regel, Gartenfl. xix. (1870) l. t. 637 = commutatum.
Ortgiesi, *Regel, l. c.* 39; (1872) 292. t. 738.—Mexic.

SPATHIPHYLLUM:—
Patini, *N. E. Br. in Gard. Chron.* (1878) II. 783.—N. Granat.
phryniifolium, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 159.—Panama.
sagittaefolium, Schott, Melet. i. 22 = Urospatha sagittaefolia.
sagittifolium, Spruce, ex Engl. in DC. Monog. Phan. ii. 279 = Urospatha Spruceana.
Schomburgkii, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 158 = Candolleanum.
Wallisi, *Regel, Gartenfl.* (1877) 323. t. 920; *et in Act. Hort. Petrop.* v. (1877) 640.—N. Granat. *vel* Venezuela.
Wallisii, Hort.—Cf. Gard. Chron. (1875). I. 558 = Stenospermation popayanense.
Wendlandii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 179; *Prod. Aroid.* 432.—Costa Rica.

SPATHIUM, Edgew. in Journ. As. Soc. Beng. xi. (1842) 145 = **Aponogeton**, Thunb. (Naiad.).
crispum, Voigt, Hort. Suburb. Calc. 694 = A. crispum.
echinatum, Voigt, l. c. = A. echinatum.
microphyllum, Voigt, l. c. = A. microphyllum.
monostachyum, Edgew. in Journ. As. Soc. Beng. xi. (1842) 148 = A. monostachyon.
undulatum, Edgew. l. c. 145 = A. crispum.

SPATHIUM, Lour. Fl. Cochinch. 217 (1790) = **Saururus**, Linn. (Piperac.).
chinense, Lour. l. c. = Saur. Loureiri.

SPATHODEA, Beauv. Fl. Owar. i. 46. t. 27 (1805). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1045
acuminata, Klotzsch, in Peters, Reise Mossamb. Bot. 191 (Dolichandrones sp.).—Afr. trop.
adenantha, *G. Don, Gen. Syst.* iv. 222.—Afr. trop.
adenophylla, DC. Prod. ix. 206 = Heterophragma adenophyllum.
alternifolia, *R. Br. Prod.* 472.—Austral.
amoena, A. DC. in DC. Prod. ix. 208 = Stereospermum hypostictum.
arcuata, Wight, Ic. t. 1340 = Dolichandrone falcata.
atrovirens, Spreng. Syst. ii. 835 = Dolichandrone falcata.
bracteosa, DC. Prod. ix. 208 = Macfadyena bracteosa.
campanulata, *Beauv. Fl. Owar.* i. 47. tt. 27–28.—Afr. trop.
campanulata, Buch.-Ham. ex DC. Prod. ix. 206 = Dolichandrone stipulata.
Candolleana, *Mart. ex DC. l. c.* 207.—Bras.
Cauda-felina, Hance, in Journ. Bot. x. (1872) 257 = Dolichandrone Cauda-felina.
Coito, DC. Prod. ix. 204 = Macfadyena Coito.
comosa, G. Don, Gen. Syst. iv. 222 = Bignonia comosa.
consanguinea, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1158, nomen.—Guian.
corymbosa, Vent. Choix, 40. t. 40 = Macfadyena corymbosa.
crispa, Bur. Monog. Bign. t. 27 = Dolichandrone Lawii.
crispa, Dalz. & Gibs. Bomb. Fl. 160 = Dolichandrone arcuata.
crispa, G. Don, Gen. Syst. iv. 222 = Dolichandrone falcata.
crispa, Wall. Cat. n. 6515 = Dolichandrone crispa.
Diepenhorstii, Miq. Fl. Ind. Bat. ii. 754 = Dolichandrone Rheedii.
Dolichandra, Steud. Nom. ed. II. ii. 616 = Macfadyena Dolichandra.
falcata, Brand. Forest. Fl. 350 = Dolichandrone Lawii.
falcata, Wall. Cat. n. 6517 = Dolichandrone falcata.
Fenzliana, *Walp. Rep.* vi. 513.—Guian.
filiformis, *DC. Prod.* ix. 209.—Austral.
flaviflora, *Miq. Stirp. Surin. Sel.* 124.—Guian.
fraxinifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 147.—Am. austr.
gigantea, Blume, Bijdr. 761 = Stereospermum hypostictum.
glandulosa, Blume, l. c. 762 = Stereospermum glandulosum.
grandiflora, Zipp. ex Span. in Linnaea, xv. (1841) 326 = Dolichandrone Rheedii.
heterophylla, *R. Br. Prod.* 472.—Austral.

SPATHODEA:—
hispida, DC. Prod. ix. 205 = Macfadyena hispida.
ignea, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 77 = Mayodendron igneum.
ilicifolia, Seem. in Bonplandia, vii. (1859) 246 = Tabebuia dentata.
indica, Pers. Syn. ii. 173 = Oroxylon indicum.
Jenischii, *Sond. in Hamb. Gartenz.* iv. (1848) 370.—Afr. trop.
Kohautiana, Walp. Rep. vi. 513 = Bignonia Unguiscati.
laevis, *Beauv. Fl. Owar.* i. 48. t. 29.—Afr. trop.
laurifolia, H. B. & K. Nov. Gen. et Sp. iii. 146 = Macfadyena laurifolia.
Lobbii, Teysm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 413 = Stereospermum hypostictum.
longiflora, Vent. in Mém. Inst. Par. (1807) 20; Pers. Syn. ii. 173 = Dolichandrone Rheedii.
Loureiriana, DC. Prod. ix. 209 = Dolichandrone Rheedii.
lutea, Benth. in Hook. Niger Fl. 461 = Dolichandrone lutea.
luzonica, Blanco, Fl. Filip. ed. II. 350 = Dolichandrone Rheedii.
macroloba, *Miq. Fl. Ind. Bat. Suppl.* 565.—Sumatra.
magnolioides, Cham. in Linnaea, vii. (1832) 659 = Tabebuia magnolioides.
Mansoana, DC. Prod. ix. 209 = Tabebuia Mansoana.
mollis, Sond. in Linnaea, xxii. (1849) 561 = Macfadyena mollis.
montana, Spreng. Syst. iv. Cur. Post. 236 = Zeyheria montana.
nilotica, Seem. in Journ. Bot. iii. (1865) 333 = campanulata.
obovata, H. B. & K. Nov. Gen. et Sp. iii. 147 = Macfadyena obovata.
obtusifolia, Cham. in Linnaea, vii. (1832) 660 = Tabebuia leucoxyla.
orinocensis, H. B. & K. Nov. Gen. et Sp. iii. 147 = Macfadyena orinocensis.
ovata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1158, nomen.—Guian.
pentandra, Hook. Bot. Mag. t. 3681 = Oroxylon indicum.
Pisoniana, DC. Prod. ix. 209 = Tabebuia Pisoniana.
platypoda, DC. l. c. 204 = Macfadyena platypoda.
podopogon, *DC. l. c.* 205.—Cuba.
portoricensis, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 293.—Porto Rico.
puberula, Klotzsch, in Peters, Reise Mossamb. Bot. 192 (Dolichandrones sp.).—Afr. trop.
Rheedii, Spreng. Syst. ii. 835 = Dolichandrone Rheedii.
rostrata, Span. in Linnaea, xv. (1841) 326 = Dolichandrone Rheedii.
Roxburghii, Spreng. Syst. ii. 835 = Heterophragma Roxburghii.
Schomburgkii, *DC. Prod.* ix. 207.—Guian.
Serratula, *DC. l. c.* 206.—Siam.
serrulata, Teijsm. & Binn. Cat. Hort. Bog. 154 = praec.
speciosa, *Brongn. Herb. Gén. Amat.* t. 4; *ex C. Morr. in Ann. Soc. Bot. Gand*, v. (1849) 213. t. 260.—Bras.
stipulata, Wall. Pl. As. Rar. iii. 20. t. 238 = Dolichandrone stipulata.
suaveolens, Benth. & Hook. f. Gen. ii. 1047, sphalm. = Stereospermum suaveolens.
tenuifolia, Boj. Hort. Maurit. 219; et ex DC. Prod. ix. 208 = *zanzibarica*.
tomentosa, Benth. in Hook. Niger Fl. 462 = Dolichandrone tomentosa.
tulipifera, G. Don, Gen. Syst. iv. 223 = campanulata.
uncata, *Spreng. Syst.* ii. 835.—Ind. occ.
uncinata, Spreng. l. c. = Macfadyena uncinata.
velutina, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 90 = Dolichandrone stipulata.
vernicosa, *Cham. in Linnaea*, vii. (1832) 661.—Bras.
xylocarpa, T. Anders. ex Brand. Forest Fl. 349 = Stereospermum xylocarpum.
zanzibarica, Boj. ex DC. Prod. ix. 208 (= Dolichandrones sp.).—Afr. trop.

SPATHODITHYROS, Hassk. in Flora, xlix. (1866) 211 = **Commelina**, Linn.
suffruticosus, Hassk. Commel. Ind. 11 = C. suffruticosa.

SPATHOGLOTTIS, Blume, Bijdr. 400 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 511.
PAXTONIA, Lindl. Bot. Reg. (1838) t. 60.
affinis, *De Vriese, Illustr. Orch.* t. 15.—Java.
angustifolia, Benth. & Hook. f. Gen. iii. 512 = plicata.
aurea, *Lindl. in Journ. Hort. Soc.* v. (1850) 34.—Malaya.
Deplanchei, *Reichb. f. in Linnaea*, xli. (1877) 86.—N. Caled.
Fortuni, Lindl. Bot. Reg. (1845) t. 19 = pubescens.
Handingiana, *Par. & Reichb. f. Otia Bot. Hamb.* 45.—Burma.
ixioides, *Lindl. in Wall. Cat.* n. 3745.—Reg. Himal.
khasyana, Griff. Notul. iii. 323 = pubescens.
lilacina, Griff. l. c. = plicata.
Lobbii, *Reichb. f. in Walp. Ann.* vi. 455.—Burma.
pacifica, *Reichb. f. in Seem. Fl. Vit.* 300.—Ins. Pacif.
parvifolia, Lindl. Bot. Reg. (1845) sub t. 19 = pubescens.
Paulinae, *F. Muell. Fragm.* vi. 95.—Austral.
Petri, *Reichb. f. in Gard. Chron.* (1877) II. 392.—Ins. Pacif.
plicata, *Blume, Bijdr.* 401. t. 76.—Malaya.
plicata, Griff. Notul. iii. 325 = aurea.
pubescens, *Lindl. in Wall. Cat.* n. 3744.—Reg. Himal.; China.
spicata, Lindl. in Journ. Linn. Soc. iii. (1859) 9 = plicata.
tomentosa, *Lindl. Bot. Reg.* (1845) sub t. 19.—Ins. Philipp.
trivalvis, Lindl. in Wall. Cat. n. 3742 = Acriopsis javanica.
unguiculata, Benth. & Hook. f. Gen. iii. 512 = Calanthes striata.
Vieillardi, *Reichb. f. in Linnaea*, xli. (1877) 85.—N. Caled.

SPATHOLOBUS, Hassk. in Flora, xxv. (1842) II. Beibl. 52. *LEGUMINOSAE*, Benth. & Hook. f. i. 535.
DREBBELIA, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 79.
acuminatus, *Benth. in Miq. Pl. Jungh.* 238.—Burma; Malaya.
africanus, *Baker, in Oliver, Fl. Trop. Afr.* ii. 188.—Afr. trop.
crassifolius, *Benth. in Miq. Pl. Jungh.* 238.—Ind. or.; Malaya.
ferrugineus, *Benth. l. c.*—Java.
gyrocarpus, *Benth. l. c.*—Malaya.
littoralis, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 52.—Java.
macropterus, *Miq. Fl. Ind. Bat. Suppl.* 303.—Sumatra.
purpureus, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 194.—Ind. or.
Roxburghii, *Benth. in Miq. Pl Jungh.* 238.—Ind. or.; Burma.
scandens, Blume, ex Miq. Fl. Ind. Bat. i. 1. 204 = ferrugineus.

SPATHOSCAPHE, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 29 = **Chamaedorea**, Willd. (Palm.).
Arenbergiana, Oerst. l. c. 30 = C. Arenbergiana.

SPATHOTECOMA, Bur. Monog. Bign. 49 (1864) = **Newbouldia**, Seem. (Bignon.).

SPATHULA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 163 = **Iris**, Tourn.
foetidissima, Fourr. l. c. = I. foetidissima.

SPATHULARIA, DC. Prod. iv. 40 (1830), err. typ. = SPATULARIA, Haw. = **Saxifraga**, Linn.

SPATHULARIA, A. St. Hil. in Mém. Mus. Par. xi. (1824) 491. t. 24, et Pl. Rem. Brés. 317. t. 28 (1824) = **Amphirrhox**, Spreng. (Violar.).
longifolia, A. St. Hil. ll. cc. 492. t. 24; 318. t. 28 = A. longifolia.

SPATHYEMA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Symplocarpus**, Salisb. (Aroid.).
angusta, Rafin. Fl. Tellur. iv. 13 = seq. ?
foetida, Rafin. in Med. Repos. N. York, v. (1808) 352 = Sym. foetidus.
lanceolata, Rafin. Fl. Tellur. iv. 13 = praec. ?
latifolia, Rafin. l. c. = praec. ?

SPATULARIA, Haw. Saxifr. Enum. 47 (1821) = **Saxifraga**, Linn.
Clusii, Haw. l. c. 48 = Sax. Clusii.
leucanthemifolia, Haw l. c. = Sax. Clusii.
stellaris, Haw. l. c. 49 = Sax. stellaris.

SPATULIMA, Rafin. Fl. Tellur. ii. 83 (1836) = **Lathyrus**, Tourn. (Legumin.).
angustifolia, Rafin. l. c. (= *Orobus sessilifolius*).—Graecia; As. Min.

SPECKLINIA, Lindl. Gen. et Sp. Orch. 8 (1830) = **Pleurothallis**, R. Br. (Orchid.).
acuminata, Lindl. Bot. Reg: sub t. 1797 = P. acuminata.
atropurpurea, Lindl. l. c. = P. atropurpurea.
ciliaris, Lindl. l. c. (1838) Misc. 31 = P. villosa.
corniculata, Steud. Nom. ed. II. ii. 489, in syn. = P. corniculata.
dichotoma, Poepp. & Endl. Nov. Gen. et Sp. i. 52.—Peruv.
elegans, Lindl. Bot. Reg. sub t. 1797 = P. elegans.
emarginata, Lindl. Gen. et Sp. Orch. 8 (excl. syn.) = P. emarginata.
flexuosa, Poepp. & Endl. Nov. Gen. et Sp. i. 52 = P. flexuosa.
floribunda, Lindl. Gen. et Sp. Orch. 9 = P. floribunda.
graminea, Poepp. & Endl. Nov. Gen. et Sp. i. 51.—Peruv.
hypophylla, Regel, Ind. Sem. Hort. Petrop. (1869) 26.—Am. trop.
Lanceola, Lindl. Gen. et Sp. Orch. 8 = P. Lanceola.
linearis, Lindl. l. c. 9 = P. linearis.
Miqueliana, H. C. Focke, in Tijdschr. Wetensch. ii. (1849) 199 = P. Miqueliana.
obovata, Lindl. Bot. Reg. (1839) Misc. 86 = P. obovata.
orbicularis, Lindl. l. c. (1838) Misc. 31 = P. orbicularis.
plantaginea, Poepp. & Endl. Nov. Gen. et Sp. i. 51. t. 89 = P. plantaginea.
pusilla, Lindl. Bot. Reg. sub t. 1797 = P. pusilla.
retusa, Lindl. l. c. = P. retusa.
scariosa, Lindl. l. c. = P. scariosa.
sertularioides, Lindl. Gen. et Sp. Orch. 8 = P. sertularioides.
violacea, Lem. Jard. Fleur. iii. (1853) Misc. 18.—Bras.

SPECULARIA, Heist. Syst. 8 (1748). *CAMPANULACEAE*, Benth. & Hook. f. ii. 562.
APENULA, Neck. Elem. i. 234 (1790).
CAMPYLOCERA, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 257.
DYSMICODON, Nutt. l. c. (1843).
LEGOUSIA, Durand, Fl. Bourg. i. 37; ii. 26 (1782).
PENTAGONIA, Vent. ex Steud. Nom. ed. II. ii. 298 (1841).
TRIODALLUS, A. DC. in DC. Prod. vii. 491 (1839).
TRIODANIS, Rafin. New Fl. Am. iv. 67 (1836).
SPECULUM, Hall. in Rupp. Fl. Jen. 31 (1745).
SPECULUM-VENERIS, Gerard. ex Meissn. Gen. Comm. 364 (1840?).
arvensis, Montand. Guide Bot. 156 = S. Speculum.
biflora, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 17; *A. Gray, in Proc. Am. Acad.* xi. (1876) 82.—Am. bor. et austr.
californica, Vatke, in Linnaea, xxxviii. (1874) 714 = biflora.
castellana, Lange, Ind. Sem. Hort. Haun. (1854) 25 = falcata.
coa, A. DC. Monog. Camp. 350 = pentagonia.
cordata, Heynh. Nom. ii. 689 = S. Speculum.
falcata, *A. DC. Monog. Camp.* 345.—Reg. Mediterr.
ghilanensis, *A. DC. l. c.* 350.—Persia.
hirsuta, Don, ex Steud. Nom. ed. II. ii. 616 = S. Speculum.
hirta, Nym. Consp. 483 = S. Speculum.
hybrida, *A. DC. Monog. Camp.* 348.—Europ.; Afr. bor.
leptocarpa, *A. Gray, in Proc. Am. Acad.* xi. (1876) 82.—Am. bor. occ.
Lindheimeri, *Vatke, in Linnaea*, xxxviii. (1874) 713.—Am. bor. occ.
Linsecomia, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 460 = leptocarpa.
ovata, Vatke, in Linnaea, xxxviii. (1874) 713 = biflora.
pentagonia, *A. DC. Monog. Camp.* 344.—As. Min.
perfoliata, *A. DC. l. c.* 351.—Am. bor. et austr.

SPECULARIA:—
Speculum, *A. DC. l. c.* 346.—Europ.; Reg. Mediterr.
stricta, Griseb. Spicil. Fl. Rumel. ii. 279 = S. Speculum.
vulgaris, Kitt. Taschenb. ed. II. 483 = hybrida?
vulgaris, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 62, 135 = S. Speculum.

SPECULUM, Hall. in Rupp. Fl. Jen. 31 (1745) = praec.

SPECULUM-VENERIS, Gerard. ex Meissn. Gen. Comm. 364 (1840?) = praec.

SPEIRANTHA, Baker, in Journ. Linn. Soc. xiv. (1875) 562. *LILIACEAE*, Benth. & Hook. f. iii. 771.
convallarioides, *Baker, l. c.* 563.—China.

SPEIRANTHES, Hassk. Cat. Hort. Bog. Alt. 47 (1844), sphalm. = **Spiranthes**, Rich. (Orchid.).

SPEIREMA, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1857) 27 = **Pratia**, Gaudich. (Campan.).
montanum, Hook. f. & Thoms. l. c. = P. montana.

SPEIRODELA, S. Wats. in Bot. Calif. ii. 190 (1880), sphalm. = SPIRODELA, Schleid. = **Lemna**, Linn.

SPENCERIA, Trimen, in Journ. Bot. xvii. (1879) 97. *ROSACEAE*.
ramalana, *Trimen, l. c.*—China.

SPENNERA, Mart. ex DC. Prod. iii. 115 (1828) = **Aciotis**, D. Don (Melastom.).
acuminifolia, Mart. ex DC. l. c. 116 = A. acuminifolia.
acuminifolia, Mart. in Flora, xxiv. (1841) II. Beibl. 37 = A. Martiana.
acutiflora, Mart. l. c. xxii. (1839) I. Beibl. 58 = A. acutiflora.
alata, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856), 125 = A. purpurascens.
annua, Mart. ex DC. Prod. iii. 115 = A. annua.
anomala, Miq. in Linnaea, xviii. (1844) 274 = A. dichotoma.
appendicularioides, Steud. ex Cogn. in Mart. Fl. Bras. xiv. III. 218 = Acisanthera recurva.
aquatica, Mart. ex DC. Prod. iii. 116 = Nepsera aquatica.
asphalti, Crueg. in Linnaea, xx. (1847) 101 = Nepsera aquatica.
brachybotrya, DC. Prod. iii. 115 = A. brachybotrya.
caulialata, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 149 = A. caulialata.
chaetodonta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
choetodon, DC. Prod. iii. 117 = Oxymeris chaetodon.
circaeifolia, DC. l. c. 116 = A. circaeifolia.
circaeoides, Benth. in Hook. Journ. Bot. ii. (1840) 297 = A. purpurascens.
circaeoides, Mart. & Schrank, ex DC. Prod. iii. 116 = A. circaeifolia.
dichotoma, Benth. in Hook. Journ. Bot. ii. (1840) 295 = A. dichotoma.
dysophylla, Benth. l. c. 296 = A. dysophylla.
ferruginea, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 142 = A. brachybotrya.
fragilis, DC. Prod. iii. 115 = A. fragilis.
glandulosa, DC. l. c. 117 (= *Pleroma glandulosum*).—Guian.
grandifolia, Miq. in Linnaea, xviii. (1844) 273 = A. purpurascens.
hydrophila, Miq. Comm. Phytog. 75 = Nepsera aquatica.
hygrophila, Triana, in Trans. Linn. Soc. xxviii. (1871) 36, sphalm. = praec.
indecora, Benth. in Hook. Journ. Bot. ii. (1840) 296 = A. laxa.
indecora, DC. Prod. iii. 116 = A. indecora.
indecora, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 147 = A. polystachya.
Kappleri, Griseb. Fl. Brit. W. Ind. 268 = A. dichotoma.
Kappleriana, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 146 = A. laxa.
lancifolia, Steud. ex Cogn. in Mart. Fl. Bras. xiv. III. 475 = A. fragilis.
latifolia, Benth. in Hook. Journ. Bot. ii. (1840) 296.—Guian.

SPENNERA:—
latifolia, DC. Prod. iii. 115 = Aciotis latifolia.
laxa, DC. l. c. 116 = A. laxa.
longifolia, Mart. ex DC. l. c. 115 = A. longifolia.
martinicensis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 147 = A. discolor.
ornata, Miq. Stirp. Surin. Sel. 56. t. 13 = A. annua.
paludosa, Mart. ex DC. Prod. iii. 115 = A. paludosa.
panicularis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 148. t. 5. f. 1 = A. caulialata.
parviflora, Benth. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 79 = Macrocentrum cristatum.
pellucida, DC. Prod. iii. 115 = A. purpurascens.
pendulifolia, DC. l. c. 116 = A. pendulifolia.
polystachya, DC. l. c. = A. polystachya.
quintuplinervia, Miq. Stirp. Surin. Sel. 57 = Clidemia japurensis.
rostellata, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 143 = A. rostellata.
rubricaulis, Mart. ex DC. Prod. iii. 115 = A. rubricaulis.
rubricaulis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 144 = A. purpurascens.
Sieberi, Steud. in Flora, xxvii. (1844) 720 = Nepsera aquatica.
sileniflora, DC. Prod. iii. 116 = A. sileniflora.
sphaeranthera, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 142 = A. sphaeranthera.
tetraptera, Miq. Comm. Phytog. 76. t. 10. f. E = A. dysophylla.
uliginosa, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 144 = A. uliginosa.
viscida, Benth. in Hook. Journ. Bot. ii. (1840) 296 = A. indecora.
viscida, Steud. in Flora, xxvii. (1844) 720 = A. laxa.
viscosa, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 146 = A. viscosa.

SPENOCARPUS, Wall. Cat. p. 236 (1831?) = **Magnolia**, Linn.

SPENOTOMA, G. Don, Gen. Syst. iii. 774 (1834), sphalm. = SPHENOTOMA, R. Br. = **Dracophyllum**, Labill. (Ericac.).

SPERANSKIA, Baill. Etud. Gén. Euph. 388 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 305.
tuberculata, *Baill. l. c.* 389.—China.

SPERGELLA, Reichb. in Moessl. Handb. ed. II. i. p. lxv (1827) = **Sagina**, Linn. (Caryoph.).
glabra, Reichb. Fl. Germ. Excurs. 794 = Sagina glabra.
glandulifera, Schur, Enum. Pl. Transs. 109 = Sagina nodosa.
macrocarpa, Reichb. Ic. Fl. Germ. v. 26 = Sagina Linnaei.
nodosa, Reichb. Fl. Germ. Excurs. 795 = Sagina nodosa.
nuriensis, Vayr. in Anal. Soc. Esp. Hist. Nat. xi. (1882) 67.—Hispan.
Saginella, Reichb. Ic. Fl. Germ. v. 26.—Helvet.
saginoides, Reichb. Fl. Germ. Excurs. 794 = Sagina Linnaei.
saxatilis, Schur, Enum. Pl. Transs. 109 = Sagina Linnaei.
subulata, Reichb. Fl. Germ. Excurs. 794 = Sagina subulata.

SPERGULA, Linn. Syst. ed. I (1735); Gen. ed. I. 133 (1737). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 152.
ARENARIA, Adans. Fam. ii. 256 (1763).
aestivosa, Koen. ex Wall. Cat. n. 6963 = Bergia aestivosa.
affinis, Hook. Ic. Pl. t. 266 = Colobanthus Billardieri.
apetala, Labill. Nov. Holl. Pl. i. 112. t. 142 = Colobanthus Billardieri.
arenarioides, *Ser. in DC. Prod.* i. 395.—Mexic.
arvensis, *Linn. Sp. Pl.* 440.—Europ.
caespitosa, Pritz. Ic. Ind. 1055, sphalm. = Arenaria caespitosa.
canadensis, D. Dietr. Syn. Pl. ii. 1598 = Spergularia rubra.
Cerviana, D. Dietr. l. c. = Arenaria Cerviana.
Chieusseana, Pomel, Nouv. Mat. Fl. Atl. 206 = arvensis.
ciliaris, *Brot. Fl. Lusit.* ii. 215.—Lusitan.
decandra, Gilib. Fl. Lituan. ii. 156 = arvensis.

SPERGULA:—
decumbens, Ell. Sketch, i. 523 = Sagina decumbens.
droseroides, Brot. Fl. Lusit. ii. 215 = Drosophyllum lusitanicum.
falcata, *Rafin. New Fl. Am.* iv. 16.—Am. bor.
filipensis, D. Dietr. Syn. Pl. ii. 1598 = Spergularia media.
fontinalis, D. Dietr. l. c. 1597 = Stellaria crassifolia.
geniculata, Pers. Syn. i. 522 = arvensis.
glabra, Willd. Sp. Pl. ii. 821 = Sagina glabra.
glandulosa, Bess. Prim. Fl. Galic. i. 298 = Sagina nodosa.
glutinosa, D. Dietr. Syn. Pl. ii. 1598 = Holosteum umbellatum.
grandis, Pers. Syn. i. 522 = Spergularia grandis.
humifusa, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 173 = Sagina humifusa.
imbricata, D. Dietr. Syn. Pl. ii. 1599 = Arenaria imbricata.
indica, Royle, Illustr. Bot. Himal. 464 = pentandra?
japonica, Sw. ex Steud. Nom. ed. I. 802 = Sagina Linnaei.
laevis, D. Dietr. Syn. Pl. ii. 1599 = Spergularia levis.
laricina, Lightf. Fl. Scot. i. 244 = Sagina subulata.
laricina, *Linn. Sp. Pl.* 441.—Sibir.
leptophylla, D. Dietr. Syn. Pl. ii. 1598 = Arenaria leptophylla.
levis, Steud. Nom. ed. II. ii. 617 = Spergularia levis.
linicola, Bor. ex Nym. Consp. 122 = arvensis.
marina, Bartl. & Wendl. f. Beitr. Bot. ii. 64 = Spergularia salina.
marina, Dufour, ex Steud. Nom. ed. II. ii. 617 = Spergularia rubra.
maxima, Dufour, ex Steud. l. c. = Spergularia rubra.
maxima, Weihe, in Arch. des [Deutschen] Apotheker-Ver. viii. (1824) 51; et in Boenn. Prod. Fl. Monast. 136 = arvensis.
media, Bartl. & Wendl. f. Beitr. Bot. ii. 64 = Spergularia media.
micrantha, Bunge, ex Ledeb. Fl. Alt. ii. 183 = Sagina Spergella.
Morisonii, Bor. in Duch. Rev. Bot. ii. (1847) 424 = vernalis.
nivalis, Lindbl. in Phys. Sällsk. Tidskr. (1838) 128; ex Fries, Mant. iii. 31 = Sagina nivalis.
nodosa, Linn. Sp. Pl. 440 = Sagina nodosa.
nodosa, Walt. Fl. Carol. 142 (err. typ. 241) = Sagina decumbens.
pallida, *Salisb. Prod.* 298.—Afr. austr.
palustris, Pers. Syn. i. 522 = Sagina nodosa.
pentandra, *Linn. Sp. Pl.* 440.—Europ.
pentandra, Reichb. Fl. Germ. Excurs. 567 = vernalis.
pilifera, DC. Fl. Fr. iv. 774 = Sagina glabra.
pilifera, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 756 = Sagina subulata.
prostrata, D. Dietr. Syn. Pl. ii. 1598 = Polycarpaea prostrata.
Pseudomorisonii, Iverus, in Bot. Notiser (1875), 81 = arvensis.
pumila, *Rafin. Fl. Ludov.* 93.—Am. bor.
purpurea, D. Dietr. Syn. Pl. ii. 1598 = Spergularia purpurea.
racemosa, D. Dietr. l. c. 1599 = Spergularia media.
radicans, D. Dietr. l. c. 1598 = Spergularia rubra.
refracta, Dethard. ex Reichb. Fl. Germ. Excurs. 567 = arvensis.
repens, Zumag. Fl. Pedem. ii. 286 = Sagina glabra.
rimarum, *J. Gay, in Ann. Sc. Nat.* Sér. II. vi. (1836) 348.—Hispan.
rubra, D. Dietr. Syn. Pl. ii. 1598 = Spergularia rubra.
rupestris, D. Dietr. l. c. = Arenaria bonariensis.
sabuletorum, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 119 = Sagina sabuletorum.
saginoides, All. Fl. Pedem. i. 118. t. 64. f. 1 = Sagina glabra.
saginoides, Linn. Sp. Pl. 441 = Sagina Spergella.
saginoides, Michx. Fl. Bor. Am. i. 276 = Sagina decumbens.
saginoides, Retz. ex Steud Nom. ed. I. 803 = Sagina subulata.
saginoides, Vahl, ex Nym. Consp. 121 = Sagina nivalis.
salina, D. Dietr. Syn. Pl. ii. 1598 = Spergularia rubra.
sativa, Boenn. Prod. Fl. Monast. 135 = arvensis.
saxatilis, Nyland. Animadv. Distrib. Pl. Fenn. 85 = Sagina Linnaei.
saxatilis, Wimm. Fl. Schles. (1832) 193 = Sagina Spergella.
segetalis, Vill. Hist. Pl. Dauph. iii. 657 = Spergularia segetalis.

SPERGULA:—
semidecandra, Turcz. ex Ledeb. Fl. Ross. i. 339 = Sagina Spergella.
serrata, Blanco, Fl. Filip. ed. II. 271 = Bergia capensis.
stricta, Sw. in Vet. Acad. Handl. Stockh. xx. (1799) 229 = Arenaria stricta.
subulata, Benth. Fl. Austral. i. 160, in syn., sphalm. = Colobanthus subulatus.
subulata, Hook. Fl. Bor. Am. i. 93 = Sagina decumbens.
subulata, Sw. in Vet. Acad. Handl. Stockh. (1789) 45. t. 1. f. 3 = Sagina subulata.
succulenta, D. Dietr. Syn. Pl. ii. 1598 = Polycarpon succulentum.
vernalis, *Willd. Fl. Berol. Prod.* 158.—Gallia.
villosa, Pers. Syn. i. 522 = Spergularia villosa.
viscosa, *Lag. in Varied. Cienc.* ii. iv. (1805) 213.—Hispan.
viscoso-pubescens, Loisel. ex D. Dietr. Syn. Pl. ii. 1598 = Spergularia viscoso-pubescens.
vulgaris, Boenn. Prod. Fl. Monast. 135 = arvensis.

SPERGULARIA, J. & C. Presl, Fl. Čech. 94 (1819).
CARYOPHYLLEAE, Benth. & Hook. f. i. 152.
ALSINELLA, Hornem. Nom. 33 (1827).
BALARDIA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 180. t. 111 (1829).
BUDA, Adans. Fam. ii. 507 (1763); Dum. Fl. Belg. 110 (1827).
CORION, Mitch. in Act. Nat. Cur. viii. (1748) App. 208, nomen prius.
DELIA, Dum. Fl. Belg. 110 (1827).
FASCICULUS, Dulac, Fl. Hautes-Pyr. 245 (1867).
HYMENOGONIUM, Rich. ex Lebel, in Mém. Soc. Sc. Cherb. xiv. (1868) 30.
LEPIGONUM, Wahlb. Fl. Gothob. 45 (1820).
MELARGYRA, Rafin. Fl. Tellur. iii. 81 (1836).
STIPULARIA, Haw. Syn. Pl. Succ. 103 (1812).
TISSA, Adans. Fam. ii. 507 (1763).
akkensis, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 55, nomen.—Marocc.
alpina, Willk. in Flora, xxxiv. (1851) 706 = rubra.
amurensis, Pomel, Nouv. Mat. Fl. Atl. 205 = rubra.
anceps, Walp. Ann. i. 83 = rubra.
andina, *Rohrb. in Linnaea*, xxxvii. (1871–73) 234.—Reg. Andin.
arvensis, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 179 = Spergula arvensis.
atheniensis, Heldr. & Sart. ex Nym. Consp. 123 = diandra.
azorica, Lebel, in Mém. Soc. Sc. Nat. Cherb. xiv. (1869) 47; Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 166 = media.
Bourgeaei, *Lebel, l. c.* 42.—Ins. Canar.
brevifolia, Walp. Ann. i. 83 = rubra.
brevinodis, Kunze, ex Kindb. Monog. Lepig. 34 = grandiflora.
campestris, Aschers. Fl. Prov. Brandenb. 94 = rubra.
campestris, Nym. Consp. 123 = diandra.
campestris, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 165.—Hispan.
canadensis, G. Don, Gen. Syst. i. 426 = media.
canariensis, *Lebel, in Mém. Soc. Sc. Nat. Cherb.* xiv. (1869) 46.—Ins. Canar.
capensis, Heynh. Nom. ii. 689 = media.
capillacea, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 163.—Hispan.
Cerviana, G. Don, Gen. Syst. i. 426 = Arenaria Cerviana.
confertiflora, *Steud. in Flora*, xxxix. (1856) 425.—Ins. J. Fernand.
depauperata, *Rohrb. in Linnaea*, xxxvii. (1871–73) 231.—Chili.
diandra, *Boiss. Fl. Orient.* i. 733.—Graecia.
diandra, Coss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 164 = rubra.
diandra, Guir. ex Willk. l. c. 165 = campestris.
Dillenii, Lebel, in Mém. Soc. Sc. Nat. Cherb. xiv. (1869) 43 = salina.
echinosperma, Čelak. in Arch. Naturw. Land. Boehm. iv. (1881) 867 = rubra.
exilis, *Fenzl, in Ledeb. Fl. Ross.* ii. 166.—Sibir.
fallax, *Lowe, in Hook. Kew Journ.* viii. (1856) 289.—Ins. Mader.
fascicularis, Nym. Consp. 122 = media.
fimbriata, *Boiss. & Reut. Diagn.* Ser. II. i. 94.—Ins. Canar.
firma, *Kunze, ex Kindb. Monog. Lepig.* 34.—Chili.
floribunda, *Rohrb. in Linnaea*, xxxvii. (1871–73) 230.—Chili.

SPERGULARIA:—
gamostyla, *Pomel, Nouv. Mat. Fl. Atl.* 202.—Alger.
glandulosa, Heynh. Nom. ii. 689 = Arenaria glandulosa.
glutinosa, G. Don, Gen. Syst. i. 426 = Holosteum liniflorum.
grandiflora, *Poepp. ex Kindb. Monog. Lepig.* 34.—Chili.
grandis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 177.—Bras.
heterosperma, Heldr. ex Nym. Consp. 122 = salina.
heterosperma, *Lebel, in Mém. Soc. Sc. Nat. Cherb.* xiv. (1869) 45.—Sicil.
laevis, Auct. = levis.
laxiflorum, Walp. Ann. i. 83 = rubra.
leiosperma, F. Schmidt, Reisen Amurl. 131 = salina.
leptophylla, *G. Don, Gen. Syst.* i. 425.—Mexic.
levis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 176.—Bras.
Liebmanniana, *Rohrb. in Linnaea*, xxxvii. (1871–73) 241.—Chili.
lignosa, Rohrb. l. c. 244 = Arenaria lignosa.
longicaulis, Pomel, Nouv. Mat. Fl. Atl. 203 = media.
longipes, Nym. Consp. 123 = rubra.
macrocarpa, Presl, Rel. Haenk. ii. 9 = grandis.
macrorrhiza, *Heynh. Nom.* ii. 689; *Gren. & Godr. Fl. Fr.* i. 276.—Ins. Corsic.
macrotheca, *Heynh. l. c.*—Calif.; Chili.
marginata, Bor. Fl. Centr. Fr. ed. III. ii. 106 = media.
marginata, Kitt. Taschenb. ed. II. 1003 = praec.?
marina, Griseb. Spicil. Fl. Rumel. i. 213; Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 165 = media.
maxima, G. Don, in Sweet, Hort. Brit. ed. III. 69 = Spergula arvensis.
media, *Presl, Fl. Sic.* p. xvii, nomen; *et ex Griseb. Spicil. Fl. Rumel.* i. 213.—Europ.
mexicana, *Hemsl. Diagn. Pl. Nov.* 22.—Mexic.
miquelonensis, *Lebel, in Bull. Soc. Bot. Fr.* xv. (1868) 58.—Terra nova (Newfoundl.).
molluginea, *G. Don, Gen. Syst.* i. 425.—Mexic.
Munbyana, *Pomel, Nouv. Mat. Fl. Atl.* 203.—Alger.
neglecta, *Syme, Engl. Bot.* ed. III. ii. 129. t. 255.—Europ.
nicaeënsis, Sarato, ex Lebel, in Mém. Soc. Sc. Nat. Cherb. xiv. (1868) 37 = purpurea.
pallida, *G. Don, Gen. Syst.* i. 425.—Afr. austr.
patens, Hochst. ex Boiss. Fl. Orient. i. 733 = diandra.
pentandra, *G. Don, Gen. Syst.* i. 425.—Europ.
platensis, *Fenzl, in Ann. Wien. Mus.* ii. (1839) 272, in obs.—Bras.
platycaulis, Bartl. ex Rohrb. in Mart. Fl. Bras. xiv. ii. 271 = grandis.
polyphylla, Rohrb. in Linnaea, xxxvii. (1871–73) 232 = rubra.
prostrata, G. Don, Gen. Syst. i. 425 = Polycarpaea prostrata.
purpurea, *G. Don, l. c.*—Hispan.
racemosa, G. Don, l. c. 426 = media.
radicans, Presl, Delic. Prag. 64 = rubra.
ramosa, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 178 = media.
remotiflora, *Steud. in Flora*, xxxix. (1856) 424.—Chili.
rubra, *J. & C. Presl, Fl. Čech.* 94.—Europ.
rupestris, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 176 = villosa.
rupestris, Fenzl, in Enum. Pl. Hueg. 9 = rubra.
rupestris, *Lebel, ex Willk. & Lange, Prod. Fl. Hisp.* iii. 166.—Europ.
rupestris, Lebel, Rech. Pl. Manche, 1848; et in Mém. Soc. Sc. Nat. Cherb. xiv. (1869) 39 = rupicola.
rupicola, *Lebel, in Mém. Soc. Sc. Nat. Cherb.* vii. (1860) 274.—Europ.
salina, *J. & C. Presl, Fl. Čech.* 95.—Europ.
salsuginea, Fenzl, in Ledeb. Fl. Ross. ii. 166 = diandra.
segetalis, *G. Don, Gen. Syst.* i. 425; *Link, Handb.* ii. 259; *Fenzl, in Ledeb. Fl. Ross.* ii. 166.—Europ.
semidecandra, Kitt. Taschenb. ed. II. 1003 = Cerastium semidecandrum?
Spergula, G. Don, Gen. Syst. i. 426 = media.
sperguloides, Heynh. Nom. ii. 690 = diandra.
succulenta, G. Don, Gen. Syst. i. 425 = Polycarpon succulentum.
tenuifolia, Pomel, Nouv. Mat. Fl. Atl. 205 = diandra.
uliginosa, Pomel, l. c. 204 = media.
villosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 178.—Bras.

SPERGULARIA:—
urbica, Nym. Consp. 122 = salina.
viscosa, G. Don, Gen. Syst. i. 425 = Spergula viscosa.
viscoso-pubescens, *Loisel. ex G. Don, l. c.* 426(sp. dub.). —Ins. Corsic.

SPERGULASTRUM, Michx. Fl. Bor. Am. i. 275 (1803) = **Stellaria**, Linn. (Caryoph.).
gramineum, Michx. l. c. 276 = St. longifolia.
lanceolatum, Michx. l. c. 275 = St. borealis.
lanuginosum, Michx. l. c. = Arenaria lanuginosa.

SPERGULUS, Brot. ex Steud. Nom. ed. II. ii. 618 (1841) = **Drosophyllum**, Link (Droser.).
droseroides, Brot. ex Steud. l. c. = D. lusitanica.

SPERLINGIA, Vahl, in Skrivt. Nat. Selsk. Kiobenh. vi. (1810) 112 = **Hoya**, R. Br. (Asclep.).
opposita, Vahl, l. c. 114 = H. Rumphii.
verticillata, Vahl, l. c. 113 = H. verticillata.

SPERMABOLUS, Teijsm. & Binn. Cat. Hort. Bog. 178 (1866). *MAGNOLIACEAE.*
fruticosus, *Teijsm. & Binn. l. c.*—Ins. Molucc.

SPERMACHITON, Llanos, Fragm. 25 (1851) = **Sporobolus**, R. Br. (Gramin.).
involutum, Llanos, l. c. = Eriochloa punctata.

SPERMACOCE, Dill. ex Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth & Hook. f. ii. 145.
Bigelovia, Spreng. Syst. i. 366 (1825).
Borreria, G. F. W. Mey. Prim. Fl. Esseq. 79. t. 1 (1818).
Chenocarpus, Neck. Elem. i. 202 (1790).
Chlorophytum, Pohl, ex DC. Prod. iv. 540 (1830).
Covolia, Neck. Elem. i. 201 (1790).
Diphragmus, Presl, Bot. Bemerk. 80 (1844).
Galiantha, Griseb. ex Lorentz, Veget. Nord. Entre Rios, 72 (1828), nomen; et in Goett. Abh. xxiv. (1879) 156.
Gruhlmania, Neck. Elem. i. 202 (1790).
Hypodematium, A. Rich. Tent. Fl. Abyss. i. 348 (1847).
Jurgensia, Rafin. Sylva Tellur. 146 (1838).
Tardavel, Adans. Fam. ii. 145 (1763).
Tessiera, DC. Prod. iv. 574 (1830).
aculeata, Willd. ex Roem. & Schult. Syst. iii. 531 = tenella.
adscendens, Fisch. ex Link, Enum. Hort. Berol. i. 132 = Mitracarpum Fischeri.
adscendens, Pav. ex DC. Prod. iv. 569 = Richardsonia adscendens.
adscendens, *Willd. ex Roem. & Schult. Syst.* iii. 532.—Madag.
affinis, Pohl, ex DC. Prod. iv. 546 = diffusa.
alata, *Aubl. Pl. Gui.* i. 60.—Guian.
ampliata, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 88.—Afr. trop.
angustifolia, Wall. Cat. n. 835 = stricta.
apiculata, Willd. ex Roem. & Schult. Syst. iii. 531 = tenella.
aquatica, *Vell. Fl. Flum.* 50; i. t. 128.—Bras.
arenosa, Pohl, ex DC. Prod. iv. 564 = Diodia arenosa.
articularis, Linn. f. Suppl. 119 = hispida.
articulata, Pohl, ex DC. Prod. iv. 564 = Diodia hyssopifolia.
aspera, Aubl. Pl. Gui. i. 59 = latifolia.
aspera, *Hemsl. Biol. Centr. Am. Bot.* ii. 58.—Mexic.
aspera, Vahl, Eclog. Am. i. 10 = ocymoides.
asperifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 132.—Mexic.
assurgens, Ruiz & Pav. Fl. Per. i. 60. t. 92 = laevis.
assurgens, Spreng. Syst. i. 402 = tenuior.
aturensis, *H. B. & K. Nov. Gen. et Sp.* iii. 345.—Venezuela.
auriculata, *F. Muell. Fragm.* iv. 42.—Austral.
avana, Wall. Cat. n. 828 = hispida.
barbata, Lam. Illustr. i. 274 = Diodia rigida.
Bartlingiana, Hemsl. Biol. Centr. Am. Bot. ii. 58 = latifolia.
bogotensis, H. B. & K. Nov. Gen. et Sp. iii. 347 = Diodia bogotensis.
brachystema, *R. Br. ex Benth. Fl. Austral.* iii. 439.—Austral.
brasiliensis, Spreng. Syst. i. 402 = tenella?
breviflora, *F. Muell. ex Benth. Fl. Austral.* iii. 442.—Austral.
Brunonis, Wall. Cat. n. 822 = Knoxia brachycarpa.
Burmanni, *DC. Prod.* iv. 555 (Quid?).—Ind. or.
buruensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 261.—Ins. Sunda.
caerulea, *Pohl, ex DC. Prod.* iv. 552.—Bras.
calyptera, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 267 = Gaillonia calycoptera.
capitata, [Moç. & Sesse,] ex DC. Prod. iv. 567 = Crusea calocephala.
capitata, *Ruiz & Pav. Fl. Per.* i. 61.—Peruv.
capitata, Willd. ex H. B. & K. Nov. Gen. et Sp. iii. 346 = tenella.
capitellata, Spreng. Neue Entdeck. iii. 46 = obscura.
capitellata, Willd. ex Roem. & Schult. Syst. iii. 530 = laevis.
Cephalotes, Willd. l. c. 531 = Mitracarpum villosum.
chaetocephala, *DC. Prod.* iv. 554.—Afr. trop.
Chapmanii, Torr. & Gray, Fl. N. Am. ii. 27 = tenuior.
ciliaris, Pav. ex DC. Prod. iv. 564 = Diodia arenosa.
coccinea, Pav. ex DC. l. c. 567 = Crusea coccinea.
coerulea, Auct. = caerulea.
coerulescens, *Aubl. Pl. Gui.* i. 57. t. 19. f. 2.—Guian.
commutata, Schult. Mant. iii. 208 = Diodia maritima.
compacta, *Hochst. in Flora*, xxv. (1842) 1. *Beibl.* 134, nomen; *et ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 239.—Afr. trop.
compressa, *Afzel. ex Hiern, l. c.* 235.—Afr. trop.
compressa, Wall. Cat. n. 6187 = Anotis Wightiana.
conferta, Schott, ex DC. Prod. iv. 563 = Diodia conferta.
cornifolia, Fisch. ex Nees, Hor. Phys. Berol. 50 = capitata.
corymbosa, Burm. f. Fl. Ind. 33 = Burmanni.
corymbosa, Linn. Sp. Pl. ed. II. 149 = *procumbens*.
corymbosa, Roth, Nov. Pl. Sp. 98 = Knoxia Heyneana.
corymbosa, *Ruiz & Pav. Fl. Per.* i. 60.—Peruv.; Bras.; Boliv.
costata, Roxb. Hort. Beng. [10]; Fl. Ind. i. 370 = Hedyotis vestita.
crassifolia, Hort. ex DC. Prod. iv. 573 = Mitracarpum stylosum.
cristata, Willd. ex Roem. & Schult. Syst. iii. 530 = Hedyotis vestita.
cymosa, *Spreng. Syst.* i. 403.—Bras.
debilis, *Benth. Fl. Austral.* iii. 441.—Austral.
decidua, Bosc, ex DC. Prod. iv. 556 = Diodia rigida?
declinata, Pav. ex DC. l. c. 562 = Diodia villosa.
decurrens, *Vell. Fl. Flum.* 50; i. t. 130.—Bras.
denticulata, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 353.—Ins. Philipp.
dibrachiata, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 87.—Afr. trop.
dichotoma, H. B. & K. Nov. Gen. et Sp. iii. 348 = Diodia glabra.
dichotoma, Willd. ex Steud. Nom. ed. II. ii. 618 = tenuior.
diffusa, H. B. & K. Nov. Gen. et Sp. iii. 343 = Mitracarpum villosum.
diffusa, *Pohl, ex DC. Prod.* iv. 544.—Bras.
diodina, Michx. Fl. Bor. Am. i. 82 = Diodia teres.
discolor, E. Mey. in Nov. Act. Nat. Cur. xii. II. (1825) 786 = Mitracarpum Humboldtianum.
distans, *H. B. & K. Nov. Gen. et Sp.* iii. 344.—Mexic.
divergens, Pohl, ex DC. Prod. iv. 568 = Richardsonia grandiflora.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 341.—Mexic.
domingensis, Wright, in Sauv. Fl.Cub. 72 = ocymoides.
echioides, H. B. & K. Nov. Gen. et Sp. iii. 344 = laevis.
elliptica, *Thunb. Pl. Bras. Dec.* iii. 28.—Bras.
elongata, Pohl, ex DC. Prod. iv. 547 = capitata.
ericaefolia, Licht. ex Roem. & Schult. Syst. iii. 281 = Anthospermum Lichtensteinii.
erioclada, *DC. Prod.* iv. 557.—Peruv.
exserta, *Benth. Fl. Austral.* iii. 441.—Austral.
exserta, Roxb. Hort. Beng. [10]; Fl. Ind. i. 368 = Knoxia corymbosa.
fasciculata, Pohl, ex DC. Prod. iv. 547 = tenella.
ferruginea, A. St. Hil. Pl. Us. Bras. t. 13 = capitata.
filifolia, Perr. & Lepr. ex DC. Prod. iv. 540 = Octodon filifolium.
filiformis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 234.—Afr. trop.
filina, Wall. Cat. n. 830 = stricta.
Fischeri, Link, Enum. Hort. Berol. i. 132 = Mitracarpum Fischeri.
flagellaris, Willd. ex Roem. & Schult. Syst. iii. 532 = stricta.
flagelliformis, Poir. Encyc. vii. 314 = stricta.
flavescens, Pohl, ex DC. Prod. iv. 548 = S. Poaya.
flexuosa, *Lour. Fl. Cochinch.* 79.—China.
frigida, Willd. ex Roem. & Schult. Syst. iii. 531 = Mitracarpum Humboldtianum.
fruticosa, Pohl, ex DC. Prod. iv. 541 = verticillata.
Galeopsidis, *DC. l. c.* 554.—Afr. trop.
galioides, Pohl, ex DC. l. c. 573 = Staelia galioides.
Gardneri, Wall. Cat. n. 834 = stricta.
gentianoides, A. St. Hil. Pl. Us. Bras. sub t. 12. p. 3 = S. Poaya.
glabra, *Michx. Fl. Bor. Am.* i. 82.—Am. bor. et austr.
glabra, Roxb. Hort. Beng. 83; Fl. Ind. i. 368 = Hedyotis glabra.
globosa, Pohl, ex DC. Prod. iv. 547 = capitata.
globosa, Schum. & Thonn. Beskr. Guin. Pl. 73 = verticillata.
gracilis, Pohl, ex DC. Prod. iv. 545 = Mitracarpum villosum.
gracilis, Ruiz & Pav. Fl. Per. i. 61. t. 92 = ocymoides.
gracillima, *Pohl, ex DC. Prod.* iv. 559.—Bras.
graminifolia, Hemsl. Biol. Centr. Am. Bot. ii. 58 = verticillata.
grandiflora, Spreng. Neue Entdeck. iii. 45 = Diodia rigida.
Haenkeana, Hemsl. Biol. Centr. Am. Bot. ii. 59 = verticillata.
havanensis, Jacq. Stirp. Am. ed. II; ex J. F. Gmel. Syst. 234 = Machaonia microphylla?
hebecarpa, *DC. Prod.* iv. 553.—Am. trop.
hebecarpa, Oliver, in Trans. Linn. Soc. xxix. (1873) 89 = Ruelliae.
hedyotidea, DC. Prod. iv. 555 = Hedyotis fruticosa.
hexagona, Willd. Sp. Pl. i. 569 = sexangularis.
hexandra, Rich. Hist. Nat. Ipec. 13, 31 = Richardsonia pilosa.
hexangularis, Aubl. Pl. Gui. i. 61 = sexangularis.
hirsuta, Willd. ex Roem. & Schult. Syst. iii. 531 = Richardsonia pilosa.
hirta, Linn. Sp. Pl. ed. II. 148 = Mitracarpum villosum.
hirta, Pav. ex DC. Prod. iv. 567 = Crusea calocephala.
hirta, Rottl. in Ges. Naturf. Fl. Neue Schr. iv. (1803) 95 = hispida.
hirta, Sw. Obs. Bot. 45 = ocymoides.
hirta, Willd. Sp. Pl. i. 571 = laevis.
hispida, *Linn. Sp. Pl.* 102.—Geront. trop.
hispida, Miq. ex Hook. f. Fl. Brit. Ind. iii. 59 = Hedyotis Auricularia.
Holmiana, Schult. ex Steud. Nom. ed. II. ii. 618 = strigosa.
humifusa, *Willd. ex Roem. & Schult. Syst.* iii. 530.—N. Granat.
hyssopifolia, H. B. & K. Nov. Gen. et Sp. iii. 342 = Diodia hyssopifolia.
hyssopifolia, Pers. Syn. i. 123 = verticillata.
hyssopifolia, *Sm. in Abbot, Insects Georgia*, i. 75. t. 38; *et in Rees, Cyclop.* xxxiii. [n. 7].—Guian.; Georgia.
inaperta, *F. Muell. Fragm.* iv. 43.—Austral.
inconspicua, *Bartl. ex DC. Prod.* iv. 557.—Peruv.
insularis, Spreng. Syst. i. 104 = Hedyotis glabra.
involucrata, Pursh, Fl. Am. Sept. i. 105 = Crusea rubra.
Kleinii, Steud. Nom. ed. II. ii. 618 = stricta.
Kotschyana, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 88.—Afr. trop.
laeta, Salisb. Prod. 60 = verticillata.
laevicaulis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 260.—Java.
laevigata, *Hemsl. Biol. Centr. Am. Bot.* ii. 59.—Mexic.
laevigata, *F. Muell. Fragm.* iv. 41.—Austral.
laevis, *Lam. Illustr.* i. 273. t. 94. f. 2.—Am. trop.
laevis, Roxb. Hort. Beng. [10]; Fl. Ind. i. 368 = Knoxia brachycarpa.
laevis, Spreng. Syst. i. 401 = ocymoides.
lanceolata, Frank, ex Presl, Bot. Bemerk. 86 = Hedyotis Frankii.
lanceolata, Link, Enum. Hort. Berol. i. 132 = tenuior.
lanigera, *Pohl, ex DC. Prod.* iv. 574.—Bras.
lasiantha, *Rafin. Fl. Ludov.* 76.—Am. bor.
lasiocarpa, R. Br. in Wall. Cat. n. 832 = stricta.

SPERMACOCE:—
lateralis, Pohl, ex DC. Prod. iv. 568 = Richardsonia grandiflora.
latifolia, *Aubl. Pl. Gui.* i. 55. t. 19.—Guian.
latifolia, Pohl, ex DC. Prod. iv. 549 = S. Poaya.
lavandulaefolia, Pohl, l. c. 548 = tenella.
leptoloba, *Benth. Fl. Austral.* iii. 440.—Austral.
leucadea, Hiern, in Oliver, Fl. Trop. Afr. iii. 237, sphalm. = seq.
leucodea, *Hochst. in Flora*, xxv. (1842) i. *Beibl.* 134, nomen.—Afr. trop.
linearis, H. B. & K. Nov. Gen. et Sp. iii. 343 = Diodia hyssopifolia.
lineata, Blume, Bijdr. 947 = Knoxia lineata.
lineata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 134 = Hedyotis lineata.
lineata, Roxb. Hort. Beng. 83; Fl. Ind. i. 369 = Hedyotis Auricularia.
linifolia, Nees, Hor. Phys. Berol. 49 = Neesiana.
linifolia, Vahl, Eclog. Am. i. 8 = laevis.
linoides, Pohl, ex DC. Prod. iv. 548 = tenella.
lithospermoides, *Bartl. ex DC. l. c.* 574.—Mexic.
longicaulis, R. Br. in Wall. Cat. n. 826 = hispida.
longiflora, D. Dietr. Syn. Pl. i. 475 = longifolia, *Aubl.*
longiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 346.—Mexic.
longifolia, *Aubl. Pl. Gui.* i. 58. t. 21.—Guian.
longifolia, Bartl. ex DC. Prod. iv. 552 = tenuior.
longifolia, *L'Hérit. ex DC. l. c.* 542.—Jamaic.
longiseta, *Hemsl. Biol. Centr. Am. Bot.* ii. 59.—Mexic.
longiseta, Pohl, ex DC. Prod. iv. 546 = tenella.
lutescens, Pohl, ex DC. l. c. 547 = tenella.
macrantha, *Reinw. ex Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 269.—Ins. Molucc.
macrophylla, Hassk. Cat. Hor. Bog. Alt. 107 = stricta.
marginata, *Benth. Fl. Austral.* iii. 440.—Austral.
megapotamica, Spreng. Syst. iv. Cur. Post. 40 = Mitracarpum Sellowianum.
membranacea, *R. Br. ex Benth. Fl. Austral.* iii. 441.—Austral.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 353.—Ins. Philipp.
microcephala, *A. Rich. Fl. Cub. Fanerog.* ii. 29.—Cuba.
microphylla, Pohl, ex DC. Prod. iv. 546 = ocymoides.
minima, Pohl, ex DC. l. c. 560 = verticillata.
mitracarpoides, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 261.—Java.
mucronata, Nees, Hor. Phys. Berol. 49 = verticillata.
multicaulis, *Benth. Fl. Austral.* iii. 440.—Austral.
muriculata, Blanco, Fl. Filip. ed. II. 44 = scaberrima.
muriculata, DC. Prod. iv. 554 = stricta.
mutilata, Blanco, Fl. Filip. ed. II. 43 = hispida.
nana, *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 369.—Ins. 'Honimoa.'
natalensis, *Hochst. in Flora*, xxvii. (1844) 555.—Afr. austr.
Neesiana, *Schult. Mant.* iii. 209.—Bras.
neglecta, Schott, ex DC. Prod. iv. 573 = Mitracarpum Sellowianum.
oaxacana, Hemsl. Biol. Centr. Am. Bot. ii. 59 = verticillata.
obscura, *DC. Prod.* iv. 556.—Ind. occ.
ocymifolia, Willd. ex Roem. & Schult. Syst. iii. 530 = Diodia rigida?
ocymoides, *Burm. f. Fl. Ind.* 34.—Cosmop. trop.
ocymoides, Herb. Madr. ex Wall. Cat. n. 6185 = stricta.
Oldenlandiae, *DC. Prod.* iv. 557.—Chili.
oronocensis, Willd. ex Roem. & Schult. Syst. iii. 531 = tenella.
ovalifolia, Hemsl. Biol. Centr. Am. Bot. ii. 59 = ocymoides.
palmetorum, *DC. Prod.* iv. 553.—Afr. trop.
papuana, *F. Muell. Papuan Pl.* i. 27.—N. Guin.
parviflora, Hemsl. Biol. Centr. Am. Bot. ii. 59; A. Gray, Syn. Fl. N. Am. i. II. 34 = ocymoides.
parviflora, Salisb. Prod. 60 = tenuior.
patula, Hemsl. Biol. Centr. Am. Bot. ii. 59 = ocymoides.
pauciflora, *Vell. Fl. Flum.* 50; i. t. 131.—Bras.
peruviana, Pers. Syn. i. 124 = corymbosa.
philippensis, *Spreng. Syst.* i. 401.—Ins. Philipp.
phyllocephala, *DC. Prod.* iv. 553.—Afr. trop.
Phyteuma, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 235.—Afr. trop.
pilosa, *DC. Prod.* iv. 553.—Afr. trop.
Poaya, *A. St. Hil. Pl. Us. Bras.* t. 12.—Bras.

SPERMACOCE:—
podocephala, Wright, in Sauv. Fl. Cub. 73; Hemsl. Biol. Centr. Am. Bot. ii. 60; A. Gray, Syn. Fl. N. Am. i. II. 34 = verticillata.
pogostoma, *Benth. Fl. Austral.* iii. 440.—Austral.
polycephala, *Bartl. ex DC. Prod.* iv. 558.—Mexic.
polygonifolia, *A. St. Hil. Voy. Distr. Diam.* i. 381.—Bras.
polyphylla, *Vell. Fl. Flum.* 50; i. t. 129.—Bras.
portoricensis, *Balb. ex DC. Prod.* iv. 552.—Ind. occ.
procera, Hemsl. Biol. Centr. Am. Bot. ii. 60 = tenella.
procumbens, Linn. Syst. ed. XII. 115 (Hedyotidis sp.?).—Ind. or.
prostrata, Aubl. Pl. Gui. i. 58. t. 22 = ocymoides.
psyllioides, H. B. & K. Nov. Gen. et Sp. iii. 346 = tenella.
pumila, *Pohl, ex DC. Prod.* iv. 543.—Bras.
purpureo-caerulea, R. Br. ex Benth. Fl. Austral. iii. 442 = laevigata.
pusilla, *Pohl, ex DC. Prod.* iv. 544, in syn.—Bras.
pusilla, Wall. in Roxb. Fl. Ind. ed. Carey, i. 379 = stricta.
pygmaea, Wright, in Sauv. Fl. Cub. 72 = verticillata.
radiata, *Sieber, ex DC. Prod.* iv. 543, in syn.—Afr. trop.
radicans, *Aubl. Pl. Gui.* i. 58. t. 20. f. 4.—Guian.
radicans, Willd. ex Spreng. Syst. i. 405 = Diodia maritima.
Radula, Spreng. Neue Entdeck. ii. 144 = Diodia sarmentosa.
Radula, Willd. ex Roem. & Schult. Syst. iii. 531 = Diodia Radula.
ramisparsa, Pohl, ex DC. Prod. iv. 544, in syn. = ocymoides.
ramosa, Wall. Cat. n. 831 = ocymoides.
ramosissima, R. Br. in Wall. Cat. n. 829 = hispida.
reclinata, Nees, Hor. Phys. Berol. 50 = verticillata.
reflexa, Pohl, ex DC. Prod. iv. 573 = Staelia reflexa.
remota, Lam. Illustr. i. 273 = tenuior.
richardsonioides, *Wright, in Sauv. Fl. Cub.* 73.—Cuba.
rigida, H. B. & K. Nov. Gen. et Sp. iii. 342 = Diodia rigida.
rigida, Salisb. Prod. 60 = hispida.
riparia, Cham. & Schlecht. in Linnaea, iii. (1828) 355 = glabra.
rotata, Poit. ex Roem. & Schult. Syst. iii. 275 = ocymoides.
Roxburghiana, Schult. Mant. iii. 199 = Knoxia brachycarpa.
Roxburghiana, Wall. Cat. n. 6186 = ocymoides.
Roxburghii, Spreng. Syst. i. 404 = Knoxia brachycarpa.
rubra, Jacq. Hort. Schoenb. iii. 3 = Crusea rubra.
rubra, *Spreng. in Schrad. Journ.* (1800) II. 199.—Hab.?
rubricaulis, *Wright, in Sauv. Fl. Cub.* 71.—Cuba.
rubrostipulata, Pohl, ex DC. Prod. iv. 547 = tenella.
Ruelliae, *DC. l. c.* 554.—Afr. trop.
Ruelliae, T. Thoms. in Speke, Journ. App. 636 = ampliata.
scaberrima, Bartl. ex DC. Prod. iv. 557 = erioclada.
scaberrima, *Blume, Bijdr.* 946.—Malaya.
scabra, Willd. Sp. Pl. i. 572 = hispida.
scabrida, Pohl, ex DC. Prod. iv. 548 = latifolia.
scandens, *Aubl. Pl. Gui.* i. 61.—Guian.
semierecta, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 371.—Sumatra.
senensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 236.—Afr. trop.
serpyllifolia, *Willd. ex Roem. & Schult. Syst.* iii. 532.—Madag.
serrulata, Beauv. Fl. Owar. i. 39. t. 23 = Diodia sarmentosa.
setosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 60.—Mexic.
setosa, Willd. ex DC. Prod. iv. 564 = Diodia rigida.
sexangularis, *Aubl. Pl. Gui.* i. 61. t. 22. f. 8.—Guian.
sexangularis, Sieber, ex Steud. Nom. ed. II. ii. 619 = laevis.
simplex, *Wright, in Sauv. Fl. Cub.* 73.—Jamaic.
sparsa, Pohl, ex DC. Prod. iv. 568 = Richardsonia grandiflora.
sphaerostigma, Oliver, in Trans. Linn. Soc. xxix. (1873) 88, in nota = ampliata.
spinosa, *Jacq. Enum. Pl. Carib.* 12; *Select. Am.* 21.—Am. trop.
spinosa, Loefl. ex Griseb. Fl. Brit. W. Ind. 349 = Diodia sarmentosa.
spinulosa, Lam. Illustr. i. 275 = spinosa.

SPERMACOCE:—
squarrosa, Poepp. ex Cham. & Schlecht. in Linnaea, iii. (1828) 363 = Mitracarpum squarrosum.
stachydea, *DC. Prod.* iv. 554.—Afr. trop.
stellata, Willd. ex Roem. & Schult. Syst. iii. 530 = verticillata.
stenophylla, *F. Muell. Fragm.* iv. 43.—Austral.
stricta, *Linn. f. Suppl.* 120; *Roxb. Hort. Beng.* 83; *Fl. Ind.* i. 370.—Geront. trop.
stricta, F. Muell. Fragm. iv. 41 = brachystema.
stricta, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 200 = ocymoides.
strictissima, Pohl, ex DC. Prod. iv. 547 = tenella.
strigosa, Hill, Veg. Syst. vii. 35 (Quid?).—Ital.; Hispan.
strigosa, Linn. Syst. ed. X. 890 = Diodia prostrata.
strigosa, Sims, Bot. Mag. t. 1558 = Crusea rubra.
strigosa, *Thunb. Pl. Bras. Dec.* iii. 29.—Bras.
strumpfioides, *Wright, in Sauv. Fl. Cub.* 73.—Cuba.
stylosa, Link, Enum. Hort. Berol. i. 132 = Mitracarpum stylosum.
subulata, Pav. ex DC. Prod. iv. 543 = Crusea subulata.
suffrutescens, Jacq. Hort. Schoenb. iii. 40. t. 322 = laevis.
suffruticosa, *R. Br. ex Benth. Fl. Austral.* iii. 443.—Austral.
suffruticosa, Hill, Veg. Syst. vii. 35 (Quid?).—Europ. austr.
suffruticosa, *Loefl. It. Hisp.* 201.—Am. trop.
suffruticosa, Spreng. Syst. i. 400 = laevis.
sumatrensis, Retz. Obs. iv. 23 = Knoxia corymbosa.
sumatrensis, Roxb. Hort. Beng. 83; Fl. Ind. i. 366 = Knoxia sumatrensis.
tenella, *H. B. & K. Nov. Gen. et Sp.* iii. 345.—Bras.
tenera, R. Br. in Wall. Cat. n. 833 = ocymoides.
tenera, *Pohl, ex DC. Prod.* iv. 543, in syn.—Bras.
tenuior, *Linn. Sp. Pl.* 102 (*excl. syn. Dill.*).—Am. trop.
tenuior, Linn. l. c., partim = glabra.
tenuior, Pav. ex DC. Prod. iv. 547 = tenella.
tenuis, *Pohl, ex DC. Prod.* iv. 543, in syn.—Bras.
tenuissima, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 234.—Afr. trop.
teres, Roxb. Hort. Beng. [10]; Fl. Ind. i. 367 = Knoxia corymbosa.
terminalis, Vell. Fl. Flum. 51; i. t. 132 = tenella.
tetracocca, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 132 = Crusea allococca.
tetragona, Hort. Berol. ex DC. Prod. iv. 553 = laevis.
tetraquetra, *A. Rich. Fl. Cub. Fanerog.* ii. 29.—Cuba.
thymocephala, Wright, in Sauv. Fl. Cub. 72 = verticillata.
tomentosa, Vell. Fl. Flum. 51; i. t. 133 = tenella.
triandra, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 134 = stricta.
tubularis, R. Br. in Wall. Cat. n. 836 = hispida.
vaginata, Willd. ex Roem. & Schult. Syst. iii. 531 = laevis.
verticillata, *Linn. Sp. Pl.* 102.—Afr. et Am. trop.
verticillata, Vell. Fl. Flum. i. t. 127 = tenella.
villosa, Sw. Prod. Veg. Ind. Occ. 29 = Mitracarpum villosum.
virgata, *Hemsl. Biol. Centr. Am. Bot.* ii. 61.—Mexic.
virgata, Link, ex Roem. & Schult. Syst. iii. 281, 531 = Mitracarpum villosum.
virginiana, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) t. 14. f. 3 = Diodia virginiana.

SPERMADICTYON, Roxb. Pl. Corom. iii. 32. t. 236 (1819) = **Hamiltonia**, Roxb. (Rubiac.).
azureum, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 225 = H. suaveolens.
mysorense, Steud. Nom. ed. II. ii. 619 = H. suaveolens.
pilosum, Spreng. Syst. iv. Cur. Post. 73 = H. pilosa.
scabrum, Spreng. l. c. = H. suaveolens.
suaveolens, Roxb. Pl. Corom. iii. 32. t. 236 = H. suaveolens.

SPERMATURA, Reichb. Consp. 141 (1828) = **Osmorhiza**, Rafin. (Umbell.).

SPERMAULAXEN, Rafin. Fl. Tellur. iii. 13 (1836); New Fl. Am. iii. 41 (1836) = **Polygonum**, Tourn.
dichotomus, Rafin. ll. cc. 14; 42 (= *P. dichotomus*).—Am. bor.

SPERMAXYRON, Steud. Nom. ed. II. ii. 619 (1841) = seq.

SPERMAXYRUM, Labill. Nov. Holl. Pl. ii. 84. t. 233 (1806) = **Olax**, Linn.
aphyllum, DC. Prod. i. 532 = O. aphylla.
Phyllanthi, Labill. Nov. Holl. Pl. ii. 84. t. 233 = O. Phyllanthi.
strictum, DC. Prod. i. 532 = O. stricta.

SPERMODON, Beauv. ex Lestib. Essai Fam. Cyp. 27 (1819) = **Rynchospora**, Vahl (Cyperac.).
confinis, Nees, in Mart. Fl. Bras. ii. 1. 119.—Bras.
curvulus, Nees, l. c. = R. tenuis.
edentulatus, Nees, in Linnaea, ix. (1834) 296 = R. junciformis.
eximius, Nees, in Seem. Bot. Voy. Herald, 222 = R. eximia.
filiformis, Nees, in Mart. Fl. Bras. ii. 1. 118 = R. longispicata.
setaceus, Nees, in Linnaea, ix. (1834) 296 = R. tenerrima.

SPERMOLEPIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 577; et in Ann. Sc. Nat. Sér. V. ii. (1864) 136. *MYRTACEAE*, Benth. & Hook. f. i. 710.
gummifera, *Brongn. & Gris, ll. cc.* 578; 137.—N. Caled.
rubiginosa, *Brongn. & Gris, ll. cc.*—N. Caled.

SPERMOLEPIS, Rafin. Neogenyt. 2 (1825) = **Apium**, Tourn. (Umbell.).
divaricata, Rafin. l. c. = A divaricatum.

SPERMOPHYLLA, Neck. Elem. i. 24 (1790) = **Ursinia**, Gaertn. (Compos.).

SPHACELE, Benth. in Bot. Reg. sub t. 1289 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1193.
PHYTOXIS, Molina, Sagg. Chile, ed. II. 145, 290 (1810).
acuminatum, *Griseb. in Goett. Abh.* xix. (1874) 238.—Reg. Argent.
alpina, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 36.—Costa Rica.
bullata, *Benth. Lab. Gen. et Sp.* 568.—N. Granat.
calycina, *Benth. l. c.* 568.—Calif.
campanulata, *Benth. in Bot. Reg.* sub t. 1289.—Chili.
clinopodioides, *Griseb. in Goett. Abh.* xxiv. (1879) 273.—Reg. Argent.
coerulea, Hort.—Cf. Gard. Chron. (1865) 934 (Quid ?).—Am. trop.
conferta, *Benth. Pl. Hartw.* 244.—N. Granat.
cordifolia, *Benth. in DC. Prod.* xii. 257.—Peruv.
floribunda, *Benth. l. c.* 254.—Hab. ?
hastata, *A. Gray, in Proc. Am. Acad.* v. (1862) 341.—Ins. Sandvic.
hastata, *Griseb. in Goett. Abh.* xix. (1874) 238.—Reg. Argent.
lamiifolia, *Benth. Lab. Gen. et Sp.* 570.—Peruv.
Lindleyi, *Benth. in Bot. Reg.* sub t. 1289 (Lindlei).—Chili.
mexicana, *S. Schau. in Linnaea*, xx. (1847) 707.—Mexic.
mutica, *Benth. Pl. Hartw.* 145.—Peruv.
paniculata, *Benth. Lab. Gen. et Sp.* 568.—Ecuador.
parviflora, *Benth. in DC. Prod.* xii. 256.—N. Granat.
Radula, *Benth. l. c.* 257.—Peruv.
salviaefolia, *Benth. Lab. Gen. et Sp.* 567.—N. Granat.
speciosa, *A. St. Hil. ex Benth. l. c.* 570.—Bras.
subhastata, *Benth. in Bot. Reg.* sub t. 1289.—Chili.
tenuiflora, *Benth. in DC. Prod.* xii. 257.—N. Granat.
tomentosa, *Benth. Lab. Gen. et Sp.* 569.—Peruv.

SPHACOPHYLLUM, Benth. in Hook. Ic. Pl. t. 1135 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 339.
Bojeri, *Benth. in Hook. Ic. Pl.* t. 1135.—Madag.
Kirkii, *Oliver, l. c.* t. 1451.—Afr. trop.

SPHAENODESMA, Schau. in DC. Prod. xi. 622 (1847), sphalm. = **Sphenodesme**, Jack (Verben.).

SPHAERALCEA, A. St. Hil. Fl. Bras. Mer. i. 209 (1825). *MALVACEAE*, Benth. & Hook. f. i. 204.
ANISODONTEA, Presl, Bot. Bemerk. 18 (1844).
MALOCOPSIS, Walp. Ann. i. 104 (1848).
MALVEOPSIS, Presl, Bot. Bemerk. 19 (1844).
MELIPHLEA, Zucc. in Abh. Akad. Muench. ii. (1847) 359.

SPHAERALCEA :—
PHYMOSIA, Desv. in Ham. Prod. Pl. Ind. Occ. 49 (1825).
SPHAEROMA, Schlecht. in Linnaea, xi. (1837) 352.
abutiloides, *G. Don, Gen. Syst.* i. 465.—Ins. Baham.; Carolina.
acerifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 228.—Am. bor.
angustifolia, *G. Don, Gen. Syst.* i. 465.—Mexic.
Berteroana, Phil. in Linnaea, xxxiii. (1864–65) 23 = obtusiloba.
bonariensis, *Griseb. in Goett. Abh.* xix. (1874) 92.—Reg. Argent.
chilensis, *C. Gay, Fl. Chil.* i. 293.—Chili.
choapina, Phil. in Linnaea, xxxiii. (1864–65) 26 = velutina.
cisplatina, *A. St. Hil. Fl. Bras. Mer.* i. 210.—Am. trop.
collina, Phil. in Linnaea, xxxiii. (1864–65) 24 = obtusiloba.
coquimbana, Phil. in Anal. Univ. Chil. (1872) 680 = obtusiloba.
Dregeana, *Harv. in Harv. & Sond. Fl. Cap.* i. 165.—Afr. austr.
elegans, *G. Don, Gen. Syst.* i. 465.—Afr. austr.
Emoryi, *Torr. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 23.—Am. bor. occ.
Fendleri, *A. Gray, Pl. Wright.* i. 21.—Am. bor. occ.
filicaulis, Hemsl. Biol. Centr. Am. Bot. i. 113 = Sida diffusa.
floribunda, Phil. in Linnaea, xxxiii. (1864–65) 25 = obtusiloba.
floribunda, *Walp. Rep.* ii. 789.—Mexic.
Galeottii, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 186 = umbellata.
Gayana, Phil. in Linnaea, xxxiii. (1864–65) 23 = obtusiloba.
glabrata, Phil. l. c. 22 = chilensis.
hastulata, *A. Gray, Pl. Wright.* i. 17.—N. Mexic.
incana, *Torr. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 23.—Am. bor. occ.
lactea, Spach, Hist. Vég. Phan. iii. 354 = Malvastrum vitifolium.
Lindheimeri, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 162.—Texas; Mexic.
longisepala, *Torr. in Bot. U. St. Expl. Exped.* ii. 255.—Am. bor. occ.
mendocina, *Phil. in Anal. Univ. Chil.* (1862) II. 392; *et in Linnaea*, xxxiii. (1864–65) 24.—Chili.
miniata, A. Gray, Gen. Illustr. ii. 70. t. 127 = Fendleri.
miniata, *Spach, Hist. Vég. Phan.* iii. 352 = cisplatina.
Munroana, *Spach, l. c.* 353.—Am. bor. occ.
nutans, *Scheidw. ex Planch. in Fl. des Serres*, vii. (1851–52) 221.—Guatemala ?
obtusiloba, *G. Don, Gen. Syst.* i. 465.—Chili.
pedata, *Torr. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 23.—Am. bor. occ.
rhombifolia, Griseb. in Goett. Abh. xix. (1874) 92 = cisplatina.
rivularis, Torr. ex A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 23 = acerifolia.
rosea, G. Don, Gen. Syst. i. 465 = umbellata.
rupestris, Phil. in Linnaea, xxxiii. (1864–65) 26 = obtusiloba.
sessiliflora, *Phil. in Anal. Univ. Chil.* (1872) 679.—Chili.
stellata, Torr. & Gray, Fl. N. Am. i. 228 = angustifolia.
sulphurea, *S. Wats. in Proc. Am. Acad.* xi. (1876) 113, 125.—Calif.
triflora, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 135.—Java.
umbellata, *G. Don, Gen. Syst.* i. 465.—Mexic.
valparidisea, Phil. in Linnaea, xxxiii. (1864–65) 23, in syn. = obtusiloba.
velutina, *Presl, Rel. Haenk.* ii. 124.—Chili.
vitifolia, Benth. & Hook. ex Hemsl. Biol. Centr. Am. Bot. i. 114 = umbellata.
Wrightii, *A. Gray, Pl. Wright.* ii. 21.—Mexic.

SPHAERANTHOIDES, A. Cunn. ex DC. Prod. v. 456 (1836) = **Pterocaulon**, Ell. (Compos.).

SPHAERANTHUS, Vaill. ex Linn. Coroll. Gen. 18 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 294.
OLIGOLEPIS, Cass. ex DC. Prod. v. 370, in syn. (1836).
POLYCEPHALOS, Forsk. Fl. Aegypt. Arab. 154 (1775).
SPRUNERA, Sch. Bip. in Flora, xxv. (1842) 1. Beibl. 134, 145.

SPHAERANTHUS :—
abyssinicus, Steetz, in Peters, Reise Mossamb. Bot. 411 = suaveolens.
africanus, Burm. f. Fl. Ind. 185. t. 60. f. 2 = Dichrocephala latifolia.
africanus, *Linn. Sp. Pl.* ed. II. 1314.—Geront. trop.
africanus, Wall. Cat. n. 3179, partim = indicus.
alatus, Blanco, Fl. Filip. ed. I. 635 = indicus.
amaranthoides, *Burm. f. Fl. Ind.* 186.—Ind. or.
angustifolius, *DC. Prod.* v. 370.—Afr. trop.
angustifolius, Sch. Bip. ex Oliver & Hiern, in Fl. Trop. Afr. iii. 333, 334 = suaveolens, Steetzii.
chinensis, Linn. Mant. i. 119 = Athroisma laciniatum.
cochinchinensis, Lour. Fl. Cochinch. 510 = africanus.
cotuloides, *DC. Prod.* v. 370.—Madag.
crispus, *Zipp. ex Span. in Linnaea*, xv. (1841) 322, nomen.—Ins. Timor. ?
elongatus, Blanco, Fl. Filip. ed. I. 636 = Pterocaulon cylindrostachyum.
erectus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 233.—Java.
glaber, DC. Prod. v. 370 = africanus.
globosus, Wall. ex DC. l. c. 369 = africanus.
gracilis, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 400.—Afr. trop.
hirtus, Willd. Sp. Pl. iii. 2395 = indicus.
indicus, Gaertn. Fruct. ii. 413. t. 164. f. 5 = suaveolens.
indicus, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 183 = africanus.
indicus, *Linn. Sp. Pl.* 927.—Geront. trop.
integrifolius, Linn. f. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 30 = Myriactis nepalensis.
Kirkii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 333.—Afr. trop.
Kotschyi, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 308 = suaveolens.
laciniatus, Wall. Cat. n. 3184 = Athroisma laciniatum.
laevigatus, Wall. ex DC. Prod. v. 369 = africanus.
microcephalus, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 326 = sphenocleoides.
microcephalus, Willd. Sp. Pl. iii. 2395 = africanus.
mollis, Roxb. Hort. Beng. 62; Fl. Ind. iii. 446 = indicus.
mossambiquensis, Steetz, in Peters, Reise Mossamb. Bot. 421 = indicus.
nubicus, *Sch. Bip. ex Schnizl. in Flora*, xxv. (1842) 1. *Beibl.* 134, 145.—Afr. trop.
ovalis, *Steetz, in Peters, Reise Mossamb. Bot.* 413.—Ind. or.
paniculatus, Cass. in Dict. Sc. Nat. l. 211 = africanus.
peduncularis, *DC. Prod.* v. 370.—Afr. austr.
peguensis, *Kurz, ex C. B. Clarke, Comp. Ind.* 97.—Burma.
polycephalus, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 95. t. 59.—Afr. trop.
senegalensis, DC. Prod. v. 370 = africanus.
sinensis, J. F. Gmel. Syst. 1280 = *chinensis*.
sphenocleoides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 333.—Afr. trop.
Steetzii, *Oliver & Hiern, l. c.* 334.—Afr. trop.
strobuliferus, *Boiss. & Noë, Diagn.* Ser. II. iii. 6.—Mesopot.
suaveolens, *DC. Prod.* v. 370.—Afr. bor.
zeylanicus, Heyne, ex Wall. Cat. n. 3180 = amaranthoides, peguensis.

SPHAERIDIOPHORA, Benth. & Hook. f. Gen. i. 494 (1865), sphalm. = seq.

SPHAERIDIOPHORUM, Desv. Journ. Bot. i. (1813) 125. t. 6 = **Indigofera**, Linn. (Legumin.).
abyssinicum, Jaub. & Spach, Illustr. v. 102 = I. linifolia.
linifolium, Desv. Journ. Bot. i. (1813) 125. t. 6 = I. linifolia.

SPHAERINE, Herb. Amaryll. 106. tt. 12, 13, 16, 18 (1837) = **Bomarea**, Mirb. (Amaryll.).
angustipetala, Benth. Pl. Hartw. 260 = B. angustipetala.
brevis, Herb. Amaryll. 108. t. 18. f. 1 = B. brevis.
coccinea, Herb. l. c. 106. t. 16. f. 1 = B. coccinea.
distichophylla, Herb. l. c. 107. t. 12. f. 3, 4 = B. distichophylla.
linifolia, Kunth, Enum. Pl. v. 786 = B. linifolia.
nervosa, Herb. Amaryll. 108. t. 13. f. 1 = B. nervosa.
secundifolia, Herb. l. c. 107. t. 12. f. 1, 2 = B. secundifolia.

SPHAERITIS, Eckl. & Zeyh. Enum. 299 (1837) = **Crassula**, Dill.
biconvexa, Eckl. & Zeyh. l. c. 300 = C. biconvexa.
incana, Eckl. & Zeyh. l. c. = C. incana.
margaritifera, Eckl. & Zeyh. l. c. 299 = C. margaritifera.
muricata, Eckl. & Zeyh. l. c. = C. cymosa.
paucifolia, Eckl. & Zeyh. l. c. 300 = C. trachysantha.
puberula, Eckl. & Zeyh. l. c. = C. subaphylla.
setigera, Eckl. & Zeyh. l. c. 301 = C. tomentosa.
stenophylla, Eckl. & Zeyh. l. c. 299 = C. cymosa.
subaphylla, Eckl. & Zeyh. l. c. 300 = C. subaphylla.
tomentosa, Eckl. & Zeyh. l. c. = C. tomentosa.
trachysantha, Eckl. & Zeyl. l. c. = C. trachysantha.
typica, Eckl. & Zeyh. l. c. 299 = C. cymosa.

SPHAERIUM, Linn. Syst. ed. I (1735) = **Coix**, Linn. (Gramin.).

SPHAEROCARDAMUM, S. Schau. in Linnaea, xx. (1847) 720. *CRUCIFERAE*, Benth. & Hook. f. i. 83.
nesliaeforme, *S. Schau. l. c.*—Mexic.

SPHAEROCARPOS, J. F. Gmel. Syst. 9 (1791) = **Globba**, Linn. (Scitam.).
Hura, J. F. Gmel. l. c. = G. versicolor.

SPHAEROCARPUS, Steud. Nom. ed. II. ii. 620 (1841), sphalm. = SPHENOCARPUS, Rich. = **Laguncularia**, Gaertn. (Combret.).

SPHAEROCARYA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 371 (1824) = **Pyrularia**, Michx. (Santal.).
edulis, Wall. l. c. = P. edulis.
leprosa, Dalz. in Hook. Kew Journ. iii. (1851) 34 = Strombosia ceylanica.
moschifera, Blume, Mus. Bot. Lugd. Bat. i. 245 = P. moschifera.
vestita, Wall. Cat. n. 7207 = P. edulis.
Wallichiana, Wight & Arn. in Edinb. N. Phil. Journ. xv. (1833) 180 = Scleropyrum Wallichianum.

SPHAEROCARYUM, Nees, ex Steud. Nom. ed. II. ii. 620 (1841) = **Isachne**, R. Br. (Gramin.).
elegans, Nees, ex Steud. l. c. = I. pulchella.

SPHAEROCEPHALA, Hill, Veg. Syst. iv. 48 (1762) = **Centaurea**, Linn. (Compos.).

SPHAEROCEPHALUS, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 67 = **Nassauvia**, Juss. (Compos.).
Lagascae, D. Don, in Phil. Mag. (1832) 389 = N. Lagascae.

SPHAEROCEPHALUS, Linn. Syst. ed. I (1735) = **Echinops**, Linn. (Compos.).

SPHAEROCHLOA, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 47. t. 5 = **Eriocaulon**, Linn.
compressa, Beauv. ex Desv. l. c. = E. compressum.
quinquangularis, Beauv. ex Desv. l. c. = E. quinquangulare.

SPHAEROCLINIUM, Sch. Bip. Tanacet. 20 (1844) = **Matricaria**, Tourn. (Compos.).
nigellaefolium, Sch. Bip. l. c. = M. nigellaefolia.

SPHAEROCODON, Benth. in Benth. & Hook. f. Gen. ii. 772 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
natalense, *Benth. in Hook. Ic. Pl.* sub t. 1190.—Afr. austr.
obtusifolium, *Benth. l. c.* t. 1190.—Afr. austr.

SPHAEROCOMA, T. Anders. in Journ. Linn. Soc. v. (1861) 16. t. 15. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 154.
Aucheri, *Boiss. Fl. Orient.* i. 739.—Persia.
Hookeri, *T. Anders. in Journ. Linn. Soc.* v. (1861) 16.—Arab.

SPHAERODENDRON, Seem. in Journ. Bot. iii. (1865) 33. t. 26 = **Cussonia**, Thunb. (Araliac.).
angolense, Seem. l. c. 34. t. 26 = C. angolensis.

SPHAEROGYNE, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 331 = **Tococa**, Aubl. (Melastom.).
cinnamomea, Linden, in Belg. Hortic. xv. (1865) 101 (Quid ?).—Costa Rica.
imperialis, Hort.—Cf. Gard. Chron. (1869) 587; (1872) 7 = seq. ?
imperialis, Linden, in Illustr. Hortic. xxiv. (1877) 109. t. 284 (Quid ?).—Peruv.
latifolia, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 331 = T. platyphylla.

SPHAEROLOBIUM, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 509. *LEGUMINOSAE*, Benth. & Hook. f. i. 469.
HUGELROEA, Steud. Nom. ed. II. i. 778 (1840).
ROEA, Hueg. Enum. Pl. Hueg. 34 (1837).
acuminatum, Benth. in Enum. Pl. Hueg. 32 = medium.
alatum, *Benth. l. c.*—Austral.
crassirameum, Meissn. in Bot. Zeit. xiii. (1855) 28 = macranthum.
daviesioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) I. 266.—Austral.
Drummondii, Turcz. l. c. 267 = macranthum.
euchilus, *Benth. Fl. Austral.* ii. 67.—Austral.
foliosum, F. Muell. Fragm. i. 166 = nudiflorum.
fornicatum, *Benth. in Enum. Pl. Hueg.* 32.—Austral.
gracile, *Benth. Fl. Austral.* ii. 64.—Austral.
grandiflorum, *Benth. in Enum. Pl. Hueg.* 32.—Austral.
linophyllum, *Benth. Fl. Austral.* ii. 64.—Austral.
macranthum, *Meissn. in Lehm. Pl. Preiss.* ii. 213.—Austral.
medium, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 14.—Austral.
medium, Meissn. in Lehm. Pl. Preiss. i. 58 = fornicatum.
minus, Labill. Nov. Holl. Pl. i. 108. t. 138 = vimineum.
nudiflorum, *Benth. Fl. Austral.* ii. 64.—Austral.
pulchellum, Meissn. in Bot. Zeit. xiii. (1855) 28 = macranthum.
racemulosum, *Benth. Fl. Austral.* ii. 65.—Austral.
scabriusculum, *Meissn. in Lehm. Pl. Preiss.* ii. 214.—Austral.
stenopterum, Meissn. l. c. i. 57 = alatum.
vimineum, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 509.—Austral.

SPHAEROMA, Schlecht. in Linnaea, xi. (1837) 352 = **Sphaeralcea**, A. St. Hil. (Malvac.).
angustifolium, Schlecht. l. c. 353 = Sphaeralcea angustifolia.
floribundum, Schlecht. l. c. = Sphaeralcea floribunda.
Julii, Harv. in Harv. & Sond. Fl. Cap. i. 166.—Afr. austr.
miniatum, Garcke, in Bot. Zeit. xi. (1853) 847, in obs. = Sphaeralcea cisplatina.
prostrata, Harv. in Harv. & Sond. Fl. Cap. i. 166.—Afr. austr.
rosea, Schlecht. in Linnaea, xi. (1837) 352 = Sphaeralcea umbellata.
umbellatum, Schlecht. l. c. = Sphaeralcea umbellata.

SPHAEROMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 401 = **Tanacetum**, Tourn. (Compos.).
argentea, Nutt. l. c. 402 = T. Nuttallii.
capitata, Nutt. l. c. = T. capitatum.

SPHAEROMORPHAEA, DC. Prod. vi. 140 (1837). *COMPOSITAE*, [Benth. & Hook. f. ii. 430.]
Centipeda, DC. l. c. = Centipeda orbicularis.
petiolaris, DC. l. c. = Epaltes australis.
Russeliana, *DC. l. c.*—Ind. or.

SPHAEROPHORA, Blume, Mus. Bot. Lugd. Bat. i. 179. f. 36 (1850) = **Morinda**, Linn. (Rubiac.).
gemella, Blume, l. c. = M. gemella.
glomerata, Blume, l. c. = M. glomerata.

SPHAEROPHORA, Sch. Bip. in Pollichia, xx-xxi. (1863) 402 = **Eremanthus**, Less. (Compos.).
bicolor, Sch. Bip. l. c. 403 = E. bicolor.

SPHAEROPHYSA, DC. Mém. Lég. 288 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 504.
caspica, DC. Prod. ii. 271 = salsula.
Kotschyana, *Boiss. Fl. Orient.* ii. 197.—Cappadoc.

SPHAEROPHYSA :—
microphylla, Jaub. & Spach, Illustr. i. 126. t. 64 = Colutea triphylla.
pycnorrhiza, Benth. & Hook. f. Gen. i. 504 = Astragalus pycnorrhizus.
salsula, *DC. Prod.* ii. 271.—Reg. Casp.; As. bor.

SPHAEROPUS, Boeck. in Flora, lvi. (1873) 89 = **Scleria**, Berg. (Cyperac.).
pygmaeus, Boeck. l. c. 90 = Sc. pygmaea.

SPHAERORHIZON, Hook. f. in Trans. Linn. Soc. xxii. (1856) 50. t. 10 = **Scybalium**, Schott (Balanoph.).
curvatum, Hook. f. l. c. 50 = Sc. depressum.
depressum, Hook. f. l. c. t. 10 = Sc. depressum.

SPHAEROSACME, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 429 (1824) = **Amoora**, Roxb. (Meliac.).
decandra, Wall. Cat. n. 1276 = A. decandra.
fragrans, Wall. ex Voigt, Hort. Suburb. Calc. 135 = A. decandra.
laxa, Wall. Cat. n. 4894 = A. cucullata.
nepalensis, M. Roem. Syn. Hesper. 98 = A. decandra.
paniculata, Wall. ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 37 = A. cucullata.
polystachya, Wall. Cat. n. 1277 = A. Rohituka.
Rohituka, Wall. l. c. n. 1278 = A. cucullata.
spectabilis, Royle, Illustr. Bot. Himal. 142 = A. spectabilis ?
spicata, Wall. Cat. n. 4895 = A. Rohituka.

SPHAEROSCHOENUS, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. 97 (1843) = **Rynchospora**, Vahl (Cyperac.).
Wallichii, Nees, l. c. = R. Wallichiana.

SPHAEROSEPALUM, Baker, in Journ. Linn. Soc. xxi. (1884) 321. *GUTTIFERAE*.
alternifolium, *Baker, l. c.*—Madag.

SPHAEROSICYOS, Hook. f. in Benth. & Hook. f. Gen. i. 824 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
LAGENARIA, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 8, partim.
Meyeri, Hook. f. in Oliver, Fl. Trop. Afr. ii. 532 = sphaericus.
sphaericus, *Hook. f. in Benth. & Hook. f. Gen.* ii. 824; *Cogn. in DC. Monog. Phan.* iii. 466.—Afr. trop. et austr.

SPHAEROSPORA, Klatt, in Linnaea, xxxii. (1863) 725 = **Acidanthera**, Hochst. (Irid.).
exscapa, Klatt, l. c. = A. tubulosa.
flexuosa, Klatt, l. c. 726 = A. flexuosa.
gigantea, Klatt, l. c. xxxiv. (1865-66) 690 = A. bicolor.

SPHAEROSPORA, Sweet, Hort. Brit. ed. I. 398 (1827) = **Gladiolus**, Tourn. (Irid.).
imbricata, Sweet, l. c. ed. II. 501 = G. segetum.
segetum, Sweet, l. c. ed. I. 398 = G. segetum.
triphylla, Sweet, l. c. ed. II. 501 = G. triphyllos.

SPHAEROSTACHYS, Miq. Syst. Pip. 375 (1843) = **Piper**, Linn.
Humboldtii, Miq. l. c. 376 = P. piluliferum.

SPHAEROSTEMA, Blume, Bijdr. 22 (1825) = **Schizandra**, Michx. (Magnol.).
axillare, Blume, l. c. 22 = Schiz. axillaris.
Blumiana, Griff. Notul. iv. 714. t. 651. f. 2; 654 = Schiz. axillaris ?
elongatum, Blume, Bijdr. 23 = Schiz. elongata.
grandiflorum, Blume, Fl. Jav. Schiz. 17 = Schiz. grandiflora.
grandiflorum, Wall. Cat. n. 4985 = Schiz. elongata.
japonicum, A. Gray, in Mem. Acad. Am. N. S. vi. (1858) 380 = seq. ?
japonicum, Sieb. & Zucc. Fl. Jap. Fam. Nat. 80 = Schiz. chinensis.
propinquum, Blume, Fl. Jav. Schiz. 16, in obs. = Schiz. propinqua.
pyrifolium, Blume, l. c. 16. t. 4.—Java.

SPHAEROSTEMMA, Reichb. Consp. 86 (1828), sphalm. = praec.

SPHAEROSTIGMA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. (1835) 49 = **Oenothera**, Linn. (Onagrar.).
alyssoides, Walp. Rep. ii. 78 = Oe. alyssoides.
andinum, Walp. l. c. 79 = Oe. andina.
bistorta, Walp. l. c. 77 = Oe. bistorta.
Boothii, Walp. l. c. = Oe. Boothii.
Chamissonis, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 49 = Oe. dentata.
cheiranthifolium, Fisch. & Mey. l. c. 50 = Oe. cheiranthifolia.
contortum, Walp. Rep. ii. 78 = Oe. strigulosa.
dentatum, Walp. l. c. = Oe. dentata.
divaricatum, C. Gay, Fl. Chil. ii. 328.—Chili.
epilobioides, Walp. Rep. ii. 78 = Oe. epilobioides.
guaraeflorum, Walp. l. c. = Oe. gauraeflora.
heterophyllum, C. Gay, Fl. Chil. ii. 327 = Oe. heterophylla, *Steud.*
hirtum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 50 = Oe. hirta.
micranthum, Walp. Rep. ii. 77 = Oe. micrantha.
minimum, Heynh. Nom. ii. 692 = Oe. hirta.
minutiflorum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 50 = Oe. minutiflora.
paradoxum, C. Gay, Fl. Chil. ii. 329 = Oe. micrantha.
parvulum, Walp. Rep. ii. 78 = Oe. strigulosa.
ramosissimum, Phil. in Linnaea, xxxiii. (1864–65) 68.—Chili.
spirale, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 50 = Oe. cheiranthifolia.
strigulosum, Fisch. & Mey. l. c. = Oe. strigulosa.
tenuifolium, C. Gay, Fl. Chil. ii. 327 = Oe. chilensis.
viridescens, Walp. Rep. ii. 77 = Oe. cheiranthifolia.

SPHAEROSTYLIS, Baill. Etud. Gén. Euph. 466. t. 21 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 327.
Tulasneana, *Baill. l. c.* 467.—Madag.

SPHAEROTELE, Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol. 95. t. 38 (1841) = **Urceolina**, Reichb. (Amaryll.).
coccinea, Link, Klotzsch & Otto, l. c. = **U.** miniata.

SPHAEROTELE, Presl, Rel. Haenk. i. 119. t. 16 (1827) = **Stenomesson**, Herb. (Amaryll.).
peruviana, Presl, l. c. 120. t. 16.—Peruv.

SPHAEROTHALAMUS, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 156. *ANONACEAE*, Benth. & Hook. f. i. 28.
insignis, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 156.—Borneo.

SPHAEROTHECA, Cham. & Schlecht. in Linnaea, ii. (1827) 605 = **Conobea**, Aubl. (Scrophul.).
scoparioides, Cham. & Schlecht. l. c. 606 = C. scoparioides.

SPHAEROTHELE, Benth. & Hook. f. Gen. iii. 733 (1883) = SPHAEROTELE, Presl = **Stenomesson**, Herb. (Amaryll.).

SPHAEROTHYLAX, Bisch. ex Krauss, in Flora, xxvii. (1844) 426. t. 1. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 113.
ANASTROPHEA, Wedd. in DC. Prod. xvii. 78 (1873).
algiformis, *Bisch. ex Krauss, in Flora*, xxvii. (1844) 426.—Afr. austr.

SPHAERULA, W. Anders. ex Hook. f. Fl. Antarct. 268 (1847) = **Acaena**, Linn. (Rosac.).

SPHAERULARIA, Heist. Syst. 9 (1748) = **Rapistrum**, Tourn. (Crucif.).

SPHALANTHUS, Jack, in Malay. Misc. ii. (1822) VII. 55 = **Quisqualis**, Linn. (Combret.).
confertum, Jack, l. c. = Q. densiflora.
ovatifolia, Steud. Nom. ed. II. ii. 621, sphalm. = praec.

SPHALLEROCARPUS, Bess. ex DC. Mém. Ombell. 60 (1829); Prod. iv. 230 (1830) = **Conopodium**, Koch (Umbell.).
Cyminum, Bess. ex DC. ll. cc. = C. Cyminum.
longilobus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 432 = Chaerophyllum Sphallerocarpus.

SPHANELLOPSIS, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 137 = **Adelobotrys**, DC. (Melastom.).
ciliata, Steud. ex Naud. l. c. = A. ciliata.

SPHEDAMNOCARPUS, Planch. ex Benth. & Hook. f. Gen. i. 256 (1862). *MALPIGHIACEAE*, Benth. & Hook. f. l. c.
angolensis, *Planch. in Oliver, Fl. Trop. Afr.* i. 279.—Afr. trop.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 110.—Madag.

SPHENANDRA, Benth. in Lindl. Introd. Nat. Syst. ed. II. 445 (1836). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 945.
viscosa, *Benth. in Lindl. Introd. Nat. Syst.* ii. 445.—Afr. austr.

SPHENANTHA, Schrad. in Linnaea, xii. (1838) 416 = **Cucurbita**, Tourn.
scabra, Schrad. l. c. = C. radicans.
scandens, Steud. Nom. ed. II. ii. 621 = C. radicans.

SPHENANTHERA, Hassk. in Versl. Kon. Akad. Wetensch. iv. (1855) 139 = **Begonia**, Tourn.
erosa, Hassk. ex Klotzsch, in Bot. Zeit. xv. (1857) 182 = B. erosa.
multangula, Klotzsch, l. c. 181 = B. multangula.
robusta, Hassk. in Versl. Kon. Akad. Wetensch. iv. (1855) 139 = B. robusta.

SPHENDAMNOCARPUS, Baker, in Journ. Linn. Soc. xx. (1813) 110, sphalm. = **Sphedamnocarpus**, Planch. (Malpig.).

SPHENISTA, Rafin. Sylva Tellur. 90 (1838) = **Hirtella**, Linn. (Rosac.).
peruviana, Rafin. l. c. 91 = H. Cosmibuena.

SPHENOCARPUS, Rich. Anal. Fr. 92 (1808) = **Laguncularia**, Gaertn. (Combret.).

SPHENOCARPUS, Wall. Cat. p. 236, sub n. 975 (1832 ?) = **Magnolia**, Linn.

SPHENOCLEA, Gaertn. Fruct. i. 113. t. 24 (1788). *CAMPANULACEAE*, Benth. & Hook. f. ii. 560.
GAERTNERA, Retz. Obs. vi. 24 (1791).
PONGATIUM, Juss. Gen. 423 (1789).
RAPINIA, Lour. Fl. Cochinch. 127 (1790).
Pongatium, DC. Prod. vii. 548 = zeylanica.
zeylanica, *Gaertn. Fruct.* i. 113. t. 24.—Cosmop. trop.

SPHENODESMA, Griff. Notul. iv. 176 (1854) = seq.

SPHENODESME, Jack, in Malay. Misc. i. (1820) I. 19. *VERBENACEAE*, Benth. & Hook. f. ii. 1159.
BRACHYNEMA, Griff. Notul. iv. 176 (1854).
DECADONTIA, Griff. l. c. 175 (1854).
ROSCOEA, Roxb. Cat. Hort. Beng. 46 (1814), nomen.
SPHAENODESMA, Schau. in DC. Prod. xi. 622 (1847).
VITICASTRUM, Presl, Bot. Bemerk. 147 (1844).
acuminata, Wight, Ic. t. 1476 = pentandra.
barbata, *Schau. in DC. Prod.* xi. 623.—Malaya.
eryciboides, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 76.—Burma.
ferruginea, Wight, Ic. t. 1474 = barbata.
Griffithiana, *Wight, l. c.* iv. III. 14.—Burma.
grossa, Kurz, For. Fl. Brit. Burma, ii. 254 = eryciboides.
Jackiana, Schau. in DC. Prod. xi. 622 = pentandra.
Jackiana, Wight, Ic. t. 1477 = Griffithiana.
microstylis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 600.—Burma.
paniculata, *C. B. Clarke, l. c.*—Ind. or.
pentandra, *Jack, in Malay. Misc.* i. (1820) I. 19.—Ind. or.; Malaya.
triflora, *Wight, Ic.* t. 1478.—Malaya.
unguiculata, *Schau. in DC. Prod.* xi. 622.—Reg. Himal.; Burma.
Wallichiana, Schau. l. c. = pentandra.

SPHENOGYNE, R. Br. in Ait. Hort. Kew. ed. II. v. 142 (1813) = **Ursinia**, Gaertn. (Compos.).
abrotanifolia, R. Br. l. c. 143 = U. abrotanifolia.
adonidifolia, DC. Prod. v. 682 = U. anthemoides.
anethifolia, Less. Syn. Comp. 243.—Afr. austr.
anethoides, DC. Prod. v. 685.—Afr. austr.
anthemoides, R. Br. in Ait. Hort. Kew. ed. II. v. 142 = U. anthemoides.
anthemoides, E. Mey. ex DC. Prod. v. 689 = U. paradoxa.

SPHENOGYNE:—
brachyloba, Kunze, in Linnaea, xx. (1847) 21.—Afr. austr.
brachypoda, Harv. in Harv. & Sond. Fl. Cap. iii. 140.—Afr. austr.
brevifolia, DC. Prod. v. 685 = *discolor*.
calendulaeflora, DC. l. c. 682.—Afr. austr.
chamaemeloides, DC. l. c. 685.—Afr. austr.
chamomillaefolia, DC. l. c. 687.—Afr. austr.
chrysanthemoides, Less. Syn. Comp. 242 = Ursinia chrysanthemoides.
ciliaris, DC. Prod. v. 687.—Afr. austr.
concolor, Harv. in Harv. & Sond. Fl. Cap iii. 142.—Afr. austr.
coronopifolia, Less. Syn. Comp. 244.—Afr. austr.
crithmifolia, R. Br. in Ait. Hort. Kew. ed. II. v. 142 = U. crithmifolia.
crocea, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1090 = U. speciosa.
dentata, R. Br. in Ait. Hort. Kew. ed. II. v. 143 = U. dentata.
discolor, Less. Syn. Comp. 243.—Afr. austr.
Dregeana, DC. Prod. v. 683.—Afr. austr.
Eckloniana, Sond. in Linnaea, xxiii. (1850) 64.—Afr. austr.
elegans, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1090 = U. speciosa.
filipes, E. Mey. ex DC. Prod. v. 687; et in Drège, Zwei Pfl. Docum. 223.—Afr. austr.
foeniculacea, Sweet, Hort. Brit. ed. I. 238 = U. foeniculacea.
gracilis, DC. Prod. v. 685 = *punctata*.
grandiflora, DC. l. c. = U. crithmifolia.
heterodonta, DC. l. c. 687.—Afr. austr.
hispida, DC. l. c. 683.—Afr. austr.
incisa, DC. l. c. 688.—Afr. austr.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 674 = *chamomillaefolia*.
leptoglossa, DC. Prod. v. 687.—Afr. austr.
leucanthemoides, Sweet, Hort. Brit. ed. I. 238 = U. leucanthemifolia.
macropoda, DC. Prod. v. 683.—Afr. austr.
microcephala, DC. l. c. 682 = U. anthemoides.
natalensis, Sch. Bip. in Flora, xxvii. (1844) 674 = U. foeniculacea.
nudicaulis, Less. Syn. Comp. 243.—Afr. austr.
odorata, R. Br. in Ait. Hort. Kew. ed. II. v. 142 = U. odorata.
paleacea, Less. Syn. Comp. 244 = U. paleacea.
pallida, DC. Prod. v. 682 = U. foeniculacea.
pauciloba, DC. l. c. 686.—Afr. austr.
pedunculatis, E. Mey. ex DC. l. c. 683 = *sericea*.
pilifera, Ker-Gawl. in Bot. Reg. t. 604; Less. Syn. Comp. 244 = U. pilifera.
pinnata, Less. l. c. 243 = U. filiformis.
punctata, Less. l. c.—Afr. austr.
pusilla, DC. Prod. v. 682.—Afr. austr.
quinquepartita, DC. l. c. 686.—Afr. austr.
rigidula, DC. l. c. 685.—Afr. austr.
scapiformis, DC. l. c. 684.—Afr. austr.
scariosa, R. Br. in Ait. Hort. Kew. ed. II. v. 142 = U. scariosa.
sericea, Less. Syn. Comp. 243.—Afr. austr.
serrata, DC. Prod. v. 688 = U. serrata.
setigera, DC. l. c. 684 = U. dentata.
Sonneratii, Cass. in Dict. Sc. Nat. l. 205 = U. serrata.
speciosa, Hort. ex Knowles & Westc. Flor. Cab. ii. (1838) 131 = U. speciosa.
subflosculosa, E. Mey. ex DC. Prod. v. 686; et in Drège, Zwei Pfl. Docum. 223 = U. paleacea.
subhirsuta, DC. Prod. v. 683.—Afr. austr.
tenuifolia, DC. l. c. 687 = U. tenuifolia.
tridentata, Harv. in Harv. & Sond. Fl. Cap. iii. 149.—Afr. austr.
trifida, Less. Syn. Comp. 244.—Afr. austr.
trifurca, Harv. in Harv. & Sond. Fl. Cap. iii. 147.—Afr. austr.
triloba, DC. Prod. v. 687 = *trifida*.
tripartita, DC. l. c. 686.—Afr. austr.
versicolor, DC. l. c. 681.—Afr. austr.
versicolor, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1090 = U. speciosa.

SPHENOPUS, Trin. Fund. Agrost. 135 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1185.
divaricatus, Reichb. Fl. Germ. Excurs. 45 = Gouani.
Gouani, *Trin. Fund. Agrost.* 135.—Reg. Mediterr.

SPHENOSCIADIUM, A. Gray, in Proc. Am. Acad. vi. (1865) 536 = **Selinum**, Linn. (Umbell.).
capitellatum, A. Gray, l. c. 537 = Selinum capitellatum.

SPHENOSTEMON, Baill. in Bull. Soc. Linn. Par. i. (1875) 53. *ILICINEAE.*
Balansae, *Baill. l. c.* 54; *et Adansonia*, xi. (1875) 307.—N. Caled.
pachycladum, *Baill. ll. cc.*—N. Caled.

SPHENOSTIGMA, Baker, in Journ. Linn. Soc. xvi. (1877) 124. *IRIDEAE*, Benth. & Hook. f. iii. 694.
CARDIOSTIGMA, Baker, in Journ. Linn. Soc. xvi. (1877) 129.
angustifolium, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 363.—Bras.
caeruleum, *Klatt, l. c.*—Bras.
geniculatum, *Klatt, l. c.*—Bras.
gracile, *Benth. & Hook. f. Gen.* iii. 695.—Bras.
lineare, Benth. & Hook. f. l. c. (Cypellae sp.).—Venezuela.
longispathum, *Benth. & Hook. f. l. c.*—Mexic.
Sellowianum, *Baker, in Journ. Linn. Soc.* xvi. (1877) 124.—Bras.
umbellatum, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 363.—Bras.

SPHENOSTYLIS, E. Mey. Comm. Pl. Afr. Austr. 148 (1835) = **Vigna**, Savi (Legumin.).
angustifolia, Sond. in Linnaea, xxiii. (1850) 33 = V. angustifolia.
marginata, E. Mey. Comm. Pl. Afr. Austr. 148 = V. marginata.

PHENOTOMA, Sweet, Fl. Austral. t. 44 (1828) = **Dracophyllum**, Labill. (Epacrid.).
capitatum, Lindl. Bot. Reg. t. 1515 = D. capitatum.
dracophylloides, Sond. in Lehm. Pl. Preiss. i. 335 = D. phlogiflorum.
Drummondii, F. Muell. Census, 109 = D. Drummondii.
gracile, Sweet, Fl. Austral. t. 44 = D. gracile.
parviflorum, F. Muell. Census, 109 = D. parviflorum.
squarrosa, G. Don, Gen. Syst. iii. 785 = D. squarrosum.

SPHERANTHUS, Hill, Veg. Syst. ii. 113 (1761), sphalm. = **Sphaeranthus**, Vaill. (Compos.).

SPHINCTACANTHUS, Benth. in Benth. & Hook. f. Gen. ii. 1118 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
Griffithii, *Benth. l. c.*; *et in Hook. Ic. Pl.* t. 1205.—Reg. Himal.

SPHINCTANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 212. *RUBIACEAE*, Benth. & Hook. f. ii. 84.
CONOSIPHON, Poepp. & Endl. Nov. Gen. et Sp. iii. 27. t. 23 [1842]; Endl. Gen. Suppl. ii. 54 (1842).
longiflorus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 503.—Cuba.
rupestris, *Benth. in Hook. Journ. Bot.* iii. (1841) 212.—Guian.

SPHINCTEROSTIGMA, Schott, Melet. 20 (1832) = **Philodendron**, Schott (Aroid.).
bipinnatifidum, Schott, l. c. = P. bipinnatifidum.

SPHINCTEROSTOMA, Stschégl. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 22 = **Andersonia**, R. Br. (Epacrid.).
axilliflorum, Stschégl. l. c. = A. colossea.
echinocephalum, Stschégl. l. c. 23 = A. patricia.

SPHINCTOLOBIUM, Vog. in Linnaea. xi. (1837) 417 = **Lonchocarpus**, H. B. & K. (Legumin.).
floribundum, Vog. l. c. 419 = L. subglaucescens.
glaucescens, Miq. Stirp. Surin. Sel. 19 = Muellera moniliformis.
nitidum, Vog. in Linnaea, xi. (1837) 419 = L. nitidus.
virgilioides, Vog. l. c. 418 = L. virgilioides.

SPHINCTOSTOMA. Benth. ex Benth. & Hook. f. Gen. ii. 772 (1876) = **Marsdenia**, R. Br. (Asclep.).

SPHINGIUM, E. Mey. Comm. Pl. Afr. Austr. 65 (1835) = **Melolobium**, Eckl. & Zeyh. (Legumin.).
canaliculatum, E. Mey. l. c. 66 = M. alpinum.
canescens, E. Mey. l. c. 67 = M. canescens.
decumbens, E. Mey. l. c. = M. decumbens.
campolobum, E. Mey. l. c. 66 = M. adenoides.
microphyllum, E. Mey. l. c. 67 = M. microphyllum.
spicatum, E. Mey. l. c. 66 = M. cernuum.
velutinum, E. Mey. l. c. 67 = M. candicans.
viscidulum, E. Mey. l. c. 66 = M. viscidulum.

SPHONDYLANTHA, Endl. Gen. 1195 (1841) = SPONDYLANTHA, Presl = **Vitis**, Tourn. (Ampel.).

SPHONDYLASTRUM, Reichb. Nom. 169 (1841) = **Myriophyllum**, Ponted. (Halorag.).

SPHONDYLIUM, [Tourn.] Adans. Fam. ii. 100 (1763) = **Heracleum**, Linn. (Umbell.).
alpinum, Scop. ex Steud. Nom. ed. II. ii. 622 = H. alpinum.
asperum, Hoffm. Gen. Umbellif. ed. I. 134 = H. asperum.
austriacum, Scop. Fl. Carn. ed. II. i. 204 = H. austriacum.
Branca, Scop. l. c. = H. Sphondylium.
conforme, Moench, Meth. 83 = H. flavescens.
decipiens, Hoffm. Gen. Umbellif. ed. I. 134 = H. pyrenaicum.
giganteum, Hoffm. l. c. 133 = H. villosum.
gummiferum, Hoffm. l. c. 135 = H. pubescens, asperum.
longifolium, Hoffm. l. c. 134 = H. Sphondylium.
marginatum, Hoffm. l. c. = H. villosum.
Panaces, Gaertn. Fruct. i. 86. t. 21. f. 4 = H. Panaces.
pubescens, Hoffm. Gen. Umbellif. ed. I. 133 = H. pubescens.
speciosum, Hoffm. l. c. 134 = H. villosum.
subvillosum, Hoffm. l. c. 133 = H. villosum.
villosum, Hoffm. l. c. = H. villosum.
vulgare, S. F. Gray, Nat. Arr. Brit. Pl. ii. 520 = H. Sphondylium.

SPHONDYLOCOCCA, Willd. ex Schult. Syst. vi. p. lxx. 799 (1820) = **Bergia**, Linn. (Elatin.).
malabarica, Willd. l. c. 799.—Ind. or.

SPHONDYLOCOCCUM, Schau. in DC. Prod. xi. 640 (1847) = **Callicarpa**, Linn. (Verben.).

SPHRAGIDIA, Thw. in Hook. Kew Journ. vii. (1855) 269. t. 10 = **Cyclostemon**, Blume (Euphorb.).
zeylanica, Thw. l. c. 270 = C. macrophyllus.

SPHYROSPERMUM, Poepp. & Endl. Nov. Gen. et Sp. i. 4. t. 8 (1835). *VACCINIACEAE*, Benth. & Hook. f. ii. 576.
buxifolium, *Poepp. & Endl. l. c.*—Peruv.
cordifolium, Benth. Pl. Hartw. 222 = Sophoclesia cordifolia.
longifolium, *Poepp. & Endl. Nov. Gen. et Sp.* i. 5.—Peruv.
majus, Griseb. in Lechl. Berb. Am. Austr. 58, nomen; et Fl. Brit. W. Ind. 148 = Sophoclesia major.
myrtifolium, *Hook. Ic. Pl.* t. 112.—Peruv.
Roraimae, *Klotzsch, in Linnaea*, xxiv. (1851) 36.—Guian.

SPICILLARIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 252 = **Petunga**, DC. (Rubiac.).
Leschenaultii, A. Rich. l. c. = P. Roxburghii.

SPICIVISCUM, Engelm. in Mem. Am. Acad. N. S. iv. (1849) 58 = **Phoradendron**, Nutt. (Loranth.).
polygynum, Karst. Fl. Columb. i. 73. t. 36 = P. polygynum.

SPICULAEA, Lindl. Swan River App. 56 (1839) = **Drakaea**, Lindl. (Orchid.).
ciliata, Lindl. l. c. = D. ciliata.

SPIELMANNIA, Cuss. ex Juss. in Dict. Sc. Nat. lv. 328 (1828) = **Trinia**, Hoffm. (Umbell.).

SPIELMANNIA, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 196. t. 15 = **Oftia**, Adans. (Myopor.).
africana, Willd. Sp. Pl. iii. 321 = O. africana.
decurrens, Moench, Meth. 479 = O. africana.

SPIELMANNIA:—
desertorum, Eckl. & Zeyh. ex Schau. in DC. Prod. xi. 526 = Oftia revoluta.
Jasminum, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 196. t. 15 = O. africana.
revoluta, E. Mey. Comm. Pl. Afr. Austr. 274 = O. revoluta.

SPIESIA, Neck. Elem. iii. 13 (1790) = **Astragalus**, Tourn. (Legumin.).

SPIGELIA, Linn. Gen. ed. I. 361 (1737). *LOGANIACEAE*, Benth. & Hook. f. ii. 790.
ARAPABACA, Adans. Fam. ii. 225 (1763).
CANALA, Pohl, Pl. Bras. Ic. ii. 62. t. 142 (1831).
COELOSTYLIS, Torr. & Gray, ex Endl. Nov. Stirp. Dec. 33 (1839).
HEINZELMANNIA, Neck. Elem. i. 371 (1790).
MONTIRA, Aubl. Pl. Gui. ii. 637. t. 257 (1775).
aenea, *Lem. Illustr. Hortic.* (1857) *Misc.* 52.—Am. centr.
ambigua, *Wright, in Sauv. Fl. Cub.* 116.—Cuba.
Anthelmia, *Linn. Sp. Pl.* 149.—Am. trop.
asperifolia, *Prog. in Mart. Fl. Bras.* vi. 1. 260.—Bras.
Beyrichiana, *Cham. & Schlecht. in Linnaea*, i. (1826) 203.—Bras.
Blanchetiana, *A. DC. in DC. Prod.* ix. 4.—Bras.
brachystachya, *Prog. in Mart. Fl. Bras.* vi. 1. 261.—Bras.
Coulteriana, *Benth. in Journ. Linn. Soc.* i. (1857) 90.—Mexic.
Flemmingiana, *Cham. & Schlecht. in Linnaea*, i. (1826) 203.—Bras.
fruticulosa, *Lam. Illustr.* i. 474.—Guian.
gentianoides, *Chapm. ex A. DC. in DC. Prod.* ix. 5.—Florida.
glabrata, *Mart. Nov. Gen. et Sp.* ii. 127. t. 193.—Bras.
gracilis, A. DC. in DC. Prod. ix. 6 = spartioides.
hamelioides, *H. B. & K. Nov. Gen. et Sp.* iii. 185 (hamellioides).—N. Granat.
hedyotidea, *A. DC. in DC. Prod.* ix. 7.—Mexic.
Humboldtiana, *Cham. & Schlecht. in Linnaea*, i. (1826) 200.—Am. trop.
humilis, *Benth. in Hook. Journ. Bot.* iii. (1841) 240.—Bras.
insignis, *Prog. in Mart. Fl. Bras.* vi. 1. 259.—Bras.
japonica, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 108 = Ophiorrhiza japonica.
laevigata, *Prog. in Mart. Fl. Bras.* vi. 1. 263.—Bras.
laurina, *Cham. & Schlecht. in Linnaea*, i. (1826) 204.—Bras.
laurina, A. DC. in DC. Prod. ix. 4, partim = glabrata.
linarioides, *A. DC. l. c.* 6.—Bras.
Lindheimeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 108.—Am. bor. occ.
loganioides, *A. DC. in DC. Prod.* ix. 4.—Florida.
longiflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 376.—Mexic.
Lonicera, Mill. Gard. Dict. ed. VIII. n. 2 = marilandica.
Lundiana, *A. DC. in DC. Prod.* ix. 3.—Bras.
macrophylla, *A. DC. l. c.* 8.—Bras.
marilandica, *Linn. Syst.* ed. XII. 734.—Am. bor.
Martiana, *Cham. in Linnaea*, viii. (1833) 15.—Bras.
mexicana, *A. DC. in DC. Prod.* ix. 7.—Mexic.
multispica, Steud. in Flora, xxvi. (1843) 764 = S. Anthelmia.
nervosa, Steud. l. c. = S. Anthelmia.
Olfersiana, *Cham. & Schlecht. in Linnaea*, i. (1826) 206.—Bras.
oppositifolia, Stokes, Bot. Mat. Med. i. 307 = S. Anthelmia.
pauciflora, A. DC. in DC. Prod. ix. 7 = pusilla.
pauciflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 376.—Mexic.
pedunculatis, D. Dietr. Syn. Pl. i. 681 = pedunculata.
pedunculata, *H. B. & K. Nov. Gen. et Sp.* iii. 185.—N. Granat.
platyphylla, Prog. in Mart. Fl. Bras. vi. 1. 256 = splendens.
polita, *Prog. l. c.* 259.—Bras.
polystachya, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1082, nomen; *et ex Prog. l. c.* 265.—Guian.
pulchella, *Mart. Nov. Gen. et Sp.* ii. 128. t. 194.—Bras.

SPIGELIA:—
pulverulenta, Mart. Nov. Gen. et Sp. ii. 126. t. 192 = Olfersiana.
pusilla, *Mart. l. c.* 130.—Bras.
quadrifolia, Stokes, Bot. Mat Med. i. 307 = marilandica.
rubiaefolia, A. DC. in DC. Prod. ix. 8 = Schlechtendaliana.
scabra, *Cham. & Schlecht. in Linnaea*, i. (1826) 202.—Bras.
scabrella, *Benth. Pl. Hartw.* 45.—Mexic.
Schlechtendaliana, *Mart. Nov. Gen. et Sp.* ii. 129.—Bras.
Schomburgkiana, Benth. in Hook. Journ. Bot. iii. (1841) 240 = Humboldtiana.
Schomburgkiana, A. DC. in DC. Prod. ix. 8 = Flemmingiana.
Selloi, Spreng. Syst. iv. Cur. Post. 59 = seq.
Sellowiana, *Cham. & Schlecht. in Linnaea*, i. (1826) 205.—Bras.
spartioides, *Cham. in Linnaea*, viii. (1833) 14.—Bras.
spartioides, Griseb. ex Wright, in Sauv. Fl. Cub. 117 = sphagnicola.
speciosa, *H. B. & K. Nov. Gen. et Sp.* iii. 186. t. 224.—Mexic.
sphagnicola, *Wright, in Sauv. Fl. Cub.* 116.—Cuba.
sphagnicosa, Wright, l. c. 74, err. typ. = praec.
splendens, *Hort. Wendl. ex Hook. Bot. Mag.* t. 5268.—Mexic.; Guatemala.
stenophylla, *Prog. in Mart. Fl. Bras.* vi. 1. 260.—Bras.
stipularis, *Prog. l. c.* 262—Bras.
texana, *A. DC. in DC. Prod.* ix. 5.—Texas.

SPILACRON, Cass. in Dict. Sc. Nat. l. 238 (1827) = **Centaurea**, Linn. (Compos.).
crupinoides, Cass. l. c. 239 = C. arenaria.

SPILANTHES, Jacq. Enum. Pl. Carib. 8 (1760).
COMPOSITAE, Benth. & Hook. f. ii. 380.
ACMELLA, Rich. in Pers. Syn. ii. 472 (1807).
ATHRONIA, Neck. Elem i. 32 (1790).
CERUCHIS, Gaertn. ex Schreb. Gen. 543 (1791).
MENDEZIA, DC. Prod. v. 532 (1836).
PYRETHRUM, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 237. t. 18.
SPILANTHUS, Linn. Mant. ii. 475 (1771).
abyssinica, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 26, nomen; et A. Rich. Tent. Fl. Abyss. i. 415 = S. Acmella.
Acmella, *Murr. Syst.* ed. XIII. 610.—Cosmop. trop. et subtrop.
affinis, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 317 = arnicoides.
africana, DC. Prod. v. 623 = S. Acmella.
alba, *L'Hérit. Stirp. Nov.* 7. t. 4.—Peruv.
alpestris, *Griseb. in Goett. Abh.* xix. (1874) 185.—Reg. Argent.
anactina, *F. Muell. Fragm.* v. 63.—Borneo; Austral.
arborea, Forst. f. in Comm. Gotting. ix. 1787–88 (1789) 66 = Petrobium arboreum.
arnicoides, *DC. Prod.* v. 620.—Bras.
Arrayana, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 408 = S. Acmella.
atriplicifolia, Houtt. Handleid. x. 536; Miq. Fl. Ind. Bat. ii. 37 = Dichrocephala latifolia.
atriplicifolia, Linn. Syst. ed. XII. iii. 236 = Isocarpha atriplicifolia.
Beccabunga, *DC. Prod.* v. 622.—Mexic.
bicolor, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 192.—Mexic.
blepharicarpa, DC. Prod. v. 621 = arnicoides.
brasiliensis, Spreng. Syst. iii. 444 = S. Acmella.
caespitosa, *DC. Prod.* v. 622.—Bras.
calva, DC. in Wight, Contrib. 19 = S. Acmella.
caulirhiza, DC. Prod. v. 623 = S. Acmella.
ciliata, *H. B. & K. Nov. Gen. et Sp.* iv. 208.—Am. austr.
commutata, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 14.—Am. bor.
costata, Benth. in Hook. Niger Fl. 436 = S. Acmella.
crocata, Sims, Bot. Mag. t. 1627 = Verbesina crocata.
debilis, H. B. & K. Nov. Gen. et Sp. iv. 210 = S. Acmella.
diffusa, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 214.—Ins. Galapag.
diffusa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 50.—Peruv.

SPILANTHES:—
doronicoides, DC. Prod. v. 620 = arnicoides.
ecliptoides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 407 = Jaegeria hirta.
exasperata, *Jacq. Ic. Pl. Rar.* iii. 15. t. 584.—Venezuela.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* iv. 208.—N. Granat.
fusca, Hort. Par. ex Lam. Encyc. i. 416 = S. Acmella.
grandiflora, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 185 = S. Acmella.
grandifolia, *Miq. Fl. Ind. Bat.* ii. 80.—Java.
grandis, DC. Prod. v. 622 = arnicoides.
helenioides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 317 = arnicoides.
hirta, *Steud. Nom.* ed. II. ii. 622.—Am. austr.
insipida, *Jacq. Enum. Pl. Carib.* 28; *Select. Am.* 215. t. 126. f. 2.—Cuba.
intermedia, *DC. Prod.* v. 624.—Am.
javanica, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 123, nomen.—Malaya.
Karvinskiana, *DC. Prod.* v. 623.—Mexic.
leiocarpa, *DC. l. c.* 626.—Peruv.
leptophylla, DC. l. c. 621 = arnicoides.
leucantha, *H. B. & K. Nov. Gen. et Sp.* iv. 210.—Ecuador.
lobata, Blanco, Fl. Filip. ed. I. 622 = S. Acmella.
longifolia, DC. Prod. v. 621 = arnicoides.
Lundii, DC. l. c. 622 = S. Acmella.
Macraei, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 317.—Chili.
macroglossa, F. Muell. Fragm. v. 63 = S. Acmella.
macropoda, DC. Prod. v. 621 = arnicoides.
macropoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 71.—Venezuela.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 529.—Bolivia.
Mariannae, DC. Prod. v. 623 = S. Acmella.
mauritiana, DC. l. c. 625 = S. Acmella.
melampodioides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 407 = S. Acmella.
melissaefolia, Salisb. Prod. 186 = S. Acmella.
multiflora, Otto, ex Sweet, Hort. Brit. ed. II. 306, nomen (Quid?).—Hab.?
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iv. 209.—Am. austr.
nitida, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 28.—Mexic.
Nuttallii, Torr. & Gray, Fl. N. Am. ii. 356 = repens.
oleracea, Linn. Syst. ed. XII. 534 = S. Acmella.
paniculata, Wall. Cat. n. 3186 = S. Acmella.
papposa, *Hemsl. Biol. Centr. Am. Bot.* ii. 193.—Mexic.
parvifolia, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 100.—Costa Rica.
peregrina, Blanco, Fl. Filip. ed. I. 622 = S. Acmella.
Poeppigii, *DC. Prod.* v. 622.—Chili.
portoricensis, *Spreng. Syst.* iii. 444.—Porto Rico.
Pseudo-acmella, Hook. & Arn. Bot. Beech. Voy. 150 = Peritylе microglossa.
Pseudo-acmella, Murr. Syst. ed. XIII. 610 = S. Acmella.
Pseudo-gummifera, Forst. f. ex DC. Prod. v. 502 = Petrobium arboreum.
pusilla, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 317 = stolonifera.
radicans, Jacq. Coll. iii. 229 = exasperata.
radicans, *Schrad. Cat. Hort. Gotting.* (1831) 5; *ex Prod.* v. 624.—Hab.?
ramosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 193.—Mexic.
repens, Hort. Par. & Berol. ex C. Koch, Ind. Sem. Hort. Berol. (1855) App. 14 = commutata.
repens, *Michx. Fl. Bor. Am.* ii. 131.—Am. bor.
repens, Spreng. ex DC. Prod. v. 625 = urens.
rhombifolia, *Zipp. ex Span. in Linnaea*, xv. (1841) 323, nomen.—Ins. Timor.
rugosa, *Blume, ex DC. Prod.* v. 625.—Java.
salivaria, Murr. in Comm. Gotting. vi. (1785) 3. t. 1 = alba.
Salzmanni, DC. Prod. v. 623 = S. Acmella.
sessilifolia, Hemsl. Biol. Centr. Am. Bot. ii. 193 = Jaegeria hirta.
sessilis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 50.—Peruv.
sphaerocephala, DC. Prod. v. 621 = arnicoides.
stenophylla, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 317 = arnicoides.

SPILANTHES:—
stolonifera, *DC. Prod.* v. 621.—Bras.
subhirsuta, *DC. l. c.* 622.—Mexic.
tenella, H. B. & K. Nov. Gen. et Sp. iv. 211 = S. Acmella.
tetrandra, Roxb. in Beats. Tracts St. Helena, App. 325 = Petrobium arboreum.
tinctoria, Lour. Fl. Cochinch. 484 = Adenostemma viscosum.
uliginosa, Sw. Prod. Veg. Ind. Occ. 110 = S. Acmella.
urens, *Jacq. Enum. Pl. Carib.* 28; *Select. Am.* 214. t. 126. f. 1.—Am. trop.
wedelioides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 318 = Eclipta elliptica.

SPILANTHUS, Linn. Mant. ii. 475 (1771) = praec.

SPILOCARPUS, Lem. Jard. Fleur. iv. (1854) Misc. 79. *BORAGINEAE*.
gracilis, *Lem. l. c.*—Cuba.

SPILOXENE, Salisb. Gen. Pl. Fragm. 44 (1866) = **Hypoxis**, Linn. (Amaryll.).
pavonina, Salisb. l. c. = H. stellata.
stellata, Salisb. l. c. = H. stellata.

SPINACEA, Schur, Enum. Pl. Transs. 574 (1866) = seq.

SPINACHIA, Hill, Hort. Kew. 421 (1768) = seq.

SPINACIA, [Tourn.] Linn. Syst. ed. I (1735). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 53.
divaricata, Turcz. ex Steud. Nom. ed. II. ii. 623, nomen = Atriplex lenticularis.
domestica, Borkh. in Rhein. Mag. i. (1793) 481 = oleracea.
fera, Linn. Sp. Pl. ed. II. 1493 = Atriplex lenticularis.
fera, Pall. Reise, i. 496 = Atriplex rosea.
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = oleracea.
inermis, Moench, Meth. 318 = oleracea.
littoralis, Jacq. Enum. Pl. Carib. 33 (Quid?).—Ind. occ.
minor, C. Koch, in Linnaea, xvii. (1843) 311 = tetrandra.
oleracea, *Linn. Sp. Pl.* 1027.—Ubique cult.
sessiliflora, Stokes, Bot. Mat. Med. iv. 537 = oleracea.
spinosa, Moench, Meth. 318 = oleracea.
tetrandra, Roxb. Hort. Beng. 72; Fl. Ind. iii. 771 = oleracea.
tetrandra, *Stev. ex Bieb. Fl. Taur. Cauc.* ii. 417.—Reg. Cauc.

SPINGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Quid?).

SPINIFEX, Linn. Mant. ii. 163 (1771). *GRAMINEAE*, Benth. & Hook. f. iii. 1109.
IXALUM, Forst. f. Prod. 92 (1786).
alterniflorus, Nees, in Lehm. Pl. Preiss. ii. 96 = longifolius.
dealbatus, *Nees & Meyen, ex Steud. Nom.* ed. II. ii. 623, nomen.—China.
elegans, *Buese, in De Vriese, Pl. Ind. Bat. Or.* 113.—Ins. Sundaic.
fragilis, R. Br. Prod. 198.—longifolius.
hirsutus, *Labill. Nov. Holl. Pl.* ii. 81. tt. 230, 231.—Austral.; Ins. Pacif.
inermis, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 292 = hirsutus.
longifolius, *R. Br. Prod.* 198.—Austral.
paradoxus, *Benth. in Hook. Ic. Pl.* tt. 1243, 1244.—Austral.
sericeus, R. Br. Prod. 198 = hirsutus.
squarrosus, *Linn. Mant.* ii. 300.—Ind. or.

SPINOVITIS, Romanet, in Compt. Rend. Par. xcii. (1881) 1096 = **Vitis**, Tourn. (Ampel.).
Davidi, Romanet, l. c. = V. Labrusca.

SPIRACANTHA, H. B. & K. Nov. Gen. et Sp. iv. 28. t. 313 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 237.
cornifera, Steud. Nom. ed. II. ii. 623, sphalm. = seq.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 29.—N. Granat.
denticulata, *Ernst, in Flora*, lvii. (1874) 213.—Venezuela.

SPIRADICLIS, Blume, Bijdr. 975 (1876). *RUBIACEAE*, Benth. & Hook. f. ii. 62.
PLEOTHECA, Wall. Cat. n. 6215 (1831 ?).
bifida, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 310.—Reg. Himal.
caespitosa, *Blume, Bijdr.* 975.—Burma ; Java.
cylindrica, *Benth. & Hook. f. Gen.* ii. 62.—Reg. Himal.

SPIRAEA, Linn. Syst. ed. I (1735). *ROSACEAE*, Benth. & Hook. f. i. 611.
ALIPENDULA, Neck. Elem. ii. 89 (1790).
ARUNCUS, Linn. Gen. ed. I. 310 (1737).
AWAYUS, Rafin. Sylva Tellur. 151 (1838).
BASILIMA, Rafin. New Fl. Am. iii. 75 (1836) ; Sylva Tellur. 152 (1838).
CHAMAEBATIARIA, Maxim. in Act. Hort. Petrop. vi. (1879) 225.
DRYMOPOGON, Rupp. Fl. Jen. ed. Hall. 124 (1745).
ELEIOSINA, Rafin. Sylva Tellur. 150 (1838).
ERIOGYNIA, Hook. Fl. Bor. Am. i. 255. t. 88 (1833).
FILIPENDULA, Tourn. ex Linn. Gen. ed. I. 145 (1737).
HOLODISCUS, Maxim. in Act. Hort. Petrop. vi. (1879) 253.
LUETKEA, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 130. t. 2.
SCHIZONOTUS, Lindl. ex Wall. Cat. n. 703 (1830).
SERICOTHECA, Rafin. Sylva Tellur. 152 (1838).
SIBIRAEA, Maxim. in Act. Hort. Petrop. vi. (1879) 116, in obs.
SORBARIA, A. Br. ex Aschers. Fl. Brandenb. 177 (1864).
THECANISIA, Rafin. New Fl. Am. ii. 38 (1836) ; Sylva Tellur. 152 (1838).
ULMARIA, [Tourn.] Hill, Hort. Kew. 213 (1768).
acutifolia, Willd. Enum. Hort. Berol. 540 = hypericifolia.
adiantifolia, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 642 = vaccinifolia.
alba, Du Roi, Harbk. Baumz. ii. 430 = salicifolia.
alpestris, *G. Don, Gen. Syst.* ii. 519.—Sibir.
alpina, *Pall. Fl. Ross.* i. I. 35, 78.—Sibir.
altaica, Pall. Reise, ii. 739 = laevigata.
altaiensis, Laxm. in Nov. Comm. Acad. Petrop. xv. (1771) 554. t. 29. f. 2 = laevigata.
ambigua, Pall. Fl. Ross. i. I. 36, in obs. = hypericifolia.
amena, Rafin. New Fl. Am. iii. 65 = salicifolia.
americana, Steud. Nom. ed. I. 805 = S. Aruncus.
amoena, Spae, in C. Morr. Ann. Soc. Bot. Gand, ii. (1846) 281. t. 72 = bella.
amurensis, Maxim. Prim. Fl. Amur. 90 (= *Physocarpa amurensis*).—Manchuria.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 9 = S. Filipendula.
angustifolia, *Otto & Dietr. Allg. Gartenz.* xiii. (1845) 249.—Hab. ?
angustiloba, Turcz. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 71 = lobata.
aquilegifolia, Pall. Reise, iii. 734 = hypericifolia.
arctica, Hort. ex Lavall. Enum. Arb. Segrez. 80 = dasyantha.
arcuata, *Hook. f. Fl. Brit. Ind.* ii. 325.—Reg. Himal.
argentea, W. Baxt. ex Loud. Hort. Brit. Suppl. ii. 674 = *nutans*.
argentea, Benth. Pl. Hartw. 82 = discolor.
argentea, C. Koch, Dendrol. i. 328 = media.
argentea, *Linn. f. Suppl.* 261.—N. Granat.
argunensis, Ledeb. Fl. Ross. ii. 18 = palmata, *Pall.*
arguta, Hoffmgg. Verz. Pfl. 114 = salicifolia.
arguta ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 494.
ariaefolia, Sm. in Rees, Cycl. xxxiii. n. 16 = discolor.
Aruncus, *Linn. Sp. Pl.* 490.—Reg. bor. temp.
Aruncus, Thunb. Fl. Jap. 211 = Astilbe japonica.
astilboides, T. Moore, in Gard. Chron. (1880) II. 114 ; Carr. in Rev. Hortic. (1884) 432 = S. Aruncus.
banatica. Janka, ex Just, Jahresb. iv. 1061 = ulmifolia.
barbata, Wall. Cat. n. 705 = Astilbe rivularis.
belgica, *Dum. Fl. Belg. Prod.* 98.—Belgium.
bella, *Sims, Bot. Mag.* t. 2426.—Reg. Himal.
Besseriana, *Sweet, Hort. Brit.* ed. II. 194.—Podolia.
bethlehemensis, Hort. ex C. Koch, Dendrol. i. 313 = salicifolia.
betulaefolia, Wats. Dendrol. Brit. i. t. 67 = corymbosa.
betulifolia, *Pall. Fl. Ross.* I. I. 33. t. 16.—As. et Am. bor.
Billardii, Hort. ex C. Koch, Dendrol. i. 312 = salicifolia.

SPIRAEA:—
blanda ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 496.
Blumei, *G. Don, Gen. Syst.* ii. 518.—Japon.
Boursierii, *Carr. in Rev. Hortic.* (1859) 520. f. 108.—Calif.
brachybotrys, Lange, in Bot. Tidsskr. xiii. (1882) 26 = canescens.
bracteata, *Rafin. New Fl. Am.* iii. 67.—Sibir.
bracteata ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 496.
brahuica, *Boiss. Fl. Orient.* ii. 690.—Ind. bor. occ. ; Beluchist.
brumalis, *Lange, in Bot. Tidsskr.* xiii. (1882) 28. (an hybr. ?).—Hab. ?
bullata, *Maxim. in Act. Hort. Petrop.* vi. (1879) 204. —Japon.
caerulescens, Poir. Encyc. vii. 350 = cantoniensis ?
caespitosa, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 418. —Am. bor. occ.
californica, Torr. in Emory, Notes Mil. Reconnois. 140 = Vauquelinia Torreyi.
callosa, Thunb. Fl. Jap. 209 = japonica.
callosa, Wall. Cat. n. 707 = bella.
camtschatica, *Pall. Fl. Ross.* i. I. 41.—Kamtschat. ; Ins. Aleut.
cana, *Waldst. & Kit. Pl. Rar. Hung.* iii. 252. t. 227. —Europ. austr.
canadensis, Fisch. ex Steud. Nom. ed. I. 805, nomen (Quid ?).—Am. bor. ?
canescens, *D. Don, Prod. Fl. Nep.* 227.—Reg. Himal.
cantoniensis, *Lour. Fl. Cochinch.* 322.—China.
capitata, Pursh, Fl. Am. Sept. i. 342 = Neillia opulifolia.
carnea, Steud. Nom. ed. I. 805 = salicifolia.
caroliniana, Marsh. Arbust. 146 = Neillia opulifolia.
carpinifolia, Willd. Enum. Hort. Berol. 540 = salicifolia.
ceanothifolia, Hornem. Hort. Hafn. ii. 466 = corymbosa.
chamaedrifolia, Blume, Bijdr. 1114 = Blumei.
chamaedrifolia, Cambess. in Ann. Sc. Nat. Sér. I. i. (1824) 362 = media.
chamaedrifolia, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. ii. 326 = gracilis.
chamaedrifolia, Jacq. Hort. Vindob. ii. 66. t. 140 = ulmifolia.
chamaedrifolia, Thunb. Fl. Jap. 210 = cantoniensis.
chamaedryfolia, Cham. & Schlecht. in Linnaea, ii. (1827) 2 = betulifolia.
chamaedryfolia, *Linn. Sp. Pl.* 489.—Europ. or. ; As. bor.
chinensis, Maxim. in Act. Hort. Petrop. vi. (1879) 193 = dasyantha.
ciliata, Rafin. New Fl. Am. iii. 66 = salicifolia.
cinerea ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 494.
coerulescens, Auct. = *caerulescens*.
conferta ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 495.
confusa, Regel & Koern. Ind. Sem. Hort. Petrop. (1857) 57 = media.
contorta, Stokes, Bot. Mat. Med. iii. 129 = S. Ulmaria.
coristantia ×, Schroed. in Gard. Chron. (1882) II. 108.
corymbosa, *Rafin. Préc. Découv.* 36.—Am. bor.
corymbosa, Roxb. Hort. Beng. 38 ; Fl. Ind. ii. 512 = cantoniensis.
crataegifolia, Link, Enum. Hort. Berol. ii. 40 = corymbosa.
crenata, Bess. ex DC. Prod. ii. 543 = Besseriana.
crenata, Cav. Desc. 503 = flabellata.
crenata, *Linn. Sp. Pl.* 489.—Europ. or. ; As. bor.
crenata, Muhl. Cat. 51 = salicifolia.
crenata, Pall. ex Bieb. Fl. Taur. Cauc. i. 392 = hypericifolia.
crenata, Thunb. Fl. Jap. 210 = Thunbergii.
crenifolia, *C. A. Mey. in Beitr. Pfl. Russ. Reich.* vi. 43.—Rossia.
cuneifolia, Borkh. Handb. Forstbot. ii. 1455 = salicifolia.
cuneifolia, Rafin. New Fl. Am. iii. 67 = Douglasii ?
cuneifolia, Wall. Cat. n. 699 = canescens.
cuneifolia, Wender. ex Steud. Nom. ed. II. ii. 624, nomen (Quid ?).—Hab. ?
cuspidata, *Hoffmgg. Verz. Pfl.* 114, nomen.—Hab. ?
dahurica, *Maxim. in Act. Hort. Petrop.* vi. (1879) 190. —As. bor.
dasyantha, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 97.—China.

SPIRAEA:—
daurica, Hort. Angl. ex Steud. Nom. ed. II. ii. 624, nomen (Quid ?).—Dahuria.
daurica, Rafin. New Fl. Am. iii. 69 = chamaedryfolia.
decumbens, *Koch, in Sturm, Deutschl. Fl.* Heft 62. —Austria.
denticulata, Rafin. New Fl. Am. iii. 71 = crenata.
denudata, J. & C. Presl, Fl. Čech. 101 = S. Ulmaria.
digitata, Willd. Sp. Pl. ii. 1061 = palmata, *Pall.*
discolor, *Pursh, Fl. Am. Sept.* i. 342.—Am. bor.
Douglasii, *Hook. Fl. Bor. Am.* i. 172.—Am. bor. occ.
dubia, *Steud. Nom.* ed. II. ii. 624.—Sibir.
dumosa, Nutt. ex Hook. Lond. Journ. Bot. vi. (1847) 217 = discolor.
elliptica, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
eximia, Hort. ex Regel, Ind. Sem. Hort. Petrop. (1866) 106 = salicifolia.
expansa, Wall. Cat. n. 702 = bella.
fastigiata, Wall. l. c. n. 708 = bella.
ferruginea, Rafin. New Fl. Am. iii. 63 = tomentosa.
Filipendula, *Linn. Sp. Pl.* 490.—Europ. ; As. bor.
filipendulina, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1091 = S. Filipendula.
fissa, Lindl. Bot. Reg. (1840) Misc. 73 = discolor.
flabellata, *Bertol. ex Guss. Pl. Rar.* 205.—Ital.
flexuosa, Fisch. ex Cambess. in Ann. Sc. Nat. Sér. I. i. (1824) 365 (err. typ. 265) = chamaedryfolia.
flexuosa, Rafin. New Fl. Am. iii. 64 = salicifolia.
flexuosa, Reichb. Fl. Germ. Excurs. 627 = decumbens.
floribunda, Hort. ex C. Koch, Dendrol. i. 306 = sorbifolia.
foliosa, Poir. Encyc. vii. 353 = media.
Fortunei, Planch. in Fl. des Serres, ix. (1853–54) 35. t. 871 = japonica.
Foxii, Hort. ex C. Koch, Dendrol. i. 329 = japonica.
Gieseleriana ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 495.
glabrata, Lange, in Bot. Tidsskr. xiii. (1882) 29 = japonica ?
glauca, Schultz, Prod. Fl. Starg. Suppl. i. 26 = S. Ulmaria.
glauca, Wall. Cat. sub n. 707 = bella.
glomerata, Rafin. New Fl. Am. iii. 63 = tomentosa.
gracilis, *Maxim. in Act. Hort. Petrop.* vi. (1879) 200. —Reg. Himal.
grandiflora, Hook. Bot. Mag. t. 4795 = Exochorda grandiflora.
grandiflora, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
grandiflora, Sweet, Hort. Brit. ed. II. 194 ; Lodd. Bot. Cab. t. 1988 = salicifolia.
Hacquetii, Fenzl, ex C. Koch, in Regel, Gartenfl. (1854) 400 = lancifolia.
heterophylla, Rafin. New Fl. Am. iii. 65 = salicifolia.
Hookeri, Hort. ex C. Koch, Dendrol. i. 332 = bella.
Hookeriana, Rafin. New Fl. Am. iii. 73 = salicifolia.
Humanni, Hort. ex C. Koch, Dendrol. i. 325 = cantonniensis.
Humboldtii, *Hort.—Cf. Gard. Chron.* (1884) I. 827.—Hab. ?
Humboldtii, *Spae, in C. Morr. Ann. Soc. Bot. Gand*, ii. (1846) 311.—Hab. ?
hydrangeaefolia, Hort.—Cf. Gard. Chron. (1877) I. 245 = splendens.
hypericifolia, Bieb. Fl. Taur. Cauc. i. 392 = crenata.
hypericifolia, *Linn. Sp. Pl.* 489 ; *DC. Fl. Fr. Suppl.* 645.—Rossia ; As. temp.
hypericifolia. Marsh. Arbust. 146 = salicifolia.
incarnata, Hort. ex C. Koch, Dendrol. i. 311 = salicifolia.
incisa, Hort. ex Reichb. Fl. Germ. Excurs. 627 ; et in Rev. Hortic. (1875) 160 = chamaedryfolia.
incisa, Thunb. Fl. Jap. 213 = Stephanandra flexuosa.
indica, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
inflexa, *Hort. ex C. Koch, in Regel, Gartenfl.* (1854) 405.—Hab. ?
integrifolia, Pall. Reise, iii. 317 = laevigata.
italica, Rafin. New Fl. Am. iii. 72 = hypericifolia ?
japonica, Cambess. in Ann. Sc. Nat. Sér. I. i. (1824) 389 = Kerria japonica.
japonica, *Linn. f. Suppl.* 262.—Reg. Himal. ; China ; Japon.
japonica, Makoy, Cat. (1836) 36 ; ex Otto & Dietr. Allg. Gartenz. v. (1837) 211 = Astilbe japonica.

SPIRAEA:—
japonica, Rafin. New Fl. Am. iii. 71 = Thunbergii.
japonica, Siebold, ex Blume, Bijdr. 1114 = cantoniensis.
kamaonensis, Hort. ex C. Koch, Dendrol. i. 331 = bella.
kamtschatica, Auct. = camtschatica.
kamtschatica, Wall. Cat. n. 704 = vestita.
Kirilowii, Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 81 = sorbifolia.
laciniata, Rafin. New Fl. Am. iii. 71, in obs. = crenata.
laevigata, *Linn. Mant.* ii. 244.—Sibir.
lanata, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
lanceolata, Borck. Handl. Forstbot. ii. 1457 = salicifolia.
lanceolata, Comm. ex Poir. Encyc. vii. 354 = cantoniensis.
lancifolia, *Hoffmgg. Preisverz. Pfl.* viii. (1825) *Nachtr.* 44; *ex Walp. Rep.* ii. 50.—Europ.
lasiocarpa, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 536 = crenifolia
latifolia, Borck. Handb. Forstbot. ii. 1871 = salicifolia.
laxiflora, Lindl. Bot. Reg. (1839) Misc. 60 = vaccinifolia.
leucantha, Lange, in Bot. Tidsskr. xiii. (1882) 30, adnot. = japonica ?
Lindleyana, Wall. Cat. n. 703 = sorbifolia.
lobata, *Jacq. Hort. Vindob.* i. 38.—Am. bor.
longigemmis, *Maxim. in Act. Hort. Petrop.* vi. (1879) 205.—China.
lucida, Dougl. ex Hook. Fl. Bor. Am. i. 172 = betulifolia.
magellanica, *Poir. Encyc.* vii. 350 (sp. dub.).—Reg. Magellan.
media, Pursh, Fl. Am. Sept. i. 342 = chamaedryfolia.
media, *F. Schmidt, Oestr. Baumz.* i. 53. t. 54.—As. bor.
Menziesii, Hook. Fl. Bor. Am. i. 173 = Douglasii.
mexicana, *Schiede, in Regel, Ind. Sem. Hort. Petrop.* (1857) 58.—Mexic.
micrantha, *Hook. f. Fl. Brit. Ind.* ii. 325.—Reg. Himal.
micropetala ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 495.
Millefolium, *Torr. in Pacif. Rail. Rep.* iv. 83. t. 5.—Calif.
mollis, *C. Koch & Bouché, in Regel, Gartenfl.* (1854) 404.—Sibir.
mongolica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 467.—China.
monogyna, Torr. in Ann. Lyc. N. York, ii. (1828) 194 = Neillia Torreyi.
montana, Hort. ex C. Koch, Ind. Sem. Hort. Berol. (1853) App. 12 = bella.
multifida, Ledeb. Fl. Ross. ii. 19 = pectinata.
multiflora ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 494.
nepalensis, Hort. ex C. Koch, Dendrol. i. 331, 332 = bella, canescens.
nepalensis, Lodd. ex Loud. Hort. Brit. 202, nomen (Quid ?).—Nepal.
nervosa, Franch. & Sav. Enum. Pl. Jap. ii. 331 = dasyantha.
Nicoudiertii, Bosse, Handb. Blumenz. ed. II. iv. 624 = crenata.
nicoviensis, Link, Enum. Hort. Berol. ii. 40 = crenifolia.
nivea, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
Nobleana, Hook. f. Bot. Mag. t. 5169 = Douglasii.
nutans, Hort. ex C. Koch, Dendrol. i. 332, 333 = canescens.
nutans, Royle, ex Steud. Nom. ed. II. ii. 624 (sp. dub.).—Ind. or.
oblongata, Wender. in Schrift. Marb. ii. (1831) 254 = chamaedryfolia.
oblongifolia, Ledeb. Fl. Ross. ii. 13, partim = hypericifolia.
oblongifolia, Regel, ex Maxim. in Act. Hort. Petrop. vi. (1879) 183 = alpina.
oblongifolia, Waldst. & Kit. Pl. Rar. Hung. iii. 261. t. 235 = media.
obovata, Rafin. ex DC. Prod. ii. 544 = salicifolia.
obovata, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 541 = hypericifolia.
obtusiloba, Opiz, Seznam, 100, in syn. = S. Ulmaria.
SPIRAEA:—
occidentalis, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 192.—Am. bor. occ.
odorata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 589 = S. Ulmaria.
opulifolia, Linn. Sp. Pl. 489 = Neillia opulifolia.
opulifolia, Pall. Reise, iii. 230 = trilobata.
orbicularis, Hoffmgg. Verz. Pfl. 114 = hypericifolia.
ostryfolia, Rafin. New Fl. Am. iii. 69 = betulifolia.
ovalifolia, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
ovata, Rafin. New Fl. Am. iii. 65 = salicifolia.
ovata, *Van Houtte, ex Regel, Gartenfl.* (1852) 229.—Hab. ?
oxyodon ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 495.
pachystachys, Hort. ex C. Koch, Dendrol. i. 314 = Douglasii.
Pallasii, G. Don, Gen. Syst. ii. 520 = grandiflora.
palmata, Murr. Syst. ed. XIII. 393 = lobata.
palmata, *Pall. Reise*, iii. 735.—Sibir.; China.
palmata, *Thunb. Fl. Jap.* 212.—Japon.
palustris, Salisb. Prod. 364 = S. Ulmaria.
paniculata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Aruncus.
paniculata, Willd. Sp. Pl. ii. 1055; G. Don, Gen. Syst. ii. 520 = salicifolia.
parvifolia, *Benth. Pl. Hartw.* 36.—Mexic.
parvifolia, Bertol. in Mem. Acad. Sc. Bologn. Ser. II. iv. (1864) 312 = gracilis.
parvifolia, Rafin. New Fl. Am iii. 63 = tomentosa.
pauciflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 414 = Neillia opulifolia.
pectinata, *Torr. & Gray, Fl. N. Am.* i. 417.—Am. bor. occ.
pikoviensis, Bess. Enum. Pl. Volh. 46 = crenifolia.
pilosa, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 282.—Turkestan.
pinnata, Blume, Cat. Gew. Buitenz. 76 = Weinmannia Blumei, *Planch.*
pinnata, Moench, Meth. 663 = sorbifolia.
pinnifolia, Walp. Rep. ii. 50, sphalm. = prunifolia.
procumbens, Kirch. in C. Koch & Fint. Wochenschr. ii. (1859) 358 = betulifolia ?
prostrata, *Maxim. in Act. Hort. Petrop.* vi. (1879) 184.—China.
prostrata, Schur, Enum. Pl. Transs. 181 = ulmifolia.
pruinosa, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
prunifolia, *Sieb. & Zucc. Fl. Jap.* i. 131.—Japon.
pubescens, *Cav. Desc.* 504.—Hab. ?
pubescens, DC. Fl. Fr. Suppl. 546 = S. Filipendula.
pubescens, Lindl. in Journ. Hort. Soc. ii. (1847) 157 = dasyantha.
pubescens, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 190.—China.
pulchella, Kunze, in Linnaea, xx. (1847) 60 = bella ?
pulverulenta, Rafin. New Fl. Am. iii. 71 = crenata.
quinqueloba, Baumg. Enum. Stirp. Transs. ii. 47 = S. Ulmaria.
racemosa ×, Schroed. in Gard. Chron. (1882) II. 108.
Reevesiana, Lindl. Bot. Reg. (1844) t. 10 = cantoniensis.
Regeliana, Hort. ex C. Koch, Dendrol. i. 314 = Douglasii.
repens, Rafin. New Fl. Am. iii. 70 = betulifolia.
reticulata, Rafin. l. c. 72 = hypericifolia ?
rhamnifolia, Wall. Cat. n. 700 = vaccinifolia.
rhodoclada, *Leresche & Levier, Deux Excurs. Bot.* 26.—Hispan.
ribifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 414 = Neillia opulifolia.
rosea, Rafin. New Fl. Am. iii. 62 = tomentosa.
rotundifolia, Hort. ex C. Koch, Dendrol. i. 327 = trilobata.
rotundifolia, *Lindl. Bot. Reg.* (1840) *Misc.* 72.—Reg. Himal.
Roxburghiana, Wall. ex Voigt, Hort. Suburb. Calc. 197 = cantoniensis.
ruberrima, Regel, Gartenfl. iii. (1854) 398 = bella.
rupestris, Hort. ex C. Koch, Dendrol. i. 326 = Blumei.
ruscifolia, Hort. ex C. Koch, l. c. 333 = canescens.
salicifolia, *Linn. Sp. Pl.* 489.—Reg. bor. trop.
salicifolia, Schang. in Pall. N. Nord. Beitr. vi. 36 = chamaedryfolia.
sanssouciana, C. Koch, Berl. Allg. Gartenz. (1857) 214 = Douglasii.
sawranica, Bess. Enum. Pl. Volh. 59 = crenata.
SPIRAEA:—
saxatilis, *Turcz. ex Bess. in Flora*, xvii. (1834) I. *Beibl.* 11.—Sibir. baical.
Schinabecki ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 495.
Schrenkiana, C. A. Mey. Ind. Sem. Hort. Petrop. ix. 96 = Spiraeanthus Schrenckianus.
sericea, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 591 = media.
serratifolia, Hort. ex C. Koch, Dendrol. i. 333 = canescens.
sibirica, Hort. ex Steud. Nom. ed. II. ii. 624 = hypericifolia.
sibirica, Rafin. New Fl. Am. iii. 64 = salicifolia.
sorbifolia, *Linn. Sp. Pl.* 490.—Reg. Himal.; As. bor.
sororia, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 600 = corymbosa.
speciosa, Hort. ex C. Koch, Dendrol. i. 325 = cantoniensis.
speciosa, *Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 41.—Java.
splendens, *Hort.—Cf. Gard. Chron.* (1877) I. 245; (1883) II. 12.—Hab. ?
stellata, Wall. Cat. sub n. 707 = bella.
stipulacea, Pursh, Fl. Am. Sept. i. 343 = Gillenia stipulacea.
stipulata, Muhl. ex Willd. Enum. Hort. Berol. 542 = Gillenia stipulacea.
subcordata, *Lenné & C. Koch, in Regel, Gartenfl.* (1854) 402.—Am. bor.
syringaeflora, Hort. ex C. Koch, Dendrol. i. 329 = japonica.
tenuissima, *Carr. in Rev. Hortic.* (1867) 140.—Hab. ?
thalictroides, Pall. Fl. Ross. i. 1. 34. t. 18 = hypericifolia.
Thunbergii, *Siebold, ex Blume, Bijdr.* 1115.—China; Japon.
thyrsiflora, C. Koch, Dendrol. i. 307 = Neillia thyrsiflora.
tobolskia, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 202, nomen.—Sibir.
tomentosa, *Linn. Sp. Pl.* 489.—Am. bor.
tomentosa, Rafin. New Fl. Am. iii. 62 = Douglasii.
trifoliata, Linn. Sp. Pl. 490 = Gillenia trifoliata.
triloba, Murr. Syst. ed. XIII. 393 = seq.
trilobata, *Linn. Mant.* ii. 224.—Sibir.; China.
triternata, Wall. Cat. n. 706 = S. Aruncus.
tuberosa, Salisb. Prod. 364 = S. Filipendula.
Ulmaria, *Linn. Sp. Pl.* 490.—Europ.; As. bor.
ulmarioides, Bory, Voy. Souterr. 124 = S. Ulmaria.
ulmifolia, *Scop. Fl. Carn.* ed. II. i. 349. t. 22.—Europ. or.; Sibir.
undulata, Borkh. Versuch Forstbot. Beschr. Holzart. 65 = salicifolia.
unguiculata, Dulac, Fl. Hautes-Pyr. 315 = S. Ulmaria.
uralensis, Sweet, Hort. Brit. ed. I. 149 = hypericifolia.
uratensis, *Franch. Pl. Davidian.* i. (1884) 107.—Mongol.
vaccinifolia, *D. Don, Prod. Fl. Nep.* 227.—Reg. Himal.
vaccinifolia, Lodd. Bot. Cab. t. 1403 = gracilis.
Vanhouttei ×, Zabel, in Wittm. Garten-Zeit. iii. (1884) 496.
venusta, *Hort. ex Otto & A. Dietr. Allg. Gartenz.* xiii. (1845) 250.—Hab. ?
venustula, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 16 = vaccinifolia.
versifolia, Rafin. New Fl. Am. iii. 68 = chamaedryfolia.
vestita, *Wall. ex G. Don, Gen. Syst.* ii. 521.—Reg. Himal.
virgata, Rafin. New Fl. Am. iii. 72 = salicifolia.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 588 = S. Filipendula.
Zuccarinii, Siebold, ex Miq. Ann. Bot. Mus. Lugd. Bat. iii. 32 = japonica.

SPIRAEANTHEMUM, A. Gray, Bot. U. St. Expl. Exped. i. 666. t. 83 (1854). *SAXIFRAGEAE*, Benth. & Hook. f. i. 650.
austro-caledonicum, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 74; *et in Ann. Sc. Nat.* Sér. V. i. (1864) 374.—N. Caled.
caledonicum, C. Muell. in Walp. Ann. vii. 910, sphalm. = praec.
densiflorum, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 74; *et in Ann. Sc. Nat.* Sér. V. i. (1864) 374.—N. Caled.

SPIRAEANTHEMUM:—
Graeffei, *Seem. Fl. Vit.* 111.—Ins. Fiji.
Katakata, *Seem. in Bonplandia*, x. (1862) 36, nomen; *et l. c.*—Ins. Fiji.
Macgillivrayi, *Seem. Fl. Vit.* 111.—Ins. N. Hebrid.
samoënse, *A. Gray, Bot. U. St. Expl. Exped.* 667. t. 83.—Ins. Samoa.
undulatum, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 342.—N. Caled.
vitiense, *A. Gray, Bot. U. St. Expl. Exped.* 669. t. 83.—Ins. Fiji.

SPIRAEANTHUS, Maxim. in Act. Hort. Petrop. vi. (1879) 226. *ROSACEAE.*
Schrenckianus, *Maxim. l. c.* 227.—Sibir. altaic.

SPIRAEOPSIS, Miq. Fl. Ind. Bat. i. 1. 719 (1855). *SAXIFRAGEAE*, Benth. & Hook. f. i. 653.
DIRHYNCHOSIA, Blume, Mél. Bot.; ex Flora, xli. (1858) 254.
celebica, *Miq. Fl. Ind. Bat.* i. 1. 719.—Ins. Celebes.

SPIRAGYNE, Neck. Elem. ii. (1790) 14 = **Gentiana**, Tourn.

SPIRALEPIS, D. Don, in Mem. Wern. Soc. v. (1826) 551 = **Leontonyx**, Cass. (Compos.).
declinata, D. Don, in Loud. Hort. Brit. 340 = Helichrysum declinatum.
glomerata, D. Don, in Mem. Wern. Soc. v. (1826) 552 = L. coloratus.
modesta, G. Don, in Loud. Hort. Brit. 340 = Helipterum gnaphaloides.
rotundifolia, D. Don, in Mem. Wern. Soc. v. (1826) 552 = Helichrysum rotundifolium.
squarrosa, D. Don, l. c. 551 = L. tomentosus.
tincta, D. Don, l. c. 552 = L. tomentosus.

SPIRANTHERA, Boj. Hort. Maurit. 226 (1837) = **Ipomoea**, Linn. (Convolv.).
peltata, Boj. l. c. = I. peltata.
pentaphylla, Boj. l. c. = I. pentaphylla.
Turpethum, Boj. l. c. = I. Turpethum.

SPIRANTHERA, Hook. Bot. Mag. sub t. 3523 (1836) = **Pronaya**, Hueg. (Pittosp.).
Fraseri, Hook. l. c. = P. elegans.

SPIRANTHERA, Rafin. Fl. Tellur. iv. 31 (1836) = **Eustrephus**, R. Br. (Liliac.).
ovata, Rafin. l. c. = E. latifolius.

SPIRANTHERA, A. St. Hil in Bull. Soc. Philom. (1823) 130. *RUTACEAE*, Benth. & Hook. f. i. 283.
TERPNANTHUS, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 152. t. 19.
odoratissima, *A. St. Hil. in Bull. Soc. Philom.* (1823) 130.—Bas.

SPIRANTHES, Rich. in Mém. Mus. Par. iv. (1818) 50. *ORCHIDEAE*, Benth. & Hook. f. iii. 596.
ARISTOTELEA, Lour. Fl. Cochinch. 522 (1790).
CYCLOPOGON, Presl, Rel. Haenk. i. 93. t. 13 (1827).
CYCLOPTERA, Endl. Ench. 113 (1841).
DOTHILIS, Rafin. Fl. Tellur. ii. 60 (1836).
GYROSTACHIS, Pers. Syn. ii. 511 (1807), nomen prius.
HELICTONIA, Ehrh. Beitr. iv. 148 (1789).
IBIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 291.
MONUSTES, Rafin. Fl. Tellur. ii. 87 (1836).
NARICA, Rafin. l. c. (1836).
SACOILA, Rafin. l. c. 86 (1836).
SARCOGLOTTIS, Presl, Rel. Haenk. i. 95. t. 15 (1827).
SAUROGLOSSUM, Lindl. Bot. Reg. t. 1618 (1833).
STENORYNCHUS, Rich. in Mém. Mus. Par. iv. (1818). 59.
STRATEUMA, Rafin. Fl. Tellur. ii. 89 (1836).
SYNASSA, Lindl. Bot. Reg. sub t. 1618 (1833).
actinosophila, *Rodrig. Orch. Nov.* ii. 286.—Bras.
acutata, *Reichb. f. & Warm. in Otia Bot. Hamb.* 84.—Bras.
aestivalis, Oakes, Cat. Verm. Pl. 28 = latifolia, *Torr.*
aestivalis, *Rich. in Mém. Mus. Par.* iv. (1818) 58.—Europ.; Afr. bor.
africana, Lindl. in Bot. Reg. sub t. 823 = Habenaria spiralis.

SPIRANTHES:—
aguacatensis, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Costa Rica.
albescens, Rodrig. Orch. Nov. i. 186 = micrantha.
Allemanii, Rodrig. l. c. ii. 288 = picta?
alpestris, *Rodrig. l. c.* i. 184.—Bras.
amoena, Spreng. Syst. iii. 708 = australis.
apiculata, *Lindl. Gen. et Sp. Orch.* 474.—Ind. occ.
aprica, *Lindl. l. c.* 469.—Bras.
argirifolia, *Rodrig. Orch. Nov.* i. 183.—Bras.
Arrabidae, *Warm. in Kjoeb. Vidensk. Meddel.* (1884) 94.—Bras.
assurgens, *Reichb. f. Beitr. Orch. Centr. Am.* 66.—Guatemala.
aurantiaca, *Hemsl. Biol. Centr. Am. Bot.* iii. 300.—Mexic.
australis, *Lindl. in Bot. Reg.* sub t. 823.—As. temp. et trop.; Australas.
autumnalis, *Rich. in Mém. Mus. Par.* iv. (1818) 59.—Europ.; As. temp.
balanophorostachya, *Reichb. f. & Warm. in Otia Bot. Hamb.* 84.—Bras.
Beckii, Lindl. Gen. et Sp. Orch. 472 = gracilis.
bicolor, *Griseb. Fl. Brit. W. Ind.* 641.—Ind. occ., Bras.
bicolor, *Lindl. in Bot. Reg.* sub t. 823.—Ins. Trinit.
bonariensis, *Lindl. Gen. et Sp. Orch.* 475.—Bras.; Reg. Argent.
bracteosa, *Lindl. Bot. Reg.* t. 1934.—Bras.
brevifolia, *Chapm. Fl. S. U. St.* 462.—Am. bor.
brevilabris, *Lindl. Gen. et Sp. Orch.* 471.—Am. bor.
calophylla, *Rodrig. Orch. Nov.* i. 182.—Bras.
camporum, *Lindl. Gen. et Sp. Orch.* 473.—Uruguay.
ceracifolia, *Rodrig. Orch. Nov.* ii. 285.—Bras.
cerina, *Lindl. Bot. Reg.* (1842) *Misc.* 20.—Guatemala.
cernua, Fleisch. & Lindem. Fl. Deutsch. Ostseeprov. 309 (Quid?).—Europ.
cernua, *Rich. in Mém. Mus. Par.* iv. (1818) 59.—Am. bor.
chilensis, *A. Rich. in C. Gay, Fl. Chil.* v. 475.—Chili.
chloreaeformis, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 33.—Mexic.
chloroleuca, *Rodrig. Orch. Nov.* i. 181.—Bras.
chlorops, *Reichb. f. Xenia Orch.* iii. 20.—Bolivia.
cinnabarina, *Hemsl. Biol. Centr. Am. Bot.* iii. 300.—Mexic.
colorans, Hemsl. l. c., sphalm. = seq.
colorata, *N. E. Br. in Gard. Chron.* (1883) i. 210.—Mexic.
colorata, *Hassk. Cat. Hort. Bog. Alt.* 47.—Java.
comosa, *Reichb. f. in Linnaea*, xviii. (1844) 408.—Mexic.
congesta, *Lindl. in Bot. Reg.* sub t. 823.—Sibir.
costaricensis, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Costa Rica.
crispata, Zoll. & Mor. Syst. Verz. Zoll. 89 = australis.
cuculligera, *Reichb. f. & Warm. in Otia Bot. Hamb.* 85.—Bras.
cuspidata, *Lindl. Gen. et Sp. Orch.* 471.—Bras.
decipiens, Hook. Fl. Bor. Am. ii. 203 = Goodyera Menziesii.
densa, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 79 = australis.
diaphana, *Lindl. Bot. Reg.* (1844) *Misc.* 12.—Mexic.?
dilatata, *Lindl. Gen. et Sp. Orch.* 474.—Uruguay.
diuretica, *Lindl. in Quart. Journ. R. Inst.* N. S. i. (1827) 54.—Chili.
elata, *Rich. in Mém. Mus. Par.* iv. (1818) 59.—Am. trop.
Eldorado, *Linden & Reichb. f. in Bot. Zeit.* xv. (1857) 157.—Bras.
ensifolia, *Reichb. f. in Bonplandia*, iv. (1856) 211.—Am. centr.
epiphyta, *Rodrig. Orch. Nov.* i. 189.—Bras.
Esmeralda, *Linden & Reichb. f. in Hamb. Gartenz.* xviii. (1862) 36.—Bras.
Eugenii, *Reichb. f. & Warm. in Otia Bot. Hamb.* 84.—Bras.
euphlebia, *Reichb. f. in Bot. Zeit.* xli. (1883) 13.—Bras.
flexuosa, Lindl. in Bot. Reg. sub t. 823 = australis.
Funckiana, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
Galeottiana, *A. Rich. l. c.*—Mexic.
Gardneri, Lindl. in Hook. Lond. Journ. Bot. i. (1842) 190 = Stenoptera ananassocomos.
gemmipara, Lindl. Syn. Brit. Fl. ed. I. 257 = Romanzoffiana.
glauca, Rafin. Fl. Tellur. ii. 88 = autumnalis.
gracilis, *Beck, Bot. N. & M. St.* 343; *Hook. Fl. Bor. Am.* ii. 202. t. 203.—Am. bor.

SPIRANTHES:—
gracilis, Hassk. Cat. Hort. Bog. Alt. 47 = Chlorosa gracilis.
graminea, *Lindl. in Benth. Pl. Hartw.* 25.—Am. bor.; Mexic.
grandiflora, Lindl. in Bot. Reg. t. 1043 = picta.
grandis, *Hassk. Cat. Hort. Bog. Alt.* 47.—Java.
gutturosa, *Reichb. f. Beitr. Orch. Centr. Am.* 67.—Am. centr.
hemichroa, *Lindl. Gen. et Sp. Orch.* 473.—Am. centr.
hirta, *Lindl. l. c.* 476.—Peruv.
homalogastra, *Reichb. f. & Warm. in Otia Bot. Hamb.* 84.—Bras.
Hostmanni, *Reichb. f. ex Griseb. Fl. Brit. W. Ind.* 640.—Ins. Trinit.; Guian.
hyemalis, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
hypnophila, *Rodrig. Orch. Nov.* i. 187.—Bras.
hysterantha, *Rodrig. l. c.* 188.—Bras.
icmadophila, *Rodrig. in Rev. Engenh.* 144; *ex eod. Orch. Nov.* ii. 286.—Bras.
inaequilatera, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 7. t. 110.—Peruv.
indica, Lindl. ex Steud. Nom. ed. II. ii. 625 = australis.
lanuginosa, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
latifolia, *A. Rich. & Gal. l. c.*—Mexic.
latifolia, *Torr. ex Lindl. Gen. et Sp. Orch.* 467.—Am. bor.
laxiflora, Rafin. Fl. Tellur. ii. 88 = tortilis.
Lindleyana, *Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol.* 26. t. 11.—Am. austr.
lineata, *Lindl. Gen. et Sp. Orch.* 471.—Bras.
lithophila, *Rodrig. Orch. Nov.* i. 190.—Bras.
Llaveana, *Lindl. in Benth. Pl. Hartw.* 73.—Mexic.
lobata, *Lindl. Bot. Reg.* (1844) *Misc.* 11.—Mexic.
longepetiolata, *Reichb. f. Beitr. Orch. Centr. Am.* 67.—Costa Rica.
longibracteata, *Rodrig. Orch. Nov.* i. 185.—Bras.
longifolia, *Lindl. Gen. et Sp. Orch.* 476; *Hassk. Cat. Hort. Bog. Alt.* 47.—Java.
longilabris, *Lindl. l. c.* 467.—Am. bor.
longispicata, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 78 = australis.
lupulina, *Hemsl. Biol. Centr. Am. Bot.* iii. 301.—Mexic.
luteo-alba, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
macrantha, *Reichb. f. in Linnaea*, xix. (1847) 378.—Bras.
macrophylla, Spreng. Syst. iii. 708 = Herminium congestum.
macropoda, *Rodrig. Orch. Nov.* i. 186.—Bras.
macrostachya, Poepp. & Endl. Nov. Gen. et Sp. ii. 8. t. 110 a–k = Stenoptera peruviana.
madrensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 301.—Mexic.
Mandonii, *Reichb. f. Xenia Orch.* iii. 21.—Bolivia.
margaritifera, Linden & Reichb. f. in Gard. Chron. (1866) 219 = S. Esmeralda.
mechoacana, *Hemsl. Biol. Centr. Am. Bot.* iii. 301.—Mexic.
micrantha, *Rodrig. Orch. Nov.* i. 183.—Bras.
minutiflora, *Reichb. f. in Bonplandia*, ii. (1854) 11.—Peruv.
minutiflora, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
monophylla, Spreng. Syst. iii. 709 = Herminium gramineum.
montana, *Hemsl. Biol. Centr. Am. Bot.* iii. 301.—Guatemala.
montana, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
multiflora, *Rodrig. Orch. Nov.* ii. 287.—Bras.
neuroptera, Reichb. f. & Warm. in Otia Bot. Hamb. 85 = simplex.
Novae-Zelandiae, Hook. f. Fl. N. Zel. i. 243 = australis.
novofriburgensis, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Bras.* 148.—Bras.
Nuil, Rich. in Mém. Mus. Par. iv. (1818) 59 = diuretica.
ochracea, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 33.—Mexic.
odorata, *Lindl. Gen. et Sp. Orch.* 467.—Am. bor.
oestivalis, Boiss. Fl. Orient. v. 91, sphalm. = aestivalis.
oestrifera, *Reichb. f. & Warm. in Otia Bot. Hamb.* 86.—Bras.

SPIRANTHES:—
orchioides, *A. Rich. Fl. Cub. Fanerog.* ii. 252.—Am. trop.
ornithocephala, *Rodrig. Orch. Nov.* ii. 288.—Bras.
orthosepala, *Reichb. f. & Warm. in Otia Bot. Hamb.* 85.—Bras.
ovalis, *Lindl. Gen. et Sp. Orch.* 466.—Texas.
paraguayensis, *Reichb. f. in Linnaea*, xxv. (1852) 230.—Paraguay.
parasitica, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
parviflora, *Hassk. Cat. Hort. Bog. Alt.* 47.—Java.
parviflora, Lindl. in Bot. Reg. sub t. 823 = australis.
pauciflora, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32; *Hemsl. Biol. Centr. Am. Bot.* iii. 302.—Mexic.
Pavonii, *Reichb. f. in Bonplandia*, iv. (1856) 211.—Peruv.
peruviana, *Presl, Rel. Haenk.* i. 93.—Peruv.
Petola, Hassk. Cat. Hort. Bog. Alt. 47 = Macodes Petola.
picta, *Lindl. in Bot. Reg.* sub t. 823.—Am. trop.
plantaginea, *Lindl. Gen. et Sp. Orch.* 468.—Peruv.
plantaginea, Spreng. Syst. iii. 708 = Microstylis congesta.
plantaginea, Torr. Fl. N. York, ii. 284 = latifolia, *Torr.*
polyantha, *Reichb. f. in Linnaea*, xviii. (1844) 408.—Mexic.
porrifolia, *Lindl. Gen. et Sp. Orch.* 467.—Calif.
prasophylla, *Reichb. f. Beitr. Orch. Centr. Am.* 65.—Guatemala.
Preslii, *Lindl. Gen. et Sp Orch.* 470.—Peruv.
pterygantha, *Reichb. f. & Warm. in Otia Bot. Hamb.* 86.—Bras.
pubens, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 33.—Mexic.
pubescens, Rodrig. Orch. Nov. i. 185 = bicolor, *Griseb.*
pudica, Lindl. Coll. Bot. t. 30; Spreng. Syst. iii. 708 = australis.
pusilla, *Miq. Fl. Ind. Bat.* iii. 722.—Sumatra.
pyramidalis, *Lindl. Gen. et Sp. Orch.* 473.—Guatemala.
quadridentata, *Lindl. in Bot. Reg.* sub t. 823.—Guian.
ramentacea, *Lindl. in Ann. Nat. Hist.* Ser. I. iv. (1840) 384.—Mexic.
Romanzoffiana, *Cham. & Schlecht. in Linnaea*, iii. (1828) 32.—As. bor. or.; Am. bor.; Hibern.
rosulata, *Lindl. Bot. Reg.* (1843) *Misc.* 60.—Guatemala.
rufescens, *Fisch. ex Klotzsch, in Otto & Dietr. Allg. Gartenz.* x. (1842) 107, in syn.—Bras.
rupestris, *Lindl. Gen. et Sp. Orch.* 474.—Reg. Argent.
rupestris, *Rodrig. Orch. Nov.* i. 189.—Bras.
saccata, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
sagittata, *Reichb. f. & Warm. in Otia Bot. Hamb.* 85.—Bras.
saltensis, *Griseb. in Goett. Abh.* xxiv. (1879) 338.—Reg. Argent.
sancta, *Reichb. f. & Warm. in Otia Bot. Hamb.* 85.—Bras.
sarcoglossa, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 31.—Mexic.
sceptrodes, *Reichb. f. in Bonplandia*, iii. (1855) 214.—Guatemala.
Schaffneri, *Reichb. f. in Linnaea*, xxviii. (1856) 382.—Mexic.
Scopulariae, *Reichb. f. in Bonplandia*, ii. (1854) 11.—Venezuela.
sellilabris, *Griseb. in Goett. Abh.* xxiv. (1879) 338.—Reg. Argent.
simplex, *A. Gray, Bot. N. U. St.* ed. V. 506.—Am. bor.
simplex, *Griseb. Fl. Brit. W. Ind.* 641.—Ind. occ.
Smithii, *Reichb. f. in Gard. Chron.* (1868) 842.—Costa Rica; Am. centr.
speciosa, *Lindl. Gen. et Sp. Orch.* 475.—Peruv.
speciosa, *A. Rich. Fl. Cub. Fanerog.* ii. 252.—Cuba.
spiralis, C. Koch, in Linnaea, xxii. (1849) 290 = autumnalis.
strateumatica, Lindl. in Bot. Reg. sub t. 823 = Zeuxine sulcata.
Stylites, *Lindl. in Journ. Linn. Soc.* i. (1857) 178.—China.
sulphurea, *Hemsl. Biol. Centr. Am. Bot.* iii. 302.—Mexic.
tenuis, *Lindl. Gen. et Sp. Orch.* 466.—Guian.
Thelymitra, *Reichb. f. Beitr. Orch. Centr. Am.* 66.—Guatemala.

SPIRANTHES:—
tortilis, Beck, Bot. N. & M. St. 343 = gracilis.
tortilis, *Rich. in Mém. Mus. Par.* iv. (1818) 59.—Ind. occ.
transversalis, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
trilineata, *Lindl. in Benth. Pl. Hartw.* 94.—Guatemala.
truncata, *Lindl. Gen. et Sp. Orch.* 470.—Bras.
uliginosa, *Rodrig. Orch. Nov.* i. 190.—Bras.
umbrosa, *Rodrig. l. c.* 188.—Bras.
unalascensis, Spreng. Syst. iii. 708 = Herminium congestum.
vaginata, *Lindl. Gen. et Sp. Orch.* 472.—Peruv.
vernalis, *Engelm. et A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 236.—Am. bor.
villosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 8.—Bras.
violacea, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 32.—Mexic.
Warmingii, *Reichb. f. & Warm. Otia Bot. Hamb.* 84.—Bras.
Weirii, *Reichb. f. in Gard. Chron.* (1870) 923.—Bras.
Wightiana, Lindl. in Wall. Cat. n. 7378 = australis.

SPIRANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

SPIRASTIGMA, L'Hérit. ex Schult. f. Syst. vii. p. lxx (1830) = **Pitcairnia**, L'Hérit. (Bromel.).

SPIREA, Pall. Reise, ii. App. 44 (1777), sphalm. = **Spiraea**, Linn. (Rosac.).

SPIREMA, Benth. in Journ. Linn. Soc. xv. (1875) 10, sphalm. = SPEIREMA, Hook. f. & Thoms. = **Pratia**, Gaudich. (Campan.).

SPIRIDANTHUS, Fenzl, in Endl. Gen. Suppl. ii. 105 (1842) = **Monolopia**, DC. (Compos.).

SPIRILLUS, J. Gay, in Compt. Rend. Acad. Sc. xxxviii. (1854) 703 = **Potamogeton**, Tourn. (Naiad.).

SPIROCARPUS, Opiz, Seznam, 93 (1852) = **Medicago**, Tourn. (Legumin.).
minimus, Opiz, l. c. = M. minima.

SPIROCHAETA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 166 = **Elephantopus**, Linn. (Compos.).
Funckii, Turcz. l. c. 167.—N. Granat.

SPIROCONUS, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 576 = **Trichodesma**, R. Br. (Borag.).
glaucus, Stev. l. c. 577 = T. molle.

SPIRODELA, Schleid. in Linnaea, xiii. (1839) 391 = **Lemna**, Linn.
atropurpurea, Montand. Guide Bot. 309 = L. polyrrhiza.
oligorrhiza, Hegelm. Lemnac. 147 = L. oligorrhiza.
polyrrhiza, Schleid. in Linnaea, xiii. (1839) 392 = L. polyrrhiza.

SPIROLOBA, Rafin. Sylva Tellur. 119 (1838) = **Pithecolobium**, Mart. (Legumin.).
bigemina, Rafin. l. c. = P. bigeminum.
fastuosa, Rafin. l. c. = Inga fastuosa.
Unguis, Rafin. l. c. = P. Unguis-cati.

SPIROLOBIUM, Orb. Sert. Patag. t. 13 (1839) = **Prosopis**, Linn. (Legumin.).
australe, Orb. l. c. 13 = P. strombulifera.

SPIRONEMA, Hochst. in Flora, xxv. (1842) 226 = **Clerodendron**, Linn. (Verben.).
myricoides, Hochst. l. c. = C. myricoides.

SPIRONEMA, Lindl. in Bot. Reg. (1840) t. 47. *COMMELINACEAE*, Benth. & Hook. f. iii. 854.
fragrans, *Lindl. l. c.*—Mexic.
orthandrum, Lindb. in Act. Soc. Sc. Fenn. x. (1871) 127. t. 4 = fragrans.
Robbinsii, Wright, in Sauv. Fl. Cub. 158 = Callisia repens.
Warszewiczianum, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 302 = Tradescantia Warszewicziana.

SPIRONEMA, Rafin. Fl. Tellur. iv. 92 (1836) = **Cassytha**, Linn. (Laurin.).
aphylla, Rafin. l. c. 93 = C. filiformis?

SPIROPODIUM, F. Muell. Fragm. i. 33 (1858) = **Pluchea**, Cass. (Compos.).
baccharoides, F. Muell. l. c. 34 = P. baccharoides.

SPIRORHYNCHUS, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 159. *CRUCIFERAE*, Benth. & Hook. f. i. 95.
sabulosus, *Kar. & Kir. l. c.* 160.—Soongar.

SPIROSPATHA, Rafin. Fl. Tellur. iv. 8 (1836) = **Homalomena**, Schott (Aroid.).
occulta, Rafin. l. c. = H. aromatica.

SPIROSPERMUM, Thou. Nov. Gen. Madag. 19 (1806). *MENISPERMACEAE*, Benth. & Hook. f. i. 39; 962.
penduliflorum, *DC. Syst.* i. 515.—Madag.

SPIROSTACHYS, Sond. in Linnaea, xxiii. (1850) 106 = **Excoecaria**, Linn. (Euphorb.).
africanus, Sond. l. c. = E. africana.
madagascarienis, Baill. Etud. Gén. Euph. 522 = E. madagascariensis.

SPIROSTACHYS, Ung.-Sternb. Vers. Syst. Salic. 100 (1866) = **Heterostachys**, Ung.-Sternb. (Chenop.).
Ritteriana, Ung.-Sternb. l. c. = H. Ritteriana.

SPIROSTACHYS, S. Wats. in Proc. Am. Acad. ix. (1874) 125. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 63.
occidentalis, *S. Wats. l. c.*—Am. bor. occ.
patagonica, *Griseb. in Goett. Abh.* xix. (1874) 86.—Reg. Argent.
vaginata, *Griseb. l. c.* 85.—Reg. Argent.

SPIROSTALIS, Rafin. Fl. Tellur. iv. 51 (1836) = **Thalia**, Linn. (Scitam.).
biflora, Rafin. l. c. = T. dealbata.

SPIROSTEMON, Griff. Notul. iv. 80 (1858) = **Parsonsia**, R. Br. (Apocyn.).
spiralis, Griff. l. c. = P. spiralis.

SPIROSTIGMA, Nees, in Mart. Fl. Bras. ix. 83 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1077.
hirsutissima, *Nees, l. c.* 84.—Bras.

SPIROSTYLIS, Nees, ex Mast. in Journ. Linn. Soc. x. (1869) 271 = **Willdenowia**, Thunb. (Restiac.).
Ecklonii, Nees, l. c. = W. Lucaeana.

SPIROSTYLIS, Presl, in Schult. f. Syst. vii. p. xvii, 163 (1829) = **Loranthus**, Linn.
Haenkei, Presl, l. c. 163 = L. spirostylis.
interruptus, Blume, l. c. 1731 = L. interruptus.

SPIROTHEROS, Rafin. in Ser. Bull. Bot. i. (1830) 221 = **Heteropogon**, Pers. (Gramin.).
melanocarpus, Rafin. l. c. = H. acuminatus.

SPIROTROPIS, Tul. in Arch. Mus. Par. iv. (1844) 113. *LEGUMINOSAE*, Benth. & Hook. f. i. 557.
Candollei, *Tul. l. c.*—Guian.
longifolia, Baill. Hist. Pl. ii. 364 = Candollei.

SPITZELIA, Sch. Bip. in Flora, xvi. (1833) 725 = **Picris**, Linn. (Compos.).
aegyptiaca, Sch. Bip. l. c. 727 = P. pilosa.
Aparine, Pomel, Nouv. Mat. Fl. Atl. 268.—Alger.
aspera, Pomel, l. c. 15.—Alger.
atlantica, Coss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 537 = P. atlantica.
aviorum, Pomel, Nouv. Mat. Fl. Atl. 16 = *S. Saharae.*
coronopifolia, Sch. Bip. in Flora, xviii. (1835) 658 = P. radicata.
cupuligera, Dur. in Duch. Rev. Bot. ii. (1846) 431 = P. pilosa.
getula, Pomel, Nouv. Mat. Fl. Atl. 267 = *S. Saharae.*

SPITZELIA :—
lyrata, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 397 = *S. Saharae*.
lyrata, Sch. Bip. in Flora, xviii. (1835) 658 = Picris radicata.
radicata, Coss. & Kral. in Bull. Soc. Bot. Fr. iv. (1857) 367 = P. radicata.
Saharae, Coss. & Kral. l. c. 397.—Afr. bor.
Sieberi, Sch. Bip. in Flora, xvii. (1834) 474 = P. radicata.
tuberosa, Pomel, Nouv. Mat. Fl. Atl. 268.—Alger.
Willkommii, Sch. Bip. ex Willk. in Linnaea, xxv. (1852) 42 = P. Willkommii.

SPIXIA, Leandr. Sacram. in Denkschr. Akad. Muench. vii. 1818–1820 (1821) 231. t. 13 = **Pera**, Mutis (Euphorb.).
barbinervis, Klotzsch, in Wiegm. Archiv, vii. (1841) 180, nomen = P. obovata.
cinerea, Poepp. ex Baill. Etud. Gén. Euph. 434, nomen = P. tomentosa.
distichophylla, Mart. in Flora, xxiv. (1841) II. Beibl. 30 = P. distichophylla.
glabrata, Mart. ex Baill. Etud. Gén. Euph. 434 = P. ferruginea.
grandiflora, Klotzsch, in Wiegm. Archiv, vii. (1841) 179, nomen = P. obovata.
heteranthera, Schrank, in Denkschr. Akad. Muench. vii. 1818–20 (1821) 242, in obs. = P. obovata.
Leandri, Mart. in Flora, xxiv. (1841) II. Beibl. 30 = P. obovata.
lucida, Poepp. ex Baill. Etud. Gén. Euph. 434, nomen = P. ferruginea.

SPIXIA, Schrank, Pl. Rar. Hort. Monac. t. 80 (1819) = **Centratherum**, Cass. (Compos.).
violacea, Schrank, l. c. = C. intermedium.

SPLITGERBERA, Miq. Comm. Phytog. 133. t. 14 (1840) = **Boehmeria**, Jacq. (Urtic.).
denudata, C. Gay, Fl. Chil. v. 362 = B. excelsa.
japonica, Miq. Comm. Phytog. 139. t. 14 = B. biloba.
macrostachya, Wight, Ic. t. 1977 = B. platyphylla.

SPODIAS, Hassk. Cat. Hort. Bog. Alt. 247 (1844), sphalm. = **Spondias**, Linn. (Anacard.).

SPODIOPOGON, Trin. Fund. Agrost. 192. t. 17 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1126.
ISCHAEMOPOGON, Griseb. Fl. Brit. W. Ind. 560 (1864).
albidus, *Benth. in Journ. Linn. Soc.* xix. (1881) 66.—Ind. or.
angustifolius, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 300.—Ind. or. ; China ; Ins. Philipp.
arcuatus, Nees, Fl. Afr. Austr. 97 = Ischaemum fasciculatum.
aureus, *Hook. & Arn. Bot. Beech. Voy.* 273.—Ins. Lu Chu.
Blumii, *Nees, ex Steud. Syn. Pl. Gram.* 373 = Ind. or. ; Malaya.
Byronis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 301.—Ins. Sandvic.
chordatus, Trin. l. c. 302 = Blumii.
ciliaris, Nees, ex Steud. Nom. ed. II. ii. 625 = Ischaemum ciliare.
conjugatus, Voigt, Hort. Suburb. Calc. 706 = Ischaemum conjugatum.
cuspidatus, Arn. ex Steud. Syn. Pl. Gram. 373 = Blumii.
depressus, *Nees, ex Steud. l. c.*—Ind. or.
foliatus, Fourn. ex Doell, in Mart. Fl. Bras. ii. III. 257 = Erianthus saccharoides.
inaequalivalvis, *Nees, ex Steud. Nom.* ed. II. ii. 625, nomen.—China.
geniculatus, Voigt, Hort. Suburb. Calc. 706 = Ischaemum aristatum.
ischaemoides, *Hook. & Arn. Bot. Beech. Voy.* 274.—Ins. Bonin.
laniger, Nees, ex Steud. Syn. Pl. Gram. 373 = angustifolius.
latifolius, Nees, Agrost. Bras. 360 = Ischaemum latifolium.
Lehmanni, Griseb. in Goett. Nachr. (1868) 91 = Pollinia Lehmanni.
notopogon, Nees, ex Steud. Syn. Pl. Gram. 373 = angustifolius.
SPODIOPOGON :—
obliquivalvis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 185 = Ischaemum ciliare.
petiolaris, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 301 = Pollinia Lehmanni.
pilosus, Nees, ex Steud. Syn. Pl. Gram. 373 = Andropogon pilosus.
pogonanthus, *Boiss. Fl. Orient.* v. 457.—Phrygia.
scrobiculatus, Nees, ex Steud. Syn. Pl. Gram. 373 = Ischaemum ciliare.
semisagittatus, Voigt, Hort. Suburb. Calc. 706 = Ischaemum semisagittatum.
sibiricus, *Trin. Fund. Agrost.* 192.—China ; Sibir.
vaginatus, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 520 = Erianthus saccharoides.
villosus, Nees, in Hook. & Arn. Bot. Beech. Voy. 242 = Ischaemum ciliare.
zeylanicus, Nees, ex Steud. Syn. Pl. Gram. 377 = Ischaemum ciliare.

SPONDIAS, Linn. Gen. ed. I. 365 (1737). *ANACARDIACEAE*, Benth. & Hook. f. i. 426.
CHRYSOMELON, Forst. ex A. Gray, Bot. U. St. Expl. Exped. i. 374, in syn. (1854).
CYTHERAEA, Wight & Arn. Prod. 173 (1834).
EVIA, Comm. ex Juss. Gen. 373 (1789).
MONBIN, Plum. ex Adans. Fam. ii. 341 (1763).
POUPARTIA, Comm. ex Juss. Gen. 372 (1789).
SHAKUA, Boj. Hort. Maurit. 82 (1837).
WARMINGIA, Engl. in Mart. Fl. Bras. xii. II. 281 (1874).
WIRTGENIA, Jungh. ex Hassk. in Flora, xxvii. (1844) 624, partim.
acida, Blume, in Flora, vii. (1824) 291 = dulcis.
acida, Soland. ex Benth. Fl. Austral. i. 492 = Pleiogynium Solandri.
acuminata, *Roxb. Hort. Beng.* 34 ; *Fl. Ind.* ii. 453 (sp. dub.).—Ind. or.
amara, Lam. Encyc. iv. 261 = mangifera.
angolensis, O. Hoffm. in Linnaea, xliii. (1880–82) 125 = Pseudospondias microcarpa.
aurantiaca, Schum. & Thonn. Besk. Guin. Pl. 225 = lutea.
axillaris, Roxb. Hort. Beng. 34; Fl. Ind. ii. 453 = lutea.
Birrea, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 152. t. 41 = Sclerocarya Birroea.
borbonica, *Baker, Fl. Maurit.* 62.—Ins. Maurit.
brasiliensis, Mart. ex Engl. in DC. Monog. Phan. iv. 244 = lutea.
Chakoua, Boj. ex Baker, Fl. Maurit. 63 = Sclerocarya castanea.
Cirouella, Tussac, Fl. Antill. iii. 37 = purpurea.
crispula, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856), 119 = purpurea
cytherea, Sonner. Voy. Ind. iii. 242 = dulcis.
cytherea, Tussac, Fl. Antill. iii. t. 28 = lutea.
dubia, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 153 = lutea.
dulcis, *Forst. f. Prod.* 34.—Cosmop. trop.
Edmonstonei, Hook. f. in Trans. Linn. Soc. xx. (1847) 230 = Bursera graveolens.
elliptica, Rottb. ex Hook. f. Fl. Brit. Ind. ii. 23 = Buchanania latifolia.
falcata, Meissn. in Flora, xxvii. (1844) 349 = Harpephyllum caffrum.
fragrans, Pav. ex Engl. in DC. Monog. Phan. iv. 247 = dulcis.
glabra, Engl. l. c. 572 (Index) = lutea.
graveolens, Macfad. Fl. Jamaic. i. 228 = lutea.
guianensis, Klotzsch, ex Engl. in DC. Monog. Phan. iv. 277 = Tapirira guianensis.
longifolia, *Roxb. Hort. Beng.* 34 ; *Fl. Ind.* ii. 453.—Ins. Maurit.
lucida, Salisb. Prod. 172 = lutea.
lutea, *Linn. Sp. Pl.* ed. II. 613.—Cosmop. trop.
lutea, Royen, ex Blume, Mus. Bot. Lugd. Bat. i. 234 = dulcis.
macrocarpa, Engl. in Mart. Fl. Bras. xii. II. 375. t. 78 = dulcis.
macrophylla, Klotzsch, ex Engl. in DC. Monog. Phan. iv. 277 = Tapirira guianensis.
macrophylla, Wall. Cat. n. 8480 = mangifera.
mangifera, *Willd. Sp. Pl.* ii. 751.—As. trop.
maxima, Engl. in DC. Monog. Phan. iv. 572 (Index) = lutea.
microcarpa, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 151 = Pseudospondias microcarpa.

SPONDIAS :—
Mombin, Jacq. Select. Am. 138 = lutea.
Mombin, Linn. Sp. Pl. ed. I. 371 = purpurea.
Myrobalanus, Jacq. Stirp. Am. t. 88 = purpurea.
Myrobalanus, Linn. Syst. ed. X. 1036 = lutea.
Myrobalanus, Vell. Fl. Flum. 197; iv. t. 185 = venulosa.
Oghigee, G. Don, Gen. Syst. ii. 79 = Odina acida.
paniculata, Roxb. ex Wight & Arn. Prod. 173 = mangifera.
parviflora, Willd. ex Schlecht. in Linnaea, xiv. (1840) 295 = Tapirira guianensis.
pinnata, Kurz, in Pegu Rep. A. 44 = mangifera.
pleiogyna, F. Muell. Fragm. iv. 78 = Pleiogynium Solandri.
Pseudo-myrobalanus, Tussac, Fl. Antill. iv. 97 = lutea.
pubescens, *Baker, Fl. Maurit.* 62.—Ins. Maurit.
pubescens, *Bouton, ex Steud. Nom.* ed. II. ii. 626.—Ins. Maurit.
purpurea, *Linn. Sp. Pl.* ed. II. 613.—Am. trop.
rugosa, Klotzsch, ex Engl. in DC. Monog. Phan. iv. 277 = Tapirira guianensis.
simplicifolia, Rottb. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 187 = Buchanania Lanzan.
sinensis, Lour. ex Steud. Nom. ed. I. 306, nomen (Quid ?).—China.
Solandri, Benth. Fl. Austral. i. 492 = Pleiogynium Solandri.
surinamensis, Klotzsch, ex Engl. in DC. Monog. Phan. iv. 277 = Tapirira guianensis.
therebintoides, Royen, ex Blume, Mus. Bot. Lugd. Bat. i. 234 = dulcis ?
tuberosa, *Arruda, in Koster, Trav. Braz.* 496 (sp. dub.).—Bras.
venulosa, *Mart. ex Engl. in DC. Monog. Phan.* iv. 245.—Bras.
viridiflora, *Fée, Cat. Méthod. Pl. Strasb.* (1836) 110 (sp. dub.).—Bras.
Wirtgenii, Hassk. Cat. Hort. Bog. Alt. 247 = Odina Wodier.
Zanzee, G. Don, Gen. Syst. ii. 79 = Pseudospondias microcarpa ?

SPONDIOIDES, Smeathm. ex Lam. Encyc. iii. 23 (1789) = **Cnestis**, Juss. (Connar.).
ferrugineum, Smeathm. ex DC. Prod. ii. 87 = C. ferruginea.
pruriens, Smeathm. ex Lam. Encyc. iii. 23 = C. corniculata.
villosa, Smeathm. l. c. 24 = Agelaea villosa.

SPONDOGONA, Rafin. Sylva Tellur. 35 (1838) = **Dipholis**, A. DC. (1844) (Sapot.).
nitida, Rafin. l. c. = D. salicifolia.

SPONDYLANTHA, Presl, Rel. Haenk. ii. 35. t. 53 (1834) = **Vitis**, Tourn. (Ampel.).
aphylla, Presl, l. c. = V. sicyoides.

SPONDYLOCOCCOS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 218 = **Callicarpa**, Linn. (Verben.).

SPONDYLOCOCCUS, Reichb. Consp. 212 d (1828) = SPHONDYLOCCA, Willd. = **Bergia**, Linn. (Elatin.).

SPONGOSTEMMA, Reichb. Ic. Fl. Germ. iv. 24 (1826) = **Scabiosa**, Tourn. (Dipsac.).

SPONGOTRICHUM, Nees, Gen. et Sp. Ast. 176 (1833) = **Olearia**, Moench (Compos.).
glandulosa, Nees, l. c. 175, 176 = O. glandulosa.

SPONIA, Comm. ex Lam. Encyc. iv. 139 (1796) = **Trema**, Lour. (Urtic.).
acuminatissima, Miq. Fl. Ind. Bat. Suppl. 410.—Sumatra.
affinis, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 329 = T. orientalis.
africana, Planch. l. c, 320 = T. africana.
amboinensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. amboinensis.
Andaresa, Comm. ex Lam. Encyc. iv. 139 = T. orientalis.

SPONIA :—
Andersonii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 336 = Parasponia Andersonii.
angustifolia, Planch. l. c. 326 = Trema angustifolia.
annulata, Teysm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 408 = Gironniera subaequalis.
argentea, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 323 = T. argentea.
argentea, Seem. Bot. Voy. Herald, 413 = T. amboinensis.
aspera, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. aspera.
Blancoi, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 327 = T. Blancoi.
bracteolata, Hochst. in Flora, xxviii. (1845) 87 = T. bracteolata.
Burmanni, Planch. in DC. Prod. xvii. 200 = T. amboinensis.
canescens, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. micrantha.
carinata, Miq. Fl. Ind. Bat. i. II. 215 = T. carinata.
Chichilea, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 334 = T. Chichilea.
Commersonii, Decne. ex Planch. l. c. 317 = T. orientalis.
crassifolia, G. Don, in Sweet, Hort. Brit. ed. III. 610 = Celtis occidentalis.
crassifolia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 340 = T. micrantha.
discolor, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. discolor.
glabrescens, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 317 = T. glabrescens.
glomerata, Hochst. in Flora, xxviii. (1845) 87 = T. glomerata.
Griffithii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 324 = T. amboinensis.
grisea, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 340 = T. micrantha.
guineensis, Planch. in DC. Prod. xvii. 197 = Celtis guineensis.
Hochstetteri, Planch. l. c. 198.—Abyss.
ilicifolia, Kurz, in Flora, lv. (1872) 448 = Aphananthe philippinensis.
imbricata, Planch. in DC. Prod. xvii. 199 = T. imbricata.
integerrima, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 144.—Panama.
laevigata, G. Don, in Sweet, Hort. Brit. ed. III. 610 = Celtis mississipiensis.
Lamarckiana, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. Lamarckiana.
Lima, Decne. l. c. = T. Lima.
lucida, Hassk. in Flora, xxv. (1842) II. Beibl. 17.—Java.
macrophylla, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. macrophylla.
macrophylla, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 331 = T. carinata.
micrantha, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. micrantha.
mollis, Decne. l. c. = T. mollis.
morifolia, Planch. in DC. Prod. xvii. 196 = T. morifolia.
nitens, Hook. f. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 325 = T. nitens.
nudiflora, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 223 = Aphananthe aspera.
occidentalis, G. Don, in Sweet, Hort. Brit. ed. III. 610 = Celtis occidentalis.
orientalis, Blume, ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 328 = T. amboinensis.
orientalis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. orientalis.
orientalis, Seem. in Bonplandia, ix. (1861) 259 = Parasponia Andersonii.
pallida, Miq. Fl. Ind. Bot. i. II. 215 = T. pallida.
pendula, Span. in Linnaea, xv. (1841) 343 = Celtis timorensis.
peruviana, Klotzsch, in Linnaea, xx. (1847) 536 = T. carinata.
politoria, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 326 = T. politoria.
pubigera, Miq. Fl. Ind. Bat. i. II. 216 = T. pubigera.
pumila, G. Don, in Sweet, Hort. Brit. ed. III. 610 = Celtis occidentalis.
rigida, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 98 = T. rigida.

SPONIA :—
rigida, Zoll. ex Planch. in DC. Prod. xvii. 181 = Celtis cinnamomea.
riparia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = Trema riparia.
rostrata, Span. in Linnaea, xv. (1841) 343, nomen (Quid?).—Ins. Timor.
rufescens, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 334 = T. rufescens.
rugosa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. rugosa.
Sampsoni, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 242 = T. timorensis.
scaberrima, Miq. Fl. Ind. Bat. i. II. 217 = T. scaberrima.
Schiedeana, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 335 = T. micrantha.
sinensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = Celtis sinensis.
Solenostigma, Steud. Nom. ed. II. ii. 626 = Celtis paniculata.
stipellata, Bello, in Anal. Soc. Esp. Hist. Nat. xii. (1883) 109.—Porto Rico.
strigosa, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 320 = T. strigosa.
strychnifolia, Teysm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 392 = Celtis laurifolia.
subserrata, Kurz, in Flora, lv. (1872) 448 = Gironniera reticulata.
tahitensis, Nadeaud, Enum. Pl. Tahiti, 42 = Parasponia Andersonii.
tetrandra, Voigt, Hort. Suburb. Calc. 294 = Celtis tetrandra.
timorensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498 = T. timorensis.
tomentosa, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 336 = T. amboinensis.
trinervia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 498.—Ins. S. Doming.
velutina, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 327 = T. amboinensis.
Vieillardi, Planch. in DC. Prod. xvii. 201.—N. Caled.
virgata, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 316 = T. timorensis.
viridis, Planch. l. c. 319 = T. aspera.
Wightii, Planch. l. c. 322 = T. orientalis.
Willdenowiana, G. Don, in Sweet, Hort. Brit. ed. III. 610 = Celtis japonica.

SPORABOLUS, Hassk. Cat. Hort. Bog. Alt. 17 (1844), sphalm. = **Sporobolus**, R. Br. (Gramin.).

SPORADANTHUS, F. Muell. ex J. Buch. in Trans. N. Z. Inst. vi. (1874) 340; x. (1878) App. 41 = **Lepyrodia**, R. Br. (Restiac.).
Traversii, F. Muell. ex J. Buch. ll. cc. = L. Traversii.

SPORLEDERA, Bernh. in Linnaea, xvi. (1842) 41 = **Ceratotheca**, Endl. (Pedalin.).
Kraussiana, Bernh. l. c. 42 = C. Kraussiana.
triloba, Bernh. l. c. = C. triloba.

SPOROBOLUS, R. Br. Prod. 169 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1148.
AGROSTICULA, Raddi, Agrost. Bras. 33. t. 1. f. 2 (1823).
BENNETIA, Rafin. in Ser. Bull. Bot. i. (1830) 220.
CRYPTOSTACHYS, Steud. Syn. Pl. Gram. 181 (1855).
DIACHYRIUM, Griseb. in Goett. Abh. xix. (1874) 256.
HELEOCHLOA, Beauv. Agrost. 23 (1812).
SPERMACHITON, Llanos, Fragm. 25 (1851).
TRIACHYRUM, Hochst. in Flora, xxiv. (1841) I. Intell. 19.
VILFA, Beauv. Agrost. 16. t. 5. f. 8 (1812).
actinocladus, *F. Muell. Fragm.* viii. 140.—Austral.
aeneus, *Kunth, Enum. Pl.* i. 213.—Bras.
affinis, *Kunth, Rév. Gram.* i. 68.—Am. austr.
affinis, *A. Rich. Tent. Fl. Abyss.* ii. 396.—Abyss.
airoides, *Torr. in Marcy, Rep.* 300.—Am. bor. occ.
albicans, *Nees, Fl. Afr. Austr.* 154.—Afr. austr.
alpicola, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 395.—Abyss.
angustifolius, *A. Rich. l. c.* 396.—Abyss.
angustus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 88 = indicus.
arabicus, *Boiss. Diagn.* Ser. I. xiii. 47.—Arab.; Ind. or.

SPOROBOLUS :—
arenaceus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 89 = asperifolius.
arenarius, Duval-Jouve, in Bull. Soc. Bot. Fl. xvi. (1869) 294 = pungens.
argutus, *Kunth, Enum. Pl.* i. 215.—Bras.
arundinaceus, Vasey, Cat. Gr. U. St. 43 = Arctagrostis latifolia.
asper, *Kunth, Rév. Gram.* i. 68.—Am. bor.
asperifolius, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 141.—Am. occ. bor. et austr.
atrovirens, *Kunth, Rév. Gram.* i. 68.—Mexic.
blephariphyllus, *A. Rich. Tent. Fl. Abyss.* ii. 398.—Abyss.
brevifolius, Nees, Fl. Afr. Austr. 160 = tenellus.
Buckleyi, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 128.—Texas.
caespitosus, *Kunth, Rév. Gram.* ii. 423. t. 125.—Ins. Adscens.
capensis, *Kunth. Enum. Pl.* i. 212.—Afr. austr.
capillaris, Miq. Anal. Bot. Ind. ii. 25 = pulchellus.
Carmichaeli, Kunth, Rév. Gram. i. 68 = Agrostis Carmichaeli.
centrifugus, *Nees, Fl. Afr. Austr.* 158.—Afr. austr.
ciliatus, *J. S. Presl, in Rel. Haenk.* i. 242.—Mexic.
commutatus, Kunth, Enum Pl. i. 214 = coromandelianus.
compressus, *Kunth, l. c.* 217.—Am. bor.
confertiflorus, *A. Rich. Tent. Fl. Abyss.* ii. 397.—Abyss.
confertus, *J. A Schmidt, Beitr. Fl. Cap. Verd. Ins.* 142.—Ins. Cap. Virid.
consimilis, *Fresen. in Mus. Senckenb.* ii. (1837) 140.—Abyss.
coromandelianus, *Link, Hort. Berol.* i. 89, in nota; *Kunth, Rév. Gram.* i. 68.—As. et Afr. trop.
cryptandrus, *A. Gray, Man. Bot. N. U. St.* ed. I. 576.—Am. bor.
cuspidatus, *Wood, Am. Bot. & Flor.* 385; *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 63.—Am. bor.
depauperatus, *Scribner, l. c.*—Am. bor.
depauperatus, Torr. ex Hemsl. Biol. Centr. Am. Bot. iii. 546 = plumbeus.
diander, *Beauv. Agrost.* 26.—As. et Austral. trop.
diffusissimus, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 90 = airoides.
discosporus, *Nees, Fl. Afr. Austr.* 158.—Afr. austr.
domingensis, *Kunth, Enum. Pl.* i. 214.—Ind. occ.
Drummondii, *Vasey, Cat. Gr. U. St.* 44.—Am. bor.
durus, *Brongn. in Duperr. Voy. Coq. Bot.* 18. t. 4.—Ins. Adscens.
effusus, *Desv. Opusc.* 60.—Bras.
elongatus, R. Br. Prod. 170 = indicus.
eminens, *J. S. Presl, in Rel. Haenk.* i. 242.—Mexic.
fastigiatus, *J. S. Presl, l. c.* 241.—Peruv.
festivus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 398.—Abyss.
filiculmis, *Vasey, Cat. Gr. U. St.* 44.—Am. bor.
fimbriatus, *Nees, Fl. Afr. Austr.* 156.—Afr. austr.
floridanus, *Chapm. Fl. S. U. St.* 550.—Florida.
gaditanus, Boiss. & Reut. Pugill. Pl. Nov. 125 = pungens.
Gigas, *Miq. Fl. Ind. Bat.* iii. 376.—Java.
glaucifolius, *Hochst. in Flora*, xxv. (1842) I. *Beibl.* 133, nomen: *et ex Steud. Syn. Pl. Gram.* 154.—Nubia; Sind.
gracillimus, *Vasey, Cat. Gr. U. St.* 44.—Am. bor.
heterolepis, *A. Gray, Man. Bot. N. U. St.* ed. I. 576.—Am. bor.
heterolepsis, Lapham, in Trans. Wisconsin St. Agric. Soc. iii. 1853 (1854) 426, sphalm. = praec.
humifusus, *Kunth, Rév. Gram.* i. 67.—Mexic.
humilis, *J. S. Presl, in Rel. Haenk.* i. 241.—Ins. Philipp.
indicus, *R. Br. Prod.* 170.—Austral.
insulanus, *Parl. in Atti 8va Ruin. Sc. Ital.* 586.—Ins. Cap. Virid.
insularis, Hook. Niger Fl. 187, sphalm. = praec.
ioclados, *Nees, Fl. Afr. Austr.* 161.—Afr. austr.; Ind. or.
Jacquemontii, *Kunth, Rév. Gram.* ii. 427. t. 127.—Ind. occ.
Jonesii, *Vasey, in Coult. Bot. Gaz.* vi. (1881) 297.—Calif.
junceus, *Kunth, Rév. Gram.* i. 68.—Am. bor.
Lamarckii, Desv. in Ham. Prod. Pl. Ind. Occ. 4 = indicus.
leptostachys, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 30.—Afr. trop.

SPOROBOLUS:—
Lindleyi, Benth. Fl. Austral. vii. 623 = pallidus.
littoralis, Kunth, Rév. Gram. i. 68 = glaucifolius?
longifolius, Wood, Class Book (1861), 775 = asper?
Ludwigii, *Hochst. in Flora*, xxix. (1846) 118.—Afr. austr.
mangaloricus, Miq. Anal. Bot. Ind. ii. 24 = minutiflorus.
marginatus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 397.—Abyss.
Matrella, *Nees, Fl. Afr. Austr.* 152.—Afr. austr.
minor, Kunth, Enum. Pl. i. 212 = indicus.
minutiflorus, *Link, Hort. Berol.* i. 88.—Ind. or.
minutus, *Link, l. c.*—Abyss.
Muehlenbergii, *Kunth, Rév. Gram.* i. 68.—Am. bor.
myrianthus, *Benth. in Hook. Niger Fl.* 565.—Afr. trop.
nervosus, *Hochst. in Flora*, xxxviii. (1855) 202.—Abyss.
noterophilus, *Hemsl. Biol. Centr. Am. Bot.* iii. 546.—Mexic.
orientalis, *Kunth, Enum. Pl.* i. 211.—Ind. or.
pallidus, Benth. Fl. Austral. vii. 623, in obs., sphalm. = arabicus.
pallidus, *Lindl. in Mitch. Journ. Trop. Austral.* 187.—Austral.
panicoides, *A. Rich. Tent. Fl. Abyss.* ii. 399.—Abyss.
parviflorus, *Kunth, Rév. Gram.* i. 68.—Austral.
pellucidus, *Hochst. in Flora*, xxxviii. (1855) 201.—Abyss.
phyllotrichus, *Hochst. l. c.*—Abyss.
piliferus, *Kunth, Enum. Pl.* i. 211.—Ind. or.; China.
plumbeus, *Hemsl. Biol. Centr. Am. Bot.* iii. 546.—Mexic.
pulchellus, *R. Br. Prod.* 170.—Austral.; Ind. or.
pungens, *Kunth, Rév. Gram.* i. 68.—Reg. Mediterr.
purgans, Griseb. Fl. Brit. W. Ind. 773 (Index), sphalm. = praec.
purpurascens, *Ham. Prod. Pl. Ind. Occ.* 5.—Am. trop.
pyramidalis, *Beauv. Fl. Owar.* ii. 36.—Afr. trop.
ramosissimus, *Kunth, Rév. Gram.* i. 269. t. 46.—Am. trop.
ramulosus, *Kunth, l. c.* 68.—Am. bor. occ.
repens, *F. S. Presl, in Rel. Haenk.* i. 241.—Mexic.
rigens, *E. Desv. in C. Gay, Fl. Chil.* vi. 295.—Chili.
robustus, *Kunth, Rév. Gram.* ii. 425. t. 126.—Afr. trop
Rueppellianus, *Fresen. in Mus. Senckenb.* ii. (1837) 139.—Abyss.
rupestris, *Kunth, Rév. Gram.* i. 267. t. 45.—Bras.
sarmentosus, *Griseb. in Goett. Abh.* xxiv. (1879) 295.—Reg. Argent.
scaber, *Phil. Fl. Atac.* 54.—Chili.
scoparius, *F. S. Presl, in Rel. Haenk.* i. 243.—Ins. Philipp.
serotinus, *A. Gray, Man. Bot. N. U. St.* ed. I. 577.—Am. bor.
spicatus, *Kunth, Rév. Gram.* i. 67.—Afr. bor.; Ind. or.
Sprengelii, *Kunth, l. c.* 68.—Bras.
stachydanthus, *A. Rich. Tent. Fl. Abyss.* ii. 394.—Abyss.
subinclusus, *Phil. Sert. Mendoc. Alt.* 49.—Chili.
subtilis, *Kunth, Rév. Gram.* ii. 421. t. 124.—Madag.
subtilis, F. Muell. Fragm. viii. 140 = pallidus.
tenacissimus, Beauv. Agrost. 26 = indicus.
tenellus, *Kunth, Enum. Pl.* i. 215.—Afr. austr.
tremulus, Kunth, Rév. Gram. i. 67 = virginicus.
tricholepis, *Coult. Man. Rocky Mount. Bot.* 411.—Am. bor.
truncatellus, *Nees, ex Steud. Syn. Pl. Gram.* 159.—Guian.
vaginaeflorus, *Vasey, Cat. Gr. U. St.* 45.—Am. bor.
verticillatus, *Nees, in Hook. Kew Journ.* ii. (1850) 101.—Ins. Philipp.
villosus, Hochst. ex Doell, in Mart. Fl. Bras. ii. II. 234 = Panicum amplexicaule.
villosus, *Kunth, Rév. Gram.* i. 68.—Guian.
virginicus, *Kunth, l. c.* 67.—Cosmop. trop.
Wolfii, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 52.—Colorado.
Wrightii, *Munro, ex Vasey, Cat. Gr. U. St.* 26, nomen.—Am. bor.

SPORTELLA, Hance, in Journ. Bot. xv. (1877) 207 = **Crataegus**, Tourn. (Rosac.).
atalantioides, Hance, l. c. = C. Pyracantha.

SPRAGUEA, Torr. in Smithson. Contrib. vi. (Pl. Frémont.) (1854) 4. t. 1. *PORTULACAEAE*, Benth. & Hook. f. i. 158.
paniculata, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 187. f. 56.—Am. bor. occ.
umbellata, *Torr. in Smithson. Contrib.* vi. (1854) 4.—N. Mexic.

SPREKELIA, Heist. Syst. 5 (1748); Desc. Brunsvig. 19 (1753). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 724.
Cybister, Herb. in Bot. Reg. (1840). t. 33 = Hippeastrum Cybister.
formosissima, *Herb. App.* [*Bot. Reg.*] 35.—Mexic.
glauca, Lindl. Bot. Reg. (1840). Misc. 49 = formosissima.
Heisteri, Trew, ex Kunth, Enum. Pl. v. 507 = formosissima.
Karwinskii, M. Roem. Syn. Ensat. 144 = formosissima.
ringens, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) 133. t. 60 = formosissima.

SPRENGALIA, 'Andr.' ex Steud. Nom. ed. II. ii. 626 (1841), sphalm. = **Sprengelia**, Sm. (Epacrid.).

SPRENGELIA, Schult. Obs. Bot. 134 (1809) = **Melhania**, Forsk. (Stercul.).
modesta, Schult. l. c. = M. abyssinica.

SPRENGELIA, Sm. in Vet. Acad. Handl. Stockh. (1794) 260; Tracts, 269. t. 2 (1798). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
POIRETIA, Cav. Ic. iv. 25. t. 343 (1797).
PONCELETIA, R. Br. Prod. 554 (1810).
Andersoni, F. Muell. Fragm. vi. 64 = Andersonia sprengelioides.
aristata, F. Muell. l. c. 63 = Andersonia aristata.
brachyanthera, F. Muell. l. c. 61 = Andersonia brachyanthera.
brachynema, F. Muell. l. c. = Andersonia heterophylla.
brachyota, F. Muell. l. c. 62 = Andersonia parvifolia.
brevifolia, F. Muell. l. c. 61 = Andersonia brevifolia.
coerulea, F. Muell. l. c. 64 = Andersonia coerulea.
colossea, F. Muell. l. c. 63 = Andersonia colossea.
depressa, F. Muell. l. c. = Andersonia depressa.
incarnata, *Sm. in Vet. Acad. Handl. Stockh.* (1794) 260; *Tracts*, 272. t. 2.—Austral.
involucrata, F. Muell. Fragm. vi. 62 = Andersonia involucrata.
latiflora, F. Muell. l. c. 61 = Andersonia latiflora.
macrantha, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 273 = incarnata.
macronema, *F. Muell. Fragm.* viii. 51.—Austral.
micrantha, F. Muell. l. c. vi. 61 = Andersonia micrantha.
montana, R. Br. Prod. 555 = incarnata.
parvifolia, F. Muell. Fragm. vi. 62 = Andersonia parvifolia.
patricia, F. Muell. l. c. 79 = Andersonia patricia.
Ponceletia, *F. Muell. l. c.* i. 39.—Austral.
ponceletioides, *Sond. in Linnaea*, xxvi. (1853–55) 254.—Austral.
propinqua, A. Cunn. ex DC. Prod. vii. 768 = incarnata.
spirophylla, F. Muell. Fragm. vi. 62 = Andersonia grandiflora.

SPRINGALIA, DC. Prod. vii. 768 (1839), err. typ. = praec.

SPRINGIA, Heurck & Muell. Arg. in Heurck, Obs. Pl. Nov. 142 (1871) = **Ichnocarpus**, R. Br. (Apocyn.).
indica, Heurck & Muell. Arg. l. c. 143 = I. ovatifolius.

SPRUCEA, Benth. in Hook. Kew Journ. v. (1853) 230. *RUBIACEAE*, Benth. & Hook. f. ii. 43.
rubescens, *Benth. in Hook. Kew Journ.* v. (1853) 230.—Bras.

SPRUNIRA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) I. Intell. 42; xxv. (1842) I. Beibl. 134, 145 = **Sphaeranthus**, Linn. (Compos.).
alata, Sch. Bip. ll. cc. = Sphaer. nubicus.

SPRUNNERA, Sch. Bip. in Walp. Rep. ii. 954 (1843) = **Codonocephalum**, Fenzl (Compos.).
inuloides, Sch. Bip. l. c. = C. inuloides.

SPYRIDIUM, Fenzl, in Enum. Pl. Hueg. 24 (1837). *RHAMNEAE*, Benth. & Hook. f. i. 382.
STENODISCUS, Reiss. in Linnaea, xxix. (1857–58) 295.
bifidum, *F. Muell. ex Benth. Fl. Austral.* i. 432.—Austral.
coactilifolium, *Reiss. in Linnaea*, xxix. (1857–58) 291.—Austral.
complicatum, *F. Muell. Fragm.* iii. 78.—Austral.
cordatum, *Benth. Fl. Austral.* i. 430.—Austral.
diffusum, Reiss. in Linnaea, xxix. (1857–58) 288 = vexilliferum.
divaricatum, *Benth. Fl. Austral.* i. 427.—Austral.
eriocephalum, *Fenzl, in Enum. Pl. Hueg.* 24.—Austral.
globulosum, *Benth. Fl. Austral.* i. 429.—Austral.
Gunnii, *Benth. l. c.*—Tasman.
halmaturinum, *F. Muell. ex Benth. l. c.* 432.—Austral.
Lawrencii, *Benth. l. c.* 430.—Tasman.
leucophractum, F. Muell. Fragm. iii. 77 = Stenanthemum leucophractum.
microcephalum, *Benth. Fl. Austral.* i. 434.—Austral.
obovatum, *Benth. l. c.* 429.—Tasman.
oligocephalum, *Benth. l. c.* 433.—Austral.
parvifolium, *F. Muell. Fragm.* iii. 79.—Austral.
pauciflorum, *Benth. Fl. Austral.* i. 432.—Austral.
phlebophyllum, *F. Muell. ex Benth. l. c.* 430.—Austral.
phylicoides, Reiss. in Linnaea, xxix. (1857–58) 286 = vexilliferum.
prostratum, Reiss. l. c. 284 = eriocephalum.
pumilum, *F. Muell. Fragm.* ix. 137.—Austral.
rotundifolium, *F. Muell. l. c.* iv. 25.—Austral.
serpyllaceum, *F. Muell. l. c.* iii. 80.—Austral.
spadiceum, *Benth. Fl. Austral.* i. 428.—Austral.
spathulatum, *F. Muell. ex Benth. l. c.* 430.—Austral.
Stuarti, Reiss. & F. Muell. in Linnaea, xxix. (1857–58) 290 = coactilifolium.
subochreatum, *Reiss. l. c.* 287.—Austral.
thymifolium, Reiss. l. c. 289 = coactilifolium.
tridentatum, *Benth. Fl. Austral.* i. 427.—Austral.
ulicinum, *Benth. l. c.* 434.—Tasman.
uncinatum, Reiss. in Linnaea, xxix. (1857–58) 289 = eriocephalum.
vexilliferum, *Reiss. l. c.* 285.—Austral.
villosum, *Benth. Fl. Austral.* i. 432.—Austral.
Waterhousii, F. Muell. Fragm. iii. 83 = Stenanthemum Waterhousii.
westringiaefolium, *Benth. Fl. Austral.* i. 431.—Austral.

SQUAMARIA, Riv. ex Linn. Gen. ed. I. (1737) 183 = **Lathraea**, Linn. (Orobanch.).
Orobanche, Scop. Fl. Carn. ed. II. i. 438 = L. Squamaria.

SQUAMATAXUS, [Nelson], Senilis, Pinaceae, 168 (1866) = **Saxegothaea**, Lindl. (Conif.).
Albertiana, [Nelson], l. c. = Sax. conspicua.

SQUAMMARIA, Riv. ex Linn. Syst. ed. I (1735) = SQUAMARIA, Riv. = **Lathraea**, Linn. (Orobanch.).

SQUILLA, Steinh. in Ann. Sc. Nat. Sér. II. vi. (1836) 276 = **Urginea**, Steinh. (Liliac.).
anthericoides, Jord. & Fourr. Ic. Fl. Eur. ii. 2 = U. anthericoides.
insularis, Jord. & Fourr. l. c. = U. Scilla.
littoralis, Jord. & Fourr. l. c. = U. Scilla.
maritima, Steinh. in Ann. Sc. Nat. Sér. II. vi. (1836) 279 = U. Scilla.
numidica, Jord. & Fourr. Ic. Fl. Eur. ii. 1 = U. Scilla.
Pancration, Steinh. in Ann. Sc. Nat. Sér. II. vi. (1836) 279 = U. Scilla.
sphaeroidea, Jord. & Fourr. Ic. Fl. Eur. ii. 1 = U. Scilla.

SRUTANTHUS, Pritz. Ic. Ind. i. 1061 (1855), sphalm. = seq.

SRUTHANTHUS, DC. Prod. iv. 671 (1830), sphalm. = STRUTHANTHUS, Mart. = **Loranthus**, Linn.

STAAVIA, Dahl, Obs. Bot. 15 (1787). *BRUNIACEAE*, Benth. & Hook. f. i. 672.
ASTROCOMA, Neck. Elem. i. 112 (1790).
LEVISANUS, Schreb. Gen. i. 149 (1789).
STAVIA, Thunb. Nov. Gen. vii. 108 (1792).

STAAVIA:—
adenandraefolia, *Eckl. & Zeyh. Enum.* 141.—Afr. austr.
capitella, Sond. in Harv. & Sond. Fl. Cap. ii. 323 = adenandraefolia.
ciliata, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 380.—Afr. austr.
ciliata, Eckl. & Zeyh. Enum. 141 = adenandraefolia.
Dregeana, Presl, Bot. Bemerk. 39 = radiata.
glaucescens, E. Mey. in Drège, Zwei Pfl. Docum. 223, nomen = glutinosa.
globosa, Sond. in Harv. & Sond. Fl. Cap. ii. 322 = ciliata.
glutinosa, *Dahl, Obs. Bot.* 17.—Afr. austr.
nuda, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 379.—Afr. austr.
nuda, Eckl. & Zeyh. Enum. 141 = adenandraefolia.
pinifolia, Willd. in Denkschr. Akad. Muench. i. (1809) 133 = radiata.
radiata, *Dahl, Obs. Bot.* 15.—Afr. austr.
rupestris, Eckl. & Zeyh. Enum. 141 = adenandraefolia.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* ii. 323.—Afr. austr.

STABEROHA, Kunth, Enum. Pl. iii. 442 (1841). *RESTIACEAE*, Benth. & Hook. f. iii. 1034.
distachya, *Kunth, l. c.* 444.—Afr. austr.
imbricata, *Kunth, l. c.* 442 = Afr. austr.
stenoptera, Kunth, l. c. 443 = imbricata.

STACHIS, Neck. Delic. Gallo-Belg. i. 255 (1768) = **Stachys**, Tourn. (Labiat.).

STACHYACANTHUS, Nees, in Mart. Fl. Bras. ix. 65 (1847). Genus dubium *ACANTHACEARUM*.—Cf. Benth. & Hook. f. Gen. ii. 1071.
Riedelianus, *Nees, l. c.* 66.—Bras.

STACHYANTHUS, DC. Prod. v. 84 (1836) = **Eremanthus**, Less. (Compos.).
argyrophyllus, Mart. & Zucc. ex Baker, in Mart. Fl Bras. vi. II. 168 = E. Martii.
Martii, DC. Prod. v. 84 = E. Martii.

STACHYARRHENA, Hook. f. in Hook. Ic. Pl. t. 1068 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 80.
longifolia, *Hook. f. in Hook. Ic. Pl.* sub t. 1068.—Bras.
spicata, *Hook. f. l. c.* t. 1068.—Bras.

STACHYCEPHALUM, Sch. Bip. ex Benth. in Hook. Ic. Pl. t. 1102 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 344.
argentinum, *Griseb. in Goett. Abh.* xxiv. (1879) 188.—Reg. Argent.
mexicanum, *Sch. Bip. ex Benth. in Hook. Ic. Pl.* t. 1102.—Mexic.

STACHYCHRYSUM, Boj. Hort. Maurit. 114 (1837) = **Piptadenia**, Benth. (Legumin.).
pterospermum, Boj. l. c. = P. chrysostachys.

STACHYCRATER, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 464 = **Osmelia**, Thw. (Samyd.).
philippinus, Turcz. l. c. 465 = O. philippinensis.
zeylanicus, Turcz. l. c. xxxvi. (1863) I. 609 = O. Gardneri ?

STACHYOBIUM, Reichb. f. in Gard. Chron. (1869) 785 = **Dendrobium**, Sw. (Orchid.).
aureum, Reichb. f. l. c. = D. binoculare.

STACHYOCNIDE, Blume, Mus. Bot. Lugd. Bat. ii. 227. t. 55 (1856) = **Pouzolzia**, Gaudich. (Urtic.).
luzonica, Blume, l. c. 228. t. 55 = P. macrura.

STACHYOPHORBE, Liebm. in Overs. Dansk. Vidensk. Selsk. (1846) 8 = **Chamaedorea**, Willd. (Palm.).
cataractarum, Liebm. l. c. = C. cataractarum.
Deckeriana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 364 = C. Deckeriana.
montana, Liebm. in Overs. Dansk. Vidensk. Selsk. (1846) 8 = oreophila.
pygmaea, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 10 = C. pygmaea.

STACHYOPOGON, Klotzsch, in Bot. Ergeb. Waldem. Reise, 49 (1862) = **Aletris**, Linn. (Haemod.).
pauciflorus, Klotzsch, l. c.—Reg. Himal.
spicatus, Klotzsch, l. c.—Reg. Himal.

STACHYS, [Tourn.] Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1208.
AMBLEIA, Spach, Hist. Vég. Phan. ix. 166 (1840), nomen.
ASPASIA, E. Mey. ex Pfeiff. Nom. i. 295 (1873).
BETONICA, Tourn. ex Linn. Syst. ed. I (1735); et Gen. ed. I. 171 (1737).
BONAMYA, Neck. Elem. i. 316 (1790).
ERIOSTACHYS, Reichb. ex Steud. Nom. ed. II. i. 588 (1840).
ERIOSTOMUM, Hoffmgg. & Link, Fl. Port. i. 105. t. 7 (1809).
GALEOPSIS, Moench, Meth. 397 (1794).
OLISIA, Spach, Hist. Vég. Phan. ix. 166 (1840).
ORTOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 136.
TETRAHITUM, Hoffmgg. & Link, Fl. Port. i. 103 (1809).
TRIXAGO, Hall. Hist. Stirp. Helv. 101 (1768); Moench, Meth. 398 (1794).
TRIXELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135.
ZIETENIA, Gled. Syst. 185 (1764).
abrotanifolia, Boiss. & Heldr. in Boiss. Fl. Orient. iv. 732 = angustifolia.
acerosa, *Boiss. Diagn.* Ser. I. vii. 57.—Persia.
aculeolata, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 18.—Afr. trop.
acutifolia, Bory & Chaub. Expéd. Mor. 168 = germanica.
acutifolia, Link, in Linnaea, ix. (1834) 575 = graeca.
adenocalyx, C. Koch, l. c. xxi. (1848) 691 = annua.
aegyptiaca, Pers. Syn. ii. 124 = affinis.
aethiopica, *Linn. Mant.* i. 82.—Afr. austr.
affinis, Bunge, Enum. Pl. Chin. Bor. 51 = palustris.
affinis, *Fresen. in Mus. Senckenb.* i. (1833) 91.—Aegypt.; Arab.
agraria, *Schlecht. & Cham. in Linnaea*, v. (1830) 100.—Texas; Mexic.
agraria, Torr. in Bot. U. St. Expl. Exped. ii. 408 = bullata.
ajugoides, *Benth. in Linnaea*, vi. (1831) 80.—Calif.
alba, Mill. Gard. Dict. ed. VIII. n. 4 = germanica.
albens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 387.—Calif.
albicaulis, Bert. ex Benth. in DC. Prod. xii. 473 = grandidentata.
albicaulis, *Lindl. Bot. Reg.* t. 1558.—Chili.
aleurites, *Boiss. & Heldr. Diagn.* Ser. I. xii. 79.—Pamphylia.
alopecuros, *Benth. Lab. Gen. et Sp.* 531.—Europ.
alpina, *Linn. Sp. Pl.* 581.—Europ.; Reg. Cauc.
altissima, Desf. Cat. Hort. Par. ed. III. 101 = italica.
ambigua, Sm. Engl. Bot. t. 2089 = sylvatica.
angustifolia, *Bieb. Fl. Taur. Cauc.* ii. 51.—Europ. austr. or.
angustifolia, Gilib. Fl. Lituan. i. 80 = palustris.
anisochila, *Vis. & Panč. in Mem. Ist. Venet.* xv. (1870) 13.—Serbia; Herceg.
annua, *Linn. Sp. Pl.* ed. II. 813.—Europ.; Reg. Cauc.
annua, Sibth. & Sm. Fl. Graec. Prod. i. 410 = recta.
annua, Walt. Fl. Carol. 161 = aspera.
arabica, *Hornem. Hort. Hafn.* 554.—Syria.
arenaria, *Vahl, Symb. Bot.* ii. 64.—Reg. Mediterr. occ.
argentea, Tausch, in Flora, xiv. (1831) 217 = germanica.
argilacea, Pomel, Nouv. Mat. Fl. Atl. 118 = Duriaei.
argolica, Boiss. Diagn. Ser. I. xii. 83 = Swainsoni.
armena, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 694 = Marrubium cephalanthum.
Artemisiae, Lour. Fl. Cochinch. 365 = Leonurus sibiricus.
arvensis, *Linn. Sp. Pl.* ed. II. 814.—Geront. temp. bor.
arvensis, Walt. Fl. Carol. 162 = aspera.
aspera, *Michx. Fl. Bor. Am.* ii. 5.—Am. bor.
atherocalyx, C. Koch, in Linnaea, xxi. (1848) 691 = recta.
Aucheri, *Benth. in DC. Prod.* xii. 487.—Persia.
aurea, *Benth. l. c.* 492.—Afr. austr.

STACHYS:—
austriaca, Heynh. Nom. ii. 697 = palustris.
baicalensis, Fisch. ex Benth. Lab. Gen. et Sp. 543 = palustris.
Balansae, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 722.—As. Min.
Balbisii, Link, Enum. Hort. Berol. ii. 110 = pubescens.
Ballota, Kuntze, Taschen-Fl. Leipz. 112 = Ballota nigra.
ballotaeformis, Vatke, in Bot. Zeit. xxxiii. (1875) 448 = Benthamiana.
barbata, Lapeyr. Hist. Abr. Pl. Pyr. 336 = heraclea.
barbigera, Viv. Fl. Cors. App. ii. 4 = heraclea.
Benthamiana, *Boiss. Fl. Orient.* iv. 734.—Persia.
betaefolia, *Pers. Syn.* ii. 124.—Hab. ?
Betonica, *Benth. Lab. Gen. et Sp.* 532.—Europ.; As. Min.
Betonica, Crantz, Stirp. Austr. ed. II. fasc. iv. 264 = annua.
Betonica, Scop. Fl. Carn. ed. II. i. 411 = recta.
betonicaeflora, *Rupr. Sert. Tiansch.* 66.—Turkestan.
betonicaefolia, Desf. in Ann. Mus. Par. x. (1807) 304 = spinulosa.
betonicaefolia, Pers. Syn. ii. 124 = heraclea.
betonicifolia, Regel, in Act. Hort. Petrop. vi. (1879) 366 = betonicaeflora.
biennis, Hort. Hafn. ex Benth. in DC. Prod. xii. 464 = lanata.
biennis, Roth, in Usteri, Ann. Bot. xiv. (1795) 27 = germanica.
biflora, *Hook. & Arn. Bot. Beech. Voy.* 155 (sp. dub.).—Calif.
Bigelovii, *A. Gray, in Proc. Am. Acad.* viii. (1872) 371.—Texas; Am. bor.
bithynica, Boiss. Diagn. Ser. I. v. 28 = germanica.
Bodeana, Bunge, Lab. Pers. 71 = tomentosa.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* ii. 309.—N. Granat.
bombycina, *Boiss. Diagn.* Ser. I. xii. 79.—Syria.
boraginoides, Benth. Lab. Gen. et Sp. 540 = Lindenii.
boraginoides, *Schlecht. & Cham. in Linnaea*, v. (1830) 100.—Mexic.
brachiata, *Boj. ex Benth. Lab. Gen. et Sp.* 547.—Madag.
brachyclada, Noé, ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 226; et in Bull. Soc. Bot. Fr. ii. (1855) 583 = annua.
Brantii, *Benth. in DC. Prod.* xii. 463.—Kurdistan.
Brantii, Huet, ex Boiss. Fl. Orient. iv. 722 = Huetii.
brasiliensis, Hort. ex Benth. Lab. Gen. et Sp. 550 = arvensis.
Bridgesii, *Benth. l. c.* 546.—Chili.
Bufonia, Thuill. Fl. Par. ed. II. i. 295 = recta.
bullata, *Benth. Lab. Gen. et Sp.* 547.—Calif.
Burchellii, *Benth. l. c.* 561.—Afr. austr.
burgsdorffioides, *Boiss. Diagn.* Ser. I. xii. 85.—Syria; Mesopot.
byzantina, C. Koch, in Linnaea, xxi. (1848) 686 = lanata.
caffra, *E. Mey. ex Benth. in DC. Prod.* xii. 495.—Afr. austr.
californica, *Benth. l. c.* 469.—Calif.
calycantha, Bieb. Fl. Taur. Cauc. iii. 393 = lavandulaefolia.
canariensis, Jacq. Ic. Pl. Rar. i. 11. t. 108 = sylvatica.
candida, Benth. Lab. Gen. et Sp. 547, partim = mollis.
candida, *Bory & Chaub. Expéd. Mor.* 167. t. 19.—Graecia.
candidissima, *Phil. in Linnaea*, xxix. (1857–58) 19.—Chili.
canescens, *Bory & Chaub. Expéd. Mor.* 167. t. 18.—Graecia.
capensis, *Presl, Bot. Bemerk.* 100.—Afr. austr.
cardinalis, Kunze, in Bot. Zeit. ii. (1844) 645 = coccinea, *Jacq.*
cassia, *Boiss. Diagn.* Ser. I. xiii. 74.—Macedon.; Syria.
caucasica, C. Koch, in Linnaea, xxi. (1848) 693 = iberica.
Chamaesideritis, Boiss. & Bal. Diagn. Ser. II. iv. 41 = citrina.
Chamissonis, *Benth. in Linnaea*, vi. (1831) 80.—Calif.
chinensis, *Bunge, ex Benth. Lab. Gen. et Sp.* 544.—China.
chonotica, *Hook. f. Fl. Antarct.* 336.—Chili.

STACHYS:—
chrysantha, *Boiss. & Heldr. Diagn.* Ser. I. vii. 56.—Graecia.
ciliaris, Boiss. l. c. xii. 78 = libanotica.
ciliata, *Dougl. ex Benth. Lab. Gen. et Sp.* 539.—Am. bor. occ.
cilicica, Boiss. & Bal. Diagn. Ser. II. iv. 42 = annua.
cinerea, Hort. ex Benth. Lab. Gen. et Sp. 536 = germanica.
circinata, *L'Hérit. Stirp. Nov.* 51. t. 26.—Hispan.; Afr. bor.
citrina, *Boiss. & Heldr. ex Benth. in DC. Prod.* xii. 490.—Graecia; As. Min.
coccinea, Hook. & Arn. Bot. Beech. Voy. 156 = bullata.
coccinea, *Jacq. Hort. Schoenb.* iii. 18. t. 284.—Texas; Mexic.
coccinea, *Ortega, Hort. Matr. Dec.* 20.—Cuba.
coerulea, *Burch. ex Benth. Lab. Gen. et Sp.* 558.—Afr. austr.
congesta, Urv. in Mém. Soc. Linn. Par. i. (1822) 324 = iberica.
cordata, Gilib. Fl. Lituan. i. 80 = sylvatica.
cordata, *Riddell, Suppl. Cat. Ohio Pl.* 15.—Am. bor.
cordato-folia, Gilib. Exercit. i. 91 = sylvatica.
cordifolia, *C. Koch, in Linnaea*, xxi. (1848) 685.—Oriens.
corsica, *Pers. Syn.* ii. 36.—Reg. Mediterr.
crenata, Rafin. Fl. Ludov. 42 (Quid ?).—Am. bor.
cretica, Forsk. Fl. Aegypt. Arab. p. xxviii = orientalis.
cretica, Linn. Sp. Pl. 581 = germanica, alpina.
cretica, *Sibth. & Sm. Fl. Graec.* vi. 47. t. 558.—Graec.; As. Min.; Syria.
cretica, Sieber, ex Boiss. Fl. Orient. iv. 718 = Tournefortii.
crispa, *Arv. Ess. Pl. Dauph.* 58 (Quid ?).—Gallia.
cuneata, *Banks, ex Benth. Lab. Gen. et Sp.* 560.—Afr. austr.
curviflora, Tausch, in Flora, xiv. (1831) 216 = alpina.
Cydni, *Kotschy, ex Tchihat. Asie Min. Bot.* ii. 173, nomen.—As. Min.
dasyanthes, Heldr. ex Boiss. Fl. Orient. iv. 721 = Heldreichii.
dasyanthes, *Rafin. Préc. Découv.* 39.—Sicil.
debilis, *Baker, in Journ. Linn. Soc.* xx. (1883) 234.—Madag.
debilis, *H. B. & K. Nov. Gen. et Sp.* ii. 309.—Ecuador.
decemdentata, Forst. f. Prod. 91 = Leucas flaccida.
decumbens, Pers. Syn. ii. 123 = mollissima.
decumbens, Reichb. Fl. Germ. Excurs. 318 = recta.
decumbens, Willd. Enum. Hort. Berol. Suppl. 41 = pubescens.
delphinensis, Jord. Cat. Jard. Bot. Grenoble (1853), 12; et in F. Schultz, Arch. Fl. Fr. et Allem. 322 = recta.
densiflora, *Benth. Lab. Gen. et Sp.* 532.—Europ. austr.
denticulata, *Burch. ex Benth. l. c.* 560.—Afr. austr.
deserti, Benth. in DC. Prod. xii. 494 = rugosa.
diffusa, Schweigg. Enum. Pl. Hort. Regiom. 69 = arenaria.
discolor, *Benth. Lab. Gen. et Sp.* 533.—Reg. Cauc.
distans, *Benth. in DC. Prod.* xii. 472.—Syria.
divaricata, Viv. Fl. Cors. App. 2 = hirta.
diversifolia, *Boiss. Diagn.* Ser. I. xii. 80.—Syria.
Dregeana, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 240.—Afr. austr.
Drummondii, *Benth. Lab. Gen. et Sp.* 551.—Texas; Mexic.
Durandiana, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 256.—Marocc.
Duriaei, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 583.—Afr. bor.
Ehrenbergii, *Boiss. Fl. Orient.* iv. 721.—Syria.
elata, C. Koch, in Linnaea, xxi. (1848) 687 = spectabilis.
elliptica, *H. B. & K. Nov. Gen. et Sp.* ii. 308.—Ecuador.
elliptica, *Phil. in Linnaea*, xxxiii. (1864–65) 193.—Chili.
elliptica, Poepp. ex Benth. in DC. Prod. xii. 473 = grandidentata.
elongata, Hort. ex Benth. Lab. Gen. et Sp. 536 = germanica.
erecta, C. Koch, in Linnaea, xxi. (1848) 691 = recta.
eriantha, *Benth. Lab. Gen. et Sp.* 549.—Mexic.
erioleuca, *Pomel, Nouv. Mat. Fl. Atl.* 305.—Alger.

STACHYS:—
excelsa, Hort. ex Benth. Lab. Gen. et Sp. 536 = germanica.
excelsa, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 194 = Keerlii.
flavescens, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 241.—Afr. austr.
floccosa, *Benth. Lab. Gen. et Sp.* 739.—Reg. Himal.
floribunda, Montbr. & Auch. in Ann. Sc. Nat. Sér. II. vi. (1836) 51 = pumila.
floridana, *Shuttl. ex Benth. in DC. Prod.* xii. 478.—Florida.
fluminensis, Vell. Fl. Flum. 241; vi. t. 1 = Leucas martinicensis.
Foeniculum, Pursh, Fl. Am. Sept. ii. 407 = Lophanthus anisatus.
foetida, Gueldenst. ex Ledeb. Fl. Ross. iii. 413 = sylvatica.
foetida, *Vitm. in Colla, Herb. Pedem.* iv. 449.—Hab.?
foliosa, Benth. in E. Mey. Comm. Pl. Afr. Austr. 241 = rugosa.
foliosa, Regel, in Act. Hort. Petrop. vi. (1879) 366 = betonicaeflora.
fragilis, Vis. Fl. Dalm. ii. 207, in obs. t. 16. f. 1 = recta.
fruticulosa, *Bieb. Fl. Taur. Cauc.* ii. 51.—Reg. Cauc.; Persia.
fusca, Moench, Meth. Suppl. 136 = alpina.
Galeottii, *Martens, in Bull. Acad. Brux.* xi. II. (1844) 192.—Mexic.
garana, Boiss. Diagn. Ser. I. xii. 76 = cretica.
gariepina, *Benth. in DC. Prod.* xii. 493.—Afr. austr.
germanica, *Linn. Sp. Pl.* 581.—Europ.; Afr. bor.
Gilliesii, *Benth. Lab. Gen. et Sp.* 740.—Chili.
glabra, Riddell, Suppl. Cat. Ohio Pl. 16 = aspera.
glaucescens, Mussin. ex Spreng. Syst. ii. 736 = sylvatica.
glutinosa, *Linn. Sp. Pl.* 581.—Reg. Mediterr.
glutinosa, Sibth. & Sm. Fl. Graec. Prod. i. 409 = recta.
graciliflora, *Presl, Bot. Bemerk.* 100.—Afr. austr.
graeca, *Boiss. & Heldr. Diagn.* Ser. I. xii. 77.—Graecia.
Grahami, Benth. Lab. Gen. et Sp. 551 = agraria.
grandidentata, *Lindl. Bot. Reg.* t. 1080.—Chili.
grandiflora, *Benth. Lab. Gen. et Sp.* 533.—As. Min.; Persia.
grandiflora, Host, Fl. Austr. ii. 169 = menthaefolia.
grandifolia, *E. Mey. Comm. Pl. Afr. Austr.* 239.—Afr. austr.
Guyoniana, *Noé, ex Munby, Cat. Pl. Alger.* 24, nomen.—Alger.
Haussknechtii, Vatke, in Bot. Zeit. xxxiii. (1875) 461 = Kotschyi.
Haynaldi, Janka, ex Nym. Consp. 580 = angustifolia.
Heldreichii, *Boiss. Fl. Orient.* iv. 721.—Graec.; Byzant.
heraclea, *All. Fl. Pedem.* i. 31. t. 84.—Europ. austr.
heraclea, Botan. Graec. ex Boiss. Fl. Orient. iv. 723 = graeca.
heraclea, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 624 = germanica.
heterophylla, Moench, Meth. Suppl. 135 = germanica.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 95.—Afr. trop.
hirsuta, *H. B. & K. Nov. Gen. et Sp.* ii. 308.—Mexic.
hirta, Georgi, Beschr. Russ. Reich. III. v. 1086 = arenaria.
hirta, *Linn. Sp. Pl.* ed. II. 813.—Reg. Mediterr.
hirtula, Pomel, Nouv. Mat. Fl. Atl. 305 = hirta.
hispanica, Mill. Gard. Dict. ed. VIII. n. 6 = germanica?
hispida, Pursh, Fl. Am. Sept. ii. 407 = aspera.
hispidula, *Hochst. in Flora*, xxviii. (1845) 66.—Afr. austr.
Huetii, *Boiss. Fl. Orient.* iv. 722.—Armenia.
humifusa, *Burch. ex Benth. Lab. Gen. et Sp.* 547.—Afr. austr.
hybrida ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 102.—Europ.
hydrophila, *Boiss. Diagn.* Ser. I. xii. 81.—Syria.
hypoleuca, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 197 = palaestina.
hypoleuca, C. Koch, in Linnaea, xxi. (1848) 688 = spectabilis.

STACHYS:—
hyssopifolia, *Michx. Fl. Bor. Am.* ii. 4.—Am. bor.
hyssopifolia, Vahl, ex Benth. Lab. Gen. et Sp. 559 = rugosa.
hyssopoides, *Burch. ex Benth. Lab. Gen. et Sp.* 558.—Afr. austr.
iberica, *Bieb. Fl. Taur. Cauc.* ii. 51.—As. Min.; Syria.
inflata, *Benth. Lab. Gen. et Sp.* 562.—Mesopot.; Persia.
inscripta, Reichb. Ic. Bot. Exot. i. 28. t. 40 = hirta.
integrifolia, *Vahl, ex Benth. Lab. Gen. et Sp.* 562.—Afr. austr.
intermedia, Ait. Hort. Kew. ed. I. ii. 301 = germanica.
intermedia, Tenore, Prod. Fl. Nap. p. xxxiv = heraclea.
italica, Benth. ex Nym. Consp. 577, partim = dasyanthes.
italica, *Mill. Gard. Dict.* ed. VIII. n. 3.—Reg. Mediterr.
italica, Orph. ex Nym. Consp. 578 = cretica.
italica, Reichb. ex Nym. l. c. = Janiana.
Iva, *Griseb. Spicil. Fl. Rumel.* ii. 143.—Macedon.
ixodes, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 738.—Persia.
Janiana, *Ces. Pass. & Gib. Comp. Fl. Ital.* 318.—Europ.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 111.—Japon.
jugalis, Burch. ex Benth. Lab. Gen. et Sp. 562 = rugosa.
Keerlii, *Benth. l. c.* 551.—Mexic.
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. v. 32.—Kurdistan.
Kraussii, *Hochst. in Flora*, xxviii. (1845) 66.—Afr. austr.
kurdica, Boiss. & Hohen. Diagn. Ser. I. v. 31 = subnuda.
labiosa, Bertol. Fl. Ital. vi. 166 = subcrenata.
laetevirens, Kotschy & Boiss. ex Boiss. Fl. Orient. iv. 734 = viscosa.
Lamarckii, *Benth. Lab. Gen. et Sp.* 562. — Afr. austr.
lamiiflora, *Rupr. Sert. Tiansch.* 67.—As. centr.
lamioides, *Benth. Pl. Hartw.* 244.—N. Granat.
lanata, Crantz, Stirp. Austr. ed. II. fasc. iv. 267 = germanica.
lanata, *Jacq. Ic. Pl. Rar.* i. 11. t. 107.—Byzant.; Reg. Cauc.; Persia.
lanata, Moench, Meth. 397 = palaestina.
lasiantha, Steud. Nom. ed. II. ii. 628, sphalm. = dasyanthes.
latifolia, Ait. Hort. Kew. ed. I. ii. 303 = circinata.
lavandulaefolia, *Vahl, Symb. Bot.* i. 42.—Reg. Cauc.; As. Min.; Persia.
laxa, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 179.—Persia.
lazica, Boiss. Fl. Orient. iv. 745 = pubescens.
leucoglossa, Griseb. Spicil. Fl. Rumel. ii. 140 = recta.
libanotica, *Benth. in DC. Prod.* xii. 462.—Syria.
Lindenii, *Benth. l. c.* 467.—Mexic.
linearifolia, C. Koch, in Linnaea, xxi. (1848) 692 = recta.
linearis, Burch. ex Benth. Lab. Gen. et Sp. 559 = rugosa.
longespicata, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 725.—As. Min.; Syria.
longiflora, *Boiss. & Bal. Diagn.* Ser. II. iv. 39.—Cilicia.
longifolia, *Benth. Lab. Gen. et Sp.* 533.—Reg. Cauc.
longifolia, Bert. ex Steud. Nom. ed. II. ii. 628 = sideritidoides.
longifolia, Rafin. in Am. Monthly Mag. (1818) 356, nomen (*sphalm.* Staelry longifolia) (Quid?).—Am. bor.
lunata, *Pall. Fl. Ross.* (*Index Tomi* ii. 1).—Rossia.
lusitanica, Bory & Chaub. Nouv. Fl. Pélop. 37 = graeca.
lusitanica, Steud. Nom. ed. I. 806 = germanica.
Lyallii, *Benth. Lab. Gen. et Sp.* 548.—Madag.
lycopsiformis, C. Koch, in Linnaea, xxi. (1848) 692 = setifera.
macedonica, D. Don, ex Steud. Nom. ed II. ii. 628; et ex Griseb. Spicil. Fl. Rumel. ii. 143 = mollis.
macilenta, *E. Mey. Comm. Pl. Afr. Austr.* 240.—Afr. austr.
Macraei, *Benth. in Linnaea*, vi. (1831) 78.—Chili.

STACHYS:—

macrocheilos, Boiss. Diagn. Ser. I. v. 30 = fruticulosa.
macrosperma, Boiss. l. c. xii. 84 = arabica.
major, *Uspensky, in Bull. Soc. Nat. Mosc.* vii. (1834) 376, nomen.—Rossia.
maritima, *Linn. Mant.* i. 82.—Europ. austr.
maritima, Urv. in Mém. Soc. Linn. Par. i. (1822) 324 = pubescens.
marrubiifolia, *Viv. Fl. Cors. App.* 2.—Ins. Corsic.; Afr. bor.
mauritiana, Pers. Syn. ii. 123 = Anisomeles malabarica.
Maweana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 626.—Marocc.
mediterranea, Vell. Fl. Flum. 241; vi. t. 2 = Leonotis nepetaefolia.
melissaefolia, *Benth. Lab. Gen. et Sp.* 538.—Reg. Himal.
menthaefolia, *Vis. in Flora*, xii. (1829) 1. *Erg.* 14.—Dalmat.
menthoides, *Kotschy & Boiss. in Fl. Orient.* iv. 727.—Armenia.
mersinaea, Boiss. Diagn. Ser. II. iv. 37 = cretica.
messeniaca, *Boiss. in DC. Prod.* xii. 475.—Graecia.
mexicana, *Benth. Lab. Gen. et Sp.* 541.—Mexic.
Meyeni, *Phil. in Anal. Univ. Chil.* (1873) 520.—Chili.
Meyenii, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 375.—Peruv.
Mialhesii, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 584.—Alger.
micrantha, *Griseb. in Goett. Abh.* xxiv. (1879) 275.—Reg. Argent.
micrantha, C. Koch, in Linnaea, xxi. (1848) 690 = annua.
modica, *Hance, in Journ. Bot.* xx. (1882) 292.—China.
mollis, *Willd. ex Benth. in Linnaea*, xi. (1837) 335; *et in DC. Prod.* xii. 475.—Marocc.
mollissima, *Willd. Hort. Berol.* 60. t. 60.—Graecia.
Montbretii, Benth. in Ann. Sc. Nat. Sér. II. vi. (1836) 48 = orientalis.
mucronata, *Sieber, ex Spreng. Syst.* ii. 733.—Ins. Creta.
multicaulis, *Benth. in DC. Prod.* xii. 486.—Persia.
multiflora, *Benth. l. c.* 492.—Afr. austr.
natalensis, *Hochst. in Flora*, xxviii. (1845) 65.—Afr. austr.
nepetaefolia, *Cav. Desc.* 74.—Hab. ?
nepetifolia, *Desf. Tabl.* ed. I. 60, nomen; *et ex Pers. Syn.* ii. 123.—Mexic.
nervosa, Gater. Desc. Pl. Montaub. 107 = annua.
neurocalycina, *Boiss. Diagn.* Ser. I. xii. 85.—Syria.
nigricans, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 238.—Afr. austr.
nitens, *Janka, in Linnaea*, xxx. (1859–60) 597.—Europ. austr.
nivea, Benth. in DC. Prod. xii. 461, in syn. = discolor.
nivea, *Labill. Ic. Pl. Syr.* iii. 5. t. 3.—Syria.
nova, Sadler, ex Reichb. Fl. Germ. Excurs. 319 = germanica.
numidica, Pomel, Nouv. Mat. Fl. Atl. 117 = circinata.
nutans, *Benth. Lab. Gen. et Sp.* 561.—Afr. austr.
Nuttallii, Shuttl. ex Benth. in DC. Prod. xii 469 = cordata.
obliqua, Waldst. & Kit. Pl. Rar. Hung. ii. 142 = orientalis.
oblongifolia, *Wall. Cat.* n. 2076; *Benth. in Wall. Pl. As. Rar.* i. 64.—Ind. or.
obscura, Boiss. & Bal. Diagn. Ser. II. iv. 43 = arabica.
obtusata, Boiss. Fl. Orient. iv. 720 = germanica.
obtusicrena, *Boiss. Diagn.* Ser. I. vii. 57.—Persia.
ochroleuca, *Phil. in Linnaea*, xxxiii. (1864–65) 192.—Chili.
ochroleuca, Pomel, Nouv. Mat. Fl. Atl. 119 = Duriaei.
officinalis, Franch. Fl. Loir-et-Cher, 483 = S. Betonica.
oligantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 233.—Madag.
olympica, Poir. Encyc. Suppl. v. 226 = lanata.
orientalis, Forsk. Fl. Aegypt. Arab. p. lxviii = affinis.
orientalis, Hort. ex Benth. Lab. Gen. et Sp. 536 = germanica.
orientalis, Linn. Sp. Pl. 582 = alpina.
orientalis, *Vahl, Symb. Bot.* ii. 64.—Europ. or.; As. Min.
palaestina, *Linn. Sp. Pl.* ed. II. 1674.—Syria.

STACHYS:—

palaestina, Vahl, Symb. Bot. ii. 64 = affinis.
pallida, Salisb. Prod. 83 = circinata.
palustris, *Linn. Sp. Pl.* 580.—Reg. bor. temp.
palustris, Walt. Fl. Carol. 162 = hyssopifolia.
paniculata, Benth. in DC. Prod. xii. 485 = Sideritis lanata.
pannonica, Láng, ex Tenore, Syll. Fl. Neap. 291 = germanica.
Parolinii, *Vis. in Mem. Ist. Venet.* i. 1842 (1843) 46. t. 4.—Graecia.
parviflora, *Benth. in DC. Prod.* xii. 490.—Ind. bor. occ.; Afghan.
parvifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 193.—Mexic.
patens, Sw. Prod. Veg. Ind. Occ. 88 = Hyptis verticillata.
patula, Griseb. Spicil. Fl. Rumel. ii. 142 = recta.
pauciflora, Benth. Lab. Gen. et Sp. 560 = affinis.
pauciflora, Vis. in Mem. Ist. Venet. i. 1842 (1843) 46. t. 2. f. 2 = orientalis.
penicillata, Heldr. & Sart. ex Nym. Consp. 577 = Heldreichii.
penicillata, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 37 = germanica.
persepolitana, *Boiss. l. c.* Ser. I. vii. 56.—Persia.
persica, *S. G. Gmel. ex C. A. Mey. Verz. Pfl. Cauc.* 94.—Persia.
peruviana, *Domb. ex Benth. Lab. Gen. et Sp.* 551.—Peruv.
phlomoides, Willd. Enum. Hort. Berol. Suppl. 41 = heraclea.
pilifera, *Benth. in DC. Prod.* xii. 487.—Persia.
pilosa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 48.—Am. bor.
pilosissima, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 194.—Mexic.
Pinardi, *Boiss. Diagn.* Ser. I. v. 29.—Caria.
pinetorum, *Boiss. & Bal. Diagn.* Ser. II. iv. 36.—Cilicia.
pisidica, Boiss. & Heldr. l. c. Ser. I. xii. 75 = germanica.
plebeia, Vatke, in Bot. Zeit. xxxiii. (1875) 449 = Benthamiana.
plumosa, Benth. in DC. Prod. xii. 492 = aurea.
plumosa, Griseb. Spicil. Fl. Rumel. ii. 139 = recta.
Poiretii, Tenore, Syll. Fl. Neap. 538 = marrubiifolia.
polystachya, Tenore, Prod. Fl. Nap. p. xxxiv = germanica.
Prattenii, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 100.—Calif.
procumbens, Lam. Fl. Fr. ii. 385 = recta.
procumbens, Poepp. ex Benth. in DC. Prod. xii. 473 = grandidentata.
procumbens, Willd. ex Benth. in Linnaea, xi. (1837) 336 = pubescens.
prostrata, Lag. ex DC. in Mém. Soc. Phys. Genèv. i. II. (1823) 455 = arenaria.
pubescens, *Tenore, Prod. Fl. Nap.* p. xxxiv.—Europ. austr.; Oriens.
pulchella, Salisb. Prod. 83 = aethiopica.
pulchra, Pomel, Nouv. Mat. Fl. Atl. 119 = Duriaei.
pumila, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 255.—Syria.
pungens, [Soland. in] Russ. l. c. = italica.
purpurea, Poir. Encyc. Suppl. v. 227 = arenaria.
pycnantha, *Benth. Pl. Hartw.* 331.—Calif.
pycnostachya, Torr. in Bot. U. St. Expl. Exped. ii. 408 = albens.
quadridentata, Bert. ex Benth. in DC. Prod. xii. 473 = Macraei.
ramosissima, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 50.—Mesopot.
ramosissima, Kochel, Pl. Banat. Rar. 26, nomen; et Bot. Reise Bannat, 82 = recta.
recta, *Linn. Mant.* i. 82.—Europ. austr.; Reg. Cauc.
recta, Parol. ex Vis. in Mem. Ist. Venet. i. 1842 (1843) 47 = Parolinii.
recta, Urv. in Mém. Soc. Linn. Par. i. (1822) 324 = angustifolia.
Reinerti, Heldr. ex Boiss. Fl. Orient. iv. 720 = germanica.
repens, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 194.—Mexic.
repens, Schur, Enum. Pl. Transs. 538 = setifera.
revoluta, Rafin. Atl. Journ. 148 (Quid ?).—Am. bor.
Riederi, Benth. in Linnaea, vi. (1831) 570 = ciliata.

STACHYS:—

rigida, Nutt. ex Benth. in DC. Prod. xii. 472 = bullata.
rosea, *Boiss. Fl. Orient.* iv. 725.—Oriens.
rosea, Hohen. in Bull. Soc. Nat. Mosc. (1838) 300 = germanica.
rosmarinifolia, Benth. Lab. Gen. et Sp. 559 = rugosa.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 82.—N. Mexic.
rotundifolia, *Moç. & Sesse, ex Benth. Lab. Gen. et Sp.* 548.—Mexic.
rubra, Donn, Hort. Cantab. ed. IV. 136 = arenaria.
rugosa, *Ait. Hort. Kew.* ed. I. ii. 303.—Afr. austr.
rugosa, Lam. Illustr. iii. 66. t. 509. f. 3 = Lamarckii.
rupestris, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 49.—Cilicia.
rupicola, *Boj. ex Benth. in DC. Prod.* xii. 477.—Madag.
Salisii, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 92 = marrubiifolia.
Salviae, Lindl. Bot. Reg. t. 1226 = Sphacele Lindleyi.
salviaefolia, Tenore, Prod. Fl. Nap. p. xxxiv; Fl. Nap. ii. 25 = italica.
satureioides, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 51.—Syria; Mesopot.
saxicola, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 257.—Marocc.
scaberula, *Vatke, in Bot. Zeit.* xxxiii. (1875) 462.—Reg. Himal.
scardica, Griseb. Reise Rumel. ii. 263 = graeca.
Schiedeana, *Schlecht. in Linnaea*, vii. (1832) 398.—Mexic.
Schimperi, *Vatke, in Linnaea*, xxxvii. (1871–73) 330.—Abyss.
scordifolia, Willd. Enum. Hort. Berol. 616 = arenaria.
scordioides, Desf. Tabl. ed. I. 219 = mollis.
scordioides, *Poir. Encyc.* vii. 373.—Afr. bor.
scyronica, Boiss. Diagn. Ser. I. v. 31 = Swainsoni.
segetum, Hagen, Chloris Boruss. 218 = palustris.
serbica, *Panč. ex Nym. Consp.* 579, nomen.—Serbia.
sericea, *Cav. Desc.* 75.—Chili.
sericea, Ledeb. Fl. Ross. iii. 412 = persica.
sericea, *Wall. Pl. As. Rar.* i. 64.—Reg. Himal.; As. bor.
serrata, *Clos, in C. Gay, Fl. Chil.* iv. 500.—Chili.
serrulata, *Burch. ex Benth. Lab. Gen. et Sp.* 549.—Afr. austr.
sessiliflora, Rafin. Atl. Journ. 148 (Quid ?).—Am. bor.
sessilifolia, *E. Mey. Comm. Pl. Afr. Austr.* 239.—Afr. austr.
setifera, *C. A. Mey. Verz. Pfl. Cauc.* 94.—As. Min.; Persia; Afghan.
sibirica, Link, Enum. Hort. Berol. ii. 109 = germanica.
sideritidoides, *Gill. ex Benth. Lab. Gen. et Sp.* 740.—Chili.
sideritidoides, C. Koch, in Linnaea, xxi. (1848) 692 = recta.
Sideritis, Vill. Hist. Pl. Dauph. ii. 375 = recta.
Sieberi, C. Koch, in Linnaea, xxi. (1848) 685 = Tournefortii.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 112.—Japon.
spathulata, *Burch. ex Benth. Lab. Gen. et Sp.* 559.—Afr. austr.
speciosa, Hook. Bot. Misc. ii. (1831) 235 = Gardoquia elliptica.
spectabilis, *Choisy, ex DC. in Mém. Soc. Phys. Genèv.* i. II. (1823) 457.—Armen.; Persia.
sphaerodonta, *Baker, in Journ. Linn. Soc.* xx. (1883) 233.—Madag.
spinosa, *Linn. Sp. Pl.* 581.—Ins. Creta.
spinulosa, Herb. Madrit. ex Willk. & Lange, Prod. Fl. Hisp. ii. 441 = heraclea.
spinulosa, *Sibth. & Sm. Fl. Graec. Prod.* i. 410.—Europ. austr. or.
splendens, Wall. Pl. As. Rar. i. 64 = melissaefolia, sericea.
Spreitzenhoferi, *Heldr. in Oestr. Bot. Zeitschr.* xxx. (1880) 344.—Ins. Cerigo.
Spruneri, *Boiss. in DC. Prod.* xii. 488.—Graecia.
stenophylla, Spreng. Nov. Prov. 40 = recta ?
stenostachya, Boiss. Diagn. Ser. II. iv. 41 = iberica.
subcrenata, *Vis. in Flora*, xii. (1829) 1. *Erg.* 15.—Dalmatia.
sublanata, Fleischm. ex Nym. Consp. 577 = germanica.
subnuda, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 50.—Kurdist.

STACHYS:—
subsessilis, *Burch. ex Benth. Lab. Gen. et Sp.* 548.—Afr. austr.
Swainsoni, *Benth. l. c.* 535.—Graecia.
Swainsoni, Sprun. ex Nym. Consp. 580 = Spruneri.
sylvatica, *Linn. Sp. Pl.* 580.—Europ.; Reg. Himal.; As. bor.
sylvatica, Nutt. Gen. Am. ii. 30 = cordata.
sylvatica, Torr. in Bot. U. St. Expl. Exped. ii. 408 = bullata.
sylvestris, Forsk. Fl. Aegypt. Arab. p. xxviii = recta.
tenuifolia, *Fisch. ex DC. in Mém. Soc. Phys. Genèv.* i. II. (1823) 454.—Rossia.
tenuifolia, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 180.—Turcia.
tenuifolia, Pall. in Link, Enum. Hort. Berol. ii. 109 = angustifolia.
tenuifolia, Willd. Sp. Pl. iii. 100 = aspera.
tetragona, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 736.—Graecia.
teucriifolia, Boiss. & Kotschy, Diagn. Ser. II. iv. 40 = distans.
Thirkei, *C. Koch, in Linnaea*, xxi. (1848) 685.—As. Min.
Thunbergii, *Benth. Lab. Gen. et Sp.* 540.—Afr. austr.
tibetica, *Vatke, in Bot. Zeitschr.* xxxiii. (1875) 447.—Reg. Himal.
tmolea, *Boiss. Diagn.* Ser. I. v. 29.—As. Min.
tomentosa, *Benth. in DC. Prod.* xii. 489.—Persia.
tomentosa, Bunge, Lab. Pers. 71 = lavandulaefolia.
tomentosa, Gater. Desc. Pl. Montaub. 107 = germanica.
Toronjilcillo, *Phil. in Linnaea*, xxxiii. (1864–65) 193.—Chili.
Tournefortii, Bory & Chaub. Nouv. Fl. Pélop. 37 = messeniaca.
Tournefortii, *Poir. Encyc. Suppl.* v. 227.—Ins. Creta.
transsilvanica, Schur, Enum. Pl. Transs. 539 = recta.
trapezuntea, Boiss. Diagn. Ser. II. iv. 38 = sylvatica.
triloba, Stokes, Bot. Mat. Med. iii. 329 = Leonurus Cardiaca.
truncata, *Kunze, ex Benth. Lab. Gen. et Sp.* 550.—Chili.
umbrosa, *Scheele, in Linnaea*, xxii. (1849) 595.—Texas.
urticaefolia, Tausch, in Flora, xiv. (1831) i. 216 = alpina.
valentina, Lag. in Varied. Cienc. ii. IV. (1805) 39 = heraclea.
velutina, Schwein. in Long, Exped. Winnep. ii. 390 = palustris.
velutina, Willd. Enum. Hort. Berol. Suppl. 41 = circinata.
venosa, *Pers. Syn.* ii. 123.—Hab.?
verbascifolia, Hort. ex Benth. Lab. Gen. et Sp. 536 = germanica.
vestita, Benth. l. c. 538 = sericea.
virgata, *Bory & Chaub. Expéd. Mor.* 166. t. 17.—Graecia.
viridis, *Boiss. & Heldr. Diagn.* Ser. II. iv. 39.—Graecia.
viscosa, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 49.—Armenia.
viticina, *Boiss. Diagn.* Ser. I. xii. 77.—Syria.
Volckmanni, *Phil. in Anal. Univ. Chil.* (1873) 519.—Chili.
Wirtgenii ×, F. Schultz, in Pollichia, xviii–xix. (1861) 117.—Europ.
Zuccarinii, Benth. Lab. Gen. et Sp. 535 = virgata.

STACHYSTEMON, Planch. in Hook. Lond. Journ. Bot. iv. (1845) 471. t. 15. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 263.
brachyphyllus, *Muell. Arg. in Linnaea*, xxxii. (1863) 76.—Austral.
polyandrus, *Benth. Fl. Austral.* vi. 62.—Austral.
vermicularis, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 472. t. 15.—Austral.

STACHYTARPHA, Link, Enum. Hort. Berol. i. 18 (1821) = seq.

STACHYTARPHETA, Vahl, Enum. i. 205 (1805). *VERBENACEAE*, Benth. & Hook. f. ii. 1145.
ABENA, Neck. Elem. i. 296 (1790).
CYMBURUS, Salisb. Parad. Lond. tt. 49, 53 (1806).
MELASANTHUS, Pohl, Pl. Bras. Ic. i. 75. tt. 60–65 (1827).
STACHYTARPHA, Link, Enum. Hort. Berol. i. 18 (1827).

STACHYTARPHETA:—
TARPHETA, Rafin. Fl. Tellur. ii. 103 (1836).
VALERIANOIDES, Medic. Phil. Bot. i. 177 (1789), nomen prius.
VERMICULARIA, Moench, Meth. Suppl. 150 (1802).
acuminata, *DC. ex Schau. in DC. Prod.* xi. 570.—Mexic.
ajugaefolia, *Schau. l. c.*—Bras.
albiflora, *DC. ex Schau. l. c.* 567.—Mexic.
angustifolia, *Vahl, Enum.* i. 205.—Guian.
aristata, Vahl, l. c. 206 = orubica.
asperrima, Walp. Rep. iv. 8, sphalm = scaberrima.
bicolor, *Hook. f. Bot. Mag.* t. 5538.—Bras.
bifurca, Benth. Pl. Hartw. 21 = Bouchea Ehrenbergii.
bogoriensis, *Zoll. & Mor. Syst. Verz. Zoll.* 52.—Java.
bracteosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 198.—N. Granat.
cajanensis, Cham. in Linnaea, vii. (1832) 243 = dichotoma.
cajanensis, *Vahl, Enum.* i. 208.—Am. trop.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 281.—Peruv.
cayennensis, Schau. in DC. Prod. xi. 562 = cajanensis.
Chamissonis, *Walp. Rep.* iv. 10.—Bras.
ciliata, Kunze, Del. Sem. Hort. Lips. (1840) 9, in Adnot. = indica.
coccinea, *Schau. in DC. Prod.* xi. 567.—Bras.
commutata, *Schau. l. c.* 570.—Bras.
crassifolia, *Schrad. in Goett. Gel. Anz.* i. (1821) 709.—Bras.
dichotoma, *Vahl, Enum.* i. 207.—Am. trop.
discolor, *Cham. in Linnaea*, vii. (1832) 251.—Bras.
elatior, *Schrad. ex Schult. Mant.* i. 172.—Bras.; Guian.
elegans, Welw. Apont. 588 = mutabilis.
Frantzii, *Polak. in Linnaea*, xli. (1877) 593.—Costa Rica.
gesnerioides, *Cham. l. c.* vii. (1832) 245.—Bras.
gibberosa, Reichb. Ic. Bot. Exot. ii. 15. t. 138 = dichotoma.
glabra, *Cham. in Linnaea*, vii. (1832) 250.—Bras.
glauca, *Walp. Rep.* iv. 11; *Schau. in DC. Prod.* xi. 569.—Bras.
goyazensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 198.—Bras.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 529.—Afr. trop.
hirsuta, Jacq. f. Eclog. Pl. ii. 10. t. 160 = canescens.
hirsutissima, Link, Enum. Hort. Berol. i. 19 = canescens.
hirta, H. B. & K. Nov. Gen. et Sp. ii. 280 = cajanensis.
hispida, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 69.—Bras.
indica, *Vahl, Enum.* i. 206.—Cosmop. trop.
integrifolia, Walp. Rep. iv. 10 = glauca.
jamaicensis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 184 = dichotoma.
jamaicensis, Vahl, Enum. i. 206 = indica.
lactea, *Schau. in DC. Prod.* xi. 562.—Bras.
lacunosa, *Mart. ex Schau. l. c.* 571.—Bras.
lanata, *Schau. l. c.*—Bras.
laxiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 199.—N. Granat.
longiflora, *Turcz. l. c.* 197.—Hab.?
Lychnitis, *Mart. ex Schau. in DC. Prod.* xi. 571.—Bras.
lythrophylla, *Schau. l. c.* 563.—Bras.
marginata, Vahl, Enum. i. 207 = indica.
Martiana, *Schau. in DC. Prod.* xi. 568.—Bras.
Maximiliani, *Schau. l. c.* 565.—Bras.
microphylla, Walp. Rep. iv. 6 = sanguinea.
mutabilis, *Vahl, Enum.* i. 208.—Ind. occ.; Venezuela.
orubica, *Vahl, l. c.*—N. Granat.
pachystachya, *Mart. ex Schau. in DC. Prod.* xi. 569.—Bras.
palustris, Schott, Cat. Hort. Vindob. (1823) = elatior.
pilosiuscula, H. B. & K. Nov. Gen. et Sp. ii. 279 = indica.
Pohliana, *Cham. in Linnaea*, vii. (1832) 248.—Bras.
polyura, *Schau. in DC. Prod.* xi. 562.—Bras.
prismatica, Vahl, Enum. i. 209 = Bouchea Ehrenbergii.

STACHYTARPHETA:—
Pseudochascanum, Walp. Rep. iv. 11 = Bouchea laetevirens.
quadrangula, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 69.—Bras.
reticulata, *Mart. ex Schau. in DC. Prod.* xi. 566.—Bras.
rhomboidalis, *Walp. Rep.* iv. 10; *Schau. l. c.* 569.—Bras.
rotundifolia, Link, ex Walp. Rep. iv. 5 = canescens.
sanguinea, *Mart. ex Schau. in DC. Prod.* xi. 564.—Bras.
scaberrima, *Cham. in Linnaea*, vii. (1832) 244.—Bras.
Schottiana, *Schau. in DC. Prod.* xi. 563.—Bras.
Sellowiana, *Schau. l. c.* 571.—Bras.
speciosa, *Pohl, ex Schau. l. c.* 568.—Bras.
squamosa, Vahl, Enum. i. 209 = Elytraria tridentata.
strigosa, *Vahl, l. c.* 207.—Mexic.; Ind. occ.
subincisa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 198.—Cuba.
triphylla, Walp. Rep. iv. 10 = rhomboidalis.
trispicata, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 70.—Bras.
umbrosa, H. B. & K. Nov. Gen. et Sp. ii. 280 = dichotoma.
urticifolia, Dalz. & Gibs. Bomb. Fl. Suppl. 68 = indica.
urticifolia, Sims, Bot. Mag. t. 1848 = dichotoma.
veronicaefolia, Cham. in Linnaea, vii. (1832) 246 = cajanensis.
villosa, Cham. l. c. 247 = commutata.
villosa, *Schau. in DC. Prod.* xi. 570.—Bras.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 197 = indica.
viscidula, *Schau. in DC. Prod.* xi. 570.—Bras.
Zuccagni, Roem. & Schult. Syst. i. 205 = mutabilis.

STACHYURUS, Sieb. & Zucc. Fl. Jap. i. 42. t. 18 (1835). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 184.
himalaicus, *Hook. f. & Thoms. ex Benth. in Journ. Linn. Soc.* v. (1861) 55.—Reg. Himal.
japonicus, Steud. Nom. ed. II. ii. 630, sphalm. = seq.
praecox, *Sieb. & Zucc. Fl. Jap.* i. 43.—Japon.

STACHYUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = **Stachys**, Tourn. (Labiat.).

STACKHOUSIA, Sm. in Trans. Linn. Soc. iv. (1798) 218. *STACKHOUSIEAE*, Benth. & Hook. f. i. 371.
PLOKIOSTIGMA, Schuch. in Linnaea, xxvi. (1853–55) 39.
TRIPTEROCOCCUS, Endl. in Enum. Pl. Hueg. 17 (1837).
aspericocca, Schuch. in Linnaea, xxvi. (1853–55) 12 = monogyna.
Brunonis, *Benth. Fl. Austral.* i. 409.—Austral.
cuneata, A. Cunn. ex Hook. f. Fl. Tasm. i. 79 = spathulata.
doryopetala, Schuch. in Linnaea, xxvi. (1853–55) 24 = viminea.
elata, F. Muell. Fragm. iii. 86 = viminea.
flava, *Hook. f. in Hook. Ic. Pl.* t. 269.—Austral.
Gunniana, Schuch. in Linnaea, xxvi. (1853–55) 18 = monogyna.
Gunnii, Hook. f. Fl. Tasm. i. 79 = monogyna.
Huegelii, *Endl. in Enum. Pl. Hueg.* 17.—Austral.
linariifolia, A. Cunn. in Field, N. S. Wales, 356 = monogyna.
lutea, Herb. Sond. ex Schuch. in Linnaea, xxvi. (1853–55) 8 = monogyna.
maculata, Sieber, ex Hook. f. in Hook. Journ. Bot. ii. (1840) 421 = spathulata.
megaloptera, *F. Muell. Fragm.* viii. 35.—Austral.
minima, *Hook. f. Fl. N. Zel.* i. 47.—N. Zel.
monogyna, *Labill. Nov. Holl. Pl.* i. 77. t. 104.—Austral.
monogyna, Labill. l. c., partim = spathulata.
monogyna, Sieber, ex Benth. Fl. Austral. i. 408 = viminea.
Muelleri, Schuch. in Linnaea, xxvi. (1853–55) 16 = monogyna.

STACKHOUSIA:—
muricata, *Lindl. Bot. Reg.* sub t. 1917.—Austral.
nuda, Lindl. l. c. = viminea.
obtusa, Lindl. l. c. = monogyna.
pubescens, *A. Rich. Sert. Astrolab.* 89. t. 33.—Austral.
pulvinaris, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 101.—Austral.
scoparia, *Benth. Fl. Austral.* i. 409.—Austral.
spathulata, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 124.—Austral.
viminea, *Sm. in Rees, Cycl.* xxxiii. n. 1.—Austral.

STADMANNIA, Lam. Illustr. ii. 443 (1793). *SAPINDACEAE*, Benth. & Hook. f. i. 406.
australis, A. Cunn. ex Hook. Bot. Mag. sub t. 4470 = Diploglottis Cunninghamii.
australis, G. Don, Gen. Syst. i. 669 = Melicocca australis.
bahiensis, *Linden, Cat.* n. 18. (1863) 10.—Bras.
bijuga, D. Dietr. Syn. Pl. ii. 1304 = Melicocca bijuga.
dentata, D. Dietr. l. c. 1305 = Melicocca dentata.
diversifolia, Spreng. Syst. ii. 243 = Doratoxylon diversifolium.
Frazeri, *Linden, Cat.* n. 17 (1861) 10.—Hab. ?
frigida, Linden, l. c. = Giesbrechtii.
Giesbrechtii, *Hort. ex Pasq. Cat. Orto Bot. Nap.* 112, nomen.—Bras. ?
integrifolia, *Linden, Cat.* n. 17 (1862) 10.—Bras.
Jonghii, *Linden, l. c.*—Hab. ?
Liboniana, *Linden, l. c.* n. 18. (1863) 10.—Bras.
macrophylla, *Linden, l. c.*—Bras. ?
olivaeformis, D. Dietr. Syn. Pl. ii. 1305 = Talisia olivaeformis.
oppositifolia, *Lam. Illustr.* ii. 443; *Encyc.* vii. 376.—Ins. Borbon.
paniculata, D. Dietr. Syn. Pl. ii. 1305 = Hypelate paniculata.
pubescens, Spreng. Syst. ii. 243 = Schleichera trijuga.
pulchella, *Linden, Cat.* n. 18. (1863) 10.—Bras.
Sideroxylon, DC. Prod. i. 615 = oppositifolia.
sorbifolia, *Linden, Cat.* n. 18. (1863) 10.—Bras.
trijuga, Spreng. Syst. ii. 243 = Schleichera trijuga.

STADTMANNIA, Walp. Ann. ii. 215 (1851–52), sphalm. = praec.

STAEBE, Hill, Veg. Syst. iii. 145 (1761); iv. 33 (1762) = **Centaurea**, Linn. (Compos.).
aethiopica, Hill, l. c. iii. 145.—Afr. bor.
Cineraria, Hill, Hort. Kew. 65 = C. Cineraria.
paniculata, Hill, l. c. 66 = C. paniculata.
ragusina, *Hill, l. c.* = C. ragusina.

STAEHELINA, Linn. Syst. ed. II. 28 (1740). *COMPOSITAE*, Benth. & Hook. f. ii. 471.
ALKIBIAS, Rafin. Fl. Tellur. iv. 119 (1836).
APLINA, Rafin. l. c. (1836).
BARBELLINA, Cass. in Dict. Sc. Nat. xlvii. 500, 501 (1827).
HIRTELLINA, Cass. l. c. 499, 511 (1827); et l. 441.
OCNERON, Rafin. Fl. Tellur. iv. 120 (1836).
ROCCARDIA, Rafin. l. c. 119 (1836).
alpina, Schramk, in Denkschr. Akad. Muench. vii. 1816–17 (1820) 59 = Saussurea alpina.
apiculata, Labill. Ic. Pl. Syr. Dec. iv. 3. t. 1 = Lobelii.
arborescens, *Linn. Mant.* i. 111.—Ins. Creta; Archip.; Graec.
baetica, *DC. Prod.* vi. 544.—Hispan.
centauroides, Burm. f. Fl. Cap. Prod. 23 = Relhania centauroides.
Chamaepeuce, Linn. Syst. ed. XII. 538 = Cnicus Chamaepeuce.
corymbosa, Thunb. Prod. Pl. Cap. 143 = Vernonia corymbosa.
corymbosa, Wight, ex DC. Prod. vi. 364 = Senecio corymbosus.
cretica, Zucc. ex DC. Prod. vi. 658 = Cnicus Chamaepeuce.
dubia, *Linn. Sp. Pl.* 840.—Europ. austr.
elegans, Walt. Fl. Carol. 202 = Liatris elegans.
fasciculata, Thunb. Prod. Pl. Cap. 143 = Lachnospermum ericifolium.
fimbriata, Forst. ex DC. Prod. vi. 153 = Craspedia uniflora.
fruticosa, *Linn. Syst.* ed. XII. 538.—Ins. Creta.
gnaphaloides, Linn. Sp. Pl. 840 = Helipterum gnaphaloides.
hastata, Vahl, Symb. Bot. i. 70 = Vernonia atriplicifolia.

STAEHELINA:—
ilicifolia, Linn. f. Suppl. 358 = Gynoxys ilicifolia.
imbricata, Berg. Desc. Pl. Cap. 233; Thunb. Prod. Pl. Cap. 143 = Metalasia uniflora.
Lobelii, *DC. in Ann. Mus. Par.* xvi. (1810) 194.—Syria.
pinnata, Lag. ex Pers. Syn. ii. 391 = Jurinea pinnata.
rosmarinifolia, Cass. in Dict. Sc. Nat. l. 439 = dubia.
solidaginoides, Willd. ex DC. Prod. v. 41 = Vernonia scorpioides.
spinosa, Vahl, Symb. Bot. i. 69 = Iphiona juniperifolia.
squarrosa, Rafin. Fl. Tellur. iv. 118 = Helipterum gnaphalioides.
syriaca, Spreng. Syst. iii. 391 = Lobelii.
tenuifolia, Presl, Fl. Sic. p. xxix = dubia ?
unifloculosa, *Sibth. & Sm. Fl. Graec. Prod.* ii. 162.—Graecia.

STAEHELINIA, Hall. Enum. Stirp. Helv. ii. 624 (1742) = **Bartsia**, Linn. (Scrophul.).
alpina, Crantz, Stirp. Austr. ed. II. 294 = B. alpina.

STAELIA, Cham. & Schlecht. in Linnaea, iii. (1828) 364. t. 3. f. 3. *RUBIACEAE*, Benth. & Hook. f. ii. 148.
caespitosa, *Griseb. in Goett. Abh.* xxiv. (1879) 158.—Reg. Argent.
galioides, *DC. Prod.* iv. 573.—Bras.
reflexa, *DC. l. c.*—Bras.
thymoides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 364.—Bras.

STAFLINUS, Rafin. New Fl. Am. iv. 28 (1836) = **Daucus**, Tourn. (Umbell.).
crinitus, Rafin. l. c. = D. crinitus.
setifolius, Rafin. l. c. = D. setifolius.

STAGMARIA, Jack, Desc. Malay. Pl. III. 12 (1823 ?) = **Gluta**, Linn. (Anacard.).
verniciflua, Jack, l. c.; et in Hook. Comp. Bot. Mag. i. (1835) 267 = G. Benghas.
verniciflua, Wall. Cat. n. 980 = Melanorrhoea Wallichii.

STAHLIA, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 255. *LEGUMINOSAE*.
maritima, *Bello, l. c.*—Porto Rico.

STALAGMITES, Spreng. Anleit. ii. II. 784 (1818) = seq.

STALAGMITIS, Murr. in Comm. Gotting. ix. 1787–88 (1789), 173 = **Garcinia**, Linn. (Guttif.).
acida, Cambess. ex Spach, Hist. Vég. Phan. v. 328 = G. cochinchinensis.
Boobicowa, G. Don, Gen. Syst. i. 621 = G. paniculata ?
Cambogia, Pers. Syn. ii. 68 = seq.
cambogioides, Murr. in Comm. Gotting. ix. 1787–88 (1789) 173 = G. spicata.
celebica, G. Don, Gen. Syst. i. 621 = G. cornea.
cochinchinensis, G. Don, l. c. = G. cochinchinensis.
Cowa, G. Don, l. c. = G. Cowa.
dulcis, Cambess. in Mém. Mus. Par. xvi. (1828) 393, 426 = G. dulcis.
elliptica, G. Don, Gen. Syst. i. 621 = G. elliptica.
erosipetala, Miq. in Journ. Bot. Néerl. i. (1861) 126 (sp. dub.).—China.
guineensis, G. Don, Gen. Syst. i. 621 (= *Xanthochymus guineensis*).—Afr. trop.
Gutta, G. Don, l. c. = G. Cambogia.
indica, G. Don, l. c. = G. indica.
javanensis, Spach, Hist. Vég. Phan. v. 328 = G. dulcis.
Kydiana, G. Don, Gen. Syst. i. 621 = G. Cowa.
lamponga, Miq. Fl. Ind. Bat. Suppl. 496 = G. cymulosa.
lanceaefolia, G. Don, Gen. Syst. i. 621 = G. lanceaefolia.
luteus, G. Don, l. c. (= *Xanthochymus luteus*).—Ind. or.
macrophylla, G. Don, l. c. (= *Xanthochymus macrophyllus*).—Ind. or.
nervosa, Miq. Fl. Ind. Bat. Suppl. 496 = G. Andersoni.
ovalifolia, G. Don, Gen. Syst. i. 621 = G. ovalifolia.
paniculata, G. Don, l. c. = G. paniculata.
pictoria, G. Don, l. c. 620 = G. Xanthochymus.
purpurea, G. Don, l. c. 621 = G. indica.
spicata, Walp. Rep. i. 396 = G. ovalifolia.
umbellata, G. Don, Gen. Syst. i. 621.—Ind. or.

STAMMARIUM, Willd. ex DC. Prod. v. 102 (1836) = **Pectis**, Linn. (Compos.).
hyssopifolium, Willd. ex DC. l. c. = P. auricularis.

STANFORDIA, S. Wats. in Bot. Calif. ii. 479 (1880) *CRUCIFERAE*.
californica, *S. Wats. l. c.*—Calif.

STANGERIA, T. Moore, in Hook. Kew Journ. v. (1853) 228. *CYCADACEAE*, Benth. & Hook. f. iii. 446.
Katzeri, *Regel, Gartenfl.* (1874) 163. t. 798.—Afr. austr.
paradoxa, *T. Moore, in Hook. Kew Journ.* v. (1853) 228.—Afr. austr.
schizodon, *Bull. Cat.* (1872) 8.—Afr. austr.

STANGGERIA, Stevens, in Proc. Linn. Soc. ii. (1854) 340, sphalm. = praec.

STANHOPEA, Frost, ex Hook. Bot. Mag. tt. 2948, 2949 (1829). *ORCHIDEAE*, Benth. & Hook. f. iii. 549.
CERATOCHILUS, Lindl. in Lodd. Bot. Cab. t. 1414 (1828).
STANHOPEASTRUM, Reichb. f. in Bot. Zeit. x. (1852) 927.
amoena, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 273 = Wardii.
atropurpurea, Lodd. ex Planch. Hort. Donat. 215 = insignis.
aurantia, Lodd. ex P. N. Don, in Donn, Hort. Cantab. ed. XIII. 608 = S. Bucephalus.
aurata, Beer, Prakt. Orch. 310 = graveolens.
aurea, Lodd. ex Lindl. Bot. Reg. (1841) Misc. 11; Endl. in Harting. Parad. Vindob. t. 31 = Wardii.
bicolor, C. Koch, Berl. Allg. Gartenz. (1857) 209, nomen (Quid ?).—Hab. ?
Bucephalus, *Lindl. Gen. et Sp. Orch.* 157.—Mexic.
calceolata, Hort. ex Reichb. f. Xenia Orch. i. 117 = eburnea.
Calceolus, *Reichb. f. l. c.*—Am. centr.
candida, *Rodrig. Orch. Nov.* i. 101.—Bras.
Cavendishii, Lindl. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 643, nomen = devoniensis ?
cirrhata, *Lindl. in Journ. Hort. Soc.* v. (1850) 37 = Nicaragua.
connata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 226.—Peruv.
costaricensis, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 424.—Costa Rica.
cymbiformis, *Reichb. f. Xenia Orch.* ii. 84. t. 124.—Am. centr.
deltoidea, *Lem. Illustr. Hortic.* ix. (1862), sub t. 340.—Mexic.
devoniensis, *Lindl. Sert. Orch.* t. 1.—Mexic.
eburnea, *Lindl. Bot. Reg.* t. 1529.—Bras.
ecornuta, *Lem. in Fl. des Serres*, ii. (1846) t. 181.—Guatemala.
expansa, P. N. Don, in Donn, Hort. Cantab. ed. XIII. 721; Hort. ex Reichb. f. Xenia Orch. i. 120 = tigrina.
flava, Lodd. ex Beer, Prakt. Orch. 312 = insignis.
florida, *Reichb. f. in Gard. Chron.* (1879) II. 615.—Am. austr.
Fregeana, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 313.—Mexic.
gibbosa, *Reichb. f. in Gard. Chron.* (1869) 1254.—Am. centr.
grandiflora, Lindl. Gen. et Sp. Orch. 158 = eburnea.
grandiflora, Reichb. f. in Walp. Ann. vi. 587 = S. Bucephalus.
graveolens, Klotzsch, ex Reichb. f. Xenia Orch. i. 118 = connata.
graveolens, *Lindl. Bot. Reg.* (1840) *Misc.* 59.—Peruv.
graveolens, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) 55. t. 54 = oculata.
guttata, Beer, Prakt. Orch. 312 = S. Bucephalus.
guttata, C. Koch, Berl. Allg. Gartenz. (1858) 364 = seq.
guttulata, Lindl. Bot. Reg. (1843) Misc. 75 = oculata.
Harrisoniae, P. N. Don, in Donn, Hort. Cantab. ed. XIII. 607 = Bifrenaria Harrisoniae.
Haselowiana, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 322 (Haseloffiana).—Peruv.
implicata, Westcott, ex Lindl. Fol. Orchid. Stanh. 6 = Martiana.
inodora, *Lodd. Cat.* n. 1147; *ex Lindl. Bot. Reg.* (1845) t. 65.—Mexic.

STANHOPEA:—
inodora, Reichb. f. Xenia Orch. ii. 157 = graveolens.
insignis, *Frost, in Hook. Bot. Mag.* tt. 2948, 2949.—Bras.
Jenischiana, Kramer, ex Reichb. f. in Bot. Zeit. x. (1852) 934 = S. Bucephalus.
Lindleyi, Zucc. in Abh. Akad. Muench. ii. (1831-36) 320 = oculata.
lyncea, P. N. Don, in Donn, Hort. Cantab. ed. XIII. 608 = devoniensis *vel* tigrina.
macrochila, *Lem. Illustr. Hortic.* vi. (1859) *Misc.* 71, cum ic.—Mexic.
maculosa, Knowles & Westc. Flor. Cab. iii. (1840) 119. t. 121 = devoniensis.
Marshii, *Reichb. f. Xenia Orch.* i. 120.—Guatemala.
Martiana, *Lindl. Bot. Reg.* (1840) *Misc.* 50.—Mexic.
nigroviolacea, Beer, Prakt. Orch. 313 = tigrina.
oculata, *Lindl. Gen. et Sp. Orch.* 158.—Mexic.
odoratissima, Hort. ex Planch. Hort. Donat. 216 = insignis.
ornatissima, *Lem. Illustr. Hortic.* ix. (1862) t. 325.—Mexic.
platyceras, *Reichb. f. in Gard. Chron.* (1868) 27.—N. Granat.
pulla, *Reichb. f. l. c.* (1877) I. 850.—Costa Rica.
quadricornis, *Lindl. Bot. Reg.* (1838) t. 5.—Am. bor. occ.
radiosa, Lem. Illustr. Hortic. vi. (1859) Misc. 72, cum ic. = saccata.
Reichenbachiana, *Roezl, ex Reichb. f. in Gard. Chron.* (1879) II. 40.—Reg. Andin.
Ruckeri, *Lindl. Bot. Reg.* (1843) sub t. 44.—Nicaragua.
Russelliana, *Lodd. Cat. ex P. N. Don, in Donn, Hort. Cantab.* ed. XIII. 721, nomen.—Bras.
saccata, *Batem. Orch. Mex. & Guatem.* t. 15.—Guatemala.
Schilleriana, *Reichb. f. Xenia Orch.* ii. 158.—Am. centr.?
serrata, Gard. Chron. (1880) I. 779 (Ind. gen.), sphalm. = cirrhata.
Shuttleworthii, *Reichb. f. in Gard. Chron.* (1876) I. 795.—N. Granat.
tigrina, *Batem. Orch. Mex. & Guatem.* t. 7.—Mexic.
tricornis, *Lindl. in Journ. Hort. Soc.* iv. (1849) 263.—Peruv.
velata, C. Morr. in Ann. Soc. Bot. Gand, iii. (1847) 333. t. 153 = Martiana.
venusta, Lindl. Bot. Reg. (1841) Misc. 11, in obs. = Wardii.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 109.—N. Granat.
Wardii, *Lodd. ex Lindl. Sert. Orch.* t. 20.—Mexic.
Warszewicziana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xx. (1852) 274.—Am. centr.
xytriophora, *Reichb. f. in Gard. Chron.* (1868) 842.—Peruv.

STANHOPEASTRUM, Reichb. f. in Bot. Zeit. x. (1852) 927 = **Stanhopea**, Frost (Orchid.).
ecornutum, Reichb. f. l. c. = Stanhopea ecornuta.

STANLEYA, Nutt. Gen. Am. ii. 21 (1818). *CRUCIFERAE*, Benth. & Hook. f. i. 80.
PODOLOBUS, Rafin. in Am. Monthly Mag. (1819) 194.
amplexifolia, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 297 = Warea amplexifolia.
cartilaginea, Spreng. Syst. ii. 909 = Sisymbrium junceum.
fruticosa, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 180 = pinnatifida.
gracilis, DC. Syst. ii. 512 = Warea cuneifolia.
heterophylla, Nutt. in Torr. & Gray, Fl. N. Am. i. 97 = pinnatifida.
integrifolia, James, in Long's Exped. ii. 17; ex Torr. in Ann. Lyc. N. York, ii. (1828) 166 = pinnatifida.
pinnatifida, *Nutt. Gen. Am.* ii. 71.—Am. bor.
tomentosa, *Parry, in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
viridiflora, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 98.—Am. bor. occ.
Washitana, DC. Syst. ii. 512 = Streptanthus maculatus?

STANNIA, Karst. Ausw. Neuer Gew. Venez. 27 (1848) = **Posoqueria**, Aubl. (Rubiac.).
formosa, Karst. l. c. t. 9 = P. formosa.
grandiflora, Karst. Fl. Columb. i. 31. t. 16.—N. Granat.
metensis, Karst. in Linnaea, xxviii. (1856) 441.—Venezuela.
panamensis, Walp. & Duchass. in Linnaea, xxiii. (1850) 755 = P. latifolia.
spectabilis, Klotzsch, ex Walp. & Duchass. l. c., in obs.—Hab.?

STAPELIA, Linn. Gen. ed. I. 64 (1737). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 784.
CARUNCULARIA, Haw. Syn. Pl. Succ. 333 (1812).
GONOSTEMON, Haw. l. c. 27 (1812).
ORBEA, Haw. l. c. 37 (1812).
PODANTHES, Haw. l. c. 32 (1812).
TRIDENTEA, Haw. l. c. 34 (1812).
TROMOTRICHE, Haw. l. c. 36 (1812).
acuminata, *Mass. Stapel.* 15. t. 17.—Afr. austr.
adscendens, Roxb. Pl. Corom. i. 28. t. 30 = Caralluma adscendens.
ambigua, *Mass. Stapel.* 13. t. 12.—Afr. austr.
Ango, A. Rich. Tent. Fl. Abyss. ii. 50 (Carallumae sp.?).—Abyss.
anguinea, Jacq. Stapel. t. 37 = picta.
angulata, *Todaro, Hort. Bot. Panorm.* i. 54. t. 13. f. 3.—Afr. austr.
anonymos, Poir. Encyc. vii. 391 = *multangula*.
aperta, *Mass. Stapel.* 23. t. 37.—Afr. austr.
arida, *Mass. Stapel.* 21. t. 33.—Afr. austr.
articulata, Ait. Hort. Kew. ed. I. i. 310 = Pectinaria articulata.
Asterias, *Mass. Stapel.* 14. t. 14.—Afr. austr.
atrata, *Todaro, Hort. Bot. Panorm.* i. 50. t. 13. f. 1.—Afr. austr.?
atropurpurea, *Salm-Dyck, Hort. Dyck.* 372.—Afr. austr.?
barbata, Mass. Stapel. 11. t. 7 = Huernia barbata.
Bayfieldii, *N. E. Br. in Gard. Chron.* (1877) I. 431. f. 66.—Hab.?
beffoniana, Hort. ex Schult. Syst. vi. 49 = variegata.
bidentata, Hort. ex Salm-Dyck, Hort. Dyck. 266, nomen = variegata.
bifolia, Hort. ex Schult. Syst. vi. 49 (Quid?).—Hab.?
bisulca, *Donn, Hort. Cantab.* ed. III. 43, nomen; *et ex Schult. Syst.* vi. 37.—Afr. austr.
brevirostris, Willd. Enum. Hort. Berol. Suppl. 14, nomen (Quid?).—Hab.?
bufonia, Jacq. Stapel. t. 35 = variegata.
bufonis, Sims, Bot. Mag. t. 1676 = normalis.
cactiformis, Hook. l. c. t. 4127 = *clavata*, Willd.
caespitosa, Mass. Stapel. 20. t. 29 = Duvalia caespitosa.
Callamulia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 77 = Boucerosia umbellata.
campanulata, Mass. Stapel. 11 = Huernia campanulata.
campanulata, Pav. ex Planch. in Fl. des Serres, viii. (1852-53) 55, sub t. 770 = Dictyanthus Pavonii.
campanulata, Sims, Bot. Mag. t. 1661 = Huernia clavigera.
canescens, *Hort. ex Haw. Syn. Succ. Pl.* 26.—Afr. austr.
caudata, Thunb. Prod. Pl. Cap. 46 = Brachystelma crispum.
chinensis, Lour. Fl. Cochinch. 165 = Hoya carnosa.
ciliata, Thunb. Prod. Pl. Cap. 46 = Diplocyatha ciliata.
ciliolata, Todaro, Ind. Sem. Hort. Panorm. (1861) 10, nomen (Quid?).—Afr. austr.
clavata, Decne. in DC. Prod. viii. 664, sphalm. = *clavigera*.
clavata, Willd. Sp. Pl. i. 1295 (Trichocaulonis sp.).—Afr. austr.
clavigera, Jacq. Stapel. t. 5 = Huernia clavigera.
clypeata, *Jacq. l. c.* t. 34.—Afr. austr.
cochinchinensis, Lour. Fl. Cochinch. 165 = Hoya cochinchinensis.
comata, *Jacq. Stapel.* t. 49.—Afr. austr.
compacta, Schult. Syst. vi. 46 = Duvalia compacta.
concinna, *Mass. Stapel.* 15. t. 18.—Afr. austr.
concolor, Salm-Dyck, Hort. Dyck. 372 (Duvaliae sp.).—Afr. austr.?
congestiflora, *Delile, in Rochet, Sec. Voy. Choa*, 345.—Afr. trop.
conspurcata, *Willd. Enum. Hort. Berol.* 284.—Afr. austr.

STAPELIA:—
cordata, *Hort. ex Haw. Syn. Pl. Succ.* 26.—Afr. austr.
Corderoyi, Hook. f. Bot. Mag. t. 6082 = Duvalia Corderoyi.
Courcelli, Hort. ex N. E. Br. in Gard. Chron. (1877) I. 140 = patentirostris.
crassa, Donn, Hort. Cantab. ed. III. 43 = Huernia tubata.
crispa, Hort. ex Haw. Syn. Pl. Succ. 31 = Huernia crispa.
Curtisii, Schult. Syst. vi. 38 = variegata.
cylindrica, Hort. ex Hook. f. Bot. Mag. t. 5930 = Echidnopsis cereiformis.
cymosa, Hort. ex Schult. Syst. vi. 49 = Duvalia hirtella.
decora, Mass. Stapel. 19. t. 26 = Piaranthus decorus.
deflexa, Hort. ex Haw. Syn. Pl. Succ. 18 = reflexa.
deflexa, Hort. Lips. (1864); ex Todaro, Hort. Bot. Panorm. i. 52 = trisulca.
deflexa, *Jacq. Stapel.* t. 20.—Afr. austr.
dejecta, *Salm-Dyck, Hort. Dyck.* 372.—Afr. austr.
dentata, Forsk. Fl. Aegypt. Arab. p. cviii. (Boucerosiae sp.).—Arab.
depressa, Hort. ex N. E. Br. in Gard. Chron. (1876) II. 809, in obs. = glabriflora.
depressa, Jacq. Stapel. t. 55 = patula.
Desmidorchis, Steud. Nom. ed. II. ii. 631 = Boucerosia Russeliana.
discolor, Todaro, Hort. Bot. Panorm. i. 49. t. 12. f. 3.—Afr. austr.?
divaricata, *Mass. Stapel.* 17. t. 22.—Afr. austr.
duodecimfida, Jacq. Stapel. t. 4 = Huernia tubata.
elegans, Mass. Stapel. 19. t. 27 = Duvalia elegans.
elongata, Sweet, Hort. Brit. ed. II. 357 = patula.
emarginata, Breit. Hort. Breit. 504, nomen (Quid?).—Hab.?
erivacaria, Roxb. Hort. Beng. 21, nomen (Quid?).—Hab.?
eruciformis, Hort. ex N. E. Br. in Gard. Chron. (1875) I. 136 = olivacea.
europaea, Guss. Fl. Sic. Prod. Suppl. i. 65 (1832); et in Atti Acc. Sc. Nap. iv. (1839) (Bot.) 87 = Boucerosia Gussoniana.
fasciculata, *Thunb. Prod. Pl. Cap.* 46.—Afr. austr.
fissirostris, *Jacq. Stapel.* t. 23.—Afr. austr.
flavicomata, *Haw. Suppl. Pl. Succ.* 8.—Afr. austr.
flavirostris, N. E. Br. in Gard. Chron. (1877) I. 558 = grandiflora.
fuscata, Hort. Parm. ex Todaro, Hort. Bot. Panorm. i. 47 = mutabilis.
fuscata, *Jacq. Stapel.* t. 46.—Afr. austr.
geminata, Mass. Stapel. 18. t. 25 = Piaranthus geminatus.
gemmifera, Hort. ex Salm-Dyck, Hort. Dyck. 266, nomen = seq.
gemmiflora, *Mass. Stapel.* 14. t. 15.—Afr. austr.
gigantea, *N. E. Br. in Gard. Chron.* (1877) I. 684. f. 112.—Afr. austr.
glabriflora, *N. E. Br. l. c.* (1876) II. 809. f. 149.—Afr. austr.?
glandulifera, Pers. Syn. i. 279 = seq.
glanduliflora, *Mass. Stapel.* 16. t. 19.—Afr. austr.
glauca, *Jacq. Stapel.* t. 44.—Afr. austr.
glomerata, Hort. ex Haw. Syn. Pl. Succ. 46 = Duvalia glomerata.
Gordoni, Mass. Stapel. 24. t. 40 = Hoodia Gordoni.
grandiflora, Curt. Bot. Mag. t. 585 = spectabilis.
grandiflora, Jacq. Stapel. t. 59 = obscura.
grandiflora, *Mass. Stapel.* 13. t. 11.—Afr. austr.
Gussoneana, Jacq. ex Lindl. Bot. Reg. t. 1731 = Boucerosia Gussoniana.
guttata, Mass. Stapel. 10 = Huernia guttata.
hamata, *Jacq. Stapel.* t. 50.—Afr. austr.
hircosa, *Jacq. l. c.* 25.—Afr. austr.
hirsuta, *Linn. Sp. Pl.* 217.—Afr. austr.
hirtella, Jacq. Stapel. t. 10 = Duvalia hirtella.
hispidula, Hornem. Hort. Hafn. i. 251 = glanduliflora.
humilis, Mass. Stapel. 10. t. 5 = Huernia humilis.
Hystrix, Hook. f. Bot. Mag. t. 5751 = Huernia Hystrix.
incarnata, Linn. f. Suppl. 171 = Boucerosia incarnata.
inodora, *Decne. in DC. Prod.* viii. 661.—Afr. austr.
involucrata, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 47 = Heterostemma tanjorense.
irrorata, Lodd. Bot. Cab. t. 127 = verrucosa.
irrorata, *Mass. Stapel.* 12. t. 9.—Afr. austr.

STAPELIA:—
italica, Tenore, ex Salm-Dyck, Hort. Dyck. 300, nomen = Boucerosia Gussoniana.
Jacquiniana, Schult. Syst. vi. 45 = Duvalia Jacquiniana.
Juvencula, *Jacq. Stapel.* t. 28.—Afr. austr.
laevigata, Hort. ex Haw. Syn. Pl. Succ. 46 = Duvalia laevigata.
laevis, Decne. in DC. Prod. viii. 658 = pedunculata.
lampadosa, Jacq. ex Salm-Dyck, Hort. Dyck. 300, nomen = Boucerosia Gussoniana.
lanifera, *Haw. Suppl. Pl. Succ.* 8.—Afr. austr.
lentiginosa, Sims, Bot. Mag. t. 506 = Huernia guttata.
lepida, *Jacq. Stapel.* t. 43.—Afr. austr.
limosa, Hort. ex Salm-Dyck, Hort. Dyck. 266 = praec.
lucida, *DC. Cat. Hort. Monsp.* (1813) 148.—Afr. austr.
lunata, Sweet, Hort. Brit. ed. II. 357 = uncinata.
macrocarpa, *A. Rich. Tent. Fl. Abyss.* ii. 50.—Abyss.
maculosa, *Jacq. Stapel.* t. 31.—Afr. austr.
mammillaris, Linn. Mant. ii. 216 = Boucerosia mammillaris.
marginata, *Willd. Enum. Hort. Berol. Suppl.* 13.—Afr. austr.
marmorata, *Jacq. Stapel.* t. 38.—Afr. austr.
Massoni, *Haw. Syn. Pl. Succ.* 18.—*Cf. N. E. Br. in Gard. Chron.* (1883) II. 761.—Afr. austr.
mastodes, Jacq. Stapel. t. 13 = Duvalia mastodes.
mastodis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
meliflua, Blanco, Fl. Filip. ed. I. 202 = Hoya diversifolia.
mixta, Donn, Hort. Cantab. ed. IV. 53, nomen = maculosa.
mixta, *Mass. Stapel.* 23. t. 38.—Afr. austr.
monstrosa, Hort. ex Steud. Nom. ed. II. ii. 631 = variegata.
moschata, Donn, Hort. Cantab. ed. III. 43 = hircosa.
multangula, Forsk. Fl. Aegypt. Arab. p. cviii (Echidnopsidis sp.?).—Arab.
multiflora, *DC. Cat. Hort. Monsp.* (1813) 149.—Afr. austr.
mutabilis, *Jacq. Stapel.* t. 29.—Afr. austr.?
namaquensis, *N. E. Br. in Gard. Chron.* (1882) II. 648.—Afr. austr.
neglecta, Todaro, Hort. Bot. Panorm. i. 47 = mutabilis.
normalis, *Jacq. Stapel.* t. 42.—Afr. austr.
obliqua, *Willd. Enum. Hort. Berol. Suppl.* 13.—Afr. austr.
obscura, *N. E. Br. in Gard. Chron.* (1877) I. 558, in obs.—Afr. austr.
ocellata, Jacq. Stapel. t. 6 = Huernia ocellata.
olivacea, *N. E. Br. in Gard. Chron.* (1875) I. 136.—Afr. austr.
ophiuncula, *Haw. Syn. Pl. Succ.* 27.—Afr. austr.
orbicularis, *Andr. Bot. Rep.* t. 439.—Afr. austr.
orbiculata, Donn, Hort. Cantab. ed. III. 43, nomen = praec.
pallens, Hort. ex Steud. Nom. ed. II. ii. 631 = pallida.
pallida, *Wendl. Coll. Bot.* 39. t. 51.—Afr. austr.
paniculata, *Willd. Enum. Hort. Berol. Suppl.* 13.—Afr. austr.
papillosa, DC. Pl. Gr. ex Desf. Tabl. ed. II. 92, nomen (Quid?).—Afr. austr.
parviflora, Mass. Stapel. 21 (Carallumae sp.).—Afr. austr.
Passerini, Todaro, Hort. Bot. Panorm. i. 47, nomen = mutabilis.
patentirostris, *N. E. Br. in Gard. Chron.* (1877) I. 140. f. 21.—Afr. austr.
patula, *Willd. Enum. Hort. Berol.* 281.—Afr. austr.
pedunculata, *Mass. Stapel.* 17. t. 21.—Afr. austr.
penduliflora, Steud. Nom. ed. II. ii. 631 = pedunculata.
picta, *Donn, Hort. Cantab.* ed. III. 43, nomen; *Bot. Mag.* t. 1169.—Hab.?
pilifera, Linn. f. Suppl. 171 = Trichocaulon piliferum.
planiflora, *Jacq. Stapel.* t. 40.—Afr. austr.
Plantii, *Hort. ex Hook. f. Bot. Mag.* t. 5692.—Afr. austr.
Plantii, MacKen, ex N. E. Br. in Gard. Chron. (1877) I. 684 = gigantea.
plicata, Hort. ex Salm-Dyck, Hort. Dyck. 267, nomen (Quid?).—Hab.?

STAPELIA:—
protensa, Hornem. Suppl. Hort. Hafn. 30 = revoluta.
pruinosa, *Mass. Stapel.* 24. t. 41.—Afr. austr.
pulchella, Hort. ex Haw. Syn. Pl. Succ. 33 = roriflua.
pulchella, *Mass. Stapel.* 22. t. 36.—Afr. austr.
pulchra, Schult. Syst. vi. 28 = roriflua.
pulla, Ait. Hort. Kew. ed. I. i. 310 = Boucerosia mammillaris.
pulvinata, Donn, Hort. Cantab. ed. IV. 53 = lanifera.
pulvinata, *Mass. Stapel.* 13. t. 13.—Afr. austr.
punctata, *Mass. l. c.* 18.—Afr. austr.
quadrangula, Blanco, Fl. Filip. ed. I. 202 = Heterostemma cuspidatum.
quadrangula, Forsk. Fl. Aegypt. Arab. 51 = Boucerosia quadrangula.
quadrangula, Schousb. Gew. Marok. 107 = Boucerosia maroccana.
quinquenervis, Schult. Syst. vi. 37 = planiflora.
radiata, Jacq. Stapel. t. 12 = Duvalia Jacquiniana.
radiata, Sims, Bot. Mag. t. 619 = Duvalia radiata.
ramosa, *Mass. Stapel.* 21. t. 32.—Afr. austr.
reclinata, Mass. l. c. 19. t. 28 = Duvalia reclinata.
reclinata, Sims, Bot. Mag. t. 1397 = Duvalia hirtella.
reflexa, *Haw. Syn. Pl. Succ.* 18.—Afr. austr.
replicata, Jacq. Stapel. t. 15 = Duvalia replicata.
reticulata, Mass. Stapel. 9. t. 2 = Huernia reticulata.
retusa, *Schult. Syst.* vi. 41.—Afr. austr.
revoluta, *Mass. Stapel.* 12.—Afr. austr.
revoluta, Sims, Bot. Mag. t. 724 = tigridia.
roriflua, *Jacq. Stapel.* t. 19.—Afr. austr.
rufa, *Mass. Stapel.* 16. t. 20.—Afr. austr.
rufescens, *Salm-Dyck, Hort. Dyck.* 373.—Afr. austr.
rugosa, *Jacq. Stapel.* t. 41.—Afr. austr.
rugosa, Wendl. Coll. Bot. ii. 41. t. 52 = Wendlandiana.
sagittata, Rottb. in Act. Lit. Univ. Hafn. i. (1778) 288 (Cynanchi sp.).—Guian.
sanguinea, Hort. ex Pasq. Cat. Orto Bot. Nap. 99, nomen (Quid?).—Hab.?
sarmentosa, Steud. Nom. ed. II. ii. 632 = Ceropegia variegata.
scutellata, Todaro, Hort. Bot. Panorm. i. 52. t. 13. f. 2 = trisulca.
serrulata, Jacq. Stapel. t. 17 = Piaranthus serrulatus.
Simsii, *Schult. Syst.* vi. 33.—Afr. austr.
sororia, Hook. f. Bot. Mag. t. 5963 = patentirostris.
sororia, Jacq. Stapel. tt. 56, 57 = patula.
sororia, *Mass. Stapel.* 23. t. 39.—Afr. austr.
spectabilis, *Haw. Syn. Pl. Succ.* 16.—Afr. austr.
stellaris, *Haw. l. c.* 19; *Suppl.* 9; *Jacq. Stapel.* t. 48.—Afr. austr.
stellata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Asterias.
stricta, *Sims, Bot. Mag.* t. 2037.—Afr. austr.
stygia, *Schult. Syst.* vi. 32 (err. typ. 562).—Afr. austr.
subulata, Forsk. Fl. Aegypt. Arab. p. cviii = Caralluma subulata.
tigridia, *Decne. in DC. Prod.* viii. 657.—Afr. austr.
tridentata, Hort. ex Schult. Syst. vi. 49, nomen (Quid?).—Hab.?
trifida, *Todaro, Hort. Bot. Panorm.* i. 45. t. 12. f. 1.—Afr. austr.
trifolia, Breit. Hort. Breit. 505, nomen (Quid?).—Hab.?
trisulca, *Donn, Hort. Cantab.* ed. III. 43; *Jacq. Stapel.* t. 33.—Afr. austr.
tsomoënsis, *N. E. Br. in Gard. Chron.* (1882) II. 168.—Afr. austr.
tubata, Jacq. Stapel. t. 3 = Huernia tubata.
tuberculata, Hort. ex Haw. Syn. Pl. Succ. 46 = Duvalia radiata.
tuberosa, Meerb. Pl. Rar. sub t. 54. f. 1 = Brachystelma tuberosum.
tubulosa, Hort. ex Steud. Nom. ed. II. ii. 632 = Huernia tubata.
umbellata, Donn, Hort. Cantab. ed. IV. 54; Roxb. Hort. Beng. 21; Pl. Corom. iii. 36. t. 241 = Boucerosia umbellata.
umbilicata, Thuret, Choix Gr. n. 2 (1869), 6, nomen; ex Todaro, Hort. Bot. Panorm. i. 47 = mutabilis.
uncinata, *Jacq. Stapel. Disp. tab.*—Afr. austr.
unguipetala, *N. E. Br. in Gard. Chron.* (1877) I. 334. f. 54.—Afr. austr.
variegata, Forsk. Fl. Aegypt. Arab. 51 = Ceropegia variegata.
variegata, *Linn. Sp. Pl.* 217; *Jacq. Misc.* i. 27. t. 4.—Afr. austr.
venusta, Mass. Stapel. 10 = Huernia venusta.
verrucosa, Hort. ex Haw. Syn. Pl. Succ. 32 = roriflua.

STAPELIA:—
verrucosa, *Mass. Stapel.* 11. t. 8.—Afr. austr.
verticillata, Hort. ex Schult. Syst. vi. 49, nomen (Quid?).—Hab.?
vetula, *Mass. Stapel.* 15. t. 16.—Afr. austr.
vetula, Sims, Bot. Mag. t. 1234 = Simsii.
virgata, Hort. ex Schult. Syst. vi. 49, nomen (Quid?).—Hab.?
Wendlandiana, *Schult. Syst.* vi. 39.—Afr. austr.
Woodfordiana, Hort. ex Haw. Syn. Pl. Succ. 42 = picta.

STAPHIDIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 325 = **Sagraea**, DC. (Melastom.).
aphananthum, Naud. l. c. 333 = Clidemia aphanantha.
attenuatum, Naud. l. c. 332 = Sag. attenuata.
Berterii, Naud. l. c. 328 = Sag. plumosa.
Bonplandii, Naud. l. c. 333 = Sag. Bonplandii.
capilliflorum, Naud. l. c. 330 = Sag. capilliflora.
cognatum, Naud. l. c. 334 = Clidemia rubra.
coriaceum, Naud. l. c. 329 = Sag. coriacea.
fasciculare, Naud. l. c. 333 = Henriettella fascicularis.
impetiolare, Naud. l. c. 329 = Calophysa pilosa.
latifolium, Naud. l. c. 326 = Sag. latifolia.
obliquum, Griseb. in Lechl. Berb. Am. Austr. 59 = Sag. obliqua.
pilosum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 334 = Sag. pilosa.
platyphyllum, Naud. l. c. 331 = Sag. platyphylla.
polystachyum, Naud. l. c. 327 = Sag. polystachya.
pulverulentum, Naud. l. c. 334 = Oxymeris pulverulenta.
rariflorum, Naud. l. c. 334 = Sag. rariflora.
rubrinerve, Naud. l. c. 328 = Sag. domingensis.
rubrum, Naud. l. c. 331 = Clidemia rubra.
scabridum, Naud. l. c. 334 = Henriettella fascicularis.
umbrosum, Naud. l. c. 327 = Sag. umbrosa.

STAPHIDIUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 305 = **Clidemia**, D. Don (Melastom.).
affine, Naud. l. c. 313 = C. novemnervia.
agreste, Naud. l. c. 324 = Leandra agrestis.
anceps, Naud. l. c. 312 = C. tiliaefolia.
astrotrichum, Naud. l. c. 324 = Miconia astrotricha.
attenuatum, Griseb. in Goett. Abh. vii. (1857) 218 = Sagraea attenuata.
Benthamianum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 324 = C. hirta.
biserratum, Naud. l. c. 315 = C. biserrata.
blepharodes, Wawra, Bot. Ergeb. Maxim. Bras. 21. t. 38 = C. blepharodes.
brachystephanum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 309 = C. brachystephana.
bracteosum, Naud. l. c. 318 = C. dependens.
chinantlanum, Naud. l. c. 318 = C. chinatlana.
chrysanthum, Naud. l. c. 310 = C. hirta.
confertiflorum, Naud. l. c. 318 = C. dependens.
conglomeratum, Naud. l. c. 321 = C. conglomerata.
cuneatum, Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 113 = Miconia cuneata.
dependens, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 319 = C. Deppeana.
dispar, Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Calophysa dispar.
divaricatum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 315 = C. serrulata.
diversifolium, Naud. l. c. 322 = Octopleura diversifolia.
elegans, Naud. l. c. 307 = C. hirta.
Galeottii, Naud. l. c. 319 = C. laxiflora.
glabratum, Naud. l. c. 324 = Miconia ibaguensis.
gracile, Naud. l. c. 314 = C. petiolaris.
Hostmannii, Naud. l. c. 310 = C. hirta.
involucratum, Naud. l. c. 323 = C. involucrata.
lanatum, Naud. l. c. 324 = Miconia lanata.
lappaceum, Naud. l. c. = Miconia lappacea.
latifolium, Griseb. in Goett. Abh. vii. (1857) 218 = Sagraea latifolia.
Lindenianum, Naud. in Ann. Sc. Nat. Sér. III. xv.i. (1852) 314 = C. petiolaris.
lucidum, Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 134 = Heterotrichum lucidum.
minutiflorum, Berg, ex Triana, l. c. 137 = Sagraea minutiflora.
mutabile, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 324 = Miconia mutabilis.
naevulum, Naud. l. c. 323 = C. naevula.
novemnervium, Naud. l. c. 312 = C. novemnervia.

STAPHIDIUM :—
octonum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 316 = Heterotrichum octonum.
oxyurum, Naud. l. c. 321 = Clidemia oxyura.
paraguayense, Naud. l. c. 324 = Oxymeris paraguayensis.
pauciflorum, Naud. l. c. 308 = C. hirta.
petiolare, Naud. l. c. 324 = C. petiolaris.
petiolatum, Naud. l. c. = C. Deppeana.
pliostemon, Naud. l. c. 313 = C. rariflora.
polyandrum, Griseb. in Lechl. Berb. Am. Austr. 59 = Miconia Benthamiana.
purpureum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 309 = C. dentata.
reversum, Naud. l. c. 324 = Oxymeris reversa.
sessiliflorum, Naud. l. c. 311 = Sagraea sessiliflora.
spicatum, Naud. l. c. 316 = C. dependens.
strigosum, Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 134 = Heterotrichum strigosum.
tiliaefolium, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 324 = C. tiliaefolia.
triste, Naud. l. c. 320 = Miconia bracteata, *Triana*.
umbrosum, Griseb. in Goett. Abh. vii. (1857) 218 = Sagraea umbrosa.
urceolatum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 316 = C. neglecta.
Wilsonii, Naud. l. c. 311 = C. erythropogon.

STAPHILEA, Medic. Pfl. Anat. 401 (1801), sphalm. = seq.

STAPHYLAEA, Linn. Syst. ed. I (1735) = seq.

STAPHYLEA, Linn. Sp. Pl. ed. I. 270 (1753). *SAPINDACEAE*, Benth. & Hook. f. i. 412.
BUMALDA, Thunb. Nov. Gen. iii. 62 (1783).
STAPHYLODENDRON, Tourn. ex Hall. Enum. Stirp. Helv. i. 423 (1742).
Bolanderi, *A. Gray, in Proc. Am. Acad.* x. (1874) 69.—Calif.
Bumalda, *DC. Prod.* ii. 2.—Japon.
chrysodonta, *Fungh. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 313.—Japon.
colchica, *Stev. in Bull. Soc. Nat. Mosc.* xxi. (1848) II. 276.—Reg. Cauc.
corymbosa, DC. Prod. ii. 3 = Turpinia occidentalis.
Emodi, *Wall. Cat.* n. 4275.—Reg. Himal.
geniculata, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 22.—Calif.
heterophylla, *Ruiz & Pav. Fl. Per.* iii. 29. t. 253.—Peruv.
indica, Burm. f. Fl. Ind. 75. t. 24. f. 2 = Leea sambucina.
montana, Roxb. ex Wight & Arn. Prod. 160 = Gymnosporia montana.
occidentalis, Sw. Prod. Veg. Ind. Occ. 55 = Turpinia occidentalis.
pinnata, *Linn. Sp. Pl.* 270.—Europ.; Reg. Cauc.
pinnatifida, Gueldenst. Reisen, i. 421 = praec.
simplicifolia, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 309 = Turpinia arguta.
trifolia, *Linn. Sp. Pl.* 270.—Am. bor.
trifoliata, Payer, Organ. t. 36 = praec.
vepretum, Roxb. ex Wight & Arn. Prod. 160 = Gymnosporia emarginata.

STAPHYLIS, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = praec.

STAPHYLLAEA, Scop. Introd. 253 (1797), sphalm. = praec.

STAPHYLLODENDRON, Scop. Fl. Carn. ed. II. ii. 223 (1772), sphalm. = seq.

STAPHYLODENDRON, Tourn. ex Hall. Enum. Stirp. Helv. i. 423 (1742) = **Staphylea**, Linn. (Sapind.).
pinnatum, Crantz, Inst. ii. 438 = Staphylea pinnata.
trifoliatum, Crantz, l. c. = Staphylea trifolia.

STAPHYLODENDRUM, Moench, Meth. 64 (1794) = praec.

STAPHYLORHODOS, Turcz. in Bull. Soc. Bot. Mosc. (1862) II. 321. Genus dubium *ROSACEARUM*, Benth. & Hook. f. i. 606.
Cotoneaster, Turcz. l. c. 322.—'N. Zel.'

STAPHYLOSYCE, Hook. f. in Benth. & Hook. f. Gen. i. 828 (1867) = **Physedra**, Hook. f. (Cucurb.).
Barteri, Hook. f. l. c. = P. Barteri.

STAPHYSAGRIA, Spach, Hist. Vég. Phan. vii. 347 (1839) = **Delphinium**, Tourn. (Ranunc.).

STARBIA, Thou. Gen. Nov. Madag. 7 (1806) = **Alectra**, Thunb. (Scrophul.).
madagascariensis, Steud. Nom. ed. II. ii. 632.—Madag.

STARKEA, Willd. Sp. Pl. iii. 2216 (1804) = **Liabum**, Adans. (Compos.).
pinnata, Nutt. Gen. Am. ii. 169 = Aplopappus spinulosus.
umbellata, Willd. Sp. Pl. iii. 2216 = L. Brownei.

STARKIA, Juss. ex Stend. Nom. ed. II. ii. 632 (1841), sphalm. = praec.

STATICE, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 88 (1737), partim. *PLUMBAGINEAE*, Benth. & Hook. f. ii. 625.
EURYCHITON, J. Grah. Cat. Pl. Bomb. Addend. (1839).
GONIOLIMON, Boiss. in DC. Prod. xii. 632 (1848).
LIMONIUM, Tourn. ex Adans. Fam. ii. 283 (1763).
PLEGORHIZA, Molina, Sagg. Chile, ed. I. 164, 351 (1782).
TAXANTHEMA, Neck. Elem. i. 115 (1790); R. Br. Prod. 426 (1810).
acerosa, Bieb. Fl. Taur. Cauc. i. 252 = Acantholimon lepturoides.
acerosa, Hohen. ex Boiss. Diagn. Ser. I. vii. 78 = Acantholimon tenuiflorum.
acerosa, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 420 = Acantholimon acerosum.
aciphylla, Jaub. & Spach, Illustr. i. 162 = Acantholimon Hohenackeri.
acutifolia, Ehrenb. ex Boiss. in DC. Prod. xii. 636 = Thouini.
acutifolia, Reichb. Ic. Fl. Germ. iii. 23 = minuta, *Linn.*
adensis, *Boiss. in DC. Prod.* xii. 670.—Arab.
aegyptiaca, Pers. Syn. i. 334 = Thouini.
airoides, Tausch, in Syll. Ratisb. ii. (1828) 255 = bellidifolia.
alata, Willd. Enum. Hort. Berol. Suppl. 15 = Thouini.
alatavica, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 259.—Turkestan.
albicans, Ledeb. Fl. Ross. iii. 465 = callicoma.
albida, Guss. Fl. Sic. Syn. i. 369 = psiloclada.
alliacea, Cav. Ic. ii. 6. t. 109 = Armeria allioides.
alliacea, Guss. ex Nym. Consp. 614 = Armeria macropoda.
alliacea, Sibth. & Sm. Fl. Graec. iii. t. 294 = Armeria majellensis.
alliacea, Willd. Sp. Pl. i. 1523 = Armeria plantaginea.
alpina, Poir. Encyc. Suppl. v. 234 = Armeria alpina.
altaica, *Hort. ex G. Don, in Loud. Hort. Brit.* 115, nomen.—Sibir.
alutacea, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) I. 367.—Rossia.
anceps, *Regel, Pl. Nov. Fedsch.* 74.—Turkestan.
androsacea, Jaub. & Spach, Illustr. i. 161, 164 = Acantholimon androsaceum.
angustifolia, Tausch, in Syll. Ratisb. ii. (1828) 254 = S. Limonium.
annulata, Spreng. Syst. i. 960 = Aegialitis annulata.
aphylla, Forsk. Fl. Aegypt. Arab. 60 = pruinosa.
aphylla, Poir. Encyc. vii. 408 = decipiens.
arabica, *Jaub. & Spach, Illustr.* i. 156. t. 85.—Arab.
Ararati, *Hort. Angl. ex Planch. in Fl. des Serres*, vii. (1851–52) 95.—Armen.
arborea, Brouss. ex Willd. Enum. Hort. Berol. 337 = fruticans.
arborescens, Brouss. Elench. Hort. Monsp. (1805) 58, nomen = fruticans.
Arbuscula, *Maxim. in Trautv. Regel, Maxim. & Winkl. Decas Pl. Nov.* 8.—Japon.
arenaria, Pers. Syn. i. 332 = Armeria plantaginea.
aretiaefolia, Fries, ex Boiss. in DC. Prod. xii. 657 = linifolia.
argentea, Pall. ex Siev. in Pall. N. Nord. Beitr. vii. 282 = callicoma.
aristata, Sibth. & Sm. Fl. Graec. Prod. i. 213 = echioides.
Armeria, Brot. Fl. Lusit. i. 486 = Armeria Welwitschii.

STATICE :—
Armeria, Hook. Fl. Bor. Am. i. 123 = Armeria sanguinolenta.
Armeria, Hook. f. Fl. Antarct. ii. 339 = Armeria macloviana.
Armeria, Linn. Sp. Pl. 274 = Armeria maritima, elongata.
Armeria, Loisel. Fl. Gall. ed. II. i. 223 = Armeria plantaginea.
Armeria, Scop. Fl. Carn. ed. II. i. 227 = Armeria alpina.
Armeria, Sm. Engl. Bot. t. 226 = Armeria pubescens.
articulata, *Loisel. Fl. Gall.* ed. II. i. 225. t. 6.—Europ. austr.
articulata, Schimp. ex Nym. Consp. 612 = cancellata.
articulata, Vis. Fl. Dalm. ii. 7 = cosyrensis.
aspera, Lam. Fl. Fr. iii. 64 = echioides.
attenuata, *Schult. Syst.* vi. 798.—Abyss.
Aucheri, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 38 = ocymifolia.
aurea, *Linn. Sp. Pl.* 276.—Sibir.
auriculaefolia, Benth. Cat. Pl. Pyren. 123 = lychnidifolia.
auriculaefolia, Brot. Fl. Lusit. i. 489 = ovalifolia.
auriculaefolia, DC. Fl. Fr. iii. 431 = Girardiana.
auriculaefolia, Poir. Encyc. vii. 401 = duriuscula.
auriculaefolia, *Vahl, Symb. Bot.* i. 25.—Europ.
auriculaefolia, Vahl, l. c., partim = oleaefolia.
auriculae-ursifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 330 = lychnidifolia.
auriculata, *Hort. Dorpat.* (1810–14); *ex Schult. Syst.* vi. 799.—Hab.?
australis, Moris, Diagn. Stirp. Sard. (1857) 1 = duriuscula.
australis, *Spreng. Syst.* i. 959.—China; Japon.; Austral.
Avei, De Not. Prosp. Fl. Ligust. 54 = echioides.
axillaris, *Forsk. Fl. Aegypt. Arab.* 58.—Arab.
bahamensis, *Griseb. Fl. Brit. W. Ind.* 389.—Ind. occ.
bahusiensis, Fries, Mant. i. 10 = S. Limonium.
Balansae, Boiss. Diagn. Ser. II. iv. 69 = pycnantha.
bayonnensis, Gren. ex Boiss. in DC. Prod. xii. 649 = occidentalis.
Beaumierana, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 66, nomen.—Marocc.
Behen, Drejer, Fl. Excurs. Hafn. 122 = S. Limonium.
bellidifolia, Auct. ex Boiss. in DC. Prod. xii. 648 = auriculaefolia.
bellidifolia, Bertol. Fl. Ital. iii. 524 = psiloclada.
bellidifolia, Colm. Cat. Pl. Catalun. 130 = virgata.
bellidifolia, *Gouan, Fl. Monsp.* 231.—Europ.
bellidifolia, Gren. & Godr. Fl. Fr. ii. 749 = caspia.
bellidifolia, Sibth. & Sm. Fl. Graec. iii. 90. t. 295 = ocymifolia.
bellidifolia, Urv. in Mém. Soc. Linn. Par. i. (1822) 291 = cordata.
Besseriana, Frivald. ex Boiss. in DC. Prod. xii. 633 = collina.
Besseriana, Schult. Syst. vi. 789 = tatarica.
bicolor, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 129.—China.
Billardieri, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 325.—Ins. Molucc.
binervosa, G. E. Sm. in Engl. Bot. Suppl. t. 2663 = auriculaefolia.
Boissieri, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 156. t. 9. f. 7.—As. Min.; Aegypt.
Bonduelli, *Lestib. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 81.—Alger.
Bourgaei, *Webb, ex Boiss. in DC. Prod.* xii. 638.—Ins. Canar.
Bovei, Jaub. & Spach, Illustr. i. 157. t. 86 = axillaris.
bracteata, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 330 = Acantholimon bracteatum.
bracteosa, *Vis. Pl. Aegypt.* 16, nomen.—Aegypt.
brasiliensis, *Boiss. in DC. Prod.* xii. 644.—Am. bor. et austr.
brassicaefolia, *Webb & Berth. Phyt. Canar.* iii. 181.—Ins Canar.
Braunii, *C. Bolle, Ind. Sem. Hort. Berol.* (1861) *App.* 4.—Ins. Cap. Virid.
Brunneri, *Webb, in Hook. Niger Fl.* 170.—Ins. Cap. Virid.
Bubanii, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 326 = occidentalis.
Bungeana, *Boiss. in DC. Prod.* xii. 642.—Mongol.
Bungei, *Claus, in Beitr. Pfl. Russ. Reich.* viii. 308.—Reg. Casp.
cabulica, *Boiss. in DC. Prod.* xii. 666.—Afghan.

STATICE:—
caesia, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 325.—Hispan.
caespitosa, Cav. Ic. i. 38 = Armeria juniperifolia.
caespitosa, Poir. Encyc. vii. 396, in obs. = Armeria chilensis.
caespitosa, Urv. in Mém. Soc. Linn. Par. vi. (1826) 606 = Armeria macloviana.
Calcarae, Todaro, ex Janka, in Termész. Füzetek, vi. (1882) 175 = dictyoclada.
californica, Boiss. in DC. Prod. xii. 643 = S. Limonium.
callicoma, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. iv. II. (1845) 212.—Sibir.
cancellata, *Bernh. ex Bertol. Fl. Ital.* iii. 525.—Dalmat.; Graecia.
cancellata, Bernh. ex Mutel, Fl. Fr. iii. 88 = furfuracea.
canescens, Host, Fl. Austr. i. 407 = Armeria canescens.
capillifolia, Poir. Encyc. vii. 397 = Armeria pinifolia.
capitata, Lam. Fl. Fr. iii. 63 = Armeria maritima.
carnosa, *Boiss. in DC. Prod.* xii. 663.—Persia.
caroliniana, Walt. Fl. Carol. 118 = S. Limonium.
caryophyllacea, Boiss. & Hohen. ex Boiss. Diagn. Ser. I. vii. 79 = Acantholimon caryophyllaceum.
caspia, Pollini, Fl. Veron. i. 418, 611 = virgata.
caspia, *Willd. Enum. Hort. Berol.* 336.—Reg. Cauc.; Sibir.
catalaunica, Willk. & Costa, ex Willk. in Linnaea, xxx. (1859–60) 122 = duriuscula.
Cephalotes, Ait. Hort. Kew. ed. I. i. 383 = Armeria latifolia.
Cephalotes, Bertol. Fl. Ital. iii. 511 = Armeria Morisii.
Cephalotes, Poir. Encyc. vii. 394 = Armeria mauritanica.
cespitosa, Auct. = caespitosa.
chilensis, *Phil. in Anal. Univ. Chil.* (1861) 58.—Chili.
chrysocephala, *Regel, in Act. Hort. Petrop.* vi. (1879) 383.—As. centr.
chrysocoma, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 429.—Soongar.
cinerea, Poir. Encyc. vii. 407 = purpurata.
collina, *Griseb. Spicil. Fl. Rumel.* ii. 300.—Graec.; As. Min.; Syria.
collina, Nym. Consp. 613 = dalmatica.
Companyonis, *Gren. & Billot, in F. Schultz, Fl. Fr. et Allem.* 338.—Gallia.
confusa, *Gren. & Godr. Fl. Fr.* ii. 743.—Reg. Mediterr.
congesta, Heldr. ex Nym. Consp. 610 = corinthiaca.
congesta, *Ledeb. Fl. Alt.* i. 427.—Sibir. altaic.
conspicua, Sims, Bot. Mag. t. 1629 = speciosa.
coralloides, Tausch, in Syll. Ratisb. ii. (1828) 255 = decipiens.
cordata, Desf. Fl. Atlant. i. 273 = virgata.
cordata, Guss. Fl. Sic. Prod. i. 382 = cosyrensis.
cordata, *Linn. Sp. Pl.* 275.—Reg. Mediterr.
cordata, Poir. Voy. Barb. ii. 142 = spathulata.
cordata, Savi, Due Cent. 82 = psiloclada.
cordata, [Suffr.] Princip. Bot. (1802) 132 = bellidifolia.
coriacea, Hort. ex Schult. Syst. vi. 778 = latifolia.
Coriaria, Pall. Tabl. Taur. 49; et ex Bieb. Fl. Taur. Cauc. i. 249 = latifolia.
corinthiaca, *Boiss. & Heldr. Diagn.* Ser. II. iv. 69.—Graecia.
corymbulosa, *Boiss. in DC. Prod.* xii. 658.—Afr. austr.
corymbulosa, *Nym. Consp.* 609.—Hispan.
Costae, Willk. in Linnaea, xxx. (1859–60) 12 = Girardiana.
cosyrensis, *Guss. Suppl. Fl. Sic. Prod.* 90.—Europ. austr.
crispa, Pers. Syn. i. 334 = mucronata.
Crouanii, Lenorm. ex Nym. Consp. 609 = S. Limonium.
cumana, *Tenore, Fl. Nap.* iii. 351.—Italia.
cuneata, Sm. ex Link, Handb. ii. 264; Hort. Loud. ex Boiss. in DC. Prod. xii. 636 = Thouini.
cuspidata, Delort. in F. Schultz, Arch. Fl. Fr. et Allem. 339 = duriuscula.
cylindrica, Steud. Nom. ed. I. 810, sphalm. = seq.
cylindrifolia, *Forsk. Fl. Aegypt. Arab.* 59.—Arab.; Afr. bor.
cymulifera, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 104.—Afr. bor.
cyrtostachya, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 328.—Afr. bor.

STATICE:—
dalmatica, *Presl, Bot. Bemerk.* 105.—Dalmat.; Graec.
davurica, Pall. Reise, iii. 320 = flexuosa.
decipiens, *Ledeb. Fl. Alt.* i. 433.—Sibir. altaic.
decumbens, *Boiss. in DC. Prod.* xii. 659.—Afr. austr.
delicatula, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 327.—Hispan.
densiflora, Girard, l. c. Sér. II. xvii. (1842) 25 = Girardiana.
densiflora, Guss. Suppl. Fl. Sic. Prod. 86 = oleaefolia.
dentata, *Hort. Pesth.* (1812); *ex Schult. Syst.* vi. 799 (Quid?).—Hab.?
denticulata, Bertol. Rar. Ital. Pl. Dec. ii. 34 = Armeria denticulata.
denudata, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1857) 37; *Gartenfl.* (1861) 159.—Hab.?
desertorum, Trautv. Pl. Imag. 17. t. 11 = tatarica.
dianthifolia, Jaub. & Spach, Illustr. i. 162 = Acantholimon venustum.
dichotoma, *Cav. Ic.* i. 37. t. 50.—Hispan.; Graec.
dichotoma, Duby, Bot. Gall. i. 388 = bellidifolia.
dichotoma, Guss. Fl. Sic. Syn. i. 372 = virgata.
dichotoma, Mutel, Fl. Fr. Suppl. 171 = auriculaefolia.
dichotoma, Reichb. Pl. Crit. viii. 7. t. 718 = oleaefolia.
dichotoma, Sibth. & Sm. Fl. Graec. Prod. i. 212 = cosyrensis.
dichotoma, Willk. in Flora, xxxv. (1852) 290 = duriuscula.
dichroantha, *Rupr. Sert. Tiansch.* 69.—As. centr.
Dickensoni, *Hort. ex Poit. in Rev. Hortic.* Sér. II. ii. (1843–44) 425.—Afr. austr.
Dicksoniana, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 643, nomen = rosea.
dictyoclada, *Boiss. in DC. Prod.* xii. 654.—Corsica.
dictyophora, Tausch, in Syll. Ratisb. ii. (1828) 254 = cancellata.
diffusa, Laterr. ex Boiss. in DC. Prod. xii. 661 = caspia.
diffusa, *Pourr. in Mém. Acad. Toul.* iii. (1788) 330.—Gall.; Lusitan.
divaricata, Cav. ex Schult. Syst. vi. 847 = reticulata.
Dodartii, Girard, in Ann. Sc. Nat. Sér. II. xvii. (1842) 31 = auriculaefolia.
Dregeana, *Presl, Bot. Bemerk.* 105; *Boiss. in DC. Prod.* xii. 659.—Afr. austr.
drepanensis, Tineo, ex Guss. Fl. Sic. Syn. ii. 805 = S. Limonium.
dschungarica, *Regel, in Act. Hort. Petrop.* vi. (1879) 386.—As. centr.
dubia, Andr. ex Guss. Suppl. Fl. Sic. Prod. 89 = dictyoclada.
Dubyei, *Gren. & Godr. Fl. Fr.* ii. 750.—Gallia; Hispan.
Dufourei, *Girard, in Ann. Sc. Nat.* Sér. II. xvii. (1842) 36.—Hispan.
Duriaei, *Girard, in Mém. Acad. Montp.* (*Sect. Sc.*) i. (1848) 187.—Afr. bor.
duriuscula, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 327.—Gall. austr.
echinata, *Hort. Gorenk.* (1812); *ex Schult. Syst.* vi. 799 (Quid?).—Hab.?
Echinus, Bieb. Fl. Taur. Cauc. i. 251 = Acantholimon Hohenackeri.
Echinus, Linn. Sp. Pl. 276 = Acantholimon androsaceum, caryophyllaceum, tenuiflorum.
echioides, Hort. ex Schult. Syst. vi. 785 = reticulata.
echioides, *Linn. Sp. Pl.* 275.—Reg. Mediterr.
echioides, Sibth. & Sm. Fl. Graec. Prod. i. 213 = rorida.
effusa, *Boiss. in DC. Prod.* xii. 646.—Lydia.
elata, *Fisch. Cat. Hort. Gorenk.* (1812) 18.—Sibir.
elegans, Coss. ex Nym. Consp. 613 = caesia.
elongata, Hoffm. Deutschl. Fl. ed. II. i. 150 = Armeria elongata.
emarginata, Schur, Enum. Pl. Transs. 558 = Gmelini.
emarginata, *Willd. Enum. Hort. Berol.* 335.—Gibralt.
equisetina, *Boiss. in DC. Prod.* xii. 658.—Afr. austr.
erinacea, Jaub. & Spach, Illustr. i. 163 = Acantholimon roseum.
eximia, *Schrenk, Enum. Pl. Nov.* i. 13.—As. centr.
fasciculata, Vent. Jard. Cels, 38 = Armeria fasciculata.
ferox, Jaub. & Spach, Illustr. i. 163 = Acantholimon Scorpius.

STATICE:—
ferulacea, *Linn. Sp. Pl.* ed. II. 396.—Reg. Mediterr. occ.
festucacea, Jaub. & Spach, Illustr. i. 163 = Acantholimon festucaceum.
filicaulis, *Boiss. Elench.* 80.—Hispan.
Fischeri, Trautv. in Act. Hort. Petrop. ii. (1873) 481 = nuda.
flexuosa, Less. in Linnaea, ix. (1834) 196 = elata.
flexuosa, *Linn. Sp. Pl.* 276.—Sibir.
foliosa, *Penny, ex W. Baxt. in Loud. Hort. Brit. Suppl.* ii. 675, nomen.—Ins. Canar.
formosa, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1103 = Armeria latifolia.
Fortuni, Lindl. Bot. Reg. (1845) t. 63 = australis.
Fradiniana, Pomel, Nouv. Mat. Fl. Atl. 128 = oleaefolia.
Franchetii, *Debeaux, in Act. Linn. Soc. Bord.* xxxi. (1877) 384.—China.
frutescens, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 643; Lem. in Fl. des Serres, iv. (1848) t. 325 = seq.
fruticans, *Webb, ex Boiss. in DC. Prod.* xii. 636.—Ins. Canar.
fruticosa, Lepech. Reise, i. 254 = suffruticosa.
furfuracea, *Lag. Gen. et Sp. Nov.* 13.—Hispan.
furfuracea, Reichb. Ic. Fl. Germ. xvii. 64 = cancellata.
Gaillardoti, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 158. t. 9. f. 9.—Syria.
genistoides, Jaub. & Spach, Illustr. i. 162 = Acantholimon genistoides.
Girardiana, *Guss. Fl. Sic. Syn.* i. 368.—Hispan.
glauca, Less. in Linnaea, ix. (1834) 196 = suffruticosa.
glauca, Pers. Syn. i. 333 = spathulata.
glauca, Willd. ex Schult. Syst. vi. 799 = Gmelini.
globulariaefolia, DC. Fl. Fr. Suppl. 379 = confusa.
globulariaefolia, *Desf. Fl. Atlant.* i. 274.—Afr. bor.
globulariaefolia, Webb, Iter Hisp. 18 = Girardiana.
globularifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 376 = salsuginosa.
globularifolia, Sibth. & Sm. Fl. Graec. iii. 90. t. 296 = Sibthorpiana.
globulifera, *Boiss. & Heldr. ex Boiss. in DC. Prod.* xii. 665.—As. Min.
glomerata, *Tausch, in Syll. Ratisb.* ii. (1828) 255.—Sardinia.
glumacea, Jaub. & Spach, Illustr. i. 162, 166 = Acantholimon glumaceum.
Gmelini, Koch, Syn. Fl. Germ. ed. II. 684 = S. Limonium.
Gmelini, Spreng. Syst. i. 957 = ovalifolia.
Gmelini, *Willd. Sp. Pl.* i. 1524.—Reg. Cauc.; Sibir.
Gougetiana, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 328.—Hispan.
gracilis, *Fisch. ex Boiss. in DC. Prod.* xii. 660.—Sibir. altaic.
gracilis, Tineo, in Guss. Fl. Sic. Syn. ii. 807 = dictyoclada.
graeca, *Boiss. in DC. Prod.* xii. 650.—Graecia; Syria.
graeca, Poir. Encyc. Suppl. v. 237 = rorida.
graminifolia, *Ait. Hort. Kew.* ed. I. i. 383.—Rossia.
graminifolia, Bess. Enum. Pl. Volh. 44 = tatarica.
grandiflora, Hort. Viln. (1811) ex Schult. Syst. vi. 777 = Armeria latifolia.
Griffithii, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 175.—Afghan.
gummifera, *Dur. ex Boiss. & Reut. Pugill. Pl. Nov.* (1852) 104.—Afr. bor.
Gussoniana, Steud. Nom. ed. II. ii. 633 = Sibthorpiana.
Halleri, Garcke, Fl. Deutschl. ed. II. 273 = Armeria elongata.
halochrysa, Fisch. ex Boiss. in DC. Prod. xii. 641 = Schrenkiana.
hirsuta, Presl, Bot. Bemerk. 105 = sinuata.
hirta, Steud. Nom. ed. I. 811 = Armeria baetica.
Histrix, Jaub. & Spach, Illustr. i. 162 = Acantholimon libanoticum.
Hoeltzeri, *Regel, in Act. Hort. Petrop.* v. (1877) 259.—Turkestan.
Hohenackeri, Jaub. & Spach, Illustr. i. 162, 167 = Acantholimon Hohenackeri.
horrida, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 331 = Acantholimon Hohenackeri.
Humboldtii, *C. Bolle, in Ind. Sem. Hort. Berol.* (1861) *App.* 4.—Ins. Canar.
humilis, Link, in Schrad. Journ. (1800) i. 61 = Armeria caespitosa.

STATICE:—
hybrida, Mont. ex Lloyd, Fl. Loire-Inf. 211 = ovalifolia.
hyssopifolia, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 329 = rorida.
iconia, *Boiss. & Heldr. in DC. Prod.* xii. 661.—Anatolia.
imbricata, *Webb; ex Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 330.—Ins. Canar.
inarimensis, Guss. Enum. Pl. Inar. 267 = minutiflora.
incana, Bieb. Fl. Taur. Cauc. i. 251; Bess. Enum. Pl. Volh. 94 = tatarica.
incana, Ledeb. Fl. Alt. i. 435 = callicoma.
incana, *Linn. Mant.* i. 59.—Tauria; Sibir.; Afr. bor.
incana, Urv. ex Boiss. in DC. Prod. xii. 672 = collina.
incana, Vis. Fl. Dalm. ii. 7 = dalmatica.
insignis, *Coss. Notes Pl. Crit.* 177.—Hispan.
intermedia, Guss. Suppl. Fl. Sic. Prod. 87 = psiloclada.
japonica, Sieb. & Zucc. Fl. Jap. Fam. Nat. ii. 72 = australis.
Jauberti, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 331 = Acantholimon roseum.
Jovibarba, *Webb, ex Boiss. in DC. Prod.* xii. 665.—Afr. trop.
juncea, *Tatarinow, in Wlangali, Reise Oestl. Kirgisenst.* 251, nomen.—Turkestan.
juniperifolia, Boiss. in DC. Prod. xii. 680 = Armeria splendens.
juniperifolia, Pall. ex Steud. Nom. ed. II. ii. 633 = *juniperina*.
juniperifolia, Vahl, Symb. Bot. i. 25 = Armeria caespitosa.
juniperina, Willd. ex Steud. Nom. ed. I. 811 = Acantholimon Hohenackeri.
Karelini, Stschégl. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 475 = Acantholimon Karelini.
kaschgarica, *Rupr. Sert. Tiansch.* 69.—Turkestan.
Kaufmanniana, *Regel, Gartenfl.* (1880) 1.—As. centr.
Kotschyi, Jaub. & Spach, Illustr. i. 162 = Acantholimon Kotschyi.
Kraussiana, *Buching. ex Krauss, in Flora*, xxviii. (1845) 73.—Afr. austr.
laeta, Ball, in Journ. Bot. xiii. (1875) 176 = ornata.
laeta, Moris, Fl. Sard. iii. 42 = duriuscula.
laeta, Salisb. Prod. 177 = mucronata.
lanceolata, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1218 = axillaris.
lanceolata, Hoffmgg. & Link, Fl. Port. i. 446 = ovalifolia.
lanceolata, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1099 = eximia.
lanceolata, Reichb. Pl. Crit. viii. 7. t. 719. f. 961 = occidentalis.
latifolia, *Sm. in Trans. Linn. Soc.* i. (1791) 250.—Bulgar.; Rossia; Reg. Cauc.
latissima, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 729.—Soongar.
Lefroyi, *Hemsl. in Journ. Bot.* xxi. (1883) 105.—Ins. Bermuda.
Legrandi, Gaut. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iv. (1877–78) 40 = Companyonis.
lepidorachis, Pomel, Nouv. Mat. Fl. Atl. 127 = sebkarum.
leptoloba, *Regel, in Act. Hort. Petrop.* vi. (1879) 385.—As. centr.
leptophylla, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 211.—Soongar.
leptostachya, *Boiss. Diagn.* Ser. I. vii. 68.—Persia.
leptostachys, *Pomel, Nouv. Mat. Fl. Atl.* 311.—Alger.
lepturoides, Jaub. & Spach, Illustr. i. 163, 170 = Acantholimon lepturoides.
Lespinassi, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 156. t. 8. f. 5.—Europ.
leucacantha, Jaub. & Spach, Illustr. i. 164 = Acantholimon Scorpius.
leucantha, Pouz. in Mém. Soc. Linn. Par. iv. (1826) 560; vi. (1827) 405; Loisel. Fl. Gall. ed. II. i. 223 = Armeria leucocephala.
lilacina, Boiss. & Bal. Diagn. Ser. II. iv. 68 = Gmelini.
limonia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Limonium.
Limonium, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 379 = ovalifolia.
Limonium, *Linn. Sp. Pl.* ed. I. 274.—Europ.; Am. bor.
Limonium, Pall. Tabl. Taur. 49 = Gmelini.

STATICE:—
Limonium, Thunb. Fl. Jap. 129 = australis.
limonoides, Bivona, ex Link, Enum. Hort. Berol. i. 295 = S. Limonium.
linearifolia, Laterr. Fl. Bord. ed. II. 189 = Armeria maritima.
linearifolia, Loisel. Fl. Gall. ed. I. i. 180 = Armeria elongata.
Lingua, *Pomel, Nouv. Mat. Fl. Atl.* 129.—Alger.
linifolia, Drège, ex Boiss. in DC. Prod. xii. 658 = equisetina.
linifolia, *Linn. f. Suppl.* 187.—Afr. austr.
littoralis, Poir. Encyc. Suppl. v. 235 = Armeria littoralis.
lobata, Linn. f. Suppl. 187 = Thouini.
longidentata, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 156. t. 8. f. 6.—Europ.
longifolia, Thunb. Prod. Pl. Cap. 54 = purpurata.
lusitanica, Poir. Voy. Barb. ii. 141 = Armeria mauritanica.
lusitanica, Poir. l. c. 173 = Armeria latifolia.
lychnidifolia, *Girard, in Ann. Sc. Nat.* Sér. II. xvii. (1842) 18. t. 3.—Europ. austr. occ.; Marocc.
lycopodioides, Girard, l. c. Sér. III. ii. (1844) 330 = Acantholimon lycopodioides.
lyrata, Bieb. Beschr. Casp. 166 = spicata.
macrophylla, *Brouss. ex Spreng. Syst.* i. 959.—Ins. Canar.
macroptera, *Webb & Berth. Phyt. Canar.* iii. 182.—Ins. Canar.
macrorrhabdos, *Boiss. Diagn.* Ser. II. iv. 67.—Afghan.
macrorrhiza, *Ledeb. Fl. Alt.* i. 434.—Sibir. altaic.
marginata, *Schott, ex Steud. Nom.* ed. II. ii. 633.—Bras.
maritima, Lam. Fl. Fr. iii. 64 = cordata, S. Limonium.
maritima, Mill. Gard. Dict. ed. VIII. n. 3 = Armeria maritima.
Maurocenia, Hort. Par. ex Dum.-Cours. Bot. Cult. ed. I. i. 661 (Quid?).—Hab.?
melia, Nym. Consp. 612 = cosyrensis.
membranacea, *Cæsern. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 117, nomen.—Rossia.
Meyeri, Boiss. in DC. Prod. xii. 645 = S. Limonium, Gmelini.
micrantha, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 159. t. 9. f. 10.—As. Min.
minuta, Desf. Fl. Atlant. i. 275 = Gougetiana.
minuta, *Falk, Beitr.* ii. 153.—Ross. austr.
minuta, *Linn. Mant.* i. 59.—Reg. Mediterr.
minuta, Mor. ex Reichb. Pl. Crit. iii. 23 = rupicola.
minuta, Tenore, ex Nym. Consp. 610, partim = Gougetiana.
minuta, Tenore, Syll. 593 = minutiflora.
minuta, Tommas. ex Nym. Consp. 612 = cancellata.
minuta, Vis. Fl. Dalm. iii. 356 = cordata.
minuta, Willk. Pl. Haloph. 137 = virgata.
minutiflora, *Guss. Suppl. Fl. Sic. Prod.* 89.—Sicil.
mongolica, *Fisch. ex Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 329.—Mongolia.
monopetala, Linn. Sp. Pl. 276 = Limoniastrum articulatum.
montana, Mill. Gard. Dict. ed. VIII. n. 2 = Armeria alpina.
montana, Soleir. ex Boiss. in DC. Prod. xii. 686 = Armeria leucocephala.
mucosa, Salzm. ex Boiss. l. c. 646 = ovalifolia.
mucronata, *Linn. f. Suppl.* 187.—Marocc.
multiceps, *Pomel, Nouv. Mat. Fl. Atl.* 128.—Alger.
myosuroides, *Regel, Pl. Nov. Fedsch.* 74.—Turkestan.
myriantha, *Schrenk, Enum. Pl. Nov.* 14.—Soongar.
nana, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 601 = auriculaefolia?
neglecta, Andrz. Rys. Bot. Wilno (1823), 48, nomen (Quid?).—Rossia.
nigricans, *Lafont, in Act. Linn. Soc. Bord.* xxvii. (1869) 155.—Europ.
nuda, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 184.—Persia.
obovata, *Ledeb. Fl. Ross.* iii. 468.—Reg. Cauc.
obtusa, Dum.-Cours. Bot. Cult. ed. I. i. 660 = cordata.
occidentalis, *Lloyd, Fl. Loire-Inf.* 212.—Europ. occ.; Marocc.
ochrantha, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 730.—Sibir. altaic.
ocimifolia, Schult. Syst. vi. 780 = seq.
ocymifolia, *Poir. Encyc. Suppl.* v. 238.—Graecia.
oleaefolia, Host, Syn. Pl. Austr. 177 = reticulata.
oleaefolia, Pourr. ex DC. Fl. Fr. iii. 422 = virgata.

STATICE:—
oleaefolia, *Scop. Delic. Insub.* i. 24. t. 10.—Reg. Mediterr. occ.
oleaefolia, Willd. Sp. Pl. i. 1525 = Dodartii.
oleifolia, Sibth. & Sm. Fl. Graec. Prod. i. 212 = virgata.
Oliveriana, Andr. ex Boiss. Fl. Orient. iv. 862 = rorida.
Oliverii, Jaub. & Spach, Illustr. i. 163, 168 = Acantholimon venustum.
Oliveriana, Andrz. ex Boiss. in DC. Prod. xii. 650 = graeca.
ornata, *Ball, in Journ. Linn. Soc.* xvi. (1878) 559.—Marocc.
Orphanidis, Boiss. Diagn. Ser. II. vi. 146 (Index), sphalm. = Armeria majellensis.
otolepis, *Schrenk, in Bull. Phys.-Math. Pétersb.* i. (1843) 362.—Soongar.
ovalifolia, *Poir. Encyc. Suppl.* v. 237.—Reg. Mediterr. occ.
Owerini, *Boiss. Fl. Orient.* iv. 870.—Reg. Cauc.
oxylepis, Boiss. in DC. Prod. xii. 647 = oleaefolia.
palmaris, Sibth. & Sm. Fl. Graec. iii. 91. t. 297 = rorida.
panormitana, Todaro, Ind. Sem. Hort. Panorm. (1857) 45.—Cf. Linnaea, xxix. (1857–58) 750 = minutiflora.
papillata, *Webb & Berth. Phyt. Canar.* iii. 177.—Ins. Canar.
parvifolia, Tinèo, in Guss. Fl. Sic. Syn. ii. 806 = dictyoclada.
patens, Fisch. ex Boiss. in DC. Prod. xii. 661 = bellidifolia.
pauciflora, Jaub. & Spach, Illustr. i. 162 = Acantholimon androsaceum.
pectinata, *Ait. Hort. Kew.* ed. I. i. 385.—Ins. Canar.
pedicellata, *Wallr. ex Boiss. in DC. Prod.* xii. 658.—Afr. austr.
peregrina, Berg. Desc. Pl. Cap. 80 (excl. syn.) = rosea.
perfoliata, *Karel. in Bull. Soc. Nat. Mosc.* (1839) 167, nomen; *et ex Boiss. in DC. Prod.* xii. 663.—Persia.
pinifolia, Brot. Fl. Lusit. i. 486 = Armeria pinifolia.
plantaginea, All. Fl. Pedem. ii. 90 = Armeria plantaginea.
plantaginea, Guss. Fl. Sic. Syn. i. 366 = Armeria Gussonei.
plantaginiflora, Jaub. & Spach, Illustr. i. 159. t. 88 = spicata.
plumosa, *Phil. Fl. Atac.* 46.—Chili.
polianthemum, *Neck. Delic. Gallo-Belg.* i. 160.—Europ.
Preauxii, *Webb & Berth. Phyt. Canar.* iii. 181.—Ins. Canar.
preciosa, *Tchertkoff, in Bull. Soc. Nat. Mosc.* vi. (1833) 15.—Rossia.
prolifera, Urv. in Mém. Soc. Linn. Par. i. (1822) 291 = rorida.
pruinosa, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 372 = caesia.
pruinosa, Forsk. ex Schult. Syst. vi. 794 = furfuracea.
pruinosa, *Linn. Mant.* i. 59.—Aegypt.
pruinosa, Webb & Berth. ex Boiss. in DC. Prod. xii. 640 = papillata.
Pseudo-armeria, Cav. Ic. i. 38 = Armeria gaditana.
Pseudo-armeria, Paxt. Mag. Bot. xi. (1844) t. 79 = Armeria latifolia.
Pseudo-limonium, Reichb. Fl. Germ. Excurs. 191 = S. Limonium.
psiloclada, *Boiss. in DC. Prod.* xii. 651.—Reg. Mediterr.
puberula, *Webb, in Bot. Reg.* t. 1450.—Ins. Canar.
pubescens, DC. Fl. Fr. Suppl. 380 = cordata.
pubescens, Koch, Syn. Fl. Germ. ed. I. 595 = cancellata.
pubescens, Reichb. Ic. Fl. Germ. ii. 79 = minuta, *Linn.*
pubescens, Salis, in Flora, xvii. (1834) II. Beibl. 13 = Armeria leucocephala.
pubescens, Sm. ex Schult. Syst. vi. 772 (cum cit. falsa) = Armeria maritima.
pubigera, Desf. Cat. Hort. Par. ed. III. 391 = Armeria maritima.
pulchella, S. F. Gray, Nat. Arr. Brit. Pl. ii. 296 = Armeria elongata.
pungens, Brot. Fl. Lusit. 487 = Armeria pungens.
pungens, Jaub. & Spach, Illustr. i. 163 = Acantholimon roseum.
purpurata, *Linn. Mant.* i. 59.—Afr. austr.

STATICE:—
purpurata, Willd. Sp. Pl. i. 1528 = rosea.
purpurea, Koch, Syn. Fl. Germ. ed. I. 595 = Armeria elongata.
pycnantha, *C. Koch, in Linnaea*, xxi. (1848) 716.—As. Min.
pygmaea, Tineo, in Guss. Fl. Sic. Syn. ii. 807 = dictyoclada.
pyramidata, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 18.—Ins. Porto-santo.
pyrrholepis, Pomel, Nouv. Mat. Fl. Atl. 131 = delicatula.
Raddiana, Boiss. in DC. Prod. xii. 653 = globulariaefolia.
ramosissima, Poir. Voy. Barb. ii. 142 = globulariaefolia.
rariflora, Drejer, Fl. Excurs. Hafn. 121 = S. Limonium.
remotispicula, *Lacaita, in Nuov. Giorn. Bot. Ital.* xvi. (1884) 168.—Italia.
reniformis, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 325 = perfoliata.
reticulata, Auct. ex Boiss. in DC. Prod. xii. 654 = dictyoclada.
reticulata, Bieb. Fl. Taur. Cauc. i. 251 = caspia.
reticulata, Gouan, Suppl. Fl. Monsp. 139, nomen = virgata.
reticulata, *Linn. Sp. Pl.* 275.—Europ.
reticulata, Siev. ex Boiss. in DC. Prod. xii. 661 = decipiens.
rorida, *Sibth. & Sm. Fl. Graec.* iii. 91. t. 298.—Graecia; Syria.
rorida, Vis. Stirp. Dalmat. 48; et ex Boiss. Fl. Orient. iv. 863 = cancellata.
rosea, Boiss. in DC. Prod. xii. 624 = Acantholimon roseum.
rosea, Pall. Reise, iii. 260 = flexuosa.
rosea, *Sm. in Rees, Cycl.* xxxiv. [n. 20].—Afr. austr.
rubella, S. T. Gmel. Reise, ii. 199. t. 34 = tatarica.
rupicola, *Badaro, ex Reichb. Fl. Germ. Excurs.* 191.—Italia.
rytidophylla, Hook. Bot. Mag. t. 4055 = rosea.
sabulosa, *Fourr. in Ann. Soc. Linn. Lyon*, N. S. xvii. (1869) 141.—Europ.
salicorniacea, *F. Muell. Fragm.* xi. 7.—Austral.
salsuginosa, *Boiss. in DC. Prod.* xii. 653.—Hispan.
sareptana, Becker, in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 12 = tomentella.
Sartorii, *Nym. Syll. Suppl.* 25.—Ins. Mycon.
scabra, Drège, ex Boiss. in DC. Prod. xii. 666 = purpurata.
scabra, *Krauss, in Flora*, xxviii. (1845) 72 = corymbulosa, *Boiss.*
scabra, Pall. ex Boiss. in DC. Prod. xii. 681 = Armeria scabra.
scabra, Pers. Syn. i. 334 = Limoniastrum articulatum.
scabra, *Thunb. Prod. Pl. Cap.* 54.—Afr. austr.
scabrida, Presl, Bot. Bemerk. 105 = equisetina.
Schrenkiana, *Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 362.—Soongar.
scoparia, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1098 = eximia.
scoparia, C. A. Mey. ex Boiss. in DC. Prod. xii. 645 = S. Limonium, Gmelini.
scoparia, Pall. ex Willd. Sp. Pl. i. 1524 = Gmelini.
scoparia, Reichb. ex Nym. Consp. 609 = S. Limonium.
Scopoliana, Bertol. Fl. Ital. iii. 528 = oleaefolia.
Scorpius, Jaub. & Spach, Illustr. i. 163 = Acantholimon Scorpius.
scorzoneraefolia, Link, in Schrad. Journ. (1800) i. 60 = Armeria plantaginea.
sebkarum, *Pomel, Nouv. Mat. Fl. Atl.* 130.—Alger.
sedoides, *Regel, in Act. Hort. Petrop.* vi. (1879) 384.—As. centr.
Semenovii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) i. 398.—Sibir.
serbica, *Nym. Syll. Suppl.* 25.—Serbia.
serotina, Reichb. Fl. Germ. Excurs. 191 = S. Limonium.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 386.—Turkestan.
sibirica, Dum.-Cours. Bot. Cult. ed. I. i. 661 = echioides.
sibirica, *Ledeb. Fl. Ross.* iii. 457.—Sibir. baical.
Sibthorpiana, *Guss. Suppl. Fl. Sic. Prod.* 87.—Sicil.
Sieberi, *Boiss. Voy. Bot. Espagne*, ii. 530, in nota.—Graecia.
sinensis, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 329 = australis.
sinuata, *Linn. Sp. Pl.* 276.—Reg. Mediterr.

STATICE:—
sisymbrifolia, Jaub. & Spach, Illustr. i. 158 = spicata.
Smithii, Tenore, Fl. Nap. iii. 350. t. 223 = virgata.
Spachii, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 322 = Acantholimon tragacanthinum, A. Scorpius.
spathulata, *Desf. Fl. Atlant.* i. 275.—Afr. bor.; Creta.
spathulata, Heldr. ex Nym. Consp. 611 = rorida.
spathulata, Hook. Brit. Fl. ed. I. 145 = auriculaefolia.
spathulata, Jan, ex Nym. Consp. 610 = oleaefolia.
spathulata, Nym. Syll. 143 = emarginata.
spathulata, Sieber, ex Boiss. in DC. Prod. xii. 650 = Sieberi.
spathulata, Willd. ex Boiss. l. c. 652 = duriuscula.
spathulata, Willd. Enum. Hort. Berol. 335 = Willdenovii.
speciosa, Forsk. Fl. Aegypt. Arab. p. xxiv = collina.
speciosa, Hort. ex Schult. Syst. vi. 796 = mucronata.
speciosa, *Linn. Sp. Pl.* 275.—Reg. Cauc.; Oriens; Sibir.
spicata, *Willd. Sp. Pl.* i. 1533.—Sibir. occ.; Reg. Cauc.; Oriens.
spinulosa, *Janka, in Termész. Füzetek*, vi. (1882) 170.—Europ.
splendens, Lag. & Rodr. in Anal. Cienc. Nat. v. (1802) 274 = Armeria splendens.
squamata, Poir. Encyc. Suppl. v. 237 = tubiflora.
Stocksii, *Boiss. in DC. Prod.* xii. 664.—Sind; Beluchist.
stricta, *Gueldenst. Reisen*, ii. 225; *ex Ledeb. Fl. Ross.* iii. 470.—Ross. austr.
suffruticosa, *Linn. Sp. Pl.* 276.—Reg. Cauc.; As. med.
supina, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 326.—Hispan.
Suworowi, *Regel, in Act. Hort. Petrop.* vii. (1880) 550.—As. centr.
tatarica, *Linn. Sp. Pl.* 275.—Europ. austr. or.; Reg. Cauc.
Taxanthema, Schult. Syst. vi. 798 = australis.
tenella, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 203.—Mongol.
Tenoreana, Guss. Enum. Pl. Inar. 268, in syn. = minutiflora.
tenuicula, Tineo, ex Guss. Suppl. Fl. Sic. Prod. 90 = reticulata.
tenuifolia, Bertol. in Moris, Stirp. Sard. Elench. ii. 8 = rupicola.
tenuifolia, Jaub. & Spach, Illustr. i. 162 = Acantholimon Hohenackeri.
terekiensis, Gueldenst. Reisen, ii. 33 = tatarica.
tetragona, Drège, ex Boiss. in DC. Prod. xii. 659 = Dregeana.
tetragona, *Thunb. Prod. Pl. Cap.* 54.—Afr. austr.
thessala, Boiss. & Heldr. Diagn. Ser. II. vi. 146 (Index), sphalm. = Armeria cariensis.
Thouini, *Viv. Fl. Lib. Spec.* 18. t. 11. f. 1.—Reg. Mediterr.
thymoides, *Girard, in Mém. Acad. Montp.* (*Sect. Sc.*) i. (1848) 189.—Alger.
Tineii, Woods, Tourist's Fl. 306 = dictyoclada.
tomentella, *Boiss. in DC. Prod.* xii. 645.—Ross. austr.
Tournefortii, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 326 = delicatula.
Tournefortii, Jaub. & Spach, Illustr. i. 165. t. 90 = Acantholimon androsaceum.
Tournefortii, Jaub. & Spach, l. c. iv. t. 388 = rorida.
tragacanthina, Jaub. & Spach, l. c. i. 164 = Acantholimon tragacanthinum.
trigona, Pall. Tabl. Taur. 49; et ex Bieb. Fl. Taur. Cauc. i. 251 = tatarica.
trigonoides, Hort. Par. ex Boiss. in DC. Prod. xii. 633 = collina.
trigonoides, Pall. Reise, iii. 637 = incana.
trinervia, Hort. Monsp. (1804); ex Schult. Syst. vi. 799 (Quid?).—Hab.?
tripteris, Hort. Par. ex Poir. Encyc. Suppl. v. 237 = Thouini.
triquetra, Boeb. in Pall. N. Nord. Beitr. vi. 261 = tatarica.
tuberculata, *Boiss. in DC. Prod.* xii. 662.—Afr. trop.
tubiflora, *Delile, Fl. Egypte*, 215. t. 25. f. 2.—Aegypt.
tubiflora, Sieber, ex Schult. Syst. vi. 798 = pruinosa.
ulicina, Willd. ex Schult. l. c. = Acantholimon libanoticum.
undulata, Bory & Chaub. Nouv. Fl. Pélop. 20. t. 11 = Armeria argyrocephala.
varia, *Hance, in Journ. Bot.* xx. (1882) 290.—China.

STATICE:—
variabilis, Salisb. Prod. 176 = Armeria latifolia.
venusta, Fenzl, ex Boiss. Fl. Orient. iv. 832 = Acantholimon venustum.
verrucosa, Willd. ex Boiss. in DC. Prod. xii. 650 = graeca.
viminea, Schrad. ex Hornem. Suppl. Hort. Hafn. 37 = virgata.
virgata, Herb. ex Nym. Consp. 611 = confusa.
virgata, Orph. ex Nym. Consp. 610 = Sieberi.
virgata, *Willd. Enum. Hort. Berol.* 336.—Reg. Mediterr.
vulgaris, Sweet, Hort. Brit. ed. I. 332 = Armeria maritima.
Willdenovii, *Poir. Encyc. Suppl.* v. 236.—Gall. austr.
Willdenowiana, Herb. ex Nym. Consp. 610 = oleaefolia.
Willdenowiana, Reichb. Pl. Crit. ii. 55. f. 293 = duriuscula.
Willdenowii, Loisel. Fl. Gall. II. i. 224 = Girardiana.
Wrightii, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 236.—Ins. Luchu.

STAUNTONIA, DC. Syst. i. 513 (1818). *BERBERIDEAE*, Benth. & Hook. f. i. 42.
angustifolia, Wall. Cat. n. 4951 = Holboellia latifolia.
Brunoniana, Wall. l. c. n. 4592 = Parvatia Brunoniana.
chinensis, *DC. Syst.* i. 514.—China.
filamentosa, Griff. Notul. iv. 334; Ic. Pl. Asiat. t. 495. f. 1 = Holboellia latifolia.
hexaphylla, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 105; *et in Arch. Mus. Par.* i. (1839) 192.—Japon.
latifolia, Wall. Cat. n. 4950 = Holboellia latifolia.
trifoliata, Griff. Notul. iv. 330; Ic. Pl. Asiat. t. 495. f. 2, 3 = Parvatia Brunoniana.

STAURACANTHUS, Link, in Schrad. Neues Journ. ii. ii. (1807) 52 = **Ulex**, Linn. (Legumin.).
aphyllus, Link, l. c. = U. genistoides.
spartioides, Webb, Otia Hisp. 27. t. 19 = U. spartioides.
spectabilis, Webb, l. c. t. 20 = U. spectabilis.

STAURANTHERA, Benth. Scroph. Ind. 57 (1835). *GESNERACEAE*, Benth. & Hook. f. ii. 1016.
ANOMORHEGMIA, Meissn. Gen. 303 (1840).
CYANANTHUS, Griff. Notul. iv. 154 (1854).
MIQUELIA, Blume, in Bull. Néerl. (1838) 94.
QUINTILIA, Endl. Gen. 1408 (1841).
Brandisii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 372.—Burma.
ecalcarata, *Benn. Pl. Jav. Rar.* 121.—Java.
grandifolia, *Benth. Scroph. Ind.* 57.—Malaya.
umbrosa, *C. B. Clarke, Comm. & Cyrt. Beng.* t. 89.—Assam.

STAURANTHUS, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 91. *RUTACEAE*, Benth. & Hook. f. i. 303.
perforatus, *Liebm. l. c.* 92.—Mexic.

STAUREGTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 170, sphalm. = STAUROGETON, Reichb. = **Lemna**, Linn.

STAURITIS, Reichb. f. in Hamb. Gartenz. xviii. (1862) 34 = **Phalaenopsis**, Blume (Orchid.).
violacea, Reichb. f. l. c. = P. violacea.

STAUROGETON, Reichb. Nom. 33 (1841) = **Lemna**, Linn.
trisulcus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 70 = L. trisulca.

STAUROGLOTTIS, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 432 = **Phalaenopsis**, Blume (Orchid.).
equestris, Schau. l. c. = P. rosea.

STAUROGYNE, Wall. Pl. As. Rar. ii. 80. t. 186 (1831) = **Ebermaiera**, Nees (Acanth.).
angustifolia, Wall. Cat. n. 4906 = E. angustifolia.
argentea, Wall. Pl. As. Rar. ii. 80. t. 186 = E. Staurogyne.

STAUROMATUM, Endl. Ench. 128 (1841), sphalm.= **Sauromatum**, Schott (Aroid.).

STAUROPHRAGMA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ix. 90 (1843). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 929.
natolicum, *Fisch. & Mey. l. c.*—As. Min.

STAUROPSIS, Reichb. f. in Hamb. Gartenz. xvi. (1860) 117. *ORCHIDEAE*, Benth. & Hook. f. iii. 572.
FIELDIA, Gaudich. in Freyc. Voy. Bot. 424. t. 36 (1826).
fasciata, *Benth. & Hook. f. Gen.* iii. 572.—Ind. or.
gigantea, *Benth. ex Pfitzer, Vergl. Morph. Orch.* 14.—Burma.
lissochiloides, *Pfitzer, l. c.*—Ins. Philipp.
pallens, Reichb. f. in Hamb. Gartenz. xvi. (1860) 117 = Phalaenopsis pallens.
philippinensis, *Reichb. f. l. c.*—Ins. Philipp.

STAUROSPERMUM, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 73 (1827) = **Mitracarpum**, Zucc. (Rubiac.).
verticillatum, Schum. & Thonn. l. c. = M. scabrum.

STAUROSTIGMA, Scheidw. in Otto & Dietr. Allg. Gartenz. xvi. (1848) 129. *AROIDEAE*, Benth. & Hook. f. iii. 987.
ASTEROSTIGMA, Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb iii. (1845) 148; Schott, in Oestr. Bot. Wochenbl. ii. (1852) 67, nomen prius.
RHOPALOSTIGMA, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 39.
concinnum, *C. Koch, in Ender. Ind. Aroid.* 77.—Bras.
Luschnathianum, *C. Koch, l. c.* 78.—Bras.
odorum, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 129.—Bras.
Pavoni, *C. Koch, in Ender. Ind. Aroid.* 78.—Peruv.
Riedelianum, *Engl. in Mart. Fl. Bras.* iii. II. 206.—Bras.
Tweedianum, *C. Koch, in Ender. Ind. Aroid.* 78.—Bras.
Vellozianum, C. Koch, l. c. = Luschnathianum.
vermitoxicum, *Engl. in DC. Monog. Phan.* ii. 517.—Rep. Argent.

STAUROTHYRAX, Griff. Notul. iv. 476 (1854); Ic. Pl. Asiat. t. 584. f. 6 = **Phyllanthus**, Linn. (Euphorb.).

STAVIA, Thunb. Nov. Gen. vii. 108 (1792), sphalm.= **Staavia**, Dahl (Bruniac.).

STAWELLIA, F. Muell. Fragm. vii. 85 (1870). *LILIACEAE*, Benth. & Hook. f. iii. 796.
dimorphantha, *F. Muell. l. c.*—Austral.

STECHMANNIA, DC. Prod. vi. 543 (1837) = **Jurinea**, Cass. (Compos.).
eriobasis, Jaub. & Spach, Illustr. ii. t. 179 = J. eriobasis.
heterophylla, Jaub. & Spach, l. c. 99. t. 181 = J. heterophylla.
leptoloba, Jaub. & Spach, l. c. 100. t. 182 = J. leptoloba.
ramosissima, Jaub. & Spach, l. c. i. 128. t. 66 = J. ramosissima.
Staehelinae, DC. Prod. vi. 543 = J. Staehelinae.

STECHYS, Boiss. Fl. Orient. iv. 694 (1879), err. typ.= **Stachys**, Linn. (Labiat.).

STEEGIA, Steud. Nom. ed. II. ii. 634 (1841), sphalm.= STEGIA, DC. = **Lavatera**, Linn. (Malvac.).

STEENHAMERA, Kostel. Allg. Med.-Pharm. Fl. iii. 838 (1834) = seq.

STEENHAMMERA, Reichb. Fl. Germ. Excurs. 337 (1831) = **Mertensia**, Roth (Borag.).
dahurica, Turcz. in Bull. Soc. Nat. Mosc. (1840) 250 = M. dahurica.
kamczatica, Turcz. l. c. = M. paniculata.
maritima, Reichb. Fl. Germ. Excurs. 337 = M. maritima.

STEENHAMMERA:—
Pallasii, Turcz. in Bull. Soc. Nat. Mosc. (1840) 247 = Mertensia sibirica.
pilosa, Turcz. l. c. 249 = M. paniculata.
pterocarpa, Turcz. l. c. 245 = M. sibirica.
rivularis, Turcz. l. c. 248 = M. rivularis.
serrulata, Turcz. l. c. 246 = M. serrulata.
sibirica, Turcz. l. c. 247 = M. sibirica.
stylosa, Turcz. l. c. 248 = M. stylosa.
virginica, Kostel. Allg. Med.-Pharm. Fl. iii. 838 = M. pulmonarioides.

STEERBECKIA, J. F. Gmel. Syst. 820 (1791) = STERBECKIA, Schreb. = **Stingana**, Aubl. (Inc. Sed.).

STEETZIA, Sond. in Linnaea, xxv. (1852) 450 = **Olearia**, Moench (Compos.).
grandiflora, Sond. l. c. 452 = O. grandiflora.
Muelleri, Sond. l. c. 453 = O. pannosa.
ovata, Sond. l. c. 452 = O. pannosa.
pannosa, Sond. l. c. 451 = O. pannosa.

STEFFENSIA, Kunth, in Linnaea, xiii. (1839) 609 = **Piper**, Linn.
acutifolia, Kunth, l. c. 650 = P. acutifolium.
aduncа, Kunth, l. c. 633 = P. aduncum.
alnoides, Kunth, l. c. 643 = P. alnoides.
ampla, Kunth, l. c. 618 = P. amplum.
angustifolia, Kunth, l. c. 622 = P. angustifolium.
anonaefolia, Kunth, l. c. 619 = P. anonaefolium.
asperifolia, Kunth, l. c. 644 = P. asperifolium.
asperiuscula, Kunth, l. c. 631 = P. hirsutum.
caracasana, Kunth, l. c. 664 = P. caracasanum.
Carpunya, Kunth, l. c. 690 = P. Chaba.
celtidifolia, Kunth, l. c. 635 = P. celtidifolium.
chimonanthifolia, Kunth, l. c. 628 = P. chimonanthifolium.
citrifolia, Kunth, l. c. 629 = P. citrifolium.
coccoloboides, Kunth, l. c. 666 = P. coccoloboides.
consanguinea, Kunth, l. c. 623 = P. consanguineum.
crassinervia, Kunth, l. c. 659 = P. crassinervium.
crocata, Kunth, l. c. 652 = P. crocatum.
diospyrifolia, Kunth, l. c. 627 = P. diospyrifolium.
elongata, Kunth, l. c. 637 = P. elongatum.
eucalyptifolia, Kunth, l. c. 620 = P. eucalyptifolium.
evonymifolia, Kunth, l. c. 624 = P. lanceolatum.
eximia, Kunth, l. c. 665 = P. obliquum.
flexuosa, Kunth, l. c. 644 = P. flexuosum.
fuliginea, Kunth, l. c. 655 = P. fuligineum.
Gaudichaudiana, Kunth, l. c. 638 = P. Gaudichaudianum.
geniculata, Kunth, l. c. 612 = P. geniculatum.
glabrata, Kunth, l. c. 632 = P. glabratum.
hirsuta, Kunth, l. c. 641 = P. hirsutum.
hydrangeifolia, Kunth, l. c. 659 = P. crassinervium.
insignis, Kunth, l. c. 667 = P. insigne.
Jacquemontiana, Kunth, l. c. 631 = P. citrifolium.
lacunosa, Kunth, l. c. 651 = P. lacunosum.
Lhotzkyana, Kunth, l. c. 658 = P. Lhotzkyanum.
Luschnathiana, Kunth, l. c. 616 = P. Luschnathianum.
mollis, Kunth, l. c. 648 = P. mollicomum.
nitida, Kunth, l. c. 611 = P. nitidum.
nubigena, Kunth, l. c. 634 = P. crocatum.
obliqua, Kunth, l. c. 663 = P. obliquum.
oblonga, Kunth, l. c. 625 = P. Chamissonis.
Olfersiana, Kunth, l. c. 645 = P. hirsutum.
Opizii, Kunth, l. c. 641 = P. Opizianum.
plantaginea, Kunth, l. c. 672 = P. plantagineum.
Poeppigii, Kunth, l. c. 646 = P. Poeppigii.
pothifolia, Kunth, l. c. 671 = P. pothifolium.
Pseudo-churumayu, Kunth, l. c. 662 = P. Pseudo-churumayu.
Radula, Kunth, l. c. 650 = P. Bredemeyeri.
rhododendrifolia, Kunth, l. c. 621 = P. rhododendrifolium.
richardiaefolia, Kunth, l. c. 668 = P. richardiaefolium.
salicariaefolia, Kunth, l. c. = P. salicariaefolium.
scabra, Kunth, l. c. 639 = P. hirsutum.
Schlechtendalii, Kunth, l. c. 673 = P. decumanum.
scopulorum, Kunth, l. c. 669 = P. scopulorum.
secunda, Kunth, l. c. 626 = P. secundum.
Sellowiana, Kunth, l. c. 657 = P. Sellowianum.
tectoniaefolia, Kunth, l. c. 660 = P. tectoniaefolium.
tiliaefolia, Kunth, l. c. 674 = P. Schiedeanum.
tuberculata, Kunth, l. c. 614 = P. tuberculatum.
tumida, Kunth, l. c. 654 = P. tumidum.
verrucosa, Kunth, l. c. 613 = P. verrucosum.
xylopioides, Kunth, l. c. 610 = P. xylopioides.

STEGANOTAENIA, Hochst. in Flora, xxvii. (1844) Beil. 4 = **Peucedanum**, Tourn. (Umbell.).
araliacea, Hochst. l. c. = P. araliaceum.

STEGANOTROPIS, Lehm. Ind. Sem. Hort. Hamb. (1826); et in Flora, x. (1827) 96 = **Centrosema**, Benth. (Legumin.).
conjugata, Lehm. ll. cc. = C. conjugatum.

STEGANOTUS, Cass. in Dict. Sc. Nat. xxxv. 396 (1825) = **Arctotis**, Linn. (Compos.).
undulatus, Cass. l. c. = A. laciniata.

STEGIA, DC. Fl. Fr. iv. 835 (1805) = **Lavatera**, Linn. (Malvac.).
Lavatera, DC. l. c. = L. trimestris.

STEGITRIS, Rafin. Sylva Tellur. 132 (1838) = **Helianthemum**, Tourn. (Cistin.).
algarviensis, Rafin. l. c. = H. ocymoides.
atriplicifolius, Rafin. l. c. = H. atriplicifolium.
calicinus, Rafin. l. c. = H. calycinum.
lasianthus, Rafin. l. c. = H. alyssoides.

STEGNOCARPUS, Torr. in Pacif. Rail. Rep. ii. 169 (1855) = **Coldenia**, Linn. (Borag.).
canescens, Torr. l. c. 169. t. 7 = C. canescens.

STEGNOSPERMA, Benth. Bot. Voy. Sulph. 17. t. 12 (1844). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 87.
CHLAMYDOSPERMA, A. Rich. Ess. Fl. Cub. 631 (1845).
cubense, *A. Rich. Ess. Fl. Cub.* 632.—Cuba.
halimifolia, *Benth. Bot. Voy. Sulph.* 17. t. 12.—Mexic.

STEGOLEPIS, Klotzsch, ex Koern. in Linnaea, xxxvii. (1871–73) 480. *RAPATEACEAE*, Benth. & Hook. f. iii. 859.
MONOTREMA, Koern. in Linnaea, xxxvii. (1871–73) 475.
aemulans, *Benth. & Hook. f. Gen.* iii. 859.—Bras.
ferruginea, *Baker, in Journ. Bot.* xx. (1882) 331.—Guian.
guianensis, *Klotzsch, ex Koern. in Linnaea*, xxxvii. (1871–73) 481.—Guian.

STEGOSIA, Lour. Fl. Cochinch. 51 (1790) = **Rottboellia**, Linn. f. (Gramin.).
cochinchinensis, Lour. l. c. = R. exaltata.

STEIGERIA, Muell. Arg. in Linnaea, xxxiv. (1865–66) 215 = **Baloghia**, Endl. (Euphorb.).
montana, Muell. Arg. l. c.—N. Caled.

STEINCHISMA, Rafin. in Ser. Bull. Bot. i. (1830) 220 = **Panicum**, Linn. (Gramin.).
divaricatum, Rafin. l. c. = P. debile?
hians, Rafin. l. c. = P. debile.

STEINERIA, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 64. t. 5 C = **Begonia**, Tourn.
ferruginea, Klotzsch, l. c. 65 = B. Hookeriana.
pulchella, Klotzsch, l. c. = B. pulchella.

STEINHEILIA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 339. t. 12. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
radians, *Decne. l. c.*—Arab.

STEINMANNIA, Opiz, Seznam, 93 (1852) = **Rumex**, Linn. (Polygonac.).
aurea, Opiz, l. c. = R. maritimus.
flavo-virens, Opiz, l. c. = R. palustris.

STEINMANNIA, Phil. f. in Anal. Univ. Chil. (1884) [reimpr.] 10. *LILIACEAE*.
graminifolia, *Phil. f. l. c.*—Chili.

STEINREITERA, Opiz, Seznam, 93 (1852) = **Thesium**, Linn. (Santalac.).
secunda, Opiz, l. c. = T. alpinum.

STEINSCHISMA, Steud. Nom. ed. II. ii. 635 (1841) = STEINCHISMA, Rafin. = **Panicum**, Linn. (Gramin.).

STEIRACTIS, DC. Prod. v. 345 (1836) = **Olearia**, Moench (Compos.).
arborescens, DC. l. c. = O. nitida.

STEIRACTIS, Rafin. Fl. Tellur. iii. 5 (1836) = **Layia**? Hook. & Arn. (Compos.).

STEIRANISIA, Rafin. Fl. Tellur. ii. 68 (1836) = **Saxifraga**, Tourn.
heterantha, Rafin. l. c. 69 = Saxif. Mertensiana.
punctata, Rafin l. c. = Saxif. umbrosa.
reflexa, Rafin. l. c. = Saxif. reflexa.

STEIREMA, Benth. & Hook. f. Gen. iii. 363 (1880), sphalm. = seq.

STEIREMIS, Rafin. Fl. Tellur. iii. 40 (1836) = **Telanthera**, R. Br. (Amarant.).
ciliata, Rafin. New Fl. Am. iv. 45 = Digera arvensis.
ficoidea, Rafin. Fl. Tellur. iii. 41 = T. ficoidea.
repens, Rafin. l. c. = T. polygonoides.
sessilifolia, Rafin. l. c. = Alternanthera sessilis?

STEIREYA, Rafin. Fl. Tellur. iv. 100 (1836) = **Trichopus**, Gaertn. (Dioscor.).
angustifolia, cordata, media, Rafin. l. c. = T. zeylanicus.

STEIROCOMA, Reichb. Nom. 88 (1841) = **Dicoma**, Cass. (Compos.).

STEIROCTIS, Rafin. Fl. Tellur. iv. 105 (1836) (Thymel.).
capitata, Rafin. l. c. = Lachnaea capitata.
uniflora, Rafin. l. c. = Cryptadenia uniflora.

STEIRODISCUS, Less. Syn. Comp. 251 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 264.
capillaceus, *Less. l. c.*—Afr. austr.
linearilobus, *DC. Prod.* vi. 74.—Afr. austr.

STEIROGLOSSA, DC. Prod. vi. 38 (1837) = **Brachycome**, Cass. (Compos.).
chamomillaefolia, DC. l. c. 39 = B. iberidifolia.
humilis, DC. l. c. = B. diversifolia.
linearifoba, DC. l. c. = B. ptychocarpa.
rigidula, DC. l. c. = B. ciliaris.

STEIRONEMA, Rafin. in Ann. Gén. Sc. Phys. vii. (1820) 193. *PRIMULACEAE*.
ciliatum, *Rafin. l. c.*—Europ.; Am. bor.
floridum, Baudo, in Ann. Sc. Nat. Sér. II. xx. (1843) 346 = lanceolatum.
heterophyllum, *Rafin. in Ann. Gén. Sc. Phys.* vii. (1820) 193.—Am. bor.
hybridum, Rafin. l. c. = heterophyllum.
lanceolatum, A. Gray, in Proc. Am. Acad. xii. (1876) 63 = heterophyllum.
longifolium, *A. Gray, l. c.*—Am. bor.
longifolium, *Rafin. in Ann. Gén. Sc. Phys.* vii. (1820) 193.—Am. bor.
radicans, *A. Gray, in Proc. Am. Acad.* xii. (1876) 63.—Am. bor.
revolutum, Rafin. in Ann. Gén. Sc. Phys. vii. (1820) 193 = longifolium, *Rafin.*

STEKHOVIA, De Vriese, in Nat. Verh. Haarl. x. (1854) 166 = **Goodenia**, Sm.
scapigera, De Vriese, l. c. 167. t. 32. f. 1–12 = G. scapigera.
viscida, De Vriese, l. c. 168 = G. viscida.

STELANTHES, Stokes, Bot. Mat. Med. ii. 339 (1812) = **Marlea**, Roxb. (1814) (Cornac.).
solitarius, Stokes, l. c. = M. begonifolia.

STELECHOCARPUS, Hook. f. & Thoms. Fl. Ind. i. 94 (1855). *ANONACEAE*, Benth. & Hook. f. i. 22.
Burakol, *Hook. f. & Thoms. l. c.*—Malaya.
montana, *Miq. Fl. Ind. Bat.* i. II. 22.—Java.

STELECHOSPERMUM, Blume, Fl. Jav. Dipter., in Adnot. (1829). Genus dubium *GUTTIFERARUM*. —Cf. Benth. & Hook. f. i. 170.
flexuosum, *Blume, l. c.*—Java.

STELEPHUROS, Adans. Fam. ii. 31 (1763) = **Phleum**, Linn. (Gramin.).

STELESTYLIS, Drude, in Mart. Fl. Bras. iii. II. 230. t. 53. *CYCLANTHACEAE*, Benth. & Hook. f. iii. 952.
coriacea, *Drude, l. c.* 231.—Bras.

STELIS, Sw. in Schrad. Journ. ii. (1799) 239. *ORCHIDEAE*, Benth. & Hook. f. iii. 490.
DIALISSA, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1844) 107.
abrupta, *Reichb. f. in Linnaea*, xli. (1877) 7.—N. Granat.
acuifera, *Lindl. Fol. Orch. Stelis*, 12.—N. Granat.
acutiflora, *Willd. Sp. Pl.* iv. 139.—Peruv.
acutissima, *Lindl. Fol. Orch. Stelis*, 14.—Peruv.
alata, *Lindl. l. c.* 18.—Venezuela.
alba, *H. B. & K. Nov. Gen. et Sp.* i. 363.—Peruv.
alismaefolia, *Lindl. Fol. Orch. Stelis*, 16.—Peruv.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* i. 362.—N. Granat.
apiculata, *Lindl. Fol. Orch. Stelis*, 11.—Peruv.
aprica, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Bras.
argentata, *Lindl. Bot. Reg.* (1842) *Misc.* 64.—Guian.
ascendens, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Peruv.
aspera, *Pers. Syn.* ii. 524.—Peruv.
atra, *Lindl. Orch. Linden.* 3.—Venezuela.
atropurpurea, Hook. Bot. Mag. t. 3975 = ciliaris.
atroviolacea, *Reichb. f. in Bonplandia*, iii. (1855) 70.—Venezuela.
attenuata, *Lindl. Fol. Orch. Stelis*, 2.—N. Granat.
aviceps, *Lindl. l. c.* 16.—Peruv.
barbicollis, *Lindl. l. c.* 9.—Peruv.
barrensis, *Lindl. l. c.* 6.—Bras.
bicornis, *Lindl. l. c.* 18.—Peruv.
biflora, Sm. in Rees, Cyclop. xxxiv. n. 13 (Bulbophylli sp.).—Reg. Himal.
biserrula, *Lindl. Fol. Orch. Stelis*, 16.—Peruv.
braccata, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 114.—N. Granat.
brevilabris, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. v. (1845) 107.—N. Granat.
Bruchmuelleri, *Reichb. f. ex Hook. f. Bot. Mag.* t. 6521 (Brueckmuelleri).—Mexic.?
caespitifica, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.
caespitosa, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Bras.
campanulifera, *Lindl. Fol. Orch. Stelis*, 3.—Bolivia.
canaliculata, *Reichb. f. in Gard. Chron.* (1872) 1718.—N. Granat.
capillaris, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Peruv.
capillipes, Reichb. f. in Bonplandia, iii. (1855) 240 = flexuosa.
carnosa, *H. B. & K. Nov. Gen. et Sp.* i. 361.—Am. austr.
catharinensis, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Peruv.
caudata, D. Don, Prod. Fl. Nep. 32 = Bulbophyllum odoratissimum.
chachapoyensis, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Peruv.
chlorantha, *Rodrig. Orch. Nov.* ii. 86.—Bras.
ciliaris, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Mexic.
columnaris, *Lindl. Fol. Orch. Stelis*, 15.—Peruv.
concinna, *Lindl. in Hook. Journ. Bot.* i. (1834) 11.—Peruv.
connata, *Presl, Rel. Haenk.* i. 103.—Peruv.
contorta, *Pers. Syn.* ii. 524.—Peruv.
cordata, *Willd. Sp. Pl.* iv. 140.—Peruv.
coriifolia, *Lindl. Fol. Orch. Stelis*, 18.—Venezuela.
costaricenris, *Reichb. f. in Bonplandia*, iii. (1855) 225.—Costa Rica.
crassifolia, *Lindl. Bot. Reg.* (1842) *Misc.* 9.—N. Granat.
cupuligera, *Reichb. f. in Bonplandia*, ii. (1854) 114.—Peruv.
cymbiformis, *Lindl. Fol. Orch. Stelis*, 11.—Venezuela.
dactyloptera, *Reichb. f. Otia Bot. Hamb.* 19.—Ecuador.
densiflora, *Lindl. Fol. Orch. Stelis*, 16.—Peruv.
depauperata, *Lindl. l. c.* 16.—Peruv.
deregularis, *Rodrig. Orch. Nov.* ii. 94.—Bras.
Dialissa, *Reichb. f. in Bonplandia*, iii. (1855) 71.—N. Granat.
discolor, *Reichb. f. l. c.* 241.—Peruv.

STELIS:—
disticha, *Poepp. & Endl. Nov. Gen. et Sp.* i. 47. t. 81.—Peruv.
drosophila, *Rodrig. Orch. Nov.* ii. 88.—Bras.
dubia, *Lindl. Fol. Orch. Stelis*, 8.—Peruv.
elatior, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Peruv.
elongata, *H. B. & K. Nov. Gen. et Sp.* i. 363.—N. Granat.
endocharis, *Reichb. f. in Linnaea*, xli. (1877) 8.—N. Granat.
Endresii, *Reichb. f. in Gard. Chron.* (1870) 1373.—Costa Rica.
eublepharis, *Reichb. f. in Bonplandia*, iii. (1855) 240.—Peruv.
euspatha, *Reichb. f. l. c.* 225.—Bolivia.
fasciculiflora, *Regel, in Act. Hort. Petrop.* i. (1871) 99.—Hab.?
Fendleri, *Lindl. Fol. Orch. Stelis*, 3.—Venezuela.
filiformis, *Lindl. l. c.* 6.—Bras.
fissa, *Lindl. l. c.* 5.—Peruv.
flacca, *Reichb. f. in Bonplandia*, iii. (1855) 210.—Peruv.
flavida, H. C. Focke, in Tijdschr. Wetensch. ii. (1849) 202 = Pleurothallis flavida.
flexuosa, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 397.—Peruv.
florea, *Lindl. Fol. Orch. Stelis*, 5.—Venezuela.
floribunda, *H. B. & K. Nov. Gen. et Sp.* i. 362.—N. Granat.
foliosa, Hook. in Ann. Nat. Hist. Ser. I. ii. (1839) 330 = Polystachya cerea.
foveata, *Lindl. Fol. Orch. Stelis*, 2.—Peruv.
fraterna, *Lindl. l. c.* 14.—Bras.
Gigas, *Rodrig. Orch. Nov.* ii. 89.—Bras.
gladiata, *Lindl. Fol. Orch. Stelis*, 9.—Peruv.
globiflora, *Reichb. f. Otia Bot. Hamb.* 19.—Ecuador.
glossula, *Reichb. f. in Gard. Chron.* (1870) 1373.—Costa Rica.
glumacea, *Lindl. Fol. Orch. Stelis*, 3.—Peruv.
graminea, *Lindl. l. c.* 11.—Peruv.
grandiflora, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Bras.
grandis, *Reichb. f. in Bonplandia*, iii. (1855) 70.—Venezuela.
grossilabris, *Reichb. f. in Gard. Chron.* (1881) II. 717.—Hab.?
gutturosa, *Reichb. f. in Bonplandia*, ii. (1854) 23.—Venezuela.
Hallii, *Lindl. in Hook. Journ. Bot.* i. (1834) 12.—Peruv.
hemicardia, *Reichb. f. in Linnaea*, xli. (1877) 8.—N. Granat.
herbiola, *Lindl. Fol. Orch. Stelis*, 10.—N. Granat.
Heylidyana, *H. C. Focke, in Tijdschr. Wetensch.* ii. (1849) 201.—Guian.
hirta, *Lindl. Fol. Orch. Stelis*, 3.—Peruv.
hirta, Sm. in Rees, Cyclop. xxxiv. n. 11 = Bulbophyllum hirtum.
humilis, *Lindl. Fol. Orch. Stelis*, 10.—Venezuela.
hylophila, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Peruv.
iminapensis, *Reichb. f. Xenia Orch.* iii. 25.—Bolivia.
intermedia, *Poepp. & Endl. Nov. Gen. et Sp.* i. 46. t. 79.—Peruv.
Jamesoni, *Lindl. in Hook. Journ. Bot.* i. (1834) 11.—Ecuador.
lamellata, *Lindl. in Hook. Ic. Pl.* t. 62.—Ecuador.
lanata, *Lindl. Fol. Orch. Stelis*, 13.—Ecuador.
lancea, *Lindl. l. c.* 3.—Peruv.
lanceolata, *Willd. Sp. Pl.* iv. 139.—Peruv.
lancilabris, *Reichb. f. Beitr. Orch. Centr. Am.* 94. t. 8.—Costa Rica.
lentiginosa, *Lindl. Fol. Orch. Stelis*, 8.—Peruv.
leucopogon, *Reichb. f. Beitr. Orch. Centr. Am.* 95. t. 9.—Costa Rica.
Liebmanni, *Reichb. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 203.—Mexic.
Lindeni, *Lindl. Orch. Linden.* 3.—Venezuela.
littoralis, *Rodrig. Orch. Nov.* ii. 85.—Bras.
lobata, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Peruv.
loxensis, *Lindl. in Benth. Pl. Hartw.* 149.—Ecuador.
lutea, *Lindl. Fol. Orch. Stelis*, 7.—Venezuela.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* i. 363.—N. Granat.
macrostachya, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 643, nomen (Quid?).—Am. centr.
major, *Reichb. f. in Bonplandia*, ii. (1854) 23.—Venezuela.

STELIS:—
maxima, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 106.—Peruv.
megantha, *Rodrig. Orch. Nov.* ii. 83.—Bras.
melanoxantha, *Reichb. f. in Bonplandia*, iii. (1855) 71.—Venezuela.
micrantha, *Rodrig. Orch. Nov.* ii. 93.—Bras.
micrantha, *Sw. in Schrad. Journ.* ii. (1799) 240.—Jamaic.
microcaulis, *Rodrig. Orch. Nov.* ii. 85.—Bras.
microglossa, *Reichb. f. in Linnaea*, xxii. (1849) 821.—Bras.
microphylla, *Lindl. Fol. Orch. Stelis*, 10.—Peruv.
microstigma, *Reichb. f. Beitr. Orch. Centr. Am.* 94. t. 8.—Costa Rica.
microtis, *Reichb. f. l. c.* 95. t. 8.—Costa Rica.
Miersii, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 396.—Bras.
Miqueliana, Lindl. Fol. Orch. Pleuroth. 17, sphalm. = Pleurothallis Miqueliana.
modesta, *Rodrig. Orch. Nov.* ii. 87.—Bras.
mononeura, *Lindl. Fol. Orch. Stelis*, 4.—Peruv.
muscifera, *Lindl. l. c.* 2.—Venezuela.
muscosa, *Lindl. l. c.* 7.—Venezuela.
myriantha, *Lindl. l. c.* 10.—Peruv.
nana, *Lindl. l. c.* 16.—Peruv.
nanegalensis, *Lindl. l. c.* 8.—Peruv.
nitens, *Reichb. f. in Bonplandia*, ii. (1854) 22.—Venezuela.
nutans, *Lindl. Fol. Orch. Stelis*, 15.—Peruv.
oblonga, *Willd. Sp. Pl.* iv. 139.—Peruv.
oblongifolia, *Lindl. Fol. Orch. Stelis*, 12.—Peruv.
obscurata, *Reichb. f. Beitr. Orch. Centr. Am.* 95. t. 8.—Costa Rica.
ochreata, *Lindl. Fol. Orch. Stelis*, 2.—Peruv.
odoratissima, Sm. in Rees, Cyclop. xxxiv. n. 12 = Bulbophyllum odoratissimum.
oligantha, *Rodrig. Orch. Nov.* ii. 93.—Bras.
omalosantha, *Rodrig. l. c.* i. 34.—Bras.
ophioglossoides, *Sw. in Schrad. Journ.* ii. (1799) 239.—Am. austr.
ovalifolia, H. C. Focke, in Tijdschr. Wetensch. ii. (1849) 202 = Pleurothallis ovalifolia.
pachystachya, *Lindl. Fol. Orch. Stelis*, 13.—N. Granat.
palmeiraënsis, *Rodrig. Orch. Nov.* ii. 92.—Bras.
papaquerensis, *Reichb. f. in Linnaea*, xxii. (1849) 822.—Bras.
parahybunensis, *Rodrig. Orch. Nov.* ii. 84.—Bras.
paraënsis, *Rodrig. Orch. Nov.* ii. 88.—Bras.
pardipes, *Reichb. f. Beitr. Orch. Centr. Am.* 96.—Costa Rica.
parviflora, Pers. Syn. ii. 524 = micrantha, *Sw.*
parvilabris, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xv. (1845) 385.—N. Granat.
parvula, *Lindl. Fol. Orch. Stelis*, 7.—Nicaragua.
pauciflora, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 396.—Bras.
peliochyla, *Rodrig. Orch. Nov.* ii. 84.—Bras.
penduliflora, *Rodrig. l. c.* 91.—Bras.
pentodonta, *Reichb. f. in Linnaea*, xxii. (1849) 821.—Am. trop.
petropolitana, *Reichb. f. in Wawra, Bot. Ergeb. Maxim. Bras.* 151.—Bras.
philargyrus, *Reichb. f. in Linnaea*, xli. (1877) 118.—Venezuela.
piperina, *Lindl. Fol. Orch. Stelis*, 17.—Peruv.
polybotrya, *Lindl. l. c.* 4.—Peruv.
polyclada, *Lindl. l. c.* 10.—Peruv.
polystachya, *Willd. Sp. Pl.* iv. 139.—Am. austr.
Porpax, *Reichb. f. in Bonplandia*, ii. (1854) 23.—Venezuela.
puberula, *Rodrig. Orch. Nov.* ii. 86.—Bras.
Pugiunculi, *Lindl. Fol. Orch. Stelis*, 17.—Peruv.
pulchella, H. B. & K. Nov. Gen. et Sp. i. 364. t. 90 = Pleurothallis pulchella.
Purdiaei, *Lindl. Fol. Orch. Stelis*, 5.—N. Granat.
purpurascens, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 18.—Mexic.
purpurea, Willd. Sp. Pl. iv. 140 = Pleurothallis linearis.
pusilla, *H. B. & K. Nov. Gen. et Sp.* i. 361.—Ecuador; Jamaic.
racemiflora, *Lodd. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 643, nomen (Quid?).—Jamaic.
racemosa, Sm. in Rees, Cyclop. xxxiv. n. 10 = Liparis odorata?
reflexa, *Lindl. Fol. Orch. Stelis*, 3.—Peruv.
revoluta, *Willd. Sp. Pl.* iv. 140.—Peruv.

STELIS:—
Ruprechtiana, *Reichb. f. in Linnaea*, xxii. (1849) 821.—Bras.
scabrida, *Lindl. in Ann. Nat. Hist.* Ser. I. v. (1840) 115.—Ins. Dominic.
Scansor, *Reichb. f. in Bonplandia*, iii. (1855) 241.—Peruv.
Serra, *Lindl. in Ann. & Mag. Nat. Hist.* Ser. I. xii. (1843) 397.—Peruv.
sesquipedalis, *Lindl. Orch. Linden.* 3.—Venezuela.
setacea, *Lindl. Fol. Orch. Stelis*, 3.—Bolivia.
smaragdina, *Rodrig. Orch. Nov.* ii. 90.—Bras.
spathulata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 46. t. 80.—Peruv.
sphaerochila, *Lindl. Fol. Orch. Stelis*, 13.—Venezuela.
spiralis, Pers. Syn. ii. 524 = Pleurothallis spiralis.
stenophylla, *Reichb. f. in Bonplandia*, iii. (1855) 70.—Peruv.
striolata, *Lindl. Fol. Orch. Stelis*, 4.—Peruv.
superbiens, *Lindl. l. c.* 8.—N. Granat.
tenuicaulis, *Lindl. l. c.* 16.—Bolivia.
tenuilabris, *Lindl. l. c.* 4.—Venezuela.
thecoglossa, *Reichb. f. Beitr. Orch. Centr. Am.* 93.—Costa Rica.
triangularis, *Rodrig. Orch. Nov.* ii. 91.—Bras.
tricardium, *Lindl. Fol. Orch. Stelis*, 14.—Peruv.
trichorrhachis, *Reichb. f. in Bonplandia*, iii. (1855) 71.—N. Granat.
tridentata, *Lindl. Fol. Orch. Stelis*, 6.—Peruv.
triplicata, *Lindl. l. c.* 2.—Peruv.
triseta, *Lindl. l. c.* 17.—Bolivia.
tristyla, *Lindl. Bot. Reg.* (1838) *Misc.* 42.—Bras.
triura, *Lindl. Orch. Linden.* 3 (an gen.?).—Venezuela.
truncata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 353.—Peruv.
tubata, Lodd. Bot. Cab. t. 1601 = Physosiphon Loddigesii.
Tweediana, *Lindl. Fol. Orch. Stelis*, 7.—Bras.
velutina, *Lindl. l. c.* 17.—Peruv.
vinosa, *Rodrig. Orch. Nov.* ii. 89.—Bras.
viridipurpurea, *Lindl. Fol. Orch. Stelis*, 3.—Bras.
vittata, *Lindl. l. c.* 13.—Peruv.
Vulcani, *Reichb. f. Otia Bot. Hamb.* 19.—Ecuador.
zonata, *Reichb. f. in Gard. Chron.* (1883) II. 556.—Guian.

STELLA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 377 = **Astragalus**, Tourn. (Legumin.).
radiata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364 = A. Stella.
sesamea, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 377 = A. sesameus.

STELLANDRIA, Brickell, in Med. Repos. N. York, vi. (1803) 327 = **Schizandra**, Michx. (Magnol.).
glabra, Brickell, l. c. = Sch. coccinea.

STELLARA, Fisch. ex Reut. in DC. Prod. xi. 39 (1847) = **Boschniakia**, C. A. Mey. (Orobanch.).
lathraeoides, Fisch. ex Reut. l. c. = B. glabra.

STELLARIA, Hall. Enum. Stirp. Helv. i. 198 (1742) = **Callitriche**, Linn. (Halor.).
autumnalis, [Weber, in] Wigg. Prim. Fl. Holsat. 2 = C. autumnalis.
intermedia, [Weber, in] l. c. = C. verna.
verna, [Weber, in] l. c. = C. verna.

STELLARIA, Linn. Sp. Pl. ed. I. 421 (1753).
CARYOPHYLLEAE, Benth. & Hook. f. i. 149.
ADENONEMA, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1836) 448 (Verz. Suppl. Fl. Alt. 36).
DICHODON, Reichb. Nom. 205 (1841).
HYLEBIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347.
LARBREA, A. St. Hil. in Mém. Mus. Par. ii. (1815) 287.
KRASCHENINIKOVIA, Turcz. ex Bess. in Flora, xvii. (1834) I. Beibl. 9.
LEUCOSTEMMA, Benth. in Wall. Cat. n. 642 (1828); et ex G. Don, Gen. Syst. i. 449 (1831).
MALACHIUM, Fries, Fl. Halland. 77 (1817).
MICROPETALON, Pers. Syn. i. 509 (1805).
MYOSANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 348.
MYOSOTON, Moench, Meth. 225 (1794).
SCHIZOTECHIUM, Reichb. Nom. 205 (1841).

STELLARIA:—
SPERGULASTRUM, Michx. Fl. Bor. Am. i. 275 (1803).
STELLULARIA, Linn. Syst. ed. VI (1748).
abortiva, *C. Gay, Fl. Chil.* i. 265.—Chili.
aculeata, Scop. Fl. Carn. ed. II. i. 316 = Arenaria grandiflora.
adulterina ×, Focke, in Bremen, Abh. i. (1867) 145.—Europ.
alpestris, Fries, Mant. i. 10 = borealis.
alpestris, Turcz. ex Ledeb. Fl. Ross. i. 387 = longipes.
alpicola, Lamotte, Prod. Fl. Centr. Fr. i. 148 = media.
alpina, S. F. Gray, Nat. Arr. Brit. Pl. ii. 660 = Cerastium alpinum.
alsinaefolia, Less. ex Ledeb. Fl. Ross. i. 380 = dichotoma.
Alsine, Afzel. & Wadsb. in Roem. Archiv, iii. (1803) 97 = media.
Alsine, Bunge, in Ledeb. Fl. Alt. ii. 156 = crassifolia.
Alsine, Hoffm. Deutschl. Fl. ed. I. 153 = uliginosa.
alsinoides, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 41.—Persia.
alsinoides, Schleich. Cat. Pl. Helv. ed. III. 29 = media.
altenoides, C. Muell. in Walp. Ann. vii. 291, sphalm. = alsinoides.
amblyosepala, *Schrenk, Enum. Pl. Nov.* ii. 54.—Soongar.
anagalloides, *Rupr. Fl. Cauc.* 222.—Reg. Cauc.
angustifolia, Hook. Journ. Bot. i. (1834) 250 = palustris.
anomala, Reichb. Fl. Germ. Excurs. 785 = Cerastium anomalum.
apetala, Opiz, ex Nym. Consp. 111 = media.
apetala, *Ucria, in Roem. Arch.* i. 1. (1796) 68.—Sicil.
aphanantha, *Griseb. in Goett. Abh.* xxiv. (1879) 27.—Reg. Argent.
aquatica, Cham. & Schlecht. in Linnaea, i. (1826) 50 = borealis.
aquatica, Pollich, Hist. Pl. Palat. i. 429 = uliginosa.
aquatica, *Scop. Fl. Carn.* ed. II. i. 319.—Europ.; As. bor.
Arenaria, Linn. Sp. Pl. 1196 = Arenaria spathulata.
aristata, *Ser. in DC. Prod.* i. 396.—Mexic.
arvalis, Phil. f. Cat. Pl. Vasc. Chil. 26 = prostrata.
arvensis, S. F. Gray, Nat. Arr. Br. Pl. ii. 660 = Cerastium arvense.
arvensis, Scop. Fl. Carn. ed. II. i. 318 = Spergula arvensis.
australis, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 581.—Java.
Baldwini, *Fenzl, ex Peyr. in Linnaea*, xxx. (1859–60) 57.—Mexic.
Barthiana, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 142 = graminea.
biflora, Host, Syn. Pl. Austr. 243 = Arenaria polygonoides.
biflora, *Linn. Sp. Pl.* 422.—Europ.
biflora, Pursh, Fl. Am. Sept. i. 317 = Arenaria lateriflora.
biflora, Rudolph, ex Ledeb. Fl. Ross. i. 387 = florida.
biflora, Wulf. in Jacq. Coll. i. 351 = Arenaria austriaca.
Boraeana, Jord. Pugill. Pl. Nov. 33 = media.
borealis, Bieb. ex Steud. Nom. ed. II. ii. 636 = petraea.
borealis, *Bigel. Fl. Bost.* ed. II. 182.—Reg. bor. et arct.
borealis, Darling. Fl. Cestr. 274 = uliginosa.
brachypetala, Bong. Vég. Ins. Sitcha, 126 = borealis.
brachypetala, *Bunge, in Ledeb. Fl. Alt.* ii. 161.—Sibir. altaic.
bracteata, Richt. in Klatt & Richt. Fl. Leipz. 386 = uliginosa.
Brauniana, *Fenzl, ex Schweinf. Beitr. Fl. Aethiop.* 60.—Afr. trop.
breviflora, Gilib. Fl. Lituan. ii. 151 = uliginosa.
brevifolia, Borkh. in Rhein. Mag. i. (1793) 523 = graminea.
brevifolia, Schum. Pl. Saell. 142 = crassifolia.
Brotherana, *Trautv. in Act. Hort. Petrop.* viii. (1883) 153.—Reg. Cauc.
bulbifera, *Benth. in Journ. Linn. Soc.* vi. (1862) 67.—Europ. centr.
bulbosa, *Wulf. in Jacq. Coll.* iii. 21.—Europ. or.; As. bor.; Reg. Himal.

STELLARIA:—
Bungeana, Fenzl, in Ledeb. Fl. Ross. i. 376 = nemorum.
caespitosa, Duf. in Act. Soc. Linn. Bord. viii. (1836) 91 = radicans?
caespitosa, Hook. f. in Hook. Journ. Bot. ii. (1840) 411 = palustris.
caespitosa, Vest, ex Steud. Nom. ed. II. ii. 636 = holostea.
calycantha, Bong. Vég. Ins. Sitcha, 127 = borealis.
Cerastium, Murr. Syst. ed. XV. 452; ex DC. Prod. i. 396 = media.
cerastoides, Linn. Sp. Pl. 422 = Cerastium trigynum.
cerastoides, Merlet, ex Steud. Nom. ed. I. 813 = Cerastium anomalum.
chinensis, *Regel, in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 283, in textu.—China.
ciliata, Gilib. Fl. Lituan. ii. 152 = S. Holostea.
ciliata, Scop. Fl. Carn. ed. II. i. 315 = Arenaria polygonoides.
ciliata, *Vahl, ex Pers. Syn.* i. 501.—Peruv.
ciliata, Willd. ex H. B. & K. Nov. Gen. et Sp. vi. 26 = ovata.
ciliatosepala, *Trautv. in Middend. Sibir. Reise*, i. II. 52. t. 8.—Sibir.
cilicica, *Boiss. & Bal. Diagn.* Ser. II. v. 59.—As. Min.; Syria.
connata, Dulac, Fl. Hautes-Pyr. 250 = S. Holostea.
cordata, Gilib. Fl. Lituan. ii. 152 = nemorum.
cordata, S. G. Gmel. ex Ledeb. Fl. Ross. i. 402 = Cerastium davuricum.
cordata, *Willd. ex Sehlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 197.—Am. austr.
crassifolia, Boland. Cat. Pl. Calif. 6, nomen = borealis.
crassifolia, Cham. & Schlecht. in Linnaea, i. (1826) 50 = humifusa.
crassifolia, *Ehrh. in Hannov. Mag.* 8tes St. (1784) 116; *et ex J. F. Gmel. Syst.* 718.—Reg. bor. et arct.
crassifolia, S. Wats. Bot. King's Exped. 38 = longipes.
crispa, *Cham. & Schlecht. in Linnaea*, i. (1826) 51.—Fret. Behring.
crispata, *Wall. ex D. Don, Prod. Fl. Nep.* 215; *Cat.* n. 633.—Reg. Himal.
cryptopetala, *Griseb. in Goett. Abh.* xxiv. (1879) 27.—Reg. Argent.
Cupaniana, Nym. Consp. 111 = media.
cuspidata, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 196 = nemorum.
davurica, Spreng. Syst. ii. 398 = petraea.
davurica, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 195.—Dahuria.
debilis, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 618.—Patagon.
decandra, Schur, Enum. Pl. Transs. 970 (Index) = media.
decipiens, *Hook. f. Fl. Antarct.* 7.—Ins. Dom. Auckland.
decumbens, *Edgew. in Trans. Linn. Soc.* xx. (1846) 35.—Reg. Himal.
depauperata, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 234.—Reg. Himal.
diandra, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 379.—Japon.
dichotoma, Georgi, Beschr. Russ. Reich. III. iv. 976 = media.
dichotoma, *Linn. Sp. Pl.* 421.—Sibir.
dichotoma, Patrin, ex Ledeb. Fl. Alt. ii. 153 = nemorum.
dichotoma, Scop. Fl. Carn. ed. II. i. 314 = bulbosa.
dichotoma, [Weber, in] Wigg. Prim. Fl. Holsat. 34 = crassifolia.
dicranoides, *Fenzl, in Ledeb. Fl. Ross.* i. 395.—Fret. Behring.
diffusa, Pall. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 195 = longifolia.
Dilleniana, Leers, Fl. Herborn. 107 = uliginosa.
Dilleniana, Moench, Enum. Pl. Hass. 214 = palustris.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 88; xv. (1842) 601.—Dahuria.
diversiflora, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 379, in nota.—Japon.
domingensis, *Spreng. Syst.* ii. 392.—Ins. S. Doming.
drymarioides, *Thw. Enum. Pl. Zeyl.* 24.—Zeylan.
dubia, Bast. Fl. Maine-et-Loire, Suppl. 24 = Cerastium anomalum.
eciliata, *Rupr. Fl. Cauc.* 297.—Reg. Cauc.

STELLARIA:—
Edwardsii, R. Br. in Parry, 1st Voy. App. p. cclxxi = longipes.
elatinoides, *Hook. f. Fl. N. Zel.* i. 25.—N. Zel.
elegans, *Meinsh. in Baer & Helm. Beitr. Kenntn. Russ. Reich.* xxvi. 146.—Sibir. or.
elegans, Ser. in DC. Prod. i. 400 = Cerastium trigynum.
Elisabethae, F. Schultz, Arch. Fl. 302 = media.
elodes, Bieb. ex Bess. Enum. Pl. Volh. 81 = crassifolia.
elongata, Nutt. Gen. Am. i. 289 = Arenaria lanuginosa.
eriantha, Turcz. ex Ledeb. Fl. Ross. i. 388 = davurica.
Eschscholtziana, *Fenzl, in Ledeb. Fl. Ross.* i. 384.—Kamtschat.
exigua, Steph. ex Steud. Nom. ed. II. ii. 636 = petraea.
falcata, Ser. in DC. Prod. i. 398 = davurica.
Fenzliana, Klotzsch, in Bot. Ergeb. Waldem. Reise, 141 = longissima.
Fenzlii, Regel, in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 264, 280 = borealis.
Fischeriana, Ser. in DC. Prod. i. 398 = florida.
flaccida, *Hook. Comp. Bot. Mag.* i. (1835) 275.—Austral.
flaccida, Peterm. Fl. Lips. Excurs. 324 = graminea?
florida, *Fisch. ex DC. Prod.* i. 399.—Camtschat.
foecunda, Hance, in Walp. Ann. ii. 95 = aquatica.
fontana, Wulf. in Jacq. Coll. i. 327 = uliginosa.
Frieseana, Turcz. ex Ledeb. Fl. Ross. i. 388 = davurica.
Friesiana, *Ser. in DC. Prod.* i. 397.—Europ. bor.
geniculata, Gueldenst. Reisen, i. 192; ex Ledeb. Fl. Ross. i. 395 (Quid?).—Reg. Cauc.
glabella, Nym. Consp. 111 = media.
glacialis, Lagg. in Oestr. Bot. Zeitzschr. xviii. (1868) 242 = uliginosa.
glandulifera, Kit. in Linnaea, xxxii. (1863) 515 = glandulosa.
glandulifera, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 141.—Ind. or.
glandulosa, *Kit. in Schult. Oestr. Fl.* ed. II. 673.—Hungar.
glareosa, Turcz. ex Steud. Nom. ed. II. ii. 636 = Cerastium trigynum.
glauca, Frivald. ex Griseb. Spicil. Fl. Rumel. i. 207 = Cerastium manticum.
glauca, E. Mey. Pl. Lab. 93 = longipes.
glauca, Salisb. Prod. 301 = S. Holostea.
glauca, Turcz. ex Ledeb. Fl. Ross. i. 388 = davurica.
glauca, *With. Bot. Arr. Brit. Pl.* ed. III. ii. 420.—Europ.; Sibir.
Gmelini, Nesterofsky, ex Fenzl, in Ledeb. Fl. Ross. i. 293 = Gypsophila uralensis.
gracilenta, *Hook. f. Fl. N. Zel.* ii. 326.—N. Zel.
gracilis, Lange, ex Hage, Obflod. in Ersler, Geogr. Tidskr. v. (1881) Hefte ix–x. 144 = crassifolia.
gracilis, Richards, in Frankl. 1st Journ. App. 738 = uliginosa.
graminea, Bigel. Fl. Bost. ed. I. 110 = longifolia.
graminea, Cham. & Schlecht. in Linnaea, i. (1826) 48 = longipes.
graminea, Grev. in Mem. Wern. Soc. iii. (1820) 429 = longipes?
graminea, James, in Trans. Am. Phil. Soc. ii. (1825) 181, nomen = Jamesiana.
graminea, *Linn. Sp. Pl.* 422.—Europ.; As. bor.
graminifolia, Gmel. ex Fenzl, in Ledeb. Fl. Ross. i. 381 = S. Holostea.
graminifolia, Salisb. Prod. 301 = graminea.
grandiflora, Gilib. Fl. Lituan. ii. 151 = graminea.
grandiflora, Woods, in Hook. Comp. Bot. Mag. ii. (1836) 264 = media.
groenlandica, Retz. Prod. Fl. Scand. ed. II. 107 = Arenaria groenlandica.
gypsophiloides, *Fenzl, in Ledeb. Fl. Ross.* i. 380.—China.
hebecalyx, *Fenzl, in Rupr. Beitr. Pfl. Russ. Reich.* ii. 26.—Lapponia.
helodes, Auct. = *elodes*.
heterophylla, Magn. in Bull. Soc. Bot. Fr. xxviii. (1881) 82 = glauca.
Holostea, *Linn. Sp. Pl.* 422.—Europ.; Reg. Cauc.
humifusa, Retz. Fl. Scand. ed. I. 84 = crassifolia.
humifusa, *Rottb. in Skrivt. Vidensk. Selsk. Kiob.* x. (1770) 447.—Reg. bor. et arct.
hypericifolia, All. Fl. Pedem. ii. 115 = uliginosa.
hypericifolia, [*Weber, in*] *Wigg. Prim. Fl. Holsat.* 34.—Europ.

STELLARIA:—
imbricata, Bong. ex Fenzl, in Ledeb. Fl. Ross. i. 385 = Eschscholtziana.
imbricata, *Bunge, in Ledeb. Fl. Alt.* ii. 159.—Sibir. altaic.
irrigua, *Bunge, Verz. Suppl. Fl. Alt.* 35.—Sibir. altaic.
Jamesiana, *Torr. in Ann. Lyc. N. York*, ii. (1828) 169.—Am. bor. occ.
Jamesii, Steud. Nom. ed. II. ii. 636, sphalm. = praec.
Jamesonii, G. Don, Gen. Syst. i. 428, sphalm. = praec.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 79.—Japon.
juncea, Fries, Summa Veg. Scand. 37 = graminea.
Kingii, *S. Wats. in Bot. King's Exp.* 39. t. 6.—Am. bor. occ.
Kotschyana, *Fenzl, ex Boiss. Fl. Orient.* i. 705.—Persia.
labradorica, Schrank, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 24 = Arenaria groenlandica.
laeta, Richards. in Frankl. 1st Journ. App. 738 = longipes.
laevis, *Rohrb. in Linnaea*, xxxvii. (1871–73) 275.—Peruv.
lanata, *Hook. f. Fl. Brit. Ind.* i. 232.—Reg. Himal.
lanceolata, *Poir. Encyc.* vii. 416.—Reg. Magellan.
lanceolata, Torr. Fl. U. St. 453 = borealis.
lanuginosa, Torr. & Gray, Fl. N. Am. i. 187 = Arenaria lanuginosa.
laricifolia, Scop. Fl. Carn. ed. II. i. 317 = Arenaria laricifolia, recurva.
lateriflora, Fleisch. in Bull. Soc. Nat. Mosc. i. (1829) 101 = crassifolia.
lateriflora, Krock. Fl. Siles. ii. 1. 52 = uliginosa.
latifolia, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 231.—Reg. Himal.
latifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 660 = Cerastium latifolium.
latifolia, Pers. Syn. i. 501 = media.
Laxmanni, Fisch. ex DC. Prod. i. 397 = glauca.
leptopetala, *Benth. Pl. Hartw.* 163.—N. Granat.
leptophylla, *Hance, in Walp. Ann.* ii. 96.—China.
ligustrina, Spreng. Syst. iv. Cur. Post. 180 = Schiedea ligustrina.
linoides, Tausch, in Flora, xix. (1836) 413 = uliginosa.
litigiosa, Magn. in Bull. Soc. Bot. Fr. xxviii. (1881) 82 = glauca.
littoralis, *Torr. in Pacif. Rail. Rep.* iv. 69.—Calif.
longicaulis, Krock. Fl. Siles. Suppl. ii. 90 (Quid?).—Europ.
longifolia, Fries, in Svensk Bot. viii. t. 506 = Friesiana.
longifolia, *Muhl. ex Willd. Enum. Hort. Berol.* 479.—Sibir.; Am. bor.
longifolia, Steud. Nom. ed. II. ii. 636, sphalm. = longissima.
longifolia, Turcz. in Bull. Soc. Nat. Mosc. (1838) 88 = graminea.
longipes, *Goldie, in Edinb. Phil. Journ.* vi. (Apr. 1822) 327.—Am. bor. et arct.
longissima, *Wall. Cat.* n. 632.—Reg. Himal.
lycopodioides, Fenzl, ex Rohrb. in Linnaea, xxxvii. (1871–73) 275 = laevis.
macropetala, *Torr. & Gray, Fl. N. Am.* i. 184.—Am. bor. occ.
magna, Gueldenst. Reisen, i. 427; ex Ledeb. Fl. Ross. i. 395 (Quid?).—Reg. Cauc.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 183.—Afr. trop.
mantica, DC. Fl. Fr. iv. 794 = Cerastium manticum.
marginata, Cham. & Schlecht. in Linnaea, i. (1826) 50 = humifusa.
media, *Cyrill. Char. Comm.* (1784) 36.—Geront. temp.
media, Sibth. Fl. Oxon. 141 = palustris.
media, Wight, ex Edgew. in Hook. f. Fl. Brit. Ind. i. 229 = crispata.
Meyeriana, *Rupr. Fl. Cauc.* 297. t. 5. f. 3.—Reg. Cauc.
micrantha, *Spruce, ex Rohrb. in Linnaea*, xxxvii. (1871–73) 280.—Ecuador.
microphylla, Schur, Enum. Pl. Transs. 116 = media.
modesta, *Fenzl, in Tijdschr. Nederl. Ind.* xiv. (1857) 165 (Quid?).—Malaya.
Moenchii, Magn. in Bull. Soc. Bot. Fr. xxviii. (1881) 82 = glauca.
moenchioides, Fenzl, ex Torr. & Gray, Fl. N. Am. i. 675 = nitens.
mollis, Fisch. ex DC. Prod. i. 397 = velutina.

STELLARIA:—
mollis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 142 = longissima.
mollis, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 196 = Cerastium Willdenowii.
mollis, Wirzén, Geogr. Pl. Prov. Casan. 43. n. 95 = borealis.
molluginea, Sweet, Hort. Brit. ed. II. 56 = Arenaria molluginea.
monogyna, D. Don, Prod. Fl. Nep. 215 = media.
monosperma, Buch.-Ham. ex D. Don, l. c. = crispata.
mosquiensis, Bieb. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 195 = longifolia.
multicaulis, Willd. Sp. Pl. ii. 714 = Cerastium trigynum.
multiflora, *Hook. Comp. Bot. Mag.* i. (1835) 275.—Austral.
muralis, Link, Enum. Hort. Berol. i. 429 = Arenaria graveolens.
Murmuria, Buch.-Ham. ex Wall. Cat. sub n. 630. p. 248 = media.
neglecta, Weihe, in Bluff & Fingerh. Comp. Fl. Germ. ed. I. i. 560 = media.
nemoralis, Salisb. Prod. 301 = nemorum.
nemorum, *Linn. Sp. Pl.* 421.—Europ.; Reg. Cauc.
nemorum, Stev. ex Fenzl, in Ledeb. Fl. Ross. i. 377 = media.
nitens, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 185, 675.—Am. bor. occ.
nitida, Hook. in Scoresby, Greenl. 411 = longipes.
nodosa, Scop. Fl. Carn. ed. II. i. 319 = Sagina nodosa.
Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 183.—Texas.
obtusa, *Engelm. in Coult. Bot. Gaz.* vii. (1882) 5.—Am. bor.
oligosperma, *Rohrb. in Linnaea*, xxxvii. (1871–73) 276.—Chili.
ovalifolia, Hook. Fl. Bor. Am. i. 97 = Arenaria physodes.
ovata, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 196.—Am. austr.
Pallasiana, Ser. in DC. Prod. i. 399 = dichotoma.
pallida, Dum. ex Nym. Consp. 111 = media.
paludosa, Laest. ex Fries, Summa Veg. Scand. 37 = crassifolia.
palustris, *Ehrh. Beitr.* v. 176, nomen; *Retz. Fl. Scand. Prod.* ed. II. 106.—Geront. temp.; Greenl.
palustris, Richards. in Frankl. 1st Journ. App. 738 = longipes.
paniculata, Edgew. in Trans. Linn. Soc. xx. (1846) 35 = crispata.
paniculata, Pall. Tabl. Taur. 51 = graminea.
parviflora, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 25.—N. Zel.
patens, D. Don, Prod. Fl. Nep. 215 = longissima.
pauciflora, *Zoll. & Mor. Syst. Verz. Zoll.* 30.—Java.
pedunculariis, Bunge, in Ledeb. Fl. Alt. ii. 157 = longipes.
pentagyna, Gaud. Fl. Helv. iii. 179 = aquatica.
pentandra, Scop. Fl. Carn. ed. II. i. 318 = Spergula pentandra.
persica, Boiss. Diagn. Ser. II. i. 85 = palustris.
petraea, *Bunge, in Ledeb. Fl. Alt.* ii. 160.—Sibir. altaic.
petraea, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 234 = decumbens.
pilosa, Dulac, Fl. Hautes-Pyr. 250 = media.
prostrata, *Baldw. in Ell. Sketch*, i. 518.—Am. bor. et austr.
Pseudo-malachium, Schur, Enum. Pl. Transs. 117 = nemorum.
pubera, *Michx. Fl. Bor. Am.* i. 273.—Am. bor.
pubescens, Willd. ex H. B. & K. Nov. Gen. et Sp. vi. 35 = Arenaria lanuginosa.
pumila, Brocchi, in Bibl. Ital. xxviii. (1822) 223 = Cerastium Thomasii.
pumila, Fenzl, Verbr. Alsin. in Tab. Synop. ad p. 18 = petraea.
pumila, S. F. Gray, Nat. Arr. Br. Pl. ii. 659 = Cerastium pumilum.
pungens, *Brongn. Voy. Coq. Bot.* t. 78.—Austral.
radians, *Linn. Sp. Pl.* 422.—Sibir.
radicans, *Lapeyr. Hist. Abr. Pl. Pyr.* 250.—Mont. Pyren.
recurvata, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 196.—N. Granat.
Reichenbachii, Wierzb. ex Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 76 = nemorum.
repens, Scop. Fl. Carn. ed. II. i. 320 = Cerastium tomentosum.

STELLARIA:—
rigida, *Bunge, in Act. Hort. Petrop.* v. (1877) 251.—Turkestan.
rotundifolia, Humb. & Bonpl. ex Steud. Nom. ed. I. 813 = cordata.
rotundifolia, *Poir. Encyc.* vii. 416.—Reg. Magellan.
Roughii, *Hook. f. Handb. N. Zeal. Fl.* 23.—N. Zel.
rubra, Scop. Fl. Carn. ed. II. i. 316 = Arenaria fasciculata, *Gouan*.
rupestris, Scop. l. c. 317. t. 18. f. 1 = Arenaria lanceolata.
ruscifolia, *Pall. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 194.—Sibir. or.; Am. bor. occ.
sabulosa, Fisch. ex DC. Prod. i. 397 = Cerastium anomalum.
saxatilis, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 215.—Reg. Himal.; As. bor.; Japon.
saxatilis, Scop. Fl. Carn. ed. II. i. 317 = Arenaria saxatilis.
Saxifraga, Bertol. Pl. Ital. Rar. iii. 55 = Arenaria Saxifraga.
scapigera, *Willd. Sp. Pl.* ii. 716.—Scotia.
Schlechtendaliana, Ser. in DC. Prod. i. 399 = dichotoma.
semidecandra, Link, Diss. Bot. Suerin (1795), 62 = Cerastium semidecandrum.
semivestita, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 230.—Reg. Himal.
serpyllifolia, Scop. Fl. Carn. ed. II. i. 319 = Arenaria serpyllifolia.
serpyllifolia, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1816) 196.—Peruv.
sikkimensis, *Hook. f. Fl. Brit. Ind.* i. 230.—Reg. Himal.
sitchana, Steud. Nom. ed. II. ii. 637 = borealis.
spatulata, *Reichb. Ic. Fl. Germ.* v. 35.—Europ.
squarrosa, Hook. Journ. Bot. i. (1834) 250 = pungens.
stenopetala, *Phil. in Linnaea*, xxxiii. (1864–65) 20.—Chili.
Stephaniana, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 194 = dichotoma.
stricta, Hook. Fl. Bor. Am. i. 96, partim = nitens.
stricta, C. Koch, in Linnaea, xv. (1841) 707 = palustris.
stricta, Richards. in Frankl. 1st Journ. App. ed. II. 15 = longipes.
stricta, Sw. ex Steud. Nom. ed. II. ii. 637 = Arenaria lapponica.
subulata, Boeb. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 195 = graminea.
subumbellata, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 233.—Reg. Himal.
sylvatica, *Maxim. ex Regel, in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 302.—Reg. Amur.
tenella, Fenzl, in Zoll. Syst. Verz. Ind. Arch. 142; et in Tijdschr. Nederl. Ind. xiv. (1857) 165 = pauciflora.
tenella, Rafin. in Am. Monthly Mag. (1818) 344; Bieb. ex Ledeb. Fl. Ross. i. 392 = longifolia.
tetragona, *Blume, Bijdr.* 63.—Java.
tetrandra, S. F. Gray, Nat. Arr. Brit. Pl. ii. 659 = Cerastium tetrandrum.
thymifolia, Wall. Cat. n. 636 = uliginosa.
tibetica, *Kurz, in Flora*, lv. (1872) 285.—Reg. Himal.
tomentosa, Link, Diss. Bot. Suerin (1795), 62 = Cerastium tomentosum.
tomentosa, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 381.—Japon.
triandra, Debray, ex Bluff Fingerh. Comp. Fl. Germ. ed. I. i. 171, nomen = Holosteum umbellatum.
trivialis, Link, Diss. Bot. Suerin (1795), 62 = Cerastium triviale.
tschachanensis, Less. ex Fenzl, in Ledeb. Fl. Ross. i. 380 = dichotoma.
uliginosa, Laest. in Vet. Akad. Handl. Stockh. (1822) 335 = borealis.
uliginosa, *Murr. Prod. Stirp. Gotting.* 55.—Reg. bor. temp. et arct.
umbellata, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 89; xv. (1842) 173.—Sibir. or.; Am. bor. occ.
umbrosa, Opiz & Rupr. in Opiz, Seznam, 93 = media.
undulata, Regel, in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 282 = chinensis.
undulata, Thunb. Fl. Jap. 185 = uliginosa.
uniflora, *Walt. Fl. Carol.* 141.—Am. bor.
velutina, *Ser. in DC. Prod.* i. 397.—Sibir.
vestita, Kurz, in Journ. Bot. xi. (1873) 194 = saxatilis.
villosa, *Poir. Encyc.* vii. 416.—Ins. Borbon.

STELLARIA:—
villosa, Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 194 = dichotoma.
virens, *Rupr. Fl. Cauc.* 222.—Reg. Cauc.
virgata, *Ser. in DC. Prod.* i. 396.—Hab.?
viscida, Bieb. Fl. Taur. Cauc. i. 342 = Cerastium anomalum.
viscosa, Albert, in Bull. Soc. Sc. Draguign. (1884), reimp. 13 (Quid?).—Gallia.
viscosa, Link, Diss. Bot. Suerin (1795), 62 = Cerastium triviale.
vulgata, Link, l. c. = Cerastium glomeratum.
Wagneriana, *Rohrb. in Linnaea*, xxxvii. (1871–73) 281.—Ecuador.
Webbiana, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 230.—Reg. Himal.
Willkommii, Roem. ex Willk. & Lange, Prod. Fl. Hisp. iii. 617 = Arenaria intricata.

STELLARIOIDES, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 369 = **Anthericum**? Linn. (Liliac.).
canaliculata, Medic. l. c.—Hab.?

STELLARIS, Dill. ex Moench, Meth. 303 (1794) = **Scilla**, Linn. (Liliac.).
bifolia, Moench, l. c. 304 = Sc. bifolia.
corymbosa, Moench, l. c. = Ornithogalum umbellatum.
erecta, Moench, l. c. 303 = Ornithogalum pyrenaicum.
latifolia, Moench, l. c. = Ornithogalum latifolium.
minima, Moench, l. c. = Gagea pusilla.
parviflora, Steinh. in Ann. Sc. Nat. Sér. II. vi. (1836) 286 = Sc. numidica.
Scilla, Moench, Meth. 304 = Urginea Scilla.

STELLASTER, Heist. Syst. 5, 19 (1748) (Quid?).

STELLERA, J. G. Gmel. ex Linn. Diss. Dass. 16 (1747); Amoen. Acad. i. 399, pro parte (1749). *THYMELAEACEAE*, Benth. & Hook. f. iii. 193.
XAIASME, Rafin. Fl. Tellur. iv. 106 (1836).
altaica, Thieb. ex Pers. Syn. i. 436 = S. Chamaejasme.
annua, Salisb. Prod. 282 = Thymelaea arvensis.
Chamaejasme, *Linn. Sp. Pl.* 559.—As. bor. et temp.
concinna, Edgew. in Trans. Linn. Soc. xx. (1846) 88 = S. Chamaejasme.
dichotoma, Fisch. ex Sweet, Hort. Brit. ed. III. 592 = S. Chamaejasme.
Ganpi, Siebold, in Verh. Batav. Gen. xii. (1830) 22; Meissn. in DC. Prod. xiv. 550 = Wikstroemia canescens.
Griffithii, Meissn. l. c. = Lessertii.
Hedyotis, Ehrenb. ex Boiss. Fl. Orient. iv. 1052 = Thymelaea Aucheri.
hypericifolia, Endl. Gen. Suppl. iv. 11. 63 = S. Chamaejasme.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 22 = Wikstroemia japonica.
lathraeoides, Fisch. ex Ledeb. Fl. Ross. iii. 323 = Boschniakia glabra.
Lessertii, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Oriens.
Passerina, Linn. Sp. Pl. 559 = Thymelaea arvensis.
persica, Endl. Gen. Suppl. iv. 11. 63 = Lessertii.
pubescens, Guss. Fl. Sic. Prod. i. 466 = Thymelaea arvensis.
spicata, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 359 = Lessertii.
stachyoides, Schrenk, in Bull. Sc. Acad. Pétersb. x. (1842) 253; et Enum. Pl. Nov. ii. 16 = Lessertii.

STELLERA, Turcz. in Bull. Soc. Nat. Mosc. (1840) 167 = **Swertia**, Linn. (Gentian.).
cyanea, Turcz. l. c. 168 = Sw. tetrapetala.

STELLINA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 318 = **Callitriche**, Linn. (Halor.).
hamulata, Bubani, l. c. = C. hamulata.

STELLIX, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 = **Psychotria**, Linn. (Rubiac.).

STELLULARIA, Benth. in Hook. Ic. Pl. t. 1318 (1880). *SCROPHULARINEAE*.
nigricans, *Benth. l. c.*—Afr. trop.

STELLULARIA, Linn. Syst. ed. VI (1748) = **Stellaria**, Linn. (Caryoph.).

STELMANIS, Rafin. Fl. Tellur. ii. 47 (1836) = **Heterotheca**, Cass. (Compos.).
scabra, Rafin. l. c. = H. Lamarckii.

STELMATION, Fourn. in Mart. Fl. Bras. vi. IV. 226 (1885). *ASCLEPIADEAE.*
myrtifolium, *Fourn. l. c.* 227.—Bras.

STELMESUS, Rafin. Fl. Tellur. ii. 19 (1836) = **Allium**, Tourn. (Liliac.).
stellatus, Rafin. l. c. = A. stellatum.

STELMOTIS, Rafin. New Fl. Am. iv. 101 (1836) = **Oldenlandia**, Linn. (Rubiac.).
glomerata, Rafin. l. c. = O. glomerata.

STEMEIENA, Rafin. Atl. Journ. 144 (1832) = **Krameria**, Loefl. (Polygal.).

STEMMACANTHA, Cass. in Bull. Soc. Philom. (1817) 12 = **Serratula**, Linn. (Compos.).
cynaroides, Cass. in Dict. Sc. Nat. l. 460 = Cnicus cynaroides.
ficifolia, Turcz. ex DC. Prod. vi. 663 = Serratula atriplicifolia.
inermis, Dulac, Fl. Hautes-Pyr. 521 = Cnicus centauroides.

STEMMADENIA, Benth. Bot. Voy. Sulph. 124. t. 44 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 707.
ODONTOSTIGMA, A. Rich. Fl. Cub. Fanerog. 86. t. 56 (1853).
bella, *Miers, Apocyn. S. Am.* 77.—Mexic.
bignoniaeflora, *Miers, l. c.*—Mexic.
Galeottiana, *Miers, l. c.* 76.—Cuba.
glabra, *Benth. Bot. Voy. Sulph.* 124. t. 44.—Honduras.
grandiflora, Miers, Apocyn. S. Am. 75 = Tabernaemontana grandiflora.
guatemalensis, Muell. Arg. in Linnaea, xxx. (1859-60) 410 = Malouetia panamensis.
insignis, *Miers, Apocyn. S. Am.* 76.—Mexic.; Venezuela.
mollis, *Benth. Bot. Voy. Sulph.* 125.—Mexic.
pubescens, *Benth. l. c.*—Mexic.

STEMMATELLA, Wedd. ex Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 82, nomen. *COMPOSITAE*, Benth. & Hook. f. ii. 359.
congesta, *Wedd. l. c.*, nomen.—Bolivia.

STEMMATIUM, Phil. in Anal. Univ. Chil. (1873) 551 = **Tristagma**, Poepp. (Liliac.).
narcissoides, Phil. l. c. = T. narcissoides.

STEMMATOPHYSUM, Steud. Nom. ed. II. ii. 637 (1841), sphalm. = seq.

STEMMATOSIPHON, Meissn. Gen. 49 (1837) = seq.

STEMMATOSIPHUM, Pohl, Pl. Bras. Ic. ii. 86. tt. 157–159 (1831) = **Symplocos**, Linn. (Styrac.).
nitens, Pohl, l. c. 88. t. 158 = Symplocos nitens.
platyphyllum, Pohl, l. c. 87. t. 157 = Symplocos platyphylla.
uniflorum, Pohl, l. c. 89. t. 159 = Symplocos uniflora.

STEMMATOSPERMUM, Beauv. Agrost. 144. t. 25. f. 5 (1812) = **Nastus**, Juss. (Gramin.).
verticillatum, Beauv. l. c. = Nastus borbonicus.

STEMMODONTIA, Cass. in Bull. Soc. Philom. (1817) 11 = **Wedelia**, Jacq. (Compos.).
scaberrima, Cass. in Dict. Sc. Nat. xlvi. 407 = W. hispida.

STEMODIA, Linn. Syst. ed. X. 1118 (1759). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 950.
DICKIA, Scop. Introd. 199 (1777).
GOMPHIPUS, Rafin. Fl. Tellur. iv. 76 (1836).
MATOUREA, Aubl. Pl. Gui. ii. 641. t. 259 (1775).

STEMODIA:—
POARIUM, Desv. in Ham. Prod. Pl. Ind. Occ. 46 (1825).
UNANUEA, Ruiz & Pav. Ic. Fl. Per. ined.; ex Benth. in DC. Prod. x. 380 (1846).
ageratifolia, *Wright, in Sauv. Fl. Cub.* 99.—Cuba.
angulata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 22.—Am. centr.
aquatica, Willd. Sp. Pl. iii. 346 = Limnophila polystachya.
arenaria, H. B. & K. Nov. Gen. et Sp. ii. 357. t. 175 = parviflora.
arvensis, Steud. Nom. ed. II. ii. 637 = viscosa.
azurea, Linden, Cat. n. 17 (1862) 6 = Otacanthus caeruleus.
balsamea, Benth. in Wall. Cat. n. 3928 = Limnophila balsamea.
bartsioides, *Benth. in Bot. Reg.* sub t. 1470.—Mexic.
Berteroana, *Benth. in DC. Prod.* x. 384.—Ins. S. Doming.
camphorata, Benth. in Wall. Cat. n. 3932 = Limnophila diffusa.
camphorata, Vahl, Symb. Bot. ii. 69 = Adenosma camphoratum.
capitata, Benth. in Wall. Cat. n. 3926 = Adenosma capitatum.
chilensis, *Benth. in Bot. Reg.* t. 1470.—Chili.
cimicina, Benth. in Wall. Cat. n. 3933 = Limnophila diffusa.
coerulea, Benth. in DC. Prod. x. 381 = Adenosma caeruleum.
coromandelina, Heyne, ex Wall. Cat. n. 3929 D = viscosa.
debilis, *Benth. Fl. Austral.* iv. 487.—Austral.
diffusa, Benth. in Wall. Cat. n. 3937 = Limnophila pygmaea.
durantifolia, *Sw. Obs. Bot.* 240.—Ind. occ.
Ehrenbergiana, *Schlecht. in Bot. Zeit.* i. (1843) 169.—Mexic.
foliosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 46.—Bras.
glabra, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 21.—Mexic.
glabra, Spreng. Syst. ii. 811 = Gratiola peruviana.
glauca, Klein, ex Nees, in Hook. Comp. Bot. Mag. ii. (1836) 310 = Ebermaiera glauca.
grandiflora, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 89 = Lindenbergia grandiflora.
gratiolaefolia, A. St. Hil. Pl. Rem. Brés. 217 = palustris.
gratioloides, Benth. in Wall. Cat. n. 3938 = Limnophila erecta.
gratioloides, *F. Muell. Fragm.* x. 89.—As.; Afr.; Austral.
grossa, *Benth. Fl. Austral.* iv. 486.—Austral.
hirsuta, Heyne, in Wall. Cat. n. 3930 = Limnophila hirsuta.
hypericifolia, Benth. in Wall. Cat. n. 3934 = Limnophila hypericifolia.
hyptoides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 8.—Bras.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* ii. 358.—Mexic.
lanata, *Ruiz & Pav. ex Benth. in DC. Prod.* x. 383.—Mexic.
lanceolata, *Benth. l. c.* 384.—Bras.
latifolia, *Regel, Ind. Sem. Hort. Petrop.* (1861) 39.—Hab.?
linophylla, *F. Muell. Fragm.* x. 88.—Austral.
littoralis, Boiss. Fl. Orient. iv. 425 = Lindenbergia urticaefolia.
lobata, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 299.—Am. trop.
lobelioides, *Lehm. in Linnaea*, xi. (1837) *Litt.* 91.—Chili.
lutea, Moon, Cat. Pl. Ceyl. 45 = Adenosma camphoratum.
lythrifolia, *F. Muell. ex Benth. Fl. Austral.* iv. 486.—Austral.
macrostachya, Benth. in Wall. Cat. n. 3925 = Lindenbergia macrostachya.
maritima, Heyne, ex Wall. Cat. n. 3931 = viscosa.
maritima, *Linn. Syst.* ed. X. 1118.—Ind. occ.
Menthastrum, Benth. in Wall. Cat. n. 3927 = Limnophila Roxburghii.
micrantha, Benth. l. c. n. 3936 = Limnophila micrantha.
microphylla, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 298.—Bras.
minuta, G. Don, Gen. Syst. iv. 542 = Vandellia scabra.

STEMODIA:—
Morgania, *F. Muell. Fragm.* x. 89.—Austral.
multifida, Spreng. Syst. ii. 811 = Conobea multifida.
muraria, Roxb. ex D. Don, Prod. Fl. Nep. 89 = Lindenbergia urticaefolia.
odoratissima, F. Muell. ex Benth. Fl. Austral. iv. 485 = Adenosma Muelleri.
palustris, *A. St. Hil. Pl. Rem. Brés.* 216.—Bras.
parviflora, *Ait. Hort. Kew.* ed. II. iv. 52.—Am. trop.
pedicellaris, *F. Muell. Fragm.* viii. 231.—Austral.
peduncularis, *Benth. in DC. Prod.* x. 382.—Mexic.
philippensis, Cham. & Schlecht. in Linnaea, iii. (1828) 5 = Lindenbergia philippensis.
punctata, F. Muell. Census, 97 = Limnophila gratissima.
pusilla, *Benth. Bot. Voy. Sulph.* 144.—Mexic.
radicans, *Griseb. Cat. Pl. Cub.* 182.—Cuba.
repens, Benth. in Wall. Cat. n. 3935 = Limnophila conferta.
ruderalis, Blanco, Fl. Filip. ed. I. 498 = Lindenbergia philippensis.
ruderalis, *Retz. Obs.* v. 25.—Ind. or.
ruderalis, Vahl, Symb. Bot. ii. 69 = Lindenbergia urticaefolia.
senegalensis, *Desf. Cat. Hort. Par.* ed. III. 107.—Afr. trop.
serrata, *Benth. in DC. Prod.* x. 381.—Ind. or.; Afr. trop.
sessiliflora, F. Muell. Census, 97 = Limnophila sessiliflora.
sessilis, Benth. in Wall. Cat. n. 3939 = Limnophila conferta.
stricta, *Cham. & Schlecht. in Linnaea*, iii. (1828) 10.—Bras.
subhastata, *Benth. in DC. Prod.* x. 381.—Bras.
suffruticosa, *H. B. & K. Nov. Gen. et Sp.* ii. 357.—Am. austr.
surinamensis, *Miq. in Linnaea*, xxii. (1849) 475.—Guian.
tenuiflora, Benth. in Wall. Cat. n. 3940 = Limnophila conferta.
tomentosa, G. Don, Gen. Syst. iv. 542 = Mazus rugosus.
trifoliata, *Reichb. Ic. Exot.* i. 3. t. 1.—Bras.
veronicoides, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 298.—Bras.
verticillaris, Link, Enum. Hort. Berol. ii. 144 = durantifolia.
viscosa, *Roxb. Pl. Corom.* ii. 33. t. 163.—Ind. or.; Afghan.
xorullensis, Spreng. Syst. ii. 810 = jorullensis.

STEMODIACRA, P. Br. Hist. Jamaic. 261. t. 22. f. 2 (1756) = praec.

STEMODOXIS, Rafin. Fl. Tellur. ii. 20 (1836) = **Allium**, Tourn. (Liliac.).
juncea, Rafin. l. c. = A. junceum.

STEMONA, Lour. Fl. Cochinch. 404 (1790). *ROXBURGHIACEAE*, Benth. & Hook. f. iii. 747.
ROXBURGHIA, [Banks, in] Roxb. Pl. Corom. i. 29 (1795).
gloriosoides, Voigt, Hort. Suburb. Calc. 650 = tuberosa.
Griffithiana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 109. t. 10.—Burma.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 92.—Japon.
sessilifolia, *Franch. & Sav. l. c.*—Japon.
tuberosa, *Lour. Fl. Cochinch.* 404.—As. trop.

STEMONACANTHUS, Nees, in Mart. Fl. Bras. ix. 53. tt. 4, 205 (1847) = **Ruellia**, Plum. (Acanth.).
angustior, Nees, l. c. 54.—Bras.
coccineus, Griseb. Fl. Brit. W. Ind. 452 = R. coccinea.
formosus, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 270 = R. formosa.
fulgidus, Griseb. Fl. Brit. W. Ind. 452 = R. fulgida.
hirsutus, Nees, in Mart. Fl. Bras. ix. 53.—Bras.
Humboldtianus, Nees, in DC. Prod. xi. 206 = R. Humboldtiana.
macrophyllus, Nees, l. c. 205 = R. macrophylla.
multiflorus, Nees, in Mart. Fl. Bras. ix. 54.—Bras.
Pearcei, Hook. f. Bot. Mag. t. 5648.—Bolivia.
radicans, Nees, in DC. Prod. xi. 207.—Guian.
salviaefolius, Nees, in Mart. Fl. Bras. ix. 54.—Bras.
Sellowianus, Nees, l. c.—Bras.
thyrsacanthoides, Nees, in DC. Prod. xi. 207.—N. Granat.
Willdenovianus, Nees, l. c.—Hab.?

STEMONE, Franch. & Sav. Enum. Pl. Jap. ii. 92 (1876) = **Stemona**, Lour. (Roxburgh.).

STEMONIX, Rafin. Atl. Journ. 165 (1833) = **Eurycles**, Salisb. (Amaryll.).
nervosus, Rafin. l. c. = E. sylvestris.

STEMONOPORUS, Thw. in Hook. Kew Journ. vi. (1854) 67. t. 2 = **Vateria**, Linn. (Dipteroc.).
acuminatus, Bedd. Fl. Sylv. t. 100 = V. acuminata.
affinis, Thw. in Hook. Kew Journ. vi. (1854) 68 = V. affinis.
canaliculatus, Thw. Enum. Pl. Zeyl. 38 = V. canaliculata.
Gardneri, Thw. in Hook. Kew Journ. vi. (1854) 68 = V. Gardneri.
lanceolatus, Thw. l. c. = V. lanceolata.
macrophyllus, Thw. Enum. Pl. Zeyl. 37 = V. ceylanica.
Moonii, Thw. l. c. 39 = V. Moonii.
nitidus, Thw. l. c. = V. nitida.
oblongifolius, Thw. in Hook. Kew Journ. vi. (1854) 68 = V. oblongifolia.
petiolaris, Thw. Enum. Pl. Zeyl. 38 = V. petiolaris.
reticulatus, Thw. l. c. = V. reticulata.
rigidus, Thw. in Hook. Kew Journ. vi. (1854) 69 = V. rigida.
Wightii, Thw. Enum. Pl. Zeyl. 37 = V. ceylanica.

STEMONURUS, Blume, Bijdr. 648 (1825) = **Gomphandra**, Wall. et **Lasianthera**, Beauv. (Olac.).
affinis, Miers, l. c. 94 = G. affinis.
africanus, Miers, l. c. 87 = Lasianthera africana.
apicalis, Miers, in Trans. Linn. Soc. xxii. (1859) 110; Thw. Enum. Pl. Zeyl. 43 = Lasianthera apicalis.
axillaris, Miers, Contrib. i. 91 = G. axillaris.
capitatus, Becc. Malesia, i. (1877) 114.—Borneo.
ceylanicus, Miers, Contrib. i. 93 = G. axillaris.
coriaceus, Miers, l. c. 87 = G. polymorpha.
crassipes, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 298 = G. crassipes.
Cumingianus, Miers, Contrib. i. 83 = G. Cumingiana, *Villar.*
foetidus, Wight, Spicil. Neilgherr. i. 21. t. 23 = Mappia tomentosa.
frutescens, Blume, Bijdr. 649.—Java.
Gardneri, Miers, Contrib. i. 89 = G. polymorpha.
grandifolius, Becc. Malesia, i. (1877) 114.—Borneo.
Heyneanus, Miers, Contrib. i. 91 = G. axillaris.
javanicus, Blume, Bijdr. 649 = Lasianthera javanica.
javanicus, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 250 = G. affinis.
lanceolatus, Becc. Malesia, i. (1877) 114 = Lasianthera lanceolata.
laxiflorus, Miers, in Ann. & Mag. Nat. Hist. Ser. II. x. (1852) 111 = G. laxiflora.
litoralis, Blume, Mus. Bot. Lugd. Bat. i. 250 = Lasianthera littoralis.
longifolius, Miers, Contrib. i. 90 = G. axillaris.
macrocarpus, Blume, Mus. Bot. Lugd. Bat. i. 250 = Lasianthera macrocarpus.
macrophyllus, Blume, l. c. = Lasianthera macrophylla.
membranaceus, Blume, l. c. = Lesianthera membranacea.
ovalifolius, M. Roem. Syn. Hesper. 24 = Lasianthera ovalifolia.
parviflorus, Blume, Mus. Bot. Lugd. Bat. i. 250 = Lasianthera parviflora.
pauciflorus, Blume, Bijdr. 648.—Java.
penangianus, Miers, Contrib. i. 90 = G. penangiana.
polymorphus, Miers, l. c. 87 = G. polymorpha.
prasinus, Blume, Mus. Bot. Lugd. Bat. i. 249 = G. affinis.
punctatus, Becc. in Malesia, i. (1877) 116, in obs.—Zeylan.?
quadrifidus, Blume, Mus. Bot. Lugd. Bat. i. 249 = G. penangiana.
scorpioides, Becc. Malesia, i. (1877) 113.—Borneo.
secundiflorus, Blume, Bijdr. 649 = Lasianthera apicalis.
tomentellus, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 298 = G. tomentella.
umbellatus, Becc. Malesia, i. (1877) 115.—Borneo.
vitiensis, Seem. Fl. Vit. 39.—Ins. Fiji.
Walkeri, Miers, Contrib. i. 93 = G. polymorpha.

STEMOPTERA, Miers, in Proc. Linn. Soc. i. (1840) 62 = **Apteria**, Nutt. (Burmann.).
lilacina, Miers, l. c. = A. setacea?

STEMOTIS, Rafin. Sylva Tellur. 152 (1838) = **Rhododendron**, Linn. (Ericac.).
album, Rafin. l. c. 153 = R. album.
coccineum, Rafin. l. c. = R. arboreum.
roseum, Rafin. l. c. = R. arboreum.

STENACHAENIUM, Benth. in Benth. & Hook. f. Gen. ii. 289 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
campestre, *Baker, in Journ. Bot.* xvi. (1878) 79.—Bras.
macrocephalum, *Benth. ex Benth. & Hook. f. Gen.* ii. 289.—Bras.
macrocephalum, Griseb. in Goett. Abh. xxiv. (1879) 183 = campestre.
megapotamicum, *Baker, in Mart. Fl. Bras.* vi. III. 105.—Bras.
Riedelii, *Baker, in Journ. Bot.* xvi. (1878) 78.—Bras.

STENACTIS, Cass. in Dict. Sc. Nat. xxxvii. 462 (1825) = **Erigeron**, Linn. (Compos.).
alpina, Cass. l. c. 485; l. 485 = E. alpinus.
ambigua, DC. Prod. v. 299 = E. strigosus.
annua, Cass. in Dict. Sc. Nat. xxxvii. 485 = E. annuus.
bellidiflora, A. Br. Ind. Sem. Hort. Berol. (1859) 10 (Quid?).—Helvet.
bellidioides, Nees, Gen. et Sp. Ast. 274 = E. bellidioides, *Benth. & Hook. f.*
Beyrichii, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. n. 2169; ex iisd. vi. 63, in obs. = E. strigosus.
delphinifolia, Cass. in Dict. Sc. Nat. xxxvii. 485; l. 483 = E. delphinifolius.
dubia, Cass. xxxvii. l. c. 485 = E. annuus.
glauca, Nees, Gen. et Sp. Ast. 275 = E. glaucus.
gnaphalioides, Cass. in Dict. Sc. Nat. l. 484 = E. gnaphalioides.
multiradiata, Lindl. in DC. Prod. v. 299 = E. multiradiatus.
pilosa, DC. l. c. = E. pilosus.
pubescens, Cass. in Dict. Sc. Nat. l. 484 = E. pubescens.
speciosa, Lindl. Bot. Reg. t. 1577 = E. speciosus.
strigosa, DC. Prod. v. 299 = E. strigosus.
verna, Nees, Gen. et Sp. Ast. 275 = E. nudicaulis.

STENANDRIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836). *ACANTHACEAE*, Benth. & Hook. f. ii. 1095.
CALDENBACHIA, Pohl, ex Nees, in DC. Prod. xi. 281 (1847).
HEMITOME, Nees, l. c. 295, in obs. (1847).
SYNANDRA, Schrad. in Goett. Gel. Anz. i. (1821) 715.
aurantiacum, *Nees, in Mart. Fl. Bras.* ix. 78.—Bras.
barbatum, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 168. t. 4.—Mexic.
chameranthemoideum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 139.—Mexic.
corymbosum, *Nees, in DC. Prod.* xi. 727.—N. Granat.
decoratum, *Nees, in Mart. Fl. Bras.* ix. 79.—Bras.
diphyllum, Nees, l. c. 75.—Bras.
droseroides, *Nees, in DC. Prod.* xi. 284.—Cuba.
dulce, *Nees, l. c.* 282.—Chili.
elegans, *Nees, l. c.* 283.—Venezuela.
hirsutum, *Nees, in Mart. Fl. Bras.* ix. 77.—Bras.
Humboldtianum, *Nees, in DC. Prod.* xi. 284.—Am. austr.
igneum, *André, in Illustr. Hortic.* xxiv. (1877) t. 266.—Peruv.
igneum, *Nees, in Mart. Fl. Bras.* ix. 78.—Bras.
Macraganum, Nees, in DC. Prod. xi. 282, sphalm. = dulce.
Macrayanum, *Nees, in Mart. Fl. Bras.* ix. 75 (an praec.?).—Chili; Peruv.
mandioccanum, *Nees, in Mart. Fl. Bras.* ix. 76.—Bras.
obtusifolium, *Nees, l. c.* 78.—Bras.
ornatum, Nees, in DC. Prod. xi. 291 = Aphelandra ornata.
Pohlii, *Nees, in Mart. Fl. Bras.* ix. 75.—Bras.
punctatum, *Griseb. Cat. Pl. Cub.* 196.—Ind. occ.
radicosum, *Nees, in DC. Prod.* xi. 283.—N. Granat.
Riedelianum, *Nees, in Mart. Fl. Bras.* ix. 76.—Bras.
rupestre, Nees, in DC. Prod. xi. 283.—Ind. occ.
scabrosum, *Nees, l. c.* 284.—Ins. S. Doming.
serpens, *Nees, in Mart. Fl. Bras.* ix. 77.—Bras.
speciosum, Nees, ex Mart. in Flora, xxiv. (1841) II. Beibl. 24 = Geissomeria longiflora.
tenellum, *Nees, in Mart. Fl. Bras.* ix. 77.—Bras.

STENANDRIUM:—
trinerve, *Nees, in Mart. Fl. Bras.* ix. 75.—Bras.
venatum, *Nees, l. c.* 79.—Bras.
villosum, *Nees, l. c.* 77.—Bras.

STENANTHEMUM, Reiss. in Linnaea, xxix. (1857–58) 295. *RHAMNEAE*, Benth. & Hook. f. i. 382.
coronatum, *Reiss. l. c.*—Austral.
humile, *Benth. Fl. Austral.* i. 436.—Austral.
leucophractum, *Reiss. in Linnaea*, xxix. (1857–58) 295.—Austral.
pimeleoides, *Benth. Fl. Austral.* i. 436.—Tasman.
pomaderroides, *Reiss. in Linnaea*, xxix. (1857–58) 295.—Austral.
tridentatum, Reiss. l. c. = Spyridium tridentatum.
Waterhousii, *Benth. Fl. Austral.* i. 436.—Austral.

STENANTHERA, R. Br. Prod. 538 (1810) = **Astroloma**, R. Br. (Epacrid.).
brachyloma, F. Muell. Fragm. iv. 99 = Brachyloma concolor.
ciliata, Lindl. Swan River App. 25 = A. longiflorum.
conostephioides, Sond. in Lehm. Pl. Preiss. i. 296 = A. conostephioides.
ericoides, F. Muell. Fragm. iv. 98 = Brachyloma ericoides.
pinifolia, R. Br. Prod. 538 = A. pinifolium.
squamuligera, F. Muell. Fragm. iv. 97 = A. Baxteri.

STENANTHIUM, Kunth, Enum. Pl. iv. 189 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 835.
angustifolium, *Kunth, l. c.* 190.—Am. bor.
frigidum, *Kunth, l. c.*—Mexic.
gramineum, Kunth, l. c. = angustifolium.
occidentale, *A. Gray, in Proc. Am. Acad.* viii. (1872) 405.—Am. bor. occ.
robustum, *S. Wats. l. c.* xiv. (1879) 278.—Am. bor.
sachalinense, *F. Schmidt, Reisen Amurl.* 188.—Ins. Sachalin.

STENANTHUS, Oerst. ex Hanst. in Linnaea, xxvi. (1853) 209 = **Columnea**, Plum. (Gesner.).
heterophyllus, Oerst. ex Hanst. l. c. = C. heterophylla.
sanguinolentus, Klotzsch, ex Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 123 = C. sanguinolenta.
serratus, Klotzsch, ex Oerst. l. c. = C. serrata.
squarrosus, Klotzsch, ex Oerst. l. c. = C. sanguinolenta.

STENAPHIA, A. Rich. Tent. Fl. Abyss. i. 9 (1847), sphalm. = **Stephania**, Lour. (Menisp.).

STENARIA, Rafin. ex Steud. Nom. ed. II. i. 776, in syn. (1840) = **Houstonia**, Gronov. (Rubiac.).

STENARRHENA, D. Don, Prod. Fl. Nep. 111 (1825) = **Salvia**, Tourn. (Labiat.).
lanata, D. Don, l. c. = Salvia lanata.

STENGELIA, Neck. Elem. ii. 258 (1790) = **Mourera**, Aubl. (Podost.).

STENGELIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26 = **Vernonia**, Schreb. (Compos.).
adoënsis, Sch. Bip. l. c. nomen = V. adoënsis.
Calvoana, Hook. f. in Journ. Linn. Soc. vii. (1864) 199 = V. Calvoana.
controversa, Sch. Bip. ex Walp. Rep. ii. 947 = V. abyssinica.
insignis, Hook. f. in Journ. Linn. Soc. vii. (1864) 199 = V. insignis.
Kotschyana, Hochst. in Flora, xxv. (1842) I. Beibl. 134, nomen = V. Kotschyana.
Schimperi, Steetz, in Peters, Reise Mossamb. Bot. 360 = V. Schimperi.

STENIA, Lindl. Bot. Reg. sub t. 1991 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 553.
STENOPOLEN, Rafin. Fl. Tellur. iv. 49 (1836), nomen prius.
Beaumontii, A. Rich. Cat. Hort. Pescat. ex Richb. f. Otia Bot. Hamb. iii. 37 = Batemannia Beaumontii.
fimbriata, Linden & Reichb. f. in Gard. Chron. (1868) 1313 = Chondrorhyncha fimbriata.
guttata, *Reichb. f. l. c.* (1880) II. 134.—Peruv.
pallida, *Lindl. Bot. Reg.* sub t. 1991; (1838) t. 20.—Guian.

STENOCAELIUM, Ledeb. & Hook. f. Gen. i. 902 (1867), sphalm = STENOCOELIUM, Ledeb. = **Seseli**, Linn. (Umbell.).

STENOCALYX, Berg, in Linnaea, xxvii. (1854) 309 = **Eugenia**, Mich. (Myrtac.).
affinis, Berg, in Mart. Fl. Bras. xiv. I. 336.—Bras.
alagoënsis, Berg, l. c. 350.—Bras.
albicans, Berg, in Linnaea, xxx. (1859-60) 698 = Calyptranthes cuprea ?
bahiensis, Berg, in Mart. Fl. Bras. xiv. I. 351.—Bras.
blastanthus, Berg, l. c. 348.—Bras.
brasiliensis, Berg, l. c. 347 = E. brasiliensis.
brunneus, Berg, l. c. 335.—Bras.
costatus, Berg, l. c. 340 = E. costata, *Cambess.*
dasyblastus, Berg, l. c. 337.—Bras.
dentatus, Berg, l. c. 338.—Bras.
Duarteanus, Berg, l. c. 342 = E. Duarteana.
dysentericus, Berg, l. c. 351 = E. dysenterica.
Gardnerianus, Berg, l. c. 350.—Bras.
gemmiflorus, Berg, l. c. 344.—Bras.
glaber, Berg, l. c. 337.—Bras.
grandiflorus, Berg, l. c. 347.—Bras.
grandifolius, Berg, l. c. 340.—Bras.
impressus, Berg, l. c. 593.—Bras.
impunctatus, Berg, l. c. 336 = E. multiflora.
lanceolatus, Berg, l. c. 341 = E. Uvalha.
Langsdorffii, Berg, l. c. 594 = E. linearifolia ?
laxus, Berg, l. c. 349 = E. laxa.
ligustrinus, Berg, in Linnaea, xxvii. (1854) 312 = E. ligustrina, *Willd.*
longipes, Berg, in Mart. Fl. Bras. xiv. I. 345.—Bras.
lucidus, Berg, l. c. 337.—Bras.
Michelii, Berg, l. c. = E. uniflora, *Linn.*
mollicoma, Berg, l. c. 350 = E. mollicoma.
mutabilis, Berg, l. c. 347.—Bras.
myrtifolius, Berg, l. c. 342 = E. myrtifolia, *Cambess.*
oblongifolius, Berg, l. c. 339.—Bras.
Patrisii, Berg, in Linnaea, xxix. (1857-58) 247 = E. Patrisii.
pistaciaefolius, Berg, in Mart. Fl. Bras. xiv. I. 343 = E. pistaciaefolia.
Pitanga, Berg, l. c. 341.—Bras.
pius, Berg, l. c. 342 = E. pia.
Plumieri, Berg, in Linnaea, xxx. (1859-60) 698.—Ind. occ.
portoricensis, Berg, l. c. xxix. (1857-58) 246 = E. portoricensis.
protractus, Berg, in Mart. Fl. Bras. xiv. I. 593.—Bras.
Pseudo-psidium, Berg, in Linnaea, xxvii. (1854) 314 = E. Pseudo-psidium.
reticulatus, Berg, in Mart. Fl. Bras. xiv. I. 340 = E. ligustrina, *Miq.*
Riedelianus, Berg, l. c. 349.—Bras.
sessilifolius, Berg, l. c. 344 = E. sessilifolia.
silvestris, Berg, l. c. 346 = E. brasiliensis.
squamiflorus, Berg, l. c. 345.—Bras.
stipularis, Berg, l. c. 348.—Bras.
strigosus, Berg, l. c. 335.—Bras.
sulcatus, Berg, l. c. 339 = E. sulcata.
ubensis, Berg, l. c. 344 = E. ubensis.
xanthoxyloides, Berg, l. c. 346 = E. xanthoxyloides.

STENOCALYX, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 393. *MALPIGHIACEAE.*
involutus, *Turcz. l. c.* 394.—N. Granat.

STENOCARPUS, R. Br. in Trans. Linn. Soc. x. (1810) 201. *PROTEACEAE*, Benth. & Hook. f. iii. 182.
AGNOSTUS, A. Cunn. in Loud. Hort. Brit. 580 (1830).
CYBELE, Salisb. in Knight, Prot. 123 (1809), nomen prius.
acacioides, F. Muell. Fragm. i. 135 = salignus.
concolor, F. Muell. l. c. iii. 147 = salignus.
Cunninghami, *R. Br. Prot. Nov.* 34.—Austral.
Cunninghami, Hook. Bot. Mag. t. 4263 = sinuatus.
dareoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 45; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 208.—N. Caled.
elegans, *Brongn. & Gris, ll. cc.* 44; 207.—N. Caled.
Forsteri, *R. Br. in Trans. Linn. Soc.* x. (1810) 201.—N. Caled.
gracilis, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 43; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 205.—N. Caled.
heterophyllus, *A. Brongn. & Gris, ll. cc.* 44; 207.—N. Caled.
intermedius, *A. Brongn. & Gris, ll. cc.* 42; 204.—N. Caled.
latifolius, *Montr. in Mém. Acad. Lyon*, x. (1860) 249.—N. Caled.

STENOCARPUS:—
laurifolius, Panch. & Sebert, Not. Bois Nouv. Caléd. 181 = laurinus.
laurinus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 43; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 205.—N. Caled.
Milnei, *Hook. Kew Journ.* vi. (1854) 359.—N. Caled.
Moorei, F. Muell. Fragm. i. 134 = salignus.
rubiginosus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 43; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 206.—N. Caled.
salignus, *R. Br. in Trans. Linn. Soc.* x. (1810) 202.—Austral.
sinuatus, *Endl. Gen. Suppl.* iv. II. 88.—Austral.
sinuosus, F. Muell. Fragm. vi. 224 = praec. ?
tremuloides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 44; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 207.—N. Caled.
villosus, *Brongn. & Gris, ll. cc.* 44; 206.—N. Caled.

STENOCEPHALUM, Sch. Bip. in Pollichia, xx-xxi. (1863) 385 = **Vernonia**, Schreb. (Compos.).
apiculatum, Sch. Bip. l. c. 387 = V. apiculata.
brevifolium, Sch. Bip. l. c. = V. megapotamica.
hexanthum, Sch. Bip. l. c. 390 = V. hexantha.
megapotamicum, Sch. Bip. l. c. 388 = V. megapotamica.
melanotrichium, Sch. Bip. l. c. = V. megapotamica.
monticolum, Sch. Bip. l. c. 386 = V. monticola.
penicillatum, Sch. Bip. l. c. 389 = V. megapotamica.
targiaefolium, Sch. Bip. l. c. = V. megapotamica.

STENOCHASMA, Griff. Notul. iii. 431 (1851) = **Amomum**, Linn. (Scitam.).
convoluta, Griff. l. c. 433 = A. Leonurus.
urceolare, Griff. l. c. 431 = A. scyphiferum.

STENOCHASMA, Miq. Pl. Jungh. 45 (1850) = **Broussonetia**, L'Hérit. (Urtic.).
ancolanum, Miq. l. c.—Sumatra.

STENOCHILUS, R. Br. Prod. 517 (1810) = **Eremophila**, R. Br. (Myopor.).
albicans, Bartl. in Lehm. Pl. Preiss. i. 351 = E. Brownii.
bignoniaeflorus, Benth. in Mitch. Journ. Trop. Austral. 386 = E. bignoniflora.
curvipes, Benth. l. c. 221 = E. maculata.
glaber, R. Br. Prod. 517 = E. Brownii.
incanus, Lindl. in Bot. Reg. (1839) Misc. 70 = E. Brownii.
longifolius, R. Br. Prod. 517 = E. longifolia.
longifolius, A. Cunn. ex Lindl. Bot. Reg. (1839) Misc. 69 (an praec.?).—Austral.
maculatus, Ker-Gawl. in Bot. Reg. t. 647 = E. maculata.
ochroleucus, A. Cunn. ex A. DC. in DC. Prod. xi. 715 = E. Brownii.
pubiflorus, Benth. in Mitch. Journ. Trop. Austral. 273 = E. longifolia.
racemosus, Endl. in Ann. Wien. Mus. ii. (1838) 202 = E. maculata.
salicinus, Benth. in Mitch. Journ. Trop. Austral. 251 = E. longifolia.
serrulatus, DC. Prod. xi. 715 = E. latifolia.
subcanescens, Bartl. in Lehm. Pl. Preiss. i. 351 = E. Brownii.
viscosus, R. Grah. in Edinb. N. Phil. Journ. (Jan.-Mar. 1829) 387 = E. Brownii.

STENOCHLOA, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 189 = **Dissanthelium**, Trin. (Gramin.).
californica, Nutt. l. c. = D. californica.

STENOCLINE, DC. Prod. vi. 218 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 312.
bracteifera, *DC. l. c.*—Madag.
chionaea, *DC. l. c.* 219.—Bras.
ericoides, *DC. l. c.*—Madag.
ferruginea, *Baker, in Journ. Linn. Soc.* xx. (1883) 187.—Madag.
filaginoides, *DC. Prod.* vi. 219.—Madag.
fruticosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 186.—Madag.
Gardneri, Baker, in Mart. Fl. Bras. vi. III. 127 = satureoides.

STENOPTERA:—
gymnocephala, *DC. Prod.* vi. 218.—Madag.
incana, *Baker, in Journ. Linn. Soc.* xx. (1883) 187.—Madag.
inuloides, *DC. in Deless. Ic. Sel.* iv. t. 53.—Madag.
lecheoides, *DC. Prod.* vi. 218.—Madag.
pusilla, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 378.—Ins. Maurit.
satureoides, *Benth. & Hook. f. Gen.* ii. 312.—Bras.

STENOCOELIUM, Ledeb. Fl. Alt. i. 298 (1829) = **Seseli**, Linn. (Umbell.).
athamantoides, Ledeb. l. c.—Sibir. altaic.
divaricatum, Turcz. in Bull. Soc. Nat. Mosc. (1838) 93; xvii. (1844) 734 = Siler divaricatum.
trichocarpum, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 80.—Soongar.

STENOCORYNE, Lindl. Bot. Reg. (1843) Misc. 53 = **Bifrenaria**, Lindl. (Orchid.).
longicornis, Lindl. l. c. = B. longicornis.

STENODISCUS, Reiss. in Linnaea, xxix. (1857-58) 295 = **Spyridium**, Fenzl (Rhamn.).
ulicinus, Reiss. l. c. = Spyr. ulicinum.

STENODON, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 146. *MELASTOMACEAE*, Benth. & Hook. f. i. 737.
gracilis, *Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 25.—Bras.
suberosus, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 146. t. 3.—Bras.

STENOGASTRA, Hanst. in Linnaea, xxvi. (1853-55) 205 = **Sinningia**, Nees (Gesner.).
concinna, Hook. f. Bot. Mag. t. 5253.—Bras.
hirsuta, Hanst. in Linnaea, xxvi. (1853-55) 205.—Bras.
multiflora ×, Lem. Illustr. Hortic. (1864) t. 390.

STENOGLOSSUM, H. B. & K. Nov. Gen. et Sp. i. 355. t. 87 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 523.
coriophorum, Lindl. Gen. et Sp. Orch. 62 = seq.
corynephorum, Spreng. Syst. iii. 742 = seq.
coryophorum, *H. B. & K. Nov. Gen. et Sp.* i. 356.—N. Granat.
subulatum, Lindl. in Hook. Ic. Pl. t. 51 = coryophorum.

STENOGLOTTIS, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209. *ORCHIDEAE*, Benth. & Hook. f. iii. 622.
calcarata, Reichb. f. in Flora, xlviii. (1865) 180 = Habenaria calcarata (in Addend.).
fimbriata, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 210.—Afr. austr.

STENOGONUM, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 170 = **Eriogonum**, Michx. (Polygon.).
salsuginosum, Nutt. l. c. = E. salsuginosum.

STENOGYNE, Benth. in Bot. Reg. sub t. 1292 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1217.
PHAEOPSIS, Nutt. ex Benth. in DC. Prod. xii. 555 (1848).
angustifolia, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 348.—Ins. Sandvic.
calaminthoides, *A. Gray, l. c.* 347.—Ins. Sandvic.
cordata, *Benth. Lab. Gen. et Sp.* 654.—Ins. Sandvic.
crenata, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 348.—Ins. Sandvic.
diffusa, A. Gray, l. c. = microphylla.
Haliakalae, Wawra, in Flora, lv. (1872) 554 = rotundifolia.
Kaaleae, *Wawra, in Flora*, lv. (1872) 556.—Ins. Sandvic.
Kamehamehae, Wawra, l. c. 532 = calaminthoides.
Kealiae, Wawra, l. c. 556 = purpurea.
macrantha, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
microphylla, *Benth. Lab. Gen. et Sp.* 655.—Ins. Sandvic.
Nelsonii, Benth. Lab. Gen. et Sp. 655 = scrophularioides.
parviflora, *H. Mann, in Proc. Am. Acad.* vii. (1867) 193.—Ins. Sandvic.
purpurea, *H. Mann, l. c.*—Ins. Sandvic.
rotundifolia, *A. Gray, in Proc. Am. Acad.* v. (1861-62) 347.—Ins. Sandvic.

STENOGYNE:—
rugosa, *Benth. in Bot. Reg.* sub t. 1292.—Ins. Sandvic.
scrophularioides, *Benth. l. c.*—Ins. Sandvic.
sessilis, *Benth. Lab. Gen. et Sp.* 654.—Ins. Sandvic.

STENOGYNE, Cass. in Dict. Sc. Nat. l. 493 (1827) = **Eriocephalus**, Linn. (Compos.).

STENOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 125 = **Calopogonium**, Desv. (Legumin.).
arboreum, Wawra, in Flora, xlvii. (1864) 243.—Bras.
brachycarpum, Benth. in Seem. Bot. Voy. Herald, 109 = C. mucunoides.
caeruleum, Benth. in Ann. Wien. Mus. ii. (1838) 125 = C. caeruleum.
galactioides, Benth. in Mart. Fl. Bras. xv. 1. 140 = C. galactioides.
glabrum, Benth. in Ann. Wien. Mus. ii. (1838) 125 = C. caeruleum.
tomentosum, Benth. l. c. = C. caeruleum.
velutinum, Benth. in Ann. Nat. Hist. Ser. I. iii. (1839) 437 = Rhynchosia Luschnathiana.

STENOLOBIUM, D. Don, in Edinb. Phil. Journ. ix. (1823) 264 = **Tecoma**, Juss. (Bignon.).
castanifolium, D. Don, l. c. = Bignonia castaneaefolia.
molle, Seem. Journ. Bot. i. (1863) 90 = T. mollis.
sambucifolium, Seem. l. c. 91 = T. stans.
stans, Seem. l. c. 88 = T. stans.

STENOLOPHUS, Cass. in Dict. Sc. Nat. xliv. 35 (1826); l. 499 (1827) = **Centaurea**, Linn. (Compos.).

STENOMERIA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 312. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 756.
decalepis, *Turcz. l. c.*—N. Granat.
pentalepis, *Turcz. l. c.* 313.—N. Granat.

STENOMERIS, Planch. in Ann. Sc. Nat. Sér. III. xviii. (1852) 319. *DIOSCOREACEAE*, Benth. & Hook. f. iii. 745.
Cumingiana, *Becc. in Nuov. Giorn. Bot. Ital.* ii. (1870) 8.—Ins. Philipp.
dioscoreaefolia, *Planch. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 320.—Ins. Philipp.

STENOMESSON, Herb. App. [Bot. Reg.] 40 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 733.
CALLITHAUMA, Herb. Amaryll. 225 (1837).
CARPODETES, Herb. App. [Bot. Reg.] 41 (1821).
CHRYSIPHIALA, [Ker-Gawl. in] Bot. Reg. sub t. 778 (1824).
CLINANTHUS, Herb. App. [Bot. Reg.] 40 (1821).
CLITANTHES, Herb. in Bot. Reg. xxv. (1839) Misc. 87.
COBURGIA, Sweet, Brit. Flow. Gard. Ser. II. t. 17 (1838).
NEAERA, Salisb. Gen. Fl. Fragm. 109 (1866).
SPHAEROTELE, Presl, Rel. Haenk. 119. t. 16 (1830).
acutum, Baker, in Saund. Refug. Bot. sub t. 308 = incarnatum.
aurantiacum, *Herb. App.* [*Bot. Reg.*] 40.—Ecuador.
breviflorum, Herb. Amaryll. 199. t. 28. f. 7, 8 = coccineum.
coccineum, *Herb. App.* [*Bot. Reg.*] 40.—Peruv.
croceum, *Herb. Amaryll.* 199. t. 28. f. 4.—Peruv.
curvidentatum, Herb. in Bot. Mag. t. 2640 = flavum.
discolor, Baker, in Saund. Refug. Bot. sub t. 308 = recurvatum.
eustephioides, Herb. in Bot. Reg. (1843) Misc. 62 = aurantiacum.
flavum, *Herb. App.* [*Bot. Reg.*] 40.—Peruv.
Hartwegii, Lindl. Bot. Reg. (1844) t. 42 = aurantiacum.
humilis, *Baker, in Saund. Refug. Bot.* sub t. 308.—Peruv.
incarnatum, *Baker, l. c.*—Peruv.
latifolium, Herb. in Bot. Mag. t. 3803 = flavum.
luteoviride, *Baker, l. c.* t. 6508.—Ecuador.
luteum, Baker, in Saund. Refug. Bot. sub t. 308 = recurvatum.
obragillensis, Baker, l. c. = recurvatum.
pauciflorum, Sweet, Hort. Brit. ed. I. 406; Herb. Amaryll. 198. t. 28. f. 3 = flavum.
Pearcei, *Baker, in Saund. Refug. Bot.* t. 308.—Peruv.
recurvatum, *Baker, l. c.*—Peruv.
rubrum, *Herb. Amaryll.* 199. t. 28. f. 6.—Peruv.
Ruizianum, Kunth, Enum. Pl. v. 642 = croceum.

STENOMESSON:—
Stricklandi, *Baker, in Gard. Chron.* (1882) II. 102.—Ecuador.
suspensum, *Baker, in Saund. Refug. Bot.* t. 22.—Peruv.
viridiflorum, *Benth. & Hook. f. Gen.* iii. 733.—Peruv.
vitellinum, Lindl. Bot. Reg. (1843) t. 2 = flavum.

STENONEMA, Hook. ex Benth. & Hook. f. Gen. i. 75 (1862) = **Dolichostylis**, Turcz. (Crucif.).

STENONIA, Baill. Etud. Gén. Euph. 578. t. 22 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 267.
Boiviniana, *Baill. l. c.* 579. t. 22.—Afr. trop.

STENONIA, Didr. in Kjoeb. Vidensk. Meddel. (1857) 146 = **Argithamnia**, Sw. (Euphorb.).
montevidensis, Didr. l. c. 148 = A. montevidensis.

STENOPETALUM, R. Br. ex DC. Syst. ii. 513 (1821); et in Mém. Mus. Par. vii. (1821) 239. *CRUCIFERAE*, Benth. & Hook. f. i. 82.
australis, C. Muell. in Walp. Ann. vii. 146, sphalm. = Menkea draboides.
croceum, Bunge, in Lehm. Pl. Preiss. i. 258 = robustum.
draboides, Hook. Ic. Pl. t. 617 = Menkea draboides.
filifolium, *Benth. Fl. Austral.* i. 78.—Austral.
gracile, Bunge, in Lehm. Pl. Preiss. i. 257 = robustum.
incisifolium, Hook. f. in Hook. Ic. Pl. t. 276 = Capsella procumbens.
lineare, *R. Br. ex DC. Syst.* ii. 513.—Austral.
minus, Bunge, in Lehm. Pl. Preiss. i. 258 = robustum.
nutans, *F. Muell. Fragm.* iii. 27.—Austral.
pedicellare, *F. Muell. ex Benth. Fl. Austral.* i. 79.—Austral.
procumbens, Hook. Ic. Pl. t. 610 = Menkea australis.
robustum, *Endl. in Enum. Pl. Hueg.* 4.—Austral.
sphaerocarpum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 35.—Austral.
velutinum, *F. Muell. Vict. Pl.* i. 49.—Austral.

STENOPHRAGMA, Celak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 445. *CRUCIFERAE*.
Thalianum, *Čelak. l. c.*—Hispan.

STENOPHYLLUM, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 391 (1873) = **Calea**, Linn. (Compos.).
senecioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 258 = C. senecioides.
Senecionis, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 391 = C. senecioides.

STENOPHYLLUS, Rafin. Neogenyt. 4 (1825) = **Scirpus**, Tourn. (Cyper.).
caespitosus, Rafin. l. c. (= *Isolepis stenophylla*).—Am. bor.

STENOPOLEN, Rafin. Fl. Tellur. iv. 49 (1836) = **Stenia**, Lindl. (1837) (Orchid.).

STENOPTERA, Presl, Rel. Haenk. i. 95. t. 14 (1827). *ORCHIDEAE*, Benth. & Hook. f. iii. 595.
PORPHYROSTACHYS, Reichb. f. Xenia Orch. i. 18 (1854).
acuta, *Lindl. Gen. et Sp. Orch.* 447.—Bras.
ananassacomos, *Reichb. f. in Bot. Zeit.* xxi. (1863) 131.—Jamaic.
cardinalis, *Lindl. Gen. et Sp. Orch.* 448.—Peruv.
goodyeroides, *Reichb. f. in Bonplandia*, iv. (1856) 211.—Peruv.
macrostachya, Reichb. f. in Bonplandia, iv. (1856) 211 = peruviana.
peruviana, *Presl, Rel. Haenk.* i. 95.—Peruv.
viscosa, *Reichb. f. in Bonplandia*, iv. (1856) 211.—Peruv.

STENORHYNCHUS, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 386 = seq.

STENORHYNCUS, Lindl. in Benth. Pl. Hartw. 92 (1842) = seq.

STENORRHYNCHIUM, Reichb. Nom. 55 (1841) = seq.

STENORRHYNCHOS, Spreng. Gen. ii. 660 (1831) = seq.

STENORRHYNCHUS, Reichb. Consp. 67 (1828) = seq.

STENORYNCHUS, Rich. in Mém. Mus. Par. iv. (1818) 59 = **Spiranthes**, Rich. (Orchid.).
actinophila, Rodr. Orch. Nov. ii. p. xv (Index) = Sp. actinophila.
albescentis, Rodr. l. c. i. p. x (Index) = Sp. micrantha.
aphyllus, Lindl. Gen. et Sp. Orch. 478 = Sp. orchioides.
argenteus, Griseb. in Goett. Abh. xxiv. (1879) 339.—Reg. Argent.
Arrabidae, Reichb. f. in Linnaea, xxii. (1849) 815 = Sp. Arrabidae.
aurantiacus, Lindl. Gen. et Sp. Orch. 479 = Sp. aurantiaca.
australis, Lindl. l. c. 477 = Sp. orchioides?
calcaratus, Rich. in Mém. Mus. Par. iv. (1818) 59 = Pelexia setacea.
ceracifolia, Rodr. Orch. Nov. ii. p. xv (Index) = Sp. ceracifolia.
cernuus, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 386.—Ecuador.
cinnabarinus, Lindl. Gen. et Sp. Orch. 479 = Sp. cinnabarina.
epiphyta, Rodr. Orch. Nov. i. p. x (Index) = Sp. epiphyta.
euphlebium, Reichb. f. in Regel, Gartenfl. (1853) 3, in obs. = Sp. euphlebia.
flavus, Spreng. Syst. iii. 710 = Corymbis flava.
hypnophila, Rodr. Orch. Nov. i. p. x (Index) = Sp. hypnophila.
hysterantha, Rodr. l. c. = Sp. hysterantha.
icmadophila, Rodr. l. c. ii. p. xv (Index) = Sp. icmadophila.
lanceolatus, Rich. in Mém. Mus. Par. iv. (1818) 59 = Sp. orchioides?
laxus, Poepp. & Endl. Nov. Gen. et Sp. ii. 7. t. 109 = Pelexia laxa.
lupulinus, Lindl. Gen. et Sp. Orch. 479 = Sp. lupulina.
macropoda, Rodr. Orch. Nov. i. p. x (Index) = Sp. macropoda.
madrensis, Reichb. f. in Bonplandia, iii. (1855) 177 = Sp. madrensis.
michuacanus, Lindl. Gen. et Sp. Orch. 480 = Sp. mechoacana.
montanus, Lindl. in Benth. Pl. Hartw. 95 = Sp. montana.
novofriburgensis, Reichb. f. in Linnaea, xxii. (1849) 815 = Sp. novofriburgensis.
nutans, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 12.—Venezuela.
orchioides, Rich. in Mém. Mus. Par. iv. (1818) 59 = Sp. orchioides.
papulosus, Lindl. Gen. et Sp. Orch. 477.—Mexic.
pauciflorus, Reichb. f. in Bonplandia, iv. (1856) 211 = Sp. pauciflora.
polystachyon, Spreng. Syst. iii. 710 (= *Chloidia vernalis*).—Jamaic.
speciosus, Rich. in Mém. Mus. Par. iv. (1818) 59 = Sp. colorata.
squamulosus, Spreng. Syst. iii. 710.—N. Granat.
sulphureus, Lindl. Gen. et Sp. Orch. 478 = Sp. sulphurea.
umbrosa, Rodr. Orch. Nov. i. p. x (Index) = Sp. umbrosa.
vaginatus, Spreng. Syst. iii. 710; Reichb. f. in Linnaea, xli. (1877) 101 = Sp. vaginata.

STENOSEMIS, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 551 (1862) = **Annesorhiza**, Cham. & Schlecht. (Umbell.).
angustifolia, E. Mey. ex Harv. & Sond. l. c.—Afr. austr.
caffra, Sond. l. c.—Afr. austr.
teretifolia, E. Mey. ex Sond. l. c. = *caffra*.

STENOSIPHON, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 326, 384. *ONAGRARIEAE*, Benth. & Hook. f. i. 792.
linifolium, Heynh. Nom. ii. 704 = virgatum.
virgatum, *Spach, in Nouv. Ann. Mus. Par.* iv. (1835) 385.—Am. bor.

STENOSIPHONIUM, Nees, in Wall. Pl. As. Rar. iii. 75 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1086.
confertum, T. Anders. in Journ. Linn. Soc. ix. (1867) 464 = diandrum.
confertum, *Nees, in DC. Prod.* xi. 105.—Ind. or.

STENOSIPHONIUM:—
diandrum, Nees, in DC. Prod. xi. 105, partim = Strobilanthes exareolatus.
diandrum, *Wight, Ic.* t. 1502.—Ind. or.
Moonianum, Wight & Gardn. ex T. Anders. in Journ. Linn. Soc. ix. (1867) 466 = Strobilanthes adenophorus.
parviflorum, *T. Anders. l. c.* 464.—Ind. or.
Russelianum, *Nees, in Wall. Pl. As. Rar.* iii. 84.—Ind. or.
Russelianum, Nees, l. c., partim = parviflorum.
Russelianum, Wight, Ic. t. 873 = confertum.
setosum, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 464.—Ind. or.
subsericeum, Nees, in Wall. Pl. As. Rar. iii. 84 = Russelianum.
zeylanicum, *T. Anders. in Thw. Enum. Pl. Zeyl.* 225.—Zeylan.

STENOSOLENIUM, Turcz. in Bull. Soc. Nat. Mosc. xiii. (1840) II. 253 = **Arnebia**, Forsk. (Borag.).
perenne, Schrenk, Enum. Pl. Nov. i. 34 = Macrotomia perennis.
saxatile, Turcz. in Bull. Soc. Nat. Mosc. xiii. (1840) II. 253.—Sibir. baical.; China.

STENOSPERMATION, Schott, Gen. Aroid. 70 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 990.
amomifolium, *Schott, Prod. Aroid.* 348.—Peruv.
angustifolium, *Hemsl. Biol. Centr. Am. Bot.* iii. 425.—Nicaragua.
marantaefolium, *Hemsl. l. c.*—Nicaragua.
Mathewsii, *Schott, Gen. Aroid.* t. 70.—Peruv.
popayanense, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 39.—N. Granat.
Spruceanum, *Schott, Gen. Aroid.* t. 70.—Bras.
Wallisi, Mast. in Gard. Chron. (1875) 558 = popayanense.

STENOSPERMATIUM, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 39 = praec.

STENOSPERMUM, Sweet, Hort. Brit. ed. II. 209 (1830) = **Metrosideros**, Banks (Myrtac.).
capitatum, Sweet, l. c. = Kunzea capitata.
ciliatum, Heynh. Nom. i. 787 = M. ciliata.
corifolium, Sweet, Hort. Brit. ed. II. 130 = Kunzea corifolia.
ericaefolium, Heynh. Nom. i. 787; ii. 338 = Kunzea ericifolia.

STENOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) II. 330. Genus dubium *GRAMINEARUM*.—Cf. Benth. & Hook. f. Gen. iii. 1228.
narduroides, *Turcz. l. c.* 331.—N. Zel.

STENOSTEPHANUS, Nees, in Mart. Fl. Bras. ix. 91 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1106.
lasiostachyus, *Nees, in DC. Prod.* xi. 311.—Venezuela.
lobeliaeformis, *Nees, in Mart. Fl. Bras.* ix. 92.—Bras.

STENOSTOMUM, Gaertn. f. Fruct. iii. 69. t. 192 (1805) = **Antirrhoea**, Comm. (Rubiac.).
acutatum, DC. Prod. iv. 460 = A. aristata.
bifurcatum, DC. l. c. = A. bifurcata.
coriaceum, Griseb. Fl. Brit. W. Ind. 334.—Ind. occ.
densiflorum, Griseb. Cat. Pl. Cub. 132 = Laugeria densiflora.
dichotomum, DC. Prod. iv. 461 = A. dichotoma, Tournefortia bicolor.
granulatum, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 507.—Cuba.
lucidum, Gaertn. f. Fruct. iii. 69 = A. lucida.
microphyllum, Griseb. Cat. Pl. Cub. 133 = Guettarda Echinodendron.
myrtifolium, Griseb. Fl. Brit. W. Ind. 334.—Ind. occ.
nitidum, DC. Prod. iv. 461.—Ind. occ.
paniculatum, Poepp. & Endl. Nov. Gen. et Sp. iii. 27. t. 23.—Peruv.?
pauciflorum, Wright, in Sauv. Fl. Cub. 67.—Cuba.
radiatum, Griseb. Cat. Pl. Cub. 132.—Ind. occ.
ramosa, Griseb. ex Benth. & Hook. f. Gen. ii. 100, sphalm. = seq.
resinosum, Griseb. Fl. Brit. W. Ind. 334 = Laugeria resinosa.
tomentosum, DC. Prod. iv. 460.—Jamaic.
viscosum, Griseb. Fl. Brit. W. Ind. 334 = A. viscosa.

STENOTAENIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 339 = **Heracleum**, Linn. (Umbell.).
Haussknechtii, Boiss. Fl. Orient. ii. 1052.—Persia.
nudicaulis, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 340.—Persia.
tordylioides, Boiss. l. c. 339.—Persia.

STENOTAPHRUM, Trin. Fund. Agrost. 175 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1108.
DIASTEMANTHE, Steud. Syn. Pl. Gram. 360 (1855).
americanum, Schrank, Pl. Rar. Hort. Monac. t. 98 = glabrum.
complanatum, Schrank, in Flora, vii. (1824) II. Beil. 26 = glabrum.
dimidiatum, Brongn. in Duperr. Voy. Coq. Bot. 127 = glabrum.
glabrum, *Trin. Fund. Agrost.* 176.—Littora tropica.
Koenigii, Schrank, in Flora, vii. (1824) II. Beil. 28 = glabrum.
lepturoides, Hensl. in Ann. Nat. Hist. Ser. I. i. (1838) 346. t. 11 = subulatum.
madagascariense, Kunth, Rév. Gram. ii. t. 211 = glabrum.
oostachyum, *Baker, in Journ. Linn. Soc.* xx. (1883) 299.—Madag.
sarmentosum, Nees, in Mart. Fl. Bras. ii. 93; Fl. Afr. Austr. 63 = glabrum.
subulatum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iii. (1835) 190.—Ins. Pacif.
Swartzianum, Nees, Fl. Afr. Austr. 62 = glabrum.
unilaterale, *Baker, in Journ. Linn. Soc.* xx. (1883) 299.—Madag.

STENOTHECA, Monn. Ess. Hierac. 71 (1829) = **Hieracium**, Tourn. (Compos.).
mariana, Monn. l. c. 72 = H. marianum.
subnuda, Monn. l. c. = H. venosum.
venosa, Monn. l. c. = H. venosum.

STENOTIUM, Presl, ex Steud. Nom. ed. II. ii. 58, in syn. (1841) = **Lobelia**, Linn. (Campan.).

STENOTROPIS, Hassk. Retzia, i. 183 (1855) = **Erythrina**, Linn. (Legumin.).
Berteroi, Hassk. in Retzia, i. 183 = E. poianthes.

STENOTUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 334 = **Aplopappus**, Cass. (Compos.).
acaulis, Nutt. l. c. = A. acaulis.
armerioides, Nutt. l. c. 335 = A. armerioides.
caespitosus, Nutt. l. c. = A. acaulis.
florifer, Torr. & Gray, Fl. N. Am. ii. 238 = Townsendia florifer.
linearifolius, Torr. & Gray, l. c. = A. linearifolius.
multicaulis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 335 = A. multicaulis.
pygmaeus, Torr. & Gray, Fl. N. Am. ii. 237 = A. pygmaeus.

STENURUS, Salisb. Gen. Pl. Fragm. 5 (1866) = **Biarum**, Schott (Aroid.).
tenuifolium, Salisb. l. c. = B. tenuifolium.

STEPHANANDRA, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1843) 739. t. 4. *ROSACEAE*, Benth. & Hook. f. i. 612.
chinensis, *Hance, in Journ. Bot.* xx. (1882) 210.—China.
flexuosa, *Sieb. & Zucc. in Abh. Akad. Muench.* iii. (1843) 739. t. 4.—Japon.
gracilis, *Franch. & Sav. Enum. Pl. Jap.* ii. 333.—Japon.
incisa, Zabel, in Berl. Garten-Zeit. iv. (1885) 511 = flexuosa.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 332.—Japon.

STEPHANANTHUS, Lehm. Ind. Sem. Hort. Hamb. (1826) 18; et in Flora, x. (1827) 96 = **Baccharis**, Linn. (Compos.).
junceus, Lehm. ll. cc. = B. juncea.

STEPHANIA, Lour. Fl. Cochinch. 608 (1790). *MENISPERMACEAE*, Benth. & Hook. f. i. 37.
CLYPEA, Blume, Bijdr. 26 (1825).
HOMOCNEMIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40.
ILIOCARPUS, Miers, l. c. (1851).
PERICHASMA, Miers, l. c. Ser. III. xviii. (1866) 21.
STYPHANIA, C. Muell. in Walp. Ann. vii. 66 (1868).

STEPHANIA:—
abyssinica, *Walp. Rep.* i. 96.—Abyss.
acuminata, *Spreng. Syst.* iv. *Cur. Post.* 316.—Java.
acuminatissima, Walp. Rep. i. 96 = praec.
appendiculata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 15.—Ind. or.
australis, A. Gray, Bot. U. St. Expl. Exped. i. 38 = discolor.
australis, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = Cissampelos Pereira.
bullulata, Miers, l. c. Ser. III. xviii. (1866) 16 = abyssinica.
Burmanni, Walp. Rep. i. 96 = Cyclea Burmanii.
capitata, *Spreng. Syst.* iv. *Cur. Post.* 316.—Java.
cauliflora, *Becc. Malesia*, i. (1877) 155.—Ins. Celebes.
concinna, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 15.—Java.
convolvulacea, Miers, l. c. Ser. II. vii. (1851) 40 = Cissampelos Pereira.
corymbosa, *Walp. Rep.* i. 96.—Malaya.
discolor, *Spreng. Syst.* iv. *Cur. Post.* 316.—Geront. trop.
elegans, *Hook. f. & Thoms. Fl. Ind.* i. 195.—Reg. Himal.
exigua, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 16.—Ins. Philipp.
fastuosa, Miers, l. c.—Afr. trop.
florulenta, *Becc. Malesia*, i. (1877) 153.—N. Guin.
Forsteri, A. Gray, Bot. U. St. Expl. Exped. i. 36 = discolor.
Gaudichaudii, A. Gray, l. c. 37 = discolor.
glabra, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = rotunda.
glandulifera, Miers, l. c. Ser. III. xviii. (1866) 15 = rotunda.
glaucescens, *Walp. Rep.* i. 96.—Ins. Timor.
gracilenta, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 15.—Ind. or.
hernandifolia, *Walp. Rep.* i. 96 = discolor.
hexandra, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 40 = rotunda.
hypoglauca, Miers, l. c. Ser. III. xviii. (1866) 15 = discolor.
intertexta, Miers, l. c. = discolor.
japonica, Miers, l. c. 14 = rotunda.
laetificata, *Oliver, Fl. Trop. Afr.* i. 47.—Afr. trop.
laevigata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 16 = abyssinica.
latifolia, Miers, l. c. 15 = discolor.
longa, Lour. Fl. Cochinch. 609 = rotunda.
longifolia, *Becc. Malesia*, i. (1877) 156.—Borneo.
Meyeriana, *Harv. Gen. S. Afr. Pl.* ed. II. 4.—Afr. austr.
obvia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 15.—Java.
pallidula, *Miers, l. c.* 16.—Austral.
praelata, *Miers, l. c.*—Afr. austr.
racemosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 282.—Ind. or.
ramuliflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xviii. (1866) 16.—Borneo.
rotunda, *Lour. Fl. Cochinch.* 608.—As. trop.
rotundata, Miq. Fl. Ind. Bat. i. II. 84 = tomentosa.
Roxburghiana, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 14 = rotunda.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 9 = abyssinica.
tetrandra, *S. Moore, in Journ. Bot.* xiii. (1875) 225.—China.
tomentosa, *Spreng. Syst.* iv. *Cur. Post.* 316.—Java.
venosa, *Spreng. l. c.* 96 = rotunda.

STEPHANIA, Willd. Sp. Pl. ii. 239 (1799) = **Steriphoma**, Spreng. (Cappar.).
cleomoides, Willd. l. c. = Steriphoma cleomoides.
elliptica, DC. Prod. i. 253 = Steriphoma ellipticum.

STEPHANIUM, Schreb. Gen. 124 (1789) = **Palicourea**, Aubl. (Rubiac.).
guianense, J. F. Gmel. Syst. 368 = P. guianensis.

STEPHANOCARPUS, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 368 = **Cistus**, Tourn.
monspeliensis, Spach, l. c. 369 = C. monspeliensis.

STEPHANOCHILUS, Coss. & Dur. ex Benth. & Hook. f. Gen. ii. 477 (1873) = **Volutarella**, Cass. (Compos.).
omphalodes, Coss. & Dur. ex Benth. & Hook. f. l. c. = V. omphalodes.

STEPHANOCOMA, Less. Syn. Comp. 56 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 461.
atriplicifolium, Turcz. ex Steud. Nom. ed. II. ii. 639 = Serratula atriplicifolia.
berckheyoides, *Less. Syn. Comp.* 56.—Afr. austr.
berkheyoides, Sch. Bip. in Flora, xxvii. (1844) 775 (= *Stobaea echinopoda*).—Afr. austr.
carduoides, Less. Syn. Comp. 56 = Berkheya carduiformis.
incana, Sch. Bip. in Flora, xxvii. (1844) 774 = Berkheya fruticosa.
rubricaulis, Sch. Bip. l. c. = B. carlinoides.

STEPHANODAPHNE, Baill. Adansonia, xi. (1875) 302. *THYMELAEACEAE*, Benth. & Hook. f. iii. 198.
Boivini, *Baill. l. c.*—Ins. Comor.
cremostachya, *Baill. l. c.* 303.—Madag.

STEPHANOGASTRA, Karst. & Triana, in Linnaea, xxviii. (1856) 425 = **Centronia**, D. Don (Melastom.).
purpurea, Karst. & Triana, l. c. 426 = C. Mutisii.

STEPHANOLIRION, Baker, in Gard. Chron. (1875) i. 234 = **Tristagma**, Poepp. (Liliac.).
narcissoides, Baker, l. c. (= *Stemmatium narcissoides*).—Chili.

STEPHANOMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 427. *COMPOSITAE*, Benth. & Hook. f. ii. 533.
HEMIPTILIUM, Torr. Bot. Emory Exped. 105 (1858).
JAMESIA, Nees, in Neuwied, Reise N. Am. ii. 442 (1844).
PTILORIA, Rafin. Atl. Journ. 145 (1832), nomen prius.
cichoriacea, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 552.—Calif.
coronaria, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 194.—Calif.
elata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 173.—Calif.
exigua, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 428.—Am. bor. occ.
heterophylla, Nutt. l. c. 427 = runcinata.
intermedia, Kellogg, in Proc. Calif. Acad. v. (1873) 39 = Microseris nutans.
lactucina, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 552.—Calif.
minor, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 427.—Am. bor. occ.
myriocladia, *Eaton, in Bot. King's Exp.* 198. t. 20.—Am. bor. occ.
paniculata, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 428.—Am. bor. occ.
Parryi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 61.—Calif.
pentachaeta, *Eaton, in Bot. King's Exp.* 199. t. 20.—Am. bor. occ.
runcinata, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 428.—Am. bor. occ.
Schottii, *Benth. & Hook. f. Gen.* ii. 533; *A. Gray, in Bot. Calif.* i. 427.—Arizona.
Thurberi, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 325.—N. Mexic.
virgata, *Benth. Bot. Voy. Sulph.* 32.—Am. bor. occ.
Wrightii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 62.—Am. bor. occ.

STEPHANOPAPPUS, Less. in Linnaea, vi. (1831) 234 = **Nestlera**, Spreng. (Compos.).
reflexus, Less. l. c. = N. biennis.

STEPHANOPHYLLUM, Guill. in Deless. Ic. Sel. iii. 61 (1837) = **Paepalanthus**, Mart. (Eriocaul.).
flagellaris, Guill. l. c. 61 = P. flagellaris.
proliferum, Guill. l. c. = P. prolifer.
viviparum, Guill. l. c. = P. viviparus.

STEPHANOPHORUM, Dulac, Fl. Hautes-Pyr. 133 (1867) = **Narcissus**, Tourn. (Amaryll.).
biflorum, Dulac, l. c. = N. biflorus.
filiforme, Dulac, l. c. 134 = N. juncifolius.
grandiflorus, Dulac, l. c. = N. Pseudo-narcissus.
infundibuliformis, Dulac, l. c. = N. Bulbocodium.
luteum, Dulac, l. c. = N. Jonquilla.
purpuraceum, Dulac, l. c. 133 = N. poëticus.

STEPHANOPHYSUM, Pohl, Pl. Bras. Ic. ii. 83. tt. 155, 156 (1831) = **Ruellia**, Plum. (Acanth.).
angustiflorum, Nees, in Mart. Fl. Bras. ix. 51.—Bras.
asperulum, Mart. & Nees, l. c. 52.—Bras.
attenuatum, Mart. in Flora, xx. (1837) ii. Beibl. 123 = Geissomeria longiflora.
Baikiei, Hook. Bot. Mag. t. 5111.—Afr. trop.
brevifolium, Pohl, Pl. Bras. Ic. ii. 84. t. 155.—Bras.
ciliare, Nees, in Mart. Fl. Bras. ix. 52.—Bras.
cordifolia, Nees, l. c. 51.—Bras.
flavum, Nees, l. c.—Bras.
Herbstii, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 269 = Ruellia Herbstii.
hirsutulum, Mart. ex Nees, in DC. Prod. xi. 205 = *asperulum*.
hirsutum, Mart. ex Nees, in Mart. Fl. Bras. ix. 50.—Bras.
longifolium, Pohl, Pl. Bras. Ic. ii. 85. t. 156 = R. amoena.
molle, Nees, in DC. Prod. xi. 203.—Peruv.
quadrifarium, Nees, l. c. 202 = Macrostegia Ruiziana ?
repens, T. Anders. in Journ. Linn. Soc. vii. (1864) 25 = *Baikiei*.
Ruizianum, Nees, in DC. Prod. xi. 203.—Peruv.; Chili.
tubulatum, Nees, l. c. 202 = R. tubulata.
ventricosum, Nees, l. c. 203 = R. ventricosa.

STEPHANOPODIUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 40. t. 246 (1842). *CHAILLETIACEAE*, Benth. & Hook. f. i. 341.
Engleri, *Baill. Adansonia*, xi. (1873–76) 109.—Bras.
peruvianum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 40. t. 246.—Peruv.

STEPHANOSIPHON, Boiv. ex C. DC. in DC. Monog. Phan. i. 446 (1878) = **Turraea**, Linn. (Meliac.).
madagascariense, Boiv. ex C. DC. l. c. = T. sericea.

STEPHANOSTACHYS, Klotzsch, ex Oerst. in Kjoeb. Vidensk. Meddel. (1858) 26 = **Chamaedorea**, Willd. (Palm.).
Casperiana, Klotzsch, ex Oerst. l. c. 27; et Oerst. L'Amérique Centrale, 14. t. 7 = C. Casperiana.
Martiana, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 29 = C. Martiana.
Tepejilote, Oerst. l. c. 28 = C. Tepejilote.
Wendlandiana, Oerst. l. c. = C. Wendlandi.

STEPHANOTELLA, Fourn. in Mart. Fl. Bras. vi. iv. 326 (1885). *ASCLEPIADEAE*.
Glaziovii, *Fourn. l. c.*—Bras.

STEPHANOTIS, Thou. Gen. Nov. Madag. 11 (1806) *ASCLEPIADEAE*, Benth. & Hook. f. ii. 773.
ISAURA, Comm. ex Poir. Encyc. Suppl. iii. 185 (1813).
JASMINANTHES, Blume, Mus. Bot. Lugd. Bat. i. 148. t. 28 (1850).
acuminata, *Brongn. in Ann. Sc. Nat.* Sér. II. vii. (1837) 30.—Madag.
chinensis, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 53.—Ins. Hongk.
floribunda, *Brongn. in Ann. Sc. Nat.* Sér. II. vii. (1837) 30.—Madag.
grandiflora, *Decne. in DC. Prod.* viii. 620.—Madag.
Isaura, *Decne. l. c.*—Madag.
longiflora, *Benth. & Hook. f. Gen.* ii. 773.—Cuba.
Maingayi, *Hook. f. Fl. Brit. Ind.* iv. 39.—Malacca.
Thouarsii, *Brongn. in Ann. Sc. Nat.* Sér. II. vii. (1837) 30.—Madag.
vincaeflora, *Benth. & Hook. f. Gen.* ii. 773.—Cuba.

STEPHANOTRICHUM, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 54 = **Clidemia**, D. Don (Melastom.).
hispidum, Naud. l. c. = Heterotrichum octonum.
octonum, Naud. ex Hemsl. Biol. Centr. Am. Bot. i. 430 = Heterotrichum octonum.

STEPHEGYNE, Korth. Verh. Nat. Gesch. Bot. 160. t. 35 (1840 ?) = **Mitragyna**, Korth. (Rubiac.).
africana, Walp. Rep. ii. 513 = M. africana.
diversifolia, Benth. & Hook. f. Gen. ii. 31; Hook. f. Fl. Brit. Ind. iii. 26.—Burma; Ins. Philipp.

STEPHEGYNE:—
parvifolia, Korth. Verh. Nat. Gesch. Bot. 161 = Mitragyna parvifolia.
parvifolia, Vidal, Sinops. t. 56. f. A = *diversifolia*.
speciosa, Korth. Verh. Nat. Gesch. Bot. 160. t. 35 = M. speciosa.
stipulata, Benth. & Hook. f. Gen. ii. 31 = M. macrophylla.
tubulosa, Benth. & Hook. f. l. c.—Ind. or.
tubulosa, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 104 = *diversifolia*.

STEPHENSONIA, Hort. ex Van Houtte, in Fl. des Serres, xv. (1862–63) 177 = **Stevensonia**, Duncan (Palm.).
grandifolia, Hort. ex Van Houtte, l. c. = Stevensonia grandifolia.

STEPTIUM, Boiss. Fl. Orient. iv. 533 (1879) = STREPTIUM, Roxb. = **Priva**, Adans. (Verben.).

STEPTORHAMPHUS, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 381 = **Lactuca**, Linn. (Compos.).
crambifolius, Bunge, l. c. = L. crambifolia.
hispidulus, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 176.—Turkestan.

STERBECKIA, Schreb. Gen. i. 360 (1789) = **Singana**, Aubl. (Inc. sed.).
guianensis, J. F. Gmel. Syst. 820 = Sing. guianensis.
lateriflora, Willd. Sp. Pl. ii. 1177 = Sing. guianensis.

STERCULIA, Linn. Diss. Dass. 29 (1747); Amoen. Acad. i. 412 (1749). *STERCULIACEAE*, Benth. & Hook. f. i. 217.
BALANGHAS, Rafin. Sylva Tellur. 72 (1838).
BRACHYCHITON, Schott, Melet. 34 (1832).
BRAXIPIS, Rafin. Sylva Tellur. 72 (1838).
CARPOPHYLLUM, Miq. Fl. Ind. Bat. Suppl. 401 (1860).
CAVALAM, Adans. Fam. ii. 357 (1763).
CHICHAEA, Presl, Rel. Haenk. ii. 140 (1836).
CLOMPANUS, Rafin. Sylva Tellur. 73 (1838).
COLARIA, Rafin. l. c. (1838).
CULHAMIA, Forsk. Fl. Aegypt. Arab. 96 (1775).
DELABECHEA, Lindl. in Mitch. Journ. Trop. Austral. 155 (1848).
ERYTHROPSIS, Lindl. in Brande, Journ. (Sept. 1827) 112, in obs.
FIRMIANA, Marsigli, in Sagg. Padov. i. (1786) 106, cum ic.
HILDEGARDIA, Schott, Melet. 33 (1832).
ICOSINIA, Rafin. Sylva Tellur. 75 (1838).
IVIRA, Aubl. Pl. Gui. 694. t. 279 (1775).
JOIRA, Steud. Nom. ed. II. ii. 639, in syn. (1841).
KARAKA, Rafin. Sylva Tellur. 72 (1838).
KAVALAMA, Rafin. l. c. (1838).
MATEATIA, Vell. Fl. Flum. ix. t. 95 (1827).
OLEOBACHIA, Hort. ex Mast. in Gard. Chron. (1880) ii. 232.
ORSOPEA, Rafin. Sylva Tellur. 73 (1838).
POECILODERMIS, Schott, Melet. 33 (1832).
POMPILA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3.
PTEROCYMBIUM, R. Br. in Benn. Pl. Jav. Rar. 219. t. 45 (1838 ?).
PTERYGOTA, Schott, Melet. 32 (1832).
SCAPHIUM, Schott, l. c. (1832).
SOUTHWELLIA, Salisb. Parad. Lond. t. 69 (1806).
THEODORIA, Neck. Elem. ii. 286 (1790).
TRICHOSIPHON, Schott, Melet. 34 (1832).
TRIPHACA, Lour. Fl. Cochinch. 577 (1790).
TRIPLOBUS, Rafin. Sylva Tellur. 110 (1838).
abyssinica, R. Br. in Salt, Abyss. App. 64 = S. Triphaca.
abyssinica, R. Br. l. c., partim = arabica.
acerifolia, *A. Cunn. in Loud. Hort. Brit.* 392.—Austral.
acerifolia, A. Cunn. l. c., partim = incana.
acerifolia, *Hemsl. Biol. Centr. Am. Bot.* i. 126.—Mexic.
acuminata, Beauv. Fl. Owar. i. 41. t. 24 = Cola acuminata.
affinis, *Mast. in Hook. f. Fl. Brit. Ind.* i. 361.—Malacca.
alata, Blanco, Fl. Filip. ed. II. 525 = Blancoi.
alata, *Roxb. Hort. Beng.* 50; *Pl. Corom.* iii. 84. t. 287.—Ind. or.
alata, Wall. Cat. n. 1127 D = guttata.

STERCULIA:—
Alexandri, *Harv. in Proc. Dubl. Univ. Zool. & Bot. Assoc.* i. (1859) 140. t. 15.—Afr. austr.
angustifolia, Jack, in Malay. Misc. i. (1820) n. 1. 21 = rubiginosa.
angustifolia, Roxb. Hort. Beng. 50; Fl. Ind. iii. 148 = S. Balanghas.
apetala, Karst. Fl. Columb. ii. 35 = carthaginensis.
arabica, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 9.—Arab.
arbutiflora, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1830) 178, nomen (Quid?).—N. Guin.
armata, Mast. in Hook. f. Fl. Brit. Ind. i. 357 = ornata, etc.
Balanghas, Ait. Hort. Kew. ed. II. v. 338 = nobilis.
Balanghas, *Linn. Sp. Pl.* 1007.—As. trop.
Barteri, *Mast. in Oliver, Fl. Trop. Afr.* i. 218.—Afr. trop.
bicolor, *Mast. in Hook. f. Fl. Brit. Ind.* i. 359.—Malacca.
Bidwilli, *Hook. ex Benth. Fl. Austral.* i. 228.—Austral.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 308.—Ins. Philipp.
Blumei, *G. Don, Gen. Syst.* i. 515.—Java.
bullata, *Panch. & Sebert, Not. Bois Nouv. Calléd.* 214.—N. Caled.
campanulata, *Wall. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 362.—Burma; Java.
Candollii, Wall. Pl. As. Rar. i. 4, in textu = populnifolia.
caribaea, *R. Br. in Benn. Pl. Jav. Rar.* 228.—Ind. occ.
caricaefolia, G. Don, Gen. Syst. i. 517 = Cola Afzelii.
carthaginensis, R. Br. in Benn. Pl. Jav. Rar. 328, partim = P. Chica.
carthaginensis, *Cav. Diss.* vi. 353.—Am. trop.
caudata, *Heward, ex Benth. Fl. Austral.* i. 230.—Austral.
ceramica, *R. Br. in Benn. Pl. Jav. Rar.* 233.—Ins. Molucc.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 486.—Madag.
Chicha, *A. St. Hil. Fl. Bras. Mer.* i. 278. t. 46.—Bras.
chrysodasys, *Miq. Fl. Ind. Bat. Suppl.* 399.—Sumatra.
cinerea, *A. Rich. Tent. Fl. Abyss.* i. 74. t. 16.—Abyss.
cinerea, Schweinf. Pl. Nilot. 15 = Hartmanniana.
citrifolia, Salisb. Prod. 387 = S. Balanghas.
coccinea, Jack, in Malay. Misc. i. (1820) n. 1. 20 = laevis.
coccinea, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 151.—Ind. or.
coccinea, Wall. Cat. n. 1122, partim = alata.
Cola, Pers. Syn. ii. 239 = Cola acuminata.
colorata, *Roxb. Pl. Corom.* i. 26. t. 25.—Ind. or.
comorensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 485.—Madag.
comosa, *Wall. Pl. As. Rar.* ii. 25.—Ins. Amboin.
convoluta, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = S. Helicteres.
cordata, *Blume, Bijdr.* 83.—Java.
cordifolia, *Blanco, Fl. Filip.* ed. I. 764.—Ins. Philipp.
cordifolia, *Cav. Diss.* v. 286. t. 144. f. 2.—Afr. trop.
crassinervia, *Miq. Fl. Ind. Bat. Suppl.* 399.—Sumatra.
crepitans, F. Muell. Ind. Perf. Linn. Sp. Pl. 37, sphalm. = Hura crepitans.
crinita, Cav. Diss. v. 285 = S. Ivira.
cuneata, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Ins. Philipp.
cuneata, Heyne, in Wall. Cat. n. 1127[3] = guttata.
cuneifolia, *Miq. Fl. Ind. Bat. Suppl.* 400.—Sumatra.
cuspidata, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Sumatra.
cymbiformis, *Blanco, Fl. Filip.* ed. II. 526 (Quid?).—Ins. Philipp.
decandra, Blanco, l. c. ed. I. 766 = Myristica intermedia.
discolor, *F. Muell. ex Benth. Fl. Austral.* i. 228.—Austral.
diversifolia, *G. Don, Gen. Syst.* i. 516.—Austral.
diversifolia, *Seem. Fl. Vit.* 23.—Ins. Fiji.
ensifolia, *Mast. in Hook. f. Fl. Brit. Ind.* i. 359.—Burma; Penang.

STERCULIA:—
erythrosiphon, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 486.—Madag.
excelsa, *Mart. in Flora*, xxiv. (1841) 11. *Beibl.* 40.—Bras.
ferruginea, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Ins. Philipp.
Fiorniana, J. F. Gmel. Syst. 1034 = plantanifolia.
Firmiana, J. F. Gmel. l. c. 1632 = praec.
foetida, *Linn. Sp. Pl.* 1008.—Geront. trop.
Forsteri, *Seem. Fl. Vit.* 23, in nota.—Ins. Tanna.
frondosa, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guian.
fulgens, *Wall. Cat.* n. 1135.—Reg. Himal.; Burma; Malaya.
geniculata, *Miq. Fl. Ind. Bat. Suppl.* 400.—Sumatra.
gilva, *Miq. l. c.* 401.—Sumatra.
glandulosa, Blanco, Fl. Filip. ed. I. 764 = Myristica simiarum.
glomerata, Blanco, l. c. = Myristica glauca?
gracilis, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 306.—Borneo.
grandiflora, Vent. Jard. Malm. sub t. 91 = Cola acuminata.
grandifolia, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Java.
guineensis, Hort. ex Steud. Nom. ed. II. ii. 640 (Quid?).—Afr. trop.?
guttata, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 148.—Ind. or.; Malaya.
Halmahairae, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 88, nomen.—N. Guin.
Hartmanniana, *Schweinf. Rel. Kotschy.* 39.—Afr. trop.
Helicteres, *Pers. Syn.* ii. 240.—N. Granat.
heterophylla, Beauv. Fl. Owar. i. t. 40 = Cola heterophylla.
heterophylla, Cunn. ex Sweet, Hort. Brit. ed. III. 82 = diversifolia, *G. Don*.
Heynei, Bedd. Fl. Sylv. t. 230 = alata.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 485.—Madag.
hyposticta, *Miq. Fl. Ind. Bat. Suppl.* 399 = Sumatra.
imberbis, *DC. Prod.* i. 483.—Guian.
incana, *Benth. Fl. Austral.* i. 228.—Austral.
insularis, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Ins. Molucc.
ipomoeaefolia, Garcke, in Peters, Reise Mossamb. Bot. 130 = S. Triphaca.
Ivira, Pers. Syn. ii. 239 = imberbis.
Ivira, *Sw. Prod. Veg. Ind. Occ.* 98.—Am. austr.
Ivira, Sw. l. c., partim = caribaea.
Jackiana, Wall. Cat. n. 1134; et in Hook. Bot. Misc. i. (1830) 287 = rubiginosa.
javanica, *R. Br. in Benn. Pl. Jav. Rar.* 230.—Java.
Joira, J. F. Gmel. Syst. 1633 (Index) = S. Ivira.
laevis, *Wall. Cat.* n. 1138; *et in Hook. Bot. Misc.* i. (1830) 287.—Burma; Malaya.
lanceaefolia, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 150.—Reg. Himal.
lanceolata, Blume, Bijdr. 82 = Blumei.
lanceolata, Buch.-Ham. in Wall. Cat. n. 1122 = coccinea.
lanceolata, *Cav. Diss.* v. 287.—China; Java.
lasiantha, Mart. Med. Mat. Bras. (1843) 18 = striata.
laurifolia, *F. Muell. Fragm.* vi. 172.—Austral.
linearicarpa, *Mast. in Hook. f. Fl. Brit. Ind.* i. 360.—Malacca.
linearis, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 305.—Borneo.
linguifolia, *Mast. in Hook. f. Fl. Brit. Ind.* i. 357.—Burma.
longifolia, Vent. Jard. Malm. sub t. 91 = Cola acuminata.
lurida, *F. Muell. ex Benth. Fl. Austral.* i. 228.—Austral.
lychnophora, *Hance, in Journ. Bot.* xiv. (1876) 243.—Cambodia.
macrocarpa, G. Don, Gen. Syst. i. 515 = Cola acuminata.
macrophylla, *Vent. Jard. Malm.* sub t. 91.—Malaya.
madagascariensis, *R. Br. in Benn. Pl. Jav. Rar.* 229.—Madag.
Maingayi, *Mast. in Hook. f. Fl. Brit. Ind.* i. 359.—Malacca.
mexicana, *R. Br. in Benn. Pl. Jav. Rar.* 227.—Mexic.

STERCULIA:—
mollis, Wall. Cat. n. 1131 = S. Balanghas.
monosperma, Vent. Jard. Malm. t. 91 = nobilis.
nitida, Vent. l. c. sub t. 91 = Cola acuminata.
nobilis, *Sm. in Rees, Cycl.* xxxiv. n. 4.—As. trop.
oblonga, *Mast. in Oliver, Fl. Trop. Afr.* i. 216.—Afr. trop.
oblongata, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Ins. Philipp.
oblongifolia, [*Moç. & Sesse,*] *ex DC. Prod.* i. 482.—Mexic.
obovata, R. Br. in Benn. Pl. Jav. Rar. 233 = S. Tragacantha.
obtusifolia, Zipp. ex Span. in Linnaea, xv. (1841) 173 = populnifolia.
ornata, *Wall. ex Voigt, Hort. Suburb. Calc.* 105, nomen; *et ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 228.—Burma.
ovalifolia, Wall. Cat. n. 1132 = lanceaefolia.
paniculata, *Wall. ex Boj. Hort. Maurit.* 36.—Ind. or.
parviflora, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 147.—Reg. Himal.; Malaya.
parvifolia, *Wall. Cat.* n. 1123.—Malaya.
peltata, G. Don, Gen. Syst. i. 517 = Pterospermum heterophyllum?
platanifolia, *Linn. f. Suppl.* 423.—China; Japon.
polyphylla, *R. Br. in Benn. Pl. Jav. Rar.* 227.—Sumatra.
populifolia, DC. Prod. i. 483 (an seq.?).—Ins. Timor.
populnifolia, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 148.—Ind. or.
propinqua, *R. Br. in Benn. Pl. Jav. Rar.* 229.—Guian.
pubescens, G. Don, Gen. Syst. i. 515 = S. Tragacantha.
pubescens, *Mast. in Hook. f. Fl. Brit. Ind.* i. 357.—Malacca.
pubescens, *F. Muell. ex Benth. Fl. Austral.* i. 228, in nota.—Austral.
punctata, [*Moç. & Sesse,*] *ex DC. Prod.* i. 483.—Mexic.
punduana, Wall. Cat. n. 2701 = Reevesia Wallichii.
purpurascens, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 306.—Borneo.
pyriformis, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 83 = platanifolia.
quadrifida, *R. Br. in Benn. Pl. Jav. Rar.* 233.—Austral.
ramiflora, *Benth. Fl. Austral.* i. 227.—Austral.
Richardiana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 486.—Madag.
robusta, *Carr. in Rev. Hortic.* (1868) 120.—China.
Roxburghiana, Wall. Cat. n. 1124 = seq.
Roxburghii, Wall. Pl. As. Rar. iii. 39 = lanceaefolia.
rubicunda, Wall. Cat. n. 1119 D, F, G = colorata.
rubiginosa, *Vent. Jard. Malm.* sub t. 91.—Burma; Malaya.
rubiginosa, Zoll. ex Miq. Fl. Ind. Bat. Suppl. 399 = chrysodasys.
rufa, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 306.—Borneo.
rugosa, *R. Br. in Benn. Pl. Jav. Rar.* 229.—N. Granat.
rupestris, *Benth. Fl. Austral.* i. 230.—Austral.
scaphigera, *Wall. Cat.* n. 1130.—Burma; Malaya.
scatigera, G. Don, Gen. Syst. i. 517, sphalm. = praec.
setigera, Delile, Voy. Méroë Bot. 61 = tomentosa?
Spangleri, *R. Br. in Benn. Pl. Jav. Rar.* 230.—Java.
spectabilis, *Miq. Fl. Ind. Bat. Suppl.* 400.—Sumatra.
stipularis, *R. Br. in Benn. Pl. Jav. Rar.* 232.—Ins. Philipp.
stipulata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 306.—Borneo.
striata, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 213.—Bras.
striatiflora, *Mast. in Hook. f. Fl. Brit. Ind.* i. 356.—Burma.
subpeltata, *Blume, Bijdr.* 83.—Java.
surinamensis, R. Br. in Benn. Pl. Jav. Rar. 229 = excelsa.
Tavia, *Baill. Adansonia*, x. (1872) 177.—Ins. Molucc.
Thwaitesii, *Mast. in Hook. f. Fl. Brit. Ind.* i. 361.—Zeylan.
tomentosa, *Guill. & Perr. Fl. Seneg. Tent.* i. 81. t. 16.—Afr. trop.
tomentosa, Thunb. Jap. Ic. Dec. iv. t. 8 = platanifolia.
Tragacantha, *Lindl. Bot. Reg.* t. 1353.—Afr. trop.
Trichosiphon, *Benth. Fl. Austral.* i. 229.—Austral.
triloba, *G. Don, Gen. Syst.* i. 517 (Quid?).—Ind. or.

STERCULIA:—
Triphaca, *R. Br. in Benn. Pl. Jav. Rar.* 228.—Afr. trop.
tubulata, *Mast. in Hook. f. Fl. Brit. Ind.* i. 362.—Malacca.
urceolata, *Sm. in Rees, Cycl.* xxxiv. n. 3.—Ins. Amboin.
urens, *Roxb. Pl. Corom.* i. 25. t. 24.—Ind. or.
versicolor, *Wall. Pl. As. Rar.* i. 48. t. 59.—Burma.
verticillata, Thonn. in Schum. & Thonn. Beskr. Guin. Pl 240 = Cola acuminata.
villifera, Steud. in Flora, xxvi. (1843) 755 = excelsa.
villosa, *Roxb. Hort. Beng.* 50; *Fl. Ind.* iii. 153.—Ind. or.
vitiensis, *Seem. Fl. Vit.* 23.—Ins. Fiji.
Wallichii, *G. Don, Gen. Syst.* i. 516 (Quid?).—Ind. or.
Zippelii, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 307.—Java.

STEREOCHILUS, Lindl. in Journ. Linn. Soc. iii. (1859) 38 = **Sarcochilus**, R. Br. (Orchid.).
hirtus, Lindl. l. c.—Reg. Himal.

STEREODERMA, Blume, Fl. Jav. Praef. 9 (1828) = = **Olea**, Tourn.
javanica, Blume, l. c. = O. maritima.

STEREOSANDRA, Blume, Mus. Bot. Lugd. Bat. ii. 176 (1856). *ORCHIDEAE*, Benth. & Hook. f iii. 607.
javanica, *Blume, l. c.*—Java.

STEREOSPERMUM, Cham. in Linnaea, vii. (1832) 720. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1047.
DIPTEROSPERMA, Hassk. Cat. Hort. Bog. Alt. 152 (1844).
HIERANTHES, Rafin. Sylva Tellur. 79 (1838).
LAGAROPYXIS, Miq. Ann. Mus. Bot. Lugd. Bat. i. 198 (1863-64).
RADERMACHERA, Zoll. & Mor. in Zoll. Syst. Verz. Ind. Archip. 53 (1854).
Arguezona, *A. Rich. Tent. Fl. Abyss.* ii. 58.—Abyss.
Banaibanai, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 314.—Ins. Philipp.
caudatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 200.—Ind. or.
chelonoides, *DC. in Bibl. Univ. Genèv.* xvii. (1838) 124; *Prod.* ix. 210.—Ind. or.; Burma.
dentatum, *A. Rich. Tent. Fl. Abyss.* ii. 58.—Abyss.
euphorioides, *DC. Prod.* ix. 211.—Madag.
filiforme, DC. in Bibl. Univ. Genèv. xvii. (1838) 124 = Spathodea filiformis.
fimbriatum, *DC. l. c.* 124; *et Prod.* ix. 211.—Burma; Malaya.
Ghorta, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 384.—Ind. or.
glandulosum, *Miq. Fl. Ind. Bat. Suppl.* 565.—Burma; Malaya.
Hasskarlii, *Teysm. ex Miq. Fl. Ind. Bat.* ii. 756 (an seq.?).—Java.
Hasskarlii, *Zoll. & Mor. in Zoll. Syst. Verz. Ind. Archip.* 54.—Malaya.
hypostictum, *Miq. Fl. Ind. Bat. Suppl.* 565.—Burma; Malaya.
integrifolium, *A. Rich. Tent. Fl. Abyss.* ii. 59.—Afr. trop.
Kunthianum, *Cham. in Linnaea*, vii. (1832) 720.—Afr. trop.
nematocarpum, *DC. Prod.* ix. 211.—Madag.
neuranthum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 91.—Burma.
pinnatum, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 151.—Ins. Philipp.
quadripinnatum, *F. Villar, l. c.* 151. t. 252.—Ins. Philipp.
Seemannii, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 314.—Ins. Philipp.
senegalense, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 200.—Afr. trop.
serrulatum, DC. in Bibl. Univ. Genèv. xvii. (1838) 124; Prod. ix. 206 = Spathodea Serratula.
sinicum, *Hance, in Journ. Bot.* xx. (1882) 16.—China.
suaveolens, *DC. ll. cc.* 124; 211.—Ind. or.; Burma.
tetragonum, *DC. ll. cc.* 124; 210.—Ind. or.

STEREOSPERMUM:—
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 383.—Burma.
zylocarpum, *Benth. & Hook. f. Gen.* ii. 1047.—Ind. or.

STEREOXYLON, Ruiz & Pav. Prod. 38. t. 6 (1794) = **Escallonia**, Mutis (Saxifr.).
ciliatum, Poir. Encyc. Suppl. v. 247 = Micromyrtus microphylla.
corymbosum, Ruiz & Pav. Fl. Per. iii. 14 = E. corymbosa.
crenulatum, Poir. ex DC. Prod. iii. 230 = Baeckea crenulata.
cuneifolium, Ruiz & Pav. Fl. Per. iii. 16 = E. discolor?
discolor, Poir. Encyc. v. 246 = E. discolor.
paniculatum, Ruiz & Pav. Fl. Per. iii. 16 = E. floribunda.
patens, Ruiz & Pav. l. c. 13 = E. myrtilloides.
pendulum, Ruiz & Pav. l. c. 16 = E. pendula.
pulverulentum, Ruiz & Pav. l. c. 15 = E. pulverulenta.
resinosum, Ruiz & Pav. l. c. 14 = E. resinosa.
revolutum, Ruiz & Pav. l. c. 15 = E. revoluta.
rubrum, Ruiz & Pav. l. c. = E. rubra.
serratum, Poir. Encyc. vii. 435 = E. serrata.
virgatum, Ruiz & Pav. Fl. Per. iii. 14 = E. virgata.

STERIGMA, DC. Syst. ii. 579 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 102.
ARTHROLOBUS, Stev. ex DC. Syst. ii. 579 (1821).
STERIGMOSTEMUM, Bieb. Fl. Taur. Cauc. Suppl. 444 (1819).
acanthocarpum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 38.—Armen.; Peruv.
brachypetalum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 386 = sulphureum.
chionophilum, Schott, Nym. & Kotschy, Analect. Bot. 111 = Anchonium Tournefortii.
contortuplicatum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 387.—Persia.
elychrisifolium, DC. Syst. ii. 581 = Anchonium Tournefortii.
longistylum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1848) 387.—Persia.
purpurascens, *Boiss. Fl. Orient.* i. 241.—Persia; Afghan.
sulphureum, *DC. Syst.* ii. 580.—As. occ. et bor.
tomentosum, *DC. l. c.* 579.—Reg. Cauc.
torulosum, *DC. l. c.* 580.—Reg. Cauc.; Armen.

STERIGMANTHE, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 252; et in Abh. Akad. Berl. (1860) 100 = **Euphorbia**, Linn.
Bojeri, Klotzsch & Garcke, ll. cc. 252; 101 = E. Bojeri.
splendens, Klotzsch & Garcke, ll. cc. 252; 100 = E. splendens.

STERIGMOSTEMON, Poir. in Dict. Sc. Nat. l. 519 (1827) = seq.

STERIGMOSTEMUM, Bieb. Fl. Taur. Cauc. Suppl. 444 (1819) = **Sterigma**, DC. (Crucif.).
incanum, Bieb. l. c. = Sterigma torulosum.
tomentosum, Bieb. l. c. = Sterigma tomentosum.

STERIPHA, Banks, ex Gaertn. Fruct. ii. 81 (1791) = = **Dichondra**, Forst. (Convolv.).
reniformis, Soland. ex Gaertn. l. c. = D. repens.

STERIPHE, Phil. in Linnaea, xxxiii. (1864-65) 141. *COMPOSITAE*, Benth. & Hook. f. ii. 251.
corymbosa, *Phil. l. c.*—Chili.

STERIPHOMA, Spreng. Syst. iv. Cur. Post. 130, 139 (1827) *CAPPARIDEAE*, Benth. & Hook. f. i. 107.
HERMUPOA, Loefl. It. Hisp. 307 (1758).
ROEMERA, Tratt. Gen. 88 (1802).
STEPHANIA, Willd. Sp. Pl. 239 (1799).
cleomoides, *Spreng. Syst.* iv. *Cur. Post.* 139.—N. Granat.
elliptica, *l. c.*—Ins. Trinit.
paradoxum, Endl. in Flora, xv. (1832) 396; Karst. Ausw. Gew. Venez. 10. t. 3 = cleomoides.
peruviana, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 267.—Peruv.

STERIS, Adans. Fam. ii. 255 (1763) = **Lychnis**, Tourn. (Caryoph.).

STERIS, Linn. Mant. i. 8 (1767) = **Hydrolea**, Linn. (Hydroph.).
aquatica, Burm. f. Fl. Ind. 73. t. 39. f. 3 = H. zeylanica.
javanica, Linn. Mant. i. 54 = H. zeylanica.
villosa, Pav. ex A. W. Benn. in Journ. Linn. Soc. xi. (1871) 275 = H. zeylanica.
villosa, Pav. ex A. W. Benn. l. c. 277 = H. multiflora.

STERNBECKIA, Pers. Syn. ii. 69, in syn. (1807), sphalm. = STERBECKIA, Schreb. = **Singana**, Aubl. (Inc. sed.).

STERNBERGIA, Waldst. & Kit. Pl. Rar. Hung. ii. 172. t. 159 (1805). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 721.
OPORANTHUS, Herb. App. [Bot. Reg.] 38 (1821).
aetnensis, Guss. Fl. Sic. Prod. i. 395; Schult. f. Syst. vii. 794 = colchiciflora.
americana, Hoffmgg. Verz. Pfl. 197 = Haylockia pusilla.
caucasica, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 27.—Reg. Cauc.
citrina, *Ker-Gawl. ex Schult. f. Syst.* vii. 795.—Graecia.
Clusiana, Ker-Gawl. ex Schult. l. c. 794 = colchiciflora.
colchiciflora, *Waldst. & Kit. Pl. Rar. Hung.* ii. 172. t. 159.—Europ. austr.
dalmatica, Herb. Amaryll. 413. t. 47. f. 2 = colchiciflora.
exigua, Ker-Gawl. ex Schult. f. Syst. vii. 795 = Tapeinanthus humilis.
exscapa, Tineo, in Guss. Fl. Sic. Syn. i. 384 (sp. dub.).—Sicil.
Fischeriana, Roem. Syn. Ensat. 46 = lutea.
Fischeriana, *Rupr. in Regel, Gartenfl.* (1868) 100. t. 576.—Reg. Caucas.
lutea, *Ker-Gawl. ex Schult. f. Syst.* vii. 795.—Reg. Mediterr.
lutea, Orph. ex Nym. Consp. 713 = citrina.
macrantha, J. Gay, ex Boiss. Fl. Orient. v. 148 = colchiciflora.
pulchella, Boiss. & Blanche, Diagn. Ser. II. iv. 97 = colchiciflora.
Schuberti, Schenk, Pl. Sp. Aegypt. 11 = colchiciflora.
sicula, Tineo, ex Guss. Fl. Sic. Syn. ii. 811 = lutea.
stipitata, Boiss. & Haussk. ex Boiss. Fl. Orient. v. 148 = colchiciflora.

STERRHYMENIA, Griseb. in Goett. Abh. xix. (1874) 231 = **Sclerophylax**, Miers (Solan.).
Cynocrambe, Griseb. l. c.—Reg. Argent.

STETHOMA, Rafin. Fl. Tellur. iv. 61 (1836) = **Dianthera**, Gronov. (Acanth.).
pectoralis, Rafin. l. c. = D. pectoralis.

STEUARTA, Catesb. ex Mill. Gard. Dict. ed. VI. 175 App. (1752), sphalm. = **Stuartia**, Linn. (Ternstr.).

STEUDELIA, Mart. Nov. Gen. et Sp. Bras. t. 168 (1827) = **Leonia**, Ruiz & Pav. (Violar.).
cymosa, Mart. l. c. t. 169 = L. cymosa.
racemosa, Mart. l. c. t. 168 = L. glycycarpa.

STEUDELIA, Presl, Steud. Nov. Pl. Gen. (1829); et Symb. Bot. i. 3. t. 2 (1830) = **Adenogramma**, Reichb. (Ficoid.).
capillaris, Eckl. & Zeyh. Enum. 283.—Afr. austr.
diffusa, Eckl. & Zeyh. l. c. 284 = A. diffusa.
galioides, Presl, Steud. Nov. Pl. Gen. (1829); et Symb. Bot. i. 3. t. 2 = A. galioides.
sylvatica, Eckl. & Zeyh. Enum. 283 = A. sylvatica.

STEUDELIA, Spreng. Neue Entdeck. iii. 59 (1822) = **Erythroxylum**, P. Br. (Lineae).
brasiliensis, Spreng. l. c. = E. ovatum.

STEUDNERA, C. Koch, in C. Koch & Fint. Wochenschr. (1862) 114. *AROIDEAE*, Benth. & Hook. f. iii. 988.
colocasiaefolia, *C. Koch, l. c.* 114.—Malaya.
discolor, Bull, in Gard. Chron. (1875) 708 = colocasiaefolia.

STEVENA, Andrz. ex DC. Syst. ii. 291 (1821) = **Alyssum**, Tourn. (Crucif.).

STEVENIA, Adams & Fisch. in Mém. Soc. Nat. Mosc. v. (1817) 84 = **Arabis**, Linn. (Crucif.).
alyssoides, Adams & Fisch. l. c.—Sibir.
cheiranthoides, DC. Syst. ii. 210 = A. incarnata.
incana, Andrz. ex DC. l. c. 291 = A. incana.

STEVENSIA, Poit. in Ann. Mus. Par. iv. (1804) 235. t. 60 = **Rondeletia**, Linn. (Rubiac.).
buxifolia, Poit. l. c. = R. buxifolia.

STEVENSONIA, J. Dunc. Cat. Hort. Maurit. 87 (1863), nomen. *PALMEAE*, Benth. & Hook. f. iii. 908.
PHOENICOPHORIUM, H. Wendl. in Illustr. Hortic. xii. (1865) Misc. 5. t. 433.
STEPHENSONIA, Hort. ex Van Houtte, Fl. des Serres, xv. (1862–63) 177.
grandiflora, J. Dunc. Cat. Pl. Maurit. 87 = seq.
grandifolia, *J. Dunc. l. c.*, in errat.—Ins. Mascar.
viridifolia, J. Dunc.—Cf. Gard. Chron. (1870) 697; Baker, Fl. Maurit. 388 = Verschaffeltia splendida.

STEVIA, Cav. Ic. iv. 32. tt. 354–356 (1797). *COMPOSITAE*, Benth. & Hook. f. ii. 242.
MUSTELIA, Spreng. Nachr. i. Bot. Gart. Halle, 28 (1801); et in Trans. Linn. Soc. vi. (1802) 152. t. 13.
NOTHITES, Cass. in Dict. Sc. Nat. xxxv. 163 (1825).
XETOLIGUS, Rafin. New Fl. Am. iv. 74 (1836).
adenophora, *Lag. Gen. et Sp. Nov.* 27.—Chili.
alpina, *Griseb. in Goett. Abh.* xix. (1874) 165.—Reg. Argent.
amabilis, *Lemmon, ex A. Gray, in Proc. Am. Acad.* xix. (1884) 1.—Arizona.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 149 = salicifolia.
anisostemma, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 167.—Peruv.
arbutifolia, Willd. ex DC. Prod. v. 115 = Ageratum arbutifolium.
aristata, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 238.—Reg. Argent.
Arnottiana, Baker, in Mart. Fl. Bras. vi. ii. 201 = Gilliesii.
Aschenborniana, *Sch. Bip. in Linnaea*, xxv. (1852) 269.—Mexic.
Berlandieri, *A. Gray, in Torr. Bot. Mex. Bound.* 73, in Adnot.—Mexic.
boliviensis, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81; *et in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
breviaristata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 238.—Reg. Argent.
breviflora, Sch. Bip. in Linnaea, xxv. (1852) 291 = aristata?
callosa, Cerv. ex Loud. Hort. Brit. 335 (Quid?).—Mexic.
callosa, Nutt. in Journ. Acad. Philad. ii. (1821) 121 = Polypteris callosa.
calycina, DC. Prod. v. 124 = Eupatorium calyculatum.
camporum, *Baker, in Mart. Fl. Bras.* vi. ii. 202.—Bras.
canescens, H. B. & K. Nov. Gen. et Sp. iv. 143 = serrata.
caracasana, *DC. Prod.* v. 119.—Venezuela.
cathartica, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 55.—Peruv.
Chamaedrys, *Griseb. in Goett. Abh.* xxiv. (1879) 167.—Reg. Argent.
cinerascens, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 209.—Bras.
Clausseni, *Sch. Bip. ex Baker, l. c.* 203.—Bras.
clinopodia, *DC. Prod.* v. 116.—Mexic.
collina, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 458.—Bras.
compacta, *Benth. Pl. Hartw.* 197.—N. Granat.
conferta, DC. Prod. v. 116 = pilosa.
congesta, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = satureiaefolia.
connata, *Lag. Elench. Hort. Madr.* (1805); *ex ej. Gen. et Sp. Nov.* 27.—Mexic.
cordifolia, *Benth. Pl. Hartw.* 39.—Mexic.
coriacea, *Lag. Gen. et Sp. Nov.* 28.—Peruv.
coronata, Sch. Bip. in Bot. Zeit. iii. (1845) 155 = urticaefolia.
coronifera, *DC. Prod.* v. 115.—Mexic.
crenata, *Benth. Pl. Hartw.* 197.—Mexic.

STEVIA:—
crenata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 205 = menthaefolia, *Sch. Bip.*
crenulata, *Baker, l. c.* 211.—Bras.
cryptantha, *Baker, l. c.* 204.—Bras.
decussata, *Baker, l. c.* 203.—Bras.
dissitifolia, *DC. Prod.* v. 121.—Mexic.
dissoluta, *Schlecht. Hort. Hal.* 16.—Mexic.
divaricata, *DC. Prod.* v. 123.—Mexic.
dodecachaeta, *DC. l. c.* 122.—Peruv.
Ehrenbergiana, *Schlecht. in Linnaea*, xvi. (1842) 370.—Mexic.
elatior, *H. B. & K. Nov. Gen. et Sp.* iv. 144.—N. Granat.
elliptica, *Hook. & Arn. Bot. Beech. Voy.* 434.—Mexic.
elongata, *H. B. & K. Nov. Gen. et Sp.* iv. 144.—N. Granat.
enarthotricha, *Lag. Gen. et Sp. Nov.* 27.—Mexic.
erecta, Walp. Rep. ii. 548, sphalm. = ovata?
Eupatoria, [Ker-Gawl.] Bot. Reg. t. 93 = purpurea.
Eupatoria, *Willd. Sp. Pl.* iii. 1775.—Mexic.
fascicularis, *Less. in Linnaea*, v. (1830) 140.—Mexic.
fastigiata, *H. B. & K. Nov. Gen. et Sp.* iv. 148.—Mexic.
febrifuga, Moç. [& Sesse,] ex DC. Prod. v. 104 = Piqueria trinervia.
ferruginea, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. ii. 211 = S. Veronicae.
fruticosa, *Griseb. in Goett. Abh.* xxiv. (1879) 167.—Reg. Argent.
Gardneriana, *Baker, in Mart. Fl. Bras.* vi. ii. 207.—Bras.
Gilliesii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 251, in textu.—Bras.
glabra, Moç. [& Sesse,] ex DC. Prod. v. 117 = connata.
glandulifera, *Schlecht. Ind. Sem. Hort. Hal.* (1839) 19.—Mexic.
glandulosa, *Hook. & Arn. Bot. Beech. Voy.* 296.—Mexic.
glutinosa, *H. B. & K. Nov. Gen. et Sp.* iv. 148.—N. Granat.
gracilipes, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 210 = Lundiana.
grandidentata, Sch. Bip. ex Klatt, in Leopoldina, xx. (1884) 75 = glandulifera.
grandidentata, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81; *et in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
gratioloides, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = S. Veronicae.
Haenkeana, *DC. Prod.* v. 122.—Mexic.
heptachaeta, *DC. l. c.*—Bras.
hirsuta, *DC. l. c.* 121.—Mexic.
hirsuta, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 239 = satureiaefolia.
hirtiflora, *Sch. Bip. in Linnaea*, xxv. (1852) 274.—Mexic.
hispidula, *DC. Prod.* v. 122.—Mexic.
hymenopappa, Moç. [& Sesse,] ex DC. Prod. v. 117 = lucida.
hyptifolia, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 434 = collina.
hyssopifolia, Cav. Desc. 190 = S. Eupatoria.
hyssopifolia, H. B. & K. Nov. Gen. et Sp. iv. 140 = laxiflora.
hyssopifolia, *Phil. Fl. Atac.* 29.—Chili.
hyssopifolia, Sims, Bot. Mag. t. 1861 = paniculata.
imbricata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 458 = Dissothrix Gardneri.
incanescens, Lag. Gen. et Sp. Nov. 27 = serrata, microcephala, uniaristata.
involucrata, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 211.—Bras.
ivaefolia, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 137.—Mexic.; N. Granat.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* iv. 144.—Mexic.
Karwinskyana, *Steud. Nom.* ed. II. ii. 641.—Mexic.
lanceolata, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = Gilliesii.
lanceolata, *Lag. Gen. et Sp. Nov.* 26.—Mexic.
lanceolata, Moç. [& Sesse,] ex DC. Prod. v. 116 = subpubescens.
lanceolata, Walp. in Linnaea, xiv. (1840) 319 = Karwinskyana.
latifolia, *Benth. Pl. Hartw.* 40.—Mexic.

STEVIA:—
lavandulaefolia, Willd. ex DC. Prod. v. 125 = Palafoxia linearis.
laxa, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = satureiaefolia.
laxiflora, *DC. Prod.* v. 122.—Mexic.
Lemmoni, *A. Gray, in Proc. A Acad.* xvii. (1882) 204.—Arizona.
leptophylla, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 205.—Bras.
leucantha, *Schlecht. Hort. Hal.* 16.—Mexic.
leuconeura, *DC. Prod.* v. 121.—Mexic.
Liebmannii, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 75.—Mexic.
linariaefolia, DC. Prod. v. 123 = satureiaefolia.
linearifolia, Steud. Nom. ed. II. ii. 641 = praec.
linearifolia, *Walp. in Linnaea*, xiv. (1840) 320.—Mexic.
linearis, Gill. ex Baker, in Mart. Fl. Bras. vi. ii. 208 = satureiaefolia.
linearis, Willd. Sp. Pl. iii. 1774 = Palafoxia linearis.
linoides, *Sch. Bip. in Linnaea*, xxv. (1852) 284.—Mexic.
longifolia, Moç. [& Sesse,] ex DC. Prod. v. 118 = viminea.
longiseta, Moç. [& Sesse,] l. c. 121 = leuconeura.
Lorentzii, *Griseb. in Goett. Abh.* xix. (1874) 164 (err. typ. 416).—Reg. Argent.
lucida, *Lag. Gen. et Sp. Nov.* 28.—Mexic.
Lundiana, *DC. Prod.* v. 122.—Bras.
macella, A. Gray, Pl. Wright. ii. 70 = micrantha.
Mandonii, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 81; *et in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
megapotamica, DC. Prod. v. 123 = satureiaefolia.
melissaefolia, *Sch. Bip. in Linnaea*, xxv. (1852) 291.—Peruv.
menthaefolia, *Phil. Fl. Atac.* 29.—Chili.
menthaefolia, *Sch. Bip. in Linnaea*, xxv. (1852) 282.—Bras.
micrantha, *Lag. Elench. Hort. Matr.* (1805); *ex ej. Gen. et Sp. Nov.* 27.—Mexic.
microcephala, *DC. Prod.* v. 118.—Mexic.
microchaeta, *Sch. Bip. in Linnaea*, xxv. (1852) 291.—Mexic.
micropappa, Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 81 = grandidentata.
microphylla, *H. B. & K. Nov. Gen. et Sp.* iv. 140.—Mexic.
minor, *Griseb. in Goett. Abh.* xix. (1874) 166.—Reg. Argent.
mollis, *Schrad. ex DC. Prod.* v. 120.—Hab.?
monardaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 147.—Mexic.
multiaristata, Spreng. Syst. iii. 449 = satureiaefolia.
multifida, [Moç. & Sesse,] ex DC. Prod. v. 121 = trifida.
myriadenia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. ii. 206.—Bras.
nepetaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 146.—Am. austr.
nervosa, *DC. Prod.* v. 117.—Mexic.
nitida, *Walp. in Linnaea*, xiv. (1840) 320.—Mexic.
oaxacana, Sch. Bip. ex Klatt, in Leopoldina, xx. (1884) 75 = lucida.
oligocephala, *DC. Prod.* v. 117.—Bras.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 115.—Bras.
origanifolia, *Walp. in Linnaea*, xiv. (1840) 321.—Mexic.
origanoides, *H. B. & K. Nov. Gen. et Sp.* iv. 147.—Mexic.
ovata, Dum.-Cours. Bot. Cult. ed. II. iv. 93 = S. Eupatoria, salicifolia.
ovata, *Lag. Elench. Hort. Matr.* (1805); *ex ej. Gen. et Sp. Nov.* 27.—Mexic.
ovata, Rafin. New Fl. Am. iv. 73 (Quid?).—Am. bor.
oxylaena, *DC. Prod.* v. 123.—Bras.
pallida, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 206 = collina.
paniculata, *Lag. Elench. Hort. Matr.* (1805); *ex ej. Gen. et Sp. Nov.* 27.—Mexic.
paniculigera, Mart. Herb. Fl. Bras. n. 772; et ex Baker, in Mart. Fl. Bras. vi. ii. 201 = urticaefolia.
pedata, Cav. Ic. iv. 33. t. 356 = Florestina pedata.
petiolata, *Sch. Bip. in Linnaea*, xxv. (1852) 291.—Peruv.
pilosa, *Lag. Gen. et Sp. Nov.* 26.—Mexic.

STEVIA:—
Plummerae, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 204.—Arizona.
podocephala, *DC. Prod.* v. 121.—Mexic.
podocephala, Schau. in Linnaea, xix. (1847) 718 = Aschenborniana.
Pohliana, *Baker, in Mart. Fl. Bras.* vi. II. 211.—Bras.
polycephala, *Baker, l. c.* 207.—Bras.
polycephala, *Bertol. Fl. Guatimal.* 32.—Guatemal.
polyphylla, *DC. Prod.* v. 123.—Am. austr.
puberula, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = Gilliesii.
puberula, *Hook. Misc. Bot.* ii. (1831) 225.—Peruv.
pubescens, *Lag. Gen. et Sp. Nov.* 26.—Mexic.
pubescens, Moç. [& Sesse,] ex DC. Prod. v. 119 = nepetaefolia.
punctata, Pers. Syn. ii. 403 = S. Eupatoria.
punctata, Sch. Bip. in Linnaea, xxv. (1852) 286 = serrata.
purpurea, Lag. Gen. et Sp. Nov. 26 = laxiflora.
purpurea, *Pers. Syn.* ii. 402.—Mexic.
pyrolaefolia, *Schlecht. in Linnaea*, xvi. (1842) 326.—Mexic.
quitensis, *H. B. & K. Nov. Gen. et Sp.* iv. 145.—Ecuador.
ranunculoides, Hemsl. Biol. Centr. Am. Bot. ii. 89 = seq.
rapunculoides, *DC. Prod.* v. 124.—Mexic.
reglensis, *Benth. Pl. Hartw.* 40.—Mexic.
Regnellii, Sch. Bip. in Linnaea, xxii. (1849) 572; xxv. (1852) 272 = S. Veronicae.
resinosa, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 457.—Bras.
rhombifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 143.—Mexic.
Riedelii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 204.—Bras.
salicifolia, *Cav. Ic.* iv. 32. t. 354.—Mexic.
salviaefolia, Hort. ex G. Don, in Loud. Hort. Brit. 335 (Quid?).—Mexic.
satureiaefolia, *Sch. Bip. in Linnaea*, xxv. (1852) 291.—Bras.
satureifolia, Cav. in Anal. Cienc. Nat. vi. (1802) 317 = Mikania amara.
scabrella, *Benth. Pl. Hartw.* 19.—Mexic.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 298.—Mexic.
Selloi, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 209 = oxylaena.
semperflorens, Tenore, ex DC. Prod. v. 117 = salicifolia.
sempervirens, Steud. Nom. ed. II. ii. 641, sphalm. = praec.
serrata, *Cav. Ic.* iv. 33. t. 355.—Am. austr.
sphacelata, Nutt. ex Torr. in Ann. Lyc. N. York, ii. (1828) 214 = Polypteris Hookeriana.
stenocephala, Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 81 = grandidentata.
stenophylla, *A. Gray, in Proc. Am. Acad.* xv. (1880) 25.—Mexic.
stricta, Hornem. Hort. Hafn. ii. 792 = purpurea.
suaveolens, Lag. Gen. et Sp. Nov. 27 = nepetaefolia.
subhirsuta, Lag. ex Reichb. Ic. Bot. Exot. ii. 33. t. 188 = purpurea?
suboctoaristata, *Lag. Gen. et Sp. Nov.* 27.—Peruv.
subpubescens, *Lag. l. c.* 28.—Mexic.
tenella, Moç. [& Sesse,] ex DC. Prod. v. 121 = micrantha.
tenuifolia, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = satureiaefolia.
tenuifolia, *Phil. in Anal. Univ. Chil.* (1862) II. 397; *et in Linnaea*, xxxiii. (1864–65) 128.—Chili.
tenuis, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 239 = S. Veronicae.
ternifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 146.—Mexic.
tomentosa, *H. B. & K. l. c.* 145.—Mexic.
tomentosa, Schau. in Linnaea, xix. (1847) 719 = nepetaefolia.
trachelioides, *DC. Prod.* v. 115.—Mexic.
trachelioides, Hook. Bot. Mag. t. 3856 = glandulifera.
trichopoda, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 64.—Mexic.
trifida, *Lag. Gen. et Sp. Nov.* 27.—Mexic.
triflora, *DC. Prod.* v. 115.—Mexic.
umbellata, Alam. ex DC. l. c. 118 = viminea.
uniaristata, *DC. l. c.* 120.—Mexic.

STEVIA:—
urticaefolia, *Thunb. Pl. Bras. Dec.* i. 13.—Bras.
vaga, *Griseb. in Goett. Abh.* xix. (1874) 165.—Reg. Argent.
Veronicae, *DC. Prod.* v. 123.—Bras.
verticillata, *Schlecht. in Linnaea*, xvi. (1842) 372.—Bras.
vigintiseta, *DC. Prod.* v. 123.—Mexic.
villosa, Moç. [& Sesse,] ex DC. l. c. 122 = viscida.
viminea, *Schrad. Hort. Gotting.* (1832); *DC. l. c.* 117.—Mexic.
violacea, Cerv. ex Loud. Hort. Brit. Suppl. ii. 675 (Quid?).—Mexic.
virgata, H. B. & K. Nov. Gen. et Sp. iv. 142 = serrata.
viscida, *H. B. & K. l. c.* 140.—Mexic.
viscosa, Hort. ex Steud. Nom. ed. II. ii. 642 = viscida.

STEVOGTIA, Neck. Elem. ii. 23 (1790) = **Convolvulus**, Tourn.
platycarpos, Rafin. Fl. Tellur. iv. 70 = Phacelia platycarpa.

STEWARTIA, Linn. in Act. Soc. Upsal. (1741) 79, sphalm. = **Stuartia**, Linn. (Ternstr.).

STHAELINA, Lag. Gen. et Sp. Nov. 24 (1816), err. typ. = **Staehelina**, Linn. (Compos.).

STIBADOTHECA, Klotzsch, in Abh. Akad. Berl. 1854 (1855), 128. t. 12 A = **Begonia**, Tourn.
ferruginea, Klotzsch, l. c. 130 = B. ferruginea.
magnifica, Klotzsch, l. c. 129 = B. ferruginea.
trachyptera, Klotzsch, l. c. 131 = B. trachyptera.
Urticae, Klotzsch, ex A. DC. in DC. Prod. xvi. I. 274 = B. trachyptera.

STIBAS, Comm. ex DC. Prod. vii. 338 (1839) = **Levenhookia**, R. Br. (Campan.).

STICHONEURON, Hook. f. in Benth. & Hook. f. Gen. iii. 747 (1883). *ROXBURGHIACEAE*, Benth. & Hook. f. l. c.
membranaceum, *Hook. f. & Thoms.* (in Herb. Kew.).—Ind. or.

STICHOPHYLLUM, Phil. Fl. Atac. 19. t. 1 D (1860) = **Pycnophyllum**, Remy (Caryophyll.).
bryoides, Phil. l. c. = P. bryoides.

STICHORCHIS, Thou. Orch. Isles Austr. Afr. Tabl. 1er (1822) = **Liparis**, Rich. (Orchid.).

STICKMANNIA, Neck. Elem. iii. 171 (1790) = **Dichorisandra**, Mikan (Commel.).
africana, Rafin. Fl. Tellur. iii. 69 = Commelina africana.
guyanensis, Rafin. l. c. = Commelina nudiflora.
longicollis, Rafin. l. c. = Commelina nudiflora.
mollis, Rafin. l. c. = D. villosula.

STICTOPHYLLUM, Edgew. in Proc. Linn. Soc. i. (1845) 253; et in Trans. Linn. Soc. xx. (1846) 78 = **Tricholepis**, DC. (Compos.).
glabrum, Edgew. in Trans. Linn. Soc. xx. (1846) 78 = T. Stictophyllum.

STIEFIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 200; Phil. Bot. ii. 40 (1791) = **Salvia**, Tourn. (Labiat.).

STIFFTIA, Mikan, Del. Bras. i. 1. t. 1 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 491.
ARISTOMENIA, Vell. Fl. Flum. 345 (1825).
AUGUSTA, Leandr. in Denkschr. Akad. Muench. vii. (1819) 235. t. 14.
GONGYLOLEPIS, Schomb. in Linnaea, xx. (1847) 759.
MOCINIA, DC. Prod. vii. 26 (1838).
SANHILARIA, Leandr. ex DC. l. c. (1838).
Benthamiana, *Baker, in Mart. Fl. Bras.* vi. III. 352.—Guian.
chrysantha, *Mikan, Del. Bras.* i. 1. t. 1.—Bras.
condensata, *Baker, in Mart. Fl. Bras.* vi. III. 351.—Guian.
insignis, Regel, Cat. Pl. Hort. Aksakov. 135 = chrysantha?
Martiana, *Baker, in Mart. Fl. Bras.* vi. III. 351.—Bras.
parviflora, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 291.—Bras.

STIFTIA, Cass. in Dict. Sc. Nat. xlvii. 499, 511 (1827), sphalm. = praec.

STIFTIA, Pohl, ex Nees, in DC. Prod. xi. 70 (1847) = **Ebermaiera**, Nees (Acanth.).
elegans, Pohl, ex Nees, l. c. 72 = E. elegans.

STIGMANTHUS, Lour. Fl. Cochinch. 146 (1790). Genus dubium *RUBIACEARUM*.—Cf. Benth. & Hook. f. ii. 29.
STIGMATANTHUS, Roem. & Schult. Syst. v. p. xv. 225 (1819).
cymosus, *Lour. Fl. Cochinch.* 146.—Cochinch.

STIGMAPHYLLON, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 48. tt. 170, 171 (1832). *MALPIGHIACEAE*, Benth. & Hook. f. i. 257.
acuminatum, *A. Juss. l. c.* 58.—Bras.
adenodon, *A. Juss. in Arch. Mus. Par.* iii. (1843) 360.—Ind. occ.
affine, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 57.—Bras.
alternans, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 316.—N. Granat.
alternifolium, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 54 = tomentosum.
angulosum, A. Juss. in Arch. Mus. Par. iii. (1843) 362.—Ind. occ.
angustifolium, Griseb. in Mart. Fl. Bras. xii. I. 41 = seq.
angustilobum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 53.—Bras.
anomalum, A. Juss. in Arch. Mus. Par. iii. (1843) 363 = urenaefolium.
aristatum, *Lindl. Bot. Reg.* t. 1659.—Am. austr.
auriculatum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 48.—Bras.
bogotense, *Triana & Planch. in Ann. Sc. Nat* Sér. IV. xviii. (1862) 320.—N. Granat.
brachiatum, *Planch. & Triana, l. c.* 316.—N. Granat.
cardiophyllum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 374.—Bras.
ciliatum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 49.—Bras.
convolvulaefolium, *A. Juss. in Arch. Mus. Par.* iii. (1843) 372.—Guian.
dichotomum, Griseb. in Linnaea, xiii. (1839) 207 = rotundifolium.
diversifolium, *A. Juss. in Arch. Mus. Par.* iii. (1843) 381.—Ind. occ.
echitoides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 318.—N. Granat.
ellipticum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 377.—Mexic.
emarginatum, *A. Juss. l. c.* 382.—Ind. occ.
Faustinum, *Wright, in Sauv. Fl. Cub.* 16.—Cuba.
Finlayanum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 373.—Ind. occ.
fulgens, *A. Juss. l. c.* 370.—Guian.
Gaudichaudianum, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 50 = vitifolium.
Gayanum, Griseb. in Linnaea, xiii. (1839) 211 = affine.
Gayanum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 56.—Bras.
hastatum, Griseb. in Linnaea, xiii. (1839) 208 = affine.
heterophyllum, *Hook. Bot. Mag.* t. 4014.—Reg. Argent.
Humboldtianum, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 56 = tomentosum.
Humboldtianum, Seem. Bot. Voy. Herald, 91 = retusum.
hypargyreum, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 318.—Panama.
hypoleucum, *Miq. in Linnaea*, xviii. (1844) 51.—Guian.
ipomoeoides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 317.—N. Granat.
irregulare, A. Juss. in Arch. Mus. Par. iii. (1843) 366 = affine.
jatrophaefolium, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 51.—Bras.
lacunosum, A. Juss. in Arch. Mus. Par. iii. (1843) 370 = puberulum.
Lalandianum, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 58; et l. c. 381 = acuminatum.
latifolium, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 128.—Guian.

STIGMAPHYLLON:—
Lindenianum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 362.—Mexic.
lineare, *Wright, ex Griseb. Cat. Pl. Cub.* 43.—Cuba.
littorale, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 55.—Bras.
macropodum, A. Juss. in Arch. Mus. Par. iii. (1843) 369 = tomentosum.
malpighioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 582.—Mexic.
Martianum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 371.—Bras.
megacarpum, Griseb. in Linnaea, xiii. (1839) 209 = tomentosum.
microphyllum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 168.—Cuba.
mucronatum, A. Juss. in Arch. Mus. Par. iii. (1843) 377 = ellipticum.
multilobum, *Miq. in Linnaea*, xix. (1847) 433.—Bras.
obtusum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 582.—Cuba.
palmatum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 362.—Ind. occ.
Paralias, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 59 = Brachypterys australis.
periplocaefolium, *A. Juss. in Arch. Mus. Par.* iii. (1843) 380.—Ind. occ.
puberulum, *Griseb. in Linnaea*, xiii. (1839) 205.—Bras.
puberum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 376.—Ind. occ.
purpureum, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 128 = fulgens.
repandum, *Griseb. in Linnaea*, xiii. (1839) 202.—Bras.
reticulatum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 378.—Cuba.
retusum, *Griseb. & Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 45.—Panama.
rhombifolium, *Wright, in Sauv. Fl. Cub.* 17.—Cuba.
Richardianum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 372.—Guian.
rotundifolium, *A. Juss. l. c.* 373.—Am. austr.
sagittatum, *A. Juss. l. c.* 362.—Ind. occ.
Sagraeanum, *A. Juss. l. c.* 379.—Cuba.
Saltzmannii, *A. Juss. l. c.* 365.—Am. austr.
sericeum, Wright, ex Griseb. Cat. Pl. Cub. 43 = diversifolium.
sinuatum, *A. Juss. in Arch. Mus. Par.* iii. (1843) 361.—Guian.
strigosum, *A. Juss. l. c.* 368.—Peruv.
ternatum, A. Juss. l. c. 376 = ellipticum.
tomentosum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 53.—Bras.
umbellatum, *Juss. ex Steud. Nom.* ed. II. ii. 642 (Quid ?).—Bras.
urenaefolium, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 52.—Bras.
velutinum, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 320.—N. Granat.
vitifolium, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 50.—Bras.

STIGMAROSA, Hook. f. & Thoms. in Fl. Brit. Ind. i. 193 (1872), sphalm. = seq.

STIGMAROTA, Lour. Fl. Cochinch. 633 (1790) = **Flacourtia**, Comm. (Bixin.).
africana, Lour. l. c. = F. Ramontchi.
edulis, Blanco, Fl. Filip. ed. II. 560 = F. sepiaria.
Jangomas, Lour. Fl. Cochinch. 634 = F. cataphracta.

STIGMATANTHUS, Roem. & Schult. Syst. v. p. xv. 220 (1819) = **Stigmanthus**, Lour. (Rubiac.).

STIGMATOCOCCA, Willd. ex Schult. Mant. iii. 3, 55 (1827). *INCERT. SEDIS.*—Cf. Benth. & Hook. f. Gen. ii. 888.
solanacea, *Willd. l. c.*—Bras.

STIGMATOPHYLLON, Meissn. Gen. 55 (1837) = **Stigmaphyllon**, A. Juss. (Malpig.).

STIGMATOPHYLLUM, Spach, Hist. Vég. Phan. iii. 153 (1834) = praec.

STIGMATOTHECA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 255 (1836–46) = **Chrysanthemum**, Tourn. (Compos.).
lacera, Sch. Bip. l. c. 257 = C. Broussonetii.
pinnatifida, Sch. Bip. l. c. 255 = C. pinnatifidum.

STILAGINELLA, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 240 = **Hyeronima**, Allem. (Euphorb.).
amazonica, Tul. l. c. 241 = H. alchorneoides.
Benthamii, Tul. l. c. 247 = H. oblonga.
Blanchetiana, Tul. l. c. 249 = H. Blanchetiana.
clusioides, Tul. l. c. 245 = H. clusioides.
ferruginea, Tul. l. c. 250 = H. ferruginea.
laxiflora, Tul. l. c. 244 = H. laxiflora.
oblonga, Tul. l. c. 248 = H. oblonga.
scabrida, Tul. l. c. 242 = H. scabrida.

STILAGO, Linn. Mant. i. 16 (1767) = **Antidesma**, Burm. (Euphorb.).
Bunius, Linn. l. c. 122 = A. Bunius.
diandra, Roxb. Pl. Corom. ii. t. 166 = A. diandrum.
lanceolaria, Roxb. Hort. Beng. 71; Fl. Ind. iii. 760 = A. diandrum.
tomentosa, Roxb. ll. cc. 71; 757 = A. Roxburghii.

STILBANTHUS, Hook. f. in Hook. Ic. Pl. t. 1286 (1879). *AMARANTACEAE*, Benth. & Hook. f. iii. 35.
scandens, *Hook. f. in Hook. Ic. Pl.* t. 1286.—Reg. Himal.

STILBE, Berg. Desc. Pl. Cap. 30. t. 4 (1767). *VERBENACEAE*, Benth. & Hook. f. ii. 1138.
LUEHEA, F. W. Schmidt, Neue u. Selt. Pfl. 23 (1793).
albiflora, *E. Mey. Comm. Pl. Afr. Austr.* 279.—Afr. austr.
cernua, Linn. f. Suppl. 441 = Campylostachys cernua.
ericoides, Lam. ex Poir. Encyc. Suppl. v. 251 = albiflora.
ericoides, *Linn. Mant.* ii. 305; *Thunb. Prod. Pl. Cap.* 29.—Afr. austr.
myrtifolia, Poir. Encyc. Suppl. v. 252 = Lonchostoma obtusiflorum.
phylicoides, *A. DC. in DC. Prod.* xii. 606.—Afr. austr.
phylicoides, Licht. ex Walp. Rep. iv. 173 (Quid ?).—Hab. ?
pinastra, Linn. Mant. ii. 305 = vestita.
procumbens, Spreng. Syst. i. 418 = Cephalanthus procumbens.
serrulata, Hochst. in Flora, xxv. (1842) 229 = Eurylobium serrulatum.
vestita, *Berg. Desc. Pl. Cap.* 30.—Afr. austr.
virgata, Poir. Encyc. Suppl. v. 252 = ericoides.

STILBOCARPA, A. Gray, Bot. U. St. Expl. Exped. i. 714 (1854). *ARALIACEAE*, Benth. & Hook. f. i. 935.
STYLBOCARPA, Decne. & Planch. in Rev. Hortic. (1854) 105, sphalm.
Lyallii, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 336.—N. Zel.
polaris, *A. Gray, Bot. U. St. Expl. Exped.* i. 714.—N. Zel.

STILINGIA, Rafin. Atl. Journ. 146 (1832), sphalm. = **Stillingia**, Linn. (Euphorb.).

STILLENGIA, Torr. in Emory, Notes Mil. Reconnois. 151 (1848), sphalm. = praec.

STILLINGFLEETIA, Boj. Hort. Maurit. 284 (1837) = **Sapium**, P. Br. (Euphorb.).
sebifera, Boj. l. c. = Sapium sebiferum.

STILLINGIA, Linn. Mant. i. 19 (1767). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 334.
DITRISYNIA, Rafin. Sylva Tellur. 64 (1838).
GYMNOSTILLINGIA, Muell. Arg. in Linnaea, xxxii. (1863) 89.
SEBORIUM, Rafin. Sylva Tellur. 63 (1838).
acutifolia, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 135.—Mexic.
africana, Baill. Adansonia, iii. (1862–63) 163 = Excoecaria africana.
Agallocha, Muell. Arg. in DC. Prod. xv. II. 1161, 1220 = Excoecaria Agallocha.
angustifolia, *Engl. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 154.—Texas.

STILLINGIA:—
annua, *Muell. Arg. in DC. Prod.* xv. II. 1160.—Mexic.
appendiculata, *Muell. Arg. in Linnaea*, xxxii. (1863) 87.—Mexic.
aquatica, *Chapm. Fl. S. U. St.* 405.—Florida.
asperococca, Muell. Arg. in DC. Prod. xv. II. 1161, 1175 = Sebastiania Chamaelea.
aucuparia, Muell. Arg. l. c. 1161, 1206 = Sapium Aucuparium.
baccata, Muell. Arg. l. c. 1161, 1211 = Sapium baccatum.
bahiensis, Baill. Adansonia, v. (1864–65) 329 = Sebastiania bahiensis.
bidentata, Baill. l. c. 324 = Sebastiania virgata.
biglandulosa, Baill. l. c. 320 = Sapium Aucuparium.
bingyrica, Muell. Arg. in DC. Prod. xv. II. 1161, 1216 = Sapium indicum.
brasiliensis, Muell. Arg. l. c. 1161, 1191 = Maprounea brasiliensis.
brevifolia, Baill. Adansonia, v. (1864–65) 328 = Sebastiania brevifolia.
campestris, Muell. Arg. in DC. Prod. xv. II. 1161, 1169 = Sebastiania corniculata.
Chamaelea, Muell. Arg. l. c. 1161, 1175 = Sebastiania Chamaelea.
cochinchinensis, Baill. Adansonia, i. (1860–61) 351 = Excoecaria Loureiroana.
Commersoniana, Baill. l. c. v. (1864–65) 330 = Gymnanthes marginata.
concolor, Baill. l. c. 327 = Actinostemon caribaeus.
coriacea, Baill. l. c. 323 = Sebastiania marginata.
corniculata, Baill. Etud. Gén. Euph. Atlas, 17 = Sebastiania corniculata.
cremostachys, Baill. Adansonia, v. (1864–65) 322 = Sapium Aucuparium.
crotonoides, Muell. Arg. in DC. Prod. xv. II. 1161, 1171 = Sebastiania corniculata.
cubana, Muell. Arg. l. c. 1161, 1212 = Excoecaria cubensis.
daphniphylla, Baill. Adansonia, v. (1864–65) 326 = Sebastiania daphniphylla.
desertorum, Klotzsch, in Wiegm. Archiv, vii. (1841) 183, nomen = Sebastiania brasiliensis.
dichotoma, *Muell. Arg. in Linnaea*, xxxii. (1863) 88.—Bras.
discolor, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 1 = Sapium discolor.
discolor, Muell. Arg. in DC. Prod. xv. II. 1161, 1885 = Sebastiania discolor.
diversifolia, Miq. Fl. Ind. Bat. Suppl. 461 = Excoecaria diversifolia.
dracunculoides, Baill. Adansonia, v. (1864–65) 321 = Sapium Aucuparium.
eglandulosa, A. Rich. Fl. Cub. Fanerog. ii. 202 = Bonania adenodon.
elliptica, *Baill. Adansonia*, iii. (1862–63) 162.—Afr. austr.
frutescens, Bosc, ex Steud. Nom. ed. I. 815 = seq.
fruticosa, Spreng. Syst. iii. 805 = Sebastiania ligustrina.
Gaudichaudii, Baill. Adansonia, v. (1864–65) 332 = Sebastiania Gaudichaudii.
glabrata, Baill. Etud. Gén. Euph. Atlas, 17 = Sebastiania multiramea.
glandulosa, Domb. ex Baill. l. c. = Adenopeltis Colliguaya.
Goudotiana, Baill. Adansonia, ii. (1861–62) 30 = Excoecaria Goudotiana.
guineensis, Benth. in Hook. Niger Fl. 501 = Excoecaria guineensis.
guyanensis, Baill. Adansonia, v. (1864–65) 332 = Maprounea guianensis.
hastata, Baill. l. c. 325 = Sebastiania ditassoides.
heterodoxa, *Muell. Arg. in Linnaea*, xxxii. (1863) 89.—Bras.
Hilariana, Baill. Adansonia, v. (1864–65) 332 = Maprounea brasiliensis.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 116 = Sapium indicum, Excoecaria acerifolia.
Hippomane, Muell. Arg. in DC. Prod. xv. II. 1161, 1204 = Sapium Aucuparium.
hypoleuca, Baill. Adansonia, v. (1864–65) 330 = Gymnanthes hypoleuca.
indica, Muell. Arg. in DC. Prod. xv. II. 1161, 1216 = Sapium indicum.
integerrima, *Baill. Adansonia*, iii. (1862–63) 162.—Afr. austr.

STILLINGIA :—
jacobinensis, Baill. Adansonia, v. (1864–65) 329 = Sebastiania jacobinensis.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 145 = Excoecaria japonica.
lanceolaria, Miq. Fl. Ind. Bat. Suppl. 461 = Excoecaria lanceolaria.
lanceolata, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 176.—Am. bor.
Lastellei, Muell. Arg. in DC. Prod. xv. II. 1162, 1218 = Excoecaria Lastellei.
Laureola, Baill. Adansonia, v. (1864–65) 327 = Sebastiania Laureola.
laurifolia, A. Rich. Fl. Cub. Fanerog. ii. 201 = Sapium Laurocerasum.
Laurocerasus, Baill. Adansonia, i. (1860–61) 351 = Sapium Laurocerasum.
ligustrina, Michx. Fl. Bor. Am. ii. 213 = Sebastiania ligustrina.
linearifolia, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 297.—Calif.
lineata, *Muell. Arg. in DC. Prod.* xv. II. 1157.—Ind. Maurit.
Luschnathiana, Muell. Arg. l. c. 1162, 1177 = Sebastiania multiramea.
macrantha, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 135.—Mexic.
madagascariensis, Baill. Adansonia, ii. (1861–62) 30 = Excoecaria madagascariensis.
marginata, Baill. l. c. v. (1864–65) 321 = Sapium marginatum.
mauritiana, Baill. l. c. ii. (1861–62) 27 = obtusifolia.
melanosticta, *Baill. l. c.* i. (1860–61) 285.—Madag.
multiramea, Muell. Arg. in DC. Prod. xv. II. 1162, 1179 = Sebastiania multramea.
myrtilloides, Baill. Adansonia, v. (1864–65) 323 = Sebastiania daphnoides.
nervosa, Baill. l. c. 328 = Sebastiania nervosa.
nutans, Geisel. Crot. Monog. 80 = Homalanthus nutans.
nutans, Vahl, ex Spreng. Syst. iii. 805.—Ins. Amicor.
obovata, Baill. Adansonia, v. (1864–65) 321 = Sapium obovatum.
obtusifolia, Muell. Arg. in DC. Prod. xv. II. 1157 = lineata.
oppositifolia, *Baill. Etud. Gén. Euph. Atlas*, 13.—Bras.
oppositifolia, Muell. Arg. in DC. Prod. xv. II. 1162, 1219 = Excoecaria crenulata.
pachystachys, Muell. Arg. in Linnaea, xxxii. (1863) 102 = Sebastiania pachystachys.
pacifica, *Muell. Arg. in DC. Prod.* xv. II. 1156.—Ins. Fiji.
paniculata, Miq. Fl. Ind. Bat. Suppl. 461 = Sapium baccatum.
patula, Muell. Arg. in DC. Prod. xv. II. 1162, 1172 = Sabastiania corniculata.
paucidentata, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 298.—Am. bor.
phyllanthiformis, Baill. Adansonia, v. (1864–65) 331 = Sebastiania Schottiana.
populnea, Geisl. Crot. Monog. 80 = Homalanthus Leschenaultianus.
prostrata, Baill. Adansonia, v. (1864–65) 324 = Sebastiania corniculata.
prunifolia, Baill. ex Muell. Arg. in DC. Prod. xv. II. 1162 = Sapium Aucuparium.
pteroclada, Baill. Adansonia, v. (1864–65) 329 = Sebastiania pteroclada.
ramosissima, Baill. l. c. 328 = Sebastiania brasiliensis.
rigida, Baill. l. c. 330 = Sebastiania rigida.
salicifolia, Klotzsch, ex Baill. Adansonia, v. (1864–65) 320 = Sapium Aucuparium.
salicifolia, Rafin. Atl. Journ. 146 (Quid ?).—Am. bor.
sanguinolenta, *Muell. Arg. in Linnaea*, xxxii. (1863) 88.—Mexic.
saxatilis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 539.—Bras.
Schottiana, Baill. Adansonia, v. (1864–65) 331 = Sebastiania Schottiana.
sebifera, Michx. Fl. Bor. Am. ii. 213 = Sapium sebiferum.
Sellowiana, Muell. Arg. in DC. Prod. xv. II. 1162, 1169 = Sebastiania corniculata.
serrata, Baill. Adansonia, v. (1864–65) 329 = Sebastiania serrata.
serrulata, Muell. Arg. in DC. Prod. xv. II. 1162, 1167 = Sebastiania serrulata.
sinensis, Muell. Arg. l. c. 1162, 1210 = Sapium sebiferum.

STILLINGIA :—
spinulosa, *Torr. in Emory, Notes Mil. Reconnois.* 151. —Am. bor.
stipulacea, Baill. Adansonia, v. (1864–65) 325 = Sebastiania stipulacea.
sylvatica, *Linn. Mant.* i. 126.—Am. bor.
tanguinia, Baill. Adansonia, ii. (1861–62) 28 = lineata.
terminalis, *Baill. l. c.* 29.—Madag.
Thuarsiana, Baill. l. c. 28.—Afr. trop.
Torreyana, S. Wats. in Proc. Am. Acad. xiv. (1879) 298 = Sebastiania Treculiana ?
trinervia, Baill. Adansonia, v. (1864–65) 328 = Sebastiania trinervia.
velutina, Muell. Arg. in DC. Prod. xv. II. 1162, 1168 = Sebastiania corniculata.
virgata, Muell. Arg. l. c. 1162, 1216 (= *Excoecaria virgata*).—Java.
Weddelliana, Baill. Adansonia, v. (1864–65) 329 = Sebastiania Weddelliana.
Widgreni, Baill. l. c. 326 = Sebastiania Widgreni.
ypanemensis, Baill. l. c. 330 = Sebastiania ypanemensis.
zelayensis, *Muell. Arg. in Linnaea*, xxxii. (1863) 87. —Mexic.

STILPNOGYNE, DC. Prod. vi. 293 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 444.
bellidioides, *DC. l. c.; et in Deless. Ic. Sel.* iv. 24. t. 54 (1839).—Afr. austr.

STILPNOPAPPUS, Mart. ex DC. Prod. v. 75 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 232.
LASIOCARPHUS, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 138 (1873).
STROPHOPAPPUS, DC. Prod. v. 75 (1836).
XIPHOCHAETA, Poepp. & Endl. Nov. Gen. et Sp. iii. 44. t. 250 (1842).
agrestis, Gardn. ex Baker, in Mart. Fl. Bras. vi. II. 136 = pratensis.
bicolor, *Mart. ex Baker, l. c.* 140.—Bras.
dentatus, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 229 = pratensis.
emarginatus, *Gardn. l. c.* vi. (1847) 424.—Bras.
ferrugineus, *Baker, in Mart. Fl. Bras.* vi. II. 138.—Bras.
glomeratus, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 423.—Bras.
patulus, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 109.—Bras.
Pohlii, *Baker, in Mart. Fl. Bras.* vi. II. 139.—Bras.
pratensis, *Mart. ex DC. Prod.* v. 76.—Bras.
procumbens, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 229.—Bras.
Regnellii, *Baker, in Mart. Fl. Bras.* vi. II. 139.—Bras.
Riedelianus, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 424.—Bras.
rufulus, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 138 = patulus.
scaposus, *DC. Prod.* v. 76.—Bras.
speciosus, Baker, in Mart. Fl. Bras. vi. II. 138 = patulus.
suffruticosus, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 228.—Bras.
tomentosus, Gardn. l. c. 230 = pratensis.
tomentosus, *Mart. ex DC. Prod.* v. 76.—Bras.
trichospiroides, *Mart. l. c.*—Bras.
villosus, *Mart. ex Baker, in Mart. Fl. Bras.* vi. II. 140. —Bras.
viridis, *Benth. ex Baker, l. c.* 137.—Bras.

STILPNOPHYLLUM, Hook. f. in Hook. Ic. Pl. t. 1147 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 33.
lineatum, *Hook. f. in Hook. Ic. Pl.* t. 1147.—Peruv.

STILPNOPHYTUM, Less. Syn. Comp. 264 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 433.
axillare, *Less. l. c.*—Afr. austr.
linifolium, *Less. l. c.*—Afr. austr.
longifolium, Less. l. c. = linifolium.
oocephalum, *DC. Prod.* vi. 93.—Afr. austr.

STIMEGAS, Rafin. Fl. Tellur. iv. 45 (1836) = **Cypripedium**, Linn. (Orchid.).
venustum, Rafin. l. c. 46 = C. venustum.

STIMENES, Rafin. Fl. Tellur. iii. 76 (1836) = **Nierembergia**, Ruiz & Pav. (Solanac.).
gracilis, Rafin. l. c. = N. gracilis.

STIMOMPHIS, Rafin. Fl. Tellur. iii. 76 (1836) = **Salpiglossis**, Ruiz & Pav. (Solan.).
linearis, Rafin. l. c. = Sal. linearis.

STIMORYNE, Rafin. Fl. Tellur. iii. 76 (1836) = **Petunia**, Juss. (Solan.).
purpurea, Rafin. l. c. = P. violacea.

STIMPSONIA, Wright, ex A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 401, in Adnot. *PRIMULACEAE*, Benth. & Hook. f. ii. 633.
chamaedryoides, *Wright, ex Gray, l. c.*—Ins. Luchu.
crispidens, Hance, in Journ. Bot. xviii. (1880) 234 (Lysimachiae sp.).—China.

STIPA, Linn. Sp. Pl. ed. II. 78 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1141.
ACHNATHERUM, Beauv. Agrost. 19. t. 6. f. 7 (1812).
ARISTELLA, Bertol. Fl. Ital. i. 690 (1833).
JARAVA, Ruiz & Pav. Prod. Fl. Per. 2. t. 1 (1794).
LASIAGROSTIS, Link, Hort. Berol. i. 99 (1827).
LASINGROSTIS, Schur, Enum. Pl. Transs. 741 (1866).
MACROCHLOA, Kunth, Rév. Gram. i. 58 (1830).
ORTHORAPHIUM, Nees, in Proc. Linn. Soc. i. (1841) 94.
PODOPOGON, Rafin. Neogenyt. 4 (1825).
PTILAGROSTIS, Griseb. in Ledeb. Fl. Ross. iv. 447 (1853).
STREPTACHNE, R. Br. Prod. 174 (1810).
TRICHOSATHERA, Ehrh. Beitr. iv. 149 (1789).
africana, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.
alpestris, Willd. ex Steud. Nom. ed. II. ii. 642 = Muehlenbergia alpestris.
altaica, Trin. ex Ledeb. Fl. Alt. i. 80 = splendens.
amethystina, *Steud. Syn. Pl. Gram.* 124.—Chili.
angulata, Steud. Nom. ed. II. ii. 642 = Zizania aquatica.
aperta, Janka, ex Čelak. in Oestr. Bot. Zeitschr. xxxiii. (1883) 319 = pennata.
appendiculata, Čelak. l. c. = pennata.
arabica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 77 = barbata.
arenaria, Brot. Fl. Lusit. i. 86 = gigantea, *Link*.
arguens, Linn. Sp. Pl. ed. II. 117 = Anthistiria arguens.
arguens, Thunb. Prod. Pl. Cap. 20 = Anthistiria imberbis.
Aristella, Gouan, Illustr. 4 = S. Calamagrostis.
Aristella, *Linn. Syst.* ed. XII. iii. App. 229.—Reg. Mediterr.; As. Min.
aristiglumis, *F. Muell. in Trans. Vict. Inst.* i. (1855) 43.—Austral.
arundinacea, *Benth. in Journ. Linn. Soc.* xix. (1881) 81.—N. Zel.
australis, *Steud. in Zoll. Syst. Verz. Ind. Archip.* 55; *Syn. Pl. Gram.* 132.—Java.
avenacea, Hook. & Arn. Bot. Beech. Voy. 403 = setigera.
avenacea, *Linn. Sp. Pl.* 78.—Am. bor.
avenacea, Spreng. ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 40 = hyalina.
avenacea, Willd. ex Steud. Nom. ed. II. ii. 643 = virescens.
barbata, *Desf. Fl. Atlant.* i. 97. t. 27.—Reg. Mediterr.
barbata, Michx. Fl. Bor. Am. i. 53 = avenacea.
bella, Phil. Sert. Mendoc. Alt. 45 = Neaei.
Bergeri, Link, in Linnaea, ix. (1834) 135 = parviflora.
Bertrandi, *Phil. l. c.* xxxiii. (1864–65) 283.—Chili.
bicolor, Cav. Ic. v. 42. t. 466. f. 2 = Neesiana.
bicolor, Nees, Agrost. Bras. 373, partim = Oryzopsis setacea.
bicolor, Pursh, Fl. Am. Sept. i. 73 = avenacea.
bicolor, *Vahl, Symb. Bot.* ii. 24.—Bras.; Chili.
Bloomeri, Boland. in Proc. Calif. Acad. iv. (1870) 168 = sibirica.
brachychaeta, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 450.—Hab. ?
brevicalyx, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 538 = Oryzopsis Fournieriana.
breviflora, *Griseb. in Goett. Nachr.* (1868) 82.—Tibet.
brevipes, *E. Desv. in C. Gay, Fl. Chil.* vi. 282.—Chili.
Bungeana, *Trin. ex Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 144.—China; Mongol.

STIPA :—
caerulescens, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 449 = Oryzopsis caerulescens.
caespitosa, Willd. ex Steud. Nom. ed. II. ii. 643 = Aristida jorullensis.
Calamagrostis, *Wahlenb. Veg. Helv.* 23.—Europ. austr.
calatajeronensis, Tineo, ex Arcang. Comp. Fl. Ital. 772 = barbata.
campylachne, Nees, in Lehm. Pl. Preiss. ii. 99 = semibarbata.
canadensis, Poir. Encyc. vii. 452 = Oryzopsis parviflora.
capensis, Thunb. Prod. Pl. Cap. 19 = tortilis ?
capillaris, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 350 = capillata.
capillaris, Lam. Illustr. i. 158 = Muehlenbergia capillaris.
capillata, Hook. Fl. Bor. Am. ii. 237 = comata.
capillata, *Linn. Sp. Pl.* ed. II. 116.—Europ. austr. ; As. bor.
Caragana, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 74 = Redowskii.
caspia, C. Koch, in Linnaea, xxi. (1848) 440 = barbata.
caudata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 75.—Chili.
chapulcensis, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 536.—Mexic.
chrysophylla, *E. Desv. in C. Gay, Fl. Chil.* vi. 278. t. 76. f. 2.—Am. bor. occ. ; Chili.
ciliata, *Scheele, in Linnaea*, xxii. (1849) 342.—Texas.
cirrosa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 536.—Mexic.
Clarazii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 237.—Patagon.
coerulea, *J. Presl, in Rel. Haenk.* i. 227.—Mexic.
coleotricha, *Steud. Syn. Pl. Gram.* 125.—Mexic.
comata, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* v. (1842) 75.—Am. bor.
commutata, Trin. & Rupr. l. c. 49 = pubescens.
compressa, *R. Br. Prod.* 175.—Austral.
conferta, Poir. Encyc. Suppl. v. 252 = Redowskii.
consanguinea, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 78 = Richteriana.
contracta, *Phil. in Linnaea*, xxxiii. (1864–65) 285.—Chili.
contracta, Willd. ex Steud. Nom. ed. II. ii. 643 = Muehlenbergia glabrata.
coronata, *Thurb. ex S. Wats. in Bot. Calif.* ii. 287.—Calif.
crinita, Gaudich. in Freyc. Voy. Bot. 407 = flavescens ?
Cumingiana, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Chili.
damascena, Boiss. Diagn. Ser. I. xiii. 45 = barbata.
dasyphylla, Czern. ex Trautv. in Act. Hort. Petrop. ix. (1884) 350, nomen = pennata.
Delilei, Steud. Syn. Pl. Gram. 126 = papposa.
demissa, *Steud. l. c.* 130.—Am. bor.
Dichelachne, Steud. Nom. ed. II. ii. 643 = Dichelachne sciurea.
diffusa, Walt. Fl. Carol. 78 = Muehlenbergia capillaris.
diffusa, Willd. ex Steud. Nom. ed. II. ii. 643 = avenacea.
Dregeana, *Steud. Syn. Pl. Gram.* 132.—Afr. austr.
Drummondii, *Steud. l. c.* 128.—Austral.
duriuscula, *Phil. in Linnaea*, xxxiii. (1864–65) 282.—Chili.
editorum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 536.—Mexic.
Ehrenbergiana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 75 = barbata.
elegantissima, *Labill. Nov. Holl. Pl.* i. 23. t. 29.—Austral.
elongata, *Roth, in Usteri, Ann. Bot.* xiv. (1795) 22.—Peruv.
elongata, *Steud. Syn. Pl. Gram.* 132.—Afr. austr.
eminens, *Cav. Ic.* v. 42. t. 467. f. 1.—Am. bor. et austr.
eminens, Nees, Agrost. Bras. 374 = setigera.
erecta, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 536.—Mexic.
erecta, Hort. Madr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 81 = capillata.
eriocaulis, *Borb. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 401.—Europ. austr.

STIPA :—
eriopus, *Benth. Fl. Austral.* vii. 570.—Austral.
eriostachya, H. B. & K. Nov. Gen. et Sp. i. 127. t. 41 = S. Jarava.
expansa, Poir. Encyc. vii. 453 = Cinna arundinacea.
expansa, Willd. ex Steud. Nom. ed. II. ii. 643 = Gymnopogon racemosus.
fernandeziana, *Phil. in Anal. Univ. Chil.* (1873) 560.—Chili.
fernandeziana, Steud. Syn. Pl. Gram. 124 = bicolor.
filiculmis, *Delile, Ind. Sem. Hort. Monsp.* (1849) 7.—Hab. ?
filifolia, *Nees, Agrost. Bras.* 379.—Reg. Argent.
filiformis, Turcz. ex Steud. Nom. ed. II. ii. 643 = mongolica.
fimbriata, H. B. & K. Nov. Gen. et Sp. i. 126 = Oryzopsis fimbriata.
flavescens, *Labill. Nov. Holl. Pl.* i. 24. t. 30.—Austral.
flavescens, Nees, in Lehm. Pl. Preiss. ii. 99 = scabra.
Fontanesii, Parl. Fl. Ital. i. 167, in obs. = gigantea.
formicarum, *Delile, Ind. Sem. Hort. Monsp.* (1849) 7.—Hab. ?
frigida, *Phil. Fl. Atac.* 54.—Chili.
gallica, Čelak. in Oestr. Bot. Zeitschr. xxxiii. (1883) 317 = pennata.
geniculata, *Phil. Sert. Mendoc. Alt.* 46.—Chili.
gigantea, *Lag. Gen. et Sp. Nov.* 3.—Reg. Mediterr. occ.
gigantea, Ledeb. Fl. Ross. iv. 448 = Lagascae.
gigantea, *Link, in Schrad. Journ.* ii. (1799) 313.—Lusitan.
Grafiana, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 116 = pennata.
Grisebachii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 537.—Mexic.
gynerioides, Phil. in Anal. Univ. Chil. (1870) 103 ; ex Hemsl. l. c. = S. Jarava.
Haussknechtii, *Boiss. Fl. Orient.* v. 501.—Persia.
hemipogon, *Benth. Fl. Austral.* vii. 569.—Austral.
heterophylla, Willd. ex Nees, Agrost. Bras. 378 = fimbriata.
hirta, *Phil. in Linnaea*, xxxiii. (1864–65) 283.—Chili.
hispanica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 77 = barbata.
Hohenackeriana, Trin. & Rupr. l. c. 80 = barbata.
holosericea, Trin. l. c. Sér. VI. i. (1831) 81 = Lagascae.
humilis, Brot. Fl. Lusit. i. 86 = tortilis.
humilis, *Cav. Ic.* v. 41. t. 466. f. 1.—Am. austr.
hyalina, *Nees, Agrost. Bras.* 378.—Reg. Argent.
hymenoides, Roem. & Schult. Syst. ii. 339 = Oryzopsis cuspidata.
ibarrensis, *H. B. & K. Nov. Gen. et Sp.* i. 125.—Ecuador.
Ichu, Kunth, Rév. Gram. i. 60 = S. Jarava.
inconspicua, J. Presl, in Rel. Haenk. i. 227 (= *Urachne Haenkeana*).—Peruv.
inebrians, *Hance, in Journ. Bot.* xiv. (1876) 212.—China.
intermedia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* v. (1842) 26.—Bras.
intricata, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 449.—Hab. ?
Jacquemontii, *Jaub. & Spach, Illustr.* iv. 60.—Reg. Himal.
Jarava, *Beauv. Agrost.* 19. t. 6. f. 3.—Mexic.
juncea, Cav. Desc. 39 = gigantea, *Lag.*
juncea, Kunze, in Flora, xxix. (1846) 631 = parviflora.
juncea, Lam. Illustr. i. 157 = capillata.
juncea, *Linn. Sp. Pl.* 78.—Reg. Mediterr.
juncea, Michx. Fl. Bor. Am. i. 54 = Oryzopsis parviflora.
juncea, Nutt. Gen. Am. i. 58 = comata.
juncea, Pursh, Fl. Am. Sept. i. 72 = spartea.
juncea, Sibth. & Sm. Fl. Graec. i. 67. t. 85 = Lagascae.
Kingii, *Boland. in Proc. Calif. Acad.* iv. (1870) 170.—Calif.
koleotricha, *Steud. Syn. Pl. Gram.* 125.—Mexic.
Kotschyana, Hochst. ex Ledeb. Fl. Ross. iv. 448 = Lagascae.
lachnophylla, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 67 (= *Urachne megalantha*).—Chili.
laeviculmis, *Nees, in Lehm. Pl. Preiss.* ii. 99.—Tasman.
Lagascae, Guss. Pl. Rar. 48 = gigantea, *Link.*
Lagascae, *Roem. & Schult. Syst.* ii. 333.—Reg. Mediterr. ; Oriens.

STIPA :—
latifolia, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 449 = Olyra latifolia.
laxa, *E. Desv. in C. Gay, Fl. Chil.* vi. 283.—Chili.
laxa, Portenschl. ex Steud. Syn. Pl. Gram. 132 = elongata, *Steud.*
Lechleriana, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
leiocarpa, *Nees, ex Steud. Syn. Pl. Gram.* 124.—Chili.
leptostachya, *Griseb. in Goett. Abh.* xxiv. (1879) 299.—Reg. Argent.
Lessingiana, Trin. & Rupr. in Mém. Acad. Pétersb. Sc. Sér. VI. Sc. Nat. v. (1842) 79 = pennata ?
leucogluma, *Steud. Syn. Pl. Gram.* 420.—Chili.
leucotricha, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 54 = setigera.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 537.—Mexic.
linearifolia, *Fourn. ex Hemsl. l. c.*—Mexic.
littorea, Burm. f. Fl. Ind. 29 = Spinifex squarrosus.
longearistata, *Steud. Syn. Pl. Gram.* 127.—Austral.
longiflora, *Steud. l. c.* 124.—Chili.
longiglumis, *Phil. in Linnaea*, xxxiii. (1864–65) 286.—Chili.
Lorentziana, *Griseb. in Goett. Abh.* xxiv. (1879) 298.—Reg. Argent.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 300.—Madag.
manicata, *E. Desv. in C. Gay, Fl. Chil.* vi. 288.—Chili.
megalantha, *Steud. Syn. Pl. Gram.* 124.—Chili.
megapotamica, *Spreng. ex Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 77.—Bras.
melanocarpa, Muhl. Desc. Gram. 183 = Heteropogon acuminatus.
melanosperma, J. Presl, in Rel. Haenk. i. 226.—Hab. ?
membranacea, Linn. Sp. Pl. 560 = Festuca bromoides.
membranacea, Pursh, Fl. Am. Sept. ii. 728 = Oryzopsis cuspidata.
Meyeriana, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 78 = barbata.
micrantha, *Cav. Ic.* v. 42 ; *R. Br. Prod.* 175.—Austral.
micrantha, Nees, ex Hook. f. Handb. Fl. N. Zel. 326 = Dichalachne sciurea.
micrantha, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 74 = ramosissima.
microsperma, Willd. ex Steud. Nom. ed. II. ii. 643 = Andropogon stipoides.
mollis, R. Br. Prod. 174 = semibarbata.
mongolica, *Turcz. ex Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Sibir. baical. ; Am. bor. occ.
mucronata, *H. B. & K. Nov. Gen. et Sp.* i. 125.—Mexic.
Muelleri, *Tate, in Trans. Roy. Soc. S. Austral.* vii. (1885) 70.—Austral.
Neaei, *Nees, ex Steud. Syn. Pl. Gram.* 126.—Patagon.
Neesiana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 27 = setigera.
nitens, *Ball, in Journ. Linn. Soc.* xvi. (1878) 712.—Marocc.
nivalis, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Nuttalliana, Steud. Nom. ed. II. ii. 643 = viridula.
occidentalis, Boland. in Proc. Calif. Acad. iv. (1870) 169, partim = comata.
occidentalis, *Thurb. ex S. Wats. in Bot. King's Exp.* 380.—Am. bor. occ.
orientalis, C. A. Mey. Verz. Pfl. Cauc. 14 = barbata.
orientalis, *Trin. ex Ledeb. Fl. Alt.* i. 83.—As. bor.
Orthoraphium, *Steud. Syn. Pl. Gram.* 131.—Ind. or.
ovata, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 73 = Oryzopsis setacea.
paleacea, Poir. Voy. Barb. ii. 101 = Anthistiria paleacea.
paleacea, Sibth. & Sm. Fl. Graec. i. 68. t. 86 = tortilis.
paleacea, Vahl, Symb. Bot. ii. 24 = barbata.
panicoides, Lam. Illustr. i. 158 = Oryzopsis setacea.
panicoides, Nees, Agrost. Bras. 376 (= *Urachne panicoides*).—Chili ; Bras.
papposa, *Delile, Ind. Sem. Hort. Monsp.* (1849) 7.—Hab. ?
papposa, Nees, Agrost. Bras. 377 = S. Jarava.
paradoxa, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 449 = Oryzopsis paradoxa.
Parishii, *Vasey, in Coult. Bot. Gaz.* vii. (1882) 33.—Calif.

STIPA:—
parviflora, *Desf. Fl. Atlant.* i. 98. t. 29.—Reg. Mediterr.
parviflora, Nutt. Gen. Am. i. 58 = viridula.
parvula, *Nees, Fl. Afr. Austr.* 169.—Afr. austr.
patens, Willd. ex Steud. Nom. ed. II. ii. 643 = Muehlenbergia stipoides.
pauciflora, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 449 = Olyra pauciflora.
pekinensis, *Hance, in Journ. Bot.* xv. (1877) 268.—China.
pellita, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 71 = Lagascae.
pennata, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 213 = barbata.
pennata, *Linn. Sp. Pl.* 78.—Europ.; As. et Am. bor.
pennatiformis, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 250 = barbata.
penniglumis, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 77 = Chrysopogon stipoides.
Pentapogon, F. Muell. Fragm. viii. 106 = Pentapogon Billardieri.
Petriei, J. Buch. Man. Indig. Grasses, N. Zeal. t. 17^2. f. 2 = setacea.
Philippii, *Steud. Syn. Pl. Gram.* 125.—Chili.
platyphylla, Czern. ex Trautv. in Act. Hort. Petrop. ix. (1884) 350, nomen = pennata.
plumosa, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 60 = barbata.
plumosa, Sieber, ex Nees, Fl. Afr. Austr. 180 = Aristida obtusa.
plumosa, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Chili.
plumosula, *Nees, ex Steud. Syn. Pl. Gram.* 127.—Chili.
Poeppigiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. Sc. Nat. v. (1842) 29.—Chili.
poganathera, *E. Desv. in C. Gay, Fl. Chil.* vi. 277.—Chili.
pratensis, *Phil. in Linnaea*, xxxiii. (1864–65) 284.—Chili.
prolifera, *Steud. Syn. Pl. Gram.* 131.—Nubia.
puberula, Steud. l. c. 128 = scabra.
pubescens, *R. Br. Prod.* 174.—Austral.
pubescens, Lag. Gen. et Sp. Nov. 3 = Lagascae.
pubinodis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* v. (1842) 50.—Tasman.
pulchella, *Munro, ex Ball, in Journ. Linn. Soc.* xxi. (1884) 237.—Patagon.
pulcherrima, C. Koch, in Linnaea, xxi. (1848) 440 = pennata.
pungens, [Nees, in] Meyen, Reise, i. 484 = S. Jarava.
purpurea, *Griseb. in Goett. Nachr.* (1868) 82.—Tibet.
pycnostachya, *Benth. Fl. Austral.* vii. 568.—Austral.
ramosissima, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 74 = micrantha.
rariflora, *Benth. in Journ. Linn. Soc.* xix. (1881) 81.—Am. antarct.
Redowskii, *Trin. in Spreng. Neue Entdeck.* ii. 53.—As. bor.
Regeliana, *Hack. in Sitzb. Akad. Wien*, lxxxix. (1884) 130.—As. centr.
Richardsoni, *Link, Hort. Berol.* ii. 245.—Am. bor.
Richteriana, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 862.—Soongar.
robusta, Nutt. ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 69 = spartea.
rostrata, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 142 (Quid?).—Ins. Galapag.
rostrata, Willd. ex Steud. Nom. ed. II. ii. 643 = Muehlenbergia stipoides.
rudis, Spreng. Syst. iv. Cur. Post. 31 = pubescens.
sareptana, Beck. in Bull. Soc. Nat. Mosc. lvii. (1882) 52 = capillata.
scabra, *Lindl. in Mitch. Journ. Trop. Austral.* 31.—Austral.
scelerata, Behr, ex Benth. Fl. Austral. vii. 568 = setacea.
Scribneri, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 125.—N. Mexic.
Sellowiana, *Nees, ex Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* v. (1842) 38.—Bras.
semibarbata, *R. Br. Prod.* 174.—Austral.
seminuda, Vahl, ex Hornem. Hort. Hafn. i. 76 = tortilis.
sericea, Michx. Fl. Bor. Am. i. 54 = Muehlenbergia capillaris.

STIPA:—
setacea, *R. Br. Prod.* 174.—Austral.
setifolia, Kunth, Enum. Pl. i. 182 (= *Urachne simplex*).—Peruv.
setigera, *J. Presl, in Rel. Haenk.* i. 226.—Am. bor. occ.
sibirica, *Lam. Illustr.* i. 158.—As. bor.; Reg. Himal.
Sibthorpii, Boiss. & Reut. ex Boiss. Fl. Orient. v. 501 = Lagascae.
spartea, Hook. Fl. Bor. Am. ii. 237 = viridula.
spartea, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 82.—Am. bor.
speciosa, *Trin. & Rupr. l. c.* Sér. VI. *Sc. Nat.* v. (1842) 45.—Calif.; Chili.
spicata, Linn. f. Suppl. 111; Thunb. Prod. Pl. Cap. 20 = Heteropogon hirtus.
spicata, Walt. Fl. Carol. 78 = Ctenium carolinianum.
Spica-venti, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 449 = Apera Spica-venti.
Spinifex, Linn. Mant. i. 84 = Spinifex squarrosus.
splendens, *Trin. in Spreng. Neue Entdeck.* ii. 54.—Sibir.
stenophylla, Czern. ex Trautv. in Act. Hort. Petrop. ix. (1884) 351, nomen = pennata.
Stillmannii, *Boland. in Proc. Calif. Acad.* iv. (1870) 169.—Calif.
Streptachne, *F. Muell. Census*, i. 133.—Austral.
striata, *Link, Hort. Berol.* i. 98.—Austral.?
stricta, *Lam. Illustr.* i. 158.—Am. bor.
stricta, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 42.—Am. bor.
subulata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 537.—Mexic.
Szowitsiana, Trin. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) 243 = barbata.
tauricola, Čelak. in Oestr. Bot. Zeitschr. xxxiii. (1883) 349 = pennata?
tenacissima, *Linn. Cent. Pl.* i. 6; *Amoen. Acad.* iv. 266.—Reg. Mediterr. occ.
tenacissima, Ucria, Hort. Panorm. (1789) 61 = tortilis.
tenella, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 452.—Hab.?
tenuiflora, *Phil. in Linnaea*, xxx. (1864–65) 281.—Chili.
tenuifolia, Sieber, ex Steud. Nom. ed. II. ii. 643 = parviflora.
tenuifolia, Steud. Syn. Pl. Gram. 128 = scabra.
tenuis, *Phil. Sert. Mendoc. Alt.* 46.—Chili.
tenuis, Willd. ex Steud. Nom. ed. II. ii. 643 = Aristida tenuis, *Kunth*.
tenuissima, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Chili.
teretifolia, *Steud. Syn. Pl. Gram.* 128.—Austral.; N. Zel.
Tirsa, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 115 = pennata.
tortilis, Buch, in Abh. Akad. Berl. 1816–17 (1819), 376 = tenacissima.
tortilis, *Desf. Fl. Atlant.* i. 99. t. 31.—Reg. Mediterr.; Oriens.; Afr. austr.
tortuosa, *E. Desv. in C. Gay, Fl. Chil.* vi. 281.—Chili.
trachysperma, *Phil. in Linnaea*, xxxiii. (1864–65) 284.—Chili.
trichophylla, *Benth. Fl. Austral.* vii. 570.—Austral.
trichotoma, Nees, Agrost. Bras. 375 (= *Urachne trichotoma*).—Uruguay.
trochlearis, [*Nees, in*] *Meyen, Reise*, i. 484, nomen.—Peruv.; Mexic.
Tuckeri, *F. Muell. Fragm.* xi. 128.—Austral.
ucranica, Steud. Nom. ed. I. 816 = seq.
ukranensis, Lam. Illustr. i. 157 = capillata.
vaginata, *Phil. in Linnaea*, xxxiii. (1864–65) 281.—Chili.
verticillata, Nees, in Spreng. Syst. iv. Cur. Post. 30 = micrantha.
villosa, Walt. Fl. Carol. 78 = Oryzopsis cuspidata.
virescens, *H. B. & K. Nov. Gen. et Sp.* i. 126. t. 680.—Mexic.
virginica, Pers. Syn. i. 99 = avenacea.
viridula, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Am. bor. occ.
Virletii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 538.—Mexic.
vulgaris, Gueldenst. Reisen, ii. 39; ex Ledeb. Fl. Ross. iv. 451 = pennata.
vulpiaeformis, *Cesati, in Att. Soc. Acc. Sc. Nap.* v. (1873) n. VII. 19.—Reg. Argent.

STIPAGROSTIS, Nees, in Linnaea, vii. (1832) 290 = **Aristida**, Linn. (Gramin.).
capensis, Nees, l. c. 291 = A. capensis.
Dregeana, Nees, Fl. Afr. Austr. 172 = A. Dregeana.
geminifolia, Nees, l. c. 173 = A. geminifolia.
obtusa, Nees, in Linnaea, vii. (1833) 293 = A. obtusa.
plumosa, Munro, ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 40 = A. plumosa.

STIPECOMA, Muell. Arg. in Mart. Fl. Bras. vi. I. 175. t. 53 (1860). *APOCYNACEAE*, Benth. & Hook. f. ii. 724.
macrocalyx, Miers, Apocyn. S. Am. 136 = Echites macrocalyx.
mucronata, Miers, l. c. 135 = Echites peltata.
ovata, Miers, l. c. 137 (Quid?).—Bras.
peltata, Miers, l. c. 134 = Echites peltata.
peltigera, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 176.—Bras.
plicata, Miers, Apocyn. S. Am. 134 = Echites peltata.
pulchra, Miers, l. c. 135 (Quid?).—Bras.
speciosa, Miers, l. c. 136 (Quid?).—Bras.

STIPELLARIA, Benth. in Hook. Kew Journ. vi. (1854) 2 = **Alchornea**, Sw. (Euphorb.).
mollis, Benth. l. c. 3 = A. mollis.
parviflora, Benth. l. c. 4 = A. parviflora.
tiliaefolia, Benth. l. c. = A. tiliaefolia.
trewioides, Benth. l. c. 3 = A. trewioides.
villosa, Benth. l. c. 4 = A. villosa.
Zollingeri, Miq. Fl. Ind. Bot. Suppl. 183, 460 = A. villosa.

STIPELLARIA, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1203 (1848) nomen. *LEGUMINOSAE*.
mollis, Klotzsch, l. c. (Quid?).—Guian.

STIPHONIA, Hemsl. Biol. Centr. Am. Bot. 218 (1880), sphalm. = STYPHONIA, Nutt. = **Rhus**, Linn. (Anacard.).

STIPULARIA, Beauv. Fl. Owar. ii. 26. t. 75 (1805). *RUBIACEAE*, Benth. & Hook. f. ii. 74.
africana, *Beauv. l. c.*—Afr. trop.
elliptica, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 80.—Afr. trop.
gabonica, *Hiern, l. c.*—Afr. trop.

STIPULARIA, Haw. Syn. Pl. Succ. 103 (1812) = **Spergularia**, J. & C. Presl (Caryoph.).
capensis, Haw. l. c. 104 = Arenaria capensis.
fallax, Edgew. in Journ. Linn. Soc. vi. (1862) 185 (Quid?).—Ind. or.
flaccida, Edgew. l. c. = Spergula pentandra.
marina, Haw. Syn. Pl. Succ. 104 = Spergularia media.
media, Haw. l. c. 103 = Spergularia media.
pentandra, Edgew. in Journ. Linn. Soc. vi. (1862) 185 = Spergula pentandra.
pubescens, Haw. Syn. Pl. Succ. 103 = Spergularia media.
rubra, Haw. l. c. = Spergularia rubra.

STIPULICIDA, Michx. Fl. Bor. Am. i. 26. t. 6 (1803). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 154.
setacea, *Michx. l. c.*—Am. bor.

STIRADOTHECA, Klotzsch, in Monatsb. Berl. Akad. (Maerz 1854) 127 = **Begonia**, Tourn.
ferruginea, Klotzsch, l. c. = B. ferruginea.
magnifica, Klotzsch, l. c. = B. magnifica.
trachyptera, Klotzsch, l. c. = B. trachyptera.

STIRLINGIA, Endl. Gen. 339 (1838). *PROTEACEAE*, Benth. & Hook. f. iii. 173.
SIMSIA, R. Br. in Trans. Linn. Soc. x. (1810) 152.
abrotanoides, *Meissn. in Lehm. Pl. Preiss.* i. 517.—Austral.
acutifolia, Endl. Gen. Suppl. iv. II. 81 = tenuifolia.
affinis, Meissn. in Lehm. Pl. Preiss. i. 516 = teretifolia.
anethifolia, Endl. Ic. t. 23 = tenuifolia.
capillifolia, Meissn. in Hook. Kew Journ. vii. (1855) 70 = simplex.
intricata, Meissn. l. c. iv. (1852) 184 = tenuifolia.
latifolia, *Steud. Nom.* ed. II. ii. 644.—Austral.
paniculata, Lindl. Swan River App. 30 = latifolia.
simplex, *Lindl. l. c.*—Austral.
tenuifolia, *Steud. Nom.* ed. II. ii. 644; *Endl. Gen. Suppl.* iv. II. 81.—Austral.
teretifolia, *Meissn. in Lehm. Pl. Preiss.* i. 515.—Austral.

STISSERA, Giseke, Prael. Linn. 207, 209 (1792) = **Curcuma**, Linn. (Scitam.).
Curcuma, Raeusch. Nom. ed. III. 2 = C. longa.
longa, Giseke, Prael. Linn. 228 = C. longa.

STISSERIA, Linn. Syst. ed. I. (1735) = **Stapelia**?, Linn. (Asclep.).

STISSERIA, Scop. Introd. 199 (1777) = **Mimusops**, Linn. (Sapot.).

STIXIS, Lour. Fl. Cochinch. 295 (1790) = **Roydsia**, Roxb. (1819) (Cappar.).
scandens, Lour. l. c.—Cochinch.

STIZA, E. Mey. Comm. Pl. Afr. Austr. 31 (1835) = **Lebeckia**, Thunb. (Legumin.).
erioloba, E. Mey. l. c. = L. pungens.
psiloloba, E. Mey. l. c. 32 = L. psiloloba.
pungens, Presl, Bot. Bemerk. 125 = L. pungens.

STIZOLOBIUM, P. Br. Hist. Jamaic. 290 (1756) = **Mucuna**, Adans. (1763) (Legumin.).
altissimum, Pers. Syn. ii. 299 = M. altissima.
bulbosum, Spreng. Syst. iii. 252 = Pachyrhizus angulatus.
domingense, Spreng. l. c. = Pachyrhizus angulatus.
ellipticum, Pers. Syn. ii. 299 = M. elliptica.
giganteum, Spreng. Syst. iv. Cur. Post. 281 = M. gigantea.
inflexum, Pers. Syn. ii. 299 = M. inflexa.
macroceras, Spreng. Syst. iv. Cur. Post. 281 = M. macroceratides.
Meyeri, Spreng. l. c. iii. 253 = M. spuria.
mite, Pers. Syn. ii. 299 = M. mitis.
molle, Spreng. Syst. iv. Cur. Post. 281 = M. mollis.
montanum, Spreng. Syst. iii. 252 = Pueraria phaseoloides.
Mutisianum, Spreng. Syst. iv. Cur. Post. 281 = M. Mutisiana.
nigricans, Pers. Syn. ii. 299 = M. nigricans.
platycarpum, Pers. l. c. = M. platycarpa.
pruriens, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 399 = M. pruriens.
scarabaeoides, Spreng. Syst. iii. 253 = Atylosia scarabaeoides.
tuberosum, Spreng. l. c. 252 = Pachyrhizus tuberosus.
urens, Pers. Syn. ii. 299 = M. urens.

STIZOLOPHUS, Cass. in Dict. Sc. Nat. xliv. 35 (1826) = **Centaurea**, Linn. (Compos.).
Balsamita, Cass. l. c. 36 = C. Balsamita.
balsamitaefolius, Cass. l. c. li. 49, 50 = C. Balsamita.
coronopifolius, Cass. l. c. 51 = C. coronopifolia.

STIZOPHYLLUM, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 197 = **Bignonia**, Linn.
adspersum, Miers, l. c. 198, nomen.—Bras.
affinis, Miers, l. c. nomen. = B. affinis.
auricomum, Miers, l. c., nomen.—Bras.
cordifolium, Miers, l. c. = B. cordifolia.
dentatum, Miers, l. c. = B. dentata.
denticulatum, Miers, l. c. 197.—Bras.
glandulosum, Miers, l. c. 198 = B. glandulosa.
insigne, Miers, l. c., nomen.—Bras.
longipes, Miers, l. c., nomen.—Bras.
occidentale, Miers, l. c. = B. occidentalis.
perforatum, Miers, l. c. = B. perforata.
punctulatum, Miers, l. c. nomen.—Bras.
velutinum, Miers, l. c. = B. velutina.

STOBAEA, Thunb. Prod. Pl. Cap. 141 (1800) = **Berkheya**, Ehrh. (Compos.).
acanthopoda, DC. Prod. vi. 521.—Afr. austr.
acarnoides, DC. l. c. 519.—Afr. austr.
adenocarpa, DC. l. c. 515 = B. fruticosa?
aristosa, DC. l. c. 518.—Afr. austr.
atractyloides, Thunb. Prod. Pl. Cap. 141 = B. carlinoides.
Bergeri, DC. Prod. vii. 37, sphalm. = seq.
Bergii, Spreng. Syst. iii. 394 = Dicoma diacanthoides.
biloba, DC. Prod. vi. 516.—Afr. austr.
bipinnatifida, Harv. in Harv. & Sond. Fl. Cap. iii. 499.—Afr. austr.
buphthalmoides, DC. Prod. vi. 517.—Afr. austr.
cardopatifolia, DC. l. c. 514.—Afr. austr.
carlinaefolia, DC. l. c. 516.—Afr. austr.

STOBAEA:—
carlinoides, E. Mey. ex DC. Prod. vi. 516 = *carlinaefolia*.
carlinoides, Thunb. Prod. Pl. Cap. 141 = Berkheya carlinoides.
cirsiifolia, DC. Prod. vi. 519.—Afr. austr.
crispa, DC. l. c. 517.—Afr. austr.
cruciata, Harv. in Harv. & Sond. Fl. Cap. iii. 498 = B. cruciata.
decurrens, Thunb. Prod. Pl. Cap. 141 = Stephanocoma berckheyoides.
discolor, DC. Prod. vi. 520.—Afr. austr.
echinacea, Harv. in Harv. & Sond. Fl. Cap. iii. 495.—Afr. austr.
echinopoda, DC. Prod. vi. 521.—Afr. austr.
epitrachys, DC. l. c. 515.—Afr. austr.
eriobasis, DC. l. c. 516.—Afr. austr.
Erysithales, DC. l. c. 519.—Afr. austr.
erysithalis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
Gaertneri, DC. Prod. vi. 522.—Afr. austr.
Gerrardii, Harv. in Harv. & Sond. Fl. Cap. iii. 495.—Afr. austr.
glabrata, Alam. ex DC. Prod. v. 660 = Jaumea mexicana.
glabrata, Thunb. Prod. Pl. Cap. 141.—Afr. austr.
glabriuscula, DC. Prod. vi. 519.—Afr. austr.
glauca, DC. l. c. 516.—Afr. austr.
glomerata, Spreng. Syst. iii. 394 = Platycarpha glomerata.
grandifolia, DC. Prod. vi. 517.—Afr. austr.
helianthiflora, DC. l. c. 520.—Afr. austr.
heterophylla, Thunb. Prod. Pl. Cap. 141.—Afr. austr.
insignis, Harv. in Harv. & Sond. Fl. Cap. iii. 496 = B. subulata.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 776 = B. grandiflora.
lanata, Thunb. Prod. Pl. Cap. 141.—Afr. austr.
M'Kenii, Harv. in Harv. & Sond. Fl. Cap. iii. 494.—Afr. austr.
membranifolia, DC. Prod. vi. 521.—Afr. austr.
microcephala, DC. l. c. 520.—Afr. austr.
multijuga, DC. l. c. 521.—Afr. austr.
onobromoides, DC. l. c. 516.—Afr. austr.
onopordifolia, DC. l. c. 521.—Afr. austr.
oppositifolia, DC. l. c. 517.—Afr. austr.
petiolata, DC. l. c. 520.—Afr. austr.
pinnata, Thunb. Prod. Pl. Cap. 141 = B. pinnata.
pinnatifida, Thunb. l. c.—Afr. austr.
platyptera, Harv. in Harv. & Sond. Fl. Cap. iii. 495.—Afr. austr.
polyacantha, DC. Prod. vi. 519.—Afr. austr.
purpurea, DC. l. c. 518 = B. purpurea.
Radula, Harv. in Harv. & Sond. Fl. Cap. iii. 491.—Afr. austr.
rhapontica, DC. Prod. vi. 519.—Afr. austr.
rigida, Thunb. Prod. Pl. Cap. 141.—Afr. austr.
rubricaulis, DC. Prod. vi. 515 = B. carlinoides.
scolymoides, DC. l. c. 520 = B. scolymoides.
seminivea, DC. l. c. 518 = B. seminivea.
sonchifolia, Harv. in Harv. & Sond. Fl. Cap. iii. 496.—Afr. austr.
speciosa, DC. Prod. vi. 518.—Afr. austr.
sphaerocephala, DC. l. c. 518.—Afr. austr.
viscosa, DC. l. c. 517.—Afr. austr.
Zeyheri, DC. l. c. 515 = B. carlinoides.
Zeyheri, Harv. & Sond. Fl. Cap. iii. 496 = B. subulata.

STOCKSIA, Benth. in Hook. Kew Journ. v. (1853) 304. *SAPINDACEAE*, Benth. & Hook. f. i. 397.
brahuica, *Benth. in Hook. Kew Journ.* v. (1853) 305.—Beluchist.

STOEBE, Linn. Coroll. Gen. 14 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 328.
EREMANTHIS, Cass. ex DC. Prod. vi. 261, in syn. (1837).
SERIPHIUM, Linn. Sp. Pl. ed. I. 928 (1753).
adpressa, *DC. ex Harv. & Sond. Fl. Cap.* iii. 285.—Afr. austr.
aethiopica, *Linn. Sp. Pl.* 831.—Afr. austr.
aethiopica, Sieber, ex DC. Prod. vi. 260 = incana.
alopecuroides, *Less. Syn. Comp.* 349.—Afr. austr.
alopecuroides, Reichb. ex DC. Prod. vi. 262 = cinerea.
artemisioides, *E. Mey, ex DC. l. c.* 263.—Afr. austr.

STOEBE:—
aspera, Thunb. Prod. Pl. Cap. 170 = Helichrysum ericaefolium.
canescens, Sch. Bip. in Pollichia, xviii–xix. (1861) 182 = Elytropappus canescens.
capitata, *Berg. Desc. Pl. Cap.* 338.—Afr. austr.
cernua, Thunb. Prod. Pl. Cap. 170 = Elytropappus Rhinocerotis.
cinerea, Sieber, ex DC. Prod. vi. 256 = Elytropappus glandulosus.
cinerea, *Thunb. Prod. Pl. Cap.* 169.—Afr. austr.
cinerea, Zeyh. ex DC. Prod. vi. 266 = Ifloga polycnemioides.
colorata, E. Mey. ex DC. l. c. 263 = spiralis.
copholepis, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 179.—Afr. austr.
cupressina, Reichb. ex DC. Prod. vi. 257 = Elytropappus Rhinocerotis.
disticha, Linn. f. Suppl. 391 (= *Trichogyne seriphioides*.—Afr. austr.
ericoides, *Berg. Desc. Pl. Cap.* 339.—Afr. austr.
fasciculata, Cass. in Dict. Sc. Nat. li. 62 = cinerea.
fasciculata, Thunb. Prod. Pl. Cap. 169 (= *Trichogyne seriphioides*).—Afr. austr.
filaginea, *Sch. Bip. ex Harv. & Sond. Fl. Cap.* iii. 285.—Afr. austr.
fusca, *Thunb. Prod. Pl. Cap.* 170.—Afr. austr.
glandulosus, Sch. Bip. in Pollichia, xviii–xix. (1861) 182 = Elytropappus glandulosus.
gnaphaloides, Linn. Mant. ii. 518 = gomphrenoides.
gnaphaloides, Thunb. Fl. Cap. ed. Schult. 726 = Metalasia pulcherrima.
gomphrenoides, *Berg. Desc. Pl. Cap.* 336.—Afr. austr.
incana, *Thunb. Prod. Pl. Cap.* 169.—Afr. austr.
laricifolia, E. Mey. ex DC. Prod. vi. 265 (= *Trichogyne laricifolia*).—Afr. austr.
leiocarpa, *Sch. Bip. in Flora*, xxvii. (1844) 694.—Afr. austr.
leucocephala, *DC. Prod.* vi. 259.—Afr. austr.
microcephala, Steud. Nom. ed. II. ii. 645, sphalm. = seq.
microphylla, *DC. Prod.* vi. 259.—Afr. austr.
mucida, E. Mey, ex DC. l. c. 256 = Elytropappus glandulosus.
muricata, Spreng. in Flora, xiii. (1830) i. Beil. 21 = Elytropappus glandulosus.
muricella, Steud. ex Sch. Bip. in Pollichia, xviii–xix. (1861) 183.—Afr. austr.
nervigera, *Sch. Bip. l. c.* 179.—Afr. austr.
nivea, Thunb. Prod. Pl. Cap. 170 = ericoides.
pallens, Sch. Bip. in Pollichia, xviii–xix. (1861) 182 = Elytropappus glandulosus.
paniculata, Cass. in Dict. Sc. Nat. li. 63 = seq.
passerinoides, *Willd. Sp. Pl.* iii. 2408.—Ins. Maurit.
phlaeoides, Harv. & Sond. Fl. Cap. iii. 282, sphalm. = seq.
phleoides, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 179.—Afr. austr.
phylicoides, *Thunb. Prod. Pl. Cap.* 169.—Afr. austr.
phyllostachya, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 179.—Afr. austr.
plumosa, Thunb. Prod. Pl. Cap. 169 = cinerea.
prostrata, *Linn. Mant.* ii. 291.—Afr. austr.
ramosissima, Sch. Bip. in Pollichia, xviii–xix. (1861) 183 = Elytropappus ambiguus.
reflexa, Linn. f. Suppl. 391 (= *Trichogyne reflexa*).—Afr. austr.
Rhinocerotis, Linn. f. Suppl. 391 = Elytropappus Rhinocerotis.
rigida, *Spreng. Syst.* iii. 441.—Afr. austr.
rugulosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 280.—Afr. austr.
scabra, Linn. f. Suppl. 391 = Elytropappus ambiguus.
spiralis, *Less. Syn. Comp.* 247.—Afr. austr.
squarrosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 282.—Afr. austr.
stenostachya, Sch. Bip. in Pollichia, xviii–xix. (1861) 183 = Elytropappus ambiguus.
subulata, Sm. in Rees, Cycl. xxxiv. n. 10 = Elytropappus ambiguus.
torta, Spreng. Neue Entdeck. iii. 42 = Disparago ericoides.
tortilis, *DC. Prod.* vi. 260.—Afr. austr.
vermiculata, DC. ex Harv. & Sond. Fl. Cap. iii. 284 = artemisioides.
virgata, Thunb. Prod. Pl. Cap. 170 = cinerea.

STOECHAS, Gueldenst. Reisen (1787); ex Ledeb. Fl. Ross. ii. 607 (1846) = **Helichrysum**, Vaill. (Compos.).
citrina, Gueldenst. l. c. = H. arenarium.

STOECHAS, Tourn. ex Linn. Syst. ed. I (1735) = **Lavandula**, Tourn. (Labiat.).
abrotanoides, Reichb. f. in Oestr. Bot. Wochenbl. vii. (1857) 161 = L. abrotanoides.
dentata, Mill. Gard. Dict. ed. VIII. n. 3; Reichb. f. l. c. = L. dentata.
officinarum, Mill. l. c. n. 1. = L. Stoechas.
pedunculata, Mill. l. c. n. 2; Reichb. f. in Oestr. Bot. Wochenbl. vii. (1857) 161 = L. pedunculata.
pinnata, Reichb. f. l. c. = L. pinnata.
rotundifolia, Reichb. f. l. c. = L. rotundifolia.

STOEHELINA, Hall. Helv. 312; ex Benth. in DC. Prod. x. 544 (1846) = **Bartsia**, Linn. (Scrophul.).

STOERKEA, Baker, in Journ. Linn. Soc. xiv. (1875) 523 = seq.

STOERKIA, Crantz, Duab. Dracon. Arb. 25 (1768) = **Dracaena**, Vand. (Liliac.).
Draco, Crantz, l. c. = D. Draco.

STOKESIA, L'Hérit. Sert. Angl. 27 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 234.
CARTESIA, Cass. in Bull. Soc. Philom. (1816) 198.
cyanea, *L'Hérit. Sert. Angl.* 28.—Am. bor.

STOLIDIA, Baill. Adansonia, ii. (1862) 359. *OLACINEAE*, Benth. & Hook. f. i. 996.
mauritiana, *Baill. l. c.*—Ins. Maurit.

STOMADENA, Rafin. Fl. Tellur. ii. 12 (1836) = **Ipomoea**, Linn. (Convolv.).
violacea, Rafin. l. c. = I. Aitoni.

STOMARRHENA, DC. Prod. vii. 738 (1839) = **Astroloma**, R. Br. (Epacrid.).
serratifolia, DC. l. c. = A. Candolleanum.
xerophylla, DC. l. c. = A. xerophyllum.

STOMATECHIUM, Spach, Hist. Vég. Phan. ix. 31 (1840), sphalm. = STOMOTECHIUM, Lehm. = **Anchusa**, Linn. (Borag.).

STOMATOCALYX, Muell. Arg. in DC. Prod. xv. II. 1142 (1866) = **Pimelodendron**, Hassk. (Euphorb.).
Griffithianus, Muell. Arg. l. c.—Malaya.

STOMOISIA, Rafin. Fl. Tellur. iv. 108 (1836) = **Utricularia**, Linn. (Lentib.).
cornuta, Rafin. l. c. = U. cornuta.

STOMOTECHIUM, Lehm. Asperif. 395 (1818) = **Anchusa**, Linn. (Borag.).
papillosum, Lehm. l. c. 396.—Afr. austr.

STONCKENYA, Rafin. in Am. Monthly Mag. iv. (1819) 193, err. typ. = HONCKENYA, Ehrh. = **Arenaria**, Rupp. (Caryoph.).

STOORIA, Neck. Elem. i. 132 (1790) = **Lobelia**, Plum. (Campan.).

STOPINACA, Rafin. Fl. Tellur. iii. 11 (1836) = **Polygonella**, Michx. (Polygon.).
parvifolia, Rafin. l. c. = P. parvifolia.

STORCKIELLA, Seem. in Bonplandia, ix. (1861) 255, 363. t. 6. *LEGUMINOSAE*, Benth. & Hook. f. i. 571.
DOGA, Baill. Adansonia, ix. (1869) 205.
Pancheri, *Baill. l. c.*—N. Caled.
vitiensis, *Seem. in Bonplandia*, ix. (1861) 255, 363. t. 6.—Ins. Fiji.

STORTHOCALYX, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 499, 660. *SAPINDACEAE*.
chryseus, *Radlk. l. c.* 661.—N. Caled.
leioneurus, *Radlk. l. c.*—N. Caled.
Pancheri, *Radlk. l. c.*—N. Caled.
sordidus, *Radlk. l. c.*—N. Caled.

STRABONIA, DC. Prod. v. 481 (1836) = **Pulicaria**, Gaertn. (Compos.).
gnaphalodes, DC. l. c. = P. gnaphalodes.

STRACHEYA, Benth. in Hook. Kew Journ. v. (1853) 306. *LEGUMINOSAE*, Benth. & Hook. f. i. 510.
tibetica, *Benth. in Hook. Kew Journ.* v. (1853) 307.—Reg. Himal.

STRAKAEA, Presl, Epim. Bot. 221 (1849) = **Bragantia**, Lour. (Aristol.).
melastomaefolia, Presl, l. c. 222 = B. corymbosa.

STRAMONIUM, Tourn. ex Hall. Enum. Stirp. Helv. ii. 489 (1742) = **Datura**, Linn. (Solan.).
arboreum, Moench, Meth. Suppl. 173 = D. suaveolens.
Datura, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = D. fastuosa.
fastuosum, Moench, Meth. 456 = D. fastuosa.
foetidum, Scop. Fl. Carn. ed. II. i. 157; All. Fl. Pedem. i. 103 = D. Stramonium.
infernale, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = D. fastuosa?
laeve, Moench, Meth. 456 = D. inermis?
Metel, Moench, l. c. = D. Metel.
spinosum, Lam. Fl. Fr. ii. 256 = D. Stramonium.
Tatula, Moench, Meth. 456 = D. Stramonium.
vulgare, Moench, l. c. = D. Stramonium.
vulgatum, Gaertn. Fruct. ii. 243. t. 132; Moench, l. c. = D. Stramonium.

STRANGALIS, Dulac, Fl. Hautes-Pyr. 195 (1867) = **Brassica**, Tourn. (Crucif.).
adpressa, Dulac, l. c. = B. adpressa.

STRANGEA, Meissn. in Hook. Kew Journ. vii. (1855) 66 = **Grevillea**, R. Br. (Proteac.).
cynanchocarpa, F. Muell. Fragm. vii. 132 = G. cynanchicarpa.
linearis, Meissn. in Hook. Kew Journ. vii. (1855) 66 = G. Strangea.

STRANGEVEIA, Baker, in Journ. Linn. Soc. xi. (1871) 429 = STRANGWEJA, Bertol. = **Hyacinthus**, Tourn. (Liliac.).

STRANGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Ardisia**, Sw. (Myrsin.).

STRANGWEIA, Baker, in Journ. Linn. Soc. xi. (1871) 423 = seq.

STRANGWEJA, Bertol. in Mem. Soc. Ital. Modena, xxi. (1835) 2. t. 1 = **Hyacinthus**, Tourn. (Liliac.).
hyacinthoides, Bertol. l. c. = H. spicatus.
spicata, Bertol. l. c. = H. spicatus.

STRANGWEYA, Benth. & Hook. f. Gen. iii. 812 (1883) = praec.

STRANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Canarium**, Rumph (Burser.).

STRANVAESIA, Lindl. Bot. Reg. t. 1956 (1837). *ROSACEAE*, Benth. & Hook. f. i. 627.
Calleryana, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 179.—China.
Davidiana, *Decne. l. c.* 179.—Tibet.
digyna, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 129 = Photinia villosa.
glauca, Baill. Hist. Pl. i. 411 = glaucescens.
glaucescens, *Lindl. Bot. Reg.* t. 1956.—Reg. Himal.
Nussia, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 178 = glaucescens.
undulata, *Decne. l. c.* 179.—China.

STRASBURGERIA, Baill. Adansonia, xi. (1876) 372. *TERNSTROEMIACEAE*.
callianthа, *Baill. l. c.*—N. Caled.

STRATEUMA, Rafin. Fl. Tellur. ii. 89 (1836) = **Zeuxine**, Lindl. (Orchid.).
zeylanica, Rafin. l. c. = Z. sulcata.

STRATEUMA, Salisb. in Trans. Hort. Soc. i. (1812) 290 = **Orchis**, Tourn.
grandis, Salisb. l. c. = O. purpurea.
militaris, Salisb. l. c. = O. Simia.

STRATIOITES, Gilib. Exercit. i. 342 (1791), sphalm. = seq.

STRATIOTES, Linn. Syst. ed. I (1735). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 454.
acoroides, Linn. f. Suppl. 268 = Enhalus Koenigii.
aculeatus, Stokes, Bot. Mat. Med. iii. 52 = aloides.
alismoides, Linn. Sp. Pl. 535 = Ottelia alismoides.
aloides, *Linn. l. c.*—Europ.
aquaticus, Pall. Reise, iii. 315 = aloides.
ensiformis, Gilib. Fl. Lituan. ii. 220 = aloides.
nymphoides, Humb. & Bonpl. in Willd. Sp. Pl. iv. 821 = Hydrocleys Commersonii.
quinquealatus, Stokes, Bot. Mat. Med. iii. 53 = Ottelia alismoides.

STRAUSSIA, A. Gray, in Proc. Am. Acad. iv. (1860) 42. *RUBIACEAE*, Benth. & Hook. f. ii. 122.
hawaiensis, *A. Gray, l. c.* 43.—Ins. Sandvic.
kaduana, *A. Gray, l. c.*—Ins. Sandvic.
mariniana, *A. Gray, l. c.*—Ins. Sandvic.

STRAVADIA, Pers. Syn. ii. 30 (1807) = seq.

STRAVADIUM, Juss. Gen. 326 (1789) = **Barringtonia**, Forst. (Myrtac.).
acuminatum, Blume, in Fl. des Serres, vii. (1851–52) 24 = B. acuminata,
acuminatum, Wall. Cat. n. 3636 = B. macrostachya.
acutangulum, Sweet, Hort. Brit. ed. I. 159 = B. acutangula.
album, Pers. Syn. ii. 30 = B. racemosa.
augustum, Wall. Cat. n. 3637 = B. augusta.
coccineum, DC. Prod. iii. 289 = B. excelsa.
cochinchinense, Blume, in Fl. des Serres, xvii. (1851–52) 24 = B. acutangula?
costatum, Blume, l. c. vii. (1851–52) 24 = B. costata.
demissum, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 81 = B. acutangula.
denticulatum, Miers, l. c. 88.—Austral.
excelsum, DC. Prod. iii. 289 = B. excelsa.
globosum, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 86 = B. acutangula.
gracile, Miers, l. c. = B. acutangula.
Horsfieldii, Miers, l. c. 85 = B. Horsfieldii.
insigne, Blume, in Fl. des Serres, xvii. (1851–52) 21, 24 = B. insignis.
integrifolium, Montr. in Mém. Acad. Lyon, x. (1860) 210.—N. Caled.
lucidum, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 88.—Java.
luzonense, Miers, l. c. 84 = B. luzonensis.
macrophyllum, Blume, in Fl. des Serres, vii. (1851–52) 24 = B. speciosa.
obtusangulum, Blume, l. c. = B. acutangula.
pubescens, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 83 = B. acutangula.
racemosum, Sweet, Hort. Brit. ed. I. 159 = B. racemosa.
Reinwardtii, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 88 = B. Reinwardtii.
reticulatum, Blume, in Fl. des Serres, vii. (1851–52) 24 = B. reticulata.
Rheedii, Blume, l. c. = B. acutangula.
rubrum, DC. Prod. iii. 289, partim = B. racemosa.
rubrum, Pers. Syn. ii. 30 = B. acutangula.
sarcostachys, Blume, in Fl. des Serres, vii. (1851–52) 24 = B. macrostachyum.
semisutum, Miers, in Trans. Linn. Soc. Sér. II. i. (1875) 89.—Ins. Pacif.
serratum, Miers, l. c. 87 = B. serrata.
spicatum, Blume, ex DC. Prod. iii. 289 = B. spicata.

STREBANTHUS, Rafin. in Ser. Bull. Bot. i. (1830) 218 = **Eryngium**, Tourn. (Umbell.).
auriculatus, Rafin. l. c. = E. Baldwini.
gracilis, Rafin. New Fl. Am. iv. 36 = E. Baldwini.
heterophylus, Rafin. l. c. 35 = E. prostratum.
humilis, Rafin. l. c. = E. Baldwini.
tenuifolia, Rafin. l. c. 36 = E. Baldwini.

STREBLANTHERA, Steud. in Flora, xxvii. (1844) 29 = **Trichodesma**, R. Br. (Borag.).
oleaefolia, A. Rich. Tent. Fl. Abyss. ii. 93.—Abyss.
trichodesmoides, Steud. in Flora, xxvii. (1844) 29 = T. calathiforme.

STREBLANTHUS, Rafin. Atl. Journ. 149 (1832) = STREBANTHUS, Rafin. = **Eryngium**, Tourn. (Umbell.).

STREBLIDIA, Link, Hort. Berol. i. 276 (1827) = **Schoenus**, Linn. (Cyperac.).
ferruginea, Link, l. c. = Sch. ferrugineus.

STREBLOCARPUS, Arn. in Ann. Sc. Nat. Sér. II. ii. (1834) 235 = **Maerua**, Forsk. (Cappar.).
angustifolia, Endl. Gen. 893 = M. oblongifolia.
Fenzlii, Parl. in Webb, Fragm. Fl. Aethiop. 24.—Abyss.
oblongifolia, Endl. Gen. 893 = M. oblongifolia.
ovalifolia, Endl. ex Walp. Rep. i. 197 = M. ovalifolia.
pubescens, Klotzsch, in Peters, Reise Mossamb. Bot. 166 = M. nervosa.
scabra, Endl. ex Walp. Rep. i. 197 = M. ovalifolia.
scandens, Klotzsch, in Peters, Reise Mossamb. Bot. 165 = M. nervosa.

STREBLOCHAETA, Benth. & Hook. f. Gen. iii. 1163 (1883), sphalm. = seq.

STREBLOCHAETE, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 421 (1851) = **Danthonia**, DC. (Gramin.).
Costlini, Hochst. ex A. Rich. l. c. = D. Streblochaeta.
nutans, Hochst. ex Steud. Syn. Pl. Gram. 245 = D. Streblochaeta.
Schimperi, Hochst. ex Steud. l. c. = D. Streblochaeta.

STREBLORHIZA, Benth. & Hook. f. Gen. i. 503. (err. typ.) (1865) = seq.

STREBLORRHIZA, Endl. Prod. Fl. Norf. 92 (1833). *LEGUMINOSAE*, Benth. & Hook. f. i. 503.
speciosa, *Endl. l. c.* 93.—Ins. Norfolk.

STREBLOSA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 245 = **Psychotria**, Linn. (Rubiac.).
polyantha, Korth. l. c. 246 = P. singalensis.
tortilis, Korth. l. c. = P. tortilis.
undulata, Korth. l. c. 247 = P. undulata.

STREBLUS, Lour. Fl. Cochinch. 614 (1790). *URTICACEAE*, Benth. & Hook. f. iii. 359.
ALBRANDIA, Gaudich. in Freyc. Voy. Bot. 509 (1826).
CALIUS, Blanco, Fl. Filip. ed. I. 698 (1837).
EPICARPURUS, Blume, Bijdr. 488 (1825).
asper, *Lour. Fl. Cochinch.* 615.—As. trop.
Brunoniana, F. Muell. Fragm. vi. 192 = Pseudomorus Brunoniana.
cordatus, Lour. Fl. Cochinch. 615 = Trophis cordata.
lactescens, Blume, Mus. Bot. Lugd.-Bat. ii. 80 = asper.
macrophyllus, Blume, l. c. (= *Diplocos macrophylla*).—Ins. Celebes.
mauritianus, Blume, l. c. = Ampalis madagascariensis.
microphylla, Kurz, For. Fl. Brit. Burma, ii. 465 = Phyllochlamys spinosa.
mitis, *Kurz, l. c.* 464.—Burma.
pendulina, F. Muell. Fragm. vi. 192 = Pseudomorus Brunoniana.
rectinervis, F. Muell. l. c. = Aphanantha philippinensis.
taxoides, Kurz, For. Fl. Brit. Burma, ii. 465 = Phyllochlamys spinosa.
zeylanica, Kurz, l. c. 464 = Taxotrophis zeylanica.

STRECKERA, Sch. Bip. in Flora, xvii. (1834) 483 = **Leontodon**, Linn. (Compos.).
Claurae, Sch. Bip. l. c. 487 = L. asperrimus.
tuberosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 197 = L. tuberosus.

STRELESKIA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 266 = **Wahlenbergia**, Schrad. (Campan.).
montana, Hook. f. l. c. 267 = W. saxicola.

STRELITSIA, Thunb. Fl. Cap. ii. 113 (1818), sphalm. = seq.

STRELITZIA, [Banks] Ait. Hort. Kew. ed. I. i. 285. t. 2 (1789). *SCITAMINEAE*, Benth. & Hook. f. iii. 656.
angusta, D. Dietr. Syn. Pl. i. 850, sphalm. = S. Augusta.

STRELITZIA:—
angustifolia, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 55 = parvifolia.
Augusta, *Thunb. Nov. Gen.* 113.—Afr. austr.
farinosa, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 55 = S. Reginae.
gigantea, J. S. Kern. Hort. Semperv. t. 589 = S. Augusta?
glauca, Rich. in Nov. Act. Nat. Cur. xv. (1831) Suppl. 17. t. 2–3 = S. Reginae.
humilis, Link, Enum. Hort. Berol. i. 150; Handb. i. 234 = S. Reginae.
juncea, Link, Enum. l. c.; Andr. ex Spreng. Syst. i. 833 = parvifolia.
lanceolata, Hort. ex Steud. Nom. ed. II. ii. 645 (Quid?).—Afr. austr.
Lemoinieri, Miellez, in Fl. des Serres, xxiii. (1880) sub t. 2370 = S. Reginae.
macrophylla, Hort. ex Steud. Nom. ed. II. ii. 645 (Quid?).—Afr. austr.
Nicolai, *Regel & C. Koch, in Gartenfl.* (1858) 265. t. 235.—Afr. austr.
ovata, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 55 = S. Reginae.
parvifolia, [*Dryand.*] *l. c.*—Afr. austr.
Principis, Andr. ex Spreng. Syst. i. 833 = parvifolia.
prolifera, Carr. in Rev. Hortic. (1869) 159 = S. Reginae.
pumila, Hort. ex Planch. in Fl. des Serres, xi. (1856) 53 = S. Reginae.
Quensoni, *Lem. Illustr. Hortic.* x. (1863) *Misc.* 20.—Afr. austr.
regalis, Salisb. Prod. 145 = S. Reginae.
Reginae, [*Banks, in*] *Ait. Hort. Kew.* ed. I. i. 285. t. 2; *Thunb. Nov. Gen.* 113.—Afr. austr.
rutilans, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) t. 53 = S. Reginae.
teretifolia, Barrow, ex Steud. Nom. ed. II. ii. 645 (Quid?).—Afr. austr.

STREMPELIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 180. t. 18. f. 2 = **Psychotria**, Linn. (Rubiac.).
guianensis, A. Rich. l. c.—Guian.

STREMPELIOPSIS, Benth. in Benth. & Hook. f. Gen. ii. 702 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
strempelioides, *Benth. l. c.*—Cuba.

STREPHIUM, Schrad. ex Nees, Agrost. Bras. 298 (1829) = **Olyra**, Linn. (Gramin.).
distichophyllum, Schrad. ex Kunth, Enum. Pl. i. 70 = O. polypodioides.
floribundum, Nees, ex Steud. Syn. Pl. Gram. 36 = O. floribunda.
pulchellum, Wright, in Sauv. Fl. Cub. 193.—Cuba.
strictiflorum, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 465 = O. strictiflora.

STREPHONEMA, Hook. f. in Benth. & Hook. f. Gen. i. 782 (1867). *LYTHRARIEAE*, Benth. & Hook. f. l. c.
Mannii, *Hook. f. l. c.*—Afr. trop.
sericea, *Hook. f. l. c.*—Afr. trop.

STREPSANTHERA, Rafin. Fl. Tellur. iv. 13 (1836) = **Anthurium**, Schott (Aroid.).
macrophyla, Rafin. l. c. = A. grandifolium.

STREPSIA, Steud. Nom. ed. II. ii. 645 (1841) = **Tillandsia**, Linn. (Bromel.).
usneoides, Steud. l. c. = T. usneoides.

STREPSILOBUS, Rafin. Sylva Tellur. 117 (1838) = **Entada**, Adans. (Legum.).
scandens, Rafin. l. c. 118 = E. scandens.

STREPSIMELA, Rafin. Sylva Tellur. 159 (1838) = **Loranthus**, Linn.
coccinea, Rafin. l. c. = L. coccineus.

STREPSIPHUS, Rafin. Fl. Tellur. ii. 91 (1836) = **Peristrophe**, Nees (Acanth.).
speciosus, Rafin. l. c. = P. speciosa.

STREPSIPHYLA, Rafin. Fl. Tellur. iii. 60 (1836) = **Drimia**, Jacq. (Liliac.).
undulata, Rafin. l. c. = Scilla undulata.
villosa, Rafin. l. c. = D. villosa.

STREPTACHNE, R. Br. Prod. 174 (1810) = **Stipa**, Linn. (Gramin.).
Calamagrostis, Dum. Obs. Gram. Belg. 135 = Stipa Calamagrostis.
cubensis, A. Rich. Fl. Cub. Fanerog. ii. 311.—Cuba.
floridana, Chapm. Fl. S. U. St. 554 = Aristida floridana.
indica, Buese, ex De Vriese, Pl. Ind. Bat. Or. 99.—Malaya.
ramosissima, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 7 = Stipa micrantha.
stipoides, R. Br. Prod. 174.—Austral.
verticillata, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 7 = Stipa micrantha.

STREPTACHNE, H. B. & K. Nov. Gen. et Sp. i. 124. t. 40 (1815) = **Aristida**, Linn. (Gramin.).
domingensis, Spreng. ex Schult. Mant. ii. 188 = Andropogon hirtiflorus.
pilosa, H. B. & K. Nov. Gen. et Sp. i. 124 = A. jorullensis.
scabra, H. B. & K. l. c. t. 40 = A. scabra.
tenuis, H. B. & K. l. c. = A. tenuis, *Kunth*.

STREPTALON, Rafin. Fl. Tellur. iii. 80 (1836) = **Hypericum**, Tourn.
dolabriforme, Rafin. l. c. = H. dolabriforme.

STREPTANTHERA, Sweet, Brit. Flow. Gard. Ser. I. iii. t. 209 (1827). *IRIDEAE*, Benth. & Hook. f. iii. 703.
cuprea, *Sweet, Hort. Brit.* ed. II. 501.—Afr. austr.
elegans, *Sweet, Brit. Flow. Gard.* Ser. I. iii. t. 209.—Afr. austr.
lineata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 390 = Sparaxis grandiflora.
tricolor, Klatt, l. c. = Sparaxis tricolor.

STREPTANTHUS, Nutt. in Journ. Acad. Philad. v. (1825) 134. t. 7. *CRUCIFERAE*, Benth. & Hook. f. i. 69.
angustifolius, A. Gray, in Proc. Acad. Philad. 1863 (1864) 57 = Arabis Drummondii.
angustifolius, Nutt. ex Torr. & Gray, Fl. N. Am. i. 76 = Arabis Holboellii.
arcuatus, Nutt. l. c. 77 = Arabis arcuata.
bracteatus, *A. Gray, Gen. Am. Bor.-or. Illustr.* i. 146.—Am. bor. occ.
brazoënsis, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 448 = Arabis petiolaris.
Breweri, *A. Gray, in Proc. Calif. Acad.* iii. (1864) 101.—Calif.
carinatus, *Wright, ex A. Gray, Pl. Wright.* ii. 11.—N. Mexic.
cordatus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 77.—Am. bor. occ.
cordatus, Torr. in Pacif. Rail. Rep. iv. 65 = tortuosus.
crassicaulis, Torr. in Stansbury, Exped. Utah, 383. t. 9 = Caulanthus crassicaulis.
diversifolius, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 363.—Calif.
flavescens, A. Gray, l. c. vi. (1864) 186 = Caulanthus procerus.
flavescens, *Hook. Ic. Pl.* t. 44.—Calif.
flavescens, Torr. in Pacif. Rail. Rep. iv. 65 = Thelypodium flavescens.
glabrifolius, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 448 = hyacinthoides.
glandulosus, *Hook. Ic. Pl.* t. 40.—Am. bor. occ.
heterophyllus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 77 = Caulanthus Coulteri.
hispidus, *A. Gray, in Proc. Calif. Acad.* iii. (1864) 101.—Calif.
Howellii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 353.—Am. bor. occ.
hyacinthoides, *Hook. Bot. Mag.* t. 3516.—Am. bor. occ.
linearifolius, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 7 = Thelypodium linearifolium.
longifolius, Benth. Pl. Hartw. 10 = Thelypodium longifolium.
maculatus, *Nutt. in Journ. Acad. Philad.* v. (1825) 134. t. 7.—Am. bor. occ.
micranthus, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 7 = Thelypodium longifolium.
obtusifolius, Hook. Bot. Mag. t. 3317 = maculatus.
petiolaris, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 7 = Arabis petiolaris.
platycarpus, *A. Gray, Pl. Wright.* ii. 10.—Am. bor. occ.

STREPTANTHUS:—
polygaloides, *A. Gray, in Proc. Am. Acad.* i. (1865) 519.—Calif.
procerus, Brewer, ex A. Gray, l. c. = Caulanthus procerus.
repandus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 77 (Arabidis sp. ?).—Calif.
sagittatus, Nutt. in Journ. Acad. Philad. vii. (1834) 12 = Thelypodium Nuttallii.
tortuosus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 152. f. 46.—Calif.
virgatus, Nutt. ex Torr. & Gray. Fl. N. Am. i. 76 = Arabis Holboellii.

STREPTIMA, Rafin. Fl. Tellur. ii. 93 (1836) = **Frankenia**, Linn.
pauciflora, Rafin. l. c. = F. pauciflora.

STREPTIUM, Roxb. Pl. Corom. ii. 25. t. 146 (1798) = **Priva**, Adans. (Verben.).
asperum, Roxb. l. c. = P. leptostachya.

STREPTOCALYX, Beer, in Flora, xxxvii. (1854) 348. *BROMELIACEAE*, Benth. & Hook. f. iii. 659.
Furstenbergi, *E. Morr. in Belg. Hortic.* (1883) 13.—Bras.
Poeppigii, *Beer, Bromel.* 141.—Bras.
Vallerandi, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 13. t. 1–2.—Bras.

STREPTOCARPUS, Lindl. Bot. Reg. t. 1173 (1828). *GESNERACEAE*, Benth. & Hook. f. ii. 1023.
angovensis, *DC. Prod.* ix. 271.—Madag.
benguellensis, *Welw. ex C. B. Clarke, in DC. Monog. Phan.* v. 150.—Afr. trop.
biflorus, Crocker, in Journ. Linn. Soc. v. (1861) 66. t. 4 = Gardenii.
Bojeri, *R. Br. in Benn. Pl. Jav. Rar.* 119.—Madag.
caulescens, *Vatke, in Linnaea*, xliii. (1880–82) 323.—Afr. trop.
Cooperi, *C. B. Clarke, in DC. Monog. Phan.* v. 150.—Afr. austr.
Fanninii, *Harv. ex C. B. Clarke, l. c.*—Afr. austr.
floribunda, Hort.—Cf. Gard. Chron. (1878) II. 282 (an hybr.?).—Hab. ?
Gardenii, *Hook. Bot. Mag.* t. 4862.—Afr. austr.
Greenii ×, Hort.—Cf. Gard. Chron. (1876) I. 74.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 324.—Madag.
Hilsenbergii, *R. Br. in Benn. Pl. Jav. Rar.* 119 (Helsingbergii).—Madag.
Kirkii, *Hook. f. Bot. Mag.* t. 6782.—Afr. trop.
lutea, *C. B. Clarke, in DC. Monog. Phan.* v. 153.—Afr. austr.
monophylla, *Welw. in Bibl. Univ. Genèv.* (1861) 202.—Afr. trop.
muscosa, *C. B. Clarke, in DC. Monog. Phan.* v. 155.—Madag.
nobilis, *C. B. Clarke, l. c.*—Afr. trop.
paniculata, *R. Br. in Benn. Pl. Rar. Jav.* 119.—Madag.
parviflorus, *E. Mey. in Drège, Zwei Pfl. Docum.* 152.—Afr. austr.
plantaginea, *Vatke, in Linnaea*, xliii. (1880–82) 323.—Ins. Comor.
polyanthus, *Hook. Bot. Mag.* t. 4850.—Afr. austr.
primuloides, *Dickie, in Journ. Linn. Soc.* ix. (1867) 126, nomen.—Hab. ?
pusillus, *Harv. ex C. B. Clarke, in DC. Monog. Phan.* v. 149.—Afr. austr.
Rexii, *Lindl. Bot. Reg.* t. 1173.—Afr. austr.
Saundersii, *Hook. Bot. Mag.* t. 5251.—Afr. austr.
Thompsonii, *R. Br. in Benn. Pl. Jav. Rar.* 119.—Madag.

STREPTOCAULON, Wight & Arn. Contrib. 64 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 744.
TRIPLOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 251.
abyssinicum, Hochst. ex Steud. Nom. ed. II. ii. 305, in syn.; 646 = Periploca linearifolia.
albicans, G. Don, Gen. Syst. iv. 162 = Periploca albicans.
Baumii, *Decne. in DC. Prod.* viii. 496.—Ins. Philipp.
calophyllum, Wight, in Wight & Arn. Contrib. 65 = Periploca calophylla.

STREPTOCAULON:—
chinense, G. Don, Gen. Syst. iv. 162 = Vallaris sinensis.
cochinchinense, G. Don, l. c. (Pergulariae sp. ?).—Cochinch.
Cumingii, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 132.—Ins. Philipp.
divaricatum, G. Don, Gen. Syst. iv. 162 = Strophanthus divergens.
extensum, Wight, in Wight & Arn. Contrib. 65 = Myriopteron paniculatum.
Griffithii, *Hook. f. Fl. Brit. Ind.* iv. 10.—Burma.
Hamiltonii, Wight, in Wight & Arn. Contrib. 64 = Decalepis Hamiltonii.
Horsfieldii, Miq. Fl. Ind. Bat. ii. 470 = Myriopteron paniculatum.
Kleinii, *Wight & Arn. Contrib.* 65.—Ind. or.
mauritianum, G. Don, Gen. Syst. iv. 162 = Tylophora laevigata.
obtusum, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 252 = Baumii.
parviflorum, G. Don, Gen. Syst. iv. 162 = Periploca parviflora.
sylvestre, *Wight, in Wight & Arn. Contrib.* 65.—Ind. or.
tomentosum, *Wight, l. c.* 64.—Burma; China.
virgatum, G. Don, Gen. Syst. iv. 162 (= *Orthosia virgata*).—Bras.
Wallichii, *Wight, in Wight & Arn. Contrib.* 65.—Malaya.

STREPTOCHAETA, Schrad. ex Nees, Agrost. Bras. 536 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1111.
LEPIDEILEMA, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 93.
spicata, *Schrad. ex Nees, Agrost. Bras.* 537.—Bras.

STREPTODESMIA, A. Gray, Bot. U. St. Expl. Exped. i. 427. t. 47 (1854) = **Adesmia**, DC. (Legumin.).
canescens, A. Gray, l. c. = Patagon.

STREPTOGLOSSA, Steetz, ex F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 491 = **Pterigeron**, A. Gray (Compos.).
Steetzii, F. Muell. l. c. = P. decurrens.

STREPTOGYNE, Beauv. Agrost. 80. t. 16. f. 8. (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1191.
crinita, *Beauv. l. c.*—Am. bor.; Ind. occ.
guianensis, Rich. ex Roem. & Schult. Syst. ii. 614 = crinita.

STREPTOLIRION, Edgew. in Proc. Linn. Soc. i. (1845) 254; et in Trans. Linn. Soc. xx. (1846) 90. t. 2. *COMMELINACEAE*, Benth. & Hook. f. iii. 852.
volubile, *Edgew. ll. cc.*—Reg. Himal.

STREPTOLOMA, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 155. *CRUCIFERAE*, Benth. & Hook. f. i. 77.
desertorum, *Bunge, l. c.* 156.—Turkest.

STREPTOPETALUM, Hochst. in Flora, xxiv. (1841) 665. *TURNERACEAE* [Benth. & Hook. f. i. 807].
Hildebrandtii, *Urb. in Eichl. Jahrb. Berl.* ii. (1883) 57.—Afr. trop.
serratum, *Hochst. in Flora*, xxiv. (1841) 666.—Abyss.

STREPTOPUS, Michx. Fl. Bor. Am. i. 200. t. 18 (1803). *LILIACEAE*, Benth. & Hook. f. iii. 769.
HEXORIMA, Rafin. in Med. Repos. N. York, v. (1808) 351.—Cf. Desv. Journ. Bot. ii. (1809) 168.
KRUHSEA, Regel, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 122.
ajanensis, *Tiling, ex Regel & Tiling, l. c.* 122.—Sibir.; Am. bor.
alpina, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = *Smilacina alpina*.
amplexicaulis, Poir. Encyc. vii. 467, sphalm. = seq.
amplexifolius, DC. Fl. Fr. iii. 174 = distortus.
borealis, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = *Smilacina borealis*.

STREPTOPUS:—
brevipes, *Baker, in Journ. Linn. Soc.* xiv. (1875) 592.—Am. bor. occ.
candidus, Wall. Cat. n. 5111 = simplex.
chinensis, Sm. in Rees, Cycl. xxxvii. sub Uvularia, n. 4 = Disporum pullum.
distortus, *Michx. Fl. Bor. Am.* i. 200; *Bess. Prim. Fl. Galic.* i. 234.—Europ.; Am. bor.
fusca, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = Smilacina fusca.
lanuginosus, Michx. Fl. Bor. Am. i. 201 = Disporum lanuginosum.
maculatus, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 170 (Dispori sp.).—Am. bor.
multiflorus, D. Dietr. Syn. Pl. ii. 1121 = Disporum pullum.
pallida, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = *Smilacina pallida*.
pedunculatus, Sm. in Rees, Cycl. xxxvii. sub Uvularia, n. 5 = Disporum pullum.
purpurea, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = Smilacina purpurea.
roseus, *Michx. Fl. Bor. Am.* ii. 201. t. 18.—Am. bor.
roseus, Wright, ex S. Wats. in Proc. Am. Acad. xiv. (1879) 269 = ajanensis.
simplex, *D. Don, Prod. Fl. Nep.* 48.—Reg. Himal.
undulatus, Rafin. in Am. Monthly Mag. (1811) 356, nomen = distortus ?

STREPTORHAMPHUS, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) II. 176, sphalm. = STEPTORHAMPHUS, Bunge = **Lactuca**, Tourn. (Compos.).

STREPTOSEMA, Presl, Bot. Bemerk. 128 (1844); in Abh. Boehm. Ges. V. iii. (1845) 558 = **Aspalathus**, Linn. (Legumin.).
Histrix, Presl, Bot. Bemerk. 128 = A. Hystrix.
verrucosa, Presl, l. c. = A. verrucosa.

STREPTOSOLEN, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 208. *SOLANACEAE*, Benth. & Hook. f. ii. 910.
Benthami, Miers, Illustr. S. Am. Pl. ii. 70 = Jamesonii.
Jamesonii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. v. (1850) 208.—N. Granat.

STREPTOSTACHYS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = **Panicum**, Linn. (Gramin.).
asperifolia, Desv. l. c. = P. Streptostachys.
hirsuta, Beauv. Agrost. t. 10. f. 11 = P. Streptostachys.

STREPTOSTIGMA, Regel, Gartenfl. (1853) 322. t. 68 = **Cacabus**, Bernh. (Solan.).
Warszewiczii, Regel, l. c.—Peruv.

STREPTOSTIGMA, Thw. in Hook. Kew Journ. vi. (1854) 298. t. 9 A = **Harpullia**, Roxb. (Sapind.).
viridiflora, Thw. l. c. = H. cupanoides.

STREPTOTHAMNUS, F. Muell. Fragm. iii. 27 (1862). *BIXINEAE*, Benth. & Hook. f. i. 972.
Becklerii, *F. Muell. l. c.* 28.—Austral.
Moorii, *F. Muell. l. c.*—Austral.

STREPTYLIS, Rafin. Fl. Tellur. iv. 122 (1836) = **Aneilema**, R. Br. (Commel.).
bracteolata, Rafin. l. c. = A. spiratum.

STRIGA, Lour. Fl. Cochinch. 22 (1790). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 968.
CAMPULEIA, Thou. Gen. Nov. Madag. 7 (1806).
MICROSYPHUS, Presl, Bot. Bemerk. 91 (1844).
PSAMMOSTACHYS, Presl, l. c. (1844).
aspera, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 362.—Afr. trop.
Campuleia, Steud. Nom. ed. II. ii. 646 = hermonthica.
coccinea, Benth. in Hook. Comp. Bot. Mag. i. (1835) 364 = lutea.
curviflora, *Benth. l. c.* 362.—Austral.
densiflora, *Benth. l. c.* 363.—Ind. or.
elegans, *Benth. l. c.*—Afr. austr.
euphrasioides, *Benth. l. c.* 364.—Ind. or.
Eustriga, Steud. Nom. ed. II. ii. 546 = lutea.

STRIGA:—
flava, *Miq. Fl. Ind. Bat.* ii. 704.—Sumatra.
Forbesii, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 364.—Afr. austr.
glabrata, Benth. l. c = euphrasioides.
hermonthica, *Benth. l. c.* 365.—Afr. bor. et trop.
hirsuta, Benth. in DC. Prod. x. 502 = lutea.
humifusa, Benth. in Hook. Comp. Bot. Mag. i. (1835) 362 = Cycnium humifusum.
humilis, Hochst. ex Benth. in DC. Prod. x. 504 = Cycnium humifusum.
latericea, *Vatke, in Linnaea*, xliii. (1880–82) 311.—Afr. trop.
lutea, *Lour. Fl. Cochinch.* 22.—As. et Afr. trop.
macrantha, *Benth. in DC. Prod.* x. 503.—Afr. trop.
Masuria, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 364.—Afr. austr.
micrantha, *A. Rich. Tent. Fl. Abyss.* ii. 130.—Abyss.
multiflora, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 363.—Austral.
orchidea, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = orobanchoides.
orobanchoides, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 361. t. 19.—Ind. or.; Afr. trop.
parviflora, *Benth. l. c.* 362.—Austral.
parvula, Miq. in Journ. Bot. Néerl. i. (1861) 113 = lutea.
phoenicea, Benth. Scroph. Ind. 40 = lutea.
pubiflora, *Klotzsch, in Peters, Reise Mossamb. Bot.* 227.—Afr. trop.
pusilla, Hochst. ex Benth. in DC. Prod. x. 503 = lutea.
Schimperiana, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 24, nomen.—Abyss.
senegalensis, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 363.—Afr. trop.
Spanogheana, *Miq. Fl. Ind. Bat.* ii. 704.—Ins. Timor. *vel* Java.
sulphurea, *Dalz. in Gibs. Bomb. Fl.* 182.—Ind. or.
Thunbergii, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 363.—Afr. austr.
zangebarica, Klotzsch, in Peters, Reise Mossamb. Bot. t. 35 = hirsuta.
zanzibarensis, *Vatke, in Linnaea*, xliii. (1880–82) 310.—Afr. trop.

STRIGILIA, Cav. Diss. vii. 358. t. 201 (1789) = **Styrax**, Linn.
acuminata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 282 = Styr. acuminatum.
alba, Miers, l. c. = Styr. album.
argentea, Miers, l. c. = Styr. argenteum.
aurea, Miers, l. c. = Pamphilia aurea.
camporum, Miers, l. c. = Styr. camporum.
chrysastera, Miers, l. c. = Styr. chrysasterum.
cordata, Poir. Encyc. vii. 469 = Styr. cordatum.
crocea, G. Don, Gen. Syst. i. 679.—Peruv.
ferruginea, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 283 = Styr. ferrugineum.
florida, Miers, l. c. 282 = Styr. floridum.
Gardneriana, Miers, l. c. = Styr. Gardnerianum.
glabrata, Miers, l. c. 283 = Styr. glabratum.
guayanensis, Miers, l. c. 282 = Styr. glabrum.
laevis, Miers, Contrib. Bot. i. t. 30 = Styr. leiophyllum.
latifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 282 = Styr. latifolium.
leiophylla, Miers, l. c. 283 = Styr. leiophyllum.
leprosa, Miers, l. c. = Styr. leprosum.
longiflora, Miers, l. c. 282 = Styr. longiflorum.
macrophylla, Miers, l. c. 283 = Styr. macrophyllum.
nervosa, Miers, l. c. 282 = Styr. reticulatum.
obliquinervia, Miers, l. c. = Styr. chrysasterum.
oblonga, Poir. Encyc. vii. 469 = Styr. oblongum.
ovata, Poir. l. c. = Styr. ovatum.
pallida, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 282 = Styr. pallidum.
parvifolia, Miers, l. c. = Styr. parvifolium.
pauciflora, Miers. l. c. 283 = Styr. pauciflorum.
Pavonii, Miers, l. c. = Styr. Pavonii.
Pohlii, Miers, l. c. 282 = Styr. Pohlii.
psilophylla, Miers, l. c. 283 = Styr. psilophyllum.
punctata, Miers, l. c. 282 = Styr. punctatum.
racemosa, Cav. Diss. vii. 358 = Styr. racemosum.
reticulata, Miers, in Ann. & Mag. Nat. Hist. Ser. III. iii. (1859) 282 = Styr. reticulatum.
subcordata, Miers, l. c. = Styr. subcordatum.
tomentosa, Miers, l. c. 283 = Styr. tomentosum.

STRIGOSELLA, Boiss. Diagn. Ser. II. iv. 22 (1853) = **Malcomia**, R. Br. Crucif.).
cabulica, Boiss. l. c. = M. cabulica.

STROBIDIA, Miq. Fl. Ind. Bat. Suppl. 614 (1860). *SCITAMINEAE*, Benth. & Hook. f. iii. 647.
sumatrana, *Miq. l. c.*—Sumatra.

STROBILA, G. Don, Gen. Syst. iv. 327 (1837) = **Arnebia**, Forsk. (Borag.).
hispidissima, G. Don, l. c. = A. hispidissima.

STROBILA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Quid?).

STROBILACANTHUS, Griseb. in Bonplandia, vi. (1858) 10. *ACANTHACEAE*, Benth. & Hook. f. ii. 1096.
lepidospermus, *Griseb. l. c.*—Panama.

STROBILANTHES, Blume, Bijdr. 781, 796 (1826). *ACANTHACEAE*, Benth. & Hook. f. ii. 1086.
ADENACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75, 196 (1832).
BUTERAEA, Nees, l. c. (1832).
DIDYPLOSANDRA, Wight, Ic. iv. iv. 3. t. 1515 (1850).
ENDOPOGON, Nees, in Wall. Pl. As. Rar. iii. 76 (1832).
GOLDFUSSIA, Nees, l. c. 196 (1832).
GUTZLAFFIA, Hance, in Hook. Kew Journ. i. (1849) 142.
LEPTACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75, 169 (1832).
MACKENZIEA, Nees, in DC. Prod. xi. 308 (1847).
PHLEBOPHYLLUM, Nees, in Wall. Pl. As. Rar. iii. 75, 102 (1832).
TRIAENANTHUS, Nees, in DC. Prod. xi. 169 (1847).
acrocephalus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 473.—Reg. Himal.
acuminatus, *T. Anders. l. c.*—Burma.
acuminatus, T. Anders. l. c., partim = longipes.
adenophorus, *Bedd. Ic. Pl. Ind. Or.* i. 53. t. 225.—Ind. or.
adenophorus, Nees, in DC. Prod. xi. 182 = lanceolatus.
adnatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 436.—Reg. Himal.
agrestis, *C. B. Clarke, l. c.* 466.—Reg. Himal.
alatus, *Blume, Bijdr.* 798.—Java.
alatus, *Nees, in DC. Prod.* xi. 194.—Reg. Himal.
alatus, Steud. Nom. ed. II. ii. 646 = Wallichii.
amabilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 476.—Ind. or.
amplectens, Nees, in Wall. Pl. As. Rar. iii. 86 = auriculatus.
anceps, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Ind. or.
ancolanus, *Miq. Fl. Ind. Bat.* ii. 799.—Sumatra.
Andersonii, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 55.—Ind. or.
angustifrons, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 466.—Reg. Himal.
anisophyllus, *T. Anders. in Cat. Pl. Hort. Bot. Calc.* 43; *et in Journ. Linn. Soc.* ix. (1867) 478.—Reg. Himal.
anisostachys, Boj. ex Nees, in DC. Prod. xi. 368 (= *Anisostachya Bojeri*).—Madag.
apricus, *T. Anders. ex Benth. Fl. Hongk.* 262.—China.
arboreus, *Span. in Linnaea*, xv. (1841) 328.—Ins. Timor.
argutus, Nees, in DC. Prod. xi. 188 = sexennis.
Arnottianus, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Zeylan.
asper, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 385 = scaber.
asper, *Wight, Ic.* t. 1518.—Ind. or.
asperrimus, Dalz. & Gibs. Bomb. Fl. 187 = Heyneanus.
asperrimus, *Nees, in DC. Prod.* xi. 183.—Zeylan.
atropurpureus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Reg. Himal.
attenuatus, Jacq. ex Nees, in DC. Prod. xi. 193 = alatus.
auriculatus, *Nees, in Wall. Pl. As. Rar.* iii. 86. t. 195.—Ind. or.; Burma.
barbatus, *Nees, l. c.* 85.—Ind. or.

STROBILANTHES:—
Beddomei, T. Anders. in Journ. Linn. Soc. ix. (1867) 482 = gracilis.
bibracteatus, *Blume, Bijdr.* 799.—Java.
biceps, Miq. Fl. Ind. Bat. ii. 804 = capitatus.
boërhaavioides, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 479.—Reg. Himal.
bolumpattianus, *Bedd. Ic. Pl. Ind. Or.* i. 46.—Ind. or.
Brandisii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 475.—Burma.
Brunonianus, *Nees, in Wall. Pl. As. Rar.* iii. 87.—Reg. Himal.
burmanicus, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 92 = Hemigraphis elegans.
callosus, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Ind. or.
callosus, Nees, l. c., partim = zeylanicus.
calycinus, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Zeylan.
campanulatus, Wight, Ic. t. 1562 = Heyneanus.
canaricus, *Bedd. Ic. Pl. Ind. Or.* i. 50.—Ind. or.
capitatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 475.—Reg. Himal.; Burma.
caudatus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 228.—Ind. or.
cerinthoides, Nees, in DC. Prod. xi. 724 = sexennis.
cernuus, *Blume, Bijdr.* 797.—Java.
Championi, T. Anders. ex Benth. Fl. Hongk. 261 = flaccidifolius.
ciliatus, T. Anders. in Journ. Linn. Soc. ix. (1867) 468, partim = warreensis.
ciliatus, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Ind. or.
collinus, *Nees, l. c.* 86.—Ins. Penang.
coloratus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 481.—Reg. Himal.
coloratus, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 312 = trifidus, calycinus.
consanguineus, T. Anders. in Thw. Enum. Pl. Zeyl. 226, partim = exareolatus.
consanguineus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 435.—Ind. or.
crassifolius, *Miq. Fl. Ind. Bat.* ii. 800.—Sumatra.
crataegifolius, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 477.—Burma.
crenulatus, Boj. ex Nees, in DC. Prod. xi. 369 = Justicia tenella.
crenulatus, Hassk. Cat. Hort. Bog. Alt. 148 = hirtus.
crinitus, T. Anders. in Journ. Linn. Soc. ix. (1867) 481 = coloratus.
crispus, T. Anders. l. c. = phyllostachyus.
crispus, *Blume, Bijdr.* 798.—Java.
cuspidatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 465.—Ind. or.
Dalhousianus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 460.—Reg. Himal.
Dalzellii, T. Anders. ex C. B. Clarke, l. c. 443 = lupulinus.
dasyspermus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 94.—Burma.
debilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 467.—Burma.
decurrens, T. Anders. in Journ. Linn. Soc. ix. (1867) 470 = adnatus.
decurrens, *Nees, in DC. Prod.* xi. 189.—Ind. or.
deflexus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 227.—Zeylan.
denticulatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 483.—Reg. Himal.
diclipteroides, *Miq. Fl. Ind. Bat.* ii. 802.—Java.
dimorphotrichus, *Hance, in Journ. Bot.* xxi. (1883) 355.—China.
discolor, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 477.—Reg. Himal.
divaricatus, *T. Anders. l. c.* 478.—Reg. Himal.
Dupeni, *Bedd. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 453.—Ind. or.
durus, T. Anders. ex C. B. Clarke, l. c. 447 = phyllostachyus.
echinatus, Nees, in Wall. Pl. As. Rar. iii. 85 = pectinatus.
Edgeworthianus, Nees, in DC. Prod. xi. 190 = auriculatus.
elatus, *Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 53.—Malaya.
elegans, *Wall. ex Voigt, Hort. Suburb. Calc.* 484, nomen.—Munipore.
elongatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 470.—Reg. Himal.

STROBILANTHES:—
eriocephalus, T. Anders. in Journ. Linn. Soc. ix. (1867) 466 = ixiocephalus.
erosus, *Nees, in DC. Prod.* xi. 180.—Java.
exareolatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 432.—Zeylan.
exsertus, *C. B. Clarke, l. c.* 445.—Zeylan.
extensus, Bedd. Ic. Pl. Ind. Or. i. 47. t. 202 = Newii.
extensus, *Nees, in DC. Prod.* xi. 195.—Reg. Himal.
Falconeri, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 484.—Burma.
farinosus, *C. B. Clarke, in Hook. f. Fl. Ind.* iv. 470.—Burma.
filiformis, *Blume, Bijdr.* 800.—Java.
fimbriatus, T. Anders. in Journ. Linn. Soc. ix. (1867) 470, partim = macrostegius.
fimbriatus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Reg. Himal.
flaccidifolius, *Nees, in DC. Prod.* xi. 194.—Ind. or.; Burma; China.
flaccidus, G. Mann, Assam For. Rep. (1876–77) par. 135 = flaccidifolius.
flavus, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 78 = Hemigraphis flava.
foetidissimus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 93.—Burma.
foliosus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 467.—Ind. or.
formosanus, *S. Moore, in Journ. Bot.* xv. (1877) 294.—Ins. Formosa.
Gardnerianus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 226.—Zeylan.
geniculatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 459.—Reg. Himal.
glabratus, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Reg. Himal.
glandulosus, *Blume, Bijdr.* 800.—Java.
glandulosus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 92.—Ins. Andaman.
glaucescens, Nees, in Wall. Pl. As. Rar. iii. 85; Cat. n. 7155 = Hemigraphis glaucescens.
glomeratus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 475.—Reg. Himal.; Burma.
glutinosus, J. Grah. Cat. Pl. Bomb. 162 = ixiocephalus.
glutinosus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Reg. Himal.
Goldfussia, Dalz. & Gibs. Bomb. Fl. Suppl. 71 (excl. syn.) = isophyllus.
gossypinus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 466.—Ind. or.
gracilis, T. Anders. l. c. 474 = geniculatus.
gracilis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 55.—Ind. or.
Grahamianus, Wight, Ic. t. 1520 = callosus.
Griffithianus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 481.—Reg. Himal.
haplanthoides, *T. Anders. l. c.* 471.—Burma.
Helferi, *T. Anders. l. c.* 472.—Burma.
helicoides, *T. Anders. in Thw. Enum. Pl. Zeyl.* 229.—Zeylan.
Helictus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 479.—Reg. Himal.
heteromallus, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 437.—Ind. or.
Heyneanus, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Ind. or.
hirsutissimus, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 313 = sexennis.
hirsutus, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 386 = Hemigraphis Decaisneana.
hirtus, *Blume, Bijdr.* 797.—Java.
homotropus, *Nees, in DC. Prod.* xi. 187.—Ind. or.
Hookeri, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Zeylan.
hypoleucus, T. Anders. in Thw. Enum. Pl. Zeyl. 226 = rhytisperma.
imbricatus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Burma.
inflatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 476.—Reg. Himal.
interruptus, Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 474 = sexennis.
involucratus, *Blume, Bijdr.* 799.—Java.
isophyllus, *T. Anders. in Cat. Pl. Hort. Bot. Calc.* 43; *et in Journ. Linn. Soc.* ix. (1867) 478.—Ind. or.
ixiocephalus, *Benth. in Flora*, xxxii. (1849) 557.—Ind. or.

STROBILANTHES:—
japonicus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 124.—China; Japon.
jeyporensis, *Bedd. Ic. Pl. Ind. Or.* i. 50.—Ind. or.
karensium, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 94 = Falconeri.
khasyanus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 471.—Reg. Himal.
Kunthianus, *T. Anders. ex Benth. Fl. Hongk.* 262, in nota.—Ind. or.
lachenensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 465.—Reg. Himal.
lactatus, Hook. Bot. Mag. t. 4366 (Ruelliae sp.?).—Bras.
laevigatus, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 467.—Reg. Himal.
laevigatus, *Wall. ex Voigt, Hort. Suburb. Calc.* 484, nomen.—Reg. Himal.
lamiifolius, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 476.—Reg. Himal.
lamioides, *T. Anders. l. c.* 485.—Burma.
lanatus, *Nees, in DC. Prod.* xi. 191.—Ind. or.
lanceifolius, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 480.—Burma.
lanceolatus, *Hook. ex Nees, in DC. Prod.* xi. 181.—Zeylan.
laxus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 228.—Zeylan.
longifolius, *Miq. Fl. Ind. Bat.* ii. 797.—Java.
longipes, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 455.—Burma.
lupulinus, T. Anders. in Journ. Linn. Soc. ix. (1867) 469 = Heyneanus, trifidus.
lupulinus, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Ind. or.
luridus, *Wight, Ic.* t. 1515–16.—Ind. or.
Maclellandi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 453.—Burma.
macrostachya, C. B. Clarke, l. c. 442 = anceps.
macrostegius, *C. B. Clarke, l. c.* 456.—Reg. Himal.
maculatus, T. Anders. in Journ. Linn. Soc. ix. (1867) 473 = petiolaris.
maculatus, *Nees, in DC. Prod.* xi. 190.—Reg. Himal.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 220.—Madag.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 448.—Malaya.
Mastersi, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 481.—Assam.
micranthus, *Wight, Ic.* t. 1519.—Ind. or.
microcarpus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 482.—Burma.
microstachyus, Benth. in Flora, xxxii. (1849) 557 = anceps.
monadelphus, *Nees, in Wall. Pl. As. Rar.* iii. 87.—Reg. Himal.
moschiferus, *Blume, Bijdr.* 800.—Java.
multidens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 461.—Reg. Himal.
mysurensis, Nees, in Wall. Pl. As. Rar. iii. 86 = Asystasia Lawiana.
Neesianus, Wight, Ic. t. 1523 = ixiocephalus.
Neesii, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 93.—Burma.
neglectus, T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 435 = consanguineus.
neilgherrensis, *Bedd. Ic. Pl. Ind. Or.* i. 45.—Ind. or.
Newii, *Bedd. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 464.—Ind. or.
nigrescens, *T. Anders. in Thw. Enum. Pl. Zeyl.* 226.—Zeylan.
nobilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 471.—Reg. Himal.
nutans, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 475.—Reg. Himal.
oliganthus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 124.—China; Japon.
oligocephalus, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 461.—Reg. Himal.
pallidus, T. Anders. in Journ. Linn. Soc. ix. (1867) 470 = decurrens.
Panichanga, *T. Anders. l. c.* 478.—Reg. Himal.
paniculatus, *T. Anders. l. c.* 483.—Zeylan.
paniculatus, Bedd. Ic. Pl. Ind. Or. i. 46. t. 199 = rubicundus.
paniculatus, Miq. Fl. Ind. Bat. ii. 802 = alatus.

STROBILANTHES:—
papillosus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 468.—Ind. or.
parabolicus, *Nees, in DC. Prod.* xi. 180.—Java.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 431.—Burma.
parviflorus, Bedd. Ic. Pl. Ind. Or. i. 45 = warreensis.
paucinervis, *T. Anders. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 470.—Burma.
pauper, *C. B. Clarke, l. c.* 463.—Reg. Himal.
pectinatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 474.—Reg. Himal.
pedunculosus, *Miq. Fl. Ind. Bat.* ii. 803.—Java.
pentstemonoides, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 477.—Reg. Himal.
pentstemonoides, T. Anders. l. c., partim = Dalhousianus.
perfoliatus, *T. Anders. l. c.* 471.—Ind. or.
Perrottetianus, *Nees, in DC. Prod.* xi. 179.—Ind. or.
petiolaris, *Nees, l. c.* 189, partim.—Reg. Himal.
petiolaris, Nees, l. c., partim = Brunonianus.
phyllocaulos, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 464.—Reg. Himal.
phyllostachyus, *Kurz, in Journ. As. Soc. Beng.* xi. (1871) II. 75.—Burma.
plumulosus, Nees, in Wall. Pl. As. Rar. iii. 86 = auriculatus.
polybotryus, *Miq. Fl. Ind. Bat.* ii. 803.—Sumatra.
polycephalus, *Miq. l. c.* 798.—Java.
polythrix, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 470.—Reg. Himal.
pterocaulis, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 93 = imbricatus.
pulcherrimus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 229.—Ind. or.
pulneyensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 438.—Ind. or.
pumilus, *Hassk. Cat. Hort. Bog. Alt.* 310.—Japon.
punctatus, Bedd. Ic. Pl. Ind. Or. i. 47, in obs. = pulneyensis.
punctatus, Nees, in DC. Prod. xi. 183 = anceps.
purpureus, Hook. ex Nees, l. c. = asperrimus.
quadrangularis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 460.—Reg. Himal.
radicans, *T. Anders. in Benth. Fl. Hongk.* 262.—China.
reflexus, Nees, in DC. Prod. xi. 194 = alatus.
remotus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 471.—Tenasserim.
reptans, Boj. ex Nees, in DC. Prod. xi. 368 = Justicia trifolioides.
rhamnifolius, *T. Anders. in Thw. Enum. Pl. Zeyl.* 226.—Zeylan.
rhombifolius, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 461.—Reg. Himal.
rhytispermus, *C. B. Clarke, l. c.* 432.—Zeylan.
roseus, *Nees, in Wall. Pl. As. Rar.* iii. 84.—Burma.
rubescens, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 479.—Reg. Himal.
rubicundus, *T. Anders. l. c.* 483.—Ind. or.
rubicundus, T. Anders. in Thw. Enum. Pl. Zeyl. 229 = paniculatus.
rufescens, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 472.—Burma.
rugosus, Wight, Ic. t. 1619 = Heyneanus.
Sabinianus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Reg. Himal.
salicifolius, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 484.—Reg. Himal.
scaber, T. Anders. in Thw. Enum. Pl. Zeyl. 227, partim = Hemigraphis flava.
scaber, *Nees, in Wall. Pl. As. Rar.* iii. 84.—Ind. or.; Burma.
scrobiculatus, *Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 445.—Ind. or.
secundus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 480.—Reg. Himal.
sessilis, *Nees, in Wall. Pl. As. Rar.* iii. 85.—Ind. or.
sessiloides, Wight, Ic. t. 1512 = praec.
sexennis, Bedd. Ic. Pl. Ind. Or. i. 48. t. 206 = homotropus.
sexennis, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Ind. or.
Simonsii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 474.—Reg. Himal.

STROBILANTHES:—
speciosus, *Blume, Bijdr.* 799.—Java.
spicatus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 484.—Reg. Himal.
stenodon, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 432.—Zeylan.
stenophyllus, *C. B. Clarke, l. c.* 472.—Reg. Himal.
subcapitatus, *C. B. Clarke, l. c.* 455.—Burma.
subnudatus, *C. B. Clarke, l. c.* 472.—Reg. Himal.
subflaccidus, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 94 = acuminatus.
sumatranus, *Miq. Fl. Ind. Bat.* ii. 802.—Sumatra.
tamburensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 454.—Reg. Himal.
Tatei, *F. Muell. in Trans. Roy. Soc. S. Austral.* v. (1882) 81.—Austral.
tener, Nees, in DC. Prod. xi. 184 = trifidus.
tetrapterus, Dalz. in Hook. Kew Journ. ii. (1850) 342 = barbatus.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 799.—Sumatra.
Thomsoni, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 478.—Reg. Himal.
Thwaitesii, *T. Anders. in Thw. Enum. Pl. Zeyl.* 227.—Zeylan.
timorensis, *Nees, in DC. Prod.* xi. 178.—Ins. Timor.
trifidus, Hook. ex Nees, l. c. 186 = calycinus.
trifidus, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Zeylan.
tristis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 470.—Ind. or.
urophyllus, *Nees, in DC. Prod.* xi. 192.—Reg. Himal.
verruculosus, *Nees, l. c.* 181.—Ind. or.
vestitus, *Nees, l. c.* 180.—Zeylan.
violaceus, *Bedd. Ic. Pl. Ind. Or.* i. 48.—Ind. or.
violaefolius, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 485.—Reg. Himal.
virgatus, *Blume, ex Zipp. in Flora*, xii. (1829) 284, nomen.—N. Guin.
viscosus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 226.—Zeylan.
Walkeri, *Arn. ex Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 312.—Zeylan.
Wallichii, *Nees, in Wall. Pl. As. Rar.* iii. 87.—Reg. Himal.
warreensis, *Dalz. in Hook. Kew Journ.* ii. (1850) 341.—Ind. or.
Wightianus, *Nees, in Wall. Pl. As. Rar.* iii. 86.—Ind. or.
Zenkerianus, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 467.—Ind. or.
zeylanicus, *T. Anders. in Thw. Enum. Pl. Zeyl.* 227.—Zeylan.

STROBILANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

STROBILANTHUS, Reichb. Handb. 190 (1837) = praec.

STROBILOCARPOS, Benth. & Hook. f. Gen. iii. 231 (1880) = seq.

STROBILOCARPUS, Klotzsch, in Linnaea, xiii. (1839) 380 = **Grubbia**, Berg. (Santal.).
diversifolius, Klotzsch, l. c. 381 = G. stricta.

STROBILORHACHIS, Klotzsch, in Otto & Dietr. Allg. Gartenz. vii. (1839) 307 = **Aphelandra**, R. Br. (Acanth.).
Blanchetiana, Nees, in Mart. Fl. Bras. ix. 85.—Bras.
glabra, Klotzsch, in Otto & Dietr. Allg. Gartenz. vii. (1839) 308 = A. prismatica.
hirta, Klotzsch, l. c. = A. lutea.
prismatica, Nees, in Mart. Fl. Bras. ix. 84 = A. prismatica.

STROBOCALYX, Sch. Bip. in Pollichia, xviii-xix. (1861) 170 = **Vernonia**, Schreb. (Compos.).
arborea, Sch. Bip. l. c. 171 = V. arborea.
Blumeana, Sch. Bip. l. c. = V. arborea.
celebica, Sch. Bip. l. c. = V. arborea.
Doniana, Sch. Bip. l. c. = V. Doniana.
elaeagnifolia, Sch. Bip. l. c. = V. elaeagnifolia.
elliptica, Sch. Bip. l. c. = V. elaeagnifolia.
glandulosa, Sch. Bip. l. c. = V. glandulosa.
insularum, Sch. Bip. l. c. 170 (= *Monosis insularum*).—Ins. Fiji.

STROBOCALYX:—
javanica, Sch. Bip. in Pollichia, xviii-xix. (1861) 171 = Vernonia arborea.
obovatus, Sch. Bip. l. c. 172 = V. scandens.
pyrrhopappa, Sch. Bip. l. c.—Ins. Philipp.
secundifolia, Sch. Bip. l. c. 171 = V. secundifolia.
solanifolia, Sch. Bip. l. c. = V. solanifolia.
Wightiana, Sch. Bip. l. c. 170 = V. Monosis.

STROBON, Rafin. Sylva Tellur. 132 (1838) = **Cistus**, Tourn., &c.
halimifolium, Rafin. l. c. = Helianthemum halimifolium.
vaginatum, Rafin. l. c. = C. vaginatus.

STROBUS, Opiz, in Lotos, iv. (1854) 94 = **Pinus**, Tourn. (Conif.).
Weymouthiana, Opiz, l. c. = P. Strobus.

STROEMERIA, Roxb. Fl. Ind. ii. 78 (1832), sphalm. = seq.

STROEMIA, Vahl, Symb. Bot. i. 19 (1790) = **Cadaba**, Forsk. (Cappar.).
capparoides, Spreng. Syst. i. 766 = C. capparoides.
dubia, D. Dietr. Syn. Pl. i. 802 = C. farinosa.
farinosa, Vahl, Symb. Bot. i. 20 = C. farinosa.
glandulosa, Vahl, l. c. = C. glandulosa.
longifolia, R. Br. in Salt, Abyss. App. 64 = C. longifolia.
ovalifolia, Zipp. ex Span. in Linnaea, xv. (1841) 164 = C. capparoides.
rotundifolia, Vahl, Symb. Bot. i. 20 = C. rotundifolia.
tetrandra, Vahl, l. c. = C. indica.
trifoliata, Roxb. Hort. Beng. 86; Fl. Ind. ii. 79 = C. trifoliata.
trifoliata, Vahl, ex Schum. & Thonn. Beskr. Guin. Pl. 114 = Euadenia trifoliata.
trinervia, D. Dietr. Syn. Pl. i. 802 = Cleome coluteoides.

STROGANOVIA, Lindl. Veg. Kingd. 355 (1847) = seq.

STROGANOWIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 386. *CRUCIFERAE*, Benth. & Hook. f. i. 88.
brachyota, *Kar. & Kir. l. c.* 387.—Sibir. altaic.
intermedia, *Kar. & Kir. l. c.* xv. (1842) 162.—Sibir. altaic.
paniculata, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 242.—Turkestan.
sagittata, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 387.—Sibir. altaic.

STROMANTHE, Sond. in Hamb. Gartenz. v. (1849) 225. *SCITAMINEAE*, Benth. & Hook. f. iii. 650.
KERCHOVEA, Jorissens, in Belg. Hortic. (1882) 201. t. 8.
MARANTOPSIS, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 97.
amabilis, *Hort. ex E. Morr. in Belg. Hortic.* xxv. (1875) 135.—Bras.
eximia, Eichl. in Berl. Phys. Abh. (1884) 81 = Maranta eximia.
Lubbersiana, E. Morr. in Belg. Hortic. xxxii. (1882) 21. t. 1 (Myrosmatis sp.).—Bras.
lutea, *Eichl. in Berl. Phys. Abh.* (1884) 81.—Venezuela.
Porteana, *A. Gris, in Ann. Sc. Nat.* Sér. IV. ix. (1858) 185.—Bras.
sanguinea, *Sond. in Hamb. Gartenz.* v. (1849) 225.—Ind. or.
Schottiana, *Eichl. in Berl. Phys. Abh.* (1884) 81.—Bras.
Selloi, A. Gris, ex Horan. Prod. Monog. Scitam. 10 = Maranta Selloi.
setosa, A. Gris, in Bull. Soc. Bot. Fr. vi. (1859) 348 = Ctenanthe setosa.
spectabilis, Lem. Jard. Fleur. iv. (1854) t. 401; et Misc. 63 = sanguinea.
Tonckat, *Eichl. in Berl. Phys. Abh.* (1884) 80.—Guian.

STROMATOCACTUS, Karw. ex Foerst. Handb. Cact. ed. II. i. 229 (1885) = **Mammillaria**, Haw. (Cact.).
Kotschoubeyi, Karw. ex Foerst. l. c. 232 (= *Anhalonium sulcatum*).—Mexic.

STROMBOCARPA, A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 243 = **Prosopis**, Linn. (Legumin.).
brevifolia, Nutt. ex A. Gray, Pl. Wright. i. 60 = P. pubescens.
cinerascens, A. Gray, l. c. 61 = P. cinerascens.
odorata, A. Gray, Bot. U. St. Expl. Exped. i. 475 = P. juliflora?
pubescens, A. Gray, Pl. Wright. i. 60 = P. pubescens.
reptans, A. Gray, Bot. U. St. Expl. Exped. i. 475 = P. reptans.
strombulifera, A. Gray, l. c. = P. strombulifera.

STROMBOCARPUS, Benth. & Hook. f. Gen. i. 591 (1867), sphalm. = praec.

STROMBODURUS, Willd. ex Steud. Nom. ed. II. ii. 647 (1841) = **Pentarrhapis**, H. B. & K. (Inc. Sed.).
gracilis, Willd. ex Steud. l. c. = P. scabra.

STROMBOSIA, Blume, Bijdr. 1154 (1826). *OLACINEAE*, Benth. & Hook. f. i. 348.
LAVALLEA, Baill. Adansonia, ii. (1862) 361.
ceylanica, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 350.—Zeylan.
grandifolia, *Hook. f. ex Benth. in Hook. Niger Fl.* 258.—Afr. trop.
javanica, *Blume, Bijdr.* 1155.—Burma; Malaya.
javanica, Thw. Enum. Pl. Zeyl. 42 = ceylanica.
lucida, *Teijsm. & Binn. Cat. Hort. Bog.* 207, nomen.—Java.
philippinensis, *Vidal, Phan. Cuming. Philipp.* 23, 102.—Ins. Philipp.

STRONGYLOCALYX, Blume, Mus. Bot. Lugd. Bat. i. 89. t. 54 (1849) = **Eugenia**, Mich. (Myrtac.).
hemisphaericus, Blume, l. c. 90 = E. hemispherica.
lanceaefolius, Blume, l. c. 89, in obs. = E. Wallichii.
leptostachyus, Blume, ex Miq. Fl. Ind. Bat. i. 1 443 = E. leptostemon.
leptostemon, Blume, Mus. Bot. Lugd. Bat. i. 89 = E. leptostemon.
oblatus, Blume, l. c., in obs. = E. oblata.
praecox, Blume, l. c., in obs. = E. Wallichii.

STRONGYLODON, Vog. in Linnaea, x. (1836) 585. *LEGUMINOSAE*, Benth. & Hook. f. i. 532.
Lastellianus, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 381.—Madag.
lucidus, *Seem. Fl. Vit.* 61.—Ins. Fiji.
macrobotrys, *A. Gray, Bot. U. St. Expl. Exped.* i. 448. t. 49.—Ins. Philipp.
madagascariensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 267.—Madag.
ruber, *Vog. in Linnaea*, x. (1836) 585.—Ins. Sandvic.

STRONGYLOMA, DC. Prod. vii. 52 (1838) = **Nassauvia**, Comm. (Compos.).
axillare, DC. l. c. = N. axillaris.
glomerulosum, DC. l. c. = N. glomerulosa.

STRONGYLOSPERMA, Less. Syn. Comp. 261 (1832) = **Cotula**, Tourn. (Compos.).
australe, Less. l. c. = C. australis.
reptans, Benth. in Enum. Pl. Hueg. 60 = C. reptans.

STROPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4; ed. II. 66 (1827) = **Chloranthus**, Sw.

STROPHADES, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 82 = **Erysimum**, Tourn. (Crucif.).
lanceolata, Boiss. l. c. 83 = E. Strophades.
linearis, Boiss. l. c. = E. oleaefolium.

STROPHANTHUS, DC. in Bull. Soc. Philom. iii. (1802) 122. t. 8. *APOCYNACEAE*, Benth. & Hook. f. ii. 714.
CERCOCOMA, Wall. Cat. n. 1623 (1831?).
CHRISTYA, Harv. in Hook. Journ. Bot. iv. (1842) 134. t. 21.
alterniflorus, *Spreng. Syst.* i. 638.—China.
brevicaudatus, *Wight, Ic.* t. 1302.—Burma; Malaya.
Bullenianus, *Mast. in Gard. Chron.* (1870) 1471. f. 257.—Afr. trop.
capensis, *A. DC. in DC. Prod.* viii. 419.—Afr. austr.
caudatus, Kurz, For. Fl. Brit. Burma, ii. 192 = Wallichii?

STROPHANTHUS:—
caudatus, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 257 = dichotomus.
chinensis, G. Don, Gen. Syst. iv. 85 = divergens.
Cumingii, *A. DC. in DC. Prod.* viii. 418.—Ins. Philipp.
dichotomus, *DC. in Bull. Soc. Philom.* iii. (1802) 123. —Burma; Malaya.
dichotomus, Wall. Cat. n. 1641 = Wallichii.
divaricatus, Hook. & Arn. Bot. Beech. Voy. 199 = divergens.
divaricatus, *Wall. Cat.* n. 1642.—Burma.
divergens, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.-Jun. 1827) 177.—China.
Griffithii, Wight, Ic. t. 1300 = dichotomus.
hirtus, Poir. in Dict. Sc. Nat. li. 150 = seq.
hispidns, *DC. in Bull. Soc. Philom.* iii. (1802) 123.— Afr. austr.
Horsfieldianus, Miq. Fl. Ind. Bat. ii. 442 = dichotomus.
Jackianus, *Wall. Cat.* n. 1643.—Ins. Penang.
Kombe, *Oliver, in Hook. Ic. Pl.* t. 1098.—Afr. trop.
laurifolius, *DC. in Bull. Soc. Philom.* iii. (1802) 123.— Afr. trop.
longicaudatus, Wight, Ic. t. 1299 = dichotomus.
pendulus, Hook. in W. Gray, Trav. West. Afr. 392. t. C = sarmentosus.
pentaphyllus, Griff. Notul. iv. 78 = dichotomus.
Petersianus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 276.—Afr. trop.
sarmentosus, *DC. in Bull. Soc. Philom.* iii. (1802) 123. t. 8. f. 1.—Afr. trop.
scandens, *Griff. Notul.* iv. 743 (an seq. ?).—Malacca.
scandens, Reinw. & Schult. Syst. iv. 412 = dichotomus.
Senegambiae, A. DC. in DC. Prod. viii. 418 = sarmentosus.
Stanleyanus, Hort. ex Hook. Bot. Mag. sub t. 4466 = Roupellia grata.
terminalis, Blume, Cat. Gew. Buitenz. 56 = dichotomus.
Wallichii, *A. DC. in DC. Prod.* viii. 418.—Ind. or.
Wightianus, *Wall. Cat.* n. 1659.—Ind. or.

STROPHIOSTOMA, Turcz. in Bull. Soc. Nat. Mosc. (1840) 258 = **Myosotis**, Tourn. (Borag.).
amoena, Rupr. Ind. Sem. Hort. Petrop. (1852); et in Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 229 = M. amoena.
propinquum, Turcz. in Bull. Soc. Nat. Mosc. (1840) 259 = M. propinqua.
sparsiflorum, Turcz. l. c. 258 = M. sparsiflora.

STROPHIS, Salisb. Gen. Pl. Fragm. 12 (1866) = **Dioscorea**, Plum.
cirrhosa, Salisb. l. c. 12 = D. cirrhosa.

STROPHIUM, Dulac, Fl. Hautes-Pyr. 247 (1867) = **Arenaria**, Rupp. (Caryoph.).
filiforme, Dulac, l. c. 248 = A. muscosa.
trinervium, Dulac, l. c. = A. trinervia.

STROPHOLIRION, Torr. in Pacif. Rail. Rep. iv. 149 (1856). *LILIACEAE*, Benth. & Hook. f. iii. 799.
MACROSCAPA, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 148.
RUPALLEYA, Morière, in Bull. Soc. Linn. Normand. viii. (1864) 313. t. 15.
californicum, *Torr. in Pacif. Rail. Rep.* iv. 149. t. 21. —Calif.

STROPHOPAPPUS, DC. Prod. v. 75 (1836) = **Stilpnopappus**, Mart. (Compos.).
bicolor, DC. l. c. = Stilpnopappus bicolor.
emarginatus, Baker, in Mart. Fl. Bras. vi. II. 141, sphalm. = Stilpnopappus emarginatus.
Riedelianus, Baker, l. c. 140, sphalm. = Stilpnopappus bicolor.

STROPHOSTYLES, Ell. Sketch, ii. 229 (1822) = **Phaseolus**, Tourn. (Legumin.).
angulosa, Ell. l. c. = P. diversifolius.
helvola, Ell. l. c. 230 = P. helvolus.
pedunculares, Ell. l. c. = P. helvolus.

STROPHOSTYLES, E. Mey. Comm. Pl. Afr. Austr. 147 (1835) = **Vigna**, Savi (Legumin.).
capensis, E. Mey. l. c. = V. capensis.

STRUCHIUM, P. Br. Hist. Jamaic. 312. t. 34 (1756) = **Sparganophorus**, Vaill. (Compos.).
africanum, Beauv. Fl. Owar. i. 81. t. 48 = Sparg. Vaillantii.
americanum, Poir. Encyc. vii. 475 = Sparg. Vaillantii.

STRUKERIA, Vell. Fl. Flum. 8; i. t. 20 (1825) = **Vochisia**, Juss.
oppugnata, Vell. ll. cc. = V. oppugnata.

STRUMARIA, Jacq. Ic. Pl. Rar. ii. 13. tt. 356–360 (1790). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 728.
CHOERADODIA, Herb. Amaryll. 87 (1837).
EUDOLON, Salisb. Gen. Pl. Fragm. 127 (1866).
HYMENETRON, Salisb. l. c. 128 (1866).
NESYNSTYLIS, Rafin. Fl. Tellur. iv. 123 (1836).
PUGIONELLA, Salisb. Gen. Pl. Fragm. 128 (1866).
STYLAGO, Salisb. l. c. 127 (1866).
angustifolia, *Jacq. Ic. Pl. Rar.* ii. 13. t. 359; *Coll. Suppl.* 48.—Afr. austr.
Baueriana, Herb. Amaryll. 394. t. 45. f. 5 = truncata.
chilensis, *Molina, Sagg. Chile*, ed. II. 130, 284.— Chili.
crispa, Ker-Gawl. in Bot. Mag. t. 1363 = Hessea crispa.
filifolia, Jacq. Ic. Pl. Rar. ii. 14. t. 361 = Hessea filifolia.
gemmata, Ker-Gawl. in Bot. Mag. t. 1620 = Hessea geminata.
liliifolia, D. Dietr. Syn. Pl. ii. 1187 = Hessea filifolia.
linguaefolia, Jacq. Ic. Pl. Rar. ii. 13. t. 356 = truncata.
rubella, *Jacq. l. c.* t. 358.—Afr. trop.
spiralis, Ait. Hort. Kew. ed. II. ii. 213 = Carpolyza spiralis.
stellaris, Ker-Gawl. in Bot. Mag. sub t. 1363 = Hessea stellaris.
strumosa, Salisb. Gen. Pl. Fragm. 127 = Hessea filifolia.
truncata, *Jacq. Ic. Pl. Rar.* ii. 13. t. 357.—Afr. trop.
undulata, Berg. ex Kunth, Enum. Pl. v. 627 = Hessea gemmata.
undulata, *Jacq. Ic. Pl. Rar.* ii. 14. t. 360.—Afr. austr.

STRUMARIUM, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 89 = **Xanthium**, Linn. (Compos.).

STRUMPFIA, Jacq. Enum. Pl. Carib. 8 (1760); Select. Am. 218 (1763). *RUBIACEAE*, Benth. & Hook. f. ii. 117.
PATSJOTTI, Adans. Fam. ii. 84 (1763).
dubia, [Spreng.] Bot. Gart. Halle, 93, nomen = seq.?
maritima, *Jacq. Enum. Pl. Carib.* 28; *Select. Am.* 218. —Ind. occ.

STRUTHANTHUS, Mart. in Flora, xiii. (1830) 102 = **Loranthus**, Linn.
acuminatus, Blume, in Schult. f. Syst. vii. 1731 = L. acuminatus.
acutifolius, Blume, l. c. = L. acutifolius.
aduncus, Blume, l. c. = L. aduncus.
affinis, Mart. in Flora, xiii. (1830) 105 = L. affinis.
alveolatus, G. Don, Gen. Syst. iii. 410 = L. alveolatus.
amplexicaulis, G. Don, l. c. = L. amplexicaulis.
anceps, G. Don, l. c. 411 = L. anceps.
andrastylus, Eichl. in Mart. Fl. Bras. v. II. 74.— Bras.
attenuatus, Eichl. l. c. 72 = L. attenuatus.
avicularius, Mart. in Flora, xiii. (1830) 105 = L. avicularis.
buxifolius, Blume, in Schult. f. Syst. vii. 1731 = L. buxifolius.
calobotrys, Eichl. in Mart. Fl. Bras. v. II. 79.— Bras.
chordocladus, Eichl. l. c. 70, in obs. = L. chordocladus.
citricola, Mart. in Flora, xiii. (1830) 105 = L. citricola.
complexus, Eichl. in Mart. Fl. Bras. v. II. 73.—Bras.
concinnus, Mart. in Flora, xiii. (1830) 105 = L. concinnus.
conduplicatus, Mart. l. c. = L. conduplicatus.

STRUTHANTHUS:—
confertus, Mart. in Flora, xiii. (1830) 104 = Loranthus confertus.
crassipes, Eichl. in Mart. Fl. Bras. v. II. 70, in obs. = L. crassipes.
cuspidatus, Mart. in Flora, xiii. (1830) 105 = L. cuspidatus.
Deppeanus, Blume, in Schult. f. Syst. vii. 1731 = L. Deppeanus.
Destructor, Blume, l. c. = L. Destructor.
dichotrianthus, Eichl. in Mart. Fl. Bras. v. II. 75.— Am. trop.
domingensis, G. Don, Gen. Syst. iii. 414 = L. domingensis.
elegans, Mart. in Flora, xiii. (1830) 105 = L. elegans, *Mart.*
ellipticus, Blume, in Schult. f. Syst. vii. 1731 = L. ellipticus.
emarginatus, G. Don, Gen. Syst. iii. 410 = L. emarginatus.
erythrocarpus, Mart. in Flora, xiii. (1830) 105 = L. erythrocarpus.
Escholzianus, Mart. l. c. 104 = L. Eschscholzianus.
Eschscholzianus, Schult. f. Syst. vii. 1613 = praec.
eugenioides, Blume, in Schult. f. l. c. 1731 = L. eugenioides.
flexicaulis, Mart. in Flora, xiii. (1830) 105 = L. flexicaulis.
floribundus, Blume, in Schult. f. Syst. vii. 1731 (Quid ?). —Hab. ?
glomeratus, Mart. in Flora, xiii. (1830) 104 = L. glomeratus.
glomeriflorus, Eichl. in Mart. Fl. Bras. v. II. 84.— Bras.
guadalupensis, G. Don, Gen. Syst. iii. 414 = L. guadalupensis.
guianensis, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 978 = L. syringaefolius.
heterophyllus, Blume, in Schult. f. Syst. vii. 1731 = L. heterophyllus.
interruptus, G. Don, Gen. Syst. iii. 413 = L. interruptus.
lancifolius, Blume, in Schult. f. Syst. vii. 1731 = L. lancifolius.
laurifolius, Blume, l. c. = L. laurifolius.
laxiflorus, G. Don, Gen. Syst. iii. 414 = L. laxiflorus.
leptostachyus, G. Don, l. c. 411 = L. leptostachyus.
ligustrifolius, Blume, in Schult. f. Syst. vii. 1731 = L. ligustrifolius.
longepedunculatus, Mart. in Flora, xiii. (1830) 105 = L. longipedunculatus.
Magdalenae, G. Don, Gen. Syst. ii. 414 = L. Magdalenae.
marginatus, Blume, in Schult. f. Syst. vii. 1731 = L. marginatus.
microphyllus, G. Don, Gen. Syst. iii. 413 = L. microphyllus.
nigricans, Eichl. in Mart. Fl. Bras. v. II. 76.— Bras.
nitens, Mart. in Flora, xiii. (1830) I. 105 = L. nitens.
nitidus, Blume, in Schult. f. Syst. vii. 1731 = L. nitidus.
oblongus, G. Don, Gen. Syst. ii. 410 = Phoradendron rubrum.
occidentalis, G. Don, l. c. = L. occidentalis.
orbicularis, Blume, in Schult. f. Syst. vii. 1731 = L. orbicularis.
orinocensis, Blume, l. c. = L. orinocensis.
ovalifolius, G. Don, Gen. Syst. iii. 411 = L. ovalifolius.
paniculatus, Blume, in Schult. f. Syst. vii. 1731 = L. paniculatus.
parvifolius, G. Don, Gen. Syst. iii. 413 = L. parvifolius.
patens, Mart. in Flora, xiii. (1830) 104 = L. patens.
Patrisii, G. Don, Gen. Syst. iii. 410 = L. Patrisii.
pauciflorus, G. Don, l. c. 414 = L. pauciflorus, *Sw.*
pauciflorus, Miq. ex Eichl. in Mart. Fl. Bras. v. II. 75 = *dichotrianthus*.
pedunculatus, G. Don, Gen. Syst. iii. 414 = L. pedunculatus.
Perrottetii, G. Don, l. c. 413 = L. Perrottetii.
phillyraeoides, Blume, in Schult. f. Syst. vii. 1731 = L. phillyreoides.
phylliraeoides, G. Don, Gen. Syst. iii. 413 = praec.
podopterus, G. Don, l. c. 410 = L. podopterus.
polyanthus, Mart. in Flora, xiii. (1830) 105 = L. polyanthos.
polyrrhizus, Mart. l. c. = L. polyrrhizos.

STRUTHANTHUS :—
polystachyus, Blume, in Schult. f. Syst. vii. 1731 = Loranthus polystachyus.
portoricensis, G. Don, Gen. Syst. iii. 414 = L. portoricensis.
psilobotrys, G. Don, l. c. = L. psilobotrys.
pterygopus, Mart. in Flora, xiii. (1830) 105 = L. pterygopus.
punctatus, Blume, in Schult. f. Syst. vii. 1731 = L. punctatus, *Ruiz & Pav.*
puracensis, Blume, l. c. = L. puracensis.
pyrifolius, Blume, l. c. = L. pyrifolius.
quercicola, Blume, l. c. = L. quercicola.
radicans, Blume, l. c. = L. radicans.
retroflexus, Blume, l. c. = L. retroflexus.
retusus, Blume, l. c. = L. Sellowianus.
rhynchophyllus, *Eichl. in Mart. Fl. Bras.* v. II. 85.—Bras.
rotundifolius, Kostel. Allg. Med.-Pharm. Flora, ii. 536 = L. rotundifolius.
rubens, Mart. in Flora, xiii. (1830) 105 = L. rubens.
rufus, Mart. l. c. = L. rufus.
salicifolius, Mart. l. c. 104 = L. salicifolius.
Salzmanni, Eichl. in Mart. Fl. Bras. v. II. 80.—Bras.
sarmentosus, Blume, in Schult. f. Syst. vii. 1731 = L. sarmentosus.
sessilis, Blume, l. c. = L. sessilis.
spicatus, G. Don, Gen. Syst. iii. 410 = L. spicatus.
squamulosus, Klotzsch, in. Schomb. Faun. Fl. Brit. Gui. 978 (= *Phthirusa squamulosa*).—Guian.
staphylinus, Mart. in Flora, xiii. (1830) 105 = L. staphylinus.
Stelis, Blume, in Schult. f. Syst. vii. 1731 = L. Stelis.
suaveolens, Blume, l. c. = L. suaveolens.
subcampestris, Mart. in Flora, xiii. (1830) 105 = L. subcampestris.
subrotundus, G. Don, Gen. Syst. iii. 413 = L. subrotundus.
syringaefolius, Mart. in Flora, xiii. (1830) 105 = L. syringaefolius.
Tagua, Blume, in Schult. f. Syst. vii. 1731 = L. Tagua.
taubatensis, Eichl. in Mart. Fl. Bras. v. II. 84.—Bras.
terniflorus, Eichl. l. c. 75 = L. terniflorus.
terniflorus, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 978 = *dichotrianthus*.
tetraqueter, Mart. in Flora, xiii. (1830) 105 = L. tetraqueter.
Theobromae, Blume, in Schult. f. Syst. vii. 1731 = L. Theobromae.
thyrsiflorus, Blume, l. c. = L. thyrsiflorus.
triceps, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1161 = L. syringaefolius.
uniflorus, G. Don, Gen. Syst. iii. 414 = L. uniflorus.
uraguensis, G. Don, l. c. 410 = L. uraguensis.
uruguayensis, Gibert, Enum. Pl. Montev. 83 = L. uraguensis.
venetus, Blume, in Schult. f. Syst. vii. 1731 = L. venetus.
virgatus, Mart. in Flora, xiii. (1830) 105 = L. virgatus.
vulgaris, Mart. l. c. xxiv. (1841) II. Beibl. 33.—Bras.

STRUTHIA, Royen, ex Linn. Gen. ed. II. 164 (1742) = **Gnidia**, Linn. (Thymel.).

STRUTHIOLA, Linn. Mant. i. 4 (1767). *THYMELAEACEAE*, Benth. & Hook. f. iii. 196.
BELVALA, Adans. Fam. ii. 285 (1763).
angustifolia, *Lam. Illustr.* i. 314 (err. typ. 214).—Afr. austr.
angustifolia, Steud. ex Meissn. in Linnaea, xiv. (1840) 469 = linearioloba.
argentea, *Lehm. Del. Sem. Hort. Hamb.* (1831) 6, 7.—Afr. austr.
aurea, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 571 = argentea.
brevifolia, Willd. ex Meissn. in Linnaea, xiv. (1840) 477 = striata.
chrysantha, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 572 = tomentosa.
chrysantha, *Licht. ex Roem. & Schult. Syst.* iii. 333.—Afr. austr.
ciliata, Andr. Bot. Rep. t. 149 = virgata.
ciliata, Lam. Illustr. i. 314 (err. typ. 214) = lucens.
dodecapetala, *Bartl. ex Meissn. in Linnaea*, xiv. (1840) 475.—Afr. austr.
Dregeana, Meissn. l. c. 472 = argentea.
Eckloniana, *Meissn. in DC. Prod.* xiv. 568.—Afr. austr.

STRUTHIOLA :—
erecta, Curt. Bot. Mag. t. 222 = linearioloba ?
erecta, *Linn. Mant.* i. 41.—Afr. austr.
formosa, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 571 = argentea.
glabra, Roem. & Schult. Syst. iii. 331 = erecta.
glabra, Zeyh. ex Meissn. in Linnaea, xiv. (1840) 468 = parviflora.
glauca, Lodd. Cat. (1807) 33 = virgata ?
glauca, Nois. ex Meissn. in DC. Prod. xiv. 570 = longiflora.
glauca, Sieber, ex Presl, Bot. Bemerk. 108 = virgata.
grandis, Bartl. ex Meissn. in Linnaea, xiv. (1840) 476 = hirsuta.
hirsuta, *Wikstr. in Vet. Akad. Handl. Stockh.* (1818) 290.—Afr. austr.
imbricata, Andr. Bot. Rep. t. 113 = striata.
imbricata, Hort. ex Meissn. in DC. Prod. xiv. 559 = Arthrosolen spicatus.
incana, Lodd. Bot. Cab. t. 11 = virgata.
juniperina, Retz. Obs. iii. 26 = linearioloba.
lanceolata, Retz. l. c. = ovata.
lateriflora, Hornem. Hort. Hafn. ii. 955 = striata.
lateriflora, Meissn. in DC. Prod. xiv. 573 = Arthrosolen spicatus.
laxa, Nois. ex Schult. Mant. iii. 260, nomen (Quid ?).—Hab. ?
linearioloba, *Meissn. in Linnaea*, xiv. (1840) 468.—Afr. austr.
littoralis, Wendl. ex Endl. Gen. Suppl. iv. II. 63 = Arthrosolen spicatus.
longiflora, *Lam. Illustr.* i. 314 (err. typ. 214) t. 78.—Afr. austr.
lucens, *Poir. Encyc.* vii. 477.—Afr. austr.
lutescens, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 568 = parviflora.
Martiana, *Meissn. l. c.* 570.—Afr. austr.
Mundtii, *Eckl. ex Meissn. l. c.* 572.—Afr. austr.
Myrsinites, Lam. Illustr. i. 314 (err. typ. 214) = ovata.
nana, Linn. f. Suppl. 128 = Gnidia nana.
oppositifolia, Cat. Pl. Bot. Gard. Liverpool (1808), 176, nomen (Quid ?).—Hab. ?
ovata, *Thunb. Prod. Pl. Cap.* 76.—Afr. austr.
parviflora, *Bartl. ex Meissn. in Linnaea*, xiv. (1840) 467.—Afr. austr.
pendula, Salisb. Prod. 282 = erecta.
pubescens, Retz. Obs. iii. 26 = virgata.
pubescens, Sims, Bot. Mag. t. 1212 = longiflora.
pubescens, Steud. ex Meissn. in Linnaea, xiv. (1840) 478 = lucens.
rigida, *Meissn. in DC. Prod.* xiv. 570.—Afr. austr.
rubra, Donn, Hort. Cantab. ed. IV. 31 = longiflora.
rubra, Steud. Nom. ed. II. ii. 648 = virgata.
scandens, Wall. ex Mast. in Hook. f. Fl. Brit. Ind. i. 582 = Cansjera Rheedii.
striata, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 572 = Mundtii.
striata, *Lam. Illustr.* i. 314 (err. typ. 214).—Afr. austr.
stricta, Donn, Hort. Cantab. ed. VI. 40 = linearioloba ?
subulata, Lam. Illustr. i. 314 (err. typ. 214) = erecta.
subulata, Vahl, ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 291 = tuberculosa ?
sulcata, F. W. Schmidt, Neue u. Selt. Pfl. 34 = striata.
superba, *Cat. Hort. Bot. Gard. Liverpool* (1808), 176, nomen (Quid).—Hab. ?
tetragona, Retz. Obs. iii. 25 = erecta.
Thomsoni, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 404.—Afr. trop.
tomentosa, *Andr. Bot. Rep.* t. 334.—Afr. austr.
tuberculosa, *Lam. Illustr.* i. 314 (err. typ. 214).—Afr. austr.
tuberculosa, Vahl, ex Wikstr. in Vet. Akad. Handl. Stockh. (1818) 287 = virgata.
Vahlii, Wikstr. l. c. 291 = tuberculosa ?
villosa, Vahl, ex Wikstr. l. c. 287 = virgata.
villosa, Wikstr. l. c. 288 = tomentosa.
virgata, Houtt. Handleid. vii. 364. t. 40. f. 2 = lucens.
virgata, *Linn. Mant.* i. 41; *Burm. f. Fl. Cap. Prod.* 12.—Afr. austr.
virgata, Wendl. Bot. Beobacht. 9. t. 2. f. 9 = striata.

STRUTIOLA, Burm. f. Fl. Cap. Prod. 12 (1768) = praec.

STRYCHNODAPHNE, Nees & Mart. in Linnaea, viii. (1833) 39 = **Ocotea**, Aubl. (Laurin.).
floribunda, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 188 = O. botryophylla.

STRYCHNODAPHNE :—
lanceolata, Nees, in Linnaea, viii. (1833) 39 = Ocotea lanceolata.
Lhotzkyi, Meissn. in DC. Prod. xv. I. 143 = O. Lhotzkii.
puberula, Nees & Mart. in Linnaea, viii. (1833) 39 = O. puberula.
salicifolia, Meissn. in Kjoeb. Vidensk. Meddel. (1870) 139 (= *Oreodaphne microbotrys*).—Bras.
suaveolens, Griseb. in Goett. Abh. xxiv. (1879) 134 = O. puberula.

STRYCHNOPSIS, Baill. in Bull. Soc. Linn. Par. i. (1885) 456. *MENISPERMACEAE*.
Thouarsii, *Baill. l. c.*—Madag.

STRYCHNOS, Linn. Syst. ed. I (1735). *LOGANIACEAE*, Benth. & Hook. f. ii. 797.
ATHERSTONEA, Pappe, Sylva Cap. ed. II. 29 (1862).
BREHMIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 25.
CHEMNICIA, Scop. Introd. 139 (1777).
CURARE, Humb. Relat. Hist. ii. 549 (1819).
IGNATIA, Linn. f. Suppl. 20 (1781).
IGNATIANA, Lour. Fl. Cochinch. 126 (1790).
LASIOSTOMA, Schreb. Gen. 75 (1789).
NARDA, Vell. Fl. Flum. 108 (1825); iii. t. 24 (1827).
ROUHAMON, Aubl. Pl. Gui. i. 93. t. 36 (1775).
TOXICARIA, Schreb. in Naturf. xix. (1783) 146.
UNGUACHA, Hochst. in Flora, xxvii. (1844) 103.
abyssinica, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 25, nomen.—Abyss.
acuminata, *Wall. Cat.* n. 1593, partim.—Burma.
acuminata, Wall. l. c., partim = rufa.
acuta, *Prog. in Mart. Fl. Bras.* vi. I. 280.—Bras.
albiflora, *Prog. l. c.* 279.—Bras.
angustifolia, *Benth. in Journ. Linn. Soc.* i. (1856) 102.—China.
Atherstonei, *Harv. Thes. Cap.* ii. 41.—Afr. austr.
axillaris, Blume, Bijdr. 1019 = monosperma.
axillaris, *Colebr. in Trans. Linn. Soc.* xii. (1819) 356. t. 15.—Reg. Himal.
axillaris, Dalz. & Gibs. Bomb. Fl. 155 = Dalzellii.
Baculum, Harv. Thes. Cap. ii. 41 = Atherstonei.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 88.—Ind. or.
Benthami, *C. B. Clarke, l. c.* 87.—Zeylan.
bicirrhosa, Lesch. ex Roxb. Fl. Ind. ed. Carey, ii. 267 = colubrina.
bicolor, *Prog. in Kjoeb. Vidensk. Meddel.* (1869) 31.—Bras.
brachiata, *Ruiz & Pav. Fl. Per.* ii. 30. t. 157.—Peruv.
brasiliensis, Benth. in Journ. Linn. Soc. i. (1856) 107, partim = Martii.
brasiliensis, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 84.—Bras.
breviflora, *A. DC. in DC. Prod.* ix. 15.—Bras.
brevifolia, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1857) 106.—Bras.
Castelnaeana, Baill. Adansonia, xii. (1879) 374 = seq.
Castelnaei, *Wedd. in Castel. Exp. Am. Sud.* v. 22; *ex Benth. in Journ. Linn. Soc.* i. (1856) 108.—Bras.
chlorantha, *Prog. in Mart. Fl. Bras.* vi. I. 273.—Costa Rica.
cinnamomea, Thw. Enum. Pl. Zeyl. 425, sphalm. = seq.
cinnamomifolia, *Thw. l. c.* 201.—Zeylan.
cirrhosa, Stokes, Bot. Mat. Med. i. 414 = colubrina.
cogens, *Benth. in Hook. Journ. Bot.* iii. (1841) 241.—Guian.
colubrina, Auct. ex DC. Prod. ix. 15 = ligustrina.
colubrina, Blume, Rumphia, i. 70 = Rheedei.
colubrina, *Linn. Sp. Pl.* 189.—Ind. or.
colubrina, Stokes, Bot. Mat. Med. i. 414 (Quid ?).—Hab. ?
colubrina, Wall. Cat. nn. 4455, 1589 = Beddomei, cinnamomifolia.
coriacea, Thw. Enum. Pl. Zel. 425 = Beddomei.
Crevauxiana, *Baill. Adansonia*, xii. (1879) 377.—Guian.
Crevauxii, G. Planch. in Journ. Pharm. et Chim. (1880), reimpr. 25 = praec.
Curare, *Benth. in Journ. Linn. Soc.* i. (1856) 108.—Am. trop.
Dalzellii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 87.—Ind. or.
darienensis, *Seem. in Bot. Voy. Herald*, 166.—Panama.

STRYCHNOS:—
densiflora, *Baill. Adansonia*, xii. (1879) 369.—Afr. austr.
depauperata, Baill. l. c. 372.—Am. trop.
dysophylla, *Benth. in Journ. Linn. Soc.* i. (1856) 103.—Afr. austr.
Erichsonii, *Schomb. Faun. Fl. Brit. Gui.* 1082, nomen; *et ex Prog. in Mart. Fl. Bras.* vi. i. 274.—Guian.
farinosa, *Blume, Rumphia*, i. 70.—Madag.
Gardneri, *A. DC. in DC. Prod.* ix. 14.—Bras.
Gaultheriana, Pierre, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 89 = malaccensis.
Geoffraeana, *Baill. Adansonia*, xii. (1879) 378.—Bras.?
glabra, *Sagot, ex Prog. in Mart. Fl. Bras.* vi. i. 275.—Guian.
Gomesiana, Casar. Atti 8ª Riun. Scient. Ital. (1841) 515 = triplinervia.
grandis, Wall. Cat. n. 4454 = Anisophyllea grandis.
Grayi, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 519.—Cuba.
Gubleri, *G. Planch. in Journ. Pharm. et Chim.* v. (1882) reimpr. 1.—Venezuela.
guineensis, Thonn. ex Didr. in Kjoeb. Vidensk. Meddel. (1854) 190 = Landolphia scandens.
hachensis, *Karst. Fl. Columb.* ii. 75. t. 138.—N. Granat.
hirsuta, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1856) 106.—Bras.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 379.—Java.
hypogyna, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 86.—Burma.
Icaja, *Baill. Adansonia*, xii. (1879) 368.—Afr. trop.
Ignatii, *Berg. Mat. Med.* i. 146.—Ins. Philipp.
innocua, *Delile, Cent. Pl. Méroë*, 58.—Nubia.
Jobertiana, *Baill. Adansonia*, xii. (1879) 367.—Bras.
lanceolaris, *Miq. Fl. Ind. Bat. Suppl.* 551.—Sumatra.
lanceolata, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1856) 105.—Bras.
laurina, Thw. Enum. Pl. Zeyl. 201 = Beddomei.
laurina, *Wall. Cat.* n. 1591.—Ind. or.; Malaya.
ligustrina, Blume, Rumphia, i. 68. t. 25 = S. Nux-vomica.
ligustrina, Zipp. ex Span. in Linnaea, xv. (1841) 325 = colubrina.
Lokua, *A. Rich. Tent. Fl. Abyss.* ii. 52.—Abyss.
lucida, *R. Br. Prod.* 469.—Austral.
lucida, Wall. Cat. n. 1590 = S. Nux-vomica, Wallichiana.
macroacanthos, *Prog. in Mart. Fl. Bras.* vi. i. 280. t. 82.—Bras.
madagascariensis, *Poir. Encyc.* viii. 696.—Madag.
madagascariensis, Spreng. ex Baker, Fl. Maurit. 235 = spinosa.
Maingayi, *C. B. Clarke, in Hook f. Fl. Brit. Ind.* iv. 88.—Malacca.
malaccensis, *Benth. in Journ. Linn. Soc.* i. (1856) 101.—Burma; Malaya.
marginata, *Benth. l. c.* 107.—Bras.
Martii, *Prog. in Mart. Fl. Bras.* vi. i. 279.—Bras.
Medeola, *Sagot, ex Prog. l. c.* 282.—Am. trop.
Melinoniana, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 256.—Guian.
micrantha, *Thw. Enum. Pl. Zeyl.* 425.—Zeylan.
minor, Benth. in Journ. Linn. Soc. i. (1856) 100 = Benthami.
minor, Blume, Rumphia, i. 70 = colubrina.
minor, *Dennst. Schluess. Hort. Malab.* 33.—Ind. or.
Mittscherlichii, Schomb. Faun. Fl. Brit. Gui. 950 = smilacina.
monosperma, *Miq. Fl. Ind. Bat.* ii. 381.—Malaya.
monosperma, Stokes, Bot. Mat. Med. i. 415 = potatorum.
multiflora, *Benth. in Journ. Linn. Soc.* i. (1856) 102.—Ins. Philipp.
muricata, Kostel. Allg. Med.-Pharm. Flora, iii. 1072 = ligustrina.
nigricans, *Prog. in Mart. Fl. Bras.* vi. i. 280. t. 79.—Bras.
nitida, G. Don, Gen. Syst. iv. 66 = S. Nux-vomica, Wallichiana.
Nux-vomica, *Linn. Sp. Pl.* 189.—Ind. or.; Burma.
oblongifolia, Hochst. in Flora, xxvii. (1844) 827 = Carissa oblongifolia.
ovalifolia, Stokes, Bot. Mat. Med. i. 412 = S. Nux-vomica.
ovalifolia, Wall. Cat. n. 1592 = laurina.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 551.—Sumatra.
panamensis, *Seem. Bot. Voy. Herald*, 166.—Panama.

STRYCHNOS:—
paniculata, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 56.—China.
parviflora, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1856) 107.—Bras.
parvifolia, *A. DC. in DC. Prod.* ix. 16.—Bras.
pedunculata, *Benth. in Journ. Linn. Soc.* i. (1856) 105.—Guian.
philippensis, Blanco, Fl. Filip. ed. II. 61 = Ignatii.
Poeppigii, *Prog. in Mart. Fl. Bras.* vi. i. 282.—Peruv.
potatorum, *Linn. f. Suppl.* 148.—Ind. or.; Burma.
Pseudo-china, Benth. in Journ. Linn. Soc. i. (1856) 104 = seq.
Pseudo-quina, *A. St. Hil. in Mém. Mus. Par.* ix. (1822) 340.—Bras.
psilosperma, *F. Muell. Fragm.* iv. 44.—Austral.
pubescens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 89.—Malacca.
randiaeformis, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 246.—Afr. austr.
Rheedei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 87.—Ind. or.
rhexioides, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1144, nomen.—Guian.
rondeletioides, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1856) 104.—Bras.
Rouhamon, *Benth. l. c.* 106.—Am. trop.
rubiginosa, *A. DC. in DC. Prod.* ix. 16.—Bras.
rufa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 89.—Burma; Malacca.
scandens, Schum. & Thonn. Beskr. Guin. Pl. 127 = Landolphia scandens.
Schomburgkiana, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1144 = pedunculata.
septemnervis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 88.—Malacca.
smilacina, *Benth. in Journ. Linn. Soc.* i. (1856) 105.—Am. trop.
spinosa, *Lam. Illustr.* ii. 38.—Madag.
subcordata, *Spruce, ex Benth. in Journ. Linn. Soc.* i. (1856) 106.—Bras.
Telankotta, J. F. Gmel. Syst. 387, sphalm. = seq.
Tetankotta, Retz. Obs. ii. 12 = potatorum.
Ticute, Steud. Nom. ed. II. ii. 648, err. typ. = seq.
Tieute, *Lesch. in Ann. Mus. Par.* xvi. (1810) 479, 480. t. 23.—Java.
tomentosa, *Benth. in Journ. Linn. Soc.* i. (1856) 104.—Guian.
toxifera, *Schomb. ex Benth. in Hook. Journ. Bot.* iii. (1841) 240; *et in Hook. Ic. Pl.* tt. 364–365.—Guian.
trinitensis, *Griseb. Fl. Brit. W. Ind.* 407.—Ind. occ.
triplinervia, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 83.—Bras.
Unguacha, *A. Rich. Tent. Fl. Abyss.* ii. 52.—Abyss.
Vacacoua, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 242.—Madag.
vestita, *Prog. in Kjoeb. Vidensk. Meddel.* (1869) 32.—Bras.
vomica, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = S. Nux-vomica.
Vuntac, Boj. Hort. Maurit. 205 = spinosa.
Wallichiana, *Benth. in Journ. Linn. Soc.* i. (1856) 102.—Reg. Himal.
Wallichiana, Steud. Nom. ed. II. ii. 648 = S. Nux-vomica, Wallichiana.
yapurensis, *G. Planch. in Journ. Pharm. et Chim.* v. (1882), reimpr. 12.—Ecuador.

STRYCHNUS, Linn. Gen. ed. I. 54 (1737) = praec.

STRYPHNODENDRON, Mart. in Flora, xx. (1837) ii. Beibl. 117. *LEGUMINOSAE*, Benth. & Hook. f. i. 590.
FOLIANTHERA, Rafin. Sylva Tellur. 120 (1838).
angustum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 375; *et in Mart. Fl. Bras.* xv. ii. 287.—Bras.
Barbatimam, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 117.—Bras.
coriaceum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 373; *et in Mart. Fl. Bras.* xv. ii. 284.—Bras.
discolor, Benth. in Hook. Journ. Bot. iv. (1842) 342 = rotundifolium.
floribundum, *Benth. l. c.* 343.—Bras.
guianense, *Benth. in Trans. Linn. Soc.* xxx. (1875) 374.—Guian.
inaequale, Benth. l. c. = microstachyum.

STRYPHNODENDRON:—
Melinonis, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 322.—Guian.
microstachyum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 82.—Bras.
obovatum, *Benth. in Trans. Linn. Soc.* xxx. (1875) 374; *et in Mart. Fl. Bras.* xv. ii. 286.—Bras.
paniculatum, Poepp. & Endl. Nov. Gen. et Sp. iii. 81. t. 291 = Piptadenia Poeppigii.
polyphyllum, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 117.—Bras.
rotundifolium, *Mart. l. c.*—Bras.

STRZELECKYA, F. Muell. in Hook. Kew Journ. ix. (1857) 308 = **Flindersia**, R. Br. (Meliac.).
dissosperma, F. Muell. l. c. = F maculosa.

STUARTIA, Linn. in Act. Soc. Upsal. (1741) 79 (Stewartia). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 185.
MALACHODENDRON, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. (1748) App. 216; Cav. Diss. v. 302. t. 158 (1788).
MALACHODENDRUM, Pers. Syn. ii. 260 (1807).
corchoroides, Forsk. Fl. Aegypt. Arab. 126 = Sida spinosa.
dentata, Nois. ex Steud. Nom. ed. I. 815, nomen (Quid?).—Hab.?
grandiflora, *Carr. in Rev. Hortic.* (1874) 399.—Japon.
grandiflora, Siebold, ex Briot, in Rev. Hortic. (1879) 430 = S. Pseudo-camellia.
Malacodendron, *Linn. Sp. Pl.* 698.—Am. bor.
marilandica, Andr. Bot. Rep. t. 397 = S. Malacodendron.
monadelpha, Maxim. in Bull. Acad. Pétersb. xi. (1867) 430 = serrata.
monadelpha, *Sieb. & Zucc. Fl. Jap.* i. 181. t. 96.—Japon.
montana, Bartr. Trav. ed. II. (1794) 332 = pentagyna.
nobilis, Salisb. Prod. 386 = S. Malacodendron.
pentagyna, *L'Hérit. Stirp. Nov.* 155. t. 74.—Am. bor.
Pseudo-camellia, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 429.—Japon.
serrata, *Maxim. l. c.* 430.—Japon.
virginica, Cav. Diss. v. 303. t. 159. f. 2 = S. Malacodendron.

STUARTINA, Sond. in Linnaea, xxv. (1852) 521. *COMPOSITAE*, Benth. & Hook. f. ii. 300.
Muelleri, *Sond. l. c.* 522.—Austral.

STUBENDORFFIA, Schrenk, ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 20 (1844). *CRUCIFERAE*, Benth. & Hook. f. i. 89.
orientalis, *Schrenk, l. c.* 21.—Soongar.

STUBENDORFIA, Walp. Rep. v. 50 (1846), sphalm. = praec.

STUPA, Aschers. Fl. Brandenb. 812 (1864) = **Stipa**, Linn. (Gramin.).

STURMIA, Gaertn. f. Fruct. iii. t. 192. f. 3 (1805) = **Antirrhoea**, Comm. (Rubiac.).
lucida, Gaertn. f. l. c. = A. lucida.

STURMIA, Hoppe, in Sturm, Deutschl. Fl. Heft. vii. (1799) = **Mibora**, Adans. (Gramin.).
minima, Hoppe, l. c. = M. verna.
verna, Pers. Syn. i. 76 = M. verna.

STURMIA, Reichb. in Moessl. Handb. ed. II. ii. 1576 (1828) = **Liparis**, Rich. (Orchid.).
abyssinica, Reichb. f. in Walp. Ann. iii. 527 = L. abyssinica.
angustilabris, F. Muell. Fragm. iv. 164 = L. reflexa.
arnoglossophylla, Reichb. f. in Bonplandia, iv. (1856) 217 = L. arnoglossophylla.
bituberculata, Reichb. f. l. c. ii. (1854) 22 = L. bituberculata, elata.
capensis, Sond. in Linnaea, xix. (1847) 71 = L. capensis.
coelogynoides, F. Muell. Fragm. ii. 71 = L. coelogynoides.
cuneilabris, Benth. Fl. Austral. vi. 273, sphalm. = L. reflexa.
elliptica, Reichb. f. in Linnaea, xxii. (1849) 833 = L. elliptica.
habenarina, F. Muell. Fragm. iv 131 = L. habenarina.

STURMIA :—
Kappleri, Reichb. f. in Linnaea, xxii. (1849) 833 = Liparis Kappleri.
liliifolia, Reichb. f. in Walp. Ann. iii. 527 = L. liliifolia.
Loeselii, Reichb. in Moesl. Handb. ed. II. ii. 1576 = L. Loeselii.
longipes, Reichb. f. in Bonplandia, iii. (1855) 250 = L. longipes.
nervosa, Reichb. f. l. c. = L. nervosa.
paludosa, Reichb. in Moessl. Handb. ed. II. ii. 1576 = Malaxis paludosa.
reflexa, F. Muell. Fragm. ii. 72 = L. reflexa.
Wageneri, Reichb. f. in Bonplandia, iii. (1855) 73 = L. Wageneri.

STURTIA, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 68 (1849) = **Gossypium**, Linn. (Malvac.).
gossypioides, R. Br. l. c. = G. Sturtii.

STUTZERIA, F. Muell. Fragm. v. 31 (1865) = **Callicoma**, Andr. (Saxifr.).
callicomoides, F. Muell. l. c. = C. Stutzeri.

STYGNANTHE, Hanst. in Linnaea, xxvi. (1853) 209 = **Columnea**, Plum. (Gesner.).
moësta, Hanst. l. c. = C. moesta.

STYLAGO, Salisb. Gen. Pl. Fragm. 127 (1866) = **Strumaria**, Jacq. (Liliac.).
rubella, Salisb. l. c. 128 = Strumaria rubella.

STYLANDRA, Nutt. Gen. Am. i. 170 (1818) = **Podostigma**, Ell. (Asclep.).
pumila, Nutt. l. c. = P. pubescens.

STYLANTHUS, Reichb. & Zoll. in Linnaea, xxviii. (1856) 312 = **Mallotus**, Lour. (Euphorb.).

STYLAPTERA, Benth. & Hook. f. Gen. iii. 202 (1880), sphalm. = seq.

STYLAPTERUS, A. Juss. in Ann. Sc. Nat. Sér, III. vi. (1846) 23. t. 1 = **Penaea**, Linn.
barbatus, A. Juss. l. c. = P. barbata.
ericoides, A. Juss. l. c. = P. ericoides.
fruticulosus, A. Juss. l. c. = P. fruticulosa.

STYLBOCARPA, Decne. & Planch. in Rev. Hortic. (1854) 105, sphalm. = **Stilbocarpa**, Decne. (Araliac.).

STYLEDIUM, Andr. Bot. Rep. t. 658 (1811?), sphalm. = **Stylidium**, Sw.

STYLESIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 377 = **Bahia**, Lag. (Compos.).
ambrosioides, Nutt. l. c. = B. ambrosioides.
puberula, Nutt. l. c. 378.—Chili.

STYLEURODON, Rafin. Fl. Tellur. ii. 14 (1836) = **Verbena**, Linn.
carolinianum, Rafin. l. c. = V. carolina.

STYLEXIA, Rafin. Fl. Tellur. iv. 121 (1836) = **Caylusea**, A. St. Hil. (Resed.).

STYLIDIUM, Lour. Fl. Cochinch. 220 (1790) = **Marlea**, Roxb. (Cornac.).
chinense, Lour. l. c. 221 = M. begonifolia.
vitiense, F. Muell. Census, 74 = M. vitiensis.

STYLIDIUM, Sw. in Ges. Naturf. Fr. Berl, Mag. i. (1807) 48. tt. 1, 2. *STYLIDIEAE*, Benth. & Hook. f. ii. [II.] 534.
ANDERSONIA, Koen. ex G. Don, Gen. Syst. iii. 721 (1834).
CANDOLLEA, Labill. in Ann. Mus. Par. vi. (1805) 453.
FORSTERIOPSIS, Sond. in Lehm. Pl. Preiss. i. 393 (1845).
VENTENATIA, Sm. Exot. Bot. ii. 13 (1806).
aciculare, Sond. in Lehm. Pl. Preiss. i. 373 = lineare.
adnatum, *R. Br. Prod.* 572.—Austral.
adpressum, *Benth. Fl. Austral.* iv. 22.—Austral.
affine, Sond. in Lehm. Pl. Preiss. i. 371 = reduplicatum.
alsinoides, *R. Br. Prod.* 572.—Austral.
amoenum, *R. Br. l. c.* 570.—Austral.

STYLIDIUM:—
androsaceum, DC. Prod. vii. 783 = guttatum.
androsaceum, Lindl. Swan River App. 29 = calcaratum.
Armeria, Labill. Nov. Holl. Pl. ii. 66. t. 216 = graminifolium.
armerium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = praec.
articulatum, *R. Br. Prod.* 570.—Austral.
assimile, *R. Br. l. c.* 569.—Austral.
Barleei, *F. Muell. Fragm.* vi. 5. t. 69.—Austral.
bellidifolium, Sond. in Lehm. Pl. Preiss. i. 376 = spathulatum.
bicolor, Lindl. Swan River App. 28 = piliferum.
brachyphyllum, *Sond. in Lehm. Pl. Preiss.* i. 386.—Austral.
breviscapum, *R. Br. Prod.* 572.—Austral.
Brunonianum, *Benth. in Enum. Pl. Hueg.* 72.—Austral.
Brunonis, Griff. Notul. iv. 275 = Kunthii.
bryoides, F. Muell. Fragm. vi. 91 = pedunculatum.
bulbiferum, *Benth. in Enum. Pl. Hueg.* 73.—Austral.
buxifolium, Hort. Belv. ex Steud. Nom. ed. I. 818, nomen (Quid?).—Austral.?
caespitosum, *R. Br. Prod.* 569.—Austral.
calcaratum, *R. Br. l. c.* 570.—Austral.
canaliculatum, *Lindl. Swan River App.* 29.—Austral.
canaliculatum, Poir. Encyc. Suppl. v. 412 = graminifolium.
capillare, *R. Br. Prod.* 570.—Austral.
caricifolium, Lindl. Swan River App. 28 = reduplicatum.
carnosum, *Benth. in Enum. Pl. Hueg.* 71.—Austral.
caulescens, Lindl. Swan River App. 29; DC. Prod. vii. 782 = amoenum.
cicatricosum, Sond. in Lehm. Pl. Preiss. i. 390 = fasciculatum.
ciliatum, *Lindl. Swan River App.* 28.—Austral.
compressum, Lindl. l. c. 29 = Brunonianum.
corymbosum, Benth. in Enum. Pl. Hueg. 72 = pulchellum.
corymbosum, *R. Br. Prod.* 571.—Austral.
crassifolium, *R. Br. l. c.*—Austral.
crossocephalum, *F. Muell. Fragm.* vi. 5.—Austral.
debile, *F. Muell. l. c.* i. 149.—Austral.
despectum, *R. Br. Prod.* 571.—Austral.
dichotomum, *DC. Prod.* vii. 783.—Austral.
Dicksoni, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 645 = pilosum.
diffusum, *R. Br. Prod.* 571.—Austral.
dispermum, *F. Muell. Fragm.* iv. 93.—Austral.
diuroides, *Lindl. Swan River App.* 29.—Austral.
divaricatum, Sond. in Lehm. Pl. Preiss. i. 385 = streptocarpum.
diversifolium, *R. Br. Prod.* 570.—Austral.
Drummondii, R. Grah. in Edinb. N. Phil. Journ. (1840–41) 208 = reduplicatum.
eglandulosum, *F. Muell. Fragm.* i. 150.—Austral.
elongatum, *Benth. Fl. Austral.* iv. 11.—Austral.
emarginatum, *Sond. in Lehm. Pl. Preiss.* i. 383.—Austral.
eriopodum, DC. Prod. vii. 784 = breviscapum.
eriorhizum, *R. Br. Prod.* 569.—Austral.
falcatum, *R. Br. l. c.* 572.—Austral.
fasciculatum, *R. Br l. c.*—Austral.
fasciculatum, Lindl. Bot. Reg. t. 1459 = adnatum.
fissilobum, *F. Muell. Fragm.* i. 154.—Austral.
Floodii, *F. Muell. l. c.* 149.—Austral.
floribundum, *R. Br. Prod.* 569.—Austral.
fruticosum, R. Br. l. c. 570 = glandulosum.
glandulosum, *Salisb. Parad. Lond.* t. 77.—Austral.
glaucum, R. Br. Prod. 569 = rupestre.
glaucum, *Labill. Nov. Holl. Pl.* ii. 64. t. 214.—Austral.
graminifolium, *Sw. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 49.—Austral.
guttatum, *R. Br. Prod.* 571.—Austral.
hebegynum, DC. Prod. vii. 782 = reduplicatum.
hirsutum, *R. Br. Prod.* 568.—Austral.
hispidum, Lindl. Swan River App. 29 = ciliatum.
Hookeri, Planch. in Fl. des Serres, vi. (1850–51) 229 = dichotomum.
imbricatum, *Benth. in Enum. Pl. Hueg.* 73.—Austral.
inundatum, R. Br. Prod. 571 = despectum.
involucratum, F. Muell. Fragm. i. 154 = breviscapum.
junceum, *R. Br. Prod.* 569.—Austral.
Kunthii, *Wall. Cat.* n. 3759.—Ind. or.; Burma.
laricifolium, *Rich. in Pers. Syn.* ii. 210.—Austral.

STYLIDIUM:—
laricifolium, Walp. Rep. ii. 704, sphalm. = reduplicatum.
laxiflorum, DC. Prod. vii. 782 = scabridum.
Lehmannianum, Sond. in Lehm. Pl. Preiss. i. 375 = rupestre.
lepidum, *F. Muell. ex Benth. Fl. Austral.* iv. 27.—Austral.
leptobotrydium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = seq.
leptobotrys, DC. Prod. vii. 783 = crassifolium.
leptocalyx, Sond. in Lehm. Pl. Preiss. i. 373 = scabridum.
leptophyllum, DC. Prod. vii. 783.—Austral.
leptorhizum, F. Muell. Fragm. i. 148.—Austral.
leptostachyum, Lindl. Swan River App. 28 = carnosum.
Lessoni, DC. Prod. vii. 337 = falcatum?
limbatum, *F. Muell. Fragm.* x. 57.—Austral.
Lindleyanum, Sond. in Lehm. Pl. Preiss. i. 390 = calcaratum.
lineare, Sieber, ex DC. Prod. vii. 333 = graminifolium.
lineare, *Sw. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 50. t. 1. f. 1.—Austral.
lineatum, *Sond. in Lehm. Pl. Preiss.* i. 376.—Austral.
lobuliflorum, *F. Muell. Fragm.* i. 153.—Austral.
longifolium, Rich. in Pers. Syn. ii. 210 = pilosum.
longitubum, *Benth. in Enum. Pl. Hueg.* 73.—Austral.
luteum, *R. Br. Prod.* 570.—Austral.
marginatum, Sond. in Lehm. Pl. Preiss. i. 379 = diversifolium.
melastachys, R. Br. Prod. 568 = graminifolium.
mitrasacmoides, F. Muell. Fragm. i. 150 = alsinoides.
mucronifolium, Hook. Bot. Mag. t. 4538 = dichotomum.
mucronifolium, Sond. in Lehm. Pl. Preiss. i. 383 = leptophyllum.
muscicola, *F. Muell. Fragm.* i. 153.—Austral.
nudum, Lindl. Swan River App. 29 = glaucum.
obtusatum, Sond. in Lehm. Pl. Preiss. i. 382 = petiolare.
pachyrhizum, *F. Muell. Fragm.* i. 152.—Austral.
pedunculatum, *R. Br. Prod.* 571.—Austral.
perminutum, F. Muell. Fragm. vi. 78 = perpusillum.
perpusillum, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 266.—Austral.
petiolare, *Sond. in Lehm. Pl. Preiss.* i. 382.—Austral.
piliferum, *R. Br. Prod.* 569.—Austral.
pilosum, *Labill. Nov. Holl. Pl.* ii. 63. t. 213.—Austral.
pilosum, Sond. in Lehm. Pl. Preiss. i. 371 = reduplicatum.
planifolium, Poir. Encyc. Suppl. v. 412 = lineare.
plantagineum, Sond. in Lehm. Preiss. i. 371 = pilosum.
polystachium, Rich. in Pers. Syn. ii. 210 = graminifolium?
Preissii, *F. Muell. Fragm.* iii. 122.—Austral.
proliferum, DC. Prod. vii. 783 = bulbiferum.
propinquum, R. Br. Prod. 572 = adnatum.
pruinosum, Sond. in Lehm. Pl. Preiss. i. 379 = diversifolium.
pubigerum, *Sond. l. c.* 383.—Austral.
pulchellum, *Sond. l. c.* 381.—Austral.
pycnostachyum, *Lindl. Swan River App.* 29.—Austral.
pygmaeum, *R. Br. Prod.* 571.—Austral.
radicans, Sond. in Lehm. Pl. Preiss. i. 381 = repens.
ramosum, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 57.—Hab.?
recurvum, R. Grah. in Bot. Mag. t. 3913 = bulbiferum.
reduplicatum, *R. Br. Prod.* 568.—Austral.
repens, *R. Br. l. c.* 571.—Austral.
rhynchocarpum, *Sond. in Lehm. Pl. Preiss.* i. 389.—Austral.
rigidulum, Sond. l. c. = streptocarpum.
robustum, *Sond. l. c.* 378.—Austral.
roseum, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 137 = tenellum.
rotundifolium, *R. Br. Prod.* 571.—Austral.
rupestre, *Sond. in Lehm. Pl. Preiss.* i. 375.—Austral.
saxifragoides, Lindl. Swan River App. 28 = ciliatum.
scabridum, *Lindl. l. c.*—Austral.

STYLIDIUM:—
scandens, *R. Br. Prod.* 570.—Austral.
scariosum, DC. Prod. vii. 783 = junceum.
schizanthum, *F. Muell. Fragm.* i. 152.—Austral.
schoenoides, DC. Prod. vii. 782 = reduplicatum.
semipartitum, F. Muell. Fragm. i. 147 = leptorhizum.
serrulatum, Rich. in Pers. Syn. ii. 210 = graminifolium.
setaceum, Labill. Nov. Holl. Pl. ii. 65 = spinulosum ?
setigerum, DC. Prod. vii. 782 = ciliatum.
sinicum, Hance, in Walp. Ann. ii. 1030 = uliginosum.
soboliferum, *F. Muell. in Hook. Kew Journ.* viii. (1856) 162.—Austral.
spathulatum, *R. Br. Prod.* 569.—Austral.
spinulosum, *R. Br. l. c.*—Austral.
squamellosum, DC. Prod. vii. 782 = caespitosum.
stipitatum, Benth. in Enum. Pl. Hueg. 72 = Levenhookia stipitata.
streptocarpum, *Sond. in Lehm. Pl. Preiss.* i. 385.—Austral.
striatum, *Lindl. Swan River App.* 28.—Austral.
subulatum, *Hook. f. Handb. N. Zeal. Fl.* 168.—N. Zel.
sulcatum, Steud. Nom. ed. I. 818; ed. II. ii. 649 = falcatum.
tenellum, R. Br. Prod. 571 = uliginosum.
tenellum, *Sw. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 51.—Ind. or.; Burma.
tenerrimum, *F. Muell. Fragm.* i. 150.—Austral.
tenerum, Spreng. Syst. iii. 749 = uliginosum.
tenue, Sond. in Lehm. Pl. Preiss. i. 380 = Brunonianum.
tenuifolium, R. Br. Prod. 570 = laricifolium.
thesioides, DC. Prod. vii. 783 = canaliculatum.
thyrsiforme, DC. l. c. = pycnostachyum.
trichopodum, *F. Muell. Fragm.* x. 86.—Austral.
uliginosum, Kurz, in Flora, lv. (1872) 303 = Kunthii.
uliginosum, *Sw. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 52.—As. et Austral. trop.
umbellatum, Labill. Nov. Holl. Pl. ii. 66. t. 217 = graminifolium ?
uniflorum, *Sond. in Lehm. Pl. Preiss.* i. 381.—Austral.
utricularioides, *Benth. in Enum. Pl. Hueg.* 73.—Austral.
verticillatum, *F. Muell. Fragm.* iv. 94.—Austral.
violaceum, *R. Br. Prod.* 569.—Austral.
Wightianum, Wall. Cat. n. 3760 (Quid?).—Ind. or.

STYLIMNUS, Rafin. in Journ. Phys. lxxxix. (1819) 100 = **Pluchea**, Cass. (Compos.).
maritimus, Rafin. ex DC. Prod. v. 452 = P. camphorata.

STYLIS, Poir. Encyc. Suppl. v. 260 (1817) = **Marlea**, Roxb. (Cornac.).
chinensis, Poir. l. c. = M. begonifolia.

STYLISMA, Rafin. Neogenyt. 2 (1825) = **Breweria**, R. Br. (Convolv.).
aquatica, Rafin. l. c.; Chapm. Fl. S. U. St. 346 = B. aquatica.
elliptica, Rafin. New Fl. Am. iv. 55 (Quid?).—Am. bor.
evolvuloides, Choisy, Conv. Rar. 160 = B. humistrata.
heterophyla, Rafin. New Fl. Am. iv. 54 (Quid?).—Am. bor.
humistrata, Chapm. Fl. S. U. St. 346 = B. humistrata.
pedunculares, Rafin. New Fl. Am. iv. 55 (Quid?).—Am. bor.
Pickeringii, A. Gray, Man. Bot. N. U. St. ed. II. 335 = B. Pickeringii.
Sherardi, Rafin. New Fl. Am. iv. 55; Fl. Tellur. iv. 83 (Quid?).—Am. bor.
tenella, Rafin. Neogenyt. 2 = B. humistrata.

STYLISMUS, Spach, Hist. Vég. Phan. ix. 94 (1840) = praec.

STYLISTA, Rafin. Sylva Tellur. 112 (1838) = **Cleome**, Linn. (Cappar.).

STYLOBASIUM, Desf. in Mém. Mus. Par. v. (1819) 37. t. 2. *ROSACEAE*, Benth. & Hook. f. i. 609.
MACROSTIGMA, Hook. Ic. Pl. t. 412 (1842).
lineare, *Nees, in Lehm. Pl. Preiss.* i. 95.—Austral.
spathulatum, *Desf. in Mém. Mus. Par.* v. (1819) 37. t. 2.—Austral.

STYLOCERAS, A. Juss. Euph. Tent. 53. t. 17 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 265.
columnare, *Muell. Arg. in DC. Prod.* xvi. 1. 10.—Bolivia.
Dombeyanum, Baill. Etud. Gén. Euph. 666; et Monog. Bux. 80 = laurifolium.
Kunthianum, *A. Juss. Euph. Tent.* 117. t. 17. f. 56.—Ecuador.
laurifolium, *H. B. & K. Nov. Gen. et Sp.* vii. 173. t. 638.—N. Granat.; Peruv.
macrostachyum, Poepp. ex Baill. Monog. Bux. 81 = Alchornea cordata.
subrotundum, Poepp. ex Baill. Adansonia, v. (1864–65) 240 = Alchornea glandulosa.

STYLOCHITON, Lepr. in Ann. Sc. Nat. Sér. II. ii. (1834) 184 (Stylochaeton). *AROIDEAE*, Benth. & Hook. f. iii. 969.
GUEINZIA, Sond. ex Schott, Aroideae, 10 (1853).
hypogeum, *Lepr. in Ann. Sc. Nat.* Sér. II. ii. (1834) 185.—Afr. trop.
lancifolium, *Kotschy & Peyr. Pl. Tinn.* 42. t. 20.—Afr. trop.
natalense, *Schott, Aroideae*, 10. t. 14.—Afr. austr.

STYLOCLINE, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 338. *COMPOSITAE* [Benth. & Hook. f. ii. 298].
ANCISTROCARPHUS, A. Gray, in Proc. Am. Acad. vii. (1868) 355.
acaulis, Kellogg, in Proc. Calif. Acad. vii. (1877) 112 = Evax caulescens.
filaginea, *A. Gray, in Proc. Am. Acad.* viii. (1873) 652.—Am. bor. occ.
gnaphaloides, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 338.—Calif.
Griffithii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 652.—Afghan.
micropoides, *A. Gray, Pl. Wright.* ii. 84.—Am. bor. occ.

STYLOCORYNA, Cav. Ic. iv. 45. t. 368 (1797) = **Randia**, Houst. (Rubiac.).
albituba, Miq. Fl. Ind. Bot. Suppl. 541.—Sumatra.
attenuata, Voigt, Hort. Suburb. Calc. 377 = Tarenna zeylanica.
auricularia, Wall. Cat. n. 8402 = Anomanthoda auriculata.
bispinosa, Griff. Notul. iv. 260 = R. longiflora.
brevifolia, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 106 (= *Webera lucens*).—Ind. or.
buruensis, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 129.—Ins. Buru.
celebica, Miq. l. c. 190.—Ins. Celebes.
coffeoides, A. Gray, in Proc. Am. Acad. iv. (1859) 309 = R. coffeoides (in Addend.).
conferta, Benth. in Hook. Niger Fl. 389 = Tarenna conferta.
coriacea, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 175.—Borneo.
corymbosa, Labill. Sert. Austro-Caled. 48. t. 48 = Olostyla corymbosa.
costata, Miq. Fl. Ind. Bat. ii. 203.—Sumatra.
cuspidata, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. iii. 4.—Afr. austr.
dasyphylla, Miq. Fl. Ind. Bat. ii. 205.—Sumatra.
densiflora, Steud. Nom. ed. II. ii. 649 = R. densiflora.
dimorphophylla, Teysm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 402 = R. densiflora.
elliptica, Thw. Enum. Pl. Zeyl. 421 = Byrsophyllum ellipticum.
Forsteniana, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 191.—Ins. Celebes.
fragilis, D. Dietr. Syn. Pl. i. 794 = seq.
fragrans, Blume, Bijdr. 982 (= *Webera fragrans*).—Malaya.
grandiflora, Benth in Hook. Niger Fl. 390 = Tarenna grandiflora.
grandifolia, Miq. Fl. Ind. Bat. Suppl. 540.—Sumatra.
Harveyi, A. Gray, in Proc. Am. Acad. iv. (1859) 309.—Ins. Fiji.
hirsuta, Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 175.—Java.
Junghuhniana, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 237.—Java?
lateriflora, Span. in Linnaea, xv. (1841) 316.—Ins. Timor.

STYLOCORYNA:—
laxiflora, Blume, Bijdr. 983 (= *Webera fragrans*).—Malaya.
lucida, Miq. Fl. Ind. Bat. Suppl. 541 (= *Webera fragrans*).—Malaya.
macrophylla, Bartl. ex DC. Prod. iv. 377.—Ins. Philipp.
macrophylla, Wall. Cat. n. 8405 = Coptosapelta flavescens.
malabarica, DC. Prod. iv. 377 = Randia malabarica.
mollissima, Walp. Rep. ii. 517 (= *Webera mollissima*).—China.
monosperma, Wight & Arn. Prod. 401.—Ind. or.
nitidula, Benth. in Hook. Niger Fl. 390 = Tarenna nitidula.
odorata, Steud. Nom. ed. II. ii. 649 (= *Webera odorata*).—Ind. or.
oppositifolia, Steud. l. c. = R. densiflora.
orophila, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 237.—Sumatra.
ovata, Miq. Fl. Ind. Bat. ii. 203.—Java.
Pandaki, DC. Prod. iv. 377 = R. malabarica.
pavettoides, Miq. Fl. Ind. Bat. ii. 354.—Ins. Banca.
penangensis, Miq. l. c. 206 (= *Webera odorata*).—Reg. Himal.; Penang.
pepericarpa, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 223.—Ins. Fiji.
polycarpa, Miq. Fl. Ind. Bat. ii. 204.—Java.
pubescens, Bartl. ex DC. Prod. iv. 377.—Ins. Philipp.
pubiflora, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 419.—Ins. Timor.
racemosa, Cav. Ic. iv. 46. t. 368 = R. racemosa.
refracta, Miq. Fl. Ind. Bat. ii. 204.—Java.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. ii. 576.—Malabar.
rigida, Wight, Ic. t. 1064 = Tarenna zeylanica.
Roxburghii, Steud. Nom. ed. II. ii. 649 (= *Webera odorata*).—Reg. Himal.; Penang.
sambucina, A. Gray, in Proc. Am. Acad. iv. (1859) 309 (Tarennae sp.).—Ins. Pacif.
setigera, Walp. Ann. ii. 793.—Sumatra.
subsessilis, A. Gray, in Mem. Am. Acad. N. S. vi. (1858) 394.—Ins. Bonin.
tomentosa, Blume, Bijdr. 983.—Java.
truncata, Wall. Cat. n. 8403 = Triseiadia truncata.
Webera, Kurz, Andam. Rep. B. 10 (= *Webera Kurzii*).—Ins. Andam. et Nicob.
Webera, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 248 = Tarenna zeylanica.
Webera, Wall. Cat. n. 8401 = Tarenna zeylanica (*et Webera odorata, stellulata, Wallichii, campanuliflora*).—As. trop.

STYLOCORYNE, Wight & Arn. Prod. 400 (1834) = **Tarenna**, Gaertn. (Rubiac.).

STYLODISCUS, Benn. Pl. Jav. Rar. 133. t. 29 (1838) = **Bischofia**, Blume (Euphorb.).
trifoliatus, Benn. l. c. = B. javanica.

STYLODON, Rafin. Neogenyt. 2 (1825) = **Verbena**, Linn.
scabrum, Rafin. l. c. = V. carolina.

STYLOGLOSSUM, Breda, Orch. Kuhl. & Hass. t. [7] (1827) = **Calanthe**, R. Br. (Orchid.).
nervosa, Breda, l. c. = C. curculigoides.

STYLOGYNE, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91 = **Ardisia**, Sw. (Myrsin.).
Martiana, A. DC. l. c. = A. latipes.

STYLOLEPIS, Lehm. Sem. Hort. Hamb. (1828) 17.—Cf. Flora, xii. (1829) 1. Erg. 55 = **Podolepis**, Labill. (Compos.).
gracilis, Lehm. l. c. = P. gracilis.

STYLOMECON, Benth. in Journ. Linn. Soc. v. (1861) 74, lapsu = HYLOMECON, Maxim. = **Stylophorum**, Nutt. (Papav.).

STYLONCERUS, Spreng. Syst. iii. 356 (1826) = **Angianthus**, Wendl. (Compos.).
cylindraceus, Steetz, in Lehm. Pl. Preiss. i. 485 = A. humifusus.
humifusus, Spreng. Syst. iii. 356 = A. humifusus.
multiflorus, Nees, in Lehm. Pl. Preiss. ii. 244 = Rutidosis Pumilo.
suberectus, Steetz, l. c. i. 436 = A. humifusus.

STYLOPAPPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 431 = **Troximon**, Nutt. (Compos.).
elatus, Nutt. l. c. 433 = T. Nuttallii.
grandiflorus, Nutt. l. c. 422 = T. grandiflorum.
laciniatus, Nutt. l. c. = T. laciniatum.
Lessingii, D. Dietr. Syn. Pl. iv. 1322 = T. humile.
pumilus, D. Dietr. l. c. = T. aurantiacum.

STYLOPHORUM, Nutt. Gen. Am. ii. 7 (1818). *PAPAVERACEAE*, Benth. & Hook. f. i. 52.
DICRANOSTIGMA, Hook. f. & Thoms. Fl. Ind. i. 255 (1855).
HYLOMECON, Maxim. in Mém. Sav. Etr. Pétersb. ix. (1858) (Prim. Fl. Amur.) 36. t. 3.
cambricum, Spreng. Syst. ii. 570 = Meconopsis cambrica.
crassifolium, Steud. Nom. ed. II. ii. 650 = Meconopsis heterophylla.
diphyllum, *Nutt. Gen. Am.* ii. 7.—Am. bor.
heterophyllum, Steud. Nom. ed. II. ii. 650 = Meconopsis heterophylla.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 11.—Japon.
lactucoides, *Baill. Hist. Pl.* iii. 114.—Reg. Himal.
nepalense, Spreng. Syst. iv. Cur. Post. 203 = Meconopsis nepaulensis.
ohiense, Spreng. Syst. ii. 570 = diphyllum.
paniculatum, G. Don, Gen. Syst. i. 135 = Meconopsis nepaulensis.
petiolatum, Nutt. Gen. Am. ii. 8 = diphyllum.
simplicifolium, Spreng. Syst. iv. Cur. Post. 203 = Meconopsis simplicifolia.

STYLOSANTHES, Sw. Prod. Veg. Ind. Occ. 108 (1788); et Fl. Ind. Occ. 1280. t. 25 (1806). *LEGUMINOSAE*, Benth. & Hook. f. i. 517.
angustifolia, *Vog. in Linnaea*, xiii. (1838) 63.—Bras.
angustissima, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1200, nomen.—Guian.
aprica, Span. in Linnaea, xv. (1841) 192 = mucronata.
Bojeri, *Vog. in Linnaea*, xii. (1838) 68.—Afr. trop.
bracteata, *Vog. l. c.* 70.—Bras.
capitata, *Vog. l. c.*—Bras.
elatior, *Sw. in Vet. Acad. Handl. Stockh.* (1789) 296. t. 11. f. 2.—Am. bor.
erecta, *Beauv. Fl. Owar.* ii. 28. t. 77.—Afr. trop.
flavicans, *Baker, in Oliver, Fl. Trop. Afr.* ii. 156.—Afr. trop.
glutinosa, *H. B. & K. Nov. Gen. et Sp.* vi. 507. t. 595.—Mexic.
gracilis, *H. B. & K. l. c.* t. 596.—N. Granat.
guianensis, *Sw. in Vet. Acad. Handl. Stockh.* (1789) 296.—Guian.
guineensis, *G. Don, Gen. Syst.* ii. 281.—Afr. trop.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 357 = erecta.
hispida, Michx. Fl. Bor. Am. ii. 75 = elatior.
hispida, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 112 = guianensis.
humilis, *H. B. & K. Nov. Gen. et Sp.* vi. 506. t. 594.—Venezuela.
humilis, Rich. ex Hemsl. Biol. Centr. Am. Bot. i. 272 = procumbens.
juncea, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 19.—Paraguay.
leiocarpa, *Vog. in Linnaea*, xii. (1838) 64.—Bras.
longiseta, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 18.—Paraguay.
montevidensis, *Vog. in Linnaea*, xii. (1838) 67.—Bras.
mucronata, *Willd. Sp. Pl.* iii. 1166.—As. et Afr. trop.
procumbens, *Sw. Prod. Veg. Ind. Occ.* 108.—Ind. occ.
rigida, *Spreng. Syst.* iii. 310.—Bras.
ruellioides, *Mart. ex Benth. in Mart. Fl. Bras.* xv. 1. 90.—Bras.
scabra, *Vog. in Linnaea*, xii. (1838) 69.—Am. trop.
setosa, Harv. in Harv. & Sond. Fl. Cap. ii. 227 = mucronata.
surinamensis, Miq. in Linnaea, xviii. (1844) 567 = guianensis.
tenuifolia, G. Don, Gen. Syst. ii. 281 = angustifolia.
viscosa, *Sw. Prod. Veg. Ind. Occ.* 108.—Am. trop.

STYLURUS, Rafin. Fl. Ludov. 27 (1817) = **Ranunculus**, Tourn.
fistulosus, Rafin. l. c. 28 = R. multifidus?

STYLURUS, Salisb. in Knight, Prot. 115 (1809) = **Grevillea**, R. Br. (1810) (Proteac.).
buxifolia, Knight, l. c. 115 = G. buxifolia.
collina, Knight, l. c. 116 = G. buxifolia.

STYLYPUS, Rafin. Neogenyt. 3 (1825) = **Geum**, Linn. (Rosac.).
vernus, Rafin. l. c. = G. vernum.

STYPA, Garcke, Fl. Deutschl. ed. II. 367 (1851) = **Stipa**, Linn. (Gramin.).

STYPANDRA, R. Br. Prod. 278 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 794.
STYPONEMA, Salisb. Gen. Pl. Fragm. 67 (1866).
caerulea, R. Br. Prod. 279, in obs. = Pasithea coerulea.
caespitosa, *R. Br. l. c.*—Austral.
coarctata, R. Br. l. c. in obs. = Excremis ramosa.
frutescens, Knowles & Westc. Flor. Cab. ii. (1838) 69 = glauca.
glauca, *R. Br. Prod.* 279.—Austral.
grandiflora, Lindl. Swan River App. 57 = glauca.
imbricata, R. Br. Prod. 279 = glauca.
propinqua, A. Cunn. in Hook. Bot. Mag. t. 3417 = glauca.
scabra, R. Br. Prod. 279 = Agrostocrinum stypandroides.
scoparia, Endl. in Lehm. Pl. Preiss. ii. 35 = glauca.
umbellata, *R. Br. Prod.* 279.—Austral.
virgata, Endl. in Lehm. Pl. Preiss. ii. 35 = glauca.

STYPHANIA, C. Muell. in Walp. Ann. vii. 66 (1868), sphalm. = **Stephania**, Lour. (Menisp.).

STYPHELIA, Sm. Spec. Bot. New Holl. 45. t. 14 (1793). *EPACRIDEAE*, Benth. & Hook. f. ii. 610.
SOLENISCIA, DC. Prod. vii. 737 (1839).
abietina, Labill. Nov. Holl. Pl. i. 48. t. 68 = Cyathodes abietina.
acerosa, Soland. ex Gaertn. Fruct. ii. 78 = Cyathodes acerosa.
acicularis, F. Muell. Census, 106 = Leucopogon acicularis.
acuminata, Spreng. Syst. i. 659 = Leucopogon acuminatus.
adscendens, *R. Br. Prod.* 537.—Austral.
affinis, Spreng. Syst. i. 658 = Leucopogon lanceolatus.
aggregata, Spreng. l. c. 657 = Acrotriche aggregata.
Allittii, F. Muell. Fragm. vi. 34 = Leucopogon Allittii.
alternifolia, Spreng. Syst. i. 655 = Leucopogon alternifolius.
amplexicaulis, Rudge, in Trans. Linn. Soc. viii. (1807) 292. t. 8 = Leucopogon amplexicaulis.
androstoma, F. Muell. Fragm. vi. 55 = Cyathodes empetrifolia.
angustifolia, DC. Prod. vii. 735 = laeta.
apiculata, Spreng. Syst. i. 656 = Leucopogon apiculatus.
appressa, Spreng. l. c. 658 = Leucopogon appressus.
assimilis, F. Muell. Census, 106 = Leucopogon assimilis.
attenuata, F. Muell. l. c. 106 = Leucopogon attenuatus.
australis, F. Muell. Fragm. vi. 43 = Leucopogon australis.
Baxteri, F. Muell. l. c. 35 = Astroloma Baxteri.
biflora, Spreng. Syst. i. 659 = Leucopogon biflorus.
Billardierii, F. Muell. Fragm. vi. 43 = Cyathodes glauca.
blepharolepis, F. Muell. l. c. 48 = Leucopogon blepharolepis.
blepharophylla, F. Muell. l. c. 34 = Leucopogon elegans.
Bossiaea, F. Muell. l. c. 47 = Leucopogon Bossiaea.
brachycephala, F. Muell. l. c. 34 = Leucopogon cucullatus.
brachyloma, F. Muell. l. c. 39 = Brachyloma Preissii.
bracteolaris, F. Muell. Census, 106 = Leucopogon bracteolaris.
brevicuspis, F. Muell. l. c. = Leucopogon brevicuspis.
breviflora, F. Muell. l. c. 107 = Leucopogon breviflorus.
Brownii, Spreng. Syst. v. 683 (Index) = Leucopogon reflexus.
Candolleana, F. Muell. Fragm. vi. 38 = Astroloma Candolleanum.
capitellata, F. Muell. l. c. 31 = Leucopogon capitellatus.

STYPHELIA:—
carinata, Spreng. Syst. i. 658, partim = Leucopogon carinatus.
ciliata, F. Muell. Fragm. vi. 42 = Brachyloma ciliatum.
collina, Labill. Nov. Holl. Pl. i. 47. t. 65 = Leucopogon collinus.
compacta, F. Muell. Census, 106 = Leucopogon compactus.
compacta, Spreng. Syst. i. 657 = Astroloma compactum.
conantha, F. Muell. Fragm. vi. 30 = Conostephium planifolium.
concinna, F. Muell. Census, 106 = Leucopogon concinnus.
concurva, F. Muell. Fragm. vi. 36 = Leucopogon concurvus.
conferta, F. Muell. Census, 106 = Leucopogon confertus.
Conostephium, F. Muell. Fragm. vi. 40 = Conostephium pendulum.
conostephoides, F. Muell. l. c. 34 = Leucopogon conostephioides.
cordata, Labill. Nov. Holl. Pl. i. 46. t. 63 = Acrotriche ovalifolia.
cordata, F. Muell. Census, 106 = Leucopogon cordatus.
cordifolia, F. Muell. Fragm. viii. 54 = Leucopogon cordifolius.
corifolia, F. Muell. Census, 105 = Leucopogon corifolius.
cornifolia, Rudge, in Trans. Linn. Soc. viii. (1807) t. 9 = Trochocarpa laurina.
corynocarpa, F. Muell. Census, 107 = Leucopogon corynocarpus.
costata, *F. Muell. in Wing's South. Sc. Record* (April, (1885).—Austral.
crassiflora, F. Muell. Fragm. vi. 40 = Leucopogon crassiflorus.
crassifolia, F. Muell. l. c. 33 = Leucopogon crassifolius.
cryptantha, F. Muell. Census, 106 = Leucopogon cryptanthus.
cucullata, Spreng. Syst. i. 656 = Leucopogon cucullatus.
cuneifolia, F. Muell. Fragm. vi. 37 = Astroloma compactum.
Cunninghami, F. Muell. l. c. 39 = Melichrus urceolatus.
cuspidata, Spreng. Syst. i. 657 = Leucopogon cuspidatus.
cymbiformis, F. Muell. Fragm. vi. 34 = Leucopogon cymbiformis.
Cymbula, Spreng. Syst. iv. Cur. Post. 67 = Leucopogon Cymbulae.
daphnoides, Sm. Spec. Bot. New Holl. 48 = Brachyloma daphnoides.
dealbata, Spreng. Syst. i. 659 = Cyathodes dealbata.
deformis, Spreng. l. c. 658 = Leucopogon deformis.
denticulata, Spreng. l. c. = Astroloma humifusum.
denudata, Spreng. Syst. iv. Cur. Post. 67 = Leucopogon microphyllus.
depressa, F. Muell. Fragm. vi. 42 = Brachyloma depressum.
depressa, Spreng. Syst. i. 655 = Acrotriche depressa.
distans, Spreng. l. c. = Leucopogon distans.
divaricata, Spreng. l. c. 658 = Acrotriche divaricata.
Drummondi, F. Muell. Fragm. vi. 37 = Astroloma Drummondii.
dura, F. Muell. Census, 107 = Leucopogon durus.
elegans, Sond. in Lehm. Pl. Preiss. i. 296 = tenuiflora.
elliptica, Sm. Spec. Bot. New Holl. 49 = Monotoca elliptica.
empetrifolia, F. Muell. in Papers Roy. Soc. Tasm. (1874) 86 = Monotoca empetrifolia.
Epacridis, F. Muell. Fragm. vi. 38 = Astroloma divaricatum.
epacrioides, Hort. ex G. Don, in Loud. Hort. Brit. 65, nomen (Quid?).—Austral.
ericoides, Sm. Spec. Bot. New Holl. 48 = Leucopogon ericoides.
erubescens, F. Muell. Fragm. vi. 33 = Leucopogon Oxycedrus.
esquamata, Spreng. Syst. i. 658 = Leucopogon esquamatus.
exarrhena, F. Muell. Fragm. vi. 31 = pusilliflora.
exolasia, F. Muell. l. c. 34 = Leucopogon exolasius.
fasciculiflora, F. Muell. l. c. viii. 55 = Acrotriche fasciculiflora.
fastigiata, Spreng. Syst. iv. Cur. Post. 67 = Leucopogon esquamata.

STYPHELIA:—
fimbriata, F. Muell. Census, 106 = Leucopogon fimbriatus.
flavescens, F. Muell. Fragm. vi. 33 = Leucopogon flavescens.
flexifolia, Spreng. Syst. i. 659 = Leucopogon flexifolius.
florulenta, F. Muell. Census, 106 = Leucopogon florulentus.
Fraseri, F. Muell. Fragm. vi. 56 = Leucopogon Fraseri.
geissoloma, F. Muell. l. c. 39 = Brachyloma concolor.
gibbosa, F. Muell. Census, 106 = Leucopogon gibbosus.
Gilbertii, F. Muell. l. c. = Leucopogon Gilbertii.
glabella, F. Muell. Fragm. vi. 32 = Leucopogon phyllostachys.
glabella, Spreng. Syst. i. 655 = Leucopogon glabellus.
glacialis, F. Muell. Census, 106 = Leucopogon glacialis.
glauca, Labill. Nov. Holl. Pl. i. 45. t. 61 = Monotoca lineata.
glaucescens, Sieber, ex Spreng. Syst. iv. Cur. Post. 67 = triflora.
gnaphalioides, F. Muell. Census, 106 = Leucopogon gnaphalioides.
Gnidium, Vent. Jard. Malm. 23 = Leucopogon Richii.
gracilis, Spreng. Syst. i. 658 = Leucopogon gracilis.
gracillima, F. Muell. Fragm. vi. 34 = Leucopogon gracillimus.
grandiuscula, F. Muell. l. c. 47 = Leucopogon atherolepis.
Hainesii, *F. Muell. l. c.* iv. 96. t. 28.—Austral.
hirsuta, F. Muell. l. c. vi. 31 = Leucopogon hirsutus.
hirtella, F. Muell. Census, 107 = Leucopogon hirtellus.
Hookeri, F. Muell. Fragm. vi. 44 = Cyathodes ascendens.
humifusa, Pers. Syn. i. 174 = Astroloma humifusum.
imbricata, Spreng. Syst. i. 656 = Leucopogon imbricatus.
interrupta, Spreng. l. c. = Leucopogon interruptus.
involucrata, Spreng. l. c. 655 = Pentachondra involucrata.
juniperina, Pers. Syn. i. 174 = Leucopogon juniperinus.
laeta, *R. Br. Prod.* 537.—Austral.
lanceolata, Sm. Spec. Bot. New Holl. 49 = Leucopogon lanceolatus.
lasionema, F. Muell. Fragm. vi. 40 = Astroloma stomarrhena.
lasiostachya, F. Muell. Census, 106 = Leucopogon lasiostachyus.
latifolia, R. Br. Prod. 537 = laeta.
ledifolia, F. Muell. Census, 107 = Monotoca ledifolia.
leptantha, F. Muell. l. c. = Leucopogon leptanthus.
leptospermoides, Spreng. Syst. i. 659 = Leucopogon leptospermoides.
Leucopogon, *F. Muell. Fragm.* iv. 97.—Austral.
Lindleyi, F. Muell. l. c. vi. 40 = Conostephium minus.
linifolia, F. Muell. l. c. 36 = Leucopogon melaleucoides.
lissanthoides, F. Muell. l. c. 33 = Leucopogon cuneifolius.
lobopogona, F. Muell. l. c. 39 = Brachyloma ericoides.
longiflora, F. Muell. l. c. viii. 54 = Astroloma longiflorum.
longifolia, *R. Br. Prod.* 537.—Austral.
Macraei, F. Muell. Fragm. vi. 46 = Leucopogon Maccraei.
macrocalyx, F. Muell. l. c. 37 = Astroloma macrocalyx.
malaica, Spreng. Syst. iv. Cur. Post. 67 = Leucopogon malayanus.
margarodes, Spreng. Syst. i. 657 = Leucopogon margarodes.
megacarpa, F. Muell. Fragm. vi. 32 = Leucopogon planifolius.
melaleucoides, *F. Muell. l. c.* iv. 97.—Austral.
Michiei, F. Muell. l. c. vi. 80 = Coleanthera myrtoides.
microcalyx, F. Muell. l. c. 37 = Astroloma microcalyx.
microdonta, F. Muell. Census, 105 = Astroloma microdontum.
micropkylla, Spreng. Syst. i. 656 = Leucopogon microphyllus.
minutiflora, F. Muell. Fragm. xi. 122 = Monotoca tamariscina.
Mitchellii, F. Muell. Census, 107 = Leucopogon Mitchellii.

STYPHELIA:—
montana, F. Muell. Fragm. vi. 45 = Lissanthe montana, Leucopogon Hookeri.
multiflora, Spreng Syst. i. 658 = Leucopogon multiflorus.
mutica, F. Muell. Fragm. vi. 45 = Leucopogon muticus.
neo-anglica, F. Muell. Census, 107 = Leucopogon neo-anglicus.
obovata, Labill. Nov. Holl. Pl. i. 48. t. 67 = Leucopogon revolutus.
obtecta, F. Muell. Census, 107 = Leucopogon obtectus.
obtusata, F. Muell. l. c. 106 = Leucopogon obtusatus.
Oldfieldii, F. Muell. l. c. = Leucopogon Oldfieldii.
Oldfieldii, F. Muell. l. c. 107 = Leucopogon ovalifolius.
oligarrhenoides, F. Muell. Fragm. ix. 47 = Monotoca oligarrhenoides.
opponens, F. Muell. l. c. vi. 48 = Leucopogon opponens.
oppositifolia, F. Muell. l. c. 32 = Leucopogon oppositifolius.
ovalifolia, Spreng. Syst. i. 656 = Acrotriche ovalifolia.
Oxycedrus, Labill. Nov. Holl. Pl. i. 49. t. 69 = Cyathodes acerosa.
ozothamnoides, F. Muell. Census, 106 = Leucopogon ozothamnoides.
pallida, Spreng. Syst. i. 658 = Astroloma pallidum.
parviflora, Andr. Bot. Rep. t. 287 = Leucopogon Richei.
parvifolia, F. Muell. in Papers Roy. Soc. Tasm. (1874) 86 = Cyathodes parvifolia.
patula, Spreng. Syst. i. 657 = Acrotriche patula.
pauciflora, Spreng. l. c. 658 = Leucopogon leptospermoides.
pendula, Spreng. l. c. 657 = Leucopogon pendulus.
pentapogona, F. Muell. Fragm. vi. 36 = Astroloma microphyllum.
phyllostachys, F. Muell. Census, 106 = Leucopogon phyllostachys.
pinifolia, Spreng. Syst. i. 659 = Astroloma pinifolium.
platyphylla, F. Muell. Fragm. vi. 37 = Astroloma tectum.
pleiosperma, F. Muell. l. c. 41 = Leucopogon pleiospermus.
pleurandroides, F. Muell. l. c. 32 = Leucopogon pleurandroides.
plumuliflora, F. Muell. l. c. 29 = Leucopogon plumuliflorus.
pluriloculata, F. Muell. l. c. 32 = Leucopogon pluriloculatus.
pogonocalyx, F. Muell. Census, 107 = Leucopogon pogonocalyx.
polymorpha, F. Muell. Fragm. vi. 31 = Leucopogon polymorphus.
polystachys, Spreng. Syst. i. 659 = Leucopogon polystachyus.
Preissii, F. Muell. Fragm. vi. 40 = Conostephium Preissii.
procumbens, Pers. Syn. i. 174 = Melichrus rotatus.
propinqua, Spreng. Syst. i. 658 = Leucopogon propinquus.
prostrata, F. Muell. Census, 105 = Astroloma prostratum.
pulchella, F. Muell. Fragm. vi. 34 = Leucopogon pulchellus.
pumila, Spreng. Syst. i. 656 = Pentachondra pumila.
pungens, F. Muell. Fragm. vi. 34 = Leucopogon propinquus.
pusilliflora, *F. Muell. l. c.* iv. 105.—Austral.
racemulosa, F. Muell. l. c. vi. 33 = Leucopogon racemulosus.
ramiflora, F. Muell. l. c. vi. 44 = Acrotriche fasciculiflora.
ramiflora, Spreng. Syst. i. 659 = Acrotriche ramiflora.
reflexa, Rudge, in Trans. Linn. Soc. x. (1811) 296. t. 19 = Leucopogon collinus?
reflexa, Spreng. Syst. i. 655 = Leucopogon reflexus.
revoluta, Spreng. l. c. 657 = Leucopogon revolutus.
Richei, Labill. Nov. Holl. Pl. i. 44. t. 60 = Leucopogon Richei.
rotata, F. Muell. Fragm. vi. 38 = Melichrus rotatus.
rotundifolia, F. Muell. l. c. 45 = Leucopogon cordifolius.
rotundifolia, Spreng. Syst. i. 655 = Leucopogon rotundifolius.

STYPHELIA:—
rubicunda, F. Muell. Fragm. vi. 31 = Leucopogon rubicundus.
rubricaulis, Spreng. Syst. i. 656 = Leucopogon revolutus.
rudis, F. Muell. Fragm. vi. 32 = Leucopogon assimilis.
rufa, F. Muell. l. c. 46. = Leucopogon rufus.
ruscifolia, Spreng. Syst. i. 656 = Leucopogon ruscifolius.
sapida, F. Muell. Fragm. vi. 42 = Lissanthe sapida.
scoparia, Sm. Spec. Bot. New Holl. 49 = Monotoca scoparia.
semiopposita, F. Muell. Fragm. vi. 49 = Leucopogon elatior.
serrulata, Labill. Holl. Pl. Nov. i. 45. t. 62 = Acrotriche serrulata.
setigera, Spreng. Syst. i. 658 = Leucopogon setiger.
Sonderi, F. Muell. Fragm. vi. 36 = Astroloma conostephioides.
splendens, Hort. ex Planch. in Fl. des Serres, x. (1854–55) 129 = Astroloma divaricatum.
sprengelioides, F. Muell. Census, 106 = Leucopogon sprengelioides.
spuria, Poir. Encyc. vii. 485 = Leucopogon ericoides.
squarrosa, F. Muell. Fragm. vi. 31 = Leucopogon squarrosus.
straminea, Spreng. Syst. i. 656 = Cyathodes straminea.
striata, Spreng. l. c. = Leucopogon striatus.
stricta, F. Muell. Census, 107 = Leucopogon strictus.
strigosa, Sm. Spec. Bot. New Holl. 49 = Lissanthe strigosa.
strongylophylla, F. Muell. Fragm. vi. 33 = Leucopogon strongylophyllus.
subulifolia, F. Muell. l. c. = Leucopogon insularis.
tamariscina, Spreng. Syst. i. 656 = Leucopogon tamariscinus.
Tameiameiae, F. Muell. Fragm. vi. 55 = Cyathodes Tameiameiae.
tecta, Spreng. Syst. i. 657 = Astroloma tectum.
tenuiflora, *Benth. Fl. Austral.* iv. 148.—Austral.
tenuifolia, Lindl. Swan River App. 25, sphalm. = praec.
tenuis, F. Muell. Census, 106 = Leucopogon tenuis.
tetragona, F. Muell. Fragm. vi. 31 = Leucopogon tetragonus.
thymifolia, F. Muell. Census, 106 = Leucopogon thymifolius.
triandra, Spreng. Syst. iv. Cur. Post. 67 = triflora.
trichocarpa, Labill. Nov. Holl. Pl. i. 47. t. 66 = Leucopogon ericoides.
triflora, *Andr. Bot. Rep.* t. 72.—Austral.
trochocarpoides, *F. Muell. Papuan Pl.* 107.—N. Guin.
tubiflora, *Sm. Spec. Bot. New Holl.* 45. t. 14.—Austral.
unilateralis, F. Muell. Census, 106 = Leucopogon unilateralis.
urceolata, F. Muell. Fragm. vi. 38 = Melichrus urceolatus.
variifolia, F. Muell. l. c. 32 = Leucopogon glabellus.
verticillata, Spreng. Syst. i. 656 = Leucopogon verticillatus.
villosa, Dryand. in Kon. & Sims, Ann. Bot. ii. (1806) 513 = Leucopogon juniperinus.
villosa, Spreng. Syst. i. 657 = Leucopogon revolutus?
virgata, Labill. Nov. Holl. Pl. i. 46. t. 64 = Leucopogon virgatus.
viridiflora, R. Br. Prod. 537 = viridis.
viridis, *Andr. Bot. Rep.* t. 312.—Austral.
Woodsii, F. Muell. Census, 107 = Leucopogon Woodsii.
xerophylla, F. Muell. Fragm. vi. 38 = Astroloma xerophyllum.

STYPHNOLOBIUM, Schott, in Wien. Zeitschr. iii. (1830) 844; ex Endl. in Linnaea, vi. (1831) Litt. 54 = **Sophora**, Linn. (Legumin.).
affine, Walp. Rep. i. 807 = Sophora affinis.
japonicum, Schott, in Wien. Zeitschr. iii. (1830) 844; ex Endl. in Linnaea, vi. (1831) Litt. 55 = Sophora japonica.

STYPHONIA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. 230 (1791); Phil. Bot. ii. 70 (1791) = **Lavandula**, Tourn. (Labiat.).

STYPHONIA, Nutt. in Torr. et Gray, Fl. N. Am. i. 220 (1838) = **Rhus**, Tourn. (Anacard.).
atra, Nutt. l. c. = Oncocarpus vitiensis.
integrifolia, Nutt. l. c. = R. integrifolia.
mollis, Nutt. l. c. = R. mollis.
serrata, Nutt. l. c. = R. integrifolia.

STYPONEMA, Salisb. Gen. Pl. Fragm. 67 (1866) = **Stypandra**, R. Br. (Liliac.).

STYRANDRA, Rafin. in Am. Monthly Mag. (1818) 266 = **Maianthemum**, Weber (Liliac.).
bifolia, Rafin. Med. Fl. ii. 84 = M. Convallaria.
canadensis, Rafin. in Am. Monthly Mag. (1818) 266 = M. Convallaria.

STYRAX, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 143 (1737). *STYRACEAE*, Benth. & Hook. f. ii. 669.
Benzoin, Hayne, Arzn. Gew. xi. t. 24 (1829).
Cyrta, Lour. Fl. Cochinch. 278 (1790).
Epigenia, Vell. Fl. Flum. 183 (1825); iv. tt. 137, 138 (1827).
Foveolaria, Ruiz & Pav. Prod. 57. t. 9 (1794), partim.
Strigilia, Cav. Diss. vii. 358. t. 201 (1789).
Tremanthus, Pers. Syn. i. 467 (1805).
Trichogamila, P. Br. Hist. Jamaic. 218 (1756).
acuminatum, *Pohl, Pl. Bras. Ic.* ii. 58. t. 138.—Bras.
agreste, *G. Don, Gen. Syst.* iv. 5.—Cochinch.
album, *Mart. ex Spreng. Syst.* ii. 285.—Bras.
alutaceum, *Seub. in Mart. Fl. Bras.* vii. 190.—Bras.
ambiguum, *Seub. l. c.* 192.—Bras.
americanum, *Lam. Encyc.* i. 82.—Am. bor.
angustifolium, *Buc'hoz, Herb. Am.* t. 26.—Hab.?
argenteum, *Presl, Rel. Haenk.* ii. 60.—Mexic.
aureum, Mart. Reise Bras. i. 551 = Pamphilia aurea.
barbatum, Willd. ex Seub. in Mart. Fl. Bras. vii. 188 = glabrum.
benjuiferum, Stokes, Bot. Mat. Med. ii. 517 = S. Benzoin.
Benzoin, *Dryand. in Phil. Trans.* lxxvii. (1787) II. 308. t. 12.—Malaya.
californicum, *Torr. in Smithson. Contrib.* vi. (1854) 4.—Calif.
camporum, *Pohl, Pl. Bras. Ic.* ii. 53. t. 136.—Bras.
Candollei, *Seub. in Mart. Fl. Bras.* vii. 189.—Bras.
chrysasterum, *Pohl, Pl. Bras. Ic.* ii. 59. t. 139.—Bras.
cordatum, *A. DC. in DC. Prod.* viii. 267.—Peruv.
cotinifolium, Salisb. Prod. 286 = officinale.
crotonoides, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 589.—Singapore.
crymophyllum, G. Don, Gen. Syst. iv. 5, sphalm. = *erymophyllum*.
ellipticum, *Jungh. & De Vriese, in De Vriese, Pl. Nov. Ind. Batav.* fasc. i. (1845) 10.—Sumatra.
erymophyllum, Pohl, Pl. Bras. Ic. ii. 57. t. 137 = glabratum.
ferrugineum, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. I. (1823) 88.—Bras.
ferrugineum, Pohl, Pl. Bras. Ic. ii. 55. t. 135 = Pohlii.
Finlaysonianum, *Wall. Cat.* n. 4403.—Siam?
floribundum, Griff. Notul. iv. 287 = serrulatum.
floridum, *Pohl, Pl. Bras. Ic.* ii. 54. t. 134.—Bras.
fulvum, Klotzsch, ex A. DC. in DC. Prod. viii. 264 = longiflorum.
Gardnerianum, *A. DC. l. c.* 263.—Bras.
glabratum, *Schott, in Spreng. Syst.* iv. *App.* 406.—Bras.
glabrescens, *Benth. Pl. Hartw.* 66.—Mexic.
glabrum, Cav. Diss. vii. 340. t. 188. f. 1 = americanum.
glabrum, *Sw. Prod. Veg. Ind. Occ.* 74.—Am. trop.
grandiflorum, J. F. Gmel. Syst. 1633, sphalm. = grandifolium.
grandiflorum, Griff. Notul. iv. 287 = serrulatum.
grandifolium, *Ait. Hort. Kew.* ed. I. ii. 75.—Am. bor.
guineense, *G. Don, Gen. Syst.* iv. 5.—Afr. trop.
guyanense, A. DC. in DC. Prod. viii. 261 = glabrum.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 589.—Reg. Himal.
japonicum, *Sieb. & Zucc. Fl. Jap.* i. 53. t. 23.—Japon.
japurense, Mart. ex Seub. in Mart. Fl. Bras. vii. 188 = glabrum.
javanicum, Blume, Bijdr. 671 = Marlea begonifolia.
Klotzschii, *Seub. in Mart. Fl. Bras.* vii. 194.—Bras.
laeve, Walt. Fl. Carol. 140 = americanum.
laevigatum, Ait. Hort. Kew. ed. I. ii. 75 = americanum.
laevigatum, Sims, Bot. Mag. t. 921 = pulverulentum.

STYRAX:—
lancifolium, *Klotzsch, ex Seub. in Mart. Fl. Bras.* vii. 191.—Bras.
latifolium, *Pohl, Pl. Bras. Ic.* ii. 60. t. 140.—Bras.
leiophyllum, *Miers, in Lindl. Veg. Kingd.* 593ª.—Bras.
leprosum, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 282.—Bras.
Limoncillo, Miers, in Journ. Linn. Soc. xvii. (1879) 292 = Symplocos martinicensis.
longiflorum, *A. DC. in DC. Prod.* viii. 264.—Bras.
macrophyllum, *Schott, in Spreng. Syst.* iv. *App.* 405, nomen; *et ex Pohl, Pl. Bras. Ic.* ii. 61. t. 141.—Bras.
Martii, *Seub. in Mart. Fl. Bras.* vii. 194.—Bras.
nervosum, A. DC. in DC. Prod. viii. 263 = ferrugineum.
Obassia, *Sieb. & Zucc. Fl. Jap.* i. 93. t. 46.—Japon.
obliquinervium, A. DC. in DC. Prod. viii. 263 = chrysasterum?
oblongum, *A. DC. l. c.* 261.—Peruv.
obtusifolium, *Griseb. Cat. Pl. Cub.* 167.—Cuba.
occidentale, Thunb. Diss. Styr. 8 = glabrum.
octandrum, L'Hérit. ex Ait. Hort. Kew. ed. I. ii. 75 = americanum.
odoratissimum, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 304 = Finlaysonianum.
officinale, *Linn. Sp. Pl.* 444.—Europ. austr.; As. Min.
officinale, Walt. Fl. Carol. 140 = grandifolium.
officinarum, Crantz, Inst. ii. 382 = officinale.
ovale, Klotzsch, ex Seub. in Mart. Fl. Bras. vii. 194 = Klotzschii.
ovatum, *A. DC. in DC. Prod.* viii. 267.—Peruv.
pallidum, *A. DC. l. c.* 261.—Guian.
parvifolium, Herb. Berol. ex A. DC. l. c. 263 = floridum.
parvifolium, *Pohl, Pl. Bras. Ic.* ii. 53. t. 133.—Bras.
pauciflorum, *A. DC. in DC. Prod.* viii. 265.—Bras.
Pavonii, *A. DC. l. c.* 266.—Peruv.
Pentlandianum, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 229.—Bolivia.
platinifolium, *Engelm. ex Torr. in Smithson. Contrib.* vi. (1854) 4, in nota.—Texas.
Pohlii, *A. DC. in DC. Prod.* viii. 264.—Bras.
polyspermum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 590.—Reg. Himal.
Porterianum, Wall. Cat. n. 4401 = serrulatum.
psilophyllum, *A. DC. in DC. Prod.* viii. 266.—Guian.
pulverulentum, *Michx. Fl. Bor. Am.* ii. 41.—Am. bor.
punctatum, *A. DC. in DC. Prod.* viii. 264.—Bras.
racemosum, *A. DC. l. c.* 265.—Peruv.
reticulatum, Mart. Reise Bras. i. 550 = ferrugineum.
rigidum, Salisb. Prod. 286 = americanum.
Rossamala, Reinw. ex Steud. Nom. ed. II. ii. 651, nomen = Marlea begonifolia?
rugosum, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 61.—Burma.
serrulatum, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 415.—Reg. Himal.; Burma; China.
subcordatum, *Moric. Pl. Nouv. Am.* 117. t. 71.—Bras.
subdenticulatum, *Miq. Fl. Ind. Bat. Suppl.* 474.—Sumatra.
suberifolium, *Hook. & Arn. Bot. Beech. Voy.* 196. t. 40.—China.
subleprosum, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1087, nomen.—Guian.
subpaniculatum, *Jungh. & De Vriese, in De Vriese, Pl. Nov. Ind. Batav.* fasc. i. (1845) 9. t. 3.—Sumatra.
tinifolium, G. Don, Gen. Syst. iv. 5 = Clethra tinifolia.
tomentosum, *Humb. & Bonpl. Pl. Aequin.* ii. 72. t. 101.—Ecuador.
tomentosum, Spreng. Syst. iv. App. 405 = macrophyllum.
varians, Seub. in Mart. Fl. Bras. vii. 189 = chrysasterum.
villosum, *Blume, Bijdr.* 671.—Java.
virgatum, Wall. Cat. n. 4400 = serrulatum.

STYROSINIA, Rafin. Fl. Tellur. ii. 95 (1836) = **Gesneria**, Linn.
coccinea, Rafin. l. c. 96 = G. aggregata.

SUAEDA, Forsk. Fl. Aegypt. Arab. 69. t. 18 B (1775). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 66.
Belovia, Moq. in DC. Prod. xiii. II. 168 (1849).
Brezia, Moq. l. c. 167 (1849).
Calvelia, Moq. l. c. (1849).
Chenopodina, Moq. l. c. 159 (1849).

SUAEDA:—
Cochliospermum, Lag. Mem. Pl Barill. 58 (1817).
Dondia, Adans. Fam. ii. 261 (1763), nomen prius.
Einadia, Rafin. Fl. Tellur. iv. 121 (1836).
Lerchea, Hall. Hort. Gotting. 21 (1743).
Schanginia, C. A. Mey. in Ledeb. Fl. Alt. i. 394 (1829).
Schoberia, C. A. Mey. l. c. 395 (1829).
Sevada, Moq. in DC. Prod. xiii. II. 154 (1849).
Trikalis, Rafin. Fl. Tellur. iii. 47 (1836).
acuminata, *Moq. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 306.—Rossia.
aestuaria, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 328 = maritima.
africana, Hort. ex Moq. in DC. Prod. xiii. II. 158 = Salsola aphylla.
albida, Pall. Illustr. Pl. 53. t. 45 = Kochia hirsuta.
altissima, *Pall. l. c.* 49. t. 42.—Europ. austr.; As. occ.
ampullacea, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 360.—Mongolia.
anonyma, J. F. Gmel. Syst. 503 (Quid?).—Reg. Arab.
arcuata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 461.—Turkest.
asphaltica, *Boiss. Fl. Orient.* iv. 938.—Palaest.
australis, *Moq. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 318.—As.; Austral.; Polynes.
baccata, *Forsk. Fl. Aegypt. Arab.* 69.—Aegypt.
baccifera, *Pall. Illustr. Pl.* 48. t. 41.—Ross. mer.
bacciformis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 328 = maritima.
Balansae, Boiss. Diagn. Ser. II. iv. 75 = setigera.
Biebersteinia, C. Koch, in Linnaea, xvii. (1843) 312 = microphylla.
calceoliformis, *Moq. Chenop. Enum.* 128.—Am. bor.
californica, Nutt. ex Moq. in DC. Prod. xiii. II. 161 = maritima.
californica, *S. Wats. in Proc. Am. Acad.* ix. (1874) 89.—Calif.
chenopodioides, Pall. Illustr. Pl. 56 = maritima.
cinerea, Schur, Enum. Pl. Transs. 570 = Kochia scoparia.
Clemente, Heynh. Nom. ii. 712 = setigera.
corniculata, *Bunge, in Act. Hort. Petrop.* vi. (1879) 429.—Tibet.; Soongar.; Sibir.
crassifolia, Heynh. Nom. ii. 712 = maritima.
crassifolia, Pall. Illustr. Pl. 54. t. 46 = Kochia hirsuta.
dendroides, *Moq. Chenop. Enum.* 126.—Reg. Cancas.
depressa, S. Wats. Bot. King's Exped. 294 = plattensis.
diffusa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 88.—Am. bor. occ.
divaricata, *Moq. Chenop. Enum.* 123.—Chili.
filiformis, Dum. Fl. Belg. 22 = maritima.
foetida, *Moq. Chenop. Enum.* 132.—Hab.?
foliosa, *Moq. in DC. Prod.* xiii. II. 156.—Peruv.
fruticosa, Delile, Fl. Aeg. Illustr. 57 = monoica.
fruticosa, Dum. Fl. Belg. 22 = fruticosa.
fruticosa, *Forsk. Fl. Aegypt. Arab.* 70.—Reg. bor. temp.
fruticosa, Hook. & Arn. Bot. Beech. Voy. 387 = Torreyana.
fruticulosa, *Moq. Chenop. Enum.* 132.—Reg. Ural.
glauca, Bunge, in Bull. Acad. Pétersb. xxv. (1879) 362 = Stauntoni.
gracilis, Moq. Chenop. Enum. 123 = setigera.
Helenae, *Moq. in DC. Prod.* xiii. II. 158.—Ins. S. Helen.
heterocarpa, *Fenzl, in Ledeb. Fl. Ross.* iii. 781.—Reg. Caucas.; Persia.
heterophylla, Bunge, in Act. Hort. Petrop. vi. (1879) 429; Boiss. Fl. Orient. iv. 943 = Afghan.; Sibir.
hirsuta, Reichb. Fl. Germ. Excurs. 580 = Kochia hirsuta.
hortensis, *Forsk. Fl. Aegypt. Arab.* 70.—Aegypt.
hyssopifolia, Pall. Illustr. Pl. 44. t. 36 = Kochia hyssopifolia.
indica, Moq. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 316 = nudiflora.
indica, Seem. Bot. Voy. Herald, 406 = australis.
intermedia, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 296.—Am. bor. occ.
Jacquinii, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 329 = maritima.
Kareliniana, *Fenzl, in Ledeb. Fl. Ross.* iii. 789.—Sibir. altaic.
Kochii, Todaro, ex Nym. Consp. 630 = maritima.
laxifolia, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) reimp. 11 = fruticosa.

SUAEDA :—
leiosperma, Moq. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 306 = altissima.
linearis, *Moq. Chenop. Enum.* 130.—Am. bor.; Cuba.
linifolia, *Pall. Illustr. Pl.* 47. t. 40.—Ross. mer. or.
longifolia, C. Koch, in Linnaea, xxii. (1849) 188 = setigera.
macrocarpa, Moq. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 309 = maritima.
maritima, *Dum. Fl. Belg.* 22.—Reg. temp. bor. et austr.
maritima, S. Wats. in Bot. King's Exped. 294 = diffusa.
maxima, C. Koch, in Linnaea, xvii. (1843) 312 = altissima.
microphylla, C. Koch. l. c. = dendroides.
microphylla, Moq. Chenop. Enum. 127 = microphylla.
microphylla, *Pall. Illustr. Pl.* 52. t. 44.—Reg. Caucas.; Sibir.
microsperma, *Fenzl, in Ledeb. Fl. Ross.* iii. 785.—Soongaria; Tibet.
minima, *Fenzl, l. c.* 789.—Sibir. altaic.
minutiflora, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 194.—Calif.
mollis, Delile, Fl. Aegypt. Illustr. 57 = vermiculata.
monantha, C. Koch, in Linnaea, xvii. (1843) 313 = setigera.
monoica, *Forsk. Fl. Aegypt. Arab.* 70.—Afr. bor.; As. occ.; Ind. or.
monoica, Schimp. ex Boiss. Fl. Orient. iv. 986 = baccata.
muricata, Pall. Illustr. Pl. 43. t. 35 = Kochia muricata.
nudiflora, *Moq. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 316.—Ind. or.
nudiflora, Moq. in DC. Prod. xiii. II. 155 = maritima.
nudiflora, Thw. Enum. Pl. Zeyl. 246 = monoica.
obtusifolia, *Trautv. in Bull. Soc. Nat. Mosc.* xl. (1867) II. 62.—Turkestan.
occidentalis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 90.—Am. bor. occ.
Pallasiana, Heynh. Nom. ii. 712 = Kochia hirsuta.
paradoxa, *Bunge, in Act. Hort. Petrop.* vi. (1879) 427.—Bokhara.
parvifolia, Moq. Chenop. Enum. 131 = microsperma.
physophora, *Pall. Illustr. Pl.* 51. t. 43.—Reg. Casp.; Persia; As. bor.
pinnatifida, Delile, Desc. Egypt. 57. n. 298 = Salsola pinnatifida.
plattensis, *Nutt. ex Moq. in DC. Prod.* xiii. II. 164, in syn.—Am. bor. occ.
platyphylla, Ehrenb. ex Boiss. Fl. Orient. iv. 986 = baccata.
prostrata, Pall. Illustr. Pl. 55. t. 47 = microsperma.
pruinosa, Lange, in Kjoeb. Vidensk. Meddel. (1861) 45 = vera.
Przewalskii, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 360.—Mongolia.
pterantha, *Bunge, in Act. Hort. Petrop.* vi. (1879) 430.—Soongar.
puberula, *Moq. Chenop. Enum.* 122.—Hab.?
pygmaea, *Moq. l. c.* 127.—Hab.?
Rosmarinus, Ehrenb. ex Boiss. Fl. Orient. iv. 951 = Seidlitzia Rosmarinus.
salsa, *Pall. Illustr. Pl.* 46. t. 39.—Reg. Caucas.; Sibir.
sedifolia, Pall. l. c. 41. tt. 32–34 = Kochia sedoides.
sedoides, Steud. Nom. ed. II. ii. 651, sphalm. = praec.
setifera, Willk. & Lange, Prod. Fl. Hisp. i. 257 = Halogeton sativus.
setigera, *Moq. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 309.—Reg. Mediterr.; As. occ.
Sieversiana, Pall. Illustr. Pl. 45. t. 38 = Kochia scoparia.
spicata, Moq. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 317 = maritima.
splendens, Gren. & Godr. Fl. Fr. iii. 30 = setigera.
Stauntonii, *Moq. Chenop. Enum.* 131.—China.
suffrutescens, *S. Wats. in Proc. Am. Acad.* ix. (1874) 88.—Am. bor. occ.
tamariscina, Lindl. in Mitch. Journ. Trop. Austral. 239 = Enchylaena microphylla.
tomentosa, Lowe, in Trans. Camb. Phil. Soc. vi. (1838), reimp. 12 = Chenolea canariensis.
Torreyana, *S. Wats. in Proc. Am. Acad.* ix. (1874) 88.—Am. bor. occ.

SUAEDA :—
tortuosa, *Moq. Chenop. Enum.* 131.—N. Zel.
transoxana, Boiss. Fl. Orient. iv. 943 = pterantha.
triandra, Henck. Adumbr. Hort. Hal. 21 = Kochia hyssopifolia.
trigyna, *Heynh. Nom.* ii. 640, in syn.—Hab.?
vera, *Forsk. Fl. Aegypt. Arab.* 69.—Reg. Mediterr.
vermiculata, *Forsk. l. c.* 70.—Afr. bor.; Arab.
villosa, Lowe, ex Moq. in DC. Prod. xiii. II. 129 = Chenolea canariensis.

SUARDA, Nocca, ex Steud. Nom. ed. II. ii. 651 (1841) = **Eugenia**, Mich. (Myrtac.).
ligustrifolia, Nocca, ex Steud. l. c. = E. baruensis.

SUARDIA, Schrank, Pl. Rar. Hort. Monac. t. 58 (1819) = **Melinis**, Beauv. (Gramin.).
picta, Schrank, l. c. = M. minutiflora.

SUBERTIA, Wood, in Proc. Acad. Nat. Sc. Philad. 1868 (1869) 172, sphalm. = SEUBERTIA, Kunth = **Brodiaea**, Sm. (Liliac.).

SUBLIMIA, Comm. ex Mart. Hist. Nat. Palm. iii. 164 (1836?) = **Hyophorbe**, Gaertn. (Palm.).
aevidaps, Comm. ex Mart. l. c. 176 = H. amaricaulis.
amaricaulis, Comm. ex Mart. l. c. 309 = H. amaricaulis.
Areca, Comm. ex Mart. l. c. 169 = Areca Catechu.
centennina, Comm. ex Mart. l. c. 174 = Acanthophoenix crinita.
palmicaulis, Comm. ex Mart. l. c. 175 = Dictyosperma album.
vilicaulis, Comm. ex Mart. l. c. 164 = H. indica.

SUBRISIA, Rafin. Sylva Tellur. 43 (1838) = **Ehretia**, Linn. (Borag.).
petiolaris, Rafin. l. c. = E. petiolaris.

SUBULARIA, Forsk. Fl. Aegypt. Arab. 117 (1775) = **Schouwia**, DC. (Crucif.).
purpurea, Forsk. l. c. = Schouwia arabica.

SUBULARIA, Ray, ex Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 83.
alpina, Willd. Sp. Pl. iii. 424 = Arabis vochinensis.
aquatica, *Linn. Sp. Pl.* 642.—Europ.; Am. bor.
monticola, *A. Br. ex Schweinf. Beitr. Fl. Aethiop.* 76.—Abyss.
repens, Pritz. Ic. Ind. 1082, sphalm. = Littorella juncea.

SUCCISA, Neck. Elem. i. 109 (1790); Wallr. Sched. Crit. 46, in syn. (1822) = **Cephalaria**, Schrad. et **Scabiosa**, Tourn. (Dipsac.).
alpina, Moench, Meth. 488 = C. alpina.
altissima, Schur, Enum. Pl. Transs. 297 = Sc. Succisa.
ambrosioides, Spreng. Syst. i. 379 = C. corniculata.
amplexicaulis, Spreng. l. c. = Sc. amplexicaulis.
angustula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 49 = Sc. Succisa.
arvensis, Rafin. Fl. Tellur. iv. 94 = Sc. arvensis.
atropurpurea, Moench, Meth. 490 = Sc. australis.
aurigerana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 51 = Sc. Succisa.
australis, Reichb. Fl. Germ. Excurs. 196 = Sc. australis.
beugesiaca, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 51 = Sc. Succisa.
bipinnata, Moench, Meth. 489 = Sc. gramuntia.
brevis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 49 = Sc. Succisa.
cagiriensis, Jeanb. & Timb. in Bull. Soc. Phys. et Nat. Toul. iii. (1875–76) 580 = Sc. Succisa.
centauroides, Spreng. Syst. i. 379 = C. corniculata.
ciliata, Rafin. Fl. Tellur. iv. 94 = Sc. ciliata.
Columbaria, Moench, Meth. 489 = Sc. Columbaria.
corniculata, Sweet, Hort. Brit. ed. I. 204 = C. corniculata.
cretacea, Reichb. Ic. Fl. Germ. xii. 23. t. 701 = C. corniculata.
cuspidata, Jord. Brev. Pl. Nov. fasc. ii. 54 = Sc. Succisa.
decurrens, Spreng. Syst. i. 379 = C. attenuata.
dentata, Jord. & Fourr. Brev. Nov. fasc. ii. 54 = Sc. Succisa.
dichotoma, Rafin. Fl. Tellur. iv. 94 = Sc. dichotoma.

SUCCISA :—
diversifolia, Spreng. Neue Entdeck. i. 279 = Scabiosa banatica.
elliptica, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 579 = Sc. Succisa.
Fuchsii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 476 = Sc. Succisa.
fuscescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 48 = Sc. Succisa.
Fussiana, Heuff. ex Schur, Enum. Pl. Transs. 293 = C. radiata.
gigantea, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875–76) 579 = Sc. Succisa.
glabrata, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 54 = Sc. Succisa.
glabrata, Sweet, Hort. Brit. ed. I. 204 = Sc. Succisa.
gracilescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 50 = Sc. Succisa.
graminifolia, Moench, Meth. Suppl. 185 = Sc. graminifolia.
humilis, Spreng. Syst. i. 379 = C. attenuata.
incisa, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 51 = Sc. Succisa.
inflexa, Schur, Enum. Pl. Transs. 298 = Sc. australis.
joppica, Spreng. Syst. i. 378 = C. joppensis.
laetevirens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 53 = Sc. Succisa.
laevigata, Spreng. Syst. i. 379 = C. corniculata.
lancifolia, Moench, Meth. 488 = C. syriaca.
leucantha, Moench, l. c. 489 = C. leucantha.
macrocalycina, Moench, l. c. = Sc. ucranica.
microcephala, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 56 = Sc. Succisa.
microcephala, Willk. in Flora, xxxiv. (1851) 740 = Sc. microcephala.
palustris, Sass, Phaner. Fl. Oesels, 57 = Sc. Succisa.
parvula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 56 = Sc. Succisa.
pentaphylla, Moench, Meth. 488 = C. transylvanica.
pinnatifida, Lange, in Kjoeb. Vidensk. Meddel. (1861) 63 = Sc. Succisa?
platyphylla, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 48 = Sc. Succisa.
praemorsa, Aschers. Fl. Brandenb. 285 = Sc. Succisa.
pratensis, Moench, Meth. 489 = Sc. Succisa.
prativaga, Jord. & Four. Brev. Pl. Nov. fasc. ii. 52 = Sc. Succisa.
procera, Jord. & Fourr. l. c. = Sc. Succisa.
propera, Jord. & Fourr. l. c. 55 = Sc. Succisa.
pyrenaica, Jord. & Fourr. l. c. 55 = Sc. Succisa.
radiata, Schur, Enum. Pl. Transs. 293 = C. radiata.
rhodanensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 53 = Sc. Succisa.
rigida, Spreng. Syst. i. 379 = C. rigida.
sabauda, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 50 = Sc. Succisa.
scabra, Spreng. Syst. i. 379 = C. scabra.
stellata, Moench, Meth. 490 = Sc. stellata.
stricta, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 52 = Sc. Succisa.
subacaulis, N. Bernardin, in Cariot. Et. Fl. ed. V. ii. 301 = Sc. Succisa.
sylvatica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 59 = Sc. Succisa.
syriaca, Spreng. Syst. i. 378. = C. syriaca.
tardans, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 54 = Sc. Succisa.
tatarica, Spreng. Syst. i. 378 = C. tatarica.
transylvanica, Spreng. l. c. = C. transylvanica.
trifida, Spreng. l. c. 379 = C. attenuata.
uralensis, Spreng. Syst. i. 378 = C. corniculata.
ustulata, Spreng. l. c. 379 = C. attenuata.
viretorum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 49 = Sc. Succisa.
vogesiaca, Jord. & Fourr. l. c. 55 = Sc. Succisa.
vulgaris, J. & C. Presl, Fl. Čech. 31 = Sc. Succisa.

SUCCISOCREPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 104 = **Crepis**, Vaill. (Compos.).
succisaefolia, Fourr. l. c. = C. succisaefolia.

SUCCOVIA, Spreng. Anleit. ii. II. 721 (1818) = **Succowia**, Medic. (Crucif.).

SUCCOWIA, Dennst. Schluess. Hort. Malab. 32 (1818) = **Hiptage**, Gaertn. (Malpig.).
fimbriata, Dennst. l. c. = H. Madablota.

SUCCOWIA, Medic. Pflanzeng. i. 64 (1792) t. 1. f. 9. *CRUCIFERAE*, Benth. & Hook. f. i. 86.
balearica, *Medic. l. c.* 65.—Reg. Mediterr.
echinata, Moench, Meth. 265 = balearica.

SUCCUTA, Desmoul. Etudes Org. Cusc. 74 (1853) = **Cuscuta**, Tourn. (Convolv.).
alba, Desmoul. l. c. = C. Epithymum.

SUCHTELENIA, Karel. ex Meissn. Gen. Comm. 188 (1840); et in Bull. Soc. Nat. Mosc. (1841) 16. t. 2. *BORAGINEAE*, Benth. & Hook. f. ii. 845.
acanthocarpa, Karel. in Bull. Soc. Nat. Mosc. (1841) 17. t. 2 = cerinthifolia.
calycina, A. DC. in DC. Prod. x. 163 = cerinthifolia.
cerinthifolia, *Karel. ex Meissn. Gen. Comm.* 188.—Reg. Transcauc.; Turkest.
uniserialis, Ledeb. Fl. Ross. iii. 169 = cerinthifolia.

SUCKLEYA, A. Gray, in Proc. Am. Acad. xi. (1876) 103. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 54.
petiolaris, *A. Gray, l. c.*—Am. bor.

SUEDA, Edgew. in Journ. Linn. Soc. vi. (1862) 192, sphalm. = **Suaeda**, Forsk. (Chenop.).

SUENSONIA, Gaudich. ex Miq. Syst. Pip. 535 (1843) = **Piper**, Linn.

SUFFRENIA, Bell. in Mém. Acad. Turin, x, xi. (1802–3) 445 = **Ammannia**, Houst. et **Rotala**, Linn. (Lythrar.).
capensis, Harv. Thes. Cap. ii. 56. t. 189 = R. filiformis.
dichotoma, Miq. Fl. Ind. Bat. i. 1. 616 = Ammannia multiflora.
filiformis, Bell. in Mém. Acad. Turin, x, xi. (1802–3) 445 = R. filiformis.

SUGEROKIA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 144 (1867) = **Heloniopsis**, A. Gray (Liliac.).
japonica, Miq. l. c. 145 = H. japonica.

SUGILLARIA, Salisb. Gen. Pl. Fragm. 18 (1866) = **Scilla**, Linn. (Liliac.).
lanceaefolia, Salisb. l. c. = S. lanceaefolia.

SUITENIA, Stokes, Bot. Mat. Med. ii. 479 (1812) = **Swietenia**, Linn. (Meliac.).

SUITRAMIA, Reichb. Nom. 173 (1841) = **Svitramia**, Cham. (Melastom.).

SUKAMINEA, Rafin. New Fl. Am. iii. 44 (1836) = **Maclura**, Nutt. (Urtic.).

SUKANA, Adans. Fam. ii. 269 (1763) = **Celosia**, Linn. (Amarant.).

SUKSDORFIA, A. Gray, in Proc. Am. Acad. xv. (1880) 41. *SAXIFRAGEAE.*
violacea, *A. Gray, l. c.* 42.—Am. bor.

SULCANUX, Rafin. Sylva Tellur. 148 (1838) = **Psychotria**, Linn. (Rubiac.).

SULIPA, Blanco, Fl. Filip. ed. I. 497 (1837) (Genus delendum).
globosa, Blanco, l. c. ed. II. 348 = Planchonia valida.
Pseudopsidium, Blanco, l. c. ed. I. 497 = Gardenia Pseudopsidium.

SULITRA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 366 = **Lessertia**, DC. (Legumin.).
africana, Medic. l. c. = L. linearis?
herbacea, Medic. Phil. Bot. i. 202 = L. linearis.
perennans, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 366 = L. perennans.
pubescens, Moench, Meth. 164 = L. linearis.

SULLA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 372 = **Hedysarum**, Tourn. (Legumin.).
amoena, Medic. l. c. 373 = H. coronarium.
coronaria, Medic. Phil. Bot. i. 209 = H. coronarium.
flexuosa, Medic. l. c. = H. flexuosum.

SULLIVANTIA, Torr. & Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 22, in nota. *SAXIFRAGEAE*, Benth. & Hook. f. i. 636.
Ohionis, *Torr. & Gray, l. c.*; *et in Hook. Lond. Journ. Bot.* i. (1842) 229.—Am. bor.
oregana, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 292.—Am. bor. occ.

SULPITIA, Rafin. Fl. Tellur. iv. 37 (1836) = **Epidendrum**, Linn. (Orchid.).
odorata, Rafin. l. c. = E. odoratissimum.

SULZERIA, Roem. & Schult. Syst. iv. p. lx. 707 (1819) = **Anabata**, Willd. (Logan.?).
odorata, Roem. & Schult. l. c. 707 = A. odorata.

SUMBAVIA, Baill. Etud. Gén. Euph. 390 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 304.
DORYXYLON, Zoll. in Tijdschr. Nederl. Ind. xiv. (1857) 172, nomen prius.
MERCADOA, Naves, in Blanco, Fl. Philipp. ed. III. t. 463 (1880?).
macrophylla, *Muell. Arg. in Flora*, xlvii. (1864) 482.—Ind. or.
rottleroides, *Baill. Etud. Gén. Euph.* 390.—Malaya.

SUMBULUS, H. Reinsch, in Jahrb. Pract. Pharm. xiii. (1846) 69 = **Ferula**, Tourn. (Umbell.).
moschatus, H. Reinsch, l. c. = F. Sumbul.

SUNANIA, Rafin. Fl. Tellur. iii. 95 (1836) = **Polygonum**, Tourn.
filiformis, Rafin. l. c. = P. virginianum.

SUNAPTEA, Griff. Notul. iv. 516 (1854) = **Vatica**, Linn. (Dipteroc.).
bantamensis, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65 = V. grandiflora.
disticha, Trimen, Syst. Cat. Pl. Ceyl. 9 = V. disticha.
grandiflora, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65 = V. grandiflora.
odorata, Griff. Notul. iv. 516 = V. grandiflora.
scabriuscula, Trimen, Syst. Cat. Pl. Ceyl. 9 = V. scabriuscula.

SUNIPIA, Buch.-Ham. ex Sm. in Rees, Cycl. xxxiv. Art. Stelis, nn. 11, 13 [1816]; Lindl. Orch. Scel. 14 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 504.
bicolor, Lindl. Gen. et Sp. Orch. 179 (= *Ione bicolor*).—Reg. Himal.
scariosa, *Lindl. in Wall. Cat.* n. 7373; *et l. c.*—Reg. Himal.; Burma.

SUPRAGO, Gaertn. Fruct. ii. 402. t. 166 (1791) = **Vernonia**, Schreb. (Compos.).
glauca, Gaertn. l. c. = V. noveboracensis.
sphaerocephala, Cass. in Dict. Sc. Nat. li. 386 = Liatris scariosa.
spicata, Gaertn. Fruct. ii. 402. t. 167 = Liatris spicata.

SUREGADA, [Roxb.] Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 206 = **Gelonium**, Roxb. (Euphorb.).
adenophora, Baill. Adansonia, i. (1860–61) 253 = G. adenophorum.
angustifolia, Baill. Etud. Gén. Euph. 396 = G. lanceolatum.
bifaria, Baill. Hist. Pl. v. 120 = G. bifarium.
bilocularis, Roxb. ex Wall. Cat. n. 7981 = G. multiflorum.
Boiviniana, Baill. Adansonia, i. (1860–61) 252 = G. Boivinianum.
ceratophora, Baill. l. c. iii. (1862–63) 154 = G. africanum.
crenulata, Baill. l. c. i. (1860–61) 252 = G. adenophorum.
glabra, Roxb. Fl. Ind. iii. 831, in obs. = G. multiflorum.
glomerulata, Baill. Etud. Gén. Euph. 396 = G. bifarium.
laurina, Baill. Adansonia, i. (1860–61) 253 = G. laurinum.
multiflora, Baill. Etud. Gén. Euph. 396 = G. multiflorum.
spicata, Baill. l. c. = G. spicatum.
zanzibariensis, Baill. Adansonia, i. (1860–61) 254 = G. zanzibariense.

SURIANA, Domb. & Cav. ex D. Don, in Edinb. N. Phil. Journ. xiii. (Apr.-Oct. 1832) 238; Meissn. Gen. Comm. 235 (1841) = **Ercilla**, A. Juss. (Phytol.).
volubilis, Domb. & Cav. l. c. = E. volubilis.

SURIANA, Plum. ex Linn. Gen. ed. I. 340 (1737). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
maritima, *Linn. Sp. Pl.* 284.—Littora trop.

SURUBEA, Jaume St. Hil. Expos. ii. 355 (1805) = SOUROUBEA, Aubl. = **Ruyschia**, Jacq. (Ternstr.).
Aubletii, G. F. W. Mey. Prim. Fl. Esseq. 120 = R. Souroubea.

SURWALA, M. Roem. Syn. Hesper. 108 (1846) = **Walsura**, Roxb. (Meliac.).
robusta, M. Roem. l. c. = W. robusta.

SUSARIUM, Phil. in Linnaea, xxxiii. (1864–65) 248 = **Symphyostemon**, Miers (Irid.).
andinum, Phil. l. c. 250 = Chamelum andinum.
nigricans, Phil. l. c. 249.—Chili.
Segethi, Phil. l. c.—Chili.

SUSIANA, Linn. Syst. ed. I (1735) = **Fritillaria**, Tourn. (Liliac.).

SUSSEA, Gaudich. Bot. Voy. Bonite, tt. 24, 25, 38 (1844–57?) = **Pandanus**, Rumph.
conoidea, Gaudich. l. c. t. 24 = P. montanus.
lagenaeformis, Gaudich. l. c. t. 25 = P. lagenaeformis.
microcarpa, Gaudich. l. c. = P. Sussea.
microstigma, Gaudich. l. c. t. 38 = P. microstigma.

SUSSODIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 104 (1825) = **Colebrookia**, Sm. (Labiat.).
oppositifolia, Buch.-Ham. l. c. = C. oppositifolia.

SUSUM, Blume, in Schult. f. Syst. vii. p. xcv. 1493 (1830). *FLAGELLARIEAE*, Benth. & Hook. f. iii. 861.
HANGUANA, Blume, Enum. Pl. Jav. 15 (1827).
VERATRONIA, Miq. Fl. Ind. Bat. 553 (1859).
anthelminthicum, *Blume, in Schult. f. Syst.* vii. 1493.—Java.
minus, *Miq. Fl. Ind. Bat. Suppl.* 598.—Sumatra.

SUTERA, Hort. ex Steud. Nom. ed. I. 820 (1821) = **Lessertia**, DC. (Legumin.).
pubescens, Hort. ex Steud. l. c. = L. perennans.

SUTERA, Roth, Bot. Bemerk. 172 (1807) = **Chaenostoma**, Benth. (Scrophul.).
brachiata, Roth, l. c. 173.—Hab.?
foetida, Roth, l. c. 172 = C. foetidum.
oppositifolia, Roth, ex Steud. Nom. ed. II. ii. 652 = C. hispidum.

SUTERA, Roth, Nov. Pl. Sp. 291 (1821). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 945.
dissecta, Walp. Rep. iii. 271 = glandulosa.
glandulosa, *Roth, Nov. Pl. Sp.* 291.—Afr. bor.; Arab.; Ind. or.
multifida, Walp. Rep. iii. 271 = Conobea multifida.
serrata, Hochst. in Flora, xxiv. (1841) 1. Intell. 43 = Stemodia serrata.

SUTERIA, DC. Prod. iv. 536 (1830) = **Pyschotria**, Linn. (Rubiac.).
brasiliensis, Mart. in Flora, xxiv. (1841) II. Beibl. 71 = P. brasiliensis.
calycina, DC. Prod. iv. 536 = P. Lindleyana.
fluminensis, Mart. in Flora, xxiv. (1841) II. Beibl. 71 = P. fluminensis.
Hookeriana, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 109 = P. Gardneriana.
macrantha, Gardn. l. c. 110 = P. macrantha.
macrocalyx, Mart. in Flora, xxiv. (1841) II. Beibl. 71 = P. megalocalyx.
nuda, Mart. l. c. = P. Gardneriana.
parviflora, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 110 = P. suterella.

SUTHERLANDIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 327 (1812). *LEGUMINOSAE*, Benth. & Hook. f. i. 503.
Darumbium, *Bert. ex Colla, in Mem. Acc Torin.* xxxvii. (1834) 55.—Chili.

SUTHERLANDIA:—
floribunda, *Carr. in Rev. Hortic.* (1870-71) 611.—Hab.?
foliolata, *Lange, Ind. Sem. Hort. Haun.* (1859) 29; *et in Linnaea*, xxx. (1859–60) 757.—Austral. ?
frutescens, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 327.—Afr. austr.
incana, E. Mey. Comm. Pl. Afr. Austr. 121 = frutescens.
microphylla, Burch. ex DC. Prod. ii. 273 = frutescens.
tomentosa, Eckl. & Zeyh. Enum. 251 = frutescens.
tomentosa, Spreng. Syst. iii. 242 = Lessertia tomentosa.
vesicaria, Spreng. l. c. = Lessertia vesicaria.

SUTHERLANDIA, J. F. Gmel. Syst. 1027 (1791) = **Heritiera**, Dryand. (Stercul.).
littoralis, J. F. Gmel. l. c. = H. littoralis.

SUTRINA, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. x. (1842) 184. *ORCHIDEAE*, Benth. & Hook. f. iii. 567.
bicolor, *Lindl. l. c.* 185.—Peruv.

SUTTONIA, A. Rich. Ess. Fl. N. Zél. 349 (1832) = **Myrsine**, Linn.
australis, A. Rich. l. c. 349. t. 38 = M. Urvillei.
divaricata, Hook. f. Fl. Antarct. 51. t. 34 = M. divaricata.
montana, Hook. f. Fl. N. Zel. ii. 334 = M. montana.
nummularia, Hook. f. l. c. i. 173 = M. nummularia.
salicina, Hook. f. Fl. Antarct. 52; et l. c. 172 = M. salicina.
saligna, Walp. Rep. vi. 450, sphalm. = praec.
tenuifolia, Hook. f. Fl. Antarct. 52.—Ins. Norfolk.

SUZYGIUM, P. Br. Hist. Jamaic. 240 (1756); Adans. Fam. ii. 244 (1763) = **Calyptranthes**, Sw. (Myrtac.).

SVIDA, Opiz, Seznam, 94 (1852) = **Cornus**, Tourn.
alba, Opiz, l. c. = C. alba.
sanguinea, Opiz, l. c. = C. sanguinea.

SVITRAMIA, Cham. in Linnaea, ix. (1834) 445. *MELASTOMACEAE*, Benth. & Hook. f. i. 786.
canescens, Walp. Rep. ii. 132 = pulchra.
pulchra, *Cham. in Linnaea*, ix. (1834) 446.—Bras.

SWAINSONA, Salisb. Parad. Lond. t. 28 (1806) *LEGUMINOSAE*, Benth. & Hook. f. i. 504.
CYCLOGYNE, Benth. in Lindl. Swan River App. 16 (1839).
DIPLOLOBIUM, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 489.
albiflora, G. Don, Gen. Syst. ii. 245 = coronillaefolia.
astragalifolia, Hort. ex G. Don, in Loud. Hort. Brit. 305 = lessertiaefolia.
brachycarpa, *Benth. Fl. Austral.* ii. 217.—Austral.
Burkei, *F. Muell. ex Benth. l. c.* 218.—Austral.
Burkittii, *F. Muell. ex Benth. l. c.*—Austral.
campylantha, *F. Muell. in Rep. Greg. Pl.* 6.—Austral.
canescens, *F. Muell. Fragm.* iii. 46.—Austral.
colutoides, *F. Muell. l. c.* x. 6.—Austral.
coronillaefolia, *Salisb. Parad. Lond.* t. 28.—Austral.
Drummondii, *Benth. Fl. Austral.* ii. 220.—Austral.
Fraseri, *Benth. l. c.* 224.—Austral.
Froebelii, Regel, Gartenfl. iii. (1854) 178 = lessertiaefolia.
galegifolia, R. Br. in Ait. Hort. Kew. ed. II. iv. 327 = coronillaefolia.
gracilis, *Benth. Fl. Austral.* ii. 220.—Austral.
grandiflora, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 74 = Greyana.
Greyana, *Lindl. Bot. Reg.* (1846) t. 66.—Austral.
laxa, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 76.—Austral.
lessertiaefolia, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 99.—Austral.
luteola, *F. Muell. Fragm.* i. 75.—Austral.
Maccullochiana, *F. Muell. l. c.* vii. 25.—Austral.
magnifica, Hort. ex Dombr. Flor. Mag. v. (1866) t. 273 = coronillaefolia ?
microphylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 410.—Austral.
monticola, *Benth. ex A. Gray, l. c.* 411.—Austral.
Murrayana, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 69.—Austral.

SWAINSONA:—
Novae-Zelandiae, *Hook. f. Handb. N. Zeal. Fl.* 51.—N. Zel.
occidentalis, *F. Muell. Fragm.* iii. 46.—Austral.
oligophylla, *F. Muell. ex Benth. Fl. Austral.* ii. 219.—Austral.
Oliverii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 152.—Austral.
oncinotropis, *F. Muell. in Melb. Chemist* (Oct. 1884).—Austral.
oroboides, *F. Muell. ex Benth. Fl. Austral.* ii. 222.—Austral.
Osbornii, T. Moore, in Gardn. Comp. (1852) t. 65 = coronillaefolia.
parviflora, *Benth. Fl. Austral.* ii. 223.—Austral.
phacifolia, *F. Muell. in S. Austral. Register* (1850); *ex Benth. Fl. Austral.* ii 221.—Austral.
phacoides, *Benth. in Mitch. Journ. Trop. Austral.* 363.—Austral.
plagiotropis, *F. Muell. Fragm.* ix. 153.—Austral.
procumbens, *F. Muell. l. c.* iii. 46.—Austral.
stenodonta, *F. Muell. l. c.* xi. 70.—Austral.
stipularis, F. Muell. in Linnaea, xxv. (1852) 393 = phacifolia.
tephrotricha, F. Muell. l. c. 392 = lessertiaefolia.
unifoliolata, *F. Muell. Fragm.* viii. 226.—Austral.
viciaefolia, F. Muell. in D. N. F. Dietr. Fl. Univ. N. S. t. 17 = lessertiaefolia.
violacea, Henders. Illustr. Bouq. t. 19 = procumbens.

SWAINSONIA, Spreng. Anleit. ii. II. 754 (1818) = praec.

SWAMMERDAMIA, DC. Prod. vi. 164 (1837) = **Helichrysum**, Vaill. (Compos.).
Antennaria, DC. l. c. = H. antennarium.
glomerata, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 120 (= *Ozothamnus glomeratus*).—N. Zel.
volubilis, Regel, Cat. Pl. Hort. Aksakov. 136, nomen.—Austral.

SWANALLOIA, Hort. ex Walp. Rep. iii. 934 (1844–45), sphalm. = **Juanulloa**, Ruiz & Pav. (Solan.).
aurantiaca, Hort. ex Walp. l. c. = J. parasitica.

SWANTIA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 358, 365 = **Vicia**, Linn. (Legumin.),
aurantia, Alef. l. c. 366 = V. aurantia.

SWARTZIA, J. F. Gmel. Syst. 360 (1791) = **Solandra**, Sw. (Solan.).
grandiflora, J. F. Gmel. l. c. = Sol. grandiflora.

SWARTZIA, Schreb. Gen. 518 (1791). *LEGUMINOSAE*, Benth. & Hook. f. i. 561.
DITHYRIA, Benth. in Hook. Journ. Bot. ii. (1840) 89.
GYNANTHISTROPHE, Poit. ex DC. Prod. ii. 424 (1825).
HOELZELIA, Neck. Elem. iii. 62 (1790).
POSSIRA, Aubl. Pl. Gui. ii. 934. t. 355 (1775).
RITTERA, Schreb. Gen. 364 (1789).
RIVERIA, H. B. & K. Nov. Gen. et Sp. vii. 266, t. 659 (1825).
ROSSINA, Steud. Nom. ed. II. ii. 652 (1841), sphalm.
TOUNATEA, Aubl. Pl. Gui. i. 549. t. 218 (1775), nomen prius.
TRISCHIDIUM, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 141. t. 4.
acuminata, *Willd. ex Vog. in Linnaea*, xi. (1837) 173.—Bras.
acutifolia, *Vog. l. c.* 174.—Bras.
alata, *Willd. Sp. Pl.* ii. 1220.—Guian.
alterna, *Benth. in Hook. Journ. Bot.* ii. (1840) 89.—Bras.
amazonica, *S. Moore, in Journ. Bot.* xvi. (1878) 132.—Bras.
apetala, *Raddi, in Mem. Moden.* xviii. *Fis.* (1820) 398.—Bras.
aptera, *DC. Mém. Légum.* 405.—Am. trop.
argentea, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 31.—Bras.
auriculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 61.—Bras.
Benthamiana, *Miq. Stirp. Surin. Sel.* 15.—Am. trop.
bifida, Steud. in Flora, xxvi. (1843) 757 = triphylla.
Blanchetii, *Benth. in Mart. Fl. Bras.* xv. II. 29.—Bras.

SWARTZIA:—
brachystachya, *DC. Mém. Légum.* 408.—Guian.
bracteosa, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 20.—Bras.
brasiliensis, Vog. in Linnaea, xi. (1837) 169 = Langsdorffii.
callistemon, Spruce, ex Benth. in Mart. Fl. Bras. xv. II. 36 = acuminata.
calophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 61. t. 267.—Peruv.
capparioides, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1104, nomen.—Guian.
cardiosperma, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 33.—Bras.
caribaea, *Griseb. Fl. Brit. W. Ind.* 212.—Ind. occ.
conferta, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 20.—Bras.
coriacea, Desv. in Ann. Sc. Nat. Sér I. ix. (1826) 424 = Dipteryx nudipes.
coriacea, Steud. Nom. ed. II. ii. 652, sphalm. = apetala.
corrugata, *Benth. in Hook. Journ. Bot.* ii. (1840) 87.—Am. trop.
crocea, *Benth. in Mart. Fl. Bras.* xv. II. 23.—Bras.
cuspidata, *Spruce, ex Benth. l. c.* 36.—Bras.
dicarpa, *Moric. ex Meissn. Gen. Comm.* 68.—Am. trop.
dipetala, Willd. ex Vog. in Linnaea, xi. (1837) 173 = dicarpa.
discolor, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 62.—Bras.
dodecandra, *Willd. Sp. Pl.* ii. 1220.—Am. austr.
elegans, *Schott, in Spreng. Syst.* iv. *App.* 407.—Bras.
eriocarpa, *Benth. in Mart. Fl. Bras.* xv. II. 17.—Guian.
erythrocarpa, Spruce, ex Benth. l. c. 28 = sericea.
Flemmingi, *Raddi, in Mem. Moden.* xviii. *Fis.* (1820) 397 (Flaemingi).—Bras.
floribunda, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 21.—Bras.
fugax, *Spruce, ex Benth. l. c.* 30.—Bras.
glabra, Vog. in Linnaea, xi. (1837) 175 = apetala.
grandiflora, Vog. l. c. 169, partim = crocea.
grandiflora, *Willd. Sp. Pl.* ii. 1220.—Am. trop.
grandifolia, *Bong. ex Benth. in Hook. Journ. Bot.* ii. (1840) 85.—Am. trop.
Hostmanni, *Benth. in Mart. Fl. Bras.* xv. II. 18.—Guian.
Langsdorffii, *Raddi, in Mem. Moden.* xviii. *Fis.* (1820) 396.—Bras.
latifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 86.—Guian.
laurifolia, *Benth. l. c.* 87.—Bras.
laxiflora, *Bong. ex Benth. l. c.* 86.—Bras.
leiocalycina, *Benth. in Mart. Fl. Bras.* xv. II. 28.—Guian.
leptopetala, *Benth. in Hook. Journ. Bot.* ii. (1840) 87.—Bras.
lomatopus, Mart. in Flora, xx. (1837) II. Beibl. 103 = multijuga, *Vog.*
longifolia, *DC. Mém. Légum.* 406.—Guian.
macrocarpa, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 38.—Bras.
macrophylla, *Willd. ex Vog. in Linnaea*, xi. (1837) 172.—Bras.
macrosperma, *Bertol. Misc. Bot.* xxi. 16.—Guatemala.
macrostachya, *Benth. in Mart. Fl. Bras.* xv. II. 24.—Bras.
madagascariensis, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 424.—Madag.
marginata, Benth. in Hook. Journ. Bot. ii. (1840) 87 = madagascariensis.
Martii, *Eichl. in Mart. Fl. Bras.* xv. II. 19.—Bras.
Mathewsii, *Benth. in Hook. Ic. Pl.* t. 1064 (Matthewsii).—Peruv.; Bolivia.
microcarpa, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 35.—Bras.
micropetala, Schott, in Spreng. Syst. iv. App. 407 = myrtifolia.
microstylis, Benth. in Hook. Journ. Bot. ii. (1840) 89 = dicarpa.
mollis, *Benth. l. c.*—Bras.
montana, Vog. in Linnaea, xi. (1837) 174 = Flemmingi.
multijuga, *A. Rich. Ess. Fl. Cub.* 457.—Cuba.
multijuga, *Vog. in Linnaea*, xi. (1837) 175.—Bras.
myrtifolia, *Sm. in Rees, Cycl.* xxxiv. n. 5.—Am. trop.
nitens, *Vog. ex Steud. Nom.* ed. II. ii. 653.—N. Granat.

SWARTZIA:—
oblonga, *Benth. in Hook. Kew Journ.* ii. (1850) 238.—Guian.
ochnacea, DC. Mém. Légum. 405. t. 58 = grandiflora.
panamensis, *Benth. in Mart. Fl. Bras.* xv. II. 38.—Am. centr.
parviflora, DC. Mém. Légum. 403. t. 60 = triphylla.
pendula, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 19.—Bras.
picta, *Spruce, ex Benth. l. c.* 25.—Bras.
pilulifera, *Benth. in Hook. Journ. Bot.* ii. (1840) 90.—Bras.
pinnata, Seem. Bot. Voy. Herald, 113 = panamensis.
pinnata, Willd. ex Vog. in Linnaea, xi. (1837) 173 = acuminata.
pinnata, *Willd. Sp. Pl.* ii. 1220.—Guian.
polyanthera, Steud. in Flora, xxvi. (1843) 757 = tomentosa.
polyphylla, *DC. Mém. Légum.* 411.—Guian.
pulcherrima, Allem. ex Benth. in Mart. Fl. Bras. xv. II. 24 = Langsdorffii.
pulchra, Vog. in Linnaea, xi. (1837) 169 = elegans.
racemosa, *Benth. in Hook. Kew Journ.* ii. (1850) 238.—Bras.
recurva, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 61.—Bras.
robiniaefolia, *Willd. ex Vog. in Linnaea*, xi. (1837) 171.—N. Granat.
rosea, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 32.—Bras.
Schomburgkii, *Benth. l. c.* 38.—Guian.
sericea, *Vog. in Linnaea*, xi. (1837) 176.—Bras.
simplex, Spreng. Syst. ii. 567 = grandiflora.
simplicifolia, Willd. Sp. Pl. ii. 1219 = grandiflora.
Sprucei, *Benth. in Mart. Fl. Bras.* xv. II. 37.—Bras.
stipellata, Benth. l. c. = Sprucei.
stipularis, Spruce, ex Benth. l. c. 38 = macrocarpa.
subspicata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1208, nomen.—Guian.
surinamensis, *Klotzsch, l. c.*, nomen.—Guian.
tomentosa, *DC. Mém. Légum.* 409. t. 59.—Guian.
Trianae, *Benth. in Mart. Fl. Bras.* xv. II. 39.—Am. trop.
triphylla, DC. Prod. ii. 423, partim = myrtifolia.
triphylla, *Willd. Sp. Pl.* 1220.—Am. trop.
velutina, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 21.—Bras.

SWEERTIA, Koch, Syn. Fl. Germ. ed. II. 558 (1844) = **Swertia**, Linn. (Gentian.).

SWEETIA, DC. Prod. ii. 381 (1825) = **Galactia**, P. Br. (Legumin.).
filiformis, DC. l. c. 382 = G. filiformis.
lignosa, DC. l. c.—Ins. S. Doming.
longifolia, DC. l. c. 381 = G. Jussiaeana.
uliginosa, G. Don, Gen. Syst. ii. 342 = Derris uliginosa.
velutina, G. Don, l. c. = Tephrosia velutina.

SWEETIA, Spreng. Syst. ii. 171, 213 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 559.
ACOSMIUM, Schott, in Spreng. Syst. iv. Cur. Post. 406 (1827).
LEPTOLOBIUM, Vog. in Linnaea, xi. 388 (1837).
bijuga, *Benth. in Journ. Linn. Soc.* viii. (1865) 261.—Bras.
brachystachya, *Benth. l. c.* 262.—Bras.
dasycarpa, *Benth. l. c.*—Bras.
elegans, *Benth. l. c.*—Bras.
fruticosa, *Spreng. Syst.* ii. 213.—Bras.
Gardneri, Benth. in Journ. Linn. Soc. viii. (1865) 263 = Rourea Gardneriana.
glabrifolia, Benth. l. c. 262 = dasycarpa.
lentiscifolia, *Spreng. Syst.* iv. *Cur. Post.* 406.—Bras.
nitens, *Benth. in Journ. Linn. Soc.* viii. (1865) 262.—Bras.
panamensis, *Benth. l. c.* 263.—Panama.
tenuifolia, *Benth. in Mart. Fl. Bras.* xv. II. 7.—Bras.

SWERTIA, All. Fl. Pedem. i. 208 (1785) = **Tolpis**, Linn. (Compos.).
barbata, All. l. c. = T. barbata.

SWERTIA, Linn. Sp. Pl. ed. I. 226 (1753). *GENTIANEAE*, Benth. & Hook. f. ii. 816.
AGATHOTES, D. Don, in Trans. Linn. Soc. xvii. (1837) 522.
ANAGALLIDIUM, Griseb. Gen. et Sp. Gent. 311 (1836).

SWERTIA:—
BLEPHARADEN, Dulac, Fl. Hautes-Pyr. 449 (1867).
HENRICEA, Lem. Lis. in Bull. Soc. Philom. (1824) 171.
MONOBOTHRIUM, Hochst. in Flora, xxvii. (1844) 27.
OPHELIA, D. Don, ex G. Don, Gen. Syst. iv. 178 (1837); et in Trans. Linn. Soc. xvii. (1837) 524.
RELLESTA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 337.
SCZUKINIA, Turcz. l. c. (1840) 165.
STELLERA, Turcz. l. c. 167 (1840).
abyssinica, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 24, nomen; *et ex Griseb. in DC. Prod.* ix. 134.—Abyss.
affinis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 126.—Ind. or.
alata, *Royle, ex D. Don, in Trans. Linn. Soc.* xvii. (1837) 523.—Reg. Himal.
albicaulis, Dougl. ex Griseb. Gen. et Sp. Gent. 330 = Frasera albicaulis.
alpestris, *Baumg. Mant.* (1846) 19.—Transsylv.
alternifolia, *Royle, Illustr. Bot. Himal.* 278. t. 67. f. 2.—Reg. Himal.
americana, Spreng. Syst. i. 861 = Halenia deflexa.
angustifolia, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 127.—Reg. Himal.
asclepiadea, H. B. & K. Nov. Gen. et Sp. iii. 175 = Halenia Pavoniana.
Aucheri, *Boiss. Diagn.* Ser. I. v. 90.—Armen.; Persia.
Balansae, *Boiss. Fl. Orient.* iv. 78.—Pontus.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 127.—Ind. or.
bimaculata, *Hook. f. & Thoms. ex C. B. Clarke, in Journ. Linn. Soc.* xiv. (1875) 449.—Reg. Himal.
brevicornis, H. B. & K. Nov. Gen. et Sp. iii. 174 = Halenia brevicornis.
brevicornis, Willd. ex Schult. Syst. vi. 132 = Halenia Pavoniana.
caerulea, Royle, Illustr. Bot. Himal. 464, (Index) = coerulea.
carinthiaca, Wulf. in Jacq. Misc. ii. 53 = Pleurogyna carinthiaca.
centrostemma, Wall. Cat. n. 4385 = Halenia elliptica.
chinensis, Franch. in Bull. Soc. Bot. Fr. xxxii. (1885) 26 = diluta.
Chirata, *Buch.-Ham. ex Wall. Cat.* n. 4372.—Reg. Himal.
ciliata, Royle, ex D. Don, in Trans. Linn. Soc. xvii. (1837) 526 = purpurascens.
clarenceana, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 16.—Afr. trop.
coerulea, *Royle, Illustr. Bot. Himal.* 278. t. 67. f. 1.—Reg. Himal.
connata, *Schrenk, Enum. Pl. Nov.* i. 37.—Sibir. altaic.
cordata, *Wall. Cat.* n. 4378.—Reg. Himal.
corniculata, Linn. Sp. Pl. 227 = Halenia sibirica.
corniculata, Michx. Fl. Bor. Am. i. 97 = Halenia deflexa.
corymbosa, *Wight, ex Griseb. Gen. et Sp. Gent.* 317; *Field. & Gardn. Sert. Pl.* t. 71.—Ind. or.
cuneata, *Wall. Cat.* n. 4380.—Reg. Himal.
decumbens, *Vahl, Symb. Bot.* i. 24.—Arab.
decussata, *Nimmo, in J. Grah. Cat. Pl. Bomb.* 249, nomen; *et ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 127.—Ind. or.
deflexa, Sm. in Rees, Cyclop. xxxiv. n. 8 = Halenia deflexa.
dichotoma, *Linn. Sp. Pl.* 227.—Sibir.
difformis, Linn. Herb. ex A. Gray, Syn. Fl. N. Am. ii. I. 125 = Frasera caroliniensis.
difformis, Linn. Sp. Pl. 226 = Sabbatia Elliotii.
dilatata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 122.—Reg. Himal.
diluta, *Benth. & Hook. f. Gen.* ii. 817.—As. or.
elegans, Wall. Cat. n. 4376 = pulchella.
erythrosticta, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 503.—China.
fastigiata, Pursh, Fl. Am. Sept. i. 101 = Frasera thyrsiflora.
filiformis, *Macfad. in Hook. Bot. Misc.* ii. (1831) 116, nomen.—Jamaic.
florida, Wall. Cat. n. 4382 = pulchella.
gracilis, H. B. & K. Nov. Gen. et Sp. iii. 176 = Halenia gracilis.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 122.—Reg. Himal.
Hookeri, *C. B. Clarke, l. c.* 127.—Reg. Himal.

SWERTIA:—
Hugelii, Griseb. in DC. Prod. ix. 133 = cuneata.
hypericoides, H. B. & K. Nov. Gen. et Sp. iii. 176 = Halenia hypericoides.
iberica, Fisch. ex Boiss. Fl. Orient. iv. 78 = punctata.
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 149 = alpestris.
intermixta, *A. Rich. Tent. Fl. Abyss.* ii. 57.—Abyss.
javanica, *Blume, Bijdr.* 848.—Java.
kamoonensis, Steud. Nom. ed. II. ii. 653, sphalm. loci nomen.—Cf. Wall. Cat. sub nn. 4379, 4380.
Kingii, *Hook. f. Ic. Pl.* t. 1442.—Reg. Himal.
lactea, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 393.—Persia; Turkest.
lahulensis, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 100 = petiolata.
longifolia, *Boiss. Diagn.* Ser. I. v. 90.—Persia.
lurida, *Royle, ex D. Don, in Trans. Linn. Soc.* xvii. (1837) 528.—Reg. Himal.
lutea, Vest, ex Reichb. Fl. Germ. Excurs. 429 = Gentiana lutea.
macrosperma, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 123.—Reg. Himal.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 206.—Afr. trop.
marginata, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 353.—Soongar.
Michauxiana, Schlecht. & Cham. in Linnaea, v. (1830) 122 = Halenia Schiedeana.
Michauxiana, Schult. Syst. vi. 130 = Halenia deflexa.
multicaulis, *D. Don, Prod. Fl. Nep.* 127.—Reg. Himal.
nervosa, *Wall. Cat.* n. 4383.—Reg. Himal.
obtusa, C. Koch, in Linnaea, xxiii. (1850) 586 = Aucheri.
obtusa, Ledeb. in Mém. Acad. Pétersb. v. (1812) 526 = perennis.
Pallasii, G. Don, Gen. Syst. iv. 176 = tetrapetala.
paniculata, *Wall. Cat.* n. 4374; *et Pl. As. Rar.* iii. 3. t. 205.—Reg. Himal.
parnassifolia, Labill. Nov. Holl. Pl. i. 72. t. 97 = Villarsia parnassifolia.
parviflora, H. B. & K. Nov. Gen. et Sp. iii. 174 = Halenia parviflora.
pauciflora, Humb. & Bonpl. ex Schult. Syst. vi. 135 = Halenia plantaginea.
peloria, Griff. Itin. Notes, 197 = Halenia elliptica.
perennis, Bieb. ex Boiss. Fl. Orient. iv. 78 = punctata.
perennis, *Linn. Sp. Pl.* 226.—Reg. bor. temp.
perfoliata, Royle, ex G. Don, Gen. Syst. iv. 176 = petiolata.
persica, Griseb. in DC. Prod. ix. 132 = Aucheri.
petiolata, *Royle, Illustr. Bot. Himal.* 277, nomen; *et ex D. Don, in Trans. Linn. Soc.* xvii. (1837) 519.—Reg. Himal.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 55.—Abyss.
pinifolia, Ruiz & Pav. ex G. Don, Gen. Syst. iv. 177 = Halenia pinifolia.
plantaginea, H. B. & K. Nov. Gen. et Sp. iii. 175 = Halenia plantaginea.
pulchella, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 127; *et ex Wall. Cat.* n. 4375.—Reg. Himal.
pumila, *Hochst. ex Hook. f. in Journ. Linn. Soc.* vii. (1864) 206.—Abyss.
punctata, *Baumg. Enum. Stirp. Transs.* i. 190.—Europ. or.; Reg. Caucas.
purpurascens, *Wall. Cat.* n. 4379.—Reg. Himal.
purpurea, Royle, Illustr. Bot. Himal. 277 = praec.?
pusilla, Pursh, Fl. Am. Sept. i. 101 = Pleurogyna carinthiaca.
quadricornis, Willd. ex Schult. Syst. vi. 134 = Halenia Pavoniana.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 56.—Abyss.
racemosa, *Wall. Cat.* n. 4377.—Reg. Himal.
recurva, Sm. in Rees, Cyclop. xxxiv. n. 9 = Halenia gracilis.
Rex, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 127.—Reg. Himal.
rotata, Linn. Sp. Pl. 226 = Pleurogyna rotata.
rotata, Pall. Fl. Ross. i. II. 98. t. 89. f. 3 = Pleurogyna carinthiaca.
rotata, *Thunb. Fl. Jap.* 115.—Japon.
Schimperi, *Griseb. in DC. Prod.* ix. 563.—Abyss.
speciosa, G. Don, Gen. Syst. iv. 176 = petiolata.
speciosa, *Wall. Cat.* n. 4384.—Reg. Himal.
stigmantha, C. Koch, in Linnaea, xxiii. (1850) 586 = punctata.
sulcata, Rottb. in Skrift. Kioeb. Selsk. x. (1770) 438. t. 1. f. 4 = Pleurogyna rotata.

SWERTIA:—
tetragona, *Edgew. in Trans. Linn. Soc.* xx. (1846) 86.—Reg. Himal.
tetrandra, *Hochst. in Flora*, xxvii. (1844) 28.—Abyss.
tetrapetala, *Pall. Fl. Ross.* i. II. 99.—Kamtschat.
tetraptera, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 503.—China.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 129.—Reg. Himal.
trichotoma, *Wall. Cat.* n. 4381.—Ind. or.
turfosa, Baumg. ex Steud. Nom. ed. II. ii. 653, nomen (Quid ?).—Transsylv.
umbellata, Mathews, ex Griseb. Gen. et Sp. Gent. 328 = Halenia gracilis.
umbellata, Ruiz & Pav. Fl. Per. iii. 21 = Halenia Pavoniana.
vacillans, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 504, in obs.—China.
zeylanica, *Walker, ex Griseb. Gen. et Sp. Gent.* 316, in syn.—Zeylan.

SWERTYA, Steud. Nom. ed. II. ii. 653 (1841), sphalm. = SWERTIA, All. = **Tolpis**, Linn. (Compos.).

SWIETENIA, Jacq. Enum. Pl. Carib. 4 (1760). *MELIACEAE*, Benth. & Hook. f. Gen. i. 338.
MAHAGONI, Adans. Fam. ii. 343 (1763).
ROIA, Scop. Introd. 226 (1777).
acutifolia, Stokes, Bot. Mat. Med. ii. 479 = S. Mahagoni.
alternifolia, Steud. Nom. ed. II. ii. 653 = Cedrela alternifolia.
angolensis, *Welw. Apont.* 587.—Afr. trop.
Chickrassia, Roxb. Hort. Beng. 33; Fl. Ind. ii. 399 = Chickrassia tabularis.
Chloroxylon, Roxb. Pl. Corom. i. 46. t. 64 = Chloroxylon Swietenia.
fabrilis, Salisb. Prod. 317 = S. Mahagoni.
febrifuga, Roxb. Pl. Corom. i. 18. t. 17 = Soymida febrifuga.
humilis, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 355. t. 7.—Mexic.
Mahagoni, *Jacq. Enum. Pl. Carib.* 20; *Linn. Sp. Pl.* ed. II. 271.—Am. austr.
Mahogani, DC. Prod. i. 625 = praec.
Mahogoni, Lam. Encyc. iii. 678 = praec.
multijuga, Schiede, in Linnaea, iv. (1829) 578 = humilis.
obtusifolia, Stokes, Bot. Mat. Med. ii. 480 = Soymida febrifuga.
rubra, Wight, ex Wall. Cat. n. 4890 = Soymida febrifuga.
senegalensis, Desv. in Lam. Encyc. iii. 679 = Khaya senegalensis.
Soymida, A. Dunc. Tent. 16 = Soymida febrifuga.
Sureni, Blume, Cat. Gew. Buitenz. 72 = Cedrela Toona.
Toon, Stokes, Bot. Mat. Med. ii. 480 = Cedrela Toona.
trilocularis, Roxb. ex Buch.-Ham. Journ. i. 184 = Chickrassia tabularis.
velutina, Wall. ex Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 65 = Chickrassia tabularis.
villosa, Wall. ex Kurz, l. c. = Chickrassia tabularis.

SWINTONIA, Griff. in Proc. Linn. Soc. i. (1846) 283. *ANACARDIACEAE*, Benth. & Hook. f. i. 421.
ANAUXANOPETALUM, Teijsm. & Binn. in Journ. Bot. Néerl. i. (1861) 368.
ASTROPETALUM, Griff. Notul. iv. 411. t. 565 (1854).
acuta, *Engl. Bot. Jahrb.* i. (1881) 44.—Borneo.
floribunda, *Griff. in Proc. Linn. Soc.* i. (1846) 283.—Burma.
glauca, *Engl. Bot. Jahrb.* i. (1881) 44.—Borneo.
Griffithii, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 75 = floribunda.
Helferi, *Hook. f. Fl. Brit. Ind.* ii. 26.—Burma.
obtusifolia, *Engl. in DC. Monog. Phan.* iv. 231.—Borneo.
Pierrei, *Hance, in Journ. Bot.* xiv. (1876) 257.—Cambodia.
Schwenckii, *Teijsm. & Binn. Cat. Hort. Bog.* 230, nomen; *et ex Hook. f. Fl. Brit. Ind.* ii. 26.—Malaya.
spicifera, *Hook. f. Fl. Brit. Ind.* ii. 27.—Malaya.

SYAGRUS, Mart. Hist. Nat. Palm. ii. 129. t. 89 (1824) = **Cocos**, Linn. (Palm.).
amara, Mart. Palmet. Orbign. 132 = C. amara.
botryophora, Mart. l. c. 133 = C. botryophora.
Chiragua, H. Wendl. in Kerch. Palm. 257.—N. Granat.

SYAGRUS:—
cocoides, Mart. Hist. Nat. Palm. ii. 130. tt. 89, 90 = Cocos Syagrus.
comosa, Mart. Palmet. Orbign. 134 = C. comosa.
macrocarpa, Rodrig. Prot.-App. Enum. Palm. Nov. 46, 48 = C. Procopiana.
majestica, Hort. ex H. Wendl. in Kerch. Palm. 257, nomen.—Hab. ?
Mikaniana, Mart. Palmet. Orbign. 133 = C. Mikaniana.
picrophylla, Rodrig. Prot.-App. Enum. Palm. Nov. 45, 48 = C. Martiana.
Sancona, Karst. in Linnaea, xxviii. (1856) 247.—N. Granat.
Tamaca, Linden, in Illustr. Hortic. xxviii. (1881) 32, nomen.—N. Granat.
Wallisi, Linden, Cat. n. 16 (1861); n. 18 (1863) 45.—N. Granat.

SYALITA, Adans. Fam. ii. 364 (1763) = **Dillenia**, Linn.

SYAMA, Jones, in As. Res. iv. (1795) 261 = **Pupalia**, Juss. (Amarant.).
lata, Pritz. Ic. Ind. i. 1082 (Quid ?).—Ind. or.

SYCAMORUS, Oliver, in Trans. Linn. Soc. xxix. (1874) 149, err. typ. = SYCOMORUS, Gasp. = **Ficus**, Tourn. (Urtic.).

SYCHINIUM, Desv. in Mém. Soc. Linn. Par. iv. (1826) 216. t. 12 = **Dorstenia**, Plum. (Urtic.).
ramosum, Desv. l. c. 217 = D. arifolia.

SYCHNOSEPALUM, Eichl. in Flora, xlvii. (1864) 395. *MENISPERMACEAE*, Benth. & Hook. f. i. 963.
DETANDRA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. 124 (1864); et xx. (1867) 17.
microphyllum, *Eichl. in Flora*, xlvii. (1864) 396.—Bras.
paraënse, *Eichl. l. c.*—Bras.
Sagotianum, *Eichl. l. c.*—Guian.

SYCIOS, Medic. Vorles. Churpf. Phys. Ges. iv. 1. (1789) 190, sphalm. = **Sicyos**, Linn. (Cucurb.).

SYCODIUM, Pomel, Nouv. Mat. Fl. Atl. 39 (1874) = **Anvillea**, DC. (Compos.).
radiatum, Pomel, l. c. = A. radiata.

SYCOMORPHE, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 35 = **Ficus**, Tourn. (Urtic.).
Roxburghii, Miq. l. c. = F. hispida.

SYCOMORUS, Gasp. Ricerch. Caprif. e Fic. 86 (1845) = **Ficus**, Tourn. (Urtic.).
antiquorum, Gasp. l. c. = F. Sycomorus.
capensis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 113. t. 3. f. B = F. capensis.
capreaefolia, Walp. Ann. iii. 418 = F. antithetophylla.
Dahro, Walp. l. c. = F. Dahro.
gnaphalocarpus, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 113 = F. gnaphalocarpa.
guineensis, Miq. l. c. 112. t. 14 = F. capensis.
gummiflua, Miq. Pl. Jungh. 64 = F. variegata.
hirsuta, Sond. in Linnaea, xxiii. (1850) 137 = F. Sonderi.
panifica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 111. t. 3. f. A = F. panifica.
rigida, Miq. l. c. 110.—Afr. trop.
riparia, Miq. l. c. 114 = F. riparia.
Schimperiana, Miq. l. c. 112 = F. Vallis-Choudae.
Thonningiana, Miq. l. c. = F. Thonningiana.
trachyphyllus, Miq. l. c. 110 = F. trachyphylla.
Vogeliana, Miq. l. c. 112 = F. Vogeliana.

SYCOPSIS, Oliver, in Trans. Linn. Soc. xxiii. (1860) 83. t. 8. *HAMAMELIDEAE*, Benth. & Hook. f. i. 666.
Griffithiana, *Oliver, l. c.*—Reg. Himal.

SYDERITIS, All. in Misc. Taur. v. (1773) 59, sphalm. = **Sideritis**, Linn. (Labiat.).

SYENA, Schreb. Gen. 39 (1789) = **Mayaca**, Aubl.
Aubletii, Schott & Endl. Melet. 24 = M. Aubletii.
fluviatilis, Willd. Sp. Pl. i. 254 = M. fluviatilis.
Mayaca, J. F. Gmel. Syst. 121 = M. fluviatilis.
Nuttalliana, Schult. Mant. i. 343.—Am. bor.

SYKESIA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 351 = **Gaertnera**, Lam. (Logan.).
Koenigii, Arn. l. c. 353 = G. Koenigii.
thyrsiflora, Arn. l. c. = G. Koenigii.
Walkeri, Arn. l. c. 354 = G. Walkeri.

SYKORAEA, Opiz, Seznam, 94 (1852) = **Campanula**, Tourn.
barbata, Opiz, l. c. = C. barbata.
hortensis, Opiz, l. c. = C. Medium.

SYLITRA, E. Mey. Comm. Pl. Afr. Austr. 114 (1835). *LEGUMINOSAE*, Benth. & Hook. f. i. 496.
angolensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 103.—Afr. trop.
biflora, *E. Mey. Comm. Pl. Afr. Austr.* 114.—Afr. austr.

SYLLEPIS, Fourn. ex Benth. & Hook. f. Gen. iii. 1125 (1883) = **Imperata**, Cyrilli (Gramin.).
polystachya, Fourn. ex Hack. in Mart. Fl. Bras. ii. III. 251 = I. exaltata.
Ruprechtii, Fourn. ex Hack. l. c. = I. brasiliensis.

SYLLISIUM, Endl. Gen. Suppl. iii. 102 (1843) = seq.

SYLLYSIUM, Meyen & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 334 = **Eugenia**, Mich. (Myrtac.).
buxifolium, Meyen & Schau. l. c. = E. pyxophylla ?

SYLVIA, Lindl. Veg. Kingd. 685 (1847), sphalm. = **Silvia**, Benth. (Scrophul.).

SYMBASIANDRA, Willd. ex Steud. Nom. ed. II. i. 767 (1840) = **Hilaria**, H. B. & K. (Gramin.).
repens, Willd. ex Steud. l. c. = H. cenchroides.

SYMBLOMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 284 = **Albertinia**, Spreng. (Compos.).
Baldwiniana, Nutt. l. c. = A. brasiliensis.

SYMBOLANTHUS, G. Don, Gen. Syst. iv. 210 (1837) = **Lisianthus**, Linn. (Gentian.).
calygonus, Griseb. in DC. Prod. ix. 80 = L. calygonus.
Kunthii, G. Don, Gen. Syst. iv. 210 = L. anomalus.
Pavonii, G. Don, l. c. = L. calygonus.
superbus, Miers, in Journ. Bot. vii. (1869) 218.—N. Granat.

SYMBRYON, Griseb. Cat. Pl. Cub. 64 (1866). *PIPERACEAE*, Benth. & Hook. f. iii. 133.
tetrastachyum, *Griseb. l. c.*—Cuba.

SYMEA, Baker, in Saund. Refug. Bot. t. 260 (1871) = **Solaria**, Phil. (Liliac.).
gillesioides, Baker, l. c. = Solaria miersioides.

SYMETHUS, Rafin. Fl. Tellur. iv. 83 (1836) = **Convolvulus**, Tourn.
siculus, Rafin. l. c. = C. siculus.

SYMMERIA, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 630. *POLYGONACEAE*, Benth. & Hook. f. iii. 105.
THURNHEYSSERA, Mart. ex Meissn. in DC. Prod. xiv. 185 (1856).
paniculata, *Benth. in Hook. Lond. Journ. Bot.* iv. 1845) 630.—Bras.

SYMMETRIA, Blume, Bijdr. 1130 (1826) = **Carallia**, Roxb. (Rhizoph.).
obovata, Blume, l. c. 1131 = C. integerrima.

SYMPACHNE, Steud. Nom. ed. II. ii. 654 (1841), sphalm. = SYMPHACHNE, Beauv. = **Eriocaulon**, Linn.

SYMPEGMA, Bunge, in Bull. Acad. Pétersb. xxv. (1879) 371. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 76.
Regelii, *Bunge, l. c.*—Turkest.

SYMPETALEIA, A. Gray, in Proc. Am. Acad. xii. (1877) 161. *LOASEAE*.
aurea, *A. Gray, l. c.*—Am. bor. occ.

SYMPHACHNE, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 47. t. 5 = **Eriocaulon**, Linn.
xyroides, Beauv. ex Desv. l. c. 48 = E. decangulare.

SYMPHIANDRA, Steud. Nom. ed. II. ii. 655 (1841), sphalm. = **Symphyandra**, DC. (Campan.).

SYMPHIONEMA, R. Br. in Trans. Linn. Soc. x. (1810) 157 = **Symphyonema**, R. Br. (Proteac.).

SYMPHITUM, Neck. in Act. Acad. Theod. Palat. ii. (1770) 479 = **Symphytum**, Tourn. (Borag.).

SYMPHONIA, Linn. f. Suppl. 49, 303 (1781). *GUTTIFERAE*, Benth. & Hook. f. i. 173.
ACTINOSTIGMA, Welw. Apont. 560 (1858).
ANEURISCUS, Presl, Symb. Bot. i. 71. t. 48 (1832).
CHRYSOPIA, Thou. Gen. Nov. Madag. 14 (1806).
acuminata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 322.—Madag.
clusioides, *Baker, in Journ. Bot.* xx. (1882) 19.—Madag.
esculenta, Arruda, ex DC. Prod. i. 563 = globulifera, Moronobea grandiflora.
esculenta, Steud. Nom. ed. II. ii. 654, partim = Platonia insignis.
eugenioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 91.—Madag.
fasciculata, *Baill. Hist. Pl.* vi. 399.—Madag.
globulifera, *Linn. f. Suppl.* 302.—Guian.; Afr. trop.
grandiflora, Spreng. Syst. iii. 334 = Moronobea grandiflora.
lepidocarpa, *Baker, in Journ. Linn. Soc.* xx. (1883) 91.—Madag.
Melleri, *Baker, l. c.* 90.—Madag.
Moronobea, Raeusch. Nom. ed. III. 191 = Moronobea grandiflora.
pauciflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 90.—Madag.

SYMPHORANTHUS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 219 = **Polypremum**, Linn. (Logan.).

SYMPHOREMA, Roxb. Pl. Corom. ii. 46. t. 186 (1798). *VERBENACEAE*, Benth. & Hook. f. ii. 1159.
ANALECTIS, Juss. in Jaume St. Hil. Expos. ii. 362 (1805).
SZEGLEEWIA, Turcz. in Bull. Soc. Nat. Mosc. (1863) ii. 212.
glabrum, *Hassk. in Flora*, xlviii. (1865) 402.—Ins. Philipp.
grossum, *Kurz, For. Fl. Brit. Burma*, ii. 254.—Burma.
involucratum, *Roxb. Pl. Corom.* ii. 46. t. 186.—Ind. or.
involucratum, Wall. Cat. n. 1740 = polyandrum.
Jackianum, Kurz, For. Fl. Brit. Burma, ii. 255 = Sphenodesme pentandra.
luzonicum, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 162.—Ins. Philipp.
luzoniense, Vidal, Sinops. t. 75. f. F = praec.
microstylis, Bedd. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 600 = Sphenodesme microstylis.
paniculatum, Heyne, ex Wall. Cat. n. 1739, partim = involucratum.
pentandrum, Kurz, For. Fl. Brit. Burma, ii. 255 = Sphenodesme Griffithiana.
polyandrum, *Wight, Ic.* t. 363.—Ind. or.
unguiculatum, Kurz, For. Fl. Brit. Burma, ii. 255 = Sphenodesme unguiculata.

SYMPHORIA, Pers. Syn. i. 214 (1805) = **Symphoricarpos**, Dill. (Caprifol.).
albus, Rafin. New Fl. N. Am. iii. 21 = Symphoricarpos racemosus.
conglomerata, Pers. Syn. i. 214 = Symphoricarpos orbiculatus.
elongata, Torr. & Gray, Fl. N. Am. ii. 3 = Symphoricarpos racemosus.
glaucescens, Spreng. Syst. i. 757 = Symphoricarpos microphyllus.
globuliflora, Sweet, Hort. Brit. ed. III. 95, in syn. sphalm. = Symphonia globulifera.
glomerata, Pursh, Fl. Am. Sept. i. 162 = Symphoricarpos orbiculatus.

SYMPHORIA:—
heterophylla, Rafin. New Fl. Am. iii. 21; Torr. & Gray, Fl. N. Am. ii. 3 = Symphoricarpos racemosus.
leucocarpa, Hort. ex DC. Prod. iv. 339 = Symphoricarpos racemosus.
microphylla, Spreng. Syst. i. 757 = Symphoricarpos microphyllus.
montana, Spreng. l. c. = Symphoricarpos microphyllus.
occidentalis, R. Br. in Richards. App. Frankl. 2nd Journ. 733 = Symphoricarpos occidentalis.
parviflora, Buek, Ind. ad DC. Prod. i. 386 = Symphoricarpos orbiculatus.
punicea, Loud. Hort. Brit. 80 = Symphoricarpos puniceus.
racemosa, Pursh, Fl. Am. Sept. i. 162 = Symphoricarpos racemosus.
rubra, Rafin. New Fl. N. Am. iii. 21.—Am. bor.
vulgaris, D. Dietr. Syn. Pl. i. 783 = Symphoricarpos orbiculatus.

SYMPHORICARPA, Neck. Elem. i. 129 (1790) = seq.

SYMPHORICARPOS, Dill. ex Juss. Gen. 211 (1789). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 4.
ANISANTHUS, Willd. ex Roem. & Schult. Syst. v. p. xiv. (1819).
DESCLIAEA, [Moç. & Sesse,] ex DC. Prod. iv. 483 (1830).
MARGARIS, DC. l. c. (1830).
SYMPHORIA, Pers. Syn. i. 214 (1805).
albus, C. Koch, Dendrol. ii. 1. 48 = racemosus.
ciliatus, Nutt. ex Torr. & Gray, Fl. N. Am. ii. 4 = mollis.
conglomeratus, Pers. Syn. i. 214 = orbiculatus.
elongatus, Presl, ex DC. Prod. iv. 339 = racemosus.
erythrocarpus, Hort. ex C. Koch, Dendrol. ii. 1. 98 = orbiculatus.
glaucescens, H. B. & K. Nov. Gen. et Sp. iii. 424. t. 295 = microphyllus.
glomeratus, C. Koch, Dendrol. ii. 1. 48 = orbiculatus.
heterophyllus, Presl, ex DC. Prod. iv. 339 = racemosus.
imberbis, Tausch, in Flora, xxi. (1838) 734 = orbiculatus.
leucocarpus, Hort. ex C. Koch, Dendrol. ii. 1. 49 = racemosus.
longiflorus, *A. Gray, in Journ. Linn. Soc.* xiv. (1873) 12.—Am. bor. occ.
mexicanus, Hort. ex C. Koch, Dendrol. ii. 1. 50 = microphyllus.
microphyllus, *H. B. & K. Nov. Gen. et Sp.* iii. 424.—Mexic.
mollis, *Nutt. ex Torr. & Gray, Fl. N. Am.* ii. 4.—Am. bor. occ.
montanus, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 249 = oreophilus.
montanus, H. B. & K. Nov. Gen. et Sp. iii. 425. t. 296 = microphyllus.
montanus, S. Wats. Bot. King's Exped. 132 = rotundifolius.
occidentalis, *Hook. Fl. Bor. Am.* i. 285.—Am. bor.
odoratus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 145 = Schoepfia fragrans.
orbiculatus, *Moench, Meth.* 503.—Am. bor.
oreophilus, *A. Gray, in Journ. Linn. Soc.* xiv. (1873) 12.—Am. bor. occ.
parviflorus, Desf. Tabl. ed. I. 114 = orbiculatus.
poluccensis, C. Koch, Dendrol. ii. 1. 149, sphalm. = *toluccensis*.
puniceus, *Steud. Nom.* ed. II. ii. 654.—Am. bor.
racemosus, *Michx. Fl. Bor. Am.* i. 107.—Am. bor.
rotundifolius, *A. Gray, Pl. Wright.* ii. 66.—Mexic.
rotundifolius, C. Koch, Dendrol. ii. 1. 48 = orbiculatus.
spicatus, Engelm. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 215 = orbiculatus.
toluccensis, Hort. ex C. Koch, Dendrol. ii. 1. 49, 50 = microphyllus, racemosus.
vulgaris, Michx. Fl. Bor. Am. i. 106 = orbiculatus.

SYMPHORICARPUS, H. B. & K. Nov. Gen. et Sp. iii. 424 (1818) = praec.

SYMPHYANDRA, A. DC. Monog. Camp. 365. t. 8 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 563.
armena, *A. DC. l. c.* 366. t. 8.—Reg. Cauc.
cretica, *A. DC. l. c.*—Ins. Creta.

SYMPHYANDRA:—
heterophylla, Schur, Enum. Pl. Transs. 431 = Wanneri.
Hofmanni, *Pant. in Magyar Növényt. Lapok*, v. (1881) 150.—Bosnia.
lazica, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 888.—Pontus.
ossetica, *A. DC. Monog. Camp.* 366. t. 8.—Reg. Cauc.
pendula, *A. DC. l. c.*—Reg. Cauc.
stylosa, Royle, Illustr. Bot. Himal. 253 = Phyteuma Thomsoni.
Wanneri, *Heuff. in Flora*, xxxvii. (1854) 292.—Transsylv.

SYMPHYDOLON, Salisb. Gen. Pl. Fragm. 142 (1866) = **Gladiolus**, Tourn. (Irid.).
angustus, Salisb. l. c. = G. angustus.
blandus, Salisb. l. c. = G. blandus.
cardinalis, Salisb. l. c. = G. cardinalis.
carneus, Salisb. l. c. = G. cuspidatus.
cuspidatus, Salisb. l. c. = G. cuspidatus.
floribundus, Salisb. l. c. = G. floribundus.
hastatus, Salisb. l. c. = G. hastatus.
Milleri, Salisb. l. c. = G. Milleri.
undulatus, Salisb. l. c. = G. undulatus.

SYMPHYLLANTHUS, Vahl, in Skrivt. Nat. Hist. Selsk. vi. (1810) 86 = **Chailletia**, DC.
glaber, Vahl, l. c. 89.—Guian.
rugosus, Vahl, l. c. 88.—Guian.

SYMPHYLLIA, Baill. Etud. Gén. Euph. 473. t. 11 (1858) = **Adenochlaena**, Baill. (Euphorb.).
mallotiformis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 156 = A. indica.
siletiana, Baill. Etud. Gén. Euph. 474. t. 11 = A. silhetensis.
silhetense, Benth. in Journ. Linn. Soc. xvii. (1878) 228 = praec.

SYMPHYLLIUM, Benth. in Journ. Linn. Soc. xvii. (1878) 228 = praec.

SYMPHYLLOCARPUS, Maxim. Prim. Fl. Amur. 151. t. 8. f. 1 (1859). *COMPOSITAE*, Benth. & Hook. f. ii. 296.
exilis, *Maxim. l. c.*—Manchuria.

SYMPHYLOMA, Steud. Nom. ed. II. ii. 654 (1841), sphalm. = SYMPHYOLOMA, C. A. Mey. = **Heracleum**, Linn. (Umbell.).

SYMPHYODOLON, Baker, in Journ. Linn. Soc. xvi. (1877) 170, sphalm. = SYMPHYDOLON, Salisb. = **Gladiolus**, Tourn. (Irid.).

SYMPHYOGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 255 = **Cynanchum**, Linn. (Asclep.).
hastatum, Turcz. l. c. 256 = C. Wilfordi.

SYMPHYOLOMA, C. A. Mey. Verz. Pfl. Cauc. 127 (1831) = **Heracleum**, Linn. (Umbell.).
graveolens, C. A. Mey. l. c.—Reg. Cauc.

SYMPHYOMERA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 116 = **Cotula**, Tourn. (Compos.).
Filicula, Hook. f. l. c. 118 = C. Filicula.

SYMPHYOMYRTUS, Schau. in Lehm. Pl. Preiss. i. 126 (1844) = **Eucalyptus**, L'Hérit. (Myrtac.).
Lehmanni, Schau. l. c. 127 = E. Lehmanni.

SYMPHYONEMA, R. Br. in Trans. Linn. Soc. x. (1810) 157 (Symphionema); Spreng. Anleit. ii. 1. 333 (1817). *PROTEACEAE*, Benth. & Hook. f. iii. 174.
abrotanoides, Sieber, in Spreng. Syst. iv. Cur. Post. 46 = paludosum.
montanum, *R. Br. in Trans. Linn. Soc.* x. (1810) 48, 157.—Austral.
montanum, Endl. Ic. t. 12 = paludosum.
paludosum, *R. Br. in Trans. Linn. Soc.* x. (1810) 48, 157.—Austral.

SYMPHYOPAPPUS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 583. *COMPOSITAE*, Benth. & Hook. f. ii. 244.
KALLOPHYLLON, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 365 (1876).

SYMPHYOPAPPUS:—
cuneatus, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 367.—Bras.
decussatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) I. 584.—Bras.
polystachyus, *Baker, in Mart. Fl. Bras.* vi. II. 368.—Bras.
reticulatus, *Baker, l. c.* 367.—Bras.
vernicosus, Sch. Bip. ex Baker, l. c. 368 = reticulatus.
viscosus, *Sch. Bip. ex Baker, l. c.* 366.—Bras.

SYMPHYOPETALON, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 54 = **Nematolepis**, Turcz. (Rutac.).
corraeoides, J. Drumm. ex Harv. l. c. = N. phebalioides.

SYMPHYOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 159 (1861) = **Cleome**, Linn.(Cappar.).
strictus, Klotzsch, l. c. = C. strigosa.

SYMPHYOSTEMON, Miers, in Proc. Linn. Soc. i. (1841) 123; et in Trans. Linn. Soc. xix. (1843) 97. *IRIDEAE*, Benth. & Hook. f. iii. 700.
PSITHYRISMA, Herb. in Bot. Reg. (1843) Misc. 85.
SUSARIUM, Phil. in Linnaea, xxxiii. (1864–65) 248.
flexuosus, Miers, in Proc. Linn. Soc. i. (1841) 122, 123 = Sisyrinchium cuspidatum.
narcissoides, Phil. f. Cat. Pl. Vasc. Chil. 283 = odoratissimus.
odoratissimus, *Miers, in Proc. Linn. Soc.* i. (1841) 122, 123.—Am. austr.

SYMPHYOTRICHUM, Nees, Gen. et Sp. Ast. 135 (1833) = **Aster**, Tourn. (Compos.).
unctuosum, Nees, l. c. = A. Novi-Belgii.

SYMPHYSIA, Presl, Epist. ad Jacq. (1827); et in Oken, Isis, xxi. (1828) 274; xxiv. (1831) 489 = **Hornemannia**, Vahl (Vaccin.).
guadalupensis, Klotzsch, in Linnaea, xxiv. (1851) 37 = H. martinicensis.
martinicensis, Presl, Epist. ad Jacq. (1827); in Oken, Isis, xxi. (1828) 274; et in Flora, xi. (1828) 110 = H. martinicensis.

SYMPHYSICARPUS, Hassk. in Flora, xl. (1857) 101 = **Heterostemma**, Wight & Arn. (Asclep.).
chrysanthus, Hassk. l. c. 102.—Java.

SYMPHYSODAPHNE, A. Rich. Fl. Cub. Fanerog. ii. 190. t. 67 (1853) = **Misanteca**, Cham. (Laurin.).
cubensis, A. Rich. l. c. = M. cubensis.

SYMPHYTUM, Tourn. ex Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 854.
abchasicum, *Trautv. in Bull. Soc. Nat. Mosc.* xliii. (1870) I. 72.—Reg. Cauc.
album, Hort. ex Steud. Nom. ed. I. 821 = officinale.
anatolicum, *Boiss. Diagn.* Ser. I. iv. 43.—Anatolia.
angustifolium, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 227 = tuberosum.
armenum, Gundelsh. ex DC. Prod. x. 66 = Onosma sericea.
asperrimum, *Donn, in Sims, Bot. Mag.* t. 929.—Reg. Cauc.; Persia.
asperrimum, Urv. in Mém. Soc. Linn. Par. i. (1822) 276 = anatolicum.
asperum, Lepech. in Nov. Act. Acad. Petrop. xiv. (1805) 442. t. 7 = asperrimum.
azureum, *Van Hall, in Tijdschr. Wetensch.* ii. (1849) 128.—Hab.?
bohemicum, F. W. Schmidt, Fl. Boëm. iii. 13. t. 263 = officinale.
borragineum, Tausch, in Flora, xix. (1836) 393 = tauricum.
brachycalyx, *Boiss. Diagn.* Ser. I. iv. 43.—Kurdistan.
brochum, Bory & Chaub. Nouv. Fl. Pélop. 13. t. 8 = bulbosum.
bulbosum, *Schimp. in Flora*, viii. (1825) 17.—Europ.
bulbosum, Schur, Enum. Pl. Transs. 468 = tuberosum.
bullatum, Hornem. Hort. Hafn. i. 179 = tauricum.
calcaratum, *Clarke, Trav.* ii. 651.—Palaest.
Carmosina, Hort. Angl. ex Steud. Nom. ed. II. ii. 654 (Quid?).—Sibir.
caucasicum, *Bieb. Fl. Taur. Cauc.* i. 128.—Reg. Cauc.
caucasicum, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 294 = Donii.

SYMPHYTUM:—
Clusii, C. C. Gmel. Fl. Bad. iv. 144 = bulbosum.
coccineum, *Hort. ex Schlecht. in Bot. Zeit.* vii. (1849) 731.—Hab.?
coeruleum, *Hort. Angl. ex DC. Prod.* x. 40.—Sibir.
Consolida, Gueldenst. Reisen, i. 420; ex Ledeb. Fl. Ross. iii. 114 = officinale.
cordatum, Bieb. Fl. Taur. Cauc. i. 130 = grandiflorum.
cordatum, *Waldst. & Kit. Pl. Rar. Hung.* i. 6. t. 7.—Hungar.
cordatum, *Willd. in Ges. Naturf. Fr. Neue Schrift.* ii. (1799) 121.—Europ. centr.
cordifolium, Baumg. Enum. Stirp. Transs. i. 126 = cordatum.
Donii, *DC. Prod.* x. 37.—Hab.?
echinatum, Ledeb. Hort. Dorpat. Suppl. (1811) 5 = asperrimum.
elatum, Tausch, in Flora, xix. (1836) 393 = officinale.
filipendulum, Bisch. l. c. ix. (1826) 562 = bulbosum.
floribundum, Shuttlew. ex Nym. Consp. 509 = mediterraneum.
foliosum, *Rehm. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 495.—Rossia.
fruticosum, Moç. ex DC. Prod. x. 121 = Antiphytum mexicanum.
grandiflorum, *DC. Prod.* x. 40.—Reg. Cauc.
Gussonei, *F. Schultz, in Arch. Fl.* (1874) p. lviii; *et in Flora*, lviii. (1875) 218.—Sicil.
hirsutum, *Rafin. Med. Fl.* ii. 95.—Am. bor.
ibericum, Stev. ex Bieb. Fl. Taur. Cauc. iii. 647 = grandiflorum.
intermedium, Fisch. Hort. Gorenk. ed. II. 27; Roem. & Schult. Syst. iv. 66 (Quid?).—Hab.?
Jacquinianum, Tausch, in Flora, xix. (1836) 392, in obs. = orientale.
kurdicum, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* ii. 5.—Kurdist.
laeve, Bess. Cat. Hort. Crem. Suppl. (1812) (Quid?).—Podolia.
macrolepis, J. Gay, ex Reichb. Fl. Germ. Excurs. 347 = bulbosum.
majus, Gueldenst. ex Ledeb. Fl. Ross. iii. 115 = asperrimum.
mediterraneum, Guss. ex F. Schultz, in Flora, lviii. (1875) 218 = Gussonei.
mediterraneum, *Koch, Syn. Fl. Germ.* ed. I. 500.—Europ. austr.
microcalyx, Opiz, Seznam, 94 = officinale.
molle, *Janka, in Termész. Füzetek*, i. (1877) 29.—Hungar.
nodosum, Schur, Enum. Pl. Transs. 468 = tuberosum.
officinale, *Linn. Sp. Pl.* 136.—Europ.
orientale, *Linn. l. c.*—Byzant.
orientale, Pall. Tabl. Taur. 47; Bieb. Fl. Taur. Cauc. i. 129 = tauricum.
orientale, Pinard, ex DC. Prod. x. 40 = brachycalyx.
ottomanum, *Frivald. in Flora*, xix. (1836) 439.—Rumelia.
Palestinae, *Boiss. Diagn.* Ser. I. xi. 94.—Syria.
pannonicum, Pers. Syn. i. 161 = cordatum.
patens, Fries, Mant. ii. 13 = orientale.
patens, Sibth. Fl. Oxon. 70 = officinale.
peregrinum, Ledeb. Ind. Sem. Hort. Dorpat. (1820) 4; et ex Spreng. Syst. i. 563 = asperrimum.
pictum, *Hort.—Cf. Gard. Chron.* (1884) I. 827.—Hab.?
punctatum, Gaud. Fl. Helv. ii. 41 = bulbosum.
racemosum, Steph. ex Roem. & Schult. Syst. iv. 752 = caucasicum.
regium, S. T. Gmel. Reise, iii. 363. t. 36. f. 1 (= *Cynoglossum umbellatum*).—Hungar.
secundum, S. T. Gmel. l. c. = Echium glomeratum.
sepulcrale, *Boiss. & Bal. ex Boiss. Fl. Orient.* iv. 174.—Pontus.
sylvaticum, *Boiss. Pl. Or. Nov. Dec.* ii. 4.—Pontus.
tanaicense, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 577.—Rossia.
tauricum, *Willd. in Ges. Naturf. Fr. Neue Schr.* ii. (1799) 121. t. 5. f. 1.—Reg. Cauc.; Armen.
tuberosum, *Linn. Sp. Pl.* 136.—Europ. austr.
uliginosum, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 227 = officinale.
uplandicum, Nym. Syll. 80 = orientale.
variegatum, Hort. ex Gard. Chron. (1880) I. 779, cum cit. falsa = officinale.
Zeyheri, Schimp. in Flora, xii. (1829) 418 = bulbosum.

SYMPIEZA, Licht. ex Roem. & Schult. Syst. iii. 8, 171 (1818). *ERICACEAE*, Benth. & Hook. f. ii. 593.
brachyphylla, *Benth. in DC. Prod.* vii. 706.—Afr. austr.
capitellata, *Licht. ex Roem. & Schult. Syst.* iii. 8, 171.—Afr. austr.
Eckloniana, *Klotzsch, in Linnaea*, viii. (1823) 656.—Afr. austr.
Kunthii, *Klotzsch, l. c.* 229.—Afr. austr.
tenuiflora, *Benth. in DC. Prod.* vii. 706.—Afr. austr.

SYMPLEURA, Miers, in Lindl. Veg. Kingd. 593 a (1847), nomen (Styrac.?).—Cf. Benth. & Hook. f. ii. 668.

SYMPLOCARPUS, Salisb. ex Nutt. Gen. Am. i. 105 (1818). *AROIDEAE*, Benth. & Hook. f. iii. 995.
ICTODES, Bigel. Med. Bot. ii. 41. t. 24 (1819).
SPATHYEMA, Rafin. in Med. Repos. N. York, v. (1808) 352.
foetidus, *Nutt. Gen. Am.* i. 106.—Am. bor.
kamtschaticus, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 168 = Lysichitum camtschatcense.

SYMPLOCOS, Jacq. Enum. Pl. Carib. 5 (1760). *STYRACEAE*, Benth. & Hook. f. ii. 668.
ALSTONIA, Linn. f. Suppl. 39 (1781).
AYPARIA, Rafin. Sylva Tellur. 154 (1838).
BARBERINA, Vell. Fl. Flum. 235 (1825); Ic. iv. t. 117 (1827).
BOBU, Adans. Fam. ii. 88 (1763).
CARLEA, Presl, Epim. Bot. 216 (1850).
CIPONIMA, Aubl. Pl. Gui. i. 566. t. 226 (1775).
CORDYLOBLASTE, Mor. in Bot. Zeit. vi. (1848) 604.
DICALIX, Lour. Fl. Cochinch. 663 (1790).
EUGENIOIDES, Linn. Fl. Zeyl. 192 (1747).
HOPEA, Linn. Mant. i. 14 (1767).
HYPOPOGON, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 246.
LODHRA, Guill. in Ann. Sc. Nat. Sér. II. xv. (1841) 161.
MONGEZIA, Vell. Fl. Flum. 229 (1825).
PALURA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 145 (1825).
PRAEALSTONIA, Miers, in Journ. Linn. Soc. xvii. (1879) 291.
PROTOHOPEA, Miers, l. c. (1879) 289.
SARIAVA, Reinw. in Syll. Ratisb. ii. (1828) 12.
SCYRTOCARPUS, Miers, in Lindl. Veg. Kingd. 593 a (1847).
SIPONIMA, A. DC. in DC. Prod. viii. 246 (1844).
STEMMATOSIPHUM, Pohl, Pl. Bras. Ic. ii. 86. tt. 157–159 (1831).
acuminata, *Bedd. Forest. Man. Bot.* 150.—Malabar.
acuminata, *Miq. Fl. Ind. Bat.* i. II. 467.—Java.
acuta, *Thw. Enum. Pl. Zeyl.* 186.—Zeylan.
adenophylla, *Wall. Cat.* n. 4427.—Malaya.
adenopus, *Hance, in Journ. Bot.* xxi. (1883) 322.—China.
affinis, *Hort. ex Pasq. Cat. Orto Bot. Nap.* 100, nomen.—Hab.?
Alstonia, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 176.—N. Granat.
anamallayana, *Bedd. Forest. Man. Bot.* 150.—Travancor.
angustata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 585.—Zeylan.
apicalis, *Thw. Enum. Pl. Zeyl.* 187.—Zeylan.
arborea, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 429.—N. Caled.
arbutifolia, Casar. Nov. Stirp. Bras. Dec. 30 = phaeoclados.
Arechea, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 176.—Am. austr.
atroviridis, Wall. ex A. DC. in DC. Prod. viii. 254 = spicata.
attenuata, Wall. Cat. n. 4426 = spicata.
bahiensis, *A. DC. in DC. Prod.* viii. 250.—Bras.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 187, 476.—Ins. Banca.
baptica, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 430.—N. Caled.
Beddomei, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 582 = reticulata.
Berterii, Miers, in Journ. Linn. Soc. xvii. (1879) 286 = martinicensis.
bractealis, *Thw. Enum. Pl. Zeyl.* 185.—Zeylan.

SYMPLOCOS:—

brevipes, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 577, in obs. = coccinea.
caerulescens, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 429.—N. Caled.
caryophylloides, *Zoll. Syst. Verz. Ind. Arch.* 136; *et in Tijdschr. Nederl. Ind.* xiv. (1857) 161.—Java.
caudata, *Wall. Cat.* n. 4413.—Reg. Himal.; China; Japon.
celastrifolia, *Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 575.—Malaya.
celastrinea, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 31.—Bras.
cerasifolia, *Wall. Cat.* n. 4434.—Malaya.
cernua, Benth. Pl. Hartw. 78 = Hartwegi.
cernua, *Humb. & Bonpl. Pl. Aequin.* i. 188. t. 53.—Peruv.
ciliata, *Benth. Pl. Hartw.* 78, in nota.—Mexic.
ciliata, *Miq. Fl. Ind. Bat.* i. II. 466.—Java.
Ciponima, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 175.—Am. trop.
ciponimoides, *Griseb. Cat. Pl. Cub.* 167.—Cuba.
citrea, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 22.—Mexic.
Claussenii, *A. DC. in DC. Prod.* viii. 251.—Bras.
coccinea, *Humb. & Bonpl. Pl. Aequin.* i. 185. t. 52.—Mexic.
congesta, *Benth. Fl. Hongk.* 211.—China.
congesta, *Zoll. Syst. Verz. Ind. Arch.* 136.—Malaya.
cordifolia, *Thw. Enum. Pl. Zeyl.* 187.—Zeylan.
coriacea, *A. DC. in DC. Prod.* viii. 248.—Peruv.
coronata, *Thw. Enum. Pl. Zeyl.* 187.—Zeylan.
costaricana, *Hemsl. Biol. Centr. Am. Bot.* ii. 301.—Costa Rica.
costata, *Choisy, ex Zoll. Syst. Verz. Ind. Arch.* 136.—Java.
crassifolia, *Benth. Fl. Hongk.* 212.—China.
crassipes, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 580.—Malacca.
crataegoides, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 145.—Reg. Himal.; Burma; China; Japon.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 518.—Cuba.
cuneata, *Thw. Enum. Pl. Zeyl.* 186.—Zeylan.
decora, *Hance, in Journ. Bot.* xii. (1874) 369.—China.
dryophila, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 578.—Reg. Himal.
elegans, *Thw. Enum. Pl. Zeyl.* 185.—Zeylan.
estrellensis, *Casar. Nov. Stirp. Bras. Dec.* 32.—Bras.
fasciculata, Roxb. ex A. DC. Prod. viii. 259 = Eurya japonica.
fasciculata, *Zoll. Syst. Verz. Ind. Arch.* 136.—Malaya.
ferruginea, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 542.—Reg. Himal.; Malaya.
flavida, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 573 = spicata.
floribunda, Wall. Cat. n. 4419 = S. Sumuntia.
foliosa, *Wight, Ic.* t. 1234.—Ind. or.
Fordii, *Hance, in Journ. Bot.* xx. (1882) 78.—China.
Gardneriana, *Wight, Ic.* t. 1231.—Ind. or.
glaucescens, *Vieill. ex Miers, in Journ. Linn. Soc.* xvii. (1879) 305.—N. Caled.
glomerata, *King, ex Gamble, Darjeeling List*, 54.—Reg. Himal.
gracilis, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 431.—N. Caled.
grandiflora, *Wall. Cat.* n. 4421.—Reg. Himal.
Hamiltoniana, Wall. Cat. n. 4420 = racemosa.
Hartwegi, *A. DC. in DC. Prod.* viii. 252.—Mexic.
hebantha, *Thw. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 586.—Zeylan.
Henscheli, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 588.—Malaya; China.
hirsuta, *A. DC. in DC. Prod.* viii. 253.—Bras.
hirsuta, Wight & Gardn. ex Thw. Enum. Pl. Zeyl. 185 = minor, latiflora.
hispidula, *Thw. Enum. Pl. Zeyl.* 186.—Zeylan.
Hohenackeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 582.—Ind. or.
Hookeri, *C. B. Clarke, l. c.* 578.—Reg. Himal.
Horsfieldiana, Miq. Fl. Ind. Bat. Suppl. 475 = ferruginea.
Itatiaiae, *Wawra, Itin. Princ. Coburgi*, i. 77. t. 4. f. B.—Bras.
iteophylla, Miq. Fl. Ind. Bat. Suppl. 187, 476 = adenophylla.
japonica, *A. DC. in DC. Prod.* viii. 255.—China; Japon.
javanica, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 64 = ferruginea.
jucunda, *Thw. Enum. Pl. Zeyl.* 186.—Zeylan.
Jurgensenii, *Hemsl. Biol. Centr. Am. Bot.* ii. 301.—Mexic.
kurgensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 576.—Ind. or.
lachnobotrya, Miq. Fl. Ind. Bat. Suppl. 475 = ferruginea.
laeta, *Thw. Enum. Pl. Zeyl.* 184.—Zeylan.
lanceaefolia, Brongn. ex Miers, in Journ. Linn. Soc. xvii. (1879) 305 = Lenormandiana.
lanceolata, *A. DC. in DC. Prod.* viii. 253.—Bras.
lancifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 133.—Assam; China; Japon.
latiflora, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 585.—Zeylan.
laurina, Wall. Cat. n. 4416 = spicata.
laxiflora, *Benth. in Trans. Linn. Soc.* xviii. (1841) 232. t. 18.—Bras.
leiostachya, Kurz, For. Fl. Brit. Burma, ii. 144, partim = nicobarica.
leiostachya, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 89.—Burma.
Lenormandiana, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 428.—N. Caled.
leptostachys, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 134 = lancifolia.
leucantha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 89.—Burma.
Limoncillo, Humb. & Bonpl. Pl. Aequin. i. 196 = martinicensis.
linearis, *Zoll. Syst. Verz. Ind. Arch.* 136, nomen.—Japon.
Loha, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 144 = spicata.
Lohu, Miers, in Journ. Linn. Soc. xvii. (1879) 298 = praec.
lucida, *Brongn. & Gris, ex Miers, l. c.* 305, in syn.—N. Caled.
lucida, Sieb. & Zucc. Fl. Jap. i. 55. t. 24 = japonica.
lucida, Wall. Cat. n. 4414 = theaefolia.
Lundii, *A. DC. in DC. Prod.* viii. 251.—Bras.
macrocarpa, *Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 582.—Ind. or.
macrophylla, *Wall. Cat. n.* 4431.—Reg. Himal.
Maingayi, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 588.—Malacca.
marginalis, *Thw. Enum. Pl. Zeyl.* 187.—Zeylan.
Martii, *A. DC. in DC. Prod.* viii. 252.—Bras.
martinicensis, *Jacq. Enum. Pl. Carib.* 24; *Linn. Sp. Pl.* ed. II. 747.—Ind. occ.
Mathewsii, *A. DC. in DC. Prod.* viii. 250 (Matthewsii).—Peruv.
microcarpa, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 303 = lancifolia.
microphylla, *Wight, Ic.* t. 1232.—Ind. or.
minor, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 586.—Zeylan.
mollis, Wall. Cat. n. 4433 = ferruginea.
monantha, *Wight, Ic.* t. 1236.—Ind. or.
montana, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 430.—N. Caled.
mucronata, *Humb. & Bonpl. Pl. Aequin.* i. 196.—N. Granat.
myrtacea, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 133.—Japon.
neriifolia, *Sieb. & Zucc. l. c.* 134.—Japon.
nervosa, A. DC. in DC. Prod. viii. 256 = racemosa.
nervosa, Wight, Ic. t. 1235 = foliosa.
nicobarica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 580.—Ins. Nicobar.
nitens, *Benth. in Trans. Linn. Soc.* xviii. (1841) 232.—Bras.
nitida, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 430.—N. Caled.
nuda, *Humb. & Bonpl. Pl. Aequin.* i. 195.—N. Granat.
oblongifolia, *Casar. Nov. Stirp. Bras. Dec.* 31.—Bras.
oblongifolia, *Vidal, Phan. Cuming. Philipp.* 33, 124.—Ins. Philipp.
obovata, *A. DC. in DC. Prod.* viii. 251.—Bras.
obovata, Wight & Gardn. ex Thw. Enum. Pl. Zeyl. 185 = obtusa.
obtusa, *Wall. Cat.* n. 4424.—Ind. or.
octopetala, *Sw. Prod. Veg. Ind. Occ.* 109.—Jamaic.
odoratissima, *Choisy, ex Zoll. Syst. Verz. Ind. Arch.* 136.—Java.
oligandra, *Bedd. Forest. Man. Bot.* 150.—Travancor.
ophirensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 579.—Malacca.
ovalis, *Wright, ex Griseb. Cat. Pl. Cub.* 168.—Cuba.
oxyphylla, *Wall. Cat.* n. 4430.—Reg. Himal.; Burma.
pallida, Franch. & Sav. Enum. Pl. Jap. i. 308 = crataegoides.
paniculata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 102.—Japon.
paniculata, Wall. ex D. Don, Prod. Fl. Nep. 145; et Cat. n. 4429 = crataegoides.
paraënsis, Endl. ex Miq. in Mart. Fl. Bras. vii. 24 = S. Ciponima.
parviflora, *Benth. in Trans. Linn. Soc.* xviii. (1841) 232.—Bras.
parvifolia, *Benth. Pl. Hartw.* 140.—Ecuador; Peruv.
pauciflora, *Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 587.—Zeylan.
pedicellata, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 89 = oxyphylla.
pendula, Thw. Enum. Pl. Zeyl. 84 = pauciflora.
pendula, *Wight, Ic.* t. 1237.—Zeylan.
pentagyna, Spreng. Neue Entdeck. ii. 129 = Vismia decipiens.
phaeoclados, *A. DC. in DC. Prod.* viii. 253.—Bras.
phyllocalyx, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 575 = japonica.
platyphylla, *Benth. in Trans. Linn. Soc.* xviii. (1841) 233.—Bras.
polycarpa, Wall. Cat. n. 4423 = spicata.
polystachya, Wall. l. c. n. 4428 = ferruginea.
prionophylla, *Hemsl. Biol. Centr. Am. Bot.* ii. 302.—Mexic.
propinqua, Hance, in Journ. Bot. vi. (1868) 329 = racemosa.
prunifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 133 = caudata.
pubescens, *Klotzsch, ex Benth. in Trans. Linn. Soc.* xviii. (1841) 233.—Bras.
pulchra, *Wight, Ic.* t. 1230.—Ind. or.
pycnantha, *Hemsl. Biol. Centr. Am. Bot.* ii. 302.—Mexic.
pycnobotrya, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 25.—Bras.
pyrifolia, *Wall. Cat.* n. 4415.—Reg. Himal.
racemosa, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 539.—Ind. or.; China.
racemosa, Wall. Cat. n. 4418 = spicata, theaefolia, S. Sumuntia.
racemosa, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 583 = rosea.
ramentacea, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 33.—Bras.
ramosissima, *Wall. Cat.* n. 4425.—Reg. Himal.
reflexa, *A. DC. in DC. Prod.* viii. 248.—Peruv.
repandula, *Miq. Fl. Ind. Bat. Suppl.* 475.—Sumatra.
reticulata, *J. Grah. Cat. Bomb. Pl.* 104.—Ind. or.
revoluta, *Casar. Nov. Stirp. Bras. Dec.* 31.—Bras.
revoluta, *A. DC. in DC. Prod.* viii. 253.—Bras.
rhamnifolia, *A. DC. l. c.*—Bras.
Ribes, *Jungh. & De Vriese, in De Vriese, Pl. Nov. Ind. Batav.* fasc. i. 11.—Java.
rigida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 581.—Malacca.
rigida, Wall. Cat. n. 4422 = racemosa.
rosea, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 51.—Travancor.
rotundifolia, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 430.—N. Caled.
rubiginosa, *Wall. Cat.* n. 4432.—Malaya.
rufescens, *Humb. & Bonpl. Pl. Aequin.* i. 192. t. 55.—N. Granat.
rufescens, Thw. Enum. Pl. Zeyl. 184 = versicolor.
salaccensis, *Miq. Fl. Ind. Bat.* i. II. 466.—Java.
salicifolia, Buch.-Ham. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 579 = pyrifolia.
salicifolia, *Griseb. Cat. Pl. Cub.* 168.—Ind. occ.
Samuntia, Miers, in Journ. Linn. Soc. xvii. (1879) 298, sphalm. = S. Sumuntia.
Schiedeana, *Schlecht. in Linnaea*, viii. (1833) 527.—Bras.
Schomburgkii, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1087, nomen.—Guian.
serrulata, *Humb. & Bonpl. Pl. Aequin.* i. 190. t. 54.—N. Granat.
sessilis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 587.—Travancor.

SYMPLOCOS:—
sinica, [*Ker-Gawl.*] *Bot. Reg.* t. 710.—China.
speciosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 302.—Mexic.
spicata, *Roxb. Hort. Beng.* 40; *Fl. Ind.* ii. 541.—As. et Austral. trop.
Stawellii, F. Muell. Fragm. v. 60 = spicata.
stravadioides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 428.—N. Caled.
subcordata, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 585 = hispidula.
subsessilis, *Choisy, ex Zoll. Syst. Verz. Ind. Arch.* 136.—Java.
subspinosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 144 = spicata.
sulcata, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 65.—Burma.
Sumuntia, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 145.—Reg. Himal.
Swinhoeana, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 226 = caudata.
tetrandra, *Mart. Syst. Mat. Med. Bras.* 49.—Bras.
theaefolia, *D. Don, Prod. Fl. Nep.* 145.—Reg. Himal.; Burma.
theophrastaefolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 134.—Japon.
Thwaitesii, *F. Muell. Fragm.* iii. 22.—Austral.
tinctoria, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 176.—Am. bor.
tomentosa, *Humb. & Bonpl. Pl. Aequin.* i. 195.—N. Granat.
uniflora, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 51 = anamallayana.
uniflora, *Benth. in Trans. Linn. Soc.* xviii. (1840) 233.—Bras.
urceolaris, *Hance, in Journ. Bot.* xiv. (1876) 307.—China.
variabilis, *Mart. ex Miq. in Mart. Fl. Bras.* vii. 30.—Bras.
Verhuelli, Jungh. & De Vriese, in De Vriese, Pl. Nov. Ind. Batav. i. 12 = ferruginea.
versicolor, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 584.—Zeylan.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 429.—N. Caled.
virgata, Royle, Illustr. Bot. Himal. 464 (Index), sphalm. = Styrax serrulatum.
xanthophylla, *Jungh. & De Vriese, in De Vriese, Pl. Nov. Ind. Batav.* i. 11.—Java.

SYMPODIUM, C. Koch, in Linnaea, xvi. (1842) 356 = **Carum**, Rupp. (Umbell.).
simplex, C. Koch, l. c. = C. elegans.

SYNACTILA, Rafin. Fl. Tellur. iv. 104 (1836) = **Passiflora**, Linn.
viridiflora, Rafin. l. c. = P. viridiflora.

SYNADENA, Rafin. Fl. Tellur. iv. 9 (1836) = **Phalaenopsis**, Blume (Orchid.).
amabilis, Rafin. l. c. = P. amabilis.

SYNADENIUM, Boiss. in DC. Prod. xv. II. 187 (1862). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 261.
arborescens, *Boiss. l. c.*—Afr. austr.
carinatum, *Boiss. l. c.*—Ins. Borbon.
Grantii, *Hook. f. Bot. Mag.* t. 5633.—Afr. trop.

SYNAECIA, Pritz. Ic. Ind. i. 1048 (1855), sphalm. = SYNOECIA, Miq. = **Ficus**, Tourn. (Urtic.).

SYNAEDRIS, Steud. Nom. ed. II. ii. 655 (1841), sphalm. = seq.

SYNAEDRYS, Lindl. Introd. Nat. Syst. ed. II. 441 (1836) = **Quercus**, Tourn. (Cupul.).
ossea, Lindl. l. c. = Q. cornea.

SYNALLODIA, Rafin. New Fl. Am. iv. 93 (1836). (Gentianac. ?)
japonica, Rafin. l. c. (Quid ?).—Japon.

SYNANDRA, Nutt. Gen. Am. ii. 29 (1818). *LABIATAE*, Benth. & Hook. f. ii. 1205.
TORREYA, Rafin. in Am. Monthly Mag. (1811) 356.
grandiflora, *Nutt. Gen. Am.* ii. 29.—Am. bor.

SYNANDRA, Schrad. in Goett. Gel. Anz. (1821) I. 715 = **Stenandrium**, Nees (Acanth.).
amoena, Schrad. l. c. 716 = Sten. igneum.

SYNANDRODAPHNE, Meissn. in DC. Prod. xv. I. 176 (1864). *LAURINEAE*, Benth. & Hook. f. iii. 159.
antillana, Meissn. l. c. (= *Oreodaphne fragrans*).—Ind. occ.
laxa, *Meissn. l. c.*—N. Granat.

SYNANDROSPADIX, Engl. Bot. Jahrb. iv. (1883) 61. *AROIDEAE*.
vermitoxicus, *Engl. l. c.* 62.—Reg. Argent.

SYNANTHERIAS, Schott, Gen. Aroid. t. 28 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 972.
sylvatica, *Schott, l. c.*—Ind. or.

SYNAPHE, Dulac, Fl. Hautes-Pyr. 90 (1867) = **Festuca**, Tourn. (Gramin.).
rigida, Dulac, l. c. = F. rigida.

SYNAPHEA, R. Br. in Trans. Linn. Soc. x. (1810) 155. *PROTEACEAE*, Benth. & Hook. f. iii. 173.
acutiloba, *Meissn. in Lehm. Pl. Preiss.* i. 528.—Austral.
brachystachya, Lindl. Swan River App. 32 = polymorpha.
decorticans, *Lindl. l. c.*—Austral.
dilatata, *R. Br. in Trans. Linn. Soc.* x. (1810) 156.—Austral.
Drummondii, Meissn. in Hook. Kew Journ. iv. (1852) 183 = dilatata.
favosa, *R. Br. in Trans. Linn. Soc.* x. (1810) 156.—Austral.
gracillima, Lindl. Swan River App. 32 = petiolaris.
petiolaris, *R. Br. in Trans. Linn. Soc.* x. (1810) 156.—Austral.
pinnata, *Lindl. Swan River App.* 32.—Austral.
polymorpha, *R. Br. in Trans. Linn. Soc.* x. (1810) 156.—Austral.
Preissii, *Meissn. in Lehm. Pl. Preiss.* i. 529.—Austral.

SYNAPISMA, Steud. Nom. ed. II. ii. 655 (1841) sphalm. = SYNASPISMA, Endl. = **Codiaeum**, Rumph (Euphorb.).

SYNAPSIS, Griseb. Cat. Pl. Cub. 187 (1866). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 943.
ilicifolia, *Griseb. l. c.*—Cuba.

SYNAPTANTHA, Hook. f. in Hook. Ic. Pl. t. 1146 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 61.
tillaeacea, *Hook. f. in Hook. Ic. Pl.* t. 1146.—Austral.

SYNAPTEA, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65 = SUNAPTEA, Griff. = **Vatica**, Linn. (Dipteroc.).

SYNAPTOLEPIS, Oliver, in Hook. Ic. Pl. t. 1074 (1870). *THYMELAEACEAE*, Benth. & Hook. f. iii. 198.
alternifolia, *Oliver, l. c.* t. 1194.—Afr. trop.
Kirkii, *Oliver, l. c.* t. 1074.—Afr. trop.

SYNARRHENA, Fisch. & Mey. in Bull. Sc. Acad. Pétersb. viii. (1841) 255 = **Mimusops**, Linn. (Sapot.).
floribunda, Fisch. & Mey. l. c. 256 = M. floribunda.
subsericea, Fisch. & Mey. l. c. 255 = M. subsericea.

SYNARTHRON, Benth. & Hook. f. Gen. ii. 477 (1873) = seq.

SYNARTHRUM, Cass. in Dict. Sc. Nat. xlviii. 448 (1827) = **Senecio**, Tourn. (Compos.).
appendiculatum, Cass. l. c. = Senecio appendiculatus.

SYNASPISMA, Endl. Gen. 1110 (1840) = **Codiaeum**, Rumph. (Euphorb.).
peltatum, Endl. l. c. = C. inophyllum.

SYNASSA, Lindl. Orch. Scel. 9 (1826); Bot. Reg. sub t. 1618 (1833) = **Spiranthes**, Rich. (Orchid.).
corymbosa, Lindl. Bot. Reg. sub t. 1618 (= *Pelexia corymbosa*).—Peruv.

SYNCARPHA, DC. in Ann. Mus. Par. xvi. (1810) 205. t. 5. f. 31 = **Helipterum**, DC. (Compos.).
gnaphaloides, DC. l. c. 207 = H. gnaphaloides.
staehelinoides, Steud. Nom. ed. II. ii. 655 = praec.

SYNCARPIA, Tenore, Ind. Sem. Hort. Neap. (1839) 12. n. 13; et in Ann. Sc. Nat. Sér. II. xiii. (1840) 381.
MYRTACEAE, Benth. & Hook. f. i. 709.
KAMPTZIA, Nees, Progr. Gratul. Kamptzia (1840); et in Nov. Act. Nat. Cur. xviii. Suppl. i. p. ix. (1841).
SINCARPIA, Tenore, in Mem. Soc. Ital. Moden. xxii. (1841) 230.
Hillii, *Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 86.—Austral.
laurifolia, *Tenore, Ind. Sem. Hort. Neap.* (1839) 12. n. 13; *et in Ann. Sc. Nat.* Sér. II. xiii. (1840) 381.—Austral.
leptopetala, *F. Muell. Fragm.* i. 79.—Austral.
Vertholenii, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 411.—Ins. Amboin.

SYNCEPHALANTHA, Bartl. Ind. Sem. Hort. Gotting. (1836) 6.—Cf. Linnaea, xii. (1838) Litt. 80. *COMPOSITAE*, Benth. & Hook. f. ii. 409.
decipiens, *Bartl. l. c.*—Mexic.

SYNCEPHALANTHUS, Benth. & Hook. f. Gen. ii. 409 (1873), err. typ. = praec.

SYNCEPHALUM, DC. Prod. vi. 282 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 326.
Bojeri, *DC. l. c.*—Madag.

SYNCHODENDRON, Boj. ex DC. Prod. v. 92 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 288.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 279.—Madag.
pauciflorum, Steud. Nom. ed. II. ii. 655, sphalm. = seq.
ramiflorum, *DC. Prod.* v. 93.—Madag.
senegalense, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 364.—Afr. austr.

SYNCLISIA, Benth. in Benth. & Hook. f. Gen. i. 36 (1862). *MENISPERMACEAE*, Benth. & Hook. f. l. c.
scabrida, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xx. (1867) 171.—Afr. trop.

SYNCODIUM, Rafin. Fl. Tellur. ii. 22 (1836) = **Scilla**, Linn. (Liliac.).
nutans, Rafin. l. c. = S. festalis.

SYNCODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111 = **Campanula**, Tourn.
cervicarium, Fourr. l. c. = C. Cervicaria.
glomeratum, Fourr. l. c. = C. glomerata.
thyrsoideum, Fourr. l. c. = C. speciosa.

SYNCOLOSTEMON, E. Mey. Comm. Pl. Afr. Austr. 230 (1837). *LABIATAE*, Benth. & Hook. f. ii. 1174.
densiflorus, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 230.—Afr. austr.
dissitiflorus, *Benth. in DC. Prod.* xii. 54.—Afr. austr.
parviflorus, *Benth. in E. Mey. Comm. Pl. Afr. Austr.* 230.—Afr. austr.
ramulosus, *Benth. l. c.*—Afr. austr.
rotundifolius, *Benth. l. c.*—Afr. austr.

SYNDESMANTHUS, Klotzsch, in Linnaea, xii. (1838) 240 = **Simocheilus**, Klotzsch (Ericac.).
articulatus, Klotzsch, l. c. 241.—Afr. austr.
capitellatus, Klotzsch, ex Benth. in DC. Prod. vii. 705 = Simocheilus oblongus.
ciliatus, Benth. l. c. 707.—Afr. austr.
fasciculatus, Klotzsch, in Linnaea, xii. (1838) 240 = Simocheilus glabellus.
glaucus, Klotzsch, l. c.—Afr. austr.
paucifolius, Benth. in DC. Prod. vii. 707.—Afr. austr.
scaber, Klotzsch, in Linnaea, xii. (1838) 241.—Afr. austr.
squarrosus, Benth. in DC. Prod. vii. 707.—Afr. austr.

SYNDESMIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 314 (1824) = **Gluta**, Linn. (Anacard.).
coarctatus, Griff. Notul. iv. 409. t. 567. f. 1 = G. velutina.
elegans, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 314 = G. elegans.
tavoyana, Wall. Cat. n. 1004 = G. tavoyana.

SYNDESMON, Hoffmgg. in Flora, xv. (1832) II. Intell. n. iv. (err. typ. n. iii.) 34 = **Anemone**, Linn. (Ranunc.).
thalictroides, Hoffmgg. l. c. = A. thalictroides.
tuberosum, Hoffmgg. l. c. = Thalictrum tuberosum.

SYNECHANTHUS, H. Wendl. in Bot. Zeit. xvi. (1858) 145. *PALMAE*, Benth. & Hook. f. iii. 912.
RATHEA, Karst. in C. Koch & Fint. Wochenschr. i. (1858) 377.
REINECKIA, Karst. l. c. 349 (1858).
angustifolius, *H. Wendl. in C. Koch & Fint. Wochenschr.* (1859) 15.—Costa Rica.
fibrosus, *H. Wendl. in Bot. Zeit.* xvi. (1858) 145.—Guatemala.
Warszewiczianus, *H. Wendl. l. c.*—Costa Rica.

SYNEDRELLA, Gaertn. Fruct. ii. 456. t. 171 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 383.
UCACOU, Adans. Fam. ii. 131 (1763), partim.
nodiflora, *Gaertn. Fruct.* ii. 456. t. 171.—Am. trop.
peduncularis, *Benth. Pl. Hartw.* 119.—Ecuador.
vialis, A. Gray, in Proc. Am. Acad. xvii. (1882) 217 = Blainvillea tampicana.

SYNEILESIS, Maxim. Prim. Fl. Amur. 165 (1859) = **Senecio**, Tourn. (Compos.).
aconitifolia, Maxim. l. c. = Senecio aconitifolius.
palmata, Maxim. in Bull. Acad. Pétersb. xix. (1874) 487 = Senecio Syneilesis.

SYNELCOSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 345 = **Tordylium**, Linn. (Umbell.).
Carmeli, Boiss. l. c. = Heracleum Carmeli.

SYNEMA, Dulac, Fl. Hautes-Pyr. 154 (1867) = **Mercurialis**, Tourn. (Euphorb.).
annuum, Dulac, l. c. = M. annua.
perenne, Dulac, l. c. = M. perennis.

SYNEXEMIA, Rafin. Neogenyt. 2 (1825) = **Phyllanthus**, Linn. (Euphorb.).
caroliniana, Rafin. l. c. = P. carolinensis.
cuneifolia, Rafin. New Fl. Am. iv. 100 (Quid?).—Am. bor.
humilis, Muell. Arg. in DC. Prod. xv. II. 399, sphalm. = P. pumilus.
obovata, Rafin. Neogenyt. 2 = P. carolinensis.
pumila, Rafin. l. c. = P. pumilus.

SYNGONIUM, Schott, in Wien. Zeitschr. iii. (1829) 780; ex Endl. in Linnaea, vi. (1831) Litt. 53. *AROIDEAE*, Benth. & Hook. f. iii. 979.
affine, *Schott, Syn. Aroid.* 67.—Am. trop.
albolineatum, *Bull. Cat.* (1871) 6.—Am. centr.
angustatum, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 178.—Nicaragua.
auritum, *Schott, in Wien. Zeitschr.* iii. (1829) 780; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Mexic.
decipiens, Schott, Syn. Aroid. 69 = Vellozianum.
gracile, Schott, l. c. 70 = affine.
Hoffmannii, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 178 = Porphyrospatha Hoffmannii.
neglectum, Schott, in Bonplandia, vii. (1859) 163 = auritum.
Oerstedianum, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 178 (Oerstedtianum).—Costa Rica.
peliocladum, *Schott, Prod. Aroid.* 202.—Costa Rica.
Plumieri, Schott, l. c. 206 = auritum.
podophyllum, *Schott, in Bot. Zeit.* ix. (1851) 85.—Mexic.
Poeppigii, Schott, Syn. Aroid. 68 = Vellozianum.
Riedelianum, Schott, l. c. 70 = Vellozianum.
Ruizii, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 418 = auritum.
salvadorense, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 178 = podophyllum.
Schottianum, Wendl. ex Schott, Prod. Aroid. 200 = Porphyrospatha Schottiana.
Vellozianum, *Schott, in Oestr. Bot. Wochenbl.* iv. (1854) 418.—Bras.
Wendlandii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 178.—Costa Rica.
Willdenowii, Schott, Syn. Aroid. 68 = Vellozianum.
xanthophidium, *Schott, Prod. Aroid.* 214.—Mexic.

SYNIMA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 501. *SAPINDACEAE*.
Cordierii, *Radlk. l. c.* 513, 539.—Austral.

SYNISOON, Baill. in Bull. Soc. Linn. Par. i. (1879) 208. *RUBIACEAE*.
Schomburgkianum, *Baill. l. c.*—Guian.

SYNMERIA, Nimmo, in J. Grah. Cat. Pl. Bomb. Addend. (1839) p. sine numero = **Habenaria**, Willd. (Orchid.).
schizochilus, Nimmo, l. c. = H. crinifera.

SYNNEMA, Benth. in DC. Prod. x. 538 (1846) = **Cardanthera**, Buch.-Ham. (Acanthac.).
avanum, Benth. l. c. = C. avana.

SYNNETIA, Sweet, Brit. Flow. Gard. Ser. I. t. 150 (1826), sphalm. = seq.

SYNNOTTIA, Baker, in Journ. Linn. Soc. xvi. (1877) 169, sphalm. = seq.

SYNNOTIA, Sweet, Brit. Flow. Gard. Ser. I. t. 150 (1826) (sphalm. Synnetia, in Ind. correct.). *IRIDEAE*, Benth. & Hook. f. iii. 709.
ANACTORION, Rafin. Fl. Tellur. iv. 34 (1836).
bicolor, *Sweet, Hort. Brit.* ed. I. 398; ed. II. 501.—Afr. austr.
galeata, *Sweet, ll. cc.*—Afr. austr.
variegata, *Sweet, Brit. Flow. Gard.* Ser. I. t. 150.—Afr. austr.
versicolor, Steud. Nom. ed. II. ii. 614, in syn., sphalm. = praec.
Walthami, Hort. Kew.—Cf. Gard. Chron. (1881) I. 370, nomen = variegata.

SYNODON, Rafin. Sylva Tellur. 95 (1838) = **Conostegia**, D. Don (Melastom.).
calyptrata, Rafin. l. c. = C. calyptrata.
montana, Rafin. l. c. = C. montana.

SYNOECIA, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 469 = **Ficus**, Tourn. (Urtic.).
diversifolia, Miq. l. c. 470. t. 9 A = F. diversifolia.
falcata, Miq. l. c. t. 9. B = F. punctata.
grandifolia, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 184 = F. oligoneura.
Guillielmi-primi, De Vriese, Synoecia, 1 = F. asperrima.
serpens, Miq. Pl. Jungh. 67 = F. punctata.
sumatrana, Miq. Fl. Ind. Bat. i. II. 329 = F. kallicarpa.

SYNOLIGA, Rafin. Fl. Tellur. ii. 15 (1836) = **Xyris**, Linn.
pauciflora, Rafin. l. c. = X. pauciflora.
subulata, Rafin. l. c. = X. subulata.

SYNOPLECTRIS, Rafin. Fl. Tellur. ii. 89 (1836) = **Neottia**, Linn. (Orchid.), etc.
picta, Rafin. l. c. 90 = Spiranthes picta.
viridis, Rafin. l. c.—Hab.?

SYNOPTERA, Rafin. Sylva Tellur. 98 (1838) = **Miconia**, Ruiz & Pav. (Melastom.).
levigata, Rafin. l. c. = M. laevigata.

SYNOSMA, Rafin. in Loud. Gard. Mag. viii. (1832) 247 = **Senecio**, Tourn. (Compos.).
reniformis, Rafin. l. c. 247 = Sen. Muehlenbergii.
suaveolens, Rafin. l. c. = Sen. suaveolens.

SYNOSTEMON, F. Muell. Fragm. i. 32 (1858) = **Phyllanthus**, Linn. (Euphorb.).
glaucus, F. Muell. l. c. 33 = P. Adami.
hirtellus, F. Muell. l. c. iii. 89 = P. hirtellus.
ramosissimus, F. Muell. l. c. i. 33 = P. ramosissimus.
rigens, F. Muell. l. c. ii. 153 = P. rigens.

SYNOUM, A. Juss. in Mém. Mus. Par. xix. (1830) 226. t. 15. *MELIACEAE*, Benth. & Hook. f. i. 335.
glandulosum, *A. Juss. l. c.* 227. t. 15.—Austral.
Muelleri, *C. DC. in DC. Monog. Phan.* i. 593.—Austral.

SYNPHYLLIUM, Griff. in Madr. Journ. Sc. Ser. I. iv. (1836) 373 = **Curanga**, Juss. (Scrophul.).
torenioides, Griff. l. c. = C. amara.

SYNPTERA, Llanos, Fragm. 98 (1851) = **Cleisostoma**, Blume (Orchid.).
subviolacea, Llanos, l. c. = C. subviolaceum.

SYNSIPHON, Regel, in Act. Hort. Petrop. vi. (1879) 490 = **Colchicum**, Linn. (Liliac.).
crociflorus, Regel, l. c. 491 = C. crociflorum, *Regel*.

SYNSTIMA, Rafin. Sylva Tellur. 48 (1838) = **Ilex**, Tourn.
acuminata, Rafin. l. c. 49 = I. ambigua.
ambigua, Rafin. ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 157 = I. ambigua.
caroliniana, Rafin. Sylva Tellur. 49 = I. ambigua.
rotundifolia, Rafin. l. c. = I. rotunda?

SYNTHERISMA, Walt. Fl. Carol. 76 (1788) = **Panicum**, Linn. (Gramin.).
aegyptiacum, Schult. ex Steud. Nom. ed. II. ii. 656 = Eleusine aegyptiaca.
ciliare, Schrad. Fl. Germ. 160 = P. sanguinale.
glabrum, Schrad. l. c. 163. t. 3. f. 6 = P. glabrum.
malabaricum, Sw. ex Schult. Syst. ii. 474 = Centotheca lappacea.
praecox, Walt. Fl. Carol. 76 = P. sanguinale.
sanguinale, Dulac, Fl. Hautes-Pyr. 77 = P. sanguinale.
serotinum, Walt. Fl. Carol. 76 = Paspalum serotinum.
villosum, Walt. l. c. 77 = P. filiforme?
vulgare, Schrad. Fl. Germ. 161 = P. sanguinale.

SYNTHLIPSIS, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 116. *CRUCIFERAE*, Benth. & Hook. f. i. 93.
Berlandieri, *A. Gray, ex Torr. Bot. Mex. Bound.* 34, in nota.—Mexic.
Greggii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 116.—N. Mexic.
heterochroma, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 321.—N. Mexic.

SYNTHYRIS, Benth. in DC. Prod. x. 454 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 963.
alpina, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 251.—Am. bor. occ.
Houghtoniana, *Benth. in DC. Prod.* x. 454.—Am. bor.
pinnatifida, *S. Wats. in Bot. King's Exp.* 227. t. 22.—Am. bor. occ.
plantaginea, *Benth. in DC. Prod.* x. 455.—Am. bor. occ.
reniformis, *Benth. l. c.* 454.—Am. bor. occ.
rotundifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 285.—Am. bor. occ.
rubra, *Benth. in DC. Prod.* x. 455.—Am. bor. occ.

SYNTRICHOPAPPUS, A. Gray, in Pacif. Rail. Rep. iv. 106 (1856). *COMPOSITAE*, Benth. & Hook. f. ii. 402.
Fremontii, *A. Gray, l. c.*—Calif.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 20.—Calif.

SYNTROPHE, G. Ehrenb. ex Muell. Arg. in Neue Denkschr. Schw. Ges. Zuerich, xvi. (1858) 227 = **Caylusea**, A. St. Hil. (Resed.).
canescens, G. Ehrenb. l. c. = C. canescens.
fruticosa, G. Ehrenb. l. c. 229 = C. canescens.

SYNZISTACHIUM, Rafin. Sylva Tellur. 89 (1838) = **Heliotropium**, Tourn. (Borag.).
peruvianum, Rafin. l. c. = H. peruvianum.

SYNZYGANTHERA, Ruiz & Pav. Prod. 137. t 50 (1794) = **Lacistema**, Sw.
purpurea, Ruiz & Pav. Syst. Veg. 273 = L. purpureum.

SYORHYNCHIUM, Hoffmgg. Verz. Pfl. 198 (1824), err. typ. = **Sisyrinchium**, Linn. (Irid.).

SYPHARISSA, Salisb. Gen. Pl. Fragm. 37 (1866) = **Urginea**, Steinh. (Liliac.).
exuviata, Salisb. l. c. 38 = U. exuviata.
filifolia, Salisb. l. c. = U. filifolia.
fragrans, Salisb. l. c. = U. fragrans.

SYPHOCAMPYLOS, Hort. Belg. ex Hook. Bot. Mag. t. 4505 (1850), sphalm. = **Siphocampylus**, Pohl (Campanul.).

SYPHOMERIS, Steud. Nom. ed. II. ii. 656 (1841), sphalm. = SIPHOMERIS, Boj. = **Grewia**, Juss. (Tiliac.).

SYRENIA, Andrz. ex DC. Syst. ii. 491 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 80.
angustifolia, Reichb. Fl. Germ. Excurs. 689 = Erysimum angustifolium, *Ehrh.*
Biebersteinii, Andrz. ex DC. Syst. ii. 493 = Erysimum cuspidatum.
cuspidata, Reichb. Fl. Germ. Excurs. 689 = Erysimum cuspidatum.
Ehrharti, Andrz. ex DC. Syst. ii. 492 (*Ehrarti*) = Erysimum angustifolium, *Ehrh.*
Ehrhartiana, Andrz. in Bess. Enum. Pl. Volh. 104 = praec.
goniocaula, Boiss. & Hohen. Diagn. Ser. I. viii. 24 = Erysimum goniocaulon.
Lamarckii, Andrz. ex DC. Syst. ii. 492 = sessiliflora.
latifolia, Andrz. ex Bess. Enum. Pl. Volh. 27 = Erysimum cuspidatum.
sessiliflora, *Ledeb. Fl. Ross.* i. 193.—Ross. mer.
siliculosa, *Andrz. ex DC. Syst.* ii. 491.—Sibir.

SYRENOPSIS, Jaub. & Spach, Illustr. i. 6. t. 3 (1842) = **Iberidella**, Boiss. (Crucif.).
stylosa, Jaub. & Spach, l. c.—As. Min.

SYRINGA, Linn. Syst. ed. I (1735). *OLEACEAE*, Benth. & Hook. f. ii. 675.
LIGUSTRINA, Rupr. Beitr. Pfl. Russ. Reich. xi. 55 (1859).
LILAC, Tourn. ex Adans. Fam. ii. 223 (1763).
LILIACUM, Renault, Fl. Dép. Orne, 100 (1800).
albiflora, Opiz, Natural. n. x. (1825) 239 = vulgaris.
amoena, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
amurensis, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 371.—Manchuria.
angustifolia, Salisb. Prod. 14 = persica.
bicolor, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
capitata, S. T. Gmel. Reise, iii. 304 = persica.
carlsruhensis, Hort. ex C. Koch, Dendrol, ii. 1. 266 = vulgaris.
chinensis, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 116 = oblata.
chinensis, Turcz. ex DC. Prod. viii. 283 = villosa.
chinensis, *Willd. Berl. Baumz.* 378.—China.
cordifolia, Stokes, Bot. Comment. i. 33 = vulgaris.
dubia, Pers. Syn. i. 9 = chinensis.
Emodi, *Wall. Cat.* n. 2831.—Reg. Himal.
Josikaea, *Jacq. f. in Flora*, xiv. (1831) 67, 399.—Transsylv.
laciniata, Mill. Gard. Dict. ed. VIII. n. 3 = persica.
latifolia, Salisb. Prod. 13 = vulgaris.
Libertii, Hort. ex Pasq. Cat. Orto Bot. Nap. 100, nomen = vulgaris ?
ligustrina, Hort. ex C. Koch, Dendrol. ii. 1. 272 = amurensis.
marliensis, Hort. l. c. 266 = vulgaris.
media, Dum.-Cours. Bot. Cult. ed. I. i. 709 = chinensis.
nigricans, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
Notgeri, Hort. l. c. = vulgaris.
oblata, *Lindl. ex Carr. in Fl. des Serres*, xiii. (1858) 126.—China.
pekinense, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 371 = amurensis.
persica, *Linn. Sp. Pl.* 9.—Reg. Cauc.; Persia.
Philemon, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
pubescens, Turcz. in Bull. Soc. Nat. Mosc. (1840) 73 = villosa.
rothomagensis, Hort. ex A. Rich. in Dict. Class. ix. 401 = chinensis.
rotundifolia, *Decne. in Nouv. Arch. Mus. Par.* Sér. II. ii. (1879) 44.—Manchuria.
speciosa, Hort. ex DC. Prod. viii. 282 = chinensis.
suspensa, Thunb. Fl. Jap. 19 = Forsythia suspensa.
versaliensis, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
villosa, *Vahl, Enum.* i. 38.—China.
vincetoxifolia, Baumg. ex Steud. Nom. ed. II. ii. 656 = S. Josikaea.
virginalis, Hort. ex C. Koch, Dendrol. ii. 1. 266 = vulgaris.
vulgaris, *Linn. Sp. Pl.* 9.—Transsylv.

SYRINGA, Tourn. ex Adans. Fam. ii. 244 (1763) = **Philadelphus**, Riv. (Saxifr.).
inodora, Moench, Meth. 678 = P. inodorus.
suaveolens, Moench, l. c. = P. coronarius.

SYRINGODEA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = **Erica**, Tourn.
abietina, D. Don, l. c. = E. abietina.
acuminata, Steud. Nom. ed. II. ii. 656 = E. Massoni.
ambigua, G. Don, Gen. Syst. iii. 824 = E. Massoni.
Archeriana, G. Don, l. c. 822 = E. Archeria.
argentiflora, G. Don, l. c. = E. purpurea.
aurea, G. Don, l. c. = E. aurea.
Bedfordiana, G. Don, l. c. 819 = E. pellucida.
Bedfordiana, G. Don, l. c. 821 = E. Massoni.
bicolor, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. bicolor.
Bowieana, G. Don, Gen. Syst. iii. 825 = E. Bowieana.
brachialis, G. Don, l. c. 820 = E. brachialis.
Broadleyana, G. Don, l. c. 818 = E. Broadleyana.
buccinaeformis, G. Don, l. c. 824 = E. buccinaeformis.
carinata, G. Don, l. c. 825 = E. Massoni.
Cephalotes, G. Don, l. c. 822 = E. Cephalotes.
cerviciflora, G. Don, l. c. 825 = E. cerviciflora.
clavaeflora, G. Don, l. c. 824 = E. clavaeflora.
clavata, G. Don, l. c. 818 = E. viridiflora.
coccinea, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. coccinea.
colorans, G. Don, Gen. Syst. iii. 821 = E. colorans.
concinna, G. Don, l. c. 823 = E. concinna.
conspicua, G. Don, l. c. 820 = E. conspicua.
coronata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. fascicularis.
costata, G. Don, Gen. Syst. iii. 818 = E. versicolor.
cruenta, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. cruenta.
curviflora, G. Don, Gen. Syst. iii. 819 = E. curviflora.
cuspidigera, G. Don, l. c. 824 = E. ignescens.
cyrillaeflora, G. Don, l. c. 821 = E. palustris.
densa, G. Don, l. c. 824 = E. ventricosa.
densiflora, Steud. Nom. ed. II. ii. 656, sphalm. = seq.
densifolia, G. Don, Gen. Syst. iii. 823 = E. Uhria.
diaphana, G. Don, l. c. 818 = E. transparens.
dichrus, G. Don, l. c. = E. dichrus.
Dickensonii, G. Don, l. c. 824 = E. pallens.
Dicksoniana, Steud. Nom. ed. II. ii. 656, sphalm. = praec.
discolor, G. Don, Gen. Syst. iii. 823 = E. discolor.
doliiformis, G. Don, l. c. 826 = E. blanda.
echiiflora, G. Don, l. c. 822 = E. coccinea.
elata, G. Don, l. c. 819 = E. conspicua.
elongata, G. Don, l. c. 826 = E. elongata.
epistomia, G. Don, l. c. 824 = E. sacciflora.
erubescens, G. Don, l. c. 819 = E. erubescens.
Ewerana, G. Don, l. c. 823 = E. Uhria.
exposita, G. Don, l. c. 826 = E. pellucida.
exsurgens, G. Don, l. c. 819 = E. exsurgens.
exudans, G. Don, l. c. 819 = E. pellucida.
fascicularis, G. Don, l. c. 825 = E. fascicularis.
filamentosa, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. filamentosa.
flammea, G. Don, Gen. Syst. iii. 819 = E. flammea.
flava, G. Don, l. c. = E. parilis.
foliacea, Benth. in DC. Prod. vii. 630, sphalm. = seq.
foliosa, G. Don, Gen. Syst. iii. 825 = E. foliacea.
gemmifera, G. Don, l. c. 820 = E. Massoni.
gilva, G. Don, l. c. 825 = E. gilva.
glabra, Steud. Nom. ed. II. ii. 656 = E. ventricosa.
glandulosa, G. Don, Gen. Syst. iii. 820 = E. pellucida.
grandiflora, G. Don, l. c. 822 = E. grandiflora.
Hibbertiana, G. Don, l. c. = E. Hibbertia.
hirsuta, G. Don, l. c. 826 = E. perspicua.
hirta, G. Don, l. c. 823 = E. speciosa.
hybrida, G. Don, l. c. 824 = E. Massoni.
ignescens, G. Don, l. c. 820 = E. ignescens.
insulsa, G. Don, l. c. 818 = E. insulsa.
lanata, G. Don, l. c. 823 = E. conspicua.
Leeana, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. Leea.
Linkii, G. Don, Gen. Syst. iii. 819 = E. protrudens.
Linnaeana, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. perspicua.

SYRINGODEA:—
linnaeoides, G. Don, Gen. Syst. iii. 819 = Erica perspicua.
lituiflora, G. Don, l. c. 820 = E. perspicua.
longifolia, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. longifolia.
lycopodiifolia, D. Don, ex Steud. Nom. ed. II. ii. 656 = E. Massoni.
mammosa, G. Don, Gen. Syst. iii. 825 = E. mammosa.
Massonii, G. Don, l. c. 820 = E. Massoni.
mutabilis, G. Don, l. c. 824 = E. cruenta.
nana, G. Don, l. c. = E. nana.
nitens, G. Don, l. c. = E. nitens.
obpyramidalis, G. Don, l. c. 821 = E. pyramidalis.
onosmaeflora, G. Don, l. c. 822 = E. onosmaeflora.
ostrina, G. Don, l. c. 824 = E. coccinea.
pallens, G. Don, l. c. 818 = E. pallens.
Patersonii, G. Don, l. c. 825 = E. abietina.
patersonioides, G. Don, l. c. = E. Massoni.
pauciflora, Steud. Nom. ed. II. ii. 656, sphalm. = E. sacciflora.
pellucida, G. Don, Gen. Syst. iii. 818 = E. pellucida.
perspicua, G. Don, l. c. 821 = E. Massoni.
phylicifolia, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. purpurea.
pinea, G. Don, Gen. Syst. iii. 821 = E. vestita.
pinifolia, G. Don, l. c. = E. longifolia.
procera, G. Don, l. c. 820 = E. ignescens.
pulchella, G. Don, l. c. 823 = E. Leea.
pulviniformis, G. Don, l. c. = E. coccinea.
pura, G. Don, l. c. = E. Leea.
purpurea, G. Don, l. c. = E. purpurea.
radiata, G. Don, l. c. 819 = E. Massoni.
refulgens, G. Don, l. c. 818 = E. versicolor.
Rollinsonii, G. Don, l. c. 826 = E. Rollissonia.
rosea, G. Don, l. c. 822 = E. vestita.
rubida, G. Don, l. c. 826 = E. Massoni.
rugosa, G. Don, l. c. 823 = E. Massoni.
sacciflora, G. Don, l. c. 825 = E. sacciflora.
Salisburiana, G. Don, l. c. 822 = E. purpurea.
sanguinea, G. Don, l. c. 824 = E. sanguinea.
sceptriformis, G. Don, l. c 826 = E. spicata.
serratifolia, G. Don, l. c. 820 = E. serratifolia.
sessiliflora, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. spicata.
simpliciflora, G. Don, Gen. Syst. iii. 820 = E. curviflora, buccinaeformis.
sordida, G. Don, l. c. = E. conspicua.
speciosa, G. Don, l. c. 824 = E. speciosa.
spicata, G. Don, l. c. 825 = E. spicata.
splendens, G. Don, l. c. 821 = E. conspicua.
spuria, G. Don, l. c. 819 = E. Massoni.
stagnalis, G. Don, l. c. = E. sulphurea.
sulphurea, G. Don, l. c. = E. sulphurea.
transparens, G. Don, l. c. 818 = E. elongata.
triphylla, G. Don, l. c. = E. triphylla.
tubiflora, G. Don, l. c. 820 = E. tubiflora.
tumida, G. Don, l. c. 825 = E. tumida.
velitaris, G. Don, l. c. = E. velitaris.
versicolor, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. versicolor.
verticillata, G. Don, Gen. Syst. iii. 821 = E. mammosa.
vestita, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. vestita.
virescens, G. Don, Gen. Syst. iii. 824 = E. virescens.
viridiflora, G. Don, l. c. = E. viridiflora.
viridis, G. Don, l. c. 822 = E. Leea.

SYRINGODEA, Hook. f. Bot. Mag. t. 6072 (1873). *IRIDEAE*, Benth. & Hook. f. iii. 693.
bicolor, *Baker, in Journ. Bot.* xiv. (1876) 67.—Afr. austr.
filifolia, *Baker, l. c.*—Afr. austr.
latifolia, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 403.—Afr. austr.
minuta, Klatt, l. c. = Pauridia hypoxidioides.
montana, *Klatt, l. c.*—Afr. austr.
pulchella, *Hook. f. Bot. Mag.* t. 6072.—Afr. austr.
rosea, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 403 = Romulea rosea.

SYRINGOSMA, Mart. ex Reichb. Consp. 134 (1828); et ex Lindl. Veg. Kingd. 601 (1847) = **Forsteronia**, G. W. F. Mey. (Apocyn.).

SYRIUM, Steud. Nom. ed. II. ii. 657 (1841), sphalm. = SIRIUM, Schreb. = **Santalum**, Linn.

SYRMATIUM, Vog. in Linnaea, x. (1835-36) 590 = **Hosackia**, Dougl. (Legumin.).
glabrum, Vog. l. c. 591 = H. glabra.
tomentosum, Vog. l. c. = H. tomentosa.

SYRRHONEMA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 124; xx. (1867) 19. Genus dubium *MENISPERMACEARUM*. Cf. Benth. & Hook. f. i. 959.
fasciculatum, *Miers, ll. cc.* 124; 20.—Ins. Fernando Po.

SYSIMBRIUM, Pall. Reise, ii. 740 (1774), err. typ. = **Sisymbrium**, Tourn. (Crucif.).

SYSIRINCHIUM, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 263, sphalm. = **Sisyrinchium**, Linn. (Irid.).

SYSPONE, Griseb. Spicil. Fl. Rumel. i. 5 (1843) = **Genista**, Linn. (Legumin.).
sagittalis, Griseb. l. c. 6 = G. sagittalis.

SYSTEMON, Regel, Ind. Sem. Hort. Petrop. (1856) 38; Gartenfl. vi. (1857) 81 = **Galipea**, Aubl. (Rutac.).
Fischeri, Regel, ll. cc.—Bras.

SYSTREPHA, Burch. Trav. i. 546 (1822) = **Ceropegia**, Linn. (Asclep.).
filiforme, Burch. l. c.—Afr. austr.
multiflora, Burch. ex Baker, in Saund. Refug. Bot. t. 10 = C. multiflora.

SYSTREPHIA, Benth. & Hook. f. Gen. ii. 779 (1876), sphalm. = praec.

SYZIGANTHUS, Steud. Syn. Pl. Cyp. 153 (1855) = **Gahnia**, Forst. (Cyperac.).
multiflorus, Steud. l. c. = G. javanica.

SYZIGIUM, Steud. Nom. ed. II. ii. 657 (1841), sphalm. = seq.

SYZYGEUM, Wight, Ic. i. 4 (1838), sphalm. = seq.

SYZYGIUM, Gaertn. Fruct. i. 166. t. 33 (1788) = **Eugenia**, Mich. (Myrtac.).
acuminatissimum, DC. Prod. iii. 261.—Java.
acuminatum, Miq. Fl. Ind. Bat. i. 1. 452 = E. acuminata.
aegiceroides, Korth. in Nederl. Kruidk. Arch. i. (1848) 203 = E. aegiceroides.
alternans, Miq. ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 500 = E. Heyneana.
alternifolium, Walp. Rep. ii. 179 = E. alternifolia.
altissimum, Wall. Cat. n. 3588 = E. Thumra.
amicorum, C. Muell. in Walp. Ann. iv. 839 = E. amicorum.
androsaemoides, Walp. Rep. ii. 179 = E. androsaemoides.
angkolanum, Miq. Fl. Ind. Bat. i. 1. 448 = E. operculata.
apodum, Miq. l. c. Suppl. 312.—Sumatra.
areolatum, DC. Prod. iii. 260 = E. areolata.
Arnottianum, Walp. Rep. ii. 180 = E. Arnottiana.
assimile, Thw. Enum. Pl. Zeyl. 116 = E. assimilis.
auriculatum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
avene, Miq. Fl. Ind. Bat. Suppl. 312.—Sumatra.
balsameum, Wall. Cat. n. 3592 = E. balsamea.
Bellutta, DC. Prod. iii. 261 = E. zeylanica.
borneense, Miq. Fl. Ind. Bat. i. 1. 453 = E. borneensis.
brachiatum, Miq. l. c. 460 = E. brachiata.
brachybotryum, Miq. l. c. 452.—Sumatra.
brachynemum, F. Muell. Fragm. iv. 59 = E. Smithii.
Brackenridgei, C. Muell. in Walp. Ann. iv. 838 = E. Brackenridgei.
brevifolium, C. Muell. l. c. 839 = E. brevifolia.
buxifolium, Hook. & Arn. Bot. Beech. Voy. 187.—China.
calophyllifolium, Walp. Rep. ii. 180 = E. calophyllifolia.
campanellum, Miq. Fl. Ind. Bat. i. 1. 451.—Java.
campanulatum, Korth. in Nederl. Kruidk. Arch. i. (1848) 203.—Borneo.
capillare, Wall. Cat. n. 3578 = E. filiformis, *Wall.*
caryophyllaeum, Gaertn. Fruct. i. 166. t. 33 = E. caryophyllaea.
caryophyllifolium, DC. Prod. iii. 260 = E. Jambolana.

SYZYGIUM:—
caryophyllifolium, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 449 = *Zippelianum*.
caudatum, Wall. Cat. n. 3591 = Eugenia cymosa.
cinerascens, Presl, Bot. Bemerk. 70 = seq.
cinereum, Wall. Cat. n. 3576 = E. brachiata.
claviflorum, Wall. l. c. 3575 = E. claviflora.
clusiaefolium, C. Muell. in Walp. Ann. iv. 839 = E. clusiaefolia.
coarctatum, Blume, ex Miq. Fl. Ind. Bat. i. 1. 437 = E. zeylanica.
concinnum, Wall. Cat. n. 3582 = E. cymosa.
confertiflorum, C. Muell. in Walp. Ann. iv. 838 = E. confertiflora (cf. Addend.).
confusum, Blume, ex Miq. Anal. Bot. Ind. i. 27 = E. microbotrya.
conicum, Korth. in Nederl. Kruidk. Arch. i. (1848) 204.—Borneo.
contractum, Wall. Cat. n. 3602 = E. pellucida.
cordatum, Hochst. in Flora, xxvii. (1844) 425 = E. cordatum, *M. Laws.*
cordifolium, Klotzsch, in Peters, Reise Mossamb. Bot. 63. t. 11 = E. cordatum, *M. Laws.*
cordifolium, Walp. Rep. ii. 179 = E. cordifolia.
corticatum, Wall. Cat. n. 3565 = E. zeylanica.
corymbosum, DC. Prod. iii. 261.—Java.
corynocarpa, C. Muell. in Walp. Ann. iv. 839 = E. corynocarpa.
costatum, Miq. Fl. Ind. Bat. i. 1. 451 = E. operculata.
costatum, Wall. Cat. n. 3556 (Quid?).—Ind. or.
cuneatum, Wall. l. c. 3598 = E. cuneata.
cuneifolium, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 144 = E. cuneifolia.
cyminiferum, Presl, Bot. Bemerk. 70 (= *Jambosa cymifera*).—Afr. austr.
cymosum, DC. Prod. iii. 259 = E. cymosa.
cymosum, Korth. in Nederl. Kruidk. Arch. i. (1848) 202 = E. pamatensis, operculata.
densiflorum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 182.—N. Caled.
densiflorum, Wall. Cat. n. 3601 = E. Arnottiana.
effusum, C. Muell. in Walp. Ann. iv. 838 = E. effusa.
ellipticum, Wall. Cat. n. 3587 = E. valdevenosa.
euneuron, Miq. Fl. Ind. Bat. Suppl. 314.—Sumatra.
excavatum, Wall. Cat. n. 3574 = E. claviflora.
expansum, Wall. l. c. 3567 = E. expansa, *Duthie.*
ficifolium, Wall. l. c. 3558 = E. tetragona.
filiforme, Wall. l. c. 3580 = E. filiformis.
firmum, Thw. Enum. Pl. Zeyl. 417 = E. grandis.
floribundum, F. Muell. Fragm. iv. 58 = E. Ventenatii.
forte, F. Muell. l. c. v. 13 = E. grandis.
fraternum, Miq. Fl. Ind. Bat. i. 1. 458.—Java.
frutescens, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
fruticosum, DC. Prod. iii. 260 = E. fruticosa.
furcatum, Blume, ex Miq. Anal. Bot. Ind. i. 23 = E. Muelleri.
Gardneri, Thw. Enum. Pl. Zeyl. 117 = E. Gardneri.
glaucescens, Miq. Anal. Bot. Ind. i. 24 = E. borneensis.
glomeratum, DC. Prod. iii. 259 = E. glomerata, *Lam.*
grande, Wall. Cat. n. 3554 = E. grandis.
gratum, Wall. l. c. 3586 = E. grata.
guineense, DC. Prod. iii. 259 = Calyptranthes guineensis.
guineense, Guill. & Perr. Fl. Seneg. Tent. 315. t. 72 = E. owariensis.
Heyneanum, Wall. Cat. n. 3599 = E. Heyneana.
hypericifolium, DC. Prod. iii. 261.—Ins. Borbon.
inophylloides, C. Muell. in Walp. Ann. iv. 838 = E. inophylloides.
inophyllum, DC. Prod. iii. 260 = E. inophylla.
Jambolanum, DC. l. c. 259 = E. Jambolana.
javanicum, Miq. Fl. Ind. Bat. i. 1. 461.—Java.
kalahiense, Korth. in Nederl. Kruidk. Arch. i. (1848) 205 = E. kalahiensis.
Korthalsianum, Miq. Fl. Ind. Bat. i. 1. 454 = E. Korthalsiana.
laevigatum, Miq. l. c. 457 = E. laevigata.
lanceolatum, Wight & Arn. Prod. 330 = E. Wightiana.
lateriflorum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 183.—N. Caled.
lateriflorum, Royle, ex Duthie, in Hook. f. Fl. Brit. Ind. ii. 499 = E. Jambolana.
latifolium, Blanco, Fl. Filip. ed. II. 294 = E. bracteata.
latifolium, DC. Prod. iii. 259.—Ins. Maurit.
laxiflorum, DC. l. c. 261 = Calyptranthes laxiflora.
leucoxylon, Korth. in Nederl. Kruidk. Arch. i. (1848) 203 = E. leucoxylon.

SYZYGIUM:—
lineare, Wall. Cat. n. 3596 = Eugenia linearis, *Duthie.*
lissophyllum, Thw. Enum. Pl. Zeyl. 117 = E. lissophylla.
longiflorum, Presl, Bot. Bemerk. 70.—Ins. Philipp.
longiflorum, Wall. Cat. n. 3572 = E. claviflora.
lucidum, Gaertn. Fruct. i. 167. t. 33 = E. lucida, *Banks.*
macranthum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 182.—N. Caled.
magnoliaefolium, DC. Prod. iii. 261.—Java.
magnoliaefolium, Korth. in Nederl. Kruidk. Arch. i. (1848) 203 = E. borneensis.
Makul, Gaertn. Fruct. i. 166. t. 33.—Zeylan.
marginatum, Korth. in Nederl. Kruidk. Arch. i. (1848) 203.—Sumatra.
Metrosideros, DC. Prod. iii. 261.—Cochinch.
micans, Brongn. & Gris, in Bull. Soc. Bot. Fr. xiii. (1866) 468.—N. Caled.
Michelii, Duthie, in Hook. f. Fl. Brit. Ind. ii. 505 = E. uniflora, *Linn.*
micranthum, Blume, ex Miq. Anal. Bot. Ind. i. 22 = E. pamatensis.
micranthum, Thw. Enum. Pl. Zeyl. 117 = E. micrantha, *Bedd.*
minutiflorum, Miq. Fl. Ind. Bat. Suppl. 311.—Sumatra.
montanum, Thw. Enum. Pl. Zeyl. 116 = E. grandis.
Muelleri, Miq. Fl. Ind. Bat. i. 1. 453 = E. Muelleri.
multipetalum, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 182.—N. Caled.
myrtifolium, DC. Prod. iii. 261 = E. myrtifolia, *Roxb.*
Neesianum, Arn. in Nov. Act. Nat. Cur. xviii. 1. (1836) 335 = E. Neesiana.
neglectum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
nelitricarpum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 53 = E. cymosa.
nervosum, DC. Prod. iii. 260 = E. operculata.
nitidum, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 221.—N. Guin.
nitidum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 183.—N. Caled.
nodosum, Miq. Fl. Ind. Bat. i. 1. 447 = E. operculata.
oblatum, Wall. Cat. n. 3569 = E. oblata.
obovatum, DC. Prod. iii. 259 = E. obovata, *Poir.*
obovatum, Korth. in Nederl. Kruidk. Arch. i. (1848) 205 = E. Muelleri.
obovatum, Wall. Cat. n. 3552 = E. operculata.
obtusatum, Blume, ex Miq. Anal. Bot. Ind. i. 25 = E. caryophyllaea.
obtusifolium, Kostel. Allg. Med.-Pharm. Fl. iv. 1532 = E. Jambolana.
occlusum, Miq. Fl. Ind. Bat. i. 1. 460 = E. occlusa.
odoratum, DC. Prod. iii. 260.—China.
oleinum, Wall. ex Walp. Rep. ii. 178 = E. myrtifolia.
oliganthum, Thw. Enum. Pl. Zeyl. 118 = E. oligantha.
ovale, Korth. in Nederl. Kruidk. Arch. i. (1848) 205.—Borneo.
owariense, Benth. in Hook. Niger Fl. 359 = E. owariensis.
palembanicum, Miq. Fl. Ind. Bat. Suppl. 313 = E. lepidocarpa.
Pancheri, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 183.—N. Caled.
Panealla, Wall. Cat. n. 3557 = E. operculata.
paniculatum, Gaertn. Fruct. i. 167. t. 33 = E. paniculata.
patens, Korth. in Nederl. Kruidk. Arch. i. (1848) 203.—Sumatra.
patens, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
politum, Wall. Cat. n. 3626 = E. zeylanica.
polyanthum, Miq. Fl. Ind. Bat. i. 1. 450.—Java.
polyanthum, Thw. Enum. Pl. Zeyl. 116 = E. operculata.
polyanthum, Walp. Rep. ii. 180 = E. polyantha, *Wight.*
Pseudo-jambolana, Miq. Fl. Ind. Bat. i. 1. 458.—Java.
pterocalyx, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. xiii. (1870-71) 386.—N. Caled.
pulchellum, Wall. Cat. n. 3566 = E. oblata.
punctulatum, Wall. l. c. 3583 = E. pyrifolia, *Duthie.*
pyrifolium, DC. Prod. iii. 261 = E. pyrifolia, *Duthie.*
quadrialatum, Teijsm. & Binn. Cat. Hort. Bog. 246.—Ins. Molucc.
racemosum, DC. Prod. iii. 261.—Java.
rameum, Wall. Cat. n. 3595 = E. tetragona.
reticulatum, Walp. Rep. ii. 179 = E. mangifolia.
revolutum, Walp. l. c. 180 = E. revoluta, *Wight.*

SYZYGIUM :—
ribesioides, Wall. Cat. n. 3553 A = Eugenia brachiata.
rigidum, Wall. l. c. 3581 = E. chlorantha.
robustum, Miq. Fl. Ind. Bat. i. 1. 1086.—Ins. Banca.
rostratum, DC. Prod. iii. 261.—Java.
rotundifolium, Arn. in Nov. Act. Nat. Cur. xviii. 1. (1836) 335 = E. rotundifolia.
rubens, Walp. Rep. ii. 180 = E. rubens.
rubescens, C. Muell. in Walp. Ann. iv. 839 = E. rubescens.
rubicundum, Wight & Arn. Prodr. 330 = E. rubicunda.
rugosum, Korth. in Nederl. Kruidk. Arch. i. (1848) 204.—Java.
salicifolium, J. Grah. Cat. Pl. Bomb. 73 = E. Heyneana.
salicifolium, Wall. Cat. n. 3597 = E. rubicunda.
sandwicensis, C. Muell. in Walp. Ann. iv. 838 = E. sandwicensis.
savaiensis, C. Muell. l. c. 839 = E. savaiensis.
scandens, Boj. Hort. Maurit. 143 = E scandens.
Schwenckii, Teijsm. & Binn. Cat. Hort. Bog. 246.—Sumatra.
sclerophyllum, Thw. Enum. Pl. Zeyl. 118 = E. sclerophylla.
scoparium, Wall. Cat. n. 3594 = E. scoparia.
sessile, Blume, ex Miq. Anal. Bot. Ind. i. 25 = E. Korthalsiana.
Sieberianum, Presl, Bot. Bemerk. 69.—Ins. Maurit.
spathulatum, Thw. Enum. Pl. Zeyl. 118 = E. olivifolia.
speciosum, Wall. Cat. n. 3568 = E. Thumra.
spicatum, DC. Prod. iii. 260 = E. zeylanica.
suavissimum, Wall. Cat. n. 3573 = E. leptantha, *Wight*.
subcapitulatum, Miq. Fl. Ind. Bat. i. 1. 450.—Java.
subdecurrens, Miq. l. c. 449.—Java.
subdecussatum, Wall. Cat. n. 3589 = E. subdecussata.
subnodosum, Miq. Fl. Ind. Bat. Suppl. 313.—Sumatra.
subtile, Miq. l. c.—Sumatra.
sylvestre, Walp. Rep. ii. 178 = E. sylvestris.
Tatna, Wall. Cat. n. 3555 = E. Tatna (in Addend.).
tenellum, Blume, ex Miq. Anal. Bot. Ind. i. 27.—Borneo.
tenue, Wall. Cat. n. 3570 = E. tenuis.
tenuiflorum, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 183.—N. Caled.
tesselatum, Korth. in Nederl. Kruidk. Arch. i. (1848) 203.—Borneo.
tetragonum, Wall. ex Wight, Illustr. Ind. Bot. ii. 16 = E. tetragona.
timorianum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 455.—Ins. Timor.
toddalioides, Walp. Rep. ii. 179 = E. toddalioides.
triplinervium, Teijsm. & Binn. Cat. Hort. Bog. 246.—Java.
truncatum, Miq. Fl. Ind. Bat. i. 1. 455.—Java.
tutuilense, C. Muell. in Walp. Ann. iv. 839 = E. tutuilensis.
umbellatum, Korth. in Nederl. Kruidk. Arch. i. (1848) 205.—Sumatra.
umbrosum, Thw. Enum. Pl. Zeyl. 118 = E. subavenis.
validum, Korth. in Nederl. Kruidk. Arch. i. (1848) 204.—Sumatra.
variifolium, Miq. Fl. Ind. Bat. Suppl. 311.—Ins. Banca.
vastum, Wall. Cat. n. 3561 = E. operculata.
venosum, DC. Prod. iii. 260 = E. frondosa.
venulosum, Wall. Cat. n. 3585 = E. venulosa.
verecundum, Wall. l. c. 3579 = E. verecunda.
vimineum, Wall. l. c. 3593 = E. cymosa.
wagapense, Brongn. & Gris, in Bull. Soc. Bot. Fr. xiii. (1866) 468.—N. Caled.
Wallichianum, Presl, Bot. Bemerk. 70 = E. operculata.
Wallichii, Walp. Rep. ii. 180 = E. Wallichii.
Wightianum, Wall. Cat. n. 3577 = E. Wightiana.
zeylanicum, DC. Prod. iii. 260 = E. zeylanica, *Wight*.
Zippelianum, Miq. Fl. Ind. Bat. i. 1. 449.—Java.

SZEGLEWIA, C. Muell. in Walp. Ann. vii. 419 (1868), sphalm. = SCZEGLEEWIA, Turcz. = **Symphorema**, Roxb. (Verben.).

SZOVITSIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 39 1835). *UMBELLIFERAE*, Benth. & Hook. f. i. 925
callicarpa, *Fisch. & Mey. l. c.*—Reg. Cauc.

SZOWITSIA, Steud. Nom. ed. II. ii. 658 (1841) = praec.

T

TABACINA, Reichb. Handb. 201 (1837) = seq.

TABACUM, Gilib. Fl. Lituan. i. 39 (1781) = **Nicotiana**, Linn. (Solan.).
latissimum, Bercht. & Opiz, Oek. Fl. Boehm. iii. 309 = N. Tabacum.
Nicotianum, Bercht. & Opiz, l. c. 307 = N. Tabacum.
ovatofolium, Gilib. Fl. Lituan. i. 39 = N. Tabacum.

TABACUS, Moench, Meth. 448 (1794) = **Nicotiana**, Linn. (Solan.).
viscidus, Moench, l. c. = N. glutinosa.

TABEBUIA, Gomez, Obs. ii. 7. t. 2 (1803). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1042.
POTAMOXYLON, Rafin. Sylva Tellur. 78 (1838).
aesculifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 494.—Mexic.
Avellanedae, *Lorentz, ex Griseb. in Goett. Abh.* xxiv. (1879) 258.—Reg. Argent.
brevipes, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 199, nomen.—Bras.
cassinoides, *DC. Prod.* ix. 213.—Bras.
citrifolia, *DC. in Bibl. Univ. Genèv.* xvii. (1838) 131; *Prod.* ix. 213.—Bras.
cordata, *Benth. Bot. Voy. Sulph.* 129.—Am. austr.
dentata, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 199.—Bras.
dracocephaloides, *Miers, l. c.*—Bras.
fallax, DC. in Bibl. Univ. Genèv. xvii. (1838) 131 = Bignonia fallax.
flavescens, *Benth. & Hook. f. ex Griseb. in Goett. Abh.* xxiv. (1879) 257.—Reg. Argent.
fluviatilis, *DC. Prod.* ix. 215.—Guian.
fuscata, *Hemsl. Biol. Centr. Am. Bot.* ii. 494.—Mexic.
Guayacan, *Hemsl. l. c.* 495.—Panama.
haemantha, *DC. Prod.* ix. 214.—Porto Rico.
heterotricha, *Hemsl. Biol. Centr. Am. Bot.* ii. 495.—S. Salvador.
ilicifolia, *DC. Prod.* ix. 213.—Madag.
japurensis, *DC. l. c.* 214.—Bras.
lanceolata, *DC. in Bibl. Univ. Genèv.* xvii. (1838) 131; *Prod.*.ix. 213.—Bras.
latifolia, DC. ll. cc. = Callichlamys riparia.
leucoxyla, *DC. ll. cc.* 131; 212.—Am. trop.
magnolioides, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 199.—Bras.
Mansoana, *DC. Prod.* ix. 214.—Bras.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 495.—Mexic.
neurophylla, *Miq. in Linnaea*, xxvi. (1853) 219.—Guian.
nodosa, *Griseb. in Goett. Abh.* xxiv. (1879) 258.—Reg. Argent.
pallida, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 199.—Ind. occ.
palustris, *Hemsl. Biol. Centr. Am. Bot.* ii. 495.—Panama.
pentaphylla, *Hemsl. l. c.*—Panama.
Pisoniana, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 199.—Bras.
pyramidata, *DC. Prod.* ix. 214.—Guian.
rosea, *DC. l. c.* 215.—Mexic.
rufinervis, DC. l. c. 213 = Callichlamys riparia.
sanguinea, DC. in Bibl. Univ. Genèv. xvii. (1838) 131; Prod. ix. 176 = Fredericea speciosa.
triphylla, *DC. Prod.* ix. 214.—Ind. occ.
uliginosa, *DC. in Bibl. Univ. Genèv.* xvii. (1838) 131; *Prod.* ix. 212.—Bras.
xanthophylla, *DC. Prod.* ix. 214.—Bras.

TABERNA, Miers, Apocyn. S. Am. 61 (1878) = **Tabernaemontana**, Plum. (Apocyn.).
cymosa, Miers, l. c. 62 = Tabernaem. cymosa.
discolor, Miers, l. c. = Tabernaem. discolor.
disparifolia, Miers, l. c. 63.—Peruv.
disticha, Miers, l. c. 64 = Tabernaem. disticha.
laurina, Miers, l. c. 63 = Tabernaem. laurifolia.
Poeppigii, Miers, l. c. = Tabernaem. Poeppigii.
Riedelii, Miers, l. c. 64 = Tabernaem. Riedelii.

TABERNAEMONTANA, Plum. ex Linn. Gen. ed. i. 66 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 706.
ANACAMPTA, Miers, Apocyn. S. Am. 64 (1878).
ANARTIA, Miers, l. c. 79 (1878).
BONAFOUSIA, A. DC. in DC. Prod. viii. 359 (1844).
CODONEMMA, Miers, Apocyn. S. Am. 72 (1878).
MERIZADENIA, Miers, l. c. 78 (1878).
PANDACA, Thou. Gen. Nov. Madag. 10 (1806).
PESCHIERA, A. DC. in DC. Prod. viii. 360 (1844).
REICHARDIA, Dennst. Schluess. Hort. Malab. 32 (1818).
REJOUA, Gaudich. in Freyc. Voy. Bot. 450. t. 61 (1826).
RHIGOSPIRA, Miers, Apocyn. S. Am. 67 (1878).
TABERNA, Miers, l. c. 61 (1878).
acapulcensis, *Miers, l. c.* 57.—Mexic.
accedens, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 107.—Bras.
acuminata, *Muell. Arg. in Linnaea*, xxx. (1859–60) 406.—Bolivia.
acutissima, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 73.—Bras.
affinis, *Muell. Arg. l. c.* 83.—Bras.
africana, *A. DC. in DC. Prod.* viii. 367.—Afr. trop.
alba, Mill. Gard. Dict. ed. VIII. n. 2 = citrifolia.
alba, Nicholson, ex Griseb. Fl. Brit. Pl. Ind. 409 = citrifolia.
alternifolia, Linn. Sp. Pl. 211 = coronaria ?
alternifolia, Roxb. Fl. Ind. ii. 24 = crispa.
Amsonia, Linn. Sp. Pl. ed. II. 308 = Amsonia Tabernaemontana.
amygdalaefolia, *Jacq. Enum. Pl. Carib.* 14.—Am. austr.
amygdalifolia, Seem. Bot. Voy. Herald, 167 = acapulcensis.
amygdalifolia, Sieber, ex A. DC. in DC. Prod. viii. 369 = persicariaefolia.
angulata, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 72.—Bras.
angustifolia, Ait. Hort. Kew. ed. I. i. 300 = Amsonia angustifolia.
apoda, *Wright, in Sauv. Fl. Cub.* 118.—Cuba.
arcuata, *Ruiz & Pav. Fl. Per.* ii. 22. t. 143.—Peruv.
aurantiaca, *Gaudich. Freyc. Voy. Bot.* 50, 55.—Ins. Rawak.
australis, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 84.—Bras.
Barteri, *Hook. Bot. Mag.* t. 5859.—Afr. trop.
Benthamiana, Muell. Arg. in Mart. Fl. Bras. vi. 1. 70 = Muelleriana.
Benthamiana, *Muell. Arg. l. c.* 80.—Bras.
Benthaminina, Miers, Apocyn. S. Am. 289, sphalm. = praec.
Berterii, *A. DC. in DC. Prod.* viii. 367.—Porto Rico.
bicolor, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 952, nomen.—Guian.
bovina, *Lour. Fl. Cochinch.* 118.—Cochinch.
bracteolaris, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 83.—Bras.
breviflora, *Muell. Arg. l. c.* 79.—Bras.
bufalina, *Lour. Fl. Cochinch.* 117.—Cochinch.
calycina, Spruce, ex Miers, Apocyn. S. Am. 73 = Muelleriana.
calycina, *Wall. Cat.* n. 1577; *et in Bot. Reg.* sub t. 1273.—Burma.
Camassi, Regel, Cat. Pl. Hort. Aksakov. 137 = Gonioma Kamassi.
catharinensis, *A. DC. in DC. Prod.* viii. 365.—Bras.
celebica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 139.—Ins. Celebes.
cerifera, *Panch. & Sebert, Not. Bois Nouv. Calèd.* 188.—N. Caled.
cerniflora, *Zipp. ex Span. in Linnaea*, xv. (1841) 325.—Ins. Timor.
cestroides, Nees, ex Mart. in Nov. Act. Nat. Cur. xi. 1. (1823) 83 = Malouetia cestroides.
cirrhosa, Blanco, Fl. Filip. ed. I. 115 = Finlaysonia obovata.
citrifolia, Forst. f. Prod. 20 = orientalis.
citrifolia, *Linn. Sp. Pl.* 210.—Ind. occ.; Mexic.
citrifolia, Lunan, Fl. Jamaic. ii. 222 = discolor.
coffeaefolia, Boj. ex Baker, Fl. Maurit. 224, sphalm. = seq.
coffeoides, *Boj. ex A. DC. in DC. Prod.* viii. 370.—Ins. Mascar. et Johann.
Collignonae, Heurck & Muell. Arg. in Heurck, Obs. Bot. 170 = graciliflora.

TABERNAEMONTANA :—

collina, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 178.—Bras.
congesta, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 71 = rubrostriolata.
coriacea, *Link, ex Roem. & Schult. Syst.* iv. 431.—Bras.
coronaria, *Willd. Enum. Hort. Berol.* 275.—Ind. or.
coronata, Paxt. Mag. Bot. xvi. (1848) t. 354, sphalm. = praec.
corymbosa, *Roxb. Hort. Beng.* [84]; *Fl. Ind.* ii. 25; *et ex Wall. in Bot. Reg.* sub t. 1273.—Ind. or.
crassa, *Benth. in Hook. Niger Fl.* 447.—Afr. trop.
crispa, *Roxb. Hort. Beng.* 20; *Fl. Ind.* ii. 24.—Ins. Andaman.
Cumingiana, *A. DC. in DC. Prod.* viii. 373.—Ins. Philipp.
cylindracea, Wall. Cat. n. 4451 = Rauwolfia serpentina.
cylindrica, Steud. Nom. ed. II. ii. 658 = praec.
cymosa, *Jacq. Enum. Pl. Carib.* 14.—N. Granat.
cymosa, Mart. ex Miers, Apocyn. S. Am. 84 = Geissospermum Martianum.
cymulosa, *Miq. Fl. Ind. Bat.* ii. 422.—Sumatra.
Decaisnei, A. DC. in DC. Prod. viii. 369 = orientalis.
densiflora, Wall. in Bot. Reg. t. 1273 = Rauwolfia densiflora.
dichotoma, *Roxb. Hort. Beng.* 20; *Fl. Ind.* ii. 21.—Ind. or.
didyma, *Vell. Fl. Flum.* 106.—Bras.
discolor, *Sw. Prod. Veg. Ind. Occ.* 52.—Ind. occ.
disticha, *A. DC. in DC. Prod.* viii. 362.—Guian.
divaricata, R. Br. ex Roem. & Schult. Syst. iv. 427.—Cf. R. Br. in Mem. Wern. Soc. i. (1811) 72 = coronaria.
divaricata, G. Don, Gen. Syst. iv. 91 = coronaria.
ebracteata, R. Br. Prod. 468 = pubescens, *R. Br.*
echinata, *Aubl. Pl. Gui.* i. 263.—Guian.
echinata, Vell. Fl. Flum. 105; iii. t. 17 = T. Hystrix.
Echites, *Linn. Syst.* ed. X. 945.—Jamaic. ?
elastica, Spreng. Syst. i. 639 = Urceola elastica.
elliptica, *Blanco, Fl. Filip.* ed. I. 115.—Ins. Philipp.
elliptica, Thunb. Fl. Jap. 111 = Amsonia elliptica.
fagraeoides, *Miq. Fl. Ind. Bat.* ii. 420.—Java.
fallax, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 84.—Bras.
fasciculata, *Poir. Encyc.* vii. 531.—Guian.
Finlaysoniana, *Wall. Cat.* n. 4452.—Siam.
flavescens, Willd. ex Roem. & Schult. Syst. iv. 798 = Malouetia flavescens.
flavicans, *Wilid. ex Roem. & Schult. l. c.* 797.—Bras.
floribunda, *Blume, Bijdr.* 1028.—Java.
frutescens, Miers, Apocyn. S. Am. 57 = laurifolia.
fuchsiaefolia, *A. DC. in DC. Prod.* viii. 365.—Bras.
Gaudichaudii, *A. DC. l. c.*—Bras.
glabrata, Mart. ex Miers, Apocyn. S. Am. 81 = oblongifolia.
globosa, Blanco, Fl. Filip. ed. I. 116 = Voacanga foetida.
globosa, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 130 = sphaerocarpa.
graciliflora, *Wall. in Bot. Reg.* sub t. 1273; *Cat.* n. 1576.—Burma.
gracilis, *Benth. in Hook. Journ. Bot.* iii. (1841) 244.—Guian.
gracilis, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 82.—Bras.
grandiflora, Hook. in W. Gray, Trav. West. Afr. 389 = africana.
grandiflora, *Jacq. Enum. Pl. Carib.* 14.—Ind. occ.
grandiflora, *Linn. Mant.* i. 53.—N. Granat.
gratissima, Lindl. in Trans. Hort. Soc. vii. 1. (1830) 55; Bot. Reg. t. 1084 = recurva.
guianensis, *Miq. in Linnaea*, xviii. (1844) 754.—Guian.
guyanensis, *Muell. Arg. in Linnaea*, xxx. (1859-60) 404.—Guian.
heterophylla, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 347, nomen.—Ins. Timor.
heterophylla, *Vahl, Eclog. Am.* ii. 22.—Guian.
Heyneana, *Wall. in Bot. Reg.* sub t. 1273; *Cat.* n. 1573.—Ind. or.
Hilariana, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 85.—Bras.
hirta, *Hook. f. Fl. Brit. Ind.* iii. 646.—Malacca.
hirtula, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 73.—Bras.
humilis, Salisb. Prod. 148 = Amsonia Tabernaemontana.

TABERNAEMONTANA :—

Hystrix, *Steud. Nom.* ed. II. ii. 658.—Bras.
indica, Willd. ex Steud. l. c. nomen (Quid ?).—Ind. or.
intercedens, Heurck & Muell. Arg. in Heurck, Obs. Bot. 166 = Heyneana.
jasminoides, H. B. & K. Nov. Gen. et Sp. iii. 225 = Malouetia jasminoides.
javanica, *Miq. Fl. Ind. Bat.* ii. 422.—Java.
Kamassi, Eckl. in S. Afr. Quart. Journ. i. (1830) 371 = Gonioma Kamassi.
lactiflua, Miers, Apocyn. S. Am. 289 (Index) = Malouetia lactiflua.
laeta, *Mart. in Flora*, xx. (1837) II. *Beibl.* 98.—Bras.
laevigata, Mart. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 77 = flavicans.
laevis, Vell. Fl. Flum. 105; iii. t. 18 = Geissospermum Vellosii.
lanceolata, *Miers, Apocyn. S. Am.* 268. t. 7.—Am. austr.
laurifolia, Blanco, Fl. Filip. ed. I. 114 = T. Pandacaqui.
laurifolia, Ker-Gawl. in Bot. Reg. t. 716 = dichotoma.
laurifolia, *Linn. Sp. Pl.* 210.—Ind. occ.
laurifolia, Schott, ex Miers, Apocyn. S. Am. 76 = Stemmadenia insignis.
laxa, Benth. in Hook. Journ. Bot. iii. (1841) 244 = Ambelania laxa.
laxiflora, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 403.—Java.
Linkii, *A. DC. in DC. Prod.* viii. 364.—Bras.
litoralis, *H. B. & K. Nov. Gen. et Sp.* iii. 228.—Mexic.
longiflora, *Benth. in Hook. Niger Fl.* 447.—Afr. trop.
longifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 243.—Guian.
lucida, *H. B. & K. Nov. Gen. et Sp.* vii. 209.—Venezuela.
Lundii, *A. DC. in DC. Prod.* viii. 365.—Bras.
lurida, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 172.—Hab. ?
macrocalyx, *Muell. Arg. in Linnaea*, xxx. (1859-60) 403.—Guian.
macrocarpa, *Jack, in Malay. Misc.* ii. (1822) n. VII. 80.—Sumatra.
macrocarpa, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 154 = Willughbeia firma.
macrophylla, Muell. Arg. in Mart. Fl. Bras. vi. 1. 75 = Phrissocarpus rigidus.
macrophylla, *Poir. Encyc. Suppl.* v. 276.—Am. austr.
malaccensis, *Hook. f. Fl. Brit. Ind.* iii. 649.—Malacca.
Martensii, *Peyr. in Linnaea*, xxx. (1859-60) 31.—Mexic.
mauritiana, *Poir. Encyc.* vii. 530.—Ins. Mascar.
membranacea, *A. DC. in DC. Prod.* viii. 370.—Ins. Nossibé.
membranifolia, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 90.—Burma.
Meyeri, *G. Don, Gen. Syst.* iv. 89.—Guian.
micrantha, *A. DC. in DC. Prod.* viii. 370.—Ins. Maurit.
micrantha, Voigt, Hort. Suburb. Calc. 527 = Wallichiana.
microcarpa, Wall. in Bot. Reg. sub t. 1273; Cat. n. 1581 = Rauwolfia microcarpa.
modesta, *Baker, in Journ. Bot.* xx. (1882) 219.—Madag.
mollis, *Hook. & Arn. Bot. Beech. Voy.* 199.—China.
monocarpa, Steud. Nom. ed. II. ii. 658, sphalm. = macrocarpa.
Muelleriana, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 181.—Bras.
multiflora, Link, ex Roem. & Schult. Syst. iv. 431 = Linkii.
multiflora, *Sm. in Rees, Cycl.* xxxv. n. 11.—Ins. Mascar.
muricata, *Link, ex Roem. & Schult. Syst.* iv. 431.—Bras.
muricata, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 76 = macrophylla.
nereifolia, *Vahl, Eclog. Am.* ii. 21.—Porto Rico.
nervosa, Desf. ex Poir. Encyc. Suppl. v. 275 = persicariaefolia.
nicobarica, *Liebm. Ind. Sem. Hort. Haun.* (1855) 27.—*Cf. Linnaea*, xxviii. (1856) 363.—Ins. Nicobar.
Noronhiana, *Boj. Hort. Maurit.* 210.—Madag.
novo-guineensis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 36.—N. Guin.

TABERNAEMONTANA :—

oblonga, *Wall. in Bot. Reg.* sub t. 1273.—Ind. or.
oblongifolia, *A. DC. in DC. Prod.* viii. 368.—Bras.
obtusa, *Sm. in Rees, Cycl.* xxxv. n. 8.—Ins. Borbon.
obtusifolia, *Poir. Encyc. Suppl.* v. 275.—Madag.
occidentalis, *Miers, Apocyn. S. Am.* 58.—Peruv.
ochracea, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 81 = muricata.
odorata, Vahl, Eclog. Am. ii. 22 = Malouetia Tamaquarina.
olivacea, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 75.—Bras.
ophiorrhizoides, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 89.—Burma.
orientalis, *R. Br. Prod.* 468.—Ind. or.; Austral.
orientalis, G. Don, Gen. Syst. iv. 92 = Melodinus orientalis.
ovalis, *Miq. Fl. Ind. Bat. Suppl.* 229, 554.—Sumatra.
pacifica, *Seem. Fl. Vit.* 160.—Ins. Fiji.
Pandacaqui, *Poir. Encyc.* vii. 529.—N. Guin.
parviflora, Boj. Hort. Maurit. 209 = mauritiana.
parviflora, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 379 = orientalis.
parviflora, Heyne, ex Wall. Cat. p. 158, re n. 1 80 = Hunteria corymbosa.
parviflora, Heyne, ex Wall. Cat. n. 4453 = Rauwolfia densiflora.
parviflora, Poir. Encyc. Suppl. v. 276 = populifolia.
parviflora, Roxb. Hort. Beng. 20; Fl. Ind. ii. 25 = Wallichiana.
pauciflora, *Blume, Bijdr.* 1028.—Java.
pauciflora, Wight, Ic. t. 393 = Wallichiana.
paucifolia, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 87.—Bras.
peduncularis, *Wall. in Bot. Reg.* sub t. 1273; *Cat.* n. 1575.—Ins. Penang.
pentasticta, *Scheff. in Flora*, lii. (1869) 309; *et in Tijdschr. Nederl. Ind.* xxxi. (1870) 22.—Malaya.
Perrottetii, *A. DC. in DC. Prod.* viii. 362.—Guian.
persicariaefolia, *Jacq. Coll.* iv. 139.—Ins. Mascar.
pilifera, Benth. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 73 = hirtula.
Poeppigii, *Muell. Arg. in Linnaea*, xxx. (1859-60) 405.—Bras.
polyantha, Blume, Bijdr. 1029 = Micrechites polyantha.
polyantha, Herb. Lugd. Bat. ex Miq. Fl. Ind. Bat. ii. 394 = Urceola elastica.
polygama, *Blanco, Fl. Filip.* ed. II. 82.—Ins. Philipp.
populifolia, *Poir. Encyc. Suppl.* v. 276.—Am. austr.
psychotriaefolia, Steud. Nom. ed. II. ii. 658 = seq.
psychotrifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 227.—Am. trop.
psychotrioides, G. Don, Gen. Syst. iv. 90 = praec.
pubescens, *R. Br. Prod.* 468.—Austral.
pubescens, *Teijsm. & Binn. Cat. Hort. Bog.* 125.—Ins. Buru.
Rauwolfiae, A. DC. in DC. Prod. viii. 364 = Salzmanni.
recurva, Hohen. ex Hook. f. Fl. Brit. Ind. iii. 646 = Heyneana.
recurva, *Roxb. Hort. Beng.* 20; *Fl. Ind.* ii. 26.—Ind. or.; Burma.
repanda, *E. Mey. ex Steud. Nom.* ed. II. ii. 658.—Guian.
reticulata, *A. DC. in DC. Prod.* viii. 366.—Bras.
rhynchophylla, *Miq. Fl. Ind. Bat.* ii. 422.—Ins. Banca.
Riedeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 139.—Ins. Celebes.
Riedelii, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 72.—Bras.
riparia, H. B. & K. Nov. Gen. et Sp. iii. 228 = Malouetia riparia.
rosea, *Tenore, Cat. Orto Nap.* (1845) 97.—Hab. ?
rostrata, *Wall. in Bot. Reg.* sub t. 1273; *Cat.* n. 1578.—Burma.
rubrostriolata, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 71.—Bras.
rupicola, *Benth. in Hook. Journ. Bot.* iii. (1841) 243.—Guian.
salicifolia, Wall. in Bot. Reg. sub t. 1273; Cat. n. 1580 = Hunteria corymbosa.
Salzmanni, *A. DC. in DC. Prod.* viii. 362.—Bras.
Sananho, *Ruiz & Pav. Fl. Per.* ii. 22. t. 144.—Peruv.
sessilifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 205.—Madag.
sessilifolia, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 952, nomen.—Guian.
sessilis, Vell. Fl. Flum. 106 = Malouetia sessilis.
solanifolia, *A. DC. in DC. Prod.* viii. 365.—Bras.

TABERNAEMONTANA:—
speciosa, *Poir. Encyc. Suppl.* v. 275.—Guian.
sphaerocarpa, *Blume, Bijdr.* 1028.—Java.
Spixiana, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 78.—Bras.
Sprucei, *Muell. Arg. l. c.* 86.—Bras.
squamosa, Sm. in Rees, Cycl. xxxv. n. 3 = mauritiana.
squamosa, Spreng. ex A. DC. in DC. Prod. viii. 327 = Urceola elastica.
stenoloba, *Muell. Arg. in Linnaea*, xxx. (1859–60) 407.—Peruv.
subcapitata, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 633 = Rauwolfia densiflora.
subcapitata, *Wall. in Bot. Reg.* sub t. 1273; *Cat.* n. 1579.—Burma.
submollis, *Mart. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 70.—Bras.
subsessilis, *Benth. in Hook. Niger Fl.* 448.—Afr. trop.
Telfairiana, *Wall. in Bot. Reg.* sub t. 1273.—Ins. Maurit.
tenuiflora, *Miq. Fl. Ind. Bat. Suppl.* 554.—Sumatra.
tenuiflora, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 76.—Bras.
tenuifolia, Willd. ex Meissn. in DC. Prod. xv. 1. 28 = Alseodaphne semecarpifolia.
ternstroemiacea, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 88.—Bras.
tetrastachya, *H. B. & K. Nov. Gen. et Sp.* iii. 227.—N. Granat.
Thurstoni, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 368.—Ins. Fiji.
umbrosa, *H. B. & K. Nov. Gen. et Sp.* iii. 226.—N. Granat.
undulata, G. F. W. Mey. Prim. Fl. Esseq. 135 = Meyeri.
undulata, Perr. ex A. DC. in DC. Prod. viii. 362 = Perrottetii.
undulata, *Vahl, Eclog. Am.* ii. 20.—Ind. occ.; Guian.
utilis, *Arn. in Edinb. N. Phil. Journ.* viii. (1830) 318.—Guian.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 168.—Peruv.
ventricosa, *Hochst. ex A. DC. in DC. Prod.* viii. 366.—Afr. austr.
verrucosa, *Blume, Bijdr.* 1029.—Java.
verticillata, Bedd. Forest. Fl. Anal. Gen. xx; Man. Bot. 159 = Rauwolfia Beddomei.
vitiensis, Seem. in Bonplandia, ix. (1861) 257 = orientalis.
Wallichiana, *Steud. Nom.* ed. II. ii. 658.—Sumatra.
Warmingii, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 107.—Bras.
Wullschlaegelii, *Griseb. Fl. Brit. W. Ind.* 409.—Jamaic.

TABRACA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Anonacea?).

TACAMAHACA, Mill. Gard. Dict. ed. VI. App. 175 (1752) = **Populus**, Linn. (Salicin.).

TACAZZEA, Decne. in DC. Prod. viii. 492 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 745.
Leptopaetia, Harv. Gen. S. Afr. Pl. ed. II. 231 (1868).
apiculata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 108.—Afr. trop.
venosa, *Decne. in DC. Prod.* viii. 493.—Abyss.

TACCA, Forst. Char. Gen. 69. t. 35 (1776). *TACCACEAE*, Benth. & Hook. f. iii. 741.
Ataccia, Presl, Rel. Haenk. 149 (1828).
artocarpifolia, *Seem. Fl. Vit.* 101.—Ins. Mascar.
aspera, Roxb. Hort. Beng. 25; Fl. Ind. ii. 169 = integrifolia.
Brownii, Seem. Fl. Vit. 100 = pinnatifida.
cristata, *Jack, in Malay. Misc.* i. (1820) n. v. 23.—Ind. or.
dubia, *Schult. f. Syst.* vii. 167.—Ins. Amboin.
Gaogao, Blanco, Fl. Filip. ed. I. 262 = pinnatifida.
guineensis, G. Don, ex Loud. Hort. Brit. 124, nomen = involucrata?
integrifolia, *Ker-Gawl. in Bot. Mag.* t. 1488.—Ind. or.
involucrata, *Schum. & Thonn. Beskr. Guin. Pl.* 177.—Afr. trop.
laevis, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 171.—Ind. or.
lancaefolia, *Zoll. & Mor. Syst. Verz. Zoll.* 91.—Java.
maculata, Seem. Fl. Vit. 103 = pinnatifida.

TACCA:—
maculata, Zipp. ex Span. in Linnaea, xv. (1841) 480, nomen (Quid?).—Ins. Timor.
madagascariensis, Boj. Hort. Maurit. 350 = pinnatifida?
montana, Schult. f. Syst. vii. 168 = palmata.
oceanica, Seem. in Journ. Bot. iv. (1866) 261 = pinnatifida.
palmata, *Blume, Enum. Pl. Jav.* i. 83.—Java.
palmatifida, *Baker, in Journ. Linn. Soc.* xv. (1876) 100.—Ins. Celebes.
Parkeri, *Seem. Fl. Vit.* 102.—Guian.
phallifera, Schult. f. Syst. vii. 167, in syn. = pinnatifida, Amorphophallus campanulatus.
pinnatifida, *Forst. Char. Gen.* 70. t. 35.—As. et Afr. trop.; Ins. Pacif.
pinnatifolia, Gaertn. Fruct. i. 43. t. 14. f. 2, sphalm. = praec.
quanzensis, Welw. Apont. 591 = pinnatifida.
Rafflesiana, Jack, ex Wall. Cat. n. 5172 = cristata.
Rumphii, *J. C. Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 442.—Ins. Philipp.
Sprucei, *Benth. in Benth. & Hook. f. Gen.* iii. 741.—Bras.
vesicaria, Blanco, Fl. Filip. ed. I. 261 = palmata.

TACCARUM, Brongn. ex Schott, Gen. Aroid. t. 65 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 986.
Endera, Regel, Gartenfl. (1872) 226. t. 732.
Lysistigma, Schott, in Bonplandia, x. (1862) 222.
cylindricum, *Arcangeli, in Nuov. Giorn. Bot. Ital.* xi. (1879) 190.—Java.
peregrinum, Engl. in DC. Monog. Phan. ii. 646 = cylindricum.
Warmingii, *Engl. in Kjoeb. Vidensk. Meddel.* (1880) 362.—Bras.
Weddellianum, *Brongn. ex Schott, Gen. Aroid.* t. 65.—Bras.

TACHIA, Aubl. Pl. Gui. i. 75. t. 29 (1775). *GENTIANEAE*, Benth. & Hook. f. ii. 813.
Myrmecia, Schreb. Gen. 74 (1789).
calygona, Griseb. Gen. et Sp. Gent. 204 = Lisianthus calygonus.
cordifolia, Mart. Nov. Gen. et Sp. ii. 119 = Leianthus longifolius.
exsertus, Mart. l. c. = Leianthus exertus.
gracilis, *Benth. in Hook. Kew Journ.* vi. (1854) 203.—Guian.
guianensis, *Aubl. Pl. Gui.* i. 75. t. 29.—Guian.
longifolia, Mart. Nov. Gen. et Sp. ii. 119 = Leianthus longifolius.
Schomburgkiana, *Benth. in Hook. Kew Journ. Bot.* vi. (1854) 204.—Guian.
Swartzia, G. Don, Gen. Syst. iv. 197 = Leianthus exsertus.

TACHIA, Pers. Syn. i. 459 (1805) = **Tachigalia**, Aubl. (Legumin.).
paniculata, Pers. l. c. 460 = Tachigalia paniculata.
punctata, Plée, ex Tul. in Arch. Mus. Par. iv. (1844) 148 = Caesalpinia monosperma.
trigona, Pers. Syn. i. 460 = Tachigalia paniculata.

TACHIADENUS, Griseb. Gen. et Sp. Gent. 200 (1839). *GENTIANEAE*, Benth. & Hook. f. ii. 804.
carinatus, *Griseb. l. c.*—Madag.
gracilis, *Griseb. in DC. Prod.* ix. 82.—Madag.
longiflorus, *Griseb. Gen. et Sp. Gent.* 202.—Madag.
platypterus, *Baker, in Journ. Bot.* xx. (1882) 172.—Madag.
trinervis, *Griseb. Gen. et Sp. Gent.* 201.—Madag.
tubiflorus, *Griseb. l. c.* 202.—Madag.

TACHIBOTA, Aubl. Pl. Gui. 287. t. 112 (1775). Genus dubium *BIXINEARUM*.—Cf. Benth. & Hook. f. i. 124.
Salmasia, Schreb. Gen. 201 (1789).
guianensis, *Aubl. Pl. Gui.* i. 287.—Guian.

TACHIGALIA, Aubl. Pl. Gui. 372. t. 143 (1775) (Tachigali). *LEGUMINOSAE*, Benth. & Hook. f. i. 582.
Cuba, Scop. Introd. 300 (1777).
Tachia, Pers. Syn. i. 459 (1805).
Tassia, Rich. ex DC. Prod. ii. 487 (1825).
Valentinia, Neck. Elem. ii. 450 (1790).
angustifolia, Miq. Stirp. Surin. Sel. 13 = paniculata.
aurea, Tul. in Arch. Mus. Par. iv. (1844) 169 = Sclerolobium aureum.

TACHIGALIA:—
bijuga, DC. Prod. ii. 487 = Sweetia bijuga.
emarginata, *G. Don, Gen. Syst.* ii. 436.—Bras.
eriocalyx, Tul. in Arch. Mus. Par. iv. (1844) 164 = paniculata.
floribunda, Steud. Nom. ed. II. ii. 659 = Moldenhawera floribunda.
glauca, *Tul. in Arch. Mus. Par.* iv. (1844) 165.—Guian.
multijuga, *Benth. in Mart. Fl. Bras.* xv. ii. 229.—Am. austr.
paniculata, *Aubl. Pl. Gui.* i. 372. t. 143. f. 1.—Guian.
Poeppigiana, Tul. in Arch. Mus. Par. iv. (1844) 168 = polyphylla.
polyphylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 60.—Bras.
ptychophysca, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. ii. 229.—Bras.
pubiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 94.—Am. trop.
purpurea, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guian.
Richardiana, *Tul. in Arch. Mus. Par.* iv. (1844) 166.—Guian.
sericea, Tul. l. c. 163 = paniculata.
trigona, Aubl. Pl. Gui. i. 372. t. 143. f. 2 = paniculata.

TACHIGALEA, Griseb. Fl. Brit. W. Ind. 501 (1861), sphalm. = TALIGALEA, Aubl. = **Amasonia**, Linn. f. (Verben.).

TACHITES, Soland. ex Gaertn. Fruct. i. 205 (1788) = **Melicytus**, Forst. (Violar.).
umbellata, Soland. ex Gaertn. l. c. = M. ramiflorus.

TACHYTES, Steud. Nom. ed. II. ii. 659 (1841), sphalm. = praec.

TACSONIA, Juss. Gen. 398 (1789). *PASSIFLOREAE*, Benth. & Hook. f. i. 811.
Odostelma, Rafin. Fl. Tellur. iv. 104 (1836).
Poggendorffia, Karst. in Linnaea, xxviii. (1856) 438; et Fl. Columb. i. 29. t. 15 (1858).
Rathea, Karst. Fl. Columb. i. 77. t. 38 (1859?).
adulterina, *Juss. in Ann. Mus. Par.* vi. (1805) 393.—N. Granat.
ampullacea, *Mast. in Mart. Fl. Bras.* xiii. 1. 539.—Ecuador.
anastomosans, *Lamb. ex DC. Prod.* iii. 335.—Peruv.
Andersoni ×, Mast. in Gard. Chron. (1875) ii. 167.
bicoronata, Mast. in Mart. Fl. Bras. xiii. 1. 541 = mixta?
bicuspidata, Karst. in Linnaea, xxx. (1859–60) 160 = Passiflora bicuspidata.
bilobata, Spreng. Syst. iii. 44 = Passiflora bilobata.
Buchanani, Lem. Illustr. Hortic. xiv. (1867) t. 519 = Passiflora vitifolia.
canaliculata, Juss. in Ann. Mus. Par. vi. (1805) 392 = Passiflora glandulosa.
candida, Poepp. & Endl. Nov. Gen. et Sp. ii. 59 = Passiflora candida.
citrifolia, Juss. in Ann. Mus. Par. vi. (1805) 392 = Passiflora citrifolia.
cumbalensis, Karst. in Linnaea, xxx. (1859–60) 161 = glaberrima.
cuneata, Benth. Pl. Hartw. 183 = Passiflora bicuspidata.
Dombeyana, M. Roem. Syn. Pepon. 197 = pedunculares.
eriantha, Benth. Pl. Hartw. 183 = mixta.
exoniensis ×, Hort. Veitch.—Cf. Gard. Chron. (1872) 1653.
flexipes, *Mast. in Journ. Linn. Soc.* xx. (1883) 26.—N. Granat.
floribunda, *Mast. in Trans. Linn. Soc.* xxvii. (1871) 628; *et in Mart. Fl. Bras.* xiii. 1. 538.—N. Granat.
Fockeana, Miq. in Linnaea, xviii. (1844) 364 = Passiflora glandulosa.
fulgens, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 127 = Passiflora coccinea, quadriglandulosa.
glaberrima, *Juss. in Ann. Mus. Par.* vi. (1805) 394.—Peruv.
glandulosa, Juss. l. c. 391 = Passiflora glandulosa.
glauca, Poepp. ex Mast. in Mart. Fl. Bras. xiii. 1. 569 = Passiflora reflexiflora.
gracilens, *A. Gray, in Bot. U. St. Expl. Exped.* i. 639.—Peruv.
grandiflora, *Forbes, Hort. Woburn.* 232.—Hab.?

TACSONIA:—
grandis, Hort. ex Mast. in Journ. Hort. Soc. N. S. iv. (1877) 127 = Van-Volxemii ?
hastata, Ruiz & Pav. ex Mast. in Trans. Linn. Soc. xxvii. (1871) 638 = Passiflora reflexiflora.
hederacea, *Mast. & André, in Journ. Linn. Soc.* xx. (1883) 29.—Ecuador.
ignea, Hort. ex Mast. in Trans. Linn. Soc. xxvii. (1871) 629 = manicata.
infundibularis, *Mast. in Journ. Bot.* xxi. (1883) 34.—Venezuela.
insignis, *Mast. in Gard. Chron.* (1873) 1112.—Am. austr.
Jamesoni, *Mast. in Mart. Fl. Bras.* xiii. 1. 537.—Peruv.
laevis, Benth. Pl. Hartw. 118 = Passiflora reflexiflora.
lanata, *Juss. in Ann. Mus. Par.* vi. (1805) 392. t. 59. f. 1.—N. Granat.
lanceolata, *Mast. in Mart. Fl. Bras.* xiii. 1. 536.—Peruv.
longifolia, Pers. Syn. ii. 223 = mixta.
Mandoni, *Mast. in Mart. Fl. Bras.* xiii. 1. 538.—Bolivia.
manicata, *Juss. in Ann. Mus. Par.* vi. (1805) 393. t. 59. f. 2.—Peruv.
Mansii, Mart. in Flora, xxii. (1839) 1. Beibl. 7 = Passiflora Mansoi.
Mathewsii, *Mast. in Mart. Fl. Bras.* xiii. 1. 539 (Matthewsii).—Peruv.
micradena, *DC. Prod.* iii. 334.—Peruv.
mixta, *Juss. in Ann. Mus. Par.* vi. (1805) 394.—Am. trop.
mollissima, Hort. ex Mast. in Gard. Chron. (1869) 388 = mixta.
mollissima, *H. B. & K. Nov. Gen. et Sp.* ii. 144.—N. Granat.
Parritae, *Mast. in Gard. Chron.* (1882) 1. 218.—N. Granat.
parvifolia, *DC. Prod.* iii. 335.—Peruv.
peduncularis, *Juss. in Ann. Mus. Par.* vi. (1805) 395.—Peruv.
pennipes, M. Roem. Syn. Pepon. 194 = pinnatistipula.
pinnatistipula, *Juss. in Ann. Mus. Par.* vi. (1805) 393.—Chili.
pubescens, DC. Prod. iii. 335 = Passiflora quadriglandulosa.
Purupuru, DC. ex Mast. in Mart. Fl. Bras. xiii. 1. 537 = pinnatistipula.
quadridentata, DC. Prod. iii. 335 = Passiflora quadriglandulosa.
quadriglandulosa, DC. l. c. = Passiflora quadriglandulosa.
quitensis, Benth. Pl. Hartw. 183 = mixta.
reflexiflora, Juss. in Ann. Mus. Par. vi. (1805) 393 = Passiflora reflexiflora.
Rohriana, DC. Prod. iii. 335 = Passiflora glandulosa.
rugosa, *Mast. in Mart. Fl. Bras.* xiii. 1. 539.—N. Granat.
sanguinea, Benth. ex Mast. l. c. 542 = Passiflora vitifolia.
sanguinea, DC. Prod. iii. 334 = Passiflora quadriglandulosa.
serrata, Karst. in Linnaea, xxx. (1859–60) 160 = mixta.
speciosa, H. B. & K. Nov. Gen. et Sp. ii. 143 = mixta.
spinescens, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1168, nomen.—Guian.
spinosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 59 = Passiflora spinosa.
splendens, *Hort. Veitch, ex Mast. in Journ. Hort. Soc.* N. S. iv. (1877) 127.—Hab. ?
Stoupyana, DC. Prod. iii. 335 = Passiflora glandulosa.
subcoriacea, Garcke, in Linnaea, xxii. (1849) 62 = Passiflora glandulosa.
Tacso, Pers. Syn. ii. 223 = mixta.
tomentosa, *Juss. in Ann. Mus. Par.* vi. (1805) 394.—Peruv.
trifoliata, *Juss. l. c.* 393.—Peruv.; Chili.
trigona, DC. Prod. iii. 334 = trifoliata.
trinervia, Juss. in Ann. Mus. Par. vi. (1805) 390. t. 58 = Passiflora trinervia.
tripartita, *Juss. l. c.* 395. t. 60.—Ecuador.
umbilicata, *Griseb. in Goett. Abh.* xix. (1874) 149.—Reg. Argent.
urceolata, *Mast. in Mart. Fl. Bras.* xiii. 1. 539.—Am. trop.
Van-Volxemii, *Hook. Bot. Mag.* t. 5571.—N. Granat.
viridiflora, Juss. in Ann. Mus. Par. vi. (1805) 389 = Passiflora viridiflora.

TAENAIS, Salisb. Gen. Pl. Fragm. 115 (1866) = **Crinum**, Linn. (Amaryll.).
ampla, Salisb. l. c. = C. giganteum.
Australasiae, Salisb. l. c. = C. flaccidum.
caricifolia, Salisb. l. c. = C. yuccaeflorum.
mucronigera, Salisb. l. c. = C. zeylanicum.
ornata, Salisb. l. c. = C. yuccaeflorum ?
yuccaeflorum, Salisb. l. c. = C. yuccaeflorum.
zeylanica, Salisb. l. c. = C. zeylanicum.

TAENIDIUM, Targ. Tozz. Cat. Veg. Mar. 80. t. 1 (1826) = **Posidonia**, Kon. (Naiad.).
acuminatum, Targ. Tozz. l. c. 84 = P. Caulini.
oceanicum, Targ. Tozz. l. c. 81 = P. Caulini.

TAENIOCARPUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 420 = **Pachyrhizus**, Rich. (Legumin.).
articulatum, Desv. l. c. 421 = P. angulatus.

TAENIOCHLAENA, Hook. f. in Benth. & Hook. f. Gen. i. 433 (1865). *CONNARACEAE*, Benth. & Hook. f. l. c.
Griffithii, *Hook. f. l. c.* 434.—Malacca.

TAENIOLA, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Tourn. (Liliac.).
vittata, Salisb. l. c. = O. vittatum.

TAENIOPETALUM, Vis. Fl. Dalm. iii. 49 (1849) = **Peucedanum**, Tourn. (Umbell.).
Neumayeri, Vis. l. c. = P. Neumayeri.
peucedanoides, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 203 = P. alsaticum.

TAENIOPHYLLUM, Blume, Bijdr. 355. t. 70 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 582.
ALWISIA, Thw. ex Lindl. in Journ. Linn. Soc. iii. (1858) 42, in obs.
acuminatissimum, *Blume, Bijdr.* 357.—Java.
Alwisii, *Lindl. in Journ. Linn. Soc.* iii. (1858) 42.—Zeylan.
asperulum, *Reichb. f. Otia Bot. Hamb.* 53.—Ins. Tahiti.
elegantissimum, *Reichb. f. l. c.*—Ins. Tahiti.
Fasciola, *Reichb. f. in Seem. Fl. Vit.* 296.—Ins. Fiji.
glandulosum, *Blume, Bijdr.* 356.—Java.
Hasseltii, *Reichb. f. Xenia Orch.* ii. 68.—Java ?
hirtum, *Blume, Bijdr.* 356.—Java.
Jerdonianum, Wight, Ic. v. t. 1756 = Saccolabium Jerdonianum.
Muelleri, *Lindl. ex Benth. Fl. Austral.* vi. 291.—Austral.
obtusum, *Blume, Bijdr.* 357.—Java.
philippinense, *Reichb. f. Otia Bot. Hamb.* 53.—Ins. Philipp.
Seemanni, *Reichb. f. in Seem. Fl. Vit.* 296.—Ins. Fiji.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.

TAENIOSAPIUM, Muell. Arg. in DC. Prod. xv. II. 1200 (1866) = **Excoecaria**, Linn. (Euphorb.).
Goudotianum, Muell. Arg. l. c. 1201 = E. Goudotiana.

TAENIOSTEMA, Spach, in Hook. Comp. Bot. Mag. ii. (1836) 289 = **Helianthemum**, Tourn. (Cistin.).
micranthum, Spach, l. c. = H. glomeratum.

TAENOSAPIUM, Benth. & Hook. f. Gen. iii. 337 (1880), err. typ. = TAENIOSAPIUM, Muell. Arg. = **Excoecaria**, Linn. (Euphorb.).

TAETSIA, Medic. Theodora, 82 (1786) = **Cordyline**, Royen (Liliac.).
ferrea, Medic. l. c. = C. terminalis.

TAFALLA, D. Don, in Edinb. N. Phil. Journ. (July–Sept. 1831) 273. *COMPOSITAE*, Benth. & Hook. f. ii. 300.
LORICARIA, Wedd. Chlor. And. i. 165. t. 27* A, B, C (1856).
THYOPSIS, Wedd. l. c. 165, in nota, t. 27 (1856).
ferruginea, *D. Don, in Edinb. N. Phil. Journ.* (July–Sept. 1831) 274.—Reg. Andin.
thyoides, *D. Don, l. c.* 273.—Reg. Andin.

TAFALLA, Ruiz & Pav. Prod. 136. t. 29 (1794) = **Hedyosmum**, Sw. (Chloranth.).
angustifolia, Ruiz & Pav. Syst. Veg. 272 = H. angustifolium.

TAFALLA:—
glauca, Ruiz & Pav. ex Spreng. Syst. iii. 866 = Hedyosmum Bonplandianum.
glauca, Ruiz & Pav. Syst. Veg. 271 = H. glaucum.
laciniata, Ruiz & Pav. l. c. 272 = H. laciniatum.
racemosa, Ruiz & Pav. l. c. 271 = H. racemosum.
scabra, Ruiz & Pav. l. c. 270 = H. scabrum *vel* glabratum.

TAGERA, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legumin.).
filiformis, Rafin. l. c. = C. Tagera.

TAGETES, Linn. Gen. ed. I. 261 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 411.
DIGLOSSUS, Cass. in Bull. Soc. Philom. (1817) 70.
ENALCIDA, Cass. l. c. (1819) 31.
SOLENOTHECA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 375.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 194.—Mexic.
apetala, *Posado, in Baill. Adansonia*, x. (1872) 186.—N. Granat.
bonariensis, Pers. Syn. ii. 459 = minuta.
campanulata, *Griseb. in Goett. Abh.* xix. (1874) 188.—Reg. Argent.
caracassana, *Humb. ex Willd. Sp. Pl.* iii. 2128.—Venezuela.
clandestina, Lag. Gen. et Sp. Nov. 28 = coronopifolia.
congesta, Hook. & Arn. Bot. Beech. Voy. 299 = micrantha.
coronopifolia, *Willd. Enum. Hort. Berol. Suppl.* 60.—Mexic.
corymbosa, Lag. Elench. (1805); ex ej. Gen. et Sp. Nov. 29 = coronopifolia.
corymbosa, Sweet, Brit. Flow. Gard. Ser. I. t. 151 = patula.
daucoides, *Schrad. Sem. Hort. Gotting.* (1833) 5.—*Cf. Linnaea*, x. (1836) *Litt.* 71.—Chili.
dianthiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 197.—Peruv.
dianthifolia, DC. Prod. v. 646, sphalm. = praec.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 72.—Peruv.
elliptica, *Sm. in Rees, Cycl.* xxxv. n. 7.—Peruv.
elongata, Willd. Sp. Pl. iii. 2127 = patula, peduncularis.
erecta, *Linn. Sp. Pl.* 887.—Mexic.
Feuillaei, Bert. ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 35. t. 30 = Lasthenia obtusifolia.
Fevillei, Steud. Nom. ed. II. ii. 12, in syn., sphalm. = praec.
filifolia, Lag. Elench. (1805); ex ej. Gen. et Sp. Nov. 28 = multifida.
filiformis, Hemsl. Biol. Centr. Am. Bot. ii. 223, sphalm. = praec.
florida, Sweet, Brit. Flow. Gard. Ser. II. t. 35 = lucida.
flosculosa, *Spreng. Syst.* iii. 571.—Reg. Argent.
foeniculacea, Desf. Cat. Hort. Par. ed. III. 171 = coronopifolia.
foeniculacea, *Poepp. ex DC. Prod.* v. 646.—Peruv.
foetidissima, *Hort. ex DC. l. c.* 645.—Mexic.
glandulifera, Schrank, Pl. Rar. Hort. Monac. t. 54 = minuta.
glandulosa, Schrank, ex Link, Enum. Hort. Berol. ii. 339 = minuta.
gracilis, *DC. Prod.* v. 645.—Peruv.
graveolens, *L'Hérit. ex DC. l. c.* 644.—Peruv.
lanceolata, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 73.—Mexic.
Lemmoni, *A. Gray, Proc. Am. Acad.* xix. (1883) 40.—Arizona.
lucida, *Cav. Ic.* iii. 33. t. 264.—Mexic.
lunulata, *Ortega, Hort. Matr. Dec.* 44.—Mexic.
macroglossa, *Polak. in Linnaea*, xli. (1877) 580.—Costa Rica.
maculata, Hort. ex Steud. Nom. ed. II. ii. 660 (Quid ?).—Hab. ?
major, Gaertn. Fruct. ii. 434. t. 172 = erecta.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 529, nomen.—Bolivia.
Michauxiana, Roem. ex Steud. Nom. ed. II. ii. 660 (Quid ?).—Hab. ?
micrantha, *Cav. Ic.* iv. 31. t. 352.—Mexic.
microglossa, *Benth. Bot. Voy. Sulph.* 118.—N. Granat.
minuta, *Linn. Sp. Pl.* 887.—Am. trop.
montana, Hort. ex DC. Prod. v. 644 = glandulifera.
multifida, *DC. l. c.* 646.—Mexic.

TAGETES:—
multiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 197.—Ecuador.
multiseta, *DC. Prod.* v. 645.—Mexic.
oligocephala, *DC. l. c.*—Mexic.
Ozolis, Spreng. ex DC. Prod. v. 647 (Quid ?).—Hab. ?
papposa, Vent. Jard. Cels, 36 = Dysodia chrysanthemoides.
Parryi, *A. Gray, in Proc. Am. Acad.* xv. (1880) 40.—Mexic.
patula, *Linn. Sp. Pl.* 887.—Mexic.
pauciloba, *DC. Prod.* v. 645.—Chili.
pectinata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 71.—Peruv.
pedunculoris, *Cav. Desc.* 201; *Lag. & Rod. in Anal. Cienc. Nat.* v. (1802) 74.—Mexic.
Pineda, *La Llave, in La Naturaleza*, vii. (1885) *Apénd.* 73.—Mexic.
Porophyllum, Vell. Fl. Flum. viii. t. 116 = minuta.
pumila Hort. Pisan. ex Pursh, Fl. Am. Sept. ii. 559 = Dysodia chrysanthemoides.
punctata, Moç. & Sesse, ex DC. Prod. v. 638 = Adenophyllum coccineum.
pusilla, *H. B. & K. Nov. Gen. et Sp.* iv. 194.—Ecuador.
remotiflora, *Kunze, in Linnaea*, xx. (1847) 23.—Mexic.
rotundifolia, *Mill. Gard. Dict.* ed. VIII. n. 4.—Mexic.
Schiedeana, *Less. in Linnaea*, ix. (1835) 271.—Mexic.
signata, *Bartl. Ind. Sem. Hort. Gotting.* (1837) 7.—Mexic.
silenoides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 272.—Peruv.
subulata, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 31 = multiseta.
subvillosa, *Lag. Elench.* (1804); *ex ej. Gen. et Sp. Nov.* 29.—Mexic.
tanacetifolia, Hort. ex Steud. Nom. ed. II. ii. 660 = foetidissima.
tenuifolia, *Cav. Ic.* ii. 54. t. 169.—Peruv.
tenuifolia, H. B. & K. Nov. Gen. et Sp. iv. 196 = patula.
terniflora, *H. B. & K. l. c.*—Ecuador.
tinctoria, *Hornsch. Del. Sem. Hort. Gryph.* (1845) 3.—Bras.
verticillata, *Lag. & Rod. in Anal. Cienc. Nat.* v. (1802) 73.—Venezuela.
Wislizenii, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 92.—Mexic.
zypaquinensis, *Humb. & Bonpl. Pl. Aequin.* ii. 18. t. 73.—N. Granat.

AGUARIA, Rafin. Sylva Tellur. 125 (1838) = **Loranthus**, Linn.
elliptica, Rafin. l. c. = L. ellipticus.
laurifolia, Rafin. l. c. = L. laurifolius.
nitida, Rafin. l. c. = L. nitidus.
punctata, Rafin. l. c. = L. punctatus.
puracensis, Rafin. l. c. = L. puracensis.
vera, Rafin. l. c. = L. Tagua.

TAINIA, Blume, Bijdr. 354 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 515.
ANIA, Lindl. Gen. et Sp. Orch. 129 (1831).
MITOPETALUM, Blume, Fl. Jav. Praef. 8 (1838).
angustifolia, *Benth. & Hook. f. Gen.* iii. 515.—Burma.
barbata, Lindl. in Gard. Chron. (1857) 68 = Eria barbata.
bicornis, *Reichb. f. in Bonplandia*, v. (1857) 54.—Ind. or.
fimbriata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 492.—Java.
latifolia, *Reichb. f. in Bonplandia*, v. (1857) 54.—Reg. Himal.; Burma.
maculata, Trimen, Cat. Pl. Ceyl. 88 (= Chrysoglossi sp.).—Zeylan.
speciosa, *Blume, Bijdr.* 354.—Malaya.
trinervis, *Reichb. f. in Bonplandia*, v. (1857) 54.—Java.

TALA, Blanco, Fl. Filip. ed. I. 484 (1837); ed. II. 338 (1845) = **Limnophila**, R. Br. (Scrophul.).
odorata, Blanco, l. c. ed. I. 484 = L. Roxburghii.

TALANELIS, Rafin. Fl. Tellur. iv. 70 (1836) = **Campanula**, Tourn.
Medium, Rafin. l. c. = C. Medium.

TALANGNINIA, Chapel. ex DC. Prod. iv. 389 (1830) = **Randia**, Houst. (Rubiac.).

TALASIUM, Spreng. Syst. iv. Cur. Post. 22 (1827) = THALASIUM, Spreng. = **Panicum**, Linn. (Gram.).

TALAUMA, Juss. Gen. 281 (1789). *MAGNOLIACEAE*, Benth. & Hook. f. i. 18.
AROMADENDRON, Blume, Bijdr. 8 (1825).
BLUMEA, Nees, in Verh. Batav. Gen. ix. (1823) 147.
BUERGERIA, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 78. t. 2 (1845).
angatensis, D. Vidal, Cat. Pl. Prov. Man. 17 (an seq. ?).—Ins. Philipp.
angatensis, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 3 = Hodgsoni.
caerulea, Jaume St. Hil. Expos. ii. 76 (*cerulea*) = Plumierii.
Candollei, *Blume, Bijdr.* 9; *Fl. Jav. Magnol.* 32.—Ins. Amboin.
Cespedesii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 23.—N. Granat.
coerulea, Steud. Nom. ed. II. ii. 660 = *caerulea*.
dubia, *Eichl. in Mart. Fl. Bras.* xiii. I. 126.—Bras.
elegans, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 70.—Java.
fragrantissima, *Hook. Ic. Pl.* tt. 208–212.—Bras.
gigantifolia, *Miq. Fl. Ind. Bat.* i. II. 15.—Sumatra.
glauca, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 70.—Sumatra.
Hodgsoni, *Hook. f. & Thoms. Fl. Ind.* i. 74.—Reg. Himal.
lanigera, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 40.—Malaya.
liliifera, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 47.—Burma.
macrocarpa, *Zucc. in Abh. Akad. Muench.* ii. (1831–36) 369. t. 1.—Mexic.
mexicana, *G. Don, Gen. Syst.* i. 85.—Mexic.
mutabilis, *Blume, Fl. Jav. Magnol.* 35.—Java.
mutabilis, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 3 = Villariana.
obovata, Benth. & Hook. f. ex Hance, in Journ. Bot. xx. (1882) 2 = Magnolia obovata.
obovata, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) *Versl.* 98.—Borneo.
ovalis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 69.—Ins. Celebes.
ovata, *A. St. Hil. Fl. Bras. Mer.* i. 26. t. 4. f. A.—Bras.
Plumierii, *DC. Prod.* i. 81.—Ind. occ.
pumila, Blume, Fl. Jav. Magnol. 38 = Magnolia pumila.
Rabaniana, *Hook. f. & Thoms. Fl. Ind.* i. 75.—Reg. Himal.
Roxburghii, G. Don, Gen. Syst. i. 85 = Magnolia pterocarpa.
rubra, *Miq. Fl. Ind. Bat.* i. II. 14.—Sumatra.
Rumphii, *Blume, Bijdr.* 10.—Ins. Molucc.
salicifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 258.—Japon.
Sebassa, *Miq. Fl. Ind. Bat. Suppl.* 153, 367.—Sumatra.
Selloi, Steud. Nom. ed. II. ii. 90, in syn., sphalm. = seq.
Sellowiana, *A. St. Hil. Fl. Bras. Mer.* i. 26. t. 4. f. B.—Bras.
Sieboldii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 257 = Magnolia obovata.
stellata, *Miq. l. c.*—Japon.
Villariana, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 307.—Ins. Philipp.
villosa, *Miq. Fl. Ind. Bat. Suppl.* 367.—Sumatra.
Vrieseana, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 70 = Magnolia Vrieseana.

TALBOTIA, Balf. in Trans. Bot. Soc. Edinb. ix. (1868) 190 = **Vellozia**, Vand. (Amaryll.).
elegans, Balf. l. c. 192 = V. elegans.

TALGUENEA, Miers, Trav. Chili, ii. 529 (1826). *RHAMNEAE*, Benth. & Hook. f. i. 385.
costata, *Miers, l. c.*—Chili.
mollis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 8; *Contrib.* i. 298.—Chili.

TALI, Adans. Fam. ii. 319 (1763) = **Connarus**, Linn.

TALIERA, Mart. Progr. Palm. Fam. et Gen. 10 (1824) = **Corypha**, Linn. (Palm.).
bengalensis, Spreng. Syst. ii. 18 = C. Taliera.
Gembanga, Schult. f. Syst. vii. 1307 = C. umbraculifera.
sylvestris, Blume, in Schult. f. l. c. = C. sylvestris.
Tali, Mart. in Schult. f. l. c. 1306 = C. Taliera.

TALIGALEA, Aubl. Pl. Gui. ii. 625. t. 252 (1775) = **Amasonia**, Linn. f. (Verben.).
campestris, Aubl. l. c. = A. erecta.
punicea, Poir. Encyc. vii. 556 = A. punicea.
umbellata, Blanco, Fl. Filip. ed. II. 337 = Ixora macrophylla.

TALINIUM, Rafin. in Am. Monthly Mag. (1818) 175 = **Talinum**, Adans. (Portul.).

TALINOPSIS, A. Gray, Pl. Wright. i. (in Smithson. Contrib. iii.) 14. t. 3 A (1852). *PORTULACEAE*, Benth. & Hook. f. i. 157.
frutescens, *A. Gray, l. c.* 15.—N. Mexic.

TALINUM, Adans. Fam. ii. 245 (1763). *PORTULACEAE*, Benth. & Hook. f. i. 157.
HELIANTHEMOIDES, Medic. Phil. Bot. i. 95 (1789).
PHEMERANTHUS, Rafin. in Med. Repos. N. York, v. (1808) 350.
acaule, Spreng. Syst. ii. 453 = Calandrinia compressa.
adscendens, Hort. Berol. ex DC. Prod. iii. 359 = Calandrinia compressa.
album, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 116 = Calandrinia alba.
aliatum, Hook. Exot. Bot. ii. t. 82, sphalm. = *ciliatum*, Hook.
Anacampseros, Moench, Meth. 232 = Anacampseros Telephiastrum.
Andrewsii, Sweet, Hort. Brit. ed. I. 170 = Calandrinia Andrewsii.
arachnoides, Ait. Hort. Kew. ed. II. iii. 149 = Anacampseros arachnoides.
Arnottii, *Hook. f. Bot. Mag.* t. 6220.—Afr. austr.
aurantiacum, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 153.—Texas.
brachypodium, *S. Wats. in Proc. Am. Acad.* xx. (1885) 355.—Mexic.
brevifolium, *Torr. in Sitgr. Rep.* 156.—Am. bor. occ.
caffrum, Eckl. & Zeyh. Enum. 282 = Portulaca caffra.
calycinum, *Engelm. in Wisliz. Tour N. Mexico*, 88.—Am. bor. occ.
caulescens, Spreng. Syst. i. 453 = Calandrinia caulescens, nitida.
ciliatum, Hook. Exot. Bot. ii. t. 82 = Calandrinia pilosiuscula.
ciliatum, Lindl. Bot. Reg. (1843) sub t. 1 = teretifolium.
ciliatum, Ruiz & Pav. Prod. 65, in obs.; et Syst. Veg. 116 = Calandrinia ciliata.
confertiflorum, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 121.—Am. bor.
crassifolium, Willd. Sp. Pl. ii. 862 = triangulare.
crenatum, *Ruiz & Pav. Prod.* 65, in obs.; *Syst. Veg.* 115.—Peruv.
cuneifolium, *Willd. Sp. Pl.* ii. 864.—Ind. or.; Arab.; Afr.
decumbens, Willd. Sp. Pl. ii. 864 = Orygia decumbens.
dichotomum, Ruiz & Pav. Syst. Veg. 118 = reflexum.
diffusum, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 70.—Chili.
filamentosum, Ait. Hort. Kew. ed. II. iii. 149 = Anacampseros filamentosa.
fruticosum, Macfad. Fl. Jamaic. ii. 169 = patens.
fruticosum, Willd. Sp. Pl. ii. 864 = triangulare.
gracile, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 70.—Chili.
grandiflorum, Hort. ex G. Don, in Loud. Hort. Brit. 188 = Calandrinia Lockharti.
humile, *Greene, in Coult. Bot. Gaz.* vi. (1881) 183.—N. Mexic.
indicum, Wight & Arn. Prod. 356 = cuneifolium.
lanceolatum, Link, Enum. Hort. Berol. ii. 2 = Anacampseros lanceolata.
Linaria, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 70.—Chili.
lineare, Hoffmgg. Verz. Pfl. 217 = Calandrinia pilosiuscula.

TALINUM:—
lineare, *H. B. & K. Nov. Gen. et Sp.* vi. 77.—Mexic.
linguaeforme, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 81 = Cotyledon linguaefolia.
lingulatum, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 116 = Calandrinia lingulata.
Menziesii, Hook. Fl. Bor. Am. i. 223. t. 70 = Calandrinia Menziesii.
mexicanum, *Hemsl. Diagn. Pl. Nov.* 23.—Mexic.
minimum, *Miers, Trav. Chil.* i. 530, nomen; *et ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 69.—Chili.
minus, *Eckl. & Zeyh. Enum.* 282.—Afr. austr.
monandrum, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 118 = Monocosmia corrigioloides.
mucronatum, *H. B. & K. Nov. Gen. et Sp.* vi. 76.—Mexic.
nanum, Nees, in Lehm. Pl. Preiss. i. 246 = Calandrinia pygmaea.
napiforme, *DC. Prod.* iii. 357.—Mexic.
nitidum, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 117 = Calandrinia nitida.
paniculatum, Gaertn. Fruct. ii. 219. t. 128 = patens.
paniculatum, Moench, Meth. 232 = triangulare.
paniculatum, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 114 = Calandrinia paniculata.
parviflorum, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 197.—Am. bor. occ.
patens, Andr. Bot. Rep. iv. t. 253 = Calandrinia Andrewsii.
patens, *Willd. Sp. Pl.* ii. 863.—Am. trop.
polyandrum, Hook. Bot. Mag. t. 4833 = Calandrinia polyandra.
polyandrum, *Ruiz & Pav. Prod.* 65, in obs.; *Syst. Veg.* 115.—Peruv.
polygaloides, *Gill. ex Arn. in Edinb. Journ. Nat. & Geog. Sc.* iii. (1831) 354–55.—Chili.
polyphyllum, Link, Enum. Hort. Berol. ii. 3 = Anacampseros polyphylla.
purpureum, Fisch. ex Sweet, Hort. Brit. ed. II. 219 (Quid ?).—Hab. ?
pygmaeum, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 407 = Calandrinia pygmaea.
racemosum, Rohrb. in Mart. Fl. Bras. xiv. II. 297 = triangulare.
reflexum, *Cav. Ic.* 1. t. 1.—Am. bor. et austr.
retusum, Willd. Enum. Hort. Berol. Suppl. 26 = Anacampseros arachnoides.
revolutum, *H. B. & K. Nov. Gen. et Sp.* vi. 76.—Venezuela.
rubens, Hort. ex Steud. Nom. ed. II. ii. 661 = Anacampseros arachnoides.
sarmentosum, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 153.—Texas.
spathulatum, A. Gray, Pl. Wright. i. 14 = reflexum.
spinescens, *Torr. in Bot. U. St. Expl. Exped.* ii. 250.—Am. bor. occ.
strictum, *Wall. Cat.* n. 6846.—Ind. or.
teretifolium, *Pursh, Fl. Am. Sept.* ii. 365.—Am. bor.
triangulare, *Willd. Sp. Pl.* ii. 862.—Am. trop.
trichotomum, Desf. Tabl. ed. I. 166 = teretifolium.
trigonum, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 71.—Chili.
umbellatum, Ruiz & Pav. Prod. 65, in obs.; Syst. Veg. 117 = Calandrinia umbellata.

TALIPULIA, Rafin. Fl. Tellur. ii. 17 (1836) = **Aneilema**, R. Br. (Commel.).

TALISIA, Aubl. Pl. Gui. i. 349. t. 136 (1775). *SAPINDACEAE*, Benth. & Hook. f. i. 401.
ACLADODEA, Ruiz & Pav. Prod. 133. t. 29 (1794).
COMATOGLOSSUM, Karst. & Triana, in Linnaea, xxviii. (1856) 436.
Acladodea, *DC. Prod.* i. 609.—Peruv.
acutifolia, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 349.—Bras.
angustifolia, *Radlk. l. c.* 345.—Bras.
carinata, *Radlk. l. c.* 348.—Guian.
cerasina, Radlk. l. c. 347 = Sapindus oblongus, cerasinus.
clathrata, *Radlk. l. c.* 349.—Bras.
coriacea, *Radlk. l. c.* 346.—Bras.
cupularis, *Radlk. l. c.* 350.—Bras.
dasyclada, *Radlk. l. c.* 348.—Bras.
esculenta, Radlk. l. c. 345 = Sapindus esculentus.
firma, *Radlk. l. c.* 346.—Bras.
glabra, DC. Prod. i. 609 = guianensis.
guianensis, *Aubl. Pl. Gui.* i. 349. t. 136.—Guian.

TALISIA:—
hemidasya, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 349.—Guian.
hexaphylla, *Vahl, Eclog. Am.* ii. 29.—Am. austr.
laxiflora, *Benth. in Hook. Kew Journ.* ii. (1850) 212 (al. gen. ?).—Bras.
longifolia, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 348; ix. (1879) 519 = Cupania longifolia.
macrophylla, *Radlk. l. c.* viii. (1878) 347.—Bras.
megaphylla, *Sagot, in Ann. Sc. Nat.* Sér. VI. xii. (1881) 196.—Guian.
mollis, Cambess. in Mém. Mus. Par. xviii. (1829) 48 = guianensis.
multinervis, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 346.—Bras.
oedipoda, *Radlk. l. c.* 347.—Bras.
olivaeformis, *Radlk. l. c.* 342.—N. Granat.
pachycarpa, *Radlk. l. c.* 350.—Guian.
pedicellaris, *Radlk. l. c.* 342.—Guian.
pilosula, *Sagot, in Ann. Sc. Nat.* Sér. VI. xii. (1881) 196.—Guian.
pinnata, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 351 = T. Acladodea.
praealta, Radlk. l. c. 345 = Cupania praealta.
pulverulenta, *Radlk. l. c.* 342.—Guian.
rosea, Vahl, Eclog. Am. ii. 30 = guianensis.
squarrosa, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 346.—Guian.
stricta, *Triana & Planch. ex Radlk. l. c.* 350.—N. Granat.
subalbens, *Radlk. l. c.* 345.—Bras.
sylvatica, Radlk. l. c. 341 = Racaria sylvatica.

TALPINARIA, Karst. Fl. Columb. i. 153. t. 76 (1859 ?) = **Pleurothallis**, R. Br. (Orchid.).
bivalvis, Karst. l. c. = P. bivalvis.

TAMALA, Rafin. Sylva Tellur. 136 (1838) = **Persea**, Plum. (Laurin.).
acuminata, Rafin. l. c. 137 = Nectandra berchemiaefolia.
borbonia, Rafin. l. c. 136 = P. carolinensis.
carolinensis, Rafin. l. c. = P. carolinensis.
palustris, Rafin. l. c. 137 = P. Lingue.

TAMARA, Roxb. ex Steud. Nom. ed. II. ii. 661 (1841) = **Nelumbium**, Juss. (Nymph.).
alba, Roxb. ex Steud. l. c. = N. speciosum.
hemisphaerica, Buch.-Ham. ex Pritz. Ic. Ind. i. 1087 = N. speciosum ?
rubra, Roxb. ex Steud. Nom. ed. II. ii. 661 = N. speciosum.

TAMARINDUS, Tourn. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 581.
indica, *Linn. Sp. Pl.* 34.—As. et Afr. trop.
Intsia, Spreng. Syst. iii. 158 = Afzelia bijuga.
occidentalis, Gaertn. Fruct. ii. 310. t. 146 = indica.
officinalis, Hook. Bot. Mag. t. 4563 = indica.
umbrosa, Salisb. Prod. 323 = indica.

TAMARISCUS, Tourn. ex Hall. Enum. Stirp. Helv. i. 419 (1742) = **Tamarix**, Linn.
decander, Lam. Fl. Fr. iii. 74 = Myricaria germanica.
gallicus, All. Fl. Pedem. ii. 87 = Tamarix gallica.
germanicus, Scop. Fl. Carn. ed. II. i. 224 = Myricaria germanica.
pentander, Lam. Fl. Fr. iii. 73 = Tamarix gallica.

TAMARIX, Linn. Syst. ed. I (1735). *TAMARISCINEAE*, Benth. & Hook. f. i. 160.
TAMARISCUS, Tourn. ex Hall. Enum. Stirp. Helv. i. 419 (1742).
TRICHAURUS, Arn. in Wight & Arn. Prod. 40 (1834).
aegyptiaca, Bertol. Misc. Bot. xiv. 15 = gallica.
affinis, *Bunge, Tent. Gen. Tamaric.* (1852) 36.—Reg. Casp.
africana, Bory & Chaub. Nouv. Fl. Pélop. 20 = tetrandra.
africana, *Poir. Voy. Barb.* ii. 139; *Desf. Fl. Atlant.* i. 269.—Reg. Mediterr.
amplexicaulis, *Ehrenb. in Linnaea*, ii. (1827) 275.—Afr. bor.
anglica, Webb, in Hook. Journ. Bot. iii. (1841) 430 = gallica.

TAMARIX:—
angustifolia, Boiss. Fl. Orient. i. 771 = florida.
angustifolia, Hohen. in Bull. Soc. Nat. Mosc. (1838) 363 = Hohenackeri.
angustifolia, Ledeb. in Eichw. Beschr. Casp.-Cauc. 12 = gracilis.
arabica, *Bunge, Tent. Gen. Tamaric.* (1852) 55.—Arab.
aralensis, *Bunge, l. c.* 59.—Reg. Casp.
arborea, *Ehrenb. ex Bunge, l. c.* 67.—Aegypt.
arceuthoides, *Bunge, in Mém. Sav. Etr. Pétersb.* vii (1851) 295.—As. bor.
articulata, *Vahl, Symb. Bot.* ii. 48. t. 32.—Arab.; Pers.; Ind. or.
articulata, Wall. Cat. n. 3756 B, C = dioica.
bachtiarica, *Bunge, ex Boiss. Fl. Orient.* i. 772.—Persia.
Balansae, *J. Gay, ex Munby, Cat.* 12, nomen.—Alger.
bounopaea, *J. Gay, l. c.*, nomen.—Alger.
Boveana, *Bunge, Tent. Gen. Tamaric.* (1852) 24.—Alger.
brachystachys, *Bunge, l. c.* 26.—Reg. Cauc.
brachystylis, *J. Gay, ex Munby, Cat.* 12, nomen.—Alger.
Bungei, *Boiss. Fl. Orient.* i. 774.—Persia.
canariensis, Willd. in Abh. Akad. Berl. 1812–13 (1816) 79 = gallica.
canescens, Desv. in Ann. Sc. Nat. Sér. I. iv. (1825) 348 = hispida.
chinensis, *Lour. Fl. Cochinch.* 182.—China.
chinensis, Sieb. & Zucc. Fl. Jap. i. 132 = juniperina.
cretica, Bunge, Tent. Gen. Tamaric. (1852) 33 = parviflora.
cupressiformis, *Ledeb. Fl. Alt.* i. 423.—Sibir. altaic.
davurica, Willd. in Abh. Akad. Berl. 1812–13 (1816) 85 = Myricaria davurica.
decandra, Pall. Fl. Ross. ii. 73. t. 80 = Myricaria longifolia.
decandra, Salisb. Prod. 173 = Myricaria germanica.
deserti, Boiss. Diagn. Ser. I. x. 9 = tetragyna.
dioica, *Roxb. Hort. Beng.* 22; *Fl. Ind.* ii. 101.—Ind. or.; Burma.
dubia, *Bunge, Tent. Gen. Tamaric.* (1852) 18.—Persia.
effusa, *Ehrenb. in Linnaea*, ii. (1827) 258.—Aegypt.
Ehrenbergii, Presl, ex Bunge, Tent. Gen. Tamaric. (1852) 54 = gallica.
elegans, Spach, Hist. Vég. Phan. v. 481 = gallica.
elongata, *Ledeb. Fl. Alt.* i. 421.—Sibir. altaic.
epacroides, Sm. in Rees, Cycl. xxxv. n. 4 = gallica.
ericoides, *Rottl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 214.—Ind. or.
Eversmanni, Presl, ex Ledeb. Fl. Ross. ii. 135 = gallica.
Faras, Edgew. in Journ. Linn. Soc. vi. (1862) 188 = articulata.
floribunda, Stev. ex Bunge, Tent. Gen. Tamaric. (1852) 44 = Hohenackeri ?
florida, *Bunge, l. c.* 37.—Persia.
Furas, Buch.-Ham. ex Royle, Illustr. Bot. Himal. 213 = articulata.
gallica, [Habl.] Beschr. Taur. 105 = tetrandra.
gallica, *Linn. Sp. Pl.* 270.—Europ.; As.; Afr.
gallica, Sieber, ex Nym. Consp. 253 = parviflora.
gallica, Thunb. Fl. Jap. 126 = chinensis.
germanica, S. T. Gmel. Reise, iv. 63 = seq. ?
germanica, Linn. Sp. Pl. 271 = Myricaria germanica.
germanica, Pall. Reise, iii. 320 = Myricaria longifolia.
germanica, Schang. in Pall. N. Nord. Beitr. vi. 49 = Myricaria davurica.
gracilis, *Willd. in Abh. Akad. Berl.* 1812–13 (1816) 81.—Sibir. altaic.
Hampeana, *Boiss. & Heldr. Diagn.* Ser. I. x. 8.—As. Min.
herbacea, Willd. in Abh. Akad. Berl. 1812–13 (1816) 84 = Myricaria germanica ?
hexandra, Hampe, in Flora, xxv. (1842) 62 = Hampeana.
hispanica, *Boiss. Diagn.* Ser. II. ii. 56.—Hispan.
hispida, *Willd. in Abh. Akad. Berl.* 1812–13 (1816) 77.—Reg. Casp.
Hohenackeri, *Bunge, Tent. Gen. Tamaric.* (1852) 44.—Reg. Cauc.
indica, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 214 = gallica.
ispahanica, *Bunge, ex Boiss. Fl. Orient.* i. 768.—Persia.
Jordanis, *Boiss. l. c.* 771.—Syria.
juniperina, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 102.—China.

TAMARIX:—
Karelini, *Bunge, l. c.* vii. (1851) 294.—Reg. Casp.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 582.—Turkestan.
Kotschyi, *Bunge, Tent. Gen. Tamaric.* (1852) 30.—Persia.
laxa, *Willd. in Abh. Akad. Berl.* 1812–13 (1816) 82.—Ross.; Sibir. altaic.
leptopetala, *Bunge, Tent. Gen. Tamaric.* (1852) 72.—Persia.
leptostachys, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 293.—Turkestan.
Lessingii, Presl, ex Bunge, Tent. Gen. Tamaric. (1852) 51 = gallica.
libanotica, Hort. ex C. Koch, Dendrol. i. 454 = gallica.
longifolia, Willd. in Abh. Akad. Berl. 1812–13 (1816) 85 = Myricaria longifolia.
longipes, Presl, ex Bunge, Tent. Gen. Tamaric. (1852) 21 = Hampeana.
macrocarpa, *Ehrenb. ex Bunge, l. c.* 79.—Aegypt.; Persia.
mannifera, *Ehrenb. ex Bunge, l. c.* 63.—Aegypt.; Arab.; Persia.
mannifera, Kotschy, ex Bunge, l. c. 52 = gallica.
mascatensis, *Bunge, l. c.* 60.—Arab.
Meyeri, Boiss. Diagn. Ser. I. x. 9 = tetragyna.
monogyna, Stokes, Bot. Mat. Med. ii. 176 = Myricaria germanica.
mucronata, Sm. in Rees, Cycl. xxxv. n. 5 = ericoides.
nilotica, Ehrenb. ex Bunge, Tent. Gen. Tamaric (1852) 54 = gallica.
Noëana, Boiss. Diagn. Ser. II. ii. 56 = tetragyna.
octandra, *Bunge, Tent. Gen. Tamaric.* (1852) 17.—Persia.
odessana, *Stev. ex Bunge, l. c.* 47.—Ross.
orientalis, Forsk. Fl. Aegypt. Arab. 206 = articulata.
palaestina, Bertol. Misc. Bot. xiv. 16 = gallica.
Pallasii, Desv. in Ann. Sc. Nat. Sér. I. iv. (1825) 349 = gallica.
paniculata, Stev. ex DC. Prod. iii. 96 = gallica.
parviflora, *DC. l. c.* 97.—Graecia.
passerinoides, *Delile, Fl. Aegypt. Illustr.* 58.—Aegypt.; Arab.
pauciovulata, *J. Gay, ex Munby, Cat.* 12, nomen.—Alger.
pentandra, Hampe, ex Bunge, Tent. Gen. Tamaric. (1852) 32 = parviflora.
pentandra, Pall. Fl. Ross. i. II. 72; Salisb. Prod. 173 = gallica.
Petteri, Presl, ex Bunge, Tent. Gen. Tamaric. (1852) 32 = parviflora.
Pharas, Buch.-Ham. ex Wall. Cat. n. 3758 = articulata.
Pharos, Steud. Nom. ed. II. ii. 661, sphalm. = praec.
plumosa, Carr. in Rev. Hortic. (1868) 358 = gallica?
polystachya, Ledeb. Fl. Ross. ii. 133 = laxa.
praecox, Portenschl. ex Bunge, Tent. Gen. Tamaric. (1852) 40 = africana.
pycnocarpa, DC. Prod. iii. 97 = passerinoides.
ramosissima, Karel. ex Boiss. Fl. Orient. i. 776 = Karelini.
ramosissima, Ledeb. Fl. Alt. i. 424 = gallica.
rosea, *Bunge, Tent. Gen. Tamaric.* (1852) 19.—Persia.
salina, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 248.—Ind. bor. occ.
salsa, Stev. ex Bunge, Tent. Gen. Tamaric. (1852) 24 = tetragyna.
senegalensis, DC. Prod. iii. 96 = gallica.
serotina, *Bunge, ex Boiss. Fl. Orient.* i. 773.—Persia.
smyrnensis, *Bunge, Tent. Gen. Tamaric.* (1852) 53.—Smyrna.
songarica, Eversm. ex Steud. Nom. ed. II. ii. 661 = Haloxylon Ammodendron.
soongorica, Pall. in Nov. Act. Acad. Petrop. x. (1797) 374 = Hololachna songorica.
speciosa, *Ball, in Journ. Linn. Soc.* xvi. (1877) 372.—Marocc.
speciosa, Hort. ex C. Koch, Dendrol. i. 453 = tetrandra.
squamosa, Steud. Nom. ed. II. ii. 661 = Myricaria germanica.
stricta, *Boiss. Diagn.* Ser. II. ii. 57.—Sind.; Beluchist.
syriaca, *Stev. ex Bunge, Tent. Gen. Tamaric.* (1852) 21, in syn.; *Boiss. Fl. Orient.* i. 767.—Syria.
Szovitsiana, *Bunge, Tent. Gen. Tamaric.* (1852) 26.—Persia.
taurica, Pall. in Nov. Act. Acad. Petrop. x. (1797) 276, in nota = tetrandra.
tenacissima, Buch.-Ham. ex Wall. Cat. n. 3757 = ericoides.

TAMARIX:—
tetragyna, Buhse, ex Bunge, Tent. Gen. Tamaric. (1852) 19 = dubia.
tetragyna, *Ehrenb. in Linnaea*, ii. (1827) 257, 258.—Aegypt.; Arab.
tetragyna, Szovits, ex Bunge, Tent. Gen. Tamaric. (1852) 18 = octandra.
tetramera, Kar. in Bull. Soc. Nat. Mosc. (1839) 155, nomen (Quid?).—Turkestan.
tetrandra, Guebh. ex Bunge, Tent. Gen. Tamaric. (1852) 52 = gallica.
tetrandra, *Pall. ex Bieb. Fl. Taur. Cauc.* i. 247.—Europ. or
tetrandra, Szovits, ex Bunge, Tent. Gen. Tamaric. (1852) 48 = odessana.
tomentosa, Sm. in Rees, Cyclop. xxxv. n. 2 = hispida.
usneoides, E. Mey. in Drège, Zwei Pfl. Docum. 225 = articulata.
variabilis, Hort. ex C. Koch, Dendrol. i. 457 = tetragyna.

TAMATAVIA, Hook. f. in Benth. & Hook. f. Gen. ii. 92 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
Melleri, *Hook. f. l. c.*; *et in Hook. Ic. Pl.* t. 1090.—Madag.

TAMBOURISSA, Sonner. Voy. Ind. Or. iii. 267. t. 134 (1782). *MONIMIACEAE*, Benth. & Hook. f. iii. 139.
AMBORA, Juss. Gen. 401 (1789).
MITHRIDATEA, Comm. ex Schreb. Gen. 783 (1791).
alternifolia, A. DC. in DC. Prod. xvi. II. 660 = quadrifida?
amplifolia, *A. DC. l. c.* 659.—Ins. Maurit.
Boivinii, *A. DC. l. c.*—Madag.
elliptica, *A. DC. l. c.* 660.—Ins. Mascar.
Ficus, *A. DC. l. c.* 659.—Java.
leptophylla, *A. DC. l. c.* 658.—Ins. Comor.
neglecta, A. DC. l. c. 659 = quadrifida.
obovata, A. DC. l. c. 660 = quadrifida.
parvifolia, *Baker, in Journ. Bot.* xx. (1882) 267.—Madag.
pedicellata, *Baker, Fl. Maurit.* 289.—Ins. Maurit.
peltata, *Baker, l. c.* 288.—Ins. Maurit.
purpurea, *A. DC. in DC. Prod.* xvi. II. 659.—Madag.
quadrifida, *Sonner. Voy. Ind. Or.* iii. 267. t. 134.—Ins. Mascar.
religiosa, *A. DC. in DC. Prod.* xvi. II. 659.—Madag.
Rota, *Baker, in Journ. Linn. Soc.* xx. (1883) 240.—Madag.
Sieberi, *A. DC. in DC. Prod.* xvi. II. 660.—Ins. Maurit.
tetragona, *A. DC. l. c.* 659.—Ins. Maurit.
trichophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 240.—Madag.
vestita, *A. DC. in DC. Prod.* xvi. II. 660.—Ins. Borbon.

TAMMSIA, Karst. Fl. Columb. i. 179. t. 89 (1861?). *RUBIACEAE*, Benth. & Hook. f. ii. 79.
anomala, *Karst. l. c.*—N. Granat.

TAMNUS, Tourn. ex Linn. Syst. ed. I (1735) = **Tamus**, Linn. (Dioscor.).

TAMONEA, Aubl. Pl. Gui. i. 441. t. 175 (1775) = **Miconia**, Ruiz & Pav. (Melastom.).
guianensis, Aubl. l. c. = M. Fothergilla.
lasiopetala, DC. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 103 = M. platyhedra.

TAMONEA, Aubl. Pl. Gui. ii. 659. t. 268 (1775). *VERBENACEAE*, Benth. & Hook. f. ii. 1147.
GHINIA, Schreb. Gen. 19 (1789).
ISCHNIA, DC. ex Meissn. Gen. 298; Comm. 206 (1840); et Prod. ix. 257 (1845).
KAEMPFERA, Houst. Reliq. 3. t. 2 (1781).
LEPTOCARPUS, Willd. in Link, Jahrb. i. III. (1820) 51.
arabica, Mirb. Hist. Pl. ed. II. xv. 233 = Priva leptostachys.
curasavica, Pers. Syn. ii. 139 = spinosa.
juncea, *Schau. in DC. Prod.* xi. 529.—Bras.
lappulacea, Poir. Encyc. vii. 568 = Priva echinata.
mutica, Sw. Prod. Veg. Ind. Occ. 94 = spicata.
scabra, *Schlecht. & Cham. in Linnaea*, v. (1830) 99.—Mexic.
spicata, *Aubl. Pl. Gui.* ii. 660. t. 268.—Guian.
spinosa, *Sw. Prod. Veg. Ind. Occ.* 94.—Ind occ.
verbenacea, Sw. Fl. Ind. Occ. ii. 1089, in syn. = spinosa.

TAMONOPSIS, Griseb. in Goett. Abh. xix. (1874) 246 = **Lantana**, Linn. (Verben.).
spicata, Griseb. l. c.—Reg. Argent.

TAMPOA, Aubl. Pl. Gui. ii. Suppl. 35 (1775). *INCERT. SEDIS.*
guyanensis, *Aubl. l. c.*—Guian.

TAMUS, Linn. Gen. ed. I. 305 (1737). *DIOSCOREACEAE*, Benth. & Hook. f. iii. 744.
TAMNUS, Tourn. ex Linn. Syst. ed. I (1735).
canariensis, Willd. ex Kunth, Enum. Pl. v. 455 (Quid?).—Ins. Canar.
communis, Link, in Buch, Beschr. Canar. Ins. 140 = edulis.
communis, *Linn. Sp. Pl.* 1028.—Europ.; Persia; Afr. bor.
cordifolia, Stokes, Bot. Mat. Med. iv. 544 = communis.
cretica, Linn. Sp. Pl. 1028 = communis.
edulis, *Lowe, in Trans. Camb. Phil. Soc.* iv. I. (1833) 12.—Ins. Mader.
Elephantipes, L'Hérit. Sert. Angl. 29 = Testudinaria Elephantipes.
Norsa, Lowe, in Trans. Camb. Phil. Soc. vi. III. (1838), reimp. 21 = edulis.
parviflora, *Kunth, Enum. Pl.* v. 454.—Ins. Teneriff.
racemosa, Gouan, Fl. Monsp. 426 = communis.
sylvestris, Hort. Berol. ex Kunth, Enum. Pl. v. 443 = Testudinaria sylvatica.

TANACETUM, Tourn. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 434.
HEMIPAPPUS, C. Koch, in Linnaea, xxiv. (1851) 340.
MATRICARIOIDES, Spach, Hist. Vég. Phan. x. 24 (1841).
OMALANTHUS, Less. Syn. Comp. 260 (1832).
OMALOTES, DC. Prod. vi. 83 (1837).
PSANACETUM, Neck. Elem. i. 89 (1790).
SPHAEROMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 401.
Achilleae, Sch. Bip. Tanacet. 58 = Chrysanthemum Achilleae.
achilleaefolium, Sch. Bip. l. c. 47 = Chrysanthemum achilleaefolium.
achilleoides, *DC. Prod.* vi. 130.—Mongol.
acutilobum, DC. l. c. 132 = Matricaria acutiloba.
aegyptiacum, Juss. ex Jacq. Hort. Vindob. iii. 46. t. 88 = Grangea maderaspatana.
alatavicum, *Herd. in Bull. Soc. Nat. Mosc.* xl. (1867) II. 129.—Sibir. ural.
albidum, DC. Prod. vi. 132 = Matricaria albida.
alpestre, Pall. ex Ledeb. Fl. Ross. ii. 602 = boreale.
alpinum, Sch. Bip. Tanacet. 61 = Chrysanthemum alpinum.
angulatum, Willd. Sp. Pl. iii. 1811 = Achillea filipendulina.
annuum, *Linn. Sp. Pl.* 844.—Europ. austr.
arcticum, Sch. Bip. Tanacet. 35 = Chrysanthemum arcticum.
argenteum, *Willd. Ach.* 51. t. 2. f. 4.—As. Min.
argyraeum, DC. Prod. vi. 134 = Schistostephium flabelliforme.
argyranthemoides, *Sch. Bip. in Pollichia*, xx-xxi. (1863) 444.—Armenia.
armenum, Sch. Bip. Tanacet. 35 = Chrysanthemum fruticulosum.
artemisioides, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 318.—Reg. Himal.
atratum, Sch. Bip. Tanacet. 62 = Chrysanthemum atratum.
Aucheri, *DC. Prod.* vii. 298.—Syria.
Aucherianum, *Sch. Bip. Tanacet.* 49.—As. Min.
Audiberti, *DC. Prod.* vi. 131.—Reg. Mediterr.
axillare, Thunb. Prod. Pl. Cap. 147 = Stilpnophytum axillare.
Balsamita, Linn. Sp. Pl. 845 = Chrysanthemum Balsamita.
balsamitoides, Sch. Bip. Tanacet. 51 = Chrysanthemum Balsamita.
balsamitum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = praec.
Barclayanum, *DC. Prod.* iv. 128.—Sibir.
bipinnatum, Sch. Bip. Tanacet. 48 = Chrysanthemum bipinnatum.
Bocconi, Sch. Bip. l. c. = Chrysanthemum Bocconi.

TANACETUM:—
boreale, *Fisch. ex Link, Enum. Hort. Berol.* ii. 313.—Podol.; Sibir.
boreale, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 401 = huronense.
Burchellii, DC. Prod. vi. 132 = Matricaria Burchellii.
camphoratum, *Less. in Linnaea*, vi. (1831) 521.—Calif.
canariense, DC. Cat. Hort. Monsp. (1813) 149 = Gonospermum canariense.
canescens, *DC. Prod.* vi. 129.—Persia.
canum, *Eaton, in Bot. King's Exp.* 179. t. 19. f. 8–14.—Am. bor. occ.
capense, *Burm. f. Fl. Cap. Prod.* 24.—Afr. austr.
capitatum, *Torr. & Gray, Fl. N. Am.* ii. 415.—Am. bor.
cappadocicum, *Sch. Bip. Tanacet.* 48.—As. Min.
Capusi, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 308.—Turkestan.
carneum, Sch. Bip. Tanacet. 58 = Chrysanthemum coronopifolium, *Willd.*
caucasicum, Sch. Bip. l. c. 60 = Chrysanthemum caucasicum.
ceratophylloides, Sch. Bip. l. c. 35 = Chrysanthemum coronopifolium, *Vill.*
chiliophyllum, Sch. Bip. l. c. 47 = Chrysanthemum millefoliatum.
chinense, A. Gray, ex Maxim. in Bull. Acad. Pétersb. xvii. (1872) 427 = Crossostephium artemisioides.
chrysanthemoides, Gaertn. Fruct. ii. 396. t. 165 = Chrysanthemum flosculosum.
cinerariaefolium, Sch. Bip. Tanacet. 58 = Chrysanthemum cinerariaefolium.
cinereum, DC. Prod. vi. 131 = Cotula cinerea.
consanguineum, DC. l. c. 133 = Schistostephium crataegifolium.
corymbosum, Sch. Bip. Tanacet. 57 = Chrysanthemum corymbosum.
cotuloides, Linn. Mant. ii. 282 = Hippia gracilis.
crataegifolium, DC. Prod. vi. 134 = Schistostephium crataegifolium.
crispum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 (Quid?).—Java.
crispum, Steud. Nom. ed. I. 825; Hoffmgg. Verz. Pfl. 116 = vulgare.
crithmifolium, Linn. Sp. Pl. 843 = Athanasia parviflora, *Thunb.*
decipiens, Sch. Bip. Tanacet. 47 = Matricaria decipiens.
densum, Sch. Bip. l. c. 48 = Chrysanthemum densum.
discoideum, Reichb. f. Ic. Fl. Germ. xvi. 51. t. 988 = Chrysanthemum discoideum.
diversifolium, *Eaton, in Bot. King's Exp.* 180. t. 19.—Am. bor. occ.
Douglasii, DC. Prod. iv. 128 = huronense.
dumosum, *Boiss. Diagn.* Ser. I. vi. 89.—Persia.
elatum, Salisb. Prod. 190 = vulgare.
elegans, Decne. in Fl. des Serres, xii. (1857) 19. t. 1191 = camphoratum.
elongatum, Sch. Bip. Tanacet. 61 = Matricaria oreades.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 320.—Reg. Himal.
fasciculatum, Sch. Bip. Tanacet. 11 = Artemisia fasciculata.
ferulaceum, *Sch. Bip. l. c.* 50.—Ins. Canar.
Fisherae, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 170.—Afghan.
flabellifolium, *Boiss. & Heldr. Diagn.* Ser. I. xi. 26.—As. Min.
flabelliforme, L'Hérit. Sert. Angl. 21 = Pentzia crenata.
frutescens, Linn. Sp. Pl. 844 = Hippia frutescens.
fruticosum, Harv. in Harv. & Sond. Fl. Cap. iii. 166, in syn., sphalm. = Matricaria multiflora.
fruticosum, Link, in Buch, Beschr. Canar. Ins. 149 = Gonospermum canariense.
fruticulosum, C. B. Clarke, Comp. Ind. 153 = gracile.
fruticulosum, *Ledeb. Fl. Alt.* iv. 58.—Reg. Himal.; Soongar.
Funkii, *Sch. Bip. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 102.—Hispan.
Gayanum, Sch. Bip. in Bonplandia, vii. (1860) 369 = Chrysanthemum Mawii.
globiferum, DC. 6me Not. Jard. Genèv. 8 = Matricaria globifera.
Gmelini, Sch. Bip. Tanacet. 35 = Chrysanthemum arcticum.

TANACETUM:—
gnaphaloides, Steud. in Flora, xiii. (1830) 542 = Helichrysum anomalum.
gossypinum, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 154.—Reg. Himal.
gracile, *Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind.* iii. 318.—Reg. Himal.
graminifolium, Reichb. f. Ic. Fl. Germ. xvi. 51 = Chrysanthemum graminifolium.
grandiflorum, Poir. Encyc. vii. 574 (= *Plagius grandiflorus*).—Afr. bor.
grandiflorum, Thunb. Prod. Pl. Cap. 147 = Matricaria grandiflora.
gratum, Wall. ex DC. Prod. v. 374 = Cyathocline lyrata.
griseum, Harv. in Harv. & Sond. Fl. Cap. iii. 168 (Schistostephii sp.).—Afr. austr.
heptalobum, DC. Prod. vi. 133 (Schistostephii sp.).—Afr. austr.
heptalobum, Sch. Bip. in Flora, xxvii. (1844) 675 = *hippiaefolium*.
Herderi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 619.—Turkestan.
heterophyllum, Boiss. Diagn. Ser. I. vi. 90 = myriophyllum.
hippiaefolium, DC. Prod. vi. 133 (Schistostephii sp.).—Afr. austr.
hispidum, DC. l. c. = Cenia hispida.
humile, Forsk. Fl. Aegypt. Arab. 148 = Grangea maderaspatana.
huronense, *Nutt. Gen. Am.* ii. 141.—Am. bor.
huronense, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 401, partim = camphoratum.
incanum, Fisch. & Mey. ex DC. Prod. vi. 129 = canescens.
incanum, Hort. ex DC. l. c. 128 = Barclayanum.
incanum, Linn. Sp. Pl. 844 = Artemisia fasciculata.
indicum, Sch. Bip. Tanacet. 50 = Chrysanthemum indicum.
integrifolium, Sch. Bip. l. c. 35 = Chrysanthemum integrifolium.
Johnstonii, *Hemsl. in Journ. Bot.* xxi. (1883) 135.—Afghan.
Kochii, Sch. Bip. Tanacet. 47 = Chrysanthemum millefoliatum.
Kotschyi, Boiss. Fl. Orient. iii. 351 = myriophyllum.
Kotzebuense, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 107; et ex DC. Prod. vi. 131 = Chrysanthemum bipinnatum.
lanuginosum, *Sch. Bip. & Herd. in Pollichia*, xx–xxi. (1863) 442.—Sibir.
larvatum, Kanitz, ex Nym. Consp. 371 = Chrysanthemum coronopifolium.
latifolium, Sch. Bip. Tanacet. 35 = Chrysanthemum lacustre.
Ledebourii, *Sch. Bip. l. c.* 47.—Reg. Casp.
leptolobum, DC. Prod. vi. 133 = Matricaria multiflora.
leptophyllum, *Sch. Bip. Tanacet.* 48.—Reg. Cauc.
Leucanthemum, Sch. Bip. l. c. 35 = Chrysanthemum Leucanthemum.
leucophyllum, *Regel, in Act. Hort. Petrop.* vii. (1880) 551.—Turkestan.
linearilobum, DC. Prod. vi. 133 = Cenia sericea.
linifolium, Thunb. Prod. Pl. Cap. 147 = Stilpnophytum linifolium.
longifolium, Thunb. l. c. = Stilpnophytum linifolium.
longifolium, *Wall. Cat.* n. 3231.—Reg. Himal.
macrophyllum, Fisch. ex Steud. Nom. ed. I. 825, nomen (Quid?).—Hab.?
macrophyllum, Sch. Bip. Tanacet. 53 = Chrysanthemum macrophyllum.
Maresii, Sch. Bip. in Bonplandia, viii. (1860) 369 = Chrysanthemum Maresii.
marginatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 177.—Borneo.
matricarioides, Ledeb. Fl. Ross. ii. 544 = Matricaria discoidea.
maximum, Sch. Bip. Tanacet. 35 = Chrysanthemum maximum.
Meyerianum, *Sch. Bip. l. c.* 47.—Sibir.
microphyllum, *DC. Prod.* vi. 131.—Hispan.
millefoliatum, *Fisch. & Mey. ex DC. l. c.* 128.—Armenia.
millefolium, Aitch. in Journ. Linn. Soc. xix. (1882) 169 = Achillea micrantha.
minimum, Sch. Bip. Tanacet. 60 = Chrysanthemum tomentosum.

TANACETUM:—
monanthos, Linn. Mant. i. 111 = Anacyclus alexandrinus.
monspeliense, Sch. Bip. Tanacet. 35 = Chrysanthemum monspeliense.
montanum, Sch. Bip. l. c. = Chrysanthemum montanum.
multicaule, Sch. Bip. l. c. 48 = Chrysanthemum multicaule.
multiflorum, Thunb. Prod. Pl. Cap. 147 = Matricaria multiflora.
Myconis, Schenk, Ind. Sem. Hort. Wirceb. (1861) = Chrysanthemum Myconis.
myriophyllum, *Willd. Sp. Pl.* iii. 1814.—Reg. Cauc.
nanum, *C. B. Clarke, Comp. Ind.* 155.—Reg. Himal.
nivale, *Sch. Bip. Tanacet.* 60.—Armenia.
niveum, Sch. Bip. l. c. 56 = Chrysanthemum praealtum.
nubigenum, *Wall. ex DC. Prod.* vi. 130.—Reg. Himal.
Nuttallii, *Torr. & Gray, Fl. N. Am.* ii. 415.—Am. bor. occ.
obtusum, Thunb. Prod. Pl. Cap. 147 = Matricaria globifera.
officinale, Gueldenst. ex Ledeb. Fl. Ross. ii. 601 = vulgare.
officinarum, Crantz, Inst. i. 273 = vulgare.
oligocephalum, Sch. Bip. Tanacet. 48 = Chrysanthemum millefoliatum.
orientale, Willd. Sp. Pl. iii. 1812 = Artemisia fasciculata.
ovatifolium, Stokes, Bot. Mat. Med. iv. 180 = Chrysanthemum Balsamita.
Pallasianum, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 55).—Manchuria.
pallens, Sch. Bip. Tanacet. 35 = Chrysanthemum pallens.
palmatifidum, *Sch. Bip. l. c.* 60.—Hispan.
palustre, Sch. Bip. l. c. 49 = Chrysanthemum palustre.
parthenifolium, Sch. Bip. l. c. 56 = Chrysanthemum praealtum.
Parthenium, Sch. Bip. l. c. 55 = Chrysanthemum Parthenium.
pauciflorum, DC. Prod. vi. 131 = Matricaria discoidea.
pauciflorum, Fisch. Hort. Gorenk. ed. II. (1812) 37 (Quid?).—Hab.?
pauciflorum, Richards. in Frankl. Narr. 1st Journ. App. ed. II. 30 = huronense.
pilosum, Berg. Desc. Pl. Cap. 244 = Hippia gracilis.
pinnatifidum, Sch. Bip. Tanacet. 52 = Chrysanthemum multifidum.
pinnatum, *Boiss. Diagn.* Ser. I. vi. 90.—Persia.
polycephalum, Sch. Bip. Tanacet. 47 (= *Pyrethrum polycephalum*).—Mesopot.
potentilloides, *A. Gray, in Proc. Am. Acad.* ix. (1874) 204.—Am. bor. occ.
Pseudo-parthenium, Sch. Bip. in Pollichia, xx–xxi. (1863) 444 = nivale.
ptarmicaeflorum, Sch. Bip. Tanacet. 50 (= *Pyrethrum ptarmicaeflorum*).—Ins. Canar.
pulchellum, *Sch. Bip. l. c.* 61.—Sibir.
pulchrum, *Sch. Bip. l. c.* 49.—Sibir. altaic.
pulverulentum, Sch. Bip. l. c. 48 = Chrysanthemum pulverulentum.
purpureum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 181 = Cyathochine lyrata.
pyrethroides, Sch. Bip. in Pollichia, xx–xxi. (1863) 441 = Chrysanthemum Richteria.
radicans, Sch. Bip. Tanacet. 48 = Chrysanthemum radicans.
robustum, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 154.—Reg. Himal.
roseum, Sch. Bip. Tanacet. 58 = Chrysanthemum coccineum.
rotundifolium, DC. Prod. vi. 133 = Schistostephium rotundifolium.
saxatile, Sch. Bip. Tanacet. 47 = Chrysanthemum millefoliatum.
Scharnhorsti, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 620.—Turkestan.
Semenovii, *Herd. in Bull. Soc. Nat. Mosc.* xl. (1867) II. 130.—As. bor.
Senecionis, *J. Gay, ex DC. Prod.* vi. 129.—Reg. Himal.
sericeum, Sch. Bip. Tanacet. 59 = Chrysanthemum argenteum.
serotinum, Sch. Bip. l. c. 35 = Chrysanthemum serotinum.
setaceum, Tausch, in Flora, xii. (1829) 11 = myriophyllum.

TANACETUM:—
sibiricum, Falk, Topogr. ii. n. 942; ex Ledeb. Fl. Alt. iv. 60 = vulgare.
sibiricum, *Linn. Sp. Pl.* 844.—Sibir.
siculum, P. G. Strobl, in Flora, lxv. (1882) 249 = vulgare.
silaifolium, Sch. Bip. Tanacet. 61 = Matricaria grandiflora.
simplicifolium, Salisb. Prod. 190 = Chrysanthemum Balsamita.
sinaicum, *Delile, ex DC. Prod.* vi. 59.—Arab.
sinense, Sch. Bip. Tanacet. 50 = Chrysanthemum sinense.
sinuatum, Sch. Bip. l. c. 35 = Chrysanthemum sinuatum.
suaveolens, Hook. Fl. Bor. Am. i. 327. t. 110 = Matricaria discoidea.
suffruticosum, Burm. ex DC. Prod. vi. 144 = Hippia gracilis.
suffruticosum, Linn. Sp. Pl. 843 = Matricaria multiflora.
suffruticosum, Thunb. Fl. Cap. ed. Schult. 641, partim = Hippia frutescens.
sulphureum, *Sch. Bip. Tanacet.* 48.—Hispan.
tauricum, Sch. Bip. l. c. = Chrysanthemum millefoliatum.
tenuifolium, *Jacquem. ex DC. Prod.* vi. 129.—Reg. Himal.
tenuisectum, Sch. Bip. in Pollichia, xx–xxi. (1863) 444 = Chrysanthemum coronopifolium, *Vill.*
tibeticum, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 154.—Reg. Himal.
tomentellum, *Kotschy, ex Boiss. Fl. Orient.* iii. 354.—Armenia.
tomentosum, DC. Prod. vi. 130 = T. Senecionis.
tomentosum, Regel & Herd. in Bull. Soc. Nat. Mosc. xl. (1867) ii. 128 = Herderi.
tomentosum, Siev. in Pall. N. Nord. Beitr. vii. 305 = fruticulosum.
tortuosum, DC. Prod. vi. 133 = Pentzia tortuosa.
Tournefortii, Sch. Bip. Tanacet. 59 = Chrysanthemum argenteum.
transiliense, *Herd. in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 129.—As. bor.
trichophyllum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 255.—Turkestan.
trifidum, *DC. Prod.* vi. 130.—Mongol.
trifidum, Franch. Pl. Davidian. i. (1884) 172 = praec.?
trifurcatum, Sch. Bip. Tanacet. 48 = Chrysanthemum trifurcatum.
uliginosum, Sibth. & Sm. Fl. Graec. Prod. ii. 167 = Cotula tridentata.
umbellatum, Gilib. Fl. Lituan. i. 171 = vulgare.
umbelliferum, *Boiss. Diagn.* Ser. II. iii. 30.—Afghan.
uniflorum, *Sch. Bip. Tanacet.* 48.—Persia.
Vahlii, *DC. Prod.* vi. 129.—Hispan.
vestitum, Thunb. Prod. Pl. Cap. 147 = Athanasia fasciculata.
virgatum, Loisel. Fl. Gall. ed. I. 551 = Chrysanthemum flosculosum.
viscosum, Wall. Cat. n. 3232 = Cyathocline lyrata.
vulgare, *Linn. Sp. Pl.* 844.—Europ.; As. bor.
Waldsteinii, Sch. Bip. Tanacet. 35 = Chrysanthemum rotundifolium.
Willkommii, Sch. Bip. ex Willk. in Flora, xxxiv. (1851) 748 = Vahlii.

TANAECIUM, Sw. Prod. Veg. Ind. Occ. 91 (1788). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1039.
albiflorum, DC. Prod. ix. 245 = T. Jaroba.
brasiliense, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. viii. (1861) 118.—Bras.
crucigerum, *Seem. in Bonplandia*, iv. (1856) 127.—Ind. occ.
exsertum, *Griseb. Fl. Brit. W. Ind.* 450.—Jamaic.
Jaroba, *Sw. Prod. Veg. Ind. Occ.* 92.—Am. trop.
lilacinum, *Seem. Bot. Voy. Herald*, 182.—Panama.
paniculatum, Sieber, ex DC. Prod. ix. 245 = Jacaranda Sagraeana.
parasiticum, Sw. Prod. Veg. Ind. Occ. 92 = Schlegelia parasitica.
pinnatum, Willd. Sp. Pl. iii. 312 = Kigelia pinnata.
praelongum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. viii. (1861) 117.—Guian.
Tripinna, Raeusch. Nom. ed. III. 178 = Colea tripinnata.

TANAKAEA, Franch. & Sav. Enum. Pl. Jap. ii. 352 (1876), err. typ. = seq.

TANAKEA, Franch. & Sav. Enum. Pl. Jap. i. 144 (1875). *SAXIFRAGEAE.*
radicans, *Franch. & Sav. l. c.*—Japon.

TANAXION, Rafin. Fl. Tellur. ii. 51 (1836) = **Erigeron**, Linn. (Compos.).
tomentosum, Rafin. l. c. = E. bahamensis.

TANDONIA, Baill. Adansonia, i. (1860–61) 184 = **Tannodia**, Baill. (Euphorb.).
cordifolia, Baill. l. c. 185 = Tannodia cordifolia.

TANDONIA, Moq. in DC. Prod. xiii. ii. 226 (1849) = **Boussingaultia**, H. B. & K. (Chenop.).
brachystachya, Moq. l. c. 227.—N. Granat.
cordifolia, Moq. l. c. 228.—N. Granat.
diffusa, Moq. l. c. 226.—Peruv.
filiformis, Moq. l. c. 227.—Peruv.
marginata, Moq. l. c. 226.—Ecuador.
obovata, Moq. l. c. 227.—N. Granat.
ramosa, Moq. l. c. = B. ramosa.

TANGARACA, Adans. Fam. ii. 143 (1763) = **Hamelia**, Jacq. (Rubiac.).

TANGHINIA, Thou. Gen. Nov. Madag. 10 (1806) = **Cerbera**, Linn. (Apocyn.).
dichotoma, G. Don, in Sweet, Hort. Brit. ed. III. 461 = Tabernaemontana dichotoma.
lactaria, G. Don, l. c. = C. Odollam.
laurifolia, G. Don, Gen. Syst. iv. 98 = C. Odollam.
maculata, G. Don, in Sweet, Hort. Brit. ed. III. 461 = Ochrosia borbonica.
Manghas, G. Don, Gen. Syst. iv. 98 = C. Odollam.
Odollam, G. Don, l. c. = C. Odollam.
venenefiua, G. Don, l. c. sphalm. = seq.
venenifera, Poir. Encyc. Suppl. v. 283 = C. Tanghin.

TANIBOUCA, Aubl. Pl. Gui. 448. t. 178 (1775) = **Terminalia**, Linn. (Combret.).
guianensis, Aubl. l. c. = Term. Tanibouca.

TANKERVILLIA, Link, Handb. i. 251 (1829) = **Phaius**, Lour. (Orchid.).
cantoniensis, Link, l. c. = P. grandifolius.

TANNODIA, Baill. Adansonia, i. (1860–61) 251. t. 7. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 304.
TANDONIA, Baill. Adansonia, i. (1860–61) 184.
cordifolia, *Baill. l. c.* 251.—Madag.

TANROUJOU, Juss. Gen. 351, in adnot. (1789) = **Hymenaea**, Linn. (Legumin.).

TANTALUS, Noronha, ex Thou. Hist. Vég. Afr. 39 (1806) = **Sarcolaena**, Thou. (Chlaen.).

TANULEPIS, Balf. f. in Journ. Linn. Soc. xvi. (1877) 17; et in Baker, Fl. Maurit. 225 (1877). *ASCLEPIADEAE.*
sphenophylla, *Balf. f. ll. cc.*—Ins. Rodriguez.

TAONABO, Aubl. Pl. Gui. i. 569. tt. 227, 228 (1775) = **Ternstroemia**, Mutis.
dentata, Aubl. l. c. 569 = Ternstr. dentata.
punctata, Aubl. l. c. 571 = Ternstr. punctata.

TAPANAVA, Adans. Fam. ii. 470 (1763) = **Pothos**, Linn. (Aroid.).
chinensis, Rafin. Fl. Tellur. iv. 14 = P. scandens.
indica, Rafin. l. c. = P. scandens.
Rheedi, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 165 = P. scandens.

TAPANAWA, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 165 = praec.

TAPANHUACANGA, Vell. ex Vand. Fl. Lusit. 9. f. 5 (1788). *RUBIACEAE*? (Quid?).
brasiliensis, Steud. Nom. ed. II. ii. 663.—Bras.

TAPEINAEGLE, Herb. in Bot. Reg. (1847) t. 22. f. 4 = **Tapeinanthus**, Herb. (Amaryll.).
humilis, Herb. l. c. = Tapeinanthus humilis.

TAPEINANTHUS, Boiss. ex Benth. in DC. Prod. xii. 436 (1848) = **Chamaesphacos**, Schrenk (Labiat.).
brahuicus, Boiss. Diagn. Ser. II. iv. 29.—Persia.
persicus, Boiss. ex Benth. in DC. Prod. xii. 436.—Persia.

TAPEINANTHUS, Herb. Amaryll. 190 (1837). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 719.
Braxireon, Rafin. Fl. Tellur. iv. 23 (1836), nomen prius.
Carregnoa, Boiss. Voy. Bot. Espagne, ii. 605 (1842).
Gymnoterpe, Salisb. Gen. Pl. Fragm. 108 (1866).
Tapeinaegle, Herb. in Bot. Reg. (1847) t. 22.
humilis, *Herb. Amaryll.* 190, 414.—Hispan.; Marocc.

TAPEINIA, Juss. Gen. 59 (1789). *IRIDEAE*, Benth. & Hook. f. iii. 699.
magellanica, *J. F. Gmel. Syst.* 108; *Ker-Gawl. Gen. Irid.* 9.—Chili.

TAPEINOCHEILOS, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 101. t. 4 (1868). *SCITAMINEAE*, Benth. & Hook. f. iii. 644.
pungens, *Miq. l. c.*—Malaya; Austral.

TAPEINOCHILUS, Benth. & Hook. f. Gen. iii. 644 (1883), sphalm. = praec.

TAPEINOPHALLUS, Baill. Dict. Bot. i. (fasc. v.), Ic. sine desc. [1877] = **Amorphophallus**, Blume (Aroid.).
Rivieri, Baill. l. c. = A. Rivieri.

TAPEINOSPERMA, Hook. f. in Benth. & Hook. f. Gen. ii. 647 (1876). *MYRSINEAE*, Benth. & Hook. f. l. c.
Lenormandii, *Hook. f. l. c.*—N. Caled.
Vieillardi, *Hook. f. l. c.*—N. Caled.

TAPEINOSTEMON, Benth. in Hook. Kew Journ. vi. (1854) 194. *GENTIANEAE*, Benth. & Hook. f. ii. 808.
capitatum, *Benth. in Hook. Kew Journ.* vi. (1854) 195.—Bras.
spenneroides, *Benth. l. c.* 194.—Bras.

TAPEINOTES, DC. Prod. vii. 544 (1839) = **Sinningia**, Nees (Gesner.).
barbata, DC. l. c. = S. Carolinae.
Carolinae, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 273 = S. Carolinae.
pusilla, DC. Prod. vii. 544 (= *Ligeria pusilla*).—Bras.
villosa, Walp. Rep. ii. 716 = S. villosa.

TAPHROGITON, Montand. Guide Bot. 346 (1868) = **Scirpus**, Tourn. (Cyperac.).
radicans, Montand. l. c. = S. radicans.
sylvaticum, Montand. l. c. = S. sylvaticus.

TAPHROSPERMUM, C. A. Mey. in Ledeb. Fl. Alt. iii. 172 (1831) = **Cochlearia**, Tourn. (Crucif.).
altaicum, C. A. Mey. l. c. 173.—Sibir. altaic.
brachycarpum, Meissn. Comm. 12 = Erysimum brachycarpon.
caucasicum, Rupr. Fl. Cauc. 84.—Reg. Cauc.
platypetalum, Schrenk, Enum. Pl. Nov. ii. 60.—Soongaria.

TAPIA, Adans. Fam. ii. 407 (1763) = **Crataeva**, Linn. (Cappar.).

TAPINA, Mart. Nov. Gen. et Sp. iii. 59. t. 225. f. 1 (1829) = **Sinningia**, Nees (Gesner.).
barbata, Mart. l. c. 60. t. 225. f. 1 = S. Carolinae.
Carolinae, Hort. ex Pasq. Cat. Ort. Bot. Nap. 101 = S. Carolinae.
pusilla, Mart. Nov. Gen. et Sp. iii. 61. t. 225. f. 2 (= *Ligeria pusilla*).—Bras.
splendens, Triana, in Belg. Hortic. vii. (1857) 198 = Episcia splendens?
variegata, Linden, Cat. n. 21. (1867) 5; et in Belg. Hortic. xvii. (1867) 105.—Peruv.
villosa, Gardn. in Hook. Ic. t. 469 = S. villosa.

TAPINANTHUS, Blume, in Schult. f. Syst. vii. 1730 (1830) = **Loranthus**, Linn.
lanceolatus, Blume, l. c. = L. lanceolatus.
sessilifolius, Blume, l. c. = L. sessilifolius.

TAPINIA, Steud. Nom. ed. II. ii. 663 (1841), sphalm. = **Tapirira**, Aubl. (Anacard.).

TAPINOCARPUS, Dalz. in Hook. Kew Journ. iii. (1851) 345 = **Theriophonum**, Blume (Aroid.).
Dalzelli, Schott, Gen. Aroid. t. 15 (*Dalzeli*) = Theriophonum crenatum.

TAPIRIA, Juss. Gen. 372 (1789), sphalm. = seq.

TAPIRIRA, Aubl. Pl. Gui. i. 470. t. 188 (1775). *ANACARDIACEAE*, Benth. & Hook. f. i. 423.
JONCQUETIA, Schreb. Gen. 308 (1789).
PEGIA, Coleb. in Trans. Linn. Soc. xv. (1827) 364.
PHLEBOCHITON, Wall. in Trans. Med. & Phys. Soc. Calc. vii. (1835) 230.
SALABERTIA, Neck. Elem. ii. 238 (1790).
TAPIRIA, Juss. Gen. 372 (1789).
acida, March. Rév. Anacard. 162 = Odina acida.
bijuga, Hook. f. ex March. l. c. = guianensis.
caffra, March. l. c. = Harpephyllum caffrum.
Colebrookiana, Hook. f. ex March. l. c. = Nothopegia Colebrookiana.
cyrtocarpa, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 222.—Mexic.
discolor, March. Rév. Anacard. 162 = Odina discolor.
extensa, *Hook. f. ex March. l. c.*—Reg. Himal.
fruticosa, March. l. c. = Odina fruticosa.
guianensis, *Aubl. Pl. Gui.* i. 470. t. 188.—Guian.
hirsuta, Hook. f. Fl. Brit. Ind. ii. 28 = extensa.
humilis, March. Rév. Anacard. 162 = Odina humilis.
Marchandii, *Engl. in Mart. Fl. Bras.* xii. II. 379.—Bras.
mexicana, *March. Rév. Anacard.* 162.—Mexic.
myriantha, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 295.—N. Granat.
obovata, March. Rév. Anacard. 162 = Odina obovata.
Pao-pombo, March. in Kjoeb. Vidensk. Meddel. (1873) 58 = guianensis.
Peckoltiana, *Engl. in Mart. Fl. Bras.* xii. II. 380.—Bras.
velutina, March. Rév. Anacard. 162 = Odina velutina.
Wodier, March. l. c. = Odina Wodier.

TAPIROCARPUS, Sagot, in Ann. Sc. Nat. Sér. VI. xiii. (1882) 292. *BURSERACEAE*.
Talisia, *Sagot, l. c.*—Guian.

TAPOGOMEA, Aubl. Pl. Gui. i. 157. tt. 60, 63 (1775) = **Cephaëlis**, Sw. (Rubiac.).
aiba, Aubl. l. c. 164 = C. alba.
axillaris, Poir. Encyc. vii. 585 = C. axillaris.
Carapichea, Poir. l. c. 587 = C. involucrata.
elata, Poir. l. c. 585 = C. elata.
glabra, Aubl. Pl. Gui. i. 165 = C. glabra.
muscosa, Poir. Encyc. vii. 587, in obs. = C. muscosa.
prunifolia, Poir. in Dict. Sc. Nat. lii. 236 = Psychotria microcephala.
punicea, Poir. Encyc. vii. 585 = C. elata.
purpurea, Aubl. Pl. Gui. i. 162 = C. purpurea.
rubra, Blanco, Fl. Filip. ed. I. 145 = Uncaria acida.
tomentosa, Aubl. Pl. Gui. i. 160. t. 61 = Psychotria tomentosa, *Muell. Arg.*
violacea, Aubl. l. c. 157 = C. violacea.

TAPOMANA, Adans. Fam. ii. 343 (1763) = **Connarus**, Linn.

TAPURA, Aubl. Pl. Gui. i. 126. t. 48 (1775). *CHAILLETIACEAE*, Benth. & Hook. f. i. 341.
ROHRIA, Schreb. Gen. 30 (1789).
africana, *Oliver, Fl. Trop. Afr.* i. 344.—Afr. trop.
amazonica, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 41. t. 246. f. 2.—Bras.
capitulifera, *Baill. Adansonia*, xi. (1873) 112.—Bras.
ciliata, *Gardn. in Hook. Ic. Pl.* t. 466.—Bras.
cubensis, *Griseb. Cat. Pl. Cub.* 56.—Cuba.
cucullata, *Benth. in Hook. Kew Journ.* v. (1853) 292.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 126. t. 48.—Guian.
latifolia, *Benth. in Hook. Kew Journ.* v. (1853) 291.—Am. trop.?

TARA, Molina, Sagg. Chile, ed. II. 153 (1810) = **Caesalpinia**, Linn. (Legumin.).
tinctoria, Molina, l. c. = C. tinctoria.

TARAKTOGENOS, Hassk. Retzia, 127 (1855). *BIXINEAE*, Benth. & Hook. f. i. 129.
Blumei, *Hassk. l. c.*—Sumatra.

TARALEA, Aubl. Pl. Gui. ii. 745. t. 298 (1775) = **Dipteryx**, Schreb. (1791) (Legumin.).
oppositifolia, Aubl. l. c. = D. oppositifolia.

TARAXACUM, Linn. Syst. ed. I (1735); Hall. Enum. Stirp. Helv. ii. 739 (1742). *COMPOSITAE*, Benth. & Hook. f. ii. 522.
CARAMANICA, Tineo, Pl. Rar. Sic. fasc. i. 3 (1846).
DENS-LEONIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 197 (1745).
LASIOPUS, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 346 (1836).
LEONTODON, Adans. Fam. ii. 112 (1763).
affine, Jord. Pugill. Pl. Nov. 113 = officinale.
alpestre, DC. Prod. vii. 148 = officinale.
alpinum, C. Koch, in Linnaea, xxiii. (1850) 676 = officinale.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. (1852) 87, nomen; Enum. Pl. Transs. 368 = palustre.
apenninum, *DC. Prod.* vii. 148.—Ital.
arcuatum, Dum. Fl. Belg. 61 = officinale.
Assemani, *Blanche, ex Boiss. Fl. Orient.* iii. 791.—As. Min.; Persia.
atlanticum, Pomel, Nouv. Mat. Fl. Atl. 9 = microcephalum.
aureum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 180 = officinale.
autumnale, Cast. Cat. Pl. Marseille, 87 = gymnanthum.
autumnale, Gueldenst. ex Ledeb. Fl. Ross. ii. 778 = Leontodon autumnalis.
bellidifolium, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 187 = officinale.
bessarabicum, Fisch. Hort. Gorenk. ed. II. (1812) 34, nomen = palustre.
bicolor, *DC. Prod.* vii. 148.—Sibir. altaic.
bithynicum, Boiss. Fl. Orient. iii. 791 = Assemani.
bithynicum, *DC. Prod.* vii. 149.—Bithyn.
boreale, *DC. l. c.* 105, in syn.—Rossia.
bulbosum, Reichb. Fl. Germ. Excurs. 270 = Crepis bulbosa.
canescens, Boiss. Fl. Orient. iii. 786 = montanum.
capense, *Sch. Bip. in Flora*, xxxi. (1848) 170.—Afr. austr.
caucasicum, DC. Prod. vii. 146 = officinale.
caulescens, Moris, in Mem. Acc. Torin. xxxvii. (1834) 108. t. 24 (= *Macrorhynchus pterocarpus*).—Chili.
ceratophorum, DC. Prod. vii. 146 = officinale.
collinum, *DC. l. c.* 149.—Sibir.
commune, Dum.-Cours. Bot. Cult. ed. II. iv. 25 = officinale.
commutatum, Jord. Pugill. Pl. Nov. 116 = officinale.
corniculatum, DC. Prod. vii. 146 = officinale.
coronopifolium, Urv. in Mém. Soc. Linn. Par. iv. (1826) 609 (= *Macrorhynchus pumilus*).—Ins. Falkland.
crepidiforme, *DC. Prod.* vii. 149.—As. Min.; Persia.
crispum, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 148 = serotinum.
Dens-leonis, Desf. Fl. Atlant. ii. 228 = officinale.
denudatum, Boiss. Diagn. Ser. II. iii. 95 = montanum.
dissectum, *Ledeb. Fl. Ross.* ii. 814.—Sibir. or.
erectum, Schrank, Baier. Fl. ii. 314 = officinale.
eriopodum, DC. Prod. vii. 147 = officinale.
erythrospermum, Andrz. ex Bess. Enum. Pl. Volh. 75, 106 = officinale.
frigidum, Schur, Enum. Pl. Transs. 368 = palustre.
fulvipilus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 527.—Afr. austr.
getulum, *Pomel, Nouv. Mat. Fl. Atl.* 264.—Alger.
Gilliesii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 31.—Reg. Argent.
glabrum, *DC. Prod.* vii. 147.—Dahuria.
glaciale, Huet, ex Nym. Consp. 437 = officinale.
glaucanthum, DC. Prod. vii. 147 = officinale.
glaucescens, Reichb. Fl. Germ. Excurs. 270 = officinale.
gymnanthum, *DC. Prod.* vii. 145.—Reg. Mediterr.
halophilum, Trautv. ex Beck. in Bull. Soc. Nat. Mosc. xl. (1867) I. 111, nomen = officinale.
hirsutum, DC. Prod. vii. 149 = Troximon humile.
Hoppeanum, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 349 = officinale.
hybernum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 410 = gymnanthum.

TARAXACUM:—
inaequilobum, *Pomel, Nouv. Mat. Fl. Atl.* 8.—Alger.
integrifolium, *C. Koch, in Linnaea*, xxiii. (1850) 676.—As. Min.
laciniatum, Martr. Pl. Crit. Tarn, 32 = officinale.
laevigatum, DC. Cat. Hort. Monsp. (1813) 149 = officinale.
lanceolatum, Poir. Encyc. v. 349 = officinale.
lanuginosum, Noé, ex Nym. Consp. 436 = serotinum.
latilobum, DC. Prod. vii. 146 = officinale?
Leontodon, Gueldenst. Reisen, ii. 56; Dum. Fl. Belg. 61 = officinale.
leptocephalum, Reichb. Fl. Germ. Excurs. 270 = palustre.
leucanthum, Ledeb. Fl. Ross. ii. 815 = bicolor.
leucospermum, Jord. Cat. Jard. Dijon (1848), 31 = officinale.
levigatum, Poir. Encyc. Suppl. iv. 420 = officinale.
libanoticum, DC. Prod. vii. 149 = serotinum.
lyratum, *DC. l. c.* 148.—Sibir. altaic.
maculatum, Jord. Pugill. Pl. Nov. 117 = officinale.
magellanicum, *Comm. ex Sch. Bip. in Flora*, xxxviii. (1855) 122.—Reg. Argent.
mexicanum, DC. Prod. vii. 146 = officinale.
microcephalum, *Pomel, Nouv. Mat. Fl. Atl.* 8.—Alger.
microcephalum, Schur, Enum. Pl. Transs. 367 = officinale.
minimum, Heldr. ex Nym. Consp. 437 = officinale.
minimum, *Terrac. in Att. Ist. Incoragg. Nap.* Ser. II. vi. (1869) 352.—Sicil.; Ital.
montanum, *DC. Prod.* vii. 145.—As. Min.; Persia.
montanum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 430 = officinale.
nigricans, Reichb. Fl. Germ. Excurs. 270 = officinale.
obovatum, DC. Rapp. Voy. ii. 83 = officinale.
officinale, [*Weber, in*] *Wigg. Prim. Fl. Holsat.* 56.—Reg. temp. bor. et austr.
officinarum, Rupr. Fl. Ingr. 619 = officinale.
Pacheri, Sch. Bip. in Flora, xxxi. (1848) 170 = apenninum.
paludosum, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 215 = palustre.
palustre, *DC. Fl. Fr.* iv. 45.—Europ.; As. bor.
parvulum, DC. Prod. vii. 149 = officinale.
phymatocarpum, *J. Vahl, in Fl. Dan.* t. 2298.—Reg. arct.
porphyranthum, *Boiss. Fl. Orient.* iii. 790.—Reg. Cauc.
pratense, Hegetschw. Fl. Schw. 762 = officinale.
procumbens, Less. in Linnaea, ix. (1834) 181 = palustre.
Pseudo-officinale, Schur, Enum. Pl. Transs. 366 = officinale.
psychrophilum, Boiss. Diagn. Ser. I. xi. 48 = crepidiforme.
pumilum, Gaud. in Ann. Sc. Nat. Sér. I. v. (1825) 103 (= *Macrorhynchus pumilus*).—Ins. Falkland.
purpurascens, Schur, Enum. Pl. Transs. 366 = officinale.
pyrenaicum, *Reut. Cat. Gr. Genèv.* (1861) 4 (Quid?).—Mont. Pyren.
pyrenaicum, Timb. ex Nym. Consp. 437 = officinale.
pyrrhopappum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 19 = serotinum.
Raii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 426 = palustre.
rubrinerve, Jord. Pugill. Pl. Nov. 115 = officinale.
runcinatum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 187 = officinale.
salinum, Dum. Fl. Belg. 61 = palustre.
salsugineum, Lamotte, in Bull. Soc. Bot. Fr. xxi. (1874) 123 = palustre.
Scorzonera, Reichb. Fl. Germ. Excurs. 270 = palustre.
Scorzonera, Trautv. Fl. Taimyr. 40 = phymatocarpum.
serotinum, *Poir. Encyc. Suppl.* iv. 420; *Sadl. Fl. Pest.* ed. I. ii. 243.—Europ. or.; As. occ.
sinense, Poir. l. c.; DC. Prod. vii. 150 = officinale.
stenocephalum, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 790.—Kurdistan.
stenolobum, *Stschégl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 180.—As. bor.
Stevenii, DC. Prod. vii. 149 = officinale.
stipitatum, Sch. Bip. Cich. 76; ex Nym. Consp. 436 = Chondrilla stipitata.
sylvaticum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 525, nomen.—Rossia.
syriacum, Boiss. Diagn. Ser. II. iii. 96 = serotinum.
taraxacoides, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 231 = officinale.

TARAXACUM:—
tenuifolium, *Koch, in Flora*, xxiii. (1840) 374.—Europ.
tomentosum, Lange, in Kjoeb. Vidensk. Meddel. (1861) 101 = serotinum.
triviale, Hegetschw. Fl. Schw. 762 = officinale.
udum, Jord. Pugill. Pl. Nov. 114 = palustre.
vulgare, Schrank, Baier. Reise, 11 = officinale.
Wallichii, DC. Prod. vii. 147 = officinale.
Wattii, *Hook. f. Fl. Brit. Ind.* iii. 402.—Reg. Himal.

TARCHONANTHUS, Linn. Coroll. Gen. 13 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 288.
SALVIASTRUM, Heist. Syst. 16 (1748).
abyssinicus, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 287 = camphoratus.
angustissimus, DC. Prod. v. 431 = minor.
camphoratus, Houtt. ex DC. l. c. 430 = Brachylaena elliptica.
camphoratus, *Linn. Sp. Pl.* 842.—Afr. austr.
dentatus, Eckl. ex DC. Prod. v. 430 = Brachylaena nereifolia.
dentatus, Thunb. Prod. Pl. Cap. 145 = Brachylaena dentata.
ellipticus, Thunb. l. c. = Brachylaena elliptica.
ericoides, Linn. f. Suppl. 360 = Eriocephalus glaber.
glaber, Linn. l. c. = Brachylaena dentata.
lanceolatus, Thunb. Prod. Pl. Cap. 145 = Brachylaena nereifolia.
litakunensis, DC. Prod. v. 431 = camphoratus.
minor, *Less. Syn. Comp.* 208.—Afr. austr.
obovatus, DC. Prod. v. 431 = minor.
procerus, Salisb. Prod. 189 = camphoratus.
racemosus, Thunb. Prod. Pl. Cap. 145 = Brachylaena racemosa.
trilobus, *DC. Prod.* v. 432.—Afr. austr.

TARDAVEL, Adans. Fam. ii. 145 (1763) = **Spermacoce**, Dill. (Rubiac.).

TARENAYA, Rafin. Sylva Tellur. 111 (1838) = **Cleome**, Linn. (Cappar.).
spinosa, Rafin. l. c. = C. spinosa.

TARENNA, Gaertn. Fruct. i. 139. t. 28 (1788). *RUBIACEAE*, Benth. & Hook. f. ii. 86.
CERISCUS, Nees, in Flora, viii. (1825) 116.
CHOMELIA, Linn. Gen. ed. I. 55 (1737).
STYLOCORYNE, Wight & Arn. Prod. 400 (1834).
WAHLENBERGIA, Blume, Cat. Gew. Buitenz. 14 (1823).
WEBERA, Schreb. Gen. 794 (1791).
angolensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 89.—Afr. trop.
conferta, *Hiern, l. c.* 90.—Afr. trop.
congensis, *Hiern, l. c.* 91.—Afr. trop.
grandiflora, *Hiern, l. c.*—Afr. trop.
macrochlamys, *Baker, in Journ. Linn. Soc.* xxi. (1885) 411.—Madag.
mossambicensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 89.—Afr. trop.
nigrescens, *Hiern, l. c.* 92.—Afr. trop.
nilotica, *Hiern, l. c.* 90.—Afr. trop.
nitidula, *Hiern, l. c.*—Afr. trop.
pallidula, *Hiern, l. c.* 91.—Afr. trop.
tetramera, *Hiern, l. c.*—Afr. trop.
zeylanica, *Gaertn. Fruct.* i. 139. t. 28. f. 3.—Zeylan.

TARIRI, Aubl. Pl. Gui. Suppl. 37 (1775) = **Picramnia**, Sw. (Simarub.).
guianensis, Aubl. l. c. = P. Tariri.
polyantha, Baill. Hist. Pl. iv. 412 = P. polyantha.

TARPHETA, Rafin. Fl. Tellur. ii. 103 (1836) = **Stachytarpheta**, Vahl (Verben.).

TARRIETIA, Blume, Bijdr. 227 (1825). *STERCULIACEAE*, Benth. & Hook. f. i. 218.
ARGYRODENDRON, F. Muell. Fragm. i. 2 (1858).
Argyrodendron, *Benth. Fl. Austral.* i. 230.—Austral.
javanica, *Blume, Bijdr.* 227.—Java.
Riedeliana, *Oliver, in Journ. Linn. Soc.* xv. (1876) 98.—Ins. Celebes.
simplicifolia, *Mast. in Hook. f. Fl. Brit. Ind.* i. 362.—Malacca.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 401.—Sumatra.
trifoliolata, *F. Muell. Fragm.* ix. 43.—Austral.

TARSINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 (Quid?).

TASMANNIA, R. Br. ex DC. Syst. i. 445 (1818) = **Drimys**, Forst. (Magnol.).
aromatica, R. Br. l. c. = D. aromatica.
dipetala, R. Br. ex DC. Prod. i. 78 = D. dipetala.
insipida, R. Br. ex DC. Syst. i. 445 = D. dipetala.
monticola, A. Rich. Sert. Astrolab. 50 = D. dipetala.
piperita, Miers, in Ann. & Mag. Nat. Hist. Ser. III. ii. (1858) 110 = D. piperita.

TASOBA, Rafin. Fl. Tellur. iii. 94 (1836) = **Polygonum**, Tourn.
arifolia, Rafin. l. c. = P. arifolium.
sagittata, Rafin. l. c. 95 = P. sagittatum.

TASSADIA, Decne. in DC. Prod. viii. 579 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 756.
Burchelii, *Fourn. in Mart. Fl. Bras.* vi. IV. 230.—Bras.
Colubrina, *Decne. in DC. Prod.* viii. 579.—Bras.
comosa, *Fourn. in Mart. Fl. Bras.* vi. IV. 229.—Bras.
Decaisneana, Miq. in Linnaea, xviii. (1844) 25 = Metastelma Decaisneana.
floribunda, *Decne. in DC. Prod.* viii. 580.—Bras.
guianensis, *Decne. l. c.* 579.—Guian.
lanceolata, *Decne. l. c.*—Bras.
leptobotrys, *Decne. l. c.*—Guian.
Martiana, *Decne. l. c.* 580.—Bras.
neovidensis, *Fourn. in Mart. Fl. Bras.* vi. IV. 229.—Bras.
obovata, *Decne. in DC. Prod.* viii. 579.—Guian.
Poeppigiana, *Decne. l. c.* 580.—Bras.
propinqua, *Decne. l. c.* 579.—Guian.
Selloana, *Fourn. in Mart. Fl. Bras.* vi. IV. 231.—Bras.
Sprucei, *Fourn. l. c.* 229.—Bras.
turriformis, *Fourn. l. c.* 231.—Bras.

TASSIA, Rich. ex DC. Prod. ii. 487 (1825) = **Tachigalia**, Juss. (Legumin.).

TATEA, F. Muell. in Trans. Roy. Soc. S. Austral. vi. (1883) 33. *VERBENACEAE*.
acaulis, F. Muell. Census, Suppl. i. 3, sphalm. = seq.
subacaulis, *F. Muell. in Trans. Roy. Soc. S. Austral.* vi. (1883) 34.—Austral.

TATEA, Seem. Fl. Vit. 125 (1866) = **Bikkia**, Reinw. (Rubiac.).
portlandioides, Seem. l. c.—N. Caled.

TATTIA, Scop. Introd. 104 (1777) = **Homalium**, Jacq. (Samyd.).
ignota, J. F. Gmel. Syst. 836 = H. Napimoga.

TAUMASTOS, Rafin. Fl. Tellur. iv. 9 (1836) = **Libertia**, Spreng. (Irid.).
compressus, Rafin. l. c. 10 = L. formosa.

TAUROSTALIX, Reichb. f. in Bot. Zeit. x. (1852) 933 = **Bulbophyllum**, Thou. (Orchid.).
herminiostachys, Reichb. f. l. c. = B. herminiostachys.

TAURRETTIA, Raeusch. Nom. ed. III. 176 (1797), sphalm. = **Tourrettia**, Juss. (Bignon.).

TAUSCHERIA, Fisch. Hort. Gorenk. ed. II. (1812) 49; et ex DC. Syst. ii. 563 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 94.
desertorum, Ledeb. Ic. Pl. Fl. Ross. ii. 14. t. 139; Royle, Illustr. Bot. Himal. 72 = lasiocarpa.
gymnocarpa, Fisch. Hort. Gorenk. ed. II. (1812) 49, nomen; et ex DC. Syst. ii. 564 = lasiocarpa.
lasiocarpa, *Fisch. l. c.*, nomen; *et ex DC. l. c.* 563.—Reg. Himal. et Casp.

TAUSCHIA, Preissler, in Flora, xi. (1828) 43 = **Hornemannia**, Vahl (Vaccin.).
hederaefolia, Preissler, l. c. 44 = H. martinicensis.

TAUSCHIA, Schlecht. in Linnaea, ix. (1834) 607. *UMBELLIFERAE*, Benth. & Hook. f. i. 882.
Coulteri, *A. Gray & Harv. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 211.—Mexic.
nudicaulis, *Schlecht. in Linnaea*, ix. (1834) 608.—Mexic.
texana, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 211 = Eulophus texanus.

TAVALLA, Pers. Syn. ii. 633 (1807) = TAFALLA, Ruiz & Pav. = **Hedyosmum**, Sw. (Chloranth.).

TAVERNARIA, Reichb. Consp. 152 (1828) = seq.

TAVERNIERA, DC. Mém. Légum. 339. t. 52 (1825?). *LEGUMINOSAE*, Benth. & Hook. f. i. 510.
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 206.—Afr. trop.
aegyptiaca, *Boiss. Diagn.* Ser. I. ix. 113.—Aegypt.
cuneifolia, Arn. in Nov. Act. Nat. Cur. xviii. I. (1836) 332 = nummularia.
cyclophylla, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 141 = Schimperi.
ephedroidea, Jaub. & Spach, Illustr. i. 122 = nummularia.
floribunda, Schweinf. Beitr. Fl. Aethiop. 17 = Schimperi.
glabra, *Boiss. Diagn.* Ser. I. ii. 90.—Persia.
glauca, *Edgew. in Journ. As. Soc. Beng.* xvi. (1874) II. 1214.—Arab.
gonoclada, Jaub. & Spach, Illustr. i. 120 = nummularia.
incana, Boiss. Diagn. Ser. I. ii. 90 = nummularia.
lappacea, *DC. Prod.* ii. 339.—Arab.
nummularia, *DC. Mém. Légum.* 340.—Persia; Mesopot.; Ind. bor. occ.
persica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 509.—Persia.
Schimperi, *Jaub. & Spach, Illustr.* v. 83. t. 474.—Abyss.
sericophylla, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 510.—Ins. Socotra.
spartea, DC. Prod. ii. 339 = nummularia.
Stocksii, *Boiss. Fl. Orient.* ii. 509.—Beluchist.

TAVOMYTA, Vitm. Summa Pl. vi. 70 (1792), sphalm. = **Tovomita**, Aubl. (Guttif.).

TAXANTHEMA, Neck. Elem. i. 115 (1790) = **Statice**, Linn. (Plumbag.).
aegyptiaca, Sweet, Hort. Brit. ed. I. 333 = S. Thouini.
alata, Sweet, l. c. = S. Thouini.
attenuata, R. Br. in Salt, Abyss. App. 64 = S. attenuata.
auriculaefolia, Sweet, Hort. Brit. ed. I. 332 = S. auriculaefolia.
australis, R. Br. Prod. 426 = S. australis.
bellidifolia, Sweet, Hort. Brit. ed. I. 332 = S. ocymifolia.
caroliniana, Sweet, l. c. = S. Limonium.
caspia, Sweet, l. c. = S. caspia.
conspicua, Sweet, l. c. 333 = S. speciosa.
cordata, Sweet, l. c. 332 = S. cordata.
dichotoma, Sweet, l. c. 333 = S. dichotoma.
echioides, Sweet, l. c. = S. rorida.
elata, Sweet, l. c. 332 = S. elata.
emarginata, Sweet, l. c. = S. emarginata.
ferulacea, Sweet, l. c. 333 = S. ferulacea.
flexuosa, Sweet, l. c. = S. flexuosa.
globulariaefolia, Sweet, l. c. 332 = S. globulariaefolia.
Gmelini, Sweet, l. c. = S. Gmelini.
graminifolia, Sweet, l. c. = S. graminifolia.
incana, Sweet, l. c. = S. incana.
latifolia, Sweet, l. c. = S. latifolia.
Limonium, Sweet, l. c. = S. Limonium.
macrophylla, Sweet, l. c. 333 = S. macrophylla.
minuta, Sweet, l. c. = S. minuta.
monopetala, Sweet, l. c. = Limoniastrum articulata.
mucronata, Sweet, l. c. = S. mucronata.
oleaefolia, Sweet, l. c. 332 = S. virgata.
pectinata, Sweet, l. c. 333 = S. pectinata.
purpurata, Sweet, l. c. 332 = S. purpurata.
reticulata, Sweet, l. c. 333 = S. reticulata.
scabra, Sweet, l. c. 332 = S. scabra.
Scoparia, Sweet, l. c. = S. Gmelini.
sinuata, Sweet, l. c. 333 = S. sinuata.
spathulata, Sweet, l. c. 332 = S. spathulata.
speciosa, Sweet, l. c. 333 = S. speciosa.
suffruticosa, Sweet, l. c. = S. suffruticosa.
tatarica, Sweet, l. c. = S. tatarica.
tetragona, Sweet, l. c. 332 = S. tetragona.
Thouini, Sweet, l. c. 333 = S. Thouini.
viminea, Sweet, l. c. = S. virgata.
virgata, Sweet, l. c. = S. virgata.
Willdenowiana, Sweet, l. c. 332 = S. Willdenovii.

TAXODIUM, Rich. in Ann. Mus. Par. xvi. (1810) 298. *CONIFERAE*, Benth. & Hook. f. iii. 429.
CUPRESPINNATA, [Nelson], Senilis, Pinaceae, 61 (1866).
GLYPTOSTROBUS, Endl. Syn. Conif. 69 (1847).
SCHUBERTIA, Mirb. in Nouv. Bull. Soc. Philom. iii. (1812) 123.
ascendens, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1833) 182 = distichum.
capense, Hort. ex Endl. Syn. Conif. 32 = Callitris arborea.
denudatum, Hort. ex Carr. Conif. ed. II. 182 = distichum.
distichum, H. B. & K. Nov. Gen. et Sp. ii. 4 = mucronatum.
distichum, *Rich. in Ann. Mus. Par.* xvi. (1810) 298. —Am. bor.
fastigiatum, Hort. ex Carr. Conif. ed. I. 145 = distichum.
gigantaum, Hort. ex Carr. l. c. 164 = Sequoia sempervirens.
giganteum, Kellogg & Behr, in Proc. Calif. Acad. ed. II. i. (1873) 51 = Sequoia gigantea.
heterophyllum, *Brongn. in Ann. Sc. Nat.* Sér. I. xxx. (1833) 184.—China.
Horsfieldii, Knight, ex Carr. Conif. ed. I. 478 = Podocarpus cupressina.
Hugeli, Laws. ex Gord. Pinet. 307 = mucronatum.
imbricatum, Hort. ex Carr. Conif. ed. II. 181 = distichum.
japonicum, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1833) 183 = Cryptomeria japonica.
japonicum, Dehnhardt, ex Gord. Pinet. 89 = heterophyllum.
juniperoides, Rich. ex Steud. Nom. ed. II. ii. 664 = Callitris arborea.
Knightii, Hort. ex C. Koch, Dendrol. ii. II. 198 = distichum.
mexicanum, Carr. Conif. ed. I. 147 = mucronatum.
microphyllum, Brongn. in Ann. Sc. Nat. Ser. I. xxx. (1833) 182 = distichum.
Montezumae, Decne. in Bull. Soc. Bot. Fr. i. (1854) 71 = mucronatum.
mucronatum, *Tenore, in Ann. Sc. Nat.* Sér. III. xix. (1853) 355.—Mexic.
nuciferum, Brongn. ex Gord. Pinet. ed. II. 126 = heterophyllum.
nutkaënse, Lamb. ex Endl. Syn. Conif. 198 = Sequoia sempervirens?
pinnatum, Hort. ex Carr. Conif. ed. I. 147 = mucronatum.
sempervirens, Lamb. Pin. 114; ed. II. 107 = Sequoia sempervirens.
sinense, Forbes, Pinet. Woburn. 179 = heterophyllum.
sinense, Nois. ex Gord. Pinet. 309 = distichum.
Washingtonianum, Winslow, in Calif. Farm. (Sept. 1854); ex Hook. Kew Journ. vii. (1855) 29 = Sequoia gigantea.

TAXOTROPHIS, Blume, Mus. Bot. Lugd. Bat. ii. 77. t. 26 (1852) [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 358.
DIPLOCOS, Bur. in DC. Prod. xvii. 215 (1873).
TOXOTROPHIS, Planch. in DC. l. c. 208 (1873).
javanica, *Blume, Mus. Bot. Lugd. Bat.* ii. 77.—Java.
microphylla, F. Muell. Fragm. vi. 193 (= *Epicarpurus microphyllus*).—N. Zel.
rectinervis, F. Muell. l. c. 192 = Aphananthe philippinensis.
Roxburghii, *Blume, Mus. Bot. Lugd. Bat.* ii. 78.—Ind. or.; Malaya.
zeylanica, *Thw. Enum. Pl. Zeyl.* 264.—Zeylan.

TAXUS, [Tourn.] Linn. Syst. ed. I (1735). *CONIFERAE*, Benth. & Hook. f. iii. 431.
VERATAXUS, [Nelson], Senilis, Pinaceae, 168 (1866).
adpressa, *Hort. ex Lindl. & Gord. in Journ. Hort. Soc.* v. (1850) 227.—Japon.
aurea, Hort. ex C. Koch, Dendrol. ii. II. 93 = baccata.
baccata, Hook. Fl. Bor. Am. ii. 167, partim = brevifolia.
baccata, *Linn. Sp. Pl.* 1040.—Reg. bor. temp.
baccata, Thunb. Fl. Jap. 275 = Cephalotaxus drupacea.
Bourcieri, Carr. in Rev. Hortic. Sér. IV. iii. (1854) 228 = brevifolia.
brevifolia, Hort. ex Gord. Pinet. 310 = adpressa.

TAXUS:—
brevifolia, *Nutt. N. Am. Sylva*, iii. 86.—Am. bor. occ.
canadensis, Bigel. in Pacif. Rail. Rep. iv. 25 = brevifolia.
canadensis, *Willd. Sp. Pl.* iv. 856.—Am. bor.
capensis, Lam. Encyc. iii. 229 = Podocarpus elongata.
chinensis, Roxb. Hort. Beng. 73 = Podocarpus chinensis.
columnaris, Hort. ex C. Koch, Dendrol. ii. II. 92 = baccata.
contorta, Griff. Notul. iv. 28 = baccata.
coriacea, Hort. ex Knight, Syn. Conif. 51 = Cephalotaxus drupacea.
cuspidata, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 108.—Japon.
disticha, Wender. ex Henk. & Hochst. Syn. Nadelh. 354 = baccata.
Dovastonii, Hort. ex Carr. Conif. ed. I. 518 = baccata.
elegantissima, Hort. ex Carr. l. c. = baccata.
elongata, Ait. Hort. Kew. ed. I. iii. 415; Thunb. Prod. Pl. Cap. 117 = Podocarpus elongata.
empetrifolia, Hort. ex Gord. Pinet. ed. II. 390 = baccata.
erecta, Hort. ex Carr. Conif. ed. I. 518 = baccata.
ericoides, Hort. ex Carr. l. c. 519 = baccata.
expansa, Hort. ex C. Koch, Dendrol. ii. II. 92 = baccata.
falcata, Thunb. Prod. Pl. Cap. 117 = Podocarpus falcata.
fastigiata, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 227 = baccata.
floridana, *Nutt. N. Am. Sylv.* iii. 92, in obs.; ed. II. ii. 155.—Florida.
Fortunei, Hort. ex Gord. Pinet. ed. II. 338 = Podocarpus koraiana.
Foxii, Hort. Angl. ex Carr. Conif. ed. I. 519 = baccata.
globosa, *Schlecht. in Linnaea*, xii. (1838) 496.—Mexic.
Harringtonia, Knight, ex Forbes, Pinet. Woburn. 217 = Cephalotaxus pedunculata.
hibernica, Hook. ex Loud. Arbor. iv. 2066 = baccata.
horizontalis, Hort. ex Carr. Conif. ed. I. 518 = baccata.
imperialis, Hort. ex Gord. Pinet. 312 = baccata.
Inukaja, Knight, Syn. Conif. 51 = Cephalotaxus pedunculata.
Jacksonii, Hort. ex C. Koch, Dendrol. ii. II. 93 = baccata.
japonica, Hook. ex Gord. Pinet. Suppl. 21 = Cephalotaxus drupacea.
japonica, Lodd. ex Gord. Pinet. 275 = Cephalotaxus pedunculata.
Lambertiana, Wall. Cat. n. 6056 = Abies Pindrow.
lancifolia, Wikstr. in Vet. Acad. Handl. Stockh. (1827) 76 = Podocarpus coriacea.
latifolia, Thunb. Prod. Pl. Cap. 117 = Podocarpus Thunbergii.
Lindleyana, A. Murr. in Edinb. N. Phil. Journ. i. (Jan.-Apr. 1855) 294 = brevifolia.
longifolia, Hort. ex Gord. Pinet. 278 = Podocarpus macrophylla.
lugubris, Salisb. Prod. 396 = baccata.
macrophylla, Banks, ex Endl. Syn. Conif. 218 = Podocarpus Thunbergii.
macrophylla, Thunb. Fl. Jap. 276 = Podocarpus macrophylla.
Makoya, Forbes, Pinet. Woburn. 218 = Podocarpus macrophylla.
marginata, Hort. ex Carr. Conif. ed. I. 518 = baccata.
Michelii, Hort. ex Carr. l. c. ed. II. 736 = baccata.
microphylla, Hort. ex Gord. Pinet. 313 = baccata.
Mitchelli, Hort. ex Gord. l. c. = baccata.
monstrosa, Hort. ex Gord. l. c. = baccata.
monstruosa, Hort. ex Parl. in DC. Prod. xvi. II. 501 = baccata.
montana, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 857 = Podocarpus taxifolia.
montana, Nutt. N. Am. Sylva, iii. 92, in obs.; ed. II. ii. 155 = floridana.
nana, Hort. ex Parl. in DC. Prod. xvi. II. 501 = baccata.
nepalensis, Jacquem. ex Parl. l. c. 500 = baccata.
nucifera, Hort. ex Parl. l. c. 439 = Glyptostrobus heterophyllus
nucifera, Linn. Sp. Pl. 1040 = Torreya nucifera.

TAXUS:—
nucifera, Telfair, ex Steud. Nom. ed. II. ii. 664 = Podocarpus chinensis.
nucifera, Wall. Tent. Fl. Nep. 57. t. 44 = baccata.
occidentalis, Nutt. N. Am. Sylv. iii. t. 108 = brevifolia.
orientalis, Bertol. Misc. Bot. xxiv. 17 = baccata.
parvifolia, Wender. Conif. 42; ex Henck. & Hochst. Syn. Nadelh. 361 = adpressa.
patagonica, Hort. ex Gord. Pinet. 300 = Saxegothaea conspicua.
pectinata, Gilib. Exercit. ii. 417 = baccata.
pendula, Hort. ex Carr. Conif. ed. I. 518 = baccata.
procumbens, Lodd. Cat. (1836) 67; ex Loud. Arbor. iv. 2067 = canadensis.
pyramidalis, Hort. ex Carr. Conif. ed. I. 519 = baccata.
recurvata, Laws. ex Gord. Pinet. 313 = baccata.
serratifolia, Nois. ex Gord. Pinet. Suppl. 43 = Phyllocladus rhomboidalis.
sinensis, Knight, ex Gord. Pinet. Suppl. 21 = Cephalotaxus pedunculata.
sparsifolia, Loud. Encyc. Trees, 940 = baccata.
spicata, Domb. ex Mirb. in Mém. Mus. Par. xiii. (1825) 76 = Podocarpus andina.
spinulosa, Sm. in Rees, Cycl. xxxv. n. 7 = Podocarpus spinulosa.
stricta, Hort. ex Gord. Pinet. 312 = baccata.
tardiva, Laws. ex Gord. l. c. 310 = adpressa.
tenuifolia, Parl. in DC. Prod. xvi. II. 509 = Podocarpus coriacea.
tomentosa, Thunb. Prod. Pl. Cap. 117 = Grubbia rosmarinifolia.
umbraculifera, Hort. ex Gord. Pinet. 312 = baccata.
variegata, Hort. ex Carr. Conif. ed. I. 518 = baccata.
verticillata, Thunb. Fl. Jap. 276 = Sciadopitys verticillata.
virgata, Wall. ex Gord. Pinet. 317 = baccata.
Wallichiana, Zucc. in Abh. Akad. Muench. iii. (1837–43) 803. t. 5 = baccata.
Washingtoni, Hort. ex Gord. Pinet. ed. II. 394 = canadensis.

TAYOTUM, Blanco, Fl. Filip. ed. I. (1837) 104; ed. II. 76 (1845). Genus dubium *APOCYNACEARUM*. —Cf. Benth. & Hook. f. ii. 690.
ngirescens, *Blanco, l. c.* ed. I. 105.—Ins. Philipp.

TCHIHATCHEWIA, Boiss. in Tchihat Asie Min. Bot. i. 292 (1860). *CRUCIFERAE*, Benth. & Hook. f. i. 93.
isatidea, *Boiss. l. c.*—Armenia.

TECLEA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 90 = **Toddalia**, Juss. (Rutac.).
nobilis, Delile, l. c. = Toddalia nobilis.

TECMARSIS, DC. Prod. v. 93 (1836) = **Pluchea**, Cass. (Compos.).
Bojeri, DC. l. c.—Madag.

TECOMA, Juss. Gen. 139 (1789). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1044.
ASTIANTHUS, D. Don, in Edinb. Phil. Journ. ix. (1823) 262.
CAMPSIDIUM, Seem. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 32.
CAMPSIS, Lour. Fl. Cochinch. 377 (1790).
DENDROPHILA, Zipp. ex Blume, Rumphia, iv. 34. t. 190 (1848).
DUCOUDRAEA, Bur. Monog. Bign. 49 (1864).
KOKOSCHKINIA, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 33.
LEUCOXYLON, Rafin. Sylva Tellur. 77 (1838).
PANDOREA, Spach, Hist. Vég. Phan. ix. (1840) 136.
PLATOLARIA, Rafin. Sylva Tellur. 78 (1838).
STENOLOBIUM, D. Don, in Edinb. Phil. Journ. ix. (1823) 264.
TECOMARIA, Spach, Hist. Vég. Phan. ix. 137 (1840).
TECOMELLA, Seem. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 30.
actinophylla, *Griseb. Cat. Pl. Cub.* 194.—Cuba.
aesculifolia, DC. Prod. ix. 221 = Tabebuia aesculifolia.
Aesculus, *DC. l. c.* 222.—Bras.
africana, G. Don, Gen. Syst. iv. 224 = Bignonia africana.

TECOMA:—
alata, DC. Rev. Bign. (1838) 24; et Prod. ix. 225 = T. Guarume.
alba, *Cham. in Linnaea*, vii. (1832) 655.—Bras.
amboinensis, *Blume, Rumphia*, iv. 35.—Ins. Amboin.
araliacea, *DC. Prod.* ix. 221.—Bras.
atrovirens, *DC. l. c.* 220.—Bras.
aurea, *DC. l. c.* 222.—Bras.
australis, *R. Br. Prod.* 471.—Austral.
austro-caledonica, *Bur. in Bull. Soc. Bot. Fr.* ix. (1862) 163.—N. Caled.
azaleaeflora, H. B. & K. Nov. Gen. et Sp. iii. 142 = rosaefolia.
azaleaefolia, G. Don, Gen. Syst. iv. 224, sphalm. = praec.
barbata, *DC. Prod.* ix. 221.—Venezuela.
Barterii, *DC. l. c.* 220.—Ind. occ.
bibracteolata, *Griseb. Cat. Pl. Cub.* 193.—Cuba.
capensis, *Lindl. Bot. Reg.* t. 1117.—Afr. austr.
Caraiba, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 14.—Bras.
Carianthera, Mart. ex DC. Prod. ix. 225 (= *Lundia intermedia*).—Bras.
ceramensis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 412.—Ins. Ceram.
chinensis, C. Koch, Dendrol. ii. 1. 307 = grandiflora.
chrysantha, *DC. Prod.* ix. 221.—Venezuela.
chrysotricha, *Mart. ex DC. l. c.* 216.—Bras.
conspicua, *DC. l. c.* 221.—Guian.
cuspidata, Blume, Rumphia, iv. 35 = Nyctocalos brunfelsiaeflorus.
cyrtantha, *Mart. ex DC. Prod.* ix. 218.—Bras.
Dendrophila, *Blume, Rumphia*, iv. 35.—N. Guin.
digitata, *H. B. & K. Nov. Gen. et Sp.* iii. 142.—Mexic.
diversifolia, G. Don, Gen. Syst. iv. 225 = australis.
dubia, *Wright, in Sauv. Fl. Cub.* 95.—Cuba.
elliptica, *DC. Prod.* ix. 220.—Bras.
eximia, *Miq. in Linnaea*, xxii. (1849) 803.—Bras.
flavescens, Mart. ex DC. Prod. ix. 216 = Tabebuia flavescens.
floribunda, A. Cunn. ex DC. Rev. Bign. (1838) 24; et Prod. ix. 225 = australis.
fulgens, *Mart. ex DC. Prod.* ix. 222.—Bras.
fulva, *G. Don, Gen. Syst.* iv. 224.—Peruv.; Bolivia.
fuscata, DC. Prod. ix. 218 = Tabebuia fuscata.
Garrocha, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 402.—Reg. Argent.
Gaudichaudi, *DC. Prod.* ix. 223.—N. Granat.
glauca, DC. l. c. = undulata.
grandiflora, *Loisel. Herb. Amat.* v. t. 286.—China; Japon.
grandiflora, Sweet, Hort. Brit. ed. I. 284 = Aeschynanthus grandiflora.
Guarume, *DC. Prod.* ix. 224.—Peruv.
Guarume, Hook. Bot. Mag. sub t. 4896, in adnot. = valdiviana.
Guayacan, Seem. Bot. Voy. Herald, 180 = Tabebuia Guayacan.
haemantha, *Griseb. Cat. Pl. Cub.* 194.—Cuba.
heptaphylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 13, 14.—Bras.
heteropoda, *DC. Prod.* ix. 219.—Peruv.
heterotricha, DC. l. c. = Tabebuia heterotricha.
Hillii, *F. Muell. Fragm.* x. 101.—Austral.
hirsuta, *DC. Prod.* ix. 222.—Ind. or.
hirta, *Mart. ex DC. l. c.*—Bras.
hypodictyon, *DC. l. c.* 217.—Bras.
hypoleuca, *Wright, in Sauv. Fl. Cub.* 94.—Cuba.
impetiginosa, *Mart. ex DC. Prod.* ix. 218.—Bras.
incisa, Sweet, Hort. Brit. ed. I. 284 = stans.
insignis, *Miq. Stirp. Surin. Sel.* 122.—Guian.
Ipe, Liais, Brés. 605 = ochracea, obtusata.
jasminoides, *Lindl. Bot. Reg.* t. 2002; *G. Don, Gen. Syst.* iv. 225.—Austral.
lasianthera, Mart. ex DC. Prod. ix. 180 = Lundia cordata.
lateriflora, Mart. in Flora, xxiv. (1841) II. Beibl. 51 = Cuspidaria lateriflora.
Latrobei, F. Muell. ex Teijsm. & Binn. Cat. Hort. Bog. 154, nomen.—Austral. ?
lepidophylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 524.—Cuba.
lepidota, *DC. Prod.* ix. 220.—Cuba.
leptophylla, *Blume, Rumphia*, iv. 35.—N. Guin.
leucophlaeos, *Mart. ex DC. Prod.* ix. 217.—Bras.
leucoxylon, *Mart. ex DC. l. c.* 219.—Ind. occ.; Guian.
longiflora, *Griseb. Cat. Pl. Cub.* 194.—Cuba.
longiflora, Mart. ex DC. Prod. ix. 218 = speciosa.
loxensis, Benth. Pl. Hartw. 354 = Delostoma nervosum.

TECOMA:—
Maximiliani, *Mart. ex DC. Prod.* ix. 222.—Bras.
meonantha, Sweet, Hort. Brit. ed. I. 284 = australis.
mexicana, Mart. ex DC. Prod. ix. 218 = Tabebuia mexicana.
Meyeriana, *DC. l. c.* 221.—Guian.
mirabilis, Hort. ex Van Houtte, in Fl. des Serres, xx. (1874) 149 = valdiviana.
mollis, *H. B. & K. Nov. Gen. et Sp.* iii. 144.—Mexic.
myriantha, *DC. Prod.* ix. 220.—Bras.
myrtifolia, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 524.—Cuba.
nigricans, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1159, nomen.—Guian.
nodosa, Griseb. in Goett. Abh. xix. (1874) 223 = Tabebuia nodosa.
Nyassae, *Oliver, in Hook. Ic. Pl.* t. 1351.—Afr. trop.
obtusata, *DC. Prod.* ix. 217.—Bras.
ochracea, *Cham. in Linnaea*, vii. (1832) 653.—Bras.
ochracea, Mart. ex DC. Prod. ix. 217 = obtusata.
ochroxantha, Kunth & Bouché, Ind. Sem. Hort. Berol. (1847) 12 = australis.
Oxleyi, A. Cunn. ex DC. Prod. ix. 225 = australis.
Pandorea, Juss. ex Steud. Nom. ed. II. ii. 664 = australis.
parviflora, Mart. ex DC. Prod. ix. 215 (= *Craterotecoma parviflora*).—Bras.
Patrisiana, *DC. l. c.* 221.—Guian.
pentaphylla, Juss. Gen. 139 = Tabebuia pentaphylla.
Petersii, Klotzsch, in Peters, Reise Mossamb. Bot. 193, in obs. = capensis.
platyantha, *Griseb. Fl. Brit. W. Ind.* 447.—Jamaic.
radicans, *Juss. Gen.* 139.—Am. bor.
rosaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 143.—Peruv.
rosea, Bertol. Fl. Guatim. 25 = Tabebuia rosea.
Salzmanni, *DC. Prod.* ix. 219.—Bras.
sambucifolia, H. B. & K. Nov. Gen. et Sp. iii. 143 = stans.
sanguinea, *Wright, in Sauv. Fl. Cub.* 96.—Cuba.
serratifolia, *G. Don, Gen. Syst.* iv. 224.—Ind. occ.
setulosa, *Griseb. Cat. Pl. Cub.* 193.—Cuba.
sinensis, Spach, Hist. Vég. Phan. ix. 135 = grandiflora.
sorbifolia, H. B. & K. Nov. Gen. et Sp. iii. 144 = mollis.
speciosa, *DC. ex Mart. in Flora*, xxiv. (1841) II. *Beibl.* 13.—Bras.
spectabilis, *Planch. & Linden, ex Planch. in Fl. des Serres*, ix. (1853-54) 233. t. 948.—Ind. occ.; Venezuela.
spiralis, *Wright, ex Griseb. Cat. Pl. Cub.* 194.—Cuba.
splendida, Spreng. Syst. ii. 834 = Esterhazya splendida.
squamellulosa, *DC. Prod.* ix. 220.—Bras.
stans, Griseb. in Goett. Abh. xxiv. (1879) 257 = T. Garrocha.
stans, *Juss. Gen.* 139.—Am. bor. et austr.
suaveolens, G. Don, Gen. Syst. iv. 224 = Stereospermum suaveolens.
subvernicosa, *DC. Prod.* ix. 219.—Bras.
Tagliabuana, *Vis. in Att. Ist. Venet. Sc.* Ser. III. iv. (1858-59) 135.—Hab. ?
Thunbergi, *Carr. in Rev. Hortic.* (1876) 440.—Hab. ?
toxophora, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 15.—Bras.
trachycarpa, *Griseb. Cat. Pl. Cub.* 192.—Cuba.
trichocalycina, *DC. Prod.* ix. 221.—Bolivia.
triphylla, Mart. ex DC. l. c. 215 = Tabebuia triphylla.
uliginosa, Mart. ex DC. l. c. 212 = Tabebuia uliginosa.
umbellata, *Sond. in Linnaea*, xxii. (1849) 562.—Bras.
undulata, *G. Don, Gen. Syst.* iv. 223.—Oriens; Ind. bor. occ.
uspallatensis, Mart. ex DC. Prod. ix. 235 = Argylia uspallatensis.
valdiviana, *Phil. in Linnaea*, xxix. (1857-58) 14.—Chili.
velutina, Lindl. in Gard. Chron. (1855) 820 = stans *vel* mollis ?
venusta, Lem. Hort. Univ. (Mai 1843) 1 = Bignonia venusta.
viminalis, *Hemsl. Biol. Centr. Am. Bot.* ii. 497.—Mexic.
xylocarpa, G. Don, Gen. Syst. iv. 225 = Stereospermum xylocarpum.
Zeyheri, Regel, Cat. Pl. Hort. Aksakov. 138 (Quid?).—Hab. ?

TECOMARIA, Spach, Hist. Vég. Phan. ix. 137 (1840) = **Tecoma**, Juss. (Bignon.).
capensis, Spach, l. c. = Tecoma capensis.
fulva, Seem. in Journ. Bot. i. (1863) 19 = Tecoma fulva.

TECOMARIA:—
Krebsii, Klotzsch, in Peters, Reise Mossamb. Bot. 193, in obs. = Tecoma capensis.
Petersii, Klotzsch, l. c. 192 = Tecoma capensis.
rosaefolia, Seem. in Journ. Bot. i. (1863) 20 = Tecoma rosaefolia.

TECOMELLA, Seem. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 30 = **Tecoma**, Juss. (Bignon.).
Bulweri, F. Muell. Fragm. iv. 147 = Diplanthera tetraphylla.
undulata, Seem. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 30 = Tecoma undulata.

TECOPHILAEA, Bert. ex Colla, in Mem. Acc. Torin. xxxix. (1836) 19. t. 55. *HAEMODORACEAE*, Benth. & Hook. f. iii. 680.
DISTREPTA, Miers, Trav. Chil. ii. 529 (1826), nomen prius.
PHYGANTHUS, Poepp. & Endl. Nov. Gen. et Sp. ii. 71. t. 200 (1838).
POEPPIGIA, Kunze, in Reichb. Consp. 212ª (1828).
albida, Miers, in Trans. Linn. Soc. xxiv. (1864) 505 = violaeflora.
cyaneo-crocea, Baker, in Journ. Linn. Soc. xvii. (1879) 496, sphalm. = seq.
cyanocrocus, *Leyb. in Bonplandia*, x. (1862) 370.—Chili.
violaeflora, *Bert. ex Colla, in Mem. Acc. Torin.* xxxix. (1836) 20.—Chili.

TECOPHILEA, Herb. Amaryll. 69 (1837) = praec.

TECTICORNIA, Hook. f. in Benth. & Hook. f. Gen. iii. 65 (1880). *CHENOPODIACEAE*, Benth. & Hook. f. l. c.
cinerea, *Hook. f. l. c.*—Austral.

TECTONA, Linn. f. Suppl. 20, 151 (1781). *VERBENACEAE*, Benth. & Hook. f. ii. 1152.
NAUTEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3.
THEKA, Adans. Fam. ii. 445 (1763).
grandis, *Linn. f. Suppl.* 151.—Ind. or.; Burma.
Hamiltoniana, *Wall. Pl. As. Rar.* iii. 68. t. 294.—Ind. or.
philippinensis, *Benth. & Hook. f. Gen.* ii. 1152.—Ins. Philipp.
ternifolia, Buch.-Ham. ex Wall. Pl. As. Rar. iii. 68 = Hamiltoniana.
Theca, Lour. Fl. Cochinch. 137 = grandis.

TECTONIA, Spreng. Anleit. ii. II. 893 (1818) = praec.

TEEDIA, Rudolphi, in Schrad. Journ. ii. (1799) 288, 289. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 936.
BORCKHAUSENIA, Roth, Catalect. fasc. ii. 56 (1800).
lucida, *Rudolphi, in Schrad. Journ.* ii. (1799) 228.—Afr. austr.
pubescens, *Burch. in Bot. Reg.* t. 214.—Afr. austr.

TEESDALEA, Aschers. Fl. Brandenb. 58 (1864) = seq.

TEESDALIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 88 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 92.
FOLIS, Dulac, Fl. Hautes-Pyr. 190 (1867).
GUELPINIA, Bast. Fl. Maine-et-Loire, Suppl. 35 (1812).
caulescens, Holl, in Hook. Journ. Bot. i. (1834) 21 = nudicaulis.
Iberis, DC. Syst. ii. 392 = nudicaulis.
irregularis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 693 = nudicaulis.
Lepidium, DC. Syst. ii. 392 = regularis.
nudicaulis, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 83.—Europ.; Reg. Mediterr.
petraea, Reichb. Fl. Germ. Excurs. 659 = Hutchinsia petraea.
regularis, *Sm. in Trans. Linn. Soc.* xi. (1815) 286.—Reg. Mediterr.; As. Min.

TEGANIUM, Schmid. Ic. Pl. 67. t. 18 (1747) = **Nolana**, Linn. (Convolv.).
procumbens, Schmid. Ic. Pl. ed. Kell. (1762) 67 = N. prostrata.

TEGANOCHARIS, Hochst. in Flora, xxiv. (1841) I. Intell. 42, sphalm. = TENAGOCHARIS, Hochst. = **Butomopsis**, Kunth (Alismac.).

TEGNERIA, Lilja, Flora oefver Sver. Odl. Vext. 76 (1839) = **Calandrinia**, H. B. & K. (Portulac.).
discolor, Lilja, l. c. 77 = C. discolor.
speciosa, Lilja, l. c. 76 = C. Menziesii.

TEICHMEYERIA, Scop. Introd. 267 (1777) = **Gustavia**, Linn. (Myrtac.).

TEICHOSTEMMA, R. Br. in Salt, Abyss. App. 65 (1814) = **Vernonia**, Schreb. (Compos.).
fruticosum, R. Br. l. c.—Abyss.

TEINOSOLEN, Hook. f. in Benth. & Hook. f. Gen. ii. 61 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
grandiflora, *Hook. f. in Herb. Kew.*—Bolivia.
macrosepala, *Hook. f. l. c.*—Bolivia.
Pearcei, *Hook. f. l. c.*—Bolivia.

TEINOSTACHYUM, Munro, in Trans. Linn. Soc. xxvi. (1868) 142. t. 3. *GRAMINEAE*, Benth. & Hook. f. iii. 1213.
attenuatum, *Munro, l. c.* 143.—Ind. or.
Griffithii, *Munro, l. c.*—Ind. or.
maculatum, *Trimen, in Journ. Bot.* xxiii. (1885) 273.—Zeylan.
schizostachyoides, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 89 = Melocanna Kurzii.
Wightii, *Munro, ex Bedd. Fl. Sylv.* ii. 233. t. 323.—Ind. or.

TEKTONA, Linn. f. Suppl. 20 (1781) = **Tectona**, Linn. f. (Verben.).

TELANTHERA, R. Br. in Tuckey, Congo, 477, in nota (1818). *AMARANTACEAE*, Benth. & Hook. f. iii. 38.
ADOKETON, Rafin. New Fl. Am. i. 65 (1836).
AMARANTESIA, Hort. ex Regel, Gartenfl. (1869) 102.
BRANDESIA, Mart. Nov. Gen. et Sp. ii. 25. t. 125 (1826).
BUCHOLZIA, Mart. l. c. 49. t. 147 (1826).
STEIREMIS, Rafin. Fl. Tellur. iii. 40 (1836).
albida, *Moq. in DC. Prod.* xiii. II. 367.—Reg. Argent.
amoena, *Regel, Gartenfl.* (1869) 101.—Bras.
andicola, *Moq. in DC. Prod.* xiii. II. 373.—Peruv.
angustata, Anderss. Eugenies Resa, Bot. 61 = strictiuscula.
arenaria, Casar. ex Seub. in Mart. Fl. Bras. v. I. 185 = Alternanthera paronychioides.
argentea, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 168 = echinocephala.
Bettzichiana, *Regel, Ind. Sem. Hort. Petrop.* (1862) 28; *Gartenfl.* (1862) 178.—Bras. ?
brasiliana, Moq. in DC. Prod. xiii. II. 382 = Mogiphanes straminea.
canescens, *Moq. l. c.* 368.—Venezuela.
capituliflora, *Moq. l. c.* 381.—N. Granat.; Peruv.
caracasana, *Moq. l. c.* 370.—Venezuela.
cinerascens, Moq. l. c. 372 = mexicana.
Crucis, *Moq. l. c.* 362.—Ind. occ.
cyclophylla, *Seub. in Mart. Fl. Bras.* v. I. 174.—Bras.
densiflora, *Moq. in DC. Prod.* xiii. II. 366.—Am. trop.
dentata, Moq. l. c. 378 = Mogiphanes brasiliensis.
diffusa, Moq. l. c. 382 = Mogiphanes ramosissima.
echinocephala, *Moq. l. c.* 373.—Ins. Galapag.
elongata, *Moq. l. c.* 372.—Bras.
eupatorioides, *Remy, in C. Gay, Fl. Chil.* v. 223.—Chili.
ficoidea, *Moq. in DC. Prod.* xiii. II. 363.—Am. trop.
filifolia, *Moq. l. c.* 368.—Ins. Galapag.
flavescens, Moq. l. c. 382 = Mogiphanes flavescens.
flavicoma, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 166.—Ins. Galapag.
floridana, Chapm. Fl. S. U. St. 383.—Florida.
frutescens, *Moq. in DC. Prod.* xiii. II. 365.—Am. trop.
glaucescens, *Moq. l. c.* 369.—Ins. Galapag.
glomerata, Moq. l. c. 376 = pubiflora.
gomphrenoides, *Moq. l. c.* 377.—Peruv.
gracilis, *Moq. l. c.* 375.—Mexic.
grandifolia, *Moq. l. c.* 376.—Mexic.
hirtula, Moq. l. c. 380 = Mogiphanes hirtula.
junciflora, *Remy, in C. Gay, Fl. Chil.* v. 224.—Chili.

TELANTHERA :—
lanceolata, *Moq. in DC. Prod.* xiii. II. 371.—N. Granat.
longipes, Moq. l. c. 370 = Alternanthera longipes.
lupulina, *Moq. l. c.* 374.—Peruv.
luzuloides, *Moq. l. c.* 378.—Bolivia.
manillensis, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 404 = frutescens.
maritima, *Moq. in DC. Prod.* xiii. II. 364.—Am. trop.; Afr. trop.
Martii, *Moq. l. c.* 374.—Bras.
martinicensis, *Moq. l. c.* 366.—Ins. Martinic.
mexicana, *Moq. l. c.* 372.—Mexic.
Meyeriana, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1856) 24.—Peruv.
microcephala, *Moq. in DC. Prod.* xiii. II. 371.—Mexic.
minutiflora, *Seub. in Mart. Fl. Bras.* v. I. 177.—Bras.
Moquini, Webb, ex Moq. in DC. Prod. xiii. II. 379 = Mogiphanes villosa.
multicaulis, Moq. l. c. 380 = Mogiphanes straminea.
nana, *Moq. l. c.* 374.—Peruv.
nodifera, *Moq. l. c.* 367.—Reg. Argent.
nudicaulis, *Moq. l. c.* 369.—Ins. Galapag.
obovata, *Moq. l. c.* 370.—Mexic.
paniculata, *Moq. l. c.* 377.—N. Granat.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 315; *Contrib.* 83.—Paraguay.
peruviana, *Moq. in DC. Prod.* xiii. II. 366.—Peruv.
philoxeroides, *Moq. l. c.* 362.—Bras.
picta, C. Koch, Wochenschr. ix. (1866) 15 = Bettzichiana.
polygonoides, *Moq. in DC. Prod.* xiii. II. 363.—Am. trop.
polygonoides, Seem. Bot. Voy. Herald, 407 = Alternanthera sessilis.
porrigens, *Moq. in DC. Prod.* xiii. II. 377.—Peruv.
praelonga, *Moq. l. c.* 369.—Bras.
puberula, *Moq. l. c.* 372.—Bras.
pubiflora, *Moq. l. c.* 375.—Am. trop.
pungens, Moq. l. c. 371 = Alternanthera achyrantha.
pycnantha, *Moq. l. c.* 376.—Mexic.
ramosissima, Moq. l. c. 381 = Mogiphanes ramosissima.
Regelii, *Seub. in Mart. Fl. Bras.* v. I. 177.—Bras.
ruía, *Moq. in DC. Prod.* xiii. II. 375.—Bras.
serpyllifolia, *Moq. l. c.* 373.—Ind. occ.
strictiuscula, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 166.—Ins. Galapag.
strigosa, *Moq. in DC. Prod.* xiii. II. 370.—Java.
tomentosa, *Moq. l. c.* 380.—Peruv.
truxillensis, *Moq. l. c.* 367.—Peruv.
versicolor, *Regel, Ind. Sem. Hort. Petrop.* (1868) 83.—Bras.
vestita, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 169.—Ins. Galapag.
villosa, *Moq. in DC. Prod.* xiii. II. 379.—Peruv.
virgata, Moq. l. c. 382 = Mogiphanes straminea ?

TELEIANDRA, Nees & Meyen, in Linnaea, viii. (1833) 46 = **Ocotea**, Aubl. (Laurin.).
glauca, Nees & Meyen, l. c. (= *Oreodaphne Teleiandra*).—Bras.

TELEIANTHERA, Endl. Gen. 301 (1837) = **Telanthera**, R. Br. (Amarant.).

TELEKIA, Baumg. Enum. Stirp. Transs. iii. 149 (1816) = **Buphthalmum**, Linn. (Compos.).
africana, Hook. f. in Journ. Linn. Soc. vii. (1864) 201 = Anisoppapus africanus.
cordifolia, DC. Prod. v. 485 = B. speciosum.
ovata, C. Koch, in Linnaea, xxiii. (1850) 712 = B. speciosum.
speciosa, Baumg. Enum. Stirp. Transs. iii. 150 = B. speciosum.
speciosissima, DC. Prod. v. 485 = B. speciosissimum.

TELELOPHUS, Dulac, Fl. Hautes-Pyr. 353 (1867) = **Seseli**, Linn. (Umbell.).
tenuifolius, Dulac, l. c. (= *Dethawia tenuifolia*).—Mont. Pyren.

TELEPHIASTRUM, Medic. Phil. Bot. i. 100 (1789) = **Anacampseros**, Linn. (Portulac.).

TELEPHIOIDES, Tourn. ex Moench, Meth. Suppl. 310 (1802) = **Andrachne**, Linn. (Euphorb.).
procumbens, Moench, l. c. = A. Telephioides.

TELEPHIUM, Tourn. ex Linn. Syst. ed. I (1785). *FICOIDEAE*, Benth. & Hook. f. i. 857.
MEROPHRAGMA, Dulac, Fl. Hautes-Pyr. 365 (1867).
album, *Gueldenst. Reise*, ii. 209.—Ross.
alternifolium, DC. Prod. iii. 366, sphalm. = oppositifolium.
alternifolium, Moench, Meth. 231 = Imperati.
glandulosum, *Bertol. Misc. Bot.* i. 18.—Assyria.
Imperati, *Linn. Sp. Pl.* 271.—Reg. Mediterr.; Oriens.
laxiflorum, DC. Prod. iii. 366 = Orygia decumbens.
madagascariense, *Baker, in Journ. Linn. Soc.* xxi. (1884) 347.—Madag.
maximum, Eichw. Skizze, 159 = Sedum maximum.
oligospermum, *Steud. ex Boiss. Fl. Orient.* i. 754.—Kurdist.
oppositifolium, *Linn. Sp. Pl.* ed. II. 388.—Afr. bor.
orientale, Boiss. Diagn. Ser. I. x. 11 = Imperati.
purpureum, Eichw. Skizze, 159 = Sedum Telephium.
repens, Lam. Fl. Fr. iii. 71 = Imperati.
sphaerospermum, *Boiss. Diagn.* Ser. I. iv. 12.—Aegypt.; Arab.
triphyllum, Eichw. Skizze, 159 = Sedum Telephium.

TELESIA, Rafin. Fl. Tellur. ii. 41 (1836) = **Zexmenia**, La Llave (Compos.).
aurea, Rafin. l. c. = Z. aurea.

TELESILLA, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1083 (1848), nomen. *ASCLEPIADEAE*.
cynanchioides, Klotzsch, l. c.—Guian.

TELESMIA, Rafin. Alsog. Am. 15 (1838) = **Salix**, Tourn.
hermaphrodita, Rafin. l. c. = S. pentandra.

TELESONIX, Rafin. Fl. Tellur. ii. 69 (1836) = **Saxifraga**, Tourn.
Jamesi, Rafin. l. c. = S. Jamesii.

TELESTRIA, Rafin. Sylva Tellur. 122 (1838) = **Bauhinia**, Linn. (Legumin.).
purpurea, Rafin. l. c. = B. purpurea.
racemosa, Rafin. l. c. = B. racemosa.

TELFAIRIA, Hook. Bot. Mag. t. 2751–2752 (1827). *CUCURBITACEAE*, Benth. & Hook. f. i. 821.
AMPELOSICYOS, Thou. Hist. Vég. Isles Afr. Austr. 68. t. 22 (1807).
JOLIFFIA, Boj. ex Delile, in Mém. Soc. Hist. Nat. Par. iii. (1827) 314. t. 6.
occidentalis, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 524.—Afr. trop.
pedata, *Hook. Bot. Mag.* t. 2751–2752.—Afr. trop.

TELFAIRIA, Newm. ex Hook. Bot. Misc. i. (1830) 291 = **Buettneria**, Loefl.
volubilis, Newm. ex Hook. l. c. = B. heterophylla.

TELINA, E. Mey. Comm. Pl. Afr. Austr. 67 (1835) = **Lotononis**, Eckl. & Zeyh. (1836) (Legumin.).
angustifolia, E. Mey. l. c. 70 = L. angustifolia.
brevifolia, Eckl. & Zeyh. ex Drège, in Linnaea, xix. (1847) 639.—Afr. austr.
Capnitidis, E. Mey. Comm. Pl. Afr. Austr. 69 = L. Capnitis.
cytisoides, E. Mey. l. c. 68 = L. cytisoides.
eriocarpa, E. Mey. l. c. = L. cytisoides.
excisa, E. Mey. l. c. 70 = L. prostrata.
genuflexa, E. Mey. l. c. 69 = L. genuflexa.
heterophylla, E. Mey. l. c. = L. azurea.
prostrata, E. Mey. l. c. = L. prostrata.
striata, E. Mey. l. c. 68 = L. carnosa.
varia, E. Mey. l. c. 70 = L. varia.
villosa, E. Mey. l. c. = L. villosa.

TELINARIA, Presl, Bot. Bemerk. 49 (1844); in Abh. Boehm. Ges. V. iii. (1845) 566 = **Cytisus**, Linn. (Legumin.).
acanthoclada, Presl, l. c. 136 = Genista acanthoclada.
albida, Presl, l. c. = Genista albida.
anglica, Presl, l. c = Genista anglica.
aprutia, Presl, l. c. = Genista Michelii ?
aspalathoides, Presl, l. c. = Genista aspalathoides.
biflora, Presl, l. c. = C. Fontanesii.
canariensis, Presl, l. c. 49 = C. canariensis.
candicans, Presl, l. c. = C. candicans.
cinerea, Presl, l. c. 136 = Genista cinerea.
congesta, Presl, l. c. 49 = Genista congesta.
Cumingii, Presl, l. c. 136 = Anarthrophyllum Cumingii.

TELINARIA :—
florida, Presl, Bot. Bemerk. 136 = Genista florida.
Friedrichsthaliana, Presl, l. c. = Genista acanthoclada.
hispanica, Presl, l. c. = Genista hispanica.
holopetala, Presl, l. c. = Genista radiata.
horrida, Presl, l. c. = Genista horrida.
humifusa, Presl, l. c. = Genista humifusa.
linifolia, Presl, l. c. 49 = Cytisus linifolius.
Lobelii, Presl, l. c. 136 = Genista aspalathoides.
pilosa, Presl, l. c. = Genista pilosa.
radiata, Presl, l. c. 135 = Genista radiata.
ramosissima, Presl, l. c. 49 = C. canariensis.
rosmarinifolia, Presl, l. c. = C. rosmarinifolius.
sagittalis, Presl, l. c. 136 = Genista sagittalis.
Salzmanni, Presl, l. c. 49 = Genista aspalathoides.
sericea, Presl, l. c. 136 = Genista sericea.
stenopetala, Presl, l. c. 49 = Genista stenopetala.
triquetra, Presl, l. c. = Genista triquetra, *L'Herit.*
umbellata, Presl, l. c. 135 = Genista umbellata.
virgata, Presl, l. c. 49 = Genista virgata.

TELINE, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 342; Phil. Bot. i. 203 (1789) = **Cytisus**, Linn. (Legumin.).
canariensis, Webb & Berth. Phyt. Canar. ii. 37 = C. canariensis.
candicans, Medic. Phil. Bot. i. 203 = C. candicans.
candidum, Steud. Nom. ed. II. ii. 665, sphalm. = praec.
congesta, Webb & Berth. Phyt. Canar. ii. 40 = Genista congesta.
linifolia, Webb & Berth. l. c. 41 = C. linifolius.
maderensis, Webb & Berth. l. c. 37 = C. maderensis.
medicagioides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 342 = C. candicans?
monspessulana, C. Koch, Dendrol. i. 30 = C. monspessulanus.
ramosissima, Webb & Berth. Phyt. Canar. ii. 38 = C. canariensis.
rosmarinifolia, Webb & Berth. l. c. 43. t. 44 = C. rosmarinifolius.
stenopetala, Webb & Berth. l. c. 39. t. 45 = Genista stenopetala.

TELIOSTACHYA, Nees, in Mart. Fl. Bras. ix. 71. t. 8 (1847) = **Lepidagathis**, Willd. (Acanth.).
alopecuroidea, Nees, in DC. Prod. xi. 263 = L. alopecuroidea?
alopecuroides, Nees, l. c. 262 = praec.
cataractae, Nees, in Mart. Fl. Bras. ix. 72.—Bras.
diffusa, Nees, l. c.—Bras.
hyssopifolia, Benth. in Hook. Niger Fl. 481 = L. hyssopifolia.
laguroidea, Nees, in DC. Prod. xi. 264 = L. laguroidea.
lanceolata, Nees, in Mart. Fl. Bras. ix. 74.—Bras.
lopecuroidea, Nees, l. c. 72, sphalm. = L. alopecuroidea.
microcarpa, Nees, l. c. 73.—Bras.

TELIPODUS, Rafin. Fl. Tellur. iii. 66 (1836) = **Philodendron**, Schott (Aroid.).
grandifolius, Rafin. l. c. = P. Simsii.

TELIPOGON, H. B. & K. Nov. Gen. et Sp. i. 335. t. 75. (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 587.
Alberti, *Reichb. f. in Linnaea*, xli. (1877) 27.—N. Granat.
andicola, *Reichb. f. in Bonplandia*, iii. (1855) 239.—Ecuador.
angustifolius, *H. B. & K. Nov. Gen. et Sp.* i. 336. t. 75.—N. Granat.; Venezuela.
antioquianus, *Reichb. f. in Linnaea*, xli. (1877) 72.—N. Granat.
Astroglossus, Reichb. f. Xenia Orch. i. 16. t. 7 = Trichoceros Astroglossus.
asuayanus, *Reichb. f. in Linnaea*, xli. (1877) 71.—Ecuador.
aureus, *Lindl. in Benth. Pl. Hartw.* 150.—Ecuador.
auritus, *Reichb. f. in Linnaea*, xli. (1877) 69.—N. Granat.
Benedicti, *Reichb. f. l. c.* 3.—Bolivia.
Boissierianus, *Reichb. f. in Bonplandia*, iv. (1856) 213.—Peruv.
Bowmani, *Reichb. f. in Linnaea*, xli. (1877) 69.—N. Granat.
Bruchmuelleri, *Reichb. f. l. c.* 28.—N. Granat.
chrysocrates, *Reichb. f. l. c.* 4.—N. Granat.
Croesus, *Reichb. f. l. c.* 70.—N. Granat.
dendriticus, *Reichb. f. Otia Bot. Hamb.* 6.—Ecuador.
dubius, *Reichb. f. in Linnaea*, xli. (1877) 104.—N. Granat.
falcatus, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 280.—N. Granat.
felinus, *Reichb. f. in Linnaea*, xli. (1877) 4.—Venezuela.
Fritillum, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 101.—Peruv.
Gustavi, *Reichb. f. in Linnaea*, xli. (1877) 105.—N. Granat.
gymnostele, *Reichb. f. l. c.* 70.—Peruv.
Hartwegii, *Reichb. f. Otia Bot. Hamb.* 6.—Ecuador.
hastatus, *Reichb. f. in Linnaea*, xli. (1877) 69.—N. Granat.
Hausmanni, Reichb. f. l. c. 70 = seq.
Hausmannianus, *Reichb. f. in Bonplandia*, ix. (1861) 213.—N. Granat.
hemimelas, *Reichb. f. in Linnaea*, xli. (1877) 72.—N. Granat.
ionopogon, *Reichb. f. l. c.* 27.—Ecuador.
Klotzscheanus, *Reichb. f. in Linnaea*, xxii. (1849) 851.—N. Granat.
latifolius, *H. B. & K. Nov. Gen. et Sp.* i. 336.—Peruv.
latifolius, Lindl. Orch. Linden. 23 = Hausmannianus.
macroglottis, *Reichb. f. in Linnaea*, xli. (1877) 71.—Ecuador.
musaicus, *Reichb. f. l. c.* 3.—N. Granat.
nitens, *Reichb. f. l. c.* 4.—N. Granat.
obovatus, *Lindl. Bot. Reg.* (1847) sub t. 27.—Peruv.
pachensis, *Reichb. f. in Linnaea*, xli. (1877) 105.—N. Granat.
pamplonensis, *Reichb. f. Xenia Orch.* i. 232.—N. Granat.
Papilio, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 101.—Peruv.
Patini, *Reichb. f. in Linnaea*, xli. (1877) 70.—N. Granat.
pogonostalix, *Reichb. f. l. c.* 72.—Ecuador.
polymerus, *Reichb. f. l. c.* 4.—N. Granat.
polyrrhizus, *Reichb. f. Otia Bot. Hamb.* 6.—Ecuador.
pulcher, *Reichb. f. in Linnaea*, xli. (1877) 105.—N. Granat.
radiatus, *Reichb. f. l. c.* 70.—Peruv.
Roezlii, *Reichb. f. l. c.* 4.—Venezuela.
tesselatus, *Lindl. in Benth. Pl. Hartw.* 150.—Peruv.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 104.—N. Granat.
Warszewiczii, *Reichb. f. Xenia Orch.* i. 231 (Warscewiczii).—Ecuador.
zephyrinus, *Reichb. f. in Linnaea*, xli. (1877) 71.—N. Granat.

TELIS, Linn. Syst. ed. I (1735) = **Trigonella**, Linn. (Legumin.).

TELLIMA, R. Br. in Frankl. Narr. 1st Journ. App. 765, in obs. (1823). *SAXIFRAGEAE*, Benth. & Hook. f. i. 637.
LITHOFRAGMA, Nutt. in Journ. Acad. Philad. vii. (1834) 26.
LITHOPHRAGMA, Torr. & Gray, Fl. N. Am. i. 583 (1840).
PLEURENDOTRIA, Rafin. Fl. Tellur. ii. 73 (1836).
affinis, *A. Gray, in Bot. Calif.* i. 198.—Am. bor. occ.
Bolanderi, *A. Gray, l. c.*—Calif.
Cymbalaria, *Steud. Nom.* ed. II. ii. 665; *Walp. Rep.* ii. 372.—Am. bor. occ.
glabra, Steud. Nom. l. c.; Walp. l. c. 371 = tenella.
grandiflora, *R. Br. in Frankl. Narr. 1st Journ. App.* 765, in obs.; *Dougl. ex Lindl. Bot. Reg.* t. 1178.—Am. bor. occ.
heterophylla, *Hook. & Arn. Bot. Beech. Voy.* 346.—Calif.
parviflora, *Hook. Fl. Bor. Am.* i. 289. t. 78 A.—Am. bor. occ.
tenella, *Steud. Nom.* ed. II. ii. 665; *Walp. Rep.* ii. 371.—Am. bor. occ.

TELMATOPHACE, Schleid. in Linnaea, xiii. (1839) 391 = **Lemna**, Linn.
arrhiza, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 70 = L. gibba.
cylindracea, Welw. ex Hegelm. Lemnaceen, 123 = Wolffia cylindracea.
gibba, Schleid. in Linnaea, xiii. (1839) 391 = Lemna gibba.
gibbosa, Montand. Guide Bot. 308 = L. gibba.
orbicularis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 70 = L. polyrhiza.
polyrhiza, Godr. Fl. Lorr. ed. I. iii. 18; ed. II. ii. 327 = L. polyrhiza.
transsilvanica, Schur, Enum. Pl. Transs. 635 = L. polyrhiza.

TELMATOPHILA, Ehrh. Beitr. iv. 147 (1789) = **Scheuchzeria**, Linn. (Naiad.).

TELMATOPHILA, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 170 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 236.
Scolymastrum, *Mart. ex Baker, l. c.*—Bras.

TELMATOSPHACE, Ball, in Journ. Linn. Soc. xvi. (1878) 685, sphalm. = TELMATOPHACE, Schleid. = **Lemna**, Linn.

TELMINOSTELMA, Fourn. in Mart. Fl. Bras. vi. IV. 218 (1885). *ASCLEPIADEAE.*
roulinioides, *Fourn. l. c.*—Bras.

TELMISSA, Fenzl, Pugill. Pl. Nov. Syr. 14 (1842) = **Sedum**, Tourn. (Crassul.).
microcarpa, Boiss. Fl. Orient. ii. 795 = Crassula microcarpa.
sedoides, Fenzl, Pugill. Pl. Nov. Syr. 15 = Crassula microcarpa.

TELOGYNE, Baill. Etud. Gén. Euph. 327 (1858) = **Trigonostemon**, Blume (Euphorb.).
indica, Baill. l. c. 328 = Trig. indicus.

TELOPEA, R. Br. in Trans. Linn. Soc. x. (1810) 197. *PROTEACEAE*, Benth. & Hook. f. iii. 184.
HYLOGYNE, Salisb. in Knight, Prot. 126 (1809), nomen prius.
oreades, *F. Muell. Fragm.* ii. 170.—Austral.
speciosissima, *R. Br. in Trans. Soc. Linn.* x. (1810) 198.—Austral.
truncata, *R. Br. l. c.*—Tasman.

TELOPEA, Soland. ex Baill. Etud. Gén. Euph. 345 (1858) = **Aleurites**, Forst. (Euphorb.).
perspicua, Soland. ex Seem. Fl. Vit. 223 = A. triloba.

TELOPOGON, Mutis, ex Spreng. Anleit. ii. I. 291 (1817) = **Telipogon**, H. B. & K. (Orchid.).

TELOXIS, Reichb. Handb. 238 (1837) = seq.

TELOXYS, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 289. t. 10 = **Chenopodium**, Tourn.
aristata, Moq. l. c. = C. aristatum.
cornuta, Torr. in Pacif. Rail. Rep. iv. 129 = C. cornutum.

TELUKRAMA, Rafin. Alsog. Am. 64 (1838) = **Cornus**, Tourn.
obovata, Rafin. l. c. (Quid?).—Am. bor.

TEMA, Adans. Fam. ii. 496 (1763) = **Panicum**, Tourn. (Gramin.).

TEMECARPUS, Proc. Linn. Soc. (1882-83) 2, err. typ. = **Semecarpus**, Linn. (Anacard.).

TEMMINCKIA, De Vriese, in Nederl. Kruidk. Arch. ii. I. (1851) 141 = **Scaevola**, Linn. (Gooden.).
Chamissoniana, De Vriese, l. c. 143 = S. Chamissoniana.
ciliata, De Vriese, l. c. 144 = S. Chamissoniana.
Gaudichaudii, De Vriese, l. c. = S. Gaudichaudii.
macrophylla, De Vriese, l. c. 146 (Quid?).—'N. Zel.'
Menziesiana, De Vriese, l. c. 144 = S. Chamissoniana.
micrantha, De Vriese, l. c. 145 = S. micrantha.
mollis, De Vriese, l. c. 147 = S. mollis.
tomentosa, De Vriese, l. c. 148 = S. tomentosa.

TEMNADENIA, Miers, Apocyn. S. Am. 207 (1878). *APOCYNACEAE.*
annularis, Miers, l. c. 216 = Prestonia annularis.
bicrura, Miers, l. c. 208 = Echites varia.
cordata, Miers, l. c. 212 = Echites cordata.

TEMNADENIA:—
corrugulata, *Miers, Apocyn. S. Am.* 215.—Bras.
Franciscea, Miers, l. c. 212 = Echites varia.
glaucescens, Miers, l. c. 214 = Mandevilla oaxacana.
lasiocarpa, Miers, l. c. 210 = Mandevilla hispida.
leptoloba, Miers, l. c. 211 = Echites leptoloba.
Lobbiana, Miers, l. c. 209 = Mandevilla hispida.
pallidiflora, Miers, l. c. 211 = Echites varia.
palustris, Miers, l. c. 213 = Echites tomentosa.
parviflora, Miers, l. c. 215 = Prestonia parviflora.
quinquangularis, Miers, l. c. 217 = Prestonia quinquangularis.
Riedelii, Miers, l. c. 216 (= *Haemadictyon Riedelii*).—Bras.
secundiflora, Miers, l. c. 211 = Echites secundiflora.
semidigyna, Miers, l. c. 213 = Echites semidigyna.
solanifolia, Miers, l. c. 214 (= *Haemadictyon solanifolium*).—Bras.
stellaris, Miers, l. c. 210 = Echites stellaris.
tenuicula, *Miers, l. c.* 216.—Bras.
tomentosa, Miers, l. c. 213 = Echites tomentosa, *Vahl.*
violacea, Miers, l. c. 208 = Echites violacea.
xanthostoma, Miers, l. c. 212 = Echites xanthostoma.

TEMNOCYDIA, Mart. ex DC. Prod. ix. 143 (1845) = **Bignonia**, Tourn.
aequinoctialis, Mart. l. c. 155 = B. aequinoctialis.
Carajura, Mart. l. c. 185 = Arrabidaea rosea.
densiflora, Mart. l. c. 181 = Lundia densiflora.
elliptica, Mart. l. c. 156 = B. Martiusiana.
flavida, Mart. l. c. 168 = B. flavida, flava.
laserpitiifolia, Mart. l. c. = B. laserpitiifolia.
spectabilis, Mart. l. c. 155 = B. aequinoctialis.

TEMNOPTERYX, Hook. f. in Benth. & Hook. f. Gen. ii. 72 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
sericea, *Hook. f. l. c.*—Afr. trop.

TEMPLETONIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 269 (1812). *LEGUMINOSAE*, Benth. & Hook. f. i. 474.
NEMATOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 20.
aculeata, *Benth. Fl. Austral.* ii. 170.—Austral.
Drummondii, *Benth. l. c.* 169.—Austral.
egena, *Benth. l. c.* 170.—Austral.
glauca, Sims, Bot. Mag. t. 2088 = retusa.
Hookeri, *Benth. Fl. Austral.* ii. 170.—Austral.
Muelleri, *Benth. l. c.* 169.—Austral.
retusa, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 269.—Austral.
sulcata, *Benth. Fl. Austral.* ii. 171.—Austral.

TEMU, Berg, in Linnaea, xxx. (1859–60) 710 = **Myrtus**, Tourn.
Cruckshanksii, Berg, l. c. 711 = Eugenia Cruckshanksii.
divaricatum, Berg, l. c. = Eugenia divaricata, *Berg.*

TEMUS, Molina, Sagg. Chile, 178 (1782) = **Myrtus**, Tourn.
moschata, Molina, l. c.—Chili.

TENAGEIA, Ehrh. Beitr. iv. 148 (1789); Reichb. Ic. Fl. Germ. ix. 22. t. 416 (1847) = **Juncus**, Tourn.
bufonia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 172 = J. bufonius.
bulbosa, Fourr. l. c. = J. supinus.
Gerardi, Fourr. l. c. = J. Gerardi.
multiflora, Fourr. l. c. = J. subulatus.
ranaria, Fourr. l. c. = J. bufonius.
sphaerocarpa, Reichb. Ic. Fl. Germ. ix. 22 = J. Tenageia.
squarrosa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 172 = J. squarrosus.
Vaillantii, Reichb. Ic. Fl. Germ. ix. 22, in obs. = J. Tenageia.

TENAGOCHARIS, Hochst. in Flora, xxiv. (1841) 369 = **Butomopsis**, Kunth (Alismac.).
alismoides, Hochst. l. c. i. Intell. 42 = B. lanceolata.
cordofana, Hochst. l. c. xxiv. (1841) 369 = B. lanceolata.
latifolia, Buchen. in Goett. Nachr. (1869) 238 = B. lanceolata.

TENARIS, E. Mey. Comm. Pl. Afr. Austr. 198 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 775.
rostrata, *N. E. Br. in Gard. Chron.* (1885) II. 39.—Afr. trop.
rubella, *E. Mey. Comm. Pl. Afr. Austr.* 198.—Afr. austr.

TENDANA, Reichb. f. in Oestr. Bot. Wochenbl. vi. (1857) 160 = **Micromeria**, Benth. (Labiat.).
Piperella, Reichb. f. l. c. = M. Piperella.

TENICROA, Rafin. Fl. Tellur. iii. 52 (1836) = **Urginea**, Steinh. (Liliac.).
fragrans, Rafin. l. c. 53 = U. fragrans.

TENOREA, Colla, Hort. Ripul. 137 (1824) = **Trixis**, P. Br. (Compos.).
Berteri, Colla, l. c. = Trixis radialis.
calyculata, Bert. ex Colla, l. c. = Trixis frutescens.

TENOREA, Gasp. Nov. Gen. Fici, 6 (1844); et Ricerche Caprif. e Fic. 81 (1845) = **Ficus**, Tourn. (Urtic.).
heterophylla, Gasp. ll. cc. 7; 81 = F. stipulata.

TENOREA, C. Koch, Dendrol. i. 670 (1869) = TENORIA, Spreng. = **Bupleurum**, Tourn. (Umbell.).

TENOREA, Rafin. Specch. i. 193 (1814) = **Zanthoxylum**, Linn. (Rutac.).
corymbosa, Rafin. l. c. = Z. corymbosum.

TENORIA, Dehnh. & Giord. Cat. Pl. Hort. Camald. ed. II. (1829) 24 = **Hygrophila**, R. Br. (Acanth.).
undulata, Dehnh. l. c. = H. spinosa.

TENORIA, Spreng. in Neue Schr. Naturf. Ges. Halle, ii. (1813) (Pl. Umb. Prod) 32 = **Bupleurum**, Tourn. (Umbell.).
arborescens, Spreng. l. c. = Heteromorpha arborescens.
baldensis, Spreng. l. c. = B. baldense.
canariensis, Spreng. in Neue Schr. Naturf. Ges. Halle, ii. (1813) 32 = Astydamia canariensis.
canescens, Schousb. ex Loud. Hort. Brit. 106 = B. gibraltaricum.
coriacea, Spreng. in Neue Schr. Naturf. Ges. Halle, ii. (1813) 32 = B. gibraltaricum.
difformis, Spreng. l. c. = B. difforme.
fruticescens, Spreng. l. c. = B. fruticescens.
fruticosa, Spreng. l. c. = B. fruticosum.
nuda, Spreng. l. c. = B. nudum.
plantaginea, Spreng. in Schult. Syst. vi. 676 = B. plantagineum.
pyrenaea, Spreng. in Neue Schr. Naturf. Ges. Halle, ii. (1813) 32 = B. angulosum.
romana, Schk. ex Spreng. in Schult. Syst. vi. 434 = Foeniculum vulgare.
spinosa, Spreng. in Neue Schr. Naturf. Ges. Halle, ii. (1813) 32 = B. spinosum.

TEPESIA, Gaertn. f. Fruct. iii. 72. t. 192 (1805) = **Hamelia**, Jacq. (Rubiac.).
dubia, Gaertn. f. l. c.—Chili.

TEPHEA, Delile, in Rochet, Sec. Voy. Choa, 340 (1846). *APOCYNACEAE.*
aequipetala, *Delile, l. c.*—Afr. trop.

TEPHIS, Adans. Fam. ii. 276 (1763) = **Polygonum**, Tourn.
frutescens, Rafin. Fl. Tellur. iii. 11 = Atraphaxis lanceolata?
parvifolia, Rafin. l. c. = Polygonella parvifolia.

TEPHRANTHUS, Neck. Elem. ii. 235 (1790) = **Phyllanthus**, Linn. (Euphorb.).

TEPHRAS, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 477 (1862) = **Galenia**, Linn. (Ficoid.).
papulosa, E. Mey. ex Harv. & Sond. l. c. = G. pruinosa.

TEPHROCACTUS, Lem. Cact. 88 (1868) = **Opuntia**, Tourn. (Cacteae).
andicolus, Lem. l. c. 88 = O. andicola.
aoracanthus, Lem. l. c. 89 = O. aoracantha.
calvus, Lem. l. c. 88 = O. Turpinii.
diadematus, Lem. l. c. = O. diademata.
platyacanthus, Lem. l. c. = O. diademata.
pusillus, Lem. l. c. = O. pusilla.
retrospinosus, Lem. l. c. = O. Parmentieri.
Turpinii, Lem. l. c. = O. Turpinii.

TEPHROSERIS, Reichb. Nom. 87 (1841) = **Senecio**, Tourn. (Compos.).
alpestris, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. alpestris.
angustifolia, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 102 = S. lugens.
aurantiaca, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. palustris.
campestris, Griseb. & Schenk, l. c.; Schur, Enum. Pl. Transs. 343 = S. campestris.
capitata, Griseb. & Schenk, l. c. = S. palustris.
cladobotrys, Griseb. & Schenk, l. c. 343 = S. cladobotrys.
Clusiana, Schur, Enum. Pl. Transs. 344 = S. alpestris.
crassifolia, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. alpestris.
crispa, Schur, Enum. Pl. Transs. 346 = S. crispatus.
Fussii, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. campestris.
longifolia, Griseb. & Schenk, l. c. 343; Schur, Enum. Pl. Transs. 344 = S. brachychaetus.
microrrhiza, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 146 = S. campestris.
palustris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = S. palustris.
papposa, Schur, Enum. Pl. Transs. 344 = S. lugens.
pratensis, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. campestris.
procera, Griseb. & Schenk, l. c. 343 = Cineraria procera.
rivularis, Schur, Enum. Pl. Transs. 347 = S. crispatus.
rupicola, Schur, l. c. 345.—Europ.
spathulaefolia, Schur, l. c. 344 = seq.
spathulifolia, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 342 = S. spathulaefolius.
stenophylla, Schur, in Oestr. Bot. Zeitschr. x. (1860) 326 = S. lugens.
transsilvanica, Schur, Enum. Pl. Transs. 345 = S. lugens?
Wolffii, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 102 = S. campestris.

TEPHROSIA, Pers. Syn. ii. 328 (1807). *LEGUMINOSAE*, Benth. & Hook. f. i. 496.
APODYNOMENE, E. Mey. Comm. Pl. Afr. Austr. 111 (1835).
BALBOA, Liebm. in Kjoeb. Vidensk. Meddel. (1856) 106.
BRISSONIA, Neck. Elem. iii. 36 (1790).
CATACLINE, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1214.
CRACCA, Linn. Fl. Zeyl. 139 (1747); et Sp. Pl. ed. I. 752 (1753), nomen prius.
EREBINTHUS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 210.
KIESERA, Reinw. in Syll. Ratisb. ii. (1828) 11.
MACRONYX, Dalz. in Hook. Kew Journ. ii. (1850) 35.
NEEDHAMIA, Scop. Introd. 310 (1777).
POGONOSTIGMA, Boiss. Diagn. Ser. I. ii. 39 (1843).
REINERIA, Moench, Meth. Suppl. 44 (1802).
REQUIENIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 91.
SEEMANNANTHA, Alef. in Bonplandia, x. (1862) 264.
XIPHOCARPUS, Presl, Symb. Bot. i. 13. t. 7 (1830).
acaciaefolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 106.—Afr. trop.
adscendens, Benth. in Mart. Fl. Bras. xv. i. 48 = purpurea.
adunca, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 432.—Am. trop.
aemula, *Harv. in Harv. & Sond. Fl. Cap.* ii. 210.—Afr. austr.
aequilata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 113.—Afr. trop.
ambigua, *D. Dietr. Syn. Pl.* iv. 1043, in syn.; *Chapm. Fl. S. U. St.* 96.—Am. bor.
amoena, Hort. Calc. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 113 = Hookeriana.
amoena, *E. Mey. Comm. Pl. Afr. Austr.* 109.—Afr. austr.
andongensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 116.—Afr. trop.
angulata, E. Mey. Comm. Pl. Afr. Austr. 109 = pallens.
angustifolia, Featherman, Bot. Rep. Louisiana Univ. (1870) 73 = onobrychoides.

TEPHROSIA:—
angustissima, *Shuttlew. ex Chapm. Fl. S. U. St.* 96.—Am. bor.
aniloides, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 258.—Porto Rico.
Anselii, *Hook. f. in Hook. Niger Fl.* 297.—Afr. trop.
anthylloides, *Hochst. ex Webb, in Hook. Kew Journ.* ii. (1850) 345; *Baker, in Oliver, Fl. Trop. Afr.* ii. 118.—Afr. trop.
Apollinea, Guill. & Perr. Fl. Seneg. Tent. 196 = anthylloides.
Apollinea, Klotzsch, in Peters, Reise Mossamb. Bot. 47 = villosa.
Apollinea, *Link, Enum. Hort. Berol.* ii. 252; *DC. Prod.* ii. 254.—Afr. trop.
arabica, *Steud. Nom.* ed. II. ii. 666.—Arab.
argentea, Pers. Syn. ii. 329 = villosa.
argentea, Wight, ex Wall. Cat. n. 5648 = fusca.
ascendens, Macfad. Fl. Jamaic. i. 257 = purpurea.
astragalina, *H. B. & K. Nov. Gen. et Sp.* vi. 464.—Ecuador.
astragaloides, *Benth. Fl. Austral.* ii. 208.—Austral.
barbigera, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 113.—Afr. trop.
Barclayana, Sweet, Hort. Brit. ed. II. 142 = mimosoides.
Barclayi, Baill. in Bull. Soc. Linn. Par. i. (1883) 388 = Dalbergia Barclayii.
Baueri, Benth. ex A. Gray, Bot. U. St. Expl. Exped. i. 408 = purpurea.
Bidwilli, *Benth. Fl. Austral.* ii. 210.—Austral.
biflora, *DC. Prod.* ii. 254.—Hab.?
biflora, Zipp. ex Span. in Linnaea, xv. (1841) 191 = purpurea.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 388.—Madag.
Bojeri, *Baill. l. c.* 389.—Madag.
brachycarpa, *F. Muell. ex Benth. Fl. Austral.* ii. 207.—Austral.
brachyloba, E. Mey. Comm. Pl. Afr. Austr. 110 = Dregeana.
brachystachya, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 99.—Nepal.
brachystachya, Zoll. ex Miq. Fl. Ind. Bat. i. 1. 299 = coarctata.
bracteolata, *Guill. & Perr. Fl. Seneg. Tent.* i. 194.—Afr. trop.
brevipes, *Benth. in Ann. Nat. Hist.* Ser. I. iii. (1839) 432.—Am. trop.
brunnea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 122.—Afr. trop.
calophylla, *Bedd. Ic. Pl. Ind. Or.* i. 36.—Ind. or.
candida, *DC. Prod.* ii. 249.—As. trop.
canescens, *E. Mey. Comm. Pl. Afr. Austr.* 109.—Afr. austr.
capensis, *Pers. Syn.* ii. 330.—Afr. austr.
capillipes, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 108.—Afr. trop.
capitulata, *Link, Enum. Hort. Berol.* ii. 251.—Ins. Sandvic.
caribaea, *DC. Prod.* ii. 251.—Am. trop.
cephalantha, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 119.—Afr. trop.
chilensis, *Trev. in Linnaea*, x. (1836) *Litt.* 73.—Chili.
chinensis, *Lindl. in Trans. Hort. Soc.* vii. (1830) 58.—China.
chrysophylla, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 49.—Mexic.
chrysophylla, *Pursh, Fl. Am. Sept.* ii. 489.—Am. bor.
cinerea, *Pers. Syn.* ii. 328.—Am. trop.
coarctata, *Miq. Fl. Ind. Bat.* i. 1. 299.—Ins. Lombok.
coccinea, Wall. Pl. As. Rar. i. 49 = tinctoria.
coerulea, Pers. Syn. ii. 329 = Dalea Mutisii.
Colonila, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 545 = purpurea.
Colutea, *Pers. Syn.* ii. 329.—Ind. occ.
Colutea, Wight, ex Wall. Cat. n. 5647 = Hookeriana.
concinna, *Baker, in Oliver, Fl. Trop. Afr.* ii. 112.—Afr. trop.
conferta, F. Muell. Census, 39 = singuliflora?
confertiflora, *Benth. in Miq. Pl. Jungh.* 208.—Java.
cordofana, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 118 = anthylloides.
coriacea, Benth. ex Harv. & Sond. Fl. Cap. ii. 208 = elongata, *E. Mey.*
coriacea, *Benth. Fl. Austral.* ii. 204.—Austral.
coronillaefolia, *DC. Prod.* ii. 252.—Ins. Borbon.

TEPHROSIA:—
coronilloides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 123.—Afr. trop.
crassa, Boj. ex Baker, in Journ. Bot. xx. (1882) 68 = purpurea.
crassifolia, *Benth. Bot. Voy. Sulph.* 80.—Mexic.
crocea, *Benth. Fl. Austral.* ii. 205.—Austral.
crotalarioides, Klotzsch, in Peters, Reise Mossamb. Bot. i. 45 = Indigofera crotalarioides.
dasyphylla, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 118.—Afr. trop.
davurica, Pers. Syn. ii. 328 = Astragalus dahuricus.
decidua, Steud. ex A. Rich. Tent. Fl. Abyss. i. 189 = subtriflora.
decora, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 123.—Afr. trop.
decumbens, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 7.—Nicaragua.
deflexa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 111.—Afr. trop.
densiflora, *Hook. f. in Hook. Niger Fl.* 296.—Afr. trop.
desertorum, *Scheele, in Linnaea*, xvii. (1843) 338.—Abyss.
dichotoma, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 415.—Ins. Philipp.
dichroocarpa, *Steud. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 31, nomen; *et ex A. Rich. Tent. Fl. Abyss.* i. 185.—Abyss.
diffusa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 210.—Afr. austr.
diffusa, Wight & Arn. Prod. 213 = purpurea.
digitata, *DC. Prod.* ii. 255.—Afr. trop.
dimorphophylla, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 116.—Afr. trop.
discolor, *E. Mey. Comm. Pl. Afr. Austr.* 111.—Afr. austr.
disperma, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 108.—Afr. trop.
dolichocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 101.—Reg. Argent.
domingensis, Pers. Syn. ii. 330 = purpurea.
Dregeana, *E. Mey. in Linnaea*, vii. (1832) 169.—Afr. austr.
drepanocarpa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 124.—Afr. trop.
dubia, Pers. Syn. ii. 330 = Lessertia diffusa.
Ehrenbergiana, Schweinf. Beitr. Fl. Aethiop. 18 = villosa.
elegans, Nutt. in Journ. Acad. Philad. vii. (1834) 105 = hispidula.
elegans, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 376.—Afr. trop.
elegans, Wall. Cat. n. 5634 = senticosa.
Elliottii, Benth. in Ann. Wien. Mus. ii. (1838) 127, in obs. = Galactia Elliottii.
elongata, *Hook. f. in Hook. Niger Fl.* 298.—Afr. trop.
elongata, *E. Mey. Comm. Pl. Afr. Austr.* 111.—Afr. austr.
emarginata, H. B. & K. Nov. Gen. et Sp. vi. 461 = toxicaria.
emeroides, *A. Rich. Tent. Fl. Abyss.* i. 190.—Abyss.
ensifolia, Harv. in Harv. & Sond. Fl. Cap. ii. 208 = elongata, *E. Mey.*
eriocarpa, *Benth. Fl. Austral.* ii. 207.—Austral.
eriosemoides, *Oliver, in Trans. Linn. Soc.* xxix. (1872) 57.—Afr. trop.
falcata, *Pers. Syn.* ii. 328.—Afr. austr.
fasciculata, *Hook. f. in Hook. Niger Fl.* 298.—Afr. trop.
ferruginea, Wall. ex Voigt, Hort. Suburb. Calc. 215 = Derris ferruginea.
filifolia, Pers. Syn. ii. 328 = Argyrolobium filiforme.
filiformis, *Pers. l. c.*—Am. austr.
filipes, *Benth. Fl. Austral.* ii. 208.—Austral.
flammea, *F. Muell. ex Benth. l. c.* 204.—Austral.
flexuosa, Chapm. ex Torr. & Gray, Fl. N. Am. i. 297 = hispidula.
flexuosa, *G. Don, Gen. Syst.* ii. 232.—Afr. trop.
Forbesii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 116, in nota.—Afr. trop.
Forrestiana, *F. Muell. Fragm.* xi. 98.—Austral.
frutescens, *DC. Prod.* ii. 256.—Honduras.
fruticosa, *DC. l. c.* 249.—Ind. or.
fulvinervis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 187.—Abyss.
fusca, *Wight & Arn. Prod.* 210.—Ind. or.
galegoides, R. Grah. ex Wall. Cat. n. 5649 = purpurea.

TEPHROSIA:—
glabrescens, Benth. Bot. Voy. Sulph., 81 = Cracca glabrescens.
glandulifera, *Benth. l. c.*—Ecuador.
glomeruliflora, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 86.—Afr. austr.
gracilipes, *Guill. & Perr. Fl. Seneg. Tent.* i. 194. t. 49.—Afr. trop.
gracilis, Nutt. Gen. Am. ii. 119 = hispidula.
Grahami, Wall. Cat. n. 5652 = tinctoria.
graminifolia, *F. Muell. ex Benth. Fl. Austral.* ii. 206.—Austral.
grandiflora, *Pers. Syn.* ii. 329.—Afr. austr.
gynothrix, Miq. in Linnaea, xviii. (1844) 29 = cinerea.
heterantha, *Griseb. in Goett. Abh.* xxiv. (1879) 101.—Reg. Argent.
heterophylla, Boj. ex Baker, in Journ. Bot. xx. (1882) 69 = Vigna angivensis.
heterophylla, Fisch. Hort. Gorenk. ed. II. (1812) 71 = tinctoria.
heterophylla, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 214.—Afr. trop.
Heyneana, Wall. Cat. n. 5631 = tinctoria.
Heyniana, Sweet, Hort. Brit. ed. II. 142 = tinctoria.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxix. (1879) 222.—Afr. trop.
hirsuta, *Schum. & Thonn. Beskr. Guin. Pl.* 377.—Afr. trop.
hirta, *Boj. Hort. Maurit.* 93, nomen.—Afr. trop.
hirta, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 546 = Hookeriana.
hispida, DC. Prod. ii. 250 = spicata.
hispidula, *Pers. Syn.* ii. 329.—Am. bor.
holosericea, Nutt. in Journ. Acad. Philad. vii. (1834) 105 = virginiana.
Hookeriana, *Wight & Arn. Prod.* 212.—Ind. or.; Malaya; Ins. Mascar.
huillensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 121.—Afr. trop.
humilis, *Guill. & Perr. Fl. Seneg. Tent.* i. 196.—Afr. trop.
hypargyrea, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 99 = tinctoria.
icthyneca, Bertol. Misc. Bot. xviii. 18 = Mundulea suberosa.
incana, Sweet, Hort. Brit. ed. II. 142; R. Grah. in Wall. Cat. n. 5644 = villosa.
indigofera, Bertol. Misc. Bot. xix. 9. t. 5 = purpurea.
inebrians, Welw. Apont. 573 = Vogelii.
intermedia, R. Grah. in Wall. Cat. n. 5632 = tinctoria.
interrupta, Hochst. ex Steud. in Flora, xxiv. (1841) 1. Intell. 31; et ex Steud. Nom. ed. II. ii. 666 = dichroocarpa.
juncea, *Benth. Fl. Austral.* ii. 208.—Austral.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 115.—Afr. trop.
Kotschyana, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 186 = bracteolata.
Kraussiana, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 87.—Afr. austr.
laevigata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 107.—Afr. trop.
Lagascana, *DC. Prod.* ii. 253.—Hab.?
lamprolobiodes, *F. Muell. Fragm.* ix. 64.—Austral.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* x. 11. (1843) 48.—Mexic.
lanceaefolia, Link, Enum. Hort. Berol. ii. 252 = purpurea.
lanceolata, R. Grah. in Wall. Cat. n. 5636 = purpurea.
lathyroides, *Guill. & Perr. Fl. Seneg. Tent.* i. 193.—Afr. trop.
leiocarpa, *A. Gray, Pl. Wright.* ii. 36.—N. Mexic.
leptoclada, *Benth. Fl. Austral.* ii. 207.—Austral.
leptostachya, DC. Prod. ii. 251 = purpurea.
leucantha, *H. B. & K. Nov. Gen. et Sp.* vi. 460.—Mexic.
Lindheimeri, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 172.—Texas.
linearis, *Pers. Syn.* ii. 330.—Afr. trop.
lineata, Schum. & Thonn. Beskr. Guin. Pl. 376 = purpurea.
littoralis, Pers. Syn. ii. 329 = cinerea.
lobata, R. Grah. in Wall. Cat. n. 5646 = purpurea.
longifolia, *Pers. Syn.* ii. 328.—Am. trop.
longipes, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 87.—Afr. austr.
lupinifolia, *DC. Prod.* ii. 255.—Afr. trop. et austr.

TEPHROSIA :—
lurida, *Sond. in Linnaea*, xxiii. (1850) 30.—Afr. austr.
luzoniensis, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 15.—Ins. Philipp.
Lyallii, *Baker, in Journ. Bot.* xx. (1882) 68.—Madag.
macrocarpa, *Benth. Fl. Austral.* ii. 209.—Austral.
macrophylla, Wall. ex Voigt, Hort. Suburb. Calc. 214 = Millettia auriculata.
macropoda, *Harv. in Harv. & Sond. Fl. Cap.* ii. 210.—Afr. austr.
madrensis, *Seem. Bot. Voy. Herald*, 280. t. 61.—Mexic.
mariana, *DC. Prod.* ii. 253.—Ins. Marian.
maxima, Pers. Syn. ii. 329 = purpurea.
melanocalyx, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 106.—Afr. trop.
mimosoides, *Pers. Syn.* ii. 330.—Ind. or.
Mitchelii, R. Grah. in Wall. Cat. n. 5641 = purpurea.
modesta, Steud. ex A. Rich. Tent. Fl. Abyss. i. 162 = Argyrolobium abyssinicum.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 463 = Cracca caribaea.
mollissima, Bertol. in Mem. Acad. Sc. Bolog. ii. (1850) 274 = spicata.
monantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 128.—Madag.
moschata, *Tussac, Fl. Antill.* ii. 22. t. 6.—Ind. occ.
mucronata, DC. Prod. ii. 251 = Lessertia mucronata.
multiflora, Featherman, Bot. Rep. Louisiana Univ. (1870) 73 = onobrychoides.
myrtifolia, *DC. Prod.* ii. 255.—Afr. austr.
nana, *Kotschy, ex Schweinf. Reliq. Kotsch.* 20. t. 16.—Afr. trop.
nematophylla, *F. Muell. Fragm.* ix. 63.—Austral.
nervosa, Pers. Syn. ii. 328 = tinctoria.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 6.—Nicaragua.
nitens, *Benth. in Seem. Bot. Voy. Herald*, 107.—Am. trop.
noctiflora, *Boj. Hort. Maurit.* 93, nomen; *et ex Baker, in Oliver, Fl. Trop. Afr.* ii. 112.—Afr. trop.
nubica, *Baker, l. c.* 125.—Afr. trop.
obcordata, *Baker, l. c.*—Afr. trop.
oblongata, *Benth. Fl. Austral.* ii. 205.—Austral.
oblongifolia, *E. Mey. Comm. Pl. Afr. Austr.* 108.—Afr. austr.
ochroleuca, Pers. Syn. ii. 329 = Cracca ochroleuca.
odorata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 404.—Ins. Socotra.
oligophylla, *Benth. Fl. Austral.* ii. 209.—Austral.
onobrychoides, *Nutt. in Journ. Acad. Philad.* vii. (1834) 104.—Am. bor. occ.
oroboides, Benth. Bot. Voy. Sulph. 81 = decumbens.
oroboides, *H. B. & K. Nov. Gen. et Sp.* vi. 462. t. 578.—Mexic.
oxygona, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 114.—Afr. trop.
pallens, *Pers. Syn.* ii. 329.—Afr. austr.
paniculata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 122.—Afr. trop.
parviflora, Wight, ex Wall. Cat. n. 5642 = purpurea.
pauciflora, *Wall. Cat.* n. 5635.—Ind. or.; Burma.
paucifolia, Nutt. Gen. Am. ii. 119 = spicata.
pedicellata, *Baker, in Oliver, Fl. Trop. Afr.* ii. 117.—Afr. trop.
penicillata, Benth. in Ann. Nat. Hist. Ser. I. iii. (1839) 431 = aduncа.
pentaphylla, Sweet, Hort. Brit. ed. II. 142; R. Grah. in Wall. Cat. n. 5650 = senticosa.
Perriniana, *DC. Prod.* ii. 251.—Am. austr.
persica, *Boiss. Diagn.* Ser. I. ii. 40.—Persia.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. 584 (Index) t. 9 = Mundulea suberosa.
phaeosperma, *F. Muell. ex Benth. Fl. Austral.* ii. 208.—Austral.
pinnata, *Pers. Syn.* ii. 330.—Afr. austr.
piscatoria, Pers. l. c. 329 = purpurea.
platycarpa, *Guill. & Perr. Fl. Seneg. Tent.* i. 195.—Afr. trop.
plicata, *Oliver, in Hook. Ic. Pl.* t. 1445.—Afr. austr.
Pogonostigma, *Boiss. Fl. Orient.* ii. 193.—Aegypt.; Arab.
polysperma, *Baker, in Oliver, Fl. Trop. Afr.* ii. 113.—Afr. trop.
polystachya, *E. Mey. Comm. Pl. Afr. Austr.* 109.—Afr. austr.

TEPHROSIA :—
polyzyga, *F. Muell. ex Benth. Fl. Austral.* ii. 206.—Austral.
porrecta, *Benth. l. c.*—Austral.
procumbens, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 54 = purpurea.
procumbens, Macfad. Fl. Jamaic. i. 256 = cinerea.
prostrata, Nutt. Gen. Am. ii. 120 = chrysophylla, *Pursh*.
Pseudo-piscatoria, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1200, nomen.—Guian.
pulchella, *Hook. f. in Hook. Niger Fl.* 299.—Afr. trop.
pulcherrima, Wight, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 112 = tinctoria.
pulchra, Coleb. in Wall. Cat. n. 5630 = Millettia pulchra.
pumila, Pers. Syn. ii. 330 = purpurea.
purpurea, *Pers. l. c.* 329.—Amphig. trop.
pusilla, *Pers. l. c.* 328.—Afr. austr.
racemosa, Sweet, Hort. Brit. ed. II. 142; Wight & Arn. Prod. 210 = Millettia racemosa.
radicans, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 121.—Afr. trop.
reflexa, *DC. Prod.* ii. 252.—Hab. ?
remotiflora, *F. Muell. ex Benth. Fl. Austral.* ii. 209.—Austral.
reptans, *Baker, in Oliver, Fl. Trop. Afr.* ii. 121.—Afr. trop.
reticulata, *Benth. Fl. Austral.* ii. 205.—Austral.
Rheedii, *DC. Prod.* ii. 255.—Ind. or.
rigida, *Baker, in Oliver, Fl. Trop. Afr.* ii. 114.—Afr. trop.
rigida, *Span. in Linnaea*, xv. (1841) 191.—Ins. Timor.
rigidula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 112.—Afr. trop.
rosea, *F. Muell. ex Benth. Fl. Austral.* ii. 211.—Austral.
rufescens, *Benth. in Linnaea*, xxii. (1849) 513.—Bras.
Rutenbergiana, *Vatke, in Bremen, Abh.* vii. (1882) 246.—Afr. trop.
Schiedeana, Schlecht. in Linnaea, xii. (1838) 299 = toxicaria.
secunda, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 115.—Afr. trop.
semiglabra, *Sond. in Linnaea*, xxiii. (1850) 29.—Afr. austr.
seminuda, *Boj. ex Baker, in Journ. Bot.* xx. (1882) 68.—Madag.
Senna, *H. B. & K. Nov. Gen. et Sp.* vi. 458.—N. Granat.
sennoides, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 292.—Ins. Celebes.
senticosa, *Pers. Syn.* ii. 330.—Ind. or.; Burma.
sericea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 107.—Afr. trop.
sericea, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 545 = purpurea.
sericea, DC. Prod. ii. 249 = Mundulea suberosa.
sericea, Hort. Bog. ex Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 76 = candida.
sericea, Pers. Syn. ii. 328 = Argyrolobium pumilum.
simplicifolia, *Franch. in Révoil, Çomali*, 28.—Afr. trop.
simplicifolia, *F. Muell. ex Benth. Fl. Austral.* ii. 206.—Austral.
singuliflora, *F. Muell. Fragm.* ix. 65.—Austral.
sophoroides, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 99.—Reg. Himal.
sphaerosperma, *Baker, in Oliver, Fl. Trop. Afr.* ii. 125.—Afr. trop.
sphaerospora, *F. Muell. in Wing, South. Sc. Record*, iii. (1883) 128.—Austral.
spicata, *Torr. & Gray, Fl. N. Am.* i. 296.—Am. bor.
spinosa, *Pers. Syn.* ii. 330.—Ind. or.; Malaya.
stipularis, *DC. Prod.* ii. 254.—Am. austr.
striata, Eckl. & Zeyh. ex Steud. Nom. ed. II. ii. 667 = capensis.
striata, Pers. Syn. ii. 328 = Lessertia pulchra.
stricta, R. Grah. in Wall. Cat. n. 5639 = purpurea.
stricta, *Pers. Syn.* ii. 329.—Afr. austr.
Stuartii, *Benth. Fl. Austral.* ii. 207.—Austral.
suberosa, DC. Prod. ii. 249 = Mundulea suberosa.
subsessilis, *R. Grah. in Wall. Cat.* n. 5969.—Ind. or.
subtriflora, *Hochst. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 117.—Abyss.

TEPHROSIA :—
Taylori, R. Grah. in Wall. Cat. n. 5637 = purpurea.
Telfairii, Sweet, Hort. Brit. ed. II. 142, nomen = Mundulea Telfairii.
tenella, *A. Gray, Pl. Wright.* ii. 36.—Mexic.
tenuis, *Wall. Cat.* n. 5970.—Ind. or.
ternata, DC. Prod. ii. 256 = Aspalathus ferruginea.
timoriensis, DC. l. c. 254 = purpurea.
tinctoria, R. Grah. in Wall. Cat. n. 5643 = purpurea.
tinctoria, *Pers. Syn.* ii. 329.—Ind. or.
tomentosa, *Pers. l. c.*—Arab.
totta, *Pers. l. c.* 328.—Afr. austr.
toxicaria, *Pers. l. c.* 329.—Am. trop.
trapezicarpa, DC. Prod. ii. 255 = Indigofera senegalensis.
trifoliata, DC. l. c. 256 = Argyrolobium sericeum.
uliginosa, Spreng. Syst. iii. 232 = Derris uliginosa.
uniflora, *Pers. Syn.* ii. 329.—Afr. trop.
uniovulata, *F. Muell. Fragm.* xi. 70.—Austral.
velutina, *Spreng. Syst.* iii. 232.—Ins. S. Doming.
venosa, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 47.—Mexic.
venustula, H. B. & K. Nov. Gen. et Sp. vi. 459 = cinerea.
vestita, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 15.—Sumatra; China.
viciaeformis, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 168.—Ind. or.
vicioides, *A. Rich. Tent. Fl. Abyss.* i. 188.—Afr. trop.
vicioides, *Schlecht. in Linnaea*, xii. (1838) 297.—Mexic.
villosa, *Pers. Syn.* ii. 329.—As. et Afr. trop.
villosa, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 54 = confertiflora.
virginiana, *Pers. Syn.* ii. 329.—Am. bor.
virginica, Bigel. Fl. Bost. ed. III. 296 = praec.
Vogelii, *Hook. f. in Hook. Niger Fl.* 296.—Afr. trop.
Wallichii, R. Grah. in Wall. Cat. n. 5640 = purpurea.

TEPHROTHAMNUS, Sch. Bip. in Pollichia, xx–xxi. (1863) 431 = **Vernonia**, Schreb. (Compos.).
calophyllus, Sch. Bip. l. c. 432.—Venezuela.
paradoxus, Sch. Bip. l. c. = V. paradoxa.
pycnanthus, Sch. Bip. l. c. 433 = V. pycnantha.

TEPHROTHAMNUS, Sweet, Hort. Brit. ed. II. 126 (1830) = **Argyrolobium**, Eckl. & Zeyh. (1836) (Legumin.).
tomentosus, Sweet, l. c. = A. Andrewsianum.

TEPION, Adans. Fam. ii. 131 (1763) = **Verbesina**, Linn. (Compos.).

TEPUALIA, Griseb. in Goett. Abh. vi. (1854) 119.
MYRTACEAE, Benth. & Hook. f. i. 710.
Philippiana, Griseb. l. c. 120 = stipularis ?
Philippii, Griseb. ex Phil. in Linnaea, xxviii. (1856) 637 = stipularis ?
stipularis, *Griseb. in Goett. Abh.* vi. (1854) 119.—Chili.

TERAMNUS, P. Br. Hist. Jamaic. 290 (1756); Sw. Prod. Veg. Ind. Occ. 105 (1788). *LEGUMINOSAE*, Benth. & Hook. f. i. 530.
Botrydium, Schott, in Wien. Zeitschr. iv. (1830) 1216; ex Endl. in Linnaea, vi. (1831) Litt. 55 = uncinatus.
clandestinus, Spreng. Syst. iii. 235 = Glycine clandestina.
flexilis, *Benth. in Journ. Linn. Soc.* viii. (1865) 265.—Reg. Himal.
grandiflorus, *Griseb. Cat. Pl. Cub.* 75.—Cuba.
hedysaroides, Spreng. Syst. iii. 235 = Glycine hedysaroides.
labialis, *Spreng. Syst.* iii. 235.—Cosmop. trop.
mollis, Benth. in Journ. Linn. Soc. viii. (1865) 265 = labialis.
obcordatus, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 382.—Madag.
oxyphyllus, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 255.—Burma.
parviflorus, Spreng. Syst. iii. 235 = labialis.
rhombifolius, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 120.—Panama.
tenuiflorus, Spreng. Syst. iii. 325 = Galactia dubia.
uncinatus, *Sw. Prod. Veg. Ind. Occ.* 105.—Am. trop.
volubilis, *Sw. l. c.*—Am. trop.
Wallichii, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 255.—Burma.

TERANA, La Llave, in La Naturaleza, vii. (1885) Apénd. 83. *COMPOSITAE.*
lanceolata, *La Llave, l. c.* 84.—Mexic.

TEREBINTHUS, [Tourn.] P. Br. Hist. Jamaic. 345 (1756) = **Pistacia**, Linn. (Anacard.).
Brownii, Jacq. Enum. Pl. Carib. 18 = Bursera gummifera.
Lentiscus, Moench, Meth. 345 = P. Lentiscus.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 356 = P. Lentiscus.

TEREBRARIA, Sesse, ex DC. Prod. iv. 458 (1830) = **Guettarda**, Linn. (Rubiac.).

TEREIANTHES, Rafin. Fl. Tellur. iii. 72 (1836) = **Reseda**, Tourn.
alba, Rafin. l. c. = R. alba.
fruticosa, Rafin. l. c. = R. alba.
glauca, Rafin. l. c. = R. glauca.
undata, Rafin. l. c. = R. alba.

TEREIANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 342 = **Reseda**, Tourn.
suffruticulosus, Fourr. l. c. = R. alba.

TEREIETRA, Rafin. Fl. Tellur. iv. 124 (1836) = **Ipomoea**, Linn. (Convolv.).
violacea, Rafin. l. c. 125 = I. violacea.

TEREIPHAS, Rafin. Fl. Tellur. iv. 95 (1836) = **Scabiosa**, Tourn. (Dipsac.).
rutefolia, Rafin. l. c. = S. rutaefolia.
urceolata, Rafin. l. c. = S. rutaefolia.

TEREMIS, Rafin. Sylva Tellur. 53 (1838) = **Lycium**, Linn. (Solan.).
elliptica, Rafin. l. c. 54 = L. barbarum.
turbinata, Rafin. l. c. = L. chinense.

TEREPIS, Rafin. Fl. Tellur. iii. 94 (1836) = **Salvia**, Tourn. (Labiat.).
Forskâlei, Rafin. l. c. = S. Forskohlei.

TERERA, [Domb.] ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 196, in obs. = **Miconia**, Ruiz & Pav. (Melastom.).
triplinervia, [Domb.] ex Naud. l. c. = M. Terera.

TERMINALIA, Linn. Mant. i. 21 (1767). *COMBRETACEAE*, Benth. & Hook. f. i. 685.
ARISTOTELIA, Comm. ex Lam. Encyc. i. 349 (1784).
BADAMIA, Gaertn. Fruct. ii. 90. t. 97 (1791).
BUCERAS, P. Br. Hist. Jamaic. t. 23 (1756), nomen prius.
BUCHENAVIA, Eichl. in Flora, xlix. (1866) 164.
BUCIDA, Linn. Syst. ed. X. 1025 (1759).
CATAPPA, Gaertn. Fruct. ii. 206. t. 127 (1791).
CHICHARRONIA, A. Rich. Ess. Fl. Cub. 529. t. 43 (1845).
CHUNCOA, Pav. ex Juss. Gen. 76 (1789).
FATREA, Juss. in Ann. Mus. Par. v. (1804) 223.
GIMBERNATEA, Ruiz & Pav. Prod. 138. t. 36 (1794).
HUDSONIA, Robins. in Lunan, Hort. Jamaic. ii. 310 (1814).
KNIPHOFIA, Scop. Introd. 327 (1777).
MYROBALANUS, Gaertn. Fruct. ii. 90. t. 97 (1791).
PAMEA, Aubl. Pl. Gui. 946. t. 359 (1775).
PENTAPTERA, Roxb. Hort. Beng. 34 (1814); Fl. Ind. ii. 437 (1832).
RESINARIA, Comm. ex Lam. Encyc. i. 349 (1784).
TANIBOUCA, Aubl. Pl. Gui. 448. t. 178 (1775).
VICENTIA, Allem. Diss. (1844).—Cf. Bot. Zeit. xii. (1854) 435.
actinophylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 22.—Bras.
acuminata, *Eichl. in Mart. Fl. Bras.* xiv. II. 92.—Bras.
acuta, Herb. Madr. ex Wall. Cat. n. 3980 C = paniculata.
acuta, Walp. Rep. ii. 61 = T. Chebula.
adamantium, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 241.—Bras.
adenopoda, *Miq. Fl. Ind. Bat. Suppl.* 327.—Sumatra.
alata, *D. Dietr. Syn. Pl.* ii. 1522.—Ind. or.
alata, Herb. Madr. ex Wall. Cat. n. 3980 B = paniculata.
alata, Roth, Nov. Pl. Sp. 379 = tomentosa.
amboinensis, Hort. ex Steud. Nom. ed. II. ii. 668 (Quid ?).—Malaya ?

TERMINALIA:—
angolensis, *O. Hoffm. in Linnaea*, xliii. (1880–82) 131.—Afr. trop.
angustifolia, Blanco, Fl. Filip. ed. I. 377 = T. Bellerica.
angustifolia, *Jacq. Hort. Vindob.* iii. 51.—Malaya.
angustifolia, Roxb. Hort. Beng. 33; Fl. Ind. ii. 437 = travancorensis.
angustifolia, Sauv. Fl. Cub. 38 = T. Buceras.
Arbuscula, *Sw. Prod. Veg. Ind. Occ.* 68.—Jamaic.
argentea, *Mart. Nov. Gen. et Sp.* i. 43.—Bras.
Arjuna, *Wight & Arn. Prod.* 314, in textu; *Bedd. Fl. Sylv.* t. 28.—Ind. or.
Arunja, Steud. Nom. ed. II. ii. 668, sphalm. = praec.
Aruta, Buch.-Ham. ex G. Don, Gen. Syst. ii. 659 = T. Chebula.
attenuata, Edgew. in Trans. Linn. Soc. xx. (1846) 46 = T. Bellerica.
australis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 240.—Bras.
avicennioides, *Guill. & Perr. Fl. Seneg. Tent.* i. 277. t. 64.—Afr. trop.
Badamia, *DC. Prod.* iii. 12.—Madag.
Badamia, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 92 = T. Catappa.
Belirica, Wall. Cat. n. 3968 = seq.
Bellerica, *Roxb. Pl. Corom.* ii. 54. t. 198.—Ind. or.; Malaya.
bengalensis, *Roxb. ex DC. Prod.* iii. 12.—Ind. or.; Malaya.
Benzoë, Pers. Syn. i. 485 = seq.
Benzoin, Linn. f. Suppl. 434 = angustifolia.
Berryi, Wight & Arn. Prod. 314 = T. Arjuna.
bialata, *Steud. Nom.* ed. II. ii. 668; *Kurz, For. Fl. Brit. Burma*, i. 456.—Ind. or.; Burma.
bialata, F. Villar, in Blanco, Fl. Filip. ed. III. Nov. App. 80 = T. Calamansanay.
biscutella, *Eichl. in Mart. Fl. Bras.* xiv. II. 88.—Bras.
Biticaria, Roxb. Hort. Beng. [90] = T. Bellerica.
Boivinii, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 95.—Madag.
brasiliensis, *Eichl. in Mart. Fl. Bras.* xiv. II. 91.—Bras.
brasiliensis, Raddi, ex Steud. Nom. ed. II. ii. 668 = januarensis.
brasiliensis, Spreng. Syst. ii. 359 = Gouania Blanchetiana.
Brownii, Fresen. in Mus. Senckenb. ii. (1837) 152. t. 9. f. 1 = cycloptera.
Buceras, *Wright, in Sauv. Fl. Cub.* 38; *Baill. Hist. Pl.* vi. 275.—Ind. occ.; Am. centr.
bursarina, *F. Muell. Fragm.* ii. 149.—Austral.
Cacoucia, Baill. Hist. Pl. vi. 275 = Cacoucia coccinea.
Calamansanay, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 310.—Ins. Philipp.
calophylla, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 102.—Madag.
capitata, Sauv. Fl. Cub. 38 = Hilariana.
Catappa, *Linn. Mant.* ii. 519.—As. trop.
Chebula, *Retz. Obs.* v. 31.—As. trop.
Chebula, Willd. ex Flem. in As. Res. xi. (1810) 181 = T. Bellerica.
Chicharronia, *Wright, in Sauv. Fl. Cub.* 38.—Cuba.
cicatricosa, *Boiss. Diagn.* Ser. I. vi. 54.—Persia.
ciliata, *Spreng. Syst.* ii. 359 (sp. dub.).—Ind. or.
circumalata, *F. Muell. Fragm.* iii. 91.—Austral.
citrina, *Roxb. ex Flem. in As. Res.* xi. (1810) 183.—Ind. or.; Malaya.
confertifolia, Steud. ex A. Rich. Tent. Fl. Abyss. i. 263 = cycloptera.
coriacea, [*Rottl.*] *ex Spreng. Fl. Hal. Mant.* 54.—Zeylan.
coriacea, Wight & Arn. Prod. 315 = tomentosa.
crenata, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 91.—Madag.
crenulata, Roth, Nov. Pl. Sp. 380; Wight & Arn. Prod. 314 = tomentosa.
cuneata, Roth, l. c. 379 = T. Arjuna.
cuneifolia, *Wall. Cat.* n. 3972.—Ind. or.
cycloptera, *R. Br. in Salt, Abyss. App.* 64.—Abyss.
dichotoma, E. Mey. Prim. Fl. Esseq. 177 = T. Taniboucа.
discolor, *F. Muell. Fragm.* iii. 92.—Austral.
discolor, *Spreng. Neue Entdeck.* ii. 111.—Bras.
disticha, Lodd. ex G. Don, in Loud. Hort. Brit. 412, nomen (Quid ?).—Ind. or.

TERMINALIA:—
edulis, Blanco, Fl. Filip. ed. II. 265 = T. Bellerica.
edulis, *F. Muell. Fragm.* ii. 151.—Austral.
eglandulosa, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 445 = T. Bellerica.
elegans, *Linden, Cat.* n. 21. (1867) 11.—Madag.
elliptica, Willd. Sp. Pl. iv. 969 = tomentosa.
erecta, Baill. Hist. Pl. vi. 266, 275 = Conocarpus erectus.
eriantha, Benth. in Hook. Kew Journ. ii. (1850) 240 = T. Taniboucа.
eriostachya, *A. Rich. Ess. Fl. Cub.* 524.—Cuba.
erythrocarpa, *F. Muell. Fragm.* ii. 150.—Austral.
erythrophylla, *Burch. Trav.* i. 400.—Afr. austr.
excelsa, *Liebm. ex Hemsl. Biol. Centr. Am. Bot.* i. 402.—Mexic.
exsculpta, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 93.—Madag.
fagifolia, *Mart. Nov. Gen. et Sp.* i. 42. t. 29.—Bras.
Fatraea, *DC. Prod.* iii. 12.—Madag.
firma, Mart. Nov. Gen. et Sp. i. 43 = lucida.
flavicans, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 99.—Madag.
foetidissima, *Griff. Notul.* iv. 685.—Malaya.
gangetica, Roxb. Hort. Beng. 23; Fl. Ind. ii. 437 = T. Chebula.
Gella, Dalz. in Hook. Kew Journ. iii. (1851) 227 = T. Bellerica.
glabra, R. Br. ex Benth. Fl. Austral. ii. 501, nomen = Muelleri.
glabra, Wight & Arn. Prod. 314 = T. Arjuna.
glabrata, *Forst. f. Prod.* 74.—Ins. Amicor.
glabrescens, *Mart. in Flora*, xx. (1837) II. *Beibl.* 124.—Bras.
glaucescens, *Planch. ex Benth. in Hook. Niger Fl.* 336.—Afr. trop.
gracilis, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 96.—Madag.
grandialata, *Eichl. in Mart. Fl. Bras.* xiv. II. 127, in obs.—Bras.
grandiflora, *Benth. Fl. Austral.* ii. 503.—Austral.
guyanensis, *Eichl. in Mart. Fl. Bras.* xiv. II. 88.—Bras.; Guian.
hemignosta, Steud. ex A. Rich. Tent. Fl. Abyss. i. 263 = cycloptera.
Hilariana, *Steud. Nom.* ed. II. ii. 668.—Bras.
Hilia, Buch.-Ham. ex Wall. Cat. n. 3970 C = citrina.
hirta, Steud. ex A. Rich. Tent. Fl. Abyss. i. 265 = Combretum Schimperianum.
horrida, Steud. Nom. ed. II. ii. 668 = Scleropyrum Wallichianum.
Hylobates, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 195.—Bras.
intermedia, Bert. ex Spreng. Syst. ii. 359 = T. Catappa.
intermedia, Span. in Linnaea, xv. (1841) 203 = microcarpa.
jamaicensis, Steud. Nom. ed. II. ii. 668, sphalm. = seq.
januarensis, *DC. Prod.* iii. 11.—Bras.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 603.—Java.
Kydiana, Roxb. ex Wall. Cat. sub n. 3974 = procera.
lanceolata, Mart. Nov. Gen. et Sp. i. 43. t. 29 = fagifolia.
lancifolia, Griff. Notul. iv. 685 = Combretum trifoliatum.
latifolia, Blanco, Fl. Filip. ed. I. 376 = T. Catappa.
latifolia, *Sw. Prod. Veg. Ind. Occ.* 68.—Jamaic.
latipes, *Benth. Fl. Austral.* ii. 501.—Austral.
laurinoides, *Teijsm. & Binn. Cat. Hort. Bog.* 237.—Java.
leiocarpa, Baill. Hist. Pl. vi. 265 = Anogeissus leiocarpa.
litoralis, *Seem. Fl. Vit.* 94.—Ins. Fiji.
lucida, *Hoffmgg. ex Mart. Nov. Gen. et Sp.* i. 43.—Bras.
macrocarpa, Steud. Nom. ed. II. ii. 668 = tomentosa.
macrophylla, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. II. 98, in syn.—Bras.
macroptera, *Guill. & Perr. Fl. Seneg. Tent.* i. 276.—Afr. trop.
macroptera, Mart. in Flora, xxiv. (1841) II. Beibl. 22 = grandialata.
macrostemon, Spreng. Syst. ii. 358 = Myrcia micrantha.
madagascariensis, Spreng. l. c. = T. Fatraea.
Maradu, Buch.-Ham. ex G. Don, Gen. Syst. ii. 660, in syn. = tomentosa.

TERMINALIA:—
Mariana, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 98. —Madag.
mauritiana, Blanco, Fl. Filip. ed. II. 264 = T. Catappa.
mauritiana, *Lam. Encyc.* i. 349.—Ins. Mascar.
melanocarpa, *F. Muell. Fragm.* iii. 92.—Austral.
microcarpa, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 457.—Ins. Timor.
microcarpa, F. Muell. Fragm. iii. 92 = Muelleri.
microphylla, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 193.—Bras.
modesta, Eichl. in Mart. Fl. Bras. xiv. II. 87 = subsericea.
modesta, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 102. —Madag.
mollis, *M. Laws. in Oliver, Fl. Trop. Afr.* ii. 417.—Afr. trop.
mollis, *Vidal, Phan. Cuming. Philipp.* 30, 112.—Ins. Philipp.
moluccana, Lam. Encyc. i. 349 = T. Catappa.
moluccana, Roxb. Hort. Beng. 33; Fl. Ind. ii. 432 = T. Bellerica.
moluccana, Wall. Cat. n. 3969 (Quid?).—Ind. or.
monaptera, Roth, Nov. Pl. Sp. 382 = paniculata.
Muelleri, *Benth. Fl. Austral.* ii. 500.—Austral.
myriocarpa, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 215.—Reg. Himal.; Burma.
Myrobalana, Roth, Nov. Pl. Sp. 378 = T. Catappa.
nitens, *Presl, Epim. Bot.* 214.—Ins. Philipp.
nitida, *Roxb. ex G. Don, Gen. Syst.* ii. 659 (sp. dub.). —Ind. or.
nitidissima, *Raeusch. Nom.* ed. III. 296.—Guian.
oblonga, *Steud. Nom.* ed. II. ii. 668.—Peruv.
oblongata, *F. Muell. Fragm.* ii. 152.—Austral.
obovata, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 241 = Hilariana.
obovata, *Steud. Nom.* ed. II. ii. 668.—Peruv.
octandra, Spreng. ex Eichl. in Mart. Fl. Bras. xiv. II. 93, nomen (Cupaniae sp.).—Bras.
odontoptera, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 217.—Guian.
ovalifolia, Rottl. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 447 = T. Arjuna.
ovata, Rottl. ex C. B. Clarke, l. c. = tomentosa.
ovatifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = T. Catappa.
oxycarpa, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 22. —Bras.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* 326.—Sumatra.
Pamaea, Steud. Nom. ed. II. ii. 668 = seq.
Pamea, *DC. Prod.* iii. 13.—Guian.
paniculata, *Roth, Nov. Pl. Sp.* 383.—Ind. or.
Papilio, *Hance, in Journ. Bot.* xv. (1877) 333.—Siam.
paraënsis, Mart. in Flora, xxiv. (1841) II. Beibl. 24 = T. Catappa.
parviflora, *Presl, Epim. Bot.* 214.—Ins. Philipp.
parviflora, Thw. Enum. Pl. Zeyl. 103 = T. Chebula.
pauciflora, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 101.—Madag.
pellucida, *Presl, Epim. Bot.* 214.—Ins. Philipp.
petiolaris, *A. Cunn. ex Benth. Fl. Austral.* ii. 502.—Austral.
phaeocarpa, *Eichl. in Mart. Fl. Bras.* xiv. II. 89.—Bras.
platyphylla, *F. Muell. Fragm.* ii. 150.—Austral.
platyptera, *F. Muell. l. c.* 151.—Austral.
polyantha, *Presl, Epim. Bot.* 213.—Ins. Philipp.
porphyrocarpa, *F. Muell. ex Benth. Fl. Austral.* ii. 501.—Austral.
procera, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 429.—Ins. Andam. et Nicob.
prunioides, *M. Laws. in Oliver, Fl. Trop. Afr.* ii. 415.—Afr. trop.
psidiifolia, Delile, Voy. Méroë, Bot. 92 = T. Arjuna.
pterocarya, *F. Muell. Fragm.* ii. 152.—Austral.
pumila, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 97. —Madag.
punctata, *Eichl. in Mart. Fl. Bras.* xiv. II. 85.—Bras.
punctata, Roth, Nov. Pl. Sp. 381 = T. Bellerica.
punctata, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. II. 98, in syn.—Bras.
pyrifolia, *Kurz, For. Fl. Brit. Burma*, i. 457; *et in Journ. As. Soc. Beng.* xlvi. (1877) II. 54.—Burma.
pyrifolia, Steud. Nom. ed. II. ii. 669 = Combretum pyrifolium.

TERMINALIA:—
reticulata, Roth, Nov. Pl. Sp. 381 = T. Chebula.
rhomboidea, *Spreng. Syst.* ii. 358.—Madag.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 616.—Ins. Samoa.
Riedelii, *Eichl. in Mart. Fl. Bras.* xiv. II. 92.—Bras.
rotata, Roxb. ex DC. Prod. iii. 12 = bengalensis.
rotundifolia, Lodd. ex G. Don, in Loud. Hort. Brit. 412, nomen (Quid?).—Ind. or.
rubrigemmis, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 102.—Madag.
Saja, Steud. Nom. ed. II. ii. 669 = myriocarpa.
salacifolia, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 663.—Afr. trop.
Schimperiana, Hochst. ex Delile, in Ferret & Galin. Voy. Abyss. iii. 133 = avicennioides.
scutifera, *Planch. ex M. Laws. in Oliver, Fl. Trop. Afr.* ii. 417.—Afr. trop.
sericea, *Burch. ex DC. Prod.* iii. 13.—Afr. austr.
sericea, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 243 = argentea.
sericocarpa, *F. Muell. Fragm.* ix. 159.—Austral.
suaveolens, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. II. 97, in syn.—Bras.
subcordata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 968 = T. Catappa.
subsericea, *Eichl. in Mart. Fl. Bras.* xiv. II. 127, in obs.—Bras.
sulcata, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 94. —Madag.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 326.—Sumatra.
Tanibouca, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 109; *Sm. in Rees, Cycl.* xxxv. n. 11.—Guian.
tarapotensis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 213.—Peruv.
Telfaireana, *Wall. Cat.* n. 3977.—Ins. Maurit.
Thozetii, *Benth. Fl. Austral.* ii. 500.—Austral.
tomentella, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 80 = T. Chebula.
tomentosa, *Mart. ex Eichl. in Mart. Fl. Bras.* xiv. II. 97, in syn.—Bras.
tomentosa, *Wight & Arn. Prod.* 314.—Ind. or.
travancorensis, *Wight & Arn. l. c.*—Ind. or.
trifoliata, *Spreng. Syst.* ii. 358.—Bras.
triopteris, Heyne, ex Wall. Cat. n. 3980 B = paniculata.
undulata, Steud. Nom. ed. II. ii. 669 = Combretum flagrocarpum.
Urjan, Royle, Illustr. Bot. Himal. 209 = T. Arjuna.
Vasivae, Spruce, ex Eichl. in Mart. Fl. Bras. xiv. II. 97 = suaveolens.
venosa, *Wall. Cat.* n. 3973.—Burma.
Vernix, *Lam. Encyc.* i. 350.—Ins. Molucc.
villosa, *Spreng. Neue Entdeck.* ii. 111.—Bras.
volucris, *R. Br. ex Benth. Fl. Austral.* ii. 498.—Austral.
zeylanica, Heurck & Muell. Arg. in Heurck, Obs. Bot. 220 = T. Chebula.

TERMINALIS, Medic. Theodora, 83 (1786) = **Dracaena**, Vand. (Liliac.).
angustifolia, Medic. l. c. = D. angustifolia.

TERMINTHOS, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 203 = **Pistacia**, Linn. (Anacard.).

TERMONTIS, Rafin. Chl. Aetn. 5 (1813) = **Linaria**, Tourn. (Scrophul.).
chalepensis, Rafin. l. c. = L. chalepensis.
cordata, Rafin. l. c.—Europ. austr.
purpurea, Rafin. l. c. 10 = L. purpurea.
villosa, Rafin. l. c. 5 = L. villosa.

TERNATEA, Tourn. ex Linn. Syst. ed. I (1735); H. B. & K. Nov. Gen. et Sp. vi. 415 (1823) = **Clitoria**, Linn. (Legumin.).
vulgaris, H. B. & K. l. c. = C. Ternatea.

TERNIOLA, Tul. in Arch. Mus. Par. vi. (1852) 189. t. 13. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 108.
DALZELLIA, Wight, Ic. v. II. 34 (1852).
LAWIA, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 112.
MNIANTHUS, Walp. Ann. iii. 443 (1853).
TULASNEA, Wight, Ic. tt. 1919–20 (1852).
foliosa, *Wedd. in DC. Prod.* xvii. 47.—Ind. or.
Lawii, *Wedd. l. c.*—Ind. or.

TERNIOLA:—
longipes, *Tul. in Arch. Mus. Par.* vi. (1852) 193. t. 13.—Ind. or.
pedunculosa, *Wedd. in DC. Prod.* xvii. 47.—Ind. or.
pulchella, *Tul. in Arch. Mus. Par.* vi. (1852) 192. t. 13. f. 4.—Ind. or.
ramosissima, *Wedd. in DC. Prod.* xvii. 47.—Ind. or.
zeylanica, *Tul. in Arch. Mus. Par.* vi. (1852) 190. t. 13. f. 3.—Zeylan.

TERNSTROEMIA, Mutis, ex Linn. f. Suppl. 39 (1781). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 182.
AMPHANIA, Banks, ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 408.
CLEYERA, Thunb. Nov. Gen. 68 (1783); Fl. Jap. 12 (1784).
DUPINIA, Scop. Introd. 195 (1777).
ERYTROCHITON, [Schlecht. in] Bot. Zeit. iv. (1846) 656.
HOFERIA, Scop. Introd. 194 (1777).
LLANOSIA, Blanco, Fl. Filip. ed. II. 319 (1845).
REINWARDTIA, Korth. Verh. Nat. Gesch. Bot. 101. t. 12 (1840?).
TAONABO, Aubl. Pl. Gui. 569. tt. 227, 228 (1775), nomen prius.
TONABEA, Juss. Gen. 262, 451 (1789).
VOELCKERIA, Klotzsch & Karst. in Endl. Gen. Suppl. iv. 66 (1847).
acuminata, *Jack, in Malay. Misc.* ii. (1822) n. VII. 26. —Sumatra.
albo-punctata, *Planch. ex Griseb. Cat. Pl. Cub.* 36.—Cuba.
aneura, Miq. Fl. Ind. Bat. Suppl. 477 = japonica.
bancana, *Miq. l. c.* 477.—Ins. Banca.
bifaria, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 143 = Eurya acuminata.
bilocularis, Roxb. Fl. Ind. ii. 522 = Saurauja tristyla.
brasiliensis, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 298. t. 59.—Bras.
brevipes, DC. in Mém. Soc. Phys. Genèv. i. (1822) 408 = elliptica.
carnosa, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 299. —Am. austr.
cernua, *Griseb. Cat. Pl. Cub.* 35.—Cuba.
clusiaefolia, *H. B. & K. Nov. Gen. et Sp.* v. 207. t. 463. f. 1.—Ecuador.
congestiflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 259.—N. Granat.
coriacea, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 352.—Malaya.
coriacea, Wall. Cat. n. 1453 = Adinandra acuminata.
corymbosa, *Sm. in Rees, Cycl.* xxxv. n. 6.—Guian.
crassifolia, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 363.—Am. austr.
crenata, Macfad. Fl. Jamaic. i. 114 = Symplocos octopetala.
crenulata, Wall. Cat. n. 3723 = Adinandra crenulata.
cuneata, *Poepp. ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 111.—Peruv.
cuneifolia, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 100.—Bras.
cuneifolia, Pav. ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 107 = elliptica.
cuspidata, Jack, in Malay. Misc. ii. (1822) n. VII. 28 (Saurauiae sp.).—Sumatra.
dasyanthera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 241.—Japon.
delicatula, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 108.—Guian.
dentata, Spreng. ex DC. l. c. i. (1821) 411 = Freziera undulata.
dentata, *Sw. Prod. Veg. Ind. Occ.* 81.—Guian.
dubia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 19 = japonica.
dumosa, Wall. Cat. n. 2245 = Adinandra dumosa.
elliptica, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 16 = sylvatica.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 81.—Ind. occ.; Mexic.
emarginata, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 104.—Zeylan.
fasciculata, Wall. Cat. subn. 1468 = Saurauja fasciculata.
flavescens, *Griseb. Cat. Pl. Cub.* 35.—Cuba.
fragrans, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 109 = japonica.
gedeënsis, Teijsm. & Binn. in Nat. Tijdschr. Neêrl. Ind. xiv. (1857) 156, sphalm. = seq.
gedehensis, *Teijsm. & Binn. l. c.* iii. (1852) 332.—Java.

TERNSTROEMIA:—
globiflora, *Ruiz & Pav. Syst.* 179.—Peruv.
globosa, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 106 = praec.
globuliflora, Steud. Nom. ed. II. ii. 669, sphalm. = globiflora.
integerrima, Wall. Cat. n. 1452 = Adinandra integerrima.
japonica, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—As. subtrop.
khasyana, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 110 = Illicium Griffithii.
lineata, DC. l. c. i. (1822) 409 = elliptica.
longipes, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1092, nomen.—Guian.
Lushia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 225 = Cleyera ochnacea.
Lushia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 109 = japonica.
macrocarpa, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 15 = penangiana.
macrocarpa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 259.—N. Granat.
macrophylla, Wall. Cat. n. 3663 = Pyrenaria acuminata.
meridionalis, *Mutis, Amer.* i. t. 9; *ex Linn. f. Suppl.* 264.—N. Granat.
meridionalis, Sw. Prod. Veg. Ind. Occ. 81 = elliptica.
micrantha, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 142, 145.—Java.
microcarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 241.—Am. austr.
multiflora, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 108 = dentata.
obovalis, *A. Rich. Ess. Fl. Cub.* 221. t. 25.—Ind. occ.
occidentalis, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 106 = globiflora.
panduana, Wall. Cat. sub n. 1470 = Saurauja punduana.
paniculata, Wall. l. c. sub n. 1469 = Saurauja napaulensis.
patens, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 109.—Sumatra.
Pavoniana, *Moric. l. c.* vii. (1836) 258. t. 13.—Peruv.
pedunculariis, DC. l. c. i. (1822) 409 = elliptica.
pedunculariis, Seem. Bot. Voy. Herald, 87 = Seemanni.
penangiana, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 110.—Malaya.
pentapetala, Jack, in Malay. Misc. i. (1820) n. v. 40 = Saurauja tristyla.
punctata, *Sw. Prod. Veg. Ind. Occ.* 81.—Guian.
quinquepartita, *Ruiz & Pav. Syst.* 180.—Peruv.
racemosa, D. Don, Prod. Fl. Nep. 225 = Saurauja napaulensis.
reticulata, Wall. Cat. n. 2246 = Adinandra integerrima.
revoluta, Splitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 99 = punctata.
Roraimae, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1092, nomen.—Guian.
rubicunda, *Klotzsch, l. c.* 1092, 1175, nomen.—Guian.
rubiginosa, Jack, in Malay. Misc. i. (1820) n. v. 39 (= *Cleyera rubiginosa*).—Sumatra.
Ruiziana, *Moric. in Mém. Soc. Phys. Genèv.* vii. (1836) 257. t. 12.—Peruv.
salicifolia, DC. l. c. i. (1822) 411 = Freziera undulata.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 362.—Guian.
Seemanni, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 260.—Panama.
sericea, Wall. Cat. n. 1454 = Adinandra villosa.
serrata, *Jack, in Malay. Misc.* ii. (1822) n. VII. 27.—Sumatra.
serrata, Roxb. Hort. Beng. 40; Fl. Ind. ii. 521 = Saurauja Roxburghii.
serrulata, Pav. ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 110 = Cleyera serrulata.
sessiliflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 577.—Bras.
siphilitica, Pav. ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 110 = Cleyera siphilitica.
suborbicularis, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1092, nomen.—Guian.
sylvatica, Cham. & Schlecht. in Linnaea, v. (1830) 220 = globiflora.
sylvatica, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 106.—Mexic.
Tepezapote, Cham. & Schlecht. l. c. vi. (1831) 420 = elliptica. ?

TERNSTROEMIA:—
Toquian, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 19.—Ins. Philipp.
trilocularis, Roxb. Hort. Beng. [93] = Saurauja tristyla.
venosa, *Spreng. Neue Entdeck.* ii. 162.—Bras.
verticillata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1092, nomen.—Guian.
vitiensis, *Seem. Fl. Vit.* 14.—Ins. Fiji.
Wightii, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 109 = japonica.

TEROBERA, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Cyp. 164 (1855) = **Cladium**, P. Br. (Cyperac.).
scirpoidea, Steud. ll. cc. = C. scirpoideum.

TEROGIA, Rafin. Fl. Tellur. iii. 56 (1836) = **Ortega**, Loefl. (Caryoph.).
dichotoma, Rafin. l. c. = O. dichotoma.

TERPNANTHUS, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 152. t. 19 = **Spiranthera**, A. St. Hil. (Rutac.).
jasminoides, Nees & Mart. l. c. 178 = S. odoratissima.

TERPNOPHYLLUM, Thw. in Hook. Kew Journ. vi. (1854) 70. t. 2 C = **Garcinia**, Linn. (Guttif.).
acuminatum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xix. (1860) 364 = G. terpnophylla.
zeylanicum, Thw. in Hook. Kew Journ. vi. (1854) 70. t. 2. f. 1 = G. terpnophylla.

TERRANEA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 11. t. 23 = **Erigeron**, Linn. (Compos.).
fernandeziana, Colla, l. c. = E. fruticosus.

TERSONIA, Moq. in DC. Prod. xiii. II. 40 (1849). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 87.
GYRANDRA, Moq. l. c. (1849).
brevipes, *Moq. l. c.*—Austral.
brevipes, Moq. l. c. (quoad ♂) = Gyrostemon ramulosus.
subvolubilis, *Benth. Fl. Austral.* v. 150.—Austral.

TERTREA, DC. Prod. iv. 481 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 108.
SCHIEDEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 186.
martinicensis, *DC. Prod.* iv. 481.—Ins. Martinic.

TERTRIA, Schrank, Fl. Monac. iii. 249 (1816) = **Polygala**, Tourn.
Chamaebuxus, Schrank, l. c. = P. Chamaebuxus.

TERUSTROEMIA, Jack, in Malay. Misc. i. (1820) n. v. 39, err. typ. = **Ternstroemia**, Mutis.

TESOTA, C. Muell. in Walp. Ann. iv. 479 (1857) = **Olneya**, A. Gray (Legumin.).

TESSARANDRA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. viii. (1851) 198 = **Linociera**, Sw. (Oleac.).
fluminensis, Miers, l. c.—Bras.

TESSARANTHIUM, Kellogg, in Proc. Calif. Acad. ii. (1862) 142 = **Frasera**, Walt. (Gentian.).
radiatum, Kellogg, l. c. 144 = F. speciosa.

TESSARIA, Ruiz & Pav. Prod. 112. t. 24 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 291.
GYNETERIA, Spreng. Anleit. ii. II. 899 (1818).
GYNHETERIA, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140.
MONOPHALACRUS, Cass. in Dict. Sc. Nat. liii. 235 (1828).
PHALACROMESUS, Cass. l. c. (1828).
POLYPAPPUS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 178.
absinthioides, *DC. Prod.* v. 457.—Chili.
absinthoides, Griseb. in Goett. Abh. xxiv. (1879) 185 = integrifolia.
ambigua, *DC. Prod.* v. 457.—Am. austr.
banksiaefolia, *Brongn. Bot. Voy. Coq.* t. 64.—Hab.?
boliviensis, *Sch. Bip. in Linnaea*, xxxiv. (1865-66) 534, nomen.—Bolivia.

TESSARIA:—
borealis, *A. Gray, Pl. Wright.* i. 102.—N. Mexic.
dentata, Ruiz & Pav. Syst. Veg. 213 = integrifolia.
integrifolia, *Ruiz & Pav. l. c.*—Bras.
legitima, DC. Prod. v. 456 = integrifolia.
mucronata, DC. l. c. = integrifolia.
redolens, Less. in Linnaea, vi. (1831) 151 (= *Monenteles redolens*).—N. Caled.

TESSENIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 318 = **Erigeron**, Linn. (Compos.).
muralis, Bubani, l. c. = E. acris.

TESSERANTHERUM, Curran, in Bull. Calif. Acad. i. n. III. (1885) 142, sphalm. = TESSARANTHIUM, Kellogg = **Frasera**, Walt. (Gentian.)

TESSERANTHIUM, Pritz. Ic. Ind. ii. 270 (1865), sphalm. = praec.

TESSIERA, DC. Prod. iv. 574 (1830) = **Spermacoce**, Linn. (Rubiac.).
lanigera, DC. l. c. = S. lanigera.
lithospermoides, DC. l. c. = S. lithospermoides.
pubescens, Miq. Stirp. Surin. Sel. 178.—Guian.

TESTUDINARIA, Salisb. in Burch. Trav. ii. 147 (1824). *DIOSCOREACEAE*, Benth. & Hook. f. iii. 744.
RHIZEMYS, Rafin. Fl. Tellur. iv. 26 (1836).
Elephantipes, *Salisb. in Burch. Trav.* ii. 148.—Afr. austr.
montana, *Burch. l. c.*—Afr. austr.
sylvatica, *Hort. Berol. ex Kunth, Enum. Pl.* v. 443.—Afr. austr.

TETA, Roxb. Hort. Beng. 24 (1814); Fl. Ind. ii. 165 (1832) = **Peliosanthes**, Andr. (Haemod.).
petiolaris, Voigt, Hort. Suburb. Calc. 650 = P. petiolaris.
viridiflora, Roxb. Hort. Beng. 24; Fl. Ind. ii. 165 = P. Teta.

TETANOSIA, Rich. ex M. Roem. Syn. Hesper. 23 (1846) = **Opilia**, Roxb. (Olacin.).
olacioides, M. Roem. l. c. = O. amentacea.

TETARIS, [Lindl. in] Chesn. Narrat. Euphr. Exped. (1868) t. ad p. 443. *BORAGINEAE*.
capitata, [*Lindl.*] *l. c.* t. ad p. 443.—Mesopot.
parviflora, [*Lindl.*] *l. c.*—Mesopot.

TETILLA, DC. Prod. iv. 667 (1830). *SAXIFRAGEAE*, Benth. & Hook. f. i. 640.
DIMORPHOPETALUM, Bert. 'Merc. Chil. (1829).'—Cf. Bull. Férussac, xx. (1830) 110; Am. Journ. Sc. Ser. I. xx. (1831) 256, nomen.
TETRAPLASIUM, Kunze, in Flora, xiv. (1831) 378.
hydrocotylaefolia, *DC. Prod.* iv. 667; et vii. 778.—Chili.

TETRACANTHUS, A. Rich. Fl. Cub. Fanerog. ii. 60 (1853). *COMPOSITAE*.
linearifolius, *A. Rich. l. c.*—Cuba.

TETRACANTHUS, Wright, ex Griseb. Cat. Pl. Cub. 153 (1866) = **Pinillosia**, Ossa (Compos.).
humilis, Wright, ex Griseb. l. c. = P. bellioides.

TETRACARPAEA, Benth. in Journ. Linn. Soc. iii. (1859) 72, sphalm. = TETRACRYPTA, Gardn. = **Anisophyllea**, (Rhizoph.).

TETRACARPAEA, Hook. f. in Hook. Ic. Pl. t. 264 (1840). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
tasmanica, *Hook. f. in Hook. Ic. Pl.* t. 264.—Tasman.

TETRACARPUM, Moench, Meth. Suppl. 240 (1802) = **Schkuhria**, Roth. (Compos.).
pinnatum, Moench, l. c. 241 = S. abrotanoides.

TETRACELLION, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 39 = TETRAPOMA, Turcz. = **Nasturtium**, R. Br. (Crucif.).
ellipsoideum, Turcz. l. c. = N. palustre.
globosum, Turcz. ex Steud. Nom. ed. II. ii. 669 = N. austriacum.

TETRACERA, Linn. Gen. ed. I. 345 (1737). *DILLENIACEAE*, Benth. & Hook. f. i. 12.
Assa, Houtt. Handleid. v. 275 (1776).
Calligonum, Lour. Fl. Cochinch. 342 (1790).
Delima, Linn. Diss. Dass. 19 (1747); Amoen. Acad. i. 403 (1749).
Delimopsis, Miq. Fl. Ind. Bat. i. II. 9 (1859).
Diploter, Rafin. Sylva Tellur. 165 (1838).
Eleiastis, Rafin. l. c. (1838).
Euryandra, Forst. Char. Gen. 81. t. 41 (1776).
Gynetra, Rafin. Sylva Tellur. 165 (1838).
Korosvel, Adans. Fam. ii. 442 (1763).
Leontoglossum, Hance, in Walp. Ann. ii. 18 (1851).
Roehlingia, Dennst. Schluess. Hort. Malab. 31 (1818).
Trachytella, DC. Syst. i. 410 (1818).
Traxilisa, Rafin. Sylva Tellur. 161 (1838).
Wahlbomia, Thunb. in Vet. Akad. Handl. Stockh. (1790) 215. t. 9.
acuminata, *DC. Syst.* i. 399.—Am. austr.
alata, *Presl, Rel. Haenk.* ii. 71.—Mexic.
alnifolia, DC. Syst. i. 401 = obtusata.
alnifolia, *Willd. Sp. Pl.* ii. 1243.—Afr. trop.
arborescens, *Jack, in Malay. Misc.* i. (1820) n. v. 45.—Sumatra.
aspera, *Willd. Sp. Pl.* ii. 1241.—Guian.
asperula, Miq. in Linnaea, xix. (1847) 133 = rotundifolia.
Assa, *DC. Syst.* i. 402.—Ind. or.; Malaya.
Blumei, *Walp. Rep.* i. 67.—Java.
Boiviniana, *Baill. Adansonia*, vii. (1866–67) 300. t. 7.—Afr. trop.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 76.—Borneo.
Breyniana, *Schlecht. in Linnaea*, viii. (1833) 174.—Bras.
Calinea, Willd. Sp. Pl. ii. 1241 = Doliocarpus Calinea.
canescens, Willd. ex Schlecht. in Linnaea, viii. (1833) 177, nomen (Myrtacea).—Bras.
carpinifolia, *Eichl. in Mart. Fl. Bras.* xiii. I. 92.—Bras.
castaneaefolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 22.—N. Granat.
ciliata, Poepp. ex Eichl. in Mart. Fl. Bras. xiii. I. 101 = Davilla cuspidulata.
costata, *Mart. ex Eichl. l. c.* 86.—Bras.
cuspidata, G. F. W. Mey. Prim. Fl. Esseq. 205 = Doliocarpus Rolandri.
Daemeliana, *F. Muell. Fragm.* v. 191.—Austral.
dichotoma, *Blume, Bijdr.* 3.—Java.
Doliocarpus, Willd. Sp. Pl. ii. 1241 = Doliocarpus Rolandri.
erecta, *Sesse & Moç. ex DC. Syst.* i. 404.—Mexic.
Euryandra, *Vahl, Symb. Bot.* iii. 71.—Malaya; N. Caled.
fagifolia, *Blume, Bijdr.* 4.—Java.
fagifolia, Willd. ex Schlecht. in Linnaea, viii. (1833) 174 = rotundifolia.
Gardneri, *Eichl. in Mart. Fl. Bras.* xiii. I. 91.—Bras.
gracilis, *Blume, Bijdr.* 4.—Java.
grandiflora, *Eichl. in Mart. Fl. Bras.* xiii. I. 92.—Bras.
Guillemini, Steud. Nom. ed. II. ii. 670 = alnifolia.
Heyneana, *Wall. Cat.* n. 6630.—Ind. or.
hispida, *Spreng. Syst.* ii. 629.—Bras.
hydrophila, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 20.—N. Granat.
hygrophila, *Kurz, For. Fl. Brit. Burma*, i. 22.—Ind. or.
jamaicensis, DC. Syst. i. 399 = volubilis.
japurensis, *Mart. & Zucc. in Abh. Akad. Muench.* i. (1832) 367.—Bras.
juncea, Hort. Angl. ex Steud. Nom. ed. II. ii. 670 (Quid?).—Austral.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 75.—Borneo.
laevigata, *Miq. Fl. Ind. Bat.* i. II. 8.—Sumatra.
laevis, *Vahl, Symb. Bot.* iii. 71.—Ind. or.; Malaya.
lasiocarpa, *Eichl. in Mart. Fl. Bras.* xiii. I. 84.—Bras.
laurifolia, Willd. ex Schlecht. in Linnaea, viii. (1833) 177 = Myrcia magnoliaefolia.
Lima, Willd. ex DC. Syst. i. 399 = Davilla rugosa.
lucida, Wall. Cat. n. 6631 = T. Euryandra.
lutea, Spreng. Neue Entdeck. ii. 164 = Ochna arborea.
macrocarpa, *Wall. Cat.* n. 6628.—Ind. or.; Malaya.

TETRACERA:—
macrophylla, Wall. ex Hook. f. & Thoms. Fl. Ind. i. 63, sphalm. = praec.
madagascariensis, *Willd. ex Schlecht. in Linnaea*, viii. (1833) 176.—Madag.
malabarica, Lam. Illustr. iii. 32 = T. Assa.
mexicana, *Eichl. in Mart. Fl. Bras.* xiii. I. 87, in obs.—Mexic.
monocarpa, *Blanco, Fl. Filip.* ed. I. 459.—Ins. Philipp.
multiflora, DC. Syst. i. 400 = Davilla Sagraeana.
multiflora, H. B. & K. ex Schlecht. in Linnaea, viii. (1833) 177, in obs. = Davilla multiflora.
nitida, *Vahl, Symb. Bot.* iii. 70.—Ins. Trinit.
Nordtiana, *F. Muell. Fragm.* v. 1.—Austral.
oblongata, *DC. Syst.* i. 399.—Bras.
obovata, DC. l. c. 401 = alnifolia.
obovata, Willd. Sp. Pl. ii. 1241 = Doliocarpus Soramia.
obtusata, *Planch. ex Oliver, Fl. Trop. Afr.* i. 12.—Afr. trop.
ovalifolia, *DC. Syst.* i. 400.—Ind. occ.; Guian.
pauciflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 88.—Madag.
Perriniana, *Spreng. Neue Entdeck.* ii. 164.—Am. austr.
plicata, Willd. ex Schlecht. in Linnaea, viii. (1833) 177 = Davilla Kunthii.
plumbea, Moric. ex Eichl. in Mart. Fl. Bras. xiii. I. 90 = Breyniana.
Poeppigiana, Schlecht. in Linnaea, viii. (1833) 174 = volubilis.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 113.—Panama.
potatoria, Afzel. ex G. Don, Gen. Syst. i. 69 = alnifolia.
Radula, *Eichl. in Mart. Fl. Bras.* xiii. I. 91. t. 23.—Bras.
rhamnifolia, *Presl, Rel. Haenk.* ii. 72.—Panama.
Rheedii, DC. Syst. i. 402 = laevis.
Riedeliana, *Eichl. in Mart. Fl. Bras.* xiii. I. 88.—Bras.
rigida, *Blume, Bijdr.* 4.—Java.
rotundifolia, *Sm. in Rees, Cycl.* xxxv. n. 7.—Am. trop.
rugosa, Guill. & Perr. Fl. Seneg. Tent. i. 2 = alnifolia.
rugosa, H. B. & K. ex Schlecht. in Linnaea, viii. (1833) 177, in obs. = Davilla Kunthii.
Rutenbergii, *Buchen. in Bremen, Abh.* vii. (1880) 12.—Madag.
salicifolia, *Presl, Rel. Haenk.* ii. 71.—Mexic.
sarmentosa, *Vahl, Symb. Bot.* iii. 70.—China.
scaberrima, *Miq. Fl. Ind. Bat.* i. II. 9.—Sumatra.
scabra, Hook. f. in Hook. Niger Fl. 203 = alnifolia.
Sellowiana, *Schlecht. in Linnaea*, viii. (1833) 175.—Bras.
senegalensis, *DC. Syst.* i. 401.—Afr. trop.
sericea, Blume, Bijdr. 3 = laevis.
sessiliflora, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 21.—N. Granat.
stricta, Willd. Sp. Pl. ii. 1241 = Doliocarpus strictus.
sumatrana, Miq. Fl. Ind. Bat. Suppl. i. 619 = fagifolia.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 265.—Guian.
Tigarea, DC. Syst. i. 403 = aspera.
tomentosa, Willd. Sp. Pl. ii. 1241 = Doliocarpus Rolandri.
triceras, Thou. ex Baill. in Bull. Soc. Linn. Par. i. (1882) 332 = madagascariensis.
trigyna, Roxb. ex Wall. Cat. n. 6629 A = T. Assa.
trigyna, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 645 = laevis.
tripetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 547.—Java.
ulmifolia, Willd. ex Schlecht. in Linnaea, viii. (1833) 177, nomen (Brockiae sp.?).—Bras.
volubilis, H. B. & K. Nov. Gen. et Sp. v. 50 = sessiliflora.
volubilis, *Linn. Sp. Pl.* 533.—Am. austr.
Wahlbomia, *DC. Syst.* i. 403.—Java.
Willdenowiana, Steud. Nom. ed. II. ii. 670 = rotundifolia.
Wuthiana, *F. Muell. Fragm.* x. 49.—Austral.

TETRACHEILOS, Lehm. Pl. Preiss. ii. 368 (1847) = **Acacia**, Willd. (Legumin.).
Meisneri, Lehm. l. c. = A. tetragonocarpa.

TETRACHNE, Nees, Fl. Afr. Austr. Gram. 375 (1841). *GRAMINEAE*, Benth & Hook. f. iii. 1171.
Dregei, *Nees, l. c.* 376.—Afr. austr.

TETRACHYRON, Schlecht. in Linnaea, xix. (1847) 744 = **Calea**, Linn. (Compos.).
manicatum, Schlecht. l. c. = C. manicata.

TETRACLEA, A. Gray, in Am. Journ. Sc. Ser. II. xvi. (1853) 98. *LABIATAE*, Benth. & Hook. f. ii. 1220.
Coulteri, *A. Gray, l. c.*—N. Mexic.

TETRACLIS, Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 271. t. 11. *EBENACEAE*, Benth. & Hook. f. ii. 666.
clusiaefolia, *Hiern, l. c.*—Madag.

TETRACME, Bunge, Del. Sem. Hort. Dorpat. (1836).—Cf. Linnaea, xii. (1838) Litt. 71. *CRUCIFERAE*, Benth. & Hook. f. i. 76.
contorta, *Boiss. Fl. Orient.* i. 317.—Afghan.
quadricornis, *Bunge, Del. Sem. Hort. Dorpat.* (1836).—*Cf. Linnaea*, xii. (1838) *Litt.* 71.—Turkestan.
recurvata, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 158.—Oriens.
secunda, *Boiss. Diagn.* Ser. II. i. 29.—Afghan.
Stocksii, *Boiss. Fl. Orient.* i. 317.—Beluchist.

TETRACOCCUS, Engelm. ex Parry, in West-Am. Scientist, i. (1885) 13. *EUPHORBIACEAE.*
dioicus, *Parry, l. c.* 13, 35.—Calif.
Engelmanni, S. Wats. in Proc. Am. Acad. xx. (1885) 373 = dioicus.

TETRACOCYNE, Turcz. in Bull. Soc. Nat. Mosc. (1863) I. 555 = **Ryania**, Vahl (Bixin.).
angustifolia, Turcz. l. c. = R. speciosa.
puberula, Turcz. l. c.—Guian.

TETRACRYPTA, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 314 = **Anisophyllea**, R. Br. (Rhizoph.).
cinnamomoides, Gardn. & Champ. l. c. = A. zeylanica.

TETRACTINOSTIGMA, Hassk. in Flora, xl. (1857) 533; Hort. Bog. Desc. 55 (1858) = **Aporosa**, Blume (Euphorb.).
lucidum, Miq. Fl. Ind. Bat. Suppl. 471.—Sumatra.
microcalyx, Hassk. in Flora, xl. (1857) 533 = A. microcalyx.

TETRACTIS, DC. Prod. i. 16 (1824), sphalm. = TETRACTYS, Spreng. (Ranunc.?).

TETRACTOMIA, Hook. f. Fl. Brit. Ind. i. 490 (1875). *RUTACEAE.*
majus, *Hook. f. l. c.* 491.—Malaya.
Roxburghii, *Hook. f. l. c.*—Malaya.

TETRACTYS, Spreng. Neue Entdeck. iii. 53 (1822). Genus dubium *RANUNCULACEARUM*.—Cf. Benth. & Hook. f. i. 3.
capensis, Spreng. l. c.—Afr. austr.

TETRADENIA, Benth. in Bot. Reg. sub t. 1300 (1830). *LABIATAE*, Benth. & Hook. f. ii. 1180.
fruticosa, *Benth. in Bot. Reg.* sub t. 1300.—Madag.

TETRADENIA, Nees, in Wall. Pl. As. Rar. ii. 61 (1831) = **Litsea**, Lam. (Laurin.).
consimilis, Nees, l. c. 64 = L. zeylanica.
foliosa, Nees, l. c. = L. zeylanica.
furfuracea, Nees, l. c. = L. zeylanica.
lanuginosa, Nees, l. c. = L. lanuginosa.
obovata, Nees, l. c. = Actinodaphne obovata.
Pseudo-Culitlawan, Nees, in Otto & Dietr. Allg. Gartenz. i. (1833) 373 = L. Pseudoculitlawan.
umbrosa, Nees, in Wall. Pl. As. Rar. ii. 64 = L. umbrosa, zeylanica.
zeylanica, Nees, l. c. = L. zeylanica.

TETRADIA, Benn. Pl. Jav. Rar. 233 (1844). *STERCULIACEAE*, Benth. & Hook. f. i. 219.
Horsfieldii, *Benn. l. c.*—Java.

TETRADIA, Thou. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 137 = **Tetrataxis**, Hook. f. (Lythrar.).
salicifolia, Thou. ex Tul. l. c. = Tetrataxis salicifolia.

TETRADICLIS, Stev. ex Bieb. Fl. Taur. Cauc. iii. 648.—Cf. 277 (1819). *RUTACEAE*, Benth. & Hook. f. i. 288.
ANATROPA, Ehrenb. in Linnaea, iv. (1829) 402.
caspia, [*Lindl. in*] *Chesn. Narrat. Euph. Exped.* (1868) t. ad p. 441.—Mesopot.
Eversmanni, *Bunge, in Linnaea*, xiv. (1840) 178.—Reg. Casp.
salsa, Claus, in Goebel, Reise, ii. 319 = Eversmanni.
salsa, *C. A. Mey. Verz. Pfl. Cauc.* 226.—Reg. Cauc.

TETRADIUM, Dulac, Fl. Hautes-Pyr. 321 (1867) = **Sedum**, Tourn. (Crassul.).
odoratum, Dulac, l. c. = S. roseum, *Scop.*

TETRADIUM, Lour. Fl. Cochinch. 91 (1790) = **Evodia**, Forst. (Rutac.).
amarissimum, Poir. Encyc. Suppl. v. 291 = Brucea sumatrana.
cymosum, Wall. ex Royle, Illustr. Bot. Himal. 157 = E. fraxinifolia.
fraxinifolium, Wall. ex Royle, l. c. = E. fraxinifolia.
trichotomum, Lour. Fl. Cochinch. 91 = E. fraxinifolia.

TETRADYMIA, DC. Prod. vi. 440 (1830). *COMPOSITAE*, Benth. & Hook. f. ii. 450.
LAGOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 416.
canescens, *DC. Prod.* vi. 440 ; *et in Deless. Ic. Sel.* iv. 27. t. 60.—Am. bor. occ.
comosa, *A. Gray, in Proc. Am. Acad.* xii. (1877) 60.—Am. bor. occ.
glabrata, *A. Gray, in Pacif. Rail. Rep.* ii. (1854) 122. t. 5.—Am. bor. occ.
inermis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 415 = canescens.
Nuttallii, *Torr. & Gray, Fl. N. Am.* ii. 447.—Am. bor. occ.
ramosissima, Torr. in Emory, Notes Mil. Reconnois. 145 = Psathyrotes ramosissima.
spinosa, *Hook. & Arn. Bot. Beech. Voy.* 360.—Am. bor. occ.
squamata, A. Gray, in Proc. Am. Acad. ix. (1874) 207 = Lepidospartum squamatum.
stenolepis, *Greene, in Bull. Calif. Acad.* i. III. (1885) 92.—Calif.

TETRAGAMESTUS, Reichb. f. in Bonplandia, ii. (1854) 21 = **Ponera**, Lindl. (Orchid.).
aureus, Reichb. f. l. c. 22 = Ornithidium ruberrimum.
isochiloides, Regel, Ind. Sem. Hort. Petrop. (1856) 20 ; et in Ann. Sc. Nat. Sér. IV. vi. (1856) 376.—N. Granat.
modestus, Reichb. f. in Bonplandia, ii. (1854) 21 = P. modesta.

TETRAGASTRIS, Gaertn. Fruct. ii. 130. t. 109 (1791) = **Hedwigia**, Sw. (Burser.).—Cf. Benth. & Hook. f. i. 326 ; iii. 318.
ossea, Gaertn. l. c.—Hab. ?

TETRAGLOCHIN, [Kunze, in] Poepp. Fragm. Syn. Pl. Chile, 26 (1833) = **Margyricarpus**, Ruiz & Pav. (Rosac.).
buxifolium, Presl, Epim. Bot. 200.—Chili.
caespitosum, Phil. in Linnaea, xxxiii. (1864–65) 63.—Chili.
microphyllum, Phil. l. c. xxviii. (1856) 687 = M. microphyllus.
strictum, [Kunze, in] Poepp. Fragm. Syn. Pl. Chile, 26 = M. alatus.
striosum, Walp. Rep. ii. 44, sphalm. = praec.

TETRAGLOSSA, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 70 = **Cleidion**, Blume (Euphorb.).
indica, Bedd. l. c. = C. javanicum ?

TETRAGONANTHUS, S. G. Gmel. Fl. Sibir. iv. 113 (1769) = **Halenia**, Borkh. (Gentian.).

TETRAGONELLA, Miq. in Lehm. Pl. Preiss. i. 245 (1844) = **Tetragonia**, Linn. (Ficoid.).
implexicoma, Miq. l. c. (*amplexicoma*) = Tetragonia implexicoma.

TETRAGONIA, Linn. Syst. ed. I (1735). *FICOIDEAE*, Benth. & Hook. f. i. 854.
DEMIDOVIA, Pall. Hort. Demid. 150. t. 1 (1781).
LUDOLFIA, Adans. Fam. ii. 244 (1763).

TETRAGONIA :—
TETRAGONELLA, Miq. in Lehm. Pl. Preiss. i. 245 (1844).
angustifolia, *Barn. in C. Gay, Fl. Chil.* ii. 470.—Chili.
Arbuscula, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 467.—Afr. austr.
calycina, *Fenzl, l. c.* 466.—Afr. austr.
chenopodioides, *Eckl. & Zeyh. Enum.* 324.—Afr. austr.
cornuta, Gaertn. Fruct. ii. 488 = expansa, *Murr.*
crystallina, *L'Hérit. Stirp. Nov.* 81. t. 39.—Peruv.
decumbens, *Mill. Gard. Dict.* ed. VIII. n. 2.—Afr. austr.
diptera, *F. Muell. Fragm.* xi. 8.—Austral.
distorta, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 467.—Afr. austr.
echinata, *Ait. Hort. Kew.* ed. I. ii. 177.—Afr. austr.
expansa, *Murr. in Comm. Gotting.* vi. (1783) 13.—Australas.
expansa, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
fruticosa, *Linn. Sp. Pl.* 480.—Afr. austr.
galenioides, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 463.—Afr. austr.
glauca, *Fenzl, l. c.* 468.—Afr. austr.
hdlimifolia, Forst. f. Prod. 39 = expansa, *Murr.*
halimoides, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 463.—Afr. austr.
Haworthii, *Fenzl, l. c.* 468.—Afr. austr.
herbacea, *Linn. Sp. Pl.* 480.—Afr. austr.
heterophylla, *Eckl. & Zeyh. Enum.* 324.—Afr. austr.
hirsuta, *Linn. f. Suppl.* 258.—Afr. austr.
implexicoma, *Hook. f. Fl. Tasm.* i. 148.—Austral.
inermis, F. Muell. in Linnaea, xxv. (1852) 384 = expansa, *Murr.*
ivaefolia, Linn. f. Suppl. 257 = Haloragis alata.
japonica, Thunb. Fl. Jap. 208 = expansa, *Thunb.*
lanceolata, *Burm. f. Fl. Cap. Prod.* 14.—Afr. austr.
linearis, *Haw. Rev. Pl. Succ.* 73.—Afr. austr.
macrocarpa, *Phil. Fl. Atac.* 19.—Chili.
maritima, *Barn. in C. Gay, Fl. Chil.* ii. 469.—Chili.
microcarpa, *Phil. Anal. Univ. Chil.* (1872) 717 ; *Fl. Atac.* 19.—Chili.
microptera, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 461.—Afr. austr.
nigrescens, *Eckl. & Zeyh. Enum.* 323.—Afr. austr.
obovata, *Haw. Rev. Pl. Succ.* 73.—Afr. austr.
pedunculata, *Phil. in Anal. Univ. Chil.* (1872) 717.—Chili.
pentandra, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 404.—Ins. Socotra.
peruviana, Hort. Chels. ex Haw. in Phil. Mag. (1827) 124 = Calandrinia phacosperma.
portulacoides, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 462.—Afr. austr.
psiloptera, *Fenzl, l. c.* 466.—Afr. austr.
purpurea, *Otto, ex Sweet, Hort. Brit.* ed. II. 234.—Hab. ?
pusilla, *Phil. in Anal. Univ. Chil.* (1872) 716.—Chili.
quadricornis, Stokes, Bot. Mat. Med. iii. 127 = expansa, *Murr.*
reduplicata, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 583.—Afr. trop.
robusta, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 467.—Afr. austr.
saligna, *Fenzl, l. c.* 469.—Afr. austr.
sarcophylla, *Fenzl, l. c.* 468.—Afr. austr.
spicata, *Linn. f. Suppl.* 258.—Afr. austr.
Tetrapteris, *Haw. Misc. Nat.* 121.—Afr. austr.
trigyna, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 77 ; *et Handb. N. Zeal. Fl.* 84.—N. Zel.
verrucosa, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 465.—Afr. austr.
villosa, Poir. Encyc. vii. 604 = hirsuta.
Zeyheri, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 465.—Afr. austr.

TETRAGONOCARPUS, Hassk. in Flora, xl. (1857) 99 = **Marsdenia**, R. Br. (Asclep.).
Teysmanni, Hassk. l. c. 100.—Malaya.

TETRAGONOLOBUS, Scop. Fl. Carn. ed. II. ii. 87 (1772) = **Lotus**, Tourn. (Legumin.).
aduncus, Griseb. Spicil. Fl. Rumel. i. 46 = L. aduncus.
aegaeus, Griseb. l. c. 45 = L. aegeus.
biflorus, Ser. in DC. Prod. ii. 215 = L. biflorus.

TETRAGONOLOBUS :—
Bouteloui, Willk. Iber. Halbinsel, 116 = Lotus siliquosus.
conjugatus, Link, Enum. Hort. Berol. ii. 264 = L. conjugatus.
edulis, Link, l. c. = L. Tetragonolobus.
glaucus, Dulac, Fl. Hautes-Pyr. 284 = L. siliquosus.
guttatus, Pomel, Nouv. Mat. Fl. Atl. 182 = L. conjugatus.
maritimus, Roth, Tent. Fl. Germ. i. 323 = L. siliquosus.
monanthos, Sacc. & Penzig, in Bot. Centralbl. v. (1881) 285.—Hab. ?
palaestinus, Boiss. Diagn. Ser. II. ii. 20.—As. Min. ; Syria.
prostratus, Moench, Meth. 164 = L. siliquosus.
purpureus, Moench, l. c. = L. Tetragonolobus.
Requienii, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 23 = L. Requieni.
Scandalida, Bernh. Syst. Verz. Erf. 243 = L. uliginosus.
siliquosus, Roth, Tent. Fl. Germ. i. 323 = L. siliquosus.
simplicifolius, Blanco, Fl. Filip. ed. II. 397 = Alysicarpus tetragonolobus.
tauricus, Bunge, ex Nym. Consp. 182 = L. siliquosus.
Wiedemanni, Boiss. Fl. Orient. ii. 176 = L. Wiedemanni.

TETRAGONOSPERMA, Scheele, in Linnaea, xxii. (1849) 166 = **Tetragonotheca**, Dill. (Compos.).
lyratifolium, Scheele, l. c. 167 = Tetragonotheca texana.

TETRAGONOTHECA, Dill. ex Linn. Gen. ed. I. 259 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 367.
BIKERA, Adans. Fam. ii. 130 (1763).
GONOTHECA, Rafin. in Med. Repos. N. York, v. (1808) 352.
HALEA, Torr. & Gray, Fl. N. Am. ii. 304 (1840).
TETRAGONOSPERMA, Scheele, in Linnaea, xxii. (1849) 166.
abyssinica, Ledeb. Ind. Sem. Hort. Dorpat. Suppl. (1824) 7 = Guizotia abyssinica.
helianthoides, *Linn. Sp. Pl.* 903.—Am. bor.
ludoviciana, *A. Gray, in E. Hall, List. Pl. Tex.* 13.—Am. bor.
parviflora, *Jacq. Enum. Pl. Carib.* 28.—Ind. occ.
texana, *A. Gray & Engelm. in Proc. Am. Acad.* i. (1848) 48.—Texas ; Mexic.

TETRAGYNE, Miq. Fl. Ind. Bat. Suppl. 463 (1860). Genus dubium *EUPHORBIACEARUM*.—Cf. Benth. & Hook. f. iii. 257.
acuminata, *Miq. l. c.* 464.—Sumatra.

TETRAHIT, Adans. Fam. ii. 190 (1763) = **Galeopsis**, Linn. (Labiat.).
grandiflorum, Gilib. Fl. Lituan. i. 83 = G. Tetrahit.
Ladanum, Moench, Meth. 394 = G. Ladanum.
leucanthum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135 = G. Tetrahit.
longiflorum, Moench, Meth. 395 = G. ochroleuca.
Navieri, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135 = G. Tetrahit.
nodosum, Moench, Meth. 395 = G. Tetrahit.
praecox, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135 = G. Tetrahit.
purpurascens, Gilib. Fl. Lituan. i. 83 = G. Tetrahit.
Reichenbachii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135 = G. Tetrahit.
sulfureum, Fourr. l. c. = G. versicolor.

TETRAHITUM, Hoffmgg. & Link, Fl. Port. i. 103 (1809) = **Stachys**, Tourn. (Labiat.).
hirtum, Hoffmgg. & Link, l. c. 104 = S. hirta.

TETRALEPIS, Steud. Syn. Pl. Cyp. 159 (1855) = **Cyathochaeta**, Nees (Cyperac.).
australis, Steud. l. c. = C. avenacea.

TETRALIX, Griseb. Cat. Pl. Cub. 8 (1866). Genus dubium *BIXINEARUM*.—Cf. Benth. & Hook. f. i. 971.
brachypetalus, *Griseb. l. c.* 9.—Cuba.

TETRALIX, Hill, Veg. Syst. iv. 18 (1762) = **Cnicus**, Linn. (Compos.).
eriophora, Hill, Herb. Brit. i. 75 = C. eriophorus.

TETRALOBUS, A. DC. in DC. Prod. viii. 667 (1844) = **Polypompholyx**, Lehm. (Lentib.).
Preissii, A. DC. l. c. = P. multifida.
pusillus, A. DC. l. c. = P. tenella.

TETRALÓPHA, Hook. f. in Hook. Ic. Pl. t. 1072 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 120.
Motleyi, *Hook. f. in Hook. Ic. Pl.* t. 1072.—Borneo.

TETRALYX, Hill, Hort. Kew. 61 (1768) = TETRALIX, Hill = **Cnicus**, Linn. (Compos.).

TETRAMELES, R. Br. in Denh. & Clapp. Trav. II. 230 (App. 25) (1826). *DATISCEAE*, Benth. & Hook. f. i. 845.
ANICTOCLEA, Nimmo, in J. Grah. Cat. Bomb. Pl. 252 (1835).
Grahamiana, Wight, Ic. t. 1956 = nudiflora.
Horsfieldii, Steud. Nom. ed. II. ii. 671 = nudiflora.
nudiflora, *R. Br. in Benn. Pl. Jav. Rar.* 79. t. 17.—Ind. or.; Malaya.
rufinervis, Miq. Fl. Ind. Bat. i. 1. 726; Pl. Jungh. 401 = nudiflora.

TETRAMERIS, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 120. t. 4 = **Comolia**, DC. (Melastom.).
Aubletii, Naud. l. c. 127 = C. Aubletii.
brachyandra, Naud. l. c. = Pterolepis glomerata, Tibouchina versicolor.
cerastifolia, Naud. l. c. 122 = Tibouchina cerastifolia.
excoriata, Naud. l. c. 125 = C. violacea.
isanthera, Naud. l. c. 123 = Tibouchina versicolor.
Martiana, Naud. l. c. 121 = Tibouchina sebastianopolitana.
nitida, Naud. l. c. 127 = Tibouchina nitida.
repanda, Naud. l. c. = Pterolepis repanda.
rotundifolia, Naud. l. c. 124 = C. denudata.
serrulata, Naud. l. c. 123 = Tibouchina nitida.
Sertularia, Naud. l. c. 126 = C. Sertularia.
simplicicaulis, Naud. l. c. 122 = Tibouchina simplicicaulis.
stenodon, Naud. l. c. 126 = C. stenodon.
trivalvis, Naud. l. c. 121. t. 4 = Macairea multinervia.
uruguayensis, Naud. l. c. 127 = Tibouchina nitida.
versicolor, Naud. l. c. = Tibouchina versicolor.
villosa, Naud. l. c. 127 = C. villosa.
violacea, Naud. l. c. 124 = C. lanceaeflora.
viscidula, Naud. l. c. 125 = C. sessilis.

TETRAMERISTA, Miq. Fl. Ind. Bat. Suppl. 534 (1860). *OCHNACEAE*, Benth. & Hook. f. i. 318.
glabra, *Miq. l. c.*—Malaya.
paniculata, *Kurz, in Journ. Bot.* xiii. (1875) 333.—Malaya.

TETRAMERIUM, Gaertn. f. Fruct. iii. 90. t. 196 (1805) = **Faramea**, Aubl. (Rubiac.).
coeruleum, Nees, ex Mart. in Nov. Act. Nat. Cur. xii. 1. (1824) 13 = F. coerulea.
jasminoides, H. B. & K. Nov. Gen. et Sp. iii. 373. t. 287 = F. jasminoides.
latifolium, Cham. & Schlecht. in Linnaea, iv. (1829) 30 = F. latifolia.
montevidense, Cham. & Schlecht. l. c. 29 = F. montevidensis.
multiflorum, Bartl. ex DC. Prod. iv. 497 = F. guayaquilensis.
occidentale, Nees, ex Mart. in Nov. Act. Nat. Cur. xii. 1. (1824) 13 = F. odoratissima.
odoratissimum, Gaertn. f. Fruct. iii. 90. t. 196 = F. odoratissima.
oleaefolium, Spreng. Syst. i. 409 = Coffea oleifolia.
paniculatum, Spreng. l. c. = Coffea paniculata.
sessilifolium, H. B. & K. Nov. Gen. et Sp. iii. 374 = F. sessilifolia.
stipulaceum, Cham. & Schlecht. in Linnaea, iv. (1829) 31 = F. stipulacea.

TETRAMERIUM, Nees, in Benth. Bot. Voy. Sulph. 147. t. 48 (1844). *ACANTHACEAE*, Benth. & Hook. f. ii. 1121.
HENRYA, Nees, in Benth. Bot. Voy. Sulph. 148. t. 49 (1844).
glandulosum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 171.—Mexic.
hispidum, *Nees, in DC. Prod.* xi. 468.—Mexic.
nervosum, *Nees, in Benth. Bot. Voy. Sulph.* 148. t. 48.—Am. centr.
ovalifolium, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 170.—Mexic.
ovatum, *Oerst. l. c.* 169.—Mexic.
platystegium, *Torr. in Bot. Mex. Bound.* 126.—Am. bor.

TETRAMERIUM:—
polystachyum, *Nees, in Benth. Bot. Voy. Sulph.* 147.—Am. austr.
racemulosum, *Nees, l. c.* 148, in obs.; *et in DC. Prod.* xi. 468.—Ind. occ.
scorpioides, *Hemsl. Biol. Centr. Am. Bot.* ii. 526.—Am. centr.

TETRAMICRA, Lindl. Gen. et Sp. Orch. 119 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 532.
LEPTOTES, Lindl. Bot. Reg. t. 1625 (1833).
bicolor, *Rolfe, in Gard. Chron.* (1883) II. 20.—Bras.
Eulophiae, *Reichb. f. ex Griseb. Cat. Pl. Cub.* 264.—Cuba.
montana, Griseb. Fl. Brit. W. Ind. 622 = Octadesmia serratifolia.
parviflora, *Lindl. ex Griseb. l. c.*—Jamaic.
platyphylla, Griseb. Cat. Pl. Cub. 264 = Bletia sarcophylla.
rigida, *Lindl. Gen. et Sp. Orch.* 119.—Ind. occ.
schoenina, *Benth. & Hook. f. Gen.* iii. 533.—Ins. S. Doming.
tenera, Griseb. ex Benth. in Journ. Linn. Soc. xviii. (1881) 314 = Eulophiae.

TETRAMOLOPIUM, Nees, Gen. et Sp. Aster. 202 (1833), partim. *COMPOSITAE*, Benth. & Hook. f. ii. 282.
ochraceum, *DC. Prod.* v. 262.—Ecuador.
phylicoides, *DC. l. c.*—Mexic.
rupestre, Nees, Gen. et Sp. Aster. 203 = Diplostephium rupestre.
tenerrimum, *Nees, l. c.* 203.—Ins. Sandvic.

TETRAMORPHAEA, DC. in Guill. Arch. Bot. ii. (1833) 331 = **Centaurea**, Linn. (Compos.).
Belangeriana, DC. Prod. vi. 609 = C. phyllocephala.
Bruguieriana, DC. l. c. = C. phyllocephala.

TETRANDRA, Miq. Fl. Ind. Bat. ii. 929 (1859) = **Tournefortia**, Linn. (Borag.).
glabra, Miq. l. c. = Tourn. Wallichii.
Wallichii, Miq. l. c. 928 = Tourn. Wallichii.
Zollingeri, Miq. l. c. = Tourn. Wallichii.

TETRANEMA, Benth. in Bot. Reg. (1843) t. 52. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 942.
mexicanum, *Benth. in Bot. Reg.* (1843) t. 52.—Mexic.

TETRANEMA, Sweet, Hort. Brit. ed. II. 149 (1830) = **Desmodium**, Desv. (Legumin.).
nutans, Sweet, l. c. = D. tiliaefolium.

TETRANTHA, Poit. ex DC. Prod. v. 696, 503 (1836) = **Riencourtia**, Cass. (Compos.).
suaveolens, Poit. ex DC. l. c. 504 = R. glomerata.

TETRANTHERA, Jacq. Hort. Schoenb. i. 59. t. 113 (1797) = **Litsea**, Lam. (Laurin.).
accedens, Blume, Mus. Bot. Lugd. Bat. i. 383 (= *Cylicodaphne accedens*).—Borneo.
acuminatissima, Blume, l. c. 381 = L. amara.
aestivalis, Spreng. Syst. ii. 267 = L. geniculata.
albicans, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 102.—Burma.
albida, Spreng. Syst. ii. 267 = Sassafras officinale.
alnoides, Miq. Pl. Jungh. 180 = L. polyantha.
amara, Nees, Syst. Laurin. 551 = L. amara.
ambigua, Blume, Mus. Bot. Lugd. Bat. i. 377.—Ins. Molucc.
ambrosa, Wall. Cat. n. 2564, sphalm. = L. umbrosa.
angulata, Nees, Syst. Laurin. 544 = L. angulata.
angusta, Wall. Cat. n. 6344 = L. amara.
angustifolia, Wall. l. c. n. 2537.—Ind. or.
angustifolia, Wall. l. c. n. 2559 = Actinodaphne angustifolia.
angustifolia, Zoll. ex Meissn. in DC. Prod. xv. 1. 182 = L. citrata.
apetala, Dalz. & Gibs. Bomb. Fl. 222 = L. tomentosa.
apetala, Herb. Mus. Par. ex Blume, Mus. Bot. Lugd. Bat. i. 382 = L. timoriana.
apetala, Roxb. Pl. Corom. ii. 25 = L. Sebifera.
areolata, Blume, Mus. Bot. Lugd. Bat. i. 381.—N. Guin.
attenuata, Wall. Cat. n. 2534 = *salicifolia*, Roxb.
bancana, Miq. Fl. Ind. Bat. i. 1. 950.—Ins. Banca.
bandongana, Miq. in Zoll. Syst. Verz. Ind. Archip. 114, 115 = L. Noronhae.

TETRANTHERA:—
Baula, Buch.-Ham. ex Nees, in Wall. Cat. n. 2540 B; et Pl. As. Rar. ii. 67 (= *Cylicodaphne nitida*).—Ind. or.
Berterii, Spreng. Syst. ii. 267.—Am. austr.
bifaria, Wall. Cat. n. 2530 = Lindera heterophylla.
bifurca, Steud. Nom. ed. II. ii. 671, sphalm. = praec.
Bindoniana, F. Muell. Fragm. v. 167.—Austral.
Blumei, Miq. Pl. Jungh. 178 = seq.
Blumii, Nees, in Wall. Pl. As. Rar. ii. 65 = Litsea ferruginea.
borneensis, Meissn. in DC. Prod. xv. 1. 184.—Borneo.
brachystachya, Blume, Mus. Bot. Lugd. Bat. i. 384.—Java.
Brawas, Blume, l. c.—Sumatra.
caduca, Buch.-Ham. ex Wall. Cat. n. 2559 = Actinodaphne angustifolia.
calicaris, Hook. f. Fl. N. Zel. i. 216 = *T. Tangao*.
californica, Hook. & Arn. Bot. Beech. Voy. 159 = Umbellularia californica.
calononeura, Miq. Fl. Ind. Bat. i. 1. 946.—Java.
calophylla, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 102 = *martabanica*.
calophylla, Miq. Pl. Jungh. 183 = L. Tetranthera.
calycaris, Hook. f. ex Meissn. in DC. Prod. xv. 1. 191 = *calicaris*.
camphoracea, Wall. ex Meissn. l. c. 25 = Cinnamomum inunctum.
capitata, Roxb. ex Nees, in Wall. Pl. As. Rar. ii. 65 = L. Sebifera.
capitulata, Miq. Pl. Jungh. 182 = L. amara.
caudata, Wall. Cat. n. 2566 (= *Daphnidium caudatum*).—Ind. or.
cauliflora, Moon, Cat. Pl. Ceyl. 69 = L. longifolia.
causticans, Pasq. in Nuov. Giorn. Bot. Ital. i. (1869) 290 = Umbellularia californica?
celastroides, Miq. ex Meissn. in DC. Prod. xv. 1. 180 = L. ligustrina.
Cervantesii, Nees, Syst. Laurin. 553 = L. glaucescens.
chartacea, Meissn. in DC. Prod. xv. 1. 186, partim.—Reg. Himal.
chartacea, Wall. Cat. n. 2531.—Nepal.
chrysantha, Blume, Mus. Bot. Lugd. Bat. i. 376 = L. minuta.
chrysophaena, Blume, l. c. 377.—Ins. Molucc.
chrysopleura, Blume, l. c. 383.—Java.
citrata, Nees, Syst. Laurin. 560 = L. citrata.
citrifolia, Spreng. Syst. ii. 266 = L. Sebifera.
cordata, Jack, in Malay. Misc. ii. (1822) n. VII. 34.—Sumatra.
cordifolia, Meissn. in DC. Prod. xv. 1. 196, sphalm. = *cordata*.
coriacea, Heyne, ex Wall. Cat. n. 2556.—Ins. Penang.
crassifolia, Blume, Mus. Bot. Lugd. Bat. i. 386.—Borneo.
Cubeba, Meissn. in DC. Prod. xv. 1. 199 = L. Cubeba.
Cuipala, D. Don, Prod. Fl. Nep. 65 = L. lanuginosa.
cuneata, Blume, Mus. Bot. Lugd. Bat. i. 375.—Ins. Celebes.
cuspidata, Blume, l. c. 383.—Java.
Daradmeda, Buch.-Ham. ex Wall. Cat. n. 2554 E = L. Sebifera.
dealbata, R. Br. Prod. 403 = L. dealbata.
Dengek, Miq. Pl. Jungh. 178 = L. fulva.
Diepenhorstii, Miq. Fl. Ind. Bat. Suppl. 363.—Sumatra.
Diglatia, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. ii. 66 = *salicifolia*, Roxb.
Diospyros, Spreng. Syst. ii. 267 = Lindera melissaefolia.
discolor, Blume, Mus. Bot. Lugd. Bat. i. 382 = L. timoriana.
diversifolia, Hassk. Cat. Hort. Bog. Alt. 90 = L. diversifolia.
diversifolia, Zoll. Syst. Verz. Ind. Arch. 114 (= *Cylicodaphne sumatrana*).—Malaya.
Doshia, D. Don, Prod. Fl. Nep. 65 = L. oblonga.
elliptica, Nees, Syst. Laurin. 550 = L. elliptica.
elongata, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. 1. 181.—Ind. or.
elongata, Wall. Cat. n. 2546 = L. elongata.
fagifolia, Wall. l. c. n. 2539 = Beilschmiedia fagifolia.
faginea, Miq. Pl. Jungh. 181 = L. amara.
Fawcettiana, F. Muell. Fragm. v. 168 = *reticulata*, Meissn.
ferruginea, R. Br. Prod. 403 = L. hexantha.
ferruginea, F. Muell. ex Meissn. in DC. Prod. xv. 1. 73 = Endiandra pubens.
firma, Blume, Mus. Bot. Lugd. Bat. i. 381.—Borneo.

TETRANTHERA :—
flexuosa, Blume, Mus. Bot. Lugd. Bat. i. 382.—Ins. Molucc.
floribunda, Champ. ex Benth. in Hook. Kew Journ. v. (1853) 199 = Litsea citrata.
floridana, Shuttl. ex Meissn. in DC. Prod. xv. 1. 181 = L. geniculata.
foliosa, Wall. Cat. n. 2563 = L. zeylanica.
Forstenii, Blume, Mus. Bot. Lugd. Bat. i. 383.—Ins. Celebes.
fruticosa, Buch.-Ham. ex Meissn. in DC. Prod. xv. 1. 189 = L. polyantha.
fruticosa, Nees, Syst. Laurin. 543 = L. amara.
fruticosa, Roxb. Hort. Beng. 73; Fl. Ind. iii. 823.—Ind. or.
fruticosa, Span. in Linnaea, xv. (1841) 346, nomen = *undulata*.
fulva, Blume, Mus. Bot. Lugd. Bat. i. 377 = L. fulva.
furfuracea, Wall. Cat. n. 2568 = L. zeylanica.
Gardneri, Thw. Enum. Pl. Zeyl. 255 = L. Gardneri.
gemelliflora, Miq. Pl. Jungh. 179.—Sumatra.
geminata, Nees, Syst. Laurin. 559 = L. Sebifera.
geniculata, Nees, l. c. 567 = L. geniculata.
glaberrima, Thw. Enum. Pl. Zeyl. 255 = L. glaberrima.
glabra, Roxb. Hort. Beng. 73, nomen.—Ind. or.
glabraria, Nees, Syst. Laurin. 569 = L. Sebifera.
glabrata, Wall. Cat. n. 2543.—Ind. or.
glauca, Wall. l. c. n. 2533 = *salicifolia*, Roxb.
glaucescens, Spreng. Syst. ii. 267 = L. glaucescens.
glaucescens, Wall. Cat. n. 2560 = Cinnamomum Cecidodaphne.
grandiflora, Wall. l. c. n. 2544 = Dodecadinia grandiflora.
grandis, Wall. l. c. n. 2552.—Ins. Penang.
Griffithii, Meissn. in DC. Prod. xv. 1. 191.—Malacca.
Gullavara, Buch.-Ham. ex Wall. Cat. n. 2593 = Actinodaphne angulifolia.
hexantha, Sieber, ex Meissn. in DC. Prod. xv. 1. 189 = L. polyantha.
Hirana, Buch.-Ham. ex Wall. Cat. n. 2554 F = L. polyantha.
hispidula, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 374 = L. Sebifera.
Hookeriana, Meissn. in DC. Prod. xv. 1. 188.—Zeylan.
hypoglauca, Miq. Pl. Jungh. 182 = Actinodaphne glomerata.
impressa, Blume, Mus. Bot. Lugd. Bat. i. 378.—Java.
insignis, Blume, l. c. 382.—Sumatra.
intermedia, Blume, l. c. 383.—Java.
itiodaphne, Nees, Syst. Laurin. 542 = L. ovalifolia.
japonica, Spreng. Syst. ii. 266 = L. japonica.
japonica, Wight, ex Wall. Cat. n. 2550 B = L. tomentosa.
khasyana, Meissn. in DC. Prod. xv. 1. 185.—Reg. Himal.
laeta, Wall. Cat. n. 2541 = L. laeta.
lanceaefolia, Roxb. Hort. Beng. 73; Fl. Ind. iii. 822 = *salicifolia*, Roxb.
lancifolia, Roxb. ex Wall. Cat. n. 2532 = L. minuta.
lanuginosa, Wall. l. c. n. 2561 = L. lanuginosa.
laurifolia, Jacq. Hort. Schoenb. i. 59 = L. Sebifera.
laurifolia, Roxb. Hort. Beng. 73; Fl. Ind. iii. 823 = *salicifolia*, Roxb.
leiantha, Kurz, For. Fl. Brit. Burma, ii. 300.—Ins. Andam.
Lhuysii, Carr. in Rev. Hortic. (1869) 368.—Japon.
ligustrina, Nees, in Wall. Pl. As. Rar. ii. 65 = L. ligustrina.
ligustrina, Thw. Enum. Pl. Zeyl. 254.—Zeylan.
litoralis, Blume, Mus. Bot. Lugd. Bat. i. 375 = L. Sebifera.
longifolia, Nees, Syst. Laurin. 528 = L. longifolia.
longifolia, Thw. Enum. Pl. Zeyl. 255 = L. laeta, glaberrima.
lucida, Hassk. Cat. Hort. Bog. Alt. 90.—Java.
lusona, Nees, ex Meissn. in DC. Prod. xv. 1. 200 = Lindera heterophylla.
luzonica, Blume, Mus. Bot. Lugd. Bat. i. 377 = L. luzonica.
macrantha, Wall. Cat. n. 2587 = L. Sebifera.
macrocalyx, Hance, in Journ. Bot. xv. (1877) 336.—China.
macrocarpa, Nees, Syst. Laurin. 565 (= *Nothaphoebe cuneata*).—Malaya.
macrophylla, Roxb. Hort. Beng. 73; Fl. Ind. iii. 822.—Ind. or.
macrophylla, Wall. Cat. n. 2549 = L. polyantha.
mappacea, Blume, Mus. Bot. Lugd. Bat. i. 381.—Sumatra.

TETRANTHERA :—
martabanica, Kurz, For. Fl. Brit. Burma, ii. 301.—Burma.
megapotamica, Spreng. Syst. iv. Cur. Post. 156.—Bras.
melastomacea, Wall. Cat. n. 2565 = Lindera melastomacea.
Merawas, Miq. Fl. Ind. Bat. Suppl. 362.—Sumatra.
micrantha, Meissn. in DC. Prod. xv. 1. 83 = *coriacea*.
Mogal, Blume, Mus. Bot. Lugd. Bat. i. 383.—Java.
mollis, Blume, l. c. 372.—Java.
mollis, Miq. Pl. Jungh. 179.—Java?
monopetala, Roxb. Pl. Corom. ii. 26 = Litsea polyantha.
multiflora, Nees, Syst. Laurin. 544 = L. Sebifera.
multiflora, Zoll. Syst. Verz. Ind. Arch. 114 = grandis.
myristicaefolia, Wall. Cat. n. 2548.—Burma.
Myrrha, Kostel. Allg. Med.-Pharm. Flora, ii. 479 = L. trinervia?
Neesiana, Schau. in Linnaea, xix. (1847) 712 = L. Neesiana.
Neesiana, Wall. Cat. n. 2558 = Lindera Neesiana.
nemoralis, Thw. Enum. Pl. Zeyl. 255 = L. nemoralis.
nervosa, Meissn. in DC. Prod. xv. 1. 187 = L. glaberrima.
nigricans, Meissn. l. c. 190.—Borneo.
nitens, Blume, Mus. Bot. Lugd. Bat. i. 375.—Java.
nitida, Roxb. Hort. Beng. 73; Fl. Ind. iii. 818 (= *Cylicodaphne nitida*).—Ind. or.
nuculanea, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 102.—Burma.
oblonga, Wall. Cat. n. 2542 = L. oblonga.
obovata, Wall. l. c. n. 2562 = Actinodaphne obovata.
obscura, Blume, Mus. Bot. Lugd. Bat. i. 386.—N. Guin.
obtusa, Nees, Syst. Laurin. 564 = L. obtusa.
ochrascens, Miq. Fl. Ind. Bat. Suppl. 363 = L. rugosa.
oleoides, Meissn. in DC. Prod. xv. 1. 195.—Ind. or.
oppositifolia, Miq. Pl. Jungh. 178 = L. fulva.
ovalifolia, Thw. Enum. Pl. Zeyl. 256 = L. ovalifolia.
oxyphylla, Wall. Cat. n. 2547 (= *Aperula oxyphylla*).—Ins. Penang.
pallens, D. Don, Prod. Fl. Nep. 68 = L. umbrosa.
pallida, Blume, Mus. Bot. Lugd. Bat. i. 377.—Celebes.
palmatinervia, Meissn. in DC. Prod. xv. 1. 191.—Ins. Fiji.
Panamanja, Nees, in Wall. Pl. As. Rar. ii. 67 = seq.
Panamonja, Buch.-Ham. ex Wall. Cat. n. 2553.—Ind. or.
Panshia, Herb. Buch.-Ham ex Wall. l. c. n. 2554 D = L. Sebifera.
Pantjara, Blume, Mus. Bot. Lugd. Bat. i. 385.—Java; Sumatra.
pentandra, Roxb. Fl. Ind. iii. 824 = L. zeylanica.
Perrottetii, Blume, Mus. Bot. Lugd. Bat. i. 384 = L. Perrottetii.
Pichurim, Spreng. in Berl. Jahrb. (1821) t. 3; ex Pritz. Ic. Ind. i. 1093 (Quid?).—Hab.?
Pickeringii, A. Gray, ex Seem. Fl. Vit. 203.—Ins. Fiji.
picta, Carr. in Rev. Hortic. (1869) 367.—Japon.
pilosa, Spreng. Syst. ii. 267 = Machilus pilosa.
platyphylla, Nees, Syst. Laurin. 570 = L. Sebifera.
polyantha, Wall. Cat. n. 2538 = L. citrata.
polycephala, Wall. ex Meissn. in DC. Prod. xv. 1. 179 = L. Sebifera.
polyneura, Hort. Bog. ex Miq. Fl. Ind. Bat. i. 1. 939 (= *Lepidadenia polyneura*).—Java?
propinqua, Blume, Mus. Bot. Lugd. Bat. i. 385.—Sumatra.
prunifolia, Wall. ex Meissn. in DC. Prod. xv. 1. 241 = Lindera Neesiana.
Pseudo-angulata, Blume, Mus. Bot. Lugd. Bat. i. 385 = L. Noronhae.
pulcherrima, Wall. Cat. n. 2567 = L. zeylanica, umbrosa.
pusilla, Blume, Mus. Bot. Lugd. Bat. i. 376 = L. minuta.
quadriflora, Roxb. Hort. Beng. 73; Fl. Ind. iii. 821.—Ind. or.
racemosa, Spreng. ex Nees, Syst. Laurin. 394 = Ocotea tristis.
rangoonensis, Meissn. in DC. Prod. xv. 1. 187.—Burma.
resinosa, Nees, Syst. Laurin. 550 = L. resinosa.
reticulata, Buch.-Ham. ex Wall. Cat. n. 2551 = Lindera reticulata.

TETRANTHERA :—
reticulata, Meissn. in DC. Prod. xv. 1. 192.—Austral.
riparia, Blume, Mus. Bot. Lugd. Bat. i. 382.—Ins. Molucc.
robusta, Nees, Syst. Laurin. 564 = Litsea robusta.
rotundifolia, Nees, ex Meissn. in DC. Prod. xv. 1. 200 = Actinodaphne chinensis.
Roxburghii, Nees, in Wall. Pl. As. Rar. ii. 65 = L. Sebifera.
rubra, Miq. ex Meissn. in DC. Prod. xv. 1. 202 = L. Noronhae.
rubra, Nees, Syst. Laurin. 564 = L. rubra.
Rumphii, Blume, Mus. Bot. Lugd. Bat. i. 382.—Ins. Molucc.
salicifolia, Roxb. ex Wall. Cat. n. 2536.—Ind. or.
salicifolia, Zoll. ex Meissn. in DC. Prod. xv. 1. 179 = L. Sebifera.
saligna, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. 1. 185 = *salicifolia*, Roxb.
saligna, Nees, in Wall. Pl. As. Rar. ii. 67 = *angustifolia*.
Sebifera, Spreng. Syst. ii. 267 = L. Sebifera.
Seemanni, Meissn. in DC. Prod. xv. 1. 192.—Ins. Fiji.
semecarpifolia, Wall. Cat. n. 6345 A = L. polyantha.
sericea, Wall. l. c. 2545.—Reg. Himal.
sikkimensis, Meissn. in DC. Prod. xv. 1. 181 = L. elongata.
subovata, Miq. Fl. Ind. Bat. Suppl. 362.—Sumatra.
Tangao, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 380.—N. Zel.
tersa, Spreng. Syst. ii. 267 = L. Sebifera.
Thomsonii, Meissn. in DC. Prod. xv. 1. 183.—Ind. or.
timoriana, Span. in Linnaea, xv. (1841) 346, nomen = L. timoriana.
tomentosa, Nees, in Wall. Pl. As. Rar. ii. 65 = L. Sebifera.
tomentosa, Roxb. ex Wall. Cat. n. 2550 A, B = L. tomentosa, *Heyne*.
trinervia, Spreng. Syst. ii. 266 = L. zeylanica, trinervia?
umbrosa, Wall. Cat. n. 2564 (*ambrosa*) = L. umbrosa.
undulata, Zipp. ex Span. in Linnaea, xv. (1841) 346, nomen.—Ins. Timor.
utilis, Meissn. in DC. Prod. xv. 1. 183.—Borneo.
varians, Blume, Mus. Bot. Lugd. Bat. i. 376.—Sumatra.
velutina, Blume, l. c.—Ind. or.
venulosa, Meissn. in DC. Prod. xv. 1. 187.—Ind. or.
verticillata, Buch.-Ham. ex Meissn. in DC. l. c. 189 = L. polyantha.
verticillata, Wall. ex Meissn. in DC. l. c. 214 = Actinodaphne angustifolia.
vestita, Nees, Syst. Laurin. 551 = L. tomentosa, *Blume*.
vestita, Wall. Cat. sub n. 2530 = *bifaria*?
villosa, Boj. Hort. Maurit. 272 = L. polyantha.
villosa, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 359 = L. Neesiana.
villosa, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 350 = L. villosa.
virens, Nees, Syst. Laurin. 536.—Ins. Penang.
vitiana, Meissn. in DC. Prod. xv. 1. 514.—Ins. Fiji.
Waitziana, Blume, Mus. Bot. Lugd. Bat. i. 384.—Java.
Wightiana, Wall. Cat. n. 2557 = L. Wightiana.
Wightii, Nees, Syst. Laurin. 523.—Ind. or.
xanthophylla, Blume, Mus. Bot. Lugd. Bat. i. 381.—Sumatra.
Zollingeriana, Miq. in Zoll. Syst. Verz. Ind. Archip. 114, 115 = *intermedia*.

TETRANTHUS, Sw. Prod. Veg. Ind. Occ. 116 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 343.
Berterii, Spreng. Syst. iii. 459 = Pinillosia tetranthoidea.
hirsutus, *Spreng. Neue Entdeck.* iii. 40.—Ins. S. Doming.
littoralis, Spreng. Syst. iii. 459 = hirsutus.
littoralis, *Sw. Prod. Veg. Ind. Occ.* 116.—Ins. S. Doming.
ruderalis, Griseb. in Goett. Abh. vii. (1857) 234 = Eleutheranthera ovata.
thymoides, Griseb. Cat. Pl. Cub. 286 = Thymopsis Wrightii.

TETRAOTIS, Reinw. in Blume, Bijdr. 892 (1826) = **Enhydra**, Lour. (Compos.).
longifolia, Reinw. l. c. = E. paludosa.
paludosa, Reinw. l. c. = E. paludosa.

TETRAPANAX, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 371 = **Fatsia**, Decne. (Araliac.).
innovans, C. Koch, l. c. = Acanthopanax innovans.
papyriferum, C. Koch, l. c. = F. papyrifera.
ricinifolium, C. Koch, l. c. = Acanthopanax ricinifolium.

TETRAPASMA, G. Don, Gen. Syst. ii. 40 (1832) = **Discaria**, Hook. (Rhamn.).
juncea, G. Don, l. c. = D. australis.

TETRAPATHAEA, Reichb. Consp. 182 (1828) = **Passiflora**, Linn.
australis, Raoul, in Ann. Sc. Nat. Sér. III. ii. (1844) 123 = P. tetrandra.

TETRAPELTIS, Wall. ex Lindl. Gen. et Sp. Orch. 212 (1832) = **Otochilus**, Lindl. (Orchid.).
fragrans, Wall. ex Lindl. l. c. 213 = O. porrecta.

TETRAPETALUM, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 1 (1865–66). *ANONACEAE*, Benth. & Hook. f. i. 955.
volubile, *Miq. l. c.*—Borneo.

TETRAPHYLAX, De Vriese, in Nat. Verh. Haarl. x. (1854) 164 = **Goodenia**, Sm.
quadrilocularis, De Vriese, l. c. 165 = G. quadrilocularis.

TETRAPHYLE, Eckl. & Zeyh. Enum. 292 (1837) = **Crassula**, Dill.
campestris, Eckl. & Zeyh. l. c. 294 = Tillaea pharnaceoides.
corallina, Eckl. & Zeyh. l. c. 293 = C. corallina.
furcata, Eckl. & Zeyh. l. c. = C. furcata.
lanceolata, Eckl. & Zeyh. l. c. 294 = C. lanceolata.
littoralis, Eckl. & Zeyh. l. c. 293 = C. littoralis.
lycopodioides, Eckl. & Zeyh. l. c. 294 = C. lycopodioides.
muscosa, Eckl. & Zeyh. l. c. = C. muscosa.
parvula, Eckl. & Zeyh. l. c. = C. parvula.
polpodacea, Eckl. & Zeyh. l. c. 293 = C. polpodacea.
propinqua, Eckl. & Zeyh. l. c. = C. propinqua.
pyramidalis, Eckl. & Zeyh. l. c. 292 = C. pyramidalis.
quadrangula, Eckl. & Zeyh. l. c. 293 = C. quadrangula.

TETRAPHYLLUM, Griff. Notul. iv. 148 (1854). *CYRTANDREAE.*
bengalense, *C. B. Clarke, in DC. Monog. Phan.* v. 137.—Ind. or.

TETRAPILUS, Lour. Fl. Cochinch. 611 (1790) = **Olea**, Tourn.
axillaris, Rafin. Sylva Tellur. 138.—Hab.?
brachiatus, Lour. Fl. Cochinch. 611 = O. maritima.

TETRAPLACUS, Radlk. in Sitzb. Akad. Muench. xv. (1885) 261. *SCROPHULARINEAE.*
platychilus, *Radlk. l. c.* 262.—Bras.

TETRAPLANDRA, Baill. in Ann. Sc. Nat. Sér. IV. ix. (1858) 200 = **Algernonia**, Baill. (Euphorb.).
Leandri, Baill. l. c. 202.—Bras.
Riedelii, Muell. Arg. in Mart. Fl. Bras. xi. II. 534.—Bras.

TETRAPLASANDRA, A. Gray, Bot. U. St. Expl. Exped. i. 727. t. 94 (1854). *ARALIACEAE*, Benth. & Hook. f. i. 947.
hawaiensis, *A. Gray, l. c.* 728. t. 94.—Ins. Sandvic.
paucidens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 4.—N. Guin.
Waimeae, *Wawra, in Flora*, lvi. (1873) 158.—Ins. Sandvic.

TETRAPLASIUM, Kunze, in Flora, xiv. (1831) 378 = **Tetilla**, DC. (Saxifr.).
petiolare, Kunze, ex Walp. Rep. ii. 265 = Tetilla hydrocotylaefolia.

TETRAPLEURA, Benth. in Hook. Journ. Bot. iv. (1842) 345. *LEGUMINOSAE*, Benth. & Hook. f. i. 590.
andongensis, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 331.—Afr. trop.
obtusangula, *Welw. ex Oliver, l. c.*—Afr. trop.
Thonningii, *Benth. in Hook. Journ. Bot.* iv. (1842) 345.—Afr. trop.

TETRAPLEURA, Parl. ex Webb, in Hook. Niger Fl. 131 (1849) = **Tornabenia**, Parl. (Umbell.).
insularis, Parl. ex Webb, l. c. = Torn. insularis.

TETRAPOGON, Desf. Fl. Atlant. ii. 388. t. 255 (1799). *GRAMINEAE*, Benth. & Hook. f. iii. 1170.
macranthus, *Benth. in Journ. Linn. Soc.* xix. (1881) 106.—Arab.
villosus, *Desf. Fl. Atlant.* ii. 388. t. 255.—Afr. bor.

TETRAPOMA, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 39 = **Nasturtium**, Linn. (Crucif.).
barbareaefolium, Turcz. l. c. 39 = N. palustre.
Crusianum, Walp. Rep. i. 154, sphalm. = seq.
Kruhsianum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 39 = N. palustre.
pyriforme, Seem. Bot. Voy. Herald, 24. t. 2.—Sibir.

TETRAPORA, Schau. in Linnaea, xvii. (1843) 238 = **Baeckea**, Linn. (Myrtac.).
glomerata, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 329 = B. pentandra.
Preissiana, Schau. in Linnaea, xvii. (1843) 238 = B. pentandra.
verrucosa, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 329 = Kunzea recurva.

TETRAPTERA, Miers, in Lindl. Veg. Kingd. 172 (1847) = **Burmannia**, Linn.

TETRAPTERA, Phil. in Bot. Zeit. xxviii. (1870) 169 = **Gaya**, H. B. & K. (Malvac.).
parviflora, Phil. l. c. = G. Gaudichaudiana.

TETRAPTERIS, Cav. Diss. ix. 433. tt. 260–262 (1790). *MALPIGHIACEAE*, Benth. & Hook. f. i. 260.
acapulcensis, *H. B. & K. Nov. Gen. et Sp.* v. 168.—Mexic.
acuminata, Raeusch. ex A. Juss. in Arch. Mus. Par. iii. (1843) 521 = mucronata.
acutifolia, *Cav. Diss.* ix. 433. t. 261.—Guian.
acutifolia, Griseb. in Linnaea, xiii. (1839) 236, partim = psilandra.
adenodon, Miq. l. c. xxii. (1849) 548 = multiglandulosa.
aequalis, *Wright, in Sauv. Fl. Cub.* 18.—Cuba.
affinis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 519.—Bras.
allopterys, *Moris, in Mem. Soc. Ital. Sc. Moden.* xxiv. II. (1850) 213.—N. Granat.
anisoptera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 520.—Bras.
argentea, *Bertol. in Nov. Comm. Acad. Bonon.* iv. (1840) 419.—Guatemala.
Benthamiana, *Griseb. in Mart. Fl. Bras.* xii. I. 88.—Bras.
Benthamii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 333.—N. Granat.
brachyptera, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 524 = magnifolia.
bracteolata, *Griseb. in Linnaea*, xiii. (1839) 238.—Bras.
buxifolia, *Cav. Diss.* ix. 434. t. 262. f. 1.—Ins. S. Doming.
calophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 525.—Guian.
chalcophylla, A. Juss. l. c. 529 = sericea.
chamaecerasifolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 7. t. 162.—Bras.
chamaedrifolia, Steud. Nom. ed. II. ii. 672, sphalm. = praec.
cinerea, *Griseb. in Linnaea*, xiii. (1839) 233.—Bras.
citrifolia, *Pers. Syn.* i. 508.—Jamaic.
citrifolia, Sw. Fl. Ind. Occ. ii. 857 = inaequalis.
complicata, *Miq. Stirp. Surin. Sel.* 82.—Guian.
Cotoneaster, *A. Juss. in Arch. Mus. Par.* iii. (1843) 528.—Mexic.
crebriflora, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 9.—Bras.
crebriflora, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 521, in obs. = Guilleminiana.
crispa, *A. Juss. l. c.* 522.—Guian.
crotonifolia, Benth. Pl. Hartw. 115 = Benthamii.
crotonifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 526.—Peruv.

TETRAPTERIS:—
cuspidata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 336.—N. Granat.
discolor, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 134 = ovalifolia.
discolor, *DC. Prod.* i. 587.—Guian.
eriocarpa, *Bertol. in Nov. Comm. Acad. Bonon.* iv. (1840) 419.—Guatemala.
fimbripetala, *A. Juss. in Arch. Mus. Par.* iii. (1843) 544.—Guian.
fraxinifolia, A. Juss. l. c. 537 = lucida.
Gaudichaudiana, A. Juss. in A. St. Hil. Fl. Bras. Mer. iii. 9 = Hiraea Gaudichaudiana.
glaberrima, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 134 = mucronata.
glabra, Griseb. in Mart. Fl. Bras. xii. I. 82 = punctulata.
glandulosa, *Griseb. l. c.* 83.—Bras.
Goudotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 333.—N. Granat.
gracilis, Walp. Ann. iii. 1152 (Index) = maranhamensis.
Guilleminiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 534.—Bras.
helianthemifolia, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 144.—Bras.
heteropetala, *Griseb. in Mart. Fl. Bras.* xii. I. 88.—Bras.
humilis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 546.—Bras.
inaequalis, *Cav. Diss.* ix. 433. t. 260.—Am. trop.
includens, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 133.—Guian.
Jamesonii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 394.—Ecuador.
Kohauti, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = citrifolia.
Lalandiana, A. Juss. in Arch. Mus. Par. iii. (1843 530 = sericea.
lancifolia, *A. Juss. l. c.* 536.—Bras.
Langsdorffiana, *A. Juss. l. c.* 539.—Bras.
lasiocarpa, A. Juss. l. c. 527 = rotundifolia.
laurifolia, *Griseb. in Mart. Fl. Bras.* xii. I. 88.—Bras.
leucanthele, Griseb. in Schomb. Faun. Fl. Brit. Gui. 1003 (= *Mascagnia leucanthele*).—Bras.; Guian.
leucosepala, *A. Juss. in Arch. Mus. Par.* iii. (1843) 531.—Bras.
ligustrifolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 12.—Bras.
longibracteata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 541.—Bras.
lucida, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 11.—Bras.
macrophylla, Poepp. ex Griseb. in Mart. Fl. Bras. xii. I. 77 = ovalifolia.
magnifolia, *Ruiz, ex Griseb. in Linnaea*, xxii. (1849) 22.—Bras.
maranhamensis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 537.—Bras.
metallicolor, A. Juss. l. c. 531 = punctulata.
mexicana, *Hook. & Arn. Bot. Beech. Voy.* 281.—Mexic.
mogoriifolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 11.—Bras.
mollis, *Griseb. in Linnaea*, xiii. (1839) 232.—Bras.
mucronata, *Cav. Diss.* ix. 434. t. 262. f. 2.—Guian.
multiflora, A. Juss. in Arch. Mus. Par. iii. (1843) 533 = mogoriifolia.
multiglandulosa, *A. Juss. l. c.* 532.—Bras.
mutabilis, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras.
nitida, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 523.—Bras.
obovata, Poepp. ex A. Juss. l. c. 314 = Hiraea crassipes.
oleifolia, *Griseb. in Mart. Fl. Bras.* xii. I. 86, in obs.—Guian.
ovalifolia, *Griseb. in Linnaea*, xiii. (1839) 237.—Bras.
paludosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 540.—Bras.
panamensis, Seem. Bot. Voy. Herald, 92 = Hiraea Barclayana.
paniculata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 245.—Porto Rico.
papyracea, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 334.—N. Granat.
pauciflora, DC. Prod. i. 587 = Diplopterys Paralias.

TETRAPTERIS :—
Poeppigiana, *Griseb. in Mart. Fl. Bras.* xii. 1. 87.—Bras.
psilandra, *Griseb. l. c.* 81.—Bras.
puberula, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 84.—Guian.
punctulata, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 10.—Bras.
racemulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 547.—Bras.
ramiflora, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 8, partim.—Bras.
ramiflora, A. Juss. l. c., partim = humilis.
renidens, *Griseb. in Mart. Fl. Bras.* xii. 1. 88.—Bras.
rotundifolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 6. t. 161.—Bras.
Schiedeana, *Cham. & Schlecht. in Linnaea*, v. (1830) 218.—Mexic.
Seemanni, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 335.—Panama.
sericea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 529.—Bras.
squarrosa, *Griseb. in Mart. Fl. Bras.* xii. 1. 87.—Bras.
Stephaniana, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 143.—Bras.
styloptera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 542.—Peruv.
suaveolens, *A. Juss. l. c.* 535.—Bolivia.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 55.—Guian.
tinifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 336.—N. Granat.
trinitensis, A. Juss. in Arch. Mus. Par. iii. (1843) 524 = discolor.
Turnerae, *Mart. ex A. Juss. l. c.* 546.—Bras.
vacciniifolia, *A. Juss. l. c.* 543.—Venezuela.
viburnifolia, Mart. ex A. Juss. l. c. 528 = rotundifolia.
xylosteifolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* iii. 7.—Bras.
Xylosteum, *Griseb. in Mart. Fl. Bras.* xii. 1. 84.—Bras.

TETRAPTERYGIUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 39 (1835). *CRUCIFERAE*, Benth. & Hook. f. i. 95.
glastifolium, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 204 = stylophorum.
glastifolium, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 39.—Armenia.
stylophorum, *Faub. & Spach, Illustr.* i. 100. t. 50.—Persia.

TETRAPTERYS, **A.** Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 261 = **Tetrapteris**, Cav.

TETRARHAPHIS, Miers, in Trans. Linn. Soc. xxi. (1853) 143 [t. 17] = **Oxytheca**, Nutt. (Polygon.).
apiculata, Miers, ex Benth. in DC. Prod. xiv. 23 = O. dendroidea.

TETRARIA, Beauv. in Mém. Inst. Fr. [Acad. Sc Par.] 1812 (1816), II. 54. *CYPERACEAE*, [Benth. & Hook. f. iii. 1063].
Buekia, Nees, in Linnaea, ix. (1835) 300.
Cyathocoma, Nees, l. c. (1835) 300.
Decalepis, Boeck. in Engl. Bot. Jahrb. v. (1884) 509.
Elynanthus, Nees, in Linnaea, vii. (1832) 520; ix. 298 (1835).
Ideleria, Kunth, Enum. Pl. ii. 310 (1837).
Lepisia, Presl, Lepis. Nov. Pl. Gen. (1829); et Symb. Bot. i. 9. t. 5 (1830).
Sclerochaetium, Nees, in Linnaea, ix. (1835) 302.
Trichoballia, Presl, Symb. Bot. i. 9 (1830).
compar, *Beauv. in Lestib. Ess. Cyp.* 36.—Afr. austr.
Thuarii, *Beauv. in Mém. Inst. Fr.* [*Acad. Sc. Par.*] 1812 (1816), II. 54.—Afr. austr.

TETRARRHENA, R. Br. Prod. 209 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1138.
acuminata, *R. Br. l. c.* 210.—Austral.
distichophylla, *R. Br. l. c.*—Austral.
Drummondiana, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 409 = laevis.
juncea, *R. Br. Prod.* 210.—Austral.
laevis, *R. Br. l. c.*—Austral.
tenacissima, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 409 = juncea.

TETRASPERMA, Steud. Nom. ed. II. ii. 673 (1841), sphalm. = TETRAPASMA, G. Don = **Discaria**, Hook. (Rhamn.).
juncea, Steud. l. c. = D. australis.

TETRASPIDIUM, Baker, in Journ. Linn. Soc. xx. (1883) 215. *SCROPHULARINEAE*.
laxiflorum, *Baker, l. c.*—Madag.

TETRASPORA, Miq. in Nederl. Kruidk. Arch. iv. (1859) 150 = **Baeckea**, Linn. (Myrtac.).
Gunniana, Miq. l. c. = B. Gunniana.

TETRASTEMON, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 317 = **Myrrhinium**, Schott (Myrtac.).
loranthoides, Hook. & Arn. l. c. 318 = M. rubriflorum.

TETRASTYLIDIUM, Engl. in Mart. Fl. Bras. xii. II. 33 (1872) *OLACIENAE*.
brasiliense, *Engl. l. c.*—Bras.

TETRASTYLIS, Barb. Rodrig. in Rev. Engenh. (Nov. 14, 1882), reimpr. 1 = **Passiflora**, Linn.
montana, Barb. Rodrig. l. c., reimpr. 3 = P. ovalis.

TETRATAXIS, Hook. f. in Benth. & Hook. f. Gen. i. 783 (1867). *LYTHRARIEAE*, Benth. & Hook. f. l. c.
Tetradia, Thou. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 137.
salicifolia, *Baker, Fl. Maurit.* 100.—Ins. Mascar.

TETRATELEIA, Sond. in Harv & Sond. Fl. Cap. i. 58 (1859) = **Cleome**, Linn. (Cappar.).
maculata, Sond. l. c. (= *Polanisia maculata*).—Afr. austr.

TETRATHECA, Sm. Bot. Nov. Holl. i. t. 2 (1793). *TREMANDREAE*, Benth. & Hook. f. i. 134.
Petrotheca, Steud. Nom. ed. II. ii. 310 (1841), sphalm.
aculeata, Steetz, in Lehm. Pl. Preiss. i. 218 = hirsuta.
affinis, *Endl. in Enum. Pl. Hueg.* 7.—Austral.
aphylla, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 97.—Austral.
baueraefolia, F. Muell. in Schuch. Syn. Trem. 29 = ciliata.
calva, Schuch. l. c. 25 = pilosa.
ciliata, *Lindl. in Mitch. Three Exped.* ii. 206.—Austral.
confertifolia, *Steetz, in Lehm. Pl. Preiss.* i. 214.—Austral.
denticulata, Sieber, ex Spreng. Syst. iv. App. 147 = pilosa.
efoliata, *F. Muell. Fragm.* x. 3.—Austral.
elongata, Schuch. Syn. Trem. 38 = setigera.
epilobioides, Steetz, in Lehm. Pl. Preiss. i. 218 = hirsuta.
ericifolia, F. Muell. Pl. Vict. i. 182, partim = pilosa.
ericifolia, *Sm. Exot. Bot.* i. 37. t. 20.—Austral.
ericoides, Planch. in Van Houtte, Fl. des Serres, x. (1854-55) 229. t. 1065 = pilosa.
filiformis, *Benth. Fl. Austral.* i. 135.—Austral.
galioides, Schnizl. Ic. t. 232 = Platytheca galioides.
glandulosa, *Labill. Nov. Holl. Pl.* i. 95. t. 123.—Austral.
glandulosa, Sm. Exot. Bot. i. 39. t. 21 = pilosa.
gracilis, Steetz, in Lehm. Pl. Preiss. i. 215 = viminea.
Gunnii, Hook. f. Fl. Tasm. i. 36 = pilosa.
Harperi, *F. Muell. Fragm.* v. 49. t. 38.—Austral.
hirsuta, *Lindl. Swan River App.* 38.—Austral.
hispidissima, *Steetz, in Lehm. Pl. Preiss.* i. 217.—Austral.
juncea, *Sm. Spec. Bot. Nov. Holl.* t. 2.—Austral.
micrantha, Schuch. Syn. Trem. 43 = pilifera.
nuda, *Lindl. Swan River App.* 38.—Austral.
oppositifolia, Pers. Syn. i. 419 = Boronia polygalifolia.
pilifera, *Lindl. Swan River App.* 38.—Austral.
pilosa, *Labill. Nov. Holl. Pl.* i. 95. t. 122.—Austral.
Preissiana, Steetz, in Lehm. Pl. Preiss. i. 219 = pilifera.
procumbens, Gunn, ex Hook. f. Fl. Tasm. i. 35 = pilosa.
pubescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 141.—Austral.

TETRATHECA :—
rubiaeoides, A. Cunn. in Field, N. S. Wales, 335 = ericifolia.
rubriseta, Lindl. Swan River App. 38 = hirsuta.
setigera, *Endl. in Enum. Pl. Hueg.* 8.—Austral.
speciosa, Hort. ex Walp. Ann. ii. 87 = Platytheca galioides.
subaphylla, *Benth. Fl. Austral.* i. 132.—Austral.
tenuiramea, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 142 = pubescens.
thymifolia, *Sm. Exot. Bot.* i. 41. t. 22.—Austral.
verticillata, Paxt. Mag. Bot. xiii. (1847) 171 = Platytheca galioides.
viminea, *Lindl. Swan River App.* 38.—Austral.
virgata, *Steetz, in Lehm. Pl. Preiss.* i. 212.—Austral.

TETRATHYLACIUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 34. t. 240 (1842). *VIOLARIEAE*, Benth. & Hook. f. i. 119.
Edmonstonia, Seem. Bot. Voy. Herald, 98. t. 18 (1853?).
macrophyllum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 34. t. 240.—Chili.

TETRATHYRIUM, Benth. Fl. Hongk. 132 (1861) = **Loropetalum**, R. Br. (Hamamel.).
subcordata, Benth. l. c. 133 = L. subcordatum.

TETRATOME, Poepp. & Endl. Nov. Gen. et Sp. ii. 46. t. 163 (1838) = **Mollinedia**, Ruiz & Pav. (Monim.).
arguta, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 321 = M. nitida.
cinerea, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 136 = M. cinerea.
elliptica, Gardn. l. c. i. (1842) 530 = M. nitida.
latifolia, Poepp. & Endl. Nov. Gen. et Sp. ii. 47 = M. latifolia.
lepidota, Poepp. & Endl. l. c. = Citriosma Poeppigii.
racemosa, Poepp. ex Schlecht. in Linnaea, xx. (1847) 114 = M. racemosa.
tomentosa, Benth. Pl. Hartw. 250 = M. tomentosa.
triflora, Poepp. & Endl. Nov. Gen. et Sp. ii. 46. t. 163 = M. racemosa.

TETRAULACIUM, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 53. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 949.
veronicaeforme, *Turcz. l. c.* 54.—Bras.

TETRAZYGIA, Rich. ex DC. Prod. iii. 172 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 762.
Antheryta, Rafin. Sylva Tellur. 95 (1838).
Chitonia, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 339.
Harrera, Macfad. Fl. Jamaic. ii. 60 (1837).
Lomanthera, Rafin. Sylva Tellur. 96 (1838).
Miconiastrum, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 341.
Naudinia, A. Rich. Ess. Fl. Cub. i. 561 (1845).
adenophora, Griseb. in Goett. Abh. ix. (1861) 53 = Miconia aurea.
albicans, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 100.—Ind. occ.
ancistrophora, Wright, in Sauv. Fl. Cub. 48 = Miconia ancistrophora.
angustiflora, Griseb. in Goett. Abh. ix. (1861) 53 = Miconia delicatula, pallens.
angustifolia, *DC. Prod.* iii. 172.—Ind. occ.
brachycentra, *Wright, in Sauv. Fl. Cub.* 49.—Cuba.
cornifolia, Griseb. in Goett. Abh. ix. (1861) 53 = Miconia cinnamomifolia.
crotonifolia, *DC. Prod.* iii. 172.—Ins. S. Doming.
discolor, *DC. l. c.*—Ind. occ.
elaeagnoides, *DC. l. c.*—Ind. occ.
elaeagnoides, Griseb. Cat. Pl. Cub. 98 = Miconia delicatula.
elaeagnoides, Griseb. Fl. Brit. W. Ind. 255 = laxiflora.
elaeagnoides, Hook. Bot. Mag. t. 4383 = Miconia pallens.
Fadyenii, Hook. Kew Journ. i. (1849) 379 = Charianthus Fadyeni.
hispida, *Macfad. Fl. Jamaic.* ii. 58.—Jamaic.
laxiflora, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 343.—Ind. occ.
semicrenata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 185 = Miconia semicrenata.
smaragdina, Griseb. in Goett. Abh. ix. (1861) 53 = Miconia smaragdina.
tetrandra, DC. Prod. iii. 172 = Miconia tetrandra.

TETRAZYGOS, Rich. ex DC. Prod. iii. 197 (1828) = **Charianthus**, D. Don (Melastom.).
ciliatus, Rich. ex DC. l. c. = C. ciliatus.

TETRAZYGUS, Triana, in Trans. Linn. Soc. xxviii. (1871) 99 = praec.

TETREILEMA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 199. Genus dubium *VERBENACEARUM*.—Cf. Benth. & Hook. f. Gen. ii. 1137.
articulatum, *Turcz. l. c.* 200.—Chili.
boliviense, *Turcz. l. c.* (bolierense).—Bolivia.

TETRODEA, Rafin. New Fl. Am. i. 81 (1836) = **Amphicarpaea**, Ell. (Legumin.).
comosa, Rafin. l. c. = A. monoica.
monoica, Rafin. l. c. = A. monoica.
subterranea, Rafin. l. c. = Voandzeia subterranea.

TETRODUS, Cass. in Dict. Sc. Nat. lv. 264, 272 (1828) = **Helenium**, Linn. (Compos.).
quadridentatus, Cass. ll. cc. = H. quadridentatum.

TETRONCIUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 17. *NAIADACEAE*, Benth. & Hook. f. iii. 1013.
CATHANTHES, Rich. in Mém. Mus. Par. i. (1815) 365.
magellanicum, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 17.—Reg. Magellan.

TETRORCHIDIUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 23. t. 227 (1842). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 288.
andinum, *Muell. Arg. in Flora*, xlvii. (1864) 538.—Peruv.
macrophyllum, *Muell. Arg. in DC. Prod.* xv. II. 1133.—Peruv.
parvulum, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 513.—Bras.
rubrivenium, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 23.—Costa Rica.
trigynum, Baill. Etud. Gén. Euph. 440 = rubrivenium.

TETRORHIZA, Rafin. Fl. Tellur. iii. 25 (1836) = TRETORRHIZA, Renealm. = **Gentiana**, Tourn.

TETRORRHIZA, Steud. Nom. ed. II. ii. 673 (1841) = praec.

TEUCRIDIUM, Hook. f. Fl. N. Zel. i. 203. t. 49 (1854). *VERBENACEAE*, Benth. & Hook. f. ii. 1157.
parvifolium, *Hook. f. Fl. N. Zel.* 203. t. 49.—N. Zel.

TEUCRION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = **Teucrium**, Tourn. (Labiat.).
chamaedryum, St. Lag. l. c. = Teucrium Chamaedrys.
scorodonium, St. Lag. l. c. = Teucrium Scorodonia.

TEUCRIUM, [Tourn.] Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1221.
BOTRYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138.
CHAMAEDRYS, Tourn. ex Hall. Enum. Stirp. Helv. ii. 630 (1742).
MELOSMON, Rafin. Fl. Tellur. iii. 85 (1836).
MONIPSIS, Rafin. l. c. (1836).
MONOCHILON, Dulac, Fl. Hautes-Pyr. 405 (1867).
POLIODENDRON, Noé, in Webb & Berth. Phyt. Canar. iii. 106. t. 173 (1835–60).
POLIUM, Tourn. ex Linn. Syst. ed. I (1735).
SCORDIUM, Cav. Ic. i. 19. t. 31 (1791).
SCORODONIA, Adans. Fam. ii. 188 (1763).
TRIXAGO, Rafin. Fl. Tellur. iii. 85 (1836).
abutiloides, *L'Hérit. Stirp. Nov.* 84.—Ins. Mader.
abyssinicum, Hochst. ex Benth. in DC. Prod. xii. 586 = T. Scordium.
Achaemenis, Schreb. Pl. Vert. Unilab. 44 = T. Polium.
aegypticum, Pers. Syn. ii. 112 = T. Polium.
africanum, *Thunb. Prod. Pl. Cap.* 95.—Afr. austr.
albidum, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 286.—Alger.
album, Poir. Encyc. Suppl. ii. 770 = T. Polium.
Alopecurus, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 585.—Afr. bor.

TEUCRIUM:—
alpestre, *Sibth. & Sm. Fl. Graec.* vi. 31. t. 538.—Ins. Creta.
alpinum, Linn. Syst. ed. X. 1094 = T. Botrys.
alyssifolium, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 104.—Lycia.
amplexicaule, *Benth. Lab. Gen. et Sp.* 687.—Hispan.
amplexicaule, Wallr. in Linnaea, xiv. (1840) 594 (Beitr. Fl. Hercyn. 224) = scordioides.
angustifolium, Salisb. Prod. 76 = fruticans.
angustifolium, Steud. Nom. ed. I. 828 = T. Polium.
angustissimum, Schreb. Pl. Vert. Unilab. 49 = T. Polium.
aragonense, Huet, ex Čelak. in Bot. Centralbl. xiv. (1883) 186 = chlorocephalum.
aragonense, Losc. & Pard. Ser. Inconf. 85 = T. Polium.
Arduini, *Linn. Mant.* i. 81.—Europ. austr.
arenarium, S. T. Gmel. Reise, i. 149. t. 26 = T. Scordium.
argutum, *R. Br. Prod.* 504.—Austral.
aroanium, *Orph. in Boiss. Diagn.* Ser. II. iv. 55.—Graecia.
arvense, Salisb. Prod. 76 = Ajuga Chamaepitys.
asiaticum, Linn. Mant. i. 80 = lancifolium.
atratum, *Pomel, Nouv. Mat. Fl. Atl.* 304.—Alger.
aureiforme, Pomel, l. c. 113 = T. Polium.
aureum, Schreb. Pl. Vert. Unilab. 43 = T. Polium.
australe, Sieber. ex Steud. Nom. ed. II. ii. 673 = Anisomeles australis.
baeticum, Boiss. & Reut. Pugill. Pl. Nov. 98 = T. Pseudo-scorodonia.
baeticum, Clem. ex Nym. Consp. 564 = fruticans.
Balansae, Ball, in Journ. Linn. Soc. xvi. (1878) 632 = mauritanicum.
balearicum, Coss. ex Marès. & Vigin. Cat. Pl. Baléar. 224 = subspinosum.
Belion, Schreb. Pl. Vert. Unilab. 47 = T. Polium.
betonicaefolium, Jacq. Coll. i. 145 = betonicum.
betonicum, *L'Hérit. Stirp. Nov.* 83. t. 40.—Ins. Mader.
bicolor, *Sm. in Rees, Cycl.* xxxv. [n. 25].—Peruv.; Chili.
Botrydium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = T. Botrys.
botryoides, Hort. ex Steud. Nom. ed. II. ii. 673 = T. Botrys.
Botrys, *Linn. Sp. Pl.* 562.—Europ.; Afr. bor.
bracteatum, *Desf. Fl. Atlant.* ii. 7. t. 120.—Reg. Mediterr. occ.
brevifolium, *Schreb. Pl. Vert. Unilab.* 27.—Graecia; Ins. Creta.
bullatum, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 260.—Marocc.
buxifolium, *Schreb. Pl. Vert. Unilab.* 42.—Hispan.; Creta.
caespitulosum, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 430 = T. Polium.
campanulatum, *Linn. Sp. Pl.* 562.—Ital.; Graecia.
canadense, *Linn. l. c.* 564.—Am. bor.
canadense, Torr. in Bot. U. St. Expl. Exped. ii. II. 409 = occidentale.
canariense, Lam. Encyc. ii. 692 = heterophyllum.
canescens, Forst. f. in Comm. Gotting. ix. (1789) 58 = betonicum.
canum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 40 = T. Chamaedrys.
capense, *Thunb. Prod. Pl. Cap.* 95.—Afr. austr.
capitatum, Linn. Sp. Pl. 566 = T. Polium.
carthaginense, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 97.—N. Granat.
caucasicum, Willd. ex Spreng. Syst. ii. 708 = T. Scordium.
Cavanillesii, Bert. ex Steud. Nom. ed. II. ii. 674 = bicolor.
Cephalotes, Pomel, Nouv. Mat. Fl. Atl. 116 = T. Polium.
chamaedrifolium, Mill. Gard. Dict. ed. VIII. n. 16 = cubense.
Chamaedrys, *Linn. Sp. Pl.* 565.—Europ.; As. bor.; Oriens.
Chamaepitys, Linn. l. c. = Ajuga Chamaepitys.
charamoniense, Cav. Desc. 82 = creticum.
chilense, *Desf. ex Steud. Nom.* ed. II. ii. 674.—Chili.
chium, J. F. Gmel. Syst. 893 = Ajuga chia.
chlorocephalum, *Čelak. in Bot. Centralbl.* xiv. (1883) 186.—Hispan.
chrysotrichium, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 96.—Europ.
Chulilia, Cav. ex Spreng. Syst. ii. 708 = T. Polium.
cinereum, *Boiss. Elench.* 79.—Hispan.

TEUCRIUM:—
cinereum, Rafin. New Fl. Am. iv. 98 (Quid ?).—Am. bor.
coeleste, Schreb. Pl. Vert. Unilab. 49 = T. Polium.
collinum, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 258.—Marocc.
collinum, Salisb. Prod. 75 = T. Botrys.
collinum, Salisb. l. c. 77 = T. Polium.
Commersonii, Spreng. Syst. ii. 710 = Salvia procurrens.
compactum, *Clem. ex Lag. Gen. et Sp. Nov.* 17.—Hispan.
compositum, Pomel, Nouv. Mat. Fl. Atl. 114 = T. Polium.
coniortodes, *Boiss. & Blanche, ex Boiss. Fl. Orient.* iv. 818.—Syria.
corbariense, Lapeyr. ex Steud. Nom. ed. I. 829 = resupinatum.
corbariense, Pourr. ex Nym. Consp. 565 = scordioides.
cordifolium, *Čelak. in Bot. Centralbl.* xiv. (1883) 221.—Europ.
corymbiferum, Desf. Fl. Atlant. ii. 8 = T. Polium.
corymbosum, *R. Br. Prod.* 504.—Austral.
creticum, Lam. Encyc. ii. 692 = brevifolium.
creticum, *Linn. Sp. Pl.* 563.—Reg. Mediterr.
crispum, Pomel, Nouv. Mat. Fl. Atl. 304 = T. Pseudo-scorodonia.
cubense, *Jacq. Enum. Pl. Carib.* 25.—Am. bor. et trop.
cuneifolium, Schimp. ex Boiss. Diagn. Ser. I. v. 44 = leucocladum.
cuneifolium, *Sibth. & Sm. Fl. Graec.* vi. 30. t. 537.—Ins. Creta.
cuneifolium, Webb, Iter Hisp. 28 = fragile.
cuspidatum, *Čelak. in Bot. Centralbl.* xiv. (1883) 222.—Europ.
cylindricum, Lam. ex Steud. Nom. ed. I. 829 = T. Polium.
cymosum, *Pers. Syn.* ii. 112.—Hispan.
cyprium, *Boiss. Diagn.* Ser. I. v. 43.—Ins. Cyprus.
Decaisnii, *Presl, Bot. Bemerk.* 100.—Arab.
decipiens, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 258.—Marocc.
densiflorum, Vis. Sem. Rar. Hort. Patav. (1847) 4.—Cf. Linnaea, xxiv. (1851) 181 = Arduini.
divaricatum, *Sieber, ex Benth. Lab. Gen. et Sp.* 682, in syn.; *Boiss. Fl. Orient.* iv. 816.—Graec.; As. Min.; Syria.
diversiflorum, Moench, Meth. Suppl. 132 = campanulatum.
elevatum, Benth. in Wall. Cat. n. 2026 = stoloniferum.
elongatum, Menyh. Kalocsa Vidék. Növényt. 146 = T. Scordium.
eriocalyx, *Čelak. in Bot. Centralbl.* xiv. (1883) 187.—Hispan.
eriocephalum, *Willk. in Linnaea*, xxv. (1852) 58.—Hispan.
euganeum, Vis. Ind. Sem. Hort. Patav. (1836); ex Walp. Rep. iii. 923 = T. Scorodonia.
fasciculatum, *Poir. Encyc. Suppl.* ii. 767.—Hab.?
filifolium, Willd. ex Benth. in Linnaea, xi. (1837) 348 = Campylanthus salsoloides.
filiforme, Poir. Encyc. Suppl. ii. 769 = Campylanthus salsoloides.
fililobum, *F. Muell. ex Benth. Fl. Austral.* v. 134.—Austral.
flavescens, Schreb. Pl. Vert. Unilab. 44 = T. Polium.
flavicans, Lam. Encyc. ii. 700 = T. Polium.
flavum, *Linn. Sp. Pl.* 565.—Reg. Mediterr.
foliosum, Pomel, Nouv. Mat. Fl. Atl. 115 = T. Polium.
Fortunei, Benth. in DC. Prod. xii. 583 = quadrifarium.
fragile, *Boiss. Elench.* 77.—Hispan.
fruticans, *Linn. Sp. Pl.* 563.—Reg. Mediterr. occ.
fulvum, Hance, in Walp. Ann. iii. 270 = quadrifarium.
Funkianum, Willk. Iber. Halbinsel, 134 = T. Polium.
garganicum, Fisch. ex Steud. Nom. ed. I. 829 (Quid ?).—Hab.?
Gasparrini, Nym. Consp. 564 = T. Scorodonia.
genevense, Crantz, Stirp. Austr. ed. II. 253 = Ajuga genevensis.
glandulosum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 23.—Mexic.
globuliferum, Salisb. Prod. 77 = abutiloides.
gnaphalodes, L'Hérit. Stirp. Nov. 84; Vahl, Symb. Bot. i. 41 = T. Polium.
graecum, Čelak. in Bot. Centralbl. xiv. (1883) 217 = divaricatum.
granatense, *Boiss. & Reut. Pugill. Pl. Nov.* 99.—Reg Mediterr. occ.
Gussonii, Nym. Consp. 567 = T. Polium.

TEUCRIUM:—
Haenseleri, *Boiss. Elench.* 79.—Hispan.
Halacsyanum, *Heldr. in Oestr. Bot. Zeitschr.* xxix. (1879) 241.—Graec.
heliotropifolium, *Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1875) 97.—Ins. Karpathos.
heterophyllum, Cav. Ic. vi. 56 = bicolor.
heterophyllum, *L'Hérit. Stirp. Nov.* 84. t. 49.—Ins. Canar.
hircanicum, *Linn. Syst.* ed. X. 1095.—Reg. Cauc.; Persia.
hirtum, Willd. ex Spreng. Syst. ii. 710 = inflatum.
hispidum, Benth. ex Wall. Cat. n. 2029 = Ajuga parviflora.
hyrcanicum, Steud. Nom. ed. II. ii. 674 = hircanicum.
hyssopifolium, Schreb. Pl. Vert. Unilab. 28 = creticum.
incanum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 184.—Afghan.
incanum, Schur, Enum. Pl. Transs. 547 = montanum.
inflatum, *Sw. Prod. Veg. Ind. Occ.* 88.—Am. austr.
integerrimum, Steud. Nom. ed. II. ii. 674 = T. Polium.
integrifolium, *Benth. Fl. Austral.* v. 133.—Austral.
intricatum, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 21.—Hispan.
Iva, Georgi, Beschr. Russ. Reich. III. v. 1076 = Ajuga Laxmanni.
Iva, Linn. Sp. Pl. 563 = Ajuga Iva.
japonicum, *Willd. Sp. Pl.* iii. 23.—Japon.
Kotschyanum, *Poech, Enum. Pl. Ins. Cyp.* 24.—As. Min.
laciniatum, *Torr. in Ann. Lyc. N. York*, ii. (1828) 231.—Am. bor. occ.
laevigatum, [Soland. in] Russ. Aleppo, ed. II. ii. 255 = Ajuga laevigata.
laevigatum, Vahl, Symb. Bot. i. 40 = cubense.
lamiifolium, Urv. in Mém. Soc. Linn. Par. i. (1822) 320 = Arduini.
lanceolatum, Benth. Lab. Gen. et Sp. 666 = corymbosum.
lanceolatum, Rafin. & Collins, in Rafin. New Fl. Am. iv. 97 (Quid?).—Am. bor.
lancifolium, *Boiss. Diagn.* Ser. II. iv. 56.—Ins. Balear.
lanigerum, Lag. Gen. et Sp. Nov. 17 = T. Polium.
lanuginosum, Hoffmgg. & Link, Fl. Port. i. 84. t. 3 = scordioides.
latifolium, Linn. Sp. Pl. 563 = fruticans.
lavandulatum, Pourr. ex Benth. Lab. Gen. et Sp. 686 = T. Polium.
Laxmanni, Murr. Syst. ed. XIII. 439 = Ajuga Laxmanni.
laxum, *D. Don, Prod. Fl. Nep.* 109.—Reg. Himal.
leucocladum, *Boiss. Diagn.* Ser. I. v. 44.—Arab.
leucophyllum, *Montb. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 55.—Cappadoc.
Leucosceptrum, Voigt, Hort. Suburb. Calc. 463 = Leucosceptrum canum.
levigatum, Rafin. New Fl. Am. iv. 96 = cubense?
Libanitis, Schreb. Pl. Vert. Unilab. 48 = pumilum.
lucidum, *Linn. Syst.* ed. X. 1095.—Reg. Mediterr.
lucidum, Sibth. & Sm. Fl. Graec. vi. 27. t. 532 = divaricatum.
lusitanicum, Benth. in DC. Prod. xii. 585, partim = lancifolium.
lusitanicum, *Lam. Encyc.* ii. 694.—Europ. austr.
lusitanicum, Schreb. Pl. Vert. Unilab. 47 = T. Polium.
macrophyllum, Lam. Encyc. ii. 692 = abutiloides.
macrostachyum, Wall. ex Benth. Lab. Gen. et Sp. 664 = Leucosceptrum canum.
macrum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 810.—Persia.
maderense, Lam. Encyc. ii. 692 = betonicum.
Majorana, *Pers. Syn.* ii. 112.—Hispan.
majoricum, *Rouy, in La Natural.* ii. (1882) (*Matér. Rév. Fl. Portug.* 11, 15).—Ins. Balear.
maritimum, Lam. Fl. Fr. ii. 414 = T. Marum.
Marum, *Linn. Sp. Pl.* 564.—Reg. Mediterr.
mascatense, *Boiss. Diagn.* Ser. I. v. 44.—Arab.
massiliense, *Linn. Sp. Pl.* ed. II. 789.—Europ. austr.
massiliense, Lour. Fl. Cochinch. 362 = stoloniferum.
mauritanicum, *Noé, ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 226; *et in Bull. Soc. Bot. Fr.* ii. (1855) 585.—Alger.
mauritanum, Linn. Sp. Pl. 563 = T. Pseudo-chamaepitys.
Mazziarii, Heldr. ex Nym. Consp. 567 = Coleus arabicus.
melissaefolium, Noronha, in Verh. Batav. Gen. v. (1790) ed. I Art. IV. 27 = stoloniferum.
melissoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 813.—Persia.

TEUCRIUM:—
microphyllum, *Desf. in Ann. Mus. Par.* x. (1807) 300.—Ins. Creta.
micropodioides, *Rouy, in La Natural.* ii. (1882) (*Matér. Rév. Fl. Portug.* 16).—Lusitan.
montanum, *Linn. Sp. Pl.* 565.—Europ.; Oriens.
Montbretii, *Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 56.—As. Min.; Syria.
moschatum, Mill. Gard. Dict. ed. VIII. n. 15 = Ajuga Iva.
mucronatum, Linn. Sp. Pl. 566 = spinosum.
multicaule, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 54.—Syria.
multiflorum, Hort. ex Benth. Lab. Gen. et Sp. 680 = T. Chamaedrys.
multiflorum, *Linn. Sp. Pl.* 564.—Hispan.
myrtifolium, Desf. Cat. Hort. Par. ed. III. 394 = T. Marum.
myrtifolium, *Poir. Encyc. Suppl.* ii. 766.—Oriens.
nepetaefolium, *Benth. in DC. Prod.* xii. 580.—China.
Nissolianum, Linn. Sp. Pl. 563 = T. Pseudo-chamaepitys.
nitidum, *Schreb. Pl. Vert. Unilab.* 35.—Mauritan.
nivale, Boiss. Diagn. Ser. I. xii. 91 = orientale.
nuchense, C. Koch, in Linnaea, xxi. (1848) 704 = T. Chamaedrys.
nudicaule, *Hook. Bot. Misc.* ii. (1831) 235.—Peruv.
occidentale, *A. Gray, Syn. Fl. N. Am.* ii. 1. 349.—Am. bor.
ochroleucum, Jord. Pugill. Pl. Nov. 137 = montanum.
Odontites, *Boiss. & Bal. Diagn.* Ser. II. iv. 57.—Cilicia.
odoratum, Lam. Fl. Fr. ii. 413 = massiliense.
odorum, Salisb. Prod. 76 = T. Marum.
officinale, Lam. Fl. Fr. ii. 414 = T. Chamaedrys.
oliganthum, *Hassk. Cat. Hort. Bog. Alt.* 310.—Japon.
Oliverianum, *Ging. ex Benth. Lab. Gen. et Sp.* 668.—Persia; Assyria.
orchideum, Lindl. Bot. Reg. t. 1255 = bicolor.
orientale, *Linn. Sp. Pl.* 562.—As. Min.
paederotoides, *Boiss. Fl. Orient.* iv. 1268 (Index).—Cf. 814.—Syria; Persia.
palmatum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 702.—Reg. Himal.; China.
palustre, H. B. & K. Nov. Gen. et Sp. ii. 306 = inflatum.
palustre, Lam. Fl. Fr. ii. 411 = T. Scordium.
paniculatum, Herb. Madr. ex Wall. Cat. n. 2025 = tomentosum.
pannonicum, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 384 = montanum.
parviflorum, *Schreb. Pl. Vert. Unilab.* 31.—As. Min.; Persia.
Pernyi, *Franch. Pl. Davidian.* i. (1884) 245.—China.
persicum, *Boiss. Diagn.* Ser. I. v. 42.—Persia.
Pestalozzae, *Boiss. l. c.* xii. 90.—Syria.
petiolare, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotra.
petiolare, Rafin. New Fl. Am. iv. 97 (Quid?).—Am. bor.
petrophilum, F. Muell. in Linnaea, xxv. (1852) 426 = corymbosum.
pictum, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 809 = procerum.
poederotoides, Boiss. & Haussk. ex Boiss. l. c. 814, err. typ. = paederotoides.
Polium, Decne. ex Presl, Bot. Bemerk. 100 = Decaisnii.
Polium, *Linn. Sp. Pl.* 566.—Reg. Mediterr.; Oriens.
Polium-trifoliatum, Vahl, Symb. Bot. i. 40 = T. Polium.
polycephalon, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 216. t. 13 = T. Polium?
polycephalum, Pomel, Nouv. Mat. Fl. Atl. 114 = T. Polium.
procerum, *Boiss. & Blanche, Diagn.* Ser. II. iv. 56.—Syria; Persia.
prostratum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotra.
prostratum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 60.—Transsylv.
pruinosum, *Boiss. Fl. Orient.* iv. 808.—Cappadoc.
Pseudiva, Gueldenst. ex Ledeb. Fl. Ross. iii. 449 = Ajuga Laxmanni.
Pseudo-chamaedrys, Wender. in Flora, ix. (1826) 358 = T. Chamaedrys.
Pseudo-chamaepitys, Georgi, Beschr. Russ. Reich. III. v. 1076 = Ajuga chia.

TEUCRIUM:—
Pseudo-chamaepitys, *Linn. Sp. Pl.* 562.—Reg. Mediterr.
Pseudo-hyssopus, Schreb. Pl. Vert. Unilab. 45 = T. Polium.
Pseudo-scorodonia, *Desf. Fl. Atlant.* ii. 5. t. 119.—Reg. Mediterr. occ.
pulverulentum, Coss. ex Barcelo, Fl. Balear. 375; et Rouy, in La Natural. ii. (1882) (Matér. Rév. Fl. Portug. 16) = T. Polium.
pumilum, *Linn. Cent. Pl.* i. 15; *Amoen. Acad.* iv. 276.—Hispan.
purpurascens, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 210 = T. Polium?
purpureum, Link, Handb. i. 456 = T. Polium.
pycnaphyllum, Schreb. Pl. Vert. Unilab. 48 = T. Polium.
pyramidale, Crantz, Stirp. Austr. ed. II. 252 = Ajuga pyramidalis.
pyrenaicum, *Linn. Sp. Pl.* 566.—Europ. austr.
quadratulum, *Schreb. Pl. Vert. Unilab.* 36.—Hispan.
quadratulum, Sibth. & Sm. Fl. Graec. vi. 25. t. 530 = microphyllum.
quadrifarium, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 108.—Reg. Himal.; China.
quadrifarium, Wight, ex Hook. f. Fl. Brit. Ind. iv. 701 = Wightii.
racemosum, *R. Br. Prod.* 504.—Austral.
ramosissimum, *Desf. Fl. Atlant.* ii. 4. t. 118.—Hispan.
regium, Auct. ex Boiss. Fl. Orient. iv. 817 = divaricatum.
regium, Hort. ex Benth. Lab. Gen. et Sp. 682 = flavum.
regium, *Schreb. Pl. Vert. Unilab.* 35.—Hispan.
reptans, Pourr. in Mém. Acad. Toul. iii. (1788) 330 = pyrenaicum.
reptans, Salisb. Prod. 75 = Ajuga reptans.
resupinatum, Balansa, ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 226 = mauritanicum.
resupinatum, *Desf. Fl. Atlant.* ii. 4. t. 117.—Reg. Mediterr. occ.
revolutum, Stokes, Bot. Mat. Med. iii. 295 = montanum.
rhombifolium, Willd. ex Spreng. Syst. ii. 732 = Hyptis canescens.
rigidum, *Benth. in DC. Prod.* xii. 578.—Assyria.
riparium, *Hochst. in Flora*, xxviii. (1845) 66.—Afr. austr.
riparium, Rafin. New Fl. Am. iv. 97 (Quid?).—Am. bor.
rosmarinifolium, Herb. Bout. ex Willk. & Lange, Prod. Fl. Hisp. ii. 469 = fruticans.
rosmarinifolium, Lam. Encyc. ii. 693 = creticum.
rosmarinifolium, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 476 = pumilum.
rotundifolium, Poir. Encyc. Suppl. ii. 767 = Salvia procurrens.
rotundifolium, Schreb. Pl. Vert. Unilab. 42 = pyrenaicum.
Royleanum, *Wall. Cat.* n. 2027.—Reg. Himal.
rupestre, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 259.—Marocc.
salicifolium, Linn. Mant. i. 80 = Ajuga salicifolia.
salviaefolium, Salisb. Prod. 76 = T. Scorodonia.
Salviastrum, Hoffmgg. & Link, Fl. Port. i. 85. t. 2 = lusitanicum.
Salviastrum, Schreb. Pl. Vert. Unilab. 38 = T. Pseudo-scorodonia.
saxatile, Cav. Ic. ii. 19. t. 121. f. 1 = buxifolium.
saxatile, Lam. Encyc. ii. 699 = pyrenaicum.
saxatile, Sieber, ex Benth. Lab. Gen. et Sp. 685 = alpestre.
saxatile, Sieber, ex Čelak. in Bot. Centralbl. xiv. (1883) 187 = Sieberi.
Schimperi, Presl, Bot. Bemerk. 100 = leucocladum.
scordioides, *Schreb. Pl. Vert. Unilab.* 37.—Reg. Mediterr.; Oriens.
Scordium, Brot. Fl. Lusit. i. 164 = scordioides.
Scordium, *Linn. Sp. Pl.* 565.—Europ.; As. temp.; Afr. bor.
Scorodonia, *Falk, Beitr.* ii. 204 (Quid?).—Sibir. altaic.
Scorodonia, *Linn. Sp. Pl.* 564.—Europ.
Scorodonia, Sibth. & Sm. Fl. Graec. Prod. i. 392 = Kotschyanum.
scrobiculatum, *C. Koch, in Linnaea*, xxi. (1848) 705.—Oriens.
secundum, Heyne, in Wall. Cat. n. 2028 = Anisomeles Heyneana.

TEUCRIUM:—
serratum, *Benth. in DC. Prod.* xii. 586.—Afghan.
sessiliflorum, *Benth. l. c.* 580.—Austral.
sessilifolium, *Clarke, Trav.* iv. 555 (sessifolium).—Thracia.
sibiricum, Linn. Sp. Pl. 564 = Nepeta ucranica.
sibiricum, Willd. ex Benth. in Linnaea, xi. (1837) 347 = Nepeta parviflora.
siculum, Guss. Fl. Sic. Syn. ii. 837 = T. Scorodonia.
Sieberi, *Celak. in Bot. Centralb.* xiv. (1883) 187.—Reg. Mediterr.
sinaicum, *Boiss. Diagn.* Ser. I. xii. 91.—Arab.
sinuatum, *Celak. in Bot. Centralb.* xiv. (1883) 218.—Kurdistan.
smyrneum, Boiss. Diagn. Ser. I. v. 42 = Kotschyanum.
Socinianum, *Boiss. Fl. Orient.* iv. 818.—Syria.
Sordium, Steud. Nom. ed. II. ii. 675, in syn., err. typ. = T. Scordium.
speciosum, Hill, Hort. Kew. 253 = canadense.
spicigerum, *C. Koch, in Linnaea*, xxi. (1848) 706.—Oriens.
spinosum, *Linn. Sp. Pl.* 566.—Reg. Mediterr.
staechadifolium, Pomel, Nouv. Mat. Fl. Atl. 113 = T. Polium.
Stocksianum, *Boiss. Diagn.* Ser. II. iv. 58.—Beluchist.
stoloniferum, *Roxb. Hort. Beng.* 44; *Fl. Ind.* iii. 3.—As. trop.
subspinosum, *Pourr. ex Willd. Enum. Hort. Berol.* 596.—Ins. Balear.
subtriphyllum, Lag. Gen. et Sp. Nov. 17 = T. Polium.
supinum, Linn. Sp. Pl. 566 = montanum.
sylvestre, Lam. Fl. Fr. ii. 412 = T. Scorodonia.
syspirense, C. Koch, in Linnaea, xxi. (1848) 704 = T. Chamaedrys.
Taylori, *Boiss. Diagn.* Ser. I. vii. 61.—Persia.
Teuthrion, Schreb. Pl. Vert. Unilab. 46 = T. Polium.
Thea, *Lour. Fl. Cochinch.* 363.—China.
thymifolium, *Schreb. Pl. Vert. Unilab.* 50.—Hispan.
thymoides, Pomel, Nouv. Mat. Fl. Atl. 112 = T. Polium.
tomentosum, *Heyne, in Wall. Cat.* n. 2025.—Ind. or.
tomentosum, Lam. Fl. Fr. ii. 407; Vill. Hist. Pl. Dauph. ii. 352 = T. Polium.
tomentosum, Moench, Meth. 382 = fruticans.
tomentosum, Wight, Ic. t. 1458 = Wightii.
trifidum, Retz. Obs. i. 21 = africanum.
trifidum, Schlecht. in Linnaea, xx. (1847) 609 = sessiliflorum.
trifidum, Wendl. Bot. Beobacht. 50 = africanum?
trifoliatum, Willd. Sp. Pl. iii. 36 = T. Polium.
tripartitum, Meyen, Reise, i. 406 = nudicaule.
undulatum, *Lour. Fl. Cochinch.* 362.—Cochinch.
undulatum, Rafin. Fl. Ludov. 42 (Quid?).—Am. bor.
valentinum, Schreb. Pl. Vert. Unilab. 46 = T. Polium.
veronicaefolium, Salisb. Prod. 76 = T. Chamaedrys.
veronicoides, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 388.—Japon.
verticillatum, *Cav. Ic.* ii. 77. t. 198.—Hispan.
vesicarium, Mill. Gard. Dict. ed. VIII. n. 17 = inflatum.
vestitum, Wall. ex Steud. Nom. ed. II. ii. 675 = quadrifarium.
villosum, Forst. f. Prod. 43 = inflatum.
vincentinum, *Rouy, in La Natural.* ii. (1882) (*Matér. Rév. Fl. Portug.* 9).—Lusit.
virescens, Pomel, Nouv. Mat. Fl. Atl. 115 = T. Polium.
virginicum, Linn. Sp. Pl. 564 = canadense.
virginicum, Thunb. Fl. Jap. 244 = japonicum.
viscidum, Blume, Bijdr. 827 = stoloniferum.
Wallichianum, Benth. in Wall. Pl. As. Rar. ii. 19 = Achyrospermum Wallichianum.
Webbianum, *Boiss. Elench.* 78.—Hispan.
Wightii, *Hook. f. Fl. Brit. Ind.* iv. 701.—Ind. or.

TEURIUM, Meerb. Afbeeld. t. 45 (1775), sphalm. = praec.

TEXIERA, Jaub. & Spach, Illustr. i. 1. t. 1 (1842). *CRUCIFERAE*, Benth. & Hook. f. i. 96.
GLASTARIA, Boiss. in Ann. Sc. Nat. Sér. II. xvi. (1841) 382.
glastifolia, *Jaub. & Spach, Illustr.* i. 1. t. 1.—As. Min.

TEXTORIA, Miq. Ann. Mus. Bot. Lugd. Bat. i. 12 (1863) = **Dendropanax**, Decne. (Araliac.).
japonica, Miq. l. c. = D. japonicum.

TEYSMANNIA, Miq. in Versl. en Med. Kon. Wetensch. Akad. vi. (1857) 193 = **Pottsia**, Hook. & Arn. (Apocyn.).
Hookeriana, Miq. l. c. 194; et Fl. Ind. Bat. ii. 456 = P. cantonensis.
laxiflora, Miq. ll. cc. 194; 455 = P. cantonensis.

TEYSMANNIA, Reichb. f. & Zoll. in Linnaea, xxviii. (1856) 657 (Teyssmania). *PALMAE*, Benth. & Hook. f. iii. 924.
altifrons, *Reichb. f. & Zoll. l. c.*—Java.

THACLA, Spach, Hist. Vég. Phan. vii. 295 (1839) = **Caltha**, Rupp. (Ranunc.).
ficarioides, Spach, l. c. = C. natans.

THACOMBAUIA, Seem. Fl. Vit. 426 (1871). *HUMIRIACEAE*.
vitiensis, *Seem. l. c.*—Ins. Fiji.

THALAMIA, Spreng. Anleit. ii. 218 (1817) = **Phyllocladus**, Rich. (Conif.).
asplenifolia, Spreng. l. c. = P. rhomboidalis.
cupressina, Spreng. Syst. iii. 890 = Dacrydium cupressinum.

THALASIUM, Spreng. Syst. iv. Cur. Post. 30 (1827) = **Panicum**, Linn. (Gramin.).
montevidense, Spreng. l. c. = P. racemosum?

THALASSIA, Banks, in Kon. & Sims, Ann. Bot. ii. (1805) 96. *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 455.
SCHIZOTHECA, Ehrenb. in Abh. Akad. Berl. i. (1832) 429.
antarctica, F. Muell. ex Aschers. in Linnaea, xxxv. (1867–68) 159 = Cymodocea antarctica.
bullata, Kunth, Enum. Pl. iii. 120 = Zostera bullata.
ciliata, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 18 = Halophila ovata.
ciliata, Kon. in Kon. & Sims, Ann. Bot. ii. (1805) 97 = Cymodocea ciliata.
Hemprichii, *Aschers. in Neumayer, Anleit. Wiss.* 361.—Litora As. trop.
stipulacea, Kon. in Kon. & Sims, Ann. Bot. ii. (1805) 97 = Halophila ovata.
testudinum, *Kon. l. c.* 96.—Mare Caribaeum.
vaginata, Decne. ex Kunth, Enum. Pl. iii. 120 = Halophila ovata.
vitrariorum, Pers. Syn. ii. 563 = testudinum.

THALESIA, Rafin. in Am. Monthly Mag. (1818) 267 = **Aphyllon**, Mitch. (Orobanch.).
uniflora, Rafin. l. c. = A. uniflorum.

THALIA, Linn. Gen. ed. I. 362 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 650.
MALACARYA, Rafin. in Am. Monthly Mag. (1819) 190.
PERONIA, [Delar. in] Red. Lil. vi. t. 342 (1812).
SPIROSTALIS, Rafin. Fl. Tellur. iv. 51 (1836).
altissima, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 917 = geniculata.
angustifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 256.—Cuba.
bicolor, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 146 = Maranta bicolor.
brachystachys, C. Koch, l. c. = Phrynium brachystachyum.
cannaeformis, Forst. f. Prod. 1 = Clinogyne dichotoma.
colorata, Vell. Fl. Flum. 5; i. t. 16 = Maranta bicolor.
composita, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 146 = Maranta Riedeliana.
concolor, Horan. Prod. Monog. Scitam. 11, sphalm. = *colorata*.
dealbata, *Fras. Thal. dealbata*, t. 1.—Am. bor.
dealbata, Hort. ex Link, Jahrb. i. III. (1820) 21 = Clinogyne dichotoma.
dichotoma, Roxb. ex Link, l. c. = Clinogyne dichotoma.
divaricata, Chapm. Fl. S. U. St. 465.—Am. bor.
erecta, *Vell. Fl. Flum.* 4; i. t. 17.—Bras.
farinosa, Brongn. in Ann. Sc. Nat. Sér. IV. xii. (1860) 195, in obs. = Ischnosiphon leucophaeus.
geniculata, *Linn. Sp. Pl.* 1193.—Am. bor.
glumacea, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 163 = Maranta glumacea.
hexantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 24. t. 132 = Maranta hexantha.

THALIA:—
latifolia, Link, Jahrb. i. III. (1820) 22 = Calathea latifolia.
leptostachya, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 258 = Maranta leptostachya.
Linkiana, Steudner, in Ind. Sem. Hort. Berol. (1857) App. 10 = Maranta Riedeliana.
Luschnathiana, C. Koch, ex Horan. Prod. Monog. Scitam. 11 = Ctenanthe Luschnathiana.
lutea, Steudner, in Ind. Sem. Hort. Berol. (1857) App. 10 = Stromanthe lutea.
marantifolia, *Vell. Fl. Flum.* 5; i. t. 15.—Bras.
multiflora, *Horkel, ex Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 76.—Bras.
nemorosa, Willd. ex Link, Jahrb. i. III. (1820) 22 = Heliconia psittacorum.
Pavonii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 77.—Peruv.
pilosa, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 146 = Stromanthe Tonckat.
pubescens, Willd. ex Link, Jahrb. i. III. (1820) 22 = Stromanthe Tonckat.
racemosa, Humb. ex Link, l. c. 21 = Stromanthe Tonckat.
rotundifolia, C. Koch, ex Horan. Prod. Monog. Scitam. 11 = Calathea rotundifolia.
sanguinea, Lem. Jard. Fleur. iii. (1852–53) t. 268 = Stromanthe sanguinea.
Selloi, C. Koch, in Steudner, Ind. Sem. Hort. Berol. (1857) App. 10; et in Berl. Allg. Gartenz. xxv. (1857) 258 = Ctenanthe Luschnathiana.
setosa, C. Koch, ll. cc. 9; 258 = Myrosma setosum.
spectabilis, Lem. Jard. Fleur. iv. (1854) t. 401; Misc. 63 = Stromanthe sanguinea.
Steudneri, C. Koch, ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 65 = Ctenanthe pilosa.
sylvestris, *Willd. ex Schult. Mant.* i. 10.—Am. austr.
tuberosa, Vell. Fl. Flum. 4; i. t. 18 = Calathea tuberosa.
unilateralis, Poepp. & Endl. Nov. Gen. et Sp. ii. 24. t. 133 = Myrosma unilaterale.

THALIANTHUS, Klotzsch, in Schomb. Reise Brit. Gui. iii. 1125 (Faun. Fl. Brit. Gui.) (1846) = **Myrosma**, Linn. f. (Scitam.).
macropus, Klotzsch, l. c. = M. cannaefolium.
macropus, Koern. in Bull. Soc. Nat. Mosc. (1862) I. 66 = Maranta Moritziana.

THALICTRUM, Tourn. ex Linn. Gen. ed. I. 164 (1737). *RANUNCULACEAE*, Benth. & Hook. f. i. 4.
PHYSOCARPUM, Bercht. & Presl, Rostl. i. Ranunc. 14 (1823).
PRATICOLA, Ehrh. Phyt. n. 15 (1786); ex ej. Beitr. iv. 146 (1789).
RUPRECHTIA, Opiz, Seznam, 86 (1852).
TRIPTERIUM, Bercht. & Presl, Rostl. i. Ranunc. 14 (1823).
abbreviatum, Liévre, in Oestr. Bot. Zeitschr. xxiii. (1873) 255 = angustifolium?
abortivum, Jord. Diagn. i. 39 = minus.
acaule, Jacquem. Voy. Bot. 3 = alpinum.
acteaefolium, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 178.—Japon.
acuminatum, Spreng. Pugill. ii. 68 = minus.
acutilobum, DC. Syst. i. 177 = foetidum.
adianthifolium, Bess. ex Eichw. Skizze, 182 = minus.
affine, Jord. Diagn. i. 42, partim = minus.
affine, Ledeb. Fl. Ross. i. 10 = simplex.
agreste, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 371 = minus.
alatum, Dulac, Fl. Hautes-Pyr. 212 = aquilegifolium.
alchemillifolium, Wall. ex D. Don, Prod. Fl. Nep. 193 = rotundifolium.
alpestre, Gaud. Fl. Helvet. iii. 502 = alpinum.
alpestre, Schur, Enum. Pl. Transs. 7 = aquilegifolium.
alpicolum, Jord. Diagn. i. 43 = simplex.
alpinum, *Linn. Sp. Pl.* 545.—Reg. bor. et arct.
altaicum, Hort. Dorpat. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 248 (Quid?).—Rossia.
altaicum, Hort. Monsp. ex Lecoy. l. c. = minus.
altissimum, Thomas, ex De Massas, in Ann. Sc. Nat. Sér. II. ix. (1838) 369 = flavum.
altissimum, Wender. in Flora, ix. (1826) 358 = glaucum.
ambigens, Jord. Diagn. i. 38 = minus.
ambiguum, Schleich. Cat. Pl. Helvet. ed. IV. 34 = minus.

THALICTRUM:—
ammophilum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 21 = angustifolium.
amoenum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 248 = minus.
amurense, Maxim. Prim. Fl. Amur. 15 = simplex.
anemonoides, Michx. Fl. Bor. Am. i. 322 = Anemone thalictroides.
angulatum, Pers. Syn. ii. 101, sphalm. = seq.
angulosum, Desf. Tabl. ed. I. 123 = glaucum.
angulosum, Hort. Par. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 251 (Quid?).—Hab.?
angustatum, Schrad. Hort. Gotting. ex Lecoy. l. c. = minus.
angustatum, Weinm. ex Lecoy. l. c. 250 = flavum.
angustifolium, *Linn. Sp. Pl.* 546.—Europ.; As. Min.
angustifolium, Pollich, Hist. Pl. Palat. ii. 101 = simplex.
angustissimum, Crantz, Stirp. Austr. ed. I. fasc. ii. 76; ed. II. 105 = angustifolium.
anonymum, Wallr. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 252 = flavum.
apiculatum, *Loud. Encyc. Pl.* 647 (Quid?).—Hab.?
apiculatum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 252 = minus.
apiifolium, Schrad. l. c. = minus.
appendiculatum, C. A. Mey. in Ledeb. Fl. Alt. ii. 356 = minus.
aquilegifolium, *Linn. Sp. Pl.* 547.—Europ.; As. bor.
arabicum, Botta, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 253 = minus.
aristatum, Willd. ex Lecoy. l. c. = revolutum.
armenum, C. Koch, ex Lecoy. l. c. = minus.
Arnaudiae, Jord. Diagn. i. 31 = minus.
arrigens, Jord. l. c. 35 = minus.
athesianum, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 288 = minus.
atropurpureum, Jacq. Hort. Vindob. iii. 34 = aquilegifolium.
auriculatum, Bess. ex Eichw. Skizze, 182 = simplex *vel* flavum.
auriculatum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 254 = simplex.
aurigeranum, Baill. & Timb. in Bull. Soc. Bot. Fr. xi. (1864) 63 = minus.
axilliflorum, Schur, Enum. Pl. Transs. 10 = simplex.
baicalense, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 85.—Sibir. baical.
banchianum, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 289 = minus.
Barthii, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 14 = minus.
Batula, Buch.-Ham. ex DC. Syst. i. 185 = rotundifolium.
Bauhini, Crantz, Stirp. Austr. ed. I. fasc. ii. 70; ed. II. 105 = angustifolium, simplex.
Bauhini, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 22 = angustifolium.
Bauhinianum, Wallr. Sched. Crit. 264 = angustifolium.
belgicum, Jord. Diagn. i. 51 = flavum.
Billotii, F. Schultz, ex Jord. l. c. 36 = minus.
Bondonii, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 121 = minus.
brachycarpum, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* vi. (1883–84) 77.—Mont. Pyren.
bracteatum, Roxb. Hort. Beng. 43; Fl. Ind. ii. 671 = Clematis Cadmia.
bracteifilum, Bertol. in Mem. Acad. Sc. Bolog. Ser. II. iii. (1863) 422 = alpinum.
bresimense, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 55 = minus.
brevepubens, Jord. Diagn. i. 33 = minus.
budense ×, Simk. in Magyar Növényt. Lapok, i. (1877) 149.—Europ.
caffrum, Eckl. & Zeyh. Enum. 1 = minus.
calabricum, *Spreng. Pugill.* i. 37.—Italia.
calabrum, Tenore, Fl. Nap. iv. 333, sphalm. = praec.
calcareum, Jord. Obs. Pl. Crit. v. (1847) 9 = minus.
calcareum, Kitt. Taschenb. ed. II. 752 (Quid?).—Europ.
callistemon, Wender. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 257 (Quid?).—Europ.?
canadense, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. bor.
capillare, Reichb. Fl. Germ. Excurs. 729 = minus.
capitatum, Jord. Diagn. i. 51 = flavum.
carolinianum, Bosc, ex DC. Syst. i. 174 = Cornuti, dasycarpum.

THALICTRUM:—
carolinianum, Walt. Fl. Carol. 157 = Anemone Walteri.
Chelidonii, *DC. Prod.* i. 11.—Reg. Himal.
Chelidonii, Hook. f. & Thoms. Fl. Ind. i. 13, partim = cultratum.
chymocarpum, Walp. Ann. ii. 5 = rhynchocarpum.
cinereum, Desf. Tabl. ed. II. 146 = glaucum?
clavatum, *DC. Syst.* i. 171.—Am. bor.
clavatum, Hook. Fl. Bor. Am. i. 2 = sparsiflorum.
Clesianum, Liévre, in Oestr. Bot. Zeitschr. xxiii. (1873) 119 = simplex.
clypeatum, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. ii. (1874) 446 = minus.
collinum, Wallr. Sched. Crit. 259 = minus.
columnare, *Costa, Supl.* (1877) 2 (sp. dub.).—Hispan.
commutatum, C. A. Mey. ex Ledeb. Fl. Ross. i. 12 = flavum?
comosum, Fries, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 260 = minus.
complex, Regel, in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 225, sphalm. = simplex.
concinnum, Willd. Enum. Hort. Berol. 584 = minus.
confertiflorum, C. A. Mey. ex Trautv. in Act. Hort. Petrop. viii. (1883) 29 = simplex.
confertum, Moench, Meth. 297 (Quid?).—Hab.?
contortum, Linn. Sp. Pl. 547 = aquilegifolium.
controversum, *Schimp. & Spenn. in Spenn. Fl. Friburg.* iii. 992.—Europ.
convexum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 262 = minus.
copticum, Jord. ex Lecoy. l. c. (Quid?).—Hab.?
cordifolium, Willd. ex Spreng. Syst. ii. 670 = podocarpum.
Cornuti, *Linn. Sp. Pl.* 545.—Am. bor.
Cornuti, Poir. Encyc. v. 319 = minus, aquilegifolium.
Cornuti, Torr. & Gray, Fl. N. Am. i. 38, partim = corynellum.
corymbosum, Liévre, in Oestr. Bot. Zeitschr. xxiii. (1873) 256 = angustifolium.
corynellum, *DC. Syst.* i. 172.—Am. bor.
Costae, *Timb. ex Debeaux, Rech. Fl. Pyrén. Orient.* i. 14.—Mont. Pyren.
crenatum, Desf. Tabl. ed. II. 146 = glaucum.
cruciatum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = minus.
Csatoi, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 15 = minus.
cultratum, *Wall. Pl. As. Rar.* ii. 26.—Reg. Himal.
cuneatum, Hort. ex Reichb. Fl. Germ. Excurs. 730 = *crenatum*.
cuneatum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 14, in textu = minus.
cuspidatum, Koch, ex Ind. Sem. Hort. Panorm. (1874) = minus.
cynapiifolium, *Fisch. ex G. Don, Gen. Syst.* i. 14.—Sibir.
cysticarpum, Wall. Pl. As. Rar. ii. 26 = Chelidonii.
Dahurici, Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 265, sphalm. = *davuricum*.
Dalingo, Buch.-Ham. ex DC. Syst. i. 175 = foliolosum.
Dalzellii, *Hook. Ic. Pl.* t. 868.—Ind. or.
dasycarpum, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* viii. 72.—Am. bor.
davuricum, Fisch. ex Ledeb. Fl. Ross. i. 11 = minus.
debile, *Buckl. in Am. Journ. Sc.* Ser. I. xlv. (1843) 175.—Am. bor. occ.
Delarbrei, Lamotte, Prod. Fl. Centr. Fr. i. 34 = minus.
densiflorum, H. B. & K. ex DC. Syst. i. 184 = glaucum.
depauperatum, Dum. in Bull. Soc. Bot. Belg. viii. (1868) 357 = minus.
despiciens, Schrad. ex Lecoy. l. c. xxiv. (1885) 266 = minus.
dichotomum, Hort. ex Steud. Nom. ed. II. ii. 266 = squarrosum.
diffusum, Schrad. ex Sweet, Hort. Brit. ed. I. 2 = minus.
dioicum, Hook. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = corynellum.
dioicum, *Linn. Sp. Pl.* 545.—Am. bor.
discolor, Willd. Enum. Hort. Berol. Suppl. 40 = glaucum.
dispermum, Hort. Dorpat. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 267 (Quid?).—Hab.?
distans, Laest. ex Nym. Consp. 6 = minus?

THALICTRUM:—
divaricatum, Herb. Berol. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = corynellum.
divaricatum, Spreng. Ind. Sem. Hort. Hal. (1807) 61 = angustifolium.
divergens, Link, Enum. Hort. Berol. ii. 92 = corynellum.
dubium, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 268 = minus.
dubium, Schum. Enum. Pl. Saell. i. 167 = simplex.
dumulosum, Jord. Diagn. i. 36 = minus.
dunense, Dum. Fl. Belg. Prod. 126 = minus.
elatum, Jacq. Hort. Vindob. iii. 49 = minus.
elegans, Jord. Obs. Pl. Crit. v. (1847) 7 = minus.
elegans, *Wall. Cat.* n. 4728.—Reg. Himal.
elongatum, Jord. ex Nym. Consp. 6 = simplex.
eminens, Jord. Obs. Pl. Crit. v. (1847) 4 = minus.
erectum, Falc. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 270 = pauciflorum, *Royle*.
exaltatum, Gaud. Fl. Helvet. iii. 515 = flavum?
exaltatum, C. A. Mey. in Ledeb. Fl. Alt. ii. 352 = simplex.
eximium, Desf. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 276 (Quid?).—Hab.?
expansum, Jord. Obs. Pl. Crit. v. (1847) 6 = minus.
falcatum, Hort. Ticin. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 271 (Quid?).—Hab.?
Falconeri, *Lecoy. l. c.* 192.—Reg. Himal.
fallacinum, Gren. ex Lecoy. l. c. 271 = minus.
Fendleri, C. L. Anders. Cat. Pl. Nevad. 117 = polycarpum.
Fendleri, *Engelm. ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 5.—Am. bor. occ.
filamentosum, *Maxim. Prim. Fl. Amur.* 13.—Reg. Amur.
filipes, Torr. & Gray, Fl. N. Am. i. 38 = clavatum.
firmum, Hort. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 272 (Quid?).—Hab.?
Fischeri, Ledeb. ex Lecoy. l. c. = minus.
fistulosum, Jacquem. ex Lecoy. l. c. = cultratum.
flaccidum, Schleich. Cat. Pl. Helvet. ed. IV. 35 = flavum?
flavo-virens, Ledeb. Fl. Ross. i. 9 = minus.
flavum, Cav. Desc. 508 = glaucum.
flavum, *Linn. Sp. Pl.* 546.—Europ.; As. bor.
flavum, Pall. Reise, iii. 519 = simplex.
flavum, Thunb. Fl. Jap. 241 = minus.
flexicaule, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 274 = minus.
flexuosum, *Bernh. Cat.* (1815); *ex Reichb. Fl. Germ. Excurs.* 728 = minus.
floribundum, Schrad. Ind. Sem. Hort. Gotting. (1832).—Cf. Linnaea, viii. (1833) Litt. 27 = minus.
foeniculaceum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 76.—China.
foetidum, Gouan, Hort. Monsp. 263 = minus.
foetidum, *Linn. Sp. Pl.* 545.—Europ.; Sibir.
foliolosum, *DC. Syst.* i. 175.—Reg. Himal.
foliolosum, Hook. f. & Thoms. Fl. Ind. i. 14, partim = Falconeri.
formosum, Steud. Nom. ed. II. ii. 676 = aquilegifolium.
Fortunei, *S. Moore, in Journ. Bot.* xvi. (1878) 130.—China.
Friesii, Rupr. in Beitr. Pfl. Russ. Reich. ii. 17 = minus.
Friesii, Wolfner, in Lotos, v. (1855) 115, 140 = minus?
frigidum, Arv. Ess. Pl. Dauph. 11 = minus.
frutetorum, Jord. Diagn. i. 33 = minus.
fulgidum, Gren. ex Nym. Consp. 5 = angustifolium.
Galeottii, *Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 131.—Mexic.
galiiforme, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 19 = angustifolium.
galioides, Andrz. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 251 (Quid?).—Podol.
galioides, Pers. Syn. ii. 101 = simplex.
galioides, Presl, ex Nym. Consp. 5 = angustifolium.
Gerberi, [Willk.] Del. Sem. Hort. Dorpat. (1880) 5 (Quid?).—Hab.?
gibbosum, *Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 132.—Mexic.
glaucescens, Willd. Enum. Hort. Berol. Suppl. 40 = minus.
glaucifolium, Wender. Ind. Sem. Hort. Mart. App. n. 12 = glaucum.
glaucophyllum, Wender. l. c. n. 13 = glaucum.

THALICTRUM:—
glaucovirens, Andrz. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 277 = flavum.
glaucum, *Desf. Tabl.* ed. I. 123; ed. II. 146.—Europ. austr.
glaucum, Sieber, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = corynellum.
globiflorum, Ledeb. Fl. Ross. i. 9 = minus.
glomerulosum, Fries, Mant. iii. 49 = minus.
glyphocarpum, Wight & Arn. Prod. 2 = javanicum.
Godroni, Jord. Diagn. i. 32 = minus.
gracile, C. A. Mey. in Ledeb. Fl. Alt. ii. 348 = minus, foetidum.
gracile, Schur, ex Lecoy. in Bull. Soc. Bot. Belg. xxix. (1885) 273 = flavum.
Grenieri, Loret, in Bull. Soc. Bot. Fr. vi. (1859) 16, in textu = minus.
Hernandezii, *Tausch, in Presl, Rel. Haenk.* ii. 69.—Mexic.
heterophyllum, Lej. Rev. Fl. Spa, 109 = flavum.
heterophyllum, Nutt. ex Torr. Bot. U. St. Expl. Exped. ii. 213 = dioicum.
heterophyllum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 20 = angustifolium?
heterophyllum, Turcz. ex Ledeb. Fl. Ross. i. 727 = simplex.
hircinum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 280 = minus.
hirsutum, Mert. ex Lecoy. l. c. 264 = corynellum.
hirsutum, Willd. l. c. 280 = foetidum.
hirtellum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 13 = minus.
hirtum, Hort. Dorpat. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 280 (Quid?).—Hab.?
Hornemanni, Schrad. ex Lecoy. l. c. = minus.
humile, Schleich. ex Mert. & Koch, Deutschl. Fl. iv. 128 = minus.
hybridum, Jord. ex Nym. Consp. 5 = flavum.
hypoleucum, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 178 = minus.
bericum, Hort. Dorpat. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 281 (Quid?).—Hab.?
incisum, Schrad. ex Lecoy. l. c. = minus.
inclinatum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 14 = minus.
iodosteum ×, Borb.—Cf. Oestr. Bot. Zeitschr. xxix. (1879) 410.—Europ.
isopyroides, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 346.—Oriens; As. bor.
Jacquinianum, Koch, Syn. Fl. Germ. ed. II. 5 = minus.
japonicum, Thunb. in Trans. Linn. Soc. ii. (1794) 337 = Coptis brachypetala.
javanicum, *Blume, Bijdr.* 2.—Ind. or.; Java.
Jordani, F. Schultz, ex Jord. Obs. Pl. Crit. v. (1847) 12 = minus, simplex.
Jordanianum, F. Schultz, Fl. Pfalz, 5 = praec.?
kamtschaticum, Nyland. Spicil. Pl. Fenn. Cent. ii. 9, in obs. = minus.
kemense, Fries, Fl. Hall. i. 94 = minus.
Kochii, Fries, Mant. iii. 46 = minus.
lacerostipellatum, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854) App. 13 = simplex.
laetum, Jord. Diagn. i. 44 = simplex.
laevigatum, Michx. Fl. Bor. Am. i. 322 = dioicum.
Laggeri, Jord. Diagn. i. 26 = minus.
lanatum, *Lecoy. in Bull. Soc. Bot. Belg.* xvi. (1877) 226.—Mexic.
laserpitiifolium, Willd. Enum. Hort. Berol. Suppl. 40 = angustifolium?
lasiostylum, Presl, Rel. Haenk. ii. 70 = Hernandezii.
lathyridifolium, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 285 = angustifolium.
latifolium, Tausch, ex Steud. Nom. ed. II. ii. 676 = flavum.
laxiflorum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 93 = aquilegifolium.
Ledebourianum, C. A. Mey. ex Rupr. Fl. Cauc. 4, in textu = minus.
leiocarpum, Fries.—Cf. Linnaea, xxix. (1857–58) 731 = baicalense.
leptophyllum, Nyland. Spicil. Pl. Fenn. Cent. ii. 8 = minus.
leucostemon, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854) App. 13 = corynellum.
leucostylum, Link, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 285 = corynellum.
Leyii, Loehr, in Flora, xxv. (1842) 107 = simplex.

THALICTRUM:—
lividum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 286 = angustifolium.
lobatum, Fisch. Cat. Hort. Dresd. ex Lecoy. l. c. 286 (Quid?).—Hab.?
longepedunculatum, Hochst. ex Steud. Nom. ed. II. ii. 676 = rhynchocarpum.
longifolium, Krock. Fl. Siles. ii. 1. 242 = angustifolium.
longistylum, *DC. Syst.* i. 171.—Am. austr.
lucidum, Guenth. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 251 (Quid?).—Siles.
lucidum, Linn. Sp. Pl. 546 = angustifolium, minus.
lumacense, Lièvre, in Oestr. Bot. Zeitschr. xxii. (1872) 287 = minus.
macilentum, Jord. Diagn. i. 31 = minus.
macranthum, Hort. Choix Pl. Liége (1880) 7; ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 288 (Quid?).—Hab.?
macrocarpum, *Gren. in Acad. Sc. Besanç. Séances Publ.* (1838) 117.—Mont. Pyren.
macrostigma, Edgew. in Trans. Linn. Soc. xx. (1846) 26 = pauciflorum.
macrostigma, Herb. Kew. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 333 = virgatum.
majus, Dum. ex Ind. Sem. Hort. Panorm. (1880) 32 (Quid?).—Hab.?
majus, Gren. in Act. Linn. Soc. Bord. viii. (1836) 2. t. 9 = macrocarpum.
majus, Jacq. Fl. Austr. v. 9. t. 430 = minus.
majus, Tenore, Syll. 265 = aquilegifolium.
marginatum, Royle, Illustr. Bot. Himal. 51 = alpinum.
maritimum, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 221 = flavum.
maritimum, Willd. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 290 = minus.
Maxwellii, Royle, Illustr. Bot. Himal. 52 = minus.
meanense, Lièvre, in Oestr. Bot. Zeitschr. xxii. (1872) 123 = minus.
medianum, Jord. Diagn. i. 48 = minus.
mediterraneum, Jord. l. c. 52 = angustifolium.
medium, Jacq. Hort. Vindob. iii. 50 = minus.
medium, Saut. Versuch, 44 = flavum.
megacarpum, Torr. in Frém. Rep. 87 = polycarpum?
menthosma, Stocks, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 291 = reniforme.
mexicanum, *DC. Syst.* i. 187 (sp. dub.).—Mexic.
mexicanum, Hort. Berol. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 292 = Hernandezii, minus.
micranthum, Dougl. ex Lecoy. l. c. = minus.
microcarpon, Loud. Hort. Brit. Suppl. ii. 676, nomen = angustifolium.
microphyllum, Anders. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 292 = minus.
microphyllum, Royle, Illustr. Bot. Himal. 51 = alpinum.
micropodum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 370 = minus.
minus, *Linn. Sp. Pl.* 546.—Europ.; As. et Afr. bor.
minus, Oed. Fl. Dan. t. 244 = simplex.
minutiflorum, C. A. Mey. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 295 = angustifolium.
minutissimum, Timb. & Gant. in Mém. Acad. Toul. Ser. VII. vii. (1875) 472 = minus.
montanum, *Schimp. & Spenn. in Spenn. Fl. Friburg.* iii. 993 (an seq.?) —Europ.
montanum, Wallr. Sched. Crit. 255 = minus.
monticolum, Jord. Diagn. i. 24 = minus.
Morisoni, C. C. Gmel. Fl. Bad. iv. 422 = flavum?
mucronatum, Ledeb. Fl. Ross. i. 8 = minus.
multiflorum, Loret, in Bull. Soc. Bot. Fr. vi. (1859) 17 = minus.
nanum, Schleich. Cat. Pl. Helvet. ed. IV. 35 = minus.
neglectum, Wallr. in Linnaea, xiv. (1840) 590 (Beitr. Fl. Hercyn. 220) = minus.
nemoralis, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 297 = minus.
Nestleri, F. Schultz, Fl. Pfalz, 4 = simplex.
neurocarpum, Royle, Illustr. Bot. Himal. 51 = reniforme.
nigrescens, Hort. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 297 = angustifolium.
nigricans, DC. Syst. i. 182; Prod. i. 14 = angustifolium.
nigricans, Jacq. Fl. Austr. v. t. 421 = flavum.
nigricans, Reichb. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 251 (Quid?).—Europ.
nitidulum, Jord. Obs. Pl. Crit. v. (1847) 17 = minus, simplex.
niveum, Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 298 = aquilegifolium.
nothum, Jord. Diagn. i. 39 = minus.

THALICTRUM:—
nudicaule, Schwein. ex Torr. & Gray, Fl. N. Am. i. 39 = clavatum.
nutans, Gilib. Fl. Lituan. ii. 284; Desf. Tabl. ed. I. 123 = minus.
nutans, Schleich. ex Mert. & Koch, Deutschl. Fl. iv. 129, 137 = flavum.
obscuratum, Jord. Diagn. i. 30 = minus.
obscurum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 299 = minus.
obtusatum, Schrad. ex Lecoy. l. c. = flavum.
obtusilobum, Tausch, Ind. Sem. Hort. Panorm. (1874) = minus.
occidentale, A. Gray, in Proc. Am. Acad. viii. (1872) 372 = dioicum.
ochotense, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 300 = minus.
ochoticum, Fisch. in Zeyh. Cat. Sem. Hort. Schwez. (1823) = minus.
odoratum, Gren. & Godr. Fl. Fr. i. 6 = minus.
olidum, Jord. Diagn. i. 22 = minus.
oligospermum, Fisch. ex Sweet, Hort. Brit. ed. I. 2 = squarrosum.
olympicum, Boiss. & Heldr. Diagn. Ser. II. i. 5 = minus.
oreites, Jord. Diagn. i. 28 = minus.
orientale, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 349.—Graec.; As. Min.
ovatum, Hort. Berol. ex C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1855) App. 15 = minus.
pallidiflorum, Hort. Monac. in Bull. Soc. Bot. Belg. xxiv. (1885) 301 = minus.
pallidum, Fisch. Del. Sem. Hort. Dorpat. 5 (Quid?). —Hab.?
pallidum, Franch. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 301 = Fortunei.
palmatum, Walt. ex Spreng. Syst. ii. 674 = Trautvetteria palmata.
paniculatum, *Bess. ex Link, Handb.* ii. 400.—Hab.?
paradoxum, Jord. Obs. Pl. Crit. v. (1847) 10 = minus.
parisiense, Jord. Diagn. i. 40 = minus.
parviflorum, Salisb. Prod. 370 = minus.
parvifolium, Bertol. in Mem. Acad. Sc. Bologn. Ser. II. v. (1865) 376 = foetidum.
parvifolium, Moench, Meth. 298 = minus.
patens, Hort. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 302 (Quid?).—Hab.?
pauciflorum, Rafin. Atl. Journ. 151 (Quid?).—Am. bor.
pauciflorum, *Royle, Illustr. Bot. Himal.* 52.—Reg. Himal.
pauciflorum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 84 = aquilegifolium.
pauciflorum, Steph. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 302 = simplex.
pauperculatum, Herm. ex DC. Syst. i. 183 = flavum.
pedunculatum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 27.—Reg. Himal.
peltatum, *DC. Prod.* i. 11.—Mexic.
persicum, Sprenger, Ind. Sem. Hort. Panorm. (1874); ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 303 = minus.
petaloideum, *Linn. Sp. Pl.* ed. II. 771.—As. bor.
petrosum, Schur, Enum. Pl. Transs. 8 = minus.
peucedanifolium, Griseb. & Schenk, in Wiegm. Archiv, xviii. 1. (1852) 312 = angustifolium.
planum, Liévre, in Oestr. Bot. Zeitschr. xxiii. (1873) 254 = angustifolium.
platycarpum, Hook. f. & Thoms. Fl. Ind. i. 13 = cultratum.
platystemon, Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 303 = aquilegifolium.
podocarpum, *H. B. & K. Nov. Gen. et Sp.* v. 38.—Am. austr.
podolicum, *Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 173.—Europ.
polycarpum, Loret, in Bull. Soc. Bot. Fr. vi. (1859) 17 = minus.
polycarpum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 288.—Am. bor. occ.
polygamum, Herb. Link, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = corynellum.
polygamum, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 172, nomen; et ex Pursh, Fl. Am. Sept. ii. 388; DC. Syst. i. 173 = corynellum, revolutum, minus.
polyphyllum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 304 = minus.
polyspermum, Arv. Ess. Pl. Dauph. 13–14 = minus.
Porphyritae, F. Schultz, in Flora, lvi. (1873) 232 = minus.
porrectum, Jord. ex Nym. Consp. 5 = flavum.

THALICTRUM:—
porrigens, Jord. Diagn. i. 48 = minus.
praecox, Jord. Obs. Pl. Crit. v. (1847) 2 = minus.
praecox, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1855) App. 15 = minus.
praecox, F. Schultz, in Pollichia, (1859) 1 = minus.
praeflorens, Jord. Diagn. i. 27 = minus.
praeruptum, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. ii. (1874) 445 = minus.
pratense, Linn. Sp. Pl. 546, in syn. = flavum.
pratense, F. Schultz, in Pollichia, xx–xxi. (1863) 102 = minus.
Princeps, Dum. in Bull. Soc. Bot. Belg. viii. (1869) 458 = flavum.
procerulum, Jord. Diagn. i. 45 = simplex.
propendens, Jord. l. c. 37 = minus.
prorepens, Jord. l. c. 50 = flavum.
pruinosum, E. Mey. in Drège, Zwei Pfl. Docum. 226 = minus.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 305.—China.
Pseudo-flavum, Schur, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 306 = flavum.
pubens, Steph. ex Lecoy. l. c. = foetidum.
puberulum, Aschenb. ex Lecoy. l. c. = Hernandezii.
pubescens, Pursh, Fl. Am. Sept. ii. 388 = corynellum, revolutum.
pubescens, Schleich. ex DC. Fl. Fr. Suppl. 633 = minus.
pubigerum, *Benth. Pl. Hartw.* 285.—Mexic.
pulchellum, Pursh, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 307 = dioicum.
punduanum, *Wall. Pl. As. Rar.* ii. 26; *Cat.* n. 3712.—Reg. Himal.
purpurascens, DC. Syst. i. 174 = dioicum.
purpurascens, Georgi, Beschr. Russ. Reich. III. iv. 1063 = minus.
purpurascens, Linn. Sp. Pl. 546 = minus, corynellum.
purpureum, Schang. in Pall. N. Nord. Beitr. vi. 42 = minus.
purpureum, Tenore, ex Steud. Nom. ed. II. ii. 676, 677 = calabricum.
pyramidale, Hort. Panorm. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 308 = minus.
pyrenaicum, Jord. Diagn. i. 29 = minus.
pyrrhocarpum, Ckodi, Ind. Sem. Hort. Valent. (1883) (Quid ?).—Hab. ?
radiatum, Royle, Illustr. Brit. Himal. 52 = saniculaeforme.
radicans, Hort. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 308 = minus.
ranunculinum, Muhl. ex Willd. Enum. Hort. Berol. 585 = Trautvetteria palmata.
rariflorum, Fries, Summa Veg. Scand. 137 = minus, simplex ?
recurvum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 309 = minus.
Regelianum, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 75 = simplex.
reniforme, *Wall. Pl. As. Rar.* ii. 26; *Cat.* n. 3716.—Reg. Himal.
repens, W. Baxt. ex Loud. Hort. Brit. Suppl. ii. 676, = minus ?
repens, C. A. Mey. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 309 = minus.
repens, Schrad. ex Lecoy. l. c. = squarrosum.
revolutum, *DC. Syst.* i. 173.—Am. bor.
revolutum, Herb. Fisch. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 264 = corynellum.
revolutum, Liévre, in Oestr. Bot. Zeitschr. xxiii. (1873) 254 = angustifolium.
rhaeticum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 76.—Europ.
rhodanense, Jord. Diagn. i. 45 = simplex.
rhynchocarpum, *Q. Dillon. & A. Rich. in Ann. Sc. Nat.* Sér. II. xiv. (1840) 262.—Afr. trop.
Richardsonii, A. Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 17, in textu = sparsiflorum.
rigidum, Fisch. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 310 = minus.
riparium, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 5 = flavum.
Rochebrunnianum, *Franch. & Sav. Enum. Pl. Jap.* ii. 264.—Japon.
rosmarinifolium, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 32 = minus.
rostellatum, *Hook. f. & Thoms. Fl. Ind.* i. 15.—Reg. Himal.
rotundifolium, *DC. Syst.* i. 185.—Reg. Himal.

THALICTRUM:—
rotundifolium, Laest. ex Nym. Consp. 6 = minus.
rotundum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 311 = minus.
rubellum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 177 = aquilegifolium.
rufinerve, Lej. & Court. Comp. Fl. Belg. ii. 207 = flavum.
rufum, *Lecoy. in Bull. Soc. Bot. Belg.* xvi. (1877) 227.—Ind. or.
rugosum, Ait. Hort. Kew. ed. I. ii. 262 = flavum, glaucum, corynellum.
rugosum, Poir. Encyc. v. 317 = angustifolium.
rugosum, Pursh, Fl. Am. Sept. ii. 388 = dasycarpum ?
rupestre, Hort. Berol. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 312 = minus.
rupestre, Madden, ex Lecoy. l. c. = saniculaeforme.
rutaefolium, *Hook. f. & Thoms. Fl. Ind.* i. 14.—Reg. Himal.
ruthenicum, Schrad. Ind. Sem. Hort. Gotting. (1832).—Cf. Linnaea, viii. (1833) Litt. 27 = minus.
rutidocarpum, *DC. Syst.* i. 172.—Am. austr.
sachalinense, *Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 152.—Ins. Sachalin.
saniculaeforme, *DC. Prod.* i. 12.—Reg. Himal.
Savatieri, Foucaud, Cat. Pl. Vasc. Charente-Inf. ii. 45 = minus.
saxatile, Schleich. ex DC. Fl. Fr. Suppl. 633 = minus.
saxatile, Vill. Prosp. 50; Hist. Pl. Dauph. iii. 714 = minus.
saxicolum, Arv. Ess. Pl. Dauph. 15 = minus.
Schimperianum, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 78 = minus.
Schultzii, Jord. Diagn. i. 32 = minus.
Schweiggeri, Spreng. Syst. ii. 671 = minus.
secundum, Edgew. in Trans. Linn. Soc. xx. (1846) 26 = pauciflorum.
Sendtneri, F. Schultz, Arch. Fl. 31 = minus.
serotinum, F. Schultz, in Pollichia, xvii. (1859) 1 = minus.
sibiricum, Ledeb. Fl. Ross. i. 11 = squarrosum.
sibiricum, Linn. Sp. Pl. 546 = minus, foetidum, isopyroides.
sibiricum, Pall. Reise, i. 219 = foetidum.
silaifolium, Jord. Diagn. i. 41 = minus.
silleanum, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 285 = minus.
silvaticum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 76.—Europ.
simplex, DC. Prod. i. 15 = flavum.
simplex, *Linn. Mant.* i. 78.—Europ. et As. bor.
sinense, *Lour. Fl. Cochinch.* 346 (sp. dub.).—China.
soboliferum, Schur, in Oestr. Bot. Zeit. x. (1860) 250, in textu = flavum.
sparsiflorum, *Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 40.—Dahuria; Am. bor. occ.
speciosum, Hort. ex Poir. Encyc. v. 315 = glaucum.
speciosum, Mill. Gard. Dict. ed. VIII. n. 2 = angustifolium.
sphaerocarpon, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 317 = angustifolium.
sphaerocarpum, Lej. & Court. Comp. Fl. Belg. ii. 208 = flavum.
sphaerospermum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 317 = angustifolium.
spurium, Timeroy, ex Jord. Obs. Pl. Crit. v. (1847) 19 = flavum.
squamiferum, *Lecoy. in Bull. Soc. Bot. Belg.* xvi. (1877) 229.—Tibet.
squarrosum, *Steph. ex Willd. Sp. Pl.* ii. 1299.—Sibir.; China.
stamineum, Linn. f. Suppl. 271 = petaloideum.
stenocarpum, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. ii. (1874) 446, in obs. = minus.
stenophyllum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 19 = angustifolium ?
stipellatum, Jord. Diagn. i. 42 = minus.
stipulaceum, Hort. ex G. Don, in Loud. Hort. Brit. 229 (Quid ?).—Europ.
strictum, Laest. ex Nym. Consp. 6 = minus, simplex.
strictum, Ledeb. Fl. Ross. i. 10 = simplex.
strigillosum, *Hemsl. Diagn. Pl. Nov.* 1.—Mexic.
styloideum, Linn. f. Suppl. 271 = foetidum.
subalpinum, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 57 = minus.
subcorymbosum ×, Borb.—Cf. Oestr. Bot. Zeitschr. xxix. (1879) 410.—Europ.
subspicatum, Jord. Diagn. i. 46 = simplex.

THALICTRUM:—
subtrigynum, Reichb. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 320 = minus.
sylvaticum, Koch, in Flora, xxiv. (1841) 426 = minus.
tenue, *Franch. in Pl. Davidian.* i. (1884) 16.—China.
tenuiculum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 320 = foetidum.
tenuifolium, Gueldenst. ex Ledeb. Fl. Ross. i. 12 = angustifolium.
tenuifolium, Schwein. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 294 = minus.
tenuifolium, Sw. Summ. Veg. Scand. 72 = simplex.
thamnophilum, Jord. Diagn. i. 34 = minus.
thibeticum, *Franch. ex Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 153.—Tibet.
Thunbergii, DC. Syst. i. 183 = minus.
Timeroyi, Jord. Obs. Pl. Crit. v. (1847) 14 = simplex.
tortuosum, Jord. Diagn. i. 38 = minus.
transsilvanicum, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 84 = minus.
tridentinum, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 219 = minus.
trifidum, Liévre, l. c. xxiii. (1873) 255 = angustifolium.
trigynum, Fisch. ex Trev. Ind. Sem. Hort. Vratisl. App. ii. (1820) 3 = squarrosum.
trispermum, Fisch. ex Link, Enum. Hort. Berol. ii. 92 = squarrosum.
triternatum, *Rupr. Fl. Caucas.* 2.—Reg. Cauc.
tuberiferum, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 227.—Japon.
tuberosum, *Linn. Sp. Pl.* 545.—Europ. occ.
udum, Jord. Diagn. i. 50 = flavum.
uncinnatum, Rehm. in Spraw. Kom. Fis. Krakow, vii. (1873) 90.—Cf. Oestr. Bot. Zeitschr. xxxiii. (1883) 239 = petaloideum.
uncinulatum, *Franch. ex Lecoy. in Bull. Soc. Bot. Belg.* xxiv. (1885) 169.—Tibet.
vaccinense, Liévre, in Oestr. Bot. Zeitschr. xxii. (1872) 125 = minus.
vaginatum, Desf. Tabl. ed. II. 274 = flavum.
vaginatum, Royle, Illustr. Bot. Himal. 52 = minus, foetidum, squamiferum.
varium, Doell, Rhein. Fl. 543 (Quid ?).—Europ.
Verloti, Arv. Ess. Pl. Dauph. 13, 14 = minus.
versiflorum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 323 = minus.
versifolium, Schrad. ex Lecoy. l. c. = minus.
vesiculosum, *Lecoy. l. c.* 130.—Am. austr.
villosum, Jacquem. ex Lecoy. l. c. 323 = Falconeri, minus.
viperinum, Andrz. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 336 = minus.
virens, Wallr. ex Koch, Syn. Fl. Germ. ed. II. 1015 (Quid ?).—Europ.
virgatum, *Hook. f. & Thoms. Fl. Ind.* i. 14.—Reg. Himal.
virgatum, Schrad. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 323 = minus.
virginicum, Drège, ex Lecoy. l. c. = dasycarpum.
virgultorum, Jord. Diagn. i. 35 = minus.
vitifolium, Schlecht. ex Ledeb. Fl. Ross. i. 741 = Trautvetteria palmata.
viviparum, Stocks, ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 323 = Chelidonii.
Vrabelyi, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 11 = minus.
vulgare, Kitt. Taschenb. ed. II. 752 = minus.
vulgatum, F. Schultz, Fl. Pfalz, 3 = minus.
Wallrothi, Boenn. Prod. Fl. Monast. 162 = flavum.
Wallrothianum, Dettd. ex Lecoy. in Bull. Soc. Bot. Belg. xxiv. (1885) 273, 324 = flavum.
Walteri, Spreng. ex Steud. Nom. ed. II. ii. 677 = Anemone Walteri.
Wrightii, *A. Gray, Pl. Wright.* ii. 7.—Mexic.

THALLIANA, Steud. Nom. ed. II. ii. 677 (1841), sphalm. = **Tralliana**, Lour. (Inc. Sed.).

THALYSIA, Linn. Syst. ed. I (1735) = **Zea**, Linn. (Gramin.).

THAMINOPHYLLUM, Harv. in Harv. & Sond. Fl. Cap. iii. 155 (1864–65). *COMPOSITAE*, Benth. & Hook. f. ii. 423.
multiflorum, *Harv. l. c.*—Afr. austr.
Mundtii, *Harv. l. c.*—Afr. austr.

THAMNEA, Soland. ex R. Br. Pl. Abel. 8 (1818), nomen; et ex Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 386. t. 38. *BRUNIACEAE*, Benth. & Hook. f. i. 671.

THAMNEA:—
depressa, *Oliver, in Journ. Linn. Soc.* ix. (1867) 332.—Afr. austr.
gracilis, *Oliver, l. c.*—Afr. austr.
hirtella, *Oliver, l. c.*—Afr. austr.
laxa, Baill. Hist. Pl. iii. 388 = Tittmannia laxa.
uniflora, *Soland. ex Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 386.—Afr. austr.

THAMNIUM, Klotzsch, in Linnaea, xii. (1838) 223 = **Scyphogyne**, Brongn. (Ericac.).
puberulum, Klotzsch, l. c. = S. puberulum.

THAMNOCALAMUS, Munro, in Trans. Linn. Soc. xxvi. (1868) 33, 157 = **Arundinaria**, Michx. (Gramin.).
Falconeri, Hook. f. ex Munro, l. c. 34 = A. falcata.
Ringala, Falc. ex Munro, l. c. 157 = A. falcata.
spathiflorus, Munro, l. c. 34 = A. spathiflora.

THAMNOCHORTUS, Berg. Desc. Pl. Cap. 353 (1767). *RESTIACEAE*, Benth. & Hook. f. iii. 1034.
aemulus, Kunth, Enum. Pl. iii. 439 = imbricatus.
argenteus, *Kunth, l. c.* 432.—Afr. austr.
bromoides, Kunth, l. c. = dichotomus.
Burchellii, *Mast. in DC. Monog. Phan.* i. 322.—Afr. austr.
caricinus, *Mast. l. c.* 327.—Afr. austr.
cernuus, *Kunth, Enum. Pl.* iii. 439.—Afr. austr.
consanguineus, Kunth, l. c. 432 = dichotomus.
debilis, Spreng. ex Steud. Nom. ed. II. ii. 677 = Restio debilis.
dichotomus, *R. Br. Prod.* 244.—Afr. austr.
dichotomus, Kunth, Enum. Pl. iii. 432 = elongatus, dichotomus.
dichotomus, Spreng. Syst. i. 187 = Restio triflorus, triticeus?
distichus, *Mast. in Journ. Linn. Soc.* x. (1869) 233.—Afr. austr.
Ecklonianus, Kunth, Enum. Pl. iii. 430 = dichotomus.
elongatus, Hochst. in Flora, xxviii. (1845) 339 = striatus.
elongatus, *Mast. in Journ. Linn. Soc.* x. (1869) 226.—Afr. austr.
erectus, *Mast. l. c.* xiv. (1875) 419.—Afr. austr.
floribundus, *Kunth, Enum. Pl.* iii. 435.—Afr. austr.
fruticosus, *Berg. Desc. Pl. Cap.* 353.—Afr. austr.
giganteus, *Kunth, Enum. Pl.* iii. 435.—Afr. austr.
gracilis, *Mast. in DC. Monog. Phan.* i. 327.—Afr. austr.
imbricatus, *Mast. in Journ. Linn. Soc.* x. (1869) 231.—Afr. austr.
micans, *Kunth, Enum. Pl.* iii. 441.—Afr. austr.
modestus, Kunth, l. c. 434 = Leptocarpus modestus.
platypteris, *Kunth, l. c.* 429.—Afr. austr.
robustus, Kunth, l. c. 436 = Cannamois Cephalotes.
scariosus, R. Br. Prod. 244 = fruticosus.
scirpiformis, *Mast. in Journ. Linn. Soc.* x. (1869) 228.—Afr. austr.
scirpoides, Kunth, Enum. Pl. iii. 438 = Cannamois scirpoides.
spicigerus, *R. Br. Prod.* 244.—Afr. austr.
spicigerus, Sieber, ex Kunth, Enum. Pl. iii. 434 = dichotomus.
striatus, *Hochst. in Flora*, xxviii. (1845) 339.—Afr. austr.
strictus, Kunth, Enum. Pl. iii. 438 = Cannamois simplex.
umbellatus, *Kunth, l. c.* 440.—Afr. austr.
virgatus, Kunth, l. c. 436 = Cannamois Cephalotes.

THAMNOSERIS, Phil. f. in Anal. Univ. Chil. (1875) 191. *COMPOSITAE*.
lacerata, *Phil. f. l. c.*—Chili.

THAMNOSMA, Torr. & Frém. in Frém. [2nd] Rep. 313 (1845). *RUTACEAE*, Benth. & Hook. f. i. 288.
RUTOSMA, A. Gray, Gen. Illustr. 143. t. 155 (1849).
montana, *Torr. & Frém. in Frém.* [2nd] *Rep.* 313; *et in Duch. Rév. Bot.* ii. (1846–47) 154.—Calif.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
texana, *Torr. in Bot. Mex. Bound.* 42.—Am. bor.

THAMNUS, Klotzsch, in Linnaea, xii. (1838) 235 = **Simocheilus**, Klotzsch (Ericac.).
multiflorus, Klotzsch, l. c. = S. obovatus.

THAMNUS, Link, Enum. Hort. Berol. ii. 426 (1822) = **Tamus**, Linn. (Dioscor.).

THANATOPHORUS, Walp. Ann. ii. 213 (1851–52), sphalm. = DANATOPHORUS, Blume = **Harpullia**, Roxb. (Sapind.).

THAPSANDRA, Griseb. Spicil. Fl. Rumel. ii. 40 (1844) = **Celsia**, Linn. (Scrophul.).
cretica, Griseb. l. c. = C. cretica.

THAPSIA, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 930.
MELANOSELINUM, Hoffm. Gen. Umb. 156 (1814).
MONIZIA, Lowe, in Hook. Kew Journ. viii. (1856) 295.
altissima, *Mill. Gard. Dict.* ed. VIII. n. 6.—Ital.
apulia, Mill. l. c. n. 4 = Elaeoselinum Asclepium.
asclepiadea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = seq.
Asclepium, Linn. Sp. Pl. 261 = Elaeoselinum Asclepium.
decipiens, *Hook. f. Bot. Mag.* t. 5670.—Ins. Mader.
decussata, Lag. Gen. et Sp. Nov. 12 = garganica.
foetida, Linn. Sp. Pl. 261 = Elaeoselinum foetidum.
garganica, *Linn. Mant.* i. 57.—Reg. Mediterr.
glomerata, Nutt. Gen. Am. i. 184 = Cymopterus glomeratus.
gummifera, Spreng. Sp. Umbellif. 31 = Laserpitium gummiferum.
Laserpitii, Spreng. l. c. 32 = Prangos ferulacea.
linearilobа, Pomel, Nouv. Mat. Fl. Atl. 314 = garganica.
marocana, Pomel, l. c. in obs. = garganica.
maxima, Mill. Gard. Dict. ed. VIII. n. 2 = villosa.
melanoselina, Masf. in Anal. Soc. Esp. Hist. Nat. x. (1881) 185 = decipiens?
meoides, Guss. Fl. Sic. Prod. i. 370 = Elaeoselinum meoides.
microcarpa, Pomel, Nouv. Mat. Fl. Atl. 312 = villosa.
minor, Hort. ex Nym. Consp. 276 = villosa.
minor, Link, ex Steud. Nom. ed. I. 831 (Quid?).—Hab.?
Moniza, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 185.—Ins. Canar.
platycarpa, Pomel, Nouv. Mat. Fl. Atl. 313 = garganica.
polygama, *Desf. Fl. Atlant.* i. 261. t. 75.—Afr. bor.
polygama, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 481 = Elaeoselinum foetidum.
praealta, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 288.—Ins. Cos.
silphia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = garganica.
stenocarpa, Pomel, Nouv. Mat. Fl. Atl. 313 = garganica.
stenoptera, Pomel, l. c. 312 = villosa.
Sylphium, Viv. Fl. Lib. Spec. 17 = garganica.
tenuifolia, Lag. Gen. et Sp. Nov. 12 = Elaeoselinum Lagascae.
transtagana, *Brot. Fl. Lusit.* i. 468.—Lusitan.
trifoliata, Linn. Sp. Pl. 262 = Thaspium trifoliatum.
trifoliata, Mill. Gard. Dict. ed. VIII. n. 5 = Thaspium barbinode.
villosa, *Linn. Sp. Pl.* 261.—Europ. austr.

THAPSIUM, Walp. Rep. ii. 403 (1843), sphalm. = **Thaspium**, Nutt. (Umbell.).

THAPSUS, Rafin. Fl. Tellur. iv. 89 (1836) = **Verbascum**, Tourn. (Scrophul.).
Blattaria, Rafin. l. c. = V. Blattaria.
floccosum, Rafin. l. c. = V. pulverulentum.
Linnaei, Opiz, Seznam, 95 = V. Thapsus.
Lychnitis, Rafin. Fl. Tellur. iv. 89 = V. Lychnitis.
montanum, Rafin. l. c. = V. montanum.
mucronatum, Rafin. l. c. = V. mucronatum.
nigrum, Rafin. l. c. = V. nigrum.
pheniceum, Rafin. l. c. = V. phoeniceum.
phlomoides, Opiz, Seznam, 96 = V. phlomoides.
pulverulentum, Rafin. Fl. Tellur. iv. 89 = V. pulverulentum.
thapsoides, Rafin. l. c. = V. thapsoides.
Schraderi, Opiz, Seznam, 96 = V. Thapsus.
sinuatum, Rafin. Fl. Tellur. iv. 89 = V. sinuatum.
Willdenowii, Opiz, Seznam, 96 = V. thapsoides.

THASPIUM, Nutt. Gen. Am. i. 196 (1818). *UMBELLIFERAE*, Benth. & Hook. f. i. 913.
UPOPION, Rafin. New Fl. Am. iv. 29 (1836).
actaeifolium, Nutt. Gen. Am. i. 196 = Archangelicum Gmelini.
atropurpureum, *Nutt. l. c.*—Am. bor.
aureum, *Nutt. l. c.*—Am. bor.
barbinode, *Nutt. l. c.*—Am. bor.
cordatum, Torr. & Gray, Fl. N. Am. i. 615 = atropurpureum.
foliosum, Royle, Illustr. Bot. Himal. 40 = Pimpinella caespitosa.
macrocarpum, Herb. Bourg. ex S. Wats. Bibl. Ind. N. Am. Bot. 429 = Peucedanum macrocarpum.
montanum, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 57 = Ligusticum montanum.
pinnatifidum, A. Gray, Man. Bot. N. U. St. ed. II. 155 = Walteri.
purpureum, Walp. Am. ii. 702 = atropurpureum.
trachypleurum, A. Gray, in Proc. Acad. Sc. Philad. 1863 (1864) 63 = Cicuta trachypleura.
trifoliatum, *A. Gray, Man. Bot. N. U. St.* ed. II. 156.—Am. bor.
verticillatum, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 233 = Angelica verticillata.
Walteri, *Shuttl. ex A. Gray, Pl. Wright.* i. 79.—Am. bor.

THAUMATOCOCCUS, Benth. in Benth. & Hook. f. Gen. iii. 652 (1883). *SCITAMINEAE*, Benth. & Hook. f. l. c.
Danielli, *Benth. l. c.*—Afr. trop.

THAUMATOPHYLLUM, Schott, in Bonplandia, vii. (1859) 31. *AROIDEAE*, Benth. & Hook. f. iii. 979.
Spruceanum, *Schott, l. c.*—Bras.

THAUMAZA, Salisb. Gen. Pl. Fragm. 15 (1866) = **Eriospermum**, Jacq. (Liliac.).
paradoxa, Salisb. l. c. = E. paradoxum.

THAUMURIA, Gaudich. in Freyc. Voy. Bot. 502 (1826) = **Parietaria**, Tourn. (Urtic.).
cretica, Gaudich. l. c. = P. cretica.

THEA, Linn. Syst. ed. I (1735) = **Camellia**, Linn. (Ternstr.).
assamica, J. W. Mast. in Journ. Agric. & Hort. Soc. Ind. iii. (1844) 63 = C. Thea.
assimilis, Seem. in Trans. Linn. Soc. xxii. (1859) 349 = C. assimilis.
Bohea, Linn. Sp. Pl. ed. II. 743 = C. Thea.
Camellia, Hoffmgg. Verz. Pfl. 117 = C. japonica.
cantoniensis, Lour. Fl. Cochinch. 339 = C. Thea.
caudata, Seem. in Trans. Linn. Soc. xxii. (1859) 348 = C. caudata.
chinensis, Sims, Bot. Mag. t. 998 = C. Thea.
cochinchinensis, Lour. Fl. Cochinch. 338 = C. Thea.
euryoides, Booth, in Trans. Hort. Soc. vii. (1830) 560 = C. euryoides.
grandifolia, Salisb. Prod. 370 = C. Thea.
imperialis, Hort. ex Steud. Nom. ed. II. ii. 678; et ex Seem. in Trans. Linn. Soc. xxii. (1859) 350 = Micromeria obovata.
japonica, Baill. Hist. Pl. iv. 229 = C. Thea.
latifolia, Lodd. ex Sweet, Hort. Brit. ed. III. 91 = C. Thea.
longifolia, Nois. ex Steud. Nom. ed. I. 832 = C. Thea.
maliflora, Seem. in Trans. Linn. Soc. xxii. (1859) 346 = C. rosaeflora.
oleosa, Lour. Fl. Cochinch. 339 = C. Thea.
parvifolia, Salisb. Prod. 370 = C. Thea.
salicifolia, Seem. in Trans. Linn. Soc. xxii. (1859) 349 = C. salicifolia.
Sasangua, Nois. ex Cels, Cat. Arbr. (1817) 35 = C. Thea.
sinensis, Linn. Sp. Pl. 515 = C. Thea.
stricta, Hayne, Arzn. Gew. vii. t. 27 = C. Thea.
viridis, Linn. Sp. Pl. ed. II. 735 = C. Thea.

THEAPHYLA, Rafin. Sylva Tellur. 139 (1838) = seq.

THEAPHYLLA, Rafin. Med. Fl. ii. 267 (1830) = **Camellia**, Linn. (Ternstr.).
anamensis, Rafin. Sylva Tellur. 139 = C. Thea?
cantoniensis, Rafin. l. c. = C. Thea.
euryoides, Rafin. l. c. = C. euryoides.

THEAPHYLLA:—
lanceolata, Rafin. Sylva Tellur. 139 = Camellia lanceolata.
laxa, Rafin. l. c. = C. Thea.
oleifera, Rafin. l. c. = C. Thea.
viridis, Rafin. l. c. = C. Thea.

THEAPHYLLUM, Nutt. ex Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 605 = **Perrottetia**, H. B. & K. (Celastr.).
celastrinum, Nutt. ex Turcz. l. c. = P. sandwicensis.

THEBESIA, Neck. Elem. ii. 313 (1790) = **Knowltonia**, Salisb. (Ranunc.).

THECA, Juss. Gen. 108 (1789) = THEKA, Adans. = **Tectona**, Linn. (Verben.).

THECACORIS, A. Juss. Tent. Euph. 12. t. 1 (1824).
EUPHORBIACEAE, Benth. & Hook. f. iii. 286.
madagascariensis, *A. Juss. l. c.* 105. t. 1.—Madag.
Manniana, *Muell. Arg. in DC. Prod.* xv. ii. 246.—Afr. trop.
stenopetala, *Muell. Arg. l. c.*—Afr. trop.
trichogyne, *Muell. Arg. in Journ. Bot.* ii. (1864) 328.—Afr. trop.

THECANISIA, Rafin. New Fl. Am. ii. 38 (1836) = **Spiraea**, Linn. (Rosac.).
angustifolia, Rafin. l. c. 39 = S. angustifolia?
discolor, Rafin. l. c. = S. discolor.
lobata, Rafin. l. c. 38 = S. lobata.
palmata, Rafin. Sylva Tellur. 152 = S. palmata?
purpurea, Rafin. New Fl. Am. ii. 38 = S. lobata?
Ulmaria, Rafin. Sylva Tellur. 152 = S. Ulmaria.

THECANTHES, Wikstr. in Vet. Akad. Handl. Stockh. (1818) 271 = **Pimelea**, Banks (Thymel.).
Cornucopiae, Wikstr. l. c. = P. Cornucopiae.
punicea, Wikstr. l. c. 272 = P. punicea.

THECOCARPUS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 93. *UMBELLIFERAE*, Benth. & Hook. f. i. 903.
meifolius, *Boiss. l. c.* 94.—Persia.

THECOSTELE, Reichb. f. in Bonplandia, v. (1857) 37. *ORCHIDEAE*, Benth. & Hook. f. iii. 539.
alata, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 144 = Zollingeri.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 37.—Java.

THEIS, Salisb. ex DC. Prod. vii. 715 (1835) = **Rhododendron**, Linn. (Ericac.).

THEKA, Adans. Fam. ii. 445 (1763) = **Tectona**, Linn. (Verben.).
grandis, Lam. Illustr. ii. 111 = Tectona grandis.
ternifolia, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 185 = Tectona Hamiltoniana.

THELA, Lour. Fl. Cochinch. 119 (1790) = **Plumbago**, Tourn.
alba, Lour. l. c. = P. zeylanica.
coccinea, Lour. l. c. = P. rosea.

THELAIA, Alef. in Linnaea, xxviii. (1856) 33 = **Pyrola**, Tourn. (Ericac.).
angustifolia, Alef. l. c. 52 = P. angustifolia.
aphylla, Alef. l. c. 39 = P. aphylla.
arenaria, Alef. in Bot. Zeit. xx. (1862) 220 = P. rotundifolia.
asarifolia, Alef. in Linnaea, xxviii. (1856) 54 = P. rotundifolia.
bracteosa, Alef. l. c. 57 = P. rotundifolia.
chlorantha, Alef. l. c. 41 = P. chlorantha.
elliptica, Alef. l. c. 47 = P. elliptica.
grandiflora, Alef. l. c. 68 = P. rotundifolia.
intermedia, Alef. l. c. 65 = P. rotundifolia.
media, Alef. in Bot. Zeit. xx. (1862) 219 = P. chlorantha?
occidentalis, Alef. in Linnaea, xxviii. (1856) 36 = P. chlorantha.
rotundifolia, Alef. l. c. 60 = P. rotundifolia.
Sartorii, Alef. l. c. 50 = P. Sartorii.
spathulata, Alef. l. c. 45 = P. picta.

THELASIS, Blume, Bijdr. 385. t. 75 (1825).
ORCHIDEAE, Benth. & Hook. f. iii. 589.
EUPROBOSCIS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 371. t. 26.
OXYANTHERA, Brongn. in Duperr. Voy. Bot. 197. t. 37 (1829).
capitata, *Blume, Bijdr.* 386.—Java.
carinata, *Blume, l. c.*—Java.
carinata, Reichb. f. in Bonplandia, v. (1857) 37 = pygmaea.
contracta, *Blume, Mus. Bot. Lugd. Bat.* ii. 187.—Java.
elegans, Blume, l. c. = Phreatia elegans.
elongata, *Blume, l. c.*—Java.
obtusa, *Blume, Bijdr.* 386.—Java.
ochreata, *Lindl. in Journ. Linn. Soc.* iii. (1859) 63.—Borneo.
pygmaea, *Lindl. l. c.*—Ind. or.
triptera, *Reichb. f. in Bonplandia*, iii. (1855) 219.—Malaya.
Zollingeri, *Reichb. f. l. c.* v. (1857) 37.—Java.

THELEOPHYTON, Moq. in DC. Prod. xiii. ii. 115 (1849) = **Atriplex**, Tourn. (Chenop.).
Billardieri, Moq. l. c. 116 = A. Billardieri.

THELEPOGON, Roth, ex Roem. & Schult. Syst. ii. 46, 788 (1817); Roth, Nov. Pl. Sp. 62 (1821).
GRAMINEAE, Benth. & Hook. f. iii. 1131.
JARDINEA, Steud. in Flora, xxxiii. (1850) 229 (Jandinea).
RHINIACHNE, Hochst. ex Steud. Syn. Pl. Gram. 360, in syn. (1855).
elegans, *Roth, ex Roem. & Schult. Syst.* ii. 788; *Nov. Pl. Sp.* 62.—Afr. trop.; Ind. or.
sanguineus, Spreng. Syst. i. 299 = Andropogon Pseudograya.

THELESPERMA, Less. in Linnaea, vi. (1831) 511.
COMPOSITAE, Benth. & Hook. f. ii. 387.
COSMIDIUM, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 361.
ambiguum, *A. Gray, in Proc. Am. Acad.* xix. (1883) 16.—Am. bor. occ.
filifolium, *A. Gray, in Hook. Kew Journ.* i. (1849) 252.—Am. bor. occ.
gracile, *A. Gray, l. c.*—Am. bor. occ.
longipes, *A. Gray, Pl. Wright.* i. 109.—Texas; Mexic.
scabiosioides, *Less. in Linnaea*, vi. (1831) 511.—Bras.
simplicifolium, *A. Gray, in Hook. Kew Journ.* i. (1849) 252.—Mexic.
subnudum, *A. Gray, in Proc. Am. Acad.* x. (1874) 72.—Am. bor. occ.
subsimplicifolium, *A. Gray, in Torr. Bot. Mex. Bound.* 90.—Am. bor. occ.

THELIGONUM, Linn. Gen. ed. I. 357 (1737) = **Thelygonum**, Linn. (Urtic.).

THELIRA, Thou. Gen. Nov. Madag. 21 (1806) = **Parinarium**, Aubl. (Rosac.).

THELMATOPHACE, Godr. Fl. Lorr. ed. I. iii. 18 (1844); ed. II. ii. 327 (1861), sphalm. = TELMATOPHACE, Schleid. = **Lemna**, Linn.

THELYCHITON, Endl. Prod. Fl. Norf. 32 (1833) = **Dendrobium**, Sw. (Orchid.).
argyropus, Endl. l. c. = Bulbophyllum argyropus.
brachypus, Endl. l. c. 33 = D. brachypus.
macropus, Endl. l. c. = D. macropus.

THELYCRANIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 394 = **Cornus**, Tourn. (Cornac.).
sanguinea, Fourr. l. c. = C. sanguinea.

THELYGONUM, Linn. Gen. ed. I. 357 (1737) (Theligonum); Schreb. Gen. ii. 644 (1791). *URTICACEAE*, Benth. & Hook. f. iii. 395.
CYNOCRAMBE, Gaertn. Fruct. i. 362. t. 75 (1788).
alsinoideum, Lam. Fl. Fr. ii. 198 = T. Cynocrambe.
Cynocrambe, *Linn. Sp. Pl.* 993.—Reg. Mediterr.; Orient; Ind. or.

THELYMITRA, Forst. Char. Gen. 97. t. 49 (1776).
ORCHIDEAE, Benth. & Hook. f. iii. 607.
MACDONALDIA, Gunn, ex Lindl. Swan River App. 50. t. 9 (1839).
angustifolia, R. Br. Prod. 314 = longifolia.
angustifolia, Hook. f. Fl. Tasm. ii. 5 = aristata.
antennifera, *Hook. f. l. c.* 4. t. 101 A.—Austral.
arenaria, Lindl. Gen. et Sp. Orch. 519 = longifolia.
aristata, *Lindl. l. c.* 521.—Austral.
Benthamiana, *Reichb. f. Beitr. Syst. Pfl.* 55.—Austral.
campanulata, Endl. in Lehm. Pl. Preiss. ii. 14 = canaliculata.
campanulata, Lindl. Swan River App. 49 = ixioides.
canaliculata, *R. Br. Prod.* 314.—Austral.
carnea, *R. Br. l. c.*—Austral.
circumsepta, *Fitzg. Austral. Orch.* t. 4.—Austral.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 271.—N. Zel.
cornicina, Reichb. f. Beitr. Syst. Pfl. 54 = canaliculata.
crinita, *Lindl. Swan River App.* 49.—Austral.
cyanea, *Lindl. ex Benth. Fl. Austral.* vi. 323.—Austral.
epipactoides, F. Muell. Fragm. v. 174 = canaliculata.
flexuosa, *Endl. Nov. Stirp. Dec.* 23.—Austral.
Forbesii, *Ridl. in H. O. Forbes, Nat. Wand. East. Archip.* 518.—Ins. Timor.
formosa, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 328.—N. Zel.
Forsteri, Sw. in Vet. Akad. Handl. Stockh. xxi. (1800) 228 = longifolia.
fuscolutea, *R. Br. Prod.* 315.—Austral.
graminea, Lindl. Swan River App. 49 = longifolia.
grandiflora, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
grandis, F. Muell. ex Benth. Fl. Austral. vi. 319 = aristata.
imberbis, *Hook. f. Fl. N. Zel.* i. 244.—N. Zel.
intermedia, *Berggr. in Minneskr. Fisiog. Sällks. Lund*, (1877) n. viii. 21.—N. Zel.
iridioides, Sieber, ex Benth. Fl. Austral. vi. 318 = ixioides.
ixioides, *Sw. in Vet. Akad. Handl. Stockh.* xxi. (1800) 228. t. 3.—Austral.
javanica, *Blume, Bijdr.* 419.—Java.
juncifolia, Lindl. Gen. et Sp. Orch. 522 = ixioides.
lilacina, F. Muell. ex Lindl. in Linnaea, xxvi. (1854-55) 242 = ixioides.
longifolia, *Forst. Char. Gen.* 98. t. 49.—Austral.
luteocilium, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
Mackibbinii, *F. Muell. in Melb. Chemist* (1881), 44.—Austral.
Macmillani, *F. Muell. Fragm.* v. 93.—Austral.
macrophylla, Lindl. Swan River App. 49 = aristata.
media, R. Br. Prod. 314 = canaliculata.
mucida, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
nemoralis, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 249.—N. Zel.
nuda, R. Br. Prod. 314 = longifolia.
ovata, F. Muell. Fragm. vi. 84 = crinita.
pardalina, F. Muell. l. c. v. 94 = villosa.
pauciflora, R. Br. Prod. 314 = longifolia.
porphyrostica, F. Muell. Fragm. v. 97 = variegata.
pulchella, *Hook. f. Fl. N. Zel.* i. 244.—N. Zel.
purpureo-fusca, *Colenso, in Trans. N. Z. Inst.* xvii. (1880) 249.—N. Zel.
rubra, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.
Smithiana, Hook. f. Fl. Tasm. ii. 4. t. 101 B = flexuosa.
stellata, *Lindl. Swan River App.* 49.—Austral.
stenopetala, *Hook. f. Fl. Antarct.* 69.—Ins. Dom. Auckland.
tigrina, *R. Br. Prod.* 315.—Austral.
uniflora, *Hook. f. Fl. Antarct.* 70.—Ins. Dom. Auckland.
variegata, *Lindl. ex Benth. Fl. Austral.* vi. 323.—Austral.
venosa, *R. Br. Prod.* 314.—Austral.
venosa, Hook. f. Fl. Tasm. ii. 4. t. 102 A = cyanea.
versicolor, Lindl. Gen. et Sp. Orch. 520 = longifolia.
villosa, *Lindl. Swan River App.* 49. t. 8 C.—Austral.
winalis, *Fitzg. in Gard. Chron.* (1882) i. 495.—Austral.

THELYPODIUM, Endl. Gen. 876 (1839). *CRUCIFERAE*, [Benth. & Hook. f. i. 81].
PACHYPODIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 96 (1838).

THELYPODIUM:—
ambiguum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 290.—Am. bor.
auriculatum, *S. Wats. l. c.* xvii. (1881–82) 321.—N. Mexic.
brachycarpum, *Torr. Bot. U. St. Expl. Exped.* ii. 231. t. 1.—Calif.
Cooperi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 246.—Calif.
flavescens, *S. Wats. in Bot. King's Exp.* 25.—Calif.
integrifolium, *Endl. in Walp. Rep.* i. 172.—Am. bor. occ.
laciniatum, *Endl. l. c.*—Am. bor. occ.
linearifolium, *S. Wats. in Bot. King's Exp.* 25.—Am. bor. occ.
longifolium, *S. Wats. l. c.*—Am. bor. occ.
neglectum, *M. Jones, in Am. Natural.* xvii. II. (1882) 875.—Calif.
Nuttallii, *S. Wats. in Bot. King's Exp.* 26.—Am. bor. occ.
petiolatum, *Hemsl. Diagn. Pl. Nov.* 2.—Mexic.
pinnatifidum, *S. Wats. in Bot. King's Exp.* 25.—Am. bor. occ.
sagittatum, *Endl, in Walp. Rep.* i. 172.—Am. bor. occ.
Wrightii, *A. Gray, Pl. Wright.* i. 7.—N. Mexic.

THELYPOGON, Spreng. Syst. iii. 682 (1826) = **Telipogon**, H. B. & K. (Orchid.).

THELYRA, DC. Prod. ii. 527 (1825) = THELIRA, Thou. = **Parinarium**, Aubl. (Rosac.).
madagascariensis, G. Don, Gen. Syst. ii. 479 = Madag.

THELYSIA, Salisb. in Trans. Hort. Soc. i. (1812) 303 = **Iris**, Tourn.
alata, Parl. Fl. Ital. iii. 317 = I. alata.
caucasica, Parl. l. c., in obs. = I. caucasica.
fumosa, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 304 = I. fumosa.
grandiflora, Salisb. in Trans. Hort. Soc. i. (1812) 303 = I. alata.
persica, Parl. Fl. Ital. iii. 317, in obs. = I. persica.

THELYTHAMNOS, Spreng. f. Tent. Suppl. Syst. 25 (1828) = **Ursinia**, Gaertn. (Compos.).
filiformis, Spreng. l. c. = U. filiformis.

THEMEDA, Forsk. Fl. Aegypt. Arab. 178 (1775) = **Anthistiria**, Linn. f. (Gramin.).
polygama, J. F. Gmel. Syst. 149 = A. Forskålii (cf. Addend).
triandra, Forsk. Fl. Aegypt. Arab. 178 = A. Forskålii.

THEMIS, Salisb. Gen. Pl. Fragm. 85 (1866) = **Brodiaea**, Sm. (Liliac.).
ixioides, Salisb. l. c. = B. ixioides.

THEMISTOCLESIA, Klotzsch, in Linnaea, xxiv. (1851) 41. *VACCINIACEAE*, Benth. & Hook. f. ii. 576.
EPISCOPIA, Mor. ex Klotzsch, in Linnaea, xxiv. (1851) 42, in syn.
buxifolia, Klotzsch, l. c. 42 = Ceratostema buxifolium.
coronilla, *Linden & André, Illustr. Hortic.* xvii. (1870) 176. t. 33.—N. Granat.
pendula, *Klotzsch, in Linnaea*, xxiv. (1851) 42.—N. Granat.

THENARDIA, H. B. & K. Nov. Gen. et Sp. iii. 209. t. 240 (1818). *APOCYNACEAE*, Benth. & Hook. f. ii. 710.
corymbosa, Benth. in Hook. Journ. Bot. iii. (1841) 246 = Forsteronia Schomburgkii.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 209. t. 240.—Mexic.
laurifolia, Benth. in Hook. Journ. Bot. iii. (1841) 246 = Forsteronia laurifolia.
scabra, *Spreng. Syst.* iv. *Cur. Post.* 65.—N. Caled.
suaveolens, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 359.—Mexic.
umbellata, *Spreng. Syst.* i. 636.—Guian.

THENARDIA, [Moç. & Sesse,] ex DC. Prod. iii. 108 (1828) = **Rhynchanthera**, DC. (Melastom.).
rosea, Moç. & Sesse, l. c. = R. mexicana.

THEOBROMA, Linn. Gen. ed. I. 351 (1737). *STERCULIACEAE*, Benth. & Hook. f. i. 225.
CACAO, Gaertn. Fruct. ii. 190. t. 122 (1791).
alba, Bernoulli, in Denkschr. Schw. Naturf. Ges. xxiv. (1871) n. III. 14 (sp. dub.).—Guian.
angustifolia, [Moç. & Sesse,] ex DC. Prod. i. 484 = speciosa.
augusta, Linn. Syst. ed. XII. 233 = Abroma augusta.
bicolor, *Humb. & Bonpl. Pl. Aequin.* i. 104. t. 30.—N. Granat.
Cacao, *Linn. Sp. Pl.* 782.—Am. trop.
caribaea, Sweet, Hort. Brit. ed. II. 67 = T. Cacao?
celtifolia, Salisb. Prod. 387 = Guazuma ulmifolia.
ferruginea, *Bernoulli, in Denkschr. Schw. Naturf. Ges.* xxiv. (1871) n. III. 13.—Bras.
glauca, *Karst. in Linnaea*, xxviii. (1856) 447 (sp. dub.).—N. Granat.
Guazuma, Linn. Sp. Pl. 782 = Guazuma ulmifolia.
guianensis, J. F. Gmel. Syst. 1151 = T. Cacao.
guyanensis, *Voigt, in Syll. Ratisb.* ii. (1828) 55 (sp. dub.).—Guian.
integerrima, Stokes, Bot. Mat. Med. iv. 83 = T. Cacao.
leiocarpa, *Bernoulli, in Denkschr. Schw. Naturf. Ges.* xxiv. (1871) n. III. 6.—Am. centr.
macrantha, *Bernoulli, l. c.* 11.—Bras.
Martiana, *D. Dietr. Syn. Pl.* iv. 1242.—Bras.
microcarpa, *Mart. in Buchn. Repert.* xxxv. (1830) 24; *Bernoulli, in Denkschr. Schw. Naturf. Ges.* xxiv. (1871) n. III. 11. t. 3.—Bras.
montana, *Goudet, ex Bernoulli, l. c.* 15 (sp. ignota).—Hab.?
nitida, *Bernoulli, l. c.*—Bras.
obovata, Klotzsch, ex Bernoulli, l. c. 14 = subincana.
ovatifolia, [Moç. & Sesse,] ex DC. Prod. i. 485 = bicolor.
pentagona, *Bernoulli, in Denkschr. Schw. Naturf. Ges.* xxiv. (1871) n. III. 6.—Am. centr.
quinquenervia, Bernoulli, l. c. 8 = speciosa.
Salzmanniana, *Bernoulli, l. c.* 7 (Saltzmanniana).—Bras.
speciosa, Willd. ex Mart. Reise Bras. iii. 1127 = macrantha.
speciosa, *Willd. ex Spreng. Syst.* iii. 332.—Bras.
Spruceana, Bernoulli, in Denkschr. Schw. Naturf. Ges. xxiv. (1871) n. III. 9 = speciosa.
subincana, *Mart. in Buchn. Repert.* xxxv. (1830) 23.—Bras.
subincana, Spruce, ex Sagot, in Ann. Sc. Nat. Sér. VI. xi. (1881) 154 = speciosa.
sylvestris, Aubl. ex Mart. in Buchn. Repert. xxxv. (1830) 24 = Martiana.
sylvestris, G. Don, Gen. Syst. i. 522 = Genipa Merianae.

THEODORA, Medic. Theod. 16 (1786) = **Schotia**, Jacq. (Legumin.).
speciosa, Medic. l. c. 16, 61 = S. speciosa.
stipulata, Eckl. & Zeyh. Enum. 261 = S. speciosa.

THEODOREA, Cass. in Dict. Sc. Nat. xlvii. (1827); liii. 463 (1828) = **Saussurea**, DC. (Compos.).
amara, Cass. l. c. liii. 464 = S. amara.
pulchella, Cass. l. c. 465 = S. japonica.

THEODOREA, Rodrig. Orch. Nov. i. 144 (1877). *ORCHIDEAE*.
gomezoides, *Rodrig. l. c.* 145.—Bras.

THEODORIA, Neck. Elem. ii. 286 (1790) = **Sterculia**, Linn.

THEOPHRASTA, Linn. Gen. ed. I. 45 (1737), partim; Juss. Gen. 150 (1789). *MYRSINEAE*, Benth. & Hook. f. ii. 649.
americana, *Linn. Sp. Pl.* 149.—Ind. occ.
americana, Sw. Obs. Bot. 59 = Jussieui.
angustifolia, Linden, Cat. n. 18 (1863) 10 (Clavijae sp.?).—Hab.?
antioquensis, Linden, l. c. n. 91 (1874) 11 (Clavijae sp.).—N. Granat.
attenuata, Linden, Cat. n. 18 (1863) 10 (Clavijae sp.?).—Hab.?
cochinchinensis, Spreng. Syst. i. 671 = Melodinus monogynus?
crassipes, Linden, Cat. n. 18 (1863) 10 (Clavijae sp.?).—Hab.?
densiflora, *Decne. in Ann. Sc. Nat.* Sér. VI. iii. (1876) 142.—Ins. S. Doming.
fusca, *Decne. l. c.* 143.—Hab.?
glycycarpa, Spreng. Syst. i. 671 = Leonia glycycarpa.

THEOPHRASTA:—
Henrici, Ham. Prod. Pl. Ind. Occ. 27 = Jussieui.
Hookeri, Linden, Cat. n. 91 (1874) 55 = Clavija Hookeri.
imperialis, *Linden, ex C. Koch & Fint. Wochenschr.* ii. (1859) 91; *Regel, Gartenfl.* (1864) 323. t. 453.—Ind. occ.
integrifolia, Pohl, ex Miq. in Mart. Fl. Bras. x. 277 = Clavija integrifolia.
Jussiaei, Hook. Bot. Mag. t. 4239 (excl. syn.) = densiflora.
Jussiéui, *Lindl. Coll. Bot.* t. 26.—Am. austr.
latifolia, Willd. ex Roem. & Schult. Syst. iv. 787 = Clavija latifolia.
longifolia, Jacq. Coll. iv. 136 = Clavija ornata.
macrophylla, Linden, Cat. n. 91 (1874) 54, nomen = Clavija grandis.
macrophylla, Link, ex Roem. & Schult. Syst. iv. 187 = Clavija macrophylla.
madagascariensis, *Willd. ex Roem. & Schult. l. c.* 787 (sp. dub.).—Madag.
minor, Linden, Cat. n. 18 (1863) 10 (Clavijae sp.?).—Hab.?
nobilis, Linden, l. c. = C. clavata.
pinnata, *Jacq. Fragm.* 49.—Venezuela.
pungens, Willd. ex Roem. & Schult. Syst. iv. 787 = Clavija ornata.
regalis, Linden, Cat. n. 21 (1867) 5 = Clavija clavata.
Rodekiana, Linden, l. c. n. 95 (1876) 12 = C. Rodekiana.
serrata, Hort. ex Hoffmgg. Verz. Pfl. Nachtr. ii. 218 = Clavija macrophylla.
serratifolia, Cat. Cels, (1828) 4; ex Colla, in Mem. Acc. Torin. xxxv. (1831) 151, 153. t. 111 = Brexia madagascariensis?
smaragdina, Hort. Linden, ex Decne. in Ann. Sc. Nat. Sér. VI. iii. (1876) 139 = Deherainia smaragdina.
spinosa, *Willd. ex C. Koch & Fint. Wochenschr.* ii. (1859) 98, 108.—Ind. occ.
tetramera, Mart. in Flora, xxiv. (1841) II. Beibl. 27 = Clavija macrophylla.
umbrosa, Linden, Cat. n. 19 (1865) 6, 15 = Clavija umbrosa.

THEOPHROSERIS, Andrae, in Bot. Zeit. xiii. (1855) 310, sphalm. = TEPHROSERIS, Reichb. = **Senecio**, Tourn. (Compos.).

THEOPYXIS, Griseb. in Goett. Abh. vi. (1856) 126 = **Lysimachia**, Tourn. (Primul.).
chilensis, Griseb. l. c. 127 = L. umbellata.

THEREBINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Oroxylum**?, Vent. (Bignoniac.).

THERESA, Clos, in C. Gay, Fl. Chil. iv. 496. t. 54 (1849) = **Perilomia**, H. B. & K. (Labiat.).
valdiviana, Clos, l. c. 497.—Chili.

THERESIA, C. Koch, in Linnaea, xxii. (1849) 232 = **Fritillaria**, Linn. (Liliac.).
Fleischeri, Klatt, in Hamb. Gartenz. xvi. (1860) 439 = F. Fleischeriana.
libanotica, Boiss. Diagn. Ser. I. xiii. 20 = F. libanotica.
persica, C. Koch, in Linnaea, xxii. (1849) 233 = F. persica.
Pinardi, Klatt, in Hamb. Gartenz. xvi. (1860) 439 = F. Pinardi.
pudica, Klatt, l. c. = F. pudica.
tulipifolia, Klatt, l. c. 438 (excl. syn.) = F. tulipifolia.

THERIOPHONUM, Blume, Rumphia, i. 127 (1835). *AROIDEAE*, Benth. & Hook. f. iii. 967.
CALYPTROCORYNE, Schott, in Oestr. Bot. Wochenschr. vii. (1857) 262.
TAPINOCARPUS, Dalz. in Hook. Kew Journ. iii. (1851) 345.
crenatum, *Blume, Rumphia*, i. 128.—Ind. or.
Dalzellii, *Schott, Aroid.* i. 15 (Dalzelii).—Ind. or.
infaustum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 260.—Ind. or.
Kleinii, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 2 = crenatum.
Wightii, Engl. in DC. Monog. Phan. ii. 608 = infaustum.
Wightii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 3.—Ind. or.
zeylanicum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 258.—Zeylan.

THERMIA, Nutt. Gen. Am. i. 282 (1818) = **Thermopsis**, R. Br. (Legumin.).
alpina, Spreng. Syst. iv. Cur. Post. 171 = Thermopsis corgonensis.
fabacea, Spreng. l. c. = Thermopsis fabacea.
laburnifolia, Spreng. l. c. = Piptanthus nepalensis.
lupinoides, Nutt. Gen. Am. i. 283 = Thermopsis lanceolata.
rhombifolia, Nutt. l. c. 282 = Thermopsis rhombifolia.

THERMINTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 133 = **Pistacia**, Linn. (Anacard.).

THERMOPHILA, Miers, in Trans. Linn. Soc. xxviii. (1872) 398 = **Salacia**, Linn. (Celastr.).
angulosa, Peyr. in Mart. Fl. Bras. xi. 1. 154 = S. multiflora.
cordata, Miers, in Trans. Linn. Soc. xxviii. (1872) 401 = S. multiflora.
emarginata, Miers, l. c. = S. multiflora.
macrophylla, Miers, l. c. 399 = S. macrophylla, *Miq.*
obovata, Miers, l. c. = S. multiflora.
rugulosa, Miers, l. c. 400 = S. multiflora.

THERMOPSIS, R. Br. in Ait. Hort. Kew. ed. II. iii. 3 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 465.
Drepilia, Rafin. New Fl. Am. ii. 52 (1836).
Scolobus, Rafin. in Journ. Phys. lxxxix. (1819) 259.
Thermia, Nutt. Gen. Am. i. 282 (1818).
alpina, Ledeb. Fl. Alt. ii. 112 = corgonensis.
alterniflora, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 18.—Turkestan.
barbata, *Benth. in Royle, Illustr. Bot. Himal.* 196. t. 32. f. 1.—Reg. Himal.
californica, *S. Wats. in Proc. Am. Acad.* xi. (1876) 126.—Am. bor. occ.
caroliniana, *M. A. Curt. in Am. Journ. Sc.* Ser. I. xliv. (1843) 80.—Am. bor.
chinensis, Benth. ex S. Moore, in Journ. Bot. xvi. (1878) 131 = fabacea.
corgonensis, *DC. in DC. Prod.* ii. 99.—Sibir. altaic.
fabacea, *DC. l. c.*—Kamtchat.
fabacea, Hook. Fl. Bor. Am. i. 128 = montana.
fabacea, Torr. Bot. Mex. Bound. 58 = californica.
fraxinifolia, *M. A. Curt. in Am. Journ. Sc.* Ser. I. xliv. (1843) 81.—Am. bor.
gorgonensis, Walp. Rep. i. 562, sphalm. = corgonensis.
inflata, *Cambess. in Jacquem. Voy. Bot.* 35.—Reg. Himal.
laburnifolia, D. Don, Prod. Fl. Nep. 239 = Piptanthus nepalensis.
lanceolata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 3.—Sibir.; Kamtchat.
lupinoides, Link, Enum. Hort. Berol. i. 401 = lanceolata.
macrophylla, *Hook. & Arn. Bot. Beech. Voy.* 329.—Calif.
macrophylla, Torr. in Pacif. Rail. Rep. iv. 81 = californica.
mollis, *M. A. Curt. ex A. Gray, in Mem. Am. Acad.* N. S. iii. (1848) 47. t. 9.—Am. bor.
montana, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 388.—Am. bor.
napaulensis, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 98 = Piptanthus nepalensis.
rhombifolia, *Richards. in Frankl. Narr. 1st Journ. App.* 737.—Am. bor.

THEROFON, Rafin. New Fl. Am. iv. 66 (1836) = **Boykinia**, Nutt. (Saxifr.).
napelloides, Rafin. l. c. = B. aconitifolia.

THEROGERON, DC. Prod. v. 283 (1836) = **Minuria**, DC. (Compos.).
denticulatus, DC. l. c. = M. denticulata.
integerrimus, DC. l. c. = M. integerrima.
tenuifolius, Sond. in Linnaea, xxv. (1852) 467 = M. Cunninghamii.

THEROLEPTA, Rafin. Neogenyt. 3 (1825) = **Marshallia**, Schreb. (Compos.).
pumila, Rafin. l. c. = M. pumila.

THEROPOGON, Maxim. in Bull. Acad. Pétersb. xv. (1871) 89. *LILIACEAE*, Benth. & Hook. f. iii. 771.
pallidus, *Maxim. l. c.* 90.—Reg. Himal.

THESIDIUM, Sond. in Flora, xl. (1857) 364. *SANTALACEAE*, Benth. & Hook. f. iii. 222.
exocarpaeoides, *Sond. l. c.* 365.—Afr. austr.
fragile, *Sond. l. c.* 364.—Afr. austr.
globosum, *A. DC. in DC. Prod.* xiv. 673.—Afr. austr.
hirtulum, Sond. in Flora, xl. (1857) 405 = globosum, strigulosum.
hirtum, Sond. l. c. 365 = globosum, strigulosum.
leptostachyum, *A. DC. in DC. Prod.* xiv. 674.—Afr. austr.
microcarpum, A. DC. l. c. = exocarpaeoides.
podocarpum, A. DC. l. c. = Thunbergii.
strigulosum, *A. DC. l. c.* 673.—Afr. austr.
Thunbergii, *Sond. in Flora*, xl. (1857) 364.—Afr. austr.

THESION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = **Thesium**, Linn. (Santal.).
micromerion, St. Lag. l. c. = Thesium Micromeria.
micromerium, St. Lag. l. c. viii. (1881) 203 = Thesium Micromeria.

THESIUM, Linn. Gen. 60 (1737). *SANTALACEAE*, Benth. & Hook. f. iii. 221.
Frisca, Spach, Hist. Vég. Phan. x. 460 (1841).
Rhinostegia, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 56.
Steinreitera, Opiz, Seznam, 93 (1852).
Xerolophus, Dulac, Fl. Hautes-Pyr. 160 (1867).
acutissimum, *A. DC. Esp. Nouv. Thes.* (1857) 4.—Afr. austr.
adpressifolium, Sond. in Flora, xl. (1857) 358 = funale.
adulterinum, Montand. Guide Bot. 286 = pratense.
alatavicum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 445.—Sibir.
alpinum, Desf. Fl. Atlant. i. 206 = humile.
alpinum, Kunth, ex A. DC. in DC. Prod. xiv. 643 = divaricatum.
alpinum, *Linn. Sp. Pl.* 207.—Europ.; Reg. Cauc.
alpinum, Thuill. Fl. Par. ed. II. 122 = humifusum.
ambiguum, Tenore, Fl. Nap. iii. 312 = pratense.
amblystachyum, *A. DC. Esp. Nouv. Thes.* (1857) 6.—Afr. austr.
amplexicaule, Linn. Mant. ii. 213 = euphorbioides.
angulosum, *A. DC. Esp. Nouv. Thes.* (1857) 2.—Afr. austr.
aphyllum, *Mart. ex A. DC. in DC. Prod.* xiv. 671.—Bras.
apiculatum, Sond. in Flora, xl. (1857) 357 = acutissimum.
asperulum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 194 = ramosum.
assimile, Sond. in Flora, xl. (1857) 360 = carinatum.
aureum, *Jaub. & Spach, Illustr.* ii. 4; iii. 151. t. 300.—Cappadoc.
australe, *R. Br. Prod.* 353.—China; Japon; Austral.
australe, A. Cunn. ex A. DC. in DC. Prod. xiv. 676 = Choretrum lateriflorum.
Basninianum, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 470.—Dahuria.
bavarum, *Schrank, Baier. Reise*, 129.—Europ.
Bergeri, *Zucc. in Abh. Akad. Muench.* ii. (1831-36) 324.—Graec.; As. Min.
bibracteatum, *Turcz. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 23.—Sibir. baical.
Billardieri, Boiss. Diagn. Ser. I. v. 48 = heterophyllum.
Boissieranum, *A. DC. in DC. Prod.* xiv. 663.—Afr. austr.
brachycephalum, Sond. in Flora, xl. (1857) 360 = capitellatum.
brachyphyllum, *Boiss. Diagn.* Ser. I. v. 48.—As. Min.
bracteatum, Gromow, ex Trautv. in Act. Hort. Petrop. ix. (1884) 152, nomen = ebracteatum?
brasiliense, *A. DC. in DC. Prod.* xiv. 671.—Bras.
brevicolle, J. & C. Presl, Fl. Čech. 54 = pratense.
brevifolium, A. DC. Esp. Nouv. Thes. (1857) 3 = leptocaule.
byzantinum, A. DC. in DC. Prod. xiv. 644 = divaricatum.
capitatum, Banks, ex A. DC. l. c. 664 = scabrum.
capitatum, *Linn. Sp. Pl.* 207.—Afr. austr.
capitellatum, *A. DC. Esp. Nouv. Thes.* (1857) 7.—Afr. austr.
capituliflorum, *Sond. in Flora*, xl. (1857) 363.—Afr. austr.
carinatum, *A. DC. Esp. Nouv. Thes.* (1857) 7.—Afr. austr.

THESIUM:—
carnosum, Wolfn. in Flora, xxxix. (1856) 561 = alpinum.
chinense, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. vii. 157 = australe.
ciliatum, *R. Br. Prod.* 353.—Afr. austr.
Colpoda, J. F. Gmel. Syst. 1636, sphalm. = seq.
Colpoon, Linn. f. Suppl. 161; Thunb. Prod. Pl. Cap. 46 = Colpoon compressum.
commutatum, *Sond. in Flora*, xl. (1857) 362.—Afr. austr.
comosum, Roth, Catalect. fasc. ii. 29 = ebracteatum.
compressum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 104.—As. Min.
confertum, Willd. ex A. DC. in DC. Prod. xiv. 659 = ericaefolium.
confine, *Sond. in Flora*, xl. (1857) 363.—Afr. austr.
corniculatum, Krauss, ex A. DC. in DC. Prod. xiv. 659 = squarrosum.
corniculatum, *E. Mey. & Drège, ex A. DC. l. c.* 662.—Afr. austr.
corymbiflorum, Sond. in Flora, xl. (1857) 354 = pinifolium.
corymbosulum, Michx. Fl. Bor. Am. i. 112 = Comandra umbellata.
corymbuligerum, *Sond. in Flora*, xl. (1857) 362.—Afr. austr.
crassifolium, *R. Br. Prod.* 353.—Afr. austr.
crassifolium, *Sond. in Flora*, xl. (1857) 355.—Afr. austr.
cucullatum, A. DC. in DC. Prod. xiv. 656 = paniculatum.
debile, *R. Br. Prod.* 353, nomen; *Spreng. Syst.* i. 830.—Afr. austr.
decumbens, Dollin. Enum. Pl. Austr. Infer. 113 = humile.
decumbens, C. C. Gmel. Fl. Bad. i. 549 = pratense.
decurrens, Blume, ex A. DC. in DC. Prod. xiv. 652 = australe.
densiflorum, *A. DC. Esp. Nouv. Thes.* (1857) 7.—Afr. austr.
diffusum, *Andrz. ex A. DC. in DC. Prod.* xiv. 644.—Podol.
divaricatum, R. Grah. ex A. DC. l. c. 658 = leptocaule.
divaricatum, *Jan, ex Mert. & Koch, Deutschl. Fl.* ii. 285.—Reg. Mediterr.
diversifolium, Sond. in Flora, xl. (1857) 359 = spicatum.
Dregeanum, *A. DC. in DC. Prod.* xiv. 657.—Afr. austr.
Dregei, Sond. in Flora, xl. (1857) 356 = gnidiaceum.
drupaceum, Labill. Nov. Holl. Pl. i. 68. t. 93 = Leptomeria Billardieri.
ebracteatum, *Hayne, in Schrad. Journ.* iii. ii. (1800) 33. t. 7.—Europ.
Ecklonianum, *Sond. in Flora*, xl. (1857) 356.—Afr. austr.
eifaliense, *Steud. Nom.* ed. II. ii. 679.—Europ.
elatius, *Sond. in Flora*, xl. (1857) 355.—Afr. austr.
elegans, Rochel, Pl. Banat. Rar. 36. t. 4 = Comandra elegans.
erectiramosum, A. DC. in DC. Prod. xiv. 655 = quinqueflorum.
ericaefolium, *A. DC. Esp. Nouv. Thes.* (1857) 3.—Afr. austr.
ericoides, *R. Br. Prod.* 353.—Afr. austr.
euphorbioides, *Berg. Desc. Pl. Cap.* 74; *Linn. Mant.* ii. 214.—Afr. austr.
euphorbioides, Jacq. ex A. DC. in DC. Prod. xiv. 655 = strictum.
euphrasioides, *A. DC. Esp. Nouv. Thes.* (1857) 8.—Afr. austr.
Exul, *A. DC. in DC. Prod.* xiv. 647.—Afr. austr.
flexuosum, *A. DC. Esp. Nouv. Thes.* (1857) 6.—Afr. austr.
foliosum, *A. DC. in DC. Prod.* xiv. 656.—Afr. austr.
fragile, Link, ex A. DC. l. c. 660 = crassifolium, *Sond.*
fragile, Linn. f. Suppl. 162 = Thesidium Thunbergii.
fragile, Spreng. ex A. DC. in DC. Prod. xiv. 656 = paniculatum.
Frisea, *Linn. Mant.* ii. 213.—Afr. austr.
Frisea, Sond. in Flora, xl. (1857) 359 = debile.
fulvipes, Griessel. l. c. xxii. (1839) 17 = intermedium.
funale, Herb. Vindob. ex A. DC. in DC. Prod. xiv. 667 = T. Frisea.
funale, *Linn. Syst.* ed. X. 944.—Afr. austr.

THESIUM:—
galioides, *A. DC. Esp. Nouv. Thes.* (1857) 5.—Afr. austr.
gallicum, Schultz, ex Steud. Nom. ed. II. ii. 679 = humifusum.
glaucum, Serres, in Bull. Soc. Bot. Fr. iv. (1857) 438 = divaricatum.
globosum, A. DC. Esp. Nouv. Thes. (1857) 4 = Thesidium globosum.
glomeruliflorum, *Sond. in Flora*, xl. (1857) 355.—Afr. austr.
gnidiaceum, *A. DC. Esp. Nouv. Thes.* (1857) 7.—Afr. austr.
graecum, Boiss. & Sprun. Diagn. Ser. I. v. 47 = Bergeri.
graecum, Zucc. in Abh. Akad. Muench. ii. (1831–36) 322 = humile.
griseum, *Sond. in Flora*, xl. (1857) 357.—Afr. austr.
heterophyllum, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 254.—As. Min.
himalense, *Royle, Illustr. Bot. Himal.* 322, nomen; *Edgew. in Trans. Linn. Soc.* xx. (1846) 88.—Reg. Himal.
hirtulum, Sond. in Flora, xl. (1857) 359 = pubescens.
hispidulum, *Lam. Illustr.* ii. 124.—Afr. austr.
hispidulum, Zeyh. ex A. DC. in DC. Prod. xiv. 659 = squarrosum.
Hockeanum, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* ii. II. 41.—Europ.
hottentottum, Sond. in Flora, xl. (1857) 363 = euphrasioides.
humifusum, *DC. Fl. Fr. Suppl.* 366.—Europ.
humifusum, Dollin. Enum. Pl. Austr. Infer. 113 = ramosum.
humifusum, Griseb. Spicil. Fl. Rumel. ii. 323 = divaricatum.
humifusum, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 295 = nevadense.
humile, *Vahl, Symb. Bot.* iii. 43.—Reg. Mediterr.
Hussenoti, Hussen. Chard. Manc. 114 = humifusum.
Hussenoti, *Schultz, ex Steud. Nom.* ed. II. ii. 679.—Gallia.
Hutterianum, *Opiz, in Bercht. & Opiz, Oek. Fl. Boehm.* ii. II. 46.—Europ.
imbricatum, Drège, ex A. DC. in DC. Prod. xiv. 662 = gnidiaceum.
imbricatum, *Thunb. Diss. Thes.* 6.—Afr. austr.
impressum, *Steud. ex A. DC. in DC. Prod.* xiv. 649.—As. Min.; Persia.
intermedium, Bess. Prim. Fl. Galic. i. 187 = bavarum.
intermedium, *Schrad. Spicil. Fl. Germ.* 27.—Europ.; Reg. Caucas.
italicum, A. DC. in DC. Prod. xiv. 644 = intermedium.
junceum, *Bernh. ex Krauss, in Flora*, xxviii. (1845) 80.—Afr. austr.
junceum, Sond. in Flora, xl. (1857) 359 = flexuosum.
juncifolium, *A. DC. Esp. Nouv. Thes.* (1857) 1.—Afr. austr.
Kotschyanum, Boiss. Diagn. Ser. I. vii. 86 = impressum.
Krebsii, *A. DC. Esp. Nouv. Thes.* (1857) 1.—Afr. austr.
laevigatum, Hort. Kew. ex A. DC. in DC. Prod. xiv. 663 = capitatum.
laxiflorum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 152.—Rossia.
leptocaule, *Sond. in Flora*, xl. (1857) 362, 404.—Afr. austr.
leptostachyum, A. DC. Esp. Nouv. Thes. (1857) 6 = Thesidium leptostachyum.
libanoticum, *Ehrenb. ex A. DC. in DC. Prod.* xiv. 648.—Syria.
linariae-folium, Gilib. Exercit. ii. 429 = intermedium.
lineare, Spreng. ex A. DC. in DC. Prod. xiv. 665 = capituliflorum.
lineatum, *Linn. f. Suppl.* 162.—Afr. austr.
lineatum, Willd. ex A. DC. in DC. Prod. xiv. 672 = Polygalae sp.?
linifolium, Schrank, Baier. Reise, 129; Baier. Fl. i. 506 = *T. Linophyllum*.
Linophyllum, DC. Fl. Fr. iii. 352 = pratense.
Linophyllum, Hayne, in Schrad. Journ. iii. (1800) I. 28. t. 6 = bavarum.
Linophyllum, Linn. Sp. Pl. 207 = bavarum, intermedium, divaricatum.
Linophyllum, Pall. Reise, ii. 458 = refractum.
Linophyllum, Patrin, ex A. DC. in DC. Prod. xiv. 646 = longifolium.
Linophyllum, Reichb. Ic. Fl. Germ. xi. 10. t. 546 = intermedium.

THESIUM:—
lobelioides, *A. DC. Esp. Nouv. Thes.* (1857) 8.—Afr. austr.
longifolium, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 100; xxv. (1852) II. 469.—Dahuria.
macranthum, *Fenzl, in Flora*, xxvi. (1843) 401.—Mesopot.
macranthum, Presl, in Sommer. Topogr. xi. 26; ex Wolfner, in Flora, xxxix. (1856) 561 = rostratum.
macrocarpon, Michx. ex A. DC. in DC. Prod. xiv. 649 = impressum.
macrocarpum, E. Mey. ex A. DC. l. c. 635 = Osyridicarpos natalensis.
macrostachyum, *A. DC. Esp. Nouv. Thes.* (1857) 6.—Afr. austr.
madagascariense, *A. DC. in DC. Prod.* xiv. 652.—Madag.
magalismontanum, *Sond. in Flora*, xl. (1857) 358.—Afr. austr.
maritimum, *C. A. Mey. Verz. Pfl. Cauc.* 40.—Reg. Casp.
microcarpum, A. DC. Esp. Nouv. Thes. (1857) 5 = Thesidium exocarpaeoides.
Micromeria, *A. DC. l. c.* 8.—Afr. austr.
micropogon, *A. DC. l. c.* 6.—Afr. austr.
monophyllum, Gilib. Exercit. ii. 428 = ebracteatum.
montanum, Ehrh. Beitr. v. 175, nomen = bavarum.
monticolum, Sond. in Flora, xl. (1857) 359 = amblystachyum.
multicaule, Fisch. & Mey. ex A. DC. in DC. Prod. xiv. 649 = Szowitsii.
multicaule, Hook. f. & Thoms. ex A. DC. l. c. 646 = pachyrhizum.
multicaule, *Ledeb. Fl. Alt.* i. 276.—Sibir. altaic.
multiflorum, A. DC. Esp. Nouv. Thes. (1857) 4 = squarrosum.
nanum, Patrin, ex A. DC. in DC. Prod. xiv. 648 = repens.
natalense, Sond. in Flora, xl. (1857) 358 = macrostachyum.
nevadense, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* i. 295.—Hispan.
nigromontanum, *Sond. in Flora*, xl. (1857) 361.—Afr. austr.
nilagiricum, Miq. Anal. Bot. Ind. iii. 15 = Wightianum.
pachyrhizum, *A. DC. in DC. Prod.* xiv. 646.—Reg. Himal.
palatinum, Roth, Catalect. fasc. ii. 29 = ramosum.
pallidum, *A. DC. Esp. Nouv. Thes.* (1857) 2.—Afr. austr.
paniculatum, *Linn. Mant.* i. 51.—Afr. austr.
paniculatum, Portenschl. ex A. DC. in DC. Prod. xiv. 659 = ericaefolium.
paniculatum, Thunb. Prod. Pl. Cap. 45 = Thunbergianum.
Parnassi, *A. DC. in DC. Prod.* xiv. 643.—Ital.; Graec.
paronychioides, Sond. in Flora, xl. (1857) 359 = junceum.
parvifolium, *A. DC. Esp. Nouv. Thes.* (1857) 3.—Afr. austr.
patentiflorum, Sond. in Flora, xl. (1857) 357 = micropogon.
phylicoides, E. Mey. ex A. DC. in DC. Prod. xiv. 672 (Phylicae sp. ?).—Afr. austr.
phyllostachyum, *Sond. in Flora*, xl. (1857) 355.—Afr. austr.
pinifolium, *A. DC. Esp. Nouv. Thes.* (1857) 2.—Afr. austr.
planifolium, A. DC. l. c. 5 = triflorum.
podocarpum, A. DC. l. c. = Thesidium Thunbergii.
pratense, *Ehrh. Beitr.* v. 175, nomen.—Europ.
pratense, Holand. ex Steud. Nom. ed. II. ii. 679 = Hussenoti.
pratense, Ledeb. Fl. Ross. iii. 542 = ramosum.
pratense, Turcz. Fl. Baic. ii. I. 79 = saxatile.
pratense, Vahl, in Fl. Dan. t. 1205 = ebracteatum.
procumbens, *C. A. Mey. Verz. Pfl. Cauc.* 40.—Reg. Caucas.
psilotoides, *Hance, in Journ. Bot.* vi. (1868) 48.—China.
pubescens, *A. DC. Esp. Nouv. Thes.* (1857) 7.—Afr. austr.
pumilum, Patrin, ex A. DC. in DC. Prod. xiv. 648 = repens.
pyrenaicum, Pourr. in Mém. Acad. Toul. iii. (1788) 331 = pratense.

THESIUM:—
quinqueflorum, *Sond. in Flora*, xl. (1857) 354.—Afr. austr.
racemosum, *Bernh. ex Krauss, l. c.* xxviii. (1845) 79.—Afr. austr.
radicans, *Hochst. l. c.* xxiv. (1841) I. *Intell.* 22, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 235.—Abyss.
radicans, Hohen. ex A. DC. in DC. Prod. xiv. 647 = Wightianum.
ramellosum, Sond. in Flora, xl. (1857) 362 = ericaefolium.
ramosum, Bess. ex A. DC. in DC. Prod. xiv. 644 = diffusum.
ramosum, Griseb. Spicil. Fl. Rumel. ii. 223 = Parnassi.
ramosum, *Hayne, in Schrad. Journ.* iii. (1800) I. 30. t. 7.—Europ.; As. or.
ramosum, Ledeb. Fl. Alt. i. 275 = refractum.
ramosum, J. & C. Presl, Fl. Čech. 54 = intermedium.
rariflorum, *Sond. in Flora*, xl. (1857) 364.—Afr. austr.
recurvifolium, Sond. l. c. 356 = lobelioides.
refractum, Kar. & Kir. ex A. DC. in DC. Prod. xiv. 644 = ramosum.
refractum, Ledeb. Fl. Ross. iii. 539, partim = saxatile.
refractum, *C. A. Mey. in Bull. Sc. Acad. Pétersb.* viii. (1841) 340.—Sibir. altaic.
repens, *Ledeb. Fl. Alt.* i. 274.—Sibir. altaic.
rigidum, Sond. in Flora, xl. (1857) 354 = lineatum.
robustum, Bernh. ex Krauss, in Flora, xxviii. (1845) 80 = strictum.
rostratum, *Mert. & Koch, Deutschl. Fl.* ii. 287.—Europ.
rugulosum, Bunge, ex A. DC. in DC. Prod. xiv. 649 = australe.
rupestre, *Ledeb. Fl. Alt.* i. 277.—Sibir. altaic.
Rytschowii, Fisch. ex A. DC. in DC. Prod. xiv. 646 = longifolium.
saxatile, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 100.—Dahuria.
scabrum, *Linn. Sp. Pl.* ed. II. 302; *Berg. Desc. Pl. Cap.* 73.—Afr. austr.
scandens, *E. Mey. in Drège, Zwei Pfl. Docum.* 226.—Afr. austr.
Schimperianum, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 22, nomen; *et ex A. Rich. Tent. Fl. Abyss.* ii. 235.—Abyss.
sedifolium, A. DC. in DC. Prod. xiv. 660 = crassifolium.
selagineum, *A. DC. Esp. Nouv. Thes.* (1857) 3.—Afr. austr.
serratum, *Kit. ex D. Dietr. Syn. Pl.* i. 878.—Pannonia.
sessiliflorum, Sekera, ex Nym. Consp. 644 = alpinum.
Souardi, Schultz, ex Steud. Nom. ed. II. ii. 680 = Hussenoti.
sparteum, *R. Br. Prod.* 353.—Afr. austr.
sparteum, Sond. in Flora, xl. (1857) 358 = macrostachyum.
spathulatum, Blume, Bijdr. 646 = Henslowia umbellata.
spathulatum, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 244 = Henslowia varians.
spicatum, *Linn. Mant.* ii. 214.—Afr. austr.
spicatum, Murr. Syst. ed. XV. 249 = funale.
spinosum, Jacq. ex A. DC. in DC. Prod. xiv. 660 = crassifolium, *Sond.*
spinosum, *Linn. f. Suppl.* 161.—Afr. austr.
spinosum, Portenschl. ex A. DC. in DC. Prod. xiv. 647 = T. Exul.
spinulosum, *A. DC. l. c.*—Afr. austr.
squarrosum, *Linn. f. Suppl.* 162.—Afr. austr.
stelleroides, *Jaub. & Spach, Illustr.* ii. 4. t. 104.—Cappadoc.
strictum, *Berg. Desc. Pl. Cap.* 73.—Afr. austr.
strictum, Dum. Fl. Belg. 16 = eifaliense.
strictum, Krauss, ex A. DC. in DC. Prod. xiv. 656 = paniculatum.
strigulosum, A. DC. Esp. Nouv. Thes. (1857) 4 = Thesidium strigulosum.
subnudum, Sond. in Flora, xl. (1857) 360 = spicatum.
Szowitsii, *A. DC. in DC. Prod.* xiv. 649.—Armenia.
tauricolum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1067.—As. Min.
tenue, *Bernh. ex Krauss, in Flora*, xxviii. (1845) 80.—Afr. austr.
tenuifolium, Saut. ex Koch, Syn. Fl. Germ. ed. II. 718 = alpinum.

THESIUM:—
tenuissimum, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 19.—Afr. trop.
teretifolium, *R. Br. Prod.* 353.—Afr. austr.
Thunbergianum, *A. DC. in DC. Prod.* xiv. 666.—Afr. austr.
transgariepinum, Sond. in Flora, xl. (1857) 356 = Zeyheri.
transsilvanicum, Schur, Enum. Pl. Transs. 589 = intermedium.
triflorum, *Linn. f. Suppl.* 162.—Afr. austr.
Turczaninowii, *Sond. in Flora*, xl. (1857) 354.—Afr. austr.
umbellatum, Linn. Sp. Pl. 208 = Comandra umbellata.
umbrosum, Opiz, in Bercht. & Opiz, Oek. Fl. Boehm. ii. II. 45 = bavarum, &c.
virens, *E. Mey. ex A. DC. in DC. Prod.* xiv. 653.—Afr. austr.
virgatum, *Lam. Illustr.* ii. 123.—Afr. austr.
virgatum, Sond. in Flora, xl. (1857) 361, 402 = juncifolium.
Vlassovianum, Fisch. ex A. DC. in DC. Prod. xiv. 646 = longifolium *vel* Barninianum.
Weickerianum, Opiz, in Bercht. & Opiz, Oek. Fl. Boehm. ii. II. 41 = alpinum.
Wightianum, *Wall. Cat.* n. 4037.—Ind. or.
Zeyheri, *A. DC. Esp. Nouv. Thes.* (1857) 8.—Afr. austr.

THESPESIA, Soland. ex Correa, in Ann. Mus. Par. ix. (1807) 290. t. 8. f. 2. *MALVACEAE*, Benth. & Hook. f. i. 208.
AZANZA, Alef. in Bot. Zeit. xix. (1861) 298.
altissima, *Spreng. Syst.* iv. *Cur. Post.* 257.—Java.
Banalo, Blanco, Fl. Filip. ed. II. 382 = macrophylla.
brasiliensis, *Spreng. Syst.* iii. 96.—Bras.
campylosiphon, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 308.—Ins. Philipp.
Danis, *Oliver, in Hook. Ic. Pl.* t. 1336.—Afr. trop.
grandiflora, *DC. Prod.* i. 456 (sp. dub.).—Porto Rico.
Lampas, Dalz. ex Dalz. & Gibs. Bomb. Fl. 19 = macrophylla.
macrophylla, *Blume, Bijdr.* 73.—As. et Afr. trop.; Ins. Malay.
populnea, *Soland. ex Correa, in Ann. Mus. Par.* ix. (1807) 290.—As. et Afr. trop.
populneoides, Kostel. Allg. Med. Pharm. Fl. v. 1861 = populnea.
sublobata, Blanco, Fl. Filip. ed. II. 382 = macrophylla.
tomentosa, *Presl, Rel. Haenk.* ii. 136.—Mexic.

THESPIDIUM, F. Muell. in Journ. Landsborough, Exped. App. (1862). *COMPOSITAE*, Benth. & Hook. f. ii. 292.
basiflorum, *F. Muell. l. c.*—Austral.

THESPIS, DC. in Guill. Arch. Bot. ii. (1833) 517. *COMPOSITAE*, Benth. & Hook. f. ii. 285.
divaricata, *DC. l. c.* 517.—Ind. or.; Burma.
erecta, DC. Prod. v. 375 = divaricata.
sinapifolia, DC. in Guill. Arch. Bot. ii. (1833) 517 = divaricata.

THEVENOTIA, DC. in Guill. Arch. Bot. ii. (1833) 331. *COMPOSITAE*, Benth. & Hook. f. ii. 466.
persica, *DC. Prod.* vi. 551; *et in Deless. Ic. Sel.* iv. t. 72.—Persia.
scabra, *Boiss. Fl. Orient.* iii. 455.—Persia.

THEVENOTIA, Linn. Gen. ed. I. 380 (1737) = **Genipa**, Tourn. (Rubiac.).

THEVETIA, Linn. Gen. ed. I. 50 = praec.

THEVETIA, Linn. Gen. ed. I. 62 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 699.
Ahouai, *A. DC. in DC. Prod.* viii. 345.—Bras.
Ahuai, Rafin. Sylva Tellur. 91 = praec.
bicornuta, *Muell. Arg. in Linnaea*, xxx. (1859–60) 392.—Paraguay.
calophylla, *Miers, Apocyn. S. Am.* 20.—Venezuela.
cornuta, Miers, l. c., sphalm. = bicornuta.
cuneifolia, *A. DC. in DC. Prod.* viii. 344.—Mexic.
Humboldtii, *Schomb. Faun. Fl. Brit. Gui.* 1144, nomen.—Guian.
linearis, Rafin. Sylva Tellur. 91; A. DC. in DC. Prod. viii. 354 = nereifolia.

THEVETIA:—
nereifolia, *Juss. ex Steud. Nom.* ed. II. ii. 680.—Am. trop.
nitida, *A. DC. in DC. Prod.* viii. 344.—Mexic.
ovata, *A. DC. l. c.*—Mexic.
plumeriaefolia, *Benth. Bot. Voy. Sulph.* 124. t. 43.—Honduras.
Yccotli, *A. DC. in DC. Prod.* viii. 343.—Mexic.

THEVETIA, Vell. Fl. Flum. 57; i. t. 151 (1825). Genus ignotum, an *RUTACEACEAE*?
pinnata, Vell. ll. cc.—Bras.

THEYGA, Molina, Sagg. Chile, ed. II. 163 (1810) = THIGA, Molina = **Laurelia**, Juss. (Monim.).

THEYODIS, A. Rich. Tent. Fl. Abyss. i. 364 (1847) = **Oldenlandia**, Linn. (Rubiac.).
octodon, A. Rich. l. c. 365 = O. hedyotoides.

THIBAUDIA, Ruiz & Pav. Fl. Per. iv. tt. 384–388, partim [1802]; et ex Jaume St. Hil. Expos. i. 362 (1805); H. B. & K. Nov. Gen. et Sp. iii. 268 (1818). *VACCINIACEAE*, Benth. & Hook. f. ii. 571.
CHUPALON, Adans. Fam. ii. 164 (1763), nomen prius.
acuminata, Dun. in DC. Prod. vii. 565 (Quid?).—Peruv.
acuminata, Hook. f. Bot. Mag. t. 5752 = Cavendishia acuminata.
acuminata, Wall. Cat. n. 6297 = Corallobotrys acuminata.
alata, Dun. in DC. Prod. vii. 562 (= *Proclesia alata*).—Peruv.
alnifolia, Dun. l. c. 564 (= *Proclesia alnifolia*).—Venezuela.
angustifolia, Hook. Ic. Pl. t. 110 = Eurygania angustifolia.
ardisiaefolia, H. B. & K. Nov. Gen. et Sp. iii. 274 = Eurygania ardisiaefolia.
auriculata, Griff. Notul. iv. 300; Ic. Pl. Asiat. t. 508 = Agapetes auriculata.
bicolor, Ruiz & Pav. ex Dun. in DC. Prod. vii. 561 = Psammisia bicolor.
bracteata, Benth. Pl. Hartw. 143 (= *Proclesia Hartwegiana*).—Ecuador.
bracteata, Ruiz & Pav. Fl. Per. iv. t. 388 (= *Proclesia bracteata*).—Peruv.
breviflora, Benth. Pl. Hartw. 224 = Psammisia breviflora.
camelifolia, Griff. Notul. iv. 301; Ic. Pl. Asiat. t. 516 = Vaccinium Griffithianum.
capitata, Benth. Pl. Hartw. 142 = Cavendishia nobilis.
caulialata, Ruiz & Pav. Fl. Per. iv. t. 386 (= *Proclesia alata*).—Peruv.
Cerander, Dun. in DC. Prod. vii. 565 (Quid?).—Hab.?
cerandra, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = praec.
cinnamomifolia, Willd. ex Walp. Ann. ii. 1087 = Cavendishia pubescens.
coarctata, Ruiz & Pav. Fl. Per. iv. t. 385 = Psammisia bicolor.
cordifolia, H. B. & K. Nov. Gen. et Sp. iii. 271 (= *Proclesia cordifolia*).—N. Granat.
coriacea, Blume, Bijdr. 860 = Vaccinium Korthalsii.
coronaria, Hook. f. Bot. Mag. t. 5575 = Themistoclesia coronilla.
corymbodendron, Ruiz & Pav. ex Dun. in DC. Prod. vii. 569 = Vaccinium corymbodendron.
crassifolia, Benth. Pl. Hartw. 65 = Cavendishia crassifolia.
crenulata, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 234.—Bolivia.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 73.—Cuba.
cuneifolia, Blume, Bijdr. 861 = Vaccinium cuneifolium.
cyathifera, Benth. Pl. Hartw. 222 = Psammisia cyathifera.
dependens, *Benth. l. c.* 220.—N. Granat.
diphylla, Dun. in DC. Prod. vii. 562 (Quid?).—Peruv.
elliptica, Blume, Bijdr. 859 = Vaccinium ellipticum.
elliptica, Ruiz & Pav. Pl. Per. iv. t. 384 (= *Siphonandra elliptica*).—Peruv.
emarginata, *Ruiz & Pav. l. c.*—Peruv.
falcata, H. B. & K. Nov. Gen. et Sp. iii. 269 = Psammisia falcata.

THIBAUDIA:—
flava, Nutt. ex Hook. f. in Hook. Bot. Mag. sub t. 4910 = Pentapterygium flavum.
floribunda, Blume, Bijdr. 859 = Vaccinium Teysmanni.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 269.—N. Granat.
formosa, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1088, nomen = Psammisia formosa.
gaultherifolia, Griff. Notul. iv. 301; Ic. Pl. Asiat. t. 512 = Vaccinium gualtheriaefolium.
glabra, Griff. ll. cc. 301; 514 = Agapetes glabra.
glandulosa, Humb. & Bonpl. Rel. Hist. 602; ex DC. Prod. vii. 556 = Gaylussacia buxifolia.
glandulosa, Willd. ex Cham. & Schlecht. in Linnaea, i. (1826) 528 = Gaylussacia decipiens.
grandiflora, Ruiz & Pav. ex Dun. in DC. Prod. vii. 553 = Ceratostema grandiflorum.
guyanensis, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1087, nomen.—Guian.
Hendersoni, *Regel, Gartenfl.* (1875) 258; *et in Act. Hort. Petrop.* iii. II. (1875) 287.—Hab.?
hirsuta, Ruiz & Pav. ex Dun. in DC. Prod. vii. 553 = Ceratostema cordifolium.
hirtiflora, *Benth. Pl. Hartw.* 224.—N. Granat.
Hookeri, Walp. Rep. vi. 412 = Cavendishia acuminata.
Jessicae, *Hook. f. Bot. Mag.* t. 5547.—Venezuela?
Jussiaei, Dun. in DC. Prod. vii. 565 (Quid?).—Peruv.
latifolia, Griseb. Fl. Brit. W. Ind. 143 = Vaccinium latifolium.
laurifolia, Blume, Bijdr. 859 = Vaccinium laurifolium.
laurifolia, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 530.—Mexic.
leucostoma, Benth. ex Meissn. in Mart. Fl. Bras. vii. 127 = Psammisia leucostoma.
longifolia, H. B. & K. Nov. Gen. et Sp. iii. 269 = Psammisia longifolia.
loranthiflora, Wall. Cat. n. 754 = Agapetes loranthiflora.
lucida, Blume, Bijdr. 860 = Vaccinium Rollisoni.
macrantha, Hook. Bot. Mag. t. 4566 = Agapetes macrantha.
macrocalyx, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 235.—Bolivia.
macrophylla, H. B. & K. Nov. Gen. et Sp. iii. 270 = Psammisia macrophylla.
madagascarensis, Spreng. Syst. ii. 294 = Agapetes madagascariensis.
Martii, *Meissn. in Mart. Fl. Bras.* vii. 173.—Bras.
melastomoides, H. B. & K. Nov. Gen. et Sp. iii. 272 (= *Proclesia melastomoides*).—N. Granat.
mellifera, G. Don, Gen. Syst. iii. 860, sphalm. = seq.
melliflora, Ruiz & Pav. Fl. Per. iv. t. 387 = Eurygania melliflora.
mexicana, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 530 = Cavendishia acuminata.
microphylla, *Lindl. in Gard. Chron.* (1848) 23.—Peruv.
microphylla, Ruiz & Pav. ex Dun. in DC. Prod. vii. 576 = Vaccinium acuminatum.
Moricandi, Dun. in DC. Prod. vii. 563 (Quid?).—Peruv.
multiflora, Klotzsch, in Linnaea, xxiv. (1851) 27 = Eurygania ovata.
myrtifolia, Griff. Notul. iv. 301; Ic. Pl. Asiat. t. 510 = Pentapterygium serpens.
myrtoidea, Blume, Bijdr. 861 = Vaccinium myrtoides.
nitida, H. B. & K. Nov. Gen. et Sp. iii. 271 = Psammisia nitida.
nutans, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1087, nomen.—Guian.
obliqua, Griff. Notul. iv. 301; Ic. Pl. Asiat. t. 515 = Agapetes setigera.
oblongifolia, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 233.—Bolivia.
ocanensis, Linden, Cat. n. 6 (1851) 4 (Quid?).—N. Granat.
panurensis, *Benth. ex Meissn. in Mart. Fl. Bras.* vii. 125.—Peruv.
penduliflora, Dun. in DC. Prod. vii. 562 = Psammisia penduliflora.
phyllireaefolia, Dun. l. c. 564 = Eurygania phillyreaefolia.
pichinchensis, *Benth. Pl. Hartw.* 223.—N. Granat.
pubescens, Benth. l. c. 224 (= *Proclesia Pseudopubescens*).—N. Granat.
pubescens, H. B. & K. Nov. Gen. et Sp. iii. 273 = Cavendishia pubescens.

THIBAUDIA:—
pulcherrima, Wall. ex Hook. Bot. Mag. t. 4303 = Agapetes variegata.
punctata, Dun. in DC. Prod. vii. 564, in syn. = punctatifolia.
punctatifolia, *Ruiz & Pav. Fl. Per.* iv. t. 387.—Peruv.
Quereme, H. B. & K. Nov. Gen. et Sp. iii. 274. t. 256 = Cavendishia Querema.
retusa, Griff. Notul. iv. 300; Ic. Pl. Asiat. t. 509 = Vaccinium retusum.
revoluta, Griff. ll. cc. 301; 513 = Vaccinium Dunalianum.
rhodoroides, Lamb. ex Dun. in DC. Prod. vii. 554 = Agapetes variegata.
rosea, *Jungh. in Flora*, xxx. (1847) 524.—Java.
rupestris, H. B. & K. Nov. Gen. et Sp. iii. 270 = Psammisia rupestris.
sarcantha, Hook. Bot. Mag. t. 5450 = Psammisia sarcantha?
scabriuscula, H. B. & K. Nov. Gen. et Sp. iii. 272 (= *Proclesia scabriuscula*).—Ecuador.
Schlimiana, *Linden, Cat.* n. 6 (1851) 4.—N. Granat.
secundiflora, Poepp. & Endl. Nov. Gen. et Sp. i. 5. t. 9 = Orthaea secundiflora.
septemnervia, Willd. ex Walp. Ann. ii. 1087 = Cavendishia pubescens.
serrata, Dun. in DC. Prod. vii. 563 (Quid?).—Peruv.
serrata, Wall. Cat. n. 6299 = Vaccinium serratum.
setigera, Wall. l. c. n. 752 = Agapetes setigera.
speciosa, Benth. Pl. Hartw. 141 = Semiramisia speciosa.
Sprengelii, Wall. Cat. n. 6298 = Vaccinium Donianum.
strobilifera, H. B. & K. Nov. Gen. et Sp. iii. 272 (= *Proclesia strobilifera*).—N. Granat.
tarapotana, Meissn. in Mart. Fl. Bras. vii. 126 = Cavendishia tarapotana.
tetragona, *Linden, Cat.* n. 6 (1851) 4.—N. Granat.
vaccinacea, Royle, Illustr. Bot. Himal. 256, sphalm. = *verticillata*.
variegata, Royle, l. c. 257 = Agapetes setigera.
variegata, Wall. Cat. n. 751 = Agapetes variegata.
varingiaefolia, Blume, Bijdr. 860 = Vaccinium varingiaefolium.
verticillata, Wall. Cat. n. 753 = Agapetes setigera.
virgata, Ruiz & Pav. ex DC. Prod. vii. 571 = Vaccinium attenuatum.
Vitis-idaea, *Griseb. in Lechl. Berb. Am. Austr.* 58.—Peruv.

THICUANIA, Rafin. Fl. Tellur. iv. 47 (1836) = **Dendrobium**, Sw. (Orchid.).
moschata, Rafin. l. c. = D. moschatum.

THIEBAUTIA, Colla, in Mém. Soc. Linn. Par. iii. (1825) 161. t. 4 = **Bletia**, Ruiz & Pav. (Orchid.).
nervosa, Colla, l. c. = B. florida.

THIELEODOXA, Cham. in Linnaea, ix. (1834) 251 = **Alibertia**, A. Rich. (Rubiac.).
elliptica, Cham. l. c. 252.—Bras.
lanceolata, Cham. l. c. 253.—Bras.

THIERSIA, Baill. Adansonia, xii. (1879) 335. *RUBIACEAE*.
insignis, *Baill. l. c.*; *et Hist. Pl.* vii. 486.—Guian.

THIGA, Molina, Sagg. Chile, ed. II. 297 (1810) = **Laurelia**, Juss. (Monim.).
chilensis, Molina, l. c. = L. aromatica.

THILACHIUM et THILAKIUM, Lour. Fl. Cochinch. 342 (1790) = **Thylachium**, Lour. (Cappar.).

THILCUM, Molina, Sagg. Chile, ed. II. 146 (1810) = **Fuchsia**, Plum. (Onagr.).
tinctorium, Molina, l. c. = F. macrostemma.

THILLAEA, Sang. in Atti Acc. Pont. Lincei, Ser. I. v. (1852) 717, sphalm. = **Tillaea**, Linn. (Crassul.).

THILOA, Eichl. in Flora, xlix. (1866) 149 = **Combretum**, Linn.
glaucocarpa, Eichl. l. c. 151 = C. glaucocarpum.
gracilis, Eichl. l. c. 152 = C. gracile.
nitida, Eichl. l. c. = C. nitidum.
paraguariensis, Eichl. l. c. 151.—Bras.
stigmaria, Eichl. l. c. 152 = C. stigmarium.

THIMUS, Neck. Delic. Gallo-Belg. i. 260 (1768) = **Thymus**, Tourn. (Labiat.).

THINOGETON, Benth. Bot. Voy. Sulph. 142 (1844) = **Cacabus**, Benth. (Solan.).
Hookeri, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 217.—Ins. Galapag.
Lobbianum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 360.—N. Granat.
maritimum, Benth. Bot. Voy. Sulph. 142 = C. maritimus.
Miersii, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 359.—Peruv.; N. Granat.

THINOUIA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 368 = **Thouinia**, Poit. (Sapind.).
compressa, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 282.—Bras.
mucronata, Radlk. l. c.—Bras.
myriantha, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 369 = Thouinia myriantha.
obliqua, Radlk. in Sitzb. Math-Phys. Akad. Muench. viii. (1878) 282.—Peruv.
scandens, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 369 = Thouinia scandens.
ternata, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 282 = Banisteria trifoliata.
ventricosa, Radlk. in Atti Congr. Bot. Firenze, 1874 (1876) 61.—Bras.

THIODIA, Benn. Pl. Jav. Rar. 192, in adnot. (1838) = **Laetia**, Linn. (Bixin.).
laetioides, Griseb. in Goett. Abh. ix. (1861) 25 = Casearia icosandra.
serrata, Endl. in Walp. Rep. i. 204.—Ind. occ.
serrata, Griseb. Fl. Brit. W. Ind. 22 = Casearia icosandra.

THIOLLIEREA, Montr. in Mém. Acad. Lyon, x. (1860) 217 = **Randia**, Linn. (Rubiac.).
artensis, Montr. l. c.—N. Caled.

THISANTHA, Eckl. & Zeyh. Enum. 302 (1837) = **Crassula**, Dill.
decumbens, Eckl. & Zeyh. l. c. = C. decumbens.
glabra, Eckl. & Zeyh. l. c. = C. glabra.
glomerata, Eckl. & Zeyh. l. c. = C. glomerata.
patens, Eckl. & Zeyh. l. c. = C. patens.
scaberula, Kunze, ex Fenzl, Adv. [2]; in Ind. Sem. Hort. Vindob. (1851).—Hab.?
strigosa, Eckl. & Zeyh. Enum. 302 = C. glomerata?
subulata, Hook. Ic. Pl. t. 590 = Tillaea subulata.

THISBE, Falc. in Lindl. Veg. Kingd. 183 (1847) = **Herminium**, Linn. (Orchid.).

THISMIA, Griff. in Proc. Linn. Soc. i. (1844) 221; et in Trans. Linn. Soc. xix. (1845) 341. t. 39. *BURMANNIACEAE*, Benth. & Hook. f. iii. 459.
MYOSTOMA, Miers, in Trans. Linn. Soc. xxv. (1866) 474. t. 57.
OPHIOMERIS, Miers, l. c. xx. (1851) 373. t. 15.
TRIBRACHYS, Champ. ex Thw. Enum. Pl. Zeyl. 325 (1864).
Aseroë, *Becc. Malesia*, i. (1878) 252.—Singapore.
Brunoniana, Duch. in Ann. Sc. Nat. Sér III. vii. (1842) 348 = Brunonis.
Brunoniana, Miers, in Trans. Linn. Soc. xx. (1851) t. 15. f. 20 = Gardneriana.
Brunonis, *Griff. in Proc. Linn. Soc.* i. (1844) 221.—Burma.
clandestina, *Miq. Fl. Ind. Bat.* iii. 616.—Java.
Gardneriana, *Hook. f. ex Thw. Enum. Pl. Zeyl.* 325.—Zeylan.
Neptunis, *Becc. Malesia*, i. (1878) 251.—Borneo.
Ophiuris, *Becc. l. c.* 252.—Borneo.

THIUM, Steud. Nom. ed. I. 833 (1821), sphalm. = TIUM, Medic. = **Astragalus**, Tourn. (Legumin.).

THLADIANTHA, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 103 (Enum. Pl. Chin. Bor. 29). *CUCURBITACEAE*, Benth. & Hook. f. i. 825.
calcarata, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 126.—Ind. or.; Burma.
cordifolia, *Cogn. in DC. Monog. Phan.* iii. 424.—Java.
dubia, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 103.—China.

THLADIANTHA:—
dubia, Hook. f. in Hook. Bot. Mag. t. 5469 = calcarata.
dubia, *Naud. ex Griseb. Ind. Sem. Hort. Gotting.* (1877) 8 (Quid?).—Hab.?
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 631.—Reg. Himal.
monoeca, *Griseb. Ind. Sem. Hort. Gotting.* (1877) 8.—Hab.?

THLAPSI, Linn. Syst. ed. I (1735), err. typ. = **Thlaspi**, Tourn. (Crucif.).

THLASIDIA, Rafin. Fl. Tellur. iv. 96 (1836) = **Scabiosa**, Tourn. (Dipsac.).
bidens, Rafin. l. c. 96 = S. bidens.

THLASPI, [Tourn.]. Linn. Gen. ed. I. 193 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 91.
CARPOCERAS, Link, Handb. ii. 289 (1831).
MINAEA, Lojac. in Nouv. Giorn. Bot. Ital. xiii. (1881) 305.
PACHYPHRAGMA, Reichb. Nom. 179 (1841).
PTEROLOBIUM, Andrz. ex DC. Syst. ii. 374 (1821).
PTEROTROPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 337.
THLASPIDEA, Opiz, Seznam, 96 (1852).
affine, *Schott & Kotschy, ex Boiss. Fl. Orient.* i. 327.—Europ. or.
africanum, Burm. f. Fl. Cap. Prod. 17 = Lepidium ruderale.
alliaceum, [Habl.] Desc. Taur. 145 = montanum.
alliaceum, *Linn. Sp. Pl.* 641.—Europ.
alpestre, C. C. Gmel. Fl. Bad. iii. 28. t. 1 = perfoliatum.
alpestre, Jacq. Enum. Stirp. Vindob. 116, 260 = alpinum.
alpestre, *Linn. Sp. Pl.* ed. II. 903.—Europ.
alpestre, Lucé, Fl. Osil. 223 = montanum.
alpestre, Vill. Fl. Delphin. 67 = virgatum.
alpinum, *Crantz, Stirp. Austr.* ed. I. fasc. i. 25; ed. II. 23.—Europ.
alpinum, Griseb. Spicil. Fl. Rumel. i. 280 = affine.
amarum, Crantz, Stirp. Austr. ed. I. fasc. i. 26; ed. II. 25 = Iberis amara.
ambiguum, *Jord. ex F. Schultz, Arch. Fl. Fr. et Allem.* 161.—Europ.
andicola, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 138.—Chili.
annuum, C. Koch, in Linnaea, xv. (1841) 258 = natolicum.
apetalum, Poir. Encyc. vii. 547 = Lepidium incisum.
arabicum, Bieb. Fl. Taur. Cauc. iii. 430 = Aethionema cappadocicum.
arabicum, Desv. Journ. Bot. iii. (1814) 167 = Aethionema cristatum.
arabicum, Vahl, Symb. Bot. ii. 76 = Schouwia arabica.
arenarium, Jord. ex F. Schultz, Arch. Fl. Fr. et Allem. 162 = alpestre.
Arnaudiae, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 60 = virgatum.
arvense, *Linn. Sp. Pl.* 641.—Europ.; As. bor.
arvernense, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 61 = alpestre.
australe, Hook. f. Fl. N. Zel. ii. 325 = Notothlaspi australe.
baicalense, DC. Syst. ii. 376 = arvense.
balcanum, *Janka, in Termesz. Füzetek*, vii. (1883) 126.—Balkan.
banaticum, *Uechtritz, in Oestr. Bot. Zeitschr.* xxv. (1875) 186.—Banat.
bellidifolium, *Griseb. Spicil. Fl. Rumel.* ii. 505.—Macedon.
beugesiacum, Jord. Diagn. i. 267 = alpestre.
bispinosum, Hornem. Hort. Hafn. ii. 604 = Lepidium spinosum.
bonariense, Poir. Encyc. vii. 543 = Lepidium bonariense.
brachypetalum, Jord. Obs. Pl. Crit. iii. (1846) 5 = virgatum.
brevicaule, *Boiss. ex Kotschy, Diagn.* Ser. II. v. 40.—Syria.
brevistylum, Jord. Obs. Pl. Crit. iii. (1846) 27 = rivale.
brevistylum, Mutel, Fl. Fr. i. 99 = Hutchinsia alpina.
bulbosum, Griseb. Spicil. Fl. Rumel. i. 280 = bellidifolium.
bulbosum, *Sprun. in Boiss. Diagn.* Ser. I. i. 74.—Graecia.
Bursa, Thunb. Fl. Jap. 259 = seq.

THLASPI :—

Bursa-pastoris, Linn. Sp. Pl. 647 = Capsella Bursa-pastoris.
Bursetta, Bergeret, ex DC. Syst. ii. 384 = praec.
Buxbaumii, Fisch. ex Hornem. Suppl. Hort. Hafn. 71 = Aethionema cappadocicum.
caerulescens, J. & C. Presl, Fl. Čech. 133 = alpestre.
calaminare, Lej. & Court. Comp. Fl. Belg. ii. 307 = alpestre.
californicum, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 365.—Calif.
campestre, Linn. Sp. Pl. 641 = Lepidium campestre.
Cardamines, Poir. Encyc. vii. 545 = Lepidium Cardamines.
cardiocarpum, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 176.—Reg. Himal.
carneum, [Soland. in] Russ. Aleppo, ed. II. ii. 257 = Aethionema cristatum.
cartilagineum, J. Mayer, in Abh. Boehm. Ges. (1786) 235. t. 7 = Lepidium crassifolium.
cataonicum, *Reut. Cat. Gr. Genèv.* (1868) 4.—*Cf. Bull. Soc. Bot. Fr.* xvi. (1869) *Rev. Bibl.* 53.—Tauria.
cepeaefolium, *Koch, in Mert. & Koch, Deutschl. Fl.* iv. 534.—Europ.
ceratocarpon, *Murr. in Comm. Gotting.* (1774) 26. t. 1.—As. Min.; Sibir.
chalepense, Poir. Encyc. vii. 547 = Lepidium chalepense.
cochleariforme, DC. Syst. ii. 381 = alpestre.
cochleariforme, Eichw. Skizze, 178 = montanum.
cochlearinum, *F. Muell. Pl. Vict.* i. 51.—Austral.
cochlearioides, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 177.—Reg. Himal.
coerulescens, Steud. Nom. ed. II. ii. 681 = alpestre.
collinum, Bieb. Fl. Taur. Cauc. ii. 99 = arvense.
commutatum, Rochel, Bot. Reise Banat. 83 = alpestre, montanum, praecox.
cordatum, Desf. in Ann. Mus. Par. xi. (1808) 382. t. 39 = Aethionema cordatum.
cornutum, [Clairv.] Man. Herb. 214 = ceratocarpum.
coronarium et *coronatum*, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1132 = Iberis umbellata.
Corongianum, Czetz, ex Nym. Consp. 63 = dacicum.
coronopifolium, Bergeret, ex Steud. Nom. ed. I. 826, in syn. = Teesdalia regularis.
corsicum, Soleir. ex Mutel, Fl. Fr. i. 101 = Lepidium hirtum.
corymbosum, Molina, Sagg. Chile, ed. II. 292 = magellanicum.
corymbosum, Reichb. Ic. Fl. Germ. ii. 3 = rotundifolium ?
cristatum, *Serres, in Bull. Soc. Bot. Fr.* iv. (1857) 434.—Europ.
cuneatum, Stokes, Bot. Mat. Med. iii. 434 = Capsella Bursa-pastoris.
cuneifolium, Griseb. ex Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 268 = montanum.
dacicum, *Heuff. in Verh. Zool.-Bot. Ges. Wien*, viii. (1858) 61.—Europ. or.
densiflorum, *Boiss. & Kotschy, ex Unger & Kotschy, Ins. Cyp.* 328; *Boiss. Fl. Orient.* i. 328.—As. Min.; Syria.
divaricatum, Poir. Encyc. vii. 543 = Lepidium divaricatum.
drabaeflorum, Fenzl, Pugill. Pl. Nov. Syr. 14 = natolicum.
Drummondi, *Benth. Fl. Austral.* i. 88.—Austral.
elegans, *Boiss. Diagn.* Ser. I. v. 82.—As. Min.
erraticum, Jord. Pugill. Pl. Nov. 12 = perfoliatum.
exile, *Phil. in Linnaea*, xxviii. (1856) 670.—Chili.
Fendleri, A. Gray, Pl. Wright. ii. 14 = alpestre.
Fischerianum, Bess. ex DC. Syst. ii. 377 = orbiculatum.
formosum, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1132 = Iberis umbellata.
Gaudinianum, Jord. Obs. Pl. Crit. iii. (1846) 14 = alpestre.
geratocarpos, Scop. Delic. Insub. i. 10, err. typ. = ceratocarpon.
glastifolium, Poir. Encyc. vii. 538 = Lepidium glastifolium.
glaucophyllum, *Barn. in C. Gay, Fl. Chil.* i. 168.—Chili.
goesingense, *Halác. in Oestr. Bot. Zeitschr.* xxx. (1880) 173.—Europ.
gracile, *Phil. in Linnaea*, xxviii. (1856) 670.—Chili.
graecum, *Jord. Obs. Pl. Crit.* iii. (1846) 30.—Pelop.
graminifolium, Poir. Encyc. vii. 546 = Lepidium graminifolium.

THLASPI :—

granatense, *Boiss. & Reut. Diagn.* Ser. II. i. 40.—Hispan.
grandiflorum, Bory & Chaub. Nouv. Fl. Pélop. 41. t. 38 = Capsella grandiflora.
Griffithianum, Boiss. Fl. Orient. i. 329 = alpestre.
hastulatum, *Stev. ex DC. Syst.* ii. 388, in syn.—Reg. Casp.
Haussknechtii, *Boiss. Fl. Orient.* i. 330 (Hausknechtii).—Cataonia.
heterophyllum, Cav. ex DC. Syst. ii. 542 = perfoliatum.
heterophyllum, DC. Fl. Fr. iv. 712 = Lepidium hirtum.
hirsutum, Lam. Fl. Fr. ii. 465 = Lepidium campestre, hirtum.
hirtum, Georgi, Beschr. Russ. Reich. III. iv. 1122 = Lepidium campestre.
hirtum, Linn. Sp. Pl. 641 = Lepidium hirtum.
Huetii, *Boiss. Diagn.* Ser. II. v. 39.—Armenia.
improperum, Jord. Diagn. i. 250 = perfoliatum.
incanum, Gilib. Fl. Lituan. ii. 55 = Lepidium campestre.
infestum, Salisb. Prod. 267 = Capsella Bursa-pastoris.
inornatum, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 145 = perfoliatum.
Ionopsidium, Janka, in Termész. Füzetek, vii. (1883) 124 = Cochlearia acaulis.
Jankae, Kern. in Oestr. Bot. Zeitschr. xvii. (1867) 35 = praecox.
Kerneri, *Huter, l. c.* xxiv. (1874) 82.—Europ. centr.
Korongianum, Czetz, ex Nym. Syll. Suppl. 37 = dacicum.
Kotschyanum, *Boiss. & Hohen. Diagn.* Ser. I. viii. 39.—Persia.
Kovátsii, *Heuff. in Flora*, xxxvi. (1853) 624.—Banat.
latifolium, *Bieb. Fl. Taur.-Cauc.* ii. 99.—Reg. Caucas.
latifolium, Opiz, Natural. n. x. (1825) 253 = arvense.
Lereschii, Reut. in Compt. Rend. Soc. Hallér. (1853–54) 17 = alpestre.
lilacinum, *Boiss. & Huet, Diagn.* Ser. II. v. 41.—As. Min.
longiracemosum, Schur, in Oestr. Bot. Zeitschr. viii. (1858) 22 = Kovátsii.
lotharingum, Jord. Diagn. i. 268 = montanum.
lutescens, Gilib. Fl. Lituan. ii. 54 = arvense.
luteum, Bivona, Cent. Pl. Sic. i. 78 = Bivonaea lutea.
macrophyllum, *Hoffm. in Comm. Soc. Phys. Med. Mosq.* i. (1805) 7.—Reg. Caucas.
magellanicum, *Pers. Syn.* ii. 189.—Reg. Magellan.
marginatum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 90 = Aethionema ovalifolium.
martiale, Jord. Diagn. i. 251 = perfoliatum.
micranthum, Boiss. & Blanche, Diagn. Ser. II. v. 41 = perfoliatum.
microphyllum, *Boiss. & Orph. l. c.* vi. 19.—Graecia.
microstylum, *Boiss. Diagn.* Ser. I. viii. 38.—Syria.
minimum, Ard. Animad. Alt. 33 = alpinum.
montanum, Bory & Chaub. Nouv. Fl. Pélop. 41 = graecum.
montanum, Huds. Fl. Angl. ed. I. 246; ed. II. 282 = alpestre.
montanum, *Linn. Sp. Pl.* 647.—Europ.
montanum, Poir. Encyc. vii. 536 = Lepidium campestre.
multifidum, Poir. l. c. 545 = Lepidium bonariense.
Mureti, *Gremli, Beitr. Fl. Schweiz*, 60.—Helvet.
Nasturtium, Bergeret, ex Steud. Nom. ed. I. 473, in syn. = Lepidium sativum.
natolicum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 180.—As. Min.; Syria.
neglectum, Crépin, in Bull. Acad. Belg. Sér. II. xiv. (1862) 87 = perfoliatum.
nemoricolum, Jord. Diagn. i. 257 = virgatum.
nemorosum, Adam, in Hoffm. Hort. Mosq. (1808) n. 3249; ex DC. Prod. i. 176 = arvense.
nevadense, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 11.—Hispan.
nudicaule, DC. Fl. Fr. iv. 708 = Teesdalia nudicaulis, regularis.
obtusatum, Pomel, Nouv. Mat. Fl. Atl. 376 = perfoliatum.
occitanicum, Jord. Obs. Pl. Crit. iii. (1846) 12 = alpestre.
ochranthum, *F. Muell. ex Benth. Fl. Austral.* i. 88.—Austral.
ochroleucum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 39.—Graec.; As. Min.
oleraceum, Poir. Encyc. vii. 547 = Lepidium oleraceum.

THLASPI :—

oppositifolium, Poir. Encyc. Suppl. v. 277 = Aethionema oppositifolium.
orbiculatum, *Stev. ex DC. Syst.* ii. 377.—Reg. Caucas.
orientale, Willd. ex DC. l. c. 562 = Aethionema cappadocicum.
oxyceras, *Boiss. Diagn.* Ser. I. viii. 37, in syn.—Syria.
papillosum, *Boiss. Fl. Orient.* i. 330.—As. Min.
Pastoraea, Janka, in Termész. Füzetek, vii. (1883) 124 = Bivonaea praecox.
peregrinum, Linn. Sp. Pl. 645 = Aethionema saxatile.
peregrinum, Poir. Encyc. vii. 541 = Aethionema cristatum.
perfoliatum, *Linn. Sp. Pl.* 641.—Europ.; As. bor.
pinnatifidum, Phil. f. Cat. Pl. Vasc. Chil. 19 = Lepidium bonariense.
pinnatum, Noulet, Fl. Sous-Pyr. 61 = Iberis pinnata.
piscidium, Poir. Encyc. vii. 546 = Lepidium piscidium.
platycarpum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vii. 57.—Natolia.
Pollichii, Poir. Encyc. vii. 547 = Lepidium graminifolium.
polymorphum, Gilib. Fl. Lituan. ii. 56 = Capsella Bursa-pastoris.
posteriflorum, Jord. Diagn. i. 250 = perfoliatum.
praecox, Schleich. ex DC. Syst. ii. 380 = alpestre.
praecox, Unger, ex Nym. Consp. 63 = rivale.
praecox, *Wulf. in Jacq. Coll.* ii. 124.—Austria.
procumbens, Lapeyr. Hist. Abr. Pl. Pyr. 366 = Capsella procumbens.
Prolongi, *Boiss. Voy. Bot. Espagne*, ii. 53. t. 14 A.—Hispan.
Psychine, Willd. Sp. Pl. iii. 443 = Psychine stylosa.
pubescens, Guss. in Ind. Sem. Hort. Boccad. (1825); ex ej. Fl. Sic. Syn. ii. 154 = Lepidium nebrodense.
pumilum, *Ledeb. Fl. Ross.* i. 164.—Reg. Caucas.
pygmaeum, Jord. Diagn. i. 252 = rivale.
Revellierii, Bor. Fl. Centr. Fr. ed. III. ii. 60 = perfoliatum.
rhaeticum, Jord. l. c. 259 = alpestre.
rivale, *Presl, Delic. Prag.* 12.—Graec.; Ital.
rosulare, *Boiss. & Balansa, Diagn.* Ser. II. v. 39.—Cappadoc.
Rothianum, Steud. Nom. ed. II. ii. 681 = Aethionema cappadocicum.
rotundifolium, *Gaud. Fl. Helv.* iv. 218.—Europ.
rotundifolium, Tineo, Pl. Rar. Sic. 46 = perfoliatum.
rubellum, Loret & Barr. Fl. Montp. i. 60 = Capsella rubella.
ruderale, All. Fl. Pedem. i. 250 = Lepidium ruderale.
Salisii, Bruegg. in Zeitschr. Ferdinand. Tirol. III. ix. (1860) 45 = alpestre.
salticolum, Jord. Diagn. i. 255 = virgatum.
samolifolium, DC. Syst. ii. 382; (Sp. fict.—Cf. Boiss. Fl. Orient. i. 330).
sativum, Crantz, Stirp. Austr. ed. I. fasc. i. 21; ed. II. 22 = Lepidium sativum.
saxatile, [Habl.] Desc. Taur. 157 = Iberis taurica.
saxatile, Linn. Sp. Pl. 646 = Aethionema saxatile.
scapiflorum, Viv. Prod. Fl. Cors. App. 3 = Lepidium humifusum.
Schrankii, J. F. Gmel. Syst. 974 = Capsella Bursa-pastoris.
spathulatum, Gater. Desc. Pl. Montaub. 115 = montanum.
spinosum, Poir. Encyc. vii. 545 = Lepidium spinosum.
stenocarpum, *Boiss. & Hohen. Diagn.* Ser. I. viii. 38, in syn.—Persia.
stenopterum, Boiss. & Reut. Diagn. Ser. I. viii. 40 = alpestre.
strictum, *Sweet, Hort. Brit.* ed. II. 27.—Persia.
stylosum, Reichb. Ic. Fl. Germ. ii. 3. t. 6 = Noccaea stylosa.
subulatum, Cav. Desc. 414 = Lepidium subulatum.
suecicum, Jord. Diagn. i. 254 = alpestre.
suffruticosum, Asso, ex Losc. & Pard. Ser. Inconf. ed. II. 38 = alpestre ?
sylvestre, Jord. Obs. Pl. Crit. iii. (1846) 9 = alpestre.
sylvium, Gaud. Fl. Helv. iv. 221 = alpinum.
Szowitsianum, *Boiss. Fl. Orient.* i. 324.—Afghan.
tasmanicum, *Hook. f. Fl. Tasman.* i. 23.—Tasman.
taygeteum, Boiss. Diagn. Ser. I. viii. 40 = graecum.
telephiifolium, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 780 = Aethionema saxatile.

THLASPI:—
tenuifolium, Lam. Fl. Fr. ii. 467 = Lepidium ruderale.
Tineanum, Huet, ex Ball, in Journ. Linn. Soc. xvi. (1877) 333 = perfoliatum.
Tinei, Nym. Syll. 205 = perfoliatum.
tuberosum, Nutt. Gen. Am. ii. 65 = Cardamine rhomboidea?
umbellatum, Crantz, Stirp. Austr. ed. I. fasc. i. 26; ed. II. 25 = Iberis umbellata.
umbellatum, *Stev. ex DC. Syst.* ii. 377.—Reg. Casp.
Verloti, Jord. Diagn. i. 256 = virgatum.
Villarsianum, Jord. Obs. Pl. Crit. iii. (1847) 26 = montanum.
violascens, *Schott & Kotschy, ex Unger & Kotschy, Ins. Cyp.* 328; *Boiss. Fl. Orient.* i. 328.—As. Min.; Syria; Ins. Cyprus.
virens, Jord. Obs. Pl. Crit. iii. (1846) 17 = alpestre.
virgatum, *Gren. & Godr. Prosp.* 8 (1846); *ex eor. Fl. Fr.* i. 144.—Mont. Pyren.
virginianum, Poir. Encyc. vii. 544 = Lepidium virginicum.
virginicum, Cav. Desc. 413 = Lepidium virginicum.
vogesiacum, Jord. ex F. Schultz, Arch. Fl. Fr. et Allem. 159 = alpestre.
vulcanorum, Lamotte, Notes Pl. Nouv. 24.—Cf. Bull. Soc. Bot. Fr. vi. (1859) 93 = virgatum.
vulgatum, Roucell, Fl. Nord. Fr. ii. 69 = Lepidium campestre.

THLASPIDEA, Opiz, Seznam, 96 (1852) = **Thlaspi**, Tourn. (Crucif.).
arvensis, Opiz, l. c. = Thlaspi arvense.

THLASPIDIUM, Spach, Hist. Vég. Phan. vi. 556 (1838) = **Lepidium**, Linn. (Crucif.).
sativum, Spach, l. c. 557 = L. sativum.

THLASPIDIUM, Tourn. ex Adans. Fam. ii. 422 (1763) = **Biscutella**, Linn. (Crucif.).
hieracifolium, Medic. Pflanzeng. i. 29 = B. didyma.
laevigatum, Medic. l. c. 30; et in Usteri, Ann. Bot. viii. (1794) 38 = B. laevigata.
saccatum, Moench, Meth. 218 = B. auriculata.
semperflorens, Andrz. ex DC. Syst. ii. 744 (Index), sphalm. = seq.
sempervirens, Andrz. ex DC. l. c. 394 = Iberis semperflorens.

THLASPIUS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = **Thlaspi**, Tourn. (Crucif.).

THLIPSOCARPUS, Kunze, in Flora, xxix. (1846) 695 = **Hyoseris**, Linn. (Compos.).
baeticus, Kunze, l. c. = H. radiata.

THOA, Aubl. Pl. Gui. ii. 874. t. 336 (1775) = **Gnetum**, Linn.
edulis, Willd. Sp. Pl. iv. 477 = G. scandens.
nigra, Hort. ex Carr. Conif. ed. I. 545 = G. nigrum.
pendula, Blanco, Fl. Filip. ed. II. 514 = G. scandens.
urens, Aubl. Pl. Gui. ii. 874 = G. urens.
urens, Poepp. ex Parl. in DC. Prod. xvi. II. 351 = G. amazonicum.

THOMASIA, J. Gay, in Mém. Mus. Par. vii. (1821) 450. tt. 21, 22. *STERCULIACEAE*, Benth. & Hook. f. i. 227.
ASTEROCHITON, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 138.
LEUCOTHAMNUS, Lindl. Swan River App. 19 (1839).
RHYNCHOSTEMON, Steetz, in Lehm. Pl. Preiss. ii. 333 (1847).
aemula, Steud. in Lehm. Pl. Preiss. i. 233 = glutinosa.
angustifolia, *Steud. l. c.* 232.—Austral.
brachystachys, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 143.—Austral.
canescens, Lindl. Swan River App. 18 = glutinosa.
cognata, *Steud. in Lehm. Pl. Preiss.* i. 232.—Austral.
cycnopotamica, Steud. l. c. 231 = grandiflora.
diffusa, G. Don, Gen. Syst. i. 527 = foliosa?
discolor, *Steud. in Lehm. Pl. Preiss.* i. 233.—Austral.
dumosa, Hort. ex Steetz, in Lehm. Pl. Preiss. ii. 356 = Rulingia parviflora.
foliosa, *J. Gay, in Mém. Mus. Par.* vii. (1821) 454. t. 22.—Austral.
Gilbertiana, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 10 = triphylla?
glabrata, Steud. in Lehm. Pl. Preiss. i. 234 = triphylla.

THOMASIA:—
glutinosa, *Lindl. Swan River App.* 18.—Austral.
grandiflora, *Lindl. l. c.*—Austral.
hypoleuca, Steud. in Lehm. Pl. Preiss. i. 234 = quercifolia.
involucrata, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 143 = Lysiosepalum Barryanum.
lasiopetaloides, Steud. in Lehm. Pl. Preiss. i. 233 = glutinosa.
laxiflora, *Benth. Fl. Austral.* i. 256.—Austral.
lucida, Steud. in Lehm. Pl. Preiss. i. 231 = grandiflora.
macrocalyx, Schlecht. in Linnaea, xx. (1847) 633 = petalocalyx.
macrocalyx, *Steud. in Lehm. Pl. Preiss.* i. 230.—Austral.
macrocarpa, *Hueg. in Endl. Nov. Stirp. Dec.* 32.—Austral.
montana, *Steud. in Lehm. Pl. Preiss.* i. 230.—Austral.
paniculata, Lindl. Swan River App. 18 = pauciflora.
pauciflora, *Lindl. l. c.*—Austral.
petalocalyx, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 35.—Austral.
pogonanthera, F. Muell. Fragm. ii. 7 = Guichenotia micrantha.
prostrata, Hort. ex Steetz, in Lehm. Pl. Preiss. ii. 356 = Rulingia parviflora.
pumila, Steud. in Lehm. Pl. Preiss. i. 233 = Guichenotia Sarotes.
purpurea, *J. Gay, in Mém. Mus. Par.* vii. (1821) 452. t. 21.—Austral.
pygmaea, *Benth. Fl. Austral.* i. 257.—Austral.
quercifolia, Bauer, ex Steetz, in Lehm. Pl. Preiss. ii. 326 = discolor.
quercifolia, *J. Gay, in Mém. Mus. Par.* vii. (1821) 459. t. 21.—Austral.
rhynchocarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 142.—Austral.
rugosa, *Turcz. l. c.* xix. (1846) II. 501.—Austral.
rulingioides, *Steud. in Lehm. Pl. Preiss.* i. 232.—Austral.
rupestris, Steud. l. c. 231 = purpurea.
salvifolia, Hook. ex Steetz, in Lehm. l. c. ii. 333 = Rulingia salvifolia.
Sarotes, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 145.—Austral.
solanacea, *J. Gay, in Mém. Mus. Par.* vii. (1821) 456. t. 21.—Austral.
stelligera, *Benth. Fl. Austral.* i. 256.—Austral.
stipulacea, Hook. Bot. Mag. t. 4111 = macrocarpa.
stipulacea, Lindl. Swan River App. 18 = triphylla.
subhastata, Steud. in Lehm. Pl. Preiss. i. 232 = pauciflora.
tenuivestita, *F. Muell. Fragm.* ii. 7.—Austral.
triflora, Walp. Ann. i. 1117 (Index), sphalm. = seq.
triloba, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) II. 500.—Austral.
triphylla, *J. Gay, in Mém. Mus. Par.* vii. (1821) 458.—Austral.
undulata, Steetz, in Lehm. Pl. Preiss. ii. 320 = purpurea.
viridis, Steud. in Lehm. l. c. i. 234 = foliosa.

THOMMASINIA, Steud. Nom. ed. II. ii. 681 (1841), sphalm. = TOMMASINIA, Bertol. = **Peucedanum**, Tourn. (Umbell.).

THOMPSONIA, R. Br. in Trans. Linn. Soc. xiii. (1820) 221 = **Deidamia**, Thou. (Passifl.).
Browniana, M. Roem. Syn. Pepon. 138 = D. Thompsoniana.

THOMPSONIA, Steud. Nom. ed. II. ii. 682 (1841), sphalm. = seq.

THOMSONIA, Wall. Pl. As. Rar. i. 83. t. 99 (1830). *AROIDEAE*, Benth. & Hook. f. iii. 971.
ALLOPHYTHION, Schott, Gen. Aroid. t. 24 (1832).
PYTHONIUM, Schott, Melet. i. 17 (1832).
Hookeri, Engl. in DC. Monog. Phan. ii. 307 = napalensis.
napalensis, *Wall. Pl. As. Rar.* i. 83. t. 99.—Ind. bor. occ.

THONNINGIA, Vahl, in Dansk. Selsk. Skrivt. vi. (1810) 124. t. 6. *BALANOPHOREAE*, Benth. & Hook. f. iii. 236.
CEPHALOPHYTON, Hook. f. ex Baker, in Journ. Linn. Soc. xx. (1883) 250.
CONOPHYTA, Schum. ex Hook. f. in Trans. Linn. Soc. xxii. (1856) 42.
HAEMATOSTROBUS, Endl. Gen. 76, in Adnot. (1836).

THONNINGIA:—
janeirensis, Liebm. in Forhandl. Skand. Naturf. Christ. (1847) 180 = Langsdorffia hypogaea.
mexicana, Liebm. l. c. = Langsdorffia hypogaea.
sanguinea, *Vahl, in Dansk. Selsk. Skrivt.* vi. (1810) 125. t. 6.—Afr. trop.

THORA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 325 = **Ranunculus**, Tourn.
venefica, Fourr. l. c. = R. Thora.

THORACOSPERMA, Klotzsch, in Linnaea, ix. (1834) 350; et xii. (1838) 229 = **Simocheilus**, Klotzsch (Ericac.).
paniculata, Klotzsch, ll. cc. = S. obovatus.

THORACOSTACHYS, Kurz, in Tijdschr. Nederl. Ind. xxvii. (1864) 224 = seq.

THORACOSTACHYUM, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) II. 75. *CYPERACEAE* [Benth. & Hook. f. iii. 1055].
bancanum, *Kurz, l. c.*—Malaya.
sumatranum, *Kurz, l. c.*—Malaya.

THORELIA, Hance, in Journ. Bot. xv. (1877) 268. *LYTHRARIAE*.
deglupta, *Hance, l. c.* 269.—Cochinch.

THORNTONIA, Reichb. Consp. 202 (1828) = **Pavonia**, Cav. (Malvac.).

THORVALDSENIA, Liebm. in Lindbl. Bot. Notiser (1844), 103. *ORCHIDEAE*.
speciosa, *Liebm. l. c.*—Mexic.

THORWALDSENIA, Bot. Zeit. iv. (1846) 396 = praec.

THOTTEA, Rottb. in Nye Dansk. Vidensk. Selsk. Skrift. ii. (1783) 529. t. 2. *ARISTOLOCHIACEAE*, Benth. & Hook. f. iii. 123.
LOBBIA, Planch. in Hook. Lond. Journ. Bot. vi. (1844) 144. t. 3.
dependens, *Klotzsch, in Monatsb. Akad. Berl.* (1859) 589.—Malaya.
grandiflora, *Rottb. in Nye Dansk. Vidensk. Selsk. Skrift.* ii. (1783) 529. t. 2.—Malaya.
macrophylla, *Becc. in Nuov. Giorn. Bot. Ital.* ii. (1870) 5.—Borneo.
rhizantha, *Becc. l. c.* 6.—Borneo.

THOUAREA, Kunth, Enum. Pl. i. 173 (1833) = **Thuarea**, Pers. (Gramin.).

THOUARSIA, Vent. ex DC. Prod. v. 318 (1836) = **Psiadia**, Jacq. (Compos.).

THOUINIA, Linn. f. Suppl. 9, 89 (1781) = **Linociera**, Sw. (Oleac.).
flavicans, Steud. Nom. ed. I. 834, sphalm. = seq.
flavida, Soyer-Willem. in Usteri, Ann. Bot. xviii. (1796) 2 = Noronhia emarginata.
ligustrina, Sw. Prod. Veg. Ind. Occ. 14 = L. ligustrina.
nutans, Linn. f. Suppl. 89 = L. purpurea.

THOUINIA, Poit. in Ann. Mus. Par. iii. (1804) 70. tt. 6, 7. *SAPINDACEAE*, Benth. & Hook. f. i. 400.
CARPIDOPTERIX, Karst. Fl. Columb. ii. 45. t. 123 (1860?).
THINOUIA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 368.
THOUINIDIUM, Radlk. in Sitzb. Akad. Muench. viii. (1878) 267, 281.
THYANA, Ham. Prod. Pl. Ind. Occ. 36 (1825).
VARGASIA, Bert. ex Spreng. Syst. ii. 283 (1825).
adenophora, Miq. in Lehm. Pl. Preiss. i. 224 = Dodonaea adenophora.
australis, A. Rich. Sert. Astrolab. ii. 31. t. 12 = Atalaya salicifolia.
canescens, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 281.—Cuba.
decandra, *Humb. & Bonpl. Pl. Aequin.* i. 198. t. 56.—Mexic.
dicarpa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 587.—Afr. trop.

THOUINIA:—
dioica, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 21. t. 4 = Schmidelia heterophylla.
discolor, *Griseb. Fl. Brit. W. Ind.* 127.—Ins. Baham.
elliptica, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 280.—Cuba.
hemiglauca, F. Muell. Fragm. i. 98 = Atalaya hemiglauca.
integrifolia, *Spreng. Neue Entdeck.* ii. 155.—Bras.
macroptera, *Casar. Nov. Stirp. Bras. Dec.* 45.—Am. trop.
mauritiana, Boj. Hort. Maurit. 56 = Hornea mauritiana.
Morisiana, Casar. Nov. Stirp. Bras. Dec. 75 = Pausandra Morisiana.
myriantha, *C. Muell. in Walp. Ann.* vii. 628.—N. Granat.
nervosa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 169.—Cuba.
ornifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 82.—Reg. Argent.
patentinervis, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 280.—Cuba.
pinnata, *Turpin, in Ann. Mus. Par.* v. (1804) 401. t. 26.—Ins. S. Doming.
polygama, *G. F. W. Mey. Prim. Fl. Esseq.* 156.—Guian.
pulverulenta, *Griseb. Cat. Pl. Cub.* 46.—Cuba.
punctata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 281.—Cuba.
rotundata, *Wright, in Sauv. Fl. Cub.* 25.—Cuba.
scandens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 284.—Bras.
serrata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 280.—Mexic.
simplicifolia, *Poit. in Ann. Mus. Par.* iii. (1804) 71. t. 6.—Ins. S. Doming.
tomentosa, *DC. Prod.* i. 612.—Ins. S. Doming.
trifoliata, *Poit. in Ann. Mus. Par.* iii. (1804) 72; v. (1804) 401, 402. t. 27.—Ins. S. Doming.
variifolia, F. Muell. Fragm. i. 45 = Atalaya variifolia.
villosa, [*Moç. & Sesse*], *ex DC. Prod.* i. 612.—Mexic.
weinmannifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 81.—Am. austr.

THOUINIA, Sm. Ic. Ined. t. 7 (1789) = **Humbertia**, Lam. (Convolv.).
spectabilis, Sm. l. c. = H. madagascariensis.

THOUINIDIUM, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 267, 281 = **Thouinia**, Poit. (Sapind.).
decandrum, Radlk. l. c. 284 = Thouinia decandra.
oblongum, Radlk. l. c.—Mexic.
pinnatum, Radlk. l. c. 283 = Thouinia pinnata.
pulverulentum, Radlk. l. c. 284 = Thouinia pulverulenta.

THOZETIA, F. Muell. ex Benth. Fl. Austral. iv. 347 (1869). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 776.
racemosa, *F. Muell. ex Benth. Fl. Austral.* iv. 347.—Austral.

THRASIA, Kunth, Syn. Pl. Aequin. i. 191 (1822), sphalm. = seq.

THRASYA, H. B. & K. Nov. Gen. et Sp. i. 120. t. 39 (1815) = **Panicum**, Linn. (Gramin.).
cultrata, Nees, in Mart. Fl. Bras. ii. 95 = P. monostachyum.
hirsuta, Nees, l. c. 94 = P. thrasyoides.
paspaloides, H. B. & K. Nov. Gen. et Sp. i. 121. t. 39 = P. Thrasya.

THRAULOCOCCUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 246 = **Sapindus**, Tourn.
erectus, Radlk. l. c. 309 = S. erectus.
simplicifolius, Radlk. l. c. = S. Thwaitesii.

THRELKELDIA, R. Br. Prod. 409 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 62.
OSTEOCARPUM, F. Muell. Second Gen. Rep. 15 (1854).
brevicuspis, *F. Muell. ex Benth. Fl. Austral.* v. 198.—Austral.
diffusa, *R. Br. Prod.* 410.—Austral.

THRELKELDIA:—
haloragoides, *F. Muell. ex Benth. Fl. Austral.* v. 198.—Austral.
proceriflora, *F. Muell. Fragm.* viii. 38.—Austral.
salsuginosa, *F. Muell. l. c.* vii. 12.—Austral.

THRICA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 428 (1821) THRINCIA, Roth = **Leontodon**, Linn. (Compos.).

THRICHOSTEMA, Linn. Coroll. Gen. 10 (1737), sphalm. = **Trichostema**, Gronov. (Labiat.).

THRINAX, Linn. f. ex Sw. Prod. Veg. Ind. Occ. 57 (1788). *PALMAE*, Benth. & Hook. f. iii. 930.
POROTHRINAX, H. Wendl. ex Griseb. Cat. Pl. Cub. 221 (1866).
argentea, *Lodd. ex Desf. Cat. Hort. Par.* ed. III. 31.—Panama.
aurantia, *Fulch. ex Schult. f. Syst.* vii. 1301.—Hab. ?
aurata, Hort. ex Kunth, Enum. Pl. iii. 254 = radiata.
barbadensis, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 320.—Ins. Barbad.
brasiliensis, Mart. l. c. t. 104 = Trithrinax brasiliensis.
Chuco, Mart. l. c. 320 = Acanthorhiza Chuco.
crinita, *Griseb. & H. Wendl. ex Kerch. Palm.* 258.—Cuba.
elegans, Hort. ex Schult. f. Syst. vii. 1301 = radiata.
excelsa, Lodd. ex Mart. Hist. Nat. Palm. iii. 320 (Quid ?).—Guian.
ferruginea, Lodd. ex Mart. l. c. (Quid ?).—Jamaic.
Garberi, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 12.—Florida.
gracilis, Hort. ex Schult. f. Syst. vii. 1301 = radiata.
graminifolia, Hort. ex H. Wendl. Cycl. Pand. Cycad. 39; Illustr. Hortic. xxxi. (1884) t. 542 = multiflora.
grandis, Hort. ex Kerch. Palm. 258 (Quid ?).—Cuba.
havanensis, *Hort.—Cf. Gard. Chron.* (1870) 37.—Ind. occ.
maritima, Lodd. ex Mart. Hist. Nat. Palm. iii. 320 (Quid ?).—Cuba.
Martii, Griseb. Cat. Pl. Cub. 221 = parviflora.
mexicana, Lodd. ex Mart. Hist. Nat. Palm. iii. 320 (Quid ?).—Mexic.
Miraguana, *Mart. Hist. Nat. Palm.* iii. 320.—Cuba.
montana, Lodd. ex Mart. l. c. (Quid ?).—Cuba.
multiflora, *Mart. l. c.* t. 103. f. 1; t. 163.—Ins. S. Doming.
parviflora, Maycock, Fl. Barbad. 146 = barbadense.
parviflora, *Sw. Prod. Veg. Ind. Occ.* 57.—Jamaic.; Ins. S. Doming.
pumila, Fulch. ex Schult. f. Syst. vii. 1301 = T. Pumilio.
Pumilio, *Lodd. ex Schult. f. l. c.*—Hab. ?
radiata, *Lodd. ex Desf. Cat. Hort. Par.* ed. III. 31.—Ins. Trinit.
rigida, *Griseb. Cat. Pl. Cub.* 221.—Cuba.
stellata, Lodd. ex Mart. Hist. Nat. Palm. iii. 320 = T. Miraguana.
tunicata, *Linden, in Illustr. Hortic.* xxviii. (1881) 32, nomen.—Cuba.
Yuraguana, A. Rich. Fl. Cub. Fanerog. ii. 278 = T. Miraguana.

THRINCIA, Roth, in Roem. Arch. i. 1. 36 (1796); Catalect. fasc. i. 97 (1797) = **Leontodon**, Linn. (Compos.).
arenaria, Martr. Fl. Tarn, 396 = L. hirtus.
caulescens, Trev. ex Spreng. Syst. iv. Cur. Post. 304 = Picris radicata.
glabra, Schleich. Cat. Pl. Helvet. ed. III. 29 = L. hirtus.
glabrata, Hoffmgg. & Link, Fl. Port. ii. 159.—Lusitan.
grumosa, Brot. Fl. Lusit. i. 325 = L. tuberosus.
hirta, Roth, Catalect. fasc. i. 98; DC. Fl. Fr. iv. 51 = L. hirtus.
hispida, DC. l. c. 52 = L. hirtus.
hispida, Roth, Catalect. fasc. i. 99 = L. saxatilis.
laevis, Lag. Gen. et Sp. Nov. 24 = L. hirtus.
leontodontoides, Wallr. in Linnaea, xiv. (1840) 662 (Beitr. Fl. Hercyn. 292).—Europ. centr.
Leyseri, Wallr. Sched. Crit. 441 = L. hirtus.
major, Peterm. in Flora, xxvii. (1844) 475 = L. hirtus.

THRINCIA:—
maroccana, Pers. Syn. ii. 368 = Leontodon maroccanus.
mauritanica, Spreng. Syst. iii. 666 = L. saxatilis, maroccanus.
Muelleri, Nym. Syll. 54 = L. Salzmanni.
nudicalyx, Lag. Gen. et Sp. Nov. 24 = L. hirtus.
nudicaulis, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28 = L. saxatilis.
orientalis, Nym. Syll. Suppl. 14 = L. hispidulus.
psilocalyx, Lag. ex Reichb. Ic. Pl. Crit. viii. 20. t. 749. f. 995. = L. hirtus.
psilostachya, Lag. ex Webb & Berth. Phyt. Canar. ii. 468 = praec.
pygmaea, Pers. Syn. ii. 369 = L. saxatilis.
Salzmanni, Nym. Consp. 470 = L. Salzmanni.
taraxacoides, Gaud. Fl. Helv. v. 49 = L. saxatilis.
tingitana, Boiss. & Reut. Pugill. Pl. Nov. 68 = L. tingitanus.
tuberosa, DC. Fl. Fr. iv. 52 = L. tuberosus.

THRIXIA, Dulac, Fl. Hautes-Pyr. 495 (1867) = **Leontodon**, Linn. (Compos.).
hirta, Dulac, l. c. 496 = L. hirtus.

THRIXSPERMUM, Lour. Fl. Cochinch. 519 (1790) = **Sarcochilus**, R. Br. (1810) (Orchid.).
acuminatissimum, Reichb. f. Xenia Orch. ii. 121 = S. acuminatissimus.
adversum, Reichb. f. l. c. 122 = S. adversus.
amplexicaule, Reichb. f. l. c. 121 = S. amplexicaulis.
anceps, Reichb. f. l. c. 122 = S. anceps.
angustifolium, Reichb. f. l. c. = S. angustifolius.
Arachnites, Reichb. f. l. c. 121 = S. Arachnites.
auriferum, Reichb. f. l. c. = S. aurifer.
australe, Reichb. f. Beitr. Syst. Pfl. 71 = S. parviflorus.
Berkeleyi, Reichb. f. in Gard. Chron. (1882) i. 557 = S. Berkeleyi.
Calceolus, Reichb. f. Xenia Orch. ii. 122 = S. Calceolus.
Ceciliae, Reichb. f. Beitr. Syst. Pfl. 71 = S. Ceciliae.
Centipeda, Lour. Fl. Cochinch. 520.—Cochinch.
compressum, Reichb. f. Xenia Orch. ii. 121 = S. compressus.
croceum, Reichb. f. l. c. 122 = S. croceus.
dilatatum, Reichb. f. l. c. = S. olivaceus.
divitiflorum, Reichb. f. in Gard. Chron. (1877) i. 750 = S. divitiflorus.
falcatum, Reichb. f. Xenia Orch. ii. 122 = S. falcatus.
Freemanii, Reichb. f. in Gard. Chron. (1877) i. 750 = S. Freemanii (in Addend.).
Godeffroyanum, Reichb. f. Xenia Orch. ii. 122.—Ins. Fiji.
gracilentum, Reichb. f. l. c. 121 = S. gracilentus.
Graeffei, Reichb. f. in Seem. Fl. Vit. 297.—Ins. Samoa.
Hartmanni, Reichb. f. in Gard. Chron. (1877) i. 716 = S. Hartmanni.
Hillii, Reichb. f. Beitr. Syst. Pfl. 71 = S. Hillii.
Hystrix, Reichb. f. in Trans. Linn. Soc. xxx. (1874) 145 = S. Hystrix.
japonicum, Reichb. f. in Bot. Zeit. xxxvi. (1878) 75 = S. japonicus.
leopardinum, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 145.—Burma.
lilacinum, Reichb. f. Xenia Orch. ii. 121 = S. lilacinus.
luniferum, Reichb. f. in Gard. Chron. (1868) 786.—Burma.
Moorei, Reichb. f. l. c. (1880) i. 104 = S. Moorei (in Addend.).
muriculatum, Reichb. f. l. c. (1881) ii. 198 = S. muriculatus (in Addend.).
obtusum, Reichb. f. Xenia Orch. ii. 121 = S. obtusus.
olivaceum, Reichb. f. l. c. 122 = S. olivaceus.
pallidum, Reichb. f. l. c. = S. pallidus.
parviflorum, Reichb. f. l. c. = S. parviflorus.
phyllorrhizum, Reichb. f. Beitr. Syst. Pfl. 71 = S. phyllorrhizus.
pictum, Reichb. f. Xenia Orch. ii. 122 = S. olivaceus.
purpurascens, Reichb. f. l. c. 121 = S. purpurascens.
rhopalorrhachis, Reichb. f. l. c. = S. rhopalorrhachis.
serraeforme, Reichb. f. l. c. 121 = S. Arachnites.
Sillemianum, Reichb. f. in Gard. Chron. (1882) i. 524 = S. Sillemianus.
spurium, Reichb. f. Xenia Orch. ii. 122 = S. spurius.
subulatum, Reichb. f. l. c. = S. subulatus.
teres, Reichb. f. l. c. 121 = S. teres.

THRIXSPERMUM :—
Teysmanni, Reichb. f. Xenia Orch. ii. 121 = Aërides Teysmanni.
tridentatum, Reichb. f. l. c. 122 = Cleisostoma tridentatus.
unguiculatum, Reichb. f. l. c. = Sarchochilus unguiculatus.
usneoides, Reichb. f. l. c. 120 = S. usneoides.
Zollingeri, Reichb. f. l. c. 121 = Aërides Zollingeri.

THRYALLIS, Linn. Sp. Pl. ed. II. 554 (1762) = **Galphimia**, Cav. (Malpig.).
brasiliensis, Linn. l. c. = G. brasiliensis.
macroptera, Mart. in Flora, xxiv. (1841) II. Beibl. 104 = Dicella macroptera.

THRYALLIS, Mart. Nov. Gen. et Sp. iii. 78. tt. 230, 231 (1829). *MALPIGHIACEAE*, Benth. & Hook. f. i. 254.
brachystachis, *Lindl. Bot. Reg.* t. 1162.—Bras.
latifolia, *Mart. Nov. Gen. et Sp.* iii. 79. t. 231.—Bras.
longifolia, *Mart. l. c.* 78. t. 230.—Bras.
rotundifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 264.—Bras.

THRYCOCEPHALUM, Steud. Nom. ed. II. ii. 682 (1841) = seq.

THRYOCEPHALON, Forst. Char. Gen. 129. t. 65 (1776) = **Kyllinga**, Rottb. (Cyper.).
nemorale, Forst. l. c. 130 = K. monocephala.

THRYPTOMENE, Endl. in Ann. Wien. Mus. ii. (1838) 192. *MYRTACEAE*, Benth. & Hook. f. i. 700.
ASTRAEA, Schau. in Linnaea, xvii. (1843) 238.
BUCHERIA, Heynh. Nom. ii. 80 (1846).
EREMOPYXIS, Baill. Adansonia, ii. (1861–62) 328.
GOMPHOTIS, Rafin. Sylva Tellur. 103 (1838).
PARYPHANTHA, Schau. in Linnaea, xvii. (1843) 235.
auriculata, *F. Muell. Fragm.* x. 24.—Austral.
australis, *Endl. in Ann. Wien. Mus.* ii. (1838) 192.—Austral.
baeckeacea, *F. Muell. Fragm.* iv. 65.—Austral.
ciliata, *F. Muell. in Woolls, Pl. Neighb. Sydney*, 23.—Austral.
denticulata, *Benth. Fl. Austral.* iii. 60.—Austral.
Drummondii, F. Muell. Census, 53 = Micromyrtus Drummondii.
Elliottii, *F. Muell. Fragm.* ix. 62.—Austral.
elobata, F. Muell. l. c. iv. 63 = Micromyrtus elobata.
ericaea, *F. Muell. l. c.* i. 12.—Austral.
flaviflora, *F. Muell. l. c.* viii. 13.—Austral.
Homalocalyx, F. Muell. l. c. iv. 63 = Homalocalyx ericaeus.
hymenonema, *F. Muell. l. c.* x. 26.—Austral.
hyporhytis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxv. (1862) II. 324.—Austral.
imbricata, F. Muell. Census, 53 = Micromyrtus imbricata.
Johnsonii, *F. Muell. Fragm.* iv. 77.—Austral.
leptocalyx, F. Muell. Census, 53 = Micromyrtus imbricata.
Maisonneuvii, *F. Muell. Fragm.* iv. 64.—Austral.
micrantha, *Hook. f. in Hook. Kew Journ.* v. (1853) 299.—Tasman.
minutiflora, F. Muell. in Woolls, Pl. Neighb. Sydney, 23 = Micromyrtus minutiflora.
Miqueliana, *F. Muell. Fragm.* i. 11.—Austral.
Mitchelliana, *F. Muell. l. c.*—Austral.
mucronulata, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 156.—Austral.
obovata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 322.—Austral.
oligandra, *F. Muell. Fragm.* i. 11.—Austral.
plicata, F. Muell. l. c. iv. 63 = Micromyrtus microphylla.
polyandra, F. Muell. l. c. 77 = Homalocalyx polyandrus.
prolifera, *Turcz. in Bull. Soc. Nat. Mosc.* xxxv. (1862) II. 324.—Austral.
racemosa, F. Muell. Census, 53 = Micromyrtus racemosus.
racemulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 156.—Austral.
saxicola, Miq. Stirp. Nov. Holl. 20 = Miqueliana.

THRYPTOMENE :—
saxicola, *Schau. in Lehm. Pl. Preiss.* i. 102.—Austral.
stenocalyx, *F. Muell. Fragm.* x. 23.—Austral.
strongylophylla, *F. Muell. ex Benth. Fl. Austral.* iii. 61.—Austral.
tenella, *Benth. l. c.* 59.—Austral.
trachycalyx, *F. Muell. Fragm.* x. 25.—Austral.
urceolaris, *F. Muell. l. c.*—Austral.

THUAREA, Pers. Syn. i. 110 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1109.
MICROTHUAREIA, Thou. Gen. Nov. Madag. 3 (1806).
ORNITHOCEPHALOCHLOA, Kurz, in Journ. Bot. xiii. (1875) 332. t. 171.
THOUAREA, Kunth, Enum. Pl. i. 173 (1833).
involucrata, Eschsch. in Mém. Acad. Pétersb. x. (1826) 291 = sarmentosa.
involuta, R. Br. Prod. 197 = sarmentosa.
latifolia, R. Br. l. c. = sarmentosa.
media, R. Br. l. c. = sarmentosa.
sarmentosa, *Pers. Syn.* i. 110.—Geront. trop.

THUESSINKIA, Korth. ex Miq. Fl. Ind. Bat. iii. 41 (1855) = **Caryota**, Linn. (Palm.).
speciosa, Korth. l. c. = C. mitis.

THUIA, Scop. Introd. 353 (1777) = **Thuya**, Linn. (Conif.)

THUIACARPUS, Benth. & Hook. f. Gen. iii. 428, (1880), sphalm. = seq.

THUIAECARPUS, Trautv. Imag. Pl. Ross. 11. t. 6 (1844) = **Juniperus**, Tourn. (Conif.).
juniperinus, Trautv. l. c. (forma monstrosa) = J. communis.

THUIOPSIS, Endl. Syn. Conif. 53 (1847) = THUJOPSIS, Sieb. & Zucc. = **Thuya**, Linn. (Conif.).

THUJA, Linn. Hort. Cliff. 449 (1737) = **Thuya**, Linn. (Conif.).

THUJOPSIS, Sieb. & Zucc. Fl. Jap. ii. 32. tt. 119, 120 (1844) = **Thuya**, Linn. (Conif.).
borealis, Hort.—Cf. Gard. Chron. (1854) 727; et ex Carr. Conif. ed. I. 113 = Cupressus nootkatensis.
cupressoides, Carr. in Jacques & Hérincq, Man. Pl. iv. 324 = Cupressus nootkatensis.
dolabrata, Hort. ex Carr. Conif. ed. II. 99 = Thuya orientalis.
dolabrata, Sieb. & Zucc. Fl. Jap. ii. 34. tt. 119, 120 = Thuya dolabrata.
laetevirens, Lindl. in Gard. Chron. (1861) 56 = Thuya dolabrata.
Standishi, Gord. Pinet. Suppl. 100 = Thuya japonica.
Tchugatskoyae, Hort. ex Carr. Conif. ed. I. 113 = Cupressus nootkatensis.
Tschugatskoi, Hort. ex C. Koch, Dendrol. ii. II. 166 = praec.

THUMBERGIA, Poit. in Rev. Hortic. Sér. II. iv. (1845–46) 409, sphalm. = **Thunbergia**, Retz. (Acanth.).

THUMUNG, Koen. in Retz. Obs. iii. 62 (1783) = **Zingiber**, Adans. (Scitam.).

THUNBERGIA, Montin, in Vet. Akad. Handl. Stockh. (1773) 288. t. 11 = **Gardenia**, Ellis (Rubiac.).
capensis, Linn. f. Suppl. 162, in syn. = G. Thunbergia.

THUNBERGIA, Retz. in Phys. Sällsk. Handl. i. (1776) 168. *ACANTHACEAE*, Benth. & Hook. f. ii. 1072.
ENDOMELAS, Rafin. Fl. Tellur. iv. 67 (1836).
HEXACENTRIS, Nees, in Wall. Pl. As. Rar. iii. 78 (1832).
MEYENIA, Nees, l. c. (1832).
PLEIMERIS, Rafin. Sylva Tellur. 21 (1838).
PLEUREMIDIS, Rafin. Fl. Tellur. iv. 67 (1836).
SCHMIDIA, Wight, Ic. t. 1848 (1852).
adenocalyx, *Radlk. in Bremen, Abh.* viii. (1883) 431.—Afr. trop.
affinis, *S. Moore, in Journ. Bot.* xviii. (1880) 5.—Afr. trop.
alata, *Boj. ex Sims, Bot. Mag.* t. 2591; *et ex Hook. Exot. Fl.* t. 177.—Afr. trop.; Ind. or. (cult.).

THUNBERGIA :—
albiflora, Gord. in Gard. Chron. (1845) 169 = alata.
angolensis, *S. Moore, in Journ. Bot.* xviii. (1880) 195.—Afr. trop.
angulata, R. Br. in Salt, Abyss. App. 65 = Saltiana.
angulata, *Hilsenb. & Boj. ex Hook. Exot. Fl.* iii. t. 166; t. 177. ff. 1–5.—Madag.
angustifolia, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. iii. 78 = fragrans.
annua, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 43, nomen; *et ex Nees, in DC. Prod.* xi. 55.—Afr. trop.
armipotens, *S. Moore, in Journ. Bot.* xviii. (1880) 195.—Afr. trop.
arnhemica, F. Muell. Fragm. ix. 73 = fragrans.
aspera, *Nees, in DC. Prod.* xi. 56.—Afr. austr.
atriplicifolia, *E. Mey. in Drège, Zwei Pfl. Docum.* 226, nomen; *et ex Nees, l. c.*—Afr. austr.
aurantiaca, Paxt. Mag. Bot. vi. (1839) 269 = alata.
Backerii, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 919 = alata.
capensis, *Retz. in Phys. Sällsk. Handb.* i. (1776) 163.—Afr. austr.
cerinthoides, *Radlk. in Bremen, Abh.* viii. (1883) 434.—Afr. trop.
chrysops, *Hook. Bot. Mag.* t. 4119.—Afr. trop.
coccinea, *Wall. Tent. Fl. Nap.* i. 48. t. 37; *R. Grah. in Edinb. Phil. Journ.* xiv. (1826) 150.—Reg. Himal.; Burma.
coerulea, Wight, ex Nees, in Wall. Pl. As. Rar. iii. 78 = Hawtayneana.
convolvulifolia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 428.—Madag.
cordata, Colla, Hort. Ripul. 139. t. 21 = fragrans.
cordifolia, Herb. Madr. ex Wall. Cat. n. 769 = Hawtayneana.
cordifolia, Nees, in DC. Prod. xi. 55 = grandiflora.
cyanea, *Boj. ex Nees, l. c.*—Madag.
Cycnium, *S. Moore, in Journ. Bot.* xviii. (1880) 194.—Afr. trop.
cynanchifolia, *Benth. in Hook. Niger Fl.* 475.—Afr. trop.
Diplocalymna, Steud. Nom. ed. II. ii. 683 = Diplocalymna volubile.
Doddsii, Paxt. Mag. Bot. xv. (1849) t. 221 = alata.
Dregeana, *Presl, Bot. Bemerk.* 94.—Afr. austr.
erecta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 18.—Afr. trop.
erecta, Herb. Madr. ex Nees, in Wall. Pl. As. Rar. iii. 78 = Hawtayneana.
fastuosa, *Poit. in Rev. Hortic.* Sér. II. iv. (1846) 409.—Mexic.
flava, Hoffmgg. Preisverz. (1841) 81.—Cf. Linnaea, xvi. (1842) Litt. 273 = fragrans.
fragrans, Drège, ex Presl, Bot. Bemerk. 94 = Dregeana.
fragrans, *Roxb. Pl. Corom.* i. 47. t. 67.—Ind. or.; Malaya; Austral.
fragrans, Wall. Cat. n. 767, partim = tomentosa.
frangulaefolia, Zipp. ex Span. in Linnaea, xv. (1841) 329 = hastata.
Fryeri, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 876 = alata.
gentianoides, *Radlk. in Bremen, Abh.* viii. (1883) 433.—Afr. trop.
geraniifolia, *Benth. in Hook. Niger Fl.* 475.—Afr. trop.
grandiflora, *Roxb. Hort. Beng.* 45; *Fl. Ind.* iii. 34.—Ind. or.; Burma.
grandiflora, Wall. Cat. n. 768, partim = laurifolia.
Harrisii, Hook. f. Bot. Mag. t. 4998 = laurifolia.
hastata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 388.—Ins. Timor.
Hawtayneana, *Wall. Tent. Fl. Nap.* 49, in obs.—Ind. or.
Hawtaynii, T. Anders. in Journ. Linn. Soc. ix. (1867) 448 = praec.
heterophylla, Wall. Cat. n. 771 = fragrans.
hirsuta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 20.—Abyss.
hirta, Sond. in Linnaea, xxiii. (1850) 89 = neglecta.
hispida, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 113.—Afr. trop.
huillensis, *S. Moore, in Journ. Bot.* xviii. (1880) 194.—Afr. trop.
humilis, Eckl. & Zeyh. ex Nees, in DC. Prod. xi. 55 = capensis.

THUNBERGIA:—
hyalina, *S. Moore, in Journ. Bot.* xviii. (1880) 195.—Afr. trop.
javanica, Gaertn. f. Fruct. iii. 22 = fragrans.
Kirkiana, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 19.—Afr. trop.
Kirkii, *Hook. f. Bot. Mag.* t. 6677.—Afr. trop.
laevis, Wall. & Nees, in Wall. Pl. As. Rar. iii. 77 = fragrans.
lancifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 19.—Afr. trop.
laurifolia, *Lindl. in Gard. Chron.* (1856) 260.—Malaya.
leucantha, Hort. ex Steud. Nom. ed. II. ii. 683 (Quid?).—Hab.?
longiflora, F. Muell. Fragm. ix. 73, in obs. = fragrans.
lutea, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 448.—Reg. Himal.
malvifolia, Buch.-Ham. ex Wall. Cat. n. 768 = grandiflora.
mysorensis, *T. Anders. ex Bedd. in Trans. Linn. Soc.* xxv. (1865) 222; *et in Journ. Linn. Soc.* ix. (1867) 448.—Ind. or.
natalensis, *Hook. Bot. Mag.* t. 5082.—Afr. austr.
neglecta, *Sond. in Linnaea*, xxiii. (1850) 89.—Afr. austr.
oblongifolia, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 125.—Afr. trop.
pendula, Hassk. Cat. Hort. Bog. Alt. 147 = coccinea.
platyphylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 217.—Madag.
Powelli, *F. Muell. Census*, 99.—Austral.
quinquenervis, Buch.-Ham. ex Nees, in Wall. Pl. As. Rar. iii. 78 = coccinea.
repens, Pers. Syn. ii. 179 = Lepidagathis repens.
reticulata, *Hochst. ex Nees, in DC. Prod.* xi. 58.—Abyss.
Roxburghia, Nees, in Wall. Pl. As. Rar. iii. 78 = fragrans.
Saltiana, *Steud. Nom.* ed. II. ii. 688.—Abyss.
scandens, Pers. ex Nees, in DC. Prod. xi. 57 = fragrans.
Schweinfurthii, *S. Moore, in Journ. Bot.* xviii. (1880) 6.—Afr. trop.
smilacifolia, *Kurz, For. Fl. Brit. Burma*, ii. 241.—Burma.
stolonifera, Blanco, Fl. Filip. ed. I. 517 = Herpestis Monnieria.
subsagittata, Blanco, l. c. 518 = fragrans.
tomentosa, *Wall. ex Nees, in Wall. Pl. As. Rar.* iii. 78.—Ind. or.
Vogeliana, *Benth. in Hook. Niger Fl.* 476.—Afr. trop.
volubilis, Pers. Syn. ii. 179 = fragrans.
Wightiana, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 448.—Ind. or.

THUNIA, Reichb. f. in Bot. Zeit. x. (1852) 764 = **Phaius**, Lour. (Orchid.).
alba, Reichb. f. l. c. = P. albus.
Bensoniae, Hook. f. Bot. Mag. t. 5694 = P. albus.
Marshalliana, Reichb. f. in Linnaea, xli. (1877) 65 = P. albus.
pulchra, Reichb. f. in Flora, lv. (1872) 276 = P. albus.
Veitchiana ×, Reichb. f. in Gard. Chron. (1885) i. 818.

THURARIA, Molina, Sagg. Chile, 159 (1782). Genus dubium *EBENACEARUM*.—Cf. Benth. & Hook. f. ii. 663.
Turaria, Spreng. Syst. ii. 260 (1825).
chilensis, *Molina, Sagg. Chile*, 159.—Chili.

THURARIA, [Nutt.] in Fras. Cat. (1813) n. 85 = **Grindelia**, Willd. (Compos.).
herbacea, [Nutt.] l. c. = G. squarrosa.

THURBERIA, Benth. in Journ. Linn. Soc. xix. (1881) 58. *GRAMINEAE*, Benth. & Hook. f. iii. 1118.
Greenia, Nutt. in Trans. Am. Phil. Soc. v. (1837) 142.
Sclerachne, Torr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) 273.
arkansana, *Benth. ex Vasey, Gr. U. St.* 16.—Am. bor. occ.
pilosa, Vasey, l. c. = arkansana.

THURBERIA, A. Gray, in Mem. Am. Acad. N. S. v. (1855) 308 = **Gossypium**, Linn. (Malvac.).
thespesioides, A. Gray, l. c. = G. Thurberi.

THURNHAUSERA, Pohl, ex G. Don, Gen. Syst. iv. 201 (1837) = **Curtia**, Cham. & Schlecht. (Gentian.).

THURNHEYSSERA, Mart. ex Meissn. in DC. Prod. xiv. 185 (1856) = **Symmeria**, Benth. (Polygon.).
Tragopyrum, Mart. ex Meissn. l. c. 186 = S. paniculata.

THURNIA, Hook. f. in Hook. Ic. Pl. tt. 1407, 1408 (1883). *JUNCACEAE*, Benth. & Hook. f. iii. 869.
Jenmani, *Hook. f. in Hook. Ic. Pl.* t. 1408.—Guian.
sphaerocephala, *Hook. f. l. c.* t. 1407.—Guian.

THURYA, Boiss. & Bal. Diagn. Ser. II. v. 63 (1856). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 978.
capitata, *Boiss. & Bal. l. c.* 64.—As. Min.

THUYA, Linn. Syst. ed. I (1735). *CONIFERAE*, Benth. & Hook. f. iii. 426.
Biota, D. Don, in Lamb. Pin. ed. II. ii. 129 (1828); Endl. Syn. Conif. 46 (1847).
Chamaecyparis, Spach, Hist. Vég. Phan. xi. 329 (1842).
Chamaepeuce, Zucc. in Endl. Ench. 139 (1841).
Platycladus, Spach, Hist. Vég. Phan. xi. 333 (1842).
Retinispora, Sieb. & Zucc. Fl. Jap. ii. 36. tt. 121–123 (1842), partim.
Thuja, Linn. Hort. Cliff. 449 (1737).
Thujopsis, Sieb. & Zucc. Fl. Jap. ii. 32. tt. 119, 120 (1844).
acuta, Moench, Meth. 692 = orientalis.
andina, Poepp. & Endl. Nov. Gen. et Sp. iii. 17. t. 220 = Libocedrus chilensis.
antarctica, Hort. ex Gord. Pinet. Suppl. 102 = orientalis.
aphylla, Burm. f. Prod. Fl. Cap. 27 = Callitris cupressoides.
aphylla, Linn. Cent. Pl. i. 32; Amoen. Acad. iv. 295 = Tamarix articulata.
argentea, Hort. ex Carr. Conif. ed. I. 94 = orientalis.
articulata, Vahl, Symb. Bot. ii. 96 = Callitris quadrivalvis.
asplenifolia, Hort. ex Carr. Conif. ed. II. 106 = plicata.
aurea, Hort. ex Carr. l. c. ed. I. 96 = orientalis.
australis, Bosc, ex Desf. Tabl. ed. II. 274, nomen; Poir. Encyc. Suppl. v. 302 = Callitris rhomboidea.
australis, Hort. ex Tenore, in Mém. Acc. Sc. Borb. Nap. iii. ii. (1832) 35 = orientalis.
californica, Hort. ex Carr. Conif. ed. II. 106 = plicata.
californica, Hort. ex C. Koch, Dendrol. ii. ii. 177 = gigantea.
canadensis, Hort. ex C. Koch, l. c. 175 = occidentalis.
caucasica, Hort. ex Gord. Pinet. Suppl. 103 = occidentalis.
chilensis, D. Don, in Lamb. Pin. ed. 8vo. 128. h. 62; Hook. Lond. Journ. Bot. ii. (1843) 199 = Libocedrus chilensis.
compacta, Standish, ex Gord. Pinet. Suppl. 103 = occidentalis.
Craigana, A. Murr. Oreg. Exp. 2. [t. 5] = Libocedrus decurrens.
cristata, Hort. ex Carr. Conif. ed. II. 97 = orientalis.
cristata, Hort. ex Carr. l. c. 111 = occidentalis.
cuneata, Domb. ex Endl. Syn. Conif. 45 = Libocedrus chilensis.
cupressoides, Hort. ex Carr. Conif. ed. II. 97 = orientalis.
cupressoides, Linn. Mant. i. 125 = Callitris cupressoides.
curviramea, Miq. Cat. Hort. Bot. Amst. 21 = Cupressus torulosa.
decora, Salisb. Prod. 398 = orientalis.
Devriesiana, Hort. Germ. ex Carr. Conif. ed. II. 141 = occidentalis.
dolabrata, *Linn. f. Suppl.* 420.—Japon.
Doniana, Hook. Lond. Journ. Bot. i. (1842) 571 (Libocedri sp.).—N. Zel.
Douglasii, Nutt. ex Parl. in DC. Prod. xvi. ii. 457 = gigantea.
dumosa, Gord. Pinet. Suppl. 102 = orientalis.
elegantissima, Hort. ex Gord. l. c. 17 = orientalis.
Ellwangeriana, Hort. ex Carr. in Rev. Hortic. (1869) 350 = occidentalis.

THUYA:—
ericoides, Hort. ex Carr. Conif. ed. II. 141 = orientalis?
ericoides, Hort. ex Gord. Pinet. Suppl. 91 = occidentalis.
excelsa, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 164 = Cupressus nootkatensis.
expansa, Laws. ex C. Koch, Dendrol. ii. ii. 182 = orientalis.
falcata, Hort. ex Carr. Conif. ed. II. 96 = orientalis.
filiformis, Lodd. ex Lindl. Bot. Reg. (1842) t. 20 = orientalis.
flagelliformis, Hort. ex Carr. Conif. ed. II. 101 = orientalis.
flagelliformis, Hort. ex Gord. Pinet. 325 = plicata.
Fortunei, Hort. ex Carr. Conif. ed. II. 94 = orientalis.
freneloides, Hort. l. c. ed. I. 93 = orientalis.
funiculata, Hort. ex Gord. Pinet. ed. II. 52 = orientalis.
gigantea, Gord. Pinet. 321 = Abies microphylla.
gigantea, *Nutt. in Journ. Acad. Philad.* vii. (1834) 52.—Am. bor. occ.
gigantea, Parl. in DC. Prod. xvi. ii. 457 = japonica.
glauca, Hort. ex Carr. Conif. ed. I. 95 = orientalis.
gracilifolia, Knight, ex Parl. in DC. Prod. xvi. ii. 459 = orientalis.
Hoveyi, Hort. ex Gord. Pinet. ed. II. 405 = occidentalis.
hybrida, Hort. ex Carr. Conif. ed. II. 102 = orientalis.
inaequalis, Desf. Tabl. ed. II. 274 = Callitris rhomboidea.
intermedia, Hort. ex Gord. Pinet. Suppl. 18 = orientalis.
japonica, Hort. ex Carr. Conif. ed. I. 94 = occidentalis.
japonica, Hort. ex Carr. l. c. ed. II. 141 = dolabrata.
japonica, *Maxim. in Bull. Acad. Pétersb.* x. (1866) 490.—Japon.
javanica, Burm. f. Fl. Ind. 202 (err. typ. 302) (Quid?).—Java.
lavandulaefolia, Poir. Encyc. Suppl. v. 303 = Taxodium heterophyllum.
lineata, Poir. l. c. = Taxodium heterophyllum.
Lobbiana, Hort. ex Gord. Pinet. 323 = gigantea.
Lobbii, Hort. ex Gord. Pinet. 323 = praec.
meldensis, Quétier, in Hérincq, Hortic. Franç. v. (1855) 231 = orientalis.
Menziesii, Dougl. ex Endl. Syn. Conif. 52 = gigantea.
minor, Hort. Angl. ex Carr. Conif. ed. II. 110 = occidentalis.
minor, Paul, ex Gord. Pinet. Suppl. 102 = orientalis.
monstrosa, Hort. ex Gord. l. c. 18 = orientalis.
nana, Hort. ex Carr. Conif. ed. I. 93; II. 110 = orientalis, occidentalis.
nepalensis, Lodd. ex G. Don, in Loud. Hort. Brit. 388; Hort. ex Carr. Conif. ed. I. 92 = orientalis.
obtusa, Mast. in Journ. Linn. Soc. xviii. (1881) 491 = Cupressus obtusa (cf. Addend.).
obtusa, Moench, Meth. 691 = occidentalis.
occidentalis, *Linn. Sp. Pl.* 1002.—Am. bor.; Sibir.
odorata, Marsh. Arbust. Am. 152 = occidentalis.
orientalis, *Linn. Sp. Pl.* 1002.—China; Japon.
pendula, Lamb. Pin. ed. 8vo. 130. t. 67. n. 64; Sieb. & Zucc. Fl. Jap. ii. 30 = orientalis.
pendulata, Hort. ex Carr. Conif. ed. I. 98 = praec.
pensilis, [Abel, in] Staunt. Embassy, 436 = Taxodium heterophyllum.
pisifera, Mast. in Journ. Linn. Soc. xviii. (1881) 489 = Cupressus pisifera (cf. Addend.).
plicata, *Donn, Hort. Cantabr.* ed. VI. 249.—Am. bor.
plicata, Lamb. Pin. ed. 8vo. 127. n. 61, partim = gigantea.
procera, Salisb. Prod. 398 = occidentalis.
pygmaea, Hort. ex Carr. Conif. ed. II. 96 = orientalis.
pyramidalis, Tenore, in Mem. Acc. Sc. Borb. Nap. iii. ii. (1832) 35. t. 2 = orientalis.
quadrangularis, *Vent. in Duham. Arb.* ed. Nov. iii. 16.—Madag.
semperaurescens, Hort. ex C. Koch, Dendrol. ii. ii. 183 = orientalis.
sibirica, Hort. ex Carr. Conif. ed. II. 106 = plicata.
sibirica, Hort. ex Gord. Pinet. Suppl. 104 = occidentalis.
sphaeroidalis, Rich. Monog. Conif. et Cyc. 45. t. 8. f. 2 = Cupressus thyoides (cf. Addend.).
sphaeroidea, Spreng. Syst. iii. 889 = praec.
Standishii, Carr. Conif. ed. II. 108 = japonica.

THUYA:—
stricta, Hort. ex Gord. Pinet. Suppl. 17 = orientalis.
tatarica, Gord. Pinet. Suppl. 104 = occidentalis.
tatarica, Lodd. ex G. Don, in Loud. Hort. Brit. 388, nomen = orientalis.
tetragona, Hook. Lond. Journ. Bot. iii. (1844) 148 = Libocedrus tetragona.
variegata, Hort. ex Carr. Conif. ed I. 94 = orientalis.
variegata. Marsh. Arbust. Am. 152 = occidentalis.
Verschaffelti, Hort. ex Parl. in DC. Prod. xvi. II. 462 = orientalis.
Vervaeneana, Van-Geert, ex Gord. Pinet. Suppl. 103 = occidentalis.
Wareana, Booth, Cat. (1838); ex Loud. Arb. Brit. iv. 2654 = plicata.
Wareana, Hort. ex Carr. Conif. ed. I. 104 = occidentalis.
Zuccarini, Hort. ex Carr. l. c. ed. II. 94 = orientalis.

THUYOPSIS, Parl. in DC. Prod. xvi. II. 460 (1868) = THUJOPSIS, Sieb. & Zucc. = **Thuya**, Linn. (Conif.).

THYANA, Ham. Prod. Pl. Ind. Occ. 36 (1825) = **Thouinia**, Poit. (Sapind.).

THYAREA, Benth. in Journ. Linn. Soc. xvii. (1878) 196 = **Thuaria**, Pers. (Gramin.).

THYELLA, Rafin. Pl. Tellur. iv. 84 (1836) = **Convolvulus**, Tourn.
obtusiloba, Rafin. l. c. = Ipomoea carnosa.
tamnifolia, Rafin. l. c. = Jacquemontia tamnifolia.

THYIA, Aschers. Fl. Brandenb. i. 886 (1864) = **Thuya**, Linn. (Conif.).

THYLACANTHA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 45 = **Angelonia**, Humb. & Bonpl. (Scrophul.).
campestris, Mart. l. c. = A. salicariaefolia.

THYLACANTHUS, Tul. in Arch. Mus. Par. iv. (1844) 175. *LEGUMINOSAE*, Benth. & Hook. f. i. 564.
corymbosus, Baill. Hist. Pl. ii. 96 = Dicymbe corymbosa.
ferrugineus, *Tul. in Arch. Mus. Par.* iv. (1844) 176.—Bras.

THYLACHIUM, Lour. Fl. Cochinch. 342 (1790) [Thilachium, em. ab DC. Prod. i. 254 (1824)]. *CAPPARIDEAE*, Benth. & Hook. f. i. 107.
africanum, *Lour. l. c.*—Afr. or.
angustifolium, *Boj. in Bouton, Rapp. Ann. Maur.* (1842) 49.—Madag.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 464.—Madag.
heterophyllum, *Juss. in Ann. Mus. Par.* xii. (1808) 71.—Madag.
laburnoides, *Baker, in Journ. Linn. Soc.* xxi. (1884) 319.—Madag.
laurifolium, *Baker, l. c.*—Madag.
lucidum, DC. Prod. i. 254 = Capparis lucida.
oppositiflorum, *DC. l. c.*—Ind. occ.
ovalifolium, Juss. in Ann. Mus. Par. xii. (1808) 71 = africanum.
ovatifolium, Steud. Nom. ed. II. ii. 683, sphalm. = praec.
panduraeforme, *Juss. in Ann. Mus. Par.* xii. (1808) 71.—Ins. Maurit.
querimbense, Klotzsch, in Peters, Reise Mossamb. Bot. 163 = africanum.
Sumangui, *Boj. Hort. Maurit.* 18; *et ex Bouton, in Rapp. Ann. Maur.* (1842) 48.—Ins. Mascar.
verrucosum, Klotzsch, in Peters, Reise Mossamb. Bot. 164 = africanum.

THYLACITIS, Renealm. ex Adans. Fam. ii. 504, 539 (1763) = **Gentiana**, Tourn.
angustifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 119 = G. acaulis.
Clusii, Fourr. l. c. = G. acaulis.
compressa, Rafin. Fl. Tellur. iii. 25 (Quid ?).—Hab. ?
grandiflora, Delarb. Fl. Auv. ed. II. 26 = G. acaulis.
leptina, Rafin. Fl. Tellur. iii. 25 = G. leptina.
maritima, Rafin. l. c. = Erythraea maritima ?

THYLACIUM, Spreng. Anleit. ii. II. 776 (1818) = **Thylachium**, Lour. (Cappar.).

THYLACOSPERMUM, Fenzl, in Endl. Gen. 967 (1840). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
BRYOMORPHA, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 172.
FLOURENSIA, Cambess. in Jacquem. Voy. Bot. t. 29 (1844).
PERIANDRA, Cambess. l. c. 27 (1844).
rupifragum, *Schrenk, Enum. Pl. Nov.* ii. 53.—Soongar.; Reg. Himal.

THYLACTITIS, Steud. Nom. ed. II. i. 671, in syn. (1840), sphalm. = THYLACITIS, Renealm. = **Gentiana**, Tourn.

THYLOCERAS, Steud. Nom. ed. II. ii. 721, in syn. (1841), sphalm. = **Styloceras**, A. Juss. (Euphorb.).

THYLOGLOSSA, Nees, in DC. Prod. xi. 389 (1847), sphalm. = TYLOGLOSSA, Hochst. = **Justicia**, Linn. (Acanth.).

THYMBRA, Linn. Sp. Pl. ed. I. 569 (1753). *LABIATAE*, Benth. & Hook. f. ii. 1190.
ambigua, Clarke, Trav. iv. 239 = spicata.
capitata, Griseb. Spicil. Fl. Rumel. ii. 127 = Thymus capitatus.
caroliniana, Walt. Fl. Carol. 162 = Macbridea pulchra.
cephalota, Desv. Opusc. 40 = Thymus Cephalotes.
ciliata, Desf. Fl. Atlant. ii. 10 = Thymus ciliatus.
ciliata, Tenore, Prod. Fl. Nap. p. lxix = Thymus Serpyllum.
hirsuta, Pers. Syn. ii. 114 = Satureia Thymbra.
hirsutissima, Vent. ex Pers. l. c. = praec.
spicata, *Linn. Sp. Pl.* 569.—Graec.; As. Min.
spicata, Pall. ex Bieb. Fl. Taur. Cauc. iv. 259 = Ziziphora taurica.
verticillata, Linn. Sp. Pl. 569 = spicata.
Zotas, Willd. ex Benth. in Linnaea, xi. (1837) 328 = Thymus ciliatus.

THYMELAEA, Tourn. ex Adans. Fam. ii. 285 (1763) = **Daphne**, Tourn. (Thymel.).

THYMELAEA, Tourn. ex Scop. Fl. Carn. ed. II. ii. 275 (1772), partim; Endl. Gen. Suppl. iv. II. 65 (1847). *THYMELAEACEAE* Benth. & Hook. f. iii. 191.
CHLAMYDANTHUS, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 355, 358.
GASTRILIA, Rafin. Fl. Tellur. iv. 105 (1836), partim.
LYGIA, Fasano, in Atti Acc. Nap. 1787 (1788) 245 (Ligia).
PAUSIA, Rafin. Fl. Tellur. iv. 105 (1836).
PIPTOCHLAMYS, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 356, 358.
alpina, All. Fl. Pedem. i. 132 = Daphne alpina.
argentea, Endl. Gen. Suppl. iv. II. 65 = T. Tartonraira.
arvensis, *Lam. Fl. Fr.* iii. 218.—Reg. Medit.; As. or. et bor.; Ind. bor. occ.
Aucheri, *Meissn. in DC. Prod.* xiv. 552.—As. Min.; Persia.
Bauhini, Endl. Gen. Suppl. iv. II. 66 = T. Sanamunda.
calycina, *Meissn. in DC. Prod.* xiv. 555.—Mont. Pyren.
candida, Scop. Fl. Carn. ed. II. i. 277 = Daphne alpina.
canescens, *Endl. Gen. Suppl.* iv. II. 65.—Marocc.; Hispan.
cilicica, *Meissn. in DC. Prod.* xiv. 552.—Cilicia.
Cneorum, Reichb. Ic. Fl. Germ. xi. t. 551 = T. Sanamunda.
Cneorum, Scop. Fl. Carn. ed. II. i. 276 = Daphne Cneorum.
coridifolia, *Endl. Gen. Suppl.* iv. II. 66.—Hispan.
dioica, *All. Auct. Fl. Pedem.* 9.—Europ.
elliptica, *Endl. Gen. Suppl.* iv. II. 66.—Hispan.
floribunda, Boiss. ex Casav. in Anal. Soc. Esp. Hist. Nat. ix. (1880) 301 = calycina.
Gnidium, All. Fl. Pedem. i. 133 = Daphne Gnidium.
Gussoni, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, iv. (1858) 121.—Sicil.
hirsuta, Endl. Gen. Suppl. iv. II. 65 = Passerina hirsuta.
Laureola, Scop. Fl. Carn. ed. II. i. 276 = Daphne Laureola.
Mezereum, Scop. l. c. = Daphne Mezereum.
microphylla, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 744; *Meissn. in DC. Prod.* xiv. 556.—Alger.

THYMELAEA:—
nitida, *Endl. Gen. Suppl.* iv. II. 65.—Reg. Mediterr.
nivalis, *Meissn. in DC. Prod.* xiv. 555.—Mont. Pyren.
orientalis, *Meissn. l. c.* 557.—'Oriens'; Hispan.
Passerina, Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 298 = arvensis.
precox, Gilib. Exercit. i. 7 = Daphne Mezereum.
pubescens, Meissn. in DC. Prod. xiv. 558 = virgata ?
Rodriguezii, Nym. Consp. 641 = velutina.
Ruizi, Losc. ex Casav. in Anal. Soc. Esp. Hist. Nat. ix. (1880) 301 = calycina.
Sanamunda, *All. Fl. Pedem.* i. 132.—Europ.
Tartonraira, *All. l. c.* 133.—Europ. austr.
thesoides, *Endl. Gen. Suppl.* iv. II. 66.—Hispan.
Thomasii, Endl. l. c. = T. Tartonraira.
tinctoria, *Endl. l. c.*—Europ. occ.
tomentosa, Endl. l. c. = T. Tartonraira.
velutina, *Meissn. in DC. Prod.* xiv. 555.—Ins. Balear.
villosa, *Endl. Gen. Suppl.* iv. II. 66.—Reg. Mediterr. occ.
virescens, *Meissn. in DC. Prod.* xiv. 554.—Alger.
virgata, *Endl. Gen. Suppl.* iv. II. 66.—Marocc.

THYMELINA, Hoffmgg. Verz. Pfl. 198 (1824) = **Gnidia**, Linn. (Thymel.).
sericea, Hoffmgg. l. c. 199, in obs. = G. sericea.
simplex, Hoffmgg. l. c. 198 = G. carinata.

THYMOPHYLLA, Lag. Gen. et Sp. Nov. 25 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 410.
LOWELLIA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 89.
Greggii, *A. Gray, l. c.* 92.—Mexic.
setifolia, *Lag. Gen. et Sp. Nov.* 25.—Mexic.

THYMOPHYLLUM, Benth. & Hook. f. Gen. ii. 410 (1873), sphalm. = praec.

THYMOPSIS, Benth. in Benth. & Hook. f. Gen. ii. 407 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.
Wrightii, *Benth. l. c.*—Cuba.

THYMOPSIS, Jaub. & Spach, Illustr. i. 72 (1843) = **Hypericum**, Tourn.
aspera, Jaub. & Spach, l. c. = H. Thymopsis.

THYMOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70, 136 = **Thymus**, Tourn. (Labiat.).
chamaedryus, St. Lag. l. c. 136 = Thymus Chamaedrys.
mastichinus, St. Lag. l. c. = Thymus Mastichina.
piperellus, St. Lag. l. c. = Thymus Piperella.
serpyllus, St. Lag. l. c. = Thymus Serpyllum.

THYMUS, Tourn. ex Linn. Syst. ed. I (1735). *LABIATAE*, Benth. & Hook. f. ii. 1186.
CORIDOTHYMUS, Reichb. f. in Oestr. Bot. Wochenbl. vii. (1857) 160.
MASTICHINA, Adans. Fam. ii. 194 (1763).
SERPYLLUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 234 (1745).
ZYGIS, Desv. in Ham. Prod. Pl. Ind. Occ. 46 (1825).
acicularis, Bess. Enum. Pl. Volh. 24 = T. Serpyllum.
acicularis, Guss. Fl. Sic. Prod. ii. 127 = striatus.
acicularis, *Waldst. & Kit. Pl. Rar. Hung.* ii. 157.—Europ.
acinoides, Schleich. ex Reichb. Fl. Germ. Excurs. 327 = Calamintha Acinos.
acinoides, Tenore, Prod. Fl. Nap. p. xxxv = Calamintha suaveolens.
Acinos, Asso, Syn. Stirp. Arag. 77 = Ziziphora acinoides.
Acinos, Brot. ex Steud. Nom. ed. II. ii. 683 = Calamintha patavina.
Acinos, Linn. Sp. Pl. 591 = Calamintha Acinos.
adscendens, Bernh. ex Link, Enum. Hort. Berol. ii. 115 = T. Serpyllum.
aegyptiacus, Bernh. ex Steud. Nom. ed. I. 835, nomen (Quid ?).—Aegypt.
aestivalis, *Kit. in Linnaea*, xxxii. (1863) 432.—Europ.
aestivus, *Reut. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 402.—Hispan.
affinis, Sieber, ex Benth. Lab. Gen. et Sp. 343 = T. Herba-barona.
affinis, Vis. in Parl. Giorn. Bot. Ital. ii. III. (1846) 4, in obs. = T. Serpyllum.
albicans, *Hoffmgg. & Link, Fl. Port.* i. 124. t. 11.—Lusitan.
albidus, Opiz, in Flora, vii. (1824) 526 = T. Serpyllum.
albinervis, Strobl, in Oestr. Bot. Zeitschr. xxxiii. (1883) 330 = T. Zygis.

THYMUS:—
albus, Link, Enum. Hort. Berol. ii. 116 = Micromeria rupestris.
algarbiensis, Lange, in Kjoeb. Vidensk. Meddel. (1863) 5 = albicans.
algeriensis, Boiss. & Reut. Pugill. Pl. Nov. (1852) 95 = ciliatus.
Allionii, Zumag. Fl. Pedem. i. 226 = Micromeria Piperella.
Almoradux, Duf. ex Boiss. Voy. Bot. Espagne, ii. 487 = tomentosus.
alpicolus, Schur, Enum. Pl. Transs. 525 = T. Serpyllum.
alpinus, Linn. Sp. Pl. 591 = Calamintha alpina.
alpinus, Pall. Tabl. Taur. 53 = Calamintha Acinos.
angulosus, Dulac, Fl. Hautes-Pyr. 401 = T. Serpyllum.
angustifolius, Pers. Syn. ii. 130 = T. Serpyllum.
angustifolius, Salisb. Prod. 86 = T. Zygis.
angustifolius, Webb, Iter Hisp. 21 = serpylloides.
angustus, Opiz, ex Déségl. in Bull. Soc. Sc. Angers, (1882) 190 = T. Serpyllum.
apricus, Opiz, Seznam, 97 = T. Serpyllum.
arenarius, Bernh. ex Reichb. Fl. Germ. Excurs. 312 = T. Serpyllum.
argaeus, Boiss. & Bal. Diagn. Ser. II. iv. 7 = T. Serpyllum.
arthooclades, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 36.—Persia.
arvensis, Schur, Enum. Pl. Transs. 972 (Index) = Calamintha Acinos.
athonicus, Bernh. ex Reichb. Fl. Germ. Excurs. 329 = Calamintha Nepeta.
attenuatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136, 70 = T. Herba-barona.
atticus, Čelak. in Flora, lxv. (1882) 564 = striatus.
aureus, Hort. ex Belg. Hortic. (1872) 356 = T. Serpyllum.
austriacus, Bernh. ex Reichb. Fl. Germ. Excurs. 312 = T. Serpyllum.
azoricus, Lodd. Bot. Cab. t. 1530 = T. Serpyllum.
azureus, Maund, ex Steud. Nom. ed. II. ii. 683, nomen (Quid ?).—Europ. austr.
Balansae, Boiss. & Kotschy, ex Boiss. Fl. Orient. iv. 556 = T. Serpyllum.
barbatus, Opiz, Boehm. Gew. (1823) 74 = T. Serpyllum.
Barrelieri, Spreng. Syst. ii. 698 = Micromeria marifolia.
Beneschianus, Opiz, Seznam, 96 = T. Serpyllum.
bidentatus, Stokes, Bot. Mat. Med. iii. 360 = Mentha Pulegium.
biflorus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 112 = Micromeria biflora.
Billardieri, *Boiss. Diagn.* Ser. II. iv. 8.—Ins. Cyprus.
biserratus, Blanco, Fl. Filip. ed. I. 478 = Hyptis brevipes.
Bleicherianus, *Pomel, Nouv. Mat. Fl. Atl.* 308.—Alger.
bonariensis, Tenore, Ind. Sem. Hort. Neap. (1839) 12. n. 14; et in Ann. Sc. Nat. Sér. II. xiii. (1840) 381 = Micromeria bonariensis.
Bovei, Benth. Lab. Gen. et Sp. 342 = T. Serpyllum.
Bowei, Steud. Nom. ed. II. ii. 683, sphalm. = praec.
brachyphyllus, Opiz, Seznam, 97 = T. Serpyllum.
bracteatus, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 8.—Hispan.
bracteatus, Opiz, Seznam, 97 = T. Serpyllum.
bracteosus, *Vis. ex Benth. Lab. Gen. et Sp.* 346.—Dalmat.
Brauneanus, Hoppe, ex Reichb. Fl. Germ. Excurs. 329 = Calamintha Nepeta.
brevicalyx, Strobl, in Oestr. Bot. Zeitschr. xxxiii. (1883) 330 = T. Serpyllum.
brevifolius, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 61; *et in Oken, Isis*, xxi. (1828) 1001.—Syria.
Broussonetii, *Boiss. Voy. Bot. Espagne*, ii. 492. t. 141.—Marocc.
Brownei, Sw. Prod. Veg. Ind. Occ. 89 = Micromeria Brownei.
caespititius, Brot. Fl. Lusit. i. 176 = T. Serpyllum.
caespitosus, Opiz, Seznam, 97 = T. Serpyllum.
Calamintha, Scop. Fl. Carn. ed. II. i. 425; Sm. Engl. Bot. t. 1676 = Calamintha officinalis.
calaminthoides, Reichb. Fl. Germ. Excurs. 329, in syn. = Calamintha officinalis.
calcicolus, Schur, Enum. Pl. Transs. 529 = T. Serpyllum.
campestris, Salisb. Prod. 85 = T. Serpyllum.
camphoratus, *Hoffmgg. & Link, Fl. Port.* i. 131.—Lusitan.
canescens, Dum. Fl. Belg. 48 = Calamintha Acinos.

THYMUS:—
canus, Stev. ex Bieb. Fl. Taur. Cauc. ii. 60 = Calamintha graveolens.
capitatus, *Hoffmgg. & Link, Fl. Port.* i. 123.—Reg. Mediterr.
capitellatus, *Hoffmgg. & Link, l. c.* 125. t. 12.—Lusitan.
cappadocicus, *Boiss. Diagn.* Ser. I. v. 17.—Cappadoc.
carniolicus, Borb. ex Déségl. in Bull. Soc. Sc. Angers, (1882) 191 = T. Serpyllum.
carnosus, *Boiss. Voy. Bot. Espagne*, ii. 490. t. 139.—Hispan.
carolinianus, Michx. Fl. Bor. Am. ii. 9 = Calamintha caroliniana.
carpathicus, Čelak. in Flora, lxv. (1882) 563 = T. Serpyllum.
caucasicus, Willd. ex Benth. in Linnaea, xi. (1837) 341 = T. Serpyllum.
Cephalotes, *Linn. Sp. Pl.* 592.—Hispan.; Lusit.
Chamaedrys, Fries, Nov. Fl. Suec. 197 = T. Serpyllum.
Chamissonis, Benth. in Linnaea, vi. (1831) 80 = Micromeria Douglasii.
Chaubardi, Boiss. & Heldr. ex Boiss. Diagn. Ser. II. iv. 6 = T. Serpyllum.
cherlerioides, Vis. in Mem. Ist. Venet. i. 1842 (1843) 44 = hirsutus.
chinensis, Hort. ex C. Koch, Dendrol. ii. 1. 314, 315 = vulgaris.
ciliatus, *Benth. Lab. Gen. et Sp.* 348.—Afr. bor.; Ins. Canar.
ciliatus, Lam. Fl. Fr. ii. 392 = T. Serpyllum.
ciliatus, Moench, Meth. Suppl. 138 = T. Mastichina.
cilicicus, *Boiss. & Bal. Diagn.* Ser. II. iv. 8.—Cilicia.
cimicinus, *Blum, ex Ledeb. Fl. Ross.* iii. 348.—Ross. austr.
citratus, Dum. Fl. Belg. 48 = T. Serpyllum.
citriodorus, Schreb. in Schweigg. & Koerte, Fl. Erlang. ii. 17 = T. Serpyllum.
clandestinus, Opiz, Boehm. Gew. (1823) 74 = aciculatis.
clandestinus, Salzm. ex Mutel, Fl. Fr. iii. 18 = Calamintha glandulosa.
clandestinus, Schur, Bot. Rundr. (1853) 63 = T. Serpyllum.
collinus, Bieb. Fl. Taur. Cauc. iii. 401 = T. Serpyllum.
collinus, Salisb. Prod. 86 = vulgaris.
coloratus, Boiss. & Reut. Pugill. Pl. Nov. (1852) 96 = ciliatus.
communis, Kitt. Taschenb. ed. II. 373 = T. Serpyllum.
comosus, Heuff. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. 1. (1852) 328 = T. Serpyllum.
comptus, *Frivald. in Flora*, xix. (1836) 489.—Rumelia.
comptus, Griseb. Spicil. Fl. Rumel. ii. 118 = odoratissimus.
concinnus, Salisb. Prod. 86 = Calamintha Acinos.
concolor, Opiz, Seznam, 97 = T. Serpyllum.
conspersus, Čelak. in Flora, lxv. (1882) 564 = striatus.
coriaceus, *Vog. in Linnaea*, xv. (1841) 74.—As. Min. ?
corsicus, Moris, Stirp. Sard. Elench. 37 = Mentha Requieni.
corsicus, Pers. Syn. ii. 131 = Calamintha corsica.
crenatus, Willd. ex Benth. in Linnaea, xi. (1837) 342 = Gardoquia affinis.
creticus, Brot. Fl. Lusit. i. 174 = capitatus.
creticus, DC. Fl. Fr. iii. 564 = Micromeria marifolia.
cristatus, Hampe, in Flora, xx. (1837) 233 = Micromeria cristata.
croaticus, Pers. Syn. ii. 130 = Micromeria croatica.
dalmaticus, Freyn, in Verh. Zool.-Bot. Ges. Wien, xxvii. (1878) 400 = T. Serpyllum.
debilis, Bunge, in Ledeb. Fl. Alt. ii. 391 = Calamintha debilis.
decumbens, Bernh. ex Reichb. Fl. Germ. Excurs. 313 = T. Serpyllum.
decussatus, *Benth. Lab. Gen. et Sp.* 482.—Arab.
deflexus, Benth. l. c. 344 = T. Serpyllum.
denticulatus, *Steud. Nom.* ed. II. ii. 684.—Cephalonia.
diffusus, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 41 = Calamintha Acinos.
diffusus, Boiss. Voy. Bot. Espagne, ii. 489 = hirtus.
diffusus, Hort. ex Benth. Lab. Gen. et Sp. 387 = Calamintha Nepeta.
diffusus, *Salzm. ex Benth. l. c.* 342.—Hispan.
Douglasii, Benth. in Linnaea, vi. (1831) 80 = Micromeria Douglasii.
duinensis, Moretti, ex Reichb. Fl. Germ. Excurs. 860 = Calamintha thymifolia.

THYMUS:—
effusus, Host, Fl. Austr. ii. 137 = T. Serpyllum.
Eisensteinianus, Opiz, ex Steud. Nom. ed. II. ii. 684 = T. Serpyllum.
elatus, Schrad. ex Reichb. Fl. Germ. Excurs. 313 = T. Serpyllum.
ellipticus, Opiz, Boehm. Gew. (1823) 74; Natural. n. 9 (1825) 103 = T. Serpyllum.
elongatus, Opiz, Seznam, 97 = T. Serpyllum.
elongatus, Schrad. ex Link, Enum. Hort. Berol. ii. 116 = tomentosus.
elwendicus, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 37.—Persia.
ericaefolius, Roth, Catalect. fasc. ii. 50 = Micromeria varia.
exiguus, Sibth. & Sm. Fl. Graec. Prod. i. 421; Fl. Graec. vi. 61. t. 575 = Calamintha graveolens.
exserens, Ehrh. ex Link, Enum. Hort. Berol. ii. 115 = T. Serpyllum.
fallax, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 33 = T. Serpyllum.
filiformis, Ait. Hort. Kew. ed. I. ii. 313 = Micromeria filiformis.
filiformis, Sieber, ex Benth. Lab. Gen. et Sp. 376 = Micromeria nervosa.
floribundus, Willd. ex Benth. in Linnaea, xi. (1837) 341 = Micromeria densiflora.
Fontanesii, Boiss. & Reut. Pugill. Pl. Nov. (1852) 95 = Monardi.
Froelichianus, Opiz, Nom. Bot. (1831) 80 = T. Serpyllum.
fruticosus, Link, Enum. Hort. Berol. ii. 116 = Micromeria marifolia.
fruticulosus, Bertol. Amoen. Ital. 101 = Micromeria approximata.
Funkii, *Coss. Notes Crit.* 125.—Hispan.
gibbosus, Stokes, Bot. Mat. Med. iii. 360 = Calamintha Acinos.
glaber, Mill. Gard. Dict. ed. VIII. n. 6 = T. Serpyllum.
glabratus, Hoffmgg. & Link, Fl. Port. i. 130. t. 15 = T. Serpyllum.
glabrescens, Benth. Lab. Gen. et Sp. 345 = T. Serpyllum.
glandulosus, Lag. ex Boiss. Voy. Bot. Espagne, ii. 487 = vulgaris.
glandulosus, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 386 = Calamintha glandulosa.
glaucus, Frivald. ex Boiss. Fl. Orient. iv. 558 = comptus.
gracilis, Willd. ex Benth. in Linnaea, xi. (1837) 341 = Micromeria xalapensis.
granatensis, *Boiss. Elench.* 74.—Hispan.
grandiflorus, Scop. Fl. Carn. ed. II. i. 424 = Calamintha grandiflora.
grandiflorus, Sims, Bot. Mag. t. 997 = Calamintha caroliniana.
gratissimus, Duf. ex Willk. & Lange, Prod. Fl. Hisp. ii. 405 = T. Serpyllum.
graveolens, Bieb. Fl. Taur. Cauc. ii. 60 = Calamintha graveolens.
graveolens, Sibth. & Sm. Fl. Graec. Prod. i. 421; Fl. Graec. vi. 61. t. 576 = teucrioides.
Guyonii, *Noé, in Bull. Soc. Bot. Fr.* ii. (1855) 580.—Alger.
Hackelianus, Opiz, Seznam, 97 = T. Serpyllum.
hayderensis, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 39.—Persia.
Herba-barona, *Loisel. Fl. Gall.* ed. I. 360. t. 9. f. 1.—Ins. Corsic.
heterophyllus, Poir. Encyc. vii. 648 = Calamintha Acinos.
heterotrichus, Griseb. Spicil. Fl. Rumel. ii. 116 = T. Serpyllum.
hirsutissimus, *Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 537.—Hungar.
hirsutissimus, Poir. Encyc. vii. 650 = Satureia Thymbra.
hirsutus, Auct. ex Benth. Lab. Gen. et Sp. 345 = T. Serpyllum.
hirsutus, *Bieb. Fl. Taur. Cauc.* ii. 59.—Graec.; As. Min.
hirtus, Link, in Buch, Beschr. Canar. Ins. 143, partim = Micromeria julianoides.
hirtus, Rafin. Caratteri, 80 = striatus.
hirtus, Sieber, ex Benth. Lab. Gen. et Sp. 386 = Micromeria marifolia.
hirtus, [Soland. in] Russ. Aleppo, ed. II. ii. 256 = Micromeria graeca.

THYMUS:—
hirtus, Viv. Fl. Lib. Spec. 30. t. 44 = Salvia aegyptiaca.
hirtus, *Willd. Enum. Hort. Berol.* 623.—Hispan.
hispanicus, Hort. ex Poir. Encyc. vii. 646 = Micromeria inodora.
Hornungianus, Opiz, Seznam, 97 = T. Serpyllum.
humifusus, Bernh. ex Link, Enum. Hort. Berol. ii. 115 = T. Serpyllum.
humifusus, Willd. ex Benth. in Linnaea, xi. (1837) 342 = Micromeria nubigena.
humillimus, *Čelak. in Flora*, lxvii. (1884) 534.—As. Min.
hyemalis, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 7.—Hispan.
ilerdensis, Gonzal. ex Costa, Supl. Fl. Catal. 63 = T. Zygis.
imbricatus, *Čelak. in Flora*, lxvii. (1884) 535.—As. Min.
imbricatus, Forsk. Fl. Aegypt. Arab. 108 = Micromeria Forskâhlei.
incanus, Sibth. & Sm. Fl. Graec. Prod. i. 421; Fl. Graec. vi. 62. t. 577 = Calamintha incana.
incanus, *Spreng. Nov. Prov.* 40.—Aegypt.
incanus, Willd. ex Benth. in Linnaea, xi. (1837) 342 = T. Serpyllum.
includens, Ehrh. ex Reichb. Fl. Germ. Excurs. 313 = T. Serpyllum.
indicus, Burm. f. Fl. Ind. 129 = Geniosporum prostratum.
inodorus, Benth. in Linnaea, vi. (1831) 77 = Micromeria varia.
inodorus, Desf. Fl. Atlant. ii. 30 = Micromeria inodora.
inodorus, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 122 = Micromeria cristata.
inodorus, Lej. Rev. Fl. Spa, 122 = T. Serpyllum.
integer, Griseb. Spicil. Fl. Rumel. ii. 116 = Billardieri.
integer, Sart. ex Nym. Consp. 594 = T. Serpyllum.
intermedius, *Schult. Obs. Bot.* 112.—Hab.?
interruptus, Opiz, Seznam, 97 = T. Serpyllum.
jalpanensis, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 38.—Persia.
Jankae, Čelak. in Flora, lxvi. (1883) 147 = T. Serpyllum.
Kollmunzerianus, Opiz, ex Benth. Lab. Gen. et Sp. 343 = T. Serpyllum.
Kosteleckianus, Opiz, Natural. n. ix. (1825) 104 = T. Serpyllum.
Kotschyanus, Boiss. & Hohen. Diagn. Ser. I. v. 16 = T. Serpyllum.
Kratzmannianus, Opiz, Seznam, 97 = T. Serpyllum.
laevigatus, Vahl, Symb. Bot. ii. 65 = T. Serpyllum.
lanceolatus, *Desf. Fl. Atlant.* ii. 30. t. 128.—Graecia; Afr. bor.
lanceolatus, Poir. Encyc. Suppl. v. 305 = Pycnanthemum lanceolatum.
lanceolatus, Sibth. & Sm. Fl. Graec. Prod. i. 419 = T. Serpyllum.
lanuginosus, Mill. Gard. Dict. ed. VIII. n. 8 = T. Serpyllum.
latifolius, Andrz. ex Trautv. in Act. Hort. Petrop. ix. (1884) 90 = T. Serpyllum.
linearis, Benth. in Wall. Pl. As. Rar. i. 31 = T. Serpyllum.
lobatus, Vog. in Linnaea, xv. (1841) 75 = villosus.
Loevyanus, Opiz, Natural. n. ix. (1825) 105 = T. Serpyllum.
longicaulis, Presl, Fl. Sic. p. xxxvii = striatus.
longiflorus, *Boiss. Elench.* 75.—Hispan.
longistylus, Opiz, Seznam, 96 = T. Serpyllum.
Loscosii, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 401.—Hispan.
lucidus, Ehrh. Beitr. vii. 150 = Ziziphora clinopodioides.
lucidus, Guss. Suppl. Fl. Sic. Prod. 198 = nitidus.
lucidus, Willd. Enum. Hort. Berol. 624 = T. Serpyllum.
lusitanicus, Boiss. Voy. Bot. Espagne, ii. 489 = villosus.
Majorana, Kuntze, Taschen-Fl. Leipz. 106 = Origanum Majorana.
majoranaefolius, Desf. Tabl. ed. II. 74 = T. Serpyllum.
Mannianus, Opiz, Seznam, 97 = T. Serpyllum.
marginatus, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 184 = T. Chamaedrys.
marginatus, Sm. in Rees, Cyclop. xxxv. n. 17 = Micromeria Piperella.

THYMUS:—
marifolius, Pourr. ex Willd. Enum. Hort. Berol. 626 = tomentosus.
marifolius, Willd. l. c. 624 = Micromeria marifolia.
Marinosci, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 684 = denticulatus.
Marinosci, Tenore, Prod. Fl. Nap. p. xxxv = striatus.
maroccanus, *Ball, in Journ. Bot.* xiii. (1875) 174.—Marocc.
Marschallianus, Viv. Fl. Cors. Diagn. 9 = T. Herba-barona.
Marschallianus, Willd. Sp. Pl. iii. 141 = T. Serpyllum.
Mastichina, *Linn. Sp. Pl.* ed. II. 827.—Hispan.; Afr. bor.
Mastichina, Sibth. & Sm. Fl. Graec. Prod. i. 422 = Micromeria nervosa.
megapotamicus, *Spreng. Syst.* iv. *Cur. Post.* 224.—Bras.
melissoides, Bernh. ex Reichb. Fl. Germ. Excurs. 327 = Calamintha patavina.
melissoides, Fisch. ex Benth. Lab. Gen. et Sp. 344 = T. Serpyllum.
melissoides, Schweigg. Enum. Pl. Hort. Regiom. 72 = Calamintha patavina?
membranaceus, *Boiss. Elench.* 75.—Hispan.
micans, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 19 = T. Serpyllum.
micranthus, Brot. Fl. Lusit. i. 176 = Micromeria graeca.
micranthus, Wierzb. ex Opiz, Nom. (1831) 77 = T. Serpyllum.
micromerioides, *Rouy, in Bull. Soc. Bot. Fr.* xxxi. (1884) 275.—Hispan.
microphyllus, Urv. in Mém. Soc. Linn. Par. i. (1822) 327 = Micromeria microphylla.
minor, Trev. Ind. Sem. Hort. Vratisl. (1819) App. i. 4 = Calamintha Nepeta.
minutus, Opiz, Seznam, 97 = T. Serpyllum.
Monardi, *Noé, in Expl. Scientif. Algér.* t. 66.—Alger.
montanus, Crantz, Stirp. Austr. ed. II. fasc. iv. 278 = Calamintha alpina.
montanus, Dum.-Cours. Bot. Cult. ed. II. iii. 32 = Satureia montana.
montanus, Waldst & Kit. Pl. Rar. Hung. i. 72. t. 71 = T. Serpyllum.
monticola, Opiz, in Lotos, iv. (1854) 103, nomen (Quid?).
Moschatella, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 33 = Calamintha officinalis.
multiflorus, *Pers. Syn.* ii. 131.—Hab.?
Munbyanus, Boiss. & Reut. Pugill. Pl. Nov. (1852) 96 = ciliatus.
neapolitanus, Herb. Kern. ex Strobl, in Oestr. Bot. Zeitschr. xxxiii. (1883) 329 = striatus.
Nepeta, Sm. Fl. Brit. ii. 642 = Calamintha Nepeta.
nepetoides, D. Don, Prod. Fl. Nep. 113 = Nepeta ruderalis.
nervosus, J. Gay, ex Benth. Lab. Gen. et Sp. 345 = T. Serpyllum.
nitens, Lamotte, Prod. Fl. Centr. Fr. ii. 595 = T. Herba-barona.
nitidus, *Guss. Fl. Sic. Syn.* ii. 97.—Sicil.
Notarisii, Zumag. Fl. Pedem. i. 226 = Micromeria thymoides.
nubigenus, H. B. & K. Nov. Gen. et Sp. ii. 316 = Micromeria nubigena.
numidicus, Benth. Lab. Gen. et Sp. 347 = T. Zygis.
numidicus, Poir. Voy. Barb. ii. 189 = lanceolatus.
nummularifolius, Loisel. Fl. Gall. ed. I. t. 9. f. 2; et ex Benth. Lab. Gen. et Sp. 732 = Calamintha alpina.
nummularius, Bieb. Fl. Taur.-Cauc. ii. 58 = T. Serpyllum.
oblongifolius, Opiz, Natural. n. x. (1825) 241 = T. Serpyllum.
obtusatus, Opiz, in Lotos, iv. (1854) 103 (Quid?).—Europ.
ocheus, Heldr. & Sart. in Boiss. Diagn. Ser. II. iv. 6 = T. Serpyllum.
odoratissimus, *Bieb. Fl. Taur.-Cauc.* iii. 405.—Ross. austr.; As. Min.
odoratissimus, Mill. Gard. Dict. ed. VIII. n. 9 = T. Serpyllum.
origanifolius, D. Don, Prod. Fl. Nep. 112 = Calamintha longicaulis.
origanifolius, Vis. in Flora, xiii. (1830) 51 = Micromeria dalmatica.

THYMUS:—
origanoides, *Webb & Berth. Phyt. Canar.* iii. 69. t. 152.—Ins. Canar.
Origanum, Kuntze, Taschen-Fl. Leipz. 106 = Origanum vulgare.
Ortmannianus, Opiz, Seznam, 97 = T. Serpyllum.
ovatus, Mill. Gard. Dict. ed. VIII. n. 7 = T. Serpyllum.
pallens, Opiz, ex Déségl. in Bull. Soc. Sc. Angers, (1882) 186 = T. Serpyllum.
pallescens, *Noé, in Expl. Scientif. Algér.* t. 66.—Alger.
pallidus, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 65, nomen = lanceolatus.
pannonicus, All. Fl. Pedem. i. 20 = T. Serpyllum.
paronychioides, *Čelak. in Flora*, lxv. (1882) 564.—Sicil.
parviflorus, Opiz, Seznam, 96 = T. Serpyllum.
parviflorus, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 386 = Mentha Requieni.
parvifolius, C. Koch, in Linnaea, xxi. (1848) 666 = T. Serpyllum.
patavinus, Jacq. Obs. Bot. iv. 7 = Calamintha patavina.
patavinus, Pall. ex Bieb. Fl. Taur.-Cauc. ii. 60 = Calamintha graveolens.
pectinatus, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 32 = odoratissimus.
Petteri, Presl, Bot. Bemerk. 100 = acicularis.
piligerus, Opiz, ex Reichb. Fl. Germ. Excurs. 312 = T. Serpyllum.
pilosus, Opiz, Boehm. Gew. (1823) 74; Natural. n. vi. (1824) 40 = T. Serpyllum.
Piperella, All. Fl. Pedem. i. 21. t. 37. f. 3 = Micromeria Piperella.
Piperella, *Linn. Syst.* ed. XII. 400.—Reg. Mediter.
Piperella, Waldst. & Kit. Pl. Rar. Hung. ii. 169 = Micromeria croatica.
piperitus, D. Don, Prod. Fl. Nep. 112 = Calamintha longicaulis.
praecox, Opiz, Natural. n. vi. (1824) 40; et in Flora, vii. (1824) Beil. 84 = T. Serpyllum.
procerus, Opiz, ex Benth. Lab. Gen. et Sp. 343 = T. Serpyllum.
procumbens, Bernh. ex Opiz, Boehm. Gew. (1823) 74 = T. Serpyllum?
pruinosus, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 253 = cappadocicus.
Pseudo-serpyllum, Reichb. ex Benth. Lab. Gen. et Sp. 344 = T. Serpyllum.
pulchellus, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 40; C. A. Mey. Verz. Pfl. Cauc. 87 = T. Serpyllum.
pulcherrimus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 140 = T. Serpyllum.
pulegioides, Linn. Sp. Pl. 592 = T. Serpyllum.
pulegioides, Reichb. Fl. Germ. Excurs. 313 = T. Serpyllum.
pulvinatus, *Čelak. in Flora*, lxvii. (1884) 533.—As. Min.
pumilus, Gueldenst. ex Ledeb. Fl. Ross. iii. 346 = T. Serpyllum.
punctatus, Tineo, ex Benth. in DC. Prod. xii. 217 = Micromeria approximata.
punctatus, Vis. in Mem. Ist. Venet. i. 1842 (1843) 43. t. 2. f. 1 = T. Serpyllum.
punctatus, Willd. Phyt. 8 = Ziziphora clinopodioides.
purpurascens, Poir. Encyc. vii. 654 = Calamintha purpurascens.
pusillus, Gueldenst. ex Ledeb. Fl. Ross. iii. 346 = T. Serpyllum.
Pusio, Dichtl, in Deutsche Bot. Monatsschr. ii. (1884) 134 = T. Serpyllum.
quadrangulus, Ehrh. ex Boiss. Fl. Orient. iv. 559 = decussatus.
rariflorus, C. Koch, in Linnaea, xxi. (1848) 666 = T. Serpyllum.
raripilus, Dichtl, in Deutsche Bot. Monatsschr. ii. (1884) 134 = T. Serpyllum.
recurvatus, *Kit. in Linnaea*, xxxii. (1863) 431.—Hungar.
reflexus, Lej. Rev. Fl. Spa, 121 = T. Serpyllum.
Reichelianus, Opiz, Seznam, 96 = T. Serpyllum.
Reinegeri, Opiz, l. c. 97 = T. Serpyllum.
repens, D. Don, Prod. Fl. Nep. 113 = Calamintha umbrosa.
repens, Gilib. Fl. Lituan. i. 74 = T. Serpyllum.
Richardi, Pers. Syn. ii. 130 = Origanum majoricum.

THYMUS:—
rigidus, Reichb. ex Bess. in Flora, xv. (1832) II. Beibl. 18 = T. Serpyllum.
robustus, Opiz, Seznam, 97 = T. Serpyllum.
Rochelianus, Čelak. in Flora, lxv. (1882) 563 = T. Chamaedrys.
Roegneri, C. Koch, in Linnaea, xxi. (1848) 666 = T. Serpyllum.
rotundifolius, Pers. Syn. ii. 131 = Zizyphora acinoides.
rotundifolius, Poir. Encyc. vii. 654 = T. Piperella.
rotundifolius, Rochel, ex Nym. Consp. 588 = Calamintha patavina.
rotundifolius, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 108 = T. Serpyllum.
sabulicola, *Coss. Notes Crit.* 125.—Hispan.
sabulicus, Walp. Ann. iii. 924, sphalm. = praec.
satureioides, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 253.—Marocc.
scaber, Presl, Fl. Sic. p. xxxvi = Satureia scabra.
scaber, Willd. ex Benth. in Linnaea, xi. (1837) 342 = Gardoquia affinis.
scharkensis, Opiz, in Lotos, iv. (1854) 103, nomen (Quid ?).—Europ.
Selloi, Spreng. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 168 = Cunila fasciculata.
senilis, Dichtl, in Deutsche Bot. Monatsschr. ii. (1884) 133 = T. Serpyllum.
serpens, Opiz, Seznam, 97 = T. Serpyllum.
serpylloides, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) 16.—Hispan.
Serpyllum, Coss. ex Willk. & Lange, Prod. Fl. Hisp. ii. 405 = bracteatus.
Serpyllum, *Linn. Sp. Pl.* 590.—Europ.; As. et Afr. bor.
serratus, Opiz, in Flora, v. (1822) 267 = T. Serpyllum.
serratus, Poir. Encyc. vii. 653 = Pycnanthemum muticum.
serrulatus, *Hochst. ex Benth. in DC. Prod.* xii. 203.—Abyss.
Sibthorpii, Benth. Lab. Gen. et Sp. 345 = lanceolatus.
silvestris, Nym. Consp. 593 = sylvestris.
simplex, Kitt. Taschenb. ed. II. 373 = T. Serpyllum.
Sintenisii, *Čelak. in Flora*, lxvii. (1884) 537.—As. Min.
sipyleus, Boiss. Diagn. Ser. I. v. 16 = T. Serpyllum.
sparsiflorus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 402 = T. Zygis.
spathulatus, Opiz, Natural. n. ix. (1825) 105; Fisch. ex Ledeb. Fl. Ross. iii. 346 = T. Serpyllum.
spicatus, W. Baxt. in Loud. Hort. Brit. Suppl. ii. 677, nomen (Quid ?).—Europ.
spinulosus, Tenore, Prod. Fl. Nap. p. xxxv = striatus.
squarrosus, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 32 = T. Serpyllum.
stenophyllus, Opiz, Seznam, 97 = T. Serpyllum.
striatus, Munby, Fl. Algér. 62 = hirtus.
striatus, Reichb. Ic. Fl. Germ. xviii. 36 = T. Serpyllum.
striatus, *Vahl, Symb. Bot.* iii. 78.—Europ. austr.; As. Min.
suaveolens, Sibth. & Sm. Fl. Graec. Prod. i. 420 = Calamintha suaveolens.
suavis, Salisb. Prod. 86 = T. Mastichina.
subcitratus, Schreb. in Schweigg. & Koerte, Fl. Erlang. ii. 18 = T. Serpyllum.
subcordatus, Vis. Fl. Dalmat. i. t. 19 = Micromeria croatica.
sublaxus, Rouy, in La Natural. ii. (1882) (Matér. Rev. Fl. Portug. 46) = vulgaris.
sublobatus, Pomel, Nouv. Mat. Fl. Atl. 308 = ciliatus.
subnudus, Spreng. Syst. ii. 698 = Micromeria Pulegium.
sudeticus, Opiz, ex Reichb. Fl. Germ. Excurs. 312 = T. Serpyllum.
sylvaticus, Bernh. Syst. Verz. Erf. 77 = Calamintha Clinopodium.
sylvestris, *Hoffmgg. & Link, Fl. Port.* i. 132. t. 16.—Hispan.
sylvestris, Schreb. in Schweigg. & Koerte, Fl. Erlang. ii. 17 = T. Serpyllum.
syriacus, *Boiss. Diagn.* Ser. I. xii. 47.—Syria.
syrticus, Spreng. Syst. ii. 697 = Salvia aegyptiaca.

THYMUS:—
taxifolius, Willd. ex Benth. in Linnaea, xi. (1837) 342 = Gardoquia taxifolia.
Teneriffae, Hort. ex Benth. Lab. Gen. et Sp. 377 = Micromeria microphylla.
Teneriffae, Poir. Encyc. vii. 650 = Micromeria Teneriffae.
tenuifolius, Mill. Gard. Dict. ed. VIII. n. 2 = T. Zygis.
teucrioides, *Boiss. & Sprun. Diagn.* Ser. I. v. 15.—Graec.
therebinthinaceus, Brouss. ex Willd. Enum. Hort. Berol. 624 = Micromeria Teneriffae.
thymbroides, Pomel, Nouv. Mat. Fl. Atl. 124 = ciliatus.
tomentosus, Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp. ii. 400 = T. Mastichina.
tomentosus, *Willd. Enum. Hort. Berol.* ii. 626.—Hispan.
Trachselianus, Opiz, Natural. n. xi. (1826) 461 = T. Serpyllum ?
Tragoriganum, Linn. Mant. i. 84 = Satureia Thymbra.
transsilvanicus, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 197 = T. Serpyllum.
tunetanus, *Pomel, Nouv. Mat. Fl. Atl.* 310.—Afr. bor.
umbrosus, Spreng. Nov. Prov. 41 = Calamintha officinalis.
variabilis, Hoffmgg. & Link, Fl. Port. i. 134. t. 15 = T. Serpyllum.
verecundus, Salisb. Prod. 86 = Calamintha patavina.
veronicaefolius, Spreng. Syst. ii. 698 = Melissa veronicaefolia.
verticillatus, Poir. Encyc. vii. 653 = Pycnanthemum lanceolatum.
villosissimus, Tausch, in Syll. Ratisb. ii. (1828) 248 = Calamintha alpina.
villosus, *Linn. Sp. Pl.* 592.—Europ. austr.
villosus, Pall. ex Bieb. Fl. Taur.-Cauc. ii. 59 = T. Serpyllum.
villosus, Sibth. & Sm. Fl. Graec. vi. 62. t. 578 = Billardieri.
villosus, Willd. ex Benth. in Linnaea, xi. (1837) 342 = T. Cephalotes.
virgatus, Tenore, Prod. Fl. Nap. p. xxxv = Micromeria graeca.
virginicus, Blanco, Fl. Filip. ed. I. 478 = Hyptis capitata.
virginicus, Linn. Mant. ii. 409 = Pycnanthemum lanceolatum.
vulgaris, Bieb. Beschr. Casp. 209 = T. Serpyllum.
vulgaris, *Linn. Sp. Pl.* 591.—Europ. austr.; As. bor.
vulgaris, Sibth. & Sm. Fl. Graec. Prod. i. 419 = hirsutus.
vulgaris, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 402 = aestivus.
Webbianus, *Rouy, in Bull. Soc. Bot. Fr.* xxxi. (1884) 275.—Hispan.
Welwitschii, *Boiss. Diagn.* Ser. II. iv. 9.—Lusitan.
Wierzbickianus, Opiz, Seznam, 97 = T. Serpyllum.
Willdenowii, Boiss. in Kelaart, Fl. Calp. 218 = hirtus.
Wondračekianus, Opiz, Seznam, 97 = T. Serpyllum.
xalapensis, H. B. & K. Nov. Gen. et Sp. ii. 316 = Micromeria Brownei.
xylorrhizus, Boiss. & Kotschy, ex Boiss. Fl. Orient. iv. 556 = T. Serpyllum.
Zattar, *Munby, Cat. Pl. Alger.* 23, nomen.—Alger.
zattarellus, Pomel, Nouv. Mat. Fl. Atl. 309 = ciliatus.
zygioides, Griseb. Spicil. Fl. Rumel. ii. 118 = striatus.
Zygis, Bieb. Beschr. Casp. 210 = odoratissimus.
Zygis, Brot. Fl. Lusit. i. 176 = sylvestris.
Zygis, *Linn. Sp. Pl.* 591.—Hispan.; Lusit.
Zygis, Pall. in Nov. Act. Acad. Petrop. x. (1797) 313 = T. Serpyllum.
Zygis, Sibth. & Sm. Fl. Graec. vi. 60. t. 574 = striatus.

THYOPSIS, Wedd. Chlor. And. i. 165, in nota, t. 27*, A B C (1856) = **Tafalla**, D. Don (Compos.).
ferruginea, Wedd. l. c. = T. ferruginea.
graveolens, Wedd. l. c. (= *Loricaria graveolens*).—N. Granat.
stenophylla, Wedd. l. c. (= *Loricaria stenophylla*).—N. Granat.

THYPHA, Costa, Introd. Fl. Catal. p. lxv. 292 (Index) (1864), sphalm. = **Typha**, Linn.

THYRIDOSTACHYUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 379 (1836) = **Ophiurus**, Gaertn. f. (Gramin.).
laeve, Nees, ex Steud. Nom. ed. II. ii. 615 = O. perforatus.

THYROCARPUS, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 225. *BORAGINEAE*, Benth. & Hook. f. ii. 847.
fulvescens, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 499.—China.
glochidiatus, *Maxim. l. c.*—China.
Sampsoni, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 225.—China.

THYROMA, Miers, Apocyn. S. Am. 22 (1878) = **Aspidosperma**, Mart. & Zucc. (Apocyn.).
bicolor, Miers, l. c. 25 = A. bicolor.
decipiens, Miers, l. c. 24 = A. decipiens.
Lhotzkiana, Miers, l. c. 25 = A. Lhotzkianum.
nitida, Miers, l. c. 24 = A. nitida.
parvifolia, Miers, l. c. 25 = A. parvifolium.
polyneura, Miers, l. c. 24 = A. polyneuron.
Riedelii, Miers, l. c. 26 = A. Riedelii.
Sellowii, Miers, l. c. 24 = A. Sellowii.
sessiliflora, Miers, l. c. 23 = A. sessiliflorum.

THYROPHORA, Neck. Elem. ii. 13 (1790) = **Gentiana**, Tourn.

THYRSACANTHUS, Nees, in Mart. Fl. Bras. ix. 97. t. 13 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1119.
ODONTONEMA, Nees, in Linnaea, xvi. (1842) 300.
amplexicaulis, *Nees, in Mart. Fl. Bras.* ix. 98.—Bras.
barlerioides, *Nees, l. c.* 97.—Bras.
bracteolatus, *Nees, in DC. Prod.* xi. 325.—N. Granat.
callistachyus, *Nees, l. c.* 326.—Mexic.
calycotrichus, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 284 = Schaueria calycotricha.
coccineus, T. Anders. l. c. = Jacobinia coccinea.
cuspidatus, *Nees, in DC. Prod.* xi. 323.—Mexic.
dissitiflorus, *Nees, in Mart. Fl. Bras.* ix. 98.—Bras.
Earlii, F. Muell. Fragm. vi. 87 = Graptophyllum Earlii.
embelioides, Pritz. Ic. Ind. i. 1103, sphalm. = *Thyrsanthus embelioides*.
Flagellum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 146.—Costa Rica.
foliaceo-bracteatus, *Oerst. l. c.*—Costa Rica.
fuchsioides, *Nees, in DC. Prod.* xi. 325.—N. Granat.
Hookerianus, *Nees, l. c.* 324.—Ecuador.
indicus, Nees, l. c. 325 = Eranthemum indicum.
interruptus, *Nees, l. c.* 326.—N. Granat.
Lemairianus, *Nees, l. c.* 729.—Hab. ?
lilacinus, Lindl. in Journ. Hort. Soc. vi. (1851) 159 = callistachys.
Lindenianus, Nees, in DC. Prod. xi. 326 = Eranthemum cuspidatum.
longifolius, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 145.—Nicaragua.
longistamineus, *Nees, in DC. Prod.* xi. 326 (an Jacobiniae sp. ?).—Peruv.
nitidus, *Nees, l. c.* 327.—Ind. occ.
pantasmensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 144.—Nicaragua.
ramosissimus, Moric. Pl. Nouv. Am. 165. t. 96; Nees, in Mart. Fl. Bras. ix. 98 = Jacobinia ramosissima (in Addend.).
ruber, *Nees, in DC. Prod.* xi. 326.—Venezuela.
rutilans, *Planch. in Fl. des Serres*, vii. (1852) 241. t. 732.—N. Granat.
Schomburgkianus, *Nees, in Hook. Lond. Journ. Bot.* iv. (1845) 636.—Guian.
sessilis, *Nees, in DC. Prod.* xi. 324.—N. Granat.
squarrosus, T. Anders. in Journ. Agri-Hort. Soc. Ind. N. S. i. (1868) 284 = Dianthera nodosa ?
strictus, *Nees, in DC. Prod.* xi. 324.—Honduras.
tubaeformis, *Nees, l. c.*—Guatemala.
variegatus, *Nees, l. c.* 325.—Guian.

THYSAMUS, Reichb. Consp. 197 (1828), sphalm. = **Thysanus**, Lour. (Inc. Sed.).

THYRSANTHEMA, Neck. Elem. i. 6 (1790) = **Chaptalia**, Vent. (Compos.).

THYRSANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 245 = **Forsteronia**, G. F. W. Mey. (Apocyn.).
Acouci, Miers, Apocyn. S. Am. 98 = F. Acouci.
adenobasis, Miers, l. c. 96 = F. adenobasis.
affinis, Miers, l. c. 101 = F. affinis.
Aubletianus, Miers, l. c. 98 = Thenardia umbellata.
Benthamianus, Miers, l. c. 95 = F. Benthamiana.
bracteatus, Miers, l. c. 102 = F. bracteata.
brasiliensis, Griseb. Fl. Brit. W. Ind. 412, in obs. = F. brasiliensis.
corylifolius, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 519 = F. corylifolia.
corymbiferus, Miers, Apocyn. S. Am. 98 = F. Schomburgkii.
corymbosus, Miers, l. c. 97 = Parsonsia corymbosa.
crebriflorus, Miers, l. c. 105.—Bras.
difformis, Miers, l. c. 99 = Trachelospermum difforme.
diospyrifolius, Miers, l. c. 96 = F. diospyrifolia.
embelioides, A. DC. in DC. Prod. viii. 387 = F. embelioides.
fasciculatus, Miers, Apocyn. S. Am. 100 = Tabernaemontana fasciculata.
Gardneri, A. DC. in DC. Prod. viii. 387 = F. Gardneri.
glabrescens, Miers, Apocyn. S. Am. 102 = F. glabrescens.
gracilis, Benth. in Hook. Journ. Bot. iii. (1841) 246.—Guian.
guyanensis, Miers, Apocyn. S. Am. 97 = F. guyanensis.
laurifolius, Miers, l. c. 94 = F. laurifolia.
leptocarpus, Griseb. in Goett. Abh. xxiv. (1879) 224 = Parsonsia leptocarpa.
Luschnatii, Miers, Apocyn. S. Am. 106 = F. Luschnathii.
macrophyllus, Miers, l. c. 96 = F. macrophylla.
meridionalis, Miers, l. c. 106 = F. meridionalis.
multinervius, Miers, l. c. 103 = F multinervia.
myrianthus, Miers, l. c. 105 = F. floribunda, *Muell. Arg.*
parviflorus, Miers, l. c. 100 = Tabernaemontana populifolia.
pilosus, A. DC. in DC. Prod. viii. 387 = F. pilosa.
placidus, Miers, Apocyn. S. Am. 101.—Bras.
populifolius, Miers, l. c. 99 = Tabernaemontana populifolia.
pubescens, Miers, l. c. 101 = F. pubescens.
pyriformis, Miers, l. c. 100.—Ind. occ.
rufus, Miers, l. c. 104 = F. rufa.
Schomburgkii, Benth. in Hook. Journ. Bot. iii. (1841) 246.—Guian.
sessilis, Miers, Apocyn. S. Am. 104 = Malouetia sessilis.
spicatus, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 520, in obs. = F. spicata.
velutinus, Miers, Apocyn. S. Am. 86 = F. multinervia.
Wilsonii, Griseb. Fl. Brit. W. Ind. 412.—Jamaic.

THYRSANTHUS, Ell. in Journ. Acad. Philad. i. (1818) 371 = **Wistaria**, Nutt. (Legumin.).
floridana, Croom, in Am. Journ. Sc. Ser. I. xxv. (1834) 75 = W. speciosa.
frutescens, Ell. in Journ. Acad. Philad. i. (1818) 371 = W. speciosa.

THYRSANTHUS, Schrank, in Denkschr. Akad. Muench. 1813 (1814) 75 = **Lysimachia**, Tourn. (Primul.).
palustris, Schrank, l. c. = L. thyrsiflora.

THYRSINE, Gled. Syst. 286 (1764) = **Cytinus**, Linn. (Balanoph.).
Hypocistis, Walp. Ann. iii. 514 = C. Hypocistis.

THYRSODIUM, Salzm. ex Benth. in Hook. Kew Journ. iv. (1852) 17. *ANACARDIACEAE*, [Benth. & Hook. f. i. 323].
giganteum, *Engl. in DC. Monog. Phan.* iv. 297.—Bras.
guianense, *Sagot, ex March. Rév. Anacard.* 160.—Guian.
Salzmannianum, Benth. in Hook. Kew Journ. iv. (1852) 18 = Schomburgkiana.
Schomburgkianum, *Benth. l. c.*—Am. trop.
Spruceanum, *Benth. l. c.* 17.—Bras.

THYRSOSMA, Rafin. Sylva Tellur. 130 (1838) = **Viburnum**, Linn. (Caprif.).
chinensis, Rafin. l. c. = V. odoratissimum.

THYSANACHNE, Presl, Thysan. Nov. Pl. Gen. (1829); et J. Presl, in Rel. Haenk. i. 252 (1830) = **Arundinella**, Raddi (Gramin.).
peruviana, J. Presl, in Rel. Haenk. i. 253 (= *Ischaemum peruvianum*).—Peruv.
scoparia, Presl, Thysan. Nov. Pl. Gen. (1829); et J. Presl, in Rel. Haenk. i. 253 = A. Deppeana.

THYSANELLA, A. Gray, ex Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 232 = **Polygonum**, Tourn.
fimbriata, A. Gray, l. c. = P. fimbriatum.

THYSANELLA, Salisb. Gen. Pl. Fragm. 67 (1866) = **Thysanotus**, R. Br. (Liliac.).

THYSANOCARPUS, Hook. Fl. Bor. Am. i. 69. t. 18. f. A (1829). *CRUCIFERAE*, Benth. & Hook. f. i. 94.
crenatus, Nutt. in Torr. & Gray, Fl. N. Am. i. 118 = laciniatus.
curvipes, *Hook. Fl. Bor. Am.* i. 69. t. 18. f. A.—Am. bor. occ.
Deppii, Nutt. ex Torr. & Gray, Fl. N. Am. i. 118 = elegans.
elegans, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 51.—Calif.
erectus, *S. Wats. in Proc. Am. Acad.* xi. (1876) 124.—Ind. occ.
laciniatus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 118.—Calif.
oblongifolius, Nutt. l. c. = pusillus.
pulchellus, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 50 = curvipes.
pusillus, *Hook. Ic. Pl.* t. 42.—Am. bor. occ.
radians, *Benth. Pl. Hartw.* 297.—Calif.
runcinatus, G. Don, Gen. Syst. i. 196 = curvipes.

THYSANOCHILUS, Falc. in Proc. Linn. Soc. i. (1839) 14. *ORCHIDEAE* (an **Cyrtopera**? Lindl.).

THYSANOLAENA, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 180. *GRAMINEAE*, Benth. & Hook. f. iii. 1120.
MYRIACHAETA, Zoll. & Mor. Verz. Pl. Zoll. 101 (1845–46).
acarifera, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 181 = T. Agrostis.
Agrostis, *Nees, in Edinb. N. Phil. Journ.* xviii. (1835) 180.—As. trop.

THYSANOSPERMUM, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 168. *RUBIACEAE*, Benth. & Hook. f. ii. 43.
diffusum, *Champ. l. c.*—Ins. Hongkong.

THYSANOTUS, R. Br. Prod. 282 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 750.
CHLAMYSPORUM, Salisb. Parad. Lond. t. 103 (1809), nomen prius.
ISANDRA, Salisb. Gen. Pl. Fragm. 67 (1866).
THYSANELLA, Salisb. l. c. (1866).
anceps, Endl. in Lehm. Pl. Preiss. ii. 37 = dichotomus.
anceps, *Lindl. Swan River App.* 58.—Austral.
Arbuscula, *Baker, in Journ. Linn. Soc.* xv. (1876) 339.—Austral.
asper, *Lindl. Swan River App.* 58.—Austral.
Banksii, R. Br. Prod. 283 = tuberosus.
Baueri, *R. Br. l. c.*—Austral.
brevipes, Endl. in Lehm. Pl. Preiss. ii. 38 = multiflorus.
chinensis, *Benth. Fl. Hongk.* 372.—Austral.; China; Ins. Philipp.
chrysanthereus, F. Muell. Fragm. v. 202 = chinensis.
deformis, R. Br. Prod. 284 = dichotomus.
dichotomus, *R. Br. l. c.*—Austral.
divaricatus, R. Br. l. c. = dichotomus.
Drummondii, *Baker, in Journ. Linn. Soc.* xv. (1876) 341.—Austral.
elatior, R. Br. Prod. 283 = tuberosus.
elongatus, R. Br. l. c. = dichotomus.
exasperatus, *F. Muell. Fragm.* i. 21.—Austral.
exiliflorus, *F. Muell. in Trans. Roy. Soc. S. Austral.* iv. (1882) 112.—Austral.
flexuosus, R. Br. Prod. 284 = dichotomus.
glaucus, *Endl. in Lehm. Pl. Preiss.* ii. 38.—Austral.
gracilis, R. Br. Prod. 283 = dichotomus.
hispidulus, R. Br. l. c. 285 = triandrus.
humilis, F. Muell. Fragm. i. 22 = Baueri.

THYSANOTUS:—
intricatus, Lindl. Bot. Reg. (1838) Misc. 59; Paxt. Mag. Bot. vi. (1839) 243 = dichotomus.
isantherus, *R. Br. Prod.* 283.—Austral.
isantherus, Lindl. in Bot. Reg. t. 655 = tuberosus.
junceus, *R. Br. Prod.* 283.—Austral.
Lindleyanus, Endl. in Lehm. Pl. Preiss. ii. 36 = dichotomus.
Manglesianus, Kunth, Enum. Pl. iv. 616 = Patersoni.
Menziesii, R. Br. Prod. 284 = Patersoni.
micranthus, Endl. in Lehm. Pl. Preiss. ii. 36 = Corynotheca dichotoma.
multiflorus, *R. Br. Prod.* 285.—Austral.
nanus, Endl. in Lehm. Pl. Preiss. ii. 38 = triandrus.
paniculatus, R. Br. Prod. 283 = tuberosus.
Patersoni, *R. Br. l. c.* 284.—Austral.
pauciflorus, *R. Br. l. c.* 285.—Austral.
proliferus, Lindl. Bot. Reg. (1838) t. 8 = multiflorus.
scaber, *Endl. in Lehm. Pl. Preiss.* ii. 37.—Austral.
sparteus, R. Br. Prod. 283 = dichotomus.
tenellus, *Endl. in Lehm. Pl. Preiss.* ii. 37.—Austral.
tenuis, Endl. l. c. = tuberosus.
tenuis, Lindl. Bot. Reg. (1838) t. 50 = dichotomus?
thyrsoideus, *Baker, in Journ. Linn. Soc.* xv. (1876) 336.—Austral.
triandrus, *R. Br. Prod.* 284.—Austral.
tuberosus, *R. Br. l. c.* 282.—Austral.
undulatorum, Behr, ex Baker, in Journ. Linn. Soc. xv. (1876) 336 = Baueri.
volubilis, R. Br. Prod. 284 = Patersoni?

THYSANTHA, Hook. Ic. Pl. t. 590 (1843) = THISANTHE, Eckl. & Zeyh. = **Tillaea**, Mich. (Crassul.).

THYSANUS, Lour. Fl. Cochinch. 284 (1790). Genus dubium *CONNARACEARUM*, Benth. & Hook. f. i. 431.
cochinchinensis, DC. Prod. ii. 91 = T. Palala.
Palala, *Lour. Fl. Cochinch.* 284.—China.

THYSSELINUM, Adans. Fam. ii. 100 (1763) = **Peucedanum**, Tourn. (Umbell.).
angustifolium, Reichb. Fl. Germ. Excurs. 453 = P. palustre.
cantabrigense, Hoffm. Gen. Umbell. ed. I. 154 = Selinum sylvestre.
Crouanorum, Bor. in Bull. Soc. Bot. Fr. xx. (1878) 30.—Gallia.
davuricum, Spreng. Syst. i. 895 = Angelica dahurica.
grandiflorum, Moench, Meth. 85 = Pleurospermum austriacum.
involucratum, Moench, l. c. (= *Pachypleurum alpinum*).—Sibir. altaic.
lucidum, Moench, Meth. Suppl. 31.—Hab.?
nigrum, Moench, Meth. 84 = Ligusticum pyrenaeum.
palustre, Hoffm. Gen. Umbell. ed. I. 154 = P. palustre.
Plinii, Spreng. Spec. Umbellif. 69 = P. palustre.
Schiwerekii, Bess. Enum. Pl. Volh. 12 = P. palustre.
silvestre, Reichb. Fl. Germ. Excurs. 453 = seq.
sylvestre, Hoffm. Gen. Umbell. ed. I. 154 = P. palustre.

TIARANTHUS, Herb. Amaryll. 202, 206 (1837) = **Pancratium**, Dill. (Amaryll.).

TIARELLA, Linn. Diss. Chen. 29 (1751); Amoen. Acad. iii. 17 (1787). *SAXIFRAGEAE*, Benth. & Hook. f. i. 637.
BLONDIA, Neck. Elem. ii. 115 (1790).
OREOTRYS, Rafin. Fl. Tellur. ii. 74 (1836).
PETALOSTEIRA, Rafin. l. c. (1836).
alternifolia, Fisch. ex Ser. in DC. Prod. iv. 50 = Tellima grandiflora.
biternata, Vent. Jard. Malm. 54. t. 54 = Astilbe decandra.
bracteata, Torr. in Ann. Lyc. N. York, ii. (1828) 204 = Heuchera bracteata.
colorans, R. Grah. in Edinb. N. Phil. Journ. (Jul.-Oct. 1829) 349 = Heuchera glabra.
cordifolia, *Linn. Sp. Pl.* 405.—As. et Am. bor.
laciniata, Hook. Fl. Bor. Am. i. 239. t. 77 = trifoliata.
Menziesii, Pursh, Fl. Am. Sept. i. 313 = Tolmiea Menziesii.
pentandra, Rafin. in Ser. Bull. Bot. i. (1830) 216, in obs. = Heuchera bracteata.
polyphylla, *D. Don, Prod. Fl. Nep.* 210.—Reg. Himal.
rhombifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 588 = trifoliata.

TIARELLA:—
stenopetala, Presl, Rel. Haenk. ii. 55 = trifoliata.
trifolia, J. F. Gmel. Syst. 1637 = trifoliata.
trifoliata, James, Cat. 181; ex S. Wats. Bibliog. Ind. Bot. N. Am. 325 = Heuchera bracteata?
trifoliata, *Linn. Sp. Pl.* 406.—As. et Am. bor.
unifolia, Retz. Obs. iii. 30 = Mitella nuda?
unifoliata, *Hook. Fl. Bor. Am.* i. 238. t. 81.—Am. bor. occ.

TIARIDIUM, Lehm. Asperif. 13 (1818); Ic. 10. t. 6 (1821) = **Heliotropium**, Tourn. (Borag.).
anisophyllum, G. Don, Gen. Syst. iv. 364 = H. indicum.
elongatum, Lehm. Asperif. 18 = H. elongatum.
heliotropioides, *Cham. in Linnaea*, iv. (1829) 453 = H. tiaridioides.
indicum, Lehm. Asperif. 14 = H. indicum.
monostachyum, G. Don, Gen. Syst. iv. 365 = H. monostachyum.
prostratum, Zipp. ex Span. in Linnaea, xv. (1 41) 335, nomen = H. prostratum?
velutinum, Lehm. Asperif. 15 = H. indicum?

TIBOUCHINA, Aubl. Pl. Gui. i. 445. t. 177 (1775).
MELASTOMACEAE, [Benth. & Hook. f. i. 743].
BRACTEARIA, DC. ex Steud. Nom. ed. II. i. 222 (1840).
CHAETOGASTRA, DC. Prod. iii. 131 (1828); Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 127.
DIPLOSTEGIUM, D. Don, in Mem. Wern. Soc. iv. (1823) 296.
HEPHESTIONIA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 36.
LASIANDRA, DC. Prod. iii. 127 (1828).
MICRANTHELLA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 347.
OREOCOSMUS, Naud. in Ann. l. c. 37 (1849).
PURPURELLA, Naud. in Ann. l. c. 301 (1849).
SAVASTANA, Scop. Introd. 213 (1777).
Ackermannii, *Cogn. in Mart. Fl. Bras.* xiv. III. 365.—Bras.
adenostemon, *Cogn. l. c.* 332.—Bras.
aegopogon, *Cogn. l. c.* 383.—Bras.
aemula, *Cogn. l. c.* 412.—Bras.
angustifolia, *Cogn. l. c.* 396.—Bras.
arborea, *Cogn. l. c.* 299.—Bras.
arenaria, *Cogn. l. c.* 329.—Bras.
Arstedii, Cogn. l. c. 297, sphalm. = Oerstedii.
aspera, *Aubl. Pl. Gui.* i. 446. t. 177.—Guian.; Bras.
asperior, *Cogn. in Mart. Fl. Bras.* xiv. III. 366.—Bras.
australis, *Cogn. l. c.* 392.—Bras.
axillaris, *Cogn. l. c.* 396.—Bras.
barbigera, *Baill. in Adansonia*, xii. (1877) 75.—Bras.
Benthamiana, *Cogn. in Mart. Fl. Bras.* xiv. III. 391.—Bras.
Bergiana, *Cogn. l. c.* 316.—Bras.
bipenicillata, *Cogn. l. c.* 385.—Am. trop.
Blanchetiana, *Cogn. l. c.* 351.—Bras.
caldensis, *Cogn. l. c.* 306.—Bras.
Candolleana, *Cogn. l. c.* 339.—Bras.
canescens, *Cogn. l. c.* 298.—Bras.
cardinalis, *Cogn. l. c.* 371.—Bras.
cerastifolia, *Cogn. l. c.* 403.—Bras.
Chamissoana, *Cogn. l. c.* 349.—Bras.
cinerea, *Cogn. l. c.* 416.—Bras.
cisplatensis, *Cogn. l. c.* 407.—Bras.
Claussenii, *Cogn. l. c.* 310.—Bras.
clidemioides, *Cogn. l. c.* 398.—Bras.
clinopodifolia, *Cogn. l. c.* 411.—Bras.
collina, *Cogn. l. c.* 318.—Bras.
cordifolia, *Cogn. l. c.* 393.—Bras.
coronata, *Cogn. l. c.* 307.—Peruv.
corymbosa, *Cogn. l. c.* 363.—Bras.
cujabensis, *Cogn. l. c.* 382.—Bras.
debilis, *Cogn. l. c.* 401.—Bras.
decemcostata, *Cogn. l. c.* 331.—Bras.
dendroides, *Cogn. l. c.* 369.—Bras.
divaricata, *Cogn. l. c.* 397.—Bras.
dubia, *Cogn. l. c.* 415.—Bras.
Eichleri, *Cogn. l. c.* 370.—Bras.
elegans, *Cogn. l. c.* 323.—Bras.
estrellensis, *Cogn. l. c.* 342.—Bras.

TIBOUCHINA:—
exasperata, *Cogn. in Mart. Fl. Bras.* xiv. III. 377.—Bras.
fissinervia, *Cogn. l. c.* 343.—Bras.
floribunda, *Cogn. l. c.* 312.—Bras.
formosa, *Cogn. l. c.* 348.—Bras.
Fothergillae, *Cogn. l. c.* 320.—Bras.
foveolata, *Cogn. l. c.* 307.—Bras.
fragilis, Baill. Hist. Pl. vii. 7 = Arthrostema fragile.
Francavillana, *Cogn. in Mart. Fl. Bras.* xiv. III. 314.—Bras.
frigidula, *Cogn. l. c.* 328.—Bras.
Gardneri, *Cogn. l. c.* 334.—Bras.
Gardneriana, *Cogn. l. c.* 319.—Bras.
Gaudichaudiana, *Baill. Hist. Pl.* vii. 7. f. 9.—Bras.
Glazioviana, *Cogn. in Mart. Fl. Bras.* xiv. III. 325.—Bras.
gracilis, *Cogn. l. c.* 386.—Bras.
grandifolia, *Cogn. l. c.* 335.—Bras.
granulosa, *Cogn. l. c.* 340.—Bras.
grossa, *Cogn. l. c.* 297.—Am. trop.
herbacea, *Cogn. l. c.* 408.—Bras.
Herincquiana, *Cogn. l. c.* 399.—Bras.
heteromalla, *Cogn. l. c.* 336.—Bras.
hieracioides, *Cogn. l. c.* 389.—Bras.
hirsutissima, *Cogn. l. c.* 356.—Bras.
holosericea, *Baill. Hist. Pl.* vii. 34.—Bras.
hospita, *Cogn. in Mart. Fl. Bras.* xiv. III. 417.—Bras.
hygrophila, *Cogn. l. c.* 413.—Bras.
inappendiculata, Baill. Adansonia, xii. (1877) 74 = mollis.
intermedia, *Cogn. in Mart. Fl. Bras.* xiv. III. 367.—Bras.
Itatiaiae, *Cogn. l. c.* 414.—Bras.
Karstenii, *Cogn. l. c.* 381.—Venezuela; N. Granat.
Langsdorffiana, *Baill. Hist. Pl.* vii. 34.—Bras.
lasiophylla, *Cogn. in Mart. Fl. Bras.* xiv. III. 297.—Guian.
lepidota, *Baill. Adansonia*, xii. (1877) 74.—Am. trop.
Lhotzkyana, *Cogn. in Mart. Fl. Bras.* xiv. III. 357.—Bras.
Lindeniana, *Cogn. l. c.* 347.—N. Granat.
longifolia, *Baill. Adansonia*, xii. (1877) 74.—Am. trop.
macrochiton, *Cogn. in Mart. Fl. Bras.* xiv. III. 313.—Bras.
macrostachya, Baill. Hist. Pl. vii. 6 = Heeria macrostachya.
Martialis, *Cogn. in Mart. Fl. Bras.* xiv. III. 346.—Bras.
Martiusiana, *Cogn. l. c.* 327.—Bras.
Mathaei, *Cogn. l. c.* 380.—Peruv.
Maximiliana, *Baill. Hist. Pl.* vii. 34.—Bras.
melastomoides, *Cogn. in Mart. Fl. Bras.* xiv. III. 377.—Bras.
minor, *Cogn. l. c.* 390.—Bras.
minutiflora, *Cogn. l. c.* 415.—Bras.
mollis, *Cogn. l. c.* 349.—Am. trop.
Moricandiana, *Baill. Adansonia*, xii. (1877) 75.—Bras.
Mosenii, *Cogn. in Mart. Fl. Bras.* xiv. III. 393.—Bras.
multiceps, *Cogn. l. c.* 326.—Bras.
multiflora, *Cogn. l. c.* 333.—Bras.
multinervia, Baill. Adansonia, xii. (1877) 75 = Macairea multinervia.
mutabilis, *Cogn. in Mart. Fl. Bras.* xiv. III. 300.—Bras.
nervulosa, *Cogn. l. c.* 315.—Bras.
nitida, *Cogn. l. c.* 404.—Am. trop.
ochypetala, *Baill. Adansonia*, xii. (1877) 75.—Peruv.
Oerstedii, *Cogn. in Mart. Fl. Bras.* xiv. III. 297 (Arstedii).—Am. centr.
organensis, *Cogn. l. c.* 308.—Bras.
ornata, *Baill. Adansonia*, xii. (1877) 74.—Am. trop.
paleacea, *Cogn. in Mart. Fl. Bras.* xiv. III. 373.—N. Granat.
pallida, *Cogn. l. c.* 316.—Bras.
papyrifera, *Cogn. l. c.* 378.—Bras.
parviflora, *Cogn. l. c.* 410.—Bras.
pauciflora, *Cogn. l. c.* 348.—Bras.
pogonanthera, *Cogn. l. c.* 381.—Bras.
pulchra, *Cogn. l. c.* 301.—Bras.
Raddiana, *Cogn. l. c.* 303.—Bras.
rediviva, *Cogn. l. c.* 399.—Bras.

TIBOUCHINA:—
Regeliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 356.—Bras.
Regnellii, *Cogn. l. c.* 305.—Bras.
Reichardtiana, *Cogn. l. c.* 361.—Bras.
reticulata, *Cogn. l. c.* 298.—Ecuador.
Riedeliana, *Cogn. l. c.* 368.—Bras.
robusta, *Cogn. l. c.* 394.—Bras.
rotundifolia, *Cogn. l. c.* 370.—Bras.
Saldanhaei, *Cogn. l. c.* 302.—Bras.
salviaefolia, *Cogn. l. c.* 364.—Bras.
scaberrima, *Cogn. l. c.* 400.—Bras.
scrobiculata, *Cogn. l. c.* 341.—Bras.
sebastianopolitana, *Cogn. l. c.* 409.—Bras.
Sellowiana, *Cogn. l. c.* 304.—Bras.
semidecandra, *Cogn. l. c.* 309.—Bras.
setulosa, *Cogn. l. c.* 338.—Bras.
simplicicaulis, *Cogn. l. c.* 406.—Bras.
Spruceana, *Cogn. l. c.* 376.—Bras.
stenocarpa, *Cogn. l. c.* 344.—Bras.
sulcata, *Cogn. l. c.* 337.—Bras.
Thereminiana, *Cogn. l. c.* 310.—Bras.
trichopoda, *Baill. Adansonia*, xii. (1877) 75.—Bras.
tuberosa, *Cogn. in Mart. Fl. Bras.* xiv. III. 395.—Bras.
urceolaris, *Cogn. l. c.* 355.—Bras.
ursina, *Cogn. l. c.* 351.—Bras.
Urvilleana, *Cogn. l. c.* 358.—Bras.
Valtherii, *Cogn. l. c.* 317.—Bras.
velutina, *Cogn. l. c.* 350.—Bras.
versicolor, *Cogn. l. c.* 405.—Bras.
verticillaris, *Cogn. l. c.* 379.—Bras.
villosissima, *Cogn. l. c.* 330.—Bras.
viminea, *Cogn. l. c.* 359.—Bras.
virgata, *Cogn. l. c.* 365.—Bras.
Weddellii, *Cogn. l. c.* 311.—Bras.

TIBUCHINA, Jaume St. Hil. Expos. ii. 173 (1805) = praec.

TICANTO, Adans. Fam. ii. 319 (1763) = **Caesalpinia**, Linn. (Legumin.).
Nuga, Medic. Theod. 52 = C. Nuga.

TICOREA, Aubl. Pl. Gui. ii. 689. t. 277 (1775).
RUTACEAE, Benth. & Hook. f. i. 285.
OZOPHYLLUM, Schreb. Gen. 452 (1791).
aculeata, Blanco, Fl. Filip. ed. I. 85 = Leea aculeata.
foetida, *Aubl. Pl. Gui.* ii. 689. t. 277.—Guian.
longiflora, *DC. in Mém. Mus. Par.* ix. (1822) 146. t. 9.—Guian.
nitida, *Presl, Symb. Bot.* i. 28.—Am. austr.
pedicellata, *DC. in Mém. Mus. Par.* ix. (1822) 145. t. 8.—Guian.

TICOREA, A. St. Hil. in Bull. Soc. Philom. (1823) 132 = **Galipea**, Aubl. (Rutac.).
bracteata, A. St. Hil. ex DC. Prod. i. 730 = G. bracteata.
febrifuga, A. St. Hil. in Mém. Mus. Par. x. (1823) 292 = G. multiflora.
jasminiflora, A. St. Hil. in Bull. Soc. Philom. (1823) 132; et in Mém. Mus. Par. x. (1823) 291 = G. multiflora.
simplicifolia, A. St. Hil. ex DC. Prod. i. 730 = G. simplicifolia.

TIEDEMANNIA, DC. Mém. Ombell. 51. t. 12 (1829); et Prod. iv. 187 (1830) = **Peucedanum**, Tourn. (Umbell.).
teretifolia, DC. Prod. iv. 187 = P. teretifolium.

TIEDMANNIA, Torr. & Gray, Fl. N. Am. i. 630 (1840), sphalm. = praec.

TIGAREA, Aubl. Pl. Gui. ii. 917. t. 350, 351 (1775) = **Doliocarpus**, Roland. (Dillen.).
aspera, Aubl. l. c. 918 = Tetracera aspera.
dentata, Aubl. l. c. 920 = D. Rolandri.

TIGAREA, Pursh, Fl. Am. Sept. i. 333. t. 15 (1814) = **Purshia**, DC. (Rosac.).
tridentata, Pursh, l. c. = P. tridentata.

TIGLIUM, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 418 = **Croton**, Linn. (Euphorb.).
Klotzschianum, Wight, Ic. t. 1914 = C. caudatus.
officinale, Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 418 = C. Tiglium.

TIGRIDIA, Juss. Gen. 57 (1789). *IRIDEAE*, Benth. & Hook. f. iii. 690.
BEATONIA, Herb. in Bot. Mag. sub t. 3779 (1840).
PARDINIA, Herb. in Lindl. Bot. Reg. (1844) Misc. 66, in obs., partim.
atrata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 136.—Mexic.
augusta, *Drapiez, in Van-Geel, Sert. Bot.* iv. (1832) t. 131; *et ex C. Morr. Hortic. Belg.* v. (1838) 279. t. 110.—Mexic.
azurea, Hort.—Cf. Fl. des Serres, ix. (1853–54) 139 = Cypella plumbea.
coelestis, Hort. Berol.—Cf. l. c. = Cypella plumbea.
conchiiflora, Sweet, Brit. Flow. Gard. Ser. I. ii. t. 128 = T. Pavonia.
curvata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 136.—Mexic.
Dugesii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 375.—Mexic.
grandiflora, Salisb. in Trans. Hort. Soc. i. (1812) 309 = T. Pavonia.
Herberti, Herb. in Bot. Mag. t. 2599 = Cypella Herberti.
Herbertiana, Steud. Nom. ed. II. ii. 686 = praec.
Houttei, Roezl, in Fl. des Serres, xxi. (1875) 49 (Hydrotaeniae sp.).—Mexic.
lutea, *Link, Klotzsch & Otto, Ic. Pl. Hort. Berol.* 85. t. 34.—Chili; Peruv.
oxypetala, R. Morris, Fl. Conspic. t. 20 = T. Pavonia.
Pavonia, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 246; *Pers. Syn.* i. 50.—Mexic.
speciosa, Poit. in Rev. Hortic. Sér. II. ii. (1843–44) 447 = T. Pavonia.
Van-Houttei, Roezl, in Fl. des Serres, xxi. (1875) t. 2174 = *Houttei*.
violacea, *Schied. & Schlecht. in Otto & Dietr. Allg. Gartenz.* vi. (1838) 233.—Mexic.

TILCO, Adans. Fam. ii. 498 (1763) = **Fuchsia**, Plum. (Onagrar.).

TILCUSTA, Rafin. Fl. Tellur. iv. 15 (1836) = **Tupistra**? Ker-Gawl. (Liliac.)
nepalensis, Rafin. l. c. = Tupistra aurantiaca?

TILDENIA, Miq. in Inst. Versl. en Med. 1842 (1843) 82 = **Peperomia**, Ruiz & Pav. (Piperac.).
mexicana, Miq. l. c. 83 = P. mexicana.
peruviana, Miq. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 483 = P. umbilicata.

TILESIA, G. F. W. Mey. Prim. Fl. Esseq. 251 (1818) = **Wulffia**, Neck. (Compos.).
capitata, G. F. W. Mey. l. c. 254 = W. stenoglossa.
spilanthoides, Griseb. Cat. Pl. Cub. 154 (= *Calydermos spilanthoides*).—Cuba.

TILESIA, Thunb. ex Steud. Nom. ed. II. ii. 686 (1841), nomen = **Gladiolus**, Tourn. (Irid.).
incarnata, Thunb. ex Steud. l. c., nomen = G. incarnatus?
spiralis, Thunb. ex Steud. l. c., nomen = G. tristis.

TILIA, [Tourn.] Linn. Syst. ed. I (1735). *TILIACEAE*, Benth. & Hook. f. i. 236.
LINDNERA, Reichb. Handb. 303 (1837).
TILIOIDES, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 213.
acuminata, Opiz, Seznam, 98 = vulgaris.
affinis, Opiz, Natural. n. xi. (1826) 461 = platyphyllos.
alba, *Ait. Hort. Kew.* ed. I. ii. 230.—Am. bor.
alba, Waldst. & Kit. Pl. Rar. Hung. i. 3. t. 3 = tomentosa.
americana, Duroi, Harbk. Baumz. ii. 467 = alba.
americana, *Linn. Sp. Pl.* 514.—Am. bor.
angulata, Hayne, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 27 = cordata?
angulata, Hayne, ex Bayer, l. c. 33 = platyphyllos.
apiculata, Court. in Nouv. Mém. Acad. Brux. ix. (1835) n. VI. 15 = platyphyllos.
argentea, Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 48 = mandshurica.
argentea, DC. Cat. Hort. Monsp. (1813) 150 = tomentosa.
asplenifolia, Hort. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 41 = platyphyllos.
aurea, Juengst. Fl. Westfalen, 194 = platyphyllos.

TILIA:—
barbata, Stokes, Bot. Mat. Med. iii. 201 = platyphyllos, vulgaris, cordata.
begonifolia, Stev. in Bull. Soc. Nat. Mosc. xxix. (1846) I. n. II. 326, in textu = rubra.
betulaefolia, Hofm. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 23 = cordata.
bicuspidata, Court. in Nouv. Mém. Acad. Brux. ix. (1835) n. VI. 13 = platyphyllos.
bohemica, Opiz, Seznam, 97, in syn. = cordata.
bohemica, Ortm. ex Nym. Consp. 131 = platyphyllos.
bohemica, Rupr. Fl. Cauc. 255, 299 = cordata?
bracteolata, Host, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 25 = cordata.
canadensis, Michx. Fl. Bor. Am. i. 306 = americana.
caroliniana, Mill. Gard. Dict. ed. VIII. n. 4 = americana.
caucasica, Rupr. Fl. Cauc. 253 (spp. plurim.).
cinerea, Rafin. Alsog. Am. 46 (Quid?).—Am. bor.
colchica, Stev. ex Rupr. Fl. Cauc. 299 = dasystyla.
communis, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 875 = platyphyllos, cordata.
corallina Sm. in Rees, Cycl. xxxv. n. 2 = platyphyllos.
cordata, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.
cordifolia, Bess. Prim. Fl. Galic. i. 343 = vulgaris.
corinthiaca, Bosc, Nouv. Cours Agric. xiii. 139 = rubra.
corylifolia, Host, Fl. Austr. ii. 59 = platyphyllos.
corymbosa, Ortm. in Opiz, Nom. 25 = platyphyllos.
crenata, Rafin. Alsog. Am. 44 (Quid?).—Am. bor.
dasystyla, Loud. Arb. Brit. i. 366 = euchlora.
dasystyla, *Stev. in Bull. Soc. Nat. Mosc.* iv. (1832) I. 260.—Orient.
euchlora, *C. Koch, in Wochenschr.* ix. (1866) 284; *Ind. Sem. Hort. Berol.* (1867) *App.* i. 9.—Hab.?
europaea, Linn. Sp. Pl. 514.—Cf. platyphyllos, vulgaris, cordata.
flaccida, Host, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 46 = platyphyllos.
flava, Wolny, ex Rochel, Pl. Banat. Rar. 2, in textu = platyphyllos.
flavescens, A. Br. in Doell, Rhein. Fl. 672 = americana.
flavescens, Kit. in Linnaea, xxxii. (1868) 548 = platyphyllos?
floribunda, A. Br. in Doell, Rhein. Fl. 672 = americana.
floribunda, Reichb. Ic. Fl. Germ. vi. 58. t. 314 = vulgaris.
foemina, Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 24 = cordata.
fulva, Rafin. Alsog. Am. 45 (Quid?).—Am. bor.
fuscata, Rafin. l. c. 45 (Quid?).—Am. bor.
glabra, Vent. in Mém. Instit. Par. iv. An xi. [1803] 9. t. 2 = americana.
glauca, Rafin. Alsog. Am. 45 (Quid?).—Am. bor.
glauca, Schwein. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 53 = alba.
grandifolia, Ehrh. Beitr. v. 158 = platyphyllos.
grandifolia, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 236 = rubra.
grandifolia, Hort. ex Steud. Nom. ed. II. ii. 687 = languida.
grata, Salisb. Prod. 367 = americana.
heterophylla, Hort. ex C. Koch, Dendrol. i. 480 = americana.
heterophylla, Vent. in Mém. Instit. Par. iv. An xi. [1803] 16. t. 5 = alba.
Hoffmanniana, Opiz, Seznam, 97 = cordata.
hollandica, Duroi, ex Steud. Nom. ed. I. 837 = americana.
hollandica, Herb. Petrop. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 30 = cordata.
hollandica, Hort. ex C. Koch, Dendrol. i. 471 = platyphyllos.
Hostii, Opiz, Seznam, 98 = platyphyllos.
hybrida, [Behlan, in] Bechst. Forstb. ed. V. 161 = vulgaris.
inciso-dentata, Hort. ex C. Koch, Dendrol. i. 480 = americana.
intermedia, DC. Prod. i. 513 = vulgaris.
intermedia, Host, Fl. Austr. ii. 61 = platyphyllos.
intermedia, Tenore, Fl. Nap. iv. 309 = cordata.
laeta, Salisb. Prod. 367.—Cf. platyphyllos, vulgaris, cordata.
languida, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 220.—Hab.?
latebracteata, Host, Fl. Austr. ii. 60 = cordata.
latifolia, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.

TILIA:—
latifolia, Salisb. Prod. 367 = americana.
laxiflora, Michx. Fl. Bor. Am. i. 306 = americana.
laxiflora, Pursh, Fl. Am. Sept. ii. 363 = alba.
leptophylla, Hort. Berol. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 58 = americana.
longifolia, Rafin. Alsog. Am. 44 (Quid?).—Am. bor.
lucida, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.
macrocarpa, Hort. ex C. Koch, l. c. 473 = platyphyllos.
macrophylla, Hort. ex Steud. Nom. ed. II. ii. 687 = alba.
macrophylla, Leveillé, in Demid. Voy. Russ. Mér. ii. 208 = cordata.
macropoda, Court. in Nouv. Mém. Acad. Brux. ix. (1835) n. VI. 10 = platyphyllos.
mandshurica, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1856) 124.—Manchur.; Japon.
mellita, Presl, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 36 = platyphyllos.
mexicana, *Benth. Pl. Hartw.* 35.—Mexic.
mexicana, Schlecht. in Linnaea, xi. (1837) 377 = americana.
Michauxii, Nutt. N. Am. Sylva, ed. II. i. 108 = alba.
microphylla, Vent. in Mém. Instit. Par. iv. An xi. [1803] 5. t. 1. f. 1 = cordata.
Miqueliana, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 434.—Japon.
mississippiensis, Bosc, Encyc. Agric. vii. 748 = americana.
mollis, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 336; Ortm. ex Opiz, Nom. 22 = platyphyllos.
mongolica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 433.—Mongolia; China.
multiflora, Hort. ex Vent. in Mém. Instit. Par. iv. An xi. [1803] 12 = americana.
multiflora, Ledeb. Fl. Ross. i. 442 = dasystyla.
mutabilis, Host, Fl. Austr. ii. 60 = platyphyllos.
neglecta, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 341 = americana.
Nickerlii, Opiz, ex Nym. Consp. 131 = platyphyllos.
nigra, Borkh. Handb. Forstb. ii. 1219 = americana.
obliqua, Opiz, Natural. n. xi. (1826) 462; Host, Fl. Austr. ii. 62 = platyphyllos.
obliquifolia, Ortm. ex Steud. Nom. ed. II. ii. 687 = platyphyllos.
oblongata, Court. in Nouv. Mém. Acad. Brux. ix. (1835) n. VI. 12 = platyphyllos.
officinarum, Crantz, Stirp. Austr. ed. I. fasc. ii. 61; ed. II. 96 = *europaea*.
ovata, Ortm. in Flora, xviii. (1835) 502 = platyphyllos.
oxycarpa, Reichb. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 37 = vulgaris.
pallida, Salisb. Prod. 367 = alba.
pallida, Wierzb. ex Reichb. Ic. Fl. Germ. vi. 58. t. 513 = platyphyllos.
pannonica, Jacq. f. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 48 = petiolaris.
parviflora, Boiss. Fl. Orient. i. 846, sphalm. = seq.
parvifolia, Ehrh. Beitr. v. 159; vii. 53 = cordata.
pauciflora, Hayne, Arzn. Gew. iii. t. 48 = platyphyllos.
pedunculares, Delile, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 58 = alba.
pekinensis, Rupr. ex Maxim. Prim. Fl. Amur. 469 = mandschurica.
petiolaris, *DC. Prod.* i. 514.—Hungar.
pilosa, Presl, ex Ortm. in Flora, xviii. (1835) 501 = platyphyllos.
platyphylla, C. A. Mey. Verz. Pfl. Cauc. 224 = rubra.
platyphyllos, Bieb. Fl. Taur. Cauc. ii. 6 = dasystyla.
platyphyllos, *Scop. Fl. Carn.* ed. II. i. 373.—Europ.
platyptera, Court. in Nouv. Mém. Acad. Brux. ix. (1835) n. VI. 11 = platyphyllos.
Podhorsciana, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 165, nomen.—Podolia.
polyantha, Schur, Enum. Pl. Transs. 131 = cordata.
praecox, A. Br. in Doell, Rhein. Fl. 673 = americana.
praecox, Host, Fl. Austr. ii. 60 = platyphyllos.
Preslii, Ortm. in Opiz, Nom. 24 = platyphyllos.
pubescens, Ait. Hort. Kew. ed. I. ii. 229 = americana.
pubescens, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.
pyramidalis, Host, Fl. Austr. ii. 61 = platyphyllos.
riparia, Rafin. Alsog. Am. 45 (Quid?).—Am. bor.
rotundifolia, Vent. in Mém. Instit. Par. iv. An xi. [1803] 12. t. 4 = tomentosa.

TILIA:—
rubra, *DC. Cat. Hort. Monsp.* (1813) 150.—Tauria.
rugosa, Host, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 22 = cordata.
semicuneata, *Rupr. Fl. Cauc.* 252, in textu.—Sibir.
septemtrionalis, Rupr. Fl. Ingr. i. 219 = cordata.
serratifolia, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.
sibirica, Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 23 = cordata.
speciosa, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.
spectabilis, Host, ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 42 = platyphyllos.
sphaerocarpa, Hort. ex C. Koch, Dendrol. i. 473 = platyphyllos.
stenopetala, Rafin. Fl. Ludov. 92 = americana?
stipulata, Gilib. Exercit. i. 329 = platyphyllos, vulgaris, cordata.
sylvestris, Desf. Tabl. ed. I. 152 = cordata.
Tecksiana, Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 25 = cordata.
tenuifolia, Host, Fl. Austr. ii. 61 = platyphyllos.
tomentosa, *Moench, Verzeich.* 137.—Europ.
translucens, Opiz, ex Nym. Consp. 131 = platyphyllos.
triflora, Puerari, in Hornem. Hort. Hafn. 493 = rubra.
triflora, Schrad. ex Bayer, in Verh. Zool.-Bot. Ges. Wien, xii. (1862) 40 = platyphyllos.
truncata, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 342 = americana.
Tučekii, Opiz, in Lotos, iv. (1854) 103 = platyphyllos.
turbinata, Hort. ex C. Koch, Dendrol. i. 473 = platyphyllos.
ulmifolia, Scop. Fl. Carn. ed. II. i. 373 = cordata.
umbellata, Rafin. Alsog. Am. 46 (Quid?).—Am. bor.
variegata, Hort. ex C. Koch, Dendrol. i. 472 = platyphyllos.
vitifolia, Host, Fl. Austr. ii. 59 = platyphyllos.
vulgaris, *Heyne, Arzn. Gew.* iii. t. 47.—Europ.

TILIACORA, Colebr. in Trans. Linn. Soc. xiii. (1822) 53. *MENISPERMACEAE*, Benth. & Hook. f. i. 36.
BAGALATTA, Roxb. ex Reichb. Consp. 86 (1828).
BRAUNEA, Willd. Sp. Pl. iv. 797 (1805), ex parte.
abnormalis, Miers, Contrib. iii. 78 = racemosa.
acuminata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 39 = racemosa.
cuspidiformis, Miers, l. c. Ser. III. xiv. (1864) 254; Contrib. iii. 77 = racemosa.
fraternaria, Miers, Contrib. iii. 77 = racemosa.
funifera, *Oliver, Fl. Trop. Afr.* i. 44.—Afr. trop.
racemosa, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 67.—Ind. or.; Java.

TILINGIA, Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 97 = **Selinum**, Linn. (Umbell.).
ajanensis, Regel & Tiling, l. c.—Sibir. or.

TILIOIDES, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 213 = **Tilia**, Tourn.

TILLAEA, Mich. ex Linn. Syst. ed. I (1735). *CRASSULACEAE*, Benth. & Hook. f. i. 657.
BULLIARDA, DC. in Bull. Soc. Philom. iii. (1801) n. 49. 1.
COMBESIA, A. Rich. Tent. Fl. Abyss. i. 307 (1847).
DISPOROCARPA, A. Rich. l. c. 307, in syn. (1847).
HELOPHYTUM, Eckl. & Zeyh. Enum. 288 (1836).
MESANCHUM, Dulac, Fl. Hautes-Pyr. 320 (1867).
abyssinica, Walp. Ann. ii. 666 = aquatica.
adscendens, Nees, in Lehm. Pl. Preiss. i. 277 = Sieberiana.
alata, Viv. Pl. Aegypt. Dec. 16 = pharnaceoides.
alsinoides, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 192.—Afr. trop.
andicola, *Phil. in Anal. Univ. Chil.* (1872) 720.—Chili.
angustifolia, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 558.—Am. bor.
aquatica, Lam. Illustr. i. 361 = Vaillantii.
aquatica, *Linn. Sp. Pl.* 128.—Europ.; Afr. trop.
aquatica, Schkuhr, in Usteri, Ann. Bot. ii. (1791) 21 = prostrata.
ascendens, Eaton, Man. Bot. N. Am. ed. V. 419 = simplex, *Nutt.*
brevifolia, *Walp. Rep.* ii. 251.—Afr. austr.
capensis, *Linn. f. Suppl.* 129.—Afr. austr.
chiloënsis, *C. Gay, Fl. Chil.* ii. 532.—Chili.

TILLAEA:—
Closiana, *C. Gay, Fl. Chil.* ii. 531.—Chili.
colorata, Nees, in Lehm. Pl. Preiss. i. 277 = Sieberiana.
connata, *Ruiz & Pav. Fl. Per.* i. 70.—Ecuador; Peruv.
cyathiformis, Steud. Nom. ed. II. ii. 687, sphalm. = *Tillandsia cyathiformis.*
cymosa, Nutt. Gen. Am. i. 110 = Diamorpha pusilla.
debilis, *Colenso, ex Hook. f. Fl. N. Zel.* i. 75.—N. Zel.
decumbens, *Willd. Sp. Pl.* i. 721.—Afr. austr.
diffusa, Willd. ex Steud. Nom. ed. II. ii. 687 = connata.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 558.—Am. bor.
Ecklonis, *Walp. Rep.* ii. 251.—Afr. austr.
elatinoides, *Walp. l. c.*—Afr. austr.
erecta, DC. Prod. iii. 406, in syn. = Sedum caespitosum.
erecta, Hook. & Arn. Bot. Beech. Voy. 24 = minima.
filiformis, *Endl. ex Walp. Rep.* ii. 252.—Afr. austr.
fluitans, *Endl. l. c.*—Afr. austr.
Hamiltonii, *T. Kirk, ex W. Ham. in Trans. N. Z. Inst.* xvii. (1885) 292.—N. Zel.
hexandra, Lapierre, in Journ. Phys. lvi. (1802) 358 = Elatine hexandra.
inanis, Steud. Nom. ed. II. ii. 687 = capensis.
intricata, Nees, in Lehm. Pl. Preiss. i. 278 = recurva.
leptopetala, Benth. Pl. Hartw. 310 = minima.
macrantha, *Hook. f. in Hook. Ic. Pl.* t. 310.—Austral.
magellanica, *Willd. ex Schult. Mant.* iii. 345.—Reg. Magellan.
minima, *Miers, Trav.* ii. 530, nomen; *et ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 338.—Chili; Calif.
moschata, *DC. Prod.* iii. 382.—N. Zel.; Ins. Antarct.; Chili.
muscosa, Forst. f. Prod. 11 = Sieberiana.
muscosa, C. Gay, Pl. Chil. ii. 529 = minima.
muscosa, *Linn. Sp. Pl.* 129.—Europ.; Afr. bor.
Ovallei, *Phil. in Anal. Univ. Chil.* (1872) 719.—Chili.
paludosa, *Schlecht. in Lechl. Berb. Am. Austr.* 54.—Chili.
pedicellosa, *F. Muell. Fragm.* xi. 118, in obs.—Austral.
peduncularis, *Sm. in Rees, Cycl.* xxxv. n. 4.—Bras.
pedunculata, Sieber, ex DC. Prod. iii. 382 = Sieberiana.
pedunculata, *Willd. ex Schult. Mant.* iii. 345.—Lusitan.
pentandra, *Royle, Illustr. Bot. Himal.* 222, nomen; *et ex Edgew. in Trans. Linn. Soc.* xx. (1846) 50.—Ind or.; Afr. trop.
perfoliata, Linn. f. Suppl. 129 = capensis.
pharnaceoides, *Hochst. ex Steud. Nom.* ed. II. ii. 687, nomen; *et ex Britten, in Oliver, Fl. Trop. Afr.* ii. 387.—Afr. trop.
prostrata, *Schkuhr, in Usteri, Ann. Bot.* xii. (1794) 6.—Europ.?
purpurata, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 472.—Austral.
radicans, *Phil. in Anal. Univ. Chil.* (1872) 718.—Chili.
recurva, *Hook. f. Fl. Tasman.* i. 146.—Austral.
reflexa, *Endl. ex Walp. Rep.* ii. 252.—Afr. austr.
rubella, Blanco, Fl. Filip. ed. I. 75 = Bergia ammannioides.
rubescens, H. B. & K. Nov. Gen. et Sp. vi. 43 = connata.
rubra, Linn. Sp. Pl. 129; Gouan, Hort. Monsp. 77 = Sedum caespitosum.
saginoides, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 473.—Soongar.
Sieberi, Steud. Nom. ed. II. ii. 687, sphalm. = seq.
Sieberiana, *Schult. Mant.* iii. 345.—Am. austr.; Austral.; N. Zel.
simplex, *Nutt. in Journ. Acad. Philad.* i. (1817) 114.—Am. bor.
simplex, *Phil. in Anal. Univ. Chil.* (1872) 719.—Chili.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 62.—N. Zel.
Solierii, *C. Gay, Fl. Chil.* ii. 530.—Chili.
subulata, *Benth. & Hook. f. Gen.* i. 658, in obs.—Afr. trop. et austr.
trichopoda, Fenzl, ex Boiss. Fl. Orient. ii. 767 = pharnaceoides.

TILLAEA:—
trichotoma, *Walp. Rep.* ii. 251.—Afr. austr.
umbellata, Willd. Sp. Pl. i. 721 = Crassula umbellata.
Vaillantii, *Willd. l. c.* 720.—Europ.; Afr. bor.
verticillaris, DC. Prod. iii. 382 = Sieberiana.
verticillaris, Hook. Ic. Pl. t. 295 = recurva.
verticillata, Steud. Nom. ed. II. ii. 687, sphalm. = Sieberiana.

TILLANDSIA, Linn. Syst. ed. I (1735). *BROMELIACEAE*, Benth. & Hook. f. iii. 669.
ACANTHOSPORA, Spreng. Anleit. ii. 1. 255 (1817); Syst. ii. 25 (1825).
ALLARDTIA, A. Dietr. in Otto & Dietr. Allg. Gartenz. xx. (1852) 241.
AMALIA, Hort. Hisp. ex Endl. Gen. 183 (1837).
ANOPLOPHYTUM, Beer, in Flora, xxxvii. (1854) 346.
BONAPARTEA, Ruiz & Pav. Fl. Per. iii. 38 (1802).
DENDROPOGON, Rafin. Neogenyt. 3 (1825).
DIAPHORANTHEMA, Beer, in Flora, xxxvii. (1854) 349; Bromel. 153 (1857).
HEXALEPIS, Rafin. Fl. Tellur. iv. 24 (1836).
MISANDRA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 102 (1819).
PHYTARRHIZA, Vis. in Mem. Ist. Venet. v. (1855) 340.
PITYROPHYLLUM, Beer, Bromel. 79 (1857).
PLATYSTACHYS, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 11; et Beer, Bromel. 80 (1857).
RENEALMIA, Linn. Gen. ed. I. 340 (1737).
STREPSIA, Steud. Nom. ed. II. ii. 645 (1841).
VRIESIA, Lindl. Bot. Reg. (1843) t. 10.
WALLISIA, E. Morr. in Belg. Hortic. xx. (1870) 97. t. 5.
acaulis, Lindl. Bot. Reg. t. 1157 = Cryptanthus undulatus.
acorifolia, *Griseb. in Goett. Nachr.* (1864) 19.—N. Granat.
acutifolia, Hort. Angl. ex Link, Enum. Hort. Berol. i. 308, nomen (Quid?).—Hab.?
aëranthos, Desf. ex Steud. Nom. ed. II. ii. 687 = dianthoidea.
aloides, *Schlecht. & Cham. in Linnaea*, vi. (1831) 55.—Mexic.
aloifolia, Hook. Exot. Fl. t. 205 = flexuosa.
amoena, Lodd. Bot. Cab. t. 76 = Billbergia speciosa.
anceps, *Lodd. l. c.* t. 771.—Ins. Trinit.
andicola, *Gill. ex Baker, in Journ. Bot.* xvi. (1878) 239.—Chili.
angustifolia, *Sw. Prod. Veg. Ind. Occ.* 57.—Ind. occ.; Am. trop.
apicroides, *Schlecht. & Cham. in Linnaea*, vi. (1831) 55.—Mexic.
argentea, *Griseb. Cat. Pl. Cub.* 254.—Cuba.
argentea, C. Koch & J. Versch. in C. Koch, Wochenschr. xi. (1868) 163 = tectorum.
augusta, Vell. Fl. Flum. 135; iii. t. 135 = Aechmea augusta.
aurantiaca, *Griseb. in Goett. Nachr.* (1864) 16.—Am. trop.
axillaris, Griseb. Fl. Brit. W. Ind. 597 = complanata.
azurea, *Presl, Rel. Haenk.* i. 124.—Peruv.
Balbisiana, *Schult. f. Syst.* vii. II. 1212.—Ind. occ.
Bartrami, Nutt. in Am. Journ. Sc. Ser. I. v. (1822) 292 = utriculata.
Bartramii, Ell. Sketch, i. 379 = recurvata, setacea.
Benthamiana, *Klotzsch, ex Beer, Bromel.* 263, nomen.—Mexic.
Berteroniana, Schult. f. Syst. vii. II. 1221 = Catopsis nutans.
bicolor, Brongn. in Duperr. Voy. Coq. Bot. 185. t. 36 = stricta.
biflora, *Ruiz & Pav. Fl. Per.* iii. 41.—Peruv.
bivittata, Linden, Cat. n. 17 (1862) 7; et Regel, Gartenfl. xiv. (1865) 2 = Cryptanthus bivittatus.
brachycaulos, *Schlecht. in Linnaea*, xviii. (1844) 422.—Mexic.
bracteata, Chapm. Fl. S. U. St. 471 = fasciculata.
bracteata, Vell. Fl. Flum. 132; iii. t. 125 = Aechmea fasciata?
bracteosa, *Klotzsch, ex Beer, Bromel.* 263.—Ins. S. Doming.
breviscapa, A. Rich. Fl. Cub. Fanerog. ii. 265 = pruinosa.
bromeliaefolia, Rudge, Pl. Gui. 32. t. 50 = Aechmea bromeliaefolia.
bryoides, *Griseb. in Goett. Abh.* xxiv. (1879) 334.—Reg. Argent.
bulbosa, *Hook. Exot. Fl.* t. 173.—Am. trop.
caerulea, *H. B. & K. Nov. Gen. et Sp.* i. 291.—Mexic.

TILLANDSIA:—
caespitosa, Gill. ex Baker, in Journ. Bot. xvi. (1878) 238 = retorta.
caespitosa, Le Conte, in Ann. Lyc. N. York, ii. (1828) 131 = setacea.
caespitosa, *Schlecht. & Cham. in Linnaea*, vi. (1831) 54.—Mexic.
calamifolia, Salisb. in Trans. Hort. Soc. i. (1812) 270 = setacea.
calothyrsus, *Poepp. ex Beer, Bromel.* 263, nomen.—Peruv.
canescens, *Sw. Prod. Veg. Ind. Occ.* 57.—Jamaic.; Mexic.
capillaris, *Ruiz & Pav. Fl. Per.* iii. 42.—Peruv.
capitata, *Griseb. Cat. Pl. Cub.* 255.—Cuba.
capituligera, *Griseb. l. c.* 254.—Cuba.
Caput-Medusae, *E. Morr. in Belg. Hortic.* xxx. (1880) 90.—Mexic.
Caraguata, D. Dietr. Syn. Pl. ii. 1059 = Caraguata Berteroniana.
cinerascens, Willd. ex Schult. f. Syst. vii. II. 1218 = disticha.
circinalis, Griseb. in Goett. Abh. xix. (1874) 272 = Duratii.
circinnata, Schlecht. in Linnaea, xviii. (1844) 430 = streptophylla.
citrina, *Burch. ex Baker, in Journ. Bot.* xvii. (1879) 235.—Bras.
clavata, D. Dietr. Syn. Pl. ii. 1060 = Caraguata lingulata.
clavata, Lam. Encyc. i. 617 = Guzmania tricolor.
coarctata, Gill. ex Baker, in Journ. Bot. xvi. (1878) 236 = bryoides.
coarctata, Willd. ex Schult. f. Syst. vii. II. 1217 = floribunda.
coerulea, Auct. = caerulea.
comata, *Vell. Fl. Flum.* 136; iii. t. 140.—Bras.
Commelyna, E. Morr. in Belg. Hortic. xx. (1870) 97 = Hamaleana.
comosa, Griseb. Fl. Brit. W. Ind. 598, 775 (Index) = Guzmania tricolor.
compacta, *Griseb. in Goett. Nachr.* (1864) 18.—Venezuela.
complanata, *Benth. Bot. Voy. Sulph.* 173.—N. Granat.
complanata, E. Morr. in Belg. Hortic. xxii. (1872) 300. t. 23 = xiphostachys.
compressa, *Bert. ex Schult. f. Syst.* vii. II. 1210.—Jamaic.
compressa, Gill. ex Baker, in Journ. Bot. xvi. (1878) 240 = Gilliesii.
concentrica, Vell. Fl. Flum. 134; iii. t. 133 (Nidularii sp.).—Bras.
conspersa, *Miq. in Linnaea*, xviii. (1844) 376.—Guian.
corallina, *C. Koch, Ind. Sem. Hort. Berol.* (1873) *App.* iv. 5.—Bras.
cordobensis, Hieron. Ic. Pl. Argent. 10 = recurvata.
crinita, Willd. ex Beer, Bromel. 152 = usneoides.
cyanea, *E. Morr. in Belg. Hortic.* xxix. (1879) 297.—Guatemala.
cyathiformis, Vell. Fl. Flum. 137; iii. t. 144 (Canistri sp.).—Bras.
decumbens, Forsk. Fl. Aegypt. Arab. 72 = Cyanotis papilionacea.
Deppeana, *Steud. Nom.* ed. II. ii. 688.—Mexic.
Devillea, Steud. l. c. = Caraguata Berteroniana.
dianthoidea, *Rossi, Cat. Pl. Hort. Modoet.* 79. t. 1.—Bras.
discolor, Hort. ex Antoine, Bromel. 43 = Karatas purpurea.
dissitiflora, *Wright, in Sauv. Fl. Cub.* 168.—Cuba.
distachia, Vell. Fl. Flum. 136; iii. t. 141 (Bilbergiae sp.).—Bras.
distachya, *Baker, in Gard. Chron.* (1880) I. 200.—Honduras.
disticha, *H. B. & K. Nov. Gen. et Sp.* i. 292.—Peruv.
disticha, Willd. ex Schult. f. Syst. vii. II. 1226 = heliconioides.
divaricata, *Benth. Bot. Voy. Sulph.* 174.—Ecuador.
Duratii, *Vis. in Nuovi Sagg. Padov.* v. (1841) 271.—Am. trop.
Ehrenbergiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 320.—Mexic.
Ehrenbergii, *Klotzsch, ex Beer, Bromel.* 264, nomen.—Mexic.
elongata, *H. B. & K. Nov. Gen. et Sp.* i. 293.—N. Granat.

TILLANDSIA:—
eminens, Lindl. & Paxt. Flow. Gard. i. (1850–51) 160 = bulbosa.
ensiformis, *Vell. Fl. Flum.* 133; iii. t. 129.—Bras.
erecta, *Gill. ex Baker, in Journ. Bot.* xvi. (1878) 239.—Chili.
erubescens, *Schlecht. in Linnaea*, xviii. (1844) 427.—Mexic.
erubescens, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxii. (1854) 153 = ionantha.
erythraea, Lindl. & Paxt. Flow. Gard. i. (1850–51) 160 = bulbosa.
excelsa, Griseb. Cat. Pl. Cub. 254 = dissitiflora.
excelsa, *Griseb. Fl. Brit. W. Ind.* 597.—Ind. occ.
exsudans, Desf. Cat. Hort. Par. ed. III. 50 = Aechmea exsudans.
farinosa, Hort. ex Schult. f. Syst. vii. II. 1229, nomen (Quid?).—Hab.?
fasciculata, *Sw. Prod. Veg. Ind. Occ.* 56.—Am. bor.; Ind. occ.
Fendleri, *Griseb. in Goett. Nachr.* (1864) 17.—Venezuela.
fenestralis, *Hort.—Cf. Gard. Chron.* (1878) I. 440.—Hab.?
filifolia, *Schlecht. & Cham. in Linnaea*, vi. (1831) 53.—Mexic.
filiformis, Lodd. ex Schult. f. Syst. vii. 1229 = usneoides?
flavescens, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 118 = vestita.
flexuosa, *Sw. Prod. Veg. Ind. Occ.* 56.—Am. trop.
floribunda, Durat. ex Vis. in Nuovi Sagg. Padov. v. (1841) 271 = Duratii.
floribunda, *H. B. & K. Nov. Gen. et Sp.* i. 292.—Peruv.
foliosa, *Griseb. Fl. Brit. W. Ind.* 597.—Am. trop.
foliosa, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 119.—Mexic.
fusca, *Baker, in Journ. Bot.* xvi. (1878) 240.—Peruv.
Gardneri, *Lindl. Bot. Reg.* (1842), sub t. 63.—Bras.
geminiflora, *Brongn. in Duperr. Voy. Coq. Bot.* 186.—Bras.
gigantea, Griseb. Fl. Brit. W. Ind. 597 = T. Regina.
gigantea, *Mart. ex Schult. f. Syst.* vii. II. 1224.—Bras.
gigantea, Ruchinger, Cat. (1876) = Duratii.
Gilliesii, *Baker, in Journ. Bot.* xvi. (1878) 240.—Chili.
gladioliflora, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 31.—Am. centr.
globosa, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 222.—Bras.
glutinosa, *Mart. ex Schult. f. Syst.* vii. II. 1225.—Bras.
gracilis, Griseb. in Goett. Nachr. (1864) 17 = procera.
gracilis, Hort. ex Sweet, Hort. Brit. ed. II. 536, nomen (Quid?).—Ind. occ.
grandis, *Schlecht. in Linnaea*, xviii. (1844) 424.—Mexic.
Hamaleana, *E. Morr. in Belg. Hortic.* xx. (1870) 97.—Ecuador.
haplostachya, *Wright, in Sauv. Fl. Cub.* 169.—Cuba.
havanensis, Beer, Bromel. 85 = fasciculata.
heliconioides, *H. B. & K. Nov. Gen. et Sp.* i. 293.—N. Granat.
heptandra, Poir. Encyc. vii. 671, sphalm. = seq.
heptantha, *Ruiz & Pav. Fl. Per.* iii. 41.—Peruv.
heterophylla, E. Morr. in Belg. Hortic. xxiii. (1873) 138 = virginalis.
hieroglyphica, *Bull, ex E. Morr. in Belg. Hortic.* xxxv. (1885) 57, in syn.—Bras.
humilis, *Presl, Rel. Haenk.* i. 125.—Chili.
imbricata, *Vell. Fl. Flum.* 134; iii. t. 131.—Bras.
inanis, Lindl. & Paxt. Flow. Gard. i. (1850–51) 159 = bulbosa.
incana, Gill. ex Baker, in Journ. Bot. xvi. (1878) 238 = virescens.
incana, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 223 = Gardneri.
incarnata, *H. B. & K. Nov. Gen. et Sp.* i. 291.—Am. trop.
incurva, *Griseb. in Goett. Nachr.* (1864) 15.—Venezuela.
incurvata, Wright, in Sauv. Fl. Cub. 169 = Deppeana.
ionantha, *Planch. in Fl. des Serres*, x. (1854–55) 101. t. 1006.—Mexic.
ixioides, *Griseb. in Goett. Abh.* xxiv. (1879) 333.—Reg. Argent.

TILLANDSIA:—
Jonghei, *C. Koch, Wochenschr.* xi. (1868) 91.—Bras.
juncea, Poir. Encyc. Suppl. v. 309; Le Conte, in Ann. Lyc. N. York, ii. (1828) 130 = setacea.
juncea, Willd. ex Steud. Nom. ed. II. ii. 688 = Agave geminiflora.
juncifolia, Regel, Gartenfl. (1874) 321 = setacea.
Karwinskyana, *Schult. f. Syst.* vii. II. 1209.—Mexic.
Kunthiana, *Gaudich. Bot. Voy. Bonite*, t. 53.—Mexic.
laevis, Vell. Fl. Flum. 133; iii. t. 126 = Pitcairnia australis?
Landbecki, Phil. in Linnaea, xxxiii. (1864–65) 248 = recurvata.
lanuginosa, Gill. ex Baker, in Journ. Bot. xvi. (1878) 237 = propinqua.
latifolia, *Meyen, Reise*, ii. 45.—Peruv.
laxa, Griseb. Fl. Brit. W. Ind. 596 = Valenzuelana.
laxa, *Griseb. in Goett. Nachr.* (1864) 18.—Ind. occ.; Venezuela.
Leiboldiana, *Schlecht. in Linnaea*, xviii. (1844) 414.—Mexic.
Lescaillei, *Wright, in Sauv. Fl. Cub.* 167.—Cuba.
ligulata, Schult. f. Syst. vii. II. 1220 = utriculata.
limbata, Schlecht. in Linnaea, xviii. (1844) 419 = flexuosa.
Lindeni, *Regel, Ind. Sem. Hort. Petrop.* 1868 (1869) 92; *E. Morr. in Belg. Hortic.* xix. (1869) 321, cum ic.—Peruv.
Lindeniana, Regel, Gartenfl. (1869) 193 = praec.
linearis, *Vell. Fl. Flum.* 133; iii. t. 128.—Bras.
lingulata, Bartr. Trav. ed. II. (1794) 59 = utriculata.
lingulata, Linn. Sp. Pl. 286 = Caraguata lingulata.
loliacea, *Mart. ex Schult. f. Syst.* vii. II. 1204.—Bras.
longibracteata, Meyen, Reise, ii. 45 = purpurea.
Lorentziana, *Griseb. in Goett. Abh.* xix. (1874) 271.—Reg. Argent.
lorifolia, *Lem. in Fl. des Serres*, ii. (Mai, 1846) sub t. 4.—Bras.
macrocnemis, Griseb. in Goett. Abh. xxiv. (1879) 332 = xiphioides.
macrostachya, Klotzsch, ex Beer, Bromel. 264, nomen = polystachya.
maculata, *Ruiz & Pav. Fl. Per.* iii. 40.—Peruv.
Malzinei, *Baker, in Kew Gard. Rep.* (1878) 59.—Mexic.
monostacha, Linn. Sp. Pl. ed. II. 410 = Guzmania tricolor.
monostachya, Bartr. Trav. ed. II. (1794) 59 = recurvata, setacea, utriculata.
monostachys, Gill. ex Baker, in Journ. Bot. xvi. (1878) 239 = recurvata.
Morelliana, Hort. ex Henfrey, in Moore & Ayres, Gard. Mag. Bot. iii. (1851) 33 = Billbergia vittata.
Moritziana, *Klotzsch, ex Beer, Bromel.* 265, nomen.—Venezuela.
Morreniana, Regel, Gartenfl. (1870) 41 = Lindeni.
mosaica, Hort.—Cf. Gard. Chron. (1874) II. 49 = *musaica*.
mucronata, *Griseb. in Goett. Nachr.* (1864) 20.—Venezuela.
multicaulis, Steud. Nom. ed. II. ii. 688 = caespitosa.
multiflora, *Benth. Bot. Voy. Sulph.* 174.—Ecuador.
musaica, Linden & André, in Illustr. Hortic. xx. (1873) 171 = Caraguata musaica.
myosura, *Griseb. in Goett. Abh.* xxiv. (1879) 333.—Reg. Argent.
Nappii, *Lor. & Niederl. in Roca, Exped. Rio-Negro*, ii. 282.—Patagon.
narthecioides, *Presl, Rel. Haenk.* i. 125.—Ecuador.
nitida, Hook. Exot. Fl. t. 218 = Catopsis nitida.
nutans, Sw. Prod. Veg. Ind. Occ. 56 = Catopsis nutans.
Nuttalliana, Schult. f. Syst. vii. II. 1220 = utriculata.
obscura, Lodd. ex G. Don, in Loud. Hort. Brit. 119, nomen (Quid?).—Am. austr.
Osyana, *Hort. Belg. ex C. Koch, in Belg. Hortic.* xii. (1862) 127.—Hab.?
paleacea, *Presl, Rel. Haenk.* i. 125.—Chili.
paniculata, *Linn. Sp. Pl.* ed. II. 410.—Ins. S. Doming.
paniculata, Schlecht. & Cham. in Linnaea, vi. (1831) 54 = Deppeana.
parviflora, *Ruiz & Pav. Fl. Per.* iii. 41.—Peruv.
patens, Willd. ex Schult. f. Syst. vii. II. 1229 = flexuosa.
paucifolia, Baker, in Gard. Chron. (1878) II. 748 = bulbosa.
pendula, *Thunb. ex Gaertn. f. Fruct.* iii. 13.—Hab.?
penduliflora, *Griseb. Fl. Brit. W. Ind.* 597.—Ins. Dominic.

TILLANDSIA:—
pinifolia, Le Conte, in Ann. Lyc. N. York, ii. (1828) 131 = setacea.
pitiphylla, Mart. ex Schult. f. Syst. vii. II. 1208 = pulchra.
platynema, *Griseb. in Goett. Nachr.* (1864) 19.—Am. trop.
pleiosticha, *Griseb. l. c.*—Venezuela.
polystachia, *Linn. Sp. Pl.* ed. II. 410.—Ind. occ.
polystachia, Vell. Fl. Flum. 136; iii. t. 138 = Aechmea brasiliensis.
polystachya, Jacq. ex Griseb. Fl. Brit. W. Ind. 595 = fasciculata.
polystachya, Muhl. Cat. ed. I. 34 = utriculata.
polytrichoides, E. Morr. in Belg. Hortic. xxx. (1880) 240 = bryoides.
procera, *Mart. ex Schult. f. Syst.* vii. II. 1224.—Bras.
propinqua, *C. Gay, Fl. Chil.* vi. 15.—Chili.
pruinosa, *Sw. Fl. Ind. Occ.* i. 594.—Ind. occ.; Venezuela.
Pseudo-ananas, Blanco, Fl. Filip. ed. I. 853 = Freycinetia insignis.
psittacina, *Hook. Bot. Mag.* t. 2841.—Bras.
pulchella, Hook. Exot. Fl. ii. t. 154 = seq.
pulchra, *Hook. l. c.* t. 154, in textu.—Am. trop.
pumila, *Griseb. in Goett. Nachr.* (1864) 16.—Venezuela.
pumila, Lindl. & Paxt. Flow. Gard. i. (1850–51) 160 = bulbosa.
punctulata, *Schlecht. & Cham. in Linnaea*, vi. (1831) 53.—Mexic.
purpurea, *Ruiz & Pav. Fl. Per.* iii. 41.—Peruv.
purpurea, Spreng. Neue Entdeck. i. 251 = Sprengeliana.
pusilla, *Gill. ex Baker, in Journ. Bot.* xvi. (1878) 237.—Chili.
quadrangularis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 119.—Mexic.
ramealis, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1121, nomen.—Guian.
ramosa, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 121.—Porto Rico.
ramosa, Sweet, Hort. Brit. ed. I. 425 = utriculata.
rectangula, *Baker, in Journ. Bot.* xvi. (1878) 238.—Reg. Argent.
recurvata, *Linn. Sp. Pl.* ed. II. 410.—Am. trop.
recurvifolia, Hook. Bot. Mag. t. 5246 = dianthoidea.
Regina, *Vell. Fl. Flum.* 136; iii. t. 142.—Am. trop.
retorta, *Griseb. in Goett. Abh.* xxiv. (1879) 334.—Reg. Argent.
rigida, Gill. ex Baker, in Journ. Bot. xvi. (1878) 239 = erecta.
rigida, Hort. ex Sweet, Hort. Brit. ed. I. 425, nomen (Quid ?).—Ind. occ.
ringens, *Griseb. Cat. Pl. Cub.* 255.—Cuba.
robusta, *Griseb. in Goett. Nachr.* (1864) 15.—Venezuela.
Roezli, *E. Morr. in Belg. Hortic.* xxvii. (1877) 272. t. 15.—Peruv.
rosea, Lindl. Bot. Reg. t. 1357 = dianthoidea.
rubida, Lindl. l. c. (1842) t. 63 = geminiflora.
rubra, Hort. ex Antoine, Bromel. 43 = Karatas purpurea.
rubra, *Ruiz & Pav. Fl. Per.* iii. 40.—Peruv.
Saundersii, *C. Koch, Ind. Sem. Hort. Berol.* (1873) *App.* iv. 6.—Bras.
saxatilis, *Vell. Fl. Flum.* 136; iii. t. 139.—Bras.
Schiedeana, Steud. Nom. ed. II. ii. 688 = vestita.
Schuchii, Beer & Fenzl, in Allg. Gartenz. xiv. (1846) 266 = Pitcairnia albiflos.
scoparia, Willd. ex Schult. f. Syst. vii. II. 1217 = straminea.
secunda, *H. B. & K. Nov. Gen. et Sp.* i. 294.—Ecuador.
Selloa, C. Koch, Ind. Sem. Hort. Berol. (1873) App. iv. 7 = setacea.
serrata, *Linn. Sp. Pl.* 286 (sp. dub.).—Ind. occ.
sessiliflora, Ruiz & Pav. Fl. Per. iii. 42 (Catopsidis sp.).—Am. trop.
setacea, Hook. Bot. Mag. t. 3275 = compressa.
setacea, Sw. Fl. Ind. Occ. i. 593.—Am. trop.
simplex, *Vell. Fl. Flum.* 133; iii. t. 130.—Bras.
sordida, Salisb. Prod. 247 = Caraguata lingulata.
spiculosa, *Griseb. in Goett. Nachr.* (1864) 17.—Venezuela.
spiculosa, Schlecht. ex E. Morr. in Belg. Hortic. xxix. (1879) 57 (Quid ?).—Mexic.
splendens, Brongn. in Fl. des Serres, ii. (Mai, 1846) t. 4.—Guian.
Sprengeliana, *Klotzsch, ex Beer, Bromel.* 266, nomen.—Bras.

TILLANDSIA:—
squamulosa, Willd. ex Schult. f. Syst. vii. II. 1209 = caerulea.
staticeflora, E. Morr. in Belg. Hortic. xxi. (1871) 177. t. 12 = filifolia.
straminea, *H. B. & K. Nov. Gen. et Sp.* i. 292.—Peruv.
streptophylla, *Scheidw. in Hortic. Belge*, iii. (1836) 252.—Mexic.
striata, Willd. ex Schult. f. Syst. vii. II. 1209 = incarnata.
stricta, Lindl. Bot. Reg. t. 1338 = dianthoidea.
stricta, *Soland. ex Sims, Bot. Mag.* t. 1529.—Bras.
strobilantha, *Poir. Encyc. Suppl.* v. 309.—Peruv.
suaveolens, *Lem. in Fl. des Serres*, ii. (Mai, 1846) sub t. 4.—Reg. Argent.
subulata, Vell. Fl. Flum. 132; iii. t. 127 = stricta.
tectorum, *E. Morr. in Belg. Hortic.* xxvii. (1877) 328. t. 18.—Peruv.
tenuifolia, Bert. ex Schult. f. Syst. vii. II. 1212 = Balbisiana.
tenuifolia, Jacq. Select. Am. 92. t. 63 = flexuosa.
tenuifolia, Linn. Sp. Pl. 286, partim = setacea.
terminalis, Vell. Fl. Flum. 137; iii. t. 143 (Nidularii sp.).—Bras.
tessellata, *Linden, Cat.* (1873) 9; *Illustr. Hortic.* (1873) 78; (1874) 123. t. 179.—Bras.
tetrantha, *Ruiz & Pav. Fl. Per.* iii. 39.—Peruv.
tetrastachia, *Vell. Fl. Flum.* 135; iii. t. 137.—Bras.
tortilis, Klotzsch, ex Beer, Bromel. 266 = pruinosa.
tortuosa, Lodd. ex Schult. f. Syst. vii. II. 1229, nomen (Quid ?).—Hab. ?
trichoides, H. B. & K. Nov. Gen. et Sp. i. 290 = usneoides.
tricholepis, *Baker, in Journ. Bot.* xvi. (1878) 237.—Bolivia.
tricolor, Schlecht. & Cham. in Linnaea, vi. (1831) 54 = punctulata.
triflora, *Vell. Fl. Flum.* 134; iii. t. 134.—Bras.
triglochinoides, *Presl, Rel. Haenk.* i. 125.—Ecuador.
tuberosa, Vell. Fl. Flum. 135; iii. t. 136 = Dyckia rariflora.
Unca, *Griseb. in Goett. Abh.* xix. (1874) 273.—Reg. Argent.
undulata, *Baker, in Journ. Bot.* xvi. (1878) 240.—Paraguay.
uniflora, H. B. & K. Nov. Gen. et Sp. i. 290 = recurvata.
unispicata, Vell. Fl. Flum. 132; iii. t. 124 = Aechmea nudicaulis.
usneoides, *Linn. Sp. Pl.* ed. II. 411.—Am. bor. et austr.
utriculata, *Linn. l. c.* ed. I. 286.—Ind. occ.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 267.—Cuba.
variabilis, *Schlecht. in Linnaea*, xviii. (1844) 418.—Mexic.
variegata, *Schlecht. l. c.* 429.—Mexic.
variegata, *Vell. Fl. Flum.* 134; iii. t. 132.—Bras.
ventricosa, *Griseb. in Goett. Nachr.* (1864) 19.—Venezuela.
ventricosa, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 222.—Bras.
vestita, Benth. Pl. Hartw. 25 = Benthamiana.
vestita, *Schlecht. & Cham. in Linnaea*, vi. (1831) 52.—Mexic.
vestita, Willd. ex Schult. f. Syst. vii. II. 1228 = Aechmea bromeliaefolia.
viminalis, *Hemsl. Biol. Centr. Am. Bot.* iii. 323.—Costa Rica.
violacea, *Klotzsch, ex Beer, Bromel.* 266.—Venezuela.
virescens, *Ruiz & Pav. Fl. Per.* iii. 43.—Peruv.
virginalis, *E. Morr. in Belg. Hortic.* xxx. (1880) 238.—Mexic.
viridiflora, Hort. ex Antoine, Bromel. 21 = viminalis.
vitellina, *Link, Klotzsch & Otto, in Otto & Dietr. Allg. Gartenz.* ix. (1841) 81.—Venezuela.
vittata, *Linden, ex Beer, Bromel.* 43.—Hab. ?
xiphioides, *Ker-Gawl. in Bot. Reg.* t. 105.—Am. trop.
xiphostachys, *Griseb. in Goett. Nachr.* (1864) 14.—Mexic.
Zahnii, Hort. Veitch, ex Hook. f. Bot. Mag. sub t. 6059 = Caraguata Zahnii.

TILLEA, Sanguin. in Att. Acc. Pont. Lincei, Ser. I. v. (1852) 506, sphalm. = **Tillaea**, Mich. (Crassul.).

TILLIA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 173 = praec.

TIMAEOSIA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 138. t. 33. (1862) = **Gypsophila**, Linn. (Caryoph.).
cerastioides, Klotzsch, l. c. 138 = G. cerastioides.
rupestris, Edgew. & Hook. f. in Fl. Brit. Ind. i. 217, sphalm. = praec.

TIMANDRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 197 = **Croton**, Linn. (Euphorb.).
dichotoma, Klotzsch, l. c. = C. dichotomus.
erythroxyloides, Klotzsch, l. c. = C. erythroxyloides.
serrata, Klotzsch, l. c. = C. serratus.

TIMANTHEA, Salisb. Prod. 208 (1796) = **Baltimora**, Linn. (Compos.).
tristis, Salisb. l. c. = B. recta.

TIMBALIA, Clos, in Bull. Soc. Bot Fr. xviii. (1871) 178 = **Crataegus**, Tourn. (Rosac.).
Pyracantha, Clos, l. c. = C. Pyracantha.

TIMBULETA, Steud. Nom. ed. II. ii. 688 (1841), err. typ. = SIMBULETA, Forsk. = **Anarrhinum**, Desf. (Scrophul.).

TIMEROYA, Benth. & Hook. f. Gen. iii. 10 (1880), sphalm. = seq.

TIMEROYEA, Montr. in Mém. Acad. Lyon, x. (1860) 247. *NYCTAGINEAE*, Benth. & Hook. f. iii. 10.
VIEILLARDIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. viii. (1861) 375.
artensis, *Montr. in Mém. Acad. Lyon*, x. (1860) 247.—N. Caled.

TIMMIA, J. F. Gmel. Syst. 524, 538 (1791) = **Cyrtanthus**, Ait. (Amaryll.).
angustifolia, J. F. Gmel. l. c. = C. angustifolius.
obliqua, J. F. Gmel. l. c. = C. obliquus.

TIMONIUS, Rumph. Amboin. iii. 216. t. 140 (1743); DC. Prod. iv. 461 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 102.
BURNEYA, Cham. & Schlecht. in Linnaea, iv. (1829) 189, partim.
ERITHALIS, Forst. f. Prod. 17 (1786).
EUPYRENA, Wight & Arn. Prod. 422 (1834).
HELOSPORA, Jack, in Trans. Linn. Soc. xiv. (1823) 127. t. 4. f. 3.
NELITRIS, Gaertn. Fruct. i. 134. t. 27 (1788).
POLYPHRAGMON, Desf. in Mém. Mus. Par. vi. (1820) 5. t. 2.
POROCARPUS, Gaertn. Fruct. ii. 473. t. 178 (1791).
PYROSTRIA, Roxb. Hort. Beng. 83 (1814); Fl. Ind. i. 388 (1832).
acuminatus, Wall. Cat. n. 6218 = Urophyllum Blumeanum.
affinis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 36.—Ins. Fiji.
Finlaysonianus, *Hook. f. Fl. Brit. Ind.* iii. 127.—Singapore.
flavescens, *Baker, Fl. Maurit.* 144.—Ins. Seychelles.
Forsteri, *DC. Prod.* iv. 461.—Ins. Societ.
fuscus, Walp. Ann. ii. 765 = Bobea fusca (in Addend.).
Gaudichaudii, *DC. Prod.* iv. 461.—Ins. Sandvic.
Jambosella, Thw. Enum. Pl. Zeyl. 153 = Koenigii.
Koenigii, *Blume, ex Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 211.—Malaya.
mutabilis, *Walp. Ann.* ii. 765.—Borneo.
platycarpus, *Montr. in Mém. Acad. Lyon*, x. (1860) 221.—N. Caled.
putaminosus, F. Muell. Fragm. iv. 92 (= *Guettardella putaminosa*).—Austral.
Rumphii, *DC. Prod.* iv. 461.—Malaya.
sapotaefolius, *A. Gray, in Proc. Am. Acad.* iv. (1860) 35.—Ins. Fiji.

TINA, Blume, Bijdr. 235 (1825) = **Harpullia**, Roxb. (Sapind.).
rupestris, Blume, l. c. = H. cupanioides.

TINA, Roem. & Schult. Syst. v. p. xxxii. 414 (1819) = **Ratonia**, DC. (Sapindac.).
cupanioides, DC. Prod. i. 614 = Cupania laevis, *T. Gelonium*.
dasycarpa, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 663.—Madag.
fulvinervis, Radlk. l. c. 662.—Madag.
Gelonium, Roem. & Schult. Syst. v. 414.—Madag.

TINA:—
isoneura, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 663.—Madag.
madagascariensis, DC. Prod. i. 614 = *T. Gelonium.*
madagascariensis, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 662 = Garuga madagascariensis.
polyphylla, Baker, in Journ. Linn. Soc. xxi. (1884) 335.—Madag.
striata, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 663.—Madag.
trijuga, Radlk. l. c. 662.—Madag.

TINAEA, Boiss. Fl. Orient. v. 58 (1884), sphalm. = TINEA, Bivona = **Habenaria**, Willd. (Orchid.).

TINAEA, Garzia, in Rel. Acc. Zel. Aci-Reale, Ann. 3–4. 24; ex Parl. Fl. Palerm. i. 138 (1845) = **Lamarkia**, Moench (Gramin.).
elegans, Garzia, ex Parl. l. c. 139 = L. aurea.

TINANTIA, Mart. & Gal. in Bull. Acad. Brux. xi. 1. (1844) 240 = **Senkenbergia**, Schau. (Nyctag.).
gypsophiloides, Mart. & Gal. l. c. = S. annulata.

TINANTIA, Schiedw. in Otto & Dietr. Allg. Gartenz. vii. (1839) 365. *COMMELINACEAE*, Benth. & Hook. f. iii. 853.
POGOMESIA, Rafin. Fl. Tellur. iii. 67 (1836), nomen prius.
anomala, *C. B. Clarke, in DC. Monog. Phan.* iii. 287.—Mexic.
erecta, Schlecht. in Linnaea, xxv. (1852) 185 = fugax.
fugax, *Scheidw. in Otto & Dietr. Allg. Gartenz.* vii. (1839) 365.—Am. trop.
latifolia, Schlecht. in Linnaea, xxv. (1852) 184 = fugax.
lineolata, Hassk. Pl. Jav. Rar. 98 = Aneilema herbaceum.
Sprucei, *C. B. Clarke, in DC. Monog. Phan.* iii. 287.—Bras.
undata, Schlecht. in Linnaea, xxv. (1852) 185 = fugax.

TINEA, Bivona, in Giorn. Sc. Sicil. (1833) 149 = **Habenaria**, Willd. (Orchid.).
cylindrica, Bivona, l. c. = H. intacta.
intacta, Boiss. Fl. Orient. v. 58 = H. intacta.

TINEA, Spreng. Neue Entdeck. ii. 165 (1821) = **Prockia**, Linn. (Tiliac.).
triplinervia, Spreng. l. c. = P. Crucis.

TINGUARRA, Parl. in Webb & Berth. Phyt. Canar. ii. 156. t. 71 (1836–50). *UMBELLIFERAE*, Benth. & Hook. f. i. 896.
cerviariaefolia, *Parl. l. c.* 157.—Ins. Canar.
montana, *Benth. & Hook. f. Gen.* i. 897.—Ins. Canar.
sicula, *Benth. & Hook. f. l. c.*—Sicil.

TINIARIA, Reichb. Handb. 236 (1837) = **Polygonum**, Tourn.
alata, Montand. Guide Bot. 270 = P. dumetorum.
carinata, Montand. l. c. = P. Convolvulus.
Convolvulus, Webb & Moq. in Webb & Berth. Phyt. Canar. iii. 221 = P. Convolvulus.
dumetorum, Opiz, Seznam, 98 = P. dumetorum.

TINNEA, Kotschy & Peyr. Pl. Tinn. 25. t. 11 (1867). *LABIATAE*, Benth. & Hook. f. ii. 1220.
RENSCHIA, Vatke, in Linnaea, xliii. (1880–82) 94.
aethiopica, *Kotschy & Peyr. Pl. Tinn.* 25. t. 11.—Afr. trop.
antiscorbutica, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 58.—Afr. trop.
erianthera, *Vatke, in Linnaea*, xliii. (1880–82) 539.—Afr. trop.
eriocalyx, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 59.—Afr. trop.
heterotypica, *S. Moore, in Journ. Bot.* xv. (1877) 69.—Afr. trop.

TINNETHAMNUS, Pritz. Ic. Ind. ii. 273 (1865) = praec.

TINNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Quid?).

TINOMISCIUM, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44. *MENISPERMACEAE*, Benth. & Hook. f. i. 34.
coriaceum, Miers, l. c. = petiolare.
elasticum, *Becc. Malesia*, i. (1877) 141.—N. Guin.
javanicum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiii. (1864) 490.—Java.
petiolare, *Miers, l. c.* Ser. II. vii. (1851) 44.—Malaya.
phytocrenoides, *Kurz, ex Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 36.—Sumatra.
pyrrhobotryum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 81.—Java.

TINOSPORA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 38. *MENISPERMACEAE*, Benth. & Hook. f. i. 34.
arfakiana, *Becc. Malesia*, i. (1877) 140.—N. Guin.
Bakis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 38; *et* Ser. III. xiii. (1864) 322.—Afr. trop.
convolvulacea, Miers, l. c. Ser. II. vii. (1851) 38 = cordifolia.
cordifolia, *Miers, l. c.*—Ind. or.
crispa, *Miers, l. c.*—Java.
flavescens, Miers, l. c. = Cocculus flavescens.
glauca, Miers, l. c. = Cocculus glaucus.
Hullsii, *F. Muell. Fragm.* v. 147.—Austral.
lacunosus, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 38 = Anamirta paniculata.
malabarica, *Miers, l. c.*—Ind. or.
nudiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 392.—Ind. or.
palminervis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiii. (1864) 317.—Burma.
reticulata, *Miers, l. c.* 321.—Ins. Philipp.
smilacina, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 52.—Austral.
sumatrana, *Becc. Malesia*, i. (1877) 139.—Sumatra.
tenera, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiii. (1864) 322.—Afr. trop.
tomentosa, *Miers, l. c.* Ser. II. vii. (1851) 38.—Ind. or.; Burma.
uliginosa, *Miers, l. c.* Ser. III. xiii. (1864) 321.—Malaya.
Walcottii, *F. Muell. ex Benth. Fl. Austral.* i. 56.—Austral.

TINUS, Linn. Syst. ed. X. 1010 (1759) = **Clethra**, Gronov. (Ericac.).
occidentalis, Linn. l. c. = C. tinifolia.
Volckameria, Crantz, Inst. ii. 376 = C. tinifolia.

TINUS, [Tourn.] Linn. Syst. ed. I (1735) = **Viburnum**, Linn. (Caprif.).
laurifolius, Borkh. in Roem. Arch. i. ii. (1797) 20 = V. Tinus.
rugosus, Spach, Hist. Vég. Phan. viii. 317 = V. rigidum.

TIPALIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Zanthoxylum**, Linn. (Rutac.).
Limonella, Dennst. l. c. = Z. Rhetsa.

TIPHA, Neck. Delic. Gallo-Belg. ii. 376 (1768) = **Typha**, Linn.

TIPHOGETON, Ehrh. Beitr. iv. 146 (1789) = **Ludwigia**, Linn. (Onagrar.).

TIPUANA, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 72. *LEGUMINOSAE*, Benth. & Hook. f. i. 546.
heteroptera, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 72.—Bras.
macrocarpa, *Benth. l. c.*—Bras.
speciosa, *Benth. l. c.*—Am. austr.

TIPULARIA, Nutt. Gen. Am. ii. 195 (1818). *ORCHIDEAE*, Benth. & Hook. f. iii. 497.
ANTHERICLIS, Rafin. in Am. Monthly Mag. (1819) 195; et in Journ. Phys. lxxxix. (1819) 261.
PLECTRURUS, Rafin. Neogenyt. 4 (1825).
camtschatica, *Spreng. Syst.* iii. 734.—Kamtschat.
discolor, *Nutt. Gen. Am.* ii. 195.—Am. bor.
Josephi, *Reichb. f. ex Lindl. in Journ. Linn. Soc.* i. (1857) 174.—Reg. Himal.

TIQUILIA, Pers. Syn. i. 157 (1805) = **Coldenia**, Linn. (Borag.).
brevifolia, Nutt. ex Torr. in Bot. Mex. Bound. 136 = C. Nuttallii.
dichotoma, Pers. Syn. i. 157 = C. dichotoma.
oregana, Torr. in Bot. U. St. Expl. Exped. ii. 411. t. 12 = C. Nuttallii.
parvifolia, Nutt. ex Benth. in Hook. Kew Journ. iii. (1851) 296 = C. Nuttallii.

TIRASEKIA, G. Don, in Sweet, Hort. Brit. ed. III. 563 (1839), sphalm. = JIRASEKIA, F. W. Schmidt = **Anagallis**, Tourn. (Primul.).

TIRICTA, Rafin. New Fl. Am. iv. 22 (1836) = **Daucus**, Tourn. (Umbell.).
daucoides, Rafin. l. c. 23 = D. Carota.

TIRTALIA, Rafin. Fl. Tellur. iv. 71 (1836) = **Ipomoea**, Linn. (Convolv.).
anceps, Rafin. l. c. = I. Turpethum.
angustifolia, Rafin. l. c. = I. angustifolia, *Jacq.*
dissecta, Rafin. l. c. = I. sinuata.
domingensis, Rafin. l. c. = I. sidaefolia.
emarginata, Rafin. l. c.—Hab.?
filicaulis, Rafin. l. c. = I. angustifolia, *Jacq.*
gemella, Rafin. l. c. = I. obscura, *Ker-Gawl.*
hispida, Rafin. l. c. = I. eriocarpa.
macrorhiza, Rafin. l. c. = I. macrorrhiza.
malabarica, Rafin. l. c. = Argyreia malabarica.
maxima, Rafin. l. c. = I. sepiaria.
pandurata, Rafin. l. c. = I. fastigiata.
quinqueloba, Rafin. l. c. = I. palmata.
sibirica, Rafin. l. c. = I. sibirica.
striata, Rafin. l. c. = I. sepiaria.
triflora, Rafin. l. c. = I. sepiaria?
triqueter, Rafin. l. c. = I. Turpethum.
umbellata, Rafin. l. c. = I. umbellata, *Linn.*
vitifolia, Rafin. l. c. = I. vitifolia.

TIRUCALLA, Rafin. Fl. Tellur. iv. 112 (1836) = **Euphorbia**, Tourn.
indica, Rafin. l. c. = E. Tiruculli.

TISSA, Adans. Fam. ii. 507 (1763) = **Spergularia**, J. & C. Presl (1819) (Caryoph.).

TITA, Scop. Introd. 219 (1777) = **Cassipourea**, Aubl. (Rhizoph.).

TITANIA, Endl. Prod. Fl. Norf. 31 (1833) = **Oberonia**, Lindl. (Orchid.).
miniata, Endl. l. c. = O. Titania.

TITELBACHIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126, sphalm. = TITTELBACHIA, Klotzsch = **Begonia**, Tourn.

TITHONA, Linn. Syst. ed. I (1735) = **Rivina**, Plum. (Phytolac.).

TITHONIA, Desf. ex Juss. Gen. 189 (1789); et in Ann. Mus. Par. i. (1802) 49. t. 4. *COMPOSITAE*, Benth. & Hook. f. ii. 374.
angustifolia, Hook. & Arn. Bot. Beech. Voy. 435 = Viguiera buddleiaeformis.
argophylla, Eaton, in Bot. King's Exp. 423 = Helianthella argophylla.
aristata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 114.—Costa Rica.
calva, Sch. Bip. in Seem. Bot. Voy. Herald, 305 = Gymnolomia calva.
decurrens, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 85 = Viguiera decurrens.
diversifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 5.—Mexic.
excelsa, DC. Prod. v. 585 = Viguiera excelsa.
glutinosa, Collie, ex Hook. & Arn. Bot. Beech. Voy. 33 = Helianthus thurifer.
helianthoides, Weinm. ex Steud. Nom. ed. II. ii. 689 = tubaeformis.
heterophylla, *Griseb. in Bonplandia*, vi. (1858) 9.—Am. centr.
ovata, Hook. Bot. Mag. t. 3901 = Zexmenia ovata.
pachycephala, DC. Prod. v. 585 = Viguiera pachycephala.
platylepis, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 368 = Gymnolomia platylepis.

TITHONIA:—
pusilla, *A. Gray, in Proc. Am. Acad.* v. (1861) 124.—Peruv.
recurrens, Hemsl. Biol. Centr. Am. Bot. ii. 176, sphalm. = Viguiera decurrens.
scaberrima, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 91 = Gymnolomia platylepis.
scabra, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 375.—Hab.?
speciosa, *Hook. ex Griseb. Cat. Pl. Cub.* 155.—Mexic.
splendens, Hort. Rendatler, ex Planch. in Fl. des Serres, viii. (1852) 19 = Dysodia grandiflora.
tagetiflora, *Desf. in Ann. Mus. Par.* i. (1802) 46. t. 4.—Mexic.
Thurberi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 655.—Mexic.
tubaeformis, *Cass. in Dict. Sc. Nat.* liv. 456.—Mexic.
tubaeformis, A. Gray, in Bot. Mex. Bound. 90 = Thurberi.
uniflora, J. F. Gmel. Syst. 1259 = tagetiflora?

TITHONIA, Raeusch. Nom. ed. III. 251 (1797) = **Rudbeckia**, Linn. (Compos.).
laciniata, Raeusch. l. c. = R. laciniata.

TITHYMALOPSIS, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 249 = **Euphorbia**, Linn.
angustifolia, Klotzsch & Garcke, l. c. = E. discoidalis.
corollata, Klotzsch & Garcke, l. c. = E. corollata, *Linn.*
sphaerorrhiza, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 44 = E. sphaerorrhiza.

TITHYMALUS, Tourn. ex Hall. Enum. Stirp. Helv. i. 189 (1742) = **Euphorbia**, Linn.
acalyphoides, Schweinf. Beitr. Fl. Aethiop. 38 = E. acalyphoides.
acenocarpus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 79 = E. akenocarpa.
acicularis, Dulac, Fl. Hautes-Pyr. 156 = E. Cyparissias.
acutifolius, Lam. Fl. Fr. iii. 90 = E. Pithyusa.
Adrianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 73 = E. Adriana.
agovensis, Schweinf. Beitr. Fl. Aethiop. 38 = E. agowensis.
agrarius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 89 = E. agraria.
aleppicus, Klotzsch & Garcke, l. c. 84 = E. aleppica.
alpinus, Klotzsch & Garcke, l. c. 70 = E. alpina.
altaicus, Klotzsch & Garcke, l. c. 79 = E. altaica.
altissimus, Klotzsch & Garcke, l. c. 80 = E. altissima.
amplexicaulis, Klotzsch & Garcke, l. c. = E. condylocarpa.
amygdaloides, Hill, Hort. Kew. 172/4 = E. amygdaloides.
Anacampseros, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 86 = E. Anacampseros.
andrachnoides, Klotzsch & Garcke, l. c. 88 = E. subcordata.
angulatus, Rafin. Fl. Tellur. iv. 115; Klotzsch & Garcke, l. c. 77 = E. angulata.
angustifolius, Gilib. Fl. Lituan. ii. 209 = E. Cyparissias.
Antiquorum, Moench, Meth. 665 = E. Antiquorum.
aphyllus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 82 = E. aphylla.
apiculatus, Klotzsch & Garcke, l. c. 94 = E. epicyparissias.
Apios, Hill, Hort. Kew. 173/3 = E. Apios.
arboreus, Lam. Fl. Fr. iii. 94 = E. dendroides.
arkansanus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 66 = E. dictyosperma.
arvalis, Klotzsch & Garcke, l. c. 82 = E. arvalis.
asperus, Klotzsch & Garcke, l. c. 78 = E. aspera.
atropurpureus, Klotzsch & Garcke, l. c. 80 = E. atropurpurea.
atro-sanguineus, Klotzsch & Garcke, l. c. 70 = E. portulacoides.
attenuatus, Klotzsch & Garcke, l. c. 69 = E. elliptica.
Aucheri, Klotzsch & Garcke, l. c. 93 = E. Aucheri.
aulacospermus, Klotzsch & Garcke, l. c. 82 = E. aulacosperma.
auriculatus, Lam. Fl. Fr. iii. 102 = E. Peplis.
azoricus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 98 = E. Pinea.
balsamiferus, Haw. Syn. Pl. Succ. 140 = E. balsamifera.

TITHYMALUS:—
Bergii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 68 = Euphorbia elliptica.
biglandulosa, Haw. Syn. Pl. Succ. 141 = E. biglandulosa.
biumbellatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 85 = E. biumbellata.
blepharophyllus, Klotzsch & Garcke, l. c. 73 = E. blepharophylla.
brachypus, Klotzsch & Garcke, l. c. 74 = E. mauritanica.
bracteatus, Haw. Syn. Pl. Succ. 138 = Pedilanthus bracteatus.
Brauni, Schweinf. in Journ. Bot. i. (1863) 295 = E. longetuberculosa.
Bridgesii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 70 = E. portulacoides.
buchtormensis, Klotzsch & Garcke, l. c. 79 = E. buchtormensis.
bupleurifolius, Haw. Syn. Pl. Succ. 138 = E. bupleurifolia.
caesius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 87 = E. Esula.
Cajogalus, Moench, Meth. 667 = E. Gerardiana.
calendulaefolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 74 = E. angulata.
calyculatus, Klotzsch & Garcke, l. c. 81 = E. calyculata.
campester, Klotzsch & Garcke, l. c. 86 = E. campestris.
capensis, Klotzsch & Garcke, l. c. 98 = E. striata.
cardiophyllus, Klotzsch & Garcke, l. c. 78 = E. cardiophylla.
carniolicus, Rafin. Fl. Tellur. iv. 115; Klotzsch & Garcke, l. c. 77 = E. carniolica.
Cassia, Klotzsch & Garcke, l. c. 67 = E. Cassia.
cebrinus, Klotzsch & Garcke, l. c. 86 = E. cerebrina.
Chamaesyce, Moench, Meth. 666 = E. Chamaesyce.
Characias, Hill, Hort. Kew. 172/4 = E. Characias.
cheiradenius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 98 = E. cheiradenia.
Chesneyi, Klotzsch & Garcke, l. c. 99 = E. Chesneyi.
chilensis, Klotzsch & Garcke, l. c. 67 = E. portulacoides.
chrysophyllus, Klotzsch & Garcke, l. c. 72 = E. chrysophylla.
cinerascens, Moench, Meth. 668 = E. segetalis.
Clementei, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 76 = E. Clementei.
cognatus, Klotzsch & Garcke, l. c. 66 = E. pilosa.
collinus, Klotzsch & Garcke, l. c. 93 = E. collina.
commutatus, Klotzsch & Garcke, l. c. 82 = E. commutata.
condylocarpus, Klotzsch & Garcke, l. c. 78 = E. condylocarpa.
confertus, Klotzsch & Garcke, l. c. 94 = E. ericoides.
consanguineus, Klotzsch & Garcke, l. c. 64 = E. consanguinea.
coralloides, Rafin. Fl. Tellur. iv. 115 = E. coralloides.
cotinifolius, Haw. Syn. Pl. Succ. 141 = E. cotinifolia.
craspedius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 87 = E. craspedia.
crispus, Haw. Rev. Pl. Succ. 61 = E. elliptica.
cuneifolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 76 = E. cuneifolia.
Cupani, Klotzsch & Garcke, l. c. 85 = E. Cupani.
cyatophorus, Moench, Meth. 667 = E. heterophylla.
cybirensis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 79 = E. cybirensis.
Cyparissas, Hill, Hort. Kew. 172/4, sphalm. = seq.
Cyparissias, Lam. Fl. Fr. iii. 96 = E. Cyparissias.
damascenus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 96 = E. tinctoria.
declinatus, Moench, Meth. Suppl. 284 = E. portlandica.
deflexus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 84 = E. deflexa.
dendroides, Hill, Hort. Kew. 172/4 = E. dendroides.
densus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 65 = E. densa.
denticulatus, Moench, Meth. 668 = E. spinosa.
depauperatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 79 = E. depauperata.
desertorum, Klotzsch & Garcke, l. c. 97 = E. desertorum.
diffusus, Lam. Fl. Fr. iii. 101 = E. spinosa.
dilatatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 89 = E. dilatata.

TITHYMALUS:—
discolor, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 97 = Euphorbia Esula.
divergens, Klotzsch, in Bot. Ergeb. Waldem. Reise, 114 = E. Rothiana.
dracunculoides, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 84 = E. dracunculoides.
dulcis, Scop. Fl. Carn. ed. II. i. 334 = E. dulcis.
dumosus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 70 = E. thamnoides.
Eckloni, Klotzsch & Garcke, l. c. 68 = E. pistiaefolia.
Elliottii, Klotzsch & Garcke, l. c. 67 = E. Ipecacuanha.
ellipticus, Klotzsch & Garcke, l. c. 69 = E. elliptica.
epicyparissias, Klotzsch & Garcke, l. c. 88 = E. epicyparissias.
epithymoides, Klotzsch & Garcke, l. c. 79 = E. epithymoides.
eriophorus, Klotzsch & Garcke, l. c. 65 = E. eriophora.
erubescens, Klotzsch & Garcke, l. c. 96 = E. macrostegia.
erythradenius, Klotzsch & Garcke, l. c. 86 = E. erythradenia.
erythrinus, Klotzsch & Garcke, l. c. 91 = E. erythrina.
erythrodon, Klotzsch & Garcke, l. c. 87 = E. erythrodon.
erythrorrhizus, Klotzsch & Garcke, l. c. 71 = E. erythrorhiza.
Esula, Hill, Hort. Kew. 172/4; Scop. Fl. Carn. ed. II. i. 338 = E. Esula.
esulaeformis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 84 = E. campestris.
exiguus, Lam. Fl. Fr. iii. 100 = E. exigua.
falcatus, Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 292; et in Abh. Akad. Berl. (1860) 83 = E. falcata.
Fendleri, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 85 = E. Fendleri?
firmus, Klotzsch & Garcke, l. c. 73 = E. Gerardiana.
flaccidus, Moench, Meth. 665 = E. mauritanica.
foliosus, Klotzsch & Garcke, in Abh. Akad. Berl. (1800) 67 = E. dumosa.
fragiferus, Klotzsch & Garcke, l. c. 79 = E. fragifera.
fruticosus, Gilib. Fl. Lituan. ii. 206 = E. palustris.
fruticosus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 78 = E. Bivonae.
furcillatus, Klotzsch & Garcke, l. c. 86 = E. furcillata.
Gaillardoti, Klotzsch & Garcke, l. c. 74 = E. Gaillardoti.
galilaeus, Klotzsch & Garcke, l. c. 83 = E. falcata.
Gayi, Klotzsch & Garcke, l. c. 88 = E. Gayi.
genistoides, Klotzsch & Garcke, l. c. 97 = E. genistoides.
Gerardi, Rafin. Fl. Tellur. iv. 115 = seq.?
Gerardianus, Steud. Nom. ed. II. ii. 689; Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 291 = E. Gerardiana.
glaberrimus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 95 = E. iteophylla.
glaucus, Moench, Meth. Suppl. 282 = E. Moenchiana.
gracilis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 92 = E. Esula.
graecus, Klotzsch & Garcke, l. c. 84 = E. graeca.
granulatus, Rafin. Fl. Tellur. iv. 115 = E. granulata.
Guyonianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 82 = E. Guyoniana.
hebecarpus, Klotzsch & Garcke, l. c. 88 = E. hebecarpa.
helioscopius, Hill, Hort. Kew. 172/3 = E. helioscopia.
herniariaefolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 87 = E. herniariaefolia.
heterophyllus, Haw. Syn. Pl. Succ. 141 = E. heterophylla.
hiberna, Rafin. Fl. Tellur. iv. 115 = E. hyberna.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 115 = E. himalayensis.
hirsutus, Lam. Fl. Fr. iii. 98 = E. pilosa.
Hochstetterianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 97 = E. monticola.
Hohenackeri, Klotzsch & Garcke, l. c. 91 = E. Orphanidis.
huachanhanus, Klotzsch & Garcke, l. c. 71 = E. huachanhana.
humilis, Klotzsch & Garcke, l. c. 67 = E. humilis.
hybernus, Hill, Hort. Kew. 172/4 = E. hyberna.
imbricatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 73 = E. imbricata.
inderiensis, Klotzsch & Garcke, l. c. 65 = E. pygmaea.
involucratus, Klotzsch & Garcke, l. c. 91 = E. involucrata, *E. Mey.*

TITHYMALUS :—
Ipecacuanhae, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 69 = Euphorbia Ipecacuanha.
ispahanicus, Klotzsch & Garcke, l. c. 89 = E. megalantha.
Kotschyanus, Klotzsch & Garcke, l. c. 96 = E. Kotschyana.
laetus, Haw. Syn. Pl. Succ. 140 = E. dendroides.
Lagascae, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 65 = E. Lagascae.
lateriflorus, Klotzsch & Garcke, l. c. 81 = E. osyridea.
Lathyris, Hill, Hort. Kew. 172/3 = E. Lathyris.
latifolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 89 = E. latifolia.
laurocerasifolius, Mill. Gard. Dict. ed. VIII. n. 2 = Pedilanthus padifolius.
leptaleus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 65 = E. graminea.
linearis, Rafin. Fl. Tellur. iv. 115 = E. linearis.
linifolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 81 = E. spinosa.
linifolius, Lam. Fl. Fr. iii. 95 = E. Esula.
literatus, Rafin. Fl. Tellur. iv. 115 = E. platyphyllos ?
longepetiolatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 68 = E. elliptica.
lucidus, Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 292; et in Abh. Akad. Berl. (1860) 89 = E. lucida.
lutescens, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 80 = E. lutescens.
macroceras, Klotzsch & Garcke, l. c. 95 = E. macroceras.
macrocladus, Klotzsch & Garcke, l. c. 97 = E. tinctoria.
macrorrhizus, Klotzsch & Garcke, l. c. 79 = E. macrorhiza.
maculatus, Moench, Meth. 666 = E. maculata.
maritimus, Lam. Fl. Fr. iii. 90 = E. Paralias.
Marschallianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 86 = E. Marschalliana.
mauritanicus, Haw. Syn. Pl. Succ. 139 = E. mauritanica.
medicagineus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 83 = E. medicaginea.
megalanthus, Klotzsch & Garcke, l. c. 91 = E. megalantha.
melapetalus, Klotzsch & Garcke, l. c. 81 = E. melapetala.
melliferus, Moench, Meth. Suppl. 282 = E. mellifera.
Meyeri, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 75 = E. erubescens.
micrantha, Rafin. Fl. Tellur. iv. 115 (Quid ?).—Hab. ?
microsciadius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 74 = E. striatella.
microsphaerus, Klotzsch & Garcke, l. c. 74 = E. microsphaera.
monticolus, Klotzsch & Garcke, l. c. 87 = E. bounophila.
multicaulis, Klotzsch & Garcke, l. c. 98 = E. sclerophylla.
Myrsinitis, Hill, Hort. Kew. 172/4 = E. Myrsinites.
myrtifolius, Mill. Gard. Dict. ed. VIII. n. 1 = Pedilanthus tithymaloides.
nevadensis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 90 = E. nevadensis.
nicaeënsis, Klotzsch & Garcke, l. c. 89 = E. nicaeënsis.
notadaenius, Klotzsch & Garcke, l. c. 78 = E. orientalis.
nudiflorus, Haw. Syn. Pl. Succ. 142 = E. nudiflora.
nummularis, Lam. Fl. Fr. iii. 101 = E. Chamaesyce.
oblongifolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 96 = E. amygdaloides.
obtusatus, Klotzsch & Garcke, l. c. 69 = E. obtusata.
obtusifolius, Klotzsch & Garcke, l. c. 74 = E. obtusifolia.
orientalis, Hill, Hort. Kew. 172/3 = E. orientalis.
ovatus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 97 = E. ovata.
pachyrrhizus, Klotzsch & Garcke, l. c. 69 = E. pachyrhiza.
Pallasii, Klotzsch & Garcke, l. c. 75 = E. Pallasii.
palustris, Hill, Hort. Kew. 172/4 = E. palustris.
panaceus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 84 = E. Terracina.
paniculatus, Rafin. Fl. Tellur. iv. 115 = E. corollata ?
papillosus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 77 = E. Duvalii.
Paralias, Hill, Hort. Kew. 172/3 = E. Paralias.
Paralias, Scop. Fl. Carn. ed. II. i. 338 = E. Gerardiana.

TITHYMALUS :—
parvulus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 93 = Euphorbia arvalis.
pendulus, Haw. Syn. Pl. Succ. 138 = E. pendula.
Peplis, Scop. Fl. Carn. ed. II. i. 340 = E. Peplis.
peploides, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 83 = E. peploides.
Peplus, Gaertn. Fruct. ii. 115 = E. Peplus.
persepolitanus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 74 = E. striatella.
Pestalozzae, Klotzsch & Garcke, l. c. 87 = E. Pestalozzae.
petiolaris, Haw. Syn. Pl. Succ. 141 = E. petiolaris.
Petitianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 96 = E. cerebrina.
Philippianus, Klotzsch & Garcke, l. c. 65 = E. Philippiana.
pictus, Haw. Syn. Pl. Succ. 143 = E. graminea.
pilosus, Hill, Hort. Kew. 172/3 = E. pilosa.
*pilulifer*us, Moench, Meth. Suppl. 283 = E. pilulifera.
pineus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 85 = E. pinea.
piscatorius, Haw. Syn. Pl. Succ. 139 = E. piscatoria.
Pithyusa, Scop. Fl. Carn. ed. II. i. 335 = E. Pithyusa.
platiphylos, Rafin. Fl. Tellur. iv. 115 = seq. ?
platyphyllos, Hill, Hort. Kew. 172/4 = E. platyphyllos.
plebejus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 86 = E. plebeia.
polycaulis, Klotzsch & Garcke, l. c. 97 = E. decipiens.
portlandicus, Hill, Hort. Kew. 172/3 = E. portlandica.
procerus, Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 291 = E. pilosa.
prunifolius, Haw. Syn. Pl. Succ. 143 = E. geniculata.
pterococcus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 72 = E. pterococca.
pumilus, Klotzsch & Garcke, l. c. 87 = E. herniariaefolia.
puniceus, Haw. Syn. Pl. Succ. 140 = E. punicea.
purpureus, Lam. Fl. Fr. iii. 98 = E. Characias.
pygmaeus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 84 = E. pygmaea.
Rapulum, Klotzsch & Garcke, l. c. 67 = E. Rapulum.
Regis-Jubae, Klotzsch & Garcke, l. c. 93 = E. Regis-Jubae.
repandus, Haw. Syn. Pl. Succ. 142 = E. repanda.
repens, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 96 = E. repens.
repetitus, Klotzsch & Garcke, l. c. 96 = E. repetita.
retusus, Klotzsch & Garcke, l. c. 83 = E. sulcata.
Reuterianus, Klotzsch & Garcke, l. c. 83 = E. Reuteriana.
revolvulus, Klotzsch & Garcke, l. c. 99 = E. corifolia.
rosea, Rafin. Fl. Tellur. iv. 115 = E. rosea ?
Rothianus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 89 = E. Rothiana.
rotundifolius, Lam. Fl. Fr. iii. 100 = E. Peplus.
rupestris, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 73 = E. rupestris.
rupestris, Lam. Fl. Fr. iii. 97 = E. Gerardiana.
rupicolus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 78 = E. rupicola.
sagittalis, Hill, Hort. Kew. 172/3, sphalm. = E. segetalis.
salicifolius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 89 = E. salicifolia.
saxatilis, Klotzsch & Garcke, l. c. 91 = E. saxatilis.
Schimperianus, Klotzsch & Garcke, l. c. 96 = E. Schimperiana.
schizoceras, Klotzsch & Garcke, l. c. 98 = E. tinctoria.
Schottii, Klotzsch & Garcke, l. c. 70 = E. Schottiana.
segetalis, Lam. Fl. Fr. iii. 90 = E. segetalis.
Seguieri, Scop. Fl. Carn. ed. II. i. 335 = E. nicaeënsis.
Selloi, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 66 = E. elodes.
semiperfoliatus, Klotzsch & Garcke, l. c. 95 = E. semiperfoliata.
serotina, Rafin. Fl. Tellur. iv. 115 = E. nicaeënsis ?
serratus, Gilib. Fl. Lituan. ii. 207 = E. Helioscopia.
serratus, Hill, Hort. Kew. 172/3 = E. serrata.
sessiliflorus, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 77 = E. oblongata.
silenifolius, Haw. Rev. Pl. Succ. 61 = E. elliptica.
sororius, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 65 = E. sororia.
spinosus, Rafin. Fl. Tellur. iv. 115; Klotzsch & Garcke, l. c. 77 = E. spinosa.
squamosus, Klotzsch & Garcke, l. c. 78 = E. aspera.
stenophyllus, Klotzsch & Garcke, l. c. 66 = E. stenophylla.
striatus, Klotzsch & Garcke, l. c. 86 = E. striatella.

TITHYMALUS :—
strictus, Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 290 = Euphorbia stricta.
subamplexicaulis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 69 = E. subamplexicaulis.
subcordatus, Klotzsch & Garcke, l. c. 88 = E. subcordata.
subcrenatus, Klotzsch & Garcke, l. c. 66 = E. portulacoides.
sylvaticus, Hill, Hort. Kew. 172/4 = E. amygdaloides.
syspirensis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 97 = E. tinctoria.
Szovitsii, Klotzsch & Garcke, l. c. 65 = E. Szovitsii.
taurinensis, Klotzsch & Garcke, l. c. 83 = E. taurinensis.
tenuifolius, Klotzsch & Garcke, l. c. 92 = E. tenuifolia.
Terracinus, Klotzsch & Garcke, l. c. 90 = E. Terracina.
Thwaitesii, Klotzsch & Garcke, l. c. 92 = E. Rothiana.
trapezoidalis, Klotzsch & Garcke, l. c. 96 = E. Terracina.
trichotomus, Klotzsch & Garcke, l. c. 81 = E. trichotoma.
trinervius, Klotzsch & Garcke, l. c. 85 = E. boetica.
tristis, Klotzsch & Garcke, l. c. 92 = E. Esula.
truncatus, Klotzsch & Garcke, l. c. 75 = E. erubescens.
tuberosus, Hill, Hort. Kew. 173/3 = E. tuberosa.
Tuckeyanus, C. Bolle, ex Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 93 = E. Tuckeyana.
Turczaninowii, Klotzsch & Garcke, l. c. 64 = E. Turczaninowii.
undulatus, Klotzsch & Garcke, l. c. 92 = E. undulata.
uniflorus, Haw. in Phil. Mag. x. (1831) 418 = E. uniflora.
variabilis, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 90 = E. variabilis.
venetus, Klotzsch & Garcke, l. c. 95 = E. Wulfenii.
verrucosus, Hill, Hort. Kew. 172/3 = E. cybirensis.
verrucosus, Rafin. Fl. Tellur. iv. 115 (Quid ?).—Hab. ?
virgatus, Haw. Syn. Pl. Succ. 139 = E. obtusifolia.
virgatus, Klotzsch & Garcke, ex Garcke, Fl. Deutschl. ed. IV. 292; et in Abh. Akad. Berl. (1860) 88 = E. virgata.
Welwitschii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 77 = E. Welwitschii.
Zeyheri, Klotzsch & Garcke, l. c. 71 = E. mauritanica.

TITRAGYNE, Salisb. Gen. Pl. Fragm. 9 (1866) = **Rohdea**, Roth (Liliac.).
japonicum, Salisb. l. c. = R. japonica.

TITTELBACHIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 (Titelbachia); et in Abh. Akad. Berl. 1854 (1855) 105. t. 10 A = **Begonia**, Tourn.
albiflora, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 = B. albiflora.
castaneaefolia, Klotzsch, l. c. = B. castaneaefolia.
complicata, Hassk. Hort. Bogor. Desc. 342 = B. Ottonis.
fuchsioides, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126; et in Abh. Akad. Berl. 1854 (1855) 106 = B. fuchsioides.
Hamiltoniana, Regel, Ind. Sem. Hort. Petrop. (1860) 45 = B. acuminata.
miniata, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126; et in Abh. Akad. Berl. 1854 (1855) 106 = B. fuchsioides.

TITTMANNIA, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 385. t. 38. *BRUNIACEAE*, Benth. & Hook. f. i. 671.
MOESSLERA, Reichb. in Moessl. Handb. ed. II. i. p. l. (1827); Consp. 160 (1828).
lateriflora, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 386.—Afr. austr.
laxa, Presl, Bot. Bemerk. 32 = lateriflora.

TITTMANNIA, Reichb. Ic. Bot. Exot. i. 27. t. 38 (1824) = **Vandellia**, Linn. (Scrophul.).
alsinoides, Spreng. Syst. ii. 800 = V. alsinoides.
angustifolia, Benth. in Wall. Cat. n. 3951 = V. angustifolia.
Colsmanni, Benth. l. c. n. 3950 = Bonnaya veronicaefolia.
elata, Benth. l. c. n. 3944 = V. elata.
erecta, Benth. l. c. n. 3947 = V. erecta.
grandiflora, Benth. l. c. n. 3949 = V. pedunculata.
grandiflora, Spreng. Syst. ii. 801 = Ilysanthes grandiflora.
hirta, Benth. in Wall. Cat. n. 3948 = V. mollis.

TITTMANNIA :—
monticola, Spreng. Syst. ii. 800 = Ilysanthes refracta.
obovata, Bunge, Enum. Pl. Chin. Bor. 49 = Mazus rugosus.
ovata, Benth. in Wall. Cat. n. 3942 = Vandellia hirsuta.
ovata, Reichb. Ic. Bot. Exot. i. 27 = V. crustacea.
pusilla, Benth. in Wall. Cat. n. 3945 = V. scabra.
scabra, Spreng. Syst. ii. 800 = V. crustacea.
scapigera, Spreng. l. c. = V. scapigera.
stachydifolia, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 156 = V. stachydifolia.
subulata, Spreng. Syst. ii. 801 = V. subulata.
sulcata, Wight, ex Wall. Cat. n. 3946 = V. scabra.
trichotoma, Benth. in Wall. Cat. n. 3943 = V. multiflora.
viscosa, Reichb. Ic. Bot. Exot. i. 26. t. 38 = V. hirsuta.

TITYRUS, Salisb. Gen. Pl. Fragm. 101 (1866) = **Narcissus**, Tourn. (Amaryll.).
bifrons, Salisb. l. c. = N. Tazetta.
compressus, Salisb. l. c. = N. compressus.
Jonquilla, Salisb. l. c. = N. Jonquilla.
primulaceus, Salisb. l. c. = N. Tazetta.
similis, Salisb. l. c. = N. Jonquilla.

TIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 373 = **Astragalus**, Tourn. (Legumin.).
hians, Moench, Meth. 162 = Oxytropis deflexa.
lineare, Moench, l. c. = A. sulcatus.
sulcatum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = A. sulcatus.

TOANABO, DC. Prod. i. 528 (1824), sphalm. = TAONABO, Aubl. = **Ternstroemia**, Linn. f.

TOBINIA, Desv. in Ham. Prod. Pl. Ind. Occ. 56 (1825) = **Zanthoxylum**, Linn. (Rutac.).
acuminatum, Desv. l. c. 57 = Z. acuminatum.
coriacea, Desv. l. c. = Z. coriaceum.
emarginata, Desv. l. c. = Z. emarginatum.
punctata, Griseb. Fl. Brit. W. Ind. 137 = Z. punctatum.
spinosa, Desv. in Ham. Prod. Pl. Ind. Occ. 57 = Z. spinosum.
ternata, Desv. l. c. = Z. ternatum.

TOCOCA, Aubl. Pl. Gui. i. 437. t. 174 (1775). *MELASTOMACEAE*, Benth. & Hook. f. i. 764.
HAPPIA, Neck. Elem. ii. 126 (1790).
SPHAEROGYNE, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 331.
acuminata, *Benth. Bot. Voy. Sulph.* 94.—N. Granat.
aristata, *Benth. in Hook. Journ. Bot.* ii. (1840) 305.—Guian.
Aubletii, D. Don, in Mem. Wern. Soc. iv. (1823) 304 = guianensis.
barbata, Benth. in Hook. Journ. Bot. ii. (1840) 305 = guianensis.
bullifera, *Mart. & Schrank, ex DC. Prod.* iii. 165.—Bras.
bullifera, Triana, in Trans. Linn. Soc. xxviii. (1871) 132, partim = occidentalis.
cardiophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 91.—Bras.
castrata, Naud. l. c. 93 = subnuda.
ciliata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 133.—Bras.
cinnamomea, *Triana, l. c.*—Bras.
cordata, *Berg, ex Triana, l. c.*—Bras.
coriacea, *S. Moore, in Journ. Bot.* xviii. (1880) 3.—Honduras.
coronata, *Benth. in Hook. Journ. Bot.* ii. (1840) 303.—Am. trop.
didymophysca, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 90 = formicaria.
egensis, *Naud. l. c.* 92.—Bras.
erythrostachya, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 132 = bullifera.
formicaria, *Mart. ex DC. Prod.* iii. 165.—Bras.
formicata, Steud. Nom. ed. II. ii. 689, sphalm. = praec.
guianensis, *Aubl. Pl. Gui.* i. 438. t. 174.—Am. trop.
heterophylla, D. Don, in Mem. Wern. Soc. iv. (1823) 305 = Calophysa heterophylla.
hirsuta, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 132 = formicaria.
hirta, *Berg, ex Triana, l. c.*—Venezuela.
lanata, *D. Don, ex Loud. Hort. Brit.* 174.—Ins. Trinit.
lancifolia, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 133.—Bras.

TOCOCA:—
latifolia, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 90.—Bras.
macrophysca, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 132.—Bras.
macroptera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 94.—Bras.
macrosperma, Mart. Nov. Gen. et Sp. iii. 148. t. 279 = Myrmidone macrosperma.
Mayeta, D. Don, in Mem. Wern. Soc. iv. (1823) 305 = Maieta guianensis, Poeppigii.
nitens, Triana, in Trans. Linn. Soc. xxviii. (1871) 133 = Weddellii.
occidentalis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 92.—Peruv.
orcheophora, *Naud. l. c.* 89.—Boliv.
parviflora, *Spruce, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 132.—Peruv.
pauciflora, *Spruce, ex Triana, l. c.* 131.—Bras.
planifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 305.—Am. trop.
platyphylla, *Benth. Pl. Hartw.* 181.—Am. trop.
pubescens, Benth. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 133 = coronata.
Roraimae, Benth. in Hook. Journ. Bot. ii. (1840) 305 (*Roreimi*) = guianensis.
sanguinea, D. Don, in Mem. Wern. Soc. iv. (1823) 305 = Miconia hispidissima.
scabriuscula, Benth. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 133 = planifolia.
spadiciflora, *Triana, l. c.* 132.—N. Granat.
stephanotricha, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 95.—Bras.
subciliata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 133.—Bras.
subnuda, *Benth. in Hook. Journ. Bot.* ii. (1840) 303.—Guian.
truncata, *Benth. l. c.* 304.—Bras.
vesiculosa, DC. Prod. iii. 166 = Calophysa vesiculosa.
Weddellii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 94.—Bras.

TOCOYENA, Aubl. Pl. Gui. i. 131. t. 50 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 83.
JOCAYENA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 82.
UCRIANA, Willd. Sp. Pl. i. 961 (1797).
acutiflora, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 82.—Bras.
brasiliensis, *Mart. l. c.*—Bras.
bullata, *Mart. l. c.* 80.—Bras.
foetida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 25. t. 129.—Bras.
havanensis, A. Rich. Fl. Cub. Fanerog. ii. 8 = Posoqueria havanensis.
hirsuta, *Moric. ex DC. Prod.* iv. 375; *et in Pl. Nouv. Am.* 85. t. 56.—Bras.
insignis, Roem. & Schult. Syst. v. 229 = Posoqueria latifolia.
latifolia, Lam. Illustr. ii. 259 = Posoqueria latifolia.
longiflora, *Aubl. Pl. Gui.* i. 131.—Guian.
longifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 411.—N. Granat.
longifolia, Poir. Encyc. vii. 692 = longiflora.
macrophylla, H. B. & K. Nov. Gen. et Sp. iii. 412 = Posoqueria latifolia.
microdon, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 82.—Bras.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iii. 411.—N. Granat.
scandens, Blume, Bijdr. 980 = Randia longiflora.
speciosa, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
velutina, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 81.—Bras.
viscidula, *Mart. l. c.* 80.—Bras.

TODAROA, Parl. in Webb & Berth. Phyt. Canar. ii. 155. t. 74 (1836-50). *UMBELLIFERAE*, Benth. & Hook. f. i. 900.
aurea, *Parl. l. c.*—Ins. Canar.
montana, Webb, ex Benth. & Hook. f. Gen. i. 897, in obs. = Tinguarra montana.

TODAROA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 28 = **Campylocentron**, Benth. (Orchid.).
micrantha, A. Rich. & Gal. l. c. = C. Schiedei.

TODDALIA, Juss. Gen. 371 (1789). *RUTACEAE*, Benth. & Hook. f. i. 300.
ASAPHES, DC. Prod. ii. 90 (1825).
ASPIDOSTIGMA, Hochst. in Flora, xxvii. (1844) 18.
BOSCIA, Thunb. Prod. Pl. Cap. 32 (1794).
CRANZIA, Schreb. Gen. 143 (1789).
DIPETALUM, Dalz. in Hook. Kew Journ. ii. (1850) 38.
DUNCANIA, Reichb. Consp. 197 (1828).
RUBENTIA, Boj. ex Steud. Nom. ed. II. ii. 475 (1841).
SCOPOLIA, Sm. Ic. Ined. ii. t. 34 (1790).
TECLEA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 90.
VEPRIS, Comm. ex A. Juss. in Mém. Mus. Par. xii. (1825) 509.
aculeata, *Pers. Syn.* i. 249.—Ind. or.; Malaya; Ins. Mascar.
ambigua, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 444.—Ins. Philipp.
angustifolia, Lam. Illustr. ii. 117 = aculeata ?
asiatica, Lam. l. c. 116 = aculeata.
bilocularis, *Wight & Arn. Prod.* 149.—Ind. or.
effusa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 443.—Ins. Philipp.
floribunda, Wall. Pl. As. Rar. iii. 17. t. 232 = aculeata.
inermis, Pers. Syn. i. 249 = paniculata.
lanceolata, *Lam. Illustr.* ii. 117.—Afr. trop.; Ins. Maurit.
megapotamica, *G. Don, Gen. Syst.* i. 805.—Bras.
micracantha, Miq. Fl. Ind. Bat. i. II. 672, sphalm. = seq.
micrantha, *Steud. Nom.* ed. II. ii. 690.—Java.
mitis, Miq. ex Hook. f. Fl. Brit. Ind. i. 492 = Zanthoxylum ovalifolium.
natalensis, *Sond. in Harv. & Sond. Fl. Cap.* i. 447.—Afr. austr.
nitida, Lam. Illustr. ii. 116 = aculeata.
nobilis, *Hook. f. ex Oliver, Fl. Trop. Afr.* i. 306.—Afr. trop.
paniculata, *Lam. Illustr.* ii. 117.—Ins. Mascar.
pilosa, *Baker, in Journ. Linn. Soc.* xxi. (1884) 329.—Madag.
rubicaulis, Roem. & Schult. Syst. v. 323 = aculeata.
schmidelioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 118.—Madag.
venosa, *G. Don, Gen. Syst.* i. 805.—Bras.
Willdenowii, *Steud. Nom.* ed. II. ii. 690.—Ins. Mascar.

TODDA-PANA, Adans. Fam. ii. 25 (1763) = **Cycas**, Linn.

TOECHIMA, Radlk. in Sitzb. Math.-Phys. Akad. Muench. ix. (1879) 501, 671. *SAPINDACEAE*.
Daemelianum, *Radlk. l. c.* 672.—Austral.
erythrocarpum, *Radlk. l. c.* 671.—Austral.
subteres, *Radlk. l. c.*—N. Guin.
tenax, Radlk. l. c. = Ratonia tenax.

TOFFIELDIA, Schrank, in Denkschr. Akad. Muench. 1813 (1814) 94, sphalm. = seq.

TOFIELDA, Pers. Syn. i. 399 (1805) = seq.

TOFIELDIA, Huds. Fl. Angl. ed. II. 157 (1778). *LILIACEAE*, Benth. & Hook. f. iii. 828.
CONRADIA, Rafin. Neogenyt. 3 (1825).
CYMBA, Dulac, Fl. Hautes-Pyr. 117 (1867).
HEBELIA, C. C. Gmel. Fl. Bad. ii. 117 (1806).
HERITERIA, Schrank, Baier. Fl. i. 133 (1789).
ISIDROGALVIA, Ruiz & Pav. Fl. Per. iii. 69. t. 302 (1802).
LEPTILIX, Rafin. Neogenyt. 3 (1825).
TRIANTHA, Nutt. Gen. Am. i. 235 (1818).
allemannica, Bluff & Finger. Comp. Fl. Germ. ed. I. i. 478 = calyculata.
alpina, Hoppe & Sternb. in Denkschr. Bot. Ges. Regensb. i. (1815) 152 = palustris.
alpina, Sm. in Trans. Linn. Soc. xii. (1817) 24 = calyculata.
anthericoides, Roth, Enum. Pl. Phaen. i. II. 109 = calyculata.
borealis, Wahlenb. Fl. Lapp. 89 = palustris.
calyculata, *Wahlenb. l. c.* 90, in obs.—Europ.; Am. bor.
capitata, Hoppe, ex Michl, in Flora, iii. (1820) 314 = palustris.

TOFIELDIA:—
cernua, *Sm. in Trans. Linn. Soc.* xii. (1817) 244.—Sibir.
coccinea, Richards. in Frankl. Narr. 1st Journ. App. 736 = calyculata.
collina, Schult. Oesterr. Fl. ed. II. i. 581 = calyculata.
falcata, *Pers. Syn.* i. 399.—Peruv.
flexuosa, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 28 = falcata.
frigida, H. B. & K. Nov. Gen. et Sp. i. 267 = falcata.
glaberrima, Macbride, ex Ell. Sketch, i. 424 = glabra.
glabra, *Nutt. Gen. Am.* i. 235.—Am. bor.
glacialis, Gaud. Fl. Helv. ii. 596 = calyculata.
glutinosa, *Pers. Syn.* i. 399.—Am. bor.
gracilis, *Franch. & Sav. Enum. Pl. Jap.* ii. 531.—Japon.
himalaica, *Baker, in Journ. Linn. Soc.* xvii. (1879) 489.—Reg. Himal.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 201.—Japon.
nepalensis, Wall. Cat. n. 5097 (Aletridis sp.).—Reg. Himal.; China.
nuda, Franch. & Sav. Fl. Jap. ii. 89, partim = stenantha.
nuda, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 176.—Japon.
nutans, *Willd. ex Schult. f. Syst.* vii. 1573, in obs.—Sibir. or.; Japon.; Am. bor.
occidentalis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 283.—Am. bor. occ.
ossifraga, Nem. ex Chaub. in Act. Linn. Soc. Bord. xix. (1853) 228 = Narthecium ossifragum.
palustris, DC. Fl. Fr. iii. 193 = calyculata.
palustris, *Huds. Fl. Angl.* ed. II. i. 175.—Reg. bor. et arct.
pubens, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 28; [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 325.—Am. bor.
pubescens, Pers. Syn. i. 399 = pubens.
pusilla, Pers. l. c. = palustris.
racemosa, Hoppe & Hornsch. in Flora, iv. (1821) 189 = calyculata.
rubra, F. Br. in Flora, iii. (1820) 469 = calyculata.
sessiliflora, Hook. Ic. Pl. t. 691 = falcata.
sibirica, Adams, in Mém. Soc. Nat. Mosc. ix. (1834) 236 = coccinea.
sordida, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 437.—Japon.
stenantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 530.—Japon.
stenopetala, Sm. in Trans. Linn. Soc. xii. (1817) 243. t. 8 f. 1 = calyculata.

TOKOYENA, Rich. ex Steud. Nom. ed. II. ii. 383, in syn. (1841), sphalm. = **Tocoyena**, Aubl. (Rubiac.).

TOLA, Wedd. ex Benth. & Hook. f. Gen. ii. 257 (1873) = **Lepidophyllum**, Cass. (Compos.).

TOLLATIA, Endl. Gen. 428 (1838) = **Layia**, Hook. & Arn. (Compos.).
chrysanthemoides, Endl. ex Walp. Rep. ii. 631 = L. chrysanthemoides.

TOLMIAEA, Buek, Ind. in DC. Prod. iii. 479 (1858), sphalm. = seq.

TOLMIEA, Hook. Fl. Bor. Am. ii. 44 (1834) = **Cladothamnus**, Bong. (Ericac.).
occidentalis, Hook. l. c. 45 = C. pyroliflorus.

TOLMIEA, Torr. & Gray, Fl. N. Am. i. 582 (1840).
SAXIFRAGEAE, Benth. & Hook. f. i. 638.
LEPTAXIS, Rafin. Fl. Tellur. ii. 75 (1836), nomen prius.
Menziesii, *Torr. & Gray, Fl. N. Am.* i. 582.—Am. bor. occ.

TOLPIS, Adans. Fam. ii. 112 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 508.
AETHONIA, D. Don, in Edinb. N. Phil. Journ. (1828–29) 309.
CALODONTA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 448.
CHATELANIA, Neck. Elem. i. 53 (1790).
DREPANIA, Juss. Gen. 169 (1789).
POLYCHAETIA, Tausch, ex Less. Syn. Comp. 129 (1832).
SCHMIDTIA, Moench, Meth. Suppl. 217 (1802).
SWERTIA, All. Fl. Pedem. i. 208 (1785).
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 459.—Abyss.

TOLPIS:—
abyssinica, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 399 = altissima.
altissima, *Pers. Syn.* ii. 377.—Reg. Mediterr.; Syria.
ambigua, Req. ex Nym. Consp. 473 = virgata.
baetica, Jord. & Fourr. Ic. Fl. Eur. i. 43 = barbata.
barbata, Bivona, Monog. 13. t. 3 = umbellata.
barbata, *Gaertn. Fruct.* ii. 372. t. 160.—Reg. Mediterr.
biaristata, Link, Handb. i. 811, sphalm. = *sexaristata*.
Bivonae, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 87 = barbata.
Calderae, *Bolle, in Bonplandia*, vii. (1859) 298.—Ins. Canar.
canariensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 84 = barbata.
capensis, Sch. Bip. in Bonplandia, ix. (1861) 172 = Hieracium capense.
concolor, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 85 = umbellata.
coronopifolia, *Bivona, Monog.* 17. t. 5.—Europ.; Ins. Canar.
cretica, Sieber, ex Nym. Consp. 473 = altissima.
crinita, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 24 = umbellata *vel* barbata.
crithmifolia, *DC. Prod.* vii. 87.—Ins. Canar.
dichroa, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 86 = barbata.
discolor, Jord. & Fourr. l. c. = barbata.
eriantha, Sch. Bip. in Bonplandia, ix. (1861) 173, in syn. = Hieracium erianthum.
farinulosa, *Walp. Ann.* ii. 955.—Ins. Cap. Virid.
filiformis, *DC. Prod.* vii. 87.—Ins. Mader.
fruticosa, *Schrank, Pl. Rar. Hort. Monac.* t. 46.—Ins. Canar.
glandulifera, *Bolle, in Bonplandia*, vii. (1859) 298.—Ins. Canar.
grandiflora, Tenore, Viagg. Basil. e Calabr. 128 = altissima.
laciniata, *Webb & Berth. Phyt. Canar.* ii. 406.—Ins. Canar.
Lagopoda, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
macrodonta, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 85 = umbellata.
macrorhiza, *DC. Prod.* vii. 87.—Ins. Mader.
macrorhiza, H. C. Wats. ex Seub. Fl. Azor. 33 = nobilis.
madagascariensis, Sch. Bip. in Bonplandia, ix. (1861) 172 = Hieracium madagascariense.
microcephala, Pomel, Nouv. Mat. Fl. Atl. 271 = umbellata.
nobilis, *Hochst. in Seub. Fl. Azor.* 33. t. 11.—Ins. Azor.
pallidiflora, Jord. & Fourr. Brev. Pl. Nouv. fasc. ii. 85 = barbata.
pectinata, *DC. Prod.* vii. 87.—Ins. Mader.
prolifera, Moench, Meth. Suppl. 220 = barbata.
pygmaea, Steud. Nom. ed. II. ii. 691 = Leontodon nudicaulis.
quadriaristata, Bivona, Monog. 9 = altissima.
quercifolia, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 398, 399 = umbellata, macrorhiza.
quinqueradiata, C. Koch, in Linnaea, xxiii. (1850) 654 = umbellata.
sexaristata, Bivona, Monog. 11. t. 2 = altissima.
staticefolia, Sch. Bip. in Bonplandia, ix. (1861) 173 = Hieracium staticefolium.
succulenta, Lowe, Man. Fl. Mad. i. 525 = fruticosa.
umbellata, *Bertol. in Mem. Soc. Emul. Genova*, ii. (1803) 133.—Reg. Mediterr.
umbellata, Bertol. l. c., partim = barbata.
virgata, Bertol. l. c. 135 = altissima.
virgata, *Bivona, Monog.* 15. t. 4.—Reg. Mediterr.
Webbii, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 409.—Ins. Canar.

TOLUIFERA, Linn. Gen. ed. II. 182 (1742) = **Myroxylon**, Linn. (Legumin.).
Balsamum, Linn. Sp. Pl. 384 = M. toluiferum.
pedicellata, Baill. Hist. Pl. ii. 383 = M. pedicellatum.
Pereirae, Baill. l. c. = M. Pereirae.
peruifera, Baill. l. c. = M. peruiferum.
pubescens, Baill. l. c. = M. pubescens.
punctata, Baill. l. c. = M. toluiferum.

TOLUIFERA, Lour. Fl. Cochinch. 262 (1790) = **Loureira**, Meissn. (Gen. ignot.).
cochinchinensis, Lour. l. c. = L. cochinchinensis, Buchanania Lanzan.

TOLUMNIA, Rafin. Fl. Tellur. ii. 101 (1836) = **Oncidium**, Sw. (Orchid.).
pulchella, Rafin. l. c. = O. pulchellum.

TOLYPANTHUS, Blume, in Schult. f. Syst. vii. II. 1731 (1830) = **Loranthus**, Linn.
coriaceus, Blume, l. c. = L. coriaceus.
involucratus, Blume, l. c. = L. involucratus.
loniceroides, Blume, l. c. = L. loniceroides.

TOLYPEUMA, E. Mey. in Drège, Zwei Pfl. Docum. 226 (1843), nomen = **Nesaea**, Comm. (Lythrar.).
floridum, E. Mey. l. c. = N. floribunda.

TOMANTEA, Steud. Nom. ed. II. ii. 691 (1841), sphalm. = seq.

TOMANTHEA, DC. Prod. vi. 564 (1837) = **Centaurea**, Linn. (Compos.).
Aucheri, DC. l. c. = C. luzeoides.

TOMANTHERA, Rafin. New Fl. Am. ii. 65 (1836) = **Gerardia**, Linn. (Scrophul.).
auriculata, Rafin. l. c. 66 = G. auriculata.
lanceolata, Rafin. l. c. = Zaluzianskya villosa.

TOMARIS, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Linn. (Polygon.).

TOMEX, Forsk. Fl. Aegypt. Arab. 32 (1775) = **Dobera**, Juss. (Salvad.).—Cf. Benth. & Hook. f. ii. 680; iii. 162, in nota.
dubia, Jacq. Hort. Vindob. ii. 90 = D. dubia.
glabra, Forsk. Fl. Aegypt. Arab. 32 = D. Roxburghii.

TOMEX, Linn. Diss. Dass. 5 (1747); Amoen. Acad. i. 389 (1749) = **Callicarpa**, Linn. (Verben.).
tomentosa, Linn. Sp. Pl. 118 = C. lanata.

TOMEX, Thunb. Nov. Gen. iii. 65 (1783) = **Litsea**, Lam. (Laurin.).
Boldo, Nees, Syst. Laurin. 719 (Index), sphalm. = seq.
Bolo, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 64 = Lindera melastomacea.
Doshia, Buch.-Ham. ex D. Don, l. c. 65 = L. oblonga.
japonica, Thunb. Nov. Gen. iii. 64 = L. japonica.
laurina, Hornem. ex Meissn. in DC. Prod. xv. I. 229 = L. zeylanica, umbrosa.
monopetala, Hoffmgg. Verz. Pfl. Nachtr. II. 221 = L. polyantha.
pubescens, Willd. ex Meissn. in DC. Prod. xv. I. 189 = L. polyantha.
Sebifera, Willd. Sp. Pl. ii. 840 = L. Sebifera.
Tetranthera, Willd. l. c. 839 = L. Sebifera.

TOMIEPHYLLUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125 = **Scrophularia**, Tourn.
caninum, Fourr. l. c. = S. canina.
tenuisectum, Fourr. l. c. = S. canina.

TOMILIX, Rafin. New Fl. Am. ii. 72 (1836) = **Macranthera**, Torr. (Scrophul.).
bracteata, Rafin. l. c. = M. fuchsioides.

TOMMASINIA, Bertol. Fl. Ital. iii. 414 (1839) = **Peucedanum**, Tourn. (Umbell.).
Kotschyi, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 302 = Dorema Aucheri.
purpurascens, Boiss. Fl. Orient. ii. 979 = Dorema Aucheri.
Szowitsii, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 302 = Dorema Aucheri.
verticillaris, Bertol. Fl. Ital. iii. 414 = P. verticillare.

TOMODON, Rafin. Fl. Tellur. iv. 22 (1836) = **Hymenocallis**, Salisb. (Amaryll.).
coronarium, Rafin. l. c. = H. crassifolia.
floridanum, Rafin. l. c. = H. crassifolia.
pratense, Rafin. l. c. = H. crassifolia?
riparium, Rafin. l. c. = H. crassifolia?
rotatum, Rafin. l. c. = H. lacera.

TOMOSTIMA, Rafin. Neogenyt. 2 (1825) = **Draba**, Linn. (Crucif.).
caroliniana, Rafin. l. c. = D. caroliniana.
hispidula, Rafin. l. c. = D. caroliniana.

TOMOSTYLIS, Montr. in Mém. Acad. Lyon, x. (1860) 201 = **Crossostyles**, Forst. (Rhizoph.).
multiflora, Montr. l. c. = C. multiflora.

TOMOTRIS, Rafin. Fl. Tellur. ii. 89 (1836) = **Corymbis**, Thou. (Orchid.).
flava, Rafin. l. c. = C. flava.
polystachia, Rafin. l. c. (= *Chloidia vernalis*).—Jamaic.

TOMOXIS, Rafin. Fl. Tellur. iii. 54 (1836) = **Ornithogalum**, Tourn. (Liliac.).
coarctata, Rafin. l. c. = O. coarctatum.
odorata, Rafin. l. c. = O. suaveolens.
virens, Rafin. l. c. = O. virens.

TONABEA, Juss. Gen. 262 (1789) = TAONABO, Aubl. = **Ternstroemia**, Linn. f.

TONDIN, G. W. Schilling, Desc. Pl. Curat. Leprae, (1778) 198; et ex J. F. Gmel. Syst. 635 (1791) (Quid?).
surinamensium, G. W. Schilling, l. c. t. 1; et ex J. F. Gmel. l. c.—Guian.

TONELLA, Nutt. ex A. Gray, in Proc. Am. Acad. vii. (1868) 378. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 941.
collinsioides, *Nutt. ex A. Gray, l. c.*—Am. bor. occ.
floribunda, *A. Gray, in Bot. Calif.* i. 556.—Calif.

TONGUEA, Endl. Gen. 1419 (1841) = **Sisymbrium**, Tourn. (Crucif.).

TONINA, Aubl. Pl. Gui. ii. 856. t. 330 (1775). *ERIOCAULEAE*, Benth. & Hook. f. iii. 1024.
HYPHYDRA, Schreb. Gen. 666 (1791).
fluviatilis, *Aubl. Pl. Gui.* ii. 857. t. 330.—Guian.

TONNINGIA, Neck. Elem. iii. 165 (1790) = **Cyanotis**, D. Don (Commel.).

TONSELLA, Schreb. Gen. 34 (1789) = TONTELEA, Aubl. = **Salacia**, Linn. (Celastr.).

TONSHIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 225 (1825) = **Saurauja**, Willd. (Ternstroem.).
polypetala, Buch.-Ham. ex D. Don, l. c. = S. napaulensis.

TONTANEA, Aubl. Pl. Gui. i. 108. t. 42 (1775) = **Coccocypselum**, P. Br. (Rubiac.).
guianensis, Aubl. l. c. = C. Tontanea.
repens, Pers. Syn. i. 134 = C. Tontanea.

TONTELEA, Aubl. Pl. Gui. i. 31. t. 10 (1775) = **Salacia**, Linn. (Celastr.).
africana, Willd. Sp. Pl. i. 194 = S. africana.
attenuata, Miers, in Trans. Linn. Soc. xxviii. (1872) 384 = S. Richardii.
Aubletiana, Miers, l. c. 383 = S. scabra.
brachypoda, Miers, l. c. 387 = S. brachypoda.
brasiliensis, Spreng. Syst. i. 177 = S. brasiliensis.
Calypso, Spreng. l. c. = S. Calypso.
calypsoides, Endl. in Walp. Rep. i. 400 = S. calypsoides.
chinensis, Spreng. Syst. i. 177 = S. Roxburghii.
corrugulata, Miers, in Trans. Linn. Soc. xxviii. (1872) 386 = S. micrantha.
crassifolia, Spreng. Syst. i. 177 = S. crassifolia.
decussata, Vahl, Enum. ii. 30 = Hippocratea decussata.
disperma, Poir. Encyc. Suppl. v. 316 = Hippocratea indica.
elliptica, Spreng. Syst. i. 178 = S. elliptica.
erythroxyloides, Endl. ex Walp. Rep. i. 400 = S. elliptica.
floribunda, Miers, in Trans. Linn. Soc. xxviii. (1872) 385 = Hippocratea acapulcensis.
glomerata, Spreng. Syst. i. 177 = S. glomerata.
grandiflora, Spreng. l. c. 178 = S. grandifolia.
Hookeriana, Miers, in Trans. Linn. Soc. xxviii. (1872) 388.—Mexic.
laevigata, Hoffmgg. ex Link, Jahrb. i. III. (1820) 68 = S. laevigata.
levigata, Poir. Encyc. Suppl. v. 316 = Hippocratea ovata.
longifolia, Miers, in Trans. Linn. Soc. xxviii. (1872) 384 = Hippocratea malpighiaefolia.

TONTELEA :—
macrophylla, Poir. Encyc. Suppl. v. 317 = Hippocratea macrophylla.
madagascariensis, Vahl, Enum. ii. 32 = Salacia madagascariensis.
malpighiaefolia, G. F. W. Mey. Prim. Fl. Esseq. 20 = Hippocratea malpighiaefolia.
micrantha, Spreng. Syst. i. 177 = S. micrantha.
multiflora, Vahl, Enum. ii. 32 = S. multiflora.
oblongifolia, Spreng. Syst. i. 178 = S. elliptica.
obtusifolia, Endl. in Walp. Rep. i. 400 = S. obtusifolia.
paniculata, Poir. Encyc. Suppl. v. 316 = Hippocratea paniculata.
parviflora, Miers, in Trans. Linn. Soc. xxviii. (1872) 386 = S. parviflora.
prinoides, Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 184 = S. prinoides.
pyriformis, Hort. ex G. Don, in Loud. Hort. Brit. 14 = S. pyriforme.
Radula, Spreng. Syst. i. 177 = S. Radula.
scabra, Vahl, Enum. ii. 29 = S. scabra.
scandens, Aubl. Pl. Gui. i. 31 = S. scabra.
senegalensis, Vahl, Enum. ii. 31 = S. senegalensis.
serrata, Endl. in Walp. Rep. i. 401 = S. serrata.
trinervia, Spreng. Syst. i. 177 = S. trinervia.
undulata, Spreng. l. c. = S. undulata.
velutina, Endl. in Walp. Rep. i. 401 = S. velutina.

TOONA, M. Roem. Syn. Hesper. 131, 139 (1846) = **Cedrela**, P. Br. (Meliac.).
ciliata, M. Roem. l. c. 139 = C. Toona.
febrifuga, M. Roem. l. c. = C. Toona.
hexandra, M. Roem. l. c. = C. Toona.
longifolia, M. Roem. l. c. 140 = C. Toona.
serrata, M. Roem. l. c. 139 = C. Toona.
sinensis, M. Roem. l. c. = C. sinensis.
velutina, M. Roem. l. c. = Chickrassia tabularis.
villosa, M. Roem. l. c. 140 = Chickrassia tabularis.

TOPEINOSTEMON, C. Muell. in Walp. Ann. v. 509 (1858), err. typ. = **Tapeinostemon**, Benth. (Gentian.).

TOPIARIS, Rafin. Sylva Tellur. 43 (1838) = **Cordia**, Linn. (Borag.).
geniculata, Rafin. l. c. = C. mirabiloides.

TOPOBEA, Aubl. Pl. Gui. i. 476. t. 189 (1775). *MELASTOMACEAE*, Benth. & Hook. f. i. 770.
DREPANANDRUM, Neck. Elem. ii. 118 (1790).
anisophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 151.—N. Granat.
calycularis, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 149.—Mexic.
fragrans, Naud. l. c. = laevigata.
glaberrima, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 150.—N. Granat.
glabrescens, *Triana, l. c.* 149.—N. Granat.
gracilis, *Triana, l. c.* 150.—N. Granat.
inflata, *Triana, l. c.*—N. Granat.
insignis, *Triana, l. c.*—N. Granat.
laevigata, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 150.—Mexic.
latifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—N. Granat.
longipes, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 147.—Peruv.
multiflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—Peruv.
paniculata, Rich. ex DC. Prod. iii. 105 = Adelobotrys adscendens.
parasitica, *Aubl. Pl. Gui.* i. 476.—Guian.
punctulata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 150.—N. Granat.
rotundifolia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 150 = Blakea rotundifolia.
sessilifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 150.—N. Granat.
setosa, *Triana, l. c.* 149.—N. Granat.
stephanochaeta, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 148.—N. Granat.
subscabrula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 150.—N. Granat.
superba, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 147.—Panama.

TOQUERA, Rafin. Sylva Tellur. 40 (1838) = **Cordia**, Linn. (Borag.).
tomentosa, Rafin. l. c. = C. Toqueve.

TORDYLIOIDES, Wall. ex DC. Prod. iv. 199 (1830) = **Heracleum**, Linn. (Umbell.).
Brunonis, Wall. ex DC. l. c. = H. Brunonis.

TORDYLIOPSIS, DC. Prod. iv. 199 (1830) = **Heracleum**, Linn. (Umbell.).
Brunonis, DC. l. c. = H. Brunonis.

TORDYLIUM, Tourn. ex Linn. Gen. ed. I. 71 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 924.
AINSWORTHIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 343.
CONDYLOCARPUS, Hoffmgg. Gen. Umbell. ed. II. 202 (1816).
HASSELQUISTIA, Linn. Cent. Pl. i. 9 (1755); Amoen. Acad. iv. 270 (1759).
SYNELCOSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 345.
absinthifolium, Pers. Syn. i. 314 = Zosima orientalis.
aegyptiacum, *Lam. Illustr.* ii. 336.—Syria; Mesopot.
americanum, Nutt. ex DC. Prod. iv. 196 = Polytaenia Nuttallii.
Anthriscus, Linn. Sp. Pl. 240 = Caucalis Anthriscus.
apulum, *Linn. l. c.* 239.—Reg. Mediterr.
apulum, Reichb. Fl. Germ. Excurs. 452 = officinale.
Arthedia, Crantz, Cl. Umb. Emend. 56 = Artedia squamata.
asperum, Gilib. Fl. Lituan. ii. 11 = Caucalis Anthriscus.
Aucheri, Jaub. & Spach, Illustr. iii. 46. t. 237 = Ormosciadium Aucheri.
austriacum, Hort. Viln. (1811); ex Schult. Syst. vi. 462 (Quid?).—Europ.
brachytaenium, *Boiss. & Heldr. Diagn.* Ser. I. x. 45.—Pamphyl.
Brunonis, Wall. Cat. n. 590 = Heracleum Brunonis.
cappadocicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 349.—Cappadoc.
concinnum, Tenore, Syll. Fl. Nap. App. iv. 11 = apulum.
cordatum, *Poir. Encyc.* vii. 712.—Ins. Cypr.; Mesopot.
cyrenaicum, Spreng. Syst. i. 895 = Apium nodiflorum.
divaricatum, Moench, Meth. 77 = syriacum.
grande, Salisb. Prod. 160 = maximum.
grandiflorum, Moench, Meth. 78 = apulum.
Hasselquistia, Crantz, Cl. Umb. Emend. 57 = aegyptiacum.
Hasselquistiae, *DC. Prod.* iv. 198.—Syria.
humile, Desf. Fl. Atlant. i. 235. t. 58 = apulum.
insulare, Clarke, Trav. iii. 337 = apulum.
intermedium, Pass. in Nuov. Giorn. Bot. Ital. iii. (1871) 167, 168 = apulum.
lanatum, *Boiss. Fl. Orient.* ii. 1030.—As. Min.
latifolium, Linn. Sp. Pl. 240 = Caucalis latifolia.
lusitanicum, Hoffmgg. & Link, Fl. Port. ii. 389 = maximum.
macropetalum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 348.—Caria.
magnum, Brot. Fl. Lusit. i. 450 = maximum.
maximum, *Linn. Sp. Pl.* 240.—Europ.; Reg. Cauc.
microspermum, Tenore, Add. (1827); ex DC. Prod. iv. 198 = officinale.
nodiflorum, Salisb. Prod. 160 = seq.
nodosum, Linn. Sp. Pl. 240 = Caucalis nodosa.
officinale, *Linn. Sp. Pl.* 239.—Europ. austr.
officinale, Reichb. Fl. Germ. Excurs. 452 = apulum.
peregrinum, Linn. Mant. i. 55 = Capnophyllum dichotomum.
persicum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1032.—Persia.
Pestalozzae, *Boiss. Diagn.* Ser. I. x. 45.—Lycia.
Secacul, Mill. Gard. Dict. ed. VIII. n. 5 = Pastinaca Secacul.
segetum, Hort. Par. ex Schult. Syst. vi. 484 = Caucalis arvensis.
Sekakul, DC. Prod. iv. 90 = *T. Secacul.*
siifolium, Scop. Fl. Carn. ed. II. i. 194 = Heracleum austriacum.
suaveolens, Delile, Fl. Egypte, Illustr. 58. n. 323 = Pastinaca Secacul.
sulcatum, Desf. Tabl. ed. II. Suppl. 274 (Quid?).—Hab.?
syriacum, *Linn. Sp. Pl.* 239.—Cilicia; Syria.
verecundum, Salisb. Prod. 160 = Caucalis Anthriscus.

TOREALA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Inga**, Scop. et **Parkia**, R. Br. (Legumin.).

TORENIA, Linn. Diss. Chen. 45 (1751); Amoen. Acad. iii. 25 (1756). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 954.
CAËLA, Adans. Fam. ii. 209 (1763).
DUNALIA, R. Br. in Salt, Abyss. App. 64 (1814).
LEGAZPIA, Blanco, Fl. Filip. ed. II. 338 (1845).
NORTENIA, Thou. Gen. Nov. Madag. 9 (1806).
PENTSTERIA, Griff. Notul. iv. 118 (1854).
PERISTEIRA, Hook. f. Fl. Brit. Ind. iv. 277–8–9 (1884), sphalm.
alata, *Benth. in DC. Prod.* x. 411.—Ins. Molucc.
alba, Buch.-Ham. ex Wall. Cat. n. 3961 = pedunculatis, Vandellia crustacea.
angustifolia, Buch.-Ham. l. c. n. 3951 A = Vandellia angustifolia.
arracanensis, *Hort. ex Paxt. Mag. Bot.* xv. (1849) 198. —Burma ?
asiatica, Benth. Scroph. Ind. 38 ; et in DC. Prod. x. 410 = pedunculatis.
asiatica, Herb. Madr. ex Wall. Cat. n. 3954 A = cordifolia.
asiatica, *Linn. Sp. Pl.* 619.—As. trop.
asiatica, Wall. Cat. n. 3953, partim = vagans.
asiatica, Wight, ex Hook. f. Fl. Brit. Ind. iv. 278 = bicolor.
auriculaefolia, Dombr. Flor. Mag. x. (1871) t. 534 = Craterostigma pumilum.
Bailloni, Godefroy, ex André, in Illustr. Hortic. xxv. (1878) 136. t. 324 = flava.
Benthamiana, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 226.—China.
bicolor, *Dalz. in Hook. Kew Journ.* iii. (1851) 38.—Ind. or.
calcarata, Griff. Notul. iv. 124 = Vandellia angustifolia.
cardiosepala, Benth. in DC. Prod. x. 409 = polygonoides.
ciliaris, *Sm. in Rees, Cycl.* xxxvi. n. 3.—Malaya.
ciliata, Hook. f. Fl. Brit. Ind. iv. 278, sphalm. = praec.
coerulea, Miq. Fl. Ind. Bat. Suppl. 562 = ciliaris.
colorans, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) t. 94 = concolor.
concolor, *Lindl. Bot. Reg.* (1846) t. 62.—China.
cordifolia, Benth. in Wall. Cat. n. 3954 = asiatica.
cordifolia, Buch.-Ham. ex Wall. l. c. n. 3949 D = Vandellia pedunculata.
cordifolia, *Roxb. Pl. Corom.* ii. 52. t. 161.—Ind. or. ; Malaya.
crustacea, Cham. & Schlecht. in Linnaea, ii. (1827) 570 = Vandellia crustacea.
diffusa, D. Don, Prod. Fl. Nep. 86 = vagans.
diffusa, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 95 = Vandellia pedunculata.
edentula, Benth. in DC. Prod. x. 410 = pedunculatis.
edentula, Griff. in Madr. Journ. Sc. Ser. I. iv. (1836) 376 = Vandellia mollis ?
erecta, Buch.-Ham. ex Wall. Cat. n. 3947 A = Vandellia erecta.
exappendiculata, Regel, Gartenfl. (1877) 34 ; et in Act. Hort. Petrop. v. (1877) 271 = pedunculatis.
fimbriata, Hook. ex R. Grah. in Edinb. N. Phil. Journ. (Jul.–Sept. 1831) 379 = Artanema fimbriatum.
flaccida, R. Br. Prod. 440 = Vandellia crustacea.
flava, *Buch.-Ham. ex Wall. Cat.* n. 3957 A, B.—As. trop.
flava, Wall. Cat. n. 3957 C = ciliaris.
Fordii, *Hook. f. Bot. Mag.* t. 6797 B.—China.
Fournieri, *Linden, ex Fourn. in Illustr. Hortic.* xxiii. (1876) 129. t. 249.—Cochinch.
globosa, Buch.-Ham. ex Wall. Cat. n. 3945 D = Vandellia scabra.
gracilis, Benth. in Wall. l. c. n. 3952 = Bonnaya veronicaefolia.
guinguenervis, Llanos, Fragm. 76 (err. typ.) = quinquenervis.
hians, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 96 = vagans.
hirsuta, *Willd. Sp. Pl.* iii. 266.—Ind. or.
hirta, Buch.-Ham. ex Wall. Cat. n. 3948 A = Vandellia mollis.
hirta, Cham. & Schlecht. in Linnaea, ii. (1827) 571 = Vandellia scabra.
hirtella, *Hook. f. Fl. Brit. Ind.* iv. 277.—Ind. or.
inflata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 192.—Japon.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 562.—Sumatra.
lanceolata, *Hook. f. Fl. Brit. Ind.* iv. 279 (sp. dub.).—Ind. or.

TORENIA:—
lepidota, Roth, Nov. Pl. Sp. 281 = Centranthera humifusa.
lobelioides, Hook. & Arn. Bot. Beech. Voy. 202 = oblonga.
longiflora, C. Morr. in Ann. Soc. Bot. Gand, ii. (1846) 441, 471 = concolor.
lucida, Buch.-Ham. ex Wall. Cat. n. 3962 = Vandellia crustacea.
minuta, Blume, Bijdr. 755 = Vandellia minuta.
molluginoides, Benth. in Wall. Cat. n. 3960 = Vandellia molluginoides.
mucronulata, *Benth. in DC. Prod.* x. 409.—Burma.
multiflora, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 96 = Vandellia multiflora.
nana, *Benth. in DC. Prod.* x. 412.—Afr. austr.
Nortenia, *Steud. Nom.* ed. II. ii. 692.—Madag. ; Bras.
oblonga, *Steud. l. c.* ; *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 227, in nota.—China.
obtusa, Blume, Bijdr. 754 = Vandellia obtusa.
obtusifolia, *Sm. in Rees, Cycl.* xxxvi. n. 5.—Ind. or.
ovata, Sweet, Hort. Brit. ed. I. 306 = Vandellia crustacea.
paniculata, *Blanco, Fl. Filip.* ed. I. 486.—Ins. Philipp.
parviflora, *Buch.-Ham. ex Wall. Cat.* n. 3958.—Amphig. trop.
pedunculatis, *Benth. in Wall. l. c.* n. 3956.—Ind. or. ; Malaya.
pedunculata, Willd. ex Steud. Nom. ed. II. ii. 692 = T. Nortenia.
plantaginea, Benth. in DC. Prod. x. 411 = Craterostigma plantagineum.
polygonoides, *Benth. Scroph. Ind.* 39.—Ind. or. ; Malaya.
pumila, Benth. in DC. Prod. x. 411 = Craterostigma pumilum.
quinquenervis, *Llanos, Fragm.* 70 (err. typ. guinguenervis).—Ins. Philipp.
racemosa, Benth. Scroph. Ind. 38 ; et in Wall. Cat. n. 3957 = ciliaris.
ramosissima, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 10.—Afr. trop.
Roxburghii, Buch.-Ham. ex Wall. Cat. n. 3856 A = Bonnaya reptans.
rubella, Buch.-Ham. l. c. n. 3962 A = Vandellia crustacea.
rubens, Benth. in DC. Prod. x. 410 = vagans.
rubens, Thw. Enum. Pl. Zeyl. 219 = hirtella.
scabra, R. Br. Prod. 440 = Vandellia crustacea.
scabra, R. Grah. in Bot. Mag. t. 3104 = Artanema fimbriatum.
Schweinfurthii, *Oliver, in Hook. Ic. Pl.* t. 1256.—Afr. trop.
sessiliflora, Benth. in Wall. Cat. n. 3959 = Vandellia sessiliflora.
Soriana, Blanco, Fl. Filip. ed. II. 340 = Vandellia Soriana.
stolonifera, *Boj. ex Benth. in DC. Prod.* x. 410.—Madag.
trichotoma, Buch.-Ham. ex Wall. Cat. n. 3943 A = Vandellia erecta.
vagans, *Roxb. Hort. Beng.* [95] ; *Fl. Ind.* iii. 96.—Ind. or. ; China.
varians, Griff. Notul. iv. 125 = Vandellia pedunculata.
varians, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 96 = Vandellia crustacea.
veronicaefolia, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 282.—Am. centr.
veronicaefolia, Willd. ex Steud. Nom. ed. II. ii. 692 = T. Nortenia.
veronicifolia, Herb. Madr. ex Wall. Cat. n. 3856 B = Bonnaya reptans.
viscosa, Sweet, Hort. Brit. ed. I. 306 = Vandellia hirsuta.

TORESIA, Pers. Syn. ii. 534 (1807) = TORRESIA, Ruiz & Pav. = **Hierochloë**, S. G. Gmel. (Gramin.).

TORFOSIDIS, Rafin. Fl. Tellur. iv. 112 (1836) = **Euphorbia**, Linn.
canariensis, Rafin. l. c. = E. canariensis.

TORICELLIA, DC. Prod. iv. 257 (1830). *CORNACEAE*, Benth. & Hook. f. i. 952.
tiliaefolia, *DC. l. c.*—Reg. Himal.

TORILIS, Adans. Fam. ii. 99 (1763) = **Caucalis**, Linn. (Umbell.).
africana, Spreng. Pugill. ii. 55 ; et in Schult. Syst. vi. 468 = C. capensis.
anthrisca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = C. Anthriscus.
Anthriscus, Bernh. Syst. Verz. Erf. 167 ; C. C. Gmel. Fl. Bad. i. 613 = C. Anthriscus.
Anthriscus, Gaertn. Fruct. 83. t. 20 ; Moench, Meth. 103 = Anthriscus vulgaris.
Anthriscus, Spreng. ex DC. Prod. iv. 219 = C. arvensis.
arvensis, Link, Enum. Hort. Berol. i. 265 = C. arvensis.
Borszczowi, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 600.—Turkestan.
bracteosa, Bianca, ex Nym. Consp. 282 = C. nodosa.
brevipes, Lowe, Man. Fl. Mad. i. 372.—Ins. Mader.
chlorocarpa, Spreng. Syst. i. 898 = C. infesta.
chrysocarpa, Boiss. & Blanche, Diagn. Ser. II. ii. 98 = C. fallax.
convexa, Dulac, Fl. Hautes-Pyr. 361 = C. Anthriscus.
crinita, Spreng. Sp. Umb. 141 = Daucus crinitus.
divaricata, Moench, Meth. Suppl. 34 = C. arvensis.
elata, Spreng. Syst. iv. Cur. Post. 119 ; DC. Prod. iv. 220 = C. Anthriscus.
erythrotricha, Reichb. f. Ic. Fl. Germ. xxi. 84 = C. erythrotricha.
Friedrichsthalii, Cesati, in Fried. Reise Neu-Griechenl. 280 = C. arvensis.
grandiflora, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 56 = *microcarpa*.
helvetica, C. C. Gmel. Fl. Bad. i. 617 = C. arvensis.
heterophylla, Guss. Fl. Sic. Prod. i. 326 = C. linearifolia.
heterotricha, Trautv. in Bull. Soc. Nat. Mosc. xxxix. (1866) 1. 329.—Turkestan.
infesta, Roth, Enum. Pl. Phan. Germ. i. 901 ; Reichb. Fl. Germ. Excurs. 447 = C. infesta.
infesta, Spreng. Pl. Umb. Prod. 24 = C. arvensis.
intermedia, Gren. in Mém. Soc. Emul. Doubs, i. 1. 'Tome 3me' (1842–43) 66 (Cat. Pl. Phan. Doubs, 34) = T. Anthriscus ?
japonica, DC. Prod. iv. 219 = C. Anthriscus.
lanuginosa, [Clairv.] Man. Herb. 78 = Anthriscus vulgaris.
leptophylla, Reichb. f. Ic. Fl. Germ. xxi. 83. t. 169 = C. leptophylla.
macrocarpa, Gaertn. Fruct. i. 83 = Chaerophyllum nodosum.
melanantha, Vatke, in Linnaea, xl. (1876) 190 = C. melanantha.
microcarpa, Bess. Enum. Pl. Volh. 43.—Rossia ; As. Min.
neglecta, Schult. Syst. vi. 484 = C. infesta.
nodosa, Gaertn. Fruct. i. 82. t. 20. f. 6 = C. nodosa.
nodosa, Kotschy, in Unger & Kotschy, Ins. Cypern, 306 = C. fallax.
obscura, Lowe, Man. Fl. Mad. i. 372.—Ins. Mader.
persica, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 102 = C. Anthriscus.
peruviana, Presl, ex DC. Prod. iv. 214 = Daucus montanus.
praetermissa, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 214 = C. Anthriscus.
purpurea, Guss. Fl. Sic. Prod. i. 325 = C. arvensis.
radiata, Moench, Meth. 103 = C. infesta.
rubella, Moench, l. c. = C. Anthriscus.
scabra, DC. Prod. iv. 219 = C. Anthriscus.
scandicina, Adans. ex Schult. Syst. vi. 524 = Anthriscus vulgaris.
setifolia, Boiss. Diagn. Ser. II. ii. 98 = Psammogeton setifolium.
stricta, Wibel, Prim. Fl. Werth. 192 = C. Anthriscus.
syriaca, Boiss. & Blanche, Diagn. Ser. II. ii. 98 = C. infesta.
tenella, Reichb. f. Ic. Fl. Germ. xxi. 84.—Europ.
tenuifolia, Lowe, Man. Fl. Mad. i. 370.—Ins. Mader.
trichosperma, Spreng. Sp. Umb. 142 = C. trichosperma.
trilobata, Viv. ex DC. Prod. iv. 121 = Pimpinella peregrina.
triradiata, Boiss. & Heldr. Diagn. Ser. I. x. 48.—Pamphylia.
tuberculata, Spreng. in Schult. Syst. vi. 426 = Pimpinella peregrina.
tumida, Moench, Meth. 102 = Chaerophyllum nodosum.
ucranica, Spreng. in Schult. Syst. vi. 485 = *microcarpa*.

TORMENTILLA, [Tourn.] Linn. Syst. ed. I (1735) = **Potentilla**, Linn. (Rosac.).
adstringens, Lindem. in Bull. Soc. Nat. Mosc. xxiii. (1850) II. 480 = P. Tormentilla.
alpina, Opiz, Boehm. Gew. (1823) 64 = P. aurea ?
Anserina, Stokes, Bot. Mat. Med. iii. 155 = P. Anserina.
dissecta, Timb. in Bull. Soc. Bot. Fr. xix. (1872) p. cxviii. = P. Tormentilla ?
divergens, Reichb. ex Bot. Zeit. i. (1843) 263 = P. Tormentilla.
erecta, Linn. Sp. Pl. 500 = P. Tormentilla.
ericetorum, Timb. in Bull. Soc. Bot. Fr. xix. (1872) p. cxviii; et in Bull. Soc. Sc. Phys. et Nat. Toul. vi. (1883–84) 114 = P. Tormentilla.
fruticosa, Stokes, Bot. Mat. Med. iii. 154 = P. fruticosa.
humifusa, G. Don, Gen. Syst. ii. 562 = P. humifusa.
montana, Schur, Enum. Pl. Transs. 188 = P. Tormentilla.
nodosa, Schur, l. c. = P. Tormentilla.
obovatifolia, Stokes, Bot. Mat. Med. iii. 152 = P. Fragariastrum.
officinalis, Curt. Fl. Lond. fasc. v. t. 35 = P. Tormentilla.
orophila, Timb. in Bull. Soc. Bot. Fr. xix. (1872) p. cxvii. = P. Tormentilla.
parviflora, Wallr. in Linnaea, xiv. (1840) 580 (Beitr. Fl. Hercyn. 210) = P. Tormentilla.
pusilla, Willd. ex Steud. Nom. ed. II. ii. 693 = P. elegans.
reptans, Linn. Sp. Pl. 500 = P. procumbens.
reptans, Stokes, Bot. Mat. Med. iii. 153 = P. reptans.
sessilifolia, Stokes, l. c. 151 = P. Tormentilla.
tuberosa, Reyn. Fl. Dép. Orne, 148 = P. Tormentilla.

TORMINALIS, Medic. Phil. Bot. i. 134 (1789) = **Pyrus**, Tourn. (Rosac.).

TORMINARIA, M. Roem. Syn. Rosifl. 130 (1847) = **Pyrus**, Tourn. (Rosac.).
Clusii, M. Roem. l. c. = P. Torminalis.
florentina, M. Roem. l. c. 131 = P. crataegifolia.
vulgaris, Schur, Enum. Pl. Transs. 207 = P. Torminalis.

TORNABENEA, Parl. ex Webb, in Hook. Kew Journ. ii. (1850) 370. *UMBELLIFERAE*, Benth. & Hook. f. i. 930.
TETRAPLEURA, Parl. in Hook. Niger Fl. 131 (1849).
Bischoffii, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 254.—Ins. Cap. Virid.
hirta, *J. A. Schmidt, l. c.* 253.—Ins. Cap. Virid.
insularis, *Parl. ex Webb, in Hook. Kew Journ.* ii. (1850) 370.—Ins. Cap. Virid.

TORNABENIA, Benth. & Hook. f. Gen. i. 930 (1867), sphalm. = praec.

TORNELIA, Gutierrez, ex Schlecht. in Linnaea, xxvi. (1853) 382 = **Monstera**, Adans. (Aroid.).
dilacerata, Schott, Prod. Aroid. 356 = M. dilacerata.
dissecta, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = M. dilacerata.
fragrans, Gutierrez, ex Schlecht. in Linnaea, xxvi. (1853) 382 = M. deliciosa.
laniata, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = M. pertusa.
Lindeni, Schott, ex Engl. in DC. Monog. Phan. ii. 259 = M. Friedrichsthalii.
Spruceana, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 40 = M. Spruceana.
subpinnata, Schott, Prod. Aroid. 357 = M. subpinnata.

TORPESIA, M. Roem. Syn. Hesper. 86, 116 (1846) = **Trichilia**, Linn. (Meliac.).
mucronata, M. Roem. l. c. 116 = Trich. heterophylla.

TORRENTIA, Vell. Fl. Flum. viii. t. 149 (1827) = **Ichthyothere**, Mart. (Compos.).
quinquenervis, Vell. l. c. = I. Cunabi.

TORRESIA, Ruiz & Pav. Prod. 125 (1794) = **Hierochloë**, S. G. Gmel. (Gramin.).
antarctica, Beauv. Agrost. 63 = H. antarctica.
biflora, Roem. & Schult. Syst. ii. 515 = Centotheca lappacea.
latifolia, Beauv. ex Roem. & Schult. l. c. = Centotheca lappacea.

TORRESIA:—
magellanica, Beauv. Agrost. 63 = Hierochloë magellanica.
redolens, Roem. & Schult. Syst. ii. 516 = H. redolens.
utriculata, Ruiz & Pav. Syst. Veg. 251 = H. utriculata.

TORREYA, Arn. in Ann. Nat. Hist. Ser. I. i. (1838) 130. *CONIFERAE*, Benth. & Hook. f. iii. 431.
CARYOTAXUS, Zucc. ex Endl. Syn. Conif. 240 (1847).
FOETATAXUS, [Nelson,] Senilis, Pinaceae 167 (1866).
bogotensis, *Linden, Cat.* n. 25 (1870) 12; *et ex Thurb. in Bull. Torrey Bot. Club*, i. (1870) 7 (sp. dub.).—N. Granat.
californica, *Torr. in N. York Journ. Pharm.* iii. (1852) 49.—Calif.
grandis, *Fortune, ex Gord. Pinet.* ed. I. 326.—China.
Humboldtii, Hort. Kew. ex Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 226 = Podocarpus taxifolia.
montana, Hort. ex Lindl. & Gord. l. c. 229 = taxifolia.
Myristica, Hook. Bot. Mag. t. 4780 = californica.
nucifera, *Sieb. & Zucc. in Abh. Akad. Muench.* iii. IV. (1846) 234.—Japon.
taxifolia, *Arn. in Ann. Nat. Hist.* Ser. I. i. (1838) 130.—Florida.

TORREYA, Croom, ex Meissn. Gen. Comm. 340 (1843) = **Croomia**, Torr. (Roxburgh.).

TORREYA, Rafin. in Am. Monthly Mag. (1818) 356 = **Synandra**, Nutt. (Labiat.).
grandiflora, Rafin. l. c.; et (1819) 194 = S. grandiflora.

TORREYA, Rafin. in Journ. Phys. lxxxix. (1819) 105 = **Cyperus**, Mich.
caespitosa, Rafin. l. c. = C. Nuttallii.
maritima, Rafin. l. c. (Pycrei sp.).—Am. bor.

TORREYA, Spreng. Neue Entdeck. ii. 121 (1821) = **Clerodendron**, Linn. (Verben.).
paniculata, Spreng. l. c.—Bras.

TORRUBIA, Vell. Fl. Flum. 139 (1825); iii. t. 150 = **Pisonia**, Plum. (Nyctag.).
opposita, Vell. ll. cc.—Bras.

TORRULIA, Steud. Nom. ed. II. ii. 693 (1841), sphalm. = praec.

TORTULA, Roxb. ex Willd. Sp. Pl. iii. 359 (1800) = **Priva**, Adans. (Verben.).
aspera, Roxb. ex Willd. l. c. = P. leptostachya.

TORULINIUM, Desv. in Ham. Prod. Pl. Ind. Occ. 15 (1825) = **Mariscus**, Ehrh. (Cyperac.).
confertum, Desv. l. c. (= *Cyperus ferax*).—Ind. occ.
ferox, Ham. l. c. (= *Cyperus ferax*).—Ind. occ.

TORYMENES, Salisb. in Trans. Hort. Soc. i. (1812) 283 = **Amomum**, Linn. (Scitam.).
officinalis, Salisb. l. c. = A. Granum-Paradisi.

TOSAGRIS, Beauv. Agrost. 29. t. 8. f. 3 (1812) = **Muehlenbergia**, Schreb. (Gramin.).
agrostidea, Beauv. l. c. = M. capillaris.

TOUCHARDIA, Gaudich. Bot. Voy. Bonite, t. 94 (1854?). *URTICACEAE*, Benth. & Hook. f. iii. 388.
candidissima, Blume, Mus. Bot. Lugd. Bat. ii. 165 = Leucosyke javanica.
latifolia, *Gaudich. Bot. Voy. Bonite*, t. 94.—Ins. Sandvic.

TOUCHIROA, Aubl. Pl. Gui. i. 384. t. 148 (1775) = **Crudia**, Schreb. (Legumin.).
aromatica, Aubl. l. c. 385 = C. aromatica.
bantamensis, Hassk. Retzia, i. 202 (= *Pryona bantamensis*).—Java.
Parivoa, Rich. ex DC. Prod. ii. 520 = C. Parivoa.
Vouarana, Rich. l. c. = Cupania Voua-rana.

TOULICHIBA, Adans. Fam. ii. 326 (1763) = **Ormosia**, Jack (Legumin.).

TOULICIA, Aubl. Pl. Gui. i. 359. t. 140 (1775). *SAPINDACEAE*, Benth. & Hook. f. i. 393.
DICRANOPETALUM, Presl, Bot. Bemerk. 24 (1844); (in Abh. Boehm. Ges. V. iii. (1845) 454).
PONAEA, Schreb. Gen. 266 (1789).
brasiliensis, *Casar. Nov. Stirp. Bras. Dec.* 45.—Bras.
bullata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 372.—Bras.
crassifolia, *Radlk. l. c.* 373.—Bras.
elliptica, *Radlk. l. c.* 371.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 359. t. 140. f. 3.—Guian.
laevigata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 372.—Bras.
megalocarpa, *Radlk. Monog. Serj.* 353.—Venezuela.
pulvinata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 371.—Guian.
stans, *Radlk. Monog. Serj.* 353.—Bras.
tomentosa, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 373.—Bras.

TOULOUCOUNA, M. Roem. Syn. Hesper. 123 (1846) = **Carapa**, Aubl. (Meliac.).
gigantea, M. Roem. l. c. = C. procera.

TOUMBOA, Naud. in Rev. Hortic. (1862) 186 = TUMBOA, Welw. = **Welwitschia**, Hook. f. (1862) (Gnetac.).

TOUNATEA, Aubl. Pl. Gui. i. 549. t. 218 (1775) = **Swartzia**, Schreb. (Legumin.).
guianensis, Aubl. l. c. 550 = S. alata.
madagascariensis, Baill. in Bull. Soc. Linn. Par. (1885) 446 = S. madagascariensis.

TOURNEFORTIA, Linn. Syst. ed. I (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 843.
ARGUSIA, Amm. ex Linn. Hort. Upsal. 36 (1748).
ARGUZIA, Rafin. Sylva Tellur. 167 (1838); Stev. in Bull. Soc. Nat. Mosc. (1851) I. 558.
MESSERSCHMIDIA, Murr. Syst. ed. XIII. 161 (1774).
MESSERSMIDIA, Linn. Hort. Upsal. 36 (1748).
OSKAMPIA, Rafin. Sylva Tellur. 123 (1838).
PITTONIA, Plum. ex Adans. Fam. ii. 177 (1763).
TETRANDRA, Miq. Fl. Ind. Bat. ii. 929 (1859).
VERRUCARIA, Medic. Malv. 103 (1787).
acclinis, F. Muell. Fragm. iv. 95 = sarmentosa.
acuminata, *DC. Prod.* ix. 520.—Ins. Mascar.
acutiflora, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 333.—Mexic.
acutifolia, Willd. ex Roem. & Schult. Syst. iv. 540 = caracasana.
alba, *Splitg. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 347.—Guian.
angulosa, Desf. Tabl. ed. II. 85, nomen (Quid ?).—Hab. ?
angustiflora, *Ruiz & Pav. Fl. Per.* ii. 25. t. 151.—Peruv.
angustifolia, Roem. & Schult. Syst. iv. 539 = T. Messerschmidia.
arborea, Blanco, Fl. Filip. ed. I. 129 = argentea.
arborescens, Lam. Illustr. i. 417 (Quid ?).—Ind. or.
argentea, *Linn. f. Suppl.* 133.—Ind. or.; Malaya; Austral.
Arguzia, Roem. & Schult. Syst. iv. 540 = sibirica.
asperrima, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 333.—Mexic.
astrotricha, *DC. Prod.* ix. 520.—Jamaic.
bicolor, *Sw. Prod. Veg. Ind. Occ.* 40.—Am. trop.
bifida, Boj. Hort. Maurit. 234 = Bojeri.
bifida, *Lam. Illustr.* i. 417.—Ins. Maurit.
Billbergiana, *Beurl. in Vet. Acad. Handl. Stockh.* 1854 (1856) 139.—Panama.
Blanchetii, *A. DC. in DC. Prod.* ix. 524.—Bras.
Bojeri, *A. DC. l. c.* 516.—Ins. Maurit.
brachiata, *A. DC. l. c.* 525.—Bras.
brasiliensis, *Poir. Encyc.* v. 357.—Bras.
breviflora, *DC. Prod.* ix. 520.—Am. trop.
calycina, *Benth. Bot. Voy. Sulph.* 139.—Mexic.
candida, *Walp. Rep.* vi. 556.—Mexic.
Candollii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 146.—Reg. Himal.
canescens, *H. B. & K. Nov. Gen. et Sp.* iii. 82.—Am. austr.
capitata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 332.—Mexic.
caracasana, *H. B. & K. Nov. Gen. et Sp.* iii. 80.—Venezuela.
caribaea, *Griseb. Fl. Brit. W. Ind.* 484.—Ind. occ.
carnosa, *Mill. Gard. Dict.* ed. VIII. n. 6.—N. Granat.

TOURNEFORTIA:—
chamaedrifolia, Sm. in Rees, Cycl. xxxvi. n. 2 = Cordia mirabiliflora.
chrysantha, *Walp. Rep.* vi. 556.—Mexic.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 84.—N. Granat.
corymbosa, Sieber, ex DC. Prod. ix. 517 = hirsutissima.
corymbosa, Willd. ex Roem. & Schult. Syst. iv. 541 = foetidissima.
cuspidata, *H. B. & K. Nov. Gen. et Sp.* iii. 83.—Ecuador.
cymosa, Griseb. in Goett. Abh. vii. (1857) 253; cf. ej. Fl. Brit. W. Ind. 483 = filiflora.
cymosa, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 145 = Heyneana.
cymosa, Jacq. Coll. i. 96 = astrotricha.
cymosa, *Linn. Sp. Pl.* ed. II. 202.—Ind. occ.
densiflora, *Mart. ex Gal. in Bull. Acad. Brux.* xi. II. (1844) 333.—Mexic.
dichotoma, [Moç. & Sesse,] ex DC. Prod. ix. 518 = foetidissima.
difformis, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 206.—Ins. Galapag.
Edgeworthii, A. DC. in DC. Prod. ix. 529 = Heliotropium zeylanicum.
elegans, *Cham. in Linnaea*, iv. (1829) 469.—Bras.
elliptica, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 332.—Mexic.
ferruginea, Lam. Illustr. i. 417 = scandens.
filiflora, *Griseb. Fl. Brit. W. Ind.* 483.—Ind. occ.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 79.—N. Granat.
foetida, Pers. Syn. i. 165 = foetidissima.
foetidissima, Descourt. ex Griseb. Fl. Brit. W. Ind. 483 = cymosa.
foetidissima, *Linn. Sp. Pl.* 140.—Ind. occ.; Mexic.
frangulaefolia, Zipp. ex Span. in Linnaea, xv. (1841) 334 = sarmentosa.
fruticosa, [Ker-Gawl.] Bot. Reg. t. 464 = T. Messerschmidia.
fruticosa, *Ortega, Hort. Matr. Dec.* 57.—Mexic.
fuliginosa, *H. B. & K. Nov. Gen. et Sp.* iii. 81.—Am. austr.
Gardneri, *A. DC. in DC. Prod.* ix. 526.—Bras.
glaberrima, *Salzm. ex DC. l. c.* 519.—Bras.
glabra, Linn. Sp. Pl. 141 = cymosa.
glabra, *Aubl. Pl. Gui.* i. 118; *Spreng. Neue Entdeck.* ii. 124.—Guian.
glabra, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 5.—Java.
gnaphalodes, *R. Br. Prod.* 496.—Am. trop.
grandifolia, *Fresen. in Mart. Fl. Bras.* viii. 56.—Bras.
Hartwegiana, *Steud. Nom.* ed. II. ii. 693.—Mexic.
heliotropioides, Hook. Bot. Mag. t. 3096 = Heliotropium anchusaefolium.
Hernandesii, *Dun. ex DC. Prod.* ix. 529.—Mexic.
Heyneana, DC. Prod. ix. 516 = Candollii.
Heyneana, *Wall. Cat.* n. 910.—Ind. or.
hirsutissima, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 140 = Billbergiana.
hirsutissima, Blanco, Fl. Filip. ed. I. 128 = sarmentosa.
hirsutissima, *Linn. Sp. Pl.* 140.—Am. trop.
hispida, *H. B. & K. Nov. Gen. et Sp.* iii. 82.—Am. austr.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 147.—Reg. Himal.
Horsfieldii, *Miq. Fl. Ind. Bat.* ii. 927.—Java.
Hostmanni, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1151, nomen.—Guian.
humilis, Linn. Sp. Pl. 141 = Heliotropium fruticosum.
incana, G. Don, Gen. Syst. iv. 368 = Meyeri.
incana, *Lam. Illustr.* i. 417.—Ind. occ.
intermedia, *Fresen. in Mart. Fl. Bras.* viii. 50.—Bras.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 147.—Reg. Himal.
laevigata, *Lam. Illustr.* i. 416.—Am. trop.
laevigata, Sm. in Rees, Cycl. xxxvi. n. 18 = volubilis.
lanceolata, *Fresen. in Mart. Fl. Bras.* viii. 55.—Bras.
latifolia, D. Dietr. Syn. Pl. i. 629 = laurifolia.
latifolia, Willd. in Roem. & Schult. Syst. iv. 540 = laevigata.
laurifolia, Poepp. ex Cham. & Schlecht. in Linnaea, iv. (1829) 465 = laevigata.
laurifolia, *Vent. Choix Pl.* 2.—Am. trop.

TOURNEFORTIA:—
laxiflora, DC. Prod. ix. 525 = floribunda.
leptostachya, *Benth. Bot. Voy. Sulph.* 140.—N. Granat.?
leucophylla, *H. B. & K. Nov. Gen. et Sp.* iii. 85.—N. Granat.
linearis, E. Mey. in Drège, Zwei Pfl. Docum. 226 (= *Heliophytum lineare*).—Afr. austr.
longifolia, *Ruiz & Pav. Fl. Per.* ii. 25. t. 150.—Peruv.
loxensis, *H. B. & K. Nov. Gen. et Sp.* iii. 81.—Ecuador.
lucida, Desf. Cat. Hort. Par. ed. III. 397 = volubilis.
lucida, *Desv. ex Ham. Prod. Pl. Ind. Occ.* 24.—Ind. occ.
macroloba, *DC. Prod.* ix. 527.—Bras.
macrophylla, Lam. Illustr. i. 416 = cymosa, foetidissima.
maculata, *Jacq. Enum. Pl. Carib.* 14.—Am. austr.
maritima, *Salzm. ex DC. Prod.* ix. 526.—Bras.
Martii, *Fresen. in Mart. Fl. Bras.* viii. 55.—Bras.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 66.—Reg. Argent.
melanochaeta, *DC. Prod.* ix. 520.—Guian.
membranacea, DC. l. c. 530 = scandens.
Messerschmidia, *Sweet, Hort. Suburb. Lond.* 31.—Ins. Canar.
mexicana, *Vatke, Ind. Sem. Hort. Berol.* (1876) *App.* 1.—Mexic.
Meyeri, *DC. Prod.* ix. 530.—Guian.
micranthos, A. DC. l. c. x. 67 = Heliotropium micranthum.
microphylla, Bert. ex Steud. Nom. ed. II. ii. 693 = scandens.
microphylla, Desv. in Ham. Prod. Pl. Ind. Occ. 24 = volubilis.
microstachya, Roem. & Schult. Syst. iv. 538 = Heliotropium microstachyum.
minuta, Bert. ex Spreng. Syst. i. 644 = incana.
mollis, Bertol. Misc. Bot. xii. 44, cum tab. (= Vangueriae sp.?).—Afr. trop.
mollis, *A. Gray, in Proc. Am. Acad.* x. (1875) 50.—Am. bor.
mollis, *F. Muell. Fragm.* i. 59.—Austral.
Monclovana, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 120.—Mexic.
monostachya, Willd. ex Roem. & Schult. Syst. iv. 540 = Heliotropium strictum.
montana, *Lour. Fl. Cochinch.* 122.—Cochinch.
mutabilis, *Vent. Choix Pl.* t. 3.—Java; Mexic.?
nitida, *H. B. & K. Nov. Gen. et Sp.* iii. 84.—N. Granat.
obscura, *A. DC. in DC. Prod.* ix. 517.—Guian.
obtusiflora, *Benth. Bot. Voy. Sulph.* 140.—Ecuador.
octostachya, *DC. Prod.* ix. 518.—N. Granat.
opaca, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 205.—Ins. Galapag.
orientalis, R. Br. Prod. 497 = sarmentosa.
ovata, *Wall. Cat.* n. 908.—Burma; China.
paniculata, *Cham. in Linnaea*, iv. (1829) 468.—Bras.
peruviana, *Poir. Encyc. Suppl.* iv. 425.—Peruv.
petiolaris, *DC. Prod.* ix. 520.—Mexic.
Pohlii, *Fresen. in Mart. Fl. Bras.* viii. 52.—Bras.
poliochros, *Spreng. Syst.* i. 644.—Jamaic.
polystachya, Poepp. ex DC. Prod. ix. 492 = Cordia Poeppigii.
polystachya, *Ruiz & Pav. Fl. Per.* ii. 24. t. 149.—Peruv.
Pseudo-heliotropium, *Lehm. in E. Otto, Hamb. Gartenz.* vi. (1850) 456.—Hab.?
psilostachya, *H. B. & K. Nov. Gen. et Sp.* iii. 78.—Am. austr.
puberula, *Baker, in Journ. Linn. Soc.* xx. (1883) 211.—Madag.
pubescens, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 198.—Ins. Galapag.
punctata, *Lem. Jard. Fleur.* iv. (1854) *Misc.* 79.—Hab.?
punctata, Spreng. Syst. i. 648 = volubilis.
pyrrhotricha, *Fresen. in Mart. Fl. Bras.* viii. 53.—Bras.
reticosa, *Wight, Ic.* t. 1386.—Ind. or.
rosmarinifolia, Steud. Nom. ed. II. ii. 693 = sibirica.
Roxburghii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 146.—Ind. or.
Royleana, DC. Prod. ix. 527 = Heliotropium zeylanicum.

TOURNEFORTIA:—
rubicunda, *Salzm. ex DC. Prod.* ix. 526.—Bras.
rufescens, *Willd. ex Roem. & Schult. Syst.* iv. 541.—Am. austr.
rufipila, *Benth. Bot. Voy. Sulph.* 140.—Mexic.
rufo-sericea, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 197.—Ins. Galapag.
rugosa, Willd. in Roem. & Schult. Syst. iv. 540 = fuliginosa.
Sagraeana, DC. Prod. ix. 522 = laurifolia.
salicifolia, *DC. l. c.* 530.—Bras.
Salzmanni, *DC. l. c.* 524.—Bras.
Sampsoni, Hance, in Journ. Bot. vi. (1868) 330 = ovata.
sarmentosa, *Lam. Illustr.* i. 416.—As. et Austral. trop.; Ins. Mascar.
scabra, *Lam. l. c.* 417.—Ins. S. Doming.
scabrida, *H. B. & K. Nov. Gen. et Sp.* iii. 83.—Am. austr.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 4.—Jamaic.
scandens, Willd. Enum. Hort. Berol. 188 = peruviana.
Schiedeana, *G. Don, Gen. Syst.* iv. 368.—Mexic.
Schomburgkii, *DC. Prod.* ix. 517.—Guian.
sericea, Vahl, Eclog. Am. i. 17 = tomentosa?
serrata, Linn. Sp. Pl. 140 = Cordia mirabiliflora.
sessilifolia, *Poir. Encyc.* v. 360.—Reg. Argent.
sibirica, *Linn. Sp. Pl.* 141.—Ross.; Sibir.
spigeliaeflora, *A. DC. in DC. Prod.* ix. 525.—Guian.
staminea, *Griseb. Fl. Brit. W. Ind.* 484.—Ind. occ.
stenoraca, *Klotzsch, in Peters, Reise Mossamb. Bot.* 250.—Afr. trop.
strigosa, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 207 = psilostachya.
suaveolens, *H. B. & K. Nov. Gen. et Sp.* iii. 81.—Peruv.
subsessilis, *Cham. in Linnaea*, viii. (1833) 119.—Bras.
subulata, Hochst. ex DC. Prod. ix. 528 = Heliotropium zeylanicum.
suffruticosa, Linn. Sp. Pl. ed. II. 202 = incana.
surinamensis, A. DC. in DC. Prod. ix. 526 = laurifolia.
synzystachya, Roem. & Schult. Syst. iv. 539 = Heliotropium synzystachyum.
syringaefolia, *Miq. Stirp. Surin. Sel.* 137. t. 41.—Guian.
syringaefolia, Vahl, Symb. Bot. iii. 23 = laurifolia.
tenuiflora, Willd. ex Roem. & Schult. Syst. iv. 540 = Declieuxia herbacea.
tenuifolia, Steud. Nom. ed. II. ii. 694, sphalm. = ternifolia.
ternata, *Wall. Cat.* n. 912.—Burma.
ternifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 84.—Am. austr.
tetragona, D. Dietr. Syn. Pl. i. 629, sphalm. = seq.
tetrandra, *Blume, Bijdr.* 845.—Ins. Molucc.
tetrandra, Wall. Cat. n. 911 = Wallichii.
tomentosa, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. trop.
trichocalycina, *DC. Prod.* ix. 517.—Mexic.
tristachya, Willd. ex Roem. & Schult. Syst. iv. 540 = psilostachya.
tuberculosa, *Cham. in Linnaea*, iv. (1829) 467.—Afr. austr.
umbellata, *H. B. & K. Nov. Gen. et Sp.* iii. 79.—Mexic.
undulata, Benth. Pl. Hartw. 20 = Hartwegiana.
undulata, *Ruiz & Pav. Fl. Per.* ii. 25.—Venezuela.
Urvilleana, *Cham. in Linnaea*, iv. (1829) 465.—Ins. Philipp.
Vauthieri, *DC. Prod.* ix. 526.—Bras.
velutina, *H. B. & K. Nov. Gen. et Sp.* iii. 79. t. 201.—Mexic.
velutina, Sm. in Rees, Cycl. xxxvi. n. 13 = argentea.
villosa, *Salzm. ex DC. Prod.* ix. 524.—Bras.
virgata, *Ruiz & Pav. Fl. Per.* ii. 25.—Peruv.
viridiflora, Gamble, Darjeeling List, 57 = Hookeri.
viridiflora, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 5; Cat. n. 907 = sarmentosa.
volubilis, *Linn. Sp. Pl.* 140.—Am. trop.
volubilis, Ruiz & Pav. Fl. Per. ii. 24 = peruviana.
vulgaris, Salzm. ex DC. Prod. ix. 524 = Salzmanni.
Walkerae, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 147.—Zeylan.
Wallichii, *DC. Prod.* ix. 527.—Malaya.
Wallichii, Thw. Enum. Pl. Zeyl. 214 = Walkerae.
Wightii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 146.—Ind. or.
zeylanica, Wight, Illustr. t. 170 = Heliotropium zeylanicum.

TOURNESOL, Adans. Fam. ii. 356 (1763) = seq.

TOURNESOLIA, Nissol. ex Scop. Introd. 243 (1777) = **Chrozophora**, Neck. (1790) (Euphorb.).

TOURNEUXIA, Coss. in Bull. Soc. Bot. Fr. vi. (1859) 395; et in Ann. Sc. Nat. Sér. IV. xviii. (1862) 211. t. 13. *COMPOSITAE*, Benth. & Hook. f. ii. 532.
variifolia, *Coss. ll. cc.* 396; 212.—Alger.

TOURNONIA, Moq. in DC. Prod. xiii. II. 225 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 77.
Hookeriana, *Moq. l. c.*—N. Granat.
leptostachys, Moq. l. c. = Boussingaultia baselloides.

TOUROLIA, Stokes, Bot. Mat. Med. iii. 104 (1812) = TOUROULIA, Aubl. = **Quiina**, Aubl. (Guttif.).

TOUROUBEA, Steud. Nom. ed. II. ii. 694 (1841), sphalm. = **Souroubea**, Aubl. (Ternstr.).

TOUROULIA, Aubl. Pl. Gui. i. 492. t. 194 (1775) = **Quiina**, Aubl. (Guttif.).
guianensis, Aubl. l. c. = Q. guianensis.
solitaria, Stokes, Bot. Mat. Med. iii. 105 = Q. guianensis.

TOURRETIA, Fouger. in Mém. Acad. Sc. Par. 1784 (1787) 205. t. 1. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1049.
DOMBEYA, L'Hérit. Stirp. Nov. 33. t. 17 (1784).
MEDICA, Cothen. Disp. Veg. 7 (1790).
lappacea, Willd. Sp. Pl. iii. 263 = volubilis.
scaber, Domb. ex DC. Prod. ix. 239 = Eccremocarpus scaber.
volubilis, *J. F. Gmel. Syst.* 940.—Peruv.

TOURRETTIA, DC. Prod. ix. 236 (1845), sphalm. = praec.

TOVARA, Adans. Fam. ii. 276 (1763) = **Polygonum**, Tourn.
virginiana, Rafin. Fl. Tellur. iii. 12 = P. virginianum.

TOVARIA, Neck. Elem. iii. 190 (1790) = **Smilacina**, Desf. (Liliac.).
davurica, Baker, in Journ. Linn. Soc. xiv. (1875) 567 = S. dahurica.
flexuosa, Baker, l. c. = S. flexuosa.
fusca, Baker, l. c. 568 = S. fusca.
japonica, Baker, l. c. 570 = S. japonica.
laxiflora, Baker, l. c. 569 = S. laxiflora.
nervulosa, Baker, l. c. = S. nervulosa.
oleracea, Baker, l. c. = S. oleracea.
oligophylla, Baker, l. c. 565.—Reg. Himal.
pallida, Baker, l. c. 566 = S. purpurea.
paniculata, Baker, l. c. 568 = S. paniculata.
purpurea, Baker, l. c. 566 = S. purpurea.
racemosa, Neck. Elem. iii. 190 = S. racemosa.
Rossii, Baker, in Journ. Linn. Soc. xvii. (1879) 387 = S. japonica.
Salvini, Baker, l. c. xiv. (1875) 567 = S. Salvini.
scilloides, Baker, l. c. = S. scilloidea.
sessilifolia, Baker, l. c. 566 = S. sessilifolia.
stellata, Neck. Elem. iii. 190 = S. stellata.
thyrsoidea, Baker, in Journ. Linn. Soc. xiv. (1875) 568 = S. thyrsoidea.
trifolia, Neck. Elem. iii. 190 = S. trifolia.

TOVARIA, Ruiz & Pav. Prod. 49. t. 8 (1794). *CAPPARIDEAE*, Benth. & Hook. f. i. 110.
BANCROFTIA, Macfad. Fl. Jamaic. i. 112 (1837).
pendula, *Ruiz & Pav. Fl. Per.* iii. 373. t. 309.—Peruv.; Ind. occ.

TOVOMIA, Pers. Syn. ii. 68 (1807) = seq.

TOVOMITA, Aubl. Pl. Gui. ii. 956. t. 364 (1775). *GUTTIFERAE*, Benth. & Hook. f. i. 173.
BEAUHARNOISIA, Ruiz & Pav. in Ann. Mus. Par. xi. (1808) 71. t. 9.
EUTHALES, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iii. 307 (1817).
MARIALVA, Vand. Fl. Lusit. 37 (1788).
MICRANTHERA, Choisy, in Mém. Soc. Hist. Nat. Par. i. (1823) 224. tt. 11, 12.

TOVOMITA:—
amazonica, *Poepp. & Endl. ex Walp. Rep.* i. 392.—Bras.; Ind. occ.
brasiliensis, *Mart. Nov. Gen. et Sp.* ii. 83.—Bras.
calyptrata, Schlecht. in Linnaea, viii. (1838) 187 = Rheedia calyptrata.
Choisyana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 281.—Guian.
clusiaefolia, *G. Don, Gen. Syst.* i. 617.—Guian.
clusioides, Cambess. in Mém. Mus. Par. xvi. (1828) 419 = Richardiana.
clusioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 166.—Cuba.
foliosa, Presl, Symb. Bot. ii. 20 = paniculata.
fructipendula, Cambess. in Mém. Mus. Par. xvi. (1828) 419 = brasiliensis.
fructipendula, *G. Don, Gen. Syst.* i. 617.—Peruv.
Goudotiana, Cambess. ex C. Muell. in Walp. Ann. vii. 355 = Ochrocarpus Goudotianus.
gracilipes, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 280.—Bras.
Grisebachiana, *Planch. & Triana, l. c.* 284.—Cuba.
guianensis, *Aubl. Pl. Gui.* ii. 957.—Guian.
hameliaefolia, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 996, nomen = Schomburgkiana?
havetioides, *Griseb. Fl. Brit. W. Ind.* 106.—Jamaic.
laurina, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 282.—Bras.
leucantha, *Planch. & Triana, l. c.* 278.—Bras.
ligustrina, Poit. ex Planch. & Triana, l. c. 270 = brasiliensis.
macrophylla, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 996, nomen.—Guian.
macrophylla, *Walp. Rep.* i. 392.—Bras.
madagascariensis, Cambess. in Mém. Mus. Par. xvi. (1828) 419 = Ochrocarpus madagascariensis.
microcarpa, *Walp. Rep.* i. 392.—Peruv.
myriandra, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 368.—Guian.
nigrescens, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 276.—Guian.
paniculata, *A. St. Hil. Fl. Bras. Mer.* i. 315.—Bras.
pentapetala, Blanco, Fl. Filip. ed. I. 432 = Ochrocarpos pentapetalus.
Plumieri, *Griseb. Fl. Brit. W. Ind.* 106.—Ins. Dominic.
pyrifolia, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 280 = macrophylla.
Richardiana, *Planch. & Triana, l. c.* 273.—Guian.
rubella, *Spruce, ex Planch. & Triana, l. c.* 275.—Bras.
Schomburgkiana, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 996, nomen (an seq.?).—Guian.
Schomburgkii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 274.—Guian.
secunda, Poepp. ex Planch. & Triana, l. c. 271 = brasiliensis.
Spruceana, *Planch. & Triana, l. c.* 277.—Am. austr.
stigmatosa, *Planch. & Triana, l. c.* 275.—N. Granat.
stylosa, *Hemsl. Diagn. Pl. Nov.* 3.—Panama.
tenuiflora, *Benth. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 272.—Bras.
turbinata, *Planch. & Triana, l. c.* 283.—N. Granat.
umbellata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 367.—Guian.
uniflora, *G. Don, Gen. Syst.* i. 617.—Guian.
Weddeliana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 277.—Bolivia.

TOVOMITOPSIS, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 261 = **Chrysochlamys**, Poepp. (Guttif.).
costaricana, Oerst. ex Planch. & Triana, l. c. 265 = C. costa-ricana.
cuneata, Planch. & Linden, l. c. 264.—N. Granat.
glauca, Oerst. ex Planch. & Triana, l. c. = C. glauca.
nicaraguensis, Oerst. ex Planch. & Triana, l. c. 266 = C. nicaraguensis.
paniculata, Planch. & Triana, l. c. 262 = Tovomita paniculata.
psychotriaefolia, Oerst. ex Planch. & Triana, l. c. 263 = C. psychotriaefolia.

TOWNSENDIA, Hook. Fl. Bor. Am. ii. 16. t. 119 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 268.
arizonica, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 85.—N. Mexic.
condensata, *Parry, in Am. Natural.* viii. (1874) 106, 213.—Am. bor. occ.

TOWNSENDIA:—
eximia, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 70.—N. Mexic.
Fendleri, *A. Gray, l. c.*—N. Mexic.
florifer, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 84.—Am. bor. occ.
Fremontii, Torr. & Gray, in Bost. Journ. Nat. Hist. v. (1847) 106 = incana.
glabella, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 86.—Am. bor. occ.
grandiflora, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 306.—Am. bor. occ.
incana, *Nutt. l. c.* 305.—Am. bor. occ.
mexicana, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 70.—Mexic.
Parryi, *Eaton, in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
Rothrockii, *A. Gray, ex Rothrock, in Wheeler, Rep.* vi. 148.—Am. bor. occ.
scapigera, *Eaton, in Bot. King's Exped.* 145. t. 17.—Am. bor. occ.
sericea, *Hook. Fl. Bor. Am.* ii. 16. t. 119.—Am. bor. occ.
spathulata, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 305.—Am. bor. occ.
strigosa, Eaton, in Bot. King's Exped. 145 = Watsoni.
strigosa, A. Gray, Bot. U. St. Expl. Exped. ii. 344 = florifer.
strigosa, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 306.—Am. bor. occ.
Watsoni, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 84.—Am. bor. occ.
Wilcoxiana, *Wood, in Bull. Torrey Bot. Club*, vi. (1875) 163.—Am. bor. occ.
Wrightii, A. Gray, in Torr. Bot. Mex. Bound. 78 = Aster Wrightii.

TOXANTHERA, Hook. f. in Hook. Ic. Pl. xv. 16. t. 1421 (1883). *CUCURBITACEAE*.
natalensis, *Hook. f. l. c.*—Afr. austr.

TOXANTHES, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 176. *COMPOSITAE*, Benth. & Hook. f. ii. 317.
ANTHOCERASTES, A. Gray, in Hook. Kew Journ. iv. (1852) 225.
major, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 64 = Scyphocoronis viscosa.
Muelleri, *Benth. Fl. Austral.* iii. 592.—Austral.
perpusilla, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 177.—Austral.

TOXANTHUS, Benth. Fl. Austral. iii. 592 (1866), sphalm. = praec.

TOXICARIA, Aepnel. ex Steud. Nom. ed. I. 434, in syn. (1821) = **Antiaris**, Lesch. (Urtic.).
macassariensis, Aepnel. ex Steud. l. c. = A. toxicaria.

TOXICARIA, Schreb. in Naturf. xix. (1783) 146 = **Strychnos**, Linn. (Logan.).
americana, Schreb. l. c. = S. Rouhamon.

TOXICODENDRON, Auct. = **Toxicodendrum**, Thunb. (Euphorb.).

TOXICODENDRON, [Tourn.] Linn. Syst. ed. I (1735) = **Rhus**, Tourn. (Anacard.).
altissimum, Mill. Gard. Dict. ed. VIII. n. 10 = R. succedanea.
arborescens, Mill. l. c. n. 9 = R. arborescens.
arboreum, Mill. l. c. n. 8 = Schmidelia Cominia.
crenatum, Mill. l. c. n. 5 = R. aromatica.
cuneatum, C. Koch, Dendrol. i. 579 = R. aromatica.
glabrum, Mill. Gard. Dict. ed. VIII. n. 3 = R. glabra.
magnum, Bertram, ex Steud. Nom. ed. II. ii. 694 = R. Toxicodendron.
pinnatifolium, Mill. Gard. Dict. ed. VIII. n. 4 = Amyris toxifera.
pubescens, Mill. l. c. n. 2 = R. Toxicodendron.
serratum, Mill. l. c. n. 7 = R. Toxicodendron.
volubile, Mill. l. c. n. 6 = R. Toxicodendron.
vulgare, Mill. l. c. n. 1 = R. Toxicodendron.

TOXICODENDRUM, Gaertn. Fruct. i. 207. t. 44 (1788) = **Allophylus**, Linn. (Sapind.).
Cobbe, Gaertn. l. c. = A. Cobbe.

TOXICODENDRUM, Thunb. in Vet. Akad. Handl. Stockh. (1796) 188. t. 7. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 280.
HYAENANCHE, Lamb. Desc. Cinch. (1797) 52. t. 10.
acutifolium, *Benth. in Journ. Linn. Soc.* xvii. (1878) 214.—Afr. austr.
capense, *Thunb. in Vet. Akad. Handl. Stockh.* (1796) 188. t. 7.—Afr. austr.

TOXICOPHLAEA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 24 = **Acokanthera**, G. Don (Apocyn.).
cestroides, A. DC. in DC. Prod. viii. 336 = A. venenata.
spectabilis, Sond. in Linnaea, xxiii. (1850) 79 = A. spectabilis.
Thunbergii, Harv. in Hook. Lond. Journ. Bot. i. (1842) 24 = A. venenata.
Thunbergii, Hort. Williams.—Cf. Gard. Chron. (1872) 327, 363 = A. spectabilis.

TOXINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Buddleia**, Houst. (Logan.).

TOXOCARPUS, Wight & Arn. Contrib. 61 (1834).
ASCLEPIADEAE, Benth. & Hook. f. ii. 746.
GONIOSTEMMA, Wight, Contrib. 62 (1834).
RHYNCHOSTIGMA, Benth. in Benth. & Hook. f. Gen. ii. 771 (1876).
SCHISTOCODON, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 362.
acuminatus, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 15.—Reg. Himal.
africanus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 109.—Afr. trop.
auriculatus, *Decne. in DC. Prod.* viii. 505.—Java.
Blumei, *Decne. l. c.* 506.—Java.
concanensis, *Hook. f. Fl. Brit. Ind.* iv. 14.—Ind. or.
crassifolius, Dalz. & Gibs. Bomb. Fl. 148 = Genianthus laurifolius.
crassifolius, Wight & Arn. Contrib. 61 = Genianthus crassifolius.
eriocarpus, *Hook. f. Fl. Brit. Ind.* iv. 15 (sp. dub.).—Ind. or.
glaucus, *Decne. in DC. Prod.* viii. 505.—Ins. Philipp.
gracilis, *Decne. l. c.*—Ins. Philipp.
Griffithii, *Decne. l. c.*—Burma; Malaya.
himalensis, *Falc. ex Hook. f. Fl. Brit. Ind.* iv. 13.—Reg. Himal.
Kleinii, *Wight & Arn. Contrib.* 61.—Ind. or.
Kurzii, *Hook. f. Fl. Brit. Ind.* iv. 15.—Ins. Andaman.
laurifolius, Wight & Arn. Contrib. 61 = Genianthus laurifolius.
longistigma, Wight & Arn. ex Steud. Nom. ed. II. ii. 694 = Roxburghii.
macrophyllus, *Decne. in DC. Prod.* viii. 505.—Java.
maritimus, *Miq. Fl. Ind. Bat.* ii. 476.—Java.
oblongifolius, *Decne. in DC. Prod.* viii. 505.—Ins. Maurit. (cult.).
Roxburghii, Griff. Notul. iv. 68; Ic. Pl. Asiat. t. 386 A. f. 4, 5 = Griffithii?
Roxburghii, *Wight & Arn. Contrib.* 61.—Ind. or.
villosus, *Decne. in DC. Prod.* viii. 506.—Java.
Wightianus, *Hook. & Arn. Bot. Beech. Voy.* 200 (sp. dub.).—Reg. Himal.; China.

TOXOPHOENIX, Schott, Nachr. Oester. Naturf. in Brasil. Anh. ii. 12 (1822) = **Astrocaryum**, G. F. W. Mey. (Palm.).
aculeatissima, Schott, l. c. 12 = A. Ayri.

TOXOPUS, Rafin. New Fl. Am. ii. 71 (1836) = **Macranthera**, Torr. (Scrophul.).
calycinus, Rafin. l. c. 72 = M. fuchsioides.
gymnanthes, Rafin. l. c. = M. Lecontii.

TOXOSIPHON, Baill. Adansonia, x. (1872) 311 = **Erythrochiton**, Nees & Mart. (Rutac.).
Lindeni, Baill. l. c. 312 = E. Lindeni.

TOXOSTIGMA, A. Rich. Tent. Fl. Abyss. ii. 86. t. 77 (1850) = **Arnebia**, Forsk. (Borag.).
luteum, A. Rich. l. c.—Abyss.
purpurascens, A. Rich. l. c. 87.—Abyss.

TOXOTROPHIS, Planch. in DC. Prod. xvii. 208 (1873), sphalm. = **Taxotrophis**, Blume (Urtic.).

TOXOTROPIS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 506 = **Corynella**, DC. (Legumin.).
Jaegeri, Turcz. l. c. 507 = C. polyantha.

TOXYLON, Rafin. in Am. Monthly Mag. iv. (1818) 188 = **Maclura**, Nutt. (Urtic.).
Maclura, Rafin. New Fl. Am. iii. 43 = M. aurantiaca.

TOZZETTIA, Parl. Nuov. Gen. e Spec. 11 (1854) = **Fritillaria**, Tourn. (Liliac.).
persica, Parl. l. c. 13 = F. persica.

TOZZETTIA, Savi, in Mem. Soc. Ital. Sc. viii. (1798) 477 = **Alopecurus**, Linn. (Gramin.).
pratensis, Savi, l. c. = A. utriculatus.
utriculata, Savi, in Usteri, Ann. Bot. xxiv. (1800) 49 = A. utriculatus.

TOZZIA, Linn. Gen. ed. I. 181 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 980.
FERECUPPA, Dulac, Fl. Hautes-Pyr. 373 (1867).
KERNERA, Schrank, Baier. Reise, 50 (1786).
alpina, *Linn. Sp. Pl.* 607.—Europ.

TRACANTHELIUM, Kit. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = **Phyteuma**, Linn. (Campan.).
canescens, Schur, l. c. = P. canescens.
salignum, Schur, Enum. Pl. Transs. 973 (Index) = P. canescens.

TRACAULON, Rafin. Fl. Tellur. iii. 13 (1836) = **Polygonum**, Tourn.
arifolium, Rafin. l. c. = P. arifolium.

TRACHELANTHUS, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 82. t. 8 C = **Begonia**, Tourn.
attenuatus, Klotzsch, l. c. 83 = B. attenuata.
rhizocarpus, Klotzsch, l. c. = B. rhizocarpa.

TRACHELANTHUS, Kunze, in Bot. Zeit. viii. (1850) 665 = **Solenanthus**, Ledeb. (Borag.).
cerinthoides, Kunze, l. c. = S. cerinthoides.
kurdicus, Boiss. Fl. Orient. iv. 271 = S. cerinthoides.

TRACHELIOPSIS, Opiz, Seznam, 98 (1852) = **Campanula**, Tourn.
latifolia, Opiz, l. c = C. latifolia.
vulgaris, Opiz, l. c. = C. Trachelium.

TRACHELIUM, Tourn. ex Linn. Syst. ed. I (1735). *CAMPANULACEAE*, Benth. & Hook. f. ii. 563.
alteratum, Bianca, ex Nym. Consp. 485 = caeruleum.
angustifolium, *Schousb. Vextr. Marokko*, 85.—Marocc.
asperuloides, *Boiss. & Orph. Diagn.* Ser. II. iii. 117 = Graecia.
azureum, Gouan, Hort. Monsp. 100 = caeruleum.
caeruleum, *Linn. Sp. Pl.* 171.—Reg. Mediterr. occ.
diffusum, Linn. f. Suppl. 143 = Prismatocarpus diffusus.
Jacquini, *Boiss. Fl. Orient.* iii. 961.—Ins. Creta.
lanceolatum, *Guss. Fl. Sic. Syn.* i. 252.—Sicil.
longifolium, Bianca, ex Guss. l. c. 577 = lanceolatum.
myrtifolium, *Boiss. Fl. Orient.* iii. 962.—As. Min.
rumelianum, *Hampe, in Flora*, xx. (1837) 234.—Graecia.
rumelicum, Boiss. Fl. Orient. iii. 961, sphalm. = praec.
serotinum, Boiss. Diagn. Ser. II. iii. 117 = tubulosum.
tenuifolium, Linn. f. Suppl. 143 = Merciera tenuifolia.
tubulosum, *Boiss. Diagn.* Ser. I. xi. 60.—Cilicia; Syria.

TRACHELOCARPUS, C. Muell. in Walp. Ann. iv. 909 (1857) = **Begonia**, Tourn.
attenuatus, C. Muell. l. c. 910 = B. attenuata.
rhizocarpus, C. Muell. l. c. = B. rhizocarpa.

TRACHELOSPERMUM, Lem. Jard. Fleur. i. (1851) t. 61. *APOCYNACEAE*, Benth. & Hook. f. ii. 720.
PARECHITES, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 193.
RHYNCHOSPERMUM, Lindl. in Journ. Hort. Soc. i. (1846) 74.
TRIADENIA, Miq. Fl. Ind. Bat. ii. 458 (1856).
axillare, *Hook. f. Fl. Brit. Ind.* iii. 668.—Reg. Himal.
difforme, *A. Gray, Syn. Fl. N. Am.* ii. 85.—Am. bor.
fragrans, *Hook. f. Fl. Brit. Ind.* iii. 667.—Reg. Himal.
gracilipes, *Hook. f. l. c.* 668.—Reg. Himal.
jasminoides, *Lem. Jard. Fleur.* i. (1851) t. 61.—China.

TRACHINEMA, Rafin. Fl. Tellur. ii. 27 (1836) = **Anthericum**, Linn. (Liliac.).
hispida, Rafin. l. c. = A. hispidum.
vespertina, Rafin. l. c. = A. falcatum.

TRACHODES, D. Don, in Trans. Linn. Soc. xvi. (1830) 182 = **Sonchus**, Tourn. (Compos.).
paniculatus, D. Don, l. c. = S. paniculatus.

TRACHOPYRON, Ger. ex Rafin. Fl. Tellur. iii. 10 (1836) = **Polygonum**, Tourn.

TRACHYANDRA, Kunth, Enum. Pl. iv. 573 (1843) = **Anthericum**, Linn. (Liliac.).
affinis, Kunth, l. c. 579 = A. Jacquinianum.
asperata, Kunth, l. c. 574 = A. asperatum.
blepharophora, Kunth, l. c. 578 = A. ciliatum, *Linn. f.*
bracteosa, Kunth, l. c. 582 = A. ciliatum, *Linn. f.*
Brehmeana, Kunth, l. c. 586 = A. Brehmeanum.
canaliculata, Kunth, l. c. 578 = A. canaliculatum.
ciliata, Kunth, l. c. 585 = A. ciliatum, *Linn. f.*
corymbosa, Kunth, l. c. 577 = A. hirsutum.
divaricata, Kunth, l. c. 580 = A. revolutum.
elongata, Kunth, l. c. 584 = A. Jacquinianum.
falcata, Kunth, l. c. 586 = A. falcatum.
fimbriata, Kunth, l. c. 583 = A. longifolium.
flexifolia, Kunth, l. c. 579 = A. flexifolium.
floribunda, Kunth, l. c. 583 = Bulbinella latifolia.
hirsuta, Kunth, l. c. 577 = A. hirsutum.
hispida, Kunth, l. c. 574 = A. hispidum.
humilis, Kunth, l. c.—Afr. austr.
Jacquinii, Kunth, l. c 578 = A. Jacquinianum.
longepedunculata, Kunth, l. c. 584 = A. longepedunculatum.
longifolia, Kunth, l. c. 582 = A. longifolium.
muricata, Kunth, l. c. 576 = A. muricatum.
paradoxa, Kunth, l. c. 576 = A. paradoxum.
pauciflora, Kunth, l. c. 584 = A. triflorum.
revoluta, Kunth, l. c. 579 = A. revolutum.
scabra, Kunth, l. c. 585 = A. scabrum.
Schultesii, Kunth, l. c. 586 = A. Schultesii.
undulata, Kunth, l. c. 582 = A. hispidum.
vespertina, Kunth, l. c. 581 = A. falcatum.

TRACHYCARPUS, H. Wendl. in Bull. Soc. Bot. Fr. viii. (1861) 429. *PALMAE*, Benth. & Hook. f. iii. 929.
excelsus, *H. Wendl. l. c.*—Japon.
Fortunei, *H. Wendl. l. c.*—China.
Griffithii, Hort. ex Rev. Hortic. (1879) 212 = khasianus.
khasianus, *H. Wendl. in Bull. Soc. Bot. Fr.* viii. (1861) 429.—Reg. Himal.
Martianus, *H. Wendl. in Kerch. Palm.* 258.—Reg. Himal.

TRACHYCARYON, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844) = **Adriana**, Gaudich. (Euphorb.).
Billardieri, Klotzsch, ex Benth. Fl. Austral. vi. 135 = A. quadripartita.
Cunninghami, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 15 = A. glabrata.
Hookeri, F. Muell. l. c. 16 = A. Hookeri.
Klotzschii, F. Muell. l. c. 15 = A. Klotzschii.
Labillardieri, Klotzsch, in Lehm. Pl. Preiss. i. 175 = A. quadripartita.

TRACHYDIUM, Lindl. in Royle, Illustr. Bot. Himal. 232 (1835). *UMBELLIFERAE*, Benth. & Hook. f. i. 884.
EREMODAUCUS, Bunge, Del. Sem. Hort. Dorpat. (1843).—Cf. Linnaea, xviii. (1844) 151.
HAPLOSCIADIUM, Hochst. in Flora, xxvii. (1844) 20.
abyssinicum, *Benth. & Hook. f. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 10.—Afr. trop.
depressum, *Boiss. Fl. Orient.* ii. 929.—Persia.
dissectum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 672.—Reg. Himal.
elbrusense, *Boiss. Fl. Orient.* ii. 930.—Persia.
hirsutulum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 672.—Reg. Himal.
Kotschyi, *Boiss. Fl. Orient.* ii. 929.—Persia.
novem-jugum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 672.—Reg. Himal.
obtusiusculum, *Benth. & Hook. f. Gen.* ii. 884; *C. B. Clarke, l. c.* 673.—Reg. Himal.
Roylei, *Lindl. in Royle, Illustr. Bot. Himal.* 232.—Reg. Himal.; Ind. or.

TRACHYLOBIUM, Hayne, in Flora, x. (1827) 743; et Arzn. Gew. xi. t. 18 (1830). *LEGUMINOSAE*, Benth. & Hook. f. i. 583.
floribundum, G. Don, Gen. Syst. ii. 459 = Hymenaea floribunda.
Gaertnerianum, Hayne, in Flora, x. (1827) 744 = Hornemannianum.
Hornemannianum, *Hayne, l. c.*—Afr. trop.
Lamarckeanum, Hayne, l. c. = Hymenaea verrucosa.
Martianum, Hayne, l. c. = Hymenaea Martiana.
mossambicense, Klotzsch, in Peters, Reise Mossamb. Bot. 21. t. 11 = Hornemannianum.
verrucosum, *Oliver, Fl. Trop. Afr.* ii. 311, in adnot.—Ins. Mascar.

TRACHYLOMIA, Nees, in Mart. Fl. Bras. ii. 1. 174 (1842) = **Scleria**, Berg. (Cyperac.).
triglomerata, Nees, l. c. = S. triglomerata.

TRACHYMARATHRUM, Tausch, in Flora, xvii. (1834) 344 = **Hippomarathrum**, Link (Umbell.).
siculum, Tausch, l. c. 344 = H. cristatum.

TRACHYMENE, DC. Prod. iv. 72 (1830) = **Platysace**, Bunge (Umbell.).
anceps, DC. Prod. iv. 73 (= *Siebera compressa*).—Austral.
Billardieri, F. Muell. Census, 62 = seq.
buxifolia, Sieber, ex DC. Prod. iv. 73 (= *Siebera Billardieri*).—Austral.
Candelabrum, Bunge, in Lehm. Pl. Preiss. i. 289 (= *Siebera juncea*).—Austral.
cirrosa, F. Muell. Census, 62 = P. cirrosa.
commutata, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 30 (= *Siebera commutata*).—Austral.
compressa, Spreng. Sp. Umbellif. 9 (= *Siebera compressa*).—Austral.
conferta, Gaudich. ex Benth. in Enum. Pl. Hueg. 54 = seq.
crassifolia, Benth. l. c. 54 (= *Siebera Billardieri*).—Austral.
deflexa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 31 (= *Siebera deflexa*).—Austral.
dissecta, F. Muell. Census, 62 (= *Siebera dissecta*).—Austral.
effusa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 31 = P. trachymenoides.
ericoides, Sieber, ex DC. Prod. iv. 73 (= *Siebera ericoides*).—Austral.
filiformis, Bunge, in Lehm. Pl. Preiss. i. 289 (= *Siebera compressa*).—Austral.
haplosciadea, F. Muell. Census, 62 (= *Siebera haplosciadea*).—Austral.
heterophylla, F. Muell. First Gen. Rep. 13 (= *Siebera heterophylla*).—Austral.
juncea, Bunge, in Lehm. Pl. Preiss. i. 287 (= *Siebera juncea*).—Austral.
lanceolata, Spreng. Syst. i. 879 (= *Siebera Billardieri*).—Austral.
linearis, Spreng. Sp. Umbellif. 7 (= *Siebera linearifolia*).—Austral.
myrtifolia, Sieber, ex DC. Prod. iv. 73 = seq.
ovalis, DC. l. c. (= *Siebera Billardieri*).—Austral.
ovata, Spreng. Sp. Umbellif. 7 = praec.
pendula, Benth. in Enum. Pl. Hueg. 54 (= *Siebera juncea*).—Austral.
platyptera, Bunge, in Lehm. Pl. Preiss. i. 287 (= *Siebera compressa*).—Austral.
ramosissima, Benth. in Enum. Pl. Hueg. 54 (= *Siebera juncea*).—Austral.
scabriuscula, Bunge, in Lehm. Pl. Preiss. i. 287 = praec.
Stephensonii, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 170 (= *Siebera Stephensonii*).—Austral.
stricta, Bunge, in Lehm. Pl. Preiss. i. 288 (= *Siebera compressa*).—Austral.
subvelutina, DC. Prod. iv. 73 (= *Siebera ericoides*).—Austral.
tenuis, DC. l. c. 73 = praec.
tenuissima, F. Muell. Census, 62 (= *Siebera tenuissima*).—Austral.
teres, Bunge, in Lehm. Pl. Preiss. i. 286 (= *Siebera juncea*).—Austral.
valida, F. Muell. Census, 62 = P. valida.

TRACHYMENE, Rudge, in Trans. Linn. Soc. x. (1811) 300. t. 21. *UMBELLIFERAE*, Benth. & Hook. f. i. 873.
CESATIA, Endl. in Ann. Wien. Mus. ii. (1838) 200.
DIDISCUS, DC. in Bot. Mag. t. 2875 (1828).

TRACHYMENE:—
DIMETOPIA, DC. Prod. iv. 71 (1830).
HEMICARPUS, F. Muell. in Hook. Kew Journ. ix. (1857) 18.
HUEGELIA, Reichb. Consp. 144 (1828).
LAMPRA, Lindl. ex DC. Prod. iv. 72 (1830).
PRITZELIA, Walp. Rep. i. 428 (1842).
australis, *Benth. Fl. Austral.* iii. 349.—Austral.
austro-caledonica, *F. Muell. Phyt. New Hebr.* 13.—N. Hebrid.
coerulea, *R. Grah. in Edinb. N. Phil. Journ.* (July-Sept. 1828) 380.—Austral.
cyanea, A. Cunn. ex DC. Prod. iv. 72 = coerulea.
cyanopetalea, *Benth. Fl. Austral.* iii. 348.—Austral.
eriocarpa, *Benth. l. c.*—Austral.
gigantea, *Ait. ex W. Baxt. in Loud. Hort. Brit. Suppl.* iii. 650.—Hab. ?
glandulosa, *Benth. Fl. Austral.* iii. 350.—Austral.
glaucifolia, *Benth. l. c.*—Austral.
hemicarpa, *Benth. l. c.* 351.—Austral.
Homei, *Seem. in Journ. Bot.* ii. (1864) 75.—N. Caled.
humilis, *Benth. Fl. Austral.* iii. 351.—Austral.
incisa, *Rudge, in Trans. Linn. Soc.* x. (1811) 300. t. 21.—Austral.
lanceolata, *Rudge, l. c.*—Austral.
ovata, *Rudge, l. c.* 300.—Austral.
pilosa, *Sm. in Rees, Cycl.* xxxix. Add. n. 2.—Austral.
procumbens, *Benth. Fl. Austral.* iii. 350.—Austral.
villosa, *Benth. Fl. Austral.* iii. 349.—Austral.

TRACHYNIA, Link, Hort. Berol. i. 42 (1827) = **Brachypodium**, Beauv. (Gramin.).
distachya, Link, l. c. 43 = B. distachyum.
rigida, Link, l. c. = Triticum asperum.
subtilis, Hort. ex Ind. Sem. Hort. Lips. (1842).—Cf. Walp. Ann. i. 948 = B. distachyum.

TRACHYNOTIA, Michx. Fl. Bor. Am. i. 63 (1803) = **Spartina**, Schreb. (Gramin.).
alterniflora, DC. Fl. Fr. Suppl. 279 = S. stricta.
alternifolia, Steud. Nom. ed. II. ii. 695, sphalm. = praec.
cynosuroides, Michx. Fl. Bor. Am. i. 64 = S. Schreberi.
fasciculata, Poir. Encyc. Suppl. ii. 443, in obs. = S. fasciculata.
juncea, Michx. Fl. Bor. Am. i. 64 = S. juncea.
patens, Poir. Encyc. Suppl. ii. 443, in obs. = S. patens.
polystachya, Michx. Fl. Bor. Am. i. 64 = S. polystachya.
stricta, DC. Fl. Fr. iii. 74 = S. stricta.

TRACHYOZUS, Reichb. Consp. 48 (1828) = **Trachys**, Pers. (Gramin.).
muricata, Steud. Syn Pl. Gram. 112 = Trachys mucronata.

TRACHYPHRYNIUM, Benth. in Benth. & Hook. f. Gen. iii. 651 (1883). *SCITAMINEAE*, Benth. & Hook. f. l. c.

TRACHYPHYTUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 533 (1840) = **Mentzelia**, Plum. (Loasac.).
albicaule, Nutt. l. c. 534 = M. albicaulis.
congestum, Nutt. l. c. = M. congesta.
gracile, Nutt. l. c. = M. albicaulis.

TRACHYPLEURUM, Reichb. Consp. 143 (1828) = **Bupleurum**, Tourn. (Umbell.).
tenuissimum, Schur, Enum. Pl. Transs. 251 = B. tenuissimum.

TRACHYPOGON, Nees, Agrost. Bras. 341 (1829), partim. *GRAMINEAE*, Benth. & Hook. f. iii. 1133.
angustifolius, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 522.—Mexic.
angustifolius, Nees, Agrost. Bras. 342 = polymorphus.
argenteus, Nees, l. c. 348 = Andropogon saccharoides.
avenaceus, Nees, l. c. 354 = Sorghum halepense.
canescens, Nees, l. c. 343 = polymorphus.
capensis, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 257 = polymorphus.
dactyloides, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 523.—Mexic.

TRACHYPOGON:—
dissolutus, Nees, in Linnaea, xix. (1847) 695 = polymorphus.
Gouini, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 523 = polymorphus.
gracilis, Anderss. in Oefv. Vet. Akad. Foerhandl. Stockh. xiv. (1857) 50 = polymorphus.
hirtus, Nees, Agrost. Bras. 346 = Heteropogon hirtus.
laguroides, Nees, l. c. 349 = Andropogon saccharoides.
ligularis, Nees, l. c. 345 = polymorphus.
macroglossus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 257 = polymorphus.
micans, Anderss. in Oefv. Vet. Akad. Foerhandl. Stockh. xiv. (1857) 47 = polymorphus.
Minarum, Nees, Agrost. Bras. 349 = Chrysopogon stipoides.
mollis, Nees, l. c. 343 = polymorphus.
Montufari, Nees, l. c. 342 = polymorphus.
Muelleri, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 523 = polymorphus.
plumosus, Nees, Agrost. Bras. 344 = polymorphus.
polymorphus, *Hack. in Mart. Fl. Bras.* ii. III. 263.—Reg. trop. et subtrop.
Preslei, Anderss. in Oefv. Vet. Akad. Foerhandl. Stockh. xiv. (1857) 50 = polymorphus.
pubescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 180 = Heteropogon pubescens.
rufus, Nees, Agrost. Bras. 346 = Andropogon rufus.
scaberrimus, Nees, l. c. 354 = Chrysopogon nutans.
Schoenanthus, Nees, in Linnaea, vii. (1832) 281 = Andropogon Nardus.
scrobiculatus, Nees, Agrost. Bras. 347 = Heteropogon acuminatus.
stipoides, Nees, l. c. 351 = Chrysopogon avenaceus.
truncatus, Anderss. in Oefv. Vet. Akad. Foerhandl. Stockh. xiv. (1857) 49 = polymorphus.
vestitus, *Anderss. l. c.* 52.—Bras.
violaceus, Anderss. l. c. 49 = polymorphus.

TRACHYRHYNCHIUM, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 113 = **Cladium**, P. Br. (Cyperac.).
iridifolium, Nees, l. c. 114 (= *Baumea Meyenii*).—Ins. Sandvic.

TRACHYRHYNGIUM, Nees, ex Kunth, Enum. Pl. ii. 314, in syn. (1837), sphalm. = praec.

TRACHYS, Pers. Syn. i. 85 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1122.
TRACHYOZUS, Reichb. Consp. 48 (1828).
TRACHYSTACHYS, A. Dietr. Sp. Pl. ii. 16 (1833).
mucronata, *Pers. Syn.* i. 85.—Ind. or.
muricata, Steud. Syn. Pl. Gram. 112, sphalm. = praec.

TRACHYSCIADIUM, Eckl. & Zeyh. Enum. 341 (1837) = **Carum**, Rupp. (Umbell.).
capense, Eckl. & Zeyh. l. c. = C. capense.

TRACHYSPERMA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Limnanthemum**, S. T. Gmel. (Gentian.).
nutans, Rafin. l. c. = L. trachyspermum.

TRACHYSPERMUM, Link, Enum. Hort. Berol. i. 267 (1821) = **Carum**, Rupp. (Umbell.).
copticum, Link, l. c. = C. copticum.
rigens, G. Don, in Loud. Hort. Brit. 103 (= *Bunium rigens*).—Afr. austr.

TRACHYSTACHYS, A. Dietr. Sp. Pl. ii. 16 (1833) = **Trachys**, Pers. (Gramin.).
geminata, A. Dietr. l. c. = Trachys mucronata.

TRACHYSTELLA, Steud. Nom. ed. II. ii. 695 (1841), sphalm. = TRACHYTELLA, DC. = **Tetracera**, Linn. (Dillen.).

TRACHYSTEMMA, Auct. ex Meissn. Gen. Comm. 188 (1840) = seq.

TRACHYSTEMON, D. Don, in Edinb. N. Phil. Journ. xiii. (1832) 239. *BORAGINEAE*, Benth. & Hook. f. ii. 855.
NORDMANNIA, Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 312.
PSILOSTEMON, DC. Prod. x. 85 (1846).

TRACHYSTEMON:—
creticum, *D. Don, ex G. Don, Gen. Syst.* iv. 809.—Archip. Graec.
orientale, *D. Don, in Edinb. N. Phil. Journ.* xiii. (1832) 239.—As. Min.

TRACHYSTIGMA, C. B. Clarke, in DC. Monog. Phan. v. 131 (1883). *CYRTANDREAE.*
Mannii, *C. B. Clarke, l. c.*—Afr. trop.

TRACHYTELLA, DC. Syst. i. 410 (1818) = **Tetracera**, Linn. (Dilleniac.).
Actaea, DC. l. c. = Tet. sarmentosa.
aspera, DC. l. c. = Tet. sarmentosa.
Calligonum, DC. l. c. = Tet. sarmentosa.

TRACHYTHECA, Nutt. ex Benth. in DC. Prod. xiv. 7 (1857) = **Eriogonum**, Michx. (Polygon.).

TRACTEMA, Rafin. Fl. Tellur. ii. 57 (1836) = **Scilla**, Linn. (Liliac.).
pumila, Rafin. l. c. = S. monophyllos.

TRADESCANTIA, Rupp. ex Linn. Syst. ed. I (1735).
COMMELINACEAE, Benth. & Hook. f. iii. 853.
APLOLEIA, Rafin. Fl. Tellur. ii. 17 (1836).
DESCANTARIA, Schlecht. in Linnaea, xxvi. (1853–55) 140.
DISGREGA, Hassk. in Flora, xlix. (1866) 215.
EPHEMERUM, Tourn. ex Moench, Meth. 237 (1794).
ETHEOSANTHES, Rafin. Neogenyt. 3 (1825).
GIBASIS, Rafin. Fl. Tellur. ii. 16 (1836).
HEMINEMA, Rafin. l. c. 17 (1836).
HETERACHTHIA, Kunze, in Bot. Zeit. viii. (1850) 1.
KNOWLESIA, Hassk. in Flora, xlix. (1866) 215.
LEIANDRA, Rafin. Fl. Tellur. ii. 17 (1836).
MANDONIA, Hassk. in Flora, liv. (1871) 260.
PHYODINA, Rafin. Fl. Tellur. ii. 16 (1836).
PYRRHEIMA, Hassk. in Flora, lii. (1869) 366.
SIDERASIS, Rafin. Fl. Tellur. iii. 67 (1836).
SKOFITZIA, Hassk. & Kanitz, in Oestr. Bot. Zeitschr. xxii. (1872) 147.
TRIPAGANDRA, Rafin. Fl. Tellur. ii. 16 (1836).
TROPITIA, Rafin. l. c. iii. 68 (1836).
acaulis, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 375 = crassifolia.
albida, Herb. Kunth, ex C. B. Clarke, in DC. Monog. Phan. iii. 290 = virginiana.
albiflora, Kunth, Enum. Pl. iv. 84 = fluminensis.
ambigua, *Mart. in Schult. f. Syst.* vii. 1170.—Bras.
amplexicaulis, *Klotzsch, ex C. B. Clarke, in DC. Monog. Phan.* iii. 304.—Mexic.
anagallidea, *Seub. in Mart. Fl. Bras.* iii. I. 249.—Bras.
Andrieuxii, *C. B. Clarke, in DC. Monog. Phan.* iii. 291. (Andreuxii).—Mexic.
anomala, Torr. Bot. Mex. Bound. 225 = Tinantia anomala.
aphylla, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 201 = Aneilema scapiflorum.
argentea, Falkenb. Bau Veget. Org. Monocot. 116 (Quid?).—Hab.?
Aubletii, J. F. Gmel. Syst. 533 = Dichorisandra Aubletiana.
axillaris, Linn. Mant. ii. 321 = Cyanotis axillaris.
axillaris, Rafin. New Fl. Am. ii. 87 = virginiana?
Balbisii, Kunth, Enum. Pl. iv. 97 = elongata.
barbata, Rafin. New Fl. Am. ii. 85 = virginiana.
barbata, Spreng. Syst. iv. Cur. Post. 138 = Cyanotis barbata.
biandra, Torr. Mex. Bound. Bot. 224 (err. typ.) = *leiandra.*
bicolor, Kunth, Enum. Pl. iv. 88 = gracilis.
bifida, Roth, Catalect. fasc. i. 42 = Tinantia fugax.
biflora, Willd. ex Schult. f. Syst. vii. 1176 = Commelina subulata.
brasiliensis, Lodd. ex Steud. Nom. ed. II. ii. 694 (Quid?).—Bras.
brevicaulis, Rafin. Atl. Journ. 150 = virginiana.
Callisia, Sw. Fl. Ind. Occ. i. 603 = Callisia repens.
canaliculata, Rafin. Atl. Journ. 150 = virginiana.
capitata, Blume, Enum. Pl. Jav. i. 6 = Cyanotis capitata.
capitata, Vell. Fl. Flum. 139; iii. t. 151 = Campelia Zanonia.
caricifolia, Hook. Bot. Mag. t. 3546 = virginiana.
carinata, *E. Mey. in Presl, Rel. Haenk.* i. 139.—Peruv.
cerinthoides, *Kunth, Enum. Pl.* iv. 83.—Bras.
ciliata, *Blume, Cat. Gew. Buitenz.* 61.—Java.
ciliata, Hort. Berol. ex Kunth, Enum. Pl. iv. 81 = virginiana.
cirrifera, Mart. Ausw. Merkw. Pfl. 10. t. 7 = virginiana.
Commelina, Neuenh. in Roem. Arch. i. II. (1797) 24 = Tinantia fugax.
Commelinii, Vell. Fl. Flum. 140; iii. t. 154 = elongata.
commelinoides, *Schult. f. Syst.* vii. 1176.—Mexic.
congesta, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 377 = elongata.
congesta, Penny, ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 650 (Quid?).—Am. bor.
consanguinea, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 301 = geniculata.
cordifolia, Griff. Priv. Journ. 208 = Streptolirion volubile.
cordifolia, *Sw. Prod. Veg. Ind. Occ.* 57.—Am. trop.
crassifolia, *Cav. Ic.* i. 54. t. 75.—Am. austr.
crassifolia, Hassk. in Miq. Pl. Jungh. 152 = Cyanotis capitata.
crassipes, Barcl. ex R. Grah. in Edinb. N. Phil. Journ. (Jan.–March, 1829) 388 = T. Crassula.
Crassula, *Link & Otto, Ic. Pl. Rar. Hort. Berol.* 13. t. 7.—Bras.
crinigera, Nees, in Linnaea, xix. (1847) 700 = virginiana.
cristata, Blanco, Fl. Philipp. ed. III. i. 293 = Commelina nudiflora.
cristata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 246 = Cyanotis papilionacea.
cristata, Linn. Syst. ed. XII. 233 = Cyanotis cristata.
cristata, Walt. Fl. Carol. 119 = virginiana.
cucullata, Roth, Nov. Pl. Sp. 189 = Cyanotis cucullata.
cumanensis, *Kunth, Enum. Pl.* iv. 96.—Venezuela.
cyanotidea, Hassk. Commel. Ind. 65 = Cyanotis villosa.
cymbispatha, *C. B. Clarke, in DC. Monog. Phan.* iii. 296.—Bolivia.
debilis, H. B. & K. Nov. Gen. et Sp. i. 262 = gracilis.
decumbens, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 118 = geniculata.
diaphana, Willd. ex Schult. f. Syst. vii. 1178, in obs. = gracilis.
dilatata, C. B. Clarke, in DC. Monog. Phan. iii. 304 = amplexicaulis.
discolor, L'Hérit. Sert. Angl. 8. t. 12 = Rhoeo discolor.
discolor, Rafin. New Fl. Am. ii. 87 = virginiana.
Disgrega, *Kunth, Enum. Pl.* iv. 97.—Mexic.
diuretica, Mart. Reise Bras. i. 281 = elongata.
divaricata, Rafin. Atl. Journ. 148 = virginiana.
divaricata, Vahl, Eclog. Am. i. 34 = Dichorisandra Aubletiana.
effusa, Mart. in Schult. f. Syst. vii. 1159 = geniculata.
Ehrenbergiana, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 305 = T. Disgrega.
elata, Lodd. Bot. Cab. t. 1513 = virginiana.
elegans, Pritz. Ic. Ind. 1110, sphalm. = Thysanocarpus elegans.
elongata, *G. F. W. Mey. Prim. Fl. Esseq.* 146.—Am. austr.
erecta, Jacq. Coll. iv. 113; Ic. Pl. Rar. ii. 13. t. 354 = Tinantia fugax.
fasciculata, Heyne, ex Roth, Nov. Pl. Sp. 189 = Cyanotis fasciculata.
filiformis, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 376 = Leptorhoës filiformis.
Fischeri, Weinm. Elench. Pl. Hort. Pawlowsk. (1824) 154; et ex Schult. f. Syst. vii. 1172 = crassifolia.
flexuosa, Rafin. Atl. Journ. 150 = virginiana.
floribunda, Kunth, Enum. Pl. iv. 89 = geniculata.
floribunda, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 377 = holosericea.
floridana, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 381.—Florida.
fluminensis, *Vell. Fl. Flum.* 140; iii. t. 152.—Bras.
formosa, Willd. Sp. Pl. ii. 20 = Cyanotis nodiflora.
fuscata, *Lodd. Bot. Cab.* t. 374.—Bras.
Galeottiana, Kunth, Enum. Pl. iv. 696 = holosericea.
gangetica, Linn. ex C. B. Clarke, in DC. Monog. Phan. iii. 216 = Aneilema vaginatum.
Gaudichaudiana, Kunth, Enum. Pl. iv. 93 = elongata.
geniculata, Blanco, Fl. Filip. ed. I. 232 = Floscopa scandens.
geniculata, *Jacq. Enum. Pl. Carib.* 18.—Am. austr.
geniculata, Lour. Fl. Cochinch. 193 = Cyanotis geniculata.
geniculata, Vell. Fl. Flum. 140; iii. t. 153 = cymbispatha.
gentianaefolia, Salisb. Prod. 216 = Campelia Zanonia.
glabra, C. B. Clarke, in DC. Monog. Phan. iii. 290, in syn. errore = virginiana.
glandulosa, *Seub. in Mart. Fl. Bras.* iii. I. 253.—Guian.
glomerata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 245 = Cyanotis cucullata.
glomerata, Willd. ex Schult. f. Syst. vii. 1175 = Floscopa glomerata.
Gonatandra, Schlecht. in Linnaea, xxiv. (1851) 659 = Campelia Zanonia.
gracilis, *H. B. & K. Nov. Gen. et Sp.* i. 261. t. 672.—Ecuador.
graminifolia, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 378 = linearis.
graminifolia, Rafin. Atl. Journ. 148 (Quid?).—Am. bor.
guianensis, Miq. in Linnaea, xviii. (1844) 374 = elongata.
Herba-ratti, Nees, l. c. xix. (1847) 699 = crassifolia.
hirsuta, *H. B. & K. Nov. Gen. et Sp.* i. 263.—N. Granat.
hirsutissima, Pohl, ex Seub. in Mart. Fl. Bras. iii. I. 254 = fuscata.
holosericea, *Kunth, Enum. Pl.* iv. 92.—Mexic.
Humboldtiana, Schult. f. Syst. vii. 1172 = crassifolia.
hypophaea, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1855) App. 9 = geniculata.
imbricata, Roxb. Hort. Beng. [23]; Fl. Ind. ii. 120 = Cyanotis cristata.
incana, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 250 = Cyanotis arachnoidea.
iridescens, Lindl. Bot. Reg. (1838) Misc. 86 = crassifolia.
Karwinskyana, Herb. Berol. ex C. B. Clarke, in DC. Monog. Phan. iii. 298 = venustula.
Karwinskyana, *Schult. f. Syst.* vii. 1165.—Mexic.
Koernickeana, Seub. in Mart. Fl. Bras. iii. I. 249 = cerinthoides.
Kunthiana, Seub. l. c. 252 = geniculata.
laevigata, Auct. = *levigata.*
lanata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 250 = Cyanotis arachnoidea.
lanuginosa, Hassk. Commel. Ind. 67 = crassifolia.
latifolia, Herb. Madr. ex Wall. Cat. n. 5223 D = Aneilema herbaceum.
latifolia, Ruiz & Pav. Fl. Per. iii. 44. t. 272 = Tinantia fugax.
laxiflora, *C. B. Clarke, in DC. Monog. Phan.* iii. 307. —Mexic.
leiandra (err. typ. *biandra*), Torr. Bot. Mex. Bound. 224–5 = Commelina leiandra.
levigata, Rafin. New Fl. Am. ii. 85 = virginiana.
linearis, Benth. Pl. Hartw. 27.—Mexic.
Loureiriana, Schult. f. Syst. vii. 1178 = Cyanotis geniculata.
Lyonii, Steud. Nom. ed. II. ii. 695; Schlecht. in Linnaea, xxv. (1852) 186 = virginiana.
malabarica, Herb. ex Schult. f. Syst. vii. 1161 = Tinantia fugax.
malabarica, Linn. Sp. Pl. ed. II. 412 = Aneilema nudiflorum.
marginata, Bouché, in Verh. Ges. Nat. Fr. Berl. i. (1819) 392 = Campelia Zanonia.
Martensiana, Kunth, Enum. Pl. iv. 697 = Callisia Martensiana.
mexicana, Hort. Belved. ex Schult. f. Syst. vii. 1179 (Quid?).—Mexic.
mexicana, Kunth, Enum. Pl. iv. 93 = elongata.
micrantha, *Torr. Bot. Mex. Bound.* 224.—Mexic.
minuta, *C. B. Clarke, in DC. Monog. Phan.* iii. 307.—Mexic.
mollis, Kunth, Enum. Pl. iv. 95 = elongata.
monandra, Sw. Prod. Veg. Ind. Occ. 57 = Callisia umbellulata.
montana, Heyne, ex Wall. Cat. n. 5203 B = Aneilema montanum.
multiflora, Kunth, Enum. Pl. iv. 96, partim = amplexicaulis.
multiflora, *Sw. Prod. Veg. Ind. Occ.* 57.—Jamaic.
mundula, Kunth, Enum. Pl. iv. 83 = fluminensis.
nana, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 374.—Mexic.

TRADESCANTIA:—
navicularis, *Ortgies, in Regel, Gartenfl.* xxvi. (1877) 130. t. 901.—Peruv.
nervosa, Linn. Mant. ii. 223 = Telipogon angustifolius.
nodiflora, Lam. Encyc. ii. 371 = Cyanotis nodiflora.
ohiensis, Rafin. Préc. Découv. 45 = virginiana.
paniculata, Heyne, ex Roth, Nov. Pl. Sp. 188 = Aneilema montanum.
paniculata, Roxb. Pl. Corom. ii. 6. t. 109 = Floscopa scandens.
papilionacea, Linn. Mant. ii. 513 = Cyanotis papilionacea.
parviflora, Ruiz & Pav. Fl. Per. iii. 43. t. 272. f. 2 = multiflora.
pellucida, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 376 = pulchella.
pilosa, Lehm. Ind. Sem. Hort. Hamb. (1827) 16; et in Nov. Act. Nat. Cur. xiv. II. (1829) 822 = virginiana.
pilosa, Willd. ex Schult. f. Syst. vii. 1155 = Cyanotis pilosa.
pilosissima, Fraser, ex Schult. f. l. c. 1175 = virginiana.
portoricensis, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 122.—Porto Rico.
procumbens, Willd. Sp. Pl. ii. 19 = multiflora.
pulchella, *H. B. & K. Nov. Gen. et Sp.* i. 262.—Mexic.
pumila, Rafin. New Fl. Am. ii. 86 (Quid?).—Am. bor.
purpurascens, *Schau. in Linnaea*, xix. (1847) 700.—Mexic.
radicans, Royle, Illustr. Bot. Himal. 403 = Cyanotis barbata.
recta, Raeusch. Nom. ed. III. 91 = Tinantia fugax.
reflexa, Rafin. New Fl. Am. ii. 87 = virginiana?
repens, *Vand. in Roem. Script.* 95.—Bras.
rhodantha, Torr. Bot. Mex. Bound. 225 = venustula.
riparia, Rafin. New Fl. Am. ii. 88 (Quid?).—Am. bor.
rosea, *Vent. Jard. Cels*, t. 24.—Am. bor.
rufa, Presl, Rel. Haenk. i. 138 = Floscopa scandens.
rupestris, Law, in J. Grah. Cat. Bomb. Pl. 223 = Cyanotis fasciculata.
rupestris, Rafin. Atl. Journ. 150 = virginiana.
scaberrima, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 120 = Aneilema protensum.
Schiedeana, Kunth, Enum. Pl. iv. 90 = geniculata.
Schlechtendalii, Kunth, l. c. 94 = elongata.
Schomburgkiana, Kunth, l. c. 663 = elongata.
Sellowiana, Kunth, l. c. 93 = elongata.
spathacea, Sw. Prod. Veg. Ind. Occ. 57 = Rhoeo discolor.
speciosa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 9 = Commelina leiandra.
speciosa, H. B. & K. Nov. Gen. et Sp. i. 263 = crassifolia.
speciosa, Linn. f. Suppl. 192 = Cyanotis nodiflora.
speciosa, Salisb. Prod. 216 = virginiana.
spicata, Knowles & Westc. Flor. Cab. ii. (1838) 159 = virginiana?
splendens, C. Koch & Bouché, in C. Koch & Fint. Wochenschr. ii. (1859) 334 = virginiana.
subaspera, Ker-Gawl. in Bot. Mag. t. 1597 = virginiana.
subscandens, *Scheidw. in Journ. Hortic. Prat.* ii. (1845) 321.—Mexic.
tenella, Kunth, Enum. Pl. iv. 83 = fluminensis.
terminalis, Blume, Enum. Pl. Jav. i. 6 = Aneilema Hamiltonianum.
thyrsiflora, Blume, l. c. = Pollia thyrsiflora.
trachyloma, C. B. Clarke, in DC. Monog. Phan. iii. 290, in syn. = virginiana.
triandra, Kunth, Enum. Pl. iv. 93 = elongata.
tricolor, Hort. ex C. B. Clarke, in DC. Monog. Phan. iii. 318 = Zebrina pendula.
triflora, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 376 = geniculata.
tuberosa, Roxb. Pl. Corom. ii. 5. t. 108 = Cyanotis tuberosa.
tumida, Lindl. Bot. Reg. (1840) t. 42 = virginiana.
umbellata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 247 = Cyanotis cristata.
umbellata, Pav. ex C. B. Clarke, l. c. 304 = amplexicaulis.
umbellata, Vahl, Eclog. iii. 7 = Tinantia fugax.
undata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 347 = Tinantia fugax.
undulata, Vahl, in Kjoeb. Skrivt. Natur. Selsk. ii. II. (1793) 27 = Tinantia fugax.
vaga, Lour. Fl. Cochinch. 193 = Cyanotis vaga.
vaga, Zoll. Syst. Verz. Ind. Arch. ii. 65 = Cyanotis cristata.

TRADESCANTIA:—
velutina, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 12.—Guatemala.
venustula, *Kunth, Enum. Pl.* iv. 87.—Mexic.
versicolor, Salisb. Prod. 216 = Rhoeo discolor.
villosa, Spreng. Syst. ii. 116 = Cyanotis villosa.
villosissima, Fras. ex C. B. Clarke, in DC. Monog. Phan. iii. 291 = virginiana.
virginiana, *Linn. Sp. Pl.* 288.—Am. bor.
virginica, Georgi, Beschr. Russ. Reich. III. iv. 888 = Trillium ovatum?
virginica, Linn. Syst. ed. X. 975 = virginiana.
virginica, Walt. Fl. Carol. 119 = rosea.
Warmingiana, Seub. ex Warm. in Kjoeb. Vidensk. Meddel. (1872) 126 = elongata.
Warszewicziana, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 11.—Guatemala.
Warszewiczii, Hort. in Rev. Hortic. Sér. IV. ix. (1860) 135 = praec.
Zanonia, Sw. Prod. Veg. Ind. Occ. 57 = Campelia Zanonia.
zebrina, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1340 = Zebrina pendula.

TRAEVIA, Neck. Elem. ii. 268 (1790) = **Trewia**, Linn. (Euphorb.).

TRAGACANTHA, Tourn. ex Linn. Syst. ed. I (1735); Mill. Gard. Dict. ed. VI (1752) = **Astragalus**, Tourn. (Legumin.).
angustifolia, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = A. angustifolius.
argentea, Mill. Gard. Dict. ed. VIII. n. 3 = A. Barba-Jovis.
aristata, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = A. aristatus.
Arnacantha, Stev. l. c. = A. Arnacantha.
aureus, Stev. l. c. = A. aureus.
breviflora, Stev. l. c. = A. eriocephalus.
caspica, Stev. l. c. = A. caspicus.
caucasica, Stev. l. c. = A. caucasicus.
compacta, Stev. l. c. = A. compactus.
cretica, Stev. l. c. = A. creticus.
Criacantha, Stev. l. c. xxix. (1856) I. 257, sphalm. = A. Arnacantha.
denudata, Stev. l. c. iv. (1832) 269 = A. microcephalus.
echinoides, Stev. l. c. = A. angustifolius.
Echinus, Stev. l. c. = A. Echinus.
eriocaulos, Stev. l. c. = A. microcephalus.
genargentea, Stev. l. c. = A. sirinicus.
glabra, Mill. Gard. Dict. ed. VIII. n. 4 = A. Clusii?
hispanica, Mill. l. c. n. 2.—Ins. Balear.
lagopodoides, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = A. lagopoides.
Lagurus, Stev. l. c. = A. Lagurus.
longifolia, Stev. l. c. = A. longifolius.
massiliensis, Mill. Gard. Dict. ed. VIII. n. 1 = A. Tragacantha.
oleaefolia, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = A. oleaefolius.
plumosa, Stev. l. c. = A. plumosus.
Pseudotragacantha, Stev. l. c. = A. Pseudotragacanthus.
ptilodes, Stev. l. c. = A. plumosus.
pycnophylla, Stev. l. c. = A. mesoleios.
sicula, Stev. l. c. = A. siculus.
triacantha, Stev. l. c. xxix. (1856) II. 153 = A. Anthacantha?
trojana, Stev. l. c. iv. (1832) 269 = A. gummifer.
vera, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374 = A. Tragacantha.
xanthura, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 269 = A. Lagurus.

TRAGANTHA, Wallr. ex Endl. Gen. 370 (1837), sphalm. = seq.

TRAGANTHES, Wallr. Sched. Crit. i. 456, in nota (1822) = **Eupatorium**, Tourn. (Compos.).
tenuifolia, Wallr. l. c. = E. foeniculaceum.

TRAGANTHUS, Klotzsch, in Wiegm. Archiv, vii. (1841) 188. t. 9; et in Hook. Lond. Journ. Bot. ii. (1843) 46 = **Bernardia**, Houst. (Euphorb.).
sidoides, Klotzsch, ll. cc. = B. sidoides.

TRAGANUM, Delile, Fl. Egypte, 204. t. 22 (1813).
CHENOPODIACEAE, Benth. & Hook. f. iii. 68.
Moquini, *Webb, ex Moq. in DC. Prod.* xiii. II. 171.—Ins. Canar.

TRAGANUM:—
nudatum, *Delile, Fl. Egypte*, 204.—Aegypt.
nudatum, Moq. in Webb & Berth. Phyt. Canar. iii. 212 = Moquini.
nudatum, Schimp. ex Boiss. Fl. Orient. iv. 985 = Halogeton alopecuroides.
nudatum, Sieber, ex Moq. in DC. Prod. xiii. II. 172 = Cornulaca monacantha.

TRAGIA, Plum. ex Linn. Gen. ed. I. 282 (1737).
EUPHORBIACEAE, Benth. & Hook. f. iii. 329.
AGIRTA, Baill. Etud. Gén. Euph. 463 (1858).
BIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 189.
CTENOMERIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 29.
LASSIA, Baill. Etud. Gén. Euph. 464. t. 4 (1858).
LEPTOBOTRYS, Baill. Etud. Gén. Euph. 478. t. 2 (1858).
LEPTORHACHIS, Klotzsch, in Wiegm. Archiv, vii. 189 (1841).
LEUCANDRA, Klotzsch, l. c. 188.
ZUCKERTIA, Baill. Etud. Gén. Euph. 495. t. 4 (1858).
adenanthera, *Baill. Adansonia*, i. (1860–61) 275.—Afr. trop.
amoena, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 414.—Bras.
angolensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 333.—Afr. trop.
angustifolia, *Benth. in Hook. Niger Fl.* 502.—Afr. trop.
angustifolia, Nutt. in Trans. Am. Phil. Soc. v. (1837) 172 = nepetaefolia.
arabica, Baill. Etud. Gén. Euph. 461 = mitis.
arborea, Comm. ex Baill. Adansonia, i. (1860–61) 267 = Acalypha arborea.
bahiensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 182.—Bras.
Bailloniana, *Muell. Arg. l. c.* 178.—Mexic.
betonicaefolia, *Muell. Arg. l. c.* 181.—Bras.
betonicaefolia, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 173.—Am. bor.
bicolor, *Miq. in Linnaea*, xxvi. (1853) 222.—Ind. or.
bicornis, Vahl, ex A. Juss. Euph. Tent. 49 = Sebastiania corniculata.
Boiviniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 183.—Madag.
bracteata, Blanco, Fl. Filip. ed. II. 481 = Pimelodendron amboinicum.
brevispica, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 262.—Texas.
burmanica, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 244.—Burma.
Camolia, Roxb. Hort. Beng. [103], nomen (Quid?).—Ind. or.
cannabina, Linn. f. Suppl. 415 = involucrata.
capensis, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 110 = Meyerianum.
capensis, *Thunb. Prod. Pl. Cap.* 14.—Afr. austr.
castaneaefolia, Juss. ex Baill. Adansonia, i. (1860–61) 267 = Acalypha marginata.
Chamaelea, Linn. Sp. Pl. 981 = Sebastiania Chamaelea.
chlorocaulon, *Baill. Etud. Gén. Euph.* 461; *et Adansonia*, v. (1864–65) 306.—Bras.
cissoides, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 406.—Bras.
colorata, Poir. Encyc. vii. 725 = Acalypha integrifolia.
cordata, Heyne, ex Wall. Cat. n. 7791 A = involucrata.
cordata, Michx. Fl. Bor. Am. ii. 176 = macrocarpa.
cordata, A. Rich. Tent. Fl. Abyss. ii. 244 = mitis.
cordata, Willd. Sp. Pl. iv. 322 = cordifolia, *Vahl.*
cordifolia, *Benth. in Hook. Niger Fl.* 501.—Afr. trop.
cordifolia, *Vahl, Symb. Bot.* i. 76.—Arabia.
corniculata, Vahl, Eclog. Am. ii. 55 = Sebastiania corniculata.
cuneata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 180.—Bras.
diffusa, Vell. Fl. Flum. x. t. 10 = volubilis.
dioica, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 95.—Ins. Socotra.
dioica, Sond. in Linnaea, xxiii. (1850) 109 = rupestris.
discolor, Muell. Arg. in DC. Prod. xv. II. 946 = innocua.
dodecandra, *Griseb. in Goett. Abh.* xix. (1874) 98.—Reg. Argent.

TRAGIA:—
elliptica, Hochst. in Flora, xxviii. (1845) 85 = Sapium indicum.
erosa, Hochst. l. c. 86 = Pyrenacantha scandens.
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 179. —Peruv.
Fendleri, *Muell. Arg. l. c.* 179.—Venezuela.
filiformis, Poir. Encyc. vii. 727 = Acalypha spiciflora.
fruticosa, Comm. ex Baill. Adansonia, i. (1860–61) 267 = Acalypha integrifolia.
furialis, Boj. Hort. Maurit. 286 = angustifolia.
Gayana, Baill. Etud. Gén. Euph. 461 = volubilis.
geraniifolia, *Klotzsch, ex Baill. l. c.*, nomen; *et ex Muell. Arg. in DC. Prod.* xv. II. 933.—Bras.
gracilis, *Griseb. in Goett. Nachr.* (1865) 176.—Cuba.
grandifolia, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 46 = Bernardia grandifolia.
haguensis, Goudot, ex Baill. Etud. Gén. Euph. 461 = volubilis.
hastata, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 407.—Bras.
hastata, Reinw. ex Hassk. Pl. Jav. Rar. 245 = Cnesmone javanica.
hexandra, Jacq. Enum. Pl. Carib. 31 = Platygyna urens.
Hildebrandtii, *Muell. Arg. in Bremen, Abh.* vii. (1880) 26.—Afr. trop.
hirsuta, *Blume, Bijdr.* 630.—Java.
hispida, Willd. Sp. Pl. iv. 323 = involucrata.
ibaguensis, Goudot, ex Muell. Arg. in DC. Prod. xv. II. 936 = volubilis.
incana, *Klotzsch, ex Baill. Etud. Gén. Euph.* 461, nomen; *et ex Muell. Arg. in DC. Prod.* xv. II. 935. —Bras.
infesta, Mart. ex D. Dietr. Syn. Pl. v. 256 = volubilis.
innocua, Blanco, Fl. Filip. ed. II. 479 = Alchornea villosa.
innocua, *Walt. Fl. Carol.* 229.—Am. bor.
integerrima, Hochst. in Flora, xxviii. (1845) 85 = Excoecaria reticulata.
integrifolia, Willd. ex Muell. Arg. in DC. Prod. xv. II. 948 = Acalypha integrifolia.
interrupta, *Spreng. Syst.* iii. 833.—Zeylan.
involucrata, *Linn. Sp. Pl.* 980.—As. trop.
japurensis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 404. —Bras.
Kirkiana, *Muell. Arg. in Flora*, xlvii. (1864) 538.—Afr. trop.
laciniata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 182.—Mexic.
lagoënsis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 409. —Bras.
laminularis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 183.—Afr. trop.
Lessertiana, *Muell. Arg. l. c.* 178.—Bras.
linearifolia, Ell. Sketch, ii. 563 = innocua.
lobata, Wall. Cat. n. 7796 = Acalypha integrifolia.
macrocarpa, *Willd. Sp. Pl.* iv. 323.—Am. bor.
macrophylla, Wall. Cat. n. 7796 = Cnesmone javanica.
Manniana, *Muell. Arg. in Flora*, xlvii. (1864) 436.—Afr. trop.
marginata, Poir. Encyc. vii. 725 = Acalypha marginata.
melochioides, *Griseb. in Goett. Abh.* xxiv. (1879) 60.—Reg. Argent.
Mercurialis, Linn. Sp. Pl. 980 = Claoxylon Mercurialis.
Mercurialis, Walt. Fl. Carol. 229 = urticaefolia.
mexicana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 181.—Mexic.
Meyeriana, *Muell. Arg. in DC. Prod.* xv. II. 938.—Afr. austr.
minor, Sond. in Linnaea, xxiii. (1850) 108 = rupestris.
Miqueliana, Muell. Arg. in DC. Prod. xv. II. 942 = bicolor.
mitis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 244.—Abyss.
moammarensis, *Baill. Etud. Gén. Euph.* 461.—Arab.
monadelpha, *Schum. & Thonn. Beskr. Guin. Pl.* 404. —Afr. trop.
monandra, Baill. Etud. Gén. Euph. 461 = volubilis.
montana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 183 = involucrata.
natalensis, Hochst. in Flora, xxviii. (1845) 85 = Excoecaria reticulata.

TRAGIA:—
natalensis, *Sond. in Linnaea*, xxiii. (1850) 107.—Afr. austr.
nepetaefolia, *Cav. Ic.* vi. 37. t. 557. f. 1.—Mexic.
nepetaefolia, Wawra, in Bot. Ergeb. Maxim. Bras. 34 = bahiensis.
Novae-Hollandiae, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 180.—Austral.
obtusata, Vahl, ex Baill. Adansonia, i. (1860–61) 267 = Acalypha integrifolia.
odorata, Steud. Nom. ed. II. ii. 696, sphalm. = Acalypha integrifolia.
ovata, *Parodi, in Anal. Soc. Cient. Argent.* xi. (1881) 51.—Reg. Argent.
pedicillaris, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 415. —Bras.
pedunculata, Beauv. Fl. Owar. i. 90. t. 54 = volubilis.
peltata, *Vell. Fl. Flum.* x. t. 6 (peliata).—Bras.
pilosa, Vell. l. c. t. 9 = Sebastiania corniculata.
pinnata, *A. Juss. Euph. Tent.* 48.—Bras.
plumosa, Desf. Tabl. ed. I. 207 = virgata.
Pohlii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 81. —Bras.
polyandra, *Vell. Fl. Flum.* x. t. 7.—Bras.
pruricus, Willd. ex Endl. Gen. Suppl. iv. III. 88 = Platygyna urens.
pungens, Muell. in DC. Prod. xv. II. 941 = cordifolia, *Vahl.*
ramosa, Torr. Bot. Mex. Bound. 200 = stylaris.
ramosa, Torr. in Ann. Lyc. N. York, ii. (1828) 245 = nepetaefolia.
reticulata, Poir. Encyc. vii. 725 = Acalypha arborea.
rugosa, Wall. Cat. n. 7794 = Cnesmone javanica.
rupestris, *Sond. in Linnaea*, xxiii. (1850) 108.—Afr. austr.
salviaefolia, Boj. ex Baill. Etud. Gén. Euph. 443 = Acalypha salviaefolia.
scandens, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 183.—Madag.
scutellariaefolia, Scheele, l. c. xxv. (1852) 587 = nepetaefolia.
Sellowiana, *Muell. Arg. l. c.* xxxiv. (1865–66) 179.—Bras.
senegalensis, *Muell. Arg. l. c.* 182.—Afr. trop.
Serra, Poepp. & Endl. Nov. Gen. et Sp. iii. 20. t. 223 = volubilis.
spathulata, *Benth. in Hook. Niger Fl.* 502.—Afr. trop.
stylaris, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 180.—Am. bor. occ.
subhastata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 20. t. 223.—Peruv.
tenuifolia, *Benth. in Hook. Niger Fl.* 502.—Afr. trop.
tenuis, Wall. Cat. n. 7787 = Acalypha brachystachya.
teucriifolia, Scheele, in Linnaea, xxv. (1852) 586 = nepetaefolia.
triandra, *Muell. Arg. in DC. Prod.* xv. II. 947.—Afr. austr.
triangularis, Vell. Fl. Flum. x. t. 8 = volubilis.
trifida, Wall. Cat. n. 7795 = involucrata.
tristis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 410.—Bras.
uberabana, *Muell. Arg. l. c.* 417.—Bras.
urens, Linn. Sp. Pl. ed. II. 1391 (nomen falsum) = innocua.
urticaefolia, Benth. Pl. Hartw. 14 = nepetaefolia.
urticaefolia, *Michx. Fl. Bor. Am.* ii. 176.—Am. bor.
villosa, Thunb. Prod. Fl. Cap. 14 = Acalypha decumbens.
virgata, *Lam. Illustr.* iii. 347.—Hab. ?
volubilis, *Linn. Sp. Pl.* 980.—Ind. occ.
volubilis, Michx. ex Baill. Etud. Gén. Euph. 460 = urticaefolia.

TRAGIOPSIS, Karst. in C. Koch & Fint. Wochenschr. ii. (1859) 5 = **Sebastiania**, Spreng. (Euphorb.).
fruticulosa, Karst. l. c. = S. corniculata.

TRAGIOPSIS, Pomel, Nouv. Mat. Fl. Atl. 139 (1874) = **Carum**, Rupp. (Umbell.).
dichotoma, Pomel, l. c. 140 = C. dichotomum.
scabriuscula, Pomel, l. c. 139 = C. dichotomum.

TRAGIUM, Spreng. Pl. Umbell. Prod. 26 (1813) = **Pimpinella**, Riv. (Umbell.).
anisoides, Steud. Nom. ed. II. ii. 696 = P. villosa.
Anisum, Link, Enum. Hort. Berol. i. 285 = P. Anisum.

TRAGIUM:—
apulum, Heynh. Nom. i. 814 = Pimpinella Tragium ?
aromaticum, Spreng. Pl. Umbell. Prod. 26 = P. aromatica.
Broteri, Spreng. l. c. 27 = P. villosa.
Buchii, Webb & Berth. Phyt. Canar. ii. 151 = P. Buchii.
canescens, Hoffm. Gen. Umbell. ed. I. 91 = P. Tragium.
Columnae, Spreng. Pl. Umbell. Prod. 26 = P. Tragium.
Cumbrae, Choisy, ex Link, in Buch, Beschr. Canar. Ins. 152 = P. Cumbrae.
Dendroselinum, Webb & Berth. Phyt. Canar. ii. 152 = P. Dendroselinum.
depressum, Sieber, ex Spreng. in Schult. Syst. vi. 392 = P. Tragium.
dichotomum, Link, Handb. i. 322 = Carum dichotomum.
eriocarpum, Schult. Syst. vi. 394 = P. eriocarpos.
glaucum, J. & C. Presl, Delic. Prag. 125 = P. Tragium.
Gussonii, J. & C. Presl, l. c. 126 = P. Gussonii.
hirtellum, Hochst. in Flora, xxiv. (1841) I. Intell. 28 = P. peregrina.
incanum, Choisy, ex DC. Prod. iv. 122 = P. Cumbrae.
palmetorum, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 696 = P. palmetorum.
peregrinum, Spreng. Pl. Umbell. Prod. 26 = P. peregrina.
petraeum, Hoffm. Gen. Umbell. ed. I. 91 = P. Tragium.
Schousboei, Spreng. in Schult. Syst. vi. 394 = P. villosa.
tauricum, Ledeb. Ind. Sem. Hort. Dorpat. (1819); et ex Link, Enum. Hort. Berol. i. 286 = P. peregrina.
tenue, Sieber, ex Link, l. c. = P. eriocarpos.
tenuifolium, Spreng. in Schult. Syst. vi. 393 = P. tenuifolia.
tortuosum, Link, Handb. i. 322 = P. villosa.
villosum, Spreng. Pl. Umbell. Prod. 26 = P. villosa.

TRAGOCERAS, Spreng. Syst. iii. 576 (1826) = seq.

TRAGOCEROS, H. B. & K. Nov. Gen. et Sp. iv. 248. t. 385 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 356.
flavicomum, *DC. Prod.* v. 533.—Mexic.
microglossum, *DC. l. c.*—Mexic.
peucedanifolium, Spreng. Syst. iii. 576 = Isostigma peucedanifolium.
Schiedeanum, *Less. in Linnaea*, ix. (1834) 269.—Mexic.
zinnioides, *H. B. & K. Nov. Gen. et Sp.* iv. 249. t. 385.—Mexic.

TRAGOPOGON, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 530.
GEROPOGON, Linn. Sp. Pl. ed. II. 1109 (1763).
acanthocarpus, *Boiss. Diagn.* Ser. I. xi. 45.—Armen.; Persia.
aculeatus, Moench, Meth. 539 = Urospermum picroides.
angustifolius, Bell. ex Willd. Sp. Pl. iii. 1494 = crocifolius.
asper, Linn. Sp. Pl. 790 = Urospermum picroides.
aureus, *Boiss. Fl. Orient.* iii. 754.—Pontus.
australis, Bourg. ex Nym. Consp. 462 = longirostris.
australis, Jord. Cat. Jard. Dijon (1848), 32 = porrifolius.
barbirostris, Bischoff, Del. Sem. Hort. Heidelb. (1846) 4.—Cf. Linnaea, xxiv. (1851) 161 = eriospermus ?
Baylei, Lecoq & Lamotte, Cat. Pl. Vasc. Fr. 246 = dubius.
bicolor, Moench, Meth. 539 = Urospermum Dalechampii.
Bischoffii ×, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 469, in obs. —Hab. ?
brachyphyllus, Nym. Consp. 462 = porrifolius.
brevifolius, C. Koch, in Linnaea, xxiii. (1850) 662 = persicus.
brevirostris, *DC. Prod.* vii. 114.—Rossia; Soongar.
buphthalmoides, Boiss. Fl. Orient. iii. 750 = persicus.
calyculatus, Jacq. Hort. Vindob. ii. 48 = Scorzonera hirsuta.
campestris, Bess. Enum. Pl. Vohl. 84 = dubius.
canus, Willd. Sp. Pl. iii. 1493 = floccosus.
capensis, Jacq. Coll. ii. 320 = Urospermum picroides.
caricifolius, Boiss. Diagn. Ser. I. vii. 4 = collinus.
carinatus, Gilib. Fl. Lituan. i. 227 = pratensis.
castellanus, *Levier, Deux Excurs. Bot.* 26.—Hispan.

TRAGOPOGON :—
caucasicus, Bess. ex DC. Prod. vii. 114 = brevirostris.
caucasicus, Fisch. & Mey. ex DC. l. c. 114 = graminifolius.
coelesyriacus, Boiss. Diagn. Ser. I. xi. 47 = longirostris.
collinus, *DC. Prod.* vii. 115.—As. Min. ; Persia.
coloratus, *C. A. Mey. Verz. Pfl. Cauc.* 61.—Reg. Cauc. ; Persia.
Crantzii ×, Dichtl, in Deutsche Bot. Monatsschr. i. (1883) 171.—Europ.
crocifolius, *Linn. Syst.* ed. X. 1191.—Europ. austr.
crocifolius, Sibth. & Sm. Fl. Graec. viii. 58. t. 779 = longirostris.
Cupani, *Guss. ex DC. Prod.* vii. 113.—Sicilia.
cuspidatus, Steud. Nom. ed. I. 843 = Troximon cuspidatum.
Dalechampii, Linn. Sp. Pl. 790 = Urospermum Dalechampii.
Dandelion, Linn. l. c. ed. II. 1111 = Krigia Dandelion.
dshimilensis, C. Koch, in Linnaea, xxiii. (1850) 663 = longirostris.
dubius, *Scop. Fl. Carn.* ed. II. ii. 95.—Europ.
elatior, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 407.—Tauria.
elatus, Salisb. Prod. 178 = dubius.
eriospermus, *Tenore, Fl. Med.* ii. 3 ; *ex ej. Fl. Nap.* iii. p. xi.—Ital.
erostris, *Boiss. & Haussk. in Fl. Orient.* iii. 754.—Kurdist.
filifolius, *Rehm. ex Boiss. l. c.*—Reg. Cauc.
floccosus, Cutand. Fl. Comp. Madrit. 439 = crocifolius.
floccosus, Tommas. ex Bischoff, Beitr. Fl. Deutschl. 97 = Tommasinii.
floccosus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 116. t. 112.—Europ. or. ; As. bor.
fritillarioides, Less. in Linnaea, vi. (1831) 101 = Picrosia longifolia.
glaber, Hill, Hort. Kew. 41 = hybridus.
glaucus, Steud. Nom. ed. I. 843 = Troximon glaucum.
Gorskianus, Reichb. f. Ic. Fl. Germ. xix. 19 = floccosus.
gracilis, *D. Don, in Mem. Wern. Soc.* iii. (1820) 414.—Reg. Himal.
graminifolius, *DC. Prod.* vii. 114.—Reg. Cauc. ; Persia.
grandiflorus, Saut. in Oestr. Bot. Wochenbl. vi. (1856) 107, in obs. = pratensis.
heterospermus, Schweigg. in Koenigsb. Arch. Naturw. i. (1812) 229 = floccosus.
hirsutus, Gouan, Fl. Monsp. 342 = Scorzonera hirsuta.
humilis, *Fisch. Hort. Gorenk.* ed. II. (1812) 34.—Reg. Cauc.
hybridus, *Linn. Sp. Pl.* 789.—Ital.
intermedius, Bess. Enum. Sem. Hort. Crem. (1821) ; ex ej. Enum. Pl. Volh. 83 = brevirostris.
jesdianus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 138.—Persia.
junceus, Wall. Cat. n. 3257 = gracilis.
Kotschyi, Boiss. Diagn. Ser. I. xi. 45 = graminifolius.
laetus, Salisb. Prod. 178 = Urospermum Dalechampii.
Lamottei, *Rouy, in Bull. Soc. Bot. Fr.* xxviii. (1881) p. lix.—Gallia.
lanatus, Linn. Sp. Pl. ed. II. 1111 = Scorzonera lanata.
latifolius, *Boiss. Diagn.* Ser. I. iv. 23.—As. Min.
leiocarpos, Saut. in Flora, xl. (1857) 178 = pratensis.
lithuanicus, Bess. ex DC. Prod. vii. 114 = brevirostris.
livescens, Bess. Enum. Pl. Volh. 30 = dubius.
longifolius, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. iii. 90.—Graecia.
longifolius, Lamotte, Prod. Fl. Centr. Fr. ii. 457 = Lamottei.
longipappus, Peterm. in Flora, xxvii. (1844) 478 = orientalis.
longirostris, *Bischoff, ex Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 469.—As. Min. ; Arab.
macropogon, C. A. Mey. in Bull. Soc. Nat. Mosc. (1838) 273 = dubius.
major, Jacq. Fl. Austr. i. 19 = dubius.
marginatus, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 136 = collinus, acanthocarpus.
minor, Billot, ex Nym. Consp. 463 = dubius.
minor, Fries, Nov. Fl. Suec. ed. II. 241 = pratensis.
minor, *Mill. Gard. Dict.* ed. VIII. n. 2.—Europ.
mutabilis, Hohen. in Bull. Soc. Nat. Mosc. (1838) 273 = collinus.
mutabilis, *Jacq. Ic. Pl. Rar.* i. 16. t. 157 ; *Misc.* ii. 316.—Reg. Cauc.

TRAGOPOGON :—
nebrodensis, *Guss. Fl. Sic. Syn.* ii. 384.—Sicil.
nervosus, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 261.—Syria.
nervulosus, Boiss. Diagn. Ser. I. xi. 47 = persicus.
novus, Geners. ex Steud. Nom. ed. I. 844 = orientalis.
olympicus, *Boiss. Diagn.* Ser. I. xi. 46.—As. Min.
orgyalis, Reut. Cat. Gr. Jard. Genèv. (1863) ; (1867) 4 = longirostris.
orientalis, *Linn. Sp. Pl.* 789.—Europ. ; As. bor.
orientalis, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 59 = pusillus.
palaestinus, Boiss. Diagn. Ser. I. xi. 46 = persicus.
parviflorus, Hornem. Hort. Hafn. ii. 970 = crocifolius.
perpusillus, *Arv. Ess. Pl. Dauph.* 39 (Quid ?).—Gallia.
persicus, *Boiss. Diagn.* Ser. I. vii. 4.—As. Min. ; Syria ; Persia.
phaenopappus, Fisch. ex Steud. Nom. ed. II. ii. 697 = coloratus.
Pichleri, *Boiss. Fl. Orient.* iii. 753.—Bithyn.
picroides, Linn. Sp. Pl. 790 = Urospermum picroides.
plantagineus, Boiss. & Huet, Diagn. Ser. II. iii. 91 = persicus.
podolicus, Bess. ex DC. Prod. vii. 114 = brevirostris.
pomifolius, J. F. Gmel. Syst. 1638, sphalm. = porrifolius.
Pommaretii, F. Schultz, in Flora, xlix. (1866) 52 = dubius.
porrifolius, *Linn. Sp. Pl.* 789.—Reg. Mediterr.
porrifolius, Pall. ex Bieb. Fl. Taur. Cauc. ii. 233 = dubius.
pratensis, *Linn. Sp. Pl.* 789.—Europ. ; As. Min. ; Ind. bor. occ.
pterocarpus, *DC. Prod.* vii. 112.—Syria.
pubescens, *Kit. in Schult. Oesterr. Fl.* ed. II. ii. 403.—Hungar.
pusillus, *Bieb. Fl. Taur. Cauc.* iii. 521.—Reg. Cauc. ; Persia.
reticulatus, *Boiss. & Huet, Diagn.* Ser. II. iii. 90.—Armenia.
revolutus, Schweigg. in Hagen, Chlor. Boruss. 288 = orientalis.
roseus, C. A. Mey. Verz. Pfl. Cauc. 61 = collinus.
roseus, Trevir. in Mag. Ges. Naturf. Fr. Berl. vii. (1816) 152 = ruber.
ruber, Eichw. Casp.-Cauc. 28 = collinus.
ruber, *S. T. Gmel. Reise*, ii. 198.—Sibir.
ruthenicus, *Bess. ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 234, nomen.—Turkestan.
Samaritani, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. v. 116.—Graecia.
sativus, Gater. Desc. Pl. Montaub. 136 = porrifolius.
Shuttleworthii, Godet, Fl. Jura, i. 401 = pratensis.
silesiacus, Krock. Fl. Siles. Suppl. ii. 224 (Quid ?).—Europ.
sinuatus, Avé-Lall. Pl. Ital. Bor. 17 = pratensis.
sonchifolius, Salisb. Prod. 178 = Urospermum picroides.
stenophyllus, Boiss. & Huet, Diagn. Ser. II. iii. 89 = reticulatus.
stenophyllus, Jord. Obs. Pl. Crit. vii. (1849) 42 = crocifolius.
talyschensis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 137 = graminifolius.
Tommasinii, *Sch. Bip. in Bischoff, Beitr. Fl. Deutschl.* 97.—Europ. austr.
tortilis, Pritz. Ic. Ind. ii. 275 = pratensis.
transsilvanicus, Schur, Enum. Pl. Transs. 361 = orientalis.
tuberosus, C. Koch, in Linnaea, xxiii. (1850) 662 = pusillus.
undulatus, Jacq. Ic. Pl. Rar. i. 16. t. 158 ; Misc. ii. 317 = orientalis.
undulatus, Thuill. Fl. Par. ed. II. 396 = pratensis.
verticillatus, Lam. Fl. Fr. ii. 77 = Urospermum Dalechampii.
villosus, [Habl.] Desc. Taur. 151 = orientalis.
villosus, *Linn. Sp. Pl.* ed. II. 1110 (sp. dub.).—Hispan.
villosus, Pall. Reise, ii. 332 = floccosus.
virginicus, Linn. Sp. Pl. 789 = Krigia amplexicaulis.
vulgaris, Gueldenst. Reisen, ii. 33 ; ex Ledeb. Fl. Ross. ii. 790 (Quid ?).—Reg. Cauc.

TRAGOPOGUM, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 109 = praec.

TRAGOPYRUM, Bieb. Fl. Taur. Cauc. iii. 284 (1819) = **Atraphaxis**, Linn. (Polygon.).
angustifolium, Endl. Gen. Suppl. iv. II. 54 = A. angustifolia.

TRAGOPYRUM :—
Aucherii, Endl. Gen. Suppl. iv. II. 54 = Atraphaxis Aucherii.
Billardieri, Endl. l. c. = A. Billardieri.
buxifolium, Bieb. Fl. Taur. Cauc. iii. 284 = A. buxifolia.
buxifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 741 = A. laetevirens.
canescens, Presl, Bot. Bemerk. 109 = A. canescens.
compactum, Presl, l. c. = A. compacta.
decipiens, Endl. Gen. Suppl. iv. II. 54 = A. decipiens.
frutescens, Radde, Reisen Ost-Sibir. 425 = A. Laxmanni ?
glaucum, Spreng. Syst. ii. 251 = A. Laxmanni.
grandiflorum, Bieb. Fl. Taur. Cauc. iii. 285 = A. grandiflora.
laetevirens, Ledeb. Fl. Alt. ii. 75 = A. laetevirens.
lanceolatum, Bieb. Fl. Taur. Cauc. iii. 285 = A. Laxmanni.
lanceolatum, Ledeb. Fl. Alt. ii. 73 = A. micrantha.
lanceolatum, Turcz. ex Meissn. in DC. Prod. xiv. 78 = A. pungens.
linearifolium, Auch. ex Meissn. l. c. 79 = A. angustifolia.
mucronatum, Hort. ex C. Koch, Dendrol. ii. 1. 361 = A. Laxmanni.
polygamum, Spreng. Syst. ii. 251 = Polygonella parvifolia.
pungens, Bieb. Fl. Taur. Cauc. iii. 285 = A. Laxmanni.
rotundifolium, Presl, Bot. Bemerk. 109 = A. spinosa.
spinosum, Presl, l. c. = A. spinosa.
suaedaefolium, Endl. Gen. Suppl. iv. II. 54 = A. suaedaefolia.
Tournefortii, Endl. l. c. = A. Tournefortii.

TRAGOSELINUM, Tourn. ex Hall. Enum. Stirp. Helv. ii. 428 (1742) = **Pimpinella**, Riv. (Umbell.).
Angelica, Lam. Fl. Fr. iii. 449 = Aegopodium Podagraria.
dissectum, Moench, Meth. 99 = P. magna.
hircinum, Moench, l. c. = P. Saxifraga.
luteum, Pomel, Nouv. Mat. Fl. Atl. 142 = P. lutea.
magnum, Moench, Meth. 99 = P. magna.
majus, Lam. Fl. Fr. iii. 448 = P. magna.
minus, Lam. l. c. 447 = P. Saxifraga.
pumilum, Lam. l. c. 448 = Trinia Kitaibelii, vulgaris.
saxifragum, Moench, Meth. 99 = P. Saxifraga.

TRAGOSMA, C. A. Mey. ex Ledeb. Fl. Ross. ii. 365 (1844) = **Cymbocarpum**, DC. (Umbell.).
foetidum, C. A. Mey. ex Ledeb. l. c. = C. anethoides.

TRAGULARIA, Koen. ex Roxb. Fl. Ind. ii. 217 (1832) = **Pisonia**, Plum. (Nyctag.).
horrida, Koen. ex Roxb. l. c. = P. aculeata.

TRAGUS, Hall. Hist. Stirp. Helv. ii. 203 (1768).
GRAMINEAE, Benth. & Hook. f. iii. 1122.
ECHISACHYS, Neck. Elem. iii. 228 (1790).
LAPPAGO, Schreb. Gen. 55 (1789).
NAZIA, Adans. Fam. ii. 31 (1763), nomen prius.
alienus, Schult. Mant. ii. 205 = racemosus.
Berteronianus, Schult. l. c. = racemosus.
biflorus, Schult. l. c. = racemosus.
brevicaulis, *Boiss. Diagn.* Ser. I. xiii. 44.—Persia.
decipiens, Boiss. Fl. Orient. v. 450 = brevicaulis.
Halleri, Doell, in Mart. Fl. Bras. ii. II. 122 = racemosus.
koelerioides, Aschers. in Verh. Bot. Ver. Brandenb. xx. (1878) p. xxx = racemosus ?
Latipes, J. Gay, ex Kunth, Enum. Pl. i. 171 = Latipes senegalensis.
muricatus, Moench, Meth. 53 = racemosus.
occidentalis, Nees, Agrost. Bras. 286 = racemosus.
orientalis, Neès, l. c. = racemosus.
racemosus, *Scop. Introd.* 73 ; *All. Fl. Pedem.* ii. 241.—Reg. Mediterr. ; Afghan.
racemosus, Steud. Nom. ed. II. ii. 697 = Deschampsia caespitosa.
senegalensis, J. Gay, ex Kunth, Enum. Pl. i. 171 = Latipes senegalensis.
tcheliensis, *Debeaux, in Act. Linn. Soc. Bord.* xxxiii. (1879) 71.—China.

TRAGUS, Panz. in Denkschr. Akad. Muench. 1813 (1814) 296 = **Festuca**, Tourn. (Gramin.).
elatior, Panz. l. c. = F. elatior.
gracilis, Panz. l. c. = Brachypodium sylvaticum.

TRAGUS :—
inermis, Panz. in Denkschr. Akad. Muench. 1813 (1814) 296 = Bromus inermis.
loliaceus, Panz. l. c. = Festuca elatior.
pinnatus, Panz. l. c. = Brachypodium pinnatum.
pratensis, Panz. l. c. = F. elatior ?
pumilus, Panz. l. c. = F. varia.
unioloides, Panz. l. c. = Bromus unioloides.
varius, Panz. l. c. = F. varia.

TRAILLIA, Lindl. ex Endl. Gen. Suppl. 1419 (1841) = **Schimpera**, Hochst. (Crucif.).

TRALLESIA, Zumag. Fl. Pedem. i. 361 (1849) = **Matricaria**, Tourn. (Compos.).
matricarioides, Zumag. l. c. = M. inodora.

TRALLIANA, Lour. Fl. Cochinch. 157 (1790). Genus dubium *CELASTRINEARUM*.—Cf. Benth. & Hook. f. i. 360.
scandens, *Lour. l. c.*—Cochinch.

TRAMBIS, Rafin. Fl. Tellur. iii. 87 (1836) = **Phlomis**, Linn. (Labiat.).
alpina, Rafin. l. c. = P. alpina.
tuberosa, Rafin. l. c. = P. tuberosa.

TRANKENIA, Thunb. Fl. Cap. ii. 250 (1818), err. typ. = **Frankenia**, Linn.

TRAPA, Linn. Gen. ed. I. 359 (1737). *ONAGRARIEAE*, Benth. & Hook. f. i. 793.
bicornis, Linn. f. Suppl. 128 = natans.
bispinosa, *Roxb. Hort. Beng.* 11; *Fl. Ind.* i. 428.—Geront. trop.
Castanea, Gilib. Fl. Lituan. ii. 127 = natans.
chinensis, Lour. Fl. Cochinch. 86 = natans.
cochinchinensis, *Lour. l. c.* 108.—Cochinchin.
hungarica, Opiz, ex Nym. Consp. 249 = natans.
incisa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 134.—Japon.
laevis, Presl, ex Nym. Consp. 249 = natans.
natans, *Linn. Sp. Pl.* 120.—Europ.; Oriens; Ind. or.
natans, Thunb. Fl. Jap. 65 = bispinosa.
quadricornis, Stokes, Bot. Mat. Med. i. 229 = natans.
quadrispinosa, Roxb. Hort. Beng. 11; Fl. Ind. i. 430 = natans.
quadrispinosa, Wall. Cat. n. 6340 = bispinosa.
verbanensis, *De Not. Cat. Sem. Ort. Bot. Roma* (1875), 28; *et in Nuov. Giorn. Bot. Ital.* viii. (1876) 42.—Ital.

TRAPAULOS, Reichb. Consp. 92 (1828), sphalm. = IRAUPALOS, Rafin. = **Viburnum**, Linn. (Caprifol.).

TRASIS, Beauv. in Lestib. Essai Fam. Cyp. 32 (1819) (Trasi) = **Cladium**, P. Br. (Cyperac.).

TRASUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 53 (1821) = **Carex**, Dill. (Cyperac.).
ampullaceus, S. F. Gray, l. c. 59 = C. ampullacea.
atratus, S. F. Gray, l. c. 56 = C. atrata.
binervis, S. F. Gray, l. c. 63 = C. binervis.
capillaris, S. F. Gray, l. c. 66 = C. capillaris.
chlorostachyos, S. F. Gray, l. c. 68 = C. Pseudocyperus.
clandestinus, S. F. Gray, l. c. 56 = C. humilis.
depauperatus, S. F. Gray, l. c. 63 = C. depauperata.
digitatus, S. F. Gray, l. c. 57 = C. digitata.
distans, S. F. Gray, l. c. 62 = C. distans.
erectus, S. F. Gray, l. c. 67 = C. sempervirens.
extensus, S. F. Gray, l. c. 61 = C. extensa.
filiformis, S. F. Gray, l. c. 59 = C. filiformis.
flavus, S. F. Gray, l. c. 60 = C. flava.
fulvus, S. F. Gray, l. c. 62 = C. fulva.
glaucus, S. F. Gray, l. c. 67 = C. glauca.
hirtus, S. F. Gray, l. c. 58 = C. hirta.
Hostianus, S. F. Gray, l. c. 61 = C. fulva.
laevigatus, S. F. Gray, l. c. 66 = C. laevigata.
limosus, S. F. Gray, l. c. 67 = C. limosa.
Oederi, S. F. Gray, l. c. 60 = C. Oederi.
pallescens, S. F. Gray, l. c. 69 = C. pallescens.
paludosus, S. F. Gray, l. c. 64 = C. paludosa.
paniceus, S. F. Gray, l. c. 63 = C. panicea.
pauciflorus, S. F. Gray, l. c. 56 = C. pauciflora.
pendulus, S. F. Gray, l. c. 65 = C. pendula.
piluliferus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 58 = Carex pilulifera.
praecox, S. F. Gray, l. c. 57 = C. praecox.
rariflorus, S. F. Gray, l. c. 68 = C. rariflora.
riparius, S. F. Gray, l. c. 64 = C. riparia.
strigosus, S. F. Gray, l. c. 65 = C. strigosa.
sylvaticus, S. F. Gray, l. c. 65 = C. sylvatica.
tomentosus, S. F. Gray, l. c. 57 = C. tomentosa.
ustulatus, S. F. Gray, l. c. 66 = C. ustulata.
vesicarius, S. F. Gray, l. c. 59 = C. vesicaria.

TRATTENIKIA, Pers. Syn. ii. 403 (1807) = **Marshallia**, Schreb. (Compos.).
angustifolia, Pers. l. c. = M. angustifolia.
lanceolata, Pers. l. c. = M. lanceolata.
latifolia, Pers. l. c. = M. latifolia.

TRATTINICKIA, Willd. Sp. Pl. iv. 975 (1806) (Trattinnickia). *BURSERACEAE*, Benth. & Hook. f. i. 326.
burseraefolia, *Mart. in Nov. Gen. et Sp.* iii. 93. t. 239.—Bras.
guianensis, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1188 = burseraefolia.
humilis, Spruce, ex Engl. in Mart. Fl. Bras. xii. II. 284 = burseraefolia.
rhoifolia, *Willd. Sp. Pl.* iv. 975.—Bras.
Ryanii, *Didr. in Kjoeb. Vidensk. Meddel.* (1857) 125.—Ins. Trinit.
Schomburgkiana, Klotzsch, ex Engl. in Mart. Fl. Bras. xii. II. 284 = seq.
Schomburgkii, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1188 = burseraefolia.

TRAUNSTEINERA, Reichb. Nom. 50 (1841); Fl. Sax. 87 (1844) = **Orchis**, Tourn.
globosa, Reichb. Fl. Sax. 87 = O. globosa.

TRAUTVETTERIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 22 (1835). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
grandis, Nutt. in Torr. & Gray, Fl. N. Am. i. 37 = palmata.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 184 = palmata.
palmata, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 22.—Am. bor. occ.; Japon.

TRAVERSIA, Hook. f. Handb. N. Zeal. Fl. 163 (1864) = **Senecio**, Tourn. (Compos.).
baccharoides, Hook. f. l. c. 164.—N. Zel.

TRAXARA, Rafin. Fl. Tellur. iv. 85 (1836) = **Lobostemon**, Lehm. (Borag.).
capitata, Rafin. l. c. 86 = L. capitatus.

TRAXILISA, Rafin. Sylva Tellur. 161 (1838) = **Tetracera**, Linn. (Dillen.).
aspera, Rafin. l. c. 162 = Tet. sarmentosa.

TRAXILUM, Rafin. Sylva Tellur. 42 (1838) = **Ehretia**, Linn. (Borag.).
asperatum, Rafin. l. c. = E. laevis.

TRECACORIS, Pritz. Ic. Ind. i. 1111 (1855), sphalm. = **Thecacoris**, A. Juss. (Euphorb.).

TRECHONAETES, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 350. *SOLANACEAE*, Benth. & Hook. f. ii. 898.
Bridgesii, Dun. in DC. Prod. xiii. I. 468 = laciniata.
laciniata, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 350.—Chili.
sativa, *Miers, l. c.* 352 (sp. dub.).—Reg. Argent.

TRECULIA, Decne. ex Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 108. t. 3. *URTICACEAE*, Benth. & Hook. f. iii. 374.
PSEUDOTRECULIA, Baill. Adansonia, xi. (1875) 292, in obs.
acuminata, *Baill. l. c.* 292.—Afr. trop.
africana, *Decne. ex Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 109.—Afr. trop.

TREICHELIA, Vatke, in Linnaea, xxxviii. (1874) 700. *CAMPANULACEAE*, Benth. & Hook. f. ii. 559.
LEPTOCODON, Sond. in Harv. & Sond. Fl. Cap. iii. 584 (1865).
longebracteata, *Vatke, in Linnaea*, xxxviii. (1874) 700.—Afr. austr.

TREISIA, Haw. Syn. Pl. Succ. 131 (1812) = **Euphorbia**, Linn.
Clava, Haw. l. c. = E. Clava.
erosa, Haw. Suppl. Pl. Succ. 66 = E. cereiformis.
Hystrix, Haw Syn. Pl. Succ. 131 = E. Hystrix.
tuberculata, Haw. Suppl. Pl. Succ. 65 = E. Clava.

TREISTERIA, Griff. Notul. iv. 109 (1854) = **Curanga**, Juss. (Scrophul.).
assamica, Griff. l. c. 111 = C. amara.
cordata, Griff. l. c. 109 = Torenia asiatica.
lanceolata, Griff. l. c. 111 (Toreniae sp.).—Burma.

TREMA, Lour. Fl. Cochinch. 562 (1790). *URTICACEAE*, Benth. & Hook. f. iii. 355.
SPONIA, Comm. ex Lam. Encyc. iv. 139 (1796).
affinis, Blume, Mus. Bot. Lugd. Bat. ii. 58 = orientalis.
africana, *Blume, l. c.*—Afr. trop.
amboinensis, *Blume, l. c.* 61.—As. et Austral. trop.
angustifolia, *Blume, l. c.* 58.—Malaya.
argentea, *Blume, l. c.*—Ins. Bonin.
aspera, *Blume, l. c.*—As. et Austral. trop.
Blancoi, *Blume, l. c.*—Ins. Philipp.
bracteolata, *Blume, l. c.*—Afr. austr.
Burmanni, Blume, l. c. 62 = amboinensis.
canescens, Blume, l. c. 58 = micrantha.
cannabina, *Lour. Fl. Cochinch.* 562 (= amboinensis ?).—Cochinchin.
carinata, *Blume, Mus. Bot. Lugd. Bat.* ii. 59.—Ins. Molucc.
Chichilea, *Blume, l. c.* 58.—Peruv.
Commersonii, Blume, l. c. 60 = orientalis.
discolor, *Blume, l. c.* 58.—Ind. or.
glabrescens, *Blume, l. c.*—Ins. Philipp.
glomerata, *Blume, l. c.* 58.—Afr. austr.
Griffithii, *Blume, l. c.* = amboinensis.
grisea, *Baker, in Journ. Linn. Soc.* xx. (1883) 263.—Madag.
imbricata, *Blume, Mus. Bot. Lugd. Bat.* ii. 63.—Ins. Molucc.
Lamarckiana, *Blume, l. c.* 58.—Ind. occ.
Lima, *Blume, l. c.*—Ind. occ.
macrophylla, *Blume, l. c.*—Am. austr.
Melinona, *Blume, l. c.* 64.—Guian.
micrantha, *Blume, l. c.* 58.—Am. trop.
mollis, *Blume, l. c.*—Ind. occ.; Venezuela.
morifolia, *Blume, l. c.* 59.—Ins. Celebes.
nitens, *Blume, l. c.* 58.—Afr. trop.
orientalis, *Blume, l. c.* 62.—Geront. trop.
pallida, *Blume, l. c.* 60.—Ins. Amboin.
politoria, *Blume, l. c.* 58.—Reg. Himal.
pubigera, *Blume, l. c.* 60.—Ins. Molucc.
rigida, *Blume, l. c.* 61.—Java.
riparia, *Blume, l. c.* 58.—Am. austr.
rufescens, *Blume, l. c.*—Jamaic.
rugosa, *Blume, l. c.*—Porto Rico.
scaberrima, *Blume, l. c.* 63.—Sumatra.
Schiedeana, Blume, l. c. 58 = micrantha.
Sieberi, Blume, l. c. = orientalis.
strigosa, *Blume, l. c.*—Afr. trop.
timorensis, *Blume, l. c.* 60.—Burma; Malaya.
velutina, Blume, l. c. 62 = amboinensis.
virgata, Blume, l. c. 59 = timorensis.
viridis, Blume, l. c. 58 = aspera.
Wightii, Blume, l. c. = orientalis.

TREMANDRA, R. Br. in Flind. Voy. App. ii. 544 (1814). *TREMANDREAE*, Benth. & Hook. f. i. 134.
diffusa, *R. Br. ex DC. Prod.* i. 344.—Austral.
Huegelii, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 647 = Tetratheca hirsuta.
oppositifolia, Steetz, in Lehm. Pl. Preiss. i. 222 = stelligera.
stelligera, *R. Br. ex DC. Prod.* i. 344.—Austral.
verticillata, Hueg. ex Walp. Ann. i. 76 = Platytheca galioides.

TREMANTHUS, Pers. Syn. i. 467 (1805) = **Styrax**, Tourn.
cordata, Pers. l. c. = S. cordatum.
ferruginea, Pers. l. c. = S. racemosum.
oblonga, Pers. l. c. = S. oblongum.
ovata, Pers. l. c. = S. ovatum.

TREMASPERMA, Rafin. Fl. Tellur. iv. 124 (1836) = **Ipomoea**, Linn. (Convolv.).
Bona-nox, Rafin. l. c. = I. Bona-nox.

TREMASTELMA, Rafin. Fl. Tellur. iv. 96 (1836) = **Scabiosa**, Tourn. (Dipsac.).
brachiata, Rafin. l. c. = S. brachiata.
Sibthorpi, Rafin. l. c. = S. brachiata.

TREMATOSPERMA, Urb. in Ber. Deutsch. Bot. Ges. i. (1883) 182. *OLACINEAE.*
cordatum, *Urb. l. c.—Cf. Eichl. Jahrb. Berl.* iii. (1884) 244.—Afr. trop.

TREMBLEYA, DC. Prod. iii. 125 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 738.
agrestis, DC. l. c. 126 = Microlicia agrestis.
calycina, *Cham. in Linnaea*, ix. (1834) 430.—Bras.
canescens, *Schnizl. Ic.* iv. t. 268. f. 1.—Hab. ?
Chamissoana, *Naud. in Ann. Sc. Nat.* Sér. III. xii. (1849) 270.—Bras.
heterostemon, *DC. Prod.* iii. 126.—Bras.
laniflora, Cogn. in Mart. Fl. Bras. xiv. III. 130 = T. Lychnitis.
Lychnitis, *DC. Prod.* iii. 12.—Bras.
neopyrenaica, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 154.—Bras.
paniculata, Naud. l. c. = heterostemon.
parviflora, Cogn. in Mart. Fl. Bras. xiv. III. 127 = heterostemon, triflora.
pentagona, *Naud. in Ann. Sc. Nat.* Sér. III. ii. (1844) 154.—Bras.
phlogiformis, *DC. Prod.* iii. 126.—Bras.
pithyoides, *Cham. in Linnaea*, ix. (1834) 428.—Bras.
pityrophylla, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 24 = Pyramia pityrophylla.
Pradosiana, *Netto, in Ann. Sc. Nat.* Sér. V. iii. (1865) 378. t. 12.—Bras.
revoluta, Naud. l. c. Sér. III. ii. (1844) 155 = calycina.
rhynanthera, Griff. Notul. iv. 677 = Melastoma malabathricum.
rosmarinoides, *DC. Prod.* iii. 125.—Bras.
Selloana, *Cogn. in Mart. Fl. Bras.* xiv. III. 133.—Bras.
stachyoides, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 154 = phlogiformis.
stenophylla, *Naud. l. c.* xii. (1849) 265.—Bras.
tridentata, *Naud. l. c.* ii. (1844) 154.—Bras.
triflora, *DC. Prod.* iii. 126.—Bras.
Warmingii, *Cogn. in Mart. Fl. Bras.* xiv. III. 133.—Bras.

TREMOTIS, Rafin. Sylva Tellur. 59 (1838) = **Ficus**, Tourn. (Urtic.).
cordata, Rafin. l. c. = F. Cunia.

TREMULA, Dum. in Van Hall, Bijdr. Wet. i. (1826) 46 = **Populus**, Linn. (Salicin.).
trepida, Rafin. Alsog. Am. 42 = P. tremuloides.
vulgaris, Opiz, Seznam, 99 = P. tremula.

TREMULARIA, Heist. Syst. 12 (1748) = **Briza**, Linn. (Gramin.).

TRENDELENBURGIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 123; et in Abh. Akad. Berl. 1854 (1855) 52. t. 3 B = **Begonia**, Tourn.
fruticosa, Klotzsch, ll. cc. = B. fruticosa.

TRENTEPOHLIA, Boeck. in Bot. Zeit. xvi. (1858) 249 = **Cyperus**, Mich.
bifoliata, Boeck, l. c. = C. Cephalotes.

TRENTEPOHLIA, Roth, Catalect. fasc. ii. 73 (1800) = **Heliophila**, Burm. f. (Crucif.).
integrifolia, Roth, l. c. 76 = H. amplexicaulis.
lepidioides, Roth, Nov. Pl. Sp. 326 (err. typ. 226) = H. diffusa.
pinnata, Roth, Catalect. fasc. ii. 73 = H. pendula.

TREPNANTHUS, Steud. Nom. ed. II. ii. 698 (1841), sphalm. = TERPNANTHUS, Nees & Mart. = **Spiranthera**, A. St. Hil. (Rutac.).

TREPOCARPUS, Nutt. ex DC. Mém. Ombell. 56 (1829); Prod. iv. 201 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 926.
ENTASIKON, Rafin. New Fl. Am. iv. 28 (1836).
Aethusae, *Nutt. ex DC. Mém. Ombell.* 56; *Prod.* iv. 201.—Am. bor.
brachycarpus, DC. ll. cc. 56; 202 = T. Aethusae.

TRESANTHERA, Karst. Fl. Columb. i. 37. t. 19 (1858) = **Rustia**, Klotzsch (Rubiac.).
condamineoides, Karst. l. c.—Venezuela.

TRESTEIRA, Hook. f. Fl. Brit. Ind. iv. 279 (1884), sphalm. = TREISTERIA, Griff. = **Curanga**, Juss. (Scrophul.).

TRETOCARYA, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 505. *BORAGINEAE.*
pratensis, *Maxim. l. c.*—China.

TRETORHIZA, Adans. Fam. ii. 503 (1763) = seq.

TRETORRHIZA, Renealm. ex Delarb. Fl. Auv. ed. II. i. 24 (1800) = **Gentiana**, Tourn.
Cruciata, Delarb. l. c. = G. Cruciata.

TREUIA, Stokes, Bot. Mat. Med. iv. 570 (1812) = **Trewia**, Linn. (Euphorb.).

TREUTLERA, Hook. f. in Hook. Ic. Pl. t. 1425 (1883). *ASCLEPIADEAE.*
insignis, *Hook. f. l. c.*—Reg. Himal.

TREVAUXIA, Steud. Nom. ed. II. ii. 698 (1841), sphalm. = TREVOUXIA, Scop. = **Luffa**, Tourn. (Cucurb.).

TREVESIA, Vis. in Giorn. Tosc. Sc. Med. Fis. e Nat. i. (1840) 72; et in Mem. Acc. Torin. Ser. II. iv. (1842) 262. *ARALIACEAE*, Benth. & Hook. f. i. 942.
PARAPANAX, Miq. Fl. Ind. Bat. Suppl. 388 (1860).
PETASULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. l. Art. IV. 3.
REYNOLDSIA, A. Gray, Bot. U. St. Expl. Exped. i. 723. t. 92/93 (1854).
eminens, *Bull, Cat.* (1884) 17.—Ins. Philipp.
insignis, Miq. Ann. Mus. Bot. Lugd. Bat. i. 220 = palmata.
littoralis, *Benth. & Hook. f. Gen.* i. 943.—Sumatra.
moluccana, *Miq. Fl. Ind. Bat.* i. 1. 748.—Ins. Amboin.
novo-guineënsis, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 26.—N. Guin.
palmata, *Vis. in Mem. Acc. Torin.* Ser. II. iv. (1842) 262.—Ind. or.; Burma.
pleiosperma, *Benth. & Hook. f. Gen.* i. 943.—Ins. Samoa.
sandwicensis, *Benth. & Hook. f. l. c.*—Ins. Sandvic.
sundaica, *Miq. Pl. Jungh.* 420.—Java.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 11.—Ins. Amboin.

TREVIA, Linn. Sp. Pl. ed. I. 1193 (1753) = **Trewia**, Linn. (Euphorb.).

TREVIRANA, Willd. Enum. Hort. Berol. 637 (1809) = **Achimenes**, P. Br. (Gesner.).
Andrieuxii, D. Dietr. Syn. Pl. iii. 568 = A. Andrieuxii.
candida, Decne. ex Lem. in Fl. des Serres, iv. (1848) t. 420 = Dicyrta candida.
coccinea, Willd. Enum. Hort. Berol. 638 = A. coccinea.
comifera, D. Dietr. Syn. Pl. iii. 568 (*conifera*) = A. comifera.
discolor, Poepp. & Endl. Nov. Gen. et Sp. iii. 8 = Diastema discolor.
divaricata, Poepp. & Endl. l. c. 9 = Diastema divaricatum.
erinoides, D. Dietr. Syn. Pl. iii. 568 = A. erinoides.
Escheri ×, Regel.—Cf. Linnaea, xxiv. (1851) 198.
grandiflora, Schiede, in Linnaea, viii. (1833) 247 = A. grandiflora.
heterophylla, Mart. Nov. Gen. et Sp. iii. 65 (err. typ. 56) = A. heterophylla.
hirsuta, D. Dietr. Syn. Pl. iii. 568 = A. hirsuta.
longiflora, D. Dietr. l. c. = A. longiflora.
maculata, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 36 = A. foliosa.
maculata, Poepp. & Endl. Nov. Gen. et Sp. iii. 8 = Diastema maculatum.
mollis, Poepp. & Endl. l. c. 9 = Diastema molle.
parviflora, Hook. & Arn. Bot. Beech. Voy. 302 = Anetanthus parviflora.
patens, Walp. Ann. i. 472, in nota = A. patens.
pulchella, Mart. Nov. Gen. et Sp. iii. 66. t. 226. f. 4 = A. coccinea.

TREVIRANA:—
scabra, Poepp. & Endl. Nov. Gen. et Sp. iii. 8 = Diastema scabrum.
tenella, D. Dietr. Syn. Pl. iii. 568 = Achimenes tenella.
tenerrima, Poepp. & Endl. Nov. Gen. et Sp. iii. 9. t. 207 C = Diastema tenerrimum.
urticaefolia, Poepp. & Endl. l. c. 8. t. 207 B = Diastema urticaefolium.

TREVIRANIA, Heynh. Nom. ii. 736 (1846) = **Psychotria**, Linn. (Rubiac.).
calycina, Heynh. l. c. = P. Lindleyana.

TREVIRANIA, Roth, in Weber, Beitr. ii. (1810) 123 = **Vandellia** ? Linn. (Scrophul.).
Gratiolae, Roth, l. c.—Hab. ?

TREVIRANIA, Spreng. Anleit. ii. 1. 400 (1817) = TREVIRANA, Willd. = **Achimenes**, P. Br. (Gesner.).

TREVOA, Miers, Trav. Chile, ii. 529 (1826). *RHAMNEAE*, Benth. & Hook. f. i. 384.
Berteroana, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 490 = Retanilla spinifera.
Berteroniana, *Steud. Nom.* ed. II. ii. 698.—Chili.
Closiana, Miers, in Ann. & Mag. Nat. Hist. Ser. III. v. (1860) 492 = Retanilla spinifera.
Frutilla, *Steud. & Hochst. ex Steud. Nom.* ed. II. ii. 698.—Chili.
quinquenervia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 158.—Chili.
tenuis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 489.—Chili.
trinervia, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 159 = seq.
trinervia, *Miers, Trav. Chile*, ii. 529.—Chili.
Weddelliana, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 491.—Bolivia.

TREVOUXIA, Scop. Introd. 152 (1777) = **Luffa**, Tourn. (Cucurb.).

TREWIA, Linn. Gen. ed. I. 152 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 318.
ROTTLERA, Willd. in Goett. Journ. Wiss. i. (1797) 7. t. 1.
africana, Baill. Adansonia, i. (1860–61) 68 = Claoxylon Barteri.
discolor, *Sm. in Rees, Cycl.* xxxvi. n. 4.—Ind. or.
hernandifolia, Roth, Nov. Pl. Sp. 374 = Macaranga indica ?
Inophyllum, Spreng. Syst. iii. 906 = Codiaeum Inophyllum.
insignis, Steud. Nom. ed. II. ii. 698 = Alchornea Blumeana.
integerrima, Stokes, Bot. Mat. Med. iv. 570 = nudiflora.
macrophylla, Blume, Bijdr. 612 = Alchornea Blumeana.
macrophylla, Roth, Nov. Pl. Sp. 373 = nudiflora.
macrostachya, Klotzsch, in Bot. Ergeb. Waldem. Reise, 117 = nudiflora.
nudiflora, *Linn. Sp. Pl.* 1193.—As. trop.
nudiflora, Wight, Ic. t. 1871, partim = polycarpa.
polycarpa, *Benth. & Hook. f. Gen.* iii. 318.—Ind. or.
pubescens, *Sm. in Rees, Cycl.* xxxvi. n. 3.—Ins. Amboin.
rusciflora, Heyne, ex Roth, Nov. Pl. Sp. 374 (Quid ?).—Ind. or.
tricuspidata, Willd. Sp. Pl. iv. 835 = Mallotus cochinchinensis.

TRIACHNE, Cass. in Bull. Soc. Philom. (1818) 48 = **Nassauvia**, Comm. (Compos.).
pygmaea, Cass. l. c. = N. pygmaea.

TRIACHYRIUM, Benth. in Journ. Linn. Soc. xix. (1881) 86, sphalm. = seq.

TRIACHYRUM, Hochst. in Flora, xxiv. (1841) 1. Intell. 19, nomen; et ex Steud. Syn. Pl. Gram. 176 (1855) = **Sporobolus**, R. Br. (Gramin.).
adoënse, Hochst. ll. cc.—Abyss.
cordofanum, Hochst. l. c. xxv. (1842) 1. Beibl. 134, nomen; et ex Steud. Syn. Pl. Gram. 176.—Nubia.
discosporum, Steud. l. c.—Afr. austr.
longifolium, Hochst. ex Steud. l. c.—Abyss.
micranthum, Steud. l. c.—Abyss.
nilagiricum, Steud. l. c. = S. piliferus.

TRIACIS, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 189 = **Turnera**, Linn.
microphylla, Griseb. l. c. = Turnera diffusa.

TRIACMA, Van Hass. ex Miq. Fl. Ind. Bat. ii. 523 (1857) = **Hoya**, R. Br. (Asclep.).

TRIACTINA, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 90. *CRASSULACEAE*, Benth. & Hook. f. i. 661.
verticillata, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 103.—Reg. Himal.

TRIADENIA, Miq. Fl. Ind. Bat. ii. 458 (1856) = **Trachelospermum**, Lem. (Apocyn.).
verrucosa, Miq. l. c. 459.—Java.

TRIADENIA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 172, 354 = **Hypericum**, Tourn.
aegyptiaca, Boiss. Fl. Orient. i. 783 = H. aegypticum.
maritima, Boiss. l. c. 784 = H. maritimum.
microphylla, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 173 = H. aegypticum.
Russeggeri, Fenzl, Pugill. Pl. Nov. Syr. 7.—Syria.
Sieberi, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 175 = H. maritimum.
thymifolia, Spach, l. c. = H. Spachianum.
Webbii, Spach, l. c. = H. aegypticum.

TRIADENUM, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Hypericum**, Tourn.
purpurascens, Rafin. in Desv. Journ. Bot. ii. (1809) 171 = H. virginicum.
purpureum, Rafin. in Med. Repos. N. York, v. (1808) 352 = H. virginicum.
virginicum, Rafin. Fl. Tellur. iii. 79 = H. virginicum.

TRIADICA, Lour. Fl. Cochinch. 610 (1790) = **Sapium**, P. Br. (Euphorb.).
chinensis, Spreng. Syst. i. 98, sphalm. = *sinensis*.
cochinchinensis, Lour. Fl. Cochinch. 610 = Excoecaria Loureiroana.
japonica, Baill. Etud. Gén. Euph. 512 = Excoecaria japonica.
sinensis, Lour. Fl. Cochinch. 610 = S. Sebifera.

TRIAENA, H. B. & K. Nov. Gen. et Sp. i. 178 [t. 61] (1815) = **Bouteloua**, Lag. (Gramin.).
racemosa, H. B. & K. l. c. 179 [t. 61] = B. Triathera.

TRIAENANTHUS, Nees, in DC. Prod. xi. 169 (1847) = **Strobilanthes**, Blume (Acanth.).
Griffithianus, Nees, l. c. = S. Griffithianus.

TRIAINA, H. B. & K. Nov. Gen. et Sp. i. t. 61 (1815), sphalm. = TRIAENA, H. B. & K. = **Bouteloua**, Lag. (Gramin).

TRIAINOLEPIS, Hook. f. in Benth. & Hook. f. Gen. ii. 126 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
africana, *Hook. f. l. c.*—Afr. trop.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 230.—Afr. trop.

TRIALLOSIA, Rafin. Fl. Tellur. iii. 60 (1836) = **Lachenalia**, Jacq. (Liliac.).
pallida, Rafin. l. c. = L. pallida.

TRIANAEA, Planch. & Linden, in Linden, Prix-Cour. n. 8 (1853) 4.—Cf. Bot. Zeit. xi. (1853) 717 = **Dyssochroma**, Miers (Solan.).
nobilis, Planch. & Linden, ll. cc.—N. Granat.

TRIANEA, Karst. in Linnaea, xxviii. (1856) 424 = **Limnobium**, Rich. (Hydroch.).
bogotensis, Karst. l. c. 425.—N. Granat.

TRIANOPTILES, Fenzl, in Endl. Gen. 113 (1836) = **Ecklonea**, Steud. (Cyperac.).
capensis, Harv. Gen. S. Afr. Pl. ed. II. 422 = E. capensis.

TRIANOSPERMA, Mart. Syst. Mat. Med. Bras. 79 (1843) = **Cayaponia**, Silva Manso (Cucurb.).
africana, Hook. f. in Oliver, Fl. Trop. Afr. ii. 568 = C. latebrosa.
angustiloba, Cogn. in Mart. Fl. Bras. vi. iv. 84 = C. angustiloba.
arguta, Mart. Syst. Mat. Med. Bras. 80, nomen.—Bras.

TRIANOSPERMA:—
attenuata, Hemsl. Biol. Centr. Am. Bot. i. 486 = Cayaponia attenuata.
Belangerii, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 201. t. 11 = C. americana.
Boykinii, M. Roem. Syn. Pepon. 45 = C. Boykinii.
Crugerii, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 193 = C. Crugeri.
diversifolia, Cogn. in Mart. Fl. Bras. vi. iv. 92. t. 28 = C. diversifolia.
ficifolia, Cogn. l. c. 89 = C. ficifolia.
ficifolia, Mart. Syst. Mat. Med. Bras. 79 = C. Martiana.
floribunda, Cogn. in Mart. Fl. Bras. vi. iv. 88. t. 26 = C. floribunda.
glandulosa, Mart. Syst. Mat. Med. Bras. 80 = C. glandulosa.
graciliflora, Griseb. Cat. Pl. Cub. 112 = C. americana.
gracillima, Cogn. in Mart. Fl. Bras. vi. iv. 91 = C. gracillima.
heterophylla, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 194 = C. heterophylla.
Hilariana, Naud. l. c. Sér. V. vi. (1866) 14 = C. ficifolia.
Lhotzkyana, Cogn. in Mart. Fl. Bras. vi. iv. 83 = C. Lhotzkyana.
Martiana, Cogn. l. c. 87. t. 25 = C. Martiana.
piauhiensis, Cogn. l. c. 83 = C. piauhiensis.
racemosa, Griseb. Cat. Pl. Cub. 112 = C. racemosa.
rigida, Cogn. in Mart. Fl. Bras. vi. iv. 94 = C. rigida.
setulosa, Cogn. l. c. 86 = C. setulosa.
simplicifolia, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 195 = C. simplicifolia.
Tayuya, Mart. Syst. Mat. Med. Bras. 80 = C. Tayuya.
Tibiricae, Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 191 = C. Tibiricae.
triangularis, Cogn. in Mart. Fl. Bras. vi. iv. 86 = C. triangularis.
trifoliolata, Cogn. l. c. 93 = C. trifoliolata.
trilobata, Cogn. l. c. 89. t. 27 = C. trilobata.

TRIANTHA, Baker, in Journ. Linn. Soc. xvii. (1879) 490 = **Tofieldia**, Huds. (Liliac.).
glutinosa, Baker, l. c. = Tof. glutinosa.
japonica, Baker, l. c. 491 = Tof. japonica.
pubens, Baker, l. c. 490 = Tof. pubens.

TRIANTHAEA, Spach, Hist. Veg. Phan. x. 39 (1841) = **Vernonia**, Schreb. (Compos.).

TRIANTHEMA, Sauv. Meth. Fol. 127 (1751); Linn. Sp. Pl. ed. I. 223 (1753). *FICOIDEAE*, Benth. & Hook. f. i. 855.
ANCISTROSTIGMA, Fenzl, in Ann. Wien. Mus. ii. (1838) 293.
DIPLOCHONIUM, Fenzl, Nov. Stirp. Dec. 57 (1839).
PAPULARIA, Forsk. Fl. Aegypt. Arab. 69 (1775).
PORTULACASTRUM, Juss. ex Medic. Phil. Bot. i. 99 (1789).
REME, Adans. Fam. ii. 245 (1763).
ROCAMA, Forsk. Fl. Aegypt. Arab. 71 (1775).
ZALEYA, Burm. f. Fl. Ind. 110. t. 31 (1768).
americana, Gill. in Edinb. Journ. Nat. & Geog. Sc. iii. (1831) 354 = Sesuvium Portulacastrum.
anceps, Thunb. Prod. Pl. Cap. 80 = Acrosanthes fistulosa.
corymbosum, E. Mey. in Drège, Zwei Pfl. Docum. 226, nomen = crystallina.
crystallina, *Vahl, Symb. Bot.* i. 32.—As. et Afr. trop.
cypseleoides, *Benth. Fl. Austral.* iii. 331.—Austral.
decandra, *Linn. Mant.* i. 70.—As. trop.
diffusa, Mill. Gard. Dict. ed. VIII. n. 2 = decandra.
dubia, Spreng. ex Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 179, in syn. = Adenocline pauciflora.
flexuosa, Schum. & Thonn. Beskr. Guin. Pl. 221 = T. Portulacastrum.
fruticosa, Vahl, Symb. Bot. i. 32 = Gymnocarpos fruticosus.
glaucifolia, F. Muell. Fragm. i. 172 = crystallina.
glinoides, Pers. Syn. i. 486 = decandra.
glossostigma, *F. Muell. in Wing, South. Sc. Record*, iii. (1883) 282.—Austral.
Govindia, Buch.-Ham. in Wall. Cat. n. 6838 = pentandra.
humifusa, Thunb. Prod. Pl. Cap. 80 = Acrosanthes humifusa.
humillima, *F. Muell. Fragm.* x. 72.—Austral.
hydaspica, *Edgew. in Journ. Linn. Soc.* vi. (1862) 203.—Ind. or.; Afr. trop.
monogyna, Linn. Mant. i. 69 = T. Portulacastrum.

TRIANTHEMA:—
obcordata, Roxb. Hort. Beng. 34; Fl. Ind. ii. 445 = T. Portulacastrum.
obcordata, Wall. Cat. n. 6837 F = pentandra.
oxycalyptra, *F. Muell. Fragm.* i. 173.—Austral.
parvifolium, *E. Mey. in Drège, Zwei Pfl. Docum.* 226, nomen; *et ex Harv. & Sond. Fl. Cap.* ii. 598.—Afr. austr.
pentandra, *Linn. Mant.* i. 70.—As. et Afr. trop.
pilosa, *F. Muell. Fragm.* i. 174.—Austral.
polyandra, Blume, Bijdr. 1137 = Sesuvium Portulacastrum.
polysperma, Hochst. ex Oliver, Fl. Trop. Afr. ii. 588 = hydaspica.
Portulacastrum, *Linn. Sp. Pl.* 223; *Moench, Meth.* 700.—Am. trop.
procumbens, Mill. Gard. Dict. ed. VIII. n. 1 = T. Portulacastrum.
rhynchocalyptra, *F. Muell. Fragm.* i. 174.—Austral.
rubens, E. Mey. in Drège, Zwei Pfl. Docum. 227, nomen = crystallina.
salsoloides, *Fenzl, ex Oliver, Fl. Trop. Afr.* ii. 588.—Afr. trop.
sedifolia, Vis. Pl. Aegypt. Dec. 19. t. 3. f. 1 = crystallina.
triquetra, Rottl. & Willd. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 181 = crystallina.
turgidifolia, *F. Muell. Fragm.* x. 83.—Austral.

TRIANTHIUM, Desv. Opusc. 69, in obs. (1831) = **Chrysopogon**, Trin. (Gramin.).

TRIANTHUS, Hook. f. Fl. Antarct. ii. 320 (1846) = **Nassauvia**, Comm. (Compos.).
ulicinus, Hook. f. l. c.—Patagon.

TRIAS, Lindl. in Wall. Cat. n. 1977 (1829); Gen. et Sp. Orch. 60 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 505.
oblonga, *Lindl. ll. cc.*—Burma.
ovata, *Lindl. Gen. et Sp. Orch.* 60.—Ind. or.

TRIASCIDIUM, Benth. & Hook. f. Gen. i. 876 (1867), sphalm. = TRISCIADIUM, Phil. = **Huanaca**, Cav. (Umbell.).

TRIASEKIA, G. Don, in Sweet, Hort. Brit. ed. III. 792 (1839), sphalm. = JIRASEKIA, F. W. Schmidt = **Anagallis**, Tourn. (Primul.).

TRIASPIS, Burch. Trav. ii. 280 (1824). *MALPIGHIACEAE*, Benth. & Hook. f. i. 259.
auriculata, *Radlk. in Bremen, Abh.* viii. (1883) 379.—Afr. trop.
flabellaria, *A. Juss. in Arch. Mus. Par.* iii. (1843) 507.—Afr. trop.
floribunda, *O. Hoffm. Sert. Madag.* 8.—Madag.
hypericoides, *Burch. Trav.* ii. 280.—Afr. austr.
lateriflora, *Oliver, Fl. Trop. Afr.* i. 281.—Afr. trop.
macropteron, *Welw. ex Oliver, l. c.*—Afr. trop.
mozambica, *A. Juss. in Arch. Mus. Par.* iii. (1843) 505.—Afr. trop.
Nelsoni, *Oliver, in Hook. Ic. Pl.* t. 1418.—Afr. austr.
odorata, *A. Juss. in Deless. Ic. Sel.* iii. 21. t. 36.—Afr. trop.
squarrosa, *Radlk. in Bremen, Abh.* viii. (1883) 377.—Afr. trop.
stipulata, *Oliver, Fl. Trop. Afr.* i. 281.—Afr. trop.

TRIATHERA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = **Bouteloua**, Lag. (Gramin.).
americana, Desv. l. c. = B. Triathera.
bromoides, Roth, ex Kunth, Enum. Pl. i. 520 = Enteropogon melicoides.
juncea, Desv. ex Beauv. Agrost. 40 = B. Triathera.
racemosa, Desv. Opusc. 72 = B. Triathera.

TRIBELES, Phil. in Linnaea, xxxiii. (1864–65) 307. *PITTOSPOREAE* [vel *SAXIFRAGEAE*], Benth. & Hook. f. i. 973.
CHALEPOA, Hook. f. in Hook. Ic. Pl. t. 1082 (1871).
australis, *Phil. in Linnaea*, xxxiii. (1864–65) 307.—Ins. Chiloe.

TRIBLEMMA, R. Br. ex DC. Prod. iii. 113 (1828) = **Bertolonia**, Raddi (Melastom.).
acaule, Steud. Nom. ed. II. ii. 698 = Eriocnema Hilariana.
leuzeanum, Steud. l. c. = B. leuzeana.
longifolium, Steud. l. c. = B. longifolia.

TRIBLEMMA :—
maculata, Mart. ex DC. Prod. iii. 114 = Bertolonia maculata.
nymphaeifolium, Mart. ex DC. l. c. 113 = B. ovata.
nymphaeifolium, Steud. Nom. ed. II. ii. 698 = B. nymphaeaefolia.
ovatum, Steud. l. c. = B. ovata.

TRIBOLACIS, Griseb. Fl. Brit. W. Ind. 297 (1860) = **Turnera**, Linn.
juncea, Griseb. l. c. = Turnera guianensis.

TRIBOLIUM, Desv. Opusc. 64 (1831) = **Lasiochloa**, Kunth (Gramin.).
hispidum, Desv. l. c. = L. hispida.

TRIBONANTHES, Endl. Nov. Stirp. Dec. 27 (1839). *HAEMODORACEAE*, Benth. & Hook. f. iii. 675.
australis, *Endl. l. c.*—Austral.
brachypetala, *Lindl. Swan River App.* 44.—Austral.
Lindleyana, Endl. in Lehm. Pl. Preiss. ii. 27 = longipetala.
longipetala, *Lindl. Swan River App.* 44.—Austral.
odora, Endl. in Lehm. Pl. Preiss. ii. 28 = brachypetala.
uniflora, *Lindl. Swan River App.* 44.—Austral.
variabilis, *Lindl. l. c.*—Austral.
violacea, Endl. in Lehm. Pl. Preiss. ii. 28 = uniflora.

TRIBRACHIA, Lindl. Bot. Reg. sub t. 832 (1824); et Coll. Bot. t. 41 (1825) = **Bulbophyllum**, Thou. (Orchid.).
biflora, Lindl. Coll. Bot. sub t. 41 (= *Stelis biflora*).—Reg. Himal.
hirta, Lindl. l. c. = B. hirtum.
odoratissima, Lindl. l. c. = B. odoratissimum.
pendula, Lindl. l. c. = B. pendulum.
purpurea, Lindl. l. c. = B. Careyanum.
racemosa, Lindl. l. c. = Liparis odorata?
reptans, Lindl. l. c. = B. reptans.

TRIBRACHIUM, Benth. & Hook. f. Gen. iii. 501 (1863), sphalm. = praec.

TRIBRACHYA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 254. *RUBIACEAE*, Benth. & Hook. f. ii. 118.
morindaeformis, *Korth. l. c.* 255.—Sumatra.

TRIBRACHYS, Champ. ex Thw. Enum. Pl. Zeyl. 325 (1864) = **Thismia**, Griff. (Burman.).
Gardneriana, Champ. ex Thw. l. c. = Thismia Gardneriana.

TRIBULA, Hill, Veg. Syst. vi. 10 (1764) = **Athamanta**, Linn. (Umbell.).
Turbith, Hill, Hort. Kew. 91 = A. Matthioli.

TRIBULARIA, Heist. Syst. 8 (1748) (Quid?).

TRIBULASTRUM, B. Juss. ex Linn. Gen. ed. II. 185 (1742) = **Neurada**, Linn. (Rosac.).

TRIBULOPIS, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 70 (1849) = **Tribulus**, Linn. (Zygoph.).
angustifolia, R. Br. l. c. = Tribulus angustifolius.
bicolor, F. Muell. Fragm. i. 47 = Tribulus bicolor.
pentandra, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 70 = Tribulus Brownii.
Solandri, R. Br. l. c. = Tribulus Solandri.
Solandri, F. Muell. Fragm. i. 47, partim = Tribulus angustifolius.

TRIBULOPSIS, F. Muell. Fragm. i. 47 (1858) = praec.

TRIBULUS, Tourn. ex Linn. Syst. ed. I (1735). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 264.
EHRENBERGIA, Mart. Nov. Gen. et Sp. ii. 72. t. 163 (1827).
HETEROZYGIS, Bunge, Suppl. Fl. Alt. 82 (1836).
KALLSTROEMIA, Scop. Introd. 212 (1777).
TRIBULOPIS, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 70 (1849).
acanthococcus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 9 = terrestris.
adscendens, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 245.—Ins. Galapag.

TRIBULUS:—
alatus, *Delile, Fl. Aeg. Illustr.* 62.—Afr. bor.; Ind. or.
alatus, Drège, ex Presl, Bot. Bemerk. 30 = pterophorus.
albus, Poir. Encyc. viii. 45 = terrestris.
angustifolius, *Benth. Fl. Austral.* i. 290.—Austral.
astrocarpus, *F. Muell. Fragm.* xii. 4.—Austral.
bicolor, *F. Muell. Pl. Vict.* i. 99.—Austral.
bicornutus, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 391 = terrestris.
bimucronatus, Kralik, in Ann. Sc. Nat. Sér. III. xi. (1849) 28, sphalm. = praec.
bimucronatus, *Viv. Pl. Aegypt. Dec.* iv. 9.—Aegypt.; Afghan.
bispinulosus, Kralik, in Ann. Sc. Nat. Sér. III. xi. (1849) 26 = bimucronatus?
brasiliensis, *Spreng. Syst.* iv. *Cur. Post.* 343.—Bras.
Brownii, *F. Muell. Pl. Vict.* i. 99.—Austral.
californicus, *S. Wats. in Proc. Am. Acad.* xi. (1876) 125.—Calif.
cistoides, *Linn. Sp. Pl.* 387.—Cosmop. trop.
cristatus, Ehrenb. ex C. Muell. in Walp. Ann. iv. 404 = Ehrenbergii.
cristatus, *Presl, Bot. Bemerk.* 30.—Afr. austr.
decolor, Macfad. Fl. Jamaic. i. 186 = maximus.
dubescens, Walp. Ann. ii. 244, err. typ. = *pubescens*.
Ehrenbergii, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* i. 249.—Afr. trop.
excrucians, Wawra & Peyr. in Sitzb. Akad. Wien, Math.-Nat. xxxviii. (1860) 557 = terrestris.
Fisheri, Kellogg, in Proc. Calif. Acad. vii. (1877) 162 = grandiflorus.
Forrestii, *F. Muell. in Wing, South. Sc. Record*, (Nov. 1885).—Austral.
grandiflorus, *Benth. & Hook. f. ex S. Wats. Ind. Bibliog. N. Am. Bot.* 149.—Am. bor. occ.
Gussonii, Todaro & Pirain. ex Parl. Fl. Ital. v. 334 = terrestris.
hirsutus, *Benth. Fl. Austral.* i. 289.—Austral.
hispidus, Presl, Bot. Bemerk. 29 = terrestris.
humifusus, Schum. & Thonn. Beskr. Guin. Pl. 215 = terrestris.
Hystrix, *R. Br. in Sturt, Exped. Centr. Austral.* ii. *App.* 69.—Austral.
inermis, Kralik, in Ann. Sc. Nat. Sér. III. xi. (1849) 25 = bimucronatus?
intermedius, Kralik, l. c. 27 = bimucronatus.
Kotschyanus, Boiss. Diagn. Ser. II. i. 111 = terrestris.
lanatus, Walp. Ann. ii. 243 = T. Hystrix.
lanuginosus, Blanco, Fl. Filip. ed. I. 350 = cistoides.
lanuginosus, Linn. Sp. Pl. 387 = terrestris.
littoralis, Ait. ex Sweet, Hort. Brit. ed. III. 766 (Quid?).—Austral.
longipetalus, Viv. Pl. Aegypt. Dec. iv. 10 = alatus.
macranthus, Hassk. in Flora, xlviii. (1865) 403 = cistoides.
macrocarpus, *F. Muell. ex Benth. Fl. Austral.* i. 289.—Austral.
macropterus, *Boiss. Diagn.* Ser. I. i. 161.—Aegypt.; Persia; Afghan.
maximus, *Linn. Sp. Pl.* 386.—Am. calid.
megistopterus, *Kralik, in Ann. Sc. Nat.* Sér. III. xi. (1849) 32.—Arab.
micans, Welw. Apont. 566 = terrestris.
microcephalus, *Bertol. f. in Mem. Acc. Sc. Bologn.* iii. (1851) 255.—Afr. trop.
minutus, *Leichh. ex Benth. Fl. Austral.* i. 291.—Austral.
mollis, Ehrenb. in Schweinf. Beitr. Fl. Aethiop. i. 29 = terrestris.
moluccanus, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 446 = cistoides.
Murex, Presl, Bot. Bemerk. 29 = terrestris.
muricatus, Stokes, Bot. Mat. Med. ii. 496 = terrestris.
occidentalis, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 69 = T. Hystrix.
parvispinus, Presl, Bot. Bemerk. 29 = terrestris.
pentandrus, Benth. Fl. Austral. i. 290 = Brownii.
pentandrus, *Forsk. Fl. Aegypt. Arab.* 81 (sp. dub.).—Arab.
persicus, Kralik, in Ann. Sc. Nat. Sér. III. xi. (1849) 31 = macropterus.
platypterus, *Benth. Fl. Austral.* i. 289.—Austral.
pterocarpus, Ehrenb. ex C. Muell. in Walp. Ann. iv. 404 = macropterus.
pterophorus, *Presl, Bot. Bemerk.* 30.—Afr. austr.
pubescens, G. Don, Gen. Syst. i. 769 = maximus.

TRIBULUS:—
ranunculiflorus, *F. Muell. Fragm.* i. 48.—Austral.
Revoili, *Franch. in Revoil, Çomali*, 22.—Afr. trop.
robustus, Boiss. & Noé, Diagn. Ser. II. i. 112 = terrestris.
sericeus, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 245.—Ins. Galapag.
sinaicus, Boiss. Diagn. Ser. I. viii. 120 = bimucronatus.
Solandri, *F. Muell. Pl. Vict.* i. 99, partim.—Austral.
Solandri, F. Muell. l. c., partim = angustifolius.
spurius, Kralik, in Ann. Sc. Nat. Sér. III. xi. (1849) 28 = bimucronatus.
subinermis, *Fisch. Cat. Hort. Gorenk.* (1808) 94.—Tibet.
terrestris, *Linn. Sp. Pl.* 387.—Amphig. trop.
terrestris, Muhl. Cat. 43 = maximus.
trijugatus, Nutt. Gen. Am. i. 277 = maximus.
trijugus, Steud. Nom. ed. II. ii. 699, sphalm. = praec.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 353.—Afr. austr.

TRICALYSIA, A. Rich. ex DC. Prod. iv. 445 (1830); et in Mém. Soc. Hist. Nat. Par. v. (1834) 224. *RUBIACEAE*, Benth. & Hook. f. ii. 95.
BUNBURYA, Meissn. ex Hochst. in Flora, xxvii. (1844) 553.
NATALANTHE, Sond. in Linnaea, xxiii. (1850) 52.
ROSEA, Klotzsch, in Monatsb. Akad. Berl. (1853) 501.
angolensis, *A. Rich. ex DC. Prod.* iv. 445; *et in Mém. Soc. Hist. Nat. Par.* v. (1834) 225.—Afr. trop.
biafrana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 122.—Afr. trop.
bracteata, *Hiern, l. c.* 120.—Afr. trop.
buxifolia, *Hiern, l. c.* 119.—Afr. trop.
congesta, *Hiern, l. c.* 120.—Afr. trop.
coriacea, *Hiern, l. c.*—Afr. trop.
crassifolia, *Benth. & Hook. f. ex Hiern, l. c.* 124.—Afr. trop.
cryptocalyx, *Baker, in Journ. Bot.* xx. (1882) 138.—Madag.
djurensis, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 123.—Afr. trop.
gabonica, *Hiern, l. c.* 122.—Afr. trop.
jasminiflora, *Benth. & Hook. f. ex Hiern, l. c.* 124.—Afr. trop.
Kirkii, *Hiern, l. c.* 124.—Afr. trop.
micrantha, *Hiern, l. c.* 122.—Afr. trop.
microphylla, *Hiern, l. c.* 123.—Afr. trop.
niamniamensis, *Schweinf. ex Hiern, l. c.* 123.—Afr. trop.
Nyassae, *Hiern, l. c.* 121.—Afr. trop.
okelensis, *Hiern, l. c.* 122.—Afr. trop.
ovalifolia, *Hiern, l. c.* 119.—Afr. trop.
pallens, *Hiern, l. c.* 121.—Afr. trop.
reticulata, *Hiern, l. c.* 121.—Afr. trop.
Sonderiana, *Hiern, l. c.* 119.—Afr. trop.
syrmanthera, *Hiern, l. c.* 120.—Afr. trop.

TRICARDIA, Torr. ex S. Wats. Bot. King's Exped. 258. t. 24 (1871). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 829.
Watsoni, *Torr. ex S. Wats. l. c.*—Am. bor. occ.

TRICARIUM, Lour. Fl. Cochinch. 557 (1790) = **Phyllanthus**, Linn. (Euphorb.).
cochinchinense, Lour. l. c. = P. cochin-chinensis.

TRICARYUM, Spreng. Syst. iii. 848 (1826) = praec.

TRICATUS, Pritz. Ic. Ind. ii. 275 (1866), sphalm. = TRICRATUS, L'Hérit. = **Abronia**, Juss. (Nyctag.).
admirabilis, Pritz. l. c. = A. umbellata.

TRICENTRUM, DC. Prod. iii. 123 (1828) = **Comolia**, DC. (Melastom.).
leptophyllum, DC. l. c. = C. leptophylla.
ovalifolium, DC. l. c. = C. denudata.

TRICERA, Schreb. Gen. 630 (1791); Sw. Fl. Ind. Occ. i. 331. t. 7 (1797) = **Buxus**, Linn. (Euphorb.).
acuminata, Griseb. in Goett. Nachr. (1865) 162 = B. acuminata.
citrifolia, Willd. Sp. Pl. iv. 338 = B. citrifolia.
cordifolia, Willd. l. c. 339 = B. cordifolia (cf. Addend.).

TRICERA :—
cubana, A. Rich. Fl. Cub. Fanerog. ii. 217 = Buxus cubana.
fasciculata, Griseb. Fl. Brit. W. Ind. 31 = B. Purdieana.
glomerata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 157 = B. glomerata.
gonoclada, Wright, ex Griseb. Cat. Pl. Cub. 282 = B. gonoclada.
laevigata, Sw. Fl. Ind. Occ. i. 333 = B. laevigata.
laevigata, Willd. Sp. Pl. iv. 338, partim = B. Vahlii.
microphylla, Griseb. in Goett. Nachr. (1865) 163 = B. Wrightii.
nepalensis, Wall. Cat. n. 7979 = Sarcococca pruniformis.
retusa, Griseb. in Goett. Nachr. (1865) 163 = B. retusa.

TRICERAIA, Willd. ex Roem. & Schult. Syst. iv. 803 (1819) = **Turpinia**, Vent. (Sapind.).
tinifolia, Humb. & Bonpl. ex Roem. Schult. l. c. = Turpinia insignis.

TRICERAS, Andrz. ex Reichb. Consp. 185 (1828) = **Matthiola**, R. Br. (Crucif.).

TRICERASTES, Presl, Rel. Haenk. ii. 88. t. 64 (1836) = **Datisca**, Linn.
glomerata, Presl, l. c. = D. glomerata.

TRICERATIA, A. Rich. Ess. Fl. Cub. 614. t. 44 (1845) = **Sicydium**, Schlecht. (Cucurb.).
bryonioides, A. Rich. l. c. 615 = S. Schiedeanum.

TRICERCANDRA, A. Gray, in Perry, Jap. Exped. ii. 318 (1857) = **Chloranthus**, Sw.
Fortuni, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 405 = C. Fortunei.
quadrifolia, A. Gray, in Perry, Jap. Exped. ii. 318 = C. japonicus.

TRICERMA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 97 = **Maytenus**, Molina (Celastr.).
crassifolium, Liebm. l. c. 98 = M. phyllanthoides.

TRICEROS, Griff. Notul. iv. 606 (1854) = **Gomphogyne**, Griff. (Cucurb.).

TRICEROS, Lour. Fl. Cochinch. 184 (1790). Genus dubium *ANACARDIACEARUM*.—Cf. Benth. & Hook. f. i. 418, 1000.
cochinchinensis, Lour. l. c.—Cochinch.
xalapensis, Spreng. Syst. i. 947 = Turpinia insignis.

TRICHACANTHUS, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 572. *ACANTHACEAE*, Benth. & Hook. f. ii. 1089.
exiguus, *Zoll. & Mor. l. c.*—Java.

TRICHACHNE, Nees, Agrost. Bras. 85 (1829) = **Panicum**, Linn. (Gramin.).
ferruginea, Nees, l. c. 90 = P. phaeothrix.
insularis, Nees, l. c. 86 = P. lanatum.
recalva, Nees, l. c. 88 = P. recalvum.
sacchariflora, Nees, l. c. 87 = P. lanatum.
Sellowii, C. Muell. in Bot. Zeit. xix. (1861) 315 = Andropogon leucostachyus.
tenuis, Nees, Agrost. Bras. 89 = P. tenerrimum.
velutinum, Nees, l. c. 90 = P. vestitum.

TRICHADENIA, Thw. in Hook. Kew Journ. vii. (1855) 196. t. 8. *BIXINEAE*, Benth. & Hook. f. i. 129.
zeylanica, *Thw. l. c.*—Zeylan.

TRICHAETA, Beauv. Agrost. 86. t. 17. f. 8 (1812) = **Trisetum**, Pers. (Gramin.).
ovata, Beauv. l. c. = Trisetum ovatum.

TRICHANDRUM, Neck. Elem. i. 84 (1790) = **Helichrysum**, Vaill. (Compos.).

TRICHANTHA, Hook. Ic. Pl. tt. 666, 667 (1844). *GESNERACEAE*, Benth. & Hook. f. ii. 1009.
major, *Hook. l. c.* t. 667.—N. Granat.
minor, *Hook. l. c.* t. 666.—N. Granat.

TRICHANTHA, Karst. & Triana, in Linnaea, xxviii. (1856) 437 = **Breweria**, R. Br. (Convolv.).
ferruginea, Karst. & Triana, l. c. 438.—N. Granat.

TRICHANTHEMIS, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 617. *COMPOSITAE*.
karataviensis, *Regel & Schmalh. l. c.*—Turkestan.

TRICHANTHERA, Ehrenb. in Linnaea, iv. (1829) 401 = **Hermannia**, Linn. (Stercul.).
modesta, Ehrenb. l. c. = H. arabica.

TRICHANTHERA, H. B. & K. Nov. Gen. et Sp. ii. 243, in obs. (1817). *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
TRIXANTHERA, Rafin. Sylva Tellur. 146 (1838).
gigantea, *Humb. & Bonpl. ex Steud. Nom.* ed. I. 708, in syn.; *Nees, in DC. Prod.* xi. 218.—N. Granat.; Panama.

TRICHANTHODIUM, Sond. & F. Muell. in Linnaea, xxv. (1852) 489 = **Gnephosis**, Cass. (Compos.).
skeriophorum, Sond. & F. Muell. l. c. 490 = G. skirrophora.

TRICHARIS, Salisb. Gen. Pl. Fragm. 24 (1866) = **Dipcadi**, Medic. (Liliac.).
serotina, Salisb. l. c. = D. serotinum.

TRICHASMA, Walp. in Linnaea, xiii. (1839) 510 = **Argyrolobium**, Eckl. & Zeyh. (Legumin.).
calycinum, Walp. l. c. = A. calycinum.
ciliatum, Walp. l. c. = A. Andrewsianum.

TRICHASTEROPHYLLUM, Willd. ex Link, Jahrb. i. III. (1820) 69 = **Helianthemum**, Tourn. (Cistin.).
hyssopifolium, Willd. ex Link, l. c. = H. glomeratum.

TRICHAURUS, Arn. in Wight & Arn. Prod. 40 (1834) = **Tamarix**, Linn.
Aucherianus, Decne. in Jacquem. Voy. Bot. 59 = Tam. passerinoides.
ericoides, Arn. ex Wight & Arn. Prod. 40 = Tam. ericoides.
pycnocarpus, Decne. in Jacquem. Voy. Bot. 59 = Tam. passerinoides.
vaginatus, Desv. ex Walp. Rep. ii. 115.—Persia.

TRICHELOSTYLIS, Lestib. Ess. Fam. Cyp. 40 (1819) = **Fimbristylis**, Vahl (Cyperac.).
angularis, Nees, in Wight, Contrib. 105, in obs. = F. angularis.
aspera, Nees, in Mart. Fl. Bras. ii. I. 80 = F. aspera.
asperrima, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 41 = F. asperrima.
autumnalis, Nees, in Linnaea, ix. (1834) 290 = F. autumnalis.
borealis, Wood, Am. Bot. & Flor. 364.—Am. bor.
capillaris, Wood, Class-Book (1861), 742 (Bulbostylidis sp.).—Am. bor.
capitata, Nees, in Linnaea, ix. (1834) 290 = F. capitata.
carinata, Wood, Class-Book (1861), 742 = Scirpus carinatus.
chaetorrhiza, Nees, in Wight, Contrib. 106 = F. junciformis.
ciliatifolia, Wood, Am. Bot. & Flor. 364 (Bulbostylidis sp.).—Am. bor.
cinnamometorum, Nees, in Linnaea, ix. (1834) 290 = F. cyperoides.
coarctata, Wood, Class-Book (1861), 742 (Bulbostylidis sp.).—Am. bor.
complanata, Nees, in Wight, Contrib. 103 = F. complanata.
contexta, Nees, in Linnaea, x. (1836) 146 (Bulbostylidis sp.).—Madag.; Afr. austr.
curvifolia, Nees, in Wight, Contrib. 105, in obs. = F. diphylla.
digitata, Dalz. ex Boeck. in Flora, lxi. (1878) 35 = F. digitata.
filiformis, Nees, in Wight, Contrib. 102 = F. pauciflora.
geminata, Nees, in Linnaea, ix. (1834) 290; et in Mart. Fl. Bras. ii. I. 80 = F. autumnalis.
globulosa, Nees, in Wight, Contrib. 105 = F. globulosa.
gracilis, Nees, in Linnaea, vii. (1832) 492 = Ficinia gracilis.
junciformis, Nees, in Wight, Contrib. 106 = F. junciformis.
Kielmeyeri, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 38 = F. complanata.
latifolia, Nees, in Wight, Contrib. 106 = F. junciformis.

TRICHELOSTYLIS :—
Ludwigii, Nees, in Linnaea, ix. (1834) 290 = Scirpus Ludwigii.
miliacea, Nees, in Wight, Contrib. 103 = Fimbristylis miliacea.
mucronulata, Torr. in Ann. Lyc. N. York, iii. (1836) 355 = F. autumnalis.
obtusifolia, Nees, in Linnaea, ix. (1834) 290 = F. obtusifolia.
pentagona, Nees, in Wight, Contrib. 105, in obs. = F. miliacea.
pentaptera, Nees, l. c. = F. pentaptera.
pilosula, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 28 = F. exilis.
quinquangularis, Nees, in Wight, Contrib. 104 = F. miliacea.
Rudgeana, Nees, in Mart. Fl. Bras. ii. I. 79 = F. complanata.
salbundia, Nees, in Wight, Contrib. 105 = F. salbundia.
scabra, Nees, l. c. 103 = F. complanata.
Schimperiana, Hochst. ex Boeck. in Linnaea, xxxvii. (1871–73) 28 = F. exilis.
stenophylla, Wood, Class-Book (1861), 742 (Bulbostylidis sp.).—Am. bor.
stricta, Nees, in Hook. Journ. Bot. ii. (1840) 394 = F. Schomburgkiana.
tenella, Nees, in Wight, Contrib. 103 = F. tenera.
tetragona, Nees, l. c. 104 = F. miliacea.
torta, Nees, l. c. 105 = F. junciformis.
Warei, Wood, Class-Book (1861), 742 (Bulbostylidis sp.).—Florida.
xyroides, Arn. ex Hook. f. Fl. Tasm. p. xlviii = F. tetragona.

TRICHERA, Schrad. Cat. Sem. Hort. Gotting. (1814); Roem. & Schult. Syst. iii. 54 (1818) = **Scabiosa**, Tourn. (Dipsac.).
ambigua, Nym. Syll. Suppl. 14 = S. ambigua.
amplexicaulis, Nym. Syll. 60 = S. amplexicaulis.
arvensis, Schrad. Cat. Sem. Hort. Gotting. (1814); Roem. & Schult. Syst. iii. 54 = S. arvensis.
bidens, Schult. Mant. iii. 29 = S. bidens.
bohemica, Nym. Consp. 347 = S. arvensis.
brevis, Schult. Mant. iii. 30 = S. papposa.
calycina, Presl, Fl. Sic. p. xxviii.—Sicil.
campestris, Schult. Mant. iii. 27 = S. arvensis.
carpathica, Nym. Syll. Suppl. 14 = S. arvensis.
carpophylax, Nym. Consp. 347 = S. sylvatica.
ciliata, Roem. & Schult. Syst. iii. 57 = S. ciliata.
collina, Nym. Syll. 60 = S. collina.
communis, Čelak. in Arch. Naturw. Landesd. Boehm. iv. (1881) 815 = S. arvensis.
cuspidata, Nym. Syll. Suppl. 14 = S. sylvatica.
diodon, Presl, Fl. Sic. p. xxviii.—Sicil.
dipsacifolia, Nym. Syll. Suppl. 14 = S. sylvatica.
diversifolia, Nym. Syll. 60 = S. banatica.
diversifolia, Schult. Mant. iii. 28 = S. banatica.
drymeia, Nym. Syll. Suppl. 14 = S. sylvatica?
dumetorum, Nym. l. c. = S. arvensis.
dumetorum, Roem. & Schult. Syst. iii. 56 = S. dumetorum.
exaltata, Nym. Consp. 347 = S. arvensis.
Fleischmanni, Nym. Syll. 59 = S. arvensis.
Godeti, Nym. Consp. 347 = S. longifolia.
hirsuta, Schult. Mant. iii. 27 = S. arvensis.
hybrida, Roem. & Schult. Syst. iii. 58 = S. integrifolia.
indivisa, Nym. Syll. Suppl. 14 = S. arvensis.
integrifolia, Roem. & Schult. Syst. iii. 59 = S. integrifolia.
lancifolia, Fuss, in Verh. Siebenb. Ver. Naturw. xiv. (1863) 195 = S. sylvatica.
legionensis, Roem. & Schult. Syst. iii. 56 = S. legionensis.
leucantha, Schrad. Cat. Sem. Hort. Gotting. (1814) 3; ex Roem. & Schult. Syst. iii. 58 = S. ciliata.
longifolia, Roem. & Schult. l. c. 56 = S. longifolia.
lyrophylla, Nym. Consp. 347 = S. macedonica.
macedonica, Nym. Syll. 60 = S. macedonica.
magnifica, Nym. Consp. 347 (= *Knautia magnifica*).—Macedon.
mollis, Nym. Syll. 60 = S. arvensis.
montana, Roem. & Schult. Syst. iii. 57 = S. montana.
mutabilis, Schrad. Cat. Sem. Hort. Gotting. (1814) = S. integrifolia.
nympharum, Nym. Syll. Suppl. 14 = S. ciliata.
ovatifolia, Roem. & Schult. Syst. iii. 56 = S. sylvatica.
pannonica, Nym. Syll. Suppl. 14 = S. sylvatica.

TRICHERA :—
puberula, Nym. Syll. Suppl. 14 = Scabiosa sylvatica.
pubescens, Schrad. Cat. Sem. Hort. Gotting. (1814); Roem. & Schult. Syst. iii. 57 = S. arvensis.
Salcedi, Roem. & Schult. Syst. iii. 55 = S. Salcedi.
subcanescens, Nym. Consp. 347 = S. sylvatica.
subscaposa, Nym. Syll. 60 = S. collina.
sylvatica, Schrad. Cat. Sem. Hort. Gotting. (1814); Roem. & Schult. Syst. iii. 55 = S. sylvatica.
Timeroyi, Nym. Syll. 60 = S. arvensis.
transsilvanica, Schur, ex Nym. Consp. 347 = S. sylvatica.
trivialis, Nym. l. c. = S. arvensis.
Urvillei, Nym. Syll. 60 = S. integrifolia.
virgata, Nym. Consp. 347 = S. arvensis.
Willichii, Schult. Mant. iii. 29 = S. plumosa.

TRICHEROSTIGMA, Boiss. in DC. Prod. xv. II. 7 (1862), sphalm. = TRICHOSTERIGMA, Klotzsch = **Euphorbia**, Linn.

TRICHILIA, P. Br. Hist. Jamaic. 278 (1756).
MELIACEAE, Benth. & Hook. f. i. 337.
ACRILIA, Griseb. Fl. Brit. W. Ind. 129 (1859).
BARBILUS, P. Br. Hist. Jamaic. 216 (1756).
ELCAJA, Forsk. Fl. Aegypt. Arab. 127 (1775).
GENIOSTEPHANUS, Fenzl, in Flora, xxvii. (1844) 312.
HALESIA, Loefl. It. Hisp. 188 (1758).
HEYNICHIA, Kunth, Ind. Sem. Hort. Berol. (1844), adn., 8.
MAFUREIRA, Bertol. Misc. Bot. ix. 5. t. 2 (1851).
MOSCHOXYLUM, A. Juss. in Mém. Mus. Par. xix. (1830) 238.
ODONTANDRA, Willd. ex Roem. & Schult. Syst. v. p. xli. 511 (1819).
ODONTOSIPHON, M. Roem. Syn. Hesper. 85, 106 (1846).
PHOLACILIA, Griseb. Fl. Brit. W. Ind. 129 (1859).
PORTESIA, Cav. Diss. vii. 369. tt. 215, 216 (1789).
ROCHETIA, Delile, in Rochet, Sec. Voy. Choa, 344 (1846).
TORPESIA, M. Roem. Syn. Hesper. 86, 116 (1846).
acuminata, *C. DC. in DC. Monog. Phan.* i. 704.—N. Granat.
affinis, A. Juss. in A. St. Hil. Fl. Bras. Mer. ii. 78 = T. Catigua.
albicans, *C. DC. in Mart. Fl. Bras.* xi. I. 218.—Bras.
alliacea, Forst. f. Prod. 33 = Dysoxylum Forsteri.
alternans, *C. DC. in DC. Monog. Phan.* i. 700.—Bras.
amplifolia, *C. DC. l. c.* 694.—Peruv.
appendiculata, *C. DC. l. c.* 696.—N. Granat.
arborescens, Spreng. Syst. iv. Cur. Post. 252 = Dysoxylum arborescens.
asterotricha, *Radlk. in Bremen, Abh.* viii. (1883) 383.—Afr. trop.
Aubletii, Steud. Nom. ed. II. ii. 700 = Guarea trichilioides.
barraënsis, C. DC. in Mart. Fl. Bras. xi. I. 222 = fluminensis?
bijuga, Labill. Sert. Austro-Caled. 54. t. 54 (= *Hartighsea Billardierii*).—N. Caled.
Blancheti, *C. DC. in Mart. Fl. Bras.* xi. I. 217.—Bras.
brachystachya, *Klotzsch, ex C. DC. in DC. Monog. Phan.* i. 650.—Guian.; Venezuela.
canjerana, *Vell. Fl. Flum.* 176; iv. t. 102.—Bras.
capensis, Pers. Syn. i. 468 = Ekebergia capensis.
capitata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 120.—Afr. trop.
Casaretti, *C. DC. in Mart. Fl. Bras.* xi. I. 217.—Bras.
cathartica, *Mart. Reise Bras.* i. 546.—Bras.
Catigna, C. DC. in DC. Monog. Phan. i. 689, sphalm. = seq.
Catigua, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* ii. 77.—Bras.
caucana, *C. DC. in DC. Monog. Phan.* i. 669.—N. Granat.
cauliflora, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 318 = Dysoxylum spectabile.
Cipo, *C. DC. in Mart. Fl. Bras.* xi. I. 214.—Guian.; Bras.
Clausseni, *C. DC. l. c.* 207.—Bras.
corcovadensis, *C. DC. l. c.* 215.—Bras.
coriacea, Heyne, ex Wall. Cat. sub n. 1265 = Walsura Piscidia.
cuneata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* ix. (1879) 642.—Guatemala.
cuspidata, *C. DC. in Mart. Fl. Bras.* xi. I. 222.—Bras.

TRICHILIA :—
discolor, *A. Juss. in Mém. Mus. Par.* xix. (1830) 236, 277.—Bras.
Distini, *C. DC. in DC. Monog. Phan.* i. 698.—Jamaic.
diversifolia, A. Juss. in Mém. Mus. Par. xix. (1830) 237, 278 = simplicifolia.
Dregeana, Harv. & Sond. Fl. Cap. i. 246 = Dregei.
Dregei, *E. Mey. in Drège, Zwei Pfl. Docum.* 227, nomen.—Afr. austr.
echinocarpa, *Walp. Ann.* ii. 227.—Guian.
Eckebergia, E. Mey. in Drège, Zwei Pfl. Docum. 227 = Ekebergia Meyeri.
elegans, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* ii. 79. t. 98.—Bras.
emarginata, *C. DC. in Mart. Fl. Bras.* xi. I. 212.—Bras.
emetica, *Vahl, Symb. Bot.* i. 31.—Afr. trop.; Arab.
euneura, *C. DC. in DC. Monog. Phan.* i. 673.—Guian.
excelsa, *Benth. in Hook. Kew Journ.* iii. (1851) 368.—Bras.
excelsa, Spreng. Syst. iv. Cur. Post. 252 = Dysoxylum excelsum.
flava, *C. DC. in Mart. Fl. Bras.* xi. I. 203. t. 59.—Bras.
flaviflora, *C. DC. l. c.* 216.—Bras.
fluminensis, *M. Roem. Syn. Hesper.* 112.—Bras.
fuscescens, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* ix. (1879) 641.—Guian.
Gardneri, *C. DC. in Mart. Fl. Bras.* xi. I. 221.—Bras.
Gaudichaudii, *C. DC. l. c.* 209.—Bras.
glabra, *Linn. Syst.* ed. X. 1020.—Am. trop.
glabra, Macfad. Fl. Jamaic. i. 172 (al. ordinis).—Jamaic.
glabra, Vell. Fl. Flum. 175; iv. t. 101 = fluminensis.
glandulosa, Sm. in Rees, Cycl. xxxvi. n. 10 = Synoum glandulosum.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. I. 205. t. 60. f. 1.—Bras.
Goudotiana, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 366.—N. Granat.
grandiflora, *Oliver, Fl. Trop. Afr.* i. 335.—Afr. trop.
Guara, Linn. Sp. Pl. ed. II. 551 = Guarea trichilioides.
guayaquilensis, *C. DC. in DC. Monog. Phan.* i. 682.—Ecuador.
guianensis, *Klotzsch, ex C. DC. l. c.* 657.—Guian.
Halesia, Loefl. ex A. Juss. in Mém. Mus. Par. xix. (1830) 236 = trifolia.
havannensis, Jacq. Enum. Pl. Carib. 20 = glabra.
heterophylla, *Willd. Sp. Pl.* ii. 554.—Madag.
Heudelotii, *Planch. ex Oliver, Fl. Trop. Afr.* i. 334.—Afr. trop.
hexandra, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 29 = Chisocheton patens.
Hieronymi, *Griseb. in Goett. Abh.* xxiv. (1879) 77.—Reg. Argent.
hirsuta, *C. DC. in Mart. Fl. Bras.* xi. I. 221. t. 63.—Bras.
hirta, *Linn. Syst.* ed. X. 1020.—Jamaic.
humilis, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. iv. 4 = Munronia javanica.
inermis, *Spreng. Neue Entdeck.* i. 285.—Bras.
insignis, *C. DC. in Mart. Fl. Bras.* xi. I. 206. t. 61. f. 1.—Bras.
jamaicensis, *C. DC. in DC. Monog. Phan.* i. 678.—Jamaic.
japurensis, *C. DC. in Mart. Fl. Bras.* xi. I. 214.—Bras.
Karstenii, *C. DC. in DC. Monog. Phan.* i. 704.—N. Granat.
Karwinskyana, *C. DC. l. c.* 663.—Mexic.
lagoënsis, *C. DC. in Mart. Fl. Bras.* xi. I. 207.—Bras.
lanceolata, *C. DC. in DC. Monog. Phan.* i. 698.—Peruv.
lepidota, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 54.—Bras.
longissima, Wall. Cat. n. 8069 = Chisocheton paniculatus.
macrocarpa, Spreng. Syst. iv. Cur. Post. 252 = Dysoxylon macrocarpum.
macrophylla, *Benth. in Hook. Kew Journ.* iii. (1851) 369.—Bras.
Martiana, *C. DC. in Mart. Fl. Bras.* xi. I. 205.—Bras.
mauritiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 589.—Ins. Maurit.
maynasiana, *C. DC. in DC. Monog. Phan.* i. 700.—Peruv.

TRICHILIA :—
membranacea, Macfad. Fl. Jamaic. i. 171 = moschata.
micrantha, *Benth. in Hook. Kew Journ.* iii. (1851) 369.—Bras.
microphylla, Walp. Ann. iv. 389 = macrophylla.
microphyllina, *C. DC. in Mart. Fl. Bras.* xi. I. 216.—Bras.
microstachya, *C. DC. l. c.* 212.—Bras.
minor, A. Rich. Ess. Fl. Cub. 301. t. 33 = glabra.
mollis, *C. DC. in Mart. Fl. Bras.* xi. I. 202.—Bras.
mollissima, Spreng. Syst. iv. Cur. Post. 252 = Dysoxylum mollissimum.
monophylla, A. Rich. Ess. Fl. N. Zél. 306 = Pittosporum tenuifolium.
montana, *H. B. & K. Nov. Gen. et Sp.* vii. 226.—N. Granat.
Moritzii, *C. DC. in DC. Monog. Phan.* i. 707.—Venezuela.
moschata, *Sw. Prod. Veg. Ind. Occ.* 67.—Jamaic.
multiflora, *Casar. Nov. Stirp. Bras. Dec.* ii. 23.—Bras.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 23.—Afr. austr.
nervosa, Vahl, Symb. Bot. i. 31 = Sandoricum indicum.
oblonga, *C. DC. in Mart. Fl. Bras.* xi. I. 218.—Bras.
octandra, Soland. ex M. Roem. Syn. Hesper. 101 = Synoum glandulosum.
odorata, *Andr. Bot. Rep.* t. 637.—Ins. S. Vincent.
Oerstediana, *C. DC. in DC. Monog. Phan.* i. 677.—Nicaragua.
oligantha, *C. DC. l. c.* 693.—Mexic.
oxyphylla, *C. DC. in Mart. Fl. Bras.* xi. I. 204.—Bras.
pallens, *C. DC. l. c.* 218.—Bras.
pallida, *Sw. Prod. Veg. Ind. Occ.* 67.—Ins. S. Doming.
paraënsis, *C. DC. in Mart. Fl. Bras.* xi. I. 213.—Bras.
parviflora, *C. DC. in DC. Monog. Phan.* i. 679.—N. Granat.
pentandra, *Blanco, Fl. Filip.* ed. I. 355.—Malaya.
peruviana, *C. DC. in DC. Monog. Phan.* i. 654.—Peruv.; Chili.
Pleeana, *C. DC. in Mart. Fl. Bras.* xi. I. 215.—Bras.
Poeppigii, *C. DC. in DC. Monog. Phan.* i. 685.—Bras.
polyantha, Poepp. ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xv. (1872) 371 = Guarea Poeppigii.
portoricensis, *Spreng. Syst.* iii. 68.—Porto Rico.
Prieureana, *A. Juss. in Mém. Mus. Par.* xix. (1830) 276.—Afr. trop.
procera, Forsyth, ex DC. Prod. i. 626 = Carapa procera.
propinqua, *C. DC. in DC. Monog. Phan.* i. 693 = Nicaragua.
Pseudostipularis, *C. DC. in Mart. Fl. Bras.* xi. I. 215.—Bras.
pteleaefolia, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* ii. 80. t. 99.—Bras.
pubescens, Klotzsch, ex C. DC. in DC. Monog. Phan. i. 650 = brachystachya.
pubescens, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Guarea pubescens.
Purdiei, *C. DC. in DC. Monog. Phan.* i. 697.—N. Granat.
quadrijuga, *H. B. & K. Nov. Gen. et Sp.* v. 215.—Porto Rico.
quinquejuga, *Vell. Fl. Flum.* 176; iv. t. 103.—Bras.
quinquevalvis, *Montr. in Mém. Acad. Lyon*, x. (1860) 191.—N. Caled.
quinquevalvis, Panch. & Sebert, Not. Bois Nouv. Caléd. 227 = Dysoxylum Lessertianum.
repanda, Vell. Fl. Flum. 176; iv. t. 104 = Casaretti.
retusa, *Oliver, Fl. Trop. Afr.* i. 334.—Afr. trop.
Richardiana, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* ii. 78.—Bras.
Riedelii, C. DC. in Mart. Fl. Bras. xi. I. 202 = echinocarpa?
rimosa, Blanco, Fl. Filip. ed. I. 297 = Heynea trijuga.
riparia, *Mart. ex C. DC. in Mart. Fl. Bras.* xi. I. 213.—Bras.
roraimana, *C. DC. in DC. Monog. Phan.* i. 670.—Guian.
rubescens, *Oliver, Fl. Trop. Afr.* i. 336.—Afr. trop.
rubra, *C. DC. in Mart. Fl. Bras.* xi. I. 203.—Bras.
rufinervis, *Blume, Bijdr.* 164.—Java.
Rueppelliana, Fresen. in Mus. Senckenb. ii. (1837) 278 = Ekebergia Ruppeliana.
Ruiziana, *C. DC. in DC. Monog. Phan.* i. 702.—Peruv.

TRICHILIA:—
scandens, *Lunan, Hort. Jamaic.* ii. 319, 320.—Jamaic.
Schiedeana, *C. DC. in DC. Monog. Phan.* i. 664.—Mexic.
schinoides, *Klotzsch, in Linnaea*, xiv. (1840) 300.—Bras.
Schomburgkii, *C. DC. in DC. Monog. Phan.* i. 695.—Guian.
sebastianopolitana, *C. DC. in Mart. Fl. Bras.* xi. 1. 216.—Bras.
Selloi, *C. DC. l. c.* 219.—Bras.
septentrionalis, *C. DC. l. c.* 220.—Bras.
silvatica, *C. DC. l. c.* 212.—Bras.
similis, Spreng. Syst. iv. Cur. Post. 252 = Dysoxylon simile.
simplicifolia, *Spreng. Syst.* iii. 69.—Ins. Martinic.
singularis, *C. DC. in Mart. Fl. Bras.* xi. 1. 217.—Bras.
Sloanei, Macfad. Fl. Jamaic. i. 168 = terminalis.
spectabilis, Forst. f. Prod. 33 = Dysoxylum spectabile.
spinosa, Willd. Sp. Pl. ii. 554 = Atalantia monophylla.
spondiodes, *Jacq. Enum. Pl. Carib.* 20; *Select. Am.* 128.—Ind. occ.
strigulosa, *Welw. ex C. DC. in DC. Monog. Phan.* i. 658.—Afr. trop.
subalata, *C. DC. in Mart. Fl. Bras.* xi. 1. 213.—Bras.
subsessilifolia, *C. DC. in DC. Monog. Phan.* i. 685.—Guian.
surinamensis, *C. DC. l. c.* 679.—Guian.
terminalis, *Jacq. Enum. Pl. Carib.* 20; *Select. Am.* 130.—Jamaic.
tetrapetala, *C. DC. in Mart. Fl. Bras.* xi. 1. 211.—Bras.
tocacheana, *C. DC. in DC. Monog. Phan.* i. 701.—Peruv.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* v. 215.—Peruv.
trachyantha, *C. DC. in DC. Monog. Phan.* i. 697.—Cuba.
trifolia, *Linn. Syst.* ed. X. 1020.—Venezuela.
trifoliata, Linn. Sp. Pl. ed. II. 551 = praec.
trifoliata, Wall. Cat. n. 8093 = Walsura piscidia.
trijuga, Vell. Fl. Flum. 176; iv. t. 105 = cathartica.
trinitensis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 279.—Ind. occ.
tripetala, Blanco, Fl. Filip. ed. I. 354 (al. ordinis).—Ins. Philipp.
tuberculata, *C. DC. in DC. Monog. Phan.* i. 711.—Panama.
Tweedieana, *C. DC. in Mart. Fl. Bras.* xi. 1. 213.—Bras.
velutina, *C. DC. l. c.* 208.—Bras.
venosa, Spreng. Syst. iii. 68 = Sandoricum indicum.
verrucosa, *C. DC. in DC. Monog. Phan.* i. 695.—N. Granat.
villosa, Wall. Cat. n. 1264 = Walsura villosa.
volubilis, Blanco, Fl. Filip. ed. II. 249 = Aglaia grandis.
Warmingii, *C. DC. in Mart. Fl. Bras.* xi. 1. 220.—Bras.
Wawrana, *C. DC. in DC. Monog. Phan.* i. 666.—Mexic.
Weddelii, *C. DC. in Mart. Fl. Bras.* xi. 1. 201.—Bras.
Welwitschii, *C. DC. in DC. Monog. Phan.* i. 659.—Afr. trop.

TRICHINIUM, R. Br. Prod. 414 (1810). *AMARANTACEAE*, Benth. & Hook. f. iii. 33.
ARTHROTRICHIUM, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 500.
GOMOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 37.
HEMISTEIRUS, F. Muell. in Linnaea, xxv. (1852) 434.
aervoides, *F. Muell. Fragm.* iii. 123.—Austral.
alopecuroideum, Lindl. Bot. Reg. (1839) t. 28 = exaltatum.
alopecuroideum, *Lindl. in Mitch. Three Exped.* ii. 13.—Austral.
angustifolium, Moq. in DC. Prod. xiii. II. 293 = macrocephalum.
arthrolasium, *F. Muell. ex Benth. Fl. Austral.* v. 239.—Austral.
astrolasium, *F. Muell. ex Benth. l. c.* 222.—Austral.
atriplicifolium, A. Cunn. ex Moq. in DC. Prod. xiii. II. 286 = obovatum.

TRICHINIUM:—
auriculaefolium, *A. Cunn. ex Moq. in DC. Prod.* xiii. II. 287.—Austral.
axillare, *F. Muell. ex Benth. Fl. Austral.* v. 232.—Austral.
Beckerianum, *F. Muell. in Linnaea*, xxv. (1852) 436.—Austral.
brachyanthum, *F. Muell. ex Benth. Fl. Austral.* v. 239.—Austral.
brachytrichum, F. Muell. Fragm. iii. 161 = helipteroides.
caespitulosum, *F. Muell. ex Benth. Fl. Austral.* v. 240.—Austral.
calostachyum, *F. Muell. ex Benth. l. c.* 236.—Austral.
candicans, Nees, in Lehm. Pl. Preiss. i. 629 = alopecuroideum.
carneum, Moq. in DC. Prod. xiii. II. 291 = Stirlingii.
carnosum, Moq. ex Benth. Fl. Austral. v. 190 = Chenolea carnosa.
Chrysurus, Meissn. ex Moq. in DC. Prod. xiii. II. 307 = Sericocoma Chrysurus.
conicum, Lindl. in Mitch. Journ. Trop. Austral. 363 = alopecuroideum.
conicum, Spreng. Syst. i. 816 = Ptilotus conicus.
corymbosum, *Gaudich. in Freyc. Voy. Bot.* 444.—Austral.
corymbosum, Spreng. Syst. i. 816 = Ptilotus corymbosus.
Cunninghamii, *Benth. Fl. Austral.* v. 238.—Austral.
declinatum, *Moq. in DC. Prod.* xiii. II. 293.—Austral.
densum, A. Cunn. ex Moq. l. c. 289 = nobile.
dissitiflorum, *F. Muell. Fragm.* iv. 89.—Austral.
distans, *R. Br. Prod.* 415.—Austral.
divaricatum, *Gaudich. in Freyc. Voy. Bot.* 445.—Austral.
Drummondii, *Moq. in DC. Prod.* xiii. II. 292.—Austral.
eriocephalum, Moq. l. c. 293 = declinatum.
erubescens, *Moq. l. c.*—Austral.
esquamatum, *Benth. Fl. Austral.* v. 229.—Austral.
exaltatum, *Benth. l. c.* 227.—Austral.
floribundum, Moq. in DC. Prod. xiii. II. 283 = sericostachyum.
Fraseri, *A. Cunn. ex Moq. l. c.* 295.—Austral.
fusiforme, *R. Br. Prod.* 415.—Austral.
fusiforme, Lindl. in Mitch. Journ. Trop. Austral. 383 = macrocephalum.
fusiforme, Nees, in Lehm. Pl. Preiss. i. 626 = Drummondii.
Gaudichaudii, Steud. Nom. ed. II. ii. 700 = corymbosum.
giganteum, A. Cunn. ex Moq. in DC. Prod. xiii. II. 296 = alopecuroideum.
gnaphalodes, A. Cunn. ex Moq. l. c. 285 = incanum.
gomphrenoides, *Moq. l. c.* 287.—Austral.
gracile, *R. Br. Prod.* 415.—Austral.
helichrysoides, *F. Muell. ex Benth. Fl. Austral.* v. 241.—Austral.
helipteroides, *F. Muell. Fragm.* iii. 122.—Austral.
holosericeum, *Moq. in DC. Prod.* xiii. II. 287.—Austral.
Hoodii, F. Muell. Fragm. viii. 232 = Ptilotus Hoodii.
humile, Nees, in Lehm. Pl. Preiss. i. 628 = Ptilotus humilis.
incanum, *R. Br. Prod.* 415.—Austral.
incanum, Moq. in DC. Prod. xiii. II. 285 = obovatum.
lanatum, Lindl. in Mitch. Three Exped. ii. 123 = obovatum.
latifolium, E. Mey. ex Moq. in DC. Prod. xiii. II. 328 = Cyathula cylindrica.
laxum, *Benth. Fl. Austral.* v. 232.—Austral.
Leucocoma, *Moq. in DC. Prod.* xiii. II. 292.—Austral.
linifolium, A. Cunn. ex Moq. l. c. = erubescens.
macrocephalum, *R. Br. Prod.* 415.—Austral.
macrocephalum, Moq. in DC. Prod. xiii. II. 290 = exaltatum.
macrocephalum, Nees, in Lehm. Pl. Preiss. i. 627 = Manglesii.
Manglesii, *Lindl. Bot. Reg.* (1839) sub t. 28.—Austral.
mucronatum, Nees, in Lehm. Pl. Preiss. i. 628 = spathulatum.
nanum, F. Muell. Fragm. iii. 161 = Ptilotus humilis.
nitidum, Moq. in DC. Prod. xiii. II. 289 = Ptilotus humilis.
nobile, *Lindl. in Mitch. Three Exped.* ii. 22.—Austral.
obovatum, *Gaudich. in Freyc. Voy. Bot.* 445.—Austral.

TRICHINIUM:—
pachocephalum, Moq. in DC. Prod. xiii. II. 294 = macrocephalum.
pallidum, Moq. l. c. 295 = alopecuroideum.
parviflorum, *Lindl. in Mitch. Three Exped.* ii. 13.—Austral.
parvifolium, *F. Muell. Rep. Babb. Exp.* 19.—Austral.
polystachyum, *Gaudich. in Freyc. Voy. Bot.* 445.—Austral.
Preissii, Nees, in Lehm. Pl. Preiss. i. 629 = alopecuroideum.
pulchellum, A. Cunn. ex Moq. in DC. Prod. xiii. II. 290 = semilanatum.
pyramidatum, *Moq. l. c.* 288.—Austral.
remotiflorum, Hook. Ic. Pl. t. 596 = Sericocoma remotiflora.
Roei, *F. Muell. ex Benth. Fl. Austral.* v. 240.—Austral.
roseum, *Moq. in DC. Prod.* xiii. II. 284.—Austral.
rotundifolium, *F. Muell. Fragm.* iii. 122.—Austral.
semilanatum, *Lindl. in Mitch. Journ. Trop. Austral.* 45.—Austral.
sericostachyum, *Nees, in Lehm. Pl. Preiss.* i. 627.—Austral.
sessilifolium, Lindl. in Mitch. Three Exped. ii. 13 = obovatum.
setigerum, A. Cunn. ex Moq. in DC. Prod. xiii. II. 290 = semilanatum.
spathulatum, *R. Br. Prod.* 415.—Austral.
spectabile, Field. & Gardn. Sert. Pl. t. 53 = Manglesii.
Stirlingii, *Lindl. Bot. Reg.* (1839) sub t. 28.—Austral.
striatum, *Moq. ex Benth. Fl. Austral.* v. 233.—Austral.
subfuscum, Moq. in DC. Prod. xiii. II. 332 = Pupalia subfusca.
variabile, F. Muell. in Linnaea, xxv. (1852) 436 = obovatum.
villosum, *Nees, in Lehm. Pl. Preiss.* i. 628.—Austral.
virgatum, A. Cunn. ex Moq. in DC. Prod. xiii. II. 286 = parviflorum.
Zeyheri, *Moq. l. c.* 296.—Afr. austr.

TRICHLIS, Hall. Hort. Gotting. 26 (1743) = **Polycarpon**, Loefl. (Caryoph.).

TRICHLORA, Baker, in Hook. Ic. Pl. t. 1237 (1877). *LILIACEAE*, Benth. & Hook. f. iii. 805.
peruviana, *Baker, l. c.*—Peruv.

TRICHLORIS, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 102. *GRAMINEAE*, Benth. & Hook. f. iii. 1166.
fasciculata, *Fourn. ex Benth. in Journ. Linn. Soc.* xix. (1881) 102.—Mexic.
pluriflora, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 560.—Mexic.

TRICHOA, Pers. Syn. ii. 634 (1807) = **Abuta**, Aubl. (Menisp.).
concolor, Endl. ex Walp. Rep. i. 95 = A. concolor.
conferta, DC. Prod. i. 103 = A. spicata.
conferta, Pers. Syn. ii. 634.—N. Granat.
guyanensis, Klotzsch, ex Eichl. in Mart. Fl. Bras. xiii. 1. 181 = A. guianensis.
racemosa, Pers. Syn. ii. 634 = A. racemosa.
spicata, Pers. l. c. = A. spicata.

TRICHOBALLIA, Presl, Symb. Bot. i. 9 (1830) = **Tetraria**, Beauv. (Cyperac.).
dactyloides, Presl, l. c., in obs.—Afr. austr.
involucrata, Presl, l. c. = Lepidosperma involucratum.
thermalis, Presl, l. c.—Afr. austr.

TRICHOBASIS, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 336. *MYRTACEAE*.
aurea, *Turcz. l. c.* 337.—Austral.

TRICHOCALYX, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 87. *ACANTHACEAE*.
obovatus, *Balf. f. l. c.* 88.—Ins. Socotra.
orbiculatus, *Balf. f. l. c.*—Ins. Socotra.

TRICHOCALYX, Schau. Myrt. Xeroc. 86, in adnot. (1840) = **Calythrix**, Labill. (Myrtac.).

TRICHOCARPUS, Neck. Elem. ii. 70 (1790) = **Prunus**, Tourn. (Rosac.).

TRICHOCARPUS, Schreb. Gen. 366 (1789) = **Sloanea**, Linn. (Tiliac.).
guianensis, Raeusch. Nom. ed. III. 158 = S. guianensis.
laurifolia, Willd. Sp. Pl. ii. 1224 = S. laurifolia.

TRICHOCARYA, Miq. Fl. Ind. Bat. i. 1. 357 (1855). *ROSACEAE*, Benth. & Hook. f. i. 607.
ANGELESIA, Korth, in Nederl. Kruidk. Arch. iii. (1855) 384.
DIEMENIA, Korth, l. c. 388 (1855).
racemosa, *Miq. Fl. Ind. Bat.* i. 1. 358.—Sumatra.
splendens, *Miq. l. c.*—Sumatra.

TRICHOCAULON, N. E. Br. in Journ. Linn. Soc. xvii. (1878) 164. *ASCLEPIADEAE.*
flavum, *N. E. Br. l. c.* 165.—Afr. austr.
piliferum, *N. E. Br. l. c.* 164.—Afr. austr.

TRICHOCENTRUM, Poepp. & Endl. Nov. Gen. et Sp. ii. 11. t. 115 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 559.
albo-coccineum, *Linden, in Belg. Hortic.* xv. (1865) 103.—Bras.
albo-purpureum, *Reichb. f. ex Rodrig. Orch. Nov.* ii. 238.—Bras.
amazonicum, *Rodrig. l. c.* i. 140.—Bras.
atropurpureum, *Linden & Reichb. f. in Gard. Chron.* (1866) 219.—Bras.
caloceras, *Endres & Reichb. f. l. c.* (1871) 1257.—Costa Rica.
candidum, *Lindl. Bot. Reg.* (1843) *Misc.* 9.—Guatemala.
capistratum, *Linden & Reichb. f. in Gard. Chron.* (1871) 1257.—Costa Rica.
Cornucopiae, *Linden & Reichb. f. l. c.* (1866) 266.—Am. trop.
funale, Lindl. Bot. Reg. (1837), sub t. 1951 = Dendrophylax funalis.
fuscum, *Lindl. l. c.* t. 1951.—Mexic.
Hoegei, *Reichb. f. in Gard. Chron.* (1881) ii. 717.—Mexic.
ionopthalmum, *Reichb. f. l. c.* (1876) ii. 100.—Bras.
iridifolium, Lodd. ex Lindl. Bot. Reg. (1838) Misc. 94 (al. gen.).—Guian.
maculatum, *Lindl. Orchid. Linden.* 24.—Bras.
orthoplectron, *Reichb. f. in Gard. Chron.* (1883) i. 562.—Am. centr. ?
Pfavii, *Reichb. f. l. c.* (1881) ii. 70.—Am. centr.
Pineli, *Lindl. l. c.* (1854) 772.—Bras.
Plectrophora, Reichb. f. in Walp. Ann. vi. 544 (Comparettiae sp.).—Guian.
Porphyrio, *Reichb. f. in Illustr. Hortic.* xxxi. (1884) 9. t. 508.—Am. centr. ?
pulchrum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 11. t. 115.—Peruv.
purpureum, *Lindl. in Gard. Chron.* (1854) 772.—Guian. ?
recurvum, *Lindl. Bot. Reg.* (1843) *Misc.* 9.—Guian.
tenuiflorum, *Lindl. in Lindl. & Paxt. Flow. Gard.* i. (1850–51) 12.—Bras.
tigrinum, *Linden & Reichb. f. in Gard. Chron.* (1869) 892.—Am. centr.

TRICHOCEPHALUM, Schur, Enum. Pl. Transs. 293 (1866) = **Dipsacus**, Linn.
pilosum, Schur, l. c. = D. pilosus.

TRICHOCEPHALUS, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 374. t. 17. f. i. = **Phylica**, Linn. (Rhamn.).
atratus, Eckl. & Zeyh. Enum. 131 = P. atrata.
callosus, Eckl. & Zeyh. l. c. 130 = P. callosa.
comosus, Eckl. & Zeyh. l. c. 132 = P. comosa.
ellipticus, G. Don, Gen. Syst. ii. 40 = Nesiota elliptica.
elongatus, Eckl. & Zeyh. Enum. 130 = Staavia ciliata.
gracilis, Eckl. & Zeyh. l. c. 131 = P. gracilis.
imberbis, Eckl. & Zeyh. l. c. 132 = P. cuspidata.
laevis, Eckl. & Zeyh. l. c. 130 = P. atrata.
littoralis, Eckl. & Zeyh. l. c. 131 = P. atrata.
ramosissimus, G. Don, Gen. Syst. ii. 40 = P. ramosissima.
rhipophorus, Eckl. & Zeyh. Enum. 131 = P. stipularis.
ruber, Eckl. & Zeyh. l. c. 131 = P. rubra.
spicatus, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 375 = P. spicata.
stipularis, Brongn. l. c. = P. stipularis.
trachyphyllus, Eckl. & Zeyh. Enum. 131 = P. trachyphylla.
verticillatus, Eckl. & Zeyh. l. c. 131 = P. verticillata.
virgata, Eckl. & Zeyh. l. c. 132 = P. virgata.

TRICHOCERAS, Spreng. Anleit. ii. 1. 291 (1817) = seq.

TRICHOCEROS, H. B. & K. Nov. Gen. et Sp. i. 337. t. 76 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 587.
ASTROGLOSSUS, Reichb. f. ex Benth. & Hook. f. Gen. iii. 588 (1883).
antennifer, *H. B. & K. Nov. Gen. et Sp.* i. 338.—Ecuador.
armillatus, *Reichb. f. in Bonplandia*, iv. (1856) 212.—Peruv.
astroglossus, *Lindl. ex Benth. & Hook. f. Gen.* iii. 588.—Peruv.
muralis, *Lindl. Gen. et Sp. Orch.* 174.—Peruv.
parviflorus, *H. B. & K. Nov. Gen. et Sp.* i. 337. t. 76.—N. Granat.
platyceros, *Reichb. f. Xenia Orch.* i. 22. t. 9. f. 3.—Peruv.
Tupaipi, *Reichb. f. in Linnaea*, xli. (1877) 33.—Peruv.

TRICHOCHAETA, Steud. Syn. Pl. Cyp. 155 (1855) = **Rynchospora**, Vahl (Cyperac.).
tenuis, Steud. l. c. 156.—Bras.

TRICHOCHLOA, Beauv. Agrost. 29. t. 8. f. 2 (1812) = **Muehlenbergia**, Schreb. (Gramin.).
alpestris, Roem. & Schult. Syst. ii. 388 = M. alpestris.
Berteroniana, Schult. Mant. ii. 209 = M. Berteroniana.
calycina, Trin. Gram. Unifl. 193 = M. debilis.
capillaris, DC. Cat. Hort. Monsp. (1813) 152; Roem. & Schult. Syst. ii. 384 = M. capillaris.
ciliata, Roem. & Schult. l. c. 386 = M. ciliata.
debilis, Roem. & Schult. l. c. 385 = M. debilis.
elegans, Roem. & Schult. l. c. 387 = M. elegans.
expansa, DC. Cat. Hort. Monsp. (1813) 151 = Cinna arundinacea.
foliosa, Trin. Fund. Agrost. 117 = M. mexicana.
glabrata, Roem. & Schult. Syst. ii. 387 = M. glabrata.
glomerata, Trin. Fund. Agrost. 117 = M. glomerata.
gracilis, Roem. & Schult. Syst. ii. 389 = M. gracilis.
implicata, Roem. & Schult. l. c. 385 = M. implicata.
longiflora, Kunth, Enum. Pl. i. 601 (Index) = seq. ?
longiseta, Trin. Fund. Agrost. 117 = M. Willdenowii.
mexicana, Trin. l. c. = M. mexicana.
microsperma, DC. Cat. Hort. Monsp. (1813) 151 = M. debilis.
mucronata, Roem. & Schult. Syst. ii. 387 = M. mucronata.
Polypogon, DC. Cat. Hort. Monsp. (1813) 152 = M. capillaris.
quadridentata, Roem. & Schult. Syst. ii. 388 = M. quadridentata.
rigida, Roem. & Schult. l. c. 386 = M. rigida.
setosa, Roem. & Schult. l. c. = M. debilis.
simplex, Schult. Mant. ii. 577 = Polypogon maritimus ?
sobolifera, Trin. Fund. Agrost. 117 = M. sobolifera.
spicata, Schult. Mant. ii. 577 = M. spicata.
stipoides, Roem. & Schult. Syst. ii. 388 = M. stipoides.
tenella, Roem. & Schult. l. c. 385 = M. tenella.
tenuiflora, Sweet, Hort. Brit. ed. I. 443 = M. Willdenowii.
virescens, Roem. & Schult. Syst. ii. 389 = M. virescens.

TRICHOCLADUS, Pers. Syn. ii. 597 (1807). *HAMAMELIDEAE*, Benth. & Hook. f. i. 667.
DAHLIA, Thunb. in Skrivt. Naturh. Selsk. Kiobenh. ii. (1792) 133. t. 4.
crinitus, *Pers. Syn.* ii. 597.—Afr. austr.
Ecklonianus, Presl, Bot. Bemerk. 75 = crinitus.
ellipticus, *Eckl. & Zeyh. Enum.* 356.—Afr. austr.
grandiflorus, *Oliver, in Hook. Ic. Pl.* t. 1480.—Afr. austr.
peltatus, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 537.—Afr. austr.
verticillatus, *Eckl. & Zeyh. Enum.* 356.—Afr. austr.

TRICHOCLINE, Cass. in Bull. Soc. Philom. (1817) 13. *COMPOSITAE*, Benth. & Hook. f. ii. 496.
AMBLYSPERMA, Benth. in Enum. Pl. Hueg. 67 (1837).
BICHENIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 236.
ERIOPUS, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 376 (1884).
INGENHUSIA, Vell. Fl. Flum. 351 (1825).
angustifolia, *Baker, in Mart. Fl. Bras.* vi. iii. 373.—Bras.

TRICHOCLINE:—
araneosa, *Baker, in Mart. Fl. Bras.* vi. iii. 374.—Bras.
arenaria, *Baker, l. c.*—Bras.
argentea, *Griseb. in Goett. Abh.* xxiv. (1879) 214.—Reg. Argent.
Cineraria, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 103.—Bras.
collina, *Baker, in Mart. Fl. Bras.* vi. iii. 373.—Paraguay.
dealbata, *Benth. & Hook. f. ex Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 359.—Chili.
denticulata, *Baker, in Mart. Fl. Bras.* vi. iii. 375.—Bras.
Eriopus, *Baker, l. c.* 376.—Bras.
exscapa, *Griseb. in Goett. Abh.* xix. (1874) 197.—Reg. Argent.
foliosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 43.—Bras.
heterophylla, *Less. in Linnaea*, v. (1830) 289.—Bras.
hieracioides, *Baker, in Mart. Fl. Bras.* vi. iii. 374.—Bras.
humilis, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 103 = Tweediei.
humilis, Less. in Linnaea, v. (1830) 288 = heterophylla.
incana, *Cass. in Dict. Sc. Nat.* lv. 216.—Bras.
macrocephala, *Less. in Linnaea*, v. (1830) 288.—Bras.
Martii, *Baker, in Mart. Fl. Bras.* vi. iii. 375.—Bras.
maxima, *Less. in Linnaea*, v. (1830) 290.—Bras.
nervosa, *Less. Syn. Comp.* 118.—Bras.
nummularia, *Baker, in Mart. Fl. Bras.* vi. iii. 375.—Bras.
pedicularifolia, *Walp. in Linnaea*, xiv. (1840) 310.—Chili.
plicata, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 103.—Chili.
polymorpha, *Baker, in Mart. Fl. Bras.* vi. iii. 373.—Bras.
scapigera, *F. Muell. Census*, 85.—Austral.
speciosa, *Less. Syn. Comp.* 117.—Bras.
tontalensis, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 49.—Reg. Argent.
Tweediei, *DC. ex Steud. Nom.* ed. II. ii. 701 (Tweedii).—Reg. Argent.

TRICHOCORONIS, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) (Pl. Fendl.) 65. *COMPOSITAE*, Benth. & Hook. f. ii. 240.
MARGACOLA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 457.
Greggii, *A. Gray, Pl. Wright.* i. 89.—Mexic.
rivularis, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 66.—N. Mexic.
Wrightii, *A. Gray, l. c.* 65.—N. Mexic.

TRICHOCREPIS, Vis. Stirp. Dalm. Rar. 18. t. 7 (1826) = **Pterotheca**, Cass. (Compos.).
bifida, Vis. l. c. 19 = P. bifida.

TRICHODESMA, R. Br. Prod. 496 (1810). *BORAGINEAE*, Benth. & Hook. f. ii. 845.
BORRAGINOIDES, Moench, Meth. 515 (1794).
FRIEDRICHSTHALIA, Fenzl, in Endl. Nov. Stirp. Dec. 53 (1839).
LEIOCARYA, Hochst. in Flora, xxvii. (1844) 30.
POLLICHIA, Medic. Bot. Beobacht. (1783) 247.
SPIROCONUS, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 576.
STREBLANTHERA, Steud. in Flora, xxvii. (1844) 29.
africanum, *R. Br. Prod.* 496.—Afr. trop.; Persia; Ind. bor. occ.
ambacensis, *Welw. Apont.* 589.—Afr. trop.
amplexicaule, *Roth, Nov. Sp. Pl.* 104.—Ind. or.
angustifolium, *Harv. Thes. Cap.* i. 26.—Afr. austr.
Aucheri, *DC. Prod.* x. 174.—Persia.
calathiforme, *Hochst. in Flora*, xxvii. (1844) 29.—Abyss.
calcaratum, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 64, nomen.—Afr. bor.
Ehrenbergii, *Schweinf. ex Boiss. Fl. Orient.* iv. 281.—Egypt.; Arab.
hirsutum, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 175 = indicum.
inaequale, Edgew. l. c. = amplexicaule.

TRICHODESMA:—
incanum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 417.—Persia.
indicum, *R. Br. Prod.* 496.—Ind. or.; Afghan; Persia.
indicum, DC. Prod. x. 172 = amplexicaule.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 154.—Reg. Himal.
Kotschyanum, Fenzl, ex A. DC. in DC. Prod. x. 172 = zeylanicum.
latisepaleum, *F. Muell. Fragm.* x. 102.—Austral.
laxiflorum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotra.
lithospermoides, *Roem. & Schult. Syst.* iv. 69.—Ind. or.
microcalyx, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotra.
molle, *A. DC. in DC. Prod.* x. 174.—Persia.
perfoliatum, Wall. Cat. n. 934 = indicum.
physaloides, *A. DC. in DC. Prod.* x. 173.—Afr. austr.
Scotti, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 81.—Ins. Socotra.
sericeum, Lindl. in Mitch. Journ. Trop. Austral. 258 = zeylanicum.
spinulosum, Voigt, Hort. Suburb. Calc. 444 = indicum.
Stocksii, *Boiss. Diagn.* Ser. II. iii. 140.—Persia; Beluchist.
strictum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 178.—Afghan.
subsessile, Wall. Cat. n. 933 = indicum.
zeylanicum, *R. Br. Prod.* 496.—As. et Austral. trop.

TRICHODIA, Griff. Notul. iv. 570 (1854) = **Paropsia**, Noronha (Passifl.).
vareciformis, Griff. l. c. 571 = P. vareciformis.

TRICHODICLIDA, Cerv. in La Naturaleza (1870), 346. *GRAMINEAE.*
linearis, *Cerv. l. c.* 347.—Mexic.
prolifera, *Cerv. l. c.* 346.—Mexic.

TRICHODIUM, Michx. Fl. Bor. Am. i. 41. t. 8 (1803) = **Agrostis**, Linn. (Gramin.).
affine, Willd. ex Steud. Nom. ed. II. ii. 701 = A. Humboldtiana.
album, Presl, Rel. Haenk. i. 244 = A. clavata.
algidum, Roem. & Schult. Syst. ii. 283 = Phippsia algida.
alpestre, Hoppe, Anl. Graes. n. 4; ex Schult. Mant. ii. 156 = A. rupestris.
alpinum, Schrad. Fl. Germ. i. 200. t. 3. f. 4 = A. rupestris.
arenarium, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 51 = A. canina.
arenosum, Schur, Enum. Pl. Transs. 736 = A. canina.
auratum, Sieber, ex Steud. Nom. ed. II. ii. 701 = A. alpina.
caninum, Schrad. Fl. Germ. i. 198 = A. canina.
clavatum, Schult. Mant. ii. 556 = A. clavata.
coerulescens, Link, ex Steud. Nom. ed. II. ii. 701 = A. canina.
decumbens, Michx. Fl. Bor. Am. i. 42 = A. decumbens.
diffusum, Link, Enum. Hort. Berol. i. 69 = A. canina, rupestris.
effusum, Link, ex Schult. Mant. iii. 554 = A. effusa.
elatum, Pursh, Fl. Am. Sept. i. 61 = A. elata.
elegans, Thore, Promen. Gascogne, 92 = A. elegans.
filiforme, Spreng. Pugill. i. 6 = A. rupestris.
flavescens, Schult. Oestr. Fl. ed. II. i. 165 = A. alpina.
glabrum, Presl, Rel. Haenk. i. 244 = A. glabra.
glaucinum, Roem. & Schult. Syst. ii. 281 = A. vulgaris.
hybridum, Roem. & Schult. l. c. 282 = A. canina.
inconspicuum, Kunze, ex Steud. Nom. ed. II. ii. 701 = A. rupestris.
intermedium, Hoppe, ex Schult. Mant. ii. 520 = A. alpina.
Kitaibelii, Schult. Oestr. Fl. ed. II. i. 165 = A. canina.
laxiflorum, Bigel. ex Michx. Fl. Bor. Am. i. 42. t. 8 = A. scabra.
laxum, Schult. Mant. ii. 157 = A. laxiflora.
montanum, Torr. Fl. U. St. 84 = A. scabra.
montanum, Willd. ex Steud. Nom. ed. II. ii. 701 = A. canina.
Muehlenbergiana, Schult. Mant. ii. 159 = A. laxiflora.
nanum, Presl, Rel. Haenk. i. 243 = A. nana.
neglectum, Schult. Oestr. Fl. ed. II. i. 166 = A. Schultesii.

TRICHODIUM:—
noveboracense, Schult. Mant. ii. 555 = Epicampes lanata.
paradisiaca, Nees, ex Steud. Syn. Pl. Gram. 163 = Agrostis chilensis.
perennans, Ell. Sketch, i. 99 = A. perennans.
pusillum, Nees & Meyen, in Mém. Acad. Pétersb. Sér. VI. vi. Sc. Nat. II. (1845) 312 = A. Meyeni.
pusillum, Van Hall, Fl. Belg. i. 56 = A. canina.
rubrum, Roem. & Schult. Syst. ii. 281 = A. vulgaris.
rubrum, Schur, Enum. Pl. Transs. 735 = A. canina.
rupestre, Schrad. Fl. Germ. i. 279 = A. alpina.
rupestre, Schur, Enum. Pl. Transs. 735 = A. rupestris.
salmanticum, Lag. Gen. et Sp. Nov. 3 = A. salmantica.
scabrum, Darl. Fl. Cestr. ed. I. i. 54 = A. perennans.
scabrum, Muhl. Desc. Gram. 61 = A. laxiflora.
Schleicheri, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 181 = A. alpina.
setaceum, Roem. & Schult. Syst. ii. 280 = A. setacea.
setifolium, Willd. ex Steud. Nom. ed. II. ii. 701 = A. setacea.
strictum, Roem. & Schult. Syst. ii. 281 = A. stricta.
subalpinum, Schur, Enum. Pl. Transs. 735 = A. canina.
tenerrimum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 83 = A. canina.
transsilvanicum, Schur, l. c. = A. canina.
trichanthum, Schult. Mant. ii. 157 = A. trichantha.
turfosum, Schur, Enum. Pl. Transs. 737 = A. canina.

TRICHODON, Benth. in Journ. Linn. Soc. xix. (1881) 112, sphalm. = TRICHOON, Roth = **Phragmites**, Trin. (1820) (Gramin.).

TRICHODRYMONIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 112 (Gesn. Centr. Am. 38) = **Episcia**, Mart. (Gesner.).
congesta, Oerst. l. c. = E. congesta.

TRICHOGALIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 396 = **Galium**, Linn. (Rubiac.).
boreale, Fourr. l. c. = G. boreale.
rotundifolium, Fourr. l. c. = G. rotundifolium.

TRICHOGAMILA, P. Br. Hist. Jamaic. 218 (1756) = **Styrax**, Linn.

TRICHOGLOTTIS, Blume, Bijdr. 359 (1825), pro parte. *ORCHIDEAE*, Benth. & Hook. f. iii. 576.
atropurpurea, *Reichb. f. in Linnaea*, xli. (1877) 30.—Ins. Philipp.
Bimae, *Reichb. f. in Bonplandia*, v. (1857) 39.—Ins. Bima.
cirrhifera, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1858) 493.—Java.
cochlearis, *Reichb. f. in Gard. Chron.* (1883) i. 142.—Sumatra.
Dawsoniana, *Reichb. f. l. c.* (1872) 699, in obs.—Burma.
fasciata, Reichb. f. in Flora, lv. (1872) 137 = Stauropsis fasciata.
Guibertii, Reichb. f. in Gard. Chron. (1872) 699, in obs. = Cleisostoma Guiberti.
lanceolaria, *Blume, Bijdr.* 360.—Java.
pallens, Lindl. in Journ. Hort. Soc. v. (1850) 34 = Phalaenopsis pallens.
philippinensis, Lindl. in Ann. & Mag. Nat. Hist. Ser. I. xv. (1845) 386 = Stauropsis philippinensis.
pusilla, *Reichb. f. in Bonplandia*, iv. (1856) 325; vi. (1857) 39 (sp. dub.).—Java.
quadricornuta, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 156 (Cleisostomatis sp.).—Ins. Nicobar.
retusa, *Blume, Bijdr.* 360.—Java.
rigida, *Blume, l. c.* 361.—Java.
tenera, Reichb. f. in Gard. Chron. (1872) 699, in obs. (Cleisostomatis sp.).—Ind. or.

TRICHOGONIA, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 459. *COMPOSITAE*, Benth. & Hook. f. ii. 243.
campestris, *Gardn. l. c.*—Bras.
Gardneri, *A. Gray, in Hook. Kew Journ.* iii. (1851) 224.—Bras.
hirtiflora, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 214.—Bras.
laxa, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 435.—Bras.
leiantha, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 215 = Gardneri.

TRICHOGONIA:—
macrolepis, *Baker, in Mart. Fl. Bras.* vi. II. 215—Bras.
Martii, *Baker, l. c.* 216.—Bras.
menthaefolia, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 434.—Bras.
multiflora, Gardn. l. c. v. (1846) 460 = villosa.
podocarpa, *Sch. Bip. in Linnaea*, xxx. (1859–60) 182.—Bras.
salviaefolia, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 460.—Bras.
Selloi, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 215 = Gardneri.
villosa, *Sch. Bip. ex Baker, l. c.* 213.—Bras.

TRICHOGYNE, Less. in Linnaea, vi. (1831) 231 = **Ifloga**, Cass. (Compos.).
cauliflora, DC. Prod. vi. 266 = I. Fontanesii.
decumbens, Less. Syn. Comp. 359.—Afr. austr.
glomerata, Harv. in Harv. & Sond. Fl. Cap. iii. 287.—Afr. austr.
laricifolia, Less. in Linnaea, vi. (1831) 231.—Afr. austr.
paronychioides, DC. Prod. vi. 266 = I. paronychioides.
polycnemoides, Steud. Nom. ed. II. ii. 701 = I. polycnemoides.
radicans, DC. Prod. vi. 265 = *reflexa*.
reflexa, Less. Syn. Comp. 360.—Afr. austr.
seriphioides, Less. l. c.—Afr. austr.
spicata, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 310 = I. Fontanesii.
verticillata, Less. Syn. Comp. 361 = I. polycnemoides.

TRICHOLAENA, Schrad. in Schult. Mant. ii. 163 (1824) = **Panicum**, Linn. (Gramin.).
arenaria, Nees, Cat. Sem. Hort. Vratisl. (1836); ex ej. Fl. Afr. Austr. 20 = P. ammophilum?
capensis, Nees, Fl. Afr. Austr. 19 = P. capense.
fragilis, A. Br. in Flora, xxiv. (1841) 275 = P. Braunii.
fusca, Schult. Mant. ii. 164 = Saccharum fuscum.
grandiflora, Hochst. in Flora, xxiv. (1841) Intell. 19, nomen; et ex A. Rich. Tent. Fl. Abyss. ii. 445 = P. insigne.
insularis, Griseb. Fl. Brit. W. Ind. 557 = P. leucophaeum.
leucantha, Hochst. ex Steud. Syn. Pl. Gram. 92 = P. leucanthum.
longiseta, Hochst. ex Steud. l. c. = P. macrotrichum.
micrantha, Schrad. in Schult. Mant. ii. 163 = P. Teneriffae.
obtecta, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 493 = P. obtectum.
rosea, Nees, Fl. Afr. Austr. i. 17 = P. Teneriffae.
saccharoides, Griseb. Fl. Brit. W. Ind. 557 = Paspalum saccharoides.
semidecumbens, Schult. Mant. ii. 164 = Saccharum spontaneum.
sphacelata, Benth. in Hook. Niger Fl. 559.—Afr. trop.
Teneriffae, Parl. in Webb & Bert. Phyt. Canar. iii. 425 = P. Teneriffae.
tonsa, Nees, Cat. Sem. Hort. Vratisl. (1836); ex ej. Fl. Afr. Austr. 16 = P. Teneriffae.
Wightii, Nees & Arn. ex Steud. Syn. Pl. Gram. 93 = P. megalanthum.

TRICHOLEPIS, DC. in Guill. Arch. Bot. ii. (1833) 515. *COMPOSITAE*, Benth. & Hook. f. ii. 475.
STICTOPHYLLUM, Edgew. in Trans. Linn. Soc. xx. (1846) 78.
amplexicaulis, *C. B. Clarke, Comp. Ind.* 240.—Ind. or.
angustifolia, *DC. Prod.* vi. 564.—Ind. or.
Candolleana, Wight & Arn. in Hook. Comp. Bot. Mag. i. (1835) 81. t. 4 = Volutarella divaricata.
elongata, *DC. Prod.* vi. 563.—Reg. Himal.
furcata, *DC. l. c.*; *et in Deless. Ic. Sel.* iv. t. 74.—Reg. Himal.
glaberrima, *DC. Prod.* vi. 564.—Ind. or.
Karensium, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 318.—Burma.
lanuginosa, Klotzsch, in Bot. Ergeb. Waldem. Reise, 79 = elongata.
montana, *Dalz. & Gibs. Bomb. Fl.* 131.—Ind. or.
nigricans, Edgew. in Trans. Linn. Soc. xx. (1846) 77 = furcata.
procumbens, Wight, Ic. Pl. t. 1139 = Volutarella divaricata.
radicans, *DC. Prod.* vi. 564.—Ind. or.

TRICHOLEPIS:—
Roylei, *Hook. f. Fl. Brit. Ind.* iii. 383.—Ind. bor. occ.
spartoides, C. B. Clarke, Comp. Ind. 241 = tibetica.
Stewartei, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 380.—Ind. bor. occ.
Stictophyllum, *C. B. Clarke, Comp. Ind.* 241.—Reg. Himal.
tibetica, *Hook. f. & Thoms. ex C. B. Clarke, l. c.*—Reg. Himal.

TRICHOLOBOS, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 290 = **Sisymbrium**, Tourn. (Crucif.).
capensis, Turcz. l. c. = S. Turczaninowii.

TRICHOLOBUS, Blume, Mus. Bot. Lugd. Bat. i. 236 (1850). *CONNARACEAE*, Benth. & Hook. f. i. 433.
cochinchinensis, *Baill. Adansonia*, ix. (1868–70) 150. —Cochinch.
connaroides, *F. Muell. Fragm.* viii. 224.—Austral.
ferrugineus, *Blume, Mus. Bot. Lugd. Bat.* i. 237.—Borneo.
fulvus, *Blume, l. c.*—Sumatra.

TRICHOLOMA, Benth. in DC. Prod. x. 426 (1846) = **Glossostigma**, Arn. (Scrophul.).
elatinoides, Benth. l. c. = G. elatinoides.

TRICHOLOPHUS, Spach, Hist. Vég. Phan. vii. 114 (1839) = **Polygala**, Tourn.
monspeliacus, Spach, l. c. 116 = P. monspeliaca.

TRICHOMARIA, 'Hook. & Arn.' ex Steud. Nom. ed. II. ii. 702 (1841), sphalm. = **Tricomaria**, Gill. (Malpigh.).

TRICHOMEMA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 195 (1821), err. typ. = seq.

TRICHONEMA, [Ker-Gawl.] in Bot. Mag. t. 575 (1802) = **Romulea**, Marat. (Irid.).
arenarium, Steud. Nom. ed. II. ii. 702 = R. arenaria.
barbatum, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 89 = R. barbata.
Bulbocodium, [Ker-Gawl.] in Bot. Mag. sub t. 575 = R. Bulbocodium.
Bulbocodium, Sm. Engl. Bot. t. 2549 = R. Columnae.
Bulbocodium, Webb & Berth. Phyt. Canar. iii. 312 = R. grandiscapa.
candidum, Tenore, Ind. Sem. Hort. Neap. (1842) 12. n. 4.—Afr. austr. ?
caulescens, [Ker-Gawl. in] Bot. Mag. t. 1392 = R. bulbocodioides.
chloroleucum, [Ker-Gawl.] l. c. sub t. 575 = R. chloroleuca.
Clusianum, Lange, in Kjoeb. Vidensk. Meddel. (1860) 75 = R. Clusiana.
coelestinum, Sweet, Hort. Brit. ed. I. 399 = Nemastylis coelestina.
collinum, Salisb. in Trans. Hort. Soc. i. (1812) 317 = R. Bulbocodium.
Columnae, Reichb. Fl. Germ. Excurs. 83 = R. Columnae.
croceum, Klatt, in Linnaea, xxxiv. (1865–66) 669 = R. Bulbocodium.
cruciatum, [Ker-Gawl.] in Bot. Mag. sub t. 575 = R. cruciata.
cupreum, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 89 = R. cuprea.
dichotomum, Klatt, in Linnaea, xxxiv. (1865–66) 666 = R. rosea.
edule, Herb. in Lindl. Bot. Reg. (1842) Misc. 86 = R. purpurascens.
elongatum, [Ker-Gawl.] in Bot. Mag. sub t. 1225 = R. elongata.
filifolium, Ker-Gawl. Irid. Gen. 82 = R. filifolia.
grandiflorum, Sprun. ex Nym. Consp. 704 = R. Bulbocodium.
grandiscapum, Webb & Berth. Phyt. Canar. iii. 312. t. 222 = R. grandiscapa.
hirsutum, Steud. Nom. ed. II. ii. 702; Klatt, in Linnaea, xxxiv. (1865–66) 665 = R. hirsuta.
humile, [Ker-Gawl.] in Bot. Mag. sub t. 575 = Geissorhiza humilis.
hypoxidiflorum, Salisb. in Trans. Hort. Soc. i. (1812) 316 = R. bulbocodioides.
latifolium, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 89 = R. latifolia.

TRICHONEMA:—
Linaresii, Gren. & Godr. Fl. Fr. iii. 238 = Romulea Requienii.
Linaresii, Klatt, in Linnaea, xxxiv. (1865–66) 660 = R. Linaresii.
longifolium, Salisb. in Trans. Hort. Soc. i. (1812) 316 = R. longifolia.
longitubum, Klatt, in Linnaea, xxxiv. (1865–66) 665 = Syringodea bicolor.
maritimum, Sibth. ex Baker, in Journ. Linn. Soc. xvi. (1877) 87 = R. Linaresii.
minimum, Tenore, Syll. Fl. Neap. 24 = R Columnae.
monadelphum, Sweet, Hort. Brit. ed. I. 399.—Afr. austr.
nivale, Boiss. & Kotschy, Diagn. Ser. II. iv. 92 = R. Bulbocodium.
ochroleucum, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 223 = R. chloroleuca.
ornithogaloides, A. Dietr. Sp. Pl. ii. 583.—Afr. austr.
parviflorum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 195 = R. Bulbocodium.
parviflorum, Steud. Nom. ed. II. ii. 702 = R. parviflora.
pudicum, [Ker-Gawl.] in Bot. Mag. sub t. 575 = R. rosea.
purpurascens, Sweet, Hort. Brit. ed. I. 399 = R. purpurascens.
Pylium, Herb. in Lindl. Bot. Reg. (1847) t. 40 = R. Bulbocodium.
quadrangulum, Sweet, Hort. Brit. ed. I. 399 = Geissorhiza quadrangula.
ramiflorum, Sweet, l. c. ed. II. 596 = R. ramiflora.
ramosum, Steud. Nom. ed. II. ii. 702 = R. arenaria.
recurvifolium, Ker-Gawl. Gen. Irid. 83 = R. bulbocodioides.
recurvum, Spreng. Syst. i. 149 = R. bulbocodioides.
reflexum, Steud. Nom. ed. II. ii. 702 = R. reflexa.
roseum, [Ker-Gawl.] in Bot. Mag. sub t. 575 = R. rosea.
simile, Steud. Nom. ed. II. ii. 702 = R. similis.
speciosum, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 223 = R. rosea.
spirale, Burch. Trav. i. 260 = R. spiralis.
subalpestre, Walp. Ann. i. 833, sphalm. = seq.
subpalustre, Herb. in Bot. Reg. (1847) t. 40. f. 1 = R. Bulbocodium.
tortuosum, Licht. Reise, iv. 51; Ker-Gawl. Gen. Irid. 83 = R. tortuosa.
uliginosum, Walp. Ann. i. 833 = R. Bulbocodium.
umbellatum, Klatt, in Linnaea, xxxiv. (1864–65) 670 = R. Bulbocodium.
vernum, Sprun. ex Nym. Consp. 705 = R. Linaresii.
Zeyheri, Steud. Nom. ed. II. ii. 702 = R. Zeyheri.

TRICHONEURA, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 148 = **Triodia**, R. Br. (Gramin.).
Hookeri, Anderss. l. c. 149 = Leptochloa Lindleyana.

TRICHOON, Roth, in Roem. Arch. i. III. (1798) 37 = **Phragmites**, Trin. (1820) (Gramin.).
Karka, Roth, l. c. = P. communis.
paniceus, Willd. ex Steud. Nom. ed. II. ii. 702 = Pennisetum setosum.

TRICHOPETALUM, Lindl. Bot. Reg. t. 1535 (1832) = **Bottionea**, Colla (Liliac.).
gracile, Lindl. l. c. = B. thysantoides.
stellatum, Lindl. l. c. = B. thysantoides.

TRICHOPHILA, Pritz. Ic. Ind. 1115 (1855), sphalm. = **Trichopilia**, Lindl. (Orchid.).

TRICHOPHORUM, Pers. Syn. i. 69 (1805) = **Eriophorum**, Linn. et **Scirpus**, Tourn. (Cyperac.).
alpinum, Pers. l. c. 70 = E. alpinum.
caespitosum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78; Enum. Pl. Trans. 694 = S. caespitosus.
comosum, A. Dietr. Sp. Pl. ii. 202 = E. comosum.
cyperinum, Pers. Syn. i. 69 = S. Eriophorum.
emergens, Norman, in 'Soc. Reg. Sc. Norveg. v.' (1868) 319 (Quid ?).—Norveg.
hudsonianum, Pers. Syn. i. 69 = E. alpinum.
hudsonicum, Steud. Nom. ed. II. ii. 702 = praec.
lineatum, Pers. Syn. i. 69 = S. lineatus.
Lockharti, G. Don, in Loud. Hort. Brit. 468 = E. Lockharti.
montevidense, G. Don, l. c. = Androtrichum polycephalum.
pendulum, Muhl. Desc. Gram. 44 = S. lineatus.

TRICHOPHYLLUM, Ehrh. Beitr. iv. 147 (1789) = **Eleocharis**, R. Br. (1810) (Cyperac.).

TRICHOPHYLLUM, Nutt. Gen. Am. ii. 166 (1818) = **Bahia**, Lag. (Compos.).
angustifolium, Less. in Linnaea, vi. (1831) 519 = Calea angustifolia.
integrifolium, Hook. Fl. Bor. Am. i. 316 = Eriophyllum caespitosum.
lanatum, Nutt. Gen. Am. ii. 168 = Eriophyllum caespitosum.
multiflorum, Nutt. in Journ. Acad. Philad. vii. (1834) 35 = Eriophyllum caespitosum.
oppositifolium, Nutt. Gen. Am. ii. 167 = oppositifolia.

TRICHOPILIA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836). *ORCHIDEAE*, Benth. & Hook. f. iii. 559.
HELCIA, Lindl. Bot. Reg. (1845) Misc. 17.
LEUCOCHYLE, Klotzsch, Ind. Sem. Hort. Berol. (1854) App. 1.
OLIVERIANA, Reichb. f. in Linnaea, xli. (1877) 111.
PILUMNA, Lindl. Bot. Reg. (1844) Misc. 73.
albida, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 362.—Venezuela.
Backhousiana, *Reichb. f. in Gard. Chron.* (1876) I. 816.—N. Granat.
callichroma, *Reichb. f. in Linnaea*, xli. (1877) 104.—N. Granat.
coccinea, Warsz. ex Lindl. & Paxt. Flow. Gard. ii. (1851–52) 79 = marginata.
costata, *Linden, Cat.* n. 22 (1869), 45.—Hab. ?
crispa, *Lindl. ex Linden, l. c.*—Am. centr.
dasyandra, *Reichb. f. Xenia Orch.* iii. 64.—N. Granat.
fragrans, *Reichb. f. in Otto, Hamb. Gartenz.* xiv. (1858) 229.—N. Granat.
Galeottiana, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 26.—Mexic.
grata, *Reichb. f. in Gard. Chron.* (1868) 1338.—Peruv. ?
hymenantha, *Reichb. f. in Bonplandia*, ii. (1854) 90. —Peruv.
Kienastiana, *Reichb. f. in Gard. Chron.* (1883) II. 166. —Hab. ?
laxa, *Reichb. f. in Otto, Hamb. Gartenz.* xiv. (1858) 229.—Bras.
lepida, Hort. ex W. G. Smith, in Flor. Mag. N. S. (1874) t. 98 = marginata.
maculata, *Reichb. f. in Bonplandia*, iii. (1855) 215.—Am. centr.
marginata, *Henfr. in Moore & Ayres, Gard. Mag. Bot.* iii. (1851) 185.—N. Granat.
mutica, *Reichb. f. in Walp. Ann.* vi. 679.—Ins. Trinit.
nobilis, *Reichb. f. Xenia Orch.* ii. 100.—N. Granat.
oicophyllax, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 97.—Hab. ?
picta, Lem. Illustr. Hortic. vi. (1859) Misc. 86; et t. 225 = Galeottiana.
Reichenheimia, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 353 = laxa.
rostrata, *Reichb. f. in Saund. Refug. Bot.* t. 100.—N. Granat.
sanguinolenta, *Reichb. f. Xenia Orch.* ii. 99.—Ecuador.
suavis, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 44.—Am. centr.
subulata, Reichb. f. in Flora, xlviii. (1865) 278 = hymenantha.
tortilis, *Lindl. Introd. Nat. Syst.* ed. II. 446; *et Bot. Reg.* t. 1863.—Mexic.
Turialvae, Batem. in Bot. Mag. t. 5550 = Galeottiana.
Turialvae, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 11.—Costa Rica.
Wageneri, *Reichb. f. in Otto, Hamb. Gartenz.* xiv. (1858) 229.—Venezuela.

TRICHOPODIUM, Lindl. Bot. Reg. sub t. 1543 (1832) = **Trichopus**, Gaertn. (Dioscor.).
angustifolium, Lindl. l. c. = Trichopus zeylanicus.
cordatum, Lindl. l. c. = Trichopus zeylanicus.
travancoricum, Bedd. Ic. Pl. Ind. Or. 68 = Trichopus zeylanicus.
zeylanicum, Thw. Enum. Pl. Zeyl. 291, 443 = Trichopus zeylanicus.

TRICHOPODIUM, Presl, Bot. Bemerk. 52 (1844) = **Dalea**, Linn. (Legumin.).
diffusum, Presl, l. c. 53 = D. diffusa.
glandulosum, Presl, l. c. 52.—Mexic.

TRICHOPTERIA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836), sphalm. = seq.

TRICHOPTERYA, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836), sphalm. = seq.

TRICHOPTERYX, Nees, Fl. Afr. Austr. 339 (1841). *GRAMINEAE*, Benth. & Hook. f. iii. 1162.
LOUDETIA, Hochst. in Flora, xxiv. (1841) I. Intell. 20; xxiv. (1841) 713.
Dregeana, *Nees, Fl. Afr. Austr.* 339.—Afr. austr.
flammea, Benth. in Journ. Linn. Soc. xix. (1881) 98, sphalm. = seq.
flammida, *Benth. l. c.* 59.—Bras.; Afr. trop.
simplex, *Benth. l. c.* 98.—Afr. austr.

TRICHOPTILIUM, A. Gray, in Torr. Bot. Mex. Bound. 97 (1858). *COMPOSITAE*, Benth. & Hook. f. ii. 415.
incisum, *A. Gray, l. c.*—Calif.

TRICHOPUS, Gaertn. Fruct. i. 44. t. 14 (1788). *DIOSCOREACEAE*, Benth. & Hook. f. iii. 745.
PODIANTHUS, Schnitzl. in Bot. Zeit. i. (1843) 739.
STEIREYA, Rafin. Fl. Tellur. iv. 100 (1836).
TRICHOPODIUM, Lindl. Bot. Reg. sub t. 1543 (1832).
piperifolius, Wall. ex Griff. in Trans. Linn. Soc. xix. (1845) 336 = Bragantia tomentosa.
zeylanicus, *Gaertn. Fruct.* i. 44. t. 14.—Ind. or.

TRICHORHIZA, Lindl. ex Steud. Nom. ed. II. ii. 702 (1841) = **Luisia**, Gaudich. (Orchid.).
teretifolia, Lindl. ex Steud. l. c.—Penang.; N. Caled.

TRICHORYNE, F. Muell. Census, 118 (1882), sphalm. = **Tricoryne**, R. Br. (Liliac.).

TRICHOSACME, Zucc. in Abh. Akad. Muench. iv. II. (1845) 11. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 765.
lanata, *Zucc. l. c.* 12.—Mexic.

TRICHOSANDRA, Decne. in DC. Prod. viii. 625 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 769.
borbonica, *Decne. l. c.* 626.—Ins. Mascar.

TRICHOSANTHA, Steud. Nom. ed. II. ii. 702 (1841), sphalm. = TRICHOSATHERA, Ehrh. = **Stipa**, Linn. (Gramin.).

TRICHOSANTHES, Linn. Gen. ed. I. 295 (1737). *CUCURBITACEAE*, Benth. & Hook. f. i. 821.
EOPEPON, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 31.
INVOLUCRARIA, Ser. in Mém. Soc. Phys. Genèv. iii. I. (1825) 25. t. 5.
PLATYGONIA, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 33. t. 5.
amara, Blanco, Fl. Filip. ed. I. 774 = Luffa acutangula.
amara, *Linn. Sp. Pl.* 1008 (sp. dub.).—Ins. S. Doming.
anaimlaiensis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 47.—Ind. or.
anamalayana, Bedd. in Trans. Linn. Soc. xxv. (1865) 217 = praec.
Anguina, *Linn. Sp. Pl.* 1008.—As. trop.
angulata, *Lam. Encyc.* i. 190 (sp. dub.).—Ind. or.
aspera, Herb. Madr. ex Wall. Cat. n. 6686 E = pubera.
asperifolia, Zipp. ex Span. in Linnaea, xv. (1841) 206, nomen = pubera.
Beccariana, *Cogn. in DC. Monog. Phan.* iii. 380.—Sumatra.
borneensis, *Cogn. l. c.* 369.—Borneo.
bracteata, *Voigt, Hort. Suburb. Calc.* 58.—Ind. or.
caudata, Willd. Sp. Pl. iv. 600 = cuspidata.
celebica, *Cogn. in DC. Monog. Phan.* iii. 385.—Ins. Celebes.
chinensis, *Ser. in DC. Prod.* iii. 315 (= cucumeroides?).—China.
cochinchinensis, M. Roem. Syn. Pepon. 96 = cucumerina.
colubrina, Jacq. f. Eclog. t. 128 = T. Anguina.
cordata, *Roxb. Hort. Beng.* 70; *Fl. Ind.* iii. 703.—Ind. or.
cordata, Wall. Cat. n. 6686 B = himalensis.
cordata, Wall. l. c. 6688 C, E = pubera.
coriacea, *Blume, Bijdr.* 935.—Sumatra.

TRICHOSANTHES:—
corniculata, Lam. Encyc. i. 191 = Ceratosanthes tuberosa.
costata, Blume, Bijdr. 933 = Gymnopetalum cochinchinense.
cucumerina, Buch.-Ham. ex Wall. Cat. n. 6688 B = pubera?
cucumerina, *Linn. Sp. Pl.* 1008.—As. et Austral. trop.
cucumerina, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 82 = cucumeroides.
cucumerina, Thunb. Fl. Jap. 332 = japonica.
cucumerina, Wall. Cat. n. 6690 E = Gymnopetalum cochinchinense.
cucumeroides, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 172.—Japon.
cuspidata, *Lam. Encyc.* i. 188.—Ind. or.
dicaelosperma, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 609.—Reg. Himal.
dioica, *Roxb. Hort. Beng.* 70; *Fl. Ind.* iii. 701.—Ind. or.
dioica, Wall. Cat. n. 6692 C = Melothria maderaspatana.
erigera, D. Dietr. Syn. Pl. v. 370, sphalm. = ovigera.
Fatoa, Buch.-Ham. in Wall. Cat. n. 6695 = Gymnopetalum cochinchinense.
foetidissima, Jacq. Coll. ii. 341 = Kedrostis foetidissima.
globosa, *Blume, Bijdr.* 936.—Java.
grandibracteata, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 98, 99 = Wallichiana.
grandiflora, *Blume, Bijdr.* 934.—Java.
grandiflora, Wall. Cat. n. 6685 = Hodgsonia heteroclita.
Hearnii, *F. Mueil. ex Benth. Fl. Austral.* iii. 315.—Austral.
heteroclita, Roxb. Hort. Beng. 70; Fl. Ind. iii. 705 = Hodgsonia heteroclita.
hexasperma, Blume, Bijdr. 935 = Hodgsonia heteroclita.
himalensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 608.—Reg. Himal.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. I. 677.—Java.
incisa, *Rottl. ex Rosenth. Syn. Pl. Diaph.* 683, nomen.—Hab.?
integrifolia, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 99.—Ind. or.; Burma.
integrifolia, Thw. Enum. Pl. Zeyl. 127 = Thwaitesii.
japonica, *Regel, Ind. Sem. Hort. Petrop.* 1868 (1869) 90.—Japon.
javanica, Miq. Pl. Jungh. 399; et Fl. Ind. Bat. i. I. 678 = Thladiantha cordifolia.
Kadam, Miq. Fl. Ind. Bat. Suppl. 331 = Hodgsonia heteroclita.
Kakidonta, Roxb. ex Wight & Arn. Prod. 350 = pubera.
Kirilowii, *Maxim. Prim. Fl. Amur.* 482.—Mongol.
laciniosa, Klein, ex Willd. Sp. Pl. iv. 601 = cucumerina.
laciniosa, Wight, ex Wall. Cat. n. 6689 A, B = pubera.
Lepiniana, *Cogn. in DC. Monog. Phan.* iii. 377.—Reg. Himal.; Ind. or.
lobata, Buch.-Ham. ex Wall. Cat. n. 6687 E = T. Anguina.
lobata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 703 = cucumerina.
longiflora, *Cogn. in DC. Monog. Phan.* iii. 374.—N. Guin.
lucioniana, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 95 = pubera.
macrocarpa, Blume, Bijdr. 935 = Hodgsonia heteroclita.
macrosiphon, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 308 = cordata.
multiloba, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 607 = Wallichiana.
multiloba, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 82.—Japon.
nervifolia, *Linn. Sp. Pl.* 1008.—Ind. or.
officinalis, Buch.-Ham. ex Wall. Cat. n. 6694 = dioica.
olida, F. Muell. in Trans. Soc. Bot. Edinb. vii. (1863) 491 = cucumerina.
ovata, *Cogn. in DC. Monog. Phan.* iii. 365.—Reg. Himal.
ovigera, *Blume, Bijdr.* 934.—Malaya.
palmata, Benth. Fl. Austral. iii. 315 = subvelutina.
palmata, Hance, in Journ. Bot. xvi. (1878) 227 = Kirilowii.
palmata, *Linn. Syst.* ed. X. 1278.—Hab.?
palmata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 704 = pubera.

TRICHOSANTHES:—
palmata, Wall. Cat. n. 6688 = cordata.
parviflora, Wight, ex Wall. l. c. 6690 E = Gymnopetalum cochinchinense.
Patara, *M. Roem. Syn. Pepon.* 97 (sp. dub.).—Ins. Tahiti.
pedatifolia, Miq. Fl. Ind. Bat. i. I. 677 = cucumerina.
pentaphylla, *F. Muell. ex Benth. Fl. Austral.* iii. 314.—Austral.
Perrottetiana, *Cogn. in DC. Monog. Phan.* iii. 362.—Ind. or.
Petala, Buch.-Ham. ex Wall. Cat. n. 6687 D = T. Anguina.
pilosa, Buch.-Ham. ex Wall. l. c. n. 6691 = cucumerina.
pilosa, *Lour. Fl. Cochinch.* 588.—Cochinch.
pilosa, Wall. Cat. n. 6687, partim = bracteata.
pomiformis, Macfad. Fl. Jamaic. ii. 143 = Cionosicys pomiformis.
pubera, *Blume, Bijdr.* 936.—Ind. or.; Malaya.
punctata, Linn. Amoen. Acad. iii. 423 = Fevillea cordifolia.
quadricirrha, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 82 = cucumeroides.
quinquangulata, *A. Gray, Bot. U. St. Expl. Exped.* i. 645.—Ins. Sulu.
reniformis, Kurz, in Journ. As. Soc. Beng. (1871) II. 57 = dicaelosperma.
reniformis, Miq. Fl. Ind. Bat. i. I. 675 = cucumerina.
Russeliana, Wall. Cat. n. 6696 = Momordica dioica.
scabra, *Lour. Fl. Cochinch.* 589.—Cochinch.
subvelutina, *F. Muell. ex Cogn. in DC. Monog. Phan.* iii. 366.—Austral.
sumatrana, *Cogn. l. c.* 373.—Sumatra.
tamnifolia, *Poir. Encyc. Suppl.* i. 386.—Porto Rico.
Tatoa, *Buch.-Ham. ex Wall. Cat.* n. 6695 (sp. dub.).—Ind. or.
Theba, Buch.-Ham. ex Wall. Cat. n. 6684 A = Hodgsonia heteroclita.
Thwaitesii, *Cogn. in DC. Monog. Phan.* iii. 387.—Zeylan.
tricuspidata, *Lour. Fl. Cochinch.* 529.—Cochinch.
tricuspis, Miq. Fl. Ind. Bat. i. I. 679 = pubera.
trifoliata, *Blume, Bijdr.* 936.—Malaya.
truncata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 608.—Reg. Himal.
tuberosa, Roxb. ex Wight & Arn. Prod. 349 = cordata.
tuberosa, Willd. Sp. Pl. iv. 601 = Ceratosanthes tuberosa.
Turolata, Buch.-Ham. ex Wall. Cat. n. 6688 A = pubera.
villosa, *Blume, Bijdr.* 934.—Java.
villosula, *Cogn. in DC. Monog. Phan.* iii. 362.—Ind. or.
Wallichiana, *Wight, in Ann. & Mag. Nat. Hist.* Ser. I. viii. (1842) 270.—Ind. or.
Wawraei, *Cogn. in DC. Monog. Phan.* iii. 384.—Singapore.
Wightiana, M. Roem. Syn. Pepon. 96 = cucumerina.

TRICHOSANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

TRICHOSATHERA, Ehrh. Beitr. iv. (1789) 149 = **Stipa**, Linn. (Gramin.).

TRICHOSCYPHA, Hook. f. in Benth. & Hook. f. Gen. i. 423 (1862). *ANACARDIACEAE*, Benth. & Hook. f. l. c.
acuminata, *Engl. Bot. Jahrb.* i. (1881) 425.—Afr. trop.
bijuga, *Engl. l. c.*—Afr. trop.
imbricata, *Engl. l. c.* 426.—Afr. trop.
longifolia, *Engl. l. c.* 425.—Afr. trop.
lucens, *Oliver, Fl. Trop. Afr.* i. 444.—Afr. trop.
Mannii, *Hook. f. in Benth. & Hook. f. Gen.* i. 423.—Afr. trop.
Oliveri, *Engl. Bot. Jahrb.* i. (1881) 425.—Afr. trop.
patens, *Engl. l. c.*—Afr. trop.
reticulata, *Engl. in DC. Monog. Phan.* iv. 307.—Afr. trop.

TRICHOSERIS, Poepp. & Endl. Nov. Gen. et Sp. t. 263 (1845), sphalm. = TROCHOSERIS, Poepp. & Endl. = **Troximon**, Nutt. (Compos.).

TRICHOSERIS, Sch. Bip. in Flora, xxii. (1839) 19 = **Pterotheca**, Cass. (Compos.).
Rueppellii, Sch. Bip. l. c. = P. bifida.

TRICHOSIA, Blume, Bijdr. Tabell. 11 (1825) = TRICHOTOSIA, Blume = **Eria**, Lindl. (Orchid.).

TRICHOSIPHON, Schott & Endl. Melet. 34 (1832) = **Sterculia**, Linn.
australe, Schott & Endl. l. c. = S. Trichosiphon.

TRICHOSMA, Lindl. Bot. Reg. (1842) t. 21. *ORCHIDEAE*, Benth. & Hook. f. iii. 518.
cylindropoda, Griff. Notul. iii. 299 = suavis.
suavis, *Lindl. Bot. Reg.* (1842) t. 21.—Reg. Himal.

TRICHOSPERMUM, Blume, Bijdr. 56 (1825). *TILIACEAE*, Benth. & Hook. f. i. 236.
BIXAGREWIA, Kurz, in Journ. Bot xiii (1875) 325. t. 160.
DICLIDOCARPUS, A. Gray, Bot. U. St. Expl. Exped. i. 200. t. 14 (1854).
javanicum, *Blume, Bijdr.* 57.—Java.
mexicanum, Baill. Hist. Pl. iv. 179 = Belotia greviaefolia.
Richii, *Seem. in Bonplandia*, ix. (1861) 254 (Ritchei); *Fl. Vit.* 27.—Ins. Fiji.

TRICHOSPIRA, H. B. & K. Nov. Gen. et Sp. iv. 27. t. 312 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 383.
biaristata, Less. in Linnaea, iv. (1829) 343 = menthoides.
menthoides, *H. B. & K. Nov. Gen. et Sp.* iv. 23. t. 312.—Am. austr.
Prieurii, DC. Prod. v. 91 = menthoides.
Pulegium, Mart. ex DC. l. c. = menthoides.

TRICHOSPORUM, D. Don, in Edinb. Phil. Journ. vii. (1822) 84 = **Aeschynanthus**, Jack (1823) (Gesner.).
albidum, Nees, in Flora, viii. (1825) 144 = A. albida.
angustifolium, Nees, l. c. = A. angustifolia.
grandiflorum, D. Don, in Edinb. Phil. Journ. vii. (1822) 85; et Prod. Fl. Nep. 125 = A. grandiflora.
ovatum, D. Don, ex C. B. Clarke, in DC. Monog. Phan. v. 41 = A. radicans.
parviflorum, D. Don, in Edinb. Phil. Journ. vii. (1822) 85; et Prod. Fl. Nep. 125 = A. ramosissima.
pulchrum, Blume, Bijdr. 764 = A. pulchra.
radicans, Nees, in Flora, viii. (1825) 144 = A. radicans.
volubile, Nees, l. c. = A. volubilis.

TRICHOSTACHYS, Hook. f. in Benth. & Hook. f. Gen. ii. 128 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
aurea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 227.—Afr. trop.
ciliata, *Hiern, l. c.*—Afr. trop.
longifolia, *Hiern, l. c.*—Afr. trop.
petiolata, *Hiern, l. c.*—Afr. trop.

TRICHOSTACHYS, Welw. Synopse, 19 (1862) = **Faurea**, Harv. (Proteac.).
speciosa, Welw. l. c. = F. speciosa.

TRICHOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 81 = **Athrixia**, Ker-Gawl. (Compos.).
asteroides, Turcz. l. c. = A. multiceps.

TRICHOSTEMA, Gronov. ex Linn. Hort. Cliff. 493 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1219.
arizonicum, *A. Gray, in Proc. Am. Acad.* viii. (1872) 371.—Arizona.
brachiatum, Lam. Illustr. iii. 71 = lineare.
brachiatum, Linn. Sp. Pl. 598 = Isanthus coeruleus.
dichotomum, *Linn. l. c.*—Am. bor.
dichotomum, Roth, Nov. Pl. Sp. 280 = lineare.
dichotomum, Torr. Bot. Mex. Bound. 134 = arizonicum.
lanatum, *Benth. Lab. Gen. et Sp.* 659.—Calif.
lanceolatum, *Benth. l. c.*—Am. bor. occ.
laxum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 387.—Calif.
lineare, Nutt. Gen. Am. ii. 39 = seq.?
lineare, *Walt. Fl. Carol.* 164.—Am. bor.
linearifolium, *Bertol. Misc. Bot.* xiii. 15. t. 2.—Am. bor.
micranthum, *A. Gray, Syn. Fl. N. Am.* ii. 348.—Calif.
oblongum, *Benth. Lab. Gen. et Sp.* 659.—Am. bor. occ.
ovatum, *Curran, in Bull. Calif. Acad.* i. III. (1885) 154.—Calif.
Parishii, Vasey, in Coult. Bot. Gaz. vi. (1881) 173 = lanatum.
pilosum, Roth, Nov. Pl. Sp. 280 = dichotomum.
Sandersoni, Pritz. Ic. Ind. ii. 276, sphalm. = Tryphostemma Sandersoni.
spiralis, Lour. Fl. Cochinch. 371 = Orthosiphon stamineus.

TRICHOSTEMMA, Cass. in Dict. Sc. Nat. xlvi. 399, 409 (1827) = **Wedelia**, Jacq. (Compos.).
hispida, Cass. l. c. 409 = W. trichostephia.

TRICHOSTEPHIUM, Cass. in Dict. Sc. Nat. lv. 266, in adnot. (1828) = seq.

TRICHOSTEPHUS, Cass. in Dict. Sc. Nat. lx. 618 (1830) = **Wedelia**, Jacq. (Compos.).
hispidus, Cass. l. c. = W. trichostephia.

TRICHOSTERIGMA, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248; et in Abh. Akad. Berl. (1860) 41 = **Euphorbia**, Linn.
californicum, Klotzsch & Garcke, ll. cc. 248; 42 = E. californica.
fulgens, Klotzsch & Garcke, ll. cc. = E. fulgens.
Hindsianum, Klotzsch & Garcke, ll. cc. = E. Hindsiana.
miserum, Klotzsch & Garcke, ll. cc. = E. misera.

TRICHOSTIGMA, A. Rich. Ess. Fl. Cub. i. 627 (1845) = **Villamilla**, Ruiz & Pav. (Phytol.).
rivinoides, A. Rich. l. c. 628 = V. octandra.

TRICHOSTOSIA, Griff. Notul. iii. 331 (1851), sphalm. = TRICHOTOSIA, Blume = **Eria**, Lindl. (Orchid.).

TRICHOTHALAMUS, Spreng. Anleit. ii. 864 (1818); Lehm. in Nov. Act. Nat. Cur. x. (1821) 585. t. 49 = **Potentilla**, Linn. (Rosac.).
albus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 370 = P. alba.
alchemilloides, Spreng. Anleit. ii. 864 = P. alchemilloides.
caulescens, Spreng. l. c. = P. caulescens.
Clusianus, Spreng. l. c. = P. Clusiana.
lignosus, Lehm. in Nov. Act. Nat. Cur. x. (1821) 585. t. 49 = P. lignosa.
lupinoides, Spreng. Anleit. ii. 864 = P. nivalis.
nitidus, Spreng. l. c. = P. nitida.
petiolulatus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 370 = P. caulescens.
tridentatus, Spreng. Anleit. ii. 864 = P. tridentata.
Valderia, Spreng. l. c. = P. Valderia.

TRICHOTOSIA, Blume, Bijdr. 342 (1825) = **Eria**, Lindl. (Orchid.).
annulata, Blume, l. c. 343 = E. annulata.
biflora, Griff. Notul. iii. 331; Ic. Pl. Asiat. t. 315 = E. biflora.
ciliata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. v. (1853) 492 = E. ciliata.
ferox, Blume, Bijdr. 342 = E. ferox.
ferox, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 184 = E. rubiginosa.
microphylla, Blume, Bijdr. 343 = E. microphylla.
pauciflora, Blume, l. c. = E. pauciflora, *Blume*.

TRICLANTHERA, Rafin. Sylva Tellur. 108 (1838) = **Crataeva**, Linn. (Cappar.).
corymbosa, Rafin. l. c. = C. magna.
falcata, Rafin. l. c. = C. falcata.

TRICLICERAS, Thonn. ex DC. Pl. Rar. Jard. Genèv. 56, in obs. (1829) = **Wormskioldia**, Thonn. (Turnerac.).
raphanoides, DC. l. c. = W. heterophylla.

TRICLINIUM, Rafin. Fl. Ludov. 79 (1817) = **Sanicula**, Tourn. (Umbell.).
marilandica, Rafin. in Am. Monthly Mag. (1818) 267 = S. marilandica.
odoratum, Rafin. Fl. Ludov. 80 = S. canadensis.

TRICLISIA, Benth. in Benth. & Hook. f. Gen. i. 39 (1862). *MENISPERMACEAE*, Benth. & Hook. f. l. c.
coriacea, *Oliver, Fl. Trop. Afr.* i. 49.—Afr. trop.
loucoubensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 458.—Madag.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 48.—Afr. trop.
patens, *Oliver, l. c.* 49.—Afr. trop.
subcordata, *Oliver, l. c.*—Afr. trop.

TRICLISPERMA, Rafin. Specch. i. 117 (1814) = **Polygala**, Tourn.
grandiflora, Rafin. l. c. = P. paucifolia.

TRICLISSA, Salisb. Gen. Pl. Fragm. 75 (1866) = **Kniphofia**, Moench (Liliac.).
Uvaria, Salisb. l. c. = K. aloides.

TRICOILENDUS, Rafin. Fl. Tellur. ii. 97 (1836) = **Indigofera**, Linn. (Legumin.).
microphylus, Rafin. l. c. = I. gracilis.

TRICOMARIA, Gill. ex Hook. Bot. Misc. iii. (1833) 157. t. 101. *MALPIGHIACEAE*, Benth. & Hook. f. i. 258.
Usillo, *Hook. & Arn. l. c.* 158.—Chili.

TRICONDYLUS, Salisb. in Knight, Prot. 121 (1809) = **Lomatia**, R. Br. (1810) (Proteac.).
ferrugineus, Knight, l. c. 123 = L. ferruginea.
myricaefolius, Knight, l. c. 122 = L. longifolia.
silaifolius, Knight, l. c. = L. silaifolia.
tinctorius, Knight, l. c. = L. tinctoria.

TRICORYNE, R. Br. Prod. 278 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 796.
acaulis, D. Dietr. Syn. Pl. ii. 1152 = Ophiopogon japonicus.
anceps, *R. Br. Prod.* 278.—Austral.
anceps, Endl. in Lehm. Pl. Preiss. ii. 36 = elatior.
caulescens, D. Dietr. Syn. Pl. ii. 1152 = Ophiopogon japonicus.
elatior, *R. Br. Prod.* 278.—Austral.
humilis, *Endl. in Lehm. Pl. Preiss.* ii. 36.—Austral.
muricata, *Baker, in Journ. Linn. Soc.* xv. (1876) 363.—Austral.
platyptera, *Reichb. f. Beitr. Syst. Pfl.* 72.—Austral.
pterocaulon, Baker, in Journ. Linn. Soc. xv. (1876) 363 = platyptera.
scabra, R. Br. Prod. 278 = elatior.
simplex, R. Br. l. c.—Austral.
tenella, R. Br. l. c. = elatior.

TRICOSCYPHA, Engl. in DC. Monog. Phan. iv. 303 (1883), sphalm. = **Trichoscypha**, Hook. f. (Anacard.).

TRICOSTULARIA, Nees, in Lehm. Pugill. viii. 50 (1844); et in Lehm. Pl. Preiss. ii. 83 (1846). *CYPERACEAE*, Benth. & Hook. f. iii. 1064.
DISCOPODIUM, Steud. Syn. Pl. Cyp. 150 (1855).
compressa, *Nees, in Lehm. Pugill.* viii. 51; *et in Lehm. Pl. Preiss.* ii. 83.—Austral.
fimbristyloides, Benth. Fl. Austral. vii. 384 = Cladium undulatum.
Neesii, *Lehm. Pugill.* viii. 52; *et in Lehm. Pl. Preiss.* ii. 83.—Austral.
paludosa, *Benth. Fl. Austral.* vii. 382.—Austral.
pauciflora, *Benth. l. c.* 383.—Austral.

TRICRATUS, L'Hérit. ex Willd. Sp. Pl. i. 807 (1797) = **Abronia**, Juss. (Nyctag.).
admirabilis, L'Hérit. ex Willd. l. c. = A. umbellata.
arenarius, Spreng. Syst. iv. Cur. Post. 53 = A. arenaria.
parviflorus, Spreng. Syst. i. 536 = Colignonia parviflora.

TRICUSPIDARIA, Ruiz & Pav. Prod. 64. t. 36 (1794). *TILIACEAE*, Benth. & Hook. f. i. 240.
CRINODENDRON, Molina, Sagg. Chile, 179 (1782).
TRICUSPIS, Pers. Syn. ii. 9 (1807).
dependens, *Ruiz & Pav. Fl. Per.* iv. t. 403; *Syst. Veg.* 112.—Peruv.
hexapetala, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 576 = dependens.
lanceolata, *Miq. in Linnaea*, xxv. (1852) 650.—Chili.
Patagua, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 51; Contrib. Bot. ii. 187. t. 82 = dependens.

TRICUSPIS, Beauv. Agrost. 77. t. 15. f. 10 (1812) = **Triodia**, R. Br. (Gramin.).
acuminata, Munro, ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 335 = Triodia acuminata.
albescens, Munro, ex A. Gray, l. c. = Triodia albescens.
ambigua, Chapm. Fl. S. U. St. 559 = Triodia ambigua.
caroliniana, Beauv. Agrost. 77 = Triodia cuprea.
cornuta, A. Gray, Man. Bot. N. U. St. ed. I. 590; Wood, Am. Bot. & Flor. 398 = Triplasis americana.

TRICUSPIS :—
latifolia, Griseb. in Goett. Abh. xix. (1874) 259.—Reg. Argent.
monstra, Munro, ex Hemsl. Diagn. Pl. Nov. 56 = Scleropogon brevifolius.
mutica, Torr. in Pacif. Rail. Rep. iv. 156 = Triodia mutica.
noveboracensis, Beauv. Agrost. 77 = Triodia cuprea.
pulchella, Torr. in Pacif. Rail. Rep. iv. 156 = Triodia pulchella.
purpurea, A. Gray, Man. Bot. N. U. St. ed. I. 589 = Triplasis purpurea.
quinquefida, Beauv. ex G. Don, in Loud. Hort. Brit. 31 = Triodia cuprea.
seslerioides, Torr. Fl. N. York, ii. 463 = Triodia seslerioides.
simplex, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 532.—Cuba.
stricta, Nutt. ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 335 = Triodia stricta.
texana, Thurb. ex S. Wats. in Proc. Am. Acad. xviii. (1883) 180 = Triodia texana.
trinerviglumis, Munro, ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 333 = Triodia trinerviglumis.

TRICUSPIS, Pers. Syn. ii. 9 (1807) = **Tricuspidaria**, Ruiz & Pav. (Tiliac.).
dependens, Pers. l. c. = Tricuspidaria dependens.

TRICYCLA, Cav. Ic. vi. 78. t. 598 (1801). *NYCTAGINEAE*, Benth. & Hook. f. iii. 7.
peruviana, Poir. Encyc. Suppl. v. 359 = Bougainvillaea peruviana.
spectabilis, Poir. l. c. = Bougainvillaea spectabilis.
spinosa, *Cav. Ic.* vi. 79. t. 598 ; *et in Anal. Cienc. Nat.* v. (1802) 63.—Reg. Argent.

TRICYRTIS, Wall. Tent. Fl. Nap. 61. t. 46 (1826). *LILIACEAE*, Benth. & Hook. f. iii. 831.
COMPSANTHUS, Spreng. Syst. iv. Cur. Post. 137 (1827).
COMPSOA, D. Don, Prod. Fl. Nep. 50 (1825), nomen prius.
elegans, Wall. Tent. Fl. Nap. ii. t. 46 = pilosa.
flava, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 434.—Japon.
formosana, *Baker in Journ. Linn. Soc.* xvii. (1879) 465.—Ins. Formosa.
hirta, *Hook. Bot. Mag.* t. 5355.—Japon.
japonica, Miq. in Versl. en Med. Kon. Akad. Wetensch. II. ii. (1868) 86 ; et Ann. Mus. Bot. Lugd. Bat. iii. 155, 201 = hirta.
latifolia, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 435.—Japon.
macropoda, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* II. ii. (1868) 86.—Japon.
pilosa, *Wall. Tent. Fl. Nap.* ii. 62 [t. 46].—Reg. Himal.

TRIDACHNE, Liebm. ex Lindl. & Paxt. Flow. Gard. iii. (1852) 45 = **Notylia**, Lindl. (Orchid.).
virens, Liebm. ex Lindl. & Paxt. l. c. = N. Huegelii.

TRIDACTYLINA, Sch. Bip. Tanacet. 48 (1844) = **Chrysanthemum**, Tourn. (Compos.).
Kirilowii, Sch. Bip. l. c. 49 (= *Pyrethrum Kirilowii*). —Sibir. baical.

TRIDACTYLITES, Haw. Saxif. Enum. 21 (1821) = **Saxifraga**, Tourn.
annua, Haw. l. c. = S. Tridactylites.
petraea, Haw. l. c. 22 = S. petraea.

TRIDALIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Quid ?).

TRIDAX, Linn. Hort. Cliff. 418 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 392.
BALBISIA, Willd. Sp. Pl. iii. 2214 (1803).
BARTOLINA, Adans. Fam. ii. 124 (1763).
CARPHOSTEPHIUM, Cass. in Dict. Sc. Nat. xliv. 62 (1826).
GALINSOGEA, H. B. & K. Nov. Gen. et Sp. iv. 252. t. 386 (1820).
MANDONIA, Wedd. in Bull. Soc. Bot. Fr. xi. (1864) 50. t. 1.
PTILOSTEPHIUM, H. B. & K. Nov. Gen. et Sp. iv. 253. tt. 387, 388 (1820).
SOGALGINA, Cass. in Bull. Soc. Philom. (1818) 31.

TRIDAX :—
angustifolia, *Spruce, ex Benth. & Hook. f. Gen.* ii. 392.—Ecuador.
balbisioides, *A. Gray, in Proc. Am. Acad.* xv. (1880) 39.—Mexic.
bicolor, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 104.—Am. bor. occ.
brachylepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 207.—Mexic.
candidissima, *A. Gray, in Proc. Am. Acad.* xv. (1880) 39.—Mexic.
coronopifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 207.—Mexic.
galardioides, Hook. & Arn. Bot. Beech. Voy. 148 = Layia galardioides.
Mandonii, *Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 82 ; *et in Linnaea*, xxxiv. (1865-66) 536, nomen.—Bolivia.
obovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 187.—Mexic.
Palmeri, *A. Gray, in Proc. Am. Acad.* xv. (1880) 38.—Mexic.
procumbens, *Linn. Sp. Pl.* 900.—Am. trop.
trifida, *A. Gray, in Proc. Am. Acad.* xv. (1880) 39.—Mexic.
trilobata, *Hemsl. Biol. Centr. Am. Bot.* ii. 208.—Mexic.

TRIDENS, Roem. & Schult. Syst. ii. 34 (1817) = **Triodia**, R. Br. (Gramin.).
ambigua, Schult. Mant. ii. 333 = Triodia ambigua.
brasiliensis, Nees, ex Steud. Syn. Pl. Gram. 248 (= *Uralepsis brasiliensis*).—Bras.
capensis, Nees, in Linnaea, vii. (1832) 324 (= *Uralepsis capensis*).—Afr. austr.
quinquefida, Schult. Mant. ii. 333 = Triodia cuprea.
verticillata, Meyen, Reise, i. 408, nomen = Diplachne verticillata.
virens, Nees, Agrost. Bras. 476 (= *Uralepsis virens*).—Bras.

TRIDENTEA, Haw. Syn. Pl. Succ. 34 (1812) = **Stapelia**, Linn.
depressa, Haw. ex Schult. Syst. vi. 850 = S. patula.
gemmiflora, Haw. Syn. Pl. Succ. 34 = S. gemmiflora.
hircosa, Schult. Syst. vi. 850 = S. hircosa.
Juvencula, Schult. l. c. 16 = S. Juvencula.
moschata, Haw. Syn. Pl. Succ. 25 = S. hircosa.
paniculata, Schult. Syst. vi. 850 = S. paniculata.
rugosa, Schult. l. c. = S. rugosa.
Simsii, Schult. l. c. = S. Simsii.
stygia, Haw. Syn. Pl. Succ. 34 = S. stygia.
vetula, Haw. l. c. 34 = S. vetula.

TRIDERMIA, Rafin. Sylva Tellur. 145 (1838) = **Grewia**, Linn. (Tiliac.).
papillosa, Rafin. l. c. = G. villosa.

TRIDESMIS, Lour. Fl. Cochinch. 576 (1790) = **Croton**, Linn. (Euphorb.).
hispida, Lour. l. c.—China.
tomentosa, Lour. l. c. = C. crassifolius.

TRIDESMIS, Spach, Hist. Vég. Phan. v. 358 (1836) ; et in Ann. Sc. Nat. Sér. II. v. (1836) 351. t. 6 = **Cratoxylon**, Blume (Hyperic.).
Billardieri, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 352. t. 6 = C. formosum.
formosa, Korth. Verh. Nat. Gesch. Bot. 179. t. 37 = C. formosum.
Jackii, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 352 = C. formosum.
ochnoides, Spach, Hist. Vég. Phan. v. 359 = C. formosum.
pruniflora, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 293 = C. pruniflorum.

TRIDESMUS, Steud. Nom. ed. II. i. 445 (1840), in syn., err. typ. = TRIDESMIS, Lour. = **Croton**, Linn. (Euphorb.).

TRIDIA, Korth. in Hoev. & De Vriese, Tijdschr. iii. (1836) 17. t. 1 = **Hypericum**, Tourn.
frankenioides, Korth. l. c. 18 = H. japonicum.

TRIDIANISIA, Baill. in Bull. Soc. Linn. Par. i. (1879) 197. *OLACINEAE*.
Chapelieri, *Baill. l. c.*—Madag.

TRIDIMERIS, Baill. Adansonia, ix. (1869) 219. *ANONACEAE*.
Hahniana, *Baill. l. c.*—Mexic.

TRIDOPHYLLUM, Neck. Elem. ii. 93 (1790) = **Potentilla**, Linn. (Rosac.).

TRIDYNIA, Rafin. ex Steud. Nom. ed. II. ii. 704 (1841) = **Lysimachia**, Tourn. (Primul.).

TRIDYRA, Steud. Nom. ed. II. ii. 84 (1841), sphalm. = praec.

TRIENDILIX, Rafin. New Fl. Am. i. 85 (1836) = **Glycine**, Linn. (Legumin.).
clandestina, Rafin. l. c. = G. clandestina.

TRIENTALIS, Rupp. ex Linn. Syst. ed. I (1735) ; Gen. ed. I. 109 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 636.
alsinaeflora, Gilib. Fl. Lituan. i. 31 = T. europaea.
americana, *Pursh, Fl. Am. Sept.* i. 252.—Am. bor.
arctica, Fisch. ex Hook. Fl. Bor. Am. ii. 121 = europaea.
borealis, Rafin. in Med. Repos. N. York, II. v. (1808) 354 = europaea.
capensis, Linn. Sp. Pl. 344 = Crassula Septas.
europaea, *Linn. l. c.*—Europ. ; As. bor. ; Am. bor. occ.
europaea, Michx. Fl. Bor. Am. i. 220 = americana.
heptagyna, Crantz, Inst. ii. 365 = Crassula Septas.
latifolia, Hook. Fl. Bor. Am. ii. 121 = europaea.

TRIEXASTIMA, Rafin. Fl. Tellur. iv. 121 (1836) = **Heteranthera**, Ruiz & Pav. (Ponted.).

TRIFAX, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4 (Quid ?).

TRIFILLIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 383 = **Medicago**, Tourn. (Legumin.).
platycarpos, Medic. l. c. = M. platycarpa.

TRIFOLIASTRUM, Moench, Meth. 123 (1794) = **Trigonella**, Linn. (Legumin.).
caeruleum, Moench, l. c. = Trigonella caerulea.

TRIFOLIUM, [Tourn.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 229 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 487.
AMARENUS, Presl, Symb. Bot. i. 47 (1830).
AMORIA, Presl, l. c. 43 (1830).
CALYCOMORPHUM, Presl, l. c. 50 (1830).
DACTIPHYLLON, Rafin. in Am. Monthly Mag. (1818) 268.
GALEARIA, Presl, Symb. Bot. i. 49 (1830).
LAGOPUS, Bernh. Syst. Verz. Erf. 229 (1800).
LOXOSPERMUM, Hochst. in Flora, xxix. (1846) 594.
LUPINASTER, Buxb. ex Heist. Syst. 9 (1748).
MICRANTHEUM, Presl, Symb. Bot. i. 47 (1830).
MICROPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362.
MISTYLLUS, Presl, Symb. Bot. i. 49 (1830).
PARAMESUS, Presl, l. c. 45 (1830).
PENTAPHYLLON, Pers. Syn. ii. 352 (1807).
SERTULA, Linn. Syst. ed. I (1735).
TRIPHYLLOIDES, Ponted. ex Moench, Meth. 509 (1794).
abbreviatum, *J. Woods, Tour. Fl.* 92.—Europ.
acaule, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 169.—Abyss.
aciculare, Nutt. in Torr. & Gray, Fl. N. Am. i. 319 = tridentatum.
adscendens, Hornem. Hort. Hafn. ii. 716 = reflexum ?
affine, Lej. ex DC. Prod. ii. 195 = medium.
affine, Presl, Symb. Bot. i. 54 = arvense.
africanum, *Ser. in DC. Prod.* ii. 200.—Afr. austr.
agrarium, Huds. Fl. Angl. ed. I. 286 = procumbens.
agrarium, *Linn. Sp. Pl.* 772.—Europ. ; Afr. bor. ; Oriens.
agrarium, Mérat, Fl. Par. 202 = patens.
agrestinum, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 153 = arvense.
aintabense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 140.—Syria.
alatum, Bivona, Stirp. Rar. iv. 14 = physodes.
albens, Fisch. ex Loud. Hort. Brit. 300 = T. Lupinaster.
albiceps, *Ehrenb. ex Sweet, Hort. Brit.* ed. II. 137.—Aegypt.

TRIFOLIUM:—
albidum, *Retz. Obs.* iv. 30.—Hab.?
albidum, Stev. ex Ledeb. Fl. Ross. i. 543 = maritimum.
albiflorum, Fisch. ex Loud. Hort. Brit. 299 = eximium.
albopurpureum, Torr. & Gray, Fl. N. Am. i. 313 = Macraei.
album, Crantz, Stirp. Austr. ed. II. fasc. v. 408 = montanum, ochroleucum.
album, Lam. Fl. Fr. ii. 603 = hybridum, repens.
album, Loisel. Fl. Gall. ed. I. 479 (err. typ. 4) = Melilotus alba.
alexandrinum, Bory & Chaub. Nouv. Fl. Pélop. 50 = latinum.
alexandrinum, *Linn. Cent. Pl.* i. 25; *Amoen. Acad.* iv. 286.—Aegypt.; As. Min.
alopecuroides, *Kit. in Linnaea*, xxxii. (1863) 619 (an seq.?).—Hungar.
alopecuroides, Pers. Syn. ii. 350, in obs. = pannonicum.
alpestre, Crantz, Stirp. Austr. ed. II. fasc. v. 407 = medium.
alpestre, *Linn. Sp. Pl.* ed. II. 1082.—Europ. austr.; As. bor.
alpicolum, *Hegetschw. Fl. Schw.* 699.—Europ.
alpinum, Georgi, Beschr. Russ. Reich. III. iv. 1191 = alpestre.
alpinum, *Linn. Sp. Pl.* 767.—Europ.
alpinum, Pall. Reise, ii. 123 = eximium.
altissimum, *Dougl. ex Hook. Fl. Bor. Am.* i. 130.—Am. bor. occ.
altissimum, Loisel. Fl. Gall. ed. I. 479 (err. typ. 4) = Melilotus officinalis.
altissimum, Torr. & Gray, in Pacif. Rail. Rep. ii. 120 = Beckwithii.
amabile, Auct. ex Lojac. in Nuov. Giorn. Bot. Ital. xv. (1883) 143 = Hemsleyi.
amabile, *H. B. & K. Nov. Gen. et Sp.* vi. 503.—Mexic.
ambiguum, *Bieb. Fl. Taur. Cauc.* ii. 208.—Reg. Cauc.
amphianthum, *Torr. & Gray, Fl. N. Am.* i. 316.—Texas.
amplectens, *Torr. & Gray, l. c.* 319.—Am. bor. occ.
ampulescens, Gilib. Fl. Lituan. ii. 89 = fragiferum.
anatolicum, Boiss. Diagn. Ser. I. ii. 31 = hybridum.
Andersonii, *A. Gray, in Proc. Am. Acad.* vi. (1865) 522.—Am. bor. occ.
andinum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 314.—Am. bor. occ.
angulatum, J. Gay, ex Reichb. Fl. Germ. Excurs. 496 = Thalii.
angulatum, *Waldst. & Kit. Pl. Rar. Hung.* i. 26. t. 27.—Europ. austr.
angustifolium, *Linn. Sp. Pl.* 769.—Reg. Mediterr.; Oriens.
anomalum, Bory & Chaub. Nouv. Fl. Pélop. 51 = physodes.
anomalum, Schrank, Pl. Rar. Hort. Monac. t. 47 = repens.
appendiculatum, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 181.—Calif.
apulum, [Hort. ex] All. in Misc. Taur. v. 76 = spumosum.
arachnoideum, Presl, Delic. Prag. 49 = Cherleri.
arenivagum, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 153 = arvense.
argutum, [Soland. in] Russ. Aleppo, ed. II. ii. 260 = spumosum.
aristatum, *Hornem. Hort. Hafn.* ii. 712.—Hab.?
aristatum, Willd. Enum. Hort. Berol. Suppl. 52, nomen; Link, Enum. Hort. Berol. ii. 262 = ligusticum.
armeniacum, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 601 = Pennyanum.
armenium, Eichw. Beschr. Casp. Cauc. 28 = pannonicum.
armenium, *Willd. Enum. Hort. Berol.* 793.—As. Min.
arrectisetum, Brot. Phyt. Lusit. i. 152. t. 63 = ligusticum.
arvense, *Linn. Sp. Pl.* 769.—Europ.; As. et Afr. bor.; Oriens.
arvense, Walt. Fl. Carol. 183 = carolinianum.
arvernense, Lamotte, Prod. Fl. Centr. Fr. i. 202 = pallescens.
arvicolum, Jord. ex Nym. Consp. 175 = arvense.
asperulum, *De Not. ex Cesati, in Bibl. Ital.* xci. (1838) 347.—Ital.
atlanticum, *Ball, in Journ. Bot.* xi. (1873) 306.—Marocc.

TRIFOLIUM:—
atropurpureum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 318 = involucratum.
Aucheri, Boiss. Fl. Orient. ii. 114 = medium.
aurantiacum, *Boiss. & Sprun. Diagn.* Ser. I. ii. 33.—Graecia.
aureum, Geners. Elench. Scepus. n. 69; ex Wahlenb. Fl. Carpat. 229 = badium.
aureum, Pollich, Hist. Pl. Palat. ii. 344 = agrarium.
aureum, Thuill. Fl. Par. ed. II. 385 = patens.
badium, Ledeb. Fl. Ross. i. 556 = rytidosemium.
badium, *Schreb. in Sturm, Deutschl. Fl.* Heft 16.—Europ.
baeticum, *Boiss. Voy. Bot. Espagne*, ii. 726.—Hispan.
baeticum, Lag. ex Ser. ex DC. Prod. ii. 193 = maritimum.
Balansae, *Boiss. Diagn.* Ser. II. v. 81.—As. Min.
Balbisianum, *Ser. in DC. Prod.* ii. 201.—Gallia.
barbatum, DC. Cat. Hort. Monsp. (1813) 152 = pannonicum.
barbigerum, *Torr. in Pacif. Rail. Rep.* iv. 79.—Calif.
Beckwithii, *Brewer, ex S. Wats. in Proc. Am. Acad.* xi. (1876) 128.—Calif.
bejariense, *Moric. Pl. Nouv. Am.* 2.—Mexic.
benghalense, Gaertn. Fruct. ii. 335 (Hedysari sp.).—Ind. or.
berytheum, Boiss. & Blanche, Diagn. Ser. II. ii. 15 = supinum.
Biasolettianum, Steud. & Hochst. ex Koch, Syn. Fl. Germ. ed. I. 173 = repens.
biceps, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 417 = Bocconi.
bicolor, Moench, Meth. 111 = hybridum.
bicorne, Forsk. Fl. Aegypt. Arab. 139 = resupinatum.
bifidum, *A. Gray, in Proc. Calif. Acad.* iii. (1863) 102.—Calif.
biflorum, Linn. Sp. Pl. 773 = Stylosanthes elatior.
bifurcatum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 353 = montanum.
bilineatum, *Fresen. in Flora*, xxii. (1839) 52.—Abyss.
Billardieri, Spreng. Syst. iii. 211 = comosum.
bithynicum, Boiss. Diagn. Ser. I. ix. 21 = medium.
Bivonae, *Guss. Fl. Sic. Prod.* ii. 512.—Sicil.
Blancheanum, Boiss. Diagn. Ser. II. ii. 13 = Desvauxii.
Bocconi, *Savi, Obs. Trif.* 37.—Europ.; Afr. bor.
Boissieri, *Guss. Fl. Sic. Syn.* ii. 858 (Boisseri).—Graec.; As. Min.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 335.—Calif.
Bonanni, *Presl, Delic. Prag.* 51.—Sicil.
Boryanum, Steud. Nom. ed. II. ii. 705 = physodes.
Bourgaei, Boiss. ex Boiss. Fl. Orient. ii. 121 = smyrneum.
brachystylos, Knaf, in Lotos, iv. (1854) 237 = pratense.
bracteatum, *Schousb. ex Willd. Enum. Hort. Berol.* 792.—Reg. Mediterr. occ.
Brandegei, *S. Wats. in Proc. Am. Acad.* xi. (1876) 130.—Am. bor. occ.
brevipes, *Phil. in Linnaea*, xxviii. (1856) 680.—Chili.
Breweri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 131.—Calif.
Brittingeri, Weitenweb. ex Opiz, Natural. n. ix. (1825) 142 = arvense.
Broteri, Link, Enum. Hort. Berol. ii. 262 = ligusticum.
brutium, *Tenore, Viagg. Calabr.* 127; *Fl. Neap. Prod. App.* v. 24.—Calabria.
bullatum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 138.—Syria.
Burchellianum, *Ser. in DC. Prod.* ii. 200.—Afr. austr.
Buxbaumii, *Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb.* i. II. (1818) 131.—Europ. austr.
caerulescens, E. Desv. ex Boiss. Diagn. Ser. II. ii. 14 = Desvauxii.
caeruleum, Auct. = coeruleum.
caespitosum, Eichw. Skizze, 166 = elegans.
caespitosum, J. F. Gmel. Syst. 1141; Willd. Sp. Pl. iii. 1359 = Thalii.
caespitosum, Sturm, Deutschl. Fl. Heft 32 = pallescens.
calabricum, *Steud. Nom.* ed. II. ii. 705.—Calabria.
calocephalum, *Fresen. in Flora*, xxii. (1839) 50.—Abyss.
calocephalum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 318 = involucratum.
caloxanthum, Griseb. Bericht. (1847) 62 = Boissieri.
calycinum, *Steud. Nom.* ed. II. ii. 705.—Sicil.
campestre, Schreb. in Sturm, Deutschl. Fl. Heft 16 = procumbens.

TRIFOLIUM:—
canescens, Sims, Bot. Mag. t. 1168 = trichocephalum.
canescens, *Willd. Sp. Pl.* iii. 1369.—As. Min.
capense, *Willd. l. c.* 1364.—Afr. austr.
capillatum, *Dum. Fl. Belg. Prod.* 101.—Belgium.
capilliforme, Delile, ex Nym. Consp. 181 = filiforme.
Carmeli, *Boiss. Diagn.* Ser. II. ii. 16.—Syria.
carolinianum, *Michx. Fl. Bor. Am.* ii. 58.—Am. bor.
carpathicum, Janka, ex Nym. Consp. 173 = pratense.
cassium, *Boiss. Diagn.* Ser. I. ix. 23.—Syria.
catanense, *Buek, Ind. ad DC. Prod.* i. p. viii.—Sicil.
caucasicum, Tausch, in Syll. Ratisb. ii. (1828) 245 = pannonicum.
caudatum, *Boiss. Diagn.* Ser. I. ix. 22.—Lydia.
cernuum, *Brot. Phyt. Lusit.* i. 150.—Lusitan.
Cherleri, *Linn. Cent. Pl.* i. 24; *Amoen. Acad.* iv. 286.—Reg. Mediterr.
chiclense, *Ball, in Journ. Linn. Soc.* xxii. (1885) 35.—Peruv.
chilense, *Hook. & Arn. Bot. Beech. Voy.* 16.—Chili.
chlorotrichum, *Boiss. & Bal. Diagn.* Ser. II. vi. 48.—Phrygia.
chrysanthum, Gaud. Fl. Helv. iv. 603 = patens.
chrysopogon, *Viv. Pl. Aegypt. Arab. Dec.* 15.—Aegypt.
ciliatum, Clarke, Trav. ii. 261 = procumbens.
ciliatum, *Nutt. in Journ. Acad. Philad.* N. S. Ser. II. i. (1847) 152.—Am. bor. occ.
ciliolatum, Benth. Pl. Hartw. 304 = ciliatum.
ciliosum, Thuill. Fl. Par. ed. II. 380 = diffusum.
cinctum, *DC. Cat. Hort. Monsp.* (1813) 152.—Gallia.
cinerascens, Kit. in Linnaea, xxxii. (1863) 619 = ochroleucum.
circumdatum, Guss. ex Steud. Nom. ed. II. ii. 705 = calabricum.
circumdatum, *Kunze, in Linnaea*, xvi. (1842) 321.—Chili.
circumscissum, Pers. Syn. ii. 349 = diffusum.
clandestinum, *Lag. Nov. Gen. et Sp.* 23.—Hispan.
Clausonis, Pomel, Nouv. Mat. Fl. Atl. 181 = Durandoi.
Clusii, Gren. & Godr. Fl. Fr. i. 414 = resupinatum.
clypeatum, Lapeyr. Hist. Abr. Pl. Pyr. 436 = maritinum.
clypeatum, *Linn. Sp. Pl.* 769.—Graecia; As. Min.
coarctatum, *Reichb. Fl. Germ. Excurs.* 494.—Italia.
coccineum, *Steud. Nom.* ed. II. ii. 705.—Chili.
coerulescens, Bieb. Fl. Taur. Cauc. iii. 509 = Trigonella azurea.
coeruleum, Bieb. l. c. = Trigonella azurea.
coeruleum, *Viv. Pl. Aegypt. Arab. Dec.* 15.—Aegypt.
coeruleum, Willd. Sp. Pl. iii. 1352; Bieb. Fl. Taur. Cauc. ii. 207 = Trigonella coerulea.
coespitosum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Thalii.
collinum, Bast. Fl. Maine-et-Loire, Suppl. 5 = Bocconi.
commutatum, Ledeb. Fl. Ross. i. 543 = maritimum.
comosum, *Labill. Ic. Pl. Syr. Dec.* v. 15.—Syria.
comosum, Linn. Sp. Pl. 767 = carolinianum.
concinnum, Phil. Fl. Atac. 14 = polymorphum.
congestum, Guss. Cat. Pl. Boccad. (1821) 81, Adnot. 12 = catanense.
congestum, Link, in Linnaea, ix. (1834–35) 584 = fragiferum.
conicum, Kit. ex Nym. Consp. 176 = striatum.
conicum, Lag. Gen. et Sp. Nov. 23 = Lagascanum.
conicum, Pers. in Savi, Obs. Trif. 41 = tenuiflorum.
constantinopolitanum, Ser. in DC. Prod. ii. 193 = alexandrinum.
controversum, Jan, ex DC. Prod. ii. 206 = filiforme.
cordifolium, Walp. Rep. i. 640, sphalm. = Origanum cordifolium.
Crosnierii, Clos, in C. Gay, Fl. Chil. ii. 68 = polymorphum.
cryptopodium, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 168.—Abyss.
cryptoscias, Griseb. Spicil. Fl. Rumel. i. 30 = uniflorum.
Cupani, Tineo, Pl. Rar. Sicil. Pugill. i. 16 = physodes.
cuspidatum, *Lour. Fl. Cochinch.* 445.—Cochinch.
Cussoni, *Hort. Berol. ex Spreng. Nov. Prov.* 41.—Hab.?
cyathiferum, *Lindl. Bot. Reg.* sub t. 1070.—Am. bor. occ.
cylindricum, Wallr. Beitr. Fl. Hercyn. 249 = tenuiflorum.

TRIFOLIUM:—
cytisoides, Pall. Reise, iii. 266 = Lespedeza juncea.
dalmaticum, *Vis. in Flora*, xii. (1829) 1. *Erg.* 21.—Europ. austr.; As. Min.
dasyphyllum, *Torr. & Gray, Fl. N. Am.* i. 315.—Am. bor. occ.
dasyurum, Presl, Symb. Bot. i. 53 = formosum.
davuricum, Laxm. in Nov. Comm. Acad. Petrop. xv. (1771) 560. t. 30 = Lespedeza trichocarpa.
decipiens, *Hornem. Hort. Hafn.* ii. 719.—Hab.?
decumbens, F. Schultz, Arch. Fl. 272 = elegans.
densiflorum, *Phil. in Linnaea*, xxviii. (1856) 619.—Chili.
dentatum, Waldst. & Kit. Pl. Rar. Hung. i. 41 = Melilotus dentata.
denudatum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 152 = gracilentum.
depauperatum, *Desv. Journ. Bot.* iv. (1814) 69.—Am. bor. occ.; Peruv.; Chili.
Desvauxii, *Boiss. & Blanche, Diagn.* Ser. II. ii. 12.—Syria.
dichotomum, Hook. & Arn. Bot. Beech. Voy. 330 = Macraei.
dichroanthum, *Boiss. Diagn.* Ser. I. ix. 20.—Palaestin.
diffusum, *Ehrh. Beitr.* vii. (1792) 165.—Europ.; Tauria.
diffusum, Thunb. Prod. Pl. Cap. 136 = capense.
dimidiatum, Salisb. Prod. 333 = T. Lupinaster.
dipsaceum, Thuill. Fl. Par. ed. II. 382 = ochroleucum.
divaricatum, *Hornem. Hort. Hafn.* ii. 715.—Europ. austr.
diversifolium, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 152 = amplectens.
dubium, *Sibth. Fl. Oxon.* 231.—Europ.
duodecimnerve, Losc. ex Willk. & Lange, Prod. Fl. Hisp. iii. 357 = glomeratum.
Durandoi, *Pomel, Nouv. Mat. Fl. Atl.* 181.—Alger.
echinatum, Bieb. Fl. Taur. Cauc. ii. 216 = supinum.
echinatum, Wierzb. ex Nym. Consp. 175 = reclinatum.
Ehrenbergii, *Hort. Berol. ex Sweet, Hort. Brit.* ed. II. 137.—Aegypt.
elegans, *Savi, Fl. Pis.* ii. 161. t. 1. f. 2.—Reg. Mediterr.; As. Min.; Afr. bor.
elongatum, Willd. Sp. Pl. iii. 1368 = armenium.
Endressi, J. Gay, ex Gren. & Godr. Fl. Fr. i. 417 = montanum.
erectum, Poir. Encyc. viii. 28 = procumbens.
erectum, Walt. Fl. Carol. 184 = Rhynchosia tomentosa.
erinaceum, Bieb. Fl. Taur. Cauc. iii. 510 = phleoides.
eriocephalum, Ledeb. Fl. Ross. i. 541 = arvense.
eriocephalum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 313.—Am. bor. occ.
eriocephalum, Parry, in Am. Natural. ix. (1875) 269, partim = longipes.
eriosphoerum, *Boiss. Diagn.* Ser. I. ix. 25.—Palaestin.
erubescens, *Fenzl, Pugill. Fl. Nov. Syr.* 5.—Oriens.
erythraeum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 420 = isthmocarpum.
eximium, *Steph. ex Ser. in DC. Prod.* ii. 203.—Sibir. ural.
expansum, Waldst. & Kit. Pl. Rar. Hung. iii. 263 = pratense.
fabaceum, *Lindl. ex Steud. Nom.* ed. II. ii. 705 (Quid?).—Calif.
filicaule, *Boiss. & Heldr. Diagn.* Ser. I. ix. 24.—Graecia.
filiforme, *Linn. Sp. Pl.* 773, pro maj. parte.—Europ.; Reg. Cauc.
filiforme, Linn. l. c., partim = dubium.
fimbriatum, Lindl. in Bot. Reg. t. 1070 = involucratum.
fistulosum, Gilib. Fl. Lituan. ii. 86 = hybridum.
flavescens, Tineo, Pl. Rar. Sicil. Pugill. i. 15 = pallidum.
flavicans, Link, Handb. ii. 170 = squarrosum.
flavum, *Presl, Fl. Sic.* p. xxi.—Sicil.
Fleischeri, *Steud. & Hochst. ex Steud. Nom.* ed. II. ii. 705; *et ex Presl, Bot. Bemerk.* 52.—As. Min.
flexicaule, Vill. Prosp. 43 = medium.
flexuosum, Jacq. Fl. Austr. iv. 45. t. 386 = medium.
fluminense, Vell. Fl. Flum. 316; vii. t. 143 = Stylosanthes guianensis.
folliculatum, Lam. Fl. Fr. ii. 599 = spumosum.
formosum, Curt. ex DC. Prod. ii. 200 = resupinatum.
formosum, Savi, Obs. Trif. 102 = elegans.
formosum, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 350.—Graecia; Oriens.

TRIFOLIUM:—
fragiferum, *Linn. Sp. Pl.* 772.—Europ.; Oriens; Reg. Himal.
frigidum, Schur, Enum. Pl. Transs. 154 = pratense.
fruticans, Linn. Sp. Pl. 770 = Psoralea bracteata.
fucatum, *Lindl. Bot. Reg.* t. 1883.—Am. bor. occ.
fulcratum, Griseb. Spicil. Fl. Rumel. i. 26 = Pignantii.
fuscum, Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 330 = agrarium.
galilaeum, Boiss. Diagn. Ser. I. ix. 26 = nervulosum.
Gambelii, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 151 = fucatum.
gemellum, Lapeyr. Hist. Abr. Pl. Pyr. 437 = Bocconei.
gemellum, *Pourr. ex Willd. Sp. Pl.* iii. 1376.—Hispan.
gemellum, Savi, in Atti Acc. Ital. i. 202; ex Bertol. Fl. Ital. viii. 152 = ligusticum.
glabellum, *Presl, Fl. Sic.* p. xxi.—Europ. austr.
glabrum, J. Gay, ex Nym. Consp. 175 = maritimum.
glaciale, Porc. in Magyar Növényt. Lapok, ix. (1885) 126 = pallescens.
glaciale, Reyner, ex J. F. Gmel. Syst. 1141; DC. Prod. ii. 197 = saxatile.
glanduliferum, *Boiss. Diagn.* Ser. I. ii. 30.—As. Min.
glareosum, Dum. Fl. Belg. 100 = pratense.
glareosum, Schleich. Cat. Pl. Helv. ed. IV. 35; Boiss. Voy. Bot. Espagne, Suppl. 727 = pallescens.
glaucum, Haussk. ex Nym. Consp. 180 = Lagrangei.
globosum, *Linn. Sp. Pl.* 767.—Byzant.; As. Min.
globosum, Ser. in DC. Prod. ii. 196 = radiosum.
glomeratum, *Linn. Sp. Pl.* 770.—Europ. occ.; Reg. Mediterr.; Oriens.
goniocarpum, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 145.—Mexic.
gracile, Thuill. Fl. Par. ed. II. 383 = arvense.
gracilentum, *Torr. & Gray, Fl. N. Am.* i. 316.—Am. bor. occ.
graecum, Griseb. Bericht. (1847) 62 = speciosum.
grandiflorum, Hook. & Arn. Bot. Beech. Voy. 16 = polymorphum.
grandiflorum, Ledeb. Fl. Alt. iii. 257 = eximium.
grandiflorum, *Schreb. in Nov. Act. Nat. Cur.* iii. (1767) 477.—Ins. Creta.
Grayi, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 189.—Calif.
guianense, Aubl. Pl. Gui. ii. 776 = Stylosanthes guianensis.
Gussoni, Tineo, Pl. Rar. Sicil. Pugill. i. 17 = speciosum.
gymnocarpum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 320.—Am. bor. occ.
hamosum, Bieb. Fl. Taur. Cauc. ii. 207 = Trigonella spicata.
Haussknechtii, *Boiss. Fl. Orient.* ii. 125.—Mesopot.
Haydenii, Porter, in Hayd. Rep. (1871) 480 = Kingii.
Haynaldi, *Meynh. Kalocsa Vidék. Növényt.* 65; *et ex Knapp, in Oestr. Bot. Zeitschr.* xxviii. (1878) 103.—Europ.
Haynaldianum, *Pantoč. in Oestr. Bot. Zeitschr.* xxviii. (1878) 382.—Europ.
hedysaroides, Pall. Reise, iii. 751 = Lespedeza juncea.
Hemsleyi, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 143.—Mexic.
heterodon, Torr. & Gray, Fl. N. Am. i. 318 = involucratum.
heterophyllum, Tratt. Arch. Gew. i. 26; Lej. Rev. Fl. Spa, 158 = pratense.
hirsutum, E. Mey. Comm. Pl. Afr. Austr. 91 = africanum.
hirsutum, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
hirtum, *All. Auct. Fl. Pedem.* 20.—Reg. Mediterr.; As. Min.
hispidissimum, Michx. ex Steud. Nom. ed. II. ii. 706 = Stylosanthes elatior.
hispidum, Desf. Fl. Atlant. ii. 200 = hirtum.
Hohenackeri, Jaub. & Spach, Illustr. ii. 55, in nota = canescens.
Humboldtianum, *A. Br. & Aschers. in Ind. Sem. Hort. Berol.* (1868) 24.—Reg. Cauc.
Humboldtii, Spreng. Syst. iii. 213 = amabile.
humile, *Ball, in Journ. Bot.* xi. (1873) 306.—Marocc.
hybridum, *Linn. Sp. Pl.* 766.—Europ. austr.; As. Min.
hybridum, Pollich, Hist. Pl. Palat. ii. 330 = elegans.
hybridum, Savi, Fl. Pis. ii. 90 = nigrescens.
hygrophilum, Boiss. Diagn. Ser. II. ii. 18 = Petrisavii.
incanum, Presl, Delic. Prag. 51 = Cherleri.
incarnatum, *Linn. Sp. Pl.* 769.—Europ. austr.
incarnatum, Steph. Enum. Stirp. Mosq. 38. n. 500 = alpestre.

TRIFOLIUM:—
indecorum, Clos, in C. Gay, Fl. Chil. ii. 67 = polymorphum.
indicum, Thunb. Prod. Pl. Cap. 136 = Melilotus indica.
intermedium, *Guss. Cat. Pl. Boccad.* (1821) 82. *Adnot.* 14.—Reg. Mediterr.
intermedium, Lapeyr. Suppl. Hist. Pl. Pyr. 115 = hybridum.
intertextum, Wender. ex Steud. Nom. ed. II. ii. 706 = Medicago intertexta?
involucratum, Dulac, Fl. Hautes-Pyr. 283 = fragiferum.
involucratum, Lam. Fl. Fr. ii. 604 = Cherleri.
involucratum, *Ortega, Hort. Matr. Dec.* 33; *Willd. Sp. Pl.* iii. 1372.—Am. bor. occ.
involucratum, Sm. in Rees, Cycl. xxxvi. n. 28 = tridentatum.
irregulare, Pourr. in Mém. Acad. Toul. iii. (1788) 331 = maritimum.
isthmocarpum, *Brot. Phyt. Lusit.* i. 148. t. 61.—Reg. Mediterr. occ.
italicum, Boiss. Fl. Orient. ii. 107, 1153 (Index) = Melilotus italica.
jaminianum, Boiss. & Reut. Diagn. Ser. II. ii. 19 = isthmocarpum.
Kingii, *S. Wats. in Bot. King's Rep.* v. 59.—Calif.
Kitaibelianum, Ser. in DC. Prod. ii. 194 = striatum.
Kochianum, Hayne, in Schrad. Neues Journ. ii. (1807) 337 = Melilotus dentata.
laevigatum, Poir. Voy. Barb. ii. 219 = strictum.
Lagascanum, *Ser. in DC. Prod.* ii. 194.—Hab.?
lagopinum, Jord. Pugill. Pl. Nov. 57 = arvense.
lagopinum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = T. Lagopus.
Lagopus, Gouan, Fl. Monsp. 105 = arvense.
Lagopus, *Pourr. ex Willd. Sp. Pl.* iii. 1365.—Hispan.; Pyren.
Lagrangei, *Boiss. Fl. Orient.* ii. 154.—Archip.; Graecia.
laguroides, Pourr. ex Link, Enum. Hort. Berol. ii. 261 = hirtum.
lanatum, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
lappaceum, *Linn. Sp. Pl.* 768.—Reg. Mediterr.; Oriens.
lasiocephalum, Link, Enum. Hort. Berol. ii. 262 = hirsutum.
latinum, *Sebast. Pl. Rom.* fasc. i. 7. t. 1. f. 2.—Europ. austr. occ.; Syria.
laxiusculum, Boiss. & Blanche, Diagn. Ser. II. ii. 13 = Desvauxii.
leiocalycinum, *Boiss. & Sprun. l. c.* Ser. I. ii. 31.—Graecia.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xi. (1876) 127.—Calif.
leptocephalum, *Wallr. in Linnaea*, xiv. (1840) 620 (*Beitr. Fl. Hercyn.* 250).—Europ. centr.
leucanthum, *Bieb. Fl. Taur. Cauc.* ii. 214.—Europ. austr.; As. Min.
leucophaeum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 418 = bracteatum.
libanoticum, Ehrenb. ex Sweet, Hort. Brit. ed. II. 137 = globosum.
ligusticum, *Balb. in Mem. Acc. Torin.* xxiii. (1818) 112, nomen; *et ex Savi, Obs. Trifol.* 38.—Reg. Mediterr.
limonium, Phil. in Linnaea, xxviii. (1856) 679 = repens.
litigiosum, Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 329 = spadiceum.
littorale, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 153 = arvense.
longepedunculatum, Steud. Nom. ed. II. ii. 706 = squarrosum.
longestipulatum, Ebel, Monteneg. 84. t. 4. f. 2 = patulum.
longestipulatum, Loisel. Fl. Gall. ed. II. ii. 122 = squarrosum.
longipedunculatum, Loisel. ex Bull. Sc. Anger (1828), 425 (err. typ.) = praec.
longipes, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 314.—Am. bor. occ.
longisetum, Boiss. & Bal. Diagn. Ser. II. vi. 47 = arvense.
lucanicum, Gasp. in Guss. Fl. Sic. Prod. ii. 494 = dalmaticum.
lucidum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 419 = strictum.

TRIFOLIUM:—
ludovicianum, Rafin. Fl. Ludov. 102 (Quid?).—Am. bor.
Lupinaster, [Habl.] Desc. Taur. 187; ex Bieb. Fl. Taur. Cauc. ii. 221 = Dorycnium suffruticosum.
Lupinaster, *Linn. Sp. Pl.* 766.—Sibir.
Lupinaster, Sibth. & Sm. Fl. Graec. Prod. ii. 95 = patulum.
lupinastrum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = T. Lupinaster.
lupulinum, Gueldenst. ex Ledeb. Fl. Ross. i. 557 = procumbens.
lupulinum, Savi, Fl. Pis. ii. 171 = Medicago lupulina.
luteolum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 179 = dubium.
luteum, Lam. Fl. Fr. ii. 604 = filiforme, procumbens.
luxurians, Hort. Par. ex Steud. Nom. ed. II. ii. 706 = repens.
Macraei, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 179.—Am. bor. occ.; Chili.
macrocalyx, Hook. Ic. Pl. t. 285 = bejariense.
macrocephalum, Poir. Encyc. Suppl. v. 336 = megacephalum.
macrocephalum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 421 = fragiferum.
macropodum, Guss. Fl. Sic. Syn. ii. 338 = Michelianum.
macropodum, *Steud. Nom.* ed. II. ii. 706.—Sicil.
macrorrhizum, Boiss. Diagn. Ser. II. v. 80 = repens.
macrorrhizum, Waldst. & Kit. Pl. Rar. Hung. i. 24 = Melilotus officinalis.
maculatum, *Host, Fl. Austr.* ii. 373.—Sicil.
magellanicum, Poir. Encyc. viii. 25 = Oxalis eriocarpa.
magnum, *Gueldenst. Reisen*, i. 426.—Transcauc.
malacanthum, Link, Enum. Hort. Berol. ii. 261 = leucanthum.
maritimum, *Huds. Fl. Angl.* ed. I. 284.—Europ. occ.; Reg. Mediterr.
marsicum, *Tenore, Syll.* 620; *ex Walp. Rep.* ii. 850.—Europ.
Matthewsii, *A. Gray, Bot. U. St. Expl. Exped.* i. 398.—Peruv.
mauritanicum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 420 = isthmocarpum.
mauritanicum, Willd. Sp. Pl. iii. 1354 = Melilotus sulcata.
mauritianum, Steud. Nom. ed. I. 849; ed. II. ii. 706 = Melilotus mauritanica.
mediterraneum, *Vell. Fl. Flum.* 317; vii. t. 144.—Bras.
medium, Griseb. Spicil. Fl. Rumel. i. 25 = patulum.
medium, *Linn. Faun. Suec.* ed. II. 558, nomen; *Huds. Fl. Angl.* ed. I. 284.—Europ.; As. bor.
meduseum, *Blanche, ex Boiss. Fl. Orient.* ii. 134.—Syria.
megacephalum, *Nutt. Gen. Am.* ii. 105.—Am. bor. occ.
megalanthum, Steud. Nom. ed. II. ii. 706 = polymorphum.
melananthum, Hook. & Arn. Bot. Beech. Voy. 331 = tridentatum.
Melilotus, Georgi, Reise, i. 227 = Melilotus alba.
Melilotus, Gueldenst. ex Ledeb. Fl. Ross. i. 538 = Melilotus officinalis.
Melilotus, Linn. Sp. Pl. ed. I. 764 = Melilotus indica.
Melilotus-caerulea, Linn. l. c. = Trigonella caerulea.
Melilotus-corniculata, Linn. l. c. 766 = Melilotus italica.
Melilotus-cretica, Linn. l. c. 765 = Trigonella cretica.
Melilotus-indica, Linn. l. c. = Melilotus mauritanica.
Melilotus-italica, Brot. Fl. Lusit. ii. 102 = Melilotus mauritanica?
Melilotus-italica, Linn. Sp. Pl. 765 = Melilotus italica.
Melilotus-mauritanica, Schousb. Vextr. Marokko, 196 = Melilotus mauritanica.
Melilotus-officinalis, Linn. Sp. Pl. 765 = Melilotus officinalis.
Melilotus-officinarum, Crantz, Inst. ii. 41 = Melilotus officinalis.
Melilotus-ornithopodioides, Linn. Sp. Pl. 766 = Trigonella ornithopodioides.
Melilotus-polonica, Linn. l. c. 765 = Melilotus polonica.
Melilotus-segetalis, Brot. Fl. Lusit. ii. 484 = Melilotus sulcata?
Melilotus-tauricum, Bieb. Fl. Taur. iii. 506 = Melilotus gracilis.
Meneghinianum, *Clem. in Mem. Acc. Sc. Torin.* Ser. II. xvi. (1857) 267.—Byzant.; Syria.
mesogitanum, Boiss. Diagn. Ser. I. ii. 34 = brutium.

TRIFOLIUM:—
messanense, Hill, Veg. Syst. xxii. 35 = seq.?
messanense, Linn. Mant. ii. 275 = Melilotus messanensis.
mexicanum, *Hemsl. Biol. Centr. Am. Bot.* i. 233.—Mexic.
Michelianum, Gaud. Fl. Helv. iv. 573 = hybridum.
Michelianum, Koch, Syn. Fl. Germ. ed. I. 174 = Meneghinianum.
Michelianum, *Savi, Fl. Pis.* ii. 159.—Europ. austr.
micranthum, Viv. Fl. Lib. Spec. 45. t. 19. f. 3 = filiforme.
microcephalum, *Pursh, Fl. Am. Sept.* ii. 478.—Am. bor. occ.
microdon, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 180.—Am. bor. occ.; Chili.
micropetalum, E. Mey. Comm. Pl. Afr. Austr. i. 90 = stipulaceum.
microphyllum, Desv. Journ. Bot. ii. (1809) 316 = pratense.
minimum, Barton, Comp. Fl. Philad. ii. 74 = procumbens.
minus, Relhan, Fl. Cantab. ed. II. 290 = dubium.
minutum, Coss. Notes Crit. 5 = cernuum.
modestum, *Boiss. Diagn.* Ser. I. ix. 27.—Syria.
Molinerii, Balb. Cat. Hort. Taur. (1813) App. i. 17 = incarnatum.
molybdocalyx, Reichb. f. Ic. Fl. Germ. xxii. 77. t. 2167 = Petrisavii.
monanthum, *A. Gray, in Proc. Am. Acad.* vi. (1865) 523.—Calif.
montanum, *Linn. Sp. Pl.* 770.—Europ. austr.; As. bor.
montanum, Linn. l. c. 772 = spadiceum.
moriferum, Boiss. Diagn. Ser. I. ix. 28 = xerocephalum.
mucronatum, *Willd. Sp. Pl.* iii. 1372.—Mexic.
multicaule, *M. Jones, in Bull. Torrey Bot. Club*, ix. (1882) 31.—Calif.
multinerve, *A. Rich. Tent. Fl. Abyss.* i. 175.—Abyss.
multistriatum, *Koch, Syn. Fl. Germ.* ed. II. 190.—Dalmat.; Graecia.
mutabile, *Portenschl. Enum. Pl. Dalm.* 16.—Europ. austr.
Mutelii, *Gren. in Acad. Sc. Besanç. Séances Publ.* (1838) 127.—Alger.
nanum, *Torr. in Ann. Lyc. N. York*, i. (1824) 35. t. 3. f. 4.—Am. bor. occ.
neglectum, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 21 = fragiferum.
neglectum, Noé, ex Nym. Consp. 178 = repens.
neolagopus, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 194.—Calif.
nervosum, *Presl, Fl. Sic.* p. xxi.—Europ. austr.
nervulosum, *Boiss. & Heldr. Diagn.* Ser. I. ix. 25.—As. Min.; Syria.
nevadense, *Boiss. Diagn.* Ser. II. ii. 17.—Hispan.
nidificum, Griseb. Spicil. Fl. Rumel. i. 32 = radiosum.
nigrescens, *Viv. Fragm. Fl. Ital.* 12. t. 13.—Europ. austr.
nigrocinctum, Boiss. & Orph. Diagn. Ser. II. vi. 47 = maritimum.
nigrolineatum, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 42; *et in Oken, Isis*, xxi. (1828) 992.—Thracia.
nigropunctatum, G. Don, Gen. Syst. ii. 181 = nigrolineatum.
nivale, Arv. Ess. Pl. Dauph. 23 (Quid?).—Gallia.
nivale, Sieber, ex Koch, Syn. Fl. Germ. ed. I. 168 = pratense.
Noëanum, Reichb. ex Koch, in Roehl. Deutschl. Fl. ed. III. v. 265 = incarnatum.
noricum, *Wulf. in Roem. Arch.* iii. (1805) 387.—Europ. austr.
nothum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 137 = repens.
Nuttallii, Steud. Nom. ed. II. ii. 707 = tridentatum.
obcordatum, Desv. Journ. Bot. iii. (1814) 76 = polymorphum.
oblongifolium, *Ser. in DC. Prod.* ii. 197.—Hispan.
obscurum, Savi, Obs. Trif. 31. f. 1 = leucanthum.
obtusiflorum, Hook. Ic. Pl. t. 281 = tridentatum.
obvallatum, Moench, Meth. 112 = Cherleri.
ochreatum, *Kunze, ex Steud. Nom.* ed. II. ii. 707.—Chili.
ochroleucum, *Huds. Fl. Angl.* ed. I. 283.—Europ.; Oriens.
ochroleucum, Rostk. & Schmidt, Fl. Seden. 302 = scabrum.
ochroleucum-magnum, Gueldenst. ex Ledeb. Fl. Ross. i. 545 = canescens.
odoratum, Schrank, Baier. Fl. ii. 286 = montanum?

TRIFOLIUM:—
officinale, Willd. Sp. Pl. iii. 1355 = Melilotus officinalis.
oliganthum, Steud. Nom. ed. II. ii. 707 = pauciflorum.
Olivierianum. Ser. in DC. Prod. ii. 197 = globosum.
olympicum, Hornem. ex Hook. Bot. Mag. t. 2790 = armenium.
ornatum, *Clarke, Trav.* iii. 336.—Ins. Cos.
ornithopodioides, Sm. Fl. Brit. iii. 782 = Trigonella ornithopodioides.
Orphanideum, Boiss. Diagn. Ser. II. ii. 17 = repens.
Ottonis, *Sprun. ex Boiss. Diagn.* Ser. I. ii. 28.—Graecia.
ovatifolium, Bory & Chaub. Nouv. Fl. Pélop. 51 = physodes.
oxaloides, Bunge, ex Nym. Consp. 177 = subterraneum.
oxypetalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 51 = carolinianum.
oxypetalum, Nym. Consp. 1036 = hirtum.
oxypetasum, Heldr. & Sart. ex Nym. l. c. 174 = hirtum.
palaestinum, *Boiss. Diagn.* Ser. I. ix. 21.—Palestin.
palavicum, Hochst. in Lorent. Wanderungen, 346 = purpureum.
pallescens, DC. Fl. Fr. Suppl. 555 = nigrescens.
pallescens, *Schreb. in Sturm, Deutschl. Fl.* Heft 15.—Europ. austr.
pallidulum, Jord. Pugill. Pl. Nov. 56 = ochroleucum.
pallidum, Bory & Chaub. Nouv. Fl. Pélop. 50. t. 28 = Pignantii.
pallidum, *Waldst. & Kit. Pl. Rar. Hung.* i. 35. t. 36.—Reg. Mediterr.
pallidum, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 418 = bracteatum.
Palmeri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 132.—Calif.
palustre, Waldst. & Kit. Pl. Rar. Hung. iii. 295 = Melilotus mauritanica?
pamphylicum, *Boiss. & Heldr. Diagn.* Ser. I. ix. 20.—As. Min.
pannonicum, *Jacq. Obs. Bot.* ii. 21.—Europ.; Oriens.
pannonicum, Sibth. & Sm. Fl. Graec. Prod. ii. 98 = armenium.
panormitanum, *Presl, Fl. Sic.* p. xxi.—Reg. Mediterr. et Casp.
paradoxum, *Steud. Nom.* ed. II. ii. 707.—Chili.
parisiense, DC. Fl. Fr. Suppl. 562 = patens.
Parnassi, *Boiss. & Sprun. Diagn.* Ser. I. ii. 30.—Graecia.
Parnassi, Orph. ex Nym. Consp. 178 = repens.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 409.—Am. bor. occ.
parviflorum, *Ehrh. Beitr.* vii. (1792) 165.—Reg. Mediterr.; As. Min.
parviflorum, Perreym. Pl. Frej. 84 = Perreymondi.
parvifolium, Bunge, ex Nym. Consp. 180 = dubium.
patens, *Schreb. in Sturm, Deutschl. Fl.* Heft 16.—Europ. austr.
patulum, *Tausch, in Syll. Ratisb.* ii. (1828) 245.—Dalmat.; Graecia.
pauciflorum, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 319.—Am. bor. occ.
pauciflorum, Urv. in Mém. Soc. Linn. Par. i. (1822) 350 = globosum.
pauciflorum, Willd. ex Steud. Nom. ed. II. ii. 707 = amabile.
pectinatum, Willd. ex Steud. l. c. = arvense.
Pennyanum, *Steud. l. c.*—Armenia.
pensylvanicum, Willd. Enum. Hort. Berol. 793 = medium.
Perreymondi, Coss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 356 = cernuum.
Perreymondi, *Gren. & Godr. Fl. Fr.* i. 422.—Hispan.
peruvianum, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* I. (1843) 12.—Peruv.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 168.—Abyss.
Petitpierreanum, Hayne, in Schrad. Neues Journ. ii. (1807) 337 = Melilotus mauritanica.
Petrisavii, *Clem. in Mem. Acc. Sc. Torin.* Ser. II. xvi. (1857) 268.—Byzant.; As. Min.
phlebocalyx, Fenzl, ex Tchihat. Asie Min. Bot. i. 29 = Cherleri.
phleoides, *Pourr. ex Willd. Sp. Pl.* iii. 1377.—Reg. Mediterr.
physanthum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 180.—Chili.
physodes, Eichw. Skizze, 166 = fragiferum.
physodes, *Stev. ex Bieb. Fl. Taur. Cauc.* ii. 217.—Europ. austr.; Oriens.

TRIFOLIUM:—
physopetalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 47 = fucatum.
pictum, Roth, Catalect. fasc. ii. 101 = hirtum.
Pignantii, *Fauché & Chaub. in Expéd. Morée Bot.* 219.—Graecia; Dalmat.
Pignati, Steud. Nom. ed. II. ii. 707, sphalm. = praec.
piliferum, Straub. ex Nym. Consp. 176 = formosum.
pilulare, *Boiss. Diagn.* Ser. I. ii. 29.—As. Min.; Syria.
platycephalum, Bisch. Del. Sem. Hort. Heidelb. (1839). —Cf. Linnaea, xiv. (1840) Litt. 132 = reflexum.
plebeium, *Boiss. Diagn.* Ser. I. ix. 23.—Syria.
plicatum, Presl, ex Sweet, Hort. Brit. ed. II. 137 = speciosum.
Plummeri, *Lemmon, ex Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 162.—Calif.
plumosum, *Dougl. ex Hook. Fl. Bor. Am.* i. 130.—Am. bor. occ.
polyanthemum, Tenore, ex DC. Prod. ii. 200 = hybridum.
polycephalum, *Ser. in DC. l. c.* 196.—Afr. austr.
polymorphum, *Poir. Encyc.* viii. 20.—Reg. Magellan.
polyphyllum, *C. A. Mey. Verz. Pfl. Cauc.* 139.—Reg. Cauc.
polyphyllum, Nutt. in Torr. & Gray, Fl. N. Am. 319 = tridentatum.
polystachyum, *Fresen. in Flora*, xxii. (1839) 50.—Abyss.
potosanum, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 144.—Mexic.
praecox, *Wallr. in Linnaea*, xiv. (1840) 621 (*Beitr. Fl. Hercyn.* 251).—Europ.
praetutianum, *Guss. Pl. Rar. Sic.* 308.—Italia.
pratense, *Linn. Sp. Pl.* 768.—Europ.; As. temp.
Preslianum, Boiss. Diagn. Ser. I. ii. 25 = arvense.
procerum, Rochel, Pl. Banat. Rar. 50 = reclinatum.
procumbens, Linn. Sp. Pl. 727, partim = dubium.
procumbens, *Linn. l. c.*—Europ.; Afr. et As. bor.
procumbens, Pollich, Hist. Pl. Palat. ii. 345 = filiforme.
prostratum, Biasol. ex Koch, Syn. Fl. Germ. ed. I. 173 = repens.
prostratum, *Burm. f. Fl. Cap. Prod.* 22.—Afr. austr.
prostratum, Griseb. Spicil. Fl. Rumel. i. 29 = Petrisavii.
prutetianum, Ser. in DC. Prod. ii. 202, sphalm. = praetutianum.
Pseudo-procumbens, C. C. Gmel. Fl. Bad. iii. 240 = procumbens.
psoralioides, Walt. Fl. Carol. 184 = Psoralea melilotoides.
purpurascens, *Fisch. ex Loud. Hort. Brit.* 300; *G. Don, Gen. Syst.* ii. 192.—Sibir.
purpurascens, Roth, Catalect. Bot. fasc. iii. 74 = diffusum.
purpureum, Gilib. Fl. Lituan. ii. 86 = pratense.
purpureum, Gueldenst. ex Ledeb. Fl. Ross. i. 546 = alpestre.
purpureum, *Loisel. Fl. Gall.* ed. I. 484. t. 14.—Europ.; Oriens; As. bor.
purpureum-majus, Gilib. Fl. Lituan. ii. 87 = alpestris.
pyrenaicum, *Pourr. in Mém. Acad. Toul.* iii. (1788) 331.—Mont. Pyren.
Quartinianum, *A. Rich. Tent. Fl. Abyss.* i. 169.—Abyss.
radiatum, *Delile, Fl. Aeg. Illustr.* 70, nomen.—Aegypt.
radicosum, *Boiss. & Hohen. Diagn.* Ser. I. ix. 27.—Persia.
radiosum, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 43; *et in Oken, Isis*, xxi. (1828) 992.—Thracia.
reclinatum, Griseb. Spicil. Fl. Rumel. i. 21 = leucanthum.
reclinatum, *Waldst. & Kit. Pl. Rar. Hung.* iii. 299. t. 269.—Hungar.
recurvum, Waldst. & Kit. l. c. ii. 179. t. 165 = vesiculosum.
reflexum, *Linn. Sp. Pl.* 766.—Am. bor.
reflexum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 356 = cernuum.
reflexum, Roem. ex Scheele, in Linnaea, xxi. (1848) 460 = amphianthum.
reflexum, Schlecht. & Cham. l. c. v. (1830) 576 = Schiedeanum.
reflexum, Ser. in DC. Prod. ii. 197 = reclinatum.
repens, *Linn. Sp. Pl.* 767.—Geront. bor. temp.
repens, Walt. Fl. Carol. 183 = carolinianum?

TRIFOLIUM:—
resupinatum, Less. in Linnaea, ix. (1834) 175 = fragiferum.
resupinatum, *Linn. Sp. Pl.* 771.—Europ. occ.; Reg. Mediterr.
resupinatum, Willk. in Flora, xxxiv. (1851) 628 = Bonani.
rhombeum, *S. Schau. in Linnaea*, xx. (1847) 740.—Mexic.
rigidum, Savi, Fl. Pis. i. t. 1. f. 1; ii. 154 = maritimum.
rigidum, Schrad. ex Steud. Nom. ed. I. 850 = villosum, *Seenus*.
rivale, Clos, in C. Gay, Fl. Chil. ii. 70 = polymorphum.
rivale, Steud. Nom. ed. II. ii. 708, in syn., sphalm. = *nivale*, Sieber.
rivulare, *Boiss. & Bal. Diagn.* Ser. II. vi. 49.—Cappadoc.
Roemerianum, Scheele, in Linnaea, xxi. (1848) 460 = amphianthum.
roseum, Presl, Delic. Prag. 50 = ochroleucum.
rotundatum, *Ser. in DC. Prod.* ii. 200.—Hab.?
rotundifolium, Bory & Chaub. Nouv. Fl. Pélop. 50 = filicaule.
rotundifolium, *Sibth. & Sm. Fl. Graec.* viii. 35. t. 747.—Gracia.
Roussaeanum, Boiss. Diagn. Ser. I. ii. 26 = Desvauxii.
Rouxii, Gren. in Mém. Soc. Emul. Doubs, Sér. III. ii. (1858) 413 = isthmocarpum.
rubellum, Jord. Pugill. Pl. Nov. (1852) 57 = arvense.
rubens, Aubry, Fl. Morb. 56; ex DC. Fl. Fr. Suppl. 556 = incarnatum.
rubens, [Habl.] Desc. Taur. 24; ex Bieb. Fl. Taur. Cauc. ii. 211 = alpestre.
rubens, *Linn. Sp. Pl.* 768.—Europ. austr.
rubicundum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 420 = isthmocarpum.
Rueppellianum, *Fresen. in Flora*, xxii. (1839) 51.—Abyss.
rupestre, *Tenore, ex Ser. in DC. Prod.* ii. 201.—Italia.
rupestre, Tenore, Prod. Fl. Nap. p. xliii = montanum.
rytidosemium, *Boiss. & Hohen. Diagn.* Ser. I. ix. 29.—Reg. Cauc.; Persia.
sabuletorum, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 153 = arvense.
sarosiense, *Hazsl. ex Neilr. Diagn. Gefaesspfl.* 35.—Hungar.
sativum, Crome, ex Boenn. Prod. Fl. Monast. 222 = pratense.
Savianum, Guss. Fl. Sic. Prod. ii. 488 = uniflorum.
saxatile, *All. Fl. Pedem.* i. 305. t. 59. f. 3.—Europ.
scabrum, *Linn. Sp. Pl.* 770.—Reg. Mediterr.; Oriens.
scabrum, Pollich, Hist. Pl. Palat. ii. 338 = striatum.
Schiedeanum, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 339.—Mexic.
Schimperi, *A. Rich. Tent. Fl. Abyss.* i. 173.—Abyss.
Schreberi, Jord. ex Reut. Cat. Pl. Genèv. ed. II. 49 = procumbens.
sclerorrhizon, Boiss. Diagn. Ser. I. ix. 28 = physodes.
scutatum, *Boiss. Diagn.* Ser. I. ii. 27.—As. Min.; Syria.
Sebastiani, *Savi, in Diar. Med. Flajani*, (1815); *ex Sebast. Rom. Pl.* fasc. ii. 14.—Ital.; Reg. Casp.
semiglabrum, Brot. Phyt. Lusit. i. 155 = Bocconi.
semipilosum, *Fresen. in Flora*, xxii. (1839) 52.—Abyss.
serrulatum, Lag. Gen. et Sp. Nov. 23 = cernuum.
setiferum, *Boiss. Diagn.* Ser. I. ii. 32.—As. Min.
sibiricum, Gueldenst. Reisen, ii. 327; ex Ledeb. Fl. Ross. i. 558 (Quid?).—Sibir.
siculum, *Presl, ex Steud. Nom.* ed. II. ii. 708.—Sicil.
simense, *Fresen. in Flora*, xxii. (1839) 49.—Abyss.
simplex, Clos, in C. Gay, Fl. Chil. ii. 69 = polymorphum.
simplicifolium, Walt. Fl. Carol. 184 = Rhynchosia tomentosa.
smyrnaeum, *Boiss. Diagn.* Ser. I. ii. 25.—As. Min.; Mesopot.
spadiceum, *Linn. Fl. Suec.* ed. II. 261.—Europ.; As. Min.
spadiceum, Thuill. Fl. Par. ed. II. 385 = procumbens.
spadiceum, Vell. Fl. Delph. 80; Hist. Pl. Dauph. iii. 491 = badium.
speciosum, Boiss. Diagn. Ser. I. ii. 33 = Boissieri.
speciosum, Bory & Chaub. Nouv. Fl. Pélop. 51 = aurantiacum.

TRIFOLIUM:—
speciosum, Marg. & Reut. Fl. Zante, 41 = patens.
speciosum, Ser. in DC. Prod. ii. 205 = comosum.
speciosum, *Willd. Sp. Pl.* iii. 1382.—Sicil.; Graec.; Oriens.
sphaerocephalum, Desf. Fl. Atlant. ii. 201 = Cherleri.
spicatum, Sibth. & Sm. Fl. Graec. viii. 32. t. 743 = Melilotus gracilis.
spinulosum, Dougl. ex Hook. Fl. Bor. Am. i. 133 = involucratum.
spumosum, *Linn. Sp. Pl.* 771.—Reg. Mediterr.
squarrosum, *Bieb. Fl. Taur. Cauc.* ii. 214.—Reg. Cauc.
squarrosum, DC. Fl. Fr. iv. 531 = panormitanum.
squarrosum, *Linn. Sp. Pl.* 768.—Europ. austr.
squarrosum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 418 = maritimum.
stellatum, Huds. Fl. Angl. ed. II. 326 = maritimum.
stellatum, *Linn. Sp. Pl.* 769.—Reg. Mediterr.; Oriens.
stellatum, Pall. Tabl. Taur. 56 = leucanthum.
stenophyllum, *Boiss. Diagn.* Ser. I. ix. 30.—Palaestin.
stenophyllum, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 151 = depauperatum.
Steudneri, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 652.—Afr. trop.
stipitatum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 149.—As. Min.
stipitatum, Clos, in C. Gay, Fl. Chil. ii. 71 = repens.
stipulaceum, Lapeyr. Suppl. Hist. Abr. Pl. Pyr. 116 = maritimum.
stipulaceum, *Thunb. Prod. Pl. Cap.* 136.—Afr. austr.
stoloniferum, *Muhl. Cat.* 70.—Am. bor.
stramineum, Guss. ex Tenore, Syll. Fl. Neap. 372 = tenuifolium.
stramineum, Presl, Fl. Sic. p. xx = incarnatum.
strepens, Crantz, Stirp. Austr. ed. II. fasc. v. 411 = agrarium.
striatum, *Linn. Sp. Pl.* 770.—Europ. austr.; As. Min.
strictum, *Linn. Cent. Pl.* i. 24; *Amoen. Acad.* iv. 285.—Europ.; Reg. Mediterr.
strictum, Schreb. in Sturm, Fl. Deutschl. Heft 15 = parviflorum.
suaveolens, Willd. Enum. Hort. Berol. i. 108; Suppl. 52 = resupinatum.
subcaulescens, A. Gray, in Ives, Rep. Colorado River, Bot. 10 = gymnocarpon.
subcaulescens, S. Wats. Bot. King's Exped. 59, partim = longipes.
subexpansum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 177 = medium.
subrotundum, *Steud. & Hochst. in Flora*, xxiv. (1841) I. *Intell.* 32; *et ex A. Rich. Tent. Fl. Abyss.* i. 172.—Abyss.
subterraneum, *Linn. Sp. Pl.* 767.—Reg. Mediterr.; Oriens.
subulatum, Gilib. Fl. Lituan. ii. 90 = montanum.
succinctum, Vis. in Flora, xii. (1829) I. Erg. 21 = cinctum.
suffocatum, *Linn. Mant.* ii. 276.—Europ. occ.; Reg. Mediterr. et Cauc.
sulphureum, C. Koch, in Linnaea, xix. (1847) 63 = armenium.
supinum, *Savi, Obs. Trif.* 46. f. 2.—Europ.; Oriens.
sylvaticum, *Gérard, ex Loisel. in Desv. Journ. Bot.* ii. (1809) 367.—Gall. austr.
tauricum, Bieb. Fl. Taur. Cauc. iii. 506 = Melilotus gracilis.
tembense, *Fresen. in Flora*, xxii. (1839) 51.—Abyss.
Tenoreanum, Boiss. & Sprun. Diagn. Ser. I. ii. 26 = tenuifolium.
Tenoreanum, Buek, Ind. ad DC. Prod. i. p. viii = montanum.
tenue, *Clem. in Atti Acc. Riun. Sc. Ital.* (1841) 521.—Bras.
tenue, *Willd. Enum. Hort. Berol. Suppl.* 52, nomen.—Hab.?
tenuiflorum, *Tenore, Prod. Fl. Nap.* p. xliv.—Europ. austr.
tenuifolium, *Tenore, l. c.*—Italia.
Thalii, *Vill. Prosp.* 43; *Hist. Pl. Dauph.* i. 289; iii. 478.—Mont. Pyren.
thymiflorum, Vill. ll. cc. 43; iii. 487 (*thimiflorum*) = saxatile.
tmoleum, Boiss. ex Boiss. Fl. Orient. ii. 141 = glanduliferum.
tomentosum, *Linn. Sp. Pl.* 771.—Reg. Mediterr.
tomentosum, Willk. ex Nym. Consp. 177 = Bonani.
transsilvanicum, Schur, Enum. Pl. Transs. 154, 157 = pratense.

TRIFOLIUM:—
triaristatum, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 54. t. 3.—Chili.
trichocephalum, *Bieb. Fl. Taur. Cauc.* ii. 212.—Reg. Cauc.
trichocephalum, Bieb. l. c., partim = canescens.
trichopterum, *Panč. in Verh. Zool.-Bot. Ges. Wien*, vi. (1856) 480.—Banat.
trichostomum, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 427.—Hab.?
tricuspidatum, *Bert. ex Steud. Nom.* ed. II. ii. 708.—Chili.
tridentatum, Hort. Berol. ex Steud. l. c. = Wormskioldii.
tridentatum, *Lindl. Bot. Reg.* sub t. 1070.—Am. bor. occ.
triste, Nutt. ex Torr. & Gray, Fl. N. Am. i. 318 = involucratum.
tumens, *Stev. ex Bieb. Fl. Taur. Cauc.* ii. 217.—Reg. Cauc.; Persia.
tumescens, Gilib. Fl. Lituan. ii. 91 = agrarium.
turgidum, Bieb. Fl. Taur. Cauc. ii. 216 = vesiculosum.
umbellatum, *A. Rich. Tent. Fl. Abyss.* i. 172.—Abyss.
umbellatum, Losc. ex Willk. & Lange, Prod. Fl. Hisp. iii. 355 = repens.
umbellatum, Ser. in DC. Prod. ii. 199 = carolinianum.
umbellulatum, *A. Rich. Tent. Fl. Abyss.* i. 172.—Abyss.
uniflorum, *Linn. Sp. Pl.* 771.—Europ. austr.
unifolium, Forsk. Fl. Aegypt. Arab. 140 = Psoralea corylifolia.
vaginatum, Schleich. Cat. Pl. Helv. 51; ex Ser. in DC. Prod. ii. 195 = pratense.
Vaillantii, Bieb. ex Fisch. Cat. Hort. Gorenk. (1808) 111, nomen = ambiguum.
Vaillantii, Loisel. in Desv. Journ. Bot. ii. (1809) 365 = Michelianum.
Vaillantii, Poir. Encyc. viii. 4 = elegans.
Vaillantii, Tenore, Prod. Fl. Nap. p. xliv = hybridum *vel* nigrescens.
variabile, Guss. ex DC. Prod. ii. 197 = leucanthum.
variegatum, Nutt. in Torr. & Gray, Fl. N. Am. i. 317 = pauciflorum.
ventricosum, *Presl, ex Steud. Nom.* ed. II. ii. 708.—Sicil.
venulosum, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 86 = repens.
vernum, *Phil. in Anal. Univ. Chil.* (1872) 687.—Chili.
versicolor, *Steud. Nom.* ed. II. ii. 708.—Chili.
vesiculosum, *Savi, Fl. Pis.* ii. 165.—Europ. austr.; Reg. Cauc.
villosum, Gilib. Fl. Lituan. ii. 89 = arvense.
villosum, *Presl, Delic. Prag.* 48.—Sicil.
villosum, *Seenus, ex Steud. Nom.* ed. I. 850.—Europ.
villosum, Tenore, Syll. Fl. Neap. 373 = pratense.
volubile, *Lour. Fl. Cochinch.* 445.—'Afr. or.'
vulgare, Gueldenst. ex Ledeb. Fl. Ross. i. 537 = Melilotus alba.
vulgare, Hayne, in Schrad. Neues Journ. ii. (1807) 336 = Melilotus officinalis.
Waldsteinianum, Sweet, Hort. Brit. ed. II. 136 = reclinatum.
Watsonii, *Lojac. in Nuov. Giorn. Bot. Ital.* xv. (1883) 186.—Calif.
Willdenovii, Spreng. Syst. iii. 208 = involucratum.
Wormskioldii, Lehm. Sem. Hort. Hamb. (1825) 17.—Cf. Linnaea, iii. (1828) Litt. 10 = involucratum.
xanthinum, *Freyn, in Bot. Centralb.* i. (1880) 308.—Graec.
Xatardii, DC. Fl. Fr. Suppl. 558 = maritimum.
xerocephalum, *Fenzl, Pugill. Pl. Nov. Syr.* 5.—As. Min.; Syria.

TRIFURCARIA, Endl. Gen. 1359 (1841) = seq.

TRIFURCIA, Herb. in Bot. Mag. sub t. 3779 (1840) = **Herbertia**, Sweet (Irid.).
caerulea, Herb. l. c. = H. caerulea.

TRIGASTROTHECA, F. Muell. in Hook. Kew Journ. ix. (1857) 16 = **Mollugo**, Linn. (Ficoid.).
molluginea, F. Muell. l. c. = M. Trigastrotheca.

TRIGELLA, Salisb. Gen. Pl. Fragm. 46 (1866) = **Cyanella**, Linn. (Haemod.).
orchidiformis, Salisb. l. c. 47 = C. orchidiformis.

TRIGLOCHIN, Riv. ex Linn. Syst. ed. I (1735).
NAIADACEAE, Benth. & Hook. f. iii. 1012.
ABBOTIA, Rafin. New Fl. Am. i. 36 (1836).
CYCNOGETON, Endl. in Ann. Wien. Mus. ii. (1838) 210.
JUNCAGO, Tourn. ex Moench, Meth. 644 (1794).
MAUNDIA, F. Muell. Fragm. i. (1858) 23.
andinum, *Phil. in Anal. Univ. Chil.* (1873) 538.—Chili.
Ani, C. Koch, in Linnaea, xxii. (1849) 273 = maritimum.
atacamensis, Phil. Fl. Atac. 49 = striatum.
atlanticum, Willd. ex Kunth, Enum. Pl. iii. 143 = bulbosum.
Barrelieri, Loisel. Fl. Gall. ed. I. 725; ed. II. i. 264 = bulbosum.
bulbosum, *Linn. Mant.* ii. 226.—Reg. Mediterr.; Afr. trop. et austr.
calcaratum, Hook. Ic. Pl. t. 416 = mucronatum.
Calcitrapa, Hook. l. c. t. 731 = centrocarpum.
centrocarpum, *Hook. l. c.* t. 728.—Austral.
chilense, Meyen, Reise, i. 354 = palustre.
ciliatum, *Ruiz & Pav. Fl. Per.* iii. 72 (sp. dub.).—Peruv.
decipiens, R. Br. Prod. 343 = striatum.
densiflorum, Domb. ex Kunth, Enum. Pl. iii. 144 = striatum.
dubium, R. Br. Prod. 343 = procerum.
elatum, Nutt. Gen. Am. i. 237 = maritimum.
filifolium, Sieber, ex Spreng. Syst. iv. Cur. Post. 142 = striatum.
flaccidum, A. Cunn. in Hook. Comp. Bot. Mag. ii. (1836) 378 = striatum.
fonticola, Phil. Fl. Atac. 49 = palustre.
himalensis, Royle, Illustr. Bot. Himal. 402 = palustre?
junceum, Gilib. Exercit. ii. 501 = palustre.
laxiflorum, *Guss. Ind. Sem. Hort. Boccad.* (1825); *Fl. Sic. Prod.* i. 451.—Sicil.; Corsic.
Lechleri, Steud. in Lechl. Berb. Am. Austr. 53 = striatum.
lineare, Endl. in Lehm. Pl. Preiss. ii. 54 = procerum.
litoreum, *Phil. in Anal. Univ. Chil.* (1873) 539.—Chili.
littorale, Phil. ex Micheli, in DC. Monog. Phan. iii. 102 = striatum.
magellanicum, Vahl, ex Kunth, Enum. Pl. iii. 142 = Tetroncium magellanicum.
maritimum, *Linn. Sp. Pl.* 105.—Reg. bor. temp.
maritimum, Thunb. Fl. Cap. ed. Schult. 340 = striatum.
Maundii, *F. Muell. Fragm.* vi. 83, in obs.—Austral.
mexicanum, H. B. & K. Nov. Gen. et Sp. i. 244 = maritimum.
minutissimum, F. Muell. Fragm. vi. 82, in obs. = centrocarpum.
montevidense, Spreng. Syst. ii. 145 = striatum.
mucronatum, *R. Br. Prod.* 343.—Austral.
nanum, F. Muell. in Trans. Vict. Inst. i. (1855) 135 = centrocarpum.
Neesii, Endl. in Lehm. Pl. Preiss. ii. 54 = mucronatum.
palustre, Desf. Fl. Atlant. i. 232 = laxiflorum.
palustre, *Linn. Sp. Pl.* 338.—Europ.; As.; Am.
palustre, Ruching. Fl. Venet. 99 = maritimum.
palustre, Schousb. Vextr. Marokko, 184 = bulbosum.
patens, Steud. Nom. ed. II. ii. 709 = bulbosum.
procerum, *R. Br. Prod.* 343.—Austral.
racemosum, Endl. in Lehm. Pl. Preiss. ii. 54 = Bulbine semibarbatum.
reflexum, Vahl, ex Kunth, Enum. Pl. iii. 142 = Tetroncium magellanicum.
Roegneri, C. Koch, in Linnaea, xxii. (1849) 272 = maritimum.
salina, Wallr. in Linnaea, xiv. (1840) 567 (Beitr. Fl. Hercyn. 197) = maritimum.
sessile, Hill, Veg. Syst. x. 35 (Quid?).—Hab.?
sexloculare, Stokes, Bot. Mat. Med. ii. 324 = maritimum.
striatum, *Ruiz & Pav. Fl. Per.* iii. 72.—Am. bor. et austr.; Afr. austr.; Austral.
triandrum, Michx. Fl. Bor. Am. i. 208 = striatum.
trichophorum, Nees, ex Endl. in Lehm. Pl. Preiss. ii. 54 = centrocarpum.
Volkmanni, *Phil. in Anal. Univ. Chil.* (1873) 539.—Chili.

TRIGLOSSUM, Fisch. Hort. Gorenk. ed. II. p. vi (1812) = **Arundinaria**, Michx. (Gramin.).
bambusinum, Fisch. l. c. p. viii = A. macrosperma.

TRIGONACHRAS, Radlk. in Sitzb. Math.-Phys. Akad. Muench, viii. (1878) 299, 304; ix. (1879) 500, 672 = **Sapindus**, Tourn.
acuta, Radlk. l. c. ix. (1879) 672 = Cupania acuta.
cultrata, Radlk. l. c. viii. (1878) 304; ix. (1879) 672 = S. cultratus.

TRIGONEA, Parl. in Giorn. l'Occhio, i. (1839) 161; ex ej. Giorn. Bot. Ital. ii. (1847) 120 = **Allium**, Tourn. (Liliac.).
sicula, Parl. ll. cc. 161; 121 = A. Dioscoridis.

TRIGONELLA, Linn. Gen. ed. I. 351 (1737).
LEGUMINOSAE, Benth. & Hook. f. i. 486.
APORANTHUS, Bromf. Fl. Vect. 117 (1856).
BOTRYOLOTUS, Jaub. & Spach, Illustr. i. 124. t. 63 (1842).
BUCERAS, Hall. ex All. Fl. Pedem. i. 313 (1785).
FALCATULA, Brot. Phyt. Lusit. 160. t. 65 (1801).
FOENUM-GRAECUM, [Tourn.] Rupp. Fl. Jen. ed. Hall. 263 (1745).
FOLLICULLIGERA, Pasq. Cat. Ort. Bot. Nap. 46 (1867).
GRAMMOCARPUS, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 18.
KENTIA, Adans. Fam. ii. 508 (1763).
MELISITUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 383.
NEPHROMEDIA, Kostel. Ind. Hort. Prag. 92 (1844).
POCOCKIA, Ser. in DC. Prod. ii. 185 (1825).
TELIS, Linn. Syst. ed. I (1735).
TRIFOLIASTRUM, Moench, Meth. 123 (1794).
aegyptiaca, Steud. Nom. ed. II. ii. 709 = stellata.
aleppica, *Boiss. & Haussk. in Boiss. Fl. Orient.* ii. 79.—Syria.
americana, Nutt. Gen. Am. ii. 120 = Hosackia Purshiana.
anatolica, Boiss. & Bal. Diagn. Ser. II. vi. 46 = polycerata.
anguina, *Delile, Fl. Egypte*, 254. t. 110. f. 2.—Afr. bor.; Mesopot.
arabica, *Delile, in Laborde, Voy. Arab. Pétr.* 84.—Arab.; Palaest.
arcuata, Boiss. & Blanch. Diagn. Ser. II. ii. 11 = berythea.
arcuata, *C. A. Mey. Verz. Pfl. Cauc.* 136.—Armen.; Persia.
arguta, Vis. Pl. Aegypt. 33 = occulta *vel* laciniata.
armata, Boiss. Fl. Orient. ii. 70 = berythea.
armata, Thunb. Prod. Pl. Cap. 137 = Melolobium collinum.
Aschersoniana, *Urb. in Verh. Bot. Ver. Brandenb.* xxiii. (1882) 67.—Aegypt.
astroites, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 40.—As. Min.; Syria.
Aucheri, Boiss. Diagn. Ser. I. ii. 20 = brachycarpa.
aurantiaca, *Boiss. l. c.* 22.—As. Min.
avium, Pritz. Ic. Ind. ii. 277, sphalm. = T. Pes-avium.
azurea, *C. A. Mey. Verz. Pfl. Cauc.* 136.—Graecia; As. Min.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. v. 79.—Graecia; As. Min.
berythea, *Boiss. & Blanche, l. c.* ii. 10.—Syria.
Besseriana, *Ser. in DC. Prod.* ii. 181.—Europ. or.; As. Min.
biflora, Griseb. Spicil. Fl. Rumel. i. 41 = lunata.
brachycarpa, *Moris, in Mem. Acc. Torin.* xxxvi. (1833) 190.—Oriens.
brahuica, Boiss. Fl. Orient. ii. 73 = polycerata.
cachemiriana, Cambess. in Jacquem. Voy. Bot. 36 = Emodi.
caelesyriaca, *Boiss. Diagn.* Ser. I. ix. 19.—Syria.
caerulea, *Ser. in DC. Prod.* ii. 181.—Europ. or.; Reg. Cauc.
caespitosa, Steud. ex A. Rich. Tent. Fl. Abyss. i. 168 = Trifolium Petitianum.
calliceras, *Fisch. ex Bieb. Fl. Taur. Cauc.* iii. 515.—Reg. Cauc.
cancellata, Desf. Tabl. ed. I. 190, nomen = striata.
capitata, *Boiss. Diagn.* Ser. I. ii. 17.—As. Min.
cariensis, *Boiss. l. c.* 21.—As. Min.
Cassia, *Boiss. l. c.* ix. 13.—Syria.
Cephalotes, *Boiss. & Bal. l. c.* Ser. II. v. 77.—Cilicia.
coerulea, Auct. = caerulea.
connata, Meyer, ex Steud. Nom. ed. II. ii. 709 = caerulea.
corniculata, *Linn. Syst.* ed. X. 1180.—Europ. austr.; As. Min.

TRIGONELLA:—
corniculata, Sibth. & Sm. Fl. Graec. viii. 44. t. 761 = Balansae.
crassifolia, *Hornem. Hort. Hafn.* ii. 723.—Aegypt.
crassipes, *Boiss. Diagn.* Ser. I. ii. 23.—As. Min.; Syria.
cretica, *Boiss. Fl. Orient.* ii. 91.—As. Min.
cylindracea, *Desv. Journ. Bot.* iii. (1814) 77.—Syria.
Desvauxii, Boiss. Diagn. Ser. II. vi. 147 (Index), sphalm. = Trifolium Desvauxii.
dicarpa, C. A. Mey. ex Boiss. Fl. Orient. ii. 77 = Noëana.
divaricata, [Clairv.] Man. Herb. 132 = monspeliaca.
dura, Vis. Pl. Aegypt. 32 = maritima.
egyptiaca, Poir. Encyc. viii. 95 = stellata.
elatior, Sibth. & Sm. Fl. Graec. Prod. ii. 108 = corniculata.
elliptica, *Boiss. Diagn.* Ser. I. ii. 20.—Persia.
elongata, Hort. Dorp. ex Link, Enum. Hort. Berol. ii. 267 = stellata.
Emodi, *Benth. in Royle, Illustr. Bot. Himal.* 197.—Reg. Himal.
ensifera, *Trautv. Del. Sem. Hort. Kiov.* (1840) 6.—*Cf. Linnaea*, xv. (1841) *Litt.* 98.—Hab.?
esculenta, Willd. Enum. Hort. Berol. 799 = corniculata.
falcata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 509.—Ins. Socotra.
fasciculata, Bertol. Misc. Bot. ii. 19 = azurea.
filiformis, Fisch. ex Ledeb. Fl. Ross. i. 532 = striata.
filipes, *Boiss. Diagn.* Ser. I. ix. 16.—Syria; Mesopot.
fimbriata, Royle, Illustr. Bot. Himal. 197 = corniculata.
Fischeriana. *Ser. in DC. Prod.* ii. 183.—As. Min.; Reg. Cauc.
flexuosa, Delile, ex Poir. Encyc. Suppl. v. 362 = anguina.
flexuosa, Fisch. ex Bieb. Fl. Taur. Cauc. iii. 515 = Fischeriana.
Foenum-graecum, [Habl.] Desc. Taur. 138 = gladiata.
Foenum-graecum, *Linn. Sp. Pl.* 777.—Europ.; Oriens.
Foenum-graecum, Sibth. & Sm. Fl. Graec. viii. 48. t. 766 = cariensis.
Foenum-graecum, Suter, Fl. Helv. ed. Hegetschw. ii. 149 = monspeliaca.
Gaillardoti, Boiss. Diagn. Ser. II. v. 78 = caelesyriaca.
geminiflora, Boiss. Fl. Orient. ii. 77 = Noëana.
geminiflora, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 219 = polycerata.
glabra, Thunb. Prod. Pl. Cap. 137 = hamosa.
gladiata, *Stev. in Fisch. Cat. Hort. Gorenk.* (1808) 112.—Reg. Mediterr.
glomerata, Hort. Par. ex Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 340 = brachycarpa.
Goalia, Buch.-Ham. ex Wall. Cat. n. 5920 C = Cyamopsis psoraloides.
gracilis, *Benth. in Royle, Illustr. Bot. Himal.* 197.—Ind. or.
graeca, *Boiss. Fl. Orient.* ii. 91.—Graecia; As. Min.
graeca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = T. Foenum-graecum.
grandiflora, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 218.—Turkestan.
Griffithii, *Boiss. Fl. Orient.* ii. 88.—Afghan.
halophila, *Boiss. Diagn.* Ser. II. v. 78.—Cilicia.
hamigera, *Clarke, Trav.* ii. 260.—Graecia.
hamosa, Bess. in Flora, xv. (1832) II. Beibl. 32, in nota = spicata.
hamosa, Delile, ex Sm. in Rees, Cycl. xxxvi. n. 7 = anguina.
hamosa, *Linn. Syst.* ed. X. 1180.—Aegypt.
hamosa, Sibth. & Sm. Fl. Graec. viii. t. 764 = Sibthorpii.
hierosolymitana, *Boiss. Diagn.* Ser. I. ix. 15.—As. Min.; Syria.
himalaica, Wall. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 88 = Emodi.
hirsuta, Thunb. Prod. Pl. Cap. 137 = Melolobium cernuum.
hybrida, Pourr. in Mém. Acad. Toul. iii. (1788) 331 = Medicago Pourretii.
incisa, Hornem. ex Fisch. & Mey. Ind. Sem. Hort. Pétrop. iv. 48 = anguina.
incisa, Royle, Illustr. Bot. Himal. 197 = polycerata.
indica, Linn. Sp. Pl. 778 = Rothia trifoliata.
isthmocarpa, *Boiss. & Bal. Diagn.* Ser. II. vi. 44.—Cappadoc.
Kotschyi, Boiss. Fl. Orient. ii. 87 = rytidocarpa.

TRIGONELLA:—
Kotschyi, *Fenzl, ex Boiss. Diagn.* Ser. I. ix. 15.—As. Min.
laciniata, Desf. Tabl. ed. I. 190 = polycerata.
laciniata, *Linn. Sp. Pl.* ed. II. 1095.—Aegypt.
lilacina, *Boiss. Diagn.* Ser. I. ix. 17.—Palaestin.
littoralis, Guss. Adnot. Cat. Pl. Boccad. (1821) 15 = maritima.
longirostris, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 133 = cariensis.
lunata, *Boiss. Diagn.* Ser. I. ii. 19.—As. Min.
macroglochin, *Dur. in Bull. Soc. Bot. Fr.* vi. (1859) 593.—Hab.?
macrorrhyncha, *Boiss. Diagn.* Ser. I. ii. 21.—Cilicia.
marginata, Hochst. & Steud. in Flora, xxiv. (1841) I. Intell. 32; et ex Steud. Nom. ed. II. ii. 709 = Trifolium subrotundum.
maritima, *Delile, in Poir. Encyc. Suppl.* v. 361.—Reg. Mediterr. or.
maritima, Tommas. ex Nym. Consp. 171 = corniculata.
media, Delile, Fl. Aegypte Illustr. 71 = polycerata.
media, Lag. ex Loud. Hort. Brit. 290; Steud. Nom. ed. II. ii. 709 (Quid?).—Hab.?
Melilotus, Link, Handb. ii. 165 = Melilotus indica.
mexicana, *G. Don, Gen. Syst.* ii. 469.—Mexic.
micrantha, *Desf. Cat. Hort. Par.* ed. III. 410.—Hab.?
microcarpa, Fresen. in Mus. Senckenb. i. (1833) 186 = stellata.
mniocarpa, Wallr. ex DC. Prod. ii. 172 = Medicago lupulina.
monantha, C. A. Mey. Verz. Pfl. Cauc. 137 = polycerata.
monspeliaca, *Linn. Sp. Pl.* 777.—Reg. Mediterr.
multiflora, *Humnicki, Cat. Pl. Luxeuil*, 18.—Gallia.
multinervis, Hochst. in Flora, xxiv. (1841) I. Intell. 32 = Trifolium multinerve.
nervosa, Klotzsch, in Bot. Ergeb. Waldem. Reise, 158 = hamosa.
nilotica, Presl, Bot. Bemerk. 52 = laciniata.
Noëana, *Boiss. Diagn.* Ser. II. ii. 11.—As. Min.; Persia.
obcordata, Wall. Cat. n. 5986 = hamosa.
occulta, *Delile, Fl. Aegypte Illustr.* 71.—Aegypt.
ornithopodioides, *DC. Fl. Fr.* iv. 550.—Europ.
ornithorynchos, Fisch. ex Hornem. Suppl. Hort. Hafn. 85 = calliceras.
orthoceras, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 399 = polycerata.
ovalis, Boiss. Elench. 36 = Medicago ovalis.
oxyrhyncha, Fisch. ex Ser. in DC. Prod. ii. 182 = calliceras.
parviflora, *Sm. in Rees, Cycl.* xxxvi. n. 16.—Helvet.
Pecten, Schenk, Pl. Sp. Aegypt. 34 = arabica.
persica, *Boiss. Diagn.* Ser. I. ii. 22.—Persia.
Pes-avium, *Bertol. Fl. Ital.* viii. 247.—Italia.
petiolaris, Viv. Fl. Lib. Sp. 49. t. 18. f. 5 = maritima.
phrygia, Boiss. & Bal. Diagn. Ser. II. vi. 45 = aurantiaca.
pinnatifida, Cav. Ic. i. 26. t. 38 = polycerata.
plagioneura, Boiss. Diagn. Ser. I. ix. 16 = cylindracea.
platycarpos, Linn. Sp. Pl. 776 = Medicago platycarpa.
plicata, *Boiss. Fl. Orient.* ii. 90.—As. Min.
polyantha, Desv. Journ. Bot. iii. (1814) 77 = stellata.
polycarpa, *Boiss. & Heldr. Diagn.* Ser. I. ix. 18.—Pamphylia.
polycerata, *Linn. Sp. Pl.* 777.—Reg. Mediterr.; Oriens.
polyceratoides, Lange, in Kjoeb. Vidensk. Meddel. (1865) 172 = polycerata.
procumbens, Reichb. Fl. Germ. Excurs. 500 = Besseriana.
prostrata, DC. Fl. Fr. Suppl. 571 = gladiata.
pubescens, *Edgew. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 88.—Ind. or.
purpurascens, Lam. Fl. Fr. ii. 590 = ornithopodioides.
radiata, *Boiss. Fl. Orient.* ii. 90.—As. Min.; Persia.
raphanina, *Boiss. l. c.* 71.—Cilicia.
retrorsa, *Boiss. l. c.* 78.—Afghan.
Retzii, Spreng. Syst. iii. 283 = arcuata.
rhytidocarpa, Boiss. & Bal. Diagn. Ser. II. vi. 44 = Emodi.
rigida, *Boiss. & Bal. l. c.* v. 79.—Cilicia.
rostrata, *Boiss. Fl. Orient.* ii. 89.—As. Min.
ruthenica, Linn. Sp. Pl. 776 = Medicago ruthenica.
ruthenica, Patrin, ex Ledeb. Fl. Alt. iii. 253 = Medicago platycarpa.

TRIGONELLA:—
rytidocarpa, *Boiss. Diagn.* Ser. I. ix. 12, in syn.—Cappadoc.
Schimperi, Hochst. in Flora, xxiv. (1841) I. Intell. 32 = Trifolium Schimperi.
sericea, Eaton & Wright, N. Am. Bot. ed. VIII. 459 = Hosackia Purshiana.
Sibthorpii, *Boiss. Diagn.* Ser. I. ix. 14.—As. Min.; Syria.
sinuata, Boiss. l. c. 18 = astroites.
smyrnea, *Boiss. l. c.* 19.—As. Min.
spicata, *Sibth. & Sm. Fl. Graec. Prod.* ii. 108.—Graecia; As. Min.
spinosa, *Linn. Sp. Pl.* 777.—Ins. Creta; Syria.
Spruneriana, *Boiss. Diagn.* Ser. I. ii. 17.—Graecia; As. Min.
stellata, *Forsk. Fl. Aegypt. Arab.* 140.—Afr. bor.; Arab.; Ins. Canar.
strangulata, *Boiss. Diagn.* Ser. I. ix. 17.—Syria.
striata, *Linn. f. Suppl.* 340.—As. Min.; Turkestan.
striata, Viv. Pl. Aeg. Dec. iv. 15 = hamosa.
suavissima, *Lindl. in Mitch. Three Exped.* i. 255.—Austral.
subinermis, Thunb. ex Harv. & Sond. Fl. Cap. ii. 79 = Melolobium collinum.
subracemosa, *Boiss. Fl. Orient.* ii. 73.—As. Min.
tenuior, Thunb. ex Harv. & Sond. Fl. Cap. ii. 79 = Melolobium collinum.
tenuis, Fisch. ex Bieb. Fl. Taur. Cauc. iii. 514 = striata.
tenuissima, Brongn. & Guill. in Ann. Sc. Nat. Sér. II. xv. (1841) 59 = suavissima.
tetrapetala, Roxb. ex Wight & Arn. Prod. 197 = Cyamopsis psoraloides.
tomentosa, Thunb. Prod. Pl. Cap. 137 = Psoralea obtusifolia.
torta, Sm. in Rees, Cycl. xxxvi. n. 7 = anguina.
torulosa, Griseb. Spicil. Fl. Rumel. i. 40 = Spruneriana.
tuberculata, Presl, Delic. Prag. 46 = gladiata.
uncata, *Boiss. & Noë, Diagn.* Ser. II. ii. 12.—Persia; Afghan.
uncinata, Ser. in DC. Prod. ii. 181 = spicata.
uncinata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 261.—As. Min.
uniflora, Munby, in Bull. Soc. Bot. Fr. xi. (1864) 45 = ornithopodioides.
velutina, *Boiss. Diagn.* Ser. I. ii. 18.—As. Min.; Syria.
villosa, Thunb. Prod. Pl. Cap. 137 = Melolobium collinum.

TRIGONIA, Aubl. Pl. Gui. i. 387. tt. 149, 150 (1775).
VOCHYSIACEAE.
HOEFFNAGELIA, Neck. Elem. iii. 68 (1790).
MAINEA, Vell. Fl. Flum. vii. t. 8 (1827).
boliviana, *Warm. in Mart. Fl. Bras.* xiii. II. 134.—Bolivia.
candida, *Warm. l. c.* 139.—Bras.
Cepo, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 115 = mollis.
crotonoides, *Cambess. l. c.*—Bras.
fasciculata, *Griseb. in Linnaea*, xxii. (1849) 29.—Bras.
floribunda, *Oerst. in Kjoeb. Vidensk. Meddel.* (1856) 38.—Nicaragua.
Glazioviana, *Warm. in Mart. Fl. Bras.* xiii. II. 129.—Bras.
hypoleuca, *Griseb. in Linnaea*, xxii. (1849) 30.—Guian.
laevis, *Aubl. Pl. Gui.* i. 390. t. 150.—Guian.
macrantha, *Warm. in Mart. Fl. Bras.* xiii. II. 129.—Peruv.
macrocarpa, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 373.—Guian.
macrostachya, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1258, nomen = villosa.
micrantha, Mart. in Flora, xx. (1837) II. Beibl. 102 = crotonoides.
microcarpa, *Sagot, ex Warm. in Mart. Fl. Bras.* xiii. II. 131.—Guian.
mollis, *DC. Prod.* i. 571.—Bras.
najadum, *Warm. in Mart. Fl. Bras.* xiii. II. 126.—Bras.
nivea, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 113.—Bras.
paniculata, *Warm. in Mart. Fl. Bras.* xiii. II. 132.—Bras.
parviflora, *Benth. in Hook. Kew Journ.* iii. (1851) 163.—Bras.; Peruv.

TRIGONIA:—
parviflora, Schott, in Spreng. Syst. iv. App. 409 = mollis.
pubescens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 114.—Bras.
rigida, *Oerst. in Kjoeb. Vidensk. Meddel.* (1856) 39.—Nicaragua.
rugosa, *Benth. Bot. Voy. Sulph.* 74.—N. Granat.
Schottiana, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 560 = mollis.
Schottiana, *Warm. in Mart. Fl. Bras.* xiii. ii. 133.—Bras.
sericea, *H. B. & K. Nov. Gen. et Sp.* v. 141.—N. Granat.
simplex, *Warm. in Mart. Fl. Bras.* xiii. ii. 125.—Bras.
Spruceana, *Benth. ex Warm. l. c.* 130.—Bras.
subcymosa, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 373.—Guian.
tomentosa, Warm. in Mart. Fl. Bras. xiii. ii. 142 = *Trigonis tomentosa*.
velutina, Pohl, ex A. W. Benn. l. c. xiii. iii. 53 = Bredemeyera velutina.
villosa, *Aubl. Pl. Gui.* i. 388. t. 149.—Guian.
xanthopila, *Garcke, in Linnaea*, xxii. (1849) 51.—Guian.

TRIGONIASTRUM, Miq. Fl. Ind. Bat. Suppl. 394 (1860). *POLYGALEAE*, Benth. & Hook. f. i. 139.
Isopteris, Wall. Cat. n. 7261 (1831).
hypoleucum, *Miq. Fl. Ind. Bat. Suppl.* 395.—Malaya.

TRIGONIDIUM, Lindl. Bot. Reg. t. 1923 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 567.
acuminatum, *Batem. in Lindl. Bot. Reg.* (1838) *Misc.* 74.—Guian.
callistele, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 178.—Hab.?
Egertonianum, *Batem. in Lindl. Bot. Reg.* (1838) *Misc.* 73.—Honduras.
latifolium, *Lindl. Bot. Reg.* t. 1923.—Bras.
macranthum, *Rodrig. Orch. Nov.* i. 105.—Bras.
monophyllum, Griseb. Fl. Brit. W. Ind. 629 = Laelia monophylla.
obtusum, *Lindl. Bot. Reg.* t. 1923.—Guian.
ringens, Lindl. l. c. (1840) Misc. 57 = Mormolyce lineolata.
Seemanni, *Reichb. f. in Seem. Bot. Voy. Herald*, 214.—Panama.
spatulatum, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 280.—N. Granat.
tenue, *Lodd. Cat.* n. 582; *ex Lindl. Bot. Reg.* (1839) *Misc.* 44.—Guian.
turbinatum, *Reichb. f. in Hamb. Gartenz.* xv. (1859) 52.—Bras.

TRIGONIS, Jacq. Enum. Pl. Carib. 3 (1760); Select. Am. 102 (1763) = **Cupania**, Plum. (Sapind.).
tomentosa, Jacq. ll. cc. 19; 102 = C. americana.

TRIGONOCARPAEA, Steud. Nom. ed. II. ii. 710 (1841), sphalm. = TRIGONOCARPUS, Wall. = **Kokoona**, Thw. (Celastr.).

TRIGONOCARPUS, Bert. ex Steud. Nom. ed. II. ii. 710 (1841) = **Chorizanthe**, R. Br. (Polygon.).
cymosus, Bert. ex C. Gay, Fl. Chil. v. 283 = C. virgata.
secundiflorus, Bert. ex Steud. Nom. ed. II. ii. 710 = C. paniculata.

TRIGONOCARPUS, Vell. Fl. Flum. 153 (1825); iv. t. 15 (1827) = **Cupania**, Plum. (Sapind.).
racemosus, Vell. ll. cc. = C. castaneaefolia.

TRIGONOCARPUS, Wall. Cat. n. 6520 (1831) = **Kokoona**, Thw. (Celastr.).
littoralis, Wall. l. c. = K. zeylanica.

TRIGONOCARYUM, Trautv. in Act. Hort. Petrop. iii. ii. (1875) 278. *BORAGINEAE*, Benth. & Hook. f. ii. 856.
prostratum, *Trautv. l. c.* 279.—Reg. Cauc.

TRIGONOCHLAMYS, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 170. t. 27. *BURSERACEAE*, Benth. & Hook. f. i. 325.
Griffithii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 170.—Malacca.

TRIGONOPTERUM, Steetz, ex Anderss. in Vet. Acad. Handl. Stockh. 1853 (1855) 183 = **Lipochaeta**, DC. (Compos.).
Ponténi, Anderss. l. c. 184 (= *Macraea lancifolia*).—Ins. Galapag.

TRIGONOSCIADIUM, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 344 = **Heracleum**, Linn. (Umbell.).
tuberosum, Boiss. l. c. 345.—Mesopot.
viscidulum, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 1051.—Assyria.

TRIGONOSPERMUM, Less. Syn. Comp. 214 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 346.
adenostemmoides, *Less. l. c.*—Mexic.
melampodioides, *DC. Prod.* v. 509.—Mexic.

TRIGONOSTEMON, Blume, Bijdr. 600 (1825) (Trigostemon); Fl. Jav. Praef. p. viii (1828). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 298.
Athroisma, Griff. Notul. iv. 477 (1854).
Enchidium, Jack, in Malay. Misc. ii. (1822) n. vii. 89, nomen prius.
Silvaea, Hook. & Arn. Bot. Beech. Voy. 211 (1836).
Telogyne, Baill. Etud. Gén. Euph. 327 (1858).
Tylosepalum, Kurz, ex Teysm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 50.
Cumingii, Muell. Arg. in Linnaea, xxxiv. (1865–66) 213 = Tritaxis Cumingii.
diplopetalus, *Thw. Enum. Pl. Zeyl.* 277.—Zeylan.
Gaudichaudi, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 213.—Cochinch.
heteranthus, *Wight, Ic.* t. 1890.—Burma.
Hookerianus, Muell. Arg. in DC. Prod. xv. ii. 1109 = semperflorens.
indicus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 214.—Malaya.
laetus, *Baill. Étud. Gén. Euph.* 341.—Burma.
laevigatus, *Muell. Arg. in Flora*, xlvii. (1864) 538.—Borneo.
Lawianus, Muell. Arg. in Linnaea, xxxiv. (1865–66) 212 = Croton Lawianus.
longifolius, *Baill. Etud. Gén. Euph.* 341. t. 11.—Burma; Malaya.
macrophyllus, Muell. Arg. in Linnaea, xxxiv. (1865–66) 213 = Ostodes macrophylla.
malaccanus, *Muell. Arg. in Flora*, xlvii. (1864) 482.—Malacca.
nemoralis, *Thw. Enum. Pl. Zeyl.* 277.—Ind. or.
semperflorens, *Muell. Arg. in DC. Prod.* xv. ii. 1110.—Reg. Himal.
serratus, *Blume, Bijdr.* 600.—Java.
zeylanicus, Muell. Arg. in Linnaea, xxxiv. (1865–66) 213 = Ostodes zeylanica.

TRIGONOTHECA, Hochst. in Flora, xxiv. (1841) 662 = **Catha**, Forsk. (Celastr.).
serrata, Hochst. l. c. = C. edulis.

TRIGONOTHECA, Sch. Bip. in Flora, xxvii. (1844) 672, in syn. = **Melanthera**, Rohr (Compos.).
natalensis, Sch. Bip. l. c. = M. Brownei.

TRIGONOTIS, Stev. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 603. *BORAGINEAE*, Benth. & Hook. f. ii. 858.
Endogonia, Lindl. Veg. Kingd. 656 (1847), nomen prius.
brevipes, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 506, in nota.—Japon.
clavata, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 603.—As. temp.
Hookeri, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 173.—Reg. Himal.
microcarpa, *Benth. ex C. B. Clarke, l. c.* 172.—Reg. Himal.
multicaulis, *Benth. ex C. B. Clarke. l. c.*—Reg. Himal.
myosotidea, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 506, in nota.—Reg. Amur.
ovalifolia, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 172.—Reg. Himal.
pedunculatis, Benth. ex S. Moore & Baker, in Journ. Linn. Soc. xvii. (1879) 384, nomen = clavata.
petiolaris, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 506.—As. or.

TRIGONOTIS:—
radicans, *Stev. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 603.—Reg. Amur.
rotundifolia, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 172.—Reg. Himal.

TRIGOSTEMON, Blume, Bijdr. 600 (1825) = **Trigonostemon**, Blume (Euphorb.).

TRIGUERA, Cav. Diss. i. 41 (1785) = **Hibiscus**, Linn. (Malvac.).
acerifolia, Cav. l. c. = H. Solandra.

TRIGUERA, Cav. Diss. ii. App. 1. t. A (1786). *SOLANACEAE*, Benth. & Hook. f. ii. 897.
ambrosiaca, *Cav. l. c.* 2.—Hispan.
baccata, J. F. Gmel. Syst. 338 = Celsia bugulifolia?
inodora, Cav. Diss. ii. App. 3 (al. gen.?).—Hispan.
Osbeckii, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 524 = Celsia bugulifolia?

TRIGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 4; ed. II. 67 (1827) = **Clematis**, Dill. (Ranunc.).
trifoliata, Noronha, ex DC. Prod. i. 6 = C. Noronhiana.

TRIGYNAEA, Schlecht. in Linnaea, ix. (1834) 328. *ANONACEAE*, Benth. & Hook. f. i. 25.
angustifolia, *Benth. in Journ. Linn. Soc.* v. (1861) 70.—Venezuela.
canescens, *Benth. l. c.*—Venezuela.
Galeottiana, *Baill. Adansonia*, viii. (1867–68) 181.—Mexic.
grandis, *Benth. in Journ. Linn. Soc.* v. (1861) 70.—Venezuela.
Matthewsi, *Benth. l. c.* 69.—Peruv.
oblongifolia, *Schlecht. in Linnaea*, ix. (1834) 329.—Bras.
Perrottetii, *Baill. Adansonia*, viii. (1867–68) 179.—Guian.
rufescens, *Baill. l. c.* 180.—Guian.

TRIGYNEIA, Reichb. Handb. 278 (1837), sphalm. = praec.

TRIKALIS, Rafin. Fl. Tellur. iii. 47 (1836) = **Suaeda**, Forsk. (Chenop.).
triflora, Rafin. l. c. = S. altissima.
trigyna, Moq. in DC. Prod. xiii. ii. 158 = S. altissima.

TRILEPIS, Nees, in Edinb. N. Phil. Journ. xvii. (Oct. 1834) 267 = **Kobresia**, Willd. (Cyperac.).
abyssinica, Boeck. in Linnaea, xxxix. (1875) 9 = Eriospora abyssinica.
Lhotzkiana, Nees, in Edinb. N. Phil. Journ. xvii. (Oct. 1834) 267 = Fintelmannia restioides.
pilosa, Boeck. in Linnaea, xxxix. (1875) 10 = Eriospora pilosa.
Royleana, Nees, in Edinb. N. Phil. Journ. xvii. (Oct. 1834) 267 = K. Royleana.

TRILEPISIUM, Thou. Gen. Nov. Madag. 22 (1806). *INCERT. SEDIS.* Cf. Benth. & Hook. f. i. 605.
madagascariense, *DC. Prod.* ii. 639.—Madag.

TRILIENA, Rafin. Sylva Tellur. 54 (1838) = **Lycium**, Linn. (Solan.).

TRILISA, Cass. in Bull. Soc. Philom. (1818) 140; et in Dict. Sc. Nat. xvi. 10 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 248.
odoratissima, *Cass. ll. cc.*—Am. bor.
paniculata, *Cass. in Dict. Sc. Nat.* lv. 310.—Am. bor.

TRILIX, Linn. Mant. ii. 153 (1771) = **Prockia**, P. Br. (Tiliac.).
Crucis, Griseb. in Goett. Abh. vii. (1857) 165 = P. Crucis.
glandulosa, Domb. ex Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 288 = Banara guianensis.
guianensis, Griseb. in Goett. Abh. vii. (1857) 165 = Banara guianensis.
hygroscopica, Liais, Brésil, 617.—Bras.
iraguensis, Griseb. in Goett. Abh. vii. (1857) 165, sphalm. = Banara ibaguensis.
lutea, Linn. Mant. ii. 247 = P. Crucis?

TRILLIDIUM, Kunth, Enum. Pl. v. 120 (1850) = **Trillium**, Linn.
Govanianum, Kunth, l. c. = Trillium Govanianum.
japonicum, Franch. & Sav. Enum. Pl. Jap. ii. 525.—Japon.

TRILLIUM, Linn. Sp. Pl. ed. I. 339 (1753). *LILIACEAE*, Benth. & Hook. f. iii. 833.
DELOSTYLIS, Rafin. in Journ. Phys. lxxxix. (1819) 102.
ESDRA, Salisb. Gen. Pl. Fragm. 60 (1866).
PHYLLANTHERUM, Rafin. in Journ. Phys. xci. (1820) 72.
TRILLIDIUM, Kunth, Enum. Pl. v. 120 (1850).
acuminatum, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Am. bor.
atropurpureum, M. A. Curt. ex Beck, Bot. N. & M. St. 361 = erectum.
amblopsis, Rafin. Med. Fl. ii. 99 (Quid?).—Am. bor.
angustifolium, Rafin. l. c. 98 (Quid?).—Am. bor.
brevipetalum, Rafin. l. c. 100 (Quid?).—Am. bor.
californicum, Kellogg, in Proc. Calif. Acad. ii. (1863) 50 = ovatum.
camtschaticum, Pall. ex Pursh, Fl. Am. Sept. i. 246 = erectum?
Catesbaei, Ell. Sketch, i. 429 = stylosum.
cernuum, Barton, Fl. N. Am. ii. 13. t. 40 = erectum.
cernuum, *Linn. Sp. Pl.* 339.—Am. bor.
cernuum, Michx. Fl. Bor. Am. i. 216 = stylosum.
discolor, Wray, ex Hook. Bot. Mag. t. 3097 = sessile.
divaricatum, Rafin. Med. Fl. ii. 102 (Quid?).—Am. bor.
erectum, *Linn. Sp. Pl.* 340.—Am. bor.
erectum, Stell. in Pall. N. Nord. Beitr. ii. 300 = obovatum.
erythrocarpum, Curt. Bot. Mag. t. 855 = grandiflorum.
erythrocarpum, *Michx. Fl. Bor. Am.* i. 216.—Am. bor.
flavum, Rafin. Med. Fl. ii. 100 (Quid?).—Am. bor.
foetidum, Salisb. Parad. Lond. t. 35 = erectum.
glaucum, Rafin. Med. Fl. ii. 102 = erectum.
Govanianum, *Wall. Cat.* n. 812; *Royle, Illustr. Bot. Himal.* 384. t. 93. f. 1.—Reg. Himal.
grandiflorum, *Salisb. Parad. Lond.* t. 1.—Am. bor.
hamosum, Rafin. Med. Fl. ii. 101 (Quid?).—Am. bor.
isanthum, Rafin. l. c. 98 (Quid?).—Am. bor.
kamtschatikum, Ledeb. Fl. Ross. iv. 121 = obovatum.
lanceolatum, Boykin, ex S. Wats. in Proc. Am. Acad. xiv. (1879) 274 = recurvatum.
latifolium, Rafin. Med. Fl. ii. 96, 101 = cernuum.
lirioides, Rafin. l. c. 100 (Quid?).—Am. bor.
longiflorum, Rafin. l. c. 97 = sessile.
maculatum, Rafin. l. c. 103 = sessile.
medium, Rafin. l. c. 102 = cernuum.
membranaceum, Rafin. l. c. 98 (Quid?).—Am. bor.
nervosum, Ell. Sketch, i. 429 = stylosum.
nivale, *Riddell, Syn. Fl. W. St.* 93.—Am. bor.
nutans, Rafin. Med. Fl. ii. 99 = erectum.
obcordatum, Rafin. l. c. 101 = grandiflorum?
obovatum, Hook. Fl. Bor. Am. ii. 180 = ovatum.
obovatum, Pursh, Fl. Am. Sept. i. 245 = erectum?
ovatum, *Pursh, l. c.*—Am. bor. occ.
pendulum, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 421 = erectum.
petiolatum, *Pursh, Fl. Am. Sept.* i. 244.—Am. bor. occ.
pictum, Pursh, l. c. = erythrocarpum.
pumilum, Pursh, l. c. 245 = pusillum.
purpureum, Kin, ex Ell. Sketch, i. 430 = erectum.
pusillum, *Michx. Fl. Bor. Am.* i. 215.—Am. bor.
rectum, D. Dietr. Syn. Pl. ii. 1212, sphalm. = *erectum*.
recurvatum, *Beck, in Am. Journ. Sc.* Ser. I. xi. (1826) 178.—Am. bor.
rhomboideum, Michx. Fl. Bor. Am. i. 215 = erectum, grandiflorum.
rivale, *S. Wats. in Proc. Am. Acad.* xx. (1885) 378.—Am. bor. occ.
roseum, Hort. ex Steud. Nom. ed. II. ii. 710 (Quid?).
rotundifolium, Rafin. Med. Fl. ii. 97 (Quid?).—Am. bor.
sessile, *Linn. Sp. Pl.* 340.—Am. bor.
Smallii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1884) 217.—Japon.
spatulatum, Rafin. Med. Fl. ii. 101 = erectum.
stylosum, *Nutt. Gen. Am.* i. 239.—Am. bor.
texanum, Buckl. in Proc. Acad. Sc. Philad. 1860 (1861) 443 = pusillum.
tinctorium, Rafin. Med. Fl. ii. 98 (Quid?).—Am. bor.

TRILLIUM:—
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xxix. (1884) 218.—Reg. Himal.; Japon.
undulatum, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 422 = erythrocarpum.
unguiculatum, Nutt. in Trans. Am. Phil. Soc. v. (1837) 154 = recurvatum.
unguiculatum, Rafin. Med. Fl. ii. 98 (Quid?).—Am. bor.
viride, Beck, in Am. Journ. Sc. Ser. I. xi. (1826) 178 = sessile.
viridescens, Nutt. in Trans. Am. Phil. Soc. v. (1837) 155 = sessile.

TRILOBULINA, Rafin. Fl. Tellur. iv. 110 (1836) = **Utricularia**, Linn. (Lentib.).
crenata, Rafin. l. c. = U. aphylla.
fibrosa, Rafin. l. c. = U. fibrosa.
striata, Rafin. l. c. = U. fibrosa.

TRILOMISA, Rafin. Fl. Tellur. ii. 91 (1836) = **Begonia**, Tourn.
undulata, Rafin. l. c. = B. undulata.

TRILOPHUS, Fisch. Hort. Gorenk. ed. II. 56 (1812) = **Menispermum**, Tourn.
Ampelisagria, Fisch. l. c. = M. dauricum.

TRILOPUS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 219 = **Hamamelis**, Linn.
dentata, Rafin. New Fl. Am. iii. 17 = H. virginiana.
estivalis, Rafin. l. c. 16 = Fothergilla Gardeni?
hyemalis, Rafin. l. c. = H. virginiana.
nigra, Rafin. l. c. = H. virginiana.
parvifolia, Rafin. l. c. 17 = H. virginiana.
riparia, Rafin. l. c. 16 = H. virginiana.
rotundifolia, Rafin. l. c. = H. virginiana.
virginica, Rafin. l. c. 15 = H. virginiana.

TRIMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Anacardium**? Linn.

TRIMEIANDRA, Rafin. Fl. Tellur. iv. 105 (1836) = **Arthosolen**, C. A. Mey. (1843) (Thymel.).
spicata, Rafin. l. c. = A. spicatus.

TRIMELOPTER, Rafin. Fl. Tellur. ii. 24 (1836) = **Ornithogalum**, Tourn. (Liliac.).
fuscatum, Rafin. l. c. = O. fuscatum.

TRIMENIA, Seem. Fl. Vit. 425. t. 99 (1871). *MONIMIACEAE*, Benth. & Hook. f. iii. 143.
weinmanniaefolia, *Seem. l. c.*—Ins. Fiji.

TRIMERANTHES, Cass. in Dict. Sc. Nat. xlix. 115 (1827) = **Siegesbeckia**, Linn. (Compos.).
dichotoma, Cass. l. c. = S. flosculosa.

TRIMERANTHUS, Karst. in Linnaea, xxx. (1859–60) 159 = **Chaetolepis**, Miq. (Melastom.).
alpestris, Karst. l. c. = C. alpestris.

TRIMERIA, Harv. Gen. S. Afr. Pl. Suppl. 417 (1858). *BIXINEAE*, Benth. & Hook. f. i. 129.
MONOSPORA, Hochst. in Flora, xxiv. (1841) 660.
RENARDIA, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 466.
alnifolia, *Harv. in Harv. & Sond. Fl. Cap.* i. 69.—Afr. austr.
trinervis, *Harv. Gen. S. Afr. Pl.* 417.—Afr. austr.

TRIMERIS, Presl, Prod. Monog. Lobel. 46 (1836) = **Lobelia**, Plum. (Campan.).
oblongifolia, Presl, l. c. = L. scaevolaefolia.

TRIMERISMA, Presl, Bot. Bemerk. 73 (1844) = **Platylophus**, D. Don (Saxifr.).
trifoliatum, Presl, l. c. = P. trifoliatus.

TRIMERIZA, Lindl. Bot. Reg. sub t. 1543 (1832) = **Bragantia**, Lour. (Aristol.).
piperina, Lindl. l. c. = B. Wallichii.

TRIMETRA, Moç. ex DC. Prod. vii. 262 (1838) = **Borrichia**, Adans. (Compos.).
ficoidea, Moc. l. c. = B. frutescens.

TRIMEZA, Salisb. in Trans. Hort. Soc. i. (1812) 308. *IRIDEAE*, Benth. & Hook. f. iii. 690.
LANSBERGIA, De Vriese, Epim. Ind. Sem. Hort. Lugd. Bat. (1846) 2.
POARCHON, Allem. Tract. Rio Jan. (1846).
REMACLEA, C. Morr. Belg. Hortic. iii. (1853) 1. t. 1.
XANTHOCROMYON, Karst. in Bot. Zeit. v. (1847) 694.
caracasana, *Benth. & Hook. f. Gen.* iii. 690.—Venezuela.
juncifolia, *Benth. & Hook. f. l. c.*—Bras.
lurida, *Salisb. in Trans. Hort. Soc.* i. (1812) 308.—Mexic.
martinicensis, Herb. in Bot. Reg. (1844) Misc. 88 = lurida.
meridensis, Herb. l. c. = lurida.

TRIMORPHA, Cass. in Dict. Sc. Nat. iii. Suppl. 65 (1816) = **Erigeron**, Linn. (Compos.).
acre, S. F. Gray, Nat. Arr. Brit. Pl. ii. 466 = E. acris.
alpina, S. F. Gray, l. c. 467 = E. alpinus.
glabrata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 402 = E. glabratus.
uniflora, S. F. Gray, Nat. Arr. Brit. Pl. ii. 467 = E. uniflorus.
vulgaris, Cass. in Dict. Sc. Nat. lv. 324 = E. acris.

TRIMORPHAEA, Cass. in Dict. Sc. Nat. xxxvii. 462 (1825) = praec.

TRIMORPHANDRA, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 190; et in Ann. Sc. Nat. Sér. V. ii. (1864) 148 = **Hibbertia**, Andr. (Dillen.).
pulchella, Brongn. & Gris, ll. cc.—N. Caled.

TRIMORPHOEA, Benth. & Hook. f. Gen. ii. 280 (1873), err. typ. = TRIMORPHAEA, Cass. = **Erigeron**, Linn. (Compos.).

TRINACTE, Gaertn. Fruct. ii. 415 (1791) = **Jungia**, Linn. (Compos.).
ferruginea, Gaertn. l. c. = J. ferruginea.

TRINAX, D. Dietr. Syn. Pl. ii. 1091 (1840), sphalm. = **Thrinax**, Linn. f. (Palm.).

TRINCHINETTIA, Endl. Gen. 1383 (1841) = **Geissopappus**, Benth. (Compos.).
caleoides, Endl. ex Walp. Rep. vi. 180 = G. caleoides.

TRINCIATELLA, Adans. Fam. ii. 112 (1763) = **Hyoseris**, Linn. (Compos.).

TRINEURIA, Presl, Bot. Bemerk. 30 (1844); (in Abh. Boehm. Ges. Wiss. V. iii. (1845) 560) = **Aspalathus**, Linn. (Legumin.).
appendiculata, Presl, Bot. Bemerk. 131 = A. ciliaris.
Chenopoda, Presl, l. c. = A. Chenopoda.
ciliaris, Presl, l. c. = A. ciliaris.
cochleariformis, Presl, l. c. = A. uniflora.
comosa, Presl, l. c. = A. comosa.
cymbaeformis, Presl. l. c. = A. uniflora.
deciduifolia, Presl, l. c. = A. nigra.
fuscescens, Presl, l. c. 131, 133 = A. nigra.
linarifolia, Presl, l. c. 131 = A. linearifolia.
marginalis, Presl, l. c. = A. marginalis.
papillosa, Presl, l. c. = A. ciliaris.
rigescens, Presl, l. c. = A. rigescens.

TRINEURON, Hook. f. Fl. Antarct. i. 23. t. 17 (1845) = **Abrotanella**, Cass. (Compos.).
nivigenum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 105 = A. nivigena.
pusillum, Hook. f. Fl. N. Zel. 131 = A. pusilla.
scapigerum, F. Muell. in Hook. Kew Journ. ix. (1857) 301 = A. scapigera.
spathulatum, Hook. f. Fl. Antarct. 24. t. 17 = A. spathulata.

TRINGA, Roxb. Hort. Beng. [81], (1814), nomen, sphalm. = TUNGA, Roxb. = **Hypolytrum**, Rich. (Cyperac.).

TRINIA, Hoffm. Gen. Umbell. ed. I. 92 (1814). *UMBELLIFERAE*, Benth. & Hook. f. i. 887.
APINELLA, Neck. Elem. i. 191 (1790).
GRAMMOPETALUM, C. A. Mey. ex Meinsh. in Linnaea, xxx. (1859–60) 514.

TRINIA:—
RUMIA, Hoffm. Gen. Umbell. ed. II. 171 (1816).
SPIELMANNIA, Cuss. ex Juss. in Dict. Sc. Nat. lv. 328 (1828).
dahurica, Turcz. ex Bess. in Flora, xvii. (1834) 1. Beibl. 14 = seselioides.
dioica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 392 = vulgaris.
Dufourei, *DC. Prod.* iv. 104.—Hispan.
elatior, Gaud. Fl. Helvet. ii. 413 = vulgaris.
glaberrima, Hoffm. Gen. Umbell. ed. I. 93 = hispida, vulgaris.
glauca, Hoffm. ex Steud. Nom. ed. II. ii. 711 = Kitaibelii.
glauca, Reichb. Pl. Crit. v. 25; Dum. Fl. Belg. Prod. 78 = vulgaris.
Henningii, *Hoffm. Gen. Umbell.* ed. I. 94.—Reg. Cauc. et Casp.
Henningii, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 127 = vulgaris.
hispida, *Hoffm. Gen. Umbell.* ed. I. 94.—Tauria; Reg. Cauc.
Hoffmanni, Bieb. Fl. Taur. Cauc. iii. 244 = hispida.
Kitaibelii, *Bieb. Fl. Taur. Cauc.* iii. 246.—Europ. or.; As. bor.
Lessingii, *Reichb. f. Ic. Fl. Germ.* xxi. 8.—Europ.
pumila, Reichb. Fl. Germ. Excurs. 473 = vulgaris.
ramosissima, Fisch. ex Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 127 = Kitaibelii.
scabra, *Boiss. & Noé, Diagn.* Ser. II. ii. 73.—As. Min.
seselioides, *Ledeb. Fl. Alt.* i. 357.—Sibir.
swellendamensis, Eckl. & Zeyh. Enum. 340 = Rhyticarpus Ecklonis.
uitenhagensis, Eckl. & Zeyh. l. c. = Rhyticarpus rugosus.
vulgaris, *DC. Prod.* iv. 103.—Europ.

TRINIUSA, Steud. Syn. Pl. Gram. 328 (1855) = **Bromus**, Dill. (Gramin.).
Danthonia, Steud. l. c. = B. macrostachys.
flavescens, Steud. l. c. = B. fasciculatus.

TRINOGETON, Walp. Rep. vi. 615 (1847), sphalm. = THINOGETON, Benth. = **Cacabus**, Bernh. (Solan.).

TRIODALLUS, A. DC. in DC. Prod. vii. 491 (1839) = seq.

TRIODANIS, Rafin. New Fl. Am. iv. 67 (1836) = **Specularia**, Heist. (Campan.).
rupestris, Rafin. l. c. = S. perfoliata.
scabra, Rafin. l. c. = S. perfoliata.

TRIODIA, R. Br. Prod. 182 (1810).—*GRAMINEAE*, Benth. & Hook. f. iii. 1175.
BRACHATERA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189.
DASYOCHLOA, Willd. ex Steud. Nom. ed. II. i. 484 (1840).
LEPTOCARYDION, Hochst. ex Benth. & Hook. f. Gen. iii. 1176 (1883).
MERISACHNE, Steud. Syn. Pl. Gram. 117 (1855).
RHOMBOLYTRUM, Link, Hort. Berol. ii. 96 (1827).
SIEGLINGIA, Bernh. Verz. Pfl. Erf. 44 (1799), nomen prius.
TRICHONEURA, Andersson in Vet. Akad. Handl. Stockh. 1853 (1855) 148.
TRICUSPIS, Beauv. Agrost. 77. t. 15. f. 10 (1812).
TRIDENS, Roem. & Schult. Syst. ii. 34 (1817).
URALEPIS, Spreng. Neue Entdeck. i. 244 (1821).
URALEPSIS, Nutt. Gen. Am. i. 62 (1818).
WINDSORIA, Nutt. Gen. Am. i. 70 (1818).
acuminata, *Benth. ex Vasey, Gr. U. St.* 35.—Am. bor. occ.
albescens, *Benth. ex Vasey, l. c.*—Texas.
alpina, Roth, Enum. Pl. Phaen. Germ. i. 1. 381 = Danthonia provincialis.
ambigua, *Benth. ex Vasey, Gr. U. St.* 35.—Am. bor.
ambigua, R. Br. Prod. 183 = Diplachne fusca.
antarctica, Benth. in Journ. Linn. Soc. xix. (1881) 111 = Catabrosa antarctica.
antarctica, *Hook. f. Fl. Antarct.* 380.—Reg. Magell.
avenacea, *H. B. & K. Nov. Gen. et Sp.* i. 156. t. 48.—Mexic.
Billardieri, Spreng. Syst. i. 330 = Festuca littoralis.
coerulescens, Desv. Opusc. 99, in obs. = cuprea.
Cunninghamii, *Benth. Fl. Austral.* vii. 606.—Austral.

TRIODIA:—
cuprea, *Jacq. Eclog. Gram.* 21. t. 16.—Am. bor.
cynosuroides, Spreng. Syst. i. 628 = Dactylis cynosuroides.
decumbens, *Beauv. Agrost.* 76. t. 15. f. 9 = Europ.; Afr. bor.
exigua, *T. Kirk, in Gard. Chron.* (1882) 1. 260; *et in Trans. N. Z. Inst.* xiv. (1882) 378.—N. Zel.
festucacea, Eichw. Skizze, 119 = Graphephorum festucaceum.
festucacea, Roth, Enum. Pl. Phaen. Germ. i. 1. 382 = Graphephorum arundinaceum.
festucoides, *Desv. Opusc.* 98.—'Am. bor.'
filiformis, *Nees, ex Steud. Syn. Pl. Gram.* 249.—Chili.
Greenii, Spreng. Syst. i. 330 = Graphephorum Elliotii.
irritans, *R. Br. Prod.* 182.—Austral.
kerguelensis, Hook. f. Fl. Antarct. 379. t. 139 = Festuca kerguelensis.
laxa, Benth. Pl. Hartw. 194, sphalm. = Triodon laxus.
litoralis, Spreng. Syst. i. 330 = Danthonia paradoxa.
melicoides, Spreng. l. c. 331 = Graphephorum melicoideum.
microdon, F. Muell. Census, 135 = Triraphis microdon.
microstachya, *R. Br. Prod.* 182.—Austral.
Mitchelli, *Benth. Fl. Austral.* vii. 606.—Austral.
mutica, *Benth. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 180.—Am. bor. occ.
novaeboracensis, Desv. Opusc. 99, in obs. = cuprea.
pallida, Spreng. Neue Entdeck. i. 246 = Glyceria pallida.
parviflora, R. Br. Prod. 182 = Diplachne parviflora.
plumosa, *Benth. in Journ. Linn. Soc.* xix. (1881) 110.—Afr. trop.
procera, *R. Br. Prod.* 182.—Austral.
pulchella, *H. B. & K. Nov. Gen. et Sp.* i. 155. t. 47.—Am. bor. occ.
pungens, *R. Br. Prod.* 182.—Austral.
pungens, Lindl. in Mitch. Journ. Trop. Austral. 340 = Mitchelli.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 181.—Mexic.
seslerioides, *Benth. ex Vasey, Gr. U. St.* 35.—Am. bor.
splendida, Steud. Syn. Pl. Gram. 249 = Deschampsia caespitosa.
stricta, *Benth. ex Vasey, Gr. U. St.* 35.—Texas.
texana, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 180.—Mexic.
tridenticulata, *Nees, ex Steud. Syn. Pl. Gram.* 249.—Reg. Argent.
trinerviglumis, *Benth. ex Vasey, Gr. U. St.* 35.—Texas.
viscida, Roem. & Schult. Syst. ii. 599 = pungens.

TRIODICA, Steud. Nom. ed. II. ii. 711 (1841), sphalm. = TRIADICA, Lour. = **Sapium**, P. Br. (Euphorb.).

TRIODON, DC. Prod. iv. 566 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 143.
EBELIA, Reichb. Nom. 74 (1841).
angulatus, *Benth. Pl. Hartw.* 70.—Mexic.
anthospermoides, *DC. Prod.* iv. 566.—Bras.
glomeratus, *DC. l. c.*—Bras.
laxus, *Benth. Pl. Hartw.* 194 (sphalm., Triodia).—Ecuador.
polymorphus, *DC. Prod.* iv. 566.—Bras.

TRIODON, Rich. in Pers. Syn. i. 60, in nota (1805) = **Rynchospora**, Vahl (Cyperac.).
setaceus, Rich. l. c. = R. tenerrima, *Nees*.

TRIODUS, Rafin. in Journ. Phys. lxxxix. (1819) 106 = **Carex**, Dill. (Cyperac.).

TRIOLENA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 328. *MELASTOMACEAE*, Benth. & Hook. f. i. 757.
hirsuta, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 81.—Am. trop.
pustulata, *Triana, l. c.*—Ecuador.
scorpioides, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 328.—Mexic.

TRIOMMA, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 171. *BURSERACEAE*, Benth. & Hook. f. i. 323.
malaccensis, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 171.—Malacca.

TRIONAEA, Medic. Malv. 45 (1787) = **Hibiscus**, Linn. (Malvac.).

TRIONUM, Linn. Syst. ed. 1 (1735); Medic. Malv. 46 (1787) = **Hibiscus**, Linn. (Malvac.).
annuum, Medic. Malv. 47 = H. Trionum.
cordifolium, Moench, Meth. Suppl. 202 = H. Trionum.
diffusum, Moench, Meth. 618 = H. Trionum.
frutescens, Medic. Malv. 47 = H. Trionum.

TRIOPTERIS, Linn. Gen. ed. I. 343 (1737). *MALPIGHIACEAE*, Benth. & Hook. f. i. 259.
acuminata, Willd. Sp. Pl. ii. 745 = Tetrapteris mucronata.
acutifolia, Willd. l. c. 744 = Tetrapteris acutifolia.
argentea, Salzm. ex Griseb. in Mart. Fl. Bras. xii. 1. 50 = Banisteria nummifera.
bifurca, Gaertn. Fruct. ii. 168 = Tetrapteris citrifolia.
brasiliensis, Poir. Encyc. viii. 105 = Banisteria ferruginea.
buxifolia, Willd. Sp. Pl. ii. 745 = Tetrapteris buxifolia.
citrifolia, Sw. Prod. Veg. Ind. Occ. 75 = Tetrapteris citrifolia.
discolor, G. F. W. Mey. Prim. Fl. Esseq. 182 = Tetrapteris discolor.
divaricata, A. Rich. ex A. Juss. in Arch. Mus. Par. iii. (1843) 412 = Banisteria divaricata.
floribunda, *Thunb. Pl. Bras. Dec.* i. 14.—Bras.
havanensis, *H. B. & K. Nov. Gen. et Sp.* v. 168. t. 451.—Cuba.
Hiraea, Gaertn. Fruct. ii. 169 = Hiraea reclinata.
inaequalis, Raeusch. Nom. ed. III. 131 = Tetrapteris citrifolia.
indica, Willd. Sp. Pl. ii. 744 = Aspidopterys Roxburghiana.
jamaicensis, Blanco, Fl. Filip. ed. I. 379 = Hiptage Madablota.
jamaicensis, *Linn. Sp. Pl.* 428.—Ind. occ.
lingulata, *Poir. Encyc.* viii. 104.—Ins. S. Doming.
lucida, *H. B. & K. Nov. Gen. et Sp.* v. 167. t. 451.—Cuba.
macrophylla, Salzm. ex Griseb. in Mart. Fl. Bras. xii. 1. 99 = Hiraea Salzmanniana.
mexicana, Walp. Rep. v. 314, sphalm. = Tetrapteris Schiedeana.
odorata, Poir. Encyc. viii. 108 = Triaspis odorata.
ovata, *Cav. Diss.* 431. t. 259.—Ins. S. Doming.
parvifolia, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 266.—Ins. Baham.
pauciflora, G. F. W. Mey. Prim. Fl. Esseq. 183 = Diplopterys Paralias.
pinnata, Poir. Encyc. viii. 108 = Flabellaria paniculata.
Poliandra, Blanco, Fl. Filip. ed. I. 380 = Berrya Ammonilla.
polycarpa, Salzm. ex A. Juss. in Arch. Mus. Par. iii. (1843) 562 = Hiraea maritima.
quadrialata, Pav. ex A. Juss. l. c. 539 = Tetrapteris Schiedeana.
reclinata, Cav. Diss. ix. 431 = Hiraea reclinata.
rigida, *Sw. Prod. Veg. Ind. Occ.* 75.—Ins. S. Doming.
sericea, Schlecht. in Linnaea, x. (1836) 243 = Gaudichaudia Schiedeana.
volubilis, Salzm. ex A. Juss. in Arch. Mus. Par. iii. (1843) 552 = Hiraea sepium.

TRIOPTERYS, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. 265 (1840) = praec.

TRIOPTOLEMEA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 102 = TRIPTOLEMEA, Mart. = **Dalbergia**, Linn. (Legumin.).

TRIOSTEON, Dill. ex Adans. Fam. ii. 500 (1763) = seq.

TRIOSTEUM, Linn. Sp. Pl. ed. I. 176 (1753). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 4.
angustifolium, *Linn. l. c.*—Am. bor.
connatum, Rafin. New Fl. Am. ii. 35 = perfoliatum.
emodorum, Spreng. Syst. iv. Cur. Post. 82 = hirsutum.
himalayanum, Wall. Cat. n. 484 = hirsutum.
hirsutum, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 538.—Reg. Himal.
hispidum, Rafin. New Fl. Am. ii. 37 (Quid?).—Am. bor.
levigatum, Rafin. l. c. (Quid?).—Am. bor.
majus, Michx. Fl. Bor. Am. i. 107 = perfoliatum.
minus, Michx. l. c. = angustifolium.
molle, Rafin. New Fl. Am. ii. 36 (Quid?).—Am. bor.
obovatum, Rafin. l. c. 37 = perfoliatum.

TRIOSTEUM:—
perfoliatum, *Linn. Sp. Pl.* 176.—Am. bor.
pinnatifidum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 476.—China.
pumilum, Rafin. New Fl. Am. ii. 36 = angustifolium.
sinuatum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 373.—Japon.
triflorum, Vahl, Symb. Bot. iii. 37 = Sabicea triflora.
villosum, Rafin. New Fl. Am. ii. 36 (Quid ?).—Am. bor.

TRIPAGANDRA, Rafin. Fl. Tellur. ii. 16 (1836) = **Tradescantia**, Linn. (Commel.).
multiflora, Rafin. l. c. = Trad. multiflora.

TRIPENTAS, Casp. in Monatsb. Berl. Akad. 1857 (1858) 43; et in Pringsh. Jahrb. i. (1858) 497, in obs. = **Hypericum**, Tourn.

TRIPETALEIA, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 731. t. 3. f. 2 = **Elliottia**, Muhl. (Ericac.).
bracteata, Maxim. in Bull. Acad. Pétersb. xi. (1867) 433 = E. bracteata (in Addend.).
paniculata, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 731 = E. paniculata (in Addend.).

TRIPETELUS, Lindl. in Mitch. Three Exped. ii. 14 (1839) = **Sambucus**, Tourn. (Caprifol.).
australasicus, Lindl. l. c. = S. xanthocarpa.

TRIPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Cupania**, Plum. (Sapind.).

TRIPHACA, Lour. Fl. Cochinch. 577 (1790) = **Sterculia**, Linn.
africana, Lour. l. c. = S. Triphaca.

TRIPHASIA, Lour. Fl. Cochinch. 152 (1790).
RUTACEAE, Benth. & Hook. f. i. 303.
Aurantiola, *Lour. l. c.* 153.—As. trop.
diacantha, M. Roem. Syn. Hesper. 37 = T. Aurantiola.
glauca, Lindl. in Mitch. Journ. Trop. Austral. 353 = Atalantia glauca.
javanica, M. Roem. Syn. Hesper. 38 = T. Aurantiola.
monophylla, DC. Prod. i. 536 = Atalantia trimera.
sarmentosa, Blume, Bijdr. 132 = Luvunga eleutherandra.
trifoliata, DC. Prod. i. 536; Blume, l. c. = T. Aurantiola.

TRIPHELIA, R. Br. ex Endl. in Enum. Pl. Hueg. 48 (1837) = **Actinodium**, Schau. (Myrtac.).
brunioides, R. Br. ex Endl. l. c. = A. Cunninghamii.

TRIPHORA, Nutt. Gen. Am. ii. 192 (1818) = **Pogonia**, Juss. (Orchid.).
gentianoides, Nutt. l. c. 193, in nota = P. gentianoides.
pendula, Nutt. l. c. = P. pendula.

TRIPHYLLOIDES, Moench, Meth. 509 (1794) = **Trifolium**, Tourn. (Legumin.).
angustifolia, Moench, l. c. 510 = Trifolium angustifolium.
clypeata, Moench, l. c. = Trifolium clypeatum.
comosa, Moench, l. c. 509 = Trifolium subterraneum.
nervosa, Moench, l. c. 510 = Trifolium alpestre.
rigida, Moench, l. c. 509 = Trifolium lappaceum.
rubens, Moench, l. c. = Trifolium rubens.

TRIPHYLLUM, Medic. Phil. Bot. i. 202 (1789) = **Medicago**, Tourn. (Legumin.).
platycarpos, Medic. l. c. = M. platycarpa.

TRIPHYSARIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 52 (1835) = **Orthocarpus**, Nutt. (Scrophul.).
versicolor, Fisch. & Mey. l. c. = O. erianthus.

TRIPINNA, Lour. Fl. Cochinch. 391 (1790) = **Colea**, Boj. (Bignon.).
africana, Voigt, Hort. Suburb. Calc. 479.—Afr. trop. *vel* Ins. Mascar.
tripinnata, Lour. Fl. Cochinch. 391 = C. tripinnata.

TRIPINNARIA, Pers. Syn. ii. 173 (1807) = **Colea**, Boj. (Bignon.).
africana, Spreng. Syst. ii. 842 (= *Tripinna africana*).—Afr. trop. *vel* Ins. Mascar.

TRIPINNARIA:—
asiatica, Spreng. Syst. ii. 842 = Colea tripinnata.
cochinchinensis, Pers. Syn. ii. 173 = C. tripinnata.
tripinnata, Steud. Nom. ed. II. ii. 712 = C. tripinnata.

TRIPLACHNE, Link, Hort. Berol. ii. 241 (1833).
GRAMINEAE, Benth. & Hook. f. iii. 1151.
nitens, *Link, l. c.*—Sicil.

TRIPLADENIA, D. Don, in Proc. Linn. Soc. i. (1839) 46 = **Kreysigia**, Reichb. (Liliac.).
Cunninghamii, D. Don, l. c. = K. multiflora.

TRIPLANDRA, Rafin. Sylva Tellur. 62 (1838) = **Croton**, Linn. (Euphorb.).
lanata, Rafin. l. c. 63 = C. lanatus.

TRIPLANDRON, Benth. Bot. Voy. Sulph. 73. t. 38 (1844) = **Clusia**, Linn. (Guttif.).
lineatum, Benth. l. c. = C. lineata.
lineatum, Seem. Bot. Voy. Herald, 88 = C. Seemanni.

TRIPLARINA, Rafin. Sylva Tellur. 104 (1838) = **Baeckea**, Linn. (Myrtac.).
camphorata, Rafin. l. c. = B. camphorata.

TRIPLARIS, Loefl. It. Hisp. 256 (1758); Linn. Syst. ed. X. 881 (1759). *POLYGONACEAE*, Benth. & Hook. f. iii. 104.
BLOCHMANNIA, Reichb. Consp. 163 (1828).
VELLASQUEZIA, Bertol. Fl. Guatim. 89. t. 11 (1840).
americana, Aubl. Pl. Gui. ii. 910. t. 347; Rottb. in Act. Lit. Univ. Hafn. i. (1778) 274. t. 3 = surinamensis.
americana, H. B. & K. Nov. Gen. et Sp. ii. 182 = felipensis.
americana, *Linn. Syst.* ed. X. 881.—Am. austr.
americana, Pav. ex Meissn. in DC. Prod. xiv. 175 = auriculata.
americana, Rob. Schomb. in Ann. Nat. Hist. Ser. I. i. (1838) 264 = Schomburgkiana.
americana, Vahl, Symb. Bot. ii. 100, partim = Vahliana.
apetala, Walp. Ann. iii. 295 = Ruprechtia apetala.
Arnottiana, *Meissn. in DC. Prod.* xiv. 176.—Ecuador.
auriculata, *Meissn. l. c.* 174.—Mexic.
Bonplandiana, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 262.—Peruv.
brachystachya, Endl. Gen. Suppl. iv. II. 55 = Ruprechtia brachystachya.
brasiliana, *Cham. in Linnaea*, viii. (1833) 139.—Bras.
caracassana, *Cham. l. c.* 137.—N. Granat.
colombiana, Meissn. in DC. Prod. xiv. 175 = Cumingiana.
coriacea, *Karst. Fl. Columb.* ii. 131. t. 169.—Venezuela.
crenata, *Casar. Nov. Stirp. Bras. Dec.* 80 (sp. dub.).—Bras.
Cumingiana, *Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iv. (1845) 148.—N. Granat.
cuneata, Endl. Gen. Suppl. iv. II. 55, sphalm. = crenata.
felipensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 263.—N. Granat.
Gardneriana, *Wedd. l. c.* 265.—Bras.
guayaquilensis, *Wedd. l. c.* 264.—Ecuador.
laurifolia, Cham. & Schlecht. in Linnaea, iii. (1828) 55 = Ruprechtia laurifolia.
Lindeniana, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 266.—N. Granat.
macrocalyx, *Casar. Nov. Stirp. Bras. Dec.* ix. 79 (sp. dub.).—Bras.
Martiana, *Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iv. (1845) 149.—Bras.
mollis, Walp. Ann. iii. 296 = Ruprechtia mollis.
Noli-tangere, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 264.—Bras.
Pachau, *Mart. Reise Bras.* iii. 1011, nomen; *et ex Meissn. in DC. Prod.* xiv. 177.—Bras.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 157; *Contrib.* 50.—Reg. Argent.
Pavonii, *Meissn. in DC. Prod.* xiv. 172.—Peruv.
peruviana, *Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iv. (1845) 149.—Peruv.
Poeppigiana, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 265.—Peruv.

TRIPLARIS:—
prognostica, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 285.—Guian.
Purdiei, *Meissn. in Mart. Fl. Bras.* v. I. 49.—N. Granat.
pyramidalis, Jacq. Select. Am. 13. t. 173. f. 5 (Quid ?).—N. Granat.
ramiflora, Jacq. l. c. 14 = Ruprechtia ramiflora.
ramiflora, Schott, ex Meissn. in Mart. Fl. Bras. v. I. 54 = Ruprechtia Lundii.
Riedeliana, *Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iv. (1845) 149.—Bras.
salicifolia, Cham. & Schlecht. in Linnaea, iii. (1828) 56 = Ruprechtia salicifolia.
salicifolia, Miq. ex Meissn. in Mart. Fl. Bras. v. I. 54 = Ruprechtia laurifolia.
scandens, Schott, ex Meissn. l. c. = Ruprechtia laurifolia.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 628.—Guian.
surinamensis, *Cham. in Linnaea*, viii. (1833) 138.—Guian.
tenuiflora, Endl. Gen. Suppl. iv. II. 55 = Ruprechtia tenuiflora.
tomentosa, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 265.—Bras.
Vahliana, *Fisch. & Mey. ex C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iv. (1845) 148.—Guian.
viridiflora, Schott, ex Meissn. in Mart. Fl. Bras. v. I. 54 = Ruprechtia Lundii.

TRIPLASANDRA, Seem. in Journ. Bot. vi. (1868) 139. *ARALIACEAE*.
oahuensis, *Seem. l. c.*—Ins. Sandvic.

TRIPLASIS, Beauv. Agrost. 81. t. 16. f. 10 (1812).
GRAMINEAE, Benth. & Hook. f. iii. 1176.
DIPLOCEA, Rafin. in Am. Journ. Sc. Ser. I. i. (1819) 252.
americana, *Beauv. Agrost.* 81. t. 16. f. 10.—Am. bor.
cornuta, Benth. in Journ. Linn. Soc. xix. (1881) 111 = americana.
purpurea, *Chapm. Fl. S. U. St.* 560.—Florida.
setacea, Griseb. in Goett. Abh. xxiv. (1879) 304 = Diplachne spicata.
sparsiflora, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 19.—Florida.

TRIPLATEIA, Bartl. in Presl, Rel. Haenk. ii. 11. t. 50 (1830) = **Hymenella**, Moç. & Sesse (Caryoph.).
diffusa, Bartl. l. c. = H. moehringioides.

TRIPLATHERA, Endl. Gen. 94 (1837) = **Bouteloua**, Lag. (Gramin.).

TRIPLECTRUM, D. Don, ex Wight & Arn. Prod. 324 (1834) = **Medinilla**, Gaudich. (Melastom.).
radicans, D. Don, ex Wight & Arn. l. c. = M. Beddomei.

TRIPLEURA, Lindl. in Wall. Cat. n. 7391 (1832); et Bot. Reg. sub t. 1618 (1833) = **Zeuxine**, Lindl. (Orchid.).
pallida, Lindl. ll. cc. = Z. sulcata.

TRIPLEUROSPERMUM, Sch. Bip. Tanacet. 31 (1844) = **Matricaria**, Tourn. (Compos.).
aromaticum, Rupr. ex Boiss. Fl. Orient. iii. 334 (= *Chamaemelum daghestanicum.*).—Reg. Casp.
bienne, Knaf, ex Nym. Consp. 374 = M. inodora.
caucasicum, J. Gay, ex Boiss. Fl. Orient. iii. 330 = M. conoclinia.
daghestanicum, Rupr. ex Boiss. l. c. 334 (= *Chamaemelum daghestanicum*).—Reg. Casp.
decipiens, Sch. Bip. ex Nym. Consp. 374 = M. decipiens.
disciforme, Sch. Bip. Tanacet. 34 = M. corymbifera.
fuscatum, Sch. Bip. in Bonplandia, viii. (1860) 369 = M. fuscata.
Hookeri, Sch. Bip. l. c. i. (1853) 151 = M. inodora.
inodorum, Sch. Bip. Tanacet. 32 = M. inodora.
maritimum, Koch, Syn. Fl. Germ. ed. II. 1026 = M. inodora.
nivale, Rupr. ex Boiss. Fl. Orient. iii. 334 (= *Chamaemelum daghestanicum*).—Reg. Casp.
uniglandulosum, Walp. Ann. i. 420 = M. trichophylla.

TRIPLIMA, Rafin. in Am. Monthly Mag. (1819) 195 = **Carex**, Dill. (Cyperac.).

TRIPLOBUS, Rafin. Sylva Tellur. 110 (1838) = **Sterculia**, Linn.
cordata, Rafin. l. c. 111 = S. cordata.

TRIPLOCENTRON, Cass. in Dict. Sc. Nat. xliv. 38 (1826) = **Centaurea**, Linn. (Compos.).
apulum, Cass. l. c. lv. 349 = C. melitensis.
melitense, Cass. l. c. xliv. 38 ; lv. 349 = C. melitensis.
solstitiale, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. solstitialis.
sulphureum, Webb, Iter Hisp. 33 = C. sulphurea.

TRIPLOCHITON, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 13 = **Hibiscus**, Linn. (Malvac.).
setosa, Alef. l. c. = H. panduraeformis.
spathacea, Alef. l. c. = H. vulpinus.

TRIPLOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 251 = **Streptocaulon**, Wight & Arn. (Asclep.).
Cumingii, Turcz. l. c. = S. Cumingii.

TRIPLOMEIA, Rafin. Sylva Tellur. 134 (1838) = **Acrodiclidium**, Nees (Laurin.).
montana, Rafin. l. c. 135 = A. jamaicense.

TRIPLORHIZA, Ehrh. Beitr. iv. (1789) 149 = **Habenaria**, Willd. (Orchid.).

TRIPLOSPERMA, G. Don, Gen. Syst. iv. 134 (1837) = **Ceropegia**?, Linn. (Asclep.).
cochinchinensis, G. Don, l. c.—Cochinch.

TRIPLOSTEGIA, Wall. ex DC. Prod. iv. 642 (1830). *DIPSACEAE*, Benth. & Hook. f. ii. 158.
glandulifera, *Wall. ex DC. l. c.* ; *et Cat.* n. 436.—Reg. Himal.
glandulosa, DC. Mém. Valér. 23. t. 5, sphalm. = praec.

TRIPODANTHERA, M. Roem. Syn. Pepon. 48 (1846) = **Gymnopetalum**, Arn. (Cucurb.).
cochinchinensis, M. Roem. l. c. = G. cochinchinense.

TRIPODION, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 348 = **Anthyllis**, Riv. (Legumin.).
lotoides, Medic. l. c. = A. tetraphylla.
tetraphyllum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 359 = A. tetraphylla.

TRIPOGON, Roth, Nov. Pl. Sp. 79 (1821). *GRAMINEAE*, Benth. & Hook. f. iii. 1169.
PLAGIOLYTRUM, Nees, in Proc. Linn. Soc. i. (1841) 95.
abyssinicus, *Nees, ex Steud. Syn. Pl. Gram.* 301.—Abyss.
bromoides, Benth. & Hook. f. Gen. iii. 1200 = abyssinicus.
bromoides, *Roth, Nov. Pl. Sp.* 79.—Ind. or.
capillatus, *Jaub. & Spach, Illustr.* iv. 47.—Ind. or.
exiguus, *Buese, in Miq. Pl. Jungh.* 353.—Java.
festucoides, Jaub. & Spach, Illustr. iv. 49 = bromoides.
filiformis, *Nees, ex Steud. Syn. Pl. Gram.* 301.—Ind. or.
Griffithii, Nees & Arn. ex Steud. l. c = bromoides.
lanatus, Hochst. ex Steud. l. c. = bromoides.
major, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 230. —Afr. trop.
minimus, Hochst. ex Steud. Syn. Pl. Gram. 301 = abyssinicus.
semitruncatus, Nees, ex Steud. l. c. = filiformis.
unidentatus, Nees, l. c. = filiformis.
zeylanicus, Nees, l. c. = bromoides.

TRIPOLION, Rafin. Fl. Tellur. ii. 46 (1836) = **Aster**, Tourn. (Compos.).
maritimum, Rafin. l. c. = A. Tripolium.
tataricum, Rafin. l. c. = A. tataricus.
tenellum, Rafin. l. c. = A. tenellus.

TRIPOLIUM, Nees, Gen. et Sp. Aster. 152 (1833) = **Aster**, Tourn. (Compos.).
andinum, Phil. in Linnaea, xxxiii. (1864-65) 129.—Chili.
angustum, Lindl. in Hook. Fl. Bor. Am. ii. 15 = A. angustus.
caricifolium, Schau. in Linnaea, xix. (1847) 721 = A. pauciflorus.
conspicuum, Lindl. ex DC. Prod. v. 254 = A. imbricatus, *Walp.* (cf. Addend.).
divaricatum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 296 = A. exilis.

TRIPOLIUM :—
flexuosum, Nees, Gen. et Sp. Aster. 155 = Aster tenuifolius.
frondosum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 296 = A. frondosus.
imbricatum, Nutt. l. c. 297 = A. imbricatus, *Walp.* (cf. Addend.).
longicaule, Duf. in Bull. Soc. Bot. Fr. vii. (1860) 327 = A. Tripolium.
maritimum, Léveillé, in Demid. Voy. Russ. Mér. ii. 193, 236, 237 = A. Tripolium.
occidentale, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 296 = A. occidentalis.
oregonum, Nutt. l. c. = A. oreganus.
paludosum, Nees, Gen. et Sp. Aster. 155 = A. paludosus.
pannonicum, Schur, Enum. Pl. Transs. 304 = A. Tripolium.
pauciflorum, Nees, Gen. et Sp. Aster. 157 = A. pauciflorus.
subulatum, Nees, l. c. 156 = A. subulatus.
uniflorum, Phil. in Linnaea, xxxiii. (1864-65) 129.—Chili.
vulgare, Nees, Gen. et Sp. Aster. 153 = A. Tripolium.

TRIPSACUM, Linn. Syst. ed. X. 1261 (1759). *GRAMINEAE*, Benth. & Hook. f. iii. 1113.
aegilopoides, Kunth, Enum. Pl. i. 602 (Index) = Rottboellia hirsuta.
aristatum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = Ischaemum aristatum.
ciliare, Rasp. l. c. = Elionurus tripsacoides.
compressum, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 466, in obs. = dactyloides.
compressum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = Hemarthria compressa.
cylindricum, Michx. Fl. Bor. Am. i. 60 = Rottboellia cylindrica.
dactyloides, *Linn. Syst.* ed. X. 1261.—Am. bor.
dactyloides, Schlecht. in Linnaea, vi. (1831) 40 = fasciculatum.
distachyum, Poir. Encyc. viii. 114 = Ischaemum rugosum.
distichum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = praec.
fasciculatum, Rasp. l. c. = Chloris radiata.
fasciculatum, *Trin. ex Steud. Nom.* ed. II. ii. 712.—Mexic.
giganteum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = Anthistiria gigantea.
granulare, Rasp. l. c. = Manisuris granularis.
hermaphroditum, Linn. Syst. ed. X. 1261 = Anthephora elegans.
hirsutum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = Rottboellia hirsuta.
lanceolatum, Rupr. ex Benth. Pl. Hartw. 347 = dactyloides.
monostachyon, Willd. Sp. Pl. iv. 202 ; Hort. Berol. 1. t. 1 = dactyloides.
mucronatum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = Trachys mucronata.
muticum, Rasp. l. c. = Ischaemum muticum.
Myuros, Rasp. l. c. = Rottboellia Myurus.
paniceum, Rasp. l. c. = Pogonatherum saccharoideum.
pubescens, Willd. ex Steud. Nom. ed. II. i. 317, in syn. ; ii. 712 = Cenchrus pubescens.

TRIPSILINA, Rafin. Fl. Tellur. iv. 103 (1836) = **Passiflora**, Linn.
fetida, Rafin. l. c. = P. foetida.

TRIPTERANTHUS, Wall. ex Miers, in Lindl. Veg. Kingd. 172 (1847) = **Burmannia**, Linn.

TRIPTERELLA, Michx. Fl. Bor. Am. i. 19. t. 3 (1803) = **Burmannia**, Linn.
alba, Schult. Mant. i. 358 = B. alba.
bicolor, Schult. l. c. 357 = B. bicolor.
caerulea, Muhl. Cat. ed. I. 4 ; et Ell. Sketch. i. 43 = B. biflora.
capitata, Michx. Fl. Bor. Am. i. 19. t. 3 = B. capitata.
dasyantha, Schult. Mant. i. 357 = B. dasyantha.
flava, Schult. l. c. = B. flava.

TRIPTERIS, Less. in Linnaea, vi. (1831) 95. *COMPOSITAE*, Benth. & Hook. f. ii. 455.
aghillana, *DC. Prod.* vi. 457.—Afr. austr.
amplectens, Harv. in Harv. & Sond. Fl. Cap. iii. 427 = arborescens.

TRIPTERIS :—
amplexicaulis, DC. Prod. vi. 457 = arborescens.
amplexicaulis, *Less. in Linnaea*, vi. (1831) 97.—Afr. austr.
arborescens, *Less. l. c.* 96.—Afr. austr.
atropurpurea, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 212 = clandestina.
Burchellii, *Hook. f. in Melliss, St. Helena*, 290. t. 45. —Ins. S. Helena.
cheiranthifolia, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) I. Intell. 26, nomen = Vaillantii.
clandestina, *Less. in Linnaea*, vi. (1831) 97.—Afr. austr.
dentata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 427.— Afr. austr.
flexuosa, *Harv. l. c.* 429.—Afr. austr.
gariepina, Sond. ex Harv. & Sond. l. c. 432 = pachypteris.
glabrata, Harv. l. c. 425 = macroptera.
glandulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 93.—Afr. austr.
herbacea, E. Mey. ex DC. Prod. vi. 458 = clandestina.
humilis, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 211.—Afr. austr.
hyoseroides, DC. Prod. vi. 458 = clandestina.
incana, *Harv. in Harv. & Sond. Fl. Cap.* iii. 426.— Afr. austr.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 767 = setifera?
leptoloba, *Harv. in Harv. & Sond. Fl. Cap.* iii. 432.— Afr. austr.
limonifolia, *DC. Prod.* vi. 458.—Afr. austr.
linearis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 426.— Afr. austr.
Lordii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 424.—Afr. trop.
macroptera, *DC. Prod.* vi. 456.—Afr. austr.
microcarpa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 427. —Afr. austr.
monocephala, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 424.—Afr. trop.
natalensis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 430. —Afr. austr.
oppositifolia, E. Mey. ex DC. Prod. vi. 456 = sinuata.
pachypteris, *Harv. in Harv. & Sond. Fl. Cap.* iii. 432. —Afr. austr.
pallescens, *DC. Prod.* vi. 456.—Afr. austr.
petiolata, *DC. l. c.* 457.—Afr. austr.
polycephala, *DC. l. c.* 458.—Afr. austr.
rigida, *Harv. in Harv. & Sond. Fl. Cap.* iii. 426.— Afr. austr.
scariosa, DC. Prod. vi. 458 = clandestina.
setifera, *DC. l. c.* 457.—Afr. austr.
sinuata, *DC. l. c.* 456.—Afr. austr.
spathulata, *DC. l. c.* 457.—Afr. austr.
spinescens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 431. —Afr. austr.
Thomii, Harv. l. c. 426 = macroptera.
tomentosa, *Less. Syn. Comp.* 90, in obs.—Afr. austr.
Vaillantii, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 260.—Arab.

TRIPTERIS, Thunb. Pl. Bras. Dec. i. 14 (1817), sphalm. = **Triopteris**, Linn. (Malpig.).

TRIPTERIUM, Bercht. & Presl, Rostl. i. Ranunculac. 14 (1823) = **Thalictrum**, Tourn. (Ranunc.).
alpestre, Schur, Enum. Pl. Transs. 974 (Index) = Th. aquilegifolium.
aquilegifolium, Bercht. & Presl, Rostl. i. Ranunculac. 14 = Th. aquilegifolium.
calabricum, Bercht. & Presl, l. c. = Th. calabricum.
contortum, Bercht. & Presl, l. c. = Th. aquilegifolium.
pauciflorum, Schur, Enum. Pl. Transs. 7 = Th. aquilegifolium.
transsilvanicum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 10 = Th. minus.

TRIPTEROCALYX, Hook. Kew Journ. v. (1853) 261 = **Abronia**, Juss. (Nyctag.).
macrantha, Torr. ex Hook. l. c. = A. Cycloptera.

TRIPTEROCARPUS, Meissn. Gen. 52 (1837) = **Bridgesia**, Bert. (Sapind.).
incisifolia, Meissn. Comm. 37 = B. incisifolia.

TRIPTEROCOCCUS, Endl. in Enum. Pl. Hueg. 17 (1837) = **Stackhousia**, Sm.
brachystigma, Schuch. in Linnaea, xxvi. (1853) 33 = S. Brunonis.
Brunonis, Endl. in Enum. Pl. Hueg. 18 = S. Brunonis.

TRIPTEROCOCCUS:—
junceus, Bunge, in Lehm. Pl. Preiss. i. 181 = Stackhousia Brunonis.
simplex, Bunge, l. c. = S. Brunonis.
spathulatus, F. Muell. ex Schuch. in Linnaea, xxvi. (1853) 29 = S. spathulata.

TRIPTEROSPERMUM, Blume, Bijdr. 849 (1826) = **Crawfurdia**, Wall. (Gentian.).
caucasicum, Owerin, ex Trautv. in Act. Hort. Petrop. viii. (1883) 454, sphalm. = *Tripleurospermum caucasicum*?
Championii, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 15 = C. japonica.
trinerve, Blume, Bijdr. 849 = C. Blumei.

TRIPTERYGIUM, Hook. f. in Benth. & Hook. f. Gen. i. 368 (1862). *CELASTRINEAE*, Benth. & Hook. f. l. c.
Bullockii, Hance, in Journ. Bot. xviii. (1880) 259 = Wilfordii.
Wilfordii, *Hook. f. in Benth. & Hook. f. Gen.* i. 368. —China; Japon.

TRIPTILION, Ruiz & Pav. Prod. 102. t. 22 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 503.
Achilleae, *DC. Prod.* vii. 51.—Chili.
andinum, *Phil. in Linnaea*, xxxiii. (1864–65) 116.—Chili.
axillare, Lag. ex Lindl. in Bot. Reg. sub t. 853 = Nassauvia axillaris.
Benaventii, *Remy, in C. Gay, Fl. Chil.* iii. 357. t. 39. —Chili.
bulbosum, *Remy, l. c.* 358.—Chili.
capillatum, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 43; *DC. Prod.* vii. 51.—Chili.
chilense, Bert. ex DC. l. c. = Achilleae.
cordifolium, Lag. in Bot. Reg. t. 853.—Chili.
diffusum, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 221.—Chili.
euphrasioides, *Bert. ex DC. Prod.* vii. 51.—Chili.
gibbosum, *Remy, in C. Gay, Fl. Chil.* iii. 356.—Chili.
glomerulosum, Lag. ex Lindl. in Bot. Reg. sub t. 853 = Nassauvia glomerulosa.
laciniatum, Willd. Sp. Pl. iii. 1626 = spinosum.
laxum, *Phil. in Linnaea*, xxxiii. (1864–65) 116.—Chili.
maritimum, Poepp. ex DC. Prod. vii. 51 = cordifolium.
pectinatum, *Phil. in Anal. Univ. Chil.* (1856) ii. 329. —Chili.
spinosum, *Ruiz & Pav. Syst. Veg.* i. 185.—Chili.
tenuifolium, *Phil. in Linnaea*, xxxiii. (1864–65) 117.—Chili.

TRIPTILIUM, DC. in Ann. Mus. Par. xix. (1812) 67 = praec.

TRIPTILODISCUS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 66 = **Helipterum**, DC. (Compos.).
pygmaeus, Turcz. l. c. = H. Dimorpholepis.

TRIPTOLEMAEA, Walp. Rep. i. 800 (1842), sphalm. = seq.

TRIPTOLEMEA, Mart. in Flora, xx. (1837) ii. Beibl. 122 = **Dalbergia**, Linn. f. (Legumin.).
glabra, Benth. in Ann. Wien. Mus. ii. (1838) 103 = D. variabilis.
latifolia, Benth. l. c. = D. variabilis.
montana, Mart. in Flora, xx. (1837) ii. Beibl. 122 = D. variabilis.
myriantha, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 102 = D. variabilis.
ovata, Benth. l. c. = D. variabilis.
pauciflora, Mart. ex Benth. l. c. = D. variabilis.
platycarpa, Benth. l. c. = D. variabilis.
pubescens, Benth. in Hook. Journ. Bot. ii. (1840) 68 = D. variabilis.
riparia, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 103 = D. riparia.

TRIQUETRA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 376 = **Astragalus**, Tourn. (Legumin.).
boetica, Medic. l. c. = A. baeticus.

TRIRAPHIS, R. Br. Prod. 185 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1177.
bromoides, *F. Muell. Fragm.* viii. 108.—Austral.
danthonioides, *F. Muell. l. c.* 125.—Austral.
diantha, F. Muell. l. c. = pungens?

TRIRAPHIS:—
microdon, *Benth. Fl. Austral.* vii. 605 (sp. dub.).—Austral.
mollis, *R. Br. Prod.* 185.—Austral.
Pumilio, *R. Br. in Denh. & Clapp. Trav. App.* 245.—Afr. trop.
pungens, *R. Br. Prod.* 185.—Austral.

TRIRAPHIS, Nees, Fl. Afr. Austr. 270 (1841) = **Danthonia**, DC. (Gramin.).
capensis, Nees, l. c. 271 = Trisetum insulare.

TRISACARPIS, Rafin. Fl. Tellur. iv. 11 (1836) = **Hippeastrum**, Herb. (Amaryll.).
falcata, Rafin. l. c. = H. calyptratum.
psittacina, Rafin. l. c. = H. psittacinum.
rubra, Rafin. l. c. = H. aulicum.

TRISANTHUS, Lour. Fl. Cochinch. 175 (1790) = **Hydrocotyle**, Tourn. (Umbell.).
cochinchinensis, Lour. l. c. 176 = H. asiatica.

TRISCENIA, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 534. *GRAMINEAE*, Benth. & Hook. f. iii. 1119.
ovina, *Griseb. l. c.*—Cuba.

TRISCHIDIUM, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 141. t. 4 = **Swartzia**, Schreb. (Legumin.).
vestitum, Tul. l. c. 142 = S. mollis.

TRISCIADIA, Hook. f. in Benth. & Hook. f. Gen. ii. 68 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
truncata, *Hook. f. l. c.* 69.—Ins. Penang.

TRISCIADIUM, Phil. in Anal. Univ. Chil. (1861) i. 61 = **Huanaca**, Cav. (Umbell.).
andinum, Phil. l. c. = H. andina.

TRISEMA, Hook. f. in Hook. Kew Journ. ix. (1857) 47. t. 1. *DILLENIACEAE*, Benth. & Hook. f. i. 14.
Trisemma, Panch. & Sebert, Not. Bois Nouv. Caléd. 211 (1874).
Vanieria, Montr. in Mém. Acad. Lyon, x. (1860) 176.
coriacea, *Hook. f. in Hook. Kew Journ.* ix. (1857) 47. —N. Caled.
Pancheri, Panch. & Sebert, Not. Bois Nouv. Caléd. 211 = coriacea.
Vieillardi, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 191; *et in Ann. Sc. Nat.* Sér. V. ii. (1864) 150.—N. Caled.

TRISEMMA, Panch. & Sebert, Not. Bois Nouv. Caléd. 211 (1874), sphalm. = praec.

TRISEPALUM, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 363 (1884). *GESNERACEAE*.
acutum, *C. B. Clarke, l. c.* 364.—Burma.
obtusum, *C. B. Clarke, l. c.* 363.—Burma.

TRISETARIA, Forsk. Fl. Aegypt. Arab. 27 (1775). *GRAMINEAE*, Benth. & Hook. f. iii. 1154.
Anomalotis, Steud. Syn. Pl. Gram. 198 (1855).
airoides, Baumg. Enum. Stirp. Transs. iii. 265 = Trisetum subspicatum.
alpestris, Baumg. l. c. 264 = Trisetum alpestre.
brevifolia, Baumg. Mant. 80 (1846) = Trisetum distichophyllum.
carpatica, Baumg. Enum. Stirp. Transs. iii. 263 = Trisetum carpathicum.
flavescens, Baumg. l. c. = Trisetum pratense.
Forskåhlii, J. F. Gmel. Syst. 177 = linearis.
linearis, *Forsk. Fl. Aegypt. Arab.* p. lx.—Aegypt.; Arab.
pratensis, Baumg. Enum. Stirp. Transs. iii. 264 = Avena pratensis.
pubescens, Schur, Enum. Pl. Transs. 974 (Index), sphalm. = Avena pubescens.
quinqueseta, *Hochst. in Flora*, xxxiii. (1855) 282.—Abyss.
sesquitertia, Baumg. Enum. Stirp. Transs. iii. 262 = Avena pubescens.
tenuis, Baumg. *l. c.* = Ventenata avenacea.

TRISETARIUM, Poir. Encyc. Suppl. v. 365 (1817), sphalm. = seq.

TRISETUM, Pers. Syn. i. 97 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1159.
Acrospelion & Acropselion, Bess. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 59.
Rebentischia, Opiz, in Lotos, iv. (1854) 104.
Rostraria, Trin. Fund. Agrost. 149 (1820).
Trichaeta, Beauv. Agrost. 86. t. 17. f. 8 (1812).
Trisetarium, Poir. Encyc. Suppl. v. 365 (1817).
agrostideum, *Fries, Mant.* iii. 180.—Europ.
agrostoides, Steud. Syn. Pl. Gram. 226 = myrianthum.
airaeforme, Steud. l. c. 229 = Deschampsia airaeformis.
airoides, Roem. & Schult. Syst. ii. 666 = subspicatum.
Alopecuros, *Roem. & Schult. Syst.* ii. 660.—Hab.?
alpestre, *Beauv. Agrost.* 88.—Europ. med.
alpinum, Roem. & Schult. Syst. ii. 663 = Avena pratensis.
andinum, Benth. Pl. Hartw. 261 = subspicatum.
andinum, *Phil. in Linnaea*, xxix. (1857–58) 93.—Chili.
andropogonoides, Steud. Syn. Pl. Gram. 228 = Avena andropogonoides.
anomalum, *Steud. l. c.* 425.—Madag.
antarcticum, Nees, in Linnaea, vii. (1832) 807 = Danthonia antarctica.
antarcticum, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 61.—N. Zel.; Chile.
arenarium, *Labill. Ic. Pl. Syr. Dec.* v. 10.—Syria; Aegypt.
argenteum, *Roem. & Schult. Syst.* ii. 665.—Europ. med.
argenteum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 77 = distichophyllum.
argentoideum, Schur, l. c. 74 = alpestre.
aristidoides, Pers. Syn. i. 98 = Avena aristidoides.
aureum, *Nees, ex Steud. Syn. Pl. Gram.* 225.—Ind. or.
aureum, *Tenore, Fl. Nap.* ii. 378.—Ital.
bambusiforme, *Fourn. ex Hemsl. in Biol. Centr. Am. Bot.* iii. 555.—Mexic.
barbatum, Nees, Fl. Afr. Austr. 345 = Dregeanum.
barbatum, *Steud. Nom.* ed. II. ii. 713; *Syn. Pl. Gram.* 229.—Chili.
barbinode, *Trin. in Linnaea*, x. (1836) 300.—Chili.
baregense, Laff. & Miégev. in Bull. Soc. Bot. Fr. xxi. (1874) 46 = agrostideum.
Berteroanum, Kunth, Rév. Gram. ii. 457. t. 142 = Deschampsia Berteroniana.
Berteronianum, Steud. Syn. Pl. Gram. 220 = Aira Gayana.
biflorum, *Hochst. in Flora*, xxxviii. (1855) 275.—Abyss.
biflorum, *Phil. in Anal. Univ. Chil.* (1873) 568.—Chili.
brachyatherum, *Phil. l. c.* 567.—Chili.
Brandegei, *Scribner, in Bull. Torrey Bot. Club*, x. (1883) 64.—Am. bor. occ.
brevifolium, Roem. & Schult. Syst. ii. 665 = distichophyllum.
Bungei, *Boiss. Fl. Orient.* v. 535.—Persia.
Burnoufii, *Req. ex Parl. Fl. Ital.* i. 263.—Ins. Corsic.
canariense, Parl. ex Webb & Berth. Phyt. Canar. iii. 399 = paniceum.
canescens, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 100 = cernuum.
carpathicum, *Roem. & Schult. Syst.* ii. 663.—Transsylv.
caudulatum, *Trin. in Linnaea*, x. (1836) 300.—Chili.
Cavanillesii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 63.—Europ. austr.; Oriens; Ind. bor. occ.
cernuum, *Trin. l. c.* 61.—Ins. Sitcha.
chiloënse, *Phil. in Linnaea*, xxix. (1857–58) 93.—Ins. Chiloë.
chloranthum, Steud. Syn. Pl. Gram. 225 = aureum, *Tenore*.
chromostachyum, *E. Desv. in C. Gay, Fl. Chil.* vi. 350.—Chili.
compressum, Fuss, Fl. Transs. 727 = Avena compressa.
condensatum, Presl, in Schult. Mant. ii. 366 = aureum, *Tenore*.
depauperatum, *Phil. in Linnaea*, xxxiii. (1864–65) 291.—Chili.
deyeuxioides, Kunth, Rév. Gram. i. 102 = Deyeuxia triflora.
distichophyllum, *Beauv. Agrost.* 88.—Europ. austr.
distichophyllum, C. A. Mey. Verz. Pfl. Cauc. 17 = rigidum.
Dregeanum, *Steud. Syn. Pl. Gram.* 227.—Afr. austr.

TRISETUM:—
Dufourei, *Boiss. & Reut. Pugill. Pl. Nov.* 122.—Hispan.
elatum, Nutt. ex A. Gray, in Proc. Acad. Sc. Philad. 1862 (1863) 337 = cernuum.
elongatum, *Kunth, Rév. Gram.* i. 101.—Mexic.
flavescens, Beauv. Agrost. 88 = pratense.
Forskălii, Beauv. l. c. = Danthonia Forskåhlei.
fraudulentum, *Steud. Syn. Pl. Gram.* 424.—Reg. Magellan.
Friesianum, Steud. l. c. 225 = agrostideum.
fuscescens, *Pomel, Nouv. Mat. Fl. Atl.* 387.—Alger.
fuscum, Roem. & Schult. Syst. ii. 664 = carpathicum.
gallecicum, *Willk. & Lange, Prod. Fl. Hisp.* i. 72.—Hispan.
Gaudinianum, Boiss. Voy. Bot. Espagne, ii. 653 = Cavanillesii.
glabratum, Schult. Mant. ii. 369 = Aira glabra.
glabrum, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 100 = Danthonia calycina.
glabrum, Dulac, Fl. Hautes-Pyr. 85 = agrostideum.
glaciale, *Boiss. Elench.* 87.—Hispan.
glomeratum, *Trin. ex Steud. Nom.* ed. II. ii. 714; *Syn. Pl. Gram.* 229.—Ins. Sandvic.
glumaceum, *Boiss. Diagn.* Ser. I. xiii. 49.—Aegypt.; Arab.
Gmelini, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 66.—Hispan.
gracile, Boiss. Voy. Bot. Espagne, ii. 654, in textu = Morisii.
gracile, *Fourn. ex Hemsl. in Biol. Centr. Am. Bot.* iii. 556.—Mexic.
groenlandicum, *Steud. Syn. Pl. Gram.* 228.—Groenland.
Hallii, *Scribner, in Bull. Torrey Bot. Club*, xi. (1884) 6.—Texas.
heteronymum, *Steud. Syn. Pl. Gram.* 229.—Chili.
hirsutum, *Phil. in Anal. Univ. Chil.* (1873) 565.—Chili.
hirtulum, Steud. Syn. Pl. Gram. 228 = Avena hirta.
hirtum, Nees, Fl. Afr. Austr. 350 = Avena hirta.
hirtum, *Trin. in Linnaea*, x. (1836) 300.—Chili.
hispanicum, *Pers. Syn.* i. 97.—Hispan.
hispidum, *Lange, in Kjoeb. Vidensk. Meddel.* (1860) 42.—Hispan.
hispidum, Trin. ex Steud. Nom. ed. II. i. 849, in syn.; ii. 714 = Koeleria hispida.
Hohenackeri, Presl, Bot. Bemerk. 120 = Gaudinia fragilis.
imberbe, Nees, Fl. Afr. Austr. 347 = Danthonia antarctica.
insulare, *Hemsl. Bot. Voy. Challenger*, i. II. 267.—Ins. St. Paul (Mare Ind.).
interruptum, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 100 = elongatum.
interruptum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 556.—Mexic.
labradoricum, *Steud. Syn. Pl. Gram.* 228.—Labrador.
lachnanthum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 416 = Avena lachnantha.
laconicum, *Boiss. & Orph. Diagn.* Ser. II. iv. 129.—Laconia.
lasiolepis, *E. Desv. in C. Gay, Fl. Chil.* vi. 346.—Chili.
laxiflorum, *Phil. in Anal. Univ. Chil.* (1873) 566.—Chili.
laxum, *Phil. l. c.* 568.—Chili.
leoninum, Steud. Syn. Pl. Gram. 228 = Danthonia antarctica?
lineare, Boiss. Diagn. Ser. I. xiii. 49 = arenarium.
litorale, *Phil. in Linnaea*, xxix. (1857–58) 92.—Chili.
lobatum, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 66 = Eatonia obtusata.
Loefflingii, Beauv. Agrost. 88, 180 = hispanicum.
Loeflingianum, Presl, Cyp. et Gram. Sic. 30 = aureum, *Tenore*.
longiaristum, *A. Rich. Tent. Fl. Abyss.* ii. 417.—Abyss.
longifolium, *Nees, Fl. Afr. Austr.* 348.—Afr. austr.
ludovicianum, *Vasey, in Bull. Torrey Bot. Club*, xii. (1885) 6.—Am. bor.
lusitanicum, Pers. Syn. i. 97 = paniceum.
luteum, Pers. l. c. = Avena lutea.
luzonense, Steud. Nom. ed. II. i. 228 = Triticum luzonense.
macilentum, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 66 = Avellinia Michelii.
macrochaetum, *Boiss. Diagn.* Ser. I. xiii. 48.—Palaestin.

TRISETUM:—
macrum, Beauv. Agrost. 88 = Ventenata macra.
malacanthum, *Steud. Syn. Pl. Gram.* 424.—Chili.
malacophyllum, *Phil. in Anal. Univ. Chil.* (1873) 566.—Chili.
malacophyllum, *Steud. Nom.* ed. II. ii. 714; *Syn. Pl. Gram.* 229.—Chili.
melitense, Steud. Syn. Pl. Gram. 225 = aureum, *Tenore*.
Michellii, Guss. Enum. Pl. Inar. 386 = Avellinia Michelii, Bromus festucoides.
micranthum, *Roem. & Schult. Syst.* ii. 659.—Europ. austr.
micratherum, E. Desv. in C. Gay, Fl. Chil. vi. 352 = Koeleria micrathera.
Miegevillii, Duval-Jouve, ex Miégev. in Bull. Soc. Bot. Fr. xxi. (1874) 47 = agrostideum.
molle, Kunth, Rév. Gram. i. 101 = subspicatum.
molle, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 64, partim = toluccense.
monticola, *Phil. in Linnaea*, xxxiii. (1864–65) 291.—Chili.
Morisii, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 66.—Ins. Sardoa.
muricatum, Spreng. Syst. i. 333, in syn. = Danthonia papillosa.
myrianthum, *C. A. Mey. in Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. 95.—Ital.; Graecia.
nanum, Steud. Syn. Pl. Gram. 227 = Diplachne nana.
Neesii, Hochst. ex Steud. l. c. = Avena Neesii.
neglectum, Lange, in Kjoeb. Vidensk. Meddel. (1860) 41 = gallecicum.
neglectum, Presl, Cyp. et Gram. Sic. 29 = aureum, *Tenore*.
neglectum, Roem. & Schult. Syst. ii. 660 = paniceum.
nitidum, *Pers. Syn.* i. 97.—Ins. Sardoa; Afr. bor.
nivosum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 556.—Mexic.
Noëanum, Boiss. Diagn. Ser. I. xiii. 48 = aureum, *Tenore*.
nudum, Pers. Syn. i. 97 = Danthonia tortuosa.
ochrostachyum, *Phil. in Linnaea*, xxxiii. (1864–65) 290.—Chili.
ovatum, *Pers. Syn.* i. 98.—Hispan.
pallidum, Roem. & Schult. Syst. ii. 657 = Danthonia tortuosa.
palustre, *Torr. Fl. U. St.* 126.—Am. bor.
paniceum, *Pers. Syn.* i. 97.—Europ.
paniculatum, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 556.—Mexic.
pappophoroides, *Steud. in Lechl. Berb. Am. Austr.* 52, nomen.—Chili.
parviflorum, Herb. Bout. ex Willk. & Lange, Prod. Fl. Hisp. i. 72 = paniceum.
parviflorum, *Pers. Syn.* i. 97.—Ital.; Afr. bor.
pensylvanicum, Roem. & Schult. Syst. ii. 658 = Avena pensylvanica.
pensylvanicum, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 66 = Eatonia obtusata.
phleoides, Kunth, Rév. Gram. i. 101 = subspicatum.
phleoides, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 65 = Koeleria phleoides.
pilosum, Roem. & Schult. Syst. ii. 662 = Avena pilosa.
Pourreti, *Roem. & Schult. l. c.* 661.—Hispan.
praecox, Dum. Obs. Gram. Belg. 122 = Aira praecox.
pratense, Dum. l. c. = Avena pratensis.
pratense, *Pers. Syn.* i. 97.—Reg. bor. temp.
Preslei, E. Desv. in C. Gay, Fl. Chil. vi. 347 = Avena Preslii.
puberulum, Tenore, Fl. Nap. iv. 17 = Avellinia Michelii.
pubescens, Roem. & Schult. Syst. ii. 662 = Avena pubescens.
pubescens, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 65 = Koeleria villosa.
pumilum, *Kunth, Rév. Gram.* i. 102.—Reg. Mediterr.; Oriens.
pungens, *Beauv. Agrost.* 88.—Hab.?
purpurascens, Torr. Fl. U. St. 127 = Avena striata, *Michx.*
rigidum, *Roem. & Schult. Syst.* ii. 662.—Reg. Cauc.; As. Min.
Rohlfsii, *Aschers. in Verh. Bot. Ver. Brandenb.* xxi. (1880) 71.—Aegypt.
Ruprechti, Steud. Syn. Pl. Gram. 226 = Avena Ruprechtii.
scabriusculum, Coss. Notes Crit. 129 = Koeleria scabriuscula.

TRISETUM:—
sericeum, *Desv. Opusc.* 100.—Bras.
sesquiflorum, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 66 = Avena sesquiflora.
sesquiterium, Beauv. Agrost. 88 = Avena pubescens.
sesquitertium, Schur, Enum. Pl. Transs. 760 = praec.
sibiricum, Rupr. in Beitr. Pfl. Russ. Reich. ii. 65 = Avena Ruprechtii.
smyrnaeum, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 66 = Koeleria biseta.
spiciforme, Dulac, Fl. Hautes-Pyr. 86 = subspicatum.
splendens, Presl, Cyp. et Gram. Sic. 30 = pratense.
splendidulum, *Steud. Syn. Pl. Gram.* 229.—Chili.
Steudelii, *Nees, in Linnaea*, vii. (1832) 308.—Afr. austr.
striatum, Pers. Syn. i. 97 = Ventenata avenacea.
subaristatum, *E. Desv. in C. Gay, Fl. Chil.* vi. 353.—Chili.
subspicatum, *Beauv. Agrost.* 88.—Reg. frigid. bor. et austr.
tenue, Baumg. ex Steud. Syn. Pl. Gram. 226 = carpathicum.
tenue, Roem. & Schult. Syst. ii. 657 = Ventenata avenacea.
Thunbergii, E. Desv. in C. Gay, Fl. Chil. vi. 350, in nota = Danthonia antarctica.
toluccense, *Kunth, Rév. Gram.* i. 101.—Mexic.
transsilvanicum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 77 = pratense.
transylvanicum, Steud. Syn. Pl. Gram. 226 = carpathicum.
Trisetaria, Roem. & Schult. Syst. ii. 658 = arenarium.
umbrosum, *Hochst. ex Steud. Syn. Pl. Gram.* 227.—Abyss.
uniflorum, Meyen, Reise, i. 484 = Agrostis Preslii.
valesiacum, Boiss. ex Nym. Consp. 814 = Cavanillesii.
variabile, *E. Desv. in C. Gay, Fl. Chil.* vi. 351.—Chili.
varium, Schur, in Oestr. Bot. Zeitschr. x. (1860) 75 = pratense.
velutinum, *Boiss. Elench.* 88.—Hispan.
villosum, Pers. Syn. i. 97 = Danthonia Thunbergii.
villosum, *Schult. Mant.* ii. 368.—Europ.
villosum, Sieber, ex Steud. Nom. ed. II. ii. 714 = Danthonia tortuosa.
virescens, *Nees, ex Steud. Syn. Pl. Gram.* 226.—Reg. Himal.
viride, Kunth, Rév. Gram. i. 101 = Deyeuxia viridis, *Fourn.*
Virletii, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 556.—Mexic.
Wilhelmsii, Ledeb. ex Schult. Mant. ii. 367 = rigidum.
Wolfii, *Vasey, in Rep. U. St. Dept. Agric.* 1874 (1875), 156 (an seq.?).—Am. bor.
Wolfii, Vasey, in Rothr. Bot. Rep. U. St. Geogr. Surv. vi. 294 = Graphephorum Wolfii.
Youngii, *Hook. f. Handb. N. Zeal. Fl.* 335.—N. Zel.

TRISIOLA, Rafin. Fl. Ludov. 144 (1817) = **Uniola**, Linn. (Gramin.).
paniculata, Rafin. l. c. = U. paniculata.

TRISSAGO, Linn. Syst. ed. I (1735) = **Teucrium**?, Tourn. (Labiat.).

TRISTACHYA, Nees, Agrost. Bras. 458 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1161.
MONOPOGON, J. Presl, Rel. Haenk. i. 324. t. 44 (1830).
arundinacea, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 417.—Abyss.
barbata, *Nees, Fl. Afr. Austr.* 269; *et ex Steud. Nom.* ed. II. ii. 714.—Arab.; Afghan.
chrysothrix, *Nees, Agrost. Bras.* 460.—Bras.
elegans, *A. Rich. Tent. Fl. Abyss.* ii. 418.—Abyss.
leiostachya, *Nees, Agrost. Bras.* 459.—Am. trop.
leucothrix, *Trin. ex Nees, l. c.* 460.—Afr. austr.
mexicana, Kunth, Enum. Pl. i. 308 = leiostachya.
microstachya, *Nees, ex Steud. Syn. Pl. Gram.* 238.—Peruv.
monocephala, *Hochst. in Flora*, xxix. (1846) 120.—Afr. austr.
simplex, Nees, Fl. Afr. Austr. 269 = Trichopteryx simplex.
somalensis, *Franch. in Révoil, Çomali*, 67.—Afr. trop.
Stocksii, Boiss. Fl. Orient. v. 552 = barbata.

TRISTAGMA, Poepp. Fragm. Syn. Pl. Chile, 8 (1833); Poepp. & Endl. Nov. Gen. et Sp. ii. 28. t. 140 (1838). *LILIACEAE*, Benth. & Hook. f. iii. 798.
STEMMATIUM, Phil. in Anal. Univ. Chil. (1873) 551.
STEPHANOLIRION, Baker, in Gard. Chron. (1875) i. 234.
dimorphopetalum, *C. Gay, Fl. Chil.* vi. 126.—Chili.
narcissoides, *Benth. & Hook. f. Gen.* iii. 799.—Chili.
nivale, *Poepp. Fragm. Syn. Pl. Chile*, 9; *Poepp. & Endl. Nov. Gen. et Sp.* ii. 28. t. 140.—Chili.

TRISTANIA, R. Br. in Ait. Hort. Kew. ed. II. iv. 417 (1812). *MYRTACEAE*, Benth. & Hook. f. i. 708.
CALLOBUXUS, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 372.
LOPHOSTEMON, Schott, in Wien. Zeitschr. iii. (1830) 772; ex Endl. in Linnaea, vi. (1831) Litt. 54.
TRISTANIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 371.
albens, Hort. Berol. ex DC. Prod. iii. 210; et ex A. Cunn. in Bot. Reg. sub t. 1839 = Syncarpia laurifolia.
angustifolia, Hook. in Mitch. Journ. Trop. Austral. 198 = Lysicarpus ternifolius.
arborescens, *Sweet, Hort. Brit.* ed. I. 156, nomen.—Austral.
burmannica, *Griff. Cantor Pl.* 17.—Burma; Malaya.
capitulata, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* x. (1863) 372.—N. Caled.
conferta, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 417.—Austral.
conferta, Griff. Notul. iv. 649 = Griffithii.
corymbosa, Hort. ex Steud. Nom. ed. II. i. 797, in syn.; ii. 714 = Eugenia myrifolia, *Sims*.
densiflora, *Carr. in Rev. Hortic.* (1881) 420.—Austral.
depressa, Hort. Angl. ex Link, Enum. Hort. Berol. ii. 273, nomen; et A. Cunn. in Bot. Reg. sub t. 1839 = suaveolens.
exiliflora, *F. Muell. Fragm.* v. 11.—Austral.
glauca, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xiii. (1866) 471, in syn.—N. Caled.
Griffithii, *Kurz, in Pegu Rep. App.* B. 50.—Burma.
insularis, *Vieill. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 300, in syn.—N. Caled.
lactiflua, *F. Muell. Fragm.* i. 82.—Austral.
laurina, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 417.—Austral.
macrophylla, A. Cunn. in Bot. Reg. t. 1839 = conferta.
macrosperma, *F. Muell. Papuan Pl.* i. 104.—N. Guin.
Maingayi, *Duthie, in Hook. f. Fl. Brit. Ind.* ii. 467.—Malaya.
merguensis, *Griff. Cantor Pl.* 18.—Burma.
myrtifolia, Ait. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 651, nomen (Quid?).—Austral.
nereifolia, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 417.—Austral.
obovata, *R. Br. in Benn. Pl. Jav. Rar.* 127. t. 27.—Ins. Banca.
persicaefolia, *A. Cunn. in Field, N. S. Wales*, 350.—Austral.
psidioides, *A. Cunn. in Bot. Reg.* sub t. 1839.—Austral.
rhytiphloia, F. Muell. Fragm. i. 81 = suaveolens.
rufescens, *Hance, in Journ. Bot.* xiv. (1876) 259.—Cambodia.
salicifolia, Link, ex Steud. Nom. ed. II. ii. 714 = suaveolens.
salicina, A. Cunn. in Bot. Reg. sub t. 1839 = nereifolia.
suaveolens, *Sm. in Rees, Cycl.* xxxvi. n. 2.—Austral.
subverticillata, H. Wendl. in Otto & Dietr. Allg. Gartenz. i. (1833) 186 = conferta.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 308 = Whitiana.
umbrosa, *A. Cunn. in Bot. Reg.* sub t. 1839.—Austral.
Whitiana, *Griff. Cantor Pl.* 18.—Malaya.
Wightiana, Duthie, in Hook. f. Fl. Brit. Ind. ii. 466 sphalm. = praec.

TRISTANIA, Poir. Encyc. Suppl. iv. 526, in obs. (1816) = **Spartina**, Schreb. (Gramin.).
arundinacea, Poir. l. c. = S. arundinacea.

TRISTANIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 371 = **Tristania**, R. Br. (Myrtac.).
Callobuxus, Brongn. & Gris, l. c. 372.—N. Caled.

TRISTANIOPSIS:—
capitulata, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 372 = Tristania capitulata.
glauca, Brongn. & Gris, l. c. xiii. (1866) 471 = Tristania glauca.
Guillainii, Vieill. ex Brongn. & Gris, in Ann. Sc. Nat. Sér. V. xiii. (1870–71) 348.—N. Caled.
Vieillardi, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 300 = Tristania insularis.

TRISTEGIS, Nees, Hor. Phys. Berol. 47, 54 (1820) = **Melinis**, Beauv. (Gramin.).
glutinosa, Nees, l. c. = M. minutiflora.

TRISTELLARIA, Reichb. Consp. 207 (1828) = seq.

TRISTELLATEIA, Thou. Gen. Nov. Madag. 47 (1806). *MALPIGHIACEAE*, Benth. & Hook. f. i. 258.
PLATYNEMA, Wight & Arn. in Edinb. N. Phil. Journ. (July, 1833) 179.
ZYMUM, Noronha, ex Thou. Vég. Iles Afr. Austr. 69 (1807).
africana, *S. Moore, in Journ. Bot.* xv. (1877) 289.—Afr. trop.
Australasiae, A. Rich. Sert. Astrolab. 159 = seq.
australis, *A. Rich. l. c.* 38. t. 15.—Austral.
Bernierana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 497.—Madag.
Bojerana, *A. Juss. l. c.* 498.—Madag.
cocculifolia, *A. Juss. l. c.* 499.—Madag.
dulcamara, *A. Juss. l. c.* 500.—Madag.
heterophylla, *A. Juss. l. c.* 495.—Madag.
madagascariensis, *Poir. Encyc. Suppl.* v. 367.—Madag.
malintana, Blanco, Fl. Filip. ed. II. 267 = australis.
montana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 496.—Madag.
novaeguineensis, Blume, ex A. Juss. l. c. 495 = australis.
pluriseta, *Baill. Adansonia*, xi. (1874) 249.—Madag.
pubescens, *Baill. l. c.*—Madag.
stenactis, *Baill. l. c.* 250.—Madag.

TRISTEMMA, Juss. Gen. 329 (1789). *MELASTOMACEAE*, Benth. & Hook. f. i. 746.
albiflorum, *Benth. in Hook. Niger Fl.* 353.—Afr. trop.
angustifolium, Blume, Bijdr. 1079 = Osbeckia chinensis.
capitatum, Triana, in Trans. Linn. Soc. xxviii. (1871) 56 = Dissotis capitata.
cornifolium, Triana, l. c. = Dissotis cornifolia.
coronatum, *Benth. in Hook. Niger Fl.* 354.—Afr. trop.
erectum, Guill. & Perr. Fl. Seneg. Tent. 312 = Dissotis capitata.
hirtum, *Beauv. Fl. Owar.* i. 94; *Vent. Choix*, sub t. 35.—Afr. trop.
incompletum, *R. Br. in Tuckey, Congo, App.* 435.—Afr. trop.
intermedium, Steud. Nom. ed. I. 853 = incompletum.
involucratum, *Benth. in Hook. Niger Fl.* 354.—Afr. trop.
littorale, *Benth. l. c.* 353.—Afr. trop.
mauritianum, J. F. Gmel. Syst. 693 = virusanum.
neglectum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 299 = Dissotis cornifolia.
Schumacheri, Guill. & Perr. Fl. Seneg. Tent. 311. t. 6. f. 6 = incompletum.
segregatum, Triana, in Trans. Linn. Soc. xxviii. (1871) 56 = Dissotis segregata.
theaefolium, Triana, l. c. = Dissotis theaefolia.
virusanum, *Juss. Gen.* 329; *Vent. Choix*, t. 35.—Ins. Mascar.

TRISTEMON, Klotzsch, in Linnaea, xii. (1838) 245 = **Scyphogyne**, Brongn. (Ericac.).
puberulus, Klotzsch, l. c. = S. puberula.
urceolatus, Klotzsch, l. c. = S. urceolata.

TRISTEMON, Rafin. Am. Monthly Mag. (1819) 192 = **Triglochin**, Rupp. (Naiad.).
conglomeratus, Rafin. Fl. Tellur. iv. 32 = Juncus effusus.
marginatus, Rafin. l. c. = Juncus marginatus.
odoratus, Rafin. l. c. = Juncus marginatus.
palustris, Rafin. in Am. Monthly Mag. (1819) 192 = Trig. palustre.
polycephalus, Rafin. Fl. Tellur. iv. 32 = Juncus scirpoides.
triander, Rafin. in Am. Monthly Mag. (1819) 192 = Trig. striatum.

TRISTEMON, Scheele, in Linnaea, xxi. (1848) 586 = **Cucurbita**, Tourn.
texanum, Scheele, l. c. = C. Pepo.

TRISTERIA, Hook. f. Fl. Brit. Ind. iv. 275 (1884), err. typ. = TREISTERIA, Griff. = **Curanga**, Juss. (Scrophul.).

TRISTERIX, Mart. in Flora, xiii. (1830) i. 108 = **Loranthus**, Linn.
aphyllus, G. Don, Gen. Syst. iii. 418 = L. aphyllus.
Reinwardtianus, Mart. in Flora, xiii. (1830) 109 = L. Reinwardtianus.
tetrandrus, Mart. l. c. = L. tetrandrus.
viridiflorus, Mart. l. c. = L. globosus.

TRISTICHA, Thou. Gen. Nov. Madag. 3 (1806). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 108.
DUFOUREA, Bory, ex Willd. Sp. Pl. v. 55 (1810).
PHILOCRENA, Bong. in Mém. Acad. Pétersb. Sér. VI. i. (1832) 80. t. 6.
alternifolia, *Thou. ex Spreng. Syst.* i. 22.—Madag.
bifaria, *Presl, Rel. Haenk.* i. 86.—Ins. Philipp.
bryoides, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 178 = hypnoides.
ceylanica, Gardn. l. c. 177 = Terniolia zeylanica.
Dregeana, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 112 = hypnoides.
fontinaloides, Welw. ex Wedd. in DC. Prod. xvii. 45 = hypnoides.
hypnoides, *Spreng. Syst.* iv. *Cur. Post.* 10.—Bras.
phasioides, *Griseb. in Goett. Abh.* xxiv. (1879) 41.—Reg. Argent.
Philocrena, Steud. Nom. ed. II. ii. 715 = hypnoides.
trifaria, *Spreng. Syst.* i. 22.—Ins. Mascar.

TRISTICHOCALYX, F. Muell. Fragm. iv. 27 (1863). *MENISPERMACEAE*, Benth. & Hook. f. i. 962.
diffusa, *Miers, Contrib.* iii. 286.—Austral.
pubescens, *F. Muell. Fragm.* iv. 27.—Austral.

TRISTIRA, Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 133. *SAPINDACEAE*.
harpullioides, *Radlk. l. c.*—Ins. Molucc.
triptera, *Radlk. l. c.*—Ins. Philipp.

TRISTYLEA, Jord. & Fourr. Ic. Fl. Eur. ii. 5 (1870) = **Saxifraga**, Tourn. (Saxifr.).
florulenta, Jord. & Fourr. l. c. = S. florulenta.

TRISTYLIUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 247 = **Cleyera**, DC. (Ternstr.).
mexicanum, Turcz. l. c. 248 = C. mexicana.

TRISYNGYNE, Baill. Adansonia, xi. (1874) 136. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 332.
Balansae, *Baill. l. c.* 137.—N. Caled.
codonandra, *Baill. l. c.* 136.—N. Caled.

TRITAENICUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 169 = **Asteriscium**, Cham. & Schlecht. (Umbell.).
nudum, Turcz. l. c.—Chili.

TRITAXIS, Baill. Etud. Gén. Euph. 342. t. 11 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 292.
Beddomei, *Benth. in Journ. Linn. Soc.* xvii. (1878) 221.—Ind. or.
Cumingii, *Benth. l. c.*—Malaya.
Gaudichaudi, *Baill. Etud. Gén. Euph.* 343.—Cochinch.
macrophylla, Muell. Arg. in Flora, xlvii. (1864) 482 = Ostodes macrophylla.
zeylanica, Muell. Arg. l. c. = Ostodes zeylanica.

TRITELANDRA, Rafin. Fl. Tellur. ii. 85 (1836) = **Epidendrum**, Linn. (Orchid.).
fuscata, Rafin. l. c. 86 = E. anceps.

TRITELEIA, Dougl. ex Lindl. Bot. Reg. sub t. 1293 (1830) = **Brodiaea**, Sm. (Liliac.).
aurea, Lindl. Bot. Reg. (1841) Misc. 76 = Milla aurea.
Berteri, Kunth, Enum. Pl. iv. 467 = Milla subbiflora.
bivalvis, Hook. in Lindl. Bot. Reg. sub t. 1293 = Milla bivalvis.
brevipes, Kunze, in Linnaea, xx. (1847) 9 = Milla brevipes.

TRITELEIA :—
conspicua, Baker, in Saund. Refug. Bot. t. 43 = Milla uniflora.
Gaudichaudiana, Kunth, Enum. Pl. iv. 467 = Milla bivalvis.
gracilis, Phil. in Anal. Univ. Chil. (1873) 550.—Chili.
graminifolia, Presl, Bot. Bemerk. 116.—Chili.
grandiflora, Lindl. Bot. Reg. sub t. 1293 = B. Douglasii.
hirtella, Kunth, Enum. Pl. iv. 465 = Milla hirtella.
laxa, Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 413. t. 15. f. 2 = B. laxa.
pedunculariis, Lindl. Bot. Reg. sub t. 1685 = B. pedunculariis.
Poeppigiana, C. Gay, Fl. Chil. vi. 117 = Milla Poeppigiana.
porrifolia, Poepp. Fragm. Syn. Pl. Chile, 10 = Milla porrifolia.
Sellowiana, Kunth, Enum. Pl. iv. 466 = Milla Sellowiana.
sessilis, Phil. in Linnaea, xxix. (1857–58) 72 = Milla sessiliflora.
uniflora, Lindl. Bot. Reg. t. 1293 = Milla uniflora.
uniflora, Van Houtte, in Fl. des Serres, ix. (1853–54) 283. t. 967 = Milla Poeppigiana.
violacea, Kunth, Enum. Pl. iv. 468 = Milla bivalvis.

TRITELEYA, Phil. in Anal. Univ. Chil. (1873), reimpr. 74, sphalm. = praec.

TRITHECA, Miq. Fl. Ind. Bat. i. 1. 614 (1855) = **Ammannia**, Houst. (Lythrar.).
pentandra, Miq. l. c. et 1089 = A. pentandra.

TRITHRINAX, Mart. Hist. Nat. Palm. ii. 149 (1837 ?). *PALMAE*, Benth. & Hook. f. iii. 925.
DIODOSPERMA, H. Wendl. in Bot. Zeit. xxxvi. (1878) 118.
acanthocoma, *Drude, in Regel, Gartenfl.* xxvii. (1878) 361. t. 959.—Bras.
aculeata, Liebm. ex Mart. Hist. Nat. Palm. iii. 320 = Acanthorhiza aculeata.
brasiliensis, *Mart. Hist. Nat. Palm.* ii. 150.—Bras.
brasiliensis, Mart. Palmet. Orbign. 44. t. 10. f. 1, partim = schizophylla.
brasiliensis, H. Wendl. in Kerch. Palm. t. 26 = acanthocoma.
campestris, *Drude & Griseb. in Goett. Abh.* xxiv. (1879) 283.—Reg. Argent.
Chuco, Walp. Ann. i. 1005 = Acanthorhiza Chuco.
compacta, Griseb. Cat. Pl. Cub. 221 = Hemithrinax compacta.
mauritiaeformis, Karst. in Linnaea, xxviii. (1856) 244 = Sabal mauritiiformis.
schizophylla, *Drude, in Mart. Fl. Bras.* iii. II. 551.—Bras.

TRITHURIA, Hook. f. Fl. Tasman. ii. 78. t. 138 (1860). *CENTROLEPIDEAE*, Benth. & Hook. f. iii. 1025.
JUNCELLA, F. Muell. 2nd Gen. Rep. 16 (1854).
occidentalis, *Benth. Fl. Austral.* vii. 199.—Austral.
submersa, *Hook. f. Fl. Tasman.* ii. 79. t. 138.—Tasman.

TRITHYROCARPUS, Hassk. in Flora, xlix. (1866) 211 = **Commelina**, Plum.
gelatinosus, Hassk. l. c. = C. gelatinosa.
oligospatha, Hassk. Commel. Ind. 25 = C. paleata.
polyspatha, Hassk. in Flora, xlix. (1866) 211 = C. obliqua.
paleatus, Hassk. l. c.; et Commel. Ind. 25 = C. paleata.

TRITICUM, Linn. Syst. ed. I (1735); Gen. ed. I. 16 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1204.
AEGILOPS, Linn. Coroll. Gen. 20 (1837).
BROMUS, Scop. Introd. 74 (1777).
CRITHODIUM, Link, in Linnaea, ix. (1834) 132. t. 3.
CRYPTOPYRUM, Heynh. Nom. ii. 174 (1846).
abyssinicum, *Steud. Syn. Pl. Gram.* 342.—Abyss.
acutum, DC. Cat. Hort. Monsp. (1813) 153 = Agropyron acutum.
acutum, Dethard. Consp. Pl. Megalop. 11 = Agropyron repens.
acutum, Fries, Summ. Veg. Scand. 249 = Agropyron obtusiusculum.

TRITICUM :—
aegilopoides, Forsk. Fl. Aegypt. Arab. 26 = Rottboellia hirsuta.
aegilopoides, A. Gray, in Proc. Acad. Philad. 1862 (1864) 79 = Agropyron caninum.
aegilopoides, Turcz. ex Griseb. in Ledeb. Fl. Ross. iv. 339 = Agropyron divergens.
Aegilops, *Beauv. Agrost.* 180, nomen.—Oriens.
aestivum, Linn. Sp. Pl. 85 = vulgare.
affine, Dethard. ex Kunth, Enum. Pl. i. 441 = Agropyron acutum.
alatum, Peterm. in Flora, xxvii. (1844) 234 = durum.
album, Gaertn. ex Steud. Nom. ed. I. 853 = vulgare.
algeriense, Desf. ex Mert. & Koch, Deutschl. Fl. i. 697 = durum.
alpinum, G. Don, ex Mitten, in Hook. Lond. Journ. Bot. vii. (1848) 533 = Agropyron repens.
amyleum, Ser. Mélang. Bot. i. 124 = dicoccum.
angustifolium, Link, Enum. Hort. Berol. i. 97 = Agropyron sibiricum.
aragonense, *Lag. in Varied. Cienc.* ii. IV. (1805) 212.—Hispan.
arenicolum, *Kern. ex Menyh. Kalocsa Vidék. Növényt.* 197.—Hungar.
Arias, Roxas Clem. in Herrera, Agric. (1818) i. 74 = T. Spelta.
aristatum, Hall. [f.] ex Steud. Nom. ed. II. ii. 715 = vulgare.
Arras, *Hochst. in Flora*, xxxi. (1848) 450.—Abyss.
arundinaceum, Schur, Enum. Pl. Transs. 806 = vulgare.
arundinaceum, Steud. Syn. Pl. Gram. 343 = Agropyron repens.
arvense, Schreb. in Koerte & Schweigg. Fl. Erlang. i. 143 = Agropyron repens.
asperrimum, Link, Handb. i. 18, sphalm. = seq.
asperum, *DC. Cat. Hort. Monsp.* (1813) 153.—Hispan.
athericum, Link, in Linnaea, xvii. (1843) 395 = Agropyron pungens.
atratum, Host, Gram. Austr. iv. 5. t. 8 = dicoccum.
attenuatum, H. B. & K. Nov. Gen. et Sp. i. 180 = Agropyron attenuatum.
Aucheri, Parl. Fl. Ital. i. 508 = Agropyron Aucheri.
baeoticum, Boiss. Diagn. Ser. I. xiii. 69 = monococcum.
barbinode, Tausch, in Flora, xvii. (1834) 447 = Brachypodium sylvaticum.
Barrelieri, Kunth, Rév. Gram. i. 145 = Brachypodium pinnatum.
Bauhini, Lag. Gen. et Sp. Nov. 6 = durum.
benghalense, Host, ex Steud. Nom. ed. II. ii. 715, nomen (Quid ?).—Hab. ?
bicorne, *Forsk. Fl. Aegypt. Arab.* 26.—Aegypt.; Syria.
biflorum, Brign. Fasc. Pl. Forojul. 18 = Agropyron caninum.
biflorum, Turcz. ex Griseb. in Ledeb. Fl. Ross. iv. 339 = Agropyron strigosum.
biunciale, Vill. Hist. Pl. Dauph. ii. 167 = Festuca Poa.
Bonaepartis, Spreng. Nachr. I. Bot. Gart. Halle, 40 = Agropyron patulum.
boreale, Turcz. Fl. Baic.-Dahur. ii. 1. 345, in obs. = Agropyron strigosum.
brachystachyon, Hornem. Hort. Hafn. i. 107 = praec. ?
brachystachyum, Lag. ex Schult. Mant. ii. 656 = durum.
brevisetum, *DC. Cat. Hort. Monsp.* (1813) 153.—Hab. ?
brizoides, Lam. Illustr. i. 212 = Demazeria sicula.
bromoides, [Weber, ex] Wigg. Prim. Fl. Holsat. 11 = Brachypodium pinnatum.
Brownei, Kunth, Rév. Gram. i. 145 = Agropyron pectinatum.
bulbosum, Steud. Syn. Pl. Gram. 346 = Agropyron bulbosum.
Bungeanum, Trin. in Bunge, Verz. Suppl. Fl. Alt. 11 = geniculatum.
Buonapartis, Roem. & Schult. Syst. ii. 768, in syn. = Agropyron patulum.
caeruleum, Ard. ex Bayle-Barelle, Monog. Agron. Cereal. 39 (Quid ?).—Hab. ?
caesium, Kunth, Rév. Gram. i. 145 = Agropyron repens.
caespitosum, DC. Cat. Hort. Monsp. (1813) 153 = Brachypodium ramosum.
caespitosum, Walp. Ann. iii. 782 = Agropyron caespitosum.

TRITICUM :—
campestre, Roem. & Schult. Syst. ii. 769 = Secale fragile.
campestre, Nym. Syll. Suppl. 74 = Agropyron glaucum.
candissimum, Bayle-Barelle, Monog. Agron. Cereal. 42; ex Metzger, Europ. Cereal. 18 = durum.
caninum, Ledeb. Fl. Alt. i. 118 = Agropyron strigosum.
caninum, Linn. Sp. Pl. 86 = Agropyron caninum.
capense, Spreng. Pugill. ii. 23 = Eragrostis Uniolae.
caucasicum, Spreng. Fl. Hal. Mant. 35 = Agropyron cristatum.
caudatum, *Gren. & Godr. Fl. Fr.* iii. 603.—Europ.
caudatum, Pers. Syn. i. 110 = Agropyron villosum.
cereale, Bernh. Syst. Verz. Erf. 46 = vulgare.
cereale, Salisb. Prod. 27 = Secale cereale.
cereale, Schrank, Baier. Fl. i. 387 = vulgare.
cerulescens, Bayle-Barelle, Monog. Agron. Cereal. 39 (Quid ?).—Hab. ?
Cevallos, *Lag. Gen. et Sp. Nov.* 6.—Hispan.
chinense, *Trin. ex Bunge, Enum. Pl. Chin. Bor.* 72.—China.
Cienfuegos, Lag. Gen. et Sp. Nov. 6 = dicoccum.
ciliare, *Trin. ex Bunge, Enum. Pl. Chin. Bor.* 72.—China.
ciliatum, Cav. Desc. 317; DC. Fl. Fr. iii. 85 = Brachypodium distachyum.
cinereum, Hort. ex Roem. & Schult. Syst. ii. 770 (Quid ?).—Hab. ?
clavatum, Seidl, ex Opiz, Natural. n. ix. (1825) 106 = vulgare.
coarctatum, Spreng. ex Steud. Nom. ed. II. ii. 716 = Gaudinia geminiflora.
cochleare, Lag. Gen. et Sp. Nov. 6 = durum.
compactum, Host, Gram. Austr. iv. 4. t. 7 = vulgare.
compositum, Linn. Sp. Pl. ed. II. 126 = vulgare.
compressum, Hort. ex Roem. & Schult. Syst. ii. 770 (Quid ?).—Hab. ?
condensatum, Kunth, Enum. Pl. i. 442 = Agropyron condensatum.
creticum, *Roem. & Schult. Syst.* ii. 769.—Ins. Creta.
crinitum, Kunth, Rév. Gram. i. 144 = Agropyron crinitum.
cristatum, Schreb. Beschr. Graeser, ii. 12. t. 23. f. 2 = Agropyron cristatum.
curvifolium, C. Muell. in Walp. Ann. vi. 1047 = Agropyron scirpeum.
cylindricum, *Ces. Pass. & Gib. Comp. Fl. Ital.* 86.—Europ.
cynosuroides, Spreng. Syst. i. 325 = Festuca cynosuroides.
dasyanthum, Ledeb. ex Spreng. Syst. i. 326 = Agropyron dasyanthum.
dasyphyllum, *Schrenk, in Bull. Sc. Acad. Pétersb.* x. (1842) 356.—Sibir. altaic.
dasystachyum, A. Gray, Man. Bot. N. U. St. ed. I. 602 = Agropyron dasystachyum.
densiflorum, Willd. Enum. Hort. Berol. 135 = Agropyron densiflorum.
desertorum, Fisch. ex Link, Enum. Hort. Berol. i. 97 = Agropyron sibiricum.
dicoccum, *Schrank, Baier. Fl.* i. 389.—Europ. (cult.).
distachyon, Brot. Fl. Lusit. i. 119 = Brachypodium distachyum.
distans, Walp. Ann. iii. 782 = Agropyron Bourgaei.
distertum, Tardent, Ess. Hist. Nat. Bessar. 37, nomen = *distachyon* ?
distichum, Schleich. ex DC. Fl. Fr. Suppl. 281 = Agropyron glaucum.
distichum, Thunb. Prod. Pl. Cap. 23 = Agropyron distichum.
divaricatum, Walp. Ann. vi. 1048 = Agropyron divaricatum.
divergens, Steud. Syn. Pl. Gram. 347 = Agropyron divergens.
diversifolium, *Steud. l. c.* 342.—Abyss.
dumetorum, [Honck.] Verz. Aller Gew. Teutschl. 363 = seq. ?
dumetorum, Schreb. in Schweig. & Koerte, Fl. Erlang. i. 143 = Agropyron repens.
duplicatum, *Steud. Syn. Pl. Gram.* 344.—Graecia.
durum, Desf. Fl. Atlant. i. 114 = sativum.
elegans, Spreng. ex Steud. Nom. ed. I. 854, nomen (Quid ?).—Hab. ?
elongatum, Host, Gram. Austr. ii. 18 = Agropyron rigidum.
elymogenes, Arndt, in Flora, xlii. (1859) 215, in textu = Agropyron strictum.
elymoides, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 440.—Abyss.

TRITICUM:—
elymoides, Hornem. Hort. Hafn. i. 107 = T. Spelta.
emarginatum, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 454.—Hab. ?
erinaceum, Hornem. Hort. Hafn. i. 106 = praec. ?
erinaceum, Hort. ex Kunth, Enum. Pl. i. 438 = vulgare.
farctum, Viv. Fl. Ital. Fragm. i. 28 = Agropyron junceum.
farrum, Bayle-Barelle, Monog. Agron. Cereal. 50. t. 4. f. 1, 2 = T. Spelta.
fastuosum, Lag. Gen. et Sp. Nov. 6 = durum.
Festuca, DC. Fl. Fr. iii. 87 = Festuca Poa.
festucoides, Bertol. Fl. Ital. i. 808 = Festuca Poa.
fibrosum, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 209.—Sibir. altaic.
filiforme, Poir. Encyc. Suppl. ii. 677 = Agropyron filiforme.
firmum, Link, Hort. Berol. ii. 188 = Agropyron repens.
flabellatum, Tausch, in Flora, xx. (1837) 117 = Festuca Tauschii.
flexum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 441 = Brachypodium flexum.
Forskalei, Roxas Clem. in Herrera, Agric. (1818) i. 74 = T. Spelta.
fragile, *Ces. Pass. & Gib. Comp. Fl. Ital.* 87.—Italia.
fragile, Link, Hort. Berol. ii. 183 = Secale fragile.
fragile, Roth, Catalecta, fasc. ii. 7 = Agropyron junceum.
Freycenetii, Hort. Lips. ex Pasq. Cat. Ort. Bot. Nap. 104, nomen (Quid ?).—Hab. ?
Gaertnerianum, Lag. Gen. et Sp. Nov. 6 = durum ?
geminatum, Spreng. Syst. i. 326 = Eleusine indica.
geniculatum, *Trin. ex Ledeb. Fl. Alt.* i. 117.—Sibir. altaic.
geniculatum, Walp. Ann. iii. 783 = Brachypodium distachyum.
genuense, *DC. Fl. Fr. Suppl.* 284.—Ital.
giganteum, Roth, Catalecta, fasc. iii. 22 = Agropyron rigidum.
glaucescens, Auct. ex Steud. Nom. ed. II. ii. 716 = Agropyron pungens.
glaucum, Bast. Fl. Maine-et-Loire, 45 = Agropyron pungens.
glaucum, Desf. Tabl. ed. I. 16 = Agropyron glaucum.
glaucum, Host, Gram. Austr. iv. 6. t. 10 = Agropyron littorale.
glaucum, [Honck.] Verz. Aller Gew. Teutschl. 358 = Agropyron glaucum ?
glaucum, Krock. Fl. Siles. Suppl. i. 194 = Agropyron rigidum ?
glaucum, Link, Hort. Berol. ii. 187 = Agropyron junceum.
glaucum, Moench, Meth. 174 = polonicum.
glaucum, Urv. in Mém. Soc. Linn. Par. iv. (1826) 601 = Agropyron repens.
glaucum, Willk. & Lange, Prod. Fl. Hisp. i. 110, sphalm. = Agropyron campestre.
Gmelini, *Trin. in Linnaea*, xii. (1838) 467.—Sibir.
gracile, Brot. Fl. Lusit. i. 121 = Brachypodium sylvaticum.
gracile, DC. Fl. Fr. iii. 84 = Brachypodium pinnatum.
gracile, Pouz. ex Gren. & Godr. Fl. Fr. iii. 608 = Agropyron glaucum.
Halleri, Viv. Fl. Ital. Fragm. 24. t. 26. f. 1 = Festuca Poa.
hamosum, Rehm. ex Boiss. Fl. Orient. v. 661 = Agropyron strigosum.
hebestachyum, Fries, in Bot. Notiser (1858), 129 ; et ex Nym. Suer. Fanerog. ii. 458 = Agropyron obtusiusculum.
Hemipoa, Delile, ex Tenore, Fl. Nap. iv. 18 ; Syll. Fl. Neap. 56 = Festuca Hemipoa.
hirsutum, Hornem. Suppl. Hort. Hafn. 13 = Agropyron cristatum.
hirsutum, Stev. ex Nym. Consp. 841 = Agropyron rigidum.
hispanicum, Viv. Fl. Ital. Fragm. 21 = Festuca Poa.
hispanicum, Willd. Sp. Pl. i. 479 = Festuca tenuiflora.
hordeaceum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 312.—Afr. bor.
hordeaceum, Steud. Syn. Pl. Gram. 430 = Agropyron orientale.
hordeiforme, Host, Gram. Austr. iv. 3. t. 5 = durum.
hordeiforme, Wall. ex Steud. Nom. ed. II. ii. 716 = Hordeum vulgare.
Hornemanni, Roxas Clem. in Herrera, Agric. (1818) i. 73 = monococcum.
Horstianum, Roxas Clem. l. c. 81 = vulgare.

TRITICUM:—
Hosteanum, Spreng. Syst. i. 323, in syn., sphalm. = praec.
hybernum, Linn. Sp. Pl. 86 = vulgare.
ichyostachyum, *Seidl, ex Opiz, Natural.* n. ix. (1825) 106.—Hab. ?
imbricatum, Bieb. Fl. Taur. Cauc. i. 88 = Agropyron cristatum.
imbricatum, Hort. Vindob. ex Roem. & Schult. Syst. ii. 756 = Agropyron sibiricum.
imbricatum, Lam. Illustr. i. 212 = Agropyron repens.
infestum, Salisb. Prod. 27 = Agropyron repens.
intermedium, Bess. Enum. Pl. Volh. 6 = Agropyron rigidum.
intermedium, Bieb. ex Kunth, Enum. Pl. i. 443 = Agropyron cristatum.
intermedium, Host, Fl. Austr. i. 180 = Agropyron glaucum.
intermedium, Nocca & Balb. Fl. Tic. i. 63 = Agropyron repens.
junceum, Linn. Cent. Pl. i. 6 ; Amoen. Acad. iv. 266 = Agropyron junceum.
Kingianum, *Endl. Prod. Fl. Norf.* 21.—Ins. Norfolk.
Kirgianum, Steud. Nom. ed. II. ii. 716, sphalm. = praec.
Koeleri, Roxas Clem. in Herrera, Agric. (1818) i. 77 = vulgare.
Kotschyanum, Boiss. ex Steud. Syn. Pl. Gram. 346 = Agropyron patulum.
Lachenalii, C. C. Gmel. Fl. Bad. i. 291 = Festuca Poa.
laevissimum, Habl. *vel* Hall. [f.] ex Steud. Nom. ed. II. ii. 716, 717 = polonicum.
lasianthum, Steud. Syn. Pl. Gram. 430 = Agropyron orientale.
latronum, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 11 = Agropyron glaucum.
laxum, Fries, Mant. iii. 13 = Agropyron acutum.
Leersianum, Roem. & Schult. Syst. ii. 755, in syn. = Agropyron repens.
ligusticum, Bertol. Fl. Ital. vi. 622 = bicorne.
ligusticum, Walp. Ann. iii. 783 = Agropyron ligusticum.
Linnaeanum, Lag. Gen. et Sp. Nov. 6 = vulgare.
litorale, Auct. ex Nym. Consp. 841 = Agropyron repens.
litorale, Host, Gram. Austr. iv. 5. t. 9 = Agropyron littorale.
litorale, Pall. Reise, iii. 287 = Elymus dasystachys.
litoreum, Brot. Phyt. Lusit. ii. 51 = Agropyron junceum.
littoreum, Schum. Enum. Pl. Saeland. i. 38 = Agropyron repens.
loliaceum, Sm. Engl. Bot. t. 221 = *Festuca loliacea*.
lolioides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 866.—Sibir. altaic.
lolioides, Pers. Syn. i. 110 = Festuca Poa.
longearistatum, Steud. Syn. Pl. Gram. 346 = Agropyron longearistatum.
luzonense, *Kunth, Enum. Pl.* i. 446.—Ins. Philipp.
macrostachyum, Le Gall, Fl. Morb. 760 = Agropyron acutum ?
macrourum, Turcz. in Bull. Soc. Nat. Mosc. (1838) 105 = Brachypodium pinnatum.
maritimum, Linn. Sp. Pl. ed. II. 127 = Cutandia maritima.
maritimum, Viv. Ann. Bot. i. II. 152 = Festuca Poa.
maritimum, Willd. Sp. Pl. i. 479, in syn. = Festuca tenuiflora.
maritimum, With. Bot. Arr. Brit. Pl. ed. II. i. 130 = *Festuca loliacea*.
maritimum, Wulf. in Jacq. Coll. iii. 34 = Festuca rigida.
maximum, Vill. Hist. Pl. Dauph. ii. 156 = polonicum ?
mexicanum, *Schrad. ex Steud. Nom.* ed. II. ii. 716.—Mexic.
missuricum, Spreng. Syst. i. 325 = Festuca spicata.
molle, Hort. ex Roem. & Schult. Syst. ii. 770 (Quid ?).—Hab. ?
monococcum, *Linn. Sp. Pl.* 86.—Graec. ; As. Min.
monostachyum, Lam. ex Roem. & Schult. Syst. ii. 742 = Brachypodium distachyum.
multiflorum, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 311.—N. Zel.
multiflorum, Steud. Nom. ed. I. 855 = Agropyron repens.
murale, Salisb. Prod. 27 = Hordeum murinum.
muricatum, Link, Enum. Hort. Berol. i. 97 = Agropyron cristatum.
Nardus, DC. Fl. Fr. iii. 87 = Festuca tenuiflora.
nigricans, *Pers. Syn.* i. 110.—Gallia.

TRITICUM:—
nodosum, Stev. ex Bieb. Fl. Taur. Cauc. iii. 96 = Agropyron junceum.
nubigenum, Steud. Syn. Pl. Gram. 342 = Agropyron nubigenum.
obtusatum, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 454 = bicorne.
obtusiflorum, DC. Cat. Hort. Monsp. (1813) 153 = Agropyron obtusiflorum.
obtusifolium, Boiss. Elench. 93 = Brachypodium ramosum.
obtusiusculum, Nym. Syll. Suppl. 74 = Agropyron obtusiusculum.
Olgae, *Regel, in Act. Hort. Petrop.* vii. (1880) 588.—Turkestan.
orientale, Bieb. Fl. Taur. Cauc. i. 86 = Agropyron orientale.
ovatum, *Rasp. in Ann. Sc. Nat.* Sér. I. v. (1825) 435 ; *Gren. & Godr. Fl. Fr.* iii. 601.—Europ.
panormitanum, Bertol. Fl. Ital. iv. 780 = Agropyron panormitanum.
patens, Brot. Fl. Lusit. i. 120 = *Festuca loliacea*.
patulum, Willd. Enum. Hort. Berol. 134 = Agropyron patulum.
pauciflorum, Schwein. in Long, Exped. Winnep. ii. 383 = Agropyron violaceum.
pectinatum, Bieb. Fl. Taur. Cauc. i. 87 = Agropyron cristatum, *J. Gaertn.*
pectinatum, R. Br. Prod. 179 = Agropyron pectinatum.
pectiniforme, Steud. Nom. ed. I. 855 = Agropyron pectiniforme.
peruvianum, Lam. Illustr. i. 212 = Agropyron peruvianum.
petraeum, Vis. & Panč. in Mem. Ist. Venet. x. (1861) 446 = Agropyron panormitanum.
phaenicoides, DC. Fl. Fr. Suppl. 284 = Brachypodium pinnatum.
pilosum, Dalz. & Gibs. Bomb. Fl. Suppl. 97 = vulgare.
pilosum, Hornem. Hort. Hafn. i. 106 (Quid ?).—Hab. ?
pilosum, Kunth, Enum. Pl. i. 442 = Agropyron pilosum.
pilosum, Seenus, Reise Istrien, 71 = Agropyron rigidum.
pinnatum, Moench, Enum. Pl. Hass. 53 = Brachypodium pinnatum.
planum, Desf. Tabl. ed. I. 16 = Agropyron patulum.
platystachyum, Lag. Gen. et Sp. Nov. 6 = durum ?
Poa, DC. Fl. Fr. iii. 86 = Festuca Poa.
poliens, Tausch, in Flora, xx. (1837) 117 = Festuca poliens.
polonicum, *Linn. Sp. Pl.* ed. II. 127.—(cult.)
polystachyum, Lag. ex Steud. Nom. ed. II. ii. 717 = Brachypodium distachyum.
Pouzolzii, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 11 = Agropyron glaucum.
proliferum, Reinw. ex Vriese, Pl. Ind. Bat. Or. 113 = Spinifex elegans.
prostratum, Linn. f. Suppl. 114 = Agropyron triticeum.
pruinosum, Hornem. Hort. Hafn. i. 107 = durum.
Pseudo-agropyrum, *Griseb. in Ledeb. Fl. Ross.* iv. 343.—Sibir. transbaic.
puberulum, Boiss. ex Steud. Syn. Pl. Gram. 345 = Agropyrum cristatum.
pubescens, Bieb. Beschr. Casp. 81 = monococcum.
pubescens, Hornem. Hort. Hafn. i. 107 (Quid ?).—Hab. ?
pubescens, Trin. in Bunge, Verz. Suppl. Fl. Alt. 11 = geniculatum.
pubiflorum, *Steud. Syn. Pl. Gram.* 429.—Reg. Magellan.
pulverulentum, Hornem. Hort. Hafn. i. 16 (Quid?).—Hab. ?
pumilum, Linn. f. Suppl. 115 = Agropyron triticeum.
pumilum, *Steud. Syn. Pl. Gram.* 344.—Sibir.
pungens, *Link, Hort. Berol.* ii. 187.—Tauria.
pungens, Pers. Syn. i. 109 = Agropyron pungens.
pycnanthum, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 10 = Agropyron littorale.
pyramidale, Delile, ex Schult. Mant. ii. 414 = vulgare.
quadratum, Mill. Gard. Dict. ed. VIII. n. 4 = vulgare.
ramificum, Link, in Linnaea, xvii. (1843) 397 = Agropyron pungens.
ramosum, Beck, ex Nym. Consp. 841 = Agropyron repens.
ramosum, *Trin. ex Ledeb. Fl. Alt.* i. 114.—Sibir. ural.
recognitum, *Steud. Syn. Pl. Gram.* 342.—Abyss.
repens, Linn. Sp. Pl. 86 = Agropyron repens.
reptans, [Clairv.] Man. Herb. 26 = praec.

TRITICUM :—
Requienii, *Ces. Pass. & Gib. Comp. Fl. Ital.* 86.—Ital.
Richardsoni, Schrad. in Linnaea, xii. (1838) 467 = Agropyron violaceum.
rigidum, Schleich. ex Mert. & Koch, Deutschl. Fl. i. 705 = Agropyron glaucum.
rigidum, Schrad. Sem. Hort. Gotting. (1803) 23 ; ex ej. Fl. Germ. i. 392 = Agropyron rigidum.
Roegnerii, Griseb. in Ledeb. Fl. Ross. iv. 339 = Agropyron Roegnerii.
Rottboellia, DC. Fl. Fr. iii. 86 (*Rottbolla*) = Demazeria loliacea.
Rouxii, Nym. Consp. 842 = Agropyron Rouxii.
rufescens, Hort. ex Steud. Nom. ed. II. ii. 717 = T. Spelta.
rupestre, Fisch. ex Link, Enum. Hort. Berol. i. 98 = Agropyron strigosum.
salinum, Salzm. ex Steud. Nom. ed. II. ii. 717, nomen (Quid ?).—Hab. ?
sardinicum, Koel. ex Spreng. Syst. i. 323 = vulgare.
Sartorii, Boiss. & Heldr. ex Nym. Consp. 840 = Agropyron junceum.
sativum, Lam. Fl. Fr. iii. 625 = durum, turgidum, vulgare.
Savignonii, Steud. Syn. Pl. Gram. 430 = seq.
Savignonii, Nym. Syll. 424 = Agropyron Savignonii.
scaberrimum, Steud. Nom. ed. II. ii. 717 = Brachypodium sylvaticum.
scaberrimum, Tausch, in Flora, xx. (1837) 118 = Brachypodium mexicanum.
scabrum, R. Br. Prod. 178 = Agropyron scabrum.
scabrum, A. Cunn. ex Hook. f. Fl. N. Zel. i. 311 = multiflorum.
scabrum, Steud. Syn. Pl. Gram. 343, sphalm. = Agropyron strictum.
Schimperi, *Hochst. in Flora*, xxiv. (1841) i. *Intell.* 20, nomen ; *et ex A. Rich. Tent. Fl. Abyss.* ii. 441.—Abyss.
Schrenkianum, Fisch. & Mey. ex Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 305 = Brachypodium Schrenkianum.
scirpeum, Guss. Fl. Sic. Prod. i. 148 = Agropyron scirpeum.
secalinum, Georgi, Bemerk. Reise Russ. Reich. i. 198 = Elymus dasystachys.
secundum, Kunth, Enum. Pl. i. 442 = Agropyron secundum.
segetale, Salisb. Prod. 27 = vulgare.
semicostatum, Steud. Syn. Pl. Gram. 346 = Agropyron semicostatum.
sepium, Lam. Fl. Fr. iii. 629 = Agropyron caninum.
sepium, Thuill. Fl. Par. ed. II. 67 = Agropyron repens.
sibiricum, Willd. Enum. Hort. Berol. 135 = Agropyron sibiricum.
siculum, *Roem. & Schult. Syst.* ii. 765.—Sicil. ; As. (cult.).
simplex, Hort. Vindob. ex Schult. Mant. ii. 422, nomen (Quid ?).—Hab. ?
sinaicum, *Steud. Syn. Pl. Gram.* 346.—Mons Sinai.
Solandri, Steud. l. c. 347 = Agropyron scabrum.
Spelta, *Linn. Sp. Pl.* 86.—Hab. ?
speltaeforme, Seidl, ex Opiz, Natural. n. ix. (1825) 106 = T. Spelta.
spinulosum, *Lag. Gen. et Sp. Nov.* 7.—Cuba.
squarrosum, Banks & Soland. ex Hook. f. in Hook. Lond. Journ. Bot. iii. (1844) 417 = Agropyron scabrum.
squarrosum, Roth, Neue Beytr. i. 128 = Agropyron patulum.
striatum, *Steud. Syn. Pl. Gram.* 346.—Ind. or.
strictum, Dethard. Consp. Pl. Megalop. 11 = Agropyron strictum.
strigosum, Less. in Linnaea, ix. (1834) 170 = Agropyron divergens.
strigosum, *Spreng. Syst.* i. 326.—Tauria.
subaristatum, Link, in Linnaea, xvii. (1847) 395 = Agropyron repens.
subsecundum, *Link, Hort. Berol.* ii. 190.—Am. bor.
subtile, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 59 = Brachypodium distachyum.
subulatum, Sm. ex Munro, in Journ. Linn. Soc. vi. (1862) 51 = Festuca Poa ?
subulatum, [Soland. in] Russ. Aleppo, ed. II. ii. 244 = Agropyron subulatum.
supinum, Schrank, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 162 = Poa littoralis.
sylvaticum, Moench, Enum. Pl. Hass. 54 = Brachypodium sylvaticum.

TRITICUM :—
sylvaticum, Salisb. Prod. 27 = Elymus europaeus.
Tauri, Walp. Ann. vi. 1048 = Agropyron Tauri.
tenellum, Linn. Syst. ed. X. 880 = Festuca Poa.
tenellum, Viv. Ann. Bot. i. II. 154 = Festuca tenuiflora.
tenue, Fisch. & Mey. ex Steud. Syn. Pl. Gram. 317 = Festuca subtilis.
tenuiculum, Loisel. in Desv. Journ. Bot. ii. (1809) 219 = Festuca Poa.
teretiflorum, Wibel, Prim. Fl. Werth. 140 = Brachypodium sylvaticum.
Thaoudar, Reut. ex Boiss. Fl. Orient. v. 673 = monococcum.
tomentosum, Bayle, Barelle, Monog. Agron. Cereal. 40. t. 2. f. 10 = durum.
Tournefortii, Walp. Ann. iii. 782 = Agropyron Aucheri.
trachycaulum, *Link, Hort. Berol.* ii. 189.—Am. bor. occ.
triaristatum, *Gren. & Godr. Fl. Fr.* iii. 602.—Europ.
trichophorum, *Link, in Linnaea*, xvii. (1843) 395.—Europ. austr.
tricoccum, Schuebl. Char. Desc. Cereal. (1818) 34 ; et in Flora, iii. (1820) 458. tt. 1, 2 = dicoccum.
triunciale, *Rasp. in Ann. Sc. Nat.* Sér. I. v. (1825) 435 ; *Gren. & Godr. Fl. Fr.* iii. 602.—Europ.
truncatum, Wallr. in Linnaea, xiv. (1840) 544 (Beitr. Fl. Hercyn. 174) = Agropyron glaucum.
Tumonia, Schrad. ex Balb. Cat. Taur. (1813) 78 ; et ex Roem. & Schult. Syst. ii. 770, nomen (Quid ?).—Hab. ?
turgidum, Linn. Sp. Pl. 86 = vulgare.
turgidum, Steud. Syn. Pl. Gram. 342 = recognitum.
unilaterale, All. Fl. Pedem. ii. 258 = Demazeria loliacea.
unilaterale, Linn. Mant. i. 35 = Festuca tenuiflora.
unioloides, Ait. Hort. Kew. ed. I. i. 122 = Demazeria sicula.
vaginans, Pers. Syn. i. 109 = Agropyron vaginans.
Vaillantianum, Roem. & Schult. Syst. ii. 775, in syn. = Agropyron repens.
variegatum, Fisch. ex Spreng. Pugill. ii. 24 = Agropyron sibiricum.
velutinum, Hook. f. Fl. Tasm. ii. 129 = Agropyron velutinum.
velutinum, Schuebl. Char. Desc. Cereal. (1818) 13 ; ex Schult. Mant. ii. 413 = vulgare.
ventricosum, *Ces. Pass. & Gib. Comp. Fl. Ital.* 86.—Europ.
venulosum, Hochst. ex Steud. Syn. Pl. Gram. 342 = recognitum.
venulosum, *Ser. Mélang. Bot.* i. 133.—Aegypt.
villosum, Beauv. Agrost. 103 = Agropyron villosum.
villosum, Host, Gram. Austr. iv. 4. t. 6 = durum.
violaceum, Hornem. in Fl. Dan. t. 2044 = Agropyron violaceum.
virescens, Aschers. ex Nym. Consp. 841 = Agropyron glaucum.
vulgare, *Vill. Hist. Pl. Dauph.* ii. 153.—Cosmop. (cult.)
Youngii, *Hook. f. Handb. N. Zeal. Fl.* 343.—N. Zel.
Zea, Host, Gram. Austr. iii. 20. t. 29 = T. Spelta.

TRITILLARIA, Sang. in Atti Acc. Pont. Lincei, Ser. I. xv. (1862) 29, sphalm. = **Fritillaria**, Tourn. (Liliac.).

TRITOMA, Ker-Gawl. in Bot. Mag. t. 744 (1804) = **Kniphofia**, Moench (Liliac.).
Burchellii, Herb. ex Sweet, Hort. Brit. ed. II. 517, nomen ; Lindl. Bot. Reg. t. 1745 = K. Burchellii.
caulescens, Carr. in Rev. Hortic. (1884) 556 = K. caulescens.
flammea, Hook. Bot. Mag. t. 4819, in syn. = Blandfordia flammea.
glauca, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 892.—Afr. austr.
grandis, Hort.—Cf. Gard. Chron. (1865) 7.—Hab. ?
Leichtlini, Carr. in Rev. Hortic. (1884) 556 = K. Leichtlinii.
Mac-Ovani, Carr. l. c. (1879) 390 = K. Macowani ?
Macowani, Hort.—Cf. Gard. Chron. (1877) II. 374, 407 = K. Macowani.
maroccana, May, in Rev. Hortic. (1878) 255 = K. pumila ?
media, Ker-Gawl. in Bot. Mag. t. 744 = K. sarmentosa.
nobilis, Carr. in Rev. Hortic. (1883) 525 = K. aloides.
pumila, Ker-Gawl. in Bot. Mag. t. 764 = K. pumila.
recurva, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 892 = *glauca*.
recurvata, Hort. l. c. ed. II. 933 = *glauca*.

TRITOMA :—
Rooperi, T. Moore, Gard. Comp. i. 113 = Kniphofia Rooperi.
Saundersii, Carr. in Rev. Hortic. (1882) 504 = K. aloides.
Uvaria, Ker-Gawl. in Bot. Mag. sub t. 744 = K. aloides.

TRITOMANTHE, Link, Enum. Hort. Berol. i. 333 (1821) = **Kniphofia**, Moench (Liliac.).
Burchellii, Steud. Nom. ed. II. ii. 718 = K. Burchellii.
media, Link, Enum. Hort. Berol. i. 333 = K. sarmentosa.
odorata, Lodd. ex Steud. Nom. ed. II. ii. 718, nomen (Quid ?).—Afr. austr.
pumila, Link, Enum. Hort. Berol. i. 333 = K. pumila.
Uvaria, Link, l. c. = K. aloides.

TRITOMIUM, Link, Handb. i. 170 (1829) = TRITOMA, Ker-Gawl. = **Kniphofia**, Moench (Liliac.).

TRITOMODON, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 548 = **Enkianthus**, Lour. (Ericac.).
japonicus, Turcz. l. c. 585.—Japon.

TRITONIA, Ker-Gawl. in Bot. Mag. t. 581 (1802).
IRIDEAE, Benth. & Hook. f. iii. 708.
AGRETTA, Eckl. Top. Verz. 23 (1827).
BELLENDENIA, Rafin. in Loud. Gard. Mag. viii. (1832) 245.
DICHONE, Laws. ex Salisb. in Trans. Hort. Soc. i. (1812) 320.
MONTBRETIA, DC. in Bull. Soc. Philom. iii. (1803) 151.
TRITONIXIA, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 354.
WAITZIA, Reichb. Consp. 60 (1828).
abyssinica, Walp. Ann. iii. 613 = Lapeyrousia abyssinica.
anigozanthiflora, *Sweet, Hort. Brit.* ed. I. 398, nomen. —Afr. austr.
aurantiaca, Eckl. Top. Verz. 29 = crocata.
aurea, Pappe, ex Hook. Bot. Mag. t. 4335 = Crocosmia aurea.
Bakeri, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 358.—Afr. austr.
Bolusii, Baker, in Journ. Bot. xiv. (1876) 337 = dubia.
capensis. *Ker-Gawl. in Bot. Mag.* t. 618.—Afr. austr.
carnea, Hort. ex Pasq. Cat. Orto Bot. Nap. 104, nomen (Quid ?).—Hab. ?
catenularis, Salisb. in Trans. Hort. Soc. i. (1812) 319 = lineata.
coccinea, Eckl. Top. Verz. 29 = crocata.
concolor, Sweet, Hort. Brit. ed. I. 398 = Ixia paniculata.
Cooperi, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 358 (Ixiae sp.).—Afr. austr.
crispa, *Ker-Gawl. in Bot. Mag.* t. 678.—Afr. austr.
crocata, *Ker-Gawl. l. c.* sub t. 581.—Afr. austr.
deusta, *Ker-Gawl. l. c.* t. 622.—Afr. austr.
dubia, *Eckl. Top. Verz.* 30, nomen ; *et ex Klatt, in Linnaea*, xxxii. (1863) 761.—Afr. austr.
fenestrata, *Ker-Gawl. in Bot. Mag.* t. 704.—Afr. austr.
flava, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 228.—Afr. austr.
fucata, Lindl. Bot. Reg. (1838) t. 35 = Antholyza fucata.
fulva, Dehnh. Cat. Pl. Hort. Camald. ed. II. 24 (Quid ?).—Afr. austr.
hyalina, Baker, in Journ. Linn. Soc. xvi. (1877) 163 = fenestrata.
kamisbergensis, *Klatt, in Linnaea*, xxxii. (1863) 760.—Afr. austr.
lacerata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 358 = crispa.
laxifolia, *Benth. & Hook. f. Gen.* iii. 708.—Afr. trop.
lineata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 228.—Afr. austr.
longiflora, Ker-Gawl. l. c. = Ixia paniculata.
Mac-Owani, Hort.—Cf. Gard. Chron. (1877) II. 407, sphalm. = Kniphofia Mac-Owani.
magniflora, Dehnh. Cat. Pl. Hort. Camald. ed. II. 24 (Quid ?).—Afr. austr.
media, Pritz. Ic. Ind. i. 1124, sphalm. = *Tritoma media*.
miniata, Ker-Gawl. in Bot. Mag. t. 609 = crocata.
nervosa, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 358 (= *Morphixia nervosa*).—Afr. austr.

TRITONIA:—
odorata, Lodd. Bot. Cab. t. 1820 = Freesia refracta.
pallida, *Ker-Gawl. in Bot. Mag.* sub t. 1275.—Afr. austr.
paniculata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 358 = Ixia paniculata.
pauciflora, *Klatt, l. c.*—Afr. austr.
pectinata, Ker-Gawl. in Bot. Mag. sub t. 678 = crispa.
Pottsii, *Benth. & Hook. f. Gen.* iii. 708.—Afr. austr.
pulchella, Dehnh. Cat. Pl. Hort. Camald. ed. II. 24 (Quid ?).—Afr. austr.
purpurea, Ker-Gawl. in Bot. Mag. sub t. 1275 = crocata.
refracta, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 228 = Freesia refracta.
Rocheana, Sweet, Hort. Brit. ed. I. 398 = seq.
Rochensis, Ker-Gawl. in Bot. Mag. t. 1503 = Ixia paniculata.
rosea, [Dryand. in] Ait. Hort. Kew. ed. II. i. 91 = capensis.
rosea, *Klatt, in Linnaea*, xxxii. (1863) 760.—Afr. austr.
sanguinea, Eckl. Top. Verz. 29 = crocata.
Schimperi, Aschers. & Klatt, in Linnaea, xxxiv. (1865–66) 697 = Acidanthera unicolor.
scillaris, *Baker, in Journ. Linn. Soc.* xvi. (1877) 163.—Afr. austr.
secunda, *Steud. Nom.* ed. II. ii. 719.—Afr. austr.
securigera, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 228.—Afr. austr.
squalida, *Ker-Gawl. in Bot. Mag.* t. 581.—Afr. austr.
striata, Ker-Gawl. l. c. sub t. 678 = Gladiolus striatus.
strictifolia, *Benth. & Hook. f. Gen.* iii. 708.—Afr. austr.
tenuiflora, Ker-Gawl. in Bot. Mag. sub t. 1275 = Ixia paniculata.
tripunctata, Dehnh. Cat. Pl. Hort. Camald. ed. II. 24 (Quid ?).—Afr. austr.
tristis, Dehnh. l. c. (Quid ?)—Afr. austr.
undulata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 163.—Afr. austr.
ventricosa, Hort. ex Pasq. Cat. Orto Bot. Nap. 104, nomen (Quid ?).—Hab. ?
viridis, *Ker-Gawl. in Bot. Mag.* t. 1275.—Afr. austr.
xanthospila, Ker-Gawl. ex Spreng. Syst. i. 154 (= *Anomatheca xanthospila*).—Afr. austr.

TRITONIXIA, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 355 = **Tritonia**, Ker-Gawl. (Irid.).
Bolusii, Klatt, l. c. 356 = Tritonia dubia.
conferta, Klatt, l. c. = Tritonia dubia.
crocata, Klatt, l. c. = Tritonia crocata.
deusta, Klatt, l. c. = Tritonia deusta.
disticha, Klatt, l. c.—Afr. austr.
dubia, Klatt, l. c. = Tritonia dubia.
hyalina, Klatt, l. c. = Tritonia fenestrata.
kamisbergensis, Klatt, l. c. = Tritonia kamisbergensis.
lineata, Klatt, l. c. 357 = Tritonia lineata.
miniata, Klatt, l. c. 356 = Tritonia crocata.
purpurea, Klatt, l. c. = Tritonia crocata.
rosea, Klatt, l. c. = Tritonia rosea.
scillaris, Klatt, l. c. 357 = Tritonia scillaris.
squalida, Klatt, l. c. 356 = Tritonia squalida.
stricta, Klatt, l. c. 357 = Tritonia scillaris.
undulata, Klatt, l. c. = Tritonia undulata.

TRITOPHUS, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 341 = **Kaempferia**, Linn. (Scitam.).

TRITRIELA, Rafin. Fl. Tellur. iii. 59 (1836) = **Ornithogalum**, Tourn. (Liliac.).

TRIUMFETTA, Plum. ex Linn. Gen. ed. I. 344 (1737). *TILIACEAE*, Benth. & Hook. f. i. 234.
BARTRAMIA, Linn. Diss. Dass. 17 (1747); Fl. Zeyl. 77 (1747); Gaertn. Fruct. ii. 137. t. 111 (1791).
PORPA, Blume, Bijdr. 117 (1825).
abutiloides, *A. St. Hil. Fl. Bras. Mer.* i. 287.—Bras.
actinocarpa, *S. Moore, in Journ. Bot.* xv. (1877) 66.—Afr. trop.
acuminata, *H. B. & K. Nov. Gen. et Sp.* v. 344.—N. Granat.
acutiloba, *DC. Pl. Rar. Jard. Genève*, 64.—Cuba.
aestuans, Royle, Illustr. Bot. Himal. 103 (Quid ?).—Reg. Himal.
althaeoides, *Lam. Encyc.* iii. 420.—Am. trop.
angulata, Lam. l. c. 421 = rhomboidea.

TRIUMFETTA:—
annua, *Linn. Mant.* i. 73.—As. et Afr. trop.
appendiculata, *F. Muell. Fragm.* iii. 7.—Austral.
Bartramia, Linn. Syst. ed. X. 1044; Roxb. Fl. Ind. ii. 463 = rhomboidea.
benguelensis, Wawra & Peyr. in Sitzb. Akad. Wien, Math.-Nat. xxxviii. (1860) 559 = flavescens.
Berterii, *Spreng. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 227.—N. Granat.
bicornuta, *Turcz. l. c.* xxxvi. (1863) i. 574.—Java.
bilocularis, Wall. Cat. n. 1081 = rotundifolia.
bogotensis, *DC. Prod.* i. 506.—Am. centr.; N. Granat.
Botteriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 260.—Mexic.
brachypetala, *Turcz. l. c.* xxxi. (1858) i. 227.—Mexic.
calycina, *Turcz. l. c.* xxxvi. (1863) i. 574.—Peruv.
calyculata, Vell. Fl. Flum. 200; v. t. 6 = semitriloba.
cana, Blume, Bijdr. 113 = obliqua ?
caudata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 353.—N. Granat.
chaetocarpa, *F. Muell. Fragm* xi. 61.—Austral.
chrysotricha, *Boj. in Bouton, Rapp. Ann. Maur.* (1842) 19.—Madag.
conspicua, *Trimen, in Journ. Bot.* xxiii. (1885) 206.—Zeylan.
cordifolia, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 91 = semitriloba.
cuneata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 84 = neglecta.
cymosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 353.—N. Granat.
denticulata, *R. Br. ex Benth. Fl. Austral.* i. 274.—Austral.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 261.—Bras.
diversifolia, E. Mey. in Drège, Zwei Pfl. Docum. 227, nomen = rhomboidea.
diversiloba, Presl, Bot. Bemerk. 20 = semitriloba.
dumetorum, Schlecht. in Linnaea, xi. (1837) 377 = bogotensis.
effusa, *E. Mey. in Drège, Zwei Pfl. Docum.* 227, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 228.—Afr. austr.
eriocarpa, A. St. Hil. Fl. Bras. Mer. i. 288 = rhomboidea.
eriophlebia, Hook. f. in Hook. Niger Fl. 235 = rhomboidea.
Fabreana, *Gaudich. in Freyc. Voy. Bot.* 478.—Ins. Marian.
flavescens, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 82.—Afr. trop.
Galeottiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 260.—Mexic.
geoides, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* i. 256.—Afr. trop.
glabra, *Spreng. Fl. Hal. Mant.* 41.—Ind. or.
glandulosa, Forsk. Fl. Aegypt. Arab. p. cxii; Lam. Encyc. iii. 421 = rhomboidea.
glandulosa, Heyne, ex Wall. Cat. n. 1077 = pilosa.
glaucescens, *R. Br. ex Benth. Fl. Austral.* i. 273.—Austral.
glechomoides, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* i. 258.—Afr. trop.
grandidens, *Hance, in Journ. Bot.* xv. (1877) 329.—China.
grandiflora, *Vahl, Eclog. Am.* ii. 34.—Ind. occ.; Mexic.
graveolens, *Blume, Bijdr.* 113.—Java.
graveolens, Zoll. & Mor. Syst. Verz. Zoll. 27 (excl. syn.) = subhirtella.
grossulariaefolia, *A. Rich. Ess. Fl. Cub.* 205.—Ind. occ.
guazumaefolia, *Boj. in Bouton, Rapp. Ann. Maur.* (1842) 18.—Madag.
havanensis, H. B. & K. Nov. Gen. et Sp. v. 345 = semitriloba.
heterophylla, Griseb. Fl. Brit. W. Ind. 96 = semitriloba.
heterophylla, *Lam. Encyc.* iii. 420.—Am. austr.
Heudelotii, *Planch. ex Mast. in Oliver, Fl. Trop. Afr.* i. 259.—Afr. trop.
hirta, *Vahl, Symb. Bot.* iii. 63.—N. Granat.
hispida, *A. Rich. Ess. Fl. Cub.* 204.—Ind. occ.
Hostmanni, Miq. in Linnaea, xxii. (1849) 466 = bogotensis ?
Hostmanniana, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1175, nomen.—Guian.

TRIUMFETTA:—
humifusa, *Hassk. Retzia*, i. 139.—Java.
indica, Buch.-Ham. ex Wall. Cat. nn. 237, 1078 D = tomentosa.
indica, Lam. Encyc. iii. 420 = annua.
Johnstoniana, *F. Muell. in Melb. Chemist* (Febr. 1885).—Austral.
Josefina, *Polak. in Linnaea*, xli. (1877) 552.—Costa Rica.
Kirkii, *Mast. in Oliver, Fl. Trop. Afr.* i. 259.—Afr. trop.
lanigera, Noronha, in Verb. Batav. Gen. v. (1790) ed. I. Art. IV. 27 = villosiuscula.
lappacea, Salisb. Prod. 366 = T. Lappula.
Lappula, *Linn. Sp. Pl.* 444.—Am. et Afr. trop.
Lappula, Schomb. Faun. Fl. Brit. Gui. 1258, nomen = heterophylla ?
Lappula, Vell. Fl. Flum. v. t. 5; text. ed. Netto, 191 = rhomboidea.
leptacantha, *F. Muell. Fragm.* xi. 62.—Austral.
Lindeniana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 229.—Mexic.
longicoma, *A. St. Hil. Fl. Bras. Mer.* i. 285.—Bras.
longicuspis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 229.—Mexic.
longipes, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1175, nomen.—Guian.
longiseta, *DC. Pl. Rar. Jard. Genève*, 64.—Cuba.
longiseta, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 92 = semitriloba.
macrocalyx, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 230 = speciosa.
macropetala, F. Muell. Fragm. iii. 8 = Corchorus echinatus.
macrophylla, Vahl, Eclog. Am. ii. 34 = althaeoides.
Martiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 228 = rhomboidea.
mauritiana, *Presl, in Oken, Isis*, xxi. (1828) 273.—Ins. Maurit.
mexicana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 230 = Belotia grewiaefolia.
micracantha, *F. Muell. Fragm.* iii. 7.—Austral.
micropetala, Lindl. Bot. Reg. t. 1058 = rhomboidea.
microphylla, Wight & Arn. Prod. 75 = pilosa ?
mollis, Schum. & Thonn. Beskr. Guin. Pl. 238 = rhomboidea.
mollissima, *H. B. & K. Nov. Gen. et Sp.* v. 345.—Am. austr.
monoica, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. i. 82 = flavescens.
nana, Boj. Hort. Maurit. 43, sphalm. = annua.
neglecta, *Wight & Arn. Prod.* 75.—Ind. or.; Afr. trop.
nemoralis, *A. St. Hil. Fl. Bras. Mer.* i. 284.—Bras.
nipaulensis, G. Don, Gen. Syst. i. 545 = annua.
obliqua, *Roth, Nov. Pl. Sp.* 224.—As. trop.
oblonga, Hornem. ex Schrank, in Syll. Ratisb. i. (1824) 213; Wall. in D. Don, Prod. Fl. Nep. 227 = pilosa.
oblongata, Link, Enum. Hort. Berol. ii. 5 = pilosa.
obovata, *Cham. & Schlecht. in Linnaea*, v. (1830) 228.—Mexic.
obscura, A. St. Hil. Fl. Bras. Mer. i. 286 = semitriloba.
octandra, Griff. Notul. iv. 512 = pilosa.
orbiculata, Roxb. Hort. Beng. [36] = rotundifolia.
Orizabae, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 261 (Orizaba).—Mexic.
orthacantha, *Welw. ex Mast. in Oliver, Fl. Trop. Afr.* i. 258.—Afr. trop.
ovata, DC. Prod. i. 507 = semitriloba.
oxyphylla, [*Moç. & Sesse,*] *ex DC. l. c.* 508.—Mexic.
paniculata, *Hook. & Arn. Bot. Beech. Voy.* 279.—Mexic.
parviflora, *Benth. Fl. Austral.* i. 274.—Austral.
pentandra, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 93. t. 19 = neglecta.
petiolaris, *Lindl. Bot. Reg.* sub t. 1058.—Peruv.
pilosa, H. B. & K. Nov. Gen. et Sp. v. 343 = bogotensis.
pilosa, *Roth, Nov. Pl. Sp.* 223.—As. et Afr. trop.
pilosa, Wall. Cat. n. 1080 = tomentosa.
pilosula, Thw. Enum. Pl. Zeyl. 401 = neglecta.
Plumieri, Gaertn. Fruct. ii. 137 = T. Lappula.
plumigera, *F. Muell. Fragm.* i. 69.—Austral.
polyandra, [Moç. & Sesse,] ex DC. Prod. i. 508 = grandiflora.
polycarpa, Wall. Cat. n. 1079 = pilosa, annua.
procumbens, Boj. Hort. Maurit. 44 = radicans.

TRIUMFETTA:—
procumbens, *Forst. f. Prod.* 35.—Austral.; Ins. Societ.
Pseudo-angulata, *Blume, Bijdr.* 111.—Java.
quinqueloba, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 574.—Peruv.
radicans, *Boj. in Bouton, Rapp. Ann. Maur.* (1842) 20.—Ins. Seychell.
rhombeaefolia, Sw. Prod. Veg. Ind. Occ. 76 = seq.
rhomboidea, *Jacq. Enum. Pl. Carib.* 22; *Select. Am.* 147. t. 90.—Reg. trop.
rhomboidea, Lindl. Coll. Bot. t. 29 = petiolaris.
riparia, Hochst. in Flora, xxvii. (1844) 295 = rhomboidea.
rotundifolia, *Lam. Encyc.* iii. 421; *Roth, Nov. Pl. Sp.* 222.—Ind. or.
rubricaulis, *H. B. & K. Nov. Gen. et Sp.* v. 342.—Venezuela.
Salzmanni, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 229 = semitriloba.
Schimperi, Hochst. in Flora, xxiv. (1841) i. Intell. 29; et ex Steud. Nom. ed. II. ii. 719 = annua.
semitriloba, Boj. Hort. Maurit. 43 = rhomboidea.
semitriloba, Griseb. Fl. Brit. W. Ind. 96, partim = heterophylla.
semitriloba, *Jacq. Enum. Pl. Carib.* 22; *Select. Am.* 147.—Reg. trop.
sepium, A. St. Hil. Fl. Bras. Mer. i. 286 = semitriloba.
setulosa, *Mast. in Oliver, Fl. Trop. Afr.* i. 259.—Afr. trop.
sinuosa, Miq. in Linnaea, xxii. (1849) 466 = heterophylla.
Sonderii, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 17.—Afr. trop. et austr.
speciosa, *Seem. Bot. Voy. Herald*, 86.—Panama; Mexic.
spicata, *Blume, Bijdr.* 111.—Java.
subhirtella, *Miq. Fl. Ind. Bat.* i. ii. 196.—Java.
suborbiculata, DC. Prod. i. 506 = rotundifolia.
suffruticosa, *Blume, Bijdr.* 113.—Java.
surinamensis, Steud. in Flora, xxvi. (1843) 755 (sp. dub.).—Guian.
Thonningiana, DC. Pl. Rar. Jard. Genève, 64 = rhomboidea.
tiliaefolia, *Vahl, ex DC. l. c.* 65.—Ind. occ.
tomentosa, *Boj. Hort. Maurit.* 43, nomen; *et in Bouton, Rapp. Ann. Maur.* (1842) 19.—Geront. trop.
tomentosa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 27 = cana.
tomentosa, Telfair, ex Wall. Cat. n. 1080 = pilosa.
trichocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 84.—Abyss.
trichocarpa, Sond. in Linnaea, xxiii. (1850) 19 = Sonderii.
trichoclada, DC. Prod. i. 507 = annua.
triclada, Link, Enum. Hort. Berol. ii. 5 = annua.
tricuspis, A. St. Hil. Fl. Bras. Mer. i. 288 = semitriloba.
trilobata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 27 = spicata?
trilocularis, Roxb. Hort. Beng. [36]; Fl. Ind. ii. 462 = rhomboidea.
ulmifolia, Desv. in Ham. Prod. Pl. Ind. Occ. 41 = semitriloba.
urticaefolia, Thunb. Fl. Javan. 19 (Quid?).—Java.
Vahlii, Poir. Encyc. Suppl. iii. 300 = rhomboidea.
velutina, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = mauritiana.
velutina, Vahl, Symb. Bot. iii. 62 = rhomboidea.
vestita, Wall. Cat. n. 1078 = tomentosa, rhomboidea.
villosiuscula, *Blume, Bijdr.* 112.—Java.
Welwitschii, *Mast. in Oliver, Fl. Trop. Afr.* i. 255.—Afr. trop.
Winneckeana, *F. Muell. in Winnecke, Rep.* (1884) 15.—Austral.

TRIUMPHETTA, Griff. Notul. iv. 512 (1854), sphalm. = praec.

TRIUNILA, Rafin. Neogenyt. 4 (1825) = **Uniola**, Linn. (Gramin.).

TRIURIS, Miers, in Proc. Linn. Soc. i. (1841) 96. *TRIURIDEAE*, Benth. & Hook. f. iii. 1002.
Hexuris, Miers, in Proc. Linn. Soc. ii. (1850) 72; et in Trans. Linn. Soc. xxi. (1852) 44.
Peltophyllum, Gardn. ll. cc. i. (1843) 176; xix. (1845) 157. t. 15.

TRIURIS:—
hyalina, *Miers, in Proc. Linn. Soc.* i. (1841) 96.—Bras.
lutea, *Benth. & Hook. f. Gen.* iii. 1002.—Bras.

TRIVALVARIA, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 19 (1865) = **Polyalthia**, Blume (Anonac.).
carnosa, Scheff. in Flora, lii. (1869) 301 = Saccopetalum carnosum.
longirostris, Becc. ex Scheff. in Ann. Jard. Buitenz. ii. (1885) 28.—N. Guin.
macrophylla, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 19 = P. macrophylla?
Stymanni, Scheff. in Flora, lii. (1869) 301.—Sumatra.

TRIVOLVULUS, Moç. & Sesse, ex Choisy, in DC. Prod. ix. 368 (1845) = **Ipomoea**, Linn. (Convolv.).
nutans, Moç. & Sesse, ex Choisy, l. c. = I. nutans.

TRIXAGO, Hall. Hist. Stirp. Helv. 101 (1768); Moench, Meth. 398 (1794) = **Stachys**, Tourn. (Labiat.).
arvensis, Hoffmgg. & Link, Fl. Port. i. 102 = S. arvensis.
colorata, Presl, Fl. Sic. p. xxxvii = S. arvensis.
cordifolia, Moench, Meth. 398 = S. arvensis.
punctata, Gilib. Fl. Lituan. i. 73 = S. arvensis.
scordifolia, Steud. Nom. ed. II. ii. 719, sphalm. = S. arvensis.

TRIXAGO, Rafin. Fl. Tellur. iii. 85 (1836) = **Teucrium**, Tourn. (Labiat.).
Botrys, Rafin. l. c. = Teucrium Botrys.

TRIXAGO, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4 = **Bartsia**, Linn. (Scrophul.).
apula, Stev. l. c. = B. Trixago.
atroviolacea, Nym. Consp. 550 = B. latifolia.
carnea, Griseb. Spicil. Fl. Rumel. ii. 12 = B. Trixago.
latifolia, Reichb. Fl. Germ. Excurs. 360 = B. latifolia.
maxima, Webb & Berth. Phyt. Canar. iii. 151 = B. Trixago.
purpurea, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4 = B. latifolia.
rhinanthina, Link, ex Spreng. Syst. ii. 773 = B. Trixago.
scordifolia, Reichb. in Flora, xiii. (1830) 131 = B. scordifolia.
versicolor, Webb & Berth. Phyt. Canar. iii. 151 = B. Trixago.
viscosa, Reichb. Fl. Germ. Excurs. 360 = B. viscosa.

TRIXANTHERA, Rafin. Sylva Tellur. 146 (1838) = **Trichanthera**, Kunth (Acanth.).
angularis, Rafin. l. c. = Trichanthera gigantea.

TRIXAPIAS, Rafin. Fl. Tellur. iv. 108 (1836) = **Utricularia**, Linn. (Lentib.).
capillacea, Rafin. l. c. = U. capillacea.

TRIXELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 135 = **Stachys**, Tourn. (Labiat.).
arvensis, Fourr. l. c. = S. arvensis.

TRIXIS, P. Br. Hist. Jamaic. 312 (1756). *COMPOSITAE*, Benth. & Hook. f. ii. 501.
Bowmania, Gardn. in Hook. Ic. Pl. t. 519 (1843).
Castra, Vell. Fl. Flum. 242 (1825).
Cleanthes, D. Don, in Trans. Linn. Soc. xvi. (1830) 194.
Dolichlasium, Lag. Amen. Nat. i. 33 (1811).
Holocheilus, Cass. in Bull. Soc. Philom. (1818) 73.
Perdicium, Linn. Pl. Afr. Rar. 22 (1762).
Platycheilus, Cass. in Dict. Sc. Nat. xxxiv. 212 (1825).
Prionanthes, Schrank, Pl. Rar. Hort. Monac. ii. t. 51 (1819).
Tenorea, Colla, Hort. Ripul. 137 (1824).
alata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 192.—Mexic.
angustifolia, *DC. Prod.* vii. 69.—Am. bor. occ.; Mexic.
antimenorrhoea, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 385 = divaricata.
auriculata, R. Grah. in Edinb. N. Phil. Journ. (Apr.–June 1827) 178; et in Bot. Mag. t. 2765 = divaricata.
Bowmanii, *Baker, in Mart. Fl. Bras.* vi. iii. 390.—Bras.
bracteata, *DC. Prod.* vii. 68.—Am. austr.

TRIXIS:—
brasiliensis, *DC. in Ann. Mus. Par.* xix. (1812) 67.—Bras.
brasiliensis, Less. in Linnaea, v. (1830) 26 = Lessingii.
cacalioides, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 187.—Peruv.
calcarea, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 460 = divaricata.
californica, Kellogg, in Proc. Calif. Acad. ii. (1863) 182 = angustifolia.
calycina, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 187.—Bras.
chionopappa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 386 = ophiorhiza.
conferta, *Benth. Pl. Hartw.* 289.—Mexic.
corymbosa, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 188.—Mexic.
corymbosa, Moç. ex DC. Prod. vii. 66 = Perezia fruticosa.
crenata, Boj. ex DC. l. c. v. 198 = Mikania dioscoreaefolia.
decurrens, *DC. l. c.* vii. 68.—Mexic.
dioscoreaefolia, Boj. ex DC. l. c. v. 198 = Mikania dioscoreaefolia.
discolor, D. Don, in Phil. Mag. (1832) 388 = divaricata.
divaricata, *Spreng. Syst.* iii. 501.—Am. trop.
Ehrenbergii, *Kunze, in Linnaea*, xvi. (1842) 317.—Mexic.
flexuosa, Less. l. c. v. (1830) 32 = mollissima.
flexuosa, Spreng. Syst. iii. 501 = divaricata.
floribunda, Boj. ex DC. Prod. v. 198 = Mikania floribunda.
frutescens, Spreng. Syst. iii. 501 = radiale.
glaberrima, *Less. Syn. Comp.* 413.—Bras.
glabra, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 297.—Mexic.
glabrata, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 388 = picroides.
glandulifera, *Benth. & Hook. f. ex Hieron. in Bol. Acad. Nac. Córdova*, iv. (1880) 50.—Chili.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. iii. 391.—Bras.
glutinosa, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 189.—Bras.
Haenkei, *Sch. Bip. in Seem. Bot. Voy. Herald*, 314.—Mexic.
havanensis, Spreng. Syst. iii. 501 = radiale.
hieracioides, DC. Prod. vii. 71 = ochroleuca.
Inula, Crantz, Inst. i. 329 = radiale.
involucrata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 193.—Mexic.
laevigata, Lag. Amen. Nat. i. 36 = radiale.
lanceolata, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 388 = glutinosa.
latifolia, Hook. & Arn. Bot. Beech. Voy. 300 = Perezia rigida.
Lessingii, *DC. Prod.* vii. 70.—Bras.
longifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 191.—Mexic.
Lundii, DC. Prod. vii. 71 = Lessingii.
magellanica, Bert. ex Steud. Nom. ed. II. ii. 75, 720 = Chaptalia exscapa.
megapotamica, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 33 = verbasciformis.
mexicana, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 27.—Mexic.
mexicana, Moç. ex Less. in Linnaea, v. (1830) 33 = radiale.
michuacana, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 28.—Mexic.
mollissima, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 299.—Bras.
myricaefolia, Boj. ex DC. Prod. v. 188 = Mikania myricaefolia.
Neaeana, *DC. Prod.* vii. 67.—Peruv.
obvallata, *Hook. & Arn. Bot. Beech. Voy.* 300. t. 65.—Mexic.
ochroleuca, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 33.—Bras.
odoratissima, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 461 = divaricata.
ophiorhiza, *Gardn. l. c.* 461.—Bras.
othonnoides, Less. in Linnaea, v. (1830) 27 = brasiliensis.
pallida, *Less. l. c.* 30.—Bras.
papillosa, *D. Don, in Phil. Mag.* (1832) 388.—Chili.
paradoxa, *Cass. Opusc.* ii. 160.—Peruv.

TRIXIS:—
patens, Sch. Bip. in Seem. Bot. Voy. Herald, 315 = Perezia patens.
picroides, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 462.—Bras.
pinnatifida, *Less. in Linnaea*, v. (1830) 29.—Bras.
pyrifolia, Boj. ex DC. Prod. v. 195 = Mikania pyrifolia.
radiale, *Lag. Amen. Nat.* i. 36, in obs.—Am. trop.
rosea, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 387 = verbasciformis.
rosea, Poepp. ex DC. Prod. vii. 57 = Leucheria andryaloides.
rosmarinifolia, *Nees, in Linnaea*, xx. (1847) 699.—Mexic.
salicifolia, D. Don, in Trans. Linn. Soc. xvi. (1830) 298 = divaricata.
Seemannii, Sch. Bip. in Seem. Bot. Voy. Herald, 315 = Perezia Seemannii.
Sellowii, Less. ex DC. Prod. vii. 71 = brasiliensis.
senecioides, Hook. Exot. Fl. ii. 101 = Leucheria senecioides.
sinuosa, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 389 = Vauthieri.
spicata, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 462.—Bras.
Sprengeliana, Gardn. l. c. 460 = divaricata.
stricta, *Less. in Linnaea*, v. (1830) 28.—Bras.
stricta, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 386 = glaberrima.
suaveolens, Mart. ex Baker, l. c. 387 = verbasciformis.
suffruticosa, S. Wats. in Bot. Calif. ii. 459 = angustifolia.
thunbergioides, Boj. ex DC. Prod. v. 198 = Mikania thunbergioides.
turbinata, Sch. Bip. in Seem. Bot. Voy. Herald, 315 = Perezia turbinata.
Vauthieri, *DC. Prod.* vii. 69.—Bras.
verbasciformis, *Less. in Linnaea*, v. (1830) 29.—Bras.

TRIXIS, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 220 = **Proserpinaca**, Linn. (Halor.).
palustris, Gaertn. Fruct. i. 115. t. 24. f. 8 = P. palustris.

TRIXIS, Sw. Prod. Veg. Ind. Occ. 7, 115 (1788); Fl. Ind. Occ. iii. 1374. t. 26 (1806) = **Clibadium**, Linn. (Compos.).
aspera, Sw. Prod. l. c. 115 = C. surinamense.
Barbasco, Auct. ex Steud. Nom. ed. II. i. 384; ii. 720 = C. Barbasco.
erosa, Sw. Prod. Veg. Ind. Occ. 115 = C. erosum.
nereifolia, Bonpl. ex Humb. Rel. Hist. i. 605 = C. neriifolium.
pedunculosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 112 (err. typ. 105) = C. pedunculosum.
scabra, Sw. Fl. Ind. Occ. iii. 1378 = C. surinamense.
terebinthinacea, Sw. Prod. Veg. Ind. Occ. 115 = C. terebinthinaceum.

TRIXOSTIS, Rafin. in Ser. Bull. Bot. i. (1830) 221 = **Aristida**, Linn. (Gramin.).
gracilis, Rafin. l. c. = A. gracilis.

TRIZEUXIS, Lindl. Coll. Bot. t. 2 (1823). *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
falcata, *Lindl. l. c.* = N. Granat.; Ins. Trinit.?

TROCHERA, Rich. in Journ. Phys. xiii. (1779) 225. t. 3 = **Ehrharta**, Thunb. (Gramin.).
bulbosa, Beauv. Agrost. 62. t. 12. f. 3 = E. bulbosa.
calicina, Beauv. l. c. 62. t. 12. f. 4 = E. calycina.
spicata, Poir. Encyc. Suppl. ii. 542, in syn., sphalm. = E. bulbosa.
striata, Rich. in Journ. Phys. xiii. (1779) 225. t. 3 = E. bulbosa.

TROCHETIA, DC. in Mém. Mus. Par. x. (1823) 106. tt. 7, 8. *STERCULIACEAE*, Benth. & Hook. f. i. 222.
Blackburniana, *Boj. Hort. Maurit.* 41, nomen; *et ex Baker, Fl. Maurit.* 29.—Ins. Maurit.
Boivini, Baill. Adansonia, x. (1871) 109 = Dombeya Boivini (in Addend.).
Candolleana, Boj. Hort. Maurit. 41 = uniflora.
contracta, Wall. Cat. n. 1162 (Quid?).—Ind. or
decanthera, *Benth. in Journ. Linn. Soc.* vi. (1862) 116.—Madag.
erythroxylon, Benth. l. c. = Melhania erythroxylon.

TROCHETIA:—
grandiflora, Boj. Hort. Maurit. 41; Lindl. Bot. Reg. (1844) Misc. 4. t. 21 = triflora.
laurifolia, *Benth. in Journ. Linn. Soc.* vi. (1862) 116.—Madag.
melanoxylon, Benth. l. c. = Melhania melanoxylon.
parviflora, *Boj. Hort. Maurit.* 41, nomen; *et ex Baker, Fl. Maurit.* 30.—Ins. Maurit.
pentaglossa, *Baker, in Journ. Linn. Soc.* xx. (1883) 102.—Madag.
Richardi, Baill. Adansonia, x. (1871) 108 = Dombeya Richardi (in Addend.).
serrata, Boj. Hort. Maurit. 41, nomen (Quid?).—Ins. Maurit.
Thouarsii, Baill. Adansonia, x. (1871) 110 = Dombeya Thouarsii (in Addend.).
triflora, *DC. in Mém. Mus. Par.* x. (1823) 109.—Ins. Maurit.
uniflora, *DC. l. c.* 107.—Ins. Borbon.

TROCHISANDRA, Bedd. Fl. Sylv. i. t. 120 (1871) = **Kurrimia**, Wall. (Celastr.).
indica, Bedd. l. c. = K. paniculata.

TROCHISCANTHES, Koch, in Nov. Act. Nat. Cur. xii. I. (1824) 103. *UMBELLIFERAE*, Benth. & Hook. f. i. 910.
PODOPETALUM, Gaud. Fl. Helv. ii. 401 (1828).
SCHNIZLEINIA, Steud. Nom. ed. II. ii. 532, 720 (1841).
nodiflorus, *Koch, in Nov. Act. Nat. Cur.* xii. I. (1824) 104.—Europ. austr.

TROCHISCANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 = praec.

TROCHOCARPA, R. Br. Prod. 548 (1810). *EPACRIDEAE*, Benth. in Hook. f. ii. 612.
DECASPORA, R. Br. Prod. 548 (1810).
Clarkei, *F. Muell. Fragm.* iv. 57.—Austral.
disticha, *Spreng. Syst.* i. 660.—Austral.
ericifolia, F. Muell. in Papers Roy. Soc. Tasm. (1874) 86 = Pentachondra ericaefolia.
glauca, Spreng. Syst. i. 660 = Cyathodes glauca.
Gunnii, *Benth. Fl. Austral.* iv. 167.—Tasmania.
involucrata, F. Muell. Fragm. vi. 57 = Pentachondra involucrata.
laurina, *R. Br. Prod.* 548.—Austral.
parviflora, *Benth. Fl. Austral.* iv. 167.—Austral.
pumila, F. Muell. Fragm. vi. 57 = Pentachondra pumila.
thymifolia, *Spreng. Syst.* i. 660.—Austral.
verticillata, F. Muell. in Papers Roy. Soc. Tasm. (1874) 86 = Pentachondra verticillata.

TROCHOCEPHALUS, Opiz, ex Bercht. Oek. Fl. Boehm. ii. 222 (1838) = **Scabiosa**, Tourn. (Dipsac.).
graminifolius, Opiz, ex Bercht. l. c. 223 = S. graminifolius.

TROCHODENDRON, Sieb. & Zucc. Fl. Jap. i. 83. tt. 39, 40 (1835). *MAGNOLIACEAE*, Benth. & Hook. f. i. 954.
GYMNANTHUS, Jungh. in. Hoev. & De Vriese, Tijdschr. vii. (1840) 308.
ralioides, *Sieb. & Zucc. Fl. Jap.* i. 84.—Japon.

TROCHOMERIA, Hook. f. in Benth. & Hook. f. Gen. i. 822 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
HETEROSICYOS, Welw. in Trans. Linn. Soc. xxvii. (1869) 33.
debilis, *Benth. & Hook. f. Gen.* i. 822.—Afr. austr.
Garcini, *Benth. & Hook. f. l. c.*—Afr. austr.
Hookeri, *Harv. Gen. S. Afr. Pl.* ed. II. 125 = Garcini.
macrocarpa, *Harv. l. c.*; *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 524.—Afr. trop. et austr.
macroura, *Hook. f. l. c.* 525.—Afr. trop.
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 113.—Madag.
pectinata, *Benth. & Hook. f. Gen.* i. 822.—Afr. austr.
polymorpha, *Cogn. in DC. Monog. Phan.* iii. 401.—Afr. trop.
sagittata, *Benth. & Hook. f. Gen.* i. 822.—Afr. austr.
stenoloba, *Cogn. in DC. Monog. Phan.* iii. 402.—Afr. trop.
vitifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 525.—Afr. trop.
Wyleyana, *Benth. & Hook. f. Gen.* i. 822.—Afr. austr.

TROCHOMERIOPSIS, Cogn. in DC. Monog. Phan. iii. 661 (1881). *CUCURBITACEAE*.
diversifolia, *Cogn. l. c.*—Madag.

TROCHOSERIS, Endl. Gen. 500 (1838) = **Troximon**, Nutt. (Compos.).
alpina, Poepp. & Endl. Nov. Gen. et Sp. iii. 56. t. 263 (= *Macrorhynchus Poeppigii*).—Chili.
pterocarpa, Poepp. & Endl. l. c. 57 (= *Macrorhynchus pterocarpus*).—Chili.

TROCHOSTIGMA, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 726. t. 2. f. 2 = **Actinidia**, Lindl. (Ternstr.).
argutum, Sieb. & Zucc. l. c. 727 = A. callosa.
Kolomikta, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 261 = A. callosa.
polygamum, Sieb. & Zucc. in Abh. Akad. Muench. iii. II. (1843) 728 = A. polygama.
repandum, Sieb. & Zucc. l. c. = A. polygama.
rufum, Sieb. & Zucc. l. c. 727 = A. callosa.
volubile, Sieb. & Zucc. l. c. 728 = A. volubilis.

TROLIUS, Gilib. Fl. Lituan. ii. 280 (1782), sphalm. = seq.

TROLLIUS, Linn. Sp. Pl. ed. I. 556 (1753). *RANUNCULACEAE*, Benth. & Hook. f. i. 7.
GAISSENIA, Rafin. in Med. Repos. N. York, v. (1808) 351.
HEGEMONE, Bunge, ex Ledeb. Fl. Ross. i. 51 (1842).
acaulis, *Lindl. Bot. Reg.* (1842) *Misc.* 56.—Reg. Himal.
aconitifolius, Hort. Berol. ex Steud. Nom. ed. I. 857; et ex Otto & Dietr. Allg. Gartenz. x. (1842) 186 = europaeus.
africanus, Wender. in Schrift. Marb. ii. (1831) 217 = asiaticus?
altaicus, *C. A. Mey. Verz. Pfl. Cauc.* 200.—Sibir. altaic.
altissimus, Crantz, Stirp. Austr. ed. I. fasc. ii. 123; Wender. in Flora, i. (1818) 578 = europaeus.
americanus, Ledeb. Fl. Ross. i. 734 = patulus.
americanus, *Muhl. in Trans. Am. Phil. Soc.* iii. (1793) 172, nomen; *et ex DC. Syst.* i. 313.—Am. bor.
asiaticus, *Linn. Sp. Pl.* 557.—Sibir.
aureus, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 936 = caucasicus.
aureus, Tausch, in Flora, xxii. (1839) 173 = patulus?
caucasicus, Ledeb. Fl. Alt. ii. 301 = altaicus.
caucasicus, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 265.—Reg. Cauc.
chinensis, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 77 = asiaticus.
connivens, Moench, Meth. 313 = europaeus.
dahuricus, Turcz. in Bull. Soc. Nat. Mosc. (1838) 86, nomen (*daharicus*); xv. (1842) 64 = Ledebourii, patulus.
decapetalus, Herb. Bosc, ex DC. Syst. i. 314 = americanus.
dschungaricus, *Regel, in Act. Hort. Petrop.* vii. (1880) 383.—Turkestan.
emarginatus, *G. Don, Gen. Syst.* i. 45.—Hab.?
europaeus, Falk, Beitr. ii. 203 = altaicus.
europaeus, *Linn. Sp. Pl.* 556; ed. II. 782.—Europ.; Reg. Cauc.
europaeus, Schlecht. in Linnaea, vi. (1831) 581 = patulus.
europaeus, Sobol. Fl. Petrop. 134 = asiaticus.
germanicus, Wender. in Schrift. Marb. ii. (1831) 217 = europaeus.
globosus, Lam. Fl. Fr. iii. 323 = europaeus.
grandiflorus, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 936 = caucasicus.
grandiflorus, Tausch, in Flora, xxii. (1839) 173 = europaeus?
humilis, Crantz, Stirp. Austr. ed. I. fasc. ii. 123; ed. II. 134 = europaeus.
japonicus, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 6 = asiaticus?
laxus, Salisb. in Trans. Linn. Soc. viii. (1807) 303 = americanus.
Ledebourii, *Reichb. Pl. Crit.* iii. 63.—Sibir.
lilacinus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 555.—Sibir. altaic.
Loddigesii, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 936 = asiaticus.
macropetalus, F. Schmidt, Reise Amurl. 218 = asiaticus.

TROLLIUS:—
medius, Wender. in Flora, i. (1818) 578 = europaeus.
minimus, Wender, l. c. 579 = europaeus.
montanus, Salisb. Prod. 373 = europaeus.
napellifolius, Roep. in Flora, iii. (1820) 105 = europaeus.
ochotensis, Tiling, in Regel & Tiling, Fl. Ajan. 38 = patulus.
palmatus, Baill. Hist. Pl. i. 22 = Calathodes palmata.
parviflorus, Ledeb. ex Reichb. Pl. Crit. iii. 63 = patulus.
patulus, Bieb. Fl. Taur. Cauc. iii. 387 = caucasicus.
patulus, *Salisb. in Trans. Linn. Soc.* viii. (1807) 303.—Reg. Cauc.; Persia; As. bor.
pentapetalus, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 391 = americanus.
pumilus, *D. Don, Prod. Fl. Nep.* 195.—Reg. Himal.
pumilus, Royle, Illustr. Bot. Himal. 54 = acaulis.
Riederianus, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 48 = patulus.
rotundiflorus, Stokes, Bot. Mat. Med. iii. 270 = europaeus.
sertiflorus, Salisb. in Trans. Linn. Soc. viii. (1807) 303 = asiaticus.
somcheticus, C. Koch, in Linnaea, xv. (1841) 247 = caucasicus.
sphaericus, Salisb. in Trans. Linn. Soc. viii. (1807) 302 = europaeus.
tauricus, Hort. ex Wender. in Flora, i. (1818) 579 = europaeus.
transsilvanicus, Schur, Enum. Pl. Transs. 26 = europaeus.
viridiflorus, Schur. l. c. = europaeus.
viridis, Mielichh. in Flora, xxii. (1839) 492 = europaeus.

TROMMSDORFFIA, Mart. Nov. Gen. et Sp. ii. 40. t. 139 (1826) = **Hebanthe**, Mart. (Amarant.).
argentata, Mart. l. c. 41 = Alternanthera argentata.
aurata, Mart. l. c. = H. aurata.
canescens, Mart. l. c. 42 = Iresine canescens.
pulverulenta, Mart. l. c. = H. pulverulenta, *Benth. & Hook. f.*

TROMMSDORFIA, Bernh. Syst. Verz. Erf. 102 (1800) = **Hypochoeris**, Linn. (Compos.).
maculata, Bernh. l. c. 140 = H. maculata.

TROMOTRICHE, Haw. Syn. Pl. Succ. 36 (1812) = **Stapelia**, Linn. (Asclep.).
ciliata, Sweet, Hort. Brit. ed. II. 358 = Diplocyatha ciliata.
fuscata, Haw. Suppl. Pl. Succ. 10 = S. fuscata.
glauca, Haw. Syn. Pl. Succ. 37 = S. glauca.
mutabilis, Sweet, Hort. Brit. ed. II. 358 = S. rufa.
obliqua, G. Don, in Loud. Hort. Brit. 96 = S. obliqua.
pruinosa, Haw. Syn. Pl. Succ. 37 = S. pruinosa.
revoluta, Haw. l. c. 36 = S. revoluta.

TROMSDORFFIA, Benth. & Hook. f. Gen. iii. 42 (1880), sphalm. = TROMMSDORFFIA, Mart. = **Hebanthe**, Mart. (Amarant.).

TROMSDORFFIA, Blume, Bijdr. 762 (1826) = **Chirita**, Buch.-Ham. (Gentian.).
elongata, Blume, l. c. 763.—Ins. Molucc.
speciosa, Blume, l. c. = C. Horsfieldii.

TROMSDORFFIA, R. Br. in Benn. Pl. Jav. Rar. 116 (1840) = **Dichrotrichum**, Reinw. (Gesner.).
elongata, R. Br. l. c. = D. ternateum.

TRONICENA, Steud. Nom. ed. II. ii. 721 (1841) = **Aeginetia**, Linn. (Orobanch.).
mirabilis, Steud. l. c. = A. Centronia.

TRONICEUS, Miq. Fl. Ind. Bat. ii. 713 (1856), in syn., sphalm. = praec.

TROOSTWYCKIA, Benth. & Hook. f. Gen. i. 434 (1862), sphalm. = seq.

TROOSTWYKIA, Miq. Fl. Ind. Bat. Suppl. 531 (1860). *CONNARACEAE*, Benth. & Hook. f. i. 434.
singularis, *Miq. l. c.*—Sumatra.

TROPAEOLUM, Linn. Gen. ed. I. 114 (1737). *GERANIACEAE*, Benth. & Hook. f. i. 274.
ANISOCENTRA, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 593.

TROPAEOLUM:—
CARDAMINDUM, Tourn. ex Adans. Fam. ii. 388 (1763).
CHYMOCARPUS, D. Don, in Trans. Linn. Soc. xvii. (1834) 13, 145.
MAGALLANA, Cav. Ic. iv. 50. t. 374 (1797).
RIXEA, C. Morr. in Ann. Soc. Bot. Gand, i. (1845) 225. t. 22.
TROPHAEUM, Linn. Syst. ed. I (1735).
aduncum, Sm. Tour, i. 165 = peregrinum.
albiflorum, *Lem. in Fl. des Serres*, iii. (1847) 241. —Peruv.; Chili.
atrosanguineum, *Gordon, in Gard. Chron.* (1845) 185. —Peruv.
aureum, C. Muell. in Walp. Ann. vii. 493, sphalm. = azureum.
azureum, Hook. Bot. Mag. t. 3985 = violaeflorum.
azureum, *Miers, Trav. Chile*, ii. 531, nomen; *Lindl. Bot. Reg.* sub t. 1547.—Chili.
Beuthii, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 161.—Bolivia.
bicolor, Steud. Nom. ed. II. ii. 721 = seq.
bicolorum, *Ruiz & Pav. Fl. Per.* iii. 76.—Peruv.
bogotense, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 425.—N. Granat.
brachyceras, *Hook. & Arn. Bot. Beech. Voy.* 14.—Chili.
brasiliense, *Casar. in Atti 3a Riun. Sc. Ital. Firenze*, (1841) 512.—Bras.
Bridgesii, *Hook. Journ. Bot.* i. (1834) 178; *Field. & Gardn. Sert. Pl.* t. 39.—Chili.
canariense, Hort. ex Lindl. & Moore, Treas. Bot. i. 212 = peregrinum.
canariense, C. R. Markham, Trav. Peru & Ind. 78 = Haynianum.
Chaixianum ×, E. Morr. Belg. Hortic. vi. (1856) 2.
chilense, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 47.—Chili.
chrysanthum, *Planch. & Linden, in Fl. des Serres*, x. (1854–55) 97.—N. Granat.
ciliatum, *Ruiz & Pav. Fl. Per.* iii. 77.—Peruv.
cirrhipes, *Hook. Ic. Pl.* t. 411.—Peruv.
coccineum, Miers, Trav. Chile, ii. 531, nomen = tricolorum.
crenatiflorum, *Hook. Bot. Mag.* t. 4245.—Peruv.
crenatum, *Karst. Fl. Columb.* i. 145. t. 72.—N. Granat.
Deckerianum, *Moritzi & Karst. in Karst. Ausw. Neu. Gew. Venez.* 38. t. 12.—Venezuela.
dentatifolium, Stokes, Bot. Mat. Med. ii. 346 = minus.
digitatum, *Karst. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 301.—Venezuela.
dipetalum, *Ruiz & Pav. Fl. Per.* iii. 77. t. 313.—Peruv.
edule, *Paxt. Mag. Bot.* ix. (1842) 127.—Chili.
elatum, Salisb. Prod. 275 = majus.
elegans, *G. Don, Gen. Syst.* i. 747.—Chili.
emarginatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 425.—Mexic.
Fintelmanni, *Wagen. ex Schlecht. in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 105.—N. Granat.
floribundum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 426.—Chili.
Funckii, *Turcz. l. c.* 425.—Venezuela.
Gaertnerianum, Haage & Schmidt, ex Regel, Gartenfl. xxx. (1884) t. 1146 = digitatum.
Haynianum, *Bernh. in Allg. Thuering. Gartenz.* (1843) n. XVIII. 73.—Peruv.
Heyneanum, Seem. Journ. Bot. i. (1863) 129 = praec.
Hookerianum, *Barn. in C. Gay, Fl. Chil.* i. 415.—Chili.
hybridum, Linn. Mant. i. 64 = majus.
incrassatum, *Steud. in Flora*, xxxix. (1856) 439.—Chili.
Jarrattii, *Paxt. Mag. Bot.* v. (1838) 29.—Chili?
Kingi, *Phil. in Anal. Univ. Chil.* (1872) 684.—Chili.
Klotzschii, *Warsz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 293, in textu.—Bolivia.
Lechleri, *Steud. in Flora*, xxxix. (1856) 440.—Chili.
leptophyllum, *G. Don, Gen. Syst.* i. 747.—Chili.
linearifolium, *Steud. in Flora*, xxxix. (1856) 440.—Chili.
Lobbianum, *Hort. Veitch, ex Hook. Bot. Mag.* t. 4097. —N. Granat.
longifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 423.—N. Granat.
majus, *Linn. Sp. Pl.* 345.—Peruv.
massiliense ×, E. Morr. Belg. Hortic. vi. (1856) 2.
minimum, Miers, Trav. Chile, ii. 531, nomen = brachyceras.

TROPAEOLUM:—
minus, *Linn. Sp. Pl.* 345.—Peruv.
Moritzianum, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* vi. (1838) 241.—Venezuela.
Morreanum, Klatt, in Hamb. Gartenz. xv. (1859) 219 = dipetalum.
mucronatum, *Meyen, Reise*, ii. 27.—Peruv.
Naudinii, *E. Morr. in Belg. Hortic.* vi. (1856) 2. t. 1. f. 3.—Hab.?
nubigenum, *Phil. in Linnaea*, xxxiii. (1864–65) 34.—Chili.
orthoceras, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 539 = brasiliense.
oxalianthum, *C. Morr. in Ann. Soc. Bot. Gand*, iv. (1848) 257. t. 207. f. 5.—Chili.
paniculatum, Meyen, Reise, i. 314, 342, nomen, sphalm. = mucronatum?
parviflorum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 423.—Venezuela.
peltophorum, *Benth. Pl. Hartw.* 127.—Peruv.
pendulum, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xviii. (1850) 377.—Am. centr.
pentaphyllum, *Lam. Encyc.* i. 612.—Reg. Argent.
peregrinum, *Linn. Sp. Pl.* 345.—N. Granat.
pilosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 424.—Venezuela.
pinnatum, *Andr. Bot. Rep.* viii. 535.—Peruv.
polyphyllum, *Cav. Ic.* iv. 65. t. 395.—Chili.
polyphyllum, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1348 = edule.
Popelarii, Drapiez, in Otto & Dietr. Allg. Gartenz. xv. (1847) 161 = albiflorum.
prostratum, Miers, Trav. Chile, ii. 531, nomen = polyphyllum.
pubescens, *H. B. & K. Nov. Gen. et Sp.* v. 251.—Peruv.
pulchellum, Salisb. Prod. 275 = minus.
quinatum, Hellen. Tropaeol. 20 = seq.
quinquelobum, Berg, in Vet. Acad. Handl. Stockh. (1765) 33. t. 1 = majus.
Reineckeanum ×, A. Dietr. in Otto & Dietr. Allg. Gartenz. xiv. (1846) 153.
repandifolium, Stokes, Bot. Mat. Med. ii. 346 = majus.
rhizophorum, *Warsz. in Otto & Dietr. Allg. Gartenz.* xx. (1882) 293.—Bolivia.
rhomboideum, *Lem. in Fl. des Serres*, iii. (1847) sub t. 236.—Chili.
Scheuerianum, E. Morr. in Belg. Hortic. vi. (1856) 1, in textu = majus.
Schillingii, [B. Verlot, in] Vilm. Fl. Pl. Terre, ed. II. 189 = majus.
Schlimmii, Linden, ex Planch. Fl. des Serres, vi. (1850–51) 47 = Wagnerianum.
sessilifolium, *Poepp. & Endl. Nov. Gen. et Sp.* i. 24. t. 38.—Chili.
Smithii, DC. Prod. i. 684 = peregrinum.
speciosum, *Poepp. & Endl. Nov. Gen. et Sp.* i. 22.—Chili.
subincrassatum, *Steud. Nom.* ed. II. ii. 721.—Chili.
tenellum, *G. Don, Gen. Syst.* i. 747.—Chili.
tenuirostre, *Steud. in Flora*, xxxix. (1856) 440.—Chili.
tricolor, Lindl. Bot. Reg. t. 1935 = seq.
tricolorum, *Sweet, Brit. Flow. Gard.* Ser. I. iii. t. 270. —Chili.
trilobum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 424.—Venezuela.
tuberosum, *Ruiz & Pav. Fl. Per.* iii. 77. t. 314.—Peruv.
umbellatum, *Hook. Bot. Mag.* t. 4337.—Ecuador.
violaeflorum, *A. Dietr. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 130.—Chili.
Wagnerianum, *Karst. in Otto & Dietr. l. c.* xvii. (1849) 306.—Venezuela.
Warmingianum, *Rohrb. in Mart. Fl. Bras.* xiv. ii. 227. —Bras.
Zanderi ×, A. Dietr. in Otto & Dietr. Allg. Gartenz. xxii. (1854) 370.

TROPEOLUM, Nocca, in Usteri, Ann. Bot. v. (1793) 48, sphalm. = praec.

TROPEXA, Rafin. Fl. Tellur. iv. 98 (1836) = **Aristolochia**, Tourn.
biloba, Rafin. l. c. = A. bilobata.

TROPHAEUM, Linn. Syst. ed. I (1735) = **Tropaeolum**, Linn. (Geran.).

TROPHIANTHUS, Scheidw. in Otto & Dietr. Allg. Gartenz. xii. (1844) 218 = **Aspasia**, Lindl. (Orchid.).
zonatus, Scheidw. l. c. = A. lunata.

TROPHIS, P. Br. Hist. Jamaic. 357 (1756); Linn. Syst. ed. X. 1289 (1759). *URTICACEAE*, Benth. & Hook. f. iii. 365.
BUCEPHALON, Plum. ex Linn. Gen. ed. I. 375 (1737).
aculeata, Roth, Nov. Pl. Sp. 368 = Streblus asper, Plecospermum spinosum.
americana, *Linn. Syst.* ed. X. 1289.—Ind. occ.
anthropophagorum, Seem. in Bonplandia, ix. (1861) 259; Fl. Vit. 258 (= Paratrophidis sp.).—Ins. Fiji.
aspera, Retz. Obs. v. 30 = Streblus asper.
coccinea, Zipp. ex Span. in Linnaea, xv. (1841) 335, nomen = Malaisia tortuosa.
cochinchinensis, Poir. Encyc. viii. 123 = Streblus asper.
cordata, Poir. l. c. 124 = Broussonetia papyrifera.
fruticosa, Hook. & Arn. Bot. Beech. Voy. 215 = Cudrania javanensis *vel* fruticosa.
glabrata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 314.—Mexic.
Heyneana, Wall. Cat. n. 4642 = Phyllochlamys spinosa.
Hilariana, Casar. Nov. Stirp. Bras. Dec. 80 = Sorocea Hilarii.
laurifolia, Willd. Sp. Pl. iv. 733 = Styloceras laurifolium.
mexicana, *Bur. in DC. Prod.* xvii. 253.—Mexic.
opaca, Forst. ex Bur. l. c. 236 = Paratrophis heterophylla.
Ramon, Schlecht. & Cham. in Linnaea, vi. (1831) 357 = americana.
scandens, Hook. & Arn. Bot. Beech. Voy. 214 = Malaisia tortuosa.
spinosa, Blume, Bijdr. 489 = Cudrania javanensis.
spinosa, Roxb. ex Willd. Sp. Pl. iv. 734 = Plecospermum spinosum.
spinosa, Roxb. Fl. Ind. iii. 762 = Phyllochlamys spinosa.
taxiformis, Spreng. Syst. iii. 902 = Phyllochlamys spinosa.
taxoides, Heyne, in Roth, Nov. Pl. Sp. 368 = Phyllochlamys spinosa.
zeylanica, Koen. ex Blume, Mus. Bot. Lugd. Bat. ii. 79 = Streblus asper.

TROPHOSPERMUM, Walp. Rep. i. 171 (1842), sphalm. = TAPHROSPERMUM, C. A. Mey. = **Cochlearia**, Tourn. (Crucif.).

TROPIDIA, Lindl. in Wall. Cat. n. 7386 (1831); et Bot. Reg. sub t. 1618 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 592.
CNEMIDIA, Lindl. Bot. Reg. sub t. 1618 (1833).
DECAISNEA, Lindl. in Wall. Cat. n. 7388 (1831).
GOVINDOOIA, Wight, Ic. vi. 34. t. 2090 (1853).
PTYCHOCHILUS, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 431. t. 12 B.
angulosa, *Blume, Orch. Archip. Ind.* 122.—Ind. or.; Burma.
assamica, Blume, l. c. 124 = curculigoides.
bambusifolia, *Trimen, Cat. Pl. Ceyl.* 90.—Zeylan.
curculigoides, *Lindl. in Wall. Cat.* n. 7386.—Reg. Himal.; Burma.
effusa, *Reichb. f. in Seem. Fl. Vit.* 295.—Ins. Fiji.
Govindooi, Blume, Orch. Archip. Ind. 122 = angulosa.
graminea, *Blume, l. c.* 124.—Malaya.
grandis, Hance, in Journ. Linn. Soc. xiii. (1873) 128 = Geodorum semicristatum.
pedunculata, *Blume, Orch. Archip. Ind.* 122.—Sumatra.
semilibera, Blume, l. c. = angulosa.
septemnervis, *Reichb. f. in Linnaea*, xxv. (1852) 230.—Ins. Philipp.
squamata, *Blume, Orch. Archip. Ind.* 123.—Borneo.

TROPIDOCARPUM, Hook. Ic. Pl. tt. 43, 52 (1836). *CRUCIFERAE*, Benth. & Hook. f. i. 82.
gracile, *Hook. Ic. Pl.* t. 43.—Am. bor. occ.
scabriusculum, Hook. l. c. t. 52 = gracile.

TROPIDOLEPIS, Tausch, in Flora, xii. (1829) 68 = **Chiliotrichum**, Cass. (Compos.).
diffusa, Tausch, l. c. = C. amelloideum.

TROPIDOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 265. Genus dubium *OLACINEARUM*.—Cf. Benth. & Hook. f. Gen. i. 995.
javanicum, *Turcz. l. c.*—Java.

TROPILIS, Rafin. Fl. Tellur. ii. 95 (1836) = **Dendrobium**, Sw. (Orchid.).
emulum, Rafin. l. c. = D. aemulum.

TROPITOMA, Rafin. New Fl. Am. ii. 19 (1836) = **Desmodium**, Desv. (Legumin.).
paradoxa, Rafin. l. c. (= *D. leucanthum*).—Am. bor.

TROPITRIA, Rafin. Fl. Tellur. iii. 68 (1836) = **Tradescantia**, Linn. (Commel.).
Crassula, Rafin. l. c. = Trad. Crassula.

TROPOCARPA, D. Don, ex Meissn. in DC. Prod. xiv. 423 (1857) = **Orites**, R. Br. (Proteac.).

TROS, Haw. Monog. Narciss. 5 (1831) = **Narcissus**, Tourn. (Amaryll.).
galanthifolia, Haw. l. c. = N. poculiformis.
poculiformis, Haw. l. c. = N. poculiformis.

TROSCHELIA, Klotzsch & Schomb. in Faun. Fl. Brit. Gui. 1066 (1843) = **Schieckia**, Meissn. (Haemod.).
orinocensis, Klotzsch, l. c. = S. orinocensis.

TROTULA, Comm. ex DC. Prod. iii. 90 (1828) = **Nesaea**, Comm. (Lythrar.).
Trianthis, Comm. ex DC. l. c. = N. triflora.

TROXIMON, Gaertn. Fruct. ii. 360 (1791). 'Genus male factum' Benth. & Hook. f. ii. 522.—Cf. seq.
Dandelion, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 268; Pers. Syn. ii. 360 = Krigia Dandelion.
lanatum, F. W. Schmidt, l. c. 269; Pers. l. c. = Scorzonera lanata.
virginicum, F. W. Schmidt, l. c. 268 = Krigia amplexicaulis.

TROXIMON, [Nutt. in] Fras. Cat. (1813), nn. 83, 84; Pursh, Fl. Am. Sept. ii. 505, 742 (1814). *COMPOSITAE*, Benth. & Hook. f. ii. 522.
AGOSERIS, Rafin. Fl. Ludov. 58 (1817).
AMMOGETON, Schrad. Ind. Sem. Hort. Gotting. (1833) 1.
CRYPTOPLEURA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 431.
KYMAPLEURA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) [in errat.] 455.
MACRORHYNCHUS, Less. Syn. Comp. 139 (1832).
STYLOPAPPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 431.
TROCHOSERIS, Endl. Gen. 500 (1838).
alpestre, *A. Gray, in Proc. Am. Acad.* xix. (1883) 70.—Am. bor. occ.
apargioides, A. Gray, in Bot. Calif. i. 438, partim = humile.
apargioides, *Less. in Linnaea*, vi. (1831) 501.—Calif.
aurantiacum, A. Gray, in Bot. Calif. i. 437, partim.—Nuttallii.
aurantiacum, *Hook. Fl. Bor. Am.* i. 300. t. 104.—Am. bor. occ.
barbellulatum, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. ii. 437.—Calif.
bulbosum, Rafin. Fl. Ludov. 57 (Quid?).—Am. bor. occ.
chilense, A. Gray, in Proc. Am. Acad. ix. (1876) 216 = heterophyllum.
ciliatum, [Nutt. in] Fras. Cat. (1813) n. 83, nomen (Quid?).—Am. bor.
cuspidatum, *Pursh, Fl. Am. Sept.* ii. 742.—Am. bor.
fritillarioides, [Froel. in] DC. Prod. vii. 251, sphalm. = Picrosia longifolia.
glaucum, [*Nutt. in*] *Fras. Cat.* (1813) n. 84, nomen; *Pursh, Fl. Am. Sept.* ii. 505, 742.—Am. bor. occ.
glaucum, Sims, Bot. Mag. t. 1667 = cuspidatum.
gracilens, *A. Gray, in Proc. Am. Acad.* xix. (1883) 71.—Am. bor. occ.
grandiflorum, *A. Gray, l. c.* ix. (1876) 216.—Am. bor. occ.
heterophyllum, *Greene, in Bull. Torrey Bot. Club*, x. (1883) 88.—Am. bor. occ.
humile, *A. Gray, in Proc. Am. Acad.* xix. (1883) 72.—Am. bor. occ.
laciniatum, *A. Gray, l. c.*—Am. bor. occ.

TROXIMON:—
marginatum, Nutt. Gen. Am. ii. 128 = cuspidatum.
Nuttallii, *A. Gray, in Proc. Am. Acad.* ix. (1876) 216.—Am. bor. occ.
odoratum, Rafin. Fl. Ludov. 58 (Quid?).—Am. bor.
pallidum, Rafin. l. c. 59 (Quid?).—Am. bor.
parviflorum, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 434.—Am. bor. occ.
pumilum, Nutt. l. c. = glaucum.
retrorsum, *A. Gray, in Proc. Am. Acad.* ix. (1876) 216.—Calif.
Roemerianum, Scheele, in Linnaea, xxii. (1849) 165 = Pinaropappus roseus.
roseum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 434 = glaucum.
taraxacifolium, Nutt. l. c. = glaucum.

TROXIRUM, Rafin. Sylva Tellur. 85 (1838) = **Peperomia**, Ruiz & Pav. (Piper.).
blandum, Rafin. l. c. = P. blanda.
filiforme, Rafin. l. c. = P. filiformis.
Pereskia, Rafin. l. c. = P. pereskiaefolia.
pulchellum, Rafin. l. c. = P. pulchella.
quadrifolium, Rafin. l. c. = P. reflexa.
reflexum, Rafin. l. c. = P. reflexa.
stellatum, Rafin. l. c. = P. stellata.
trifolium, Rafin. l. c. = P. trifolia.
verticillatum, Rafin. l. c. = P. pulchella.

TROXISTEMON, Rafin. Fl. Tellur. iv. 23 (1836) = **Hymenocallis**, Salisb. (Amaryll.).
distichus, Rafin. l. c. 123 = H. littoralis.
Dryandri, Rafin. l. c. (*driandrum*) = H. littoralis.
fragrans, Rafin. l. c. 23 = H. caribaea?
littorale, Rafin. l. c. = H. littoralis.
mexicanus, Rafin. l. c. 123 = H. lacera?

TROZELIA, Rafin. Sylva Tellur. 54 (1838) = **Acnistus**, Schott (Solan.).
umbellata, Rafin. l. c. = A. umbellatus.

TRUJANOA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 1 (1825). Genus dubium *EUPHORBIACEARUM*.
mexicana, D. Dietr. Syn. Pl. i. 804 = pinnata.
pinnata, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. ii. 1.—Mexic.

TRUNCARIA, DC. Prod. iii. 106 (1828) = **Miconia**, Ruiz & Pav. (Melastom.).
caryophyllaea, DC. l. c. = M. caryophyllacea.

TRYALLIS, C. Muell. in Walp. Ann. vii. 469 (1868–71), sphalm. = **Thryallis**, Mart. (Malpig.).

TRYGONANTHUS, Endl. ex Steud. Nom. ed. II. ii. 69, in syn. (1841) = **Loranthus**, Linn.

TRYMALIUM, Fenzl, in Enum. Pl. Hueg. 20 (1837). *RHAMNEAE*, Benth. & Hook. f. i. 382.
albicans, *Reiss. in Lehm. Pl. Preiss.* ii. 280.—Austral.
angustifolium, *Reiss. l. c.* 284.—Austral.
Behrii, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 274 = Spyridium subochreatum.
biauritum, Reiss. & F. Muell. l. c. 281 = Pomaderris obcordata.
bifidum, F. Muell. l. c. 282 = Spyridium bifidum.
Billardieri, *Fenzl, in Enum. Pl. Hueg.* 25.—Austral.
bilobatum, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 279 = Pomaderris obcordatum.
brevifolium, *Reiss. in Lehm. Pl. Preiss.* ii. 281.—Austral.
capsulare, G. Don, in Sweet, Hort. Brit. ed. III. 137 = Colubrina asiatica.
Daltoni, *F. Muell. Fragm.* ix. 135.—Austral.
daphnifolium, Reiss. in Lehm. Pl. Preiss. ii. 283 = ledifolium.
daphnoides, Reiss. in Linnaea, xxix. (1857–58) 278 = Spyridium spathulatum.
diversifolium, F. Muell. Second Ind. Vict. Pl. 11 = Spyridium vexilliferum.
eupatorioides, Reiss. & F. Muell. in Linnaea, xxix. (1857–58) 270 = Spyridium parvifolium.
expansum, Steud. in Lehm. Pl. Preiss. i. 185 = Billardieri.
flabellare, *F. Muell. ex Reiss. in Linnaea*, xxix. (1857–58) 281.—Austral.
floribundum, Steud. in Lehm. Pl. Preiss. i. 185 = Billardieri.

TRYMALIUM:—
globulosum, Fenzl, in Enum. Pl. Hueg. 25 = Spyridium globulosum.
halmaturinum, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 283 = Spyridium halmaturinum.
helianthemifolium, Reiss. l. c. 271 = Pomaderris ledifolia.
hermannioides, Reiss. l. c. 275 = Spyridium parvifolium.
ledifolium, *Fenzl, in Enum. Pl. Hueg.* 24.—Austral.
leucophractum, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 295 = Stenanthemum leucophractum.
Leucopogon, F. Muell. ex Reiss. l. c. 274 = Spyridium subochreatum.
majoranifolium, Fenzl, in Enum. Pl. Hueg. 21 = Spyridium spadiceum.
microphyllum, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 273 = Spyridium Lawrencii.
obovatum, Reiss. l. c. 278 = Spyridium obovatum.
oligocephalum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 460 = Spyridium oligocephalum.
parvifolium, F. Muell. ex Reiss. in Linnaea, xxix. (1857–58) 275 = Spyridium parvifolium.
phlebophyllum, F. Muell. ex Reiss. l. c. 272 = Spyridium phlebophyllum.
polycephalum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 460 = Spyridium subochreatum.
rosmarinifolium, Reiss. in Lehm. Pl. Preiss. ii. 283 = ledifolium.
serpyllaceum, Reiss. & F. Muell. in Linnaea, xxix. (1857–58) 280 = Spyridium serpyllaceum.
spadiceum, Fenzl, in Enum. Pl. Hueg. 26 = Spyridium spadiceum.
spathulatum, F. Muell. in Trans. Vict. Inst. i. (1855) 122 = Spyridium spathulatum.
stenophyllum, Reiss. in Linnaea, xxix. (1857–58) 282 = Spyridium bifidum.
Stuarti, F. Muell. ex Reiss. l. c. 289, 290 = Spyridium coactilifolium.
subochreatum, F. Muell. in Trans. Vict. Inst. i. (1855) 122 = Spyridium subochreatum.
thomasioides, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 459 = Spyridium spadiceum.
uncinatum, F. Muell. Second Ind. Vict. Pl. 11 = Spyridium eriocephalum.
velutinum, Reiss. in Linnaea, xxix. (1857–58) 276 = Spyridium obovatum.
vexilliferum, F. Muell. ex Reiss. l. c. 286 = Spyridium vexilliferum.
Wayae, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* v. (1882) 80; vi. (1883) 109.—Austral.
Wayii, F. Muell. & Tate, l. c. vi. (1883) 109 = praec.
Wendlandianum, G. Don, in Sweet, Hort. Brit. ed. III. 137 = Pomaderris ferruginea.
westringiaefolium, Reiss. in Lehm. Pl. Preiss. ii. 284 = Spyridium westringiaefolium.
Wichurae, *Nees, ex Reiss. l. c.* ii. 281.—Austral.

TRYMATOCOCCUS, Poepp. & Endl. Nov. Gen. et Sp. ii. 30. t. 142 (1838). *URTICACEAE*, Benth. & Hook. f. iii. 366.
africanus, *Baill. Adansonia*, xi. (1875) 300.—Afr. trop.
amazonicus, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 30. t. 142.—Bras.

TRYOCEPHALUM, Endl. Gen. 119 (1836) = THRYOCEPHALON, Forst. = **Kyllinga**, Rottb. (Cyperac.).

TRYPHANE, Reichb. Ic. Fl. Germ. v. 28 (1841); Nom. 205 (1841) = **Arenaria**, Rupp. (Caryoph.).
cespitosa, Reichb. l. c. = A. verna.
Facchinii, Reichb. l. c. 29.—Europ.
Gerardi, Reichb. l. c. = A. verna.
hispida, Reichb. l. c. = A. hispida.
obtusa, Reichb. l. c. 32 = A. polygonoides.
recurva, Reichb. l. c. 29 = A. recurva.
verna, Reichb. l. c. 28 = A. verna.

TRYPHERA, Blume, Bijdr. 549 (1825) = **Mollugo**, Linn. (Ficoid.).
prostrata, Blume, l. c. = M. hirta.

TRYPHIA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Holothrix**, Rich. (Orchid.).
Mac-Owaniana, Reichb. f. Otia Bot. Hamb. 108 = H. Mac-Owaniana.
major, Sond. in Linnaea, xix. (1847) 82 = H. secunda.

TRYPHIA:—
nivea, Zeyh. ex Harv. Gen. S. Afr. Pl. 323 = Holothrix Lindleyana?
orthoceras, Harv. Thes. Cap. ii. 4 = H. orthoceras.
parviflora, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209 = H. parviflora.
secunda, Lindl. Bot. Reg. sub t. 1701 = H. Lindleyana.

TRYPHOSTEMMA, Harv. Thes. Cap. i. 32. t. 51 (1859). *PASSIFLOREAE*, Benth. & Hook. f. i. 811.
Hanningtonianum, *Mast. in Hook. Ic. Pl.* t. 1484.—Afr. trop.
natalense, Mast. in Trans. Linn. Soc. xxvii. (1871) 639 = Sandersoni.
Sandersoni, *Harv. Thes. Cap.* i. 33.—Afr. austr.
zanzibaricum, *Mast. in Oliver, Fl. Trop. Afr.* ii. 508.—Afr. trop.

TRYPTOMENE, Walp. Rep. ii. 157 (1843), sphalm. = **Thryptomene**, Endl. (Myrtac.).

TSCHUDYA, DC. Prod. iii. 155 (1828) = **Oxymeris**, DC. (Melastom.).
asperiuscula, DC. l. c. = O. asperiuscula.
berbiceana, Griseb. in Goett. Abh. ix. (1861) 47 = Miconia Acinodendron.
ibaguensis, Griseb. l. c. = Miconia ibaguensis.
ionopogon, Mart. in Flora, xxiv. (1841) ii. Beibl. 37 = O. inopogon.
lanata, Griseb. Fl. Brit. W. Ind. 250 = Miconia lanata.
pulverulenta, DC. Prod. iii. 155 = O. pulverulenta.
robusta, Sagot, in Ann. Sc. Nat. Sér. VI. xv. (1883) 329 = Miconia trichodes.
rufescens, DC. Prod. iii. 155 = O. rufescens.
spondylantha, Griseb. Fl. Brit. W. Ind. 251 = Miconia spondylantha.
strigillosa, Griseb. l. c. 250 = Platycentrum clidemioides.
verticillata, DC. Prod. iii. 155 = O. rufescens.

TSIA, Adans. Fam. ii. 450 (1763) = **Camellia**, Linn. (Ternstr.).

TSIANA, J. F. Gmel. Syst. i. 9 (1791) = **Costus**, Linn. (Scitam.).
speciosa, J. F. Gmel. l. c. = C. speciosus.

TSIEMTANI, Adans. Fam. ii. 303 (1763) = **Rumphia**, Linn. (Anacard.).

TSJERUCANIRAM, Adans. Fam. ii. 80 (1763) = **Cansjera**? Juss. (Olacin.).

TSJINKIA, Adans. Fam. ii. 401 (1763) = **Lagerstroemia**, Linn. (Lythrar.).

TSUBAKI, Adans. Fam. ii. 399 (1763) = **Camellia**, Linn. (Ternstr.).

TSUGA, Carr. Conif. ed. I. 185 (1855). *CONIFERAE*, Benth. & Hook. f. iii. 440.
MICROPEUCE, Gord. Pinet. Suppl. 13 (1862).
Balfouriana, *McNab, in Journ. Linn. Soc.* xix. (1882) 211.—Am. bor. occ.
Brunoniana, *Carr. Conif.* ed. I. 188.—Reg. Himal.
canadensis, *Carr. l. c.* 189.—Am. bor.
caroliniana, *Engelm. in Coult. Bot. Gaz.* vi. (1881) 223.—Am. bor.
diversifolia, *Mast. in Journ. Linn. Soc.* xviii. (1881) 514.—Japon.
Douglasii, Carr. Conif. ed. I. 192 = Pseudotsuga Douglasii.
Hookeriana, *Carr. l. c.* ed. II. 252.—Calif.
Lindleyana, Roezl, Cat. Grain. Conif. Mex. 8 = Pseudotsuga Douglasii.
Mertensiana, *Carr. Conif.* ed. II. 250.—Am. bor. occ.
Pattoniana, *Engelm. in Gard. Chron.* (1879) ii. 756.—Am. bor. occ.
Roezlii, *Carr. in Rev. Hortic.* (1870) 217.—Calif.
Sieboldii, *Carr. Conif.* ed. I. 186.—Japon.
Tsuja, A. Murr. in Proc. Hort. Soc. Lond. ii. (1862) 508 = Sieboldii.

TSUSIOPHYLLUM, Maxim. in Mém. Acad. Pétersb. Sér. VII. xvi. (1871) n. ix. 12. *ERICACEAE*, Benth. & Hook. f. ii. 602.
Tanakae, *Maxim. l. c.*—Japon.

TSUTSUSI, Adans. Fam. ii. 164 (1763) = **Rhododendron**, Linn. (Ericac.).

TUAMINA, Alef. in Bonplandia, ix. (1861) 102 = **Vicia**, Tourn. (Legumin.).
Michauxii, Alef. l. c. = V. Michauxii.

TUBANTHERA, Comm. ex DC. Prod. ii. 30 (1825) = **Colubrina**, Rich. (Rhamn.).
Katapa, Rafin. Sylva, Tellur. 154 = C. asiatica.

TUBERARIA, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 364 = **Helianthemum**, Tourn. (Cistin.).
annua, Spach, l. c. 365 = H. guttatum.
brevipes, Willk. Ic. Pl. Hisp. ii. 79. t. 116 = H. guttatum.
bupleurifolia, Willk. l. c. 77. t. 115 = H. bupleurifolium.
discolor, Pomel, Nouv. Mat. Fl. Atl. 218 = H. discolor.
echioides, Willk. Ic. Pl. Hisp. ii. 81. t. 118 = H. echioides.
globulariaefolia, Willk. l. c. 71. t. 111 = H. globulariaefolium.
glomerata, Willk. l. c. t. 117 = H. glomeratum.
guttata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 340 = H. guttatum.
inconspicua, Willk. Ic. Pl. Hisp. ii. 78. t. 116 = H. guttatum.
macrocephala, Willk. l. c. 80. t. 117.—Hispan.
patula, Pomel, Nouv. Mat. Fl. Atl. 354 = H. patulum.
perennis, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 365 = H. Tuberaria.
variabilis, Willk. Ic. Pl. Hisp. ii. 78. tt. 112–114 = H. variabile.
villosissima, Pomel, Nouv. Mat. Fl. Atl. 218 = H. villosissimum.
viscida, Pomel, l. c. = H. glutinosum?
vulgaris, Willk. Ic. Pl. Hisp. ii. 69. t. 110 = H. vulgare.

TUBEROSA, Heist. Syst. 5 (1748) = **Polianthes**, Linn. (Amaryll.).
amica, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 430 = P. tuberosa.

TUBEROSTYLES, Benth. & Hook. f. Gen. ii. 241 (1873), err. typ. = seq.

TUBEROSTYLIS, Steetz, in Seem. Bot. Voy. Herald, 142. t. 29 (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 241.
Rhizophorae, *Steetz, l. c.*—Panama.

TUBIFLORA, J. F. Gmel. Syst. 27 (1791) = **Elytraria**, Michx. (1803) (Acanth.).
carolinensis, J. F. Gmel. l. c. = E. virgata.

TUBILIUM, Cass. in Bull. Soc. Philom. (1817) 153 = **Pulicaria**, Gaertn. (Compos.).
angustifolium, Cass. in Dict. Sc. Nat. lvi. 20 = P. inuloides.
siculum, Choisy, ex DC. Prod. v. 477 = P. sicula.

TUBOCYTISUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 358 = **Cytisus**, Linn. (Legumin.).
biflorus, Fourr. l. c. = C. biflorus.
capitatus, Fourr. l. c. = C. capitatus.
elongatus, Fourr. l. c. = C. biflorus.
supinus, Fourr. l. c. xvii. (1869) 195 = C. supinus.

TUBOPADUS, Pomel, Mat. Fl. Atl. 8 (1860) = **Prunus**, Tourn. (Rosac.).
prostratus, Pomel, l. c. = P. prostrata.

TUCKERMANNIA, Klotzsch. in Wiegm. Archiv, vii. (1841) 248 = **Corema**, D. Don (Empetr.).
Conradi, Klotzsch, l. c. = C. Conradi.

TUCKERMANNIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 363 = **Leptosyne**, DC. (Compos.).
maritima, Nutt. l. c. = L. maritima.

TUCKEYA, Gaudich. Bot. Voy. Bonite, t. 26 (1844–49) = **Pandanus**, Linn.
Candelabrum, Gaudich. l. c. f. 10–20 = P. Candelabrum.

TULA, Adans. Fam. ii. 500 (1763). *RUBIACEAE?* Adansoni, *Roem. & Schult. Syst.* iv. 355.—Peruv.

TULAKENIA, Rafin. Fl. Tellur. iv. 120 (1836) = **Jurinea**, Cass. (Compos.).
blanda, Rafin. l. c. = J. mollis.
cyanoides, Rafin. l. c. = J. cyanoides.
mollis, Rafin. l. c. = J. mollis.

TULASNEA, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 142. *MELASTOMACEAE*, Benth. & Hook. f. i. 739.
foliosa, *Naud. l. c.* 143.—Bras.
gracillima, *Naud. l. c.* t. 2.—Bras.

TULASNEA, Wight, Ic. v. tt. 1919–20 (1852) = **Terniola**, Tul. (Podostem.).
ceylanica, Wight, l. c. t. 1919 = Terniola zeylanica.
foliosa, Wight, l. c. = Terniola foliosa.
Lawii, Wight, l. c. (*Lauri*) = Terniola Lawii.
pedunculata, Wight, l. c. t. 1919 = Terniola pedunculosa.
pedunculosa, Wedd. in DC. Prod. xvii. 47, sphalm. = praec.
ramosissima, Wight, Ic. t. 1920 = Terniola ramosissima.
zeylanica, Wedd. in DC. Prod. xvii. 46 = *ceylanica*.

TULBACHIA, D. Dietr. Syn. Pl. ii. 1030 (1840), err. typ. = seq.

TULBAGHIA, Linn. Mant. ii. 148 (1771) (Tulbagia); Neck. Elem. iii. 188 (1790). *LILIACEAE*, Benth. & Hook. f. iii. 798.
OMENTARIA, Salisb. Gen. Pl. Fragm. 87 (1866).
acutiloba, *Harv. Thes. Cap.* ii. 51. t. 180.—Afr. austr.
aequinoctialis, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 246.—Afr. trop.
affinis, Link, Enum. Hort. Berol. i. 310 = alliacea.
alliacea, Ker-Gawl. in Bot. Mag. t. 806 = capensis.
alliacea, *Linn. f. Suppl.* 193; *Thunb. Prod. Pl. Cap.* 60.—Afr. austr.
australis, Link, ex Steud. Nom. ed. I. 857 (nom. delend. sec. ej. ed. II. ii. 722).
brachystemma, Kunth, Enum. Pl. iv. 483 = alliacea.
Camerooni, *Baker, in Journ. Bot.* xvi. (1878) 321.—Afr. trop.
capensis, *Linn. Mant.* ii. 223.—Afr. austr.
cepacea, *Linn. f. Suppl.* 194 (excl. syn.).—Afr. austr.
cepacea, Linn. f. l. c., partim = capensis.
cernua, *Avé-Lall. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 25.—Afr. austr.
Dregeana, *Kunth, Enum. Pl.* iv. 483.—Afr. austr.
Heisteri, Fabr. Enum. Pl. Hort. Helmst. 4 = Agapanthus umbellatus.
hypoxidea, *Sm. in Rees, Cycl.* xxxvi. n. 3; *Schult. f. Syst.* vii. 995.—Afr. austr.
inodora, Gaertn. Fruct. i. 57 = alliacea.
Ludwigiana, Harv. in Bot. Mag. t. 3547 = alliacea.
narcissifolia, Salisb. Prod. 219 = alliacea.
pulchella, Avé-Lall. in Bull. Phys.-Math. Acad. Pétersb. iii. (1844) 204 = capensis.
violacea, *Harv. in Bot. Mag.* t. 3555.—Afr. austr.

TULEXIS, Rafin. Fl. Tellur. iv. 42 (1836) = **Brassavola**, R. Br. (Orchid.).
bicolor, Rafin. l. c. 43 = B. tuberculata.

TULIPA, Linn. Syst. ed. I (1735); Gen. ed. I. 91 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 818.
LIRIACTIS, Rafin. Fl. Tellur. ii. 97 (1836).
LIRIOPOGON, Rafin. l. c. 35 (1836).
ORITHYIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 336 (1836).
PODONIX, Rafin. Fl. Tellur. iv. 28 (1836).
acrocarpa, Jord. ex Baker, in Journ. Linn. Soc. xiv. (1874) 293 = gallica.
acuminata, *Vahl, ex Hornem. Hort. Hafn.* i. 328.—Hab.?
acutiflora, DC. ex Baker, in Journ. Linn. Soc. xiv. (1874) 283 = Didieri.
acutiflora, Poir. Encyc. viii. 134 = T. Oculus-solis.
agenensis, [DC. in] Red. Lil. t. 60*, Add. = T. Oculus-solis.
Alberti, *Regel, Gartenfl.* (1877) 257; *et in Act. Hort. Petrop.* v. (1877) 264.—Turkestan.
aleppensis, Boiss. ex Regel, Gartenfl. (1873) 296; et Act. Hort. Petrop. ii. (1873) 450 = T. Oculus-solis.

TULIPA:—
aleppica, Baker, in Journ. Linn. Soc. xiv. (1874) 278 = T. Oculus-solis.
alpestris, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 120 = australis.
alpina, J. Gay, ex Baker, in Journ. Linn. Soc. xiv. (1874) 289 = pulchella.
altaica, Gebler, ex Schult. f. Syst. vii. 385, 553 = uniflora.
altaica, *Pall. ex Spreng. Syst.* ii. 63.—Sibir. altaic.
apula, Guss. ex Tenore, Fl. Neap. Syll. App. v. 12 = praecox.
aristata, *Regel, in Act. Hort. Petrop.* vi. (1879) 506.—China.
armena, Boiss. Diagn. Ser. II. iv. 99 = Gesneriana.
atheniensis, Orphan. in Bull. Congr. Bot. Pétersb. 1869 (1870) 116 = Orphanidea.
aurea, Rafin. Atl. Journ. 153 (Quid?).—Am. bor.
australis, *Link, in Schrad. Journ.* ii. (1799) 317.—Lusit.; Gall.
Beccariana, *Bicchi, Agg. Fl. Lucch.* (1860) 21, nomen; *et Desc. Nuov. Sp. Tulip.* 2, cum ic.—Ital.
Behmiana, *Regel, in Act. Hort. Petrop.* vi. (1879) 505.—Turkestan.
bicolor, Rafin. Atl. Journ. 153 (Quid?).—Am. bor.
Biebersteiniana, *Schult. f. Syst.* vii. 382.—Reg. Cauc.; Podol.
biflora, Becker, ex Nym. Consp. 723 = Biebersteiniana.
biflora, Donn, Hort. Cantab. ed. V. 75 = Clusiana.
biflora, Hort. Lund. ex Schult. f. Syst. vii. 384 = patens.
biflora, *Pall. Reise*, iii. 727.—Reg. Cauc.
Billietiana, Jord. in Ann. Soc. Linn. Lyon, N. S. v. (1858) 11 = Didieri.
bithynica, *Griseb. Spicil. Fl. Rumel.* ii. 382, in syn.—As. Min.
boeotica, *Boiss. & Heldr. Diagn.* Ser. II. iv. 99.—Graecia.
Boissieri, Regel, Gartenfl. (1873) 296 = montana.
Bonarotiana, Reboul, Nonn. Sp. Tulip. 7 = strangulata.
Borszczowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) 1. 438.—Turkestan.
brachystemon, *Regel, Gartenfl.* (1882) 323.—Turkestan.
Breyniana, Ker-Gawl. in Bot. Mag. t. 717 = australis.
Breyniana, Linn. Sp. Pl. 306 = Baeometra columellaris.
Browniana, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1172 = patens.
Brujniana, Roxb. Hort. Beng. 24, nomen = *Breyniana*?
bucharica, Merklin, ex Boiss. Fl. Orient. v. 193 = sogdiana.
Buhseana, Boiss. Diagn. Ser. II. iv. 98 = humilis.
campsopetala, *Delaun. ex Loisel. Herb. Gén. Amat.* iii. t. 172.—Hab.?
caucasica, Orsin. ex Nym. Consp. 723 = australis.
caucasica, *Otto, ex Steud. Nom.* ed. II. ii. 722.—Reg. Cauc.
Celsiana, [DC. in] Red. Lil. t. 38 = australis.
Celsiana, Henning, in Mém. Soc. Nat. Mosc. vi. (1823) 70 = Biebersteiniana.
chrysantha, *Boiss. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 279.—Persia; Ind. bor. occ.
Clusiana, [*DC. in*] *Red. Lil.* t. 37.—Europ.; Oriens.
Clusiana, Orph. ex Nym. Consp. 723 = Orphanidea.
Clusiana, Shepherd, ex Schult. f. Syst. vii. 1684 = stellata.
connivens, *Levier, Orig. Tul. Sav. Ital.* 12.—Italia.
cornuta, [Delile, in] Red. Lil. t. 445 = acuminata.
coronaria, Salisb. Prod. 236 = Gesneriana.
cretica, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 19.—Ins. Creta.
crispatula, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 211.—Persia.
crocata, Orph. in Bull. Congr. Bot. Pétersb. 1869 (1870) 116 = Orphanidea.
cuspidata, *Regel, Gartenfl.* (1884) 66.—Alger.
cuspidata, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 17.—Persia.
Cypriani, Hort. ex Steud. Nom. ed. II. ii. 722 = Clusiana.
dasystemon, *Regel, in Act. Hort. Petrop.* vi. (1879) 507.—Turkestan.
Didieri, *Jord. Obs. Pl. Crit.* i. (1846) 34.—Europ.
Didieri, Passer. in Nuov. Giorn. Bot. Ital. iii. (1871) 168 = Passeriniana.
edulis, *Baker, in Journ. Linn. Soc.* xiv. (1874) 295.—China; Japon.

TULIPA:—
Eichleri, *Regel, Gartenfl.* (1874) 193.—Turkestan.
elegans, *Hort. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 286.—Hab.?
Elwesi, Regel, Gartenfl. (1884) t. 1147 = cuspidata, *Regel*.
erythronioides, *Baker, in Journ. Bot.* xiii. (1875) 292.—China.
etrusca, *Levier, in Bull. Soc. Sc. Nat. Neuchât.* xiv. (1884) 262.—Ital.
eunanthiae, Orph. in Boiss. Diagn. Ser. II. iv. 100 = boeotica.
florentina, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 290 = sylvestris.
foliosa, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 80.—Armenia.
Foxiana, Reboul, Select. Sp. Tulip. 2 = praecox.
fragrans, *Munby, in Bull. Soc. Bot. Fr.* xiii. (1866) 256.—Alger.
Fransoniana, Parl. Nuov. Gen. e Spec. 15 = Didieri.
fulgens, *Hort. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 285.—Hab.?
gallica, *Delaun. ex Loisel. Herb. Gén. Amat.* t. 160.—Gallia.
Gesneriana, Bell. App. Fl. Pedem. 226 = Didieri.
Gesneriana, [Ker-Gawl. in] Bot. Reg. t. 380 = praecox.
Gesneriana, *Linn. Sp. Pl.* 306.—Europ. or.; Oriens.
Gesneriana, Reboul, Sp. Tulip. (1822) 6 = connivens.
glaucophylla, Fisch. ex Regel, in Act. Hort. Petrop. ii. (1873) 448 = Borszczowi.
graminifolia, Baker, in Journ. Bot. xiii. (1875) 230 = edulis.
Greigi, *Regel, Gartenfl.* (1878) 290. t. 773.—Turkestan.
Grisebachiana, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 265 = Orphanidea.
Grisebachii, Borb. l. c. xxxiii. (1883) 202 = praec.
Hageri, *Heldr. in Regel, Gartenfl.* (1874) 97. t. 790.—Graecia.
heliophthalma, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = T. Oculus-solis.
heteropetala, Ledeb. Ic. Pl. Fl. Ross. i. 21. t. 85 = uniflora.
heterophylla, *Baker, in Journ. Linn. Soc.* xiv. (1874) 295.—Mont. Altaic.
hexagonata, Borb.—Cf. Oestr. Bot. Zeitschr. xxxiii. (1883) 202 = praecox.
hispanica, Willd. ex Schult. f. Syst. vii. 380, in obs. = Clusiana.
hortensis, Gaertn. Fruct. i. 64 = Gesneriana.
hortulanorum, Wender. in Otto & Dietr. Allg. Gartenz. vi. (1838) 71 = suaveolens.
humilis, *Herb. in Lindl. Bot. Reg.* (1844) *Misc.* 30.—Persia.
hungarica, *Borb.—Cf. Oestr. Bot. Zeitschr.* xxxiii. (1883) 202.—Hungar.
iliensis, *Regel, Gartenfl.* (1879) 162. t. 975.—Turkestan.
Julia, C. Koch, in Linnaea, xxii. (1849) 225 = montana.
Kaufmanniana, *Regel, Gartenfl.* (1877) 194; *et in Act. Hort. Petrop.* v. (1877) 265.—Turkestan.
Kesselringi, *Regel, ll. cc.* 34; 637.—Turkestan.
Kolpakowskiana, *Regel, in Act. Hort. Petrop.* v. (1877) 266.—Turkestan.
Korolkowi, *Regel, l. c.* iii. 11. (1875) 295.—Turkestan.
Krauseana, *Regel, l. c.* vi. (1879) 508.—Turkestan.
laciniata, Fisch. ex Bellerm. in Ges. Naturf. Fr. Berl. Mag. vii. (1816) 67 = Gesneriana.
laciniata, Rafin. Fl. Tellur. ii. 35 = praec.?
lanata, *Regel, in Act. Hort. Petrop.* viii. (1884) 647. t. 4.—Bokhara.
Lehmanniana, Merck. ex Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 513 = chrysantha.
linifolia, *Regel, in Act. Hort. Petrop.* viii. (1884) 648.—Bokhara.
Lortetii, Jord. in Ann. Soc. Lyon, N. S. v. (1858) 9 = T. Oculus-solis.
Lownei, *Baker, in Journ. Linn. Soc.* xiv. (1874) 294.—Syria.
lurida, *Levier, Orig. Tul. Sav. Ital.* 15 (sp. dub.).—Europ.
maculata, Dulac, Fl. Hautes-Pyr. 107 = Clusiana.
maculata, *Hort. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 289.—Hab.?
maculata, Roth, Nov. Pl. Sp. 196 = australis.

TULIPA:—
maleolens, *Reboul, App. Nonnull. Sp. Tulip.* 1.—Ital.
Marshalliana, Andrz. ex Baker, in Journ. Linn. Soc. xiv. (1874) 290 = sylvestris.
Martelliana, *Levier, in Bull. Soc. Sc. Nat. Neuchât.* xiv. (1884) 245.—Europ.
mauriana, Jord. & Fourr. Ic. Fl. Eur. i. 9 = Didieri.
mauritiana, Jord. in Ann. Soc. Lyon, N. S. (1858) 12 = Didieri.
media, Agardh, ex Schult. f. Syst. vii. 379 = acuminata.
microgyna, Baker, in Journ. Linn. Soc. xiv. (1874) 292 = Biebersteiniana.
Minervae, Orph. ex Baker, l. c. 294 = Orphanidea.
montana, *Lindl. in Bot. Reg.* t. 1106.—Oriens.
montana, Rafin. Atl. Journ. 153 (Quid?).—Am. bor.
neglecta, Reboul, Select. Sp. Tul. 7 = strangulata.
Oculus-solis, Heldr. ex Baker, in Journ. Linn. Soc. xiv. (1874) 286 = boeotica.
Oculus-solis, [Ker-Gawl.] Bot. Reg. t. 204 = praecox.
Oculis-solis, Koch, Syn. Fl. Germ. ed. I. 707 = Didieri.
Oculus-solis, *St. Amans, in Rec. Soc. Agr. Agen.* i. 75.—Europ. austr.; Oriens.
odoratissima, Vis. Ort. Bot. Padov. 149 = suaveolens?
orientalis, *Levier, in Bull. Soc. Sc. Nat. Neuchât.* xiv. (1884) 263.—Europ. austr.; Reg. Cauc.
ornithogaloides, Fisch. ex Bess. in Flora, xvii. (1834) 1. Beibl. 25 = Lloydia triflora.
Orphanidea, *Boiss. ex Heldr. in Regel, Gartenfl.* (1862) 309. t. 373.—Graecia.
Orphanidesii, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1874) 294, sphalm. = praec.
Ostrowskiana, *Regel, Gartenfl.* (1884) 34.—Turkestan.
oxypetala, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) II. 81.—Tauria.
Passeriniana, *Levier, in Bull. Soc. Sc. Nat. Neuchât.* xiv. (1884) 270.—Europ.
patens, *Agardh. ex Schult. f. Syst.* vii. 384.—Sibir.
persica, Willd. ex Kunth, Enum. Pl. iv. 226 = patens.
planifolia, Jord. in Ann. Soc. Linn. Lyon, N. S. v. (1858) 13 = Didieri.
platystigma, *Jord. in Billot, Annot.* 32.—Gallia.
polychroma, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 18.—Persia.
praecox, Cav. Desc. 448 = Clusiana.
praecox, *Tenore, Fl. Nap.* i. 170.—Europ. austr.; Syria.
primulina, *Baker, in Gard. Chron.* (1882) II. 8.—Alger.
pubescens, *Willd. Enum. Hort. Berol. Suppl.* 17.—Hab.?
pudica, Rafin. Atl. Journ. 153 (Quid?).—Am. bor.
pulchella, Boiss. ex Baker, in Journ. Linn. Soc. xiv. (1874) 284 = montana.
pulchella, *Fenzl, in Kotschy, Reise Cilicisch. Taurus*, 379.—Cilicia.
pumila, Moench, Meth. 301 = sylvestris.
pumila, Tausch, in Flora, xii. (1829) 1. Erg. 47 = Biebersteiniana.
Raddii, Reboul, Nonnull. Sp. Tulip. 5 = praecox.
Regeli, *Elwes.—Cf. Gard. Chron.* (1879) I. 397.—Hab.?
repens, Fisch. ex Sweet, Brit. Flow. Gard. Ser. II. t. 97 = sylvestris.
retroflexa, *Hort. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 282.—Hab.?
rubroalba, Brot. Fl. Lusit. i. 520 = Clusiana.
saxatilis, *Sieber, ex Spreng. Syst.* ii. 63.—Ins. Creta.
scabriscapa, *Strangw. in Bot. Reg.* t. 1990.—Ital.
Schrenkii, Regel, Gartenfl. (1873) 297; et in Act. Hort. Petrop. ii. (1873) 452 = Gesneriana.
serotina, *Reboul, Sel. Sp. Tulip.* 6.—Ital.
sibirica, Patrin, ex Kunth, Enum. Pl. iv. 226 = patens.
Sibthorpiana, Sibth. & Sm. Fl. Graec. Prod. i. 229 = Fritillaria Sibthorpiana.
silvestris, Herb. Bout. ex Willk. & Lange, Prod. Fl. Hisp. i. 219 = australis.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 514.—Turkestan.
Sommierii, *Levier, Orig. Tul. Sav. Ital.* 15.—Europ.
spathulata, Bertol. Fl. Ital. iv. 85, 86 = Gesneriana.
stellata, *Hook. Bot. Mag.* t. 2762.—Reg. Himal.; Persia.
stenopetala, Delaun. ex Loisel. Herb. Gén. Amat. iii. t. 171 = acuminata.

TULIPA:—
strangulata, Heldr. ex Baker, in Journ. Linn. Soc. xiv. (1874) 286 = boeotica.
strangulata, *Reboul, Nonnull. Sp. Tulip.* 6.—Etruria.
strangwaysiana, Reboul, ex Baker, in Journ. Linn. Soc. xiv. (1874) 281 = praecox.
stricta, Stokes, Bot. Mat. Med. ii. 239 = Gesneriana.
stylosa, Fisch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 28 = uniflora.
suaveolens, *Roth, Catalect.* fasc. i. 45.—Rossia austr.
sylvestris, Bieb. Fl. Taur. Cauc. i. 270 = Biebersteiniana.
sylvestris, Falk, ex Kunth, Enum. Pl. iv. 226 = altaica.
sylvestris, Gouan, ex Baker, in Journ. Linn. Soc. xiv. (1874) 293 = australis.
sylvestris, Laxm. ex Schult. f. Syst. vii. 384 = patens.
sylvestris, *Linn. Sp. Pl.* 305.—Europ.
systola, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 17.—Persia.
tetraphylla, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 296.—Turkestan.
thianschanica, *Regel, l. c.* vi. (1879) 508.—As. centr.
Thirkeana, C. Koch, in Linnaea, xxii. (1849) 226 = Biebersteiniana.
transtagana, Brot. Fl. Lusit. i. 519 = australis.
tricolor, Becker, ex Nym. Consp. 723 = Biebersteiniana.
tricolor, C. Koch, in Tchihat. Asie Min. Bot. ii. 532 = Biebersteiniana.
tricolor, Ledeb. Ic. Pl. Fl. Ross. ii. 13. t. 135 = patens.
triphylla, *Regel, in Act. Hort. Petrop.* v. (1877) 636.—Turkestan.
turcarum, Levier, in Bull. Soc. Sc. Nat. Neuchât. xiv. (1884) 271 = suaveolens.
turcica, Griseb. Spicil. Fl. Rumel. ii. 382 = bithynica.
turcica, Kunth, Enum. Pl. iv. 221 = acuminata.
turcica, Roth, Catalect. fasc. i. 45 = sylvestris.
turkestanica, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 296.—Turkestan.
undulata, Jacquem. ex Baker, in Journ. Linn. Soc. xiv. (1874) 279 = chrysantha.
undulatifolia, *Boiss. Diagn.* Ser. I. v. 57.—As. Min.
unguiculata, Rafin. Fl. Tellur. ii. 35 (Quid?).—Hab.?
uniflora, *Bess. ex Baker, in Journ. Linn. Soc.* xiv. (1874) 295.—Sibir.
variopicta, Reboul, Select. Sp. Tulip. 7 = strangulata.
violacea, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 211.—Persia.
volgensis, Bieb. ex Eichw. Casp.-Cauc. 2 = suaveolens.
wolgensis, Auct. ex Steud. Nom. ed. II. ii. 723 = Celsiana.

TULIPASTRUM, Spach, Hist. Vég. Phan. vii. 481 (1839) = **Magnolia**, Linn.
americanum, Spach, l. c. 483 = M. acuminata.

TULIPIFERA, Herm. ex Mill. Gard. Dict. ed. VI. (1752) = **Liriodendron**, Linn. (Magnol.).
Liriodendrum, Mill. l. c. ed. VIII. = L. tulipifera.

TULISMA, Rafin. Fl. Tellur. ii. 98 (1836) = **Gesneria**, Linn.
verticillata, Rafin. l. c. = G. maculata.

TULLIA, Leavenw. in Am. Journ. Sc. Ser. I. xx. (1830) 343. t. 5 = **Pycnanthemum**, Michx. (Labiat.).
pycnanthemoides, Leavenw. l. c. = P. Tullia.

TULOCARPUS, Hook. & Arn. Bot. Beech. Voy. 298. t. 63 (1841) = **Guardiola**, Cerv. (Compos.).
mexicanus, Hook. & Arn. l. c. 299. t. 63 = G. Tulocarpus.

TULOCLINIA, Rafin. Fl. Tellur. iv. 119 (1836) = **Metalasia**, R. Br. (Asclep.).
imbricata, Rafin. l. c. = M. uniflora.

TULOPHOS, Rafin. Fl. Tellur. iii. 71 (1836) = **Brodiaea**, Sm. (Liliac.).
grandiflora, Rafin. l. c. = B. Douglasii.
laxa, Rafin. l. c. = B. laxa.

TULORIMA, Rafin. Fl. Tellur. ii. 69 (1836) = **Saxifraga**, Tourn.
Cymbalaria, Rafin. l. c. = S. Cymbalaria.

TULOTIS, Rafin. Fl. Tellur. ii. 37 (1836) = **Habenaria**, Willd. (Orchid.).
fuscescens, Rafin. l. c. = H. herbiola.
herbiola, Rafin. l. c. = H. herbiola.

TUMALIS, Rafin. Fl. Tellur. iv. 114 (1836) = **Euphorbia**, Linn.
Bojeri, Rafin. l. c. = E. Bojeri.

TUMBOA, Welw. cf. Gard. Chron. (1861) 75 = **Welwitschia**, Hook. f. (1862) (Gnetac.).
Bainesii, Hook. f. in Gard. Chron. (1861) 1008 = W. mirabilis.
strobilifera, Welw. l. c. (1862) 71 = W. mirabilis.

TUMELAIA, Rafin. Fl. Tellur. iv. 105 (1836) = **Daphne**, Tourn. (Thymel.).

TUNGA, Roxb. Fl. Ind. ed. Carey, i. 187 (1820) = **Hypolytrum**, Rich. (Cyperac.).
diandra, Roxb. l. c. 188 = H. latifolium, *Rich.*
laevigata, Roxb. l. c. = Lipocarpha argentea.
triceps, Roxb. l. c. 187 = Lipocarpha sphacelata.

TUNICA, Hall. Enum. Stirp. Helv. i. 381 (1742); Scop. Fl. Carn. ed. II. i. 298 (1772). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 145.
DIANTHELLA, Clauson, ex Pomel, Mat. Fl. Atl. 9 (1860).
FIEDLERIA, Reichb. Nom. 206 (1841); Ic. Fl. Germ. vi. t. 246 (1844).
KOHLRAUSCHIA, Kunth, Fl. Berol. i. 108 (1838).
arabica, *Boiss. Diagn.* Ser. I. viii. 62.—Arab.
arenaria, Scop. Fl. Carn. ed. II. i. 301 = Dianthus monspessulanus
arenicola, Nym. Consp. 100 = T. Saxifraga.
artegioides, Auct. = ortegioides.
barbata, Scop. Fl. Carn. ed. II. i. 298 = Dianthus barbatus.
bicolor, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 10 = T. Saxifraga.
brachypetala, Jaub. & Spach, Illustr. i. 10 = pachygona.
Carthusianorum, Scop. Fl. Carn. ed. II. i. 299 = Dianthus Carthusianorum.
Caryophyllus, Scop. l. c. 301 = Dianthus Caryophyllus.
ciliata, Dulac, Fl. Hautes-Pyr. 260 = T. Saxifraga.
compressa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 50.—Afr. bor.
cretica, *Fisch. & Mey. l. c.* 49.—Ins. Creta.
dianthoides, *Fisch. & Mey. l. c.* 48.—Ins. Creta.
dianthoides, Rafin. Fl. Tellur. ii. 54 = Silene dianthoides.
erecta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 10 = T. Saxifraga.
Gasparrini, *Nym. Consp.* 100.—Sicil.
glumacea, *Boiss. Fl. Orient.* i. 517.—Dalmat.; Graecia.
graminea, *Boiss. Diagn.* Ser. I. viii. 60.—Graecia.
hispidula, *Boiss. & Heldr. l. c.* 62.—As. Min.
illyrica, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 49.—Europ. austr.
Morrisii, Walp. Ann. ii. 101 = Dianthus Morrisii.
ochroleuca, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 49.—Graecia.
olympica, *Boiss. Diagn.* Ser. I. viii. 61.—As. Min.
Orfanidesiana, Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 253 = thessala.
ortegioides, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 36 (artegioides).—As. Min.
pachygona, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 50.—As. Min.; Persia.
pamphylica, *Boiss. & Bal. Diagn.* Ser. II. vi. 27.—Pamphylia.
plumaria, Scop. Fl. Carn. ed. II. i. 300 = Dianthus plumarius.
prolifera, *Scop. l. c.* 299.—Europ.; Reg. Cauc.
rigida, Rafin. Fl. Tellur. ii. 54 = Dianthus rigidus?
rigida, Reichb. Ic. Fl. Germ. v. 42 = T. Saxifraga.
Saxifraga, *Scop. Fl. Carn.* ed. II. i. 300.—Europ.; Reg. Cauc.
Sibthorpii, *Boiss. Diagn.* Ser. I. viii. 61.—Graecia.
stricta, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 50.—As. Min.; Persia; Sibir.
syriaca, *Boiss. Diagn.* Ser. I. viii. 63.—Syria.
thessala, *Boiss. l. c.*—Graecia.
velutina, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 66.—Europ. austr.; Reg. Cauc.

TUNICA:—
virginea, Scop. Fl. Carn. ed. II. i. 302 = Dianthus virgineus.
xerophila, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 10 = T. Saxifraga.
xylorrhiza, Boiss. in Ann. Sc. Nat. Sér. IV. i. (1854) 246 = Gypsophila ortegioides.

TUPA, G. Don, Gen. Syst. iii. 700 (1834) = **Lobelia**, Plum. (Campan.).
acuminata, A. DC. in DC. Prod. vii. 396 = L. acuminata.
arguta, G. Don, Gen. Syst. iii. 700 = Siphocampylus giganteus.
assurgens, A. DC. in DC. Prod. vii. 394 = L. assurgens.
atropurpurea, Vis. in Nuov. Sagg. Padov. vi. (1847) 95 = L. fulgens?
Berterii, A. DC. in DC. Prod. vii. 392 = L. serrata.
Besseriana, A. DC. l. c. 393 = L. polyphylla.
blanda, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 308.—Chili.
bracteosa, A. DC. in DC. Prod. vii. 393 = L. polyphylla.
Bridgesii, A. DC. l. c. 394 = L. Bridgesii.
caudata, Griseb. Fl. Brit. W. Ind. 386.—Ind. occ.
Cavanillesiana, G. Don, Gen. Syst. iii. 700 = L. mucronata, *Cav.*
cirsiifolia, A. DC. in DC. Prod. vii. 395 = L. cirsiifolia.
columnaris, Vatke, in Linnaea, xxxviii. (1874) 725 = L. columnaris.
conglobata, A. DC. in DC. Prod. vii. 395 = L. conglobata.
coquimbana, Phil. ex Vatke, in Linnaea, xxxviii. (1874) 726 = L. polyphylla.
costaricana, Planch. & Oerst. in Kjoeb. Vidensk. Meddel. (1857) 154.—Costa Rica.
crassicaulis, Hook. Bot. Mag. t. 4505.—Hab.?
Deckenii, Aschers. in Bot. Zeit. xxvii. (1869) 71; Decken, Reisen Ost-Afrika, iii. 74 = L. Deckenii.
digitalifolia, Griseb. Fl. Brit. W. Ind. 387.—Ind. occ.
domingensis, Vatke, in Linnaea, xxxviii. (1874) 728.—Ins. S. Doming.
ensifolia, A. DC. in DC. Prod. vii. 396.—Ins. S. Doming.
Feuillei, G. Don, Gen. Syst. iii. 700 = L. Tupa.
flavescens, A. DC. in DC. Prod. vii. 395 = L. conglobata.
hyssopifolia, A. DC. l. c. 392 = L. hyssopifolia.
ignescens, Payer, Organog. t. 149 = L. fulgens?
imberbis, Griseb. in Mem. Am. Acad. N. S. viii. (1862) 516.—Ind. occ.
infesta, Griseb. Fl. Brit. W. Ind. 387.—Ind. occ.
Kerstenii, Vatke, in Linnaea, xxxviii. (1874) 725 = L. Deckenii.
laxiflora, Planch. & Oerst. in Kjoeb. Vidensk. Meddel. (1857) 154 = L. laxiflora.
Martagon, Griseb. Fl. Brit. W. Ind. 386.—Ind. occ.
montana, Phil. in Anal. Univ. Chil. (1873) 506.—Chili.
montana, Wright, ex Griseb. Cat. Pl. Cub. 159.—Cuba.
mucronata, A. DC. in DC. Prod. vii. 392 = L. mucronata, *Cav.*
obovata, G. Don, Gen. Syst. iii. 700 = L. incana.
ovata, G. Don, l. c.—Peruv.
ovata, Phil. in Anal. Univ. Chil. (1873) 507.—Chili.
persicaefolia, A. DC. in DC. Prod. vii. 395 = L. persicaefolia.
polyphylla, G. Don, Gen. Syst. iii. 700 = L. polyphylla.
portoricensis, Vatke, in Linnaea, xxxviii. (1874) 727.—Porto Rico.
purpurea, G. Don, Gen. Syst. iii. 700 = L. purpurea.
racemosa, A. DC. in DC. Prod. vii. 394 = L. racemosa, *Sims.*
Rhynchopetalum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 9 = L. Rhynchopetalum.
robusta, A. DC. in DC. Prod. vii. 294 = L. robusta.
salicifolia, G. Don, Gen. Syst. iii. 700 = Siphocampylus giganteus.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 10 = L. Gibberoa.
secunda, G. Don, Gen. Syst. iii. 700 = L. secunda.
sonchifolia, Griseb. Fl. Brit. W. Ind. 388 = L. sonchifolia.
stricta, A. DC. in DC. Prod. vii. 395 = L. stricta.
subdentata, A. DC. l. c. 393 = L. polyphylla.

TUPEIA, Blume, Fl. Jav. Loranth. tt. 27, 28 (1830?) = **Henslowia**, Blume (Santal.).
microphylla, Chalon, in Mém. Soc. Hainaut, Sér. III. v. (1870) 324 = Lepidoceras Kingii.
myrtifolia, Chalon, l. c. = Loranthus myrtifolius.
racemosa, Korth. in Bull. Néerl. (1838) 157 = H. Reinwardtiana.
Reinwardtiana, Blume, Fl. Jav. Loranth. t. 28; et ex Korth. in Verh. Batav. Gen. xvii. (1839) 195 = H. Reinwardtiana.
squamifera, Chalon, in Mém. Soc. Hainaut, Sér. III. v. (1870) 324 = Lepidoceras squamifer.
umbellata, Blume, Fl. Jav. Loranth. t. 27; et ex Korth. in Verh. Batav. Gen. xvii. (1839) 195 = H. umbellata.

TUPEIA, Cham. & Schlecht. in Linnaea, iii. (1828) 208. *LORANTHACEAE*, Benth. & Hook. f. iii. 216.
antarctica, *Cham. & Schlecht. l. c.*—N. Zel.
Cunninghami, Miq. l. c. xviii. (1844) 85 = antarctica.
pubigera, Miq. l. c. 86 = antarctica.
undulata, Colenso, in Trans. N. Z. Inst. xvi. (1884) 329 = antarctica.

TUPELO, Adans. Fam. ii. 80 (1763) = **Nyssa**, Gronov. (Cornac.).

TUPIDANTHUS, Hook. f. & Thoms. in Hook. Bot. Mag. t. 4908 (1856). *ARALIACEAE*, Benth. & Hook. f. i. 947.
calyptratus, *Hook. f. & Thoms. in Hook. Bot. Mag.* t. 4908.—Reg. Himal.
Pueckleri, *C. Koch, in C. Koch & Fint. Wochenschr.* ii. (1859) 348, cum f.—Java.

TUPISTRA, Ker-Gawl. in Bot. Mag. t. 1655 (1814). *LILIACEAE*, Benth. & Hook. f. iii. 772.
CAMPYLANDRA, Baker, in Journ. Linn. Soc. xiv. (1875) 582. t. 20.
MACROSTIGMA, Kunth, Ind. Sem. Hort. Berol. (1848) 12.—Cf. Ann. Sc. Nat. Sér. III. xi. (1849) 220; Enum. Pl. v. 319 (1850).
PLATYMETRA, Noronha, ex Salisb. Gen. Pl. Fragm. 9 (1866).
TILCUSTA, Rafin. Fl. Tellur. iv. 15 (1836).
aurantiaca, *Wall. Cat.* n. 5194.—Reg. Himal.
Macrostigma, *Baker, in Hook. Bot. Mag.* t. 6280.—Reg. Himal.
nutans, *Wall. in Bot. Reg.* t. 1223.—Reg. Himal.
singapureana, *Wall. Cat.* n. 5195.—Singapore.
squalida, Baker, in Journ. Linn. Soc. xiv. (1875) 580, partim = T. Macrostigma.
squalida, *Ker-Gawl. in Bot. Mag.* t. 1655.—Ins. Amboina.
Stoliczana, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 199.—Burma.

TURARIA, Spreng. Syst. ii. 260 (1825) = **Thuraria**, Molina (Inc. Sed.).

TURBINA, Rafin. Fl. Tellur. iv. 81 (1836) = **Ipomoea**, Linn. (Convolv.).
corymbosa, Rafin. l. c. = I. Burmanni.

TURBITH, Tausch, in Flora, xvii. (1834) 343 = **Athamanta**, Linn. (Umbell.).
annua, Tausch, l. c. = A. cretensis.
Matthioli, Tausch, l. c. = A. Matthioli.

TURCZANINOVIA, DC. Prod. v. 257 (1836) = **Aster**, Tourn. (Compos.).
fastigiata, DC. l. c.; Coll. Mém. ix. 20. t. 4 = A. fastigiatus.

TURETTA, Vell. Fl. Flum. 233 (1825) (Quid?).
racemosa, Vell. l. c. 234; v. t. 115.—Bras.

TURGENIA, Hoffm. Gen. Umbell. 59 (1814) = **Caucalis**, Linn. (Umbell.).
brachyacantha, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 52 = C. latifolia.
foeniculacea, Fenzl, Pugill. Pl. Nov. Syr. 18.—As. Min.; Persia.
heterocarpa, DC. Prod. iv. 218.—Persia.
latifolia, Hoffm. Gen. Umbell. 59 = C. latifolia.
multiflora, DC. Prod. iv. 218 = C. latifolia.
tuberculata, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 52 = C. latifolia.

TURGENIOPSIS, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 53 = **Caucalis**, Linn. (Umbell.)
foeniculacea, Boiss. l. c.—Syria.

TURGOSEA, Haw. Rev. Pl. Succ. 14 (1821) = **Crassula**, Dill.
aloides, Haw. l. c. 16 = C. alooides.
capitella, Haw. l. c. 17 = C. capitella.
linguaefolia, Haw. l. c. 14 = C. linguaefolia.
obovata, Haw. l. c. 16 = C. obovata.
pertusa, Haw. l. c. 14 = C. pertusa.
thyrsiflora, Haw. l. c. 17 = C. thyrsiflora.
tomentosa, Haw. l. c. 14 = C. tomentosa.
turrita, Haw. l. c. 17 = C. turrita.

TURIA, Forsk. Fl. Aegypt. Arab. 165 (1775) = **Luffa**, Tourn. (Cucurb.).
cordata, J. F. Gmel. Syst. 403 = L. aegyptiaca.
cylindrica, J. F. Gmel. l. c. = L. aegyptiaca.
Gijef, Forsk. Aegypt. Arab. 166 = Corallocarpus Gijef.
Leloja, Forsk. l. c. 165 (Quid?).—Arab.
Moghadd, Forsk. l. c. 166 = Coccinia Moimoi.
sativa, Forsk. l. c. p. cxxi. = L. aegyptiaca.

TURNERA, Plum. ex Linn. Gen. ed. I. 87 (1737). *TURNERACEAE*, Benth. & Hook. f. i. 806.
BOHADSCHIA, Presl, Rel. Haenk. ii. 98. t. 68 (1831).
PUMILEA, P. Br. Hist. Jamaic. 188 (1756).
TRIACIS, Griseb. in Mem. Am. Acad. N. S. viii. (1860) (Pl. Wright. Cub.) 189.
TRIBOLACIS, Griseb. Fl. Brit. W. Ind. 297 (1860).
acaulis, *Griseb. Cat. Pl. Cub.* 114.—Cuba.
acuta, Spreng. Syst. i. 940 = ulmifolia.
acuta, *Willd. ex Schult. Syst.* vi. 678.—Bras.
alba, Liebm. in Ann. Sc. Nat. Sér. III. ix. (1848) 318 = ulmifolia.
albicans, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 135.—Bras.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = ulmifolia.
annularis, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 124.—Bras.
aphrodisiaca, L. F. Ward, Virgin. Med. Monthly (April 1876), 49; ex Journ. Bot. xviii. (1880) 20 = diffusa.
apifera, DC. Prod. iii. 346, sphalm. = opifera.
arcuata, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 146.—Bras.
arenaria, Spruce, ex Urban, l. c. 115 = melochioides.
aspera, Poir. Encyc. viii. 144 = Piriqueta villosa.
aturensis, D. Dietr. Syn. Pl. ii. 1008 = Piriqueta villosa.
aurantiaca, *Benth. in Hook. Journ. Bot.* iv. (1842) 116.—Guian.
aurea, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 226 = Piriqueta aurea.
Benthamiana, Schomb. Faun. Fl. Brit. Gui. 1166 = velutina.
Berneriana, Tul. in Ann. Sc. Nat. Sér. V. ix. (1868) 322 = Piriqueta Berneriana.
betonicaefolia, Salisb. Prod. 174 = Piriqueta villosa.
Blanchetiana, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 130.—Bras.
brasiliensis, *Willd. ex Schult. Syst.* vi. 678.—Bras.
callosa, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 105.—Mexic.
calyptrocarpa, *Urban, l. c.* 128.—Bras.
capensis, Harv. in Harv. & Sond. Fl. Cap. ii. 599 = Piriqueta capensis.
capitata, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 215.—Bras.
caroliniana, S. Wats. Bibliog. Ind. N. Am. Bot. 391 = Piriqueta fulva.
carpinifolia, H. B. & K. Nov. Gen. et Sp. vi. 125 = acuta.
cearensis, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 100.—Bras.
chamaedrifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 221.—Bras.
Chamaedrys, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1166, nomen = ulmifolia.
cistoides, Baill. Hist. Pl. iv. 287 = ulmifolia.
cistoides, Linn. Sp. Pl. ed. II. 387 = Piriqueta villosa.
cistoides, Pursh, Fl. Am. Sept. i. 206 = Piriqueta fulva.
Clausseniana, *Urban, in Eichl. Fahrb. Berl.* ii. (1883) 89.—Bras.
coerulea, [Moç. & Sesse,] ex DC. Prod. iii. 346 = ulmifolia.
corchorifolia, Willd. ex Schult. Syst. vi. 678 = odorata.

TURNERA:—
corchoroides, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1166, nomen = ulmifolia.
coriacea, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 144. —Bras.
cuneifolia, Juss. ex Urban, l. c. 139 = seq.
cuneiformis, Juss. ex Poir. Encyc. viii. 142 = ulmifolia.
curassavica, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 118.—N. Granat.
decipiens, Baill. Adansonia, x. (1872) 246 = dichotoma.
Desvauxii, DC. Prod. iii. 348 = hirta.
dichotoma, *Gardn. ex Hook. Ic. Pl.* t. 522.—Bras.
diffusa, *Willd. ex Schult. Syst.* vi. 679.—Bras.
dolichostigma, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 106.—Bras.
Duarteana, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 223 = Piriqueta Duarteana.
elegans, [Otto, ex] Nees, Hor. Phys. Berol. 36 = ulmifolia.
elliptica, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 107. —Bras.
frutescens, Aubl. Pl. Gui. i. 290. t. 113. f. 2 = rupestris.
frutescens, Mill. ex Steud. Nom. ed. I. 858 = ulmifolia.
genistoides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 220. t. 121.—Bras.
glabra, DC. Prod. iii. 347 = Piriqueta fulva.
Glaziovii, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 93. —Bras.
guianensis, *Aubl. Pl. Gui.* i. 291. t. 114.—Guian.
hebepetala, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 127.—Bras.
helianthemoides, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 224 = Piriqueta villosa.
hermannioides, *Cambess. l. c.* 218.—Bras.
hexandra, Spreng. Syst. iv. Cur. Post. 124 = odorata.
Hilaireana, Urban, in Eichl. Jahrb. Berl. ii. (1883) 108 = lanceolata.
Hindsiana, *Benth. Bot. Voy. Sulph.* 101.—Ecuador.
Hindsiana, Hemsl. Biol. Centr. Am. Bot. i. 474 = panamensis.
hirsuta, Bert. ex DC. Prod. iii. 347 = Piriqueta villosa.
hirsutissima, Wright, in Sauv. Fl. Cub. 55 = T. Pumilea.
hirta, *Desv. in Ham. Prod. Pl. Ind. Occ.* 33.—Guian.
hirta, Willd. ex Schult. Syst. vi. 678 = Piriqueta villosa.
Humboldtii, Spreng. Syst. i. 941 = guianensis.
humifusa, Endl. ex Walp. Rep. ii. 230 = diffusa.
incana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 217. —Bras.
integrifolia, Willd. ex Schult. Syst. vi. 679 = Piriqueta fulva.
lamiifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 213.—Bras.
lanceolata, *Cambess. l. c.* 214.—Bras.
longiflora, *Cambess. l. c.* 216.—Bras.
longipes, *Triana, ex Urban, in Eichl. Jahrb. Berl.* ii. (1883) 99.—Venezuela.
lucida, *Urban, l. c.* 137.—Bras.
lutescens, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 217 = opifera.
macrophylla, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 95.—Bras.
Melochia, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvii. (1873) 187.—N. Granat.
melochioides, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 219.—Bras.
microphylla, Desv. in Ham. Prod. Pl. Ind. Occ. 33 = diffusa.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 126 = ulmifolia.
nana, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 219.— Bras.
nervosa, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 108.—Paraguay.
oblongifolia, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 215 = lanceolata.
obtusifolia, Sm. in Rees, Cycl. xxxvi. n. 3 = ulmifolia.
odorata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 107.—Guian.
odorata, Vahl, ex DC. Prod. iii. 346 = ulmifolia.
opifera, Benth. in Hook. Journ. Bot. iv. (1842) 115 = ulmifolia.
opifera, *Mart. Reise Bras.* i. 552.—Bras.
ovata, Bello & Espinosa, in Anal. Soc. Esp. Hist. Nat. x. (1881) 275 = Piriqueta ovata.

TURNERA:—
panamensis, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 92.—Panama.
paniculata, Willd. in Schult. Syst. vi. 678 = Melochia graminifolia.
parviflora, Benth. in Hook. Journ. Bot. iv. (1842) 116 = odorata.
pernambucensis, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 133.—Bras.
peruviana, Willd. ex Schult. Syst. vi. 679 = ulmifolia.
pinifolia, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 220. —Bras.
pinnatifida, Juss. ex Poir. Encyc. viii. 144 = sidoides.
Pohliana, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 104. —Bras.
procumbens, Gardn. ex Hook. Ic. Pl. sub t. 522 = genistoides.
pumila, Hill, Veg. Syst. xiv. 6, sphalm. = seq.
Pumilea, *Linn. Syst.* ed. X. 965.—Am. trop.
Pumilea, Poir. Encyc. viii. 143 = diffusa.
racemosa, Jacq. Hort. Vindob. iii. 49 = Piriqueta racemosa.
ramosissima, Spruce, ex Urban, in Eichl. Jahrb. Berl. ii. (1883) 116 = melochioides.
refracta, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1166, nomen = ulmifolia.
Riedeliana, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 110.—Bras.
rosea, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 225 = Piriqueta rosea.
rugosa, Willd. Sp. Pl. i. 1504 = Piriqueta villosa.
rupestris, *Aubl. Pl. Gui.* i. 289. t. 113. f. 1.—Guian.
salicifolia, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 227 = serrata.
salicifolia, Seem. Bot. Voy. Herald, 130 = panamensis.
Schomburgkiana, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 132.—Guian.
sedoides, DC. Prod. iii. 347 = sidoides.
sericea, H. B. & K. Nov. Gen. et Sp. vi. 127 = ulmifolia.
serrata, *Vell. Fl. Flum.* 128; iii. t. 108.—Bras.
setosa, Sm. in Rees, Cycl. xxxvi. n. 6 = sidoides.
sidaefolia, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 227 = Piriqueta sidifolia.
sidoides, *Linn. Mant.* i. 58.—Bras.
sidoides, Vell. Fl. Flum. 127; iii. t. 107 (Quid?).— Bras.
stachydifolia, *Urban & Rolfe, in Eichl. Jahrb. Berl.* ii. (1883) 122.—Bras.
stipularis, *Urban, l. c.* 131.—Bras.
subglabra, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 984, nomen = aurantiaca.
subulata, Sm. in Rees, Cycl. xxxvi. n. 2 = ulmifolia.
surinamensis, Miq. in Linnaea, xviii. (1844) 748 = ulmifolia.
tomentosa, H. B. & K. Nov. Gen. et Sp. vi. 125 = odorata.
tomentosa, Willd. ex Schult. Syst. vi. 678 = Piriqueta villosa.
tomentosa, Wood, Am. Bot. & Flor. 129 = Piriqueta fulva.
trigona, *Urban, in Eichl. Jahrb. Berl.* ii. (1883) 111. —Bras.
trioniflora, Sims, Bot. Mag. t. 2106 = ulmifolia.
ulmifolia, *Linn. Sp. Pl.* 271.—Am. trop.
velutina, *Benth. in Hook. Journ. Bot.* iv. (1842) 116. —Guian.
velutina, Presl, Rel. Haenk. ii. 44 = ulmifolia.
villosa, Raeusch. Nom. ed. III. 86 = Piriqueta villosa.
virgata, Willd. ex Schult. Syst. vi. 678 = ulmifolia.
viscosa, Sauvalle, Fl. Cub. 55 = Piriqueta viscosa.
Weddelliana, *Urban & Rolfe, in Eichl. Jahrb. Berl.* ii. (1883) 90.—Paraguay.
xanthotricha, Shuttlew. ex Urban, l. c. 71 = Piriqueta fulva.
zanthotricha, Shuttlew. ex Urban, l. c. = praec.

TURPINIA, Humb. & Bonpl. Pl. Aequin. i. 113. t. 33 (1808) = **Barnadesia**, Mutis (Compos.).
laurifolia, Humb. & Bonpl. l. c. (= *Fulcaldea laurifolia*).—Peruv.

TURPINIA, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 24 (1824) = **Vernonia**, Schreb. (Compos.).
tomentosa, Lex. l. c. = V. Monosis.

TURPINIA, Pers. Syn. ii. 314 (1807) = **Poiretia**, Vent. (Legumin.).
montevidensis, Spreng. Syst. iv. Cur. Post. 293 = P. psoraloides.
punctata, Pers. Syn. ii. 314 = P. punctata.

TURPINIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Rhus**, Tourn. (Anacard.).
aromatica, Rafin. ex Desv. Journ. Bot. ii. (1809) 170 = R. aromatica.
glabra, Rafin. in Med. Repos. N. York, v. (1808) 352 = R. glabra.
pubescens, Rafin. l. c. = R. pubescens?
suaveolens, Rafin. l. c. = R. aromatica.

TURPINIA, Vent. Choix, 31. t. 31 (1803). *SAPINDACEAE*, Benth. & Hook. f. i. 413.
Dalrympelea, Roxb. Hort. Beng. 17 (1814); Pl. Corom. iii. 76. t. 279 (1819).
Eyrea, Champ. in Hook. Kew Journ. iii. (1851) 331.
Lacepedea, H. B. & K. Nov. Gen. et Sp. v. 142. t. 144 (1821).
Ochranthe, Lindl. Bot. Reg. t. 1819 (1836).
Triceraia, Willd. ex Roem. & Schult. Syst. iv. 803 (1819).
arguta, *Seem. Bot. Voy. Herald*, 371.—China.
corymbosa, G. Don, Gen. Syst. ii. 3 = occidentalis.
glandulosa, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 250.—Porto Rico.
insignis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 296.—Mexic.
latifolia, Wall. Cat. n. 4939 = pomifera.
martabanica, Wall. Cat. n. 4278 = pomifera.
megaphylla, *Tul. in Ann. Sc. Nat.* Sér. III. vi. (1846) 361.—N. Granat.
microcarpa, Wight & Arn. Prod. 156 = pomifera.
montana, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 182 = pomifera.
nepalensis, Wall. Cat. n. 4277 = pomifera.
occidentalis, *G. Don, Gen. Syst.* ii. 3.—Ind. occ.
paniculata, *Vent. Choix*, 31.—Ind. occ.; Mexic.
pinnata, *Hemsl. Biol. Centr. Am. Bot.* i. 216.— Mexic.
pomifera, *DC. Prod.* ii. 3.—As. trop.
Schiedeana, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 296.—Mexic.
sphaerocarpa, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 42.—Java.

TURPITHUM, Rafin. Fl. Tellur. iv. 71 (1836) = **Ipomoea**, Linn. (Convolv.).
australe, Rafin. l. c. = I. Turpethum.
indicum, Rafin. l. c. = I. Turpethum.

TURRAEA, Linn. Mant. ii. 150 (1771). *MELIACEAE*, Benth. & Hook. f. i. 331.
Rutaea, M. Roem. Syn. Hesper. 93 (1846).
Scyphostigma, M. Roem. l. c. 80 (1846).
Stephanosiphon, Boiss. ex C. DC. in DC. Monog. Phan. i. 446 (1878).
abyssinica, *Hochst. in Flora*, xxiv. (1841) I. *Intell.* 29. —Abyss.
alata, Wight & Arn. Prod. 117; Wight, Ic. t. 90 = Naregamia alata.
americana, *Vell. Fl. Flum.* 175; iv. t. 100.—Bras.
Billardieri, Juss. in Mém. Mus. Par. xix. (1830) 218 = pubescens.
Boivini, *Baill. Adansonia*, xi. (1873–76) 253.— Madag.
Brownii, *C. DC. in DC. Monog. Phan.* i. 442.— Austral.
concinna, *Benn. Pl. Jav. Rar.* i. 182.—Java.
decandra, Blanco, Fl. Filipp. ed. I. 347 = Dysoxylum Blancoi.
floribunda, *Hochst. in Flora*, xxvii. (1844) 297.— Afr. austr.
Fockei, *Buchen. in Bremen, Abh.* vii. (1880) 14.— Madag.
glabra, Cav. Diss. vii. 360 = maculata.
graciliflora, Schlecht. in Linnaea, xxv. (1852) 217 = heterophylla.
herbacea, Poir. Encyc. viii. 147 = Schwenkia brasiliensis.
heterophylla, Harv. & Sond. Fl. Cap. i. 245 = floribunda.
heterophylla, *Sm. in Rees, Cycl.* xxxvi. n. 6.—Afr. trop.

TURRAEA :—
Hildebrandtii, *O. Hoffm. Sert. Pl. Madag.* 10.—Madag.
indica, *C. DC. in DC. Monog. Phan.* i. 439.—Ind. or.
Kindtii, *Buchen. in Bremen, Abh.* vii. (1880) 15.—Madag.
lanceolata, Cav. Diss. vii. 361. t. 205. f. 1 = Quivisia lanceolata.
lobata, Lindl. Bot. Reg. (1843) Misc. 61 = heterophylla.
maculata, *Sm. Ic. Ined.* t. 11.—Madag.
mombassana, *Hiern, ex C. DC. in DC. Monog. Phan.* i. 439.—Afr. trop.
nilotica, *Kotschy & Peyr. Pl. Tinn.* 12. t. 6.—Afr. trop.
obtusifolia, *Hochst. in Flora*, xxvii. (1844) 296 (err. typ. 962).—Afr. austr.
octandra, *Blanco, Fl. Filipp.* ed. I. 349.—Ins. Philipp.
Pervillii, *Baill. Adansonia*, xi. (1873–76) 252.—Afr. trop.
pinnata, *Span. in Linnaea*, xv. (1841) 183.—Ins. Timor.
pinnata, Wall. Pl. As. Rar. ii. 21. t. 119 = Munronia Wallichii.
procera, *C. DC. in DC. Monog. Phan.* i. 444.—Afr. trop.
producta, *Baill. Adansonia*, xi. (1873–76) 254.—Madag.
propinqua, Hook. f. in Hook. Niger Fl. 254 = Vogelii.
pubescens, Benth. Fl. Austral. i. 379 = Brownii.
pubescens, *Hellen. in Vet. Acad. Nya Handl. Stockh.* ix. (1788) 308. t. 10. f. 1.—China.
pumila, *Benn. Pl. Jav. Rar.* i. 183.—Java.
quercifolia, G. Don, Gen. Syst. i. 678 = heterophylla.
Richardi, *Baill. Adansonia*, xi. (1873–76) 252.—Madag.
rigida, Vent. Choix, 48 = Quivisia chilosantha.
sericea, *Sm. Ic. Ined.* t. 12.—Madag.
stricta, Steud. Nom. ed. I. 859; ed. II. ii. 725 = Quivisia chilosantha.
tetramera, Benn. Pl. Jav. Rar. 184 = Quivisia tetramera.
tomentosa, Cav. Diss. vii. 361. t. 205 = sericea.
tricaraeopsis, *Baill. Adansonia*, xi. (1873–76) 252.—Afr. trop.
trichostylis, Miq. Fl. Ind. Bat. Suppl. 502 = Leptonychia glabra.
villosa, *Benn. Pl. Jav. Rar.* i. 182.—Java.
virens, Blanco, Fl. Filipp. ed. II. 243 = Dysoxylum Blancoi.
virens, Hellen. in Vet. Acad. Nya Handl. Stockh. ix. (1788) 308. t. 10. f. 1 = Atalantia monophylla.
virens, *Linn. Mant.* ii. 237 (err. typ. 732).—Ind. or.
Vogelii, *Hook. f. in Hook. Niger Fl.* 253.—Afr. trop.
Wakefieldii, *Oliver, in Hook. Ic. Pl.* t. 1489.—Afr. trop.
Zollingeri, *C. DC. in DC. Monog. Phan.* i. 441.—Java.

TURRAEANTHUS, Baill. Adansonia, xi. (1874) 261. *MELIACEAE.*
longipes, *Baill. l. c.* 262.—Afr. trop.
Mannii, *Baill. l. c.* 261.—Afr. trop.

TURRETIA, DC. Prod. ix. 236 (1845) = seq.

TURRETTIA, Poir. Encyc. vii. 719 (1806) = **Tourretia**, Fouger. (Bignon.).

TURRIGERA, Decne. in DC. Prod. viii. 590 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 751.
halophila, *Griseb. in Goett. Abh.* xxiv. (1879) 232.—Reg. Argent.
inconspicua, *Decne. in DC. Prod.* viii. 590.—Bras.

TURRITA, Wallr. Sched. Crit. 351, in nota (1822) = **Arabis**, Linn. (Crucif.).
major, Wallr. l. c. 352 = A. Turrita.
pendula, Wallr. l. c. = A. pendula.

TURRITIS, Tourn. ex Linn. Syst. ed. I (1735) = **Arabis**, Linn. (Crucif.).
accedens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. hirsuta.
alpestris, Schleich. Cat. Pl. Helv. ed. IV. 36 = A. ciliata.

TURRITIS :—
alpina, Braune, Salisb. Fl. ii. t. 2 = Arabis bellidifolia, pumila.
alpina, Linn. Syst. ed. XII. 443 = A. hirsuta.
arenosa, Lapeyr. Hist. Abr. Pl. Pyr. i. 387 = A. arenosa.
arvensis, R. Br. in Ait. Hort. Kew. ed. II. iv. 108 = Moricandia arvensis.
auriculata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = A. auriculata.
bellidifolia, All. Fl. Pedem. i. 270 = A. bellidifolia.
brachycarpa, Torr. & Gray, Fl. N. Am. i. 79 = A. Drummondii.
Brassica, Leers, Fl. Herborn. 147 = A. brassicaeformis.
Breutelii, Reichb. ex Lange, Consp. Fl. Groenl. i. 50 = A. Breutelii.
caerulea, All. Fl. Pedem. i. 270 = A. caerulea.
cenisia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. alpestris.
ciliata, Hoepfn. Mag. Naturk. Helv. iv. (1789) 29, 38 = A. ciliata.
ciliata, Tausch, in Flora, xix. (1836) 407 = A. hirsuta.
collisparsa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. hirsuta.
curtisiliqua, Fries, ex DC. Syst. ii. 223 = A. hirsuta.
diffusa, Hook. Fl. Bor. Am. i. 41 = Sisymbrium salsugineum.
Dregeana, Sond. in Linnaea, xxiii. (1850) 2 = A. perfoliata.
elongata, Raeusch. Nom. ed. III. 189 = A. Turrita.
enneaphyllos, Scop. ex Steud. Nom. ed. I. 268, 859 = Cardamine enneaphylla.
falcata, Turcz. in Bull. Soc. Nat. Mosc. (1840) 63.—Sibir.
Gerardi, Bess. Prim. Fl. Galic. ii. 87 = A. Gerardi.
Gerardi, Presl, ex Nym. Consp. 33 = A. hirsuta.
Gerardiana, Ram. ex DC. Syst. ii. 222 = A. hirsuta.
glabra, Linn. Sp. Pl. 666 = A. perfoliata.
Grahamii, Lehm. Del. Sem. Hort. Hamb. (1831) 7.—Cf. Linnaea, vi. (1831) Litt. 74 = A. Holboellii.
hirsuta, Brot. ex Nym. Consp. 33 = A. lusitanica.
hirsuta, Linn. Sp. Pl. 666 = A. hirsuta.
hirtella, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. hirsuta.
hispida, Hemsl. Biol. Centr. Am. Bot. i. 35, sphalm. = Sisymbrium hispidulum.
hispidula, DC. Syst. ii. 213 = Sisymbrium hispidulum.
humilis, Schleich. ex Koch, Syn. Fl. Germ. ed. I. 40 = A. muralis.
idanensis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. hirsuta.
laevigata, Muhl. ex Willd. Sp. Pl. iii. 543 = A. laevigata.
lasiophylla, Hook. & Arn. Bot. Beech. Voy. 321 = Sisymbrium reflexum.
Loeselii, R. Br. in Ait. Hort. Kew. ed. II. iv. 109 = Sisymbrium Loeselii.
lyrata, Rafin. in Am. Monthly Mag. ii. (1817) 44 = A. canadensis.
macrocarpa, Nutt. in Torr. & Gray, Fl. N. Am. i. 78 = A. perfoliata.
minor, Schleich. ex Willd. Enum. Hort. Berol. 685 = A. incana, muralis.
mollis, Hook. Fl. Bor. Am. i. 40 = A. Holboellii?
mollissima, Hoepfn. Mag. Naturk. Helv. iv. (1789) 29, 38 = A. saxatilis.
multiflora, Lapeyr. Hist. Abr. Pl. Pyr. 386 = A. hirsuta.
muralis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. muralis.
nemorensis, Wolf. in Hoffm. Deutschl. Fl. ed. II. ii. 58 = A. hirsuta.
oblongata, Rafin. in Am. Monthly Mag. ii. (1817) 44 = A. hirsuta.
ochroleuca, Lam. Fl. Fr. ii. 490 = A. Turrita.
ovata, Pursh, Fl. Am. Sept. ii. 438 = A. hirsuta.
patula, Ehrh. Beitr. vii. (1792) 159 = A. auriculata.
patula, R. Grah. in Edinb. N. Phil. Journ. (July–Oct. 1829) 350 = A. Holboellii.
pauciflora, Grimm, in Nov. Act. Nat. Cur. iii. (1767) App. 348 = A. brassicaeformis.
pendula, Desf. Tabl. ed. II. 152 = A. Turrita.
perfoliata, Neck. Delic. Gallo-Belg. 283 = A. perfoliata.
planisiliqua, Pers. Syn. ii. 205 = A. hirsuta.
praecox, Sm. in Rees, Cycl. xxxvi. n. 5 = A. hirsuta.
procera, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = A. Gerardi.
propera, Fourr. l. c. 333 = A. hirsuta.

TURRITIS :—
pubescens, Desf. Fl. Atlant. ii. 92 = Arabis pubescens.
pubigera, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 333 = A. Gerardi.
purpurea, Lam. Fl. Fr. ii. 491 = A. verna.
Raii, Vill. Prosp. 39; Hist. Pl. Dauph. iii. 326. t. 38 = A. stricta.
Raji, J. & C. Presl, Fl. Čech. 140 = A. hirsuta.
recta, Mert. & Koch, Deutschl. Fl. iv. 744 (Index) = A. auriculata.
retrofracta, Hook. Fl. Bor. Am. i. 41 = A. Holboellii.
rigida, Wall. Cat. n. 4783 = A. perfoliata.
rigidula, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = A. Gerardi.
rupestris, Hoppe, ex Roehl. Deutschl. Fl. ed. II. 358; et ex Braune, in Flora, ii. (1819) 566 = A. ciliata.
sagittata, Bertol. Pl. Genuens. 185 = A. Gerardi.
sagittata, Jahn, ex Schrank, Pl. Rar. Hort. Monac. t. 99 = A. brassicaeformis.
salsuginosa, DC. Syst. ii. 212 = Sisymbrium salsugineum.
sanguinea, Claus, in Beitr. Pfl. Russ. Reich. viii. 57.—Rossia.
saxatilis, Kit. in Linnaea, xxxii. (1863) 495 = A. saxatilis?
serpyllifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = A. serpyllifolia.
setosa, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 93 = Diplotaxis setosa.
spathulata, Nutt. in Torr. & Gray, Fl. N. Am. i. 78 = A. Drummondii?
stenopetala, Willd. Enum. Hort. Berol. Suppl. 46 = A. hirsuta.
stricta, All. Auct. Fl. Pedem. 18 = A. Allionii.
stricta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = A. stricta.
stricta, R. Grah. in Edinb. N. Phil. Journ. (July–Oct. 1829) 350 = A. Drummondii.
stricta, Host, Fl. Austr. ii. 268 = A. perfoliata.
verna, Lam. Fl. Fr. ii. 490 = A. alpina.

TURSENIA, Cass. in Dict. Sc. Nat. xxxvii. 480 (1825) = **Baccharis**, Linn. (Compos.).
humifusa, Cass. l. c. = B. humifusa.
sinuata, Cass. l. c. = B. sinuata.

TUSSACA, Rafin. Préc. Découv. 42 (1814); et in Journ. Phys. lxxxix. (1819) 261 = **Goodyera**, R. Br. (Orchid.).
reticulata, Rafin. Préc. Découv. 43 = G. pubescens.
secunda, Rafin. l. c. = G. repens.

TUSSACIA, Klotzsch, ex Beer, Bromel. 99 (1857) = **Catopsis**, Griseb. (Bromel.).
apicroides, Beer, l. c.—Mexic.
fulgens, Klotzsch, ex Beer, l. c. 100 = Tillandsia aurantiaca.
nitida, Beer, l. c. = C. nitida.
sessiliflora, Beer, l. c. 101.—Peruv.
vitellina, Klotzsch, ex Beer, l. c. 99.—Venezuela.

TUSSACIA, Reichb. Ic. Exot. i. 28. t. 41 (1827) (Tussaca); et in Moessl. Handb. i. p. xxxvii (1827). *GESNERACEAE*, Benth. & Hook. f. ii. 1006.
CHRYSOTHEMIS, Decne. in Rev. Hortic. Sér. III. iii. (1849) 242.
Friedrichsthaliana, *Hanst. in Linnaea*, xxxiv. (1865–66) 337.—Guatemala.
pulchella, *Reichb. Ic. Bot.* i. 28. t. 41; *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 113.—Panama.
rupestris, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 363.—Guian.
sanguinea, *Heynh. Nom.* ii. 747.—Am. austr.
semi-clausa, *Hanst. ex André, in Illustr. Hortic.* xvii. (1870) 140. t. 28.—Bras.
villosa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 363.—Guian.

TUSSACIA, Willd. ex Roem. & Schult. Syst. vii. p. x (1829) (Quid?).
elegans, Willd. l. c. 57.—Am. austr.

TUSSILAGO, [Tourn.] Linn. Syst. ed. I (1735); Linn. Gen. ed. I. 252 (1737), partim. *COMPOSITAE*, Benth. & Hook. f. ii. 438.
FARFARA, Gilib. Fl. Lituan. i. 177 (1781).
alba, Linn. Sp. Pl. 866 = Petasites albus.

TUSSILAGO:—
albicans, Sw. Prod. Veg. Ind. Occ. 113 = Chaptalia albicans.
alpestris, Hegetschw. Fl. Schw. 842 = Homogyne alpina?
alpina, Georgi, Reise, i. 230 = Gerbera Anandria.
alpina, Linn. Sp. Pl. 865 = Homogyne alpina.
alpina, Scop. Fl. Carn. ed. II. ii. 156 = Homogyne discolor.
Anandria, Linn. Sp. Pl. 865 = Gerbera Anandria.
androgyna, Gilib. Fl. Lituan. i. 176 = Petasites officinalis.
aquatica, Georgi, Reise, i. 230 = Petasites laevigatus.
Bellidiastrum, Linn. Hort. Upsal. 259. t. 3. f. 2 = Gerbera Anandria.
bicolor, Herb. Willd. ex DC. Prod. vii. 44 = Chaptalia runcinata.
bifolia, Lavy, Vég. Orig. 13; ex DC. Prod. vii. 208 (Quid?).—Hab.?
bohemica, Hoppe, Taschenb. (1803) 60; ex DC. Prod. v. 205 = Petasites laevigatus.
Cacalia, Scop. Fl. Carn. ed. II. ii. 156 = Adenostyles viridis.
caryophyllata, Vell. Fl. Flum. viii. t. 142 (Quid?).—Bras.
corymbosa, R. Br. in Parry, 1st Voy. App. 269 = Petasites frigidus.
dentata, Linn. Sp. Pl. ed. II. 1213 = Chaptalia dentata.
discolor, Jacq. Fl. Austr. iii. 27 = Homogyne discolor.
elongata, Salisb. Prod. 196 = Petasites officinalis.
exscapa, Pers. Syn. ii. 456 = Chaptalia exscapa.
Farfara, *Linn. Sp. Pl.* 865.—Europ.; As. bor. occ.; Ind. or.
fragrans, Vill. in Act. Soc. Hist. Nat. Par. i. (1792) 72 = Petasites fragrans.
frigida, Linn. Sp. Pl. 865 = Petasites frigidus.
frigida, Vill. Hist. Pl. Dauph. iii. 175; Suter, Fl. Helv. ii. 180 = Petasites niveus.
glabrescens, Steud. Nom. ed. I. 859 = Petasites albus.
Gmelini, Turcz. ex DC. Prod. vii. 271 = Gerbera Anandria.
hastifolia, Kit. in Linnaea, xxxii. (1883) 372 = Petasites niveus?
hermaphrodita, Gilib. Fl. Lituan. i. 175 = Petasites officinalis.
hybrida, Linn. Sp. Pl. 866 = Petasites officinalis.
integerrima, Vell. Fl. Flum. viii. t. 140 (Quid?).—Bras.
integrifolia, Michx. ex Willd. Sp. Pl. iii. 1964 = Chaptalia tomentosa.
intermedia, Hinterhuber, ex Trachsel, in Flora, xii. (1829) 32 = Petasites officinalis.
japonica, Linn. Mant. i. 113 = Senecio Kaempferi.
Kablikianus, Opiz, Seznam, 73, in syn. = Petasites officinalis.
laevigata, Willd. Sp. Pl. iii. 1969 = Petasites laevigatus.
lobata, Hort. ex Steud. Nom. ed. II. ii. 726 = Petasites palmatus.
lutea, J. F. Gmel. Syst. 1225 = Petasites albus.
lyrata, Pers. Syn. ii. 456 = Chaptalia nutans.
lyrata, Willd. Sp. Pl. iii. 1963 = Gerbera Anandria.
macrophylla, Wall. Cat. n. 2989 = Gerbera macrophylla.
micrantha, Andrz. Rys Botan. ii. (1830) 27 (Quid?).—Rossia.
nana, J. F. Gmel. Syst. 1224 = Gerbera Anandria.
nivea, Vill. in Act. Soc. Hist. Nat. Par. i. (1792) 73 = Petasites niveus.
nutans, Linn. Syst. ed. X. 1214 = Chaptalia nutans.
oblongifolia, Bertol. Misc. Bot. vi. 28.—Am. bor.
odorata, Vill. ex DC. Prod. v. 208 = Adenostyles viridis.
palmata, Ait. Hort. Kew. ed. I. iii. 188 = Petasites palmatus.
paradoxa, Retz. Obs. ii. 24 = Petasites niveus.
paradoxa, Roth, ex Steud. Nom. ed. I. 860 = Petasites spurius.
patens, Vell. Fl. Flum. viii. t. 144 (Quid?).—Bras.
Petasites, Linn. Sp. Pl. 866 = Petasites japonicus.
piloselloides, Pers. Syn. ii. 456 = Chaptalia piloselloides.
pumila, Sw. Prod. Veg. Ind. Occ. 113 = Chaptalia sinuata.
racemosa, Tausch, in Syll. Ratisb. ii. (1828) 251 = Petasites fragrans.
radiata, Gilib. Fl. Lituan. iii. 177 = T. Farfara.
radiata, J. F. Gmel. Syst. 1226 = Petasites laevigatus.

TUSSILAGO:—
ramosa, Hoppe, in Flora, x. (1827) 649 = Petasites albus.
reflexa, Vell. Fl. Flum. viii. t. 141 (Quid?).—Bras.
rheifolia, Warsz. in Otto & Dietr. Allg. Gartenz. xx. (1852) 293 (Quid?).—Bolivia.
riparia, Salisb. Prod. 196 = Petasites officinalis.
rubella, J. F. Gmel. Syst. 1225 (Quid?).—Sibir.
rubra, Link, in Buch, Beschr. Canar. Ins. 149, 176 = Senecio cruentus, *DC.*
ruderalis, Salisb. Prod. 195 = T. Farfara.
rupestris, Wall. Cat. n. 2990 = T. Farfara.
sagittata, Herb. Banks, ex Pursh, Fl. Am. Sept. ii. 531 = Petasites sagittatus.
sarmentosa, Pers. Syn. ii. 456 = Chevreulia stolonifera.
saxatilis, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 138 = Petasites frigidus?
sibirica, J. F. Gmel. Syst. 1225 = Gerbera Anandria.
sinuata, Pers. Syn. ii. 456 = Chaptalia sinuata.
spuria, Retz. Obs. i. 29. t. 2 = Petasites spurius.
spuria, Schrank, ex Steud. Nom. ed. I. 860 = Petasites niveus.
suaveolens, Desf. Tabl. ed. I. 97 = Petasites fragrans.
sylvestris, Jacq. Fl. Austr. v. App. t. 12 = Homogyne sylvestris.
tomentosa, Ehrh. Beitr. iii. 65 = Petasites spurius.
trifurcata, Forst. f. in Comm. Gotting. ix. (1789) 39 = Senecio trifurcatus.
vaccina, Vell. Fl. Flum. viii. t. 143 (Quid?).—Bras.
vulgaris, Lam. Fl. Fr. ii. 71 = T. Farfara.

TUTUCA, Molina, Sagg. Chile, ed. II. 135 (1810). Genus dubium *ERICACEARUM*.
chilensis, Molina, l. c.—Chili.
fistulosa, Molina, l. c. 288.—Chili.

TWEEDIA, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 291 = **Oxypetalum**, R. Br. (Asclep.).
birostrata, Hook. & Arn. l. c. = O. Hookeri.
Brunonis, Hook. & Arn. l. c. 292 = O. Brunonis.
coerulea, D. Don, ex Sweet, Brit. Flow. Gard. Ser. II. t. 407.—Reg. Argent.
floribunda, Hort. ex Lem. in Fl. des Serres, iv. (1848) 347 = O. solanoides.
macrolepis, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 291 = O. macrolepis.
pubescens, Hort. ex Lem. in Fl. des Serres, iv. (1848) t. 347 = O. solanoides.
rosea, Hort. ex Lem. l. c. = O. solanoides.
versicolor, Hook. Bot. Mag. t. 3630 = *coerulea*.

TYDAEA, Decne. Rev. Hortic. Ser. III. ii. (1848) 468 = **Isoloma**, Decne. (Gesner.).
amabilis, Planch. & Linden, in Linden, Cat. n. 10 (1855) 4; et in Fl. des Serres, x. (1855) 287. t. 1070.—N. Granat.
Ceciliae, André, Illustr. Hortic. xxiii. (1876) 178. t. 260.—N. Granat.
elegans, Planch. & Linden, in Linden, Cat. n. 10 (1855) 4 = I. elegans.
gigantea ×, Planch. Fl. des Serres, x. (1854–55) 21. t. 975–76.
Hansteini ×, Ortgies, in Regel, Gartenfl. vii. (1858) 65. t. 218.
Hillii, Regel, Gartenfl. iii. (1854) 73, 74.—Hab.?
Karsteniana, Hort. ex Hanst. in Linnaea, xxxiv. (1865–66) 439 = I. picta?
Lindeni, André, Illustr. Hortic. xx. (1873) 183. t. 147 = seq.
Lindeniana, Regel, Gartenfl. xvii. (1868) 257. t. 589.—Am. trop.
magnifica, Hanst. in Linnaea, xxxiv. (1865–66) 440 = Achimenes magnifica.
ocellata, Regel, Gartenfl. iii. (1854) 74 = I. ocellatum.
Ortgiesii ×, Hort. Van Houtte, ex Planch. in Fl. des Serres, xi. (1856) 213. t. 1181–1182.
pardina, Linden & André, Illustr. Hortic. xx. (1873) 212. t. 152.—N. Granat.
picta, Decne. in Rev. Hortic. Sér. III. ii. (1848) 468; et in Regel, Gartenfl. iii. (1854) 73, 75 = I. picta?
pyramidalis, Regel, Gartenfl. x. (1861) 193. t. 328.—Hab.?
venosa, Hort. ex Lescuyer, in L'Hortic. Franç. Sér. III. ii. (1861) 248. t. 20.—N. Granat.
Warszewiczii, Regel, Gartenfl. iii. (1854) 73, 74.—N. Granat.

TYDEA, C. Muell. in Walp. Ann. v. 427 (1858–60), sphalm. = praec.

TYLACANTHA, Endl. Gen. 672 (1839) = THYLACANTHA, Nees & Mart. = **Angelonia**, Humb. & Bonpl. (Scrophul.).

TYLANTHUS, Reiss. in Endl. Gen. 1101 (1840) = **Phylica**, Linn. (Rhamn.).
abietinus, Presl, Bot. Bemerk. 38 = P. abietina.
atratus, Presl, l. c. 37 = P. atrata.
brevifolius, Presl, l. c. 38 = P. brevifolia.
callosus, Presl, l. c. 37, 38 = P. callosa.
comosus, Presl, l. c. 37 = P. comosa.
debilis, Presl, l. c. 38 = P. debilis.
diosmoides, Presl, l. c. = P. diosmoides.
distichus, Presl, l. c. = P. disticha.
emirnensis, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 128.—Madag.
empetroides, Presl, Bot. Bemerk. 38 = P. disticha, rubra.
eriophorus, Presl, l. c. = P. eriophora.
excelsus, Presl, l. c. 37, 38 = P. excelsa.
gracilis, Presl, l. c. 37 = P. gracilis.
imberbis, Presl, l. c. = P. imberbis.
lanceolatus, Presl, l. c. 38 = P. eriophora.
litoralis, Presl, l. c. 37 = P. atrata.
parviflorus, Presl, l. c. 38 = P. parviflora.
retrorsus, Presl, l. c. 37 = P. retrorsa.
rosmarinifolius, Presl, l. c. 38 = P. rosmarinifolia.
secundus, Presl, l. c. = P. eriophora.
Thunbergianus, Presl, l. c. = P. Thunbergiana.
virgatus, Presl, l. c. 37 = P. virgata.
Willdenowianus, Presl, l. c. 38 = P. Willdenowiana.

TYLLOMA, D. Don, in Trans. Linn. Soc. xvi. (1830) 238 = **Chaetanthera**, Ruiz & Pav. (Compos.).
glabratum, DC. Prod. vii. 32.—Chili.
lanatum, Phil. in Linnaea, xxxiii. (1864–65) 112.—Chili.
limbatum, D. Don, in Trans. Linn. Soc. xvi. (1830) 236, 301 = C. limbata.
pusillum, D. Don, in Phil. Mag. (1832) 391 = C. pusilla.
renifolium, Wedd. Chlor. And. i. 28. t. 8 A.—Chili.
splendens, Wedd. l. c. 27.—Chili.

TYLOCHILUS, Nees, in Verh. Gartenb. Ges. Berl. viii. (1832) 194. t. 3 = **Cyrtopodium**, R. Br. (Orchid.).
flavus, Nees, l. c. 195 = C. flavum.

TYLODERMA, Miers, in Trans. Linn. Soc. xxviii. (1872) 413 = **Hippocratea**, Linn. (Celastr.).
praecelsa, Miers, l. c. 414.—Panama.

TYLODONTIA, Griseb. Cat. Pl. Cub. 175 (1866) = **Astephanus**, R. Br. (Asclep.).
cubensis, Griseb. l. c.—Cuba.

TYLOGLOSSA, Hochst. in Flora, xxv. (1842) i. Beibl. 144 = **Justicia**, Houst. (Acanth.).
acuminata, Hochst. l. c. xxvi. (1843) 73 = Adhatoda acuminata.
adenostachya, Nees, in Mart. Fl. Bras. ix. 117.—Bras.
chrysotrichoma, Nees, l. c. 116.—Bras.
ciliata, Nees, l. c.—Bras.
flexuosa, Nees, l. c.—Bras.
genistiformis, Nees, l. c. 115.—Bras.
glandulosa, Nees, l. c. 117.—Bras.
Kotschyi, Hochst. in Flora, xxvi. (1843) 74.—Afr. trop.
major, Hochst. l. c. 73 = J. major.
minor, Hochst. l. c. = J. minor.
palustris, Hochst. l. c. xxv. (1842) i. Beibl. 144; xxvi. (1843) 72 = J. palustris.
pubescens, Hochst. l. c. xxviii. (1845) 71.—Afr. austr.
rostrata, Hochst. ex Nees, in DC. Prod. xi. 397 = J. Schimperiana.
Schimperi, Hochst. in Flora, xxvi. (1843) 73 = J. Schimperiana.
sexangularis, Hochst. in Flora, xxv. (1842) i. Beibl. 144 = Dicliptera sexangularis.

TYLOMIUM, Presl, Prod. Monog. Lobel. 31 (1836) = **Lobelia**, Plum. (Campan.).
assurgens, Presl, l. c. 32 = L. assurgens.
conglobatum, Presl, l. c. = L. conglobata.
flavescens, Presl, l. c. = L. conglobata.
robustum, Presl, l. c. = L. robusta.

TYLOPHORA, R. Br. in Mem. Wern. Soc. i. (1809) 28. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 770.
AMBLYOGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 310.

TYLOPHORA:—
Belostemma, Wall. in Wight, Contrib. 52 (1834).
Homolostyles, Wall. ex Wight, l. c. 50 (1834).
Hybanthera, Endl. Prod. Fl. Norf. 59 (1833).
Iphisia, Wight, Contrib. 52 (1834).
albida, *Zipp. ex Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
aristolochioides, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 129.—Japon.
asthmatica, *Wight & Arn. in Wright, Contrib.* 51.—Ind. or.; Malaya.
barbata, *R. Br. Prod.* 460.—Austral.
Belostemma, *Benth. in Benth. & Hook. f. Gen.* ii. 771.—Reg. Himal.
bifida, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 134.—Ins. Philipp.
biglandulosa, *F. Muell. Fragm.* ix. 71, 169.—Ins. Norfolk.
Bojeriana, *Decne. in DC. Prod.* viii. 607.—Madag.
Brackenridgei, *A. Gray, in Proc. Am. Acad.* v. (1862) 334.—Ins. Fiji.
brevipes, *F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 134.—Ins. Philipp.
caffra, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 542 (err. typ. 442).—Afr. austr.
calcarata, *Benth. Fl. Austral.* iv. 335.—Austral.
capparidifolia, *Wight & Arn. in Wright, Contrib.* 51.—Ind. or.
carnosa, Dalz. & Gibs. Bomb. Fl. 150 = Dalzellii.
carnosa, Wall. ex Wight, Contrib. 49 = tenuis.
chlorantha, *Miq. Fl. Ind. Bat.* ii. 486.—Java.
cirrosa, *Aschers. ex Schweinf. Beitr. Fl. Aethiop.* 132.—Afr. trop.
cissoides, *Blume, Bijdr.* 1061.—Java.
congesta, *Decne. in DC. Prod.* viii. 609.—Java.
cordifolia, *Thw. Enum. Pl. Zeyl.* 196.—Zeylan.
crassifolia, *Zipp. ex Decne. in DC. Prod.* viii. 612.—Ins. Timor.
cuspidata, *Zipp. ex Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 274. t. 10.—Ins. Timor.
Dalzellii, *Hook. f. Fl. Brit. Ind.* iv. 43.—Ind. or.
enervis, *F. Muell. Fragm.* ix. 70.—Ins. Dom. Howe.
erecta, *F. Muell. ex Benth. Fl. Austral.* iv. 334.—Austral.
exilis, *Colebr. in Trans. Linn. Soc.* xii. (1818) 358.—Ind. or.; Malaya.
fasciculata, *Buch.-Ham. ex Wight, Contrib.* 50.—Ind. or.
fasciculata, Thw. Enum. Pl. Zeyl. 197 = T. Iphisia.
flava, *Trimen, in Journ. Bot.* xxiii. (1885) 239.—Zeylan.
flexuosa, *R. Br. Prod.* 460.—Austral.
floribunda, *Benth. Fl. Austral.* iv. 335.—Austral.
floribunda, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 128.—Japon.
globifera, *Hook. f. Fl. Brit. Ind.* iv. 41.—Ins. Andaman; Malacca.
Govanii, *Decne. in DC. Prod.* viii. 610.—Reg. Himal.
grandiflora, *R. Br. Prod.* 460.—Austral.
Helferi, *Hook. f. Fl. Brit. Ind.* iv. 40.—Burma.
heterophylla, A. Rich. Tent. Fl. Abyss. ii. 41 = Vincetoxicum heterophyllum.
himalaica, *Hook. f. Fl. Brit. Ind.* iv. 42.—Reg. Himal.
hirsuta, *Wight, Contrib.* 49.—Reg. Himal.
hispida, *Decne. in DC. Prod.* viii. 610.—China.
incana, *Brunn. in Flora*, xxiii. (1840) II. *Beibl.* 26.—Afr. trop.
Iphisia, *Decne. in DC. Prod.* viii. 610.—Ind. or.
Jacquemontii, Decne. l. c. 607 = hirsuta.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 129.—Japon.
laevigata, *Decne. in DC. Prod.* viii. 612.—Ins. Maurit.
laevis, *Decne. l. c.* 611.—Hab.?
lanceolata, *Zipp. ex Span. in Linnaea*, xv. (1841) 324, nomen.—Ins. Timor.
longifolia, *Wight, Contrib.* 50.—Ind. or.
lutescens, *Decne. in Herb. Gén. Amat. et Hort. Univ.* 5 (1844).—Hab.?
lycioides, *Decne. in DC. Prod.* viii. 608.—Afr. austr.
macrantha, Hance, in Journ. Bot. xx. (1882) 79 = Dregea volubilis.
macrantha, *Hook. f. Fl. Brit. Ind.* iv. 40.—Ind. or.
macrobotrya, *Miq. Fl. Ind. Bat.* ii. 488.—Java?
macrophylla, *Benth. Fl. Austral.* iv. 334.—Austral.
membranifolia, Thw. Enum. Pl. Zeyl. 424 = zeylanica.
micrantha, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 377.—Ins. Timor.

TYLOPHORA:—
micrantha, Thw. Enum. Pl. Zeyl. 197 = zeylanica.
microstachys, *Hook. in Hook. Ic. Pl.* t. 1450.—Reg. Himal.
mollissima, *Wall. Cat.* n. 8202; *Wight, Contrib.* 49 = hirsuta.
ovata, *Hook. ex Steud. Nom.* ed. II. ii. 726.—China.
paniculata, *R. Br. Prod.* 460.—Austral.
parviflora, Wight, Ic. iv. n. 1274, textu = pauciflora.
pauciflora, Herb. Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iv. 44 = Dalzellii.
pauciflora, Hohen. ex Hook. f. l. c. 40 = T. Iphisia.
pauciflora, *Wight & Arn. in Wight, Contrib.* 49.—Ind. or.
Perrottetiana, *Decne. in DC. Prod.* viii. 609.—Ins. Philipp.
pubescens, Wall. ex Hook. f. Fl. Brit. Ind. iv. 45 = asthmatica.
punctata, Kostel. Allg. Med. Pharm. Fl. iii. 1084 = Cosmostigma racemosa.
purpurea, *Wall. ex Wight, Contrib.* 51.—Burma.
rotundifolia, *Buch.-Ham. ex Wight, l. c.* 50.—Ind. or.
rupestris, *Blume, Bijdr.* 1062.—Java.
samoënsis, *A. Gray, in Proc. Am. Acad.* v. (1862) 334.—Ins. Samoa.
sublanceolata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 128.—Japon.
sylvatica, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 273.—Afr. trop.
syringaefolia, *E. Mey. Comm. Pl. Afr. Austr.* 198; *Drège, ex Brunn. in Flora*, xxiii. (1840) II. *Beibl.* 27.—Afr. austr.
Tanakae, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* i. 321.—Japon.
tenerrima, *Wight, Contrib.* 50.—Reg. Himal.
tenuis, *Blume, Bijdr.* 1062.—Java.
tenuissima, Wight, Contrib. 49 = tenuis.
tomentosa, G. Don, Gen. Syst. iv. 130 = asthmatica?
velutina, G. Don, l. c. 128 = Marsdenia velutina.
villosa, *Blume, Bijdr.* 1061; *Miq. Fl. Ind. Bot.* ii. 486.—Java.
vomitoria, Voigt, Hort. Suburb. Calc. 539 = asthmatica.
Wallichii, *Hook. f. Fl. Brit. Ind.* iv. 45.—Singapore.
Woollsii, *Benth. Fl. Austral.* iv. 335.—Austral.
zeylanica, *Decne. in DC. Prod.* viii. 608.—Ind. or.

TYLOSEPALUM, Kurz, ex Teysm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 50 = **Trigonostemon**, Blume (Euphorb.).
aurantiacum, Kurz, ex Teysm. & Binn. l. c.—Ins. Banca.

TYLOSTYLIS, Blume, Fl. Jav. Praef. 6 (1828) = **Eria**, Lindl. (Orchid.).

TYLOTHRASYA, Doell, in Mart. Fl. Bras. ii. II. 295. t. 37 (1877) = **Panicum**, Linn. (Gramin.).
petrosa, Doell, l. c. 296 = P. petrosum.

TYMPANANTHE, Hassk. in Flora, xxx. (1847) 757 = **Dictyanthus**, Decne. (Asclep.).
suberosa, Hassk. l. c. 758 = D. Pavonii.

TYNANTHUS, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 193. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1031.
Schizopsis, Bur. ex Baill. Adansonia, v. (1865) 371. tt. 10, 11.
cognatus, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 193.—Bras.
confertiflorus, Miers, l. c., nomen (Quid?).—Bras.
elegans, *Miers, l. c.* 193.—Bras.
fasciculatus, *Miers, l. c.*—Bras.
gibbus, Miers, l. c., nomen (Quid?).—Bras.
Gondotiana, *Bur. in Baill. Adansonia*, viii. (1867–68) 274.—Bras.
labiatus, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 193.—Bras.
laxiflorus, *Miers, l. c.*, nomen.—Peruv.
petiolatus, *Miers, l. c.*, nomen.—Bras.
strictus, Miers, l. c., nomen (= *Pyrostegia lauta*).—Bras.

TYNUS, J. S. Presl, Rostl. ii. 93 (1823) = **Viburnum**, Linn. (Caprifol.).
ellipticus, J. S. Presl, l. c. = V. tinoides.
lauriformis, J. S. Presl, l. c. = V. Tinus.
lucidus, J. S. Presl, l. c. = V. Tinus.
nudus, J. S. Presl, l. c. = V. nudum.
rugosus, J. S. Presl, l. c. = V. rigidum.

TYPHA, Linn. Syst. ed. I (1735); Gen. ed. I. 281 (1737). *TYPHACEAE*, Benth. & Hook. f. iii. 955.
Massula, Dulac, Fl. Hautes-Pyr. 47 (1867).
abyssinica, Reichb. f. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 89 = angustata.
aequalis, Schnizl. Typh. 25 = angustata.
ambigua, Schur, ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 76 = latifolia.
americana, Rich. ex Rohrb. l. c. 97 = domingensis.
angustata, *Bory & Chaub. Nouv. Fl. Pélop.* 4.—Reg. Mediterr. or.; Oriens.
angustifolia, All. Fl. Pedem. ii. 273 = Laxmanni.
angustifolia, Aubl. Pl. Gui. ii. 847 = domingensis.
angustifolia, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 338 = javanica.
angustifolia, Eckl. & Zeyh. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 100 = capensis.
angustifolia, Hook. f. Fl. Tasm. ii. 38 = Muelleri.
angustifolia, *Linn. Sp. Pl.* 971.—Europ.; Am. bor.
angustifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 226 = angustata.
angustissima, Griff. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 92 = Laxmanni.
australis, Schum. & Thonn. Beskr. Guin. Pl. 401 = angustifolia.
baetulona.—Cf. Bull. Soc. Bot. Fr. xi. (1864) Bibl. 230, sphalm. = *bethulona*.
Balansae, Reut. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 90 = stenophylla.
bethulona, Costa, Intr. Fl. Catal. (1864) 251 = latifolia.
Brownii, Kunth, Enum. Pl. iii. 92 = angustifolia?
Bungeana, Presl, Epim. Bot. 239 = Laxmanni.
capensis, *Rohrb. in Verh. Bot. Ver. Brandenb.* xi. (1869) 96.—Afr. austr.
caucasica, Lehm. ex Rohrb. l. c. 92 = Laxmanni.
communis, Noronha, ex Thou. Prod. Phyt. 2; in Mélang. (1811), nomen (Quid?).—Madag.
crassa, Rafin. Atl. Journ. 148 (Quid?).—Am. bor.
daenatica, Ehrenb. ex Steud. Nom. ed. II. ii. 727 = seq.
damiattica, Ehrenb. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 88 = angustata.
domingensis, *Pers. Syn.* ii. 532.—As. trop.
Ehrenbergii, Schur, ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 88 = angustata.
elata, Bor. Fl. Centr. Fr. ed. III. ii. 733 = latifolia.
elatior, Boenn. Prod. Fl. Monast. 274 = angustifolia.
elatior, Bor. in Guill. Arch. Bot. ii. (1833) 399 = latifolia.
elatior, *Rafin. in Med. Repos. N. York*, v. (1808) 354, nomen; *Atl. Journ.* 148.—Am. bor.
elephantina, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 566.—Reg. Mediterr.; Ind. or.
elephantina, Schimp. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 95 = Schimperi.
elephantina, Thw. Enum. Pl. Zeyl. = javanica.
elliptica, C. C. Gmel. Fl. Bad. iii. 603 = Laxmanni.
Engelmanni, A. Br. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 76 = latifolia.
essequeboënsis, G. F. W. Mey. ex Rohrb. l. c. 97 = domingensis.
gigantea, Schur, ex Kunth, Enum. Pl. iii. 92 = domingensis.
glauca, *Godr. Fl. Lorr.* ed. I. iii. 20; ed. II. ii. 332.—Gallia.
gracilis, Jord. Obs. Pl. Crit. vii. (1849) 43; et Grain. Rec. Jard. Grenoble (1849), 24 = Laxmanni.
gracilis, Rafin. New Fl. Am. ii. 91 (Quid?).—Am. bor.
gracilis, Reichb. Ic. Fl. Germ. ix. 2. t. 520. f. 744 = angustifolia.
gracilis, Schur, Enum. Pl. Trans. 637 = angustata.
Haussknechtii, *Rohrb. in Verh. Bot. Ver. Brandenb.* xi. (1869) 99.—Armenia.
intermedia, Schleich. Cat. Pl. Helv. 59 = Laxmanni?
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 206. t. 1 = latifolia.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 160.—Japon.
javanica, *Schnitzl. ex Zoll. Cat.* n. 77.—As. trop.; Ins. Mascar.
juncea, Stev. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 92 = Laxmanni.
juncifolia, Čelak. in Lotos, xvi. (1866) 149 = stenophylla.
juncifolia, Montand. in Friche-Joset, Syn. Fl. Jura, 325 = Laxmanni.
latifolia, Forst. f. Prod. 64 = angustifolia.

TYPHA:—
latifolia, Krauss, in Flora, xxviii. (1845) 343 = capensis.
latifolia, *Linn. Sp. Pl.* 971.—Europ.; Am. bor.
latifolia, G. F. W. Mey. Prim. Fl. Esseq. 262 = domingensis.
Laxmanni, *Lepech. in Nov. Act. Acad. Petrop.* xii. (1801) 84.—Europ.; As. bor.
macranthelia, Webb & Berth. Phyt. Canar. iii. 291 = angustifolia.
major, Curt. Fl. Lond. fasc. iii. t. 61 = latifolia.
Martini, Jord. Cat. Gr. Jard. Grenoble (1850) = Laxmanni.
maxima, Schur, ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 97 = domingensis.
media, Barbieri, ex Rohrb. l. c. 90 = stenophylla.
media, Bory & Chaub. Exp. Sc. Morée, 11 = angustata.
media, C. C. Gmel. Fl. Bad. iii. 602 = angustifolia.
media, Pollini, Fl. Veron. iii. 100 = latifolia?
minima, Hofm. Deutschl. Fl. ed. II. ii. 251; Funk, ex Willd. Sp. Pl. iv. 198 = Laxmanni.
minor, Curt. Fl. Lond. fasc. iii. t. 62 = angustifolia.
minor, Sm. Fl. Brit. iii. 960, quoad syn. = Laxmanni.
minuta, Schrenk, ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 90 = stenophylla.
Muelleri, *Rohrb. l. c.* 95.—Austral.; N. Zel.
nana, Avé-Lall. Pl. Ital. Bor. 19 = Laxmanni.
orientalis, Presl, Epim. Bot. 239 = Shuttleworthii.
pendula, Fisch. ex Sond. Fl. Hamb. 508 = latifolia.
Poitiaei, Poit. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 92 = Laxmanni.
remotiuscula, Schur, Enum. Pl. Transs. 637 = latifolia.
Schimperi, *Rohrb. in Verh. Bot. Ver. Brandenb.* xi. (1869) 95.—Abyss.
Shuttleworthii, *Koch & Sond. in Koch, Syn. Pl. Germ.* ed. II. 786.—Europ.
Shuttleworthii, Lehm. Pl. Preiss. ii. 1 = Muelleri.
spiralis, Rafin. Atl. Journ. 148 (Quid?).—Am. bor.
stenophylla, *Fisch. & Mey. in Bull. Phys.-Math. Pétersb.* iii. (1845) 209.—Ital.; Rossia; Reg. Casp.
stenophylla, Haussk. ex Rohrb. in Verh. Bot. Ver. Brandenb. xi. (1869) 99 = Haussknechtii.
tenuifolia, H. B. & K. Nov. Gen. et Sp. i. 82 = domingensis.
transsilvanica, Schur, Enum. Pl. Transs. 637 = Shuttleworthii.
truxillensis, H. B. & K. Nov. Gen. et Sp. i. 82 = domingensis.

TYPHALEA, Neck. Elem. ii. 412 (1790) = **Pavonia**, Cav. (Malvac.).
spinifex, Presl, Bot. Bemerk. 19 = P. spinifex.

TYPHOIDES, Moench, Meth. 201 (1794) = **Phalaris**, Linn. (Gramin.).
arundinacea, Moench, l. c. 202 = P. arundinacea.

TYPHONIUM, Schott, in Wien. Zeitschr. iii. (1829) 72; ex Endl. in Linnaea, vi. (1831) Litt. 53. *AROIDEAE*, Benth. & Hook. f. iii. 967.
HETEROSTALIS, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 261.
alismifolium, *F. Muell. Fragm.* viii. 186.—Austral.
angustilobum, *F. Muell. l. c.* x. 66.—Austral.
Brownii, *Schott, Aroid.* 11. t. 15.—Austral.
bulbiferum, *Dalz. in Hook. Kew Journ.* iv. (1852) 113.—Ind. or.
cochinchinense, Blume, Rumphia, i. 135 = Theriophonum Wightii.
crassifolium, Ledeb. Fl. Ross. iv. 10 = Helicophyllum crassifolium.
crenatum, Schott, Melet. i. 17 = Theriophonum crenatum.
cuspidatum, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 367.—As. trop.
discolor, *Hassk. Cat. Hort. Bog. Alt.* 54.—Java.
divaricatum, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 367; *et ex Blume, Rumphia*, i. 130. t. 36.—As. trop.
diversifolium, *Wall. Cat.* n. 8933 A.—Ind. or.
fallax, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 260.—Java.
flagelliferum, Griff. Notul. iii. 144 = cuspidatum.
flagelliforme, Blume, Rumphia, i. 134 = cuspidatum.
foliolosum, Engl. in DC. Monog. Phan. ii. 618 = diversifolium.
giganteum, *Engl. Bot. Jahrb.* iv. (1883) 66.—China.
gracile, *Schott, Aroid.* i. 12.—Ind. or.
hastiferum, Miq. Fl. Ind. Bat. iii. 194; et in Bot. Zeit. xiv. (1856) 563 = cuspidatum.
Huegelianum, Schott, Aroid. i. 13. t. 19 = diversifolium.

TYPHONIUM:—
javanicum, Miq. Fl. Ind. Bat. iii. 193; et in Bot. Zeit. xiv. (1856) 563 = divaricatum.
liliifolium, *F. Muell. ex Schott, Prod. Aroid.* 107.—Austral.
minutum, Schott, in Wien. Zeitschr. iii. (1829) 72; ex Endl. in Linnaea, vi. (1831) Litt. 54 = Theriophonum Wightii?
Mottleyana, *Schott, Prod. Aroid.* 106.—Ind. or.; Malaya.
orixense, Schott, in Wien. Zeitschr. iii. (1829) 72; ex Endl. in Linnaea, vi. (1831) Litt. 54 = trilobatum.
pedatum, Engl. in DC. Monog. Phan. ii. 613 = fallax.
pedatum, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 262.—Burma.
Reinwardtiana, De Vriese & Miq. ex Miq. Fl. Ind. Bat. iii. 195 = cuspidatum.
Roxburghii, Baker, in Saund. Refug. Bot. t. 283 = trilobatum.
Roxburghii, *Schott, Aroid.* i. 12. t. 17.—Ind. or.
siamense, Engl. in DC. Monog. Phan. ii. 615 = trilobatum.
sylvaticum, Voigt, Hort. Suburb. Calc. 686 = Synantherias silvaticus.
trilobatum, *Schott, in Wien. Zeitschr.* iii. (1829) 72; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 54.—Ind. or.; Malaya.
triste, Griff. Notul. iii. 145 = trilobatum.
tuberculigerum, *Schott, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 123.—Japon.

TYPHONODORUM, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 70. *AROIDEAE*, Benth. & Hook. f. iii. 977.
Lindleyanum, *Schott, l. c.*—Madag.
madagascariense, *Engl. Bot. Jahrb.* i. (1881) 188.—Madag.

TYPHYLA, Heist. Syst. 12 (1748) 12 (Quid?).

TYRIA, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1850) = **Bernardia**, Houst. (Euphorb.).
myricaefolia, Scheele, in Linnaea, xxv. (1852) 581 = B. myricaefolia.

TYRIA, Klotzsch, in Linnaea, xxiv. (1851) 21 = **Macleania**, Hook. (Vaccin.).
Salapa, Klotzsch, l. c. = M. Salapa.

TYRIMNUS, Cass. in Bull. Soc. Philom. (1818) 168; et in Dict. Sc. Nat. xli. 314, 335 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 470.
leucographus, *Cass. ll. cc.* 168; 335.—Europ. austr.

TYSSACIA, Steud. Nom. ed. II. ii. 727 (1841), sphalm. = **Tussacia**, Reichb. (Gesner.).

TYTONIA, G. Don, Gen. Syst. i. 749 (1831) = **Hydrocera**, Blume (Geran.).
madagascariensis, G. Don, l. c. 750 = Impatiens madagascariensis.
natans, G. Don, l. c. 749 = H. angustifolia.

U

UAPACA, Baill. Etud. Gén. Euph. 595 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 282.
GYMNOCARPUS, Thou. ex Baill. Etud. Gén. Euph. 595 (1858).
benguelensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 332.—Afr. trop.
Bojeri, *Baill. Adansonia*, xi. (1874) 176.—Madag.
clusiacea, *Baker, in Journ. Linn. Soc.* xviii. (1881) 278.—Madag.
clusioides, *Baker, l. c.* xxi. (1885) 441.—Madag.
densifolia, *Baker, l. c.* xx. (1883) 252.—Madag.
ferruginea, *Baill. Etud. Gén. Euph.* 596.—Madag.
guineensis, *Muell. Arg. in Flora*, xlvii. (1864) 517.—Afr. trop.
Heudelotii, *Baill. Adansonia*, i. (1860–61) 81.—Afr. trop.
Kirkiana, *Muell. Arg. in Flora*, xlvii. (1864) 517.—Afr. trop.
myricaefolia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 440.—Madag.
nitida, *Muell. Arg. in Flora*, xlvii. (1864) 517.—Afr. trop.
Thouarsii, *Baill. Etud. Gén. Euph.* 598.—Madag.

UBIAEA, J. Gay, in A. Rich. Tent. Fl. Abyss. i. 447 (1847) = **Landtia**, Less. (Compos.).
Schimperi, J. Gay, l. c. = L. Schimperi.

UBIUM, J. F. Gmel. Syst. 839 (1791) = **Dioscorea**, Plum.
quadrifarium, J. F. Gmel. l. c. (Quid?).—Hab.?

UCACEA, Cass. in Dict. Sc. Nat. xxvii. 9 (1823) = **Blainvillea**, Cass. (Compos.).

UCACOU, Adans. Fam. ii. 131 (1763) = **Synedrella**, Gaertn. (Compos.).

UCRIA, Targ. ex Pfeiff. Nom. ii. 1519 (1874) (nomen auctoris lapsu pro genere habitum).—Cf. Benth. & Hook. f. Gen. iii. 964.

UCRIANA, Willd. Sp. Pl. i. 961 (1798) = **Tocoyena** Aubl. (Rubiac.).
Hamelliae, D. Dietr. Syn. Pl. i. 798 (Adenosacmes sp.).—Nepal.
hirsuta, D. Dietr. l. c. = T. hirsuta.
Humboldtii, Spreng. Syst. i. 761 = T. longifolia.
insignis, Willd. ex Spreng. Syst. i. 761 = Posoqueria latifolia.
longiflora, D. Dietr. Syn. Pl. i. 798 = T. longiflora.
longifolia, D. Dietr. l. c. = T. longifolia.
longifolia, Spreng. Syst. i. 761 = Augusta lanceolata.
macrophylla, D. Dietr. Syn. Pl. i. 798 = Posoqueria latifolia.
Mussaendae, D. Dietr. l. c. (= *Menestoria Mussaendae*).—Nepal.
Mutisii, Spreng. Syst. i. 761 = T. Mutisii.
racemosa, Schum. & Thonn. Beskr. Guin. Pl. 107 = Oxyanthus tubiflorus.
rigida, D. Dietr. Syn. Pl. i. 798 (= *Menestoria rigida*).—Nepal.
speciosa, Willd. Sp. Pl. i. 961 = T. longiflora.
Wallichii, D. Dietr. Syn. Pl. i. 798 = Mussaenda Roxburghii?

UDORA, Nutt. Gen. Am. ii. 242 (1818) = **Elodea**, Michx. (Hydroch.).
australis, F. Muell. Second Gen. Rep. 16 = Hydrilla verticillata.
brasiliensis, Endl. & Mart. ex Seub. in Mart. Fl. Bras. iii. 1. 99.—Am. austr.
canadensis, Nutt. Gen. Am. ii. 242 = E. canadensis.
cordofana, Hochst. in Flora, xxv. (1842) Beibl. i. 133.—Afr. trop.
granatensis, Spreng. Syst. iv. Cur. Post. 25 = E. granatensis.
guianensis, Steud. Nom. ed. II. ii. 727 = E. guianensis.
guyanensis, Chat. Anat. Comp. (1856) 26. t. 10. f. 4–6 = praec.?
lithuanica, Reichb. Fl. Germ. Excurs. 139 = Hydrilla verticillata.
occidentalis, Koch, Syn. Fl. Germ. ed. I. 669 = Hydrilla verticillata.
orinocensis, Spreng. Syst. iv. Cur. Post. 25 = E. orinocensis.
pomeranica, Reichb. Ic. Fl. Germ. vii. 31 = Hydrilla verticillata.
verticillata, Nutt. Gen. Am. ii. 242 = E. canadensis.
verticillata, Spreng. Syst. i. 170 = Hydrilla verticillata.

UEBELINIA, Hochst. in Flora, xxiv. (1841) 1. Intell. 30; II. 664. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 148.
abyssinica, *Hochst. ll. cc.* 30; 665.—Abyss.
rotundifolia, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 397.—Afr. trop.
spathulaefolia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 46 = abyssinica.

UGNI, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 579 = **Myrtus**, Tourn.
Berteroi, Phil. f. Cat. Pl. Vasc. Chil. 79 = M. Selkirkii.
Bridgesii, Berg, in Linnaea, xxvii. (1854) 389, in obs.—Hab.?
Candollei, Berg, l. c. 388 = M. Candollii.
Friedrichsthalii, Berg, l. c. = Eugenia Friedrichsthalii.
Krausei, Phil. ex Berg, in Bot. Zeit. xvi. (1858) 250 = M. Krausei.

UGNI :—
Molinae, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 579 = Myrtus Molinae.
montana, Berg, in Linnaea, xxvii. (1854) 392 = M. montana.
myricoides, Berg, l. c. 391 = M. myricoides.
Oerstedii, Berg, l. c. 389 = M. Oerstedii, Eugenia irazuensis.
Philippii, Berg, l. c. 387 (*Philipii*) = M. Molinae.
Poeppigii, Berg, l. c. 386.—Chili.
Selkirkii, Berg, l. c. 392 = M. Selkirkii.
Warscewiczii, Berg, l. c. 390 = Eugenia Warszewiczii.

UGONA, Adans. Fam. ii. (22) (1763) = **Hugonia**, Linn. (Linac.).

UHDEA, Kunth, Ind. Sem. Hort. Berol. (1847) 13.—Cf. Ann. Sc. Nat. Sér. III. ix. (1848) 316 = **Moñtanoa**, La Llave (Compos.).
bipinnatifida, Kunth, l. c. = M. bipinnatifida.

UITENIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = **Erioglossum**, Blume (Sapind.).
stilaginea, Noronha, ex Miq. Fl. Ind. Bat. i. II. 574 = E. edule.

ULANTHA, Hook. Bot. Mag. sub t. 2990, et Index (1830) = **Chloraea**, Lindl. (Orchid.).
grandiflora, Hook. l. c.—Ins. S. Vincent. (cult.).

ULEX, Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 483.
LEONHARDIA, Opiz, in Lotos, vii. (1857) 88.
NEPA, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 286.
STAURACANTHUS, Link, in Schrad. Neues Journ. ii. II. (1808) 52.
africanus, *Webb, in Ann. Sc. Nat.* Sér. III. xvii. (1852) 289.—Afr. bor. occ.
aphyllus, Link, ex Nym. Consp. 149 = genistoides.
argenteus, *Welw. ex Webb, Otia Hisp.* 44.—Lusitan.
armoricanus, Mabille, in Act. Linn. Soc. Bord. xxv. (1864) 524 = europaeus.
australis, Funk, ex Webb, Otia Hisp. 42 = Willkommii.
australis, Roxas Clem. Ensay. Vid, 291 = parviflorus.
australis, Welw. ex Webb, Otia Hisp. 41 = Welwitschianus.
autumnalis, Thore, Ess. Chlor. Land. 299 = nanus.
baeticus, *Boiss. Elench.* 80.—Reg. Mediterr. occ.
baeticus, Willk. ex Webb, Otia Hisp. 42 = Willkommii.
Boivini, Coss. ex Nym. Consp. 149 = Cossonii.
Boivini, *Webb, Iter Hisp.* 51.—Reg. Mediterr. occ.
Boivini, Welw. ex Webb, Otia Hisp. 32 = Escayracii.
Boivini, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 443 = Webbianus.
Bourgaeanus, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 289 = baeticus.
brachyacanthus, *Boiss. Diagn.* Ser. II. ii. 9.—Hispan.
canescens, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 159.—Hispan.
capensis, Linn. Sp. Pl. 741 = Mundtia spinosa.
compositus, Moench, Meth. 289 = europaeus.
Cossonii, *Nym. Syll.* 278.—Hispan.
densus, *Welw. ex Webb, Otia Hisp.* 43.—Lusitan.
ericetorum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 449 = Welwitschianus.
erinaceus, *Welw. ex Webb, Otia Hisp.* 44.—Lusitan.
Escayracii, *Nym. Syll.* 278.—Lusitan.
europaeus, Brot. Fl. Lusit. ii. 78 = nanus.
europaeus, *Linn. Sp. Pl.* 241.—Europ. occ.
europaeus, Savi, Alb. Tosc. i. 228 = parviflorus ?
floridus, Salisb. Prod. 329 = europaeus.
Funkii, Webb, Otia Hisp. 42 ; Nym. Consp. 148 = Willkommii.
Gallii, *Planch. in Ann. Sc. Nat.* Sér. III. xi. (1849) 213. t. 9.—Europ. occ.
genistoides, *Brot. Fl. Lusit.* ii. 78.—Lusitan.
genistoides, Roxas Clem. Ensay. Vid, 291 = spartioides.
genistoides, Salzm. ex Lindl. Bot. Reg. t. 1452 = Boivini.
grandiflorus, Pourr. in Mém. Acad. Toul. iii. (1788) 333 = europaeus.
hibernicus, G. Don, Gen. Syst. ii. 148 = europaeus.
hispanicus, Hort.—Cf. Gard. Chron. (1882) i. 668 = Genista hispanica.
hispanicus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 426 = Genista hirsuta.
ianthocladus, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 290 = baeticus.
intermedius, Le Gall, in Congr. Sc. Fr. xvi. (1850) 189 ; Fl. Morb. 815 = Gallii.

ULEX :—
Jussiaei, *Webb, in Ann. Sc. Nat.* Sér. III. xvii. (1852) 291.—Lusitan.
lanuginosus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 424 = Genista hirsuta.
lucidus, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 469, sphalm. = seq.
luridus, Nym. Syll. 278 = Webbianus.
lusitanicus, *Mariz, in Bol. Soc. Broter.* ii. (1884) 115.—Lusitan.
lusitanicus, Pourr. ex Colm. sec. Willk. & Lange, Prod. Fl. Hisp. iii. 449 = Welwitschianus.
major, Thore, Ess. Chlor. Land. 299 = europaeus.
Megalorites, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 469 = Boivini.
micranthus, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 236 = Willkommii.
minor, Roth, Catalect. fasc. i. 83 = nanus.
mitis, Hort. ex G. Don, in Loud. Hort. Brit. 280 = europaeus.
nanus, *Forst. in Symons, Syn.* 160 ; *Sm. Fl. Brit.* 757.—Europ. occ.
nanus, Hort.—Cf. Gard. Chron. (1882) i. 668 = Genista hispanica.
opistolepis, Webb, in Ann. Sc. Nat. Sér. III. xvii. (1852) 291 = europaeus.
parviflorus, Losc. & Pard. Ser. Inconf. ed. II. 91 = recurvatus.
parviflorus, *Pourr. in Mém. Acad. Toul.* iii. (1788) 334.—Europ. occ.
provincialis, Le Gall, Fl. Morb. 128 = Gallii.
provincialis, Loisel. in Desv. Journ. Bot. ii. (1809) 361 = parviflorus.
recurvatus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 449 = Welwitschianus.
recurvatus, *Willk. l. c.* 448.—Hispan.
Salzmanni, *Willk. l. c.* 469.—Hispan.
scaber, Kunze, in Flora, xxix. (1846) 696. t. 2 = baeticus.
sparsiflorus, Lange, in Kjoeb. Vidensk. Meddel. (1881) 104 (Quid ?).—Hispan.
spartioides, *Nym. Syll.* 278.—Lusitan.
spectabilis, *Welw. ex Webb, Otia Hisp.* 27.—Lusitan.
strictus, Hort. ex Del. Sem. Hort. Heidelb. (1852) 8.—Cf. Linnaea, xxvi. (1853-55) 739 (an seq. ?).—Hab. ?
strictus, Mackay, in Trans. Roy. Irish Acad. xiv. (1824) 166 = europaeus.
Vaillantii, Nym. Syll. 278 = Boivini.
vernalis, Thore, Ess. Chlor. Land. 299 = europaeus.
Webbianus, *Coss. Notes Crit.* ii. 32.—Hispan.
Welwitschianus, Coss. ex Mariz, in Bol. Soc. Broter. ii. (1884) 116 = baeticus.
Welwitschianus, *Planch. in Ann. Sc. Nat.* Sér. III. xi. (1849) 216.—Lusitan.
Willkommii, *Webb, l. c.* xvii. (1852) 290.—Hispan.

ULINA, Opiz, Seznam, 100 (1852) = **Inula**, Linn. (Compos.).
germanica, Opiz, l. c. = I. germanica.
hirta, Opiz, l. c. = I. hirta.
salicina, Opiz, l. c. = I. salicina.

ULLOA, Pers. Syn. i. 218 (1805) = **Juanulloa**, Ruiz & Pav. (Solan.).
parasitica, Pers. l. c. = J. parasitica.

ULLUCUS, Caldas, in Seman. Nuov. Granad. (1809) 185. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 77.
GANDOLA, Moq. in DC. Prod. xiii. II. 224 (1849).
MELLOCA, Lindl. in Gard. Chron. (1847) 685.
Kunthii, Moq. in Bibl. Univ. Genèv. xi. (1849) 80 = tuberosus.
tuberosus, *Caldas, in Seman. Nuov. Granad.* (1809) 185.—Reg. Andin.

ULMARIA, [Tourn.] Hill, Hort. Kew. 213 (1768) = **Spiraea**, Linn. (Rosac.).
Aruncus, Hill, l. c. 214 = S. Aruncus.
denudata, Opiz, Seznam, 100 = S. Ulmaria.
Filipendula, Hill, Hort. Kew. 214 = S. Filipendula.
glauca, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 370 = S. Ulmaria.
obtusiloba, Opiz, Seznam, 100 = S. Ulmaria.
palustris, Moench, Meth. 663 = S. Ulmaria.
pentapetala, Gilib. Fl. Lituan. ii. 236 = S. Ulmaria.
rubra, Hill, Hort. Kew. 214.—Am. bor.
Spiraea-Ulmaria, Hill, l. c. = S. Ulmaria.
trifoliata, Hill, l. c. = Gillenia trifoliata.
variegata, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 863 = S. Ulmaria.

ULMUS, [Tourn.] Linn. Syst. ed. I (1735). *URTICACEAE*, Benth. & Hook. f. iii. 351.
CHAETOPTELEA, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 76.
MICROPTELEA, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 358.
Abelicea, Sibth. & Sm. Fl. Graec. Prod. i. 172 = Zelkova cretica.
acuta, Dum. Fl. Belg. 25 = pedunculata.
alata, *Michx. Fl. Bor. Am.* i. 173.—Am. bor.
alba, Kit. ex Willd. Berl. Baumz. 518 = pedunculata.
alba, Rafin. Fl. Ludov. 115 = americana.
americana, Ait. Hort. Kew. ed. I. i. 319 = fulva.
americana, *Linn. Sp. Pl.* 226.—Am. bor.
americana, Planch. in DC. Prod. xvii. 155, partim = racemosa.
angustifolia, Moench, Verz. Ausl. Baeume, 137 = campestris.
antarctica, Hort. ex C. Koch, Dendrol. ii. I. 418 = montana.
aquatica, Rafin. Fl. Ludov. 165 = Planera aquatica.
aurea, Hort. ex C. Koch, Dendrol. ii. I. 412 = campestris.
batavina, Hort. ex C. Koch, l. c. 414 = montana.
Berandii, Hort. ex C. Koch, l. c. 409 = campestris.
betulaefolia, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. iii. 1376 = campestris.
betuloides, Hort. ex Steud. Nom. ed. II. ii. 727 = praec. ?
campestris, *Linn. Sp. Pl.* 225.—Europ. ; Oriens.
campestris, Linn. l. c., partim = montana.
campestris, O. F. Muell. in Fl. Dan. t. 632 = pedunculata.
campestris, Walt. Fl. Carol. 111 = Planera aquatica.
carpinifolia, Borkh. in Rhein. Mag. i. (1793) 498 = campestris.
cebennensis, Audib. ex Planch. in DC. Prod. xvii. 160 = montana.
chinensis, Desf. Tabl. ed. I. 212 ; Pers. Syn. i. 291 = campestris.
ciliata, Ehrh. Beitr. v. 161 ; vi. 88 = pedunculata.
cinerea, Hort. ex Planch. in DC. Prod. xvii. 160 = montana.
cornubiensis, Hort. ex C. Koch, Dendrol. ii. I. 409 = campestris.
corylacea, Dum. Fl. Belg. 25 = montana ?
corylifolia, Bor. Fl. Centr. Fr. ed. III. ii. 577 = montana.
corylifolia, Host, Fl. Austr. i. 329 = campestris.
crassifolia, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 169.—Am. bor.
crispa, Willd. Enum. Hort. Berol. 295 = fulva.
Davidiana, *Planch. in Compt. Rend. Par.* lxxiv. (1872) i. 1498, nomen ; *et in DC. Prod.* xvii. 158.—China.
dentata, Rafin. New Fl. Am. iii. 39 (Quid ?).—Am. bor.
dimidiata, Rafin. l. c. (Quid ?).—Am. bor.
effusa, Sibth. Fl. Oxon. 87 = montana.
effusa, Willd. Prod. Fl. Berl. 94 = pedunculata.
elliptica, C. Koch, in Linnaea, xxii. (1849) 599 = campestris.
erosa, *Roth, Nov. Pl. Sp.* 183.—Ind. or.
erosa, Wall. Cat. n. 3546 = Wallichii.
excelsa, Borkh. Handb. Forstbot. i. 839 = montana.
exoniensis, Hort. ex Loud. Arbor. Brit. iii. 1399 = montana.
expansa, Rota, in Bot. Zeit. xiii. (1855) 469 = montana.
floridana, Chapm. Fl. S. U. St. 416 = americana.
foliacea, Gilib. Exercit. ii. 395 = campestris.
Fordii, Hort. ex Loud. Arbor. Brit. iii. 1399 = montana.
forficata, Presl, in Verh. Ges. Vaterl. Mus. Boehm. (1841) 76 = montana ?
fulva, *Michx. Fl. Bor. Am.* i. 172.—Am. bor.
fungosa, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vi. 384 = campestris.
gigantea, Hort. ex C. Koch, Dendrol. ii. I. 417 = montana.
glabra, Huds. Fl. Angl. ed. I. 95 = montana.
glabra, Mill. Gard. Dict. ed. VIII. n. 4 = campestris.
glaucescens, *Franch. Pl. Davidian.* i. (1884) 267.—China.
glutinosa, Willd. Enum. Hort. Berol. Suppl. 14 = montana ?
hollandica, Duroi, Harbk. Baumz. ii. 505 = pedunculata.
hollandica, Mill. Gard. Dict. ed. VIII. n. 5 = campestris.

ULMUS :—
Hookeriana, Planch. in DC. Prod. xvii. 162 = lancifolia.
horizontalis, Hort. ex Loud. Arbor. Brit. iii. 1398 = montana.
humilis, Amman, ex Steud. Nom. ed. I. 862 = pumila.
integrifolia, Roxb. in Willd. Sp. Pl. i. 1326 = Holoptelea integrifolia.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 28 = parvifolia.
Keaki, Siebold, l. c. = Zelkova acuminata.
Keia-ki, Planch. in DC. Prod. xvii. 163 = praec.
laevigata, Royle, Illustr. Bot. Himal. 341 = Wallichii.
laevis, Pall. Fl. Ross. i. 75. t. 48. f. F = pedunculata.
lancifolia, *Roxb. Hort. Beng.* 86 ; *Fl. Ind.* ii. 66.—Reg. Himal. ; Burma.
latifolia, Moench, Meth. 333 = montana.
leucocarpa, Schur, Enum. Pl. Transs. 603 = montana.
leucosperma, Schur, l. c. = praec.
longifolia, Rafin. New Fl. Am. iii. 38 (Quid ?).—Am. bor.
macrocarpa, *Hance, in Journ. Bot.* vi. (1868) 332.—China.
major, Hohen. in Bull. Soc. Nat. Mosc. (1838) 261 = campestris.
major, Sm. Engl. Bot. t. 2542 = montana.
mexicana, *Planch. in DC. Prod.* xvii. 156.—Mexic.
microphylla, Pers. Syn. i. 291 = pumila.
minor, Mill. Gard. Dict. ed. VIII. n. 6 = campestris.
modiolina, Dum.-Cours. Bot. Cult. ed. II. vi. 384 = campestris.
mollifolia, Marsh. Arbust. 156 = americana.
montana, *With. Bot. Arr. Brit. Pl.* ed. III. ii. 279 ; *Sm. Engl. Fl.* ii. 22.—Europ. ; As. bor.
nana, Borkh. Handb. Forstbot. i. 839 = campestris.
nemoralis, Ait. Hort. Kew. ed. I. i. 320 = Zelkova crenata.
nemoralis, Boiss. ex Planch. in DC. Prod. xvii. 156 = campestris.
nemorosa, Borkh. Handb. Forstbot. i. 846 = campestris.
nigra, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. iii. 1398 = montana.
nitens, Moench, Meth. 333 = campestris.
nuda, Ehrh. Beitr. v. 160 ; vi. 86 = campestris.
obovata, Rafin. New Fl. Am. iii. 38 (Quid ?).—Am. bor.
octandra, Schkuhr, Handb. 178. t. 67 = pedunculata.
ontariensis, Hort. ex Steud. Nom. ed. II. ii. 728 = montana.
opaca, Nutt. N. Am. Sylv. i. 51 = crassifolia.
parvifolia, *Jacq. Hort. Schoenb.* iii. 6. t. 262.—As. temp.
pedunculata, *Fouger. in Mém. Acad. Sc. Par.* 1784 (1787), 215. t. 2.—Europ. ; Reg. Cauc.
pendula, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. 1398 = montana.
pendula, Willd. Berl. Baumz. 519 ; Enum. Hort. Berol. Suppl. 14 = fulva.
pinguis, Rafin. Fl. Ludov. 115 (Quid ?).—Am. bor.
planereoides ×, Carr. in Rev. Hortic. (1875) 286, 287.
planifolia, Hort. ex Loud. Arbor. Brit. iii. 1377 = campestris.
polygama, Juss. ex Poir. Encyc. iv. 611 ; Rich. in Michx. Fl. Bor. Am. ii. 248 = Zelkova crenata.
procera, Salisb. Prod. 391 = campestris.
pubescens, Walt. Fl. Carol. 112 = fulva.
pumila, Hort. ex Planch. in DC. Prod. xvii. 161 = parvifolia.
pumila, *Linn. Sp. Pl.* 226.—As. bor.
pumila, Pall. Reise, iii. 106 = campestris.
pumila, Walt. Fl. Carol. 111 = alata.
purpurea, Hort. ex Planch. in DC. Prod. xvii. 158 = campestris.
racemosa, Borkh. Handb. Forstbot. i. 851 = pedunculata.
racemosa, *D. Thomas, in Am. Journ. Sc.* Ser. I. xix. (1831) 170.—Am. bor.
reticulata, Dum. Fl. Belg. 25 = pedunculata.
rotundifolia, Carr. in Rev. Hortic. (1868) 374 = macrocarpa ?
rubra, Hort. ex Loud. Arbor. Brit. iii. 1398 = montana.
rubra, Michx. f. Hist. Arb. Am. iii. 278. t. 6 = fulva.
rugosa, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. iii. 1398 = montana.

ULMUS :—
sarniensis, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. iii. 1376 = campestris.
sativa, Mill. Gard. Dict. ed. VIII. n. 3 ; Duroi, Harbk. Baumz. ii. 502 = campestris.
sativa, Pistohlk, in [Hupel,] Nord. Miscellan. xvii. ; ex Ledeb. Fl. Ross. iii. 648 = pedunculata.
scabra, Duroi, Harbk. Baumz. ii. [543] Zusatz = campestris.
scabra, Mill. Gard. Dict. ed. VIII. n. 2 = montana ?
scampstoniensis, Hort. ex Steud. Nom. ed. II. ii. 728 = campestris.
sparsa, Dum. Fl. Belg. 25 = campestris.
striata, Bosc, Encyc. Agric. vii. 623, nomen (Quid ?).—Am. bor.
stricta, Lindl. Syn. Brit. Fl. ed. II. 227 = campestris.
suberosa, Moench, Verz. Ausl. Baeume, 136 ; Ehrh. Beitr. vi. 87 = campestris.
surculosa, Stokes, Bot. Mat. Med. ii. 35 = campestris.
tetrandra, Schkuhr, Handb. 178. [t. 57] = campestris.
tiliaefolia, Host, Fl. Austr. i. 329 = campestris.
tomentosa, Borkh. Handb. Forstbot. i. 856 = montana.
tortuosa, Host, Fl. Austr. i. 330 = campestris.
transbaicalensis, Steud. Nom. ed. I. 862 = pumila.
turkestanica, *Regel, Gartenfl.* (1884) 28, in textu.—Turkestan.
urticaefolia, *Audib. Cat. Hort. Tonn.* (1817) 23 = montana.
viminalis, Lodd. Cat. (1836) ; ex Loud. Arbor. Brit. iii. 1376 = campestris.
virgata, Roxb. Hort. Beng. 21 ; Fl. Ind. ii. 67 = parvifolia.
viscosa, Desf. ex Steud. Nom. ed. I. 862 = montana ?
vulgaris, Pall. Reise, iii. 314 ; Dum. Fl. Belg. 25 = campestris.
Wallichiana, Planch. in Compt. Rend. Par. lxxiv. (1872) 1. 1498 = seq.
Wallichii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 277–8.—Reg. Himal.
Webbiana, Lee, ex C. Koch, Dendrol. ii. 1. 412 = campestris.

ULOMA, Rafin. Fl. Tellur. ii. 62 (1836) = **Colea**, Boj. (Bignon.).
Telfairia, Rafin. l. c. = C. Telfairii.

ULOPTERA, Fenzl, in Flora, xxvi. (1843) 460 = **Ferula**, Tourn. (Umbell.).
macrocarpa, Fenzl, l. c. (= *Ferulago macrocarpa*).—Mesopot.

ULOSPERMUM, Link, Enum. Hort. Berol. i. 267 (1821) = **Capnophyllum**, Gaertn. (Umbell.).
dichotomum, Link, l. c. = C. dichotomum.
siculum, Link, Handb. i. 343 = Hippomarathrum siculum.
tauricum, Link, l. c. = Sanicula crithmifolia.

ULOSTOMA, D. Don, in G. Don, Gen. Syst. iv. 196 (1837) = **Gentiana**, Tourn.
filamentosa, D. Don, l. c. = G. filamentosa.

ULRICIA, Jacq. ex Steud. Nom. ed. I. 862 (1821) = **Lepechinia**, Willd. (Labiat.).
pyramidata, Jacq. ex Steud. l. c. = L. spicata.

ULTICONA, Rafin. Sylva Tellur. 55 (1838) = **Hebecladus**, Miers (1845) (Solan.).
aspera, Rafin. l. c. = H. asperus.
biflora, Rafin. l. c. = H. biflorus.
viridiflora, Rafin. l. c. = H. viridiflorus.

ULUXIA, Juss. in Dict. Sc. Nat. x. 103 (1818) = **Columellia**, Ruiz & Pav.

ULVA, Hall. Enum. Stirp. Helv. i. 242 (1742) = **Carex**, Dill. (Cyper.).

UMARI, Adans. Fam. ii. 342 (1763) = **Geoffraea**, Linn. (Legumin.).

UMBELLULARIA, Nutt. N. Am. Sylv. 87 (1842).
LAURINEAE, Benth. & Hook. f. iii. 162.
DRIMOPHYLLUM, Nutt. N. Am. Sylv. 85 (1842).
SCIADIODAPHNE, Reichb. Nom. 70 (1841).

UMBELLULARIA :—
californica, *Nutt. N. Am. Sylv.* 87.—Am. bor. occ.
parvifolia, *Hemsl. Biol. Centr. Am. Bot.* iii. 77.—Mexic.

UMBILICARIA, Pers. Syn. i. 510 (1805) = **Cotyledon**, Tourn. (Crassul.).

UMBILICUS, DC. in Bull. Soc. Philom. (1801) 49 = **Cotyledon**, Tourn. (Crassul.).
affinis, Schrenk, Enum. Pl. Nov. 72.—Soongar.
Aizoon, Fenzl, Pugill. Pl. Nov. Syr. 15.—As. Min.
alpestris, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 354.—Soongar.
botryoides, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 308 = C. Umbilicus.
chloranthus, Heldr. & Sart. ex Boiss. Fl. Orient. ii. 768.—Europ. or. ; As. Min.
chrysanthus, Boiss. & Heldr. Diagn. Ser. I. x. 15.—As. Min.
deflexus, Pomel, Nouv. Mat. Fl. Atl. 324.—Alger.
elymaiticus, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 770.—Persia.
erectus, DC. Fl. Fr. iv. 384 = C. lusitanica.
erubescens, Maxim. Prim. Fl. Amur. 114.—Manchur.
fimbriatus, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 241 = C. fimbriata.
gaditanus, Boiss. Diagn. Ser. I. vi. 58.—Alger.
glaber, Regel & Winkler, in Act. Hort. Petrop. vi. (1879) 302.—Turkestan.
globulariaefolius, Fenzl, Pugill. Pl. Nov. Syr. 15 = C. globulariaefolius.
Haussknechtii, Boiss. & Reut. ex Boiss. Fl. Orient. ii. 774.—Mesopot.
Heylandianus, Webb & Berth. Phyt. Canar. i. 176.—Ins. Canar.
hispidus, DC. Prod. iii. 399 = C. Mucizonia.
horizontalis, DC. l. c. 400 = C. Umbilicus.
horizontalis, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 258 = *Schmidtii*.
intermedius, Boiss. Fl. Orient. ii. 769.—As. Min. ; Persia.
Lampusae, Kotschy, in Unger & Kotschy, Ins. Cypern, 312 = Sedum Lampusae.
leucanthus, G. Don, Gen. Syst. iii. 113 = C. leucantha.
libanoticus, DC. Prod. iii. 399 = C. libanotica.
Lievenii, G. Don, Gen. Syst. iii. 113 = C. Lievenii.
Lievenii, Ledeb. Fl. Ross. ii. 173.—Persia ; Sibir.
linearifolius, Franch. in Ann. Sc. Nat. Sér. VI. xvi. (1883) 290.—Turkestan.
lineatus, Boiss. Diagn. Ser. I. x. 14.—Palaestin.
lineolatus, Boiss. Fl. Orient. ii. 769 = *intermedius*.
linifolius, Rupr. Sert. Tiansch. 46.—As. centr.
luteus, Decne. in Jacquem. Voy. Bot. 62. t. 73 = C. Oreades.
luteus, Webb & Berth. Phyt. Canar. i. 177.—Ins. Canar.
malacophyllus, DC. Prod. iii. 400 = C. malacophylla.
mexicanus, Schlecht. in Linnaea, xiii. (1839) 409 = C. mexicana.
micranthus, Pomel, Nouv. Mat. Fl. Atl. 325.—Alger.
microstachyum, Kotschy, in Unger & Kotschy, Ins. Cypern, 311 = Sedum microstachyum.
multicaulis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 93 = C. Lievenii.
oppositifolius, Ledeb. Fl. Ross. ii. 176.—Reg. Cauc.
Oreades, Decne. in Jacquem. Voy. Bot. 62 = C. Oreades.
oxypetalus, Boiss. Diagn. Ser. I. iii. 15 = *tropaeolifolius*.
pallidus, Schott & Kotschy, in Oestr. Bot. Wochenbl. vii. (1857) 229 = *chrysanthus*.
paniculatus, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 583.—Kokania.
parviflorus, Boiss. Fl. Orient. ii. 768. n. 2 = *Spruneri-anus*.
parviflorus, Boiss. l. c. n. 3 = *chloranthus*.
parviflorus, DC. Prod. iii. 400 = C. parviflora.
patens, Pomel, Nouv. Mat. Fl. Atl. 324.—Alger.
patulus, Pomel, l. c. 323.—Alger.
pendulinus, DC. Pl. Grass. t. 162 = C. Umbilicus.
persicus, Boiss. Diagn. Ser. I. iii. 14.—Persia.
Pestalozzae, Boiss. Diagn. Ser. II. v. 96.—As. Min.
platyphyllus, Schrenk, Enum. Pl. Nov. 71.—Persia ; Soongar.
pubescens, G. Don, Gen. Syst. iii. 111 = Sedum pilosum.
pulvinatus, Rupr. Sert. Tiansch. 47.—Turkestan.

UMBILICUS:—
radicans, Klotzsch, in Bot. Ergeb. Waldem. Reise, 143 = Sedum rosulatum.
radiciflorus, Steud. ex Boiss. Fl. Orient. ii. 772 = *libanoticus*.
ramosissimus, Maxim. Prim. Fl. Amur. 472 = Cotyledon fimbriata.
samius, DC. Prod. iii. 399 = C. samia.
Schmidtii, Bolle, in Bonplandia, vii. (1859) 245.—Ins. Canar.
sedoides, DC. Prod. iii. 400 = C. sedoides.
Semenovii, Regel & Herd. in Bull. Soc. Nat. Mosc. xxxix. (1866) II. 65.—As. temp.
semiensis, J. Gay, ex A. Rich. Tent. Fl. Abyss. i. 308 = C. simensis.
Sempervivum, DC. Prod. iii. 399 = C. Sempervivum.
sessilis, Dulac, Fl. Hautes-Pyr. 326 = C. sedoides.
serratus, DC. Prod. iii. 400 = C. serrata.
simplex, C. Koch, in Linnaea, xix. (1847) 41 = C. Umbilicus.
spathulatus, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 92 = C. spathulata.
spinosus, DC. Prod. iii. 400 = C. spinosa.
Sprunerianus, Boiss. Diagn. Ser. I. vi. 57.—Graecia.
stamineus, Ledeb. Fl. Ross. ii. 174.—Sibir.
subulatus, G. Don, Gen. Syst. iii. 112 = Sedum subulatum.
thyrsiflorus, DC. Prod. iii. 400 = C. leucantha.
tropoeolifolius, Boiss. Diagn. Ser. I. iii. 14.—Persia.
tuberculosus, DC. ex Steud. Nom. ed. II. ii. 729 = C. tuberculosa.
turkestanicus, Regel & Winkl. in Act. Hort. Petrop. vi. (1879) 301.—Turkestan.
Winkleri, Willk. in Ber. Deutsch. Bot. Ges. i. (1883) 269.—Europ.

UMSEMA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Pontederia**, Linn.
mucronata, Rafin. l. c. = P. cordata.
obtusifolia, Rafin. l. c. = P. cordata.

UNANNEA, Steud. Nom. ed. II. ii. 729 (1841), sphalm. = seq.

UNANUEA, Ruiz & Pav. Ic. Fl. Per. Ined.; ex Benth. in DC. Prod. x. 380 (1846) = **Stemodia**, Linn. (Scrophul.).
febrifuga, Ruiz & Pav. l. c.; ex Benth. l. c. 382 = S. suffruticosa.

UNCARIA, Burch. Trav. i. 536 (1822) = **Harpagophytum**, DC. (Pedal.).
procumbens, Burch. l. c. = H. procumbens.

UNCARIA, Schreb. Gen. i. 125 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 31.
AGYLOPHORA, Neck. Elem. i. 145 (1790).
OUROUPARIA, Aubl. Pl. Gui. i. 177. t. 68 (1775), nomen prius.
UNCINARIA, Reichb. Nom. 230 (1841).
acida, [Hook. f. & Thoms.] ex Hook. f. Fl. Brit. Ind. iii. 31 = dasyoneura.
acida, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 520.—Ins. Penang.
aculeata, Willd. in Usteri, Delect. Opusc. Bot. ii. (1793) 200 = guianensis.
africana, *G. Don, Gen. Syst.* iii. 471.—Afr. trop.
appendiculata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 222.—N. Guin.
attenuata, *Korth. Verh. Nat. Gesch. Bot.* 170. t. 34.—Malaya.
brevispina, Maing. ex Hook. f. Fl. Brit. Ind. iii. 31 = dasyoneura.
callophylla, *Korth. Verh. Nat. Gesch. Bot.* 170.—Borneo.
canescens, *Korth. l. c.* 172.—Malaya.
cirrhiflora, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 520 (Quid?).—Malaya.
dasyoneura, *Korth. Verh. Nat. Gesch. Bot.* 169.—Malaya; Zeylan.
elliptica, R. Br. in Wall. Cat. n. 6104 = dasyoneura.
eumececarpa, Korth. ex Miq. Fl. Ind. Bat. ii. 143 = pedicellatum.
eurhyncha, *Miq. l. c. Suppl.* 539.—Ins. Banca.
ferrea, *DC. Prod.* iv. 348.—Malaya.
ferrea, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 105 = Hookeri.
ferruginea, DC. Prod. iv. 348 = pedicellata.
ferruginea, Kurz, For. Fl. Brit. Burma, ii. 69 = sclerophylla.

UNCARIA:—
florida, *Vidal, Phan. Cuming. Philipp.* 176.—Ins. Philipp.
Gambier, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 517.—Malaya.
Gambier, Thw. Enum. Pl. Zeyl. 138 = dasyoneura.
Gambier, Wall. Cat. n. 6103 C, partim = attenuata.
glabrata, *DC. Prod.* iv. 348.—Java.
glabrata, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 105 = florida.
guianensis, *J. F. Gmel. Syst.* 370.—Guian.
Hallii, Korth. Verh. Nat. Gesch. Bot. 165. t. 33 = pedicellata.
homomalla, *Miq. Fl. Ind. Bat.* ii. 343.—Ind. or.; Malaya.
Hookeri, *Vidal, Phan. Cuming. Philipp.* 177.—Ins. Philipp.
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 151 = ferrea.
inermis, Willd. in Usteri, Delect. Opusc. Bot. ii. (1793) 199. t. 3 = Mitragyna africana.
insignis, DC. Prod. iv. 348 = sclerophylla.
jasminiflora, *Wall. Cat.* n. 6103 C, F.—Malaya.
laevigata, *Wall. l. c.* 6111.—Reg. Himal.; Burma.
lanosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 131.—Malaya.
Lobbii, *Hook. f. Fl. Brit. Ind.* iii. 33.—Singapore.
macrophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 132.—Reg. Himal.
nemorosa, *Korth. Verh. Nat. Gesch. Bot.* 166.—Sumatra.
ovalifolia, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 519.—Malaya.
ovata, *R. Br. in Wall. Cat.* n. 6112, partim.—Malaya.
paucinervis, *Teysm. & Binn. Cat. Hort. Bog.* 117.—Java.
pedicellata, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 520.—Malaya.
pilosa, *Roxb. ll. cc.* 86; 520.—Reg. Himal.; Burma.
pilosa, Wall. Cat. n. 6108 C = homomalla.
pteropoda, *Miq. Fl. Ind. Bat.* ii. 343.—Malaya.
rhynchophylla, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 108 = Nauclea rhynchophylla.
Roxburghiana, *Korth. Verh. Nat. Gesch. Bot.* 172.—Malaya.
Roxburghii, Korth. Obs. Naucl. Ind. 17 = praec.
sclerophylla, DC. in Deless. Ic. Sel. iii. 48. t. 81 = attenuata.
sclerophylla, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 520.—Burma; Malaya.
sclerophylla, Wall. Cat. n. 6105 = pedicellata.
sessilifolia, Roxb. in Wall. l. c. n. 6107 = macrophylla.
sessilifructus, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 520.—Reg. Himal.; Burma.
setiloba, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 223.—Java.
speciosa, Wall. Cat. n. 6106 = sclerophylla.
spinosa, Raeusch. Nom. ed. III. 55 = guianensis.
surinamensis, *Miq. in Linnaea*, xix. (1847) 129.—Guian.
tomentosa, *DC. Prod.* iv. 349.—N. Granat.
Wallichii, *Korth. Obs. Naucl. Ind.* 17, nomen (Quid?).—Ind. or.

UNCARIOPSIS, Karst. in Linnaea, xxx. (1859–60) 152; Fl. Columb. i. 181. t. 90 (1861?). *RUBIACEAE*, Benth. & Hook. f. ii. 67.
suaveolens, *Karst. ll. cc.*—N. Granat.

UNCIFERA, Lindl. in Journ. Linn. Soc. iii. (1859) 39. *ORCHIDEAE*, Benth. & Hook. f. iii. 579.
acuminata, *Lindl. l. c.* 40.—Reg. Himal.
heteroglossa, *Reichb. f. in Gard. Chron.* (1878) II. 234.—Hab.?
obtusifolia, *Lindl. in Journ. Linn. Soc.* iii. (1859) 40.—Reg. Himal.

UNCINA, C. A. Mey. in Ledeb. Fl. Alt. iv. (1833) 234, sphalm. = **Uncinia**, Pers. (Cyperac.).

UNCINARIA, Reichb. Nom. 230 (1841) = **Uncaria**, Schreb. (Rubiac.).

UNCINIA, Pers. Syn. ii. 534 (1807). *CYPERACEAE*, Benth. & Hook. f. iii. 1072.
affinis, Colenso, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 392 = riparia.
alopecuroides, Colenso, in Trans. N. Z. Inst. xv. (1883) 335 = australis.
angusta, Nees, in Linnaea, ix. (1835–36) 305 = Douglasii.

UNCINIA:—
angustata, Boeck. in Linnaea, xli. (1877) 347, in syn. = Douglasii.
australis, R. Br. Prod. 241, in obs. = leptostachya.
australis, *Pers. Syn.* ii. 534.—N. Zel.; Ins. Pacif.
Banksii, Boott, in Hook. f. Fl. N. Zel. i. 287 = riparia.
bella, Phil. in Linnaea, xxx. (1859–60) 204 = macrophylla.
bractata, Colenso, in Trans. N. Z. Inst. xvi. (1884) 341 = australis.
brevicaulis, *Thou. Fl. Trist. Acugna*, 35. t. 6.—Ins. Tristan da Cunha.
breviculmis, Carmich. in Trans. Linn. Soc. xii. (1819) 508 = brevicaulis.
breviseta, Torr. in Ann. Lyc. N. York, iii. (1836) 428 = Carex filifolia (cf. Addend.).
caespitosa, *Colenso, ex Boott, in Hook. f. Fl. N. Zel.* i. 287.—N. Zel.
Cheesemanniana, Boeck. in Engl. Bot. Jahrb. v. (1884) 521 = compacta.
chlorostachya, Phil. in Linnaea, xxxiii. (1864–65) 275 = phleoides.
compacta, *R. Br. Prod.* 241.—Reg. temp.; Austral.
compacta, A. Rich. Voy. Astrolab. Bot. 118 = australis.
Cumingii, Nees, in Linnaea, ix. (1835–36) 305 = phleoides.
debilior, F. Muell. Fragm. viii. 151 = filiformis.
De-la-Costa, Steud. in Lechl. Berb. Am. Austr. 52 = macloviana.
digyna, Hochst. ex Steud. Syn. Pl. Cyp. 244 = Carex parasitica (in Addend.).
distans, Colenso, ex Boott, in Hook. f. Fl. N. Zel. i. 286 = leptostachya.
divaricata, Boott, in Hook. f. l. c. = compacta.
Douglasii, *Boott, in Hook. f. Fl. Antarct.* ii. 369.—Ins. J. Fernand.
Durvillei, Steud. Syn. Pl. Cyp. 243 = trichocarpa.
erinacea, *Pers. Syn.* ii. 534.—Am. austr.
europaea, J. Gay, in Flora, x. (1827) 26 = Carex microglochin.
fernandesiana, Nees, ex Boeck. in Linnaea, xli. (1877) 347, in syn. = Douglasii.
ferruginea, Boott, in Hook. f. Fl. N. Zel. i. 288. t. 64 B = australis.
filifolia, Nees, in Neuwied, Reise Nord-Amer. ii. 450 = Carex filifolia? (cf. Addend.).
filiformis, Boott, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 392 = riparia.
filiformis, *Colenso, ex Boott, in Hook. f. Fl. N. Zel.* i. 286.—Austral.; N. Zel.
Galeottii, Boott, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 400 = mexicana.
gracilis, Decne. in Bot. Voy. Astrolab. et Zélée, ii. t. 6. fig. B = tenuis.
gracilis, Thou. Fl. Trist. Acugna, 35. t. 5 = brevicaulis.
Hookeri, Boott, in Hook. f. Fl. Antarct. 91. t. 51 = rupestris.
Hookeri, Colenso, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 395 = compacta.
horizontalis, Colenso, in Trans. N. Z. Inst. xv. (1883) 334 = caespitosa.
jamaicensis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 272 = mexicana.
jamaicensis, *Pers. Syn.* ii. 534.—Am. trop.
Kingii, *R. Br. ex Boott, in Hook. f. Fl. Antarct.* ii. 370. t. 145.—Chili.
lasiocarpa, Steud. ex Boeck. in Linnaea, xli. (1877) 349 = phleoides.
laxiflora, Petrie, in Trans. N. Z. Inst. xvii. (1885) 271 = riparia.
Lechleriana, *Steud. Syn. Pl. Cyp.* 244.—Reg. Magellan.
Lehmanni, Nees, in Linnaea, x. (1836) 206 = Carex Lehmannii.
leptostachya, Phil. in Linnaea, xxxiii. (1864–65) 274 = trichocarpa.
leptostachya, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 116 (*Choix Pl. N. Zél.* 12. t. 5 B).—N. Zel.
Lindleyana, Kunth, Enum. Pl. ii. 526 = australis.
longearistata, Steud. Syn. Pl. Cyp. 243 = erinacea.
longifolia, E. Desv. in C. Gay, Fl. Chil. vi. 226. t. 72. f. 1 = phleoides.
longifolia, Kunth, Enum. Pl. ii. 527 = erinacea.
longifolia, Phil. ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 399 = phleoides.
longispica, Boeck. in Flora, xli. (1858) 650 = phleoides.

UNCINIA:—
macloviana, *Gaudich. in Freyc. Voy. Bot.* 412.—Ins. Falkland.
macrolepis, Decne. [et Hombron] in Bot. Voy. Astrolab. et Zélée, 13. t. 6. f. A = Lechleriana.
macrophylla, *Steud. Syn. Pl. Cyp.* 244.—Chili.
macrostachya, E. Desv. in C. Gay, Fl. Chil. vi. 229. t. 72. f. 2 = multifaria.
mexicana, *Steud. Syn. Pl. Cyp.* 243.—Mexic.
microglochin, Spreng. Syst. iii. 830 = Carex microglochin.
montana, *Phil. in Anal. Univ. Chil.* (1865) II. 322.—Chili.
Montteana, Phil. in Linnaea, xxx. (1859–60) 205 = phleoides.
Moseleyana, Boeck. in Flora, lxi. (1878) 170 = compacta.
multifaria, *Nees, ex Boott, in Hook. f. Fl. Antarct.* ii. 369.—Chili.
nepalensis, Nees, in Wight, Contrib. 129 = Carex linearis.
nervosa, Boott, in Hook. f. Fl. Tasman. ii. 102. t. 153 = compacta.
nigra, Colenso, in Trans. N. Z. Inst. xvii. (1885) 253 = australis.
obtusata, *Colenso, l. c.* xvi. (1884) 341.—N. Zel.
phalaroides, Boott, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 396 = macrophylla.
phalaroides, Nees, in Mart. Fl. Bras. ii. 1. 202 = Carex phalaroides.
Philippii, Hohen. ex Steud. Syn. Pl. Cyp. 243 = erinacea.
phleoides, C. A. Mey. in Bull. Acad. Brux. ix. (1842) 249 = mexicana.
phleoides, *Pers. Syn.* ii. 534.—Chili.
phyllostachya, Nees, in Mart. Fl. Bras. ii. 1. 201 = Carex seticulmis.
purpurata, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 272.—N. Zel.
rigida, Boeck. in Flora, lxv. (1882) 64 = brevicaulis.
rigida, Petrie, in Trans. N. Z. Inst. xvii. (1885) 271 = rubra.
rigidula, Steud. Syn. Pl. Cyp. 245 = australis.
riparia, *R. Br. Prod.* 241.—Austral.; N. Zel.
riparia, Hook. f. in Hook. Lond. Journ. Bot. iii. (1844) 417 = rupestris.
rubra, *Colenso, ex Boott, in Hook. f. Fl. N. Zel.* i. 287. t. 64 A.—N. Zel.
rupestris, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 117; (*Choix Pl. N. Zél.* 13. t. 5. f. A).—N. Zel.
scaberrima, Nees, in Linnaea, ix. (1835–36) 305 = australis.
scabra, Colenso, ex Boott, in Hook. f. Fl. N. Zel. i. 285 = leptostachya.
Sellowiana, Nees, in Mart. Fl. Bras. ii. 1. 201 = Carex Sellowiana.
Sinclairii, *Boott, ex Hook. f. Handb. N. Zeal. Fl.* 309. —N. Zel.
spartea, *Nees, in Linnaea*, x. (1836) 205.—Afr. austr.
spartea, Spreng. Syst. iii. 830 = praec.?
Sprengelii, Nees, in Linnaea, x. (1836) 205 = Carex Sprengelii.
tenella, *R. Br. Prod.* 241.—Austral.
tenuis, *Poepp. ex Kunth, Enum. Pl.* ii. 525.—Chili.
trichocarpa, E. Desv. in C. Gay, Fl. Chil. vi. 227. t. 72. f. 3 = phleoides.
trichocarpa, *C. A. Mey. Cyp. Nov.* 11. t. 4.—Chili.
Urvillei, Steud. Syn. Pl. Cyp. 243 = trichocarpa.

UNCINUS, Raeusch. Nom. ed. III. 63 (1797), sphalm. = ONCINUS, Lour. = **Melodinus**, Forst. (Apocyn.).

UNEDO, Hoffmgg. & Link, Fl. Port. i. 415 (1809) = **Arbustus**, Tourn. (Ericac.).
edulis, Hoffmgg. & Link, l. c. = A. Unedo.

UNGERIA, Nees, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 34 = **Cyperus**, Mich.
monocephala, Nees, ex C. B. Clarke, l. c. = C. Cephalotes.

UNGERIA, Schott & Endl. Melet. 27. t. 4 (1832). *STERCULIACEAE*, Benth. & Hook. f. i. 219.
floribunda, *Schott & Endl. l. c.*—Ins. Norfolc.

UNGERNIA, Bunge, in Bull. Soc. Nat. Mosc. xlix. (1875) II. 271. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 721.
flava, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 149.—Persia.

UNGERNIA:—
Oldhami, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) 76. —Japon.
trisphaera, *Bunge, in Bull. Soc. Nat. Mosc.* xlix. (1875) II. 273.—Persia.

UNGNADIA, Endl. Atakt. t. 36 (1834). *SAPINDACEAE*, Benth. & Hook. f. i. 398.
heptaphylla, Scheele, in Linnaea, xxii. (1849) 352 = speciosa.
heterophylla, Scheele, l. c. xxi. (1848) 589 = speciosa.
speciosa, *Endl. Atakt.* t. 36; *Nov. Stirp. Dec.* 86.—Texas.

UNGUACHA, Hochst. in Flora, xxvii. (1844) 103 = **Strychnos**, Linn. (Logan.).
simiarum, Hochst. l. c. = S. Unguacha.

UNIFOLIUM, Hall. Enum. Stirp. Helv. i. 288 (1742) = **Maianthemum**, Weber (Liliac.).
quadrifidum, All. Fl. Pedem. i. 124 = M. Convallaria.

UNIOLA, Linn. Gen. ed. I. 353 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1192.
CHASMANTHIUM, Link, Hort. Berol. i. 159 (1827).
NEVROCTOLA, Rafin. Neogenyt. 4 (1825).
TRISIOLA, Rafin. Fl. Ludov. 144 (1817).
TRIUNILA, Rafin. Neogenyt. 4 (1825).
bipinnata, Linn. Sp. Pl. ed. II. 104 = Eragrostis cynosuroides.
borbonica, *Steud. Syn. Pl. Gram.* 281.—Ins. Borbon.
capensis, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 360 = Eragrostis Uniolae.
ciliata, Beauv. Agrost. 75 (Quid?).—Hab.?
distichophylla, Labill. Nov. Holl. Pl. i. 21. t. 24 = Distichlis maritima.
flexuosa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 99 = Distichlis maritima.
gracilis, *Michx. Fl. Bor. Am.* i. 71.—Am. bor.
indica, Spreng. Syst. i. 349 = Eragrostis unioloides.
intermedia, *Bosc, ex Beauv. Agrost.* 75.—Am. bor.
Jardinii, *Steud. ex Jard. Herb. Côte Occ. Afr.* 8, nomen; *Steud. Syn. Pl. Gram.* 281.—Afr. trop.
lappacea, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 358 = Centotheca lappacea.
latifolia, *Michx. Fl. Bor. Am.* i. 71.—Am. bor.
maritima, Michx. l. c. = Distichlis maritima.
mucronata, Linn. Sp. Pl. ed. II. 104 = Eragrostis mucronata, *Benth. & Hook. f.* (in Addend.).
multiflora, Nutt. in Trans. Am. Phil. Soc. v. (1837) 148 = Distichlis maritima.
nitida, Baldw. ex Ell. Sketch, i. 167 = sessiliflora.
ornithoryncha, *Steud. Syn. Pl. Gram.* 280.—Am. bor.
paniculata, *Linn. Sp. Pl.* 71.—Am. bor.
paniculata, Roth, Neue Beytr. i. 98 = latifolia.
prostrata, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = Distichlis prostrata.
pungens, Beauv. Agrost. 75 = Sporobolus pungens.
pungens, Rupr. ex Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 234 = Distichlis maritima.
racemiflora, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Ins. S. Doming.
scoparia, Steud. Syn. Pl. Gram. 281 = Poa scoparia.
sessiliflora, *Poir. Encyc.* viii. 185.—Am. bor.
simplex, Trin. ex Steud. Nom. ed. II. ii. 730 (Quid?). —Hab.?
sparta, *Trin. in Linnaea*, x. (1836) 307.—Cuba.
spicata, Linn. Sp. Pl. 71 = Distichlis maritima.
stricta, Torr. in Ann. Lyc. N. York, i. (1824) 155 = Distichlis maritima.
thalassica, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 359 = Distichlis maritima.
virgata, Bartr. ex Schult. Syst. ii. 595 = gracilis.
virgata, Griseb. Fl. Brit. W. Ind. 531 = racemiflora.

UNISEMA, Rafin. in Med. Repos. N. York, v. (1819) 192; et in Journ. Phys. lxxxix. (1819) 261 = **Pontederia**, Linn.
acutifolia, Rafin. Med. Fl. ii. 107 = P. cordata.
deltifolia, Rafin. l. c. 105.—Am. bor.
heterophylla, Rafin. l. c. 108.—Am. bor.
lancifolia, Rafin. Atl. Journ. 178 = P. cordata.
latifolia, Rafin. Med. Fl. ii. 107.—Am. bor.
media, Rafin. l. c.—Am. bor.
mucronata, Rafin. l. c. = P. cordata.
obliquata, Rafin. l. c.—Am. bor.
obtusifolia, Rafin. in Med. Repos. N. York, v. (1819) 350 = P. cordata.

UNISEMA:—
orbiculata, Rafin. Med. Fl. ii. 108 = Pontederia rotundifolia.
pedunculatis, Rafin. New Fl. Am. ii. 75 = P. cordata.
Purshiana, Rafin. Med. Fl. ii. 107 = P. cordata.
rotundifolia, Rafin. l. c. 108.—Am. bor.
sagittata, Rafin. Fl. Ludov. 18 (= *Sagittaria cerulea*). —Am. bor.

UNJALA, Blume, Cat. Gew. Buitenz. 56 (1823) = **Heptapleurum**, Gaertn. (Araliac.).
bipinnata, Reinw. ex Blume, l. c.—Java.
decomposita, Reinw. ex Blume, l. c.—Java.
longifolia, Reinw. ex Blume, l. c. = H. longifolium.
Rheedii, Reinw. ex Blume, l. c. (*Rheedii*) = H. tomentosum, polybotryum, scandens, rigidum.
serrata, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 89 = H. serratum.

UNONA, Linn. f. Suppl. 270 (1781). *ANONACEAE*, Benth. & Hook. f. i. 24.
ANCANA, F. Muell. Fragm. v. 27. t. 35 (1865).
DESMOS, Lour. Fl. Cochinch. 352 (1790).
MEIOGYNE, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 12 (1865–66).
MELODORUM, Lour. Fl. Cochinch. 351 (1790).
acuminata, *DC. Syst.* i. 488.—Guian.
acuta, *Zoll. in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 11.—Java.
acutiflora, Dun. Monog. Anon. 116. t. 22 = Xylopia acutiflora.
aethiopica, Dun. l. c. 113 = Xylopia aethiopica.
Alphonsii, *Steud. Nom.* ed. II. ii. 730.—Cuba.
Alphonsii, Wall. Cat. n. 6426 = U. Dasymaschala.
altissima, Russ. ex Wall. Cat. n. 6442 = Polyalthia longifolia.
ambongoënsis, Baill. Adansonia, viii. (1867–68) 350.—Madag.
Amherstiana, Wall. ex DC. Mém. Anon. 28; Wall. Cat. n. 6424 = discolor.
Ancana, *F. Muell. Fragm.* v. 27.—Austral.
aromatica, Dun. Monog. Anon. 112 = concolor.
asiatica, Steud. Nom. ed. II. ii. 730 = pannosa.
biglandulosa, Blume, Bijdr. 16 = discolor.
Boivini, Baill. in Bull. Soc. Linn. Par. i. (1882) 339 = Popowia Boivini.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 11.—Borneo.
Brandisana, *Pierre, Fl. For. Cochinch.* t. 19.—Burma.
Cabog, Blanco, Fl. Filip. ed. I. 466 = U. Desmos.
caffra, E. Mey. ex C. Muell. in Walp. Ann. vii. 53 = Guatteria caffra.
camphorata, Blanco, Fl. Filip. ed. I. 468 = Uvaria dulcis.
Cananga, *Spreng. Syst.* iv. *Cur. Post.* 215.—Java.
Candolleana, Steud. Nom. ed. II. ii. 730 = U. Dasymaschala.
carminativa, Arruda, Diss. (1810) 48 = Xylopia sericea.
cauliflora, Blanco, Fl. Filip. ed. II. 323 = Melodorum fulgens.
cauliflora, Hook. f. & Thoms. Fl. Ind. i. 137 = Polyalthia cinnamomea.
cerasoides, Baill. Hist. Pl. i. 272 = Polyalthia cerasoides.
Chapelieri, Baill. in Bull. Soc. Linn. Par. i. (1882) 338 = Polyalthia Chapelieri.
chinensis, DC. Syst. i. 495 = discolor.
chloroxantha, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 17 (= *Monoon chloroxanthum*).—N. Guin.
clavata, Reinw. ex Blume, Fl. Jav. Anon. 66 = Xylopia javanica.
cochinchinensis, DC. Syst. i. 495 = U. Desmos.
coelophloea, *Scheff. in Flora*, lii. (1869) 300.—Malaya.
concinna, *Pierre, Fl. For. Cochinch.* t. 28.—Cochinch.
concolor, *Willd. Sp. Pl.* ii. 1271.—Guian.
cordifolia, Roxb. ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 59 (sphalm. pro Uvaria) = discolor.
Corinthi, Baill. Hist. Pl. i. 272 = Polyalthia Korinti.
corniculata, Blanco, Fl. Filip. ed. I. 469 = Artabotrys suaveolens.
corticosa, *Pierre, Fl. For. Cochinch.* t. 20.—Cochinch.
costata, *Miq. Fl. Ind. Bat. Suppl.* 376.—Sumatra.
crassipetala, *Dun. Monog. Anon.* 101. t. 24.—Guian.
crinita, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 61.—Malacca.
cubensis, *Steud. Nom.* ed. II. ii. 730.—Cuba.

UNONA:—
Dasymaschala, *Blume, Fl. Jav. Anon.* 55. t. 27. —Burma; Malaya.
debilis, *Pierre, Fl. For. Cochinch.* t. 29.—Cochinch.
dehiscens, Blanco, Fl. Filip. ed. I. 466 = Xylopia Blancoi.
desmantha, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 61.—Reg. Himal.; Malaya.
Desmos, *Raeusch. Nom.* ed. III. 161; *Dun. Monog. Anon.* 112.—Reg. Himal.; Malaya.
discolor, *Vahl, Symb. Bot.* ii. 63. t. 36.—As. trop.
discolor, Wall. Cat. n. 6420 D, E = U. Desmos.
discreta, *Linn. f. Suppl.* 270.—Guian.
diversifolia, *Span. in Linnaea*, xv. (1841) 163.—Ins. Timor.
dumetorum, *Dun. Monog. Anon.* 116.—Cochinch.
dumosa, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 670.—Ind. or.; Malaya.
Dunalii, Hook. f. & Thoms. Fl. Ind. i. 131, partim = discolor.
Dunalii, *Wall. Cat.* n. 6425.—Burma; Malaya.
elegans, *Thw. Enum. Pl. Zeyl.* 398.—Zeylan.
elliptica, Blume, Bijdr. 15 = Polyalthia elliptica.
erecta, *Pierre, Fl. For. Cochinch.* t. 31.—Cochinch.
esculenta, Dun. Monog. Anon. 107 = Artabotrys odoratissimus.
fasciculata, Steud. Nom. ed. II. ii. 730 = Polyalthia simiarum.
ferruginea, *Oliver, Fl. Trop. Afr.* i. 35.—Afr. trop.
Fitzgeraldii, F. Muell. Fragm. vi. 1 = Fitzgeraldia mitrastigma.
flagellaris, *Becc. in Nuov. Giorn. Bot. Ital.* iii. (1871) 186.—Borneo.
fluminensis, Vell. Fl. Flum. v. t. 121 = Xylopia sericea.
fulgens, Labill. Sert. Austro-Caled. 57. t. 56 = Polyalthia nitidissima.
fulva, Wall. Cat. n. 6427 = U. Desmos.
furfuracea, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 205 (*Mém. Anon.* 29).—Madag.
fuscata, *DC. Syst.* i. 488.—Guian.
Gerrardi, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 540.—Madag.
glauca, *Zipp. in Alg. Konst- en Letter-Bode*, i. (1829) 297, nomen.—Malaya.
glaucocarpa, Baill. in Bull. Soc. Linn. Par. i. (1882) 339 = Popowia glaucocarpa.
grandiflora, Leschen. ex DC. Prod. i. 90 = Uvaria purpurea.
Greveana, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 339.—Madag.
hamata, Blume, Bijdr. 15 = Artabotrys odoratissimus.
Hancei, *Pierre, Fl. For. Cochinch.* t. 32.—Cochinch.
Harmandii, *Pierre, l. c.* t. 24.—Cochinch.
heterocarpa, *Miq. Fl. Ind. Bat.* i. II. 41.—Java.
hirsuta, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 469. —Afr. trop.
humilis, Blume, Bijdr. 17 = Uvaria humilis.
jucunda, *Pierre, Fl. For. Cochinch.* t. 25.—Cochinch.
Kentii, Blume, Bijdr. 16 = Melodorum Kentii.
laevigata, *Wall. Cat.* n. 6428 (= Roxburghiana?).—Ind. or.
latifolia, *Dun. Monog. Anon.* 115.—Ins. Molucc.
latifolia, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. i. 60 = Brandisiana.
Lawii, *Hook. f. & Thoms. Fl. Ind.* i. 132.—Ind. or.
Leichhardtii, F. Muell. Fragm. iii. 41 = Melodorum Leichhardtii.
lepidota, *Oliver, Fl. Trop. Afr.* i. 36.—Afr. trop.
leptopetala, Dun. Monog. Anon. 114 = Cananga odorata.
Lessertiana, Dun. l. c. 107. t. 26 = discolor.
ligularis, *Dun. l. c.* 110.—Ins. Amboin.
littoralis, Blume, Bijdr. 16 = Uvaria littoralis.
longiflora, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 668.—Reg. Himal.
longiflora, Steud. Nom. ed. II. ii. 730, sphalm.—Cf. longifolia.
longifolia, Dun. Monog. Anon. 109 = Polyalthia longifolia.
longifolia, Steud. Nom. ed. II. ii. 730, sphalm. = Uvaria longifolia.
lucida, Boj. Hort. Maurit. 6 = Uvaria lucida.
lucida, DC. Syst. i. 498 = Xylopia longifolia.
lucidula, *Oliver, Fl. Trop. Afr.* i. 35.—Afr. trop.
luensis, *Pierre, Fl. For. Cochinch.* t. 30.—Cochinch.
macrocarpa, Baill. in Bull. Soc. Linn. Par. i. (1882) 339 = Popowia macrocarpa.

UNONA:—
macrocarpa, Vahl, ex DC. Syst. i. 489 = Uvaria Chamae.
macrophylla, Blume, Bijdr. 17 = Goniothalamus macrophyllus.
Marenteria, DC. Syst. i. 487 = Uvaria Marenteria.
maritima, Baill. in Bull. Soc. Linn. Par. i. (1882) 339 = Polyalthia maritima.
Mesnyi, *Pierre, Fl. For. Cochinch.* t. 17.—Malacca; Siam.
micrantha, *Miq. Fl. Ind. Bat.* i. II. 43.—Java.
modesta, *Pierre, Fl. For. Cochinch.* t. 28.—Cochinch.
mollissima, Zipp. ex Span. in Linnaea, xv. (1841) 162 = Uvaria timorensis.
monilifera, DC. Prod. i. 91, sphalm. = discreta.
moniliformis, Steud. Nom. ed. II. ii. 731 (sphalm. pro Uvaria monilifera) = discolor.
monosperma, Pierre, Fl. For. Cochinch. sub t. 18 = Cananga monosperma.
Musaria, *Dun. Monog. Anon.* 100.—Ins. Molucc.
Narum, Dun. l. c. 99 = Uvaria Narum.
nitens, F. Muell. Fragm. iii. 2 = Polyalthia nitidissima.
nitidissima, Dun. Monog. Anon. 109. t. 23 = Polyalthia nitidissima.
obovata, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 469.—Afr. trop.
obtusiflora, *DC. ex Dun. Monog. Anon.* 102.—Guian.
odorata, Blanco, Fl. Filip. ed. II. 325 = U. Desmos.
odorata, Dun. Monog. Anon. 108 = Cananga odorata.
odoratissima, Steud. Nom. ed. II. ii. 730 (sphalm. pro Uvaria) = Artabotrys odoratissima.
Oliveriana, Baill. Adansonia, viii. (1867–68) 307.—Afr. trop.
ossea, Blanco, Fl. Filip. ed. I. 467 = discolor.
ovata, Vahl, ex DC. Syst. i. 489 = Uvaria ovata.
oxypetala, DC. l. c. 496 = Xylopia acutiflora?
pachypetala, Spreng. Syst. ii. 637 = crassipetala.
pannosa, *Dalz. in Hook. Kew Journ.* iii. (1851) 207.—Ind. or.
parviflora, Steud. Nom. ed. II. ii. 730 = Xylopia Dunaliana.
parvifolia, *Oliver, Fl. Trop. Afr.* i. 36.—Afr. trop.
pedunculosa, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 204 (Mém. Anon. 28) = U. Desmos.
penduliflora, Dun. Monog. Anon. 100. t. 28 = Cymbopetalum penduliflorum.
pilosa, Boiv. ex Baill. Adansonia, viii. (1867–68) 322 = Popowia pilosa.
polycarpa, DC. Syst. i. 499 = Xylopia polycarpa.
praecox, *Hook. f. & Thoms. Fl. Ind.* i. 136.—Reg. Himal.
pycnantha, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 60 (Polyalthiae sp.).—Malacca.
pygmaea, Walp. Rep. i. 79, sphalm. = Asimina pygmaea.
Richardiana, Baill. in Bull. Soc. Linn. Par. i. (1882) 339 = Polyalthia Richardiana.
rigida, *Zipp. ex Span. in Linnaea*, xv. (1841) 163, nomen.—Ins. Timor.
Roxburghiana, Wall. Cat. sub n. 6428 A = Oxymitra fornicata.
Roxburghiana, Wall. l. c. B = discolor.
selanica, *DC. Prod.* i. 92.—Ins. Molucc.
sempervivens, Baill. Hist. Pl. i. 272 (Polyalthiae sp.). —As. trop.
setigera, Blanco, Fl. Filip. ed. II. 323 = Uvaria purpurea.
simiarum, Baill. ex Pierre, Fl. For. Cochinch. t. 23 = Polyalthia simiarum.
stenopetala, *Hook. f. & Thoms. Fl. Ind.* i. 136.—Singapore.
suaveolens, Blume, Bijdr. 17 = Artabotrys suaveolens.
subbiglandulosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 11.—Borneo.
subcordata, Blume, Bijdr. 15 = Polyalthia subcordata.
subglandulosa, Walp. Ann. vii. 54, sphalm. = subbiglandulosa.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 377.—Sumatra.
sylvatica, *Dun. Monog. Anon.* 115.—Cochinch.
Thorelii, *Pierre, Fl. For. Cochinch.* t. 22.—Cochinch.
tomentosa, Willd. Sp. Pl. ii. 1271 = U. Desmos.
tripelata, Blanco, Fl. Filip. ed. II. 324, sphalm. = seq.
tripetala, DC. Prod. i. 90 = seq.
tripetaloidea, *Dun. Monog. Anon.* 104.—Java.
tristis, *Pierre, Fl. For. Cochinch.* t. 27.—Cochinch.
uncinata, Dun. Monog. Anon. 105. t. 12 = Artabotrys odoratissimus.
undulata, *Dun. l. c.* 111.—Afr. trop.

UNONA:—
undulata, Wall. Pl. As. Rar. iii. 42. t. 265 = discolor.
uniflora, Baill. Adansonia, viii. (1868) 327, sphalm. = Duguetia Candollei.
velutina, Blume, ex Hook. f. & Thoms. Fl. Ind. i. 130, sphalm. = Cananga odorata.
velutina, *Hance, in Journ. Bot.* xv. (1877) 328.—Cambodia.
verrucosa, *Becc. in Nuov. Giorn. Bot. Ital.* iii. (1871) 185.—Borneo.
violacea, *Dun. Monog. Anon.* 105. t. 25.—Mexic.
virgata, Blume, Bijdr. 14 = Cananga virgata.
viridiflora, *Bedd. Ic. Pl. Ind. Or.* i. 34.—Ind. or.
viridiflora, *Splitg. in Nederl. Kruidk. Arch.* i. (1848) 224.—Guian.
xylopioides, *Dun. Monog. Anon.* 117. t. 21.—Venezuela.
zeylanica, *Hook. f. & Thoms. Fl. Ind.* i. 132.—Zeylan.

UNXIA, Bert. ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 37. t. 32 = **Blennosperma**, Less. (Compos.).
anthemidifolia, Phil. f. Cat. Pl. Vasc. Chil. 156, 187, sphalm. = seq.
anthemifolia, Bert. ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 38 = B. chilense.

UNXIA, H. B. & K. Nov. Gen. et Sp. iv. 279. tt. 401, 402 (1820) = **Villanova**, Lag. (Compos.).
achillaeoides, Less. in Linnaea, v. (1830) 159 = V. achillaeoides.
alternifolia, Spreng. Syst. iii. 620 = V. alternifolia.
anemonifolia, H. B. & K. Nov. Gen. et Sp. iv. 279. t. 402 = V. oppositifolia.
dissecta, Hook. Bot. Misc. ii. (1831) 227 = V. dissecta.
pratensis, H. B. & K. Nov. Gen. et Sp. iv. 279. t. 401 = V. oppositifolia.

UNXIA, Linn. f. Suppl. 56 (1781) = **Melampodium**, Linn. (Compos.).
achillaeoides, Hemsl. Biol. Centr. Am. Bot. ii. 145 = M. achillaeoides.
camphorata, Linn. f. Suppl. 368 = M. camphoratum.
digyna, Steetz, in Seem. Bot. Voy. Herald, 154. t. 30 = M. digynum.
hirsuta, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 112 (err. typ. 105) = M. camphoratum.

UPATA, Adans. Fam. ii. 201 (1763) = **Avicennia**, Linn. (Verben.).

UPOPION, Rafin. New Fl. Am. iv. 29 (1836) = **Thaspium**, Nutt. (Umbell.).
cordatum, Rafin. l. c. 31 = T. trifoliatum.
heterophylum, Rafin. l. c. = T. trifoliatum.
lobatum, Rafin. l. c. 30.—Am. bor.
pinnatum, Rafin. l. c. = T. trifoliatum.
trifoliatum, Rafin. l. c. = T. trifoliatum.

UPUDALIA, Rafin. Fl. Tellur. iv. 66 (1836) = **Daedalacanthus**, T. Anders. (1864) (Acanth.).
pulchella, Rafin. l. c. = D. nervosus.

URACHNE, Trin. Fund. Agrost. 109 (1820) = **Oryzopsis**, Michx. (Gramin.).
acutigluma, Steud. Syn. Pl. Gram. 121.—Japon.?
aemulans, Steud. in Lechl. Berb. Am. Austr. 52.—Chili.
asperifolia, Trin. Gram. Unifl. 174 = O. asperifolia.
brevicaudata, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. iii. (1835) 127 = O. parviflora.
canadensis, Torr. & Gray, ex Trin. & Rupr. l. c. v. (1842) 17 = O. parviflora.
chilensis, Trin. l. c. iii. (1835) 123 (= *Nassella chilensis*). —Chili.
coerulescens, Trin. Fund. Agrost. 110 = O. coerulescens.
comosa, Trin. Gram. Unifl. 175 (= *Piptatherum miliaceum*).—Europ. austr.
depressa, Steud. Syn. Pl. Gram. 123.—Chili.
Fieldingii, C. Muell. in Bot. Zeit. xix. (1861) 342.—Hab.?
frutescens, Link, Hort. Berol. i. 93 (= *Piptatherum miliaceum*).—Europ. austr.
fusca, Steud. Syn. Pl. Gram. 123.—Chili.
gigantea, Steud. l. c.—Chili.
grandiflora, Steud. & Hochst. ex Steud. l. c. 124 = Stipa megalantha.
grandiflora, Trin. Gram. Unifl. 174 = O. holciformis.
Haenkeana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 20.—Peruv.

URACHNE:—
holciformis, C. Koch, in Linnaea, xxi. (1848) 439 = Oryzopsis holciformis.
laevis, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 20.—Peruv.
lanata, Trin. l. c. iii. (1835) 126 = O. cuspidata.
leucosperma, Link, Hort. Berol. i. 94 = O. asperifolia.
Linnaei, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 13 = O. paradoxa.
longiflora, Steud. Nom. ed. II. ii. 731.—Chili.
macrathera, Steud. Syn. Pl. Gram. 123 = *trichotoma*.
major, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 21.—Chili?
megalantha, Steud. Nom. ed. II. ii. 731 = Stipa megalantha.
melanosperma, Link, Hort. Berol. i. 94 = O. melanocarpa.
Meyeniana, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 20.—Peruv.
micrantha, Trin. & Rupr. l. c. 16 = O. micrantha.
molinioides, Steud. Syn. Pl. Gram. 122 = O. pubiflora.
multiflora, Link, Hort. Berol. i. 93 (= *Piptatherum miliaceum*).—Europ.
mutica, Steud. Nom. ed. II. ii. 731 = O. asperifolia.
obtusa, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 22.—Peruv.
panicoides, Trin. l. c. iii. (1835) 124.—Chili; Bras.
paradoxa, Link, Hort. Berol. i. 92 = O. paradoxa.
parviflora, Trin. Fund. Agrost. 110 (= *Piptatherum miliaceum*).—Europ.
pauciflora, Boiss. Fl. Orient. v. 507, sphalm. = praec.
pubiflora, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 21.—Peruv.
purpurea, Steud. Syn. Pl. Gram. 122 = O. coerulescens.
racemosa, Trin. Gram. Unifl. 174 = O. melanocarpa?
ramosa, Steud. Nom. ed. II. ii. 731; Syn. Pl. Gram. 123 = *chilensis*.
ramosissima, Trin. Gram. Unifl. 173 = Stipa micrantha.
setacea, Walp. Ann. iii. 730, sphalm. = seq.
setosa, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. iii. (1835) 124 = O. setacea.
simplex, Trin. & Rupr. l. c. v. (1842) 22.—Chili.
sinaica, Steud. Syn. Pl. Gram. 122 = O. holciformis.
songarica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 15 = O. holciformis.
sphacelata, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 230 (= *Piptatherum sphacelatum*).—Persia.
stipoides, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. (1842) 25 (= *Piptochaetium bicolor*).—Bras.
Thomasii, Steud. Syn. Pl. Gram. 122 (= *Piptatherum miliaceum*).—Europ.
trichotoma, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. iii. (1835) 124.—Bras.
virescens, Trin. Fund. Agrost. 110 (= *Piptatherum virescens*).—Europ. or.; As. Min.

URAGOGA, Linn. Gen. ed. I. 378, et Index (1737) = **Psychotria**, Linn. (Rubiac.).
arbutifolia, Baill. Adansonia, xii. (1879) 292.—N. Caled.
asthmatica, Baill. Hist. Pl. vii. 376 = Cephaëlis asthmatica (in Addend.).
baladensis, Baill. Adansonia, xii. (1879) 287.—N. Caled.
Balansae, Baill. l. c. 223.—N. Caled.
brachylaena, Baill. l. c. 250.—N. Caled.
calliantha, Baill. l. c. 229.—N. Caled.
calorhamnus, Baill. l. c. 286.—N. Caled.
Calothyrsus, Baill. l. c. 249.—N. Caled.
canalensis, Baill. l. c. 254.—N. Caled.
cardiochlamys, Baill. l. c. 287.—N. Caled.
coptosperma, Baill. l. c. 289.—N. Caled.
costularia, Baill. l. c. 248.—Venezuela.
emetica, Baill. Hist. Pl. vii. 371 = P. emetica.
Fagueti, Baill. Adansonia, xii. (1879) 224.—N. Caled.
furcellata, Baill. ex Vatke, in Bremen, Abh. ix. (1885) 118 = P. furcellata.
Gabriellae, Baill. Adansonia, xii. (1879) 228.—N. Caled.
goniocarpa, Baill. l. c. 285.—N. Caled.
granatensis, Baill. Hist. Pl. vii. 370 = P. Ipecacuanha?
Grosourdieana, Baill. Adansonia, xii. (1879) 227 = P. Grosourdyeana (in Addend.).
guianensis, Baill. Hist. Pl. vii. 376 = Cephaëlis guianensis (in Addend.).

URAGOGA:—
Guilloveana, Baill. Adansonia, xii. (1879) 329.—N. Guin.
Hombroniana, Baill. l. c. 333.—Ins. Marian.
Ipecacuanha, Baill. Hist. Pl. vii. 281 = Psychotria Ipecacuanha.
jasminiflora, Baill. Adansonia, xii. (1879) 226.—N. Caled.
ligustrina, Baill. l. c. 253.—N. Caled.
lyciiflora, Baill. l. c. 290.—N. Caled.
lycioides, Baill. l. c. 228.—N. Caled.
macroglossa, Baill. l. c. 251.—N. Caled.
Marcgravii, Baill. Hist. Pl. vii. 372 = P. Marcgravii.
macropoda, Baill. l. c. 376 = Geophila macropoda.
microglossa, Baill. Adansonia, xii. (1879) 252.—N. Caled.
Micromyrtus, Baill. l. c. 291.—N. Caled.
monanthos, Baill. l. c. 293 = N. Caled.
muscosa, Baill. Hist. Pl. vii. 376 = Cephaëlis muscosa.
Nathaliae, Baill. Adansonia, xii. (1879) 248.—N. Caled.
Nekouana, Baill. l. c. 287.—N. Caled.
noxia, Baill. Hist. Pl. vii. 372 = P. noxia.
officinalis, Baill. l. c. 376 = P. officinalis.
oleoides, Baill. Adansonia, xii. (1879) 252.—N. Caled.
Pancheri, Baill. l. c. 222.—N. Caled.
Paramaracarpus, Baill. l. c. 293.—N. Caled.
parasitica, Baill. l. c. 328 = P. parasitica.
Poissoniana, Baill. l. c. 230.—N. Caled.
Remyana, Baill. l. c. 253.—Ecuador.
reniformis, Baill. Hist. Pl. vii. 376 = Geophila reniformis.
rosmarinifolia, Baill. Adansonia, xii. (1879) 291.—N. Caled.
ruelliaefolia, Baill. Hist. Pl. vii. 372 = P. ruelliaefolia.
rupicola, Baill. Adansonia, xii. (1879) 288.—N. Caled.
sagittalis, Baill. l. c. 249.—N. Caled.
Simira, Baill. Hist. Pl. vii. 374 = P. parviflora.
Spachiana, Baill. Adansonia, xii. (1879) 285.—N. Caled.
stricta, Baill. l. c. 250.—N. Caled.
subuniflora, Baill. l. c. 292.—N. Caled.
tinctoria, Baill. Hist. Pl. vii. 374 = Palicourea tinctoria.
tortilis, Baill. Adansonia, xii. (1879) 325 = P. tortilis.
trichopodantha, Baill. l. c. 294.—N. Caled.
trisulcata, Baill. l. c. 289.—N. Caled.
undata, Baill. Hist. Pl. vii. 371 = P. undata.
viburnifolia, Baill. Adansonia, xii. (1879) 330.—Mexic.
Vieillardi, Baill. l. c. 224.—N. Caled.

URALEPIS, Spreng. Neue Entdeck. i. 244 (1821) = seq.

URALEPSIS, Nutt. Gen. Am. i. 62 (1818) = **Triodia**, R. Br. (Gramin.).
alba, Steud. Syn. Pl. Gram. 248 = Diplachne alba.
alopecuroides, Steud. l. c. = T. plumosa.
ambigua, Kunth, Rév. Gram. i. 108 = T. ambigua.
arenaria, Steud. Syn. Pl. Gram. 248 = Diplachne arenaria.
aristulata, Nutt. Gen. Am. i. 63 = Triplasis purpurea.
avenacea, Kunth, Rév. Gram. i. 108 = T. avenacea.
brasiliensis, Steud. Syn. Pl. Gram. 248.—Bras.
brevicuspidata, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 93 = Diplachne dubia.
capensis, Kunth, Enum. Pl. i. 319.—Afr. austr.
capensis, Steud. Syn. Pl. Gram. 248.—Afr. austr.
ciliata, Steud. l. c. 247.—Afr. trop.
composita, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 94 = Diplachne fascicularis.
cornuta, Ell. Sketch, i. 580 = Triplasis americana.
cuprea, Kunth, Rév. Gram. i. 108 = T. cuprea.
densiflora, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 94 = T. stricta.
Drummondii, Steud. Syn. Pl. Gram. 247 = Diplachne fusca.
Eleusine, Steud. l. c. 248 = Diplachne Eleusine.
elongata, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 89 = T. trinerviglumis.
flaccida, Doell, in Mart. Fl. Bras. ii. III. 95.—Bras.
fusca, Steud. Syn. Pl. Gram. 247 = Diplachne fusca.
livida, Steud. l. c. 248 = Diplachne livida.
mutica, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 569 = T. mutica.
pallida, Kunth, Rév. Gram. i. 108 = Glyceria pallida.
pilosa, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 94 = T. acuminata.
pilosa, Buckl. l. c. 95 = T. mutica.
poaeoides, Buckl. l. c. = Sclerochloa californica.

URALEPSIS:—
pulchella, Kunth, Rév. Gram. i. 108 = Triodia pulchella.
pungens, Kunth, l. c. = Aeluropus arabicus.
purpurea, Nutt. Gen. Am. i. 62 = Triplasis purpurea.
quadridentata, Doell, in Mart. Fl. Bras. ii. III. 240.—Bras.
Tricuspis, Steud. Nom. ed. II. i. 564, in syn. = T. cuprea.
verticillata, Steud. Syn. Pl. Gram. 248 = Diplachne verticillata.
virens, Kunth, Enum. Pl. i. 319.—Bras.

URANANTHUS, Benth. Pl. Hartw. 46 (1840) = **Eustoma**, Salisb. (Gentian.).
chironioides, Benth. l. c. 46 = E. silenifolium.
glaucifolius, Benth. l. c. = E. silenifolium.
pallidus, Liebm. Ind. Sem. Hort. Hafn. (1845); ex Sond. in E. Otto, Allg. Gartenz. iv. (1848) 368 = E. silenifolium.
Russelianus, Benth. Pl. Hartw. 46 = E. Russelianum.

URANDRA, Thw. in Hook. Kew Journ. vii. (1855) 211 = **Lasianthera**, Beauv. (Olacin.).
apicalis, Thw. l. c. = L. apicalis.

URANIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 100; et Desv. in Mém. Soc. Linn. Par. iv. (1826) 309, sphalm. = **Uraria**, Desv. (Legumin.).

URANIA, Schreb. Gen. 212 (1789) = **Ravenala**, Adans. (Scitam.).
amazonica, Mart. Reise Bras. iii. 946; Hort. ex Lem. Illustr. Hortic. vii. (1860) t. 239 = R. guyanensis.
guyanensis, Rich. in Nov. Act. Nat. Cur. xv. (1831) Suppl. 21 = R. guyanensis.
madagascariensis, Raeusch. Nom. ed. III. 91 = R. madagascariensis.
Ravenalia, Rich. in Nov. Act. Nat. Cur. xv. (1831) Suppl. 19 = R. madagascariensis.
speciosa, Willd. Sp. Pl. ii. 7 = R. madagascariensis.

URANOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
prostrata, Fourr. l. c. 128 = V. prostrata.
Teucrium, Fourr. l. c. = V. Teucrium.

URANTHERA, Naud. in Ann. Sc. Nat. Sér. III. iii. (1845) 189 = **Acisanthera**, P. Br. (Melastom.).
alsinaefolia, Naud. l. c. = A. alsinaefolia.
dicranophora, Naud. l. c. xii. (1849) 283 = A. quadrata.
Hostmanni, Naud. l. c. 284 = A. recurva.
recurva, Naud. l. c. iii. (1845) 189 = A. recurva.
variabilis, Naud. l. c. 190. t. 7. f. 18 = A. variabilis.

URARIA, Desv. Journ. Bot. i. (1813) 122. t. 5. f. 19. *LEGUMINOSAE*, Benth. & Hook. f. i. 521.
DOODIA, Roxb. Hort. Beng. 99 (1814); Fl. Ind. iii. 365 (1832).
acuminata, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 235, 236 = picta?
alopecuroides, Sweet, Hort. Brit. ed. II. 149; Wight, Ic. t. 290 = U. Lagopus.
arborea, G. Don, Gen. Syst. ii. 287 = U. Lagopus.
barbata, Desv. in Mém. Soc. Linn. Par. iv. (1826) 309 = Desmodium barbatum.
campanulata, Wall. Cat. n. 5685 = Lourea campanulata.
cercidifolia, Desv. in Mém. Soc. Linn. Par. iv. (1826) 309, sphalm. = seq.
cercifolia, Desv. Journ. Bot. i. (1813) 123. t. 5. f. 19 = lagopoides.
comosa, DC. Prod. ii. 324 = crinita.
cordata, Wall. Cat. n. 5679 = seq.
cordifolia, *Wall. Pl. As. Rar.* i. 33.—Burma.
cornosa, Span. in Linnaea, xv. (1841) 193 = crinita.
crinita, *Desv. Journ. Bot.* i. (1813) 123.—As. trop.
cylindracea, *Benth. Fl. Austral.* ii. 237.—Austral.
desmodoides, R. Grah. in Wall. Cat. n. 5683 = hamosa.
hamosa, Wall. Cat. n. 5681 A, C = U. Lagopus, lagopoides.
hamosa, *Wall. l. c.* B.—Ind. or.; Malaya.
lagocephala, DC. Prod. ii. 324 = Desmodium barbatum.
lagopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = U. Lagopus.
lagopodioides, Desv. in Mém. Soc. Linn. Par. iv. (1826) 309 = lagopoides?
lagopodioides, Wall. Cat. n. 5676 = U. Lagopus.
lagopoides, *DC. Prod.* ii. 324.—As. et Austral. trop.
Lagopus, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 100.—Reg. Himal.; Burma.

URARIA:—
lanceolata, R. Grah. in Wall. Cat. n. 5682 = hamosa.
leptostachya, Wall. l. c. 5684 = hamosa.
leucantha, Zipp. ex Span. in Linnaea, xv. (1841) 193 = picta.
linearis, Hassk. in Flora, xxv. (1842) II. Beibl. 61; Cat. Hort. Bog. Alt. 273 = picta.
macrostachya, Wall. Pl. As. Rar. ii. t. 110 = crinita.
obcordata, Miq. Fl. Ind. Bat. Suppl. 305 = Desmodium obcordatum.
paniculata, *Hassk. Cat. Hort. Bog. Alt.* 273.—Java.
picta, *Desv. Journ. Bot.* i. (1813) 123. t. 5. f. 19.—Geront. trop.
picta, Wight, Ic. t. 411 = crinita.
prunellaefolia, *R. Grah. in Wall. Cat.* n. 5686.—Reg. Himal.
repanda, Wall. Cat. n. 5677.—Burma.
retrofracta, Wall. l. c. 5678 = Mecopus nidulans.
retusa, Wall. l. c. 5680 = lagopoides.
styracifolia, Wight & Arn. Prod. 222 = Desmodium retroflexum.

URASPERMUM, Nutt. Gen. Am. i. 192 (1818) = **Osmorhiza**, Rafin. (Umbell.).
Claytoni, Nutt. l. c. 193 = O. longistylis.
hirsutum, Bigel. Fl. Bost. ed. III. 120 = O. brevistylis.

URATEA, J. F. Gmel. Syst. 677 (1791) = OURATEA, Aubl. = **Gomphia**, Schreb. (Ochnac.).

URBANIA, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 10 = **Lyperia**, Benth. (Scrophul.).
lyperiaeflora, Vatke, in Linnaea, xliii. (1880–82) 306 = Lindenbergia sinaica.
lyperiaeflora, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 10.—Afr. trop.

URCEOLA, Roxb. in As. Res. v. (1798) 169. *APOCYNACEAE*, Benth. & Hook. f. ii. 716.
CHAVANNESIA, A. DC. in DC. Prod. viii. 444 (1844).
brachysepala, *Hook. f. Fl. Brit. Ind.* iii. 659.—Malacca.
elastica, *Roxb. in As. Res.* v. (1798) 169.—Malaya.
esculenta, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 658.—Burma.
lucida, *Benth. & Hook. f. Gen.* ii. 716.—Burma; Ins. Penang.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 658.—Singapore.
malaccensis, *Hook. f. l. c.*—Malacca.
torulosa, *Hook. f. l. c.* 659.—Malacca.

URCEOLA, Vand. Fl. Lusit. 8. t. 1. f. 4 (1788) (Quid?).
Vandelli, Roem. & Schult. Syst. iii. 99.—Bras.

URCEOLARIA *capensis*, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 396, sphalm. = Utricularia capensis.

URCEOLARIA, Herb. App. [Bot. Reg.] 28 (1821) = **Urceolina**, Reichb. (Amaryll.).
pendula, Herb. l. c. = Urceolina pendula.

URCEOLARIA, Molina, Sagg. Chile, ed. II. 136, 277 (1810) = **Sarmienta**, Ruiz & Pav. (Gesner.).
chilensis, Molina, ll. cc. = S. repens.

URCEOLARIA, Willd. ex Cothen. Disp. Veg. Meth. 10 (1790) = **Schradera**, Vahl (Rubiac.).
exotica, J. F. Gmel. Syst. 390 = S. capitata.

URCEOLINA, Reichb. Consp. 61 (1828). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 732.
COLLANIA, Schult. f. Syst. vii. II. p. liii, 893 (1830).
LEPERIZA, Herb. App. [Bot. Reg.] 41 (1821).
PENTLANDIA, Herb. in Bot. Reg. (1839) t. 68.
SPHAEROTELE, Link, Klotzsch & Otto, Ic. Pl. Rar. Hort. Berol. 95. t. 38 (1841).
URCEOLARIA, Herb. App. [Bot. Reg.] 28 (1821).
aurea, Lindl. in Gard. Chron. (1864) 627, 890 = pendula.
fulva, Herb. Amaryll. 194 = pendula.
latifolia, *Benth. & Hook. f. Gen.* iii. 732.—Peruv.
lobata, Pritz. Ic. Ind. 1134, sphalm. = Urena lobata.
miniata, *Benth. & Hook. f. Gen.* iii. 732.—Peruv.
pendula, *Herb. Amaryll.* 193.—Peruv.

URECHITES, Muell. Arg. in Bot. Zeit. xviii. (1860) 22. *APOCYNACEAE*, Benth. & Hook. f. ii. 727.
Andrieuxii, *Muell. Arg. in Linnaea*, xxx. (1859–60) 442.—Mexic.
Jaegeri, *Muell. Arg. l. c.* 443.—Ins. S. Doming.
jamaicensis, Miers, Apocyn. S. Am. 112 = Echites jamaicensis.
Karwinskii, *Muell. Arg. in Linnaea*, xxx. (1859–60) 440.—Mexic.
Neriandra, Miers, Apocyn. S. Am. 111 = Echites Andrewsii.
suberecta, *Muell. Arg. in Linnaea*, xxx. (1859–60) 444.—Ins. S. Doming.

URENA, Dill. ex Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 205.
aculeata, Mill. Gard. Dict. ed. VIII. n. 2 = lobata.
americana, Linn. Pl. Surin. 11; Amoen. Acad. viii. 259; Linn. f. Suppl. 308 = lobata.
Armitiana, *F. Muell. Fragm.* x. 78.—Austral.
Blumei, *Hassk. in Hoev. & De Vriese, Tijdschr.* xii. (1845) 78.—Java.
campestris, *Vell. Fl. Flum.* 286; vii. t. 47.—Bras.
cana, Wall. Cat. n. 1930 B = lobata.
chinensis, Osbeck, Voy. ed. Angl. i. 363 (Quid?).—China.
cordata, Herb. Heyne, ex Mast. in Hook. f. Fl. Brit. Ind. i. 330 = Pavonia glechomifolia.
diversifolia, Schum. & Thonn. Beskr. Guin. Pl. 308 = lobata.
foetida, L'Hérit. ex DC. Prod. i. 444 = Pavonia paniculata.
glabra, R. Br. in Salt, Abyss. App. 65 = Pavonia glechomifolia.
grandiflora, [Moç. & Sesse,] ex DC. Prod. i. 442 = lobata.
Haenkeana, Walp. Rep. i. 297 = lobata.
Hamiltoniana, Wall. Cat. n. 2700 = repanda.
hastata, *Vell. Fl. Flum.* 285; vii. t. 43.—Bras.
heteromorpha, *Montr. in Mém. Acad. Lyon*, x. (1860) 180.—N. Caled.
heterophylla, Presl, Rel. Haenk. ii. 127 = lobata.
heterophylla, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 111 (Quid?).—Guian.
heterophylla, Schrad. Cat. Sem. Hort. Gotting. (1835). —Cf. Linnaea, xi. (1837) Litt. 90 = phyllomorpha.
heterophylla, Sm. in Rees, Cycl. xxxvii. n. 86 = sinuata.
hirta, *Vell. Fl. Flum.* 286; vii. t. 45.—Bras.
hispida, Boj. ex Baker, in Journ. Bot. xx. (1882) 45 = Pavonia Bojeri.
Lappago, Sm. in Rees, Cycl. xxxvii. n. 11; DC. Prod. i. 441 = sinuata.
leptocarpa, Linn. f. Suppl. 508 = Pavonia Typhalaea.
leucantha, Poepp. ex Garcke, in Eichl. Jahrb. Berl. i. (1881) 211 = Pavonia leucantha.
lobata, *Linn. Sp. Pl.* 692.—Cosmop. trop.
Loureirii, Meissn. ex Steud. Nom. ed. II. ii. 732 = Malachra Urena.
malacoides, Hill, Veg. Syst. xv. 20 (Quid?).—China.
Manihot, Klotzsch, in Bot. Zeit. iv. (1846) 102, nomen = sinuata.
mauritiana, Sieber, ex Colla, Hort. Ripul. 142 = lobata.
mediterranea, *Vell. Fl. Flum.* 285; vii. t. 42.—Bras.
Meyeri, F. Muell. Fragm. x. 79 = Pavonia Columella.
microcarpa, DC. Prod. i. 442 = lobata.
mollis, R. Br. in Salt, Abyss. App. 65 = Pavonia Kraussiana.
monopetala, *Lour. Fl. Cochinch.* 418.—Cochinch.
morifolia, DC. Prod. i. 442 = sinuata.
multifida, Cav. Diss. vi. 336. t. 184. f. 2 = sinuata.
muricata, DC. Prod. i. 442 = sinuata.
obtusata, Guill. & Perr. Fl. Seneg. Tent. i. 48 = lobata.
ovalifolia, Forsk. Fl. Aegypt. Arab. 124 = Hibiscus micranthus.
palmata, Roxb. Hort. Beng. [97]; Fl. Ind. iii. 182 = lobata.
paradoxa, H. B. & K. Nov. Gen. et Sp. v. 278 = sinuata.
phyllomorpha, *Steud. Nom.* ed. II. ii. 732.—Java.
pilosa, Burch. ex DC. Prod. i. 438 = Pavonia Columella.
polyflora, Lour. Fl. Cochinch. 417 = Malachra Urena.
praemorsa, Salisb. Prod. 382 = Pavonia praemorsa.
procumbens, *Linn. Sp. Pl.* 692.—China.
repanda, Blume, Bijdr. 64 = Blumei.

URENA:—
repanda, *Roxb. Hort. Beng.* 51; *Fl. Ind.* iii. 182.—Ind. or.; Burma.
reticulata, Cav. Diss. vi. 335. t. 183. f. 2 = lobata.
Ribesia, Sm. in Rees, Cycl. xxxvii. n. 5 = lobata.
ricinocarpa, *Eckl. & Zeyh. Enum.* 37.—Afr. austr.
rigida, *Wall. Cat.* n. 1929, partim.—Burma; Malaya.
rigida, Wall. l. c., partim = repanda.
scabriuscula, DC. Prod. i. 441 = lobata.
sidoides, *Vell. Fl. Flum.* 286; vii. t. 46.—Bras.
Sieberi, Colla, Hort. Ripul. 142. t. 30 = lobata.
sinuata, *Linn. Sp. Pl.* 692.—Reg. trop.
sinuata, Sw. Obs. Bot. 264 = lobata.
speciosa, Wall. Pl. As. Rar. i. 23. t. 26 = repanda.
stellata, Spreng. Neue Entdeck. ii. 163 = Pavonia Typhalaea.
stellipila, *Lem. Jard. Fleur.* iii. (1852) t. 281.—Bras.
subtriloba, *Schrank, Pl. Rar. Hort. Monac.* t. 79.—Bras.
Swartzii, DC. Prod. i. 442 = lobata.
Swartzii, Macfad. Fl. Jamaic. i. 60 = sinuata.
tomentosa, *Blume, Bijdr.* 65.—Java.
tomentosa, Wall. Cat. n. 1933 H = sinuata.
tricuspis, Cav. Diss. vi. 334. t. 183. f. 1 = lobata.
trilobata, Vell. Fl. Flum. vii. 269. t. 44 = lobata.
Typhalaea, Linn. Mant. ii. 258 = Pavonia Typhalaea.
viminea, Cav. Diss. vi. 335. t. 184. f. 1 = lobata.
virgata, Guill. & Perr. Fl. Seneg. Tent. i. 48 = lobata.
Xanthium, *Vell. Fl. Flum.* 284; vii. t. 41 (sp. dub.).—Bras.

URERA, Gaudich. in Freyc. Voy. Bot. 496 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 383.
CALOSTIMA, Rafin. Fl. Tellur. iii. 47 (1836).
SCEPOCARPUS, Wedd. in DC. Prod. xvi. I. 98 (1869).
acuminata, *Gaudich. in Freyc. Voy. Bot.* 497.—Ins. Maurit. et Timor.
acuminata, Miq. in Mart. Fl. Bras. iv. I. 190 = alceaefolia.
alceaefolia, *Gaudich. in Freyc. Voy. Bot.* 497.—Am. trop.
amberana, *Baker, in Journ. Bot.* xx. (1882) 267.—Madag.
amplissima, Blume, Mus. Bot. Lugd. Bat. ii. t. 22 = Laportea amplissima.
armigera, Miq. in Mart. Fl. Bras. iv. I. 193 = baccifera.
aurantiaca, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 201.—Bras.
baccifera, *Gaudich. in Freyc. Voy. Bot.* 497.—Am. austr.
cameroonensis, *Wedd. in DC. Prod.* xvi. I. 97.—Afr. trop.
capitata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 201.—Peruv.
caracasana, Griseb. Fl. Brit. W. Ind. 154 = alceaefolia.
Commersoniana, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 202 = Laportea crenulata.
corallina, *Wedd. in DC. Prod.* xvi. I. 90.—Costa Rica.
crassifolia, *Wedd. in Arch. Mus. Par.* ix. (1856) 161. —Ins. Dominic.
crenulata, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 203 = Laportea crenulata.
densiflora, Miq. in Mart. Fl. Bras. iv. I. 190 = alceaefolia.
denticulata, Miq. l. c. 192 = baccifera.
elata, Griseb. Fl. Brit. W. Ind. 154 = sinuata.
excelsa, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 178 = Laportea Gigas.
expansa, *Griseb. Fl. Brit. W. Ind.* 155.—Jamaic.
ficifolia, Gaudich. in Freyc. Voy. Bot. 497 = Obetia ficifolia.
Gaudichaudiana, Hensl. in Ann. Nat. Hist. Ser. I. i. (1838) 341 = Fleurya aestuans.
Gaudichaudiana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 177 = Laportea Gaudichaudiana.
gigantea, Gaudich. in Freyc. Voy. Bot. 497 = Laportea crenulata.
girardinioides, Seem. Bot. Voy. Herald, 194 = laciniata.
glabra, Wedd. in Arch. Mus. Par. ix. (1856) 149 = sandwicensis.
grandifolia, *Parodi, in Anal. Soc. Cient. Argent.* iv. (1877) 310.—Paraguay.

URERA:—
horrida, Miq. in Mart. Fl. Bras. iv. 192 = baccifera.
Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 479.—Madag.
Hypselodendron, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 203.—Abyss.
Jacquinii, Wedd. l. c. 200 = alceaefolia.
javaenensis, Gaudich. in Freyc. Voy. Bot. 497 = Laportea crenulata.
Kaalae, *Wawra, in Flora*, lvii. (1874) 542.—Ins. Sandvic.
laciniata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 203.—Panama.
lamiifolia, Gaudich. in Freyc. Voy. Bot. 497 = Boehmeria nivea.
Leichardiana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 178 = Laportea photiniphylla.
longifolia, *Wedd. l. c.* 177.—Madag.
madagascariensis, Gaudich. in Freyc. Voy. Bot. 497 = Obetia madagascariensis.
membranacea, *Wedd. in Arch. Mus. Par.* ix. (1856) 103 (Quid?).—Hab.?
microcarpa, *Wedd. l. c.* 156.—Jamaic.
microstigma, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 203 = Laportea microstigma.
mitis, Miq. in Mart. Fl. Bras. iv. i. 191 = alceaefolia.
oblongifolia, Benth. in Hook. Niger Fl. 515 = obovata.
obovata, *Benth. l. c.* 516.—Afr. trop.
oligoloba, *Baker, in Journ. Linn. Soc.* xx. (1883) 265 —Madag.
palmata, Gaudich. in Freyc. Voy. Bot. 497 = Girardinia palmata.
parietariaefolia, *Gaudich. l. c.* (Quid?).—Hab.?
platycarpa, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 202 = Laportea platycarpa.
Punu, *Wedd. l. c.* 200.—Bras.
Radula, *Baker, in Journ. Linn. Soc.* xviii. (1881) 279. —Madag.
rotundifolia, Wedd. in Ann. Sc. Nat. Ser. IV. i. (1854) 177 = Laportea Gigas.
sandwicensis, *Wedd. l. c.* 178.—Ins. Sandvic.
scandens, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 90; *Contrib.* 38.—Paraguay.
Schimperi, Wedd. in Arch. Mus. Par. ix. (1856) 158 = U. Hypselodendron.
Schimperiana, Wedd. l. c. 582 = praec.
simplex, *Wedd. in DC. Prod.* xvi. i. 90.—N. Granat.
sinuata, Blume, Bijdr. 605 = Laportea crenulata.
sinuata, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 201.—Panama.
sphaerophylla, *Baker, in Journ. Linn. Soc.* xxi. (1885) 445.—Madag.
spicata, Wedd. in Arch. Mus. Par. ix. (1856) 155 = Gyrotaenia spicata.
stimulans, Blume, ex Miq. Fl. Ind. Bat. i. ii. 231 = Laportea stimulans.
stimulans, Wedd. in DC. Prod. xvi. i. 98 = Laportea microstigma.
subpeltata, Miq. in Mart. Fl. Bras. iv. i. 189 = alceaefolia.

URETIA, Rafin. Fl. Tellur. iii. 40 (1836) = **Digera**, Forsk. (Amarant.).
alternifolia, Rafin. l. c. = D. arvensis.

URGINEA, Steinh. in Ann. Sc. Nat. Sér. II. i. (1834) 321. t. 14. *LILIACEAE*, Benth. & Hook. f. iii. 810.
Fusifilum, Rafin. Fl. Tellur. ii. 27 (1836).
Montassa, Salisb. Gen. Pl. Fragm. 36 (1866).
Physodia, Salisb. l. c. 37 (1866).
Pilasia, Rafin. Fl. Tellur. iii. 53 (1836).
Squilla, Steinh. in Ann. Sc. Nat. Sér. II. vi. (1836) 276.
Sypharissa, Salisb. Gen. Pl. Fragm. 37 (1866).
Tenicroa, Rafin. Fl. Tellur. iii. 52 (1836).
alooides, Bolus, in Journ. Linn. Soc. xviii. (1881) 395 = Notosceptrum alooides.
altissima, *Baker, l. c.* xiii. (1873) 221.—Afr. trop.
angolensis, *Baker, in Journ. Bot.* xii. (1874) 364.—Afr. trop.
anthericoides, *Steinh. in Ann. Sc. Nat.* Sér. II. i. (1834) 328.—Marocc.
Beccarii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 223. —Abyss.
brevipes, *Baker, in Journ. Bot.* xii. (1874) 364.—Afr. trop.

URGINEA:—
chlorantha, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 248.—Afr. trop.
ciliata, *Baker, in Journ. Linn. Soc.* xiii. (1873) 218. —Afr. austr.
comosa, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 247.—Afr. trop.
congesta, *Wight, Ic.* t. 2065.—Ind. or.
coromandeliana, Wight, l. c. t. 2064 = indica.
exuviata, *Steinh. in Ann. Sc. Nat.* Sér. II. i. (1834) 330.—Afr. austr.
filifolia, *Steinh. l. c.*—Afr. austr.
fragrans, *Steinh. l. c.*—Afr. austr.
fugax, *Steinh. l. c.* 328. t. 14. f. 1.—Reg. Mediterr.
grandiflora, *Baker, in Journ. Linn. Soc.* xiii. (1873) 223.—Oriens.
Hesperia, *Webb & Berth. Phyt. Canar.* iii. 399 —Ins. Canar.
indica, *Kunth, Enum. Pl.* iv. 333.—Ind. or.; Burma; Afr. trop.
japonica, Hort. Par. ex Rev. Hortic. Sér. II. ii. (1843–44) 374 = Scilla japonica.
marginata, *Baker, in Journ. Linn. Soc.* xiii. (1873) 218.—Afr. austr.
maritima, Baker, l. c. 221 = U. Scilla.
mascarenensis, *Baker, in Journ. Bot.* xii. (1874) 363. —Madag.
micrantha, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 294.—Afr. trop.
nematodes, *Baker, in Journ. Linn. Soc.* xiii. (1873) 218.—Afr. austr.
nigritana, *Baker, l. c.* 224.—Afr. trop.
Pancration, Nym. Consp. 729 = U. Scilla.
Petitiana, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 294. —Abyss.
physodes, *Baker, in Journ. Linn. Soc.* xiii. (1873) 217. —Afr. austr.
porphyrostachys, *Baker, ex Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 411.—Ins. Socotra.
psilostachya, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 247.—Afr. trop.
pusilla, *Baker, in Journ. Linn. Soc.* xiii. (1873) 217. —Afr. austr.
Quartiniana, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 284.—Abyss.
rigidifolia, *Baker, in Journ. Bot.* xvi. (1878) 323.—Afr. austr.
Scilla, *Steinh. in Ann. Sc. Nat.* Sér. II. i. (1834) 321. —Afr. austr.
secunda, *Baker, in Journ. Linn. Soc.* xiii. (1873) 222. —Afr. austr.
senegalensis, Kunth, Enum. Pl. iv. 334 = indica.
simensis, *Schweinf. Beitr. Fl. Aethiop.* 291.—Abyss.
undulata, *Steinh. in Ann. Sc. Nat.* Sér. II. i. (1834) 330.—Reg. Mediterr.
zambesiaca, *Baker, in Journ. Linn. Soc.* xiii. (1873) 223.—Afr. trop.

URGINIA, Kunth, Enum. Pl. iv. (1843) 331 = praec.

URINARIA, Medic. Malv. 80 (1787) = **Phyllanthus**, Linn. (Euphorb.).
erecta, Medic. l. c. = P. Niruri.

URMENETEA, Phil. Fl. Atac. 26. t. 3 (1860). *COMPOSITAE*, Benth. & Hook. f. ii. 487.
atacamensis, Phil. l. c. 27.—Chili.

URNECTIS, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
hirta, Rafin. l. c. = S. nigricans.

UROCARPUS, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 54 = **Asterolasia**, F. Muell. (Rutac.).
phebalioides, J. Drumm. ex Harv. l. c. 55 = A. phebalioides.

UROCHLAENA, Nees, Fl. Afr. Austr. 437 (1841). *GRAMINEAE*, Benth. & Hook. f. iii. 1181.
pusilla, *Nees, l. c.* 438.—Afr. austr.

UROCHLOA, Beauv. Agrost. 52. t. 11. f. 1 (1812) = **Panicum**, Linn. (Gramin.).
aristatum, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = P. aristatum.
cimicina, Kunth, Rév. Gram. i. 31 = P. cimicinum.
fasciculata, Kunth, l. c. = P. fasciculiflorum.

UROCHLOA:—
glabra, Brongn. Voy. Coq. Bot. ii. 121 = Panicum glumare.
mauritiana, Boj. Hort. Maurit. 363 = P. javanicum.
panicoides, Beauv. Agrost. 52. t. 11 = P. javanicum.
paniculata, Benth. in Hook. Niger Fl. 558.—Afr. trop.
paspaloides, J. Presl, in Rel. Haenk. i. 319 = P. ambiguum, *Trin.*
pubescens, Kunth, Rév. Gram. i. 31 = P. javanicum.
semialata, Kunth, l. c. = P. semialatum.
uniseta, J. Presl, in Rel. Haenk. i. 319 = Setaria uniseta.

URODESMIUM, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 338. *MELASTOMACEAE*, Benth. & Hook. f. i. 748.
huberioides, *Naud. l. c.*—Bras.

URODON, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 16 = **Pultenaea**, Sm. (Legumin.).
capitatus, Turcz. 17 = P. Urodon.
dasyphyllus, Turcz. l. c. xxvi. (1853) i. 268 = P. Urodon.

UROMORUS, Bur. in DC. Prod. xvii. 236 (1873) = **Paratrophis**, Blume (Urtic.).
anthopophagorum, Bur. l. c.—Ins. Fiji.
philippinensis, Bur. l. c. 237 = P. philippinensis.
tahitensis, Bur. l. c. = Pseudomorus Brunoniana.

UROPAPPUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 424 = **Microseris**, D. Don (Compos.).
grandiflorus, Nutt. l. c. 425 = M. linearifolia.
heterocarpus, Nutt. l. c. = M. Lindleyi.
Lindleyi, Nutt. l. c. = M. Lindleyi.
linearifolius, Nutt. l. c. = M. linearifolia.

UROPEDIUM, Lindl. Orch. Linden. 28 (1846) = **Selenipedium**, Reichb. f. (1854) (Orchid.).
Lindenii, Lindl. l. c. = S. caudatum.

UROPETALON, [Ker-Gawl. in] Bot. Reg. t. 156 (1816) = **Dipcadi**, Medic. (Liliac.).
Bourgaei, Nym. Consp. 731 = D. serotinum.
ciliare, Eckl. & Zeyh. ex Harv. Thes. Cap. ii. 45 = D. ciliare.
concanense, Dalz. in Hook. Kew Journ. ii. (1850) 143 = D. concanense.
crispum, Burch. [ex Ker-Gawl.] in Bot. Reg. sub t. 156 = D. crispum.
depressum, A. Rich. Tent. Fl. Abyss. ii. 325.—Abyss.
erythraeum, Boiss. Fl. Orient. v. 286 = D. erythraeum.
fesoghlense, Solms, in Schweinf. Beitr. Fl. Aethiop. 204.—Afr. bor.
fulvum, Sweet, Hort. Brit. ed. I. 419 = D. serotinum.
glaucum, Burch. ex [Ker-Gawl. in] Bot. Reg. t. 156; et Trav. i. 548 = D. glaucum.
hyacinthoides, Spreng. Syst. iv. Cur. Post. 135 = D. hyacinthoides.
hydsuricum, Baker, in Journ. Linn. Soc. xi. (1871) 397 = seq.
hysudricum, Edgew. in Trans. Linn. Soc. xx. (1846) 88 = D. hydsuricum.
longifolium, Lindl. in Bot. Reg. t. 974 = D. longifolium.
maritimum, Boj. Hort. Maurit. 346, nomen.—Madag.
minimum, Steud. in A. Rich. Tent. Fl. Abyss. ii. 325 = D. minimum.
montanum, Dalz. in Hook. Kew Journ. ii. (1850) 142 = D. montanum.
serotinum, [Ker-Gawl. in] Bot. Reg. sub t. 156 = D. serotinum.
serotinum, Webb & Berth. Phyt. Canar. iii. 341 = D. erythraeum.
tacazzeanum, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 325 = D. tacazzeanum.
umbonatum, Baker, in Saund. Refug. Bot. t. 17 = D. umbonatum.
unicolor, Stocks, in Hook. Kew Journ. iv. (1852) 180 = D. unicolor.
viride, [Ker-Gawl. in] Bot. Reg. sub t. 156 = D. filamentosum.
Welwitschii, Baker, in Saund. Refug. Bot. t. 16 = D. Welwitschii.

UROPETALUM, Burch. Trav. i. 548 (1822); Spreng. Syst. ii. 78 (1825) = praec.

UROPHYLLON, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Tourn. (Liliac.).
caudatum, Salisb. l. c. 36 = O. caudatum.
niveum, Salisb. l. c. = O. niveum.

UROPHYLLUM, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 173, sphalm. = **Urospatha**, Schott (Aroid.).

UROPHYLLUM, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 184 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 71.
Axanthes, Blume, Bijdr. 1002 (1826).
Axanthiopsis, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 195.
Cymelonema, Presl, Epim. Bot. 210 (1849).
Maschalanthe, Blume, Fl. Jav. Praef. p. vi. (1828).
Wallichia, Reinw. ex Blume, in Cat. Gew. Buitenz. 11 (1823).
Afzelii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 73.—Afr. trop.
arboreum, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 194 = glabrum.
biloculare, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 313 = Diplospora Kurzii.
Blumeanum, *Hook. f. Fl. Brit. Ind.* iii. 99.—Malaya.
borneense, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 238.—Borneo.
callicarpoides, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 72.—Afr. trop.
coriaceum, Miq. Fl. Ind. Bat. Suppl. 542 = Randia densiflora.
corymbosum, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 194.—Java; Sumatra.
ellipticum, *Thw. Enum. Pl. Zeyl.* 419.—Zeylan.
elongatum, *Miq. Fl. Ind. Bat.* ii. 225.—Borneo.
glabrum, *Jack, ex Roxb. Fl. Ind. ed. Carey*, ii. 186.—Burma; Malaya.
glabrum, Wall. Cat. n. 8316 = Blumeanum.
Griffithianum, *Hook. f. Fl. Brit. Ind.* iii. 98.—Malaya.
hirsutum, *Hook. f. l. c.*—Malaya.
hirtellum, *Benth. in Hook. Niger Fl.* 397.—Afr. trop.
insulare, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 73.—Afr. trop.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 133.—Borneo.
longifolium, *Hook. f. Fl. Brit. Ind.* iii. 99.—Burma.
Lyallii, *Baker, in Journ. Linn. Soc.* xx. (1883) 165.—Madag.
macrophyllum, *Korth. in Nederl. Kruidk. Arch.* II. ii. (1851) 194.—Java.
memecyloides, *Vidal, Phan. Cuming. Philipp.* 63.—Ins. Philipp.
micranthum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 74.—Afr. trop.
micranthum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 237.—Java.
moluccanum, *Miq. l. c.* 238.—Ins. Sunda.
parviflorum, *Wall. Cat.* n. 8320 = streptopodium.
paucinerve, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 74.—Afr. trop.
peltistigma, *Miq. Fl. Ind. Bat. Suppl.* 542.—Sumatra.
polyneurum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 239.—Borneo.
porphyraceum, *Baill. Hist. Pl.* vii. 378.—Java.
repandulum, Miq. Fl. Ind. Bat. ii. 355 = glabrum.
rubens, *Benth. in Hook. Niger Fl.* 396.—Afr. trop.
streptopodium, *Wall. Cat.* n. 8317, partim.—Malaya.
streptopodium, Wall. l. c., partim = Blumeanum.
strigosum, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 194.—Java; Borneo.
strigosum, Kurz, For. Fl. Brit. Burma, ii. 53 = longifolium.
tomentosum, *Miq. Fl. Ind. Bat.* ii. 224.—Ins. Malaya.
umbellulatum, *Miq. Fl. Ind. Bat. Suppl.* 542.—Sumatra.
villosum, *Jack & Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 184.—Malaya.
viridiflorum, *Schum. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 74.—Afr. trop.
zeylanicum, *Thw. Enum. Pl. Zeyl.* 418.—Zeylan.

UROSKINNERA, Lindl. in Gard. Chron. (1857) 36. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 942.
hirtiflora, *Hemsl. Diagn. Pl. Nov.* 34.—Mexic.
spectabilis, *Lindl. in Gard. Chron.* (1857) 36.—Guatemala.

UROSPATHA, Schott, Aroid. 3. tt. 7–10 (1853). *AROIDEAE*, Benth. & Hook. f. iii. 996.
Urophyllum, C. Koch, in Berl. Allg. Gartenz. xxv. (1857) 173, sphalm.

UROSPATHA:—
affinis, Schott, Aroid. i. 3. t. 8 = caudata.
angustiloba, *Engl. in Mart. Fl. Bras.* iii. II. 123. t. 23.—Bras.
caudata, *Schott, Aroid.* i. 3. t. 8.—Am. austr.
decipiens, *Schott, in Bonplandia*, v. (1857) 128.—Bras.
desciscens, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 99.—Bras.
dubia, *Schott, in Bonplandia*, v. (1857) 128.—Ind. occ.
Friedrichsthalii, *Schott, Aroid.* i. 3. t. 7.—Nicaragua.
grandis, *Schott, in Bonplandia*, v. (1857) 128.—Panama.
Hostmannii, *Schott, l. c.*—Afr. trop.
Langsdorffiana, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 253.—Bras.
maculata, Linden.—Cf. Gard. Chron. (1865) 365 (Quid?).—Bras.
marmorea, Linden.—Cf. l. c. (Quid?).—Bras.
Meyeri, *Schott, Aroid.* i. 4.—Guian.
Poeppigiana, *Schott, l. c.*; *et* t. 10.—Bras.
quinquenervis, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 254 = Spruceana.
Riedeliana, *Schott, l. c.* 253.—Bras.
sagittaefolia, *Schott, Aroid.* i. 4; *et in Bonplandia*, v. (1857) 128.—Bras.
Spruceana, *Schott, in Bonplandia*, v. (1857) 128.—Bras.

UROSPERMUM, Auct. ex Steud. Nom. ed. II. ii. 733 (1841), sphalm. = URASPERMUM, Nutt. = **Osmorhiza**, Rafin. (Umbell.).

UROSPERMUM, Scop. Introd. 122 (1777). *COMPOSITAE*, Benth. & Hook. f. ii. 531.
Arnopogon, Willd. Sp. Pl. iii. 1496 (1803).
asperum, DC. Fl. Fr. iv. 63 = picroides.
capense, Spreng. Syst. iii. 662 = picroides.
Dalechampii, *F. W. Schmidt, Samml. Phys. Aufs.* i. (1795) 276.—Europ. austr.
picroides, *F. W. Schmidt. l. c.* 275.—Europ. austr.

UROSTELMA, Bunge, Enum. Pl. Chin. Bor. 44; (in Mém. Sav. Etr. Pétersb. ii. (1835) 118) = **Metaplexis**, R. Br. (Asclep.).
chinense, Bunge, ll. cc. = M. chinensis.

UROSTIGMA, Gasp. Nov. Gen. Fici, 7 (1844) = **Ficus**, Tourn. (Urtic.).
abutilifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 551 = F. abutilifolia.
acamptophyllum, Miq. Fl. Ind. Bat. Suppl. 439 = F. acamptophylla.
accedens, Miq. Fl. Ind. Bat. i. II. 347 = F. superba.
acrocarpum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 557 = F. acrocarpa.
aegeirophyllum, Miq. l. c. 565 = F. infectoria.
affine, Miq. l. c. 564 = F. religiosa.
aggregatum, Miq. l. c. 559 = F. aggregata.
altissimum, Miq. in Zoll. Syst. Verz. Ind. Archip. 90 = F. altissima.
amazonicum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 541 = F. amazonica.
amblyphyllum, Miq. l. c. 569 = F. retusa.
Ampelos, Dalz. & Gibs. Bomb. Fl. 315 = F. gibbosa.
anacardiifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 587 = F. anacardiifolia.
angustifolium, Miq. l. c. 539 = F. angustifolia.
annulatum, Miq. in Zoll. Syst. Verz. Ind. Archip. 90 = F. annulata.
apiculatum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 570 = F. apiculata.
apiocarpum, Miq. Fl. Ind. Bat. Suppl. 440 = F. apiocarpa.
arbutifolium, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 412 = F. arbutifolia.
Arnottianum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 564 = F. Arnottiana.
atrovirens, Regel, Ind. Sem. Hort. Petrop. (1858) 37 = F. tristis.
atrox, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 550 = F. atrox.
baccatum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 327 = F. baccata.
Backhousii, Miq. in Journ. Bot. Néerl. i. (1861) 240.—Austral.
balicum, Miq. Fl. Ind. Bat. i. II. 348.—Malaya.
bengalense, Gasp. Nov. Gen. Fici, 7; Ricerche, 82. t. 7. f. 14–21 = F. benghalensis.
benjaminum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 583 = F. benjamina.
bibracteatum, Regel, Ind. Sem. Hort. Petrop. (1858) 37 = F. bibracteata.

UROSTIGMA:—
bicorne, Miq. Pl. Jungh. 47 = Ficus pilosa.
Binderianum, Kotschy, in Sitzb. Akad. Wien, Math. Nat. li. (1865) 353 = F. platyphylla.
Binnendykii, Miq. Fl. Ind. Bat. i. II. 341 = F. Binnendijkii.
biverrucellum, Miq. Fl. Ind. Bat. Suppl. 436 = F. annulata.
Bonplandianum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 323 = F. Bonplandiana.
botryapioides, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 538 = F. botryapioides.
brachypodium, Miq. l. c. 552 = F. eugenioides.
bracteatum, Miq. l. c. 576 = F. bracteata.
Burkei, Miq. l. c. 555 = F. Burkei.
callophyllum, Miq. Fl. Ind. Bat. i. II. 349 = F. callophylla.
calyptoceras, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 527 = F. calyptroceras.
calyptratum, Miq. l. c. 553 = F. calyptrata.
canaliculatum, Miq. l. c. 579 = F. glabella.
casapiense, Miq. l. c. 548 = F. casapiensis.
catalpaefolium, Miq. l. c. 551 = F. umbellata.
catappaefolium, Miq. l. c. 533 = F. catappaefolia.
caulobotryum, Miq. l. c. 568 = F. Tjakela.
caulocarpum, Miq. l. c. 568 = F. caulobotrya.
cestrifolium, Miq. l. c. 541 = F. cestrifolia.
ceylonense, Miq. l. c. 570 = F. Tjakela.
chiriquianum, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 412 = F. chiriquiana.
chrysopthalmum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 575 = F. chrysophthalma.
chrysotrix, Miq. in Zoll. Syst. Verz. Ind. Archip. 90, 96 = F. pilosa.
circumscissum, Miq. Pl. Jungh. 292 = F. elastica.
clusiaefolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 543 = F. clusiaefolia.
clusioides, Miq. l. c. 579 = F. clusioides.
complicatum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 325 = F. complicata.
concinnum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 570 = F. glabella.
connivens, Miq. l. c. 573 = F. tomentosa.
conocarpum, Miq. Fl. Ind. Bat. i. II. 350 = F. annulata.
consanguineum, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 408 = F. consanguinea.
consociatum, Miq. in Zoll. Syst. Verz. Ind. Archip. 91 = F. consociata.
cordifolium, Dalz. & Gibs. Bomb. Fl. 242 = F. Arnottiana.
cordifolium, Gasp. Nov. Gen. Fici, 7 = F. cordifolia.
cordifolium, Miq. in Hook. Lond. Journ. Bot. v. (1847) 564 = F. Rumphii.
costaricanum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 322 = F. costaricana.
cotinifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 530 = F. cotinifolia.
courtallense, Miq. l. c. 564 = F. Arnottiana.
crassinervium, Miq. l. c. 535 = F. crassinervia.
crassirameum, Miq. Pl. Jungh. 48 = F. procera.
crocatum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 531 = F. crocata.
cuneatum, Miq. l. c. 585 = Erythroxylum burmanicum.
Cunninghamii, Miq. l. c. 560 = F. Cunninghamii.
cycloneuron, Miq. Fl. Ind. Bat. Suppl. 438.—Sumatra.
cyclophyllum, Miq. in Mart. Fl. Bras. iv. I. 91 = F. cyclophylla.
cystopodum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 532 (err. typ. 352) = F. cystopoda.
Dalhousiae, Miq. l. c. 571 = F. Dalhousiae.
dasycarpum, Miq. l. c. 574 = F. mysorensis.
Dekdekena, Miq. l. c. 558 = F. Dekdekena.
depressum, Miq. Fl. Ind. Bat. i. II. 351 = F. pruniformis.
depressum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 576 = F. annulata.
Diepenhorstii, Miq. Fl. Ind. Bat. Suppl. 439.—Sumatra.
diospyrifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 586 = F. diospyrifolia.
doliarium, Miq. l. c. 527 = F. doliaria.
drupaceum, Miq. l. c. 581 = F. drupacea.
elasticum, Miq. l. c. 578 = F. elastica.
elegans, Miq. in Hook. Niger Fl. 521 = F. elegans.
ellipticum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 528 = F. elliptica.
enorme, Miq. l. c. 544 = F. enormis.
eriobotryoides, Miq. l. c. 587 = F. eriobotryoides.
erythrostictum, Miq. l. c. 540 = F. erythrosticta.

UROSTIGMA :—
eugeniaefolium, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 329 = Ficus eugeniaefolia.
eugenioides, Miq. in Journ. Bot. Néerl. i. (1861) 238 = F. eugenioides.
euneuon, Miq. Fl. Ind. Bat. i. II. 353 = F. nervosa.
euomphalum, Miq. in Mart. Fl. Bras. iv. I. 103 = F. euomphala.
excelsum, Miq. Fl. Ind. Bat. i. II. 350 = F. excelsa.
eximium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 527 = F. eximia.
fagifolium, Miq. l. c. 544 = F. fagifolia.
fagineum, Miq. l. c. 537 = F. faginea.
fazokelense, Miq. l. c. 552 = F. fazokelensis.
ferrugineum, Miq. l. c. 534 = F. ferruginea.
flavescens, Miq. Pl. Jungh. 48 = F. annulata.
Fraseri, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 561 = F. Cunninghamii.
fuligineum, Miq. l. c. 545 = F. hirsuta.
fulvum, Miq. l. c. 528 = F. holosericea ?
fuscescens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 329 = F. fuscescens.
galactophorum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 533 = F. galactophora.
Gamelleira, Miq. in Mart. Fl. Bras. iv. I. 94 = F. Gomelleira, doliarium.
gardeniaefolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 546 = F. hirsuta.
Gardnerianum, Miq. l. c. 530 = F. Gardneriana.
geminum, Miq. l. c. 547 = F. gemina.
giganteum, Miq. l. c. 530 = F. gigantea.
giganteum, Miq. in Zoll. Syst. Verz. Ind. Archip. 90, 96 = F. stupenda.
glabellum, Miq. Fl. Ind. Bat. i. II. 340 = F. glabella.
glaberrimum, Miq. l. c. = F. glaberrima.
glabrum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 536 = F. eximia.
glanduliferum, Miq. l. c. 550 (non Ficus glandulifera).—Bras.
glaucophyllum, Gasp. Nov. Gen. Fici, 7 = F. cordata.
glaucum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 322 = F. glauca.
globosum, Miq. Fl. Ind. Bat. i. II. 335 = F. globosa, *Blume*.
glumosum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 552 = F. glumosa.
glycicarpum, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 409 = F. glycicarpa.
Gomelleira, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 531 = F. Gomelleira.
griseum, Miq. l. c. 588 = F. grisea, *Vahl*.
guatemalanum, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 411 = F. guatemalana.
gummiferum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 529 = F. gummifera, *Miq*.
haematocarpum, Miq. l. c. 584 = F. benjamina.
Hartwegii, Miq. l. c. 545 = F. Hartwegii.
Hasseltii, Miq. Pl. Jungh. 46 = F. pubinervis.
hirsutum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 530 = F. hirsuta.
Hochstetteri, Miq. l. c. 555 = F. Hochstetteri.
holosericeum, Miq. l. c. 531 = F. holosericea.
Huegeli, Miq. l. c. 586 = F. macrophylla.
ilicinum, Sond. in Linnaea, xxiii. (1850) 136 = F. ilicina.
infectorium, Miq. Fl. Ind. Bat. i. II. 339 = F. infectoria.
infectorium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 566 = F. Tjakela.
infestum, Miq. l. c. 550.—Bras.
ingens, Miq. l. c. 554 = F. Schimperiana.
intramarginale, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 320 = F. intramarginalis.
involucratum, Miq. Fl. Ind. Bat. i. II. 334 = F. involucrata.
involutum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 320 = F. involuta.
javanicum, Miq. Fl. Ind. Bat. i. II. 334 = F. cordifolia.
Karet, Miq. l. c. 348 = F. elastica.
Kotschyanum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 553 = F. Kotschyana.
Kunthii, Miq. in Mart. Fl. Bras. iv. I. 96 = F. Kunthii.
lacciferum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 575 = F. altissima.
lachnocaulon, Miq. in Journ. Bot. Néerl. i. (1861) 238 = F. platypoda.
laevigatum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 539 = F. laevigata.
Lambertianum, Miq. l. c. 565 = F. infectoria.
lancifolium, Miq. l. c. 538 = F. lancifolia.

UROSTIGMA :—
lapathifolium, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 319 = Ficus lapathifolium.
laurifolium, Griseb. in Goett. Abh. vii. (1857) 194 = F. lentiginosa.
laurifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 539 = F. laurifolia.
Leichhardtii, Miq. in Journ. Bot. Néerl. i. (1861) 235 = F. platypoda.
lentiginosum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 323 = F. lentiginosa.
leucocarpum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 576 = F. infectoria.
leucostictum, Miq. l. c. 535 = F. leucosticta.
leucotrichum, Miq. in Journ. Bot. Néerl. i. (1861) 234 = F. leucotricha.
Liebmannianum, Miq. in Seem. Bot. Voy. Herald, 195 = F. Liebmanniana.
littorale, Miq. in Zoll. Syst. Verz. Ind. Archip. 91 = F. retusa.
longifolium, Miq. in Mart. Fl. Bras. iv. I. 95 = F. longifolia.
longipes, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 321 = F. longipes.
lucescens, Miq. Fl. Ind. Bat. i. II. 341 = F. infectoria.
Luschnathianum, Miq. in Mart. Fl. Bras. iv. I. 101 = F. Luschnathiana.
luteum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 554 = F. lutea.
macrocarpum, Walp. Ann. i. 684, sphalm. = F. acrocarpa.
macrophyllum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 560 = F. macrophylla.
magnificum, Regel, Gartenfl. (1858) 253; Ind. Sem. Hort. Petrop. (1858) 37.—Afr. trop.
Manok, Miq. in Zoll. Syst. Verz. Ind. Archip. 90, 96 = F. globosa, *Blume*.
Mathewsii, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 549 = F. Mathewsii.
Maximilianum, Miq. l. c. 529 = F. Maximiliana.
microcarpum, Miq. l. c. 583 = F. retusa.
modestum, Miq. l. c. 586 = F. nervosa.
monadenum, Miq. Fl. Ind. Bat. Suppl. 437 = F. sumatrana.
Moritzianum, Miq. in Zoll. Syst. Verz. Ind. Archip. 91, 97 = F. glabella.
Muelleri, Miq. in Journ. Bot. Néerl. i. (1861) 235 = F. Muelleri.
mysorense, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 574 = F. mysorensis.
myxaefolium, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 410 = F. myxaefolia.
natalense, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 556 = F. natalensis.
neglectum, Miq. Fl. Ind. Bat. i. II. 347 = F. benjamina.
nervosum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 585 = F. nervosa.
nesophilum, Miq. in Journ. Bot. Néerl. i. (1861) 237 = F. glabella.
Neumanni, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 534 = F. Neumanni.
nitidum, Gasp. Nov. Gen. Fici, 7 = F. nitidum.
nitidum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 582 = F. retusa.
nudum, Miq. l. c. 584 = F. benjamina.
nummularia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 330 (sp. delend.).—Mexic.
nymphaefolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 527 = F. nymphaeifolia.
obliquum, Miq. l. c. 563 = F. obliqua.
obtusifolium, Miq. l. c. 569 = F. obtusifolia.
obversum, Miq. l. c. 573 = F. tomentosa.
odoratum, Miq. Pl. Jungh. 49 = F. elastica.
Oerstediana, Miq. in Seem. Bot. Voy. Herald, 196. t. 36 = F. Oerstediana.
oligoneuron, Miq. Fl. Ind. Bat. Suppl. 438 = F. oligoneura.
onustum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 575 = F. globosa.
organense, Miq. l. c. 542 = F. organensis.
ottoniaefolium, Miq. l. c. 557 = F. ottoniaefolia.
ovale, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 324 = F. ovalis.
ovatum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 553 = F. ovata.
ovoideum, Miq. l. c. 581, (excl. syn.) = F. retusa.

UROSTIGMA :—
padifolium, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 324 = Ficus padifolia.
pallidum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 530 = F. pallida.
paraënse, Miq. l. c. 534 = F. paraënsis.
parvifolium, Miq. l. c. 570 = F. glabella.
pedunculatum, Miq. l. c. 540 = F. pedunculata, [*Dryand.*].
peracutum, Miq. Fl. Ind. Bat. i. II. 343 = F. pruniformis.
perseaefolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 567 = F. infectoria.
pertusum, Miq. l. c. 548 = F. pertusa.
petiolare, Miq. l. c. 527 = F. petiolaris.
pilosum, Miq. in Zoll. Syst. Verz. Ind. Archip. 90 = F. pilosa.
pisiferum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 580 = F. retusa.
pisocarpum, Miq. Fl. Ind. Bat. i. II. 344 = F. pisocarpa.
planicostatum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 586 = F. planicostata.
platypodium, Miq. l. c. 561 = F. platypoda.
Pohlianum, Miq. in Mart. Fl. Bras. iv. I. 100 = F. Pohliana.
politum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 553 = F. polita.
populifolium, Miq. l. c. 552 = F. populifolia.
populneum, Miq. l. c. 537 = F. populnea.
Princeps, Miq. l. c. 533 = F. Princeps.
prinoides, Miq. l. c. 540 = F. prinoides.
procerum, Miq. Fl. Ind. Bat. i. II. 336 = F. procera.
prolixum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 560 = F. prolixa.
protensum, Griseb. in Bonplandia, vi. (1858) 4 = F. protensa.
pruniforme, Miq. in Zoll. Syst. Verz. Ind. Archip. 91, 97 = F. pruniformis.
Pseudo-benjaminum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 566 = F. Tsiela.
Pseudo-rubrum, Miq. Fl. Ind. Bat. i. II. 343 = F. indica.
Pseudo-Tjiela, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 566 = F. Tsiela.
psychotriaefolium, Miq. l. c. 561 = F. Cunninghamii.
puberulum, Miq. l. c. 562 = F. puberula.
pyrifolium, Miq. Fl. Ind. Bat. i. II. 338.—Java.
reflexum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 569 = F. reflexa.
religiosum, Gasp. Richerche Caprif. 82. t. 7. f. 1–5 = F. religiosa.
retusum, Gasp. Nov. Gen. Fici, 7; Miq. in Hook. Lond. Journ. Bot. vi. (1847) 581 = F. retusa.
rhododendrifolium, Miq. l. c. 579 = F. rhododendrifolia.
rigidum, Miq. l. c. 578 = F. rigida.
Rolandri, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 329, in nota = F. Rolandri.
rubescens, Miq. Fl. Ind. Bat. i. II. 338 = F. indica.
rubicundum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 553 = F. rubicunda.
rubiginosum, Gasp. Nov. Gen. Fici, 7 = F. rubiginosa.
rubrum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 559 = F. rubra.
rufum, Miq. l. c. 528 = F. rufa.
Rumphii, Miq. in Zoll. Syst. Verz. Ind. Archip. 90 = F. Rumphii.
salicifolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 556 = F. salicifolia.
Salzmannianum, Miq. l. c. 547 = F. Salzmanniana.
Stae-crucis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 328, in nota = F. Sancti-crucis.
sapidum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 327 = F. sapida.
Saussureanum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 533 = F. Saussureana.
saxophilum, Miq. Fl. Ind. Bat. i. II. 333 = F. saxophila.
scandens, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 330 = F. pumila.
Schiedeanum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 539 = F. ligustrina.
Schimperi, Miq. l. c. 555 = F. Hochstetteri.
Schumacheri, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 328, in nota = F. Schumacheri.
simile, Regel, Ind. Sem. Hort. Petrop. (1858) 36.—Am. austr.

UROSTIGMA :—
sphaeridiophorum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 543 = Ficus sphaeridiophora.
splendens, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 587 = F. splendens.
squamellosum, Miq. in Journ. Bot. Néerl. i. (1861) 239 = F. macrophylla.
stipulosum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 568 = F. infectoria.
strictum, Miq. Pl. Jungh. 50 = F. stricta.
subapiculatum, Miq. in Mart. Fl. Bras. iv. 1. 101 = F. subapiculata.
subcordatum, Miq. Fl. Ind. Bat. i. II. 349 = F. subcordata.
subcuspidatum, Miq. in Zoll. Syst. Verz. Ind. Archip. 91, 97 = F. pilosa.
subglaucinum, Miq. in Journ. Bot. Néerl. i. (1861) 240.—Austral.
subnymphaeifolia, Miq. Cat. Hort. Amst. 107 = F. subnymphaeifolia.
subpanduraeforme, Miq. l. c. = F. heterophylla.
subtriplinervium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 542 = F. subtriplinervia.
sulcipes, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 413 = F. sulcipes.
sumatranum, Miq. Pl. Jungh. 49 = F. sumatrana.
sundaicum, Miq. Fl. Ind. Bat. i. II. 339 = F. indica.
superbum, Miq. Pl. Jungh. 46 = F. superba.
syringaefolium, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 535 = F. syringaefolia.
tecolutense, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 324 = F. tecolutensis.
terebratum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 558 = F. terebrata.
Thonningii, Miq. in Hook. l. c. 557 = F. microcarpa.
Thunbergii, Miq. l. c. 556 = F. cordata.
timorense, Miq. l. c. 569 = F. infectoria.
Tjakela, Miq. l. c. 567 = F. Tjakela.
Tjiela, Miq. l. c. 580 = F. indica.
Tjiela, Thw. Enum. Pl. Zeyl. 265, partim = F. Trimeni.
trigonatum, Griseb. in Goett. Abh. vii. (1857) 194; Miq. Ann. Mus. Bot. Lugd. Bat. iii. 297 = F. crassinervia.
trigonum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 548 = F. trigona.
tomentellum, Miq. l. c. 531 = F. tomentella.
tomentosum, Miq. l. c. 573 = F. tomentosa.
trichocarpum, Miq. Fl. Ind. Bat. i. II. 338 = F. trichocarpa.
truncatum, Miq. in Zoll. Syst. Verz. Ind. Archip. 91, 97 = F. truncata.
turbinatum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 328 = F. turbinata.
Tweedianum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 547. t. 21 = F. Tweediana.
validum, Miq. Fl. Ind. Bat. i. II. 336 = F. annulata.
velutinum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 531 = F. velutina.
venustum, Miq. l. c. 538 = F. venusta.
verrucosum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 321 = F. verrucosa, *Hemsl.*
virgatum, Miq. Fl. Ind. Bat. i. II. 342 (stirps dub.).—Java.
vitellinum, Miq. in Journ. Bot. Néerl. i. (1861) 237 = F. puberula.
Vogelii, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 553 = F. Vogelii.
volubile, Dalz. & Gibs. Bomb. Fl. 242 = F. gibbosa.
Warszewiczii, Miq. in Versl. en Med. Kon. Akad. Wetensch. xiii. (1862) 410 = F. Warczewiczii.
Wightianum, Miq. in Hook. Lond. Journ. Bot. vi. (1847) 566 = F. infectoria.
xanthophyllum, Miq. l. c. 554 = F. lutea.
xylophyllum, Miq. l. c. 577 = F. xylophylla.
Zollingerianum, Miq. in Zoll. Syst. Verz. Ind. Archip. 91, 97 = F. obscura.

UROSTYLIS, Meissn. Gen. 207; Comm. 133 (1839) = **Layia**, Hook. & Arn. (Compos.).
gaillardioides, Meissn. l. c.; Comm. 133 = L. galardioides.

URSINIA, Gaertn. Fruct. ii. 462. t. 174 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 456.
CHRONOBASIS, DC. ex Benth. & Hook. f. Gen. ii. 457 (1873).

URSINIA :—
LEPTOTIS, Hoffmgg. Verz. Pfl. 72 (1824).
OLIGAERION, Cass. in Dict. Sc. Nat. ii. Suppl. 75 (1816).
SPERMOPHYLLA, Neck. Elem. i. 24 (1790).
SPHENOGYNE, R. Br. in Ait. Hort. Kew. ed. II. v. 142 (1813).
THELYTHAMNOS, Spreng. f. Tent. Suppl. Syst. 25 (1828).
abrotanifolia, *Spreng. Syst.* iii. 609.—Afr. austr.
abyssinica, *Sch. Bip. ex Walp. Rep.* vi. 182.—Abyss.
affinis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 151.—Afr. austr.
albicaulis, *DC. Prod.* v. 689.—Afr. austr.
annua, *Less. in Linnaea*, vi. (1831) 165, in Adnot.—Afr. austr.
anthemoides, *Gaertn. Fruct.* ii. 463, in obs.; *Poir. Encyc.* viii. 257.—Afr. austr.
apiculata, *DC. Prod.* v. 689.—Afr. austr.
cakilefolia, *DC. l. c.* 690.—Afr. austr.
calenduliflora, *Benth. & Hook. f. Gen.* ii. 456.—Afr. austr.
chrysanthemoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 152.—Afr. austr.
crithmifolia, *Spreng. Syst.* iii. 609.—Afr. austr.
crithmoides, Poir. Encyc. viii. 258 = pilifera.
dentata, *Poir. l. c.* 256.—Afr. austr.
filicaulis, *DC. Prod.* v. 689.—Afr. austr.
filiformis, *Griessel. in Linnaea*, v. (1830) 414.—Afr. austr.
foeniculacea, *Poir. Encyc.* viii. 258.—Afr. austr.
indecora, *DC. Prod.* v. 690.—Afr. austr.
leucanthemifolia, *Poir. Encyc.* viii. 258.—Afr. austr.
montana, *DC. Prod.* v. 688.—Afr. austr.
nana, *DC. l. c.* 690.—Afr. austr.
odorata, *Spreng. Syst.* iii. 609.—Afr. austr.
paleacea, *Moench, Meth.* 608.—Afr. austr.
paradoxa, *Gaertn. Fruct.* ii. 462. t. 174. f. 4.—Afr. austr.
pedunculosa, *DC. Prod.* v. 689.—Afr. austr.
pilifera, *Gaertn. Fruct.* ii. 463, in obs.; *Poir. Encyc.* viii. 256.—Afr. austr.
pinnata, Benth. & Hook. f. Gen. ii. 456 = filiformis.
pygmaea, *DC. Prod.* v. 690.—Afr. austr.
radicans, *DC. l. c.* 689.—Afr. austr.
scariosa, *Poir. Encyc.* viii. 257.—Afr. austr.
serrata, *Spreng. Syst.* iii. 609.—Afr. austr.
speciosa, *DC. Prod.* v. 690.—Afr. austr.
tenuifolia, *Poir. Encyc.* viii. 259.—Afr. austr.
tenuiloba, DC. Prod. v. 689 = foeniculacea.
tridentata, *Benth. & Hook. f. Gen.* ii. 456.—Afr. austr.

URTICA, [Tourn.] Linn. Syst. ed. I (1735). *URTICACEAE*, Benth. & Hook. f. iii. 381.
ADICEA, Rafin. Cat. 13 (1824).
ILDEFONSIA, Mart. ex Steud. Nom. ed. II. i. 802 (1840).
RUTICA, Neck. Elem. ii. 202 (1790).
SELEPSION, Rafin. Fl. Tellur. iii. 48 (1836).
acerifolia, Zenker, Pl. Ind. Dec. i. tt. 3, 4 = Girardinia palmata.
acuminata, Poir. Encyc. Suppl. iv. 224 = Urera acuminata.
adoënsis, Hochst. in Flora, xxiv. (1841) 1. Intell. 21 = Girardinia condensata.
aestuans, Linn. Sp. Pl. ed. II. 1396 = Fleurya aestuans.
aestuans, Sieber, ex Steud. Nom. ed. II. ii. 734 = Urera alceaefolia.
affinis, Hook. & Arn. Bot. Beech. Voy. 69 = Fleurya interrupta.
alata, J. F. Gmel. Syst. 268, sphalm. = elata.
alba, Blume, Bijdr. 499 = Leucosyke alba.
alba, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 165 = Leucosyke sumatrana.
alba, Rafin. Fl. Ludov. 114 (Quid?).—Am. bor.
alba, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 165 = Leucosyke celebica.
alba, Zipp. ex Blume, l. c. = Leucosyke ochroneura.
albido-punctata, Steud. Nom. ed. II. ii. 734 = Pipturus argenteus.
alceaefolia, Poir. Encyc. Suppl. iv. 227 = Urera alceaefolia.
alienata, Jacq. ex Blume, Mus. Bot. Lugd. Bat. ii. 231 = Pouzolzia confinis, indica.
alienata, Linn. Syst. ed. XII. 622 = Pouzolzia indica.

URTICA :—
alienata, Roth, ex Blume, Mus. Bot. Lugd. Bat. ii. 231 = Pouzolzia Rothiana.
alnifolia, Bert. ex Griseb. Fl. Brit. W. Ind. 709 = Trema micrantha.
americana, Deless. ex Gaudich. in Freyc. Voy. Bot. 498 = Laportea americana.
amoena, Wall. Cat. n. 4626 (Conocephali sp.).—Ind. or.
andicola, Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 198 = echinata.
angulata, Blume, Bijdr. 494 = Pilea stipulosa.
angustata, Blume, l. c. 499 = Debregeasia velutina.
angustifolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 60 = Boehmeria macrophylla.
angustifolia, Fisch. ex Hornem. Suppl. Hort. Hafn. 107 = dioica.
anisophylla, Wall. Cat. n. 4594 = Pilea anisophylla.
appendiculata, Wall. Cat. n. 4604 = Villebrunea integrifolia.
aquatica, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 291 = dioica.
aquatica, Moon, Cat. Pl. Ceyl. 62 = Boehmeria malabarica.
arborea, Linn. f. Suppl. 417 = Gesnouinia arborea.
arborea, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cypholophus macrocephalus.
arborescens, Blanco, Fl. Filip. ed. II. 483 = Pipturus asper.
arborescens, *Link, Enum. Hort. Berol.* ii. 386.—Ins. Philipp.
arborescens, Poir. ex Wedd. in DC. Prod. xvi. 1. 59 (Leucosykes sp.?).—Hab.?
ardens, Blume, Bijdr. 504 = Laportea microstigma?
ardens, Link, Enum. Hort. Berol. ii. 385 = parviflora.
argentea, Forst. f. Prod. 65 = Pipturus argenteus.
arguta, H. B. & K. Nov. Gen. et Sp. ii. 39 = Pilea arguta.
armata, Kunth, ex Wedd. in DC. Prod. xvi. 1. 101 = Girardinia palmata.
armigera, Presl, Bot. Bemerk. 110 = Urera baccifera.
articulata, Juss. ex Wedd. in Arch. Mus. Par. ix. (1856) 180 = Pilea articulata.
arvensis, Poepp. ex Miq. in Mart. Fl. Bras. iv. 1. 200 = Pilea hyalina.
atlantica, Blume, Mus. Bot. Lugd. Bat. ii. 149 = urens.
atrovirens, *Req. ex Loisel. in Mém. Soc. Linn. Par.* vi. (1827) 432.—Europ. austr.
atrox, Leschen. ex Gaudich. in Freyc. Voy. Bot. 498 = Laportea peltata.
aucklandica, *Hook. f. Fl. Antarct.* 68.—Ins. Auckland.
australis, *Hook. f. l. c.*—N. Zel.
azorica, Hochst. ex Seub. Fl. Azor. 28 = membranacea.
baccifera, Blanco, Fl. Filip. ed. I. 695 = Pipturus asper.
baccifera, Linn. Sp. Pl. ed. II. 1398 = Urera baccifera.
balearica, Linn. Syst. ed. X. 1265 = pilulifera.
ballotaefolia, *Wedd. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 197.—N. Granat.
Berlandiera, Blume, Mus. Bot. Lugd. Bat. ii. 151 = chamaedryoides.
Berteroana, Phil. in Linnaea, xxxiii. (1864–65) 235 = echinata.
betulaefolia, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 59 = Pilea hederacea.
bicolor, Roxb. Hort. Beng. 67; Fl. Ind. iii. 589 = Debregeasia hypoleuca.
bifida, Hassk. Cat. Hort. Bog. Alt. 79 = Boehmeria biloba.
biloba, Hort. ex Regel, Cat. Pl. Hort. Aksakov. 144 (Quid?).—Hab.?
blanda, Wall. ex Wedd. in Arch. Mus. Par. ix. (1856) 365 = Boehmeria platyphylla.
Blitum, Salzm. ex Wedd. l. c. 569 = Phenax vulgaris.
Bollae, Kanitz, in Flora, lv. (1872) 17 = dioica.
bonariensis, Pers. Syn. ii. 552 = spatulata.
Bovista, Riddell, ex Wedd. in Arch. Mus. Par. ix. (1856) 61 = chamaedryoides.
bracteata, Steud. in Flora, xxxiii. (1850) 258 = magellanica.
Breweri, *S. Wats. in Proc. Am. Acad.* x. (1875) 348.—Am. bor. occ.
bulbifera, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 214 = Laportea bulbifera.

URTICA:—
bullata, Blume, Mus. Bot. Lugd. Bat. ii. 145 = grandidentata.
bullosa, Hochst. ex Steud. in Flora, xxxiii. (1850) 259 = Girardinia bulbosa.
cærulea, Blume, Bijdr. 495 = Cypholophus Blumeanus.
caffra, Thunb. Prod. Pl. Cap. 31 = Fleurya capensis ?
callitrichoides, H. B. & K. Nov. Gen. et Sp. ii. 40 = Pilea muscosa.
calophylla, *Schlecht. in Linnaea*, xxvii. (1854) 488 (Quid ?).—Hab. ?
canadensis, Linn. Sp. Pl. 985 = Laportea canadensis.
candicans, Burm. f. Fl. Ind. 197 (err. typ. 297) = Boehmeria nivea.
candidissima, Blume, Bijdr. 498 = Leucosyke javanica.
cannabina, *Linn. Sp. Pl.* 984.—As. bor.; Persia.
capensis, Linn. f. Suppl. 417 = Leidesia capensis.
capensis, E. Mey. ex Wedd. in Arch. Mus. Par. ix. (1856) 582 = Fleurya capensis.
capitata, Blanco, Fl. Filip. ed. II. 483 = Fleurya interrupta.
capitata, Linn. Sp. Pl. 985 = Leucosyke alba.
capitata, Schwein. ex Blume, Mus. Bot. Lugd. Bat. ii. 214 = Boehmeria cylindrica.
capitellata, Poir. Encyc. Suppl. iv. 227 = Leucosyke alba.
caracasana, *Griseb. in Goett. Abh.* xxiv. (1879) 87.—Reg. Argent.
caracasana, Jacq. Hort. Schoenb. iii. 71. t. 386 = Urera alceaefolia.
caravellana, Schrank, Pl. Rar. Hort. Monac. t. 82 = Fleurya aestuans.
carolinensis, Deless. ex Gaudich. in Freyc. Voy. Bot. 498 = Laportea carolinensis.
cataractae, Steud. Nom. ed. II. ii. 734 = Pilea trinervia.
caudata, Burm. f. Fl. Ind. 198 (err. typ. 298) = Boehmeria platyphylla.
caudata, Vahl, Symb. Bot. ii. 96 = membranacea.
caudata, Wall. Cat. n. 4600 = Pouzolzia caudata.
caudigera, Wall. Cat. n. 4585 = Boehmeria platyphylla.
celebica, Blume, Bijdr. 492 = Boehmeria platyphylla.
cellulosa, Spreng. Neue Entdeck. iii. 21 = Pilea ciliaris.
chamaedryoides, *Pursh, Fl. Am. Sept.* i. 113.—Am. bor.
Chesneyana, Blume, Mus. Bot. Lugd. Bat. ii. 153 = pilulifera.
Chiple, Buch.-Ham. ex Wall. Cat. n. 4616 = Pouzolzia viminea.
Churta, Buch.-Ham. ex Wall. l. c. 4608 = Laportea crenulata.
ciliaris, Linn. Syst. ed. X. 1266 = Pilea ciliaris.
ciliata, Sw. in Vet. Acad. Handl. Stockh. (1787) 32 = Pilea Swartzii.
cilindrica, Crantz, Inst. i. 147 = Boehmeria cylindrica.
cinerascens, Blume, Bijdr. 497 = Pipturus argenteus.
cinerascens, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 204 = Pouzolzia cinerascens.
cinerascens, *Poir. Encyc. Suppl.* iv. 226.—Ind. or.
cinerascens, Wall. Cat. n. 4591 = Boehmeria platyphylla.
collina, Bert. ex Spreng. Syst. iii. 839 = Pilea Chamaedrys.
comosa, Buch.-Ham. ex Wall. Cat. sub n. 4616 = Boehmeria sidaefolia.
condensata, Hochst. ex Steud. in Flora, xxxiii. (1850) 260 = Girardinia condensata.
conglobata, Blume, Bijdr. 500 = Pouzolzia conglobata.
conglobata, Pav. ex Wedd. in Arch. Mus. Par. ix. (1856) 102 (Phenacis sp.).—Hab. ?
contracta, *Blume, Bijdr.* 505 (Villebruniae sp. ?).—Java.
convexa, Hornem. ex Spreng. Syst. iii. 839 = pilulifera.
corallina, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 295 = Urera corallina.
cordata, Gaudich. in Freyc. Voy. Bot. 498, in syn. = Fleurya aestuans.
cordatifolia, *Steud. in Flora*, xxxiii. (1850) 258.—China.
cordifolia, Moench, Meth. 351 = pilulifera.
corylifolia, Juss. ex Poir. Encyc. Suppl. iv. 226 = Fleurya aestuans.
corymbosa, Savigny, in Lam. Encyc. iv. 640 = Pilea corymbosa.
crassifolia, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 222 = Boehmeria platyphylla.

URTICA:—
crassifolia, Willd. Sp. Pl. iv. 349 = Pilea densiflora.
crenata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cypholophus macrocephalus.
crenulata, Roxb. Hort. Beng. 67; Fl. Ind. iii. 591 = Laportea crenulata.
crenulata, Sw. in Vet. Akad. Handl. Stockh. viii. (1785) 35; ex ej. Fl. Ind. Occ. i. 319 = Pilea cuneifolia.
cubensis, *Klotzsch, ex Herd. Ind. Sem. Hort. Petrop.* (1858) 38.—Cuba.
cuneata, A. Rich. Fl. Cub. Fanerog. ii. 224 = Fleurya umbellata.
cuneifolia, Sw. in Vet. Acad. Handl. Stockh. viii. (1787) 70 = Pilea cuneifolia.
cuneiformis, Herb. Mus. Par. ex Blume, Mus. Bot. Lugd. Bat. ii. 56 = Pilea insignis.
cuneiformis, Savigny, in Lam. Encyc. iv. 640 = Pilea cuneiformis.
cuneiformis, Steud. Nom. ed. II. ii. 734 = Pilea cuneifolia.
cuspidata, Willd. Sp. Pl. iv. 349 = Pilea lucens.
cylindrica, Linn. Sp. Pl. 984 = Boehmeria cylindrica.
cylindrica, Wall. ex Wedd. in DC. Prod. xvi. 1. 61 = Boehmeria platyphylla.
cylindrica, Wedd. l. c. = Pilea cuneifolia.
cymosa, Hassk. Pl. Jav. Rar. 200 = Fleurya cymosa.
Darwinii, Hook. f. Fl. Antarct. 343 = magellanica.
dasyphylla, Poepp. ex Wedd. in Arch. Mus. Par. ix. (1856) 178 = Pilea globosa.
dauciodora, Pav. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 223 = Pilea dauciodora.
debilis, Endl. Prod. Fl. Norfolk. 37 = Parietaria debilis.
decumana, Roxb. Hort. Beng. 67; Fl. Ind. iii. 587 = Laportea decumana.
depressa, Sw. in Vet. Acad. Handl. Stockh. viii. (1787) 64 = Pilea depressa.
dichotoma, Blume, Bijdr. 499 = Debregeasia dichotoma.
dichroa, Poepp. ex Wedd. in DC. Prod. xvi. 1. 61 = Pilea nummulariaefolia.
diffusa, Sw. in Vet. Acad. Handl. Stockh. viii. (1785) 30 = Pilea diffusa.
dioecia, Hill, Veg. Syst. xxvi. 23 = seq.
dioica, *Linn. Sp. Pl.* 984.—Reg. bor. temp.
dioica, Thunb. Fl. Jap. 69 = Thunbergiana.
dioica, Vell. Fl. Flum. x. t. 18 = spatulata.
diplotricha, Phil. in Linnaea, xxxiii. (1864–65) 234 = dioica.
distachya, Spreng. Syst. iii. 840 = Boehmeria cylindrica ?
divaricata, Forsk. Fl. Aegypt. Arab. 160 = Fleurya aestuans.
divaricata, Linn. Sp. Pl. 985 = Laportea canadensis.
divergens, G. F. W. Mey. Prim. Fl. Esseq. 264 = Fleurya aestuans.
diversifolia, Blume, Bijdr. 498 = Maoutia rugosa.
diversifolia, Link, Enum. Hort. Berol. ii. 385 = Girardinia palmata.
Dodartii, Linn. Syst. ed. X. 1265 = pilulifera.
dubia, Forsk. Fl. Aegypt. Arab. 121 = membranacea.
echinata, *Benth. Pl. Hartw.* 252.—Am. austr.
Eckloniana, Blume, Mus. Bot. Lugd. Bat. ii. 142 = dioica.
elata, Sw. Prod. Veg. Ind. Occ. 37 = Urera sinuata.
elevata, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 15 = morifolia.
elongata, Blume, Bijdr. 494 = Pilea trinervia.
elongata, J. F. Gmel. Syst. 269 = Boehmeria platyphylla.
elongata, *Link, Enum. Hort. Berol.* ii. 385.—Ins. Philipp.
elongata, *Vahl, in West, Bidr. Ste-Croix*, 306.—Ind. occ.
evitata, Wall. Cat. n. 4588 = Laportea terminalis.
excelsa, Bert. ex Steud. in Flora, xxxiii. (1850) 261 = Boehmeria excelsa.
expansa, Sw. Adnot. Bot. 60; et Griseb. Fl. Brit. W. Ind. 155 = Urera expansa.
falcicrenata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 293 = subincisa.
fasciculata, Savigny, in Lam. Encyc. iv. 640 = Pilea pumila, *A. Gray*.
fastigiata, *Blume, Mus. Bot. Lugd. Bat.* ii. 151.—Chili.
fernandeziana, Steud. Nom. ed. II. ii. 734 = Parietaria debilis.
ferocissima, Sweet, Hort. Brit. ed. I. 363 = Girardinia heterophylla.
ferox, Blanco, Fl. Filip. ed. II. 484 = Laportea Gaudichaudiana.

URTICA:—
ferox, *Forst. f. Prod.* 66.—N. Zel.
ficifolia, Savigny, in Lam. Encyc. iv. 638 = Obetia ficifolia.
filiformis, Walt. Fl. Carol. 230 = Boehmeria cylindrica ?
Finlaysoniana, *Wall. Cat.* n. 4644.—Siam ?
flabellata, *H. B. & K. Nov. Gen. et Sp.* ii. 40.—Ecuador.
floribunda, H. B. & K. l. c. 38 = Pilea multiflora.
foliolosa, Wall. ex Wedd. in Arch. Mus. Par. ix. (1856) 103 = Pouzolzia scabra.
foliosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 142.—Japon.
freireaeformis, Steud. Nom. ed. II. ii. 734 = Parietaria debilis.
frutescens, *Poir. Encyc. Suppl.* iv. 226 (Quid ?).—Am. austr.
frutescens, Thunb. Fl. Jap. 70 = Villebrunea frutescens.
fruticosa, Banks, ex Miq. Fl. Ind. Bat. i. ii. 283 = Botrymorus paniculata.
fruticosa, Wight, Ic. t. 694 = Pouzolzia indica.
furialis, Boj. ex Blume, Mus. Bot. Lugd. Bat. ii. 159 = Obetia ficifolia.
galeopsifolia, *Jacq. f. ex Blume, l. c.* 145.—Hab. ?
galeopsifolia, Wierzb. ex Opiz, Natural. n. ix. (1825) 107 = dioica.
Gaudichaudii, Steud. Nom. ed. II. ii. 734 = Fleurya interrupta.
gemina, Lour. Fl. Cochinch. 558 = Acalypha australis.
gibbosa, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 54 = Pilea trinervia.
gigantea, Poir. Encyc. Suppl. iv. 225 = Laportea crenulata.
gigantea, Teysm. ex Miq. Fl. Ind. Bat. i. ii. 282 = Laportea amplissima.
Gigas, A. Cunn. ex Wedd. in Arch. Mus. Par. ix. (1856) 129 = Laportea Gigas.
Girardinia, Steud. Nom. ed. II. ii. 734 = Girardinia heterophylla.
glaberrima, Blume, Bijdr. 493 = Pilea smilacifolia.
glabrata, Clem. ex Vis. Fl. Dalm. i. 217 = dioica.
glandulifera, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 293.—Mexic.
globosa, Pav. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 208 = Pilea globosa.
globulifera, *Roxb. Hort. Beng.* 67; *Fl. Ind.* iii. 593.—Ind. or.
glomerata, Klein, ex Willd. Sp. Pl. iv. 361 = Pouzolzia indica.
glomerulaeflora, *Steud. Nom.* ed. II. ii. 735; *et in Flora*, xxxiii. (1850) 257.—Ins. J. Fernand.
Goglado, Buch.-Ham. in Wall. Cat. n. 4593 = Pilea smilacifolia.
gracilenta, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 122.—Mexic.
gracilescens, Blume, Mus. Bot. Lugd. Bat. ii. 151 = chamaedryoides.
gracilis, Ait. Hort. Kew. ed. I. iii. 341 = dioica.
gracilis, Rafin. Atl. Journ. 16 = chamaedryoides.
grandidentata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 296 = Urera baccifera.
grandidentata, *Miq. Pl. Jungh.* 27.—Java.
grandidentata, Moris, Stirp. Sard. Elench. fasc. ii. 9 = atrovirens.
grandifolia, *Linn. Syst.* ed. X. 1266 = Pilea grandis.
grandifolia, Sw. ex Griseb. Fl. Brit. W. Ind. 159 = Pilea pubescens.
grandifolia, Thunb. ex Wedd. in DC. Prod. xvi. 1. 213 = Boehmeria platyphylla.
grandis, Hook. & Arn. Bot. Beech. Voy. 95 = Boehmeria stipularis.
grossa, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Fleurya grossa.
grossa, Wall. Cat. n. 4615 = Distemon indicum.
grossecrenata, Mart. ex Miq. in Mart. Fl. Bras. iv. 1. 199 = Pilea pubescens.
Hamiltoniana, Wall. Cat. n. 4590 = Pilea scripta, Boehmeria platyphylla.
hastata, Forst. ex Spreng. Syst. iii. 836 = inermis.
Haussknechtii, *Boiss. Fl. Orient.* iv. 1146.—Cataonia.
hederacea, Savigny, in Lam. Encyc. iv. 642 = Pilea hederacea.
heracleifolia, Juss. ex Wedd. in Arch. Mus. Par. ix. (1856) 107 = Obetia ficifolia.
herniariifolia, Willd. Sp. Pl. iv. 359 = Pilea muscosa.
herniarioides, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 64 = Pilea muscosa.

URTICA:—
heterophylla, Vahl, Symb. Bot. i. 76 = Girardinia heterophylla.
heterophylla, Wight, Ic. t. 687 = Girardinia palmata.
himalayensis, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 11 = parviflora.
hippurioides, Griff. Itin. Notes, 362 = Pouzolzia pentandra.
hirsuta, Pav. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 220 = Pilea hirsuta.
hirsuta, Vahl, Symb. Bot. i. 77 = Fleurya aestuans?
hirta, Blume, Bijdr. 495 = Pouzolzia hirta.
hirta, Sw. Fl. Ind. Occ. i. 285 = Phenax urticaefolius.
hispida, DC. Fl. Fr. Suppl. 355 = dioica.
hispida, Muell. Arg. in DC. Prod. xv. II. 880, sphalm. = Acalypha Poiretii.
hispidula, Cariot, Etude des Fleurs, ii. 505 = dioica.
holosericea, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 183.—Am. bor. occ.
horrida, H. B. & K. Nov. Gen. et Sp. ii. 41 = Urera baccifera.
horrida, Link, Enum. Hort. Berol. ii. 385 = Girardinia heterophylla.
humilis, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 69 = Rousselia lappulacea.
hygrophila, Miq. Pl. Jungh. 28 = Pilea hygrophila.
hyperborea, *Jacquem. ex Wedd. in Arch. Mus. Par.* ix. (1856) 68.—Reg. Himal.
Hypselodendron, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 260 = Urera Hypselodendron.
Ildefonsia, *Steud. Nom.* ed. II. ii. 735.—Bras.
inaequalis, Juss. ex Poir. Encyc. Suppl. iv. 222 = Pilea inaequalis.
incana, Blume, Bijdr. 497 = Pipturus velutinus.
incisa, Heyne, ex Wall. Cat. sub n. 4603 = Girardinia heterophylla.
incisa, *Poir. Encyc. Suppl.* iv. 224.—Austral.; N. Zel.
inermis, *Spreng. Syst.* iii. 836.—Ins. Pacif.
iners, Forsk. Fl. Aegypt. Arab. 160 = urens.
iners, Nees & Mart. ex Wedd. in Arch. Mus. Par. ix. (1856) 271 = Pilea Maximiliani.
integerrima, Blume, in Flora, vii. (1824) 291 = Phytocrene macrophylla.
integrifolia, Savigny, in Lam. Encyc. iv. 637 = pilulifera.
interrupta, Jacq. ex Blume, Mus. Bot. Lugd. Bat. ii. 220 = Boehmeria Jacquiniana.
interrupta, Linn. Sp. Pl. 985 = Fleurya interrupta.
interrupta, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = ovalifolia.
involucrata, Roxb. Hort. Beng. 67; Fl. Ind. iii. 592 = Macaranga involucrata.
involucrata, Sims, Bot. Mag. t. 2481 = Pilea pubescens.
involuta, Steud. Nom. ed. II. ii. 735 = Macaranga involucrata.
japonica, Blanco, Fl. Filip. ed. I. 694 = Pouzolzia viminea.
japonica, Linn. f. Suppl. 418 = Boehmeria platyphylla.
japonica, Thunb. Fl. Jap. 70 (non Urticacea).—Japon.
javaënsis, Poir. ex Gaudich. in Freyc. Voy. Bot. 497 = Laportea crenulata.
javanica, Blume, Bijdr. 503 = Fleurya interrupta.
kioviensis, Bogowitch, in Bull. Soc. Nat. Mosc. (1843) n. II. p. 324 = dioica.
laciniata, Goudot, ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 203 = Urera laciniata.
laetevirens, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 236.—Japon.
lamiifolia, Juss. ex Pers. Syn. ii. 555 = Boehmeria nivea.
lamiifolia, Poir. ex Wedd. in Arch. Mus. Par. ix. (1856) 103 (Debregeasiae sp.).—Hab.?
lanceolata, A. Rich. Fl. Cub. Fanerog. ii. 225 = Pilea nudicaulis.
lanceolata, Savigny, in Lam. Encyc. iv. 639 = Pilea lanceolata.
lappulacea, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 69 = Rousselia lappulacea.
latifolia, *Moon, Cat. Pl. Ceyl.* 62, nomen.—Zeylan.
latifolia, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 113 = Fleurya aestuans.
laurifolia, *Poir. Encyc. Suppl.* iv. 224.—Reg. Magellan.
laurina, Sm. in Rees, Cyclop. xxxvii. n. 76 (Villebruneae sp.).—Ins. Amboin.
laxa, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 58 = Pilea laxa.
leptophylla, H. B. & K. Nov. Gen. et Sp. ii. 39 = magellanica.

URTICA:—
leptostachya, Sm. in Rees, Cycl. xxxvii. n. 65 = Droguetia ovata.
leptostachys, Juss. ex Pers. Syn. ii. 554 = Droguetia ovata.
leucophaea, Blume, Bijdr. 493 = Pilea leucophaea.
leucophylla, Wall. Cat. n. 4067 = Debregeasia Wallichiana.
leucotrys, Rafin. Fl. Ludov. 114 (Quid?).—Am. bor.
limoniodora, Pav. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 216 = Pilea citriodora.
Lindeniana, A. Rich. Fl. Cub. Fanerog. ii. 225 = Pilea serrulata.
lineata, *Roxb. Hort. Beng.* [103]; *Fl. Ind.* iii. 589.—Ins. Molucc.
Linkiana, Heynh. Nom. i. 832 = Girardinia heterophylla.
lobata, E. Mey. ex Blume, Mus. Bot. Lugd. Bat. ii. 144 = Meyeri.
lobulata, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Meyeri.
lomatocarpa, Hochst. ex Steud. in Flora, xxxiii. (1850) 260 = Fleurya interrupta.
longifolia, Burm. f. Fl. Ind. 197 (err. typ. 297) = Debregeasia velutina.
longifolia, Willd. Sp. Pl. iv. 348 = Pilea verbascifolia.
longipes, Mart. ex Miq. Fl. Bras. iv. I. 201 = Pilea hyalina.
Lowei, Seub. Fl. Azor. 28 = morifolia.
lucens, Savigny, in Lam. Encyc. iv. 641 = Pilea lucens.
lucida, Dum.-Cours. Bot. Cult. ed. II. vi. 369 = Pilea densiflora.
lucida, Sw. in Vet. Akad. Handl. Stockh. (1785) 31 = Pilea lucida.
lucifuga, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 285 = incisa.
lusitanica, Brot. Fl. Lusit. i. 205 = membranacea.
lutescens, Blume, Bijdr. 494 = Cypholophus rufescens.
Lyallii, *S. Wats. in Proc. Am. Acad.* x. (1875) 348.—Am. bor. occ.
macrophylla, Hort. ex Schlecht. in Linnaea, xxvii. (1854) 488 = calophylla.
macrophylla, Thunb. Fl. Jap. 69 = Boehmeria platyphylla.
macrostachya, Wall. ex D. Don, Prod. Fl. Nep. 60 = Boehmeria macrophylla.
maculata, Savigny, in Lam. Encyc. iv. 644 (Pileae sp.).—Ins. S. Doming.
madagascariensis, Juss. ex Poir. Encyc. Suppl. iv. 227 = Obetia madagascariensis.
magellanica, *Juss. ex Poir. l. c.* 223.—Reg. Magellan.
major, Kanitz, in Bot. Zeit. xx. (1862) 190; xxi. (1863) 54 = dioica.
malabarica, Wall. Cat. n. 4610 = Boehmeria malabarica.
manillensis, Walp. ex Bur. in DC. Prod. xvii. 256 = Fatoua pilosa.
marginata, Poepp. ex Wedd. in Arch. Mus. Par. ix. (1856) 238 = Pilea marginata.
mariana, Mill. Gard. Dict. ed. VIII. n. 7 = Boehmeria cylindrica.
Masafuerae, *Phil. in Bot. Zeit.* xiv. (1856) 647; *et in Anal. Univ. Chil.* (1856) 167.—Ins. Masafuer.
melanocarpa, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 210 = Cypholophus melanocarpus.
melastomoides, Hort. ex Spreng. Syst. iii. 842 = Leucosyke alba.
melastomoides, H. B. & K. Nov. Gen. et Sp. ii. 38 = Pilea multiflora.
melastomoides, Poir. Encyc. Suppl. iv. 223 = Pilea trinervia.
membranacea, Barbieri, ex Blume, Mus. Bot. Lugd. Bat. ii. 149 = urens.
membranacea, Bourg. ex Wedd. in DC. Prod. xvi. I. 54 = morifolia.
membranacea, *Poir. ex Savigny, in Lam. Encyc.* iv. 638.—Reg. Mediterr.
membranacea, Sw. ex Wedd. in Arch. Mus. Par. ix. (1856) 103 = Urera membranacea.
mexicana, Blume, Mus. Bot. Lugd. Bat. ii. 140 = dioica.
mexicana, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 291.—Mexic.
Meyeniana, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 422 = Laportea stimulans?
Meyeri, *Wedd. in Arch. Mus. Par.* ix. (1856) 64.—Afr. austr.
micrantha, *Kunth & Bouché, in Ann. Sc. Nat.* Sér. III. vii. (1847) 182.—Hab.?

URTICA:—
microphylla, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 1147 = Haussknechtii.
microphylla, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 66 = Pilea microphylla.
Millettii, *Hook. & Arn. Bot. Beech. Voy.* 214 (sp. dub.).—China.
minima, *Roxb. Hort. Beng.* [103]; *Fl. Ind.* iii. 585.—Ins. Molucc.
minor, Lam. Fl. Fr. ii. 194 = urens.
minutifolia, *Griseb. in Goett. Abh.* xix. (1874) 112.—Reg. Argent.
mitis, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Fleurya capensis.
mitis, Vell. Fl. Flum. x. t. 19 = Urera alceaefolia.
mollis, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 222 = Boehmeria platyphylla.
mollis, *Steud. Nom.* ed. II. ii. 735; *et in Flora*, xxxiii. (1850) 258.—Chili.
mollissima, Blume, Bijdr. 501 = Pipturus mollissimus.
mollissima, Sm. in Rees, Cyclop. xxxvii. n. 54 = Pouzolzia laevigata.
moluccana, *Blume, Bijdr.* 492.—Ins. Molucc.
monilifera, Sm. ex Wedd. in Arch. Mus. Par. ix. (1856) 445 = Pipturus rotundifolius.
monoica, Gilib. Exercit. ii. 450 = urens.
montana, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 227 = Boehmeria monticola.
morifolia, *Poir. Encyc. Suppl.* iv. 223.—Ins. Canar.
morifolia, Sm. in Rees, Cycl. xxxvii. n. 20 = ballotaefolia.
moroides, A. Cunn. ex Wedd. in Arch. Mus. Par. ix. (1856) 142 = Laportea moroides.
multiflora, Poir. Encyc. Suppl. iv. 223 = Pilea multiflora.
muralis, *Vahl, Symb. Bot.* i. 77.—Arab.
muricata, Heyne, ex Wall. Cat. n. 4612 = Debregeasia velutina.
Mutisiana, Spreng. Syst. iii. 840 = Pilea multiflora.
Myrapi, Korth. ex Blume, Mus. Bot. Lugd. Bat. ii. 227 = Boehmeria monticola.
nana, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 60 = Pouzolzia indica.
nana, *Hook. Journ. Bot.* i. (1834) 232.—N. Granat.
naucliflora, Roxb. Hort. Beng. 67; Fl. Ind. iii. 592 = Conocephalus suaveolens.
neglecta, Guss. Ind. Sem. Hort. Boccad. (1825) 11 = membranacea.
nemorosa, H. B. & K. Nov. Gen. et Sp. ii. 42 = Fleurya aestuans.
nicaraguensis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 292.—Nicaragua.
nitida, Hort. Berol. ex Walp. Ann. i. 647 = Pilea densiflora.
nitida, Vell. Fl. Flum. x. t. 20 = Urera baccifera.
nivea, Linn. Sp. Pl. 985 = Boehmeria nivea.
nudicaulis, Sw. in Vet. Akad. Handl. Stockh. (1785) 34 = Pilea nudicaulis.
nummularifolia, Sw. l. c. viii. (1787) 63. t. 1. f. 2 = Pilea nummulariaefolia.
nutans, Poepp. ex Wedd. in Arch. Mus. Par. ix. (1856) 196 = Pilea nutans.
obesa, Wall. Cat. n. 4587 = Pilea Wightii.
oblongata, Koch, ex Maly, Fl. N. Steierm. 63 = urens.
obnoxia, Hassk. Cat. Hort. Bog. Alt. 305 = Laportea microstigma?
obtusata, Vahl, ex Wedd. in Arch. Mus. Par. ix. (1856) 262 = Pilea obtusata.
ochroneura, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 165 = Leucosyke ochroneura.
orbiculata, Rich. ex Steud. Nom. ed. II. ii. 735 (Pileae sp.?).—Hab.?
oreophila, Miq. Pl. Jungh. 27 = Pilea trinervia.
Orizabae, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 292 = chamaedryoides.
ovalifolia, *Blume, Bijdr.* 504.—Java.
ovalifolia, Buching. in Flora, xxviii. (1845) 88 = Fleurya capensis.
ovalifolia, Stokes, Bot. Mat. Med. iv. 372 = urens.
palmata, Forsk. Fl. Aegypt. Arab. 159 = Girardinia heterophylla.
palustris, Juss. ex Pers. Syn. ii. 552 = Boehmeria cylindrica.
paniculata, Gaudich. in Freyc. Voy. Bot. 497 = Fleurya ruderalis.
paniculata, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 589 = Wallichiana.
parasitica, Blume, Bijdr. 506 = Pellionia javanica?

URTICA:—
parasitica, Forsk. Fl. Aegypt. Arab. 160 = muralis.
Parietaria, Linn. Sp. Pl. 985 = Pilea ciliaris.
Parietaria, Savigny, in Lam. Encyc. iv. 639 = Pilea articulata.
parietariaefolia, Bert. ex Steud. Nom. ed. II. ii. 735 = Parietaria debilis.
parietariaefolia, Deless. ex Presl, Rep. Bot. 162 = Urera parietariaefolia.
parva, Blume, Bijdr. 502 = Pouzolzia indica.
parviflora, *Roxb. Hort. Beng.* 67 ; *Fl. Ind.* iii. 581.—Reg. Himal.
parvifolia, Buch.-Ham. ex Wall. Cat. sub n. 4616 = Pouzolzia viminea.
parvula, Blume, Mus. Bot. Lugd. Bat. ii. 150 = urens.
pauciflora, Hochst. ex Steud. in Flora, xxxiii. (1850) 258 = Droguetia diffusa.
pedunculares, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Fleurya capensis.
pedunculares, Sm. in Rees, Cycl. xxxvii. n. 6 = Pilea trinervia.
pedunculata, Wedd. in Arch. Mus. Par. ix. (1856) 118 = Fleurya capensis.
pellucida, Labill. Sert. Austro-Caled. 79. t. 80 = Pipturus velutinus.
peltata, Blume, Bijdr. 496 = Laportea peltata.
pendula, Willd. Sp. Pl. iv. 358 = Pilea rupipendia.
penduliflora, Wall. Cat. n. 4595 = Boehmeria macrophylla.
pentandra, Roxb. Hort. Beng. 67; Fl. Ind. iii. 583 = Pouzolzia pentandra.
peploides, Steud. Nom. ed. II. ii. 735 = Pilea peploides.
petiolaris, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 215 = Pilea stipulosa.
petiolata, Steud. Nom. ed. II. ii. 735 = Fleurya aestuans.
photiniphylla, A. Cunn. ex Wedd. in Arch. Mus. Par. ix. (1856) 139 = Laportea photiniphylla.
pilosa, Lour. Fl. Cochinch. 558 = Acalypha Poiretii.
pilosiuscula, Blume, Bijdr. 491 = Boehmeria platyphylla.
pilulifera, *Linn. Sp. Pl.* 983.—Reg. Mediterr.; Oriens; Ind. or.
Pittoni, Blume, Mus. Bot. Lugd. Bat. ii. 147 = membranacea.
platyphylla, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 60 = Boehmeria platyphylla.
platyphylla, Wedd. in Arch. Mus. Par. ix. (1856) 86 = dioica.
polystachya, Wall. Cat. n. 4584 = Boehmeria polystachya.
portulacina, Spreng. Syst. iii. 838 = Pilea muscosa.
portulacoides, Spreng. Neue Entdeck. iii. 21 = Pilea muscosa.
procera, Muhl. ex Willd. Sp. Pl. iv. 353 = dioica.
procridioides, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Pouzolzia procridioides.
propinqua, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 293 = chamaedryoides.
Pseudo-dioica, Steud. Nom. ed. II. ii. 735; et in Flora, xxxiii. (1850) 257 = dioica.
pubescens, Ledeb. Fl. Alt. iv. 240, in nota = dioica.
pubescens, Roxb. ex Wall. Cat. n. 4614, sphalm. = *frutescens*.
pulchella, *Hort. Angl. ex Link, Enum. Hort. Berol.* ii. 385.—Ind. or.
pulcherrima, Roxb. Hort. Beng. 67; Fl. Ind. iii. 588 = Sarcochlamys pulcherrima.
pulcherrima, Wall. Cat. n. 4596 = Boehmeria macrophylla.
pulegifolia, Poir. Encyc. Suppl. iv. 224 = Pilea pulegifolia.
Pulus, Steud. Nom. ed. II. ii. 735 = Laportea microstigma?
pumila, Euphras. ex Griseb. Fl. Brit. W. Ind. 159 = Pilea hyalina.
pumila, Linn. Sp. Pl. 984 = Pilea pumila.
pumila, Schwein. ex Blume, Mus. Bot. Lugd. Bat. ii. 213 = Boehmeria cylindrica.
punctata, Buch.-Ham. ex Wall. Cat. sub n. 4616 = Pouzolzia viminea.
punctata, H. B. & K. Nov. Gen. et Sp. ii. 38 = Pilea punctata.
purpurascens, Nutt. in Trans. Am. Phil. Soc. v. (1837) 169 = chamaedryoides.
pusilla, Poir. Encyc. Suppl. iv. 224 = Australina pusilla.
pustulata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 294 = Laportea canadensis.

URTICA:—
Puya, Buch.-Ham. ex Wall. Cat. n. 4605 = Maoutia Puya.
quadrifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 263 = Pilea quadrifolia.
quadristipulata, Dulac, Fl. Hautes-Pyr. 150 = urens.
quinquenervis, Wall. Cat. n. 4601 = Pouzolzia Bennettiana.
racemosa, Burm. ex Wedd. in Arch. Mus. Par. ix. (1856) 112 = Fleurya aestuans.
racemosa, Gaudich. in Freyc. Voy. Bot. 497 = Fleurya interrupta.
radicans, Bolla, in Verh. Ver. Naturk. Pressburg, i. (1856) Abh. 9; Sitzb. 24 = dioica.
radicans, Pohl, ex Miq. in Mart. Fl. Bras. iv. 1. 199 = Pilea pubescens.
radicans, Sw. in Vet. Akad. Handl. Stockh. (1785) 29 = Pilea radicans.
radicans, Wight, Ic. t. 1974 = Pilea Wightii.
Radula, E. Mey. in Drège, Zwei Pfl. Docum. 228 = Australina capensis.
reclinata, Hort. Par. ex Wedd. in Arch. Mus. Par. ix. (1856) 219 = Pilea crassifolia.
recurvata, Otto, ex Blume, Mus. Bot. Lugd. Bat. ii. 147 = membranacea.
repanda, Blume, Bijdr. 501 = Pipturus repandus.
repanda, Kolmann, ex Planch. in DC. Prod. xvii. 199 = Trema amboinensis.
repens, Sw. in Vet. Acad. Handl. Stockh. viii. (1787) 61 = Pilea repens.
repens, Wall. Cat. n. 4619 = Pouzolzia indica.
reticulata, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 70 = Pilea reticulata.
rhodopleura, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 168 = Villebrunea rhodopleura.
rhombea, Linn. f. Suppl. 417 = Pilea rhombea.
Riedlei, Steud. Nom. ed. II. ii. 736 = Pilea lucens.
rigida, G. Don, ex Benth. in Hook. Niger Fl. 519 = Boehmeria rigida.
rivularis, Schlecht. ex Wedd. in Arch. Mus. Par. ix. (1856) 541 = Droguetia diffusa.
rotundifolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 60 = Boehmeria platyphylla.
rotundifolia, Savigny, in Lam. Encyc. iv. 644 = Pipturus rotundifolius.
Roxburghii, Steud. Nom. ed. II. ii. 736 = Villebrunea frutescens.
rubescens, Blume, Bijdr. 506 = Villebrunea scabra.
rubra, Rafin. Fl. Ludov. 113 (Quid?).—Am. bor.
rubra, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 203 = Pouzolzia viminea.
rubricaulis, *Herd. ex Regel, Ind. Sem. Hort. Petrop.* (1860) 34.—Hab.?
rubricaulis, Hornem. ex Blume, Mus. Bot. Lugd. Bat. ii. 150 = stachyoides?
rubricaulis, *Span. in Linnaea*, xv. (1841) 344.—Ins. Timor.
ruderalis, Forst. f. Prod. 66 = Fleurya ruderalis.
rufa, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 59 = Pilea rufa.
rugosa, Reinw. ex Blume, Mus. Bot. Lugd. Bat. ii. 223 = Boehmeria platyphylla.
rugosa, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 60 = Pilea rugosa.
rugosissima, Reinw. in Blume, Bijdr. 490 = Boehmeria platyphylla.
rugosula, Wall. Cat. n. 4597 = Boehmeria rugulosa.
Rumphii, Kostel. Allg. Med. Pharm. Fl. ii. 400 (Laporteae sp.?).—Ins. Molucc.
rupestris, *Guss. Adnot. Cat. Pl. Boccad.* (1821) 15.—Sicil.
rupipendia, Savigny, in Lam. Encyc. iv. 640 = Pilea rupipendia.
salicifolia, Roxb. Hort. Beng. 67 = Debregeasia hypoleuca.
salicifolia, Vahl, ex Wedd. in Arch. Mus. Par. ix. (1856) 334 = Lecanthus Wightii, Elatostema papillosum.
sandwicensis, Wedd. l. c. 67 = Hesperocnide sandwicensis.
sanguinea, Blume, Bijdr. 501 = Pouzolzia viminea.
saxicola, Mor. ex Wedd. in Arch. Mus. Par. ix. (1856) 105 (al. ord.).—Hab.?
scabra, Blume, Bijdr. 504 (al. ord.).—Java.
scabra, Zoll. ex Blume, Mus. Bot. Lugd. Bat ii. 167 = Villebrunea scabra.
scabrella, Hornem. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 29 = U. Ildefonsia.

URTICA:—
scabrella, Roxb. Hort. Beng. 67; Fl. Ind. iii. 581 = Boehmeria platyphylla.
Scalpe, Comm. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 198 = spatulata.
Schimperiana, Hochst. ex Steud. in Flora, xxxiii. (1850) 259 = Fleurya aestuans.
scripta, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 59 = Pilea scripta.
semicrenata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 167 = Villebrunea semicrenata.
semidentata, Juss. ex Poir. Encyc. Suppl. iv. 222 = Pilea semidentata.
semiserrata, Sm. ex Wedd. in Arch. Mus. Par. ix. (1856) 569 = Phenax vulgaris.
serpyllacea, H. B. & K. Nov. Gen. et Sp. ii. 37 = Pilea serpyllifolia.
Serra, *Blume, Mus. Bot. Lugd. Bat.* ii. 140.—Mexic.
serrulata, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 65 = Pilea serrulata.
sessiliflora, Blanco, Fl. Filip. ed. I. 696 = Fleurya interrupta.
sessiliflora, Sw. in Vet. Akad. Handl. Stockh. (1785) 33 = Pilea sessiliflora.
sessilifolia, Savigny, in Lam. Encyc. iv. 642 = Pilea sessilifolia.
sessilifolia, Wedd. in Arch. Mus. Par. ix. (1856) 105, sphalm. = *sessiliflora*, Blanco.
sicula, Gasp. ex Guss. Fl. Sic. Syn. ii. 580 = dioica.
simensis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 260.—Abyss.
sinuata, Blume, Bijdr. 605 = Laportea crenulata.
smilacifolia, Wall. ex Voigt, Hort. Suburb. Calc. 279 = Pilea smilacifolia.
spatulata, *Sm. in Rees, Cyclop.* xxxvii. n. 17.—Am. austr.
sphaerocephala, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 589 = Leucosyke alba.
spicata, Thunb. Fl. Jap. 69; Blume, Bijdr. 492 = Boehmeria platyphylla.
spicigera, Steud. Nom. ed. II. ii. 736 = Fleurya interrupta.
spinosa, Blume, Bijdr. 507 = Taxotrophis javanica.
spiralis, Domb. ex Wedd. in Ann. Sc. Nat. Sér. III. xviii. (1852) 232 = Myriocarpa densiflora.
spirealis, *Blume, Mus. Bot. Lugd. Bat.* ii. 152.—Mexic.
squamigera, Wall. Cat. n. 4592 = Chamabainia cuspidata.
stachydifolia, Kunth & Bouché, Ind. Sem. Hort. Berol. (1846) 11 = chamaedryoides.
stachyoides, *Webb & Berth. Phyt. Canar.* iii. 259. t. 210.—Ins. Canar.
stachyophylla, Webb, ex Wedd. in Arch. Mus. Par. ix. (1856) 73 = stachyoides.
stimulans, Linn. f. Suppl. 418 = Laportea stimulans.
stipulacea, *Bertol. Misc. Bot.* ii. 24.—As. Min.
stipulosa, Miq. Pl. Jungh. 28 = Pilea stipulosa.
stolonifera, Sw. in Vet. Acad. Handl. Stockh. viii. (1787) 62 = Pilea stolonifera.
strigosa, Blume, Bijdr. 495 = Pouzolzia indica.
subcordata, Comm. ex Steud. Nom. ed. II. ii. 736 = Pilea lucens.
subincisa, *Benth. Pl. Hartw.* 293.—Mexic.
submitis, Boiss. Fl. Orient. iv. 1146 = dioica.
subperforata, Wall. ex Wedd. in Arch. Mus. Par. ix. (1856) 383 = Boehmeria subperforata.
succulenta, Salzm. ex Wedd. in Arch. Mus. Par. ix. (1856) 177 = Pilea hyalina.
suffruticosa, Roxb. Hort. Beng. 67; Fl. Ind. iii. 584 = Pouzolzia indica.
superba, Wall. Cat. n. 4625 (Conocephali sp.).—Ind. or.
sycophylla, Bory, Voy. i. 281 = Obetia ficifolia.
sylvatica, Blume, Bijdr. 506 = Villebrunea sylvatica.
tasmanica, F. Muell. First Gen. Rep. 18 = Australina Muelleri.
tenacissima, Roxb. Hort. Beng. 67; Fl. Ind. iii. 590 = Boehmeria nivea.
tenuis, Steud. in Flora, xxxiii. (1850) 257 = Pouzolzia indica.
ternata, Heyne, ex Wall. Cat. n. 4599 = Pouzolzia Wightii.
ternifolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 59 = Boehmeria platyphylla.
tetraphylla, Steud. in Flora, xxxiii. (1850) 260 = Pilea tetraphylla.
Thunbergiana, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 214.—Japon.

URTICA :—
thymifolia, H. B. & K. Nov. Gen. et Sp. ii. 37 = Pilea globosa.
tiliaefolia, H. B. & K. l. c. 42 = Urera alceaefolia.
tomentosa, Juss. ex Wedd. in Arch. Mus. Par. ix. (1856) 367 = Boehmeria platyphylla.
trachycarpa, Wedd. l. c. 95 = holosericea.
triandra, Blume, Bijdr. 496 = Pouzolzia parvifolia.
trianthemoides, Sw. in Vet. Akad. Handl. Stockh. viii. (1787) 68 = Pilea muscosa.
trilobata, Savigny, in Lam. Encyc. iv. 639 = Pilea trilobata.
trinervia, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 582 = Pilea trinervia.
triplinervia, Poir. Encyc. Suppl. iv. 224 = Pilea lucens.
triplinervia, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 220 = Pouzolzia indica.
triplinervis, Buch.-Ham. ex Wall. Cat. sub n. 4590 = Pilea scripta.
triplinervis, Juss. ex Pers. Syn. ii. 552 = Pilea lucens.
triplinervis, Wall. Cat. n. 4617 = Pouzolzia indica.
tuberculata, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 159 = Fleurya aestuans.
tuberosa, Roxb. Hort. Beng. 67; Fl. Ind. iii. 583 = Pouzolzia tuberosa.
ulmifolia, H. B. & K. Nov. Gen. et Sp. ii. 42 = Urera alceaefolia.
ulmifolia, Vahl, ex Wedd. in Arch. Mus. Par. ix. (1856) 569 = Phenax vulgaris.
umbellata, Blanco, Fl. Filip. ed. I. 696 = Laportea Gaudichaudiana.
umbellata, Bory, Voy. iii. 173 = Pilea insignis.
umbrosa, Wall. Cat. n. 4598 = Pilea umbrosa.
uragera, Steud. Nom. ed. II. ii. 736 = Boehmeria platyphylla.
urens, Bert. ex Steud. Nom. ed. II. ii. 736 = spathulata.
urens, *Linn. Sp. Pl.* 984.—Geront. temp.
urentissima, Comm. ex Pers. Syn. ii. 552 = spatulata.
urophylla, Wall. ex Blume, Mus. Bot. Lugd. Bat. ii. 166 = Villebrunea integrifolia.
utilis, Hort. ex De Vriese, Pl. Ind. Bat. Or. 146 = Boehmeria nivea.
variegata, Spreng. Syst. iii. 839 = Pilea sessiliflora.
venosa, Wall. Cat. n. 4602 A = Boehmeria polystachya.
venosa, Wall. l. c. B = Boehmeria rugulosa.
verbascifolia, Savigny, in Lam. Encyc. iv. 641 = Pilea verbascifolia.
verna, Rafin. Atl. Journ. 16 = chamaedryoides.
veronicaefolia, F. G. Dietr. Vollst. Lexik. Gaertn. x. 296 (Quid?).—Jamaic.
verrucosa, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 295 = Urera alceaefolia.
verrucosa, Moon, Cat. Pl. Ceyl. 62, nomen = Debregeasia velutina.
verticillata, Vahl, Symb. Bot. i. 76 = urens.
vesicaria, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 587 = Pouzolzia vesicaria.
villosa, Blanco, Fl. Filip. ed. I. 695 = Pouzolzia indica.
villosa, Salzm. ex Miq. in Mart. Fl. Bras. iv. 1. 197 = Fleurya aestuans.
villosa, *Thunb. Fl. Jap.* 70.—Japon.
viminea, Wall. Cat. n. 4616 = Boehmeria sidaefolia, Pouzolzia viminea.
virgata, Forst. f. Prod. 66 = Boehmeria platyphylla.
virulens, Royle, Illustr. Bot. Himal. 333, 466 (Index), sphalm. = seq.
virulenta, Wall. Cat. n. 4586 = parviflora.
Wallichiana, *Steud. Nom.* ed. II. ii. 736.—Ins. Molucc.
Whitlowi, Muhl. in Green, Cat. Pl. N. York; ex Wedd. in Arch. Mus. Par. ix. (1856) 123 = Laportea canadensis.

URUCU, Adans. Fam. ii. 381 (1763) = **Bixa**, Linn.

URUPARIA, Rafin. Sylva Tellur. 148 (1838) = OUROUPARIA, Aubl. = **Uncaria**, Schreb. (Rubiac.).
versicolor, Rafin. l. c. = Uncaria guianensis.

URVILLAEA, DC. Prod. i. 602 (1824) = seq.

URVILLEA, H. B. & K. Nov. Gen. et Sp. v. 105. t. 440 (1821). *SAPINDACEAE*, Benth. & Hook. f. i. 392.
affinis, *Schlecht. in Linnaea*, xviii. (1844) 38 (err. typ. 54).—Bras.

URVILLEA:—
Berteriana, *DC. Prod.* i. 602.—N. Granat.
dasycarpa, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 265.—Mexic.
euryptera, *Griseb. in Goett. Abh.* xxiv. (1879) 79.—Reg. Argent.
euryptera, Griseb. l. c., partim = Serjania hebecarpa.
ferruginea, Lindl. Bot. Reg. t. 1077 = Serjania cuspidata.
glabra, Benth. in Hook. Kew Journ. iii. (1851) 192, in obs. = Serjania tenuis.
glabra, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 353.—Bras.
glabrata, Steud. Nom. ed. II. i. 736, sphalm. = praec.
intermedia, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 263.—Bras.
laevis, *Radlk. in Atti, Congr. Bot. Firenze*, 1874 (1876), 63.—Bras.
mexicana, *A. Gray, Pl. Wright.* i. 38.—Mexic.
pubescens, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1179, nomen (Quid?).—Guian.
rufescens, *Cambess. in A. St. Hil. Fl. Bras. Mer.* i. 354.—Bras.
Schomburgkii, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1180, nomen = Serjania paucidentata.
Seriana, Griseb. Fl. Brit. W. Ind. 123 = ulmacea.
stipitata, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 264.—Bras.
tenera, *Wawra, in Oestr. Bot. Zeitschr.* xxix. (1879) 215.—Bras.
tridentata, Miq. in Linnaea, xix. (1847) 434 = Serjania dentata.
triphylla, *Radlk. Monog. Serj.* 47, 73.—Bras.
ulmacea, *H. B. & K. Nov. Gen. et Sp.* v. 105. t. 440.—Venezuela.
uniloba, Radlk. Monog. Serj. (1875) 173 = ulmacea.
villosa, *Radlk. in Sitzb. Math.-Phys. Akad. Muench.* viii. (1878) 265.—Bras.

USIONIS, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
aquatica, Rafin. l. c. = S. cinerea.
aurita, Rafin. l. c. = S. aurita.
Dicksoniana, Rafin. l. c. = S. phylicifolia.
foliolosa, Rafin. l. c. = S. foliolosa.
fuscata, Rafin. l. c. = S. petiolaris.
grisea, Rafin. l. c. = S. petiolaris.
humilis, Rafin. l. c. 17 = S. humilis.
pedicillaris, Rafin. l. c. 14 = S. myrtilloides.
petiolaris, Rafin. l. c. = S. petiolaris.
rosmarinifolia, Rafin. l. c. = S. repens.
uliginosa, Rafin. l. c. = S. aurita.

USTERIA, Cav. Ic. ii. 15. t. 116 (1793) = **Maurandia**, Ortega (Scrophul.).
antirrhiniflora, Poir. Encyc. Suppl. v. 405 = Antirrhinum maurandioides.
scandens, Cav. Ic. ii. 15 = M. semperflorens.

USTERIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Acalypha**, Linn. (Euphorb.).
racemosa, Dennst. l. c. = A. paniculata.

USTERIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 480 = **Scilla**, Linn. (Liliac.).
disperma, Medic. in Usteri, Ann. Bot. ii. (1791) 11 = S. hispanica.
hyacinthiflora, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 480 = S. festalis.
secunda, Medic. in Usteri, Ann. Bot. ii. (1791) 12 = S. festalis.

USTERIA, Willd. in Roem. & Usteri, Mag. (1790) 151; et ex Ges. Naturf. Fr. Berl. Schr. x. (1792) 51. t. 2. *LOGANIACEAE*, Benth. & Hook. f. ii. 796.
MONODYNAMIS, J. F. Gmel. Syst. 10 (1791).
guianensis, Steud. Nom. ed. II. ii. 736, sphalm. = seq.
guineensis, *Willd. in Ges. Naturf. Fr. Berl. Schr.* x. (1792) 55.—Afr. trop.
volubilis, *Afzel. Gen. Guin.* [27], cum ic. in titulo.—Afr. trop.

USUBIS, Burm. f. Fl. Ind. 89. t. 32. f. 1 (1768) = **Allophylus**, Linn. (Sapind.).
triphylla, Burm. f. l. c. = A. Cobbe.

UTANIA, G. Don, Gen. Syst. iv. 663 (1837) = **Fagraea**, Thunb. (Logan.).
morindaefolia, G. Don, l. c. = F. morindaefolia.

UTANICA, Steud. Nom. ed. II. ii. 736 (1841), sphalm. = praec.

UTEA, Jaume St. Hil. Expos. ii. 203 (1805) = OUTEA, Aubl. = **Macrolobium**, Schreb. (Legumin.).

UTERVERIA, Bertol. Pl. Nov. Hort. Bonon. ii. 8 (1839) = **Capparis**, Tourn.
Breynia, Bertol. l. c. 10 = C. Breynia.
comosa, Bertol. l. c. 9 = C. comosa.
Cynophallophora, Bertol. l. c. = C. Cynophallophora.
Eustachiana, Bertol. l. c. = C. Eustachiana.
frondosa, Bertol. l. c. 8 = C. frondosa.
linearis, Bertol. l. c. 10 = C. linearis.
tenuisiliqua, Bertol. l. c. = C. tenuisiliqua.
verrucosa, Bertol. l. c. 9 = C. verrucosa.

UTLERIA, Bedd. ex Benth. & Hook. f. Gen. ii. 743 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.
salicifolia, *Bedd. ex Hook. f. Fl. Brit. Ind.* iv. 7.—Ind. or.

UTRICULARIA, Linn. Syst. ed. I. (1735). *LENTIBULARIEAE*, Benth. & Hook. f. ii 987.
AKENTRA, Benj. in Linnaea, xx. (1847) 319.
ASKOFAKE, ENSKIDE, HAMULIA, LEPIACTIS, MEGOPIZA, MEIONULA, MELONEURA, NELIPUS, PERSONULA, PLECTOMA, PLESISA, STOMOISIA, TRILOBULINA, TRIXAPIAS, VESICULINA, Rafin. Fl. Tellur. iv. 108–110 (1836).
DIUROSPERMUM, Edgew. in Proc. Linn. Soc. i. (1847) 351.
LENTIBULARIA, Adans. Fam. ii. 208 (1763).
acuta, Benj. in Linnaea, xx. (1847) 314 = caerulea.
adpressa, *Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 159.—Bras.
affinis, *Wight, in Hook. Kew Journ.* i. (1849) 373; *et Ic.* t. 1580/1.—Ind. or.
alata, *Benj. in Bot. Zeit.* iii. (1845) 212.—Ind. or.
alba, Hoffmgg. ex Link, Jahrb. i. III. (1820) 54 = obtusa.
albiflora, *R. Br. Prod.* 431.—Austral.
albiflora, *Griff. Notul.* iv. 168.—Burma.
albo-coerulea, *Dalz. in Hook. Kew Journ.* iii. (1851) 279.—Ind. or.
alpina, *Georgi, Beschr. Russ. Reich.* III. iv. 655.—Sibir. or.
alpina, *Jacq. Enum. Pl. Carib.* 11.—Am. trop.
ambigua, A. DC. in DC. Prod. viii. 9 = exoleta.
amethystina, *Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 162.—Am. trop.
andicola, *Benj. in Linnaea*, xx. (1847) 308.—Mexic.
angulosa, *Poir. Encyc.* viii. 273.—Guian.
angustifolia, *Benj. in Linnaea*, xx. (1847) 311.—Bras.
anomala, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 153.—Am. trop.
antirrhinoides, Wall. Cat. n. 1498 B = bifida.
aphylla, *Ruiz & Pav. Fl. Per.* i. 20. t. 31. f. d.—Peruv.
appressa, Benj. in Mart. Fl. Bras. x. 244 = adpressa.
arcuata, *Wight, in Hook. Kew Journ.* i. (1849) 372; *et Ic.* t. 1571/1.—Ind. or.
arenaria, *A. DC. in DC. Prod.* viii. 20.—Afr. trop.
aurea, *Lour. Fl. Cochinch.* 26.—Cochinch.
australis, R. Br. Prod. 430 = flexuosa.
barbata, *R. Br. l. c.* 432.—Austral.
Baueri, *R. Br. l. c.* 431.—Austral.
Benjaminiana, *Oliver, in Journ. Linn. Soc.* iv. (1860) 176.—Guian.
bicolor, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 162.—Bras.
bifida, Boj. ex A. DC. in DC. Prod. viii. 25 = hians.
bifida, *Linn. Sp. Pl.* 18.—As. et Austral. trop.
bifida, Macrae, ex A. DC. in DC. Prod. viii. 21 = racemosa.
bifida, Wight, Ic. t. 1584 f. 2 = verticillata.
biflora, Bonpl. ex A. DC. in DC. Prod. viii. 10 = obtusa.
biflora, *Lam. Illustr.* i. 50.—Am. bor.
biflora, Nutt. Gen. Am. i. 13 = longirostris.
biflora, Roxb. Fl. Ind. ed. Carey, i. 144 = exoleta.
biflora, Wall. Cat. n. 1498 = bifida.
biloba, *R. Br. Prod.* 432.—Austral.
bipartita, Chapm. Fl. S. U. St. 283 = fibrosa.
bipartita, *Ell. Sketch*, i. 22.—Am. bor.
bisquamata, *Schrank, in Flora*, vii. (1824) II. *Beil.* 9.—Afr. austr.

UTRICULARIA :—
Blanchetii, *A. DC. in DC. Prod.* viii. 13.—Bras.
Blumei, Miq. Fl. Ind. Bat. ii. 994 = flexuosa.
Botecudorum, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 151.—Bras.
brachiata, *Oliver, in Journ. Linn. Soc.* iii. (1859) 187.—Reg. Himal.
brachypoda, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1578/1 = affinis.
Bremii, Heer, ex Koelliker, Verz. Phan. Gew. Zuer. 142 = minor.
brevicaulis, Benj. in Linnaea, xx. (1847) 303 = bifida.
breviscapa, *Wright, ex Griseb. Cat. Pl. Cub.* 161.—Cuba.
caerulea, Benth. Fl. Hongk. 256 = affinis.
caerulea, A. DC. in DC. Prod. viii. 19 = racemosa.
caerulea, Heyne, ex Wall. Cat. sub n. 1493 = reticulata.
caerulea, *Linn. Sp. Pl.* 18.—Ind. or.
calumpitensis, *Llanos, Fragm.* 11.—Ins. Philipp.
calycifida, *Benj. in Linnaea*, xx. (1847) 493.—Guian.
campestris, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 333 = racemosa.
capensis, *Spreng. Syst.* i. 50.—Afr. austr.
capillacea, Thw. Enum. Pl. Zeyl. 171 = Wallichiana.
capillacea, Wall. Cat. n. 6399 = scandens.
capillacea, *Willd. Sp. Pl.* i. 113.—Ind. or.
capillaris, *D. Don, Prod. Fl. Nep.* 85.—Nepal.
capillaris, Hoffmgg. ex A. Dietr. Sp. Pl. i. 466 = filiformis.
ceratophylla, Michx. Fl. Bor. Am. i. 12 = inflata.
cernua, Hoffmgg. ex Benj. in Mart. Fl. Bras. x. 237 = oligosperma.
Chamissonis, G. Weber, ex Benj. l. c. 241 = globulariaefolia.
chrysantha, *R. Br. Prod.* 432.—Austral.
clandestina, *Nutt. ex A. Gray, Man. Bot. N. U. St.* ed. I. 287.—Am. bor.
coccinea, *Benj. in Linnaea*, xx. (1847) 490.—Am. trop.
coerulea, Auct. = caerulea.
Colensoi, *Hook. f. Fl. N. Zel.* i. 206.—N. Zel.
colorata, *Benj. in Mart. Fl. Bras.* x. 240.—Bras.
complanata, Wall. Cat. n. 1497 = racemosa.
compressa, *R. Br. Prod.* 431.—Austral.
conferta, *Hassk. in Versl. en Kon. Akad. Wetensch.* iv. (1855) 137; *Retzia*, i. 90.—Java?
conferta, Wight, in Hook. Kew Journ. i. (1849) 372; et Ic. t. 1575 = caerulea.
confervaefolia, Jacks. ex D. Don, Prod. Fl. Nep. 84 = flexuosa.
cornuta, *Michx. Fl. Bor. Am.* i. 12.—Am. bor. et austr.
crenata, Vahl, Enum. i. 203 = aphylla.
cucullata, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 152.—Bras.
cyanea, *R. Br. Prod.* 431.—Austral.
cymbantha, [*Welw. ex*] *Oliver, in Journ. Linn. Soc.* ix. (1867) 147.—Afr. trop.
decipiens, Dalz. in Hook. Kew Journ. iii. (1851) 279 = affinis.
dentata, G. Weber, ex Benj. in Mart. Fl. Bras. x. 248 = polyschista.
denticulata, *Benj. in Linnaea*, xx. (1847) 492.—Mexic.
diantha, A. DC. in DC. Prod. viii. 21 = bifida.
diantha, Schult. Mant. i. 169 = exoleta.
dichotoma, *Labill. Nov. Holl. Pl.* i. 11. t. 8.—Austral.
diflora, Roxb. Hort. Beng. 4 = exoleta.
diploglossa, *Welw. ex Oliver, in Journ. Linn. Soc.* ix. (1867) 147.—Afr. trop.
dubia, *Benj. in Linnaea*, xx. (1847) 494.—N. Granat.
dubia, Rosellini, ex Ces. Pass. & Gib. Comp. Fl. Ital. 416 = vulgaris?
Ecklonii, *Spreng. Syst.* iv. *Cur. Post.* 336.—Afr. austr.
elegans, Wall. Cat. n. 1502 = exoleta.
emarginata, *Benj. in Linnaea*, xx. (1847) 489.—Mexic.
Endresii, *Reichb. f. in Gard. Chron.* (1874) II. 582.—Costa Rica.
erectiflora, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 159.—Bras.
exilis, *Oliver, in Journ. Linn. Soc.* ix. (1867) 154.—Afr. trop.
exoleta, *R. Br. Prod.* 430.—As. et Austral. trop.
extensa, Hance, in Walp. Ann. iii. 3 = flexuosa.
fasciculata, Roxb. Hort. Beng. 4; Fl. Ind. i. 143 = flexuosa.
fibrosa, Chapm. Fl. S. U. St. 283 = biflora.

UTRICULARIA :—
fibrosa, *Walt. Fl. Carol.* 64.—Am. bor.
filicaulis, Wall. Cat. n. 1501 = racemosa.
filiformis, *Roem. & Schult. Syst.* i. 199.—Bras.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* ii. 225.—Venezuela.
firmula, *Welw. ex Oliver, in Journ. Linn. Soc.* ix. (1867) 152.—Afr. trop.
flaccida, *A. DC. in DC. Prod.* viii. 17.—Bras.
flava, R. Br. Prod. 432 = chrysantha.
flexuosa, *Vahl, Enum.* i. 198.—As. et Austral. trop.
Fockeana, *Miq. Stirp. Surin. Sel.* 113.—Guian.
foliosa, *Linn. Sp. Pl.* 18.—Am. trop.
foliosa, Salzm. ex A. DC. in DC. Prod. viii. 26 = Genlisea filiformis.
fontana, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 163.—Bras.
fornicata, Le Conte, in Ann. Lyc. N. York, i. (1824) 76. t. 6. f. 6 = gibba.
foveolata, *Edgew. in Proc. Linn. Soc.* i. (1847) 351.—Bengal.
fulva, *F. Muell. in Trans. Phil. Soc. Vict.* iii. (1858) 63.—Austral.
furcata, Pers. Syn. i. 18 = aphylla.
furcellata, *Oliver, in Journ. Linn. Soc.* iii. (1859) 189.—Reg. Himal.
fusiformis, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 9.—Bras.
Gayana, *A. DC. in DC. Prod.* viii. 9.—Chili.
geminiloba, *Benj. in Mart. Fl. Bras.* x. 242.—Bras.
geminiscapa, Benj. in Linnaea, xx. (1847) 305 = clandestina.
genliseoides, *Benj. in Mart. Fl. Bras.* x. 250.—Bras.
gibba, Le Conte, in Ann. Lyc. N. York, i. (1824) 76 = minor.
gibba, *Linn. Sp. Pl.* 18.—Am. bor.
gibbosa, Hill, Veg. Syst. xx. 45 = praec.
glandulosa, G. Weber, ex Benj. in Mart. Fl. Bras. x. 254 = Genlisea violacea.
globulariaefolia, *Mart. ex Benj. l. c.* 241.—Bras.
glochidiata, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1581 = orbiculata.
Gomezii, *A. DC. in DC. Prod.* viii. 13.—Bras.
gracilis, H. B. & K. Nov. Gen. et Sp. ii. 225 = obtusa.
gracilis, Lepr. ex Oliver, in Journ. Linn. Soc. ix. (1867) 147 = exoleta.
Grafiana, Koch, in Flora, xxx. (1847) 265 = intermedia.
graminifolia, Banks & Soland. ex Oliver, in Journ. Linn. Soc. iii. (1859) 179 = affinis?
graminifolia, R. Br. Prod. 432 = cyanea.
graminifolia, J. Grah. Cat. Pl. Bomb. 165 = reticulata.
graminifolia, Vahl, Enum. i. 195 = caerulea.
grandiflora, Pers. Syn. i. 18 = alpina.
Greenei, Oakes, in Hovey, Mag. Hortic. vii. (1841) 180 = resupinata.
Griffithii, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1576 = affinis.
guianensis, *Splitg. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 323.—Guian.
guyanensis, *A. DC. in DC. Prod.* viii. 11.—Guian.
Harlandi, *Oliver, ex Benth. Fl. Hongk.* 257.—Ins. Hongkong.
heterosepala, *Benj. in Linnaea*, xx. (1847) 310.—Ins. Philipp.
hians, *A. DC. in DC. Prod.* viii. 25.—Madag.
hirta, *Klein, ex Link, Jahrb.* i. III. (1820) 55.—Ind. or.
hirtella, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 156.—Bras.
hispida, *Lam. Illustr.* i. 50.—Guian.
Hookeri, *Lehm. Pugill.* viii. 47.—Austral.
Humboldtii, *Rob. Schomb. in Verh. Befoerd. Gartenb. Preuss.* xv. (1841) 141. t. 3.—Guian.
humilis, Heyne, ex Wall. Cat. n. 1495 = reticulata.
humilis, Vahl, Enum. i. 203 = bifida.
hydrocarpa, *Vahl, l. c.* 200.—Guian.
ibarensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 427.—Madag.
inaequalis, Benj. in Linnaea, xx. (1847) 304 = flexuosa.
inaequalis, Benj. in Mart. Fl. Bras. x. 243 = subulata.
inaequalis, A. DC. in DC. Prod. viii. 666 = Hookeri.
inflata, *Walt. Fl. Carol.* 64.—Am. bor.
inflexa, Forsk. Fl. Aegypt. Arab. 9 = stellaris.
integra, Le Conte, in Ann. Lyc. N. York, i. (1824) 76 = biflora.
intermedia, *Hayne, in Schrad. Journ.* (1800) I. 18; *Roth, Catalect.* fasc. ii. (1800) p. ante 1.—Europ.; As. et Am. bor.

UTRICULARIA :—
intricata, Griff. ex Oliver, in Journ. Linn. Soc. iii. (1859) 179 = affinis.
Jamesoniana, *Oliver, l. c.* iv. (1860) 169.—Ecuador.
juncea, *Vahl, Enum.* i. 202.—Am. bor. et austr.
kumaonensis, *Oliver, in Journ. Linn. Soc.* iii. (1859) 189.—Reg. Himal.
laciniata, Mart. ex Benj. in Mart. Fl. Bras. x. 251 = Polypompholyx laciniata.
laciniata, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 157.—Bras.
lagoënsis, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 1.—Bras.
lasiocaulis, *F. Muell. in Melb. Chemist*, (1885) 50.—Austral.
lateriflora, *R. Br. Prod.* 431.—Austral.
latilabiata, Benj. in Linnaea, xx. (1847) 315 = Hookeri.
latiloba, Benth. in Enum. Pl. Hueg. 82 = Polypompholyx multifida.
laxa, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 160.—Bras.
Lehmanni, *Benj. in Bot. Zeit.* iii. (1845) 213.—Afr. trop.
leptantha, Benj. in Linnaea, xx. (1847) 313 = pusilla.
leptoplectra, *F. Muell. in Melb. Chemist* (1885), 50.—Austral.
lilacina, *Griff. Notul.* iv. 168.—Burma.
limosa, *R. Br. Prod.* 432.—Austral.
linarioides, *Welw. ex Oliver, in Journ. Linn. Soc.* ix. (1867) 151.—Afr. trop.
linearifolia, Benj. in Linnaea, xx. (1847) 306 = Hookeri.
lingulata, *Baker, in Journ. Linn. Soc.* xx. (1883) 216.—Madag.
lingulata, G. Weber, ex Benj. in Mart. Fl. Bras. x. 246 = nana.
livida, *E. Mey. Comm. Pl. Afr. Austr.* 281.—Afr. austr.
longecalcarata, *Benj. in Linnaea*, xx. (1847) 314.—Afr. austr.
longeciliata, A. DC. in DC. Prod. viii. 23 = Polypompholyx laciniata.
longifolia, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 545.—Bras.
longirostris, Le Conte, in Ell. Sketch, i. 21 = fibrosa.
longissima, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 973, nomen.—Guian.
Lundii, *A. DC. in DC. Prod.* viii. 14.—Bras.
macrocarpa, Wall. Cat. n. 1494 = stellaris.
macroceras, A. DC. in DC. Prod. viii. 666 = Menziesii.
macrolepis, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1580/2 = Wallichiana.
macroptera, G. Brueckn. ex Garcke, Fl. Deutschl. ed. III. 274 = minor.
macrorhiza, Le Conte, in Ann. Lyc. N. York, i. (1824) 73 = foliosa.
madagascariensis, *A. DC. in DC. Prod.* viii. 20.—Madag.
major, Koen. ex Vahl, Enum. i. 200, in nota = hydrocarpa.
major, *Schmidel, Ic. Pl. ed. Bischof*, 80. t. 21. ff. a-l.—Europ.
Mannii, *Oliver, in Journ. Linn. Soc.* ix. (1867) 149.—Afr. trop.
maxima, *G. Weber, ex Benj. in Mart. Fl. Bras.* x. 249.—Bras.
media, Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat. Sér. II. xi. (1839) 155 = subulata.
media, Schum. Enum. Pl. Saell. i. 9 = intermedia.
Menziesii, *R. Br. Prod.* 431.—Austral.
micrantha, *Benj. in Linnaea*, xx. (1847) 495.—N. Granat.
microcarpa, F. G. Dietr. Vollst. Lexik. Gaertn. x. 305 = pusilla.
micropetala, *Sm. in Rees, Cycl.* xxxvii. n. 58.—Afr. trop.
millefolia, Nutt. ex Tuckerm. in Am. Journ. Sc. Ser. I. xlv. (1843) 28 = intermedia.
minima, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 6.—Bras.
minor, *Linn. Sp. Pl.* 18.—Europ.; As. et Am. bor.
minor, Pursh, Fl. Am. Sept. i. 15 = gibba.
minor, Thuill. Fl. Par. ed. II. 12 = intermedia.
minutissima, Boj. ex A. DC. in DC. Prod. viii. 25 = perpusilla.
minutissima, *Vahl, Enum.* i. 204.—Malacca.
modesta, A. DC. in DC. Prod. viii. 17.—Bras.

UTRICULARIA:—
monantha, *Benj. in Mart. Fl. Bras.* x. 244.—Bras.
monanthos, *Hook. f. Fl. Tasm.* i. 299.—Tasman.; N. Zel.
montana, Jacq. Select. Am. 7. t. 6 = alpina.
montana, *Poir. Encyc.* viii. 268.—Ins. Martinic.
multicaulis, *Oliver, in Journ. Linn. Soc.* iii. (1859) 188.—Reg. Himal.
multifida, R. Br. Prod. 432 = Polypompholyx multifida.
muscosa, *Benj. in Linnaea*, xx. (1847) 489.—Guian.
myriocista, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 150.—Am. trop.
nana, *A. St. Hil. & Girard, l. c.* 156.—Bras.
natans, Salzm. ex A. St. Hil. & Girard, l. c. 154 = pallens.
neglecta, Lehm. Ind. Scholar. Hamb. (1828) 38 = major.
nelumbifolia, *Gardn. in Hook. Ic. Pl.* t. 505–506.—Bras.
neottioides, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 157.—Bras.
nephrophylla, *Benj. in Mart. Fl. Bras.* x. 246.—Bras.
nervosa, *G. Weber, ex Benj. l. c.* 247.—Bras.
nivea, Vahl, Enum. i. 203 = racemosa.
Novae-Zelandiae, *Hook. f. Fl. N. Zel.* i. 206.—N. Zel.
obovata, Miq. in Linnaea, xix. (1847) 144 = amethystina.
obovata, G. Weber, ex Benj. in Mart. Fl. Bras. x. 253 = Genlisea repens.
obtusa, *Sw. Prod. Veg. Ind. Occ.* 14.—Am. trop.
obtusata, Sw. Fl. Ind. Occ. i. 41 = praec.
obtusiloba, *Benj. in Linnaea*, xx. (1847) 312.—Ind. or.; China.
occidentalis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 95.—Am. bor. occ.
ochroleuca, *R. Hartm. in Bot. Notiser* (1857), 30.—Europ.
oligocista, *Benj. in Linnaea*, xx. (1847) 491.—N. Granat.
oligosperma, *A. St. Hil. Voy. Distr. Diam.* ii. 427.—Bras.
olivacea, *Wright, ex Griseb. Cat. Pl. Cub.* 161.—Cuba.
oppositiflora, R. Br. Prod. 430 = dichotoma.
oppositifolia, Spreng. Syst. i. 50 = dichotoma.
orbiculata, *Wall. Cat.* n. 1500.—As. trop.
oryzetorum, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 331 = reticulata.
palatina, G. Weber, ex Benj. in Mart. Fl. Bras. x. 239.—Bras.
pallens, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 153.—Guian.
Parkeri, *Baker, in Journ. Linn. Soc.* xx. (1883) 216.—Madag.
Parkeriana, *A. DC. in DC. Prod.* viii. 9.—Am. trop.
parviflora, R. Br. Prod. 431 = lateriflora.
pauciflora, *Blume, Bijdr.* 739.—Java.
paucifolia, *Benj. in Linnaea*, xx. (1847) 309.—Ind. or.
pectinata, *Splitg. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 325.—Guian.
pedicellata, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1578/2 = caerulea.
peltata, *Spruce, ex Oliver, in Journ. Linn. Soc.* iv. (1860) 172.—Am. trop.
perminuta, F. Muell. Fragm. vi. 160 = violacea.
perpusilla, *A. DC. in DC. Prod.* viii. 25.—Madag.
personata, Le Conte, in Ann. Lyc. N. York, i. (1824) 77. t. 6 = cornuta.
picta, *Warm. in Kjoeb. Vidensk. Meddel.* (1874) 10.—Bras.
polygaloides, Edgew. in Proc. Linn. Soc. i. (1847) 351 = reticulata.
polyschista, *Benj. in Mart. Fl. Bras.* x. 248.—Bras.
porphyrophylla, *Wright, ex Griseb. Cat. Pl. Cub.* 162.—Cuba.
praelonga, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 161.—Bras.; Guian.
prehensilis, *E. Mey. Comm. Pl. Afr. Austr.* 282.—Afr. austr.
Preissii, A. DC. in DC. Prod. viii. 666 = Hookeri.
protrusa, *Hook. f. Fl. N. Zel.* i. 206.—N. Zel.
pterosperma, Edgew. in Proc. Linn. Soc. i. (1847) 352 = exoleta.

UTRICULARIA:—
puberula, *Benj. in Mart. Fl. Bras.* x. 248.—Bras.
pubescens, *Sm. in Rees, Cycl.* xxxvii. n. 53.—Afr. trop.
pulchella, C. B. Lehm. in Flora, xxvi. (1843) 786 = minor.
pumila, *Benj. in Linnaea*, xx. (1847) 313.—Am. bor.
pumila, Walt. Fl. Carol. 64 = gibba, biflora.
punctata, *Wall. Cat.* n. 2121.—Burma; Malaya.
punctifolia, *Benj. in Linnaea*, xx. (1847) 493.—N. Granat.
purpurascens, J. Grah. Cat. Bomb. Pl. 165 = caerulea.
purpurea, *Walt. Fl. Carol.* 64.—Am. bor.
purpurea, Willd. ex Benj. in Linnaea, xx. (1847) 309 = paucifolia.
purpureo-caerulea, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 154.—Bras.
pusilla, J. Grah. Cat. Bomb. Pl. 165 = orbiculata.
pusilla, *Vahl, Enum.* i. 202.—Am. trop.
pygmaea, *R. Br. Prod.* 432.—Austral.
quinqueradiata, Spruce, ex Oliver, in Journ. Linn. Soc. iv. (1860) 171 = inflata.
racemosa, *Wall. Cat.* n. 1496.—As. trop.
racemosa, Wight, Ic. t. 1584. f. 1 = rosea.
ramosa, *Vahl, Enum.* i. 204.—Ind. or.
reclinata, *Hassk. in Versl. Kon. Akad. Wetensch.* iv. (1855) 161; *et Retzia*, i. 92.—Java.
recurva, *Lour. Fl. Cochinch.* 26.—Cochinch.
reflexa, *Oliver, in Journ. Linn. Soc.* ix. (1867) 146.—Afr. trop.
reniformis, *A. St. Hil. Voy. Minas Geraes*, i. 224.—Bras.
resupinata, *B. D. Greene, ex Hitchcock, Cat. Pl. Massach.* (1835) 107, nomen; *et in Bigel. Fl. Bost.* ed. III. 10.—Am. bor.
reticulata, *Sm. Exot. Bot.* ii. 119. t. 119.—Ind. or.
rhodocnemis, Mart. ex Benj. in Mart. Fl. Bras. x. 237 = oligosperma.
rigida, *Benj. in Linnaea*, xx. (1847) 303.—Afr. trop.
rosea, *Edgew. in Proc. Linn. Soc.* i. (1847) 352.—Ind. or.
rosulata, *Benj. in Linnaea*, xx. (1847) 310.—Ins. Philipp.
Roxburghii, Spreng. Syst. i. 52 = exoleta.
saccata, Le Conte, in Ell. Sketch, i. 21 = purpurea.
sacciformis, *Benj. in Linnaea*, xx. (1847) 302.—Ind. or.
saharunporensis, Royle, ex Oliver, in Journ. Linn. Soc. iii. (1859) 176 = exoleta.
Salzmanni, *A. St. Hil. & Girard, in Ann. Sc. Nat.* Sér. II. xi. (1839) 151.—Bras.
Sandersoni, *Oliver, in Journ. Linn. Soc.* ix. (1867) 155.—Afr. trop.
sanguinea, *Oliver, l. c.* 153.—Afr. trop.
scandens, *Benj. in Linnaea*, xx. (1847) 309, partim.—Ind. or.
scandens, Benj. l. c., partim = Wallichiana.
sclerocarpa, *Wright, in Sauv. Fl. Cub.* 91.—Cuba.
secunda, *Benj. in Linnaea*, xx. (1847) 308.—Mexic.
Selloi, *G. Weber, ex Benj. in Mart. Fl. Bras.* x. 250.—Bras.
setacea, Michx. Fl. Bor. Am. i. 12 = subulata.
setacea, Wall. Cat. n. 6398 = hirta, racemosa.
similis, Lehm. Pugill. viii. 666 = Hookeri.
simplex, *R. Br. Prod.* 431.—Austral.
simplex, *Wright & Sauv. Fl. Cub.* 91.—Ind. occ.
sinuata, *Benj. in Linnaea*, xx. (1847) 491.—Mexic.
Smithiana, Wight, in Hook. Kew Journ. i. (1849) 373; *et Ic.* t. 1577 = caerulea.
spartea, *Baker, in Journ. Linn. Soc.* xx. (1883) 216.—Madag.
spatulata, Splitg. ex De Vriese, in Nederl. Kruidk. Arch. i. (1848) 324 = amethystina.
speciosa, R. Br. Prod. 430 = dichotoma.
spectabilis, Madauss. ex Garcke, Fl. Deutschl. ed. III. 274 = major.
spiralis, *Sm. in Rees, Cycl.* xxxvii. n. 5.—Afr. trop.
spirandra, *Wright, ex Griseb. Cat. Pl. Cub.* 161.—Cuba.
spiricaulis, Miq. ex Oliver, in Journ. Linn. Soc. iii. (1859) 180 = reticulata.
Spruceana, *Benth. ex Oliver, l. c.* iv. (1860) 173.—Bras.
squamosa, *Benj. in Bot. Zeit.* iii. (1845) 212.—Nepal.

UTRICULARIA:—
squamosa, Wight, in Hook. Kew Journ. i. (1849) 373; et Ic. t. 1579 = caerulea.
stellaris, *Linn. f. Suppl.* 86.—As.; Austral.; Afr. trop.
stolonifera, *Benj. in Mart. Fl. Bras.* x. 245.—Bras.
striata, Le Conte, in Ann. Lyc. N. York, i. (1824) 75 = fibrosa.
striata, Tuckerm. in Am. Journ. Sc. Ser. I. xlv. (1843) 29 = clandestina.
striatula, *Sm. in Rees, Cycl.* xxxvii. n. 17.—Zeylan.
stricta, *G. F. W. Mey. Prim. Fl. Esseq.* 14.—Guian.
suaveolens, Afzel. ex Benj. in Linnaea, xx. (1847) 303 = rigida.
subsimilis, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 334.—N. Zel.
subulata, *Linn. Sp. Pl.* 18.—Am. bor. et trop.
sumatrana, *Miq. Fl. Ind. Bat.* ii. 998.—Sumatra.
superba, G. Weber, ex Benj. in Mart. Fl. Bras. x. 252 = Genlisea ornata.
tenella, R. Br. Prod. 432 = Polypompholyx tenella.
tenuifolia, *Benj. in Linnaea*, xx. (1847) 304.—Ind. occ.
tenuis, *Cav. Ic.* v. 24. t. 440. f. 2.—Chili.
tertia, Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat. Sér. II. xi. (1839) 158 = pusilla.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 12 = stellaris.
tortilis, *Welw. ex Oliver, in Journ. Linn. Soc.* ix. (1867) 150.—Afr. trop.
tremula, Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat. Sér. II. xi. (1839) 155 = subulata.
tribracteata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 18.—Abyss.
trichophylla, *Spruce, ex Oliver, in Journ. Linn. Soc.* iv. (1860) 173.—Am. trop.
tricolor, *A. St. Hil. Voy. Distr. Diam.* ii. 418.—Bras.
triloba, *Benj. in Mart. Fl. Bras.* x. 248.—Bras.
trinervia, *Benj. l. c.* 247.—Bras.
tubulata, *F. Muell. Fragm.* ix. 48.—Austral.
uliginoides, Wight, in Hook. Kew Journ. i. (1849) 372; et Ic. t. 1573 = caerulea.
uliginosa, Benth. in Hook. l. c. iv. (1852) 299 = caerulea.
uliginosa, Vahl, Enum. i. 203 = reticulata.
uniflora, R. Br. Prod. 431 = dichotoma.
uniflora, Spruce, ex Oliver, in Journ. Linn. Soc. iv. (1860) 173 = Spruceana.
unifolia, Ruiz & Pav. Fl. Per. i. 20. t. 31 = alpina.
vaga, *Griseb. Cat. Pl. Cub.* 161.—Cuba.
versicolor, Benj. in Linnaea, xx. (1847) 488 = amethystina.
verticillata, *Benj. l. c.* 312.—Malacca.
verticillata, Salzm. ex A. St. Hil. & Girard, in Ann. Sc. Nat. Sér. II. xi. (1839) 151 (*verticiliata*) = Salzmanni.
violacea, *R. Br. Prod.* 431.—Austral.
viscosa, *Spruce, ex Oliver, in Journ. Linn. Soc.* iv. (1860) 172.—Bras.
volubilis, *R. Br. Prod.* 430.—Austral.
volubilis, Hook. ex Benj. in Linnaea, xx. (1847) 309 = scandens.
vulgaris, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 84 = flexuosa.
vulgaris, *Linn. Sp. Pl.* 18.—Europ.; As. et Am. bor.
vulgaris, Vell. Fl. Flum. 18; i. t. 44 = oligosperma.
Wallichiana, Benj. in Bot. Zeit. iii. (1845) 213 = bifida.
Wallichiana, *Wight, Ic.* t. 1572/1.—Ind. or.
Wallichii, Wight, in Hook. Kew Journ. i. (1849) 372 = praec.
Welwitschii, *Oliver, in Journ. Linn. Soc.* ix. (1867) 152.—Afr. trop.

UVARIA, Linn. Diss. Dass. 21 (1747); Amoen. Acad. i. 404 (1749). *ANONACEAE*, Benth. & Hook. f. i. 23.
FITZALANIA, F. Muell. Fragm. iv. 33 (1863).
MARENTERIA, Thou. Gen. Nov. Madag. 18 (1806).
NARUM, Adans. Fam. ii. 365 (1763).
PYRAGMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.
WARIA, Aubl. Pl. Gui. i. 604. t. 243 (1775).
Abermoa, F. G. Dietr. Vollst. Lexik. Gaertn. x. 311 = Guatteria Abermoa.
acrantha, Miq. Fl. Ind. Bat. Suppl. 368 = ovalifolia.
acuminata, *Oliver, Fl. Trop. Afr.* i. 21.—Afr. trop.
acuminatissima, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 6—Borneo.

UVARIA:—
acuta, Teysm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 395 = Unona acuta.
aethiopica, Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 9 = Xylopia aethiopica.
Alphonsii, C. Muell. in Walp. Ann. iv. 67 = Unona Dasymaschala.
Amuyon, Blanco, Fl. Filip. ed. I. 463 = Melodorum fulgens.
amygdalina, *A. Gray, Bot. U. St. Expl. Exped.* i. 31. —Ins. Fiji.
angolensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 23.—Afr. trop.
argentea, *Blume, Fl. Jav. Anon.* 24.—Java.
aromatica, *Lam. Encyc.* i. 596 = Unona concolor.
Asterias, *S. Moore, in Journ. Bot.* xv. (1877) 289.—Afr. trop.
astrosticta, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 8.—Sumatra.
aurita, *Blume, Fl. Jav. Anon.* 15.—Java.
axillaris, Roxb. Hort. Beng. 94; Fl. Ind. ii. 667 = Cananga odorata.
Badajamba, Roxb. Hort. Beng. 43 (= *Guatteria Badajamba*).—Ind. or.
badiiflora, Hance, in Walp. Ann. ii. 19 = microcarpa.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 388.—Ins. Banca.
bicolor, Roxb. Hort. Beng. 94; Fl. Ind. ii. 662 = Melodorum bicolor.
bicolor, Wall. Cat. n. 6466 = Melodorum Wallichii.
Blumeana, Steud. Nom. ed. II. ii. 737 = hirsuta.
bracteata, *Roxb. Hort. Beng.* 43; Fl. Ind. ii. 660.—Ind. or.; Burma.
brasiliensis, Vell. Fl. Flum. 238; v. t. 122 (Cymbopetali sp.).—Bras.
Burahol, *Blume, Bijdr.* 14.—Java.
caffra, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 8.—Afr. austr.
calamistrata, *Hance, in Journ. Bot.* xx. (1882) 77.—China.
callicarpa, *Baill. Adansonia*, viii. (1867–68) 347.—Madag.
canangioides, Reichb. f. & Zoll. ex Miq. Fl. Ind. Bat. Suppl. 378 = Polyalthia Jenkinsii.
cauliflora, Wall. Cat. n. 6476 = Polyalthia cauliflora.
cerasoides, Roxb. Pl. Corom. i. 30. t. 33 = Polyalthia cerasoides.
ceylanica, Heynh. Nom. i. 832 = zeylanica.
Chamae, *Beauv. Fl. Owar.* ii. 43. t. 83. f. 2.—Afr. trop.
ciliata, Heyne, ex Wight & Arn. Prod. 10 = Miliusa indica.
clavata, Walp. Rep. i. 80 = Anaxagorea javanica.
Commersoniana, *Baill. Adansonia*, viii. (1867–68) 346.—Madag.
concava, *Teysm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 406.—Java.
connivens, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 465.—Afr. trop.
conoidea, Lem. Jard. Fleur. ii. (1852) Misc. 12 = Asimina triloba.
cordata, *Schum. & Thonn. Beskr. Guin. Pl.* 255.—Afr. trop.
cordata, Wall. Cat. n. 6486 = macrophylla.
cordifolia, Roxb. Hort. Beng. 94; Fl. Ind. ii. 662 = Unona discolor.
coriacea, Vahl, Symb. Bot. iii. 72 = zeylanica.
costata, Wall. Cat. n. 6480 = Polyalthia costata.
cristata, *R. Br. ex Oliver, Fl. Trop. Afr.* i. 23.—Afr. trop.
cylindrica, *Schum. & Thonn. Beskr. Guin. Pl.* 256.—Afr. trop.
cyrtopoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 3.—Ins. Amboin.
dasychlamys, *Miq. l. c.* 2.—Sumatra.
decandra, Ruiz & Pav. ex G. Don, Gen. Syst. i. 100 = Guatteria decandra.
dioica, Roxb. Hort. Beng. 94; Fl. Ind. ii. 659 = Miliusa Roxburghiana.
discolor, C. Muell. in Walp. Ann. iv. 67 = Unona Dasymaschala.
diversifolia, *Span. in Linnaea*, xv. (1841) 168.—Ins. Timor.
dulcis, *Dun. Mon. Anon.* 90. t. 13.—Burma; Malaya.
ebracteolata, *Presl, Rel. Haenk.* ii. 77.—Ins. Luzon.
elegans, Wall. Cat. n. 6474 A = Melodorum elegans.
elliptica, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 202 = Bocagea elliptica.

UVARIA:—
elongata, *Roxb. Hort. Beng.* 43, nomen (Quid?).—Ind. or.
esculenta, Roxb. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 201 = Artabotrys odoratissimus.
excelsa, Vahl, ex Dun. Monog. Anon. 132 (= *Oxandra laurifolia*).—Ind. occ.
excelsa, *Wall. Cat.* n. 6477.—Ind. or.; Malaya.
farcta, Wall. Cat. n. 6460 = Cananga odorata.
fasciculata, Steud. Nom. ed. II ii. 738 = Polyalthia simiarum.
febrifuga, Humb. & Bonpl. ex DC. Syst. i. 498 = Unona xylopioides.
ferruginea, *Buch.-Ham. ex Hook. f. & Thoms. Fl. Ind.* i. 96.—Burma.
ferruginea, Koen. ex Wall. Cat. n. 6485 D = purpurea.
flava, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 419.—Ins. Banca.
flexuosa, Heyne, ex Wall. Cat. n. 6473 B = U. Narum.
fluminensis, *Vell. Fl. Flum.* 238; v. t. 121.—Bras.
foetida, *Ruiz & Pav. ex G. Don, Gen. Syst.* i.93.—Peruv.
fornicata, Roxb. Hort. Beng. 94; Fl. Ind. ii. 662 = Oxymitra fornicata.
fulgens, Wall. Cat. n. 6482 = Melodorum fulgens.
fulva, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. i. 79 = Melodorum rubiginosum.
furfuracea, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 338.—Madag.
furfuracea, Walp. Rep. i. 79 = Unona furfuracea.
fusca, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 466.—Afr. trop.
Gaertneri, *DC. Syst.* i. 482.—Zeylan.
gamopetala, *Zoll. in Linnaea*, xxix. (1857–58) 310.—Java.
gigantea, Wall. Cat. n. 6469 A, B = Goniothalamus giganteus.
glabra, *Span. in Linnaea*, xv. (1841) 162.—Ins. Timor.
glauca, *Hassk. Cat. Hort. Bog. Alt.* 175.—Java?
globosa, *Hook. f. in Hook. Niger Fl.* 210.—Afr. trop.
Goezeana, *F. Muell. Fragm.* vii. 125.—Austral.
Gomeziana, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 203 = bracteata.
gracilis, *Hook. f. in Hook. Niger Fl.* 210.—Afr. trop.
grandiflora, Roxb. Hort. Beng. 43; Fl. Ind. ii. 665 = purpurea.
grandiflora, Wall. Cat. n. 6485 E = Hamiltonii.
guatterioides, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 202.—Guian.
Hahniana, *Baill. Adansonia*, viii. (1868) 347.—Mexic.
hamata, *Dun. Monog. Anon.* 106. t. 27.—Zeylan.
hamata, Roxb. Fl. Ind. ii. 666 = Artabotrys odoratissimus.
Hamiltonii, *Hook. f. & Thoms. Fl. Ind.* i. 96.—Reg. Himal.
Hasselti, *Blume, Fl. Jav. Anon.* 46.—Java.
heterocarpa, *Blume, l. c.* 41.—Java.
heteroclita, Roxb. Hort. Beng. 43; Fl. Ind. ii. 663 = Kadsura Roxburghiana.
heteropetala, *F. Muell. Fragm.* iii. 1.—Austral.
Heyneana, Wall. Cat. n. 6463 = Mitrephora Heyneana.
Heyneana, Wight & Arn. Prod. 8 = zeylanica.
Hiran, Buch.-Ham. ex Wall. Cat. n. 6443 B (= *Guatteria Badajamba*).—Ind. or.
hirsuta, *Jack, in Malay. Misc.* i. (1820) n. v. 46.—Malaya.
hirsuta, Vell. Fl. Flum. 238; v. t. 124 = Trigynaea oblongifolia.
Holtzei, *F. Muell. in Wing, South. Sc. Record*, iii. (July 1883).—Austral.
hortensis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 28 = Cananga odorata.
humilis, *Blume, Bijdr.* 17.—Java.
japonica, Linn. Sp. Pl. 536 = Kadsura japonica.
javana, Dun. Monog. Anon. 91. t. 14 = dulcis.
javanica, Steud. Nom. ed. II. ii. 738 = praec.
javanica, Thunb. Fl. Javan. 19 = Unona tripetaloidea.
Kirkii, *Oliver, ex Hook. f. Bot. Mag.* t. 6006.—Afr. trop.
Laha, Miq. Fl. Ind. Bat. Suppl. 369 = Ellipeia Laha.

UVARIA:—
lamponga, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 22.—Sumatra.
lanceolata, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 3.—Ins. Celebes.
lanceolata, Sw. Prod. Veg. Ind. Occ. 87 (= *Oxandra virgata*).—Ind. occ.
Lanotan, Blanco, Fl. Filip. ed. I. 464 = Goniothalamus giganteus.
lanuginosa, *Wall. Cat.* n. 6454.—Ind. or.
Larep, *Miq. Fl. Ind. Bat. Suppl.* 370.—Sumatra.
latifolia, Blume, Fl. Jav. Anon. 37. t. 15 = Melodorum latifolium.
laurifolia, Sw. Fl. Ind. Occ. ii. 1001 (= *Oxandra laurifolia*).—Ind. occ.
ligularis, Lam. Encyc. i. 597 = Unona ligularis.
littoralis, *Blume, Fl. Jav. Anon.* 26.—Ind. or.
Lobbiana, *Hook. f. & Thoms. Fl. Ind.* i. 100.—Malaya.
longiflora, Steud. Nom. ed. II. ii. 738, sphalm. = Polyalthia longifolia.
longifolia, Blume, Bijdr. 13 = Unona latifolia.
longifolia, Ruiz & Pav. ex G. Don, Gen. Syst. i. 100 = Guatteria Pavonii.
longifolia, Sonner. Voy. Indes, iii. 260 = Polyalthia longifolia.
lucida, *Boj. ex Sweet, Hort. Brit.* ed. II. 14; *Benth. in Trans. Linn. Soc.* xxiii. (1862) 465.—Afr. trop.
lurida, Dalz. & Gibs. Bomb. Fl. 3 = *U. Narum*, Wall.
lurida, *Hook. f. & Thoms. Fl. Ind.* i. 101.—Reg. Himal.
lutea, Roxb. Pl. Corom. i. 32. t. 36 = Alphonsea lutea.
lutea, Wall. Cat. n. 6462 = zeylanica.
lutea, Wight & Arn. Prod. 8 = Alphonsea zeylanica, Mitrephora Heyneana.
mabiformis, Griff. Notul. iv. 709 = Melodorum pisocarpum.
macrantha, *Hassk. Retzia*, i. 116.—Java.
macrocarpa, Vahl, ex DC. Syst. i. 489 = U. Chamae.
macrocarpa, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 142.—Bras.
macrophylla, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 663. —As. trop.
macropoda, *Hook. f. & Thoms. Fl. Ind.* i. 101.—Zeylan.
mangattensis, Dennst. Schluess. Hort. Malab. 31 = Alphonsea zeylanica.
manubriata, Wall. Cat. n. 6456 = Melodorum manubriatum.
Marenteria, *Baill. Adansonia*, viii. (1868) 304; *Hist. Pl.* i. 200.—Madag.
Melodorum, Raeusch. Nom. ed. III. 162 = Unona sylvatica.
membranacea, *Benth. Fl. Austral.* i. 51.—Austral.
micrantha, *Hook. f. & Thoms. Fl. Ind.* i. 103.—Burma; Malaya.
microcarpa, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 257.—China.
mollis, Wall. Cat. n. 6475 = Unona pannosa.
moluccana, Kostel. Allg. Med. Pharm. Fl. v. 1707 = Unona Musaria.
monilifera, Gaertn. Fruct. ii. 156 = Unona discolor, discreta.
monopetala, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 8. t. 2 = Hexalobus senegalensis.
monosperma, Lam. Encyc. i. 596 = Guatteria Ouregou.
monosperma, Vell. Fl. Flum. 238; v. t. 123 = Guatteria nigrescens.
montana, *Blume, Fl. Jav. Anon.* 45.—Java.
montana, Dun. ex Wall. Cat. sub n. 6462 = Alphonsea lutea.
montana, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 28 = montana.
monticola, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 5.—Borneo.
multiflora, *Teysm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 394.—Ins. Bantam.
Musaria, A. DC. Mém. Anon. 29 = Unona Musaria.
Narum, *A. DC. l. c.* 29.—Ind. or.
Narum, Wall. Cat. n. 9 (sp. div.).—Ind. or.
neglecta, *A. Rich. Ess. Fl. Cub.* 44.—Cuba.
nervosa, Wall. Cat. n. 6479 = Melodorum rubiginosum.
nicaraguensis, *Baill. Hist. Pl.* i. 195.—Nicaragua.
nitida, *Roxb. Hort. Beng.* 94; *Fl. Ind.* ii. 667.—Ins. Molucc.
nutans, Wall. Cat. n. 6481 = Phaeanthus nutans.

UVARIA:—
obovata, Heyne, ex Wall. Cat. n. 6471 = Goniothalamus cardiopetalus.
obovata, Torr. & Gray, Fl. N. Am. i. 45 = Asimina grandiflora.
obtusa, Blume, Bijdr. 13 = Mitrephora obtusa.
ochroleuca, *Zoll. in Linnaea*, xxix. (1856) 307.—Java.
odorata, Lam. Encyc. i. 595 = Cananga odorata.
odoratissima, Roxb. Hort. Beng. 43; Fl. Ind. ii. 666 = Artabotrys odoratissimus.
olens, Buch.-Ham. ex Wall. Cat. sub n. 6441 = Miliusa velutina.
ophthalmica, Roxb. ex G. Don, Gen. Syst. i. 93 = Phaeanthus nutans.
ossea, Blanco, Fl. Filip. ed. II. 322 = Unona discolor.
ovalifolia, *Blume, Fl. Jav. Anon.* 27.—Java.
ovata, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 205.—Afr. trop.
oxyantha, Wall. Cat. n. 6478 = Xylopia oxyantha.
paracaroënsis, Dennst. Schluess. Hort. Malab. 31 = zeylanica.
Parveana, Zoll. ex Miq. Fl. Ind. Bat. i. II. 48 (= *Guatteria Parveana*).—Sumatra.
parviflora, Hook. f. & Thoms. Fl. Ind. i. 103 (Cyathostemmatis sp.).—Ins. Penang.
parviflora, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 9. t. 3. f. 1 = Xylopia parviflora.
parviflora, Torr. & Gray, Fl. N. Am. i. 45 = Asimina parviflora.
pauciovulata, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 51.—Malacca.
pilosa, Roxb. Fl. Ind. ii. 665 = hirsuta.
platypetala, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 257 = purpurea.
polyantha, Wall. Cat. n. 6467 = Melodorum polyanthum.
polyprena, *Blume, Fl. Jav. Anon.* 35.—Java.
prismatica, Wall. Cat. n. 6455 = Melodorum prismaticum.
ptychocalyx, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 4.—Malaya.
purpurea, *Blume, Bijdr.* 11.—As. trop.
pygmaea, Torr. & Gray, Fl. N. Am. i. 45 = Asimina pygmaea.
pyrifolia, Reinw. ex Blume, Fl. Jav. Schizand. 16 (= *Sphaerostema pyrifolium*).—Java.
reticulata, Blume, Fl. Jav. Anon. 50. t. 24 = Mitrephora reticulata.
rhodantha, Hance, in Walp. Ann. ii. 19 = purpurea.
Rosenbergiana, Scheff. in Ann. Jard. Buitenz. i. (1876) 2 = ovalifolia?
Roxburghiana, Wall. Cat. n. 6423 A = Oxymitra fornicata.
rubiginosa, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 202 = Melodorum rubiginosum.
rufa, *Blume, Fl. Jav. Anon.* 19.—Java.
rufa, Wall. Cat. n. 6455 = Melodorum prismaticum.
rufescens, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 202 = macrophylla.
rugosa, *Blume, Bijdr.* 12.—Java.
Russelii, Wall. Cat. n. 6464 = Alphonsea lutea.
scabra, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 3.—Ins. Molucc.
scabrida, *Oliver, Fl. Trop. Afr.* i. 21.—Afr. trop.
sclerocarpa, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 203 = Saccopetalum sclerocarpum.
semecarpifolia, *Hook. f. & Thoms. Fl. Ind.* i. 97.—Malaya.
sesquipedalis, Colebr. ex Wall. Cat. sub n. 6446 = Goniothalamus sesquipedalis.
sessiliflora, *Zoll. in Linnaea*, xxix. (1857–58) 312.—Malaya?
sessilis, Vell. Fl. Flum. 238; v. t. 125 = Duguetia bracteosa.
setigera, Blanco, Fl. Filip. ed. II. 323 = purpurea.
simiarum, Buch.-Ham. ex Wall. Cat. sub n. 6440 = Polyalthia simiarum.
sinensis, Blanco, Fl. Filip. ed. I. 465 = Artabotrys odoratissimus.
solanifolia, *Presl, Rel. Haenk.* ii. 76.—Ins. Luzon.
sorzogonensis, *Presl, l. c.*—Ins. Luzon.
spectabilis, *DC. Syst.* i. 484.—Guian.
sphaerocarpa, *Blume, Bijdr.* 12.—Java.
sphenocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 98.—Zeylan.
subcordata, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 9.—Ins. Molucc.?

UVARIA:—
suberosa, Roxb. Pl. Corom. i. 31. t. 34 = Polyalthia suberosa.
subrepanda, *Wall. Cat.* n. 6483.—Singapore.
sumatrana, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 51.—Malaya.
sumatrana, Kurz, Andam. Rep. 29 = micrantha.
synsepala, *Miq. in Journ. Bot. Néerl.* i. (1861) 128.—China.
timoriensis, *Blume, Fl. Jav. Anon.* 21.—Ins. Timor.
tomentosa, Roxb. Pl. Corom. i. 31. t. 35 = Saccopetalum tomentosum.
tomentosa, Vahl, ex DC. Syst. i. 504 (= *Guatteria rufa*).—Ind. occ.
tomentosa, Wall. Cat. n. 6454 = Melodorum lanuginosum.
trichomalla, Blume, Fl. Jav. Anon. 42. t. 18 = hirsuta.
trifoliata, Gaertn. Fruct. ii. 156. t. 114. f. 2 = Gaertneri.
trigyna, *Mart. Fl. Bras.* xiii. 1. 40.—Bras.
triloba, Torr. & Gray, Fl. N. Am. i. 45 = Asimina triloba.
tripetala, Lam. Encyc. i. 597 = Unona tripetaloidea.
tripetala, Roxb. Hort. Beng. 43; Fl. Ind. ii. 667 = Phaeanthus nutans.
uncata, Lour. Fl. Cochinch. 349 = Artabotrys odoratissimus.
uncata, Vahl, ex DC. Syst. i. 492 = Unona discolor.
undulata, Lam. Encyc. i. 595 = Cananga adorata.
undulata, Roxb. Hort. Beng. 43 = Unona discolor.
Uregu, Raeusch. Nom. ed. III. 160 = Guatteria Ouregou.
velutina, Blume, Bijdr. 13 = hirsuta.
velutina, Dun. Monog. Anon. 91 = Miliusa velutina.
ventricosa, Roxb. Hort. Beng. 43; Fl. Ind. ii. 658 = Alphonsea ventricosa.
verrucosa, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 3.—Borneo.
villosa, Roxb. Hort. Beng. 43; Fl. Ind. ii. 664 = Miliusa velutina.
virgata, Blume, Fl. Jav. Anon. 43 = Cananga virgata.
virgata, Sw. Fl. Ind. Occ. ii. 999 (= *Oxandra virgata*).—Ind. occ.
viridiflora, *Ruiz & Pav. ex G. Don, Gen. Syst.* i. 100.—Peruv.
viridiflora, *Splitg. in Nederl. Kruidk. Arch.* i. (1849) 224.—Guian.
Vogelii, *Hook. f. in Hook. Ic. Pl.* t. 767.—Afr. trop.
zeylanica, Aubl. Pl. Gui. ii. 605. t. 243 = Unona concolor.
zeylanica, Deless. ex DC. Syst. i. 483 = dulcis.
zeylanica, Domb. ex DC. l. c. 504 = Guatteria eriopoda.
zeylanica, Lam. Encyc. i. 596 = U. Narum.
zeylanica, *Linn. Sp. Pl.* 536.—Ind. or.
Zippeliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 4.—Ins. Amboin.

UVA-URSI, Tourn. ex Moench, Meth. 470 (1794) = **Arctostaphylos**, Adans. (Ericac.).
alpina, S. F. Gray, Nat. Arr. Brit. Pl. ii. 401 = A. alpina.
buxifolia, S. F. Gray, l. c. 400 = A. Uva-ursi.
procumbens, Moench, Meth. 470 = A. Uva-ursi.

UVEDALIA, R. Br. Prod. 440 (1810) = **Mimulus**, Linn. (Scrophul.).
linearis, R. Br. l. c. = M. Uvedaliae.
nana, E. Mey. ex Benth. in DC. Prod. x. 412 = Torenia nana.

UVIRANDRA, Jaume St. Hil. Expos. i. 52 (1805) = OUVIRANDRA, Thou. = **Aponogeton**, Thunb. (Naiad.).

UVULANA, Rafin. in Am. Monthly Mag. (1811) 356, sphalm. = **Uvularia**, Linn. (Liliac.).
angulata, Rafin. l. c. 356, nomen (Quid?).—Am. bor.

UVULARIA, Linn. Gen. ed. I. 93 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 880.
Baimo, Rafin. Fl. Tellur. iv. 27 (1836).
Oakesia, S. Wats. in Proc. Am. Acad. xiv. (1879) 269.
Uvulana, Rafin. in Am. Monthly Mag. (1811) 356.
acutifolia, Rafin. in Am. Monthly Mag. (1817) 359, nomen (Quid?).—Am. bor.
amplexicaulis, Mill. Gard. Dict. ed. VIII. n. 1 = Streptopus distortus.

UVULARIA:—
amplexifolia, Linn. Sp. Pl. 304 = Streptopus distortus.
Betua, Buch.-Ham. ex D. Don, in Proc. Linn. Soc. i. (1839) 45 = Disporum calcaratum.
calcarata, Wall. Cat. n. 5087 = Disporum calcaratum.
chinensis, Ker-Gawl. in Bot. Mag. t. 916 = Disporum pullum.
cirrhosa, Thunb. Fl. Jap. 136. t. 2 = Fritillaria verticillata.
distorta, Pers. Syn. i. 360 = Streptopus distortus.
flava, Sm. Exot. Bot. i. 97. t. 50 = perfoliata.
floridana, *Chapm. Fl. S. U. St.* 487.—Florida.
grandiflora, *Sm. Exot. Bot.* i. 99. t. 51.—Am. bor.
grandifolia, Pursh, Fl. Am. Sept. i. 231, sphalm. = praec.
Hamiltoniana, Wall. Cat. n. 5088 = Disporum calcaratum.
hirta, Thunb. Fl. Jap. 135 = Tricyrtis hirta.
lanceolata, Ait. Hort. Kew. ed. I. i. 434 = perfoliata.
lanuginosa, Pers. Syn. i. 360 = Disporum lanuginosum.
Leschenaultiana, Wall. Cat. n. 5089 = Disporum Leschenaultianum.
multiflora, Reinw. ex Kunth, Enum. Pl. iv. 207 = Disporum pullum.
oppositifolia, Schlecht. ex Kunth, l. c. 254 = Fritillaria camtchatcensis.
parviflora, Wall. in As. Res. xiii. (1820) 378 = Disporum pullum.
perfoliata, *Linn. Sp. Pl.* 304.—Am. bor.
Pitsutu, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 50 = Disporum pullum.
puberula, *Michx. Fl. Bor. Am.* i. 199.—Am. bor.
puberula, Sm. in Rees, Cycl. xxxvii. n. 5 = Disporum Menziesii.
rosea, Pers. Syn. i. 360 = Streptopus roseus.
sessiliflora, J. F. Gmel. Syst. 1642, sphalm. = seq.
sessilifolia, *Linn. Sp. Pl.* 304.—Am. bor.
sessilis, Thunb. Fl. Jap. 135 = Disporum sessile.
Smithii, Hook. Fl. Bor. Am. ii. 174. t. 189 = Disporum Menziesii.
umbellata, Wall. in As. Res. xiii. (1820) 379 = Disporum pullum.
viridescens, Maxim. Prim. Fl. Amur. 273 = Disporum smilacinum.

UWAROWIA, Bunge, in Bull. Sc. Acad. Pétersb. vii. (1840) 278 = **Verbena**, Linn.
chrysanthemifolia, Bunge, l. c. 279 = V. sulphurea.
sulphurea, Schau. in DC. Prod. xi. 551 = V. sulphurea.

V

VACCARIA, Medic. Phil. Bot. i. 96 (1789) = **Saponaria**, Linn. (Caryoph.).
arvensis, Link, Handb. ii. 240 = S. Vaccaria.
grandiflora, Jaub. & Spach, Illustr. iii. 40. t. 231 = S. Vaccaria.
inclusa, Ledeb. Fl. Ross. i. 776 = S. Vaccaria?
oxyodonta, Boiss. Diagn. Ser. II. i. 68 = S. Vaccaria.
parviflora, Moench, Meth. 63 = S. Vaccaria.
perfoliata, Sweet, Hort. Brit. ed. II. 51 = S. Vaccaria.
pyramidata, Medic. Phil. Bot. i. 96 = S. Vaccaria.
segetalis, Garke, ex Aschers. Fl. Brandenb. i. 84 = S. Vaccaria.
sessiliflora, Steud. Nom. ed. II. ii. 739, sphalm. = seq.
sessilifolia, Sweet, Hort. Brit. ed. II. 51 = S. Vaccaria.
vulgaris, Host, Fl. Austr. i. 518 = S. Vaccaria.

VACCINIUM, Linn. Syst. ed. I (1735). *VACCINIACEAE*, Benth. & Hook. f. ii. 573.
Acosta, Lour. Fl. Cochinch. 276 (1790).
Batodendron, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 261.
Cavinium, Thou. Gen. Nov. Madag. 11 (1806).
Epigynium, Klotzsch, in Linnaea, xxiv. (1851) 49.
Metagonia, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 263.
Picrococcus, Nutt. l. c. 262 (1843).
Polycodium, Rafin. in Am. Monthly Mag. (1818) 266.
Vitis-Idaea, Tourn. ex Moench, Meth. 47 (1794).
Acosta, *Raeusch. Nom.* ed. III. 109 (sp. dub.).—Cochinch.

VACCINIUM:—
acuminatissimum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 36.—Malaya.
acuminatum, Benth. Pl. Hartw. 222 = staphelioides.
acuminatum, *H. B. & K. Nov. Gen. et Sp.* iii. 264.—N. Granat.
acuminatum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 214 = Corallobotrys acuminata.
adenandrum, Decne. Bot. Voy. Vénus, 23 = cereum.
affine, *Klotzsch, in Linnaea*, xxiv. (1851) 61.—Peruv.
affine, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 173 = Donianum.
agathosmoides, *Wedd. Chlor. And.* ii. 179.—N. Granat.
alaternoides, *H. B. & K. Nov. Gen. et Sp.* iii. 265.—Peruv.
alaternoides, Soland. ex Seem. Fl. Vit. 146 = cereum.
alatum, Domb. ex Dun. in DC. Prod. vii. 562 (= *Proclesia alata*).—Peruv.
albiflorum, Hook. Bot. Mag. t. 3428 = corymbosum.
album, Lam. Encyc. i. 73 = canadense.
album, Linn. Sp. Pl. 350 = Lonicera ciliata.
album, Pursh, Fl. Am. Sept. i. 285 = stamineum.
amoenum, Ait. Hort. Kew. ed. I. ii. 12 = corymbosum.
angulosum, Dulac, Fl. Hautes-Pyr. 460 = V. Myrtillus.
angustifolium, Ait. Hort. Kew. ed. I. ii. 11 = pensylvanicum.
angustifolium, *Benth. Pl. Hartw.* 45.—Mexic.
arboreum, Lesch. ex Dun. in DC. Prod. vii. 555 = Leschenaultii.
arboreum, *Marsh. Arbust. Am.* 157.—Am. bor.
arbutoides, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 453.—Reg. Himal.
Arctostaphylos, *Linn. Sp. Pl.* 351.—Armen.; Reg. Cauc.
Arctostaphylos, Willd. Sp. Pl. ii. 353 = maderense.
ardisioides, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 452.—Burma.
assimile, *Wright, in Sauv. Fl. Cub.* 85.—Cuba.
attenuatum, *Dun. in DC. Prod.* vii. 571.—Peruv.
auriculatum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 214 = Agapetes auriculata.
bancanum, *Miq. Fl. Ind. Bat. Suppl.* 587.—Burma; Malaya.
bicolor, F. Muell. Sel. Pl. Indust. Cult. 249 = Psammisia bicolor.
brachycerum, Michx. Fl. Bor. Am. i. 234 = Gaylussacia brachycera.
brachystachyum, Benth. Pl. Hartw. 65 = confertum.
bracteatum, *Thunb. Fl. Jap.* 156.—Japon.
brasiliense, Spreng. Nov. Prov. 42 = Gaylussacia Pseudo-vaccinium.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 29.—Japon.
buxifolium, Gilib. Fl. Lituan. i. 4 = V. Vitis-Idaea.
buxifolium, *Hook. f. in Hook. Ic. Pl.* t. 891.—Borneo.
buxifolium, Salisb. Parad. Lond. t. 4 = Gaylussacia brachycera.
caespitosum, *Michx. Fl. Bor. Am.* i. 234.—Am. bor.
callosum, Griseb. ex Wedd. Chlor. And. ii. 177 = floribundum.
calycinum, Sm. in Rees, Cyclop. xxxvi. n. 7 = penduliflorum.
campanulatum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 85 = Agapetes campanulata.
canadense, *Kalm, ex Richards. in Frankl. 1st Journ.* ed. II. 12.—Am. bor.
cantabricum, Huds. Fl. Angl. ed. I. 143 = Daboëcia polifolia.
caracasanum, *H. B. & K. Nov. Gen. et Sp.* iii. 266.—Venezuela.
caracasanum, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 530 = leucanthum.
carnosum, Pers. Syn. i. 479 = crassifolium.
cereum, *Forst. f. Prod.* 28.—Ins. Amicor.
cereum, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 360 = penduliflorum.
Chamissonis, Bong. in Mém. Acad. Pétersb. ii. (1833) 151 = ovalifolium.
chinense, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 297 = bracteatum.
ciliatum, Gilib. Exercit. i. 11 = uliginosum.
ciliatum, *Thunb. Fl. Jap.* 156.—Japon.
confertum, *H. B. & K. Nov. Gen. et Sp.* iii. 265.—Mexic.
consanguineum, *Klotzsch, in Linnaea*, xxiv. (1851) 64.—Panama.

VACCINIUM:—
Constablaei, A. Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 42 = corymbosum.
cordatum, *Hemsl. Biol. Centr. Am. Bot.* ii. 274.—Mexic.
coriaceum, *Hook. f. in Hook. Ic. Pl.* t. 892.—Borneo.
coriaceum, Miq. Fl. Ind. Bat. ii. 1062 = Korthalsii.
corymbiferum, *Miq. l. c. Suppl.* 588.—Sumatra.
corymbodendron, *Dun. in DC. Prod.* vii. 569.—Peruv.
corymbosum, Frank, ex Dun. l. c. 566 = Gaylussacia frondosa.
corymbosum, *Linn. Sp. Pl.* 350.—Am. bor.
crassifolium, *Andr. Bot. Rep.* t. 105.—Am. bor.
crenulatum, *Dun. in DC. Prod.* vii. 570.—Peruv.
croceum, Steud. Nom. ed. II. ii. 739, in syn., sphalm. = cereum.
cubense, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 171.—Cuba.
cuneifolium, *Miq. Fl. Ind. Bat.* ii. 1062.—Java.
cuspidatum, *Planch. ex Wedd. Chlor. And.* ii. 180, in obs.—N. Granat.
cylindraceum, *Sm. in Rees, Cyclop.* xxxvi. n. 23.—Ins. Azores.
cyrtodon, *Miq. Fl. Ind. Bat.* ii. 1062.—Java.
decamerocarpon, Dun. in DC. Prod. vii. 566, (excl. syn.) = Gaylussacia frondosa.
dempoënse, *Fawcett, in Journ. Bot.* xxiii. (1885) 254.—Sumatra.
dendrophilum, *Benth. Pl. Hartw.* 219.—Ecuador.
densiflorum, *Benth. l. c.* 222.—N. Granat.
dentatum, Sm. in Rees, Cyclop. xxxvi. n. 31 = cereum.
dependens, Ruiz & Pav. ex G. Don, Gen. Syst. iii. 859 = Gaylussacia dependens.
dichotomum, Boj. ex Dun. in DC. Prod. vii. 570 = secundiflorum.
didymanthum, *Dun. l. c.* 575.—Peruv.
diffusum, Ait. Hort. Kew. ed. I. ii. 11 = arboreum.
discolor, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 531.—Mexic.
disocarpum, A. Gray, Syn. Fl. N. Am. ii. 1. 23 = corymbosum.
disomorphum, Michx. Fl. Bor. Am. i. 231 = corymbosum.
Donianum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 161 = bracteatum.
Donianum, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 174.—Reg. Himal.
dumosum, Andr. Bot. Rep. t. 112 = Gaylussacia dumosa.
Dunalianum, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 175 (Dunallianum).—Reg. Himal.
elevatum, Banks & Soland. ex Wikstr. in Vet. Akad. Handl. Stockh. (1823) 281 = stamineum.
Elliotii, Chapm. Fl. S. U. St. 260 = virgatum.
ellipticum, *Miq. Fl. Ind. Bat.* ii. 1060.—Java.
elongatum, Wats. Dendrol. Brit. t. 125 B = corymbosum.
emirnense, *Hook. Ic. Pl.* t. 131.—Madag.
empetrifolium, H. B. & K. Nov. Gen. et Sp. iii. 263 = penaeoides.
epacridifolium, Benth. Pl. Hartw. 221 = penaeoides.
eriocladum, *Dun. in DC. Prod.* vii. 571.—Mexic.
erythrinum, *Hook. Bot. Mag.* t. 4688.—Java.
erythrocarpum, *Michx. Fl. Bor. Am.* i. 227.—Am. bor.
exaristatum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 86 = bancanum.
fasciculatum, *Boj. ex Dun. in DC. Prod.* vii. 570.—Madag.
fissum, *Schrank, in Denkschr. Bot. Ges. Regensb.* i. II. (1818) 15.—Labrador.
floribundum, *H. B. & K. Nov. Gen. et Sp.* iii. 266.—Peruv.
floribundum, Miq. Fl. Ind. Bat. ii. 1060 = Teysmanni.
floridanum, Schwein. ex Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 262 = stamineum.
Forbesii, Fawcett, in H. O. Forbes, Nat. Wand. East. Archip. 506 = dempoënse.
Forbesii, Hook. Ic. Pl. t. 345 = emirnense.
formosum, *Andr. Bot. Rep.* t. 97.—Am. bor.
fragrans, Wall. ex Voigt, Hort. Suburb. Calc. 333 = Diospyros vaccinioides.
frondosum, Ait. Hort. Kew. ed. I. ii. 11 = Gaylussacia dumosa.
frondosum, Linn. Sp. Pl. 351 = Gaylussacia frondosa.
fuscatum, Ait. Hort. Kew. ed. I. ii. 11 = corymbosum.

VACCINIUM:—
galegiformis, C. Koch, Dendrol. ii. 99, sphalm. = *galiformis*.
galezans, Michx. Fl. Bor. Am. i. 232 = virgatum.
galiformis, Sm. in Rees, Cyclop. xxxvi. n. 16 = virgatum.
geminiflorum, *H. B. & K. Nov. Gen. et Sp.* iii. 267.—Mexic.
gilvum, *Meerb. Pl. Sel.* t. 18.—Hab.?
glabrum, Wats. Dendrol. Brit. t. 125 D = Gaylussacia resinosa?
glauco-album, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 453.—Reg. Himal.
glaucum, Lam. Encyc. i. 73; Michx. Fl. Bor. Am. i. 231 = Gaylussacia frondosa.
grandiflorum, Herb. Par. ex Dun. in DC. Prod. vii. 553 = Ceratostema grandiflorum.
grandiflorum, Wats. Dendrol. t. 125 A = corymbosum.
Griffithianum, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 174.—Reg. Himal.; China.
guadalupense, Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1827) 63 = Hornemannia martinicensis.
gualtheriaefolium, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 453.—Reg. Himal.
gualtherioides, Bigel. in New Engl. Med. Journ. v. (1816) 335 = uliginosum.
halleriaefolium, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 156, nomen.—Am. bor.
Hasseltii, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 40.—Java; Sumatra.
hatamense, *Becc. Malesia*, i. (1878) 210.—N. Guin.
hirsutum, *Buckl. in Am. Journ. Sc.* Ser. I. xlv. (1843) 175.—Am. bor.
hirsutum, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 170 = Agapetes setigera.
hirtellum, Ait. Hort. Kew. ed. II. ii. 357 = Gaylussacia dumosa.
hirtum, *Thunb. Fl. Jap.* 155.—Japon.
hispidulum, Linn. Sp. Pl. 352 = Chiogenes serpyllifolia.
hispidulum, Wangenh. Beitr. Nordamer. Holzart. 188. t. 30. f. 67 = Oxycoccos macrocarpus.
Humboldtii, *Klotzsch, in Linnaea*, xxiv. (1851) 57.—N. Granat.
humifusum, R. Grah. in Edinb. N. Phil. Journ. (Apr.–June, 1831) 193 = Gaultheria Myrsinites.
humile, Willd. Enum. Hort. Berol. Suppl. 21 = pensylvanicum.
Idsuroei, *Franch. & Sav. Enum. Pl. Jap.* ii. 425.—Japon.
Imrayi, Hook. Ic. Pl. t. 292 = Hornemannia martinicensis.
intermedium ×, Ruthe, Fl. Brandenb. 377. t. 1.—Europ.
iteophyllum, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 223.—China.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 28.—Japon.
javanicum, *Hook. Ic. Pl.* t. 740.—Java.
jesoënse, Miq. Ann. Mus. Bot. Lugd. Bat. i. 28 = V. Vitis-Idaea.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 40.—Java.
Kruhsianum, Fisch. ex Herd. in Act. Hort. Petrop. i. (1871–72) 320 = uliginosum.
Kunthianum, *Klotzsch, in Linnaea*, xxiv. (1851) 56.—Mexic.
laevigatum, *Boj. ex Dun. in DC. Prod.* vii. 571.—Madag.
lanceolatum, DC. ex Dun. l. c. 570 = ovatum.
latifolium, *Benth. & Hook. f. Gen.* ii. 575.—Ind. occ.
latifolium, Hort. Angl. ex Steud. Nom. ed. II. ii. 739 = virgatum.
laurifolium, *Miq. Fl. Ind. Bat.* ii. 1061.—Java.
ledifolium, Pohl, Pl. Bras. Ic. ii. 36. t. 123 = Gaylussacia ledifolia.
Lenneanum, *Warsz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292, nomen.—Bolivia.
leptanthum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 37.—Java; Sumatra.
Leschenaultii, *Wight, Ic.* t. 1188.—Ind. or.
leucanthum, *Schlecht. in Linnaea*, viii. (1833) 524.—Mexic.
leucostomum, *Lindl. in Gard. Chron.* (1848) 7.—Peruv.
ligustrinum, Linn. Sp. Pl. 351 = Lyonia paniculata.
ligustrinum, Michx. Fl. Bor. Am. i. 233; Pursh, Fl. Am. Sept. i. 288 = virgatum.

VACCINIUM:—
littoreum, *Miq. Fl. Ind. Bat. Suppl.* 587.—Sumatra.
longeracemosum, *Franch. & Sav. Enum. Pl. Jap.* ii. 425.—Japon.
longiflorum, *Wikstr. in Vet. Akad. Handl. Stockh.* (1823) 278.—Ins. Azor.
loranthiflorum, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 446 = Agapetes loranthiflora.
lucidum, Miq. Fl. Ind. Bat. ii. 1061 = Rollisoni.
Macgillivrayi, *Seem. in Journ. Bot.* ii. (1864) 77.—N. Hebrid.
Macraeanum, Klotzsch, in Linnaea, xxiv. (1851) 59 = cereum.
macrocarpon, Ait. Hort. Kew. ed. I. ii. 13 = Oxycoccus macrocarpus.
macrophyllum, Domb. ex Dun. in DC. Prod. vii. 564 = Thibaudia punctatifolia.
macrostemon, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 85 = Agapetes macrostemon.
maderense, *Link, Enum. Hort. Berol.* i. 375.—Ins. Mader.
malaccense, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 172.—Malaya.
marginatum, *Dun. in DC. Prod.* vii. 570.—Peruv.
marianum, Wats. Dendrol. Brit. t. 124 = corymbosum.
marylandicum, Hort. Angl. ex Steud. Nom. ed. II. ii. 739 = corymbosum.
melliflorum, F. Muell. Sel. Pl. Indust. Cult. 249, in obs. = Eurygania melliflora.
membranaceum, Dougl. ex Hook. Fl. Bor. Am. ii. 32 = myrtilloides.
meridionale, *Sw. Prod. Veg. Ind. Occ.* 62.—Jamaic.
Merkii, Fisch. ex Herd. in Act. Hort. Petrop. i. (1871–72) 323 = Arctostaphylos alpina.
Meyenianum, Klotzsch, in Linnaea, xxiv. (1851) 59 = penduliflorum.
micranthum, *Dun. in DC. Prod.* vii. 568.—Mexic.
microcarpum, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 334, err. typ. = Oxycoccus palustris.
microphyllum, Reinw. ex Blume, Bijdr. 851 = Diplycosia microphylla.
miniatum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 85 = Agapetes miniata.
minutiflorum, Wats. Dendrol. Brit. t. 125 C = pensylvanicum.
Miquelii, *Boerl. in Veth, Midden-Sumatra*, iv. *Nat. Hist.* 22.—Sumatra.
montanum, *Pohl, Pl. Bras. Ic.* ii. 38. t. 125.—Bras.
Moritzianum, *Klotzsch, in Linnaea*, xxiv. (1851) 63.—N. Granat.
Mortinia, *Benth. Pl. Hartw.* 221.—Ecuador.
mucronatum, Linn. Sp. Pl. 350 = Nemopanthus fascicularis.
multiflorum, Dun. in DC. Prod. vii. 572 = pensylvanicum.
muscicola, *Hook. Ic. Pl.* t. 717.—Ecuador.
Myrsinites, *Lam. Encyc.* i. 73.—Am. bor.
myrtifolium, Michx. Fl. Bor. Am. i. 229 = crassifolium.
myrtifolium, Willd. ex Klotzsch, in Linnaea, xxiv. (1851) 58 = Humboldtii.
myrtilloides, Ell. Sketch, i. 500 = virgatum.
myrtilloides, *Hook. Fl. Bor. Am.* ii. 32, partim; *A. Gray, Man. Bot. N. U. St.* ed. V. 291.—Am. bor.
myrtilloides, Michx. Fl. Bor. Am. i. 234 = pensylvanicum.
myrtilloides, S. Wats. Bot. King's Exped. 209 = V. Myrtillus.
myrtillum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = V. Myrtillus.
Myrtillus, Cham. & Schlecht. in Linnaea, i. (1826) 525 = ovalifolium.
Myrtillus, *Linn. Sp. Pl.* 349.—Reg. bor. temp. et arct.
myrtoides, *Miq. Fl. Ind. Bat.* ii. 1062.—Ins. Molucc.; Sumatra.
neilgherrense, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 173.—Ind. or.
nemorosum, Salisb. Prod. 291 = V. Vitis-Idaea.
nigrum, Desf. Tabl. ed. II. 274, nomen (Quid?).—Hab. ?
nitidum, *Andr. Bot. Rep.* t. 480.—Am. bor.
Nummularia, *Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 451.—Reg. Himal.
oblongifolium, Hort. ex Dun. in DC. Prod. vii. 576 = Oxycoccos macrocarpus.
obovatum, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 174 = Agapetes obovata.

VACCINIUM:—
obtusum, Pursh, Fl. Am. Sept. i. 290 = Gaultheria Myrsinites?
obversum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 220.—Java.
occidentale, *A. Gray, in Bot. Calif.* i. 451.—Calif.
odontocerum, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 214 = Agapetes angulata.
odontocerum, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 173 = Agapetes variegata.
Oldhami, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 161 = bracteatum, ciliatum.
orientale, Sw. in Weber & Mohr, Beitr. i. 6 = V. Acosta.
Ottonis, Klotzsch, in Linnaea, xxiv. (1851) 61 = floribundum.
ovalifolium, *Sm. in Rees, Cyclop.* xxxvi. n. 2.—Am. bor. occ.
ovatum, *Pursh, Fl. Am. Sept.* i. 290.—Am. bor. occ.
Oxycoccos, Linn. Sp. Pl. 351 = Oxycoccos palustris.
pachyphyllum, *Hemsl. Biol. Centr. Am. Bot.* ii. 275.—Costa Rica.
padifolium, Hochst. ex Steud. Nom. ed. II. ii. 740 = longiflorum.
padifolium, Sm. in Rees, Cycl. xxxvi. n. 22 = maderense.
pallidum, Ait. Hort. Kew. ed. I. ii. 10 = corymbosum.
palustre, Salisb. Prod. 291 = Oxycoccos palustris.
paradisearum, *Becc. Malesia*, i. (1878) 209.—N. Guin.
parviflorum, Andr. Bot. Rep. t. 125 = Gaylussacia resinosa.
parvifolium, *Sm. in Rees, Cyclop.* xxxvi. n. 3.—Am. bor. occ.
penaeoides, *H. B. & K. Nov. Gen. et Sp.* iii. 264.—Ecuador.
penduliflorum, *Gaudich. in Freyc. Voy. Bot.* i. 454. t. 68.—Ins. Sandvic.
pennsylvanicum, Torr. Fl. N. U. St. 416 = vacillans.
pensylvanicum, *Lam. Encyc.* i. 74 (pensilvanicum).—Am. bor.
pensylvanicum, Mill. Gard. Dict. ed. VIII. n. 3 = Royena glabra.
pernettyoides, *Griseb. ex Wedd. Chlor. And.* ii. 179.—Peruv.
polystachyum, Benth. Pl. Hartw. 140 = floribundum.
praestans, *Lamb. in Trans. Linn. Soc.* x. (1811) 264. t. 9.—Kamschat.
propinquum, Salisb. Prod. 291 = Oxycoccus macrocarpus.
prunifolium, Lodd. Cat. ex Loud. Hort. Brit. 156; et Loud. Encyc. Pl. 1546 = ovatum.
puberulum, *Klotzsch, ex Meissn. in Mart. Fl. Bras.* vii. 128.—Guian.
pubescens, Wormsk. ex Hornem. in Fl. Dan. t. 1516 = uliginosum.
pulchellum, Fisch. ex Herd. in Act. Hort. Petrop. i. (1871–72) 313 = V. Vitis-Idaea.
pumilum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 85.—Burma.
punctatum, Lam. Fl. Fr. iii. 396 = V. Vitis-Idaea.
punctifolium, Stokes, Bot. Mat. Med. ii. 363 = V. Vitis-Idaea.
Ramonii, *Griseb. Cat. Pl. Cub.* 52.—Cuba.
ramosissimum, *Dun. in DC. Prod.* vii. 571.—Peruv.
ramulosum, Willd. Enum. Hort. Berol. Suppl. 20 = pensylvanicum.
reflexum, *Hook. f. Bot. Mag.* t. 5781.—Bolivia.
reflexum, *Klotzsch, in Linnaea*, xxiv. (1851) 60.—Madag.
resinosum, Ait. Hort. Kew. ed. I. ii. 12 = Gaylussacia resinosa.
reticulatum, Sm. in Rees, Cyclop. xxxvi. n. 30 = cereum.
retusum, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 451.—Reg. Himal.
Rollisoni, *Hook. Bot. Mag.* t. 4612.—Java.
rotundifolium, Wight, Illustr. Ind. Bot. ii. 119. t. 139 = Leschenaultii.
Roylei, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 214 = Agapetes setigera.
rubrum, Dulac, Fl. Hautes-Pyr. 460 = V. Vitis-Idaea.
rubrum, Gilib. Fl. Lituan. i. 4 = uliginosum.
rugosum, Hook. f. & Thoms. ex Hook. Bot. Mag. sub t. 5198 = Pentapterygium rugosum.
rugosum, Hort. Angl. ex Steud. Nom. ed. II. ii. 740, nomen (Quid?).—Hab. ?
salicifolium, *Steud. Nom.* ed. II. ii. 740.—Am. bor.

VACCINIUM:—
salicinum, Aschers. in Flora, xliii. (1860) 370 = pensylvanicum.
salicinum, *Cham. & Schlecht. in Linnaea*, i. (1826) 525.—Unalaschk.
salicinum, Lodd. ex Loud. Hort. Brit. 156 = salicifolium.
salignum, Hook. f. Illustr. Himal. Pl. t. 15 A = Agapetes saligna.
salvifolium, Hort. Angl. ex Steud. Nom. ed. II. ii. 740, nomen (Quid?).—Hab. ?
scabrum, Pohl, Pl. Bras. Ic. ii. 37. t. 124 = Gaylussacia Pseudo-gaultheria.
Schlechtendalii, *G. Don, Gen. Syst.* iii. 856.—Mexic.
secundiflorum, *Hook. Ic. Pl.* t. 134.—Madag.
secundum, *Klotzsch, in Linnaea*, xxiv. (1851) 62.—Am. centr.
sednense, Pers. Syn. i. 478 = uliginosum.
serpens, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 171 = Pentapterygium serpens.
serratum, *Wight, l. c.*—Reg. Himal.
sibiricum, Hort. ex Steud. Nom. ed. II. ii. 740, nomen (Quid?).—Sibir.
Sieboldi, Miq. Ann. Mus. Bot. Lugd. Bat. i. 29 = ciliatum.
sikkimense, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 451.—Reg. Himal.
Smallii, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 398 = hirtum.
smilacifolium, Griseb. Fl. Brit. W. Ind. 144 = Hornemannia smilacifolia.
spicatum, Poir. Encyc. Suppl. i. 280 = V. Acosta.
splendens, Rupr. in Beitr. Pfl. Russ. Reich. iv. 56 = uliginosum.
Sprengelii, Hort.—Cf. Gard. Chron. (1883) I. 473 = V. Myrsinites.
Sprengelii, Wall. Cat. n. 6296 = Agapetes obovata.
stamineum, *Linn. Sp. Pl.* 350.—Am. bor.
staphelioides, *Planch. ex Wedd. Chlor. And.* ii. 180.—N. Granat.
stenophyllum, Steud. Nom. ed. II. ii. 740 = angustifolium.
suaveolens, Boj. ex Dun. in DC. Prod. vii. 570 = emirnense.
subcrenulatum, *Klotzsch, ex Meissn. in Mart. Fl. Bras.* vii. 128.—Guian.
sumatranum, *Jack, in Malay. Misc.* ii. (1822) n. VII. 18.—Sumatra.
tenellum, Ait. Hort. Kew. ed. I. ii. 12 = virgatum.
tenellum, Pursh, Fl. Am. Sept. i. 288 = pensylvanicum.
terniflorum, *Dun. in DC. Prod.* vii. 575.—Peruv.
Teysmanni, *Miq. Fl. Ind. Bat.* ii. 1062.—Java.
Thibautii, *Regel, Gartenfl.* xv. (1866) 357; *et in Ind. Sem. Hort. Petrop.* 1866 (1867) 96.—Am. trop.
thymifolium, *Klotzsch, in Linnaea*, xxiv. (1851) 62.—N. Granat.
timorense, *Fawcett, in H. O. Forbes, Nat. Wand. East. Archip.* 509.—Ins. Timor.
tomentosum, Pursh, ex A. Gray, Syn. Fl. N. Am. ii. 1. 19 = Gaylussacia frondosa.
trinerve, Domb. ex Dun. in DC. Prod. vii. 563 (= *Proclesia bracteata*).—Peruv.
uliginosum, *Linn. Sp. Pl.* 350.—Reg. bor. et arct.
ursinum, M. A. Curt. in Am. Journ. Sc. Ser. I. xliv. (1843) 82 = Gaylussacia ursina.
vacillans, *Soland. ex A. Gray, Man. Bot. N. U. St.* ed. I. 261.—Am. bor.
variegatum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 84 = Agapetes macrantha.
varingiaefolium, *Miq. Fl. Ind. Bat.* ii. 1061.—Java.
venosum, *Wight, in Calc. Journ. Nat. Hist.* viii. (1847) 172.—Reg. Himal.
venustum, Ait. Hort. Kew. ed. I. ii. 11 = Gaylussacia frondosa.
verticillatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 83.—Burma.
verticillatum, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 170 = Agapetes setigera.
villosum, *Sm. in Rees, Cyclop.* xxxvi. n. 29.—Mexic.
virgatum, *Ait. Hort. Kew.* ed. I. ii. 12.—Am. bor.
virgatum, Bigel. Fl. Bost. ed. II. 152 = vacillans.
virgineum, J. F. Gmel. Syst. 1643 (Index) = Oxycoccus palustris?
Vitis-Idaea, *Linn. Sp. Pl.* 351.—Reg. bor. et arct.
Wallichianum, Wight, in Calc. Journ. Nat. Hist. viii. (1847) 171 = Agapetes setigera.
Watsoni, Sweet, Hort. Brit. ed. II. 330 = Gaylussacia dumosa.

VACCINIUM:—
Wrightii, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 398 = bracteatum.
Wrightii, Griseb. Cat. Pl. Cub. 52 = assimile.
Zollingeri, Miq. Fl. Ind. Bat. ii. 1061 = laurifolium.

VACHELLIA, Wight & Arn. Prod. 272 (1834) = **Acacia**, Tourn. (Legumin.).
Farnesiana, Wight & Arn. l. c. = A. Farnesiana.

VACHENDORFIA, Adans. Fam. ii. 47 (1763) = **Wachendorfia**, Burm. f. (Haemod.).

VADA-KODI, Adans. Fam. ii. 701 (1763) = **Justicia**, Houst. (Acanth.).

VAGARIA, Herb. Amaryll. 226 (1837). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 735.
VAGINARIA, Kunth, Enum. Pl. v. 693 (1850).
parviflora, *Herb. Amaryll.* 226.—Syria.

VAGINARIA, Kunth, Enum. Pl. v. 693 (1850) = praec.

VAGINARIA, Pers. Syn. i. 70 (1805) = **Fuirena**, Rottb. (Cyperac.).
Richardi, Pers. l. c. = F. scirpoidea.

VAGNERA, Adans. Fam. ii. 496 (1763) = **Smilacina**, Desf. (Liliac.).

VAHEA, Lam. Illustr. ii. 292 (1798?) = **Landolphia**, Beauv. (1806) (Apocyn.).
angustifolia, Miq. Fl. Ind. Bat. ii. 394.—Sumatra.
comorensis, Boj. Hort. Maurit. 207; et ex A. DC. in DC. Prod. viii. 328.—Ins. Comor.
crassipes, Radlk. in Bremen, Abh. viii. (1883) 399.—Madag.
florida, F. Muell. Extra-trop. Pl. Indian ed. (1880) 344 = L. florida.
gummifera, Lam. Illustr. 292. t. 169 = Urceola elastica.
Heudelotii, F. Muell. Extra-trop. Pl. Indian ed. (1880) 344 = L. Heudelotii.
madagascariensis, Boj. Hort. Maurit. 207 (excl. syn.); et ex A. DC. in DC. Prod. viii. 327.—Madag.
madagascariensis, Boj. l. c., partim = Urceola elastica.
owariensis, F. Muell. Extra-trop. Pl. Indian ed. (1880) 344 = L. owariensis.
senegalensis, A. DC. in DC. Prod. viii. 328.—Afr. trop.

VAHLBERGELLA, Blytt, Norges Fl. iii. 1070 (1876) = WAHLBERGELLA, Fries = **Lychnis**, Tourn. (Caryoph.).

VAHLIA, Dahl, Obs. Bot. 21 (1787) = **Dombeya**, Cav. (Sterculiac.).
fastuosa, Dahl, l. c. (*tastuosa*).—Ins. Maurit.

VAHLIA, Thunb. Nov. Gen. ii. 36 (1782). *SAXIFRAGEAE*, Benth. & Hook. f. i. 637.
BISTELLA, Delile, in Caill. Voy. Méroé (Cent. Pl. Afr.) 97. t. 63 (1826).
RUSSELIA, Linn. f. Suppl. 24 (1781).
capensis, *Thunb. Nov. Gen.* ii. 36.—Afr. trop. et austr.
cordofana, Hochst. in Flora, xxiv. (1841) i. Intell. 43 = oldenlandioides.
macrantha, Klotzsch, in Peters, Reise Mossamb. Bot. 175 = oldenlandioides.
Oldenlandiae, DC. Prod. iv. 54 = oldenlandioides.
oldenlandioides, *Roxb. Hort. Beng.* 86; *Fl. Ind.* ii. 89.—As. et Afr. trop.
ramosissima, A. DC. ex DC. Prod. iv. 54 = oldenlandioides.
sessiliflora, DC. l. c. = viscosa.
silenoides, A. DC. ex DC. l. c. = oldenlandioides.
tomentosa, *A. DC. ex DC. l. c.*—Afr. trop.
verbasciflora, Welw. ex Oliver, Fl. Trop. Afr. ii. 384, in obs. = capensis.
viscosa, *Roxb. Hort. Beng.* 86; *Fl. Ind.* ii. 89.—As. et Afr. trop.
Weddenii, Reichb. Ic. Bot. Exot. i. 62. t. 91 = viscosa.

VAHLODEA, Fries, in Bot. Notiser (1842), 141, 178 = **Deschampsia**, Beauv. (Gramin.).
atropurpurea, Fries, l. c. 178; et Summa Veg. Scand. 243 = D. atropurpurea.

VAILLANTIA, Tourn. ex Linn. Syst. ed. I (1735) (Valantia); Gled. in Mém. Acad. Berl. v. (1751) 127. *RUBIACEAE*, Benth. & Hook. f. ii. 148.
aculeata, Tenore, Fl. Neap. Prod. App. iv. 33 = muralis.
alpina, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 86, nomen = Galium vernum.
Aparine, J. F. Gmel. Syst. 241 = Galium tricorne.
Aparine, Lam. Fl. Fr. iii. 383 = Galium Aparine.
Aparine, Linn. Sp. Pl. 1051 = Galium saccharatum.
aprica, Tausch, in Flora, xii. (1829) 647 = Galium apricum.
aristata, Boiss. & Heldr. Diagn. Ser. I. vi. 65 = Galium apricum.
articulata, Linn. Sp. Pl. 1052 = Galium articulatum.
aspera, Hort. ex Roem. & Schult. Syst. iii. 219 = Galium pedemontanum.
Bauhini, Schur, Enum. Pl. Transs. 287 = Galium vernum.
chersonensis, Willd. Sp. Pl. iv. 951 = Galium Cruciata.
ciliata, Opiz, ex J. & C. Presl, Fl. Čech. 34; et Natural. vi. (1824) 45 = Galium Cruciata.
ciliata, [Soland. in] Russ. Aleppo, ed. II. ii. 266 = hispida.
cordata, Pers. Syn. i. 129 = Galium articulatum.
Cruciata, Linn. Sp. Pl. 1052 = Galium Cruciata.
Cucullaria, Linn. Cent. Pl. i. 32; Amoen. Acad. iv. 296 = Callipeltis Cucullaria.
cucullata, Hill, Veg. Syst. vii. 27 = praec.?
filiformis, Ait. Hort. Kew. ed. I. iii. 428 = Galium filiforme.
glabra, Linn. Sp. Pl. ed. II. 491 = Galium vernum.
granularis, Roem. & Schult. Syst. iii. 249, sphalm. = *granulata*, Spreng.
granulata, Roth, ex Willd. Sp. Pl. iv. 949 = Galium pedemontanum.
granulata, Spreng. in Schrad. Journ. (1800) ii. 200 = Galium saccharatum?
hirsuta, Gilib. Exercit. i. 25 = Galium Cruciata.
hispida, Georgi, Beschr. Russ. Reich. III. iv. 1366 = Galium Cruciata.
hispida, *Linn. Syst.* ed. X. 1307.—Reg. Mediterr.
hispida, Lumnitz. Fl. Poson. 454 = Galium pedemontanum.
hispida, Pall. Tabl. Taur. 59 = Galium coronatum.
hispida, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = *lanata*.
humifusa, Willd. Sp. Pl. iv. 949 = Galium coronatum.
hypocarpia, Linn. Syst. ed. X. 1307 = Relbunium hypocarpium.
incrassata, Pomel, Nouv. Mat. Fl. Atl. 78 = hispida.
lanata, Delile, ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 (Quid?).—Aegypt.
muralis, Bieb. Fl. Taur. Cauc. ii. 435 = Galium articulatum.
muralis, *Linn. Sp. Pl.* 1051.—Europ. austr.
pedemontana, Bell. in Mém. Acad. Sc. Turin, v. 1790–1791 (1793) 252 = Galium pedemontanum.
persica, C. Koch, in Linnaea, xxiv. (1851) 473 = Galium coronatum.
pusilla, Stev. in Mém. Soc. Nat. Mosc. vii. (1829) 274 = Galium articulatum.
quadriflora, Moench, Meth. 640 = muralis.
rediviva, Hoffm. Hort. Mosq. (1808) n. 3387; ex Bieb. Fl. Taur. Cauc. iii. 640 = Galium coronatum.
rupestris, Lam. Fl. Fr. iii. 384 = Galium murale.
saccharata, C. C. Gmel. Fl. Bad. iii. 781 = Galium saccharatum.
Scopolii, Bess. Enum. Pl. Volh. 51, 89 = Galium vernum.
spuria, Pers. Syn. i. 129 = Galium tricorne.
taurica, Pall. in Nov. Act. Acad. Petrop. x. (1797) 315 = Galium coronatum.
tricornis, Roth, Neue Beytr. i. 142 = Galium tricorne.
triflora, Lam. Fl. Fr. iii. 384 = Galium tricorne.
verna, G. Don, in Loud. Hort. Brit. 411 = Galium vernum.

VAINILLA, Salisb. Parad. Lond. ii. ii. sub t. 82 (1807) = **Vanilla**, Juss. (Orchid.).

VALANTIA, Linn. Syst. ed. I (1735) = **Vaillantia**, Tourn. (Rubiac.).

VALARUM, Schur, Enum. Pl. Transs. 54 (1866) = **Sisymbrium**, Tourn. (Crucif.).
officinale, Schur, l. c. = S. officinale.

VALCARCELIA, Lag. ex Lindl. Introd. Nat. Syst. ed. II. 157 (1836). Genus ignotum *LEGUMINOSARUM*.—Cf. Benth. & Hook. f. i. 465.

VALCARCELLA, Steud. Nom. ed. II. 741 (1841), sphalm. = praec.

VALDESIA, Ruiz & Pav. Prod. 67 (1794) = **Blakea**, P. Br. (Melastom.).
latifolia, Ruiz & Pav. Fl. Per. iv. t. 407 = B. latifolia.
ovalis, Ruiz & Pav. Syst. Veg. 121 = B. ovalis.
repens, Ruiz & Pav. l. c. = B. repens.
rosea, Ruiz & Pav. Fl. Per. iv. t. 408 = B. rosea.
subconnata, Tafalla, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 148 = B. subconnata.

VALDIA, Plum. ex Adans. Fam. ii. 157 (1763) = **Clerodendron**, Linn. (Verben.).

VALDIVIA, Remy, in C. Gay, Fl. Chil. iii. 43. t. 29 (1847). *SAXIFRAGEAE*, Benth. & Hook. f. i. 644.
Gayana, *Remy, l. c.* 44.—Chili.

VALENTIANA, Rafin. Speech. i. 87 (1814); ex DC. Prod. iv. 340. Genus dubium *CAPRIFOLIACEARUM*.—Cf. Benth. & Hook. f. ii. 2.
volubilis, *Rafin. l. c.*—Abyss.

VALENTINIA, Neck. Elem. ii. 450 (1790) = **Tachigalia**, Aubl. (Legumin.).

VALENTINIA, Sw. Prod. Veg. Ind. Occ. 63 (1788) = **Casearia**, Jacq. (Samyd.).
coriacea, Griseb. Cat. Pl. Cub. 9.—Cuba.
ilicifolia, Sw. Prod. Veg. Ind. Occ. 63.—Ind. occ.
sylvestris, Raeusch. Nom. ed. III. 109 = Erystathes sylvestris.

VALENZUELA, Mutis, ex Caldas, in Seman. Nuev. Gran. iii. (1810) 21 = seq.?

VALENZUELIA, Bert. ex Cambess. in Nouv. Ann. Mus. Par. iii. (1834) 236. t. 14. *SAPINDACEAE*, Benth. & Hook. f. i. 394.
GUINDILIA, Gill. in Hook. Bot. Misc. iii. (1833) 170.
trinervis, *Bert. ex Cambess. in Nouv. Ann. Mus. Par.* iii. (1834) 236.—Chili.

VALERANDA, Neck. Elem. ii. 33 (1790) = **Orphium**?, E. Mey. (Gentian.).

VALERIANA, Tourn. ex Linn. Syst. ed. I (1735). *VALERIANEAE*, Benth. & Hook. f. ii. 154.
ARETIASTRUM, Spach, Hist. Veg. Phan. x. 304 (1841).
OLIGACOCE, Willd. ex DC. Prod. iv. 632 (1830).
PHU, Rupp. Fl. Jen. ed. Hall. 214 (1745).
VEXIBIA, Rafin. Neogenyt. 3 (1825).
acuminata, Royle, Illustr. Bot. Himal. 241 = Hardwickii.
adscendens, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 173.—Ecuador.
aequiloba, *Clos, in C. Gay, Fl. Chil.* iii. 228.—Chili.
affinis, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 123.—Mexic.
affinis, Opiz, Natural. n. x. (1825) 243 = dioica.
alba, [Mazziari], in Ionios Anthol. ii. (1834) 450 = Centranthus ruber.
alliariaefolia, *Vahl, Enum.* ii. 11.—Europ. or.; As. occ.
alpestris, *Stev. in Mém. Soc. Nat. Mosc.* v. (1814) 342.—Reg. Cauc.
alpigena, Schur, Enum. Pl. Transs. 290 = montana.
alpina, Adam, in Weber & Mohr, Beitr. i. 45 = alpestris.
alpina, *Vell. Fl. Flum.* 28; i. t. 68.—Bras.
alternifolia, Ledeb. Fl. Alt. i. 52 = officinalis.
alternifolia, Wall. ex Hook. f. Fl. Brit. Ind. iv. 291 = Calorhabdos Brunonianus.
altissima, Hornem. Hort. Hafn. ii. 950 = sambucifolia.
altissima, Mikan, ex Bess. Enum. Pl. Volh. 4 = exaltata.
alypifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 325.—Ecuador.
amoena, Salisb. Prod. 56 = tripteris.

VALERIANA:—
Andonaegui, *Phil. in Anal. Univ. Chil.* (1872) 733.—Chili.
angustifolia, Bieb. Fl. Taur. Cauc. i. 24 = Centranthus longiflorus.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 4 = Centranthus angustifolius.
angustifolia, Sibth. & Sm. Fl. Graec. i. 22. t. 29 = Centranthus Sibthorpii.
angustifolia, Tausch, ex Host, Fl. Austr. i. 36 = officinalis.
angustiloba, *DC. Prod.* iv. 641.—Oriens.
annua, S. F. Gray, Nat. Arr. Brit. Pl. ii. 478 = Centranthus Calcitrapa.
apula, *Pourr. in Mém. Acad. Toul.* iii. (1788) 332.—Europ.
aquatica, Schlecht. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 214 = Leschenaultii.
aretioides, H. B. & K. Nov. Gen. et Sp. iii. 324 = Phyllactis aretioides.
arizonica, *A. Gray, in Proc. Am. Acad.* xix. (1883) 81.—Arizona.
armeriaefolia, *Schlecht. in Linnaea*, xxviii. (1856) 485.—Peruv.
Arnottiana, Steud. Nom. ed. II. ii. 741 = Hookeriana.
Arnottiana, Wight, Ic. t. 1045, 1046 = Hardwickii.
asarifolia, *Dufr. Hist. Valér.* 44.—Ins. Creta.
astrephioides, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 53 = Astrephia laxa.
auriculata, Tardent, Ess. Hist. Nat. Bessar. 65 = Valerianella Auricula?
barbareaefolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 121.—Mexic.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 214.—Ind. or.
Berterii, Colla, in Mem. Acc. Torin. xxxviii. (1835) 5 = V. Papilla.
Berteroniana, Steud. & Hochst. ex Steud. Nom. ed. II. ii. 741 = Bridgesii.
Bertiscea, *Panč. Elench. Pl. Vasc. Crna-Gora*, 42.—Europ.
Bonplandiana, *Wedd. Chlor. And.* ii. 19.—Ecuador.
brachiata, Pers. Syn. i. 37 = pinnatifida.
bracteata, Benth. Pl. Hartw. 195 = Phyllactis bracteata.
bracteata, Boiss. Fl. Orient. iii. 91 = leucophaea.
bracteata, Royle, Illustr. Bot. Himal. 241 = pyrolaefolia.
bracteosa, *Phil. in Linnaea*, xxxiii. (1864–65) 101.—Chili.
bractescens, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 57 = Phyllactis bracteata.
Bridgesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 365.—Chili.
Brunoniana, Wall. Cat. n. 405 = Calorhabdos Brunoniana.
Brunoniana, Wight & Arn. Prod. 443 = Leschenaultii.
bulbosa, Pall. Reise, i. 148 = tuberosa.
bulbosa, *Wedd. Chlor. And.* ii. 24.—Bolivia.
Bustillosi, *Phil. in Linnaea*, xxxiii. (1864–65) 104.—Chili.
caespitosa, *Rupr. Sert. Tiansch.* 51.—As. centr.
Calcitrapa, Linn. Sp. Pl. 31 = Centranthus Calcitrapa.
calthaefolia, Sieber, ex Steud. Nom. ed. I. 868, nomen (Quid?).—Hab.?
Candolleana, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 112 = scandens.
capensis, *Thunb. Prod. Pl. Cap.* 7.—Afr. austr.
capitata, *Pall. ex Link, Jahrb.* i. III. (1820) 66.—Reg. arct.
Cardamines, Bieb. Fl. Taur.-Cauc. i. 24 = sisymbriifolia.
carnosa, *Sm. Pl. Ic. Ined.* t. 52.—Reg. Magellan.
Catacata, *Azara, Voy.* ii. 490.—Reg. Andin.
celtica, *Linn. Sp. Pl.* 32.—Europ.
celtica, Vill. Hist. Pl. Dauph. ii. 285 = saliunca.
cephalantha, *Schlecht. in Linnaea*, xxviii. (1856) 483.—Peruv.
ceratophylla, *H. B. & K. Nov. Gen. et Sp.* iii. 333.—Mexic.
chaerophylla, Pers. Syn. i. 37 = Astrephia chaerophylloides.
chaerophylloides, Sm. Pl. Ic. Ined. t. 53 = Astrephia chaerophylloides.
chamaedryfolia, Cham. & Schlecht. in Linnaea, iii. (1828) 129 = Valerianopsis chamaedryfolia.
chinensis, Linn. Sp. Pl. 33 = Boerhaavia repanda.
ciliata, Torr. & Gray, Fl. N. Am. ii. 49 = edulis.
clarionaefolia, *Phil. in Linnaea*, xxviii. (1856) 702.—Chili.

VALERIANA:—
Clematitis, *H. B. & K. Nov. Gen. et Sp.* iii. 327.—N. Granat.
coarctata, *Ruiz & Pav. Fl. Per.* i. 40. t. 68. f. a.—Peruv.
colchaguensis, *Phil. in Linnaea*, xxxiii. (1864–65) 104.—Chili.
collina, Wallr. in Linnaea, xiv. (1840) 537 (Beitr. Fl. Hercyn. 167) = officinalis.
compacta, Opiz, Natural. n. x. (1825) 244 = dioica.
connata, *Ruiz & Pav. Fl. Per.* i. 39. t. 67. f. c.—Peruv.
cordata, *Griseb. in Goett. Abh.* vi. (1854) 126; *Phil. in Linnaea*, xxviii. (1856) 697.—Chili.
cordifolia, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 54 = Phyllactis cordifolia.
Cornucopiae, Linn. Sp. Pl. 31 = Fedia Cornucopiae.
coronata, Mill. Gard. Dict. ed. VIII. n. 11 = Valerianella discoidea.
coronata, Salisb. Prod. 56 = Valerianella coronata.
crassifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 328.—N. Granat.
crassipes, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 56 = Phyllactis crassipes.
Crinii, Orph. in Boiss. Diagn. Ser. II. ii. 119 = montana.
crispa, Hook. & Arn. Bot. Beech. Voy. 27 = pterocarpa.
crispa, Ruiz & Pav. Fl. Per. i. 41 = Astrephia crispa.
cuspidata, Bertol. ex DC. Prod. iv. 636 = montana.
dacica, Porc. in Magyar Növenyt. Lapok, ix. (1885) 127 = tripteris.
daghestanica, *Rupr. ex Boiss. Fl. Orient.* iii. 87.—Reg. Cauc.
decussata, Bonpl. ex Wedd. Chlor. And. ii. 19 = Bonplandiana.
decussata, *Ruiz & Pav. Fl. Per.* i. 42. t. 70. f. b.—Peruv.
densa, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 55 = Phyllactis densa.
densiflora, *Benth. Pl. Hartw.* 39.—Mexic.
dentata, All. Fl. Pedem. i. 4 = Valerianella dentata.
denudata, *Benth. Pl. Hartw.* 20.—Mexic.
dichotoma, [Moç. & Sesse], ex DC. Prod. iv. 640 = ceratophylla.
dinorrhiza, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 54 = Phyllactis dinorrhiza.
dioica, Bess. Prim. Fl. Galic. i. 48 = elongata.
dioica, *Linn. Sp. Pl.* 31.—Europ.
dioica, Pursh, Fl. Am. Sept. ii. 727 = sylvatica.
Dioscoridis, *Sibth. & Sm. Fl. Graec. Prod.* i. 21.—Graecia; As. Min.
discoidea, Willd. Sp. Pl. i. 184 = Valerianella discoidea.
divaricata, Hinter, in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 333 = exaltata.
diversifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 115.—Japon.
dubia, Bunge, in Ledeb. Fl. Alt. i. 52 = officinalis.
dubia, Turcz. ex Steud. Nom. ed. II. ii. 741 = heterophylla.
echinata, Linn. Syst. ed. X. 861 = Valerianella echinata.
edulis, *Nutt. in Torr. & Gray, Fl. N. Am.* ii. 48.—Am. bor. occ.
effusa, *Griseb. in Goett. Abh.* xix. (1874) 162.—Reg. Argent.
elata, D. Don, Prod. Fl. Nep. 159 = Hardwickii.
elegans, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 212.—Reg. Himal.
elegans, *Clos, in C. Gay, Fl. Chil.* iii. 236.—Chili.
elongata, Geners. Elench. Scepus. n. 32; ex Wahlenb. Fl. Carpat. 11 = tripteris.
elongata, *Jacq. Enum. Pl. Vindob.* 7, 205.—Austr.
Engleriana, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 54 = Phyllactis pinnatifida.
eriocephala, Boiss. Fl. Orient. iii. 1031 (Index), sphalm. = Valerianella hirsutissima.
erysimoides, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 16.—Peruv.
exaltata, Baumg. Enum. Stirp. Transs. 56 = sambucifolia.
exaltata, *Mikan, f. ex Pohl, Tent. Fl. Bohem.* i. 41.—Rossia.
excelsa, Poir. Encyc. viii. 301 = officinalis.
exscapa, *Griseb. in Lechl. Berb. Am. Austr.* 57, nomen.—Peruv.
exscapa, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 3 = longiflora, *Willk.*
ferax, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 55 = Phyllactis ferax.

VALERIANA:—
ficariaefolia, *Boiss. Fl. Orient.* iii. 89.—Persia.
flaccidissima, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 228.—Japon.
floribunda, *Phil. in Linnaea*, xxviii. (1856) 701.—Chili.
florida, Salisb. Prod. 55 = Centranthus ruber.
foliosa, *Phil. in Linnaea*, xxviii. (1856) 698.—Chili.
Fonki, *Phil. l. c.* 698.—Chili.
fragilis, *Clos, in C. Gay, Fl. Chil.* iii. 226.—Chili.
Galeottianum, Mart. & Gal. in Bull. Acad. Brux. xi. 1. (1844) 124 = Phyllactis pratensis.
Gesneri ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 114.—Europ.
glauca, Lapeyr. Hist. Abr. Pl. Pyr. 19, in syn. = apula.
glauca, *Poepp. ex DC. Prod.* iv. 639.—Chili.
globifera, Pers. Syn. i. 37 = seq.
globiflora, *Ruiz & Pav. Fl. Per.* i. 43. t. 65. f. b.—Peruv.
globulariaefolia, Ramond, ex DC. Fl. Fr. iv. 236 = apula.
globularis, *A. Gray, in Proc. Am. Acad.* v. (1861) 322.—Peruv.
gracilipes, *Clos, in C. Gay, Fl. Chil.* iii. 231.—Chili.
gracilis, *Benth. Pl. Hartw.* 196.—Ecuador.
grandiflora, Wall. Cat. n. 403, sphalm. = Veronica himalensis.
grandifolia, *Phil. in Linnaea*, xxviii. (1856) 700.—Chili.
Grisiana, *Wedd. Chlor. And.* ii. 19.—Peruv.
Hardwickiana, Schult. Mant. i. 259 = Hardwickii.
Hardwickii, D. Don, Prod. Fl. Nep. 159 = Wallichii.
Hardwickii, Thw. Enum. Pl. Zeyl. 159 = Moonii.
Hardwickii, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 166.—Reg. Himal.; Malaya.
hebecarpa, *DC. Prod.* iv. 638.—Chili.
heterophylla, Baumg. Enum. Stirp. Transs. 57 = tripteris.
heterophylla, Loisel. Fl. Gall. ed. I. 21 = apula.
heterophylla, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. VII. 153, nomen; xviii. (1845) I. 321.—Sibir. baical.
hirtella, *H. B. & K. Nov. Gen. et Sp.* iii. 327.—Ecuador.
hispida, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 56 = Phyllactis hispida.
hispida, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 172.—N. Granat.
hispidula, Boiss. Diagn. Ser. I. iii. 56 = officinalis.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 84 = Hardwickii.
Hookeri, Shuttl. in Flora, xx. (1837) 450 = sitchensis.
Hookeriana, *Wight & Arn. Prod.* 444.—Ind. or.
Hoppii, Reichb. Ic. Fl. Germ. xii. 34 = tripteris.
Hornschuchiana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 357.—Chili.
hortensis, Lam. Fl. Fr. iii. 359 = V. Phu.
Humboldtii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 365.—Chili.
hyalinorhiza, H. B. & K. Nov. Gen. et Sp. iii. 331 = Humboldtii.
hyalinorrhiza, *Ruiz & Pav. Fl. Per.* i. 41. t. 67. f. b.—Chili; Peruv.
inconspicua, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 55 = Phyllactis inconspicua.
incrassata, Bory & Chaub. Nouv. Fl. Pélop. 2 = Valerianella eriocarpa.
intermedia, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 88, 89 = montana.
intermedia, Vahl, Enum. ii. 9 = tripteris.
interrupta, *Ruiz & Pav. Fl. Per.* i. 42. t. 67. f. a.—Peruv.
isophylla, Arv. Ess. Pl. Dauph. 34 (Quid?).—Gallia.
italica, Lam. Illustr. i. 92 = Dioscoridis.
Jaeschkii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 212.—Reg. Himal.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 115.—Japon.
Jatamansi, Jones, in As. Res. ii. (1790) 416; iv. (1795) 451 = Wallichii.
Jatamansi, Wall. Cat. n. 431 = Nardostachys Jatamansi.
javanica, Blume, Bijdr. 919 = Hardwickii.
Jetamansi, Dufr. Hist. Valér. 43, sphalm. = *V. Jatamansi.*
laciniata, Ruiz & Pav. Fl. Per. i. 42 = Astrephia chaerophylloides.
laciniata, Salisb. Prod. 56 = Centranthus Calcitrapa.

VALERIANA:—
laciniosa, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 121.—Mexic.
laeta, Salisb. Prod. 56 = V. Phu.
laevicaulis, Stokes, Bot. Mat. Med. i. 95 = V. Phu.
laevigata, Willd. ex Steud. Nom. ed. I. 869, nomen (Quid ?).—Am. austr.
lapathifolia, *Vahl, Enum.* ii. 11.—Reg. Magellan.
lasiocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 160.—Reg. Argent.
latifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 124.—Mexic.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 328.—N. Granat.
laxa, *Phil. in Linnaea*, xxviii. (1856) 701.—Chili.
laxiflora, *DC. Prod.* iv. 638.—Chili.
laxiflora, Phil. in Anal. Univ. Chil. (1872) 734 = sparsiflora.
lepidota, *Clos, in C. Gay, Fl. Chil.* iii. 230.—Chili.
Leschenaultii, *DC. Prod.* iv. 640.—Ind. or.
leucocarpa, *DC. l. c.* 638.—Chili.
leucophaea, *DC. l. c.* 644.—As. Min.; Persia.
Lindleyana, Wall. Cat. n. 404 = Picrorhiza Kurroa.
lobata, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 53 = Astrephia lobata.
Locusta, Linn. Sp. Pl. 33 = Valerianellae spp. variae.
Locusta, Ucria, Hort. Reg. Panorm. 46; ex Guss. Fl. Sic. Prod. i. 24 = Fedia Cornucopiae.
longiflora, *Regel & Schmalh. in Act. Hort. Petrop.* vii. (1880) 384.— Turkestan.
longiflora, *Willk. in Flora*, xxxiv. (1851) 733.—Hispan.
longifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 330.—N. Granat.
lucida, Hort. Par. ex DC. Prod. iv. 641 = officinalis.
lutescens, *Phil. in Linnaea*, xxviii. (1856) 699.—Chili.
lyrata, *Phil. l. c.* xxxiii. (1864–65) 103.—Chili.
lyrata, *Vahl, Enum.* ii. 4.—Peruv.
macrophylla, Bieb. Fl. Taur.-Cauc. i. 25 = alliariaefolia.
macrorhiza, Poepp. ex DC. Prod. iv. 635 = Phyllactis macrorrhiza.
macrosiphon, Vilm. Fl. Pl. Terre, ed. I. 916 = Centranthus macrosiphon.
magellanica, *Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 54.—Reg. Magellan.
magellanica, Lam. Illustr. i. 93 = carnosa.
magna, *Clos, in C. Gay, Fl. Chil.* iii. 227.—Chili.
major, Pall. Reise, iii. 318 = officinalis.
Mandoniana, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 54 = Phyllactis Mandoniana.
mexicana, *DC. Prod.* iv. 640.—Mexic.
microphylla, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 50 = Beddomei.
microphylla, *H. B. & K. Nov. Gen. et Sp.* iii. 325.—Peruv.
micropterina, *Wedd. Chlor. And.* ii. 26.—Bolivia.
Mikaniae, *Lindl. in Journ. Hort. Soc.* iii. (1848) 316.—Guatemala.
mixta, Linn. Sp. Pl. 34 = Valerianella dentata.
monandra, Vill. Hist. Pl. Dauph. ii. 280 = Centranthus angustifolius.
monorrhiza, Dufr. Hist. Valér. 44 = tuberosa.
montana, Bieb. Fl. Taur.-Cauc. i. 25 = alpestris.
montana, Geners. Elench. Scepus. n. 31; ex Wahlenb. Fl. Carpat. 11 = dioica.
montana, Georgi, Beschr. Russ. Reich. III. iv. 662 = capitata.
montana, *Linn. Sp. Pl.* 32.—Europ.
Montteana, *Phil. in Linnaea*, xxx. (1859–60) 191.—Chili.
Moonii, *Arn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 213.—Zeylan.
Morisoni, Mert. & Koch, Deutschl. Fl. i. 397, in syn., sphalm. = Valerianella dentata.
multiceps, Wallr. in Linnaea, xiv. (1840) 539 (Beitr. Fl. Hercyn. 169) = exaltata.
multifida, Roem. & Schult. Syst. i. 363, in syn. = Valerianella membranacea.
muriculata, *Phil. in Linnaea*, xxxiii. (1864–65) 102.—Chili.
Mutisiana, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 54 = Phyllactis Mutisiana.
Napus, *Lindl. Bot. Reg.* (1840) *Misc.* 76.—Mexic.
nasturtiifolia, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 4 = dioica.
nivalis, *Wedd. Chlor. And.* ii. 23.—Bolivia.

VALERIANA:—
oblongifolia, *Ruiz & Pav. Fl. Per.* i. 40. t. 65. f. a.—Peruv.
obovata, Clos, in C. Gay, Fl. Chil. iii. 221 = obtusifolia.
obovata, Schult. Mant. i. 265 = edulis ?
obtusifolia, *DC. Prod.* iv. 635.—Chili.
officinalis, *Linn. Sp.* 31.—Europ.; As. bor.
olenaea, *Boiss. & Heldr. Diagn.* Ser. II. ii. 118.—Graecia.
oligantha, *Boiss. & Heldr. l. c.* Ser. I. x. 72.—Graecia.
olitoria, Willd. Fl. Berol. Prod. 19 = Valerianella olitoria.
orbiculata, Sibth. & Sm. Fl. Graec. Prod. i. 21 = Centranthus Calcitrapa.
oreocharis, *Phil. in Linnaea*, xxviii. (1856) 699.—Chili.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 112 = Valerianopsis organensis.
origanifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 171.—Peruv.
paniculata, *Ruiz & Pav. Fl. Per.* i. 41. t. 70. f. a.—Peruv.
Papilla, *Bert. ex DC. Prod.* iv. 638.—Chili.
parviflora, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 57 (= *Porteria parviflora*).—Venezuela.
parviflora, Miegév. in Bull. Soc. Bot. Fr. xiv. (1867) 145 = Centranthus Calcitrapa.
pauciflora, Hook. Fl. Bor. Am. i. 292. t. 101 = sitchensis.
pauciflora, *Michx. Fl. Bot. Am.* i. 18.—Am. bor.
Pavonii, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 16. t. 215.—Peruv.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864–65) 101.—Chili.
peltata, *Clos, in C. Gay, Fl. Chil.* iii. 229.—Chili.
petrophila, *Bunge, in Ledeb. Fl. Alt.* i. 54.—Afghan.; Sibir. altaic.
Phaseoli, *A. Br. in Ind. Sem. Hort. Berol.* (1851) 13 (*App.* 2).—Mexic.
phellocarpa, *Phil. in Linnaea*, xxxiii. (1864–65) 100.—Chili.
Phu, Lapeyr. Hist. Abr. Pl. Pyr. 18 = montana.
Phu, *Linn. Sp. Pl.* 32.—Europ.; Reg. Cauc.
pilosa, *Phil. in Linnaea*, xxviii. (1856) 702.—Chili.
pilosa, *Ruiz & Pav. Fl. Per.* i. 39. t. 66. f. a.—Peruv.
pilosiuscula, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 122.—Mexic.
pinnata, Gilib. Exercit. ii. 454 = officinalis.
pinnatifida, *Ruiz & Pav. Fl. Per.* i. 40. t. 69. f. b.—Peruv.
plantaginea, *H. B. & K. Nov. Gen. et Sp.* iii. 329.—Ecuador.
polemonioides, *H. B. & K. l. c.* 332.—Ecuador.
polybotrya, Hoeck, in Engl. Bot. Jahrb. iii. (1882) 55 = Phyllactis polybotrya.
polygama, Bess. ex DC. Prod. iv. 637 = elongata.
polystachya, Sm. Pl. Ic. Ined. t. 51 = Valerianopsis polystachya.
pratensis, Steud. Nom. ed. II. ii. 742 = Phyllactis pratensis.
procera, *H. B. & K. Nov. Gen. et Sp.* iii. 329.—Mexic.
procurrens, Wallr. in Linnaea, xiv. (1840) 540 (Beitr. Fl. Hercyn. 170) = officinalis.
pterocarpa, *Hook. & Arn. Bot. Beech. Voy.* 28, in nota.—Chili.
pubescens, Germann, in Bot. Zeit. Regensb. iv. (1805) 359 = Valerianella uncinata.
pubescens, *Phil. Fl. Atac.* 26.—Chili.
pulchella, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 123.—Mexic.
pulchella, *Phil. in Linnaea*, xxxiii. (1864–65) 102.—Chili.
pumila, Willd. Sp. Pl. i. 184 = Valerianella membranacea.
pusilla, Royle, Illustr. Bot. Himal. 241 = dioica.
pycnantha, *A. Gray, in Proc. Am. Acad.* v. (1861) 322.—Peruv.
pygmaea, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 172.—Ecuador.
pyramidalis, *H. B. & K. Nov. Gen. et Sp.* iii. 331.—Ecuador.
pyrenaica, *Linn. Sp. Pl.* 33.—Mont. Pyren.
pyrolaefolia, *Decne. in Jacquem. Voy. Bot.* 85. t. 93.—Reg. Himal.
quadrangularis, *H. B. & K. Nov. Gen. et Sp.* iii. 326.—Peruv.
radiata, Willd. Sp. Pl. i. 184 = Valerianella radiata.
radicalis, *Clos, in C. Gay, Fl. Chil.* iii. 215.—Chili.
ramosissima, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 122.—Mexic.

VALERIANA:—
Regnellii, *Sond. in Linnaea*, xxii. (1849) 573.—Bras.
regularis, *Clos, in C. Gay, Fl. Chil.* iii. 225.—Chili.
remota, *Ball, in Journ. Linn. Soc.* xxii. (1885) 42.—Peruv.
repens, Host, Fl. Austr. i. 35 = officinalis.
repens, *Wall. Cat.* n. 434; *et ex DC. Prod.* iv. 640.—Nepal.
rhizantha, *A. Gray, in Proc. Am. Acad.* v. (1861) 322.—Peruv.
rigida, Ruiz & Pav. Fl. Per. i. 39. t. 65. f. c = Phyllactis rigida.
ringens, Salisb. Prod. 56 = Fedia Cornucopiae.
rotundifolia, Sibth. & Sm. Fl. Graec. t. 31 = Centranthus Calcitrapa.
rotundifolia, Vill. Hist. Pl. Dauph. ii. 283 = montana.
rotundiloba, *Clos, in C. Gay, Fl. Chil.* iii. 233.—Chili.
Roylei, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 83.—Reg. Himal.
rubra, Linn. Sp. Pl. 31 = Centranthus ruber.
rumicoides, *Wedd. Chlor. And.* ii. 22.—N. Granat.
rupestris, Pall. Reise, iii. 266 = Patrinia rupestris (in Addend.).
rupestris, *Phil. in Anal. Univ. Chil.* (1872) 734.—Chili.
rupestris, Uspenski, ex Ledeb. Fl. Ross. ii. 426 = Patrinia sibirica.
rupicola, Lag. in Varied. Cienc. ii. iv. (1805) 212 = apula.
rupicola, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 16. t. 217.—Chili.
ruthenica, Willd. Sp. Pl. i. 181 = Patrinia sibirica.
salicariaefolia, Vahl, Enum. ii. 16 = Valerianopsis salicariaefolia.
saliunca, *All. Fl. Pedem.* i. 3. t. 70. f. 1.—Europ. austr.
sambucifolia, Eichw. Skizze, 152 = exaltata.
sambucifolia, *Mikan f. ex Pohl, Tent. Fl. Bohem.* i. 41.—Europ.
samolifolia, Bert. ex Colla, in Mem. Acc. Torin. xxxviii. (1835) 4. t. 21 = Valerianella samolifolia.
Sanguisorbae, Pers. Syn. i. 37 = seq.
sanguisorbaefolia, *Cav. Ic.* v. 34.—Chili.
saxatilis, Georgi, Beschr. Russ. Berol. III. iv. 662 = capitata.
saxatilis, *Linn. Sp. Pl.* 33.—Europ.
saxatilis, Vill. Hist. Pl. Dauph. ii. 286 = celtica.
saxicola, *C. A. Mey. Verz. Pfl. Cauc.* 49.—Reg. Cauc.
scandens, Forsk. Fl. Aegypt. Arab. 12 = Boerhaavia scandens.
scandens, *Loefl. It. Hisp.* 235.—Am. trop.
sciophila, Schur, Enum. Pl. Transs. 289 = sambucifolia.
scorpioides, *DC. Prod.* iv. 635.—Mexic.
scrophulariaefolia, Pourr. in Mém. Acad. Toul. iii. (1788) 331 = montana.
sedifolia, Urv. in Mém. Soc. Linn. Par. iv. (1826) 612 = Phyllactis sedifolia.
sedoides, Hombron & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 53 = Phyllactis sedifolia.
senecioides, *Phil. in Anal. Univ. Chil.* (1872) 735.—Chili.
serrata, *Ruiz & Pav. Fl. Per.* i. 40. t. 68. f. c.—Peruv.
sibirica, Linn. Sp. Pl. 34 = Patrinia sibirica.
sibirica, Willd. Sp. Pl. i. 181 = Patrinia rupestris (in Addend.).
siifolia, C. Koch, in Linnaea, xix. (1847) 32 = Dioscoridis.
simplex, *Clos, in C. Gay, Fl. Chil.* iii. 224.—Chili.
simplicifolia, Kab.—Cf. Verh. Bot. Ver. Brandenb. vi. (1864) 110 = dioica.
sinensis, J. F. Gmel. Syst. 92 = *chinensis*.
sisymbriifolia, Schur, Enum. Pl. Transs. 290 = tripteris.
sisymbriifolia, Urv. in Mém. Soc. Linn. Par. i. (1822) 260 = Dioscoridis.
sisymbriifolia, *Vahl, Enum.* ii. 7.—As. Min.; Persia.
sitchensis, *Bong. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 145.—Am. bor. occ.
sorbifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 332.—Mexic.
sparsiflora, *Clos, in C. Gay, Fl. Chil.* iii. 218.—Chili.
spathulata, *Phil. in Anal. Univ. Chil.* (1862) II. 397.—Chili.
spathulata, Ruiz & Pav. Fl. Per. i. 40 t. 68. f. b = Phyllactis spathulata.

VALERIANA:—
speluncaria, *Boiss. Fl. Orient.* iii. 88.—Cilicia.
sphaerocarpa, *Phil. in Anal. Univ. Chil.* (1872) 730. —Chili.
Spica, Vahl, Enum. ii. 13 = Wallichii.
Steveni, Ledeb. Fl. Ross. ii. 436 = Centranthus Calcitrapa.
stolonifera, Czern. in Bull. Soc. Nat. Mosc. xviii. (1845) II. 133, in nota = officinalis.
Stracheyi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 212.—Reg. Himal.
striata, Royle, Illustr. Bot. Himal. 241, nomen (Quid ?). —Cashmere.
stricta, *Clos, in C. Gay, Fl. Chil.* iii. 235.—Chili.
subdentata, Opiz, ex Mert. & Koch, Deutschl. Fl. i. 394 = dioica.
subincisa, *Benth. Pl. Hartw.* 39.—Mexic.
supina, DC. Fl. Fr. iv. 237 = saliunca.
supina, *Linn. Mant.* i. 27.—Austria.
sylvatica, *Banks, ex Richards. in Frankl. 1st Journ.* ed. II. *App.* 2.—Am. bor.
sylvatica, F. W. Schmidt, Fl. Böem. ii. 2 = dioica.
sylvestris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 478 = dioica.
sylvestris, Groschke, in Keyserl. & Dersch. Beschr. Prov. Kurland, (1805) 110 = officinalis.
tenera, Wall. ex DC. Prod. iv. 640 = Hardwickii.
tenuifolia, Ruiz & Pav. Fl. Per. i. 39. t. 65. f. d = Phyllactis tenuifolia.
tenuissima, Schur, Enum. Pl. Transs. 288 = officinalis.
toluccana, *DC. Prod.* iv. 640.—Mexic.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* iii. 327.—Ecuador.
transsilvanica, Schur, Enum. Pl. Transs. 290 = tripteris.
triloba, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 115.—Japon.
trinervis, Viv. Fl. Lib. Spec. 67; Fl. Cors. 3 = Centranthus nervosus.
triphylla, *H. B. & K. Nov. Gen. et Sp.* iii. 326.—N. Granat.
tripteris, *Linn. Sp. Pl.* 32.—Europ.
tuberosa, *Linn. l. c.* 33.—Europ. austr.; As. bor.
tuberosa, Sprun. ex Nym. Consp. 336 = Dioscoridis.
uncinata, Bieb. Fl. Taur. Cauc. i. 26 = Valerianella uncinata.
undulata, Steud. Nom. ed. II. ii. 742, sphalm. = Veronica Anagallis ?
urticaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 330.—N. Granat.; Peruv.
vaga, *Clos, in C. Gay, Fl. Chil.* iii. 227.—Chili.
vaginata, *H. B. & K. Nov. Gen. et Sp.* iii. 331.—Mexic.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 700.—Chili.
velutina, *Clos, in C. Gay, Fl. Chil.* iii. 233.—Chili.
veronicaefolia, Willd. ex Steud. Nom. ed. I. 870 = urticaefolia ?
verticillata, *Clos, in C. Gay, Fl. Chil.* iii. 223.—Chili.
vesicaria, Mill. Gard. Dict. ed. VIII. n. 10 = Valerianella vesicaria.
Villarsii, Timb. in Mém. Acad. Toul. Sér. IV. vi. (1856) 104 (Quid ?).—Gallia.
villosa, Moon, Cat. Pl. Ceyl. 5 = Moonii.
villosa, Thunb. Fl. Jap. 32 = Patrinia villosa.
villosa, Wall. Cat. n. 433; et ex DC. Prod. iv. 640 = Wallichii.
violifolia, Griff. Notul. iv. 213; Ic. Pl. Asiat. t. 465 = Wallichii.
virescens, *Clos, in C. Gay, Fl. Chil.* iii. 222.—Chili.
virgata, *Ruiz & Pav. Fl. Per.* i. 42. t. 66. f. b.—Peruv.
Volkmanni, *Phil. in Anal. Univ. Chil.* (1872) 733.—Chili.
volubilis, [Moç. & Sesse], ex DC. Prod. iv. 634 = scandens.
Wallichii, *DC. l. c.* 640.—Reg. Himal.

VALERIANELLA, Tourn. ex Hall. Enum. Stirp. Helv. 666 (1742); Moench, Meth. 493 (1794), partim. *VALERIANACEAE*, Benth. & Hook. f. ii. 156.
BETCKEA, DC. Mém. Valer. 18 (1829); Prod. iv. 642 (1830).
DUFRESNIA, DC. l. c. 8. t. 3 (1829); et Prod. l. c. 624 (1830).
FEDIA, Gaertn. Fruct. ii. 36. t. 86. f. 3 (1791), partim.

VALERIANELLA:—
LOCUSTA, Riv. ex Medic. Phil. Bot. i. 55 (1789).
MASEMA, Dulac, Fl. Hautes-Pyr. 476 (1867).
ODONTOCARPA, Neck. Elem. i. 123 (1790).
PLECTRITIS, DC. Mém. Valer. 13 (1829); Prod. iv. 651 (1830).
POLYPREMUM, Adans. Fam. ii. 152 (1763).
abyssinica, Fresen. in Mus. Senckenb. ii. (1837) 116 = dentata.
affinis, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 837.—Ins. Socotra.
Amarella, *Krok, in Vet. Akad. Handl. Stockh.* v. (1864). n. 1. 31, 55.—Texas.
ambigua, J. Lange, ex Krok, l. c. 56 = fallax.
amblyotis, *Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc.* (1838) 263.—Reg. Cauc.; Persia.
angulosa, DC. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 654, nomen (Quid ?).—Hab. ?
anomala, *A. Gray, in Proc. Am. Acad.* xix. (1883) 83. —Am. bor. occ.
aphanoptera, *A. Gray, l. c.*—Am. bor. occ.
Aucheri, Boiss. Diagn. Ser. I. iii. 58 = Szovitsiana.
Auricula, DC. Fl. Fr. Suppl. 492 = rimosa.
Boissieri, *Krok, in Vet. Akad. Handl. Stockh.* v. (1864) n. 1. 32, 81.—Phrygia.
brachystephana, *Bertol. Fl. Ital.* i. 193.—Ital.
bracteata, *Lowe, Man. Fl. Mad.* i. 407.—Ins. Mader.
caerulea, Eaton & Wright, N. Am. Bot. ed. VIII. 469 = olitoria.
calcarata, Stev. ex Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 51 = sclerocarpa.
campanulata, Bivona, Stirp. Rar. Sicil. Manip. iv. 18 = eriocarpa.
capitata, *Boiss. & Bal. Diagn.* Ser. II. vi. 92.—Phrygia.
carinata, *Loisel. Not. Pl. Fr.* 149.—Europ.; Oriens.
chenopodifolia, *DC. Prod.* iv. 629.—Am. bor.
chlorodonta, *Coss. & Dur. ex Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 740.—Alger.
chlorostephana, Boiss. & Bal. Diagn. Ser. II. vi. 93 = Boissieri.
chlorostephana, *Pomel, Nouv. Mat. Fl. Atl.* 67.—Alger.
coerulea, Auct. vide caerulea.
congesta, *Lindl. Bot. Reg.* t. 1094.—Am. bor. occ.
conjungens, *Bunge, ex Boiss. Fl. Orient.* iii. 109.—Persia.
corniculata, *C. A. Mey. Verz. Pfl. Cauc.* 49.—Reg. Cauc.
Cornucopiae, Loisel. Fl. Gall. ed. I. 23 = Fedia Cornucopiae.
coronata, *DC. Fl. Fr.* iv. 241.—Europ.; Oriens.
coronata, DC. Prod. iv. 628 = discoidea.
corymbifera, *Boiss. Fl. Orient.* iii. 107.—Pamphyl.
costata, *DC. Prod.* iv. 626.—Europ.
cucullata, Dufour, ex Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 40 = eriocarpa.
cymbaecarpa, *C. A. Mey. Verz. Pfl. Cauc.* 49.—As. Min.; Persia.
dactylophylla, *Boiss. & Hohen. Diagn.* Ser. I. x. 75.—Syria; Persia.
dasycarpa, Link, Enum. Hort. Berol. i. 63 = dentata.
dentata, Loisel. Fl. Gall. ed. I. 21 = rimosa.
dentata, *Pollich, Hist. Pl. Palat.* i. 30.—Europ.; As. bor.
dichotoma, Gilib. Exercit. ii. 456 = olitoria.
diodon, *Boiss. Diagn.* Ser. I. iii. 57.—As. Min.
diplusodon, *Boiss. l. c.* x. 74.—Syria.
discoidea, *Loisel. Not. Pl. Fr.* 148.—Graec.; As. Min.
divaricata, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 61.—Hispan.
Dufresnia, *Bunge, ex Boiss. Fl. Orient.* iii. 109.—As. Min.; Pers.; Afghan.
echinata, *DC. Fl. Fr.* iv. 242.—Europ. austr.; As. Min.
eriocarpa, *Desv. Journ. Bot.* ii. (1809) 314. t. 11. f. 2.—Europ.; Afr. bor.
eriocephala, Hampe, in Flora, xxv. (1842) 64 = hirsutissima.
exscapa, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 251 = Hohenackeria bupleurifolia.
Fagopyrum, Walp. Rep. ii. 527 = chenopodifolia.
fallax, *Coss. & Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. v. (1856) 32, nomen; *Krok, in Vet. Akad. Handl. Stockh.* v. (1864) n. 1. 56.—Ital.; Afr. bor.
gibbosa, DC. Prod. iv. 626 = costata.

VALERIANELLA:—
gjoelbaschiensis, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 109.—Lycia.
glomerata, *Boiss. & Bal. Diagn.* Ser. II. vi. 93.—As. Min.
gracilis, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 29 = membranacea.
gracilis, Haberl. ex Heynh. Nom. i. 835 (Quid ?).—Tauria.
hamata, DC. Fl. Fr. Suppl. 494 = coronata.
heterocarpa, Bisch. Hort. Heidelb. (1851) 6.—Cf. Linnaea, xxv. (1852) 317 = membranacea.
hirsutissima, *Link, in Linnaea*, ix. (1834) 580.—Graec.; As. Min.
hispidiuscula, Mérat, Rev. Fl. Par. 227 = dentata.
Huetii, *Boiss. Fl. Orient.* iii. 101.—Armenia.
incrassata, Nym. Syll. 62 = eriocarpa.
Kotschyi, *Boiss. Diagn.* Ser. I. iii. 60.—Syria.
lasiocarpa, *Stev. ex Betcke, Animad. Bot. Valer.* 26. —Reg. Cauc.; Mesopot.; Oriens.
lasiocephala, Stev. ex Betcke, l. c. 21 = coronata.
laticuspis, Bertol. Fl. Ital. i. 856 = membranacea.
laxa, *Boiss. & Bal. Diagn.* Ser. II. ii. 121.—As. Min.; Syria.
laxa, Presl, Repert. Bot. Syst. i. 11, in nota = Astrephia laxa.
laxiflora, Dufr. Hist. Valér. 58. t. 3 = rimosa.
leiocarpa, Auct. ex Steud. Nom. ed. II. ii. 743 = dentata.
leptocarpa, *Pomel, Nouv. Mat. Fl. Atl.* 293.—Alger.
lingulata, Presl, Bot. Bemerk. 88 = dentata.
Locusta, Betcke, Animad. Bot. Valer. 10 = olitoria.
longiflora, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 43.—Turkestan.
longiflora, *Walp. Rep.* ii. 527.—Am. bor. occ.
lutea, Moench, Meth. 493 = Patrinia sibirica.
macrocera, *A. Gray, in Proc. Am. Acad.* xix. (1883) 83. —Am. bor. occ.
macrocyathus, Pomel, Nouv. Mat. Fl. Atl. 69 = eriocarpa.
membranacea, *Loisel. Not. Pl. Fr.* 150.—Europ.
microcarpa, Loisel. l. c. 151 = dentata.
microcephala, Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 77 = divaricata.
microcyathus, Pomel, Nouv. Mat. Fl. Atl. 68 = eriocarpa.
microstephana, Boiss. & Bal. Diagn. Ser. II. vi. 92 = lasiocarpa.
minuta, Steud. Nom. ed. II. i. 626, sphalm. = seq.
mixta, Dufr. Hist. Valér. 58. t. 3 = dentata.
monodon, C. Koch, in Linnaea, xvii. (1843) 34 = cymbaecarpa.
Morisonii, DC. Prod. iv. 627 = dentata.
multidentata, Losc. & Pard. Ser. Inconf. Pl. Arag. ed. Willk. 49 = coronata.
muricata, Bieb. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 654, nomen (Quid ?).—Reg. Cauc.
muricata, Heldr. ex Nym. Consp. 339 = eriocarpa.
mutica, Kern. in Oestr. Bot. Zeitschr. xx. (1870) 357 = membranacea.
Nuttallii, *Walp. Rep.* ii. 527.—Am. bor. occ.
obtusiloba, *Boiss. Diagn.* Ser. I. iii. 59.—Graec.; As. Min.
olitoria, *Pollich, Hist. Pl. Palat.* i. 30.—Europ.; Oriens.
orientalis, *Boiss. & Bal. Diagn.* Ser. II. ii. 120.—As. Min.
otodonta, Pomel, Nouv. Mat. Fl. Atl. 68 = dentata.
oxyrhyncha, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 51.—As. Min.; Persia.
parviflora, Dougl. ex Torr. & Gray, Fl. N. Am. ii. 50 = congesta.
patellaria, Wood, Class-Book (1861), 405 = Woodsiana.
persica, *Boiss. Diagn.* Ser. I. iii. 58.—Persia.
petrophila, Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 98 = Valeriana petrophila.
plagiocyathus, Pomel, Nouv. Mat. Fl. Atl. 69 = eriocarpa.
plagiostephana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 52.—Reg. Cauc.; Persia.
platyloba, *Dufr. Hist. Valér.* 59. t. 3. f. 1.—Reg. Mediterr.
praecox, Willk. in Linnaea, xxx. (1859–60) 104 = carinata.
pteroloba, *Steud. Nom.* ed. II. ii. 743.—Iberia.
pterophila, Steud. l. c. = Valeriana petrophila.
puberula, DC. Prod. iv. 627 = dentata.
puberula, Reichb. ex Nym. Consp. 339 = eriocarpa.
pubescens, Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 48 = uncinata.

VALERIANELLA:—
pubescens, Mérat, Nouv. Fl. Par. (1812) 13 = dentata.
pulchella, Fisch. ex Krok, in Vet. Akad. Handl. Stockh. v. (1864) n. 1. 75 = membranacea.
pumila, DC. Fl. Fr. Suppl. 494 = membranacea.
punctata, Hort. Patav. ex Heynh. Nom. i. 836, nomen (Quid ?).—Hab. ?
pusilla, Miégev. in Bull. Soc. Bot. Fr. x. (1863) 27 = olitoria.
quadrangularis, Steud. Nom. ed. II. i. 626 = carinata.
radiata, *Dufr. Hist. Valér.* 57.—Am. bor.
remota, Boiss. & Hohen. Diagn. Ser. I. x. 75 = sclerocarpa.
rhombicarpa, Aiken, ex Torr. & Gray, Fl. N. Am. ii. 51 = olitoria.
rimosa, *Bast. in Desv. Journ. Bot.* iii. (1814) 20.—Hispan.
rotata, Roem. & Schult. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 654 (Quid ?).—Hab. ?
rugulosa, Fisch. Hort. Gorenk. ed. II. (1812) 43 = eriocarpa.
samolifolia, *A. Gray, in Proc. Am. Acad.* xix. (1883) 83.—Am. bor. occ.
sclerocarpa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 53.—Reg. Cauc.; Persia.
Sheilii, Boiss. Fl. Orient. iii. 109 = membranacea.
sicula, Link, in Linnaea, ix. (1834) 579 = discoidea.
Soyeri, *Buching. ex Boiss. Diagn.* Ser. I. x. 74.—Graecia.
sphaerocarpa, DC. Prod. iv. 628 = membranacea.
stenocarpa, *Krok, in Vet. Acad. Handl. Stockh.* v. (1864) n. 1. 64.—Texas.
stephanodon, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 741.—Alger.
Stocksii, *Boiss. Fl. Orient.* iii. 99.—Persia; Beluchist.
striata, Stev. in Fisch. Cat. Hort. Gorenk. ed. II. (1812) 43 = olitoria.
Szovitsiana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 48.—Arab.; Syria; Persia.
trichocarpa, Steud. Nom. ed. II. ii. 743, sphalm. = turgida.
tridentata, Betcke, Animadv. Bot. Valer. 12; Reichb. Fl. Germ. Excurs. 198 = rimosa.
tridentata, Gasp. in Il Progresso, iv. (1833) 169 = membranacea.
trigonocarpa, DC. Prod. iv. 628 = turgida.
triplaris, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 111.—Persia; Afghan.
triquetra, Shuttl. in Flora, xx. (1837) 211 = chenopodifolia.
truncata, *Betcke, Animadv. Bot. Valer.* 22; *DC. Prod.* iv. 627.—Europ. austr.; Oriens.
truncata, Pett. ex Nym. Consp. 339 = eriocarpa.
tuberculata, *Boiss. Diagn.* Ser. I. iii. 59.—Syria; Persia.
turgida, *Betcke, Animadv. Bot. Valer.* 14; *DC. Prod.* iv. 626.—Reg. Mediterr.; Tauria.
turkestanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 43.—Turkestan.
umbilicata, Wood, Class-Book (1861), 405 = Woodsiana.
uncinata, *Dufr. Hist. Valér.* 60.—Reg. Cauc.; Persia.
vesicaria, *Moench, Meth.* 493.—Reg. Mediterr.; Persia.
Woodsiana, *Walp. Rep.* ii. 527.—Am. bor.

VALERIANOIDES, Medic. Phil. Bot. i. 177 (1789) = **Stachytarpheta**, Vahl (Verben.).
indica, Medic. l. c. 178 = S. indica.
jamaicensis, Medic. l. c. = S. indica.

VALERIANOPSIS, C. A. Muell. in Mart. Fl. Bras. vi. IV. 345 (1885). *VALERIANACEAE.*
angustifolia, *C. A. Muell. l. c.* 346.—Bras.
chamaedryfolia, *C. A. Muell. l. c.* 349.—Bras.
Eichleriana, *C. A. Muell. l. c.* 348.—Bras.
foliosa, *C. A. Muell. l. c.* 347.—Bras.
organensis, *C. A. Muell. l. c.* 348.—Bras.
polystachya, *C. A. Muell. l. c.* 350.—Bras.
salicariaefolia, *C. A. Muell. l. c.* 347.—Bras.

VALIKAHA, Adans. Fam. ii. 84 (1763) = **Memecylon**, Linn. (Melastom.).

VALISNERIA, Scop. Introd. 68 (1777), sphalm. = **Vallisneria**, Mich. (Hydroch.).

VALKERA, Stokes, Bot. Mat. Med. i. 447 (1812) = **Walkera**, Schreb. (Inc. Sed.).

VALLANTIA, A. Dietr. Syn. Pl. i. 420 (1839) = **Vaillantia**, Tourn. (Rubiac.).

VALLARIS, Burm. f. Fl. Ind. 51 (1768). *APOCYNACEAE*, Benth. & Hook. f. ii. 710.
EMERICIA, Roem. & Schult. Syst. iv. 33, 401 (1819).
PELTANTHERA, Roth, Nov. Pl. Sp. 132 (1821).
anceps, Wall. Cat. n. 1622 (al. ordin. ?).—Burma.
assamensis, Griff. Notul. iv. 77 = Heynii ?
clavata, *G. Don, Gen. Syst.* iv. 79.—Ins. Molucc.
controversa, Spreng. Syst. iv. Cur. Post. 65 = Lepistemon Wallichii.
dichotoma, Wall. Cat. n. 1621; Griff. Notul. iv. 77 = Heynii.
divaricata, G. Don, Gen. Syst. iv. 79 = Strophanthes divaricata.
Heynii, *Spreng. Syst.* i. 635.—Ind. or.; Burma.
indica, J. F. Gmel. Syst. 391 = V. Pergulana.
lancifolia, *Hook. f. Fl. Brit. Ind.* iii. 651.—Malacca.
laxiflora, Blume, Bijdr. 1043 = Pottsia cantonensis.
Maingayi, *Hook. f. Fl. Brit. Ind.* iii. 651.—Malacca.
ovalis, Miq. Fl. Ind. Bat. ii. 427 = V. Pergulana.
Pergulana, *Burm. f. Fl. Ind.* 51.—Ind. or.; Malaya.
Pergularia, Hook. Ic. Pl. t. 153, sphalm. = praec.
sinensis, *G. Don, Gen. Syst.* iv. 79.—China.

VALLARIS, Rafin. Fl. Tellur. iv. 114 (1836) = **Euphorbia**, Linn.
fimbriata, Rafin. l. c. 115.—Hab. ?
Ipecacuana, Rafin. l. c. = E. Ipecacuanha.
missurica, Rafin. l. c. (Quid ?).—Am. bor.
portulacoides, Rafin. l. c. = E. portulacoides.
uniflora, Rafin. l. c. = E. uniflora.

VALLEA, Mutis, ex Linn. f. Suppl. 42 (1781). *TILIACEAE*, Benth. & Hook. f. i. 238.
Calomala, Blanco, Fl. Filip. ed. I. 439 = Elaeocarpus nitidus.
cernua, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 236.—N. Granat.
cordifolia, *Ruiz & Pav. Syst. Veg.* 132.—Peruv.
glabra, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 48.—Peruv.
macrophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 238.—N. Granat.
ovata, *Turcz. l. c.* 237.—N. Granat.
pubescens, *H. B. & K. Nov. Gen. et Sp.* v. 350.—N. Granat.
pyrifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 237.—Ecuador.
Schlimii, *Turcz. l. c.* xxxvi. (1863) 1. 576.—N. Granat.
stipularis, *Linn. f. Suppl.* 266.—N. Granat.

VALLESIA, Ruiz & Pav. Prod. 28. t. 5 (1794) *APOCYNACEAE*, Benth. & Hook. f. ii. 696.
chiococcoides, H. B. & K. Nov. Gen. et Sp. iii. 233. t. 241 = cymbaefolia.
cymbaefolia, *Ortega, Hort. Matr. Dec.* 58.—Am. trop.
dichotoma, Ruiz & Pav. Fl. Per. ii. 26. t. 151 = cymbaefolia.
glabra, Link, Enum. Hort. Berol. i. 207 = cymbaefolia.
hypoglauca, *Ernst, in Journ. Bot.* viii. (1870) 375.—Venezuela.
mexicana, *Muell. Arg. in Linnaea*, xxx. (1859–60) 393.—Mexic.
pubescens, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 195.—Ins. Galapag.
punctata, *Spreng. Neue Entdeck.* iii. 33.—Bras.

VALLISNERIA, Mich. ex Linn. Gen. ed. I. 300 (1737). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 451 [bis].
PHYSKIUM, Lour. Fl. Cochinch. 662 (1790).
aethiopica, *Fenzl, in Flora*, xviii. (1844) 311, nomen; *et in Sitzb. Akad. Wien, Math.-Nat.* li. (1865) *Abth.* ii. 139.—Afr. trop.
alternifolia, Roxb. Hort. Beng. 71; Fl. Ind. iii. 750 = Lagarosiphon Roxburghii.
americana, Michx. Fl. Bor. Am. ii. 220 = spiralis.
bulbosa, Poir. Encyc. viii. 321 = Sagittaria sagittifolia.
Jacquini, Savi, Oss. (1816) 12 = spiralis.
Jacquiniana, Spreng. Syst. iii. 900 = praec.
Linnei, Bercht. & Presl, Rostl. i. Hydrochar. 7 = spiralis.
Micheliana, Spreng. Syst. iii. 900 = seq.
Michelii, Savi, Oss. (1816) 11 = spiralis.

VALLISNERIA:—
nana, R. Br. Prod. 345 = spiralis.
numidica, *Pomel, Nouv. Mat. Fl. Atl.* 386.—Alger.
octandra, Roxb. Pl. Corom. ii. 34 = Blyxa Roxburghii.
Physcium, *Juss. ex Spreng. Syst.* iii. 900.—Cochinch.
pusilla, Barbieri, ex Bertol. Fl. Ital. x. 299, 300 = spiralis.
spiralis, *Linn. Sp. Pl.* 1015.—Reg. temp. et calid.
spiraloides, Roxb. Hort. Beng. 71; Fl. Ind. iii. 750 = spiralis.
verticillata, Roxb. ll. cc. 71; 751 = Hydrilla verticillata.

VALLOTA, Herb. App. [Bot. Reg.] 29 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 729.
elata, M. Roem. Syn. Ensat. 110 = purpurea.
eximia, Hort. ex Flor. Mag. iv. (1865) t. 225 = purpurea.
grandiflora, Carr. Rev. Hortic. (1870) 50 = purpurea.
purpurea, *Herb. App.* [*Bot. Reg.*] 29.—Afr. austr.

VALLOTA, Steud. Nom. ed. II. ii. 744 (1841), sphalm. = VALOTA, Adans. = **Panicum**, Linn. (Gramin.).

VALORADIA, Hochst. in Flora, xxiv. (1841) 1. Intell. 23; xxv. (1842) 239 = **Ceratostigma**, Bunge (Plumbag.).
abyssinica, Hochst. ll. cc. 23; 239 = C. abyssinicum.
patula, Hochst. l. c. xxiv. (1842) 240.—Abyss.
plumbaginoides, Boiss. in DC. Prod. xii. 695 = C. plumbaginoides.

VALOTA, Adans. Fam. ii. 495 (1763) = **Panicum**, Linn. (Gramin.).

VALOTA, Dum. Anal. Fam. 58 (1829) = **Vallota**, Herb. (Amaryll.).

VALSONICA, Scop. Ann. iv. Hist. Nat. 97 (1770), sphalm. = **Watsonia**, Mill. (Irid.).

VALTETA, Rafin. Sylva Tellur. 53 (1838) = **Iochroma**, Benth. (1845) (Solanac.).
fuchsioides, Rafin. l. c. = I. fuchsioides.
gesneroides, Rafin. l. c. = I. gesnerioides.

VALVINTERLOBUS, Dulac, Fl. Hautes-Pyr. 459 (1867) = **Wahlenbergia**, Schrad. (Campan.).
filiformis, Dulac, l. c. = W. hederacea.

VANALPHIMIA, Lesch. ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 419 = **Saurauja**, Willd. (Ternstr.).
Djinote, Lesch. ex DC. l. c. 424 = S. crenulata.
lanceolata, Lesch. ex DC. l. c. 421 = S. lanceolata.

VANALPHMIA, Steud. Nom. ed. II. ii. 744 (1841), sphalm. = praec.

VANCOUVERIA, C. Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 351. *BERBERIDEAE*, Benth. & Hook. f. i. 44.
SCULERIA, Rafin. Fl. Tellur. ii. 51 (1836).
chrysantha, *Greene, in Bull. Calif. Acad.* i. 11. (1885) 66.—Oregon.
hexandra, *C. Morr. & Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 351.—Am. bor. occ.

VANDA, Jones, in As. Res. iv. (1795) 302; R. Br. in Bot. Reg. t. 506 (1820). *ORCHIDEAE*, Benth. & Hook. f. iii. 578.
alpina, *Lindl. Fol. Orch. Vanda*, 10.—Reg. Himal.
Batemanni, Lindl. Bot. Reg. (1846) t. 59 = Stauropsis lissochiloides.
Bensoni, *Batem. in Bot. Mag.* t. 5611; *Veitch & Reichb. f. in Gard. Chron.* (1867) 180, cum f. —Burma.
bicaudata, Thw. Enum. Pl. Zeyl. 429 (= Cottoniae sp.).—Zeylan.
bicolor, *Griff. Notul.* iii. 354; *Ic. Pl. Asiat.* t. 330.—Reg. Himal.
Boxallii, *Reichb. f. in Gard. Chron.* (1881) 1. 366.—Ins. Philipp.
brunnea, *Reichb. f. Xenia Orch.* ii. 138.—Burma.
Cathcarti, Lindl. Fol. Orch. Vanda, 8; Hook. f. Illustr. Himal. Pl. t. 23 = Arachnantha Cathcartii.
clitellaria, *Reichb. f. in Flora*, lv. (1872) 277.—Ins. Marian.

VANDA :—
coerulea, *Griff. ex Lindl. Bot. Reg.* (1847) sub t. 30. —Reg. Himal.
coerulescens, *Griff. Notul.* iii. 352; *Ic. Pl. Asiat.* t. 331.—Burma.
coerulescens, [Lindl.] in Journ. Hort. Soc. vi. (1857) p. viii, lapsu = coerulea.
concolor, *Blume, Rumphia*, iv. 49.—China.
congesta, Lindl. Bot. Reg. (1839) Misc. 61 = Acampe congesta.
crassiloba, *Teijsm. & Binn. Cat. Hort. Bog.* 48, nomen. —Ins. Buru.
cristata, *Lindl. in Wall. Cat.* n. 7328; *et Gen. et Sp. Orch.* 216.—Reg. Himal.
cruenta, Lodd. Cat. ex Sweet, Hort. Brit. ed. I. 388 (Quid?).—China.
Cumingii, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 654 = lamellata.
Denisoniana, *Benson & Reichb. f. in Gard. Chron.* (1869) 528.—Burma.
densiflora, *Lindl. in Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 21, sub t. 42.—Burma.
falcata, Beer, Prakt. Orch. 317 = Angraecum falcatum.
fasciata, Gardn. ex Lindl. Fol. Orch. Acampe, 2 = Acampe Wightiana.
fimbriata, Gardn. ex Thw. Enum. Pl. Zeyl. 305 (= *Cleisostoma acaule*).—Zeylan.
furva, Lindl. Bot. Reg. (1844) Misc. 39, partim = concolor.
furva, *Lindl. Gen. et Sp. Orch.* 215.—Ins. Molucc.
fuscoviridis, Lindl. in Gard. Chron. (1848) 351 = furva.
gigantea, Lindl. in Wall. Cat. n. 7326 = Stauropsis gigantea.
Griffithii, Lindl. in Lindl. & Paxt. Flow. Gard. ii. (1851–52) 22 = alpina.
Guiberti, Linden, ex Reichb. f. in Bot. Zeit. xx. (1862) 375 = Cleisostoma Guiberti.
hastifera, *Reichb. f. in Linnaea*, xli. (1877) 30.—Hab.?
helvola, *Blume, Rumphia*, iv. 49.—Java.
Hindsii, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 237.—N. Guin.
Hookeri, Hort.—Cf. Gard. Chron. (1882) II 375, 558 = seq.
Hookeriana, *Reichb. f. in Bonplandia*, iv. (1856) 324.—Malaya.
insignis, *Blume, Rumphia*, iv. 49. t. 192.—Malaya.
lamellata, *Lindl. Bot. Reg.* (1838) *Misc.* 66.—Ins. Philipp.
lilacina, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 325.—Siam.
limbata, *Blume, Rumphia*, iv. 49.—Java.
Lindlayana, Griff. Notul. iii. 353, err. typ. = seq.
Lindleyana, Griff. ex Lindl. & Paxt. Flow. Gard. ii. (1851–52) 22 = Stauropsis gigantea.
lissochiloides, Lindl. Gen. et Sp. Orch. 216 = Stauropsis lissochiloides.
longifolia, Lindl. in Wall. Cat. n. 7322; et in Lindl. & Paxt. Flow. Gard. ii. (1851–52) 21 = Acampe longifolia.
Lowei, Lindl. in Gard. Chron. (1847) 239 = Arachnanthe Lowei.
multiflora, Lindl. Coll. Bot. t. 38 = Acampe longifolia.
paniculata, R. Br. in Bot. Reg. sub t. 506; et Not. ad calcem tomi vi. [3] = Sarcanthus paniculatus.
Parishii, *Reichb. f. Xenia Orch.* ii. 138.—Burma.
parviflora, *Lindl. Bot. Reg.* (1844) *Misc.* 45.—Ind. or.; Burma.
pauciflora, Breda, Orch. Kuhl. Hass. [t. 13] (Sarcochili sp.).—Java.
pedunculariis, Lindl. Gen. et Sp. Orch. 216 = Cottonia macrostachya.
pulchella, Wight, Ic. t. 1671 = Saccolabium calceolare.
pusilla, Teijsm. & Binn. in Tijdschr. Nederl. Ind. v. (1853) 493 = Trichoglottis pusilla.
recurva, Hook. Exot. Fl. iii. t. 187 = Sarcanthus rostratus.
rostrata, Lodd. Bot. Cab. t. 1008 = Sarcanthus rostratus.
Roxburghii, *R. Br. in Bot. Reg.* t. 506.—Ind. or.; Burma.
Sanderiana, *Reichb. f. in Gard. Chron.* (1882) I. 588.—Ins. Philipp.
scripta, Spreng. Syst. iii. 719 = Grammatophyllum speciosum.
spathulata, *Spreng. l. c.*—Ind. or.
Stangeana, *Reichb. f. in Bot. Zeit.* xvi. (1858) 351.—Assam.

VANDA :—
Storiei, Storie, ex Reichb. f. in Gard. Chron. (1880) II. 296 = Renanthera Storiei.
striata, Reichb. f. Xenia Orch. ii. 137 = cristata.
suaveolens, Blume, Rumphia, iv. 49 = tricolor.
suavis, *Lindl. in Gard. Chron.* (1848) 351.—Java.
suavis, F. Muell. Fragm. vii. 135 = Hindsii.
Sulingi, Blume, Mus. Bot. Lugd. Bat. i. 62 = Renanthera Sulingi.
teres, *Lindl. in Wall. Cat.* n. 7324.—Ind. or.; Burma.
teretifolia, Lindl. Coll. Bot. t. 6 = Sarcanthus teretifolius.
tessellata, Hook. ex G. Don, in Loud. Hort. Brit. 372; Lodd. ex Paxt. Mag. Bot. vii. (1840) 265 = Roxburghii.
tesselloides, Reichb. f. in Walp. Ann. vi. 864 = Roxburghii.
testacea, Reichb. f. in Gard. Chron. (1877) II. 166 = parviflora.
trichorhiza, Hook. Exot. Fl. i. t. 72 = Luisia trichorhiza.
tricolor, Hook. Bot. Mag. t. 4432 = suavis.
tricolor, *Lindl. Bot. Reg.* (1847) sub t. 59.—Java.
undulata, Lindl. in Journ. Linn. Soc. iii. (1859) 42 (Stauropsidis sp.).—Reg. Himal.
unicolor, Steud. Nom. ed. II. ii. 744 = lamellata, Roxburghii.
Vidalii, Boxall, ex Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 240 = lamellata.
violacea, *Lindl. Bot. Reg.* (1841) *Misc.* 12.—Ins. Philipp.
Vipanii, *Reichb. f. in Gard. Chron.* (1882) II. 134.—Burma.
Wightiana, Lindl. ex Wight, Ic. t. 1670 = Acampe Wightiana.
Wightii, *Reichb. f. in Walp. Ann.* vi. 932.—Ind. or.

VANDALEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 331 = **Sisymbrium**, Tourn. (Crucif.).
rupestricola, Fourr. l. c. = S. austriacum.

VANDEA, Griff. Notul. iii. 352 (1851), sphalm. = **Vanda**, R. Br. (Orchid.).

VANDELLIA, Linn. Mant. i. 12 (1767). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 955.
ANAGALLOIDES, Krock. Fl. Siles, ii. I. 398 (1790).
DICEROS, Blume, Bijdr. 751 (1826).
ELLOBUM, Blume, l. c. 746 (1826).
HORNEMANNIA, Link & Otto, Ic. Pl. Sel. 9. t. 3 (1820?).
ILYOGETON, Endl. Gen. 684 (1839).
LINDERNIA, All. Misc. Taur. iii. (1775) 178. t. 5.
MITRANTHUS, Hochst. in Flora, xxvii. (1844) 103.
PYXIDARIA, Rupp. Fl. Jen. ed. Hall. 237 (1745).
TITTMANNIA, Reichb. Ic. Bot. Exot. i. 27. t. 38 (1824).
VRIESEA, Hassk. in Flora, xxv. (1842) II. Beibl. 27.
alba, Benth. Scroph. Ind. 35 = crustacea.
alsinoides, *Benth. in DC. Prod.* x. 415.—Austral.
angustifolia, *Benth. Scroph. Ind.* 37.—As. trop.
arridens, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 226 = pedunculata.
Brownii, Benth. in DC. Prod. x. 413 = crustacea.
caespitosa, *Benth. l. c.* 415.—Java.
Chamaedrys, G. Don, Gen. Syst. iv. 549 = Gratiola Chamaedrys.
clausa, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 60 = Bonnaya clausa.
cordifolia, G. Don, Gen. Syst. iv. 549 = pedunculata.
corymbosa, *Baker, in Journ. Bot.* xx. (1882) 221.—Madag.
crustacea, *Benth. Scroph. Ind.* 35.—Cosmop. trop.
cymulosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 117.—Japon.
diffusa, Blanco, Fl. Filip. ed. I. 505 = Torenia parviflora.
diffusa, *Linn. Mant.* i. 89.—Am. trop.
diffusa, Willd. Sp. Pl. iii. 343 = pedunculata.
elata, *Benth. Scroph. Ind.* 36.—Burma.
Ellobum, *Benth. in DC. Prod.* x. 417.—Java.
erecta, *Benth. Scroph. Ind.* 36.—Europ.; Geront. trop. et subtrop.
erecta, Benth. l. c., partim = multiflora.
glandulifera, *Benth. in DC. Prod.* x. 415.—Java.
hirsuta, *Buch.-Ham. ex Benth. Scroph. Ind.* 36.—As. trop.
hirta, Buch.-Ham. ex Wall. Cat. sub n. 3942 = mollis.

VANDELLIA :—
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 280.—Reg. Himal.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 118.—Japon.
latifolia, *Benth. in DC. Prod.* x. 415.—Java.
laxa, Benth. Scroph. Ind. 36 = scabra.
limosa, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 394.—China.
lobelioides, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 61 = Ilysanthes lobelioides.
lobelioides, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 120.—Afr. trop.
Malurea, Raeusch. Nom. ed. III. 180 = Stemodia pusilla.
Matourea, Steud. Nom. ed. I. 870, sphalm. = praec.
minima, Royle, ex Benth. Scroph. Ind. 37 = sessiliflora.
minuta, *Miq. Fl. Ind. Bat.* ii. 691.—Java.
mollis, *Benth. Scroph. Ind.* 37.—Reg. Himal.; China.
molluginoides, *Benth. l. c.* 35.—Burma.
monnerioides, Buch.-Ham. ex Wall. Cat. sub n. 3945 = scabra.
montana, *Benth. in DC. Prod.* x. 415.—Java.
multiflora, Blanco, Fl. Filip. ed. I. 505 = Torenia asiatica.
multiflora, *G. Don, Gen. Syst.* iv. 549.—Ind. or.
multiflora, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 148 = hirsuta.
nervosa, *Benth. in DC. Prod.* x. 417.—Java.
nummularifolia, *D. Don, Prod. Fl. Nep.* 86.—Reg. Himal.; China.
oblonga, Benth. Scroph. Ind. 35 = Torenia oblonga.
obovata, Walp. Rep. iii. 294 = Mazus rugosus.
obtusa, *Miq. Fl. Ind. Bat.* ii. 691.—Java.
ovata, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 394 = hirsuta.
pachypoda, *Franch. & Sav. Enum. Pl. Jap.* ii. 457.—Japon.
palustris, Benth. in DC. Prod. x. 417, sphalm. = *pratensis*.
pedunculata, *Benth. Scroph. Ind.* 37.—As. trop.
plantaginea, F. Muell. in Trans. Vict. Inst. iii. (1859) 62 = Hemiarrhena plantaginea.
pratensis, Vahl, Eclog. Am. ii. 48 = Beyrichia villosa?
prostrata, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 966, nomen.—Guian.
pubescens, *Benth. in DC. Prod.* x. 415.—Austral.
Pyxidaria, Maxim. in Bull. Acad. Pétersb. xx. (1875) 449 = erecta.
racemosa, *Spreng. Neue Entdeck.* i. 262.—Bras.
repens, Thou. ex Steud. Nom. ed. II. ii. 744 = diffusa.
Roxburghii, G. Don, Gen. Syst. iv. 549 = pedunculata.
scabra, *Benth. Scroph. Ind.* 36.—As. et Afr. trop.
scapigera, *Benth. in DC. Prod.* x. 415.—Austral.
senegalensis, *Benth. l. c.* 416.—Afr. trop.
sessiliflora, *Benth. Scroph. Ind.* 37.—Reg. Himal.
Soriana, *Blanco, Fl. Filip.* ed. I. 506.—Ins. Philipp.
sphaerocarpa, *Benth. in DC. Prod.* x. 417.—Java.
stachydifolia, *Walp. Rep.* iii. 294.—China.
stemodioides, *Miq. Fl. Ind. Bat. Suppl.* 563.—Ins. Banca.
strigosa, G. Don, Gen. Syst. iv. 549 = Gratiola strigosa.
subcrenulata, *Miq. in Journ. Bot. Néerl.* i. (1861) 113. —China.
subulata, *Benth. in DC. Prod.* x. 415.—Austral.
sulcata, Steud. Nom. ed. II. ii. 744 = scabra.
urticifolia, *Hance, in Journ. Bot.* vi. (1868) 301.—China.
varians, G. Don, Gen. Syst. iv. 549 = crustacea.

VANDESIA, Salisb. in Trans. Hort. Soc. i. (1812) 332 = **Bomarea**, Mirb. (Amaryll.).
edulis, Salisb. l. c. = B. Salsilla.

VANESSA, Rafin. Fl. Tellur. iii. 57 (1836) = **Manettia**, Mutis (Rubiac.).
cordifolia, Rafin. l. c. = M. cordifolia.

VANGUERIA, Juss. Gen. 206 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 111.
LAGYNIAS, E. Mey. in Drège, Zwei Pfl. Docum. 197 (1843).
MEYNA, Roxb. ex Link, in Jahrb. i. III. (1820) 32.
RYTIGYNIA, Blume, Mus. Bot. Lugd. Bat. i. 178 (1850).

VANGUERIA:—
VAVANGA, Rohr, in Skrivt. Nat. Selsk. ii. 1. (1792) 207, 208. t. 7.
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 353.—Abyss.
atroviridis, Wall. Cat. n. 8412 = Plectronia glabra.
Barnimiana, Schweinf. Pl. Nilot. 30. t. 11 = abyssinica.
canthioides, *Benth. in Hook. Niger Fl.* 408.—Afr. trop.
Commersonii, Jacq. Hort. Schoenb. i. 20 = madagascariensis.
concolor, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 150.—Afr. trop.
crassa, Schweinf. ex Hiern, l. c. 145 (= *Canthium crassum*).—Afr. trop.
cymosa, Gaertn. f. Fruct. iii. 75. t. 193 = madagascariensis.
dicocca, Miq. Fl. Ind. Bat. ii. 251 = Plectronia didyma.
edulis, Miq. Fl. Ind. Bat. ii. 313 = spinosa.
edulis, Vahl, Symb. Bot. iii. 36 = madagascariensis.
emirnensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 168.—Madag.
euonymoides, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 150.—Afr. trop.
infausta, *Burch. Trav.* ii. 258.—Afr. trop. et austr.
lasiantha, *Sond. in Harv. & Sond. Fl. Cap.* iii. 14.—Afr. austr.
latifolia, *Sond. l. c.* 15.—Afr. austr.
leiophylla, *Miq. Fl. Ind. Bat.* ii. 250.—Java?
leucophloea, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 153 = Fadogia leucophloea.
lucidula, Miq. Fl. Ind. Bat. Suppl. 544 = Plectronia didyma.
macrocalyx, *Sond. in Linnaea*, xxiii. (1850) 59.—Afr. austr.
madagascariensis, *J. F. Gmel. Syst.* 367.—Afr. trop.
membranacea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 151.—Afr. trop.
Miqueliana, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 153, in obs. = spinosa.
mollis, Wall. Cat. n. 8411 = spinosa.
oligantha, *Miq. Fl. Ind. Bat.* ii. 251.—Ins. Sunda.
palembanica, Miq. Fl. Ind. Bat. Suppl. 544 = Diplospora singularis.
parvifolia, *Sond. in Linnaea*, xxiii. (1850) 58.—Afr. austr.
pauciflora, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 151.—Afr. trop.
pubescens, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 314 = spinosa.
senegalensis, *Benth. & Hook. f. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 149.—Afr. trop.
spinosa, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 536.—As. trop.
spirostylis, Miq. Fl. Ind. Bat. ii. 250 = Plectronia didyma.
stellata, Blanco, Fl. Filip. ed. I. 167 = spinosa.
tetraphylla, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 152.—Afr. trop.
tomentosa, Hochst. in Flora, xxv. (1842) 238 = infausta.
umbellulata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 150.—Afr. trop.
velutina, *Hiern, l. c.* 151.—Afr. trop.
velutina, Hook. Bot. Mag. t. 3014 = infausta.
venosa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 353 = madagascariensis.
venosa, *Sond. in Harv. & Sond. Fl. Cap.* iii. 14.—Afr. austr.
verrucosa, Sieber, ex DC. Prod. iv. 455 (Quid?).—Ins. Maurit.
Zeyheri, Sond. in Harv. & Sond. Fl. Cap. iii. 15 = Fadogia Zeyheri.

VANGUIERIA, Pers. Syn. i. 205 (1805) = praec.

VANHALLIA, Schult. f. Syst. vii. p. xviii. 166 (1829) = **Bragantia**, Lour. (Aristol.).
tomentosa, Schult. f. l. c. 166 = B. tomentosa.

VAN-HOUTTEA, Lem. in Bull. Soc. Hort. Orléans, i. (1845) 346 = **Houttea**, Decne. (Gesner.).

VANIERIA, Lour. Fl. Cochinch. 564 (1790). Genus dubium *URTICACEARUM*.—Cf. Benth. & Hook. f. Gen. iii. 351.
alternifolia, Stokes, Bot. Mat. Med. iv. 381 = *cochinchinensis*.

VANIERIA:—
chinensis, Lour. Fl. Cochinch. 564 (= *Procris cantoniensis*).—China.
cochinchinensis, Lour. l. c. 565 (= *Procris cochinchinensis*).—China.

VANIERIA, Montr. in Mém. Acad. Lyon, x. (1860) 176 = **Trisema**, Hook. f. (Dillen.).
tomentosa, Montr. l. c. = T. coriacea?

VANILLA, Plum. ex Mill. Gard. Dict. ed. VI. (1752); Juss. Gen. 66 (1789); Sw. in Nov. Act. Soc. Sc. Upsal. vi. (1799) 66. t. 5; et in Schrad. Journ. (1799) ii. 208. *ORCHIDEAE*, Benth. & Hook. f. iii. 590.
MYROBOMA, Salisb. Parad. Lond. t. 82 (1807).
VANILLOPHORUM, Neck. Elem. iii. 134 (1790).
acutifolia, Lodd. Cat. ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 655, nomen (Quid?).—Venezuela.
africana, *Lindl. in Journ. Linn. Soc.* vi. (1862) 137.—Afr. trop.
albida, *Blume, Cat. Gew. Buitenz.* 100; *et Bijdr.* 422.—Malaya.
anaromatica, Griseb. Fl. Brit. W. Ind. 638 = aromatica.
angustifolia, *Willd. Sp. Pl.* iv. 121.—Java.
aphylla, *Blume, Bijdr.* 422.—Java.
aphylla, Wight, Ic. t. 931 = Wightii.
aromatica, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 66; *et in Schrad. Journ.* (1799) ii. 208.—Am. trop.
aromatica, Willd. Sp. Pl. iv. 121, partim = planifolia.
axillaris, *Mill. Gard. Dict.* ed. VIII. n. 2.—Mexic.
barbellata, Reichb. f. in Flora, xlviii. (1865) 274 = claviculata.
bicolor, *Lindl. Bot. Reg.* (1838) *Misc.* 37.—Guian.
calopogon, *Reichb. f. Otia Bot. Hamb.* 40.—Ins. Philipp.
Chamissonis, *Klotzsch, in Bot. Zeit.* iv. (1846) 564.—Bras.?
claviculata, Lindl. in Mem. Am. Acad. N. S. viii. (1861) 219, nomen = Wrightii.
claviculata, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) 66; *et in Schrad. Journ.* (1799) ii. 209.—Ind. occ.
Epidendrum, Mirb. Hist. Pl. ed. II. ix. 249 = aromatica.
Fasciola, *Spreng. Pugill.* ii. 83 = Taeniophyllum Fasciola.
grandiflora, Lindl. Gen. et Sp. Orch. 435 = V. Pompona.
grandifolia, *Lindl. in Journ. Linn. Soc.* vi. (1862) 138.—Afr. trop.
gratiosa, Griseb. Cat. Pl. Cub. 267 = Wrightii.
Griffithii, *Reichb. f. in Bonplandia*, ii. (1854) 88.—Ind. or.
guianensis, *Splitg. in Ann. Sc. Nat.* Sér. II. xv. (1841) 279.—Guian.
hamata, *Klotzsch, in Bot. Zeit.* iv. (1846) 563.—Peruv.
Humblotii, *Reichb. f. in Gard. Chron.* (1885) i. 726; *et in Flora*, lxviii. (1885) 378.—Afr. trop.
inodora, Schiede, in Linnaea, iv. (1829) 574 = aromatica.
lutea, Wright, ex Griseb. Cat. Pl. Cub. 267 = palmarum.
lutescens, Moq. ex Dupuis, in Rev. Hortic. Sér. IV. v. (1856) 121. f. 24 = V. Pompona.
majajensis, Blanco, Fl. Filip. ed. II. 593 = planifolia.
Methonica, *Reichb. f. & Warsz. in Bonplandia*, ii. (1854) 97.—N. Granat.
mexicana, Mill. Gard. Dict. ed. VIII. n. 1 = aromatica, planifolia.
Moonii, *Thw. Enum. Pl. Zeyl.* 312.—Zeylan.
odorata, *Presl, Rel. Haenk.* 101.—Ecuador.
ovalis, Blanco, Fl. Filip. ed. II. 448 = aromatica.
palembanica, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 243.—Sumatra.
palmarum, *Lindl. Gen. et Sp. Orch.* 436.—Bras.
Parishii, *Reichb. f. Otia Bot. Hamb.* 39.—Burma.
parvifolia, *Rodrig. Orch. Nov.* ii. 271.—Bras.
Pfaviana, *Reichb. f. in Gard. Chron.* (1883) ii. 230.—Mexic.
phaeantha, *Reichb. f. in Flora*, xlviii. (1865) 274.—Ind. occ.
Phalaenopsis, *Reichb. f. ex Van Houtte, Fl. des Serres*, xvii. (1867–68) 97. t. 1769–1770.—Ins. Seychelles.
planifolia, *Andr. Bot. Rep.* t. 538.—Ind. occ.

VANILLA:—
planifolia, Griseb. Fl. Brit. W. Ind. 638 = phaeantha.
Poitaei, *Reichb. f. in Linnaea*, xli. (1877) 66.—Ins. S. Doming.
Pompona, *Schiede, in Linnaea*, iv. (1829) 573.—Mexic.
pterosperma, Lindl. ex Wall. Cat. n. 7402 = Galeola Hydra.
Roscheri, *Reichb. f. in Linnaea*, xli. (1877) 65.—Afr. trop.
rubiginosa, Griff. Notul. iii. 246 = Galeola Hydra?
Ruiziana, *Klotzsch, in Bot. Zeit.* iv. (1846) 563.—Peruv.
sativa, Schiede, in Linnaea, iv. (1829) 573 = planifolia.
speciosa, Boxall, ex Naves, in Blanco, Fl. Philipp. ed. III. Nov. App. 248, nomen (Quid?).—Ins. Philipp.
surinamensis, Reichb. f. in Nederl. Kruidk. Arch. iv. (1859) 321 = guianensis.
sylvestris, Schiede, in Linnaea, iv. (1829) 573 = planifolia.
viridiflora, Blume, Bijdr. 422 = planifolia.
Walkeriae, *Wight, Ic.* t. 932.—Ind. or.
Wightii, *Lindl. ex Wight, l. c.* sub t. 931.—Ind. or.
Wrightii, *Reichb. f. in Flora*, xlviii. (1865) 273.—Ind. occ.

VANILLOPHORUM, Neck. Elem. iii. 134 (1790) = praec.

VANILLOSMA, Spach, Hist. Vég. Phan. x. 39 (1841); xiv. 424 (1848) = **Piptocarpha**, R. Br. (Compos.).
acuminatum, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 128 = P. leprosa.
albidum, Mart. ex Baker, l. c. = P. leprosa.
argyrophyllum, Mart. ex Baker, l. c. 121 = P. lucida.
bicolor, Mart. ex Baker, l. c. 131 = P. Lundiana.
candicans, Mart. ex Baker, l. c. 128 = P. leprosa.
cinereum, Mart. ex Baker, l. c. 23 = Vernonia discolor.
firmum, Mart. ex Sch. Bip. in Pollichia, xx–xxi. (1863) 424 = P. rotundifolia.
multiflorum, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 21, 22 = Vernonia oppositifolia.
obtusum, Mart. ex Sch. Bip. in Pollichia, xx–xxi. (1863) 415 = P. pyrifolia.
pyrifolium, Mart. ex Sch. Bip. l. c. = P. pyrifolia.
rufulum, Mart. ex Sch. Bip. l. c. 417 = P. axillaris.

VANILLOSMOPSIS, Sch. Bip. in Pollichia, xviii–xix. (1861) 166; xx–xxi. (1863) 397. *COMPOSITAE*, Benth. & Hook. f. ii. 226.
ALBERTINIA, DC. Prod. v. (1836) 82, partim.
albertinioides, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 168.—Bras.
arborea, *Baker, in Mart. Fl. Bras.* vi. ii. 16.—Bras.
bicolor, Sch. Bip. in Pollichia, xviii–xix. (1861) 168 = Eremanthus bicolor.
brasiliensis, *Sch. Bip. l. c.* xx–xxi. (1863) 401.—Bras.
Candolleana, Sch. Bip. l. c. xviii–xix. (1861) 168 = erythropappa.
capitata, *Sch. Bip. l. c.* 167, partim.—Bras.
capitata, Sch. Bip. l. c., partim = discolor.
capitata, Sch. Bip. l. c. xx–xxi. (1863) 399, partim = arborea.
discolor, *Benth. & Hook. f. Gen.* ii. 226; *Baker, in Mart. Fl. Bras.* vi. ii. 17.—Bras.
erythropappa, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 167.—Bras.
glomerata, Sch. Bip. l. c. = erythropappa.
Pohlii, *Baker, in Mart. Fl. Bras.* vi. ii. 18.—Bras.
polycephala, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 168.—Bras.
saligna, Sch. Bip. l. c. = polycephala.
syncephala, Sch. Bip. l. c. = Lychnophora reticulata.

VANTANEA, Aubl. Pl. Gui. i. 572. t. 229 (1775).
HUMIRIACEAE, Benth. & Hook. f. i. 246.
HELLERIA, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 38. t. 7.
LEMNISCIA, Schreb. Gen. 358 (1789).
guianensis, *Aubl. Pl. Gui.* i. 572. t. 229.—Guian.
guianensis, Poir. Encyc. viii. 334 = parviflora.
minor, *Benth. in Hook. Kew Journ.* v. (1853) 99.—Guian.
obovata, *Benth. l. c.*—Bras.
ovalifolia, *Benth. l. c.*—Bras.

VANTANEA:—
paniculata, *Urban, in Mart. Fl. Bras.* xii. II. 450.—Bras.
parviflora, *Lam. Journ. Hist. Nat. Par.* i. (1792) 145.—Guian.

VARASIA, Phil. Fl. Atac. 35. t. 5 (1860) = **Gentiana**, Linn.
podocarpa, Phil. l. c. 36 = G. podocarpa.
sessilis, Phil. in Anal. Univ. Chil. (1861) I. 534; et in Linnaea, xxxiii. (1864–65) 179.—Chili.

VARECA, Gaertn. Fruct. i. 290. t. 60 (1788) = **Casearia**, Jacq. (Samyd.).
zeylanica, Gaertn. l. c. = C. esculenta.

VARECA, Roxb. Hort. Beng. 18 (1814); Fl. Ind. i. 647 (1832) = **Alsodeia**, Thou. (Violar.).
heteroclita, Roxb. ll. cc. 18; 648 = A. Roxburghii.
lanceolata, Roxb. ll. cc. 86; 648 = A. lanceolata.
moluccana, Roxb. ll. cc. 18; 647.—Ins. Molucc.

VARENNEA, DC. Prod. ii. 522 (1825) = **Eysenhardtia**, H. B. & K. (Legumin.).
polystachya, DC. l. c. = E. amorphoides.

VARGASIA, Bert. ex Spreng. Syst. ii. 283 (1825) = **Thouinia**, Poit. (Sapind.).
glabra, Bert. l. c. 388.—Ins. S. Doming.
tomentosa, Bert. l. c.—Ins. S. Doming.

VARGASIA, DC. Prod. v. 676 (1836) = **Galinsoga**, Ruiz & Pav. (Compos.).
caracasana, DC. l. c.—Venezuela.

VARGASIA, Ersnt, Várgas Consid. como Bot. 23 (1877). *TERNSTROEMIACEAE.*
tremadena, *Ernst, l. c.*—Venezuela.
viridiflora, *Ernst, l. c.*—Venezuela.

VARILLA, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 106. *COMPOSITAE*, Benth. & Hook. f. ii. 363.
mexicana, *A. Gray, l. c.*—Mexic.
texana, *A. Gray, Pl. Wright.* i. 123.—Texas.

VARINGA, Rafin. Sylva Tellur. 58 (1838) = **Ficus**, Tourn. (Urtic.).
repens, Rafin. l. c. = F. pumila.

VARRONIA, P. Br. Hist. Jamaic. 172 (1756); Linn. Syst. ed. X. 916 (1759) = **Cordia**, Linn. (Borag.).
abyssinica, DC. Prod. ix. 469 = C. Myxa.
alba, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. alba.
alnifolia, Hornem. Hort. Hafn. i. 227 = C. alnifolia.
angustifolia, West, Bidr. Ste-Croix, 202 = C. angustifolia.
Bonplandii, Desv. Journ. Bot. i. (1808) 275 = C. Bonplandii.
bullata, Linn. Syst. ed. X. 916 = C. globosa.
calyptrata, DC. Prod. ix. 469 = C. alba.
canescens, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 203.—Ins. Galapag.
chamaedryoides, Willd. ex Roem. & Schult. Syst. iv. 801, in obs. = C. chamaedryoides.
corymbosa, Desv. Journ. Bot. i. (1808) 275 = C. ulmifolia.
crenata, Ruiz & Pav. Fl. Per. ii. 23 = C. lantanoides.
crenulata, Pav. ex DC. Prod. ix. 492 = C. crenulata.
curassavica, Bert. ex DC. l. c. 491; Mayc. Fl. Barb. 84 = C. cylindristachya.
curassavica, Jacq. Enum. Pl. Carib. 14 = C. curassavica.
curassavica, Lam. Illustr. i. 419 = C. angustifolia.
curassavica, Vell. Fl. Flum. 79; ii. t. 85 = C. salicina.
cylindristachya, Ruiz & Pav. Fl. Per. ii. 23 = C. cylindristachya.
dasycephala, Desv. Journ. Bot. i. (1808) 274 = C. dasycephala.
dasycephala, Spreng. ex DC. Prod. ix. 496 = C. globosa.
dichotoma, Ruiz & Pav. Fl. Per. ii. 23 = C. bifurcata.
divaricata, [Willd.] Roem. & Schult. Syst. iv. 802 = C. divaricata, dentata.
ferruginea, Lam. Illustr. i. 418 = C. ferruginea.
flava, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 201.—Ins. Galapag.
floribunda, Desv. Journ. Bot. i. (1808) 267 = C. floribunda.
geniculata, Pers. Syn. i. 167 = C. mirabiloides.

VARRONIA:—
globosa, Desv. Journ. Bot. i. (1808) 276 = Cordia ulmifolia.
globosa, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. globosa.
globosa, Vell. Fl. Flum. 79; ii. t. 84 (Quid?).—Bras.
globulifera, Willd. ex Cham. in Linnaea, iv. (1829) 491 = C. Bonplandii.
grandiflora, Desv. Journ. Bot. i. (1808) 273 = C. grandiflora.
guasumaefolia, Desv. l. c. 276 (*guasumaefolia*) = C. urticifolia.
guianensis, Desv. l. c. 270 = C. guianensis.
Humboldtii, Willd. ex Roem. & Schult. Syst. iv. 801, in obs. = C. floribunda.
humilis, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. ulmifolia.
integerrima, Stokes, Bot. Mat. Med. i. 424 = C. Loureiri.
integrifolia, Desv. Journ. Bot. i. (1808) 271 = C. integrifolia.
lanceolata, Desv. l. c. 277 = C. lanceolata.
lanceolata, Willd. ex Cham. in Linnaea, iv. (1829) 492 = C. mariquitensis.
lantanoides, Willd. ex Cham. l. c. = C. grandiflora.
leucophlyctis, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 203 = C. leucophlyctis.
Lima, Desv. Journ. Bot. i. (1808) 278 = C. Lima.
linearis, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 204 = C. linearis, *Hook.*
linearis, Pav. ex DC. Prod. ix. 493 = C. linearis, *A. DC.*
lineata, Linn. Syst. ed. X. 916 = C. ulmifolia.
macrocephala, Desv. Journ. Bot. i. (1808) 273 = C. macrocephala.
macrocephala, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 78 = C. Neowediana.
macrostachia, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. macrostachya.
macrostachya, Ruiz & Pav. Fl. Per. ii. t. 147 = C. cylindristachya.
martinicensis, Aubl. Pl. Gui. i. 232 = C. Aubletii.
martinicensis, Bert. ex DC. Prod. ix. 483 = C. salvifolia.
martinicensis, Desf. Tabl. ed. I. 71 = C. ferruginea.
martinicensis, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. martinicensis.
martinicensis, Lam. Illustr. i. 419 = C. curassavica.
martintiscensis, Crantz, Inst. ii. 380, sphalm. = *martinicensis*, Jacq.
microcephala, Willd. ex Roem. & Schult. Syst. iv. 801, in obs. = C. microcephala.
microphylla, Desv. Journ. Bot. i. (1808) 275 = C. microphylla.
mirabiloides, Jacq. Enum. Pl. Carib. 14; Select. Am. 41 = C. mirabiloides.
mollis, Desf. Tabl. ed. I. 220 = Melochiae species?
mollis, Vest, ex DC. Prod. ix. 491 = C. cylindristachya.
monosperma, Jacq. Hort. Schoenb. i. 18 = C. ulmifolia.
obliqua, Ruiz & Pav. Fl. Per. ii. 24 = C. peruviana.
paniculata, Wikst. in Vet. Akad. Handl. Stockh. (1827) 59 = C. ulmifolia.
parviflora, Ortega, Hort. Matr. Dec. 86 = C. bifurcata.
pilulifera, Willd. ex Roem. & Schult. Syst. iv. 801, in obs. = C. globosa.
polycephala, Lam. Illustr. i. 418 = C. ulmifolia.
pyrifolia, Willd. ex Roem. & Schult. Syst. iv. 802 = C. ferruginea.
revoluta, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 204 = C. revoluta.
rotundifolia, A. DC. in DC. Prod. ix. 469 = C. rotundifolia.
rugosa, Willd. ex Roem. & Schult. Syst. iv. 801, in obs. = C. canescens.
salvifolia, Balb. Cat. Hort. Taur. (1813) 79 = C. cylindristachya.
scaberrima, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 202.—Ins. Galapag.
scaberrima, Bert. ex Spreng. Syst. i. 654 = C. Lima.
Scouleri, Anderss. in Vet. Akad. Handl. Stockh. 1853 (1855) 204 = C. Scouleri.
sinensis, Lour. Fl. Cochinch. 138 = C. Loureiri.
sphaerocephala, Willd. ex Roem. & Schult. Syst. iv. 801, in syn. = C. dasycephala.
spicata, Salzm. ex DC. Prod. ix. 489 = C. cylindristachya.
tomentosa, Lam. Illustr. i. 419 = C. tomentosa.
ulmifolia, Dum.-Cours. Bot. Cult. ed. I. ii. 148, in obs. = C. monosperma.

VARTHEIMIA, Benth. & Hook. f. Gen. ii. 333 (1873), sphalm. = seq.

VARTHEMIA, DC. Prod. v. 473 (1836) = **Iphiona**, Cass. (Compos.).
arabica, Boiss. Diagn. Ser. I. vi. 74 = I. arabica.
arabica, Thoms. in Speke, Journ. App. 638 = Pulicaria Grantii.
candicans, Boiss. Fl. Orient. iii. 212.—Egypt.
conyzoides, Boiss. Diagn. Ser. II. iii. 10.—Arab.
iphionoides, Boiss. & Blanche, l. c. 9 = I. iphionoides.
Kotschyi, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 161, 307 = Vicoa auriculata.
montana, Boiss. Fl. Orient. iii. 212 = *conyzoides.*
persica, DC. Prod. v. 473 = I. persica.

VASARGIA, Steud. Nom. ed. II. 745 (1841), sphalm. = VARGASIA, DC. = **Galinsoga**, Ruiz & Pav. (Compos.).
caracasana, Steud. l. c. (= *Vargasia caracasana*).—Venezuela.

VASCOA, DC. Mém. Légum. 186 (1825); Prod. ii. 119 (1825) = **Rafnia**, Thunb. (Legumin.).
acuminata, E. Mey. in Linnaea, vii. (1832) 148 = R. perfoliata.
amplexicaulis, DC. Mém. Légum. 187; Prod. ii. 119 = R. amplexicaulis.
perfoliata, DC. ll. cc. = R. perfoliata.

VASCOA, DC. Prod. i. 337, in obs. (1824) = **Mundtia**, H. B. & K. (Polygal.).

VASCONCELLEA, A. St. Hil. Mém. Réséd. ii. 12 (1837) = **Carica**, Linn. (Passifl.).
aurantiaca, Regel, Gartenfl. xxv. (1876) 184.—N. Granat.
Boissieri, A. DC. in DC. Prod. xv. I. 415 = C. Boissieri.
candicans, A. DC. l. c. 417 = C. candicans.
cauliflora, A. DC. l. c. 415 = C. cauliflora.
cestriflora, A. DC. l. c. 418.—N. Granat.
chilensis, Planch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 259.—Chili.
glandulosa, A. DC. in DC. Prod. xv. I. 417 = C. glandulosa.
Goudotiana, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 192.—N. Granat.
heterophylla, A. DC. in DC. Prod. xv. I. 417 = C. heterophylla.
Hookeri, A. DC. l. c. 418.—Hab.?
lanceolata, A. DC. l. c. 416 = C. lanceolata (in Addend.).
Manihot, Triana & Planch. in Ann. Sc. Nat. Sér. V. xvii. (1873) 191.—N. Granat.
microcarpa, A. DC. in DC. Prod. xv. I. 418 = C. microcarpa.
monoica, A. DC. l. c. = C. monoica.
parviflora, A. DC. l. c. 417.—Peruv.
peltata, A. DC. l. c. 416 = C. peltata.
pubescens, A. DC. l. c. 419.—Peruv.
quercifolia, A. St. Hil. Mém. Réséd. ii. 13 = C. quercifolia (in Addend.).

VASCONCELLIA, Mart. in Flora, xxiv. (1841) II. Beibl. 12 (Herb. Fl. Bras. 252) = **Arrabidaea**, DC. (Bignon.).
acutifolia, Mart. ex DC. Prod. ix. 185 = A. acutifolia.
desertorum, Mart. l. c. = A. desertorum.
fasciculata, Mart. l. c. = A. fasciculata.
florida, Mart. l. c. 184 = A. florida.
Sego, Mart. in Flora, xxiv. (1841) II. Beibl. 12 = A. Agnus-castus.
subincana, Mart. ex DC. Prod. ix. 184 = Bignonia subincana.
tuberculata, Mart. l. c. = A. tuberculata.

VASCONCELLOSIA, Caruel, in Nuov. Giorn. Bot. Ital. viii. (1876) 22. *PASSIFLOREAE.*
hastata, *Caruel, l. c.*—Bras.

VASCONELLA, Regel, Gartenfl. xxv. (1876) 184 = VASCONCELLEA, A. St. Hil. = **Carica**, Linn. (Passifl.).

VASEYA, Thurb. in Proc. Acad. Nat. Sc. Philad. (1863) 79 = **Muehlenbergia**, Schreb. (Gramin.).
comata, Thurb. l. c. = M. comata.

VASIVAEA, Baill. Adansonia, x. (1872) 193. *TILIACEAE*.
alchorneoides, *Baill. l. c.* 194.—Am. trop.

VASQUEZIA, Phil. Fl. Atac. 31. t. 5 (1860) = **Villanova**, Lag. (Compos.).
biternata, Phil. l. c. 31.— Chili.

VATAIREA, Aubl. Pl. Gui. ii. 755. t. 302 (1775) = **Pterocarpus**, Linn. (Legumin.).
guianensis, Aubl. l. c.—Guian.

VATERIA, Linn. Gen. ed. I. 153 (1737). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 193.
HEMIPHRACTUS, Turcz. in Bull. Soc. Nat. Mosc. (1859) I. 262.
STEMONOPORUS, Thw. in Hook. Kew Journ. vi. (1854) 67. t. 2.
acuminata, *Hayne, Arzn. Gew.* xi. sub t. 5.—Ind. or.
acuminata, Thw. Enum. Pl. Zeyl. 403 = jucunda.
affinis, *Thw. l. c.*—Zeylan.
canaliculata, *Thw. l. c.* 403.—Zeylan.
ceylanica, *Wight, Illustr.* i. 88.—Zeylan.
cordifolia, Thw. Enum. Pl. Zeyl. 404 = Monoporandra cordifolia.
disticha, Thw. l. c. = Vatica disticha.
elegans, Thw. l. c. = Monoporandra elegans.
flexuosa, *Lour. Fl. Cochinch.* 334.—China.
Gardneri, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
indica, Gaertn. f. Fruct. iii. 53. t. 189 = acuminata.
indica, *Linn. Sp. Pl.* 515.—Ind. or.
jucunda, *Thw. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 314.—Zeylan.
lanceaefolia, A. DC. in DC. Prod. xvi. II. 625, sphalm. = nitida.
lanceaefolia, Roxb. Fl. Ind. ii. 601 = Vatica lanceaefolia.
lanceolaria, Roxb. Hort. Beng. 42 = Vatica lanceaefolia.
lanceolata, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
lanceolata, Wight & Arn. Prod. 84; Wall. Cat. n. 4405 = Vatica lanceaefolia.
lancifolia, Thw. Enum. Pl. Zeyl. 404 = nitida.
malabarica, Blume, Mus. Bot. Lugd. Bat. ii. 29 = indica.
Moonii, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
nervosa, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 206.—Zeylan.
nitida, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
oblongifolia, *Thw. l. c.*—Zeylan.
papuana, Dyer, ex Hemsl. Bot. Voy. Challenger, i. III. 123 = Vatica papuana.
pauciflora, Walp. Rep. v. 126 = Vatica pauciflora.
petiolaris, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
Rassak, Walp. Rep. v. 126 = Vatica Rassak.
reticulata, *Thw. Enum. Pl. Zeyl.* 403.—Zeylan.
rigida, *Thw. l. c.*—Zeylan.
Roxburghiana, Wight, ex Arn. in Ann. Nat. Hist. Ser. I. iii. (1839) 155 = Vatica Roxburghiana.
scabriuscula, Thw. Enum. Pl. Zeyl. 404 = Vatica scabriuscula.
seychellarum, *Dyer, in Baker, Fl. Maurit.* 526.—Ins. Seychelles.
Wightii, Thw. Enum. Pl. Zeyl. 403 = ceylanica.
zeylanica, Walp. Rep. v. 126 = ceylanica.

VATICA, Linn. Mant. ii. 152 (1771). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 192.
ISAUXIS, Reichb. Nom. 210 (1841).
PTERANTHERA, Blume, Mus. Bot. Lugd. Bat. ii. 30. 1852 (1856).
RETINODENDRON, Korth. Verh. Nat. Gesch. Bot. 55. t. 8 (1839).
SEIDLIA, Kostel. Allg. Med. Pharm. Flora, v. 1945 (1836).
SUNAPTEA, Griff. Notul. iv. 56 (1854).
SYNAPTEA, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 65.
acuminata, A. DC. in DC. Prod. xvi. II. 622 = Vateria jucunda.
affinis, *Thw. Enum. Pl. Zeyl.* 404.—Zeylan.
africana, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 15. —Afr. trop.
apteranthera, Blanco, Fl. Filip. ed. II. 281 = Shorea Mangachapoi.
astrotricha, *Hance, in Journ. Bot.* xiv. (1876) 241.—Cambodia.
bancana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 348.—Ins. Banca.
bantamensis, *Benth. & Hook. f. Gen.* i. 192.—Ins. Bantam.
Canaca, Buch.-Ham. ex Wall. Cat. n. 4405 B = lanceaefolia.
canaliculata, A. DC. in DC. Prod. xvi. II. 621 = Vateria canaliculata.
chinensis, *Linn. Mant.* ii. 242.—Ind. or.
cordifolia, Thw. Enum. Pl. Zeyl. 404 (sp. dub.).—Zeylan.
disticha, *A. DC. in DC. Prod.* xvi. II. 620.—Zeylan.
eximia, Miq. Fl. Ind. Bat. Suppl. 486 = Shorea eximia.
faginea, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 301.—Burma.
Gardneri, A. DC. in DC. Prod. xvi. II. 622 = Vateria Gardneri.
grandiflora, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 301.—Burma; Malaya.
Helferi, *Dyer, l. c.* 302.—Burma.
laccifera, Wight & Arn. Prod. 84 = Shorea Talura.
lanceaefolia, *Blume, Mus. Bot. Lugd. Bat.* ii. 31.—Reg. Himal.; Burma.
lanceolata, A. DC. in DC. Prod. xvi. II. 621 = Vateria lanceolata.
laurifolia, Steud. Nom. ed. II. ii. 745, sphalm. = Shorea laccifera.
lepidota, Korth. Verh. Nat. Gesch. Bot. 73 = Shorea lepidota.
Maingayi, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 302.—Malacca.
Mangachapoi, Blanco, Fl. Filip. ed. I. 401 = Shorea Mangachapoi.
melanoxylon, *Benth. & Hook. f. Gen.* i. 192.—Borneo.
Moonii, A. DC. in DC. Prod. xvi. II. 622 = Vateria Moonii.
nitida, A. DC. l. c. = Vateria nitida.
oblonga, A. DC. l. c. 621 = Vateria oblongifolia.
oblongifolia, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 160.—Borneo.
obscura, *Trimen, in Journ. Bot.* xxiii. (1885) 203.—Zeylan.
obtusa, Steud. Nom. ed. II. ii. 745 = Shorea obtusa.
ovalis, Korth. Verh. Nat. Gesch. Bot. 73 = Shorea ovalis.
pallida, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 302.—Ins. Penang.
papuana, *Dyer, in Journ. Bot.* xvi. (1878) 100.—N. Guin.
pauciflora, *Blume, Mus. Bot. Lugd. Bat.* ii. 31. t. 7.—Sumatra.
petiolaris, A. DC. in DC. Prod. xvi. II. 621 = Vateria petiolaris.
Rassak, *Blume, Mus. Bot. Lugd. Bat.* ii. 31.—Borneo.
reticulata, A. DC. in DC. Prod. xvi. II. 620 = Vateria reticulata.
rigida, A. DC. l. c. 621 = Vateria rigida.
robusta, Steud. Nom. ed. II. ii. 745 = Shorea robusta.
Roxburghiana, Blume, Mus. Bot. Lugd. Bat. ii. 31. t. 7 = chinensis.
scabriuscula, *A. DC. in DC. Prod.* xvi. II. 620.—Zeylan.
Scaphula, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 301.—Burma.
Scaphula, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 21 = Shorea Mangachapoi.
Schouteniana, Scheff. in Tijdschr. Nederl. Ind. xxxii. (1871) 408 = bancana.
sinensis, J. F. Gmel. Syst. 748 = *chinensis*.
stipulosa, Miq. Fl. Ind. Bat. Suppl. 486 (Shoreae sp.). —Sumatra.
sublacunosa, Miq. l. c. = Shorea eximia.
Thwaitesii, A. DC. in DC. Prod. xvi. II. 621 = Vateria affinis.
trigyna, Griff. Notul. iv. 514 = Dipterocarpus Griffithii.
Tumbugaia, Wight & Arn. Prod. 84 = Shorea Tumbuggaia.
venulosa, *Blume, Mus. Bot. Lugd. Bat.* ii. 32.—Borneo.
Wightii, A. DC. in DC. Prod. xvi. II. 620 = Vateria ceylanica.
Zollingeriana, *A. DC. l. c.* 618.—Malaya.

VATKEA, Hildeb. & O. Hoffm. in Verh. Bot. Ver. Brandenb. xxii. (1880) 45, 78 = **Martynia**, Houst. (Pedalin.).
diandra, O. Hoffm. in Linnaea, xliii. (1880–82) 554 (in indice) = M. diandra.

VAUANTHES, Haw. Rev. Pl. Succ. 18 (1821) = **Grammanthes**, DC. (1828) (Crassul.).
chloraeflora, Haw. l. c. 19 = G. gentianoides.
chloraefolia, Steud. Nom. ed. II. 746, sphalm. = praec.

VAUPELLIA, Griseb. Fl. Brit. W. Ind. 460 (1861) = **Pentarhaphia**, Lindl. (Gesner.).
calycina, Griseb. l. c. = P. calycina.

VAUQUELINIA, Correa, ex Humb. & Bonpl. Pl. Aequin. i. 140. t. 40 (1808). *ROSACEAE*, Benth. & Hook. f. i. 615.
corymbosa, *Correa, ex Humb. & Bonpl. l. c.*—Mexic.
corymbosa, Torr. Bot. Mex. Bound. 64 = Torreyi.
Karwinskyi, *Maxim. in Act. Hort. Petrop.* vi. (1879) 236.—Mexic.
Torreyi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 147. —Mexic.

VAUTHIERA, A. Rich. Ess. Fl. N. Zél. 106. t. 20 (1832) = **Cladium**, P. Br. (Cyperac.).
australis, R. Rich. l. c. 107. t. 20.—N. Zel.

VAVAEA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 212. *MELIACEAE*, Benth. & Hook. f. i. 331.
amicorum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 212.—Ins. Amicor.
Harveyi, *Seem. Fl. Vit.* 35.—Ins. Fiji.
vitiensis, Seem. l. c. = amicorum.

VAVANGA, Rohr, in Skrivt. Nat. Selsk. ii. I. (1792) 207, 208. t. 7 = **Vangueria**, Juss. (Rubiac.).
chinensis, Rohr, l. c. 207 = Vangueria madagascariensis.
edulis, Vahl, l. c. 208 = Vangueria madagascariensis.

VAZQUEZIA, Pritz. Ic. Ind. ii. 282 (1865), err. typ. = VASQUEZIA, Phil. = **Villanova**, Lag. (Compos.).

VEATCHIA, A. Gray, in Bull. Calif. Acad. i. n. I. (1884) 4; et in Proc. Am. Acad. xx. (1885) 290. *ANACARDIACEAE*.
cedrosensis, *A. Gray, ll. cc.* 4; 291.—Calif.

VEATCHIA, Kellogg, in Proc. Calif. Acad. ii. (1863) 11 = **Brodiaea**, Sm. (Liliac.).
crystallina, Kellogg, l. c. = B. lactea.

VEDELA, Adans. Fam. ii. 502 (1763) (Quid?).

VEGELIA, Neck. Elem. iii. 385 (1790) = WEIGELA, Thunb. = **Diervilla**, Tourn. (Caprifol.).

VEITCHIA, Lindl. in Gard. Chron. (1861) 265 = **Picea**, Link (Conif.).
japonica, Lindl. l. c. = P. Alcockiana.

VEITCHIA, H. Wendl. in Seem. Fl. Vit. 270. t. 81 (1868). *PALMAE*, Benth. & Hook. f. iii. 887.
Canterburyana, F. Muell. Fragm. vii. 101 = Hedyscepe Canterburyana.
Joannis, *H. Wendl. in Seem. Fl. Vit.* 271.—Ins. Fiji.
spiralis, *H. Wendl. l. c.* 270.—N. Hebrid.
Storckii, *H. Wendl. l. c.*—Ins. Fiji.
subglobosa, *H. Wendl. l. c.* 272.—Ins. Fiji.

VELAEA, DC. Mém. Ombell. 61 (1829); Prod. iv. 230 (1830) = **Arracacia**, Bancr. (Umbell.).
decumbens, Benth. Pl. Hartw. 38 = A. decumbens.
toluccensis, DC. Prod. iv. 231 = A. tolucensis.

VELAGA, Adans. Fam. ii. 398 (1763); Gaertn. Fruct. ii. 245. t. 133 (1791) = **Pterospermum**, Schreb. (Stercul.).
globosa, Gaertn. l. c. 246 = Lagerstroemia indica.
locarpa, Gaertn. l. c. 245 = P. Heyneanum.

VELARUM, Reichb. in Moessl. Handb. ed. II. ii. 1103, 1165 (1828) = **Sisymbrium**, Tourn. (Crucif.).
officinale, Reichb. l. c. 1165 = S. officinale.
polyceratum, Reichb. l. c. = S. polyceratium.
supinum, Reichb. l. c. = S. supinum.

VELASQUEZIA, Pritz. Ic. Ind. 1147 (1855), sphalm. = VELLASQUEZIA, Bertol. = **Triplaris**, Linn. (Polygon.).

VELEZIA, Linn. Sp. Pl. ed. I. 332 (1753). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 144.
arabica, *Hochst. & Steud. ex Steud. Nom.* ed. II. ii. 746, nomen.—Arab.
clavata, Urv. in Mém. Soc. Linn. Par. i. (1822) 284 = quadridentata.
fasciculata, *Boiss. Diagn.* Ser. I. viii. 92.—Syria.
hispida, *Boiss. et Bal. l. c.* II. v. 57.—Phrygia.
latifolia, Eschsch. in Mém. Acad. Pétersb. x. (1823) 286 = Frankenia grandifolia.
quadridentata, *Sibth. & Sm. Fl. Graec. Prod.* i. 283; *Fl. Græc.* iv. 81. t. 391.—Graec.; As. Min.
rigida, *Linn. Sp. Pl.* 332.—Reg. Mediterr.; Oriens.

VELLA, Linn. Coroll. Gen. 12 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 85.
annua, Linn. Sp. Pl. 641 = Carrichtera Vellae.
aspera, Pers. Syn. ii. 185 = Boleum asperum.
glabrescens, *Coss. Illustr. Fl. Atlant.* i. 71. t. 48.—Alger.
hispida, Vahl, ex DC. Syst. ii. 641 = Boleum asperum.
integrifolia, Salisb. Prod. 265 = V. Pseudocytisus.
Pseudocytisus, *Linn. Sp. Pl.* 641.—Hispan.
spinosa, *Boiss. Elench.* 14.—Hispan.
tatarica, Pall. Reise, iii. 521 = Euclidium tataricum.
tenuissima, Pall. l. c. 740 = Euclidium tataricum.

VELLASQUEZIA, Bertol. Fl. Guatim. 39. t. 11 (1840) = **Triplaris**, Loefl. (Polygon.).
melanodendron, Bertol. l. c. 40. t. 11.—Guatemala.

VELLEA, D. Dietr. ex Steud. Nom. ed. II. ii. 746 (1841), sphalm. = VELAEA, DC. = **Arracacia**, Bancr. (Umbell.).

VELLEIA, Sm. in Trans. Linn. Soc. iv. (1798) 217. *GOODENOVIEAE*, Benth. & Hook. f. ii. [II.] 538.
EUTHALES, R. Br. Prod. 579 (1810).
MENOCERAS, Lindl. Veg. Kingd. 695 (1847).
arguta, R. Br. Prod. 580 = paradoxa.
connata, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 18.—Austral.
cycnopotamica, *F. Muell. Fragm.* vi. 7.—Austral.
Daviesii, *F. Muell. l. c.* x. 10.—Austral.
dichotoma, Auct. ex DC. Prod. vii. 518 = lyrata.
discophora, *F. Muell. Fragm.* x. 10.—Austral.
lanceolata, Lindl. Swan River App. 26 = Goodenia filiformis.
lyrata, *R. Br. Prod.* 580.—Austral.
macrocalyx, *De Vriese, in Mitch. Journ. Trop. Austral.* 258.—Austral.
macrophylla, *Benth. Fl. Austral.* iv. 47.—Austral.
macroplectra, *F. Muell. Fragm.* xii. 22.—Austral.
montana, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 265.—Austral.
panduriformis, *A. Cunn. ex Benth. Fl. Austral.* iv. 46.—Austral.
paradoxa, *R. Br. Prod.* 580.—Austral.
perfoliata, *R. Br. l. c.* 581.—Austral.
pubescens, *R. Br. l. c.*—Austral.
spathulata, *R. Br. l. c.* 580.—Austral.
spathulata, A. Juss. in Ann. Mus. Par. xviii. (1811) 17. t. 1 = lyrata.
trinervis, *Labill. Nov. Holl. Pl.* i. 54. t. 77.—Austral.

VELLEJA, Schrad. Neues Journ. iii. 1. (1809) 118 = praec.

VELLERUCA, Pomel, Mat. Fl. Atl. 12 (1860) = **Eruca**, Tourn. (Crucif.).
longistyla, Pomel, l. c. = E. longistyla.
setulosa, Pomel, l. c. = E. setulosa.
vesicaria, Pomel, l. c. = E. vesicaria.

VELLEYA, Walp. Ann. i. 471 (1848–49) = **Velleia**, Sm. (Gooden.).

VELLOSIA, Spreng. Syst. iii. 338 (1826) = seq.

VELLOZIA, Vand. Fl. Lusit. 32. t. 2 (1788). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 739.
SCHNIZLEINIA, Steud. ex Hochst. in Flora, xxvii. (1844) 31.
TALBOTIA, Balf. in Trans. Bot. Soc. Edinb. ix. (1868) 190.
XEROPHYTA, Juss. Gen. 50 (1789).
abietina, *Mart. Nov. Gen. et Sp.* i. 14. t. 6.—Bras.
albiflora, *Pohl, Pl. Bras. Ic.* i. 121. t. 96.—Bras.
aloaefolia, *Mart. Nov. Gen. et Sp.* i. 15. t. 7.—Bras.
alutacea, Pohl, Pl. Bras. Ic. i. 130 = variabilis?

VELLOZIA:—
asperula, *Mart. Nov. Gen. et Sp.* i. 15. t. 8.—Bras.
barbaceniaefolia, *Seub. in Mart. Fl. Bras.* iii. 1. 79.—Bras.
brevifolia, *Seub. l. c.* 84. t. 10. f. 2.—Bras.
breviscapa, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
candida, *Mikan, Delect. Fl. et Faun. Bras.* fasc. ii. [t. 7].—Bras.
capillaris, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 264.—Afr. trop.
caruncularis, *Mart. ex Seub. in Mart. Fl. Bras.* iii. 1. 78. t. 8. f. 1.—Bras.
cinerascens, *Mart. ex Seub. l. c.* 74.—Bras.
coerulescens, Hort. Belg. ex Gumbl. in Gard. Chron. (1874) II. 623, 657 = Barbacenia purpurea.
compacta, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
crassicaulis, Mart. l. c. 292 = albiflora.
cryptantha, *Seub. in Mart. Fl. Bras.* iii. 1. 80.—Bras.
dasypus, *Seub. l. c.* 81.—Bras.
elegans, *Talbot, ex Balf. in Trans. Bot. Soc. Edinb.* ix. (1868) 79, nomen; *Oliver, ex Hook. f. Bot. Mag.* t. 5803.—Afr. austr.
epidendroides, *Mart. in Schult. f. Syst.* vii. 292.—Bras.
flavicans, Mart. l. c. 293 = squamata.
glabra, Mikan, ex Spreng. Syst. iii. 338 = phalocarpa.
glauca, *Pohl, Pl. Bras. Ic.* i. 125. t. 100.—Bras.
glochidea, *Pohl, l. c.* 129.—Bras.
gracilis, *Seub. in Mart. Fl. Bras.* iii. 1. 81.—Bras.
graminea, *Pohl, Pl. Bras. Ic.* i. 118. t. 93.—Bras.
hemisphaerica, *Seub. in Mart. Fl. Bras.* iii. 1. 80.—Bras.
incurvata, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
intermedia, *Seub. in Mart. Fl. Bras.* iii. 1. 78.—Bras.
lanata, *Pohl, Pl. Bras. Ic.* i. 130.—Bras.
leptophylla, *Seub. in Mart. Fl. Bras.* iii. 1. 84.—Bras.
macrantha, *Lem. Illustr. Hortic.* xii. (1865) *Misc.* 44.—Bras.
maritima, *Vell. Fl. Flum.* 219; v. t. 79.—Bras.
minima, *Pohl, Pl. Bras. Ic.* i. 119. t. 94.—Bras.
ornata, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
papillosa, Pohl, Pl. Bras. Ic. i. 128 = asperula.
phalocarpa, *Pohl, l. c.* 123. t. 98.—Bras.
plicata, *Mart. Nov. Gen. et Sp.* i. 16. t. 9.—Bras.
punctulata, *Seub. in Mart. Fl. Bras.* iii. 1. 82.—Bras.
pusilla, *Pohl, Pl. Bras. Ic.* i. 122. t. 97.—Bras.
resinosa, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
scabra, *Spreng. Syst.* iii. 338.—Bras.
Sellovii, *Seub. in Mart. Fl. Bras.* iii. 1. 75.—Bras.
Spekei, *Baker, in Trans. Linn. Soc.* xxix. (1875) 156.—Afr. trop.
squalida, *Mart. in Schult. f. Syst.* vii. 292.—Bras.
squamata, Auct. ex Steud. Nom. ed. II. ii. 746 = Veratrum Sabadilla.
squamata, *Pohl, Pl. Bras. Ic.* i. 124. t. 99.—Bras.
squarrosa, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 264.—Afr. trop.
stenophylla, *Welw. ex Baker, l. c.* 265.—Afr. trop.
sulphurea, *Pohl, Pl. Bras. Ic.* i. 120. t. 95.—Bras.
Talboti, Balf. in Trans. Bot. Soc. Edinb. ix. (1868) 190 = elegans.
taxifolia, *Mart. ex Seub. in Mart. Fl. Bras.* iii. 1. 75.—Bras.
tenella, Mart. in Schult. f. Syst. vii. 293 = graminea.
tertia, Spreng. Neue Entdeck. ii. 108 = candida.
tomentosa, *Pohl, Pl. Bras. Ic.* i. 130.—Bras.
tragacantha, *Mart. ex Seub. in Mart. Fl. Bras.* iii. 1. 75.—Bras.
triquetra, *Pohl, Pl. Bras. Ic.* i. 129.—Bras.
tubiflora, H. B. & K. Nov. Gen. et Sp. vii. 155 = Barbacenia tubiflora (in Addend.).
variabilis, *Mart. in Schult. f. Syst.* vii. 293.—Bras.
velutina, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 265.—Afr. trop.
verruculosa, *Mart. in Schult. f. Syst.* vii. 293.—Bras.

VELLOZOA, Lem. Jard. Fleur. iv. (1853), in obs. ad t. 390 = praec.

VELPEAULIA, Gaudich. Bot. Voy. Bonite, t. 109 (1844–52) = **Dolia**, Lindl. (Convolv.).
alibrexioides, Gaudich. l. c.—Hab. ?

VELTHAEIMIA, Thunb. Fl. Cap. ii. 278 (1818), sphalm. = seq.

VELTHEIMIA, Gled. in Mém. Acad. Berl. 1769 (1771) 66. *LILIACEAE*, Benth. & Hook. f. iii. 811.
abyssinica, [DC. in] Red. Lil. iv. t. 186 = Kniphofia abyssinica.
bracteata, *Harv. ex Baker, in Journ. Linn. Soc.* xi. (1871) 411.—Afr. austr.
capensis, [DC. in] Red. Lil. iv. t. 193 = viridifolia.
glauca, *Jacq. Hort. Schoenb.* i. 40. t. 77.—Afr. austr.
intermedia, *Sweet, Hort. Brit.* ed. I. 409.—Afr. austr.
media, Donn, Hort. Cantab. ed. IV. 74 = Kniphofia sarmentosa.
pumila, Willd. Sp. Pl. ii. 182 = Kniphofia pumila.
repens, [Ker-Gawl.] Recens. Pl. 18 = Kniphofia sarmentosa.
sarmentosa, Willd. Enum. Hort. Berol. 380 = Kniphofia sarmentosa.
speciosa, Roth, Nov. Pl. Sp. 190 = Kniphofia aloides.
undulata, Moench, Meth. 631 = viridifolia.
Uvaria, Willd. Sp. Pl. ii. 182 = Kniphofia aloides.
viridifolia, *Jacq. Hort. Schoenb.* i. 41. t. 78.—Afr. austr.

VELTIS, Adans. Fam. ii. 116 (1763) = **Centaurea**, Linn. (Compos.).

VEMONIA, Edgew. in Journ. As. Soc. Beng. xvi. (1847) II. 1216, err. typ. = **Vernonia**, Schreb. (Compos.).

VENANA, Lam. Illustr. ii. 99. t. 131 (1800) = **Brexia**, Noronha (1806) (Saxifr.).
madagascariensis, Lam. l. c. = B. madagascariensis.

VENATRIS, Rafin. Fl. Tellur. ii. 47 (1836) = **Aster**, Tourn. (Compos.).
falcata, Rafin. l. c. = A. falcatus.
linifolius, Rafin. l. c. = A. acris.
pauciflorus, Rafin. l. c. = A. pauciflorus.
salicifolius, Rafin. l. c. = A. diffusus?
tenuifolius, Rafin. l. c. = A. tenuifolius.

VENDREDIA, Baill. Hist. Pl. viii. 264 (1882) = **Rhetinodendron**, Meissn. (Compos.).
Berterii, Baill. l. c. 265 = R. Berteroi.

VENEGASIA, DC. Prod. vi. 43 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 397.
PARTHENOPSIS, Kellogg, in Proc. Calif. Acad. v. (1875) 100.
carpesioides, *DC. Prod.* vi. 43.—Calif.

VENEGAZIA, Benth. & Hook. f. Gen. ii. 397 (1873), sphalm. = praec.

VENIDIUM, Less. in Linnaea, vi. (1831) 91. *COMPOSITAE*, Benth. & Hook. f. ii. 459.
ANTROSPERMUM, Sch. Bip. in Flora, xxvii. (1844) 773.
CLEITRIA, Schrad. Ind. Sem. Hort. Gotting. (1831).—Cf. Linnaea, viii. (1833) Litt. 23.
angustifolium, *DC. Prod.* vi. 492.—Afr. austr.
arctotoides, Hort. Hamb. ex Harv. & Sond. Fl. Cap. iii. 463 = hirsutum.
arctotoides, *Less. Syn. Comp.* 33.—Afr. austr.
aureum, *DC. Prod.* vi. 494.—Afr. austr.
calendulaceum, Less. Syn. Comp. 32 = decurrens.
canescens, DC. Prod. vi. 493 = decurrens.
cinerarium, DC. l. c. = perfoliatum.
decurrens, *Less. Syn. Comp.* 32.—Afr. austr.
discolor, *Less. l. c.* 31.—Afr. austr.
erosum, *Harv. in Harv. & Sond. Fl. Cap.* iii. 462.—Afr. austr.
fugax, *Harv. l. c.* 464.—Afr. austr.
hirsutum, *Harv. l. c.* 463.—Afr. austr.
hispidulum, DC. Prod. vi. 493 = Kraussii.
hispidulum, *Less. Syn. Comp.* 34.—Afr. austr.
Kraussii, *Sch. Bip. in Flora*, xxvii. (1844) 773.—Afr. austr.
macrocephalum, *DC. Prod.* vi. 494.—Afr. austr.
macrospermum, *DC. l. c.* 492.—Afr. austr.
microcephalum, *DC. l. c.* 493.—Afr. austr.
perfoliatum, *Less. Syn. Comp.* 30.—Afr. austr.
plantagineum, Less. l. c. = semipapposum.

VENIDIUM:—
puberulum, DC. Prod. vi. 493 = hispidulum.
ramosissimum, *Sch. Bip. in Flora*, xxvii. (1844) 772. —Afr. austr.
scabrum, Less. Syn. Comp. 29 = semipapposum.
Schraderi, *DC. Prod.* vi. 494.—Afr. austr.
semipapposum, *DC. l. c.* 491.—Afr. austr.
spathuligerum, *DC. l. c.* 493.—Afr. austr.
speciosum, *Regel, Ind. Sem. Hort. Petrop.* (1857) 43. —Afr. austr.
subacaule, *DC. Prod.* vi. 493.—Afr. austr.
subcalvum, DC. l. c. 492 = semipapposum.
Weyleyi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 463.— Afr. austr.

VENIERA, Salisb. Gen. Pl. Fragm. 101 (1866) = **Narcissus**, Tourn. (Amaryll.).
tenuior, Salisb. l. c. = N. gracilis.

VENILIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125 = **Scrophularia**, Tourn.
vernalis, Fourr. l. c. = S. vernalis.

VENTENATA, Koel. Desc. Gram. 272 (1802).
GRAMINEAE, Benth. & Hook. f. iii. 1159.
HETERANTHUS, Dum. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 183.
MALYA, Opiz, Seznam, 62 (1852).
avenacea, *Koel. Desc. Gram.* 274.—Europ.
Blanchei, *Boiss. Fl. Orient.* v. 539.—Syria.
bromoides, Koel. Desc. Gram. 273 = avenacea.
dubia, Coss. Expl. Algér. 104 = avenacea.
macra, *Balansa, ex Boiss. Fl. Orient.* v. 540.—As. Min.; Mesopot.
quercetorum, *Boiss. & Bal. Diagn.* Ser. II. iv. 130.— Phrygia.
subenervis, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 305.—Phrygia.
triflora, F. Schultz, ex Nym. Consp. 815 = avenacea.

VENTENATIA, Beauv. Fl. Owar. i. 29. t. 17 (1805) = **Oncoba**, Forsk. (Bixin.).
glauca, Beauv. l. c. 30 = O. glauca.

VENTENATIA, Cav. Ic. iv. 79 (1797), err. typ. = VINTENATIA, Cav.

VENTENATIA, Sm. Exot. Bot. ii. 13. tt. 66, 67 (1805) [1806 ?] = **Stylidium**, Sw. (Campan.).
major, Sm. l. c. 13. t. 66 = S. graminifolium.
minor, Sm. l. c. 15. t. 67 = S. lineare.

VENTENATIA, Tratt. Gen. Pl. Disp. 86 (1802) = **Pedilanthus**, Neck. (Euphorb.).
bracteata, Tratt. l. c. 87 = P. bracteatus.

VENTILAGO, Gaertn. Fruct. i. 223. t. 49 (1788).
RHAMNEAE, Benth. & Hook. f. i. 375.
ENRILA, Blanco, Fl. Filip. ed. I. 709 (1837).
bombaiensis, *Dals. in Hook. Kew Journ.* iii. (1851) 36.—Ind. or.
bracteata, Heyne, ex Wall. Cat. n. 4269 = madraspatana.
buxoides, *Baill. Adansonia*, xi. (1874) 268.— N. Caled.
calyculata, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 124.—Ind. or.
cernua, *Tul. l. c.* 123.—Ins. Rawak.
cinerascens, G. Don, Gen. Syst. ii. 28 = Rhamnus cinerascens.
denticulata, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 417 = calyculata.
ecorollata, *F. Muell. in Wing, South. Sc. Record* (July 1883).—Austral.
lanceata, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 121. —Ins. Seychelles.
leiocarpa, *Benth. in Journ. Linn. Soc.* v. (1861) 77.— As. or.
leptadenia, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 124.—Madag.
lucens, *Miq. Fl. Ind. Bat. Suppl.* 330.—Sumatra.
macrantha, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 123 = calyculata.
maderaspatana, Benth. in Hook. Kew Journ. iv. (1852) 42 = leiocarpa.
maderaspatana, *Gaertn. Fruct.* i. 223. t. 49.—Ind. or.

VENTILAGO:—
maderaspatana, Roxb. Pl. Corom. i. 55. t. 76 = calyculata.
Maingayi, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 631. —Malaya.
monoica, Blanco, Fl. Filip. ed. II. 124 = maderaspatana.
oblongifolia, *Blume, Bijdr.* 1144.—Java.
silhetiana, M. Laws. in Hook. f. Fl. Brit. Ind. i. 631, in syn. = calyculata.
Smithiana, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 125 = calyculata.
sororia, *Hance, in Journ. Bot.* xiv. (1876) 243.— China.
sulphurea, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 125 = calyculata.
viminalis, *Hook. in Mitch. Journ. Trop. Austral.* 369. —Austral.
vitiensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 274.— Ins. Fiji.

VEPRECELLA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 312. t. 15. *MELASTOMACEAE*, Benth. & Hook. f. i. 752.
hispida, *Baker, in Journ. Linn. Soc.* xxi. (1884) 342. —Madag.
laxiflora, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 314.—Madag.
lutea, *Naud. l. c.* 313.—Madag.
macrophylla, *Naud. l. c.* 314.—Madag.
nigrescens, *Naud. l. c.* 313.—Madag.
schizocarpa, *Baker, in Journ. Bot.* xx. (1882) 112.— Madag.
vestita, *Baker, in Journ. Linn. Soc.* xx. (1883) 146.— Madag.

VEPRIS, Comm. ex A. Juss. in Mém. Mus. Par. xii. (1825) 509 = **Toddalia**, Juss. (Rutac.).
inermis, Comm. l. c. 538 = T. paniculata.
lanceolata, G. Don, Gen. Syst. i. 806 = T. lanceolata.
obovata, G. Don, l. c. = T. paniculata.
querimbensis, Klotzsch, in Peters, Reise Mossamb. Bot. 87 = T. lanceolata.
simplicifolia, Endl. Prod. Fl. Norfolk. 89 = Acronychia Endlicheri.

VERATAXUS, [Nelson,] Senilis, Pinaceae, 168 (1866) = **Taxus**, Tourn (Conif.).
adpressa, Carr. Conif. ed. II. 731 = T. baccata.

VERATRONIA, Miq. Fl. Ind. Bat. iii. 553 (1859) = **Susum**, Blume (Flagell.).
malayana, Miq. l. c. = S. anthelminticum.

VERATRUM, [Tourn.] Linn. Syst. ed. I (1737).
LILIACEAE, Benth. & Hook. f. iii. 834.
ACEDILANTHUS, Benth. & Hook. f. Gen. iii. 834 (1883), sphalm.
ACELIDANTHUS, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 94) (1847).
ANEPSA, Rafin. Fl. Tellur. ii. 31 (1836).
album, *Linn. Sp. Pl.* 1044.—Europ.; As. bor.
album, Michx. Fl. Bor. Am. ii. 249 = viride.
album, S. Wats. Bot. King's Exped. 344 = californicum.
americanum, Mill. Gard. Dict. ed. VIII. n. 4 = Helonias bullata.
angustifolium, Pursh, Fl. Am. Sept. i. 242 = Stenanthium angustifolium.
californicum, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 103.—Am. bor. occ.
caricifolium, Schlecht. Ind. Sem. Hort. Hal. (1838) 8 = Schoenocaulon officinale.
Dubouzeti, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 17 = Anthericum Rossii.
Eschscholtzii, A. Gray, in Ann. Lyc. N. York, iv. (1837) 119 = album.
fimbriatum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 391.—Calif.
flavum, Herb. ex Schult. f. Syst. vii. 1565 = Chamaelirium carolinianum.
frigidum, Schlecht. & Cham. in Linnaea, vi. (1831) 46 = Stenanthium frigidum.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 44. t. 96 = Smilacina purpurea.
intermedium, *Chapm. Fl. S. U. St.* 489.—Florida.

VERATRUM:—
Lobelianum, Bernh. in Schrad. Neues Journ. ii. II. (1807) 356 = album.
luteum, Linn. Sp. Pl. 1044 = Chamaelirium carolinianum.
Maackii, *Regel, in Mém. Acad. Pétersb.* Sér. VII. iv. (1861) IV. 154.—Sibir. or.
malayanum, Jack, in Malay. Misc. i. (1820) v. n. II. 25 = Susum anthelminticum.
Maximowiczii, *Baker, in Journ. Linn. Soc.* xvii. (1879) 472.—Japon.
nigrum, *Linn. Sp. Pl.* 1044.—Europ.; As. bor.
officinale, Schlecht. & Cham. in Linnaea, vi. (1831) 45 = Schoenocaulon officinale.
oxysepalum, Turcz. in Bull. Soc. Nat. Mosc. (1840) 79 = album.
parviflorum, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 166 = album.
parviflorum, *Michx. Fl. Bor. Am.* ii. 250.—Am. bor.
polygamum, Gilib. Exercit. ii. 463 = album.
purpureum, Salisb. Prod. 214 = nigrum.
Sabadilla, *Retz. Obs.* i. 31 (err. typ. 29) (sp. dub.).— Mexic.
stamineum, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 230.—Japon.
tenuifolium, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 380 = Schoenocaulon officinale.
virescens, Mart. & Gal. l. c. 381 = Zygadenus mexicanus.
virginicum, Ait. Hort. Kew. ed. II. v. 426 = Melanthum virginicum.
viride, *Ait. l. c.* ed. I. iii. 422.—Am. bor.
viride, Roehl. Deutschl. Fl. ed. II. ii. 237 = album.
Woodii, *Robbins, ex A. Gray, Man. Bot. N. U. St.* ed. II. 477.—Am. bor.

VERBASCUM, Tourn. ex Linn. Syst. ed. I (1737).
SCROPHULARINEAE, Benth. & Hook. f. ii. 928.
BLATTARIA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 247 (1745).
FLOMOSIA, Rafin. Fl. Tellur. iv. 89 (1836).
LASIAKE, Rafin. l. c. (1836).
LEIOSANDRA, Rafin. l. c. (1836).
LYCHNITIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125.
THAPSUS, Rafin. Fl. Tellur. iv. 89 (1836).
abietinum, Borb. in Verh. Bot. Ver. Brandenb. xvii. (1875) 60 = nigrum.
acuminatum, Schrad. ex Steud. Nom. ed. I. 871 (Quid ?).—Hab. ?
Adeliae, *Heldr. in Boiss. Diagn.* Ser. II. iii. 145.— Ins. Cyclad.
adenocaulon, *Boiss. & Bal. l. c.* vi. 128.—Cappadoc.
adenophorum, *Boiss. & Bal. l. c.* Ser. I. iv. 64.—Caria.
adenophorum, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 442 = glandulosum.
adulterinum ×, Koch, Syn. Fl. Germ. ed. I. 512.— Europ.
aethiopicum, *Ehrenb. ex Sweet, Hort. Brit.* ed. II. 381. —Mons Sinai.
alatum, Lam. Fl. Fr. ii. 259 = V. Thapsus.
album, Mill. Gard. Dict. ed. VIII. n. 3 = V. Lychnitis.
alceoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 306.—Kurdistan.
alepense, *Benth. in DC. Prod.* x. 241.—Syria.
alopecuroides, *Bivona, ex Guss. Fl. Sic. Syn.* i. 265.— Sicilia.
Alopecuros, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 440 = eriophorum.
alopecurus, Thuill. Fl. Par. ed. II. 110 = nigrum.
alpestre, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 317 = eriorrhabdon.
alpestre, *Hegetschw. Fl. Schw.* 13.—Europ.
alpigenum, C. Koch, in Linnaea, xxii. (1849) 724 = pyramidatum.
alpinum, Salisb. Prod. 105 = Ramondia pyrenaica.
alpinum, Turra, Fl. Ital. Prod. 64 = nigrum.
alyssifolium, *Boiss. Diagn.* Ser. I. iv. 58.—Armenia.
Amanum, *Boiss. Fl. Orient.* iv. 310.—Syria.
ambiguum, Lej. Rev. Fl. Spa, 229 = adulterinum.
amoenum, Salisb. Prod. 105 = Celsia Arcturus.
angustifolium, *Tenore, Fl. Nap.* iii. 218. t. 122.—Ital.
angustius, Schrank, in Hoppe, Taschenb. (1809) 90; ex Schrad. Monog. Verb. i. 18; ii. 57 = V. Thapsus.
annuum, Mill. Gard. Dict. ed. VIII. n. 11 = virgatum.
Antinori, *Boiss. & Heldr. Diagn.* Ser. II. iii. 144.— As. Min.

VERBASCUM:—
antiochium, *Boiss. & Heldr. Diagn.* Ser. I. xii. 9.—Syria.
Arcturus, Linn. Sp. Pl. 178 = Celsia Arcturus.
argentatum, *Delile, ex Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 441.—Hab.?
argenteum, *Tenore, Fl. Nap.* iii. 220. t. 215. f. 1.—Ital.
argyrostachyon, Tenore, Relaz. Viagg. Abruzz. 52; Syll. Fl. Neap. 107 = phlomoides.
armeniacum, Boiss. Fl. Orient. iv. 361 = Celsia aurea.
armenum, *Boiss. & Kotschy, ex Boiss. l. c.* 304.—Armenia.
asperulum, Boiss. Diagn. Ser. I. iv. 57 = cheiranthifolium.
astrophyllum ×, Boiss. & Heldr. Diagn. Ser. I. xii. 13.—As. Min.
atheniense, Hort. Angl. ex Steud. Nom. ed. II. ii. 747 (Quid?).—Graecia?
auriculatum, Sibth. & Sm. Fl. Graec. Prod. i. 149 = mucronatum.
auritum ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 129.—Gallia.
australe, Schrad. Monog. Verb. i. 28 = phlomoides.
austriacum, Schott, ex Roem. & Schult. Syst. iv. 341 = nigrum.
bactrianum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 422.—Turkestan.
banaticum, Schrad. Monog. Verb. ii. 28 = nigrum.
Barnadesii, Vahl, Symb. Bot. ii. 39 = Celsia Bernadesii.
barradense, Boiss. Fl. Orient. iv. 318 = sinaiticum.
basianicum, *Boiss. et Haussk. ex Boiss. l. c.* 341.—Mesopot.
Bastardi, Roem. & Schult. Syst. iv. 335 = thapsiforme.
beryteum, *Boiss. Diagn.* Ser. I. xii. 28.—Syria.
betonicaefolium, Desf. in Ann. Mus. Par. xi. (1808) 54 = nigrum?
bicolle, Roem. & Schult. Syst. iv. 325 = V. Thapsus.
bicolle, Schrank, Baier. Fl. i. 465 = thapsiforme?
bicolor, Badaro, Osserv. 3; in Brugnat. Giorn. Fis. Dec. II. vii. (1824) 365 = Boerhavi.
Biebersteinii, Bess. Ind. Sem. Hort. Crem. (1821); et in Enum. Pl. Volh. 53 = V. Lychnitis.
bipinnatifidum, Sims, Bot. Mag. t. 1777 = pinnatifidum.
Bischoffii ×, G. F. Koch, in Pollichia, (1849) 23.—Europ.
biscutellaefolium, *Benth. in DC. Prod.* x. 241.—Armenia.
bithynicum, *Boiss. Diagn.* Ser. I. iv. 63.—Bithynia.
Blancheanum, *Boiss. Fl. Orient.* iv. 345.—Syria.
Blattaria, *Linn. Sp. Pl.* 178.—Europ.; As. bor.
Blattaria, J. A. Schmidt, in Journ. Bot. vi. (1868) 243 = V. Thapsus.
Blattaria, Vell. Fl. Flum. 100; iii. t. 3 = virgatum.
blattariforme, Griseb. in Wiegm. Archiv, xviii. (1852) 321 = V. Blattaria.
blattarioides, Gaud. Fl. Helv. ii. 127 = Gaudini.
blattarioides, Lam. Encyc. iv. 225 = virgatum.
blattarioides, Wahlenb. ex Griseb. Spicil. Fl. Rumel. ii. 43 = xanthophoeniceum.
blattarium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = V. Blattaria.
Boerhavi, *Linn. Mant.* i. 45.—Reg. Mediterr.
Boerhavii, Frivald. ex Nym. Consp. 526 = phlomoides.
Boerhavii, Pall. Reise, iii. 540 = ovalifolium.
bohemicum, Borb. in Termész. Füzetek, ix. (1885) 276 = phlomoides.
bombyciferum, Boiss. Diagn. Ser. I. iv. 52 = V. Lagurus.
bombyciferum, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 165 = longifolium.
bombyciforme, Boiss. Fl. Orient. iv. 303 = V. Lagurus.
bracteatum, Kickx, ex Dum. Fl. Belg. Prod. 38 = V. Thapsus.
bracteatum, J. & C. Presl, Delic. Prag. 214 = V. Lychnitis.
bracteolatum, *Delile, Ind. Sem. Hort. Monsp. Suppl.* (1840) 4; *ex Godr. Fl. Juven.* ed. II. 95.—Hab.?
brassicaefolium, Crantz, Inst. ii. 323 = Celsia Arcturus.
Brauneanum ×, Wirtg. Fl. Preuss. Rheinprov. 315, 318.—Europ.
brevidens, Benth. in DC. Prod. x. 235 = lasianthum.
Brockmuelleri ×, Ruhm. in Eichl. Jahrb. Berl. (1881) 248.—Europ.
bugulaefolium, Lam. Encyc. iv. 226 = Celsia bugulifolia.

VERBASCUM:—
bulbosum, Ehrenb. ex Boiss. Fl. Orient. iv. 345 = beryteum.
caesareum, *Boiss. Diagn.* Ser. I. xii. 7.—Palaest.
calabricum, Thom. & Rich. ex Bertol. Fl. Ital. ii. 601 = angustifolium.
calaurium, Heldr. ex Boiss. Fl. Orient. iv. 313 = plicatum.
calvescens, Schur, Enum. Pl. Transs. 481 = phlomoides.
calvum, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 338.—Kurdist.
calycinum, *Ball, in Journ. Bot.* xiii. (1875) 172.—Marocc.
calyculatum ×, Chaub. in St. Amans, Fl. Agen. 87.—Europ.
campestre, *Boiss. & Heldr. Diagn.* Ser. I. xii. 5.—Anatolia.
candelabrum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 717.—Sibir. altaic.
candidissimum, DC. Fl. Fr. Suppl. 413 = mucronatum.
canescens, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 470 = V. Thapsus.
Capusi, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 222.—Turkestan.
caucasicum, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 25 = rubiginosum.
caucasicum, Hort. ex Schrad. Monog. Verb. ii. 48 = pyramidatum.
caucasicum, Schult. Obs. Bot. 42 = ferrugineum.
Ceccarinianum, Boiss. & Heldr. ex Boiss. Fl. Orient. iv. 322 = sinuatum.
Cedreti, *Boiss. Diagn.* Ser. I. xii. 19.—Syria.
Celsiae, Boiss. Voy. Bot. Espagne, ii. 444 = virgatum.
celsioides, *Benth. in DC. Prod.* x. 229.—Reg. Himal.
ceratophyllum, Schrad. Monog. Verb. ii. 7 = pinnatifidum.
cerinum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 21.—Lycaonia.
cestroides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 331.—Mesopot.
Chaixi, Vill. Prosp. 22; Hist. Pl. Dauph. ii. 491 = nigrum.
cheiranthifolium, *Boiss. Diagn.* Ser. I. iv. 56.—As. Min.
chium, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 309 = glomeratum.
chryserium, *Schrad. Monog. Verb.* ii. 9.—Palaestin.
chrysochaete, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 89.—Lycia.
chrysorrhacos, *Boiss. Diagn.* Ser. I. iv. 58.—Caria.
chrysostallostemum, Steud. Nom. ed. II. ii. 748, sphalm. = *crystallostemum*.
cilicicum, *Boiss. Diagn.* Ser. II. iii. 149.—Cilicia.
cisalpinum, Biroli, in Brugnat. Giorn. Fis. Dec. II. iv. (1821) 284 = phoeniceum.
claudiopolitanum ×, Simk. in Magyar Novényt. Lapok, ii. (1878) 150.—Europ.
Claytoni, Michx. Fl. Bor. Am. i. 148 = V. Blattaria.
coeruleum, Schanz, Cat. Sem. Hort. Vindob. (1822); ex Schrad. Monog. Verb. ii. 13 = speciosum.
collinum, Salisb. Prod. 104 = phoeniceum.
collinum, Schrad. Monog. Verb. i. 35 = V. Thapsus.
commutatum ×, Kern. ex Nym. Consp. 531.—Europ.
commutatum, *Tenore, Relaz. Viagg. Abruzz.* 121, nomen; *Syll.* 108.—Europ. austr.
comosum ×, Simk. in Termész. Füzetek, ii. (1878) 36.—Europ.
compactum, Bieb. Fl. Taur. Cauc. i. 159 = ovalifolium.
compactum, Hort. ex Reichb. Fl. Germ. Excurs. 384 = montanum.
condensatum, Schrad. Monog. Verb. i. 31. t. 3 = phlomoides.
conocarpum, *Moris, Elench. Fl. Sard.* ii. 7.—Ins. Sardoa.
cordatum, *Desf. Fl. Atlant.* i. 186.—Alger.
cordifolium, Stokes, Bot. Mat. Med. i. 381 = nigrum.
Cossonianum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 583.—Marocc.
cotoneum, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.). i. (1853). 440 = glomeratum.
crassifolium, DC. Fl. Fr. iii. 601 = montanum.
crassifolium, *Hoffmgg. & Link, Fl. Port.* i. 213. t. 26.—Lusitan.
crenatifolium, *Boiss. Diagn.* Ser. I. iv. 51.—Bulgar.
crispum, Ehrenb. ex Boiss. Fl. Orient. iv. 341 = Schimperianum.
crystallostemon, [Weinm.]. Cat. Sem. Hort. Dorpat. (1818); ex Schrad. Monog. Verb. ii. 45 = virgatum.
cupreum, Sims. Bot. Mag. t. 1226 = phoeniceum.

VERBASCUM:—
cuspidatum, Schrad. Monog. Verb. i. 23 = V. Thapsus.
cylindrocarpum, *Griseb. Spicil. Fl. Rumel.* ii. 46.—Macedon.
daenense, *Boiss. Diagn.* Ser. I. vii. 38.—Persia.
damascenum, *Boiss. l. c.* xii. 19.—Syria.
danubiale ×, Simk. in Termész. Füzetek, ii. (1878) 36.—Europ.
decalvans ×, Borb. l. c. ix. (1885) 278.—Europ.
decurrens, Stokes, Bot. Mat. Med. i. 378 = V. Thapsus.
Delileanum, *Franch. Etud. Verb.* 68.—Europ.
delphicum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 146.—Graecia.
densiflorum, Bertol. Pl. Rar. Ital. Dec. iii. 52 = thapsiforme.
dentatum, Lapeyr. Hist. Abr. Pl. Pyr. i. 114 = nigrum.
dentifolium, *Delile, Ind. Sem. Hort. Monsp.* (1836) 28.—*Cf. Ann. Sc. Nat.* Sér. II. vii. (1837) 287.—Alger.
denudatum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 23.—Lycia.
denudatum ×, Pfund, Monog. Verb. Prod. 51; et Berch. & Opiz, Oek. Fl. Boehm. iii. 369, 397.—Europ.
detersile, *Boiss. & Heldr. Diagn.* Ser. I. xii. 15.—Lycia.
digitalifolium, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 342.—Mesopot.
dimorphum ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 140.—Europ.
diphyon ×, Franch. l. c. 193.—Europ.
divaricatum ×, Kitt. Taschenb. ed. II. 352.—Europ.
diversifolium, *Hochst. in Lorent, Wanderungen*, 335.—Mesopot.
dubium, *Roem. & Schult. Syst.* iv. 330.—Lusitania.
elegans, *Boiss. & Heldr. Diagn.* Ser. I. xii. 12.—As. Min.
elegans, Salisb. Prod. 105 = Celsia orientalis.
elongatum, Moench, Meth. 446 = Boerhavi.
elongatum, Willd. Enum. Hort. Berol. 223 = V. Thapsus.
epixanthinum, *Boiss. & Heldr. Diagn.* Ser. I. vii. 39.—Graecia.
erianthum, *Benth. in DC. Prod.* x. 235.—Afghan.
eriophorum, *Godr. in Mém. Acad. Montp.* (*Sect. Médic*). i. (1853) 440.—Hab.?
eriorrhabdon, *Boiss. in Tchihat. Asie Min. Bot.* ii. 4.—As. Min.
eriostachyum, Buek, Ind. ad DC. Prod. iii. 492, sphalm. = glabratum.
erythrocaule, Schur, Enum. Pl. Transs. 483 = glabratum.
euphraticum, *Benth. in DC. Prod.* x. 241.—Cappadoc.
Euryale ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 147.—Europ.
farinosum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 542 = pulverulentum.
fasciculatum, Ehrenb. ex Sweet, Hort. Brit. ed. II. 381 = sinaiticum.
ferrugineum, Andr. Bot. Rep. t. 162 = phoeniceum.
ferrugineum, *Mill. Gard. Dict.* ed. VIII. n. 10.—Europ. austr.
firmum, Dulac, Fl. Hautes-Pyr. 384 = V. Lychnitis.
flagriforme ×, Pfund. Monog. Verb. Prod. 49; Opiz, ex Bercht. & Opiz, Oek. Fl. Boehm. iii. 370, 399.—Europ.
floccosum, Waldst. & Kit. Pl. Rar. Hung. i. 81 = pulverulentum.
foetidum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 141 (faetidum).—Graecia.
foliosum ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 142.—Europ.
formosum, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 25, nomen; ex Schrad. Monog. Verb. ii. 37 = ovalifolium.
formosum, Fisch. ex Schrank, Pl. Rar. Hort. Monac. t. 22 = spectabile.
Freynianum ×, Borb. in Oestr. Bot. Zeitschr. xxvi. (1876) 88.—Europ.
Gaillardoti, Boiss. Diagn. Ser. II. vi. 128 = sinuatum.
galilaeum, *Boiss. Diagn.* Ser. I. xii. 8.—Syria; Palaestin.
gallicum, Willd. Sp. Pl. i. 1005 = nigrum.
garganicum, *Tenore, Relaz. Viagg. Abruzz.* 53.—Italia.
Gaudini ×, Doell, ex Nym. Consp. 530.—Europ.

VERBASCUM:—
geminatum ×, Freyn, in Oestr. Bot. Zeitschr. xxvi. (1876) 370.—Europ.
geminiflorum, *Hochst. in Lorent, Wanderungen*, 336. —Mesopot.
georgicum, *Benth. in DC. Prod.* x. 228.—As. Min.; Persia.
Germaniciae, *Haussk. ex Boiss. Fl. Orient.* iv. 334.— Cataonia.
germanum ×, Franch. Fl. Loir-et-Cher, 409.—Europ.
giganteum, Willk. in Linnaea, xxv. (1852) 51 = V. Thapsus.
glabratum, *Frivald. in Flora*, xix. (1836) 440.—Banat.
glabrum, *Delarbre, in Journ. Santé Bord.* ii. 166; *ex Schrad. Monog. Verb.* ii. 55.—Gallia.
glabrum, Mill. Gard. Dict. ed. VIII. n. 8 = V. Blattaria.
glabrum, Willd. Enum. Hort. Berol. 225 = virgatum.
glanduligerum, *Velen. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 424.—Europ.
glandulosum, *Delile, Ind. Sem. Hort. Monsp.* (1849). —*Cf. Ann. Sc. Nat.* Sér. III. xii. (1849) 367.— Europ.
glandulosum, Thore, ex Roem. & Schult. Syst. iv. 348 = virgatum.
globiflorum, *Boiss. & Noé, Diagn.* Ser. II. iii. 148.— Galatia.
glomeratum, *Boiss. l. c.* Ser. I. iv. 52.—As. Min.
gnaphalodes, *Bieb. Fl. Taur. Cauc.* iii. 152.—Tauria; Reg. Cauc.
Godronii ×, Bor. Fl. Centr. Fr. ed. III. ii. 472.— Gallia.
gossypinum, *Bieb. Fl. Taur. Cauc.* iii. 152.—Iberia.
graciliflorum, *Delile, ex Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 443.—Hab.?
graecum, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. iii. 148.—Graecia.
graecum, Pall. ex Bieb. Fl. Taur. Cauc. i. 161 = sinuatum.
granatense, *Boiss. Voy. Bot. Espagne*, ii. 441.— Hispan.
grandiflorum, *Gueldenst. Reisen*, ii. 65; *ex Ledeb. Fl. Ross.* iii. 203.—Ross. austr.
grandiflorum, Mill. Gard. Dict. ed. VIII. n. 5 = phlomoides.
grandiflorum, Poir. Encyc. Suppl. iii. 715 = montanum.
grandiflorum, Schrad. Hort. Gotting. 19. t. 13 = virgatum.
grossum, Schleich. Cat. Pl. Helv. ed. IV. 36 (Quid?). —Europ.
Guicciardii, *Heldr. ex Boiss. Diagn.* Ser. II. vi. 127.— Graecia.
Gussonii, Tineo, in Guss. Fl. Sic. Syn. ii. 797 = sinuatum.
haemorrhoidale, Ait. Hort. Kew. ed. I. i. 236 = pulverulentum.
Haenseleri, *Boiss. Voy. Bot. Espagne*, ii. 442.—Reg. Mediterr. occ.
Hausmanni ×, Čelak, in Arch. Naturw. Landesd. Boehm. ii. (1873) 317.—Europ.
Haynaldianum ×, Borb. in Oestr. Bot. Zeitschr. xxv. (1875) 213.—Europ.
Heldreichii, *Boiss. Diagn.* Ser. II. iii. 147.—Graecia.
heteromallum, Panč. ex Boiss. Fl. Orient. iv. 305 = malacotrichum.
Henriquezii, *Lange, ex Henriq. in Soc. Georgr. Lisboa, Sec. Bot.* (1183) 80.—Lusitan.
heterophlomos ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 136.—Europ.
heterophyllum, Griseb. Spicil. Fl. Rumel. ii. 49 = garganicum.
heterophyllum, J. Mill. Ic. t. 67 (Quid?).—Hab.?
heterophyllum, Moretti, in Brugnat. Giorn. Fis. Dec. II. v. (1822) 111 = pulverulentum.
Heuffelii, Neilr. Aufz. Ung. Slav. Gefaesspfl. Nachtr. 53 = longifolium.
Hinkei, Frivald. in Flora, xix. (1836) 440 = nigrum.
Hohenackeri, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 42.—Reg. Cauc.
Hookerianum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 584.—Marocc.
Hornemanni, Wierzb. in Reichb. Ic. Fl. Germ. xx. 17 = glabratum.
Hornemannii, *Bess. ex Steud. Nom.* ed. I. 872; ed. II. ii. 748.—Europ.
humile, Bernh. ex Steud. l. c. 872 = nigrum.

VERBASCUM:—
humile, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 241.—Thracia.
Humnickii ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 110.—Europ.
hybridum ×, Brot. Fl. Lusit. i. 170.—Lusitan.
hybridum, Lej. Fl. Spa, ii. 299 = nigrum.
hyoseridifolium, Boiss. Fl. Orient. iv. 325 = seq.
hyoserifolium, Schrad. Monog. Verb. ii. 31. t. 3. f. 1 = speciosum.
hypoleucum, Boiss. & Heldr. Diagn. Ser. II. iii. 147 = nigrum.
incanum, Gaud. Fl. Helv. ii. 121 = Schottianum.
indicum, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 236 = V. Thapsus.
infidelium, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 342.—Cataonia.
insigne, Boiss. & Bal. Diagn. Ser. II. vi. 129 = Wiedemannianum.
insulare, Boiss. & Heldr. l. c. Ser. I. xii. 14 = eriorrhabdon.
integrifolium, Rochel, Pl. Banat. Rar. 57, in textu = speciosum.
interjectum, *Pfund, Monog. Verb. Prod.* 49.—Europ.
intermedium, Léman, ex Roem. & Schult. Syst. iv. 326 = V. Thapsus.
intermedium, Rupr. ex Pfund, Monog. Verb. Prod. 54 = nigrum?
isauricum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 22.— Phrygia.
italicum, Moric. Fl. Venet. i. 116 = phlomoides.
Jernacha, Hochst. ex Benth. in DC. Prod. x. 227 = phlomoides.
Juratzkae ×, Dichtl, in Deutsche Bot. Monatsschr. ii. (1884) 134.—Europ.
Kanitzianum, *Simk. & Walz. in Magyar Növényt. Lapok*, ii. (1878) 148, 149.—Europ.
khorassanicum, *Boiss. Fl. Orient.* iv. 319.—Persia.
Kickxianum, Dum. Fl. Belg. Prod. 38 = V. Thapsus.
Killasii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 101.—Europ.
Klotzschianum ×, Wirtg. Fl. Preuss. Rheinprov. 321. —Europ.
kochiaeforme, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 335.—Persia.
Kochianum ×, Wirtg. Fl. Preuss. Rheinprov. 320.— Europ.
Koechelii, Fenzl, ex Boiss. Fl. Orient. iv. 344 = lyratifolium.
Kotschyi, Boiss. & Hohen. Diagn. Ser. I. iv. 56 = syriacum.
laetum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 338. —Mesopot.
laevigatum, Enum. Sem. Hort. Prag. (1822); ex Schrad. Monog. Verb. ii. 45 = virgatum.
Lagurus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 42.—Byzant.
Lagurus, Noé, ex Nym. Consp. 526 = crenatifolium.
Lamottei ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 124.—Gallia.
lanatum, Gilib. Fl. Lituan. i. 40 = V. Thapsus.
lanatum, Schrad. Monog. Verb. ii. 28. t. 2. f. 1 = nigrum.
lanceolatum, Kit. in Linnaea, xxxii. (1863) 442 = virgatum.
lasianthum, *Boiss. ex Benth. in DC. Prod.* x. 234.— As. Min.
laxiflorum, Presl, Delic. Prag. 76 = pulverulentum.
laxiflorum, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 90.—Caria.
leianthum, *Benth. in DC. Prod.* x. 239.—Cilicia.
leiocaulon, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 28 = glabratum.
leiostachyon, Griseb. Spicil. Fl. Rumel. ii. 43 = glabratum.
Lemaitrei, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xxii. (1868) 13.—Europ.
leptocladum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 10.— Pamphylia.
leptocladum, Panč. Elench. Pl. Crna-Gora, 69 = glabratum.
leptostachyon, DC. Fl. Fr. Suppl. 415 = simplex.
leucanthemum, Duf. ex Gren. & Godr. Fl. Fr. ii. 552 = V. Lychnitis.
leuconeurum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 27.— Phrygia.
leucophyllum, *Griseb. Spicil. Fl. Rumel.* ii. 46.— Macedon.

VERBASCUM:—
Linnaei, Opiz, ex Bercht. & Opiz, Oek. Fl. Boehm. iii. 366, 371; Pfund, ex Nym. Consp. 527 = V. Thapsus.
longifolium, DC. Fl. Fr. Suppl. 414 = speciosum.
longifolium, *Tenore, Prod. Fl. Nap.* p. xvi.—Europ. austr.
longiracemosum ×, Chaub. in St. Amans, Fl. Agen. 89.—Europ.
luridum, Salisb. Prod. 104 = V. Blattaria.
luteum, Mill. Gard. Dict. ed. VIII. n. 4 = Boerhavi.
lychnitideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = V. Lychnitis.
Lychnitis, *Linn. Sp. Pl.* 177.—Europ.; Reg. Cauc.
Lychnitis, Schultz, Prod. Fl. Starg. Suppl. i. 14 = V. Thapsus.
lycium, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 89. —Lycia.
lydium, *Boiss. Diagn.* Ser. I. iv. 62.—Lydia.
lyratifolium, [*Roechel*] *Koch, ex Benth. in DC. Prod.* x. 242.—Cilicia.
lyratum, Lam. Encyc. iv. 223 = Celsia cretica.
lyratum, Pourr. ex Nym. Consp. 530 = nigrum.
macilentum ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 176.—Europ.
macranthum, Hoffmgg. & Link, Fl. Port. i. 215. t. 27 = phlomoides.
macrocarpum, *Boiss. Diagn.* Ser. I. xii. 6.—Persia.
macrophyllum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 161.—Persia.
macrophyllum, C. Koch, in Linnaea, xxii. (1849) 728 = nigrum.
macrophyllum, Kotschy, ex Boiss. Fl. Orient. iv. 315 = galilaeum.
macrosepalum, Boiss. & Kotschy, ex Boiss. l. c. 317 = eriorrhabdon.
macrostachyum, Griseb. Spicil. Fl. Rumel. ii. 58, in textu = longifolium.
macrurum, Tenore, Prod. Fl. Nap. App. v. 9 = thapsiforme.
magellense, *Tenore, Fl. Nap.* iii. 218.—Ital.
majale, DC. Fl. Fr. Suppl. 415 = Boerhavi.
malacotrichum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 142.—Graecia.
mallophorum, *Boiss. & Heldr. l. c.* Ser. I. vii. 39.— Graecia.
Martini ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 172.—Europ.
medum, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 23.—Persia.
megaphlomos, Boiss. & Heldr. Diagn. Ser. II. iii. 144 = speciosum.
mesopotamicum, *Boiss. Diagn.* Ser. I. xii. 24.— Mesopot.
messanense, Tineo, in Guss. Fl. Sic. Syn. ii. 887 = thapsiforme.
micranthum, Moretti, in Brugnat, Giorn. Fis. Dec. II. v. (1822) 111 = V. Lychnitis.
microcalycinum, Fenzl, ex Boiss. Fl. Orient. iv. 338 = leianthum.
microcarpum, *Benth. in DC. Prod.* x. 230.—Mesopot.
millelacuum, Boiss. in Tchihat. Asie Min. Bot. ii. 3 = pycnostachyum.
mixtum ×, Ram. ex DC. Fl. Fr. iii. 603.—Europ.
mixtum, Thomas, Cat. Pl. Suiss. 31; et ex Gaud. Fl. Helv. ii. 121 = Schottianum.
Moenchii, Schultz, Prod. Fl. Starg. Suppl. i. 14 = V. Lychnitis.
molle, C. Koch, in Linnaea, xvii. (1843) 284 = georgicum.
monspessulanum, Pers. Syn. i. 215 = nigrum.
montanum, Griseb. Spicil. Fl. Rumel. ii. 49 = phlomoides.
montanum, *Schrad. Hort. Gotting.* 18. t. 12.—Europ.
monticolum, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 325 = cheiranthifolium.
mosellanum ×, Wirtg. Fl. Preuss. Rheinprov. 316, 319.—Europ.
mucronatum, *Lam. Encyc.* iv. 218.—As. Min.; Creta.
Muhlenbeckii, Godr. Bon Cult. Nancy (1846); ex Gren. & Godr. Fl. Fr. ii. 558 = V. Blattaria.
Myconi, Linn. Sp. Pl. 178 = Ramondia pyrenaica.
myrianthum, *Boiss. Fl. Orient.* iv. 330.—As. Min.
myriocarpum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 21.— Pamphylia.
napifolium, *Boiss. Diagn.* Ser. I. iv. 60.—Caria.
neglectum, Guss. Suppl. Fl. Sic. Prod. 59 = V. Thapsus.

VERBASCUM:—
Neilreichii ×, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 367.—Europ.
Neilreichii, Schur, Enum. Pl. Transs. 484 = speciosum.
nemorosum, Noé, ex Nym. Consp. 526 = crenatifolium.
nemorosum, Schrad. Monog. Verb. i. 32 = phlomoides.
nevadense, *Boiss. Voy. Bot. Espagne*, ii. 443.—Hispan.
nigrum, *Linn. Sp. Pl.* 178.—Europ.; As. bor.
nigrum, Pall. ex Bieb. Fl. Taur. Cauc. i. 160 = orientale.
Nisus ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 150.—Europ.
nitidum, Stokes, Bot. Mat. Med. i. 382 = V. Blattaria.
niveum, *Tenore, Prod. Fl. Nap.* p. xvi.—Italia.
nothum ×, Koch, Syn. Fl. Germ. ed. I. 512.—Europ.
Nouelianum ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 117.—Europ.
Novae-Hollandiae, Roem. Hort. Turic. ex Steud. Nom. ed. II. ii. 748 (Quid?).—Austral.?
numidicum, *Pomel, Nouv. Mat. Fl. Atl.* 95.—Alger.
obovatum, *Hochst. in Lorent, Wanderungen*, 336.—Mesopot.
officinarum, Crantz, Inst. ii. 323 = V. Thapsus.
olympicum, *Boiss. Diagn.* Ser. I. iv. 54.—Bithynia.
olympicum, *Bunyard, in Gard. Chron.* (1885) II. 19 (an praec.?).—Graecia?
orchideum, Host, Fl. Austr. i. 288 = nigrum.
oreophilum, *C. Koch, in Linnaea*, xxii. (1849) 726.—Armenia.
orgyale, *Boiss. & Heldr. Diagn.* Ser. I. xii. 15.—Lycia.
orientale, All. Fl. Pedem. i. 106 = Celsia orientalis.
orientale, *Bieb. Fl. Taur. Cauc.* i. 160.—Reg. Cauc.
orientale, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 44 = nigrum.
Osbeckii, Linn. Sp. Pl. 179 = Celsia bugulifolia.
ovalifolium, *Donn, Hort. Cantab.* ed. IV. 42, nomen; *Sims, Bot. Mag.* t. 1037.—Reg. Cauc.
ovatum, Schrad. Hort. Gotting. 21. t. 15 = nigrum.
oxycarpum, C. Koch, in Linnaea, xxii. (1849) 729 = eriorrhabdon.
pallidum, Nees, in Flora, ii. (1819) 296 = V. Thapsus.
pannosum, Vis. & Panč. in Mem. Ist. Venet. xii. (1864) 475 = longifolium.
parisiense, Thuill. Fl. Par. ed. II. 110 = nigrum.
parviflorum, *Lam. Encyc.* iv. 222.—As. Min.
parvoflore, Gilib. Fl. Lituan. i. 41 = V. Lychnitis.
peregrinum, Fisch. ex Steud. Nom. ed. I. 872; ed. II. ii. 748 (Quid?).—Hab.?
Pestalozzae, *Boiss. Diagn.* Ser. I. xii. 25.—Lycia.
petiolare, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 327.—Kurdistan.
phlomoides, Auct. ex Benth. in DC. Prod. x. 237 = V. Lychnitis.
phlomoides, *Linn. Sp. Pl.* 1194.—Europ.; As. Min.
phlomoides, Russ. ex Benth. in DC. Prod. x. 233 = glomeratum.
phlomoides, Sibth. & Sm. Fl. Graec. iii. 19. t. 224 = Boerhavi.
phlomoides, Spreng. Fl. Hal. 73 = montanum.
phlomoides, Thuill. Fl. Par. ed. II. 109 = pulverulentum.
phlomoides, Urv. in Mém. Soc. Linn. Par. i. (1822) 281 = lasianthum.
phoeniceum, *Linn. Sp. Pl.* 178.—Europ.; As. bor.
phoeniceum, Wallr. Ann. Bot. 29 = Wallrothii.
phyllostachyum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 331.—Kurdist.
pilematophorum, C. Koch, in Linnaea, xxii. (1849) 721 = pycnostachyum.
pilosum ×, Doell, in Flora, xxxii. (1849) 465.—Europ.
Pinardi, *Boiss. Diagn.* Ser. I. iv. 62.—Caria.
pinnatifidum, *Vahl, Symb. Bot.* ii. 39.—Graecia; Tauria; As. Min.
pisidicum, Boiss. & Heldr. Diagn. Ser. I. xii. 16 = cheiranthifolium.
plantagineum, Moris, Stirp. Sard. Elench. i. 33 = V. Thapsus.
plantaginifolium, Schur, Enum. Pl. Transs. 484 = phoeniceum.
plicatum, *Sibth. & Sm. Fl. Graec.* iii. 21. t. 226.—Graecia.
polystachyum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 716 = songaricum.
protractum, Fenzl, in Tchihat. Asie Min. Bot. ii. 6 = V. Lychnitis.

VERBASCUM:—
prusianum, *Boiss. Diagn.* Ser. I. vii. 37.—Banat.
Pseudo-blattaria, Schleich. Cat Pl. Helv. ed. IV. 36 = V. Blattaria.
Pseudo-lychnitis, Schur, Enum. Pl. Transs. 483 = V. Lychnitis.
Pseudo-nigrum, Bogenhard, in Flora, xxiv. (1841) 150 = nigrum.
Pseudo-phoeniceum ×, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 403.—Europ.
pterocaulon ×, Nym. Consp. 528.—Europ.
ptychophyllum, *Boiss. Diagn.* Ser. I. xii. 8.—Syria.
pulverulentum, Nicotra, Prod. Fl. Mess. 349, partim = sinuatum.
pulverulentum, Salisb. Prod. 104; Bieb. Fl. Taur. Cauc. i. 160 = V. Lychnitis.
pulverulentum, Spreng. Fl. Hal. 74 = phlomoides.
pulverulentum, *Vill. Prosp.* 22; *Hist. Pl. Dauph.* 490.—Europ. austr.
pulvinatum, Thuill. Fl. Par. ed. II. 109 = pulverulentum.
pumilum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 20.—As. Min.
punalense, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 161.—Persia.
puniceum, Schrad. Hort. Gotting. 20. t. 14 = phoeniceum.
pycnostachyum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 11.—As. Min.
pycnostachyum, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 64, nomen = Cossonianum?
pyramidale, *Host, Fl. Austr.* i. 287.—Austria.
pyramidale, March, ex Georgi, Beschr. Russ. Reich. III. iv. 782 = seq.?
pyramidatum, *Bieb. Fl. Taur. Cauc.* i. 161.—Reg. Cauc.
racemiferum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 332.—Assyria.
ramigerum, *Link, ex Schrad. Monog. Verb.* i. 37.—Germania.
ramosissimum, DC. Fl. Fr. Suppl. 416 = thapsiforme.
ramosissimum, Lloyd, ex Bor. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 12 = V. Blattaria.
ramosissimum, Poir. Encyc. Suppl. iii. 718 = V. Lychnitis.
Rascheyanum, Boiss. Diagn. Ser. I. xii. 18 = Cedreti.
Regelianum ×, Wirtg. Fl. Preuss. Rheinprov. 320.—Europ.
Reissekii ×, Kern. ex Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 195.—Europ.
repandum, Rochel, Bot. Reise Banat, 86 = V. Lychnitis.
repandum, Willd. Enum. Hort. Berol. 226 = V. Blattaria.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 (Quid?).—Java.
rhombifolium, Stokes, Bot. Mat. Med. i. 378 = V. Lychnitis.
rigidulum, *Delile, ex Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 442.—Hab.?
rigidum, Boiss. & Heldr. Diagn. Ser. II. iii. 143 = plicatum.
ringens ×, Boiss. & Heldr. l. c. I. xii. 13.—Lycaon.
Rolletianum, Hort. ex Roem. & Schult. Syst. iv. 333 = phlomoides.
rotundifolium, Tenore, Prod. Fl. Nap. p. lxvi = Boerhavi.
rubiginosum, Diard, Cat. Pl. St. Calais, 151 = V. Blattaria.
rubiginosum ×, Waldst. & Kit. Pl. Rar. Hung. ii. 216. t. 197.—Europ.; Reg. Cauc.
rubricaule, *Boiss. & Heldr. Diagn.* Ser. I. xii. 23.—Phrygia.
ruderale, Pfeiff. in Bot. Zeit. iii. (1845) 329 = nigrum?
rugulosum, Willd. Enum. Hort. Berol. 224 = phlomoides.
saccatum, C. Koch, in Linnaea, xvii. (1843) 283 = ovalifolium.
salutans, Delile, Ind. Sem. Hort. Monsp. (1847) 8 = bracteolatum.
salviaefolium, *Boiss. Diagn.* Ser. I. iv. 55.—Caria.
Samaritanii, Heldr. in Boiss. Diagn. Ser. II. vi. 127 = epixanthinum.
samniticum, Tenore, Relaz. Viagg. Abruzz. 54 = phlomoides.
Sartorii, Boiss. & Heldr. Diagn. Ser. I. vii. 38 = phlomoides.

VERBASCUM:—
saxatile, Salisb. Prod. 104 = phlomoides.
scabrum, Presl, Fl. Sic. p. xxxv = sinuatum.
scaposum, *Boiss. Diagn.* Ser. I. xii. 26.—Syria.
Schiedeanum, Koch, Syn. Fl. Germ. ed. II. 592 = nigrum.
Schimperianum, *Boiss. Diagn.* Ser. I. xii. 11.—Arab.
Schmidli ×, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 296.—Europ.
Schneideri, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 317 = eriorrhabdon.
Schottianum, Koch, Syn. Fl. Germ. ed. II. 591 = Wirtgeni.
Schottianum, Schrad. Monog. Verb. ii. 13 = speciosum.
Schraderi, G. F. W. Mey. Chlor. Hannov. 326 = V. Thapsus.
Schultzianum ×, Wirtg. Fl. Preuss. Rheinprov. 316, 319.—Europ.
sclavonicum, Steud. Nom. ed. II. ii. 749, sphalm. = *slavonicum*.
semialbum ×, Chaub. in St. Amans, Fl. Agen. 88.—Europ.
semidecurrens, Kit. ex Nym. Consp. 526 = phlomoides.
semilanatum ×, Borb. Vizsg. (1877) 68; ex Freyn, in Termész. Füzetek, iii. IV. (1879) 279.—Europ.
seminigrum, Fries, Nov. Fl. Suec. 68 = V. Thapsus.
seytunense, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 325 = cheiranthifolium.
simplex, Hoffmgg. & Link, Fl. Port. i. 216 = dubium, thapsoides.
simplex, *Labill. Pl. Syr. Dec.* iv. 10. t. 5.—Syria.
sinaiticum, *Benth. in DC. Prod.* x. 236.—Arab.
sinuatum, [Habl.] Desc. Phys. Taur. 156 = pinnatifidum.
sinuatum, Lemaître, ex Bor. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 13 = Lemaitrei.
sinuatum, *Linn. Sp. Pl.* 178.—Reg. Mediterr.; Oriens.
sinuatum, Schur, Enum. Pl. Transs. 482 = orientale.
slavonicum, Kit. in Rochel, Pl. Banat. Rar. 26 = phlomoides.
Smithii, Cf. Flora, ii. (1819) 628 = pulverulentum.
smyrnaeum, *Boiss. Diagn.* Ser. I. iv. 61.—Lydia.
songaricum, *Schrenk, Enum. Pl. Nov.* i. 26.—Soongaria.
speciosum, Opiz, Seznam, 102 = *denudatum*, Pfund.
speciosum, *Schrad. Hort. Gotting.* 22. t. 16.—Europ. austr.; As. Min.
spectabile, *Bieb. Fl. Taur. Cauc.* iii. 158.—Tauria.
spectabile, Salisb. Prod. 104 = V. Thapsus.
sphenandroides, *C. Koch, in Linnaea*, xxii. (1849) 731.—As. Min.
spinosum, *Linn. Cent. Pl.* ii. 10; *Amoen. Acad.* iv. 307.—Ins. Creta.
splendidum ×, Boiss. Diagn. Ser. I. iv. 53.—As. Min.
spurium ×, Koch, Syn. Fl. Germ. ed. I. 511.—Europ.
stachydifolium, Boiss. & Heldr. Diagn. Ser. I. xii. 25 = sphenandroides.
stachydiforme, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 159.—Reg. Cauc.
stenocarpum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 18.—Pamphylia.
stenophyllum, Boiss. & Heldr. l. c. 17 = cheiranthifolium.
Steveni, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 160.—Oriens.
Stevenii, [Trev.] Cat. Hort. Vindob. (1821); ex Schrad. Monog. Verb. 15; Spreng. ex Hoffmgg. Verz. Pfl. Nachtr. 226 = gnaphalodes.
strictum, *Clarke, Trav.* ii. 262.—Ins. Rhodus.
subalpinum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii-xxiv. (1880) 100.—Europ.
subalpinum, Schur, Enum. Pl. Transs. 481 = V. Thapsus.
sublyratum ×, Borb. in Termész. Füzetek, ix. (1885) 278.—Europ.
subnivale, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 305.—Cataonia.
sulcatum, *Desf. Cat. Hort. Par.* ed. III. 396.—Hab.?
syriacum, *Schrad. Monog. Verb.* ii. 6. t. 1. f. 1.—Syria.
Szovitsianum, *Boiss. Fl. Orient.* iv. 333.—Persia; Reg. Cauc.
tabsoides, Krock, Fl. Siles. Suppl. i. 272, sphalm. = thapsoides.
talyschense, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 160.—Sibir.
Tapsus, Neck, Delic. Gallo-Belg. i. 121 = V. Thapsus.
taraxacifolium, Boiss. Diagn. Ser. I. iv. 59 = parviflorum.

VERBASCUM:—
Tauri, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 145.—Cilicia.
tauricum, Hook. Bot. Mag. t. 3799 = rubiginosum.
tenellum, *Hegetschw. Fl. Schw.* 14.—Helvet.
Ternacha, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 108 = phlomoides.
Thapsi ×, Linn. Sp. Pl. ed. II. 1669.—Europ.
thapsiforme, Guss. Fl. Sic. Prod. i. 260 = V. Thapsus.
thapsiforme, *Schrad. Monog. Verb.* i. 21.—Europ.; As. Min.
thapsoides, Balb. ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 282 = montanum.
thapsoides, Comolli, Prod. Fl. Com. 40 = thapsiforme.
thapsoides, Host, Syn. Pl. Austr. 115 = speciosum.
thapsoides, *Linn. Syst.* ed. XII. 169.—Europ.
thapsoides, Schrank, ex Schrad. Monog. Verb. 19 = V. Thapsus.
thapsoides, Vill. Hist. Pl. Dauph. ii. 490 = phlomoides.
thapsum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = V. Thapsus.
Thapsus, Brot. Fl. Lusit. i. 270 = crassifolium.
Thapsus, *Linn. Sp. Pl.* 177.—Europ.; Oriens; Reg. Himal.
Thapsus, Plenk, Pl. Offic. ii. 10. t. 109 = phlomoides.
Thapsus, Trev. Prosp. Fl. Eug. 24 = thapsiforme.
Thomaeanum ×, Wirtg. Fl. Preuss. Rheinprov. 321.—Europ.
thyrsoideum, Host, Fl. Austr. i. 289 = nigrum.
Tiberiadis, *Boiss. Diagn.* Ser. I. xii. 26.—Syria.
tolosanum ×, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. i. (1872–73) 132.—Europ.
tomentosulum ×, Freyn, in Oestr. Bot. Zeitschr. xxvii. (1877) 397.—Europ.
tomentosum, Lam. Fl. Fr. ii. 260 = phlomoides.
Tommasinianum ×, Freyn, in Termész. Füzetek, iii. iv. (1879) 279 = *semilanatum*.
Tournefortii, Crantz, Inst. ii. 323 = Celsia bugulifolia.
tripolitanum, *Boiss. Diagn.* Ser. I. xii. 9.—Syria.
triste, Sibth. & Sm. Fl. Graec. Prod. i. 151 = phoeniceum.
turkestanicum, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 221.—Turkestan.
turriculatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 28 = Blumea appendiculata?
undulatum, Bieb. Fl. Taur. Cauc. i. 161 = sinuatum.
undulatum, Lam. Encyc. iv. 221 = plicatum, ptychophyllum.
urticaefolium, Lam. l. c. 220 = nigrum.
ustulatum, Čelak, in Arch. Naturw. Landesd. Boehm. ii. (1873) 318 = commutatum.
valentinum, *Burnat & Barbey, Voy. Bot. Baléares*, 50.—Hispan.
vernale, Wierzb. ex Rochel, Bot. Reise Banat, 86 = nigrum.
versicolor, Wender. ex Steud. Nom. ed. II. ii. 749 (Quid?).—Hab. ?
versiflorum, *Schrad. Monog. Verb.* i. 36.—Bohem.
vestitum, *Franch. ex Borb. in Termész. Füzetek*, ix. (1885) 277.—Europ.
vidavense ×, Simk. l. c. v. (1881) 49.—Europ.
villosum, Bernh. ex Steud. Nom. ed. I. 873 = nigrum.
viminale, Guss. Pl. Rar. 101. t. 21 = phlomoides.
virens, Host, Fl. Austr. i. 290 = orientale.
virens, Reichb. ex Nym. Consp. 530 = nigrum?
virgatum, Spreng. Syst. i. 621 = V. Blattaria.
virgatum, *Stokes, in With. Bot. Arr. Brit. Pl.* ed. II. i. 227.—Europ.
viscidulum, Pers. Syn. i. 215 = virgatum.
Visianianum, Reichb. f. Ic. Fl. Germ. xx. 13 = V. Thapsus.
vulcanicum, *Boiss. & Heldr. Diagn.* Ser. I. xii. 6.—Lycaonia.
Wallichianum, Benth. in DC. Prod. x. 231 (sp. dub.).—Ind. or.
Wallrothii, *Bluff & Fingerh. Comp. Fl. Germ.* ed. I. i. 288.—Europ.
Weldenii, Braun, ex Moretti, in Brugnat. Giorn. Fis. Dec. II. v. (1822) 43 = V. Lychnitis.
Wiedemannianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 51.—As. Min.
Wierzbickii, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 167 = nigrum.
Wilhelmsianum, C. Koch, in Linnaea, xxii. (1849) 724 = orientale.

VERBASCUM:—
Wirtgeni ×, Franch. in Mém. Soc. Acad. Maine-et-Loire, xxii. (1868) 160.—Europ.
xanthophoeniceum, *Griseb. Spicil. Fl. Rumel.* ii. 42.—Banat.
Zuccarinii, Nym. Consp. 531 = graecum.

VERBENA, Linn. Syst. ed. I (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1146.
BILLARDIERA, Moench, Meth. 369 (1794).
BURSERIA, Loefl. It. Hisp. 194 (1758).
GLANDULARIA, J. F. Gmel. Syst. 920 (1791).
OBLETIA, Rozier, in Journ. Phys. i. (1773) 367.
PATYA, Neck. Elem. i. 296 (1790).
SHUTTLEWORTHIA, Meissn. Gen. 290 (1840).
STYLEURODON, Rafin. Fl. Tellur. ii. 104 (1836).
STYLODON, Rafin. Neogenyt. 2 (1825).
UWAROWIA, Bunge, in Bull. Sc. Acad. Pétersb. vii. (1840) 278.
affinis, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 322 = litoralis.
alata, *Cham. in Linnaea*, vii. (1832) 258.—Bras.
alopecurioides, Hort. ex Steud. Nom. ed. II. ii. 749 = stricta.
Alopecurus, *Cav. Desc.* 68.—Am. bor.
alpigena, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 378 = Lippia scorodonioides.
alternifolia, *Hort. ex Schau. in DC. Prod.* xi. 544.—Chili.
americana, *Mill. Gard. Dict.* ed. VIII. n. 10.—Panama.
amoena ×, Paxt. Mag. Bot. vii. (1840) t. ad p. 3.
Andrieuxii, *Schau. in DC. Prod.* xi. 553.—Mexic.
angustifolia, *Michx. Fl. Bor. Am.* ii. 14.—Am. bor.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 15 = Stachytarpheta angustifolia.
aphylla, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 161. t. 46.—Chili.
araucana, *Phil. in Anal. Univ. Chil.* (1873) 520.—Chili.
aristata, Vahl, Eclog. Am. ii. 2 = Stachytarpheta orubica.
arisonica, A. Gray, in Proc. Am. Acad. xix. (1883) 95 = remota.
Arraniana, Hort. ex Steud. Nom. ed. II. ii. 749 = incisa.
asparagoides, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 165.—Chili.
aspera, *Gill. & Hook. l. c.* 163.—Chili.
aspera; Kunze, ex Schau. in DC. Prod. xi. 540 = corymbosa.
Aubletia, *Jacq. Hort. Vindob.* ii. 82.—Am. bor.
barbata, *R. Grah. in Edinb. N. Phil. Journ.* (Oct.–Dec. 1827) 176.—Mexic.
Berterii, *Schau. in DC. Prod.* xi. 551.—Chili.
bipinnatifida, *Schau. l. c.* 553.—Am. bor.
biserrata, H. B. & K. Nov. Gen. et Sp. ii. 275 = Carolina.
bonariensis, *Linn. Sp. Pl.* 20.—Reg. trop.
bracteata, *Cav. ex Lag. & Rodr. in Anal. Cienc. Nat.* iv. (1801) 260.—Am. bor.; N. Mexic.
bracteosa, Michx. Fl. Bor. Am. ii. 13 = praec.
brasiliensis, Vell. Fl. Flum. 17; i. t. 40 = litoralis.
bryoides, *Phil. Fl. Atac.* 40.—Chili.
Buchnera, Vell. Fl. Flum. 17; i. t. 42 = lobata.
caerulea, Vatke, in Ind. Sem. Hort. Berol. (1876) App. 1 = xutha.
caespitosa, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 165.—Chili.
calcicola, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 378 = clavata.
canescens, Chapm. Fl. S. U. St. 307 = bracteata.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 274. t. 136.—Mexic.
cannabifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 152.—Japon.
capensis, Thunb. Prod. Pl. Cap. 96 = bonariensis, Lippia asperifolia.
capitata, Forsk. Fl. Aegypt. Arab. 10 = Lippia nodiflora.
caracasana, H. B. & K. Nov. Gen. et Sp. ii. 275 = litoralis.
carnea, *Medic. Beobacht.* (1783) 131.—Am. bor.
Carolina, *Linn. Syst.* ed. X. 852; *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
caroliniana, Hort. ex Schau. in DC. Prod. xi. 545 = carnea.
caroliniana, *Michx. Fl. Bor. Am.* ii. 14.—Am. bor.

VERBENA:—
caroliniana, Willd. Sp. Pl. i. 119 = Carolina.
Carroo, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 112.—Patagon.
caudata, Salisb. Prod. 71 = Stachytarpheta indica.
cayennensis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = Stachytarpheta cajanensis.
chamaedryfolia, *Juss. in Ann. Mus. Par.* vii. (1806) 73.—Am. trop.
Chamissonis, Walp. Rep. iv. 18 = intermedia.
ciliata, *Benth. Pl. Hartw.* 21.—Mexic.
cinerascens, *Schau. in DC. Prod.* xi. 545.—Chili.
citriodora, Cav. Desc. 68 = Lippia citriodora.
clavata, *Ruiz & Pav. Fl. Per.* i. 21. t. 33.—Peruv.
colchaguensis, *Phil. in Anal. Univ. Chil.* (1873) 521.—Chili.
corymbosa, Cham. in Linnaea, vii. (1832) 255 = lobata.
corymbosa, *Ruiz & Pav. Fl. Per.* i. 22. t. 33.—Chili.
crithmifolia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 169.—Chili.
cuneata, Willd. ex Spreng. Syst. ii. 752 = Lippia nodiflora.
cuneifolia, Rafin. in Med. Repos. N. York, II. v. (1808) 360 = stricta.
cuneifolia, *Ruiz & Pav. Fl. Per.* i. 22.—Peruv.
cuneiformis, Spreng. Syst. ii. 750 = praec.
Cunha, Vell. Fl. Flum. 17; i. t. 41 = phlogiflora.
curassavica, Linn. Sp. Pl. 19 = Tamonea spinosa.
delicatula, *Mart. & Zucc. in Otto & Dietr. Allg. Gartenz.* ii. (1834) 245.—Mexic.
dentata, Vis. Ort. Bot. Padova, 150 = Lippia chamaedryfolia.
deserticola, *Phil. Fl. Atac.* 40.—Chili.
Diceras, Bert. ex Steud. Nom. ed. II. ii. 750 = sulphurea.
dichotoma, Ruiz & Pav. Fl. Per. i. 23 = Stachytarpheta dichotoma.
diffusa, Poir. Encyc. viii. 550 = urticifolia.
dissecta, Poepp. ex Schau. in DC. Prod. xi. 551 = sulphurea.
dissecta, *Willd. ex Spreng. Syst.* ii. 750.—Chili.
Doniana, Steud. Nom. ed. II. ii. 750 = venosa.
Drummondii, Hort. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 680 = V. Aubletia.
Echegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 66.—Chili.
echinata, Phil. Sert. Mendoc. Alt. 33 = seriphioides.
Ehrenbergiana, *Schau. in DC. Prod.* xi. 548.—Mexic.
elegans, *H. B. & K. Nov. Gen. et Sp.* ii. 273.—Mexic.
elliptica, Willd. ex Spreng. Syst. ii. 751 = Lippia canescens.
ellipticifolia, Stokes, Bot. Mat. Med. i. 38 = Stachytarpheta indica.
elongata, Salisb. Prod. 71 = bonariensis.
ephedroides, *Cham. in Linnaea*, vii. (1832) 261.—Bras.
erinacea, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 164. t. 48.—Chili.
erinoides, *Lam. Illustr.* i. 57.—Peruv.
erinoides, Poepp. ex Schau. in DC. Prod. xi. 551 = Berterii.
exilis, *Schau. l. c.* 553.—Mexic.
fasciculata, *Benth. Bot. Voy. Sulph.* 153.—Peruv.
filicaulis, *Schau. in DC. Prod.* xi. 549.—Bras.
flava, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 170.—Chili.
fluminensis, Vell. Fl. Flum. 17; i. t. 38 = Bouchea Pseudo-gervao.
foetida, *Phil. in Linnaea*, xxix. (1857–58) 19.—Chili.
Forskålaei, Vahl, Symb. Bot. iii. 6 = Priva leptostachya.
fragrans, Salisb. Prod. 71 = Lippia citriodora.
fruticosa, *Mill. Gard. Dict.* ed. VIII. n. 14.—Mexic.
geraniaefolia, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1198 = tenera.
glabrata, H. B. & K. Nov. Gen. et Sp. ii. 276 = litoralis.
glabrata, *Phil. Sert. Mendoc. Alt.* 32.—Chili.
glandulosa, Moris, in Ann. Stor. Nat. iv. (1830) 39 = hispida.
glauca, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 163.—Chili.
globiflora, L'Hérit. Stirp. Nov. 23. t. 12 = Lippia asperifolia.
globiflora, Nocca, ex Spreng. Syst. ii. 751 = Lippia reptans.
globulifera, Spreng. Syst. ii. 751, sphalm. = Lippia asperifolia.
gracilis, Cham. in Linnaea, vii. (1832) 262 = intermedia.

VERBENA:—
gracilis, Desf. Cat. Hort. Par. ed. III. 393 = canescens.
grandiflora, *Ortega, Hort. Matr. Dec.* 2.—Cuba.
grandiflora, Steud. Nom. ed. I. 873 = V. Aubletia.
gratissima, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 160 = Lippia lycioides.
gynobasis, *Wedd. Chlor. And.* ii. 156.—Peruv.
hastata, *Linn. Sp. Pl.* 20.—Am. bor.
heterophylla, Willd. ex Spreng. Syst. ii. 749 = inflata.
hirsuta, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 321 = Carolina.
hirta, *Spreng. Syst.* ii. 749.—Reg. Argent.
hispida, *Ruiz & Pav. Fl. Per.* i. 22. t. 34.—Am. austr.
humifusa, *Cham. in Linnaea*, vii. (1832) 271.—Bras.
hybrida ×, Hort. ex Vilm. Fl. Pl. Terre, Suppl. 195.
hyderabadensis, Rottl. ex Wall. Cat. n. 6318 = Bouchea hyderabadensis.
incarnata, Rafin. Atl. Journ. 153 (Quid?).—Am. bor.
incisa, *Hook. Bot. Mag.* t. 3628.—Panama.
indica, Jacq. Obs. iv. 7. t. 86 = Stachytarpheta angustifolia.
indica, Linn. Syst. ed. X. 851 = Stachytarpheta indica.
inflata, *H. B. & K. Nov. Gen. et Sp.* ii. 273.—Ecuador.
integrifolia, Michx. ex Walp. Rep. iv. 13 = angustifolia.
intermedia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 166.—Chili.
intermedia ×, Penny, ex G. Don, in Loud. Hort. Brit. Suppl. i. 602.
jamaicensis, Linn. Sp. Pl. 19 = Stachytarpheta indica.
jamaicensis, Vell. Fl. Flum. 16; i. t. 37 = Stachytarpheta dichotoma.
javanica, Burm. f. Fl. Ind. 12 = Lippia javanica.
juncea, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 162 = Baillonia juncea.
juniperina, *Lag. Gen. et Sp. Nov.* 19.—Chili.
Lamberti, Sims, Bot. Mag. t. 2200 = V. Aubletia.
lanata, Willd. ex Walp. Rep. iv. 48 = Lippia canescens.
lanceolata, Willd. ex Spreng. Syst. ii. 748 = litoralis.
lancifolia, Steud. Nom. ed. II. ii. 750 = Stachytarpheta indica.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864–65) 194.—Chili.
lantanoides, Willd. ex Spreng. Syst. ii. 763 = Lippia geminata.
lappulacea, Linn. Sp. Pl. 19 = Priva echinata.
lasiostachys, Link, Enum. Hort. Berol. ii. 122 = prostrata.
lavandulaefolia, *Phil. in Anal. Univ. Chil.* (1873) 521.—Chili.
ligustrina, Lag. Gen. et Sp. Nov. 18 = Lippia lycioides.
lilacina, *Greene, in Bull. Calif. Acad.* i. IV. (1885) 210.—Ins. Cedros.
Lindleyana, Walp. Ann. iii. 232 = Buddleia Lindleyana.
lipozygioides, *Walp. Rep.* iv. 16.—Chili.
litoralis, *H. B. & K. Nov. Gen. et Sp.* ii. 276. t. 137.—Am. trop.
lobata, *Vell. Fl. Flum.* 18; i. t. 43.—Bras.
lobelioides, R. Grah. in Edinb. N. Phil. Journ. xxix. (1840) 175 = Priva laevis.
longiflora, Lam. Illustr. i. 57 = V. Aubletia.
longifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 323.—Mexic.
Lorentzii, *Niederl. ex Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 370.—Reg. Argent.
Lucaeana, Walp. Rep. iv. 23 = xutha.
macrostachya, F. Muell. Fragm. i. 60 = officinalis.
Mahoneti, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 938 = tenera.
marrubioides, *Cham. in Linnaea*, vii. (1832) 266.—Bras.
Matthesii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 196.—Texas.
maysorensis, Wight, ex Wall. Cat. n. 6318 = Bouchea hyderabadensis.
megapotamica, Spreng. Syst. iv. Cur. Post. 230 = phlogiflora.
melindres, Gill. in Lindl. Bot. Reg. t. 1184 = chamaedryfolia.
melindroides, Cham. in Linnaea, vii. (1832) 270 = chamaedryfolia.

VERBENA:—
melissoides, Sweet, ex Steud. Nom. ed. II. ii. 750 = chamaedryfolia.
mendocina, *Phil. Sert. Mendoc. Alt.* 33.—Chili.
menthaefolia, Benth. Pl. Hartw. 21 = officinalis.
mexicana, Linn. Sp. Pl. 19 = Priva hispida.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 272. t. 133.—Am. austr.
microphylla, *Phil. in Linnaea*, xxix. (1857–58) 21.—Chili.
minima, *Meyen, Reise*, i. 451.—Peruv.
mollis, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 323 = Carolina.
mollis, Rafin. Atl. Journ. 146 (Quid?).—Am. bor.
montevidensis, Spreng. Syst. ii. 747 = stellarioides.
moranensis, Willd. ex Spreng. l. c. 750 = elegans.
multifida, Ruiz & Pav. Fl. Per. i. 21 = erinoides.
mutabilis, Jacq. Coll. ii. 334 = Stachytarpheta mutabilis.
mysorensis, Wight, ex Wall. Cat. n. 6318 A = Bouchea hyderabadensis.
Negundo, Royle, Illustr. Bot. Himal. 466 (Index), sphalm. = Vitex Negundo.
nervosa, Scheele, in Linnaea, xvii. (1843) 352 = Buchnera elongata.
Nivenii, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 938 = teucrioides.
nodiflora, Linn. Sp. Pl. 20 = Lippia nodiflora.
nodiflora, Petagn. ex Spreng. Syst. ii. 752 = Lippia repens.
nudiflora, *Nutt. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 195.—Ins. Sandvic.
Oblaetia, Retz. in Vet. Akad. Nya Handl. Stockh. (1778) 143 = V. Aubletia.
Obletia, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 194. t. 7 = praec.
odorata, Desf. ex Steud. Nom. ed. I. 898, in syn. = Lippia asperifolia.
odorata, Meyen, Reise, i. 480 = erinoides.
officinalis, *Linn. Sp. Pl.* 20.—Amphig. temp.
orchioides, Hort. ex Walp. Rep. iv. 36 = Priva laevis.
origenes, *Phil. in Linnaea*, xxix. (1857–58) 20.—Chili.
orubica, Linn. Sp. Pl. 18 = Stachytarpheta orubica.
ovata, *Cham. in Linnaea*, vii. (1832) 263.—Bras.
paniculata, Lam. Encyc. viii. 548 = hastata.
parviflora, Ruiz, ex Walp. Rep. iv. 25 = canescens.
patagonica, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 114.—Patagon.
pauciflora, Walp. Rep. vi. 687 = seq.
paucifolia, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 324 = Carolina.
paucifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 196.—Texas.
phlogiflora, *Cham. in Linnaea*, vii. (1832) 266.—Bras.
pinnatifida, Lam. Illustr. i. 57 = hastata.
pinnatisecta, *Schau. in DC. Prod.* xi. 549.—Bras.
platensis, Spreng. Syst. ii. 748 = teucrioides.
pogonostoma, Klotzsch, ex Schau. in DC. Prod. xi. 550 = seq.
pogostoma, *Klotzsch, ex Walp. Rep.* iv. 31.—Reg. Argent.
polycephala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 194.—Chili.
polycnemoides, Lag. Gen. et Sp. Nov. 18 = minima.
polystachya, H. B. & K. Nov. Gen. et Sp. ii. 274 = Carolina.
porrigens, *Phil. in Linnaea*, xxxiii. (1864–65) 195.—Chili.
prismatica, Linn. Sp. Pl. 19 = Bouchea Ehrenbergii.
procumbens, Forsk. Fl. Aegypt. Arab. 10 = supina.
prostrata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 41.—Am. bor.
Pseudo-gervao, A. St. Hil. Pl. Us. Brés. t. 40 = Bouchea Pseudo-gervao.
Pseudo-juncea, *C. Gay, Fl. Chil.* v. 19.—Chili.
pulchella, Sweet, Brit. Flow. Gard. Ser. I. iii. t. 295 = tenera.
pulcherrima, Vilm. Fl. Pl. Terre, ed. I. 937 = erinoides.
quadrangularis, Vell. Fl. Flum. 16; i. t. 39 = bonariensis.
radicans, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 170.—Chili.
radicans, Moench, Meth. 369 = supina.
recta, H. B. & K. Nov. Gen. et Sp. ii. 277 = Carolina.
remota, *Benth. Pl. Hartw.* 21 = canescens.
repens, Bertol. Rar. Ital. Pl. Dec. ii. 27 = Lippia nodiflora.

VERBENA:—
repens, Spreng. ex Steud. Nom. ed. II. ii. 750 = bracteata.
reptans, Loisel. ex Spreng. Pugill. ii. 70 = Zapania reptans.
rhinanthifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 323.—Mexic.
ribifolia, *Walp. Rep.* iv. 29.—Chili.
rigens, Michx. Fl. Bor. Am. ii. 14 = stricta.
rigida, Spreng. Syst. iv. Cur. Post. 230 = venosa.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 755 = canescens.
rubra, Salisb. Prod. 71 = V. Aubletia.
rugosa, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. iv. t. 318 = venosa.
rugosa, *Mill. Gard. Dict.* ed. VIII. n. 18.—Mexic.
rugosa, Muhl. ex Willd. Enum. Hort. Berol. 633 = angustifolia.
Sabini, Sweet, Brit. Flow. Gard. Ser. II. iv. sub t. 347 = erinoides.
sagittalis, *Cham. in Linnaea*, vii. (1832) 259.—Bras.
salviaefolia, Hook. & Arn. Bot. Beech. Voy. 42 = Lippia chilensis.
sarmentosa, Willd. Enum. Hort. Berol. 632 = Lippia nodiflora.
scaberrima, Cham. in Linnaea, vii. (1832) 267 = venosa.
scabra, Muhl. ex Spreng. Syst. ii. 752 = Lippia lanceolata.
scabra, Vahl, Eclog. Am. ii. 2 = urticifolia.
scoparia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 161. t. 47.—Chili.
scoparia, Tausch, in Flora, xix. (1836) 390 = stricta?
scordioides, Cham. in Linnaea, vii. (1832) 269 = teucrioides.
scrobiculata, *Griseb. in Goett. Abh.* xxiv. (1879) 275.—Reg. Argent.
selaginoides, Kunth, ex Walp. Rep. iv. 15 = alternifolia.
Selloi, Spreng. Syst. ii. 750 = erinoides.
seriphioides, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 164.—Chili.
setosa, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 321 = officinalis.
simplex, Lehm. Ind. Sem. Hort. Hamb. (1825) 17.—Cf. Linnaea, iii. (1828) Litt. 10 = angustifolia.
sororia, D. Don, Prod. Fl. Nep. 104 = officinalis.
spartioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 195.—Chili.
spathulata, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 162.—Chili.
spicata, Gilib. Fl. Lituan. i. 92 = officinalis.
spuria, Linn. Sp. Pl. 20 = officinalis.
squamosa, Jacq. Hort. Schoenb. i. 3. t. 5 = Elytraria tridentata.
squarrosa, Roth, Catalect. fasc. iii. 3 = bracteata.
stellarioides, *Cham. in Linnaea*, vii. (1832) 264.—Bras.
stenophylla, Scheele, in Linnaea, xvii. (1843) 352 = Buchnera rosea.
stoechadifolia, Linn. Sp. Pl. 19 = Lippia stoechadifolia.
stricta, *Vent. Jard. Cels*, 53. t. 53.—Am. bor.
strigosa, *Cham. in Linnaea*, vii. (1832) 256.—Bras.
strigosa, Hook. Comp. Bot. Mag. i. (1835) 176 = xutha.
Struthionum, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 116.—Reg. Argent.
subfruticosa, Aubl. Pl. Gui. i. 16 = Lippia stoechadifolia.
suffruticosa, Steud. Nom. ed. II. ii. 750 = praec.
sulphurea, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. iii. t. 221.—Am. austr.
supina, *Linn. Sp. Pl.* 21.—Geront. trop.
tenera, *Spreng. Syst.* ii. 750.—Reg. Argent.
tenuis, Steud. Nom. ed. II. ii. 751 = intermedia.
tenuispicata, *Stapf, in Denkschr. Akad. Wien*, l. (1885) 34.—Persia.
teucriifolia, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 322.—Mexic.
teucrioides, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 167.—Chili.
thymifolia, *Lag. Gen. et Sp. Nov.* 19.—Paraguay.
thymoides, *Cham. in Linnaea*, vii. (1832) 257.—Bras.
trichotoma, Moench, Meth. Suppl. 131 = bonariensis.

VERBENA:—
tridactylites, *Lag. Gen. et Sp. Nov.* 19.—Ins. Falkland.
tridens, *Lag. l. c.*—Paraguay.
trifida, *H. B. & K. Nov. Gen. et Sp.* ii. 273. t. 134.—Mexic.
trifurcata, *Phil. in Linnaea*, xxix. (1857–58) 21.—Chili.
triphylla, L'Hérit. Stirp. Nov. 21 = Lippia citriodora.
tuberosa, R. Grah. in Edinb. N. Phil. Journ. xxix. (1840) 174 = Priva laevis.
Tweedieana, Niven, ex Hook. Bot. Mag. t. 3541 = phlogiflora.
uniflora, *Phil. in Linnaea*, xxix. (1857–58) 20.—Chili.
urticifolia, *Linn. Sp. Pl.* 20.—Am. temp. et trop.
valerianoides, *H. B. & K. Nov. Gen. et Sp.* ii. 277.—N. Granat.
venosa, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 167.—Reg. Argent.
veronicaefolia, H. B. & K. Nov. Gen. et Sp. ii. 275 = Carolina.
veronicaefolia, Sm. in Rees, Cycl. xxxvi. n. 28 = chamaedryfolia.
virgata, Ruiz & Pav. Fl. Per. i. 20 = Lippia urticoides.
Wrightii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 337.—Mexic.
xutha, *Lehm. Del. Sem. Hort. Hamb.* (1834).—*Cf. Linnaea*, x. (1835–36) *Litt.* 115.—Am. bor.

VERBESINA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 379.
Abesina, Neck. Elem. i. 33 (1790).
Ancistrophora, A. Gray, in Mem. Am. Acad. N. S. vi. (1857) 457.
Ditrichum, Cass. in Bull. Soc. Philom. (1817) 33.
Hamulium, Cass. l. c. (1820) 173; et in Dict. Sc. Nat. xx. 260 (1821).
Hingstonia, Rafin. in Med. Repos. N. York, II. v. (1808) 352.
Locheria, Neck. Elem. i. 41 (1790).
Ochronelis, Rafin. Atl. Journ. 153 (1832).
Phaethusa, Gaertn. Fruct. ii. 425. t. 169 (1791).
Platypteris, H. B. & K. Nov. Gen. et Sp. iv. 200 (1820).
Saubinetia, Remy, in C. Gay, Fl. Chil. iv. 282. t. 49 (1849).
Tepion, Adans. Fam. ii. 131 (1763).
Ximenesia, Cav. Ic. ii. 60. t. 178 (1793).
abscondita, *Klatt, in Leopoldina*, xx. (1884) 93.—Mexic.
abyssinica, A. Rich. Tent. Fl. Abyss. i. 409 = Coreopsis abyssinica.
Acmella, Linn. Sp. Pl. 901 = Spilanthes Acmella.
acuminata, *DC. Prod.* v. 614.—Venezuela.
aestuans, *Lam. Encyc.* iii. 262.—Am. trop.
alata, *Linn. Sp. Pl.* 901.—Am. trop.
alata, Neuenh. in Roem. Arch. i. ii. (1797) 31 = Rudbeckia alata.
alba, Linn. Sp. Pl. 902 = Eclipta erecta.
Ancistrophora, A. Gray, in Proc. Am. Acad. xix. (1883) 14 = Wrightii, *Griseb.*
aquatica, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 28 = Sesuvium Portulacastrum.
arborea, *H. B. & K. Nov. Gen. et Sp.* iv. 202.—Ecuador.
argentea, *Bertol. Fl. Guatimal.* 35.—Guatemala.
argentea, *Gaud. in Freyc. Voy. Bot.* 463.—Ins. Marian.
Arnottii, *Baker, in Mart. Fl. Bras.* vi. iii. 215.—Paraguay.
aspilioides, *Griseb. in Goett. Abh.* xxiv. (1879) 194.—Reg. Argent.
asteroides, Linn. Sp. Pl. 902 = Amellus Lychnitis.
atriplicifolia, Poir. Encyc. viii. 460 = Montañoa arborescens.
aurea, DC. Prod. v. 613 = Zexmenia aurea.
auriculata, *DC. l. c.* 617.—Mexic.
auriculata, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 315 = subcordata.
australis, *Baker, in Mart. Fl. Bras.* vi. iii. 215.—Bras.
biflora, Linn. Sp. Pl. ed. II. 1272 = Wedelia biflora.
biflora, Wall. Cat. n. 3204 E = Wedelia Wallichii.
bipinnatifida, *Baker, in Mart. Fl. Bras.* vi. iii. 213.—Bras.
Bosvallea, Linn. f. Suppl. 379 = Glossocardia linearifolia.
Boswellia, Roxb. Hort. Beng. [101]; Fl. Ind. iii. 443 = praec.

VERBESINA:—
brasiliana, Pers. Syn. ii. 472 = Adenostemma brasilianum.
buphthalmoides, Link & Otto, Ic. Pl. Sel. 105. t. 49 = Spilanthes arnicoides.
calendulacea, Linn. Sp. Pl. 902 = Wedelia calendulacea.
canescens, Gaudich. in Freyc. Voy. Bot. 463 = Wedelia Chamissonis.
Capitaneja, *Nees, in Linnaea*, xix. (1847) 729.—Mexic.
ceanothifolia, Hook. & Arn. Bot. Beech. Voy. 299 = Zexmenia ceanothifolia, *Benth. & Hook. f.*
ceanothifolia, Willd. Sp. Pl. iii. 2225 = Zexmenia ceanothifolia, *Sch. Bip.*
chinensis, Linn. Sp. Pl. 901 = Anisopappus chinensis.
ciliata, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 391.—Afr. trop.
Clausseni, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. iii. 211 = glabrata.
Clausseni, *Sch. Bip. ex Baker, l. c.* 212.—Bras.
coahuilensis, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 106.—Am. bor. occ.
connata, Gaudich. in Freyc. Voy. Bot. 464 = Lipochaeta connata.
conyzoides, [Moç. & Sesse,] ex DC. Prod. v. 616 = virgata.
conyzoides, *Trew, Pl. Rar.* 8. t. 6.—Hab.?
Coreopsis, Michx. Fl. Bor. Am. ii. 134 = Actinomeris squarrosa.
Coulteri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 13.—Mexic.
crocata, *Less. Syn. Comp.* 232.—Mexic.
Cumingii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
debilis, Spreng. Neue Entdeck. ii. 137 = Eclipta pusilla.
decurrens, *Vell. Fl. Flum.* viii. t. 114.—Bras.
dentata, *H. B. & K. Nov. Gen. et Sp.* iv. 205.—Ecuador.
dichotoma, Murr. in Comm. Gotting. (1779) 15. t. 4 = Blainvillea rhomboidea.
dichotoma, Reinw. ex Miq. Fl. Ind. Bat. ii. 75 = Wollastonia moluccana.
dichotoma, Sieber, ex Steud. Nom. ed. II. ii. 751 = Synedrella nodiflora.
dichotoma, Wall. Cat. n. 3204 = Blainvillea rhomboidea, Wedelia biflora.
dissita, *A. Gray, in Proc. Am. Acad.* xx. (1885) 299.—Calif.
diversifolia, *DC. Prod.* v. 615.—Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* iv. 204.—Ecuador.
encelioides, *Benth. & Hook. f. ex A. Gray, in Bot. Calif.* i. 350.—Am. bor. et austr.
floribunda, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 407.—Bras.
foliacea, Spreng. Syst. iii. 578 = Eleutheranthera ovata.
Fraseri, *Hemsl. Biol. Centr. Am. Bot.* ii. 187.—Mexic.
fruticosa, Burm. ex Spreng. Syst. iii. 602 = Wulffia stenoglossa?
fruticosa, Desf. Cat. Hort. Par. ed. III. 185; et ex DC. Prod. v. 610 = Zexmenia monocephala.
fruticosa, Linn. Syst. ed. X. 1226 = ilicifolia.
fruticosa, Willd. Sp. Pl. iii. 577 = tridentata.
gigantea, *Jacq. Ic. Pl. Rar.* i. 17. t. 175.—Am. trop.
glabrata, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 315.—Bras.
Grayi, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 188.—Mexic.
Grisebachii, *Baker, in Mart. Fl. Bras.* vi. iii. 214.—Reg. Argent.
guianensis, *Baker, l. c.* 211.—Guian.
hastata, Kellogg, ex Curran, in Bull. Calif. Acad. i. iii. (1885) 140 = venosa.
hastulata, Hook. & Arn. Bot. Beech. Voy. 87 = Lipochaeta hastulata.
helianthoides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 424 = glabrata.
helianthoides, Griseb. in Goett. Abh. xxiv. (1879) 194 = Grisebachii.
helianthoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 316 = Arnottii.
helianthoides, *H. B. & K. Nov. Gen. et Sp.* iv. 204.—Am. bor. et austr.
helianthoides, Michx. Fl. Bor. Am. ii. 135 = Actinomeris helianthoides.

VERBESINA:—
heterophylla, *A. Gray, in Proc. Am. Acad.* xix. (1883) 12.—Am. bor.
heterosperma, Griseb. in Goett. Abh. xxiv. (1879) 193 = encelioides.
Hookerii, Klatt, in Leopoldina, xx. (1884) 92 = Arnottii.
Humboldtii, Spreng. Syst. iii. 577 = helianthoides.
hypoleuca, *A. Gray, in Proc. Am. Acad.* xv. (1880) 37.—Mexic.
ilicifolia, *Poir. Encyc.* viii. 459.—Ins. S. Doming.
involucrata, *A. Rich. Tent. Fl. Abyss.* i. 409.—Abyss.
laciniata, Nutt. Gen. Am. ii. 170 = virginica.
laciniata, Walt. Fl. Carol. 213 (Quid?).—Am. bor.
lanceolata, Poir. Encyc. viii. 460 = Blainvillea rhomboidea.
lancifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 406 = diversifolia.
lavarum, Gaudich. in Freyc. Voy. Bot. 464 = Lipochaeta lavarum.
Lavenia, Linn. Sp. Pl. 902 = Adenostemma viscosum.
Lavenia, Roxb. Hort. Beng. 62; Fl. Ind. iii. 442 = Blainvillea rhomboidea.
leprosa, *Klatt, in Leopoldina*, xx. (1884) 93.—Ins. Martinic.
lineata, A. Rich. Tent. Fl. Abyss. i. 410 = Coreopsis Prestinaria.
linifolia, *Linn. Syst.* ed. X. 1226.—Jamaic.
lobata, Gaudich. in Freyc. Voy. Bot. 464 = Lipochaeta lobata.
longifolia, A. Gray, in Proc. Am. Acad. xix. (1883) 12 = Actinomeris longifolia.
longipes, *Hemsl. Biol. Centr. Am. Bot.* ii. 188.—Mexic.
macrantha, A. Rich. Tent. Fl. Abyss. i. 408 = Coreopsis macrantha.
mameana, *André, in Rev. Hortic.* (1885) 14, 16. f. 5.—Ecuador.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
megapotamica, *Spreng. Syst.* iii. 578.—Bras.
membranifolia, Poir. Encyc. Suppl. iii. 18 = Wulffia stenoglossa.
mexicana, Cerv. ex DC. Prod. v. 609 = Perymenium Cervantesii.
microcephala, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 98.—Mexic.
microcephala, Sch. Bip. ex A. Gray, in Proc. Am. Acad. xix. (1883) 12 = pinnatifida.
microptera, DC. Prod. v. 616 = virginica.
mollis, Alam. ex DC. l. c. 588 = Viguiera buddleiaeformis.
mollis, *H. B. & K. Nov. Gen. et Sp.* iv. 203.—Mexic.
moluccana, Blume, Bijdr. 912 (= *Wollastonia moluccana*).—Ins. Molucc.
montana, Blume, l. c. 911 (= *Wollastonia montana*).—Java.
montevidensis, Spreng. Syst. iii. 578 = Aspilia buphthalmiflora.
monticola, Hook. f. in Journ. Linn. Soc. vii. (1864) 200 = Coreopsis monticola.
mutica, Linn. Sp. Pl. ed. II. 1273 = Chrysanthellum procumbens.
mutica, Sieber, ex Steud. Nom. ed. II. ii. 751 = Wulffia stenoglossa.
neriifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 188.—Mexic.
nicaraguensis, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 97 = pinnatifida.
nicotianaefolia, *Baker, in Mart. Fl. Bras.* vi. iii. 212.—Bras.
nodiflora, Linn. Cent. Pl. i. 28; Amoen. Acad. iv. 290 = Synedrella nodiflora.
nudicaulis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 12.—Am. bor.
oaxacana, *DC. Prod.* v. 614.—Mexic.
occidentalis, *Walt. Fl. Carol.* 213.—Am. bor.
Oerstediana, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 96.—Costa Rica.
olivacea, *Klatt, in Leopoldina*, xx. (1884) 93.—Mexic.
oppositiflora, Poir. Encyc. viii. 460 = Wulffia stenoglossa.
ovata, A. Gray, in Proc. Am. Acad. xix. (1883) 13 = Actinomeris ovata.
ovata, Hort. Par. ex Poir. Encyc. viii. 454 = Heliopsis bupthalmoides.

VERBESINA:—
ovatifolia, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 189.—Mexic.
pallens, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 97.—Mexic.
paniculata, Poir. Encyc. viii. 456 = virginica.
pauciflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 189.—Mexic.
persicifolia, *DC. Prod.* v. 614.—Mexic.
Phaetusa, Cass. ex DC. l. c. 615 = occidentalis.
pinnata, *Clark, ex DC. l. c.*—Hab. ?
pinnatifida, *Cav. Ic.* i. 67. t. 100.—Mexic.
pinnatifida, Sw. Prod. Veg. Ind. Occ. 114 = gigantea.
platyptera, *Benth. & Hook. f. Gen.* ii. 350, in obs.—Am. trop.
podocephala, A. Gray, Pl. Wright. ii. 92 = Zexmenia podocephala.
polycephala, DC. Prod. v. 616 = virginica.
populifolia, *Hill, Hort. Kew.* 31.—China.
prostrata, *Hook. & Arn. Bot. Beech. Voy.* 195.—China.
prostrata, Linn. Sp. Pl. 902 = Eclipta erecta.
Pseudo-acmella, Linn. l. c. 901 = Spilanthes Acmella.
pterocaula, [*Moç. & Sesse,*] *ex DC. Prod.* v. 616.—Mexic.
pusilla, Poir. Encyc. viii. 459 = Eclipta pusilla.
repanda, Pers. Syn. ii. 471 = Anthemis repanda.
Rueppellii, A. Rich. Tent. Fl. Abyss. i. 410 = Coreopsis chrysantha.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 205.—Mexic.
Sartori, Hort. ex André, in Rev. Hortic. (1885) 15 = pinnatifida.
sativa, Roxb. ex Sims, Bot. Mag. t. 1017; et Fl. Ind. iii. 441 = Guizotia abyssinica.
Saubinetia, *Klatt, in Leopoldina*, xx. (1884) 92.—Chili.
scaberrima, Wall. Cat. n. 3208 = Wedelia Wallichii.
scabra, Benth. Pl. Hartw. 41 = Actinomeris tetraptera.
scabra, Phil. Sert. Mendoc. Alt. 28 = encelioides.
scandens, *Roxb. Hort. Beng.* 62; *Fl. Ind.* iii. 441.—Ind. or.
Schomburgkii, *Sch. Bip. in Schomb. Faun. Fl. Brit. Gui.* 1078, nomen.—Guian.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 306.—Mexic.
semiserrata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 214 = sordescens.
sericea, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 14.—Mexic.
serrata, *Cav. Ic.* iii. 7. t. 214.—Mexic.; Cuba.
Siegesbeckia, Michx. Fl. Bor. Am. ii. 134 = occidentalis.
sinensis, J. F. Gmel. Syst. 1256 = *chinensis*.
sinuata, Ell. Sketch, ii. 411 = virginica.
solitaria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 (Wedeliae sp.).—Java.
Soratae, *Sch. Bip. in Linnaea*, xxxiv. (1865-66) 528, nomen.—Bolivia.
sordescens, *DC. Prod.* v. 613.—Bras.
sororia, *A. Gray, in Proc. Am. Acad.* xv. (1880) 37.—Mexic.
spicata, Lour. Fl. Cochinch. 507 = Eclipta spicata.
stricta, A. Gray, in Proc. Am. Acad. xix. (1883) 13 = Actinomeris stricta.
strigulosa, Gaudich. in Freyc. Voy. Bot. 463 = Wedelia aristata.
subcordata, *DC. Prod.* v. 614.—Bras.
sublobata, *Benth. Pl. Hartw.* 76.—Mexic.
subpanduraeformis, Klatt, in Leopoldina, xx. (1884) 92 = subcordata.
succulenta, Hook. & Arn. Bot. Beech. Voy. 87 = Lipochaeta succulenta.
tetraptera, A. Gray, in Proc. Am. Acad. xix. (1883) 13 = Actinomeris tetraptera.
texana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862), 458 = virginica.
tomentosa, *DC. Prod.* v. 614.—Am. austr.
tridentata, *Spreng. Syst.* iii. 577.—Am. austr.
triplinervia, *Vis. in Nuov. Sagg. Acc. Sc. Fis. Padov.* v. (1840) 264.—Mexic.
triradiata, *Vell. Fl. Flum.* viii. t. 115.—Bras.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* iv. 203.—N. Granat.
urticaefolia, Blume, Bijdr. 911 = Wedelia urticaefolia.
urticaefolia, Span. ex Miq. Fl. Ind. Bat. ii. 73 = Wedelia aristata.
venosa, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 110.—Ins. Cedros.
verbascifolia, *Walp. in Bot. Zeit.* ix. (1851) 63.—Venezuela.

VERBESINA:—
veris, A. Rich. Tent. Fl. Abyss. i. 407 = Coreopsis Prestinaria.
vigueroides, *Baker, in Mart. Fl. Bras.* vi. III. 215.—Paraguay.
villosa, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 370 = virginica.
virgata, *Cav. Ic.* iii. 38. t. 275.—Mexic.
virginica, *Linn. Sp. Pl.* 901.—Am. bor.
Warei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 12.—Am. bor.
Wrightii, *A. Gray, l. c.*—Am. bor.
Wrightii, *Griseb. Cat. Pl. Cub.* 155.—Cuba.

VERBIASCUM, Fenzl, Diagn. Pl. Or. 47 (1880), sphalm. = **Verbascum**, Tourn. (Scrophul.).

VEREA, Willd. Sp. Pl. ii. 471 (1799) = seq.

VEREIA, Andr. Bot. Rep. t. 21 (1797) = **Kalanchoë**, Adans. (Crassul.).
acutiflora, Andr. l. c. t. 560 = K. acutiflora.
aegyptiaca, Steud. Nom. ed. II. ii. 751 = K. crenata.
alternans, Spreng. Syst. ii. 260 = K. alternans.
ceratophylla, D. Dietr. Syn. Pl. ii. 1328 = K. ceratophylla.
crenata, Andr. Bot. Rep. t. 21 = K. Afzeliana.
floribunda, D. Dietr. Syn. Pl. ii. 1328 = K. floribunda, *Wight & Arn.*
grandiflora, D. Dietr. l. c. = K. grandiflora, *Wight & Arn.*
heterophylla, D. Dietr. l. c. = K. glandulosa.
laciniata, Willd. Enum. Hort. Berol. 433 = K. laciniata.
lanceolata, Spreng. Syst. ii. 260 = K. lanceolata.
nudicaulis, Spreng. l. c. = K. crenata.
pinnata, Spreng. l. c. = Bryophyllum calycinum.
rotundifolia, D. Dietr. Syn. Pl. ii. 1328 = K. rotundifolia.
spathulata, D. Dietr. l. c. 1327 = K. spathulata.

VERHUELLIA, Miq. Syst. Pip. 47 (1843). *PIPERACEAE*, Benth. & Hook. f. iii. 131.
MILDEA, Griseb. Cat. Pl. Cub. 63 (1866).
brasiliensis, Miq. Syst. Pip. 48 = Peperomia brasiliensis.
cordifolia, *C. DC. in DC. Prod.* xvi. I. 391.—Cuba.
elegans, *Miq. Syst. Pip.* 48.—Ind. occ.
Knoblocheriana, *C. DC. in DC. Prod.* xvi. I. 391.—Afr. trop.
serpens, Griseb. Fl. Brit. W. Ind. 164 = elegans.
serpens, *Miq. Syst. Pip.* 49.—Jamaic.

VERINEA, Pomel, Mat. Fl. Atl. 1 (1860) = **Asphodelus**, Tourn. (Liliac.).
fistulosa, Pomel, l. c. 2 = A. fistulosus.
tenuifolia, Pomel, l. c. = A. tenuifolius.

VERLANGIA, Neck. Elem. ii. 125 (1790); Rafin. Sylva Tellur. 33 (1838) = **Argania**, Roem. & Schult. (Sapot.).
Argan, Rafin. l. c. 34 = A. Sideroxylon.
indica, Rafin. l. c. = Flacourtia sepiaria.
sicula, Rafin. l. c. = A. Sideroxylon.

VERLOTIA, Fourn. in Mart. Fl. Bras. vi. IV. 324 (1885). *ASCLEPIADEAE*.
dracontea, *Fourn. l. c.* 325.—Bras.
heterophylla, *Fourn. l. c.*—Bras.
macrocalyx, *Fourn. l. c.*—Bras.
suberosa, *Fourn. l. c.* 324.—Bras.
virgultorum, *Fourn. l. c.*—Bras.
Weddellii, *Fourn. l. c.* 326.—Bras.

VERMICULARIA, Moench, Meth. Suppl. 150 (1802) = **Stachytarpheta**, Vahl (1805) (Verben.).
decurrens, Moench, l. c. = S. indica.
lancifolia, Moench, l. c. = S. indica.

VERMIFUGA, Ruiz & Pav. Prod. 114. t. 24 (1794) = **Flaveria**, Juss. (Compos.).
corymbosa, Ruiz & Pav. Syst. Veg. 216 = F. Contrayerba.

VERMONETA, Comm. ex Juss. Gen. 343, in obs. (1789) = **Homalium**, Jacq. (Samyd.).
decadenia, Comm. ex Juss. l. c. = H. paniculatum.

VERMONIA, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 172, err. typ. = **Vernonia**, Schreb. (Compos.).

VERMONTEA, Steud. Nom. ed. I. 111, 874 (1821), sphalm. = **Homalium**, Jacq. (Samyd.).
decandra, Steud. l. c. 111 = H. paniculatum.

VERNASOLIS, Rafin. Atl. Journ. 152 (1832) = **Gaillardia** ?, Fouger. (Compos.).
auriculata, glauca, heterophylla, Rafin. l. c. (Quid ?).—Am. bor.

VERNICIA, Lour. Fl. Cochinch. 586 (1790) = **Aleurites**, Forst. (Euphorb.).
montana, Lour. l. c. = A. triloba.

VERNISECKIA, Steud. Nom. ed. II. ii. 752 (1841) = WERNISECKIA, Scop. = **Humiria**, Aubl.

VERNIX, Adans. Fam. ii. 347 (1763) = **Rhus**, Tourn. (Anacard.).

VERNONELLA, Sond. in Linnaea, xxiii. (1850) 62 = **Vernonia**, Schreb. (Compos.).
africana, Sond. l. c. = Vernonia Vernonella.

VERNONIA, Schreb. Gen. ii. 541 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 227.
ACHYROCOMA, Cass. in Dict. Sc. Nat. xxvi. 21 (1823); lvii. 341 (1828).
ACILEPIS, D. Don, Prod. Fl. Nep. 169 (1825).
AMBASSA, Steetz, in Peters, Reise Mossamb. Bot. 364 (1863).
ASCARICIDA, Cass. in Dict. Sc. Nat. iii. Suppl. 38 (1816).
BACCHAROIDES, Moench, Meth. 578 (1794).
BECHIUM, DC. Prod. v. 70 (1836).
BEHEN, Hill, Veg. Syst. iv. 41 (1762).
BRACHEILEMA, R. Br. in Salt, Abyss. App. 65 (1814).
CANDIDEA, Tenore, in Atti Acc. Nap. iv. (1839) 104. tt. 1, 2.
CENTRAPALUS, Cass. in Bull. Soc. Philom. (1817) 10; et in Dict. Sc. Nat. vii. 382 (1817).
CHELIUSIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26.
CLAOTRACHELUS, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 565.
CRITONIOPSIS, Sch. Bip. in Pollichia, xx–xxi. (1863) 340.
CRYSTALLOPOLLEN, Steetz, in Peters, Reise Mossamb. Bot. 363 (1863).
CYANOPIS, Blume, Fl. Jav. Praef. p. vi. (1828); et in DC. Prod. v. 69 (1836).
CYANOPSIS, Endl. Ench. 232 (1841).
CYANTHILLIUM, Blume, Bijdr. 889 (1826).
DISTEPHANUS, Cass. in Bull. Soc. Philom. (1817) 151.
FLUSTULA, Rafin. Sylva Tellur. 116 (1838).
GYMNANTHEMUM, Cass. in Bull. Soc. Philom. (1817) 10.
HOLOLEPIS, DC. in Ann. Mus. Par. xvi. (1810) 189. t. 6.
ISOMERIA, D. Don, ex DC. Prod. v. 24 (1836).
ISONEMA, Cass. in Bull. Soc. Philom. (1817) 152.
KERINGA, Rafin. Sylva Tellur. 144 (1838).
LEIBOLDIA, Schlecht. in Linnaea, xix. (1847) 742.
LEPIDAPLOA, Cass. in Bull. Soc. Philom. (1817) 66; et in Dict. Sc. Nat. xxvi. 16 (1823).
LEUCOMERIS, Blume, ex DC. Prod. v. 21 (1836).
LINZIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26.
LLERASIA, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 37.
LYSISTEMMA, Steetz, in Peters, Reise Mossamb. Bot. 340 (1863).
MONOSIS, DC. Prod. v. 77 (1836).
PLECTRECA, Rafin. Fl. Tellur. iv. 119 (1836).
POLYDORA, Fenzl, in Flora, xxvii. (1844) 312.
PUNDUANA, Steetz, in Peters, Reise Mossamb. Bot. 345 (1863).
STENGELIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26.
STENOCEPHALUM, Sch. Bip. in Pollichia, xx–xxi. (1863) 385.
STROBOCALYX, Sch. Bip. l. c. xviii–xix. (1861) 170.
SUPRAGO, Gaertn. Fruct. ii. 402. t. 166 (1791).

VERNONIA:—
Teichostemma, R. Br. in Salt, Abyss. App. 65 (1814).
Tephrothamnus, Sch. Bip. in Pollichia, xx–xxi. (1863) 431.
Trianthaea, Spach, Hist. Vég. Phan. x. 39 (1841).
Turpinia, La Llave & Lex. Nov. Veg. Desc. fasc. i. 24 (1824).
Vernonella, Sond. in Linnaea, xxiii. (1850) 62.
Webbia, DC. Prod. v. 72 (1836).
Xipholepis, Steetz, in Peters, Reise Mossamb. Bot. 344 (1863).
abbreviata, DC. Prod. v. 25 = cinerea.
abyssinica, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) 1. *Intell.* 26, nomen; *et in Walp. Rep.* ii. 947.—Abyss.
accedens, *Miq. Fl. Ind. Bat.* ii. 18.—Java.
Achyrocoma, *Less. in Linnaea*, iv. (1829) 313.—Am. bor.?
Acilepis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 68.—Nicaragua.
acuminata, DC. Prod. v. 32 = volkameriaefolia.
acuminata, *Less. in Linnaea*, vi. (1831) 663.—Ind. occ.
acuminata, Willd. ex Less. l. c. iv. (1829) 284 = canescens.
acutangula, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 225.—Bras.
acutangula, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 123 = Piptocarpha macropoda.
acutifolia, Hook. Bot. Mag. t. 3062 = sericea.
adamantium, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 222.—Bras.
adenophylla, *Mart. ex DC. Prod.* v. 17.—Bras.
adenosticta, Fenzl, ex Walp. Rep. vi. 98 = amygdalina.
adoënsis, *Sch. Bip. ex Walp. l. c.* ii. 946.—Abyss.
adproximata, Less. in Linnaea, vi. (1831) 655 = diffusa.
affinis, *H. B. & K. Nov. Gen. et Sp.* iv. 39.—Peruv.
affinis, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 119, nomen (Quid?).—Malaya.
Alamani, *DC. Prod.* v. 61.—Mexic.
alata, Heyne, ex DC. l. c. 448 = Laggera alata.
albicans, DC. in Wight, Contrib. 6 = cinerea, chinensis.
albicaulis, Vahl, ex Pers. Syn. ii. 404 = longifolia.
albicola, Steud. Nom. ed. II. ii. 752, sphalm. = *alpicola*.
albiflora, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 224 = geminata.
albiflora, Rafin. Herb. 29 (Quid?).—Am. bor.
alpestris, *Baker, in Mart. Fl. Bras.* vi. II. 55.—Bras.
alpicola, Mart. ex DC. Prod. v. 17 = Piptolepis buxoides.
altissima, *Nutt. Gen. Am.* ii. 134.—Am. bor.
altissima, Walp. Rep. ii. 540 = angustifolia.
amara, Stokes, Bot. Mat. Med. iv. 165 = Saussurea amara.
ambigua, *Kotschy & Peyr. Pl. Tinn.* 35. t. 17 B.—Afr. trop.
ammophila, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 227.—Bras.
amygdalina, *Delile, Cent. Pl. Méroë*, 41.—Afr. trop.
anceps, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 233.—Zeylan.
Andersoni, C. B. Clarke, Comp. Ind. 27 = Cumingiana.
Andrieuxii, DC. Prod. v. 16 = Liabum Andrieuxii.
andromachioides, Less. in Linnaea, vi. (1831) 397 = Liabum andromachioides.
angulifolia, *DC. Prod.* v. 29.—Afr. austr.
angustifolia, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 236 = nudiflora.
angustifolia, *Michx. Fl. Bor. Am.* ii. 94.—Am. bor.
anisochaetoides, *Sond. in Linnaea*, xxiii. (1850) 61.—Afr. austr.
anthelmintica, *Willd. Sp. Pl.* iii. 1634.—Reg. Himal.
aphanantha, *Baker, in Journ. Linn. Soc.* xx. (1883) 176.—Madag.
apiculata, *Mart. ex DC. Prod.* v. 51.—Bras.
apocynifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 175.—Madag.
appendiculata, *Less. in Linnaea*, vi. (1831) 636.—Madag.
approximata, Baker, in Mart. Fl. Bras. vi. II. 23 = diffusa.
arabica, Decne. ex Boiss. Fl. Orient. iii. 154 = atriplicifolia.

VERNONIA:—
araneosa, *Baker, in Mart. Fl. Bras.* vi. II. 32.—Bras.
ararana, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 227.—Bras.
araripensis, *Gardn. l. c.* 222.—Bras.
arborea, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1824) 218.—As. trop.
arborescens, *Sw. Fl. Ind. Occ.* ii. 1320.—Ind. occ.
Arbuscula, *Less. in Linnaea*, vi. (1831) 664.—Ins. Maurit.
arbutilifolia, *Baker, in Journ. Bot.* xx. (1882) 169, in obs.—Madag.
arctioides, *Less. in Linnaea*, vi. (1831) 400.—Mexic.
arcuata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 220 = linearis.
arenaria, Gardn. l. c. 221 = grisea.
arenaria, *Mart. ex DC. Prod.* v. 54.—Bras.
argentea, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 108 = Hyalis argentea.
argentea, *Less. in Linnaea*, vi. (1831) 672.—Bras.
arguta, *Baker, in Journ. Linn. Soc.* xx. (1883) 175.—Madag.
argyropappa, Buek, Index ad DC. Prod. ii. Praef. p. v = Salzmanni.
argyrophylla, *Less. in Linnaea*, vi. (1831) 627.—Bras.
argyrotrichia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 96.—Bras.
ariaefolia, Mart. ex DC. Prod. v. 30 = argyrophylla.
aristata, *Less. in Linnaea*, iv. (1829) 313.—Hab.?
aristata, Reinw. ex De Vriese, Pl. Ind. Bat. Or. 130 = moluccensis.
aristata, Sch. Bip. in Flora, xviii. (1844) 667 = natalensis.
arkansana, *DC. Prod.* vii. 264.—Am. bor.
Arnottiana, Thoms. ex C. B. Clarke, Comp. Ind. 20 = salvifolia.
Aschenborniana, *S. Schau. in Linnaea*, xix. (1847) 714.—Mexic.
Aschersonii, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 161.—Afr. trop.
aspera, Buch.-Ham. in Trans. Linn. Soc. xiv. (1824) 219, partim = divergens, Roxburghii.
aspera, DC. Prod. v. 31 = Roxburghii.
aspera, Less. in Linnaea, vi. (1831) 643 = divergens?
assans, Mart. ex DC. Prod. v. 38 = scabra.
asteriflora, *Mart. l. c.* 29.—Bras.
asterotrichia, Benth. ex Baker, in Mart. Fl. Bras. vi. II. 124 = Piptocarpha opaca.
asterotrichia, Poepp. & Endl. Nov. Gen. et Sp. iii. 41. t. 247 = Piptocarpha asterotrichia.
atriplicifolia, *Jaub. & Spach, Illustr.* iv. 94. t. 359.—Arab.
attenuata, *DC. Prod.* v. 33.—Reg. Himal.; Burma.
aurea, *Mart. ex DC. l. c.* 58.—Bras.
auriculata, *Griseb. in Goett. Abh.* xxiv. (1879) 163.—Reg. Argent.
aurifodinarum, Mart. ex DC. Prod. v. 48 = obscura.
axillaris, *Less. in Linnaea*, iv. (1829) 256.—Bras.
axillaris, Sch. Bip. in Pollichia, xx–xxi. (1863) 418 = Piptocarpha macropoda.
axilliflora, Mart. ex Less. in Linnaea, iv. (1829) 297 = V. Cotoneaster.
baccharoides, *H. B. & K. Nov. Gen. et Sp.* iv. 40.—N. Granat.
bahamensis, *Griseb. Fl. Brit. W. Ind.* 352.—Ins. Baham.
Bainesii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 272.—Afr. trop.
Bakeri, *Vatke, in Bremen, Abh.* ix. (1885) 119.—Madag.
Baldwini, *Torr. in Ann. Lyc. N. York*, ii. (1828) 211.—Am. bor.
barbata, *Less. in Linnaea*, iv. (1829) 287.—Bras.
barbinervis, *Sch. Bip. in Seem. Bot. Voy. Herald*, 297.—Mexic.
bardanoides, *Less. in Linnaea*, vi. (1831) 669.—Bras.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1883) 173.—Madag.
Beddomei, *Hook. f. Fl. Brit. Ind.* iii. 231.—Ind. or.
Benthamiana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 282.—Afr. trop.
Berteriana, DC. Prod. v. 52 = arborescens.
Beyrichii, *Less. in Linnaea*, iv. (1829) 275.—Bras.
Biafrae, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 270.—Afr. trop.
bifrons, Vahl, ex Less. in Linnaea, v. (1830) 261 = Anastraphia ilicifolia.
Bipontini, Vatke, l. c. xxxix. (1875) 478 = Leopoldi.

VERNONIA:—
Blanchetiana, DC. Prod. v. 20 = Piptocarpha Lundiana.
blanda, *DC. l. c.* 32.—Burma.
blandula, C. B. Clarke, Comp. Ind. 26 = praec.
Blumeana, DC. Prod. v. 22 = arborea.
blumeoides, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 198.—Afr. trop.
Bojeri, *Less. in Linnaea*, vi. (1831) 676.—Afr. trop.
Bolleana, *Sch. Bip. in Seem. Bot. Voy. Herald*, 297.—Mexic.
borneensis, *Miq. Fl. Ind. Bat.* ii. 16.—Borneo.
Bottae, *Jaub. & Spach, Illustr.* iv. 92.—Arab.
brachiata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 67.—Costa Rica.
brachylepis, *Griseb. in Goett. Abh.* xxiv. (1879) 164.—Reg. Argent.
brachyscypha, *Baker, in Journ. Linn. Soc.* xx. (1883) 178.—Madag.
bracteata, *Wall. Cat.* n. 2921.—Reg. Himal.
bracteolata, DC. Prod. v. 62 = subsessilis.
brasiliensis, Less. in Linnaea, vi. (1831) 682 = Albertinia brasiliensis.
brasiliensis, Mart. ex DC. Prod. v. 39 = polyanthes.
brasiliensis, Sch. Bip. in Pollichia, xviii–xix. (1861) 161 = Vanillosmopsis brasiliensis.
brevifolia, *Less. in Linnaea*, iv. (1829) 285.—Bras.
brevifolia, Rafin. New Fl. Am. iv. 77 (Quid?).—Am. bor.
brevipetiolata, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 85.—Bras.
brunioides, Less. in Linnaea, iv. (1829) 248 = Lychnophora brunioides.
Buchingeri, *Sch. Bip. ex Steetz, in Peters, Reise Mossamb. Bot.* 359.—Abyss.
buddleiaefolia, *Mart. ex DC. Prod.* v. 45.—Bras.
bullata, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 67.—Nicaragua.
bupleurifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 87 = obtusata.
Burchelliana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 209 = Piptolepis Oleaster.
Burtoni, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 281.—Afr. trop.
buxifolia, *Less. in Linnaea*, iv. (1829) 313.—Ins. S. Doming.
buxoides, Less. l. c. 247 = Piptolepis buxoides.
caducissima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 73 = Salzmanni.
callicephala, Sch. Bip. ex Baker, l. c. 32 = V. Radula.
Calvoana, *Hook. f. Bot. Mag.* t. 5698.—Afr. trop.
calycina, *Wall. Cat.* n. 2924.—Burma.
Camara, *Liais, Brés.* 627.—Bras.
campestris, DC. Prod. v. 43 = desertorum.
campestris, Mart. ex DC. l. c. 45 = grandiflora.
cana, Edgew. in Journ. As. Soc. Beng. xvi. II. (1847) 1216 = atriplicifolia?
Candolleana, *Wight & Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 346.—Ind. or.
Candollei, *DC. Prod.* v. 21.—Bras.
Candollei, *Vatke, in Bremen, Abh.* ix. (1885) 119.—Madag.
canescens, *H. B. & K. Nov. Gen. et Sp.* iv. 35.—Peruv.
canescens, Sch. Bip. in Flora, xxvii. (1844) 667 = pinifolia.
capitata, Less. in Linnaea, iv. (1829) 270 = Vanillosmopsis capitata.
capitellata, DC. Prod. v. 20 = Piptocarpha leprosa.
capituliflora, *Miq. Fl. Ind. Bat.* ii. 19.—Ins. Sunda.
carbonaria, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 28 = megapotamica.
carduoides, *Baker, l. c.* 34.—Bras.
celebica, DC. Prod. v. 21 = arborea.
centaurioides, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 536, nomen.—Bolivia.
centriflora, Link & Otto, Ic. Pl. Sel. 117. t. 55 = scorpioides.
Cephalotes, *DC. Prod.* v. 57.—Bras.
cernua, Boj. ex DC. l. c. vii. 24 (= *Moquinia Bojeri*).—Afr. trop.
chalybaea, *Mart. ex DC. l. c.* v. 54.—Bras.
Chamaedrys, *Less. in Linnaea*, iv. (1829) 259.—Bras.
Chamaepeuces, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 31.—Bras.
Chamissonis, *Less. in Linnaea*, iv. (1829) 304.—Bras.
chiliantha, *Walp. l. c.* xiv. (1840) 508.—Peruv.; Chili.
chinensis, *Less. l. c.* vi. (1831) 105, 674.—Malaya; China.

VERNONIA:—
chromolepis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 224 = rubricaulis.
chrysophylla, Gardn. l. c. vi. (1847) 417 = obovata.
cincta, *Griseb. in Goett. Abh.* xxiv. (1879) 162.—Reg. Argent.
cinerascens, *Sch. Bip. in Schweinf. Beitr. Fl. Aethiop.* 162.— Ind. bor. occ.; Abyss.
cinerea, *Less. in Linnaea*, iv. (1829) 291.—Amphig. trop.
cirsiiflora, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 36 = bardanoides.
cirsiifolia, Steud. Nom. ed. II. ii. 753 = Pacourina cirsiifolia.
clarenceana, Hook. f. in Journ. Linn. Soc. vi. (1862) 11 = Conyza clarenceana.
clavata, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 220.—Bras.
clivorum, *Hance, in Journ. Bot.* vii. (1869) 164.—China.
Cockburniana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 837.—Ins. Socotra.
coelestina, Schrad. ex DC. Prod. v. 62 = pauciflora.
cognata, *Less. in Linnaea*, vi. (1831) 670.—Bras.
collina, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 213 = mucronulata.
compacta, *Gardn. l. c.* 216.—Bras.
compactiflora, *Mart. ex Baker, in Mart. Fl. Bras.* vi. II. 44.—Bras.
complanata, Sch. Bip. ex Baker, l. c. 104 = densiflora.
complicata, *Wright, ex Griseb. Cat. Pl. Cub.* 143.—Cuba.
condensata, *Baker, in Journ. Bot.* xiii. (1875) 202.—Bras.
conferta, *Benth. in Hook. Niger Fl.* 427.—Afr. trop.
conferta, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 124 = Piptocarpha Riedelii.
congesta, Benth. in Hook. Lond. Journ. Bot. i. (1842) 487 = Inula Cappa.
conyzoides, DC. in Wight, Contrib. 6 = cinerea.
conyzoides, Thw. Enum. Pl. Zeyl. 160 = Hookeriana.
conyzoides, Wight, Ic. t. 829 = Candolleana.
corcovadensis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 218 = polyanthes.
cordata, *H. B. & K. Nov. Gen. et Sp.* iv. 39.—Mexic.
cordiaefolia, *H. B. & K. l. c.* 38.—N. Granat.
cordifolia, Willd. ex Less. in Linnaea, iv. (1829) 267 = cordata.
cordigera, *Mart. ex DC. Prod.* v. 58.—Bras.
coriacea, *Less. in Linnaea*, vi. (1831) 661.—Bras.
corymbiformis, *DC. Prod.* v. 62.—Mexic.
corymbosa, *Less. in Linnaea*, vi. (1831) 647.—Afr. austr.
corymbosa, Schwein. in Long, Exped. Winnep. ii. 394 = fasciculata.
corymbulosa, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 113 = viscidula.
Cotoneaster, *Less. in Linnaea*, iv. (1829) 298.—Bras.
Coulonii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 93.—Bras.
crenata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 218 = ferruginea.
crenulata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 96 = ignobilis.
crinita, Rafin. New Fl. Am. iv. 77 (Quid?).—Am. bor.
crotonoides, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 166.—Bras.
cubensis, Griseb. Cat. Pl. Cub. 144 = hieracioides.
cuiabensis, *Baker, in Mart. Fl. Bras.* vi. II. 37.—Bras.
Cumingiana, *Benth. in Hook. Kew Journ.* iv. (1852) 232.—Reg. Himal.; China.
cuneata, Less. in Linnaea, vi. (1831) 644 = scandens.
cuneifolia, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 215.—Bras.
cuspidata, Buek, Ind. ad DC. Prod. ii. p. v = volkameriaefolia.
cuspidata, Steud. Nom. ed. II. ii. 753 = arborea?
cyanonioides, Walp. in Linnaea, xiv. (1840) 509 = cinerea.
cyanopidea, Hochst. ex Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 161 = Aschersonii.
cyanopioides, Walp. Rep. ii. 541, sphalm. = cinerea.

VERNONIA:—
cylindrica, A. Rich. Tent. Fl. Abyss. i. 374 = podocoma.
cylindrica, *Sch. Bip. ex Walp. Rep.* ii. 945.—Afr. trop.
cymosa, *Blume, Bijdr.* 894.—Java.
daphnoides, *Walp. in Linnaea*, xiv. (1840) 510.—Bras.
davalliaefolia, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 422 = obscura.
debilis, Mart. ex DC. Prod. v. 54 = V. Cotoneaster.
decantha, Sch. Bip. in Linnaea, xx. (1847) 512 = ehretifolia.
decumbens, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 115.—Bras.
decussata, Mart. ex DC. Prod. v. 58 = pycnostachya.
delapsa, *Baker, in Journ. Linn. Soc.* xx. (1883) 172.—Madag.
demulans, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 195.—Afr. trop.
dendigulensis, DC. Prod. vii. 263 = Candolleana.
densevillosa, Mart. ex DC. l. c. v. 43 = obovata.
densiflora, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 114.—Bras.
denticulata, DC. Prod. v. 36 = Beyrichii.
Deppeana, *Less. in Linnaea*, vi. (1831) 398.—Mexic.
desertorum, *Mart. ex DC. Prod.* v. 43.—Bras.
dichocarpa, *Less. in Linnaea*, iv. (1829) 269.—Mexic.
dichotoma, *Sch. Bip. l. c.* xxx. (1859–60) 173.—N. Granat.
diffusa, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 408 = cinerea.
diffusa, *Less. in Linnaea*, iv. (1829) 272.—Bras.
discolor, *Less. l. c.* 274.—Bras.
dissoluta, *Baker, in Journ. Linn. Soc.* xx. (1883) 174.—Madag.
divaricata, Less. in Linnaea, iv. (1829) 306 = arborescens.
divaricata, Sw. Fl. Ind. Occ. iii. 1319 = arborescens, acuminata.
divergens, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 172, nomen; *Benth. ex C. B. Clarke, Comp. Ind.* 14.—Ind. or.; Burma.
diversifolia, *Boj. ex DC. Prod.* v. 26.—Madag.
domingensis, DC. l. c. 30 = buxifolia.
Doniana, *DC. l. c.* 23.—Afr. trop.
Dregeana, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. austr.
dura, *Mart. ex DC. Prod.* v. 59.—Bras.
Echinops, Mart. ex Baker, in Mart. Fl. Bras. iv. II. 167 = Eremanthus sphaerocephalus.
echioides, *Less. in Linnaea*, iv. (1829) 278.—Bras.
echitifolia, *Mart. ex DC. Prod.* v. 60.—Bras.
edulis, Steud. Nom. ed. II. ii. 753 = Pacourina edulis.
Ehrenbergiana, *Sch. Bip. in Linnaea*, xx. (1847) 513.—Mexic.
ehretiaefolia, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 420 = varroniaefolia.
ehretifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 39.—Guian.
elaeagnifolia, *DC. Prod.* v. 22.—Burma.
elaeagnoides, H. B. & K. Nov. Gen. et Sp. iv. 42 = Piptocarpha elaeagnoides.
elaeagnoides, Sch. Bip. in Flora, xxvii. (1844) 668 = Kraussii.
Elaeagnus, Sch. Bip. in Pollichia, xviii–xix. (1861) 166 = Eremanthus Elaeagnus.
elegans, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 421.—Bras.
elliptica, DC. in Wight, Contrib. 5 = elaeagnifolia.
elongata, Willd. ex Less. in Linnaea, iv. (1829) 302 = gracilis.
emarginata, Wikstr. in Vet. Acad. Handl. Stockh. (1827) 73 = rigida.
eminens, *Bisch. Cat. Sem. Hort. Heidelb.* (1847).—*Cf. Ann. Sc. Nat.* Sér. III. ix. (1848) 320.—Afr. bor.
ensifolia, Mart. ex DC. Prod. v. 53 = glabrata.
eremophila, *Mart. ex DC. l. c.* 54.—Bras.
ericaefolia, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 236 = brevifolia.
ericoides, Less. in Linnaea, iv. (1829) 247 (excl. syn.) = Piptolepis ericoides.
erigeroides, DC. Prod. v. 25 = cinerea.
Erigerontis, Mart. ex DC. l. c. 43 = simplex.
erioclada, *DC. l. c.* 44.—Bras.
eriolepis, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 224.—Bras.

VERNONIA:—
eriosematoides, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 254 = Inula Cappa.
erubescens, Hochst. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 280 = Aschersonii.
Ervendbergii, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 203.—Mexic.
erythrophila, *DC. Prod.* v. 57.—Bras.
erythrophiloides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 52 = praec.
estrellensis, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 215 = puberula.
eupatorifolia, *DC. Prod.* v. 37.—Bras.
eupatorioides, *Blume, Bijdr.* 894.—Java.
exaltata, Hort. ex Kunth, Ind. Sem. Hort. Berol. (1848) 14 = Verbesina sericea.
exilis, *Miq. in Journ. Bot. Néerl.* i. (1861) 98.—China.
exsucca, Mart. ex DC. Prod. v. 85 = Eremanthus exsuccus.
extensa, *DC. l. c.* 33.—Reg. Himal.
extraxillaris, DC. l. c. 56 = linearis.
fagifolia, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 216.—Bras.
farinosa, *Baker, in Mart. Fl. Bras.* vi. II. 84.—Bras.
fasciculata, Blume, Bijdr. 893, in nota = cinerea.
fasciculata, *Michx. Fl. Bor. Am.* ii. 94.—Am. bor.
fastigiata, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 282.—Afr. trop.
ferruginea, *Less. in Linnaea*, iv. (1829) 271.—Bras.
filifolia, Mart. ex DC. Prod. v. 44 = brevifolia.
filigera, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 288.—Afr. trop.
fimbrillifera, Less. in Linnaea, iv. (1829) 313 (= *Decaneurum fimbrilliferum*).—Ins. Borbon.
firma, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 290.—Afr. trop.
firmula, Mart. ex DC. Prod. v. 59 = dura.
flagellaris, Vatke, in Linnaea, xxxix. (1875) 479 = urticaefolia.
flavescens, Less. l. c. vi. (1831) 657 = scorpioides.
flexuosa, *Sims, Bot. Mag.* t. 2477.—Bras.
floccosa, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 225.—Bras.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iv. 38.—Am. trop.
florida, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 212 = nitidula.
flotowioides, *Baker, in Mart. Fl. Bras.* vi. II. 81.—Bras.
foliosa, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 210.—Bras.
foliosa, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 161.—Mexic.
Fortunei, Sch. Bip. in Flora, xxxv. (1852) 48 = solanifolia.
fragrans, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 24.—Mexic.
Francavillana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 296.—Afr. trop.
frangulaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 37.—N. Granat.
frondosa, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 294.—Afr. trop.
fruticosa, Sw. Fl. Ind. Occ. iii. 1323 = rigida.
fruticulosa, *Mart. ex DC. Prod.* v. 53.—Bras.
fulta, *Griseb. in Goett. Abh.* xxiv. (1879) 164.—Reg. Argent.
fusco-pilosa, *Baker, in Journ. Linn. Soc.* xx. (1883) 179.—Madag.
galamensis, Less. in Linnaea, iv. (1829) 314 = pauciflora.
Gardneri, *Thw. Enum. Pl. Zeyl.* 161.—Zeylan.
Garnieriana, *Klatt, in Linnaea*, xxxvii. (1871–73) 508.—Madag.
Gaudichaudiana, DC. Prod. v. 21 = Piptocarpha umbellulata.
geminata, H. B. & K. Nov. Gen. et Sp. iv. 35 = canescens.
geminata, *Less. in Linnaea*, iv. (1829) 303.—Bras.
geminiflora, Poepp. in Poepp. & Endl. Nov. Gen. et Sp. iii. 42 = Salzmanni.
gerberaeformis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 285.—Afr. trop.
Gerrardi, *Harv. Thes. Cap.* ii. 36.—Afr. austr.
glabra, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 194.—Afr. trop.

VERNONIA:—
glabrata, *Less. in Linnaea*, iv. (1829) 294.—Bras.
glandulosa, *DC. Prod.* v. 22.—Madag.
glauca, Willd. Sp. Pl. iii. 1633 = noveboracensis.
Glazioviana, *Baker, in Mart. Fl. Bras.* vi. II. 41.—Bras.
glomerata, Sch. Bip. in Bot. Zeit. iii. (1845) 155 = Vanillosmopsis erythropappa.
glomerulifera, Baker, in Mart. Fl. Bras. vi. II. 123 = seq.
glomeruliflora, Walp. in Linnaea, xiv. (1840) 311 = Piptocarpha macropoda.
glutinosa, *DC. Prod.* v. 18.—Madag.
gnaphaliifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 34.—Cuba.
gnaphalioides, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 78.—Bras.
gochnatioides, *Hook. & Arn. ex DC. Prod.* vii. 264.—Reg. Argent.
Gomphrena, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 253.—China.
gracilis, *H. B. & K. Nov. Gen. et Sp.* iv. 34.—Am. trop.
gracilis, Poir. Encyc. Suppl. v. 465 = Liatris gracilis.
graminifolia, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 421.—Bras.
grandiflora, *Less. in Linnaea*, vi. (1831) 660.—Bras.
grandifolia, DC. Prod. v. 39 = serrata.
grandis, Boj. ex DC. l. c. 68 (= *Decaneurum grande*).—Madag.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 92.—Afr. trop.
gratiosa, *Hance, in Journ. Bot.* xx. (1882) 290.—China.
Greggii, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 204.—Mexic.
grisea, *Baker, in Mart. Fl. Bras.* vi. II. 61.—Bras.
Grisebachii, *Sch. Bip. in Journ. Bot.* i. (1863) 231.—Cuba.
guineensis, *Benth. in Hook. Niger Fl.* 427.—Afr. trop.
Haenkeana, *DC. Prod.* v. 37.—Peruv.
hakeaefolia, Less. in Linnaea, iv. (1829) 249 = Lychnophora hakeaefolia.
halimifolia, Mart. ex DC. Prod. v. 150 = Eupatorium halimifolium.
Halimodendron, Mart. ex DC. l. c. 85 = Eremanthus pycnocephalus.
havanensis, *DC. l. c.* 37.—Cuba.
hebeclada, DC. l. c. 36 = Westiniana.
hecatantha, DC. l. c. 53 = coriacea.
Helferi, *Hook. f. Fl. Brit. Ind.* iii. 231.—Burma.
helophila, *Mart. ex DC. Prod.* v. 50.—Bras.
heptantha, *DC. l. c.* 42.—Peruv.
heterophylla, Poir. Encyc. Suppl. v. 465 = Liatris scariosa.
hexantha, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 26.—Bras.
hieracioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1862) 511.—Cuba.
Hilairiana, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 113 = petiolaris.
Hildebrandtii, Vatke, in Oestr. Bot. Zeitschr. xxv. (1875) 323 = Gongrothamnus Hildebrandtii.
hirsuta, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. austr.
hirsutiflora, Poir. Encyc. viii. 502 = Liatris pycnostachya ?
hirtiflora, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 70.—Bras.
Hochstetteri, *Sch. Bip. ex Hochst. in Flora*, xxiv. (1841) I. *Intell.* 26.—Abyss.
holosericea, *Mart. ex DC. Prod.* v. 43.—Bras.
Hookeriana, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 346.—Zeylan.
hoveaefolia, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 423.—Bras.
hymenolepis, *A. Rich. Tent. Fl. Abyss.* i. 378.—Afr. trop.
hymenolepis, Vatke, in Linnaea, xxxix. (1875) 477 = filigera.
Hypochaeris, *DC. Prod.* v. 45.—Bras.
hypoleuca, *DC. l. c.* 27.—Mexic.
icosantha, DC. l. c. 49 = arborescens.
ignobilis, *Less. in Linnaea*, vi. (1831) 658.—Bras.
ilicifolia, Boj. ex Benth. & Hook. f. Gen. ii. 333 = Iphiona ilicifolia.

VERNONIA:—
imbricata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 209 = Piptolepis imbricata.
immunis, *Griseb. in Goett. Abh.* xxiv. (1879) 163.—Reg. Argent.
imponens, Mart. ex DC. Prod. v. 46 = sericea.
inaequiserrata, *Sch. Bip. in Journ. Bot.* i. (1863) 232.—Cuba.
incana, DC. Prod. v. 82, sphalm. = Eremanthus incanus.
incana, *Less. in Linnaea*, iv. (1829) 277.—Bras.
incanescens, Mart. ex DC. Prod. v. 82 = Vanillosmopsis capitata.
inconspicua, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 65 = helophila.
indica, Steud. Nom. ed. II. ii. 753, sphalm. = Veronica Anagallis.
indica, *Wall. ex C. B. Clarke, Comp. Ind.* 16.—Ind. or.
infundibularis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 285.—Afr. trop.
insignis, *Oliver & Hiern, l. c.* 292.—Afr. trop.
interjecta, *Baker, in Mart. Fl. Bras.* vi. II. 28.—Bras.
intermedia, *DC. Prod.* v. 27.—Bras.
inulaefolia, Baker, in Journ. Linn. Soc. xx. (1883) 180 = Bakeri.
inulaefolia, *Steud. Nom.* ed. II. ii. 753; *et ex A. Rich. Tent. Fl. Abyss.* i. 375.—Abyss.
inuloides, *DC. Prod.* v. 62.—Mexic.
involucrata, Less. in Linnaea, iv. (1829) 246 = pedunculata.
iodopappa, Sch. Bip. in Bot. Zeit. iii. (1845) 155 = crotonoides.
isotrichia, *DC. Prod.* v. 18.—Bras.
isotrichia, Mart. in Flora, xx. (1837) II. Beibl. 126 = Piptocarpha umbellulata.
jaceaeoides, A. Rich. Tent. Fl. Abyss. i. 376 = purpurea.
Jamesii, *Torr. & Gray, Fl. N. Am.* ii. 58.—N. Mexic.
javanica, DC. Prod. v. 22 = arborea.
jugalis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 270.—Afr. trop.
karaguensis, *Oliver & Hiern, in Trans. Linn. Soc.* xxix. (1873) 91.—Afr. trop.
Karstenii, *Sch. Bip. in Linnaea*, xxx. (1859–60) 169.—N. Granat.
Karvinskiana, *DC. Prod.* v. 62.—Mexic.
Karwinskiana, Hort. ex Kunth, Ind. Sem. Hort. Berol. (1848) 14 = Verbesina sericea.
Kingii, C. B. Clarke, Comp. Ind. 12 = clivorum.
Kirkii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 274.—Afr. trop.
Koestlini, Hochst. in Flora, xxiv. (1841) I. Intell. 26 = Hochstetteri.
Kotschyana, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. trop.
Kotschyi, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 287 = praec. ?
Kraussii, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. austr. et trop.
Kroneana, *Miq. in Journ. Bot. Néerl.* i. (1861) 98.—China.
Kurzii, C. B. Clarke, Comp. Ind. 24 = solanifolia.
lacunosa, *Mart. ex DC. Prod.* v. 56.—Bras.
laevigata, *Mart. ex DC. l. c.*—Bras.
lanceolaris, *DC. l. c.* 37.—Mexic.
lanuginosa, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 219 = scorpioides.
lappoides, *Baker, in Mart. Fl. Bras.* vi. II. 35.—Bras.
latifolia, Lem. Illustr. Hortic. ii. (1855) Misc. 39 = argyrophylla.
laurifolia, *DC. Prod.* v. 30.—Bras.
laxa, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 214.—Bras.
laxiflora, Less. in Linnaea, vi. (1831) 646 = cinerea.
Lechleri, *Sch. Bip. in Bonplandia*, iii. (1855) 236.—Peruv.
Leiboldiana, *Schlecht. in Linnaea*, xix. (1847) 742.—Mexic.
leiocarpa, *DC. Prod.* v. 34.—Mexic.
Leopoldi, *Vatke, in Linnaea*, xxxix. (1875) 478.—Afr. trop.
lepidota, *Griseb. Cat. Pl. Cub.* 145.—Cuba.
leprosa, Less. in Linnaea, iv. (1829) 252 = Piptocarpha leprosa.

VERNONIA:—
leptoclada, *Sch. Bip. in Journ. Bot.* i. (1863) 233.—Cuba.
leptophylla, DC. in Wight, Contrib. 6 = cinerea.
leptospermoides, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 143 = Piptolepis ericoides.
leptospermoides, Mart. ex DC. Prod. v. 17 = Piptolepis leptospermoides.
Lessingiana, Mart. ex DC. l. c. 60 = barbata.
lessingioides, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 92.—Bras.
Lettermanni, *Engelm. ex A. Gray, in Proc. Am. Acad.* xvi. (1881) 78.—Am. bor. occ.
leucophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 176.—Madag.
liatroides, *DC. Prod.* v. 34.—Mexic.
liatroides, A. Gray, in Proc. Am. Acad. v. (1861) 181, (excl. syn.) = Ervendbergii.
ligulaefolia, *Mart. ex DC. Prod.* v. 45.—Bras.
lilacina, *Mart. ex DC. l. c.* 48.—Bras.
linariaefolia, DC. l. c. 28 = psilophylla.
linarifolia, DC. l. c. 57, sphalm. = linearifolia.
Lindbergii, *Baker, in Mart. Fl. Bras.* vi. II. 105.—Bras.
Lindheimeri, *Gray & Engelm. in Proc. Am. Acad.* i. (1848) 46.—Texas.
linearifolia, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 237 = intermedia.
linearifolia, *Less. in Linnaea*, iv. (1829) 287.—Bras.
linearis, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 237 = rubricaulis.
linearis, *Spreng. Syst.* iii. 437.—Bras.
linifolia, Blume, Bijdr. 893 = cinerea.
lithospermoides, *Baker, in Mart. Fl. Bras.* vi. II. 66.—Bras.
Livingstoniana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 295.—Afr. trop.
Lobbii, *Hook. f. Fl. Brit. Ind.* iii. 231.—Burma.
lobelioides, Wall. Cat. n. 2927 = Ainsliaea pteropoda.
longeracemosa, Mart. ex DC. Prod. v. 42 = scorpioides.
longicaulis, DC. l. c. 33 = saligna.
longifolia, *Pers. Syn.* ii. 404.—Ind. occ.
lucida, *Less. in Linnaea*, iv. (1829) 260.—Bras.
Lundiana, Less. l. c. vi. (1831) 631 = Piptocarpha Lundiana.
Luschnathiana, Walp. l. c. xiv. (1840) 312 = Piptocarpha Lundiana.
Lyallii, *Baker, in Journ. Linn. Soc.* xx. (1883) 174.—Madag.
macranthele, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 83 = elegans.
macrocephala, *Less. in Linnaea*, iv. (1829) 298.—Bras.
macrocephala, A. Rich. Tent. Fl. Abyss. i. 377 = adoënsis.
macrophylla, Boj. ex DC. Prod. v. 27 = appendiculata.
macrophylla, *Less. in Linnaea*, vi. (1831) 668.—Bras.
macropoda, DC. Prod. v. 19 = Piptocarpha macropoda.
madagascariensis, *Less. in Linnaea*, vi. (1831) 644.—Madag.
malabarica, *Hook. f. Fl. Brit. Ind.* iii. 230.—Ind. or.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 534, nomen.—Bolivia.
Mannii, Hook. f. in Journ. Linn. Soc. vii. (1864) 198 = Inula Mannii.
Mansoana, *Baker, in Mart. Fl. Bras.* vi. II. 84.—Bras.
marginata, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 278.—Afr. trop.
marginata, Rafin. Atl. Journ. 146 (Quid ?).—Am. bor.
Mariana, *Mart. ex Baker, in Mart. Fl. Bras.* vi. II. 107.—Bras.
Marrubiastrum, Mart. ex Baker, l. c. 77 = lacunosa.
Martiana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 210 = Piptolepis Martiana.
megapotamica, *Spreng. Syst.* iii. 437.—Bras.
melanopappa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 28 = megapotamica.
Melleri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 282.—Afr. trop.
membranacea, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 217 = ruficoma.
menthaefolia, *Less. in Linnaea*, iv. (1829) 268.—Cuba.
Merana, *Baker, in Journ. Bot.* xx. (1882) 139.—Madag.
mespilifolia, *Less. in Linnaea*, vi. (1831) 641.—Afr. austr.

VERNONIA:—
Metziana, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 234 = divergens.
mexicana, *Less. in Linnaea*, vi. (1831) 680.—Mexic.
micans, Benth. Pl. Hartw. 196 = Piptocarpha elaeagnoides.
micradenia, *DC. Prod.* v. 38.—Am. austr.
micrantha, H. B. & K. Nov. Gen. et Sp. iv. 37 = canescens.
microcephala, *DC. Prod.* v. 39.—Am. trop.
microdonta, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 76 = glabrata.
Miersiana, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 115 = Salzmanni.
misera, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 278.—Afr. trop.
missionis, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 422.—Bras.
mollis, *H. B. & K. Nov. Gen. et Sp.* iv. 36.—Am. austr.
mollissima, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 237.—Reg. Argent.
moluccensis, *Miq. Fl. Ind. Bat.* ii. 19.—Ins. Molucc.
monocephala, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 418.—Bras.
monocephala, *Harv. in Harv. & Sond. Fl. Cap.* iii. 53.—Afr. austr.
Monosis, Benth. ex C. B. Clarke, Comp. Ind. 24 = arborea.
Monosis, *Sch. Bip. in Linnaea*, xx. (1847) 507.—Mexic.
montana, Hook. f. Fl. Brit. Ind. iii. 234, sphalm. = cinerea.
montevidensis, *Nees, ex Otto & Dietr. Allg. Gartenz.* i. (1833) 229.—Reg. Argent.
monticola, *Mart. ex DC. Prod.* v. 18.—Bras.
moquinioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 177.—Madag.
moquinioides, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 116 = mucronulata.
Moramballae, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 278.—Afr. trop.
Moritziana, *Sch. Bip. in Linnaea*, xx. (1847) 511.—Venezuela.
mossambiquensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 292.—Afr. trop.
mucronifolia, *DC. Prod.* v. 55.—Bras.
mucronulata, *Less. in Linnaea*, iv. (1829) 266.—Bras.
multiflora, Less. l. c. vi. (1831) 642 = divergens?
multiflosculosa, Mart. ex DC. Prod. v. 50 = macrophylla.
muricata, *DC. l. c.* 55.—Bras.
myriantha, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 198.—Afr. trop.
myriocephala, *DC. Prod.* v. 40.—Peruv.?
myriocephala, *A. Rich. Tent. Fl. Abyss.* i. 374.—Abyss.
natalensis, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. austr.
neilgherryensis, Thw. Enum. Pl. Zeyl. 160 = setigera.
nemoralis, *Thw. l. c.* 161.—Zeylan.
neriifolia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 217 = ruficoma.
nigrescens, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 28 = megapotamica.
nigritiana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 288.—Afr. trop.
nilgherryensis, DC. Prod. v. 32 = divergens.
nitens, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 221.—Bras.
nitens, Steud. Nom. ed. I. 875 = arborescens.
nitidula, *Less. in Linnaea*, iv. (1829) 260.—Bras.
notata, Less. l. c. 256 = Piptocarpha notata.
noveboracensis, *Willd. Sp. Pl.* iii. 1632.—Am. bor.
nudicaulis, *Less. in Linnaea*, vi. (1831) 637.—Madag.
nudiflora, *Less. l. c.* iv. (1829) 258.—Bras.
Nyassae, *Oliver, in Hook. Ic. Pl.* t. 1349.—Afr. trop.
oaxacana, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 74.—Mexic.
obconica, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 286.—Afr. trop.
oblonga, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 211 = Piptocarpha oblonga.
oblonga, Sch. Bip. in Pollichia, xx-xxi. (1863) 416 = Piptocarpha axillaris.
oblongifolia, Pohl, ex Mart. in Flora, xxiv. (1841) II. Beibl. 107 = Riedelii, geminata.
obovata, *Less. in Linnaea*, iv. (1829) 279.—Bras.
obscura, *Less. l. c.* 296.—Bras.
obtusata, *Less. in Linnaea*, vi. (1831) 662.—Bras.
obtusifolia, *Less. l. c.* iv. (1829) 308.—Bras.
ochroleuca, *Baker, in Journ. Linn. Soc.* xx. (1883) 179.—Madag.
octangularis, Meyen, Reise, i. 444 = Baccharis Meyeniana.
octantha, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 87.—Bras.
odoratissima, H. B. & K. Nov. Gen. et Sp. iv. 41 = scabra.
Oleaster, DC. Prod. v. 17 = Piptolepis Oleaster.
Oleaster, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 210 = Piptolepis Gardneri.
oligactoides, *Less. in Linnaea*, vi. (1831) 648.—Bras.
oligocephala, Edgew. in Trans. Linn. Soc. xx. (1846) 63 = revoluta.
oligocephala, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 420 = aurea.
oligocephala, Sch. Bip. ex Walp. Rep. ii. 947 = Kraussii.
oligolepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 56.—Bras.
oligophylla, *Michx. Fl. Bor. Am.* ii. 94.—Am. bor.
onopordioides, *Baker, in Mart. Fl. Bras.* vi. II. 36.—Bras.
opaca, Benth. in Hook. Journ. Bot. ii. (1840) 39 = Piptocarpha opaca.
oppositifolia, *Less. in Linnaea*, iv. (1829) 273.—Bras.
Ottonis, *Sch. Bip. l. c.* xx. (1847) 508.—Cuba.
ovalifolia, *Torr. & Gray, Fl. N. Am.* ii. 59 = noveboracensis.
ovata, *Less. in Linnaea*, iv. (1829) 294.—Bras.
oxylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 70.—Bras.
oxyphylla, DC. Prod. v. 19 = Piptocarpha oxyphylla.
pachyclada, *Baker, in Journ. Linn. Soc.* xx. (1883) 178.—Madag.
pacchensis, *Benth. Pl. Hartw.* 134.—Peruv.
pallens, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 161.—Mexic.
paludosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 113.—Bras.
palustris, Gardn. l. c. v. (1846) 215 = serrata.
pandurata, *Hort. Vindob. ex Link, Enum. Hort. Berol.* ii. 296.—Afr. trop.
paniculata, *DC. Prod.* v. 23.—Mexic.
paradoxa, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 432, in syn.—Venezuela.
Parishii, *Hook. f. Fl. Brit. Ind.* iii. 240.—Burma.
parviflora, *Klatt, in Linnaea*, xxxvii. (1871–73) 508.—Madag.
parviflora, Reinw. ex Blume, Bijdr. 893 = cinerea.
patens, *H. B. & K. Nov. Gen. et Sp.* iv. 41.—Am. austr.
patens, Less. in Linnaea, iv. (1829) 276, partim = polyanthes.
patula, Mart. ex DC. Prod. v. 55 = remotiflora.
pauciflora, Boj. ex DC. Prod. v. 18 = glutinosa.
pauciflora, *Less. in Linnaea*, iv. (1829) 292.—Afr. trop.
pauciflora, Poir. Encyc. Suppl. v. 465 = Liatris pauciflora.
Paulina, Mart. ex Steud. Nom. ed. II. ii. 754 = psilophylla.
pectinata, *DC. Prod.* v. 31.—Ind. or.
pectiniformis, *DC. in Wight, Contrib.* 6.—Ind. or.
pectoralis, *Baker, in Journ. Bot.* xx. (1882) 139.—Madag.
pedunculata, *DC. Prod.* v. 16.—Bras.
peguensis, *C. B. Clarke, Comp. Ind.* 13.—Burma.
pellita, *H. B. & K. Nov. Gen. et Sp.* iv. 36.—Venezuela.
pellucida, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 128 = Piptocarpha pellucida.
peninsularis, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 233.—Ind. or.
pentacantha, DC. Prod. v. 30 = seq.
pentaconta, *DC. l. c.* 696.—Bras.
Perrottetii, *Sch. Bip. ex Walp. Rep.* ii. 947.—Afr. trop.
persicifolia, Desf. Cat. Hort. Par. ed. III. 400 = sericea.
Petersii, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 90.—Afr. trop.
petiolaris, *DC. Prod.* v. 37.—Bras.
Petitiana, A. Rich. Tent. Fl. Abyss. i. 373 = pauciflora.
philippinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 312.—Ins. Philipp.
phylicaeformis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 252.—Peruv.
physalifolia, DC. in Wight, Contrib. 6 = cinerea.
Pinaster, Less. in Linnaea, iv. (1829) 249 = Lychnophora ericoides.
pinguis, *Griseb. in Goett. Abh.* xxiv. (1879) 165.—Reg. Argent.
pinifolia, *Less. in Linnaea*, iv. (1829) 257.—Afr. austr.
piptocarphoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 177.—Madag.
plantaginifolia, Less. in Linnaea, iv. (1829) 251 = Eremanthus plantaginifolius.
plantaginoides, DC. Prod. v. 27, in syn. = squarrosa.
platensis, *Less. in Linnaea*, iv. (1829) 312.—Bras.
platycephala, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 212 = Albertinia brasiliensis.
plumbaginifolia, *Fenzl, in Flora*, xxvii. (1844) 312.—Afr. trop.
podocoma, *Sch. Bip. ex Schweinf. & Aschers. in Beitr. Fl. Aethiop.* 287, nomen; *et Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 296.—Afr. trop.
Poeppigiana, DC. Prod. v. 20 = Piptocarpha Poeppigiana.
Poeppigiana, DC. l. c. 55 = Salzmanni.
Pohliana, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 113 = cuneifolia.
polyantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.
polyanthes, *Less. in Linnaea*, vi. (1831) 651.—Bras.
polycephala, DC. Prod. v. 39 = ferruginea.
polygalaefolia, *Less. in Linnaea*, vi. (1831) 628.—Madag.
polygalaefolia, Licht. ex Less. Syn. Comp. 200 = Iphiona polygalifolia.
polymorpha, Vatke, in Linnaea, xxxix. (1875) 476 = abyssinica, tigrensis, adoënsis, mossambiquensis.
polyphylla, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 63.—Bras.
polytricholepis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 415.—Madag.
populifolia, *Spreng. Syst.* iii. 434.—Ins. Maurit.
Poskeana, *Vatke & Hildeb. in Oestr. Bot. Zeitschr.* xxv. (1875) 324.—Afr. trop.
praealta, Michx. Fl. Bor. Am. ii. 95, partim; et DC. Prod. v. 63, partim = altissima.
praealta, Willd. Sp. Pl. iii. 1633 = noveboracensis.
procumbens, Boj. ex DC. Prod. v. 18 = polygalaefolia.
profuga, *De Not. Ind. Sem. Hort. Genuen.* (1840).—*Cf. Linnaea*, xv. (1841) *Litt.* 92.—Amer.?
prolifera, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 408.—Ins. Timor.
proteaeformis, Less. in Linnaea, iv. (1829) 249 = Lychnophora ericoides.
Proteopsis, DC. Prod. v. 16 = Proteopsis argentea.
Pseudomyrtus, A. St. Hil. Voy. Distr. Diam. i. 94, 367 = Piptolepis buxoides.
psilophylla, *DC. Prod.* v. 28.—Bras.
psilostachya, *DC. l. c.* 43.—Bras.
psittacorum, DC. l. c. 36 = polyanthes.
pterocaulon, *Baker, in Journ. Bot.* xvi. (1878) 77.—Reg. Argent.
pteropoda, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 283.—Afr. trop.
puberula, *Less. in Linnaea*, vi. (1831) 649.—Bras.
pulverulenta, *Baker, in Mart. Fl. Bras.* vi. II. 42.—Bras.
pumila, *Kotschy & Peyr. Pl. Tinn.* 37. t. 17 A.—Afr. trop.
punctata, *Sw. ex Wikstr. in Vet. Acad. Handl. Stockh.* (1827) 72 = longifolia.
puncticulata, DC. Prod. vii. 264 = pectiniformis.
punduana, DC. l. c. v. 32 = volkameriaefolia.
pungens, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1847) 418.—Bras.
purpurascens, *Sch. Bip. ex Walp. Rep.* ii. 945.—Mexic.
purpurea, *Sch. Bip. ex Walp. l. c.* 946.—Afr. trop.
pycnantha, *Benth. Pl. Hartw.* 134.—Peruv.
pycnocephala, Mart. ex DC. Prod. v. 85 = Eremanthus pycnocephalus.
pycnostachya, *DC. l. c.* 58.—Bras.
pyrifolia, DC. l. c. 19 = Piptocarpha pyrifolia.
pyrifolia, *D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 238.—Am. austr.
quadriflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 173.—Madag.

VERNONIA:—
Quartiniana, *A. Rich. Tent. Fl. Abyss.* i. 379.—Abyss.
quercifolia, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 194 = senegalensis.
quindecimflora, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 535, nomen.—Bolivia.
quinqueflora, *Less. l. c.* vi. (1831) 656.—Bras.
racemosa, *Delponte, in Mem. Acc. Torin.* Ser. II. xiv. (1854) 396.—Ins. S. Doming.
Radula, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 428 = carduoides.
Radula, *Mart. ex DC. Prod.* v. 52.—Bras.
ramiflora, Less. in Linnaea, iv. (1829) 255 = Piptocarpha Sellowii.
reflexa, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 223.—Bras.
Reinwardtiana, *De Vriese & Miq. ex Miq. Fl. Ind. Bat.* ii. 15.—Ins. Celebes.
remotiflora, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 112 (err. typ. 105).—Bras.
resinosa, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 419 = fruticulosa.
reticulata, Gardn. l. c. v. (1846) 226 = bardanoides.
reticulata, Willd. ex Less. in Linnaea, v. (1829) 284 = canescens.
revoluta, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1824) 218.—Reg. Himal.
rhaponticoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 180.—Madag.
Rheedii, *Kostel. Allg. Med. Pharm. Fl.* ii. 641 (Quid?). —Ind. or.
rhodolepis, *Baker, in Journ. Bot.* xx. (1882) 139.—Madag.
rhomboidea, Edgew. in Trans. Linn. Soc. xx. (1846) 62 = cinerea.
Riedeliana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 213 = Stilpnopappus bicolor.
Riedelii, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 64.—Bras.
rigida, *Sw. Fl. Ind. Occ.* iii 1322.—Ind. occ.
rigida, Wall. Cat. n. 2925 = teres.
rigiophylla, DC. Prod. v. 15 = teres.
rigiophylla, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 118.—Bras.
riparia, Hook. f. Fl. Brit. Ind. iii. 230, sphalm. = teres.
rosea, *Mart. ex DC. Prod.* v. 59.—Bras.
rosmarinifolia, *Less. in Linnaea*, iv. (1829) 286.—Bras.
Rothii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 293.—Afr. trop.
rotundifolia, Less. in Linnaea, iv. (1829) 254 = Piptocarpha rotundifolia.
Roxburghii, *Less. l. c.* vi. (1831) 674.—Reg. Himal.
ruboides, *Less. l. c.* iv. (1829) 290.—Bras.
rubricaulis, *Humb. & Bonpl. Pl. Aequin.* ii. 66. t. 99. —Am. trop.
rubricaulis, Less. in Linnaea, iv. (1829) 299, partim = squarrosa.
rubriramea, *Mart. ex DC. Prod.* v. 38 —Bras.
Rueppellii, *Sch. Bip. ex Walp. Rep.* ii. 946.—Abyss.
ruficoma, *Schlecht. ex Mart. in Flora*, xxiv. (1841) II. *Beibl.* 110, nomen; *et ex Baker, in Mart. Fl. Bras.* vi. II. 105.—Bras.
rufogrisea, *A. St. Hil. Voy. Distr. Diam.* ii. 306, 453. —Bras.
Rugeliana, Shuttl. ex A. Gray, Pl. Wright. i. 82 = noveboracensis.
rugosa, *Blume, Bijdr.* 894.—Java.
rugulosa, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 83.—Bras.
rupestris, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 114.—Bras.
Sagraeana, DC. Prod. v. 55 = rigida.
salicifolia, Gill. ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 237 = rubricaulis.
salicifolia, Less. in Linnaea, iv. (1829) 249 = Lychnophora salicifolia.
salicifolia, *Sch. Bip. l. c.* xx. (1847) 507.—Mexic.
saligna, *DC. Prod.* v. 33.—Reg. Himal.; Burma.
salvifolia, *Wight, Ic.* t. 1079.—Ind. or.
Salvinae, *Hemsl. Biol. Centr. Am. Bot.* ii. 73.—Guatemala
Salzmanni, *DC. Prod.* v. 55.—Bras.
Sancti-Antonii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 66 = adamantium.
Sarmentiana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 221 = arenaria.
saxicola, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 81.—Bras.

VERNONIA:—
scaberrima, Nutt. Gen. Am. ii. 134 = angustifolia.
scabra, *Pers. Syn.* ii. 404.—Bras.
scandens, *DC. Prod.* v. 32.—Reg. Himal.; Burma.
scandens, Less. ex Baker, in Mart. Fl. Bras. vi. II. 124 = Piptocarpha Riedelii.
scapigera, *Baker, l. c.* 55.—Bras.
scapigera, Less. in Linnaea, iv. (1829) 250 = Eremanthus scapigerus.
scariosa, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 346. —Zeylan.
scariosa, *Baker, in Journ. Bot.* xx. (1882) 169.—Madag.
scariosa, Poir. Encyc. viii. 502 = Liatris scariosa.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 204.—Mexic.
Schiedeana, *Less. in Linnaea*, vi. (1831) 399, 663.—Mexic.
Schimperi, *DC. Prod.* vii. 264.—Arab.
Schimperi, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) I. Intell. 26 = inulaefolia.
Schlechtendaliana, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 41 = ligulaefolia.
Schomburgkiana, Sch. Bip. in Linnaea, xx. (1847) 509 = ehretifolia.
Schultzii, *Karst. ex Sch. Bip. l. c.* xxx. (1859–60) 168. —N. Granat.
Schweinfurthii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 285.—Afr. trop.
Schweinitzii, Steud. Nom. ed. II. ii. 755 = fasciculata.
schwenkiaefolia, *Mart. ex DC. Prod.* v. 44.—Bras.
sclareaefolia, *Sch. Bip. in Linnaea*, xxx. (1859–60) 170.—N. Granat.
scorpioides, *Pers. Syn.* ii. 404.—Am. trop.
sechellensis, *Baker, Fl. Maurit.* 162.—Ins. Seychelles.
secunda, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 93.—Bras.
secundifolia, *Boj. ex DC. Prod.* v. 22.—Madag.
Seemanniana, *Steetz, in Seem. Bot. Voy. Herald*, 139. —Panama.
Sellowii, *Less. in Linnaea*, iv. (1829) 301.—Bras.
senecionea, Mart. ex DC. Prod. v. 54 = cognata.
senegalensis, Desf. Cat. Hort. Par. ed. III. 400 = pauciflora.
senegalensis, *Less. in Linnaea*, iv. (1829) 265; vi. (1831) 641.—Afr. trop.
senescens, Mart. ex DC. Prod. v. 59 = barbata.
sericea, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 112 (err. typ. 105).—Bras.
serrata, *Less. in Linnaea*, iv. (1829) 275.—Bras.
serratuloides, *H. B. & K. Nov. Gen. et Sp.* iv. 33.—Mexic.
serrulata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 22 = puberula.
sessiliflora, Willd. ex Less. in Linnaea, iv. (1829) 309 = remotiflora.
sessilifolia, *Less. l. c.* vi. (1831) 659.—Bras.
setigera, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 345. —Zeylan.
shirensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 291.—Afr. trop.
simplex, *Less. in Linnaea*, iv. (1829) 280.—Bras.
Sinclairi, *Benth. Bot. Voy. Sulph.* 109.—Mexic.
sinulata, *Miq. Fl. Ind. Bat.* ii. 18.—Java.
Smithiana, *Less. in Linnaea*, vi. (1831) 638.—Afr. trop.
solanifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 486.—Burma; China.
solidaginifolia, Boj. ex DC. Prod. v. 30 = Gongrothamnus solidaginifolius.
somalensis, *Franch. in Revoil, Çomali*, 34.—Afr. trop.
sordida, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 107 = mariana.
Sorocabae, Sch. Bip. ex Baker, l. c. 58 = oligactoides.
sororia, DC. Prod. v. 40 = scorpioides.
sparsiflora, *Baker, in Journ. Linn. Soc.* xx. (1883) 172. —Madag.
spathulaefolia, Mart. ex DC. Prod. v. 83 = Eremanthus plantaginifolius.
spathulata, Hochst. ex Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 162 = cinerascens.
spathulata, Sch. Bip. ex Aschers. in Schweinf. l. c., in nota = atriplicifolia.
speciosa, Less in Linnaea, iv. (1829) 290 = Stilpnopappus patulus.

VERNONIA:—
sphaeroidea, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 283 = Baldwini.
Spixiana, *Mart. ex DC. Prod.* v. 53.—Bras.
splendens, Less. in Linnaea, iv. (1829) 252 = Piptocarpha lucida.
Sprengeliana, *Sch. Bip. in Journ. Bot.* i. (1863) 232. —Ins. S. Doming.
squamosa, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 419 = buddleiaefolia.
squamulosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 44.—Reg. Argent.
squarrosa, Baill. Hist. Pl. viii. 296 = Serratula Scordium.
squarrosa, Less. in Linnaea, vi. (1831) 627 = teres.
squarrosa, *Less. l. c.* 678.—Bras.
staavioides, Less. l. c. iv. (1829) 249 = Lychnophora staavioides.
Stachyanthus, Mart. ex DC. Prod. v. 84 = Eremanthus Martii.
staehelinoides, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 75 = linearis.
stahelinoides, *Harv. Thes. Cap.* ii. 36.—Afr. austr.
Steetziana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 273.—Afr. trop.
Steetzii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 297.—Mexic.
stellaris, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 23.—Mexic.
stenocephala, *Oliver, in Hook. Ic. Pl.* t. 1349.—Afr. trop.
stenophylla, *Less. in Linnaea*, vi. (1831) 667.—Ins. S. Doming.
stictophylla, *Wright, in Sauv. Fl. Cub.* 74.—Cuba.
stoechadifolia, Sch. Bip. ex Walp. Rep. vi. 98 = Perrottetii.
Stoechas, *Mart. ex Baker, in Mart. Fl. Bras.* vi. II. 49.—Bras.
streptoclada, *Baker, in Journ. Linn. Soc.* xxi. (1885) 416.—Madag.
stricta, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 219.—Bras.
suaveolens, *H. B. & K. Nov. Gen. et Sp.* iv. 38.—N. Granat.
subcordata, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 226 = helophila.
subrepanda, Pers. Syn. ii. 404 = scorpioides.
subsessilis, *DC. Prod.* v. 62.—Ind. or.
subsessilis, Walp. in Linnaea, xiv. (1840) 312 = dura?
subsimplex, *Miq. Fl. Ind. Bat. Suppl.* 535.—Sumatra.
subsquarrosa, DC. Prod. v. 41 = geminata.
subulata, *Baker, in Mart. Fl. Bras.* vi. II. 108.—Bras.
subverticillata, *Sch. Bip. ex Baker, l. c.* 99.—Bras.
Sutherlandi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 52. —Afr. austr.
syncephala, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 64.—Bras.
talaumifolia, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 25.—Reg. Himal.
tanalensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 271.—Madag.
tarchonanthifolia, *Sch. Bip. in Linnaea*, xx. (1847) 507.—Mexic.
Tenoreana, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 92.—Afr. trop.
teres, *Wall. Cat.* n. 2926.—Reg. Himal.; Burma.
tereticaulis, DC. Prod. v. 20 = Piptocarpha tereticaulis.
terniflora, *Less. in Linnaea*, vi. (1831) 633.—Madag.
Teysmanniana, *Miq. Fl. Ind. Bat.* ii. 14.—Java.
theophrastifolia, *Schweinf. ex Oliver & Hiern*, in *Oliver, Fl. Trop. Afr.* iii. 294.—Afr. trop.
Thomae, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 66.—Ins. S. Thomas.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iii. 232.—Burma.
Thomsoniana, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 91.—Afr. trop.
Thwaitesii, *C. B. Clarke, Comp. Ind.* 11.—Zeylan.
tigrensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 290.—Afr. trop.
toluccana, *DC. Prod.* v. 30.—Mexic.
tomentella, *Mart. ex DC. l. c.* 59.—Bras.
tomentosa, Ell. Sketch, ii. 288 = noveboracensis.
tomentosa, Sinning, in Nees & Sinning, Amoen. Hort. Bonn. ii. (1824) 10. t. 5 = arborescens.
tournefortioides, H. B. & K. Nov. Gen. et Sp. iv. 34 = scorpioides.

VERNONIA :—
tragiaefolia, *DC. Prod.* v. 60.—Bras.
travancorica, *Hook. f. Fl. Brit. Ind.* iii. 240.—Ind. or.
triantha, *S. Schau. in Linnaea*, xix. (1847) 714.—Mexic.
tricephala, *Gardn. in Hook. Lond. Journ. Bot.* v. (1846) 223.—Bras.
trichantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 416.—Madag.
trichocarpha, Spreng. Syst. iii. 437 = Lychnophora trichocarpha.
tricholepis, DC. Prod. v. 54 = remotiflora.
tricholepis, Griseb. Fl. Brit. W. Ind. 354 = araripensis.
trifloscculosa, *H. B. & K. Nov. Gen. et Sp.* iv. 40.—Mexic.
triplinervia, Spreng. Syst. iii. 435 = Baccharis triplinervis.
turbinata, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 90. t. 56. f. B.—Afr. trop.
Tweedieana, *Baker, in Mart. Fl. Bras.* vi. II. 99.—Reg. Argent.
umbellata, Less. in Linnaea, iv. (1829) 255 = Piptocarpha ramiflora.
uncinata, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 277.—Afr. trop.
undulata, *Oliver & Hiern, l. c.* 276.—Afr. trop.
undulata, Steud. Nom. ed. II. ii. 753, in syn., sphalm. = Veronica Anagallis.
uniflora, *Sch. Bip. in Linnaea*, xx. (1847) 506.—Mexic.
unionis, *Sch. Bip. ex Walp. Rep.* ii. 945.—Abyss.
urticaefolia, *A. Rich. Tent. Fl. Abyss.* i. 378.—Abyss.
vagans, DC. Prod. v. 32 = scandens.
Vahliana, Less. in Linnaea, iv. (1829) 306 = longifolia.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 33.—Cuba.
varroniaefolia, *DC. Prod.* v. 56.—Bras.
Vauthierana, DC. l. c. 21 = Piptocarpha Vauthieriana.
venosissima, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 30.—Bras.
vepretorum, *Mart. ex DC. Prod.* v. 59.—Bras.
verbascifolia, *Less. in Linnaea*, iv. (1829) 310.—Bras.
verbascoides, *Walp. l. c.* xiv. (1840) 313.—Bras.
verbenacea, DC. Prod. v. 35 = ignobilis.
Vernonella, *Harv. in Harv. & Sond. Fl. Cap.* iii. 53.—Afr. austr.
verpretorum, Steud. Nom. ed. II. ii. 755, sphalm. = vepretorum.
vestita, *Baker, in Mart. Fl. Bras.* vi. II. 83.—Bras.
vialis, *DC. Prod.* v. 25.—Madag.
villosissima, Less. in Linnaea, iv. (1829) 249 = Lychnophora villosissima.
vindex, Mart. ex DC. Prod. v. 160 = Eupatorium vindex.
violacea, *Oliver & Hiern, ex Oliver, in Trans. Linn. Soc.* xxix. (1873) 91.—Afr. trop.
virens, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 71.—Bras.
virgulata, *Mart. ex DC. Prod.* v. 42.—Bras.
viscidula, *Less. in Linnaea*, iv. (1829) 289.—Bras.
Vogeliana, *Benth. in Hook. Niger Fl.* 427.—Afr. trop.
Vogeliana, Thoms. in Speke, Journ. App. 637 = Thomsoniana.
volkameriaefolia, Bedd. Fl. Sylv. t. 225 = travancorica.
volkameriaefolia, *DC. Prod.* v. 32.—Reg. Himal.; Burma.
voluta, *Baker, in Journ. Linn. Soc.* xxi. (1885) 415.—Madag.
vulgaris, Hook. f. Fl. Brit. Ind. iii. 240, sphalm. = volkameriaefolia.
Walpersiana, Steud. Nom. ed. II. ii. 755 = dura ?
Warmingiana, *Baker, in Mart. Fl. Bras.* vi. II. 68.—Bras.
Westiniana, *Less. in Linnaea*, vi. (1831) 650.—Bras.
Wightiana, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 345.—Zeylan.
Wightiana, Wall. Cat. n. 2923 = Laggera alata.
Wrightii, *Sch. Bip. in Journ. Bot.* i. (1863) 234.—Cuba.
xanthophylla, *Mart. ex DC. Prod.* v. 58.—Bras.
zanzibarensis, *Less. in Linnaea*, vi. (1831) 637.—Afr. trop.

VERNONIA :—
zeylanica, *Less. l. c.* iv. (1829) 344.—Zeylan. ; Madag.
Zollingeriana, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 119 = cinerea.
zollingerianoides, *Sch. Bip. l. c.*, nomen.—Java ?
Zuccariniana, *Mart. ex DC. Prod.* v. 55.—Bras.

VERONICA, [Tourn.] Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 964.
AGERELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 129.
AIDELUS, Spreng. Syst. iv. Cur. Post. 17 (1827).
ALLOPLEIA, Rafin. Med. Fl. ii. 109 (1830).
AZURINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
BECCABUNGA, Fourr. l. c. (1869).
CALISTACHYA, Rafin. in Med. Repos. N. York, II. v. (1808) 60, 352.
CARDIA, Dulac, Fl. Hautes-Pyr. 387 (1867).
COCHLIDIOSPERMUM, Reichb. Consp. 121 (1828).
COERULINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
CYMBOPHYLLUM, F. Muell. in Hook. Kew Journ. viii. (1856) 202.
DERWENTIA, Rafin. Fl. Tellur. iv. 55 (1836).
DIPLOPHYLLUM, Lehm. in Ges. Naturf. Fr. Berl. Mag. viii. (1814) 310.
EUSTACHIA, Rafin. Cat. 14 (1824).
HEBE, Comm. ex Juss. Gen. 105 (1789).
HEDYSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
LEPTANDRA, Nutt. Gen. Am. i. 7 (1818).
LIMNASPIDIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
ODICARDIS, Rafin. Fl. Tellur. iv. 55 (1836).
OMPHALOSPORA, Bartl. Ord. 171 (1830).
PANOXIS, Rafin. Med. Fl. ii. 109 (1830).
PETRODORA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
POCILLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 129.
PONARIA, Rafin. Med. Fl. ii. 109 (1830).
PSEUDOLYSIMACHION, Opiz, Seznam, 80 (1852).
PYGMAEA, Hook. f. Handb. N. Zeal. Fl. 217 (1864).
URANOSTACHYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
VERONICASTRUM, Heist. ex Fabr. Hort. Helmst. 111 (1759).
VERONICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128.
abrotanifolia, Link, ex Steud. Nom. ed. II. ii. 755 (Quid ?).—Sibir.
abrotanifolia, Stokes, Bot. Comment. i. 58 = multifida.
abyssinica, *Fresen. in Bot. Zeit.* ii. (1844) 356.—Abyss.
acinifolia, *Linn. Sp. Pl.* ed. II. 19.—Europ.; As.
acinifolia, F. W. Schmidt, Fl. Boëm. i. 31 = arvensis.
acinifolia, Willd. Fl. Berol. Prod. 11 = praecox.
acuminata, Link, Jahrb. i. III. (1820) 34 = spicata.
acuta, Mart. Hort. Erlang. (1814) 12 = longifolia.
acutiflora, Benth. in DC. Prod. x. 460 = ligustrifolia.
acutiflora, *Lapeyr. ex Roem. & Schult. Syst.* i. 112; *Suppl. Hist. Pl. Pyr.* 7.—Mont. Pyren.
acutifolia, Gilib. Exercit. i. 119 = V. Anagallis.
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 208.—Afr. trop.
agrestis, Gouan, ex Nym. Consp. 548 = didyma.
agrestis, *Linn. Sp. Pl.* 13.—Europ.; As. et Afr. bor.
alba, [Mazziari,] in Ionios, Anthol. ii. (1834) 442 = agrestis.
alba, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = prostrata.
albanica, C. Koch, in Linnaea, xxii. (1849) 701 = amoena.
aleppica, *Boiss. Diagn.* Ser. II. iii. 169.—As. Min.; Assyr.
Allionii, F. W. Schmidt, Fl. Boëm. i. 6 = officinalis.
Allionii, *Vill. Prosp.* 20; *Hist. Pl. Dauph.* ii. 8.—Europ. austr.
alpestris, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 170 = serpyllifolia.
alpina, Georgi, Reise, i. 195 = ciliata.
alpina, *Linn. Sp. Pl.* 11.—Reg. bor. et arct.
alpiphila, Arv. Ess. Pl. Dauph. 55 (Quid ?).—Gallia.

VERONICA :—
altaica, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 19 = spuria.
alternifolia, Lej. Fl. Spa, ii. 286 = spicata.
ambigua, Lucé, Topog. Nachr. Oesel, 4 = V. Anagallis.
ambigua, Mart. Hort. Erlang. (1814) 13 = longifolia.
americana, *Schwein. ex Benth. in DC. Prod.* x. 468.—Am. bor.
amethystina, Willd. Enum. Hort. Berol. 17 = spuria.
amoena, *Bieb. Fl. Taur. Cauc.* i. 14.—Reg. Cauc.
amoena, Heldr. ex Nym. Consp. 548 = glauca.
amplexicaulis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 352.—N. Zel.
Anagallis, Bong. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 157 = americana.
Anagallis, *Linn. Sp. Pl.* 12 (Anagallis-aquatica).—Reg. bor. temp.
anagalloides, *Guss. Pl. Rar.* 5. t. 3.—Graecia; Oriens.
Anarrhinum, Cat. Hort. Dresd.; ex Reichb. Fl. Germ. Excurs. 371 = longifolia.
Andersonii ×, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 3.
angulata, Opiz, Natural. n. ix. (1825) 112, nomen = latifolia.
angustata, Opiz, l. c. 111, nomen = spicata.
angustifolia, Bernh. Ehrenp. 35 = austriaca.
angustifolia, A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 458 = ligustrifolia.
angustifolia, Fisch. ex Link, Enum. Hort. Berol. i. 19 = spuria.
angustifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 306 = scutellata.
angustifolia, A. Rich. Ess. Fl. N. Zél. 187 = parviflora.
angustifolia, Steud. Nom. ed. II. ii. 756 = tubiflora.
anisophylla, C. Koch, in Linnaea, xvii. (1843) 287 = orientalis.
anomala, *Armstr. in Trans. N. Z. Inst.* iv. (1872) 291.—N. Zel.
apennina, *Tausch, in Syll. Ratisb.* ii. (1828) 247.—Italia.
aphylla, *Linn. Sp. Pl.* 11.—Europ. austr.
aquatica, Benquerel, in Neuch. Bull. v. (1859–61) 449, nomen = V. Anagallis ?
aquatica, S. F. Gray, Nat. Arr. Brit. Pl. ii. 306 = V. Anagallis.
arborea, *F. Buch. in Trans. N. Z. Inst.* vi. (1874) 242.—N. Zel.
arbustula, Hassk. Allg. Regist. Flora, 864, sphalm. = seq.
arbustulosa, Hoffmgg. Preisverz. (1841) 85; et ex Reichb. Pl. Crit. x. 2 = fruticulosa.
arenaria, *A. Cunn. ex Benth. in DC. Prod.* x. 463.—Austral.
arguta, Benth. l. c. 478 = notabilis.
arguta, *R. Br. Prod.* 435.—Austral.
arguta, Moretti, in Syll. Ratisb. i. (1824) 103 = spicata.
arguta, Schrad. Comm. Veron. Spic. 22 = longifolia.
arguteserrata, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 626 = biloba.
aristata, Walp. Rep. iv. 797, sphalm. = Vernonia natalensis.
armena, *Boiss. & Huet, Diagn.* Ser. II. iii. 166.—Armenia.
Armstrongii, *T. Kirk, in Trans. N. Z. Inst.* xi. (1879) 464.—N. Zel.
arvensis, *Linn. Sp. Pl.* 13.—Europ.; As. et Afr. bor.
arvensis, Thunb. Fl. Jap. 20 = agrestis.
ascendens, *G. Don, ex Sweet, Hort. Brit.* ed. II. 397.—Hab. ?
aspera, Pritz. Ic. Ind. i. 1153, sphalm. = longifolia.
Assoana, *Willk. in Linnaea*, xxx. (1859–60) 120.—Hispan.
atlantica, Ball, in Journ. Bot. xiii. (1875) 174 = cuneifolia.
atlantica, Pers. Syn. i. 12 = rosea.
Aucheri, *Boiss. Diagn.* Ser. I. iv. 80.—Persia.
aurifolia, Dicks. ex Roem. & Usteri, Mag. iv. 10tes St. 190 (Quid ?).—Britan.
australis, Schrad. Comm. Veron. Spic. 24 = spicata.
austriaca, *Linn. Syst.* ed. X. 849.—Europ.; Reg. Cauc.
azurea, Link, Enum. Hort. Berol. i. 22 = longifolia.
Bachofenii, Heuff. in Flora, xviii. (1835) I. 253 = grandis.
Barrelieri, Schott, ex Roem. & Schult. Syst. i. 94 = spicata.

VERONICA:—
bartsiaefolia, Boiss. Fl. Orient. iv. 464 = biloba.
Bastardi, Bor. Fl. Centr. Fr. ed. III. ii. 487 = V. Teucrium.
Baumgartenii, *Roem. & Schult. Syst.* i. 100.—Banat.
Beccabunga, Aitch. in Journ. Linn. Soc. xix. (1882) 180 = V. Anagallis.
Beccabunga, Auct. Am. ex A. Gray, Syn. Fl. Am. ii. 1. 287 = americana.
Beccabunga, *Linn. Sp. Pl.* 12.—Reg. bor. temp.
Bellardi, All. Fl. Pedem. i. 77 = arvensis.
bellidioides, *Linn. Sp. Pl.* 11.—Europ. austr.
Benthami, *Hook. f. Fl. Antarct.* i. 60.—Ins. Dom. Auckland.
Benthami, C. Koch, ex Boiss. Fl. Orient. iv. 439 = peduncularis.
bibarbata, Stokes, Bot. Mat. Med. i. 31 = V. Chamaedrys.
Bidwillii, *Hook. Ic. Pl.* t. 814.—N. Zel.
Biebersteinii, Hassk. Allg. Regist. Flora, 865 = *Verbascum Biebersteinii*.
Biebsteinii, Richter, in Denkschr. Akad. Wien, l. (1885) 24 = multifida.
bihariensis, Kern. in Oestr. Bot. Zeitschr. xxxiii. (1873) 371 = austriaca.
Billardieri, Kotschy, ex Benth. in DC. Prod. x. 480 = Kotschyana.
Billardierii, Vahl, Enum. i. 70 = orientalis.
biloba, Bourg. ex Boiss. Fl. Orient. iv. 464 = campylopoda.
biloba, *Linn. Mant.* ii. 172.—Reg. Himal. et Cauc.; Sibir.
biserrata, Schur, Enum. Pl. Transs. 497 = grandis.
Bodardi, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = V. Cymbalaria.
bombycina, *Boiss. & Kotschy*, *Diagn.* Ser. II. iii. 171.—Syria.
Bonarota, Linn. Sp. Pl. 11 = Paederota Bonarota.
borealis, Laest. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 336 = serpyllifolia.
Brachtii, Opiz, Natural. n. ix. (1825) 110 = longifolia.
brachyphylla, Schult. Mant. i. 96 = spicata.
brachysepala, Sch. Bip. in Pollichia, xvi-xvii. (1859) 43 = V. Teucrium.
brachystyla, Rehm. in Spraw. Kom. Fis. Krakow, vii. (1873) 89 = V. Teucrium.
bracteata, Opiz, Natural. n. x. (1825) 244 = longifolia.
bracteata, Schult. Mant. i. 116 (Quid ?).—Hab. ?
bracteata, *Willd. ex Steud. Nom.* ed. II. ii. 756.—Hungar. ?
brevicaulis, Tausch, in Syll. Ratisb. ii. (1828) 247 = V. Teucrium.
brevifolia, Bieb. Fl. Taur. Cauc. i. 6 = spuria.
brevifolia, Link, Enum. Hort. Berol. i. 21 = spicata.
brevi-pedunculata, Gilib. Fl. Lituan. i. 119 = arvensis.
brevistyla, *Moris, in Mem. Acc. Torin.* Ser. II. ii. (1840) 301.—Ins. Sardoa.
Brownei, Roem. & Schult. Syst. i. 118 = arguta.
Brunoniana, Wall. Cat. n. 405 = Calorhabdos Brunoniana.
Buchanani, *Hook. f. Handb. N. Zeal. Fl.* 211.—N. Zel.
Bungabecca, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 241 = serpyllifolia.
Bungei, *Boiss. Fl. Orient.* iv. 463.—Persia.
Buxbaumi, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 187 = pectinata.
Buxbaumiana, Pall. ex Bieb. Fl. Taur. Cauc. i. 9 = gentianoides.
Buxbaumii, Tenore, Fl. Nap. i. 7. t. 1 = Tournefortii.
buxifolia, *Benth. in DC. Prod.* x. 462.—N. Zel.
caesaraugustiana, *Echeand. Fl. Caesaraug.* (1861) 21.—Hispan.
caespitosa, *Boiss. Diagn.* Ser. I. iv. 79.—Graecia; As. Min.; Syria.
callistachya, *Ledeb. ex Loud. Hort. Brit. Suppl.* iii. 555 (Quid ?).—Podolia.
Calverti, Boiss. Fl. Orient. iv. 450 = telephiifolia.
calycida, Fries, ex Reichb. Fl. Germ. Excurs. 365 = agrestis.
calycina, *R. Br. Prod.* 435.—Austral.
calycina, A. Cunn. in Bot. Mag. sub t. 3461 = plebeia.
campylopoda, *Boiss. Diagn.* Ser. I. iv. 80.—Oriens.
camtschatica, J. F. Gmel. Syst. 29 = kamtchatica.
cana, *Wall. Cat.* n. 401; *Benth. Scroph. Ind.* 45.—Reg. Himal.; Japan; Formosa.
candida, Lodd. Cat. (1823); ex Schult. Mant. i. Add. II. 228 (Quid ?).—Hab. ?

VERONICA:—
candidissima, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 932 = incana.
canescens, Bast. Fl. Maine-et-Loire, Suppl. 21 = V. Teucrium.
canescens, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 503.—N. Zel.
canescens, Presl, ex Schult. Mant. i. Add. II. 229, in syn. = prostrata.
canescens, Schrad. Comm. Veron. Spic. 19 = incana.
canterburiensis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 355.—N. Zel.
capensis, Burm. f. Fl. Cap. Prod. 1 = serpyllifolia.
capensis, *Fenzl, in Linnaea*, xvii. (1843) 332.—Afr. austr.
capitata, *Royle, ex Benth. Scroph. Ind.* 45.—Reg. Himal.
capitulata, Poir. Encyc. Suppl. v. 466 = spuria.
cardiocarpa, Walp. Rep. iii. 335 = biloba.
cariensis, Boiss. Diagn. Ser. I. iv. 75 = cuneifolia.
carnea, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 357.—N. Zel.
carnea, DC. in Mém. Soc. Phys. Genèv. i. (1822) 459 = longifolia.
carnea, Opiz, ex Schult. Mant. i. Add. II. 229, in obs. = prostrata.
carnea, Vitm. Summa Pl. i. 34 = alpina.
carnosula, Hook. f. Bot. Mag. t. 6587 = pinguifolia.
carnosula, *Hook. f. Handb. N. Zeal. Fl.* 210.—N. Zel.
carnosula, Lam. Illustr. i. 47 = peregrina.
caroliniana, Poir. Encyc. viii. 520 = Mitreola petiolata.
caroliniana, Walt. Fl. Carol. 61 = peregrina.
cartilaginea, Ledeb. Fl. Alt. i. 28, in adnot. = spuria.
catarractae, *Forst. f. Prod.* 3.—N. Zel.
caucasica, *Bieb. Fl. Taur. Cauc.* i. 13.—Reg. Cauc.
čechica, Opiz, in Lotos, iv. (1854) 105 (Quid ?).—Bohem.
ceratocarpa, *C. A. Mey. Verz. Pfl. Cauc.* 106.—Reg. Casp.
cespitosa, Boiss. Diagn. Ser. I. iv. 79 = caespitosa.
Chaixii, Lapeyr. Suppl. Hist. Abr. Pl. Pyr. 6 = V. Teucrium.
chamaedrya, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = V. Chamaedrys.
chamaedryoides, Bory & Chaub. Nouv. Fl. Pélop. i. t. 1 = V. Chamaedrys.
Chamaedrys, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 211 = melissaefolia.
Chamaedrys, *Linn. Sp. Pl.* 13.—Europ.; As. Min.; Syria.
chamaepithyoides, Desf. Tabl. ed. I. 50 = verna.
chamaepithyoides, Lam. Illustr. i. 47 = digitata.
Chamaepitys, *Griseb. Spicil. Fl. Rumel.* ii. 25.—Byzant.; As. Min.
Chamaepitys, Pers. Syn. i. 13 = verna.
Chamedris, Neck. Delic. Gallo-Belg. i. 9 = V. Chamaedrys.
chathamica, *T. Buch. in Trans. N. Z. Inst.* vii. (1875) 338.—N. Zel.
Chaubardi, *Boiss. & Reut. Diagn.* Ser. II. iii. 174.—Graecia.
Cheesmani, *Benth. in Hook. Ic. Pl.* t. 1366. f. A.—N. Zel.
chillensis, H. B. & K. Nov. Gen. et Sp. ii. 390 = peregrina.
cibiniensis, Schur, Enum. Pl. Transs. 498 = longifolia.
ciliaris, Hoffm. in Comm. Gotting. xv. (1804) 121. t. 5 = longifolia.
ciliata, *Fisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 56.—Reg. Himal.; As. bor.
ciliata, Schloss. & Vukot. Syll. Fl. Croat. 90 = acinifolia.
cinerea, *Boiss. & Bal. Diagn.* Ser. II. vi. 131.—As. Min.
cinerea, Rafin. New Fl. Am. iv. 39 (Quid ?).—Am. bor.
circaeoides, G. Don, ex Loud. Hort. Brit. 7 (Quid ?).—Helvet.
Clarkeana, Schult. Mant. i. Add. II. 193 = alpina.
Clusii, Schott, ex Roem. & Schult. Syst. i. 94 = spicata.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 209.—N. Zel.
collina, Opiz, Natural. n. ix. (1825) 108 = triphyllos.
colocensis, *Menyh. Kalocsa Vidék. Növényt.* 134.—Hungar.
commutata, Seidl, ex Bercht. Oek. Fl. Boehm. i. 31 = longifolia.

VERONICA:—
comosa, *Richter, in Denkschr. Akad. Wien*, l. (1885) 24.—Persia.
complicata, Hoffm. Phytog. Blaetter, i. (1803) 98 = longifolia.
condensata, Hort. ex Reichb. Fl. Germ. Excurs. 372 = longifolia.
conferta, Boiss. Fl. Orient. iv. 459 = glaberrima.
confusa, Roem. & Schult. Syst. i. 95 = longifolia.
coniosperma, Wallr. in Linnaea, xiv. (1840) 533 (Beitr. Fl. Hercyn. 163) = acinifolia.
connata, Rafin. Med. Fl. ii. 110 = scutellata.
cordata, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = prostrata.
corymbosa, Hort. ex G. Don, in Loud. Hort. Brit. 6, nomen (Quid ?).—Hab. ?
Crantzii, Schult. Obs. Bot. 3 = orchidea.
crassifolia, Kit. ex Roem. & Schult. Syst. i. 16 = orchidea.
crassifolia, Wierzb. ex Heuff. in Flora, xviii. (1835) 251 = spicata.
crassifolia, *Zeyh. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 23.—Hungar.
crenata, Lodd. Cat. (1823); ex Schult. Mant. i. Add. II. 228 (Quid ?).—Hab. ?
crenulata, Hoffm. Phytog. Blaetter, i. (1803) 95 = longifolia.
cretica, Willd. ex Link, Jahrb. i. III. (1820) 41 = thymifolia.
crinita, Kit. in Schult. Oestr. Fl. ed. II. i. 26 = V. Teucrium.
crispa, Hort. ex Reichb. Fl. Germ. Excurs. 368 = latifolia.
crispa, Lodd. Cat. (1823); ex Schult. Mant. i. Add. II. 228 (Quid ?).—Hab. ?
Crista-galli, *Stev. in Trans. Linn. Soc.* xi. (1815) 408. t. 31.—Reg. Cauc.
cristata, Bernh. Ehrenp. 14 = orchidea.
cuneata, Guss. Enum. Pl. Inar. 241. t. 3. f. 2 = V. Cymbalaria.
cuneifolia, *D. Don, in Fellows, Discov. in Lycia*, 291; *et in Ann. & Mag. Nat. Hist.* Ser. I. vii. (1841) 457.—As. Min.
cupressoides, *Hook. f. Handb. N. Zeal. Fl.* 212.—N. Zel.
Cusickii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 288.—Am. bor. occ.
cuspidata, Willd. ex Link, Jahrb. i. III. (1820) 36 = longifolia.
cycnorum, Miq. in Lehm. Pl. Preiss. i. 342 = calycina.
Cymbalaria, *Bodard, Mém. Véronique Cymbalaire*, 3; *ex Bertol. Amoen. Ital.* 56.—Reg. Mediterr.; As. Min.
cymbalariaefolia, Vahl, Enum. i. 81 = V. Cymbalaria.
cymbalarifolia, J. F. Gmel. Enum. Stirp. Tubing. 6 = hederaefolia.
cymbalarifolia, F. W. Schmidt, Fl. Boëm. i. 26 = Tournefortii.
cymbalarioides, *Blanche, ex Boiss. Fl. Orient.* iv. 468.—Syria; Mesopot.
Dabneyi, *Hochst. ex Seub. Fl. Azor.* 39. t. 8. f. 1.—Ins. Azor.
daurica, *Stev. in Mém. Soc. Nat. Mosc.* v. (1817) 339.—Dahuria.
decorosa, *F. Muell. in Linnaea*, xxv. (1852) 430.—Austral.
decumbens, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 352.—N. Zel.
decussata, *Moench, Verz.* 137.—Ins. Falkland.
decussata, [Soland. in] Ait. Hort. Kew. ed. I. i. 20 = elliptica.
deflectans, Krock. Fl. Siles. Suppl. i. 42 = V. Chamaedrys.
deltigera, *Wall. Cat.* n. 402; *Benth. Scroph. Ind.* 45.—Reg. Himal.; Afghan.
deltoidea, Spreng. Syst. iv. Cur. Post. 17 = plebeia.
densiflora, *Ledeb. Fl. Alt.* i. 34.—Sibir. altaic.
densifolia, *F. Muell. Fragm.* ii. 137.—Austral.
dentata, F. W. Schmidt, Fl. Boëm. i. 20 = V. Teucrium.
dentosa, Hort. ex Reichb. Fl. Germ. Excurs. 372 = longifolia.
depauperata, Waldst. & Kit. Pl. Rar. Hung. iii. 272 = aphylla.
depressa, *Kit. in Schult. Oestr. Fl.* ed. II. i. 32.—Hungar.
depressa, Schur, Enum. Pl. Transs. 495 = officinalis.
Derwentia, *Andr. Bot. Rep.* t. 531.—Austral.

VERONICA:—
dianthifolia, A. Cunn. in Loud. Hort. Brit. 467 = arenaria.
dichrus, Schott & Kotschy, in Oestr. Bot. Wochenbl. vii. (1857) 205 = cuneifolia.
didyma, *Tenore, Prod. Fl. Nap.* p. vi.—Europ.; Afr. bor.; Oriens.
Dieffenbachii, *Benth. in DC. Prod.* x. 459.—N. Zel.
diffusa, *Hook. f. in Hook. Ic. Pl.* t. 645.—N. Zel.
diffusa, Rafin. New Fl. Am. iv. 38 (Quid?).—Am. bor.
digitata, Lam. Fl. Fr. ii. 445 = triphyllos.
digitata, *Vahl, Symb. Bot.* i. 2.—Gall. austr.; Hispan.
Dillenii, Crantz, Stirp. Austr. ed. II. fasc. iv. 352 = verna.
diosmaefolia, Knowles & Westc. Flor. Cab. iii. (1839) 65. t. 106 = formosa.
diosmifolia, *R. Cunn. in Bot. Mag.* sub t. 3461.—N. Zel.
distans, *R. Br. Prod.* 435.—Austral.
divaricata, *Boiss. & Bal. Diagn.* Ser. II. iii. 173.—Cilicia.
divaricata, Tausch, in Flora, iv. (1821) 561 = V. Chamaedrys.
diversifolia, Pantoč. in Verh. Ver. Natur. Heilk. Presb. ii. (1871–72) 70 = prostrata.
diversifolia, Steud. Nom. ed. I. 876 = virginica.
Drummondi, Benth. in DC. Prod. x. 478 = distans.
Dryadis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 54 = officinalis.
Drymeja, Schur, Enum. Pl. Transs. 495 = pedunculatis.
dubia, Chaix, ex Lapeyr. Suppl. Hist. Pl. Pyr. 6 = V. Teucrium.
dubia, DC. Fl. Fr. iii. 462 = V. Chamaedrys.
elata, Host, Fl. Austr. i. 3 = longifolia.
elatior, Ehrh. ex Willd. Enum. Hort. Berol. 17 = longifolia.
elbursensis, Boiss. Diagn. Ser. I. xii. 46 = biloba.
elegans, DC. Cat. Hort. Monsp. (1813) 154 = spuria.
elliptica, *Forst. f. Prod.* 3.—N. Zel.; Reg. Magell.
elongata, Benth. in DC. Prod. x. 478 = plebeia.
epacridea, *Hook. f. Handb. N. Zeal. Fl.* 213.—N. Zel.
ericetorum, Opiz, Natural. n. ix. (1825) 111, nomen = V. Chamaedrys.
erinoides, Boiss. & Sprun. Diagn. Ser. I. iv. 78 = thessalica.
euphrasiaefolia, *Link, Jahrb.* i. III. (1820) 41.—Persia.
exaltata, *Maund, Bot. Gard.* iv. (1834–35) n. 543.—Sibir.
excelsa, Desf. Cat. Hort. Par. ed. III. 391 = longifolia.
exilis, Schott, ex Boiss. Fl. Orient. iv. 464 = biloba.
falcata, Mart. Hort. Erlang. (1814) 10 = longifolia.
fallacina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 42 = V. Cymbalaria.
filiformis, DC. Fl. Fr. Suppl. 388 = Tournefortii.
filiformis, *Sm. in Trans. Linn. Soc.* i. (1791) 195.—As. Min.
finaustrina, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 28 = Benthami.
fistulosa, Krock. Fl. Siles. Suppl. i. 20 = longifolia.
flexuosa, Host & Moretti, in Bibl. Ital. xii. (1818) 369 = longifolia.
floribunda, R. Br. ex Walp. Rep. iii. 346 = parviflora.
florida, Salisb. Prod. 92 = V. Teucrium.
florida, F. W. Schmidt, Fl. Boëm. i. 23 = V. Chamaedrys.
fluminensis, *Vell. Fl. Flum.* 11; i. t. 25.—Bras.
foliosa, Waldst. & Kit. Pl. Rar. Hung. ii. 106 = spuria.
Fonki, *Phil. in Linnaea*, xxix. (1857–58) 110.—Chili.
fontana, Willd. ex Link, Jahrb. i. III. (1820) 41 = serpyllifolia.
fontinalis, Salisb. Prod. 91 = V. Beccabunga.
formosa, *R. Br. Prod.* 434.—Tasmania.
fragilis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 446.—Persia.
Friesiana, Knaf, in Flora, xxix. (1846) 305 = agrestis.
Friesii, *Nem. ex Chaub. in Act. Linn. Soc. Bord.* xix. (1853) 232.—Gallia.
Froelichiana, Reichb. Fl. Germ. Excurs. 367 = V. Chamaedrys.
frutescens, Scop. Fl. Carn. ed. II. i. 19 = fruticulosa.
fruticans, Jacq. Enum. Stirp. Vindob. 2, 200 = fruticulosa.
fruticosa, Wulf. ex Steud. Nom. ed. II. ii. 757 = seq.

VERONICA:—
fruticulosa, *Linn. Sp. Pl.* ed. II. 15.—Europ.; Groenland.
gadensis, *Gueldenst. Reisen*, i. 426; *ex Ledeb. Fl. Ross.* iii. 256.—Reg. Cauc.
galactites, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 232 = spuria.
galathica, *Boiss. Fl. Orient.* iv. 448.—Reg. Cauc.
galeopsifolia, Gilib. Fl. Lituan. i. 104 = spicata.
Gebhardiana, Vest, ex Schult. Mant. i. 104 = alpina.
geniculata, Host, Fl. Austr. i. 5 = longifolia.
gentianifolia, Sweet, Hort. Brit. ed. II. 397 = gentianoides.
gentianoides, *Vahl, Symb. Bot.* i. 1.—Reg. Cauc.; As. Min.
glaberrima, *Boiss. & Bal. Diagn.* Ser. II. iii. 172.—Cilicia.
glabra, Ehrh. ex Schrad. Comm. Veron. Spic. 25 = longifolia.
glabriuscula, Opiz, Natural. n. ix. (1825) 111, nomen = V. Chamaedrys.
glandulifera, Freyn, in Oestr. Bot. Zeitschr. xxvi. (1876) 371 = V. Cymbalaria?
glandulifera, Opiz, Natural. n. ix. (1825) 111 = spicata.
glandulosa, *Hochst. ex Benth. in DC. Prod.* x. 482.—Abyss.
glauca, *Sibth. & Sm. Fl. Graec. Prod.* i. 9.—Graecia.
glauca, Ung. ex Nym. Consp. 548 = peloponnesiaca.
glauco-caerulea, Armst. in Trans. N. Z. Inst. xiii. (1881) 353 = pimeleoides.
gorumsensis, Boiss. & Kotschy, ex Boiss. Fl. Orient. iv. 458 = acinifolia.
Gouani, Moretti, in Brugnat. Giorn. Fis. Dec. II. v. (1822) 41, in obs. = Ponae.
gracilis, *R. Br. Prod.* 435.—Austral.
graeca, Sprun. ex Benth. in DC. Prod. x. 484 = glauca.
graminea, [Mazziari,] in Ionios, Anthol. ii. (1834) 442 = V. Anagallis.
grandiflora, *D. Don, in Fellows, Discov. in Lycia*, 292; *et in Ann. & Mag. Nat. Hist.* Ser. I. vii. (1841) 458.—Caria.
grandiflora, Gaertn. in Comm. Acad. Petrop. xiv. (1770) 531. t. 18. f. 1 = kamtchatica.
grandiflora, Wall. Cat. n. 403 = himalensis.
grandifolia, Vest, ex Schult. Mant. i. 103 = alpina.
grandis, *Fisch. ex Spreng. Neue Entdeck.* ii. 122.—Europ.
Grayi, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 354.—N. Zel.
Griffithii, Benth. in DC. Prod. x. 485 = biloba.
grossa, Mart. Hort. Erlang. (1814) 12 = longifolia.
Guentheri, Opiz, Natural. n. ix. (1825) 111, nomen = officinalis.
Gunnii, Benth. in DC. Prod. x. 477 = calycina.
Haastii, *Hook. f. Handb. N. Zeal. Fl.* 213.—N. Zel.
Haussknechtii, *Boiss. Fl. Orient.* iv. 438.—Mesopot.
Hectori, *Hook. f. Handb. N. Zel. Fl.* 212.—N. Zel.
hederaefolia, *Linn. Sp. Pl.* 13.—Europ.; Oriens; As. bor.
hederaefolia, Miq. ex Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 510 = agrestis.
hederifolia, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 190 = hederaefolia.
herniarioides, Pourr. ex Lapeyr. Suppl. Hist. Pl. Pyr. 4 = alpina.
Henningii, Opiz, ex Walp. Rep. iii. 363, in nota, sphalm. = *Nenningii*.
heterophylla, Salisb. Ic. Stirp. Rar. 7 = orientalis.
heterophylla, Steud. Nom. ed. II. ii. 757 = latifolia.
heterophyllos, Boeber, ex Ledeb. Fl. Ross. iii. 256, sphalm. = *tetraphyllos*.
Hillebrandi, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 49 = distans.
himalensis, *D. Don, Prod. Fl. Nep.* 92.—Reg. Himal.
hirsuta, Hopkirk, Fl. Glott. 9 = officinalis.
hirsuta, Lucé, Topog. Nachr. Oesel, 5 = arvensis.
hirsuta, Vukot. ex Nym. Consp. 548 = acinifolia.
hirsutissima, Opiz, Natural. n. ix. (1825) 111, nomen = V. Chamaedrys.
hispanica, Meerb. Pl. Rar. t. 11 = pinnata.
hispanica, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 601 = Assoana.
hispidula, *Boiss. & Huet, Diagn.* Ser. II. iii. 173.—Armenia.
Hookeriana, *Walp. Rep.* iii. 341.—N. Zel.

VERONICA:—
hortensis, Opiz, Natural. n. ix. (1825) 111, nomen = V. Chamaedrys.
hospita, Mert. & Koch, Deutschl. Fl. i. 332 = Tournefortii.
Hostii, Moretti, in Bibl. Ital. xii. (1818) 369 = longifolia.
Hulkeana, *F. Muell. ex Hook. f. Handb. N. Zeal. Fl.* 213.—N. Zel.
humifusa, Dicks. ex With. Bot. Arr. Brit. Pl. ed. III. ii. 15 = serpyllifolia.
humile, *Janka, in Termész. Füzetek*, iv. (1880) 294.—Europ.
hybrida, Bieb. Fl. Taur. Cauc. i. 8 = orchidea.
hybrida, Georgi, Beschr. Russ. Reich. Nachtr. 256 = longifolia.
hybrida, Linn. Sp. Pl. 11 = spicata.
hyssopifolia, Schur, Enum. Pl. Transs. 493 = austriaca.
imbricata, Woerlein, in Ber. Bot. Ver. Landshut, viii. (1882) 199.—Europ.
imperfoliatā, Benth. in DC. Prod. x. 463 = perfoliata.
incana, *Linn. Sp. Pl.* 10.—Europ. austr. or.; As. bor.
incana, F. W. Schmidt, Fl. Boëm. i. 5 = longifolia.
incana, Waldst. & Kit. Pl. Rar. Hung. iii. 271. t. 244 = spicata.
incarnata, Lodd. ex Loud. Hort. Brit. 6 = spuria.
incarnata, Stokes, Bot. Comm. i. 85 = longifolia.
incisa, Opiz, Natural. n. ix. (1825) 111, nomen = latifolia.
incisa, [Soland. in] Ait. Hort. Kew. ed. I. i. 19 = spuria.
indica, Roxb. ex A. Dietr. Sp. Pl. i. 522 = V. Anagallis.
integrifolia, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = V. Teucrium.
integrifolia, Schrank, Prim. Fl. Salisb. 20 = alpina.
intermedia, Schwein. in Am. Journ. Sc. Ser. I. viii. (1824) 268 = americana.
intermedia, Sternb. ex Schult. Mant. i. 110 = V. Chamaedrys.
intermedia, *Terrac. ex Ann. Acc. Asp. Napol.* Ser. III. vi. (1866).—*Cf. Bull. Soc. Bot. Fr.* xvi. (1869) *Bibl.* 57.—Europ. austr.
irregularis, Lapeyr. Hist. Abr. Pl. Pyr. i. 6 = nummularia.
irrigans, *T. Kirk, in Trans. N. Z. Inst.* ii. (1870) 94.—N. Zel.
ixodes, *Boiss. & Bal. Diagn.* Ser. II. iii. 172.—Cappadoc.
Jacquini, Baumg. Enum. Stirp. Transs. i. 26 = austriaca.
japonica, Steud. Nom. ed. II. ii. 757 = virginica.
javanica, Blume, Bijdr. 742.—Reg. Himal.; Malaya.
Jungbaueri, Opiz, Natural. n. ix. (1825) 110 = longifolia.
kamtchatica, *Linn. f. Suppl.* 83.—Kamtchat.; Ins. Aleut.
kermesina, Loud. Encyc. Pl. 1546 = speciosa.
Kirkii, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 356.—N. Zel.
Koeleri, *Roem. & Schult. Syst.* i. 127 (sp. dub.).—Hab.?
Kosteleckii, Opiz, Natural. n. ix. (1825) 109 = longifolia.
Kotschyana, *Benth. in DC. Prod.* x. 480.—Cilicia.
kurdica, *Benth. l. c.* 473.—As. Min.; Persia.
labiata, R. Br. Prod. 434 = V. Derwentia.
laciniata, Moench, Meth. 435 = austriaca.
laciniata, [Soland. in] Ait. Hort. Kew. ed. I. i. 19 = spuria.
laeta, Hort. ex Reichb. Fl. Germ. Excurs. 370 = spicata.
laeta, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 414.—Sibir.
laevis, *Benth. in DC. Prod.* x. 461.—N. Zel.
laevis, Lam. Fl. Fr. ii. 444 = peregrina.
lamiifolia, Hayne, in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 132 = V. Chamaedrys.
lanceolata, *Benth. in DC. Prod.* x. 462.—N. Zel.
lanceolata, Opiz, Natural. n. ix. (1825) 111, nomen = officinalis.
lanosa, Royle, ex Benth. Scroph. Ind. 45 = deltigera.
lanuginosa, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 293.—Reg. Himal.
Lappago, Schrank, Baier. Fl. i. 218 = hederaefolia.
latifolia, Linn. Herb. ex Nym. Consp. 545 = V. Teucrium.
latifolia, *Linn. Sp. Pl.* 13.—Europ.; Reg. Cauc.

VERONICA:—
Lavaudiana, *Raoul, Choix Pl. Nouv. Zél.* 16. t. 10.—N. Zel.
laxa, *Benth. Scroph. Ind.* 45.—Reg. Himal.; Japan.
laxiflora, *Lej. Rev. Fl. Spa*, 4.—Belgium.
leiocarpa, Boiss. Diagn. Ser. I. xii. 45 = stenobotrys.
leptostachya, Hort. ex Reichb. Fl. Germ. Excurs. 371 = spuria.
leucantha, Helm, in Mém. Soc. Nat. Mosc. ii. (1809) 104. t. 8 = spuria.
Lewisii, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 357.—N. Zel.
ligustrifolia, *A. Cunn. in Bot. Mag.* sub t. 3461.—N. Zel.
lilacina, Towns. in Bull. Soc. Bot. Fr. xxv. (1878) 16 = bellidioides.
limosa, Lej. Rev. Fl. Spa, 2 = V. Beccabunga.
linariaefolia, Pall. ex Link, Jahrb. i. III. (1820) 35 = spuria.
linariaefolia, *Vis. in Att. Ist. Venet.* Ser. III. iv. (1858–59) 134.—Hab.?
Linastrum, Arv. Ess. Pl. Dauph. 55 (Quid?).—Gallia.
Lindleyana, Paxt. Mag. Bot. xii. (1846) 247 = salicifolia.
Lindleyana, Wall. Cat. n. 404 = Picrorhiza Kurroa.
linifolia, *Hook. f. Handb. N. Zeal. Fl.* 214.—N. Zel.
litoralis, *Vell. Fl. Flum.* 10; i. t. 24.—Bras.
liwanensis, C. Koch, in Linnaea, xxii. (1849) 698 = telephiifolia.
loganioides, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 359.—N. Zel.
longebracteata, Link, Enum. Hort. Berol. i. 20 = spicata.
longe-pedunculata, Gilib. Fl. Lituan. i. 118 = agrestis.
longiflora, *Roem. & Schult. Syst.* i. 95.—Hab.?
longifolia, Hoffm. Phytog. Blaetter, i. (1803) 92 = spicata?
longifolia, *Linn. Sp. Pl.* 10.—Europ.; As. Min.; Sibir.
longifolio-repens, Gilib. Fl. Lituan. i. 117 = V. Anagallis.
longipes, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = V. Cymbalaria.
longistyla, Ball, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 47 = aphylla.
lucens, T. Kirk, ex Armst. in Trans. N. Z. Inst. xiii. (1881) 355 = canterburiensis.
lucida, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = serpyllifolia.
Luetkeana, *Rupr. Sert. Tiansch.* 62.—Turkestan.
lutetiana, Schult. Mant. i. 109 = V. Teucrium.
luxurians, Ledeb. Fl. Alt. i. 27, in Adnot. = longifolia.
Lyallii, *Hook. f. Fl. N. Zel.* i. 196.—N. Zel.
lycopodioides, *Hook. f. Handb. N. Zeal. Fl.* 211.—N. Zel.
lysimachioides, *Boiss. Diagn.* Ser. II. iii. 165.—Lydia.
macrantha, *Hook. f. Handb. N. Zeal. Fl.* 213.—N. Zel.
macrocalyx, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 353.—N. Zel.
macrocarpa, Hort. ex Reichb. Fl. Germ. Excurs. 371 = longifolia.
macrocarpa, Turcz. ex Steud. Nom. ed. II. ii. 758 = ciliata.
macrocarpa, *Vahl, Symb. Bot.* iii. 4.—N. Zel.
macrodonta, Borb. in Oestr. Bot. Zeitzschr. xxxiv. (1884) 73 = prostrata.
macropoda, *Boiss. Diagn.* Ser. I. iv. 79.—Persia.
macrostachya, Vahl, Enum. i. 71 = polifolia.
macrostemon, *Bunge, ex Ledeb. Fl. Alt.* i. 35.—Reg. Himal.; As. bor.
macroura, *Hook. f. ex Benth. in DC. Prod.* x. 459.—N. Zel.
Maddeni, Edgew. ex Hook. f. Fl. Brit. Ind. iv. 296 = javanica.
major, Presl, ex Schult. Mant. i. Add. II. 229, in syn. = serpyllifolia.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 19.—Afr. trop.
marilandica, Linn. Sp. Pl. 14 = Polypremum procumbens.
marilandica, Murr. in Comm. Gotting. (1782) 11. t. 3 = peregrina.
maritima, Linn. Sp. Pl. 10 = longifolia.
maxima, Mill. Gard. Dict. ed. VIII. n. 11 = latifolia.
maxima, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 179 = melissaefolia.

VERONICA:—
media, Lej. Rev. Fl. Spa, 4 = spicata.
media, Schrad. Comm. Veron. Spic. 23 = longifolia.
melancholica, Desf. ex Sweet, Hort. Brit. ed. II. 397 = seq.
melincolica, Hort. Par. ex Schult. Mant. i. 117 (Quid?).—Hab.?
melissaefolia, *Poir. Encyc.* viii. 526.—As. Min.; Reg. Cauc.
menthaefolia, Schott, ex Roem. & Schult. Syst. i. 94 = spicata.
Menziesii, Benth. in DC. Prod. x. 461 = elliptica.
mesotypa, *Link, Handb.* i. 522.—Europ.
Michauxii, *Lam. Illustr.* i. 44.—Persia.
micrantha, Hoffmgg. & Link, Fl. Port. i. 286 = V. Chamaedrys.
micrantha, Schur, Enum. Pl. Transs. 978 (Index) = arvensis.
microcarpa, *Boiss. Diagn.* Ser. I. iv. 76.—Armen.; Persia.
microcoma ×, Borb. in Oestr. Bot. Zeitschr. xxix. (1879) 134.—Europ.
microphylla, Kit. in Schult. Oestr. Fl. ed. II. i. 20 = serpyllifolia.
microtheca, Boiss. & Bal. Diagn. Ser. II. vi. 131 = biloba.
minima, *Echeand. Fl. Caesaraug.* (1861) 21.—Hispan.
minima, C. Koch, in Linnaea, xxii. (1849) 700 = hispidula.
minor, Vahl, ex Mert. & Koch, Deutschl. Fl. i. 317 = V. Anagallis.
minuta, Krock. Fl. Siles. Suppl. i. 47 (Quid?).—Europ.
minuta, *C. A. Mey. Verz. Pfl. Cauc.* 105.—Reg. Cauc.
missurica, *Rafin. in Am. Monthly Mag.* (1818) 175.—Am. bor.
mollis, Rafin. New Fl. Am. iv. 38 (Quid?).—Am. bor.
mollis, Willd. ex Link, Jahrb. i. III. (1820) 39 = spuria.
mollis, Zea, ex Roem. & Schult. Syst. i. 112 = V. Chamaedrys.
mollissima, Hornem. Suppl. Hort. Hafn. 3 = praec.
mollissima, Link, Handb. i. 527; Hort. ex Steud. Nom. ed. II. ii. 758 = V. Chamaedrys.
monstrosa, Hort. ex Schult. Mant. i. 100 = orchidea.
monstrosa, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = officinalis.
montana, *Linn. Cent. Pl.* i. 3; *Amoen. Acad.* iv. 263.—Europ.
montana, Pall. Reise, ii. 522 = austriaca.
monticola, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 354.—N. Zel.
monticola ×, Trautv. in Bull. Acad. Pétersb. x. (1866) 398.—Reg. Cauc.
montioides, Boiss. Diagn. Ser. I. vii. 43 = V. Anagallis.
Muelleri, *J. Buch. in Trans. N. Z. Inst.* xiv. (1882) 351.—N. Zel.
Muelleri, Mart. ex Walp. Rep. iii. 361 = longifolia.
Muelleri, Vest, ex Schrank, in Flora, vii. (1824) II. Beibl. 8 = seq.
Muelleriana, Vest, ex Schult. Mant. i. 107 = officinalis.
multicaulis, Bernh. ex Walp. Rep. iii. 362 = longifolia.
multifida, Colm. Cat. Pl. Catalun. 118 = Assoana.
multifida, Jacq. Fl. Austr. iv. 15 = austriaca.
multifida, *Linn. Sp. Pl.* 13.—As. Min.; As. bor.
murorum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 508.—Japon.
muscosa, Hort. ex Schult. Mant. i. 117 = orchidea.
myrtifolia, Banks & Soland. ex Benth. in DC. Prod. x. 460 = macrocarpa.
neglecta, Kit. ex Schult. Oestr. Fl. ed. II. i. 17 = spicata.
neglecta, F. W. Schmidt, Fl. Boëm. i. 12 = serpyllifolia.
neglecta, Vahl, Enum. i. 59 = incana.
nemorosa, *Kastropp, in Oestr. Bot. Zeitschr.* xvii. (1867) 107.—Europ.
nemorum, Pall. ex Link, Jahrb. i. III. (1820) 42, in textu = peduncularis.
Nenningii, Opiz, Natural. n. ix. (1825) 110 = longifolia.
nigricans, C. Koch, in Linnaea, xvii. (1843) 288 = peduncularis.
Nimrodi, *Richter, ex Stapf, in Denkschr. Akad. Wien*, l. (1885) 91.—Mesopot.

VERONICA:—
nitens, Host, Fl. Austr. i. 7 = spicata.
nitens, Schult. Mant. i. 97 = longifolia.
nitida, Ehrh. ex Roem. & Schult. Syst. i. 87 = spuria.
nitida, Hort. ex Poir. Encyc. Suppl. v. 467 = V. Teucrium.
nitida, Link, Enum. Hort. Berol. i. 20 = longifolia.
nitidula, Reichb. ex Steud. Nom. ed. II. ii. 755, in syn. = agrestis.
nitidula, Reichb. ex Walp. Rep. iii. 336 = didyma.
nivalis, Benth. in DC. Prod. x. 477, sphalm. = Hookeriana.
nivalis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 55; x. (1859) 139 = alpina.
nivea, Hook. f. in Hook. Ic. Pl. t. 640 = Hookeriana.
nivea, *Lindl. Bot. Reg.* (1842) *Misc.* 42.—Austral.
nodiflora, Pritz. Ic. Ind. i. 1155 = Verbena nodiflora.
Nočana, Boiss. Diagn. Ser. II. iii. 168 = orientalis.
notabilis, *F. Muell. ex Benth. Fl. Austral.* iv. 511.—Austral.
nova, Koel. ex Roem. Collect. 306 = Koeleri.
Novae-Hollandiae, Poir. Encyc. viii. 526 = distans?
nudicaulis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 415.—Persia; Soongar.
nudicaulis, Lam. Fl. Fr. ii. 437 = aphylla.
Nummularia, *Gouan, Illustr.* 1. t. 1. f. 2.—Mont. Pyren.
nummulariaefolia, Nym. Consp. 547 = V. Nummularia.
nummulariaefolia, Thuill. Fl. Par. ed. II. 6 = serpyllifolia.
nutans, Bong. in Mém. Acad. Pétersb. ii. (1833) 157 = alpina.
obliqua, *Kit. in Linnaea*, xxxii. (1863) 445 (Quid?).—Hungar.
obovata, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 502.—N. Zel.
obscura, Dum. Fl. Belg. 35 = agrestis.
obtusiuscula, Opiz, Natural. n. ix. (1825) 110, nomen = spicata.
ocymifolia, Thuill. Fl. Par. (1824) 10 = praecox.
odora, Hook. f. Fl. Antarct. i. 62. t. 41 = buxifolia.
officinalis, *Linn. Sp. Pl.* 11.—Europ.; As. Min.; As. bor.
officinarum, Crantz, Stirp. Austr. ed. II. fasc. iv. 336 = praec.
olympica, Pall. ex Link, Jahrb. i. III. (1820) 40 = gentianoides.
Onoei, *Franch. & Sav. Enum. Pl. Jap.* ii. 457.—Japon.
opaca, Fries, Novit. Fl. Suec. ed. II. 3 = agrestis.
oppositifolia, Gilib. Exercit. i. 107 = spicata.
orbicularis, Fisch. ex Trautv. in Bull. Acad. Pétersb. x. (1866) 397 = telephiifolia.
orbiculata, Kern. ex Nym. Consp. 545 = prostrata.
orchidea, *Crantz, Stirp. Austr.* ed. II. fasc. iv. 333.—Reg. Danub. et Cauc.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 10; *Ait. Hort. Kew.* ed. I. i. 23.—As. Min.; Persia.
orientalis, Willd. Enum. Hort. Berol. i. 22 = austriaca.
Orsiniana, Tenore, App. v. 4; et Succ. Relaz. 43 = prostrata.
osiliensis, Lucé, Topog. Nachr. Oesel, 4 = V. Anagallis.
ossetica, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 180 = caucasica.
ovalifolia, Schur, Enum. Pl. Transs. 496 = spicata.
oxycarpa, Boiss. Diagn. Ser. I. vii. 44 = V. Anagallis.
oxyphylla, Stev. ex Bess. Enum. Pl. Volh. 48 = longifolia.
Paederotae, *Boiss. Diagn.* Ser. I. iv. 78.—Persia.
pallens, Host, Fl. Austr. i. 6 = spicata.
pallescens, Gater. Desc. Pl. Montaub. 27 = peregrina.
pallida, Hornem. Hort. Hafn. i. 17 = gentianoides.
palmatiloba, Miégev. in Bull. Soc. Bot. Fr. xiv. (1867) 146 = verna.
paludosa, Lej. Fl. Spa, i. 22 = spicata.
palustris, Salisb. Prod. 92 = V. Anagallis.
paniculata, Linn. Syst. ed. X. 849 = spuria.
paniculata, Willd. Sp. Pl. i. 71 = V. Teucrium.
pannonica, Mill. Gard. Dict. ed. VIII. n. 5 = orchidea.
panormitana, Tineo, in Guss. Suppl. Fl. Sic. Prod. 4 = V. Cymbalaria.
parmularia, Poit. & Turp. Fl. Par. 16. t. 14 = scutellata.
parviflora, *Vahl, Symb. Bot.* iii. 4.—N. Zel.
parviflora, Vahl, Enum. i. 72, sphalm. = seq.
parvifolia, Vahl, l. c. 362 = orientalis.

VERONICA:—
pauciflora, Kit. ex Link, Jahrb. i. III. (1820) 42 = Baumgartenii.
pectinata, Georgi, Beschr. Russ. Reich. III. iv. 652 = orientalis.
pectinata, *Linn. Mant.* i. 24.—Oriens.
pectinata, Opiz, ex Schult. Mant. i. Add. II. 229 = prostrata.
pedunculariis, *Bieb. Fl. Taur. Cauc.* i. 11.—As. Min.; Reg. Cauc.
pedunculariis, Bory & Chaub. Nouv. Fl. Pélop. 1 = Chaubardi.
pedunculariis, Stev. ex Reichb. Fl. Germ. Excurs. 369, in obs. = caucasia.
pedunculata, Labill. Ic. Pl. Syn. Dec. v. 8 = syriaca.
pedunculata, Vahl, Enum. i. 77 = V. Chamaedrys.
peloponnesiaca, *Boiss. & Orph. in Boiss. Fl. Orient*, iv. 462.—Graecia.
peregrina, *Linn. Sp. Pl.* 14.—Am. bor.
perfoliata, *R. Br. Prod.* 434.—Austral.
perfoliata, Rafin. New Fl. Am. iv. 37 (Quid?).—Am. bor.
perpusilla, Boiss. Diagn. Ser. I. vii. 43 = nudicaulis.
persica, Hort. ex Poir. Encyc. viii. 542 = Tournefortii.
persicifolia, Schott, ex Link, Enum. Hort. Berol. i. 21 = longifolia.
peruviana, Willd. ex A. Dietr. Sp. Pl. i. 534 = peregrina.
Petersii, Opiz, in Weitenw. Beitr. i. (1836) 67 (err. typ. 76) = *Pettersii*.
petiolaris, Lapeyr. Hist. Abr. Pl. Pyren. i. 5 = longifolia.
petiolata, Opiz, Natural. n. ix. (1825) 111, nomen = V. Chamaedrys.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 127.— Abyss.
petraea, Baumg. Enum. Stirp. Transs. i. 21 = Baumgartenii.
petraea, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 250.—Reg. Cauc.
Pettersii, Opiz, Natural. n. ix. (1825) 109; et x. (1825) 245 = longifolia.
phoenicantha, C. Koch, in Linnaea, xxii. (1849) 690 = pedunculariis.
pilocarpa, Link, Enum. Hort. Berol. i. 25 = austriaca.
pilosa, Linn. Sp. Pl. ed. II. 1663 = V. Chamaedrys.
pimeleoides, *Hook. f. Fl. N. Zel.* i. 195.—N. Zel.
pinguifolia, *Hook. f. Handb. N. Zeal. Fl.* 210.—N. Zel.
pinnata, *Linn. Mant.* i. 24.—Sibir. altaic.
pinnata, Schur, Enum. Pl. Transs. 494 = multifida.
pinnatifida, Lam. Illustr. i. 47 = verna.
pinnatifida, Pohl, Tent. Fl. Bohem. i. 17 = austriaca.
pinnatifida, Salisb. Prod. 90 = pinnata.
pinnatifida, Willd. ex Schult. Mant. i. 117 (Quid?).—Hab.?
plebeia, *R. Br. Prod.* 435.—Austral.
plena, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = officinalis.
plicata, Pohl, Tent. Fl. Bohem. i. 15. f. 1 = V. Chamaedrys.
polifolia, *Benth. in DC. Prod.* x. 473.—As. Min.; Syria.
polita, Fries, Novit. Fl. Suec. ed. II. 1 = didyma.
polyanthos, Thuill. Fl. Par. ed. II. 9 = arvensis.
polygonoides, Lam. Illustr. i 47 = verna.
polymorpha, Willd. Enum. Hort. Berol. Suppl. 2 = austriaca.
polystachya, Link, Enum. Hort. Berol. i. 23 = longifolia.
Ponae, *Gouan, Illustr.* i. 1. t. 1. f. 1.—Europ.
praealta, Lodd. Cat. (1823); Sweet, Hort. Brit. ed. II. 397 (Quid?).—Hab.?
praecox, *All. Auct. Fl. Pedem.* 5. t. 1. f. 1.—Reg. Medit.
praetutiana, Moretti, in Bibl. Ital. lxxii. (1833) 217 = officinalis.
prasifolia, Desf. Cat. Hort. Par. ed. III. 82 = longifolia.
pratensis, Crantz, Stirp. Austr. ed. II. fasc. iv. 344 = prostrata.
precox, Rafin. Atl. Journ. 79 (Quid?).—Am. bor.
procerior, Presl, ex Schult. Mant. i. Add. II. 229 = verna.
procumbens, *Clarke, Trav.* i. 747, nomen.—Tauria.
prostrata, *Linn. Sp. Pl.* ed. II. 22.—Europ.
prostrata, Sibth. & Sm. Fl. Graec. Prod. i. 7 = V. Teucrium.
Pseudo-arvensis, Tineo, Pl. Rar. Sicil. 34 = arvensis.

VERONICA:—
Pseudo-chamaedrys, Jacq. Fl. Austr. i. 37. t. 60 = V. Teucrium.
Pseudo-incana, Schur, Enum. Pl. Transs. 496 = spicata.
pubescens, *Banks & Soland. ex Benth. in DC. Prod.* x. 461.—N. Zel.
pulchella, Bast. Fl. Maine-et-Loire, 144 = agrestis.
pulchella, *Bernh. ex Roem. & Schult. Syst.* i. 126.—Hab.?
pulchella, Salisb. Prod. 92 = V. Chamaedrys.
pulchra, G. Don, in Loud. Hort. Brit. 7 = arenaria.
pumila, All. Fl. Pedem. i. 75 = alpina.
pumila, *Clarke, Trav.* iv. 554, 559.—Bulgaria.
punctata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 93 = V. Anagallis.
purpurea, Steud. Nom. ed. II. ii. 759 = virginica.
Purshii, *G. Don, Gen. Syst.* iv. 573 (an generis?).—Am. bor.
pusilla, Benth. in DC. Prod. x. 468 = V. Anagallis.
pusilla, Hohen. & Boiss. ex Benth. in DC. Prod. x. 490 = nudicaulis.
pusilla, *Kit. in Linnaea*, xxxii. (1863) 449 (Quid?).—Hungar.
pygmaea, Schrank, Prim. Fl. Salisb. 20 = alpina.
pyrenaica, All. Fl. Pedem. i. 73 = Allionii.
quadrifolia, Schkuhr, Handb. i. 9 = longifolia.
quinquefida, Gilib. Fl. Lituan. i. 120 = triphyllos.
quinquefolia, Stokes, Bot. Mat. Med. i. 28 = virginica.
racemifoliata, *Perez-Lara, in Anal. Soc. Esp. Hist. Nat.* xi. (1882) 401.—Hispan.
rakaiensis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 356.—N. Zel.
ramosa, Pall. ex Link, Jahrb. i. III. (1820) 38 = spuria.
Raoulii, *Hook. f. Handb. N. Zeal. Fl.* 214.—N. Zel.
recta, Benth. in DC. Prod. x. 474 = austriaca.
reniformis, Pursh, Fl. Am. Sept. i. 10 = Purshii.
reniformis, *Rafin. in Med. Repos. N. York*, v. (1808) 360; *Journ. Bot.* i. (1808) II. 228.—Am. bor.
repens, *All. ex Roem. & Schult. Syst.* i. 110 = Allionii.
repens, Boiss. Voy. Bot. Espagne, ii. 469 = apennina.
repens, *DC. Fl. Fr.* iii. 727.—Ins. Corsic.
repens, Gilib. Fl. Lituan. i. 108 = officinalis.
reticulata, C. Koch, in Linnaea, xxii. (1849) 702 = ceratocarpa.
Reuteriana, *Boiss. Diagn.* Ser. I. xii. 46.—Mesopot.
rigens, Mart. Hort. Erlang. (1814) 13 = longifolia.
ringens, Steud. Nom. ed. II. ii. 759, sphalm. = praec.
riparia, Seidl, ex Bercht. Oek. Fl. Boehm. i. 30 = longifolia.
romana, All. Fl. Pedem. i. 79 = verna.
romana, Linn. Sp. Pl. 14 = peregrina.
romana, F. W. Schmidt, Fl. Boëm. i. 31 = arvensis.
rosea, *Desf. Fl. Atlant.* i. 13.—Alger.
rotundifolia, Gilib. Exercit. i. 120 = V. Beccabunga.
rotundifolia, Lucé, Fl. Osil. 3 = serpyllifolia.
rotundifolia, Ruiz & Pav. Fl. Per. i. 6 = Sibthorpia pichinchensis.
rotundifolia, Schrank, Prim. Fl. Salisb. 21 = alpina.
rotundifolio-erecta, Gilib. Fl. Lituan. i. 116 = V. Anagallis.
rubella, Willd. ex Link, Jahrb. i. III. (1820) 38 = spuria.
rubicunda, Ledeb. Fl. Alt. i. 28, in Adnot. = spuria.
rubrifolia, *Boiss. Diagn.* Ser. I. xii. 46.—Persia.
ruderalis, Vahl, Enum. i. 66 = serpyllifolia.
Rudolphiana, Hayne, in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 133 = V. Chamaedrys.
rugosa, Hort. ex Reichb. Fl. Germ. Excurs. 370 = spicata.
rupestris, Aitch. & Hemsl. in Journ. Linn. Soc. xix. (1882) 180. t. 25 = deltigera.
rupestris, Salisb. Prod. 91 = fruticulosa.
rupestris, *Tardent, Ess. Hist. Nat. Bessarab.* 49.—Ross. austr.
ruthenica, Roem. & Schult. Syst. i. 96 = longifolia.
ruthenica, Spreng. ex Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 22, 23 = praec.?
salicifolia, A. Cunn. ex Hook. f. Handb. N. Zeal. Fl. 207 = macrocarpa.
salicifolia, *Forst. f. Prod.* 3.—N. Zel.
salicornioides, *Hook. f. Handb. N. Zeal. Fl.* 212.—N. Zel.
salina, Schur, Enum. Pl. Transs. 492 = anagalloides.
Sartoriana, *Boiss. & Heldr. Diagn.* Ser. II. iii. 171.—Graecia.
satureiaefolia, Poit. & Turp. Fl. Par. 18. t. 17 = prostrata.

VERONICA:—
saturejoides, *Vis. Fl. Dalm.* ii. 168. t. 33. f. 2 *.—Dalmat.
saxatilis, Scop. Fl. Carn. ed. II. i. 11 = fruticulosa.
scardica, Griseb. Spicil. Fl. Rumel. ii. 31 = V. Beccabunga.
Schmidtiana, *Regel, Ind. Sem. Hort. Petrop.* (1864) 22.—Ins. Sachalin.
Schmidtii, Pohl, ex Schult. Mant. i. 108 = V. Chamaedrys.
Schmidtii, Roem. & Schult. Syst. i. 115 = V. Teucrium.
Schreberi, Baumg. Fl. Lips. 6 = longifolia.
Schreberiana, Reichb. Fl. Germ. Excurs. 371 = praec.
sciaphila, *Menyh. Kalocsa Vidék. Növenyt.* 131.—Hungar.
scrophularioides, Wall. Cat. sub n. 401 = cana.
scutellata, *Linn. Sp. Pl.* 12.—Reg. bor. temp.
secundiflora, C. Koch, in Linnaea, xvii. (1843) 288 = pedunculariis.
segetalis, Pers. in Usteri, Ann. Bot. xiv. (1795) 38 = praecox.
sempervirens, Lam. Fl. Fr. ii. 436 = Ponae.
serotina, Hort. ex Roem. & Schult. Syst. i. 89 = longifolia.
serpyllifolia, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 598 = apennina.
serpyllifolia, *Linn. Sp. Pl.* 12.—Reg. bor. temp.
serpylloides, *Regel, in Act. Hort. Petrop.* vi. (1879) 345.—Turkestan.
serrulata, Willd. ex Link, Jahrb. i. III. (1820) 38 = spuria.
sessiliflora, *Bunge, in Ledeb. Fl. Atl.* i. 32.—Sibir. altaic.
sessilifolia, Opiz, Natural. n. ix. (1825) 111, nomen = spicata, V. Chamaedrys.
setigera, D. Don, in Mem. Wern. Soc. iii. (1821) 297 = officinalis.
sibirica, Linn. Sp. Pl. ed. II. 12 = virginica.
sibthorpiaefolia, *Chaub. in Act. Linn. Soc. Bord.* xix. (1853) 233.—Gallia.
Sieboldiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 119.—Japon.
simensis, Fresen. in Bot. Zeit. ii. (1844) 356 = violaefolia.
simplex, Opiz, ex Schult. Mant. i. Add. II. 229 = hederaefolia, arvensis.
Simpsoni, *Phil. in Anal. Univ. Chil.* (1873) 526.—Chili.
spadana, Lej. Fl. Spa, i. 22 = officinalis.
sparsiflora, Rafin. Atl. Journ. 79 (Quid?).—Am. bor.
spathulata, *Benth. in DC. Prod.* x. 477.—N. Zel.
speciosa, *R. Cunn. in Bot. Mag.* sub t. 3461.—N. Zel.
spicata, *Linn. Sp. Pl.* 10.—Europ.; As. bor.
spicato-racemosa, Gilib. Fl. Lituan. i. 107 = serpyllifolia.
spuria, *Linn. Sp. Pl.* 10.—Europ.
spuria, Schkuhr, Handb. i. 9 = longifolia.
squamosa, J. & C. Presl, Fl. Čech. 2 = spicata.
stachyoides, Schur, Enum. Pl. Transs. 497 = spicata.
stenobotrys, *Boiss. & Blanche, Diagn.* Ser. II. iii. 166.—Cilicia; Syria.
stenophylla, Steud. Nom. ed. II. ii. 760 = parviflora.
Stephaniana, Roem. & Schult. Syst. i. Mant. 96 = spuria.
Stelleri, *Pall. ex Link, Jahrb.* i. III. (1820) 40.—Ins. Aleut.
Sternbergiana, Bernh. Ehrenp. 21 = spicata.
stolonifera, Lehm. Sem. Hort. Hamb. (1820); (1824) 20; et Pl. Preiss. i. 342 = calycina.
stricta, Banks & Soland. ex Benth. in DC. Prod. x. 459 = salicifolia.
stricta, Lodd. Cat. (1823); ex Schult. Mant. i. Add. II. 228 (Quid?).—Hab.?
strictiflora, Steud. Nom. ed. I. 879 = nummularia.
subacaulis, Lam. Illustr. i. 44 = aphylla.
subarctica, Schur, Enum. Pl. Transs. 495 = officinalis.
subbiscutata, Crantz, Stirp. Austr. ed. II. fasc. iv. 343 = montana.
subincana, *Hoffm. in Comm. Gotting.* xv. (1804) 125. t. 7.—Europ.
sublaciniata, G. Don, Gen. Syst. iv. 567 = spuria.
succulenta, All. Fl. Pedem. i. 78 = verna.
surculosa, *Boiss. & Bal. Diagn.* Ser. II. iii. 170.—Cilicia.
syriaca, *Roem. & Schult. Syst.* i. 116.—As. Min.; Syria.

VERONICA:—
syspirensis, C. Koch, in Linnaea, xxii. (1849) 698 = kurdica.
taurica, Willd. Sp. Pl. i. 70 = orientalis.
telephiifolia, *Vahl, Enum.* i. 65.—Reg. Cauc.
tenella, All. Fl. Pedem. i. 75 = serpyllifolia.
tenella, Jan, ex Ces. Passer. & Gib. Comp. Fl. Ital. 350 = anagalloides.
tenella, Viv. Fl. Cors. Spec. Nov. 3 = repens.
tenerrima, F. W. Schmidt, in Mayer, Samml. Phys. Aufs. i. (1791) 198 = V. Anagallis.
tenuifolia, Asso, Syn. Stirp. Arag. 2. t. 1 = Assoana.
tenuifolia, Bieb. Fl. Taur. Cauc. i. 13 = multifida.
tenuifolia, Desf. ex Steud. Nom. ed. I. 879 (Quid?).—Hab.?
tenuis, Ledeb. Fl. Alt. i. 38 = anagalloides.
ternifolia, Opiz, ex Schult. Mant. i. Add. II. 229, in syn. = V. Teucrium.
tetragona, *Hook. Ic. Pl.* t. 580.—N. Zel.
tetraphyllos, Boeber. ex Georgi, Beschr. Russ. Reich. III. iv. 653 (Quid?).—Ross. austr.
tetrasticha, *Hook. f. Handb. N. Zeal. Fl.* 212.—N. Zel.
teucrioides, Boiss. & Heldr. Diagn. Ser. II. iii. 169 = thymifolia.
teucriphylla, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = V. Teucrium.
Teucrium, Brot. Fl. Lusit. i. 13 = V. Chamaedrys.
Teucrium, Crantz, Stirp. Austr. ed. II. fasc. iv. 337, partim = alpina.
Teucrium, *Linn. Sp. Pl.* ed. II. 16.—Europ.; As. bor.
Teucrium, Pollich, Hist. Pl. Palat. i. 12 = latifolia.
Teucrium, Savi, Due Cent. 5 = prostrata.
thessalica, *Benth. in DC. Prod.* x. 480.—Graecia.
Thunbergii, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 402 = laxa.
thymifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 6.—Graecia; Creta.
thymifolia, Willd. ex Link, Jahrb. i. III. (1820) 41 = euphrasiaefolia.
thyrsiflora, Wall. Cat. n. 6393 = Scrophularia calycina.
ticinensis, Pollini, Fl. Veron. iii. App. 768 = longifolia.
Tournefortii, *C. C. Gmel. Fl. Bad.* i. 39.—Europ.; As. temp.; Afr. bor.
Tournefortii, Vill. Prosp. 20; Hist. Pl. Dauph. ii. 9 = officinalis.
Townsendi, Gremli, ex Dalla Torre, Atl. Alpenfl. 174 = bellidioides.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 55 = longifolia.
Traversii, *Hook. f. Handb. N. Zeal. Fl.* 208.—N. Zel.
trichadena, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 42 = V. Cymbalaria.
trichocarpa, Schult. Mant. i. 106 = austriaca.
trifida, Gilib. Fl. Lituan. i. 121 = verna.
triloba, Opiz, Natural. 108; et in Lotos, iv. (1854) 157 = hederaefolia.
triphyllos, Gouan, ex Nym. Consp. 547 = verna.
triphyllos, *Linn. Sp. Pl.* 14.—Europ.; Afr. bor.; As. Min.
triquetra, Kit. ex Walp. Rep. iii. 363 = spuria.
trisepala, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 324.—N. Zel.
tubiflora, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 53.—Dahuria.
tymphrestea, Boiss. & Sprun. Diagn. Ser. I. iv. 77 = thymifolia.
uliginosa, Rafin. in Am. Monthly Mag. (1818) 175 = scutellata.
umbrosa, Bieb. Fl. Taur. Cauc. i. 11 = peduncularis.
umbrosa, Reichb. ex Schult. Mant. i. Add. II. 229, in syn. = hederaefolia.
undulata, Wall. in Roxb. Fl. Ind. ed. Carey, i. 147 = V. Anagallis.
upsaliensis, Hort. ex Schult. Mant. i. 95 = spuria.
urticaefolia, Jacq. Fl. Austr. i. 37 = latifolia.
urticifolia, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 190 = praec.
Vaillantii, Reichb. ex Nym. Consp. 544 = spicata.
variegata, Hort. ex Vis. in Att. Ist. Venet. Sc. Ser. III. iv. (1858–59) 135 = versicolor.
Verbenae, Hort. ex Schult. Mant. i. 95 = spuria.
verna, *Linn. Sp. Pl.* 14.—Europ.; Oriens; As. bor.
vernicosa, *Hook. f. Handb. N. Zeal. Fl.* 208.—N. Zel.
versicolor, Drej. in Fl. Dan. t. 2221 = agrestis.

VERONICA:—
versicolor, *Vis. in Att. Ist. Venet. Sc.* Ser. III. iv. (1858–59) 134 (an hybr.?).—Hab.?
verticillata, Gilib. Fl. Lituan. i. 97 = longifolia.
Vestiana, Schult. Mant. i. 111 = melissaefolia.
villosa, Schrad. Comm. Veron. Spic. 31 = longifolia.
villosa, Steud. Nom. ed. I. 879 = scutellata.
villosa, Wormsk. ex Roem. & Schult. Syst. i. 101 = alpina.
villosiuscula, Krock. Fl. Siles. Suppl. i. 31 = serpyllifolia.
violaefolia, *Hochst. ex Benth. in DC. Prod.* x. 488.—Abyss.
virgata, Krock. Fl. Siles. Suppl. 22 (Quid?).—Europ.
virgata, Willd. ex Link, Jahrb. i. III. (1820) 39 = spuria.
virgata, Menyh. Kalocsa Vidék. Növényt. 134 = hederaefolia.
virginica, *Linn. Sp. Pl.* 9.—Am. bor.
viridis, Presl, ex Schult. Mant. i. Add. II. 229, in syn. = prostrata.
viscida, Waldst. ex Roem. & Schult. Syst. i. 124 = praecox.
viscosa, *Boiss. Diagn.* Ser. I. xii. 47.—Syria; Mesopot.
viscosa, Kit. ex A. Dietr. Sp. Pl. i. 529 = praecox.
viscosa, Willd. ex Link, Jahrb. i. III. (1820) 34 = spicata.
vitifolia, Sm. ex Walp. Rep. iii. 367 = plebeia.
vogerensis, Walp. Ann. iii. 196 = wogerensis.
vulgaris, Opiz, Natural. n. ix. (1825) 111, nomen = officinalis, V. Chamaedrys.
Waldsteiniana, Schott, ex Schult. Mant. i. 117 (Quid?).—Europ.
Wolffiana, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 55 = serpyllifolia.
wogerensis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 126.—Abyss.
Wormskioldi, Roem. & Schult. Syst. i. 101 = alpina.
xalapensis, H. B. & K. Nov. Gen. et Sp. ii. 389 = peregrina.
yedoënsis, *Franch. & Sav. Enum. Pl. Jap.* ii. 458.—Japon.

VERONICASTRUM, Heist. ex Fabr. Hort. Helmst. 111 (1759) = **Veronica**, Tourn. (Scrophul.).
album, Moench, Meth. 437 = Veronica virginica.
alpinum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = Veronica alpina.
dentatum, Opiz, Seznam, 102 = Veronica Teucrium.
diversifolium, Moench, Meth. Suppl. 158 = Veronica virginica.
incisum, Moench, l. c. = Veronica spuria.
laciniatum, Moench, l. c. = Veronica spuria.
latifolium, Opiz, Seznam, 103 (err. typ. 203) = Veronica latifolia.
prostratum, Opiz, l. c. 102 = Veronica prostrata.
rubellum, Moench, Meth. 437 = Veronica virginica.
serpyllifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = Veronica serpyllifolia.

VERONICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
Chamaedrys, Fourr. l. c. = Veronica Chamaedrys.
urticaefolia, Fourr. l. c. = Veronica latifolia.

VERREAUXIA, Benth. Fl. Austral. iv. 105 (1869).
GOODENOVIEAE, Benth. & Hook. f. ii. [II.] 540.
paniculata, *Benth. Fl. Austral.* iv. 105.—Austral.
Reinwardtii, *Benth. l. c.*—Austral.

VERRUCARIA, Medic. Malv. 103 (1787) = **Tournefortia**, Linn. (Borag.).
volubilis, Medic. l. c. 104 = T. volubilis.

VERRUCULARIA, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 327. *MALPIGHIACEAE*, Benth. & Hook. f. i. 255.
glaucophylla, *Juss. l. c.*—Bras.

VERSCHAFFELTIA, H. Wendl. in Illustr. Hortic. xii. (1865) Misc. 5. *PALMAE*, Benth. & Hook. f. iii. 908.
REGELIA, Hort. ex H. Wendl. in Illustr. Hortic. xii. (1865) Misc. 6.
melanochaetes, H. Wendl. ex André, l. c. xviii. (1871) 54. t. 54 = Roscheria melanochoetes.
splendida, *H. Wendl. l. c.* xii. (1865) Misc. 6.—Ins. Seychelles.

VERTICILLARIA, Ruiz & Pav. Prod. 81. t. 15 (1794) = **Rheedia**, Linn. (Guttif.).
acuminata, Ruiz & Pav. Syst. Veg. 140 = R. acuminata.
peruviana, G. Don, Gen. Syst. i. 615 = R. acuminata.

VERTICORDIA, DC. in Dict. Class. xi. 400 (1826).
MYRTACEAE, Benth. & Hook. f. i. 697.
CHRYSORRHOË, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 357.
DIPLACHNE, R. Br. ex Desf. in Mém. Mus. Par. v. (1819) 272.
acerosa, *Lindl. Swan River App.* 6.—Austral.
brachypoda, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 158 = multiflora.
brachystylis, F. Muell. Fragm. i. 164 = habrantha.
Brownii, *DC. Prod.* iii. 209.—Austral.
callitricha, Meissn. in Journ. Linn. Soc. i. (1856) 39 = monadelpha.
carinata, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 19 = Drummondii.
cespitosa, Turcz. l. c. xx. (1847) I. 157 = densiflora.
chrysantha, *Endl. in Ann. Wien. Mus.* ii. (1838) 195.—Austral.
chrysostachys, *Meissn. in Journ. Linn. Soc.* i. (1856) 41.—Austral.
compta, Endl. in Ann. Wien. Mus. ii. (1838) 194 = insignis.
conferta, Benth. Fl. Austral. iii. 22 = fastigiata.
Cunninghamii, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* ii. (1841) 207.—Austral.
cuspidata, A. Cunn. ex Schau. l. c. 212 = pennigera.
demissa, F. Muell. ex Benth. Fl. Austral. iii. 25 = oxylepis.
densiflora, *Lindl. Swan River App.* 6.—Austral.
Drummondii, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* ii. (1841) 208.—Austral.
Endlicheriana, Schau. in Lehm. Pl. Preiss. i. 101 = Preissii.
fastigiata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 327.—Austral.
fimbrilepis, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 158.—Austral.
fimbripetala, Turcz. l. c. xxii. (1849) II. 19 = Huegelii.
Fontanesii, *DC. Prod.* iii. 209.—Austral.
Forrestii, *F. Muell. in Wing, South. Sc. Record*, iii. (March, 1883).—Austral.
Gilbertii, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 160 = chrysantha.
grandiflora, *Endl. in Ann. Wien. Mus.* ii. (1838) 195.—Austral.
grandis, *Drumm. in Hook. Kew Journ.* v. (1853) 119.—Austral.
habrantha, *Schau. in Lehm. Pl. Preiss.* i. 99.—Austral.
Harveyi, *Benth. Fl. Austral.* iii. 22.—Austral.
heliantha, Lindl. Swan River App. 6 = grandiflora.
helichrysantha, *F. Muell. ex Benth. Fl. Austral.* iii. 21.—Austral.
hirta, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 327 = Preissii.
Huegelii, *Endl. in Enum. Pl. Hueg.* 46.—Austral.
Hughani, *F. Muell. Fragm.* xi. 10.—Austral.
humilis, *Benth. Fl. Austral.* iii. 26.—Austral.
insignis, *Endl. in Enum. Pl. Hueg.* 47.—Austral.
integrisepala, F. Muell. Fragm. iv. 57; et ex Benth. Fl. Austral. iii. 15 = Darwinia verticordina.
Jamiesonii, *F. Muell. in Wing, South. Sc. Record*, iii. (March, 1883).—Austral.
Lehmanni, *Schau. in Lehm. Pl. Preiss.* i. 99.—Austral.
lepidophylla, *F. Muell. Fragm.* i. 228.—Austral.
Lindleyi, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 210 = Drummondii.
minutiflora, *F. Muell. Fragm.* iv. 58.—Austral.
monadelpha, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 158.—Austral.
multiflora, *Turcz. l. c.* 159.—Austral.
nitens, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* ii. (1841) 223.—Austral.
nobilis, Meissn. in Journ. Linn. Soc. i. (1856) 39 = grandiflora.
oculata, *Meissn. l. c.* 41.—Austral.
ovalifolia, *Meissn. l. c.* 40.—Austral.
oxylepis, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 327.—Austral.

VERTICORDIA:—
pectinata, *Turcz. in Bull. Acad. Pétersb.* x. (1852) 327.—Austral.
penicillaris, *F. Muell. Fragm.* i. 226.—Austral.
pennigera, *Endl. in Enum. Pl. Hueg.* 46.—Austral.
pentandra, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 157 = picta.
pholidophylla, *F. Muell. Fragm.* i. 227.—Austral.
picta, *Endl. in Ann. Wien. Mus.* ii. (1838) 194.—Austral.
platystigma, A. Cunn. ex Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 214 = Huegelii.
polytricha, *Benth. Fl. Austral.* iii. 25.—Austral.
Preissii, *Schau. in Lehm. Pl. Preiss.* i. 101.—Austral.
Roei, Endl. in Ann. Wien. Mus. ii. (1838) 194 = insignis.
serrata, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* ii. (1841) 222.—Austral.
setigera, Lindl. Swan River App. 7 = pennigera.
Sieberi, Dies. ex Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 201 = Fontanesii.
spicata, *F. Muell. Fragm.* i. 226.—Austral.
stelluligera, *Meissn. in Journ. Linn. Soc.* i. (1857) 38.—Austral.
stylosa, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 160 = Huegelii.
umbellata, Turcz. l. c. 159 = habrantha.
Wilhelmii, *F. Muell. in Trans. Vict. Inst.* i. (1855) 122.—Austral.

VERULAMIA, DC. ex Poir. Encyc. viii. 543 (1808) = **Pavetta**, Linn. (Rubiac.).
corymbosa, DC. ex Poir. l. c. = P. Baconia.

VERUTINA, Cass. in Dict. Sc. Nat. xliv. 38 (1826); lviii. 8 (1829) = **Centaurea**, Linn. (Compos.).
heterophylla, Cass. l. c. lviii. 9 = C. Verutum.

VERZINUM, Rafin. Sylva Tellur. 23 (1838) = **Cytisus**, Linn. (Legumin.).
arboreum, Rafin. l. c. = C. arboreus.
patens, Rafin. l. c. = C. patens.

VESALEA, Mart. & Gal. in Bull. Acad. Brux. xi. 1. (1844) 242 = **Abelia**, R. Br. (Caprifol.).
floribunda, Mart. & Gal. l. c. = A. floribunda.
hirsuta, Mart. & Gal. l. c. = A. speciosa.

VESICARIA, Tourn. ex Adans. Fam. ii. 420 (1763).
CRUCIFERAE, Benth. & Hook. f. i. 73.
ALYSSOIDES, Tourn. ex Adans. Fam. ii. 419 (1763).
CISTOCARPIUM, Spach, Hist. Vég. Phan. vi. 471 (1838).
CLASTOPUS, Boiss. Fl. Orient. i. 261 (1867).
PHYSOPTYCHIS, Boiss. l. c. 260 (1867).
alpina, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 102.—Am. bor. occ.
andicola, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 138 = arctica.
angustifolia, A. Gray, Pl. Wright. ii. 13, partim = Gordoni.
angustifolia, Nutt. in Torr. & Gray, Fl. N. Am. i. 101 = gracilis.
angustifolia, Scheele, in Linnaea, xxi. (1848) 584 = recurvata.
arctica, Barn. in C. Gay, Fl. Chil. i. 161 = mendocina.
arctica, Coulter, in Hayden, Rep. (1872) 760 = ludoviciana?
arctica, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 9 = Fendleri.
arctica, *Richards. in Frankl. Narr.* 1*st Journ. App.* 743; *et App.* ed. II. 26.—Am. bor. et arct.
arenosa, Richards. ll. cc. = arctica.
argentea, *Schau. in Linnaea*, xx. (1847) 720.—Mexic.
argyraea, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 146.—Texas.
auriculata, *Engelm. & Gray, l. c.* v. (1847) 240.—Texas.
Barrelieri, Parl. Enum. Sem. Hort. Bot. Florent. (1863) 25 = graeca.
Berlandieri, *A. Gray, in S. Wats. Bibliog. Ind. N. Am. Bot.* i. 75.—Texas.
brevistyla, Torr. & Gray, Fl. N. Am. i. 101 = grandiflora.
canescens, *Schrank, in Syll. Ratisb.* ii. (1828) 63.—Podolia.

VESICARIA:—
cochlearioides, Spreng. Syst. ii. 872 = Cochlearia flava.
corymbosa, Nym. Syll. 202 = Alyssum corymbosum.
cretica, Poir. Encyc. viii. 570 = Alyssum creticum.
dasycarpa, Poir. l. c. 571 = Alyssum dasycarpum.
deltoidea, Poir. l. c. 572 = Aubrietia deltoidea.
densiflora, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 145.—Texas.
dentata, Sm. in Rees, Cycl. xxxvii. n. 1 = Coluteocarpus reticulatus.
didymocarpa, Hook. Fl. Bor. Am. i. 49 = Physaria didymocarpa.
digitata, *C. A. Mey. Verz. Pfl. Cauc.* 180.—Reg. Cauc.
edentula, Poir. Encyc. viii. 572 = Alyssum gemonense.
Engelmanni, *A. Gray, Gen. Am. Bor. Or. Illustr.* i. 162. t. 70.—Texas.
Fendleri, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 9.—N. Mexic.
frigida, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 296.—Venezuela.
fruticulosa, Desv. Journ. Bot. iii. (1814) 171 = Alyssum corymbosum?
gemonensis, Poir. Encyc. viii. 571 = Alyssum gemonense.
Geyeri, Hook. Lond. Journ. Bot. vi. (1847) 70. t. 5 = Physaria Geyeri.
glabrescens, *Boiss. Diagn.* Ser. II. v. 32.—Cilicia.
globosa, Desv. Journ. Bot. iii. (1814) 171, 184 = ludoviciana.
gnaphalodes, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 161.—Persia.
Gordoni, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 149.—Am. bor. occ.
gracilis, *Hook. Bot. Mag.* t. 3533.—Am. bor. occ.
graeca, *Reut. in Cat. Hort. Genèv.* (1858); *ex Boiss. Fl. Orient.* i. 262.—Ital.; Graec.
grandiflora, *Hook. l. c.* t. 3464.—Texas; Mexic.
incana, Desv. Journ. Bot. iii. (1814) 171 = Alyssum incanum.
Kingii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 353.—Calif.
lanuginosa, Poir. Encyc. viii. 572 = Clypeola eriocarpa.
lasiocarpa, *Hook. Bot. Mag.* sub t. 3464.—Texas.
Lescurii, A. Gray, Man. Bot. N. U. St. ed. II. 38 = Alyssum Lescurii.
Lindheimeri, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 145.—Texas.
ludoviciana, *DC. Syst.* ii. 297.—Am. bor. occ.
mendocina, *Phil. in Linnaea*, xxxiii. (1864–65) 12.—Chili.
microcarpa, Vis. Fl. Dalm. ii. t. 32. f. 2 = Alyssum gemonense.
montana, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864), 58.—Am. bor. occ.
montevidensis, Eichl. in Mart. Fl. Bras. xiii. 1. 302 = mendocina.
Newberryi, A. Gray, Colorado Expl. Surv. (1861) Bot. 6 = Physaria Newberryi.
Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 101.—Am. bor. occ.
occidentalis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 353.—Calif.
pallida, *Torr. & Gray, Fl. N. Am.* i. 668.—Am. bor.
paniculata, *Desv. Journ. Bot.* iii. (1814) 171.—Ins. Creta.
physiphora, *Andrz. ex DC. Syst.* ii. 300.—Hab.?
polyantha, Schlecht. in Bot. Zeit. xi. (1853) 620 = gracilis?
pulchella, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 15 = Engelmanni.
purpurea, *A. Gray, Pl. Wright.* ii. 14.—N. Mexic.
recurvata, *Engelm. in Bost. Journ. Nat. Hist.* vi. (1850) 147.—Am. bor. occ.
repanda, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 101.—Am. bor. occ.
reticulata, Lam. ex Poir. Encyc. viii. 572 = Coluteocarpus reticulatus.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1882) 320.—Mexic.
Shortii, *Torr. & Gray, Fl. N. Am.* i. 102.—Am. bor. occ.
sinuata, Cav. Desc. 404; Poir. Encyc. viii. 570 = Alyssum sinuatum.
stenophylla, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 149 = Fendleri.
utriculata, *DC. Fl. Fr.* iv. 696; *Poir. Encyc.* viii. 571.—Europ.
utriculata, Puccin. Syn. Fl. Lucens. 335, partim = graeca.
vestita, *Desv. Journ. Bot.* iii. (1814) 171.—Persia.

VESICULINA, Rafin. Fl. Tellur. iv. 109 (1836) = **Utricularia**, Linn. (Lentib.).
albiflora, Rafin. l. c. = U. albiflora?
compressa, Rafin. l. c. = U. compressa.
gibba, Rafin. l. c. = U. gibba.
graminifolia, Rafin. l. c. = U. cyanea.
purpurea, Rafin. l. c. = U. purpurea.
pygmea, Rafin. l. c. = U. pygmaea.
saccata, Rafin. l. c. = U. purpurea.
setacea, Rafin. l. c. = U. subulata.
tenella, Rafin. l. c. = Polypompholyx tenella.

VESLINGIA, Heist. Syst. 11 (1748); Fabric. Hort. Helmst. 363 (1759) = **Aizoon**, Linn. (Ficoid.).
cauliflora, Moench, Meth. Suppl. 299 = A. canariense.
Heisteri, Fabric. ex Willd. Sp. Pl. ii. 1053 = A. canariense.

VESLINGIA, Vis. in Nuov. Sagg. Acc. Sc. Padov. v. (1840) 269 = **Guizotia**, Cass. (Compos.).
scabra, Vis. l. c. = G. abyssinica.

VESPUCCIA, Parl. Nuov. Gen. e Spec. 55 (1854) = **Hydrocleys**, Rich. (Alismac.).
Humboldtii, Parl. l. c. 56 = H. Commersonii.

VESTIA, Willd. Enum. Hort. Berol. 208 (1809)
SOLANACEAE, Benth. & Hook. f. ii. 906.
foetida, Hoffmgg. Verz. Pfl. 119 = lycioides.
lycioides, *Willd. Enum. Hort. Berol.* 208.—Chili.

VETIVERIA, Thou. ex Virey, in Journ. Pharm. Sér. I. xiii. (1827) 499 = **Andropogon**, Linn. (Gramin.).
arundinacea, Griseb. Fl. Brit. W. Ind. 559 = A. squarrosus.
muricata, Griseb. l. c. 560 = A. squarrosus.
odorata, Virey, in Journ. Pharm. Sér. I. xiii. (1827) 499 = A. squarrosus.

VETRIX, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
Forbiana, Rafin. l. c. = S. purpurea.
Helix, Rafin. l. c. = S. purpurea.
Lambertiana, Rafin. l. c. = S. purpurea.
linearifolia, Rafin. l. c. 17 = S. purpurea?
purpurea, Rafin. l. c. 13 = S. purpurea.
sicula, Rafin. l. c. = S. purpurea?

VEXIBIA, Rafin. Neogenyt. 3 (1825) = **Valeriana**, Tourn.

VEXILLARIA, Hoffmgg. Verz. Pfl. 119 (1824); Benth. in Ann. Wien. Mus. ii. (1838) 117 = **Centrosema**, DC. (Legumin.).
arborea, Hoffmgg. ex Benth. l. c. 115 = Clitoria Hoffmanseggii.
brasiliana, Hoffmgg. Verz. Pfl. 119 = C. brasilianum.
glycinoides, Benth. in Ann. Wien. Mus. ii. (1838) 117 = Clitoria glycinoides.
grandiflora, Benth. l. c. = C. platycarpum.
laurifolia, Benth. l. c. = Clitoria cajanifolia.
rubiginosa, Benth. l. c. = Clitoria glycinoides.
triquetra, Hoffmgg. ex Benth. l. c. 122 = C. triquetrum (in Addend.).
villosa, Benth. l. c. 117 = Clitoria glycinoides.

VEXILLARIA, Rafin. in Am. Monthly Mag. (1818) 268 = **Clitoria**, Linn. (Legumin.).
Mariana, Rafin. l. c. = C. Mariana.

VIALIA, Vis. Ind. Sem. Hort. Patav. (1840).—Cf. Linnaea, xv. (1841) Litt. 103 = **Melhania**, Forsk. (Stercul.).
macrophylla, Vis. l. c. = M. macrophylla.

VIBEXIA, Rafin. Atl. Journ. 144 (1832) = **Sophora**, Linn. (Legumin.).
sericea, Rafin. l. c. = S. sericea.

VIBO, Medic. Phil. Bot. i. 178 (1789) = **Emex**, Neck. (Polygon.).
spinosa, Medic. l. c. = E. spinosa.

VIBONES, Rafin. Fl. Tellur. iii. 45 (1836) = **Rumex**, Tourn. (Polygon.).
auratus, Rafin. l. c. = R. sanguineus.

VIBORGIA, Moench, Meth. 132 (1794) = **Cytisus**, Linn. (Legumin.).
austriaca, Moench, l. c. 133 = C. austriacus.
capitata, Moench, Meth. Suppl. 43 = C. capitatus.
hirsuta, Moench, Meth. 132 = C. biflorus.
purpurea, Moench, Meth. Suppl. 43 = C. purpureus.
supina, Moench, Meth. 132 = C. supinus.

VIBORGIA, Spreng. Anleit. ii. II. 748 (1818) = **Wiborgia**, Thunb. &c.

VIBORGIA, Spreng. Nachr. I. Bot. Gart. Halle, 41 (1801) = WIBORGIA, Roth = **Galinsoga** Ruiz & Pav. (Compos.).

VIBORQUIA, Ortega, Hort. Matr. Dec. 66 (1798) = **Eysenhardtia**, H. B. & K. (Legumin.).
polystachya, Ortega, l. c. = E. amorphoides.

VIBURNUM, Linn. Syst. ed. I (1735); Gen. ed. I. 86 (1737). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 3.
IRAUPALOS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 87.
LENTAGO, Rafin. l. c. (1820) 87.
MICROTINUS, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 293.
OPULUS, Tourn. ex Linn. Gen. ed. I. 86 (1737).
OREINOTUS, Oerst. in Kjoeb. Vidensk. Meddel. (1860) 281.
SOLENOTINUS, Spach, Hist. Vég. Phan. viii. 305 (1839).
THYRSOSMA, Rafin. Sylva Tellur. 130 (1838).
TINUS, [Tourn.] Linn. Syst. ed. I (1735).
TYNUS, J. S. Presl, Rostl. ii. 93 (1823).
acerifolium, Bong. in Mém. Acad. Pétersb. ii. (1833) 144 = pauciflorum.
acerifolium, Gueldenst. ex Ledeb. Fl. Ross. ii. 385 = orientale.
acerifolium, *Linn. Sp. Pl.* 268.—Am. bor.
acuminatum, Hort. ex Steud. Nom. ed. I. 880 = dentatum.
acuminatum, Wall. Cat. n. 465; et ex DC. Prod. iv. 325 = punctatum.
acutifolium, *Benth. Pl. Hartw.* 59.—Mexic.
adenophyllum, Wall. ex Voigt, Hort. Suburb. Calc. 398, nomen (Quid?).—Hab.?
alnifolium, Marsh. Arbust. Am. 162 = lantanoides.
alpinum, Macfad. Fl. Jamaic. ii. 201 = glabratum.
alternifolium, *Zoll. & Mor. Syst. Verz. Zoll.* 59.—Java.
amblodes, Rafin. Alsog. Am. 55 (Quid?).—Am. bor.
americanum, Mill. Gard. Dict. ed. VIII. n. 8 = V. Opulus.
atro-cyaneum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 7.—Reg. Himal.
Awabuki, Hort. Berol. ex C. Koch, Wochenschr. x. (1867) 109 = odoratissimum.
ayavacense, *H. B. & K. Nov. Gen. et Sp.* iii. 428.—Peruv.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 268.—Japon.
burejaeticum, *Regel & Herd. in Gartenfl.* (1862) 407.—China.
burejanum, Herd. in Bull. Soc. Nat. Mosc. liii. (1878) I. 11 = praec.
capitatum, Rafin. Alsog. Am. 56 (Quid?).—Am. bor.
capitellatum, Wight & Arn. Prod. 388 = coriaceum.
cassinoides, *Linn. Sp. Pl.* ed. II. 384.—Am. bor.
cassinoides, Willd. Sp. Pl. i. 1491 = obovatum, *Walt.*
chinense, Hook. & Arn. Bot. Beech. Voy. 190 = Premna integrifolia.
chinense, Zeyh. ex Steud. Nom. ed. I. 880 = odoratissimum.
Colebrookeanum, *Wall. Cat.* n. 460; *et ex DC. Prod.* iv. 325.—Reg. Himal.
cordifolium, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 175, partim = furcatum.
cordifolium, *Wall. Cat.* n. 462; *et ex DC. Prod.* iv. 327.—Reg. Himal.
coriaceum, *Blume, Bijdr.* 656.—Reg. Himal.; Malaya.
corylifolium, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 174.—Reg. Himal.
costaricanum, *Hemsl. Biol. Centr. Am. Bot.* ii. 2.—Costa Rica.
cotinifolium, *D. Don, Prod. Fl. Nep.* 141.—Reg. Himal.
cuspidatum, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 171 = tomentosum.
cuspidatum. Thunb. Fl. Jap. 125 = Hydrangea cuspidata.
cylindricum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 142 = coriaceum.
davuricum, Maxim. in Mém. Sav. Etr. Pétersb. ix. (1859) 135 = burejaeticum.
davuricum, *Pall. Fl. Ross.* i. II. 30.—Dahuria.
delicatulum, *Vatke, in Bot. Zeit.* xxx. (1872) 718.—Montenegro.
densiflorum, *Chapm. Fl. S. U. St. Suppl.* 624.—Florida.

VIBURNUM:—
densum, *Benth. Pl. Hartw.* 59.—Mexic.
dentatum, *Linn. Sp. Pl.* 268.—Am. bor.
dentatum, Thunb. Fl. Jap. 122 = tomentosum.
dichotomum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 142 = Viscum articulatum.
dilatatum, *Thunb. Fl. Jap.* 124.—China; Japon.
discolor, *Benth. Pl. Hartw.* 83.—Guatemala.
divaricatum, *Benth. l. c.* 132.—Peruv.
dubium, *Steud. Nom.* ed. II. ii. 762.—Hab.?
edule, Rafin. in Med. Repos. N. York, II. v. (1808) 354; Pursh, Fl. Am. Sept. i. 203 = V. Opulus.
elatum, *Benth. Pl. Hartw.* 59.—Mexic.
elegans, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 36 = monogynum?
ellipticum, *Hook. Fl. Bor. Am.* i. 280.—Am. bor. occ.
erosum, A. Gray, in Perry, Jap. Exped. ii. 313 = dilatatum.
erosum, *Thunb. Fl. Jap.* 124.—China; Japon.
erubescens, *Wall. Cat.* nn. 459, 7474; *et ex DC. Prod.* iv. 329.—Reg. Himal.; Ind. or.
farinosum, Stokes, Bot. Mat. Med. ii. 168 = V. Lantana.
ferrugineum, Rafin. Alsog. Am. 52 (Quid?).—Am. bor.
foetens, *Decne. in Jacquem. Voy. Bot.* 75. t. 84.—Reg. Himal.
foetidum, *Wall. Cat.* n. 466; *Pl. As. Rar.* i. 49.—Reg. Himal.; Burma.
Forbesii, *Fawc. in Forbes, Wander. Natural.* 506.—Ins. Timor.
Fordiae, *Hance, in Journ. Bot.* xxi. (1883) 321.—China.
fragrans, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 107.—China.
fragrans, Loisel. Herb. Amat. vii. 466 = dubium.
furcatum, *Blume, ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 175, in syn.; *et ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 483.—China; Japon.
fuscum, *Hemsl. Biol. Centr. Am. Bot.* ii. 2.—Mexic.
glabratum, *H. B. & K. Nov. Gen. et Sp.* iii. 428.—Am. trop.
glabrum, Willd. ex Schult. Syst. vi. 639 = glabratum.
glandulosum, Salisb. Prod. 172 = V. Opulus.
glaucum, Rafin. Alsog. Am. 53 (Quid?).—Am. bor.
glomeratum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 483.—China.
grandiflorum, Wall. Cat. n. 464; et ex DC. Prod. iv. 329 = nervosum.
grandifolium, Sm. in Rees, Cyclop. xxxvii. n. 14 = lantanoides.
Griffithianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 4.—Reg. Himal.
Hanceanum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 487.—China.
Hartwegi, *Benth. Pl. Hartw.* 84.—Mexic.; Guatemala.
Hasseltii, *Miq. Fl. Ind. Bat.* ii. 123.—Java.
hebanthum, Thw. Enum. Pl. Zeyl. 136, partim = coriaceum.
hebanthum, *Wight & Arn. Prod.* 388.—Ind. or.
heterophylum, Rafin. Alsog. Am. 52 (Quid?).—Am. bor.
hirtum, Thunb. Fl. Jap. 124 = Hydrangea hirta.
hyemale, Salisb. Prod. 172 = V. Tinus.
integerrimum, Wall. Cat. n. 457; et ex DC. Prod. iv. 324 = sambucinum.
involucratum, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 5.—Florida.
involucratum, Wall. Cat. n. 458; et ex DC. Prod. iv. 327 = stellulatum.
Irabutha, Blume, ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 177 = odoratissimum.
Jacquemontii, *Planch. Hort. Donat.* 29.—Reg. Himal.
japonicum, Spreng. Syst. i. 934 = Cornus japonica.
Junghuhnii, *Miq. Fl. Ind. Bat.* ii. 123.—Java.
Kanbok, *Siebold, in Verh. Batav. Gen.* xii. (1830) 70.—Japon.
Keteleeri, Carr. in Rev. Hortic. (1863) 269 = macrocephalum.
laevigatum, Ait. Hort. Kew. ed. I. i. 371 = obovatum, *Walt.*
laevigatum, Willd. Berl. Baumz. 401 = nudum.
lanceolatum, Hill, Hort. Kew. 457; Veg. Syst. xiii. 64 = obovatum, *Walt.*
Lantana, *Linn. Sp. Pl.* 268.—Europ.; Reg. Cauc.; Afr. bor.
Lantana, Wall. ex D. Don, Prod. Fl. Nep. 141 = nervosum.
lantanoides, *Michx. Fl. Bor. Am.* i. 179.—Am. bor.
lantanoides. Miq. Ann. Mus. Bot. Lugd. Bat. i. 265 = furcatum.

VIBURNUM:—
lasiophyllum, *Benth. Pl. Hartw.* 189.—N. Granat.
latifolium, Hort. ex Schult. Syst. vi. 630 = rigidum.
laurifolium, C. Koch, Dendrol. ii. I. 63 = V. Tinus.
laurifolium, Rafin. Alsog. Am. 52 = nudum.
lauriforme, Lam. Fl. Fr. iii. 363 = V. Tinus.
lentaginum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = V. Lentago.
Lentago, Duroi, Harbk. Baumz. ii. 485 = prunifolium.
Lentago, *Linn. Sp. Pl.* 268.—Am. bor.
lentagoides, Rafin. in Med. Repos. N. York, II. v. (1808) 354 = V. Lentago.
lentum, S. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 136 = V. Lantana.
lobatum, Lam. Fl. Fr. iii. 363 = V. Opulus.
longifolium, Lodd. ex C. Koch, Dendrol. ii. I. 67 = pubescens.
longifolium, Rafin. Alsog. Am. 56 (Quid?).—Am. bor.
lucidum, Mill. Gard. Dict. ed. VIII. n. 5 = V. Tinus.
lutescens, *Blume, Bijdr.* 655.—Java.
lutescens, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 176 = Colebrookeanum.
luzonicum, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 310.—Ins. Philipp.
macrocephalum, *Fortune, in Journ. Hort. Soc.* ii. (1847) 244; *et in Lindl. Bot. Reg.* (1847) t. 43.—China.
macrodon, Rafin. Atl. Journ. 17 (Quid?).—Am. bor.
macrophyllum, *Thunb. Fl. Jap.* 125.—Japon.
macrophyllum, Van Hall, in Fl. Jard. Roy. Pays-Bas, ii. (1859) 97 = Buergeri.
maculatum, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 266 = V. Lantana.
medium, Hort. Belv. ex Steud. Nom. ed. I. 880. nomen (Quid?).—Hab.?
membranaceum, *Hemsl. in Biol. Centr. Am. Bot.* ii. 2.—Mexic.
membranaceum, Rafin. Alsog. Am. 51 (Quid?).—Am. bor.
microcarpum, *Schlecht. & Cham. in Linnaea*, v. (1830) 170.—Mexic.
microphyllum, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Mexic.
molle, *Michx. Fl. Bor. Am.* i. 180.—Am. bor.
monogynum, *Blume, Bijdr.* 655.—Java.
Mullaha, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 141 = stellulatum.
multratum ×, Hort. ex C. Koch, Dendrol. ii. I. 54.
nervosum, *D. Don, Prod. Fl. Nep.* 141.—Reg. Himal.
nervosum, Hook. & Arn. Bot. Beech. Voy. 190 = sempervirens.
nitidum, Ait. Hort. Kew. ed. I. i. 371 = nudum.
nudum, *Linn. Sp. Pl.* 268.—Am. bor.
obliquatum, Rafin. Alsog. Am. 54 (Quid?).—Am. bor.
obovatum, *Ruiz & Pav. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1860) 286 (err. typ. 386).—Peruv.
obovatum, *Walt. Fl. Carol.* 116.—Am. bor.
odoratissimum, *Ker-Gawl. in Bot. Reg.* t. 456.—Reg Himal.; China.
oleaefolium, Tausch, in Flora, xxi. (1838) 733 = nudum.
opuloides, Muhl. Cat. 32 = V. Opulus.
Opulus, *Linn. Sp. Pl.* 268.—Europ.; As. et Am. bor.
orientale, *Pall. Fl. Ross.* i. II. 31.—As. Min.
Oxycoccos, Pursh. Fl. Am. Sept. i. 203 = V. Opulus.
pallidum, Salisb. Prod. 172 = V. Lantana.
palustre, Rafin. Alsog. Am. 56 = V. Opulus?
parviflorum, *Mart. & Gal. in Bull. Acad. Brux.* xi. I. (1844) 243.—Mexic.
pauciflorum, *La Pylaie, ex Torr. & Gray, Fl. N. Am.* ii. 17.—Am. bor.
pauciflorum, Rafin. Alsog. Am. 58 (Quid?).—Am. bor.
phlebotrichum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 173.—Japon.
pinchinchense, *Benth. Pl. Hartw.* 188.—Ecuador.
pinnina, Rafin. Alsog. Am. 57 = V. Opulus.
plicatum, A. Gray, in Perry, Jap. Exped. ii. 313 = furcatum.
plicatum, Thunb. in Trans. Linn. Soc. ii. (1794) 332 = tomentosum.
polycarpum, Wall. Cat. n. 455; et ex DC. Prod. iv. 328 = cotinifolium.
premnaceum, Wall. ll. cc. 461; 325 = foetidum.
primina, Rafin. in Med. Repos. N. York, ii. v. (1808) 354 = V. Opulus.
prunifolium, *Linn. Sp. Pl.* 268.—Am. bor.
pubescens, *Pursh, Fl. Am. Sept.* i. 202.—Am. bor.
pubigerum, Wight & Arn. Prod. 389 = erubescens.

VIBURNUM:—
pubinerve, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 265 = phlebotrichum.
punctatum, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 142.—Reg. Himal.
punctatum, Rafin. ex DC. Prod. iv. 326 = cassinoides.
punicifolium, Desf. Hist. Arb. i. 345 = nudum.
pyrifolium, Poir. Encyc. viii. 653 = prunifolium.
pyrifolium, Pursh, Fl. Am. Sept. i. 201 = cassinoides.
Rafinesquianum, Schult. Syst. vi. 630 = pubescens.
repandum, Rafin. Alsog. Am. 53 (Quid ?).—Am. bor.
reticulatum, *Ruiz & Pav. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1860) 286 (err. typ. 386).—Peruv.
retusum, Rafin. Alsog. Am. 53 (Quid ?).—Am. bor.
rhombifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Mexic.
rigidum, *Vent. Jard. Malm.* 98.—Ins. Canar.
rosaceum, Hort. ex Steud. Nom. ed. II. ii. 762 = seq.
roseum, Steud. l. c. ed. I. 880 = V. Opulus.
rotundifolium, Rafin. Alsog. Am. 54 (Quid ?).—Am. bor.
rufidulum, Rafin. l. c. 56 (Quid ?).—Am. bor.
rugosum, Pers. Syn. i. 326 = rigidum.
sambucinum, *Reinw. ex Blume, Bijdr.* 656.—Malaya.
Sandankwa, *Hassk. Retzia*, i. 37.—China.
scabrellum, Chapm. Fl. S. U. St. 172 = molle.
scandens, Linn. f. Suppl. 184 = Hydrangea virens.
schensianum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 480.—China.
sempervirens, *C. Koch, Hort. Dendr.* 300; *Wochenschr.* x. (1867) 109.—China.
serratum, Thunb. Fl. Jap. 124 = Hydrangea hortensis.
setigerum, *Hance, in Journ. Bot.* xx. (1882) 261.—China.
sibiricum, Nois. ex Steud. Nom. ed. I. 880, nomen.—(Quid ?).—Sibir. ?
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 267.—Japon.
Simonsii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 177.—Reg. Himal.
sinensis, Zeyh. ex Colla, Hort. Ripul. 145 = odoratissimum.
squamatum, Muhl. ex Willd. Enum. Hort. Berol. 327 = cassinoides.
stellato-pilosum, *Polak. in Linnaea*, xli. (1877) 564.—Costa Rica.
stellato-tomentosum, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Costa Rica.
stellatum, *Hemsl. l. c.*—Mexic.
stellulatum, *Wall. Cat.* n. 463; *et ex DC. Prod.* iv. 327.—Reg. Himal.
stenocalyx, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Mexic.
strictum, Link, Enum. Hort. Berol. i. 288 = rigidum.
strictum, Sweet, Hort. Brit. ed. I. 195 = V. Tinus.
strigosum, Regel, Cat. Pl. Hort. Aksakov. 145, nomen (Quid ?).—Hab. ?
sulcatum, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Mexic.
sumatranum, *Miq. Fl. Ind. Bat. Suppl.* 537.—Sumatra.
sundaicum, Miq. Fl. Ind. Bat. ii. 121 = lutescens.
suspensum, *Lindl. in Journ. Hort. Soc.* viii. (1853) 130.—Hab. ?
tiliaefolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Mexic.
tinoides, *Linn. f. Suppl.* 184.—Am. austr.
tinoides, Mutis, ex H. B. & K. Nov. Gen. et Sp. iii. 428 = ayavacense ?
Tinus, *Linn. Sp. Pl.* 267.—Reg. Mediterr.
tomentosum, Hance, in Journ. Bot. viii. (1870) 273 = Hanceanum.
tomentosum, Lam. Fl. Fr. iii. 363 = V. Lantana.
tomentosum, Rafin. in Med. Repos. N. York, II. v. (1808) 354 = pubescens.
tomentosum, *Thunb. Fl. Jap.* 123.—China; Japon.
trilobum, Marsh. Arbust. Am. 162 = V. Opulus.
triphyllum, *Benth. Pl. Hartw.* 188.—N. Granat.
urceolatum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 172.—Japon.
venulosum, Benth. Fl. Hongk. 142 = sempervirens.
villosum, Rafin. in Med. Repos. N. York, II. v. (1808) 361 = pubescens.
villosum, *Sw. Prod. Veg. Ind. Occ.* 54.—Am. trop.
virens, Thunb. Fl. Jap. 123 = Hydrangea virens.
Wendlandi, *Hemsl. Biol. Centr. Am. Bot.* ii. 3.—Costa Rica.
Wightianum, Hohen. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 5 = punctatum.
Wightianum, Wall. Cat. n. 3729; Pl. As. Rar. ii. 29 = erubescens.

VIBURNUM:—
Wrightii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 267 = phlebotrichum.
zeylanicum, Gardn. ex Oerst. in Kjoeb. Vidensk. Meddel. (1860) 300 = coriaceum.
Zippelii, *Miq. Fl. Ind. Bat.* ii. 122.—N. Guin. ?

VICARYA, Stocks, in Proc. Linn. Soc. i. (1848) 367, nomen (Malvac.).

VICARYA, Wall. ex Voigt, Hort. Suburb. Calc. 544 (1845) = **Myriopteron**, Griff. (Asclep.).
cristata, Wall. l. c. = M. paniculatum.

VICATIA, DC. Prod. iv. 243 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 883.
coniifolia, *DC. l. c.*—Reg. Himal.
millefolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 671.—Reg. Himal.
Stewartii, *C. B. Clarke, l. c.*—Reg. Himal.

VICENTIA, Allem. Diss. cum tab. (1844).—Cf. Bot. Zeit. xii. (1854) 435 = **Terminalia**, Linn. (Combret.).
acuminata, Allem. l. c.—Bras.

VICIA, Tourn. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 524.
ABACOSA, Alef. in Bonplandia, ix. (1861) 102.
ATOSSA, Alef. l. c. (1861) 100.
COPPOLERIA, Todaro, in Atti Acc. Sc. Palermo, N. S. i. (1845) 14.
CRACCA, [Riv.] Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359.
CUJUNIA, Alef. in Bonplandia, ix. (1861) 101.
ENDIUSA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 359.
ERVILIA, Link, Enum. Hort. Berol. ii. 240 (1822).
ERVUM, Tourn. ex Linn. Gen. ed. I. 217 (1737).
FABA, Tourn. ex Linn. Syst. ed. I (1735).
HYPECHUSA, Alef. in Bot. Zeit. xviii. (1860) 164.
OROBELLA, Presl, in Weitenw. Beitr. ii. (1837) 27. t. 4.
PARALLOSA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 359.
RHYNCHIUM, Dulac, Fl. Hautes-Pyr. 290 (1767).
SELLUNIA, Alef. in Oestr. Bot. Zeitschr. ix. (1859) 358.
SWANTIA, Alef. l. c. (1859) 365.
TUAMINA, Alef. in Bonplandia, ix. (1861) 102.
VICILLA, Schur, Enum. Pl. Transs. 170 (1866).
VICIOIDES, Moench, Meth. 135 (1794).
WIGGERSIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 6, 33 (1801).
abbreviata, *Fisch. ex Spreng. Pugill.* i. 50.—Reg. Cauc.
abbreviata, C. A. Mey. Verz. Pfl. Cauc. 146 = cassubica.
abyssinica, Alef. in Bonplandia, ix. (1861) 72 = sativa.
acerosa, *Clos, in C. Gay, Fl. Chil.* ii. 137.—Chili.
acutifolia, *Ell. Sketch*, ii. 225.—Am. bor.
agrigentinum, Link, Handb. ii. 190 = V. Bivonea.
aintabensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 577.—Syria; Mesopot.
alba, Medic. in Vorles. Churpf. Phys. Gen. ii. (1787) 359 = sativa ?
alba, Moench, Meth. 148 = sativa.
albicans, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 33 = atropurpurea.
alpestris, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 93.—Reg. Cauc.; As. Min.
alpina, Pall. Reise, ii. 568 = multicaulis.
alternans, *Kit. in Linnaea*, xxxii. (1863) 631.—Banat.
alternifolia, Kit. l. c. = alternans.
altissima, *Desf. Fl. Atlant.* ii. 163.—Reg. Mediterr. occ.
ambigua, Guss. Fl. Sic. Prod. ii. 435 = varia.
americana, *Muhl. ex Willd. Sp. Pl.* iii. 1096.—Am. bor.
amoena, *Fisch. ex Ser. in DC. Prod.* ii. 355.—Sibir.
amphicarpa, Dorthes, in Journ. Phys. xxxv. (1789) 131 = sativa.
amphicarpos, Gouan, Herbor. 48 = Lathyrus setifolius.
andicola, H. B. & K. Nov. Gen. et Sp. vi. 498 = graminea.
andina, *Phil. in Anal. Univ. Chil.* (1872) 692.—Chili.
angulata, Bout. ex Willk. in Bot. Zeit. v. (1847) 429 = calcarata.
angulosa, Gater. Desc. Pl. Montaub. 129 = bithynica.

VICIA:—
angustifolia, *Clos, in C. Gay, Fl. Chil.* ii. 135.—Chili.
angustifolia, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 249 = lathyroides.
angustifolia, Linn. Amoen. Acad. iv. 105; Reich. Fl. Moeno-Franc. ii. 44 = sativa.
angustifolia, Moench, Meth. 148 = calcarata.
anisodonta, Boiss. & Huet, Diagn. Ser. II. ii. 41 = alpestris.
anomala, Boenn. Prod. Fl. Monast. 218 = sepium.
aphylla, C. Sm. in Buch, Beschr. Canar. Ins. 157 = Lathyrus aphyllus.
aquitanica, Clav. in Act. Linn. Soc. Bord. xxxvii. (1883) p. vi, 554 = atropurpurea.
araucana, *Phil. in Anal. Univ. Chil.* (1872) 693.—Chili.
arenivaga, Lamotte, Prod. Fl. Centr. Fr. i. 214 = lathyroides.
argentea, *Lapeyr. Hist. Abr. Pl. Pyr.* 417.—Hispan.
aristata, Dulac, Fl. Hautes-Pyr. 287 = V. Orobus.
armena, Boiss. Diagn. Ser. I. ii. 102 = persica.
articulata, *Hornem. Enum. Pl. Hort. Hafn.* (1807) 41; *Hort. Hafn.* ii. 690.—Hab. ?
articulata, Willd. Enum. Hort. Berol. 764 = monanthos.
assyriaca, *Boiss. Diagn.* Ser. I. ix. 123.—Armenia.
astragaloides, Willd. ex Ledeb. Fl. Ross. i. 637 = Astragalus striatellus.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 193.—Alger.
atropurpurea, *Desf. Fl. Atlant.* ii. 164.—Algeria.
atropurpurea, Steud. Nom. ed. II. ii. 763 = Nissoliana.
Aucheri, Jaub. & Spach, Illustr. i. 80. t. 41 = variegata.
aurantia, *Boiss. Fl. Orient.* ii. 578.—As. Min.
australis, Tenore, Fl. Nap. v. 112 = Gerardi.
azurea, C. Koch, in Linnaea, xix. (1847) 66 = alpestris.
azurea, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 191 = V. Cracca.
Bacla, *Moench, Meth.* 148.—Hab. ?
bactra, *Zuccagni, in Roem. Collect.* 151.—Hab. ?
baetica, *Lange, ex Willk. & Lange, Prod. Fl. Hisp.* iii. 306.—Hispan.
baetica, Schousb. ex Willk. l. c. 298 = lutea.
Balansae, *Boiss. Fl. Orient.* ii. 569.—Reg. Caucas.
Barbazitae, *Guss. & Tenore, Ind. Sem. Hort. Neap.* (1839) 12. n. 15.—*Cf. Ann. Sc. Nat.* Sér. II. xiii. (1840) 381.—Graecia; Sicil.
benghalensis, Hort. ex Link, Enum. Hort. Berol. ii. 239 = Nissoliana.
benghalensis, Linn. Sp. Pl. 736 = atropurpurea ?
Berteroana, *Phil. in Linnaea*, xxviii. (1856) 622.—Chili.
bicolor, *Willd. Enum. Hort. Berol. Suppl.* 51, nomen; *et ex Hornem. Suppl. Hort. Hafn.* 81.—Hab. ?
bidentata, Hook. Bot. Misc. ii. (1831) 215 = graminea.
Biebersteini, Guss. Fl. Sic. Syn. ii. 297 = pubescens.
Biebersteinii, Bess. ex Bieb. Fl. Taur. Cauc. iii. 472 = grandiflora.
Biebersteinii, C. A. Mey. Verz. Pl. Cauc. 147 = hyrcanica.
biennis, *Linn. Sp. Pl.* 736.—Sibir.
biflora, *Desf. Fl. Atlant.* ii. 166. t. 197.—Afr. bor.
biflora, C. A. Mey. Verz. Pfl. Cauc. 147 = calcarata.
biflora, Webb & Berth. Phyt. Canar. ii. 105. t. 65 B = atropurpurea.
bifoliolata, *Rodrig. in Bull. Soc. Bot. Fr.* xxv. (1878) 239.—Ins. Minorc.
bijuga, *Guill. ex Hook. Bot. Misc.* iii. (1833) 197.—Chili.
bipartita, *Moench, Meth.* 149.—Hab. ?
bithynica, *Linn. Syst.* ed. X. 1166.—Reg. Mediterr.
Bivonae, Ser. in DC. Prod. ii. 357 = V. Pseudo-cracca.
Bivonea, *Rafin. Préc. Découv.* 37.—Sicilia.
Bobartii, E. Forst. in Trans. Linn. Soc. xvi. (1830) 442 = sativa.
boetica, Fisch. ex Ser. in DC. Prod. ii. 365, nomen (Quid ?).—Hab. ?
boetica, Schousb. ex Nym. Consp. 209 = lutea.
bohemica, *F. F. Gmel. Syst.* 1110.—Bohem.
Boissieri, Heldr. & Sart. in Boiss. Diagn. Ser. II. ii. 40 = villosa.
branchitropis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 412.—Sibir. altaic.
Broteriana, Ser. in DC. Prod. ii. 357 = atropurpurea.
bythynica, Schrank, Baier. Fl. ii. 262 = bithynica.
caesarea, *Boiss. & Bal. Diagn.* Ser. II. vi. 69.—Cappadoc.
calcarata, *Desf. Fl. Atlant.* ii. 166.—Reg. Medit.; Persia.
campestris, Schur, Enum. Pl. Transs. 169 = sativa.

VICIA :—

canadensis, Zuccagni, in Roem. Collect. 151 = sativa.
canariensis, *Alef. in Bonplandia*, ix. (1861) 72.—Ins. Canar.
Candolleana, Tenore, Fl. Nap. v. 111 = Gerardi.
canescens, *Labill. Ic. Pl. Syr. Dec.* i. 17. t. 7.—Syria.
canescens, Sibth. & Sm. Fl. Graec. Prod. ii. 71 = Gerardi.
capensis, Berg. Desc. Pl. Cap. 215 = Lessertia pulchra.
cappadocica, *Boiss. Diagn.* Ser. II. vi. 68.—Cappadoc.
capreolata, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838), reimpr. 23.—Ins. Mader.
carnea, Kotschy, in Unger & Kotschy, Cypern, 386 = Michauxii.
caroliniana, *Walt. Fl. Carol.* 182.—Am. bor.
Cassia, *Boiss. Diagn.* Ser. I. ix. 119.—Syria; Ins. Cyprus.
cassubica, Bieb. Fl. Taur. Cauc. iii. 466 = truncatula.
cassubica, Lapeyr. Hist. Abr. Pl. Pyr. ii. 417 = Gerardi.
cassubica, *Linn. Sp. Pl.* 735.—Europ.
cassubica, Oeder, in Fl. Dan. t. 98 = V. Orobus.
chaetocalyx, *Webb & Berth. Phyt. Canar.* ii. 106.—Ins. Canar.
chilensis, D. Dietr. Syn. Pl. iv. 1107 = pallida, *Hook. & Arn.*
chlorantha, Heuff. ex Nym. Consp. 209 = truncatula.
ciceroidea, *Boiss. Diagn.* Ser. I. vi. 47.—Persia.
ciliaris, *Phil. in Linnaea*, xxviii. (1856) 621.—Chili.
ciliaris, *Sibth. & Sm. Fl. Graec. Prod.* ii. 71.—As. Min.
ciliata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 20 = lutea.
cilicia, Boiss. Diagn. Ser. I. ix. 122 = canescens.
cinerea, Bieb. Fl. Taur. Cauc. iii. 470 = calcarata.
cirrhosa, *C. Sm. ex Webb et Berth. Phyt. Canar.* ii. 103.—Ins. Canar.
Closiana, Phil. f. Cat. Pl. Vasc. Chil. 62 = angustifolia.
Clymenum, *S. G. Gmel. Reise*, iii. 306. t. 33. f. 1.—Sibir.
consentina, Reichb. ex Nym. Consp. 207 = V. Pseudocracca.
consentina, Spreng. Pugill. ii. 74 = variegata.
consobrina, *Pomel, Nouv. Mat. Fl. Atl.* 192.—Alger.
conspicua, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 22.—Ins. Mader.
cordata, Wulf. ex Hoppe, in Sturm, Dentschl. Fl. Heft 32 = sativa.
Coris, Thouin, ex Mart. Hort. Erlang. (1814) 156, nomen (Quid ?).—Hab. ?
cornigera, *Chaub. in St. Amans, Fl. Agen.* 294.—Gallia.
corsica, *Nym. Syll.* 310, in syn.; *Janka, in Termész. Füzetek*, ix. (1885) 137.—Ins. Corsic.
Cosentini, Guss. Fl. Sic. Prod. ii. 426 = sativa.
costata, *Ledeb. Fl. Alt.* iii. 346.—Sibir. altaic.
Cracca, Benth. in Royle, Illustr. Bot. Himal. 200 = tenuifolia.
Cracca, *Linn. Sp. Pl.* 735.—Europ.; As. et Am. bor.
craccoides, Rafin. in Am. Monthly Mag. (1819) 194 (*craceoides*) = caroliniana.
cretica, *Boiss. & Heldr. Diagn.* Ser. I. ix. 118.—Graecia.
cuneata, *Guss. Fl. Sic. Prod.* ii. 428.—Sicil.
cuspidata, *Boiss. Diagn.* Ser. I. ii. 104.—As. Min.; Syria.
cypria, *Kotschy, in Unger & Kotschy, Cypern*, 384.—Ins. Cyprus.
dasycarpa, Tenore, Viagg. Abruzz. (1829) 81 = varia.
davurica, Willd. ex Ledeb. Fl. Ross. i. 674 = megalotropis.
debilis, *Perez Lara, in Anal. Soc. Esp. Hist. Nat.* xi. (1882) 402.—Hispan.
debilis, Steud. Nom. ed. II. ii. 763 = graminea.
Dennesiana, *H. C. Wats. in Godman, Nat. Hist. Azor.* 154, 155.—Ins. Azor.
dentata, Fisch. ex Hornem. Suppl. Hort. Hafn. 80 = dumetorum.
dentata, *Gill. ex Hook. Bot. Misc.* iii. (1833) 197.—Reg. Argent.
depauperata, Clos, in C. Gay, Fl. Chil. ii. 132 = acerosa.
desertorum, Link, Enum. Hort. Berol. ii. 238 = dumetorum.
Diomedis, Gren. ex Nym. Consp. 210 = sativa.
disperma, *DC. Cat. Hort. Monsp.* (1813) 154.—Gall. austr.
disticha, Jacq. Enum. Pl. Carib. 27 (Quid ?).—Ind. occ.

VICIA :—

djimilensis, C. Koch, ex Boiss. Fl. Orient. ii. 569 = Balansae.
Douglasii, Torr. in Am. Journ. Sc. Ser. I. iv. (1822) 66 (*Douglassii*) = V. Cracca ?
Drymeja, Schur, Enum. Pl. Transs. 167 = sepium.
dubia, Schult. Obs. Bot. 149 = articulata.
dumetorum, Benth. in Royle, Illustr. Bot. Himal. 200 = rigidula.
dumetorum, *Linn. Sp. Pl.* 734.—Europ.; Sibir.
dumetorum, Thuill. Fl. Par. ed. II. 366 = sepium.
dumicola, Dulac, Fl. Hautes-Pyr. 288 = dumetorum.
Durandi, *Boiss. Diagn.* Ser. I. ix. 116.—Mauritan.
ecirrhosa, *Rupr. ex Boiss. Fl. Orient.* ii. 583.—Reg. Caucas.
elegans, Guss. Fl. Sic. Prod. ii. 438 = tenuifolia.
equina, Steud. Nom. ed. I. 881; Reichb. Fl. Germ. Excurs. 532 = V. Faba.
erecta, *Perez Lara, in Anal. Soc. Esp. Hist. Nat.* xi. (1882) 402.—Hispan.
erviformis, *Boiss. Voy. Bot. Espagne*, ii. 191.—Reg. Medit. occ.
erviformis, Winkl. ex Willk. & Lange, Prod. Fl. Hisp. iii. 306 = baetica.
Ervilia, *Willd. Sp. Pl.* iii. 1103.—Europ.; Afr. bor.
Ervilla, Medic. Beobacht. (1783) 179 = praec. ?
ervoides, Hampe, in Flora, xxv. (1842) 61 = pubescens.
esculenta, Salisb. Prod. 339 = V. Faba.
excisa, Schur, Enum. Pl. Transs. 169 = sativa.
exigua, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 272.—Am. bor. occ.
exigua, Torr. & Gray, in Pacif. Rail. Rep. ii. 162 = pulchella.
exstipulata, S. G. Gmel. Reise, iii. 307. t. 33. f. 2 = Lathyrus Aphaca.
Faba, *Linn. Sp. Pl.* 737.—Hab. ?
Fagonii, Lapeyr. Hist. Abr. Pl. Pyr. 419 = pyrenaica.
ferruginea, [Bess.], ex Mart. Hort. Erlang. (1814) 156, nomen (Quid ?).—Hab. ?
filicaulis, *Webb & Berth. Phyt. Canar.* ii. 102.—Ins. Canar.
fimbriata, Auch. ex Boiss. Fl. Orient. ii. 237 = Astragalus conduplicatus.
Fischeri, Steud. Nom. ed. II. ii. 763 = dumetorum.
flavida, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 20 = lutea.
Fleischeri, Hochst. & Steud. ex Boiss. Fl. Orient. ii. 592 = palaestina.
floridana, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 292.—Florida.
fodinarum, *Phil. in Anal. Univ. Chil.* (1872) 692.—Chili.
foetens, Gilib. Fl. Lituan. ii. 103 = sylvatica.
Fontanesiana, Tenore, ex Nym. Consp. 207 = Gerardi.
Forsteri, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 172 = sativa.
frutescens, Gilib. Fl. Lituan. ii. 104 = cassubica.
fulgens, *Battand. in Bull. Soc. Bot. Fr.* xxxii. (1885) 338.—Afr. bor.
galeata, *Boiss. Diagn.* Ser. I. ii. 103.—Syria.
galegifolia, Andr. Bot. Rep. t. 319 = Swainsona coronillaefolia.
gallobrovincialis, Poir. Encyc. Suppl. v. 471 = Gerardi.
Gayi, *Phil. in Linnaea*, xxviii. (1856) 623.—Chili.
gemella, *Crantz, Stirp. Austr.* ed. II. fasc. v. 389.—Europ.; Oriens.
geminiflora, *Trautv. in Act. Hort. Petrop.* iii. 1. (1875) 42.—Dahuria.
Gerardi, Jacq. Fl. Austr. iii. 16 = cassubica.
Gerardi, A. St. Hil. in Bull. Soc. Agric. Orléans, i. (1810) 281 (Not. 43) = V. Cracca.
Gerardi, *Vill. Hist. Pl. Dauph.* i. 256.—Europ. austr.
Gerardi, Willd. Fl. Berol. Prod. 239 = tenuifolia.
gigantea, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 93.—China.
gigantea, *Hook. Fl. Bor. Am.* i. 157.—Am. bor. occ.
glabra, Schleich. Cat. Pl. Helv. ed. IV. 37 = sativa.
glauca, *Presl, Delic. Prag.* 37.—Reg. Mediterr.
globosa, *Retz. Obs.* iii. 39.—Hab. ?
Graccha, Lepech. Reisen, iii. 42, sphalm. = V. Cracca.
gracilis, *Loisel. Fl. Gall.* ed. I. 460.—Europ.; Syria.
gracilis, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 34 = Loweanum.
gracilis, [Soland. in] Russ. Aleppo, ed. II. ii. 259 (Quid ?).—Syria.
graminea, *Sm. in Rees, Cyclop.* xxxvii. n. 27.—Chili; Bras.

VICIA :—

grandiflora, *Scop. Fl. Carn.* ed. II. ii. 65.—Europ. austr.; Oriens.
grandiflora, Sm. in Rees, Cyclop. xxxvii. n. 7 = Menziesii.
grata, *Phil. in Linnaea*, xxviii. (1856) 622.—Chili.
gregaria, *Boiss. & Heldr. Diagn.* Ser. I. ix. 121.—As. Min.
Griffithii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 178.—Ind. bor. occ.
Halleri, Crantz, Inst. ii. 30 = sylvatica ?
hedysaroides, Fisch. ex Ledeb. Fl. Ross i. 672 = amoena.
helvetica, Hort. Prag. ex Ser. in DC. Prod. ii. 361 = sativa.
heterophylla, Presl, Delic. Prag. 37 = sativa.
heterophylla, Reichb. Fl. Germ. Excurs. 531 = narbonensis.
Hexameri, Alef. in Bot. Zeit. xvii. (1859) 231 = caroliniana.
hirsuta, Fisch. ex Ser. in DC. Prod. ii. 365, nomen (Quid ?).—Hab. ?
hirsuta, *S. F. Gray, Nat. Arr. Brit. Pl.* ii. 614.—Europ.; As. et Afr. bor.
hirsutissima, *Cyrilli, ex Mart. Hort. Erlang.* (1814) 154, nomen; *et ex Tenore, Fl. Nap.* v. 119.—Ital.
hirta, Balb. ex Pers. Syn. ii. 308 = lutea.
hirta, Tenore, Fl. Med. i. 578 = hirsutissima.
Hookeri, *G. Don, Gen. Syst.* ii. 319.—Chili.
Hookeriana, Presl, Repert. Bot. Syst. i. 8, in nota = Hookeri.
Hookeriana, Walp. Rep. i. 715 = gigantea, *Hook.*
humilis, *H. B. & K. Nov. Gen. et Sp.* vi. 498.—Mexic.
hungarica, *Heuff. in Oestr. Bot. Zeitschr.* viii. (1858) 26.—Hungar.
hybrida, Georgi, Beschr. Russ. Reich. III. iv. 1169 = pannonica.
hybrida, Huds. Fl. Angl. ed. II. 319 = lutea.
hybrida, *Linn. Sp. Pl.* 737.—Europ.; Syria; Afr. bor.
hypoleuca, Boiss. Diagn. Ser. II. v. 92 = alpestris.
hyrcanica, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 53.—Reg. Caucas.
imbricata, Gilib. Fl. Lituan. ii. 104 = V. Cracca.
incana, Lam. Fl. Fr. ii. 560 = atropurpurea.
incana, Thuill. Fl. Par. ed. II. 367 = V. Cracca.
incana, Vill. Hist. Pl. Dauph. iii. 449 = Gerardi.
incisa, Bieb. Fl. Taur. Cauc. iii. 471 = sativa.
incisa, Orph. ex Boiss. Fl. Orient. ii. 574 = V. Barbazitae.
inconspicua, *Phil. in Linnaea*, xxviii. (1856) 623.—Chili.
incurva, Roth, Beytr. ii. 98 = Lathyrus incurvus.
intermedia, Viv. Fl. Lib. Spec. 42. t. 19. f. 1 = sativa.
iranica, *Boiss. Fl. Orient.* ii. 581.—Persia.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 385 = pallida, *Turcz.*
Jauberti, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 71 = hyrcanica.
jeniseensis, Willd. ex Ledeb. Fl. Ross. i. 672 = amoena.
Keatingi, Steud. Nom. ed. II. ii. 764 = americana.
Kingii, Hook. f. Fl. Antarct. ii. 258 = graminea.
Kitaibeliana, Schur, Enum. Pl. Transs. 168 = grandiflora.
Kitaibelii, Reichb. ex Nym. Consp. 206 = V. Cracca.
kokanica, *Regel & Schmalh. in Regel, Pl. Nov. Fedsch.* 20.—Turkestan.
Kotschyana, *Boiss. Diagn.* Ser. I. vi. 47.—Persia.
laeta, Cesati, in Friderichst. Reise Griech. 280 = V. Barbazitae.
laevigata, Sm. Engl. Bot. t. 483 = lutea.
Lagopus, *Pomel, Nouv. Mat. Fl. Atl.* 193.—Alger.
lanceolata, *Phil. in Linnaea*, xxxiii. (1864–65) 51.—Chili.
lanciformis, Lange, in Kjoeb. Vidensk. Meddel. (1865) 183 = sativa.
lathyroides, All. Fl. Pedem. i. 326 = sativa.
lathyroides, *Linn. Sp. Pl.* 736.—Europ.; As. Min.; Afr. austr.
latifolia, Moench, Meth. 149 = narbonensis.
laxiflora, Boiss. Fl. Orient. ii. 586 = tenuifolia.
laxiflora, Brot. Phyt. Lusit. i. 125. t. 52 = gracilis.
Leavenworthii, *Torr. & Gray, Fl. N. Am.* i. 271.—Am. bor. occ.

VICIA :—
Lens, Coss. & Germ. Fl. Par. ed. I. 143 = Lens esculenta.
Lenticula, Janka, in Termész. Füzetek, ix. (1885) 138 = Lens Lenticula.
lentiformis, Schur, Enum. Pl. Transs. 168 = lutea.
leptocarpa, *Bianca, in Atti Acc. Gioen. Sc. Nat.* Ser. II. xiv. (1858) 51.—Sicil.
leptophylla, *Rafin. Caratteri*, 71.—Sicil.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 168 = lutea.
leucantha, Bivona, Stirp. Rar. Sicil. i. [9] = V. Bivonea.
leucophaea, *Greene, in Coult. Bot. Gaz.* vi. (1881) 217.—N. Mexic.
leucosperma, Moench, Meth. 148 = sativa.
Leyboldi, *Phil. in Linnaea*, xxxiii. (1864–65) 51.—Chili.
lilacina, *Ledeb. Fl. Alt.* iii. 348.—Sibir. altaic.
linearifolia, *Hook. & Arn. Bot. Beech. Voy.* 20.—Chili.
lineata, Bieb. Fl. Taur. Cauc. iii. 473 = pannonica.
littoralis, Jacq. Enum. Pl. Carib. 27 = Tephrosia cinerea.
littoralis, Salzm. in Flora, iv. (1821) 110 = V. Pseudo-cracca.
littoralis, Tenore, Fl. Nap. iv. 103 = varia.
Loiseleurii, Boiss. Fl. Orient. ii. 595, sphalm. = Meyeri.
longifolia, *Poir. Encyc.* viii. 557.—Syria.
longifolia, Schur, Enum. Pl. Transs. 169 = sativa.
Loweana, *Steud. Nom.* ed. II. ii. 764.—Ins. Mader.
ludoviciana, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 271.—Am. bor.
luganensis, Schleich. Cat. Pl. Helv. ed. III. 31 = sativa.
lunata, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 594.—Lydia.
lutea, *Linn. Sp. Pl.* 736.—Reg. Medit.; Oriens.
lutea, Pall. ex Bieb. Fl. Taur. Cauc. ii. 162 = grandiflora.
lutescens, Muhl. Cat. 68 = acutifolia.
Macraei, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833). 195.—Chili.
macrocarpa, *Bertol. Fl. Ital.* vii. 511.—Europ. austr.
maculata, *Pomel, Nouv. Mat. Fl. Atl.* 191.—Alger.
maculata, Presl, Fl. Sic. p. xxiii = sativa.
magellanica, *Hook. f. Fl. Antarct.* ii. 257.—Reg. Magellan.
magnifolia, *Clos, in C. Gay, Fl. Chil.* ii. 135.—Chili.
Martini, Gren. ex Nym. Consp. 210 = sativa.
Matthewsii, *A. Gray, Bot. U. St. Expl. Exped.* i. 419.—Peruv.
media, *Host, Fl. Austr.* ii. 335.—Europ.
megalosperma, Bieb. Fl. Taur. Cauc. ii. 161 = peregrina.
megalotropis, *Ledeb. Fl. Alt.* iii. 344.—Sibir. altaic.
melanops, *Sibth. & Sm. Fl. Grace. Prod.* ii. 72.—Europ. austr.
Menziesii, *Spreng. Syst.* iii. 267.—Ins. Sandvic.
mexicana, *Hemsl. Diagn. Pl. Nov.* 47.—Mexic.
Meyeri, *Boiss. Fl. Orient.* ii. 595.—Reg. Casp.
Michauxii, Schrank, ex Steud. Nom. ed. I. 882, nomen (Quid?).—Am. bor.?
Michauxii, *Spreng. Fl. Hal. Mant.* 48.—Syria; Persia.
Michelli, Desv. Journ. Bot. iv. (1814) 269, sphalm. = *Mitchelli*.
micrantha, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 197 = Hookeri.
micrantha, Lowe, in Trans. Camb. Phil. Soc. iv. (1833) 33 = Loweana.
micrantha, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 271.—Am. bor. occ.
microphylla, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 343.—Graecia.
militans, Crantz, Stirp. Austr. ed. II. fasc. v. 388 = cassubica.
minima, Gilib. Fl. Lituan. ii. 106 = lathyroides.
minuta, Moench, Meth. Suppl. 50 = bithynica.
minutiflora, D. Dietr. Syn. Pl. iv. 1107 = micrantha.
Mitchelli, Rafin. Préc. Découv. 37 = hirsuta.
modesta, *Phil. in Linnaea*, xxxiii. (1864–65) 49.—Chili.
mollis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 177.—Reg. Himal.
mollis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 576.—Syria.
monadelpha, Roth, Catalecta, fasc. ii. 97 = narbonensis.

VICIA :—
monantha, Retz. Obs. iii. 39 = calcarata.
monanthos, *Desf. Fl. Atlant.* ii. 165.—Reg. Mediterr.
monanthos, Ucria, in Roem. Arch. i. 1. (1796) 69 (Quid?).—Sicil.?
monanthos, Viv. Fl. Lib. Spec. 42 = peregrina.
Monardi, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 131.—Alger.
monosperma, C. Koch, in Linnaea, xv. (1841) 721 = cassubica.
montana, Froel. ex Koch, Syn. Fl. Germ. ed. II. 215 = sepium.
Montbreti, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* x. 59.—Mesopot.
montevidensis, Benth. in Mart. Fl. Bras. xv. 1. 110 = obscura.
montevidensis, Vog. in Linnaea, xiii. (1839) 34 = dentata.
Morisiana, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 172, in obs. = macrocarpa.
mucronata, *Clos, in C. Gay, Fl. Chil.* ii. 139.—Chili.
multicaulis, *Ledeb. Fl. Alt.* iii. 345.—Sibir. altaic.
multicaulis, Wallr. in Linnaea, xiv. (1840) 625 (Beitr. Fl. Hercyn. 255) = lathyroides.
multifida, Wallr. Ann. Bot. 85 = monanthos.
multiflora, Lam. Fl. Fr. ii. 560 = V. Cracca, Gerardi.
multiflora, Pollich, Hist. Pl. Palat. ii. 307 = cassubica.
Musquineæ, Bosc, ex Steud. Nom. ed. I. 882, nomen (Quid?).—Europ.
nana, Kellogg, in Proc. Calif. Acad. vii. 1. (1877) 89 = Lathyrus nevadensis.
nana, *Vog. in Linnaea*, xiii. (1839) 37.—Reg. Argent.
narbonensis, *Linn. Sp. Pl.* 737.—Reg. Mediterr.; Oriens.
nebrodensis, Huet, ex Nym. Consp. 210 = cuneata.
nemoralis, Steud. Nom. ed. I. 882 = sativa.
nigra, Steud. l. c. = sativa.
nigricans, *Hook. & Arn. Bot. Beech. Voy.* 20.—Chili.
Nissoliana, Gouan, Herbor. 51 (excl syn.) = pannonica.
Nissoliana, *Linn. Sp. Pl.* 735.—Europ.
nodosa, Fisch. ex Ser. in DC. Prod. ii. 365, nomen (Quid?).—Hab.?
nodosa, Gaertn. Fruct. ii. t. 151 = globosa?
Noëana, *Boiss. & Reut. ex Boiss. Fl. Orient.* ii. 572.—Syria.
Nordmanniana, Ower. ex Trautv. in Act. Hort. Petrop. viii. (1883) 251 = Balansae?
notata, Gilib. Fl. Lituan. ii. 105 = sativa.
obcordata, Reichb. Fl. Germ. Excurs. 530 = sativa.
obscura, *Vog. in Linnaea*, xiii. (1839) 36.—Bras.
occulta, Lehm. ex G. Don, in Sweet, Hort. Brit. ed. III. 184, nomen (Quid?).—Europ.
ochroleuca, Gilib. Fl. Lituan. ii. 102 = pisiformis.
ochroleuca, *Tenore, Prod. Fl. Nap.* p. xlii.—Italia.
olbiensis, Reut. ex Timb. in Bull. Soc. Bot. Fr. xiii. (1866) p. cli = cuneata.
onobrychioides, *Linn. Sp. Pl.* 735.—Reg. Mediterr.
onusta, Wallr. in Linnaea, xiv. (1840) 623 (Beitr. Fl. Hercyn. 253) = tenuifolia.
oregana, Nutt. in Torr. & Gray, Fl. N. Am. i. 270 = americana.
oroboides, *Wulf. in Jacq. Coll.* iv. 323.—Europ. austr.
Orobus, *DC. Fl. Fr. Suppl.* 577.—Europ.
oxyphylla, Schur, Enum. Pl. Transs. 167 = sepium.
palaestina, *Boiss. Diagn. Ser.* I. ix. 116.—Syria; Mesopot.
pallens, *Reichb. Fl. Germ. Excurs.* 529 (Quid?).—Helvet.
pallida, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 196.—Chili.
pallida, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 178 = sativa.
pallida, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 789.—Dahuria; Reg. Himal.
pannonica, *Crantz. Stirp. Austr.* ed. II. fasc. v. 393.—Europ. austr.; Oriens.
paposana, *Phil. Fl. Atac.* 15.—Chili.
parviflora, *Cav. in Anal. Cienc. Nat.* iv. (1801) 73.—Marocc.
parviflora, Hook. & Arn. Bot. Beech. Voy. 20 = Hookeri.
parviflora, Lapeyr. Hist. Abr. Pl. Pyr. 418 = hirsuta.
parviflora, Loisel. Fl. Gall. ed. I. 460 = disperma.
parviflora, Michx. Fl. Bor. Am. ii. 69 = caroliniana.
patagonica, *Hook. f. Fl. Antarct.* ii. 259.—Patagon.
patula, Moench, Meth. 147 = dumetorum.

VICIA :—
pauciflora, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 196 = graminea.
pauciflora, *Presl, Fl. Sic.* p. xxii.—Sicil.
paucifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 173.—Afr. trop.
pectinata, *Lowe, Man. Fl. Mad.* i. 205.—Ins. Mader.
pedunculariis, *Rojas Clem. Pl. Espont. Titag.* 16.—Hispan.
pedunculata, Gueldenst. Reisen, i. 195; ex Ledeb. Fl. Ross. i. 679 (Quid?).—Rossia.
pedunculata, *Peyr. in Linnaea*, xxx. (1859–60) 80.—Mexic.
pedunculata, *Shuttlew. ex Roux, in Bull. Soc. Bot. et Hort. Prov.* iii. (1881) 124.—Hab.?
pellucida, Jacq. Hort. Schoenb. ii. 50 = Lessertia falciformis.
peregrina, *Linn. Sp. Pl.* 737.—Reg. Mediterr.; Oriens; Ind. bor. occ.
perennis, DC. Cat. Hort. Monsp. (1813) 155 = atropurpurea.
persepolitana, Boiss. Diagn. Ser. I. vi. 48 = Michauxii.
persica, *Boiss. l. c.* ii. 103.—As. Min.
picta, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 41.—Armenia.
pilisiensis, Aschers. & Janka, in Aschers. & Kanitz, Cat. Corm. Serb. 105 (= *Orobus ochroleucus*).—Hungar.
pilosa, Bieb. Fl. Taur. Cauc. ii. 161 = sativa.
pimpinelloides, Mauri, Pl. Rom. Cent. xiii. 35 = sativa.
Pinardi, *Boiss. Diagn.* Ser. I. ix. 120.—As. Min.
pinetorum, *Boiss. & Sprun. l. c.* ii. 104.—Graecia.
Piscidia, Forst. ex DC. Prod. ii. 379 (= *Orobus Piscidia*).—N. Caled.
pisiformis, *Linn. Sp. Pl.* 734.—Europ.
platycarpos, *Roth, Bot. Abh.* 10. t. 1.—Hab.?
plumosa, Martr. Fl. Tarn, 179 = varia.
podocarpa, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 571 = sericocarpa.
polymorpha, Godr. Fl. Lorr. ed. I. i. 179 = sativa.
polyphylla, Desf. Fl. Atlant. ii. 162 = tenuifolia.
polyphylla, Jan, ex Nym. Consp. 207 = Gerardi.
polyphylla, Moç. & Sesse, ex G. Don, Gen. Syst. ii. 318 = Sessei.
polyphylla, Reichb. Fl. Germ. Excurs. 527 = varia.
polyphylla, Sibth. & Sm. Fl. Graec. vii. 86. t. 699 = Sibthorpii.
polyphylla, Waldst. & Kit. Pl. Rar. Hung. iii. 282. t. 254 = villosa.
polysperma, Tenore, Fl. Neap. Prod. App. v. 22 = altissima.
pontica, Willd. Sp. Pl. iii. 1094 = V. Cracca.
porphyrea, *Reichb. Fl. Germ. Excurs.* 532, in obs.—Hab.?
pratensis, Wallr. in Linnaea, xiv. (1840) 626 (Beitr. Fl. Hercyn. 256) = sepium?
Pseudo-cracca, *Bertol. Rar. Ital. Pl. Dec.* iii. 58.—Italia.
Pseudo-cracca, Fritze, ex Willk. & Lange, Prod. Fl. Hisp. iii. 306 = baetica.
Pseudo-cracca, Mérat, Fl. Par. ed. III. t. 2 = varia.
Pseudo-lutea, *Schur, in Verh. Naturf. Ver. Bruenn*, xv. 11. (1877) 189.—Europ.
Pseudo-orobus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 41.—Dahuria.
Pseudo-sepium, *Nym. in Linnaea*, xviii. (1844) 648.—Sicil.
Pseudo-villosa, Schur, Enum. Pl. Transs. 166 = varia.
pubescens, *Link, Handb.* ii. 190; *Boiss. Fl. Orient.* ii. 596.—Reg. Mediterr.
pulchella, *H. B. & K. Nov. Gen. et Sp.* vi. 499.—Am. bor. occ.
punctata, Schleich. ex Pers. Syn. ii. 307 = onobrychioides.
purpurascens, DC. Cat. Hort. Monsp. (1813) 155 = pannonica.
purpurea, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 93 = alpestris.
pusilla, Muhl. ex Willd. Sp. Pl. iii. 1106 = gemella.
pygmaea, Link, Handb. ii. 191 = sativa.
pyrenaica, *Pourr. in Mém. Acad. Toul.* iii. (1788) 333.—Mont. Pyren.
quinquenervia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 50.—Japon.
Rapunculus, Debeaux, in Act. Linn. Soc. Bord. xxxi. (1877) 144 = amoena.
repens, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 344.—Ins. Cyclad.

VICIA:—
remrevillensis, Hussenot, Chard. Nanc. 98; ex Godr. Fl. Lorr. ed. II. i. 196 = sativa.
retusa, *Steud. Nom.* ed. II. ii. 764.—Chili.
Reuteriana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 73 = villosa.
Reverchoni, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 291.—Texas.
rigida, Herbich, Stirp. Rar. Bucov. 50 = cassubica.
rigidula, *Royle, Illustr. Bot. Himal.* 200.—Reg. Himal.
Rosani, *Tenore, Fl. Nap.* v. 115.—Ital.
rotundifolia, Gilib. Fl. Lituan. ii. 106 = sepium.
salaminia, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. ii. 39.—Graecia.
Salisii, J. Gay, ex Nym. Consp. 212 = pubescens.
Sallei, Timb. in Bull. Soc. Bot. Fr. xiii. (1866) p. cxlix = cuneata.
sativa, Guss. Fl. Sic. Prod. ii. 425 = macrocarpa.
sativa, *Linn. Sp. Pl.* 736.—Europ.; Afr. bor.; Oriens.
scepusiensis, *Kit. in Linnaea*, xxxii. (1863) 629.—Europ.
segetalis, Thuill. Fl. Par. ed. II. 367 = sativa.
Seguenzae, Huet, ex Nym. Consp. 208 = sicula.
Selloi, Vog. in Linnaea, xiii. (1839) 33 = graminea.
semiglabra, *Rupr. ex Boiss. Fl. Orient.* ii. 583.—Reg. Caucas.
separia, Dulac, Fl. Hautes-Pyr. 287 = sepium.
sepium, *Linn. Sp. Pl.* 737.—Europ.; As. bor.; Reg. Himal.
sericea, Moench, Meth. 147 = Nissoliana.
sericocarpa, *Fenzl, Pugill. Pl. Nov. Syr.* 4.—As. Min.; Persia.
serrata, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 80 = grandiflora.
serratifolia, Jacq. Fl. Austr. v. 30 = narbonensis.
Sessei, *G. Don, Gen. Syst.* ii. 318.—Mexic.
sessiliflora, *Clos, in C. Gay. Fl. Chil.* ii. 129.—Chili.
setifolia, H. B. & K. Nov. Gen. et Sp. vi. 500 = graminea.
Sibthorpii, *Boiss. Diagn.* Ser. I. ix. 122.—Graecia.
sicula, *Guss. Fl. Sic. Syn.* ii. 292.—Ins. Sicil.
singarensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 594.—Mesopot.
sitchensis, Bong. in Mém. Acad. Pétersb. ii. (1833) 129 = gigantea, *Hook.*
smyrnaea, *Boiss. Diagn.* Ser. II. ii. 38.—As. Min.
Solisi, *Phil. in Anal. Univ. Chil.* (1865) II. 326.—Chili.
sordida, Salisb. Prod. 339 = sepium.
sordida, Waldst. & Kit. Pl. Rar. Hung. ii. 133 = grandiflora.
sparsiflora, *Tenore, Fl. Nap.* v. 110.—Ital.
sparsifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 270 = americana.
Spruneri, Boiss. Diagn. Ser. I. ix. 119 = cretica.
spuria, Rafin. Caratteri, 72 = hybrida.
Stabiana, Tenore, Fl. Nap. v. 112 = Gerardi.
stenophylla, *Bianca, in Atti Acc. Gioen. Sc. Nat.* Ser. II. xiv. (1858) 56.—Sicil.
stenophylla, Schur, Enum. Pl. Transs. 169 = sativa.
stenophylla, *Vog. in Linnaea*, xiii. (1839) 35.—Reg. Argent.
stipulacea, Pursh, Fl. Am. Sept. ii. 739 = Lathyrus polymorphus.
striata, Bieb. Fl. Taur. Cauc. ii. 162 = pannonica.
subrotunda, Schur, Enum. Pl. Transs. 167 = sepium.
subserrata, *Phil. in Linnaea*, xxviii. (1856) 624.—Chili.
subulata, Moç. & Sesse, ex Steud. Nom. ed. II. ii. 765, nomen (Quid?).—Mexic.
subvillosa, *Boiss. Fl. Orient.* ii. 580.—Persia.
sylvatica, Benth. in Royle, Illustr. Bot. Himal. 200 = pallida, *Turcz.*
sylvatica, *Linn. Sp. Pl.* 734.—Europ.
sylvatica, Nutt. Gen. Am. ii. 97 = americana.
syriaca, Medic. Beobacht. (1783) 178 = seq.?
syriaca, Weinm. Hort. Dorp. (1810) 162 = calcarata.
syrtica, *Duby, Bot. Gall.* ed. II. i. 151.—Hispan.
Talpa, Raym. ex Poir. Encyc. viii. 563 = pyrenaica.
Tanakae, Franch. & Sav. Enum. Pl. Jap. i. 103; ii. 326 = V. Pseudo-orobus.
tenera, *R. Grah. in Wall. Cat.* n. 5928.—Reg. Himal.
Tenoreana, Martr. Pl. Crit. Tarn. 20 = gracilis, *Loisel.*
tenuifolia, *Roth, Tent. Fl. Germ.* i. 309.—Europ.; Oriens; As. bor.
tenuifolia, Tenore, Fl. Nap. v. 113 = varia.
tephrosioides, *Vog. in Linnaea*, xiii. (1839) 32.—Reg. Argent.

VICIA:—
Terronii, Steud. Nom. ed. II. ii. 765, sphalm. = hirsuta.
tetragona, *Desf. Cat. Hort. Par.* ed. III. 410.—Hab.?
tetrasperma, Moench, Meth. 148 = gemella.
tetrasperma, Moris, Fl. Sard. i. 567 = pubescens.
thessala, Sprun. ex Boiss. Fl. Orient. ii. 574 = V. Barbazitae.
Thouini, Mart. Hort. Erlang. (1814) 156, nomen (Quid?).—Hab.?
Timbali, Loret, in Loret & Barr. Fl. Monsp. 187, 804 = cuneata.
tomentosa, Hort. ex Steud. Nom. ed. I. 883; ed. II. ii. 765 = Coursetia tomentosa.
torulosa, Boiss. Fl. Orient. ii. 589 = caesarea.
torulosa, Desf. Tabl. ed. I. 196, nomen (Quid?).—Hab.?
torulosa, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 173 = sativa.
trichocalyx, Moris, Stirp. Sard. Elench. Fasc. iii. 7 = atropurpurea.
trichomera, Alef. in Bonplandia, ix. (1861) 70 = truncatula.
tricolor, Schur, Enum. Pl. Transs. 167 = sepium.
tricolor, Sebast. & Mauri, Fl. Rom. Prod. 245. t. 4 = melanops.
tricuspidata, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 158.—Tauria.
tridentata, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 93.—China.
tridentata, Gater. Desc. Pl. Montaub. 129 = lutea.
tridentata, Schwein. in Long, Exped. Winnep. ii. 392 = americana.
trifida, D. Dietr. Syn. Pl. iv. 1112 = americana.
triflora, Sieber, in Ind. Sem. Hort. Pesth. (1825); ex Tenore, Syll. Fl. Neap. 357 (Quid?).—Hab.?
triflora, Tenore, Prod. Fl. Nap. p. xlii = calcarata.
trijuga, *A. Br. & Bouché, in Ind. Sem. Hort. Berol.* (1853) *App.* 13.—Hab.?
truncata, Nutt. in Torr. & Gray, Fl. N. Am. i. 270 = americana.
truncata, *Phil. in Linnaea*, xxxiii. (1864–65) 50.—Chili.
truncatula, *Fisch. Cat. Hort. Gorenk.* ed. II. (1812) 72; *Bieb. Fl. Taur. Cauc.* iii. 473.—Bannat.; Reg. Caucas.; Persia.
ucrainica, *Andrz. ex Trautv. in Act. Hort. Petrop.* iii. 1. (1874) 44 (sp. dub).—Ross.
uncinata, Desf. ex Nym. Consp. 210 = sativa.
uncinata, Reichb. Fl. Germ. Excurs. 530 = pannonica.
unguiculata, *Clav. in Act. Linn. Soc. Bord.* xxxvii. (1883) p. vi, 552.—Gallia.
unijuga, *A. Br. in Ind. Sem. Hort. Berol.* (1853) *App.* 12.—Sibir.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 681.—Chili.
varia, *Host, Fl. Austr.* ii. 332.—Europ.; Oriens; Afr. bor.
variegata, Gilib. Fl. Lituan. ii. 103 = dumetorum.
variegata, Sibth. & Sm. Fl. Graec. Prod. ii. 69 = villosa.
variegata, *Willd. Sp. Pl.* iii. 1096.—As. Min.; Persia.
venosa, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 395.—Sibir.
venulosa, *Boiss. & Hohen. Diagn.* Ser. I. ix. 123.—Persia.
versicolor, Salisb. Prod. 339 = V. Cracca.
vestita, Boiss. Elench. 39 = lutea.
vestita, *Clos, in C. Gay, Fl. Chil.* ii. 134.—Chili.
vicina, *Clos, l. c.* 138.—Chili.
villosa, Brot. Fl. Lusit. ii. 150 = atropurpurea.
villosa, *Roth, Tent. Fl. Germ.* ii. II. 182.—Europ.; Oriens.
violacea, Dulac, Fl. Hautes-Pyr. 287 = onobrychioides.
virginica, Mart. Hort. Erlang. (1814) 156, nomen (Quid?).—Hab.?
vulcanica, Huet, ex Nym. Consp. 207 = V. Pseudo-cracca.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 617 = V. Faba.
vulgaris, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 378 = sativa.

VICILLA, Schur, Enum. Pl. Transs. 170 (1866) = **Vicia**, Tourn. (Legumin.).
cassubica, Schur, l. c. = Vicia cassubica.
Orobus, Schur, l. c. 171 = Vicia Orobus.
pisiformis, Schur, l. c. 170 = Vicia pisiformis.
sylvatica, Schur, l. c. = Vicia sylvatica.

VICIOIDES, Moench, Meth. 135 (1794) = **Vicia**, Tourn. (Legumin.).
hirsuta, Moench, l. c. 137 = Vicia pannonica.
hybrida, Moench, l. c. 136 = Vicia hybrida.
lutea, Moench, l. c. = Vicia lutea.
sepium, Moench, l. c. = Vicia sepium.
striata, Moench, l. c. 137 = Vicia pannonica.
sylvatica, Moench, l. c. 136 = Vicia sylvatica.
uncinata, Moench, l. c. = Vicia pannonica.

VICOA, Cass. in Ann. Sc. Nat. Sér. I. xvii. (1829) 418. *COMPOSITAE*, Benth. & Hook. f. ii. 334.
PENTANEMA, Cass. in Bull. Soc. Philom. (1818) 74.
appendiculata, DC. Prod. v. 474 = auriculata.
auriculata, *Cass. in Ann. Sc. Nat.* Sér. I. xvii. (1829) 418.—Ind. or.; Burma.
aurita, DC. Prod. v. 474 = auriculata.
cernua, *Dalz. in Dalz. & Gibs. Bomb. Fl.* 126.—Ind. or.
chrysopsioides, Benth. & Hook. f. Gen. ii. 335 (*chrysophthalmoides*) = Pulicaria petiolaris.
divaricata, *Benth. & Hook. f. Gen. l. c.*; *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 362.—Afr. trop.
indica, DC. in Wight, Contrib. 10 = auriculata.
Kotschyi, *Benth. & Hook. f. Gen.* ii. 335.—Afr. trop.
multicaule, *Benth. & Hook. f. l. c.*—Persia.
radiata, Benth. & Hook. f. l. c. = seq.
vestita, *Benth. & Hook. f. l. c.*—Ind. or.

VICTORIA, Lindl. Monog. 3 (1837). *NYMPHAEACEAE*. Benth. & Hook. f. i. 47.
amazonica, Sowerby, in Ann. & Mag. Nat. Hist. Ser. II. vi. (1850) 310 = regia.
Amazonum, Klotzsch, in Bot. Zeit. v. (1847) 245 = regia.
Cruziana, Orbign. in Ann. Sc. Nat. Sér. II. xiii. (1840) 57 = regia.
Fitzroyana, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1388 = Nymphaea gigantea.
regalis, Schomb. in Mag. Zool. & Bot. xi. (1837) t. 12 = regia.
regia, *Lindl. Monog.* 3; *et Bot. Reg.* (1838) *Misc.* 9.—Guian.
Regina, J. E. Gray, in Mag. Zool. & Bot. ii. (1837) p. iv; Schomb. in Proc. Bot. Soc. Lond. i. (1839) 44 = regia.
Reginae, Hook. Kew Journ. ii. (1850) 314, in nota = praec.

VICTORIPERREA, Homb. & Jacquinot, ex Decne. Bot. Voy. Astrol. & Zél. 83 (1853) = **Freycinetia**, Gaudich. (Pandan.).
impavida, Homb. l. c. 83.—Hab.?

VIDALIA, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 17 (1880) = **Kayea**, Wall. (Guttif.).
Garciae, F. Villar, l. c. 18.—Ins. Philipp.
lepidota, F. Villar, l. c. = K. lepidota.
Navesii, F. Villar, l. c.—Ins. Philipp.

VIEILLARDIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. viii. (1861) 375 = **Timeroyea**, Montr. (Nyctag.).
austro-caledonica, Brongn. et Gris, l. c. 376.—N. Caled.

VIEILLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) 196 = **Castanospermum**, A. Cunn. (Legumin.).
grandiflora, Montr. l. c.—N. Caled.

VIELLARDIA, Benth. & Hook. f. Gen. 556 (1865); et iii. 10 (1880), sphalm. = VIEILLARDIA, Montr., et VIEILLARDIA, Brongn. & Gris.

VIERAEA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 225 (1835–41). *COMPOSITAE*, Benth. & Hook. f. ii. 334.
VIERIA, Webb & Berth. l. c. t. 84 (1835–41).
laevigata, *Webb & Berth. l. c.* 226 [t. 84].—Ins. Canar.

VIEREYA, Steud. Nom. ed. II. ii. 774 (1841), sphalm. = VIREYA, Blume = **Rhododendron**, Linn. (Ericac.); et VIREYA, Rafin. = **Alloplectus**, Mart. (Gesner.).

VIERIA, Webb & Berth. Phyt. Canar. Ic. t. 84 (1836–41) = **Vieraea**, Sch. Bip. (Compos.).

VIEUSSEUXIA D. Delar. Desc. Pl. Nov. 31. t. 5 (1766) = **Moraea**, Mill. (Irid.).
angustifolia, Eckl. Top. Verz. 12.—Afr. austr.
aristata, D. Delar. Desc. Pl. Nov. 33 = M. edulis.
aristata, Houtt. Handleid. xii. 105. t. 80 = M. tricuspis.

VIEUSSEUXIA :—
aurantiaca, Eckl. Top. Verz. 13 = Moraea aurantiaca.
Bellenderi, Sweet, Hort. Brit. ed. I. 395 = M. Pavonia.
bituminosa, Eckl. Top. Verz. 14 = M. viscaria.
Brehmii, Eckl. l. c. 12.—Afr. austr.
ciliata, Eckl. l. c. 11 = M. papilionacea.
collina, Eckl. l. c. 12 = Homeria collina.
crispa, Eckl. l. c. = M. fimbriata, crispa.
curvata, Eckl. l. c. 13.—Afr. austr.
edulis, Link, Enum. Hort. Berol. i. 56 = M. edulis.
Freuchenia, Steud. Nom. ed. II. ii. 765 = M. ramosa.
fugax, D. Delar. Desc. Pl. Nov. 33 = M. fugax.
fugax, Roem. & Schult. Syst. i. 489 = M. edulis.
geniculata, Eckl. Top. Verz. 12.—Afr. austr.
glaucopis, DC. in Ann. Mus. Par. ii. (1803) 141 = M. glaucopis.
gracilis, Eckl. Top. Verz. 12 = M. gracilis.
graminifolia, Eckl. l. c. 11 = M. polyanthos.
grandiflora, Eckl. l. c. 14 = M. grandiflora.
intermedia, Eckl. l. c. 12.—Afr. austr.
iridioides, [Delar. in] Red. Lil. vi. t. 340 = M. Candolleana.
iridioides, Link, Enum. Hort. Berol. i. 56 = M. iridioides.
juncea, Eckl. ex Steud. Nom. ed. II. ii. 765.—Afr. austr.
longifolia, Eckl. Top. Verz. 14 = M. edulis.
lurida, Sweet, Hort. Brit. ed. I. 395 = M. lurida.
lutea, Eckl. Top. Verz. 14.—Afr. austr.
martinicensis, DC. in Ann. Mus. Par. ii. (1803) 138 = Trimeza lurida.
multiflora, Eckl. Top. Verz. 12 = M. multiflora.
multifolia, Eckl. l. c.—Afr. austr.
mutila, Eckl. l. c.; A. Dietr. Sp. Pl. ii. 494 = Iris mutila.
mutila, Klatt, in Linnaea, xxxiv. (1865–66) 621 = M. tripetala.
nervosa, Eckl. Top. Verz. 12 = M. papilionacea.
pauciflora, Eckl. l. c. = M. pauciflora.
Pavonia, DC. in Ann. Mus. Par. ii. (1803) 139 = M. Pavonia.
plumaria, Eckl. Top. Verz. 14 = M. iriopetala.
polystachya, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 131 = M. polystachya.
pulchra, Eckl. Top. Verz. 13.—Afr. austr.
ramosa, Eckl. l. c. 12 = M. ramosa.
rivularis, Eckl. l. c. 11.—Afr. austr.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 305 = M. diversifolia.
serrata, Eckl. ex Steud. Nom. ed. II. ii. 765.—Afr. austr.
setacea, Eckl. Top. Verz. 13 = M. setacea.
spiralis, D. Delar. Desc. Pl. Nov. 31. t. 5 = M. Pavonia.
tabularis, Eckl. Top. Verz. 13.—Afr. austr.
tenuifolia, Eckl. ex Steud. Nom. ed. II. ii. 765.—Afr. austr.
tenuis, Roem. & Schult. Syst. i. 491 = M. tenuis.
tricuspis, Fisch. Cat. Hort. Gorenk. ex Spreng. Syst. i. 165 = M. tricuspis.
tridentata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 305 = M. diversifolia.
tripetaloides, DC. in Ann. Mus. Par. ii. (1803) 138 = M. tripetala.
tristis, Eckl. Top. Verz. 11 = M. tristis.
unguicularis, Roem. & Schult. Syst. i. 491 = M. unguicularis.
villosa, Spreng. Syst. i. 166 = M. Pavonia.
viscaria, Eckl. Top. Verz. 12 = M. viscaria.

VIGIA, Vell. Fl. Flum. ix. t. 128 (1827) = **Plukenetia**, Linn. (Euphorb.).
serrata, Vell. l. c. (= *Fragariopsis scandens*).—Bras.

VIGIERA, Benth. & Hook. f. Gen. i. 644 (1865), err. typ. = seq.

VIGIERIA, Vell. Fl. Flum. 76 (1825); ii. tt. 73, 74 (1827) = **Escallonia**, Linn. (Saxifr.).
hispida, Vell. l. c. 76; ii. t. 73.—Bras.
laevis, Vell. l. c. 77; ii. t. 74.—Bras.

VIGINEIXIA, Pomel, Nouv. Mat. Fl. Atl. 12 (1874) = **Picris**, Linn. (Compos.).
Balansae, Pomel, l. c. (= *Helminthia Balansae*, Coss. & Dur.).—Afr. bor.

VIGNA, Savi, Mem. Phas. iii. 7 (1826?). *LEGUMINOSAE*, Benth. & Hook. f. i. 539.
CALLICYSTHUS, Endl. Prod. Fl. Norf. 90 (1833).
OTOPTERA, DC. Prod. ii. 240 (1825).
PLECTROTROPIS, Schum. & Thonn. Beskr. Guin. Pl. 338 (1827).
SCYTALIS, E. Mey. Comm. Pl. Afr. Austr. 144 (1835).
SPHENOSTYLIS, E. Mey. l. c. 148 (1835).
STROPHOSTYLES, E. Mey. l. c. 147 (1835).
Afzelii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 202.—Afr. trop.
ambacensis, *Welw. ex Baker, l. c.* 201.—Afr. trop.
andongensis, *Baker, l. c.* 197.—Afr. trop.
angivensis, *Baker, in Journ. Bot.* xx. (1882) 69.—Madag.
angustifolia, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 240.—Afr. austr.
angustifolia, Hook. f. in Hook. Niger Fl. 311 = capensis.
anomala, Walp. Rep. i. 779 = retusa.
Benthami, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 262.—Afr. trop.
brachycarpa, *Kurz, in Journ. As. Soc. Beng.* xliii. II. (1874) 185.—Burma.
brachystachys, Benth. Bot. Voy. Sulph. 86 = glabra.
brasiliensis, Mart. ex Benth. in Mart. Fl. Bras. xv. I. 194 = glabra.
Burchellii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 239.—Afr. austr.
calcarata, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 247 = Phaseolus calcaratus.
capensis, *Walp. in Linnaea*, xiii. (1839) 533.—Geront. trop.
carinalis, Benth. Bot. Voy. Sulph. 86 = capensis.
Catjang, *Walp. in Linnaea*, xiii. (1839) 533.—Cosmop. trop.
coerulea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 203.—Afr. trop.
comosa, *Baker, l. c.* 202.—Afr. trop.
crinita, *A. Rich. Tent. Fl. Abyss.* i. 217.—Abyss.
decipiens, *Harv. in Harv. & Sond. Fl. Cap.* ii. 241 (excl. syn.).—Afr. austr.
decipiens, Harv. l. c., partim = triloba.
dolichoides, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 206.—Reg. Himal.; Java.
Donii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 202.—Afr. trop.
frutescens, *A. Rich. Tent. Fl. Abyss.* i. 218.—Abyss.
gangetica, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 185 = glabra.
glabra, *Savi, Mem. Phas.* iii. 8.—Cosmop. trop.
golungensis, *Baker, in Oliver, Fl. Trop. Afr.* ii. 199.—Afr. trop.
gracilis, *Hook. f. in Hook. Niger Fl.* 311.—Afr. trop.
hastata, Blume, ex Miq. Fl. Ind. Bat. i. I. 187 = V. Catjang.
hastifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 200.—Afr. trop.
helicopus, Walp. in Linnaea, xiii. (1839) 534 = glabra.
heterophylla, *A. Rich. Tent. Fl. Abyss.* i. 218.—Abyss.
hirsuta, Feay, ex Wood, Bot. & Fl. 96 = glabra?
hirta, Hook. Ic. Pl. t. 637 = capensis.
hispida, *Walp. in Linnaea*, xiii. (1839) 534.—Afr. austr.
holosericea, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 200.—Afr. trop.
huillensis, *Welw. ex Baker, l. c.* 204.—Afr. trop.
Kotschyi, *Schweinf. Reliq. Kotschy.* (1868) 29. t. 22.—Afr. trop.
lanceolata, *Benth. in Mitch. Journ. Trop. Austral.* 350.—Austral.
lancifolia, *A. Rich. Tent. Fl. Abyss.* i. 221.—Abyss.
linearifolia, Hook. f. in Hook. Niger Fl. 309 = reticulata.
lobata, *Endl. ex Miq. Fl. Ind. Bat.* i. I. 188.—Afr. austr.; Ins. Timor.?
lobatifolia, *Baker, in Oliver, Fl. Trop. Afr.* ii. 199.—Afr. trop.
lobelifolia, Baker, l. c. 613, sphalm. = praec.
lucens, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 207.—Burma.
lutea, A. Gray, Bot. U. St. Expl. Exped. i. 452 = retusa.
luteola, Benth. in Mart. Fl. Bras. xv. I. 194. t. 50. f. 2 = glabra.
marginata, *Benth. ex Harv. & Sond. Fl. Cap.* ii. 240.—Afr. austr.
membranacea, *A. Rich. Tent. Fl. Abyss.* i. 219.—Abyss.

VIGNA:—
monachalis, Endl. ex Hassk. Cat. Hort. Bog. Alt. 279 = V. Catjang.
multiflora, *Hook. f. in Hook. Niger Fl.* 308.—Afr. trop.
nigritia, Hook. f. l. c. 310 = glabra.
nilotica, *Hook. f. l. c.* 311.—Afr. trop.
oahuensis, A. Gray, Bot. U. St. Expl. Exped. i. 450 = owahuensis.
oblonga, *Benth. Bot. Voy. Sulph.* 86.—Am. trop.; Afr. trop.
oblongifolia, *A. Rich. Tent. Fl. Abyss.* i. 220.—Abyss.
opisotricha, A. Rich. l. c. = Phaseolus Mungo.
ornata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 203.—Afr. trop.
owahuensis, *Vog. in Linnaea*, x. (1836) 585.—Ins. Sandvic.
paraguariensis, *Benth. in Mart. Fl. Bras.* xv. I. 195.—Am. austr.
Parkeri, *Baker, in Journ. Bot.* xx. (1882) 69.—Madag.
parviflora, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 201.—Afr. trop.
parvifolia, *Planch. ex Baker, l. c.* 206.—Afr. trop.
phaseoloides, *Baker, l. c.* 200.—Afr. trop.
pilosa, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 207.—Ind. or.
pubigera, *Baker, in Oliver, Fl. Trop. Afr.* ii. 201.—Afr. trop.
radicans, *Welw. ex Baker, l. c.* 198.—Afr. trop.
repens, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 205.—Burma.
reticulata, *Hook. f. in Hook. Niger Fl.* 310.—Afr. trop.
retusa, *Walp. in Linnaea*, xiii. (1839) 534.—Cosmop. trop.
rotundifolia, *Hassk. Cat. Hort. Bog. Alt.* 279.—Java.
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 451.—Ins. Sandvic.
scabra, Sond. in Linnaea, xxiii. (1850) 32 = capensis.
Schimperi, *Baker, in Oliver, Fl. Trop. Afr.* ii. 201.—Abyss.
sinensis, Endl. ex Hassk. Pl. Jav. Rar. 386 = V. Catjang.
suberecta, *Benth. in Mitch. Journ. Trop. Austral.* 388.—Austral.
syndwicensis, C. Muell. in Walp. Ann. iv. 563, sphalm. = sandwicensis.
tenuis; *D. Dietr. Syn. Pl.* iv. 1197.—Afr. austr.
tenuis, *Franch. in Revoil, Çomali*, 29.—Afr. trop.
Thonningii, Hook. f. in Hook. Niger Fl. 311 = capensis.
tranquebarica, D. Dietr. Syn. Pl. iv. 1198 = V. Catjang.
triloba, *Walp. in Linnaea*, xiii. (1839) 534.—Afr. trop. et austr.
tuberosa, A. Rich. Tent. Fl. Abyss. i. 217. t. 42 = capensis.
unguiculata, Walp. Rep. i. 779 = V. Catjang.
venulosa, *Baker, in Oliver, Fl. Trop. Afr.* ii. 203.—Afr. trop.
vexillata, A. Rich. Ess. Fl. Cub. 440 = capensis.
villosa, Hook. & Arn. Bot. Beech. Voy. 81 = owahuensis.
villosa, Savi, Mem. Phas. iii. 9 = glabra.
Wightii, *Benth. ex Bedd. in Trans. Linn. Soc.* xxv. (1865) 215.—Ind. or.

VIGNALDIA, A. Rich. Tent. Fl. Abyss. i. 357 (1847) = **Pentas**, Benth. (Rubiac.).
luteola, Schweinf. Beitr. Fl. Aethiop. 282 = Mussaenda luteola.
occidentalis, Hook. f. in Journ. Linn. Soc. vii. (1864) 197 = P. occidentalis.
Quartiniana, A. Rich. Tent. Fl. Abyss. i. 357 = P. carnea.
Schimperi, A. Rich. l. c. 358 = P. Schimperiana.
Schimperiana, Hiern, in Oliver, Fl. Trop. Afr. iii. 45 = praec.

VIGNANTHA, Schur, Enum. Pl. Trans. 705 (1866) = **Carex**, Dill. (Cyperac.).
acuta, Schur, Enum. Pl. Transs. 706 = C. acuta.
dacica, Schur, l. c. 705 = C. caespitosa.
Dreyeri, Schur, l. c. 706 = C. caespitosa.
Moenchiana, Schur, l. c. = C. acuta.
mucronata, Schur, l. c. 705 = C. mucronata.
prolixa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 176 = C. acuta.

VIGNANTHA :—
rigida, Schur, Enum. Pl. Transs. 705 = Carex caespitosa.
stricta, Schur, l. c. 707 = C. stricta.
touranginiana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 176 = C. acuta.
vulgaris, Schur, Enum. Pl. Transs. 706 = C. vulgaris.

VIGNAUDIA, Schweinf. Beitr. Fl. Aethiop. 282 (1867) sphalm. = VIGNALDIA, A. Rich. = **Pentas**, Benth. (Rubiac.).

VIGNEA, Beauv. in Lestib. Essai Fam. Cyp. 22 (1819); et in Flora, iv. (1821) 17 = **Carex**, Dill. (Cyperac.).
acroandra, Schur, Enum. Pl. Transs. 699 = C. acroandra.
acuta, Reichb. in Moessl. Handb. ed. II. iii. 1626 = C. acuta.
altissima, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 79 = C. muricata.
ammophila, Reichb. in Moessl. Handb. ed. II. iii. 1615 = C. divisa.
appressa, Reichb. l. c. 1621 = C. paniculata.
aquatilis, Reichb. l. c. 1624 = C. aquatilis.
arenaria, Reichb. l. c. 1618 = C. arenaria.
argyroglochin, Reichb. l. c. 1612 = C. leporina.
axillaris, Reichb. l. c. 1617 = C. axillaris.
Bertolonii, Reichb. l. c. 1608 = C. divisa.
bicolor, Reichb. l. c. 1624 = C. bicolor.
Boenninghauseniana, Reichb. l. c. 1616 = C. axillaris.
bracteosa, Reichb. l. c. 1619 = C. bracteosa.
brizoides, Reichb. l. c. 1611 = C. brizoides.
caespitosa, Reichb. l. c. 1625 = C. caespitosa.
canescens, Reichb. l. c. 1612 = C. canescens.
capitata, Reichb. l. c. 1606 = C. capitata.
cephalophora, Reichb. l. c. 1607 = C. cephalophora.
cerina, Reichb. Fl. Germ. Excurs. 140^7 = C. vulgaris.
chlorocarpa, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 193 = C. vulgaris.
chordorrhiza, Reichb. in Moessl. Handb. ed. II. iii. 1608 = C. chordorrhiza.
compacta, Reichb. l. c. 1623 = C. vulgaris.
conjugata, Reichb. l. c. 1622 = C. brunnea.
crassinervis, Schur, Enum. Pl. Transs. 704 = C. vulgaris.
curta, Opiz, Seznam, 103 (err. typ. 203) = C. canescens.
curvula, Opiz, l. c.; Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 78 = C. Schreberi.
cyperoides, Peterm. Fl. Lips. Excurs. 49 = C. cyperoides.
dacica, Fuss, Fl. Transs. 681 = C. caespitosa.
Davalliana, Reichb. in Moessl. Handb. ed. II. iii. 1605 = C. Davalliana.
dioica, Reichb. l. c. 1604 = C. dioica.
disticha, Peterm. Fl. Bienitz, 16 = C. disticha.
divisa, Reichb. in Moessl. Handb. ed. II. iii. 1615 = C. divisa.
divulsa, Reichb. l. c. 1619 = C. divulsa.
Drejeri, Fuss, Fl. Transs. 680 = C. caespitosa.
echinata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 176 = C. echinata.
elongata, Reichb. in Moessl. Handb. ed. II. iii. 1617 = C. elongata.
festucacea, Reichb. l. c. 1614 = C. straminea.
foenea, Reichb. l. c. = C. straminea.
foetida, Reichb. l. c. 1607 = C. foetida.
Gaudiniana, Pritz. Ic. Ind. i. 1161 = C. Gaudiniana.
Gebhardi, Reichb. in Moessl. Handb. ed. II. iii. 1613 = C. elongata.
glareosa, Reichb. l. c. 1626 = C. glareosa.
grypos, Reichb. l. c. 1608 = C. echinata.
guestphalica, Reichb. Fl. Germ. Excurs. 140^7 = C. muricata.
Heleonastes, Reichb. in Moessl. Handb. ed. II. iii. 1609 = C. Heleonastes.
hydrophila, Reichb. Fl. Germ. Excurs. 140^7 = C. echinata.
incurva, Reichb. in Moessl. Handb. ed. II. iii. 1607 = C. incurva.
intermedia, Reichb. l. c. 1618 = C. disticha.
juncoides, Opiz, Seznam, 104 = C. juncoides.
lagopina, Reichb. in Moessl. Handb. ed. II. iii. 1609 = C. lagopina.
lagopodioides, Reichb. l. c. 1614 = C. lagopodioides (cf. Addend.).
leiorhyncha, Pritz. Ic. Ind. i. 1161 = C. leiorhyncha.
leporina, Reichb. in Moessl. Handb. ed. II. iii. 1613 = C. leporina.
loliacea, Reichb. l. c. 1610 = C. loliacea.
lucorum, Reichb. l. c. 1614 = C. lucorum.

VIGNEA :—
maritima, Reichb. in Moessl. Handb. ed. II. iii. 1623 = Carex maritima.
microstachya, Reichb. l. c. = C. microstachya.
microstylis, Reichb. Fl. Germ. Excurs. 140^6 = C. microstyla.
Moenchiana, Reichb. in Moessl. Handb. ed. II. iii. 1626 = C. acuta.
mucronata, Reichb. Fl. Germ. Excurs. 140^7 = C. mucronata.
multiflora, Reichb. in Moessl. Handb. ed. II. iii. 1621 = C. vulpinoidea.
muricata, Reichb. l. c. 1611 = C. muricata.
neglecta, Peterm. Fl. Bienitz, 17 = C. caespitosa.
nemorosa, Reichb. in Moessl. Handb. ed. II. iii. 1620 = C. vulpina.
Nenningiana, Opiz, Seznam, 103 (err. typ. 203) = C. Nenningiana (in Addend.).
norvegica, Reichb. in Moessl. Handb. ed. II. iii. 1610 = C. norvegica.
Ohmuelleriana, Pritz. Ic. Ind. i. 1161 = C. arenaria.
pacifica, Reichb. ex Nym. Consp. 777 = C. vulgaris.
paniculata, Reichb. in Moessl. Handb. ed. II. iii. 1622 = C. paniculata.
paradoxa, Reichb. l. c. 1621 = C. paradoxa.
Persoonii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 = C. canescens.
psammophila, Schur, Enum. Pl. Transs. 700 = C. brizoides.
Pseudo-arenaria, Reichb. Ic. Fl. Germ. viii. t. 208 = C. arenaria.
Pseudo-brizoides, Schur, Enum. Pl. Transs. 700 = C. Pseudo-brizoides.
Pseudo-Schreberi, Schur, l. c. = C. brizoides.
pulicaris, Reichb. in Moessl. Handb. ed. II. iii. 1606 = C. pulicaris.
pulla, Reichb. l. c. 1624 = C. saxatilis.
remota, Reichb. l. c. 1616 = C. remota.
repens, Reichb. l. c. 1618 = C. repens.
repens, Schur, Enum. Pl. Transs. 701 = C. arenaria.
retroflexa, Reichb. in Moessl. Handb. ed. II. iii. 1610 = C. rosea.
rigida, Fuss, Fl. Transs. 681 = C. vulgaris.
rivularis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 78 = C. divisa.
rosea, Reichb. in Moessl. Handb. ed. II. iii. 1611 = C. rosea.
salina, Reichb. l. c. 1623 = C. salina.
saxatilis, Reichb. l. c. 1624 = C. saxatilis.
Schreberi, Reichb. l. c. 1614 = C. Schreberi.
scirpoides, Reichb. l. c. 1611 = C. echinata.
scoparia, Reichb. l. c. 1613 = C. scoparia.
setifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 175 = C. divisa.
squarrosa, Reichb. in Moessl. Handb. ed. II. iii. 1606 = C. squarrosa.
stellulata, Reichb. l. c. 1610 = C. echinata.
stenophylla, Reichb. l. c. 1607 = C. stenophylla.
stipata, Reichb. l. c. 1620 = C. stipata.
striata, Reichb. l. c. 1625 = C. lacistoma.
stricta, Reichb. l. c. 1624 = C. stricta.
strictifolia, Opiz, Seznam, 103 (err. typ. 203) = C. acuta.
tenuissima, Schur, Enum. Pl. Transs. 979 (Index) = C. muricata.
teretiuscula, Reichb. in Moessl. Handb. ed. II. iii. 1621 = C. paniculata.
trinervis, Reichb. l. c. 1625 = C. trinervis.
tripartita, Reichb. l. c. 1609 = C. curvula.
virens, Reichb. l. c. 1611 = C. muricata.
vulgaris, Fuss, Fl. Transs. 680 = C. vulgaris.
vulpina, Reichb. in Moessl. Handb. ed. II. iii. 1620 = C. vulpina.

VIGOLINA, Poir. Encyc. viii. 613 (1808) = **Galinsoga**, Ruiz & Pav. (Compos.).
Acmella, Poir. l. c. = G. parviflora.
Armella, Steud. Nom. ed. II. i. 656, in syn. sphalm. = praec.

VIGUIERA, H. B. & K. Nov. Gen. et Sp. iv. 224. t. 379 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 375.
BAHIOPSIS, Kellogg, in Proc. Calif. Acad. ii. (1863) 35.
HARPALIUM, DC. Prod. v. 583 (1836).
LEIGHIA, Cass. in Dict. Sc. Nat. xxv. 435 (1822).

VIGUIERA :—
acuminata, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 91 = Oyedaea acuminata.
anchusaefolia, *Baker, in Mart. Fl. Bras.* vi. III. 222.—Bras.
arenaria, *Baker, l. c.* 226.—Bras.
asperrima, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 401.—Bras.
aspilioides, *Baker, in Mart. Fl. Bras.* vi. III. 228.—Bras.
attenuata, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 400 = Aspilia asperrima.
blepharolepis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 5.—N. Mexic.
Bonplandiana, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 399 = Oyedaea Bonplandiana.
bracteata, *Gardn. l. c.* 404.—Bras.
brevipes, DC. Prod. v. 579 = helianthoides.
buddleiaeformis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 177.—Mexic.
calva, *Benth. & Hook. f. Gen.* ii. 375.—Bolivia.
canescens, *DC. Prod.* v. 579.—Mexic.
cordifolia, *A. Gray, Pl. Wright.* i. 107.—Mexic.
decurrens, *A. Gray, in Proc. Am. Acad.* xix. (1883) 5.—Am. bor.
deltoidea, *A. Gray, l. c.* v. (1859) 161.—Calif.
densifolia, *Baker, in Mart. Fl. Bras.* vi. III. 219.—Paraguay.
dentata, Spreng. Syst. iii. 615 = helianthoides.
discolor, *Baker, in Mart. Fl. Bras.* vi. III. 228.—Bras.
dissitifolia, *Baker, l. c.* 225.—Bras.
elegans, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 398 = Aspilia foliacea.
elliptica, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 194 = Aspilia Burchellii.
excelsa, *Benth. & Hook. f. Gen.* ii. 375; *et ex Hemsl. Biol. Centr. Am. Bot.* ii. 177.—Mexic.
filifolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 219.—Bras.
floribunda, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 401 = Aspilia floribunda.
foliacea, Spreng. Syst. iii. 616 = Aspilia foliacea.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 224.—Bras.
Ghiesbreghtii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 6.—Mexic.
Gilliesii, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 39.—Chili.
glabra, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 398 = Aspilia glabra.
glandulosa, Bertol. Misc. Bot. vii. 45 = Baldwinia uniflora.
gracilis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 402 = Aspilia gracilis.
grammatoglossa, *DC. Prod.* v. 580.—Mexic.
grandiflora, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 404.—Bras.
helianthoides, *H. B. & K. Nov. Gen. et Sp.* iv. 226. t. 379.—Mexic.; Ind. occ.
hirsuta, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 397 = Aspilia foliacea.
hispida, *Baker, in Mart. Fl. Bras.* vi. III. 220.—Bras.
Humboldtiana, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 398 = Oyedaea Humboldtiana.
imbricata, *Baker, in Mart. Fl. Bras.* vi. III. 220.—Bras.
Kunthiana, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 399.—Bras.
laciniata, *A. Gray, in Torr. Bot. Mex. Bound.* 89.—Calif.
lanata, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 218.—Calif.
laxa, *Baker, in Mart. Fl. Bras.* vi. III. 222.—Bras.
laxa, DC. Prod. v. 580 = helianthoides.
linearis, *Sch. Bip. ex Hemsl. in Biol. Centr. Am. Bot.* ii. 178.—Mexic.
macrophylla, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 92 = Oyedaea macrophylla.
macrorhiza, *Baker, in Mart. Fl. Bras.* vi. III. 225.—Bras.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
microcline, DC. Prod. v. 579 = helianthoides?
mollis, *Griseb. in Goett. Abh.* xix. (1874) 183.—Reg. Argent.
nervosa, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 403.—Bras.

VIGUIERA:—
nivea, Benth. & Hook. f. Gen. ii. 378, in nota sub Encelia = tephrodes.
nonneaefolia, *Baker, in Mart. Fl. Bras.* vi. III. 222.—Bras.
nudicaulis, *Baker, l. c.* 228.—Bras.
oblonga, Benth. & Hook. f. Gen. ii. 374, in nota sub Oyedaea = Aspilia oblonga.
oblongifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 402.—Bras.
obtusifolia, *Baker, in Mart. Fl. Bras.* vi. III. 226.—Bras.
oppositipes, *DC. Prod.* v. 580.—Mexic.
ovalifolia, *Baker, in Mart. Fl. Bras.* vi. III. 226.—Bras.
pachycephala, *Hemsl. Biol. Centr. Am. Bot.* ii. 178.—Mexic.
Parishii, *Greene, in Bull. Torrey Bot. Club*, ix. (1882) 15.—Arizona.
peruviana, *A. Gray, in Proc. Am. Acad.* iv. (1859) 124.—Peruv.
pilosa, *Baker, in Mart. Fl. Bras.* vi. III. 223.—Bras.
platyphylla, *Baker, l. c.* 227.—Bras.
Poeppigii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 6.—Am. bor.
Pohliana, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 202 = Aspilia laevissima.
prostrata, DC. Prod. v. 579 = Helianthus tracheliifolius.
pusilla, Less. ex Baker, in Mart. Fl. Bras. vi. III. 194 = Aspilia pusilla.
Radula, *Baker, l. c.* 223.—Bras.
ramosissima, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 402 = Aspilia floribunda.
reflexa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 196 = Aspilia reflexa.
reticulata, *S. Wats. in Am. Natural.* vii. (1873) 301.—Am. bor. occ.
retifolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 223.—Bras.
rigida, Hort.—Cf. Gard. Chron. (1881) II. 396 = Helianthus rigidus.
robusta, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 403.—Bras.
rugosa, *Benth. & Hook. f. Gen.* ii. 375; *et ex Hemsl. Biol. Centr. Am. Bot.* ii. 178.—Mexic.
Sagraeana, DC. Prod. v. 579 = helianthoides.
scabra, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 227 = robusta.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 305.—Mexic.
sessilifolia, *DC. Prod.* v. 579.—Mexic.
sphaerocephala, *Hemsl. Biol. Centr. Am. Bot.* ii. 179.—Mexic.
stenophylla, *Griseb. in Goett. Abh.* xxiv. (1879) 193.—Reg. Argent.
subincisa, *Benth. Bot. Voy. Sulph.* 27.—Calif.
tenuifolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 399.—Bras.
tephrodes, *A. Gray, in Proc. Am. Acad.* xvii. (1882) 218.—Calif.
texana, Torr. & Gray, Fl. N. Am. ii. 318 = helianthoides.
tomentosa, *A. Gray, in Proc. Am. Acad.* v. (1862) 161.—Calif.
triplinervis, Less. ex Baker, in Mart. Fl. Bras. vi. III. 191 = Aspilia glabra.
triquetra, DC. Prod. v. 579 = helianthoides.
tuberosa, *Benth. & Hook. f. Gen.* ii. 388.—Bolivia.
tuberosa, *Griseb. in Goett. Abh.* xxiv. (1879) 193.—Reg. Argent.
tucumanensis, *Griseb. l. c.* xix. (1874) 183.—Reg. Argent.
urticiformis, *Hemsl. Biol. Centr. Am. Bot.* ii. 179.—Mexic.
vernonioides, *Baker, in Mart. Fl. Bras.* vi. III. 223.—Bras.
wedelioides, *Baker, in Journ. Bot.* xx. (1882) 226.—Madag.

VILARIA, Guett. Mém. Minéral. Dauph. i. Pref. 108; ii. t. 19 (1779) = **Berardia**, Vill. (Compos.).
subacaulis, Guett. l. c. 170 = B. subacaulis.

VILFA, Adans. Fam. ii. 495 (1763) = **Agrostis**, Linn. (Gramin.).

VILFA, Beauv. Agrost. 16. t. 5. f. 8 (1812) = **Sporobolus**, R. Br. (Gramin.).
actinoclada, F. Muell. Fragm. vi. 84 = S. actinocladus.

VILFA:—
acuminata, Trin. Sp. Gram. iii. t. 348.—Bras.
acutigluma, Steud. in Lechl. Berb. Am. Austr. 56.—Peruv.
adusta, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 80.—Bras.
aemula, Beauv. Agrost. 16 = Deyeuxia Forsteri.
aenea, Trin. Sp. Gram. i. t. 23 = Sporobolus aeneus.
affinis, Steud. Nom. ed. II. ii. 766 = S. affinis, *Kunth.*
africana, Beauv. Agrost. 16 = S. capensis.
agrostidea, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 88 = S. cryptandrus.
airoides, Trin. ex Steud. Nom. ed. II. ii. 766 = S. airoides.
alba, Beauv. Agrost. 16 = Agrostis alba.
alba, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 89 = Eatonia obtusata.
albicans, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 101 = S. albicans.
algida, Trin. Gram. Unifl. 159 = Phippsia algida.
alopecuroides, Beauv. Agrost. 16 = Polypogon monspeliensis.
alpicola, Steud. Syn. Pl. Gram. 154 = S. alpicola.
angustifolia, Steud. l. c. 160 = S. angustifolius.
arabica, Steud. l. c. 421 = S. arabicus.
arachnoidea, Beauv. Agrost. 147 = Muehlenbergia expansa.
arenaria, Trin. Gram. Unifl. 160 = S. pungens.
arguta, Nees, Agrost. Bras. 395 = S. argutus.
arkansana, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 64 = S. argutus.
articulata, Beauv. Agrost. 16 = Chaeturus fasciculatus.
arundinacea, Trin. Gram. Unifl. 157 = Arctagrostis latifolia.
aspera, Beauv. Agrost. 16 = S. asper.
asperifolia, Meyen, Reise, i. 349, 408 = S. asperifolius.
atrorubens, H. B. & K. Nov. Gen. et Sp. vii. t. 685 = S. atrovirens.
atrovirens, H. B. & K. l. c. i. 138 = S. atrovirens.
australis, Beauv. Agrost. 16 = Agrostis australis.
barbata, Beauv. l. c. = S. virginicus.
Berteroana, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 100 = Ins. S. Doming.
Billardierii, Beauv. Agrost. 16 = Deyeuxia Billardieri.
brachystachys, Presl, Fl. Sic. p. xliv = Agrostis brachystachys.
brevifolia, Nees, in Linnaea, vii. (1832) 294 = S. tenellus.
Brongniartii, Steud. Nom. ed. II. ii. 766 = S. durus.
caespitosa, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 72 = S. caespitosus.
capensis, Beauv. Agrost. 16 = S. capensis.
capillaris, Arn. ex Miq. Anal. Bot. Ind. ii. 25 = S. pulchellus.
capillaris, Beauv. Agrost. 147 = Agrostis delicatula.
Carmichaeli, Steud. Nom. ed. II. ii. 767 = Agrostis Carmichaeli.
centrifuga, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 57 = S. centrifugus.
ciliata, Beauv. Agrost. 16 = Festuca Thunbergii.
ciliata, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 63 = S. ciliatus.
clandestina, Nees, ex Steud. Nom. ed. II. ii. 767 = S. asper.
coarctata, Beauv. Agrost. 16 = Agrostis alba.
commutata, Trin. Gram. Unifl. 156 = S. commutatus.
composita, Beauv. Agrost. 16 = Agrostis composita.
compressa, Beauv. l. c. = S. compressus.
conferta, Steud. Syn. Pl. Gram. 160 = S. confertus.
confertiflora, Steud. l. c. 155 = S. confertiflorus.
confinis, Steud. l. c. 160 = S. affinis, *A. Rich.*
consimilis, Steud. Nom. ed. II. ii. 767 = S. consimilis.
coromandelina, Beauv. Agrost. 16 = S. coromandelianus.
crinita, Beauv. l. c. = Dichelachne crinita.
cruciata, Beauv. l. c. = Chloris cruciata.
cryptandra, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 69 = S. cryptandrus.
cuspidata, Torr. ex Hook. Fl. Bor. Am. ii. 238 = S. cuspidatus.
cylindrica, Beauv. Agrost. 16 = Deyeuxia cylindrica.
debilis, Beauv. l. c. = Deyeuxia Forsteri.
decipiens, Beauv. l. c. = Deyeuxia scabra.
decumbens, Beauv. l. c. = Agrostis alba?
densa, Beauv. l. c. = Agrostis verticillata.
depauperata, Torr. ex Hook. Fl. Bor. Am. ii. 257. t. 236 = S. depauperatus.

VILFA:—
diandra, Trin. Gram. Unifl. 154 = Sporobolus orientalis.
dianthera, Steud. Syn. Pl. Gram. 162 = Agrostis dianthera.
discospora, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 59 = S. commutatus.
dispar, Beauv. Agrost. 16 = Agrostis elata?
distichophylla, Trin. Gram. Unifl. 160 = Agrostis distichophylla.
divaricata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 146 = Agrostis vulgaris?
domingensis, Trin. in Spreng. Neue Entdeck. ii. 59 = S. domingensis.
Drummondii, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 106 = S. Drummondii.
dulcis, Beauv. Agrost. 16 = Agrostis verticillata.
dura, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 54 = S. virginicus?
elatior, Nees, Agrost. Bras. 396 = S. Sprengelii.
elegans, H. B. & K. Nov. Gen. et Sp. i. 139 = Agrostis Humboldtiana.
elongata, Beauv. Agrost. 16 = S. indicus.
eminens, Steud. Nom. ed. II. ii. 767 = S. eminens.
erosa, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 86 = S. diander.
exilis, Trin. l. c. 89 = S. indicus.
eximia, Nees, ex Trin. l. c. 77.—Bras.
familiaris, Steud. in Lechl. Berb. Am. Austr. 56.—Peruv.
fasciculata, H. B. & K. Nov. Gen. et Sp. i. 139 = Agrostis fasciculata.
fastigiata, [Nees, in] Meyen, Reise, i. 484 = S. fastigiatus.
festiva, Steud. Syn. Pl. Gram. 158 = S. festivus.
filiculmis, Thurb. ex Vasey, Cat. Gr. U. St. 44 = S. filiculmis.
fimbriata, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 69 = S. fimbriatus.
frondosa, Beauv. Agrost. 16 = Muehlenbergia mexicana.
frondosa, Presl, Cyp. et Gram. Sic. 23 = Agrostis vulgaris.
fulvescens, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 76.—Am. bor.
Funckii, Steud. Syn. Pl. Gram. 159.—Venezuela.
gaditana, Steud. l. c. 154 = S. pungens.
Gaudichaudii, Steud. Nom. ed. II. ii. 767.—Ins. Falkland.
gavana, Steud. in Lechl. Berb. Am. Austr. 56.—Peruv.
geniculata, Nees, ex Steud. Syn. Pl. Gram. 156.—Ind. or.
gigantea, Beauv. Agrost. 16 = Agrostis alba.
gigantea, Turcz. ex Griseb. in Ledeb. Fl. Ross. iv. 435 = Arctagrostis latifolia.
Gigas, Steud. Syn. Pl. Gram. 160 = S. Gigas.
glauca, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 52 = *helvola*.
glaucescens, Presl, Cyp. et Gram. Sic. 23 = Agrostis sicula.
glaucifolia, Steud. Syn. Pl. Gram. 154 = S. glaucifolius.
glomerata, J. Presl, in Rel. Haenk. i. 239 = Agrostis glomerata.
gracilis, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 104.—Am. bor.
gracilis, Trin. l. c. 74 = *subsetacea*.
gracillima, Thurb. in Bot. Calif. ii. 268 = S. gracillimus.
grandiflora, Nees, ex Steud. Syn. Pl. Gram. 158.—Chili.
helvola, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 52.—Arab.; Abyss.
heterolepis, A. Gray, in Ann. Lyc. N. York, iii. (1835) 233 = S. heterolepis.
heterolepsis, Lapham, in Trans. Wisconsin St. Agric. Soc. iii. 1853 (1854) 426, sphalm. = praec.
hispida, Beauv. Agrost. 16 = Agrostis vulgaris.
Hookeri, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 106 = S. asper.
humifusa, H. B. & K. Nov. Gen. et Sp. i. 137 = S. humifusus.
humilis, Steud. Nom. ed. II. ii. 767 = S. humilis.
hybrida, Beauv. Agrost. 16 = Agrostis canina.
indica, Trin. ex Steud. Nom. ed. II. ii. 767 = S. indicus.
insularis, Steud. Syn. Pl. Gram. 156 = S. insularis.
intermedia, Trin. Gram. Unifl. 156.—Ind. occ.
involuta, Beauv. Agrost. 16 = Agrostis involuta.
ioclados, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 65 = S. ioclados.
Jacquemontii, Trin. l. c. 92 = S. Jacquemontii.

VILFA :—
juncea, Trin. Gram. Unifl. 157 = Sporobolus junceus.
laterifolia, Beauv. Agrost. 16 = Muehlenbergia mexicana.
laxiflora, Beauv. l. c. = Agrostis laxiflora.
lenta, Beauv. l. c. = Panicum filiforme.
Lindleyi, Steud. Syn. Pl. Gram. 162 = S. Lindleyi.
linearis, Beauv. Agrost. 16 = Cynodon Dactylon.
littoralis, Beauv. l. c. = S. virginicus.
lobata, Beauv. l c. = Deyeuxia quadriseta.
longifolia, Torr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 107 = S. asper.
Ludwigii, Steud. Syn. Pl. Gram. 155 = S. Ludwigii.
lutosa, Beauv. Agrost. 16 = Polypogon littoralis.
luxurians, Steud. in Lechl. Berb. Am. Austr. 56.—Peruv.
macra, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. 1840) 79.—Am. bor.
macusaniensis, Steud. in Lechl. Berb. Am. Austr. 56.—Peruv.
magellanica, Beauv. Agrost. 16 = Agrostis magellanica.
mangalorica, Hochst. ex Miq. Anal. Bot. Ind. ii. 24 = S. minutiflorus.
marginata, Steud. Syn. Pl. Gram. 155 = S. marginatus.
maritima, Beauv. Agrost. 16 = Agrostis alba.
Matrella, Nees, Agrost. Bras. 400 = S. Matrella.
mauritiana, Steud. Syn. Pl. Gram. 157.—Ins. Maurit.
mexicana, Beauv. Agrost. 16 = Muehlenbergia mexicana.
minima, Trin. ex Steud. Syn. Pl. Gram. 162 = Agrostis minima.
minima, Vasey, in Monthly Rep U. St. Dep. Agric. 1874 (1875) 155 ; et in Rothr. Rep. Bot. U. St. Surv. vi. 282 = S. Wolfii.
minuta, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 84 = S. minutus.
minutiflora, Trin. Gram. Unifl. 158 = S. minutiflorus.
Mitchellii, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 53.—Austral.
monandra, Trin. Gram. Unifl. 159 = Phippsia algida.
montana, Beauv. Agrost. 16 ; Hook. f. in Journ. Linn. Soc. vii. (1864) 228 = Deyeuxia montana.
Muehlenbergii, Steud. Nom. ed. II. ii. 767 = S. Muehlenbergii.
muricata, J. Presl, in Rel. Haenk. i. 240 = Agrostis muricata.
murina, Sieber, ex Steud. Syn. Pl. Gram. 157 = S. virginicus.
myriantha, Steud. l. c. 158 = S. myrianthus.
natalensis, Steud. l. c. 154.—Afr. austr.
neglecta, Beauv. ex Steud. Nom. ed. II. ii. 767 = Agrostis alba.
nigra, S. F. Gray, Nat. Arr. Brit. Pl. ii. 145 = Agrostis vulgaris.
noterophila, Rupr. in Bull. Acad. Brux. ix. II. (1842) 227 = S. noterophila.
Novae-Hollandiae, Beauv. Agrost. 181 = Deyeuxia Forsteri.
nutans, Beauv. l. c. = Ichnanthus pallens.
orientalis, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 65 = S. orientalis.
Orizabae, Rupr. in Bull. Acad. Brux. ix. II. (1842) 227. —Mexic.
pallida, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 62 = S. arabicus.
palustris, Beauv. Agrost. 16 = Agrostis alba.
panicea, Beauv. l. c. = Gastridium australe.
panicoides, Beauv. ex Steud. Nom. ed. II. ii. 768 = Panicum Lamarckii.
paniculata, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 67.—Afr. trop.
parviflora, Beauv. Agrost. 16 = S. parviflorus.
parvula, Steud. Nom. ed. II. ii. 768 = S. minutus.
patula, Beauv. Agrost. 16 = Agrostis alba.
pilifera, Trin. Gram. Unifl. 157 = S. piliferus.
pilosa, Beauv. Agrost. 16 = Calamagrostis tenella.
plebeia, Beauv. l. c. = Deyeuxia plebeia.
plumbea, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 98 = S. plumbeus.
Preslii, Steud. Nom. ed. II. ii. 768 = S. ciliatus.
procera, Beauv. Agrost. 16 = Agrostis procera.
pulchella, Presl, Cyp. et Gram. Sic. 23 = Agrostis elegans.
pulchella, Trin. Gram. Unifl. 160 = S. pulchellus.
pumila, Beauv. Agrost. 16 = Agrostis vulgaris.
pungens, Beauv. l. c. = S. pungens.
pungens, Trin. Sp. Gram. i. t. 47 = S. virginicus.

VILFA :—
purpurascens, Beauv. Agrost. 16 = Sporobolus purpurascens.
pyramidalis, Trin. ex Steud. Nom. ed. II. ii. 768 = S. pyramidalis.
racemosa, Beauv. Agrost. 16 = Muehlenbergia glomerata.
ramosissima, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 58 = S. ramosissimus.
ramulosa, H. B. & K. Nov. Gen. et Sp. i. 137 = S. ramulosus.
rara, Beauv. Agrost. 16 = Dichelachne sciurea.
repens, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 102 = S. repens.
retrofracta, Beauv. Agrost. 16 = Deyeuxia Forsteri.
Retzii, Steud. Nom. ed. II. ii. 768 = S. diander.
rhizomatosa, Steud. Syn. Pl. Gram. 421.—Madag.
Richardi, Steud. l. c. 153.—Ind. occ.
Richardsonis, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 103.—Am. bor.
Riehlii, Steud. Syn. Pl. Gram. 154.—Am. bor.
rigens, Trin. Sp. Gram. iii. t. 250 = S. rigens.
rigida, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 89 = Ammophila longifolia ?
rigidifolia, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 80.—Ind. or.
robusta, Trin. l. c. 54 = S. robustus.
Roxburghiana, Nees, ex Steud. Nom. ed. II. ii. 768 = S. commutatus.
Roxburghii, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 59 = S. commutatus.
rupestris, Trin. l. c. 66 = S. rupestris.
rupestris, Trin. Gram. Panic. 22 = S. indicus.
Ruppelliana, Steud. Nom. ed. II. ii. 768 = S. Rueppellianus.
sabeana, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 90 = S. coromandelianus *vel* argutus.
scabra, Beauv. Agrost. 16 = Deyeuxia scabra.
Schiedeana, Trin. in Mém. Acad. Pétersb Sér. VI. Sc. Nat. v. II. (1840) 73.—Am. bor.
sciurea, Beauv. Agrost. 16 = Dichelachne sciurea.
scoparia, Steud. Nom. ed. II. ii. 768 = S. scoparius.
serotina, Torr. ex Trin. Sp. Gram. iii. t. 251 = S. serotinus.
setacea, Beauv. Agrost. 16 = Agrostis setacea.
setulosa, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 55.—Arab.
sicula, Steud. Nom. ed. II. ii. 768 = Festuca sicula.
Spartinae, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 82.—Texas.
spicata, Beauv. Agrost. 16 = S. spicatus.
Sporobolus, Trin. Gram. Unifl. 160 = S. Sprengelii.
squarrosa, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 100.—Am. bor.
stachydantha, Steud. Syn. Pl. Gram. 156 = S. stachydanthus.
stellata, Beauv. Agrost. 16 = Cynodon Dactylon.
stolonifera, Beauv. l. c. = Agrostis stolonifera.
stolonifera, Presl, Cyp. et Gram. Sic. 22 = Agrostis verticillata.
subpyramidata, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 61.—Texas.
subsetacea, Trin. l. c. 133, in nota.—Am. bor.
subtilis, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 88 = S. subtilis.
sylvatica, Beauv. Agrost. 182 = Agrostis alba.
tenacissima, H. B. & K. Nov. Gen. et Sp. i. 138 = S. indicus.
tenera, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 87 = S. serotinus.
tenuissima, Schult. Mant. ii. 479 = S. minutiflorus.
tremula, Trin. Gram. Unifl. 155 = S. tremulus.
trichodes, H. B. & K. Nov. Gen. et Sp. i. 139 = Agrostis trichodes.
tricholepis, Torr. in Pacif. Rail. Rep. iv. 155 = S. tricholepis.
Triniana, Steud. Syn. Pl. Gram. 156 = S. rupestris ?
truncatella, Steud. l. c. 159 = S. truncatella.
utilis, Torr. in Pacif. Rail. Rep. v. 365 = S. depauperatus.
vaginiflora, Torr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 56 = S. vaginaeflorus.
varians, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 89 (Quid ?).—Am. bor.
verticillata, Beauv. Agrost. 16 = Agrostis verticillata.
verticillata, Steud. Syn. Pl. Gram. 158 = S. verticillatus.
Villarsii, Beauv. Agrost. 16 = Agrostis verticillata.
villifera, Steud. Syn. Pl. Gram. 159 = S. ciliatus.

VILFA :—
villosa, Trin. Sp. Gram. iii. t. 252 = Sporobolus villosus.
vinealis, Beauv. Agrost. 182 = Agrostis canina.
Vinzenti, Steud. Syn. Pl. Gram. 155.—Texas.
virginica, Beauv. Agrost. 16 = S. virginicus.
vulgaris, Beauv. l. c. = Agrostis vulgaris.
vulgaris, Wight, ex Steud. Syn. Pl. Gram. 168 = Agrostis Wightii.
xerampelina, Trin. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. v. II. (1840) 103.—Hab. ?

VILLAMILLA, Ruiz & Pav. [Fl. Per. et Chil. Ic. Ined. t. 402] ex Moq. in DC. Prod. xiii. II. 10 (1849).
PHYTOLACCACEAE, Benth. & Hook. f. iii. 81.
TRICHOSTIGMA, A. Rich. Ess. Fl. Cub. ii. 306 (1845).
octandra, *Hook. f. in Benth. & Hook. f. Gen.* iii. 81.—Am. trop.
peruviana, *Hook. f. l. c.*—Peruv.
tinctoria, *Ruiz & Pav.* [*Fl. Per. et Chil. Ic.* Ined. t. 402] *ex Moq. in DC. Prod.* xiii. II. 10.—Peruv.

VILLANOVA, Lag. Gen. et Sp. Nov. 31 (1816).
COMPOSITAE, Benth. & Hook. f. ii. 404.
CHLAMYSPERMA, Less. Syn. Comp. 256 (1832).
UNXIA, H. B. & K. Nov. Gen. et Sp. iv. 279. tt. 401, 402 (1820).
VASQUEZIA, Phil. Fl. Atac. 31. t. 5 (1860).
achillaeoides, *Less. Syn. Comp.* 256.—Mexic.
alternifolia, *Lag. Gen. et Sp. Nov.* 21.—Peruv.
anemonefolia, Less. Syn. Comp. 256 = oppositifolia.
chrysanthemoides, *A. Gray, Pl. Wright.* ii. 96.—N. Mexic.
dissecta, *DC. Prod.* v. 75.—Peruv.
oppositifolia, *Lag. Gen. et Sp. Nov.* 21.—Mexic.
pinnatifida, *Lag. l. c.*—Reg. Magellan.
pratensis, *Benth. & Hook. f. Gen.* ii. 405.—Mexic.
titicacensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 296.—Peruv.

VILLANOVA, Ortega, Hort. Matr. Dec. 47 (1789) = **Parthenium**, Linn. (Compos.).
bipinnatifida, Ortega, l. c. = P. Hysterophorus.

VILLANOVA, Pourr. ex Cutanda, Comp. Fl. Madr. 595 (1861) = **Securinega**, Comm. (Euphorb.).
buxifolia, Pourr. l. c. = S. buxifolia.

VILLARESIA, Ruiz & Pav. Prod. 35 (1794).
OLACINEAE, Benth. & Hook. f. i. 353.
CHARIESSA, Miq. Fl. Ind. Bat. i. 794 (1855).
CITRONELLA, D. Don, in Edinb. N. Phil. Journ. xiii. (1832) 243.
PLEUROPETALON, Blume, Mus. Bot. Lugd. Bat. i. 248 (1850).
VILLAREZIA, Roem. & Schult. Syst. p. xxvii. 331, sphalm. (1819).
Congonha, Miers, *in Ann. & Mag. Nat. Hist.* Ser. III. ix. (1862) 112 ; *Contrib.* ii. 117. t. 69.—Bras.
cuspidata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ix. (1862) 113.—Bras.
dichotoma, *Miers, l. c.* 114.—Bras.
emarginata, *Ruiz & Pav. Syst. Veg.* 64.—Peruv.
Gonghona, C. Muell. in Walp. Ann. vii. 569, sphalm. = V. Congonha.
grandiflora, *Fisch. ex Regel, Gartenfl.* (1856) 61 ; (1857) 1. t. 180.—Bras.
grandifolia, Fisch. & Mey. ex Engl. in Mart. Fl. Bras. xii. II. 54 = megaphylla.
macrocarpa, Scheff. in Ann. Jard. Buitenz. i. (1876) 13 = Gonocarpum macrocarpum.
megaphylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ix. (1862) 114.—Bras.
Moorei, *F. Muell. ex Benth. Fl. Austral.* i. 396.—Austral.
mucronata, Reiss. in Mart. Fl. Bras. xii. II. 75. t. 22 = V. Congonha.
mucronata, Ruiz & Pav. Fl. Per. iii. 9. t. 231 = emarginata.
paniculata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ix. (1862) 114.—Bras.
pungens, *Miers, l. c.* 112.—Chili.
ramiflora, *Miers, l. c.* 116.—Bras.
scandens, Hassk. Retzia, i. 152 = Chailletia timorensis.
Smythii, *F. Muell. Fragm.* v. 156.—Austral.
virescens, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. ix. (1862) 115.—Bras.

VILLAREZIA, Roem. & Schult. Syst. v. p. xxvii. 331 (1819), sphalm. = praec.

VILLARIA, Bally, Notice Hist. sur Villar (1858).—Cf. Bull. Soc. Bot. Fr. v. (1858) 309 = **Villarsia**, Vent. (Gentian.).

VILLARIA, DC. Prod. vi. 542 (1837) = VILARIA, Guett. = **Berardia**, Vill. (Compos.).

VILLARIA, Rolfe, in Journ. Linn. Soc. xxi. (1884) 311. *RUBIACEAE.*
littoralis, *Vidal, Phan. Cuming. Philipp.* 180.—Ins. Philipp.
philippinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 311.—Ins. Philipp.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 180.—Ins. Philipp.

VILLARSIA, J. F. Gmel. Syst. 447 (1791) = **Limnanthemum**, S. T. Gmel. (Gentian.).
aquatica, J. F. Gmel. l. c. = L. lacunosum.

VILLARSIA, Neck. Elem. ii. 110 (1790) = **Cabomba**, Aubl. (Nymph.).

VILLARSIA, Vent. Choix, t. 9 (1803), partim. *GENTIANEAE*, Benth. & Hook. f. ii. 819.
RENEALMIA, Houtt. Handleid. viii. 335. t. 47. f. 1 (1777).
VILLARIA, Bally, Notice Hist. sur Villar (1858).—Cf. Bull. Soc. Bot. Fr. v. (1858) 309.
albiflora, *F. Muell. Fragm.* ii. 21.—Austral.
calthifolia, *F. Muell. l. c.* vi. 140.—Austral.
cambodiana, *Hance, in Journ. Bot.* xv. (1877) 335.—Cambodia.
capitata, *Nees, in Lehm. Pl. Preiss.* i. 365.—Austral.
chilensis, Lodd. Bot. Cab. t. 1994 = parnassifolia?
communis, A. St. Hil. Voy. Distr. Diam. ii. 413 = Limnanthemum Humboldtianum.
congestiflora, *F. Muell. Fragm.* vi. 141.—Austral.
cordata, Ell. Sketch, i. 230 = Limnanthemum lacunosum.
crenata, F. Muell. Fragm. iv. 127 = Limnanthemum crenatum.
Crista-galli, Hook. Fl. Bor. Am. ii. 70; Griseb. Gen. et Sp. Gent. 338 = Menyanthes Crista-galli.
cristata, Spreng. Syst. i. 582 = Limnanthemum cristatum.
eglandulosa, Griff. Ic. Pl. Asiat. t. 385 = Limnanthemum indicum.
exaltata, G. Don, Gen. Syst. iv. 169 = parnassifolia.
exaltata, F. Muell. Fragm. vi. 140 = reniformis.
excelsa, Loisel. Herb. Gén. Amat. v. t. 292 = parnassifolia.
exigua, F. Muell. Fragm. vi. 137 = Limnanthemum exiguum.
exiliflora, F. Muell. l. c. v. 46 = Limnanthemum exiliflorum.
geminata, R. Br. Prod. 457 = Limnanthemum geminatum.
glandulosa, Griff. Notul. iv. 89 = Limnanthemum indicum.
grandifolia, *Fisch. & Mey. in Ind. Sem. Hort. Petrop.* xi. *Suppl.* 78.—Bras.
Gunnii, Hook. f. Fl. Tasm. ii. 368 = Liparophyllum Gunnii.
Humboldtiana, H. B. & K. Nov. Gen. et Sp. iii. 187 = Limnanthemum Humboldtianum.
hydrocharoides, F. Muell. Fragm. vi. 139 = Limnanthemum hydrocharoides.
Hydrophyllum, Roem. & Schult. Syst. iv. 181 = Limnanthemum Hydrophyllum.
indica, Vent. Choix, sub t. 9 = Limnanthemum indicum.
indica, Wall. Cat. n. 4352 = Limnanthemum cristatum.
involucrata, Hook. Ic. Pl. t. 725 = capitata.
lacunosa, Vent. Choix, sub t. 9 = Limnanthemum lacunosum.
lasiosperma, *F. Muell. Fragm.* vi. 137.—Austral.
latifolia, *Benth. in Enum. Pl. Hueg.* 81.—Austral.
macrophylla, Roem. & Schult. Syst. iv. 180 = Limnanthemum indicum.
microphylla, A. St. Hil. Voy. Distr. Diam. ii. 413 = Limnanthemum microphyllum.
minima, F. Muell. Fragm. iv. 128 = Limnanthemum minimum.
minor, *D. Don, ex G. Don, Gen. Syst.* iv. 168.—Ind. or.
nymphaeifolia, Fras. in Hook. Bot. Misc. i. (1830) 257 = Limnanthemum indicum.

VILLARSIA:—
Nymphoides, Vent. Choix, sub t. 9 = Limnanthemum peltatum.
orbiculata, G. Don, Gen. Syst. iv. 169 = Limnanthemum indicum?
ovata, *Vent. Choix*, t. 9.—Afr. austr.
parnassifolia, R. Br. ex Griseb. in DC. Prod. ix. 136 = reniformis.
parnassifolia, *R. Br. Prod.* 457.—Austral.
parviflora, G. Don, Gen. Syst. iv. 169, sphalm. = seq.
parvifolia, Wall. Cat. n. 4351 = Limnanthemum parvifolium.
peltata, Roem. & Schult. Syst. iv. 178 = Limnanthemum peltatum.
platiphylla, A. St. Hil. Voy. Distr. Diam. ii. 413 = Limnanthemum Humboldtianum.
pumila, Hook. Fl. Bor. Am. ii. 70. t. 157; Griseb. Gen. et Sp. Gent. 338 = Hesperochiron californicus.
reniformis, *R. Br. Prod.* 437.—Austral.
reniformis, Nees, in Lehm. Pl. Preiss. i. 364 = albiflora.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. iii. 1031 = Limnanthemum indicum?
rotundifolia, Hort. ex Griseb. in DC. Prod. ix. 136 = parnassifolia.
sarmentosa, Sims, Bot. Mag. t. 1328 = Limnanthemum geminatum.
senegalensis, G. Don, Gen. Syst. iv. 169 = Limnanthemum indicum.
Simsii, G. Don, l. c. = Limnanthemum indicum.
Swartzii, G. Don, l. c. = Limnanthemum indicum.
trachysperma, Ell. Sketch, i. 230 = Limnanthemum trachyspermum.
trachysperma, F. Muell. Fragm. vi. 136 = Limnanthemum indicum.
violifolia, *F. Muell. l. c.* 138.—Austral.

VILLEBRUNEA, Gaudich. Bot. Voy. Bonite, tt. 91, 92 (1844–66). *URTICACEAE*, Benth. & Hook. f. iii. 390.
OREOCNIDE, Miq. Pl. Jungh. i. 39. (1851).
appendiculata, Wedd. in DC. Prod. xvi. 1. 235^{20} = integrifolia.
crenulata, *Gaudich. Bot. Voy. Bonite*, t. 92.—Hab.?
frutescens, *Blume, Mus. Bot. Lugd. Bat.* ii. 168.—Reg. Himal.; China; Japon.
glomerata, *Blume, l. c.* 167.—Java.
integrifolia, *Gaudich. Bot. Voy. Bonite*, t. 91.—Ind. or.; Malaya.
murina, *Blume, Mus. Bot. Lugd. Bat.* ii. 168.—N. Guin.
pilosula, Blume, l. c. = glomerata.
rhodopleura, *Blume, l. c.* (al. gen.?).—N. Guin.
rubescens, Blume, l. c. 167 = scabra.
rufescens, *Blume, l. c.* 166.—N. Guin.
scabra, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 196.—Java.
semicrenata, *Blume, Mus. Bot. Lugd. Bat.* ii. 167.—Ins. Amboin.
sylvatica, Blume, l. c. 167. f. 15 = integrifolia.
trinervis, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 195.—Ins. Philipp.

VILLEMETIA, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 206. t. 9 = WILLEMETIA, Maerkl. = **Chenolea**, Thunb. (Chenop.).

VILMORINIA, DC. Prod. ii. 239 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 499.
multiflora, *DC. l. c.*—Ins. S. Doming.

VILSHENICA, Thou. ex Reichb. Consp. 212^{d} (1828) (Quid?).

VIMEN, P. Br. Hist. Jamaic. 369. t. 22. f. 5 (1756) (Quid?).

VIMEN, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
acuminata, Rafin. l. c. 14 = S. cinerea.
Andersoniana, Rafin. l. c. = S. nigricans.
bicolor, Rafin. l. c. = S. phylicifolia.
cinerea, Rafin. l. c. = S. cinerea.
cotinifolia, Rafin. l. c. = S. nigricans.
discolor, Rafin. l. c. = S. discolor.
Fosteriana, Rafin. l. c. = S. nigricans.

VIMEN:—
hastata, Rafin. Alsog. Am. 14 = Salix hastata.
involucea, Rafin. l. c. (Quid?).—Hab.?
malifolia, Rafin. l. c. = S. hastata.
Muhlenbergiana, Rafin. l. c. = S. tristis?
myricoides, Rafin. l. c. = S. cordata.
parvifolia, Rafin. l. c. = S. repens.
phylicifolia, Rafin. l. c. = S. chlorophylla?
prinoides, Rafin. l. c. = S. brachystachys.
prostrata, Rafin. l. c. = S. repens?
pyrenaica, Rafin. l. c. = S. pyrenaica.
repens, Rafin. l. c. = S. repens.
retusa, Rafin. l. c. = S. retusa.
Russeliana, Rafin. l. c. = S. fragilis.
silesiaca, Rafin. l. c. = S. silesiaca.
Starkeana, Rafin. l. c. = S. livida.
Uva-ursi, Rafin. l. c. = S. Cutleri.
violacea, Rafin. l. c. (Quid?).—Hab.?
Wulfeniana, Rafin. l. c. = S. phylicifolia?

VIMINARIA, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 507. *LEGUMINOSAE*, Benth. & Hook. f. i. 469.
denudata, *Sm. l. c.*—Austral.
juncea, Hoffmgg. Verz. Pfl. 200 = denudata.
lateriflora, Link, Enum. Hort. Berol. i. 403 = Jacksonia scoparia.
Preissii, Meissn. in Lehm. Pl. Preiss. i. 57 (Quid?).—Austral.

VINCA, Linn. Syst. ed. I (1735). *APOCYNACEAE*, Benth. & Hook. f. ii. 703.
CATHARANTHUS, G. Don, Gen. Syst. iv. 95 (1837).
LOCHNERA, Reichb. Consp. 134 (1828).
PERVINCA, Tourn. ex Adans. Fam. ii. 172 (1763).
acutiflora, Bertol. ex Koch, Syn. Fl. Germ. ed. II. 557 = minor.
acutiflora, Bertol. Fl. Ital. ii. 751 = difformis.
alba, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = Tabernaemontana coronaria.
Bottae, Jaub. & Spach, Illustr. ii. 104. t. 186 = libanotica.
difformis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 333.—Lusitan.
elegantissima, Hort. ex Pasq. Cat. Ort. Bot. Nap. 107, nomen (Quid?).—Hab.?
ellipticifolia, Stokes, Bot. Mat. Med. i. 495 = minor.
erecta, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 330.—Turkestan.
foetida, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = Voacanga foetida.
grandiflora, Salisb. Prod. 146 = major.
Gulielmi-Waldemarii, Klotzsch, in Bot. Ergeb. Waldem. Reise, 89 = rosea.
herbacea, *Waldst. & Kit. Pl. Rar. Hung.* i. 8. t. 9.—Europ. or.; As. Min.
humilis, Salisb. Prod. 146 = minor.
intermedia, Tausch, in Flora, xix. (1836) 386 = minor.
lancea, *Boj. ex A. DC. in DC. Prod.* viii. 382.—Madag.
libanotica, *Zucc. in Denkschr. Akad. Muench.* iii. (1840) 247. t. 8.—As. Min.; Persia.
lutea, Linn. Cent. Pl. ii. 12; Amoen. Acad. iv. 309 = Gonolobus carolinensis.
major, Brot. Fl. Lusit. i. 280 = difformis.
major, *Linn. Sp. Pl.* 209.—Reg. Mediterr.
media, Hoffmgg. & Link, Fl. Port. i. 376. t. 70 = difformis.
minor, *Linn. Sp. Pl.* 209.—Europ.
minor, Sm. Fl. Graec. Prod. i. 164 = herbacea.
ovatifolia, Stokes, Bot. Mat. Med. i. 497 = major.
parviflora, Retz. Obs. ii. 14 = pusilla.
pubescens, Urv. in Mém. Soc. Linn. Par. i. (1822) 282 = major.
pumila, Clarke, Trav. iv. 555 = herbacea.
pusilla, *Murr. in Comm. Gotting.* iii. 1772 (1773) 66. t. 2. f. 1.—Ind. or.
rosea, *Linn. Syst.* ed. X. 944.—Cosmop. trop.
Schottii, Mikan, ex Steud. Nom. ed. I. 884 (Quid?).—Hab.?
sessilifolia, A. DC. in DC. Prod. viii. 383 = libanotica.
speciosa, Salisb. Prod. 147 = rosea.
sterilis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 (Nerii sp.).—Java.
trichophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 204.—Madag.

VINCENTIA, Boj. in Hook. Bot. Misc. i. (1830) 293. t. 62 = **Grewia**, Linn. (Tiliac.).
triflora, Boj. = G. triflora.

VINCENTIA, Gaudich. in Freyc. Voy. Bot. 417 (1826) = **Cladium**, P. Br. (Cyperac.).
anceps, Kunth, Enum. Pl. ii. 315 = C. iridifolium.
angustifolia, Gaudich. in Freyc. Voy. Bot. 417.—Ins. Sandvic.
gladiata, Boeck. in Linnaea, xxxviii. (1874) 250 = C. Sinclairii.
latifolia, Kunth, Enum. Pl. ii. 314 = C. iridifolium.
scirpoidea, Boeck. in Linnaea, xxxviii. (1874) 250 = C. scirpoideum.

VINCETOXICUM, Rupp. Fl. Jen. ed. Hall. 25 (1745). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 761.
ALEXITOXICON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67.
AMPHISTELMA, Griseb. Fl. Brit. W. Ind. 417 (1861).
BLYTTIA, Arn. in Mag. Zool. & Bot. ii. (1837) 420.
BUNBURIA, Harv. Gen. S. Afr. Pl. ed. I. 416 (1838).
CYATHELLA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 332. t. 12.
CYNOCTONUM, E. Mey. Comm. Pl. Afr. Austr. 215 (1837).
HAPLOSTEMMA, Endl. Gen. Suppl. iii. 75 (1843).
LYONIA, Ell. Sketch, i. 316 (1817).
ORTHOSIA, Decne. in DC. Prod. viii. 526 (1844).
SCHIZOSTEPHANUS, Hochst. ex Benth. & Hook. f. Gen. ii. 762 (1876).
SEUTERA, Reichb. Consp. 131 (1828).
acanthocarpos, Walt. Fl. Carol. 104 = Gonolobus hirsutus.
acuminatum, *C. Morr. & Decne. in DC. Prod.* viii. 524.—Japon.
albidum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 38 = officinale.
album, Aschers. Fl. Brandenb. i. 421 = officinale.
alpicolum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 39 = officinale.
alpinum, Schott, ex Boiss. Fl. Orient. iv. 56 = fuscatum.
ambiguum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 366.—Japon.
amplexicaule, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 162.—Japon.
amplifolium, C. Koch, in Linnaea, xxiii. (1850) 592 = nigrum.
anatolicum, *Boiss. Diagn.* Ser. I. iv. 39.—As. Min.; Persia; Afghan.
aristolochioides, *Franch. & Sav. Enum. Pl. Jap.* ii. 445.—Japon.
Arnottianum, Wight, Ic. t. 1614 = Cynanchum Arnottianum.
atratum, *Morr. & Decne. in Bull. Acad. Brux.* (1836) 17.—China; Japon.
atrum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 40 = nigrum.
beugesiacum, Jord. & Fourr. l. c. 38 = officinale.
Brandtii, *Franch. & Sav. Enum. Pl. Jap.* ii. 440.—Japon.
canescens, Decne. in DC. Prod. viii. 523 = Cynanchum glaucum.
carnosum, *Benth. Fl. Austral.* iv. 331.—Austral.
chinense, *S. Moore, in Journ. Bot.* xiii. (1875) 228.—China.
contiguum, Gren. & Godr. Fl. Fr. ii. 480 = officinale.
corymbosum, *Benth. & Hook. f. Gen.* ii. 762.—Ind. or.
croaticum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 37 = officinale.
dumeticolum, Jord. & Fourr. l. c. 38 = officinale.
elegans, *Benth. Fl. Austral.* iv. 330.—Austral.
floribundum, *Franch. & Sav. Enum. Pl. Jap.* ii. 444.—Japon.
Fradini, Pomel, Nouv. Mat. Fl. Atl. 82 = officinale.
fruticulosum, *Decne. in DC. Prod.* viii. 525.—Arab.
funebre, Boiss. & Kotschy, Diagn. Ser. I. xi. 79 = medium.
fuscatum, *Reichb. Ic. Fl. Germ.* xvii. 17.—Europ. or.; As. Min.
gonocarpos, Walt. Fl. Carol. 104 = Gonolobus carolinensis.
Guilleminianum, Decne. in DC. Prod. viii. 526 (= *Orthosia virgata*).—Bras.
heterophyllum, *Vatke, in Linnaea*, xl. (1876) 212.—Afr. trop.

VINCETOXICUM:—
hirundinaria, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 404 = officinale.
Huteri, *Vis. & Aschers. in Oestr. Bot. Zeitschr.* xix. (1869) 67.—Europ.
inamoenum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 361.—Manchur.
japonicum, *C. Morr. & Decne. in Bull. Acad. Brux.* (1836) 17.—Japon.
kenouriense, Wight, Contrib. 58 = officinale.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* ii. 440.—Japon.
Kunthii, *Hemsl. Biol. Centr. Am. Bot.* ii. 328.—Mexic.
latifolium, C. Koch, in Linnaea, xxiii. (1850) 591 = medium.
latifolium, *Stev. in Bull. Soc. Nat. Mosc.* xxx. (1857) I. 331.—Tauria.
laxum, C. Koch, in Linnaea, xxiii. (1850) 590, in obs.; Gren. & Godr. Fl. Fr. ii. 480 = officinale.
leptocladum, *Decne. in DC. Prod.* viii. 526.—Am. austr.
leptolepis, *Benth. Fl. Austral.* iv. 331.—Austral.
linifolium, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 79.—Ins. Socotra.
luridum, Stocks, ex Hook. f. Fl. Brit. Ind. iv. 22 = Cynanchum Arnottianum.
luteolum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 39 = officinale.
luteum, *Hoffmgg. & Link, Fl. Port.* i. 381.—Lusitan.
luteum, Sieber, ex Boiss. Fl. Orient. iv. 52 = Cynanchum glaucum.
macrophyllum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 162.—Japon.
mandschuricum, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 227.—Manchuria.
medium, *Decne. in DC. Prod.* viii. 523.—Reg. Cauc.; Persia.
medium, Kostel. Allg. Med.-Pharm. Fl. iii. 1093 = nigrum.
melanthum, *Decne. in DC. Prod.* viii. 526.—Bras.
melanthum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 40 = nigrum.
mexicanum, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 115.—Mexic.
Meyeri, *Benth. & Hook. f. Gen.* ii. 762.—Afr. austr.
microphyllum, *Decne. in DC. Prod.* viii. 525.—Am. trop.
minus, Walp. Rep. vi. 483 = fuscatum.
mongolicum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 356.—Mongol.
montanum, Decne. in DC. Prod. viii. 525 = Cynanchum Arnottianum *vel* Tylophora Govanii.
multinerve, *Franch. & Sav. Enum. Pl. Jap.* i. 319.—Japon.
nigrescens, Jord. ex Nym. Consp. 496 = nigrum.
nigricans, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 133 = melanthum.
nigrum, *Moench, Meth. Suppl.* 313.—Europ. austr.; As. Min.
nikoënse, *Franch. & Sav. Enum. Pl. Jap.* ii. 445.—Japon.
nivale, *Boiss. & Heldr. Diagn.* Ser. I. vii. 20.—Graecia.
ochroleucum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 38 = officinale.
officinale, *Moench, Meth.* 717.—Europ.; Reg. Cauc.
ovatum, *Benth. Fl. Austral.* iv. 330.—Austral.
palustre, *A. Gray, Syn. Fl. N. Am.* ii. I. 102.—Am. bor.
parviflorum, *Decne. in DC. Prod.* viii. 525.—As. Min.
pauciflorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 127.—Japon.
petrophilum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 39 = officinale.
pisidicum, Boiss. & Heldr. Diagn. Ser. I. xi. 79 = fuscatum.
pubens, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. i. (1872–73) 97 = officinale?
puberulum, Timb. l. c. 386 = officinale.
pumilum, *Decne. in DC. Prod.* viii. 525.—Persia.
purpurascens, *C. Morr. & Decne. in Bull. Acad. Brux.* (1836) 17.—Japon.
pyrenaicum, Timb. & Jeanb. ex Nym. Consp. 496 = officinale.
racemosum, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. i. (1872–73) 97 = officinale?
Rehmanni, *Boiss. Fl. Orient.* iv. 53.—Reg. Cauc.
roseum, *Benth. & Hook. f. Gen.* ii. 762.—Dahuria.
rubellum, *Franch. & Sav. Enum. Pl. Jap.* ii. 442.—Japon.

VINCETOXICUM:—
ruscinonensis, Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. i. (1872–73) 97 = officinale?
Rutenbergianum, *Vatke, in Bremen, Abh.* ix. (1885) 126.—Afr. trop.
salinarum, *Benth. & Hook. f. Gen.* ii. 762.—Cuba.
scoparium, *A. Gray, Syn. Fl. N. Am.* ii. I. 102.—Ind. occ.; Mexic.
sepium, *Decne. in DC. Prod.* viii. 526.—Mexic.
sibiricum, *Decne. l. c.* 525.—Sibir.; China.
Sieboldii, *Franch. & Sav. Enum. Pl. Jap.* ii. 444.—Japon.
speciosum, *Boiss. & Heldr. Diagn.* Ser. I. iv. 39.—Graecia.
stelliflorum, Boiss. & Kotschy, l. c. Ser. II. iii. 120 = parviflorum.
sublanceolatum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 367.—Japon.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 444.—Japon.
Thirkeanum, C. Koch, in Linnaea, xxiii. (1850) 591 = speciosum.
tmoleum, *Boiss. Diagn.* Ser. I. iv. 38.—As. Min.
triste, Griseb. Spicil. Fl. Rumel. ii. 68 = speciosum.
undulatum, Heldr. in Boiss. Diagn. Ser. II. vi. 121 = nivale.
Verreyi, *Franch. & Sav. Enum. Pl. Jap.* ii. 438.—Japon.
versicolor, *Decne. in DC. Prod.* viii. 524.—China.
volubile, *Maxim. Prim. Fl. Amur.* 195.—Manchuria.
vulgare, Bernh. Syst. Verz. Erf. 311 = officinale.
Wilfordi, Franch. & Sav. Enum. Pl. Jap. ii. 445 = Cynanchum Wilfordi.

VINDICTA, Rafin. Fl. Tellur. ii. 51 (1836) = **Jeffersonia**, Bart. (Berber.).
begonifolia, Rafin. l. c. = J. binata.

VINSONIA, Gaudich. Bot. Voy. Bonite, tt. 17, 23, 31 (1844–66) = **Pandanus**, Rumph.
acuminata, Gaudich. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 40 = P. acuminatus.
consanguinea, Gaudich. ex Balf. f. l. c. 65 = P. utilis.
drupacea, Gaudich. Bot. Voy. Bonite, t. 31 = P. purpurascens.
elegans, Gaudich. l. c. t. 17 = P. sylvestris.
humilis, Gaudich. l. c. t. 17. f. 12, 13 = P. sylvestris.
lucida, Gaudich. l. c. t. 17. f. 14, 15 = P. sylvestris.
macrostigma, Gaudich. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 65 = P. utilis.
media, Gaudich. ex Balf. f. l. c. = P. utilis.
palustris, Gaudich. Bot. Voy. Bonite, t. 17. f. 18–23 = P. utilis.
Pervilleana, Gaudich. l. c. t. 31. f. 1–7 = P. Bouchéanus.
propinqua, Gaudich. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 65 = P. utilis.
purpurascens, Gaudich. Bot. Voy. Bonite, t. 17 = P. purpurascens.
stephanocarpa, Gaudich. l. c. t. 23. f. 2–5, 7, 8 = P. utilis.
striata, Gaudich. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 65 = P. utilis.
sylvestris, Gaudich. Bot. Voy. Bonite, t. 17. ff. 16, 17 = P. Barkleyi.
Thouarsii, Gaudich. ex Balf. f. in Journ. Linn. Soc. xvii. (1878) 65 = P. utilis.
utilis, Gaudich. Bot. Voy. Bonite, t. 17. f. 1–5 = P. utilis.

VINTENATIA, Cav. Ic. iv. 28. t. 349. f. 1 (1797), *postea* VENTENATIA, l. c. 79 = **Astroloma**, R. Br. et **Melichrus**, R. Br. (Epacrid.).
humifusa, Cav. l. c. = A. humifusum.
procumbens, Cav. l. c. = M. rotatus.

VINTERA, Humb. & Bonpl. Pl. Aequin. i. t. 58 (1805) = WINTERA, Humb. & Bonpl. = **Drimys**, Forst. (Magnol.).

VINTICENA, Steud. Nom. ed. II. ii. 769 (1841) = **Grewia**, Linn. (Tiliac.).
triflora, Steud. l. c. = G. triflora.

VIOLA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 267 (1737). *VIOLARIEAE*, Benth. & Hook. f. i. 117.
CHRYSION, Spach, Hist. Vég. Phan. v. 509 (1836).
DISCHIDIUM, Reichb. Handb. 269 (1837).

VIOLA:—
Erpetion, DC. ex Sweet, Brit. Flow. Gard. ii. t. 170 (1826).
Ion, Medic. Malv. 102 (1787).
Jonium, Reichb. ex Steud. Nom. ed. II. i. 815 (1840).
Lophion, Spach, Hist. Vég. Phan. v. 516 (1836).
Mnemion, Spach, l. c. 510 (1836).
abortiva, Jord. Pugill. Pl. Nov. 14 = hirta.
abyssinica, *Steud. ex Oliver*, *Fl. Trop. Afr.* i. 105. —Abyss.
acaulis, Roem. & Schult. Syst. v. 389, in syn. = cornuta.
acuminata, Ledeb. Fl. Ross. i. 252 = canina.
acuta, Bigel. Fl. Bost. ed. II. 95 = primulifolia.
adriatica, *Freyn, in Flora*, lxvii. (1884) 679.—Europ. austr.
adulterina, Godr. Thèse Hybrid. 18 = alba.
adulterina ×, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 108.—Europ.
aduncа, Sm. in Rees, Cyclop. xxxvii. n. 63 = canina.
aetolica, *Boiss. & Heldr. Diagn.* Ser. II. vi. 24.—Graecia.
affinis, Le Conte, in Ann. Lyc. N. York, ii. (1828) 138 = cucullata.
agrestis, Jord. Obs. Pl. Crit. ii. (1846) 15 = tricolor.
alata, *Burgersdyk, in Miq. Pl. Jungh.* 121.—Java.
alba, *Bess. Prim. Fl. Galic.* i. 171.—Europ.
albiflora, Link, Enum. Hort. Berol. i. 241 = canadensis.
algida, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 391 = Lysipomia reniformis.
alleghanensis, Roem. & Schult. l. c. 360 = sagittata.
Allionii, Pio, Viola Spec. Bot. Med. 20 = arenaria.
alpestris, *Hegetschw. Fl. Schw.* 245.—Helvet.
alpestris, Jord. Obs. Pl. Crit. ii. (1846) 32, in textu = tricolor.
alpina, All. Fl. Pedem. ii. 98 = cenisia.
alpina, Huds. Fl. Angl. ed. I. 331 = canina.
alpina, *Jacq. Enum. Stirp. Vindob.* 159, 291.—Carinth.
alpina, Ruiz & Pav. ex Ging. in DC. Prod. i. 300 = pygmaea.
alpina, Salisb. Prod. 130 = calcarata.
altaica, [*Ker-Gawl.*] *Bot. Reg.* t. 54; *Pall. ex Sims, Bot. Mag.* t. 1776.—As. Min.; Reg. Cauc.; Sibir.
ambigua, Barc. Fl. Balear. 53 (Quid?).—Ins. Balear.
ambigua, Waldst. & Kit. Pl. Rar. Hung. ii. 208 = campestris.
amoena, T. F. Forst. in Symons, Syn. 198 = lutea.
amoena, Le Conte, in Ann. Lyc. N. York, ii. (1828) 144 = blanda, *Willd.*
angustifolia, Banks, ex Ging. in DC. Prod. i. 307 = Ionidium heterophyllum.
angustifolia, *Phil. in Linnaea*, xxviii. (1856) 612.—Chili.
apetala, Gilib. Fl. Lituan. ii. 124 = mirabilis.
apetala, Roxb. Hort. Beng. 18; Fl. Ind. i. 650 = Patrinii.
apetala, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 61 = canina.
appendiculata, Steud. Nom. ed. II. ii. 770 = occulta.
aprica, Spreng. Syst. i. 800 = arenaria.
arborea, *Forsk. Fl. Aegypt. Arab.* p. cxx.—Arab.
arborea, Linn. Syst. ed. X. 1239 = Corynostylis Loeflingii.
arborescens, *Linn. Sp. Pl.* 935.—Reg. Mediterr. occ.
arborescens, Loefl. ex Ging. in DC. Prod. i. 289 = Corynostylis Loeflingii.
arcuata, *Blume, Bijdr.* 58.—Java.
arenaria, *DC. Fl. Fr.* iv. 806.—Europ.; As. bor.; Am. bor.
arenicola, Chabert, in Bull. Soc. Bot. Fr. xviii. (1871) 196 = sylvestris.
arguta, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 391.—Am. austr.
armena, *Boiss. & Huet, Diagn.* Ser. II. v. 48.—Reg. Cauc.; Persia.
arvatica, Jord. Pugill. Pl. Nov. 24 = tricolor.
arvensioides, *Strobl, in Oestr. Bot. Zeitschr.* xxvii. (1877) 224.—Italia.
arvensis, Murr. Prod. Stirp. Gotting. 73 = tricolor.
asarifolia, Pursh, Fl. Am. Sept. ii. 732 = cucullata.
asarifolia, Schreb. ex Muhl. in Trans. Am. Phil. Soc. iii. (1793) 178 = canina.
aspera, Ging. in DC. Prod. i. 295 = serpens.
Asterias, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 145. t. 99.—Chili.
atlantica, Pomel, Nouv. Mat. Fl. Atl. 215 = tricolor.

VIOLA:—
atropurpurea, *Leybold, in Anal. Univ. Chil.* (1858) 159.—Chili.
atropurpurea, Rafin. ex Ging. in DC. Prod. i. 291 = pedata.
attenuata, Steud. Nom. ed. I. 884 = Ionidium oppositifolium.
attenuata, Sweet, Hort. Brit. ed. III. 54 = lanceolata.
Aubletii, G. F. W. Mey. Prim. Fl. Esseq. 123 = Corynostylis Hybanthus.
aurantiaca, *Leybold, in Flora*, xlvii. (1864) 41.—Chili.
aurasiaca, Pomel, Nouv. Mat. Fl. Atl. 342 = gracilis.
aurea, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 185. f. 54.—Calif.
Auricula, *Leybold, in Flora*, xlvii. (1864) 40.—Chili.
auriculata, Leybold, in Anal. Univ. Chil. (1863) II. 672; ex Phil. f. Cat. Pl. Vasc. Chil. 20 = praec.
austriaca, A. & J. Kern. in Ber. Nat. Verh. Innsbr. iii. (1872) 71 = odorata.
Averyi, Kellogg, in Pacif. Rural Press (May 1879); ex Curran, in Bull. Calif. Acad. i. III. (1885) 131 = Patrinii.
badensis ×, Wiesb. in Oestr. Bot. Zeitschr. xxiv. (1874) 225.—Europ.
Balbisii, Re, Fl. Segus. 73 = arenaria.
balsaminoides, Gardn. in Hook. Ic. Pl. t. 217 = cerasifolia.
banatica, Kit. ex Roem. & Schult. Syst. v. 382 = tricolor.
barbata, Muhl. Cat. 25; Willd. ex Roem. & Schult. Syst. v. 356 = cucullata.
Barroetana, *Schaffn. ex Hemsl. Diagn. Pl. Nov.* 20.—Mexic.
Baumgarteniana, Schur, Enum. Pl. Transs. 87 = alpina.
bavarica, *Schrank, Baier. Fl.* i. 491.—Bavaria.
Beckwithii, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 119. t. 1.—Am. bor. occ.
begoniaefolia, *Benth. Pl. Hartw.* 161.—Peruv.
bella, Gren. ex Gren. & Godr. Fl. Fr. i. 184 = tricolor.
Beraudii, Bor. Fl. Centr. Fr. ed. III. ii. 76 = sepincola.
Berteroniana, Steud. in Flora, xxxix. (1856) 422 = V. Portalesia.
Bertoloni, Pio, Viola Spec. Bot. Med. 34 = gracilis.
Bertolonii, *De Salis, in Ann. Sc. Nat.* Sér. II. v. (1836) 116.—Corsica.
Besseri, *Rupr. Fl. Cauc.* 151.—Reg. Cauc.
betonicaefolia, Boj. ex Baker, Fl. Maurit. 10 = Patrinii.
betonicifolia, *Sm. in Rees, Cycl.* xxxvii. n. 7.—Austral; Ins. Norfolk.
bicolor, Gilib. Fl. Lituan. ii. 123; Pursh, Fl. Am. Sept. i. 175 = tricolor.
Bielziana, *Schur, Enum. Pl. Transs.* 86.—Transsylv.
biflora, *Linn. Sp. Pl.* 936.—Reg. bor. temp.
Billotiana, F. Schultz, ex Nym. Consp. 77 = seq.
Billotii, F. Schultz, in Flora, xxiii. (1840) 120 = persicifolia.
Bisseti, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) I. 5.—Japon.
biturigensis, Bor. ex Déségl. in Mém. Soc. Acad. Maine-et-Loire, xiv. (1863) 104 (Quid?).—Gallia.
blanda, Salisb. Prod. 130 = cornuta?
blanda, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 166. —Montenegr.
blanda, *Willd. Hort. Berol.* t. 24.—Am. bor.
Blumii, Korth. in Miq. Ann. Mus. Bot. Lugd. Bat. iii. 76 = serpens.
borealis, Weinm. in Linnaea, x. (1835–36) 66 = Selkirkii.
brachyceras, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 301 = blanda.
brachypetala, *C. Gay, Fl. Chil.* i. 212.—Chili.
brachysepala, *Maxim. Prim. Fl. Amur.* 50.—Reg. Amur.
brevicaulis, Raeusch. Nom. ed. III. 258 (Quid?).—Siles.
Brooksii, *Kellogg, in Calif. Hortic.* ix. (1879) 281; *ex Bot. Calif.* ii. 433.—Calif.
Broussonetiana, *Roem. & Schult. Syst.* v. 371.—Ins. Canar.
Bruneauniana, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 946 = odorata.
Bubanii, Timb. in Congr. Sc. Fr. xix. (1852) 280 = tricolor.
Buchaniana, DC. ex D. Don, Prod. Fl. Nep. 206 = serpens.
bulbosa, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 334.—China.
Bungeana, *Rupr. Fl. Cauc.* 147, in textu.—Reg. Cauc.

VIOLA:—
Burgersdijkii, *Oudem. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 77.—Java.
bustillosia, *C. Gay, Fl. Chil.* i. 211.—Chili.
buxifolia, Poir. Encyc. viii. 646 = Ionidium buxifolium.
caespitosa, D. Don, Prod. Fl. Nep. 205 = Patrinii.
caespitosa, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 701.—Hispan.
calabra, Huet, ex Nym. Consp. 77 = arenaria.
calamaria, Steud. Nom. ed. II. ii. 770, sphalm. = seq.
calaminaria, Lej. Rev. Fl. Spa, 49 = tricolor.
calcarata, *Linn. Sp. Pl.* 935.—Europ.
calcarata, Munby, Fl. Algér. 99 = gracilis.
calcarata, Sibth. & Sm. Fl. Graec. Prod. i. 147 = Clementiana.
calcarata, Vill. Hist. Pl. Dauph. i. 238; ii. 666; Robs. Brit. Fl. 107 = tricolor.
Calceolaria, Linn. Sp. Pl. ed. II. 1327 = Ionidium Ipecacuanha.
Caleyana, *G. Don, Gen. Syst.* i. 329.—Austral.
callosa, *Benth. Hartw.* 161, in nota.—Peruv.
calthaefolia, *Poir. Pl. Encyc.* viii. 627.—Hab.?
calycina, *Boiss. & Heldr. Diagn.* Ser. I. viii. 53.—Pisidia.
campestris, *Bieb. Fl. Taur. Cauc.* i. 171.—Europ.; As. Min.
canadensis, *Linn. Sp. Pl.* 936.—Am. bor.
canariensis, Berth. in Mém. Soc. Linn. Par. v. (1827) 420 = teydea.
canescens, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 450 = serpens.
canina, Cham. & Schlecht. in Linnaea, ii. (1827) 146 = sylvestris.
canina, *Linn. Sp. Pl.* 935.—Reg. bor. temp.
cano-barbata, *Leybold, in Flora*, xlix. (1866) 285.—Chili.
capensis, Thunb. Prod. Pl. Cap. 40 = Ionidium capense.
capillaris, *Pers. Syn.* i. 256.—N. Granat.
cassinensis, *Strobl, in Oestr. Bot. Zeitschr.* xxvii. (1877) 228.—Ital.
caulescens, Molina, ex Phil. f. Cat. Pl. Vasc. Chil. 20 = rubella?
celtica, Sauzé & Maillard, ex Nym. Consp. 78 = canina.
cenisia, Lag. & Rodr. in Anal. Cienc. v. (1802) 270 = nevadensis.
cenisia, *Linn. Sp. Pl.* ed. II. 1325.—Europ. austr.
cenisia, Schangin, in Pall. N. Nord. Beitr. vi. 103 = tricolor.
cerasifolia, *A. St. Hil. Pl. Rem. Brés.* 277.—Bras.
chamaedrifolia, Ruiz & Pav. ex Ging. in DC. Prod. i. 306 = rubella.
Chamaedrys, *Leybold, in Flora*, xlvii. (1864) 41.—Chili.
Chamissoniana, *Ging. in Linnaea*, i. (1826) 408.—Ins. Sandvic.
cheiranthifolia, *Humb. & Bonpl. Pl. Aequin.* i. 111. t. 32.—Ins. Canar.
chelmea, *Boiss. Diagn.* Ser. II. i. 54.—Graecia.
chinensis, G. Don, Gen. Syst. i. 322 = Patrinii.
Christii, *Wolf, in Bull. Soc. Murith.* x. (1881) 43.—Helvet.
chrysantha, Fisch. ex Steud. Nom. ed. II. ii. 770 = altaica.
chrysantha, *Hook. Ic. Pl.* t. 49.—Am. bor. occ.
chrysantha, *Phil. in Linnaea*, xxxiii. (1864–65) 15.—Chili.
chrysantha ×, Schrad. ex Reichb. Fl. Germ. Excurs. 709.—Europ.
ciliata, Muhl. Cat. ed. I. 25 = sagittata.
ciliata, *Schlecht. in Linnaea*, xii. (1838) 202.—Mexic.
ciliata, Willd. ex Roem. & Schult. Syst. v. 360 = cucullata.
cinerascens, Kern. in Oestr. Bot. Zeitschr. xviii. (1868) 20 = arenaria.
cinerea, *Boiss. Diagn.* Ser. I. i. 7.—Arab.; Persia.
circaeoides, Steud. Nom. ed. I. 885 = Ionidium circaeoides.
clandestina, Pursh, Fl. Am. Sept. i. 173 = blanda.
claytonioides, Steud. Nom. ed. I. 885 = Ionidium claytonioides.
Clementiana, *Boiss. Diagn.* Ser. II. i. 55.—Macedon.
collina, *Bess. Enum. Pl. Volh.* 10.—Europ.
Commersonii, *DC. ex Ging. in DC. Prod.* i. 297.—Reg. Magellan.
Comollia, *Messara, Prod. Fl. Valtell.* 208.—Ital.
concolor, Forst. in Trans. Linn. Soc. vi. (1802) 309 = Ionidium concolor.
conferta, A. St. Hil. Pl. Rem. Brés. 279 = cerasifolia.

VIOLA:—
confinis, Jord. ex Nym. Consp. 80 = tricolor.
confusa, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 260 = serpens.
congener, Le Conte, in Ann. Lyc. N. York, ii. (1828) 140 = cucullata.
congesta, *Gill. in Hook. Bot. Misc.* iii. (1833) 144. t. 97.—Peruv.
Conilii, *Franch. & Sav. Enum. Pl. Jap.* ii. 285.—Japon.
consanguinea, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 166.—Dalmat.
consimilis, Jord. Pugill. Pl. Nov. 18 = odorata.
conspersa, Reichb. Pl. Crit. i. 44 = canina.
contempta, Jord. Pugill. Pl. Nov. 24 = tricolor.
corallensis, *Phil. in Linnaea*, xxxiii. (1864–65) 16.—Chili.
corchorifolia. *Domb. ex Ging. in DC. Prod.* i. 304.—Peruv.
cordata, Walt. Fl. Carol. 219 = cucullata.
cordata, Willd. Enum. Hort. Berol. 264 = Broussonetiana.
cordifolia, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 114 = arenaria.
cordifolia, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 62 = cucullata.
cornuta, Desf. Fl. Atlant. ii. 313 = gracilis.
cornuta, *Linn. Sp. Pl.* ed. II. 1325.—Europ. austr.
corralensis, *Phil. in Linnaea*, xxxiii. (1864–65) 16.—Chili.
corsica, Jord. in Mém. Soc. Acad. Maine-et-Loire, i. (1857) 86 (nomen) = seq. ?
corsica, Nym. Syll. 228 = Bertoloni ii.
Cotyledon, *Ging. in DC. Prod.* i. 300.—Chili.
crassifolia, *Fenzl, Pugill. Pl. Nov. Syr.* 12.—As. Min.
crassiuscula, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 16 = nevadensis.
cretica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 51.—Ins. Creta.
cucullata, *Ait. Hort. Kew.* ed. I. iii. 288.—Am. bor.
cucullata, Bigel. ex Ging. in DC. Prod. i. 294 = palustris.
cucullata, Moç. & Sessé, ex Ging. l. c. 289 = Corynostylis Berterii.
cucullifolia, Wender. ex Steud. Nom. ed. II. ii. 771 (Quid ?).—Hab. ?
cuneata, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 290.—Am. bor.
Cunninghamii, *Hook. f. Fl. N. Zel.* i. 16.—Tasman.; N. Zel.
Curtisii, E. Forst. in Engl. Bot. Suppl. t. 2693 = tricolor.
cyanea, Čelak. in Oestr. Bot. Zeitschr. xxii. (1872) 349 = odorata.
dactyloides, *Roem. & Schult. Syst.* v. 351.—Sibir.
debilis, Michx. Fl. Bor. Am. i. 150 = striata.
debilis, Pursh, Fl. Am. Sept. i. 174 = canina.
declinata, *Waldst. & Kit. Pl. Rar. Hung.* iii. 248.—Hungar.
decora, Kellogg, in Pacif. Rural Press (May 1879); ex Curran, in Bull. Calif. Acad. i. III. (1885) 131 = Beckwithii.
decumbens, J. F. Gmel. Syst. 413 = *brevicaulis*.
decumbens, *Linn. f. Suppl.* 397.—Afr. austr.
decumbens, Moench, Meth. Suppl. 73 = calcarata.
degener, Schweigg. ex Mert. & Koch, Deutschl. Fl. ii. 264 = sylvestris.
Dehnhardtii, Tenore, Ind. Sem. Hort. Neap. (1830) 12; ex ej. Syll. 117 = odorata.
delphinantha, *Boiss. Diagn.* Ser. I. i. 7.—Graecia.
delphinensis, Jord. Obs. Pl. Crit. vii. (1849) 3 = sciaphila.
delphinifolia, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 136.—Am. bor.
Demetria, Prolongo, ex Boiss. Voy. Bot. Espagne, ii. 73 = tricolor.
deminuta, Schur, Enum. Pl. Transs. 87 = tricolor.
dentata, Pursh, Fl. Am. Sept. i. 172 = sagittata.
derelicta, Jord. in Billot, Annot. Fl. Fr. et Allem. 101 = tricolor.
Deseglisei, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 82 = tricolor.
diandra, Murr. Syst. ed. XIII. 669 = Corynostylis diandra.
dichotoma, [Moç. & Sessé,] ex Ging. in DC. Prod. i. 297 = scandens.
dichroa, *Boiss. Diagn.* Ser. II. v. 48.—Armen.
diffusa, *Ging. in DC. Prod.* i. 298.—Reg. Himal.; China.

VIOLA:—
diffusa, Vell. Fl. Flum. viii. t. 163 = cerasifolia.
digitata, Pursh, Fl. Am. Sept. i. 171 = pedata.
digitata, Willd. ex Roem. & Schult. Syst. v. 352 = dactyloides.
dissecta, Ledeb. Fl. Alt. i. 255 = pinnata.
distans, *Wall. Cat.* n. 4022.—Ind. or.; Malaya.
Dombeyana, *DC. ex Ging. in DC. Prod.* i. 305.—Am. austr.
Domeikoana, *C. Gay, Fl. Chil.* i. 220.—Chili.
Douglasii, Steud. Nom. ed. II. ii. 771 = chrysantha, *Hook.*
Drymeia, Schur, Enum. Pl. Transs. 79 = hirta.
dubia, Wall. Cat. n. 1450 = serpens ?
dumetorum, Jord. Pugill. Pl. Nov. 16 = alba.
dumetorum, *Phil. in Linnaea*, xxviii. (1856) 612.—Chili.
ebracteolata, *Fenzl, Illustr. Taur.* 49.—Syria; Persia.
edulis, Spach, Hist. Vég. Phan. v. 508 = palmata.
Einseleana, F. Schultz, ex Nym. Consp. 77 = Schultzii.
elata, Gueldenst. Reisen, i. 421; ex Ledeb. Fl. Ross. i. 258 (Quid ?).—Rossia.
elatior, *Fries, Mant.* iii. 126.—Europ.; As. bor.; Oriens.
elatior, Wimm. Fl. Schles. (1841) 46 = canina.
elegantula, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 167 = gracilis.
elongata, Huet, ex Nym. Consp. 80 = gracilis.
elongata, *Poir. Encyc.* viii. 644.—'Am.'
emarginata, Le Conte, in Ann. Lyc. N. York, ii. (1828) 142 = sagittata.
emirnensis, Boj. ex Oliver, Fl. Trop. Afr. i. 105 = abyssinica.
enneasperma, Linn. Sp. Pl. 937 = Ionidium heterophyllum.
epipsila, Fries, ex Nym. Consp. 79 = suecica.
epipsila, Ledeb. Ind. Sem. Hort. Dorpat. (1820) 5 = palustris.
erecta, Gilib. Fl. Lituan. ii. 124 = elatior.
erecta, Roth, Nov. Pl. Sp. 165 = Ionidium heterophyllum.
ericetorum, Schrad. ex Link, Enum. Hort. Berol. i. 240 = sylvestris.
eriocarpa, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 75 = pubescens.
esculenta, Ell. Sketch, i. 300 = palmata.
etbaica, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 665.—Afr. trop.
Eugeniae, Parl. in Nuov. Giorn. Bot. Ital. vii. (1875) 68 = calcarata.
excelsa, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 391 = Corynostylis Hybanthus.
excisa, Hance, in Journ. Bot. vi. (1868) 296 = verecunda.
exigua, *Vell. Fl. Flum.* viii. 164.—Bras.
exilis, *Phil. in Linnaea*, xxviii. (1856) 672.—Chili.
Falconeri, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 185.—Reg. Himal.
fennica, Nyland. Spicil. Pl. Fenn. Cent. i. 28 = palustris.
fennica, Steph. ex Nym. Consp. 79 = uliginosa.
ferrariensis, Campana, Cat. Pl. Ferrar.; ex Pio, Viola Spec. Bot. Med. 22 = canina.
filicaulis, *Hook. f. Fl. N. Zel.* i. 16.—N. Zel.
filiformis, Ruiz & Pav. ex DC. Prod. i. 310 = Ionidium glutinosum.
fimbriata, *Steud. in Flora*, xxxix. (1856) 423.—Chili.
fimbriatula, Sm. in Rees, Cyclop. xxxvii. n. 16 = sagittata.
Fischeri, Sweet, Hort. Brit. ed. II. 44 = Gmeliniana.
flabellifolia, Lodd. Bot. Cab. t. 777 = pedata.
flagelliformis, *Hemsl. Diagn. Pl. Nov.* 20.—Mexic.
flavicans, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 292.—Bolivia.
flavicornis, E. Forst. ex Nym. Consp. 77 = sylvestris.
flavicornis, Sm. Engl. Fl. i. 304 = canina.
flavida, Jord. Pugill. Pl. Nov. 25 = tricolor.
flavida, *Schur, in Verh. Naturf. Ver. Bruenn*, xv. II. (1877) 119.—Europ.
flexuosa, Moretti, ex Roem. & Schult. Syst. v. 370 = persicifolia.
floribunda, Jord. Pugill. Pl. Nov. 19 = odorata.
Flos-Idae, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 5.—Reg. Argent.
Flos-Mariae, *Hieron. l. c.* 8.—Reg. Argent.
foliosa, Čelak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 477 = odorata.
formosa, Vukot. in Rad Jugos. Akad. Zagreb. lxvi. (1883) 20 = sylvestris.
Foucaudi, Sav. Cat. Pl. Charente-Inf. 49 = tricolor.

VIOLA:—
Foudrasi, Jord. Obs. Pl. Crit. vii. (1849) 4 = hirta.
fragrans, *Sieber, Reise Kreta*, ii. 320.—Ins. Creta.
fragrans, Wiesb. ex Sabransky, in Oestr. Bot. Zeitschr. xxxiv. (1884) 133 = *Wiesbaurii*.
fraterna, *Hegetschw. Fl. Schw.* 241.—Helvet.
frigida, *Phil. Fl. Atac.* 9.—Chili.
frutescens, Roth, Nov. Pl. Sp. 166 = Ionidium suffruticosum.
frutescens, Ruiz & Pav. ex Ging. in DC. Prod. i. 308 = Ionidium oppositifolium.
fusiformis, Sm. in Rees, Cyclop. xxxvii. n. 9 = Gmeliniana.
Ganderi, Hausm. ex Nym. Consp. 79 = campestris.
garganica, Strobl, in Oestr. Bot. Zeitschr. xxvii. (1877) 224 = tricolor.
gibbosa, Rafin. in Am. Monthly Mag. (1811) 356, nomen = hastata.
glabella, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 142.—Am. bor. occ.
glacialis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 49. t. 165.—Chili.
glandulifera, *Hook. Ic. Pl.* t. 67.—Ecuador.
glandulifera, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 452 = serpens.
glandulosa, Domb. ex Ging. in DC. Prod. i. 297 = maculata.
glauca, Bieb. Fl. Taur. Cauc. iii. 165 = arenaria.
glaucescens, Oudem. in Miq. Ann. Mus. Bot. Lugd. Bat. iii. 74 = distans.
glechomoides, *Leybold, in Flora*, xlvii. (1864) 40.—Chili.
glutinosa, Poir. Encyc. viii. 647 = Ionidium glutinosum.
Gmeliniana, *Roem. & Schult. Syst.* v. 354.—Sibir.
gracilescens, Jord. Obs. Pl. Crit. ii. (1846) 20 = tricolor.
gracilis, Bivona, Stirp. Rar. Sicil. iv. 25 = calcarata.
gracilis, R. Br. ex Ging. in DC. Prod. i. 305 = hederacea.
gracilis, Clem. Sert. Orient. 16 = Clementiana.
gracilis, Martr. ex Nym. Consp. 88 = tricolor.
gracilis, Reichb. Fl. Germ. Excurs. 709 = declinata.
gracilis, *Sibth. & Sm. Fl. Graec. Prod.* i. 146.—Graecia.
gracillima, *A. St. Hil. Pl. Rem. Brés.* 275.—Bras.
Grahami, *Benth. Pl. Hartw.* 35.—Mexic.
grandiflora, Griseb. Spicil. Fl. Rumel. i. 238 = Clementiana.
grandiflora, 'Linn.' Herb. ex Steud. Nom. ed. II. ii. 771 = altaica.
grandiflora, Linn. Mant. ii. 518 = tricolor.
grandifolia, Wall. Cat. n. 1447 = Patrinii.
granulosa, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 292.—Bolivia.
Grayi, Franch. & Sav. Enum. Pl. Jap. ii. 288 = sylvestris.
Griffithiana, Boiss. Diagn. Ser. II. i. 54 = serpens.
Grisebachiana, *Vis. in Mem. Ist. Venet.* x. (1861) 436.—Serbia.
Gronovii, Willd. ex Roem. & Schult. Syst. v. 359, in textu = primulifolia.
grypoceras, A. Gray, in Perry, Jap. Exped. ii. 308 = sylvestris.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 133 = Ionidium heterophyllum.
gymnocarpa, Janka, in Linnaea, xxx. (1859–60) 559 = collina.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1872) 377.—Am. bor. occ.
Hamiltoniana, D. Don, Prod. Fl. Nep. 206 = serpens.
hastata, *Michx. Fl. Bor. Am.* ii. 149.—Am. bor.
hebecarpa, Schur, Enum. Pl. Transs. 79 = hirta.
hederacea, *Labill. Nov. Holl. Pl.* i. 66. t. 91.—Austral.
Heldreichiana, *Boiss. Diagn.* Ser. I. viii. 53.—As. Min.
helvetica ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 72.—Europ.
heterophylla, Bertol. Rar. Ital. Pl. Dec. iii. 53 = gracilis, calcarata.
heterophylla, Muhl. Cat. 25 = palmata.
heterophylla, Poir. Encyc. viii. 646 = Ionidium heterophyllum.
heterosepala, Boiss. & Heldr. Diagn. Ser. II. vi. 23 = aetolica.
hirsuta, Crantz, Inst. ii. 430; Roem. & Schult. Syst. v. 361 = hirta.
hirta, *Linn. Sp. Pl.* 934.—Europ.; As. bor.

VIOLA:—
hirta, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 50 = hastata.
hirtaeformis ×, Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 190.—Europ.
hirtipes, *S. Moore, in Journ. Linn. Soc.* xvii. (1879) 379. t. 16.—China.
hirto-alba, Gren. & Godr. Fl. Fr. i. 176 = hirta.
hispida, Lam. Fl. Fr. ii. 679 = rothomagensis.
hispida, Lapeyr. Hist. Abr. Pl. Pyr. 123 = lutea.
Hladnikii, Reichb. ex Nym. Consp. 77 = Schultzii.
Holubyi, *Schur, in Verh. Naturf. Ver. Bruenn*, xv. II. (1877) 107.—Europ.
Hookeri, *Thoms. ex Hook. f. Fl. Brit. Ind.* i. 183.—Reg. Himal.
Hookeriana, *H. B. & K. Nov. Gen. et Sp.* v. 369.—Mexic.
Hornemanniana, Roem. & Schult. Syst. v. 370 = canina.
hortensis, Schur, Enum. Pl. Transs. 82 = odorata.
Huidobrii, *C. Gay, Fl. Chil.* i. 214.—Chili.
Humboldtii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 121.—N. Granat.
humifusa, *Vukot. in Rad Jugos. Akad. Zagreb.* lxvi. (1883) 16.—Europ.
humilis, *H. B. & K. Nov. Gen. et Sp.* v. 369.—Mexic.
Hybanthus, Aubl. Pl. Gui. ii. 811. t. 319 = Corynostylis Hybanthus.
Hybanthus, Linn. Sp. Pl. ed. II. 1325 = Corynostylis Loeflingii.
hybrida, Krock. Fl. Siles. Suppl. i. 291 (Quid?).—Europ.
hybrida, Schur, Enum. Pl. Transs. 79 = odorata.
hybrida, Wulf. ex Roem. & Schult. Syst. v. 357 = uliginosa.
hydrocotyloides, *Armst. in Trans. N. Z. Inst.* xiv. (1882) 360.—N. Zel.
hymettia, Boiss. & Heldr. Diagn. Ser. II. i. 57 = tricolor.
ignobilis, *Rupr. Fl. Cauc.* 148.—Reg. Cauc.
imberbis, Ledeb. Fl. Alt. i. 257 = Selkirkii.
incisa, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 302.—Sibir.
incomparabilis, *Schur, in Verh. Naturf. Ver. Bruenn*, xv. II. (1877) 99.—Europ.
incompta, Jord. Obs. Pl. Crit. vii. (1849) 11 = odorata.
inconspicua, *Blume, Cat. Gew. Buitenz.* 57; *Bijdr.* 58.—Java.
ingrica, *Bieb. ex Rupr. in Beitr. Pfl. Russ. Reich.* iv. 48.—Rossia.
inodora, Gilib. Fl. Lituan. ii. 123 = palustris.
insularis, Gren. & Godr. Fl. Fr. i. 178 = sylvestris.
insularis, Gren. & Godr. l. c. 185 = Bertolonii.
integerrima, *Phil. in Linnaea*, xxviii. (1856) 672.—Chili.
integerrima, *Spreng. Pugill.* i. 21.—Hab.?
integrifolius, Dulac, Fl. Hautes-Pyr. 179 = cenisia.
intermedia, Krock. Fl. Siles. Suppl. i. 289 (Quid?).—Europ.
intermedia, Lej. Rev. Fl. Spa, 51 = rothamagensis.
intricata, *Arv. in Bull. Soc. Dauph.* vii. (1880) 264.—Gallia.
Ipecacuanha, Linn. Mant. ii. 484 = Ionidium Ipecacuanha.
ircutiana, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 298.—Sibir.
italica, Voigt, ex Vilm. Fl. Pl. Terre, ed. III. 1206 = odorata.
Itoubou, Aubl. Pl. Gui. ii. 808 = Ionidium Ipecacuanha.
Jacquiniana, Steud. Nom. ed. I. 885 = Ionidium Jacquinianum.
japonica, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 357 (sp. dub.).—Java (cult.).
japonica, *Langsd. ex Ging. in DC. Prod.* i. 295.—Japon.
Jaubertiana, *Marés & Vigin. Cat. Pl. Baléar.* 37. t. 2.—Ins. Balear.
Jooi, *Janka, in Oestr. Bot. Wochenbl.* vii. (1857) 198.—Transsylv.
jucunda, Jord. Pugill. Pl. Nov. 20 = odorata.
Julia ×, Bruegg. in Zeitschr. Ferdinand. Tirol. III. ix. (1860) 62.—Europ.
Juressi, Link, ex Steud. Nom. ed. I. 885 (Quid?).—Hab.?
kalksburgensis ×, Wiesb. in Oestr. Bot. Zeitschr. xxvii. (1877) 153.—Europ.
kamtschatica, Ging. in Linnaea, i. (1826) 406 = Selkirkii.

VIOLA:—
kauaaënsis, *A. Gray, Bot. U. St. Expl. Exped.* i. 85.—Ins. Sandvic.
kauensis, A. Gray, in Proc. Am. Acad. ii. (1852) 325 = praec.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 153.—Japon.
Kerneri ×, Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 189.—Europ.
Kitaibeliana, Roem. & Schult. Syst. v. 383 = tricolor.
Kochii, *Kirschl. in Mém. Strasb.* iii. n. II. (1840) 12. t. 1.—Europ.
Krokeri, J. F. Gmel. Syst. 412 = arenaria.
kunawurensis, *Royle, Illustr. Bot. Himal.* 75. t. 18. f. 3.—Reg. Himal.
labiosa, *Sm. in Rees, Cycl.* xxxvii. n. 48.—Austral.
labradorica, Schrank, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 12 = canina.
laciniosa, *A. Gray, in Perry, Jap. Exped.* ii. 308.—Japon.
lactea, Koch & Ziz, Cat. Pl. Palat. 6; Fries, Nov. Fl. Suec. 86 = pratensis.
lactea, Reichb. Fl. Germ. Excurs. 707 = persicifolia.
lactea, *Sm. Fl. Brit.* i. 247.—Europ.
lamiifolia, Pritz. Ic. Ind. 1163 = *lancifolia*, Thore.
lancea, Poir. Encyc. viii. 635 = pratensis.
lanceifolia, Schum. & Thonn. Beskr. Guin. Pl. 134 = Ionidium enneaspermum?
lanceolata, *Linn. Sp. Pl.* 934.—Am. bor.
lanceolata, Pall. Reise, iii. 223 = Gmeliniana.
lancifolia, Bess. Prim. Fl. Galic. i. 173 = canina.
lancifolia, Bieb. Fl. Taur.-Cauc. iii. 165 = Ruppii.
lancifolia, Steud. Nom. ed. II. ii. 772, sphalm. = Corynostylis Hybanthus.
lancifolia, Thore, Chlor. Land. 357 = lactea.
Langsdorfii, *Fisch. ex Ging. in DC. Prod.* i. 296.—Sibir. or.; Am. bor. occ.
latistipula, *Hemsl. Diagn. Pl. Nov.* 21.—Mexic.
laurifolia, Sm. in Rees, Cycl. xxxvii. n. 58 = Corynostylis Hybanthus.
Lechlerii, *Griseb. in Goett. Abh.* vi. (1854) 116.—Chili.
Leconteana, G. Don, Gen. Syst. i. 324 = blanda.
Lejeunii, Jord. Pugill. Pl. Nov. 27 = tricolor.
lepida, Jord. l. c. 28 = tricolor.
leptoceras, Schur, Enum. Pl. Transs. 87 = tricolor.
leucantha, Schur, l. c. 82 = odorata.
Lewisiana, Ging. in DC. Prod. i. 298 = striata.
Leyboldiana, *Phil. in Linnaea*, xxxiii. (1864–65) 16.—Chili.
libanotica, *Boiss. Diagn.* Ser. I. viii. 52.—Syria.
lilascens, Heldr. ex Boiss. Fl. Orient. i. 466 = tricolor.
linariifolia, Vahl, Eclog. Am. ii. 18 = Ionidium strictum.
Lindeniana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 386.—N. Granat.
lineariaefolia, Poir. Encyc. viii. 648 = Ionidium strictum.
linearifolia, Vahl, in West, Bidr. Ste-Croix, 303 = Ionidium strictum.
lineata, Ortega, Hort. Matr. Dec. 49 = Ionidium lineatum.
linguaefolia, Nutt. in Torr. & Gray, Fl. N. Am. i. 141 = Nuttallii.
linifolia, Juss. in Poir. Encyc. viii. 647 = Ionidium linifolium.
litoralis, *Phil. Fl. Atac.* 9.—Chili.
litoralis, Spreng. Nov. Prov. 43 = pratensis.
livida, Kit. ex Roem. & Schult. Syst. v. 363 = arenaria.
Lloydii, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 80 = tricolor.
lobata, *Benth. Pl. Hartw.* 298.—Calif.
Loefflingiana, Steud. Nom. ed. I. 885 = Ionidium Ipecacuanha.
longepedunculata, *Franch. & Sav. Enum. Pl. Jap.* ii. 286.—Japon.
longicaulis, Krock. Fl. Siles. Suppl. i. 295 (Quid?).—Europ.
longifolia, R. Br. ex Ging. in DC. Prod. i. 294 = betonicifolia.
longifolia, Poir. Encyc. viii. 649 = Noisettia longifolia.
longipes, Nutt. in Torr. & Gray, Fl. N. Am. i. 140 = canina.
longiscapa, DC. ex G. Don, Gen. Syst. i. 322 = betonicifolia.
longiviola, Pourr. ex Roem. & Schult. Syst. v. 380 = arborescens.
lucorum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 698 = canina.

VIOLA:—
ludovicea, Jan, ex Ging. in DC. Prod. i. 303 = prostrata.
lusitana, Brot. Fl. Lusit. i. 306 = lactea.
lutea, Comm. ex Ging. in DC. Prod. i. 297 = maculata.
lutea, Huds. Fl. Angl. ed. I. 331; Sm. Fl. Brit. i. 248 = tricolor.
lutea, Lam. Fl. Fr. ii. 680 = biflora.
lutea, *Vell. Fl. Flum.* viii. t. 162.—Bras.
luteola, Jord. Pugill. Pl. Nov. 27 = tricolor.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 16.—N. Zel.
macedonica, *Boiss. & Heldr. Diagn.* Ser. II. i. 56.—Macedon.
macrantha, Huet, ex Nym. Consp. 81 = calcarata.
macroceras, *Bunge, in Ledeb. Fl. Alt.* i. 256.—Sibir.
macrostipulata, Schur, Enum. Pl. Transs. 84 = Schultzii.
maculata, *Cav. Ic.* vi. 20. t. 539.—Chili.
maculata, Patrin, ex Ging. in DC. Prod. i. 293 = variegata.
maderensis, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 36.—Ins. Mader.
madonia, Presl, ex Reichb. Pl. Crit. iii. 60 = calcarata.
magellanica, *Forst. f. in Comm. Gotting.* ix. (1789) 41. t. 8.—Reg. Magellan.
magellensis, Porta & Rigo, ex Strobl, in Oestr. Bot. Zeitschr. xxvii. (1877) 228 = cenisia.
magniflora, Molina, Sagg. Chile, ed. II. 283 = magellanica.
marina, Vilm. Fl. Pl. Terre, ed. III. 205 = Campanula Medium.
maritima, Spreng. ex Mert. & Koch, Deutschl. Fl. ii. 272 = tricolor.
martia, Gilib. Fl. Lituan. ii. 122; S. T. Gmel. Reise, i. 65 = odorata.
Martii ×, Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 190.—Europ.
Massoni, Banks, ex Ging. in DC. Prod. i. 308 = Ionidium capense.
Mauritii, *Teplouchlow, in Bull. Soc. Oural. Amat. Sc. Nat.* vii. III. 37.—Rossia.
maviensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 150.—Ins. Sandvic.
media, Chaillet, ex Roem. & Schult. Syst. v. 382 = tricolor.
meduanensis, Bor. Fl. Centr. Fr. ed. III. ii. 80 = tricolor.
melanosperma, Bert. ex Ging. in DC. Prod. i. 308 = Ionidium oppositifolium.
mendanca, Vell. Fl. Flum. viii. t. 160 = Schweiggeria fruticosa.
mentita, Jord. in Billot, Annot. Fl. Fr. et Allem. 101 tricolor.
Metziana, Hohen. ex Hook. f. Fl. Brit. Ind. i. 184 = distans.
micrantha, *Albert, in Bull. Soc. Sc. Draguign.* (1884) (reimpr. 6).—Gallia.
micrantha, Bertol. Misc. Bot. i. 15 = tricolor.
micrantha, Presl, Delic. Prag. 27 = parvula.
micrantha, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 183 = canina.
micranthella, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 291.—Bolivia.
microceras, *Rupr. Beitr. Pfl. Russ. Reich.* ii. 22. t. 1. f. 2.—Sibir.
microceras, Schur, Enum. Pl. Trans. 80 = collina.
microphylla, *Phil. in Linnaea*, xxviii. (1856) 611.—Chili.
microphylla, Schur, in Oestr. Bot. Zeitschr. viii. (1858) 394 = alpina.
microphylla, Willd. ex Roem. & Schult. Syst. v. 391 = Ionidium glutinosum.
microphyllos, *Poir. Encyc.* viii. 628.—Reg. Magellan.
Miersii, Bert. ex Steud. Nom. ed II. ii. 772 = V. Asterias.
Minae, Strobl, in Oestr. Bot. Zeitschr. xxvii. (1877) 225 = calcarata.
minuta, *Bieb. Fl. Taur. Cauc.* i. 173.—Reg. Cauc.
mirabilis, *Linn. Sp. Pl.* 936.—Europ.; As. bor.
mixta, *Kern. in Oestr. Bot. Zeitschr.* xviii. (1868) 21.—Europ.
modesta, Ball, in Journ. Bot. xi. (1873) 300 = tezensis.
modesta, *Fenzl, Pugill. Pl. Nov. Syr.* 12.—Syria.
mongolica, *Franch. Pl. Davidian.* i. (1884) 42.—China.
monopetala, Roem. & Schult. Syst. v. 400 = Ionidium filiforme.
Montagnei, *C. Gay, Fl. Chil.* i. 222.—Chili.
montana, DC. Fl. Fr. iv. 807 = elatior.
montana, Kellogg, in Proc. Calif. Acad. i. ed. II. (1873) 55 = lobata.

VIOLA:—
montana, Ledeb. Fl. Alt. i. 261 = pratensis.
montana, Linn. Sp. Pl. 935 = canina.
montana, Plan. Ens. Fl. Gallega, 206 = persicifolia.
monticola, Jord. Obs. Pl. Crit. ii. (1846) 37 = tricolor.
Mucusii, Orph. ex Nym. Consp. 80 = tricolor.
Muhlenbergiana, Ging. in DC. Prod. i. 297 = canina.
Muhlenbergii, Torr. Fl. U. St. i. 256 = canina.
multicaulis, Jord. Pugill. Pl. Nov. 15 = odorata.
multifida, Mill. Gard. Dict. ed. VIII. n. 5 = pedata.
multifida, Willd. ex Roem. & Schult. Syst. v. 352 = pinnata.
Munbyana, Boiss. & Reut. Pugill. Pl. Nov. 15 = gracilis.
muscoides, *Phil. in Linnaea*, xxviii. (1856) 672.—Chili.
mutabilis, Rochel, Pl. Banat. Rar. (Index) = tricolor.
mysurensis, Heyne, ex Wall. Cat. n. 1446 = Patrinii.
nana, Godr. in Mém. Soc. Sc. Nat. Cherb. xix. (1875) 166 = tricolor.
Nannei, *Polak. in Linnaea*, xli. (1877) 547.—Costa Rica.
nebrodensis, Presl, Delic. Prag. 26 = calcarata.
neglecta, Bieb. Fl. Taur. Cauc. i. 172 = sylvestris.
neglecta, F. W. Schmidt, Fl. Boëm. iii. 55 = canina.
nemausensis, Jord. Obs. Pl. Crit. ii. (1846) 18 = tricolor.
nemoralis, Jord. Pugill. Pl. Nov. 21 = sylvestris.
nemoralis, Kuetz. in Linnaea, vii. (1832) 46 = Ruppii.
nepaulensis, DC. ex D. Don, Prod. Fl. Nep. 205 = Patrinii.
nephrodes, Rafin. in Am. Monthly Mag. (1811) 356, nomen (Quid?).—Am. bor.
nevadensis, *Boiss. Elench.* 26.—Hispan.
Nicolai, Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 4 = Orphanidis.
nigricans, Schur, ex Nym. Consp. 78 = alba.
nilagirica, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 556 = Patrinii.
nitens, Host, ex Reichb. Fl. Germ. Excurs. 705 = uliginosa.
nivalis, *Benth. Pl. Hartw.* 160.—Bolivia.
nivalis, *Phil. in Linnaea*, xxviii. (1856) 611.—Chili.
nivalis, *Vest, ex Roem. & Schult. Syst.* v. 365.—Hab.?
Notoniana, Wall. Cat. n. 1449 = distans.
nubigena, *Leybold, in Flora*, xlvii. (1864) 40.—Chili.
nummularia, Vukot. in Rad Jugos. Akad. Zagreb. lxvi. (1883) 16 = nummularifolia?
nummulariaefolia, Vill. Prosp. 26 = seq.?
nummularifolia, *All. Fl. Pedem.* ii. 98. t. 9. f. 4.—Europ.
nummularifolia, Panč. Verz. Serb. Phan. n. 438 = Grisebachiana.
nummularifolia, Schleich. ex Link, Enum. Hort. Berol. i. 240 = arenaria.
nummularifolia, F. W. Schmidt, Fl. Boëm. iii. 47 = canina.
Nuttallii, Benth. Pl. Hartw. 298 = aurea.
Nuttallii, *Pursh, Fl. Am. Sept.* i. 174.—Am. bor. occ.
obliqua, Ait. Hort. Kew. ed. I. iii. 288 = cucullata.
obliqua, Pursh, Fl. Am. Sept. i. 172 = blanda.
obliquifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 337.—Venezuela.
obscura, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 10 = collina.
occulta, Hort. Argent. ex Otto, in Ind. Sem. Hort. Hamb. (1853) 8 = tricolor.
occulta, *Lehm. l. c.* (1829).—*Cf. Linnaea*, v. (1830) *Litt.* 46.—As. occ. et bor.
ocellata, *Torr. & Gray, Fl. N. Am.* i. 142.—Calif.
ochroleuca, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 69 = striata.
odontocalycina, *Boiss. Fl. Orient.* i. 461.—Armen.
odora, Neck. Delic. Gallo-Belg. ii. 367 = seq.
odorata, *Linn. Sp. Pl.* 934.—Europ.; Afr. et As. bor.
odorata, Thunb. Fl. Jap. 326 = japonica, *Langsd.*
oetolica, Boiss. Fl. Orient. i. 464, err. typ. = aetolica.
officinalis, Crantz, Inst. ii. 431 = odorata.
olimpia, Beggiat. in Att. Soc. Ital. Sc. Nat. viii. (1865) 174. t. 1 = olympica?
olonnensis, Genev. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 178 = tricolor.
olympica, *Boiss. Diagn.* Ser. II. i. 55.—As. Min.
Opizii, Knaf, ex Nym. Consp. 78 = collina.
oppositifolia, Linn. Sp. Pl. ed. II. 1327 = Ionidium oppositifolium.
orbelica, *Panč. Elem.* 16.—Europ. austr.

VIOLA:—
orbiculata, *Geyer, ex A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 11.—Am. bor. occ.
Orbignyana, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 353.—Chili.
orchidiflora, Rudge, Pl. Gui. i. 11. t. 10 = Noisettia longifolia.
Oreades, Bieb. Fl. Taur. Cauc. iii. 167 = altaica.
Orphanidis, *Boiss. Fl. Orient.* i. 464.—Macedon.
orthoceras, *Ledeb. Fl. Ross.* i. 258.—Reg. Cauc.
ovata, Nutt. Gen. Am. i. 148 = sagittata.
ovata, Rafin. ex Ging. in DC. Prod. i. 293 = primulifolia.
pachyrhiza, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 84.—Japon.
pachyrrhiza, *Boiss. & Hohen. Diagn.* Ser. I. vi. 21.—Kurdistan.
Paillouxi, Jord. Obs. Pl. Crit. ii. (1846) 36 = tricolor.
Pallasii, Fisch. ex Steud. Nom. ed. I. 886 = altaica.
pallens, Forst. ex Roem. & Schult. Syst. v. 359 = blanda.
pallescens, Jord. Obs. Pl. Crit. ii. (1846) 10 = tricolor.
pallida, Salisb. Prod. 130 = hirta.
pallidiflora, Huet, ex Nym. Consp. 81 = calcarata.
palmaesis, Knowles & Westc. Flor. Cab. ii. (1838) 165 = palmensis.
palmaris, Buch.-Ham. ex Ging. in DC. Prod. i. 298 = distans.
palmata, *Linn. Sp. Pl.* 933.—Am. bor.
palmata, Patrin, ex Ging. in DC. Prod. i. 293 = dactyloides.
palmensis, *Webb & Berth. Phyt. Canar.* i. 112, in syn.—Ins. Canar.
paludosa, [Clairv.] Man. Herb. 67 = seq.
palustris, *Linn. Sp. Pl.* 934.—Reg. bor. temp.
papilionacea, Pursh, Fl. Am. Sept. i. 173 = cucullata.
papillaris, Boiss. & Hohen. Diagn. Ser. I. viii. 52 = spathulata.
paradoxa, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 28.—Ins. Mader.
pariaetariaefolia, Banks, ex Ging. in DC. Prod. i. 308 = Ionidium oppositifolium.
Parkeri, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 261 = diffusa.
parmensis, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 946 = odorata.
parviflora, Kit. ex Roem. & Schult. Syst. v. 383 = tricolor.
parviflora, Linn. f. Suppl. 360 = Ionidium glutinosum.
parvifolia, *Benth. Pl. Hartw.* 161.—Ecuador.
parvifolia, Roem. & Schult. Syst. v. 391 = Ionidium glutinosum.
parvula, Opiz, Boehm. Gew. (1823) 32; Natural. n. vi. (1824) 45 = hirta.
parvula, *Tineo, Pl. Rar. Sicil. Pugill.* i. 5.—Europ. austr.; Syria.
Patrinii, *Ging. in DC. Prod.* i. 293.—Rossia; As. bor.; Reg. Himal.
pedata, *Linn. Sp. Pl.* 933.—Am. bor.
pedatifida, G. Don, Gen. Syst. i. 320 = delphinifolia?
peduncularis, Langsd. ex Ging. in DC. Prod. i. 297 = Langsdorfii.
pedunculata, *Torr. & Gray, Fl. N. Am.* i. 141.—Am. bor. occ.
pedunculata, Torr. in Pacif. Rail. Rep. iv. 68, partim = aurea.
pensylvanica, Michx. Fl. Bor. Am. ii. 149 = pubescens.
pentadactyla, *Fenzl, Pugill. Pl. Nov. Syr.* 12.—Syria.
peregrina, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 82 = tricolor.
perennis, Miegév. in Bull. Soc. Bot. Fr. xii. (1865) 340 = odorata?
perennis, Vilm. Fl. Pl. Terre, ed. I. 949 = tricolor.
perexigua, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 326.—N. Zel.
permixta ×, Jord. Obs. Pl. Crit. vii. (1849) 6.—Europ.
perplexa, Gremli, Excurs. Schweiz, ed. II. 106 = odorata.
persicaefolia, Kirschl. in Mém. Strasb. iii. n. ii. (1840) 13 = persicifolia?
persicariaefolia, Poir. Encyc. viii. 628 = stipularis.
persicifolia, Ledeb. Fl. Alt. i. 260 = elatior.
persicifolia, *Roth, Tent. Fl. Germ.* ii. i. 271.—Europ.
Pesneaui, Lloyd, Fl. Ouest Fr. ed. III. 43 = rothomagensis.
phalacrocarpa, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 318.—Japon.

VIOLA:—
philippica, *Cav. Ic.* vi. 19. t. 529. f. 2.—Ins. Philipp.
Philippii, Leybold, in Anal. Univ. Chil. (1859) 681 = microphylla.
phyteumaefolia, DC. ex G. Don, Gen. Syst. i. 322 = betonicifolia.
pichinchensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 335.—Ecuador.
picta, *Gillot, in Bull. Soc. Bot. Fr.* xxxii. (1885) 241.—Gallia.
pilosa, Blume, Bijdr. 57 = serpens.
pilosa, Pers. Syn. i. 255 = rothomagensis.
pinnata, *Linn. Sp. Pl.* 934.—Europ.; As. bor.
pinnata, Richards. in Frankl. 1st Journ. App. 734 = pedata.
pinnatifida, *All. Misc. Taur.* iii. t. 5.—Europ.
podolica, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 124, nomen (Quid?).—Europ.
poëtica, *Boiss. & Sprun. Diagn.* Ser. I. vi. 21.—Graecia.
polychroma, *Kern. Sched. Fl. Exsicc.* ii. 89.—Austria.
polygalaefolia, Poir. Encyc. viii. 647 = Ionidium polygalaefolium.
polypoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 555.—Chili.
porphyrea, Uechtritz, ex Garcke, Fl. Deutschl. ed. XIII. 51 = sciaphila.
Portalesia, *C. Gay, Fl. Chil.* i. 209. t. 6.—Chili.
portulacea, *Leybold, in Flora*, xlviii. (1865) 381.—Chili.
praecox, Heuff. ex Fuss, Fl. Transs. 81 = alba.
praemorsa, Dougl. in Bot. Reg. t. 1254 = aurea.
pratensis, *Mert. & Koch, Deutschl. Fl.* ii. 267.—Europ.
primulifolia, *Linn. Sp. Pl.* 934.—Am. bor.
primulifolia, Lour. Fl. Cochinch. 513 = Patrinii.
primulifolia, Pursh, Fl. Am. Sept. i. 173 = sagittata.
primulifolia, Schult. Obs. Bot. 48 = uliginosa.
prionantha, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 82 = Patrinii.
procera, Pall. ex Ledeb. Fl. Ross. i. 251 = elatior.
procumbens, Gilib. Fl. Lituan. ii. 123 = canina.
procumbens, J. F. Gmel. Syst. 413 = decumbens.
propera, Jord. Obs. Pl. Crit. vii. (1849) 6 = collina.
propinqua, Jord. Pugill. Pl. Nov. 18 = odorata.
prostrata, *Roem. & Schult. Syst.* v. 382.—Ital.; Ins. Canar.
Provostii, Bor. Fl. Cent. Fr. ed. III. ii. 82 = tricolor.
prunellaefolia, Fisch. ex Ging. in DC. Prod. i. 293 = Patrinii.
prunellaefolia, *H. B. & K. Nov. Gen. et Sp.* v. 370.—N. Granat.
prunifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 391 = Ionidium anomalum.
psammophila, *Phil. in Linnaea*, xxxiii. (1864–65) 14.—Chili.
Pseudo-canina, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 113 = canina.
Pseudo-gracilis, Strobl, in Oestr. Bot. Zeitschr. xxvii. (1877) 227 = gracilis.
Pseudo-lutea, Schur, Enum. Pl. Transs. 87 = tricolor.
Pseudo-suavis, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 103 = odorata.
pteropoda, *Hemsl. Diagn. Pl. Nov.* 20.—Mexic.
puberula, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 102.—Hispan.
pubescens, *Ait. Hort. Kew.* ed. I. iii. 290.—Am. bor.
pulchella, Salisb. Prod. 130 = canina.
pumila, Chaix, in Vill. Hist. Pl. Dauph. i. 339 = pratensis.
pumila, Fries, ex Nym. Consp. 78 = canina.
pumila, Schur, in Verh. Naturf. Ver. Bruenn, xv. ii. (1877) 114 = arenaria.
punctata, Humb. ex Roem. & Schult. Syst. v. 391 = Polygala glandulosa.
punctata, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 67 = canina.
punicea, Ruiz & Pav. ex Ging. in DC. Prod. i. 304 = corchorifolia.
purpurascens, F. W. Schmidt, Fl. Boëm. iii. 49 = arenaria.
purpurea, Kellogg, in Proc. Calif. Acad. i. ed. II. (1873) 55 = aurea.
purpurea, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) i. 310, in textu.—Reg. Cauc.
pusilla, *Hook. & Arn. ex Hook. Bot. Misc.* iii. (1833) 145.—Chili.
pusilla, Schleich. Cat. Pl. Helv. ed. IV. 61 = arenaria.
pusillima, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 291.—Bolivia.

VIOLA:—
pycnophylla, *Franch. & Sav. Enum. Pl. Jap.* ii. 285. —Japon.
pygmaea, *Juss. ex Poir. Encyc.* viii. 630.—Peruv.
pyrenaica, *Ram. ex DC. Fl. Fr.* iv. 803.—Mont. Pyren.
pyrenaica, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 697 = Willkommii.
pyrolaefolia, Poir. Encyc. viii. 636 = maculata.
Raddeana, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) II. 463, 501.—Reg. Amur.; China.
radicans, DC. ex Ging. in DC. Prod. i. 297 = canina.
ramosissima, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 336.—Ecuador.
ranunculifolia, Juss. ex Poir. Encyc. viii. 626 = palmata.
recta, Garcke, Fl. Deutschl. ed. III. 47 = persicifolia.
Reichenbachiana, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 78 = sylvestris.
renifolia, *A. Gray, in Proc. Am. Acad.* viii. (1870) 288.—Am. bor.
reniformis, *R. Br. ex Ging. in DC. Prod.* i. 305.—Austral.
reniformis, Wall. Cat. n. 1443 = biflora.
repens, Buch.-Ham. ex Ging. in DC. Prod. i. 296 = serpens.
repens, Domb. ex Ging. l. c. 305 = Dombeyana.
repens, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 70 = striata.
repens, Turcz. in Bull. Soc. Nat. Mosc. (1838) 88 = palustris.
reptans, Presl, ex Reichb. Fl. Germ. Excurs. 705 = odorata.
resupinata, Moench, Meth. 230 = cucullata.
reticulata, *Schlecht. in Linnaea*, xii. (1838) 201.—Mexic.
revoluta, Heuff. ex Reichb. Fl. Germ. Excurs. 705 = campestris.
rhombifolia, *Leybold, in Anal. Univ. Chil.* (1859) 680. —Chili.
riparia, Hartm. f. Handb. 128; ex Nym. Syll. Suppl. 41 = uliginosa.
riphaea, J. F. Gmel. Syst. 413; Roem. & Schult. Syst. v. 368, in textu = arenaria.
Riviniana, Reichb. Pl. Crit. i. 81 = sylvestris.
Rogoviczii, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 125, nomen (Quid?).—Rossia.
rostrata, *Muhl. Cat.* 26; *Pursh, Fl. Am. Sept.* i. 174. —Am. bor.
rosulata, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 49. t. 166. —Chili.
rothomagensis, *Desf. Tabl.* ed. I. 153.—Europ.
rotundifolia, Hook. Lond. Journ. Bot. vi. (1847) 73 = sarmentosa.
rotundifolia, *Michx. Fl. Bor. Am.* ii. 150.—Am. bor.
Roxburghiana, Voigt, Hort. Suburb. Calc. 76 = Patrinii.
Royleana, Wall. Cat. n. 1448 = serpens.
rubella, *Cav. Ic.* vi. 20. t. 351. f. 1.—Chili.
rubella, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 144 = V. Portalesia.
rubra, Aubl. ex Steud. Nom. ed. II. ii. 773 = hirta.
rupestris, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1791) 60 = arenaria.
Ruppii, *All. Fl. Pedem.* ii. 99.—Europ.
Ruppii, J. & C. Presl, Fl. Čech. 53 = pratensis.
Ruppii, Reichb. Fl. Germ. Excurs. 707 = canina.
ruralis, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 81 = tricolor.
sabulosa, Bor. Not. 3; Dum. in Bull. Soc. Bot. Belg. vii. (1868) 353 = tricolor.
sagittaefolia, Salisb. Prod. 130 = seq.
sagittata, *Ait. Hort. Kew.* ed. I. iii. 287.—Am. bor.
Sagoti, Jord. Obs. Pl. Crit. ii. (1846) 34 = tricolor.
salina, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 299 = Selkirkii.
sarmentosa, Bieb. Fl. Taur.-Cauc. i. 172 = odorata.
sarmentosa, Burgersdijk, in Miq. Pl. Jungh. 120 = Burgersdijkii.
sarmentosa, *Dougl. in Hook. Fl. Bor. Am.* i. 80.—Am. bor. occ.
saxatilis, Hoffmgg. ex Roem. & Schult. Syst. v. 380 = saxicola.
saxatilis, F. W. Schmidt, Fl. Boëm. iii. 60 = tricolor.
saxicola, *Roem. & Schult. Syst* v. 380.—Reg. Mediterr. occ.
saxigena, Schur, Enum. Pl. Transs. 84 = arenaria.
scabra, Braun, in Flora. iii. (1820) 469 = hirta?
scandens, Bert. ex DC. Prod. i. 289 = Corynostylis Berterii.

VIOLA:—
scandens, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 391; *H. B. & K. Nov. Gen. et Sp.* v. 371. t. 493. —Mexic.
scanica, Fries, Summa Veg. Scand. 35, 153 = palustris.
scardica, Nym. Consp. 81 = cenisia.
scaturiginosa, Wallr. Sched. Crit. 97 = uliginosa.
Schleideniana, *Bogenh. Taschenb. Fl. Jena*, 162.—Europ.
Schmidtiana, Roem. & Schult. Syst. v. 363 = canina.
Schultzii, *Billot, ex F. W. Schultz, in Flora*, xxiii. (1840) 121.—Europ.
sciaphila, Joo, ex Janka, in Oestr. Bot. Wochenbl. vii. (1857) 198 = Jooi.
sciaphila, *Koch, Syn. Fl. Germ.* ed. II. 90.—Europ.
scorpiuroides, *Coss. in Bull. Soc. Bot. Fr.* xix. (1872) 80.—Tripoli.
scotophylla, Jord. Obs. Pl. Crit. vii. (1849) 9 = odorata.
Scouleri, Dougl. ex Steud. Nom. ed. II. ii. 773 = canadensis.
scrotiformis, *DC. ex Ging. in DC. Prod.* i. 299.—Afr. austr.
segetalis, Jord. Obs. Pl. Crit. ii. (1846) 12; Schur, Enum. Pl. Transs. 85 = tricolor.
Selkirkii, *Pursh, ex Goldie, in Edinb. Phil. Journ.* vi. (Apr. 1822) 324.—Reg. bor. et frigid.
semperflorens, Hort. ex Steud. Nom. ed. I. 887 = odorata.
Sempervivum, *C. Gay, Fl. Chil.* i. 226.—Chili.
sepincola, *Jord. Obs. Pl. Crit.* vii. (1849) 8.—Europ.
septemloba, Le Conte, in Ann. Lyc. N. York, ii. (1828) 141 = palmata.
sequoiensis, Kellogg, in Proc. Calif. Acad. ii. (1863) 185 = lobata.
serpens, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 449.—Ind. or.; Malaya; China.
serpyllifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 391.—Am. austr.
setosa, *Sm. in Rees, Cycl.* xxxvii. n. 60.—Chili.
Sheltonii, *Torr. in Pacif. Rail. Rep.* iv. 67. t. 2.—Calif.
sicula, Schur, Enum. Pl. Transs. 86 = Bielziana.
Sieberi, *Hook. Comp. Bot. Mag.* i. (1835) 274.—Austral.
Sieberiana, Spreng. Syst. iv. Cur. Post. 96 = hederacea.
Sieboldi, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 320.—Japon.
silvatica, Fries, ex Hartm. f. in Bot. Notiser (1841), 81 = sylvestris.
somchetica, C. Koch, in Linnaea, xv. (1841) 251 = uliginosa.
soongorica, Fisch. ex Ledeb. Fl. Ross. i. 257 = tricolor.
sororia, Willd. Hort. Berol. i. 72 = cucullata.
spathulata, *Willd. ex Roem. & Schult. Syst.* v. 353.—Reg. Casp.; Persia.
speciosa, *Pantoč. in Oestr. Bot. Zeitschr.* xxiii. (1873) 79.—Serbia.
speciosa, Schrad. ex Reichb. Fl. Germ. Excurs. 708 = altaica.
spectabilis, C. Richt. in Oestr. Bot. Zeitschr. xxxv. (1885) 419 = odorata.
Sprengeliana, Steud. Nom ed. I. 887 = Ionidium concolor.
spuria ×, Čelak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 478.—Europ.
stagnina, Kit. in Schult. Oestr. Fl. ed. II. i. 426 = persicifolia.
stellata, Miers, Trav. ii. 531 = V. Asterias.
Steveni, Bess. ex Faucon. Excurs. Bot. Valais, 96 = odorata.
stipularis, Cav. Ic. vi. 21 = capillaris, setosa.
stipularis, *Sw. Prod. Veg. Ind. Occ.* 117.—Ind. occ.
Stocksii, *Boiss. Fl. Orient.* i. 453.—Afghan.; Beluch.
stolonifera, *Rodrig. in Bull. Soc. Bot. Fr.* xxv. (1878) 238.—Ins. Balear.
striata, *Ait. Hort. Kew.* ed. I. iii. 290.—Am. bor.
striata, Hook. Lond. Journ. Bot. vi. (1847) 72 = glabella.
stricta, Hornem. Hort. Hafn. ii. 958, Add. = Ruppii.
stricta, Muhl. in Trans. Am. Phil. Soc. iii. (1793) 178; et ex Pursh, Fl. Am. Sept. i. 175 = Ionidium concolor.
stricta, Poir. Encyc. viii. 648 = Ionidium strictum.
suaveolens, Perrier & Songeon, in Billot, Annot. Fl. Fr. et Allem. 183 = ambigua.
suaveolens, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 107 = odorata?
suaveolens ×, Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 190.—Europ.
suavis, Bieb. Fl. Taur. Cauc. iii. 164 = odorata.

VIOLA:—
suavis, Fisch. ex Ging. in DC. Prod. i. 294 = Gmeliniana.
suavissima, Jord. Pugill. Pl. Nov. 21 = odorata.
subcarnea, Jord. l. c. 17 = odorata.
subdimidiata, *A. St. Hil. Pl. Rem. Brés.* 276.—Bras.
suberosa, Desf. Fl. Atlant. ii. 313 = arborescens.
subtilis, Jord. in Billot, Annot. Fl. Fr. et Allem. 101 = tricolor.
sudetica, Willd. Enum. Hort. Berol. Suppl. 12 = tricolor.
suecica, *Fries, Summa Veg. Scand.* 556.—Europ. bor.
suffruticosa, Linn. Sp. Pl. 937 = Ionidium heterophyllum.
sulfurea, Cariot, Etud. des Fleurs, ed. VII. ii. 81 = odorata.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 389.—Sumatra.
summa, Vell. Fl. Flum. viii. t. 161 = Anchietea salutaris.
sylvatica, Auct. = seq.
sylvestris, *Lam. Fl. Fr.* ii. 680.—Europ.; As. temp.
Symei, Baker, in Thirsk, Bot. Exch. Rep. (1859) 8 = tricolor.
taurica, *C. A. Mey. ex Regel, Ind. Sem. Hort. Petrop.* (1860) 44.—Tauria.
tenella, Muhl. Cat. 25; Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 78 = tricolor.
tenella, Poir. Encyc. viii. 644 = parvula?
tenella, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen; et in Am. Monthly Mag. (1819) 191 (Quid?).—Am. bor.
tenerrima, Wiesb. ex Nym. Consp. 78 = hirta.
tenuis, Benth. in Hook. Lond. Journ. Bot. i. (1842) 482 = diffusa.
tetrapetala, Molina, Sagg. Chile, ed. II. 133 = maculata.
teucriifolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 391 = rubella.
teydea, *Berth. in Mém. Soc. Linn. Par.* v. (1827) 420. t. 7.—Ins. Canar.
tezensis, *Ball, in Journ. Linn. Soc.* xvi. (1877) 350.—Marocc.
thesiifolia, Juss. ex Poir. Encyc. viii. 649 = Ionidium heterophyllum.
thessala, Boiss. & Sprun. Diagn. Ser. I. viii. 51 = alba.
thessala, Heldr. ex Nym Consp. 78 = odorata.
thianschanica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 427.—As. centr.
Thibaudieri, *Franch. & Sav. Enum. Pl. Jap.* ii. 290. —Japon.
Thomasiana, Songeon & Perrier, in Billot, Annot. Fl. Fr. et Allem. 183 = campestris.
Thomsoni, *Oudem. in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 74.—Ind. or.
Timbali, Jord. Pugill. Pl. Nov. 22 = tricolor.
tolosana, Timb. Et. Hist. Bot. Viola (1854), 6 = sepincola.
tracheliifolia, Ging. in Linnaea, i. (1826) 409 = Chamissoniana.
transsilvanica, Schur, in Oestr. Bot. Zeitschr. x. (1860) 184 = Jooi.
tricolor, *Linn. Sp. Pl.* 935.—Europ.; As. et Am. bor.
tridentata, *Sm. in Rees, Cycl.* xxxvii. n. 76.—Reg. Magellan.
trifida, Spreng. Pugill. i. 22 = hastata.
triloba, Schwein. in Am. Journ. Sc. Ser. I. v. (1822) 57 = palmata.
trinervis, Korth. in Nederl. Kruidk. Arch. i. (1848) 357 = inconspicua.
tripartita, G. Don, ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 657, nomen = seq.?
tripartita, Ell. Sketch, i. 302 = hastata.
truncata, Meyen, Reise, i. 314 = volcanica.
tuberosa, Auct. ex Steud. Nom. ed. II. ii. 773 = arborescens.
turfosa, Kirschl. ex F. Schultz, in Flora, xxiii. (1840) 121 = Schultzii.
turkestanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 12.—Turkestan.
uliginosa, *Bess. Prim. Fl. Galic.* i. 169; *Schrad. Neues Journ.* iv. (1810) 80.—Europ.
uliginosa, Muhl. Cat. 25 = canina.
umbraticola, *H. B. & K. Nov. Gen. et Sp.* v. 370.—Mexic.
umbrosa, Fries, Novit. 271 = Selkirkii.
umbrosa, Hoppe, in Flora, xi. (1828) 648 = collina.
umbrosa, Saut. in Flora, xxii. (1839) 259 = sciaphila.
uniflora, *Linn. Sp. Pl.* 936.—Sibir.

VIOLA:—
uniflora, Vand. ex Roem. Script. 147 (Quid ?).—Bras.
vaginata, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 324.—Japon.
Valderia, All. Fl. Pedem. ii. 98 = cenisia.
Valderia, Reichb. Fl. Germ. Excurs. 709 = gracilis.
variata, Jord. in Billot, Annot. Fl. Fr. et Allem. 166 = tricolor.
variegata, *Fisch. ex Link, Enum. Hort. Berol.* i. 240. —Dahur.; China.
variegata, Vukot. in Rad Jugos. Akad. Zagreb. xliv. (1878) 198 = tricolor ?
vegetalis, C. Muell. in Walp. Ann. vii. 215, sphalm. = *segetalis*.
verecunda, *A. Gray, in Mem. Acad. Am.* N. S. vi. (1858) 382.—Japon.
veronicaefolia, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 121. —N. Granat.
versicolor, Salisb. Prod. 130 = tricolor.
verticillata, *Cerv. in Anal. Cienc. Nat.* vi. (1803) 195. —Chili.
verticillata, Ortega, Hort. Matr. Dec. 50 = Ionidium polygalaefolium.
vicina, Martr. Fl. Tam, 80 = sylvestris.
Villarsiana, Roem. & Schult. Syst. v. 388 = tricolor.
villosa, Walt. Fl. Carol. 219 = cucullata.
vindobonensis ×, Wiesb. in Oestr. Bot. Zeitschr. xxx. (1880) 191.—Europ.
vinealis, Bor. Fl. Centr. Fr. ed. III. ii. 75 = odorata.
violacea, Wiesb. ex Dichtl, in Deutsche Bot. Monatsschr. iii. (1885) 45 = alba.
virescens, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 77 = alba.
virgata, Gren. in Gren. & Godr. Fl. Fr. i. 184 = tricolor.
vivariensis, Jord. Obs. Pl. Crit. i. (1846) 17 = tricolor.
volcanica, *Gill. in Hook. Bot. Misc.* iii. (1833) 145. t. 98.—Peruv.
vulgaris, Rupr. Fl. Ingr. 135 = sylvestris.
Wahlenbergii, Beurl. ex Nym. Consp. 77 = canina.
Walkerii, Wight, Illustr. Ind. Bot. i. 42 = Patrinii.
Wallichiana, Ging. in DC. Prod. i. 300 = biflora.
Wiedemanni, *Boiss. Fl. Orient.* i. 457.—Anatolia.
Wiesbauriana, Dichtl, in Deutsche Bot. Monatsschr. iii. (1885) 45 = seq.
Wiesbaurii ×, Sabransky, in Oestr. Bot. Zeitschr. xxxiv. (1884) 131. in nota.—Europ.
Wightiana, Wall. Cat. n. 4021 = serpens.
Willkommii, *De Roem. ex Willk. in Linnaea*, xxv. (1852) 10.—Hispan.
Willkommii, Teplouchow, in Bull. Soc. Oural. Amat. Sc. Nat. vii. II. 24 (Quid ?).—Rossia.
Wrightiana, *Watt, in Journ. Linn. Soc.* xviii. (1881) 379.—Reg. Himal.
yezoënsis, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 325.—Japon.
Zongia, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 300. —Madag.
Zoysii, Wulf, in Jacq. Coll. iv. 297 = calcarata.

VIOLAEOIDES, Michx. ex DC. Prod. i. 280 (1824) = **Noisettia**, H. B. & K. (Violar.).

VIONAEA, Neck. Elem. i. 107 (1790) = **Leucadendron**, Berg. (Proteac.).

VIORNA, Reichb. Handb. 277 (1837) = **Clematis**, Dill. (Ranunc.).
urnigera, Spach, Hist. Vég. Phan. vii. 270 = C. Viorna.

VIRAEA, 'Vahl,' ex Benth. & Hook. f. Gen. ii. 512 (1873), sphalm. = VIREA, Adans. = **Leontodon**, Linn. (Compos.).

VIRAYA, Gaudich. in Freyc. Voy. Bot. 466. t. 89 (1826) = **Waitzia**, Wendl. (Compos.).
Podolepis, Gaudich. l. c. = W. Podolepis.

VIRCHOWIA, Schenk, Sel. Sem. Wirceb. Adn. Crit. (1852) 1; ex Urban, in Ber. Deutsch. Bot. Ges. ii. (1884) 432. *SCROPHULARINEAE*.
cubensis, *Bartl. & Schenk, l. c.*—Hab. ?

VIRDIKA, Adans. Fam. ii. (19) (1763) = **Albuca**, Linn. (Liliac.).

VIREA, Adans. Fam. ii. 112 (1763) = **Leontodon**, Linn. (Compos.).
asplenioides, Vahl, ex Hornem. Hort. Hafn. 759 = Picris asplenioides.
autumnalis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 430 = L. autumnalis.
hastilis, Gaertn. Fruct. ii. 365. t. 159. f. 3 = L. hastilis.
hispida, S. F. Gray, Nat. Arr. Brit. Pl. ii. 429 = L. hispidus.
incana, S. F. Gray, l. c. = L. incanus.
pyrenaica, S. F. Gray, l. c. = L. pyrenaicus.
scabra, Scop. Delic. Insubr. ii. 25. t. 13 = Picris asplenioides.

VIRECTA, Afzel. ex Sm. in Rees, Cycl. xxxvii., partim (1817). *RUBIACEAE*, Benth. & Hook. f. ii. 55.
PHYTEUMOIDES, Smeathm. ex DC. Prod. iv. 414 (1830).
angustifolia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 48.—Afr. trop.
carnea, Baill. Hist. Pl. vii. 331 = Pentas carnea.
elatior, DC. Prod. iv. 415 = procumbens ?
fasciculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 136 = Ophiorrhiza fasciculata.
lanceolata, Baill. Hist. Pl. vii. 380 = Mussaenda luteola.
lutea, *G. Don, Gen. Syst.* iv. 52.—Afr. trop.
multiflora, *Sm. in Rees, Cycl.* xxxvii. n. 4.—Afr. trop.
paniculata, *G. Don, Gen. Syst.* iv. 521.—Afr. trop.
procumbens, *Sm. in Rees, Cycl.* xxxvii. n. 2.—Afr. trop.
prostrata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 136 = Ophiorrhiza Harrisiana.
setigera, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 48.—Afr. trop.
suffruticosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 136 = Ophiorrhiza fasciculata.

VIRECTA, Linn. f. Suppl. 17 (1781) = **Sipanea**, Aubl. (Rubiac.).
biflora, Linn. f. l. c. 134 = S. biflora.
dichotoma, Spreng. Syst. i. 702 = S. pratensis.
erythraeoides, Steud. Nom. ed. II. ii. 774 = S. erythraeoides.
glomerata, Spreng. Syst. i. 702 = S. glomerata.
pratensis, Vahl, Eclog. Am. ii. 11 = S. pratensis.
radicans, Steud. Nom. ed. II. ii. 774 = S. biflora.
virens, Vahl, Symb. Bot. ii. 38 = S. biflora.

VIREYA, Blume, Bijdr. 854 (1826) = **Rhododendron**, Linn. (Ericac.).
alba, Blume, l. c. 855 = R. album.
celebica, Blume, l. c. = R. malayanum.
javanica, Blume, l. c. 854 = R. javanicum.
retusa, Blume, l. c. 856 = R. retusum.
tubiflora, Blume, l. c. 855 = R. malayanum.

VIREYA, Rafin. Specch. i. 194 (1814) = **Alloplectus**, Mart. (1829) (Gesner.).
sanguinolenta, Rafin. l. c. = A. sanguineus.

VIRGA, Hill, Veg. Syst. v. 12 (1763) = **Dipsacus**, Linn.
pilosa, Hill, Hort. Kew. 75 = D. pilosus.

VIRGA-AUREA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 181 (1745) = **Solidago**, Vaill. (Compos.).

VIRGARIA, Rafin. ex DC. Prod. v. 243 (1836) = **Aster**, Tourn. (Compos.).
concolor, Rafin. ex DC. l. c. = A. concolor.

VIRGAUREA, Rupp. Fl. Jen. ed. Hall. 178 (1745) = **Erigeron**, Linn. (Compos.).

VIRGILIA, Lam. Illustr. ii. 454. t. 326 (1793). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
ANDRASTIS, Rafin. ex Benth. in Ann. Mus. Wien. ii. (1838) 86.
APHORA, Neck. Elem. iii. 50 (1790).
argentea, Poir. Encyc. viii. 679 = Ammodendrum Sieversii.
aurea, Lam. Illustr. ii. 454 = Calpurnia lasiogyne.
capensis, *Lam. l. c.*—Afr. austr.
genistoides, Poir. Encyc. viii. 679 = Cyclopia genistoides.

VIRGILIA:—
grandis, E. Mey. Comm. Pl. Afr. Austr. 1 (Quid ?).—Afr. austr.
intrusa, R. Br. in Ait. Hort. Kew. ed. II. iii. 4 = Calpurnia intrusa.
lasiogyne, D. Dietr. Syn. Pl. ii. 1501 = Calpurnia lasiogyne.
lutea, Michx. f. Hist. Arb. Am. iii. 266. t. 3 = Cladrastis tinctoria.
robinioides, DC. Prod. ii. 98 = Calpurnia robinioides.
rubiginosa, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 98 (Quid ?).—Ins. Guadaloup.
secundiflora, Cav. Ic. v. 1 = Sophora secundiflora.
sylvatica, DC. Prod. ii. 98 = Calpurnia sylvatica.

VIRGILIA, L'Hérit. Diss. [1788 ?].—Cf. Dryand. Cat. Bibl. Banks, iii. 313 = **Gaillardia**, Fouger. (Compos.).
grandiflora, [Nutt.] in Fras. Cat. (1813) n. 87 = G. bicolor.
heloides, L'Hérit. Diss. [1788 ?]; et ex DC. Prod. v. 652 = G. bicolor.

VIRGULARIA, Ruiz & Pav. Prod. 92. t. 19 (1794) = **Gerardia**, Linn. (Scrophul.).
alpestris, Mart. Nov. Gen. et Sp. iii. 10. t. 205 = G. brachyphylla.
campestris, Mart. l. c. 7. t. 203 = Esterhazya splendida.
genistifolia, Mart. l. c. 11 = G. genistifolia.
lanceolata, Ruiz & Pav. Syst. Veg. 161.—Peruv.
Mansonii, Lhotzky, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 276 = Esterhazya splendida.
montana, Mart. Nov. Gen. et Sp. iii. 9. t. 204 = Esterhazya splendida.
revoluta, Ruiz & Pav. Syst. Veg. 161.—Peruv.
sebifera, Pritz. Ic. Ind. 1167, sphalm. = *Virola sebifera*.
splendida, Mart. Nov. Gen. et Sp. iii. 11 = Esterhazya splendida.

VIRGULUS, Rafin. Fl. Tellur. ii. 46 (1836) = **Aster**, Tourn. (Compos.).
concolor, Rafin. l. c. = A. concolor.

VIRLETIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 402 (1873) = **Bahia**, Lag. (Compos.).

VIROLA, Aubl. Pl. Gui. ii. 904. t. 345 (1775) = **Myristica**, Linn.
sebifera, Aubl. l. c. = M. sebifera.

VISCAGO, Hall. Enum. Stirp. Helv. i. 373 (1742) = **Silene**, Linn. (Caryoph.).
aspera, Hornem. Hort. Hafn. 409 = S. apetala ?
baccifera, Vest, Man. Bot. 668 = Cucubalus bacciferus.
Behen, Hornem. Hort. Hafn. 409; Baumg. Enum. Stirp. Transs. i. 395 = S. inflata.
catholica, Hornem. l. c. 410 = S. catholica.
chlorantha, Moench, Meth. 704 = S. chlorantha.
clavata, Moench, l. c. = S. italica.
Fabaria, Hornem. Hort. Hafn. 409 = S. Fabaria.
fimbriata, Hornem. l. c. = S. fimbriata.
fusca, Moench, Meth. 704 = S. catholica.
gigantea, Moench, l. c. 705 = S. gigantea.
glutinosa, Baumg. Enum. Stirp. Transs. i. 395 = S. viscosa.
italica, Hornem. Hort. Hafn. 410 = S. italica.
maritima, Hornem. in Fl. Dan. t. 857 = S. maritima.
mollissima, Baumg. Enum. Stirp. Transs. i. 396 = S. italica.
mollissima, Moench, Meth. Suppl. 303 = S. mollissima.
multiflora, Hornem. Hort. Hafn. 410; Baumg. Enum. Stirp. Transs. i. 396 = S. multiflora.
nutans, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. nutans.
Otites, Hornem. Hort. Hafn. 411 = S. Otites.
paradoxa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. paradoxa.
parviflora, Hornem. Hort. Hafn. 411 = S. Otites.
pilosa, Hornem, l. c. 410 = S. italica.
polygama, Stokes, Bot. Mat. Med. ii. 533 = S. Otites.
sibirica, Hornem. Hort. Hafn. 410 = S. sibirica.
stellata, Reichb. Ic. Desc. Pl. t. 27 = S. stellata.
tartarica, Hornem. Hort. Hafn. 410 = S. tatarica.
unilateralis, Moench, Meth. 704 = S. tatarica.
viscosa, Moench, Meth. Suppl. 303 = S. viscosa.
vulgaris, Mert. & Koch, Deutschl. Fl. iii. 325 = Lychnis Viscaria.
wolgensis, Hornem. Suppl. Hort. Hafn. 48 = S. wolgensis.

VISCARIA, Riv. ex Rupp. Fl. Jen. ed. Hall. 126 (1745) = **Lychnis**, Tourn. (Caryoph.).
acutata, Walp. Ann. vii. 290, sphalm. = *oculata*.
alba, Hort. ex G. Don, in Loud. Hort. Brit. 186 = L. Viscaria.
albiflora, Sweet, Hort. Brit. ed. II. 54; ed. III. 66 = L. Viscaria.
alpina, G. Don, Gen. Syst. i. 415 = L. alpina.
aspera, Poir. Encyc. Suppl. iii. 537 = L. Coeli-rosa.
atropurpurea, Griseb. Spicil. Fl. Rumel. i. 166 = L. atropurpurea.
Burridgii, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 957 = L. Coeli-rosa.
cardinalis, Hort. ex Vilm. l. c. ed. III. 302 = L. Coeli-rosa.
Dunnetti, Hort. ex Vilm. l. c. 1220 = L. Coeli-rosa.
elegans, Hort. ex Vilm. l. c. 301 = L. Coeli-rosa.
helvetica, G. Don, Gen. Syst. i. 415 = L. alpina.
intermedia, Ledeb. Fl. Ross. i. 329.—Sibir.
Lagrangei, Coss. in Bull. Soc. Bot. Fr. xx. (1873) 243 = L. Lagrangei (in Addend.).
magellanica, G. Don, Gen. Syst. i. 415 = L. magellanica.
neglecta, G. Don, l. c. = L. Viscaria.
oculata, Lindl. Bot. Reg. (1843) t. 53 = L. Coeli-rosa.
purpurea, Wimm. Fl. Schles. (1841) 67 = L. Viscaria.
Sartorii, Boiss. Diagn. Ser. II. i. 79.—Graecia.
Sigeri, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 299 = Silene nivalis?
suecica, Sweet, Hort. Brit. ed. III. 66 = L. alpina.
viscosa, Aschers. Fl. Brandenb. i. 85 = L. viscosa.
vulgaris, Roehl. Deutschl. Fl. ed. II. ii. 275 = L. Viscaria.

VISCOIDES, Jacq. Select. Am. 73. t. 51 (1763) = **Psychotria**, Linn. (Rubiac.).
pendulum, Jacq. l. c. = P. parasitica.

VISCUM, Tourn. ex Linn. Gen. ed. I. 284 (1737). *LORANTHACEAE*, Benth. & Hook. f. iii. 213.
affine, Pohl, ex DC. Prod. iv. 281 = Phoradendron rubrum.
album, *Linn. Sp. Pl.* 1023.—Europ.; As. temp.
album, Thunb. Fl. Jap. 64 = Kaempferi.
album, Walt. Fl. Carol. 241 = Phoradendron flavescens.
album, Webb, Iter Hisp. 42 = cruciatum.
ambiguum, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 356 = Eubrachion Arnottii.
anceps, E. Mey. in Drège, Zwei Pfl. Docum. 229 = articulatum.
anceps, Spreng. Syst. i. 487 = Phoradendron hexastichon?
angulatum, *Heyne, ex DC. Prod.* iv. 283.—As. et Austral. trop.
angustifolium, Bert. ex Spreng. Syst. i. 487 = Phoradendron rubrum.
angustifolium, DC. Prod. iv. 281 = Phoradendron angustifolium.
antarcticum, Forst. f. Prod. 70 = Tupeia antarctica.
aoraiense, Nadeaud, Enum. Pl. Tahiti, 64 = articulatum.
aphyllum, Griff. Notul. iv. 634 = articulatum.
apodum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 439.—Madag.
articulatum, *Burm. f. Fl. Ind.* 311.—As. et Austral. trop.
articulatum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 133 = japonicum.
articulatum, Pohl, ex DC. Prod. iv. 284 = Phoradendron tunaeforme.
attenuatum, DC. Prod. iv. 284 = articulatum.
austriacum, Wiesb. ex Dichtl, in Deutsche Bot. Monatsschr. ii. (1884) 154 = album.
benghalense, Roxb. ex Wight & Arn. Prod. 379 = monoicum.
Berterianum, DC. Prod. iv. 281 = Phoradendron Berterianum.
Bolleanum, Seem. Bot. Voy. Herald, 295. t. 63 = Phoradendron Bolleanum.
brachystachyum, DC. Prod. iv. 280 = Phoradendron brachystachyum.
buxifolium, Lam. Encyc. iii. 56 = Dendrophthora buxifolia.
buxifolium, Spreng. Syst. i. 488 = torulosum.
capense, Bory, Voy. i. 320 = triflorum.
capense, *Linn. f. Suppl.* 426.—Afr. austr.
capillare, Blume, Bijdr. 667 (Psiloti sp.—CRYPT.).
capitellatum, *Sm. in Rees, Cycl.* xxxvii. n. 18.—Ind. or.

VISCUM:—
caucasicum, Stend. Nom. ed. I. 888 = Arceuthobium Oxycedri.
chilense, *Hook. & Arn. Bot. Beech. Voy.* 28.—Chili.
chrysostachyum, Presl, Epim. Bot. 254 = Phoradendron chrysostachyum.
clavatum, Benth. Pl. Hartw. 189 = Phoradendron clavatum.
compressum, Poir. Encyc. Suppl. ii. 861 = articulatum.
confertum, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 764 = monoicum.
continuum, *E. Mey. in Drège, Zwei Pfl. Docum.* 229, nomen.—Afr. austr.
cordifolium, *Presl, Epim. Bot.* 253.—Panama.
cornifolium, A. Cunn. ex Oliver, in Journ. Linn. Soc. vii. (1864) 103 = Notothixos cornifolius.
cornifolium, Presl, Epim. Bot. 254 = Phoradendron Schottii?
crassifolium, Pohl, ex DC. Prod. iv. 280 = Phoradendron crassifolium.
Crassulae, *Eckl. & Zeyh. Enum.* 357.—Afr. austr.
crassuloides, Presl, Epim. Bot. 252 = Phoradendron crassuloides.
cruciatum, *Sieber, ex Spreng. Syst.* i. 488, in syn.; *Boiss. Voy. Bot. Espagne*, ii. 274.—Hispan.
cryptophlebium, *Baker, in Journ. Bot.* xx. (1882) 245.—Madag.
Cumingianum, Presl, Epim. Bot. 255 = Ginalloa Cumingiana.
cuneifolium, *Baker, in Journ. Linn. Soc.* xxi. (1885) 438.—Madag.
cupressoides, Macfad. Fl. Jamaic. ii. 197 = Dendrophthora cupressoides.
cupulatum, DC. Prod. iv. 285 = Dendrophthora cupulata.
dichotomum, Bert. ex Spreng. Syst. i. 488 = Phoradendron Berterianum.
dichotomum, D. Don, Prod. Fl. Nep. 147 = articulatum.
dichotomum, Gilib. Exercit. ii. 394 = album.
dimidiatum, *Miq. in Linnaea*, xviii. (1844) 58.—Guian.
dimidiatum, Sagot, ex Eichl. in Mart. Fl. Bras. v. II. 126 = Phoradendron Schottii.
distichum, *Endl. Prod. Fl. Norfolk.* 61.—Ins. Norfolk.
domingense, Spreng. Syst. i. 487 = Dendrophthora domingensis.
echinocarpum, *Baker, in Journ. Linn. Soc.* xx. (1883) 248.—Madag.
ellipticum, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 106 = Phoradendron ellipticum.
elongatum, Wall. Cat. n. 495; DC. Prod. iv. 284 = articulatum.
ensifolium, Pohl, ex DC. l. c. 281 = Phoradendron ensifolium.
Eucleae, *Eckl. & Zeyh. Enum.* 357.—Afr. austr.
Euphorbiae, E. Mey. in Drège, Zwei Pfl. Docum. 229 = V. Crassulae.
falcatum, Cham. & Schlecht. in Linnaea, v. (1830) 172 = Phoradendron angustifolium.
falcatum, Hook. Ic. Pl. t. 368 = Phoradendron falcatum.
falcatum, Wall. Cat. n. 492; DC. Prod. iv. 278 = monoicum.
falcifrons, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 356.—Uraguay.
flagelliforme, Descourt. Fl. Antill. v. 204 = Dendrophthora gracilis.
flavens, Sw. Prod. Veg. Ind. Occ. 32 = Phoradendron chrysocladon.
flavescens, Comm. ex DC. Prod. iv. 286 = Myzodendron punctulatum.
flavescens, Pursh, Fl. Am. Sept. i. 114 = Phoradendron flavescens.
floccosum, Thw. Enum. Pl. Zeyl. 418 = Notothixos floccosus.
florianum, *Andersson. in Vet. Akad. Handl. Stockh.* 1853 (1855) 219.—Ins. Galapag.
Fockeanum, Miq. in Linnaea, xviii. (1844) 60 = Phoradendron Schottii.
Fockeanum, Sagot, ex Eichl. in Mart. Fl. Bras. v. II. 89, 90 (= *Oryctanthus botryostachys*) = Loranthus ruficaulis.
fragile, Wall. Cat. n. 498; DC. Prod. iv. 284 = articulatum.
galapageium, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 216.—Ins. Galapag.

VISCUM:—
geminatum, *Korth. in Verh. Batav. Gen.* xvii. (1839) 259.—Borneo.
glandulosum, *Miq. in Linnaea*, xviii. (1844) 60.—Guian.
glaucum, Eckl. & Zeyh. in South. Afr. Journ. (1830) 375; ex eor. Enum. 357 = rotundifolium.
globuliflorum, *Presl, Epim. Bot.* 251.—Peruv.
gracile, DC. Prod. iv. 285 = Dendrophthora gracilis.
granulosum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 438.—Madag.
grossum, *Wight, in Wall. Cat.* n. 6879.—Ind. or.
guyanense, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1161 (= *Phthirusa guyanensis*).—Guian.
Helferi, Presl, Epim. Bot. 256 = Ginalloa Helferi.
Henslovii, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 216.—Ins. Galapag.
heteranthum, Wall. Cat. n. 488; DC. Prod. iv. 279 = Henslowia heterantha.
hexastichum, DC. l. c. 282 = Phoradendron hexastichon.
Heyneanum, DC. l. c. 278 = orientale.
incanum, Hook. Ic. Pl. t. 73 = Notothixos incanus.
interruptum, DC. Prod. iv. 282 = Phoradendron interruptum.
jamaicense, Macfad. Fl. Jamaic. ii. 195 = Phoradendron trinervium.
japonicum, *Thunb. in Trans. Linn. Soc.* ii. (1794) 329.—As.; Afr.; Austral. trop. et subtrop.
Kaempferi, *DC. Prod.* iv. 285.—Japon.
Kunthianum, DC. l. c. 283 = Phoradendron rubrum.
lanceolato-ellipticum, Pohl, ex DC. l. c. 282 = Phoradendron interruptum.
latifolium, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 142 = Henslowia heterantha.
latifolium, Lam. Encyc. iii. 57 = Phoradendron hexastichon.
latifolium, Sw. Fl. Ind. Occ. i. 268 = Phoradendron Schottii.
laurifolium, Presl, Epim. Bot. 255 = Phoradendron laurifolium.
laxum, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 16.—Hispan.
leptostachyum, DC. Prod. iv. 280 = Dendrophthora leptostachya.
leucarpum, Rafin. Fl. Ludov. 79 = Phoradendron flavescens.
leucocarpum, DC. Prod. iv. 280, in syn. = praec.
Liga, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 355.—Chili.
ligulatum, Blume, Bijdr. 667 (Psiloti sp.—CRYPT.).
ligustrinum, *Presl, Epim. Bot.* 252.—Mexic.
Lindsayi, *Oliver, ex Hook. f. Handb. N. Zeal. Fl.* 108.—N. Zel.
linifolium, *Presl, Epim. Bot.* 251.—Mexic.
lophiocladum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 437.—Madag.
macrophyllum, Macfad. Fl. Jamaic. ii. 195 = Phoradendron chrysocladon.
macrophyllum, Spreng. Syst. i. 488 = Phoradendron hexastichon?
macrostachyon, Jacq. Coll. ii. 109. t. 5. f. 3 = Dendrophthora macrostachya.
macrostachyon, Spreng. Syst. i. 487 = Dendrophthora leptostachya.
Mangiferae, Wight, in Wall. Cat. n. 6878 = capitellatum.
martinicense, DC. Prod. iv. 280 = Phoradendron hexastichon?
microphyllum, Pohl, ex DC. l. c. 283 = Phoradendron rubrum.
microstachyum, Presl, Epim. Bot. 252 = Phoradendron rubrum?
minimum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 581.—Afr. austr.
moniliforme, Blume, Bijdr. 667 = articulatum.
moniliforme, Wight & Arn. Prod. 380 = japonicum.
monoicum, *Roxb. Hort. Beng.* [105]; *Fl. Ind.* iii. 763.—Ind. or.; Burma.
monoicum, Wight, ex Wall. Cat. n. 6875 = verruculosum.
monstrosum, Bert. ex DC. Prod. iv. 284 = Dendrophthora opuntioides.
montanum, Zoll. & Mor. Syst. Verz. Zoll. 39 = orientale.
mucronatum, DC. Prod. iv. 282 = Phoradendron rubrum.
multicostatum, *Baker, in Journ. Linn. Soc.* xx. (1883) 248.—Madag.

VISCUM :—
myriophlebium, *Baker, in Journ. Linn. Soc.* xx. (1883) 248.—Madag.
myrtilloides, Willd. Sp. Pl. i. 739 = Phoradendron rubrum.
navicellatum, Korth. in Verh. Batav. Gen. xvii. (1839) 256 = orientale.
nepalense, Spreng. Syst. iv. Cur. Post. 47 = articulatum.
nervosum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 388 = triflorum.
nitidum, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 105 = Phoradendron nitidum.
obliquum, *Presl, Epim. Bot.* 253.—Mexic.
oblongifolium, *DC. Prod.* iv. 283.—Ins. Guadaloup.
oblongifolium, Rafin. New Fl. Am. iii. 23 = Phoradendron rubrum?
obovatum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 579. —Afr. austr.
obscurum, *Thunb. Prod. Pl. Cap.* 31 (sp. dub.).—Afr. austr.
obtusatum, Wall. Cat. n. 494; DC. Prod. iv. 279 = ovalifolium.
obtusissimum, *Miq. in Linnaea*, xviii. (1844) 602.—Bras.
ochroleucum, Rafin. in New Fl. Am. iii. 23 = Phoradendron flavescens.
oliganthum, Presl, Bot. Bemerk. 75 = obscurum?
Opuntia, Thunb. Fl. Jap. 64 = japonicum.
opuntioides, Heyne, ex DC. Prod. iv. 284; Roxb. Fl. Ind. iii. 764 = articulatum.
opuntioides, Linn. Sp. Pl. 1023 = Dendropthora opuntioides.
orbiculatum, *Wight, Ic.* t. 1016.—Ind. or.
orientale, DC. Prod. iv. 278 = cruciatum.
orientale, *Willd. Sp. Pl.* iv. 737.—As. et Austral. trop.
ovalifolium, *Wall. Cat.* n. 489; *DC. Prod.* iv. 278.—Burma; China; Malaya.
Oxycedri, DC. Fl. Fr. iv. 274 = Arceuthobium Oxycedri.
Pamattonis, Korth. in Verh. Batav. Gen. xvii. (1839) 255 = orientale.
pauciflorum, *Linn. f. Suppl.* 246 (sp. dub.).—Afr. austr.
pauciflorum, E. Mey. in Drège, Zwei Pfl. Docum. 229 = obscurum.
penninervium, Steud. Nom. ed. II. ii. 775, sphalm. = seq.
pennivenium, DC. Prod. iv. 282 = Phoradendron pennivenium.
pentanthum, *Baker, in Journ. Linn. Soc.* xx. (1883) 249.—Madag.
Perrottetii, DC. Prod. iv. 280 = Phoradendron Perrottetii.
philippense, Llanos, Fragm. 52 = articulatum.
piperoides, DC. Prod. iv. 281 = Phoradendron Schottii.
platycaulon, Bert. ex A. Gray, Bot. U. St. Expl. Exped. i. 744 = articulatum.
platyphyllum, Spreng. Syst. iv. Cur. Post. 47 = Henslowia heterantha.
portulacoides, *Presl, Epim. Bot.* 252.—Ecuador.
pubigerum, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 208; R. Cunn. ex Miq. in Linnaea, xviii. (1844) 86 = Tupeia antarctica.
purpureum, Linn. Sp. Pl. 1023 = Loranthus uniflorus.
quadrangulare, DC. Prod. iv. 283 = Phoradendron rubrum?
quernum, Auct. ex Reichb. Fl. Germ. Excurs. 203 = Loranthus europaeus.
racemosum, Aubl. Pl. Gui. ii. 895 = Phoradendron hexastichon.
Radula, *Baker, in Journ. Linn. Soc.* xxi. (1885) 439. —Madag.
ramosissimum, *Wall. Cat.* n. 6876.—Ind. or.
ramosissimum, Wight, Ic. t. 1017 = angulatum.
Randiae, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 278.—Porto Rico.
Reichenbachianum, Seem. Bot. Voy. Herald, 294. t. 62 = Phoradendron Reichenbachianum.
rhytidocarpum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 438.—Madag.
robustum, *Eckl. & Zeyh. Enum.* 358.—Afr. austr.
rotundifolium, Bory, Voy. i. 320 = triflorum.
rotundifolium, Eckl. & Zeyh. Enum. 357 = obscurum.
rotundifolium, *Linn. f. Suppl.* 426.—Afr. austr.
Roxburghianum, Korth. in Verh. Batav. Gen. xvii. (1839) 257 = orientale.
rubrum, Burm. f. Fl. Ind. 311 (Hederae sp.).—Ind. or.
rubrum, Griseb. ex Eichl. in Mart. Fl. Bras. v. II. 115 = Phoradendron corynarthron.

VISCUM :—
rubrum, Linn. Sp. Pl. 1023 = Phoradendron rubrum.
rugosum, Rafin. New Fl. Am. iii. 24 = Loranthus uniflorus?
salicifolium, Presl, Epim. Bot. 254 = Phoradendron salicifolium.
salicornoides, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 208 = articulatum.
sauruoides, DC. Prod. iv. 280 (excl. syn.) = Phoradendron Schottii.
Schiedeanum, DC. l. c. 670 = Phoradendron angustifolium.
Schottii, Pohl, ex DC. l. c. 281 = Phoradendron Schottii.
serotinum, Rafin. New Fl. Am. iii. 22 (Quid?).—Am. bor.
Sertularia, Presl, Bot. Bemerk. 75 = articulatum.
spathulifolium, Thw. Enum. Pl. Zeyl. 136 = Ginalloa spathulifolia.
squamigerum, Benth. Pl. Hartw. 190 = Phoradendron squamigerum.
stellatum, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 142 = album.
stenophyllum, Spreng. Syst. i. 487 = Phoradendron angustifolium.
subaureum, F. Muell. ex Oliver, in Journ. Linn. Soc. vii. (1864) 103 = Notothixos subaureus.
surinamense, *G. W. Schilling, Desc. Pl. Leprae*, 201. t. 3.—Guian.
taenioides, Comm. ex Thou. Obs. Pl. Afr. in Mélang. 43 = japonicum.
tereticaule, DC. Prod. iv. 280 = Phoradendron Schottii.
teretifolium, Hemsl. Biol. Centr. Am. Bot. iii. 85, sphalm. = praec.
terrestre, Linn. Sp. Pl. 1023 = Lysimachia stricta.
tetragonum, DC. Prod. iv. 282 = Phoradendron rubrum.
thymifolium, *Presl, Epim. Bot.* 251.—Afr. austr.
tomentosum, DC. Prod. iv. 670 = Phoradendron tomentosum.
toruloides, Eichl. in Mart. Fl. Bras. v. II. 107, in nota = torulosum.
torulosum, *DC. Prod.* iv. 279.—Ecuador.
trachycarpum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 439.—Madag.
tricostatum, *E. Mey. in Drège, Zwei Pfl. Docum.* 229, nomen; *et ex Harv. in Harv. & Sond. Fl. Cap.* ii. 580.—Afr. austr.
triflorum, *DC. Prod.* iv. 279.—Ins. Maurit.
trigonum, D. Dietr. Syn. Pl. i. 546 = Phoradendron rubrum.
trinervium, Lam. Encyc. iii. 57 = Phoradendron trinervum.
trinervium, Spreng. Syst. i. 488 = Phoradendron rubrum.
tuberculatum, *A. Rich. Tent. Fl. Abyss.* i. 338.—Abyss.
tunaefolium, Steud. Nom. ed. II. ii. 775, sphalm., in syn. = seq.
tunaeforme, DC. Prod. iv. 284 = Phoradendron tunaeforme.
umbellatum, *Blume, Bijdr.* 666.—Java.
undulatum, Pohl, ex DC. Prod. iv. 282 = Phoradendron undulatum.
vaginatum, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 740 = Arceuthobium vaginatum.
vaginatum, Spreng. Syst. i. 487 = Dendrophthora cupulata.
velutinum, DC. Prod. iv. 281 = Phoradendron velutinum.
venosum, DC. l. c. 279 = triflorum.
verrucosulum, Steud. Nom. ed. II. ii. 776, sphalm. = verruculosum.
verrucosum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 581. —Afr. austr.
verruculosum, *Wight & Arn. Prod.* 379.—Ind. or.
verticillatum, Lam. Encyc. iii. 57 (Rhipsalidis sp.?).—Jamaic.
verticillatum, Linn. Sp. Pl. 1023 = Phoradendron trinervium.
verticillatum, Nutt. Gen. Am. ii. 235, partim = Phoradendron flavescens.
verticillatum, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 764 = orientale.
verticilliflorum, Royle, Illustr. Bot. Himal. 235 = orientale?
villosum, Nutt. in Torr. & Gray, Fl. N. Am. i. 654 = Phoradendron flavescens.
Wallichianum, *Wight & Arn. Prod.* 379.—Ind. or.
Wightianum, Wall. Cat. n. 6877 = japonicum.
Wightianum, *Wight & Arn. Prod.* 380.—Ind. or.

VISCUM :—
Willdenowianum, *Molkenb. in Miq. Pl. Jungh.* 108.—Palaestin.
xanthophyllum, A. Cunn. ex Oliver, in Journ. Linn. Soc. vii. (1864) 103 = Notothixos cornifolius.

VISENA, Schult. Syst. vi. 736 (1820), sphalm. = seq.

VISENIA, Houtt. Handleid. viii. 308 (1777) = **Melochia**, Dill. (Stercul.).
alba, Endl. ex Walp. Rep. i. 351.—Java.
Berteriana, Spreng. Syst. iii. 30 = Waltheria glabra.
caracasana, Spreng. l. c. = M. caracasana.
concatenata, Spreng. l. c. 31 = M. corchorifolia.
corchorifolia, Spreng. l. c. 30 = M. corchorifolia.
corymbosa, Spreng. l. c. 29 = M. corymbosa (in Addend.).
crenata, Spreng. l. c. 30 = M. tomentosa.
depressa, Spreng. l. c. = M. depressa.
graminifolia, Spreng. l. c. iv. Cur. Post. 248 = M. graminifolia.
hirsuta, Spreng. l. c. iii. 30 = M. hirsuta, *Cav.*
indica, J. F. Gmel. Syst. 515 = M. odorata.
inflata, Spreng. Syst. iii. 30 = M. lupulina.
javanica, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 302 = M. odorata.
lupulina, Spreng. Syst. iii. 31 = M. lupulina.
madagascarensis, Spreng. l. c. 30.—Madag.
mollis, Spreng. l. c. = M. mollis.
nodiflora, Spreng. l. c. = M. nodiflora.
polystachya, Spreng. l. c. = M. venosa.
sericea, Spreng. l. c. iv. Cur. Post. 248 = M. venosa.
serrata, Spreng. l. c. iii. 30 = M. hirsuta, *Cav.*
supina, Spreng. l. c. 31 = M. corchorifolia.
tomentosa, Miq. Anal. Bot. Ind. iii. 4 = M. arborea.
Ulmariae, Spreng. Syst. iv. Cur. Post. 248 = M. ulmarioides.
umbellata, Houtt. Handleid. viii. 309. t. 46. f. 3 = M. odorata.
umbellata, Wight, Ic. 509 = M. arborea.
velutina, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 212 (= *Riedlea velutina*).—Ind. or.
venosa, Spreng. Syst. iii. 30 = M. venosa.

VISIANIA, DC. Prod. viii. 289 (1844) = **Ligustrum**, Tourn. (Oleac.).
glomerata, Miq. Fl. Ind. Bat. ii. 549 = L. glomeratum.
grandiflora, DC. Prod. viii. 289 = L. nepalense.
paniculata, DC. l. c. = L. glomeratum.
phyllothyrsa, Miq. Fl. Ind. Bat. Suppl. 558.—Sumatra.
pubinervis, Miq. Fl. Ind. Bat. ii. 548 = L. robustum.
robusta, DC. Prod. viii. 289 = L. robustum.
robusta, Wight, Ic. t. 1242 = L. Roxburghii.
sumatrana, Miq. Fl. Ind. Bat. ii. 549.—Sumatra.
undulata, Miq. l. c. 548 = L. undulatum.

VISIANIA, Gasp. Nov. Gen. Fici, 9 (1844) = **Ficus**, Tourn. (Urtic.).
elastica, Gasp. l. c. = F. elastica.

VISINIA, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 382, sphalm. = seq.

VISMIA, Vand. Fl. Lusit. 51. t. 3. f. 24 (1788).
HYPERICINEAE, Benth. & Hook. f. i. 166.
ACROSSANTHES, Presl, Bot. Bemerk. 22 (1844).
CASPIA, Pison. ex Scop. Introd. 276 (1777).
acuminata, Kunth, ex Spreng. Syst. iii. 350 = glabra.
acuminata, *Pers. Syn.* ii. 86.—Guian.
affinis, *Oliver, Fl. Trop. Afr.* i. 161.—Afr. trop.
alternifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 579 (Psorospermi sp.?).—Sumatra.
angusta, Miq. in Linnaea, xviii. (1844) 27 = ferruginea.
arborescens, Choisy, Prod. Hyp. 36 = Cratoxylon Blancoi.
baccifera, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 300 = dealbata.
baccifera, *H. G. Reich. in Mart. Fl. Bras.* xii. I. 204.—Bras.
Billbergiana, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 117.—Panama.
brasiliensis, *Choisy, Prod. Hyp.* 35. t. 2.—Bras.
Caparosa, H. B. & K. Nov. Gen. et Sp. v. 182 = acuminata.
cayennensis, *Pers. Syn.* ii. 86.—Am. trop.
cochinchinensis, Spreng. Syst. iii. 350 = Cratoxylon polyanthum.
confertiflora, *Spruce, ex H. G. Reich. in Mart. Fl. Bras.* xii. I. 205.—Bras.

VISMIA :—
cuspidata, Steud. Nom. ed. II. i. 787, in syn. = ferruginea.
dealbata, *H. B. & K. Nov. Gen. et Sp.* v. 184. t. 454. —N. Granat.
decipiens, *Cham. & Schlecht. in Linnaea*, iii. (1828) 116.—Bras.
ferruginea, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 117 = Billbergiana.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* v. 183.—Am. trop.
frondosa, *Oliver, Fl. Trop. Afr.* i. 161.—Afr. trop.
glabra, *Ruiz & Pav. Syst. Veg.* 183.—Am. trop.
guianensis, *DC. Prod.* i. 542.—Am. trop.
guianensis, Seem. Bot. Voy. Herald, 88 = viridiflora.
guineensis, *Choisy, Prod. Hyp.* 36.—Afr. trop.
guttifera, Pers. Syn. ii. 86; Salzm. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 382 = baccifera.
Hilairii, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 334 = magnoliaefolia.
japurensis, *H. G. Reich. in Mart. Fl. Bras.* xii. i. 209. —Bras.
laccifera, Mart. Reise Bras. i. 552 = brasiliensis.
laevis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 303.—N. Granat.
lanceolata, Pers. Syn. ii. 86 = glabra.
lasiantha, Klotzsch, ex H. G. Reich. in Mart. Fl. Bras. xii. i. 198 = brasiliensis.
latifolia, *Choisy, Prod. Hyp.* 36.—Guian.
latifolia, H. B. & K. Nov. Gen. et Sp. v. 183 = ferruginea *vel* dealbata.
laurifolia, *Pers. Syn.* ii. 86.—Bras.
lauriformis, *Choisy, Prod. Hyp.* 35.—N. Granat.
laxiflora, *H. G. Reich. in Mart. Fl. Bras.* xii. i. 203.—Guian.
leonensis, *Hook. f. in Hook. Niger Fl.* 243.—Afr. trop.
Lindeniana, *Decne. ex Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 381.—N. Granat.
longifolia, A. St. Hil. Fl. Bras. Mer. i. 326. t. 68 = brasiliensis.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 184.—Venezuela.
macrophylla, Salzm. ex Griseb. Fl. Brit. W. Ind. 111 = ferruginea.
magnoliaefolia, *Cham. & Schlecht. in Linnaea*, iii. (1828) 118.—Bras.
Martiana, *H. G. Reich. in Mart. Fl. Bras.* xii. i. 204.—Bras.
mexicana, *Schlecht. in Linnaea*, x. (1828) 245.—Mexic.
micrantha, Mart. Reise Bras. i. 552 = parviflora.
micrantha, *A. St. Hil. Fl. Bras. Mer.* i. 327.—Bras.
obtusa, *Spruce, ex H. G. Reich. in Mart. Fl. Bras.* xii. i. 207.—Bras.
panamensis, Duchass. & Walp. in Linnaea, xxiii. (1850) 748 = ferruginea.
parviflora, *Cham. & Schlecht. l. c.* iii. (1828) 119.—Bras.
petiolata, Choisy, Prod. Hyp. 36 = laurifolia.
ramuliflora, Miq. Stirp. Surin. Sel. 88 = cayennensis.
reticulata, Choisy, Prod. Hyp. 34 = latifolia.
rubescens, *Oliver, Fl. Afr. trop.* i. 160.—Afr. trop.
rufescens, *Pers. Syn.* ii. 86.—Guian.
Sellowiana, Klotzsch, ex H. G. Reich. in Mart. Fl. Bras. xii. i. 198 = brasiliensis.
sessilifolia, Choisy, Prod. Hyp. 34 = rufescens.
sessilifolia, *DC. Prod.* i. 542.—Guian.
Sieberiana, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 999, nomen.—Guian.
tomentosa, *Ruiz & Pav. Syst. Veg.* 183.—Peruv.
viridiflora, *Duchass. ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 302.—N. Granat.

VISNAGA, Gaertn. Fruct. i 92. t. 21 (1788) = **Ammi**, Tourn. (Umbell.).
daucoides, Gaertn. l. c. = A. Visnaga.
meoides, Rafin. New Fl. Am. iv. 28 = A. anethifolium?
vera, Rafin. l. c. = A. Visnaga.

VISNEA, Linn. f. Suppl. 36 (1781). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 182.
MOCANERA, Juss. Gen. 318 (1789).
Mocanera, *Linn. f. Suppl.* 251.—Ins. Canar.

VISNEA, Steud. ex Endl. Gen. 173 (1836) = **Barbacenia**, Vand. (Amaryll.).

VISSADALI, Adans. Fam. ii. 145 (1763) = **Knoxia**, Linn. (Rubiac.).

VISTNU, Adans. Fam. ii. 245 (1763) = **Evolvulus**, Linn. (Convolv).

VITELLARIA, Gaertn. f. Fruct. iii. 131. t. 205 (1805) = **Lucuma**, Juss. (Sapot.).
Casaretti, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 326 = L. Casaretti.
curvifolia, Radlk. l. c. = L. curvifolia.
glabrescens, Radlk. l. c. = L. glabrescens.
grandiflora, Radlk. l. c. = L. grandiflora.
littoralis, Radlk. l. c. = L. littoralis.
mammosa, Radlk. l. c. 321, 323, 325 = L. mammosa.
marginata, Radlk. l. c. 326 = L. marginata.
multiflora, Radlk. l. c. 320, 326 = L. multiflora.
nervosa, Radlk. l. c. 326 = L. Rivicoa.
paradoxa, Gaertn. f. Fruct. iii. 131. t. 205 = L. paradoxa.
pauciflora, Radlk. in Sitzb. Math.-Phys. Akad. Muench. xii. (1882) 320, 326 = L. pauciflora.
procera, Radlk. l. c. 325, 326 = L. procera.
pulverulenta, Radlk. l. c. 326 = L. pulverulenta.
retusa, Radlk. l. c. = L. retusa.
Rivicoa, Radlk. l. c. 323, 326 = L. Rivicoa.
serpentaria, Radlk. l. c. 320, 326 = L. serpentaria.
sphaerocarpa, Radlk. l. c. 326 = L. sphaerocarpa.
Valenzuelana, Radlk. l. c. 320, 326 = L. Valenzuelana.
venosa, Radlk. l. c. 326 = L. venosa.

VITENIA, Noronha, ex Cambess. in Mém. Mus. Par. xviii. (1829) 10.—Cf. UITENIA, Noronha = **Erioglossum**, Blume (Sapind.).
edulis, Steud. Nom. ed. II. ii. 776 = E. edule.
stilaginea, Noronha, ex Cambess. in Mém. Mus. Par. xviii. (1829) 10.—Hab.?

VITEX, Tourn. ex Linn. Syst. ed. I (1735). *VERBENACEAE*, Benth. & Hook. f. ii. 1154.
CASARETTOA, Walp. Rep. iv. 91 (1844).
CHRYSOMALLUM, Thou. Gen. Nov. Madag. 8 (1806).
EPHIALIS, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1838) 461.
LIMIA, Vand. Fl. Lusit. 42. t. 3. f. 21 (1788).
NEPHRANDRA, Willd. in Cothen. Disp. Veg. 8 (1790).
PSILOGYNE, DC. in Bibl. Univ. Genèv. xvii. (1838) 132 (Rev. Bign. 16).
PYROSTOMA, G. F. W. Mey. Prim. Fl. Esseq. 219 (1818).
WALLROTHIA, Roth, Nov. Pl. Sp. 317 (1821).
acuminata, *R. Br. Prod.* 512.—Austral.
Agnus, Stokes, Bot. Mat. Med. ii. 413 = V. Agnus-castus.
Agnus-castus, Kurz, For. Fl. Brit. Burma, ii. 269, partim = trifolia.
Agnus-castus, *Linn. Sp. Pl.* 638.—Reg. Mediterr.; Oriens.
alata, Roxb. Hort. Beng. 46; Fl. Ind. iii. 72 = pedunculariis.
alata, Schau. in DC. Prod. xi. 685 = limonifolia.
alata, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 203; *Heyne, ex Roth, Nov. Pl. Sp.* 316.—Ind. or.
altissima, *Linn. f. Suppl.* 294.—Ind. or.
altissima, Naves, ex F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 160 = littoralis, *Decne.*
appendiculata, Rottl. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 584 = alata.
arborea, Desf. Cat. Hort. Par. ed. III. 391 = V. Negundo.
arborea, Roxb. Hort. Beng. 46; Fl. Ind. iii. 73 = pubescens.
articulata, Steud. Nom. ed. II. ii. 777 = pubescens.
avicennioides, *A. Rich. Fl. Cub. Fanerog.* ii. 149.—Cuba.
axillaris, *Wall. Cat.* n. 1760.—Burma.
Babula, Buch.-Ham. ex Wall. l. c. n. 1745 = heterophylla.
bahiensis, *Schau. in DC. Prod.* xi. 687.—Bras.
Balbisii, *Spreng. Syst.* ii. 756.—Ins. S. Doming.
beraviensis, *Vatke, in Linnaea*, xliii. (1880–82) 535.—Madag.
bicolor, Willd. Enum. Hort. Berol. 660 = V. Negundo.
bignonioides, *H. B. & K. Nov. Gen. et Sp.* ii. 246.—Venezuela.
Bojeri, *Schau. in DC. Prod.* xi. 694.—Madag.
bombacifolia, Wall. Cat. n. 1749 = glabrata.
bracteata, Herb. Horsf. ex Miq. Fl. Ind. Bat. ii. 862 = pubescens.
brasiliensis, *Juss. ex Steud. Nom.* ed. I. 888.—Bras.
brasiliensis, Mart. ex Schau. in DC. Prod. xi. 690 = Sellowiana.

VITEX:—
californica, Benth. Bot. Voy. Sulph. 10, sphalm. = Vitis californica.
canescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 101.—Burma.
canescens, *Wall. Cat.* n. 1757 (= an praec.?).—Burma.
cannabifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 152 (*Fl. Jap. Fam. Nat.* ii. 28).—Japon.
capitata, *Vahl, Eclog. Am.* ii. 50. t. 18.—Am. trop.
caribaea, Benth. Bot. Voy. Sulph. 10, sphalm. = Vitis californica.
chinensis, Mill. Gard. Dict. ed. VIII. n. 5 = V. Negundo.
chrysocarpa, *Planch. in Hook. Niger Fl.* 486.—Afr. trop.
chrysoclada, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 268.—Afr. trop.
Chrysomallum, *Steud. Nom.* ed. II. ii. 777.—Madag.
Cienkowskii, *Kotschy & Peyr. Pl. Tinn.* 27. n. 42. t. 12.—Afr. trop.
Cofassus, *Reinw. ex Blume, Bijdr.* 813.—Ins. Molucc.
compressa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 224.—N. Granat.
coriacea, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 586.—Malacca.
cuneata, *Schum. & Thonn. Beskr. Guin. Pl.* 289.—Afr. trop.
Cunninghami, Schau. in DC. Prod. xi. 691 = glabrata.
cymosa, *Bert. ex Spreng. Syst.* ii. 757.—Bras.
Dalrympleana, F. Muell. Fragm. iv. 128 = Gmelina macrophylla.
dentata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 265. —Afr. trop.
digitata, Wight, ex Steud. Nom. ed. II. ii. 777 = pubescens.
divaricata, *Sw. Prod. Veg. Ind. Occ.* 93.—Ind. occ.; Guian.
diversifolia, *Kurz, Andam. Rep. App.* A 45; B 14.—Ins. Andaman.
Doniana, *Sweet, Hort. Brit.* ed. I. 323.—Afr. trop.
elegans, *Griff. Notul.* iv. 740.—Burma (cult.).
ferruginea, Boj. ex Schau. in DC. Prod. xi. 694 = Bojeri.
ferruginea, *Schum. & Thonn. Beskr. Guin. Pl.* 288.—Afr. trop.
Finlaysoniana, Wall. Cat. n. 6314 = vestita.
flavens, *H. B. & K. Nov. Gen. et Sp.* ii. 246.—Peruv.
floridula, *Duchass. & Walp. in Walp. Ann.* iii. 240.—Panama.
gamosepala, *Griff. Notul.* iv. 178.—Malacca.
Gardneriana, *Schau. in DC. Prod.* xi. 687.—Bras.
geniculata, *Blanco, Fl. Filip.* ed. I. 514.—Ins. Philipp.
gigantea, *H. B. & K. Nov. Gen. et Sp.* ii. 245.—Ecuador.
glaberrima, Zipp. ex Span. in Linnaea, xv. (1841) 330 = littoralis, *Decne.*
glabrata, *R. Br. Prod.* 512.—As. et Austral. trop.
glabrata, F. Muell. Fragm. v. 35 = littoralis, *A. Cunn.*
gracilis, Salisb. Prod. 107 = V. Negundo.
grandiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 223.—N. Granat.
Haynga, *Roxb. Hort. Beng.* 46, nomen.—Ind. or.
heptaphylla, *A. Juss. in Ann. Mus. Par.* vii. (1806) 77.—Ins. Dominic.
heterophylla, Blume, ex Miq. Fl. Ind. Bat. ii. 862 = pubescens.
heterophylla, *Roxb. Hort. Beng.* 46; *Fl. Ind.* iii. 75. —Assam; Burma; Malaya.
heterophylla, Schau. in DC. Prod. xi. 686, partim = urceolata.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 534. —Afr. trop.
hypoleuca, *Schau. in DC. Prod.* xi. 690.—Bras.
ibarensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 226. —Madag.
ilicifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 148.—Cuba.
inaequifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 223 = pubescens.
incisa, Lam. Encyc. ii. 612 = V. Negundo.
incisa, Wall. Cat. n. 1746, partim = trifolia.
indica, Mill. Gard. Dict. ed. VIII, in errat. = trifolia.
integerrima, Mill. l. c. n. 3 = trifolia.
integra, Medic. Beobacht. (1782) 313 = V. Agnus-castus.
involucratus, Presl, Bot. Bemerk. 148 = Sphenodesme unguiculata.
laciniata, Hort. ex Schau. in DC. Prod. xi. 684 = V. Negundo.

VITEX:—
laciniosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 225.—Bras.
laevigata, Herb. Madr. ex Wight & Arn. Prod. 433 = Psychotria connata.
lanceolata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 224.—Java.
lanigera, *Schau. in DC. Prod.* xi. 695.—Madag.
lasiophylla, Benth. Bot. Voy. Sulph. 155 = mollis.
latifolia, Lam. Encyc. ii. 613 = pubescens.
latifolia, Mill. Gard. Dict. ed. VIII. n. 2 = V. Agnus-castus.
latifolia, Wight, ex Steud. Nom. ed. II. ii. 777 = altissima.
Leichhardtii, F. Muell. Fragm. iii. 58 = Gmelina Leichhardtii.
lepidota, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 225.—Bras.
Leucoxylon, Blanco, Fl. Filip. ed. I. 516 = V. Negundo.
Leucoxylon, *Linn. f. Suppl.* 293.—Ind. or.
Leucoxylon, Schau. in DC. Prod. xi. 692, partim = glabrata.
Leucoxylon, Span. ex Miq. Fl. Ind. Bat. ii. 863 = littoralis, *Decne.*
Lignum-vitae, *A. Cunn. ex Schau. in DC. Prod.* xi. 692.—Austral.
limonifolia, *Wall. Cat.* n. 1754.—Burma; Siam.
Lindeni, *Hook. f. Bot. Mag.* t. 6230.—N. Granat.
littoralis, *A. Cunn. in Ann. Nat. Hist.* Ser. I. i. (1838) 461.—N. Zel.
littoralis, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 401 (cf. timoriensis).—Ins. Timor.
Loureiri, *Hook. & Arn. Bot. Beech. Voy.* 206. t. 48.—China.
Loureirii, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 585 = urceolata.
lupinifolia, Salisb. Prod. 106 = V. Agnus-castus.
macrophylla, R. Br. Prod. 512 = Gmelina macrophylla.
madiensis, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 134.—Afr. trop.
melicopea, F. Muell. Fragm. v. 35 = acuminata.
Melleri, *Baker, in Journ. Linn. Soc.* xx. (1883) 227.—Madag.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 245.—Mexic.
moluccana, *Blume, Bijdr.* 813.—Ins. Molucc.
Mombassae, *Vatke, in Linnaea*, xliii. (1880–82) 533.—Afr. trop.
montevidensis, *Cham. l. c.* vii. (1832) 373.—Bras.
Morava, Buch.-Ham. ex Wall. Cat. n. 1752 = peduncularis.
multiflora, Miq. in Linnaea, xviii. (1844) 739 = divaricata.
multinervis, *Schau. in DC. Prod.* xi. 688.—Bras.
Negundo, *Linn. Sp. Pl.* 638.—As. trop. et subtrop.
Negundo, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 86 = pubescens.
obovata, *E. Mey. Comm. Pl. Afr. Austr.* i. 273.—Afr. austr.
orinocensis, *H. B. & K. Nov. Gen. et Sp.* ii. 247.—Venezuela.
ovata, Thunb. Fl. Jap. 257 = trifolia.
pachyclada, *Baker, in Journ. Linn. Soc.* xx. (1883) 227.—Madag.
pallida, Wall. Cat. n. 1751 = glabrata.
paludosa, *Vatke, in Linnaea*, xliii. (1880–82) 534.—Afr. trop.
paniculata, Lam. Encyc. ii. 612 = V. Negundo.
parviflora, *A. Juss. in Ann. Mus. Par.* vii. (1806) 76.—Ins. Philip.
peduncularis, *Wall. Cat.* n. 1753.—Ind. or.; Burma.
Pervillei, *Baker, Fl. Maurit.* 256.—Ins. Seychelles.
Petersiana, *Klotzsch, in Peters, Reise Mossamb. Bot.* 264.—Afr. trop.
phillyreaefolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 226.—Madag.
pinnata, Linn. Sp. Pl. 638 = Aglaia odorata.
polygama, *Cham. in Linnaea*, vii. (1832) 371.—Bras.
pteropoda, *Miq. Fl. Ind. Bat. Suppl.* 567.—Malaya.
pubescens, Heyne, ex Wall. Cat. n. 1755 = altissima.
pubescens, *Vahl, Symb. Bot.* iii. 85.—As. trop.
punctata, *Schau. in DC. Prod.* xi. 687.—Ins. Molucc.
repens, Blanco, Fl. Filip. ed. I. 513 = trifolia.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. iii. 826 = V. Leucoxylon?
rigens, *Griseb. Cat. Pl. Cub.* 216.—Cuba.
robusta, Lebas, in Rev. Hortic. (1869) 30 = V. Agnus-castus.
rotundifolia, Linn. f. Suppl. 294 = trifolia.

VITEX:—
rufescens, *A. Juss. in Ann. Mus. Par.* vii. (1806) 77.—Bras.
saligna, Roxb. Hort. Beng. 46; Fl. Ind. iii. 75 = V. Leucoxylon.
Sampsoni, *Hance, in Journ. Bot.* vi. (1868) 115.—China.
scabra, *Wall. Cat.* n. 1758.—Burma.
Schomburgkiana, *Schau. in DC. Prod.* xi. 690.—Guian.
Sellowiana, *Cham. in Linnaea*, vii. (1832) 108.—Bras.
sexdentata, Wall. Cat. n. 1759 = Caryopteris grata.
simplicifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 586.—Malacca.
simplicifolia, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 133.—Afr. trop.
sinuata, Medic. in Act. Acad. Theod. Palat. iv. Phys. (1780) 202. t. 8 = V. Agnus-castus?
spicata, *Lour. Fl. Cochinch.* 390.—China; Cochinch.
Strickeri, *Vatke & Hildebr. in Linnaea*, xliii. (1880–82) 532.—Afr. trop.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 567.—Sumatra.
Taruma, Mart. Syst. Mat. Med. Bras. 55 = montevidensis.
tettensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 264.—Afr. trop.
timoriensis, Walp. Rep. iv. 84 = littoralis, *Decne.*
Tracyana, F. Muell. Fragm. v. 61 = Clerodendron Tracyanum.
trichantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 434.—Madag.
triflora, *Vahl, Eclog. Am.* ii. 49.—Guian.
trifolia, *Linn. Sp. Pl.* 638.—As. et Austral. trop.
triphylla, Royle, Illustr. Bot. Himal. 299 = praec.
umbrosa, G. Don, ex Sabine, in Trans. Hort. Soc. v. (1824) 455 = Doniana.
umbrosa, *Sw. Prod. Veg. Ind. Occ.* 93.—Ind. occ.
undulata, Wall. Cat. n. 1756 = heterophylla.
uniflora, *Baker, in Journ. Bot.* xx. (1882) 221.—Madag.
urceolata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 585.—Malacca.
variifolia, Salisb. Prod. 107 = trifolia.
Vauthieri, *DC. ex Schau. in DC. Prod.* xi. 690.—Bras.
verticillata, Lam. Fl. Fr. ii. 363 = V. Agnus-castus.
vestita, *Wall. Cat.* n. 1750.—Malaya.
vitiensis, *Seem. Fl. Vit.* 190.—Ins. Fiji.
Wimberleyi, *Kurz, For. Fl. Brit. Burma*, ii. 271.—Ins. Andaman.
zanzibarensis, *Vatke, in Linnaea*, xliii. (1880–82) 533.—Afr. trop.
Zeyheri, *Sond. ex Schau. in DC. Prod.* xi. 693.—Afr. austr.
zeylanica, Burm. f. Fl. Ind. 138 (Stereospermi sp. ?).—Zeylan.
zeylanica, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 223 = altissima.

VITICASTRUM, Presl, Bot. Bemerk. 147 (1844) = **Sphenodesme**, Jack (Verben.).
racemosum, Presl, l. c. 148 = S. barbata.

VITICELLA, Dill. ex Moench, Meth. 296 (1794) = **Clematis**, Dill. (Ranunc.).
campaniflora, Bercht. & Presl, Rostl. i. Ranunc. 11 = C. campaniflora.
crispa, Bercht. & Presl, l. c. = C. crispa.
deltoidea, Moench, Meth. 297 = C. Viticella.
florida, Bercht. & Presl, Rostl. i. Ranunc. 10 = C. florida.

VITICELLA, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 220 = **Hydrophyllum**, Linn.

VITICENA, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 57, sphalm. = VINTICENA, Steud. = **Grewia**, Linn. (Tiliac.).

VITIPHOENIX, Becc. in Ann. Jard. Buitenz. ii. (1885) 91, 126. *PALMAE.*
filifera, *Becc. l. c.*—Ins. Fiji.

VITIS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 56 (1737). *AMPELIDEAE*, Benth. & Hook. f. i. 387.
ADENOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 417.
ALLOSAMPELA, Rafin. Med. Fl. ii. 122 (1830).
AMPELOPSIS, [Rich. in] Michx. Fl. Bor. Am. i. 159 (1803), ex parte.
BOTRIA, Lour. Fl. Cochinch. 153 (1790).

VITIS:—
CAUSONIS, Rafin. Med. Fl. ii. 122 (1830).
CAYRATIA, Juss. ex Guill. in Dict. Class. Hist. Nat. iv. 346 (1823).
CISSUS, Linn. Diss. Dass. 6 (1747); Amoen. Acad. i. 389 (1749).
COLUMELLA, Lour. Fl. Cochinch. 85 (1790).
GONOLOMA, Rafin. Sylva Tellur. 86 (1838).
INGENHOUSSIA, Dennst. Schluess. Hort. Malab. 33 (1818).
IRSIOLA, P. Br. Hist. Jamaic. 147. t. 4. f. 2 (1756).
ITUTERION, Rafin. Sylva Tellur. 86 (1838).
KEMOXIS, Rafin. l. c. (1838).
NEKEMIAS, Rafin. l. c. 87 (1838).
PEDASTIS, Rafin. l. c. (1838).
QUINARIA, Rafin. Med. Fl. ii. 122 (1830).
RINXOSTYLIS, Rafin. Sylva Tellur. 87 (1838).
SAELANTHUS, Forsk. Fl. Aegypt. Arab. 33 (1775).
SPINOVITIS, Romanet, in Compt. Rend. Par. xcii. (1881) 1096.
SPONDYLANTHA, Presl, Rel. Haenk. ii. 35. t. 53 (1834).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 112.—Abyss.
acapulcensis, *H. B. & K. Nov. Gen. et Sp.* vii. 230.—Mexic.
acerifolia, Rafin. Med. Fl. ii. 130 = riparia, rotundifolia.
acetosa, *F. Muell. Pl. Vict.* i. 94, in textu.—Austral.
acida, *Chapm. Fl. S. U. St.* 70.—Am. bor.
acida, Thunb. Fl. Japan. 11 (Quid?).—Java.
aconitifolia, Hance, in Journ. Linn. Soc. xiii. (1873) 77 = serjanaefolia.
acris, *F. Muell. Fragm.* ii. 75.—Austral.
aculeata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 88.—Ins. Timor.
acuminata, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 37 = caribaea.
acuminata, *Seem. in Bonplandia*, ix. (1861) 255.—Ins. Fiji.
adenantha, *Baker, in Oliver, Fl. Trop. Afr.* i. 403.—Afr. trop.
adenocaulis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 79.—Abyss.
adnata, *Wall. Cat.* n. 5998.—As. et Austral. trop.
adstricta, Hance, in Journ. Bot. xx. (1882) 258 = vinifera.
aegirophylla, *Boiss. Fl. Orient.* i. 956.—Turkestan.
aestivalis, *Michx. Fl. Bor. Am.* ii. 230.—Am. bor.
aestivalis, Torr. in Pacif. Rail. Rep. vii. 9 = arizonica.
africana, *Spreng. Syst.* i. 778.—Afr. trop.
Afzelii, *Baker, in Oliver, Fl. Trop. Afr.* i. 389.—Afr. trop.
albida, *Baker, in Mart. Fl. Bras.* xiv. II. 212.—Bras.
alexandrina, Fisch. Hort. Gorenk. ed. II. (1812) 53 = vinifera?
amara, Rafin. Med. Fl. ii. 132 = cordifolia.
amboinensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 76.—Ins. Amboin.
amplexa, *Baker, in Oliver, Fl. Trop. Afr.* i. 403.—Afr. trop.
amurensis, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 266 = vinifera.
anamalayana, *Bedd. in Trans. Linn. Soc.* xxv. (1865) 213.—Ind. or.
andongensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 406.—Afr. trop.
anemonifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 80.—Ins. Timor.
angolensis, *Baker, in Oliver, Fl. Trop. Afr.* i. 410.—Afr. trop.
angulata, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 176, in nota = repanda.
angulata, Rafin. Med. Fl. ii. 133 = rotundifolia.
angustifolia, Benth. Fl. Hongk. 54 = corniculata.
angustifolia, *Wall. Cat.* n. 6033.—Reg. Himal.; Malaya; China.
angustissima, *F. Muell. Fragm.* i. 141.—Austral.
antarctica, *Benth. Fl. Austral.* i. 447; *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 83.—Austral.
apiana, Rafin. Med. Fl. ii. 140 = vinifera.
apiifolia, Hort. ex Steud. Nom. ed. II. ii. 777 = vinifera.
apyrena, Hort. ex Roem. & Schult. Syst. v. 300 = vinifera.
aquosa, Wall. Cat. n. 6000 = repanda.
arachnoidea, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 37 = caribaea.
aralioides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 411.—Afr. trop.

VITIS:—
araneosa, *Dalz. & Gibs. Bomb. Fl.* 41.—Ind. or.
araneosa, *Le Conte, in Proc. Acad. Philad.* 1852–53 (1854) 272.—Am. bor.
araneosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 94.—Sumatra.
arborea, *Linn. Sp. Pl.* 203.—Oriens; Am. bor.
arcuata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 414.—Afr. trop.
arguta, *Baker, in Oliver, Fl. Trop. Afr.* i. 392.—Afr. trop.
arizonensis, [Parry,] in Rep. Commiss. Agric. 1870 (1871) 416 = seq.
arizonica, *Engelm. in Am. Natural.* ii. (1869) 321; ix. (1875) 268.—Am. bor.
articulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 76.—Borneo.
asarifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 396.—Afr. trop.
asperifolia, *Baker, in Mart. Fl. Bras.* xiv. II. 207.—Bras.
assamica, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 648.—Assam.
assimilis, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 302 = lanceolaria.
atroviridis, Wall. Cat. n. 6040 = Gynostemma pedatum.
augustissima, F. Muell. Census, 24, err. typ. = angustissima.
auriculata, *Wall. Cat.* n. 6031.—Reg. Himal.; Burma; Java.
Bainesii, *Hook. Bot. Mag.* t. 5472.—Afr. trop.
barbata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 478.—Reg. Himal.; Burma.
Barteri, *Baker, in Oliver, Fl. Trop. Afr.* i. 390.—Afr. trop.
Baudiniana, F. Muell. Fragm. iv. 136 = antarctica.
Berlanderi, *Planch. in Compt. Rend. Par.* xci. (1880) 425.—Texas; N. Mexic.
bicolor, Rafin. Med. Fl. ii. 140; Le Conte, in Proc. Acad. Philad. 1852–53 (1854) 272 = aestivalis.
bifida, Rafin. l. c. 128 = aestivalis?
bipinnata, Torr. & Gray, Fl. N. Am. i. 243 = arborea.
biternata, *Baker, in Journ. Bot.* xx. (1882) 51.—Madag.
blanda, Rafin. Med. Fl. ii. 128 = V. Labrusca.
blanda, *Sweet, Hort. Brit.* ed. I. 73.—Am. bor.
Blumeana, F. Muell. Sel. Pl. Indust. Cult. 255 = pallida.
bombycina, *Baker, in Oliver, Fl. Trop. Afr.* i. 399.—Afr. trop.
bororensis, *Baker, in Oliver, Fl. Trop. Afr.* i. 411.—Afr. trop.
brachypoda, *F. Muell. Fragm.* ix. 125.—Austral.
bracteata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = discolor?
bracteata, Rafin. Med. Fl. ii. 125; Le Conte, in Proc. Acad. Sc. Philad. 1852–53 (1854) 271 = V. Labrusca?
bracteolata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 483.—Reg. Himal.
bryoniaefolia, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 85 = vinifera.
Burchellii, *Baker, in Mart. Fl. Bras.* xiv. II. 201.—Bras.
caesia, *Sabine, in Trans. Hort. Soc.* v. (1824) 447.—Afr. trop.
californica, *Benth. Bot. Voy. Sulph.* 10 (sphalm. Vitex).—Calif.
californica, [Parry] in Rep. Commiss. Agric. 1870 (1871) 415 = arizonica.
callosa, Rafin. Med. Fl. ii. 125 = rotundifolia.
campestris, *Baker, in Mart. Fl. Bras.* xiv. II. 200.—Bras.
campestris, [Nutt.] in Fras. Cat. (1813) n. 88 = riparia?
campylocarpa, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 302.—Assam; Burma.
cana, Rafin. Med. Fl. ii. 140 = V. Labrusca?
canarensis, *Dalz. in Hook. Kew Journ.* iii. (1851) 123.—Ind. or.
candicans, *Engelm. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 166.—Am. bor.
canina, Rafin. Med. Fl. ii. 127 = V. Labrusca.
cantoniensis, *Seem. Bot. Voy. Herald*, 370.—Malacca; China.
capensis, *Burm. f. Fl. Cap. Prod.* 7; *Thunb. Prod. Pl. Cap.* 44.—Afr. austr.
capreolata, *D. Don, Prod. Fl. Nep.* 188.—Reg. Himal.
cardiophylla, F. Muell. Fragm. ii. 73 = repens.

VITIS:—
caribaea, *DC. Prod.* i. 634.—Am. bor.; Ind. occ.
caribaea, Hook. & Arn. Bot. Beech. Voy. 327 = californica.
carnosa, Wall. Cat. n. 6018 = trifolia.
Catawba, Hort. ex C. Koch, Dendrol. i. 551 = V. Labrusca.
cavicaulis, *Baker, in Oliver, Fl. Trop. Afr.* i. 400.—Afr. trop.
cebennensis, Jord. ex Nym. Consp. 136 = vinifera.
cerasiformis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 251.—Java.
cerifera, Teijsm. & Binn. Cat. Hort. Bog. 168, sphalm. = praec.
Chantinii, *Lécard, ex Carr. in Rev. Hortic.* (1881) 457.—Afr. trop.
chloroleuca, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 406.—Afr. trop.
chontalensis, *Seem. in Journ. Bot.* vii. (1869) 332.—Nicaragua.
ciliata, Rafin. Med. Fl. ii. 129 = V. Labrusca?
cinerea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 (*cineracea*) = flexuosa.
cinnamomea, Wall. Cat. n. 5989 C = tomentosa.
cinnamomea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 483.—Malaya.
cirrhosa, *Thunb. Prod. Pl. Cap.* 44.—Afr. austr.
clematidea, *F. Muell. Fragm.* ii. 74.—Austral.
coccinea, *Baker, in Mart. Fl. Bras.* xiv. II. 216.—Bras.
coffeaecarpa, *Teijsm. & Binn. Cat. Hort. Bog.* 168, nomen.—Java.
Coignetiae, *Pulliat, ex Planch. in Journ. Vigne Amér.* (1883) 186.—Japon.
columbina, Rafin. Med. Fl. ii. 131 = riparia.
Commersoni, *Baker, Fl. Maurit.* 54.—Ins. Maurit.
compositifolia, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 659.—Penang; Malacca.
concinna, *Baker, in Oliver, Fl. Trop. Afr.* i. 413.—Afr. trop.
concolor, Rafin. Med. Fl. ii. 130 = riparia.
congesta, *Baker, in Oliver, Fl. Trop. Afr.* i. 412.—Afr. trop.
constricta, *Baker, l. c.* 409.—Afr. trop.
coralloides, Hook. f. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 659 = elegans.
cordata, Dum.-Cours. Bot. Cult. ed. II. iv. 619 = indivisa.
cordata, Wall. Cat. n. 6001; Voigt, Hort. Suburb. Calc. 28 = repens.
cordifolia, Darlingt. Fl. Cestr. 151 = riparia?
cordifolia, *Lam. Illustr.* ii. 134.—Am. bor.
cordifolia, Michx. Fl. Bor. Am. ii. 231 = praec.?
cordifolia, Roth, ex Roem. & Schult. Syst. v. 318 = lanata.
coriacea, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 78.—Ins. Celebes.
coriacea, Shuttl. ex C. Koch, Dendrol. i. 550 = candicans.
corinthiaca, Rafin. Med. Fl. ii. 141 = vinifera.
corniculata, *Benth. Fl. Hongk.* 54.—Ins. Hongkong.
cornifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 390.—Afr. trop.
corylifolia, *Baker, l. c.* 396.—Afr. trop.
costata, Wall. Cat. n. 6011 = discolor?
crassifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 391.—Afr. trop.
crassiuscula, *Baker, l. c.* 406.—Afr. trop.
crenata, *Thunb. ex Sw. in Vet. Akad. Nya Handl. Stockh.* (1825) 428.—Ind. occ.
crenata, Wall. Cat. n. 6021 = trifolia.
Cumingiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 416.—Ins. Philipp.
Currori, *Baker, in Oliver, Fl. Trop. Afr.* i. 401.—Afr. trop.
curvipoda, *Baker, l. c.* 413.—Afr. trop.
cuspidata, Lynch, in Proc. Linn. Soc. (1875–80) 38 = inconstans.
cuspidifera, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 91, in adnot.—Reg. Himal.
cylindrica, Rafin. Med. Fl. ii. 141 = vinifera.
cymosa, *Blume, Bijdr.* 195.—Java.
cymosa, Wall. Cat. n. 6017 = japonica.
cyphopetala, *Baker, in Oliver, Fl. Trop. Afr.* i. 407.—Afr. trop.
dapsilis, [Roxas Clem. in] Herrera, Agric. (1818) i. 327 = orientalis.
debilis, *Baker, in Oliver, Fl. Trop. Afr.* i. 403.—Afr. trop.

VITIS:—
dentata, *Link, Enum. Hort. Berol.* i. 236.—Hab.?
denticulata, Rafin. Ann. Nat. 11 (Quid?).—Am. bor.
dichotoma, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 77.—Java.
Diepenhorstii, *Miq. l. c.* 77, in textu.—Sumatra.
diffusa, *Miq. l. c.* 83.—Malaya.
diffusiflora, *Baker, in Oliver, Fl. Trop. Afr.* i. 390.—Afr. trop.
digitata, Rafin. Med. Fl. ii. 125 = V. Labrusca?
dimidiata, Rafin. l. c. 129 = riparia.
discolor, *Dalz. in Hook. Kew Journ.* ii. (1850) 39.—Ind. or.; Malaya.
dissecta, *Baker, in Oliver, Fl. Trop. Afr.* i. 414.—Afr. trop.
divaricata, Wall. Cat. n. 5994 = lanata, indica.
diversifolia, Wall. Cat. n. 5996 = discolor.
Duarteana, *Baker, in Mart. Fl. Bras.* xiv. II. 204.—Bras.
dubia, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 661.—Reg. Himal.
Durandii, Lécard, ex Carr. in Rev. Hortic. (1881) 456 = rufescens.
elegans, C. Koch, in Ind. Sem. Hort. Berol. (1855) App. 16 = heterophylla.
elegans, *Kurz, in Tijdschr. Nederl. Ind.* xxviii. (1865) 166.—Singapore.
elliptica, *Hemsl. Biol. Centr. Am. Bot.* i. 208.—Mexic.
elongata, *Wall. Cat.* n. 6016.—Ind. or.
enneaphylla, *Eichl. in Mart. Fl. Bras.* xiv. II. 217.—Bras.
erioclada, *Wight & Arn. Prod.* 130.—Ind. or.
eriopoda, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 95.—Sumatra.
erosa, *Baker, in Mart. Fl. Bras.* xiv. II. 210.—Am. trop.
erythroclada, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 301.—Burma.
erythrodes, *Fresen. in Mus. Senckenb.* ii. (1837) 284.—Abyss.
Faidherbii, *Lécard, ex Carr. in Rev. Hortic.* (1881) 457.—Afr. trop.
farinosa, Rafin. Med. Fl. ii. 140 = vinifera.
farinosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 394.—Afr. trop.
ferruginea, Rafin. Med. Fl. ii. 128 = V. Labrusca.
ficifolia, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 86 = V. Labrusca.
Figariana, *Baker, in Oliver, Fl. Trop. Afr.* i. 415.—Afr. trop.
filiformis, Lam. Illustr. ii. 135 = pentaphylla.
flavicans, *Baker, in Oliver, Fl. Trop. Afr.* i. 413.—Afr. trop.
flexuosa, Hort.—Cf. Gard. Chron. (1882) II. 491 = V. Labrusca.
flexuosa, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332.—Ind. or.; China; Japon.
floribunda, *Baker, in Journ. Bot.* xx. (1882) 50.—Madag.
floridana, Rafin. Med. Fl. ii. 133 = vulpina?
fluminicola, Steud. Nom. ed. II. ii. 777 = repanda.
Forsteniana, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 86.—Ins. Amboin.
fulva, Rafin. Med. Fl. ii. 124 = aestivalis?
furcata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 646.—Malaya.
Gardneri, *M. Laws. l. c.* 656.—Zeylan.
geniculata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 81.—Java.
Gibertii, *Baker, in Mart. Fl. Bras.* xiv. II. 211.—Uruguay.
gigantea, *Bedd. in Trans. Linn. Soc.* xxv. (1865) 212.—Ind. or.
glaberrima, Wall. Cat. n. 5991, partim = hastata.
glaberrima, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 476.—Malaya.
glabrata, Heyne, ex Roth, Nov. Pl. Sp. 156 = latifolia.
glandulosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 479.—Reg. Himal.
glareosa, Rafin. Med. Fl. ii. 126 = aestivalis.
glauca, Wall. Cat. n. 5990 = repens.
glauca, *Wight & Arn. Prod.* 126.—Ind. or.
glaucophylla, *Baker, in Oliver, Fl. Trop. Afr.* i. 392.—Afr. trop.
glomerata, Rafin. Med. Fl. ii. 142 = caribaea.
glyptocarpa, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 645.—Zeylan.

VITIS:—
gongylodes, *Baker, in Mart. Fl. Bras.* xiv. II. 209.—Bras.; Peruv.
gorgonobotrys, *Webb, in Hook. Kew Journ.* ii. (1850) 313.—Afr. trop.
gracilis, *Baker, in Oliver, Fl. Trop. Afr.* i. 404.—Afr. trop.
gracilis, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 477.—Singapore.
Grantii, *Baker, in Oliver, Fl. Trop. Afr.* i. 400.—Afr. trop.
granulosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 80.—Peruv.
grisea, *Baker, in Oliver, Fl. Trop. Afr.* i. 395.—Afr. trop.
Hancockii, *Hance, in Journ. Bot.* xx. (1882) 4.—China.
Hardyi, *Lécard, ex Carr. in Rev. Hortic.* (1881) 458.—Afr. trop.
hastata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 85, in adnot.—Ind. or.; Java.
havanica, Regel, Cat. Pl. Hort. Aksakov. 146 = caribaea?
hederacea, *Ehrh. Beitr.* vi. (1791) 85.—Am. bor.
hederacea, Wall. Cat. n. 6019 = obtecta, capreolata.
Helferi, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 662.—Burma.
helvola, [Roxas Clem. in] Herrera, Agric. (1818) i. 327 = orientalis.
heptaphylla, *Linn. Mant.* ii. 212.—Ind. or.
heracleifolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 398.—Afr. trop.
heterophylla, *Thunb. Fl. Jap.* 103.—China; Japon.
heterophylla, Wall. Cat. n. 6036 = Rheedei.
Heyneana, Roem. & Schult. Syst. v. 318; DC. Prod. i. 634 = lanata.
Heyneana, *Wall. Cat.* n. 5988.—Ind. or.
hiemalis, Auct. = *hyemalis*.
himalayana, *Brandis, For. Fl.* 100.—Ind. or.
hispida, *Eckl. & Zeyh. Enum.* 56.—Afr. austr.
Hochstetteri, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 83.—Abyss.
Hookeri, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 661.—Reg. Himal.
humulifolia, Hort.—Cf. Gard. Chron. (1882) II. 491 = heterophylla.
hyemalis, Dum.-Cours. Bot. Cult. ed. II. iv. 619 = cordifolia?
hyemalis, Rafin. Med. Fl. ii. 125 = rotundifolia.
hypoglauca, *F. Muell. Pl. Vict.* i. 94.—Austral.
ibuensis, *Baker, in Oliver, Fl. Trop. Afr.* i. 402.—Afr. trop.
imperialis, *Miq. Fl. Ind. Bat. Suppl.* 518.—Sumatra.
inaequalis, Wall. Cat. n. 6010 = discolor.
incarnata, *Teijsm. & Binn. Cat. Hort. Bog.* 168, nomen.—Sumatra.
incisa, Jacq. Hort. Schoenb. iv. 14 = riparia.
incisa, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 243.—Texas.
inconstans, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 91.—Japon.
indica, Hook. & Arn. Bot. Beech. Voy. 260 = lanata.
indica, *Linn. Sp. Pl.* 202.—Ind. or.
indica, Sw. Obs. Bot. 95 (excl. syn.) = caribaea.
indica, Thunb. Fl. Jap. 103 = flexuosa.
indica, Thw. Enum. Pl. Zeyl. 63 = erioclada.
indivisa, *Willd. Berl. Baumz.* ii. 538.—Am. bor.
integrifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 391.—Afr. trop.
integrifolia, Rafin. Fl. Ludov. 89 (Quid?).—Am. bor.
intermedia, Muhl. Cat. 26 = aestivalis.
intricata, *Baker, in Oliver, Fl. Trop. Afr.* i. 404.—Afr. trop.
inundata, *Baker, in Mart. Fl. Bras.* xiv. II. 205.—Bras.
ipomoeifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 395.—Afr. trop.
irawadyana, Steud. Nom. ed. II. ii. 778 = japonica.
Isabella, Otto & Dietr. Allg. Gartenz. x. (1842) 353 = vinifera?
japonica, Thunb. Fl. Jap. 104.—China; Japon.; Malaya.
jatrophoides, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 400.—Afr. trop.
javalensis, *Seem. in Journ. Bot.* vii. (1869) 332.—Nicaragua.
javanica, *Spreng. Syst.* iv. *Cur. Post.* 90.—Java.

VITIS:—
juncea, *Baker, in Oliver, Fl. Trop. Afr.* i. 401.—Afr. trop.
Kanguruh, Hort. ex DC. Prod. i. 629 = antarctica.
Kleinii, Wall. Cat. n. 6008 = glauca, latifolia.
Labrusca, *Linn. Sp. Pl.* 203.—Am. bor.
Labrusca, Linn. l. c., partim = caribaea.
Labrusca, Lour. Fl. Cochinch. 155 = triloba?
Labrusca, Scop. Fl. Carn. ed. II. i. 169 = vinifera.
Labrusca, Thunb. Fl. Jap. 103 = Coignetiae.
Labrusca, Walt. Fl. Carol. 242 = aestivalis.
labruscoides, Muhl. Cat. 26 = aestivalis.
labruscoides, Muhl. & Rafin. in Rafin. Med. Fl. ii. 127 = V. Labrusca.
laciniosa, Linn. Sp. Pl. 203 = vinifera.
laciniosa, Marsh. Arbust. Am. 165 = aestivalis.
laeta, Wall. Cat. n. 6002 = repanda.
laevigata, F. Muell. Sel. Pl. Indust. Cult. 255 = auriculata.
lanata, Roxb. ex Wall. Cat. n. 5995 C, D = barbata.
lanata, *Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 660.—Ind. or.; China.
lanata, Wall. Cat. n. 5995 A? = tomentosa.
lanceolaria, *Wall. Cat.* n. 6013, partim.—Ind. or.; Malaya.
lanceolaria, Wall. l. c. H = rumicisperma.
lanceolaria, Wight, Ic. t. 28 = sulcata.
Landuk, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 90.—Java.
latifolia, Buch.-Ham. ex Wall. Cat. n. 5994 G = barbata.
latifolia, Lam. ex Steud. Nom. ed. II. ii. 778 = glauca.
latifolia, Rafin. Med. Fl. ii. 126 = V. Labrusca.
latifolia, *Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 661.—Ind. or.
latifolia, Vahl, ex Steud. Nom. ed. II. ii. 778 = adnata.
Lecardii, *Carr. in Rev. Hortic.* (1881) 456.—Afr. trop.
leeoides, Maxim. in Bull. Acad. Pétersb. xix. (1874) 158 = cantoniensis.
lenticellata, *Baker, in Journ. Linn. Soc.* xx. (1883) 122.—Madag.
leonensis, *Baker, in Oliver, Fl. Trop. Afr.* i. 398.—Afr. trop.
leucostaphyla, Dennst. Schluess. Hort. Malab. 33 = lanceolaria.
leucostyla, Steud. Nom. ed. II. ii. 778, sphalm. = praec.
Linnaei, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 176 = repanda.
Linnaei, *Wall. Cat.* n. 5987.—Ind. or.
Linsecomii, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 451 = aestivalis.
lonchiphylla, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 646.—Zeylan.
longepedunculata, *Teijsm. & Binn. Cat. Hort. Bog.* 168.—Ins. Ceram.
longifolia, Rafin. Med. Fl. ii. 129 = cordifolia?
Lowii, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 165.—Borneo.
lucida, Fisch. ex Steud. Nom. ed. II. ii. 778 = antarctica.
luteola, Rafin. Med. Fl. ii. 127 = V. Labrusca.
macrophylla, Royle, Illustr. Bot. Himal. 145 = himalayana.
macropus, *Hook. Bot. Mag.* t. 5479.—Afr. trop.
macrostachya, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 94.—Malaya.
Mannii, *Baker, in Oliver, Fl. Trop. Afr.* i. 412.—Afr. trop.
Mappia, *Baker, Fl. Maurit.* 54.—Ins. Maurit.
maritima, Rafin. Med. Fl. ii. 142 = caribaea.
maura, Rafin. l. c. 141 (Quid?).—Marocc.
Metziana, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 84.—Sumatra.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* i. 203.—Mexic.
microdiptera, *Baker, in Journ. Linn. Soc.* xviii. (1881) 266.—Madag.
microdonta, *Baker, in Journ. Bot.* xx. (1882) 50.—Madag.
Miqueliana, *Baker, in Mart. Fl. Bras.* xiv. II. 210.—Am. trop.
modesta, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 84.—Sumatra.
mollis, Baker, in Oliver, Fl. Trop. Afr. i. 413 = serpens.
mollis, *Wall. Cat.* n. 6025.—Reg. Himal.; Malaya.
mollissima, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 482.—Malaya.
monosperma, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = adnata.

VITIS:—
montana, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 652.—Reg. Himal.
montana, Rafin. Med. Fl. ii. 130 = riparia.
monticola, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 450 = aestivalis.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 180.—Afr. trop.
Motleyi, *Hook. f. in Trans. Linn. Soc.* xxiii. (1860) 165.—Borneo.
mucronata, Wall. ex Steud. Nom. ed. II. ii. 778 = lanceolaria.
multiloba, Rafin. Med. Fl. ii. 125 = aestivalis.
multistriata, *Baker, in Oliver, Fl. Trop. Afr.* i. 410.—Afr. trop.
muricata, Wall. Cat. n. 6015 = lanceolaria.
muscadina, Rafin. Med. Fl. ii. 132 = rotundifolia.
mustangensis, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 451 = candicans.
mutabilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 75.—Java.
neilgherrensis, Wight, Ic. t. 965 = himalayana.
nepalensis, Royle, Illustr. Bot. Himal. 149 = capreolata.
nervosa, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 650.—Reg. Himal.
neurosa, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 204 = planicaulis.
nitens, *F. Muell. Fragm.* ii. 73.—Austral.
nitida, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 662.—Malaya.
nodosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 87.—Malaya.
novemfolia, *Wall. Cat.* n. 6030.—Singapore.
nymphaeifolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 393.—Afr. trop.
obliqua, Rafin. Med. Fl. ii. 128 = V. Labrusca?
oblonga, *Benth. Fl. Austral.* i. 447.—Austral.
obovata, *Baker, in Mart. Fl. Bras.* xiv. II. 207, in obs.—Guatemala.
obovata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 658.—Reg. Himal.
obovata, Rafin. Med. Fl. ii. 135 = V. Labrusca.
obtecta, *Wall. Cat.* n. 6026.—Reg. Himal.; Burma; China.
obtusata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 414.—Afr. trop.
obtusifolia, Buch.-Ham. ex Wall. Cat. sub n. 6004 = tomentosa.
occidentalis, Bartram, ex Le Conte, in Rep. Commiss. Patents (Agric.), 1857 (1858) 229 = V. Labrusca.
ochracea, *Teijsm. & Binn. Cat. Hort. Bog.* 388.—Ins. Celebes.
odoratissima, Donn, Cat. Hort. Cantab. ed. IX. 71 = riparia.
opaca, *F. Muell. ex Benth. Fl. Austral.* i. 450.—Austral.
oppositifolia, Buch.-Ham. ex Steud. Nom. ed. II. ii. 778 = tomentosa.
orientalis, Boiss. Fl. Orient. i. 956 = arborea.
orientalis, [*Roxas Clem. in*] *Herrera, Agric.* (1818) i. 327.—Hispan. (cult.).
oxyodonta, *Baker, in Journ. Bot.* xx. (1882) 50.—Madag.
oxyphylla, *A. Rich. Tent. Fl. Abyss.* i. 114.—Abyss.
oxyphylla, Wall. Cat. n. 6035 = bracteolata.
Pagnuccii, *Romanet, in Congr. Géog. Toul.* (1884); *et in Journ. Vigne Amér.* (1885) 283, nomen.—China.
pallida, *Wight & Arn. Prod.* 125.—As. et Afr. trop.
palmata, *Baker, in Mart. Fl. Bras.* xiv. II. 216.—Bras.
palmata, Le Conte, in Proc. Acad. Sc. Philad. 1852–53 (1854) 274 = V. Labrusca.
palmata, Vahl, Symb. Bot. iii. 42 = aestivalis.
palmatifida, *Baker, in Oliver, Fl. Trop. Afr.* i. 397.—Afr. trop.
paniculata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 507.—Ins. Socotra.
paniculata, Heyne, ex Wall. Cat. n. 6022 = japonica, repanda.
pannosa, *Baker, in Mart. Fl. Bras.* xiv. II. 206.—Bras.
pannosa, *Baker, in Oliver, Fl. Trop. Afr.* i. 402.—Afr. trop.
papillata, *Hance, in Journ. Bot.* xvi. (1878) 226.—China.
papuana, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 74.—N. Guin.
Parkeri, *Baker, in Mart. Fl. Bras.* xiv. II. 209.—Guian.

VITIS:—
parviflora, Baker, in Oliver, Fl. Trop. Afr. i. 392, sphalm. = seq.
parvifolia, Roxb. Hort. Beng. 18; Fl. Ind. i. 662 = flexuosa.
patens, Colmeiro, Enum. Pl. Hisp.-Lusit. i. 545, sphalm. = seq.
patiens, [Roxas Clem. in] Herrera, Agric. (1818) i. 327 = orientalis.
paucidentata, *Baker, in Oliver, Fl. Trop. Afr.* i. 410. —Afr. trop.
pedata, *Vahl, ex Wall. Cat.* n. 6027.—As. trop.
pedicellata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 650. —Reg. Himal.
peduncularis, *Wall. Cat.* n. 6024.—Malaya.
peltata, Rafin. Med. Fl. ii. 133 = rotundifolia, riparia?
pendula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 407.—Afr. trop.
penninervis, *F. Muell. Fragm.* vi. 177.—Austral.
pentagona, Buch.-Ham. ex Wall. Cat. n. 5994 F = lanata.
pentagona, *Voigt, Hort. Suburb. Calc.* 28.—Burma.
pentaphylla, Guill. & Perr. Fl. Seneg. Tent. i. 135 = multistriata.
pentaphylla, Miq. Ann. Mus. Bot. Lugd. Bat. i. 91 = serjanaefolia.
pentaphylla, *Thunb. Fl. Jap.* 105.—Japon.
pergamacea, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 73.— Ins. Celebes.
persica, *Boiss. Fl. Orient.* i. 955.—Persia; Afghan.
Piasezkii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 461.—China.
pinnata, *Vahl, Symb. Bot.* iii. 43.—Hab.?
pisicarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 79.— N. Guin.
planicaulis, *Hook. f. Bot. Mag.* t. 5685.—Reg. Himal.
platanifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 398. —Afr. trop.
Pohlii, *Baker, in Mart. Fl. Bras.* xiv. II. 205.—Bras.
Poiretiana, Rafin. Med. Fl. ii. 134 = rotundifolia, riparia?
polita, Miq. Ann. Mus. Bot. Lugd. Bat. i. 95 = Pterisanthes polita.
polydactyla, *Miq. l. c.* 80.—Sumatra.
polystachya, *Wall. Cat.* n. 6028, partim.—Burma; Malaya.
polystachya, Wall. l. c., partim, et n. 9031 = nitida.
polythyrsa, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 89.— Sumatra.
populifolia, Lindh. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 165 = rupestris.
populifolia, Rafin. Med. Fl. ii. 131 = riparia.
populnea, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 83.— Afr. trop.
producta, *Baker, in Oliver, Fl. Trop. Afr.* i. 389.— Afr. trop.
prolifera, Rafin. Med. Fl. ii. 134 = V. Labrusca.
pruriens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 408.—Afr. trop.
psoralifolia, F. Muell. Fragm. ii. 75 = trifolia.
pterisantha, Miq. Ann. Mus. Bot. Lugd. Bat. i. 94 = Pterisanthes cissoides.
pterophora, *Baker, in Mart. Fl. Bras.* xiv. II. 213.— Bras.
pubescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 90.— Mexic.
pubiflora, *Miq. l. c.* 74.—Sumatra.
pulcherrima, *Eichl. in Mart. Fl. Bras.* xiv. II. 217.— Bras.
pullaria, *Le Conte, in Proc. Acad. Sc. Philad.* 1852–53 (1854), 273.—Am. bor.
punctata, Rafin. Med. Fl. ii. 140 (Quid?).—Hab.?
Purani, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 188 = flexuosa.
quadrangularis, *Wall. Cat.* n. 5992.—As. et Afr. trop.
quadricornuta, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 85. —Java.
quinquefolia, Lam. Illustr. ii. 135 = hederacea.
quinquefolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = japonica.
racemifera, *Jack, in Malay. Misc.* ii. (1822) n. VII. 94. —Sumatra.
radiata, Zipp. ex Span. in Linnaea, xv. (1841) 184, nomen = adnata.
Rafflesiae, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 76.— Sumatra.
repanda, *Wight & Arn. Prod.* 125.—Ind. or.
repens, Wall. Cat. n. 5999 B = adnata.
repens, *Wight & Arn. Prod.* 125.—Ind. or.; Austral.

VITIS:—
reticulata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 655. —Zeylan.
Rheedei, *Wight & Arn. Prod.* 127.—Ind. or.
rhombifolia, *Baker, in Mart. Fl. Bras.* xiv. II. 207.— Am. trop.
riparia, A. Gray, in Mem. Am. Acad. N. S. iv. (1849) 30 = arizonica.
riparia, *Michx. Fl. Bor. Am.* ii. 231.—Am. bor.
riparia, Wall. Cat. n. 6038 = repanda.
Romaneti, *Romanet, in Compt. Rend. Par.* (1881) 1096.—China.
rosea, Royle, Illustr. Bot. Himal. 149 = repanda.
rostrata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 85.— Borneo.
rotundifolia, *Michx. Fl. Bor. Am.* ii. 231.—Am. bor.; Mexic.
Roxburghii, *Wight & Arn. Prod.* 127.—Ind. or.
Royleana, Hort. ex C. Koch, Dendrol. i. 556 = capreolata.
rubifolia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 480.— Reg. Himal.
rubiginosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 394.—Afr. trop.
rubra, Desf. Tabl. ed. I. 140 = riparia.
rufa, Rafin. Med. Fl. ii. 140 = V. Labrusca?
rufescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 83.— Afr. trop.
rufula, *Miq. l. c.* 94.—Sumatra.
rugosa, Hook. f. & Thoms. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 650 = nervosa.
rugosa, Rafin. Med. Fl. ii. 127 = V. Labrusca.
rugosa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 480 = lanata, indica.
rumicisperma, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 661.—Reg. Himal.
rupestris, *Scheele, in Linnaea*, xxi. (1848) 591.— Texas.
saccharina, Rafin. Med. Fl. ii. 140 = vinifera?
sagittifolia, M. Laws. in Hook. f. Fl. Brit. Ind. i. 645 = hastata.
salmonea, *Baker, in Oliver, Fl. Trop. Afr.* i. 394.— Afr. trop.
salutaris, *Baker, in Mart. Fl. Bras.* xiv. II. 211.— Am. trop.
saponaria, *Seem. in Bonplandia*, ix. (1861) 254.— Austral.; Ins. Pacif.
sarcocephala, *Schweinf. ex Oliver, in Hook. Ic. Pl.* t. 1253.—Afr. trop.
saxatilis, Rafin. Med. Fl. ii. 124 = aestivalis.
scabra, *Baker, in Mart. Fl. Bras.* xiv. II. 202.— Bras.
scabricaulis, *Baker, l. c.* 213.—Bras.
scabriuscula, Wall. ex Steud. Nom. ed. II. ii. 778 (Quid?).—Ind. or.
Schimperiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 112.—Abyss.
Selloana, *Baker, in Mart. Fl. Bras.* xiv. II. 204.— Bras.
semicordata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 481 = himalayana.
semiglabra, *Sond. in Linnaea*, xxiii. (1850) 24.— Afr. austr.
serjanaefolia, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 159.—Japon.
serotina, Rafin. Med. Fl. ii. 126 = aestivalis?
serpens, *Baker, in Oliver, Fl. Trop. Afr.* i. 412.—Afr. trop.
serratifolia, Wight & Arn. Prod. 128 = lanceolaria.
serrulata, Wall. Cat. n. 6034 = capreolata.
sessilifolia, *Baker, in Mart. Fl. Bras.* xiv. II. 205.— Bras.
setosa, *Wall. Cat.* n. 6009.—Ind. or.
sicyoides, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 83.— Am. trop.
Sieboldii, Hort. ex C. Koch, Dendrol. i. 549 = V. Labrusca.
sikkimensis, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 650.—Reg. Himal.
silvestris, Roth, Enum. Pl. Phan. Germ. i. 746 = vinifera.
Simsiana, *Baker, in Mart. Fl. Bras.* xiv. II. 214.— Bras.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 125 = heterophylla.
sinuata, G. Don, Gen. Syst. i. 711 = aestivalis.
sinuosa, Bosc, ex [Roxas Clem. in] Herrera, Agric. (1818) i. 326 = vinifera?

VITIS:—
Smithiana, *Baker, in Oliver, Fl. Trop. Afr.* i. 390.— Afr. trop.
spectabilis, *Kurz, in Journ. Bot.* xii. (1874) 196.— Reg. Himal.
spinosa, *Baker, in Mart. Fl. Bras.* xiv. II. 204.— Bras.
stenoloba, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 408.—Afr. trop.
stercoracea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = indica.
sterculifolia, *F. Muell. ex Benth. Fl. Austral.* i. 450. —Austral.
stipulacea, *Baker, in Oliver, Fl. Trop. Afr.* i. 408.— Afr. trop.
striata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 79.— Chili.
subaphylla, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 506.—Ins. Socotra.
subciliata, *Baker, in Oliver, Fl. Trop. Afr.* i. 409.— Afr. trop.
suberecta, *Baker, in Mart. Fl. Bras.* xiv. II. 206.— Bras.
suberosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* i. 392.—Afr. trop.
subrhomboidea, *Baker, in Mart. Fl. Bras.* xiv. II. 208. —Bras.
succinea, Rafin. Med. Fl. ii. 140 = vinifera?
succisa, Hance, ex Walp. Ann. ii. 231 = flexuosa.
sulcata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 661.— Ind. or.
sulcicaulis, *Baker, in Mart. Fl. Bras.* xiv. II. 208.— Bras.
sylvestris, Bartram, ex Le Conte, in Rep. Commiss. Patents (Agric.), 1857 (1858), 229 = V. Labrusca.
sylvestris, Blume, Bijdr. 194 = flexuosa.
sylvestris, C. C. Gmel. Fl. Bad. i. 543 = vinifera.
taurina, Walt. Fl. Carol. 242 = V. Labrusca.
tenuicaulis, *Baker, in Oliver, Fl. Trop. Afr.* i. 404.— Afr. trop.
tenuifolia, Le Conte, in Proc. Acad. Sc. Philad. 1852–53 (1854) 271 = V. Labrusca.
tenuifolia, Wight & Arn. Prod. 129 = japonica.
ternata, *Baker, in Mart. Fl. Bras.* xiv. II. 216.—Bras.
ternata, Roth, ex Roem. & Schult. Syst. v. 319 = tomentosa.
tetragyna, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 78.— Ins. Celebes.
Teysmanni, *Miq. l. c.* 82.—Java.
Thomsoni, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 657. —Reg. Himal.
Thonningii, *Baker, in Oliver, Fl. Trop. Afr.* i. 407.— Afr. trop.
Thunbergii, Regel, Gartenfl. (1864) 34. t. 424 = vinifera.
Thunbergii, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1843) 198 = V. Labrusca.
thyrsiflora, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 88.— Sumatra.
tiliacea, *Hemsl. Biol. Centr. Am. Bot.* i. 204.—Mexic.
tiliaefolia, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 320 = caribaea.
tomentosa, *Heyne, ex Roth, Nov. Pl. Sp.* 157.—Ind. or.
trichophora, Wall. Cat. n. 6039 = Gynostemma pedatum.
tricuspidata, Lynch, in Journ. Linn. Soc. xvi. (1879) 308 = inconstans.
tricuspis, *Baker, in Mart. Fl. Bras.* xiv. II. 209.— Bras.
trifida, Roth, Nov. Pl. Sp. 157 = tomentosa.
trifolia, *Linn. Sp. Pl.* 203.—As. et Austral. trop.
trifoliata, *Baker, in Mart. Fl. Bras.* xiv. II. 212.— Am. trop.
trifoliata, Thunb. Fl. Javan. 11 (Quid?).—Java.
triloba, Roth, Nov. Pl. Sp. 157 = tomentosa.
triphylla, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = Vitex trifolia.
triternata, *Baker, in Journ. Linn. Soc.* xx. (1883) 123. —Madag.
truncata, Blume, Bijdr. 195 = flexuosa.
truncata, Wall. Cat. n. 6032 = tenuifolia.
tuberculata, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 656. —Burma; Java.
tuberculata, Wall. Cat. n. 6014 = rumicisperma.
tuberosa, *Hemsl. Biol. Centr. Am. Bot.* i. 204.— Mexic.
turbinata, Rafin. Med. Fl. ii. 140 = vinifera?
Tweedieana, *Baker, in Mart. Fl. Bras.* xiv. II. 214.— Reg. Argent.

VITIS:—
ulmifolia, *Baker, in Mart. Fl. Bras.* xiv. II. 213.—Peruv.
ursina, Rafin. Med. Fl. ii. 124 = aestivalis.
uvifera, *Baker, in Oliver, Fl. Trop. Afr.* i. 391.—Afr. trop.
Veitchi, Lynch, in Journ. Linn. Soc. xvii. (1879) 308 = inconstans.
verrucaria, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = nodosa.
verrucosa, Rafin. Med. Fl. ii. 133 = rotundifolia.
Vicaryana, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 204.—Reg. Himal.
vinifera, *Linn. Sp. Pl.* 202.—Oriens.; Ind. bor. occ.
vinifera, Marsh. Arbust. Am. 166 = aestivalis.
virginiana, Hort. ex Poir. Encyc. viii. 608 = riparia.
virginiana, Poir. l. c. 808 = aestivalis.
vitiensis, *Seem. Mission Viti*, 434; *Fl. Vit.* 44.—Ins. Fiji.
Vogelii, *Baker, in Oliver, Fl. Trop. Afr.* i. 409.—Afr. trop.
vulpina, Bartram, ex Le Conte, in Rep. Commiss. Patents (Agric.). 1857 (1858) 229 = V. Labrusca.
vulpina, Jacq. Hort. Schoenb. iv. 13. t. 425 = aestivalis.
vulpina, *Linn. Sp. Pl.* 203 = riparia, rotundifolia.
vulpina, Muhl. Cat. 26 = cordifolia.
Wallichii, DC. Prod. i. 634 = flexuosa.
Wallichii, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 302 = pallida.
Warmingii, *Baker, in Mart. Fl. Bras.* xiv. II. 201.—Bras.
Welwitschii, *Baker, in Oliver, Fl. Trop. Afr.* i. 363.—Afr. trop.
Wightiana, Wall. Cat. n. 6003 = repanda.
zeylanica, Russ. ex Wall. Cat. sub n. 5993 D = latifolia.

VITIS-IDAEA, Tourn. ex Moench, Meth. 47 (1794) = **Vaccinium**, Linn. (Vaccin.).
alba, Moench, l. c. = Lonicera ciliata.
Kablikiana, Opiz, Seznam, 104 = Vacc. *intermedium*.
Myrtillus, Moench, Meth. 47 = Vacc. Myrtillus.
punctata, Moench, l. c. = Vacc. Vitis-Idaea.
punctifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 406 = Vacc. Vitis-Idaea.

VITMANIA, Turra, ex Cav. Ic. iii. 53, in obs. (1794) = **Oxybaphus**, Vahl (Nyctag.).
viscosa, Turra, ex Steud. Nom. ed. I. 140, in syn.; 889 = O. viscosus.

VITMANNIA, Vahl, Symb. Bot. iii. 51. t. 60 (1794) = **Samadera**, Gaertn. (Simarub.).
elliptica, Vahl, l. c. = S. indica.
lucida, Steud. Nom. ed. II. ii. 779 = S. lucida.
polyandra, Steud. l. c. = Brownlowia lanceolata.

VITMANNIA, Wight & Arn. Prod. 166 (1834) = **Noltea**, Reichb. (Simarub.).
africana, Wight & Arn. l. c. = N. africana.

VITTADENIA, Steud. Nom. ed. II. ii. 779 (1841), sphalm. = seq.

VITTADINIA, A. Rich. Ess. Fl. Nouv.-Zél. 250 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 281.
EURYBIOPSIS, DC. Prod. v. 260 (1836).
MICROGYNE, Less. Syn. Comp. 190 (1832).
arenaria, *A. Gray, in Proc. Am. Acad.* v. (1861) 120.—Ins. Sandvic.
australis, *A. Rich. Ess. Fl. Nouv.-Zél.* 251.—Austral.; N. Zel.
brachycomoides, *F. Muell. Fragm.* v. 86.—Austral.
Chamissonis, *A. Gray, in Proc. Am. Acad.* v. (1861) 119.—Ins. Sandvic.
consanguinea, *A. Gray, l. c.* 120.—Ins. Sandvic.
conyzoides, *A. Gray, l. c.* 120.—Ins. Sandvic.
cuneata, DC. Prod. v. 281 = australis.
dentata, DC. l. c. = australis?
hispidula, F. Muell. ex A. Gray, in Proc. Am. Acad. v. (1861) 118 = scabra.
humilis, *A. Gray, l. c.*—Ins. Sandvic.
lobata, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 959 = australis?
macrorhiza, *A. Gray, in Proc. Am. Acad.* v. (1861) 118.—Austral.
multifida, *Griseb. in Goett. Abh.* xix. (1874) 171.—Reg. Argent.
Remyi, *A. Gray, in Proc. Am. Acad.* v. (1861) 119.—Ins. Sandvic.

VITTADINIA:—
scabra, *DC. Prod.* v. 281.—Austral.
scabra, Hook. f. Fl. Tasman. i. 181 = australis.
spathulata, *F. Muell. ex Sond. in Linnaea*, xxv. (1852) 457.—Austral.
tenerrima, *A. Gray, in Proc. Am. Acad.* v. (1861) 119.—Ins. Sandvic.
trifurcata, *Griseb. in Goett. Abh.* xxiv. (1879) 178.—Reg. Argent.
triloba, DC. Prod. v. 281 = australis.

VITTMANNIA, Turra, ex Endl. Gen. 311 (1837) = **Oxybaphus**, Vahl (Nyctag.).

VIVIANIA, Cav. in Anal. Cienc. Nat. vii. (1804) 211. t. 49. *GERANIACEAE*, Benth. & Hook. f. i. 275.
CAESAREA, Cambess. in Mém. Mus. Par. xviii. (1829) 373. t. 18.
CISSAROBRYON, Kunze, ex Poepp. Fragm. Syn. Pl. Chile, 29 (1833).
LINOSTIGMA, Klotzsch, in Linnaea, x. (1836) 438.
MACRAEA, Lindl. in Brand. Journ. xxv. (1828) 104.
XEROPETALON, Hook. Bot. Misc. i. (1830) 174.
aristulatum, *Phil. in Anal. Univ. Chil.* (1872) 687.—Chili.
australis, *Phil. l. c.* 683.—Chili.
brevipedunculata, *Phil. in Linnaea*, xxxiii. (1864–65) 33.—Chili.
calycina, Griseb. in Goett. Abh. xix. 104 = Wendtia calycina.
crenata, *G. Don, ex Hook. Bot. Misc.* iii. (1833) 149.—Chili.
grandiflora, G. Don, ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 657, sphalm. = seq.
grandifolia, *Hook. & Arn. Bot. Misc.* iii. (1833) 149.—Chili.
marifolia, *Cav. in Anal. Cienc. Nat.* vii. (1804) 212.—Chili.
ovata, *Phil. in Linnaea*, xxviii. (1856) 677.—Chili.
parviflora, G. Don, ex W. Baxt. in Loud. Hort. Brit. Suppl. iii. 657, sphalm. = seq.
parvifolia, *Klotzsch, in Linnaea*, x. (1836) 435.—Chili.
petiolata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 149.—Chili.
rosea, Klotzsch, in Linnaea, x. (1836) 435 = marifolia.
spinescens, *Presl, Symb. Bot.* ii. 7.—Chili.
tenuicaulis, *Barn. in C. Gay, Fl. Chil.* i. 402. t. 12.—Chili.

VIVIANIA, Colla, in Mém. Soc. Linn. Par. iv. (1826) 25 = **Billiottia**, DC. (Rubiac.).
psychotrioides, Colla, l. c. = B. psychotrioides.

VIVIANIA, Rafin. Specch. i. 117 (1814); ex DC. Prod. iv. 458 = **Guettarda**, Linn. (Rubiac.).
domingensis, Rafin. l. c. = G. coriacea?

VIVIANIA, Willd. ex Less. in Linnaea, iv. (1829) 318; DC. Prod. v. 95 (1836) = **Liabum**, Adans. (Compos.).
bicolor, Willd. ex Less. l. c. = L. melastomoides.

VLAMINGIA, De Vriese, in Lehm. Pl. Preiss. i. 398 (1845) = **Ionidium**, Vent. (Violar.).
australasica, De Vriese, l. c. 399 = I. calycinum.

VLECKIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Lophanthus**, Benth. (1829) (Labiat.).
anisata, Rafin. Fl. Tellur. iii. 89 = L. anisatus.
bracteata, Rafin. New Fl. Am. iv. 94 = L. anisatus?
chinensis, Rafin. Fl. Tellur. iii. 89.—L. chinensis.
multiflora, Rafin. l. c. = Nepeta lavandulacea?
nepetoides, Rafin. in Med. Repos. N. York, v. (1808) 352 = L. nepetoides.
scrophularifolia, Rafin. Fl. Tellur. iii. 89 = L. scrophulariaefolius.
urticaefolia, Rafin. l. c. = L. urticifolius.

VOACANGA, Thou. Gen. Nov. Madag. 10 (1806). *APOCYNACEAE*, [Benth. & Hook. f. ii. 707].
ANNULARIA, Hochst. in Flora, xxiv. (1841) 670.
CYCLOSTIGMA, Hochst. ex Endl. Gen. Suppl. ii. 56 (1842); et in Flora, xxvii. (1844) 828.
DICRUS, Reinw. in Blume, Cat. Hort. Buitenz. 47 (1823).
ORCHIPEDA, Blume, Bijdr. 1026 (1826).
PIPTOLAENA, Harv. in Hook. Lond. Journ. Bot. i. 25 (1842).
POOTIA, Miq. in Versl. en Med. Kon. Akad. Wetensch. vi. (1857) 192; Fl. Ind. Bat. ii. 417 ('1856').

VOACANGA:—
Cumingiana, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 313.—Ins. Philipp.
Dregei, *E. Mey. Comm. Pl. Afr. Austr.* 189.—Afr. austr.
foetida, *Rolfe, in Journ. Bot.* xxi. (1883) 202.—Java.
grandifolia, *Rolfe, l. c.*—Java.
Thouarsii, *Roem. & Schult. Syst.* iv. 439.—Madag.

VOANDZEIA, Thou. Gen. Nov. Madag. 23 (1806). *LEGUMINOSAE*, Benth. & Hook. f. i. 539.
CRYPTOLOBUS, Spreng. Syst. iii. 152 (1826).
GEOLOBUS, Rafin. New Fl. Am. i. 81 (1836).
subterranea, *Thou. Gen. Nov. Madag.* 23.—Afr. trop. et austr.

VOCHISIA, Juss. Gen. 424 (1789). *VOCHYSIACEAE*, Benth. & Hook. f. i. 976.
CUCULLARIA, Schreb. Gen. 6 (1789).
SALMONIA, Scop. Introd. 209 (1777).
STRUKERIA, Vell. Fl. Flum. i. t. 20 (1827).
VOCHY, Aubl. Pl. Gui. i. 18 (1775), nomen prius.
VOCHYA, Vell. ex Vand. Fl. Lusit. I. t. 1. f. 1 (1788); et in Roem. Script. 69. t. 6 (1796).
VOCHYSIA, Poir. Encyc. viii. 681 (1808).
acuminata, *Bong. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iii. II. (1839) 5 (sp. dub.).—Am. trop.
alpestris, *Mart. Nov. Gen. et Sp.* i. 145. t. 87.—Bras.
arcuata, *Garcke, in Linnaea*, xxii. (1849) 58.—Guian.
bifalcata, *Warm. in Mart. Fl. Bras.* xiii. II. 84.—Bras.
calophylla, *Spruce, ex Warm. l. c.* 98.—Bras.
cayennensis, *Warm. l. c.* 80.—Guian.
cinnamomea, *Pohl, Pl. Bras. Ic.* ii. 29.—Bras.
citrifolia, *Poir. Encyc. Suppl.* v. 491.—Bras.
costata, *Warm. in Mart. Fl. Bras.* xiii. II. 100.—Guian.
crassifolia, *Warm. l. c.* 77.—Guian.
cuiabensis, *Liais, Brésil*, 601.—Bras.
cuneata, Pohl, Pl. Bras. Ic. ii. 26 = thyrsoidea.
curvata, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1099, nomen (Quid?).—Guian.
dasyantha, *Warm. in Mart. Fl. Bras.* xiii. II. 95.—Bras.
densiflora, *Spruce, ex Warm. l. c.* 101.—Bras.
discolor, *Warm. l. c.* 81.—Bras.
divergens, *Pohl, Pl. Bras. Ic.* ii. 19.—Bras.
elliptica, *Mart. Nov. Gen. et Sp.* i. 141. t. 84.—Bras.
elongata, *Pohl, Pl. Bras. Ic.* ii. 25.—Bras.
emarginata, *Vahl, ex Poir. Encyc.* viii. 682.—Bras.
excelsa, A. Dietr. Sp. Pl. i. 103 = guianensis.
fastigiata, *Warm. in Mart. Fl. Bras.* xiii. II. 91.—Bras.
ferruginea, *Mart. Nov. Gen. et Sp.* i. 151. t. 92.—Bras.
ferruginea, Pohl, Pl. Bras. Ic. ii. 31 = pyramidalis.
floribunda, *Mart. Nov. Gen. et Sp.* i. 149. t. 91.—Bras.
Fontanesii, Zucc. ex Warm. in Mart. Fl. Bras. xiii. II. 104 = emarginata.
Gardneri, *Warm. l. c.* 70.—Bras.
glaberrima, *Warm. l. c.* 78.—Guian.
glabrescens, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1099, nomen (Quid?).—Guian.
Glazioviana, *Warm. in Mart. Fl. Bras.* xiii. II. 86.—Bras.
grandis, *Mart. Nov. Gen. et Sp.* i. 146. t. 88.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 18. t. 6 (Vochy guianensis).—Guian.
gummifera, *Mart. ex Warm. in Mart. Fl. Bras.* xiii. II. 82.—Bras.
Haenkeana, *Mart. Nov. Gen. et Sp.* i. 147. t. 89.—Am. trop.
herbacea, *Pohl, Pl. Bras. Ic.* ii. 27.—Bras.
laurifolia, *Warm. in Mart. Fl. Bras.* xiii. II. 96.—Bras.
lucida, *Presl, Rel. Haenk.* ii. 12.—Am. trop.
magnifica, *Warm. in Mart. Fl. Bras.* xiii. II. 85.—Bras.
maliformis, Klotzsch, ex Warm. l. c. 24 = Callisthene fasciculata.
micrantha, *Pohl, Pl. Bras. Ic.* ii. 20.—Bras.
oblongifolia, *Warm. in Mart. Fl. Bras.* xiii. II. 84.—Bras.
obscura, *Warm. l. c.* 78.—Bras.

VOCHISIA :—
opaca, Pohl, ex Warm. in Mart. Fl. Bras. xiii. II. 91, in obs. = tucanorum.
oppugnata, *Warm. l. c.* 87.—Bras.
parviflora, *Spruce, ex Warm. l. c.* 75.—Bras.
petraea, *Warm. l. c.* 63.—Bras.
pruinosa, *Pohl, Pl. Bras. Ic.* ii. 22.—Bras.
pumila, *Pohl, l. c.* 21.—Bras.
punctata, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 102.—Bras.
pygmaea, *Bong. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* iii. II. (1839) 7. t. 4.—Am. trop.
pyramidalis, *Mart. Nov. Gen. et Sp.* i. 148. t. 90.—Bras.
quadrangulata, *Warm. in Kjoeb. Vidensk. Meddel.* (1867) 39 ; *et in Mart. Fl. Bras.* xiii. II. 97.—Bras.
racemosa, *Poir. Encyc.* viii. 682.—Guian.
rectiflora, *Warm. in Mart. Fl. Bras.* xiii. II. 96.—Bras.
rotundifolia, *Mart. Nov. Gen. et Sp.* i. 140. t. 83.—Bras.
rufa, *Mart. l. c.* 144. t. 86.—Bras.
Schomburgkiana, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1013. nomen (Quid ?).—Guian.
Schomburgkii, *Warm. in Mart. Fl. Bras.* xiii. II. 78.—Guian.
Selloi, *Warm. l. c.* 86.—Bras.
sericea, Pohl, Pl. Bras. Ic. ii. 28 = rufa.
sessilifolia, *Warm. in Mart. Fl. Bras.* xiii. II. 67.—Bras.
speciosa, *Warm. l. c.* 79.—Guian.
splendens, *Spruce, ex Warm. l. c.* 101.—Bras.
Sprucei, *Warm. l. c.* 99.—Peruv.
tetraphylla, *DC. Prod.* iii. 27.—Guian.
thyrsoidea, *Pohl, Pl. Bras. Ic.* ii. 24.—Bras.
tomentosa, DC. Prod. iii. 26 = ferruginea.
tucanorum, *Mart. Nov. Gen. et Sp.* i. 142. t. 85.—Bras.
venulosa, *Warm. in Mart. Fl. Bras.* xiii. II. 74.—Bras.
verticillata, *A. Dietr. Syn. Pl.* i. 22.—Bras.
vismiaefolia, *Spruce, ex Warm. in Mart. Fl. Bras.* xiii. II. 99.—Peruv.

VOCHY, Aubl. Pl. Gui. i. 18 (1775) = praec.

VOCHYA, Vell. ex Vand. Fl. Lusit. 1. t. 1. f. i (1788) ; et in Roem. Script. 69. t. 6 (1796) = praec.

VOCHYSIA, Poir. Encyc. viii. 681 (1808) = praec.

VOELCKERIA, Klotzsch & Karst. in Endl. Gen. Suppl. iv. 66 (1850) = **Ternstroemia**, Linn.

VOGELIA, J. F. Gmel. Syst. 107 (1791) = **Burmannia**, Linn.
capitata, J. F. Gmel. l. c. = B. capitata.

VOGELIA, Lam. Illustr. ii. 147. t. 149 (1792). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 628.
africana, *Lam. l. c.* 149.—Afr. austr.
arabica, Boiss. in DC. Prod. xii. 696 = indica.
indica, *Gibs. ex Wight, in Calc. Journ. Nat. Hist.* vii. (1847) 17 ; *Ic.* t. 1075.—Ind. or. ; Arab.
pendula, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 76.—Ins. Socotra.
perfoliata, Stocks, ex Wight, in Calc. Journ. Nat. Hist. vii. (1847) 17 = indica.

VOGELIA, Medic. Pflanzeng. 32. t. 1. f. 6 (1792) = **Neslia**, Desv. (Crucif.).
paniculata, Hornem. Hort. Hafn. ii. 594 = N. paniculata.
sagittata, Medic. Pflanzeng. 32 ; Bernh. Syst. Verz. Erf. 190 = N. paniculata.

VOGLERA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 480 (1800) = **Genista**, Linn. (Legumin.).
germanica, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = G. germanica.
hispanica, Fourr. l. c. = G. hispanica.
spinosa, Gaertn. Mey. & Scherb. Fl. Wett. ii. 500 = G. germanica.

VOHIRIA, Juss. Gen. 141 (1789) = **Voyria**, Aubl. (Gentian.).

VOIGTIA, Klotzsch, in Hayne, Arzn. Gew. xiv. adnot. ad t. 15 (1846) = **Bathysa**, Presl (Rubiac.).
australis, Klotzsch, l. c.—Bras.

VOIGTIA, Roth, in Roem. & Usteri, Mag. iv. St. x. (1790) 17 = **Andryala**, Linn. (Compos.).
tomentosa, Roth, l. c. = A. Rothia.

VOIGTIA, Spreng. Syst. iii. 673 (1826) = **Barnadesia**, Mutis (Compos.).
laurifolia, Spreng. l. c. (= *Fulcaldea laurifolia*).—Peruv.

VOLKAMERIA, P. Br. Hist. Jamaic. 214. t. 21. f. 1 (1756) = **Clethra**, Gronov. (Ericac.).

VOLKAMERIA, Linn. Syst. ed. I (1735) ; Gen. ed. I. 347 (1737) = **Clerodendron**, Linn. (Verben.).
Acerbiana, Vis. Pl. Aegypt. 23. t. 4. f. 1 = C. Acerbiana.
aculeata, Linn. Sp. Pl. 637 = C. aculeatum.
alternifolia, Burm. f. Fl. Ind. 137 = C. calamitosum.
angulata, Lour. Fl. Cochinch. 389 = C. paniculatum.
angustifolia, Andr. Bot. Rep. t. 554 = C. heterophyllum.
angustifolia, Poir. Encyc. viii. 688 = C. angustifolium.
Buchanani, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 60 = C. foetidum.
buxifolia, Willd. Enum. Hort. Berol. 658 = C. buxifolium.
capensis, Burm. f. Fl. Cap. Prod. 17 = Capparis Volkameriae.
capitata, Willd. Sp. Pl. iii. 384 = C. capitatum ?
Casopanguil, Blanco, Fl. Filip. ed. II. 356 = C. intermedium.
coccinea, Herb. Amat. ex Schau. in DC. Prod. xi. 669 = C. squamatum.
Commersonii, Poir. Encyc. viii. 688 = C. Commersonii.
cordifolia, Hochst. in Flora, xxv. (1842) 227, in nota = C. cordifolium.
dentata, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 61 = C. squamatum.
diversifolia, Vahl, Symb. Bot. ii. 75 = C. paniculatum.
farinosa, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 64 = C. venosum.
foetida, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 103 = C. foetidum.
fragrans, Vent. Jard. Malm. 70 = C. fragrans.
grandiflora, Blanco, Fl. Filip. ed. I. 512 = C. macrostegium.
herbacea, Roxb. Hort. Beng. 46 = C. serratum.
heterophylla, Poir. Encyc. viii. 687 = C. heterophyllum.
inermis, Blanco, Fl. Filip. ed. I. 511 = C. intermedium.
inermis, Linn. Sp. Pl. 637 = C. nereifolium.
inermis, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 239 = Geniostoma rupestre.
infortunata, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 59 = C. infortunatum.
japonica, Thunb. Fl. Jap. 255 = C. fragrans.
Kaempferi, Jacq. Ic. Pl. Rar. iii. t. 500 = C. squamatum.
laurifolia, Hort. ex Steud. Nom. ed. I. 890 = C. viscosum.
ligustrina, Jacq. Coll. Suppl. 118. t. 5. f. 1 = C. ligustrinum.
Madaeera, Roxb. Hort. Beng. 46, nomen = C. Madaeera.
multiflora, Burm. f. Fl. Ind. 137 = C. phlomoides.
nereifolia, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 64 = C. nereifolium.
obovata, Roxb. ll. cc. [95] ; 62 = C. obovatum.
odorata, [Buch.-Ham.] Roxb. Hort. Beng. 46 = Caryopteris Wallichiana.
odoratissima, Fisch. ex Steud. Nom. ed. I. 890.—Hab. ?
Petasites, Lour. Fl. Cochinch. 388.—China.
pumila, Lour. l. c. = C. pumilum.
scandens, Linn. f. Suppl. 292.—Zeylan.
serrata, Linn. Mant. i. 90 = C. serratum.
spinosa, Juss. in Ann. Mus. Par. fasc. 37. p. 76 ; ex Pers. Syn. ii. 144.—Peruv.
tomentosa, Vent. Jard. Malm. t. 84 = C. tomentosum.
urticifolia, Roxb. Hort. Beng. 46 ; Fl. Ind. iii. 61 = C. squamatum.
verticillata, Ruiz & Pav. ex Walp. Rep. iv. 73.—Peruv.

VOLKMANNIA, Jacq. Hort. Schoenb. iii. 48. t. 338 (1798) = **Clerodendron**, Linn. (Verben.).
japonica, Jacq. l. c. = C. fragrans.

VOLUTARELLA, Cass. in Dict. Sc. Nat. xliv. 36 (1826). *COMPOSITAE*, Benth. & Hook. f. ii. 476.
AMBERBOA, Less. Syn. Comp. 8 (1832).
AMBERBOI, Adans. Fam. ii. 117 (1763), nomen prius.
CYANASTRUM, Cass. in Dict. Sc. Nat. xliv. 39 (1826).
CYANOPSIS, Cass. in Bull. Soc. Philom. (1817) 200.
LACELLIA, Viv. Fl. Lib. Spec. 58. t. 22. f. 2 (1824).
OLIGOCHAETA, C. Koch, in Linnaea, xvii. (1843) 42.
PARAPHYSIS, Steud. Nom. ed. II. ii. 267 (1841).
PHAENOPAPPUS, DC. ex Steud. Nom. ed. II. ii. 313 (1841).
STEPHANOCHILUS, Coss. & Dur. ex Benth. & Hook. f, Gen. ii. 477 (1873).
VOLUTARIA, Cass. in Bull. Soc. Philom. (1816) 200.
abyssinica, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 198.—Abyss.
bicolor, *Cass. in Dict. Sc. Nat.* l. 250.—Afr. bor.
crupinoides, Ball, in Journ. Linn. Soc. xvi. (1878) 525 = bicolor.
divaricata, *Benth. & Hook. f. Gen.* ii. 476.—Ind. or. ; Afghan.
erucifolia, *Cass. in Dict. Sc. Nat.* lviii. 453.—Hab. ?
leucantha, *Ball, in Journ. Linn. Soc.* xvi. (1878) 526.—Marocc.
Lippii, *Cass. in Dict. Sc. Nat.* xliv. 39.—Am. austr. ; Ins. Canar.
muricata, *Benth. & Hook. f. Gen.* ii. 476 ; *Ball, in Journ. Linn. Soc.* xvi. (1878) 526.—Marocc.
omphalodes, *Benth. & Hook. f. l. c.* 477.—Alger.

VOLUTARIA, Cass. in Bull. Soc. Philom. (1816) 200 = **Volutarella**, Cass. (1826) (Compos.).
Lippii, Cass. l. c. = Volutarella Lippii.

VOLUTELLA, Forsk. Fl. Aegypt. Arab. 84 (1775) = = **Cassytha**, Linn. (Laurin.).
aphylla, Forsk. l. c. = C. filiformis.

VOLVULUS, Medic. in Staatsw. Vorles. Churpf. Phys. Oek. Ges. i. (1791) 202 = **Calystegia**, R. Br. (Convolv.).
sepium, Medic. Phil. Bot. ii. 42 = C. sepium.

VORMIA, Adans. Fam. ii. 284 (1763) = **Selago**, Linn.

VOSACAN, Adans. Fam. ii. 130 (1763) = **Helianthus**, Linn. (Compos.).

VOSAKAN, Adans. Fam. ii. 619 (Index) (1763) = praec.

VOSSIA, Wall. & Griff. in Journ. As. Soc. Beng. v. (1836) 572. t. 23. *GRAMINEAE*, Benth. & Hook. f. iii. 1131.
EREMOCHLOA, Buese, in Miq. Pl. Jungh. 357 (1854).
cuspidata, *Griff. Ic. Pl. Asiat.* t. 153.—Ind. or.
hordeoides, *Munro, in Harv. Gen. S. Afr. Pl.* ed. II. 442.—Afr. austr.
procera, *Wall. & Griff. in Journ. As. Soc. Beng.* v. (1836) 573. t. 23.—Ind. or. ; Afr. trop.
speciosa, *Benth. in Journ. Linn. Soc.* xix. (1881) 70.—Ind. or.

VOTOMITA, Aubl. Pl. Gui. i. 90. t. 35 (1775). Genus dubium *RUBIACEARUM*.—Cf. Benth. & Hook. f. ii. 29.
GLOSSOCOMA, Endl. Gen. 799, in syn. (1839).
GUILLEMINIA, Neck. Elem. ii. 132 (1790).
guianensis, *Aubl. Pl. Gui.* i. 90. t. 35.—Guian.

VOUACAPOUA, Aubl. Pl. Gui. Suppl. 9. t. 373 (1775) = **Andira**, Lam. (1783) (Legumin.).
americana, Aubl. l. c. = A. excelsa.

VOUAPA, Aubl. Pl. Gui. i. 25. t. 7 (1775) = **Macrolobium**, Schreb. (1789) (Legumin.).
acaciaefolia, Baill. Hist. Pl. ii. 109 = M. acaciaefolium.
angustifolia, Benth. in Hook. Kew Journ. ii. (1850) 239 = M. chrysostachyum.
bifolia, Aubl. Pl. Gui. i. 25. t. 7 = M. Vuapa.
chrysostachya, Miq. Stirp. Surin. Sel. 11 = M. chrysostachyum.
crassifolia, Baill. Adansonia, vi. (1865) 179 (= *Anthonota crassifolia*).—Afr. trop.
demonstrans, Baill. l. c. 180. t. 3 = M. demonstrans.
elegans, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 40 = M. Vuapa.
explicans, Baill. Adansonia, vi. (1865) 181 = M. Heudelotii.
macrophylla, Baill. l. c. 178 = M. Palisoti.

VOUAPA:—
phaseolaecarpa, Hayne, Arzn. Gew. xi. t. 20 = Cynometra racemosa.
Simira, Aubl. Pl. Gui. i. 27. t. 8 = Macrolobium Simira.
staminea, DC. Prod. ii. 511 = M. Vuapa.
violacea, [Lam.] Poir. Encyc. viii. 700 = M. Simira.

VOUARANA, Aubl. Pl. Gui. Suppl. 12. t. 374 (1775) = **Cupania**, Plum. (Sapind.).
guianensis, Aubl. l. c. = C. Vouarana.

VOUAY, Aubl. Pl. Gui. ii. App. 99 (1775) = **Geonoma**, Willd. (Palm.).

VOUCAPOUA, Steud. Nom. ed. II. ii. 780 (1841), sphalm. = VOUACAPOUA, Aubl. = **Andira**, Lam. (Legumin.).

VOYARA, Aubl. Pl. Gui. Suppl. 26 (1775) (Quid?).
montana, Aubl. l. c.—Guian.

VOYRA, Reichb. Consp. 133 (1828) = seq.

VOYRIA, Aubl. Pl. Gui. i. 208. t. 83 (1775). *GENTIANEAE*, Benth. & Hook. f. ii. 806.
CIMINALIS, Rafin. Fl. Tellur. iii. 19 (1836).
DISADENA, Miq. Stirp. Surin. Sel. 150 (1850).
HUMBOLDTIA, Neck. Elem. ii. 16 (1790).
LEIANTHOSTEMON, Miq. Stirp. Surin. Sel. 147 (1850).
LEIPHAIMOS, Cham. & Schlecht. in Linnaea, vi. (1831) 387.
LITA, Schreb. Gen. 795 (1791).
PNEUMONANTHOPSIS, Miq. Stirp. Surin. Sel. 150 (1850).
acuminata, *Benth. in Hook. Journ. Bot.* ii. (1840) 46.—Guian.
angustiflora, Steud. in Flora, xxvi. (1843) 765 = aurantiaca.
angustiloba, *Spruce, ex Benth. in Hook. Kew Journ.* vi. (1854) 197.—Bras.
aphylla, *Pers. Syn.* i. 284.—Am. trop.
araguensis, *Karst. in Linnaea*, xxviii. (1856) 415.—Venezuela.
aurantiaca, *Splitg. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 135.—Guian.
brachyloba, *Griseb. Cat. Pl. Cub.* 181.—Cuba.
breviflora, *Lam. Illustr.* i. 491.—Guian.
caerulea, *Aubl. Pl. Gui.* i. 211. t. 83. f. 2.—Guian.
calycina, *Splitg. ex A. DC. in DC. Prod.* ix. 562, in obs.—Guian.
chionea, *Benth. in Hook. Kew Journ.* vi. (1854) 197.—Bras.
clavata, *Splitg. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 133.—Guian.
coerulea, Auct. = caerulea.
corymbosa, *Splitg. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 136.—Guian.
disadenantha, *Griseb. Cat. Pl. Cub.* 181.—Cuba.
flavescens, *Griseb. in DC. Prod.* ix. 85.—Bras.
leucantha, *Miq. in Tijdschr. Wetensch.* ii. (1849) 124.—Guian.
mexicana, *Griseb. Gen. et Sp. Gent.* 208.—Mexic.
nivea, *Miq. in Linnaea*, xix. (1847) 140.—Guian.
nivea, Spruce, ex Prog. in Mart. Fl. Bras. vi. 1. 223 = chionea.
nuda, *Splitg. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 134.—Guian.
obconica, *Prog. in Mart. Fl. Bras.* vi. 1. 221.—Bras.
pallida, Garcke, in Linnaea, xxii. (1849) 65 = aphylla.
parviflora, *Miq. in Tijdschr. Wetensch.* ii. (1849) 123.—Guian.
rosea, *Aubl. Pl. Gui.* i. 209. t. 83. f. 1.—Guian.
simplex, *Griseb. in Seem. Bot. Voy. Herald*, 170.—Panama.
spathacea, *Lam. Illustr.* i. 491; *Poir. Encyc.* viii. 771.—Guian.
spathacea, Splitg. in Hoev. & De Vriese, Tijdschr. vii. (1840) 137 = calycina.
Spruceana, *Benth. in Hook. Kew Journ.* vi. (1854) 197.—Bras.
sulphurea, *Prog. in Mart. Fl. Bras.* vi. 1. 221.—Bras.
tenella, *Guild. ex Hook. Bot. Misc.* i. (1830) 47.—Ins. S. Vincent.
tenera, Hook. l. c. t. 25 B, sphalm. = praec.
tenuiflora, *Griseb. in DC. Prod.* ix. 85.—Guian.

VOYRIA:—
trinitensis, *Griseb. Fl. Brit. W. Ind.* 425.—Ins. Trinit.
uniflora, Pers. Syn. i. 284 = aphylla.

VOYRIELLA, Miq. Stirp. Surin. Sel. 146 (1850). *GENTIANEAE*, Benth. & Hook. f. ii. 806.
parviflora, *Miq. l. c.* 147.—Guian.

VRENA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 27, err. typ. = **Urena**, Dill. (Malvac.).

VRIESEA, Beer, Bromel. 91 (1857) = VRIESIA, Lindl. = **Tillandsia**, Linn. (Bromel.).

VRIESEA, Hassk. in Flora, xxv. (1842) II. Beibl. 27 = **Vandellia**, Linn. (Scrophul.).
monticola, Hassk. l. c. = Ilysanthes refracta.
nervosa, Hassk. l. c. 28 = Vand. nervosa.
sphaerocarpa, Hassk. l. c. 27 = Vand. sphaerocarpa.

VRIESIA, Lindl. Bot. Reg. (1843) t. 10 = **Tillandsia**, Linn. (Bromel.).
aloëfolia, Beer, Bromel. 95 = T. flexuosa.
amethystina, E. Morr. Belg. Hortic. xxxiv. (1884) 330. t. 15–16.—Bras.
anceps, Lem. in Fl. des Serres, v. (1849) sub t. 432 = T. anceps.
Barilleti, E. Morr. Belg. Hortic. xxxiii. (1883) 33. t. 3.—Ecuador.
bellula, Linden, Cat. n. 109 (1883) 7; et in Gard. Chron. (1884) I. 140 = heliconioides.
Billbergiae, Lem. Illustr. Hortic. xvi. (1869) Misc. 91.—Mexic.
billbergioides, E. Morr. ex Antoine, Bromel. 17.—Bras.
bituminosa, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 347 = T. platynema.
Botteri, Hort. ex Regel, Gartenfl. xxxiii. (1884) 293 = T. xiphostachys.
brachystachys, Regel, Gartenfl. xv. (1866) 258. t. 518 = *carinata*.
bracteosa, Beer, Bromel. 263 = T. bracteosa.
carinata, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 349.—Bras.
chrysostachis, E. Morr. Belg. Hortic. xxxi. (1881) 87.—Peruv.
conferta, Gaudich. Bot. Voy. Bonite, t. 65 = T. ensiformis.
corallina, Regel, Gartenfl. xix. (1870) 354. t. 671 = T. corallina.
Duvaliana, E. Morr. Belg. Hortic. xxxiv. (1884) 105. tt. 7, 8.—Bras.
ensiformis, Beer, Bromel. 92 = T. ensiformis.
Falkenbergii, Hort. Bull.—Cf. Gard. Chron. (1880) I. 759 = T. heliconioides.
fenestralis, Linden & André, in Illustr. Hortic. xxii. (1875) 124. t. 215.—Bras.
geniculata, Wawra, Bot. Ergeb. Reise Bras. 156. t. 25 = T. Regina.
gigantea, Gaudich. Bot. Voy. Bonite, t. 70.—Bras.
gigantea, Lem. Illustr. Hortic. xiv. (1867) t. 516 = T. Regina.
gladioliflora, Antoine, in Wien. Illustr. Garten-Zeit. v. (1880) 97. t. 1; et Bromel. 23. t. 15 = T. gladioliflora.
glaucophylla, Hook. Bot. Mag. t. 4415.—N. Granat.
Glaziouana, Lem. Illustr. Hortic. xiv. (1867) Misc. 43 = T. Regina.
glutinosa, Lindl. in Gard. Chron. (1856) 388 = T. glutinosa.
gracilis, Gaudich. Bot. Voy. Bonite, t. 67 = T. procera.
guttata, Linden & André, in Illustr. Hortic. xxii. (1875) 43. t. 200.—Bras.
heliconioides, Lindl. Bot. Reg. (1843) t. 10, in obs.; et ex E. Morr. in Illustr. Hortic. xxx. (1883) t. 490 = T. heliconioides.
hieroglyphica, E. Morr. Illustr. Hortic. xxxi. (1884) 41. t. 514.—Bras.
imbricata, Beer, Bromel. 94 = T. imbricata.
incurvata, Gaudich. Bot. Voy. Bonite, t. 68 = T. caespitosa.
incurvata, E. Morr. Belg. Hortic. xxxii. (1882) 52. t. 2 = *inflata*.
inflata, Wawra, Itin. Princ. Coburgi, i. 161.—Bras.
Itatiaiae, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 221.—Bras.
Jonghei, E. Morr. Belg. Hortic. xxviii. (1878) 257 = T. Jonghei.

VRIESIA:—
Krameri, E. Morr. Belg. Hortic. xxxiv. (1884) 369 = Tillandsia psittacina.
Lindeni, Lem. Illustr. Hortic. xvi. (1869) t. 610.—Peruv.
maculata, Beer, Bromel. 98 = T. maculata.
Malzinei, E. Morr. Belg. Hortic. xxiv. (1874) 313. t. 14 = T. Malzinei.
Morreni, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 219.—Bras.
Morreniana, Hort. ex E. Morr. Belg. Hortic. xxxii. (1882) 287 = T. psittacina.
musaica, Cogn. & March. Pl. Ornament. ii. (1874) t. 39.—N. Granat.
paraibica, Wawra, Itin. Princ. Coburgi, i. 160.—Bras.
Philippocoburgi, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 219.—Bras.
platynema, Gaudich. Bot. Voy. Bonite, t. 66 = T. platynema.
Platzmanni, E. Morr. Belg. Hortic. xxv. (1875) 349. t. 23.—Bras.
Princeps, E. Morr. ex Regel, Gartenfl. xxvi. (1877) 158.—Bras.
prodigiosa, Lem. Illustr. Hortic. xvi. (1869) Misc. 92.—Mexic.
psittacina, Lindl. Bot. Reg. (1843) t. 10 = T. psittacina.
ramosa, Beer, Bromel. 265 = T. utriculata.
recurvata, Gaudich. Bot. Voy. Bonite, t. 69.—Bras.
Regina, Beer, Bromel. 97 = T. Regina.
Rodigasiana, E. Morr. Illustr. Hortic. xxix. (1882) 171. t. 467.—Bras.
rubra, Beer, Bromel. 98 = T. rubra.
sanguinolenta, Cogn. & March. Pl. Ornament. ii. (1874) t. 52; et ex André, in Illustr. Hortic. xxii. (1875) 44.—N. Granat.
Saundersi, E. Morr. Belg. Hortic. xxv. (1875) 350, nomen.—Hab.?
scalaris, E. Morr. l. c. xxix. (1879) 301; xxx. (1880) 309. t. 15.—Bras.
simplex, Beer, Bromel. 97 = T. simplex.
speciosa, Hook. Bot. Mag. t. 4382 = T. splendens.
splendens, Lem. in Fl. des Serres, vi. (1850–51) 162 = T. splendens.
tenuifolia, Beer, Bromel. 96 = T. flexuosa.
tessellata, E. Morr. Belg. Hortic. xxxii. (1882) 381. t. 14–16 = T. tessellata.
utriculata, Regel, in Ind. Sem. Hort. Petrop. (1869) 34 = T. utriculata.
viminalis, E. Morr. Belg. Hortic. xxviii. (1878) 257. t. 14–15 = T. viminalis.
Warmingi, E. Morr. l. c. xxxiv. (1884) 260. t. 12–13.—Bras.
Wawranea, Antoine, Bromel. 1. t. 1, 2.—Bras.
Xiphion, Platzmann, ex Antoine, l. c. 25 = T. Jonghei.
xyphostachys, Hook. Bot. Mag. t. 5287 = T. xiphostachys.

VROLICKIA, Steud. Nom. ed. II. ii. 780 (1841), sphalm. = seq.

VROLIKIA, Spreng. Syst. iii. 149 (1826) = **Heteranthia**, Nees (Scrophul.).
polygaloides, Spreng. l. c. 157 = H. decipiens.

VRTICA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = **Urtica**, Tourn.

VRYDAGZENIA, Benth. & Hook. f. Gen. iii. 598 (1883), err. typ. = seq.

VRYDAGZYNEA, Blume, Orch. Archip. Ind. 71. tt. 17, 19, 20 (1858). *ORCHIDEAE*, Benth. & Hook. f. Gen. iii. 598.
albida, *Blume, l. c.* 75.—Malaya.
elongata, *Blume, l. c.* 74.—N. Guin.
gracilis, *Blume, l. c.* 72.—Malaya.
nuda, *Blume, l. c.* 74.—Java.
purpurea, *Blume, l. c.* 73.—Java.
sandwicensis, *Benth. & Hook. f. Gen.* iii. 599.—Ins. Sandvic.
uncinata, *Blume, Orch. Archip. Ind.* 76.—Java.
vitiensis, *Reichb. f. Otia Bot. Hamb.* 51.—Ins. Fiji.

VULNERARIA, Tourn. ex Hall. Enum. Stirp. Helv. ii. 569 (1742) = **Anthyllis**, Riv. (Legumin.).
Anthyllis, Scop. Fl. Carn. ed. II. ii. 55 = A. Vulneraria.
argentea, Lam. Fl. Fr. ii. 651 = A. Barba-Jovis.

VULNERARIA :—
Barba-Jovis, Link, Handb. ii. 194 = Anthyllis Barba-Jovis.
cornicina, Link, l. c. 195 = A. cornicina.
Dillenii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 359 = A. Vulneraria.
hamosa, Link, Handb. ii. 195 = A. hamosa.
heterophylla, Moench, Meth. 146 = A. Vulneraria.
lagopodium, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 348 = A. Vulneraria.
lotoides, Link, Handb. ii. 195 = A. lotoides.
montana, Scop. Fl. Carn. ed. II. ii. 56 = A. montana.
nudiflora, Lam. Fl. Fr. ii. 650 = A. Gerardi.
polyphylla, Link, Handb. ii. 195 = A. Vulneraria.
rubriflora, Link, l. c. = A. Vulneraria.
rustica, Lam. Fl. Fr. ii. 649 = A. Vulneraria.
sericea, Link, Handb. ii. 194 = A. sericea.
tetraphylla, Moench, Meth. 146 = A. tetraphylla.
vesicaria, Lam. Fl. Fr. ii. 650 = A. tetraphylla.

VULPIA, C. C. Gmel. Fl. Bad. i. 8 (1805) = **Festuca**, Tourn. (Gramin.).
aetnensis, Tineo, Pl. Rar. Sicil. (1846) 22 = F. ciliata.
agrestis, Duval-Jouve, in Rev. Sc. Nat. (Juin, 1880) 49 = F. bromoides.
alopecura, Dum. Obs. Gram. Belg. 100 = F. Alopecuros.
ambigua, A. G. More, in Journ. Linn. Soc. v. (1861) 190 = F. Myuros.
antucensis, Trin. in Linnaea, x. (1836) 303 = F. antucensis.
attenuata, Parl. Giorn. Bot. i. II. (1844) 349 = F. attenuata.
Brauniana, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 419 = Agropyrum scabrum.
brevis, Boiss. & Kotschy, Diagn. Ser. II. 139 = F. inops.
bromoides, S. F. Gray, Nat. Arr. Brit. Pl. ii. 124 = F. bromoides.
Broteri, Boiss. & Reut. Pugill. Pl. Nov. (1852) 128 = F. Myuros.
Browniana, Nees, in Lehm. Pl. Preiss. ii. 107 = Agropyrum scabrum.
ciliata, Link, Hort. Berol. i. 147 ; Dum. Obs. Gram. Belg. 100 = F. ciliata.
cynosuroides, Coss. ex Battand. & Trab. Fl. Alger. (1884) 92 = F. cynosuroides.
delicatula, Dum. Obs. Gram. Belg. 100 = F. delicatula.
geniculata, Link, Hort. Berol. i. 148 = F. clavata.
gypsophila, Nym. Consp. 821 = F. gypsophila.
hirtiglumis, Boiss. & Haussk. in Boiss. Fl. Orient. v. 629.—Mesopot.
hispidula, De Not. ex Parl. Fl. Ital. i. 429 = F. ligustica.
incrassata, Parl. in Ann. Sc. Nat. Sér. II. xv. (1841) 298 = F. clavata.
inops, Hack. in Flora, lxiii. (1880) 476 = F. inops.
Lachenalii, Heynh. Nom. i. 853 = F. Poa.
ligustica, Link, Hort. Berol. i. 148 = F. ligustica.
ligustica, Reichb. Ic. Fl. Germ. i. 25 = F. clavata.
Linneana, Parl. in Giorn. Bot. Ital. i. I. (1844) 346 = F. bromoides.
longiseta, Hack. Cat. Gram. Portugal, 24 = F. bromoides.
longivaginata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = F. Myuros.
maritima, S. F. Gray, Nat. Arr. Brit. Pl. ii. 124 = F. bromoides.
megastachya, Nees, Fl. Afr. Austr. 441.—Afr. austr.
membranacea, Dum. Obs. Gram. Belg. 100 = F. bromoides.
Michelii, Reichb. Fl. Germ. Excurs. 140[3] = Avellinia Michelii.
microstachya, Munro, in Benth. Pl. Hartw. 342 = F. microstachya.
muralis, Nees, in Linnaea, xix. (1847) 694 = F. muralis.
murorum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 123 = F. Myuros.
Myuros, Reichb. Fl. Germ. Excurs. 37 = F. ciliata.
Myurus, C. C. Gmel. Fl. Bad. i. 8 = F. Myuros.
Nardus, Dum. Fl. Belg. 159 = F. tenuiflora.
panormitana, Parl. Fl. Palerm. i. 188 = F. bromoides.
patens, Boiss. Diagn. Ser. I. xiii. 62 = F. pectinella.
pectinata, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 419 = Agropyrum pectinatum.
pectinella, Boiss. Fl. Orient. v. 631 = F. pectinella.
pilosa, C. C. Gmel. Fl. Bad. i. 9, in obs. = F. Myuros.

VULPIA :—
Pseudomyurus, Reichb. Fl. Germ. Excurs. 37 = Festuca Myuros.
quadriflora, Trin. ex Steud. Nom. II. ii. 780 = F. quadriflora.
reclinata, Dum. in Bull. Soc. Bot. Belg. xii. (1873) 204 = F. Myuros ?
rectiseta, Nees, in Lehm. Pl. Preiss. ii. 107 = Agropyrum scabrum.
scabra, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 419 = Agropyron scabrum.
sciuroides, C. C. Gmel. Fl. Bad. i. 9, in obs. = F. bromoides.
setacea, Parl. in Ann. Sc. Nat. Sér. II. xv. (1841) 297 = F. setacea.
sicula, Link, Hort. Berol. ii. 272 = F. sicula.
stipoides, Dum. Obs. Gram. Belg. 100.—Belgium.
subciliata, Link, in Linnaea, xvii. (1843) 402 = F. subciliata.
tenella, Heynh. Nom. i. 854 = F. tenella, *Willd.*
tenuicula, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 27 = Avellinia tenuicula.
tenuiflora, Heynh. Nom. i. 854 = F. tenuiflora.
tenuis, Parl. in Ann. Sc. Nat. Sér. II. xv. (1841) 299 = F. tenuis, *Nym.*
ulochaeta, Nees, ex Steud. Syn. Pl. Gram. 305 = F. ulochaeta.
uniglumis, Dum. Obs. Gram. Belg. 101 = F. bromoides.

VVARIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28, err. typ. = **Uvaria**, Linn. (Anonac.).

VYENOMUS, Presl, Bot. Bemerk. 32 (1844) (in Abh. Boehm. Ges. V. iii. (1845) 462) = **Euonymus**, Linn. (Celastr.).
pendula, Presl, l. c. = E. pendulus.

W

WACCHENDORFIA, Burm. f. Fl. Cap. Prod. 2 (1768), sphalm. = seq.

WACHENDORFIA, Burm. Wachend. (1757) ; et ex Linn. Syst. ed. X. 864 (1759). *HAEMODORACEAE*, Benth. & Hook. f. iii. 673.
PEDILONIA, Presl, Pedil. Nov. Pl. Gen. (1829) ; et in Flora, xii. (1829) 568.
brevifolia, Soland. ex Ker-Gawl. in Bot. Mag. t. 1066 = hirsuta.
Breyniana, *D. Dietr. Syn. Pl.* i. 170 = hirsuta.
corymbosa, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.
elata, Salisb. Prod. 45 = thyrsiflora.
graminea, Thunb. Prod. Pl. Cap. 12 = graminifolia.
graminifolia, *Linn. f. Suppl.* 101.—Afr. austr.
Herberti, Sweet, Hort. Brit. ed. I. 400 = paniculata.
Hibberti, Loud. Hort. Brit. 18, sphalm. = praec.
hirsuta, Linn. Syst. ed. XII. 735 = Dilatris corymbosa.
hirsuta, *Thunb. Prod. Pl. Cap.* 12.—Afr. austr.
humilis, Salisb. Prod. 45 = paniculata.
lugubris, Salisb. l. c. (Quid ?).—Hab. ?
orinocensis, *H. B. & K. Nov. Gen. et Sp.* i. 319.—Venezuela.
pallida, Salisb. Prod. 45.—Afr. austr.
paniculata, *Linn. Syst.* ed. X. 864.—Afr. austr.
tenella, *Thunb. Prod. Pl. Cap.* 12.—Afr. austr.
thyrsiflora, *Linn. Syst.* ed. X. 864.—Afr. austr.
umbellata, Linn. Mant. ii. 320 = Dilatris corymbosa.
villosa, Andr. Bot. Rep. t. 398 = hirsuta.
villosa, Linn. Syst. ed. X. 864 (Quid ?).—Hab. ?

WACHENDORFIA, Loefl. It. Hisp. 177 (1758) = **Callisia**, Linn. (Commel.).

WADAPUS, Rafin. Fl. Tellur. iii. 77 (1836) = **Gomphrena**, Linn. (Amarant.).

WADDINGTONIA, Phil. Fl. Atac. 41. t. 5 (1860) = **Petunia**, Juss. (Solan.).
floribunda, Phil. l. c. = P. parviflora.

WADEA, Rafin. Sylva Tellur. 56 (1838) = **Cestrum**, Linn. (Solan.).
latifolia, Rafin. l. c. = C. latifolium.

WAGATEA, Dalz. in Hook. Kew Journ. iii. (1851) 90. *LEGUMINOSAE*, Benth. & Hook. f. i. 568.
spicata, *Dalz. l. c.*—*Cf.* 89.—Ind. or.

WAGENERIA, Klotzsch, in Abh. Akad. Berl. 1854 (1855) 112. t. 10 C = **Begonia**, Tourn.
brasiliensis, Klotzsch, in Monatsb. Berl. Acad. (Maerz, 1854) 126 ; et in Abh. Akad. Berl. 1854 (1855), 119 = B. polygonifolia.
convolvulacea, Klotzsch, ll. cc. 126 ; 116 = B. convolvulacea.
deflexa, Klotzsch, ll. cc. 126 ; 113 = B. scandens.
dichotoma, Klotzsch, ll. cc. 126 ; 117 = B. dichotoma.
fagifolia, Klotzsch, ll. cc. 126 ; 113 = B. fagifolia.
glabra, Klotzsch, ll. cc. 126 ; 115 = B. scandens.
glandulosa, Klotzsch, ex A. DC. in DC. Prod. xv. I. 339 = B. glandulosa.
hispida, Klotzsch, ex A. DC. l. c. 363 = B. hispida.
Huegelii, Klotzsch, in Ind. Sem. Hort. Berol. (1855) App. 2 = B. Huegelii.
inciso-serrata, A. DC. in Mart. Fl. Bras. iv. I. 374, sphalm. = B. inciso-serrata.
longiceps, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 = B. longipes.
longipes, Klotzsch, in Abh. Akad. Berl. 1854 (1855), 117 = B. longipes.
lucida, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 ; et in Abh. Akad. Berl. 1854 (1855), 114 = B. scandens.
montana, Klotzsch, ll. cc. 126 ; 115 = B. scandens.
Moritziana, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 = B. meridensis.
reniformis, Klotzsch, l. c. ; et in Abh. Akad. Berl. 1854 (1855), 118 = B. longipes.
rugosa, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 = B. rugosa.
scabrida, Klotzsch, ex A. DC. in DC. Prod. xv. I. 367 = B. scabrida.
scandens, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126, nomen = B. scandens ?
Schottiana, Klotzsch, l. c. = B. convolvulacea.
tomentosa, Klotzsch, l. c. = B. tomentosa.
vitifolia, Klotzsch, l. c. ; et in Abh. Akad. Berl. 1854 (1855), 116 = B. vitifolia.

WAGNERIA, Klotzsch, in Monatsb. Berl. Akad. (Maerz, 1854) 126 = praec.

WAGNERIA, Lem. Illustr. Hortic. iv. (1857) sub t. 115 = **Diervilla**, Tourn. (Caprifol.).
Middendorffiana, Lem. l. c. = D. Middendorfiana.

WAHABIA, Fenzl, in Flora, xxvii. (1844) 312, nomen = **Barleria**, Linn. (Acanth.).
longiflora, Fenzl, l. c., nomen = B. acanthoides.

WAHLBERGELLA, Fries, in Bot. Notiser (1843), 143 = **Lychnis**, Tourn. (Caryoph.).
affinis, Fries, l. c. = L. affinis.
albiflora, Rupr. in Hofm. Noerdl. Ural. ii. Anhang, 30 = L. apetala ?
angustiflora, Rupr. Beitr. Pfl. Russ. Reich. ii. 24 = L. affinis.
apetala, Fries, Summa Veg. Scand. 155 = L. apetala.
involucrata, Rupr. Beitr. Pfl. Russ. Reich. ii. 24 (= *Gasterolychnis involucrata*).—Reg. Ural.
macropetala, Rupr. in Hofm. Noerdl. Ural. ii. Anhang, 30 = L. apetala ?
triflora, Fries, Summa Veg. Scand. 155, in obs. = L. affinis.
uniflora, Fries, in Bot. Notiser (1843), 143.—Europ. ?
uralensis, Rupr. in Hofm. Noerdl. Ural. ii. Anhang, 17, 31 (= *Gasterolychnis uralensis*).—Reg. Ural.
Vahlii, Rupr. Beitr. Pfl. Russ. Reich. ii. 24 = L. affinis.

WAHLBOMIA, Thunb. in Vet. Akad. Handl. Stockh. (1790) 215. t. 9 = **Tetracera**, Linn. (Dilleniac.).
indica, Thunb. l. c. = T. Wahlbomia.

WAHLENBERGIA, Blume, Cat. Gew. Buitenz. 14 (1823) = **Tarenna**, Gaertn. (Rubiac.).
fragrans, Blume, l. c. 15 (= *Webera fragrans*).—Malaya.

WAHLENBERGIA, R. Br. in Wall. Cat. p. 262.—Cf. n. 4342 (1831) = **Chailletia**, DC.

WAHLENBERGIA, Schrad. Cat. Hort. Gotting. (1814); et in Comm. Gotting. vi. (1823–27) 123. *CAMPANULACEAE*, Benth. & Hook. f. ii. 555.
AIKINIA, Salisb. ex A. DC. Monog. Camp. 135, 141 (1830).
CERVICINA, Delile, Fl. Egypte, 150. t. 5. f. 2 (1813).
EDRAIANTHUS, A. DC. in DC. Prod. vii. 448 (1839).
HECALE, Rafin. Fl. Tellur. ii. 79 (1836).
HEDRAEANTHUS, Griseb. Spicil. Fl. Rumel. ii. 292 (1844).
PETALOSTIMA, Rafin. Fl. Tellur. ii. 79 (1836).
PILOREA, Rafin. l. c. 80 (1836).
SCHULTESIA, Roth, Enum. Pl. Phan. Germ. i. 690 (1827).
STRELESKIA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 226.
VALVINTERLOBUS, Dulac, Fl. Hautes-Pyr. 459 (1867).
acaulis, *E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr.* (1837) 9, nomen; *A. DC. in DC. Prod.* vii. 430.—Afr. austr.
adpressa, *Sond. in Harv. & Sond. Fl. Cap.* iii. 583.—Afr. austr.
agrestis, A. DC. Monog. Camp. 145 = gracilis.
albomarginata, Hook. Ic. Pl. t. 818 = saxicola.
androsacea, *A. DC. Monog. Camp.* 150. t. 19. f. 1.—Afr. austr.
angustifolia, *A. DC. l. c.* 162.—Ins. S. Helen.
annularis, *A. DC. in DC. Prod.* vii. 437.—Afr. austr.
arenaria, A. DC. l. c. 436 = androsacea.
arguta, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 15.—Afr. trop.
arida, *Griseb. in Goett. Abh.* xix. (1874) 200.—Reg. Argent.
arvatica, Sweet, Hort. Brit. ed. II. 593 = hederacea.
Banksiana, *A. DC. Monog. Camp.* 154.—Afr. austr.
Berteroi, *Hook. & Arn. in Hook. Journ. Bot.* i. (1834) 279.—Ins. J. Fernand.
bilocularis, A. DC. in DC. Prod. vii. 439 = undulata.
Bojeri, *A. DC. l. c.* 435.—Ins. Madag.
Bowkeriae, *Sond. in Harv. & Sond. Fl. Cap.* iii. 577.—Afr. austr.
brasiliensis, *Cham. in Linnaea*, viii. (1833) 318.—Bras.
Burchellii, *A. DC. in DC. Prod.* vii. 438.—Ins. S. Helen.
caffra, *A. DC. l. c.* 432.—Afr. austr.
caledonica, *Sond. in Harv. & Sond. Fl. Cap.* iii. 579.—Afr. austr.
californica, Kellogg, in Proc. Calif. Acad. ii. (1863) 158 = Campanula linnaeifolia.
calycina, *Schlecht. in Lechl. Berb. Am. Austr.* 58, nomen.—Peruv.
campanuloides, Vatke, in Linnaea, xxxviii. (1874) 700 = W. Cervicina.
capensis, *A. DC. Monog. Camp.* 136. t. 18.—Afr. austr.
capillacea, *A. DC. l. c.* 156.—Afr. austr.
capillaris, G. Don, in Sweet, Hort. Brit. ed. III. 419 = gracilis.
capillifolia, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen (Quid?).—Afr. austr.
cartilaginea, *Hook. f. Handb. N. Zeal. Fl.* 170.—N. Zel.
cernua, *A. DC. Monog. Camp.* 148.—Afr. austr.
cernua, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 9; et ex A. DC. in DC. Prod. vii. 439 = Meyeri.
Cervicina, *A. DC. Monog. Camp.* 156.—Afr. bor.
Chamissoniana, G. Don, Gen. Syst. iii. 740 = undulata.
chondrophylla, Buek, in Eckl. & Zeyh. Enum. 381 = Lobelia aspera.
ciliolata, A. DC. in DC. Prod. vii. 436 = cernua.
claviculata, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 9, nomen; et ex A. DC. in DC. Prod. vii. 432 = exilis.
clematidea, Schrenk, Enum. Pl. Nov. 38 = Codonopsis ovata.
clivosa, A. DC. Monog. Camp. 161 = angustifolia.
corymbosa, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen; et ex A. DC. in DC. Prod. vii. 432 = Lightfootia capillaris.
costata, *A. DC.* in DC. Prod. vii. 427.—Afr. austr.
crispa, *Nees, Del. Sem. Hort. Vratisl.* (1850) 3.—*Cf. Linnaea*, xxiv. (1851) 237.—Armenia.
dalmatica, *A. DC. Monog. Camp.* 134.—Dalmat.

WAHLENBERGIA:—
debilis, *Buek, in Eckl. & Zeyh. Enum.* 378.—Afr. austr.
decipiens, *A. DC. in DC. Prod.* vii. 427.—Afr. austr.
dehiscens, A. DC. Monog. Camp. 145 = gracilis.
denticulata, A. DC. l. c. 152. t. 16 = Lightfootia capillaris.
denudata, *A. DC. l. c.* 147.—Afr. austr.
dicentrifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 430.—Reg. Himal.
dichotoma, A. DC. in DC. Prod. vii. 437 = androsacea.
diffusa, A. DC. Monog. Camp. 137 = Microcodon lineare.
divaricata, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen; et ex A. DC. in DC. Prod. vii. 432 = paniculata, ingrata.
divergens, *A. DC. in DC. Prod.* vii. 427.—Afr. austr.
diversifolia, A. DC. Monog. Camp. 139 = procumbens.
Dregeana, A. DC. in DC. Prod. vii. 428 = Ecklonii.
Dunantii, A. DC. Monog. Camp. 152 = paniculata.
Ecklonii, *Buek, in Eckl. & Zeyh. Enum.* 380.—Afr. austr.
Elatines, Link, Handb. i. 631 = Campanula Elatines.
elongata, Schrad. Cat. Hort. Gotting. (1814); ex Roth, Nov. Sp. Pl. 399 = capensis.
emirnensis, *A. DC. in DC. Prod.* vii. 432.—Madag.
ensifolia, *A. DC. Monog. Camp.* 162.—Ins. Borbon.
epacridea, *Sond. in Harv. & Sond. Fl. Cap.* iii. 584.—Afr. austr.
Erinus, Link, Handb. i. 631 = Campanula Erinus.
etbaica, *Vatke, in Linnaea*, xxxviii. (1874) 700.—Afr. trop.
exilis, *A. DC. Monog. Camp.* 151. t. 16.—Afr. austr.
fernandeziana, *A. DC. l. c.* 160.—Ins. J. Fernand.
flaccida, *A. DC. l. c.* 138.—Afr. austr.
flaccida, Presl, Symb. Bot. 29. t. 18 = Campanula garganica.
foliosa, A. DC. Monog. Camp. 160 = linifolia.
garganica, G. Don, Gen. Syst. iii. 738 = Campanula garganica.
gracilis, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen; et ex A. DC. in DC. Prod. vii. 435 = cernua.
gracilis, *Schrad. Blumenb.* 38, in obs.; *A. DC. Monog. Camp.* 142.—Geront. trop. et temp. austr.
Grahamae, *Hemsl. Bot. Voy. Challenger*, i. III. 46.—Ins. J. Fernand.
graminifolia, *A. DC. Monog. Camp.* 130.—Italia.
grandiflora, Schrad. Cat. Hort. Gotting. (1814); et ej. Blumenb. 38, in obs. = Platycodon grandiflorum.
hederacea, *Reichb. Pl. Crit.* v. 47; *A. DC. Monog. Camp.* 140.—Europ.
Hilsenbergii, *A. DC. in DC. Prod.* vii. 429.—Madag.
hirsuta, *Steud. Nom.* ed. II. ii. 782.—Chili.
hispidula, Eckl. & Zeyh. Enum. 381 = paniculata.
hispidula, Schrad. Blumenb. 38, in obs.; A. DC. Monog. Camp. 155 = Microcodon depressum.
hispidula, Schrad. ex A. DC. Monog. Camp. 128 = Microcodon sparsiflorum.
homallanthina, *A. DC. in DC. Prod.* vii. 425.—Sibir. or.
huillana, *A. DC. in Ann. Sc. Nat.* Sér. V. vi. (1866) 333.—Afr. trop.
humilis, *A. DC. Monog. Camp.* 147.—Afr. trop.
inconspicua, *A. DC. in DC. Prod.* vii. 430.—Afr. austr.
indica, A. DC. Monog. Camp. 146 = gracilis.
ingrata, *A. DC. in DC. Prod.* vii. 432.—Afr. austr.
inhambanensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 303.—Afr. trop.
integrifolia, A. DC. in DC. Prod. vii. 428 = stellarioides.
Kitaibelii, *A. DC. Monog. Camp.* 131.—Banat.; Transsylv.
Krebsii, Cham. in Linnaea, viii. (1833) 195 = Zeyheri.
Kruhsiana, Fisch. ex A. DC. Monog. Camp. 126 = homallanthina.
Larrainii, A. DC. in DC. Prod. vii. 438, sphalm., in syn. = fernandeziana.
lavandulaefolia, A. DC. Monog. Camp. 144 = gracilis.
linarioides, *A. DC. l. c.* 158.—Am. austr.
linearis, A. DC. l. c. 137 = Microcodon lineare.
linearis, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen (Quid?).—Afr. austr.
linifolia, *A. DC. Monog. Camp.* 162.—Ins. S. Helen.
lobelioides, Link, Handb. i. 632; A. DC. l. c. 157 = pendula.
lurida, Schouw. Hort. Haun.—Cf. Linnaea, xxiv. (1851) 161 = Codonopsis lurida.

WAHLENBERGIA:—
madagascariensis, *A. DC. Monog. Camp.* 139.—Madag.
Mannii, *Vatke, in Linnaea*, xxxviii. (1874) 700.—Afr. trop.
marginata, A. DC. Monog. Camp. 143 = gracilis.
Massonii, A. DC. l. c. 153 = paniculata.
Meyeri, *A. DC. in DC. Prod.* vii. 439.—Afr. austr.
micrantha, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 9; et ex A. DC. in DC. Prod. vii. 411 = Laurentia micrantha.
montana, *A. DC. in DC. l. c.* 430.—Afr. austr.
multicaulis, Benth. in Enum. Pl. Hueg. 75 = gracilis.
namaquana, Sond. in Harv. & Sond. Fl. Cap. iii. 582 = annularis.
nudicaulis, *A. DC. Monog. Camp.* 149.—Afr. austr.
nudicaulis, Eckl. & Zeyh. Enum. 380 = androsacea.
nutabunda, *A. DC. Monog. Camp.* 151.—Ital.
oocarpa, *Sond. in Harv. & Sond. Fl. Cap.* iii. 576.—Afr. austr.
oppositifolia, *A. DC. in DC. Prod.* vii. 429.—Afr. austr.
ovata, *D. Don, Prod. Fl. Nep.* 156.—Nepal.
oxyphylla, *A. DC. in DC. Prod.* vii. 427.—Afr. austr.
paniculata, Eckl. & Zeyh. Enum. 380 = Ecklonii.
paniculata, *Schrad. Blumenb.* 38, in obs.; *A. DC. Monog. Camp.* 153.—Afr. austr.
parviflora, A. DC. in DC. Prod. vii. 437 = androsacea.
patula, *A. DC. l. c.* 436.—Afr. austr.
pauciflora, *A. DC. l. c.* 437.—Afr. austr.
peduncularis, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 22.—Reg. Himal.
pendula, *Schrad. Cat. Hort. Gotting.* (1814) 3, nomen; *et ej. Blumenb.* 38, in obs.—Ins. Mader. et Canar.
pentamera, A. DC. in DC. Prod. vii. 428 = Ecklonii.
perotifolia, Wight & Arn. Prod. 405 = Cephalostigma Schimperi.
peruviana, *A. Gray, in Proc. Am. Acad.* v. (1861) 152.—Peruv.
pilosa, *Buek, in Eckl. & Zeyh. Enum.* 381.—Afr. austr.
pilosa, DC. in DC. Prod. vii. 431 = polyclada.
polyclada, *A. DC. l. c.* 789.—Afr. austr.
polyclada, Hook. f. in Journ. Linn. Soc. vi. (1862) 15 = Mannii.
Preissii, De Vriese, in Lehm. Pl. Preiss. ii. 241 = gracilis?
procumbens, *A. DC. Monog. Camp.* 140. t. 15.—Afr. austr.
prostrata, *E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr.* (1837) 9, nomen; *et ex A. DC. in DC. Prod.* vii. 431.—Afr. austr.
Pumilio, *A. DC. Monog. Camp.* 134.—Dalmatia.
purpurea, A. DC. in DC. Prod. vii. 425 = Codonopsis purpurea.
pusilla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 2.—Abyss.
quadrifida, A. DC. Monog. Camp. 144 = gracilis.
ramulosa, *E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr.* (1837) 8, nomen; *et ex A. DC. in DC. Prod.* vii. 432.—Afr. austr.
reflexa, *Steud. Nom.* ed. II. ii. 782.—Chili.
repens, Schrad. Blumenb. 37 = procumbens.
rigida, *Bernh. in Flora*, xxvii. (1844) 820.—Afr. austr.
riparia, *A. DC. Monog. Camp.* 146.—Afr. trop.
robusta, Sond. in Harv. & Sond. Fl. Cap. iii. 584 = rigida.
rotundifolia, A. DC. in DC. Prod. vii. 425 = Codonopsis rotundifolia.
Roxburghii, *A. DC. Monog. Camp.* 162.—Ins. S. Helen.
Roylei, A. DC. in DC. Prod. vii. 425 = Codonopsis ovata.
rudis, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen; et ex A. DC. in DC. Prod. vii. 431 = paniculata.
Rutenbergiana, *Vatke, in Bremen, Abh.* ix. (1885) 123.—Madag.
saxicola, *A. DC. Monog. Camp.* 144.—Austral.; N. Zel.
Saxifraga, A. DC. in DC. Prod. vii. 427 = Ecklonii.
Schimperi, Hochst. ex Hemsl. in Oliver, Fl. Trop. Afr. iii. 472 = Cephalostigma hirsutum.
semiglabra, A. DC. in DC. Prod. vii. 437 = androsacea.
Sieberi, A. DC. Monog. Camp. 144 = gracilis.
silenoides, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 3.—Abyss.
simplicicaulis, De Vriese, in Lehm. Pl. Preiss. ii. 241 = gracilis.

WAHLENBERGIA:—
sphaerocarpa, A. DC. in DC Prod. vii. 436 = androsacea.
spicata, E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr. (1837) 8, nomen; A. DC. in DC. Prod. vii. 441 = Lightfootia juncea.
spinulosa, *A. DC. Monog. Camp.* 155.—Afr. austr.
stellarioides, *Cham. in Linnaea*, viii. (1833) 196.—Afr. austr.
striata, A. DC. in DC. Prod. vii. 439 = undulata.
stricta, Sweet, Hort. Brit. ed. II. 593 = gracilis.
swellendamensis, Buek, in Eckl. & Zeyh. Enum. 381 = Ecklonii.
tenuifolia, *A. DC. Monog. Camp.* 133.—Croatia.
tenuis, A. DC. in DC. Prod. vii. 440 = debilis.
thalictrifolia, A. DC. l. c. 425 = Codonopsis thalictrifolia.
tuberosa, *Hook. f. Bot. Mag.* t. 6155.—Ins. J. Fernand.
turbinata, A. DC. in DC. Prod. vii. 427 = Ecklonii.
undulata, *A. DC. Monog. Camp.* 148.—Afr. austr.
variabilis, *E. Mey. in Drège, Cat. Pl. Exsicc. Afr. Austr.* (1837) 8, nomen; *et ex A. DC. in DC. Prod.* vii. 436.—Afr. austr.
Verreauxii, A. DC. in DC. Prod. vii. 436 = androsacea.
vincaeflora, Decne. in Rev. Hortic. Sér. III. iii. (1849) 41 = gracilis.
viridis, A. DC. in DC. Prod. vii. 424 = Codonopsis viridis.
Wyleyana, *Sond. in Harv. & Sond. Fl. Cap.* iii. 583.—Afr. austr.
Zeyheri, *Buek, in Eckl. & Zeyh. Enum.* 379.—Afr. austr.

WAHLENBERGIA, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 387 (1827) = **Enhydra**, Lour. (Compos.).
globularis, Schum. in Schum. & Thonn. l. c. = E. fluctuans.
glomerata, DC. Prod. v. 497, in syn., sphalm. = praec.

WAILESIA, Lindl. in Journ. Hort. Soc. iv. (1849) 261 = **Dipodium**, R. Br. (Orchid.).
paludosa, Reichb. f. in Bonplandia, ii. (1854) 92 = D. paludosum.
picta, Lindl. in Journ. Hort. Soc. iv. (1849) 261 = D. pictum.

WAITZIA, Reichb. Consp. 60 (1828) = **Tritonia**, Ker-Gawl. (Irid.).
aequimaculata, Heynh. Nom. i. 854 (= *Montbretia aequimaculata*).—Hab. ?
anigozanthaeflora, Heynh. l. c. = T. anigozanthiflora.
concolor, Heynh. l. c. = Ixia paniculata.
crispa, Heynh. l. c. = T. crispa.
crocata, Hoffm. Verz. Pfl. Nachtr. iii. 14 = T. crocata.
deusta, Heynh. Nom. i. 854 = T. deusta.
fenestrata, Heynh. l. c. = T. fenestrata.
flava, Heynh. l. c. = T. flava.
fucata, Heynh. l. c. = Antholyza fucata.
lineata, Heynh. l. c. = T. lineata.
longiflora, Heynh. l. c. = Ixia paniculata.
miniata, Heynh. l. c. = T. crocata.
odorata, Heynh. l. c. = Freesia refracta.
pallida, Heynh. l. c. 855 = T. pallida.
pectinata, Heynh. l. c. = T. crispa.
refracta, Heynh. l. c. = Freesia refracta.
Rochensis, Heynh. l. c. = Ixia paniculata.
rosea, Heynh. l. c. = T. capensis.
squalida, Heynh. l. c. = T. squalida.
striata, Heynh. l. c. = Gladiolus striatus.
tenuiflora, Heynh. l. c. = Ixia paniculata.
viridis, Heynh. l. c. = T. viridis.
xanthospila, Heynh. l. c. (= *Anomatheca xanthospila*).—Afr. austr.

WAITZIA, Wendl. Coll. Pl. ii. 13. t. 42 (1808). *COMPOSITAE*, Benth. & Hook. f. ii. 307.
MORNA, Lindl. Bot. Reg. t. 1941 (1837); et (1838) t. 9.
PTEROCHAETA, Steetz, in Lehm. Pl. Preiss. i. 455 (1845).
VIRAYA, Gaudich. in Freyc. Voy. Bot. 466. t. 89 (1826).
acuminata, Steetz, in Lehm. Pl. Preiss. i. 453 = corymbosa.
aurea, *Steetz, l. c.* 452.—Austral.
brachyrrhyncha, F. Muell. in Linnaea, xxv. (1852) 407 = Helipterum incanum.

WAITZIA:—
brevirostris, Steetz, in Lehm. Pl. Preiss. i. 451 = Steetziana.
citrina, Steetz, l. c. = Steetziana.
corymbosa, *Wendl. Coll. Pl.* ii. 13. t. 42.—Austral.
dasycarpa, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 77 = Steetziana.
discolor, Turcz. l. c. (1851) I. 194 = corymbosa.
grandiflora, *Naud. in Rev. Hortic.* (1865) 185.—Austral.
nivea, Benth. Fl. Austral. iii. 636 = odontolepis.
odontolepis, *Turcz in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 77.—Austral.
paniculata, *F. Muell. ex Benth. Fl. Austral.* iii. 637.—Austral.
Podolepis, *Steetz, in Lehm. Pl. Preiss.* i. 450.—Austral.
Steetziana, *Lehm. ex Steetz, l. c.* 454.—Austral.
sulphurea, Steetz, l. c. 453 = Steetziana.
tenella, Hook. Bot. Mag. t. 5342 = Steetziana.

WALAFRIDA, E. Mey. Comm. Pl. Afr. Austr. 272 (1837) = **Selago**, Linn.
nitida, E. Mey. l. c.—Afr. austr.
trimera, Hochst. in Flora, xxviii. (1845) 70.—Afr. austr.

WALAFRIDIA, Endl. Gen. 642 (1839) = praec.

WALCOTTIA, F. Muell. Fragm. i. 241 (1859) = **Lachnostachys**, Hook. (Verben.).
eriobotrya, F. Muell. l. c. = L. Walcottii.

WALCUFFA, J. F. Gmel. Syst. 1029 (1791) = **Dombeya**, Cav. (Stercul.).
Pentapetes, Bruce, ex Steud. Nom. ed. I. 891 = D. Bruceana.
torrida, J. F. Gmel. Syst. 1029 = D. Bruceana.

WALDEMARIA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 99 (1862) = **Rhododendron**, Linn. (Ericac.).
argentea, Klotzsch, l. c. = R. grande.

WALDHEIMIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 125 = **Allardia**, Decne. (Compos.).
glabra, Regel, in Act. Hort. Petrop. vi. (1879) 308, 309 = A. glabra.
Korolkowi, Regel & Schmalh. l. c. 308, 310.—Turkestan.
nivea, Regel, l. c. 308, 309 = A. nivea.
Stracheyana, Regel, l. c.—Reg. Himal.
tomentosa, Regel, l. c. 308, 310 = A. tomentosa.
tridactylites, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 126 = A. glabra.

WALDSCHMIDIA, [Weber, in] Wigg. Prim. Fl. Holsat. 19 (1780) = **Limnanthemum**, S. T. Gmel. (Gentian.).
nymphoides, [Weber, in] Wigg. l. c. 20 = L. nymphoides.

WALDSCHMIDTIA, Bluff & Fingerh. Comp. Fl. Germ. i. 265 (err. typ. 465) (1825) = praec.

WALDSCHMIDTIA, Scop. Introd. 100 (1777) = **Crudia**, Schreb. (1789) (Legumin.).

WALDSTEINIA, Willd. in Ges. Naturf. Fr. Neue Schr. ii. (1799) 105. t. 4. *ROSACEAE*, Benth. & Hook. f. i. 619.
COMAROPSIS, Rich. in Nestl. Potent. 16. t. 1 (1816).
Doniana, Tratt. Ros. Monog. iii. 109 = fragarioides.
fragarioides, *Tratt. l. c.* 107.—Am. bor.
geoides, *Willd. in Ges. Naturf. Fr. Neue Schr.* ii. (1799) 106.—Hungar.; Tauria.
lobata, *Torr. & Gray, Fl. N. Am.* i. 426.—Am. bor.
Moenchii, *Tratt. Ros. Monog.* iii. 106.—Hab. ?
sibirica, *Tratt. l. c.* 108.—Sibir. baical.
trifolia, Rochel, ex Koch, in Linnaea, xiii. (1839) 337, 338 = sibirica.
trifoliata, Steud. Nom. ed. II. ii. 783, sphalm. = praec.

WALKERA, Schreb. Gen. 378 (1789) = **Gomphia**, Schreb. (Ochnac.).
integrifolia, DC. Prod. i. 737.—Guian.
serrata, Willd. Sp. Pl. i. 1145 = G. angustifolia.
zeylonensis, Hort. ex DC. in Ann. Mus. Par. xvii. (1811) 421 = G. angustifolia.

WALKERIA, Mill. ex Ehret, in Phil. Trans. liii. (1763) 130 = **Nolana**, Linn. (Convolv.).

WALKUFFA, Bruce, ex Steud. Nom. ed. I. 891 (1821) = WALCUFFA, J. F. Gmel. = **Dombeya**, Cav. (Stercul.).

WALLACEA, Spruce, ex Benth. & Hook. f. Gen. i. 320 (1862). *OCHNACEAE*, Benth. & Hook. f. l. c.
insignis, *Spruce, ex Benth. & Hook. f. l. c.* 321.—Bras.

WALLENIA, Sw. Prod. Veg. Ind. Occ. 31 (1788). *MYRSINEAE*, Benth. & Hook. f. ii. 643.
PETESIOIDES, Jacq. Stirp. Am. 17 (1763).
angularis, Jacq. Hort. Schoenb. i. t. 30 = seq.
angulata, *Jacq. l. c.* i. 13.—Hab. ?
angustifolia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 87 = Ardisia semicrenata.
clusiifolia, *Griseb. Fl. Brit. W. Ind.* 394.—Ind. occ.
laurifolia, Balb. ex Griseb. Fl. Brit. W. Ind. 394 = clusiifolia.
laurifolia, *Sw. Prod. Veg. Ind. Occ.* 31.—Ind. occ.
laxiflora, Mart. Nov. Gen. et Sp. iii. 89. t. 237 = Conomorpha laxiflora.
myrianthos, Reichb. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 102 = Cybianthus myrianthos.
nemoralis, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 21.—Bras.
venosa, *Griseb. Fl. Brit. W. Ind.* 394.—Jamaic.

WALLERIA, J. Kirk, in Trans. Linn. Soc. xxiv. (1864) 497. t. 52. *LILIACEAE*, Benth. & Hook. f. iii. 831.
angolensis, Baker, l. c. Ser. II. i. (1878) 262 = Mackenzii.
Mackenzii, *J. Kirk, l. c.* Ser. I. xxiv. (1864) 497. t. 52. f. 2.—Afr. trop.
nutans, J. Kirk, l. c. t. 52. f. 1 = Mackenzii.

WALLIA, Alef. in Bonplandia, ix. (1861) 335 = **Juglans**, Linn.
cinerea, Alef. l. c. 336 = J. cinerea.
fraxinifolia, Alef. l. c. = Pterocarya caucasica.
nigra, Alef. l. c. = J. nigra.
pyriformis, Alef. l. c. = J. pyriformis.
regia, C. DC. in DC. Prod. xvi. II. 136, in syn., sphalm. = J. regia.

WALLICHIA, DC. in Mém. Mus. Par. x. (1823) 104. t. 6 = **Eriolaena**, DC. (Stercul.).
quinquelocularis, Steud. Nom. ed. II. ii. 783 = E. quinquelocularis.
spectabilis, DC. in Mém. Mus. Par. x. (1823) 104. t. 6 = E. spectabilis.

WALLICHIA, Reinw. ex Blume, in Cat. Gew. Buitenz. 11 (1823) = **Urophyllum**, Wall. (Rubiac.).
arborea, Reinw. l. c. = U. glabrum.
porphyracea, Mart. ex Rosenth. Syn. Pl. Diaphor. 333 = U. porphyraceum.

WALLICHIA, Roxb. Pl. Corom. iii. 91. t. 295 (1819). *PALMAE*, Benth. & Hook. f. iii. 916.
HARINA, Buch.-Ham. in Mem. Wern. Soc. v. (1826) 317.
WRIGHTEA, Roxb. Hort. Beng. 68 (1814); Fl. Ind. iii. 621 (1832).
caryotoides, *Roxb. Pl. Corom.* iii. 91. t. 295.—Reg. Himal.
caryotoides, Wall. Cat. n. 8596 B = densiflora.
caudata, Mart. Hist. Nat. Palm. iii. 315 = Didymosperma caudatum.
densiflora, *Mart. l. c.* 190.—Reg. Himal.
disticha, *T. Anders. in Journ. Linn. Soc.* xi. (1871) 6.—Reg. Himal.
excelsa, Karst. ex Linden, in Illustr. Hortic. xxviii. (1881) 32, nomen (Quid ?).—N. Granat.
Horsfieldii, Blume, Rumphia, ii. 112, in obs. = Didymosperma Horsfieldii.
nana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 488 = Didymosperma nanum.
oblongifolia, Griff. l. c. 486 = densiflora.
Oranii, Blume, Rumphia, ii. 113 = Didymosperma porphyrocarpum.
porphyrocarpa, Mart. Hist. Nat. Palm. iii. 190, 315 = Didymosperma porphyrocarpum.

WALLICHIA:—
Reinwardtiana, Miq. Pl. Jungh. 157 = Didymosperma Reinwardtiana.
tremula, Mart. Hist. Nat. Palm. iii. 315 = Didymosperma tremula.
Yomae, Kurz, For. Fl. Brit. Burma, ii. 533 = disticha.
zebrina, Hort. Williams.—Cf. Gard. Chron. (1880) i. 759 (Quid?).—Hab.?

WALLINIA, Moq. in DC. Prod. xiii. ii. 143 (1849) = **Lophiocarpus**, Turcz. (Chenop.).
polystachya, Moq. l. c. = L. polystachyus.

WALLISIA, E. Morr. in Belg. Hortic. (1870) 97. t. 5 = **Tillandsia**, Linn. (Bromel.).
Hamaleana, E. Morr. l. c. = T. Hamaleana.

WALLISIA, Regel, Gartenfl. xxiv. (1875) 230. t. 839 = **Lisianthus**, Aubl. (Gentian.).
Princeps, Regel, l. c. = L. Princeps.

WALLROTHIA, Roth, Nov. Pl. Sp. 317 (1821) = **Vitex**, Tourn. (Verben.).
articulata, Roth, l. c. 317 = V. pubescens.
Leucoxylon, Roth, l. c. 319 = V. Leucoxylon.

WALLROTHIA, Spreng. Pugill. ii. 52 (1815) = **Seseli**, Linn. (Umbellif.).
divaricata, Presl, Delic. Prag. 134 = Carum alpinum.
splendens, Spreng. in Schult. Syst. vi. 557.—Mont. Pyren.
tenuifolia, DC. Prod. iv. 162 = *splendens*.
tuberosa, Spreng. Pugill. ii. 52 = Carum alpinum.

WALPERSIA, Harv. & Sond. Fl. Cap. ii. 26 (1861).
LEGUMINOSAE, Benth. & Hook. f. i. 473.
burtonioides, *Harv. & Sond. l. c.*—Afr. austr.

WALPERSIA, Meissn. ex Krauss, in Flora, xxvii. (1844) 357 = **Rhynchosia**, Lour. (Legumin.).
villosa, Meissn. ex Krauss, l. c. = R. sigmodes.

WALPERSIA, Reiss. in Endl. Gen. 1100 (1840) = **Phylica**, Linn. (Rhamnac.).
capitata, Presl, Bot. Bemerk. 37, 38 = P. spicata.
curvifolia, Presl, l. c. 38 = P. reclinata.
Dregeana, Presl, l. c. = P. stipularis.
hirtifolia, Presl, l. c.—Afr. austr.
rhipophora, Presl, l. c. 37 = P. stipularis.
rigida, Presl, l. c. 38 = P. rigida.
spicata, Reiss. in Endl. Gen. 1100 = P. spicata.
squarrosa, Presl, Bot. Bemerk. 38 = P. plumosa.
stipularis, Reiss. in Endl. Gen. 1100 = P. stipularis.

WALSURA, Roxb. Hort. Beng. 32 (1814); Fl. Ind. ii. 386 (1832). *MELIACEAE*, Benth. & Hook. f. i. 336.
Monocyclis, Wall. ex Voigt, Hort. Suburb. Calc. 135 (1845).
Surwala, M. Roem. Syn. Hesper. 108 (1846).
Antila, Wight & Arn. Prod. 120 = ternata.
Gardneri, *Thw. Enum. Pl. Zeyl.* 61.—Zeylan.
hypoleuca, *Kurz, Rep. Veget. Andam.* ed. II. 33, nomen; *et in Journ. As. Soc. Beng.* xli. (1872) ii. 296.—Ins. Andaman.
lanceolata, Wall. Cat. n. 4886 = Aglaia Roxburghiana.
neurodes, Hiern, in Hook. f. Fl. Brit. Ind. i. 564.—Malacca.
oxycarpa, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 200.—Ins. Andaman.
pinnata, *Hassk. Retzia*, i. 147.—Java.
piscidia, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 387.—Ind. or.
pubescens, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 297 = Heynea trijuga.
robusta, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 386.—Reg. Himal.; Burma.
ternata, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 389.—Ind. or.
Thwaitesii, C. DC. in DC. Monog. Phan. i. 635 = W. Piscidia.
trichostemon, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 60.—Burma; Siam.
trijuga, Kurz, in Journ. As. Soc. Beng. xliv. (1875) ii. 148 = Heynea trijuga.
tubulata, *Hiern, in Hook. Fl. Brit. Ind.* i. 563.—Reg. Himal.

WALSURA:—
villosa, Kurz, in Journ. As. Soc. Beng. xxxix. ii. (1870) 72 = trichostemon.
villosa, *Wall. Cat.* n. 1264; *Wight & Arn. Prod.* 120.—Burma.

WALTERIANA, Fraser, ex Endl. Gen. 1413 (1841) = **Cliftonia**, Banks (Cyrill.).

WALTHERIA, Linn. Gen. ed. I. 203 (1737).
STERCULIACEAE, Benth. & Hook. f. i. 224.
Astropus, Spreng. Neue Entdeck. iii. 64 (1822).
Lophanthus, Forst. Char. Gen. 27. t. 14 (1776).
africana, Schum. & Thonn. Beskr. Guin. Pl. 296 = indica.
albicans, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 214.—Bras.
americana, Linn. Sp. Pl. 673 = indica.
angustifolia, Linn. Syst. ed. X. 1140; H.B. & K. Nov. Gen. et Sp. v. 332 = indica.
arborescens, Cav. Diss. vi. 316 = indica.
Astropus, *Spreng. Syst.* iii. 31.—Bras.
boliviensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 571.—Bolivia.
brachypetala, Turcz. l. c. xxxi. (1858) i. 215 = ferruginea.
bracteosa, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 37.—Bras.
brevipes, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 213.—Mexic.
capitata, *Vell. Fl. Flum.* 274; vii. t. 4.—Bras.
caroliniana, Walt. Fl. Carol. 175 = Piriqueta fulva.
carpinifolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 38.—Bras.
cinerascens, *A. St. Hil. Fl. Bras. Mer.* i. 152.—Bras.
communis, *A. St. Hil. l. c.* 155.—Bras.
corchorifolia, Pers. Syn. ii. 216 = indica.
cordata, *Sm. in Rees, Cycl.* xxxvii. n. 8.—Ind. occ.
debilis, Boj. Hort. Maurit. 39, nomen (Quid?).—Ins. Comor.
detonsa, *A. Gray, Pl. Wright.* ii. 24.—N. Mexic.
Douradinha, *A. St. Hil. Pl. Us. Bras.* t. 36.—Bras.
elliptica, Cav. Diss. vi. 316. t. 171. f. 2 = indica.
erioclada, DC. Prod. i. 493 = indica.
excelsa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 215.—Bras.
ferruginea, *A. St. Hil. Fl. Bras. Mer.* i. 150.—Bras.
fruticosa, *Rottb. in Act. Lit. Univ. Hafn.* i. (1778) 296.—Guian.
glabra, *Poir. Encyc.* viii. 325.—Ins. Guadeloup.
glabriuscula, A. St. Hil. Fl. Bras. Mer. i. 155 = communis.
glomerata, *Presl, Rel. Haenk.* ii. 152.—Panama.
gracilis, A. St. Hil. Fl. Bras. Mer. i. 154 = communis.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 295 = indica.
Haenkeana, D. Dietr. Syn. Pl. iv. 756 = Preslii.
hirsuta, *Presl, Rel. Haenk.* ii. 152.—Mexic.
indica, *Linn. Sp. Pl.* 673.—Cosmop. trop.
intricata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 214.—Cuba.
involucrata, *Benth. in Hook. Journ. Bot.* iv. (1842) 125.—Guian.
laevis, D. Wolf, ex Schrank, Pl. Rar. Hort. Monac. t. 55 = glabra.
lanata, A. St. Hil. Fl. Bras. Mer. i. 154 = communis.
lanceolata, *R. Br. ex Mast. in Oliver, Fl. Trop. Afr.* i. 235.—Afr. trop.
lantanaefolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 38.—Bras.
longifolia, *DC. Prod.* i. 493.—Ind. occ.
longiramea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 571.—Ind. or.
lophantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.
Lophanthus, *Forst. f. Prod.* 47.—Ins. Marques.
macropoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 216.—Bras.
maritima, *A. St. Hil. Fl. Bras. Mer.* i. 153.—Bras.
microphylla, Cav. Diss. vi. 317. t. 170. f. 2 = indica.
microphylla, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 144 = Coldenia procumbens.
monogynia, *Vell. Fl. Flum.* 274; vii. t. 6.—Bras.
obtusa, Willd. ex Steud. Nom. ed. II. ii. 783 = indica.
ovata, Cav. Diss. vi. 317. t. 171. f. 1 = indica.
paniculata, *Benth. in Hook. Journ. Bot.* iv. (1842) 126.—Guian.

WALTHERIA:—
pauciflora, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 235 = indica.
pedunculata, Willd. ex Steud. Nom. ed. II. ii. 783 = indica.
pentagynia, *Vell. Fl. Flum.* 274; vii. t. 5.—Bras.
Preslii, *Walp. Rep.* i. 340.—Mexic.
pyrolaefolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 190.—Ins. Sandvic.
reticulata, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 231.—Ins. Galapag.
rotundifolia, Presl, Rel. Haenk. ii. 151 = Preslii.
rotundifolia, *Schrank, in Syll. Ratisb.* ii. (1828) 65.—Bras.
sericea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 214.—Ecuador.
surinamensis, *Turcz. l. c.* xxxvi. (1863) i. 571.—Guian.
terminalis, Vell. Fl. Flum. 275; vii. t. 7 = Turnera ulmifolia.
viscosissima, *A. St. Hil. Fl. Bras. Mer.* i. 150.—Bras.

WANGENHEIMIA, F. G. Dietr. Vollst. Lexik. Gaertn. x. 536 (1810) = **Gilibertia**, Ruiz & Pav. (Araliac.).
umbellata, F. G. Dietr. l. c. 537 = G. umbellata.

WANGENHEIMIA, Moench, Meth. 200 (1794).
GRAMINEAE, Benth. & Hook. f. iii. 1170.
disticha, *Moench, l. c.*—Hispan.
Lima, Trin. Fund. Agrost. 132 = disticha.

WARBURTONIA, F. Muell. Fragm. i. 229 t. 9 (1859) = **Hibbertia**, Andr. (Dillen.).
potentillina, F. Muell. l. c. 230. t. 9 = H. grossulariaefolia.

WARCZEWICZELLA, Reichb. f. in Bot. Zeit. x. (1852) 635 = **Zygopetalum**, Hook. (Orchid.).
amazonica, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 97 = Z. amazonicum.
aromatica, Reichb. f. in Walp. Ann. vi. 654 = Z. aromaticum.
candida, Reichb. f. in Bot. Zeit. x. (1852) 636 = Z. candidum.
cochlearis, Reichb. f. l. c. 714 = Z. cochleare.
discolor, Reichb. f. l. c. 636 = Z. discolor.
Lueddemanniana, Reichb. f. in Hamb. Gartenz. xvi. (1860) 179 = Z. Lueddemanniana.
marginata, Reichb. f. in Bot. Zeit. x. (1852) 636 = Z. marginatum.
picta, Reichb. f. in Gard. Chron. (1883) ii. 8 = Z. pictum.
velata, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 97 = Z. velatum.
vittata, Gard. Chron. (1878) ii. 627, sphalm. = praec.
Wailesiana, E. Morr. Belg. Hortic. xxviii. (1878) 183. t. 10. f. 1 = Z. Wailesianum.

WARCZEWITZIA, Skinner, in Lindl. & Paxt. Flow. Gard. i. (1850–51) 45 = **Catasetum**, Rich. (Orchid.).

WAREA, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 127 = **Biswarea**, Cogn. (Cucurb.).
tonglensis, C. B. Clarke, l. c. 129 = B. tonglensis.

WAREA, Nutt. in Journ. Acad. Philad. vii. (1834) 83. t. 10. *CRUCIFERAE*, Benth. & Hook. f. i. 80.
amplexifolia, *Nutt. l. c.*—Am. bor.
cuneifolia, *Nutt. l. c.* 84.—Am. bor.

WARIA, Aubl. Pl. Gui. i. 604. t. 243 (1775) = **Uvaria**, Linn. (Anonac.).
zeylanica, Aubl. l. c. 604 = Uvaria zeylanica.
zeylanica, Aubl. l. c. t. 243 = Unona discolor.

WARIONIA, Benth. & Coss. in Bull. Soc. Bot. Fr. xix. (1872) 165. *COMPOSITAE*, Benth. & Hook. f. ii. 474.
Saharae, *Benth. & Coss. l. c.* 166.—Afr. bor.

WARMINGIA, Engl. in Mart. Fl. Bras. xii. ii. 281 (1874) = **Spondias**, Linn. (Anacard.).
pauciflora, Engl. l. c. = S. purpurea.

WARMINGIA, Reichb. f. Otia Bot. Hamb. 87 (1881). *ORCHIDEAE*, Benth. & Hook. f. iii. 477.
Eugenii, *Reichb. f. l. c.*—Bras.

WARNERA, Mill. Gard. Dict. ed. VIII. (1768) = **Hydrastis**, Ellis (Ranunc.).
canadensis, Mill. l. c. = H. canadensis.
diphylla, Rafin. Med. Fl. i. 253 = H. canadensis.
tinctoria, Rafin. l. c. = H. canadensis.

WARREA, Lindl. Bot. Reg. (1843) Misc. 14. *ORCHIDEAE*, Benth. & Hook. f. iii. 545.
bahiensis, Hort. ex Reichb. f. in Bot. Zeit. x. (1852) 931 = Paradisanthus bahiensis.
bidentata, *Lindl. Bot. Reg.* (1844) *Misc.* 76; *et in Lindl. & Paxt. Flow. Gard.* i. (1850-51) 73.—N. Granat.
candida, Lindl. & Paxt. l. c. 32 = Zygopetalum candidum.
cinerea, Benth. in Journ. Linn. Soc. xviii. (1881) 321, sphalm. = Aganisia cyanea (in Addend.).
cyanea, Lindl. Bot. Reg. (1844) Misc. 2; (1845) t. 28 = Aganisia cyanea (in Addend.).
digitata, Lem. Illustr. Hortic. iii. (1856) Misc. 70; iv. (1857) t. 152 = Zygopetalum Wailesianum.
discolor, Lindl. in Journ. Hort. Soc. iv. (1849) 265 = Zygopetalum discolor.
Lindeniana, Henfrey, in Moore & Ayres, Gard. Mag. Bot. i. (1850) 177 = bidentata.
marginata, Reichb. f. in Bot. Zeit. x. (1852) 636 = Zygopetalum marginatum.
quadrata, Lindl. in Gard. Chron. (1853) 647 = Zygopetalum marginatum.
tricolor, *Lindl. Bot. Reg.* (1843) *Misc.* 14.—Bras.
unijugata, Regel, Ind. Sem. Hort. Petrop. (1855) 22 = tricolor.
Wailesiana, Lindl. in Journ. Hort. Soc. iv. (1849) 264 = Zygopetalum Wailesianum.

WARSCEWICZELLA, Reichb. f. Xenia Orch. i. 61, in obs. (1858) = WARCZEWICZELLA, Reichb. f. = **Zygopetalum**, Hook. (Orchid.).

WARSZEWICZELLA, Benth. & Hook. f. Gen. iii. 543 (1880) = praec.

WARSZEWICZIA, Klotzsch, in Monatsb. Akad. Berl. (1853) 496. *RUBIACEAE*, Benth. & Hook. f. ii. 48.
coccinea, *Klotzsch, l. c.* 497.—Am. trop.
macrophylla, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 72 = coccinea.
maynensis, Wedd. l. c. = coccinea.
peltata, *Wedd. l. c.* 73.—N. Granat.
Poeppigiana, Klotzsch, in Monatsb. Akad. Berl. (1853) 497 = coccinea.
pulcherrima, Klotzsch, l. c. = coccinea.
Schomburgkiana, Klotzsch, l. c. = coccinea.
splendens, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 72 = coccinea.

WARTHEMIA, Boiss. Diagn. Ser. I. vi. 74 (1845) = VARTHEMIA, DC. = **Iphiona**, Cass. (Compos.).

WARTMANNIA, Muell. Arg. in Linnaea, xxxiv. (1865-66) 218 = **Homalanthus**, A. Juss. (Euphorb.).
stillingiaefolia, Muell. Arg. l. c. 219 = H. stillingifolius.

WASHINGTONIA, H. Wendl. in Bot. Zeit. xxxvii. (1879) 68. *PALMAE*, Benth. & Hook. f. iii. 923.
filifera, *H. Wendl. l. c.*—Calif.
robusta, H. Wendl. in Berl. Garten-Zeit. ii. (1883) 198; et in Rev. Hortic. (1885) 401. f. 73 = filifera?

WASHINGTONIA, Winsl. in Calif. Farm. (Sept. 1854); ex Hook. Kew Journ. vii. (1855) 29 = **Sequoia**, Endl. (Conif.).
americana, Hort. ex Gord. Pinet. Suppl. 106 = S. gigantea.
californica, Winsl. in Calif. Farm. (Sept. 1854); ex Hook. Kew Journ. vii (1855) 29 = S. gigantea.
gigantea, Hort. ex Carr. Conif. ed. II. 217 = S. gigantea.

WATSONIA, Mill. Gard. Dict. ed. VII (1759). *IRIDEAE*, Benth. & Hook. f. iii. 705.
BEILIA, Eckl. Top. Verz. 43 (1827).
LOMENIA, Pourr. in Mém. Acad. Toul. iii. (1788) 74. t. 5.
MERIANA, Trew, Pl. Sel. Ehret. 11. t. 40 (1750).
NEUBERIA, Eckl. Top. Verz. 37 (1827).
VALSONICA, Scop. Ann. IV. Hist. Nat. 97 (1770).
aletroides, *Ker-Gawl. in Bot. Mag.* t. 533.—Afr. austr.
amoena, Pers. Syn. i. 43 = Gladiolus hirsutus.
angusta, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 230.—Afr. austr.
atrosanguinea, Klatt, in Linnaea, xxxii. (1863) 738 = angusta.
brevifolia, *Ker-Gawl. in Bot. Mag.* t. 601.—Afr. austr.
campanulata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 354 = Ixia columellaris.
coccinea, *Herb. ex Baker, in Journ. Linn. Soc.* xvi. (1877) 158.—Afr. austr.
compacta, Lodd. Bot. Cab. t. 1577 = Micranthus alopecuroideus.
cylindrica, *Baker, in Journ. Bot.* xiv. (1876) 336.—Madag.
densiflora, *Baker, l. c.*—Afr. austr.
dubia, Eckl. Top. Verz. 36 = W. Meriana.
erubescens, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 631 = humilis.
filifolia, E. Mey. ex Baker, in Journ. Linn. Soc. xvi. (1877) 159 = punctata.
fulgens, Pers. Syn. i. 42 = W. Meriana.
fulgida, Salisb. in Trans. Hort. Soc. i. (1812) 323 = angusta.
humilis, *Mill. Gard. Dict.* ed. VIII. n. 2.—Afr. austr.
humilis, Pers. Syn. i. 42 = Antholyza Merianella.
hyacinthoides, Pers. l. c. 43 = brevifolia.
iridifolia, Eckl. Top. Verz. 35, nomen (Quid?).—Afr. austr.
iridifolia, Ker-Gawl. in Bot. Mag. t. 600 = W. Meriana.
Jacquini, Pers. Syn. i. 42 = aletroides.
laccata, Ker-Gawl. in Bot. Mag. sub t. 631; Pers. Syn. i. 43 = humilis.
Lamarkii, *A. Dietr. Sp. Pl.* ii. 570.—Afr. austr.
litura, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 353 = W. Meriana.
longifolia, Steud. Nom. ed. II. ii. 784 = Gladiolus Ecklonii.
lucens, Pers. Syn. i. 42 = Antholyza lucidior.
lucidior, Eckl. Top. Verz. 36 = Antholyza lucidior.
maculata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 352 = humilis.
marginata, *Ker-Gawl. in Bot. Mag.* t. 608.—Afr. austr.
Meriana, *Mill. Gard. Dict.* ed. VIII. n. 1; *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 230.—Afr. austr.
Merianella, Eckl. Top. Verz. 36 = Gladiolus hirsutus.
minuta, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 353.—Afr. austr.
natalensis, Eckl. Top. Verz. 34 = Gladiolus psittacinus.
palustris, Pers. Syn. i. 43 = Dierama ensifolium.
pellucida, Eckl. Top. Verz. 36 = coccinea.
pilosa, Klatt, in Trans. S. Afr. Phil. Soc. iii. (1885) 200 = Antholyza Merianella.
plantaginea, Ker-Gawl. in Bot. Mag. t. 553 = Micranthus alopecuroideus.
pottbergensis, Eckl. Top. Verz. 36, nomen (Quid?).—Afr. austr.
praecox, Pers. Syn. i. 42 = Antholyza revoluta.
punctata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 229.—Afr. austr.
racemosa, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 354 = Ixia columellaris?
recurva, Pers. Syn. i. 43 = Antholyza revoluta.
retusa, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 354.—Afr. austr.
revoluta, Pers. Syn. i. 42 = Antholyza revoluta.
rosea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1804) 230.—Afr. austr.
roseo-alba, Ker-Gawl. in Bot. Mag. t. 537 = W. Meriana.
rubens, Ker-Gawl. l. c. t. 1072 = punctata.
spicata, Ker-Gawl. in Kon. & Sims, Ann. Bot. I. (1804) 229 = Micranthus alopecuroideus.
striata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 352 = rosea.
stricta, *Spreng. Nov. Prov.* 43.—Afr. austr.
strictiflora, *Ker-Gawl. in Bot. Mag.* t. 1406.—Afr. austr.
subulata, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 353 = punctata.

WATSONIA:—
tabularis, Eckl. Top. Verz. 35, nomen (Quid?).—Afr. austr.
tigrina, Eckl. l. c. 36, nomen (Quid?).—Afr. austr.
triticea, Spreng. Syst. i. 150 = punctata, Micranthus alopecuroideus.
tubulosa, Drège, ex Baker, in Journ. Linn. Soc. xvi. (1877) 158 = humilis.
tubulosa, Eckl. Top. Verz. 36 = marginata.
tubulosa, Pers. Syn. i. 42 = aletroides.

WATTAKAKA, Hassk. in Flora, xl. (1857) 99 = **Dregea**, E. Mey. (Asclep.).
pubescens, Miq. Fl. Ind. Bat. ii. 496.—Java.
viridiflora, Hassk. in Flora, xl. (1887) 99 = D. volubilis.

WEBBIA, DC. Prod. v. 72 (1836) = **Vernonia**, Schreb. (Compos.).
aristata, DC. l. c. 73 = V. natalensis.
elaeagnoides, DC. l. c. = V. Kraussii.
hirsuta, DC. l. c. = V. hirsuta.
nudicaulis, DC. l. c. = V. Dregeana.
oligocephala, DC. l. c. = V. Kraussii.
pinifolia, DC. l. c. 72 = V. pinifolia.
serratuloides, DC. l. c. = V. Perrottetii.
Smithiana, DC. l. c. = V. Smithiana.

WEBBIA, Sch. Bip. in Walp. Rep. ii. 970 (1843) = **Conyza**, Linn. (Compos.).
Kraussii, Sch. Bip. l. c. 971 = C. obscura.

WEBBIA, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 356 = **Hypericum**, Tourn.
canariensis, Webb & Berth. Phyt. Canar. i. 48 = H. canariense.
floribunda, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 356 = H. floribundum.
heterophylla, Spach, Hist. Vég. Phan. v. 409 = H. canariense.
platypetala, Spach, l. c. 410, sphalm. = seq.
platysepala, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 356 = H. canariense.

WEBERA, Cramer, Disp. Syst. 38 (1803) = **Plectronia**, Linn. (Rubiac.).

WEBERA, J. F. Gmel. Syst. 820 (1791) = **Bellucia**, Neck. (Melastom.).
quinquenervia, J. F. Gmel. l. c. = B. Aubletii.

WEBERA, Schreb. Gen. 794 (1791) = **Tarenna**, Gaertn. (Rubiac.).
asiatica, Bedd. Forest. Man. Bot. 133 = T. zeylanica.
attenuata, Hook. f. Fl. Brit. Ind. iii. 104.—China.
auriculata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 537 = Anomanthodia auriculata.
bispinosa, Kurz, For. Fl. Brit. Burma, ii. 49 = Randia longiflora.
campaniflora, Hook. f. Fl. Brit. Ind. iii. 106.—Ind. or.
canarica, Hook. f. l. c.—Ind. or.
cerifera, Moon, Cat. Pl. Ceyl. 19 = T. zeylanica.
coriacea, Wall. ex Steud. Nom. ed. II. ii. 784 = *grandifolia*.
corymbosa, Sm. ex Wight & Arn. Prod. 429 = Ixora parviflora.
corymbosa, Willd. Sp. Pl. i. 1224 = T. zeylanica.
costata, Hook. f. Fl. Brit. Ind. iii. 103.—Malaya.
Cumingiana, Vidal, Phan. Cuming. Philipp. 178.—Ins. Philipp.
cymosa, Willd. Sp. Pl. i. 1224 = Plectronia didyma.
Dallachiana, F. Muell. ex Benth. Fl. Austral. iii. 412.—Austral.
densiflora, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 536 = Randia densiflora.
disperma, Hook. f. Fl. Brit. Ind. iii. 102.—Reg. Himal.
expandens, F. Muell. Fragm. vi. 25.—Austral.
fasciculata, Kurz, For. Fl. Brit. Burma, ii. 49 = Randia fasciculata.
fragrans, Hook. f. Fl. Brit. Ind. iii. 103.—Malaya.
glomeriflora, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 311 = T. zeylanica.
grandifolia, Hook. f. Fl. Brit. Ind. iii. 105.—Malaya.
Helferi, Hook. f. l. c. 107.—Burma.
hispidula, Baker, in Journ. Bot. xx. (1882) 137.—Madag.

WEBERA :—
Kurzii, Hook. f. Fl. Brit. Ind. iii. 105.—Ins. Andam. et Nicobar.
lanceolata, Moon, Cat. Pl. Ceyl. 19 = Plectronia didyma.
longiflora, Kurz, For. Fl. Brit. Burma, ii. 48 = Randia longiflora.
longifolia, Hook. f. Fl. Brit. Ind. iii. 105.—Ins. Penang.
lucens, Hook. f. l. c. 106.—Ind. or.
luzoniensis, Vidal, Phan. Cuming. Philipp. 179.—Ins. Philipp.
macrophylla, Roxb. Hort. Beng. 85; Fl. Ind. i. 697 = *odorata*.
Maingayi, Hook. f. Fl. Brit. Ind. iii. 103.—Malacca.
mollis, Hook. f. l. c. 104.—Malaya.
mollissima, Benth. ex Hance, in Journ. Linn. Soc. xiii. (1873) 105.—China.
monosperma, Hook. f. Fl. Brit. Ind. iii. 103.—Ind. or.
myrtifolia, Kurz, For. Fl. Brit. Burma, ii. 49.—Burma.
nilagirica, Hook. f. Fl. Brit. Ind. iii. 106.—Ind. or.
odorata, Roxb. Hort. Beng. 15; Fl. Ind. i. 699.—Reg. Himal.; Penang.
oppositifolia, Roxb. ll. cc. 85; 698 = Randia densiflora.
pavettoides, Benth. & Hook. f. Gen. ii. 95, in nota sub Kraussia.—Afr. trop.
pumila, Hook. f. Fl. Brit. Ind. iii. 103.—Reg. Himal.
scandens, Roxb. Hort. Beng. 15; Fl. Ind. i. 698 = Randia longiflora.
sechellensis, Baker, Fl. Maurit. 139.—Ins. Seychelles.
siamensis, Kurz, For. Fl. Brit. Burma, ii. 48 = Randia longiflora.
stellulata, Hook. f. Fl. Brit. Ind. iii. 104.—Malaya.
subsessilis, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 166.—Ins. Bonin.
sumatrana, Boerl. in Veth, Midden-Sumatra, iv. II. Flora, 21.—Sumatra.
tetrandra, Wall. Cat. n. 8265 = Plectronia Rheedii.
tetrandra, Willd. Sp. Pl. i. 1224 = Plectronia parviflora.
thyrsoidea, Roth, Nov. Pl. Sp. 149 = Wendlandia Notoniana.
truncata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 538 = Trisciadia truncata.
Wallichii, Hook. f. Fl. Brit. Ind. iii. 105.—Ins. Penang.

WEDDELLINA, Tul. in Ann. Sc. Nat. Sér. III. xi (1849) 113. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 109.
squamulosa, *Tul. l. c.*—Guian.
uaupensis, *Benth. in Benth. & Hook. f. Gen.* iii. 109.—Bras.

WEDELA, Steud. Nom. ed. II. ii. 784 (1841), sphalm. = VEDELA, Adans. = **Ardisia**, Sw. (Myrsin.).

WEDELIA, Jacq. Enum. Pl. Carib. 8 (1760); Select. Am. 217 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 370.
ANTHEMIOPSIS, Boj. ex DC. Prod. v. 547, 548 (1836).
MENOTRICHE, Steetz, in Peters, Reise Mossamb. Bot. 472 (1863).
NIEBUHRIA, Neck. Elem. i. 30 (1790).
STEMMODONTIA, Cass. in Bull. Soc. Philom. (1817) 11.
TRICHOSTEMMA, Cass. in Dict. Sc. Nat. xlvi. (1827) 399, 409.
TRICHOSTEPHIUM, Cass. l. c. lv. 266, in obs. (1828).
TRICHOSTEPHUS, Cass. l. c. lx. 618 (1830).
WOLLASTONIA, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 414 (Herb. Tim. Desc. 86).
abyssinica, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 327.—Afr. trop.
acapulcensis, *H. B. & K. Nov. Gen. et Sp.* iv. 215.—Mexic.
acuminata, *DC. Prod.* v. 541.—Cuba.
affinis, *DC. l. c.*—Ins. S. Thomas (Ind. occ.).
africana, *Beauv. Fl. Owar.* ii. 19. t. 69.—Afr. trop.
alagoënsis, *Baker, in Mart. Fl. Bras.* vi. III. 187.—Bras.
alutacea, Pohl, ex Baker, l. c. 200 = Aspilia asperrima.

WEDELIA:—
ambigua, *DC. Prod.* v. 541.—Ins. Guadaloup.
aristata, *Less. in Linnaea*, vi. (1831) 160.—Ins. Molucc.
asperrima, *Benth. Fl. Austral.* iii. 539.—Austral.
aurea, D. Don, ex Hook. Bot. Mag. t. 3384 = Zexmenia aurea.
bengalensis, Rich. in Pers. Syn. ii. 490 = calendulacea.
biflora, C. B. Clarke, Comp. Ind. 137 = Wallichii.
biflora, *DC. in Wight, Contrib.* 18.—Geront. trop.
brachycarpa, *Baker, in Mart. Fl. Bras.* vi. III. 181.—Paraguay.
brachylepis, *Griseb. in Goett. Abh.* xxiv. (1879) 190.—Reg. Argent.
buphtalmiflora, *Lor. in Roca, Exped. Rio-Negro*, ii. 237.—Patagon.
buphthalmoides, *Griseb. in Goett. Abh.* vii. (1857) 235.—Ind. occ.
calendulacea, *Less. Syn. Comp.* 222.—As. et Austral. trop.
calendulacea, Rich. in Pers. Syn. ii. 490 = hispida.
calycina, Rich. l. c. = affinis *vel* acapulcensis.
calycina, Spreng. Syst. iii. 580 = ambigua
caracasana, DC. Prod. v. 541 = Zexmenia caracasana.
caribaea, Spreng. Syst. iii. 580 = cruciana.
carnosa, *Pers. Syn.* ii. 490.—Am. trop.
Chamissonis, *Less. in Linnaea*, vi. (1831) 161.—Ins. Guahan.
chrysostephana, *Griseb. in Goett. Abh.* xxiv. (1879) 190.—Reg. Argent.
cordata, Hook. & Arn. Bot. Beech. Voy. 435 = Gymnolomia patens.
cordifolia, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 261 = Calea hispida.
crenata, Rich. in Pers. Syn. ii. 490 = carnosa.
cruciana, *Rich. in Pers. l. c.*—Ind. occ.
Cunninghami, DC. Prod. v. 540 = urticifolia.
discoidea, Less. in Linnaea, vi. (1831) 728 = Eleutheranthera ovata.
ecliptoides, F. Muell. Pl. Vict. sub t. 39 = Eclipta platyglossa.
Ehrenbergii, *Less. in Linnaea*, vi. (1831) 729.—Ins. S. Doming.
elongata, *Vatke, in Bremen, Abh.* ix. (1885) 121.—Madag.
equatorialis, *Benth. & Hook. f. Gen.* ii. 350, in obs.—Am. trop.
filipes, *Hemsl. Biol. Centr. Am. Bot.* ii. 170.—Guatemala.
Forsteriana, *Endl. Prod. Fl. Norf.* 51.—Ins. Norfolk.
fructicosa, Jacq. Enum. Pl. Carib. 28 = frutescens.
frutescens, *Hook. f. ex Benth. in Journ. Linn. Soc.* xiii. (1873) 439 (an tenuicaulis?).—Ins. Galapag.
frutescens, *Jacq. Select. Am.* 217. t. 130.—N. Granat.
frutescens, Sieber, ex Griseb. Fl. Brit. W. Ind. 372 = Sieberi.
frutescens, Spreng. ex DC. Prod. v. 541 = reticulata.
Gaudichaudi, *DC. Prod.* v. 542.—Peruv.
goyazensis, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 288.—Bras.
gracilis, *Rich. in Pers. Syn.* ii. 490.—Ind. occ.
grandiflora, *Benth. Bot. Voy. Sulph.* 115.—Ecuador.
helianthoides, H. B. & K. Nov. Gen. et Sp. iv. 216 = Zexmenia helianthoides, *Benth. & Hook. f.*
hispida, *H. B. & K. l. c.* 215.—Mexic.
Hookeriana, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 289.—Bras.
Horsfieldiana, *Benth. & Hook. f. Gen.* ii. 371.—Java.
humilis, *Schweigg. Enum. Pl. Hort. Regiom.* 79.—Hab.?
Jacquini, Rich. in Pers. Syn. ii. 490 = reticulata.
laevissima, Less. ex Baker, in Mart. Fl. Bras. vi. III. 202 = Aspilia laevissima.
lanceolata, *DC. Prod.* v. 541.—Ins. Porto Rico.
latifolia, *DC. l. c.* 542.—Am. austr.
linearifolia, *Baker, in Mart. Fl. Bras.* vi. III. 182.—Bras.
linearis, *Zoll. in Flora*, xxx. (1847) 534.—Java.
longifolia, *Mart. ex Baker, in Mart. Fl. Bras.* vi. III. 182.—Bras.
Lundii, *DC. Prod.* v. 539.—Bras.
macrodonta, *DC. l. c.*—Bras.
Menotriche, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 377.—Afr. trop.
minor, Hornem. Hort. Hafn. 855 = Melampodium paludosum.
modesta, *Baker, in Mart. Fl. Bras.* vi. III. 185.—Bras.
mollis, *Pers. Syn.* ii. 490.—Am. austr.

WEDELIA:—
mossambicensis, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 97.—Afr. trop.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 63.—Afr. austr.
oblongifolia, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 256 = Calea Candolleana.
oligocephala, *Baker, l. c.* 182.—Bras.
ovatifolia, Willd. Enum. Hort. Berol. Suppl. 61 = Melampodium paludosum.
pallida, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 288 = macrodonta.
paludicola, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 50.—Cuba.
paludosa, *DC. Prod.* v. 538.—Bras.
paniculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 69.—Ecuador.
parviflora, *Rich. in Pers. Syn.* ii. 490.—Ins. Guadaloup.
pedunculosa, DC. Prod. v. 484, sphalm. = paludosa.
perfoliata, Willd. Sp. Pl. iii. 2335 = Melampodium perfoliatum.
phyllocephala, *Hemsl. Biol. Centr. Am. Bot.* ii. 170.—Guatemala.
pilosa, *Baker, in Mart. Fl. Bras.* vi. III. 181.—Bras.
Pohlii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 183 = Vauthieri.
populifolia, *Hook. & Arn. Bot. Beech. Voy.* 435.—Nicaragua.
pratensis, *Vatke, in Bremen, Abh.* ix. (1885) 121.—Madag.
psammophila, *Poepp. et Endl. Nov. Gen. et Sp.* iii. 50.—Bras.
puberula, *DC. Prod.* v. 540.—Bras.
pulchella, *H. B. & K. Nov. Gen. et Sp.* iv. 214.—Am. austr.
pumila, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 194 = Aspilia procumbens.
radiosa, *Ker-Gawl. in Bot. Reg.* t. 610.—Bras.
reticulata, *DC. Prod.* v. 540.—Ins. Porto Rico.
reticulata, Griseb. Cat. Pl. Cub. 154 = Zexmenia reticulata.
rudis, Benth. ex Baker, in Mart. Fl. Bras. vi. III. 189 = Zexmenia rudis.
scaberrima, *Benth. in Ann. Nat. Hist.* Ser. I. ii. (1839) 110.—Guian.
scaberrima, Hook. f. Fl. Brit. Ind. iii. 307, sphalm., in syn. = Wallichii.
scaberrima, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 255 = Calea elongata.
scandens, C. B. Clarke, Comp. Ind. 136 (excl. syn.) = biflora.
scandens, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 125 = subvelutina.
Schultziana, *Miq. Fl. Ind. Bat.* ii. 69.—Ins. Bali.
Serra, Pohl, ex Baker, in Mart. Fl. Bras. vi. III. 184 = macrodonta.
serrata, *Rich. in Pers. Syn.* ii. 490.—Ins. S. Doming.
sessilifolia, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 196 = Aspilia reflexa.
sessilifolia, Pohl, ex Baker, l. c. 180 = paludosa.
Sieberi, *Griseb. Fl. Brit. W. Ind.* 372, in obs.—Ind. occ.
spilanthoides, *F. Muell. Fragm.* v. 64.—Austral.
strigosa, Hook. & Arn. Bot. Beech. Voy. 435 = Aspilia strigosa.
subflexuosa, Hook. & Arn. l. c. = Gymnolomia subflexuosa.
subvelutina, *DC. Prod.* v. 540.—Bras.
tenuicaulis, Hook. f. in Trans. Linn. Soc. xx. (1847) 213 = Blainvillea tenuicaulis.
titicacensis, Meyen & Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 269 = Villanova titicacensis.
Trichostemma, Steud. Nom. ed. II. ii. 703, in syn., sphalm. = seq.
Trichostephia, *DC. Prod.* v. 540.—Bras.
urticaefolia, *DC. in Wight, Contrib.* 18.—Ind. or.
Vauthieri, *DC. Prod.* v. 539.—Bras.
verbesinoides, *F. Muell. ex Benth. Fl. Austral.* iii. 538.—Austral.
veronicaefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 53.—Cuba.
villosa, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 289, partim.—Bras.
villosa, Gardn. l. c., partim = alagoënsis.
Wallichii, *Less. in Linnaea*, vi. (1831) 162.—Reg. Himal.; Java.
Zollingeriana, Sch. Bip. ex C. B. Clarke, Comp. Ind. 136 = calendulacea.

WEDELIA, Loefl. It. Hisp. 180 (1758) = **Allionia**, Loefl. (Nyctag.).
incarnata, Linn. Syst. ed. X. 890.—Am. bor. et austr.
malachroides, Benth. Bot. Voy. Sulph. 114 = incarnata.
violacea, Linn. Syst. ed. X. 890.—Venezuela.

WEDELIOPSIS, Planch. ex Benth. in Hook. Niger Fl. 351 (1849) = **Dissotis**, Benth. (Melastom.).

WEHLIA, F. Muell. Fragm. x. 22 (1876). *MYRTACEAE.*
coarctata, *F. Muell. l. c.* 23.—Austral.
staminosa, *F. Muell. in Wing, South. Sc. Record*, iii. (1884) 282.—Austral.
thryptomenoides, *F. Muell. Fragm.* x. 22.—Austral.

WEIGELA, Thunb. in Vet. Acad. Handl. Stockh. (1780) 137. t. 5 = **Diervilla**, Tourn. (Caprifol.).
alba, Carr. in Rev. Hortic. (1861) 331 = D. florida.
amabilis, Hort. ex Carr. l. c. Sér. IV. ii. (1853) 305 = D. florida.
biformis, Hort. ex C. Koch, Dendrol. ii. 1. 42 = D. coraeënsis.
candida, Carr. in Rev. Hortic. (1879) 130.—Hab. ?
coraeënsis, Thunb. in Trans. Linn. Soc. ii. (1794) 331 = D. coraeënsis.
Desboisii, Carr. in Rev. Hortic. (1861) 332 = D. coraeënsis.
excelsa, Carr. l. c. (1873) 279.—Hab. ?
fallax, Miq. Fl. Ind. Bat. ii. 128.—Java.
floribunda, C. A. Mey. in Bull. Phys.-Math. Pétersb. xiii. (1855) 219 = D. floribunda.
florida, A. DC. in Ann. Sc. Nat. Sér. II. xi. (1839) 241 = D. florida.
Groenewegenii, Carr. in Rev. Hortic. (1861) 332 = D. coraeënsis.
hortensis, C. A. Mey. in Bull. Phys.-Math. Pétersb. xiii. (1855) 219 = D. hortensis.
intermedia, Hort. ex C. Koch, Dendrol. ii. 1. 42 = D. coraeënsis.
Isolinae, Van Houtte, in Fl. des Serres, xiv. (1861) t. 1445 = D. rosea.
japonica, Thunb. Fl. Jap. 90 = D. japonica.
Lavallei ×, Lescuyer, in Hérincq, L'Hortic. Franç. (1870) 145. [t. 5].
Lowii ×, Hort. ex Hérincq, l. c. 232.
Metelerkampi, Hort. ex C. Koch, Dendrol. ii. 1. 42 = D. coraeënsis.
Middendorfiana, Hort. ex Carr. in Rev. Hortic. Sér. IV. ii. (1853) 306 = D. Middendorfiana.
pauciflora, A. DC. in Ann. Sc. Nat. Sér. II. xi. (1839) 241 = D. florida.
rosea, Lindl. in Journ. Hort. Soc. i. (1846) 65, 189. t. 6 = D. florida.
splendens, Hort. ex Carr. in Fl. des Serres, viii. (1852–53) 292 = D. sessilifolia.
Stelzneri, Van Houtte, l. c. xiv. (1861) 142 = D. rosea ?

WEIGELIA, Pers. Syn. i. 176 (1805) = praec.

WEIGELTIA, A. DC. in Trans. Linn. Soc. xvii. (1834) 102 = **Cybianthus**, Mart. (Myrsin.).
Blanchetii, A. DC. in DC. Prod. viii. 115.—Bras.
capitellata, Miq. in Mart. Fl. Bras. x. 300 = C. myrianthos.
detergens, Mart. Pl. Med. Bras. Ined. t. 43 = C. detergens.
Gardneri, A. DC. in DC. Prod. viii. 115.—Bras.
longifolia, Benth. ex Miq. in Mart. Fl. Bras. x. 299 = C. longifolius.
myrianthos, A. DC. in Trans. Linn. Soc. xvii. (1834) 102 = C. myrianthos.
obovata, Mart. in Flora, xxiv. (1841) II. Beibl. 21 = C. obovatus.

WEIGELTIA, Reichb. Consp. 155 (1828), nomen. *LEGUMINOSAE.*

WEIHEA, Eckl. Top. Verz. 22 (1827) = **Geissorhiza**, Ker-Gawl. (Irid.).
elatior, Eckl. l. c. = G. excisa.
excisa, Eckl. l. c. = G. excisa.

WEIHEA, Reichb. Consp. 212[b] (1828) = **Burtonia**, R. Br. (Legumin.).

WEIHEA, Spreng. Syst. ii. 559 (1825). *RHIZOPHOREAE*, Benth. & Hook. f. i. 681.
ANSTRUTHERIA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 344. t. 4.
RICHAEIA, Thou. Gen. Nov. Madag. 25 (1806).
africana, *Benth. ex Oliver, Fl. Trop. Afr.* ii. 410.—Afr. trop.
Afzelii, *Oliver, l. c.* 411.—Afr. trop.
ceylanica, Hensl. in Hook. f. Fl. Brit. Ind. ii. 441 = zeylanica.
lanceolata, *Baill. Adansonia*, iii. (1862–63) 39.—Ins. Comor.
leptoclada, *Baill. l. c.*—Madag.
madagascariensis, *Spreng. Syst.* ii. 594.—Madag.
microphylla, *Baill. Adansonia*, iii. (1862–63) 39.—Madag.
myriocarpa, *Baill. l. c.*—Madag.
ovata, *Baill. l. c.*—Ins. Comor.
phoeotricha, *Baill. l. c.*—Madag.
plumosa, *Oliver, Fl. Trop. Afr.* ii. 411.—Afr. trop.
sessiliflora, *Baker, in Journ. Linn. Soc.* xxi. (1884) 341.—Madag.
zeylanica, *Baill. Adansonia*, iii. (1862–63) 38.—Zeylan.

WEILBACHIA, Klotzsch & Oerst. in Abh. Akad. Berl. 1854 (1855) 119. t. 11 A = **Begonia**, Tourn.
pustulata, Klotzsch, l. c. 121 = B. pustulata.
reptans, Klotzsch & Oerst. l. c. 120 = B. reptans.

WEINGAERTNERIA, Bernh. Syst. Verz. Erf. 23, 51 (1800) = **Corynephorus**, Beauv. (1812) (Gramin.).
canescens, Bernh. l. c. 51 = C. canescens.

WEINGARTNERIA, Benth. in Journ. Linn. Soc. xix. (1881) 95, sphalm. = praec.

WEINMANNIA, Linn. Syst. ed. X. 1005 (1759). *SAXIFRAGEAE*, Benth. & Hook. f. i. 653.
ARNOLDIA, Blume, Bijdr. 868 (1826).
LEIOSPERMUM, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 91.
PTEROPHYLLA, D. Don, l. c. 93 (1830).
WINDMANNIA, P. Br. Hist. Jamaic. 212 (1756).
affinis, *A. Gray, Bot. U. St. Expl. Exped.* i. 674.—Ins. Fiji.
alnifolia, Dombey, ex Engl. in Linnaea, xxxvi. (1869–70) 597 = heterophylla.
alpestris, Macfad. Fl. Jamaic. ii. 183 = pinnata.
arborea, Sm. in Rees, Cycl. xxxviii. sub n. 3 = tinctoria.
Arnoldia, A. Gray, Bot. U. St. Expl. Exped. i. 676, in obs. = Blumei.
auriculata, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 87 = dryadifolia.
australis, A. Cunn. in Field, N. S. Wales, 353 = Acrophyllum venosum.
Balbisiana, *H. B. & K. Nov. Gen. et Sp.* vi. 51.—Peruv.
Benthami, F. Muell. Fragm. v. 180 = Geissois Benthami.
betulina, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 357 = sylvicola.
Biagiana, *F. Muell. Fragm.* v. 16.—Austral.
bifida, *Poepp. ex Engl. in Linnaea*, xxxvi. (1869–70) 608.—Peruv.
biglandulosa, A. Cunn. in Hook. Ic. Pl. t. 301 = Anodopetalum biglandulosum.
Blumei, A. Gray, Bot. U. St. Expl. Exped. i. 676, in obs. = sundana.
Blumei, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 470.—Java.
Boiviniana, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 154 (*Biviniana*) = tinctoria.
Bojeriana, *Tul. l. c.* 155.—Madag.
brachystachya, Willd. ex Engl. in Linnaea, xxxvi. (1869–70) 606 = microphylla.
caripensis, H. B. & K. Nov. Gen. et Sp. vi. 58 = pinnata.
chilensis, DC. Prod. iv. 665 = Caldcluvia paniculata.
cinerea, *Ruiz & Pav. Fl. Per.* iv. t. 332.—Peruv.
comorensis, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 153.—Ins. Comor.
cordata, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 87.—Am. austr.
corymbosa, Ruiz & Pav. Fl. Per. iv. t. 330 = Caldcluvia paniculata.
crassifolia, *Ruiz & Pav. l. c.* t. 331.—Am. austr.
crenata, *Presl, Rel. Haenk.* ii. 51.—Peruv.

WEINMANNIA:—
decora, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 154.—Madag.
Denhami, *Seem. Fl. Vit.* 109.—N. Hebrid.
dentata, Ruiz & Pav. Fl. Per. iv. t. 334 = trichosperma.
dichotoma, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 73.—N. Caled.
dioica, Spreng. f. Tent. Suppl. Syst. 13 = Hippobromus alatus.
discolor, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 104.—Bras.
diversifolia, Moric. ex Ser. in DC. Prod. iv. 10 = cinerea ?
dryadifolia, *Moric. ex Ser. l. c.* 9.—Peruv.
elliptica, *H. B. & K. Nov. Gen. et Sp.* vi. 50.—Peruv.
eriocarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 156.—Madag.
fagaroides, *H. B. & K. Nov. Gen. et Sp.* vi. 54.—Peruv.
floribunda, *Baker, in Journ. Linn. Soc.* xx. (1883) 138.—Madag.
fraxinea, *Sm. ex D. Don, in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 93.—Ins. Molucc.
fraxinifolia, *Baker, in Journ. Linn. Soc.* xxi. (1884) 339.—Madag.
fraxinifolia, *Miq. Fl. Ind. Bat.* i. 1. 1095.—Java.
fuchsioides, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 357 = sylvicola.
glabra, Lam. Encyc. vii. 578 = tinctoria.
glabra, Linn. f. Suppl. 228 = pinnata.
glabra, Sieber, ex Engl. in Linnaea, xxxvi. (1869–70) 618 = hirta.
glomerata, *Presl, Rel. Haenk.* ii. 52.—Peruv.
guyanensis, *Klotzsch, ex Engl. in Linnaea*, xxxvi. (1869–70) 605.—Guian.
Haenkeana, *Engl. l. c.* 632.—Peruv.
heterophylla, *H. B. & K. Nov. Gen. et Sp.* vi. 53.—N. Granat.
heterophylla, Ruiz & Pav. Fl. Per. iv. t. 331 = cordata.
Hildebrantii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 475.—Madag.
hirsuta, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 89, sphalm. = seq.
hirta, *Sw. Prod. Veg. Ind. Occ.* 63.—Jamaic.
hirtella, *H. B. & K. Nov. Gen. et Sp.* vi. 56.—N. Granat.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. 1. 718.—Java.
Humblotii, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 475.—Madag.
humilis, *Engl. in Linnaea*, xxxvi. (1869–70) 620.—Bras.
intermedia, *Schlecht. & Cham. in Linnaea*, v. (1830) 555.—Mexic.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 280.—Bras.
Kunthiana, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 86 = ovata.
lachnocarpa, *F. Muell. Fragm.* viii. 7.—Austral.
Lansbergiana, *Engl. in Linnaea*, xxxvi. (1869–70) 628.—Venezuela.
Lantziana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 472.—Madag.
latifolia, *Presl, Rel. Haenk.* ii. 51.—Peruv.
latifolia, Willd. ex Engl. in Linnaea, xxxvi. (1869–70) 602 = macrophylla.
laurina, *H. B. & K. Nov. Gen. et Sp.* vi. 51.—Am. austr.
Lechleriana, *Engl. in Linnaea*, xxxvi. (1869–70) 630.—Peruv.
lentiscifolia, *Presl, Rel. Haenk.* ii. 52.—Peruv.
lucens, *Baker, in Journ. Bot.* xx. (1882) 70.—Madag.
Macgillivrayi, *Seem. Fl. Vit.* 109.—N. Hebrid.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* vi. 52.—Ecuador.
macrostachya, DC. ex Ser. in DC. Prod. iv. 10 = tinctoria.
madagascariensis, DC. ex Ser. l. c. 9 = mauritiana.
marginata, *Moric. ex Ser. l. c.* 10.—Peruv.
mauritiana, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 88.—Ins. Maurit.
microphylla, *H. B. & K. Nov. Gen. et Sp.* vi. 54.—Peruv.
microphylla, Ruiz & Pav. Fl. Per. iv. t. 334 = parvifolia.
minutiflora, *Baker, in Journ. Linn. Soc.* xxi. (1884) 339.—Madag.
nitida, D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 89 = hirta.

WEINMANNIA:—
oppositifolia, Willd. ex Engl. in Linnaea, xxxvi. (1869–70) 600 = ovata.
organensis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 104.—Bras.
ovalis, DC. ex Engl. in Linnaea, xxxvi. (1869–70) 600 = ovata.
ovalis, Ruiz & Pav. Fl. Per. iv. t. 333 = elliptica.
ovata, *Cav. Ic.* vi. 45. t. 566.—Peruv.
ovata, Ruiz & Pav. Fl. Per. iv. t. 333 = dryadifolia.
paniculata, Cav. Ic. vi. 44 = Caldcluvia paniculata.
paniculata, F. Muell. Fragm. ii. 83 = Ackama Muelleri.
paniculosa, F. Muell. l. c. 175 = Ackama Muelleri.
parviflora, *Forst. f. Prod.* 26.—Ins. Tahiti.
parviflora, Pers. Syn. i. 438 = Clethra guianensis.
parvifolia, *Ruiz, ex D. Don, in Edinb. N. Phil. Journ.* (Apr.–June, 1830) 89.—Peruv.
paulliniaefolia, *Pohl, ex Ser. in DC. Prod.* iv. 11.—Bras.
pentaphylla, *Ruiz & Pav. Fl. Per.* iv. t. 330.—Peruv.
pinnata, *Linn. Syst.* ed. X. 1005.—Am. trop.
pinnata, Ruiz, ex Engl. in Linnaea, xxxvi. (1869–70) 611 = cinerea.
polyphylla, *Moric. ex Ser. in DC. Prod.* iv. 11.—Peruv.
polystachia, Ruiz, ex D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 90 = subsessiliflora.
producta, *Moric. ex Ser. in DC. Prod.* iv. 11.—Peruv.
pubescens, *H. B. & K. Nov. Gen. et Sp.* vi. 56.—Venezuela.
pubescens, Ruiz, ex D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 89 = reticulata.
racemosa, *Linn. f. Suppl.* 227; *Forst. f. Prod.* 29.—N. Zel.
reticulata, *Ruiz & Pav. Fl. Per.* iv. t. 332.—Peruv.
rhodoxylon, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 152.—Ins. Marian.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 675. t. 85.—Ins. Fiji.
Riviniana, Engl. in Linnaea, xxxvi. (1869–70) 635, sphalm. = tinctoria.
rosaefolia, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 358; A. Gray, Bot. U. St. Expl. Exped. i. 671 = Ackama rosaefolia.
rubifolia, *Benth. Fl. Austral.* iii. 445.—Austral.
rugosa, Benth. Pl. Hartw. 132 = latifolia.
Rutenbergii, *Engl. in Bremen, Abh.* vii. (1880) 16.—Madag.
sambucina, Ruiz, ex D. Don, in Edinb. N. Phil. Journ. (Apr.–June, 1830) 90 = cinerea.
samoënsis, *A. Gray, Bot. U. St. Expl. Exped.* i. 677.—Ins. Pacif.
serrata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 73.—N. Caled.
sessiliflora, Pav. ex Ser. in DC. Prod. iv. 11 = producta.
sorbifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 57.—N. Granat.
spatiosa, Hook. f. Fl. N. Zel. i. 80, err. typ. = seq.
speciosa, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 358 = racemosa.
spiraeoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 677.—Ins. Fiji.
Spruceana, *Engl. in Linnaea*, xxxvi. (1869–70) 607.—Ecuador.
subcordata, *Moric. ex Ser. in DC. Prod.* iv. 8.—Peruv.
subsessiliflora, *Ruiz & Pav. Fl. Per.* iv. t. 334.—Am. austr.
sulcata, *Engl. in Linnaea*, xxxvi. (1869–70) 624.—Ecuador.
sundana, *Miq. Fl. Ind. Bat.* i. 1. 718.—Java.
sylvicola, *Soland. ex A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839) 357.—N. Zel.
ternata, *Engl. in Linnaea*, xxxvi. (1869–70) 603.—Peruv.
tinctoria, *Sm. in Rees, Cyclop.* xxxviii. n. 2.—Ins. Mascar.
tomentosa, *Linn. f. Suppl.* 227.—N. Granat.
Trianaea, *Wedd. Chlor. And.* ii. 211. t. 72.—N. Granat.
trichosperma, *Cav. Ic.* vi. 45. t. 567.—Peruv.; Chili.
trifoliata, Lam. Encyc. vii. 579 = mauritiana.
trifoliata, Linn. f. Suppl. 227 = Platylophus trifoliatus.
trigyna, *Baker, in Journ. Bot.* xx. (1882) 109.—Madag.
venosa, Knowles & Westc. Flor. Cab. ii. (1838) 81 = Acrophyllum venosum.
vitiensis, *Seem. Fl. Vit.* 110.—Ins. Fiji.

WEINREICHIA, Reichb. Consp. 152 (1828) = **Pterocarpus**, Linn. (Legumin.).

WEITENWEBERA, Opiz, Seznam, 104 (1852) = **Campanula**, Tourn.
Cervicaria, Opiz, l. c. 105 = C. Cervicaria.
glomerata, Opiz, l. c. = C. glomerata.

WELDENIA, Schult. f. in Flora, xii. (1829) 3. t. 1 A. *COMMELINACEAE*, Benth. & Hook. f. iii. 856.
LAMPRA, Benth. Pl. Hartw. 95 (1842).
RUGENDASIA, Schiede, ex Schlecht. Hort. Halens. 14 (1841).
candida, *Schult. f. in Flora*, xii. (1829) 3. t. 1 A.—Mexic.
Schultesii, Schlecht. Hort. Halens. 14 = candida.

WELEZIA, Neck. Elem. iii. 394 (1790), sphalm. = **Velezia**, Linn. (Caryoph.).

WELFIA, H. Wendl.—Cf. Gard. Chron. (1869) 1236; et ex André, Illustr. Hortic. xviii. (1871) 93. t. 62. *PALMAE*, Benth. & Hook. f. iii. 915.
Georgii, *H. Wendl. ex André, l. c.* 94.—Costa Rica.
regia, *H. Wendl.—Cf. Gard. Chron.* (1869) 1236; *et ex André, l. c.* 93. t. 62.—N. Granat.

WELLINGTONIA, Lindl. in Gard. Chron. (1853) 819, 823 = **Sequoia**, Endl. (Conif.).
californica, Sargent, Rep. For. N. Am. (U. St. Census, ix.) 184 = S. gigantea.
gigantea, Lindl. in Gard. Chron. (1853) 819, 823 = S. gigantea.

WELLINGTONIA, Meissn. Gen. Comm. 207, in nota (1840) = **Meliosma**, Blume (Sabiac.).
Arnottiana, Meissn. l. c. = M. Arnottiana.

WELLSTEDIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 407. *BORAGINEAE*.
socotrana, *Balf. f. l. c.* 408.—Ins. Socotra.

WELWITSCHIA, Hook. f. in Gard. Chron. (1862) 71; et in Trans. Linn. Soc. xxiv. (1863) 6. tt. 1–14. *GNETACEAE*, Benth. & Hook. f. iii. 418.
TOUMBOA, Naud. in Rev. Hortic. (1862) 186.
TUMBOA, Welw.—Cf. Gard. Chron. (1861) 75, nomen prius.
Bainesii, Carr. Conif. ed. II. 783 = mirabilis.
mirabilis, *Hook. f. in Trans. Linn. Soc.* xxiv. (1863) 7.—Afr. trop.

WELWITSCHIA, Reichb. Handb. 194 (1837) = **Gilia**, Ruiz & Pav. (Polemon.).

WENDEROTHIA, Schlecht. in Linnaea, xii. (1838) 330 = **Canavalia**, DC. (Legumin.).
discolor, Schlecht. l. c. 331 = C. villosa.
glabra, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 193.—Mexic.
hirsuta, Mart. & Gal. l. c. 192.—Mexic.
pilosa, Mart. & Gal. l. c. 191.—Mexic.

WENDIA, Hoffm. Gen. Pl. Umbell. ed. I. 136 (1814) = **Heracleum**, Linn. (Umbell.).
alpina, Schur, Enum. Pl. Transs. 268 = H. alpinum.
austriaca, Schur, l. c. = H. austriacum.
Chorodanum, Hoffm. Gen. Pl. Umbell. ed. I. 139. t. 1 = H. Chorodanum.
siifolia, Schur, Enum. Pl. Transs. 268 = H. austriacum.

WENDLANDIA, Bartl. Ord. 211 (1830); et ex DC. Prod. iv. 411. *RUBIACEAE*, Benth. & Hook. f. ii. 50.
SESTINIA, Boiss. & Hohen. Diagn. Ser. I. iii. 55 (1843).
angustifolia, *Wight, ex Hook. f. Fl. Brit. Ind.* iii. 40.—Ind. or.
appendiculata, Wall. ex Hook. f. l. c. 38 = puberula.
bicuspidata, Wight & Arn. Prod. 403 = Notoniana.
bifaria, Wall. Cat. n. 6278 = Urophyllum hirsutum.
budleioides, Wall. in Wight & Arn. Prod. 402 = tinctoria.
cinerea, DC. Prod. iv. 412; Wall. Cat. n. 6268 = exserta.
coriacea, *DC. l. c.*—Reg. Himal.
corymbosa, DC. l. c. 413; Wall. Cat. n. 6276 = Greenea Jackii.

WENDLANDIA:—
dasythyrsa, *Miq. Fl. Ind. Bat.* ii. 159.—Java.
densiflora, *DC. Prod.* iv. 412 (dentiflora).—Java.
exserta, Blanco, Fl. Filip. ed. II. 104 = paniculata.
exserta, *DC. Prod.* iv. 411.—Ind. or.
glabrata, *DC. l. c.*—As. trop.
glomerulata, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 310.—Burma.
Heyneana, Wall. Cat. n. 6274; et ex Wight & Arn. Prod. 403 = glabrata.
Junghuhniana, *Miq. Fl. Ind. Bat.* ii. 159.—Java.
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. iii. 55.—Kurdist.
laevigata, Miq. Fl. Ind. Bat. ii. 346 = glabrata.
Lawii, *Hook. f. Fl. Brit. Ind.* iii. 40.—Ind. or.
Lawsoniae, *DC. Prod.* iv. 413.—Cochinch.
ligustrina, *Wall. Cat.* n. 6272.—Burma.
longifolia, DC. Prod. iv. 412. = Adenosacme longifolia.
luzoniensis, DC. l. c. = paniculata.
malayana, G. Don, Gen. Syst. iii. 519 = Adenosacme longifolia.
multiflora, Bartl. ex DC. Prod. iv. 412 = paniculata.
nitens, *Wall. Cat.* n. 6271.—Burma.
Notoniana, *Wall. l. c.* n. 6273; *et ex Wight & Arn. Prod.* 403.—Ind. or.
paniculata, *DC. Prod.* iv. 411.—Ind. or.; Malaya.
pendula, *DC. l. c.* 412.—Reg. Himal.
pilosa, G. Don, Gen. Syst. iii. 519 = Bertiera africana.
proxima, DC. Prod. iv. 411 = tinctoria.
puberula, *DC. l. c.* 412.—Reg. Himal.; Burma.
racemosa, G. Don, Gen. Syst. iii. 519 = Bertiera macrocarpa.
rufescens, *Miq. Fl. Ind. Bat.* ii. 160.—Java.
scabra, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 310 = puberula.
scandens, G. Don, Gen. Syst. iii. 520 = Rondeletia scandens.
secunda, Griff. Notul. iv. 266 = Greenea Wightiana.
spicata, DC. Prod. iv. 412 = Greenea Jackii.
sulcata, *G. Don, Gen. Syst.* iii. 519.—Afr. trop.
sumatrana, Miq. Fl. Ind. Bat. ii. 159 = glabrata.
tenuiflora, Miq. ex Hook. f. Fl. Brit. Ind. iii. 39 = glabrata.
tetrandra, DC. Prod. iv. 412 = Hedyotis macrophylla.
Teysmanniana, *Miq. Fl. Ind. Bat.* ii. 346.—Sumatra.
thyrsoidea, Steud. Nom. ed. II. ii. 786 = Notoniana.
tinctoria, *DC. Prod.* iv. 411.—Reg. Himal.; Burma.
trichantha, *Miq. Fl. Ind. Bat.* ii. 347.—Java.
uvariifolia, Hance, in Journ. Bot. viii. (1870) 73 = paniculata.
virgata, G. Don, Gen. Syst. iii. 519 = Pouchetia africana.
Wallichii, *Wight & Arn. Prod.* 402, in obs.—Reg. Himal.
Wightiana, Wall. Cat. n. 6277 = Greenea Wightiana.

WENDLANDIA, Willd. Sp. Pl. ii. 275 (1799) = **Cocculus**, DC. (1818) (Menisp.).
caroliniana, Nutt. Gen. Am. i. 241 = C. carolinus.
populifolia, Willd. Sp. Pl. ii. 275 = carolinus.

WENDTIA, Ledeb. Fl. Ross. ii. 328 (1844) = WENDIA, Hoffm. = **Heracleum**, Linn. (Umbellif.).

WENDTIA, Meyen, Reise, i. 307 (1834). *GERANIACEAE*, Benth. & Hook. f. i. 275.
HYPERUM, Presl, Epim. Bot. 211 (1850).
MARTINIERA, Guill. in Deless. Ic. Sel. iii. 23. t. 40 (1837).
aphanifolia, Griseb. in Goett. Abh. xxiv. (1879) 71.—Reg. Argent.
argentea, Griseb. l. c. 70 = calycina.
calycina, *Griseb.* l. c.—Reg. Argent.
gracilis, *Meyen, Reise*, i. 308.—Chili.
Poeppigiana, *Klotzsch, in Linnaea*, x. (1835–36) 432.—Chili.
Reynoldsii, *Endl. in Walp. Rep.* i. 460.—Chili.

WENSEA, Wendl. Coll. Pl. iii. 24. t. 84 (1819) = **Pogostemon**, Desf. (Labiat.).
pyramidata, Wendl. l. c. 26. t. 84 = P. plectrantoides.

WERNERIA, H. B. & K. Nov. Gen. et Sp. iv. 189. tt. 368, 369 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 451.
ORESIGONIA, Willd. ex DC. Prod. vi. 323 (1837).
ORIBASIA, Moç. & Sesse, ex DC. l. c. (1837).
africana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 422.—Abyss.

WERNERIA:—
apiculata, Sch. Bip. in Bonplandia, iv. (1856) 55, nomen = pygmaea.
aretioides, *Wedd. Chlor. And.* i. 86.—Bolivia.
brachypappa, *Phil. in Anal. Univ. Chil.* (1873) 501.—Chili.
brachypappa, Sch. Bip. in Bonplandia, iv. (1856) 53, 55, nomen = pygmaea.
caespitosa, *Wedd. Chlor. And.* i. 83.—Peruv.
calyculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 204.—Ecuador.
canaliculata, Sch. Bip. in Bonplandia, iv. (1856) 52, 53, 55, nomen = staticaefolia, pumila.
carnulosa, *A. Gray, in Proc. Am. Acad.* v. (1861) 140.—Peruv.
caulescens, *Griseb. in Goett. Abh.* xxiv. (1879) 208.—Reg. Argent.
celmisioides, Sch. Bip. ex Wedd. Chlor. And. i. 82 = staticaefolia.
cherlerioides, Sch. Bip. l. c. 84 = pygmaea.
ciliata, Wedd. ex Sch. Bip. in Linnaea, xxxiv. (1865–66) 530 = ciliolata.
ciliolata, *A. Gray, in Proc. Am. Acad.* v. (1861) 140.—Peruv.
cochlearis, *Griseb. in Goett. Abh.* xxiv. (1879) 208.—Reg. Argent.
cortusifolia, Griseb. l. c. xix. (1874) 194 = Senecio wernerioides.
dactylophylla, *Sch. Bip. in Bonplandia*, iv. (1856) 53, 55, nomen; *et ex Wedd. Chlor. And.* i. 87.—Peruv.
densa, *Benth. Pl. Hartw.* 211.—N. Granat.
digitata, *Wedd. Chlor. And.* i. 86.—Peruv.
disticha, *H. B. & K. Nov. Gen. et Sp.* iv. 193. t. 369.—Ecuador.
Ellisii, *Hook. f. Fl. Brit. Ind.* iii. 357.—Reg. Himal.
frigida, DC. Prod. vi. 323, sphalm. = rigida.
glandulosa, *Wedd. Chlor. And.* i. 85.—Peruv.
graminifolia, Benth. Pl. Hartw. 211 = pygmaea.
graminifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 192. t. 368. f. 1.—Ecuador.
Haenkei, Sch. Bip. ex Wedd. Chlor. And. i. 83, in obs. = caespitosa.
heteroloba, *Wedd. l. c.* 88.—Peruv.
humilis, *H. B. & K. Nov. Gen. et Sp.* iv. 191.—Ecuador.
Lechleri, *Sch. Bip. in Lechl. Berb. Am. Austr.* 57.—Peruv.
Mandoniana, *Wedd. ex Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 80, nomen.—Bolivia.
melanandra, *Wedd. Chlor. And.* i. 88.—Bolivia.
minima, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 277 = pygmaea.
Mocinniana, *DC. Prod.* vi. 324.—Mexic.
nana, *Benth. & Hook. f. Gen.* ii. 451.—Reg. Himal.
nubigena, *H. B. & K. Nov. Gen. et Sp.* iv. 193.—Ecuador.
nuda, A. Gray, in Proc. Am. Acad. v. (1861) 139 = Orbignyana.
Orbignyana, *Wedd. Chlor. And.* i. 85.—Bolivia; Peruv.
pinnatifida, *Remy, in C. Gay, Fl. Chil.* iv. 216.—Chili.
plantaginifolia, *Wedd. ex Sch. Bip. in Bull. Soc. Bot. Fr.* xii. (1865) 80, nomen.—Bolivia.
pumila, *H. B. & K. Nov. Gen. et Sp. Am.* iv. 192. t. 368. f. 2.—Ecuador.
pygmaea, *Gill. ex Hook. ex Arn. in Hook. Journ. Bot.* iii. (1841) 348.—Chili; Peruv.
rhizoma, Remy, in C. Gay, Fl. Chil. iv. 215 = pygmaea.
rigida, *H. B. & K. Nov. Gen. et Sp.* iv. 191.—Ecuador.
setosa, Wedd. ex Sch. Bip. in Linnaea, xxxiv. (1865–66) 530 = strigosissima.
solivaefolia, *Sch. Bip. in Bonplandia*, iv. (1856) 53, 55, nomen; *et ex Wedd. Chlor. And.* i. 87.—Peruv.
spathulata, *Wedd. l. c.* 85.—Bolivia.
staticaefolia, *Sch. Bip. in Bonplandia*, iv. (1856) 53, 55, nomen; *et ex Wedd. l. c.* 81.—Peruv.
strigosissima, *A. Gray, in Proc. Am. Acad.* v. (1861) 140.—Peruv.
villosa, *A. Gray, l. c.* 139.—Peruv.

WERNISEKIA, Scop. Introd. 273 (1777) = **Humiria** Aubl.

WERRINUWA, Heyne, Tracts on India, 49 (1814) = **Guizotia**, Cass. (Compos.).

WESTIA, Vahl, in Skrivt. Nat. Selsk. vi. (1810) 117. *LEGUMINOSAE*, Benth. & Hook. f. i. 583.
grandiflora, *Vahl, l. c.* 118.—Afr. trop.
parviflora, *Vahl, l. c.* 119.—Afr. trop.

WESTONIA, Spreng. Syst. iii. 152, 230 (1826) = **Rothia**, Pers. (Legumin.).
humifusa, Spreng. l. c. 230 = R. trifoliata.

WESTRINGIA, Sm. in Vet. Akad. Handl. Stockh. (1797) 171; et Tracts, 279. t. 3 (1798). *LABIATAE*, Benth. & Hook. f. ii. 1219.
angustifolia, R. Br. Prod. 501 = rigida.
brevifolia, *Benth. Lab. Gen. et Sp.* 459.—Tasman.
capitata, Bartl. in Lehm. Pl. Preiss. i. 362 = Microcorys capitata.
cephalantha, *F. Muell. Fragm.* vi. 110.—Austral.
cinerea, R. Br. Prod. 501 = rigida.
Dampieri, *R. Br. l. c.*—Austral.
eremicola, *A. Cunn. ex Benth. Lab. Gen. et Sp.* 459.—Austral.
glabra, *R. Br. Prod.* 501.—Austral.
grandifolia, *F. Muell. ex Benth. Fl. Austral.* v. 128.—Austral.
grevillina, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 49 = rigida.
longifolia, *R. Br. Prod.* 501.—Austral.
longifolia, Lindl. Bot. Reg. t. 1481 = eremicola.
rigida, *R. Br. Prod.* 501.—Austral.
rosmarinacea, Andr. Bot. Rep. t. 214 = rosmariniformis.
rosmariniformis, Labill. ex Benth. in DC. Prod. xii. 570 = Dampieri.
rosmariniformis, *Sm. in Vetensk. Akad. Handl. Stockh.* (1797) 171; *et Tracts*, 277. t. 3.—Austral.
rubiaefolia, *R. Br. Prod.* 501.—Tasman.
senifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 49.—Austral.
serpyllifolia, Bartl. in Lehm. Pl. Preiss. i. 362 = Microcorys purpurea?
triphylla, Ait. ex W. Baxt. in Loud. Hort. Brit. Suppl. ii. 681, nomen (Quid?).—Austral.
violacea, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 49 = glabra.

WETRIA, Baill. Etud. Gén. Euph. 409 (1858) = **Alchornea**, Sw. (Euphorb.).
trewioides, Baill. l. c. = A. Blumeana.

WETTINIA, Poepp. ex Endl. Gen. 243 (1837). *PALMAE*, Benth. & Hook. f. iii. 902.
augusta, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 39. t. 153.—Peruv.
illaqueans, Spruce, in Journ. Linn. Soc. iii. (1859) 191 = Catoblastus maynensis.
maynensis, Spruce, l. c. = Catoblastus maynensis.
Poeppigii, Kunth, Enum. Pl. iii. 109 = augusta.

WHIPPLEA, Torr. in Pacif. Rail. Rep. iv. 90 (1856). *SAXIFRAGEAE*, Benth. & Hook. f. i. 644.
modesta, *Torr. l. c.*—Calif.
utahensis, S. Wats. in Am. Natural. vii. (1873) 300 = Fendlera utahensis (in Addend.).

WHITEFIELDIA, Nees, in DC. Prod. xi. 220 (1847), sphalm. = **Whitfieldia**, Hook. (Acanth.).

WHITEHEADIA, Harv. Gen. S. Afr. Pl. ed. II. 396 (1868). *LILIACEAE*, Benth. & Hook. f. iii. 807.
bifolia, Baker, in Journ. Linn. Soc. xiii. (1873) 226 = latifolia.
latifolia, *Harv. Gen. S. Afr. Pl.* ed. II. 396.—Afr. austr.

WHITFIELDIA, Hook. Bot. Mag. t. 4155 (1845). *ACANTHACEAE*, Benth. & Hook. f. ii. 1085.
lateritia, *Hook. l. c.*—Afr. trop.
longifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 27.—Afr. trop.

WHITIA, Blume, Cat. Gew. Buitenz. 16 (1823) = **Cyrtandra**, Forst. (Gesner.).
carnosa, Blume, Bijdr. 775 = C. carnosa.
longifolia, Blume, Cat. Gew. Buitenz. 86 = C. oblongifolia.
oblongifolia, Blume, l. c. 17 = C. oblongifolia.
pilosa, Zipp. ex C. B. Clarke, in DC. Monog. Phan. v. 206 = C. oblongifolia.

WHITLAVIA, Harv. in Hook. Lond. Journ. Bot. v. (1846) 311 = **Phacelia**, Juss. (Hydroph.).
gloxinioides, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1226 = P. Whitlavia.
grandiflora, Harv. in Hook. Lond. Journ. Bot. v. (1846) 312 = P. Whitlavia.
minor, Harv. l. c. = P. Whitlavia.

WHITLEYA, D. Don, in News Lit. and Fashion, iii. 108; ex Sweet, Brit. Flow. Gard. Ser. I. t. 125 (1825) = **Scopolia**, Jacq. (Solan.).
stramonifolia, D. Don, ll. cc. = S. lurida.

WHITNEYA, A. Gray, in Proc. Am. Acad. vi. (1865) 549. *COMPOSITAE*, Benth. & Hook. f. ii. 400.
dealbata, *A. Gray, l. c.* 550.—Calif.

WIASEMSKYA, Klotzsch, in Bot. Zeit. iv. (1847) 594, absque desc. *RUBIACEAE* (Quid?).

WIBELIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 1. 97 (1801) = **Crepis**, Vaill. (Compos.).
chondrilloides, Hoppe & Hornsch. Tageb. 278; et ex Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 304 = C. chondrilloides.
foetida, Sch. Bip. Cich. 64 = C. foetida.
graveolens, Gaertn. Mey. & Scherb. Fl. Wett. iii. 1. 144 = C. foetida.

WIBELIA, Pers. Syn. i. 210 (1805) = **Paypayrola**, Aubl. (Violar.).
brasiliensis, Spreng. Neue Entdeck. iii. 63 = P. brasiliensis.
guianensis, Pers. Syn. i. 210 = P. guianensis.

WIBELIA, Roehl. Deutschl. Fl. ed. II. ii. 426 (1813) = **Chondrilla**, Tourn. (Compos.).
apargioides, Roehl. l. c. = C. Peltidium, stipitata.
hieracioides, Roehl. l. c. 427 = Crepis tectorum?
setosa, Roehl. l. c. 582 = Crepis setosa.

WIBORGIA, Roth, Catalect. fasc. ii. 112 (1800) = **Galinsoga**, Ruiz & Pav. (Compos.).
Acmelea, Roth, l. c. = G. parviflora.
armella, Steud. Nom. ed. II. i. 656, in syn., sphalm. = praec.
brachystephana, Heynh. Nom. ii. 767 = G. brachystephana.
oblongifolia, Hook. Bot. Misc. ii. (1831) 226 = G. oblongifolia.
parviflora, H. B. & K. Nov. Gen. et Sp. iv. 256 = G. parviflora.
triloba, Spreng. Nachtr. 1. Bot. Gart. Halle, 41 = Tridax trilobata.
urticaefolia, H. B. & K. Nov. Gen. et Sp. iv. 257. t. 389 = Sabazia urticaefolia.

WIBORGIA, Thunb. Nov. Gen. x. 137 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 477.
LOETHAINIA, Heynh. Nom. i. 883 (1840), sine char.
PELTARIA, Burm. ex DC. Prod. ii. 420 (1825).
angustifolia, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 365 = tenuifolia.
armata, Harv. in Harv. & Sond. Fl. Cap. ii. 91 = spinescens.
cuspidata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 364.—Afr. austr.
flexuosa, *E. Mey. Comm. Pl. Afr. Austr.* 28.—Afr. austr.
floribunda, E. Mey. l. c. = obcordata.
fusca, Eckl. & Zeyh. Enum. 194 = obcordata.
fusca, *Thunb. Nov. Gen.* x. 138.—Afr. austr.
fusca, Willd. ex Walp. in Linnaea, xiii. (1839) 464 = Rafnia cuneifolia.
grandiflora, E. Mey. Comm. Pl. Afr. Austr. 31 = Lebeckia sessilifolia.
heteroclados, E. Mey. l. c. 29 = spinescens.
incurvata, *E. Mey. l. c.* 31.—Afr. austr.
lanceolata, E. Mey. l. c. 29 = sericea.
monoptera, E. Mey. l. c. 30 = spinescens.
obcordata, *Thunb. Nov. Gen.* x. 137.—Afr. austr.
obcordata, Willd. ex Steud. Nom. ed. II. ii. 787 = Loddigesia oxalidifolia.
oblongata, E. Mey. Comm. Pl. Afr. Austr. 31 = fusca.
parviflora, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 364, sphalm. = parvifolia.
parvifolia, *E. Mey. Comm. Pl. Afr. Austr.* 31.—Afr. austr.
sericea, Eckl. & Zeyh. Enum. 194 = obcordata.
sericea, *Thunb. Nov. Gen.* x. 139.—Afr. austr.

WIBORGIA:—
spinescens, *Eckl. & Zeyh. Enum.* 194.—Afr. austr.
tenuifolia, *E. Mey. in Linnaea*, vii. (1832) 153.—Afr. austr.
tetraptera, E. Mey. Comm. Pl. Afr. Austr. 29 = tenuifolia.

WICHURAEA, Nees, in Lehm. Pl. Preiss. ii. 290 (1847) = **Cryptandra**, Sm. (Rhamn.).
arbutiflora, Nees, l. c. = C. arbutiflora.
tubulosa, Nees, l. c. = C. arbutiflora.

WICHURAEA, M. Roem. Syn. Ensat. 277 (1847) = **Bomarea**, Mirb. (Amaryll.).
acicularis, M. Roem. l. c. 280 = B. glaucescens.
andimarcana, M. Roem. l. c. 279 = B. andimarcana.
dulcis, M. Roem. l. c. 278 = B. glaucescens.
glaucescens, M. Roem. l. c. 279 = B. glaucescens.
involucrosa, M. Roem. l. c. 278 = B. glaucescens.
parvifolia, M. Roem. l. c. 279 = B. glaucescens.
puberula, M. Roem. l. c. = B. glaucescens.

WICHUREA, Benth. & Hook. f. Gen. i. 383 (1862), sphalm. = WICHURAEA, Nees = **Cryptandra**, Sm. (Rhamn.).

WICKSTROEMIA, Reichb. Consp. 209 (1828), sphalm. = **Wikstroemia**, Schrad. (Thymel.).

WIDDRINGTONIA, Endl. Cat. Hort. Vindob. i. 209 (1842); Syn. Conif. 31 (1842) = **Callitris**, Vent. (Conif.).
Commersonii, Endl. Syn. Conif. 34.—Madag.
cupressoides, Endl. Cat. Hort. Vindob. i. 209 = C. cupressoides.
ericoides, Knight, Syn. Conif. 13 = Thuya sphaeroidalis.
glauca, Carr. Conif. ed. II. 61 = Thuya orientalis.
juniperoides, Endl. Syn. Conif. 32 = C. arborea.
natalensis, Endl. l. c. 34.—Afr. austr.
Wallichiana, Gord. Pinet. Suppl. 107 = seq.
Wallichii, Endl. Syn. Conif. 34.—Afr. austr.

WIEDEMANNIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. iv. 51 (1837). *LABIATAE*, Benth. & Hook. f. ii. 1210.
erythrotricha, Benth. in DC. Prod. xii. 503 = orientalis.
multifida, *Benth. l. c.*—Armenia.
orientalis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 52.—Armenia.

WIEGMANNIA, Hochst. & Steud. ex Steud. Nom. ed. II. ii. 787 (1841) = **Maerua**, Forsk. (Cappar.).
arabica, Hochst. & Steud. l. c. = M. crassifolia.

WIEGMANNIA, Meyen, Reise, ii. 139 (1835) = **Kadua**, Cham. & Schlecht. (Rubiac.).
glauca, Meyen, l. c. = K. cordata.

WIELANDIA, Baill. Etud. Gén. Euph. 568. t. 22 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 270.
elegans, *Baill. l. c.* 569. t. 22.—Ins. Seychelles.
oblongifolia, Baill. l. c. = Savia oblongifolia.

WIERZBICKIA, Reichb. Nom. 205 (1841) = **Arenaria**, Rupp. (Caryoph.).
laricifolia, Reichb. Ic. Fl. Germ. v. 30. t. 212 = A. laricifolia.
liniflora, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = A. laricifolia.
macrocarpa, Reichb. Ic. Fl. Germ. v. 30. t. 211 = A. laricifolia.
striata, Reichb. l. c. 30 = A. capillacea, etc.

WIESTIA, Boiss. Fl. Orient. v. 559 (1884) = **Boissiera**, Hochst. (Gramin.).

WIESTIA, Sch. Bip. in Herb. & Winckl. Jahrb. Pharm. iv. (1841) 154, in nota = **Lactuca**, Tourn. (Compos.).
acuminata, Sch. Bip. l. c. = L. acuminata.
canadensis, Sch. Bip. l. c. = L. canadensis.
elongata, Sch. Bip. l. c. = L. canadensis.
floridana, Sch. Bip. l. c. = L. floridana.
graminifolia, Sch. Bip. l. c. = L. graminifolia.
spectabilis, Sch. Bip. l. c. = L. Scariola?
tatarica, Sch. Bip. l. c. = L. tatarica.
tenerrima, Sch. Bip. l. c. = L. tenerrima.
tuberosa, Sch. Bip. l. c. = L. tuberosa.
virosa, Sch. Bip. l. c. = L. virosa.

WIGANDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 177 = seq.

WIGANDIA, H. B. & K. Nov. Gen. et Sp. iii. 126 (1818). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 831.
COHIBA, Rafin. Fl. Tellur. iii. 75 (1836).
californica, Hook. & Arn. Bot. Beech. Voy. 364 = Eriodictyon glutinosum.
caracasana, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 1225 = urens, *H. B. & K.*
caracasana, *H. B. & K. Nov. Gen. et Sp.* iii. 128.—Mexic.; Venezuela.
crispa, *H. B. & K. l. c.* 129.—Ecuador.
herbacea, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 117.—Reg. Argent.
Kunthii, Choisy, in Ann. Sc. Nat. Sér. I. xxx. (1823) 246 = urens, *H. B. & K.*
leucocephala, Sch. Bip. in Pollichia, xviii-xix. (1861) 180 = Stoebe leucocephala.
macrophylla, Cham. & Schlecht. in Linnaea, vii. (1832) 382 = urens, *H. B. & K.*
pruritiva, Spreng. Syst. i. 865 = crispa.
scorpioides, *Choisy, in Ann. Sc. Nat.* Sér. I. xxx. (1833) 249.—Mexic.; Nicarag.
urens, *Choisy, l. c.* 246.—Mexic.
urens, *H. B. & K. Nov. Gen. et Sp.* iii. 127.—Mexic.
Vigieri, *Carr. in Rev. Hortic.* (1866) 340.—Hab.?

WIGANDIA, Neck. Elem. i. 95 (1790) = **Disparago**, Gaertn. (Compos.).
disparaginoides, Less. Syn. Comp. 362 = D. lasiocarpa.

WIGGERSIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 1. 6, 33 (1801) = **Vicia**, Tourn. (Legumin.).
cuspidata, Alef. in Bonplandia, ix. (1861) 69 = V. cuspidata.
lathyroides, Gaertn. Mey. & Scherb. Fl. Wett. iii. 1. 34 = V. lathyroides.
lutea, Gaertn. Mey. & Scherb. l. c. 36 = V. lutea.
minima, Alef. in Bonplandia, ix. (1861) 69 = V. lathyroides.
sepium, Gaertn. Mey. & Scherb. Fl. Wett. iii. 1. 34 = V. sepium.
sylvatica, Gaertn. Mey. & Scherb. l. c. = V. sylvatica.

WIGHTIA, Spreng. ex DC. Prod. v. 67 (1836) = **Centratherum**, Cass. (Compos.).
formosa, Spreng. l. c. = C. molle.

WIGHTIA, Wall. Pl. As. Rar. i. 71. t. 81 (1830). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 938.
borneensis, *Hook. f. in Hook. Ic. Pl.* t. 1444.—Borneo.
gigantea, *Wall. Pl. As. Rar.* i. 71.—Reg. Himal.
madagascariensis, Boj. ex Decne. in DC. Prod. viii. 500 = Pentopetia androsaemifolia.

WIGMANNIA, Walp. Rep. vi. 27 (1847), sphalm. = WIEGMANNIA, Meyen = **Kadua**, Cham. & Schlecht. (Rubiac.).

WIKSTROEMIA, Endl. Prod. Fl. Norf. 47 (1833). *THYMELAEACEAE*, Benth. & Hook. f. iii. 193.
CAPURA, Linn. Mant. ii. 149 (1771).
DIPLOMORPHA, Meissn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 289.
androsaemifolia, *Decne. in Jacquem. Voy. Bot.* 146.—Java.
australis, *Endl. Prod. Fl. Norf.* 47.—Ins. Norfolk.
buxifolia, A. Gray, in Journ. Bot. iii. (1865) 304 = W. Uva-ursi.
Candolleana, *Meissn. in DC. Prod.* xiv. 544.—Java.
canescens, *Meissn. in Denkschr. Bot. Ges. Regensb.* iii. (1841) 288.—Reg. Himal.; Zeylan.; China.
Chamaedaphne, *Meissn. in DC. Prod.* xiv. 547.—China.
chinensis, Meissn. l. c. 546 = canescens?
coriacea, Seem. Fl. Vit. 206 = viridiflora.
Cunninghamii, *Meissn. in DC. Prod.* xiv. 545.—Ins. Norfolk.
elongata, *A. Gray, in Journ. Bot.* iii. (1865) 303.—Ins. Sandvic.
foetida, A. Gray, l. c. 302 = viridiflora.

WIKSTROEMIA:—
Forsteri, Decne. in Jacquem. Voy. Bot. 145 = viridiflora.
hypericifolia, Meissn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 287 = Stellera Chamaejasme.
inamoena, Meissn. in DC. Prod. xiv. 547 = canescens.
indica, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 357 = viridiflora.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 134.—Japon.
Junghuhniana, *Miq. Fl. Ind. Bat.* i. 1. 879.—Java.
Monnula, *Hance, in Journ. Bot.* xvi. (1878) 13.—China.
nutans, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 195.—Ins. Hongkong.
ovalifolia, *Decne. in Jacquem. Voy. Bot.* 146.—Afr. austr.
ovata, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 357.—Ins. Philipp.
ovata, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 182 = viridiflora.
pauciflora, Franch. & Sav. Enum. Pl. Jap. i. 406 = canescens.
phillyreaefolia, *A. Gray, in Journ. Bot.* iii. (1865) 304.—Ins. Sandvic.
retusa, *A. Gray, l. c.* 303.—Ins. Luchu.
rotundifolia, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 357.—Ins. Pacif.
salicifolia, Decne. in Jacquem. Voy. Bot. 144. t. 149 = canescens.
sandwichensis, *Meissn. in DC. Prod.* xiv. 545.—Ins. Sandvic.
Shuttleworthiana, Meissn. l. c. 544 = seq.
Shuttleworthii, Meissn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 287 = viridiflora.
sikokiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 481.—Japon.
Spanoghii, *Decne. in Jacquem. Voy. Bot.* 146.—Ins. Timor.; China?
subspicata, *Meissn. in DC. Prod.* xiv. 547.—Afr. austr.
tenuiramis, *Miq. Fl. Ind. Bat. Suppl.* 141, 354.—Ins. Banca.
Uva-ursi, *A. Gray, in Journ. Bot.* iii. (1865) 304.—Ins. Sandvic.
virgata, Meissn. in Denkschr. Bot. Ges. Regensb. iii. (1841) 289 = canescens.
viridiflora, *Meissn. l. c.* 286.—As. trop.; Austral.

WIKSTROEMIA, Schrad. in Goett. Gel. Anz. (1821) ii. 710 (Wickstroemia) = **Laplacea**, H. B. & K. (Ternstr.).
fructicosa, Schrad. l. c. 711 = L. semiserrata.
fruticosa, Spreng. in Vet. Akad. Handl. Stockh. (1821) 168, in obs. = praec.

WIKSTROEMIA, Spreng. in Vet. Acad. Handl. Stockh. (1821) 167. t. 3; Syst. iii. 434 (1826) = **Eupatorium**, Tourn. (Compos.).
glandulosa, Spreng. ll. cc. = E. Dalea.

WILBERFORCIA, Hook. f. ex Planch. in Hook. Ic. Pl. t. 796 (1848) = **Breweria**, R. Br. (Convolv.).

WILBRANDIA, Presl, in Oken, Isis, xxi. (1828) 273. *BORAGINEAE*.
paniculata, *Berl. l. c.*—Ins. Martinic.

WILBRANDIA, Silva Manso, Enum. 30 (1836). *CUCURBITACEAE*, Benth. & Hook. f. i. 831.
drastica, Mart. Syst. Mat. Med. Bras. 80 = Riedeli.
ebracteata, *Cogn. in Mart. Fl. Bras.* vi. iv. 33.—Bras.
fluminensis, Wawra, in Oestr. Bot. Zeitschr. xiii. (1863) 109 = Riedeli.
hibiscoides, *Silva Manso, Enum.* 30.—Bras.
linearis, *Cogn. in Mart. Fl. Bras.* vi. iv. 34.—Bras.
longibracteata, *Cogn. l. c.* 32.—Bras.
Riedeli, *Silva Manso, Enum.* 50.—Bras.
sagittifolia, Griseb. in Goett. Abh. xix. (1874) 145 = villosa.
scabra, Mart. Syst. Mat. Med. Bras. 81 = Riedeli.
verticillata, Cogn. in Mart. Fl. Bras. vi. iv. 30 = Riedeli.
villosa, *Cogn. l. c.* 34. t. 7.—Bras.

WILCKEA, Scop. Introd. 170 (1777) = **Vitis**? Linn. (Ampel.).

WILCKIA, Scop. Introd. 317 (1777) = **Malcomia**, R. Br. (1812) (Crucif.).
africana, F. Muell. Nat. Pl. Vict. i. 33 = M. africana.
maritima, Scop. Introd. 317 = M. maritima.

WILDENOWIA, Thunb. in Vet. Akad. Handl. Stockh. xi. (1790) 26. t. 2, sphalm. = **Willdenowia**, Thunb. (Restiac.).

WILHELMSIA, C. Koch, in Linnaea, xxi. (1848) 400 = **Koeleria**, Pers. (Gramin.).
caucasica, C. Koch, l. c. = K. phleoides.

WILHELMSIA, Reichb. Consp. 206 (1828) = **Arenaria**, Rupp. (Caryoph.).

WILIBALDA, Sternb. in Flora, ii. (1819) 6 = **Coleanthus**, Seid. (Gramin.).
subtilis, Roth, Enum. Pl. Phan. Germ. i. 92 = C. subtilis.

WILKESIA, A. Gray, in Proc. Am. Acad. ii. (1852) 160. *COMPOSITAE*, Benth. & Hook. f. ii. 393.
Gymnoxiphium, *A. Gray, l. c.*—Ins. Sandvic.

WILKIA, F. Muell. Census, 5 (1882), sphalm. = WILCKIA, Scop. = **Malcomia**, R. Br. (Crucif.).

WILKIEA, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 64 = **Kibara**, Endl. (Monim.).
calyptrocalyx, Muell. l. c. = K. macrophylla.
Hugeliana, A. DC. in DC. Prod. xvi. II. 669 = Mollinedia Huegeliana.
macrophylla, A. DC. l. c. = K. macrophylla.

WILLDENOVIA, J. F. Gmel. Syst. 362 (1791) = **Rondeletia**, Linn. (Rubiac.).
Schreberi, J. F. Gmel. l. c.—Hab.?

WILLDENOVIA, Thunb. Fl. Cap. i. 312 (1807) = **Willdenowia**, Thunb. (Restiac.).

WILLDENOWA, Cav. Ic. i. 61. t. 89 (1791) = **Adenophyllum**, Pers. (Compos.).
glandulosa, Cav. l. c. 89 = A. coccineum.

WILLDENOWIA, Steud. Nom. ed. I. 892 (1821) = praec.

WILLDENOWIA, Steud. Nom. ed. I. 892 (1821) = **Rondeletia**, Linn. (Rubiac.).

WILLDENOWIA, Thunb. in Vet. Akad. Handl. Stockh. xi. (1790) 26. t. 2 (Wildenowia); Willd. Sp. Pl. iv. 717 (1805). *RESTIACEAE*, Benth. & Hook. f. iii. 1036.
ANTOCHORTUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836).
NEMATANTHUS, Nees, in Linnaea, v. (1830) 661.
SPIROSTYLIS, Nees, ex Mast. in Journ. Linn. Soc. x. (1869) 271.
arescens, *Kunth, Enum. Pl.* iii. 454.—Afr. austr.
brevis, *Nees, ex Mast. in Journ. Linn. Soc.* x. (1869) 269.—Afr. austr.
compressa, Steud. in Flora, xii. (1829) 134 = Chrysitrix capensis.
compressa, Thunb. in Vet. Akad. Handl. Stockh. xi. (1790) t. 2. f. 3 = Cannamois Cephalotes.
cuspidata, *Mast. in Journ. Linn. Soc.* x. (1869) 271.—Afr. austr.
Ecklonii, Kunth, Enum. Pl. iii. 456 = striata.
fimbriata, *Kunth, l. c.* 455.—Afr. austr.
humilis, *Nees, ex Mast. in Journ. Linn. Soc.* x. (1869) 272.—Afr. austr.
Lucaeana, *Kunth, Enum. Pl.* iii. 455.—Afr. austr.
neglecta, Steud. Syn. Pl. Cyp. 263 = striata.
striata, Spreng. Syst. i. 188 = Hypodiscus Willdenovia.
striata, *Thunb. in Vet. Akad. Handl. Stockh.* xi. (1790) 57. t. 2. f. 1.—Afr. austr.
sulcata, *Mast. in Journ. Linn. Soc.* x. (1869) 270.—Afr. austr.
teres, *Thunb. in Vet. Akad. Handl. Stockh.* xi. (1790) 28. t. 2. f. 2.—Afr. austr.
tristachya, Spreng. ex Kunth, Enum. Pl. iii. 410 = Restio bifidus.

WILLEMETIA, Brongn. in Ann. Sc. Nat. Ser. I. x. (1827) 370. t. 16. f. 1 = **Noltea**, Reichb. (Rhamn.).
africana, Brongn. l. c. 371 = N. africana.
scandens, Eckl. & Zeyh. Enum. 130 = N. africana.

WILLEMETIA, Maerkl. in Schrad. Journ. iii. 1. (1800) 329 = **Chenolea**, Thunb. (Chenop.).
arenaria, Maerkl. l. c. 330 = Kochia arenaria.
caspica, Lehm. ex Bunge, in Mém. Sav. Etr. Pétersb. vi. (1857) 279 (= *Echinopsilon caspicus*).—Reg. Casp.
eriophora, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 209 = Kochia eriophora.
hirsuta, Moq. l. c. 210 = Kochia hirsuta.
lanata, Maerkl. in Schrad. Journ. iii. 1. (1800) 330 = Kochia hyssopifolia.
muricata, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 208 = Kochia muricata.
sedoides, Moq. l. c. 210 = Kochia sedoides.

WILLEMETIA, Neck. Elem. i. 50 (1790) = **Chondrilla**, Tourn. (Compos.).
apargioides, Cass. in Dict. Sc. Nat. xlviii. 428 = C. Peltidium, stipitata.
hieracioides, Monn. Ess. Hierac. 80 = C. stipitata.
prenanthoides, Gren. & Godr. Fl. Fr. ii. 315 = C. prenanthoides.
stipitata, Cass. in Dict. Sc. Nat. xlviii. 428 = C. stipitata.
tuberosa, Fisch. & Mey. ex DC. Prod. vii. 150.—Reg. Casp.

WILLIAMIA, Baill. Etud. Gén. Euph. 559. t. 27 (1858) = **Phyllanthus**, Linn. (Euphorb.).
pruinosa, Baill. l. c. 560. t. 27 = P. discolor.

WILLIBALDA, Steud. Nom. ed. II. ii. 787 (1841), sphalm. = WILIBALDA, Sternb. = **Coleanthus**, Seid. (Gramin.).

WILLICHIA, Mutis, ex Linn. Mant. ii. 553 (1737) = **Sibthorpia**, Linn. (Scrophul.).
callitrichoides, Spreng. Syst. iv. Cur. Post. 26 = Hydranthelium callitrichoides.
repens, Mutis, ex Linn. Mant. ii. 558 = S. pichinchensis?

WILLISELLUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 736 (1821) = **Elatine**, Linn.

WILLKOMMIA, Sch. Bip. ex Nym. Consp. 357 (1879) = **Senecio**, Tourn. (Compos.).
minuta, Sch. Bip, l. c. = S. minutus.

WILLOUGHBEIA, Hook. f. Fl. Brit. Ind. iii. 623 (1883), sphalm. = **Willughbeia**, Roxb. (Apocyn.).

WILLUGBAEYA, Neck. Elem. i. 82 (1790) = **Mikania**, Willd. (Compos.).

WILLUGHBEIA, Klotzsch, in Peters, Reise Mossamb. Bot. 281 (1861) = **Landolphia**, Beauv. (Apocyn.).
cordata, Klotzsch, l. c. 283 = L. florida?
Petersiana, Klotzsch, l. c. 281 = L. Petersiana.
senensis, Klotzsch, l. c. 282.—Afr. trop.

WILLUGHBEIA, Roxb. Pl. Corom. iii. 77. t. 280 (1819). *APOCYNACEAE*, Benth. & Hook. f. ii. 691.
ANCYLOCLADUS, Wall. Pl. As. Rar. iii. 45 (1832).
apiculata, *Miq. Fl. Ind. Bat. Suppl.* 551.—Sumatra.
Burbidgei, [Dyer,] in Kew Gard. Rep. (1880) 44 = firma.
celebica, *Blume, Bijdr.* 1024.—Ins. Celebes.
ceylanica, *Thw. Enum. Pl. Zeyl.* 191.—Zeylan.
coriacea, *Wall. Cat.* n. 1620.—Malaya.
drupacea, Blanco, Fl. Filip. ed. I. 132 = Ardisia humilis.
edulis, *Roxb. Hort. Beng.* 20; *Fl. Ind.* ii. 57.—Reg. Himal.; Burma; Malaya.
firma, *Blume, Mus. Bot. Lugd. Bat.* i. 154.—Malaya.
flavescens, [*Dyer,*] *in Kew Gard. Rep.* (1881) 45; *et ex Hook. f. Fl. Brit. Ind.* iii. 624.—Singapore.
grandiflora, *Dyer, ex Hook. f. Fl. Brit. Ind.* iii. 625.—Malaya.
Gudara, Steud. Nom. ed. II. ii. 787 = edulis.

WILLUGHBEIA:—
javanica, *Blume, Bijdr.* 1024.—Java.
martabanica, Wall. Cat. n. 1619; Pl. As. Rar. iii. 45. t. 272 = edulis.
multilocularis, *Blanco, Fl. Filip.* ed. I. 131.—Ins. Philipp.
oblonga, *Dyer, ex Hook. f. Fl. Brit. Ind.* iii. 625.—Malacca.
pubescens, Mart. Reise Bras. ii. 789 = Hancornia speciosa.
rufescens, *Dyer, ex Hook. f. Fl. Brit. Ind.* iii. 626.—Malacca.
tenuiflora, *Dyer, l. c.* 625.—Malacca.
Treacheri, [*Dyer,*] *in Kew Rep.* (1880) 44, nomen.—Borneo.
umbrosa, *Blume, Mus. Bot. Lugd. Bat.* i. 154.—Java.

WILLUGHBEIA, Scop. in Schreb. Gen. 162 (1789) = **Ambelania**, Aubl. (Apocyn.).
acida, J. F. Gmel. Syst. 434 = A. acida.
auriculata, Spreng. Syst. iv. Cur. Post. 71 = Fagraea auriculata.
elliptica, Spreng. l. c. = Fagraea elliptica.
fragrans, Spreng. l. c. = Fagraea fragrans.
guianensis, J. F. Gmel. Syst. 434 (= *Pacouria guianensis*).—Guian.
obovata, Spreng. Syst. iv. Cur. Post. 71 = Fagraea obovata.
racemosa, Spreng. l. c. = Fagraea racemosa.
scandens, Willd. Sp. Pl. i. 1231 (= *Pacouria guianensis*).—Guian.
volubilis, Spreng. Syst. iv. Cur. Post. 71 = Fagraea racemosa.
zeylanica, Spreng. Syst. i. 672 = Fagraea ceilanica.

WILSONIA, R. Br. Prod. 490 (1810). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 880.
Backhousii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 275.—Austral.
humilis, *R. Br. Prod.* 490.—Austral.
rotundifolia, *Hook. Ic. Pl.* t. 410.—Austral.

WILSONIA, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 172. t. 49 = **Dipyrena**, Hook. (Verben.).
glaberrima, Gill. & Hook. l. c. 173 = D. glaberrima.

WIMMERIA, Nees, ex Meissn. in DC. Prod. xv. 1. 65 (1864) = **Beilschmiedia**, Nees (Laurin.).
thomaea, Nees, l. c. = B. thomaea.

WIMMERIA, Schlecht. in Linnaea, vi. (1831) 427. *CELASTRINEAE*, Benth. & Hook. f. 369.
concolor, *Cham. & Schlecht. l. c.* 428.—Mexic.
confusa, *Hemsl. Diagn. Pl. Nov.* 6.—Mexic.
discolor, *Cham. & Schlecht. in Linnaea*, vi. (1837) 428.—Mexic.
integerrima, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 451 = Zinowiewia integerrima.
pallida, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 379 = confusa.
persicifolia, *Radlk. l. c.*—Mexic.
pubescens, *Radlk. l. c.* 378.—Mexic.
serrulata, *Radlk. l. c.* 379.—Reg. Argent.

WINCHIA, A. DC. in DC. Prod. viii. 326 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 695.
atroviridis, Kurz, For. Fl. Brit. Burma, ii. 170 = Chilocarpus atroviridis.
calophylla, *A. DC. in DC. Prod.* viii. 326.—Burma.
cirrhifera, Gardn. ex Thw. Enum. Pl. Zeyl. 191 = Willughbeia ceylanica.

WINDMANNIA, P. Br. Hist. Jamaic. 212 (1756) = **Weinmannia**, Linn. (Saxifr.).

WINDSORIA, Nutt. Gen. Am. i. 70 (1818) = **Triodia**, R. Br. (Gramin.).
ambigua, Nutt. l. c. = T. ambigua.
pallida, Torr. Cat. Pl. N. York, 91 = Glyceria pallida.
poaeformis, Nutt. Gen. Am. i. 70 = T. cuprea.
stricta, Nutt. in Trans. Am. Phil. Soc. v. (1837) 147.—Am. bor.

WINKLERIA, Reichb. Nom. 236 (1841) = **Mertensia**, Roth (Borag.).

WINTERA, Murr. Syst. ed. XIV. 507 (1784) = **Drimys**, Forst. (Magnol.).
aromatica, Murr. l. c. = D. Winteri.
axillaris, J. F. Gmel. Syst. 852; Willd. Sp. Pl. ii. 1239 = D. axillaris.
granadensis, J. F. Gmel. l. c. 853; Willd. l. c. 1240 = seq.
granatensis, Murr. Syst. ed. XIV. 507 = D. Winteri.

WINTERANA, Linn. Syst. ed. X. 1045 (1759) = seq.

WINTERANIA, Linn. Hort. Cliff. 488 (1737) = **Canella**, P. Br.
Canella, Linn. Syst. ed. X. 1045 = C. alba.
lanceolata, Poir. Encyc. viii. 799 = Drimys aromatica.

WINTERLIA, Dennst. Schluess. Hort. Malab. 27 (1818) = **Limonia**, Linn. (Rutac.).
aromatica, Dennst. l. c. = L. Winterlia.

WINTERLIA, Moench, Meth. 74 (1794) = **Ilex**, Tourn.
glabra, C. Koch, Dendrol. ii. 1. 225 = I. glabra.
triflora, Moench, Meth. 74 = I. glabra.

WINTERLIA, Spreng. Syst. i. 519 (1825) = **Ammannia**, Linn. (Lythrar.).
uliginosa, Spreng. l. c. 788 = A. pentandra.

WIRTGENIA, Jungh. ex Hassk. in Flora, xxvii. (1844) 624 = **Spondias**, Linn. et **Odina**, Roxb. (Anacard.).
decandra, Jungh. l. c. 625 = S. Mangifera.
octandra, Jungh. l. c. 624 = Odina Wodier.

WIRTGENIA, Nees, ex Doell, in Mart. Fl. Bras. ii. II. 40 (1877) = **Paspalum**, Linn. (Gramin.).
paspaloides, Nees, ex Doell, l. c. = P. malacophyllum.

WIRTGENIA, Sch. Bip. in Flora, xxv. (1842) 433 = **Aspilia**, Thou. (Compos.).
abyssinica, Sch. Bip. ex Walp. Rep. vi. 146 = A. abyssinica.
frutescens, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 287 = Wedelia abyssinica.
Kotschyii, Sch. Bip. in Flora, xxv. (1842) 435 = A. Kotschyana.
longeradiata, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 287 = A. Kotschyana?
mexicana, Sch. Bip. in Seem. Bot. Voy. Herald, 304 = A. mexicana.
multiflora, Fenzl, in Flora, xxvii. (1844) 312 = A. multiflora.
Schimperi, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 412 = A. Schimperi.
texana, Sch. Bip. in Seem. Bot. Voy. Herald, 304 = Wedelia hispida.

WISENIA, J. F. Gmel. Syst. 515 (1791) = VISENIA, Houtt. = **Melochia**, Dill. (Stercul.).

WISLIZENIA, Engelm. in Wisliz. Tour N. Mexico, 99 (1848). *CAPPARIDEAE*, Benth. & Hook. f. i. 106.
Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1873) 622. —Am. bor. occ.
refracta, *Engelm. in Wislis. Tour N. Mexico*, 99.—Texas.

WISNERIA, M. Micheli, in DC. Monog. Phan. iii. 82 (1881). *ALISMACEAE*, Benth. & Hook. f. iii. 1007.
filifolia, *Hook. f. in Benth. & Hook. f. Gen.* iii. 1007. —Madag.
Schweinfurthii, *Hook. f. l. c.*—Afr. trop.
triandra, *M. Micheli, in DC. Monog. Phan.* iii. 82.—Ind. or.

WISSADULA, Medic. Malv. 24 (1787). *MALVACEAE*, Benth. & Hook. f. i. 204.
divergens, *Benth. in Benth. & Hook. f. Gen.* i. 204.—Am. trop.
excelsior, Presl, Rel. Haenk. ii. 118 = zeylanica.
gymnostachya, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 202 = spicata.
heterosperma, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 182 = rostrata.
hirsuta, *Presl, Rel. Haenk.* ii. 118.—Bras.
Jamesonii, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 202 = spicata.

WISSADULA:—
Leschenaultiana, Mast. in Hook. f. Fl. Brit. Ind. i. 325 = rostrata.
mucronulata, *A. Gray, ex Torr. Bot. Mex. Bound.* 39. —Am. bor. occ.
nudiflora, *Benth. Bot. Voy. Sulph.* 69.—Am. centr.
periplocifolia, Griseb. Cat. Pl. Cub. 25, partim = mucronulata.
periplocifolia, Thw. Enum. Pl. Zeyl. 27 = zeylanica.
rostrata, *Planch. in Hook. Niger Fl.* 229.—Cosmop. trop.
rostrata, Planch. l. c., partim = zeylanica.
scabra, *Presl, Rel. Haenk.* ii. 117. t. 69. f. 1–14. —Mexic.
spicata, *Presl, l. c.* 117.—Am. trop.
zeylanica, *Medic. Malv.* 25.—Cosmop. trop.

WISTARIA, Nutt. Gen. Am. ii. 115 (1818) (Wisteria); Spreng. Syst. iii. 255 (1826). *LEGUMINOSAE*, Benth. & Hook. f. i. 499.
BRADBURYA, Rafin. Fl. Ludov. 104 (1817).
DIPLONYX, Rafin. l. c. 108, ex desc. (1817).
KRAUNHIA, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen prius.
THYRSANTHUS, Ell. in Journ. Acad. Philad. i. (1817) 371.
australis, *F. Muell. in Journ. Bot.* xxii. (1884) 290.—Austral.
Barkhausiana, *Hort. Cels, ex Pasq. Cat. Orto Bot. Nap.* 108.—Hab.?
brachybotrys, *Sieb. & Zucc. Fl. Jap.* i. 92.—Japon.
chinensis, *DC. Prod.* ii. 390.—China.
consequana, Loud. ex G. Don, in Loud. Hort. Brit. 315 = chinensis.
dubia, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 324.—China.
floribunda, DC. Prod. ii. 390 = chinensis.
frutescens, Poir. in Lam. Illustr. iii. 674 = speciosa.
japonica, Sieb. & Zucc. Fl. Jap. i. 88 = Millettia japonica.
macrostachya, Nutt. in Torr. & Gray, Fl. N. Am. i. 283 = speciosa.
megasperma, F. Muell. Fragm. i. 13 = Millettia megasperma.
multijuga, Van Houtte, in Fl. des Serres, xix. (1869–70) 126 = chinensis.
pallida, Dalz. & Gibs. Bomb. Fl. 61 = Milletia racemosa.
racemosa, Dalz. & Gibs. l. c. = Millettia racemosa.
sinensis, Sweet, Hort. Brit. ed. I. 121 = chinensis.
speciosa, *Nutt. Gen. Am.* ii. 116.—Am. bor.

WISTERIA, Nutt. Gen. Am. ii. 115 (1818) = praec.

WITHANIA, Pauq. Diss. Bellad. (1824) 14. *SOLANACEAE*, Benth. & Hook. f. ii. 893.
DEPREA, Rafin. Sylva Tellur. 57 (1838).
EPLATEIA, Rafin. l. c. (1838).
HYPNOTICUM, Rodrig. ex Meissn. Gen. Comm. 184 (1840).
OPSAGO, Rafin. Sylva Tellur. 54 (1838).
PHYSALOIDES, Moench, Meth. 473 (1794).
PUNEERIA, Stocks, in Journ. As. Soc. Bombay, iii. (1849) 55.
SCLEROMPHALOS, Griff. Notul. iv. 96 (1854).
adpressa, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 64, nomen.—Afr. bor.
anonacea, Dun. in DC. Prod. xxii. 1. 463 = Athenaea anonacea.
arborescens, *Dun. l. c.* 455.—Hab.?
aristata, *Pauq. Diss. Bellad.* (1824) 15.—Ins. Canar.
coagulans, *Dun. in DC. Prod.* xiii. 1. 685.—Afghan.
Coronopus, Torr. Bot. Mex. Bound. 155 = Chamaesarachia Coronopus.
frutescens, *Pauq. Diss. Bellad.* (1824) 15.—Hispan.
hirsuta, Dun. in DC. Prod. xiii. 1. 463 = Athenaea hirsuta.
Martiana, Dun. l. c. 462 = Athenaea Martiana.
micrantha, Dun. l. c. 460 = Athenaea micrantha.
mollis, *Dun. l. c.* 460.—Bras.
Morisoni, Dun. l. c. 454 = somnifera.
novo-friburgensis, *Dun. l. c.* 462.—Bras.
oocarpa, *Dun. l. c.* 463.—Bras.
orinocensis, *Dun. l. c.* 455.—Venezuela.
picta, Dun. l. c. 458 = Athenaea picta.
pogogena, Dun. l. c. 459 = Athenaea pogogena.
Pohliana, Dun. l. c. 461 = Athenaea Pohliana.
pulvinata, *Dun. l. c.* 457.—Bras.

WITHANIA:—
pyrifolia, *Dun. in DC. Prod.* xiii. 1. 461.—Bras.
ramosa, *Dun. l. c.* 456.—Honduras.
Riebeckii, *Schweinf. ex Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotra.
Schottiana, Dun. in DC. Prod. xiii. 1. 461 = Athenaea Schottiana.
somnifera, *Dun. l. c.* 453.—Reg. Mediterr.; Oriens.
sordida, Dun. l. c. 456 = Chamaesaracha sordida.
suberosa, Endl. Gen. 666 = frutescens.
subtriflora, *Dun. in DC. Prod.* xiii. 1. 455.—Peruv.
villosa, Pauq. Diss. Bellad. (1824) 15 = frutescens.
xalapensis, Dun. in DC. Prod. xiii. 1. 455 = Athenaea xalapensis.

WITHARIA, Reichb. Consp. 126 (1828), sphalm. = praec.

WITHERINGIA, L'Hérit. Sert. Angl. 1, 33. t. 1 (1788) = **Bassovia**, Aubl. (Solan.).
acuminata, Dun. in DC. Prod. xiii. 1. 403.—Bras.
acutifolia, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 (= *Saracha acutifolia*).—Am. austr.
alata, Miers, l. c. = Saracha alata.
allogona, Miers, l. c. = Saracha allogona.
angustifolia, Dun. Syn. Solan. 2; H. B. & K. Nov. Gen. et Sp. iii. 17 = Solanum cotopaxense.
aspera, Spreng. Neue Entdeck. i. 266.—Bras.
auriculata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 22 = Saracha auriculata.
Berteroana, Remy, in C. Gay, Fl. Chil. v. 65 = Solanum Berterianum.
biflora, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Hebecladus biflorus.
Candollii, Miers, l. c = Saracha laxa.
chenopodioides, Remy, in C. Gay, Fl. Chil. iv. 69 = Solanum nigrum.
ciliata, H. B. & K. Nov. Gen. et Sp. iii. 15 = Brachistus ciliatus.
ciliata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha ciliata.
conica, Steud. Nom. ed. II. ii. 788.—Bras.
conspersa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1853) 92 = Saracha conspersa.
contorta, Miers, l. c. = Saracha contorta.
crassifolia, Dun. Hist. Solan. 108 = Solanum crassifolium.
crispa, Remy, in C. Gay, Fl. Chil. v. 66 = Solanum crispum.
dentata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha dentata.
diffusa, Miers, l. c. = Saracha laxa.
diploconos, Mart. Nov. Gen. et Sp. iii. 76. t. 229 = Cyphomandra diploconos.
divaricata, Mart. l. c. 75. t. 228 = Cyphomandra divaricata.
diversifolia, Klotzsch, in Walp. Rep. iii. 29 = Brachistus diversifolius.
dumetorum, Dun. Syn. Solan. 1; H. B. & K. Nov. Gen. et Sp. iii. 16 = Brachistus dumetorum.
flexuosa, Remy, in C. Gay, Fl. Chil. v. 70 = Solanum flexuosum *Phil. f.* (in Addend.).
Fournieri, Heurck & Muell. Arg. in Heurck, Obs. Bot. 91.—Mexic.
furcata, Remy, in C. Gay, Fl. Chil. iv. 67 = Solanum furcatum.
Gaudichaudiana, Remy, l. c. 71 = Solanum Gaudichaudi.
Gayana, Remy, l. c. 67 = Solanum Gayanum.
glabrata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha glabrata.
glandulosa, Miers, l. c. = Saracha glandulosa.
hirsuta, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 541 = B. Gardneri.
Jaltomata, Miers, in Ann. & Mag. Nat. Hist. Ser. xi. (1853) 92 = Saracha Jaltomata.
lanceolata, Dun. in DC. Prod. xiii. 1. 404.—Bras.
laxa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha laxa.
macrophylla, Dun. Syn. Solan. 1; H. B. & K. Nov. Gen. et Sp. iii. 14 = Brachistus macrophyllus.
maritima, Remy, in C. Gay, Fl. Chil. iv. 72 = Solanum maritimum.
Menziesii, Dun. in DC. Prod. xiii. 1. 159 = Solanum Menziesii.
microphylla, Griseb. in Lechl. Berb. Am. Austr. 58. —Peruv.
mollis, Dun. in DC. Prod. xiii. 1. 459 = Athenaea picta.

WITHERINGIA :—
mollis, H. B. & K. Nov. Gen. et Sp. iii. 15 = Brachistus mollis.
montana, Dun. Syn. Solan. 24 = Solanum montanum.
multifida, Dun. l. c. 3 = Solanum multifidum.
pendula, Roem. & Schult. Syst. iii. 522 = aspera.
phyllantha, Dun. Syn. Solan. 2 = Solanum phyllanthum.
picta, Mart. Nov. Gen. et Sp. iii. 74. t. 227 = Athenaea picta.
pinnata, Remy, in C. Gay, Fl. Chil. iv. 72 = Solanum pinnatum.
pinnatifida, Dun. Syn. Solan. 2 = Solanum pinnatifidum.
pogonandra, Lem. Illustr. Hortic. vii. (1860) t. 242; et in Rev. Hortic. Sér. IV. ix. (1860) 587.—Mexic.
procumbens, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha procumbens.
propinqua, Miers, l. c. = Saracha propinqua.
purpurea, Lodd. Bot. Cab. t. 1892.—Chili.
rhomboidea, Dun. Syn. Solan. 13; H. B. & K. Nov. Gen. et Sp. iii. 15 = Brachistus rhomboideus.
riparia, H. B. & K. l. c. 16 = Brachistus riparius.
rubra, Remy, in C. Gay, Fl. Chil. iv. 68 = Solanum nigrum.
ruderalis, Remy, l. c. 69 = Solanum ruderale.
salicifolia, Hook. Bot. Misc. ii. (1831) 231 = Capraria biflora.
Selloviana, Sendt. in Mart. Fl. Bras. x. 123 = Saracha Sellowiana.
solanacea, L'Hérit. Sert. Angl. 1, 33. t. 1 = Brachistus solanaceus.
solanacea, Miers, ex Benth. & Hook. f. Gen. ii. 891, in nota = Bassovia solanacea.
stramonifolia, H. B. & K. Nov. Gen. et Sp. iii. 13 = Brachistus stramonifolius.
Tomatillo, Remy, in C. Gay, Fl. Chil. v. 64 = Solanum Tomatillo.
tomentosa, Remy, l. c. 71 = Solanum tomentosum, *Phil. f.*
umbellata, Dun. Syn. Solan. 2 = Solanum umbellatum.
umbellata, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 92 = Saracha umbellata.
vestita, Miers, l. c. = Saracha vestita.
villosa, Miers, l. c. = Saracha villosa.
viscosa, Miers, l. c. = Saracha viscosa.
Zuccagniana, Miers, l. c. = Saracha Zuccagniana.

WITHERINGIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 145 = **Athenaea**, Sendt. (Solan.).
anonacea, Miers, l. c. 146 = A. anonacea.
hirsuta, Miers, l. c. = A. hirsuta.
Martiana, Miers, l. c. = A. Martiana.
micrantha, Miers, l. c. 145 = A. micrantha.
pogogena, Miers, l. c. = A. pogogena.
Pohliana, Miers, l. c. 146 = A. Pohliana.
Schottiana, Miers, l. c. = A. Schottiana.

WITSENIA, Thunb. Gen. Nov. ii. 34. t. 17 (1782) (Witsena); Prod. Pl. Cap. i. 7 (1794). *IRIDEAE*, Benth. & Hook. f. iii. 701.
capitata, Klatt, in Linnaea, xxxiv. (1865–66) 546 = Aristea fruticosa.
corymbosa, Ker-Gawl. in Bot. Mag. t. 895 = Aristea corymbosa (in Addend.).
fruticosa, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 237 = ramosa.
magellanica, Pers. Syn. i. 42 = Tapeinia magellanica.
maura, *Thunb. Nov. Gen.* ii. 34.—Afr. austr.
partita, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1804) 237 = Klattia partita.
pumila, Vahl, Enum. ii. 48 = Tapeinia magellanica.
pyramidalis, Pers. Syn. i. 42 = Lapeyrousia compressa.
ramosa, *Vahl, Enum.* ii. 47.—Afr. austr.
spicata, E. Mey. ex Baker, in Journ. Bot. xiv. (1876) 267 = Aristea racemosa.
tomentosa, Salisb. in Trans. Hort. Soc. i. (1812) 312 = maura.

WITTELSBACHIA, Mart. Nov. Gen. et Sp. i. 80. t. 55 (1824) = **Cochlospermum**, Kunth (Bixin.).
Gossypium, Mart. l. c. 82 = C. Gossypium.
insignis, Mart. l. c. 81. t. 55 = C. insigne.
orinocensis, Mart. l. c. 83 = C. orinoccense.
vitifolia, Mart. l. c. 82 = C. vitifolium.

WITTSTEINIA, F. Muell. Fragm. ii. 136 (1861); iii. 166. *VACCINIACEAE*, Benth. & Hook. f. ii. 577.
vacciniacea, *F. Muell. l. c.* 136.—Austral.

WOEHLERIA, Griseb. in Goett. Abh. ix. (1861) 11. *AMARANTACEAE*, Benth. & Hook. f. iii. 39.
serpyllifolia, *Griseb. l. c.*—Cuba.

WOLFFIA, Horkel, ex Schleid. in Linnaea, xiii. (1839) 389. *LEMNACEAE*, Benth. & Hook. f. iii. 1001.
BRUNIERA, Franch. in Billotia (1864), 25.
GRANTIA, Griff. ex Voigt, Hort. Suburb. Calc. 692 (1845); et Notul. iii. 223 (1851).
HORKELIA, Reichb. ex Bartl. Ord. Nat. 76 (1830).
arrhiza, Wimm. Fl. Schles. (1857) 140 = Michelii.
brasiliensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xii. (1849) 170.—Bras.
columbiana, *Karst. Bot. Unters.* i. (1865) 103.—Mexic.
conguensis, Welw. ex Trimen, in Journ. Bot. iv. (1866) 223, nomen = Welwitschii.
cylindracea, *Hegelm. Lemnac.* 123.—Afr. trop.
Delilii, Miq. in Nederl. Kruidk. Arch. iii. (1855) 429 = Michelii.
Delilii, *Schleid. in Linnaea*, xiii. (1839) 390.—Aegypt.
denticulata, *Hegelm. Lemnac.* 133.—Afr. austr.
gladiata, *Hegelm. l. c.* t. 3. f. 24.—Mexic.
hyalina, Hegelm. l. c. 128 = Delilii.
lingulata, *Hegelm. l. c.* 132. t. 4. f. 31, 32.—Bras.
Michelii, *Schleid. Beitr. Bot.* 233.—Geront. temp.; Austral.
microscopica, *Kurz, in Journ. Linn. Soc.* ix. (1867) 265.—Ind. or.
oblonga, *Hegelm. Lemnac.* 131.—Chili.
punctata, Griseb. Fl. Brit. W. Ind. 512 = brasiliensis.
repanda, *Hegelm. in Journ. Bot.* iii. (1865) 114.—Afr. trop.
Schleideni, Miq. in Nederl. Kruidk. Arch. iii. (1855) 428 = Michelii.
Welwitchsii, *Hegelm. in Journ. Bot.* iii. (1865) 113.—Afr. trop.

WOLFIA, Dennst. Schluess. Hort. Malab. 38 (1818). *SCITAMINEAE.*
spectabilis, *Dennst. l. c.* (an Renealmiae sp.?).—Ind. or.

WOLFIA, Kunth, Enum. Pl. iii. 4 (1841), sphalm. = **Wolffia**, Horkel (Lemnac.).

WOLFIA, Spreng. Syst. i. 808 (1825) (Quid?).
brasiliensis, Spreng. l. c.—Bras.

WOLKENSTEINIA, Regel, Ind. Sem. Hort. Petrop. (1864) 19; Gartenfl. xiv. (1865) 131. t. 471 = **Gomphia**, Schreb. (Ochnac.).
Theophrasta, Regel, ll. cc. = G. gigantophylla.

WOLLASTONIA, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 414; (Herb. Tim. Desc. 86) = **Wedelia**, Jacq. (Compos.).
asperrima, Decne. l. c. = Wedelia asperrima.
biflora, Dalz. & Gibs. Bomb. Fl. 128 = Wedelia urticaefolia.
biflora, DC. Prod. v. 546 = Wedelia biflora.
canescens, DC. l. c. 547 = Wedelia Chamissonis.
ecliptoides, F. Muell. Pl. Vict. ii. sub t. 39 = Eclipta platyglossa.
elongata, DC. Prod. v. 548 = Wedelia elongata.
Forsteriana, DC. l. c. = Wedelia Forsteriana.
glabrata, DC. Coll. Mém. ix. t. 7.—Ins. Timor.
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 72 = Wedelia Horsfieldiana.
insularis, DC. Prod. v. 548.—Ins. Pacif.
javana, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 70.—Java.
javanica, Miq. Fl. Ind. Bat. ii. 1078 = praec.
Junghuhniana, Miq. Pl. Jungh. 502.—Java.
lantanoides, DC. Prod. v. 548.—Ind. occ.
moluccana, DC. l. c. 547.—Ins. Molucc.
montana, DC. l. c.—Java.
pedunculatis, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 69.—Java.
procumbens, DC. Prod. v. 548 = Chrysogonum procumbens.
prostrata, DC. l. c. 549.—Bras.

WOLLASTONIA :—
prostrata, Hook. & Arn. Bot. Beech. Voy. 195.—Mexic.
scaberrima, C. B. Clarke, Comp. Ind. 138 = Wedelia Wallichii.
scabriuscula, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 414 = Wedelia biflora.
scabriuscula, Sch. Bip. ex Miq. Fl. Ind. Bat. ii. 70 = Wedelia Schultziana.
serrulata, Miq. l. c. 74.—Sumatra.
strigulosa, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 414 = Wedelia aristata.
urticaefolia, Hassk. Cat. Hort. Bog. Alt. 99 = Wedelia urticaefolia.
zanzibarensis, DC. Prod. v. 547 = Wedelia biflora.
Zollingeriana, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 123, nomen.—Ins. Sunda.

WOODFORDIA, Salisb. Parad. Lond. t. 42 (1806). *LYTHRARIEAE*, Benth. & Hook. f. i. 778.
floribunda, *Salisb. l. c.*—As. et Afr. trop.
fruticosa, Kurz, in Journ. As. Soc. Beng. xl. (1871) 56 = floribunda.
tomentosa, Bedd. Fl. Sylv. Madr. 117 = floribunda.
uniflora, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 334.—Afr. trop.

WOODVILLEA, DC. Prod. v. 318 (1836) = **Erigeron**, Linn. (Compos.).
calendulacea, DC. l. c. = E. glaucus.

WOOLLSIA, F. Muell. Fragm. viii. 52, 55 (1872) = **Lysinema**, R. Br. (Epacrid.).
pungens, F. Muell. l. c. 55 = L. pungens.

WORMIA, Rottb. in Nye Samml. Vidensk. Selsk. Skrift. ii. (1783) 532. *DILLENIACEAE*, Benth. & Hook. f. i. 13.
CAPELLIA, Blume, Bijdr. 5 (1825).
LENIDIA, Thou. Gen. Nov. Madag. 17 (1806).
alata, *R. Br. ex DC. Syst.* i. 434.—Austral.; N. Guin.
apetala, *Gaudich. in Freyc. Voy. Bot.* 476. t. 99.—Ins. Molucc.
artocarpifolia, *Baker, in Journ. Linn. Soc.* xx. (1883) 88.—Madag.
augusta, Steud. Nom. ed. II. ii. 789 = Dillenia pentagyna.
biflora, *Seem. Fl. Vit.* 3.—Ins. Fiji.
bracteata, Hook. f. & Thoms. Fl. Ind. i. 68 = Dillenia bracteata.
Burbidgei, *Hook. f. Bot. Mag.* t. 6531.—Borneo.
castanaefolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 78.—N. Guin.
coromandelina, Spreng. Syst. ii. 631 = Dillenia pentagyna.
dentata, DC. Syst. i. 434 = triquetra.
excelsa, Hook. f. & Thoms. Fl. Ind. i. 67 = suffruticosa.
excelsa, *Jack, in Malay. Misc.* ii. (1822) n. VII. 69.—Sumatra.
ferruginea, *Baill. Adansonia*, vi. (1865-66) 268.—Ins. Seychelles.
floribunda, Steud. Nom. ed. II. ii. 789 = Dillenia floribunda.
grandiflora, *Miq. Fl. Ind. Bat. Suppl.* 619.—Sumatra.
integra, Hook. f. & Thoms. Fl. Ind. i. 68 = Dillenia retusa.
madagascariensis, *DC. Syst.* i. 433.—Madag.
membranifolia, *Seem. Fl. Vit.* 4.—Ins. Fiji.
oblonga, *Wall. Cat.* n. 951.—Malaya.
obovata, Spreng. Syst. iv. Cur. Post. 213 = Dillenia aurea.
ochreata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 77.—Ins. Celebes.
pteropoda, *Miq. l. c.*—Ins. Molucc.
pulchella, *Jack, in Malay. Misc.* ii. (1820) VII. 70.—Malaya.
retusa, Hook. f. & Thoms. Fl. Ind. i. 68 = Dillenia retusa.
revoluta, Hort. Bog. ex Miq. Journ. Bot. Néerl. i. (1861) 364 = suffruticosa.
scabrella, Spreng. Syst. iv. Cur. Post. 213 = Dillenia scabrella.
subsessilis, Miq. Fl. Ind. Bat. Suppl. 619 = suffruticosa.
suffruticosa, *Griff. Notul.* iv. 706.—Malaya.
triquetra, *Rottb. in Nye Samml. Vidensk. Selsk. Skrift.* ii. (1783) 532. t. 1.—Zeylan.

WORMIA, Vahl, in Skrift. Nat. Selsk. Kjoeb. vi. (1810) 104 = **Ancistrocladus**, Wall. (Dipteroc.).
hamata, Vahl, l. c. = A. Vahlii.

WORMSKIOLDIA, Thonn. in Schum & Thonn. Beskr. Guin. Pl. 165 (1827). *TURNERACEAE*, Benth. & Hook. f. i. 807.
SCHUMACHERIA, Spreng. Gen. 232 (1830).
TRICLICERAS, Thonn. ex DC. Pl. Rar. Jard. Genèv. 56 (1829).
TRICLISIA, Thonn. ex DC. l. c. 50 (1829).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 299 = Streptopetalum serratum.
Biviniana, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 324.—Madag.
brevicaulis, *Urban, in Eichl. Jahrb. Bot. Berl.* ii. (1883) 51.—Afr. trop.
diversifolia, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 36. t. 11 = heterophylla.
glandulifera, *Klotzsch, in Peters, Reise Mossamb. Bot.* 146. t. 26.—Afr. trop.
heterophylla, *Schum. & Thonn. Beskr. Guin. Pl.* 165. —Afr. trop.
Hildebrandtii, Urban & Rolfe, ex Urban, in Eichl. Jahrb. Bot. Berl. ii. (1883) 57 = Streptopetalum Hildebrandtii.
lobata, *Urb. l. c.* 52.—Afr. trop.
longepedunculata, *Mast. in Oliver, Fl. Trop. Afr.* ii. 502.—Afr. trop.
pilosa, Schweinf. ex Urban, in Eichl. Jahrb. Bot. Berl. ii. (1883) 54 = heterophylla.
serrata, Hochst. in Flora, xxiv. (1841) 666 = Streptopetalum serratum.
tanacetifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 147.—Afr. trop.

WORRNIA, J. F. Gmel. Syst. 1646 (Index), sphalm. = **Wormia**, Rottb. (Dilleniac.).

WREDOWIA, Eckl. Top. Verz. 16 (1827) = **Aristea**, Soland. (Irid.).
pulchella, Steud. Nom. ed. II. ii. 789, sphalm. = seq.
pulchra, Eckl. Top. Verz. 16 = A. Wredowia.

WRENCIALA, A. Gray, Bot. U. St. Expl. Exped. i. 180, in obs. (1854) = **Plagianthus**, Forst. (Malvac.).

WRIGHTEA, Roxb. Hort. Beng. 68 (1814); Fl. Ind. iii. 621 (1832) = **Wallichia**, Roxb. (Palm.).
caryotoides, Roxb. ll. cc. = Wallichia caryotoides.

WRIGHTEA, Tussac, Fl. Antill. sub t. 78 (1808) = **Meriania**, Sw. (Melastom.).
rosea, Tussac, l. c. = Meriania leucantha.

WRIGHTIA, R. Br. in Mem. Wern. Soc. i. (1809) 73. *APOCYNACEAE*, Benth. & Hook. f. ii. 712.
BALFOURIA, R. Br. in Mem. Wern. Soc. i. (1809) 70.
angustifolia, *Thw. Enum. Pl. Zeyl.* 193.—Zeylan.
antidysenterica, R. Br. in Mem. Wern. Soc. i. (1809) 73 = zeylanica.
antidysenterica, J. Grah. Cat. Bomb. Pl. 114 = Holarrhena antidysenterica.
calycina, *A. DC. in DC. Prod.* viii. 406.—Java.
Candollei, *Vidal, Phan. Cuming. Philipp.* 186 (Cadollei).—Ins. Philipp.
coalita, Buch.-Ham. ex Pritz. Ic. Ind. i. 1175 (= *Periploca arborea*).—Ind. or.
coccinea, *Sims, Bot. Mag.* t. 2696.—Reg. Himal.; Ins. Philipp.
Coraia, Wall. Cat. n. 1615 = tomentosa.
Cunninghamii, *Benth. Fl. Austral.* iv. 317.—Austral.
dubia, *Spreng. Syst.* i. 638.—Ind. or. ?
flavido-rosea, Trimen, Syst. Cat. Fl. Pl. Ceyl. 54 = seq.
flavo-rosea, *Trimen, in Journ. Bot.* xxiii. (1885) 238. —Zeylan.
Hamiltoniana, Wall. Cat. n. 4461 = tomentosa.
javanica, *A. DC. in DC. Prod.* viii. 405.—Java.
laciniata, *A. DC. l. c.* 406.—Hab. ?
laevis, *Hook. f. Fl. Brit. Ind.* iii. 654.—Malacca.
madagascariensis, *Boj. ex A. DC. in DC. Prod.* viii. 408.—Madag.
mollissima, Hort. Pondich. ex A. DC. l. c. 406 = tinctoria.
mollissima, Wall. Cat. n. 1627 = tomentosa.
multiflora, Zipp. ex Span. in Linnaea, xv. (1841) 325 = tinctoria.
ovata, *A. DC. in DC. Prod.* viii. 405.—Ins. Philipp.
Piscidia, G. Don, Gen. Syst. iv. 86 = Melodinus monogynus.
pubescens, Blume, Bijdr. 1045 = javanica.
pubescens, *R. Br. in Mem. Wern. Soc.* i. (1809) 73.—Malaya; Austral.

WRIGHTIA:—
pubescens, Roth, Nov. Pl. Sp. 120, 397 = tomentosa.
religiosa, *Benth. & Hook. f. Gen.* ii. 713.—Siam.
Rothii, G. Don, Gen. Syst. iv. 86 = tinctoria.
saligna, *F. Muell. ex Benth. Fl. Austral.* iv. 316.—Austral.
Spanogheana, *Miq. Fl. Ind. Bat.* ii. 434.—Ins. Timor.
timorensis, *Miq. l. c.* 433.—Ins. Timor.
tinctoria, *R. Br. in Mem. Wern. Soc.* i. (1809) 73.—Ind. or.
tomentosa, *Roem. & Schult. Syst.* iv. 414.—Ind. or.
Wallichii, A. DC. in DC. Prod. viii. 405 = tomentosa.
zeylanica, *R. Br. in Mem. Wern. Soc.* i. (1809) 73.—Zeylan.

WRIGHTIA, Soland. ex Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 124 = **Meriania**, Sw. (Melastom.).
alba, Soland. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 65 = M. leucantha.
coccinea, Soland. ex Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 126 = M. leucantha.
nana, Soland. ex Naud. l. c. = M. leucantha.
superba, Soland. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 65 = M. purpurea.

WRIXONIA, F. Muell. Fragm. x. 18 (1876). *LABIATAE*.
prostantheroides, *F. Muell. l. c.*—Austral.

WUERSCHMITTIA, Sch. Bip. ex Hochst. in Flora, xxiv. (1841) 1. Intell. 27 = **Melanthera**, Rohr (Compos.).
abyssinica, Sch. Bip. ex Hochst. l. c. = M. abyssinica.

WUERTHIA, Regel, in Bot. Zeit. ix. (1851) 595 = **Ixia**, Linn. (Irid.).
elegans, Regel, l. c. 596 = I. columellaris.

WULFENIA, Jacq. Misc. ii. 62 (1781). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 963.
Ageria, Sm. in Trans. Linn. Soc. vi. (1802) 96 = Paederota Ageria.
Amherstiana, *Benth. Scroph. Ind.* 46.—Reg. Himal.
Bonarota, Sm. in Trans. Linn. Soc. vi. (1802) 96 = Paederota Bonarota.
carinthiaca, *Jacq. Misc.* ii. 62. t. 8. f. 1; *Ic. Pl. Rar.* i. 1. t. 2.—Carinth.
chamaedrifolia, Host, Fl. Austr. i. 19 = Paederota Bonarota.
intermedia, Wall. Cat. n. 408 = Rhyncoglossum obliquum.
lutea, Host, Fl. Austr. i. 19 = Paederota Ageria.
Notoniana, Wall. Cat. n. 409 = Klugia Notoniana.
obliqua, Wall. Cat. n. 407 = Rhyncoglossum obliquum.
orientalis, *Boiss. Diagn.* Ser. I. iv. 75.—Syria.
reniformis, *Dougl. ex Benth. Scroph. Ind.* 46; *et in Hook. Fl. Bor. Am.* ii. 102. t. 71.—Am. bor. occ.

WULFFIA, Neck. Elem. i. 35 (1790). *COMPOSITAE*, Benth. & Hook. f. ii. 367.
CHATIAKELLA, Cass. in Dict. Sc. Nat. xxix. 491 (1823).
CHYLODIA, Rich. ex Cass. l. c. (1823).
CRODISPERMA, Poit. ex Cass. l. c. xlvi. 403 (1827).
TILESIA, G. F. W. Mey. Prim. Fl. Esseq. 251 (1818).
Blanchetii, DC. Prod. v. 564.—Bras.
capitata, Sch. Bip. in Linnaea, xxi. (1848) 246 = stenoglossa.
elongata, Miq. Stirp. Surin. Sel. 193 = stenoglossa.
havanensis, DC. Prod. v. 563 = stenoglossa.
longifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 293 = stenoglossa.
maculata, DC. Prod. v. 563 = stenoglossa.
oblongifolia, DC. l. c. 564 = stenoglossa.
platyglossa, DC. l. c. 563 = stenoglossa.
quitensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 182.—Ecuador.
Salzmanni, DC. Prod. v. 563 = stenoglossa.
scandens, *DC. l. c.* 564.—Peruv.
stenoglossa, *DC. l. c.* 563.—Am. trop.
strigosa, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 174 = stenoglossa.
suffruticosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 292 = stenoglossa.

WULLSCHLAEGELIA, Reichb. f. in Bot. Zeit. xxi. (1863) 131. *ORCHIDEAE*, Benth. & Hook. f. iii. 594.
aphylla, *Reichb. f. l. c.*—Jamaic.
calcarata, *Benth. in Journ. Linn. Soc.* xviii. (1881) 342.—Bras.

WUNDERLICHIA, Riedel, ex Benth. & Hook. f. Gen. ii. 489 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 489.
Glaziovii, *Baker, in Journ. Bot.* xx. (1882) 225.—Bras.
insignis, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 285. —Bras.
mirabilis, *Riedel, ex Baker, in Mart. Fl. Bras.* vi. III. 343.—Bras.

WURFBAEINIA, Steud. Nom. ed. II. ii. 789 (1841), sphalm. = seq.

WURFBAINIA, Giseke, Prael. Linn. 199, 206 (1792) = **Amomum**, Linn. (Scitam.).
uliginosa, Giseke. l. c. 199 = A. uliginosum.

WURMBAEA, Steud. Nom. ed. II. ii. 789 (1841), sphalm. = seq.

WURMBEA, Thunb. Nov. Gen. i. 18. t. 1 (1781). *LILIACEAE*, Benth. & Hook. f. iii. 823.
SKIZIMA, Rafin. Fl. Tellur. ii. 32 (1836).
bullata, Willd. Hort. Berol. i. 8 = Aletris aurea.
campanulata, Willd. Sp. Pl. ii. 265 = capensis.
capensis, *Thunb. Nov. Gen.* i. 18.—Afr. austr.
dioica, *F. Muell. Fragm.* x. 119.—Austral.
Drummondii, *Benth. Fl. Austral.* vii. 28.—Austral.
inusta, *Banks, ex Ker-Gawl. in Bot. Mag.* sub t. 694. —Afr. austr.
Kraussii, *Baker, in Journ. Linn. Soc.* xvii. (1879) 487. —Afr. austr.
longiflora, Willd. Sp. Pl. ii. 266 = capensis.
pumila, Willd. l. c. 265 = capensis.
purpurea, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 694; [Dryand. in] Ait. Hort. Kew. ed. II. ii. 326 = capensis.
purpurea, Drège, ex Kunth, Enum. Pl. iv. 156 = Dipidax ciliata.
pygmaea, *Benth. Fl. Austral.* vii. 28.—Austral.
remota, Banks, ex Ker-Gawl. in Bot. Mag. sub t. 694 = capensis.
revoluta, Banks, l. c. = capensis.
stigmosa, *Banks, l. c.*—Afr. austr.
tenella, *Benth. Fl. Austral.* vii. 28.—Austral.
tenuis, *Baker, in Journ. Linn. Soc.* xvii. (1879) 436.—Afr. trop.
truncata, Schlecht. in Linnaea, i. (1826) 93 = capensis.
tubulosa, *Benth. Fl. Austral.* vii. 28.—Austral.

WURMSCHNITTIA, Benth. in Journ. Linn. Soc. xiii. (1873) 441, sphalm. = WUERSCHMITTIA, Sch. Bip. = **Melanthera**, Rohr (Compos.).

WURTZIA, Baill. Adansonia, i. (1860–61) 186. t. 7 = **Phyllanthus**, Linn. (Euphorb.).
tetracocca, Baill. l. c. 187 = P. virens.

WYDLERIA, DC. Coll. Mém. v. 36. t. 7 (1829); Prod. iv. 103 (1830) = **Carum**, Rupp. (Umbell.).
chilensis, Fisch. & Trautv. Ind. Sem. Hort. Petrop. vii. 58.—Chili.
humilis, Phil. in Linnaea, xxxiii. (1864–65) 92.—Chili.
portoricensis, DC. Coll. Mém. v. 36. t. 7 = C. Petroselinum.

WYETHIA, Nutt. in Journ. Acad. Philad. vii. (1834) 39. t. 5. *COMPOSITAE*, Benth. & Hook. f. ii. 374.
ALARÇONIA, DC. Prod. v. 537 (1836).
MELARHIZA, Kellogg, in Proc. Calif. Acad. i. ed. II. (1873) 37.
amplexicaulis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 352.—Am. bor. occ.
angustifolia, *Nutt. l. c.*—Am. bor. occ.
arizonica, *A. Gray, in Proc. Am. Acad.* viii. (1873) 655.—Am. bor. occ.
californica, *Nutt. ex Walp. Rep.* ii. 608, nomen.—Calif.

WYETHIA:—
coriacea, *A. Gray, in Proc. Am. Acad.* xi. (1876) 77.—Calif.
glabra, *A. Gray, l. c.* vi. (1865) 543.—Calif.
helenioides, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 353.—Calif.
helianthoides, *Nutt. in Journ. Acad. Philad.* vii. (1834) 38.—Am. bor. occ.
longicaulis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 4.—Calif.
mollis, *A. Gray, l. c.* vi. (1865) 544.—Calif.
ovata, *Torr. & Gray, ex Torr. in Emory, Notes Mil. Reconnois.* 143.—Calif.
reticulata, *Greene, in Bull. Calif. Acad.* i. 1. (1884) 9.—Calif.
robusta, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 352 = angustifolia.
scabra, *Hook. Lond. Journ. Bot.* vi. (1847) 245.—Am. bor. occ.

WYLIA, Hoffm. Gen. Umb. ed. I. 3. t. 2 (1814) = **Scandix**, Tourn. (Umbell.).
australis, Hoffm. l. c. 5. t. 2. f. 1 = S. australis.
grandiflora, Hoffm. l. c. 15. t. 2. f. 3 = S. grandiflora.
iberica, Hoffm. l. c. i. 19. t. 2. f. 4 = S. iberica.
radians, Hoffm. l. c. 11. t. 2. f. 2 = S. grandiflora.

X

XAIASME, Rafin. Fl. Tellur. iv. 106 (1836) = **Stellera**, Linn. (Thymel.).
bicolor, Rafin. l. c. = S. Chamaejasme.

XALKITIS, Rafin. New Fl. Am. iv. 71 (1836) = **Aster**, Tourn. (Compos.).
ciliata, Rafin. l. c. = A. ciliatus, *Walt.*

XAMACRISTA, Rafin. Sylva Tellur. 127 (1838) = **Cassia**, Linn. (Legumin.).
triflora, Rafin. l. c. = C. Chamaecrista.

XAMESIKE, Rafin. Fl. Tellur. iv. 115 (1836) = **Euphorbia**, Linn.
scordifolia, Rafin. l. c. = E. scordifolia.
vulgaris, Rafin. l. c. = E. Chamaesyce.
xamobala, Rafin. l. c. = E. Chamaesyce?

XANTHAEA, Reichb. Nom. 132 (1841) = **Erythraea**, Renealm. (Gentian.).

XANTHANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 189 = **Anthoxanthum**, Linn. (Gramin.).

XANTHE, Schreb. Gen. 710 (1791) = **Clusia**, Linn. (Guttif.).
oblonga, Raeusch. Nom. ed. III. 292 = C. macrocarpa.
parviflora, Willd. Sp. Pl. iv. 877 = C. macrocarpa.
scandens, Willd. l. c. = C. microcarpa.

XANTHIDIUM, Delpino, Studi sopra Artemis. 62 (1871) = **Franseria**, Cav. (Compos.).
ambrosioides, Delpino, l. c. 63 = F. ambrosioides.
artemisioides, Delpino, l. c. = F. artemisioides.
discolor, Delpino, l. c. = F. discolor.
rhombophyllum, Delpino, l. c. 64.—Peruv.
tenuifolium, Delpino, l. c. 62 = F. tenuifolia.

XANTHISMA, DC. Prod. v. 94 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 253.
CENTAURIDIUM, Torr. & Gray, Fl. N. Am. ii. 246 (1840).
Drummondii, Hook. f. Bot. Mag. sub t. 6275 = texanum.
texanum, *DC. Prod.* v. 95; *Torr. in Marcy, Rep.* (1852) 303. t. 10.—Texas.

XANTHIUM, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 286 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 355.
ACANTHOXANTHIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 110.
LAPPARIA, Heist. Syst. 11 (1748).
STRUMARIUM, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 89.

XANTHIUM:—
abyssinicum, Wallr. Beitr. Bot. i. (1842) 230 = X. Strumarium.
ambrosioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 310.—Chili.
americanum, Walt. Fl. Carol. 231 = canadense.
anatolicum, Boiss. & Heldr. Diagn. Ser. I. xi. 32, in nota = X. Strumarium.
annuum, Georgi, Reise, ii. 713 = X. Strumarium.
antiquorum, Wallr. Beitr. Bot. i. (1842) 229 = X. Strumarium.
arenarium, *Lasche, in Bot. Zeit.* xiv. (1856) 411.—Am.?
armatum, Humb. & Bonpl. ex Wallr. Beitr. Bot. i. (1842) 242 = spinosum.
brachyacanthum, Wallr. l. c. 244 = spinosum.
brasilicum, Vell. Fl. Flum. x. t. 23 = X. Strumarium.
brevirostre, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 405.—Abyss.
brevirostre, Wallr. Beitr. Bot. i. (1842) 235 = X. Strumarium.
canadense, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor. et austr.
canadense, Sabbat. ex Nym. Consp. 475 = spinosum.
catharticum, H. B. & K. Nov. Gen. et Sp. iv. 274 = spinosum.
Cavanillesii, Schouw, Ind. Sem. Hort. Haun. (1849) 14.—Cf. Ann. Sc. Nat. Sér. III. xii. (1849) 357 = canadense.
chinense, Mill. Gard. Dict. ed. VIII. n. 4 = X. Strumarium.
cordifolium, Stokes, Bot. Mat. Med. iv. 380 = X. Strumarium.
cuneatum, Moench, Meth. Suppl. 300 = canadense.
discolor, Wallr. Beitr. Bot. i. (1842) 232 = X. Strumarium.
echinatum, Murr. in Comm. Gotting. vi. (1783–84) 32. t. 4 = canadense.
eriocarpon, Wallr. Beitr. Bot. i. (1842) 242 = ambrosioides.
fruticosum, Linn. f. Suppl. 413 = Franseria artemisioides.
fuscescens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 36 = X. Strumarium.
homothalamum, *Spreng. Neue Entdeck.* i. 259.—Bras.
humile, Heynh. Nom. i. 858 (Quid?).—Am. austr.?
inaequilaterum, DC. Prod. v. 523 = X. Strumarium.
indicum, Koen. ex Roxb. Hort. Beng. 67; Fl. Ind. iii. 601 = X. Strumarium.
italicum, Moretti, in Brugnat. Giorn. Fis. Dec. II. v. (1822) 326 = orientale.
laevigatum, Muhl. ex Wallr. Beitr. Bot. i. (1842) 231 = X. Strumarium.
longirostre, Wallr. l. c. 237 = canadense.
macrocarpum, DC. Fl. Fr. Suppl. 356 = canadense.
maculatum, Rafin. in Am. Monthly Mag. (1818) 344; Am. Journ. Sc. Ser. I. i. (1819) 151 = canadense.
monoicum, Gilib. Fl. Lituan. i. 170 = X. Strumarium.
numidicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 37 = X. Strumarium.
occidentale, Bertol. Lucub. 38 = canadense.
orientale, Blume, Bijdr. 915 = X. Strumarium.
orientale, Cav. Ic. iii. 11. t. 221 = canadense.
orientale, *Linn. Sp. Pl.* ed. II. 1400.—China; Japon.; Zeylan.
oviforme, Wallr. Beitr. Bot. i. (1842) 240 = orientale.
parvifolium, *DC. Prod.* v. 524.—Am. austr.
pensylvanicum, *Wallr. Beitr. Bot.* i. (1842) 236.—Am. bor.
pungens, Wallr. l. c. 231 = X. Strumarium.
Revelieri, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 36 = orientale.
riparium, Lasche, in Bot. Zeit. xiv. (1856) 412 = orientale.
Roxburghii, Wallr. Beitr. Bot. i. (1842) 233 = X. Strumarium.
saccharatum, Wallr. l. c. 238 = orientale.
sphaerocephalum, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 503 = X. Strumarium.
spinosum, *Linn. Sp. Pl.* 987.—Cosmop.
Strumarium, Ell. Sketch, ii. 479; Bigel. Fl. Bost. ed. II. 342 = canadense.
Strumarium, *Linn. Sp. Pl.* 987.—Cosmop.
vulgare, Lam. Fl. Fr. ii. 56 = X. Strumarium.
xanthocarpum, Wallr. Beitr. Bot. i. (1842) 241 = spinosum.

XANTHO, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 191 = **Lasthenia**, Cass. (Compos.).
glabrata, Remy, l. c. 192 = L. glabrata.

XANTHOCEPHALUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140. *COMPOSITAE*, Benth. & Hook. f. ii. 249.
GRINDELIOPSIS, Sch. Bip. in Bonplandia, vi. (1858) 356.
GUENTHERA, Regel, Ind. Sem. Hort. Petrop. (1857) 42; et Gartenfl. (1858) 44.
XANTHOCOMA, H. B. & K. Nov. Gen. et Sp. iv. 310. t. 412 (1820).
Alamani, *Benth. & Hook. f. Gen.* ii. 249.—Mexic.
Benthamianum, *Hemsl. Biol. Centr. Am. Bot.* ii. 110.—Mexic.
Bonplandianum, Benth. Pl. Hartw. 16 = Benthamianum.
Bonplandianum, DC. Prod. vi. 44 = centauroides.
centauroides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 140.—Am. austr.
conoideum, *Hemsl. Biol. Centr. Am. Bot.* ii. 110.—Mexic.
dentatum, *Schau. ex Hemsl. l. c.* 111.—Mexic.
gymnospermoides, *Benth. & Hook. f. Gen.* ii. 249.—Mexic.
humile, *Benth. & Hook. f. l. c.*; *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 111.—Mexic.
microglossum, *Spruce, ex Benth. & Hook. f. Gen.* ii. 249.—Bras.?
sericocarpum, *A. Gray, in Proc. Am. Acad.* xv. (1880) 31.—Mexic.
suffruticosum, *DC. Prod.* vi. 44.—Mexic.
Wrightii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 632.—Am. bor. occ.

XANTHOCERAS, Bunge, (Enum. Pl. Chin. Bor. 11) in Mém. Sav. Etr. Pétersb. ii. (1834) 85. *SAPINDACEAE*, Benth. & Hook. f. i. 408.
sorbifolia, *Bunge, l. c.*—China.

XANTHOCERCIS, Baill. Adansonia, ix. (1870) 293. *LEGUMINOSAE.*
madagascariensis, *Baill. l. c.* 294.—Madag.

XANTHOCHRYSUM, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 199 = **Helichrysum**, Vaill. (Compos.).
filifolium, Turcz. l. c. = H. filifolium.

XANTHOCHYMUS, Roxb. Pl. Corom. ii. 51. t. 196 (1798) = **Garcinia**, Linn. (Guttif.).
dubius, Roxb. ex Steud. Nom. ed. II. ii. 790 (Quid?).—Ind. or.
dulcis, Roxb. Hort. Beng. 42; Fl. Ind. ii. 631 = G. dulcis.
elegans, Zipp. ex Span. in Linnaea, xv. (1841) 178.—Ins. Timor.
guineensis, G. Don, in Loud. Hort. Brit. 320, nomen (Quid?).—Afr. trop.
javanensis, Blume, Bijdr. 216 = G. dulcis.
luteus, Lodd. ex G. Don, in Loud. Hort. Brit. 320, nomen (Quid?).—Ind. or.
macrophyllus, Lodd. l. c. (Quid?).—Ind. or.
madagascariensis, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 305 = G. madagascariensis.
ovalifolius, Bedd. Fl. Syn. Gen. p. xxi = G. spicata.
ovalifolius, Roxb. Hort. Beng. 42; Fl. Ind. ii. 632 = G. ovalifolia.
pictorius, Roxb. Pl. Corom. ii. 51. t. 196 = G. Xanthochymus.
purpureus, Lodd. ex Loud. Hort. Brit. 320 (Quid?).—Ind. or.
quadrifarius, Oliver, Fl. Trop. Afr. i. 168 = G. quadrifaria.
spicatus, Wight & Arn. Prod. 102 = G. ovalifolia.
tinctorius, DC. Prod. i. 562 = G. Xanthochymus.

XANTHOCOMA, H. B. & K. Nov. Gen. et Sp. iv. 310. t. 412 (1820) = **Xanthocephalum**, Willd. (Compos.).
dentata, Schau. in Linnaea, xix. (1847) 723.—Mexic.
gymnospermoides, Benth. & Hook. f. Gen. ii. 250, lapsu = Xanthocephalum gymnospermoides.
humilis, H. B. & K. Nov. Gen. et Sp. iv. 311. t. 112 = Xanthocephalum humile.

XANTHOCROMYON, Karst. in Bot. Zeit. v. (1847) 694 = **Trimeza**, Salisb. (Irid.).
Herberti, Karst. l. c. = T. lurida.

XANTHOGALUM, Avé-Lall. in Fisch. & Mey. Ind. Sem. Hort. Petrop. viii. 73 (1842) = **Dorema**, D. Don (Umbell.).
purpurascens, Avé-Lall. l. c. = D. Aucheri.

XANTHOLEPIS, Willd. ex Less. in Linnaea, iv. (1829) 338 = **Cacosmia**, H. B. & K. (Compos.).
tomentosa, Willd. ex Less. l. c. = C. rugosa.

XANTHOLINUM, Reichb. Handb. 306 (1837) = **Linum**, Tourn.
arboreum, Reichb. l. c. 307 = L. arboreum.
campanulatum, Reichb. l. c. = L. campanulatum.
capitatum, Reichb. l. c. = L. capitatum.
flavum, Reichb. l. c. = L. flavum.
nodiflorum, Reichb. l. c. = L. nodiflorum.
tauricum, Reichb. l. c. = L. tauricum.

XANTHONANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119; viii. (1881) 189 = **Anthoxanthum**, Linn. (Gramin.).
odoratum, St. Lag. l. c. = A. odoratum.

XANTHOPHTHALMUM, Sch. Bip. Tanacet. 17 (1844) = **Chrysanthemum**, Tourn. (Compos.).
macrocephalum, Sch. Bip. in Bonplandia, viii. (1860) 369 = C. trifurcatum.
segetum, Sch. Bip. Tanacet. 17 = C. segetum.
trifurcatum, Sch. Bip. in Bonplandia, viii. (1860) 369 = C. trifurcatum.

XANTHOPHYLLON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = **Zanthoxylum**, Linn. (Rutac.).
Clava-Herculis, St. Lag. l. c. = Z. Clava-Herculis.
clavatum, St. Lag. l. c. = X. Clava-Herculis.

XANTHOPHYLLUM, Roxb. Hort. Beng. [28] (1814); Pl. Corom. iii. 81. t. 284 (1819). *POLYGALEAE*, Benth. & Hook. f. i. 139.
JACKIA, Blume, Bijdr. 60 (1825).
KAULFUSSIA, Dennst. Schluess. Hort. Malab. 30 (1818).
MACINTYRIA, F. Muell. Fragm. v. 8 (1865).
SKAPHIUM, Miq. Fl. Bat. Ind. Suppl. 357 (1860).
acuminatissimum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 276.—Sumatra.
adenopodum, *Miq. Fl. Ind. Bat. Suppl.* 393.—Sumatra.
adenotus, *Miq. l. c.*—Sumatra.
affine, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 271.—Burma; Malaya.
ancolanum, *Miq. Fl. Ind. Bat. Suppl.* 394.—Sumatra.
angustifolium, Wight, Illustr. Ind. Bot. i. 50 = flavescens.
Arnottianum, Wight, l. c. = flavescens.
borneense, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 277.—Borneo.
cordatum, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 274.—Borneo.
eglandulosum, *Griff. Notul.* iv. 537.—Burma.
ellipticum, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* i. 276.—Malaya.
eurhynchum, *Miq. l. c.* 277.—Sumatra.
excelsum, *Blume, ex Miq. Fl. Ind. Bat.* i. II. 129.—Java.
flavescens, *Roxb. Hort. Beng.* [88]; *Fl. Ind.* ii. 222.—Ind. or.; Malaya.
flavescens, Wall. Cat. n. 4198 = affine.
glaucescens, *Miq. Fl. Ind. Bat. Suppl.* 394.—Sumatra.
glaucum, *Wall. Cat.* n. 4199.—Burma.
Griffithii, *Hook. f. ex A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 210.—Burma; Malaya.
insigne, *A. W. Benn. l. c.* 211.—Malacca.
Korthalsianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 277.—Borneo.
lateriflorum, *Miq. l. c.* 318.—Sumatra.
longifolium, *D. Dietr. Syn. Pl.* ii. 1277.—Java.
Macintyrii, *F. Muell. Fragm.* v. 57.—Austral.
Maingayi, *Hook. f. ex A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 210.—Malacca.
obscurum, *A. W. Benn. l. c.* 211.—Singapore.
palembanicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 317.—Sumatra.
paniculatum, Miq. Fl. Ind. Bat. Suppl. 393 = flavescens.

XANTHOPHYLLUM:—
Roxburghianum, Wight, Illustr. Ind. Bot. i. 50 = flavescens.
rufum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 210.—Malacca.
stipitatum, *A. W. Benn. l. c.*—Malacca.
sumatranum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 275.—Sumatra.
undulatum, Wight, Illustr. Ind. Bot. i. 50 = flavescens.
virens, Roxb. Hort. Beng. [28]; Fl. Ind. ii. [221] = flavescens.
virescens, D. Dietr. Syn. Pl. ii. 1277 = excelsum.
vitellinum, *D. Dietr. l. c.*—Java.

XANTHOPHYTUM, Reinw. ex Blume, Cat. Gew. Buitenz. 57 (1823); et Bijdr. 989. *RUBIACEAE*, Benth. & Hook. f. ii. 53.
calycinum, *Benth. & Hook. f. l. c.*—Ins. Pacif.
fruticulosum, *Reinw. ex Blume, Cat. Gew. Buitenz.* 57; *et Bijdr.* 989.—Java.
spicatum, *Blume, ex DC. Prod.* iv. 413.—Java.

XANTHOPSIS, C. Koch, in Linnaea, xxiv. (1851) 422 = **Centaurea**, Linn. (Compos.).
xanthocephala, C. Koch, l. c. = C. xanthocephala.

XANTHORIZA, Woodhouse, in Med. Repos. N. York, v. (1802) 160, sphalm. = seq.

XANTHORRHIZA, Marsh. Arbust. Am. 167 (1785) = **Zanthorhiza**, L'Hérit. (Ranunc.).

XANTHORRHOEA, Sm. in Trans. Linn. Soc. iv. (1798) 219. *JUNCACEAE*, Benth. & Hook. f. iii. 865.
arborea, *R. Br. Prod.* 288.—Austral.
australis, *R. Br. l. c.*—Austral.
bracteata, *R. Br. l. c.*—Austral.
Brunonis, Endl. in Lehm. Pl. Preiss. ii. 39 = Preissii.
Drummondii, Harv. in Hook. Kew Journ. vii. (1855) 57 = Preissii.
gracilis, *Endl. in Lehm. Pl. Preiss.* ii. 39.—Austral.
hastilis, *R. Br. Prod.* 288.—Austral.
hastilis, Sm. in Trans. Linn. Soc. iv. (1798) 219 = media.
humilis, Steud. Nom. ed. II. ii. 790, sphalm. = X. Pumilio.
macronema, *F. Muell. Fragm.* iv. 112.—Austral.
media, *R. Br. Prod.* 288.—Austral.
minor, *R. Br. l. c.*—Austral.
pecoris, F. Muell. Fragm. iv. 110 = Preissii.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 39.—Austral.
Pumilio, *R. Br. Prod.* 288.—Austral.
quadrangulata, *F. Muell. Fragm.* iv. 111.—Austral.
quadrangulata, Tate, in Trans. Roy. Soc. Austral. vi. (1882–83) 168 = Tateana.
resinosa, Pers. Syn. i. 370 = media.
simiplana, *F. Muell. Fragm.* iv. 111.—Austral.
Tateana, *F. Muell. in Zeitschr. Oesterr. Apoth. Ver.* xxiii. (1883) 294.—Austral.

XANTHOSELINUM, Schur, Enum. Pl. Transs. 264 (1866) = **Peucedanum**, Tourn. (Umbell.).
alsaticum, Schur, l. c. = P. alsaticum.

XANTHOSIA, Rudge, in Trans. Linn. Soc. x. (1811) 301. t. 22. *UMBELLIFERAE*, Benth. & Hook. f. i. 874.
CRUCIELLA, Leschen. ex DC. Prod. ii. 74 (1830).
LEUCOLAENA, R. Br. App. Flind. Voy. 557 (1814).
PENTAPELTIS, Bunge, in Lehm. Pl. Preiss. i. 292 (1845).
SCHOENOLAENA, Bunge, l. c. 289 (1845).
Atkinsoniana, *F. Muell. Fragm.* ii. 127.—Austral.
candida, *Steud. Nom.* ed. II. ii. 790; *Bunge, in Lehm. Pl. Preiss.* i. 291.—Austral.
ciliata, *Hook. Ic. Pl.* t. 726.—Austral.
dissecta, *Hook. f. in Hook. l. c.* t. 302.—Austral.
fruticulosa, *Benth. Fl. Austral.* iii. 361.—Austral.
glabrata, Bunge, in Lehm. Pl. Preiss. i. 290 = pusilla.
hederifolia, *Benth. Fl. Austral.* iii. 359.—Austral.
hirsuta, DC. Prod. iv. 74 = pilosa.
Huegelii, *Steud. Nom.* ed. II. ii. 791; *Bunge, in Lehm. Pl. Preiss.* i. 291.—Austral.
juncea, *Benth. Fl. Austral.* iii. 359.—Austral.
leiophylla, F. Muell. ex Klatt, in Linnaea, xxix. (1857–58) 710 = dissecta.
montana, Sieber, ex DC. Prod. iv. 74 = pilosa.
pannosa, Steud. Nom. ed. II. ii. 791 = pilosa.
peduncularis, *Benth. Fl. Austral.* iii. 362.—Austral.

XANTHOSIA:—
peltigera, *Steud. Nom.* ed. II. ii. 791; *Benth. Fl. Austral.* iii. 359.—Austral.
pilosa, *Rudge, in Trans. Linn. Soc.* x. (1811) 301. t. 22. f. 1.—Austral.
pinnatisecta, F. Muell. ex Klatt, in Linnaea, xxix. (1857–58) 711 = dissecta.
pusilla, *Bunge, in Lehm. Pl. Preiss.* i. 291.—Austral.
rhomboidea, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 32 = Actinotus rhomboideus.
rotundifolia, *DC. Prod.* iv. 75.—Austral.
singuliflora, *F. Muell. Fragm.* iv. 184.—Austral.
tenuior, *Benth. Fl. Austral.* iii. 359.—Austral.
tridentata, *DC. Prod.* iv. 75.—Austral.
vestita, *Benth. Fl. Austral.* iii. 363.—Austral.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 32 = pusilla.

XANTHOSOMA, Schott, Melet. i. 19 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 976.
ACONTIAS, Schott, Melet. i. 19 (1832).
PHYLLOTAENIUM, André, in Illustr. Hortic. xix. (1872) 3. t. 88.
appendiculatum, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 417 = atrovirens.
atrovirens, *C. Koch & Bouché, in Ind. Sem. Hort. Berol.* (1854) *App.* 3.—Venezuela.
auriculatum, *Regel, Ind. Sem. Hort. Petrop.* 1868 (1869) 81.—Bras.
Barilleti, *Carr. in Rev. Hortic.* (1882) 259. f. 58.—Bras.?
belophyllum, *Kunth, Enum. Pl.* iii. 44; *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 369.—Am. trop.
blandum, Schott, in Bonplandia, x. (1862) 346 = X. Mafaffa.
caracasanum, Schott, Syn. Aroid. 60 = belophyllum.
Caracu, *C. Koch & Bouché, in Ind. Sem. Hort. Berol.* (1854) *App.* 3.—Am. austr.
conspurcatum, *Schott, Syn. Aroid.* 61.—Guian.
cubense, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 33, nomen; *Engl. in DC. Monog. Phan.* ii. 485.—Cuba.
edule, Hort. ex Schott, Syn. Aroid. 70 = Syngonium affine.
edule, *Schott, Melet.* i. 19.—Guian.
gracile, Miq. Hort. Amstelod. (1853); et ex Schott, Syn. Aroid. 70 = Syngonium affine.
hastatum, *Eggers, Fl. Ste-Croix* (1879), 99 (sp. dub.).—Ind. occ.
hastifolium, *C. Koch, in Ind. Sem. Hort. Berol.* (1854) *App.* 2.—Bras.
helleborifolium, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 33.—Am. trop.
Hoffmanni, *Schott, l. c.*—Costa Rica.
Holtonianum, *Schott, l. c.* xix. (1859) 39.—N. Granat.
ianthinum, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854) App. 2 = violaceum.
Jacquini, *Schott, Melet.* i. 19.—Venezuela.
Lindeni, *S. Moore, in Flor. & Pom.* (1872) 4; *Engl. in DC. Monog. Phan.* ii. 482.—N. Granat.
Mafaffa, *Schott, Araceen*, ii. (1855) 5.—Bras.
Maximilianum, *Schott, in Bonplandia*, x. (1862) 322.—Bras.
mexicanum, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 15.—Mexic.
mirabile, *Mast. in Gard. Chron.* (1874) 258. f. 53 (an gen.?).—Am. trop.
mucronatum, Schott, Syn. Aroid. 63 = Alocasia macrorhiza.
pedatum, *Hemsl. Biol. Centr. Am. Bot.* iii. 418.—Guatemala.
pentaphyllum, *Schott, in Oestr. Bot. Zeitzschr.* xv. (1865) 33, nomen; *Engl. in DC. Monog. Phan.* ii. 484.—Bras.
peregrinum, Griseb. Fl. Brit. W. Ind. 511 = atrovirens.
pilosum, *C. Koch & August. in Ind. Sem. Hort. Berol.* (1855) *App.* 2.—Costa Rica.
platylobum, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 33, nomen; *Engl. in Mart. Fl. Bras.* iii. II. 196. t. 45.—Bras.
Plumieri, *Schott, l. c.*—Ind. occ.
Poeppigii, Schott. in Oestr. Bot. Wochenbl. iv. (1854) 417 = X. Mafaffa.
pubescens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 89. t. 299.—Peruv.
Riedelianum, *Schott, in Oestr. Bot. Zeitschr.* xv. (1865) 33.—Bras.
robustum, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 370.—Mexic.

XANTHOSOMA:—
roseum, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 178.—Costa Rica.
sagittifolium, C. Koch, in Ind. Sem. Hort. Berol. (1854) App. 2 = X. Mafaffa.
sagittifolium, Liebm. in Kjoeb. Vidensk. Meddel. (1849) 15 = Jacquini.
sagittifolium, *Schott, Melet.* i. 19.—Am. trop.
striolatum, *Schott, Syn. Aroid.* 61.—Bras.
subandinum, Schott, l. c. 60 = X. Mafaffa.
sylvestre, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 114.—Ins. Porto Rico.
undipes, C. Koch, in Bonplandia, iv. (1856) 4 = Jacquini.
utile, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854) App. 2 = sagittifolium.
variegatum, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 33 = helleborifolium.
violaceum, *Schott, Ind. Sem. Hort. Berol.* (1853) 370. —Ind. occ.
Wallisi, *Hort.—Cf. Gard. Chron.* (1869) 587 (sphalm. *Wallichi*).—N. Granat.
Wendlandii, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 33, nomen = Hoffmanni.
xanthorrhizon, C. Koch, in Bonplandia, iv. (1856) 4 = sagittifolium.

XANTHOSTEMON, F. Muell. in Hook. Kew Journ. ix. (1857) 17. *MYRTACEAE*, Benth. & Hook. f. i. 711.
Draparnaudia, Montr. in Mem. Acad. Lyon, x. (1860) 205.
Fremya, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 372.
Salisia, Panch. ex Brongn. & Gris, l. c. 373 (1863).
chrysanthus, *F. Muell. ex Benth. Fl. Austral.* iii. 268. —Austral.
eucalyptoides, F. Muell. Fragm. i. 81 = Metrosideros eucalyptoides.
Pancheri, *Panch. & Sebert, Not. Bois N. Caléd.* 253.—N. Caled.
paradoxus, *F. Muell. in Hook. Kew Journ.* ix. (1857) 18.—Austral.
ruber, Panch. & Sebert, Not. Bois N. Caléd. 252 = Metrosideros rubra.
Verdugonianus, *Naves, ex F. Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 82.—Ins. Philipp.

XANTHOXYLON, Spreng. Anleit. ii. II. 655 (1818) = seq.

XANTHOXYLUM, J. F. Gmel. Syst. 509 (1791) = **Zanthoxylum**, Linn. (Rutac.).

XANTIUM, Gilib. Fl. Lituan. i. 170 (1781), sphalm. = **Xanthium**, Linn. (Compos.).

XANTOLIS, Rafin. Sylva Tellur. 35 (1838) = **Sideroxylon**, Dill. (Sapot.).
tomentosa, Rafin. l. c. = S. tomentosum.

XANTOPHTALMUM, Sang. in Att. Acc. Pont. Lincei, Ser. I. xvii. (1862) 41, 199 = **Chrysanthemum**, Tourn. (Compos.).

XARITONIA, Rafin. Fl. Tellur. iv. 9 (1836) = **Oncidium**, Sw. (Orchid.).
elegans, Rafin. l. c. = O. quadripetalum.

XATARDIA, Meissn. Gen. 145 (1838). *UMBELLIFERAE*, Benth. & Hook. f. i. 902.
Petitia, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 219.
scabra, *Meissn. Gen.* 145.—Mont. Pyren.

XATARTA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 176 = praec.

XATATIA, Bubani, in Nuov. Ann. Sc. Nat. Bologn. ix. (1843) 92, sphalm. = **Xatardia**, Meissn. (Umbell.).
pyrenaica, Bubani, l. c. = Xatardia scabra.

XAVERIA, Endl. Gen. Suppl. iv. III. 30 (1850) = **Anemonopsis**, Sieb. & Zucc. (Ranunc.).
macrophylla, Endl. ex Walp. Ann. ii. 12 = A. macrophylla.

XEILYATHUM, Rafin. Fl. Tellur. ii. 62 (1836) = **Oncidium**, Sw. (Orchid.).
altissimum, Rafin. l. c. = O. altissimum.

XENIATRUM, Salisb. Gen. Pl. Fragm. 58 (1866) = **Clintonia**, Rafin. (Liliac.).
umbellatum, Salisb. l. c. = C. umbellata.

XENISMIA, DC. Prod. v. 509 (1836) = **Oligocarpus**, Less. (Compos.).
acanthosperma, DC. l. c.—Afr. austr.

XENOCARPUS, Cass. in Dict. Sc. Nat. lix. 108 (1829) = **Cineraria**, Linn. (Compos.).
geifolius, Cass. l. c. 109 = Senecio geifolius.

XENOCHLOA, Licht. in Roem. & Schult. Syst. ii. 29, 501 (1817). Genus *GRAMINEARUM* delendum.—Cf. Benth. & Hook. f. iii. 1096.
arundinacea, Licht. l. c. 501 (= *Phragmites Xenochloa*). —Afr. austr.

XENOPHONTA, Benth. & Hook. f. Gen. ii. 484 (1873), err. typ. = seq.

XENOPHONTIA, Vell. Fl. Flum. 346 (1825); Ic. viii. t. 85 (1827) = **Barnadesia**, Mutis (Compos.).
caryophylla, Vell. ll. cc. 347; t. 85 = B. rosea.
obovata, Regel, Cat. Pl. Hort. Aksakov. 147 (Quid?). —Hab. ?

XENOPHYA, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 124 (1863–64). *AROIDEAE*, Benth. & Hook. f. iii. 972.
brancaefolia, *Schott, l. c.*—N. Guin.

XENOPOMA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399 = **Micromeria**, Benth. (1829) (Labiat.).
bolivianum, Griseb. in Goett. Abh. xxiv. (1879) 272 = M. boliviana.
eugenioides, Griseb. l. c. xix. (1874) 237 = M. eugenioides.
obovatum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399 = M. obovata.
odorum, Griseb. in Goett. Abh. xix. (1874) 236 = M. odora.
verticillatum, Griseb. l. c. 237 = Bystropogon mollis.

XEODOLON, Salisb. Gen. Pl. Fragm. 18 (1866) = = **Scilla**, Linn. (Liliac.).
revolutum, Salisb. l. c. = S. revoluta.

XERACINA, Rafin. Sylva Tellur. 98 (1838) = **Adelobotrys**, DC. etc. (Melastom.).
aquatica, Rafin. l. c. = Nepsera aquatica.
scandens, Rafin. l. c. = A. scandens.
villosa, Rafin. l. c. = Desmoscelis villosa.

XERALIS, Rafin. Sylva Tellur. 82 (1838) = **Charachera**, Forsk. (Inc. sed.).

XERALSINE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = **Arenaria**, Rupp. (Caryoph.).
fasciculata, Fourr. l. c. = A. fasciculata.
mucronata, Fourr. l. c. = A. mucronata.

XERANDRA, Rafin. Fl. Tellur. iii. 43 (1836) = **Iresine**, P. Br. (Amarant.).
celosioides, Rafin. l. c. = I. celosioides.
elatior, Rafin. l. c. = I. elatior.

XERANTHEMUM, Tourn. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 463.
Harrisonia, Neck. Elem. i. 84 (1790).
Xeroloma, Cass. in Dict. Sc. Nat. lix. 120 (1829).
angustifolium, Lam. Encyc. iii. 241 = Helichrysum paniculatum.
Annettae, Kalen. in Bull. Soc. Nat. Mosc. (1834) 194 = squarrosum.
annuum, Asso, Syn. Pl. Arag. 118 = inapertum.
annuum, *Linn. Sp. Pl.* 957.—Europ. austr.
annuum, Scop. Fl. Carn. ed. II. ii. 180 = cylindraceum.
argenteum, Thunb. Prod. Pl. Cap. 152 = Helichrysum argenteum.
australe, Pomel, Nouv. Mat. Fl. Atl. 33 = inapertum.
bellidioides, Forst. f. Prod. 55 = Helichrysum bellidioides.
bracteatum, Vent. Jard. Malm. 2 = Helichrysum bracteatum.
caespitosum, Thou. Fl. Tristan d'Acugn. 39. t. 8 (*cespitosum*) = Chevreulia stolonifera.
canescens, Linn. Pl. Afr. Rar. 20; Amoen. Acad. vi. 100 = Helipterum canescens.

XERANTHEMUM:—
chinense, *Lour. Fl. Cochinch.* 498.—China.
ciliatum, Linn. Sp. Pl. 859 = Cullumia decurrens.
corymbosum, Lam. Encyc. iii. 242 = Helichrysum fulgidum.
crucifolium, Hill, Veg. Syst. ii. 95, sphalm. = *cruciifolium*.
cylindraceum, Colm. Apunt. Fl. Castill. 85 = inapertum.
cylindraceum, *Sm. Fl. Graec. Prod.* ii. 172.—Europ. austr.; As. Min.
cylindricum, Spreng. Syst. iii. 485 = praec.
erectum, Presl, Delic. Prag. 106 = inapertum.
ericoides, Lam. Encyc. iii. 240 = Helichrysum ericoides.
erucifolium, Linn. Sp. Pl. 858 = Serratula xeranthemoides.
fasciculatum, Andr. Bot. Rep. t. 242 = Helichrysum sesamoides.
ferrugineum, Lam. Encyc. iii. 237 = Helipterum ferrugineum.
foetidum, Moench, Meth. 582 = inapertum.
fragrans, *Andr. Bot. Rep.* t. 561.—Afr. austr.
frigidum, Labill. Ic. Pl. Syr. Dec. ii. 9 = Helichrysum frigidum.
fulgidum, Linn. f. Suppl. 365 = Helichrysum fulgidum.
herbaceum, Andr. Bot. Rep. t. 487 = Helichrysum squamosum.
heterophyllum, Lam. Encyc. iii. 239 = Helichrysum sesamoides.
humile, Andr. Bot. Rep. t. 652 = Helichrysum humile.
imbricatum, Burm. f. Fl. Cap. Prod. 25 = Metalasia uniflora.
imbricatum, Linn. Pl. Rar. Afr. 20; Amoen. Acad. vi. 100 = Helipterum canescens.
inapertum, Bieb. Fl. Taur. Cauc. ii. 302 = cylindraceum.
inapertum, *Mill. Gard. Dict.* ed. VIII. n. 2; *Willd. Sp. Pl.* iii. 1902.—Reg. Mediterr. et Cauc.
incanum, Crantz, Inst. i. 290 = Helipterum canescens.
incomptum, Cass. in Dict. Sc. Nat. lix. 115 = inapertum.
inodorum, Moench, Meth. 582 = annuum.
lancifolium, Thunb. Prod. Pl. Cap. 152 = Helichrysum lancifolium.
leucanthemifolium, Burm. f. Fl. Cap. Prod. 26 = Conyza pinnatilobata.
longepapposum, *Fisch. & Mey. in Nouv. Mém. Soc. Nat. Mosc.* iv. (1835) 337.—As. Min.; Persia.
modestum, *Ball, in Journ. Bot.* xi. (1873) 368.—Marocc.
moniliforme, Houtt. ex DC. Prod. vi. 172 = Helichrysum ericoides.
Neumayeri, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 266.—Europ. austr.
nobile, *Grauer, Dec.* 16.—Afr. austr.
orientale, Mill. Gard. Dict. ed. VIII. n. 3 = Chardinia xeranthemoides.
ornatum, Cass. in Dict. Sc. Nat. lix. 114 = annuum.
paniculatum, Linn. Sp. Pl. 859 = Helichrysum paniculatum.
pinifolium, Lam. Encyc. iii. 240 = Helichrysum humile.
polyfolium, Crantz, Inst. i. 290 = Helichrysum retortum.
proliferum, Linn. Sp. Pl. 858 = Phaenocoma prolifera.
pungens, Lam. Encyc. iii. 236 = Siebera pungens.
pungens, Steud. Nom. ed. II. ii. 791 = Serratula xeranthemoides.
radiatum, Lam. Fl. Fr. ii. 48 = annuum.
radicans, Thunb. Prod. Pl. Cap. 153 = Helichysum retortum.
ramosum, *Burm. f. Fl. Cap. Prod.* 26 (Quid?).—Afr. austr.
recurvatum, Linn. f. Suppl. 366 = Helichrysum recurvatum.
regale, Salisb. Prod. 193 = Phaenocoma prolifera.
retortum, Linn. Sp. Pl. 858 = Helichrysum retortum.
retortum, Lour. Fl. Cochinch. 498 = Helichrysum cochinchinense.
rigidum, Andr. Bot. Rep. t. 387 = Helichrysum striatum.
serpyllifolium, Lam. Encyc. iii. 238 = Helipterum canescens.
sesamoides, J. Gay, ex Willk. & Lange, Prod. Fl. Hisp. ii. 128 = cylindraceum.
sesamoides, Linn. Sp. Pl. 859 = Helichrysum sesamoides.

XERANTHEMUM:—
speciosissimum, Linn. Sp. Pl. 858 = Helipterum speciosissimum.
speciosum, L'Hérit. ex DC. Prod. vi. 530 = Chardinia xeranthemoides.
speciosum, Thunb. Prod. Pl. Cap. 153 = Helipterum speciosissimum.
spinosum, Linn. Sp. Pl. 859 = Dicoma Burmanni.
spirale, Andr. Bot. Reg. t. 262 = Helipterum variegatum.
squamosum, Thunb. Prod. Pl. Cap. 152 = Helichrysum squamosum.
squarrosum, *Boiss. Diagn.* Ser. I. vi. 101.—As. Min.; Persia.
squarrosum, Lam. Encyc. iii. 241 = Metalasia umbellata.
Staehelina, DC. Prod. vi. 212, sphalm. = Helipterum virgatum.
Staehelina, *Linn. Syst.* ed. XII. 546; *Thunb. Prod. Pl. Cap.* 153.—Afr. austr.
stoloniferum, Linn. f. Suppl. 366 = Helichrysum stoloniferum.
striatum, *Thunb. Prod. Pl. Cap.* 152.—Afr. austr.
tenuifolium, Burm. f. Fl. Cap. Prod. 26 = Helichrysum mucronatum.
variegatum, Berg, Desc. Pl. Cap. 271 = Helichrysum vestitum.
vermiculatum, Lam. Encyc. iii. 240 = Metalasia uniflora.
vestitum, Linn. Sp. Pl. 858 = Helichrysum vestitum.
vestitum, Steud. Nom. ed. II. ii. 791 = Helipterum variegatum.
virgatum, Berg, Desc. Pl. Cap. 275 = Helipterum virgatum.

XERANTHIUM, Lepech. Reise Russ. i. 240 (1774) = **Xanthium**, Linn. (Compos.).

XERANTHUS, Miers, Trav. Chil. ii. 529 (1826), nomen = **Grahamia**, Gill. (Portulac.).
salicosus, Miers, l. c. = G. bracteata.

XERIS, Medic. in Staatsw. Vorles. Churpf. Phys-Oek. Ges. i. (1791) 256 = **Iris**, Tourn.

XEROBIUS, Cass. in (Bull. Soc. Philom. (1817) 153; fide Pfeiff. Nom. ii. 1629) Dict. Sc. Nat. lix. 127 (1829) = **Egletes**, Cass. (Compos.).
lanatus, Cass. l. c. 128 = E. bellidiflora.

XEROBOTRYS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 267 = **Arctostaphylos**, Adans. (Ericac.).
argutus, Nutt. l. c. 268 = A. tomentosa.
cordifolius, Nutt. l. c. = A. tomentosa.
glaucus, Nutt. l. c. = A. glauca.
tomentosus, Nutt. l. c. 267 = A. tomentosa.
venulosus, Nutt. l. c. 268 = A. Hookeri.

XEROCARPA, Spach, Hist. Vég. Phan. ix. 583 (1840) = **Scaevola**, Linn. (Gooden.).

XEROCARPUS, Guill. & Perr. Fl. Seneg. Tent. 169. t. 44 (1832) = **Rothia**, Pers. (Legumin.).
abyssinicus, Schweinf. Beitr. Pl. Aethiop. 21 = Lotononis Leobordea.
hirsutus, Guill. & Perr. Fl. Seneg. Tent. 170 = R. hirsuta.

XEROCHLAMYS, Baker, in Journ. Bot. xx. (1882) 45. *CHLAENACEAE.*
pilosa, *Baker, l. c.*—Madag.

XEROCHLOA, R. Br. Prod. 196 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1108.
barbata, *R. Br. l. c.* 197.—Austral.
imberbis, *R. Br. l. c.*—Austral.
laniflora, *Benth. Fl. Austral.* vii. 502.—Austral.
latifolia, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 2; *Cat. Hort. Bog. Alt.* 21.—Java.

XEROCLADIA, Harv. in Harv. & Sond. Fl. Cap. ii. 278 (1861-62). *LEGUMINOSAE*, Benth. & Hook. f. i. 591.
Zeyheri, *Harv. l. c.*—Afr. austr.

XEROCOCCUS, Oerst. in Kjoeb. Vidensk. Meddel. (1852) 52. *RUBIACEAE*, Benth. & Hook. f. ii. 70.
congestus, *Oerst. l. c.* 52.—Costa Rica.

XERODERA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 324 = **Ranunculus**, Tourn.
graminea, Fourr. l. c. = R. gramineus.

XEROGONA, Rafin. Fl. Tellur. iv. 103 (1836) = **Passiflora**, Linn.
biloba, Rafin. l. c. = P. capsularis.

XEROLOLOPHUS, Dulac, Fl. Hautes-Pyr. 160 (1867) = **Thesium**, Linn. (Santal.).
alpinus, Dulac, l. c. 161 = T. alpinum.
pratense, Dulac, l. c. = T. pratense.
trinervis, Dulac, l. c. = T. intermedium.

XEROLOMA, Cass. in Dict. Sc. Nat. lix. 120 (1829) = **Xeranthemum**, Tourn. (Compos.).
foetidum, Cass. l. c. 121 = Xeranth. cylindraceum.
inapertum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 98 = Xeranth. inapertum?

XEROMALON, Rafin. New Fl. Am. iii. 11 (1836) = **Amelanchier**, Medic. (Rosac.).
obovatum, Rafin. l. c. 12 = A. canadensis.

XEROMPHIS, Rafin. Sylva Tellur. 21 (1838) = **Randia**, Houst. (Rubiac.).
Retzi, Rafin. l. c. = A. dumetorum.

XERONEMA, Brongn. & Gris, in Bull. Soc. Bot. Fr. xi. (1864) 316. *LILIACEAE*, Benth. & Hook. f. iii. 781.
SCLERONEMA, Brongn. & Gris, in Ann. Sc. Nat. Sér. V. ii. (1864) 166.
Moorei, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 317.—N. Cal.

XEROPAPPUS, Wall. Cat. n. 2980 (1830?) = **Dicoma**, Cass. (Compos.).
lanuginosus, Wall. l. c. = D. tomentosa.

XEROPETALON, Hook. Bot. Misc. i. (1830) 174 = **Vivianía**, Cav. (Geran.).

XEROPETALUM, Delile, Cent. Pl. Méroé, 84 (1826) = **Dombeya**, Cav. (Stercul.).
Brucei, Hochst. in Flora, xxiv. (1841) 1. Intell. 29, nomen = D. Bruceana.
hirsutum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 78 = D. Schimperiana.
minus, Endl. Nov. Stirp. Dec. 37 = D. multiflora.
multiflorum, Endl. l. c. 36 = D. multiflora.
quinquesetum, Delile, Cent. Pl. Méroé, 84 = D. multiflora?
rotundifolium, Hochst. in Flora, xxvii. (1844) 295 = D. rotundifolia.
tiliaceum, Endl. Nov. Stirp. Dec. 37 = D. tiliacea.

XEROPETALUM, Reichb. Consp. 154 (1828) = **Dillwynia**, Sm. (Legumin.).
cinerascens, Steud. Nom. ed. II. ii. 791 = D. cinerascens.

XEROPHYLLUM, Michx. Fl. Bor. Am. i. 210 (1803). *LILIACEAE*, Benth. & Hook. f. iii. 826.
asphodeloides, *Nutt. Gen. Am.* i. 235.—Am. bor.
asphodeloides, Spreng. Syst. ii. 144 = setifolium.
Douglasii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 284.—Am. bor. occ.
gramineum, Nutt. Gen. Am. i. 235 = Stenanthium angustifolium.
Sabadilla, D. Don, ex Sweet, Hort. Brit. ed. III. 712 = Veratrum Sabadilla.
setifolium, *Michx. Fl. Bor. Am.* i. 211.—Am. bor.
tenax, Nutt. Gen. Am. i. 235 = setifolium.

XEROPHYSA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 150 = **Astragalus**, Tourn. (Legumin.).
physodes, Stev. l. c. = A. physodes.

XEROPHYTA, Juss. Gen. 50 (1789) = **Vellozia**, Vand. (Amaryll.).
abietina, Spreng. Syst. iv. Cur. Post. 137 = V. abietina.
capillaris, Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 264 = V. capillaris.
cinerascens, Mart. ex Schult. f. Syst. vii. 292 = V. cinerascens.
clavata, Baker, in Journ. Bot. xiii. (1875) 233.—Afr. austr.

XEROPHYTA:—
dasylirioides, Baker, in Journ. Bot. xiii. (1875) 235.—Madag.
elegans, Baker, l. c. 234 = V. elegans.
equisetoides, Baker, l. c. 233.—Afr. austr.
madagascariensis, J. F. Gmel. Syst. 530.—Madag.
Melleri, Baker, in Journ. Bot. xiii. (1875) 234.—Afr. trop.
minima, Baker, l. c. 235 = V. minima.
minuta, Baker, l. c. 234.—Afr. austr.
neglecta, Schult. f. Syst. vii. 289.—Ins. Franciae.
pectinata, Baker, in Journ. Bot. xx. (1882) 270.—Madag.
pinifolia, Lam. ex Poir. Encyc. viii. 804 = *madagascariensis.*
plicata, Spreng. Syst. iv. Cur. Post. 137 = V. plicata.
Pohliana, Schult. f. Syst. vii. Add. 1666 = V. minima.
retinervis, Baker, in Journ. Bot. xiii. (1875) 233.—Afr. austr.
Schnitzleinia, Baker, l. c. 235.—Abyss.
Selloi, Baker, l. c. 236 = V. Sellovii.
sessiliflora, Baker, l. c. xx. (1882) 271.—Madag.
Spekei, Baker, l. c. xiii. (1875) 234.—Afr. trop.
spinulosa, Ridl. in Journ. Linn. Soc. xx. (1883) 333.—Madag.
squarrosa, Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 264 = V. squarrosa.
stenophylla, Baker, l. c. 265 = V. stenophylla.
taxifolia, Mart. ex Schult. f. Syst. vii. 291 = V. pusilla?
tragacantha, Mart. ex Schult. f. l. c. 290 = V. tragacantha.
triquetra, Baker. in Journ. Bot. xiii. (1875) 236 = V. triquetra.
velutina, Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 265 = V. velutina.
viscosa, Baker, in Journ. Bot. xiii. (1875) 235.—Afr. austr.

XEROSIPHON, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 55 = **Gomphrena**, Linn. (Amarant.).
gracilis, Turcz. l. c. = G. angustiflora.

XEROSOLLYA, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 362 = **Sollya**, Lindl. (Pittosp.).
Gilbertii, Turcz. l. c. = S. parviflora?

XEROSPERMUM, Blume, Rumphia, iii. 99 (1850). *SAPINDACEAE*, Benth. & Hook. f. i. 405.
acuminatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 95.—Borneo.
glabratum, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 300, 310; ix. (1879) 516 = Noronhianum.
laevigatum, *Radlk. l. c.* 305, 310 (Sapindi sp.?).—Ind. or.
lanceolatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 95.—Borneo.
muricatum, *Radlk. l. c.*—Ind. or.
Noronhianum, *Blume, Rumphia*, iii. 100.—Malaya.

XEROTES, R. Br. Prod. 252 (1810). *JUNCACEAE*, Benth. & Hook. f. iii. 863.
LOMANDRA, Labill. Nov. Holl. Pl. i. 92 (1804), nomen prius.
aemula, R. Br. Prod. 262 = multiflora.
arenaria, R. Br. l. c. = longifolia.
asparagoides, Endl. in Lehm. Pl. Preiss. ii. 49 = pauciflora.
Banksii, *R. Br. Prod.* 263.—Austral.
Brownii, F. Muell. Fragm. viii. 206 = multiflora.
caespitosa, *Benth. Fl. Austral.* vii. 104.—Austral.
collina, *R. Br. Prod.* 260.—Austral.
decomposita, R. Br. l. c. 262 = multiflora.
denticulata, R. Br. l. c. 261 = filiformis.
distans, R. Br. l. c. 262 = multiflora.
Drummondii, *F. Muell. ex Benth. Fl. Austral.* vii. 99.—Austral.
dura, *F. Muell. in Trans. Vict. Inst.* i. (1855) 42.—Austral.
echinata, A. Cunn. ex Benth. Fl. Austral. vii. 112 = Acanthocarpus Preissii.
effusa, *Lindl. in Mitch. Three Exped.* ii. 101.—Austral.
elongata, *Benth. Fl. Austral.* vii. 106.—Austral.
Endlicheri, *F. Muell. Fragm.* viii. 205.—Austral.
filamentosa, Brongn. in Duper. Voy. Coq. Bot. 183. t. 35 = leucocephala.
filiformis, *R. Br. Prod.* 261.—Austral.
fimbriata, F. Muell. Fragm. viii. 211 = Chamaexeros fimbriata.

XEROTES:—
flexifolia, *R. Br. Prod.* 260.—Austral.
fluviatilis, R. Br. l. c. 262 = longifolia.
fragrans, F. Muell. ex Sond. in Linnaea, xxviii. (1856) 219 = effusa.
glauca, *R. Br. Prod.* 260.—Austral.
gracilis, R. Br. l. c. 261 = filiformis.
graminea, Endl. in Lehm. Pl. Preiss. ii. 48 = pauciflora.
hastilis, *R. Br. Prod.* 263.—Austral.
Hystrix, R. Br. l. c. 262 = longifolia.
juncea, *F. Muell. in Trans. Vict. Inst.* i. (1855) 135.—Austral.
latifolia, Steud. Nom. ed. II. ii. 65, in syn., sphalm. = longifolia.
laxa, R. Br. Prod. 261 = filiformis.
leucocephala, *R. Br. l. c.* 260.—Austral.
longifolia, *R. Br. l. c.* 262.—Austral.
media, R. Br. l. c. = multiflora.
micrantha, *Endl. in Lehm. Pl. Preiss.* ii. 49.—Austral.
montana, R. Br. Prod. 262 = longifolia.
mucronata, R. Br. l. c. 260 = Acanthocarpus Preissii ?
mucronata, Sieber, ex Benth. Fl. Austral. vii. 106 = glauca.
multiflora, *R. Br. Prod.* 262.—Austral.
odora, *Endl. in Lehm. Pl. Preiss.* ii. 50.—Austral.
Ordii, *F. Muell. Fragm.* xi. 23.—Austral.
pauciflora, *R. Br. Prod.* 261.—Austral.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 50.—Austral.
purpurea, *Endl. l. c.* 49.—Austral.
rigida, *R. Br. Prod.* 261.—Austral.
rigida, Endl. in Lehm. Pl. Preiss. ii. 50 = Sonderi.
rupestris, *Endl. l. c.*—Austral.
sericea, *Endl. l. c.* 51.—Austral.
Serra, Endl. l. c. 49 = Chamaexeros Serra.
Sonderi, *F. Muell. Fragm.* viii. 206.—Austral.
sororia, *F. Muell. First Gen. Rep.* 15; *Benth. Fl. Austral.* vii. 100.—Austral.
spartea, *Endl. in Lehm. Pl. Preiss.* ii. 51.—Austral.
suaveolens, *Endl. l. c.* 50.—Austral.
tenuifolia, R. Br. Prod. 261 = filiformis.
tenuifolia, Endl. in Lehm. Pl. Preiss. ii. 49 = Endlicheri.
Thunbergii, F. Muell. Fragm. viii. 208 = filiformis.
turbinata, *Endl. in Lehm. Pl. Preiss.* ii. 51.—Austral.
typhina, Lindl. in Mitch. Three Exped. ii. 41 = leucocephala.
umbrosa, Endl. in Lehm. Pl. Preiss. ii. 50 = suaveolens.

XEROTHAMNUS, DC. Prod. v. 311 (1836) = **Osteospermum**, Linn. (Compos.).
Ecklonianus, DC. l. c. = O. scabrum.

XEROTIS, Hoffmgg. Verz. Pfl. Nachtr. ii. 38 (1826) = **Xerotes**, R. Br. (Juncac.).

XEROTIUM, Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 343 (1825) = **Filago**, Linn. (Compos.).
gallicum, Bluff & Fingerh. l. c. 344 = F. gallica.
minimum, Bluff & Fingerh. l. c. 345 = F. minima.
montanum, Bluff & Fingerh. l. c. 344 = F. minima.
subulatum, Dulac, Fl. Hautes-Pyr. 514 = F. gallica.

XESTAEA, Griseb. in Linnaea, xxii. (1849) 35 = **Schultesia**, Mart. (Gentian.).
lisianthoides, Griseb. l. c. 36 = S. lisianthoides.

XETOLA, Rafin. Fl. Tellur. iv. 95 (1836) = **Cephalaria**, Schrad. (Dipsac.).
transylvanica, Rafin. l. c. = C. transylvanica.
trenta, Rafin. l. c. = C. transylvanica.

XETOLIGUS, Rafin. New Fl. Am. iv. 74 (1836) = **Stevia**, Cav. (Compos.).
brevifolius, Rafin. l. c.—Am. bor.
salicifolius, Rafin. l. c. = S. salicifolia.

XILOPIA, Juss. Gen. 284 (1789) = **Xylopia**, Linn. (Anonac.).

XIMENESIA, Cav. Ic. ii. 60. t. 178 (1793) = **Verbesina**, Linn. (Compos.).
australis, Hook. & Arn. ex DC. Prod. vii. 291 = V. australis.
Cavanillesii Spreng. Syst. iii. 605 = Iostephane heterophylla.

XIMENESIA:—
cordata, H. B. & K. Nov. Gen. et Sp. iv. 228 = Encelia cordata.
encelioides, Cav. Ic. ii. 60. t. 178 = Verbesina encelioides.
foetida, Spreng. Syst. iii. 606 = Encelia foetida.
heterophylla, H. B. & K. Nov. Gen. et Sp. iv. 227. t. 280 = Encelia heterophylla.
hirta, Mart. ex DC. Prod. v. 578 = Encelia mexicana.
hispida, H. B. & K. Nov. Gen. et Sp. iv. 227 = Encelia hispida.
microptera, DC. Prod. v. 627 = V. australis.
procumbens, Boj. ex DC. l. c. 561 = Aspilia Bojeri.

XIMENIA, Plum. ex Linn. Gen. ed. I. 361 (1737). *OLACINEAE*, Benth. & Hook. f. i. 346.
HEYMASSOLI, Aubl. Pl. Gui. 324. t. 125 (1775).
PIMECARIA, Rafin. Alsog. Am. 64 (1838).
ROTTBOELIA, Scop. Introd. 233 (1777).
aculeata, Crantz, Inst. ii. 381 = americana.
aegyptiaca, Juss. Gen. 260 = Balanites Roxburghii.
aegyptiaca, Linn. Sp. Pl. 1194 = Balanites aegyptiaca.
Agihalid, Mill. Gard. Dict. ed. VIII. n. 2 = Balanites aegyptiaca.
americana, *Linn. Sp. Pl.* 1193.—Cosmop. trop.
arborea, Royle, Illustr. Bot. Himal. 154 = Zygophyllum arboreum.
arborescens, Tussac, ex Walp. Rep. i. 377, sphalm. = americana.
borneensis, Baill. Adansonia, xi. (1873–76) 271 = Scorodocarpus borneensis.
caffra, *Sond. in Linnaea*, xxiii. (1850) 21.—Afr. austr.
coriacea, *Engl. in Mart. Fl. Bras.* xii. II. 10.—Bras.
elliptica, Forst. f. Prod. 27 = americana.
elliptica, Spreng. Syst. ii. 217, partim = verrucosa.
exarmata, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 22.—Austral.
ferox, Poir. Encyc. viii. 805 = Balanites aegyptiaca.
fluminensis, M. Roem. Syn. Hesper. 22 = americana.
inermis, Linn. Sp. Pl, ed. II. 497 = americana.
lanceolata, DC. Prod. i. 533 = Acronychia laurifolia.
laurina, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 89 = americana.
loranthifolia, Span. in Linnaea, xv. (1841) 177 = americana.
montana, Macfad. Fl. Jamaic. i. 121 = americana.
multiflora, Jacq. Enum. Pl. Carib. 19 = americana.
oblonga, Lam. ex Hemsl. Biol. Centr. Am. Bot. i. 185 = americana.
oblongifolia, *Steud. Nom.* ed. II. ii. 792.—Austral.
olacoides, Wight & Arn. Prod. 89 = Opilia amentacea.
parviflora, *Benth. Pl. Hartw.* 7.—Mexic.
rigidum, Royle, Illustr. Bot. Himal. 154, 466 = Zygophyllum rigidum.
Russeliana, Wall. Cat. n. 6784 = americana.
spinosa, Salisb. Prod. 276 = americana.
subscandens, *Griff. Notul.* iv. 691.—Ind. or.
verrucosa, *M. Roem. Syn. Hesper.* 23.—Ins. S. Doming.

XIPHIDIUM, Loefl. It. Hisp. 179 (1758); Aubl. Pl. Gui. 33. t. 11 (1775). *HAEMODORACEAE*, Benth. & Hook. f. iii. 675.
albidum, Lam. Illustr. i. 131 = caeruleum.
album, Willd. Sp. Pl. i. 248 = caeruleum.
angustifolium, Willd. ex Link, Jahrb. i. III. (1820) 73 = Wachendorfia orinocensis.
coeruleum, *Aubl. Pl. Gui.* i. 33. t. 11.—Am. trop.
floribundum, Sw. Prod. Veg. Ind. Occ. 17 = caeruleum.
giganteum, Lindl. Bot. Reg. (1846) t. 66 = coeruleum.
rubrum, *D. Don, in Edinb. N. Phil. Journ.* (Oct. 1832) 235.—Peruv.
xanthorrhizon, *Wright, ex Griseb. Cat. Pl. Cub.* 252.—Cuba.

XIPHION, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Iris**, Tourn.
acoroides, Alef. in Bot. Zeit. xxi. (1863) 297 = I. Pseudacorus.
Aitchisoni, Baker, in Journ. Bot. xiii. (1875) 108 = I. Aitchisoni.
alatum, Baker, l. c. ix. (1871) 108 = I. alata.
Aucheri, Baker, l. c. 110 = I. fumosa.
brachycuspis, Alef. in Bot. Zeit. xxi. (1863) 297 = I. setosa.

XIPHION:—
caucasicum, Baker, in Journ. Bot. ix. (1871) 109 = Iris caucasica.
Danfordiae, Baker, l. c. xiv. (1876) 265 = I. Danfordiae.
diversifolium, Klatt, in Linnaea, xxxiv. (1865–66) 572 = Moraea diversifolia.
Donianum, Alef. in Bot. Zeit. xxi. (1863) 297 = I. ensata.
filifolium, Klatt, in Linnaea, xxxiv. (1865–66) 571 = I. filifolia.
fimbriatum, Alef. in Bot. Zeit. xxi. (1863) 297 = I. japonica.
flaccidum, Alef. l. c. = I. versicolor.
flexuosum, Alef. l. c. = I. sibirica.
foetidissimum, Parl. Nuov. Gen. e Spec. 45 = I. foetidissima.
Fontanesii, Baker, in Gard. Chron. (1876) 559 = I. Fontanesii.
gramineum, Schrank, in Flora, vii. (1824) II. Beibl. 17 = I. graminea.
Gueldenstedtii, Schrank, l. c. 18 = I. Gueldenstaedtiana.
Histrio, Hook. f. Bot. Mag. t. 6033 = I. Histrio.
Jacquini, Schrank, in Flora, vii. (1824) II. Beibl. 17 = I. xiphioides.
junceum, Klatt, in Linnaea, xxxiv. (1865–66) 570 = I. juncea.
Kolpakowskianum, Baker, in Bot. Mag. t. 6489 = I. Kolpakowskiana.
Krelagi, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 361 = I. reticulata.
latifolium, Mill. Gard. Dict. ed. VIII. n. 3; Alef. in Bot. Zeit. xxi. (1863) 297 = I. xiphioides.
lusitanicum, Alef. l. c. = I. juncea.
melliftuum, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 419, nomen (Quid?).—Hab. ?
Monnieri, Alef. in Bot. Zeit. xxi. (1863) 297 = I. Monnieri.
nothum, Alef. l. c. = I. spuria.
ochroleucum, Alef. l. c. = I. ochroleucum.
ochroleucum, Schrank, in Flora, vii. (1824) II. Beibl. 18 = I. Gueldenstaedtiana.
orientale, Schrank, l. c. 19 = I. sibirica.
palaestinum, Baker, in Journ. Bot. ix. (1871) 108 = I. palaestina.
Pallasii, Alef. in Bot. Zeit. xxi. (1863) 297 = I. ensata.
papilionaceum, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 174 = Moraea papilionacea.
persicum, Mill. Gard. Dict. ed. VIII. n. 1 = I. persica.
planifolium, Mill. l. c. n. 4 = I. alata.
pratense, Parl. Nuov. Gen. e Spec. 45 = I. sibirica.
Pseudo-acorus, Schrank, Fl. Monac. i. t. 99 = I. Pseudacorus.
reticulatum, Klatt, in Linnaea, xxxiv. (1865–66) 572 = I. reticulata.
ruthenicum, Alef. in Bot. Zeit. xxi. (1863) 297 = I. ruthenica.
sibiricum, Schrank, in Flora, vii. (1824) II. Beibl. 19 = I. sibirica.
Sisyrinchium, Baker, in Journ. Bot. ix. (1871) 42 = I. Sisyrinchium.
sordidum, Salisb. in Trans. Hort. Soc. i. (1812) 303 = I. juncea.
spurium, Alef. in Bot. Zeit. xxi. (1863) 297 = I. spuria.
stenogynum, Alef. l. c. 297 = I. Gueldenstaedtiana.
Stocksii, Baker, in Gard. Chron. (1876) 723 = I. Stocksii.
tenuifolium, Schrank, in Flora, vii. (1824) II. Beibl. 19, 20 = I. tenuifolia.
tingitanum, Baker, in Journ. Bot. ix. (1871) 13 = I. tingitana.
tingitanum, Hook. f. Bot. Mag. t. 5981 = I. filifolia.
triflorum, Alef. in Bot. Zeit. xxi. (1863) 297 = I. ensata.
tripetalum, Alef. l. c. = I. tripetala.
versicolor, Alef. l. c. = I. versicolor.
verum, Schrank, in Flora, vii. (1824) II. Beibl. 16 = I. Xiphium.
virginicum, Alef. in Bot. Zeit. xxi. (1863) 297 = I. versicolor.
vulgare, Mill. Gard. Dict. ed. VIII. n. 2 = I. Xiphium.

XIPHIUM, Mill. Gard. Dict. ed. VII. (1759) = praec.

XIPHIZUSA, Reichb. f. in Bot. Zeit. ix. (1852) 919 = **Bulbophyllum**, Thou. (Orchid.).
chloroptera, Reichb. f. l. c. 919 = B. meridense.
gladiata, Reichb. f. l. c. = B. meridense.
Weddelii, Reichb. f. l. c. = B. Weddelii.

XIPHOCARPUS, Presl, Symb. Bot. i. 13. t. 7 (1830) = **Tephrosia**, Pers. (Legumin.).
candidus, Endl. Gen. 1273; Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 76 = T. candida.
martinicensis, Presl, Symb. Bot. i. 14. t. 7 = T. candida.
minor, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 76.—Java.

XIPHOCHAETA, Poepp. & Endl. Nov. Gen. et Sp. iii. 44. t. 250 (1842) = **Stilpopappus**, Mart. (Compos.).
aquatica, Poepp. & Endl. l. c. = S. viridis.

XIPHOCOMA, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 270, in textu; xxv. (1852) I. 537. t. 7 = **Ranunculus**, Tourn.
dasycarpa, Stev. l. c. (1852) 539. t. 7. f. 3 = R. dasycarpus.
heterophylla, Stev. l. c. f. 2 = R. orientalis.
leptalea, Stev. l. c. 541 = R. leptaleus.
orientalis, Stev. l. c. 538 = R. orientalis.
tenuifolia, Stev. l. c. 540. t. 7. f. 4 = R. orientalis.

XIPHOLEPIS, Steetz, in Peters, Reise Mossamb. Bot. 344 (1863) = **Vernonia**, Schreb. (Compos.).
aspera, Steetz, l. c. = V. Roxburghii.
silhetensis, Steetz, l. c. 345 = V. bracteata.

XIPHOPHYLLUM, Ehrh. Beitr. iv. 148 (1789) = **Cephalanthera**, Rich. (Orchid.).

XIPHOSIUM, Griff. in Calc. Journ. Nat. Hist. v. 364 (1845) = **Eria**, Lindl. (Orchid.).
acuminatum, Griff. l. c. = E. carinata.
roseum, Griff. l. c. = E. rosea.

XIPHOTHECA, Eckl. & Zeyh. Enum. 166 (1836) = **Priestleya**, DC. (Legumin.).
axillaris, Eckl. & Zeyh. l. c. 167 = Amphithalea cuneifolia.
lanceolata, Eckl. & Zeyh. l. c. = P. sericea?
polycarpa, Eckl. & Zeyh. l. c. 166 = P. tecta.
rotundifolia, Eckl. & Zeyh. l. c. = P. tecta.
sericea, Eckl. & Zeyh. l. c. 167 = P. sericea.
tomentosa, Eckl. & Zeyh. l. c. 166 = P. villosa.
villosa, Eckl. & Zeyh. l. c. = P. villosa.

XOLANTHA, Rafin. Caratteri, 73 (1810) = **Helianthemum**, Tourn. (Cistin.).
racemosa, Rafin. l. c. = H. variabile.

XOLANTHES, Rafin. Sylva Tellur. 132 (1838) = **Helianthemum**, Tourn. (Cistin.).
guttatus, Rafin. l. c. = H. guttatum.
racemosa, Rafin. l. c. = H. variabile.

XOLEMIA, Rafin. Fl. Tellur. iii. 22 (1836) = **Gentiana**, Tourn.
acuminata, Rafin. l. c. (Quid?).—Hab.?
albiflora, Rafin. l. c. = G. albiflora.
axillaris, Rafin. l. c. = G. axillaris.
Catesbei, Rafin. l. c. = G. Saponaria.
clausa, Rafin. l. c. = G. Andrewsii.
Collinsiana, Rafin. l. c. = G. Collinsiana.
Elliotea, Rafin. l. c. = G. Elliottea.
enervis, Rafin. l. c. (Quid?).—Hab.?
fistulosa, Rafin. l. c. (Quid?).—Hab.?
heterophyla, Rafin. l. c. = G. ochroleuca.
latifolia, Rafin. l. c. (Quid?).—Hab.?
longiflora, Rafin. l. c. = G. altaica.
obovata, Rafin. l. c. (Quid?).—Hab.?
palustris, Rafin. l. c. = Erythraea ramosissima.
rotundifolia, Rafin. l. c. = G. bavarica.
rubella, Rafin. l. c. (Quid?).—Hab.?
serpentaria, Rafin. l. c. = G. ochroleuca.
Shortiana, Rafin. l. c. = G. Shortiana.
striata, Rafin. l. c. (Quid?).—Hab.?
terniflora, Rafin. l. c. (Quid?).—Hab.?
trachiloma, Rafin. l. c. (Quid?).—Hab.?

XOLISMA, Rafin. in Am. Monthly Mag. (1819) 193 = **Andromeda**, Linn. (Ericac.).

XUARESIA, Pers. Syn. i. 176 (1805) = seq.

XUAREZIA, Ruiz & Pav. Prod. 24. t. 4 (1794) = **Capraria**, Tourn. (Scrophul.).
biflora, Ruiz & Pav. Fl. Per. ii. 13 = C. biflora.

XURIS, Adans. Fam. ii. (20) (1763) = **Xyris**, Gronov.
fistulosa, Rafin. Fl. Tellur. ii. 14 (Quid?).—Hab.?

XYLADENIUS, Desv. in Ham. Prod. Pl. Ind. Occ. 41 (1825) = **Banara**, Aubl. (Samyd.).
glandulosus, Desv. ex Ham. l. c. = B. guianensis.

XYLANTHEMA, Neck. Elem. i. 67 (1790) = **Cnicus**, Linn. (Compos.).

XYLIA, Benth. in Hook. Journ. Bot. iv. (1842) 417. *LEGUMINOSAE*, Benth. & Hook. f. i. 594.
dolabriformis, *Benth. in Hook. Journ. Bot.* iv. (1842) 417.—Burma; Malaya.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 353.—Madag.
longipes, *Baill. l. c.* 354.—Madag.

XYLOBIUM, Lindl. Bot. Reg. sub t. 897 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 547.
ONKERIPUS, Rafin. Fl. Tellur. iv. 42 (1836).
PENTULOPS, Rafin. l. c. (1836).
brachypus, *Hemsl. Biol. Centr. Am. Bot.* iii. 252.—Nicaragua.
concavum, *Hemsl. l. c.*—Guatemala.
elongatum, *Hemsl. l. c.*—Am. centr.
longifolium, *Lindl. ex Spreng. Syst.* iii. 733.—N. Granat.
squalens, *Lindl. Bot. Reg.* sub t. 897.—Bras.
stachyobiorum, *Hemsl. Biol. Centr. Am. Bot.* iii. 252.—Nicaragua.

XYLOCALYX, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 84. *SCROPHULARINEAE.*
asper, *Balf. f. l. c.*—Ins. Socotra.

XYLOCARPUS, Koen. in Naturf. xx. (1784) 2; et ex A. Juss. in Mém. Mus. Par. xix. (1830) 243 = **Carapa**, Aubl. (Meliac.).
Antila, Buch.-Ham. ex Wall. Cat. n. 4893 = Walsura ternata.
Carapa, Spreng. Syst. ii. 213 = C. guianensis.
carnulosus, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 582.—Java.
Forstenii, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 62.—Ins. Celebes.
Granatum, Koen. in Naturf. xx. (1784) 2 = C. moluccensis.
guineensis, M. Roem. Syn. Hesper. 124 = C. procera, moluccensis.
moluccensis, M. Roem. l. c. = C. moluccensis.
obovatus, A. Juss. in Mém. Mus. Par. xix. (1830) 244 = C. moluccensis.
procerus, Steud. Nom. ed. II. ii. 793 = C. procera.
Toulocouna, Steud. l. c. 793 = C. procera.

XYLOCOCCUS, Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 258 = **Arctostaphylos**, Adans. (Ericac.).
bicolor, Nutt. l. c. 259 = A. bicolor.

XYLOLAENA, Baill. in Bull. Soc. Linn. Par. i. (1884) 410 = XYLOOLAENA = **Scleroolaena**, Baill. (Chlaen.).

XYLOMELUM, Sm. in Trans. Linn. Soc. iv. (1798) 214. *PROTEACEAE*, Benth. & Hook. f. iii. 179.
angustifolium, *Kipp. ex Meissn. in Hook. Journ. Bot.* (1852) 209.—Austral.
occidentale, *R. Br. Prot. Nov.* 31.—Austral.
pyriforme, *Knight, Prot.* 105; *R. Br. in Trans. Linn. Soc.* x. (1811) 189.—Austral.
salicinum, *A. Cunn. ex R. Br. Prot. Nov.* 31.—Austral.
Scottianum, F. Muell. Fragm. v. 174 = salicinum.

XYLON, Linn. Gen. ed. I. 53 (1737) = **Bombax**, Linn. (Malvac.).

XYLON, [Tourn.] Medic. Malv. 43 (1787) = **Gossypium**, Linn. (Malv.).
hirsutum, Medic. l. c. = G. herbaceum.
indicum, Medic. l. c. = G. herbaceum.
leoninum, Medic. l. c. = G. herbaceum.
religiosum, Moench, Meth. 616 = G. herbaceum.
strictum, Medic. Malv. 43 = G. herbaceum.

XYLOOLAENA, Baill. Dict. ii. 2, in obs. (1879) = **Scleroolaena**, Baill. (Chlaen.).
Richardi, Baill. in Bull. Soc. Linn. Par. i. (1884) 412 = S. Richardi.

XYLOPHYLLA, Linn. Mant. ii. 147 (1771); Schreb. Gen. 200 (1789) = **Phyllanthus**, Linn. (Euphorb.).
angustifolia, Sw. Prod. Veg. Ind. Occ. 28 = P. angustifolius.
Arbuscula, Sw. l. c. = P. speciosus.
artensis, Montr. in Mém. Acad. Lyon, x. (1860) 250 (Exocarpi sp.).—N. Caled.
asplenifolia, Salisb. Prod. 390 = P. latifolius.
elongata, Jacq. Hort. Schoenb. iii. 53 = P. angustifolius.
ensifolia, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 249 = Exocarpus xylophylloides.
falcata, Sw. Prod. Veg. Ind. Occ. 28 = P. epiphyllanthus.
latifolia, Linn. Mant. ii. 221 = P. latifolius.
latifolia, Sims, Bot. Mag. t. 1021 = P. speciosus.
latifolia, Sw. Prod. Veg. Ind. Occ. 28 = P. montanus.
linearis, Steud. Nom. ed. II. ii. 793 = P. linearis.
longifolia, Linn. Mant. ii. 221 = Exocarpus ceramicus.
Lucena, Roth, Nov. Pl. Sp. 185 = Fluggia Leucopyrus.
media, Lodd. Cat. ex G. Don, in Loud. Hort. Brit. 391, nomen.—Jamaic.
montana, Sims, Bot. Mag. t. 2652 = P. angustifolius.
montana, Sw. Prod. Veg. Ind. Occ. 28 = P. montanus.
obovata, Willd. Enum. Hort. Berol. 329 = Fluggia microcarpa.
obtusata, Thunb. Pl. Bras. Dec. i. 12 = P. obtusatus.
ramiflora, Ait. Hort. Kew. ed. I. i. 376 = Securinega ramiflora.
speciosa, Sweet, Hort. Brit. ed. I. 360 = P. speciosus.

XYLOPIA, Linn. Syst. ed. X. 1250 (1759). *ANONACEAE*, Benth. & Hook. f. i. 28.
BULLIARDA, Neck. Elem. ii. 321 (1790).
COELOCLINE, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 208.
HABZELIA, A. DC. l. c. 207 (1832).
KROCKERIA, Neck. Elem. ii. 317 (1790).
PARARTABOTRYS, Miq. Fl. Ind. Bot. Suppl. 154 (1860).
PATONIA, Wight, Illustr. Ind. Bot. i. 18 (1838).
XYLOPICRUM, P. Br. Hist. Jamaic. 250 (1756).
acuminata, *Dun. Mon. Anon.* 122. t. 16.—Ind. occ.
acutiflora, Benth. in Trans. Linn. Soc. xxiii. (1862) 478 = Dunaliana.
acutiflora, *A. Rich. Ess. Fl. Cub.* 55, in adnot.—Afr. trop.
aethiopica, *A. Rich. l. c.* 53.—Afr. trop.
africana, *Oliver, Fl. Trop. Afr.* i. 30.—Afr. trop.
amplexicaulis, *Baill. Adansonia*, iv. (1863–64) 142.—Madag.
aromatica, Baill. Hist. Pl. i. 278 = Unona concolor.
barbata, *Hoffm. ex Mart. in Mart. Fl. Bras.* xiii. i. 43.—Bras.
Blancoi, *Vidal, Sinops. Atlas*, p. xii. t. 5. f. C.—Ins. Philipp.
brasiliensis, *Spreng. Neue Entdeck.* iii. 50.—Bras.
buxifolia, *Baill. Adansonia*, iv. (1863–64) 143.—Madag.
Cacanes, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 155.—Bras.
caudata, *Hook. f. & Thoms. Fl. Ind.* i. 125.—Malaya.
Championii, *Hook. f. & Thoms. l. c.* 126.—Zeylan.
cubensis, *A. Rich. Ess. Fl. Cub.* 36.—Cuba.
dicarpa, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 85.—Singapore.
Dunaliana, *Vallot, in Bull. Soc. Bot. Fr.* xxix. (1883) 219.—Afr. trop.
elliptica, *Maing. ex Hook. f. Fl. Brit. Ind.* i. 86.—Malacca.
emarginata, *Mart. Fl. Bras.* xiii. i. 42.—Bras.
ferruginea, *Baill. Hist. Pl.* i. 225.—Malacca.
frutescens, *Aubl. Pl. Gui.* ii. 602. t. 292.—Am. trop.

XYLOPIA :—
frutescens, Gaertn. Fruct. i. 339. t. 69. f. 7 = muricata.
frutescens, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = Anona cinerea.
fusca, *Maing. ex Hook. f. Fl. Brit. Ind.* i. 83.—Malacca.
glabra, *Linn. Syst.* ed. X. 1250.—Jamaic.
grandiflora, *A. St. Hil. Fl. Bras. Mer.* i. 40. t. 8.—Bras.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 340.—Madag.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 6.—Jamaic.
javanica, *Steud. Nom.* ed. II. ii. 793.—Java.
Lamarckii, *Baill. Adansonia*, iv. (1863-64) 142.—Madag.
Langsdorfiana, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 133.—Bras.
Lastelliana, *Baill. Adansonia*, iv. (1863-64) 144.—Madag.
ligustrifolia, *Dun. Mon. Anon.* 121. t. 18.—Ecuador.
longifolia, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 210.—Panama.
lucida, *Baill. Hist. Pl.* i. 278.—Bras.
Maccreai, F. Muell. Fragm. vi. 176 = Melodorum Maccreai.
macrantha, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 38.—N. Granat.
magna, *Maing. ex Hook. f. Fl. Brit. Ind.* i. 84.—Malacca.
Maingayi, *Hook. f. & Thoms. l. c.* 85.—Malacca.
malayana, *Hook. f. & Thoms. Fl. Ind.* i. 125.—Malaya.
martinicensis, *Spreng. Syst.* ii. 636.—Ins. Martinic.
micrantha, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 27.—N. Guin.
muricata, *Linn. Syst.* ed. X. 1250.—Jamaic.
muricata, Vell. Fl. Flum. ix. t. 127 = frutescens.
nigricans, *Hook. f. & Thoms. Fl. Ind.* i. 125.—Zeylan.
nitida, *Dun. Mon. Anon.* 122. t. 20.—Guian.
obtusifolia, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 83.—Malacca.
obtusifolia, *A. Rich. Ess. Fl. Cub.* 39.—Cuba.
ochrantha, *Mart. Fl. Bras.* xiii. 1. 43.—Bras.
odoratissima, *Welw. ex Oliver, Fl. Afr. Trop.* i. 31.—Afr. trop.
orientalis, Spreng. Syst. ii. 636 = Unona selanica.
oxyantha, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 83.—Singapore.
pallescens, *Baill. Adansonia*, xi. (1873-76) 178.—N. Caled.
Pancheri, *Baill. l. c.* 177.—N. Caled.
parviflora, Benth. in Trans. Linn. Soc. xxiii. (1862) 479 = Dunaliana.
parviflora, *Spruce, in Journ. Linn. Soc.* v. (1861) 6.—Bras.
parviflora, *Vallot, in Bull. Soc. Bot. Fr.* xxix. (1882) 219.—Afr. trop.
parvifolia, *Hook. f. & Thoms. Fl. Ind.* i. 125.—Zeylan.
parvifolia, Schlecht. in Linnaea, ix. (1834) 327 = brasiliensis.
Pierrei, *Hance, in Journ. Bot.* xv. (1877) 328.—Cambodia.
polycarpa, *Oliver, Fl. Trop. Afr.* i. 32.—Afr. trop.
prinoides, *Dun. Mon. Anon.* 122. t. 15.—Guian.
pustulata, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 85.—Malacca.
pygmaea, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 151.—Bras.
Richardi, *Boiv. ex Baill. Adansonia*, iv. (1863-64) 145.—Ins. Mascar.
rubescens, *Oliver, Fl. Trop. Afr.* i. 30.—Afr. trop.
ruscifolia, Humb. & Bonpl. ex DC. Syst. i. 500 = ligustrifolia.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* v. 63.—Am. austr.
sericea, *A. St. Hil. Pl. Us. Bras.* t. 33.—Bras.
setosa, Poir. Encyc. viii. 812 = frutescens.
Spruceana, *Benth. ex Spruce, in Journ. Linn. Soc.* v. (1861) 5.—Bras.
Thomsoni, *Oliver, Fl. Trop. Afr.* i. 31.—Afr. trop.
Tooropiana, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 392.—Malaya.
trunciflora, *Cham. & Schlecht. in Linnaea*, vi. (1831) 417.—Mexic.
undulata, *Beauv. Fl. Owar.* i. 27. t. 16.—Afr. trop.
Vieillardi, *Baill. Adansonia*, viii. (1867-68) 202.—N. Caled.
Vielana, *Pierre, Fl. For. Cochinch.* t. 34.—Cochinch.

XYLOPICRON, Adans. Fam. ii. 365 (1763) = **Xylopia**, Linn. (Anonac.).
glabrum, Crantz, Inst. ii. 202 = Xylopia glabra.
muricatum, Crantz, l. c. = Xylopia muricata.

XYLOPICRUM, P. Br. Hist. Jamaic. 250 (1756) = praec.

XYLOPLEURUM, Spach, Hist. Vég. Phan. iv. 378 (1835) = **Oenothera**, Linn. (Onagrar.).
Berlandieri, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 370 = O. Delessertiana.
Drummondii, Spach, l. c. 371 = O. speciosa.
hirsutum, Spach, l. c. 370 = O. hirsuta.
Nuttallii, Spach, l. c. 371 = O. speciosa.
obtusifolium, Spach, l. c. 372 = O. speciosa.

XYLORHIZA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 297 = **Aster**, Tourn. (Compos.).
glabriuscula, Nutt. l. c. = A. Xylorhiza.
villosa, Nutt. l. c. 298 = A. Xylorhiza.

XYLORHIZA, Salisb. Gen. Pl. Fragm. 89 (1866) = **Allium**, Tourn. (Liliac.).
angulosa, Salisb. l. c. = A. angulosum.
senescens, Salisb. l. c. = A. senescens.

XYLOSMA, Forst. f. Prod. 72 (1786). *BIXINEAE*, Benth. & Hook. f. i. 128.
BESSERA, Spreng. Pugill. ii. 90 (1815).
CRAEPALOPRUMNON, Karst. Fl. Columb. 123. t. 61, 62 (1859?).
HISINGERA, Hellen. in Vet. Akad. Handl. Stockh. (1792) 32. t. 2.
KOELERA, Willd. Sp. Pl. iv. 750 (1805).
LIMACIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. iv. 383 (1818).
MYROXYLON, Forst. Char. Gen. 125 (1776).
RUMEA, Poit. in Mém. Mus. Par. i. (1815) 62. t. 4.
Benthami, *Griseb. in Goett. Abh.* ix. (1861) 27.—Am. trop.
buxifolium, *A. Gray, in Mem. Am. Acad.* N. S. viii. (1860) 155.—Cuba.
calophyllum, *Griseb. in Goett. Abh.* ix. (1861) 27.—Bras.
ciliatifolium, *Eichl. in Mart. Fl. Bras.* xiii. 1. 449.—Bras.
cinereum, *Hemsl. Biol. Centr. Am. Bot.* i. 57.—Mexic.
conicarpum, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 232 = suaveolens.
controversum, *Clos, l. c.* 231.—Reg. Himal.
Cumingii, *Clos, l. c.*—Ins. Philipp.
digynum, *Eichl. in Mart. Fl. Bras.* xiii. 1. 447.—Bras.
elegans, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 97.—N. Granat.
ellipticum, *Hemsl. Biol. Centr. Am. Bot.* i. 57.—Mexic.
ellipticum, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 343.—Madag.
flexuosum, *Hemsl. Biol. Centr. Am. Bot.* i. 57.—Mexic.
fragrans, *Decne. in Bot. Voy. Vénus*, t. 25.—Hab.?
hawaiense, *Seem. Fl. Vit.* 7.—Ins. Sandvic.
Hillebrandi, *Wawra, in Flora*, lvi. (1873) 171.—Ins. Sandvic.
infestum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1860) 155.—Cuba.
integrifolium, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 230 = orbiculatum.
intermedium, *Griseb. in Goett. Abh.* ix. (1861) 27; *Triana & Planch. l. c.* xvii. (1861) 100.—Veraguas.
japonicum, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 381 = racemosum.
lanceolatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 553.—Mexic.
latifolium, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 194.—Ind. or.
Lepinei, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 233 = suaveolens.
leprosipes, *Clos, l. c.* 230.—Java.
longifolium, *Clos, l. c.*—Reg. Himal.
luzonensis, *Clos, l. c.* 231.—Ins. Philipp.
molle, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 98.—N. Granat.
monospora, *Harv. Thes. Cap.* ii. 52.—Afr. austr.
nitidum, *A. Gray, ex Griseb. in Fl. Brit. W. Ind.* 21.—Am. trop.
obovatum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 97.—N. Granat.
orbiculatum, *Forst. f. Prod.* 72.—Ins. Amicor.

XYLOSMA :—
ovatum, *Benth. Fl. Austral.* i. 108.—Austral.
panamensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 554.—Panama.
prunifolium, *Griseb. in Goett. Abh.* ix. (1861) 27; *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 98.—N. Granat.
pubescens, *Griseb. l. c.* xxiv. (1879) 20.—Reg. Argent.
racemosum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 155.—Ind. or.
Salzmanni, *Eichl. in Mart. Fl. Bras.* xiii. 1. 448.—Bras.
schaefferioides, *A. Gray, in Mem. Am. Acad.* N. S. viii. (1860) 155.—Cuba.
Seemannii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 99.—Panama.
senticosum, *Hance, in Journ. Bot.* vi. (1868) 328.—China.
spiculiferum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 96.—N. Granat.
suaveolens, *Forst. f. Prod.* 72.—Ins. Pacif.
velutinum, *Triana & Karst. in Karst. Fl. Columb.* i. 123. t. 97.—N. Granat.

XYLOSTEON, Tourn. ex Adans. Fam. ii. 157, 501 (1763) = **Lonicera**, Linn. (Caprif.).
alpigenum, Fuss, Fl. Transs. 284 = L. alpigena.
alpinum, Dum.-Cours. Bot. Cult. ed. II. iv. 336 = L. alpigena.
arboreum, Webb, Iter Hisp. 42 = L. arborea.
asperifolium, Decne. in Jacquem. Voy. Bot. 77. t. 85 = L. asperifolia.
caeruleum, Dum.-Cours. Bot. Cult. ed. II. iv. 336 = L. caerulea.
campaniflorum, Lodd. Bot. Cab. t. 1361 = L. nigra.
canadense, Duham. ex DC. Prod. iv. 337 = L. caerulea.
ciliatum, Pursh, Fl. Am. Sept. 161 = L. ciliata.
ciliosum, Steud. Nom. ed. II. ii. 793 = L. ciliosa.
coeruleum, Auct. = *caerulea*.
cordatum, Moench, Meth. 502 = L. tatarica.
dumetorum, Moench, l. c. = L. Xylosteum.
gibbiflorum, Rupr. & Maxim. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 136 = L. chrysantha.
ibericum, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 44 = L. iberica.
involucratum, Richards. in Frankl. Narr. 1st Journ. App. 733 = L. involucrata.
Karelini, Rupr. Sert. Tiansch. 50 = L. Karelini.
ligustrinum, D. Don, Prod. Fl. Nep. 140 = L. ligustrina.
Loureirii, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 172 = L. acuminata.
Maackii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 369 = L. Maackii.
Maximowiczii, Rupr. l. c. (1856) 136 = L. Maximowiczii.
mexicanum, H. B. & K. Nov. Gen. et Sp. iii. 426. t. 297 = L. gibbosa.
Naisoca, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 140 = L. ligustrina.
nigrum, Medic. Pfl. Anat. 97; Roehl. Deutschl. Fl. ed. II. ii. 176 = L. nigra.
oblongifolium, Goldie, in Edinb. Phil. Journ. vi. (Apr. 1822) 323 = L. caerulea.
puniceum, Rafin. New Fl. Am. iii. 21 = Symphoricarpos puniceus.
purpurascens, Jacquem. Voy. Bot. 79. t. 87 = L. purpurascens.
pyrenaicum, Dum.-Cours. Bot. Cult. ed. II. iv. 336 = L. pyrenaica.
scandens, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 140 = L. macrantha.
Solonis, Eaton, Man. Bot. N. Am. ed. VI. 398; ed. VII. 518 = L. caerulea.
spinosum, Decne. in Jacquem. Voy. Bot. 78. t. 86 = L. spinosa.
tartaricum, Medic. Pfl. Anat. 97 = L. tatarica.
tataricum, Michx. Fl. Am. Bor. i. 106 = L. ciliata.
villosum, Michx. l. c. = L. caerulea.
vulgare, Roehl. Deutschl. Fl. ed. II. ii. 176 = L. Xylosteum.

XYLOSTEUM, Rupp. Fl. Jen. ed. Hall. 250 (1745) = praec.

XYLOTHECA, Hochst. in Flora, xxvi. (1843) 69 = **Oncoba**, Forsk. (Bixin.).
Kraussiana, Hochst. l. c. = O. Kraussiana.

XYNOPHYLLA, Montr. in Mém. Acad. Lyon, x. (1860) 250 = **Exocarpus**, Labill. (Santal.).
artensis, Montr. l. c.—N. Caled.

XYOMALOBIUM, Weale, in Journ. Linn. Soc. xiii. (1873) 50, err. typ. = **Xysmalobium**, R. Br. (Asclep.).

XYPHANTHUS, Rafin. Fl. Ludov. 103 (1817) = **Erythrina**, Linn. (Legumin.).
hederifolius, Rafin. l. c.—Am. bor.

XYPHIDIUM, Neck. Elem. iii. 168 (1790) = **Xiphidium**, Aubl. (Haemod.).

XYPHIDIUM, Steud. Nom. ed. II. ii. 793 (1841) = XIPHION, Tourn. = **Iris**, Tourn.
sordidum, Steud. l. c. = I. juncea.

XYPHION, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 418 = XIPHION, Tourn. = **Iris**, Tourn.

XYPHOSTYLIS, Rafin. Fl. Tellur. iv. 52 (1836) = **Canna**, Linn. (Scitam.).
angustifolia, Rafin. l. c. = C. angustifolia.
lutea, Rafin. l. c. = C. lutea.

XYRIDANTHE, Lindl. Swan River App. 23 (1839) = **Helipterum**, DC. (Compos.).
stricta, Lindl. l. c. = H. strictum.

XYRIDION, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 163 = **Iris**, Tourn.
acoroideum, Klatt, in Bot. Zeit. xxx. (1872) 500 = I. Pseudacorus.
aureum, Klatt, l. c. 501 = I. aurea.
flexuosum, Klatt, l. c. 500 = I. sibirica.
foetidissimum, Klatt, l. c. = I. foetidissima.
gramineum, Klatt, l. c. = I. graminea.
Gueldenstaedtianum, Klatt, l. c. 501 = I. Gueldenstaedtiana.
halophilum, Klatt, l. c. 500 = I. Gueldenstaedtiana.
laevigatum, Klatt, l. c. = I. laevigata.
Monnieri, Klatt, l. c. = I. Monnieri.
nothum, Klatt, l. c. = I. spuria.
ochroleucum, Klatt, l. c. = I. ochroleuca.
Pseudacorus, Klatt, l. c. = I. Pseudacorus.
Reichenbachianum, Klatt, l. c. = I. spuria.
setosum, Klatt, l. c. = I. setosa.
sibiricum, Klatt, l. c. = I. sibirica.
spurium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 163 = I. spuria.
stenogynum, Klatt, in Bot. Zeit. xxx. (1872) 500 = I. Gueldenstaedtiana.
tridentatum, Klatt, l. c. = I. tridentata.
ventricosum, Klatt, l. c. = I. ventricosa.
violaceum, Klatt, l. c. = I. laevigata.

XYRIDIUM, Steud. Nom. ed. II. ii. 793 (1841) = praec.

XYRIS, Gronov. ex Linn. Gen. ed. I. 11 (1737).
XYRIDEAE, Benth. & Hook. f. iii. 842.
JUPICA, Rafin. Fl. Tellur. ii. 15 (1836).
KOTSJILETTI, Adans. Fam. ii. 60 (1763).
RAMOTHA, Rafin. Fl. Tellur. ii. 15 (1836).
SCHISMAXON, Steud. in Bot. Zeit. xvi. (1856) 391.
SYNOLIGA, Rafin. Fl. Tellur. ii. 15 (1836).
XYROIDES, Thou. Nov. Gen. Madag. 4 (1806).
acuminata, Miq. ex Steud. Syn. Pl. Cyp. 284 = communis.
aemula, Endl. in Lehm. Pl. Preiss. ii. 54 = flexifolia.
altissima, Lodd. Bot. Cab. t. 1900 = Bobartia indica.
ambigua, *Beyr. ex Kunth, Enum. Pl.* iv. 13.—Am. bor.
americana, *Aubl. Pl. Gui.* i. 40.—Am. trop.
americana, Poepp. ex Kunth, Enum. Pl. iv. 12 = laxifolia.
americana, Vahl, Enum. ii. 205 = glabrata.
anceps, *Lam. Illustr.* i. 132.—Ind. or.; Malaya.
anceps, Pers. Syn. i. 56 = caroliniana.
anceps, Vahl, Enum. ii. 204 = complanata.
aphylla, *Vell. Fl. Flum.* 29; i. t. 72 (sp. dub.).—Bras.
arescens, Kunth, Enum. Pl. iv. 3 = hymenachne.
asperata, Kunth, l. c. 6 = trachyphylla.
asperula, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 57.—Bras.
asperula, *Seub. in Mart. Fl. Bras.* iii. 1. 219.—Bras.
asterocephala, Seub. l. c. = involucrata.

XYRIS:—
bahiensis, Steud. Syn. Pl. Cyp. 287 = anceps.
Baldwiniana, *Schult. Mant.* i. 351.—Am. bor.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 608.—Malaya.
bicarinata, *Griseb. Cat. Pl. Cub.* 223.—Cuba.
blepharophylla, Kunth, Enum. Pl. iv. 5 = pterygoblephara.
blepharophylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 56.—Bras.
bracteata, R. Br. Prod. 256 = operculata?
brasiliensis, Spreng. Nov. Prov. 48 = Fimbristylis monostachya.
brevifolia, Ell. Sketch, i. 52 = Elliottii.
brevifolia, *Michx. Fl. Bor. Am.* i. 23.—Am. bor.
brevifolia, *A. St. Hil. Voy. Distr. Diam.* i. 390.—Bras.
Brownei, *Kunth, Enum. Pl.* iv. 22.—Austral.
bulbosa, Kunth, l. c. 11 = flexuosa.
caerulea, Lam. Illustr. i. 132 = americana.
calocephala, *Miq. Fl. Ind. Bat.* iii. 527.—Java.
canadensis, Schnizl. Ic. i. t. 47 = caroliniana.
capensis, *Baker, in Journ. Linn. Soc.* xx. (1883) 276.—Madag.
capensis, *Thunb. Prod. Pl. Cap.* 12.—Afr. austr.
capensis, Willd. ex Kunth, Enum. Pl. iv. 16 = humilis.
Capito, *Hance, in Journ. Bot.* xiv. (1876) 262.—Cambodia.
caroliniana, *Walt. Fl. Carol.* 69.—Am. bor.
castanea, Klotzsch, ex Seub. in Mart. Fl. Bras. iii. 1. 220 = rigida.
caulescens, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1064, nomen.—Guian.
ciliata, *Thunb. Pl. Bras. Dec.* iii. 27.—Bras.
communis, *Kunth, Enum. Pl.* iv. 12.—Am. trop.
complanata, *R. Br. Prod.* 256.—Austral.
compressa, Willd. ex A. Dietr. Sp. Pl. ii. 372 = anceps.
conocephala, *Wright, in Sauv. Fl. Cub.* 159 (an torta?).—Cuba.
consanguinea, *Kunth, Enum. Pl.* iv. 8.—Bras.
denticulata, R. Br. Prod. 256 = pauciflora.
desertorum, Pohl, ex Seub. in Mart. Fl. Bras. iii. 1. 214 = rupicola.
difformis, *Chapm. Fl. S. U. St.* 500.—Florida.
elata, *Chapm. l. c.* 501.—Florida.
Elliottii, *Chapm. l. c.* 500.—Am. bor.
elongata, Rudge, in Trans. Linn. Soc. x. (1811) 289. t. 15 = complanata.
eriophora, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 1117 = Fimbristylis lanata.
eriophylla, *Reichb. ex Kunth, Enum. Pl.* iv. 4.—Guian.
Falx, *Griseb. Cat. Pl. Cub.* 223.—Cuba.
filiformis, *Lam. Illustr.* i. 132.—Afr. trop.
fimbriata, Beyr. ex Kunth, Enum. Pl. iv. 12 = caroliniana.
fimbriata, *Ell. Sketch*, i. 52.—Am. bor.
flabelliformis, *Chapm. Fl. S. U. St.* 499.—Florida.
flexifolia, *R. Br. Prod.* 256.—Austral.
flexuosa, Muhl. Cat. 5 (1813); ex Ell. Sketch, i. 51 = platylepis.
Fontanesiana, Kunth, Enum. Pl. iv. 10 = anceps.
glabrata, *Griseb. Fl. Brit. W. Ind.* 525.—Am. trop.
gracilis, *R. Br. Prod.* 256.—Austral.
gracillima, *F. Muell. Fragm.* viii. 203.—Austral.
graminosa, *Pohl, ex Mart. in Flora*, xxiv. (1841) II. *Beibl.* 55.—Bras.
grandiceps, *Griseb. Cat. Pl. Cub.* 223.—Cuba.
guianensis, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 896, nomen = communis.
guianensis, *Steud. Syn. Pl. Cyp.* 285.—Guian.
gymnoptera, Griseb. Cat. Pl. Cub. 223 = communis.
humilis, *Kunth, Enum. Pl.* iv. 15.—Ins. Mascar.
hymenachne, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 55.—Bras.
Hystrix, *Seub. in Mart. Fl. Bras.* iii. 1. 219.—Bras.
indica, *Linn. Sp. Pl.* 42.—Ind. or.; Malaya.
indica, Pursh, Fl. Am. Sept. i. 33 = flexuosa.
indica, Vell. Fl. Flum. 29; i. t. 71 = communis.
indica, Willd. ex Kunth, Enum. Pl. iv. 19 = anceps.
insularis, *Steud. Syn. Pl. Cyp.* 288 (sp. dub.).—Madag.
involucrata, *Nees, in Hook. Journ. Bot.* ii. (1840) 397.—Guian.
iridifolia, *Chapm. Fl. S. U. St.* 501.—Am. bor.
juncea, Baldw. in Ell. Sketch, i. 53 = Baldwiniana.
juncea, R. Br. Prod. 256 = gracilis.
Jupicai, Michx. Fl. Bor. Am. i. 23 = caroliniana.

XYRIS:—
Jupicai, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = americana.
lacera, *R. Br. Prod.* 257.—Austral.
lacerata, *Pohl, ex Seub. in Mart. Fl. Bras.* iii. 1. 216.—Bras.
laevis, R. Br. Prod. 256 = complanata.
laevis, Nees, ex Benth. Fl. Austral. vii. 380 = Mesomelaena sphaerocephala.
lanata, *R. Br. Prod.* 257.—Austral.
lanuginosa, *Seub. in Mart. Fl. Bras.* iii. 1. 214.—Bras.
lappacea, Heyne, ex Mart. in Wall. Pl. As. Rar. iii. 30 = anceps?
laxiflora, *F. Muell. Fragm.* viii. 203.—Austral.
laxifolia, Mart. in Flora, xxiv. (1841) II. Beibl. 53 = communis.
lomatophylla, *Mart. l. c.* 57.—Bras.
longifolia, *Mart. l. c.* 54.—Bras.
macrocephala, *Vahl, Enum.* ii. 204.—Guian.
malaccensis, Steud. Syn. Pl. Cyp. 287 = complanata.
Martiana, Seub. in Mart. Fl. Bras. iii. 1. 216 = blepharophylla.
melanocephala, *Miq. Fl. Ind. Bat.* iii. 528.—Java; Sumatra.
metallica, *Klotzsch, ex Seub. in Mart. Fl. Bras.* iii. 1. 213.—Bras.
microcephala, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 119 (sp. dub.).—Java.
Minarum. *Seub. in Mart. Fl. Bras.* iii. 1. 215.—Bras.
minima, Steud. Syn. Pl. Cyp. 288 = humilis.
montivaga, *Kunth, Enum. Pl.* iv. 7.—Bras.
navicularis, *Griseb. Cat. Pl. Cub.* 223.—Cuba.
nilagiriensis, Steud. Syn. Pl. Cyp. 288 = capensis.
nitida, Willd. ex A. Dietr. Sp. Pl. ii. 372 = anceps.
nubigena, *Kunth, Enum. Pl.* iv. 3.—Bras.
oligantha, Steud. Syn. Pl. Cyp. 288 = pauciflora.
operculata, R. Br. Prod. 257 = Brownei.
operculata, *Labill. Nov. Holl. Pl.* i. 14. t. 10.—Austral.
oryzetorum, Miq. ex Steud. Syn. Pl. Cyp. 288 = pauciflora.
pallida, Mart. in Flora, xxiv. (1841) II. Beibl. 57 = anceps.
paludosa, R. Br. Prod. 256 = pauciflora.
paraënsis, *Poepp. ex Kunth, Enum.* Pl. iv. 9.—Bras.; Guian.
pauciflora, Wall. Cat. n. 6083 = Wallichii.
pauciflora, *Willd. Phytogr.* i. 2. t. 1.—As. et Austral. trop.
Perrottetii, Steud. Syn. Pl. Cyp. 319 = anceps.
pilosa, *Kunth, Enum. Pl.* iv. 6.—Bras.
plantaginea, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 514.—Bras.
platicaulis, Poir. Encyc. viii. 820 = anceps.
platylepis, *Chapm. Fl. S. U. St.* 501.—Am. bor.
pterygoblephara, *Steud. Syn. Pl. Cyp.* 285.—Bras.
pubescens, *Poir. Encyc.* viii. 819 (sp. dub.).—Ind. occ.
pumila, Pohl, ex Seub. in Mart. Fl. Bras. iii. 1. 217 = savanensis.
Purshii, Sweet, Hort. Brit. ed. II. 546 = torta *vel* flexuosa.
pusilla, R. Br. Prod. 256 = pauciflora.
restiacea, Mart. in Flora, xxiv. (1841) II. Beibl. 57 = ciliata.
rhombipetala, Wright, in Sauv. Fl. Cub. 160 = ambigua.
rigida, *Kunth, Enum. Pl.* iv. 15.—Bras.
robusta, *Mart. in Wall. Pl. As. Rar.* iii. 30; *et in Cat.* n. 6087.—Reg. Himal.
rupicola, *Kunth, Enum. Pl.* iv. 5.—Bras.
savanensis, *Miq. in Linnaea*, xviii. (1844) 605.—Am. trop.
scabra, R. Br. Prod. 256 = complanata.
scabra, Engelm. in Bost. Journ. Nat. Hist. v. (1845) 235 = flexuosa.
schizachne, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 56.—Bras.
schoenoides, Mart. in Wall. Pl. As. Rar. iii. 30 = capensis.
Sellowiana, Kunth, Enum. Pl. iv. 13 = macrocephala.
semifuscata, Boj. ex Baker, in Journ. Linn. Soc. xx. (1883) 277 = capensis.
serotina, *Chapm. Fl. S. U. St.* 500.—Florida.
setacea, *Chapm. l. c. Suppl.* 658.—Florida.
sororia, *Kunth, Enum. Pl.* iv. 15.—Am. bor.
spectabilis, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 54.—Bras.

XYRIS:—
spiralis, Pohl, ex Seub. in Mart. Fl. Bras. iii. 1. 213 = tortula.
stricta, *Chapm. Fl. S. U. St.* 500.—Florida.
strobilifera, Kunth, Enum. Pl. iv. 4 = ciliata.
subulata, *Ruiz & Pav. Fl. Per.* i. 46. t. 7. f. 6.—Peruv.
surinamensis, Miq. in Linnaea, xvii. (1843) 58 = communis.
surinamensis, *Spreng. f. Tent. Suppl. Syst.* 2 (sp. dub.).—Guian.
tenella, *Kunth, Enum. Pl.* iv. 9.—Bras.
tenuifolia, Chapm. Fl. S. U. St. 502 = Baldwiniana.
tenuis, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen (Quid ?).—Am. bor.
teretifolia, R. Br. Prod. 257 = lacera ?
torta, *Sm. in Rees, Cyclop.* xxxix. n. 11.—Am. bor.
tortula, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 55.—Bras.
trachyphylla, *Mart. l. c.* 56.—Bras.
vaginata, Spreng. Syst. i. 183 = Abolboda brasiliensis.
vivipara, *H. B. & K. Nov. Gen. et Sp.* i. 255 (sp. dub.).—Venezuela.
Walkeri, Wight, ex Kunth, Enum. Pl. iv. 19 = complanata.
Wallichii, *Kunth, l. c.* 16.—Reg. Himal. ; Burma.

XYROIDES, Thou. Gen. Nov. Madag. 4 (1806) = praec.

XYROPLEURUM, Auct. ex Steud. Nom. ed. II. ii. 794 (1841); err. typ. = XYLOPLEURUM, Spach = **Oenothera**, Linn. (Onagrar.).

XYSMALOBIUM, R. Br. in Mem. Wern. Soc. i. (1809) 38. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 752.
albens, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus albens.
appendiculatum, D. Dietr. l. c. = Gomphocarpus appendiculatus.
concolor, D. Dietr. l. c. = Gomphocarpus concolor.
coronarium, D. Dietr. l. c. = Gomphocarpus coronarius.
crispum, D. Dietr. l. c. = Gomphocarpus crispus.
dealbatum, D. Dietr. l. c. = Gomphocarpus dealbatus.
gomphocarpoides, *D. Dietr. l. c.* (sp. dub.).—Afr. austr.
grandiflorum, R. Br. in Mem. Wern. Soc. i. (1809) 39 = Gomphocarpus grandiflorus.
Heudelotianum, *Decne. in DC. Prod.* viii. 520.—Afr. trop.
humile, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus humilis.
involucratum, *Decne. in DC. Prod.* viii. 520.—Afr. austr.
lapathifolium, Decne. l. c. 519 = undulatum.
ligulatum, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus dealbatus.
linguaeforme, *Harv. ex Weale, in Journ. Linn. Soc.* xiii. (1873) 50.—Afr. austr.
marginatum, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus marginatus.
orbiculare, *D. Dietr. l. c.*—Afr. austr.
pedunculatum, *Harv. Thes. Cap.* ii. 8. t. 112.—Afr. austr.
prunelloides, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) 1. 254.—Afr. austr.
reflectens, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus reflectens.
rigidum, D. Dietr. l. c. = Gomphocarpus rigidus.
sessile, *Decne. in DC. Prod.* viii. 519.—Afr. austr.
undulatum, *R. Br. in Mem. Werm. Soc.* i. (1809) 39.—Afr. austr.
vexillare, D. Dietr. Syn. Pl. ii. 902 = Gomphocarpus dealbatus.
viridiflorum, D. Dietr. l. c. = Gomphocarpus viridiflorus.

XYSTIDIUM, Trin. Fund. Agrost. 102 (1820) = **Perotis**, Ait. (Gramin.).
barbatum, J. Presl, in Rel. Haenk. i. 228 = P. latifolia.
maritimum, Trin. Fund. Agrost. 102. t. 2 = P. rara.

XYSTRIS, Schreb. Gen. i. 138 (1789) (Quid ?).
Schreberi, J. F. Gmel. Syst. 390.—Hab. ?

Y

YAMALA, Rafin. Fl. Tellur. ii. 75 (1836) = **Heuchera**, Linn. (Saxifr.).
cylindrica, Rafin. l. c. = H. cylindrica.

YANGAPA, Rafin. Sylva Tellur. 20 (1838) = **Gardenia**, Ellis (Rubiac.).
flava, Rafin. l. c. = G. coronaria.

YANGUA, Spruce, in Journ. Linn. Soc. iii. (1859) 197 = **Cybistax**, Mart. (Bignon.).
tinctoria, Spruce, l. c. 198.—Peruv.

YERMOLOFFIA, Bélang. Voy. Ind. or. t. absque num. (1838 ?) = **Lagochilus**, Bunge (Labiat.).
hispida, Bélang. l. c. = L. hispidus, *Boiss.*
insignis, Bélang. l. c. = L. insignis.

YERMOLOFIA, Endl. Gen. 625 (1838) = praec.

YGRAMELA, Rafin. Atl. Journ. 199 (1833) = **Limosella**, Linn. (Scrophul.).
maritima, Rafin. l. c. = L. maritima.

YMNOSTEMA, Neck. Elem. i. 132 (1790) = **Lobelia**, Plum. (Campan.).

YMNOSTEMMA, Steud. Nom. ed. II. ii. 794 (1841) sphalm. = praec.

YOANIA, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 68. *ORCHIDEAE*, Benth. & Hook. f iii. 617.
japonica, *Maxim. l. c.* 69.—Reg. Himal. ; Japon.

YODES, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 298, err. typ. = **Iodes**, Blume (Olacin.).

YOUNGIA, Cass. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 88 = **Crepis**, Vaill. (Compos.).
acaulis, DC. Prod. vii. 193 = C. acaulis.
affinis, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 566 = Prenanthes affinis.
ambigua, DC. Prod. vii. 193 = C. japonica.
aspera, Steud. Nom. ed. II. ii. 393, in syn. = Chondrilla aspera.
chinensis, DC. Prod. vii. 194 = Lactuca Fischeriana.
chrysantha, Maxim. Prim. Fl. Amur. 181 = Lactuca denticulata.
debilis, DC. Prod. vii. 194 = Lactuca debilis.
dentata, DC. l. c. 193 (excl. syn.) = Lactuca denticulata.
diversifolia, Ledeb. Fl. Ross. ii. 837 = C. tenuifolia.
fastigiata, DC. Prod. vii. 193 = C. japonica.
flexuosa, Ledeb. Fl. Ross. ii. 838 = C. nana.
fuscipappa, Thw. Enum. Pl. Zeyl. 168 = C. fuscipappa.
glauca, Edgew. in Trans. Linn. Soc. xx. (1846) 79 = C. flexuosa.
gracilis, Miq. in Journ. Bot. Néerl. i. (1861) 106 = C. japonica ?
hastata, DC. Prod. vii. 194 = Lactuca denticulata.
humilis, DC. l. c. 194.—Japon.
humilis, Miq. in Journ. Bot. Néerl. i. (1861) 106.—China.
integra, A. Gray, in Mem. Am. Acad. N. S. vi. (1859) 396 = C. integra.
integrifolia, Cass. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 89 = C. japonica.
japonica, DC. Prod. vii. 194 = C. japonica.
Kochiana, Ledeb. Fl. Ross. ii. 837 = Phaecasium lampsanoides.
lanceolata, DC. Prod. vii. 193 = C. integra.
lyrata, Cass. in Ann. Sc. Nat. Sér. I. xxiii. (1831) 88 = C. japonica.
mauritiana, DC. Prod. vii. 192 = C. japonica.
multiflora, DC. l. c. 194 = C. japonica.
napifolia, DC. l. c. 193 = C. japonica.
Paosia, Steud. Nom. ed. II. ii. 794, sphalm. = seq.
Poosia, DC. Prod. vii. 193 = C. japonica.
procumbens, DC. l. c. = Launaea nudicaulis.
pygmaea, Ledeb. Fl. Ross. ii. 838 = C. nana.
runcinata, DC. Prod. vii. 192 = C. japonica.
serotina, Maxim. Prim. Fl. Amur. 180 = Lactuca denticulata.
sonchifolia, Maxim. l. c. = Lactuca denticulata.
Stenoma, Ledeb. Fl. Ross. ii. 837 = C. Stenoma.
striata, DC. Prod. vii. 193 = C. japonica.
Thunbergiana, DC. l. c. 192 = C. japonica.

YPOMAEA, Robin, Voy. Louis. iii. 402 (1807), sphalm. = **Ipomoea**, Linn. (Convolv.).

YUCCA, Dill. ex Linn. Gen. ed. I. 99 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 778.
CODONOCRINUM, Willd. ex Schult. f. Syst. vii. 718 (1829).
acaulis, *H. B. & K. Nov. Gen. et Sp.* i. 289.—Venezuela.
acrotricha, Schiede, in Linnaea, iv. (1829) 230 = Dasylirion acrotrichum.
acuminata, Hort. ex Carr. in Rev. Hortic. Sér. IV. viii. (1859) 398 = flexilis.
acuminata, Sweet, Brit. Flow. Gard. Ser. I. t. 195 = gloriosa.
acutifolia, *Truff. in Rev. Hortic.* (1869) 320.—Hab. ?
agavoides, Hort. ex Carr. l. c. Sér. IV. vii. (1858) 580 = Treculeana.
albo-spica, Hort. ex Van Houtte, in Fl. des Serres, xvii. (1867–68) 110. f. 1612 = constricta.
aletriformis, Haw. in Phil. Mag. (1831) 415 (al. gen.).—Afr. austr.
aloifolia, *Linn. Sp. Pl.* 319.—Am. bor. ; Ind. occ.
aloifolia, Torr. in Pacif. Rail. Rep. iv. 147 = Whipplei.
angustifolia, Carr. in Rev. Hortic. Sér. IV. ix. (1860) 20. f. 3, 4 = constricta.
angustifolia, Hort. ex Carr. l. c. viii. (1859) 398 = flexilis.
angustifolia, Karw. ex G. Don, in Sweet, Hort. Brit. ed. III. 707, nomen = stenophylla.
angustifolia, Pursh, Fl. Am. Sept. i. 227 = glauca.
antwerpensis, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 228 = filamentosa.
arcuata, Haw. Suppl. Pl. Succ. 33 = aloifolia.
argospatha, *Verlot, in Rev. Hortic.* (1868) 393.—Hab. ?
armata, Steud. Nom. ed. II. ii. 795 = aloifolia.
aspera, Regel, Ind. Sem. Hort. Petrop. (1858) 24 ; Gartenfl. (1859) 14 = Treculeana.
Atkinsi, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 221 = aloifolia.
baccata, *Torr. Bot. Mex. Bound.* 221.—Am. bor. occ. ; Mexic.
Barrancasecca, Hort. ex Pasq. Cat. Orto Bot. Nap. 108, nomen (Quid ?).—Hab. ?
Boerhaavii, *Baker, in Gard. Chron.* (1870) 1217.—Mexic.
Boscii, Desf. Tabl. ed. II. 28 = Agave geminiflora.
brevifolia, *Engelm. in S. Wats. Bot. King's Exped.* 496.—Calif.
brevifolia, Schott, ex Torr. Bot. Mex. Bound. 221 = Schottii.
californica, Lem. Illustr. Hortic. x. (1863) sub t. 372 = Whipplei.
canaliculata, Hook. Bot. Mag. t. 5201 = Treculeana.
circinata, Baker, in Gard. Chron. (1870) 1088 = baccata.
concava, Haw. Suppl. Pl. Succ. 34 = filamentosa.
concava, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 226 = Treculeana.
conspicua, Haw. Suppl. Pl. Succ. 32 = aloifolia.
conspicua, *Hort. ex Regel, Gartenfl.* xx. (1871) 149.—Hab. ?
constricta, *Buckl. in Proc. Acad. Sc. Philad.* 1862 (1863), 8.—Mexic.
contorta, Hort. ex Carr. in Rev. Hortic. Sér. IV. vii. (1858) 579, 580 = rupicola, Treculeana.
cornuta, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 226 = Treculeana.
crassifila, Engelm. in Trans. Acad. Sc. St. Louis, iii. (1873) 44 = baccata.
crenulata, Haw. Suppl. Pl. Succ. 33 = aloifolia.
De-Smetiana, *Baker, in Gard. Chron.* (1870) 1217.—Mexic.
Draco, Carr. in Rev. Hortic. Sér. IV. viii. (1859) 389 = Dracaena Draco ?
Draconis, Linn. Amoen. Acad. iii. 407 = Dracaena Draco.
Draconis, Linn. Sp. Pl. 319 = aloifolia.
Ehrenbergii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 538, nomen.—Hab. ?
elata, *Engelm. in Coult. Bot. Gaz.* vii. (1881) 17.—Am. bor. occ.
Ellacombei, Hort. ex Baker, in Saund. Refug. Bot. t. 317 = gloriosa.
Engelmanni, Mast. in Gard. Chron. (1880) II. 43, sphalm. = Whipplei.
ensifolia, Baker, in Saund. Refug. Bot. t. 318 = flexilis *vel* gloriosa.

YUCCA:—
exigua, Baker, in Saund. Refug. Bot. t. 314 = filamentosa.
Eylesii, Hort. Peacock, ex Baker, in Journ. Linn. Soc. xviii. (1880) 224 = flexilis.
falcata, Hort. Peacock, ex Baker, l. c. = flexilis.
filamentosa, *Linn. Sp. Pl.* 319.—Am. bor.
filamentosa, Wood, in Proc. Acad. Sc. Philad. (1868) 167 = baccata.
filifera, Hort. ex Engelm. in Trans. Acad. Sc. St. Louis, iii. (1873) 46; Chabaud. in Rev. Hortic. (1876) 432. f. 97 = baccata.
flaccida, Haw. Suppl. Pl. Succ. 34 = filamentosa.
flexilis, *Carr. in Rev. Hortic.* Sér. IV. viii. (1859) 398. f. 89.—Mexic.
fragilifolia, Baker, in Gard. Chron. (1870) 1088 = baccata.
funifera, Lem. Illustr. Hortic. xiii. (1866) Misc. 99.—Mexic.
Ghiesbreghtii, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 222 = guatemalensis.
gigantea, *Lem. Illustr. Hortic.* vi. (1859) *Misc.* 91.—Mexic.
glauca, Nois. ex Sims, Bot. Mag. t. 2662 = filamentosa.
glauca, [*Nutt.*] *in Fras. Cat.* (1813) n. 89.—Am. bor.
glaucescens, Haw. Suppl. Pl. Succ. 35 = filamentosa.
gloriosa, *Linn. Sp. Pl.* 319.—Am. bor.
gracilis, Link, ex Sweet, Hort. Brit. ed. II. 597 (Quid ?).—Mexic.
graminifolia, Wood, in Proc. Acad. Sc. Philad. (1868) 167 = Whipplei.
graminifolia, Zucc. in Cat. Hort. Monac. (1837); et in Abh. Akad. Muench. iii. (1837–43) 226 = Dasylirion graminifolium.
guatemalensis, *Baker, in Saund. Refug. Bot.* t. 313.—Guatemala; Mexic.
Haruckeriana, Crantz, Duab. Drac. Arbor. 29 = aloifolia.
horrida, Humb. ex Steud. Nom. ed. II. ii. 795 = spinosa.
integerrima, Stokes, Bot. Mat. Med. ii. 267 = gloriosa.
japonica, Hort. ex Carr. in Rev. Hortic. Sér. IV. viii. (1859) 488 = gloriosa.
laetevirens, Karw. ex G. Don, in Sweet, Hort. Brit. ed. III. 707, nomen (Quid ?).—Mexic.
Lenneana, *Baker, in Journ. Linn. Soc.* xiv. (1875) 538.—Hab. ?
longifolia, [Engelm.] ex Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 8 = Treculeana.
longifolia, Hort. ex Carr. in Rev. Hortic. Sér. IV. iii. (1859) 398 = flexilis.
longifolia, Karw. ex Schult. f. Syst. vii. 1715 = Nolina longifolia.
lutescens, Carr. in Rev. Hortic. Sér. IV. vii. (1858) 579 = rupicola.
macrocarpa, *Engelm. in Coult. Bot. Gaz.* vi. (1881) 224.—Arizona.
mexicana, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 224 = flexilis.
Mooreana, Hort. Peacock, ex Baker, l. c. 222 = guatemalensis.
nobilis, Hort. Peacock, ex Baker, l. c. 224 = flexilis.
obliqua, Haw. Syn. Pl. Succ. 69 = gloriosa.
orchioides, Carr. in Rev. Hortic. (1861) 369. t. 89 = filamentosa.
Parmentieri, Hort. ex Carr. l. c. Sér. IV. viii. (1859) 389 = aloifolia.
parviflora, Torr. Bot. Mex. Bound. 221 = Hesperaloë yuccaefolia.
patens, André, Illustr. Hortic. xvii. (1870) 120 = gloriosa.
Peacockii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 223.—Mexic.
pendula, Sieber, ex Carr. in Rev. Hortic. Sér. IV. viii. (1859) 488. t. 104 = gloriosa.
periculosa, Baker, in Gard. Chron. (1870) 1088 = baccata.
pitcairnifolia, Karw. ex G. Don, in Sweet, Hort. Brit. ed. III. 707, nomen (Quid ?).—Mexic.
plicata, Hort. ex C. Koch, Dendrol. ii. II. 345 = gloriosa.
plicatilis, Hort. ex C. Koch, l. c. = gloriosa.
polyphylla, Baker, in Gard. Chron. (1870) 1088 = constricta.
pruinosa, Baker, l. c. 1122 = gloriosa.
puberula, Haw. in Phil. Mag. (1828) 186 = filamentosa.
puberula, Torr. Bot. Mex. Bound. 221 = Schottii.
purpurea, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 221 = aloifolia.

YUCCA:—
quadricolor, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 221 = aloifolia.
recurva, Haw. Syn. Pl. Succ. 69 = gloriosa.
recurvata, Hort. ex Carr. in Rev. Hortic. Sér. IV. vii. (1858) 580 = Treculeana.
recurvifolia, Salisb. Parad. Lond. t. 31 = gloriosa.
revoluta, Hort. ex Carr. in Rev. Hortic. Sér. IV. vii. (1858) 580 = Treculeana.
Roezlii, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 222 = guatemalensis.
rufocincta, Haw. Suppl. Pl. Succ. 37 = gloriosa.
rupicola, *Scheele, in Linnaea*, xxiii. (1850) 143.—Texas.
scabrifolia, Baker, in Gard. Chron. (1870) 1088 = baccata.
Schottii, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1873) 46.—Arizona.
semicylindrica, Baker, in Gard. Chron. (1870) 1217 = flexilis.
serratifolia, *Karw. ex Schult. f. Syst.* vii. 1716.—Mexic.
serrulata, Haw. Syn. Pl. Succ. 70 = aloifolia.
spinosa, *H. B. & K. Nov. Gen. et Sp.* i. 289.—Mexic.
stenophylla, Hort. ex Carr. in Rev. Hortic. Sér. IV. viii. (1859) 398 = flexilis.
stenophylla, *Steud. Nom.* ed. II. ii. 795.—Mexic.
striata, Hort.—Cf. Gard. Chron. (1865) 365 (Quid ?).—Hab. ?
stricta, Sims, Bot. Mag. t. 2222 = glauca.
superba, Haw. Suppl. Pl. Succ. 36 = gloriosa.
superba, Roxb. Hort. Beng. 25 = Furcraea tuberosa.
tenuifolia, Haw. Suppl. Pl. Succ. 34 = aloifolia.
Toneliana, *Lem. Illustr. Hortic.* xii. (1865) Misc. 52.—Mexic.
tortifolia, Lindheimer, ex Torr. in Bot. Mex. Bound. 221 = rupicola.
tortilis, Hort. ex Carr. in Rev. Hortic. Sér. IV. vii. (1858) 579 = rupicola.
tortulata, Baker, in Gard. Chron. (1870) 1122 = gloriosa.
Treculeana, *Carr. in Rev. Hortic.* Sér. IV. vii. (1858) 580; *et ex Hérincq, Hortic. Franç.* (1864) 235. t. 16.—Mexic.
tricolor, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 221 = aloifolia.
undulata, Hort. ex Carr. in Rev. Hortic. Sér. IV. vii. (1858) 580 = Treculeana.
Whipplei, *Torr. in Bot. Mex. Bound.* 222, in textu.—Calif.
yucatana, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1873) 37.—Mexic.

YULANIA, Spach, Hist. Vég. Phan. vii. 462 (1839) = **Magnolia**, Linn.
conspicua, Spach, l. c. 464 = M. Yulan.
japonica, Spach, l. c. 466 = M. obovata.
Lenneana ×, Lem. Illustr. Hortic. i. (1854) t. 37 = M. *Lennei*.

Z

ZACINTHA, [Tourn.] Linn. Syst. ed. I (1735); Gaertn. Fruct. ii. 358. t. 157 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 511.
pendula, Raeusch. Nom. ed. III. 229 = verrucosa.
verrucosa, *Gaertn. Fruct.* ii. 358. t. 157. f. 7.—Europ. austr.; As. Min.

ZACINTHA, Vell. Fl. Flum. 276; vii. t. 9 (1825) = **Clavija**, Ruiz & Pav. (Myrsin.).
nutans, Vell. ll. cc. = C. integrifolia ?

ZACYNTHA, Adans. Fam. ii. 112 (1763) = **Zacintha**, Tourn. (Compos.).

ZAGA, Rafin. Fl. Tellur. ii. 33 (1836) = **Abrus**, Linn. (Legumin.).
latifolia, Rafin. l. c. = A. precatorius.
parvifolia, Rafin. l. c. = A. precatorius.

ZAHLBRUCKERA, Steud. Nom. ed. II. ii. 795 (1841), sphalm. = **Zahlbrucknera**, Reichb. (Saxifrag.).

ZAHLBRUCKNERA, Pohl, ex Nees, in Mart. Fl. Bras. ix. 16, etc. (1847) (Acanthac.).
achyranthoidea, Pohl, ex Nees, l. c. 73 (= *Teliostachya microcarpa*).—Bras.
conferta, Pohl, ex Nees, l. c. 21 = Hygrophila conferta.
fruticosa, Pohl, ex Nees, l. c. 16 = Ebermaiera mandioccana.
longifolia, Pohl, ex Nees, l. c. 21 = Hygrophila costata.
lutea, Pohl, ex Nees, in DC. Prod. xi. 210 (= *Arrhostoxylum glabrum*).—Bras.
Maranhonis, Pohl, ex Nees, in Mart. Fl. Bras. ix. 25 = Calophanes Maranhonis.
micrantha, Pohl, ex Nees, l. c. 22 = Hygrophila costata.
oblongifolia, Pohl, ex Nees, in DC. Prod. xi. 86 = Hygrophila costata.
pilosa, Pohl, ex Nees, l. c. 113 (= *Homotropium erythrorhizum*).—Bras.
repens, Pohl, ex Nees, l. c. 66 = Nelsonia campestris.
salicifolia, Pohl, ex Nees, l. c. 214 = Ruellia costata.

ZAHLBRUCKNERA, Reichb. Fl. Germ. Excurs. 551 (1832). *SAXIFRAGEAE*, Benth. & Hook. f. i. 636.
OREOSPLENIUM, Zahlb. ex Endl. Gen. 814, 1446 (1841).
paradoxa, *Reichb. Fl. Germ. Excurs.* 551.—Europ.

ZALA, Lour. Fl. Cochinch. 405 (1790) = **Pistia**, Linn.
asiatica, Lour. l. c. = P. Stratiotes.

ZALACCA, Rumph. Amboin. v. 57. t. 2 (1747). *PALMAE*, Benth. & Hook. f. iii. 932.
SALACCA, Reinw. in Syll. Ratisb. ii. (1828) 3.
SALAKKA, Reinw. ex Blume, Cat. Gew. Buitenz. 112 (1823).
affinis, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 9.—Malaya.
Blumeana, Mart. Hist. Nat. Palm. iii. 201. t. 123 = edulis.
conferta, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 16.—Malaya.
edulis, *Blume, in Schult. f. Syst.* vii. 1334; *Mart. in Wall. Pl. As. Rar.* iii. 14.—Burma; Malaya.
glabrescens, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 14.—Ins. Penang.
macrostachya, Griff. l. c. 13 = edulis.
Rumphii, Wall. Pl. As. Rar. iii. t. 223, 224 = edulis.
secunda, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 12.—Assam.
Wagneri, Hort. ex H. Wendl. in Kerch. Palm. 258, nomen (Quid ?).—Hab. ?
Wallichiana, Mart. Hist. Nat. Palm. iii. 200. tt. 118, 119, 136 = edulis.

ZALEIA, Steud. Nom. ed. II. ii. 795 (1841) = seq.

ZALEIJA, Burm. f. Fl. Ind. 110. t. 31 (1768) = **Trianthema**, Sauv. (Ficoid.).
decandra, Burm. l. c. = T. decandra.

ZALITEA, Rafin. New Fl. Am. iv. 98 (1836) = **Euphorbia**, Linn.
linearis, Rafin. l. c. 99 (Quid ?).—Am. bor.

ZALLIA, Roxb. Fl. Ind. iii. 74 (1832) = ZALEIJA, Burm. f. = **Trianthema**, Sauv. (Ficoid.).

ZALMARIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 85 = **Rondeletia**, Linn. (Rubiac.).

ZALUCANIA, Steud. Nom. ed. II. ii. 795 (1841), sphalm. = **Zaluzania**, Pers. (Compos.).

ZALUZANIA, Comm. ex Gaertn. f. Fruct. iii. 74. t. 192 (1805) = **Bertiera**, Aubl. (Rubiac.).

ZALUZANIA, Pers. Syn. ii. 473 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 362.
CHILIOPHYLLUM, DC. Prod. v. 554 (1836).
CHRYSOPHANIA, Kunth, ex Less. Syn. Comp. 224 (1832).
FERDINANDA, Lag. Gen. et Sp. Nov. 31 (1816).
HYBRIDELLA, Cass. in Bull. Soc. Philom. (1817) 12; et in Dict. Sc. Nat. xxii. 86 (1821).
angusta, Benth. & Hook. f. Gen. ii. 362, sphalm. = augusta.
asperrima, *Sch. Bip. in Flora*, xlvii. (1864) 218.—Mexic.
augusta, *Sch. Bip. l. c.* xliv. (1861) 562.—Mexic.
cinerascens, Sch. Bip. l. c. xlvii. (1864) 218, 219 = Gymnolomia cinerascens.

ZALUZANIA:—
Coulteri, *Hemsl. Diagn. Sp. Nov.* 33.—Mexic.
ensifolia, Sch. Bip. in Flora, xlvii. (1864) 216 = Gymnolomia ensifolia.
globosa, *Sch. Bip. l. c.* xliv. (1861) 564.—Mexic.
indica, Reinw. ex Blume, Mus. Bot. Lugd. Bat. i. 239 = Geniostoma rupestre.
megacephala, Sch. Bip. in Flora, xliv. (1861) 563 = augusta.
mollissima, *A. Gray, in Proc. Am. Acad.* xv. (1880) 35. —Mexic.
montagnaefolia, *Sch. Bip. in Flora*, xliv. (1861) 563. —Mexic.
myriophylla, Sch. Bip. l. c. 565 = globosa.
oppositifolia, Sch. Bip. l. c. 562 = Calea scabrifolia.
pinnatilobata, Sch. Bip. l. c. xlvii. (1864) 218, 219 = Gymnolomia pinnatilobata.
squarrosa, Sch. Bip. l. c. 216, 217 = Gymnolomia squarrosa.
triloba, *Pers. Syn.* ii. 473.—Mexic.
trilobata, Hoffmgg. Verz. Pfl. Nachtr. II. 231 = praec.

ZALUZIANSKIA, Benth. & Hook. f. Gen. ii. 944 (1876), sphalm. = seq.

ZALUZIANSKYA, F. W. Schmidt, Neue u. Selt. Pfl. 11 (1793). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 944.
NYCTERINIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 239 (1834).
Benthamiana, *Walp. Rep.* iii. 309.—Afr. austr.
capensis, *Walp. l. c.* 307.—Afr. austr.
coriacea, *Walp. l. c.* 306.—Afr. austr.
dentata, *Walp. l. c.* 307.—Afr. austr.
divaricata, *Walp. l. c.* 308.—Afr. austr.
longiflora, *Walp. l. c.* 307.—Afr. austr.
lychnidea, *Walp. l. c.*—Afr. austr.
maritima, *Walp. l. c.*—Afr. austr.
natalensis, *Bernh. in Flora*, xxvii. (1844) 834.—Afr. austr.
ovata, *Walp. Rep.* iii. 307.—Afr. austr.
peduncularis, *Walp. l. c.* 308.—Afr. austr.
pumila, *Walp. l. c.* 307.—Afr. austr.
pusilla, *Walp. l. c.* 308.—Afr. austr.
selaginoides, *Walp. l. c.*—Afr. austr.
spathacea, *Walp. l. c.* 306.—Afr. austr.
villosa, *F. W. Schmidt, Neue u. Selt. Pfl.* 11.—Afr. austr.

ZAMARIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 85 = **Randia**, Houst. (Rubiac.).

ZAMIA, Linn. Sp. Pl. ed. II. 1659 (1763). *CYCADACEAE*, Benth. & Hook. f. iii. 447.
AULACOPHYLLUM, Regel, Gartenfl. xxv. (1876) 140.
acuminata, *Oerst. ex Dyer, in Hemsl. Biol. Centr. Am. Bot.* iii. 194.—Nicaragua.
Altensteinii, Heynh. Nom. i. 862 = Encephalartos Altensteinii.
amplifolia, Hort. Bull, ex Mast. in Gard. Chron. (1878) II. 810 = Wallisii.
angustifolia, *Jacq. Coll.* iii. 263.—Ins. Baham.
angustissima, Miq. in Verh. Nederl. Inst. III. iv. (1851) 184 = angustifolia.
atropurpurea, Parment. ex Miq. Prod. Syst. Cycad. 23 = Ceratozamia mexicana.
aurea, Hort. ex Miq. in Tijdschr. Wetensch. i. (1848) 297 = Encephalartos horridus.
Baraquiniana, *Hort.—Cf. Gard. Chron.* (1868) 349; *et Regel, Gartenfl.* xxvii. (1878) 7.—Hab. ?
Beeldsnijderiana, Beelds. Cat. t. 1; ex Pritz. Ic. Ind. i. 1179 (Quid ?).—Hab. ?
boliviana, A. DC. in DC. Prod. xvi. II. 540 = Brongniartii.
Brongniartii, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1850 ?) 249. t. 4.—Bolivia; Bras.
caffra, Thunb. Prod. Pl. Cap. 92 = Encephalartos caffer.
calocoma, Miq. in Fl. des Serres, vii. (1851–52) Misc. 141 = Microcycas calocoma.
caracasana, Lodd. Cat. ex Miq. in Linnaea, xvii. (1843) 738 = Loddigesii.
Chigua, *Seem. Bot. Voy. Herald*, 201. t. 43.—N. Granat.
concinna, Hort. Booth, ex Regel, Gartenfl. xxvii. (1878) 9 = pumila.
concinna, Hort. Par. ex H. Wendl. Ind. Palm. 52, nomen (Quid ?).—Hab. ?
corallipes, J. Versch. Cat. n. 17 (1873–74) 5, 7 = Macrozamia spiralis.

ZAMIA:—
Corsoniana, G. Don, in Loud. Gard. Mag. xviii. (1842) 371 = Cycas Rumphii.
crassifolia, *Hort. ex T. Moore, Fl. & Pom.* (1877) 14, 15.—Hab. ?
cycadifolia, S. Brunn. in Flora, viii. (1825) 733 = Macrozamia Fraseri.
cycadifolia, *Dyer, in Hemsl. Biol. Centr. Am. Bot.* iii. 195.—Mexic.
cycadifolia, Jacq. Fragm. 27 = Encephalartos cycadifolius.
cycadifolia, Lodd. ex Miq. in Linnaea, xvii. (1843) 719 = Encephalartos caffer.
Cycadis, Linn. f. Suppl. 443 = Encephalartos caffer.
cylindrica, Hort. Booth, ex Regel, Gartenfl. xxvii. (1878) 9 = pumila.
cylindrica, Liebm. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 195 = Leiboldii.
debilis, *Ait. Hort. Kew.* ed. I. iii. 478.—Ind. occ.
debilis, Linn. f. ex Salisb. Prod. 400 = laeta.
debilis, Lodd. Bot. Cat. t. 155 = angustifolia.
dentata, Hort. Angl. ex Link, Enum. Hort. Berol. ii. 432, nomen (Quid ?).—Hab. ?
dentata, Voigt, in Syll. Ratisb. ii. (1828) 53 = tenuis.
elegans, Hort. ex Yates, in Proc. Linn. Soc. ii. (1849) 18, 19 = Encephalartos caffer.
elliptica, Lodd. ex Miq. in Linnaea, xvii. (1843) 723 = Encephalartos caffer.
elongata, Heynh. Nom. i. 862 = Encephalartos elongatus.
eriolepis, Hort. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 195 = Leiboldii.
Fischeri, *Miq. in Hort. Houttean.* 20; *et in Tijdschr. Wetensch.* i. (1848) 201.—Am. caled.
floridana, A. DC. in DC. Prod. xvi. II. 544 = angustifolia.
Fraseri, Van Houtte, ex Regel, Gartenfl. xxv. (1876) 229 = Macrozamia spiralis.
Friderici-Guilielmi, Hort. Parment. ex Miq. Prod. Syst. Cycad. 22 = Dioon edule.
furfuracea, *Ait. Hort. Kew.* ed. I. iii. 477.—Mexic.
fusca, Hort. Paull. ex Regel, Gartenfl. xxv. (1876) 261 = Cerarozamia Katzeriana.
Galeotti, De Vriese, in Hoev. & De Vriese, Tijdschr. xii. (1845) 24 = Ceratozamia mexicana.
gleina, Hort. ex Miq. in Linnaea, xvii. (1843) 729 = Encephalartos horridus.
grandis, Hort.—Cf. Gard. Chron. (1865) 365 (Quid ?). —Hab. ?
Gutierrezi, *Sauv. Anal. Acad. Cienc. Nat. Haban.* (1868) (reimp. 2).—Cuba.
horrida, Jacq. Fragm. 27 = Encephalartos horridus.
horrida, Lodd. ex Miq. in Linnaea, xvii. (1843) 725 = Encephalartos lanuginosus.
humilis, Salisb. Prod. 400 = pumila.
integrifolia, *Ait. Hort. Kew.* ed. I. iii. 478.—Ind. occ.
integrifolia, A. Gray, ex A. DC. in DC. Prod. xvi. II. 544 = angustifolia.
integrifolia, Pursh, Fl. Am. Sept. ii. 648 = pumila.
integrifolia, Rich. Conif. et Cycad. 191. t. 27 = media.
Kickxii, *Miq. Monog. Cycad.* 71. t. 8.—Cuba.
laeta, *Salisb. Prod.* 400 (sp. dub.).—Hab. ?
lanuginosa, Jacq. Fragm. 28 = Encephalartos lanuginosus.
latifolia, *Lodd. Cat.* n. 187; *ex Miq. in Linnaea*, xvii. (1843) 733.—Honduras.
Lawsoniana, *Dyer, in Hemsl. Biol. Centr. Am. Bot.* iii. 195.—Mexic.
Lehmanniana, Eckl. & Zeyh. ex Eckl. in Otto & Dietr. Allg. Gartenz. i. (1833) 158; et ex Lehm. Pugill. vi. 7 = Encephalartos Lehmanni.
Leiboldii, *Miq. in Linnaea*, xix. (1847) 425.—Mexic.
Lindeni, *Regel, ex André, in Illustr. Hortic.* xxii. (1875) 23. t. 195.—Ecuador.
Lindleyana, H. Wendl. Ind. Palm. 53 = Z. Chigua.
Lindleyi, Warsz. in Otto & Dietr. Allg. Gartenz. xix. (1851) 146 = Z. Chigua.
linearifolia, Linden, in Illustr. Hortic. xxviii. (1881) 32, nomen (Quid ?).—Cuba.
linifolia, Hort. Pawl. ex Regel, Gartenfl. xxvii. (1878) 12 = angustifolia.
Loddigesii, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 72.—Mexic.
longifolia, Jacq. Fragm. 28 = Encephalartos longifolius.
Mackenii, Hort. ex Miq. in Verh. Kon. Akad. Wetensch. II. iii. (1869) 55 = Macrozamia Pauli-Guilielmi.

ZAMIA:—
Macleani, Regel, Gartenfl. xxv. (1876) 371, sphalm. = *Maeleni*.
macrophylla, Parment. ex Miq. Prod. Syst. Cycad. 23 = Ceratozamia mexicana.
Maeleni, Miq. in Linnaea, xviii. (1844) 97 = Dioon edule.
manicata, *Linden, ex Regel, Gartenfl.* (1878) 8; *et in Illustr. Hortic.* xxviii. (1881) 32, nomen.—Bras.
media, *Jacq. Hort. Schoenb.* iii. 77. tt. 397, 398.—Ind. occ.
media, Sims, Bot. Mag. t. 1838 = pumila.
mexicana, Miq. Prod. Cycad. 13, 28 = Loddigesii.
Miqueli, Hort. Adelaide, ex Regel, Gartenfl. xxv. (1876) 230 = Macrozamia Miquelii.
montana, *A. Br. in Monatsb. Akad. Berl.* (1875) 376. —N. Granat.
multifoliolata, *A. DC. in DC. Prod.* xvi. II. 545.—Cuba.
muricata, *Willd. Sp. Pl.* iv. 847.—Am. trop.
nana, Hort. ex Miq. in Linnaea, xvii. (1843) 729 = Encephalartos horridus.
obliqua, *A. Br. in Monatsb. Akad. Berl.* (1875) 376; *Regel, in Illustr. Hortic.* xxiv. (1877) 140. t. 289.—N. Granat.
occidentalis, Lodd. Cat. n. 177; ex Miq. in Linnaea, xvii. (1843) 711 = Encephalartos pungens.
Ottonis, *Miq. in Linnaea*, xvii. (1843) 740.—Cuba.
pallida, *Salisb. Prod.* 401 (sp. dub.).—Hab. ?
parasitica, Poepp. ex Eichl. in Mart. Fl. Bras. iv. I. 414 = Poeppigiana.
picta, *Dyer, in Hemsl. Biol. Centr. Am. Bot.* iii. 194. —Mexic. ?
Poeppigiana, *Mart. & Eichl. in Mart. Fl. Bras.* iv. I. 414. t. 109.—Peruv.
Potemkini, *Hort. ex Miq. in Verh. Kon. Akad. Wetensch.* II. iv. (1870) 30.—Hab. ?
prasina, Bull, Cat. (1882) 20; ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 194 = latifolia.
Princeps, Hort. Bull, ex Dyer, l. c. = Z. Chigua.
prunifera, Lodd. Cat. ex G. Don, in Loud. Hort. Brit. 403, nomen = Encephalartos pruniferus.
Pseudo-parasitica, *Yates, in Seem. Bot. Voy. Herald*, 202, 253.—Panama.
pumila, Hort. ex Regel, Gartenfl. xxvii. (1878) 11 = pygmaea.
pumila, *Linn. Sp. Pl.* ed. II. 1659.—Ind. occ.
pungens, Ait. Hort. Kew. ed. I. iii. 478 = Encephalartos pungens.
pungens, Hort. ex Miq. in Linnaea, xvii. (1843) 725 = Encephalartos lanuginosus.
pungens, Linn. f. ex Salisb. Prod. 401 = pallida.
pungens, Lodd. ex Miq. in Linnaea, xvii. (1843) 725 = Encephalartos elongatus.
pygmaea, *Sims, Bot. Mag.* t. 1741.—Ind. occ.
repanda, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 403, nomen.—Hab. ?
Roezli, Regel, in Linden, Cat. n. 90 (1873) 10; et in Illustr. Hortic. xx. (1873) 126. t. 133–134 = Pseudo-parasitica.
serrulata, Lodd. Cat. ex Miq. in Linnaea, xvii. (1843) 738 = Loddigesii.
Sieboldi, Cf. Gard. Chron. (1878) II. 594, sphalm. = Leiboldii.
Skinneri, *Warsz. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 146.—Am. centr.
spartea, *A. DC. in DC. Prod.* xvi. II. 539.—Mexic.
spinosa, Lodd. Cat. n. 168; ex Miq. in Linnaea, xvii. (1843) 721 = Encephalartos Altensteinii.
spinosissima, Hort. ex Miq. l. c. = praec.
spinulosa, Heynh. Nom. i. 862 = Encephalartos Lehmanni.
spinulosa, Hort. ex Miq. in Linnaea, xvii. (1843) 721 = Encephalartos Altensteinii.
spiralis, R. Br. Prod. 348 = Macrozamia spiralis, Fraseri.
spiralis, Salisb. Prod. 401 = Macrozamia spiralis.
stricta, Griseb. Cat. Pl. Cub. 217 = multifoliolata.
stricta, Miq. in Verh. Nederl. Inst. III. iv. (1851) 183 = angustifolia.
tenuifolia, Fisch. ex Miq. in Linnaea, xix. (1847) 429 = Fischeri.
tenuis, *Willd. Sp. Pl.* iv. 846.—Ins. Baham.
terrestris, Hort. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 195 = Loddigesii.
tonkinensis, *Linden & Rodigas, in Illustr. Hortic.* xxxii. (1885) 27. t. 547.—Tonkin.

ZAMIA :—
tricuspidata, Hort. in Verh. Ver. Bef. Gartenb. v. 186 = Encephalartos horridus.
tridentata, Willd. Sp. Pl. iv. 846 = Encephalartos tridentatus.
unidentata, Hort. ex Miq. in Linnaea, xvii. (1843) 715 = Encephalartos tridentatus.
variegata, *Warsz. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 253, nomen.—Hab. ?
vernicosa, Hort.—Cf. Gard. Chron. (1868) 349 (Quid?). —Hab. ?
Verschaffeltii, *Miq. in Arch. Néerl.* v. (1870) 82; *et in Verh. Kon. Akad. Wetensch.* II. iv. (1870) 31. —Mexic.
vestita, *Van Houtte, Cat.* 10; *ex Miq. in Tijdschr. Wetensch.* iii. (1849) 300.—Am. centr.
villosa, Gaertn. Fruct. i. t. 3 = Encephalartos cycadifolius.
villosa, Herb. Van Royen, ex A. DC. in DC. Prod. xvi. II. 546 = Encephalartos caffer.
Wallisii, *Hort. Veitch, ex A. Br. in Monatsb. Akad. Berl.* (1875) 376.—N. Granat.
Yatesii, Miq. in Verh. Nederl. Inst. III. iv. (1851) 182 = angustifolia.

ZAMIOCULCAS, Schott, Syn. Aroid. 71 (1856). *AROIDEAE*, Benth. & Hook. f. iii. 993.
GONATOPUS, Engl. in DC. Monog. ii. 208 (1879).
Boivini, *Decne. in Bull. Soc. Bot. Fr.* xvii. (1870) 321. —Afr. trop.
Loddigesii, *Schott, Syn. Aroid.* 71.—Afr. trop.

ZAMZELA, Rafin. Sylva Tellur. 90 (1838) = **Hirtella**, Linn. (Rosac.).
racemosa, Rafin. l. c. = H. americana ?
rugosa, Rafin. l. c. = H. rugosa.

ZANICHELIA, Gilib. Exercit. ii. 419 (1792), sphalm. = seq.

ZANICHELLIA, Roth, Tent. Fl. Germ. ii. II. 420 (1793), err. typ. = seq.

ZANNICHALLIA, Reut. in Compt. Rend. Soc. Hallér. (1854–56) 129, sphalm. = seq.

ZANNICHELLA, Linn. Syst. ed. I. (1735) = seq.

ZANNICHELLIA, Mich. ex Linn. Syst. ed. I (1735); Gen. ed. I. 278 (1737). *NAIADACEAE*, Benth. & Hook. f. iii. 1016.
PELTA, Dulac, Fl. Hautes-Pyr. 43 (1867).
aculeata, Schur, in Oestr. Bot. Zeitschr. xx. (1870) 203 = palustris.
brachystemon, J. Gay, ex Reut. Cat. Gr. Jard. Genèv. (1854) 4 = palustris.
contorta, Cham. in Linnaea, ii. (1827) 231 = palustris.
cylindrocarpa, Koern. ex C. Muell. in Walp. Ann. vi. 3 = Lepilaena cylindrocarpa.
dentata, Willd. Sp. Pl. iv. 181 = palustris.
digyna, J. Gay, ex Bréb. Fl. Norm. ed. II. 252; ed. III. 289 = palustris.
geniculata, Gilib. Exercit. ii. 419 = palustris.
gibberosa, Reichb. in Moessl. Handb. ed. II. iii. 1591 = palustris.
graminifolia, Gilib. Exerc. Bot. in Schola Princip. Univ. Viln. (1772) 31 = palustris ?
intermedia, Nielsen, in Bot. Tidsskr. Ser. II. i. (1872) 205 = seq. ?
intermedia, Torr. in Beck, Bot. N. & M. St. 385 = palustris.
laevis, *Presl, Bot. Bemerk.* 112.—Afr. austr.
indica, Rottl. ex Royle, Illustr. Bot. Himal. 404 (Quid?). — Ind. or.
macrostemon, J. Gay, ex Willk. & Lange, Prod. Fl. Hisp. i. 26 = palustris.
major, Boenn. ex Reichb. in Moessl. Handb. ed. II. iii. 1591 = palustris.
marina, Nielsen, in Bot. Tidsskr. Ser. II. i. (1872) 204 = palustris.
maritima, Nolte, Novit. Fl. Holsat. 75 = palustris.
palustris, *Linn. Sp. Pl.* 969.—Europ.; Am. bor.
pedicellata, Buch.-Ham. ex Wall. Cat. n. 5185; Fries, Mant. iii. 133 = palustris.
pedunculata, Fuss, Ber. Phan. Fl. Siebenb. (1854) 13 = Ruppia rostellata.

ZANNICHELLIA :—
pedunculata, Reichb. in Moessl. Handb. ed. II. iii. 1591 = palustris.
peltata, Bertol. Fl. Ital. x. 10 = palustris.
polycarpa, Nolte, Novit. Fl. Holsat. 75 = palustris.
Preissii, Lehm. Pl. Preiss. ii. 3 = Lepilaena Preissii.
pumila, Le Gall, Fl. Morb. 568 (Quid ?).—Gallia.
radicans, Wallm. in Liter. Bot. Ges. Regensb. xi. (1841) 20 = palustris.
repens, Boenn. Prod. Fl. Monast. 272 = palustris.
Rosenii, Wallm. in Liter. Bot. Ges. Regensb. xi. (1841) 18 = palustris.
stylaris, *Presl, Bot. Bemerk.* 112.—Afr. austr.
tenuis, Reut. Cat. Gr. Jard. Genèv. (1854) 4; et in Compt. Rend. Soc. Hallér. (1854–56) 129 = palustris.
tuberosa, *Lour. Fl. Cochinch.* 543.—Cochinch.
vaginalis, Delile, ex Kunth, Enum. Pl. iii. 126 = Althenia filiformis.

ZANONIA, Linn. Coroll. Gen. 19 (1737). *CUCURBITACEAE*, Benth. & Hook. f. i. 839.
angulata, Wall. Pl. As. Rar. ii. 28 = indica.
capricornica, F. Muell. Fragm. vii. 61 = Alsomitra capricornica.
cissoides, Wall. Pl. As. Rar. ii. 28 = Gynostemma pedatum.
clavigera, Wall. l. c. = Alsomitra clavigera.
heterosperma, Wall. l. c. 29 = Gomphogyne heterosperma.
Hookeri, F. Muell. Census, 76 = Alsomitra Hookeri.
indica, *Linn. Sp. Pl.* ed. II. 1457.—Ind. or.; Malaya.
integerrima, Wall. ex Cogn. in DC. Monog. Phan. iii. 931 = Alsomitra clavigera.
laxa, Wall. Pl. As. Rar. ii. 29 = Gynostemma Wightianum.
macrocarpa, *Blume, Bijdr.* 937.—Java.
oblonga, Wall. ex Steud. Nom. ed. II. ii. 796, nomen (Quid ?).—Ind. or.
pedata, Miq. Fl. Ind. Bat. i. 1. 683 = Gynostemma pedatum.
sarcophylla, Wall. Pl. As. Rar. ii. 28 = Alsomitra sarcophylla.
Stephensiana, F. Muell. Fragm. viii. 181 = Alsomitra Stephensiana.
timorana, Span. in Linnaea, xv. (1841) 205 = Alsomitra timorana.
timorensis, Walp. Rep. ii. 194, sphalm. = praec.
Wightiana, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 356 B = Gynostemma Wightianum.

ZANONIA, [Plum.] Cram. Disp. Syst. 75 (1803) = **Campelia**, Rich. (Commel.).
bibracteata, Cram. l. c. = C. Zanonia.

ZANTEDESCHIA, C. Koch, Ind. Sem. Hort. Berol. (1854) App. 9, partim = **Schismatoglottis**, Zoll. & Mor. (Aroid.).
alba, C. Koch, l. c. = Homalomena alba.
calyptrata, C. Koch, l. c. = S. calyptrata.
cordata, C. Koch, l. c. = Homalomena aromatica.
foetida, C. Koch, l. c. = Homalomena aromatica.
rubens, C. Koch, l. c. = Homalomena rubescens.
virosa, C. Koch, l. c. = Colocasia virosa.

ZANTEDESCHIA, Spreng. Syst. iii. 765 (1826) = **Richardia**, Kunth (Aroid.).
aethiopica, Spreng. l. c. = R. africana.
albo-maculata, Baill. in Bull. Soc. Linn. Par. i. (1880) 254; et ex Engl. Bot. Jahrb. iv. (1883) 64 = R. albomaculata.
angustiloba, Engl. l. c. = R. angustiloba.
aromatica, Spreng. Syst. iii. 765 = Homalomena aromatica.
asperata, C. Koch, in Ind. Sem. Hort. Berol. (1853) App. 5 = Philodendron asperatum.
hastata, Engl. Bot. Jahrb. iv. (1883) 64 = R. hastata.
macrocarpa, Engl. l. c.—Afr. austr.
melanoleuca, Engl. l. c. = R. melanoleuca.
occulta, Spreng. Syst. iii. 765 = Homalomena aromatica.
Rehmanni, Engl. Bot. Jahrb. iv. (1883) 63.—Afr. austr.

ZANTHORHIZA, L'Hérit. Stirp. Nov. 79. t. 38 (1784). *RANUNCULACEAE*, [Benth. & Hook. f. i. 9.]
XANTHORRHIZA, Marsh. Arbust. 167 (1785).

ZANTHORHIZA :—
apiifolia, *L'Hérit. Stirp. Nov.* 79. t. 38.—Am. bor.
simplicissima, Marsh. Arbust. 168 = apiifolia.
tinctoria, Woodh. in Med. Repos. N. York, v. (1802) 160 = apiifolia.

ZANTHOXILON, Franch. & Sav. Enum. Pl. Jap. i. 72 (1875), sphalm. = seq.

ZANTHOXYLON, Walt. Fl. Carol. 52, 243 (1788) = seq.

ZANTHOXYLUM, Linn. Hort. Cliff. 487 (1737). *RUTACEAE*, Benth. & Hook. f. i. 297.
AUBERTIA, Bory, Voy. i. 356. t. 18 (1804).
BLACKBURNIA, Forst. Char. Gen. t. 6 (1776).
CURTISIA, Schreb. Gen. 199 (1789).
DORATIUM, Soland. ex Jaume St. Hil. Expos. ii. 267 (1805).
FAGARA, Linn. Syst. ed. X. 897 (1759).
LACURIS, Buch.-Ham. ex Wall. Cat. n. 7119 (1849).
LANGSDORFIA, Leand. Sacram. in Denkschr. Akad. Muench. (1819) 229.
MIOPTRILA, Rafin. Am. Man. Mulb. 37 (1839).
OCHROXYLUM, Schreb. Gen. 826 (1791).
PENTANOME, [Moç. & Sesse,] ex DC. Prod. i. 725, in syn.
POHLANA, Leand. Sacram. in Denkschr. Akad. Muench. (1819) 229.
PSEUDOPETALON, Rafin. Fl. Ludov. 107 (1817).
PTEROTA, P. Br. Hist. Jamaic. 146. t. 5 (1756).
TENOREA, Rafin. Specch. i. 193 (1814).
TIPALIA, Dennst. Schluess. Hort. Malab. 31 (1818).
TOBINIA, Desv. in Ham. Prod. Pl. Ind. Occ. 56 (1825).
XANTHOXYLON, Spreng. Anleit. ii. II. 655 (1818).
acanthopodium, *DC. Prod.* i. 727.—Reg. Himal.
accedens, Miq. Fl. Ind. Bat. i. II. 671 = Evodia accedens.
aculeatissimum, *Engl. in Mart. Fl. Bras.* xii. II. 176. —Bolivia.
aculeatum, Macfad. Fl. Jamaic. i. 191 = acuminatum.
acuminatum, Macfad. l. c. 192 = coriaceum.
acuminatum, *Sw. Fl. Ind. Occ.* i. 575.—Jamaic.
acutifolium, *Engl. in Mart. Fl. Bras.* xii. II. 170.—Bras.
affine, *H. B. & K. Nov. Gen. et Sp.* vi. 3.—Mexic.
ailanthoides, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 138.—Formosa; Japon.
alatum, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 768.—Reg. Himal.; China.
alatum, *Steud. Nom.* ed. II. ii. 796.—Afr. austr.
alatum, Wall. Cat. n. 1209, partim = acanthopodium.
album, *Vahl, Eclog. Am.* iii. 47.—Am. austr.
alveolatum, Shuttl. ex S. Wats. Bibl. Ind. N. Am. Bot. 156 = Z. Clava-Herculis.
americanum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
amoyense, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 277.—N. Granat.
andamanicum, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 199.—Ins. Andaman.
araliaceum, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 274 = Casimiroa edulis.
arenarium, *Engl. in Mart. Fl. Bras.* xii. II. 164. —Bras.
argenteum, Lodd. Cat. ex G. Don, in Loud. Hort. Brit. 398, nomen (Quid ?).—Hab. ?
armatum, DC. Prod. i. 727, sphalm. = alatum.
Arnottianum, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 146.—Ins. Bonin.
aromaticum, Griseb. Fl. Brit. W. Ind. 138 = caribaeum.
aromaticum, Miq. Fl. Ind. Bat. i. II. 670 = Evodia glabra.
aromaticum, Willd. Sp. Pl. iv. 755 = Z. Clava-Herculis.
articulatum, *Engl. in Mart. Fl. Bras.* xii. II. 161. —Bras.
Aubertia, *DC. Prod.* i. 725.—Ins. Borbon.
Aubertianum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = praec.
australasicum, A. Juss. in Mém. Mus. Par. xi. (1825) 503 = Geijera parviflora.
australe, *G. Don, in Loud. Hort. Brit.* 398.—Austral.
Avicennae, *DC. Prod.* i. 726.—China.
Bajarnandia, *Wall. Cat.* n. 7118 (sp. dub.).—Assam.
Balansae, *Baill. Adansonia*, xi. (1875) 300.—Ins. Loyalt.

ZANTHOXYLUM:—
Blackburnia, *Benth. Fl. Austral.* i. 363.—Austral.
bombacifolium, *A. Rich. Ess. Fl. Cub.* 329. t. 35.—Cuba.
borbonicum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 440.—Ins. Borbon.
brachyacanthum, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 65.—Austral.
Bretschneideri, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 73.—China.
Budrunga, *DC. Prod.* i. 728 (= seq. ?).—Ind. or.
Budrunga, *Wall. Cat.* n. 1211.—Reg. Himal.; Burma.
Bungeanum, Maxim. in Bull. Acad. Pétersb. xvi. (1871) 212 = Bungei.
Bungei, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 209 = alatum.
Bungei, *Planch. l. c.* Sér. III. xix. (1853) 82.—China.
camphoratum, *Planch. & Linden, l. c.*—Venezuela.
capense, *Harv. in Harv. & Sond. Fl. Cap.* i. 446.—Afr. austr.
caribaeum, Gaertn. Fruct. i. 333 = americanum.
caribaeum, *Lam. Encyc.* ii. 39.—N. Granat.
carolinianum, Gaertn. Fruct. i. 333 = caribaeum.
carolinianum, Lam. Encyc. ii. 39 = Z. Clava-Herculis.
Catesbianum, Rafin. Med. Fl. ii. 114 = Z. Clava-Herculis.
cauliflorum, Steud. Nom. ed. I. 897 = americanum.
Chiloperone, *Mart. ex Engl. in Mart. Fl. Bras.* xii. ii. 170.—Bras.
ciliatum, *Engl. l. c.* 180.—Venezuela.
cinereum, *Engl. l. c.* 172.—Bras.
Clava-Herculis, DC. Prod. i. 727 = caribaeum.
Clava-Herculis, Lam. Encyc. ii. 38 = americanum.
Clava-Herculis, *Linn. Sp. Pl.* 270.—Ins. Carib.
Clava-Herculis, Lour. Fl. Cochinch. 659 = Avicennae.
clavatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = Z. Clava-Herculis.
Coco, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 168.—Reg. Argent.
connaroides, Wight & Arn. Prod. 148 = Heynea trijuga.
coriaceum, A. Rich. Ess. Fl. Cub. 326. t. 34 = emarginatum.
coriaceum, *Walp. Rep.* i. 521.—Jamaic.
coriaceum, Wright, ex A. Gray, Pl. Wright. i. 31 = Z. Clava-Herculis.
corymbosum, *DC. Prod.* i. 726.—Ind. occ.
crenatum, Wall. Cat. n. 1216 = Z. Budrunga.
cribrosum, *Spreng. Syst.* i. 946.—Ins. S. Doming.
cuiabense, *Engl. in Mart. Fl. Bras.* xii. ii. 163.—Bras.
Culantrilo, *H. B. & K. Nov. Gen. et Sp.* vi. 2.—Am. trop.
cuspidatum, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 329.—China.
Danielli, Benn. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 201. t. 5 (Evodiae sp.).—China.
digynum, Engelm. ex A. Gray, Pl. Wright. i. 31 = Z. Clava-Herculis.
dipetalum, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 319.—Ins. Sandvic.
dumosum, *A. Rich. Ess. Fl. Cub.* 327.—Cuba.
duplicipunctatum, *Wright, ex Griseb. Cat. Pl. Cub.* 49.—Cuba.
elegans, *Engl. in Mart. Fl. Bras.* xii. ii. 159.—Bras.
elegans, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 141.—Bras.
elephantiasis, Macfad. Fl. Jamaic. i. 193 = caribaeum.
emarginatum, *Sw. Fl. Ind. Occ.* i. 572.—Ind. oc.
emarginellum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 22.—Japon.
euneuron, Miq. Fl. Ind. Bat. Suppl. 532 = Evodia euneura.
Finlaysonianum, *Wall. Cat.* n. 7114.—Siam ?
flavum, *Vahl, Eclog. Am.* iii. 48.—Am. austr.
floribundum, Wall. Cat. n. 1206 = Toddalia aculeata.
floridanum, Nutt. Sylva, iii. 14. t. 85 = caribaeum.
fraxineum, Willd. Sp. Pl. iv. 757 = americanum.
fraxinifolium, Marsh. Arbust. 167 = americanum.
fraxinifolium, Walt. Fl. Carol. 243 = Z. Clava-Herculis.
furfuraceum, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 272.—N. Granat.
Gardneri, *Engl. in Mart. Fl. Bras.* xii. ii. 165.—Bras.
Ghisbreghtii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 274.—Mexic.

ZANTHOXYLUM:—
glandulosum, *Teijsm. & Binn. Cat. Hort. Bog.* 234, nomen.—Java.
gracile, *Wright, in Sauv. Fl. Cub.* 19.—Cuba.
grandifolium, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 273.—N. Granat.
guianense, *Klotzsch, in Schomb. Faun. Fl. Brit. Gui.* 1189, nomen.—Guian.
Hamiltonianum, *Wall. Cat.* n. 7117.—Reg. Himal.
hastile, Steud. Nom. ed. II. ii. 796, sphalm. = *hostile*.
hermaphroditum, *Willd. Sp. Pl.* iv. 756.—Guian.
heterophyllum, *Sm. in Rees, Cycl.* xxxix. n. 14.—Ins. Mascar.
hirsutum, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 450 = Z. Clava-Herculis.
horridum, *DC. Prod.* i. 728.—Japon.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 597.—Java.
hostile, Wall. Cat. n. 1210 = alatum, acanthopodium.
hyemale, *A. St. Hil. Pl. Us. Bras.* t. 37.—Bras.
javanicum, Walp. Rep. i. 519 = Picrasma javanica.
juglandifolia, *Willd. Sp. Pl.* iv. 756.—Ins. S. Doming.
juniperinum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 77.—Am. trop.
kauaense, *A. Gray, Bot. U. St. Expl. Exped.* i. 354.—Ins. Sandvic.
khasianum, *Hook. f. Fl. Brit. Ind.* i. 494.—Reg. Himal.
Kibada, Siebold, in Verh. Batav. Gen. xii. (1830) 48 = alatum ?
Lamarkianum, Cham. in Linnaea, v. (1830) 58 = Evodia triphylla.
lanceolatum, Poir. Encyc. Suppl. ii. 293 = Z. Clava-Herculis.
Langsdorffii, Mart. in Flora, xx. (1837) ii. Beibl. 97 = rhoifolium.
lanuginosum, Kostel. Allg. Med.-Pharm. Flora, v. 1783 = Evodia latifolia.
latespinosum, *Engl. in Mart. Fl. Bras.* xii. ii. 173.—Bras.
latifolium, G. Don, Gen. Syst. i. 804 = Evodia latifolia.
lentiscifolium, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 329 = Z. Avicennae.
Leprieurii, *Guill. & Perr. Fl. Seneg. Tent.* 141.—Afr. trop.
Limoncello, *Planch. & Oerst. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 312.—N. Granat.
limonifolium, *Wall. ex Voigt, Hort. Suburb. Calc.* 185, nomen.—Reg. Himal.
longifolium, Wall. Cat. n. 7115 = myriacanthum.
lucidum, Miq. Fl. Ind. Bat. Suppl. 532 = Evodia lucida.
lucidum, *A. St. Hil. Fl. Bras. Mer.* i. 92, in nota.—Bras.
lucidum, Wall. Cat. n. 1212 = ovalifolium.
macrocarpum, C. Muell. in Walp. Ann. vii. 527, sphalm. = microcarpum.
macrophyllum, Miq. Fl. Ind. Bat. i. ii. 670 = Evodia accedens.
macrophyllum, Nutt. Sylva, iii. 10 = Z. Clava-Herculis.
macrophyllum, *Oliver, Fl. Trop. Afr.* i. 304.—Afr. trop.
macrophyllum, Ryan, ex Eggers, Fl. St. Croix (1879), 38 = Z. Ochroxylum.
macrospermum, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 276.—N. Granat.
mantchuricum, Benn. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 200 = schinifolium.
Marambong, Miq. Fl. Ind. Bat. Suppl. 533 = Evodia Roxburghiana.
martinicense, *DC. Prod.* i. 726.—Ins. Martinic.
maviense, *H. Mann, in Proc. Bost. Soc. Nat. Hist.* x. (1866) 319.—Ins. Sandvic.
Mayu, *Bert. in Ann. Sc. Nat.* Sér. I. xxi. (1830) 345.—Ins. J. Fernand.
melanacanthum, *Planch. l. c.* Sér. III. xix. (1853) 82; *et ex Oliver, Fl. Trop. Afr.* i. 305.—Afr. trop.
melanostictum, *Schlecht. & Cham. in Linnaea*, v. (1830) 231.—Mexic.
microcarpum, *Griseb. Fl. Brit. W. Ind.* 138.—Ins. Dominic.
microphyllum, Desf. Cat. Hort. Par. ed. III. 256, nomen (Quid ?).—Hab. ?
minutiflorum, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 278.—N. Granat.
mite, Willd. Enum. Hort. Berol. 1013 = americanum.
monogynum, *A. St. Hil. in Bull. Soc. Philom.* (1823) 129.—Bras.

ZANTHOXYLUM:—
montanum, *Blume, Bijdr.* 248.—Java.
myriacanthum, *Wall. Cat.* n. 1214.—Malaya.
Naranjillo, *Griseb. in Goett. Abh.* xxiv. (1879) 76.—Reg. Argent.
natalense, Hochst. in Flora, xxvii. (1844) 304 = Cnestis natalensis.
nemorale, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 92.—Bras.
nigrum, *Mart. l. c.* xxii. (1839) i. *Beibl.* 11.—Bras.
nilagiricum, Miq. Anal. Bot. Ind. iii. 6 = Evodia Roxburghiana.
nitidum, Bunge, Enum. Pl. Chin. Bor. 13 = Bungei.
nitidum, *DC. Prod.* i. 727.—China.
nitidum, A. St. Hil. Fl. Bras. Mer. i. 77 = lucidum.
nitidum, Wall. Cat. n. 1207 = Toddalia aculeata.
Novae-Zeelandiae, A. Rich. Ess. Fl. N. Zél. 291 = Hedycarya arborea.
oblongum, Wall. Cat. n. 1218 = Z. Rhetsa.
obscurum, *Engl. in Mart. Fl. Bras.* xii. ii. 169.—Bras.
obtusifolium, G. Don, Gen. Syst. i. 804 = Evodia obtusifolia.
obtusifolium, *Poir. Encyc. Suppl.* ii. 293.—Ind. or.
Ochroxylum, *DC. Prod.* i. 725.—Am. trop.
oppositifolium, DC. l. c. 728 = Boronia alata.
ovalifolium, *Wight, Illustr.* i. 169.—Ind. or.
oxyphyllum, *Edgew. in Trans. Linn. Soc.* xx. (1846) 42.—Reg. Himal.
paniculatum, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 12.—Ins. Rodriguez.
parviflorum, *Benth. Fl. Austral.* i. 363.—Austral.
pauciflorum, *Engl. in Mart. Fl. Bras.* xii. ii. 164.—Bras.
Peckoltianum, *Engl. l. c.* 155.—Bras.
peltophorum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 442 = rhoifolium.
Pentanome, *DC. Prod.* i. 725.—Mexic.
Perrottetii, DC. l. c. 726 = rhoifolium ?
petiolare, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 140.—Bras.
phyllopterum, *Wright, in Sauv. Fl. Cub.* 19.—Cuba.
pimpinelloides, *DC. Prod.* i. 726.—Ins. S. Doming.
piperitum, Benn. in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 200 = Avicennae.
piperitum, *DC. Prod.* i. 725.—China; Japon.
piperitum, Hook. & Arn. Bot. Beech. Voy. 261 = Arnottianum.
pistacifolium, *Griseb. Cat. Pl. Cub.* 49.—Cuba.
planispinum, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 138 = alatum.
Pohlianum, *Engl. in Mart. Fl. Bras.* xii. ii. 166.—Bras.
polyacanthum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 443 = stelligerum.
polygamum, Schum. & Thonn. Beskr. Guin. Pl. 433 = senegalense.
praecox, *A. St. Hil. Fl. Bras. Mer.* i. 76.—Bras.
pteleaefolium, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 330 = Evodia triphylla.
Pterota, *H. B. & K. Nov. Gen. et Sp.* vi. 3.—Ind. occ.
Pterota, Seem. Bot. Voy. Herald, 94 = Z. Limoncello.
pterotum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = Z. Pterota.
pubescens, A. St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 141 = rhoifolium.
pumilum, Zipp. ex Span. in Linnaea, xv. (1841) 185, nomen (Quid ?).—Ins. Timor.
punctatum, *Vahl, in West, Bidr. Ste-Croix*, 310.—Ind. occ.
quinduense, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 275.—N. Granat.
ramiflorum, Michx. Fl. Bor. Am. ii. 235 = americanum.
Reguellianum, *Engl. in Mart. Fl. Bras.* xii. ii. 174.—Bras.
Rhetsa, *DC. Prod.* i. 728.—Ind. or.
rhetsum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = praec.
rhizinoides, *Blume, Bijdr.* 248.—Java.
rhoifolium, *Lam. Encyc.* ii. 39.—Am. trop.
ricinoides, Walp. Rep. i. 520, sphalm. = rhizinoides.
Riedelianum, *Engl. in Mart. Fl. Bras.* xii. ii. 162.—Bras.
rigidum, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 756.—Am. austr.
Roxburghianum, Cham. in Linnaea, v. (1830) 58 = Evodia Roxburghiana.
rubescens, *Planch. ex Hook. Niger Fl.* 270.—Afr. trop.

ZANTHOXYLUM :—
rufescens, *Desf. Cat. Hort. Par.* ed. III. 407.—Hab. ?
rugosum, *A. St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 140.—Bras.
Ruizianum, Klotzsch, ex Engl. in Mart. Fl. Bras. xii. II. 170 = obscurum.
Rumphianum, Cham. in Linnaea, v. (1830) 58 = Evodia latifolia.
sapindifolium, *Wall. Cat.* n. 1215.—Burma.
sapindoides, DC. Prod. i. 728 = emarginatum.
scandens, *Blume, Bijdr.* 249.—Java.
schinifolium, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 137.—China; Japon.
Selloi, *Spreng. Syst.* i. 947.—Bras.
senegalense, *DC. Prod.* i. 726.—Afr. trop.
sepiarium, Wight, Illustr. Ind. Bot. i. 169 = ovalifolium.
Serra, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 440 = Saurauja napaulensis.
serrulatum, *Blume, Bijdr.* 249.—Java.
simplicifolium, Vahl, Eclog. Am. iii. 45 = Z. Ochroxylum.
simulans, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 208 = Bungei.
sorbifolium, A. St. Hil. Fl. Bras. Mer. i. 75 = rhoifolium.
Spinifex, *DC. Prod.* i. 728.—Ind. occ.; Venezuela.
spinosum, *Sw. Fl. Ind. Occ.* i. 574.—Ind. occ.
spondiaefolium, *Wall. Cat.* n. 1217.—Burma.
Sprucei, *Engl. in Mart. Fl. Bras.* xii. II. 167.—Peruv.
stelligerum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 442.—Bras.
stenopterum, *Wright, in Sauv. Fl. Cub.* 19.—Cuba.
stipitatum, *Engl. in Mart. Fl. Bras.* xii. II. 161.—Bolivia.
subserratum, *Engl. l. c.* 160.—Bras.
Sumach, *Griseb. Fl. Brit. W. Ind.* 138, in obs.—Ins. Guadeloup.
Sumach, Macfad. Fl. Jamaic. i. 195 = Brunellia comocladifolia.
syncarpum, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 279.—N. Granat.
taediosum, *A. Rich. Ess. Fl. Cub.* 330.—Cuba.
tenuifolium, *Engl. in Mart. Fl. Bras.* xii. II. 160.—Bras.
terebinthoides, *Bertol. f. in Mem. Acad. Sc. Bologn.* v. (1854) 470.—Afr. trop.
ternatum, *Sw. Fl. Ind. Occ.* i. 570.—Ind. occ.
tetraspermum, *Wight & Arn. Prod.* 148.—Ind. or.
texanum, *Buckl. in Bull. Torrey Bot. Club*, x. (1883) 90.—Texas.
Thunbergii, *DC. Prod.* i. 726.—Afr. austr.
tidorense, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 246.—Ins. Molucc.
Timbor, Wall. Cat. n. 7116 = acanthopodium.
timoriense, *Span. in Linnaea*, xv. (1841) 185.—Ins. Timor.
Tingoassuiba, *A. St. Hil. Fl. Bras. Mer.* i. 78.—Bras.
tomentellum, *Hook. f. Fl. Brit. Ind.* i. 493.—Reg. Himal.
torvum, *F. Muell. Fragm.* vii. 140.—Austral.
tragodes, *DC. Prod.* i. 725.—Ins. S. Doming.
tricarpum, Hook. Fl. Bor. Am. i. 118 = americanum.
tricarpum, Michx. Fl. Bor. Am. ii. 235 = Z. Clava-Herculis.
triflorum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 597 = Melicope indica ?
trifoliatum, Linn. Sp. Pl. 270 = Acanthopanax aculeatum.
triphyllum, G. Don, Gen. Syst. i. 804 = Evodia triphylla.
triphyllum, Wight, Ic. t. 204 = Evodia Roxburghiana.
triplinerve, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 597.—Ins. Philipp.
tuberculatum, *Engl. in Mart. Fl. Bras.* xii. II. 166.—Bras.
undulatum, Wall. Cat. n. 1208 = Toddalia lanceolata.
varians, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 214.—Ins. Fiji.
velutinum, Benth. Pl. Hartw. 167 = rigidum.
violaceum, Wall. Cat. n. 1213 = oxyphyllum.
Wallichianum, Steud. Nom. ed. II. ii. 797 = ovalifolium.
zeylanicum, DC. Prod. i. 728 = Evodia Roxburghiana.

ZANTHYRSIS, Rafin. New Fl. Am. iii. 84 (1836) = **Sophora**, Linn. (Leguminos.).
paniculata, Rafin. l. c. = S. tomentosa.

ZANTORRHIZA, Steud. Nom. ed. II. ii. 796 (1841) = **Zanthorhiza**, L'Herit. (Ranunc.).

ZAPAMIA, Steud. Nom. ed. II. ii. 796 (1841), sphalm. = seq.

ZAPANIA, Lam. Illustr. i. 58. t. 17. n. 42 (1791) = **Lippia**, Houst. (Verben.).
angustifolia, Poir. Encyc. viii. 841 = Stachytarpheta angustifolia.
arabica, Poir. l. c. 844 = Priva leptostachya.
aristata, Mirb. Hist. Nat. Pl. ed. II. xv. 240 = Stachytarpheta orubica.
bracteosa, Poir. Encyc. viii. 843 = Verbena bracteata.
cajenensis, Mirb. Hist. Nat. Pl. ed. II. xv. 241 = Stachytarpheta cajenensis.
canescens, Gibert, Enum. Pl. Montev. 44 = L. canescens.
citriodora, Lam. Illustr. i. 59 = L. citriodora.
corymbosa, Poir. Encyc. viii. 844 = Verbena corymbosa.
cuneifolia, Torr. in Ann. Lyc. N. York, ii. (1828) 234 = L. cuneifolia.
curassavica, Lam. Illustr. i. 59 = Tamonea spinosa.
dichotoma, Mirb. Hist. Nat. Pl. ed. II. xv. 240 = Stachytarpheta dichotoma.
Echinus, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 71. t. 5 = Hyptis Sideritis.
geminata, Gibert, Enum. Pl. Montev. 44 = L. geminata.
globiflora, Juss. in Ann. Mus. Par. vii. (1806) 72 = L. asperifolia.
hispida, Zuccagni, in Roem. Collect. 118 = Priva hispida.
indica, Lam. Illustr. i. 59 = Stachytarpheta indica.
jamaicensis, Lam. l. c. = Stachytarpheta dichotoma.
javanica, Lam. l. c. = L. javanica.
lanceolata, Juss. in Ann. Mus. Par. vii. (1806) 72 = L. lanceolata.
lantanoides, Lam. Illustr. i. 58 = L. asperifolia.
lappulacea, Lam. l. c. 59 = Priva echinata.
marginata, Mirb. Hist. Nat. Pl. ed. II. xv. 240 = Stachytarpheta indica.
mexicana, Lam. Illustr. i. 59 = Priva hispida.
mutabilis, Lam. l. c. = Stachytarpheta mutabilis.
nodiflora, Lam. l. c. = L. nodiflora.
nudiflora, Spreng. Pugill. ii. 70, in obs., sphalm. = prace.
odorata, Pers. Syn. ii. 140 = L. asperifolia.
odoratissima, Scop. Delic. Insub. i. 34 = L. asperifolia.
orubica, Mirb. Hist. Nat. Pl. ed. II. xv. 241 = Stachytarpheta orubica.
prismatica, Lam. Illustr. i. 59 = Bouchea Ehrenbergii.
reclinata, Lam. l. c. 58 = L. stoechadifolia.
repens, Bertol. Rar. Ital. Pl. Dec. iii. 27 = L. nodiflora.
reptans, Spreng. Pugill. ii. 70, in obs. = L. reptans.
scaberrima, Juss. ex Pers. Syn. ii. 140.—Hab. ?
squamosa, Mirb. Hist. Nat. Pl. ed. II. xv. 241 = Elytraria tridentata.
staechadifolia, Juss. in Ann. Mus. Par. vii. (1806) 72 = L. stoechadifolia.
strigosa, Mirb. Hist. Nat. Pl. ed. II. xv. 241 = Stachytarpheta strigosa.
suberosa, Spreng. Pugill. ii. 69 = L. reptans.
urticifolia, Poir. Encyc. Suppl. v. 520 = Stachytarpheta dichotoma.
virgata, Poir. Encyc. viii. 845 = L. urticoides.

ZAPPANIA, Zuccagni, in Roem. Collect. 117 (1809), sphalm. = praec.

ZARA, Benth. & Hook. f. Gen. iii. 964 (1883), err. typ. = ZALA, Lour. = **Pistia**, Linn.

ZARABELLIA, Cass. in Dict. Sc. Nat. lix. 240 (1829) = **Melampodium**, Linn. (Compos.).
rhomboidea, Cass. l. c. = M. paludosum.

ZARABELLIA, Neck. Elem. i. 10 (1790) = **Berkheya**, Ehrh. (Compos.).

ZARCOA, Llanos, in Bot. Zeit. xv. (1857) 423 = **Bridelia**, Willd. (Euphorb.).
phillipica, Llanos, l. c. = Glochidion philippense.

ZATARENDIA, Rafin. Fl. Tellur. iii. 86 (1836) = **Origanum**, Tourn. (Labiat.).
egyptiaca, Rafin. l. c. = O. Maru.

ZATARIA, Boiss. Diagn. Ser. I. v. 18 (1844). *LABIATAE*, Benth. & Hook. f. ii. 1186.
bracteata, Boiss. l. c. Ser. II. iv. 12 = multiflora.
humilis, Benth. in DC. Prod. xii. 183 = Nepeta bracteata.
multiflora, *Boiss. Diagn.* Ser. I. v. 18.—Persia.

ZAUSCHERIA, Steud. Nom. ed. II. ii. 797 (1841), sphalm. = seq.

ZAUSCHNERIA, Presl, Rel. Haenk. ii. 28. t. 52 (1831). *ONAGRARIEAE*, Benth. & Hook. f. i. 788.
californica, *Presl, l. c.*—Calif.
mexicana, Presl, l. c. 29 = californica.

ZAZINTHA, Hall. in Rupp. Fl. Jen. 209 (1745), sphalm. = **Zacintha**, Tourn. (Compos.).

ZEA, Linn. Gen. ed. I. 279 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1114.
MAYS, Tourn. ex Gaertn. Fruct. i. 6. t. 1 (1788).
THALYSIA, Linn. Syst. ed. I (1735).
alba, Mill. Gard. Dict. ed. VIII. n. 2 = Z. Mays.
altissima, Gmel. ex Steud. Nom. ed. I. 898 = Z. Mays.
americana, Mill. Gard. Dict. ed. VIII. n. 1 = Z. Mays.
amylacea, Sturtev.—Cf. Gard. Chron. (1883) II. 443 = Z. Mays.
Caragua, Steud. Nom. ed. II. ii. 797, sphalm. = Z. Curagua.
cryptosperma, Bonaf. Hist. Nat. Maïs, 30. t. 5 bis = Z. Mays.
Curagua, *Molina, Sagg. Chil.* 129, 354.—Chili.
erythrolepis, *Bonaf. Hist. Nat. Maïs*, 30. t. 5.—Am. bor. (cult.).
everta, Sturtev.—Cf. Gard. Chron. (1883) II. 443 = Z. Mays.
hirta, *Bonaf. in Ann. Sc. Nat.* Sér. I. xvii. (1829) 158.—Calif.
indentata, Sturtev.—Cf. Gard. Chron. (1883) II. 443 = Z. Mays.
indurata, Sturtev. l. c. = Z. Mays.
japonica, *Van Houtte, Fl. des Serres*, xvi. (1865–67) 121. t. 1673–74.—Japon.
macrosperma, *Klotzsch, in Bot. Zeit.* ix. (1851) 718.—Peruv.
Maïs, Vell. Fl. Flum. x. t. 3 = seq.
Mays, *Linn. Sp. Pl.* 971.—Paraguay.
minor, Gmel. ex Steud. Nom. ed. I. 898 = Z. Mays.
mucronata, Poit. ex Vilm. in Rev. Hortic. Sér. II. i. (1843) 246, in nota – rostrata.
praecox, Steud. Nom. ed. I. 898 = Z. Mays.
rostrata, *Bonaf. in Ann. Soc. Agric. Lyon*, v. (1842) 197; *et ex Vilm. in Rev. Hortic.* Sér. II. i. (1843) 245.—Peruv.
saccharata, Sturtev.—Cf. Gard. Chron. (1883) II. 443 = Z. Mays.
segetalis, Salisb. Prod. 28 = Z. Mays.
vaginata, Sturtev.—Cf. Gard. Chron. (1883) II. 443 = Z. Mays.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 3 = Z. Mays.

ZEBRINA, Schnizl. in Bot. Zeit. vii. (1849) 870. *COMMELINACEAE*, Benth. & Hook. f. iii. 856.
leiandra, *C. B. Clarke, in DC. Monog. Phan.* iii. 318.—Mexic.
pendula, *Schnizl. in Bot. Zeit.* vii. (1849) 870.—Mexic.

ZEDOARIA, Rafin. Fl. Tellur. iv. 53 (1836) = **Amomum**, Linn. (Scitam.).

ZEDUBA, Ham. ex Meissn. Gen. Comm. 280 (1842) = **Calanthe**, R. Br. (Orchid.).

ZEHNERIA, Endl. Prod. Fl. Norfolk. 69 (1833) = **Melothria**, Linn. (Cucurb.).
affinis, Endl. ex Walp. Rep. ii. 196 = M. punctata.
angolensis, Hook. f. in Oliver, Fl. Trop. Afr. ii. 559 = M. angolensis.
Baueriana, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 624 = M. mucronata.
Baueriana, Endl. Prod. Fl. Norfolk. 69 = M. Baueriana.
cerasiformis, Stocks, in Hook. Kew Journ. iv. (1852) 149 = Blastania fimbristipula.
connivens, Miq. Fl. Ind. Bat. i. I. 656 = M. heterophylla.

ZEHNERIA :—
cordata, Sond. in Harv. & Sond. Fl. Cap. ii. 486 = Melothria cordata.
Cunninghamii, F. Muell. in Hook. Kew Journ. viii. (1856) 50 = M. Cunninghamii.
cuspidata, Endl. ex Walp. Rep. ii. 196 = M. punctata.
debilis, Sond. in Harv. & Sond. Fl. Cap. ii. 488 = Trochomeria debilis.
deltoidea, Hook. f. in Oliver, Fl. Trop. Afr. ii. 560 = M. Mannii.
deltoidea, Miq. Pl. Jungh. 397 = M. Rauwenhoffii.
dilatata, Endl. ex Walp. Rep. ii. 196 = M. cordata.
Ecklonis, Endl. ex Walp. l. c. = M. punctata.
erythrocarpa, F. Muell. in Hook. Kew Journ. viii. (1856) 51 = Bryonopsis laciniosa.
exasperata, Miq. Fl. Ind. Bat. i. 1. 655 = M. punctata.
filiformis, Miq. l. c. = M. heterophylla.
Garcini, Sond. in Harv. & Sond. Fl. Cap. ii. 487 = Trochomeria Garcini.
Garcini, Stocks, in Hook. Kew Journ. iv. (1852) 149 = Blastania Garcini.
hastata, Miq. Pl. Jungh. 398; Fl. Ind. Bat. i. 1. 656 = M. heterophylla.
hederacea, Sond. in Harv. & Sond. Fl. Cap. ii. 487 = M. hederacea.
Hookeriana, Arn. in Hook. Journ. Bot. iii. (1841) 275 = M. perpusilla.
Hookeriana, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 80 = M. japonica.
longepedunculata, Hochst. ex Cogn. in DC. Monog. Phan. iii. 612 = M. longepedunculata.
longepedunculata, A. Rich. Tent. Fl. Abyss. i. 287 = M. scrobiculata.
lucida, Hook. f. in Oliver, Fl. Trop. Afr. ii. 559 = M. lucida.
macrocarpa, Sond. in Harv. & Sond. Fl. Cap. ii. 488 = Trochomeria macrocarpa.
maysurensis, Arn. in Hook. Journ. Bot. iii. (1841) 275 = M. mucronata.
membranacea, Endl. ex Walp. Rep. ii. 196 = M. punctata.
micrantha, Hook. f. in Oliver, Fl. Trop. Afr. ii. 560 = M. minutiflora.
micrantha, F. Muell. Fragm. i. 182 = M. Muelleri.
microsperma, Hook. f. in Oliver, Fl. Trop. Afr. ii. 559 = M. microsperma.
mucronata, Miq. Fl. Ind. Bat. i. 1. 656 = M. mucronata.
mysorensis, Wight, in Ann. & Mag. Nat. Hist. Ser. I. viii. (1842) 268 = M. mucronata.
obtusiloba, Sond. in Harv. & Sond. Fl. Cap. ii. 487 = M. obtusiloba.
pectinata, Sond. l. c. = Trochomeria pectinata.
racemosa, Hook. f. in Oliver, Fl. Trop. Afr. ii. 561 = M. racemosa.
scabra, Sond. in Harv. & Sond. Fl. Cap. ii. 486 = M. punctata.
scrobiculata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 287 = M. scrobiculata.
suavis, Endl. ex Walp. Rep. ii. 196 = M. punctata.
umbellata, Thw. Enum. Pl. Zeyl. 125 = M. heterophylla.
velutina, Arn. in Hook. Journ. Bot. iii. (1841) 275; Endl. ex Walp. Rep. ii. 196 = M. velutina.
velutina, A. Rich. Tent. Fl. Abyss. i. 287 = M. tomentosa.
Wyleyana, Sond. in Harv. & Sond. Fl. Cap. ii. 488 = Trochomeria Wyleyana.

ZELEA, Hort. ex Tenore, Atti 3ª Reun. Sc. Ital. (1841) 504 = **Carapa**, Aubl. (Meliac.).
splendens, Hort. ex Tenore, l. c. = C. procera.

ZELKOVA, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 356. *URTICACEAE*, Benth. & Hook. f. iii. 353.
ABELICEA, Reichb. Consp. 84 (1828); Baill. Hist. Pl. vi. 185 (1875).
HEMIPTELEA, Planch. in Compt. Rend. Acad. Par. lxxiv. (1872) 131.
Abelicea, Boiss. Fl. Orient. iv. 1159 = cretica.
acuminata, *Planch. in Compt. Rend. Acad. Par.* lxxiv. (1872) 1496; *et in DC. Prod.* xvii. 166.—Japon.
crenata, *Spach, in Ann. Sc. Nat.* Sér. II. xv. (1841) 358.—Reg. Cauc.
cretica, *Spach, Hist. Vég. Phan.* xi. 121.—Ins. Creta.
Keaki, Maxim. in Bull. Acad. Pétersb. xviii. (1873) 288 = acuminata.
stipulacea, *Franch. & Sav. Enum. Pl. Jap.* ii. 489 (err. typ. 984).—Japon.

ZELMIRA, Rafin. Fl. Tellur. iv. 50 (1836) = **Calathea**, G. F. W. Mey. (Scitam.).
violacea, Rafin. l. c. = C. violacea.

ZELONOPS, Rafin. Fl. Tellur. ii. 102 (1836) = **Phoenix**, Linn. (Palm.).
pusilla, Rafin. l. c. = P. farinifera.

ZENKERIA, Arn. in Mag. Zool. & Bot. ii. (1838) 548 = **Apuleia**, Mart. (Legumin.).
dalbergioides, Arn. l. c. = A. praecox.
Lundii, F. Didr. in Kjoeb. Viddensk. Meddel. (1853) 88 = A. praecox.

ZENKERIA, Reichb. Nom. 236 (1841) = **Parmentiera**, DC. (Bignon.).

ZENKERIA, Trin. in Linnaea, xi. (1837) 150. t. 3. *GRAMINEAE*, Benth. & Hook. f. iii. 1155.
AMPHIDONAX, Nees, ex Steud. Syn. Pl. Gram. 196 (1855).
elegans, *Trin. in Linnaea*, xi. (1837) 150. t. 3.—Ind. or.
obtusiflora, *Benth. in Journ. Linn. Soc.* xix. (1881) 93.—Zeylan.

ZENOBIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 158. *ERICACEAE*, Benth. & Hook. f. ii. 587.
dealbata, D. Don, ex Steud. Nom. ed. II. i. 89, in syn.; ii. 797 = speciosa.
floribunda, DC. Prod. vii. 598 = Pieris floribunda.
racemosa, DC. l. c. = Leucothoë racemosa.
speciosa, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 158.—Am. bor.

ZENOPOGON, Link, Handb. ii. 481 (1831) = **Anthyllis**, Riv. (Legumin.).
cytisoides, Link, l. c. 481, 146 = A. cytisoides.
Genistae, Link, ll. cc. = A. Genistae.
Herrmanniae, Link, ll. cc. = A. Hermanniae.

ZEOCRITON, Beauv. Agrost. 114. t. 21. f. 2 (1812) = **Hordeum**, Tourn. (Gramin.).
commune, Beauv. l. c. 182 = H. Zeocriton.
complanatum, Beauv. l. c. 115 = H. complanatum.
distichum, Beauv. l. c. = H. distichon.
Hystrix, Beauv. l. c. = H. maritimum.
maritimum, Beauv. l. c. = H. maritimum.
murinum, Beauv. l. c. = H. murinum.
nodosum, Beauv. l. c. = H. pratense.
pratense, Beauv. l. c. = H. secalinum.
rigidum, Beauv. l. c. = H. maritimum.
secalinum, Beauv. l. c. = H. capense.
strictum, Beauv. l. c. = H. bulbosum.
vulgare, S. F. Gray, Nat. Arr. Brit. Pl. ii. 90 = H. Zeocriton.

ZEPHYRA, D. Don, in Edinb. N. Phil. Journ. (Oct. 1832) 236. *HAEMODORACEAE*, Benth. & Hook. f. iii. 680.
DICOLUS, Phil. in Anal. Univ. Chil. (1873) 550; (Desc. Pl. Nuev. ii. 74).
amoena, Miers, in Trans. Linn. Soc. xxiv. (1864) 503 = elegans.
elegans, *D. Don, in Edinb. N. Phil. Journ.* (Oct. 1832) 236.—Peruv.; Chili.

ZEPHYRANTHES, Herb. App. [Bot. Reg.] 36 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 723.
ARGYROPSIS, M. Roem. Syn. Ensat. 125 (1847).
ARVIELA, Salisb. Gen. Pl. Fragm. 135 (1866).
HABRANTHUS, Herb. in Bot. Mag. t. 2464 (1824); et in Bot. Reg. t. 1345 (1830), partim.
MESOCHLOA, Rafin. Fl. Tellur. iv. 10 (1836).
PLECTRONEMA, Rafin. l. c. (1836).
POGONEMA, Rafin. l. c. (1836).
PYROLIRION, Herb. App. [Bot. Reg.] 37 (1821).
Ackermannia, M. Roem. Syn. Ensat. 124 = verecunda.
acuminata, Herb. in Bot. Reg. sub t. 1345 = mesochloa.
Akermanniana, Herb. ex G. Don, in Sweet, Hort. Brit. ed. III. 671, nomen (Quid?).—Mexic.
Andersoniana, *Benth. & Hook. f. Gen.* iii. 724.—Reg. Argent.
Andersonii, Nichols. Dict. Gard. ii. 107 = praec.
Atamasco, *Herb. App.* [*Bot. Reg.*] 36.—Amer.
aurea, *Benth. & Hook. f. Gen.* iii. 724.—Peruv.

ZEPHYRANTHES:—
aurea, S. Wats. in Proc. Am. Acad. xviii. (1883) 161 = longifolia.
bifolia, M. Roem. Syn. Ensat. 125 = rosea.
candida, *Herb. in Bot. Mag.* t. 2607.—Reg. Argent.
carinata, *Herb. l. c.* t. 2594.—Am. centr.; Jamaic.
carnea, D. Dietr. Syn. Pl. ii. 1175 = rosea.
chloroleuca, *Herb. App.* [*Bot. Reg.*] 36.—Afr. austr.?
chlorosolen, D. Dietr. Syn. Pl. ii. 1176 = Cooperia Drummondii.
citrina, *Baker, in Bot. Mag.* t. 6605.—Guian.
Commersoniana, *Herb. Amaryll.* 174. t. 29. f. 3.—Reg. Argent.
concolor, *Benth. & Hook. f. Gen.* iii. 724, in nota.—Mexic.
depauperata, *Herb. Amaryll.* 171.—Chili.
Drummondi, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 328 = Cooperia pedunculata.
flava, *Ind. Kew.* i. 102 (perperam citata, post 1885 publici juris facta).
flavescens, Herb. in Lindl. in Bot. Reg. sub t. 1345 = mesochloa.
gracilifolia, *Nichols. Dict. Gard.* ii. 107.—Reg. Argent.
gracilis, *Herb. Amaryll.* 172. t. 29. f. 1.—Peruv.
Grahamiana, Herb. l. c. 175. t. 29. f. 2 = verecunda.
grandiflora, Lindl. Bot. Reg. t. 902 = carinata.
Herbertiana, D. Dietr. Syn. Pl. ii. 1176 = Cooperia Drummondii.
lilacina, Liebm. Ind. Sem. Hort. Haun. (1844).—Cf. Linnaea, xviii. (1844) 509 = verecunda?
Lindleyana, *Herb. Amaryll.* 174. t. 35. f. 5.—Mexic.
longifolia, *Hemsl. Diagn. Pl. Nov.* 55.—Mexic.
lutea, Herb. App. [Bot. Reg.] 36 = Hippeastrum chilense.
macrosiphon, *Baker, in Gard. Chron.* (1881) ii. 70.—Mexic.
mesochloa, *Herb. in Bot. Reg.* sub t. 1345; et t. 1361.—Reg. Argent.
minima, *Herb. Amaryll.* 172. t. 24. f. 3.—Bras.
minuta, D. Dietr. Syn. Pl. ii. 1176 = Amaryllis minuta.
nervosa, D. Dietr. l. c. = verecunda.
nervosa, Herb. Amaryll. 172 = tubispatha.
nervosa, Mart. & Gal. in Bull. Acad. Brux. x. ii. (1843) 114 = Lindleyana.
nivea, D. Dietr. Syn. Pl. ii. 1176 = candida.
pallida, M. Roem. Syn. Ensat. 124 = verecunda.
pudica, D. Dietr. Syn. Pl. ii. 1176 = Amaryllis Belladonna.
pumila, *M. Roem. Syn. Ensat.* 122.—Hab.?
purpurea, Phil. in Linnaea, xxix. (1857–58) 65 = Hippeastrum roseum.
pusilla, D. Dietr. Syn. Pl. ii. 1176 = Haylockia pusilla.
rosea, *Lindl. in Bot. Reg.* t. 821.—Cuba.
sessilis, Herb. Amaryll. 175 = verecunda.
spofforthiana ×, Herb. in Bot. Reg. t. 1746.
striata, Herb. in Bot. Mag. t. 2593 = verecunda.
texana, *Herb. l. c.* sub t. 3482.—Texas.
Treatiae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 300.—Florida.
tubispatha, *Herb. App.* [*Bot. Reg.*] 36.—Am. trop.
verecunda, *Herb. in Bot. Mag.* t. 2583.—Mexic.
versicolor, *Nichols. Dict. Gard.* ii. 107.—Uruguay.

ZERDANA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 84. *CRUCIFERAE*, Benth. & Hook. f. i. 80.
anchonioides, *Boiss. l. c.*—Persia.

ZERNA, Panz. in Denkschr. Akad. Muench. 1813 (1814) 296 = **Bromus**, Dill. et **Festuca**, Tourn. (Gramin.).
angustifolia, Bess. in Mém. Soc. Nat. Mosc. vi. (1823) 203 = B. erectus.
aspera, Panz. in Denkschr. Akad. Muench. 1813 (1814) 297 = B. asper.
bromoides, Panz. l. c. = Festuca bromoides.
distachyos, Panz. l. c. = Brachypodium distachyum.
erecta, Panz. l. c. = B. erectus.
gigantea, Panz. l. c. = B. giganteus.
ligustica, Panz. l. c. = B. rubens, scoparius, Festuca clavata.
macrostachys, Panz. l. c. = B. macrostachys.
madritensis, Panz. l. c. = B. madritensis.
Myuros, Panz. l. c. = Festuca Myuros.
sterilis, Panz. l. c. = B. sterilis.
tectorum, Panz. l. c. = B. tectorum.

ZERUMBET, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 329 = **Zingiber**, Adans. (Scitam.).
Zingiber, Lestib. l. c. = Zingiber Zerumbet.

ZERUMBET, Wendl. Sert. Hannov. fasc. iv. 3. t. 19 (1798) = **Alpinia**, Linn. (Scitam.).
speciosum, Wendl. l. c. = A. nutans.

ZETOCAPNIA, Link & Otto, Ic. Pl. Rar. i. t. 18 (1828), sphalm. = COETOCAPNIA, Link & Otto = **Bravoa**, Lex. (Amaryll.).
geminiflora, Link & Otto, l. c. = B. geminiflora.

ZEUGITES, P. Br. Hist. Jamaic. 341 (1756); Schreb. Gen. 810 (1791). *GRAMINEAE*, Benth. & Hook. f. iii. 1191.
DESPRETZIA, Kunth, Rév. Gram. ii. 485. t. 157 (1830).
GALEOTTIA, Rupr. ex Gal. in Bull. Acad. Brux. ix. II. (1842) 247.
KROMBHOLZIA, Rupr. in Bull. Acad. Brux. ix. II. (1842) 247.
SENITES, Adans. Fam. ii. 39 (1763).
americana, Rupr. in Bull. Acad. Brux. ix. II. (1842) 247 = mexicana.
americana, *Willd. Sp. Pl.* iv. 204.—Jamaic.
colorata, *Griseb. Fl. Brit. W. Ind.* 536, in obs.—Venezuela.
Galeottiana, *Hemsl. Biol. Centr. Am. Bot.* iii. 577.—Mexic.
Hartwegi, *Fourn. ex Hemsl. l. c.*—Guatemala.
jamaicensis, Raeusch. Nom. ed. III. 270 = americana.
latifolia, *Hemsl. Biol. Centr. Am. Bot.* iii. 577.—Mexic.
mexicana, *Trin. ex Steud. Nom.* ed. II. ii. 798.—Mexic.
Munroana, *Hemsl. Biol. Centr. Am. Bot.* iii. 577.—Guatemala.

ZEUXINE, Lindl. Orchid. Scel. 9 (1826) (Zeuxina); Reichb. Consp. 67 (1828). *ORCHIDEAE*, Benth. & Hook. f. iii. 599.
ADENOSTYLIS, Blume, Bijdr. 414 (1825).
HAPLOCHILUS, Endl. Ench. 113 (1841); Gen. Suppl. ii. 20 (1842).
MONOCHILUS, Wall. ex Lindl. Gen. et Sp. Orch. 486 (1840).
PSYCHECHILUS, Breda, Orch. Kuhl. et Hass. [t. 9] (1827).
STRATEUMA, Rafin. Fl. Tellur. ii. 89 (1836).
TRIPLEURA, Lindl. in Wall. Cat. n. 7391 (1832); et Bot. Reg. sub t. 1618 (1833).
africana, *Reichb. f. in Flora*, l. (1867) 103.—Afr. trop.
bracteata, Wight, Ic. t. 1724, bis. = sulcata.
brevifolia, Wight, Ic. t. 1725 = sulcata.
clandestina, *Blume, Orch. Archip. Ind.* 70.—Java.
emarginata, Lindl. Gen. et Sp. Orch. 485 = sulcata.
flava, *Trimen, Cat. Pl. Ceyl.* 90.—Ind. or.; Malaya.
Godefroyi, *Reichb. f. Otia Bot. Hamb.* 34.—Cambodia.
goodyeroides, *Lindl. Gen. et Sp. Orch.* 486.—Reg. Himal.; Burma.
gracilis, *Blume, Orch. Archip. Ind.* 69.—Java.
integerrima, Lindl. Gen. et Sp. Orch. 486 = sulcata.
longilabris, *Trimen, Cat. Pl. Ceyl.* 90.—Ind. or.
membranacea, Lindl. Gen. et Sp. Orch. 486 = sulcata.
moniliformis, Griff. Notul. iii. 397 = Cheirostylis pusilla.
nervosa, *Trimen, Cat. Pl. Ceyl.* 90.—Reg. Himal.
procumbens, Blume, Orch. Archip. Ind. 68 = sulcata.
purpurascens, *Blume, l. c.* 71.—Sumatra; Borneo.
regia, *Trimen, Cat. Pl. Ceyl.* 90.—Zeylan.
robusta, Wight, Ic. t. 1726 = sulcata.
sulcata, *Lindl. Gen. et Sp. Orch.* 485.—Ind. or.; Malaya; China.
Tripleura, Lindl. in Journ. Linn. Soc. i. (1857) 186 = sulcata.

ZEXMENIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 13 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 373.
LASIANTHAEA, DC. Prod. v. 607 (1836).
LASIANTHUS, Zucc. ex DC. l. c. (1836).
TELESIA, Rafin. Fl. Tellur. ii. 41 (1836).
aurea, *Benth. & Hook. f. Gen.* ii. 371, in nota sub Wedelia.—Mexic.
brevifolia, *A. Gray, Pl. Wright.* i. 112.—Mexic.
caracasana, *Benth. & Hook. f. Gen.* ii. 371, in nota sub Wedelia.—Venezuela.
ceanothifolia, *Sch. Bip. in Seem. Bot. Voy. Herald*, 305.—Mexic.
costaricensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 95.—Am. centr.

ZEXMENIA:—
crocea, *A. Gray, Pl. Wright.* i. 114.—Mexic.
elegans, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 373.—Hab. ?
fasciculata, *Hemsl. Biol. Centr. Am. Bot.* ii. 172.—Mexic.
Ghiesbrechtii, *A. Gray, Pl. Wright.* i. 113.—Mexic.
gnaphalioides, *A. Gray, in Proc. Am. Acad.* xv. (1880) 36.—Mexic.
goyazensis, *Benth. & Hook. f. Gen.* ii. 373.—Bras.
Grayi, Sch. Bip. in Seem. Bot. Voy. Herald, 305 = Verbesina Grayi.
Greggii, *A. Gray, Pl. Wright.* i. 113.—Mexic.
helianthoides, *Benth. & Hook. f. Gen.* ii. 373.—Ecuador.
helianthoides, *A. Gray, Pl. Wright.* i. 113.—Mexic.
hispida, A. Gray, in Proc. Am. Acad. xix. (1883) 10 = Wedelia hispida.
hispidula, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 458 = Blainvillea tampicana.
imbricata, *Sch. Bip. in Seem. Bot. Voy. Herald*, 306.—Mexic.
lantanifolia, *Sch. Bip. l. c.*—Mexic.
Lindenii, *Sch. Bip. l. c.*—Mexic.
longipes, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 95.—Costa Rica.
macrocephala, *Benth. & Hook. f. Gen.* ii. 373; *Hemsl. Biol. Centr. Am. Bot.* ii. 173.—Mexic.
microcephala, *Hemsl. l. c.*—Mexic.
monocephala, *Heynh. Nom.* i. 863; ii. 772; *Sch. Bip. in Seem. Bot. Voy. Herald*, 306.—Mexic.
nicaraguensis, C. Muell. in Walp. Ann. v. 226, sphalm. = costaricensis.
ovata, *Benth. & Hook. f. Gen.* ii. 373.—Mexic.
podocephala, *A. Gray, Syn. Fl. N. Am.* i. II. 286.—Arizona.
reticulata, *Benth. & Hook. f. Gen.* ii. 373.—Cuba.
rudis, *Baker, in Mart. Fl. Bras.* vi. III. 188.—Bras.
salmeoides, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 381 = Otopappus curviflorus.
Salvinii, *Hemsl. Biol. Centr. Am. Bot.* ii. 173.—Guatemala.
scaberrima, *Benth. & Hook. f. Gen.* ii. 373.—Guian.
scandens, *Hemsl. Biol. Centr. Am. Bot.* ii. 174.—Mexic.
Seemannii, *A. Gray, Pl. Wright.* i. 114.—Mexic.
serrata, *La Llave, in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 13.—Mexic.
stenantha, *Hemsl. Biol. Centr. Am. Bot.* ii. 174.—Mexic.
strigosa, *Sch. Bip. in Seem. Bot. Voy. Herald*, 306.—Mexic.
tagetiflora, *D. Don, ex Sweet, Hort. Brit.* ed. II. 309, nomen.—Mexic.
texana, A. Gray, Pl. Wright. i. 112 = Wedelia hispida.
trachylepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 175.—Mexic.
umbellata, Benth. & Hook. f. Gen. ii. 373 = ceanothifolia.
villosa, *Polak, in Linnaea*, xli. (1877) 579.—Costa Rica.
wedelioides, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 328.—Peruv.
zinnioides, *Hemsl. Biol. Centr. Am. Bot.* ii. 175.—Mexic.

ZEYHERA, DC. in Bibl. Univ. Genèv. xvii. (1838) 123; Prod. ix. 211 = **Zeyheria**, Mart. (Bignon.).

ZEYHERA, Less. Syn. Comp. 199 (1832) = ZEYHERIA, Spreng. f. = **Geigeria**, Griessel. (Compos.).

ZEYHERIA, Mart. Nov. Gen. et Sp. ii. 65. t. 159 (1826). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1043.
barbata, Miq. in Flora, xxv. (1842) 430 = Tecoma barbata.
digitata, Miq. l. c. 431 = Tecoma Meyeriana.
fluviatilis, Miq. l. c. 428 = Tabebuia fluviatilis.
montana, *Mart. Nov. Gen. et Sp.* ii. 66. t. 159.—Bras.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 250.—Guian.
tuberculosa, Bur. ex Verlot, in Rev. Hortic. (1868) 154 = montana.
Velloziana, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 201, in obs. = montana.

ZEYHERIA, Spreng. f. Tent. Suppl. Syst. 26 (1828) = **Geigeria**, Griessel. (Compos.).
acaulis, Spreng. f. l. c. = G. africana.

ZICHIA, Steud. Nom. ed. II. ii. 798 (1841), sphalm. = seq.

ZICHYA, Hueg. Bot. Archiv, t. 1 (1837) = **Kennedya**, Vent. (Legumin.).
angustifolia, Lindl. Bot. Reg. (1839) sub t. 52 = K. coccinea.
coccinea, Benth. in Ann. Wien. Mus. ii. (1838) 123 = K. coccinea.
dilatata, Pritz. Ic. Ind. i. 1181 = K. coccinea.
glabrata, Benth. in Ann. Wien. Mus. ii. (1838) 123 = K. glabrata.
glabrata, Meissn. in Lehm. Pl. Preiss. i. 92 = K. coccinea.
heterophylla, G. Don, in Sweet, Hort. Brit. ed. III. 187 = K. coccinea.
inophylla, Benth. in Ann. Wien. Mus. ii. (1838) 123 = K. coccinea.
Latrobeana, Meissn. in Lehm. Pl. Preiss. i. 94 = Glycine Latrobeana.
microphylla, W. Baxt. in Loud. Hort. Brit. Suppl. iii. 659, nomen = K. microphylla ?
Molly, Hueg. Bot. Archiv, t. 1 = K. coccinea.
pannosa, Paxt. Mag. Bot. viii. (1841) 147 = K. coccinea.
sericea, Benth. in Ann. Wien. Mus. ii. (1838) 123 = K. coccinea.
tricolor, Lindl. Bot. Reg. (1839) t. 52 = K. coccinea.
villosa, Lindl. l. c. (1841) Misc. 41 = K. coccinea.

ZIEGERA, Rafin. Sylva Tellur. 94 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
levigata, Rafin. l. c. = M. laevigata.

ZIERIA, Sm. in Trans. Linn. Soc. iv. (1798) 216. *RUTACEAE*, Benth. & Hook. f. i. 291.
arborescens, Sims, Bot. Mag. sub t. 1395 = Smithii.
aspalathoides, *A. Cunn. ex Benth. Fl. Austral.* i. 305.—Austral.
canescens, R. Br. ex Benth. l. c. = pilosa.
commutata, *Schult. Mant.* iii. 324.—Austral.
cytisoides, *Sm. in Rees, Cycl.* xxxix. n. 4.—Austral.
Frankliniae, Millig. ex Kipp. in Proc. Linn. Soc. ii. (1852) 201 = Acradenia Frankliniae.
Fraseri, *Hook. in Mitch. Journ. Trop. Austral.* 339.—Austral.
furfuracea, *R. Br. ex Benth. Fl. Austral.* i. 306.—Austral.
granulata, *C. Moore, ex Benth. l. c.* 307.—Austral.
hirsuta, DC. Prod. i. 723 = pilosa.
involucrata, *R. Br. ex Benth. Fl. Austral.* i. 306.—Austral.
laevigata, *Sm. in Rees, Cycl.* xxxix. n. 2.—Austral.
lanceolata, R. Br. ex Sm. l. c. n. 1 = Smithii.
macrophylla, Bonpl. Desc. Pl. Rar. Malm. 64 = Smithii.
microphylla, *Bonpl. l. c.*—Austral.
microphylla, Sieber, ex Schult. Mant. iii. 324 = commutata.
obcordata, *A. Cunn. in Field, N. S. Wales*, 330.—Austral.
octandra, *Sweet, ex G. Don, Gen. Syst.* i. 794.—Austral.
pauciflora, Sm. in Rees, Cycl. xxxix. n. 3 = pilosa.
pilosa, *Rudge, in Trans. Linn. Soc.* x. (1811) 293. t. 17. f. 2.—Austral.
revoluta, A. Cunn. in Field, N. S. Wales, 330 = laevigata.
Smithii, *Andr. Bot. Rep.* t. 606.—Austral.
trifoliata, Loisel. Herb. Amat. iii. t. 190 = Smithii.
veronicea, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 11.—Austral.

ZIERIDIUM, Baill. Adansonia, x. (1872) 303. *RUTACEAE*.
gracile, *Baill. l. c.* 304.—N. Caled.

ZIERVOGLIA, Neck. Elem. i. 254 (1790) = **Cynanchum**, Linn. (Asclep.).

ZIETENIA, Gled. Syst. 185 (1764) = **Stachys**, Tourn. (Labiat.).
lavandulifolia, Link, Enum. Hort. Berol. ii. 110 = S. lavandulaefolia.
orientalis, Gled. in Mém. Acad. Berl. (1766) 3 = S. lavandulaefolia.

ZIGADENUS, Michx. Fl. Bor. Am. i. 213. t. 22 (1803) = **Zygadenus**, Michx.

ZIGMALOBA, Rafin. Sylva Tellur. 120 (1838) = **Acacia**, Willd. (Legumin.).
sulcata, Rafin. l. c. = A. sulcata.

ZILLA, Forsk. Fl. Aegypt. Arab. 121 (1775). *CRUCIFERAE*, Benth. & Hook. f. i. 98.
Chamaerapistrum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 381 = Physorhynchus brahuicus.
macrocarpa, D. Dietr. Syn. Pl. iii. 678 = myagroides.
macroptera, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 670.—Alger.
microcarpa, Vis. in Bibl. Ital. lxxxiii. (1836) 67 = myagroides.
myagroides, *Forsk. Fl. Aegypt. Arab.* 121.—Aegypt.; Arab.
schouwioides, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 382 = Physorhynchus brahuicus.

ZINGIBER, Adans. Fam. ii. 66 (1763). *SCITAMINEAE*, Benth. & Hook. f. iii. 646.
CASSUMUNAR, Colla, Nov. Scit. Gen. Comm. Taur. 9 (1830).
DIETRICHIA, Giseke, Prael. Linn. 208 (1792).
DYMCZEWICZIA, Horan. Prod. Scitam. 26 (1862).
JAEGERA, Giseke, Prael. Linn. 203 (1792).
LAMPUJANG, Koen. in Retz. Obs. iii. 62 (1783).
THUMUNG, Koen. l. c. (1783).
ZERUMBET, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 329.
amaricans, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 28; *Blume, Enum. Pl. Jav.* 43.—Java.
Anthorrhiza, Hort. Hafn. ex Horan, Prod. Scitam. 27 = Z. Cassumunar.
aquosum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 28; *Blume, Enum. Pl. Jav.* 43.—Java.
aromaticum, Noronha, l. c. = officinale.
barbatum, *Wall. Pl. As. Rar.* i. 46. t. 55.—Burma.
Blancoi, *Hassk. in Flora*, xlvii. (1864) 20.—Ins. Philipp.
capitatum, *Roxb. in As. Res.* xi. (1810) 348.—As. trop.
Cassumunar, *Roxb. l. c.* 347.—As. trop.
cernuum, *Dalz. in Hook. Kew Journ.* iv. (1852) 342.—Ind. or.
chrysanthum, *Rosc. Scitam.* t. 86.—Reg. Himal.
Clarkii, *King, ex Benth. & Hook. f. Gen.* iii. 646.—Reg. Himal.
Cliffordiae, Andr. Bot. Rep. ix. t. 555 = Z. Cassumunar.
coloratum, *N. E. Br. in Gard. Chron.* (1879) II. 166.—Borneo.
confine, *Miq. in Journ. Bot. Néerl.* i. (1861) 94.—China.
corallinum, *Hance, in Journ. Bot.* xviii. (1880) 301.—China.
corynostachyum, Steud. Nom. ed. II. ii. 798 = Amomum corynostachyum.
cylindricum, *Moon, Cat. Pl. Ceyl.* 1, nomen; *Thw. Enum. Pl. Zeyl.* 315.—Zeylan.
dubium, *Afzel. Remed. Guin.* ii. 9.—Afr. trop.
elatum, Roxb. Hort. Beng. 1; Fl. Ind. i. 57 = capitatum.
Ensal, Gaertn. Fruct. i. 35 = Elettaria major.
flavescens, Link, ex A. Dietr. Sp. Pl. i. 54 = chrysanthum?
Galanga, Stokes, Bot. Mat. Med. i. 72 = Alpinia Galanga.
globosum, Stokes, l. c. = Alpinia globosa.
gracile, *Jack, in Malay. Misc.* i. (1820) n. I. 1.—Ins. Penang.
gramineum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 28; *Blume, Enum. Pl. Jav.* 45.—Java.
inflexum, *Blume, l. c.* 43.—Java.
insipidum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = Elettaria paludosa.
integrilabrum, *Hance, in Journ. Bot.* xx. (1882) 80.—China.
ligulatum, *Roxb. in As. Res.* xi. (1810) 348.—Ind. or.
ligulatum, Wall. Cat. n. 6566 = chrysanthum.
luridum, Salisb. in Trans. Hort. Soc. i. (1812) 284 = Z. Cassumunar.
macrostachyum, *Dalz. in Hook. Kew Journ.* iv. (1852) 342.—Ind. or.
marginatum, *Roxb. in As. Res.* xi. (1810) 349.—Reg. Himal.
medium, Stokes, Bot. Mat. Med. i. 72 = Alpinia alba.

ZINGIBER:—
Meleguetta, Gaertn. Fruct. i. 34. t. 12. f. 11 = Amomum angustifolium.
Mijooka, Siebold, in Verh. Batav. Gen. xii. (1830) 18 = Z. Mioga.
minus, Gaertn. l. c. 35 = Amomum Cardamon.
Mioga, *Rosc. in Trans. Linn. Soc.* viii. (1807) 348.—Japon.
missionis, Wall. Cat. n. 6565 = officinale.
montanum, Link, ex A. Dietr. Sp. Pl. i. 52 = Z. Cassumuna.
nigrum, Gaertn. Fruct. i. 35 = Alpinia Allughas.
Nimmonii, *Dalz. in Hook. Kew Journ.* iv. (1852) 341.—Ind. or.
odoriferum, *Blume, Enum. Pl. Jav.* 44.—Java.
officinale, *Rosc. in Trans. Linn. Soc.* viii. (1807) 348.—Trop. cult.
ovoideum, *Blume, Enum. Pl. Jav.* 43.—Java.
panduratum, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 55.—Burma.
pardocheilum, *Wall. ex Voigt, Hort. Suburb. Calc.* 562, nomen.—Burma.
Parishii, *Hook. f. Bot. Mag.* t. 6019.—Burma.
purpureum, Rosc. in Trans. Linn. Soc. viii. (1807) 348 = Z. Cassumunar.
Railleti, *T. Durand, in Bull. Soc. Linn. Par.* i. (1883) 401.—Hab.?
ramosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 28 = Alpinia Pinanga.
roseum, *Rosc. in Trans. Linn. Soc.* viii. (1807) 348.—Ind. or.
rubens, *Roxb. in As. Res.* xi. (1810) 348.—Ind. or.
Sjooka, *Siebold, in Verh. Batav. Gen.* xii. (1830) 18.—Japon.
spectabile, *Griff. Notul.* iii. 413.—Malaya.
spurium, Koen. in Retz. Obs. iii. 60 = Z. Zerumbet.
squarrosum, *Roxb. in As. Res.* xi. (1810) 348.—Burma.
squarrosum, Wight, Ic. t. 2004 = Wightianum.
sylvestre, Gaertn. Fruct. i. 34 = Alpinia alba.
truncatum, Stokes, Bot. Mat. Med. i. 68 = Z. Zerumbet.
uncinatum, Stokes, l. c. 73 = Renealmia racemosa.
Wightianum, *Thw. Enum. Pl. Zeyl.* 315.—Ind. or.
xantorrhizon, Hort. Hafn. ex Steud. Nom. ed. II. ii. 799 = Z. Cassumunar.
Zerumbet, *Rosc. ex Sm. Exot. Bot.* ii. 105. t. 112; *et in Trans. Linn. Soc.* viii. (1807) 348.—As. trop.

ZINNIA, Linn. Syst. ed. X. 1221 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 357.
CRASSINA, Scepin, Sched. Acid. Veget. 42 (1758).
DIPLOTHRIX, DC. Prod. v. 611 (1836).
LEJICA, DC. Prod. v. 534, in syn. (1836).
LEPIA, Hill, Veg. Syst. ii. 97 (1761).
SANVITALIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 357 (1873).
acerosa, *A. Gray, Pl. Wright.* i. 105, 106.—Texas; Mexic.
ambigua, Hort. Berol. ex Salm-Dyck, Hort. Dyck. (1834) 293 (Quid?).—Hab.?
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 251.—Mexic.
anomala, *A. Gray, Pl. Wright.* i. 106.—Texas.
aurea, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 971 = Haageana?
bicolor, *Hemsl. Biol. Centr. Am. Bot.* ii. 153.—Mexic.
bicuspis, [Moç. & Sesse,] ex DC. Prod. v. 535 = Tragoceros zinnioides.
bidens, Retz. Obs. v. 28 = Glossocardia linearifolia.
Darwinii, Haage & Schmidt.—Cf. Gard. Chron. (1875) II. 782 = Glossogyne pinnatifida.
discolor, Hort. Hal. ex Heynh. Nom. i. 864, nomen (Quid?).—Hab.?
elegans, *Jacq. Coll.* iii. 152.—Mexic.
florida, Salisb. Prod. 205 = pauciflora.
floridana, Rafin. New Fl. Am. iv. 70 (Quid?).—Am. bor.
fulva, Schrank, in Denkschr. Bot. Ges. Regensb. i. (1815) 146 = pauciflora.
Ghiesbreghtii, *Verlot, in Rev. Hortic.* (1862) 368.—Mexic.
grandiflora, Hort. ex DC. Prod. v. 536 = hybrida.
grandiflora, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 348.—Am. bor. occ.
Haageana, *Regel, Gartenfl.* x. (1861) 355.—Am. trop.
hybrida, Hort. Par. ex Poir. Encyc. viii. 862 = verticillata.

ZINNIA:—
hybrida, *Roem. & Usteri, Mag. Bot.* i. 1tes St. (1787) 49; *Sims, Bot. Mag.* t. 2123.—Am. austr.
intermedia, Engelm. in Wisliz. Tour N. Mexic. 107 = pauciflora.
juniperifolia, *A. Gray, Pl. Wright.* i. 105.—Mexic.
leptopoda, DC. Prod. v. 535 = pauciflora.
linearifolia, Moench, Meth. Suppl. 261 = pauciflora.
linearis, *Benth. Pl. Hartw.* 17.—Mexic.
lutea, Gaertn. Fruct. ii. 459. t. 172 = pauciflora.
maritima, *H. B. & K. Nov. Gen. et Sp.* iv. 251.—Mexic.
mendocina, *Phil. Sert. Mendoc. Alt.* 27.—Chili.
mexicana, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 971 = Haageana?
multiflora, Linn. Sp. Pl. ed. II. 1269 = pauciflora.
pauciflora, *Linn. l. c.*—Am. bor. et austr.
peruviana, Linn. Syst. ed. X. 1221 = pauciflora.
pumila, *A. Gray, in Mem. Am. Acad.* N. S. iv. (1849) 81.—Am. bor. occ.
revoluta, Cav. Ic. iii. 26 = pauciflora.
Roezlii, *Hort.—Cf. Gard. Chron.* (1872) 1392.—Mexic.
tenuiflora, Jacq. Ic. Pl. Rar. iii. 15. t. 590 = pauciflora.
verticillata, *Andr. Bot. Rep.* iii. t. 189.—Mexic.
verticillata, Cav. Desc. 209 = pauciflora.
violacea, Cav. Ic. i. 57 = elegans.

ZINOWIEWIA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 275. *CELASTRINEAE*, Benth. & Hook. f. i. 364.
integerrima, *Turcz. l. c.* 276.—Mexic.

ZIPANIA, Pers. Syn. ii. 140 (1807), sphalm. = ZAPANIA, Scop. = **Lippia**, Linn. (Verben.).

ZIPPELIA, Blume, in Schult. f. Syst. vii. 1614 (1830). *PIPERACEAE*, Benth. & Hook. f. iii. 128.
begoniaefolia, *Blume, l. c.* 1651.—Java.
lappacea, Benn. Pl. Jav. Rar. 76 = begoniaefolia.

ZIPPELIA, Reichb. ex Endl. Gen. Suppl. ii. 6 (1842) = **Brugmansia**, Blume (Cytinac.).

ZIZANIA, Gronov. ex Linn. Gen. ed. II. 455 (1742). *GRAMINEAE*, Benth. & Hook. f. iii. 1115.
ELYMUS, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 210.
FARTIS, Adans. Fam. ii. 37 (1763).
HYDROPYRUM, Link, Hort. Berol. i. 252 (1827).
MELINUM, Link, Handb. i. 96 (1829).
ZIZANIOPSIS, Doell & Aschers. in Mart. Fl. Bras. ii. 12. t. 3 (1871).
aquatica, *Linn. Sp. Pl.* 991.—Am. bor.
aristata, Kunth, Rév. Gram. i. 8 = Hygroryza aristata.
bonariensis, *Balansa & Poitrass. in Bull. Soc. Hist. Nat. Toul.* xii. (1878) 228.—Reg. Argent.
ciliata, Spreng. Syst. ii. 136 (Quid?) = Leersia hexandra.
clavulosa, Michx. Fl. Bor. Am. i. 75 = aquatica.
dahurica, *Turcz. ex Steud. Syn. Pl. Gram.* 4.—Dahuria.
effusa, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 52 = aquatica.
fluitans, Michx. Fl. Bor. Am. i. 75 = Hydrochloa carolinensis.
latifolia, Turcz. in Bull. Soc. Nat. Mosc. (1838) 105; xxix. (1856) I. 2 = aquatica.
lenticularis, Michx. ex Steud. Nom. ed. I. 898, nomen (Quid?).—Am. bor.
microstachya, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. ii. (1839) 183 = miliacea.
miliacea, *Michx. Fl. Bor. Am.* i. 74.—Am. bor.; Ind. occ.
natans, Bosc, ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. Sc. Nat. ii. (1839) 186 = Hydrochloa carolinensis?
nutans, Steud. Nom. ed. II. ii. 799, sphalm. = praec.
palustris, Linn. Mant. ii. 295 = aquatica.
Retzii, Spreng. Syst. ii. 136 = aristata.
subtilis, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 452 = Coleanthus subtilis.
terrestris, Linn. Sp. Pl. 991 (Quid? al. ord.?).—Ind. or.

ZIZANIOPSIS, Doell & Aschers. in Mart. Fl. Bras. ii. II. 12. t. 3 (1871) = **Zizania**, Gronov. (Gramin.).
microstachya, Doell & Aschers. l. c. 13 = Zizania miliacea.

ZIZIA, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 128 = **Carum**, Rupp. (Umbell.).
aurea, Koch, l. c. 129 = Thaspium aureum.
cordata, Koch, ex DC. Prod. iv. 100 = Thaspium trifoliatum.
integerrima, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 493 = Pimpinella integerrima.
parvifolia, Rafin. New Fl. Am. iv. 31 = Thaspium trifoliatum.
pinnatifida, Buckl. in Am. Journ. Sc. Ser. I. xlv. (1843) 175 = Thaspium Walteri.

ZIZIPHORA, Linn. Hort. Cliff. 305 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1198.
FALDERMANNIA, Trautv. in Bull. Sc. Acad. Pétersb. vi. (1840) 185.
HEDYOSMOS, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) App. 211.
acinoides, *Linn. Sp. Pl.* 22.—Hispan.
acinoides, Pall. Reise, iii. 362 = clinopodioides.
acutifolia, *Montbr. & Auch. ex Benth. in Ann. Sc. Nat.* Sér. II. vi. (1836) 43.—Syria.
alpina, Mill. Gard. Dict. ed. VIII. n. 4 = capitata.
bracteolata, Schult. Mant. i. 179 = Micromeria bracteolata.
Brantii, C. Koch, in Linnaea, xvii. (1843) 294 = clinopodioides.
canescens, Benth. Lab. Gen. et Sp. 321 = clinopodioides.
capitata, Falk, Beitr. ii. 98 = clinopodioides.
capitata, *Linn. Sp. Pl.* 21.—Reg. Mediterr.; Oriens.
cleonioides, *Boiss. Diagn.* Ser. I. v. 14.—As. Min.
clinopodioides, *Lam. Illustr.* i. 63; *Bieb. Fl. Taur. Cauc.* i. 17.—As. Min.; Persia; Sibir.
compacta, Frivald. in Flora, xviii. (1835) 336 = capitata.
Cunila, Desf. Tabl. ed. I. 55 = clinopodioides.
dasyantha, Bieb. Fl. Taur. Cauc. i. 18 = clinopodioides.
fasciculata, C. Koch, ex Boiss. Fl. Orient. iv. 586 = clinopodioides.
glabella, Roem. & Schult. Syst. i. 209 = Calamintha glabella.
glomerata, Frivald. ex Ledeb. Fl. Ross. iii. 370 = capitata.
Gundelsheimeri, C. Koch, in Linnaea, xvii. (1843) 293 = clinopodioides.
hispanica, *Linn. Cent. Pl.* i. 3; *Amoen. Acad.* iv. 262.—Reg. Mediterr.
hispida, Schult. Mant. i. 179 = Hedeoma hispida.
javanica, *Blume, Bijdr.* 822.—Java.
lanceolata, Boiss. Fl. Orient. iv. 588 = taurica.
Mariana, Roem. & Schult. Syst. i. 208 = Cunila Mariana.
media, Link, Enum. Hort. Berol. i. 17 = clinopodioides.
Mussini, Adam, in Weber & Mohr, Beitr. i. (1805) 43 = clinopodioides.
nummularia, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 33 = clinopodioides.
odoratissima, Lodd. ex Schult. Mant. i. 178 = clinopodioides.
persica, Bunge, Lab. Pers. 39 = tenuior.
pulegioides, Desf. Tabl. ed. I. 55, nomen; ed. II. 66 = Hedeoma pulegioides.
pungens, Bunge, ex Ledeb. Fl. Atl. i. 23 = Nepeta pungens.
Puschkini, Adam, in Weber & Mohr, Beitr. i. (1805) 42 = clinopodioides.
serpyllacea, Bieb. Fl. Taur. Cauc. i. 18 = clinopodioides.
serpyllacea, Tenore, Syll. Fl. Neap. 16 = tenuior.
siliquosa, Lour. Fl. Cochinch. 27 = Justicia diffusa?
spicata, Cav. Desc. 20 = hispanica.
spicata, Lag. & Rodr. in Anal. Cienc. Nat. iv. (1801) 259 = tenuior.
taurica, *Bieb. Fl. Taur. Cauc.* i. 414; iii. 21.—As. Min.; Persia.
taurica, C. Koch, in Linnaea, xvii. (1843) 294 = tenuior.
tenuior, Falk, Beitr. ii. 98 = clinopodioides.
tenuior, *Linn. Sp. Pl.* 21.—As. occ.
thymifolia, Salisb. Prod. 72 = acinoides.
thymoides, Roem. & Schult. Syst. i. 209 = Thymus Serpyllum.
verticillata, Steud. Nom. ed. I. 899 = Cunila verticillata.

ZIZIPHUS, Tourn. ex Linn. Syst. ed. I. (1735) = **Zizyphus**, Tourn. (Rhamn.).

ZIZYPHON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70, 137 = **Zizyphus**, Tourn. (Rhamn.).
jujubum, St. Lag. l. c. = Ziziphus Jujuba.

ZIZYPHORA, Dum. Anal. Fam. 22 (1829), sphalm. = **Ziziphora**, Linn. (Labiat.).

ZIZYPHUS, Tourn. ex Linn. Syst. ed. I (1735) (Ziziphus); Adans. Fam. ii. 304 (1763). *RHAMNEAE*, Benth. & Hook. f. i. 375.
DECORIMA, Rafin. Sylva Tellur. 31 (1838).
JUJUBA, Burm. Thes. Zeyl. 131. t. 61 (1737).
MANSANA, J. F. Gmel. Syst. 580 (1791).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 136 = Z. Jujuba.
acuminata, *Benth. Bot. Voy. Sulph.* 78.—Mexic.
acuminata, Royle, Illustr. Bot. Himal. 168, nomen = oxyphylla.
adelensis, *Delile, in Rochet, Sec. Voy. Choa*, 341.—Afr. trop.
africana, Mill. Gard. Dict. ed. VIII. n. 4 = Z. Spina-Christi.
agrestis, *Roem. & Schult. Syst.* v. 341.—China.
albens, Roxb. Hort. Beng. 17; Fl. Ind. i. 607 = Z. Oenoplia.
angulata, *Lam. Encyc.* iii. 320.—Hab.?
apetala, *Hook. f. Fl. Brit. Ind.* i. 635.—Reg. Himal.
Aubletii, D. Dietr. Syn. Pl. i. 810 = Paliurus ramosissimus.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 5.—Persia.
Baclei, DC. Prod. ii. 20 = mucronata.
Baenchia, Buch.-Ham. ex Wall. Cat. n. 4249 (Quid?).—Ind. or.
Bhunder, Royle, Illustr. Bot. Himal. 168, nomen = horrida?
bubalina, Licht. ex Roem. & Schult. Syst. v. 334 = mucronata.
Buraea, Buch.-Ham. ex Wall. Cat. n. 4240 G = rugosa.
Burmanniana, D. Dietr. Syn. Pl. i. 810 = Berchemia Burmanniana.
calophylla, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 366.—Malaya.
capensis, Thunb. ex Poir. Encyc. Suppl. iii. 193 = Scutia Commersonii.
Caracutta, Buch.-Ham. ex Roxb. Hort. Beng. 17; Fl. Ind. i. 612 = Z. Xylopyrus.
celtidifolia, DC. Prod. ii. 20 = Z. Oenoplia.
chinensis, Spreng. Syst. i. 770 = sativa.
Cinnamomum, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 380.—N. Granat.
commutata, Roem. & Schult. Syst. v. 336 = Celtis aculeata.
cotinifolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 87.—Bras.
cuneata, Wall. Cat. n. 7268 = Z. Xylopyrus.
Dalanta, *Blanco, Fl. Filip.* ed. II. 121.—Ins. Philipp.
domingensis, Duham. Arb. ed. Nov. iii. 56 = Ceanothus reclinatus.
elegans, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 366.—Singapore.
elliptica, Roxb. Hort. Beng. 17; Fl. Ind. i. 610 = Z. Xylopyrus.
emarginata, Sw. Fl. Ind. Occ. iii. 1954 = Scutia ferrea.
exserta, *DC. Prod.* ii. 21.—Ins. Philipp.
ferruginea, Heyne, ex Wall. Cat. n. 4246 B, partim = Z. Oenoplia.
flavescens, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 367 = Berchemia flavescens.
flexuosa, Wall. l. c. 365 = sativa.
floribunda, Wall. l. c. 368 = Berchemia floribunda.
funiculosa, *Buch.-Ham. ex Wall. Cat.* n. 4234.—Ind. or.; Malaya.
Gardenii, Steud. Nom. ed. II. ii. 800, sphalm. = seq.
Gardneri, Reiss. in Endl. Nov. Stirp. Dec. 30 = Z. Joazeiro.
glabra, Roxb. Hort. Beng. 17; Fl. Ind. i. 614 = rugosa.
glabrata, *Heyne, ex Roth, Nov. Pl. Sp.* 159.—Ind. or.
Glaziovii, *Warm. in Kjoeb. Vidensk. Meddel.* (1880) 373.—Bras.
globularis, *Wall. Cat.* n. 4248.—Burma.
guatemalensis, *Hemsl. Diagn. Pl. Nov.* 6.—Guatemala.
hamosa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 369 = Sageretia hamosa.
havanensis, H. B. & K. Nov. Gen. et Sp. vii. 57 = Sarcomphalus havanensis.

ZIZYPHUS:—
helvola, *Sond. in Harv. & Sond. Fl. Cap.* i. 476.—Afr. austr.
heterogenea, Poir. Encyc. Suppl. iii. 193 = Z. Spina-Christi.
heterogenea, Russ. ex Wall. Cat. sub n. 4239 = Z. Xylopyrus.
heteroneura, *Griseb. in Bonplandia*, vi. (1858) 3.—Panama.
horrida, *Roth, Nov. Pl. Sp.* 159.—Ind. or.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 643 = rugosa.
Iguanea, Lam. Encyc. iii. 318 = Celtis aculeata.
illucida, C. Muell. in Walp. Ann. vii. 587, sphalm. = lucida.
incurva, *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 614.—Ind. or.
insularis, *C. Sm. in Tuckey, Congo*, 250, nomen.—Ins. Cap. Verd.
javanensis, *Blume, Bijdr.* 1142.—Java.
Joazeiro, *Mart. Reise Bras.* ii. 581.—Bras.
Jujuba, *Lam. Encyc.* iii. 318.—Ind. or.; Malaya.
latifolia, Blanco, Fl. Filip. ed. II. 121 = Z. Oenoplia.
latifolia, Roxb. Hort. Beng. 17; Fl. Ind. i. 607 = rugosa.
lineata, Willd. Sp. Pl. i. 1102 = Berchemia lineata.
Linnaei, *M. Laws. in Hook. f. Fl. Brit. Ind.* i. 635.—Zeylan.
lotoidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = Z. Lotus.
Lotus, Aitch. Cat. Pl. Punjab, 33 = rotundifolia.
Lotus, Blanco, Fl. Filip. ed. II. 121 = Z. Oenoplia.
Lotus, *Lam. Encyc.* iii. 317.—Reg. Mediterr.
Loureiriana, D. Dietr. Syn. Pl. i. 811 = Berchemia Loureiriana.
lucida, Moench, Meth. Suppl. 101 = Rhamnus prinoides.
lucida, *Moon, Cat. Pl. Ceyl.* 17, nomen; *Thw. Enum. Pl. Zeyl.* 74.—Zeylan.
lycioides, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 168.—Am. bor.; Mexic.
mauritiana, Lam. Encyc. iii. 319 = Z. Jujuba.
melanogona, Boj. Hort. Maurit. 69 = Z. Jujuba?
melastomoides, A. Cunn. ex Steud. Nom. ed. II. ii. 800, nomen (Celtidis sp.).—Austral.
microphylla, Roxb. Hort. Beng. 17; Fl. Ind. i. 613 = rotundifolia.
mirtoides, Ortega, Hort. Matr. Dec. 119 = Condalia microphylla.
mirzaporiensis, Royle, Illustr. Bot. Himal. 168, nomen (Quid?).—Reg. Himal.
Mistol, *Griseb. in Goett. Abh.* xix. (1874) 99.—Reg. Argent.
mitis, A. Rich. Tent. Fl. Abyss. i. 137 = mucronata.
mucronata, *Willd. Enum. Hort. Berol.* 251.—Afr. trop. et austr.
myrtoides, Auct. = *mirtoides*.
Nabeca, Edgew. in Journ. Linn. Soc. vi. (1862) 200 sphalm. = seq.
Napeca, Lam. Encyc. iii. 320 = Z. Spina-Christi.
Napeca, Linn. Herb. ex M. Laws. in Hook. f. Fl. Brit. Ind. i. 635 = Linnaei.
Napeca, Roxb. Fl. Ind. i. 612 = Z. Oenoplia.
Napeca, *Willd. Sp. Pl.* i. 1104.—Zeylan.?
Nats'me, Siebold, in Verh. Batav. Gen. xii. (1830) 48 = sativa.
nitida, Roxb. Hort. Beng. 17; Fl. Ind. i. 609 = sativa.
nummularia, DC. Prod. ii. 21, in syn.; Wight & Arn. Prod. 162 = rotundifolia.
obliqua, Heyne, ex Roth, Nov. Pl. Sp. 160 = rugosa.
obtusifolia, *A. Gray, Gen. Illustr.* ii. 170.—Texas.
Oenoplia, *Mill. Gard. Dict.* ed. VIII. n. 3.—As. et Austral. trop.
officinarum, Medic. Bot. Beobacht. (1782) 333 = sativa.
oppositifolia, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 370 = Sageretia oppositifolia.
orbicularis, Roem. & Schult. Syst. v. 338 = Z. Xylopyrus.
ornata, *Miq. Fl. Ind. Bat.* i. 1. 642.—Ins. Banca.
orthacantha, DC. Prod. ii. 21 = Z. Jujuba.
oxycarpa, *DC. l. c.*—Malaya.
oxyphylla, *Edgew. in Trans. Linn. Soc.* xx. (1846) 43.—Reg. Himal.
Paliurus, Willd. Berl. Baumz. 415; Host, Fl. Austr. i. 306 = Paliurus Spina-Christi (cf. Addend.).
pallens, Wall. Cat. n. 4247 = Z. Oenoplia.

ZIZYPHUS:—
paniculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 190 = incurva.
paniculata, Roth, Nov. Pl. Sp. 161 = rugosa.
Parryi, *Torr. Bot. Mex. Bound.* 46.—Calif.
parvifolia, *Delile, Cent. Pl. Méroé*, 89.—Aegypt.
pedicellata, Wall. Cat. n. 4243 = Z. Oenoplia.
peruviana, Lam. Encyc. iii. 320 = Maytenus octogona.
platyphylla, *Reiss. in Mart. Fl. Bras.* xi. 1. 87.—Bras.
Poiretiana, D. Dietr. Syn. Pl. i. 811 = Berchemia racemosa.
Poiretii, *G. Don, Gen. Syst.* ii. 26.—Ins. S. Doming.
pomaderroides, Fenzl, in Enum. Pl. Hueg. 20 = Alphitonia excelsa.
pubiflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 479.—Ins. Timor.
quadrilocularis, *F. Muell. Fragm.* iii. 57.—Austral.
racemosa, Wall. Cat. n. 7878 D = Bridelia stipularis.
ramosissima, Spreng. Syst. i. 771 = Paliurus ramosissimus.
reticulata, *DC. Prod.* ii. 20.—Am. austr.
Rignonii, *Delponte, in Mem. Acc. Tor.* xiv. (1854) 404.—Hab.?
rotundata, DC. Prod. ii. 21 = Z. Jujuba.
rotundifolia, *Lam. Encyc.* iii. 319.—Persia; Ind. or.
rotundifolia, Roth, Nov. Pl. Sp. 160 = Z. Xylopyrus.
Roxburghiana, Voigt, Hort. Suburb. Calc. 146 = rugosa.
rufa, Mart. in Flora, xxiv. (1841) II. Beibl. 65 = Rhamnus ruffa?
rufula, Miq. Fl. Ind. Bat. i. 1. 643 = Z. Oenoplia.
rugosa, *Lam. Encyc.* iii. 319.—Ind. or.; Burma.
ruminata, Buch.-Ham. ex Wall. Cat. n. 4236 = Z. Xylopyrus.
sativa, *Gaertn. Fruct.* i. 202.—Reg. Mediterr.; As. temp.
scandens, Roxb. Hort. Beng. 17; et in Wall. Cat. n. 7269 = Z. Oenoplia.
sinensis, Lam. Encyc. iii. 317 = sativa.
soporifera, Roem. & Schult. Syst. v. 340 = sativa.
Sororia, Roem. & Schult. l. c. 337 = Z. Jujuba.
sphaerocarpa, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 119.—Ins. Mascar.
Spina-Christi, Georgi, Beschr. Russ. Reich. III. iv. 795 = Paliurus aculeatus.
Spina-Christi, *Willd. Sp. Pl.* i. 1105.—Afr. bor.; Oriens.
spinosa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 70 = praec.
strychnifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 380.—N. Granat.
subquinquenervia, *Miq. Fl. Ind. Bat. Suppl.* 330.—Sumatra.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 2 = Z. Lotus.
thyrsiflora, *Benth. Bot. Voy. Sulph.* 78.—Ecuador.
timoriensis, *DC. Prod.* ii. 20.—Ins. Timor.
tomentosa, Poir. Encyc. Suppl. iii. 192 = Poiretii.
tomentosa, Roxb. Hort. Beng. 17; Fl. Ind. i. 611 = rugosa.
trinervia, Poir. Encyc. Suppl. iii. 192 = exserta.
trinervia, Roth, Nov. Pl. Sp. 168 = Z. Jujuba.
trinervia, Roxb. Hort. Beng. 17; Fl. Ind. i. 606 = glabrata.
tripartita, Roem. & Schult. Syst. v. 342 = Rhus albida.
umbellata, Poir. Encyc. Suppl. iii. 193 = Karwinskia Humboldtiana.
undulata, *Reiss. in Mart. Fl. Bras.* xi. 1. 88.—Bras.
venulosa, Wall. Cat. n. 4235 = funiculosa.
virgata, D. Dietr. Syn. Pl. i. 810 = incurva.
volubilis, Willd. Berl. Baumz. 415 = Berchemia volubilis.
vulgaris, Lam. Encyc. iii. 316 = sativa.
wynadensis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 42.—Ind. or.
Xylopyrus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 136 = Z. Jujuba.
Xylopyrus, *Willd. Sp. Pl.* i. 1104.—Ind. or.
Zeyheriana, *Sond. in Harv. & Sond. Fl. Cap.* i. 476.—Afr. austr.
zonulatus, Blanco, Fl. Filip. ed. II. 120 = Z. Xylopyrus.

ZIZZIA, Roth, Man. Bot. ii. 896 (1830) (Crucif.).
pyrenaica, Roth, l. c. = Draba pyrenaica.
tortuosa, Roth, l. c. 897 = Alyssum tortuosum.

ZODUBA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 30 (1825) = **Calanthe**, R. Br. (Orchid.).
Masuca, Buch.-Ham. l. c. = C. Masuca.

ZOËGEA, Linn. Mant. i. 15 (1767). *COMPOSITAE*, Benth. & Hook. f. ii. 477.
aleppica, *Jacq. Coll.* i. 89.—Hab.?
aristata, DC. Prod. vi. 562 = purpurea.
capensis, Linn. f. Suppl. 382 = Relhania pedunculata.
crinita, *Boiss. Diagn.* Ser. II. iii. 63.—Persia; Afghan.
deluensis, Bunge, ex Boiss. Fl. Orient. iii. 698 = crinita.
Leptaurea, *Linn. Mant.* i. 117.—Syria; Mesopot.
mianensis, *Bunge, ex Boiss. Fl. Orient.* iii. 697.—Persia.
purpurea, *Fresen. in Mus. Senckenb.* i. (1833) 86. t. 5.—Persia; Afghan.; Arab.

ZOLLERNIA, Maximil. & Nees, in Nov. Act. Nat. Cur. xiii. II. (1827) praef. p. xiii. t. C, D. *LEGUMINOSAE*, Benth. & Hook. f. i. 560.
COQUEBERTIA, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1833) 111.
discolor, Vog. in Linnaea, xi. (1837) 166 = falcata.
falcata, *Maximil. & Nees, in Nov. Act. Nat. Cur.* xiii. II. (1827) praef. p. xvi.—Bras.
Houlletiana, Tul. in Arch. Mus. Par. iv. (1844) 190 = ilicifolia.
ilicifolia, *Vog. in Linnaea*, xi. (1837) 166.—Bras.
latifolia, *Benth. in Mart. Fl. Bras.* xv. II. 9.—Bras.
securidacifolia, *Benth. l. c.*—Bras.
splendens, *Maximil. & Nees, in Nov. Act. Nat. Cur.* xiii. II. (1827) praef. p. xv.—Bras.
Vogelii, Tul. in Arch. Mus. Par. iv. (1844) 192 = ilicifolia.

ZOLLIKOFERIA, DC. Prod. vii. 183 (1838) = **Launaea**, Cass. (Compos.).
acanthodes, Boiss. Fl. Orient. iii. 827.—Persia.
angustifolia, Coss. & Dur. in Bull. Soc. Bot. Fr. ii. (1855) 254 = arabica.
arabica, Boiss. Diagn. Ser. I. vii. 12.—Afr. bor.
Candolleana, Coss. ex Pomel, Nouv. Mat. Fl. Atl. 8 = L. chondrilloides.
Cassiniana, Boiss. Fl. Orient. iii. 822.—Aegypt.; Arab.
chondrilloides, DC. Prod. vii. 183 = L. chondrilloides.
fallax, Boiss. Fl. Orient. iii. 824.—Aegypt.; Persia; Afghan.
glomerata, Boiss. l. c. 826 = L. glomerata.
leucodon, Fisch. & Mey. ex Boiss. l. c. 824 = *fallax*.
longiloba, Boiss. & Reut. Pugill. Pl. Nov. (1852) 70 = *resedifolia*.
massavensis, Boiss. Fl. Orient. iii. 825.—Afr. bor.; Arab.
mucronata, Boiss. Diagn. Ser. I. vii. 12 = L. chondrilloides.
nudicaulis, Boiss. Fl. Orient. iii. 824 = L. nudicaulis.
polyclada, Boiss. l. c. 827.—Persia; Beluchist.
prenanthoides, Hochst. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 459 = Sonchus prenanthoides.
pumila, DC. Prod. vii. 183.—Hispan.; Arab.
quercifolia, Coss. & Kral. in Bull. Soc. Bot. Fr. iv. (1857) 369 = Sonchus quercifolius.
resedifolia, Coss. Notes Crit. 120 = Sonchus chondrilloides.
spinosa, Boiss. Fl. Orient. iii. 826 = L. spinosa.
stenocephala, Boiss. l. c. 823.—Beluchist.
Stocksiana, Boiss. l. c. 825.—Beluchist.
tenuiloba, Boiss. Diagn. Ser. I. xi. 50.—Syria; Arab.

ZOLLIKOFERIA, Nees, in Bluff & Fingerh. Comp. Fl. Germ. ed. I. ii. 305 (1825) = **Chondrilla**, (Compos.).
hieracioides, Nees, l. c. = C. Peltidium.
Peltidium, Gaud. Fl. Helvet. v. 143 = C. Peltidium.

ZOLLINGERIA, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 303. *SAPINDACEAE*.
macrocarpa, *Kurz, l. c.*—Burma.
macrocarpa, F. Villar, in Blanco, Fl. Filip. ed. III. Nov. App. 53 = Tristira triptera.
triptera, Rolfe, in Journ. Linn. Soc. xxi. (1884) 309 = Tristira triptera.

ZOLLINGERIA, Sch. Bip. in Flora, xxxvii. (1854) 273 = **Rhychospermum**, Reinw. (Compos.).
scandens, Sch. Bip. l. c. 275 = R. verticillatum.

ZOMICARPA, Schott, Syn. Aroid. 33 (1856). *AROIDEAE*, Benth. & Hook. f. iii. 970.
Pythonium, *Schott, l. c.*—Bras.
Riedelianum, *Schott, Gen. Aroid.* t. 23. f. 18, 34.—Bras.
Steigeriana, *Ferd. Maxim. ex Schott, in Bonplandia*, x. (1862) 86.—Bras.

ZOMICARPELLA, N. E. Br. in Gard. Chron. (1881) II. 266. *AROIDEAE*, Benth. & Hook. f. iii. 970.
maculata, *N. E. Br. l. c.*—N. Granat.

ZONABLEPHIS, Rafin. Sylva Tellur. 143 (1838) = **Acanthus**, Tourn.
polistachys, Rafin. l. c. 144 = A. polystachyus.

ZONANTHUS, Griseb. in Journ. Linn. Soc. vi. (1862) 145. *GENTIANEAE*, Benth. & Hook. f. ii. 813.
cubensis, *Griseb. l. c.*—Cuba.

ZONARIA, Steud. Nom. ed. II. i. 101 (1840), sphalm. = **Zornia**, J. F. Gmel. (Legumin.).

ZOOPHORA, Bernh. Syst. Verz. Erf. 308 (1800) = **Orchis**, Tourn.
atropurpurea, Bernh. l. c. 311 = O. militaris.
rubella, Bernh. l. c. = O. militaris.

ZOOPHTHALMUM, P. Br. Hist. Jamaic. 295. t. 31 (1756) = **Mucuna**, Adans. (1763) (Legumin.).

ZORNIA, J. F. Gmel. Syst. 1076, 1096 (1791). *LEGUMINOSAE*, Benth. & Hook. f. i. 518.
MYRIADENUS, Desv. Journ. Bot. iii. (1813) 121. t. 4.
andicola, *Griseb. in Goett. Abh.* xxiv. (1879) 106.—Reg. Argent.
angustifolia, Sm. in Rees, Cycl. xxxix. n. 1 = diphylla.
barbata, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 325.—Peruv.
biarticulata, G. Don, Gen. Syst. ii. 280 = diphylla.
brachycarpa, Walp. Rep. i. 726, sphalm. = *trachycarpa*.
bracteata, *J. F. Gmel. Syst.* 1096.—Am. bor.; Afr. austr.
brasiliensis, *Vog. in Linnaea*, xii. (1838) 62.—Bras.
capensis, Pers. Syn. ii. 318 = bracteata.
ceylanica, Heynh. Nom. i. 865 = diphylla.
chaetophora, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 56 = diphylla.
conjugata, Sm. in Rees, Cycl. xxxix. n. 3 = diphylla.
diphylla, *Pers. Syn.* ii. 318.—Cosmop. trop.
disperma, R. Grah. in Wall. Cat. n. 5663 = Geissaspis cristata.
dyctiocarpa, DC. Prod. ii. 317 = diphylla.
echinocarpa, *Benth. in Mart. Fl. Bras.* xv. 1. 85.—Bras.
elegans, Pers. Syn. ii. 318 = Desmodium elegans.
flemmingioides, *Moric. Pl. Nouv. Am.* 127. t. 76.—Bras.
Gardneriana, *Moric. l. c.* 129. t. 77.—Bras.
gemella, Vog. in Linnaea, xii. (1838) 61 = diphylla.
gibbosa, Span. l. c. xv. (1841) 192 = diphylla.
glabella, Span. l. c. = diphylla.
glabra, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 325.—Bras.
glauca, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 228 = Phylacium bracteosum.
glochidiata, Reichb. ex DC. Prod. ii. 316 = diphylla.
gracilis, DC. l. c. 316 = diphylla.
graminea, Span. in Linnaea, xv. (1841) 192 = diphylla.
havanensis, *A. Rich. Ess. Fl. Cub.* 423.—Cuba.
heterophylla, Sm. in. Rees, Cycl. xxxix. n. 5 = bracteata.
inermis, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 325.—Hab.?
laevis, Schlecht. & Cham. in Linnaea, v. (1830) 582 = diphylla.
latifolia, DC. Prod. ii. 317 = diphylla.
linearis, E. Mey. Comm. Pl. Afr. Austr. 122 = bracteata.
lutescens, Steud. Nom. ed. I. 900 = Desmodium lutescens.
maranhamensis, *G. Don, Gen. Syst.* ii. 280.—Bras.
microphylla, Desv. in Mém. Soc. Linn. Par. iv. (1826) 325 = diphylla.
myriadena, *Benth. in Mart. Fl. Bras.* xv. 1. 85.—Bras.
nuda, Vog. in Linnaea, x. (1836) 587 = diphylla.

ZORNIA:—
ovata, Vog. in Linnaea, xii. (1838) 58 = diphylla.
perforata, Vog. l. c. 59 = diphylla.
pubescens, H. B. & K. Nov. Gen. et Sp. vi. 515 = diphylla.
pulchella, Pers. Syn. ii. 318 = Desmodium pulchellum.
reticulata, Sm. in. Rees, Cycl. xxxix. n. 2 = diphylla.
sericea, Moric. Pl. Nouv. Am. 126. t. 75 = diphylla.
Sloanei, *Griseb. Fl. Brit. W. Ind.* 709.—Cuba.
strobilifera, Pers. Syn. ii. 319 = Flemingia strobilifera.
surinamensis, Miq. in Ann. & Mag. Nat. Hist. Ser. I. xi. (1843) 14 = diphylla.
tenuifolia, *Moric. Pl. Nouv. Am.* 132. t. 79.—Bras.
tetraphylla, Michx. Fl. Bor. Am. ii. 76. t. 41 = bracteata.
thymifolia, H. B. & K. Nov. Gen. et Sp. vi. 514 = diphylla.
trachycarpa, Vog. in Linnaea, xii. (1838) 60 = diphylla.
venusta, Wall. ex Steud. Nom. ed. II. ii. 801, nomen (Quid ?).—Ind. or.
virgata, *Moric. Pl. Nouv. Am.* 131. t. 78.—Bras.
Walkeri, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 330 = diphylla.
zeylonensis, Pers. Syn. ii. 318 = diphylla.

ZORNIA, Moench, Meth. 410 (1794) = **Lallemantia**, Fisch. (Legumin.).
canescens, Moench, l. c. 411 = L. canescens.
linearifolia, Moench, Meth. Suppl. 139 = Dracocephalum Ruyschiana.
nutans, Moench, Meth. 411 = Dracocephalum nutans.
partita, Moench, Meth. Suppl. 139 = Dracocephalum austriacum.
parviflora, Moench, Meth. 411 = Dracocephalum thymiflorum.
peltata, Moench, l. c. = L. peltata.

ZOROXUS, Rafin. New Fl. Am. iv. 88 (1836) = **Polygala**, Tourn.

ZOSIMA, Hoffm. Gen. Umbell. ed. I. 145. t. 4 (1814). *UMBELLIFERAE*, Benth. & Hook. f. i. 924.
absinthifolia, Link, Enum. Hort. Berol. i. 274 (*apsinthifolia*); DC. Prod. iv. 195 = orientalis.
anethifolia, DC. l. c. 196 = Ducrosia anethifolia.
dichotoma, *Boiss. Diagn.* Ser. I. vi. 63.—Persia.
frigida, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1038.—Persia.
humilis, Fenzl, ex Boiss. l. c. 1062 (= *Pastinaca zozimioides*).—Cilicia.
lasiocarpa, *Boiss. l. c.* 1038.—Afghan.
Nuttallii, D. Dietr. Syn. Pl. ii. 973 = Polytaenia Nuttallii.
orientalis, *Hoffm. Gen. Umbell.* ed. I. 148. t. 4.—As. occ.
radians, Boiss. & Hohen. Diagn. Ser. I. x. 43 = orientalis.
tragioides, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 341.—Persia.

ZOSIMA, Phil. Sert. Mendoc. Alt. 29 (1871) = **Philibertia**, H. B. & K. (Asclep.).
violacea, Phil. l. c. = P. solanoides.

ZOSIMIA, Bieb. Fl. Taur. Cauc. iii. 229 (1819) = **Zosima**, Hoffm. (Umbellif.).

ZOSTER, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = seq.

ZOSTERA, Linn. Diss. Dass. 27 (1747); Amoen. Acad. i. 411 (1749). *NAIADACEAE*, Benth. & Hook. f. iii. 1017.
ALGA, Lam. Fl. Fr. iii. 539 (1778).
angustifolia, Loser, in Oestr. Bot. Zeitschr. xiii. (1863) 383 = nana.
angustifolia, Reichb. Ic. Fl. Germ. vii. 3 = marina.
bullata, Delile, Fl. Egypte, 289. t. 53. f. 6 = Halophila stipulacea ?
ciliata, Forsk. Fl. Aegypt. Arab. 157 = Cymodocea ciliata.
emarginata, Ehrenb. & Hempr. ex Aschers. in Linnaea, xxxv. (1867–68) 167 = nana.
marina, R. Br. Prod. 338 = tasmanica.
marina, *Linn. Sp. Pl.* 968.—Europ.
marina, Targ. Tozz. Cat. Veg. Marin. (1826) 87 = Cymodocea aequorea.
marina, Vis. Stirp. Dalm. Spec. 51 = Posidonia Caulini.

ZOSTERA:—
maritima, Gaertn. Fruct. i. 76. t. 19 = marina.
mediterranea, DC. Fl. Fr. iii. 154 = Cymodocea aequorea.
minor, Nolte, ex Reichb. Ic. Fl. Germ. vii. 2 = nana.
Muelleri, Irmisch, ex Aschers. in Linnaea, xxxv. (1867–68) 168 = nana.
nana, *Roth, Enum. Pl. Phaen. Germ.* i. 8.—Reg. temp.
nodosa, Guss. Pl. Sic. Syn. ii. 565 = nana.
nodosa, Ucria, in Roem. Arch. i. 1. (1796) 70 = Cymodocea aequorea.
Noltii, Hornem. in Fl. Dan. t. 2041 = nana.
oceanica, Linn. Mant. i. 123 = Posidonia Caulini.
oceanica, [Weber,] in Wigg. Prim. Fl. Holsat. 65 = marina.
pumila, Le Gall, in Congr. Sc. Fr. xvi. 1. (1850) 96, 144 = nana.
serrulata, Bertol. Fl. Ital. x. 5 = marina.
serrulata, Targ. Tozz. Cat. Veg. Marin. (1826), 85 = Cymodocea aequorea.
stenophylla, Rafin. in Am. Monthly Mag. (1818) 175 = marina.
stipulacea, Forsk. Fl. Aegypt. Arab. 158 = Halophila ovata.
tasmanica, *Martens, ex Aschers. in Linnaea*, xxxv. (1867–68) 168.—Austral.
tridentata, Solms, in Schweinf. Beitr. Fl. Aethiop. 196 = Halodule australis.
trinervis, Stokes, Bot. Mat. Med. iv. 319 = marina.
uninervis, Forsk. Fl. Aegypt. Arab. 157 = Halodule australis.
uninervis, Vahl, ex Reichb. Fl. Germ. Excurs. 137 = nana.

ZOSTEROSPERMUM, Beauv. in Lestib. Essai Fam. Cyp. 28 (1819) = **Rynchospora**, Vahl (Cyperac.).
gracile, Desv. ex Ham. Prod. Pl. Ind. Occ. 14 = R. glauca ?

ZOSTEROSTYLIS, Blume, Bijdr. 418. t. 32 (1825) = **Cryptostylis**, R. Br. (Orchid.).
Arachnites, Blume, l. c. 419. t. 32 = C. Arachnites.
filiformis, Miq. Fl. Ind. Bat. iii. 721 = C. filiformis.
Walkerae, Wight, Ic. t. 1748 = C. Arachnites.
zeylanica, Lindl. Gen. et Sp. Orch. 446 = C. Arachnites.

ZOUCHIA, Rafin. Fl. Tellur. iv. 22 (1836) = **Pancratium**, Dill. (Amaryll.).
illyrica, Rafin. l. c. = P. illyricum.

ZOZIMA, DC. Prod. iv. 195 (1830), sphalm. = **Zosima**, Hoffm. (Umbellif.).

ZOZIMIA, DC. Prod. iv. 195 (1830) = praec.

ZOYSIA, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 440. *GRAMINEAE*, Benth. & Hook. f. iii. 1124.
MATRELLA, Pers. Syn. i. 73 (1805).
OSTERDAMIA, Neck. Elem. iii. 218 (1790).
aristata, C. Muell. in Bot. Zeit. xiii. (1855) 272 = pungens.
Brownii, C. Muell. l. c. = pungens.
Griffithiana, C. Muell. l. c. 273 = pungens.
japonica, Steud. Syn. Pl. Gram. 414 = pungens.
macrantha, Desv. Opusc. 54 = pungens.
macrostachya, Franch. & Sav. Enum. Pl. Jap. ii. 608 = pungens ?
pungens, *Willd. in Ges. Naturf. Fr. Neue Schr.* iii. (1801) 441.—As. or.; Austral.; Ins. Mascar.; Maurit.
rigida, Willd. ex Kunth, Rév. Gram. i. 156 = Leptothrium rigidum.
sedoides, C. Muell. in Bot. Zeit. xiii. (1855) 273 = pungens.
setacea, Nees, ex Steud. Nom. ed. II. ii. 801 = pungens.
sinica, Hance, in Journ. Bot. vii. (1869) 168 = pungens.
tenuifolia, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. Sc. Nat. ii. (1836) 96 = pungens.
tremula, Beauv. Agrost. 148 = Sporobolus virginicus.

ZSCHOKKEA, Muell. Arg. in Mart. Fl. Bras. vi. 1. 20. t. 6, 7 (1860) = **Lacmellia**, Karst. (Apocyn.).
arborescens, Muell. Arg. l. c. 22.—Bras.
floribunda, Muell. Arg. l. c. 23 = L. floribunda.
gracilis, Muell. Arg. l. c. 21.—Bras.

ZSCHOKKEA:—
guyanensis, Muell. Arg. in Linnaea, xxx. (1859–60) 391 = Lacmellia floribunda.
microcarpa, Muell. Arg. in Mart. Fl. Bras. vi. 1. 23.—Bras.
monosperma, Muell. Arg. l. c. 22.—Bras.
ramosissima, Muell. Arg. l. c. 21.—Bras.

ZSCHOKKIA, Benth. & Hook. f. Gen. ii. 694 (1876), sphalm. = praec.

ZUCCA, Comm. ex Juss. Gen. 398 (1789) = **Momordica**, Tourn. (Cucurb.).
Commersoniana, Ser. in DC. Prod. iii. 319 = M. cochinchinensis.

ZUCCAGNIA, Cav. Ic. v. 2. t. 403 (1799). *LEGUMINOSAE*, Benth. & Hook. f. i. 587.
angulata, *Hook. & Arn. Bot. Beech. Voy.* 22.—Chili.
eremophila, *Phil. Fl. Atac.* 17. t. 1 B.—Chili.
microphylla, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 38.—Chili.
mucronata, *Phil. Fl. Atac.* 17.—Chili.
punctata, *Cav. Ic.* v. 2. t. 403.—Chili.

ZUCCAGNIA, Thunb. Nov. Gen. ix. 127 (1798) (Zuccangnia) = **Dipcadi**, Medic. (Liliac.).
viridis, Thunb. l. c. = D. filamentosum.

ZUCCARINIA, Blume, Bijdr. 1006 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 97.
macrophylla, *Blume, l. c.* 1007.—Java.

ZUCCARINIA, Maerckl. in Ann. Ges. Wetterau, ii. (1811) 252 (Gen. dub.).
verbenacea, Maerckl. l. c. 253 (Quid ?).—Hab. ?

ZUCCARINIA, Spreng. Syst. iv. Cur. Post. 50, 81 (1827) = **Jackia**, Wall. (Rubiac.).
ornata, Spreng. l. c. 81 = J. ornata.

ZUCCHELIA, Buek, Index, in DC. Prod. iii. 508 (1858), sphalm. = seq.

ZUCCHELLIA, Decne. in DC. Prod. viii. 492 (1844) = **Raphionacme**, Harv. (Asclep.).
angolensis, Decne. l. c.—Afr. trop.

ZUCKERTIA, Baill. Etud. Gén. Euph. 495. t. 4 (1858) = **Tragia**, Plum. (Euphorb.).
cordata, Baill. l. c. 496 = T. Bailloniana.

ZUELANIA, A. Rich. Ess. Fl. Cub. 88. t. 12 (1845) = **Casearia**, Jacq. (Samyd.).
crenata, Griseb. Cat. Pl. Cub. 9.—Cuba.
icosandra, Clos, in Ann. Sc. Nat. Sér. IV. viii. (1857) 242, in obs. = C. icosandra.
laetioides, A. Rich. Ess. Fl. Cub. 89 = C. icosandra.
tremula, Griseb. Cat. Pl. Cub. 9.—Cuba.

ZUGILUS, Rafin. Fl. Ludov. 159 (1817) = **Ostrya**, Mich. (Cupul.).
virginica, Rafin. l. c. 159 = O. virginica.

ZULATIA, Neck. Elem. ii. 117 (1790) = **Miconia**, Ruiz & Pav. (Melastom.).
alata, Rafin. Sylva Tellur. 97 = M. alata.
grandiflora, Rafin. l. c. = Rhynchanthera grandiflora.
levigata, Rafin. l. c. = M. laevigata.

ZURLOA, Tenore, Atti 3ª Riun. Sc. Ital. 503 (1841) = **Carapa**, Aubl. (Meliac.).
insignis, Tenore, l. c. = C. procera.
splendens, Tenore, Cat. Orto Bot. Nap. (1845) 99 = C. procera.

ZWAARDEKRONIA, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 252 = **Psychotria**, Linn. (Rubiac.).
lurida, Korth. l. c. = Chasalia curviflora.

ZWACKHIA, Sendt. in Reichb. Ic. Fl. Germ. xviii. 65. t. 115, 1316 (1858). *BORAGINEAE*, Benth. & Hook. f. ii. 862.
aurea, *Sendt. l. c.*—Banat.

ZWARDEKRONIA, Hook. f. Fl. Brit. Ind. iii. 177 (1880), err. typ., in syn. = ZWAARDEKRONIA, Korth. = **Psychotria**, Linn. (Rubiac.).

ZWINGERA, Neck. Elem. iii. 385 (1790) = **Nolina**, Linn. (Convolv.).

ZWINGERA, Schreb. Gen. ii. 802 (1791) = **Simaba**, Aubl. (Simarub.).
amara, Willd. Sp. Pl. ii. 569 = Simaruba amara.
Aruba, Spreng. Syst. ii. 319 = Simaba guianensis.
bahiensis, Steud. Nom. ed. II. ii. 802 = Simaba ferruginea?
bicolor, Steud. l. c. = Simaba bicolor.
ferruginea, Spreng. Syst. ii. 319 = Simaba ferruginea.
floribunda, Spreng. l. c. = Simaba floribunda.
orinocensis, Spreng. l. c. = Simaba orinocensis.
quassioides, Spreng. Syst. iv. Cur. Post. 163 = Picrasma quassioides.
suaveolens, Spreng. Syst. ii. 319.—Bras.
trichilioides, Spreng. l. c. = Simaba trichilioides.
undulata, Steud. Nom. ed. II. ii. 802 = Simaba undulata.

ZYGADENUS, Michx. Fl. Bor. Am. i. 213. t. 22 (1803) (Zigadenus). *LILIACEAE*, Benth. & Hook. f. iii. 835.
AMIANTANTHIUM, Kunth, Enum. Pl. iv. 179 (1843).
AMIANTHUM, Rafin. Fl. Tellur. iv. 27 (1836).
ANTICLEA, Kunth, Enum. Pl. iv. 191 (1843).
CHITONIA, Salisb. Gen. Pl. Fragm. 51 (1866).
CHROSPERMA, Rafin. Neogenyt. 3 (1825).
ENDOCLES, Salisb. Gen. Pl. Fragm. 51 (1866).
GEISELERIA, Kunth, in Abh. Berl. Akad. (1842) 3.
GOMPHOSTYLIS, Rafin. Fl. Tellur. ii. 30 (1836).
MONADENUS, Salisb. Gen. Pl. Fragm. 51 (1866).
angustifolius, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 280.—Am. bor.
bracteatus, Sweet, Hort. Brit. ed. I. 429; Schult. f. Syst. vii. 1559 = elegans.
canadensis, Hort. ex Baker, in Journ. Linn. Soc. xvii. (1879) 482 = elegans.
chloranthus, Richards. in Frankl. Narr. 1st Journ. App. 736 = elegans.
commutatus, Schult. f. Syst. vii. 1560 = elegans.
Douglasii, Torr. in Pacif. Rail. Rep. vii. 20 = Fremontii.
elegans, *Pursh, Fl. Am. Sept.* i. 241.—Am. bor.
Fremontii, *Torr. ex S. Wats. Bot. King's Exped.* 343.—Calif.
frigidus, D. Don, in Edinb. N. Phil. Journ. (Oct. 1832) 233 = Stenanthium frigidum.
fuscatus, *Rafin. Fl. Tellur.* ii. 30 (Quid?).—Hab.?
glaberrimus, Hook. & Arn. Bot. Beech. Voy. 160 = Fremontii.
glaberrimus, *Michx. Fl. Bor. Am.* i. 214. t. 22.—Am. bor.
glaucus, Nutt. ex Baker, in Journ. Linn. Soc. xvii. (1879) 480 = Fremontii.
glaucus, Nutt. in Journ. Acad. Philad. vii. (1834) 56 = elegans.
hybridus, Endl. ex Kunth, Enum. Pl. iv. 196 = Melanthium virginicum.
japonicus, Miq. in Verh. Kon. Akad. Wetensch. II. ii. (1868) 88 = Veratrum Maackii.
leimanthoides, *A. Gray, Man. Bot. N. U. St.* ed. II. 476.—Am. bor.
mexicanus, *Hemsl. Biol. Centr. Am. Bot.* iii. 382.—Mexic.
monoecus, Kunth, Enum. Pl. iv. 196 = Veratrum parviflorum.
muscitoxicum, *Regel, Gartenfl.* xxxii. (1883) 164.—Am. bor.
Nuttallii, *A. Gray, Man. Bot. N. U. St.* ed. II. 476, in obs.—Calif.
Nuttallii, S. Wats. in Bot. King's Exped. 343, partim = venenosus.
paniculatus, *S. Wats. l. c.*—Am. bor. occ.
porrifolius, *Greene, in Bull. Torrey Bot. Club*, viii. (1881) 123.—N. Mexic.
sachalinensis, F. Schmidt, Reisen Amurl. 88 = Stenanthium sachalinense.
sibiricus, *A. Gray, in Ann. Lyc. N. York*, iv. (1837) 112.—Sibir.
speciosus, Dougl. ex Hook. Fl. Bor. Am. ii. 177 = elegans.
venenosus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 279.—Am. bor. occ.

ZYGADENUS:—
virginicus, Endl. ex Kunth, Enum. Pl. iv. 195 = Melanthium virginicum.
volcanicus, *Benth. Pl. Hartw.* 96.—Guatemala.
volcanicus, S. Wats. in Proc. Am. Acad. xviii. (1883) 165 = mexicanus.

ZYGIA, Benth. & Hook. f. Gen. ii. 1188 (1876), err. typ. = ZYGIS, Desv. = **Micromeria**, Benth. (Labiat.).

ZYGIA, P. Br. Hist. Jamaic. 279. t. 22 (1789) = **Pithecolobium**, Mart. (Legumin.).

ZYGIA, Walp. Rep. i. 928 (1842); Benth. in Hook. Lond. Journ. Bot. iii. (1844) 92 = **Albizzia**, Durazz. (Legumin.).
Brownei, Schweinf. Rel. Kotsch. 10. t. 10 = A. Welwitschiii.
Brownei, Walp. Rep. i. 928 = A. Brownei.
fastigiata, E. Mey. Comm. Pl. Afr. Austr. 165 = A. fastigiata.
Petersiana, Bolle, in Peters, Reise Mossamb. Bot. t. 1 = A. Petersiana.
Sassa, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 93, in obs. = A. fastigiata.
virgultosa, E. Mey. ex Walp. Rep. i. 928 (Ingae sp.).—Afr. austr.

ZYGIS, Desv. in Ham. Prod. Pl. Ind. Occ. 46 (1825) = **Micromeria**, Benth. (Labiat.).
aromatica, Desv. l. c. = M. obovata.

ZYGODIA, Benth. in Benth. & Hook. f. Gen. ii. 716 (1876). *APOCYNACEAE*, Benth. & Hook. f. l. c.
axillaris, *Benth. in Hook. Ic. Pl.* t. 1184.—Afr. trop.
myrtifolia, *Benth. l. c.* xii. 73.—Afr. trop.
subsessilis, *Benth. l. c.*—Afr. trop.

ZYGOGLOSSUM, Reinw. ex Blume, Cat. Gew. Buitenz. 100 (1823), nomen; et in Syll. Ratisb. ii. (1828) 4 = **Cirrhopetalum**, Lindl. (Orchid.).
umbellatum, Reinw. ll. cc. 100; 5 = C. Thouarsii.

ZYGOGYNUM, Baill. Adansonia, vii. (1867) 298. *MAGNOLIACEAE*.
pomiferum, *Baill. l. c.* x. (1872) 334.—N. Caled.
stipitatum, *Baill. l. c.*—N. Caled.
Vieillardi, *Baill. l. c.* vii. (1867) 298.—N. Caled.

ZYGOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 573 = **Ratonia**, DC. (Sapind.).
rufescens, Turcz. l. c. 574 = R. Zygolepis.

ZYGOMENES, Salisb. in Trans. Hort. Soc. i. (1812) 271 = **Cyanotis**, D. Don (Commel.).
abyssinica, Hassk. in Schweinf. Beitr. Fl. Aethiop. 211 = C. hirsuta.
axillaris, Salisb. in Trans. Hort. Soc. i. (1812) 271 = C. axillaris.
caespitosa, Kotschy & Peyr. Pl. Tinn. t. 22[d] = C. longifolia.
parasitica, Hassk. in Schweinf. Beitr. Fl. Aethiop. 214 = C. parasitica.
pauciflora, Hassk. l. c. 213 = C. pauciflora.
polyrrhiza, Hassk. l. c. 214 = C. polyrrhiza.

ZYGOMERIS, [Moç. & Sesse,] ex DC. Prod. ii. 315 (1825) = **Amicia**, H. B. & K. (Legumin).
flava, [Moç. & Sesse,] ex DC. l. c. = A. Zygomeris.

ZYGOON, Hiern, in Oliver, Fl. Trop. Afr. iii. 113 (1877). *RUBIACEAE*.
graveolens, *Hiern, l. c.* 114.—Afr. trop.

ZYGOPELTIS, Fenzl, ex Endl. Gen. Suppl. ii. 107 (1842) = **Heldreichia**, Boiss. (Crucif.).

ZYGOPETALON, Reichb. Consp. 69 (1828) = seq.

ZYGOPETALUM, Hook. Bot. Mag. t. 2748 (1827). *ORCHIDEAE*, Benth. & Hook. f. iii. 542.
BOLLEA, Reichb. f. in Bot. Zeit. x. (1852) 667.
CHAUBARDIA, Reichb. f. l. c. (1852) 671.
COCHLEANTHES, Rafin. Fl. Tellur. iv. 45 (1836).
GALEOTTIA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 25.
HUNTLEYA, Batem. ex Lindl. Bot. Reg. (1837) sub t. 1991.

ZYGOPETALUM:—
KEFERSTEINIA, Reichb. f. in Bot. Zeit. x. (1852) 638.
MENADENIUM, Rafin. Fl. Tellur. iv. 45 (1836).
PESCATORIA, Reichb. f. in Bot. Zeit. x. (1852) 667.
PROMENAEA, Lindl. Bot. Reg. (1843) Misc. 13.
WARCZEWICZELLA, Reichb. f. in Bot. Zeit. x. (1852) 635.
ZYGOSEPALUM, Reichb. f. in Walp. Ann. vi. 665 (1863).
africanum, Hook. Bot. Mag. t. 3812 = Odontoglossum bictoniense.
amazonicum, *Reichb. f. in Walp. Ann.* vi. 655.—Am. trop.
aromaticum, *Reichb. f. in Bot. Zeit.* x. (1852) 668.—Panama.
Backhousianum, *Reichb. f. in Gard. Chron.* (1877) ii. 456, in syn.—Ecuador?
bellum, *Reichb. f. l. c.* (1878) i. 492, in nota.—N. Granat.
bicallosum, *Reichb. f. Otia Bot. Hamb.* 9.—Ecuador.
bidentatum, *Reichb. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 251.—Mexic.
brachypetalum, *Lindl. Bot. Reg.* (1844) *Misc.* 9.—Bras.
brachystalix, *Reichb. f. in Walp. Ann.* vi. 660.—Ins. Trinit.
Burkei, *Reichb. f. in Gard. Chron.* (1883) ii. 684.—Guian.
Burtii, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 251.—Costa Rica.
candidum, *Reichb. f. in Walp. Ann.* vi. 656.—Bras.
cerinum, *Reichb. f. l. c.* 651.—Panama.
Clayi ×, Reichb. f. in Gard. Chron. (1877) i. 684.
cochleare, *Lindl. in Bot. Reg.* t. 1857.—Am. trop.
cochleata, Paxt. Mag. Bot. iv. (1838) 279 (Index), err. typ. = praec.
coeleste, *Reichb. f. in Linnaea*, xli. (1877) 5.—N. Granat.
coloratum, *Linden & Reichb. f. in Walp. Ann.* vi. 662.—N. Granat.
conchaceum, Hoffmgg. ex Reichb. f. l. c. 653 = cochleare.
coronarium, *Reichb. f. in Linnaea*, xli. (1877) 108.—N. Granat.
crinitum, *Lodd. Bot. Cab.* t. 1687.—Bras.
Dayanum, *Reichb. f. in Gard. Chron.* (1872) 1618, in syn.—N. Granat.
discolor, *Reichb. f. in Walp. Ann.* vi. 655.—Costa Rica.
euglossum, Reichb. f. in Gard. Chron. (1876) ii. 808, in syn. = intermedium.
expansum, *Reichb. f. l. c.* (1878) i. 168.—Ecuador?
flabelliforme, Reichb. f. in Walp. Ann. vi. 652 = cochleare.
flaveolum, *Linden & Reichb. f. in Walp. l. c.* 662.—N. Granat.
forcipatum, Hort.—Cf. Gard. Chron. (1885) ii. 70 = laminatum.
forcipatum, *Reichb. f. l. c.* (1883) ii. 360.—Hab.?
fragrans, Linden, Cat. n. 6 (1851) 9 = marginatum.
Gairianum, *Reichb. f. in Gard. Chron.* (1879) i. 684, in syn.—Hab.?
Gautieri, *Lem. Illustr. Hortic.* xiv. (1867) t. 535.—Bras.
Gemma, *Reichb. f. in Gard. Chron.* (1874) i. 406, in syn.—Hab.?
gramineum, *Lindl. Bot. Reg.* (1844) *Misc.* 10.—N. Granat.
grandiflorum, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 251.—Mexic.
guttatum, *Reichb. f. in Walp. Ann.* vi. 659.—Hab.?
intermedium, Hort. Petrop. ex Regel, Ind. Sem. Hort. Petrop. (1855) 22, 23 = Mackaii.
intermedium, *Lodd. Cat.* n. 1136; *ex Lindl. Bot. Reg.* (1844) *Misc.* 9.—Bras.
ionoleucum, *Reichb. f. in Bot. Zeit.* xxiii. (1865) 99.—N. Granat.
Kegelii, *Reichb. f. l. c.* x. (1852) 668.—Guian.
Klabochii, *Reichb. f. in Gard. Chron.* (1885) ii. 391.—N. Granat.
Klabochorum, *Reichb. f. l. c.* (1879) i. 684.—Hab.?
lacteum, *Reichb. f. l. c.* (1872) 1290.—Panama.
Lalindei, *Reichb. f. l. c.* (1874) ii. 33, in nota.—N. Granat.
lamellosum, *Reichb. f. l. c.* (1875) ii. 225, in nota.—N. Granat.
laminatum, *Reichb. f. l. c.* (1885) ii. 70.—Hab.?
Lawrenceanum, *Reichb. f. l. c.* (1878) ii. 266, in nota.—Hab.?

ZYGOPETALUM:—
Lehmanni, *Reichb. f. in Gard. Chron.* (1879) II. 424, in syn.—Reg. Andin.
lentiginosum, *Reichb. f. in Walp. Ann.* vi. 659.—Bras.
Lueddemannianum, *Reichb. f. l. c.* 653.—Am. trop.
Mackaii, *Hook. Bot. Mag.* t. 2748.—Bras.
mandibulare, Reichb. f. in Gard. Chron. (1877) I. 684 = maxillare.
marginatum, *Reichb. f. in Walp. Ann.* vi. 654.—Am. centr.
Massangei, *Hort.—Cf. Gard. Chron.* (1876) I. 603.—Hab.?
maxillare, *Lodd. Bot. Cab.* t. 1776; *Paxt. Mag. Bot.* iv. (1838) 271.—Bras.
Meleagris, *Benth. in Journ. Linn. Soc.* xviii. (1880) 321; *Benth. & Hook. f. Gen.* iii. 546, in nota.—Bras.
micranthum, Rodrig. Orch. Nov. i. 109 = Paradisanthus paranaënsis.
microphytum, *Rodrig. l. c.*—Bras.
micropterum, *Reichb. f. in Gard. Chron.* (1881) II. 134, in nota.—Hab.?
microtus, Hoffmgg. ex Reichb. f. in Walp. Ann. vi. 661 = crinitum.
Moritzii, *Reichb. f. l. c.* 658.—Am. centr.
Mosenianum, *Rodrig. Orch. Nov.* i. 108.—Bras.
Murrayanum, *Gardn. ex Hook. Bot. Mag.* t. 3674.—Bras.
Mystacinum, *Reichb. f. in Gard. Chron.* (1881) I. 530, in nota.—N. Granat.
obtusatum, *Reichb. f. l. c.* (1878) II. 300.—Hab.?
pardinum, *Reichb. f. in Walp. Ann.* vi. 662.—Reg. And.
Patini, *Reichb. f. in Gard. Chron.* (1874) II. 34, in nota.—Hab.?
pentachromum ×, Reichb. f. l. c. (1885) I. 532.
pictum, *Reichb. f. l. c.* (1883) II. 8, in nota.—Am. trop.
pubescens, Hoffmgg. Verz. Orch.—Cf. Bot. Zeit. i. (1843) 835 = crinitum.
pulvinare, *Reichb. f. in Linnaea*, xli. (1877) 107.—N. Granat.
quadratum, Pfitzer, Vergl. Morph. Orch. 58 = marginatum.
Rivieri, Carr. in Rev. Hortic. (1873) 191, cum ic. = intermedium.
Roezlii, Reichb. f. in Gard. Chron. (1877) I. 620 = intermedium.
Rollissonii, *Reichb. f. in Walp. Ann.* vi. 659.—Bras.
rostratum, *Hook. Bot. Mag.* t. 2819.—Guian.
Russelianum, *Reichb. f. in Gard. Chron.* (1878) II. 524, in syn.—Hab.?
sanguinolentum, *Reichb. f. in Walp. Ann.* vi. 658.—Venezuela.
Sedeni ×, Reichb. f. in Gard. Chron. (1874) II. 290.
Sellowii, *Reichb. f. in Walp. Ann.* vi. 660.—Bras.
stapelioides, *Reichb. f. l. c.* 658.—Bras.
stenochilum, Lodd. Bot. Cab. t. 1923 = crinitum.
taurinum, *Reichb. f. in Linnaea*, xlii. (1877) 5.—N. Granat.
tricolor, Lindl. Bot. Reg. (1846) t. 64 = Acacallis cyanea.
triste, *Rodrig. Orch. Nov.* i. 108.—Bras.
triumphans, *Reichb. f. in Walp. Ann.* vi. 652.—N. Granat.
velatum, *Reichb. f. l. c.* 655.—N. Granat.
velutinum, Hoffmgg. Verz. Orch.—Cf. Bot. Zeit. i. (1843) 835 = intermedium.
violaceum, *Reichb. f. in Walp. Ann.* vi. 650.—Guian.
Wailesianum, *Reichb. f. l. c.* 656.—Bras.
Wallisii, *Reichb. f. in Gard. Chron.* (1869) 710, in syn.; *et Van Houtte, in Fl. des Serres*, xviii. (1869) 17. t. 1828.—Ecuador.
Wendlandi, *Reichb. f. Beitr. Orch. Centr. Am.* 74.—Costa Rica.
xanthinum, *Reichb. f. in Walp. Ann.* vi. 659.—Bras.

ZYGOPHYLLON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.

ZYGOPHYLLUM, Linn. Syst. ed. I (1735). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 266.
AGOPHYLLUM, Neck. Elem. ii. 227 (1790).
AGROPHYLLUM, Endl. Gen. 1164 (1841).
FABAGO, Tourn. ex Adans. Fam. ii. 507 (1763).
ROEPERA, A. Juss. in Mém. Mus. Par. xii. (1825) 454. t. 5. f. 3.

ZYGOPHYLLUM:—
aestuans, *Linn. Sp. Pl.* ed. II. 552 (Quid?).—Guian.
alatum, E. Mey. in Drège, Zwei Pfl. Docum. 230 = fulvum.
album, *Linn. Sp. Pl.* ed. II. 552.—Cilicia; Afr. bor.; Ins. Canar.
altaicum, Steph. ex Ledeb. Fl. Ross. i. 484 = pterocarpum.
ammophilum, F. Muell. in Linnaea, xxv. (1852) 376 = Billardierii.
apiculatum, *F. Muell. l. c.* 373.—Austral.
arboreum, Jacq. Enum. Pl. Carib. 20; Select. Am. 130 = Guaiacum arboreum.
arboreum, Royle, Illustr. Bot. Himal. 154 (Quid?).—Reg. Himal.
atriplicoides, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 41.—Armen.; Persia.
aurantiacum, F. Muell. in Linnaea, xxv. (1852) 376, in nota = fruticulosum.
australasicum, Miq. in Lehm. Pl. Preiss. i. 164 = Nitraria Schoberi.
Billardierii, *DC. Prod.* i. 705.—Austral.
brachypterum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 397.—Persia; Soongar.
capense, Lam. Encyc. ii. 443 = microphyllum.
carnosum, Wall. Cat. n. 1192 (Quid?).—Ind. or.
coccineum, Lepech. It. ed. Germ. i. 310 = Eichwaldi.
coccineum, *Linn. Sp. Pl.* 386.—Afr. bor.; Arab.; Sind.
Commelini, Eckl. & Zeyh. Enum. 97 = sessilifolium, flexuosum.
connaroides, Wight & Arn. ex Hiern, in Hook. f. Fl. Brit. Ind. i. 565 = Heynea trijuga.
cordatum, Burm. f. Fl. Cap. Prod. 12 (Quid?).—Afr. austr.
cordifolium, *Linn. f. Suppl.* 232.—Afr. austr.
cornutum, *Coss. in Bull. Soc. Bot. Fr.* ii. (1855) 364.—Alger.
crenatum, F. Muell. in Linnaea, xxv. (1852) 374 = glaucescens.
cuneifolium, *Eckl. & Zeyh. Enum.* 97.—Afr. austr.
debile, *Cham. in Linnaea*, v. (1830) 45.—Afr. austr.
decumbens, *Delile, Fl. Egypte*, 221. t. 27. f. 3.—Aegypt.
desertorum, Forsk. Fl. Aegypt. Arab. 87. t. 11 = coccineum.
dichotomum, *Licht. ex Cham. in Linnaea*, v. (1830) 48.—Afr. austr.
divaricatum, *Eckl. & Zeyh. Enum.* 97.—Afr. austr.
Dregeanum, Presl, Bot. Bemerk. 30 = simplex.
Dregeanum, *Sond. in Harv. & Sond. Fl. Cap.* i. 365.—Afr. austr.
dumosum, *Boiss. Diagn.* Ser. I. viii. 125.—Arab.
Eichwaldi, *C. A. Mey. in Eichw. Pl. Casp.-Cauc.* 15.—Reg. Casp.
eurypterum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 49.—Syria; Persia; Afghan.
fabagifolium, Baill. Hist. Pl. iv. 417 = fruticulosum.
fabagineum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 137 = seq.
Fabago, *Linn. Sp. Pl.* 385.—Hispan.; Afr. bor.; As. occ.
Fabago, Thunb. Fl. Cap. ed. Schult. 543 = foetidum.
fasciculatum, Licht. ex Cham. in Linnaea, v. (1830) 48, nomen (al. gen.).—Afr. austr.
flexuosum, *Eckl. & Zeyh. Enum.* 97.—Afr. austr.
foetidum, Eckl. & Zeyh. l. c. 96 = debile.
foetidum, E. Mey. in Drège, Zwei Pfl. Docum. 230 = Meyeri.
foetidum, *Schrad. & Wendl. Sert. Hanov.* 17. t. 9.—Afr. austr.
Fontanesii, *Webb & Berth. Phyt. Canar.* i. 17.—Ins. Canar.; Afr. bor.
fruticulosum, *DC. Prod.* i. 705.—Austral.
fulvum, *Linn. Sp. Pl.* 386.—Afr. austr.
fulvum, Mill. Gard. Dict. ed. VIII. n. 4 = sessilifolium.
furcatum, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 106.—Sibir. altaic.
garipense, E. Mey. in Drège, Zwei Pfl. Docum. 230 = microcarpum.
Geslini, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 705.—Alger.
glaucescens, *F. Muell. Pl. Vict.* i. 228.—Austral.
glaucum, *E. Mey. in Drège, Zwei Pfl. Docum.* 230, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 362.—Afr. austr.
glaucum, F. Muell. in Trans. Vict. Inst. i. (1855) 29 = glaucescens.

ZYGOPHYLLUM:—
horridum, Cham. in Linnaea, v. (1830) 46 = microphyllum.
horridum, Eckl. & Zeyh. Enum. 98 = retrofractum.
Howittii, *F. Muell. Fragm.* iii. 150.—Austral.
incrustatum, *E. Mey. in Drège, Zwei Pfl. Docum.* 230, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 362.—Afr. austr.
insuave, Eckl. & Zeyh. Enum. 96 = uitenhagense.
insuave, Sims, Bot. Mag. t. 372 = foetidum.
iodocarpum, *F. Muell. in Linnaea*, xxv. (1852) 372.—Austral.
Karelini, *Fisch. ex Karel. in Bull. Soc. Nat. Mosc.* (1839) 149, nomen; *Boiss. Fl. Orient.* i. 910.—Reg. Casp.
lanatum, Willd. Sp. Pl. ii. 564 = Seetzenia africana?
latifolium, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 198.—Soongar.
Lehmannianum, Bunge, in Arb. Naturf. Ver. Riga, i. II. (1847) 202 = Karelini.
leptopetalum, *E. Mey. in Drège, Zwei Pfl. Docum.* 230, nomen; *et ex Harv. & Sond. Fl. Cap.* i. 363.—Afr. austr.
Lichtensteinianum, *Cham. in Linnaea*, v. (1830) 47.—Afr. austr.
Lichtensteinianum, Eckl. & Zeyh. Enum. 96 = Z. Morgsana, foetidum.
limosum, Eckl. & Zeyh. l. c. = sessilifolium.
macrophyllum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 16.—Turkestan.
macropterum, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 102.—Persia; Soongar.
maculatum, *Ait. Hort. Kew.* ed. I. ii. 60.—Afr. austr.
maritimum, Eckl. & Zeyh. Enum. 96 = fulvum, uitenhagense.
Melongena, *Bunge, in Ledeb. Fl. Alt.* ii. 104.—Sibir. altaic.
Meyeri, *Sond. in Harv. & Sond. Fl. Cap.* i. 364.—Afr. austr.
microcarpum, Eckl. & Zeyh. Enum. 98 = retrofractum.
microcarpum, *Licht. ex Cham. in Linnaea*, v. (1830) 46.—Afr. austr.
microcarpum, E. Mey. in Drège, Zwei Pfl. Docum. 230 = simplex.
micronatum, Opiz, Natural. n. vi. (1824) 42 = Z. Fabago?
microphyllum, G. Don, Gen. Syst. i. 771, in syn., sphalm. = microcarpum.
microphyllum, Eckl. & Zeyh. Enum. 98 = simplex.
microphyllum, *Linn. f. Suppl.* 232.—Afr. austr.
miniatum, *Cham. in Linnaea*, v. (1830) 49.—Turkestan.
Morgsana, *Linn. Sp. Pl.* 385.—Afr. austr.
mucronatum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 438.—China.
Nitraria, Linn. in Amoen. Acad. iii. 409 = Nitraria Schoberi?
Novae-Zeelandiae, Steud. Nom. ed. II. ii. 802, sphalm. = *Zanthoxylum Novae-Zeelandiae*?
obtusum, Vicary, in Proc. Linn. Soc. i. (1848) 367 = simplex.
orbiculatum, *Welw. ex Oliver, Fl. Trop. Afr.* i. 285.—Afr. trop.
ovigerum, *Fisch. & Mey. ex Karel. in Bull. Soc. Nat. Mosc.* (1839) 142, nomen; *Boiss. Fl. Orient.* i. 914.—Reg. Casp.
parvifolium, G. Don, Gen. Syst. i. 771 = microcarpum.
pinnatum, Cham. in Linnaea, v. (1830) 48 = macropterum.
portulacoides, Cham. l. c. 50 = Miltianthus portulacoides.
portulacoides, Forsk. Fl. Aegypt. Arab. 88 = simplex.
Potanini, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 438.—Soongar.
prismaticum, E. Mey. in Drège, Zwei Pfl. Docum. 230, nomen = prismatocarpum.
prismatocarpum, *Sond. in Harv. & Sond. Fl. Cap.* i. 357.—Afr. austr.
prismatothecum, *F. Muell. in Linnaea*, xxv. (1852) 375.—Austral.
proliferum, Forsk. Fl. Aegypt. Arab. 87 = album.
propinquum, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 283 = coccineum.
prostratum, E. Mey. in Drège, Zwei Pfl. Docum. 230 = microcarpum.
prostratum, Thunb. Prod. Pl. Cap. 189 = Seetzenia africana.
pterocarpum, *Bunge, in Ledeb. Fl. Alt.* ii. 103.—Sibir. altaic.

ZYGOPHYLLUM:—
pygmaeum, *Eckl. & Zeyh. Enum.* 97.—Afr. austr.
Retama, *Gill. ex Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 166.—Chili.
retrofractum, Eckl. & Zeyh. Enum. 96 = foetidum.
retrofractum, Jacq. Hort. Schoenb. t. 354 = flexuosum.
retrofractum, *Thunb. Prod. Pl. Cap.* 80.—Afr. austr.
rigescens, F. Mey. in Drège, Zwei Pfl. Docum. 230 = cuneifolium.
rigidum, Royle, Illustr. Bot. Himal. 154 (Quid?).—Ind. bor. occ.
Rosowii, *Bunge, in Linnaea*, xvii. (1843) 5.—Mongol.
semiteres, Eckl. & Zeyh. Enum. 97 = fulvum.
sessilifolium, *Linn. Sp. Pl.* 385.—Afr. austr.
sessilifolium, Sims, Bot. Mag. t. 2184 = fulvum.
Sieversianum, Steph. ex Ledeb. Fl. Ross. i. 484 = macropterum.
simplex, *Linn. Mant.* i. 68.—Afr. bor. et austr.; As. occ.
spatulatum, *D. Don, Gen. Syst.* i. 771.—Afr. austr.
spinosum, *Linn. Sp. Pl.* 386.—Afr. austr.
stellulatum, *C. Sm. in Tuckey, Congo*, 250, nomen.—Ins. Cap. Virid.
steropterum, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 308.—Soongar.
subtrijugum, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 105.—Sibir. altaic.
succulentum, Salisb. Prod. 319 = Z. Morgsana.
terminale, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 437 = apiculatum.

ZYGOPHYLLUM:—
Thunbergianum, Eckl. & Zeyh. Enum. 97 = Lichtensteinianum.
tridentatum, [Moç. & Sesse,] ex DC. Prod. i. 706 = Larrea mexicana.
turcomanicum, *Fisch. ex Karel. in Bull. Soc. Nat. Mosc.* (1839) 149, nomen; *Boiss. Fl. Orient.* i. 914.—Reg. Casp.
uitenhagense, *Sond. in Harv. & Sond. Fl. Cap.* i. 361.—Afr. austr.
Webbianum, Coss. in Bull. Soc. Bot. Fr. ii. (1855) 365, in obs. = Fontanesii.

ZYGOSEPALUM, Reichb. f. in Walp. Ann. vi. 665 (1863) = **Zygopetalum**, Hook. (Orchid.).
Kegelii, Reichb. f. l. c. 666 = Zygopetalum Kegelii.
rostratum, Reichb. f. l. c. = Zygopetalum rostratum.

ZYGOSPERMUM, Thw. ex Baill. Etud. Gén. Euph. 620. t. 27 (1858) = **Phyllanthus**, Linn. (Euphorb.).
zeylanicum, Thw. ex Baill. l. c. = P. cyanospermus.

ZYGOSTATES, Lindl. Bot. Reg. sub t. 1927 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 569.
DACTYLOSTYLES, Scheidw. in Otto & Dietr. Allg. Gartenz. vii. (1839) 405.
cornuta, *Lindl. Bot. Reg.* sub t. 1927.—Bras.
Greeniana, *Reichb. f. in Gard. Chron.* (1869) 988.—Bras.
lunata, *Lindl. Bot. Reg.* sub t. 1927.—Bras.
pellucida, *Reichb. f. in Walp. Ann.* vi. 564.—Bras.

ZYGOSTELMA, Benth. in Benth. & Hook. f. Gen. ii. 740 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. l. c.

ZYGOSTELMA, Fourn. in Mart. Fl. Bras. vi. IV. 232 (1885). *ASCLEPIADEAE*.
calcaratum, *Fourn. l. c.*—Bras.

ZYGOSTIGMA, Griseb. Gen. et Sp. Gent. 150 (1839). *GENTIANEAE*, Benth. & Hook. f. ii. 810.
australe, *Griseb. l. c.*—Bras.
uniflorum, *Griseb. l. c.* 151.—Reg. Argent.

ZYMUM, Noronha, ex Thou. Hist. Vég. Isles Afr. Austr. 69. t. 23 (1806) = **Tristellateia**, Thou. (Malpig.).
madagascarense, Juss. ex Spreng. Syst. ii. 329 = T. madagascariensis.
tristellatum, Noronha, ex Thou. Hist. Vég. Isles Afr. Austr. 69. t. 23 = T. madagascariensis.

ZYRPHELIS, Cass. in Ann. Sc. Nat. Ser. I. xvii. (1829) 420 = **Mairia**, Nees (Compos.).
amoena, Cass. l. c. 421 = M. taxifolia.
microcephala, Nees, Gen. et Sp. Aster. 252 = M. microcephala.
taxifolia, Nees, l. c. 250 = M. taxifolia.

ZYZOPHYLLUM, Salisb. Prod. 319 (1796), err. typ. = **Zygophyllum**, Linn. (Rutac.).

ZYZYGIUM, Brongn. Enum. Genr. 123 (1843), sphalm. = SYZYGIUM, Gaertn. = **Eugenia**, Mich. (Myrtac.).

APPENDIX

ADDENDA ET EMENDANDA GRAVIORA

HACTENUS NOTATA

VOLUME I

(MENDA NON COMMEMORATA LECTOR BENEVOLENS IPSE CORRIGET)

a. = *adde*. d. = *dele*. l. = *lege*. p. = *pone*.

PAG.

ABELMOSCHUS.
1³. l. *aculeatus*, Walp. . . . = H. surattensis, cannabina.
angustifolius, Walp. Rep. i. 308 = H. angustifolius.
cryptocarpus, Walp. . . . = H. Abelmoschus.
cubensis, Walp. . . . = H. Abelmoschus.

ABIES.
2³. l. *Khutrow*, Royle, Illustr. Bot. Himal. 353; Loud. etc.
3¹. Pindrow, *Royle* (*l. c.*) 350. t. 86; *Spach*, etc.

ABILDGAARDIA.
3². l. *polygama*, Spreng. . . . = Pharus angustifolius.

ABRONIA.
3³. a. TRIPTEROCALYX, Hook. Kew Journ. v. (1853) 261.

ABRUS.
4¹. a. HULTHENIA, Blume, ex Miq. Fl. Ind. Bat. i. 1. 160 (1855).

ABSINTHIUM.
4². l. *camphoratum*, Roehl. Deutschl. Fl. ed. II. ii. 451; Bess. etc.
grandiflorum, Roehl. (l. c.) = *Mutellina*, Steud.
a. *mutellinum*, Roehl. (l. c.) 450 =

ABUTILON.
4³. a. BOGENHARDIA, Reichb. Nom. 200 (1841).
asiaticum, *G. Don, Gen. Syst.* i. 503.—Reg. trop.
5¹. l. australe, *Presl, Bot. Bemerk.* 20; *Garcke*, etc.
a. Berlandieri, *A. Gray, in Proc. Am. Acad.* xx. (1885) 358, in obs.—Mexic.
l. ceratocarpum, *C. Gay, Fl. Chil.* i. 331.—Chili.
depauperatum, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 230.—Ins. Galapag.
5³. l. *oxyphyllum*, Presl, Bot. Bemerk. 20; Edgew. etc.

ACACIA.
6¹. a. PHYLLODOCE, Link, Handb. ii. 132 (1831).
6². a. adenanthera, *Zeyh. ex Steud. Nom.* ed. I. 532. —Hab.?
7¹. a. *biglobosa*, Royle, Illustr. Bot. Himal. 182 = Parkia Roxburghii, africana.
l. *brevipes*, A. Cunn. . . . t. 3358, etc.
7². l. *Cebil*, Griseb. . . . 136 = Piptadenia Cebil.
8¹. l. *decussata*, Tenore, . . . (1845) 77, etc.
8³. l. *gigantea*, Noronha, etc.

PAG.

ACACIA:—
8³. a. *Kaeringa*, Royle, Illustr. Bot Himal. 153 = A. leucophyloea.
10². post lineam 49 insere, *melanoceroides*, Beurl. l. c. = praec.?
12². a. *stipulacea*, Royle, (l. c.) (Quid?).—Ind. or.
13¹. *tamariscifolia*, Steud. Nom. ed. II. ii. 149, sphalm. = tamarindifolia.
l. *tomentella*, Zipp. etc.
tortilis, Hayne, Arzn. Gew. x.
a. post lin. 68 insere, *tortuosa*, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 124 = Prosopis dulcis.

ACALYPHA.
14³. l. *boehmerioides*, Miq. . . . 459 = fallax.
15¹. *Commersoniana*, Baill. . . . = integrifolia.
15³. *hispida*, Blume, . . . = fallax.
indica, Hort. ex Steud. Nom. ed. II. i. 10; Herb. Linn. ex Muell. Arg. in DC. Prod. xv. II. 872 = fallax.
leptostachya, H. B. & K. etc.
16¹. a. *Macafeeana*, Hort.—Cf. Gard. Chron. (1878) I. 10 = Wilkesiana.
musaica, Hort. (l. c.) (1877) I. 520; (1878) I. 10 = Wilkesiana.
17¹. *tenera*, Royle, Illustr. Bot. Himal. 327 (Quid?). —Reg. Himal.

17². l. ACAMPTOPAPPUS . . . . = **Aplopappus**, Cass. (Compos.).

ACANTHODIUM.
17³. l. *grossum*, Nees, . . . 274 = B. grossa.
hirtum, Hochst. . . . = B. linearifolia.

ACANTHOLIPPIA.
18². l. *salsoloides*, Griseb. . . . = L. salsoloides.

ACANTHOMINTHA.
l. lanceolata, *Curran*, . . . i. 1. (1884), etc.

ACANTHOPHOENIX.
18³. l. crinita, *H. Wendl.* etc.
rubra, *H. Wendl.* etc.

19³. l. ACANTHYLLIS, Pomel.

ACARNA, Boehm. in Ludw. Def. Gen. Pl. 195 (1760); All. etc.
gummifera, Willd. . . . = Atractylis gummifera.
macrocephala, Willd. . . . = Atr. macrocephala.
macrophylla, Willd. . . . = Atr. macrophylla.

PAG.

ACER.
20². a. *lactescens*, Steud. Nom. ed. II. i. 11 = platanoides.
Leopoldi, Hort. ex C. Koch, Dendrol. i. 528 = A. Pseudo-platanus.
liburnicum, Hort. ex C. Koch, (l. c.) 539 = monspessulanum.
20³. *lobatum*, Hort. ex Steud. Nom. ed. II. ii. 11 = platanoides.
l. *lobatum*, Rafin. . . . 48 = Negundo aceroides?
a. *microphyllum*, Hort. ex C. Koch, Dendrol. i. 536 = italum.
nanum, Steud. Nom. ed. II. i. 11 = campestre.
Pavia, Hort. ex C. Koch, Dendrol. i. 541 = dasycarpum.
21¹. *pulverulentum*, Hort. ex C. Koch, (l. c.) 535 = campestre.
splendens, Hort. ex C. Koch, (l. c.) 543 = rubrum.
tomentosum, Hort. ex Steud. Nom. ed. II. i. 12 = dasycarpum.
l. *trifoliatum*, Rafin. . . . 48 = Negundo aceroides?

ACERAS.
21². l. *angustifolia*, Lindl. . . . = Herminium longicrure.
longicruris, C. Wright, . . . = Herminium longicrure.

ACETOSA.
21³. a. *arvensis*, Montand. Guide Bot. 269 = R. Acetosella.

ACHAETA.
l. *geniculata*, Fourn. . . . 553 = Koehleria cristata.

ACHILLEA.
22¹. l. alpina, *Linn.* . . . 899.—Europ.
22³. a. *firma*, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 430 (Quid?).—Gallia.
23³. l. pectinata, *Willd.* etc.
24¹. *sylvatica*, Tenore, Ind. Sem. Hort. Neap. (1827); Fl. Nap. etc.

ACHIMENES.
24². a. *alba*, Hort. Padov. ex Walp. Ann. ii. 1067 = Niphaea oblonga.
atrosanguinea, Hort. ex Regel, in Flora, xxxii. (1849) 180 = amoena.
l. magnifica, *Lindl.* . . . 116.—Am. centr.
a. *Kleei*, E. Otto, ex Regel, in Flora, xxxii. (1849) 182 = hirsuta.
24³. l. *picta*, Benth. . . . t. 4126 = Isoloma picta.
viscida, Lindl. & Paxt. Flow. Gard. ii. (1851) 59 = Heppiella atrosanguinea.

PAG.

ACHRAS.

25¹. l. *Caimito*, Ruiz & Pav. . . . 18 = Lucuma Caimito.
a. *occidentalis*, Hort. Cels, ex Tenore, Cat. Ort. Nap. (1845) 77 (Quid ?).—Hab. ?
l. *Pohlmaniana*, F. Muell. . . . 184 = Sideroxylon Pohlmanianum.
a. *Tchicomame*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 95 = A. Sapota.

ACHYRANTHES.

25². d. CODIVALIA, et ECLOTORIPA, Rafin. (nunc Pupalia).

ACHYROCLINE.

26³. l. *satureoides*, Gardn. . . . = Stenocline satureoides.

26³. l. ACHYROCHOMA, Cass. in Dict. Sc. Nat. xxvi. 21 (1823) ; lvii. 341 (1828) =

l. ACHYROMA, Wendl. Bot. Beobacht. ; etc.

ACHYROPAPPUS.

26³. a. *ambiguus*, Steud. Nom. ed. II. i. 437, in syn. = *Achyrophorus ambiguus*.

ACHYROPHORUS.

27¹. l. *albiflorus*, Sch. Bip.
cryptocephalus, Sch. Bip. . . . 57 = Werneria ambigua.
a. *Gayii*, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 116 = H. chrysantha.
d. *helveticus*, Less.
l. *Hohenackeri*, Sch. Bip. (l. c.) 57 = Werneria ambigua.

27². a. *taraxacoides*, Steud. Nom. ed. II. ii. 226, in syn. = H. Scorzonerae.
l. *taraxacoides*, Walp. (l. c.) 336 = H. taraxacoides.
uniflorus, F. W. Schmidt, . . . Aufs.

27³. l. **ACICALYPTUS.**

ACIDANTHERA.

27³. a. SOLENANTHUS, Steud. teste Klatt, ex Baker, in Journ. Linn. Soc. xvi. (1877) 159.
capensis, Benth. & Hook. f. Gen. iii. 706 = Tritonia capensis.

28¹. a. *pauciflora*, Benth. & Hook. f. (l. c.) = Tritonia pauciflora.

28¹. l. ACINODENDRUM, Linn. . . . = **Micania**, Ruiz & Pav. (1794) (Melastom.).

ACINOS.

28². a. *Siso*, Siebold, in Verh. Batav. Gen. xii. (1830) 32 = Perilla arguta ?

ACIPHYLLA.

28³. l. GINGIDIUM, F. Muell. in Trans. Phil. Soc. etc. quoad glacialis, procumbens, simplicifolia, dele *F. Muell. ex*

ACMENA.

29³. a. *pendula*, G. Benn. Gather. Natural. 327, nomen.—Austral.

ACOKANTHERA.

30¹. a. oblongifolia, *Benth. & Hook. f. Gen.* ii. 696.—Afr. trop.
Schimperi, *Benth. & Hook. f.* (*l. c.*).—Abyssin.

30¹. a. ACOME, Baker, in Journ. Bot. xx. (1882) 18, sphalm. = **Cleome**, Linn. (Cappar.).

ACORUS.

32¹. a. CALAMUS, Auct. ex Linn. Gen. ed. I. 104 (1737).
l. *cochinchinensis*, Schott, . . . 22 = A. Calamus.

32¹. l. ACOSTA, Lour. . . . = **Vaccinium**, Linn.

32³. insere, ACRIVIOLA, Boerh. ex Ludw. Def. Gen. Pl. 192 (1747) = **Tropaeolum**, Linn.

ACRIDOCARPUS.

32³. l. orientalis, *A. Juss.* . . .—Arab.

ACRODICLIDIUM.

33². l. *Ita-uba*, Meissn. . . . = Endiandra Ita-uba.

PAG.

ACROELYTRUM.

33². a. *Urvillei*, Steud. ex Miq. Fl. Ind. Bat. iii. 400 = L. gracile.

33³. l. ACROLOBUS, Klotzsch, in Abh. etc.

34¹. l. ACROSSANTHES, Presl, Bot. Bemerk. 22 (1844) = **Vismia**, Vand. (Hypericin.).

34¹. l. ACROTAPHROS, Steud. ex Hochst. in Flora, xxix. (1846) 599 ; et ex A. Rich. etc.

ACROTREMA.

34². l. *Wightianum*, Wall. . . . = Arnottianum.
a. *Wightianum*, Wight & Arn. Prod. 6 = costatum.

ACTAEA.

34². l. *dahurica*, Turcz. . . . = Cimicifuga dahurica (in Addend.).
davurica, Franch. . . . = praec.

34³. *pterosperma*, Fisch. & Mey. . . . = Cimicifuga dahurica (in Addend.).
racemosa, Walt. ex Pursh, Fl. Am. Sept. ii. 373, etc.

ACTINELLA.

35¹. a. *californica*, Benth. & Hook. f. Gen. ii. 415, in nota = Baeria coronaria.
l. *Richardsonii*, A. Gray, in Bot. Calif. i. 394, partim = biennis.
Richardsonii, *Nutt.* . . . 379.—Am. bor. occ.

ACTINOLEPIS.

35³. l. *anthemoides*, A. Gray, . . . 378 = Baeria coronaria.
coronaria, A. Gray, . . . 197 = Baeria coronaria.
lanosa, A. Gray, l. c. = Eriophyllum lanosum.
mutica, A. Gray, . . . 197 = Baeria mutica.
tenella, A. Gray, l. c. = Baeria tenella.
Wallacei, A. Gray, l. c. = Eriophyllum Wallacei.

ACTINOMERIS.

35³. l. *alternifolia*, DC. . . . 575 = *helianthoides*, Nutt. . . . 181 = Verbesina helianthoides.

36¹. *Wrightii*, A. Gray, . . . 85 = Verbesina Wrightii.

36¹. post lineam 19 insere

ACTINOPHLOEUS, Becc. in Ann. Jard. Buitenz. ii. (1885) 126, in obs. *PALMAE.*
ambiguus, *Becc. l. c.*—N. Guinea.
propinquus, *Becc. l. c.*—Ins. Aru.

ACTINOPHYLLUM.

36¹. l. *pedicellatum*, Pav. . . . 195, etc.

ACTINOSPORA.

36¹. l. *dahurica*, Turcz. . . . = C. dahurica.
pterosperma, Walp. . . . = C. dahurica.

ACTINOSTEMON.

36². l. *acuminatus*, Baill. . . . = caribaeus.
angustifolius, Klotzsch, . . . = caribaeus.

36². l. ACTINOSTIGMA, Welw. . . . [1859] = **Symphonia**, Linn. f. (Guttif.).

ACTINOTUS.

36². l. leucocephalus, *Benth.* etc.

ADELOSTIGMA.

37³. a. CANCELLARIA, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 3[illegible] (1877).

ADENANDRA.

37³. l. GLANDULIFOLIA, Wendl. Collect. 35. t. 10 (1808).

38². a. ADENOBIUM, Steud. Nom. ed. II. ii. 346 (1841), sphalm. = ADENOBASIUM, Presl.

ADENOGYRUS.

39². l. *Braunii*, Klotzsch, ll. cc. (= *Phoberos Mundtii*). —Afr. austr.

ADENONCOS.

39³. l. *virens*, Blume, etc.

PAG.

ADENOPHYLLUM.

40¹. a. *Cavanillesii*, Steud. Nom. ed. II. i. 534, sphalm. = coccineum.

40¹. a. ADENOPOGON, Welw. Synopse, 27 (1862) = **Swertia**, Linn. (Gentian.).
stellaroides, Welw. l. c.—Afr. trop.

ADENOROPIUM.

40². a. *coccinea*, Steud. Nom. ed. II. i. 799, in syn. = J. coccinea.

d. **ADENOSTEMMA**, Forst.

41¹. d. *glandulosum*, Benth. (vide infra).

a. ADENOSTEMMA, Hook. f. Fl. Brit. Ind. i. 242 (1874), sphalm. = ODONTOSTEMMA, Benth. = **Arenaria**, Rupp. (Caryoph.).
glandulosum, Hook. f. l. c. = A. Benthami.

ADENOSTEPHANUS.

41¹. l. *austro-caledonicus*, Brongn. & Gris, . . . 229 = Kermadecia austro-caledonica.
Bleasdalii, Benth. . . . 417 = Roupala Bleasdalii.
incanus, Klotzsch, etc.
laxiflorus, Meissn. etc.

ADESMIA.

41². a. HETEROLOMA, Desv. ex Reichb. Consp. 152 (1828).

42¹. l. glabriuscula, *Vog.* . . . 592.—Chili.

42². *radicans*, Bert. . . . l. c. = glabriuscula.

ADHATODA.

42³. l. *chrysotrichoma*, Nees, . . . 408 = Justicia chrysotrichoma.

43¹. l. *Gilliesii*, Nees, l. c. 395 = Justicia Gilliesii.
orchioides, Nees, l. c. 393 = Justicia orchioides.

43². *Tweediana*, Nees, l. c. 395 = Justicia Tweediana.

ADONIS.

44². a. *Walziana* ×, Simk. in Magyar Növényt. Lapok, ii. (1878) 151.—Hungar.

ADRIANA.

44³. a. MEIALISA, Rafin. Sylva Tellur. 63 (1838).

44³. l. ADUPLA, Bosc, . . . (1805) = **Mariscus**, Ehrh. (Cyperac.).

AECHMEA.

45¹. a. CHEVALIERIA, Gaudich. Bot. Voy. Bonite, tt. 61, 62 (1852 ?).

AEGOPOGON.

47². a. ATHEROPHORA, Willd. ex Steud. Nom. ed. II. i. 167 (1840).

AELUROPUS.

47². a. *niloticus*, Edgew. ex Aitch. Cat. Punj. Pl. 169, sphalm. = niliacus.

48¹. l. **AËRANTHES**, . . . *ORCHIDEAE*, Benth. & Hook. f. iii. 576.
Arachnitis, *Lindl.* . . . sub t. 817, etc.
filiformis, *Griseb.* etc.

AËRANTHUS.

48¹. l. *jamaicensis*, Reichb. f. . . . = Angraecum jamaicense.

AËRIDES.

48³. l. *latifolium*, Thw. . . . 429 = Doritis latifolia.

AERVA.

49². l. *brachiata*, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 291; etc.
incana, Mart. (l. c.).
Monsonia, *Mart.* (*l. c.*).
pubescens, *Mart.* (*l. c.*).
setacea, Mart. (l. c.).

AESCHYNOMENE.

50². a. *aculeata*, Royle, Illustr. Bot. Himal. 191, nomen (Quid ?).—Ind. or.
cassioides, *Desv. in Mém. Soc. Linn. Par.* iv. (1826) 327.—Guian.

50³. falcata, *Desv.* (*l. c.*) 323.—Ind. or.

PAG.

AESCHYNOMENE:—
51¹. microphylla, *Desv.* (*l. c.*).—Ind. or.
l. *miniata*, Ortega, . . . = Piscidia Sesbania.
a. monosperma, *Desv.* (*l. c.*).—Bras.
muricata, Desv. (l. c.) 326 = Adesmia muricata.
onobrychifolia, Desv. (l. c.) = Adesmia bicolor.
l. *patagonica*, Hort. ex Link, Enum. Hort. Berol. ii. 249; etc.
a. *pimpinellaefolia*, Desv. (l. c.) = Adesmia muricata.
l. *platycarpa*, Michx. . . . = Sesbania platycarpa.

AESCULUS.
51². l. Pavia, Boerh. ex Linn. Syst. ed. I (1735); etc.
51³. a. *octandra*, Marsh. Arbust. 4 = flava.

AETHIONEMA.
52¹. l. Diastrophis (non diastrophis).

52². a. AETHULIA, A. Gray, Syn. Fl. i. II. 92, in syn. (1884), sphalm. = **Ethulia**, Linn. (Compos.).

AFZELIA.
52³. l. Intsia, Thou. . . . (1806).
a. Macrolobium, Zipp. ex Miq. Fl. Ind. Bat. i. I. 80 (1855).

AGANISEA.
53¹. a. *cinerea*, Benth. in Journ. Linn. Soc. xviii. (1880) 321; Benth. & Hook. f. Gen. iii. 544, sphalm. = cyanea.
cyanea, *Benth. & Hook. f. Gen.* iii. 545.—N. Granat.
graminea, Benth. & Hook. f. l. c. = Zygopetalum gramineum.

53³. l. AGARISTA, DC. . . . (1836) = **Leptosyne**, DC. (Compos.).
calliopsidea, DC. l. c. = L. calliopsidea.

AGARISTA.
53³. l. *eriophylla*, G. Don, l. c. = Gaultheria ferruginea.

54¹. a. AGASTACHYS, Ehrh. Beitr. iv. 146 (1789) = **Carex**, Dill. (Cyperac.).

AGATHAEA.
54¹. l. *spathulata*, Fisch. . . . Sem. Hort. Petrop. etc.

AGATHOSMA.
54². l. acerosa, *Eckl. & Zeyl.*
54³. apiculata, *G. F. W. Mey.*
bifida, *Bartl. & Wendl. f.* . . .—Afr. austr.
55¹. *hybrida*, Bartl. & Wendl. f. . . . = pubescens.
mollis, Bartl. & Wendl. f. . . . = pubescens.

AGAVE.
57². a. sebastiana, *Greene, in Bull. Calif. Acad.* i. III. (1885) 214.—Calif.

AGERATUM.
58¹. l. domingense, *Spreng.*

58³. l. AGGEIANTHUS, Wight,

AGLAIA.
59¹. l. *inequalis*, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1885) 409;
59². *Sipannas*, Miq. . . . Suppl. 506.

AGLAODORUM.
59². l. *Griffithii*, . . . = Aglaonema Griffithii.

59². l. **AGLAONEMA**, Schott, in Wien. Zeitschr. iii. (1829) 892; ex Endl. in Linnaea, vi. (1831) Litt. 53.
l. *acutispathum*, N. E. Br. . . . 39 = modestum.
a. Griffithii, *Schott, Syn. Aroid.* 123.—Malacca.
l. *integrifolium*, Schott, . . . 20 = oblongifolium.
marantifolium, Blume, . . . 66 = oblongifolium.
nitidum, Kunth, . . . 56 = oblongifolium.
oblongifolium, *Schott, in Wien. Zeitschr.* iii. (1829) 892; *ex Endl. in Linnaea*, vi. (1831) *Litt.* 53.—Malaya.
palustre, Teijsm. & Binn. . . . 399 = Griffithii.

AGONIS.
59³. l. glabra, *Turcz. in Bull. Phys. Math.* etc.

AGRIMONIA.
60¹. l. *Eupatoria*, Thw.
60². *javanica*, Jungh. Java, i. 664; et ex Miq. etc.
zeylanica, Moon, Cat. Pl. Ceyl. 37 = Alchemilla indica.

PAG.

AGROPYRON.
60³. a. Cynopoa, Ehrh. Phytophyl. n. 72 (1780); Beitr. iv. (1789) 148.
Kratzmannia, Opiz, Seznam, 56 (1852).
l. attenuatum, *Roem. & Schult.*
Caldesii, *Goiran*, etc.
campestre, *Gren. & Godr.*
61¹. a. Goiranicum, *Vis. ex Goiran, Pl. Vasc. Nov. Veron.* (1874) 21.—Italia.
l. Roegnerii, *Boiss.* . . . (Raegnerii).
61². a. *trachycaulon*, Hort. ex Steud. Syn. Pl. Gram. 344 = Richardsoni.
l. *violaceum*, Vasey, . . . = Richardsoni.

AGROSTEMMA.
61². a. *infesta*, Salisb. Prod. 304 = L. Agrostemma.

AGROSTIS.
61³. a. *abyssinica*, Ehrenb. & Hempr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 54 = Sporobolus robustus.
aequalis, Ehrenb. & Hempr. (l. c.) 52 (= *Vilfa helvola*).—Arab.
l. *altissima*, Tuckerm.
62¹. l. brachystachys, *Schult.* . . . ii.
62². Carmichaeli, *Schult.* . . . ii.
62³. a. collina, *Schleich. Cat. Pl. Helv.* ed. IV. 6 (Quid?).—Helv.
l. composita, *Poir. Encyc. Suppl.*
congesta, *Schum. & Thonn. Beskr. Guin. Pl.* 46.—Afr. trop.
l. *crinita*, R. Br. . . . = Dichelachne crinita.
cylindrica, R. Br. . . . = Deyeuxia cylindrica.
63¹. *divaricata*, Hort. ex DC. Cat. Hort. Monsp. (1813) 93, etc.
a. *elatior*, Ehrenb. & Hempr. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 55 (= *Vilfa setulosa*).—Arab.
extensa, *Schum. & Thonn. Beskr. Guin. Pl.* 45.—Afr. trop.
63². l. *glutinosa*, Fisch. ex Schrank, Pl. Rar. Hort. Monac. t. 58, etc.
63³. *koelerioides*, E. Desv. . . . 317 = Deyeuxia anomala?
64¹. a. *Novae-Angliae*, Tuckerm. in Am. Journ. Sc. Ser. I. xlv. (1843) 44 = elata.
64². l. *patula*, Gaud. in Alpina, iii. (1808) 14;
64³. *purpurea*, Gaud. (l. c.) 11;
Royleana, Trin. . . . 125 = Deyeuxia Royleana (in Addend.).
65¹. l. *sciurea*, R. Br. . . . = Dichelachne sciurea.
sesquiflora, E. Desv. . . . 318 = Deyeuxia sesquiflora.
a. *sesquivalvis*, Benth. in Journ. Linn. Soc. xix. (1881) 91, sphalm. = prae
65². a. *subspicata*, Arv. Ess. Pl. Da[illegible]ph. 67 (Quid?).—Gallia.
65³. l. *zeylanica*, Klein, . . . 83 = Panicum sparsicomum.

AICHRYSON.
66². l. *pulchellum*, C. A. Mey. . . . 55 = S. pulchellum.

AILANTHUS.
66³. a. *simplex*, Panch. ex Brongn. & Gris, in Ann. Sc. Nat. Sér. V. iii. (186[illegible]) 231 = Soulamea Pancheri.

AINSLIAEA.
66³. l. *glumacea*, Sch. Bip. . . .; Klatt, in Sitzb. Akad. Muench. (1878) I. 97, etc.

AIRA.
67¹. a. Caryophyllea, Opiz, Seznam, 27 (1852).
d. Periballia, Trin. (*nunc* **Deschampsia**).
a. Salmasia, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317.
67². l. *atropurpurea*, Wahlenb. . . . 37 = Deschampsia atropurpurea.
a. *brachyphylla*, Steud. Nom. ed. II. i. 766, in syn. = Sporobolus tenellus.
l. *chinensis*, Retz. . . . 10 = Panicum megalanthum.
67³. *distichophylla*, Spreng. Nov. Prov. 47.
filiformis, *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 326.
68¹. *hirsuta*, Schleich. ex Gaud. in Alpina, iii. (1808) 48.
Kingii, Hook. f. . . . t. 13[illegible]5.

PAG.

AIRA:—
68². *neutralis*, Walp. Ann. i. 932, sphalm. = *australis*, Raoul.
68³. *spicata*, Linn. . . . 63 = Panicum indicum.
spicata, Linn. . . . 64 = Trisetum subspicatum.
subtriflora, Lag. . . . ii. IV. (1805).
a. *subviolaceum*, Ehrenb. ex Trin. in Mém. Acad. Pétersb. Sér. VI. v. II. (1840) 84 = Sporobolus minutus.
l. *viatica*, Griff. . . . 54 = Panicum semialatum.

AJUGA.
69³. l. Bugula, Tourn. ex Linn. Syst. ed. I (1735).
a. Bulga, Linn. l. c. (1735).
70¹. a. *caespitosa*, Schleich. Cat. Pl. Helv. ed. IV. 6 = pyramidalis.
l. *Chamaepitys*, Guss.
Chamaepitys, *Schreb.*
chia, *Schreb.*
humilis, Miq. . . . 114 = Lamium humile.
a. *laxa*, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 109 = Teucrium laxum.
70². *microphylla*, Hegetschw. Fl. Schw. 561 = reptans?
l. *rupestris*, Schleich. Cat. Pl. Helv. ed. IV. 6 = pyramidalis.

AKENTRA.
70³. l. *inflata*, Benj. l. c. = U. Benjaminiana.

ALAFIA.
70³. a. landolphioides, *Benth. & Hook. f. Gen.* ii. 708, in obs.—Afr. trop.

ALANGIUM.
70³. l. *cordifolium*, Zoll. & Mor. . . . 63 = Marlea virgata.

ALATERNUS.
71¹. l. *sempervirens*, Roehl. Deutschl. Fl. ed. II. ii. 176 =

ALBERTA.
71¹. a. isosepala, *Baker, in Journ. Bot.* xix. (1882) 138.—Madag.
l. *laurifolia*, Baker, . . . 271 = minor.

71². l. ALBINEA, Hombr. & Jacquinot, Bot. Voy. Astrol. et Zél. Dicot. Phan. t. 4 (1845); et ex Decne. etc.
oresigenesa, . . . = P. criniferum.

ALBUCA.
72². l. Virdika, Adans.

ALBUCEA.
72³. a. *neapolitana*, Montand. Guide Bot. 315 = O. nutans.

ALCHEMILLA.
73². a. *splendens* ×, Christ, ex Gremli, Excurs. Fl. Schw. ed. II. 179.—Helvet.

73². a. ALCHIMILLA, Tourn. ex Hall. Enum. Stirp. Helv. i. 184 (1742) = praec.

74². a. ALEPYRUM, Hieron. Centrolep. 103 (1873) = **Gaimardia**, Gaudich. (Centrol.).

ALISMA.
75³. d. Baldellia, Parl. (*nunc* **Echinodorus**).
Nectalisma, Fourr. (*nunc* **Elisma**).

ALKANNA.
76³. l. syriaca, *Boiss. & Hohen.*

ALLARDIA.
77². l. incana, *Klatt*, . . . (1878) I. 88.

ALLIUM.
77³. d. Periloba, Rafin. . . . 87 (*nunc* **Nothoscordum**).
78¹. l. atropurpureum, *Waldst. & Kit.* . . . 17.—Hungar.; Reg. Cauc.; As. centr.
78². *borbonicum*, Regel, . . . 218 = Nothoscordum fragrans.
a. *calcareum*, Montand. Guide Bot. 317 = ochroleucum.
79¹. l. Cyrilli, *Tenore*, . . . Graecia;

PAG.

ALLIUM:—

79³. *Gardinia*, Bert. . . . 664 (= *Triteleia violacea*). —Chili.
a. *graminifolium*, Montand. Guide Bot. 320 = rotundum.
80¹. l. Grayi, *Regel*,
80². a. *lilacinum*, Royle, Illustr. Bot. Himal. 392 = rubens.
longistamineum, *Royle* (*l. c.*) (Quid?).— Reg. Himal.
80³. l. *montanum*, F. W. Schmidt, . . . = angulosum.
81². a. *ovatum*, Arv. Ess. Pl. Dauph. 65 (Quid?).— Gallia.
81³. *pratense*, Montand. Guide Bot. 317 = fallax.
Rakkjoo, Siebold, in Verh. Batav. Gen. xii. (1830) 16 = Bakeri.
l. reticulatum, [*Nutt. in*] *Fras. Cat.* (1813) n. 1, nomen; et
82¹. a. *rubrifolium*, Arv. Ess. Pl. Dauph. 65 (Quid?).— Gallia.
82². l. stellatum, [*Nutt. in*] *Fras. Cat.* (1813) n. 2; et

ALLMANIA.

83². a. BELUTTA, Rafin. Fl. Tellur. iii. 39 (1836).

ALLOPLECTUS.

84². a. *splendens*, G. Don, ex Loud. Encyc. Pl. ii. 1402 = dichrous.

ALNUS.

84³. a. DUSCHEKIA, Opiz, Seznam, 38 (1852).
85¹. *elongata*, Royle, Illustr. Bot. Himal. 343, nomen (Quid?).—Reg. Himal.
l. microphylla, *Arv. Add. Monog. Pilos.* 20.— *Cf.*
a. *obtusifolia*, Royle, Illustr. Bot. Himal. 343, nomen (Quid?).—Himal.
ovata, Lange, ex Kjellm. in Nordensk. Vega-Exped. Vidensk. Iatt. i. 555 = viridis.

ALOCASIA.

85². a. SCHIZOCASIA, Schott, in Bonplandia, x. (1862) 148.
illustris, Bull, Cat. (1873) 4; Flor. Mag. (1874) 107 = Colocasia antiquorum.
85³. l. Regina, *N. E. Br.* . . . 711; *et in Illustr. Hortic.* (1885) 11. t. 544.
a. singaporensis, *Linden, ex Regel, Gartenfl.* (1865) 252 (Quid?).—Singapore.

ALOËXYLUM.

88¹. l. = Aq. Agallocha (bis in syn.).

ALOMIA.

88¹. d. ORSINIA, Bertol. (*nunc* **Clibadium**).
l. *Armani*, Baker, . . . 191 = Clibadium rotundifolium.

ALONA.

88¹. l. *tomentosa*, Lindl. . . . 46 = Dolia tomentosa.

ALOPECURUS.

88². a. CERDOSURUS, Ehrh. Beitr. iv. 148 (1789).
88³. l. *caesius*, Schur, . . . syn. = aristulatus.
fulvus, Sm. . . . t. 1467 = aristulatus.
89¹. *otiporensis*, Griff. . . . 94 = Phalaris minor.
paludosus, Beauv. . . . 481 = aristulatus.

ALOYSIA.

89². a. *triphylla*, Royle, Illustr. Bot. Himal. 299 = Lippia citriodora.

ALPHITONIA.

89³. l. *zizyphoides*, A. Gray, . . . 278 = excelsa.

ALPHONSEA.

89³. a. *verrucosa*, Baill. Hist. Pl. i. 215 = ventricosa.

ALPINIA.

89³. d. BOJERIA, Rafin. (*nunc* **Amomum**).
l. CENOLOPHON, Blume, . . . Jav. i. 60 (1827).
alata, *A. Dietr.* . . . China; Java.
90². a. vittata, *Hort. Bull. ex Dallière, Pl. Ornament.* i. t. 29.—Hab.?

ALSEODAPHNE.

90³. l. semecarpifolia, *Nees.*

PAG.

ALSINASTRUM.

90³. a. *americanum*, Torr. & Gray, ex Rupr. Fl. Ingr. 197, in obs. = E. americanus.

ALSINE.

92¹. a. *subnivalis*, Hegetschw. Fl. Schw. 422 (Quid?). —Europ.

ALSODEIA.

92³. a. DRIPAX, Noronha, ex Thou. Hist. Vég. Afr. ii. 58 (1807).
93¹. l. Physiphora, *Mart.*

ALSTROEMERIA.

94¹. l. *exserens*, Meyen, Reise, i. 311.
94². *montana*, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 183;
pulchella, Sims, . . . t. 2353.
pulchra, Sims,

ALTENSTEINIA.

94³. l. nubigena, *Reichb. f.* . . .—Ecuador.
95¹. *pilifera*, H. B. & K. . . . 73 (Stenopterae sp.).— Ecuador.

ALTERNANTHERA.

95¹. a. EVERION, Rafin. Sylva Tellur. 124 (1838).
l. *argentata*, Moq. . . . 352 = Hebanthe parviflora?
dubia, H. B. & K. . . . 209 = Iresine canescens.
frutescens, R. Br. . . . syn. = Telanthera frutescens?
glabra, Moq. . . . 361 = nodiflora?
a. *grandiflora*, Moq. ex Benth. & Hook. f. Gen. iii. 41 = Hebanthe decipiens.

ALTHAEA.

95³. l. *Froloviana*, Fisch. . . . 467 = ficifolia.
Kraussiana, Buching. . . . xxvii.

ALTINGIA.

96¹. l. chinensis, *Oliver*, etc.
caerulea, Poir. . . . = excelsa.

ALVARDIA.

96². a. *arborea*, Welw. Apont. 590 = seq.

96². a. ALWISIA, Thw. ex Lindl. in Journ. Linn. Soc. iii. (1838) 42, in obs. = **Taeniophyllum**, Blume (Orchid.).
minuta, Thw. ex Lindl. l. c. = T. Alwisii.

ALYSICARPUS.

96². l. *cylindricus*, Desv. in Mém. . . . = rugosus.

ALYSSOPSIS.

96³. l. *sagittata*, Boiss. . . . = deflexa.

ALYSSUM.

97¹. l. *Besseri*, *Sdev.*
97³. filifolium, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 54.

ALYXIA.

99¹. l. *actinophylla*, A. Cunn. . . . = Alstonia verticillosa.

AMAIOUA.

99². a. *fusifera*, *genipoides*, *oocarpa*, *pallens*, *petiolaris*, Spruce, ex Benth. & Hook. f. Gen. ii. 82 = Duroiae spp.?
l. *saccifera*, Mart. . . . 91 = Duroia saccifera.
surinamensis, Steud. . . . 763 = Duroia surinamensis.

AMALIA, Reichb.

99². a. *acuminata*, *albida*, *anceps*, *aurantiaca*, *autumnalis*, *cinnabarina*, *flava*, *fulva*, *furfuracea*, *Galeottiana*, *gigantea*, *grandiflora*, *majalis*, *pedunculares*, *Perrinii*, *purpurea*, *rubescens*, *superbiens*, Heynh. Nom. ii. 29, omnes Laeliae homonymae.

AMANOA.

99⁵. l. *brasiliensis*, Baill. . . . (= *Gonatogyne brasiliensis*).—Bras.

AMARANTHUS.

100¹. a. *Berchtholdii*, Hort. ex Mart. in Nov. Act. Nat. Cur. xiii. (1826) 290 (Quid?).—Hab.?
100². l. *emarginatus*, Salzm.
100³. a. *miniatus*, Hort. Avign. ex Hook. f. Fl. Brit. Ind. iv. 721 = A. Blitum.

PAG.

AMARYLLIS.

101³. l. *andicola*, Poepp. . . .—Chili.
a. *Carnarvonia* ×, DC. Pl. Rar. Hort. Genev. t. 9.
102². *Harrisonii*, Bury, Hexand. Pl. 27 = Hippeastrum solandriflorum.
Johnsoni, Bury, (l. c.) t. 1 = Hippeastrum solandriflorum.
marmorata ×, Hort. ex C. Morr. Ann. Bot. Gand, i. (1845) 148. t. 16.
Mendeli, Hort. ex Flor. Mag. (1875) t. 167 = Hippeastrum Reginae.
102³. *picta*, Bury, Hexand. Pl. t. 5 = Hippeastrum solandriflorum.

AMBERBOA.

103². l. *xanthocephala*, DC. l. c. 561 = Centaurea xanthocephala.

AMBLYOPOGON.

104². a. *aristata*, Boiss. Diagn. Ser. II. iii. 66 = Cent. aristata.
integrifolius, Boiss. (l. c.) = Cent. integrifolia.
microcephala, Boiss. (l. c.) = Cent. microcephala.

AMBROSIA.

105¹. a. purpurascens, *Tenore*, *Ind. Sem. Hort. Neap.* (1825) 1.—Hab.?

AMBULIA.

105¹. a. *ebracteata*, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 268 = L. hirsuta.

AMELANCHIER.

105². l. PERAPHYLLUM, Nutt. (*nunc* genus proprium).

105³. l. AMERIMNON, P. Br. . . . = **Dalbergia**, Linn. f. (Legumin.).

105³. a. AMERIMNUM, Scop. Introd. 305 (1777) = praec.

AMIANTHIUM.

106¹. in syn. lege Z. muscitoxicus (bis).

AMMANNIA.

106². a. DITHECA, Miq. Fl. Ind. Bat. i. 615 (1855).
EUTELIA, R. Br. ex DC. Prod. iii. 76, in syn. (1828).
MIRKOOA, Wight, Ic. i. t. 217, in textu (1840).
NESOEA, Wight, (l. c.) (1840).
auriculata, Rafin. Atl. Journ. 146 (Quid?).— Am. bor.

AMMI.

107². a. *paucifolium*, Steud. Nom. ed. II. ii. 592, in syn., sphalm. = indicum.
l. *trachycarpum*, C. A. Mey. ex Boiss.

AMOGETON.

108¹. l. = **Aponogeton**, Linn. (Naiad.).

AMOMUM.

108¹. a. BOJERIA, Rafin. Fl. Tellur. iv. 50 (1836).
l. DIRACODES, Blume, . . . (1827).
GEANTHUS, Reinw. in Blume, Cat. Gew. Buitenz. 29 (1823); et
108². echinatum, *Willd. Sp. Pl.*
medium, Lour. . . . 4 = Alpinia alba.

AMORPHA.

109². a. BONAFIDIA, Neck. Elem. iii. 46 (1790).
l. canescens, [*Nutt. in*] *Fras. Cat.* (1813) n. 4, nomen.
nana, [*Nutt. in*] . . . n. 5, nomen.

AMORPHOPHALLUS.

109³. a. DUNALIA, Montr. in Act. Soc. Linn. Bord. xxvi. (1866) 576.
l. *difformis*, Blume, . . . 149 = Anchomanes Hookeri.
Konjac, *C. Koch*, . . . 166.—Cochinch.; Japon.
Lacourii, Linden & André, . . . 90. t. 316 = Pseudodracontium Lacourii.
Peyri, *Linden.*—*Cf.*
Rivieri, Dur. . . . 15 = A. Konjac.

AMPHIBOLIS.

110³. d. GRAUMUELLERA, Reichb. (*nunc* **Cymodocea**).

PAG.

AMPHIDONAX.
111¹. l. *tenella*, Wight & Arn. . . . 197 = Z. elegans.

111³. l. AMPHISTELMA, Griseb. . . . (1861). *salinarum*, Wright, . . . 175 = V. salinarum.

AMPHITHALIA.
112¹. l. *spinosa*, Steud. . . . = Coelidium spinosum (in Addend.).

AMYGDALUS.
112³. l. *arabica*, Olivier, Voy. Atlas, t. 47.—Arab.

AMYRIS.
113¹. l. *acuminata*, DC. Prod. ii. 78, in syn. (= *Icica acuminata*).—Guian.

ANABASIS.
114¹. a. *ramosissima*, Boiss. ex Benth. & Hook. f. Gen. iii. 70 = Haloxylon ramosissimum.

ANACAMPSEROS.
114². l. *paucidens*, Haw. . . . 242 = S. paucidens.

ANACHARIS.
115¹. l. *callitrichoides*, Rich. . . . 2 = E. callitrichoides.
chilensis, Planch. l. c. = E. chilensis.
Matthewsii, Planch. . . . 73 = E. chilensis.

ANADENIA.
115³. l. *Manglesii*, R. Grah. . . . 337 = G. glabrata.

ANAGALLIS.
116¹. l. *caerulea*, Schreb. Fl. Lips. (1771) 5; Lam. *parviflora*, Loisel. . . . 401.

ANALECTIS.
116². a. *speciosa*, Vahl, in Danske Nat. Selsk. Skrivt. vi. (1810) 94 = S. involucratum.

ANAMIRTA.
116². l. *Baueriana*, Endl. . . . = paniculata.
flavescens, Miq. . . . = paniculata.

116². l. ANAMOMIS, Griseb. . . . (1860).

ANANTHERIX.
116³. l. *Nuttalliana*, G. Don,
pumila, Nutt. . . . l. c. = Podostigma pubescens.

ANAPHALIS.
117¹. a. MARGARITARIA, Opiz, Seznam, 63 (1852).
l. aristata, *DC. l. c.* 274.

ANARRHINUM.
117². l. *pedicellatum*, T. Anders. . . . 26 = Schweinfurthia pedicellata.

ANAXAGOREA.
118². a. canescens, *Benth. & Hook. f. Gen.* ii. 814, 815. —Afr. trop.
119¹. l. *ciliata*, Cham. . . . 400 = Moritzia ciliata.
a. *confertiflora*, DC. Prod. x. 55 = Lycopsis confertiflora.
l. *dasyantha*, Cham. l. c. 437 = Moritzia dasyantha.

ANCHUSA.
119¹. l. flava, *Forsk.* . . . 40.—Arab.
119². l. *Sellowiana*, Cham. . . . 115 = Moritzia Sellowiana.

119³. l. ANCISTROCARPHUS, A Gray, . . . 355 = **Stylocline**, Nutt. (Compos.).
filagineus, A. Gray, . . . 356 = S. filaginea.

ANCYLOGYNE.
120². l. *longiflora*, Hook. . . . t. 5588 = S. longiflora.
munita, Nees, l. c. = S. munita.

ANDERSONIA.
120². l. Homalostoma (non homalostoma).
120³. *subsessilis*, Panch. . . . 67 = Leucopogon subsessilis.

ANDERSONIA, Roxb.
120³. a. *Rohitoca*, Griff. Notul. iv. 507, err. typ. = seq.

ANDRACHNE.
121¹. l. *fruticosa*, Linn. Sp. Pl. 1014 (= *Melanthesopsis fruticosus*).—China.
121². *rotundifolia*, Eichw. . . . = A. Telephioides.
Telephioides, *Linn.* (non telephioides).

VOL. II.

PAG.

ANDRIAPETALUM.
121². l. *polystachyum*, Schott,

ANDRIEUXIA.
121². l. *quitensis*, Benth. . . . 206 = Gymnolomia quitensis.

ANDROMACHIA.
122¹. l. *sessiliflora*, H. B. & K. etc.
sessilifolia, DC. Prod. v. 96, sphalm. = praec.

ANDROMEDA.
122¹. l. *campanulata*, Miq. . . . = Enkianthus campanulatus.
122². a. *cordata*, Royle, Illustr. Bot. Himal. 258 = Pieris ovalifolia ?
l. *coriifolia*, Thunb. Pl. Bras., etc.
122³. l. *japonica*, Thunb. . . . = Pieris ovalifolia.
littoralis, Steud. . . . 88 (Agauriae sp.).—Malaya.
123¹. l. *revoluta*, Bouton, . . . 603 = (Agauriae sp.).—Madag.
a. *rupestris*, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 49 = Gaultheria antipoda.

ANDROPOGON.
123³. l. anthistirioides, et *anthistirioides*.
124¹. *aristulatus*, Hochst.
125³. d. *guineensis*, Steud.
126¹. l. Iwarancusa, *Jones, in As. Res.* iv. (1795) 109. —Ind. or.; Afr. trop.
126². *montanus*, Roxb.
126³. *nutans*, Linn.
peduncularis, *Baker*, . . . 444.—Ins. Borbon.
127¹. *pilosus*, Klein, . . . 920 = Ischaemum pilosum.
Pseudograyi, Pseudohirtus, Pseudoischaemum.
127². d. *Schoenanthus*, Benth.
127³. a. *stypticus*, Welw. Synopse, 27 (Quid ?).—Afr. trop.
l. *speciosus*, Steud. . . . 375 (Rottbolliae sp.).—Reg. Himal.
128¹. a. *Trachypogon*, Royle, Illustr. Bot. Himal. 417, nomen (Quid ?).—Reg. Himal.
tristis, Royle, (l. c.) nomen (Quid ?).—Reg. Himal.
128². l. virginicus, *Linn.*

ANDROSACE.
128². a. aretioides, *Hegetschw. Fl. Helv.* 187.—Helvet.
129¹. *nidulans*, Royle, Illustr. Bot. Himal. 310, nomen (Quid ?).—Reg. Himal.
reptans, Royle, (l. c.) nomen (Quid ?).—Reg. Himal.

ANDROSAEMUM.
129². l. *Cambessedesii*, Cass. . . . 131 = H. Cambessedesii.

ANDROTRICHUM.
129². a. *montevidense*, Steud. Nom. ed. II. i. 587, in syn. = seq.

ANDRYALA.
129³. a. *rosea*, Steud. Nom. ed. II. ii. 779, in syn., sphalm. = A. Rothia.

ANEILEMA.
130³. l. *tacazzeanum* (bis).

ANEMONE.
131². d. *ambigua*, (sphalm. pro Pulsatilla).
132². l. *thalictroides*, Linn. . . . 542 = Thalictrum anemonoides.
132³. a. *Zichyi*, Schur, in Oestr. Bot. Zeitschr. (1863) 317 = pratensis.

132³. l. ANEMONOPSIS, Pritz. . . . = ANEMOPSIS, Hook. = **Houttuynia**, Thunb. (Piperac.).

132³. a. ANEMONOSPERMOS, Boerh. ex Boehm. in Ludw. Def. Gen. Pl. 192 (1760); Adans.

ANETANTHUS.
133¹. l. alata, *Benth. & Hook. f.*

ANGELICA.
133². a. PHYSOLOPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) II. 728.
133³. *decursiva*, Franch. & Sav. Enum. Pl. Jap. i. 187 = Peucedanum decursivum.

PAG.

ANGELONIA.
134². l. *lobelioides*, Mart. Reise Bras. i. 397, nomen;

ANGIANTHUS.
134³. l. Pleuropappus, *Benth.* (non pleuropappus).

134³. l. ANGOSTURA, Roem. & Schult. . . . (1819) = **Cusparia**, Humb. (Rutac.).
Cuspare, Roem. & Schult. l. c. = C. febrifuga.

ANGRAECUM.
135¹. a. GUSSONEA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 67.
crenatum, *Reichb. f. ex Williams, Man. Orchid.* ed. VI. 115.—Afr. trop.
crystallinum, *Henshall, Pract. Treat. Orchid.* 106 (chrystalinum), nomen.—Afr. trop.
l. falcatum, *Lindl. Gen. et Sp. Orch.* 237, in obs.
135². jamaicense, *Reichb. f. & Wullschl.*
a. *Leonii*, André in Rev. Hortic. (1885) 294 = Humblotii.
Leonis, Hort. . . . Humblotii.

ANGULOA.
136¹. l. *squalida*, Poepp. & Endl. . . . 74 = Lycomormium squalidum.

ANGURIA.
136². d. Warmingii, *Cogn. l. c.*—Bras.

ANHALONIUM.
136². l. *areolosum*, Lem. . . . 35 = M. areolosa.
fissipedum, Monv, . . . = *sulcatum*.
Kotchubeyi, Lem. . . . = M. sulcata.

ANIA.
136³. l. *angustifolia*, Lindl. . . . n. 3740 = Tainia angustifolia.

136³. l. **ANIBA**, Aubl. Pl. Gui. 327 (1775). *LAURINEAE.*
guianensis, *Aubl. l. c.*—Guian.

136³. insere, ANIL, March. ex Ludw. Defin. Pl. 117 (1737) = **Indigofera**, Linn. (Legumin.).

ANISANTHUS.
137¹. l. *abyssinicus*, Klatt, . . . xv. (non xii).
huillensis, Klatt, . . . xv.

ANISEIA.
137¹. l. aurea, *Kellogg*, . . . 83 = I. aurea (in Addend.).

139². l. **ANNESORHIZA.**
a. caffra, *Benth. & Hook. f. Gen.* i. 902.—Afr. austr.
l. capensis, *Cham. & Schlecht. in Linnaea*, i. (1826) 399.—Afr. austr.

ANOECTOCHILUS.
140². l. *sandwicensis*, Lindl. . . . 500 = Vrydagzynea sandwicensis.

ANOMATHECA.
140³ l. *calamifolia*, Klatt, . . . xv.
xanthospila, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 171 = Freesia xanthospila.

ANOMOCARPUS.
140³. l. *tenuifolius*, Miers, . . . = C. tenuifolia.

ANONA.
141¹. l. *conophylla*, C. Muell. in Walp. Ann. vii. 58, sphalm. = ionophylla.
141². a. *Kraussiana*, Hochst. in Flora, xxvi. (1843) 69 = Oncoba Kraussiana.
l. *polycarpa*, Oliver, Fl. Trop. Afr. i. 32 = Xylopia polycarpa.
141³. a. *uncata*, Baill. Hist. Pl. i. 232 = Artabotrys odoratissimus.

143¹. l. ANSERINA, Dum. . . . **Chenopodium**, Tourn.
a. *betaefolia*, Montand. Guide Bot. 264 = C. polyspermum.
candicans, Montand. (l. c.) 262 = C. album.
ficifolia, Montand. l. c. 264 = C. ficifolium.
foetida, Montand. l. c. = C. Vulvaria.
muralis, Montand. (l. c.) 263 = C. murale.
opulifolia, Montand. l. c. = C. opulifolium.
stramonifolia, Montand. l. c. = C. hybridum.
urbica, Montand. l. c. = C. urbicum.

PAG.

ANTENNARIA.

143¹. l. *andina*, A. Gray, . . . 138 = Mniodes andina.
linearifolia, Wedd. . . . 150 = Leontopodium linearifolium.
magellanica, Sch. Bip. . . . 117 = chilensis.
143². l. *monoica*, Wedd. . . . 150 = Leontopodium monoicum.

143³. l. ANTHËEISCHIMA.

ANTHEMIS.

144¹. l. LYONNETIA, Cass. . . . xxxiv. 166 (1825).
144². a. *comosa*, Peterm. Fl. Bienitz, 42 (Quid ?).—Europ.
coronata, Peterm. (l. c.) 43 (Quid ?).—Europ.
144³. l. *discoidea*, Willd.
145². parthenoides, *Bernh.*
145³. *salvifolia*, Boj. . . . 3 = Sphacophyllum madagascariense.

ANTHERICUM.

146². l. LILIAGO, [Caesalp.] Linn. Class. Pl. 22 (1747); Salisb., etc.
d. BLEPHANTHERA, Rafin. (*nunc* **Bulbine**).
147². l. *pomeridianum*, Ker-Gawl. in [Quart.] Journ. Roy. Inst. i. (1816) 181; et Bot. Reg.
147³. *Rossii*, Hook. f.
a. *striatellum*, Steud. Nom. ed. II. ii. 228, sphalm. = Allium striatellum.
l. *subtrigynum*, Jacq. . . . = Zygadenus muscitoxicus.

ANTHISTIRIA.

148². l. anathera, *Nees, in Royle, Illustr. Bot. Himal.* 417.—Ind. or.
Forskălii, *Kunth*, . . . 162.—Arab.
148³. a. *semiberbis*, Nees, Fl. Afr. Austr. 125 = ciliata.

ANTHOCEPHALUS.

148³. a. PLATANOCEPHALUS, Vaill. ex Crantz, Inst. i. 349 (1766).

149¹. post ANTHOCHLAMYDAM, pone

ANTHOCHLOA, Nees & Meyen, Reise, ii. 14 (1845); et in Lindl. Introd. Nat. Syst. ed. II. 450 (1836). *GRAMINEAE*, Benth. & Hook. f. iii. 1189.
lepida, *Nees & Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 165.—Peruv.; Boliv.
lepidula, Nees & Meyen, Reise, ii. 14 = praec.
rupestris, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 347.—Bolivia.

ANTHOCOMA.

149¹. l. *flavescens*, Zoll. & Mor. . . . 570 = Cymaria acuminata.

ANTHODON.

149². l. *Selloanum*, Schult. . . . 411 = S. brasiliensis.
a. *Wrightianum*, Miers, ex Triana & Planch. in Ann. Sc. Nat. Sér. V. xvi. (1872) 372, in nota = H. malpighiaefolia ?

ANTHOLYZA.

149². l. *bicolor*, Gasparr. in Ann. Civ. i. (1833) 118, etc.

ANTHOSPERMUM.

150¹. l. *Crocyllis*, Sond. . . . 32 = Crocyllis anthospermoides.

ANTHURIUM.

152¹. a. *laciniatum*, C. Koch, Ind. Sem. Hort. Berol. (1855) 4 = podophyllum.

153¹. a. post **ANTHURIUM** pone
ANTHYLLIS, Adans. Fam. ii. 271 (1763) = **Polycarpaea**, Lam. et **Polycarpon**, Loefl. (Caryoph.).

ANTHYLLIS.

153². a. rubriflora, *Hegetschw. Fl. Schw.* 693.—Helvet.
153³. lin. 1ma l. **ANTHYLLIS.**

ANTIARIS.

153³. a. TOXICARIA, Aepnel. ex Steud. Nom. ed. I. 434, in syn. (1821).

ANTICHARIS.

153³. a. scoparia, *Hiern, ex Benth. & Hook. f. Gen.* ii. 928.—Afr. austr.

PAG.

ANTIDESMA.

154¹. a. *glabella*, Koen. ex Benn. Pl. Jav. Rar. 192 (Quid ?).—Java.

155¹. **ANTIRRHOEA** (ponitur post **ANTIRRHINUM**).

155². l. *borbonica*, J. F. Gmel. ex C. DC. in DC. Monog. Phan. i. 43 = Quivisia Sieberi.
a. borbonica, *J. F. Gmel. Syst.* i. 244.—Ins. Borbon.

ANTIRRHINUM.

155¹. a. SACCULARIA, Kellogg, in Proc. Calif. Acad. ii. (1863) 17.
155³. l. *elatiniflorum*, Wahlenb. in Bergg. Resor, ii. Bihang, 37;
glandulosum, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 144 = vexillocalyculatum.
a. *glutinosum*, Brot. Phyt. Lusit. i. 129 = Linaria filifolia.
156². l. Nuttallianum, *Benth.*

ANTITAXIS.

157¹. l. *longifolia*, Miers, . . . 357 = Pycnarrhena longifolia.

157¹. a. ANTOCHLOA, Nees & Meyen, in Lindl. Introd. Nat. Syst. ed. II. 450 (1836) = **Anthochloa**, Nees & Meyen (Gramin.).

157². l. ANTONIANA, Tuss. Fl. Antill. ii. 62. t. 16 (1818) = **Faramea**, Aubl. (Rubiac.).

ANTROLEPIS.

157². l. *leucocephala*, Welw.

ANYCHIA.

157². d. BUINALIS, Rafin. (*nunc* **Paronychia**).

APALOCHLAMYS.

157³. a. *spectabilis*, Steud. Nom. ed. II. i. 255, in syn., sphalm. = Cassinia spectabilis.

APERA.

158³. a. *Liebmanni*, Benth. & Hook. f. Gen. iii. 1153, sphalm. = Deyeuxia Liebmanni.

APERULA.

159¹. a. *lucida*, Blume, . . . 366 (= *Litsea lucida*).—Java.
l. *oxyphylla*, Blume, l. c.—Ins. Penang.

159³. l. APHANOCHAETA, A. Gray, . . . (1857) = **Pentachaeta**, Nutt. (Compos.).
exilis, A. Gray, l. c. 100 = P. exilis.

APHELANDRA.

159³. d. HEMITOME, Nees, et SYNANDRA, Schrad. (*nunc* **Stenandrium**).
l. Hydromestus, *Hemsl.*
160¹. l. mediaurata, *Cogn. & March. in Dallière*, pumila, *Bull, Cat.* n. 143 (1878) 4; *Bot. Mag.* t. 6467.—Bras.

APHYLLOCLADUS.

160². l. *spartioides*, Wedd. l. c. = H. spartioides.

APHYLLON.

160³. l. GYMNOCAULIS, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 180.

APIUM.

161². l. CICLOSPERMUM, Lag. . . . (1821).
a. *decumbens*, Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 45 (Quid ?).—Reg. Magell.

APLOPAPPUS.

162¹. l. ACAMPTOPAPPUS, A. Gray,
a. ERICAMERIA, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 318.
162². l. ericoides, *DC.* . . . 346.
162³. l. *junceus*, Greene, . . . IV. (1884).
nevadensis, Kellogg, . . . 9 = acaulis.

APODYTES.

164³ a. HEMILOBIUM, Welw. Synopse, 20 (1862).

APOLLONIAS.

165¹. l. *Barbusana*, R. Br.

PAG.

166¹. l. APOZIA, Willd. ex Benth. in Linnaea, xi. (1837) 328.
chamaedryoides, *herniariodes*, *serpyllacea*, Willd. ex Benth. l. c.

APPENDICULA.

166¹. l. membranacea, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 491; *et in Nederl. Kruidk.*

ARABIS.

168¹. d. PILOSELLA, Kostel. (*nunc* **Sisymbrium**).
168³. d. *Dregeana*, C. Muell. in Walp. Ann. vii. 103 = perfoliata.
168³. a. *Ermani*, Kjellm. in Nordensk. Vega-Exped. Vidensk. Iakttl. i. 537 = Parrya Ermani.
169². a. *parryoides*, Kjellm. (l. c.) = Parrya Ermani.

ARACHNITES.

170³. a. *limbata*, Link, Handb. i. 247 = O. limbata.

l. ARACHNOPOGON, Berg, . . . = **Hypochoeris**, Linn. (Compos.).
l. *heterospermum*, Berg, l. c. = H. glabra.

171¹. a. ARAGALLUS, Neck. Elem. iii. 12 (1790) = **Astragalus**, Tourn. (Legumin.).

171¹. a. ARAGO, Endl. Ench. 340 (1841), sphalm. = seq.

171¹. a. **ARAGOA**, H. B. & K. Nov. Gen. et Sp. iii. 154 (1818). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 975.
abietina, *H. B. & K. Nov. Gen. et Sp.* iii. 156. t. 217.—N. Granat.
cupressina, *H. B. & K. l. c.* 156. t. 216.—N. Granat.
juniperina, H. B. & K. ex Steud. Nom. ed. II. i. 118 = cupressina.
lycopodioides, *Benth. ex Oliver, in Hook. Ic. Pl.* t. 1325.—N. Granat.

ARALIA.

171¹. a. GINSENG, Wood, Am. Bot. & Flor. 142 (1871).
171². *elegans*, Linden, ex Decne. & Planch. in Rév. Hortic. Sér. IV. iii. (1854) 108 = Oreopanax elegans.
171³. *parasitica*, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 138 = Pentapanax parasiticum.
Reginae, *Linden, ex André, Illust. Hortic.* (1879) 25. t. 337.—N. Caled.
172¹. l. *trigyna*, Gaudich. . . . = Panax Gaudichaudii.

ARCTIUM.

173³. l. nemorosum, *Lej. in Court. Mag. Hortic.* 289; *ex Lej. & Court. Comp. Fl. Belg.* iii. 129.—Europ.

ARCTOSTAPHYLOS.

174¹. l. alpina, *Spreng.* . . . Reg. bor. et arct.

ARCTOTIS.

174². l. ANEMONOSPERMOS, Boerh. ex Boehm. in Ludw. Def. Gen. Pl. 192 (1760); Adans.
a. *acaulis*, Brot. Fl. Lusit. i. 401 = Cryptostemma calendulaceum.

ARDISIA.

175². a. STRANGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4.
175³. l. *Blumei*, DC.
176¹. l. *fuliginosa*, Blume, . . . 692 = Pimelandra fuliginosa.
177¹. l. scrobiculata, Seem. . . . 150.—N. Hebr.; N. Caled.
177². l. *Teysmanniana*, Scheff. . . . 253 = Pimelandra Teysmanniana.

ARECA.

177². l. Alicae, *F. Muell. in Gard. Chron.* (1879) I. 760; *et in Gartenfl.*
177³. a. *gigantea*, Reinw. Reise Ind. Arch. 500 = Gulubia moluccana.
minor, W. Hill, ex F. Muell. Fragm. viii. 235 = Kentia minor.

ARENARIA.

178². a. DUFOUREA, Gren. in Act. Soc. Linn. Bord. ix. (1837) 25.
Allionii, Raeusch. Nom. ed. III. 130, nomen (Quid ?).—Italia.

PAG.

ARENARIA:—

179[1]. l. Cerviana, *Cham. & Schlecht.*

180[1]. a. *massiliensis*, Fenzl, ex Gren. & Godr. Fl. Fr. i. 262 = A. Gouffeia.
miquelonensis, La Pylaie, ex Lebel, in Bull. Soc. Bot. Fr. xv. (1868) 58 = Sperugalaria miquelonensis.

181[1]. l. sedoides, *Froel. ex Reichb. Fl. Germ. Excurs.* 794, in syn.—Tyrol.

181[2]. a. *Spergula*, Borkh. in Rhein. Mag. i. (1793) 538 = Spergula arvensis.
sperguloides, *Mutel, Fl. Fr.* 165.—Europ.
l. *triandra*, Schrank, . . . t. 30.

181[3]. l. *viscida*, Hall. f. ex Loisel.

ARGITHAMNIA.

182[3]. a. ODOTALON, Rafin. Sylva Tellur. 66 (1838).

ARGOSTEMMA.

183[1]. a. javanicum, *Wawra, Itin. Princ. Coburgi*, i. 117. t. 39. f. D.—Jav.

183[2]. l. ARGUSIA, Amm. ex Linn. Hort. Upsal. 36 (1748) = seq.

183[2]. l. ARGUZIA, Rafin. Sylva Tellur. 167 (1836); Stev. in Bull. Soc. Nat. Mosc. (1851) i. 558 = **Tournefortia**, Linn. (Borag.).

ARGYLIA.

183[3]. a. OXYMITUS, Presl, Bot. Bemerk. 92 (1844).
tenuifolia, Presl, (l. c.) 93, in obs. = canescens.

ARGYREIA.

183[3]. l. *festiva*, Wall. . . . 76 = Lettsomia festiva.

ARGYROLOBIUM.

184[2]. a. LOTOPHYLLUS, Link, Handb. ii. 156 (1831), nomen prius.

ARISAEMA.

185[3]. a. *Dioscoridis*, Royle, Illust. Bot. Himal. 407 = Arum Dioscoridis.
l. nepenthoides, *Mart. in Flora*, xv. (1831) II. 458, 459; *et ex*
speciosum, *Mart. (l. c.)*; *et ex*
utile, *Hook. f. ex Schott, Prod. Aroid.* 30.

ARISARUM.

186[1]. a. multicarinatum, *Wall. ex Royle, Illust. Bot. Himal.* 406.—Reg. Himal.

ARISTEA.

186[1]. a. corymbosa, *Benth. & Hook. f. Gen.* iii. 701.—Afr. austr.

ARISTIDA.

181[2].–188[1]. l. passim, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1843).

186[2]. l. barbicollis, *Trin. & Rupr.* . . . 152.—Afr. austr.

186[3]. cyanantha, *Steud. Nom.* ed. II. ii. 131; *Syn. divaricata*, Lag. Gen. & Sp. Nov. 3.

187[1]. l. hirtigluma, *Steud.* . . . VI. *Sc. Nat.* v. (1842) 171.—Afr. trop.
hystricula, *Edgew. in Journ. Linn. Soc.* vi. (1862) 208.—Ind. or.

187[2]. a. longiflora, *Schum. & Thonn. Beskr. Guin. Pl.* 48.—Afr. trop.

187[2]. a. *mallica*, Edgew. in Journ. Linn. Soc. vi. (1862) 209 = funiculata.

188[1]. a. submucronata, *Schum. & Thonn. Beskr. Guin. Pl.* 47.—Afr. trop.
l. *trichodes*, Steud. Syn. Pl. Gram. 140 = Cumingiana.

ARISTOLOCHIA.

188[2]. l. *aethiopica*, Welw. . . . 548 = albida.
a. arborea, *Linden, Cat.* n. 13 (1858) 6; *Hook. f. Bot. Mag.* t. 5295.—N. Granat.
l. *Bonplandii*, Tenore, in Rendic. Acc. Sc. Nap. i. (1842) 345; et ex

188[3]. a. *dodecandra*, Rafin. Med. Fl. i. 62 = A. Serpentaria?

189[1]. a. hyperborea, *Paxt. Mag. Bot.* vi. (1839) 53.—Hab.?

190[1]. l. ringens, *Vahl*, . . . —Ind. occ.

190[2]. insere ARISTOTELIA, Comm. ex Lam. Encyc. i. 349 (1784) = **Terminalia**, Linn. (Combret.).

PAG.

ARMENIACA.

190[3]. l. *Mume*, Siebold, in Verh. Batav. Gen. xii. (1830) 69; De Vriese,

ARMERIA.

191[1]. l. *Pseudo-armeria*, Willk. & Lange, Prod. Fl. Hisp. ii. 365 = latifolia.

191[3]. a. ARMOLA, Montand. Guide Bot. 261 (1868) = **Atriplex**, Tourn. (Chenop.).
campestris, Montand. l. c. = Atr. patula.
deltoidea, Montand. l. c. = Atr. hastata.
mixta, Montand. l. c. = Atr. patula.

ARMORACIA.

191[3]. a. *americana*, Hook. & Arn. Brit. Fl. ed. VI. 28, in nota = Nasturtium lacustre.
natans, Hook. & Arn. l. c. = Nasturtium natans.

ARNEBIA.

192[1]. a. *tetrastigma*, Auct. ex Steud. Nom. ed. II. ii. 56 = tinctoria.

ARNICA.

192[1]. l. *austriaca*, [Hoppe] Roehl. Deutschl. Fl. ed. II. ii. 462 =

ARRACACIA.

193[3]. a. arguta, *Benth. & Hook. f. ex S. Wats. Bibliog. Ind. Bot. N. Am.* 419.—Am. bor.

193[3]. insere ARRASCHKOOLIA, Sch. Bip. ex Hochst. in Flora, xxv. (1842) i. Beibl. 144 = ARASCHCOOLIA, Sch. Bip. = **Geigeria**, Griessel. (Compos.).

ARRHENATHERUM.

193[3]. l. kentuckense, *Torr. Fl. U. St.* 131, in nota.

ARRHOSTOXYLUM.

194[1]. a. *grandiflorum*, Nees, in DC. Prod. xi. 158 = R. grandiflora.

ARROZIA.

194[1]. l. *micrantha*, Schrad. l. c. = L. micrantha.

ARTEMISIA.

195[3]. a. DRACUNCULUS, [Rupp.] Ledeb. Fl. Ross. ii. 566, in syn., cum cit. falsa (1846).

196[1]. l. Absinthium, *Linn.* . . . —Geront. temp.

196[2]. d. *campestris*, Ledeb.

196[3]. a. *doonense*, Royle, Illust. Bot. Himal. 250 = A. Absinthium.

197[2]. a. *Ibukijomogi*, Siebold, in Verh. Batav. Gen. xii. (1830) 70 = vulgaris.

197[3]. l. kohatica, *Klatt.* . . . (1878) i. 91.

198[3]. l. *pedemontana*, Balb. in Mém. Acad. Taur. (1810) reimpr. 21. t. 3;

199[1]. l. Schlagintweitiana, *Klatt*, . . . (1878) i. 90.

199[3]. a. *trifida*, Pav. ex Delpino, Studi Artemis. 64 (= *Xanthidium rhombophyllum*).—Peruv.

ARTHRAXON.

200[1]. l. PSILOPOGON, Hochst. in Flora, xxiv. (1841) i. Intell. 20.

ARTHROSOLEN.

200[3]. a. TRIMEIANDRA, Rafin. Fl. Tellur. iv. 105 (1836).

ARTHROSTEMA.

201[1]. a. *Humboldtianum*, Kunth, Ind. Sem. Hort. Berol. (1844) 11.—Cf. Linnaea, xviii. (1844) 504 = Monochaetum Humboldtianum.

ARTOCARPUS.

201[3]. l. *exculpta*, Hort. . . . 84 = Ficus exculpta.

ARTROLOBIUM.

202[1]. a. *parviflorum*, Desv. in Mém. Soc. Linn. Par. iv. (1826) 304 = Coronilla parviflora.
repandum, Desv. (l. c.) = Coronilla repanda.

ARUBA.

202[2]. l. *guianensis*, Aubl. l. c. = S. guianensis.

ARUM.

202[2]. l. *bulbiferum*, Roxb. . . . = Amorphophallus bulbifer.

202[3]. a. *compressum*, G. Don, in Edinb. Phil. Journ. xi. (1824) 344 (= *Corynophallus Afzelii*?).—Afr. trop.

PAG.

ARUM:—

202[3]. a. *crinitum*, Ait. . . . = Helicodiceros crinitus.

203[1]. *hastatum*, Royle, Illustr. Bot. Himal. 406, nomen (Quid?).—Ind. or.
insignifolium, Royle, l. c. nomen (Quid?).—Ind. or.
l. *integrifolium*, Link, . . . 394 = Aglaonema oblongifolium.

ARUNDINA.

204[1]. l. pulchella, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 492; *et in speciosa*, Blume, . . . 73 = densa.

ARUNDINELLA.

204[2]. l. *flammida*, Trin. . . . 267 = Trichopteryx flammida.
a. *gigantea*, Dalz. & Gibs. Bomb. Fl. 293 (Quid?).—Ind. or.

204[3]. l. holcoides, *Trin.* . . . 71.—Burma.
a. *miliacea*, Nees, ex Royle, Illustr. Bot. Himal. 417, nomen (Quid?).—Reg. Himal.
l. nervosa, *Nees*, . . . 237.—China.

ARUNDO.

205[1]. l. *australis*, Cav. . . . = Phragmites communis.

205[2]. *festucoides*, Desf. . . . = Ampelodesma tenax.
madagascariensis, *Kunth*, . . . 48.—Geront. trop.
a. *montana*, Gaud. in Alpina, iii. (1808) 19; Agrost. Helv. i. 91 = Deyeuxia acutifolia.

205[3]. *nigricans*, Mérat, Fl. Par. ed. II. 33 = Phragmites communis.
Pseudo-arundinacea, Schleich. ex Gaud. in Alpina, iii. (1808) 19 = Deyeuxia acutiflora.

206[1]. l. ARUNGANA, Pers. . . . = **Haronga**, Thou. . . . (1806).
crenata, Pers. . . . = Psorospermum cordiaefolium.
mollusca, Pers. . . . = Psorospermum cuspidatum.

ARYTERA.

206[2]. l. *littoralis*, Blume, . . . 170 = Ratonia littoralis (in Addend.).

ASCLEPIAS.

207[2]. a. *amoena*, Ruiz & Pav. ex Hemsl. Biol. Centr. Am. Bot. ii. 325 = neglecta.
l. *acornuta*, Kellogg, . . . 54 = Gomphocarpus cordifolius.

208[1]. a. filiformis, *Benth. & Hook. f. Gen.* ii. 753, in nota sub Schizoglosso.—Afr. austr.
l. *filiformis*, Jacq. . . .—Afr. austr.
fragrans, Rafin. . . . = A. syriaca (syn. *Cornuti*).

208[2]. lanceolata, *Walt.* . . . 105.—Am. bor.
longicornis, Kellogg, . . . 65 = Gomphocarpus cordifolius.

208[3]. a. *odoratissima*, Breon, ex Brongn. in Ann. Sc. Nat. Sér. II. vii. (1837) 30 = Stephanotis floribunda.
l. *paupercula*, Michx. . . . 118 = lanceolata.

209[1]. *Pseudosarsa*, Roxb.
scandens, Beauv. . . . 56 = Daemia scandens.

ASCOLEPIS.

209[3]. l. eriocauloides, *Nees*,

ASCYRUM.

209[3]. l. HYPERICOIDES, Plum. ex Adans.

210[1]. *nummularifolium*, Banks, ex Pursh, Fl. Am. Sept. ii. 273 =

ASPALATHUS.

210[2]. d. NEFRAKIS, Rafin. (*nunc* **Brya**).

ASPARAGUS.

212[3]. l. *cuscutoides*, Burch. . . . 606 = Schizobasis cuscutoides.
davuricus, *Fisch.* . . . 340.

213[2]. a. *ramosus*, Royle, Illustr. Bot. Himal. 447 (Index) sphalm. = racemosus.

213[3]. sarmentosus, [*Delile, in*] *Red. Liliac.* t. 460, quoad fig.; *Herb. Linn. ex Baker, in Journ. Linn. Soc.* xvi. (1875) 625.—Afr. austr.
l. sarmentosus, *Linn.* . . . 314.—Zeylan.

ASPERELLA.

214[1]. a. californica, *Benth. ex Vasey, Cat. Gr. U. St.* 46.—Calif.
l. Hystrix, *Humb. in Roem. & Ust. Mag.* iii. VII. (1790) 5.—Am. bor.

PAG.

ASPERULA.

214³. l. involucruta, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 21; *et in*

ASPHODELINE.

215². l. liburnica, *Reichb.*

ASPHODELUS.

216¹. a. pyramidalis, *Royle, Illustr. Bot. Himal.* 392, nomen.—Reg. Himal.

ASPIDOSPERMA.

216². a. THYROMA, Miers, Apocyn. S. Am. 22 (1878).

ASPILIA.

217¹. d. abyssinica, *Oliver & Hiern* (=seq.).
l. *Gayana*, Benth. & Hook. f. . . . 372 = helianthoides.
Prieureana, Benth. & Hook. f. . . . 372 = Kotschyana.

ASTELIA.

217³. l. grandis, *Hook. f.* . . . i. ed. II. (1871).
trinervia, *T. Kirk*, . . . i. ed. II. (1871).

ASTER.

218². a. COELESTINA, Hill, Hort. Kew. 8 (1769).
GALATELLA, Cass. in Dict. Sc. Nat. xxxvii. 463 (1825).
d. HOMOSTYLIUM, Nees (*nunc* **Microglossa**).
a. *acer*, Roehl. Deutschl. Fl. ed. II. ii. 462 = acris.

218³. l. andinus, *Nutt.* . . . 290.—Reg. Andin.

219¹. l. Bellidiastrum, *Meyen, Reise*, i. 311, nomen; *Nees*, etc.

219³. a. confusus, *Harv. ex Harv. & Sond. Fl. Cap.* iii. 75.—Afr. austr.

220¹. l. diplostephioides, *Benth. ex C. B. Clarke, Comp. Ind.* 45.

220³. l. Gerlachii, *Hance*,
glabriusculus, Torr. & Gray, . . . = A. Xylorrhiza.

221¹. *Heterochaeta*, C. B. Clarke.
imbricatus, *Walp.* . . . 574.—Chili.

222³. l. Pseudamellus, Pseudodiffusus, Pseudodumosus, etc.

223¹. a. *Rivesii*, Hort.—Cf. Linnaea, xxiv. (1851) 235 = Felicia Reevesii.

223². l. *scaposus*, Klatt, . . . (1878) I. 84.
scorzonerifolius, Regel, . . . 308 = Andersoni, *A. Gray*.

224¹. l. *tenuis*, Kellogg, . . . 114 = exilis?
a. tuberosus, *Less. ex Baker, in Mart. Fl. Bras.* vi. III. 22.—Bras.

224². l. Xylorrhiza, *Torr. & Gray*,

ASTEROCHAETE.

225². l. *arundinacea*, Kunth, . . . = Schoenus arundinaceus.
nitens, Kunth, . . . = Cyclocampe nitens.

ASTRAGALUS.

226². a. ASTRAGALOIDES, Adans. Fam. ii. 323 (1763).

226³. PHYLLOLOBUM, Fisch. ex Spreng. Nov. Prov. 33 (1818).
d. SPIESIA, Neck. (*nunc* **Oxytropis**).
a. TRAGACANTHA, Tourn. ex Linn. Syst. ed. I. (1735); Mill. Gard. Dict. ed. VI. (1752).

227². l. anemophilus, *Greene, . . . Acad.* i. IV. (1885).

227³. l. Arnacantha, *Bieb.*

228³. Candolleanus, *Royle*, . . . 199.—Reg. Himal.

229². *clandestinus*, Roth, ex Spreng. in Schrad. Journ. (1800) II. 195 =

229³. *crassicarpus*, [Nutt. in] Fras. Cat. (1813) n. 6, nomen; et

230¹. *dasyglottis*, Fisch. . . . = hypoglottis.

230². Dipelta, *Bunge*,

231¹. fastidiosus, *Greene*, . . . i. IV. (1885).

233¹. Kentrophyta, *A. Gray*,

233³. a. *longipes*, Arv. Ess. Pl. Dauph. 23 (Quid?).—Gallia.

236¹. l. Pattersoni, *A. Gray*, . . . 235.
a. *phacaceus*, Arv. (l. c.) 24 (Quid?).—Gallia.

236². prusianus, *Boiss.* . . . 88.—As. Min.
l. Pseudaustralis, Pseudocaspius, *Pseudocicer*, Pseudonobrychis, *Pseudopentaglottis*, Pseudostella.

236³. *Pseudotragacantha*, (bis).

237¹. *rotundifolius*, J. & C. Presl, Fl. Čech. 151 =

238². a. subverticillaris, *R. Grah. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 62, in obs.—Hab.?

PAG.

ASTRANTIA.

240¹. l. *pallida*, J. & C. Presl, Fl. Čech. 62 = major.

ASTREPHIA.

240¹. d. OLIGACOCE, Willd. (*nunc* **Valeriana**).

ASTROCARPUS.

240². a. SESAMOIDES, Tourn. ex Reichb. Consp. 186 (1828).
l. *Sesamoides*, Colm.
Sesamoides, *DC.*

ASTROCARYUM.

240³. a. *pictum*, Hort. ex Balf. f. in Baker, Fl. Maurit. 388 = Stevensonia grandifolia.

ASTRONIUM.

241². a. MYRACRODRUON, Allem. in Trab. Comm. Sc. Expl. Cearà; Secç. Bot. 3. tt. 1, 2 (1862).

241². l. gracile et macrocalyx . . . 45.

241². l. ASTROPHYTUM, Lem. Cact. Gen. Nov. et Sp. 3

ASYSTASIA.

241³. a. bella, *Benth. & Hook. f. Gen.* ii. 1095.—Afr. austr.

242¹. l. *Thyrsacanthus*, Klotzsch,

ATALAYA.

242². a. *australis*, F. Muell. ex Baill. Hist. Pl. v. 419 = multiflora.

242². l. ATAXIA, R. Br. in Parry, 1st Voy. App. 193 (non 202).

ATHAMANTA.

243². l. *sicula*, Linn. l. c. = Tinguarra sicula.

ATHANASIA.

243³. l. parviflora, *J. F. Gmel. Syst.* 1207; *Thunb.*

ATHENAEA.

244¹. a. *viscosa*, Benth. & Hook. f. Gen. ii. 891, in nota (Physalidis sp.).—Mexic.

244¹. l. **ATHEROLEPIS**, Hook. f.

ATHRIXIA.

244². l. rosmarinifolia, *Oliver & Hiern*,

ATRAPHAXIS.

245². a. ACARNA, Boehm. in Ludw. Def. Gen. Pl. 195 (1760); All. Fl. Pedem. i. 153 (1785).

ATRIPLEX.

245³. a. ARMOLA, Montand. Guide Bot. 261 (1868).

246¹. l. *Babingtonii*, Woods, Tourists' Fl. 316 =

246². *deltoidea*, Bab. Prim. Fl. Sarn. 32;

247³. a. *ucenica*, Arv. Ess. Pl. Dauph. 59 (Quid?).—Gallia.

ATROPA.

248². l. *spinosa*, Meyen, . . . 469 =

ATTALEA.

248². l. Humboldtiana, *Spruce, in Journ. Linn. Soc.* xi. (1869) 163.

248³. *racemosa*, Spruce (l. c.) 166 =

AUBRIETIA.

249¹. a. *Campbellii*, Hort.—Cf. Gard. Chron. (1864) 1133; Lem. Illustr. Hortic. xii. (1865) t. 455 = deltoidea.

AUDIBERTIA.

249². l. *Dorrii*, Kellogg, . . . 190 = incana.

AUGEA.

249². a. PIOTES, Soland. ex Britt. in Journ. Bot. xxii. (1884) 147.

249³. l. AULACOPHYLLUM, Regel, . . . 140 =

AULAYA.

250¹. l. capensis, *Harv. l. c.* 250.—Afr. austr. (non = H. capensis).
scarlatina, . . . = H. scarlatina.

AUSTRALINA.

251.³. l. *flaccida*, Wedd. . . . ix. (1856).
Muelleri, *Wedd.* . . . ix. (1856).

PAG.

AVENA.

252¹. a. PREISSIA, Opiz, Seznam, 79 (1852).

252³. *deusta*, Ball, in Journ. Linn. Soc. xvi. (1878) 719, in obs. = barbata.
l. *elatior*, Linn. . . . = Arrhenatherum avenaceum.

253¹. *hybrida*, Peterm. Fl. Bienitz, 13 =

253². *Loeflingiana*, Cav. . . . 1 = Trisetum Cavanillesii.
longifolia, Thore, Promen. Gascogne, 92 =
mysorensis, Spreng. . . . 337 = Enteropogon melicoides.

253³. *pilosa*, J. Presl, . . . 253 = Preslii.

254². l. *spicata*, Linn. . . . = Danthonia spicata.
a. *spicata*, Willk. & Lange, Prod. Fl. Hisp. i. 71 = Trisetum subspicatum.

254³. a. AVENASTRUM, Jess. Deutschl. Graes. 214 (1863). (Gramin.)
caryophyllum, Jess. l. c. 217 = Aira caryophyllaea.
elatius, Jess. l. c. 215 = Arrhenatherum avenaceum.
flavescens, Jess. l. c. = Trisetum flavescens.
planiculme, Jess. l. c. 216 = Avena planiculmis.
praecox, Jess. l. c. 217 = Aira praecox.
pratense, Jess. l. c. 216 = Avena pratensis.
pubescens, Jess. l. c. = Avena pubescens.

AVERRHOA.

254³. a. *minima*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 101 (Quid?).—China.

255². insere, AXINOPUS, Kunth, Enum. Pl. i. 48 (1833), sphalm. = AXONOPUS, Beauv. = **Paspalum**, Linn. (Gramin.).

AXYRIS.

255². l. *villosa*, Moq. . . . 117 = Kirilowia villosa.

AZALEA.

256¹. a. *alticlarensis* ×, Lindl. Bot. Reg. (1842) t. 27.
balsaminaeflora, Hort. ex Carr. in Rev. Hortic. (1882) 432, cum tab.—Hab.?
l. *bullata*, Forst. ex Hook. Ic. Pl. t. 30.

256². a. *Laetitiae*, Herb. in Bot. Reg. (1845) t. 51.
marginata ×, Paxt. Mag. Bot. ii. (1842) 121.
pulchra, Paxt. (l. c.) i. (1841) 126.
rosea, Hort. ex Loisel. in Duham. Arb. ed. Nov. v. 224. t. 64 = R. nudiflorum.

AZORELLA.

257². l. reniformis, *Pers. Syn.*

BABIANA.

257³. l. densiflora, Ecklonii, lineolata, parviflora, punctata, quadripartita, scabra, et undulato-venosa, . . . xv.
a. *longiflora*, Steud. Nom. ed. II. i. 683, in syn. = Ixia paniculata.

BACCAUREA.

258¹. l. pyrrhodasya, *Muell. Arg.*

BACCHARIS.

260¹. l. Myrsinites, *Pers.*

261². a. tomentosa, *Thunb. Pl. Bras. Dec.* iii. 38.—Bras.

BACTRIS.

262¹. a. leptocarpa, *Trail, ex im Thurn, in Timehri*, iii. (1884) 253.—Guian.

262². megalocarpa, *Trail, ex im Thurn*, (*l. c.*) i. (1882) 242; iii. (1884) 256.—Guian.

BADUSA.

263¹. l. corymbifera, *A. Gray*,

BAECKEA.

263¹. a. DROSODENDRON, M. Roem. Syn. Hesper. 140 (1846).

264¹. a. BAEOCHORTUS, Ehrh. Beitr. iv. 149 (1789) = **Carex**, Dill. (Cyperac.).

BAEOMETRA.

264¹. a. EPIONIX, Rafin. Fl. Tellur. ii. 81 (1836).

BAHIA.

264³. l. Woodhousei, *A. Gray*,

BAHIOPSIS.

264³. l. *lanata*, Kellogg, l. c. = V. lanata.

PAG.

BALANITES.

265[1]. a. *arboreum*, Royle, Illustr. Bot. Himal. 447 (Index) (= *Zygophyllum arboreum*).—Reg. Himal.
rigidum, Royle, (l. c.) (= *Zygophyllum rigidum*).—Reg. Himal.

265[3]. l. BALDELLIA, Parl. . . . = **Echinodorus**, Rich. (Alismac.).
ranunculoides, Parl. . . . = E. ranunculoides.

BALLOTA.

266[1]. l. *frutescens*, Steud. Nom. ed. II. ii. 155 = integrifolia.

266[2]. a. *royleoides*, Benth. . . . 520 = Lasiocorys arabica.
rupestris, Vis. . . . 216 = hirsuta?

BALSAMEA.

266[2]. a. *Commiphora*, Engl. l. c. 41, nomen. = C. Agallocha.
l. *caudata*, Engl. l. c. 42 =
a. *erythraea*, Engl. Bot. Jahrb. i. (1881) 42 = C. erythraea.
l. *Harveyi*, Engl. l. c. = C. Harveyi.
a. *Kataf*, Engl. l. c. = C. Kataf.
mossambicensis, Engl. l. c. = C. mossambicensis.
pubescens, Engl. l. c. = C. pubescens.
Playfairii, Engl. l. c., nomen. = C. Myrrha.

BALSAMODENDRUM.

267[1]. a. *Commiphora*, Royle, Illustr. Bot. Himal. 175 = C. Agallocha.

BALTIMORA.

267[2]. l. Scolospermum, *Steetz*,

BAMBUSA.

268[1]. a. *Matake*, *Metake*, Siebold, in Verh. Batav. Gen. xii. (1830) 4; *Hatsik*, *Jataki*, *Kantsik*, *Kinmeitsik*, *Kurotake*, *Moosoo*, *Narihiratke*, *Nigamatake*, *Sinotake*, *Suwootsik*, *Tootsik*, Siebold, (l. c.) 5; *Ginmeisasa*, *Hoteitsik*, *Kumasasa*, *Sikak'take*, Siebold, (l. c.) 6.—Japon.

BANARA.

268[3]. l. Xyladenius, Desv. . . . Pl.

269[1]. *brevifolia*, . . . Scolopia Dasyanthera.
ibaguensis, *Tul.* . . . 290.

BANISTERIA.

269[2]. a. *brachiata*, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 118 = Billbergiana.

BANKSIA.

270[3]. l. Isostylis, Spach, Hist. Vég. Phan. x. 402 (1840).

BARBACENIA.

272[1]. l. *squamata*, Lindl. . . . 64 = Vellozia squamata.
a. tubiflora, *Benth. & Hook. f. Gen.* iii. 740.—Bras.

BARKHAUSIA.

273[1]. l. *Clausonis*, Pomel, . . . = C. Clausonis.

273[2]. *juvenalis*, Delile, . . .—Cf. Ann. Sc. Nat. Sér. II. vii. (1837) 286 . . .

273[3]. *Ròylei*, DC. . . . 157 = Pterotheca bifida.

BARLAEA.

273[3]. l. *calcarata*, Reichb. f. l. c. = H. calcarata (in Addend.).

BARLERIA.

273[3]. a. alba, *Lodd. Bot. Cab.* t. 360.—Ins. Maurit.

274[1]. elegans, *S. Moore, in Journ. Bot.* xviii. (1880) 269.—Afr. austr.

274[2]. l. limnogeton, *S. Moore*, . . . Afr. trop.

274[3]. a. *polyphylla*, Steud. Nom. ed. II. i. 386, sphalm. = Barbieria polyphylla.
l. *pyramidalis*, Lam. . . . Prod. xi.
a. *pyramidata*, Lam. ex Steud. Nom. ed. II. i. 209, in syn., sphalm. = praec.

BARRINGTONIA.

275[3]. a. Commersona, Sonner. Voy. tt. 8, 9 (1782). Michelia, Linn. Syst. ed. I (1735).
acutangula, Blume, Bijdr. 1097 = insignis.
acutangula, Korth. in Nederl. Kruidk. Arch. i. (1848) 206 = reticulata.
acutangula, Zoll. ex Miq. Fl. Ind. Bat. i. 1. 488 = serrata.

PAG.

BARRINGTONIA:—

275[3]. l. augusta, *Kurz*,
d. Butonica, *Forst.*

276[1]. a. calyptrata, *R. Br. ex Benth. Fl. Austral.* iii. 288, in nota.—Austral.
l. *elongata*, Korth. . . . 206 = macrocarpa.
a. *Levequii*, Jardin, in Mém. Soc. Sc. Cherb. v. (1857) 296, 311 = speciosa.
l. luzonensis, *Vidal*, . . . 43, 113.
luzoniensis, Vidal, l. c. 13 = praec.
a. *racemosa*, Hort. ex Miq. Fl. Ind. Bat. i. 1. 488 = insignis.
l. speciosa, *Forst. Char. Gen.* 76. t. 38; *Linn. f.* . . . 312.—Ins. Pacif.

276[1]. l. BARROTIA, Gaudich. . . . (1848?).

276[2]. a. BARTERIA, Welw. in Bibl. Univ. Genèv. (1861) 200 = **Brasenia**, Schreb. (Nymphac.).
africana, Welw. l. c. = Brasenia Schreberi.

BASELLA.

277[2]. d. *cananifolia*, (err. typ. loco *crassifolia*).

BASILIMA.

277[2]. l. *pygmea*, Rafin. l. c. 76 = S. sorbifolia?

BASSIA.

277[3]. d. *Erskineana*, F. Muell. (vide infra.).

BASSIA.

277[3]. l. *albescens*, Griseb. . . . 164 = Labourdonnaisia albescens.
a. Erskineana, *F. Muell. in Melb. Chem. & Drug.* (Apr. 1885).—N. Guin.

278[1]. *Junghuhniana*, De Vriese, Pl. Reinw. 62 = Payena Junghuhniana.
retusa, Rich, ex A. Gray, in Proc. Am. Acad. v. (1861–62) 327 = Isonandra Richii.

BASSOVIA.

278[1]. a. *anonacea*, Miers, in Ann. & Mag. Nat. Hist. Ser. II. xi. (1853) 94, in obs. = Athenaea anonacea.
hirsuta, Miers, (l. c.) = Athenaea hirsuta.
Martiana, Miers, (l. c.) = Athenaea Martiana.
micrantha, Miers, l. c. = Athenaea micrantha.
mollis, Miers, l. c. = Withania mollis.
novo-friburgensis, Miers, l. c. = Withania novo-friburgensis.
oocarpa, Miers, l. c. = Withania oocarpa.
pogogena, Miers, l. c. = Athenaea picta.
Pohliana, Miers, l. c. = Athenaea Pohliana.
pyrifolia, Miers, (l. c.) = Withania pyrifolia.
Schottiana, Miers, (l. c.) = Athenaea Schottiana.

BATATAS.

278[2]. l. *incurva*, Benth. . . . = I. carnosa.

BATEMANNIA.

278[3]. l. *Meleagris*, Reichb. f. . . . = Zygopetalum Meleagris.

BATRACHIUM.

278[3]. a. *argutum*, Opiz, Seznam, 21 = R. aquatilis.

279[1]. l. *ololeucos*, Bossch, . . . 6 =

279[2]. *trichophyllum*, Bossch, . . . 5;

BATSCHIA.

279[2]. a. *fimbriata*, [Nutt. in] Fras. Cat. n. 11, nomen = L. angustifolium.

BAUHINIA.

279[3]. l. armata, *Thunb. Pl. Bras. Dec.* iii. 33.

280[3]. a. *inermis*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 102 (Quid?).—Archip. Ind.

BEAUFORTIA.

282[2]. a. Manglesia, Lindl. Swan River App. t. 3 (1839).

BECKMANNIA.

282[3]. l. Bruchmannia, Nutt.

BEGONIA.

283[2]. l. Augustia, . . . Berl. 1854 (1855).

283[3]. a. *Ameliae* ×, André, in Rev. Hortic. (1885) 512. f. 89, 90.
l. anemonoides, *Azara, Voy.* ii. 503.—Reg. Andin.

PAG.

BEGONIA:—

284[1]. a. *Chelsoni* ×, Hort. ex T. Moore, Flor. et Pom. (1871) 234, cum fig.

284[2]. l. Cumingiana, *A. DC.* . . . 320.—Ins. Philipp.

285[2]. incarnata, *Link & Otto, Ic. Pl. Rar.*
a. *Ingramii* ×, T. Moore & Ayres, Gard. Mag. Bot. ii. (1850) 153.
intermedia ×, Veitch, ex Van Houtte, in Fl. des Serres, xix. (1873) 39, cum fig.
Leopoldii ×, Lem. Illustr. Hortic. vi. (1859) t. 205.

285[3]. l. Lindleyana, *Warsz.*
magnifica, Warsz. ex Klotzsch, in Abh. Akad. Berl. 1854 (1855) 129 =
Martiana, *Link & Otto, Ic. Pl. Rar.*

286[3]. platanifolia, *Schott*,
a. *prestoniensis* ×, T. Moore, ex Planch. in Fl. des Serres, viii. (1852–53) 189. t. 819.

287[1]. l. *rubro-nervia*, Hort. . . . 40 = rubro-venia.
rufa, *Thunb. Pl. Bras. Dec.* i. 11.

287[2]. semperflorens, *Link & Otto*, . . . 9. t. 5.

287[3]. a. stipularis, *Spreng. in Schrad. Journ.* (1800) ii. 195.—Hab.?
Teysmanniana, *Miq. Fl. Ind. Bat.* i. 1. 1092, in obs.—Sumatra.

288[1]. *Worthiana* ×, Carr. in Rev. Hortic. (1874) 117. f. 15.
zeylanica, *Van Houtte, ex Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 204.—Zeylan.

BEHEN, Moench.

288[2]. l. *angustifolius*, Link, Handb. 245; Schott, etc.
a. *fabarius*, Link, l. c. = S. Cucubalus.
fimbriatus, Link, l. c. = S. fimbriata.
lanuginosus, Link, (l. c.) = S. lanuginosa.
maritimus, Link, l. c. = S. maritima.
Pumilio, Link, (l. c.) = S. Pumilio.
uniflorus, Link, (l. c.) = S. maritima.

BEILSCHMIEDIA.

288[2]. a. Wimmeria, Nees, ex Meissn. in DC. Prod. xv. 1. 65 (1864).
gaboonensis, *Benth. & Hook. f. Gen.* iii. 158.—Afr. trop.
Mannii, *Benth. & Hook. f.* (*l. c.*).—Afr. trop.
minutiflora, *Benth. & Hook. f.* (*l. c.*).—Afr. trop.

289[1]. l. BELLENDENIA, Rafin. in Loud. Gard. Mag. viii. (1832) 245 =

BELLUCIA.

290[1]. l. Webera, J. F. Gmel. . . . (1791).

BELOPERONE.

290[3]. l. longepetiolata, *Nees.*

BERARDIA.

291[3]. l. *phylicoides*, Brongn. l. c. = Raspalia phylicoides.

BERBERIS.

292[2]. a. Chirisigui, *Azara, Voy.* ii. 514.—Boliv.

292[3]. l. laurina, *Thunb. Pl. Bras. Dec.*

293[2]. *Roemeriana*, Scheele, . . . 154, 352 =

294[3]. BERINIA, . . . l. 30 (non 50).
l. *andryaloides*, Brignol. . . . 30 (non 50).

BERKHEYA.

295[1]. a. *incana*, Spreng. ex Sch. Bip. in Flora, xxvii. (1844) 774 (= *Stobaea adenocarpa*).—Afr. austr.
l. *Zeyheri*, Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 429 = subulata.

BERNIERA.

295[3]. l. *nepalensis*, DC. l. c. = G. nepalensis.

BESLERIA.

296[3]. d. Drupina, Linn. (*nunc* **Curanga**).

297[1]. l. crassifolia, *Schott, in Flora*, iv. (1821) 198.—Bras.
grandifolia, *Schott*, (*l. c.*) 197.—Bras.
mollissima, *Spreng. in Schrad. Journ.* (1800) ii. 196.—Am. austr.

297[2]. *serrulata*, Jacq. . . . 21 = Drymonia serrulata.

PAG.
297³. l. BETCKEA, DC. . . . (1829) et Prod. iv. 642 (1830) = Valerianella, Tourn. (Rubiac.).
major, Fisch. & Mey. . . . V. samolifolia.
samolifolia, DC. . . . V. samolifolia.
samolifolia, Gill. . . . = *Gilliesii*.

BETONICA.
298¹. l. *pradica*, Zanted. in Comm. Ateneo Brescia, 1815 (1818), 88

BETULA.
298². l. *antarctica*, Forst. f.
299¹. a. *resinifera*, Royle, Illustr. Bot. Himal. 343, 345 = utilis ?

BEZANILLA.
299³. l. *chilensis*, Remy, . . . 110 = P. chilensis.

BIASOLETTA.
300¹. l. *tuberosa*, Koch, . . . xix.

300². l. **BICORNELLA**, Lindl. . . . (1835) . . . Benth. & Hook. f. iii. 624.

BIDENS.
300³. l. *Bandana*, Wall.
Bardana, Steud. Nom. ed. II. ii. 202, sphalm. = praec.
301¹. graveolens, *Mart. Reise Bras.* i. 286.
301². *pulcherrima*, Sch. Bip. . . . nomen = Cosmos diversifolius.
301³. a. *serratula*, Jardin, in Mém. Soc. Sc. Cherb. v. (1857) 296, sphalm. = serrata.
301³. l. *tuberosa*, Sch. Bip. . . . = Viguiera tuberosa, *Benth. & Hook. f.*

BIFORA.
302¹. a. ANIDRUM, Neck. Elem. i. 188 (1790), nomen prius.

BIGELOWIA.
302³. l. Bigelovii, *A. Gray, l. c.* 642; xvii.
Douglasii, A. Gray, l. c. 645 = *oppositifolia*, A. Gray, . . . 32 = Eupatorium mygindaefolium.

BIGNONIA.
303¹. d. HIERANTHES, Rafin. (*nunc* **Stereospermum**).
ODISCA, Rafin. (*nunc* **Colea**).
l. *alba*, Aubl. . . . = Macfadyena bracteosa.
a. *alliacea*, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 138 = Sagraeana.
303². l. binata, *Thunb. Pl. Bras. Dec.* iii. 35,
303³. *Coito*, Vell. . . . = Macfadyena Coito.
304¹. l. elliptica, *Thunb.* (*l. c.*) 34.
Ferdinandi, Welw. . . . 584 (= *Ferdinandoa superba*).—Afr. trop.
304². a. *lanata*, R. Br. ex Fresen. in Flora, xxi. (1838) 607 = discolor.
304³. l. *myriantha*, Cham. in Linnaea, vii. (1832) 684.
305¹. *populifolia*, DC. . . . 159 = Melloa populifolia.
305³. *Telfairiae*, Boj. . . . t. 2976 = Colea Telfairiae.

BIKKIA.
306¹. a. *tetrandra*, A. Gray, in Proc. Am. Acad. iv. (1860) 307 = grandiflora.

306². l. BILLARDIERIA, Vahl,

BILLBERGIA.
307¹. l. speciosa, *Thunb. Pl. Bras. Dec.* iii. 30.

307². a. BINIA, Noronha, ex Thou. Gen. Nov. Madag. 8 (1806) = **Noronhia**, Stadm. (Oleac.).

BIOPHYTUM.
307³. l. *Umbraculum*, Welw. . . . 590 = sensitivum.

BIPINNULA.
307³. a. *biplumata*, Reichb. f. Xenia Orch. 62, in obs. = bonariensis.

BIXA.
309¹. a. ORLEANIA, Commel. ex Boehm. in Ludw. Def. Gen. Pl. 380 (1760).

BLAKEA.
310³. l. Pyxidanthus, *Triana*,

BLENNOSPERMA.
311². l. UNXIA, Bert. ex Colla,

PAG.
312¹. post **BLEPHARISTEMMA**, Benth. pone
BLEPHARIZONIA, Greene, in Bull. Calif. Acad. i. n. iv. (1885) 279. *COMPOSITAE.*
laxa, *Greene, l. c.*—Calif.
plumosa, *Greene, l. c.*—Calif.

BLEPHAROCALYX.
312¹. l. *cisplatensis*, Griseb. . . . = Eugenia cisplatensis.

BLETIA.
312³. d. DILOMILIS, Rafin. (*nunc* **Octadesmia**).
HECABE, Rafin. (*nunc* **Phaius**).
l. fulgens, *Linden & Reichb. f.*
313¹. l. *guineensis*, G. Don, in Edinb. Phil. Journ. xi. (1824) 344 (*guinense*), nomen = Eulophia guineensis.
a. *Jongheana*, Reichb. f. in Flora, lv. (1872) 158 = Laelia Jongheana.
313². l. Wageneri, *Reichb. f.*

BLETILLA.
313². a. *striata*, Reichb. f. in Bot. Zeit. xxxvi. (1878) 75 = Bletia hyacinthina.

BLIGHIA.
313². a. BONANNIA, Rafin. Specchio, xv. 115 (1814).

313². a. BLISMUS, Montand. Guide Bot. 346 (1868), sphalm. = BLYSMUS, Panz. = **Scirpus**, Tourn. (Cyperac.).

BLOOMERIA.
313³. a. montana, *Greene, in Bull. Calif. Acad.* i. iv. (1885) 281.—Calif.

BLUFFIA.
313³. l. *Eckloniana*, Nees, . . . 447 = P. semialatum.

BLUMENBACHIA.
314³. a. contorta, *Benth. & Hook. f. ex Hieron. Pl. Diaphor.* 120.—Reg. Argent.
contorta, Hook. f. Bot. Mag. t. 6134 = *lateritia*, Benth. & Hook. f. ex Hieron. (l. c.) = Loasa compressa.

BLYSMUS.
315¹. a. *distichus*, Montand. Guide Bot. 346 = S. Caricis.

BLYXA.
315¹. a. HYDROTROPHUS, C. B. Clarke, in Journ. Linn. Soc. xiv. (1873) 8. t. 1.

BOBEA.
315². a. ferruginea, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) II. 211, nomen.—Malaya.
Forsteri, Korth. l. c. 211, 212 = Timonius Forsteri.
fusca, *Korth. l. c.* 213.—Malaya.
Gaertneri, Korth. l. c. = Timonius Forsteri.
setosa, *Korth.* (*l. c.*).—Malaya.

BOEA.
316¹. l. *flocculosa*, C. B. Clarke,

BOEHMERIA.
316³. l. Didymogyne, *Wedd.*
Janagi-itsigo, Siebold,

BOERHAAVIA.
317³. a. DANTIA, Lipp. ex Choisy, in DC. Prod. xiii. II. 449 (1849).
318¹. l. plicata, *Boj. . . . Maurit.* (1841) 43.
318². *spicata*, Choisy, . . . 456 = Senkenbergia erecta.

BOJERIA.
318³. a. lanceolata, *Benth. & Hook. f. Gen.* ii. 334.—Afr. austr.

BOLDUS.
319². l. *chilensis*, Schult. f. l. c. 56 = P. Boldus.

BOLTONIA.
319³. l. incisa, *Benth.* . . . 173.—As. or.
indica, *Benth.* . . . 174.—Malaya; China.

BOMBYCELLA.
320³. l. *bicolor*, Welw. . . . 589 = H. micranthus.

BONAMIA.
321¹. l. *Menziesii*, A. Gray, l. c. = Breweria Menziesii.

PAG.
BONATEA.
321². a. *benghalensis*, Griff. in Calc. Journ. Nat. Hist. vi. (1844) 382 = H. digitata.

322¹. l. BONPLANDIA, Willd. . . . = **Cusparia**, Humb. (Rutac.).
Angostura, Rich. . . . 10 = C. febrifuga.
Candolleana, Spreng. l. c. = C. Candolleana.
cuneifolia, Spreng. l. c. = C. cuneifolia.
Erythrochiton, Spreng. l. c. = C. pentandra.
Fontanesiana, Spreng. l. c. = C. Fontanesiana.
heterophylla, Spreng. l. c. = C. heterophylla.
Lasiostemon, Spreng. l. c. = C. silvestris.
pentagyna, Spreng. l. c. = C. pentagyna.
resinosa, Spreng. l. c. = Rauia resinosa.
trifoliata, Spreng. l. c. = C. febrifuga.

BORAGO.
322³. l. *cretica*, Willd. . . . 778 = Trachystemon creticum.
orientalis, Linn. . . . 138 = Trachystemon orientale.

BORRERIA.
324³. l. *simplex*, Griseb. . . . 350 = S. simplex.
strictissima, DC. . . . 547 = S. strumpfioides.

326². l. BOTRYANTHE, Klotzsch, . . . vii.

BOTRYANTHUS.
326². l. *Kerneri*, Marches. in Boll. Adriat. Soc. Sc. Trieste, vii. (1882) 266 = *speciosa*, Marches. (l. c.) 267 =

BOURRERIA.
327³. l. domingensis, *Griseb. Fl. Brit. W. Ind.* 482.

BOUVARDIA.
328³. a. *semperflorens* ×, Joly, in Wien. Garten-Zeit. (1885) 149. t. 1.

BOWLESIA.
329¹. l. *diversifolia*, Meyen, Reise, i. 447 =

BRACHIARIA.
329². l. *eruciformis*, Griseb. l. c. = P. eruciforme.

BRACHYCOME.
330². l. *xanthocomoides*, Less. Syn. Comp. 192; et in

BRACHYLOMA.
331¹. a. leucomallon, *Hanst. in Linnaea*, xxix. (1857–58) 528.—N. Granat.
Moritzianum, *Hanst. l. c.* 524.—N. Granat.
rhodomallon, *Hanst.* (*l. c.*) 526.—Venezuela.
ventricosum, *Hanst.* (*l. c.*) 528.—Hab. ?

BRACHYPODIUM.
331². d. HEMIBROMUS, Steud. (*nunc* **Glyceria**).

BRACHYRAMPHUS.
332¹. a. *Schimperi*, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 283, 307 = Sonchus Spachii.

BRACHYRHYNCHOS.
332¹. l. *eupatorioides*, DC. . . . 439 = S. eupatorioides.
a. *pinceus*, Sch. Bip. in Flora, xxvii. (1844) 702, sphalm. = *junceus*.

BRACHYSPATHA.
332². l. *Konjac*, C. Koch, . . . (1855) 166, etc.

BRAGANTIA, Vand.
333². l. *Vandelli*, Mart. Reise Bras. 280.

BRANDESIA.
333³. l. *elongata*, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 313 = *porrigens*, Mart. (l. c.) 314 = *serpyllifolia*, Mart. (l. c.) =

BRASENIA.
333³. a. BARTERIA, Welw. in Bibl. Univ. Genèv. (1861) 20.

BRASSAVOLA.
334¹. l. grandiflora, *Lindl. l. c.* 16.—Ind. occ.
334². a. stricta, *Hort.—Cf. Gard. Chron.* (1885) I. 501. f. 92 (sp. dub.).—Hab. ?

PAG.

BRASSIA.

334³. l. spathacea, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 279.

BRASSICA.

334³. a. CONIROSTRUM, Dulac, Fl. Hautes-Pyr. 195 (1867).
l. *antarctica*, Molina, . . . 292 = Sisymbrium magellanicum.
apula, Janka, l. c. 180 = Diplotaxis apula.
335¹. a. *corsica*, Jord. Esp. Vég. Affin. 7 = insularis.
335². l. *foliosa*, Hook. f. & Anders. in Hook. f. Fl. Brit. Ind. i. 157 = alba.
fragilis, Sieber, . . . = Diplotaxis Sieberi.
magellanica, Pers. . . . 207 = Sisymbrium magellanicum.
335³. *nigra*, Koch, . . . 713 = sinapoides.
oxyrrhina, *Willk. & Lange, Prod. Fl. Hisp.* iii. 854.
a. *patens*, Murr. ex Moench, Meth. 251 = Sisymbrium patens?
l. Pseudoerucastrum, *Brot.*
336¹. a. sinapioides, *Roth, Man.* ii. 957; *et ex Mert. & Koch, Deutschl. Fl.* iv. 713.—Orb. vet. (Cosmop. cult.).
Sinapis, Noul. Fl. Sous-Pyr. 32 = sinapioides.
l. *Sinapis*, Vis. Fl. Dalm. iii. 136 = B. Sinapistrum.

BREWERIA.

337². a. intermedia, *Hochst. in Flora*, xxvii. (1844) *Bes. Beil.* 8.—Arab.
l. latifolia, *Hochst.* (*l. c.*); *Benth.*
337³. *minima*, A. Gray, . . . 228 = Convolvulus pentapetaloides.
virgata, *Hochst.* (*l. c.*); *Vatke*,

BREYNIA.

338¹. a. patens, *Rolfe, in Journ. Bot.* xx. (1882) 359.—Ins. Formosa.

BRICKELLIA.

338². a. reticulata, *A. Gray, Pl. Wright.* i. 84.—Mexic.

BRIZOPYRUM.

339³. a. *mucronatum*, Aitch. Cat. Punj. Pl. 169 (Aeluropi sp.?).—Ind. or.
mucronatum, Nees, in Wall. Cat. n. 8898 = Eragrostris mucronata, *Benth. & Hook. f.*
l. *obtusiflorum*, Fourn. . . . 579 = Uniola paniculata.
a. *unioloides*, Munro, in Journ. Linn. Soc. vi. (1861) 45 = Eragrostis Uniolae.

BRODIAEA.

340¹. a. TULOPHOS, Rafin. Fl. Tellur. iii. 71 (1836).

BROMELIA.

340². a. Forsstroemii, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856), 110, in nota.—Ins. Guadeloup.
340³. *Pigna*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 103 = Ananas sativa.
Mai-pouri, Perr. (l. c.) = Ananas sativa.
strobilina, *Beurl.* (*l. c.*).—Panama.

BROMIDIUM.

341¹. l. *anomalum*, Doell, . . . 103 = Deyeuxia anomala.

BROMUS.

341¹. a. CALLIAGROSTIS, Ehrh. Beitr. iv. 146 (1789).
341². l. *bohemicus*, F. W. Schmidt, in Meyer, Samml. Phys. Aufs. i. (1791) 189 =
341³. *decumbens*, Koel. . . . = Triodia decumbens.

BROSIMUM.

344². l. Galactodendron, *D. Don, in Edinb. N. Phil. Journ.* (Oct.–Dec. 1829) 171.—
spurium, Sw. . . . 12 = Pseudolmedia havanensis.

BROTERA.

344². l. *Leprieurii*, Guill. & Perr. l. c. = M. abyssinica.

BROWALLIA.

345¹. l. *humifusa*, Forsk. . . . = Cycnium humifusum.
Jamesonii, Benth. . . . = Streptosolen Jamesonii.

BRUCEA.

345². d. NIMA, Buch.-Ham. (*nunc* **Picrasma**).

PAG.

BRUNFELSIA.

345³. a. MARTIA, Lacerda, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 26 (1862).

346³. l. **BRUNSVIGIA**, Heist. Disc. Brunsvig. 1 (1753); Beschr.

BRYA.

347¹. a. NEFRAKIS, Rafin. Sylva Tellur. 69 (1838).

BRYANTHUS.

347¹. a. PHYLLODOCE, Salisb. Parad. Lond. t. 36 (1806).

BRYONIA.

348². a. *Tajuja*, [Mart.] Hort. Reg. Monac. Seminifer, (1836).—Cf. Linnaea, xii. (1838) Litt. 86 = seq.

BUBON.

348³. l. *hypoleucus*, Meissn. . . . 536 = S. hypoleucus.
pimpinellifolius, Eckl. & Zeyh. . . . . 353 (Annesorhizae sp.).
a. *tortuosus*, Hoffmgg. & Link, Fl. Port. ii. 452 = Pimpinella villosa.

BUCHNERA.

349³. l. MICALIA, Rafin. (*nunc* **Escobedia.**).
350¹. l. *humifusa*, Vahl, . . . 81 = Cycnium humifusum.

BUCHOLZIA, Mart.

350³. l. *lupulina*, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 315 =
philoxeroides, Mart. (l. c.) 316 =

BUCKLEYA.

350³. a. lanceolata, *Miq. Cat. Mus. Bot. Lugd. Bat.* 79. —China; Japon.
Quadriala, Benth. & Hook. f. Gen. iii. 227 = lanceolata.

BUDDLEIA.

351¹. a. TOXINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4.

352¹. l. BUEKIA, Nees, . . . 300 = **Tetraria**, Beauv. (Cyperac.).
punctoria, Nees, l. c. (= *Elynanthus punctorius*) (in Addend.).

BUETTNERIA.

352². a. WATSONIA, Boehm. in Ludw. Def. Gen. Pl. 278 (1760).
l. *dasyphylla*, J. Gay, in Mém. Mus. Par. x. (1823) 200;
hermanniaefolia, J. Gay, (l. c.) 204;
353³. a. *triplinervia*, Bivona, ex Baill. Adansonia, x. (1871) 106 = Ropalocarpus triplinervius.

353². l. BUGULA, Tourn. ex Linn. Syst. ed. I (1735); et ex

BULBOCODIUM.

354¹. l. *autumnale*, Lapeyr.
dentatum, Schur, . . . 166 = ruthenicum.
Fontanesii, Heynh. . . . 81 = Merendera filifolia.
lusitanicum, Heynh. . . . 81 = Merendera Bulbocodium.
persicum, Regel, . . . t. 1058 = Merendera persica.
robustum, Bunge, . . . 491 = Merendera robusta.
vernum, *Linn. Sp. Pl.* 294.—Europ.

BULBOPHYLLUM.

354². a. SARCOPODIUM, Lindl. in Lindl. & Paxt. Flow. Gard. i. (1850) 155, partim.
l. biflorum, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 488; *et* . . . 397.
355¹. *lasiochilum*, Par. . . . 153 = Cirrhopetalum lasiochilum.
355². l. membranaceum, *Teysm. & Binn.* (*ut supra*) 481; *et* . . . 397.

BULBOSTYLIS, Kunth.

356³. a. *stenophyllus*, Nees, ex Boeck. in Linnaea, xxxvi. (1869–70) 752.—Am. bor.?

356³. insere BULGA, Linn. Syst. ed. I (1735) = BUGULA, Tourn. = Ajuga, Linn. (Labiat.).

PAG.

BULLIARDA.

356². l. *brevifolia*, Eckl. & Zeyh. . . . 290 = T. brevifolia.
elatinoides, Eckl. & Zeyh. (l. c.) = T. elatinoides.
356³. *trichotoma*, Eckl. & Zeyh. (l. c.) = T. trichotoma.

BUMELIA.

356³. a. ROBERTIA, Scop. Introd. 154 (1777).
d. SPONDOGONA, Rafin. (*nunc* **Dipholis**).
a. *Berterii*, Balb. ex Griseb. Cat. Pl. Cub. 56 = Schefferia Berterii.
confertiflora, [Nutt. in] Fras. Cat. (1813) n. 12, nomen (Quid?).—Am. bor.

BUPHTHALMUM.

359¹. l. *stenophyllum*, Link, . . . = Odontospermum stenophyllum.

BUPLEURUM.

359². a. *brassicaefolium*, Arv. Monog. Pilos. 50 (Quid?). —Gallia.
l. *difforme*, Linn. . . . 238 = Rhyticarpus difformis.

BURMANNIA.

361². a. *bifaria*, Royle, Illustr. Bot. Himal. 373 = coelestis?

BURSA.

361³. l. *Pastoris*, [Weber, in]

BURSERA.

362¹. l. Aloëxylon, bicolor, bipinnata, cuneata, jorullensis, penicillata, Schiedeana, (omnes) *Engl. Bot. Jahrb.* i. (1881) 44; *et*
Galeottiana, grandifolia, Karsteniana, lancifolia, ovalifolia, (omnes) *Engl. Bot. Jahrb.* i. (1881) 43; *et*
obovata, Turcz. . . . 614 = Protium Copal.
Sellowii, Turcz. . . . 613 = Protium Icicariba.

BURSINOPETALUM.

362². a. *Gardnerianum*, Baill. Adansonia, iii. (1862–63) 83 = M. arborea.

BUTOMOPSIS.

363¹. l. lanceolata, *Kunth*, . . . 165.—Geront. trop.; Austral.

BUTONICA.

363¹. l. *Rumphiana*, Miers, . . . = Barringt. speciosa.

BUXUS.

363². l. cordifolia, *Spreng. l. c.*—Ind. occ.
a. *fruticosa*, Borkh. in Rhein. Mag. i. (1793) 544 = sempervirens.
l. *Harlandi*, Hance, . . . 123 = sempervirens.
japonica, Muell. Arg. . . . 23 = microphylla.
microphylla, *Sieb. & Zucc.* . . . 142.—China; Japon.
stenophylla, Hance, . . . 331 = sempervirens.

BYBLIS.

363². a. DROSANTHUS, R. Br. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 305.
DROSOPHORUS, R. Br. ex Planch. (l. c.) 306 (1848).
l. *caerulea*, R. Br. . . . = liniflora.
filifolia, Planch. . . . = liniflora.

CACALIA.

365². l. *Arbuscula*, Thunb. . . . = Othonna Arbuscula.
candicans, Vell. . . . = S. candicans.
365³. a. *flava*, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 333 = Gynura nepalensis.
366¹. l. *ilicifolia*, H. B. & K. . . . 163 = Gynoxys ilicifolia.

CACHRYS.

367². l. *pterochlaena*, DC. . . . 237 = Hippomarathrum siculum?

CACOUCIA.

367³. a. *coccinea*, M. Laws. in Oliver, Fl. Trop. Afr. ii. 434 = velutina.
racemosa, Steud. Nom. ed. II. i. 836, sphalm. = Lumnitzera racemosa.

CACTUS.

367³. a. *Ayrampe*, Azara, Voy. ii. 526.—Reg. Andin.
Berteri, Colla, in Mem. Acc. Torin. xxxvii. (1834) 77. t. 17. f. 2 = Echinocactus Berterii.

PAG.

CACTUS:—
368[1]. l. *ferox*, [Nutt. in] Fras. Cat. (1813) n. 23, nomen; Nutt.
368[2]. l. *fragilis*, [Nutt. in] Fras. Cat. (1813) n. 24, nomen; Nutt.
a. *horridus*, Colla, (ut supra) 76. t. 17. f. 1 = Echinocactus horridus.
369[2]. *viviparus*, [Nutt. in] Fras. Cat. (1813) n. 22, nomen; Nutt.

CADABA.
369[3]. a. *farinosa*, Fenzl, in Webb, Fragm. Fl. Aethiop. 24 (=*Streblanthus Fenzlii*).—Afr. trop.

CADIA.
369[3]. a. MOZAMBE, Rafin. Sylva Tellur. 112 (1838).

370[1]. a. CAËLA, Adans. Fam. ii. 209 (1763) = **Torenia**, Linn. (Scrophul.).

CAESALPINIA.
370[2]. l. *Berterii*, Spreng. . . . 370 = bijuga.
370[3]. a. *laevigata*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 104 (Quid?).—Ins. Philipp.
371[1]. l. *pectinata*, Cav. Desc. 467 = tinctoria.
a. *pilosa*, Ruiz, ex G. Don, Gen. Syst. ii. 434 = Hoffmanseggia pilosa.

CAIOPHORA.
371[3]. l. *lateritia*, Benth. . . . iii. t. 119 = B. lateritia?

372[1]. l. **CALACANTHUS**, T. Anders.

CALADIUM.
373[2]. l. *petiolatum*, Hook. . . . t. 3728

CALAMAGROSTIS.
374[3]. l. *Humboldtiana*, Steud. . . . 250 = Deyeuxia stricta.
montana, Host, Gram. iv. t. 46.
375[1]. *pumila*, Hook. f. . . . 176 = Leptochloa Lindleyana.
375[2]. *sylvatica*, Host, . . . 48; DC. . . . Suppl. 253 = varia.

CALAMINTHA.
375[3]. l. Clinopodium, *Spenn. Handb.* ii. 429; *Benth. Fenzlii*, Vis. . . . 7 = Mentha Pulegium.
376[1]. a. *pauciflora*, Lange, in Kjoeb. Vidensk. Meddel. (1863) 10 = officinalis.

CALAMOSAGUS.
376[2]. l. in syn. K. polystachya (bis) et K. scaphigera.

CALAMUS.
377[3]. a. Regis-Daniae, *Kamphoev. ex Rink, Nicob. Ins.* 132, nomen.—Ins. Nicobar.

CALANTHE.
379[2]. l. *speciosa*, Vieill. . . . 44 = Phaius grandifolius.

379[2]. a. CALANTHEA, Miers, in Proc. Roy. Hort. Soc. iv. (1864) 161 = **Capparis**, Tourn.
pulcherrima, Miers, l. c. = Capp. pulcherrima.

CALATHEA.
379[3]. a. Baraquini, *Regel, Gartenfl.* xviii. (1869) 99.—Bras.
Baraquiniana, Regel, l. c. xxviii. (1879) 301 = praec.
380[1]. l. *macilenta*, Lindl. . . . t. 1210 = Myrosma cannaefolium.
Myrosma, Koern. . . . 87 = Myrosma cannaefolium.
a. *strobilacea*, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) 1. 111, sphalm. = praec.
380[2]. varians, *Koern.* (*l. c.*) 136.—Am. trop.

CALCEOLARIA.
381[1]. l. fuchsiaefolia, *Hemsl. in The Garden* (1879), 258.—Hab.?
381[3]. *nudicaulis*, Meyen, Reise, i. 355, nomen; et
382[1]. *verbascifolia*, Bert. . . . nomen = andina.

382[1]. insere CALCEOLARIA, Loefl. It. Hisp. 183 (1758) = **Ionidium**, Vent. (1803) (Violar.).

CALCITRAPA.
382[2]. a. *Jacea*, Peterm. Fl. Bienitz, 48 = Cent. Jacea.
pratensis, Peterm. (l. c.) 49 = Cent. pratensis.
tinctoria, Roehl. Deutschl. Fl. ed. II. ii. 436, in obs. = Carthamus tinctorius.

PAG.

CALEA.
383[1]. l. *spectabilis*, Labill. . . . 186 = Cassinia spectabilis.
Suriani, Cass. . . . 33 = Neurolaena lobata.

383[2]. a. CALEATIA, Mart. ex Steud. Nom. ed. II. ii. 76, in syn. (1841) = **Lucuma**, Molina (Sapotac.).
Caimito, Mart. l. c. = L. Caimito.

CALIMERIS.
384[2]. l. *flexuosa*, Royle, Illustr. Bot. Himal. 251. t. 58. f. 2 =

CALINEA.
384[2]. a. *Bentegeati*, Walp. & Duchass. ex Eichl. in Mart. Fl. Bras. xiii. 1. 71 (=*Pinzona calineoides*).—Ins. Guadeloup.

CALIPHRURIA.
384[3]. l. *subedentata*, Baker,

CALLA.
384[3]. a. CALLAION, Rafin. New Fl. Am. ii. 90 (1836).
l. *nitida*, Jack, . . . 24 = Aglaonema oblongifolia.
oblongifolia, Roxb. . . . 516 = Aglaonema oblongifolia.

385[1]. a. CALLAION, Rafin. New Fl. Am. ii. 90 (1836) = **Calla**, Linn. (Aroid.).
bispatha, *brevis*, *heterophylla*, *palustris*, Rafin. l. c. = Calla palustris.

385[1]. a. CALLIAGROSTIS, Ehrh. Beitr. iv. (1789) 146 = **Bromus**, Dill. (Gramin.).

CALLICOMA.
387[1]. l. Stutzeri, *F. Muell.* . . . 31

CALLIGONUM.
387[1]. d. TRAXILISA, Rafin. (*nunc* **Tetracera**).
387[3]. a. WACHENDORFIA, Loefl. It. Hisp. 177 (1758).

388[3]. a. CALLITHRONUM, Ehrh. Beitr. iv. (1789) 148 = **Cephalanthera**, Rich. (Orchid.).

CALLITRICHE.
389[1]. l. *intermedia*, Hoppe,
389[2]. *spuria*, Rafin. . . . 354 = verna.

389[2]. l. **CALLITRIS**, Endl. Cat. Hort. Vindob. i. 209 (1842); et

389[3]. a. CALLOBUXUS, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 372 = **Tristania**, R. Br. (Myrtac.).

CALODENDRUM.
390[2]. l. capense, *Thunb.*

CALODRACON.
390[2]. l. *Sieberi*, Planch. . . . 132 = Cord. terminalis.

CALOGYNE.
390[3]. l. *Distylis*, F. Muell.

CALONYCTION.
390[3]. l. *comospermum*, Boj.

CALOPHANES.
391[1]. l. Hildebrandtii, *S. Moore*, . . . (1880) 8.—
391[2]. thunbergiaeflora, *S. Moore*, . . . (1880) 8.—

CALOPHYLLUM.
391[3]. l. *Bintagor*, Roxb.

CALOSCORDUM.
392[3]. a. *exsertum*, Herb. l. c. (1847) sub t. 5 = Allium Bakeri.
l. *neriniflorum*, Herb. l. c. (1844) Misc. 67.

CALTHA.
394[1]. a. *napalensis*, Royle, Illustr. Bot. Himal. 22 = palustris.
himalayanus, Royle, (l. c.) 44 = *himalensis*.
pusilla, Herb. Lamb. ex Pursh, Fl. Am. Sept. ii. 390 = natans.

CALYCADENIA.
394[2]. l. *plumosa*, Kellogg, . . . 49 = Blepharizonia plumosa.

PAG.

CALYCERA.
394[3]. a. *andina*, Miers, Trav. Chil. ii. 531; et in Ann. & Mag. Nat. Hist. Ser. III. iv. (1860) 186 (=*Nastanthus agglomeratus*).—Chili.
l. herbacea, *Cav.*
a. lanata, *Benth. & Hook. f. Gen.* iii. 102.—Chili.

CALYCOPTERIS.
395[2]. a. Joan, *Siebold, in Verh. Batav. Gen.* xii. (1830) 23.—Japon.

CALYPTRANTHES.
396[3]. *nitida*, Raeusch. . . . 144 = Eugenia lucida, *Banks*.

CALYSTEGIA.
398[2]. l. *villosa*, Kellogg, . . . 17 = Convolvulus villosus.

CAMASSIA.
399[2]. l. CYANOTRIS, Rafin. . . . Mag. (1818) 356; QUAMASIA, Rafin. in Am. Monthly Mag. (1818) 265;

CAMELINA.
399[3]. a. incana, *F. & C. Presl, Fl. Čech.* 134.—Europ.

CAMELLIA.
400[1]. l. *oleifolia*, Wall.

400[2]. a. CAMERARIA, Boehm. in Ludw. Def. Gen. Pl. 56 (1760) = **Hemerocallis**, Linn. (Liliac.).

CAMERARIA.
400[2]. a. havanensis, *Benth. & Hook. f. Gen.* ii. 705.—Cuba.

CAMOENSIA.
400[2]. a. GIGANTHEMUM, Welw. Apont. 585 [1859], nomen prius.

CAMPANULA.
401[3]. l. *clivosa*, Banks, . . . = Wahlenbergia angustifolia.
404[2]. *nitida*, [Soland.] . . . 346 = pyramidalis?
405[1]. *porosa*, Thunb. . . . 39 = Samolus porosus.
406[1]. a. *Smithii* ×, T. Moore, Flor. & Pomol. (1875) 209.
406[3]. *thalictriflora*, Steud. Nom. ed. II. ii. 271, sphalm. = seq.
l. *thalictrifolia*, Spreng.

CAMPANUMOEA.
407[2]. l. *lanceolata*, Sieb. & Zucc. . . . 91 = Codonopsis lanceolata.

407[3]. l. CAMPHORATA, Tourn. ex Ludw. Def. Gen. Pl. 230 (1747) = **Camphorosma**, Linn. (Chenop.).

CAMPHOROSMA.
407[3]. l. CAMPHORATA, Tourn. ex Ludw. Def. Gen. Pl. 230 (1747).

CAMPNOSPERMA.
408[1]. a. CYRTOSPERMUM, Benth. in Hook. Kew Journ. iv. (1852) 13.
DREPANOSPERMUM, Benth. in Benth. & Hook. f. Gen. i. 425 (1862).

CAMPSIDIUM.
408[2]. l. *chilense*, Reiss. & Seem. . . . 147 = T. valdiviana.

410[1]. a. CANANGIUM, Baill. Hist. Pl. i. 213, in nota (1868) = praec.

CANARIUM.
410[2] a. STRANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4.

CANAVALIA.
411[1]. d. *incurva*, DC.
l. *Moneta*, Welw. . . . 62 = obtusifolia.

CANNA.
412[2]. a. *Annaei*, Hort. Par.—Cf. Rev. Hortic. (1861) 470. f. 111 = indica.
413[1]. a. textoria, *Noronha, Prod. Phytol.* (*in Thou. Mélang.*) (1811).—Madag.

413[1]. l. CANNABINA, Tourn. ex Linn. Amoen. Acad. i. 415 (1749) = **Datisca**, Linn.

PAG.

CANSCORA.

413[2]. l. FLEMINGIA, Roxb. ex Wall. Cat. sub nn. 4 61, 4364 (1830?).

CANTHAROSPERMUM.

413[3]. a. *paucifolium*, Royle, Illustr. Bot. Himal. 192, sphalm. = *pauciflorum*.

413[3]. l. CANTHIOPSIS, Seem. . . . (1866)

CANTHIUM.

413[3]. l. *acuminatum*, D. Dietr. . . . 779 (= *Gynopachis acuminata*).—Java.

414[3]. a. *sinense*, Steud. Nom. ed. II. i. 663, in syn. = *chinense*.

CAPELLENIA.

415[1]. l. *moluccana*, Teysm. & Binn. . . . l. c. = E. moluccanum.

415[3]. l. CAPNOREA, Rafin. . . . = **Hesperochiron**, S. Wats. (1871) (Hydrophyll.).
nana, Rafin. l. c. = H. californicus.

CAPPARIS.

415[3]. a. CALANTHEA, Miers, in Proc. Roy. Hort. Soc. iv. (1864) 161.
d. TRICLANTHERA, Rafin. (*nunc* **Crataeva**).

CAPRIFOLIUM.

418[2]. l. *germanicum*, Roehl. Deutschl. Fl. ed. II. ii. 176 =
perfoliatum, Roehl. (l. c.) ed. II. ii. 176 =

CAPSELLA.

418[3]. l. *Berengeriana*, Hort. . . . 234 = elliptica.
elliptica, *C. A. Mey*. . . . 199.—Reg. temp.
procumbens, Fries, . . . 14 = elliptica.

419[2]. l. **CAPURA**, Blanco, . . . 264 (1837).
d. OTOPHORA, Blume, (*nunc* genus proprium).
a. *alata*, Teijsm. & Binn. Cat. Hort. Bogor. 214 = Otophora alata.
a. multijuga, *Benth. & Hook. f. Gen.* i. 405.—Borneo.
spectabilis, Teijsm. & Binn. l. c. = Otophora spectabilis.
Zollingerianus, Teijsm. & Binn. l. c. 215 = Otophora Zollingeriana.

CARAGANA.

419[3]. *sylvatica*, Hoffmgg. Nachtr. Preisverz. (1842) 28.—Cf. Linnaea, xvi. (1842) Litt. 282 = frutescens?

CARALLIA.

420[2]. l. *Symmetria*, Blume,

420[2]. l. CARAMANICA, Tineo, Pl. Rar. Sic. fasc. i. 8 (1846).

CARDAMINE.

421[1]. l. *bracteata*, S. Moore, . . . 130 = Eutrema Wasabi.
chelidonioides, S. Moore, . . . 130 = C. Tanakae.

421[3]. *Heyneana*, Welw. . . . 676 (*Hayneana*).

421[3]. javanica, *Miq*. . . . 17. t. 10.

422[2]. a. rotundifolia, *Michx. Fl. Bor. Am.* ii. 30.—Am. bor.

423[2]. a. CARDIOPHYLLUM, Ehrh. Beitr. iv. (1789) 148 = **Listera**, R. Br. (Orchid.).

CARDIOSPERMUM.

423[3]. a. RHODIOLA, Lour. Fl. Cochinch. 627 (1790).
l. *triphyllum*, Vell. . . . 25 = Urvillea triphylla.

CARDUNCELLUS.

424[1]. a. *lanatus*, Moris, Fl. Sard. ii. 439 = Carthamus lanatus.

CARDUUS.

424[2]. l. *arenarius*, DC. . . . 457 = Decandollei.

424[3]. *Candollei*, DC. Prod. vi. 625 = Decandollei.

425[1]. Decandollei, *Moretti, in Brugnat. Giorn. Fis. Dec.* II. v. (1822) 45.—Europ.

425[3]. *litigiosus*, Nocca & Balb. . . . 12 = Decandollei.
montanus, Suffr. . . . 625 = Decandollei.

426[2]. *Sanctae-Balmae*, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 426 = Decandollei.

PAG.

427[1]. insere CARELIA, Cav. in Anal. Cienc. Nat. vi. (1802) 317 = **Mikania**, Willd. (1803) (Compos.).
saturiaefolia, Cav. l. c. = M. amara.

CAREX.

427[1]. a. AGASTACHYS, Ehrh. Beitr. iv. (1789) 146.
BAEOCHORTUS, Ehrh. (l. c.) 149 (1789).
CHORDORRHIZA, Ehrh. (l. c.) 148 (1789).
l. PHYLLOSTACHYS, Torr. in Ann. Lyc. N. York, iii. (1836) 404, in obs.

428[2]. a. *bengalensis*, Thw. Enum. Fl. Zeyl. 355 = indica.
l. *Benthamiana*, Boott, . . . Himal. 412 = notha.

428[3]. *brevicaulis*, Thou. Fl. Trist. d'Acugn. 35 = Uncinia brevicaulis.

429[1]. *Brownii*, Tuckerm. . . . 15 (non 21).

430[3]. *divisa*, Oed. . . . = Oederi.

431[1]. *excurva*, Boott, . . . 57 = nemostachys.

431[2]. *ferruginea*, Stev. . . . Nat.
filifolia, *Nutt*. . . . 204.—Am. bor.
flacca, Schreb. Spicil. App. n. 669.

431[3]. fusca, *All*. . . . 269.—Europ.

432[2]. Halleriana, *Asso, Stirp. Arag.* 133. t. 9.
a. *hamosa*, Thou. Fl. Trist. Acugn. 35 = Uncinia brevicaulis.

433[2]. l. lagopodioides, *Schkuhr*, . . . 20.

434[1]. *lithosperma*, Linn. . . . XII. 618; Murr. Syst. ed. XIII. 706.
a. *longiaristata*, Boott, ex Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 350 = indica.

435[1]. *Moritzii*, Steud. in Zoll. Verz. Ind. Archip. ii. 60; Miq. Fl. Ind. Bat. iii. 350 = indica.

435[2]. *Nenningiana*, Opiz, Seznam, 103 (err. typ. 203) (Quid?).—Europ.

436[2]. parasitica, *Kunze, in Schkuhr, Riedgr. Suppl.* i. 1. 83, in nota.—Abyss.
pendula, Schreb. Spicil. 62 = glauca.

437[1]. Pseudo- et *Pseudo-* (passim).

437[2]. psilochloa, *Wall. Cat.* n. 3534.—Burma.

438[3]. l. *simpliuscula*, Wahlenb. . . . 141 = Kobresia caricina.

439[1]. a. stramentitia, *Boott, ex Boeck. in Linnaea*, xl. (1876) 351.—Reg. Himal.
l. *strictifolia*, Opiz, Seznam, 103 (err. typ. 203) = acuta.

439[3]. a. *tenuissima*, Schur, Enum. Pl. Transs. 704 = muricata.
l. tribuloides, *Wahlenb*. . . . 145.—Am. bor.

440[3]. a. *xanthosperma*, Dewey, in Am. Journ. Sc. Ser. II. xlii. (1866) 334 = flaccosperma.

CARICA.

441[1]. l. VASCONCELLEA, A. St. Hil. . . . 12.
a. hastata, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 77.—Bras.
lanceolata, *Benth. & Hook. f. ex Hieron. Pl. Diaph.* 122.—Reg. Argent.
l. Posoposa, *Linn*.
a. quercifolia, *Benth. & Hook. f. ex Hieron.* (*l. c.*).—Reg. Argent.

CARISSA.

441[2]. l. *oblongifolia*, Hochst. . . . 827 = Acokanthera oblongifolia.
Schimperi, A. DC. . . . 675 = Acokanthera Schimperi (in Addend.).
a. *sepiaria*, Royle, Illustr. Bot. Himal. 15, nomen = C. spinarum.

CARLUDOVICA.

442[1]. l. *angustifolia*, Ruiz & Pav. l. c. 292 = Cyclanthus Plumierii.
a. *angustifolia*, Seem. ex Griseb. Fl. Brit. W. Ind. 513 = Oerstedii?
a. gracilis, *Liebm. ex Griseb. Fl. Brit. W. Ind.* 513.—Ind. occ.; Mexic.

442[2]. jamaicensis, *Lodd. ex Sweet, Hort. Brit.* ed. III. 630 (sp. dub.).—Ind. occ.
palmaefolia, Sweet, (l. c.) = Plumerii.
Plumerii, Griseb. in Goett. Abh. vii. (1857) 259 = gracilis.
l. *Plumieri*, Griseb. . . . 513 = Plumerii.

CARPINUS.

443[3]. a. *lacera*, Royle, Illustr. Bot. Himal. 449, sphalm. = *Corylus lacera*.

PAG.

CARPOPOGON.

444[2]. l. *anguineum*, Roxb. ex Sweet, Hort. Brit. ed. I. 119, nomen; et

CARRICHTERA.

444[3]. l. Vellae, *DC*.

445[2]. l. **CARUM**, Rupp.
a. TRAGIOPSIS, Pomel, Nouv. Mat. Fl. Atl. 139. (1874).

445[3]. l. Lomatocarum, *Boiss*.
a. *nigrum*, Royle, Illustr. Bot. Himal. 229 (Quid?).—Reg. Himal.

446[1]. l. *Panatjan*, Baill. . . . = Pimpinella Pruatjan.

CARYA.

446[2]. l. texana, *Buckl. in Proc. Acad. Sc. Philad.* 1860 (1861) 444; *C. DC*.

CARYOPHYLLUS, Linn.

447[1]. l. *pterocarpus*, Vieill. . . . 256 = E. pterocarpa.
a. *rugosus*, Blume, ex Miq. Fl. Ind. Bat. i. 1. 437 = E. zeylanica.

CARYOSPERMUM.

447[1]. l. arborescens, *F. Muell. Fragm.* vi. 202; viii. 279.

CARYOTA.

447[1]. a. THUESSINKIA, Korth. ex Miq. Fl. Ind. Bat. iii. 41 (1855).

447[2]. l. mitis, *Lour*. . . . 569; *ed. Willd*. 697.

CASEARIA.

447[3]. l. GUIDONIA, Plum. ex Adans. Fam. ii. 305 (1763).

448[1]. *Cheela*, Royle, Illustr. Bot. Himal. 170 = tomentosa.

448[3]. *pauciflora*, Royle, (l. c.) = tomentosa.
piscidia, Buch.-Ham. ex Royle, (l. c.) = tomentosa.

449[2]. l. CASPIA, Pison. ex Scop. . . . = **Vismia**, Vand.

CASSIA.

449[3]. l. GRIMALDIA, Schrank, in Allg. Bot. Biblioth. iv. (1805) 184.
a. *acacalis*, Royle, in Journ. As. Soc. Beng. i. (1832) 462 (err. typ. 452) = C. Absus.

445[1]. *artemisiaefolia*, R. Br. in Sturt, Centr. Austral. ii. App. 77 = artemisioides.
l. *aversiflora*, Hook. . . . t. 2638 =

450[2]. *canaliculata*, R. Br. . . . ii. App. 77 =

450[3]. l. *conspicua*, G. Don, in Edinb. Phil. Journ. xi. (1824) 344; Vog.

451[3]. a. *glaucescens*, A. Cunn. ex R. Br. in Sturt, Centr. Austral. ii. App. 77 = artemisioides?

453[1]. l. phyllodinea, *R. Br*. . . . ii. *App*. 78.
platypoda, R. Br. . . . ii. App. 78 =

454[1]. Sturtii, *R. Br*. . . . ii. *App*. 77.

CASSIPOUREA.

455[2]. l. *gummifera*, Tul. . . . 123 = Dactylopetalum sessiliflorum.
lanceolata, Tul. l. c. 121 = Weihea lanceolata.
leptoclada, Tul. l. c. 125 = Weihea leptoclada.
madagascariensis, DC. l. c. = Weihea madagascariensis.
microphylla, Tul. . . . 124 = Weihea microphylla.
myriocarpa, Tul. . . . 162 = Weihea myriocarpa.
ovata, Tul. . . . 121 = Weihea ovata.
phaeotricha, Tul. l. c. 122 = Weihea phaeotricha.
serrata, Benth. . . . 223 = macrophylla.

CASSYTHA.

455[2]. a. SPIRONEMA, Rafin. l. c. (1836).

CASTANEA.

456[1]. a. *cochinchinensis*, Steud. Nom. ed. II. i. 624, in syn. = sativa.
sinensis, Perr. in Mém. Soc. Linn. Par. iii. (1825) 108 = sativa?

CATALPA.

458[2]. l. cordifolia, *Jaume St. Hil*. . . . 5.—Am. bor.
speciosa, Warder, . . . 1 = cordifolia.

CATAPODIUM.

458[3]. a. *filiforme*, Nees, in Royle, Illustr. Bot. Himal. 417 = F. filiformis, *Steud*.

PAG.

CATASETUM.

459¹. l. purum, *Nees & Sinning, Amoen. Bot. Bonn.* ii. (1824) i. t. 1.

CATESBAEA.

459². l. macracantha, *Wright,*

CATHA.

459³. l. *campestris,* Presl, . . . 34 = Putterlickia pyracantha.

CATTLEYA.

461³. a. violacea, *Reichb. f. in Gard. Chron.* (1877) II. 424, in nota.—Guian.

CAUCALIS.

462². a. grandiflora, *Benth. & Hook. f. Gen.* i. 929.—Oriens.

l. *grandiflora,* Linn. . . . =
siifolia, Roehl. Deutschl. Fl. ed. II. ii. 214 =

CAULINIA, Willd.

462¹. l. *composita,* Buch.-Ham. . . . n. 5184 = Cerato-phyllum demersum.

CAYAPONIA.

463². a. ARKEZOSTIS, Rafin. New Fl. Am. iv. 100 (1836).

CEANOTHUS.

464¹. a. DIPLISCA, Rafin. Sylva Tellur. 31 (1838).

l. *azureus,* Kellogg, . . . 54 =
californicus, Kellogg, . . . 54 =

a. *divergens,* Poepp. ex Endl. Gen. 1099 = Sageretia divergens.

464². *riparius,* Poepp. ex Endl. (l. c.) = Sageretia riparia.

464³. l. thyrsiflora, *Eschscholtz,*

a. *venosus,* Royle, Illustr. Bot. Himal. 169, nomen (Quid ?).—Reg. Himal.

CECROPIA.

465¹. a. *peltata,* Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856), 148 = Humboldtiana.

CEDRELA.

465¹. l. *Cedro,* M. Roem. Syn. Hesper. 136, in syn. =

465². *Rosmarinus,* Lour. . . . = Baeckea cochinchinensis.

CEDRUS.

465². l. *Deodara,* Loud. Hort. Brit. 338; Arbor. Libani, *Loud. Hort. Brit.* 338.—

CELASTRUS.

465³. a. EUONYMOIDES, Isnard, ex Medic. Phil. Bot. i. 173 (1789).

l. *campestris,* Eckl. & Zeyh. . . . = Putterlickia pyracantha.
Colpoon, Hort. . . . = Putterlickia pyracantha.

466¹. *integer,* Thunb. . . . = Putterlickia pyracantha.

466². *linearis,* Linn. f. . . . (Gymnosporiae sp.).—Afr. austr.
obtusus, Thunb. . . . = Putterlickia pyracantha.

466³. *pyracanthus,* Linn. . . . 197 = Putterlickia pyracantha.
quadrangulatus, Schrad. . . . 716 = Elaeodendron quadrangulatum.

467¹. a. *venustus,* Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 44 (Quid ?).—Reg. Magell.

l. *verrucosus,* E. Mey. . . . 453 (Putterlickiae sp.).

CELOSIA.

467³. l. *gnaphalioides,* Linn. f. . . . 161 = Pfaffia gnaphalioides.
laxa, *Schum. & Thonn.*

CELSIA.

468¹. a. DITOXIA, Rafin. Préc. Découv. 39 (1814); et in Desv. Journ. Bot. iv. (1814) 270.

CELTIS.

468³. a. *alpina,* Royle, Illustr. Bot. Himal. 341 = australis ?

l. *biflora,* Ruiz, ex Klotzsch, l. c. 539.

469¹. l. dichotoma, *Ruiz, ex Klotzsch, in Linnaea,* xx. (1847) 539.

a. *Inglisii,* Royle, (l. c.) = australis ?

469². *Muku,* Siebold, in Verh. Batav. Gen. xii. (1830) 28 = Aphananthe aspera.

469³. l. pubescens, *Spreng. Syst.* i. 931.
spinosa, Ruiz, ex Klotzsch, in Linnaea, xx. (1847) 539 =

PAG.

CENCHRUS.

470². l. *macrostachyus,* Hochst. . . . 109 = Pennisetum Prieurii.

470³. l. CENOLOPHON, Blume, . . . i. 60 (1827).

CENTAUREA.

471². l. acromaura, *Hoffmgg. Nachtr. Preisverz.* (1842) 30.—*Cf. Linnaea,* xvi. (1842) *Litt.* 283.

471³. armena, *Boiss.*

472². cerinthifolia, *Sibth. & Sm.*

a. *chlorosperma,* Arv. Ess. Pl. Dauph. 39 = *serratula.*

473¹. l. cyanoides, *Wahlenb. in Bergg. Resor,* ii. *Bihang,* 65; *et in Isis,* xxi.
Debeauxii, Gren. & Godr.

474¹. horrida, *Badaro, . . . Fis.* Dec. II. vii. (1824) 367

474³. a. *microcephala,* Steud. Nom. ed. II. ii. 568, in syn., sphalm. = aspera.

475². l. napulifera, *Rochel, ex Frivald. in Magyar Tud. Társ. Evkön.* ii. (1835) 250

475³. Picris. *Pall.*
Pseudo-behen, Pseudo-collina; Pseudo-crupina, *Pseudo-philostizus, Pseudo-phrygia, Pseudo-scabiosa, Pseudo-uniflora.*

476². a. *serratula,* Arv. Ess. Pl. Dauph. 38 (Quid ?).—Gallia.

476³. *spinoso-ciliata,* Seenus, Reise Istrien, 65 = nigra.
stenocarpa, Arv. Ess. Pl. Dauph. 39 (Quid ?).—Gallia.

477³. insere CENTINODIUM, Montand. Guide Bot. 270 (1868) = **Polygonum,** Tourn.
axillare, Montand. l. c. = P. aviculare.

CENTRATHERUM.

478². a. frutescens, *Benth. ex C. B. Clarke, Comp. Ind.* 5.—Java.

CENTROPOGON.

479¹. a. barbatus, *Benth. & Hook. f. Gen.* ii. 547.—Peruv.; N. Granat.
erianthus, *Benth. & Hook. f.* (*l. c.*).—Ecuador.

479². Hartwegi, *Benth. & Hook. f.* (*l. c.*).—Ecuador.
popayanensis, *Benth. & Hook. f.* (*l. c.*).—N. Granat.

CENTROSEMA.

479³. a. triquetrum, *Spruce, ex Benth. & Hook. f. Gen.* i. 528.—Peruv.

CENTUNCULUS.

480¹. a. *indicus,* Royle, Illustr. Bot. Himal. 310 = tenellus.

CEPHAËLIS.

480². d. IPECACUANHA, Arruda (*nunc* **Psychotria**).

l. EVEA, Aubl. . . . i. 100.
TAPOGOMEA, Aubl. . . . i. 157.

a. asthmatica, *Baill. Hist. Pl.* vii. 376, in adnot.—Hab. ?
guianensis, *Baill.* (l. c.).—Guian.

CEPHALANTHERA.

481¹. a. LORCHOPHYLLUM, Ehrh. Beitr. iv. (1789) 148.
XIPHOPHYLLUM, Ehrh. l. c. (1789).

CEPHALARIA.

481². a. XETOLA, Rafin. Fl. Tellur. iv. 95 (1836).
graminifolia, Steud. Nom. ed. II. ii. 651, in syn., sphalm. = Scabiosa graminifolia.

481³. l. CEPHALIDIUM, A. Rich. . . . 290 =
a. *citrifolium,* A. Rich. l. c. = A. morindaefolius.

CEPHALOSCHOENUS.

482². a. *longirostris,* Nees, in Nov. Act. Nat. Cur. xix. Stippl. i. (1843) 101, in obs. = seq.

l. *longisetis,* Nees, in Linnaea, ix. (1834) 296.

CERASTIUM.

483². a. DOERRIENA, Borkh. in Rhein. Mag. i. (1793) 528.

d. DUFOUREA, Gren. (*nunc* **Arenaria**).

a. *acutifolium,* Dufour, Lettr. Mont. Maudit. 372 = alpinum.

PAG.

CERASTIUM:—

484¹. *dahuricum,* Schrank, Pl. Rar. Hist. Monac. t. 75 = davuricum ?

485¹. *microcarpum,* Arv. Ess. Pl. Dauph. 22 (Quid ?).—Gallia.

485². *Ramondi,* Fenzl, ex Benth. & Hook. f. Gen. i. 149 = Arenaria purpurascens.

CERASUS.

486². l. *Hixa,* C. Sm. . . . = P. lusitanica.

CERATANDRA.

487¹. a. *atrata,* Steud. Nom. ed. II. ii. 415, sphalm. = chloroleuca.

CERATOSTEMA.

488². a. *coronaria,* Hort. Linden, ex Hook. f. Bot. Mag. sub t. 5575 = Themistoclesia coronilla.

CERATOZAMIA.

488³. l. *boliviana,* Brongn. . . . 9 = Zamia Brongniarti.
muricata, Miq. . . . nomen = Zamia muricata.

CERBERA.

489¹. d. NEIOSPERMA, Rafin. (*nunc* **Ochrosia**).

l. *chinensis,* Spreng. l. c. (Rauwolfiae sp.).—China.
fruticosa, Roxb. . . . Ker-Gawl. . . . =

489³. insere CERDOSURUS, Ehrh. Beitr. iv. (1789) 148 = **Alopecurus,** Linn. (Gramin.).

CERESIA.

489³. a. *biflora,* Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 301 = Paspalum biflorum.

CEREUS.

491¹. a. Hermannianus, *Suring. in Versl. en Meddel. Kon. Akad.* III. ii. (1885) 194.—Ind. occ.

491². l. *lumbricoides,* Lem. . . . 60 = Rhipsalis sarmentosa.

491³. a. *Martini,* Dupuis, in Rev. Hortic. Sér. IV. ix. (1860) 657. f. 123 = Martinianus.

l. obtusus, *Haw.* . . . 70.—Afr. austr.

493¹. *variabilis,* Pfeiff. . . . 105 = obtusa.

CERINTHE.

493². a. *pyrenaica,* Arv. Ess. Pl. Dauph. 52 (Quid ?).—Gallia.

CESTRUM.

495³. a. *macrophyllum,* Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1858), 140 = Billbergianum.

496¹. l. thyrsoideum, *H. B. & K.*

CHAENOSTOMA.

497². l. SUTERA, Roth, . . . (1807).

a. *gilanicum,* Steud. Nom. ed. II. ii. 175, in syn., sphalm. = macrospermum.

CHAEROPHYLLUM.

498². l. Sphallerocarpus, *Kar. & Kir.*

CHAETACHME.

498². l. *nitida,* Planch. & Harv. . . . 16 = aristata.

CHAETANTHERA.

498¹. l. *dealbata,* Hook. & Arn. . . . 104 = Trichocline dealbata.

CHAETARIA.

499¹. l. *Forskolii,* Nees, . . . = A. barbicollis.

CHAETOCALYX.

499². l. acutifolia, *Benth.*
latisiliqua, *Hemsl. Biol. Centr. Am. Bot.* i. 268.

CHAETOSPORA, R. Br.

500³. l. *Burmanni,* Schrad. . . . 4 = S. Burmanni.

501¹. d. *pterosperma,* H. B. & K.

CHAETOSPORA, H. B. & K. (post praec. ponitur).

500². a. *pterosperma,* Nees, in Mart. Fl. Bras. ii. I. 128 = praec.

CHAETOTHYLAX.

501². a. eustachiana, *Benth. & Hook. f. Gen.* ii. 1110.—Ind. occ.

PAG.

CHAILLETIA.

501². l. Bojeri, *Tul.* . . . Madag.
501³. *gelonioides*, Hook. f. . . . 570 = sumatrana.
reflexa, Oliver, Fl. Trop. Afr. i. 343, sphalm. = deflexa.
sumatrana, *Miq.* . . . 328.—Ind. or. ; Malaya.

CHALCANTHUS.

501³. l. *renifolius*, . . . = H. renifolia.

502¹. l. CHALEPOA, Hook. f. in Hook. Ic.

502². l. CHAMAECLEMA, Boerh. ex Boehm. in Ludw. Def. Gen. Pl. 115 ; Moench,

CHAMAECYPARIS.

502². l. *breviramea*, Maxim. . . . = Cupressus obtusa.
Keteleri, Stand. . . . = Cupressus obtusa.
502³. *nana*, Hort. . . . = Cupressus thyoides.
pendula, Maxim. . . . = Cupressus pendula.
Pseudo-squarrosa, Parl. . . . = Cupressus thyoides.
pumila, Hort. . . . = Cupressus thyoides.
sphaeroidea, Spach. . . . = Cupressus thyoides.
squarrosa, Hort. . . . = Cupressus thyoides.
variegata, Hort. . . . = Cupressus thyoides.

CHAMAEDOREA.

503¹. l. *Ghiesbreghtii*, Hort. . . . 239, nomen = Gaussia Ghiesbreghtii.

CHAMAEMELUM, Vis.

504¹. l. *auriculatum*, Boiss. . . . 23 = M. auriculatum.

504². insere CHAMAEMORUS, Ehrh. Beitr. iv. (1789) 147 = **Rubus**, Tourn. (Rosac.).

CHAMAEPEUCE.

504³. a. *cretica*, Steud. Nom. ed. II. ii. 630, in syn., sphalm. = mutica.
l. *Diacantha*, DC. l. c. = Cn. Diacantha.

CHAMAERAPHIS.

504³. a. PSEUDORAPHIS, Griff. Notul. iii. 29 (1851).

CHAMAEROPS.

505¹. l. *Biroo*, Siebold, in Verh. Batav. Gen. xii. (1830) 11 ;
excelsa, Thunb. . . . 130 =

505². insere CHAMAESCHOENUS, Ehrh. Beitr. iv. (1789) 147 = **Scirpus**, Tourn. (Cyperac.).

CHAMAESPHACOS.

505². a. TAPEINANTHUS, Boiss. ex Benth. in DC. Prod. xii. 436 (1848).

CHAMITIS.

506¹. l. *complicata*, Soland. ex Forst. f. in Comm. Gotting. ix. (1789) 45 ;

506². l. CHAPTALIA, Royle, . . . Himal. 251.

CHAPTALIA.

506². d. piloselloides, *Baker*, . . . —Bras.

CHASALIA.

507¹. l. *amicorum*, A. Gray, . . . 43 = Ixora carnea.

CHAVICA.

508². l. *petiolata*, C. DC. . . . 389 = P. reticulatum.

510². l. CHEIRISANTHERA, Hort. ex Lindl. & Paxt. Flow. Gard. ii. (1851) 59 ;
atrosanguinea, Hort. ex Lindl. & Paxt. l. c. ;

CHELIDONIUM.

510³. l. *hybridum*, Linn. . . . = Roemeria violacea.
511¹. *violaceum*, Lam. . . . = Roemeria violacea.

CHENOPODIUM.

512¹. l. *Atriplicis*, Linn. f. . . . = album ?

CHILOCHLOA.

514³. l. *dentata*, Trin. . . . = Prionanthium rigidum.

CHIMARRHIS.

515¹. a. *cyrrhosa*, Steud. Nom. ed. II. ii. 88, in syn., sphalm. = cymosa.

CHIONANTHUS.

515³. l. *Ghaeri*, Gaertn. . . . 190 = Scirpodendron costatum.

PAG.

CHIONANTHUS.

515³. l. *angustifolia*, Rafin. ante *chinensem*, Maxim. ponenda.

CHIONOLAENA.

516¹. l. Arbuscula, *DC.*
larandulacea, Hemsl.
lavandulifolia, *Benth. & Hook. f.*

CHIRONIA.

517². a. purpurascens, *Benth. & Hook. f. Gen.* ii. 805.—Afr. austr.

CHLAMYDOSTYLUS.

518¹. l. *Medusa*, Baker, l. c. = N. Medusa.

CHLORAEA.

518³. l. *densa*, A Rich. . . . 454 = membranacea.

CHLORIS.

519². a. LANGSDORFFIA, Fisch. ex Regel, Ind. Sem. Hort. Petrop. (1863) 26.

520². l. *Roxburghiana*, Schult. . . . 339.

CHLOROPHORA.

520³. a. excelsa, *Benth. & Hook. f. Gen.* iii. 363.—Afr. trop.

520³. insere CHLOROPHYLLUM, Liais, Brés. 613 (1872), sphalm. = **Chrysophyllum**, Linn. (Sapotac.).

CHLOROPHYTUM.

521¹. l. Schidospermum, *Baker*,

CHONDRILLA.

522². l. PELTIDIUM, Zollikof. in Nat. Anz. Schw. Ges. iv. (1820) 13.

522³. l. CHONDROLAENA, Nees, . . . = **Prionanthium**, Desv. (Gramin.).
phalaroides, Nees, . . . P. rigidum.

CHONDRORHYNCHA.

522³. l. fimbriata, *Reichb. f. in Gard. Chron.* (1879) II., 648, in nota ;

523². insere CHORDORRHIZA, Ehrh. Beitr. iv. (1789) 148 = **Carex**, Dill. (Cyperac.).

CHORIZANTHE.

524¹. a. insignis, *Curran, in Bull. Calif. Acad.* i. IV. (1885) 275.—Calif.

CHRISTOPHORIANA.

525². l. *spicata*, Borkh. in Rhein. Mag. i. (1793) 458 ; Moench,

CHRYSANTHELLUM.

525³. l. HINTERHUBERA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 42.

CHRYSANTHEMUM.

526¹. d. OTOSPERMUM, Willk. (*nunc* **Matricaria**).
l. *artemisiaefolium*, Klatt, . . . (1878) I.
526³. *Herminii*, Hoffmgg. & Link, . . . ii. 331 ; t. 102.
larvatum, Griseb. . . . 266 = coronopifolium.
527². *setabense*, Dufour, . . . = paludosum.
sylvaticum, Hoffmgg. & Link, Fl. Port. ii. 329 =
trichophyllum, Boiss. . . . 10 = Matricaria tricho phylla.
Turreanum, Vis. Stirp. Dalm. Rev.
viscidum, Steud. Nom. ed. II. i. 755, in syn. = viscosum.

527². insere CHRYSASTRUM, Willd. ex Wedd. Chlor. And. i. 211, in nota (1855) = **Liabum**, Adans. (Compos.).
sagittatum, Willd. ex Wedd. l. c. (= *Munnozia sagittata*).—Reg. Andin.

CHRYSOCOMA.

528². a. GUNDLACHIA, A. Gray, in Proc. Am. Acad. xvi. (1881) 100.

529³. l. **CHRYSOPHTHALMUM**, Sch. Bip.

CHRYSOPHYLLUM.

530². a. *philippense*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 109 (Quid ?).— Ins. Philipp.

PAG.

CHRYSOPOGON.

530². a. DIPOGON, Willd. in Steud. Nom. ed. II. i. 518 (1840).

CHRYSURUS.

531³. l. **Lamarkia**, Moench,

531³. insere CHRYSIS, Renealm. ex DC. Prod. v. 585 (1836) = **Helianthus**, Linn. (Compos.).

CIMICIFUGA.

535². d. PITYROSPERMA, Sieb. & Zucc. . . . iii. IV.
a. dahurica, *Maxim. Prim. Fl. Amur.* 28.—Reg. Amur. ; China.
l. japonica, *Spreng.*

CINCHONA.

536². l. *macrophylla*, Karst. . . . 151 = Remija macrophylla.
536³. *prismatostylis*, Karst. . . . 7 = Remija primatostylis.

CINERARIA.

537². l. *aurita*, Andr. . . . = Senecio cruentus.
a. *cladobotrys*, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) 343, in obs. = Senecio cladobotrys.
537³. l. *cruenta*, Mass. . . . = Senecio cruentus.
a. elegantissima, *Hoffmgg. Nachtr. Preisverz.* (1842) 26.—*Cf. Linnaea*, xvi. (1842) *Litt.* 280 (an hybr. ?).—Hab. ?
l. *Fussii*, Nym. Syll. 2 =
538¹. *lanata*, Link, . . . = Senecio cruentus.
538². a. *longifolia*, Griseb. & Schenk, (ut supra) 343 = Senecio cladobotrys.
539¹. l. *Webberiana* ×, Paxt. . . . 125.

CINNAMOMUM.

539³. l. Maruba, *Siebold, in Verh. Batav. Gen.* xii. (1830) 24 ; *Meissn.* . . . —Japon.

CIPURA.

540². l. *Northiana*, Endl. . . . 6 = Marica Northiana.

CIRSIUM.

541². l. *acanthifolium*, Arv. Monog. Pilos. 52 ; et in Bull.
a. *angustatum*, Peterm. Fl. Bienitz, 56 (Quid ?).—Europ.
bifrons, Arv. Monog. Pilos. 52 (Quid ?).—Gallia.
bracteosum, Arv. Add. Monog. Pilos. 18 (Quid ?).—Gallia.
541³. *capitatum*, Arv. Monog. Pilos. Suppl. 36 = *variegatum*.
carduophilum, Arv. Ess. Pl. Dauph. 38, in obs. (Quid ?).—Gallia.
cenisium, Arv. Monog. Pilos. Suppl. 38 (Quid ?).—Gallia.
cirsiophilum, Arv. Ess. Pl. Dauph. 38, in obs. (Quid ?).—Gallia.
542¹. l. *dentatum*, DC. . . . 654 = Carthamus dentatus.
542³. a. *laceratum*, Arv. Monog. Pilos. Suppl. 37 (Quid ?).—Gallia.
543². *orophilum*, Arv. Ess. Pl. Dauph. 37 (Quid ?).—Gallia.
544². *variegatum*, Arv. Monog. Pilos. 51 (Quid ?).—Gallia.

CISSAMPELOS.

544³. l. *discolor*, DC. . . . = Stephania discolor.
glabra, Buch.-Ham. . . . = Stephania discolor, rotunda.
hernandifolia, Willd. . . . = Stephania discolor.
545¹. *umbellata*, E. Mey. . . . = Stephania discolor.

CISSUS.

546¹. l. *hederacea*, Royle, Illustr. Bot. Himal. 149 =
547¹. *quinquefolia*, Borkh. in Rhein. Mag. i. (1793) 595 ; Desf.
rosea, Royle, Illustr. Bot. Himal. 145, 149. t. 26. f. 1 = V. repanda.
547². a. *Tweediana*, Hieron. Pl. Diaph. 66 = V. Tweedieana.

CISTANCHE

547². a. violacea, *Hoffmgg. & Link, Fl. Pont.* i. 320.—Reg. Mediterr.

CISTUS.

548¹. l. *corsicus*, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 416.
549². *strigosus*, Fisch. ex Link, Enum. Hort. Berol. ii. 75.

PAG.

CITRIOBATUS.

550². l. pauciflorus, *A. Cunn.*

CITRULLUS.

550³. a. COLOCYNTHIS, [Tourn.] Linn. Syst. ed. I (1735).

CITRUS.

551¹. l. *Daidai*, Siebold, in Verh. Batav. Gen. xii. (1830) 59
Kunep, Siebold,

551². *Papeda*, Miq.
a. *Sabou*, Siebold, (l. c.) (Quid?).—Japon.
Toomikan, Siebold, (l. c.) (Quid?).—Japon.
Sarcodactylus, Siebold, (l. c.); Nooten, Fleurs Fr. Java, t. [29] = C. Medica.

CLADIUM.

551³. a. VAUTHIERA, A. Rich. Fl. N. Zel. 106. t. 20 (1830).

552². insere CLADOSPERMA, Griff. Notul. iii. 165 (1851) = **Pinanga**, Blume (Palm.).

CLARKIA.

553². l. *Eiseneana*, Kellogg, . . . 94 = pulchella.

CLAUSENA.

553³. l. COOKIA, Sonner. Voy. Ind. iii. 258. t. 130 (1782).

CLAVIJA.

554¹. a. *pungens*, Decne. in Ann. Sc. Nat. Sér. VI. iii. (1876) 145, in obs. = ornata.

554³. l. CLEACHNE, Roland. ex Rottb. in Act. Lit. Univ. Hafn. i. (1778) 285 =

CLEISTES.

555³. a. *bella*, Reichb. f. & Warm. Otia Bot. Hamb. 81 = P. bella.
caloptera, Reichb. f. & Warm. l. c. = P. caloptera.
l. *lutea*, Lindl. . . . 409 = P. grandiflora.
a. *Mantiqueirae*, Reichb. f. & Warm. Otia Bot. Hamb. 81 = P. Mantiqueirae.
paludosa, Benth. ex Griseb. Fl. Brit. W. Ind. 637 = P. tenuis.
l. *parviflora*, Lindl. . . . 410 = P. parviflora.

CLEMATIS.

556¹. l. SIEBOLDIA, Hoffmgg. Preisverz. (1842) Nachtr. 28.—Cf. Linnaea, xvi. (1842) Litt. 281.
a. *atroviolacea* ×, Carr. in Rev. Hortic. (1879) 350.

556². l. depauperata, *Colenso*, . . . (1868) ed. I. *Essays*, n. IV. 18; ed. II. (1871) 248.

556³. d. *glauca*, Claus,

557¹. l. *Jackmanni* ×, Hort.—Cf. Gard. Chron. (1864) 773, 823; et ex Lem. Illustr. Hortic. xi. (1864) t. 414;

558¹. l. *sibirica*, Mill. . . . 12 = C. Flammula.
d. *tubulosa*, Hook.

CLEOME.

558³. d. MOZAMBE, Rafin. (*nunc* **Cadaba**).
PODOGYNE, Hoffmgg. (*nunc* **Gynandropsis**).
l. *aphylla*, Thunb. . . . = Cadaba juncea.

559¹. a. dumosa, *Baker, in Journ. Bot.* xx. (1882) 18. —Madag.

559². micrantha, *Baker*, (*l. c.*).—Madag.

560¹. l. semitetrandra, *Sond.* . . . (semitetranda).
stenophylla, *Klotzsch*, . . . 1164 = Guian.; Bras.

CLERODENDRON.

561². l. *hemiderma*, F. Muell. . . . 61 = Glossocarya hemiderma.
Kaempferi, Siebold, in Verh. Batav. Gen. xii. (1830) 31 (err. typ. 51); Fisch.
Linnaei, F. Muell. . . . 151 = Glossocarya hemiderma.

562¹. a. violaceum, *Zeyh. Hort. Schwetzing.* 54; *ex Link, Enum. Hort. Berol.* ii. 127.—Hab.?

562². post CLETHROSPERUM pone
CLEVELANDIA, Greene, in Bull. Calif. Acad. i. n. IV. (1885) 182 (Scrophul.).
Bildingi, *Greene, l. c.* = Calif.

PAG.

CLIBADIUM.

563¹. l. *neriifolium*, DC. . . . 507 = Espeletia neriifolia.

CLIDEMIA.

563². l. *Epibaterium*, DC. . . . = Ossaea Epibaterium.

563³. *Xantholasia*, DC. = Oxymeria Xantholasia.
Xantholasia, Gardn.

CLINOGYNE.

565². a. ILYTHURIA, Rafin. Fl. Tellur. iv. 51 (1836).
Blumei, *Benth. & Hook. f. Gen.* iii. 650.—Java.

CLITORIA.

566³. a. *philippensis*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 110 (Quid?).—Ins. Philipp.

CLUSIA.

567². a. Galactodendron, *Desv. in Ann Sc. Nat.* Sér. II. xviii. (1842) 315. t. 8.—Venezuela.

567³. *longifolia*, Rich. . . . 113 = Tovomita clusiaefolia.

CLUYTIA.

568². l. *hirta*, Linn. f. . . . = Lachnostylis capensis.

568³. a. *vaccinioides*, Muell. Arg. in DC. Prod. xv. II. 436 = heterophylla.

CLYPEA.

568³. l. *hernandifolia*, Wight & Arn. . . . = S. discolor.
venosa, Blume, l. c. = S. rotunda.

CNEMIDOSTACHYS.

569². l. *Vahlii*, Spreng.

CNIDIUM.

572¹. a. *tortuosum*, Benth. & Hook. f. Gen. i. 902 = Foeniculum tortuosum.

COBAEA.

572¹. a. campanulata, *Hemsl. in The Garden*, xvii. (1880) 352.—N. Granat.
l. gracilis, *Hemsl. l. c.*—Costa Rica.

572². a. Trianae, *Hemsl.* (*l. c.*) 353.—N. Granat.

COCCOCYPSELUM.

573¹. l. *violaceum*, Desf. . . . 404 = campanuliflorum.

COCCOLOBA.

573². a. GUAIABARA, Plum. ex Boehm. in Ludw. Def. Gen. Pl. 402 (1760).

COCCULUS.

574¹. d. LEGNEPHORA, Miers, (*nunc* **Pericampylus**).

574². l. coriaceus, *Blume*,
a. diversifolius, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 10.—Japon.
l. *Forsteri*, DC. 517 = Stephania discolor.
a. *indicus*, Royle, Illustr. Bot. Himal. 61 = Anamirta paniculata?
l. *longifolius*, Decne. . . . 84 = Pycnarrhena longifolia.

574³. *planiflorus*, Wall. . . . = Pycnarrhena planiflorus.
planifolius, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 44, sphalm. = *planiflorus*.

COCHLEARIA.

575¹. l. IONOPSIDIUM, Reichb. Ic. Fl. Germ. vii. 26. t. 649 (1829) (an genus proprium?).
flava, *Roxb.* . . . nomen; *et ex Royle, Illustr. Bot. Himal.* 72.—Ind. or.

575². polysperma, *Costa*, . . . 86.—Hispan.

575³. a. *Wasabi*, Siebold, in Verh. Batav. Gen. xii. (1830) 54 = Eutrema Wasabi.

COCHLIODA.

575³. a. rosea, *Benth. & Hook. f. Gen.* iii. 560.—Ecuador.
sanguinea, *Benth. & Hook. f. l. c.*—Peruv. Ecuador.

COCOS.

576². a. *indica*, Royle, Illustr. Bot. Himal. 395 = nucifera.

CODIAEUM.

576³. l. Inophyllum, *Muell. Arg.*
a. *undulatum*, Cogn. & March. in Dallière, Pl. Ornament. i. (1873) t. 15 = Croton undulatus
Weismanni, *Cogn. & March.* (*l. c.*) t. 22.—Hab.?

PAG.

CODONOPHORA.

577². a. *Selloi*, Hanst. ex Benth. & Hook. f. Gen. ii. 1003 = praec.

CODONOPSIS.

577². a. *cashmeriana*, Royle, Illustr. Bot. Himal. 450 (Index), sphalm. = Campanula cashmiriana.

COELIDIUM.

578¹. a. spinosum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 25.—Afr. austr.

COELOGYNE.

578². d. *angustifolia*, Wight,

579¹. l. *Lowii*, Paxt.
rhombophora, Reichb. f. . . . 116 = Pholidota rhombophora.

COFFEA.

579³. a. *herbacea*, Aubl. ex Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 176, in syn. = Faramea herbacea.

COIX.

581¹. l. *crypsoides*, C. Muell. . . . 334 = Polytoca barbata. in syn. passim = C. Lacryma-Jobi.

COLAX.

581¹. l. *jugosus*, Lindl. . . . 51 = Lycaste jugosa.
Puydtii, Linden & André, . . . 369 = Lycaste Puydtii.

COLBERTIA.

581². l. *coromandeliana*, DC.

COLCHICUM.

581². a. EUDESMIS, Rafin. Fl. Tellur. iii. 80 (1836).

COLDENIA.

582¹. l. *angolensis*, Welw. . . . 591 = procumbens.

COLEA.

582¹. a. ODISCA, Rafin. Fl. Tellur. 80 (1838).

582². l. Telfairiae, *Boj.*

COLEANTHERA.

582². l. myrtoides, *Stschégl.*

COLEBROOKIA.

582³. a. SUSSODIA, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 104 (1825).

COLEUS.

583¹. l. *africanus*, Benth. . . . 54 = Solenostemon ocymoides.
a. *Marshallii* ×, T. Moore, in Gard. Chron. (1868) 376; et in Floral World (1878), 353.
Veitchii ×, Hort. ex Dombr. Flor. Mag. vi. (1867) t. 345.

COLLADONIA, DC.

583³. l. *anisoptera*, Boiss. . . . 946 = P. anisopetala?
microcarpa, Boiss. . . . 946.—Syria.

COLLETIA.

584¹. l. *Ephedra*, Bert. . . . = Trevoa Frutilla.
Ephedra, Vent. . . . 16 = Retanilla Ephedra.

584². *obcordata*, Vent. . . . 92 = Retanilla obcordata.

COLLINSIA.

584³. l. *divaricata*, Kellogg, . . . 36 = sparsiflora.

COLLOMIA.

584³. l. *coccinea*, Lehm. . . . t. 1622 = Gilia coccinea.

585¹. *grandiflora*, Dougl. . . . t. 1174 = Gilia grandiflora.

COLOBANTHUS.

585¹. l. Benthamianus, *Fenzl*, . . . 49.—Ins. Antarct.

585². *pulvinatus*, F. Muell. . . . = Benthamianus.
subulatus, Hook. f. . . . 13 = Benthamianus.

COLOCASIA.

585². a. *himalensis*, Royle, Illustr. Bot. Himal. 407 = antiquorum.
l. humilis, *Hassk. in Hoev.* . . . 160; *Cat. Hort. Bog. Alt.* 56.
indica, *Engl. in DC. Monog. Phan.* ii. 494.—Malaya.
neo-guineensis, André, . . . 68 = Schismatoglottis neo-guineensis.

PAG.

COLOCASIA:—

585[2]. l. *odorata*, Hook. Bot. Mag. t. 3935 = Alocasia macrorhiza.
prunipes, C. Koch. & Bouché, . . . = antiquorum.

585[3]. l. *ramosa*, Schott, . . . = Alocasia cucullata.
rapiformis, Kunth, . . . 40 = Alocasia rapiformis.
vivipara, Thw. . . . 336 = Remusatia vivipara.

COLPODIUM.

586[1]. l. *latifolium*, R. Br. . . . p. clxxxvi = Arctagrostis latifolia.

COLPOON.

586[1]. l. Rhoiacarpos, A. DC. in DC. Prod.

586[1]. insere COLUBRINA, Brun. ex Montand. Guide Bot. 268 (1868) = **Polygonum**, Tourn.
intorta, Montand. l. c. = P. Bistorta.
vivipara, Montand. l. c. = P. vivipara.

COLUBRINA.

586[1]. d. Diplisca, Rafin. (*nunc* **Ceanothus**).

COLUMNEA.

586[3]. a. *bicolor*, Rafin. Fl. Tellur. ii. 57 = hirsuta.
587[1]. l. *scabrosa*, Soland. . . . = Rhabdothammus Solandri.
587[2]. *trifoliata*, Link, . . . = Stemodia trifoliata.
Warszewicziana, *Klotzsch & Hanst.*

COLURIA.

587[2]. l. Laxmannia, Fisch. Hort. Gorenk. ed. II. (1812) 67.

COMATOGLOSSUM.

588[2]. l. *strictum*, Karst. & Triana, l. c. 437 = T. stricta.

COMBRETUM.

589[1]. l. *ferrugineum*, Hoffmgg. ex Mart. in Flora, xxii. (1839) i. Beibl. 64, in obs. =

COMMELINA.

590[3]. a. *angustifolia*, Kellogg, ex Curran, in Bull. Calif. Acad. i. iii. (1885) 150 = dianthifolia ?

COMPARETTIA.

594[1]. l. *saccata*, Poepp. & Endl. . . . 72 = Scelochilus saccatus.

CONANTHERA.

594[2]. a. *biflora*, Phil. f. Cat. Pl. Vasc. Chil. 282, in syn., sphalm. = bifolia.

CONCEVEIBA.

594[2]. l. *africana*, Muell. Arg. . . . 530 = Neoboutonia Mannii.
Martiana, Baill. . . . 221 = Alchornea Martiana.
terminalis, Muell. Arg. . . . 167 = Gavarretia terminalis.

CONOCARPUS.

596[3]. d. Rhizaeris, Rafin. (*nunc* **Laguncularia**).

CONOPODIUM.

597[3]. l. Sphallerocarpus, Bess. ex DC. Mém. Ombell. 60 (1829); et Prod.

CONRADIA.

599[1]. l. *exserta*, Mart. . . . = P. catalpiflora.

CONRINGIA.

599[1]. l. *orientalis*, Dum. . . . 123 = perfoliata.
perfoliata, *Link*, . . . 172.—Europ.; Oriens.

CONVALLARIA.

599[2]. d. Globeria, Rafin. (*nunc* **Liriope**).
l. *bifolia*, Linn. . . . = Maianthemum Convallaria.

CONVOLVULUS.

599[3]. d. Pentanthus, Rafin. (*nunc* **Jacquemontia**).
600[2]. l. *angulatus*, Salzm. . . . Genèv. viii. (1838) 59 =
600[3]. *bifidus*, Vell. . . . = Jacquemontia grandiflora.
601[2]. *cirrhosus*, R. Br. . . . 475 = arvensis.
congestus, R. Br. . . . 64 = glomeratus, *Choisy*.
601[3]. *cymosus*, Roem. & Schult. . . . = Ipomoea Senegambiae.
602[1]. elongatus, *Willd.* . . .—Ins. Canar.; Afr. bor.
602[2]. a. *fruticulosus*, Bolle, in Bonplandia, ix. (1861) 54 = Perraudieri (in Addend.).
l. *fulcratus*, Greene, . . . i. iv.

PAG.

CONVOLVULUS:—

602[3]. a. hirtus, *Mill. Gard. Dict.* ed. VIII. n. 10.—Jamaic.
603[1]. l. *Imperati*, Vahl, . . . = Ipomoea carnosa.
603[3]. macrostegius, *Greene*, . . . i. iv.
mauritanicus, Boiss. . . . t. 122 = elongatus.
604[2]. a. *ovatifolius*, Hook. & Arn. Bot. Beech. Voy. 90 = Jacquemontia sandwicensis, et Breweriae sp. (= *Bonamia Menziesii*).
604[3]. Perraudieri, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 58.—Ins. Canar.
606[2]. tenuis, *Tenore, Ind. Hort. Sem. Neap.* (1839) 11. n. 3, nomen.—Hab. ?
l. translucens, *Hance, in Journ.*

CONYZA.

607[1]. l. *ambigua*, DC. . . . = Erigeron linifolius.
607[2]. *aspera*, Bory,
607[3]. a. *calophylla*, Willd. ex Sch. Bip. in Pollichia, xx–xxi. (1863) 432 (= *Tephrothamnus calophyllus*).—Venezuela.
608[1]. l. *dichotoma*, Pers. . . . = Allagopappus dichotomus.
608[2]. a. *florida*, Steud. Nom. ed. II. i. 583, in syn., sphalm. = falcata ?
608[3]. l. *inuloides*, [Dryand.] . . . = Allagopappus dichotomus.
609[1]. *Mursa*, Steud. Nom. ed. II. i. 414, sphalm. = seq.
Musra, Buch.-Ham. ex Wall. Cat. n. 3095.
609[3]. *salicina*, Kellogg, . . . 36 = Baccharis viminea.

COOPERIA.

610[3]. l. Sceptranthes, R. Grah. . . . (1836).
chlorosolen, Herb. . . . = Drummondii.

COPERNICIA.

611[1]. a. *campestris*, Burmeist. Reise La Plata Staaten, ii. 48 = Trithrinax campestris.
l. hospita, *Mart. l. c.* 243.

COPISMA.

611[1]. l. *trichodes*, E. Mey. . . . = R. totta.

CORALLORHIZA.

612[2]. a. *occidentalis*, La Pilaye, in Mém. Soc. Linn. Par. iv. (1826) 502 = innata.

CORCHORUS.

613[1]. a. *prostratus*, Royle, Illustr. Bot. Himal. 103, nomen (Quid ?).—Ind. or.
sloanoides, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iv. (1840) 23 = Entelea arborescens.

CORDIA.

613[2]. a. *americana*, Steud. Nom. ed. II. ii. 309, in syn. = Patagonula americana.
613[3]. l. *geraschanthoides*, Rich. . . . = C. Gerascanthus.
Geraschanthus, Sw. . . . 478 = geraschanthoides.
614[1]. *Geraschanthus*, Jacq. . . . = C. Gerascanthus.
614[3]. *Perrottetii*, A. DC. . . . 482 = Premna tomentosa.
a. Perrottetii, *Wight, Ic.* t. 1381.—Ind. or.
615[1]. d. ulmifolia, *Juss.*

CORDYLINE.

615[3]. a. Euphyleia, Rafin. Fl. Tellur. iv. 16 (1836).
amabilis ×, Cogn. & March. in Dallière, Pl. Ornament. i. (1873) t. 49.
Baptisti ×, Cogn. & March. (l. c.) t. 37.
Hendersoni ×, Cogn. & March. (l. c.) t. 31.
616[1]. l. *spectabilis*, Kunth & Bouché, . . . (1847) 11 =
616[2]. a. *Weismanni* ×, Cogn. & March. in Dallière, Pl. Ornament. i. (1873) t. 14.

COREOPSIS.

616[3]. l. Heterodonta, Nutt. ex Benth. & Hook. f. Gen. ii. 385 (1873).
617[1]. *hirta*, Rafin. . . . 15 = Rudbeckia triflora ?

CORETHROGYNE.

617[3]. a. *cana*, Greene, in Bull. Calif. Acad. i. iv. (1885) 223 = Diplostephium canum.

CORMUS.

618[3]. a. *florentina*, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 157 = P. crataegifolia.
trilobata, Decne. l. c. = P. trilobata.

PAG.

CORNUS.

619[2]. d. florentina, *Decne.* (*quae* Cornus).
a. *nervosa*, Royle, Illustr. Bot. Himal. 234, nomen (Quid ?).—Ind. or.
d. trilobata, *Decne.* (*quae* Cornus).

619[3]. post **COROKIAM** pone
CORONA, Fisch. ex R. Grah. in Edinb. N. Phil. Journ. xxi. (1836) 157 = **Fritillaria**, Tourn. (Liliac.).
leucantha, Fisch. ex R. Grah. l. c. 158 = F. ruthenica.
verticillata, Fisch. ex R. Grah. l. c. 159 = F. verticillata.

CORONILLA.

620[2]. l. *Securigera*, Salisb. . . . 341 = praec.

CORONOPUS.

620[2]. l. *Raddii*, Savi, . . . 33 = S. Raddii.

CORREA.

620[2]. a. *Harrisii* ×, Paxt. Mag. Bot. vii. (1840) 79.
longiflora ×, Paxt. (l. c.) 195.

CORTIA.

621[1]. l. Lindleyii, *DC.* . . . 187 (Lindleii).

CORVISARTIA.

621[1]. l. *indica*, DC. in Royle, Illustr. Bot. Himal. 234; et

CORYANTHES.

621[1]. l. *elegantissima*, Mast.

CORYDALIS.

621[2]. a. Odoptera, Rafin. Cat. 15 (1824).
622[2]. *parviflora*, Royle, Illustr. Bot. Himal. 374 (Quid ?).—Ind. or.
Pseudocava, *Pseudo-pumila.*

623[2]. l. **CORYNANTHE**, Welw. Apont. 590, nomen [1859]; et in

CORYNELLA.

623[2]. a. *domingensis*, Steud. Nom. ed. II. ii. 461, in syn. = paucifolia.

CORYTHOLOMA.

624[2]. l. *gracile*, Brongn. . . . 466 = G. punctata.

COSMANTHUS.

624[2]. a. *Purshii*, Wood, Class-Book (1861), 566 = P. Purshii.

COSTUS.

625[1]. a. Pyxa, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 3.

COTINUS.

625[3]. a. *radicans*, Borkh. in Rhein. Mag. i. (1793) 592 = R. Toxicodendron.
Toxicodendron, Borkh. l. c. = R. Toxicodendron.

COTONEASTER.

625[3]. l. *emarginata* ×, Hoffmgg. Nachtr. Preisverz. (1842) 29;—Cf. Linnaea, xvi. (1842) Litt. 282;

COTULA.

626[2]. l. Featherstonii, *F. Muell. Veg. Chath. Isl.* 27, in syn.—
626[3]. Gymnogyne, *F. Muell.*
627[1]. tridentata, *Benth. & Hook. f.* . . . 429.

COTYLEDON.

627[2]. a. *calyculata*, Soland. ex Sims, Bot. Mag. sub t. 1409 = praec.
627[3]. l. *hybrida*, Hort. Par. ex DC. Pl. Grass. t. 65; et libanotica, *Labill.*

COUMA.

628[3]. a. oblonga, *Benth. ex Muell. Arg. in Linnaea*, xxx. (1859–60) 390.—Venezuela.

COUSSAPOA.

630[3]. l. dealbata, . . . 17 (non 16).

COVELLIA.

631[3]. a. *ulmifolia*, Gasp. Nov. Gen. Fici (1844), 10 = F. ulmifolia.

PAG.

CRAMBE.

632³. l. Tataria, *Sebéok, in Jacq.*
a. *tatarica*, Willd. Sp. Pl. iii. 418 = praec.

CRANIOLARIA.

633¹. l. *unibracteata*, Nees & Mart. . . . 67 = Martynia spathacea.

CRASSULA.

633⁸. a. conspicua, *Harv. Rev. Pl. Succ.* 204.—Afr. austr.
634¹. l. decumbens, *Harv. in Harv. & Sond. Fl. Cap.* ii. 353.—Afr. austr.

CRATAEGUS.

635¹. a. OXYACANTHA, Medic. Phil. Bot. i. 150 (1789).
635². l. brachyacantha, *Sargent & Engelm.*
a. *Carrierei* ×, Carr. in Rev. Hortic. (1883) 108, cum ic.
caucasica, C. Koch, in Verh. Ver. Bef. Gartenb. II. i. (1853) 286 = C. Oxyacantha, pectinata.
Cosansasi, Herb. Lugd. Bat. ex C. Koch, Wochenschr. v. (1862) 396 = sanguinea.
635⁵. a. *flabellata*, Bosc, ex M. Roem. Syn. Rosifl. 136 = coccinea.
germanica, Hort. ex C. Koch, in Verh. Ver. Bef. Gartenb. II. i. (1853) 225 = Pyrus germanica.
l. grandiflora, *C. Koch*, (*l. c.*) 226; *et in*
a. *Lambertiana*, C. Koch, (l. c.) 229 = mexicana.
Loddigesiana, C. Koch, (l. c.) 228 = mexicana.
636¹. *media*, Bechst. Forstbot. 150 = C. Oxyacantha.
microphylla, *C. Koch*, (*l. c.*) 288.—Reg. Cauc.
peruviana, *C. Koch*, (*l. c.*) 297.—Peru.
l. *pontica*, C. Koch, . . . II. i. (1853)
636². a. *rivularis*, Brew. & S. Wats. in Bot. Calif. i. 189 = Douglasii.
sicula, *C. Koch*, (*l. c.*) 288.—Sicil.
songarica, C. Koch, (l. c.) 287 = laciniata.
636³. *suecica*, Auct. ex Steud. Nom. ed. II. ii. 423, in syn. = Pyrus intermedia.
l. *uniflora*, Muenchh. . . . v. II. (1770) 566 =

CRATAEVA.

636⁸. a. TRICLANTHERA, Rafin. Sylva Tellur. 108 (1838).

CRAWFURDIA.

637². a. *speciosa*, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 442, partim = campanulacea.

CREMANIUM.

637³. l. *Sieberi*, Griseb.

CREPIS.

639². l. bulbosa, *Tausch*, . . . (1828).
639³. a. *curiensis* ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 110.—Europ.
l. djimilensis, *Boiss. Fl. Orient.* iii. 833.—Reg. Cauc.
640¹. a. *helvetica* ×, Bruegg. (l. c.).—Europ.
640². l. *hieracioides*, Ledeb. . . . 126 = lyrata.
hieracioides, *Waldst. & Kit. Pl. Rar. Hung.* i. 42. t. 70; *Willd.*
hispidissima, Bartl. in Bartl. & Wendl. f. Beitr. ii. 125
640³. a. *Muretiana* ×, Bruegg. (l. c.).—Europ.
l. *Novae-Zelandiae*, Hook. f. . . . 164 = Sonchus Novae-Zelandiae.
641². Stenoma, *Turcz.*
641³. a. *turicensis* ×, Bruegg. (l. c.).—Europ.

CRINUM.

642². l. *anomalum*, Herb. in Bot. Mag. sub t. 2121; et
642⁵. a. Binsii, *Hoffmgg. Preisverz.* (1841) 77.—*Cf. Linnaea*, xvi. (1842) *Litt.* 267.—Bras.
643¹. *himalense*, Royle, Illustr. Bot. Himal. 374 = amoenum?
643². *Roxburghii*, Dalz. & Gibs. Bomb. Fl. 275 = defixum.

644⁴. post **CROCKERIAM** insere
CROCOCYLUM, Steud. Nom. ed. II. i. 803, sphalm. = CROCOXYLON, Eckl. & Zeyh. = **Elaeodendron**, Jacq. (Tiliac.).

PAG.

CROCUS.

646¹. l. vitellinus, *Wahlenb. in Bergg. Resor*, ii. *Bihang*, 59; *et*

CROTALARIA.

647¹. a. *arida*, Royle, Illustr. Bot. Himal. 191 = C. Burhia?
647². *cannabinus*, Royle, (l. c.) 82 = juncea?
647³. l. cytisifolia, *Hoffmgg. Nachtr. Preisverz.* (1842) 25.—*Cf. Linnaea*, xvi. (1842) *Litt.* 279.—Peruv.

CROTON.

651². d. MEIALISA, Rafin. (*quae* **Adriana**).
ODOTALON, Rafin. (*nunc* **Argithamnia**).
l. PALANOSTIGMA, Mart.
651³. a. appendiculatus, *Hort. Veitch. ex Rev. Hortic.* (1877) 87. f. 13.—Polynes.
652¹. *Bergmani* ×, Chantrier, ex André, in Illust. Hortic. xxvii. (1880) 102. t. 389.
Camaza, Perr. in Mém. Soc. Linn. Par. iii. (1825) 112 = C. Tiglium?
652⁸. l. crassifolius, *Geisel.* . . . 19.—China.
654¹. a. imperialis, *Hort.—Cf. Gard. Chron.* (1875) II. 192; *et ex T. Moore, Flor. & Pomol.* (1876) 209.—N. Hebrid.
interruptus, Hort. (l. c.) (1871) f. 31 = Codiaeum variegatum.
Johannis, Veitch, ex Regel, Gartenfl. xxii. (1873) 251 = Codiaeum variegatum.
655¹. *multicolor*, Veitch, ex André, Illust. Hortic. xix. (1872) 120 = Codiaeum variegatum?
musaicus ×, André, in Rev. Hortic. (1882) 240.
655². l. Palanostigma, *Klotzsch*,
655³. *philippensis*, Lam.
656¹. a. *polystachyus*, Hook. & Arn. Bot. Beech. Voy. 270 = Cumingii.
656². l. *ricinocarpus*, Linn. . . . = Leidesia capensis.
656³. *Siraki*, Siebold, in Verh. Batav. Gen. xii. (1830) 30; Sieb. & Zucc.
657³. a. *Weismanni*, Cogn. & March. in Dallière, Pl. Ornament. i. (1873) t. 22 = Codiaeum Weismanni.

CRUCIANELLA.

658¹ a. MAPPA, Habl. ex Ledeb. Fl. Ross. ii. 403 (1844). RUBEOLA, [Tourn.] Ludw. Def. Pl. 2 (1737).
658². l. *stylosa*, Trin. . . . = Phuopsis stylosa.

658². l. CRUCIATA, [Tourn.] Ludw. Def. Pl. 2 (1737) =

CRUPINA.

659¹. l. *alpestris*, Arv. Add. Monog. Pilos. 19 = vulgaris.

CRYPTANGIUM.

660². a. LERISCA, Schlecht. in Bot. Zeit. iii. (1845) 476.

CRYPTOCARYA.

661¹. a. kawaiensis, *Benth. & Hook. f. Gen.* iii. 158.—Ins. Sandvic.

CTENIUM.

662³. l. *americanum*, Spreng.
sechellense, Baker, . . . = Enteropogon melicoides.

CUCUBALUS.

663². a. SCRIBAEA, Borkh. in Rhein. Mag. i. (1793) 591.
663³. l. *saxifragus*, Schreb. Ic. Pl. Desc. Dec. i. 9. t. 5; Linn.

CUCUMIS.

664¹. l. *ambiguus*, Fenzl,
664². *Bicirrha*, Forst.
chrysocarpus, Welw. . . . 63, sphalm. = seq.
cognatus, *Fenzl*,
a. *himalensis*, Royle, Illustr. Bot. Himal. 218, nomen (Quid?).—Reg. Himal.
664³. *Melopepo*, Steud. Nom. ed. II. ii. 303, in syn. = C. Melo.
665¹. *verrucosus*, Hort. ex Steud. Nom. ed. I. 244 = C. Melo.

CUCURBITELLA.

665³. a. PRASOPEPON, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 26.

PAG.

CUMINUM.

666³. a. *aethiopicum*, Royle, Illustr. Bot. Himal. 230 = Ammi majus.
regium, Royle, (l. c.) = Ammi majus.

CUNONIA.

667¹. d. OOSTERDYKIA, Burm. Rar. Pl. Afr.
l. *celebica*, Blume, . . . 868 = Spiraeopsis celebica.

CUPANIA.

667³. l. *cupanioides*, Cambess. . . . 29 = laevis.

CUPRESSUS.

671¹. l. *breviramis*, F. Muell. . . . [ed. II.] 64, in obs. = obtusa.
a. *Corneyana*, Knight, Syn. Conif. 19 = torulosa.
671². l. obtusa, *C. Koch*, . . . 168.—Japon.
pendens, F. Muell. . . . [ed. II.] 64, in obs. = obtusa.
pisifera, *C. Koch*, . . . 170.—Japon.
squarrosa, Laws. . . . = pisifera.
thyoides, *Linn.* . . . 1003.—Am. bor.

CURCULIGO.

671³. l. latifolia, [*Dryand.*]
a. *petiolata*, Royle, Illustr. Bot. Himal. 376 = orchioides?

CURCUMA.

672¹. l. KUA, Rheede, ex Medic.
a. *Kuchoor*, Royle, Illustr. Bot. Himal. 357, 359 = caesia.

CURTIA.

672². a. HACKELIA, Pohl, ex Griseb. in DC. Prod. ix. 56 (1845).

674³. l. CUSSAMBIUM, Rumph. ex Lam. Encyc. ii. 230 (1786); Buch.-Ham.

CUSSONIA.

675¹. a. *quercifolia*, Hort. Cels, ex Colla, Hort. Ripul. 43 = triptera.

676². l. CYANOTRIS, Rafin. . . . (1818) 356;

676³. l. CYATHIDIUM, Lindl. . . . (1835).

677¹. post **CYATHOGYNEM** pone
CYATHOMISCUS, Turcz. in Bull. Soc. Nat. Mosc. (1863) I. 562, in obs. = **Marianthus**, Hueg. (Pittosp.).

CYATHULA, Phil.

677². l. *angustifolia*, Moq. l. c. 328 = Sericocoma angustifolia.

CYCAS.

677³. l. circinalis, *Linn.*
d. *circinalis*, Thou.
a. *circinnalis*, Auct. = circinalis.
hypoleuca, Presl, . . . (al. ord.).

CYCLANTHUS.

678³. a. *bifolius*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 112, sphalm. = seq.

CYCLEA.

678³. d. PERAPHORA, Miers (*nunc* **Lophophyllum**).

CYCLOSTIGMA, Phil.

680². l. *tenue*, Phil. l. c. = L. tenuis.

CYCNOCHES.

680³. a. *Lindleyi*, Hort. ex Linden, Pescatorea (1860), sub t. 22 = Lueddemannia Pescatorei.

CYMBIDIUM.

682¹. d. OLGASIS, Rafin. (*nunc* **Oncidium**).
TRICHORHIZA, Lindl. (*nunc* **Limia**).
682². l. *grandiflora*, Sw. l. c. 76 = Pogonia grandiflora.
683¹. *stapelioides*, Link & Otto, . . . 111. t. 52.

683³. l. CYMELONEMA, Presl, . . . = **Urophyllum**, Wall. (Rubiac.).
memecyloides, Presl, l. c. = U. memecyloides.

CYMODOCEA.

683³. d. HALODULE, Endl. (genus proprium).
a. *australis*, Trimen, Cat. Pl. Ceyl. 99 = Halodule australis.
l. ciliata, *Ehrenb.* . . . 3.—Mar. Ind. et Pacif.
isoëtifolia, *Aschers. l. c.*—Mar. Ind. et Pacif.

PAG.

CYNANCHUM.

684[1]. a. SARCOCYPHULA, Harv. Thes. Cap. ii. 58. t. 191 (1863).
684[2]. l. *aphyllum*, Vell. . . . = *Amphistelma*
685[1]. a. *kunawurense*, Royle, Illustr. Bot. Himal. 274, nomen (Quid ?).—Ind or.
l. *molle*, Steud. . . . 905 = Anisotoma cordifolia ?
685[2]. l. *pendulum*, Poir. . . . 429 = Daemia extensa.

CYNOCTONUM.

686[2]. l. *molle*, E. Mey. . . . 216 = Anisotoma cordifolia ?
a. *tuberculatum*, Blume, ex Miq. Fl. Ind. Bat. ii. 478.—Java.

CYNOGLOSSUM.

686[3]. l. *Berteri*, Colla, . . . 153 = Selkirkia Berteroi.
687[1] *glomeratum*, [Nutt. in] Fras. Cat. (1813) n. 18, nomen ; et ex
687[2]. *javanicum*, Lehm. in Neue Schr. Naturf. Ges. Halle, iii. II. (1817) 21 = *macrophyllum*, Royle, Illustr. Bot. Himal. 305 = Paracaryum glochidiatum.
688[1]. *zeylanicum*, Thunb. ex Lehm. (ut supra) 20 =

CYNOSORCHIS.

688[3]. a. *arnottioides*, Reichb. f. in Bonplandia, iii. (1853) 213.—Ins. Borbon.
Commersoni, Reichb. f. (l. c.), in obs.—Hab. ?
parviflora, Reichb. f. (l. c.).—Ins. Borbon.

CYNOSURUS.

688[3]. l. *capillaceus*, Beauv. . . . = Leptochloa capillacea.
capillaris, Hort. . . . = praec.

CYPERORCHIS.

689[2]. a. *Wallichii*, Blume, Orch. Ind. Archip. 92 = Cymbidium cyperifolium ?

CYPERUS.

689[2]. d. OPETIOLA, Gaertn. (*nunc* **Mariscus**).
TORULINIUM, Desv. (*nunc* **Mariscus**).
a. UNGERIA, Nees, ex C. B. Clarke, in Journ. Linn. Soc. xxi. (1884) 34.
689[3]. l. *aequalis*, Vahl, . . . 320 = prolifer.
a. *Amane*, Siebold, in Verh. Batav. Gen. xii. (1830) 384 = Fimbristylis aestivalis.
691[3]. l. *consocius*, Steud. . . . 299, 324 = *ferax*.
693[2]. a. *frondosus*, Salisb. Prod. 30 = alternifolius.
l. *Ginge*, Welw. . . . 586 = flabelliformis.
694[1]. a. *horizontalis*, Salisb. Prod. 30 = Schoenus compressus.
695[2]. l. *macreilema*, Steud. . . . 298, 324 = Mariscus albescens.
696[1]. a. *niger*, Salisb. Prod. 30 = Schoenus nigricans.
696[2]. *ovatus*, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 168 = Mariscus uniflorus.
697[1]. *politus*, Welw. ex Ridl. in Trans. Linn. Soc. Ser. II. ii. (18) 125 = flavescens.
697[2]. l. prolifer, *Lam.* . . . 147.—Afr. austr. ; Ins. Mascar.
Pseudo-bromoides, *Pseudo-giganteus*, Pseudo-strigosus, *Pseudo-vegetus*,
697[3]. *puncticulatus*, Aitch. in Journ. Linn. Soc. xix. (1882) 189 = Renschii (non Reinschii).
698[1]. a. *riparius*, Salisb. Prod. 30 = pannonicus.
698[2]. *scirpinus*, Salzm. ex Steud. in Flora, xxxiii. (1850) 229, nomen ; et Syn. Pl. Cyp. 108 = Fimbristylis bahiensis.
698[3]. l. *soongoricus*, Kar. & Kir. . . . = difformis.
a. *sparsiflorus*, Baldw. in Trans. Am. Phil. Soc. Philad. ii. (1825) 168 = Mariscus glomeratus.
699[3]. l. *umbellatus*, Benth.
700[2]. l. Wolfii, *Wood, in Bull. Torrey Bot. Club*, vi. (1876) 72 ; *et in*

CYPHOMATTIA.

701[1]. l. *lanata*, Boiss. l. c. = R. lanata.

701[2]. a. CYPRIPEDILUM, Aschers. Fl. Brandenb. 700 (1864) = seq.

CYPRIPEDIUM.

701[2]. a. *Ainsworthii*, Reichb. f. in Gard. Chron. (1879) I. 748 = *Selenipedium Ainsworthii*.
albo-purpureum, Reichb. f. (l. c.) (1877) II. 38 = *Selenipedium albo-purpureum*.
Arthurianum ×, Reichb. f. (l. c.) (1874) II. 676.

PAG.

CYPRIPEDIUM :—

701[2]. a. *Ashburtoniae* ×, Reichb. f. (l. c.) (1871) 1647. f. 348.
calanthum ×, Reichb. f. (l. c.) (1880) II. 652.
calophyllum ×, Reichb. f. (l. c.) (1881) I. 169.
calurum ×, Reichb. f. (l. c.) 41 ; et T. Moore, Flor. & Pomol. (1884) 145, cum ic.
Chantini, Hort.—Cf. Gard. Chron. (1882) II. 750 = insigne.
chloroneurum ×, Reichb. f. (l. c.) (1880) II. 525.
conchiferum ×, Reichb. f. (l. c.) (1881) I. 330.
Crossianum ×, Reichb. f. (l. c.) (1873) 877.
Dauthieri ×, Reichb. f. (l. c.) (1882) I. 218, nomen ; Kittel, in Berl. Garten-Zeit. (1885) 272, 273 = *Harrisianum*.
Dominianum, Reichb. f. (l. c.) (1870) 1181 = *Selenipedium Dominianum*.
discolor ×, Reichb. f. (l. c.) (1882) I. 218.
elatum, Poit. ex Perr. in Mém. Soc. Linn. Par. iii. (1825) (Quid ?).—Guian.
euryandrum ×, Reichb. f. (l. c.) (1875) II. 772.
701[3]. *gemmiferum* ×, Reichb. f. (l. c.) (1881) I. 814.
Godefroyae, *Hort. ex Godefr.-Lebeuf, Orchidoph.* (1883) 830, cum ic.—Malaya.
l. *hirsutum*, Mill. . . . 3 = pubescens.
humile, *Salisb.* . . . (1791).
a. *Leeanum* ×, Reichb. f. in Gard. Chron. (1884) I. 50, 140.
leucorrhodum ×, Reichb. f. (l. c.) (1885) I. 270.
lucidum ×, Reichb. f. (l. c.) (1877) II. 521.
macropterum ×, Reichb. f. (l. c.) (1882) II. 552.
marmorophyllum ×, Reichb. f. (l. c.) (1876) II. 130.
Maulei, Hort.—Cf. Gard. Chron. (1882) II. 750 = insigne.
Meirax ×, Reichb. f. (l. c.) (1880) II. 524.
melanophthalmum ×, Reichb. f. (l. c.) 525.
Morganiae ×, Reichb. f. (l. c.) (1880) II. 134.
nitens ×, Reichb. f. (l. c.) (1878) I. 398.
l. *palmifolium*, Lindl. . . . 527 = Selenipedium palmifolium.
paulistanum, Rodrig. . . . 203 = Selenipedium vittatum.
a. *porphyreum* ×, Reichb. f. in Gard. Chron. (1878) I. 366 = Selenipedium porphyrium.
porphyrospilum ×, Reichb. f. (l. c.) (1879) II. 489.
pycnopterum ×, Reichb. f. (l. c.) (1876) I. 622.
radiosum ×, Reichb. f. (l. c.) (1885) II. 424 (*radissum*).
Reichenbachii, Bull, Cat. n. 48 (1869–70) 4 = Selenipedium longifolium.
702[1]. *Sedeni*, Reichb. f. in Gard. Chron. (1873) 1431 = *Selenipedium Sedeni*.
selligerum ×, [Reichb. f.] in Rev. Hortic. (1877) 88 ; Gard. Chron. (1880) I. 776.
stenophyllum, Reichb. f. in Gard. Chron. (1876) I. 461 = *Selenipedium stenophyllum*.
superciliare ×, Reichb. f. (l. c.) II. 795.
Swanianum ×, Reichb. f. (l. c.) (1876) II. 36.
tessellatum ×, Reichb. f. (l. c.) (1875) II. 614.
vernixium ×, Reichb. f. (l. c.) (1879) I. 398.
vexillarium ×, Reichb. f. (l. c.) (1870) 1373.
Williamsianum ×, Reichb. f. (l. c.) (1882) I. 218.

702[1]. post CYRBASIUM pone
CYRETHRUM, Boiss. Diagn. Ser. I. xi. 25 (1879), err. typ. = PYRETHRUM, Hall. = **Chrysanthemum**, Tourn. (Compos.).

CYRTANTHUS.

704[1]. l. sanguineus, *Walp.*
striatus, Herb. . . . t. 2534.

CYRTOPODIUM.

705[1]. l. plantagineum, *Benth.*

CYSTACANTHUS.

705[2]. l. turgidus, *Nichols.*

705[3]. l. CYSTOGYNE, Gasp. Nov. Gen. Fici (1844); et Ricerch.
leucosticta, Gasp. ll. cc.

PAG.

CYTINUS.

706[1]. l. HYPOCISTIS, [Tourn.] Adans.

CYTISUS.

706[2]. l. *cinereus*, Vis. Fl. Dalmat. iii. 270, in obs. ; Vukot.
707[1]. l. *Laburnum*, Linn. . . . = Laburnum anagyroides.
a. longissimus, *Hoffmgg. Preisverz.* (1841) 83. —*Cf. Linnaea*, xvi. (1841) *Litt.* 276.—Hab. ?
708[1]. l. *Weldeni*, Vis. . . . = Petteria ramentacea.
virgatus, Steud. Nom. ed. II. i. 478 = grandiflorus.

CZACKIA.

708[1]. a. *liliiflora*, Montand. Guide Bot. 316 = P. Liliastrum.

708[2]. post DACTIPHYLLUM pone
DACTYLADENIA, Welw. Apont. 572 [1859] = **Griffonia**, Hook. f. (1865) (Rosac.).
floribunda, Welw. l. c. = G. Bellayana.

708[3]. l. DACTYLANTHERA, Welw. . . . [1859] = **Symphonia**, Linn. f. ? (Guttif.) (*vel* Garcinia punctata, *Oliver*).

708[3]. a. DACTYLETHRIA, Ehrh. Beitr. iv. (1789) 148 = **Digitalis**, Tourn. (Scrophul.).

DACTYLIS.

708[3]. a. *disticha*, Ball, in Ann. & Mag. Nat. Hist. xi. (1843) 348 = Wangenheimia disticha.

DAEDALACANTHUS.

709[3]. a. UPUDALIA, Rafin. Fl. Tellur. iv. 66 (1836), nomen prius.

DAEMIA.

709[3]. l. *aethiopica*, Decne. . . . 544 = extensa.
guineensis, G. Don, . . . = extensa.
Kempeana, F. Muell. . . . 94 = Pentatropis Kempeana.
scandens, G. Don, . . . 94 = extensa.

DAEMONORHOPS.

710[1]. a. *plumosus*, Hort. Bull.—Cf. Gard. Chron. (1870) 1086. f. 206 (an Sagus sp. ?).—Hab. ?

DAHLIA.

710[2]. a. lutea, *Hort.*—*Cf. Gard. Chron.* (1879) II. 521. f. 83.—Hab. ?

DAIS.

710[2]. l. *disperma*, Forst. f. . . . 33 = Phaleria disperma.
octandra, Linn. . . . 2 = Phaleria octandra.

DALBERGIA.

711[3]. l. Thomsoni, *Benth.*

DALEA.

712[1]. l. aurea, [*Nutt. in*] *Fras. Cat.* (1813) n. 29, *nomen* ; *et ex ej.*
712[2]. *enneandra*, [Nutt. in] (l. c.) n. 30 ; et ex

DALRYMPELEA.

714[1]. a. *nepalensis*, Royle, Illustr. Bot. Himal. 166 = T. pomifera ?

DAMMARA, Rumph.

714[2]. a. *Brownii*, Hort. Angl. ex Lem. Illustr. Hortic. ii. (1855) Misc. 60 = A. australis ?

DANTHONIA.

715[3]. l. *flaccida*, T. Kirk, . . . 224 = Agrostis avenoides.
716[2]. l. *Thouarsii*, Nees, . . . 337 = tortuosa.
Thuarii, Desv. . . . 99 = praec.

DAPHNE.

716[3]. d. GASTRILIA, Rafin. (*nunc* **Thymelaea**).
a. *canescens*, Wikstr. in Vet. Akad. Handl. Stockh. (1818) 308 = Thymelaea canescens.
717[1]. l. *euphorbioides*, Pusch. ex Wikstr. (l. c.) 307
717[2]. *osyroides*, Vahl, ex Wikstr. (l. c.) 328
a. *pubescens*, Roehl. Deutschl. Fl. ed. II. ii. 247 = alpina ?
717[3]. *tinctoria*, Wikstr. (l. c.) 308 = Passerina tinctoria.

DAPHNIDIUM.

717[3]. l. *bibracteatum*, Nees, . . . 615 = Litsaea tenuiramis ?

PAG.

DASISTOMA.

719[1]. a. *flava*, Wood, Class-Book (1861), 529 = G. flava. *grandiflora*, Wood, Am. Bot. & Flor. 231 = G. grandiflora, *Benth.*

719[3]. l. **DATISCA**, Linn.
a. CANNABINA, Tourn. ex Linn. Amoen. Acad. i. 415 (1749).

DATURA.

720[1]. l. dubia, *Bianca, in Atti Acc. Gioen.* Ser. II. i. (1844) 330; *et*

DAUCUS.

720[2]. a. POMELIA, Durando, ex Pomel, Mat. Fl. Atl. 7 (1860).

DECAISNEA.

722[3]. a. SLACKIA, Griff. Itin. Notes, 187 (1848).

DECAISNEA, Lindl.

722[3]. l. *angulosa*, Lindl. l. c. = T. angulosa.

723[1]. post DECALOPHIUM pone

DECAMERIA, Welw. Apont. 579, in obs. [1859] = **Gardenia**, Ellis (Rubiac.).
Jovis-tonantis, Welw. l. c. = G. Jovis-tonantis.

DECANEURUM.

723[1]. l. *frutescens*, DC. l. c. 66.—Java.

DELPHINIUM.

725[2]. a. *Actaea*, Royle, Illustr. Bot. Himal. 452 (Index), sphalm., nomen delendum.

725[3]. distichum, *Geyer, ex Hook. Lond. Journ. Bot.* vi. (1847) 68.—Am. bor. occ.

726[1]. l. *flammeum*, Kellogg, . . . 22 = cardinale.

DEMAZERIA.

727[1]. a. tuberculosa. *Battand. & Trab. Fl. Algér.* (1884) 100.—Alger.

727[2]. l. DENDRIUM, Desv. . . . iii. (1814)

DENDROBIUM.

727[3]. l. alpestre, *Lindl. in Royle*, . . . 370. t. 88. f. 2.

728[3]. a. *denudatum*, Royle, Illustr. Bot. Himal. 371, sphalm. = denudans.

729[1]. *grandiflorum*, Sw. Adnot. Bot. 54 = Stanhopea Bucephalus.

DENDROMECON.

732[2]. l. *Harfordii*, Kellogg, . . . 102 = rigidum.

DENDROPHTHOË.

733[1]. l. *verticillata*, Scheff. . . . 27 = L. verticillatus, *Benth. & Hook. f.*

DENDROPHTHORA.

733[1]. a. *squamigera*, Benth. & Hook. f. Gen. iii. 214 = Phoradendron squamigerum.

DEROEMERA.

734[2]. a. *unifolia*, Reichb. f. in Bonplandia, iii. (1855) 213.—Abyss.

DERRIS.

734[2]. a. NOTHODERRIS, Blume, ex Miq. Fl. Ind. Bat. i. 1. 148 (1855).

734[3]. nobilis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 245.—Afr. trop.
oblonga, *Hance, in Journ. Bot.* xvii. (1879) 10.—China.

DESCHAMPSIA.

735[1]. l. *grandis*, Bess. . . . = caespitosa.
montana, Parl. . . . = caespitosa.

735[2]. a. *versicolor*, Arv. Ess. Pl. Dauph. 68 = caespitosa.
l. *Wahlenbergiana*, Scheele, l. c. = caespitosa.

DESMANTHUS.

735[2]. a. ACUAN, Medic. Theod. 62 (1786).
flavescens, Steud. Nom. ed. II. ii. 418, sphalm. (= *Desmochaete flavescens*) = Pupalia lappacea.

735[3]. l. DESMAZERIA, . . . = **Demazeria**,

DESMODIUM.

736[1]. d. HETEROLOMA, Desv. (*nunc* **Adesmia**).

PAG.

DESMODIUM.

736[3]. l. *diffusum*, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 100; et Prod. ii. 335 = laxiflorum.
diffusum, *DC. Prod.* ii. 336.—Ind. or.; Burma.

737[1]. a. heterocarpum, *Endl. in Ann. Wien. Mus.* i. (1836) 185.—Ins. Societ.
lagopodioides, *Endl.* (*l. c.*).—N. Caled.

737[2]. l. macrophyllum, *Desv.* . . . nomen; *et in Mém. Linn. Soc. Par.* iv. (1826) 306.—Ind. or.

737[3]. a. *peripterum*, Zipp. ex Miq. Fl. Ind. Bat. i. 1. 172 = Pseudarthria viscida.

DESMONCUS.

738[3]. a. *macrantha*, im Thurn, in Timehri, iii. (1884) 259, sphalm. = seq.

DEUTZIA.

739[3]. a. *californica*, Hort. ex Carr. in Rev. Hortic. (1866) 339 = Philadelphus Gordonianus.

DEYEUXIA.

740[3]. a. Royleana, *Trimen, Syst. Cat. Pl. Ceyl.* 108.—Ind. or.

DIANELLA.

741[3]. l. RHUACOPHILA, Blume, . . . (1827).
a. *philippensis*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 112 (Quid?).—Ins. Philipp.

DIANTHERA.

741[3]. a. JUNGIA, Boehm. in Ludw. Def. Gen. Pl. 92 (1760).

742[1]. *ensiformis*, Wood, Am. Bot. Flor. 234 = crassifolia.
humilis, Wood, (l. c.) 235 = ovata.

DIANTHUS.

744[1]. l. Faurei, *Arv. Monog. Pilos.* 49; *et in*

745[1]. *obcordatus*, Reut. . . . 281 = Tunica glumacea. *Oreades*, Ball,

745[2]. *ponticus*, Wahlenb. in Bergg. Resor, ii. Bihang, 29; et in Isis,

745[3]. pusillus, *F. W. Schmidt*, . . . (1790).
a. *saxatilis*, F. W. Schmidt, in Neue Abh. Boehm. Ges. i. (1790) 28 (Quid?).—Europ.

746[1]. *splendidissimus*, Hoffmgg. Nachtr. Preisverz. (1842) 27.—Cf. Linnaea, xvi. (1842) Litt. 280 = barbatus.

746[2]. l. DIAPEDIUM, Kon. in Kon. & Sims, Ann. Bot. ii. (1805) 189 = **Dicliptera**, Juss. (Acanth.).
a. *chinense*, Kon. l. c. = Dicl. chinensis.
cuspidatum, Kon. l. c. = Dicl. micranthes.
frondosum, Kon. l. c. = Dicl. frondosa.
martinicense, Kon. l. c. = Dicl. martinicensis.
pubescens, Kon. l. c. = Dicl. pubescens.

DIASTEMA.

747[1]. l. caracasanum, *Klotzsch & Hanst. l. c.* xxvii. (1854) 720; *et*

DICERATELLA.

748[1]. l. floccosa, *Boiss.*

748[2]. l. DICEROS, Blume, Bijdr. 751

DICHOSTYLIS.

750[1]. a. *Torresiana*, Gaudich. in Freyc. Voy. Bot. 413 = Fimbristylis globosa.

750[2]. l. **DICHROLEPIS**, Welw. . . . [1859] (an **Eriocaulon**? Linn.).

DICHROMENA.

751[1]. a. Reverchoni, *S. H. Wright, in Bull. Torrey Bot. Club*, ix. (1882) 86.—Texas.

753[2]. a. DICROCEPHALA, Royle, Illustr. Bot. Himal. 249 (1835), sphalm. = **Dichrocephala**, L'Hérit. (Compos.).

DICTAMNUS.

753[2]. a. *altaica*, Fisch. ex Royle, Illustr. Bot. Himal. 156 = D. Fraxinella.

DICTYOCARYUM.

753[3]. l. Wallisii, *Hort. Bull, ex Seem. in Gard. Chron.* (1870) 698;

PAG.

755[1]. l. **DIDYMELES**, Thou. . . .—Cf. Benth. & Hook. f. iii. 397.

DIDYMOCARPUS.

755[3]. a. *venosus*, Royle, Illustr. Bot. Himal. 294 (Quid?). —Reg. Himal.

DIDYMOSPERMA.

756[1]. a. distichum, *Hook. f. in Kew Gard. Rep.* 1882 (1884), 61.—Reg. Himal.

756[3]. lin. 1[ma] l. **DIEFFENBACHIA.**
nobilis, Hort. Bull, ex Cogn. & March. in Dallière, Pl. Ornament. i. (1873) t. 4;

DIERVILLA.

757[1]. l. rosea, *Planch. in Fl. des Serres*, ix. (1854) 247, in obs.; *Mast.*

DIGITALIS.

757[2]. a. DACTYLETHRIA, Ehrh. Beitr. iv. 148 (1789).

DIGITARIA.

758[3]. a. *sanguinalis*, Royle, Illustr. Bot. Himal. 418, sphalm. = P. sanguinale.

DIMACRIA.

760[1]. l. *fulgens*, Colv. ex Sweet, Hort. Brit. ed. I. 77;

DIMORPHOSTACHYS.

761[1]. l. *Drummondii*, Fourn. . . . l. c. = Paspalum lentiginosum.

761[1]. l. DINEBA, Beauv. . . . = **Dinebra**, Jacq. (Gramin.).

DIODIA.

762[1]. l. *simplex*. Sw. . . . 29 = Spermacoce simplex.

762[2]. post **DIODIAM**, pone
DIODIOIDES, Loefl. It. Hisp. 201 (1758) = **Spermacoce**, Dill. (Rubiac.).

DIOSCOREA.

763[3]. a. illustrata, *Bull, Cat.* (1873) 6; *Gard. Chron.* (1873) 1730, 1731.—Bras.

764[2]. *sagittata*, Royle, Illustr. Bot. Himal. 378, 379 = glabra.

766[3]. a. DIOSPERMUM, Hook. f. Fl. Brit. Ind. iii. 227, 228 (1881), err. typ. = **Oiospermum**, Less. (Compos.).

DIOSPYROS.

766[2]. a. *amara*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 112 (Quid?).—China.

767[2]. l. mabacea, *F. Muell.*

767[3]. *nigra*, Perr. (ut supra) 113; Blanco,

DIPHOLIS.

769[2]. a. SPONDOGONA, Rafin. Sylva Tellur. 35 (1838).

DIPLADENIA.

770[1]. a. HOMALADENIA, Miers, Apocyn. S. Am. 163 (1878).

DIPLOCHONIUM.

771[1]. l. = **Trianthema**, Sauv.
a. *uvarioides*, Edgew. in Trans. Linn. Soc. vi. (1862) 204, sphalm. = T. hydaspica.

773[1]. l. DIPLOPYRAMIS, Welw. Apont. 591 [1859] = **Oxygonum**, Burch. (Polygon.).
aethiopica, Welw. l. c. (= *Ceratogonon atriplicifolium*).—Afr. trop.

773[1]. insere DIPLORRHIZA, Ehrh. Beitr. iv. (1789) 147 = **Habenaria**, Willd. (Orchid.).

DIPLOTAXIS.

773[3]. l. Barrelieri, *DC.*

DIPSACUS.

775[1]. a. TRICHOCEPHALUM, Schur, Enum. Pl. Transs. 293 (1866).
fallax ×, Simk. in Termész. Füzetek, ii. (1878) 150.—Europ.
levis, Royle, Illustr. Bot. Himal. 245, nomen (Quid?).—Ind. or.

DISCARIA.

778[2]. l. *articulata*, Miers, . . . errat. = Retanilla articulata.

PAG.

DISEMMA.

779[3]. l. *caerulescens*, Seem. . . . 366.—Ins. Amicor.

DISSORHYNCHIUM.

781[1]. l. *muricatum*, Schau. l. c. = H. muricata, *Vidal*.

DISSOTIS.

781[1]. a. HEDUSA, Rafin. Sylva Tellur. 101 (1838). KADALIA, Rafin. (l. c.) (1838).

DISTEGANTHUS.

781[2]. a. *Morrenianus*, Linden, ex Linden & André, in Illustr. Hortic. (1874) 120 = Ronnbergia Morreniana.

784[1]. a. DODONAEA, Boehm. in Ludw. Def. Gen. Pl. 212 (1760) = **Ptelea**, Linn. (Rutac.).

DOLICHANDRA.

785[1]. l. *cynanchoides*, Cham. l. c. 658 = M. Dolichandra.
a. *Fenzliana*, Miq. in Linnaea, xviii. (1844) 251 = Spathodea Fenzliana.

DOLICHLASIUM.

785[1]. l. *glanduliferum*, Lag. . . . = T. glandulifera.

DOLICHOS.

785[3]. a. *decipiens*, Burch. ex Harv. & Sond. Fl. Cap. ii. 241 = Vigna triloba.

DOMBEYA.

787[3]. l. HILSENBERGIA, Boj. in Rapp. Soc. Hist. Nat. Maurice, xii. (1841) 44; et in
a. Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 496.—Madag.
l. cannabina, *Hilsenb. & Boj. in Rapp. Soc. Hist. Nat. Maurice*, xii. (1841) 45.—Madag.
a. crassipes, *Baill. l. c.* 495.—Madag.
l. elliptica, *Boj.* (*ut supra*) 46; *et in* erythroclada, *Boj. ll. cc.* 47; 191.
hirsuta, *Boj.* (*ut supra*) 46; *et in*
788[1]. modesta, *Hook.* . . . t. 4578.—Madag.
a. *opulifolia*, Boj. (l. c.) 49 = viburniflora.
l. platanifolia, *Boj.* (*l. c.*) 46; *et in* Pseudo-populus, *Baill.*
a. Richardi, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 500.—Madag.
l. spectabilis, *Boj.* (*ut supra*) 46; *et in*
a. Thouarsii, *Baill.* (*l. c.*).—Madag.
l. triumfettaefolia, *Boj.* (*l. c.*) 47; *et in*
a. viburniflora, *Boj.* (*l. c.*) 47; *et in*

788[1]. l. DOMBEYA, L'Hérit. . . . = **Tourretia**, Fouger. *lappacea*, L'Hérit. l. c. = T. volubilis.

DONATIA.

788[2]. a. ORITES, Banks & Soland. ex Hook. f. Fl. Antarct. 282 (1847).

DOODIA.

789[1]. l. *simplicifolia*, Roxb. l. c. 366 = U. prunellaefolia?

DOREMA.

789[2]. a. XANTHOGALUM, Avé-Lall. in Fisch. & Mey. Ind. Sem. Hort. Petrop. viii. 73 (1842).

DORIA.

789[3]. l. *nodulosa*, Sch.-Bip. in Flora, xxvii. (1844) 769, sphalm. = O. perfoliata.
ramosa, DC. . . . 472 = O. ramosa.
a. *undulata*, DC. Prod. vi. 483, sphalm. = *undulosa*.

DORSTENIA.

790[3]. l. Barnimiana, *Schweinf.*

791[1]. a. *rugosa*, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. i. (1839) 215 = Elatostema rugosum.

DORYCNIUM.

791[2]. a. ORTHOLOTUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 363.

DRACAENA.

795[2]. a. *amabilis* ×, Hort.—Cf. Gard. Chron. (1871) 841; Flor. & Pomol. (1874) 75.
aurantiaca ×, T. Moore, in Gard. Chron. (1876) i. 689.
Baptistii ×, Hort. ex Gard. Chron. (1874) i. 92; Flor. & Pomol. (1875) 52.

PAG.

DRACAENA:—

795[2]. a. *Chelsoni*, Veitch.—Cf. Gard. Chron. (1868) 349 = Cordyline terminalis.
cuprea ×, T. Moore, in Gard. Chron. (1876) i. 690.

795[3]. *Goldieana* ×, Hort. Bull, ex Flor. & Pomol. (1873) 187.—Afr. trop.
Hendersoni ×, Veitch.—Cf. Gard. Chron. (1874) i. 531; Flor. & Pomol. (1875) 53.
hybrida ×, Hort.—Cf. Gard. Chron. (1875) i. 551; Flor. & Pomol. (1876) 17.
jaspidea ×, Linden, Cat. n. 93 (1875) 37.

796[1]. *lentiginosa* ×, Hort.—Cf. Gard. Chron. (1867) 461; Illustr. Hortic. (1870) 179.
Lindeni ×, Hort. Linden, ex André, in Illustr. Hortic. xxvii. (1880) 85. t. 384.
Macleayi, Hort. ex Regel, Gartenfl. (1871) 149 = Cordyline terminalis.
Massangeana, Hort. ex E. Morr. in Belg. Hortic. x xi. (1881) 327 = fragrans.
metallica, Bull, Cat. (1872) 5; Flor. Mag. N. S. (1872) t. 24 = Cordyline terminalis.
Reginae ×, Hort.—Cf. Gard. Chron. (1868) 26; T. Moore, in Flor. & Pomol. (1872) 63.
Regis ×, André, in Illustr. Hortic. xxvi. (1879) 153. t. 360.
Salviati ×, Linden, Cat. n. 93 (1875) 37.

796[2]. a. *serrulata*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 113 = angustifolia?
Thomsoni ×, Hort.—Cf. Gard. Chron. (1880) ii. 656, 690; T. Moore, in Flor. Mag. N. S. (1881) t. 441.

DRACOCEPHALUM.

796[3]. a. *cuspidatum*, [Nutt. in] Fras. Cat. (1813) n. 28, nomen (Quid?).—Am. bor.

797[1]. *punctatum*, Steud. Nom. ed. II. ii. 153, in syn. sphalm. = D. Moldavica.

DRACOPHYLLUM.

797[3]. l. *cosmelioides*, Panch. . . . 156 = gracile.
d. *gracile*, Brongn. & Gris, etc.
a. *rosmariniforme*, R. Cunn. ex A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 49 = Urvilleanum.

DRACUNCULUS.

797[3]. a. *paludosus*, Montand. Guide Bot. 309 = Calla palustris.

DREPANOCARPUS.

798[2]. a. *lanatus*, Steud. Nom. ed. II. ii. 610, in syn., sphalm. = seq.

798[3] l. **DRIESSENIA**, . . . (1842?). bulbosa, *Triana*,

DRIMYSPERMUM.

799[2]. l. *Burmanni*, Decne. l. c. 40 = P. octandra.
Forsteri, Meissn. l. c. = P. disperma.

DROSERA.

800[1]. a. *muscipula*, Royle, Illustr. Bot. Himal. 75 = peltata.

800[2]. l. peltata, *Sm.* . . . 1546.—As. trop.; Austral.
d. *peltata*, Wight & Arn.

DRYANDRA.

801[1]. l. *cordata*, Thunb. l. c. = A. cordata.

801[2]. *oleifera*, Lam. . . . = Aleurites cordata.
Vernicia, Correa, . . . = A. cordata.
a. *verrucosa*, Royle, Illustr. Bot. Himal. 328, 452 = A. cordata.

DRYMARIA.

801[2]. l. diandra, *Blume, Bijdr.* 62.—Malaya?

DRYMOPHLOEUS.

802[1]. l. *puniceus*, Becc.

DRYPETES.

802[2]. l. *bengalensis*, Spreng. . . . = Fluggea microcarpa.
crocea, Poit. ex Spreng. Syst. iii. 902 = Xylosma nitidum.
crocea, *Poit. in Mém.* . . . t. 8; *Muell. Arg. in DC. Prod.* xv. ii. 456.—Ind. occ.
a. *laevigata*, Griseb. ex Eggers, Fl. St. Croix, (1879) 90 = Gymnanthes polyandra.

DUBOISIA.

802[3]. l. Hopwoodii, *F. Muell. Fragm.* x. 20.

PAG.

DUBRUEILIA.

802[3]. l. *microphylla*, Gaudich. . . . = P. muscosa.

DUFOUREA.

803[2]. l. *glabra*, H. B. & K.
a. *velutina*, Mart. & Gal. in Bull. Acad. Brux. xii. ii. (1845) 259.—Mexic.

DUHAMELIA.

803[3]. a. *pauciflora*, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 267 = H. ventricosa.

803[2]. l. DUNALIA, R. Br. in Salt, Abyss. App. 64 (1814) =

DYPSIS.

805[3]. l. *nodifera*, Mart. l. c. 312 = Phloga polystachia.

DYSODIA.

805[3]. a. *chrysanthemifolia*, Steud. Nom. ed. II. ii. 660, in syn., sphalm. = seq.
l. grandiflora, *DC.*

DYSOXYLUM.

806[2]. l. Blancoi, *Vidal*,

EBERMAIERA.

808[1]. l. nitida, *S. Moore, in Gard. Chron.*

808[2]. l. ECCOILOPUS, Steud. . . . = **Miscanthus**, Anderss.
andropogonoides, . . . = M. cotulifer.

ECHINACEA.

809[3]. a. Porteri, *Wood, Am. Bot. & Flor.* [445], in Add. —Am. bor.

ECHINOCACTUS.

812[2]. l. rinconensis, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 18.—
a. *rufispinus*, Bertol. in Ann. Stor. Nat. iii. (1830) 223 = Melocactus rufispinus.
l. Sickmanni, [*Lehm.*] *Del. Sem. Hort. Hamb.* (1837); *ex Linnaea*, xii. (1838) *Litt.* 83. —Am. austr.

812[3]. tubiflorus, *Hort. Angl.*

ECHINOCHLOA.

813[3]. a. *aristata*, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 299 = P. aristatum?
hispidula, Nees, in Royle, Illustr. Bot. Himal. 416, 420 = P. Crus-galli.

ECHINODORUS.

814[1]. a. BALDELLIA, Parl. Nuov. Gen. e Spec. 57 (1854).

ECHINOLOBIUM.

814[2]. a. *alpinum*, Desv. in Mém. Soc. Linn. Par. iv. (1826) 310 = H. alpinum.
argenteum, Desv. l. c. 312 = H. candidum.
biarticulatum, Desv. l. c. 310 = Desmodium biarticulatum.
carnosum, Desv. l. c. 313 = H. pallidum.
caucasicum, Desv. l. c. 312 = H. obscurum.
coronarium, Desv. l. c. 313 = H. coronarium.
flexuosum, Desv. l. c. = H. flexuosum.
fruticosum, Desv. l. c. 311 = H. fruticosum.
humile, Desv. l. c. 313 = H. humile.
incarnatum, Desv. l. c. 311 = H. incarnatum.
nitidum, Desv. l. c. 312 = H. nitidum.
obscurum, Desv. l. c. 310 = H. obscurum.
pallidum, Desv. l. c. 313 = H. pallidum.
sennoides, Desv. l. c. = Ormocarpum sennoides.
spinosissimum, Desv. l. c. = H. spinosissimum.
tauricum, Desv. l. c. 311 = H. tauricum.
trichocarpum, Desv. l. c. 312 = H. trichocarpum.
varium, Desv. l. c. = H. varium.

ECHINOSPERMUM.

816[2]. a. HACKELIA, Opiz, in Bercht. Oekon. Fl. Boehm. ii. 147 (1838).

816[3]. l. laevigatum, *Kar. & Kir.*

ECHITES.

817[2]. a. RHAPTOCARPUS, Miers, Apocyn. S. Am. 151 (1878).
l. bicolor, *Miq.*

817[3]. a. *Brownei*, Griseb. Fl. Brit. W. Ind. 414 = torulosa.

818[2]. guayaquilensis, *Miers, Apocyn. S. Am.* 233.—Ecuador.

819[1]. l. *nutans*, Anders. . . . *Haemadictyon*

PAG.

ECHIUM.

820[1]. d. Isorium, Rafin. (*nunc* **Lobostemon**).

EGLETES.

823[1]. a. *bellidifolia*, Steud. Nom. ed. II. ii. 420, in syn., sphalm. = praec.

EHRETIA.

823[1]. a. Lutrostylis, G. Don, Gen. Syst. iv. 391 (1837).

824[2]. l. EICHLERIA, Hartog, . . . = **Labourdonnaisia**, Boj. (Sapotac.).

ELAEAGNUS.

825[1]. l. *edulis*, Siebold, ex Carr. in Rev. Hortic. (1869) 300 = multiflora.
a. *Ghuwacen*, Royle, Illustr. Bot. Himal. 323 = umbellata ?
l. *Perrottetii*, Schlecht. . . . 613 = latifolia.
a. *philippensis*, Perr. in Mém. Soc. Linn. Par. iii. (1825) 114 = latifolia.
Rhamnoides, Royle, Illustr. Bot. Himal. 453 (Index) = Hippophaë Rhamnoides.

ELAEIS.

825[2]. a. *guianensis*, Steud. Nom. ed. II. ii. 249, in syn., sphalm. = guineensis.

ELAEOCARPUS.

825[3]. l. ceylanicus, *Mast. in Hook. f. Fl. Brit. Ind.* i. 406.—Zeylan.

826[3]. l. ELAEOCHYTRIS, Fenzl, . . . = **Ferula**, Tourn. (Umbell.).
cachroides, Fenzl, . . . = F. Schlechtendalii.
meifolia, Fenzl, . . . = F. meifolia.

ELAEODENDRON.

826[3]. a. *dichotomum*, Royle, Illustr. Bot. Himal. 167 = glaucum.

827[1]. l. *papillosa*, Hochst. . . . 305 = capense.

829[1]. l. ELECTRA, Panz. . . . 1813 (1814)
calycina, Panz. (l. c.), in obs. = S. brevifolius.

ELEGIA.

829[1]. d. Lamprocaulos, Mast. (genus retinendum).

ELEOCHARIS.

829[3]. a. Trichophyllum, Ehrh. Beitr. iv. (1789) 147.
austro-caledonica, Vieill. in Ann. Sc. Nat. Sér. IV. xvi. (1861) 38 = plantaginea.

830[3]. l. Robbinsii, *Oakes, in Hovey. Mag. Hortic.* vii.

ELEUSINE.

832[1]. a. brevifolia, *Wall. Cat.* n. 3815.—Ind. or.
l. *dura*, Lam. . . . = Sclerochloa dura.

832[2]. a. *Luco*, Welw. Apont. 591 = coracana.
stolonifera, *R. Br.* . . . 63, nomen.—Abyss.
textilis, Welw. in Murray, Journ. Trav. (1858) i. 31 = indica.
l. verticillata, *Roxb.* etc.

ELISMA.

833[1]. a. Nectalisma, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 156.

ELLIOTTIA.

833[2]. a. bracteata, *Benth. & Hook. f. Gen.* ii. 598.—Japon.
paniculata, *Benth. & Hook. f.* (*l. c.*)—Japon.

ELLIPEIA.

833[3]. l. *ferruginea*, Hook. f. & Thoms. . . . 52 = Uvaria ferruginea.
Laha, *Miq.*

ELODEA.

833[3]. a. *canadensis*, Fras. f. ex Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 168 = H. virginicum.
petiolata, Le Conte, ex Spach, (l. c.) 162 = H. paludosum.

ELODEA.

834[1]. a. Luchia, Steud. Nom. ed. II. ii. 75 (1841).
Serpicula, Pursh, Fl. Am. Sept. i. 33 (1817).
l. granatensis, *Humb. & Bonpl.*

834[2]. l. ELUTHERIA, P. Br. . . . = **Trichilia**, P. Br. (Meliac.).

PAG.

ELUTHERIA.

834[2]. a. *nobilis*, Triana & Planch. in Ann. Sc. Nat. Sér. V. xv. (1872) 376 = microphylla.

ELYMUS.

834[3]. l. dahuricus, *Turcz.* . . . ; *et* xxix. (1856) i. 61.—Dahuria.

835[1]. *glaucus*, Regel, . . . 585 = dasystachys.

ELYNANTHUS.

835[3]. a. *punctorius*, Benth. & Hook. f. Gen. iii. 1064.—Afr. austr.

EMBELIA.

836[2]. l. *abyssinica*, Baker, . . . 497 – Schimperi.

ENCELIA.

838[3]. a. *procumbens*, Steud. Nom. ed. II. ii. 249, in syn. = parvifolia.

ENCEPHALARTOS.

838[3]. l. *Acantha*, Mast. . . . 810 = cycadifolius.

839[1]. l. ENCHIDIUM, Jack, . . . 257 = **Trigonostemon**, Blume (1825).

839[2]. l. ENCHYSIA, Presl, . . . = **Laurentia**, Michx.

ENCLIANDRA.

839[2]. l. *parviflora*, Zucc. l. c. 337 = F. Eucliandra.

ENDIANDRA.

839[3]. a. Ita-uba, *Benth. in Benth. & Hook. f. Gen.* iii. 154.—Bras.
Navalium, *Benth. l. c.*—Bras.

840[1]. l. ENDOGONIA, Lindl. . . . = **Trigonotis**, Stev. (1851).

ENGELHARDTIA.

841[1]. a. *aurifolia*, Steud. Nom. ed. II. ii. 412, sphalm. = praec.
l. Oreomunea, *C. DC.*
a. *Oreomunnea*, C. DC. in DC. Prod. xvi. II. 142, err. typ. = praec.

842[2]. l. ENULA, Caesalp. ex Boehm. in Ludw. Def. Gen. Pl. 184 (1760) ; Neck.

EPACRIS.

842[3]. a. incarnata, *Hoffmgg. Preisverz.* ed. X. 81.—*Cf. Linnaea*, xvi. (1842) *Litt.* 274.—Hab. ?

EPICARPURUS.

844[3]. l. *microphyllus*, Raoul, . . . 14 = Paratrophis heterophylla.

EPILENDRUM.

845[1]. a. Laeliopsis, Lindl. in Paxt. Flow. Gard. iii. (1852–53) 155. t. 105.

845[2]. *anisatum*, Lindl. Fol. Orch. Epid. 55 = Juergensenii.

846[2]. l. *cochleatum*, Curt. . . . t. 152

847[3]. a. Juergensenii, *Reichb. f. in Gard. Chron.* (1880) II. 524, *in obs.*—Mexic.

851[2] l. EPIGYNIUM, Klotzsch, . . . (1851).
Leschenaultii, et *serratum* (etiam 1851).

EPIGYNUM.

851[3]. l. Legouixia, Heurck & Muell. Arg. in Heurck, Obs. Bot. 145 (1871).

EPILOBIUM.

852[2]. a. *decussatum*, Royle, Illustr. Bot. Himal. 211, nomen (Quid ?).—Reg. Himal.

852[3]. l. *flaccidum*, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 32 =

853[1]. *Herbertianum*, Royle, Illustr. Bot. Himal. 211, nomen (Quid ?).—Reg. Himal.
heterophyllum, Hegetsch. . . . Beitr. Crit. Schw. 345 =
a. *hybridum*, Arv. Ess. Pl. Dauph. 28 (Quid ?).—Gallia.
hyssopifolium, Arv. (l. c.) 29 (Quid ?).—Gallia.
intermedium, *Bory, ex Thore, Promen. Gasc.* 24 (Quid ?).—Gallia.
l. *intermedium*, Hegetsch. . . . Beitr. Crit. Schw. 345 =
a. *laeve*, Royle, Illustr. Bot. Himal. 211 = montanum ?

PAG.

EPILOBIUM :—

853[3]. l. *pauciflorum*, Schrank, in Denkschr. Bot. Ges. Regensb. i II. (1818) 15 =
a. *pendulum*, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 31 = nummularifolium.

854[1]. l. *ramiflorum*, Hegetsch. . . . Beitr. Crit. Schw. 345 =

854[2]. a. *scutellariaefolium*, Arv. Ess. Pl. Dauph. 29 (Quid ?).—Gallia.
l. *tetragonum*, Aitch.

854[3]. thymifolium, *R. Cunn. ex A. Cunn.* etc.

EPIPACTIS.

855[1] a. Limonias, Ehrh. Beitr. iv. (1789) 147.
l. atrorubens, *Roehl. Deutschl. Fl.* ed. II. ii. 481 ; *Schult.*
babianifolia, Roxb. . . . nomen = Pogonia babianifolia.

855[2]. a. *Gutta-sanguinis*, Arv. Ess. Pl. Dauph. 64 (Quid ?).— Gallia.

855[3]. *trinervia*, Roxb. . . . 455 = Pogonia trinervia.
l. *viridiflora*, Roehl. (l. c.) ; Reichb.

EPIPHYLLUM.

856[1]. a. *cruentum*, Hort.—Cf. Flor. Mag. N. S. (1880) t. 413 - Phyllocactus Ackermanni.

EPISCIA.

856[3]. l. *subacaulis*, Griseb. . . . 462 = Napeanthus subacaulis.

856[3]. l. EPISIPHIS, Rafin. . . . **Hypericum**, Tourn.

EPISTEPHIUM.

856[3]. a. *Cruegeri*, Reichb. f. ex Griseb. Fl. Brit. W. Ind. 637 = parviflorum.
l. *racemosum*, Rodrig. l. c. = sclerophyllum.

ERAGROSTIS.

857[2]. a. *Amherstiana*, Nees, in Royle, Illustr. Bot. Himal. 417, nomen (Quid ?).—Ind. or.
l. *aurea*, Steud. in Flora, xxix. (1846) 20 ; et

857[3]. *caudata*, Tourn. . . . 573 = capillaris.
cuspidata, Link, . . . = Vahlii.
elegantula, Nees, in Royle, Illustr. Bot. Himal. 417 =

858[1]. Frankii, *Steud.*
infirma, Steud.

858[2]. *mexicana*, Link, . . . 190 = capillaris.
mucronata, *Benth. & Hook. f. Gen.* iii. 1187.—Ind. or.
nigra, *Nees, in Royle, Illustr. Bot. Himal.* 417; *et ex*

858[3]. *rhachitricha*, Hochst. ex Steud. Syn. Pl. Gram. 267 =

859[1]. *rufescens*, Schult. Mant. ii. 319 = Vahlii.
solida, Nees, . . . = Vahlii.
spectabilis, Steud. . . . 564 = pectinacea ?
a. *thalassica*, Steud. Nom. ed. II. ii. 111, in. syn. = Poa thalassica.

859[2]. l. unioloides, *Nees*, . . . 264.—As. trop.

ERANTHEMUM.

859[2]. d. Upudalia, Rafin. (*nunc* **Daedalacanthus**).

ERECHTITES.

860[3]. l. *bathurstiana*, DC.

EREMURUS.

862[3]. a. soogdiana, *Benth. & Hook. f. Gen.* iii. 787.—Sibir.

ERIA.

864[2]. l. vestita, *Lindl.* . . . 76

ERIANTHUS.

865[1]. l. versicolor, *Nees, in Royle, Illustr. Bot. Himal.* 416, nomen ; *et ex*
a. *vulpinus*, Nees, (l. c.) 417, nomen (Quid ?).—Reg. Himal.

ERICA.

865[1]. a. Ericoides, Ludw. Def. Pl. 38 (1737); ed. II. 34 (1747).

866[1]. *Burnettii* ×, Hort. ex Planch. in Fl. des Serres, viii. (1852–53) 261. t. 845.
Cavendishiana ×, Paxt. Mag. Bot. xiii. (1847) 3.
Cavendishii ×, Paxt. l. c. t. [2] = praec.

867[1]. *Douglasiae* ×, T. Moore, in Moore & Ayres, Mag. Bot. iii. (1851) 9. t. [1] f. 1.

PAG.

ERICA:—
868[3]. *laxifolia*, Steud. Nom. ed. II. ii. 7, in syn., sphalm. = taxifolia.
869[1]. *Marnockiana* ×, T. Moore, (ut supra) f. 2.
minima, Pritz. Ic. Ind. i. 417 = hirtiflora.
869[2]. *Murrayana* ×, Turnb. ex Paxt. Mag. Bot. xi. (1844) 77.
nigrescens, Steud. Nom. ed. II. ii. 7, in syn. = nigrita.
870[1]. *pauciflora*, Steud. Nom. ed. II. ii. 656, in syn., sphalm. = sacciflora.
871[2]. *simulata* ×, T. Moore, (ut supra) f. 3.
Spenceriana ×, Planch. in Fl. des Serres, xxii. (1877) 111. t. 2323.

873[1]. a. ERICOIDES, Ludw. Def. Pl. 38 (1737); ed. II. 34 (1747) = **Erica**, Tourn.

ERIGERON.
873[3]. l. *asteroides*, Hort. Angl. . . . 324 = glaucus.
875[2]. *mixtus* ×, Arv. Add. Monog. Pilos. 19.—Cf.

ERINUS.
877[1]. a. pulchellus, *Jarosz, Pl. Nov. Cap.* 20.—Afr. austr.
l. *villosus*, Thunb. . . . = Zaluzianskya Benthamiana.

ERIOBOTRYA.
877[1]. a. *elliptica*, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. ii. 370 = petiolata.

ERIOCAULON.
878[2]. a. *hexangulare*, Royle, Illustr. Bot. Himal. 409, sphalm. = sexangulare.
879[1]. l. nepalense, *Presc. ex Bong.*
879[3]. *viride*, Koern. l. c. 637 = nepalense.
Wallichianum, Mart. . . . 249 = sexangulare.
880[1]. d. *Wallichianum*, Thw.

ERIOCHAETA.
880[1]. l. *densiflora, reversa, secundiflora*, Fig. & De Not. . . . (omnes) = P. penicellatum.

ERIOPHORUM.
882[2]. a. *Cannabina*, Royle, Illustr. Bot. Himal. 415 = comosum, Spodiopogon angustifolius.
depauperatum, Arv. Ess. Pl. Dauph. 66 (Quid?). —Gallia.

ERIOPHYLLUM.
883[1]. l. *multicaule*, A. Gray, . . . 24 = Actinolepis multicaulis.

883[2]. insere ERIOPUS, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 376 (1884) = **Trichocline**, Cass. (Compos.).
acaulis, Sch. Bip. ex Baker, l. c. = T. Eriopus.

ERIOSEMA.
883[3]. l. Guienzii, *Sond.* . . . (1850).
heterophyllum, *Benth. l. c.* xxii. (1849) 520. —Bras.

885[1]. insere ERIOSTOMA, Boiv. ex Baill. Adansonia, xii. (1878) 208 = **Hypobathrum**, Blume (Rubiac.).
albicaulis, Boiv. ex Baill. l. c. = H. albicaule.

ERUCAGO.
889[1]. l. *glandulosa*, Roehl. Deutschl. Fl. ed. II. ii. 355 =

ERUCARIA.
889[2]. l. *latifolia*, DC. l. c. 675 = Brassica adpressa.

ERVUM.
889[3]. l. *cyaneum*, Boiss. & Hohen. . . . 115 = Lens cyanea.
dalmaticum, Presl, . . . 31 = Lens nigricans.

ERYSIMUM.
894[1]. l. Strophades, *Boiss.*

ERYTHRAEA.
895[1]. l. *Pickeringii*, Oakes, . . . Mag. Hortic. vii. (1841) 179 =

ERYTHRINA.
895[2]. a. alba, *Cogn. & March. in Dallière, Pl. Ornament.* i. (1873) t. 56.—Hab.?
895[3]. horrida, *Eggers, Fl. St. Croix*, (1879) 45.— Ins. S. Croix; Virgin.
896[1]. *suberosa*, Welw. Synopse, 30. n. 73 = suberifera.

PAG.

ERYTHRONIUM.
896[3]. a. *hyacinthoides*, Royle, Illustr. Bot. Himal. 453 (Index), sphalm. = Scilla indica.

ERYTHRORCHIS.
897[1]. a. *Lindleyana*, Reichb. f. in Bonplandia, v. (1857) 37 = G. Lindleyana.

ERYTHROXYLUM.
897[3]. l. discolor *Boj. in Rapp. Soc. Hist. Nat. Maurice*, xii. (1841) 38; *et*
jossinoides, *Boj.* (*l. c.*) 38; *et*
lanceum, *Boj.* (*l. c.*) 39; *et*
898[1]. myrtoides, *Boj.* (*l. c.*) 39; *et*
platyclados, *Boj.* (*l. c.*) 40; *et*

ESCALLONIA.
898[2]. l. *bifida*, Link & Otto, Ic. Pl. Rar. 45. t. 23 =

ESCOBEDIA.
899[2]. a. MICALIA, Rafin. Fl. Tellur. ii. 104 (1836).

ETHULIA.
900[2]. l. *alata*, Sond. . . . 60 = Epaltes alata.

EUCLIDIUM.
904[1]. l. ORNITHORHYNCHIUM, Roehl. . . . (1812).

EUCNIDE.
904[1]. a. *cordata*, Kellogg, ex Curran, in Bull. Calif. Acad. i. IV. (1885) 137 = M. cordata.

EUGEISSONA.
904[3]. minor *et* insignis, *transponuntur.*

EUGENIA.
905[1]. l. SIPHONEUGENA, Berg,
905[2]. a. aromatica, *Sond. ex Berg, in Mart. Fl. Bras.* xiv. II. 379, in syn.—Bras.
905[3]. *Billardiana*, Steud. Nom. ed. II. ii. 177, sphalm. = Myrcia Billardiana.
906[2]. l. confertiflora, *A. Gray*,
907[2]. frondosa, *Wall. Cat.* sub n. 3590.
908[2]. a. *lasiopus*, Mart. ex DC. Prod. iii. 253 = Myrcia lasiopus.
910[1]. polyantha, *Miq. in Linnaea*, xviii. (1844) 741. —Guian.
911[1]. l. Schaueriana, *Miq. in Linnaea*, xxii. (1849) 174; *et*
911[2]. a. Tatna, *Wall. Cat.* sub n. 3555 (Quid?)— Ind. or.
911[3]. trunciflora, *Rich. ex Berg, in Linnaea*, xxx. (1859-60) 702, in syn.—Ind. occ.

912[2]. l. EULALIA, Trin. . . . = **Miscanthus**, Anderss.

EULOPHIA.
913[2]. a. *vera*, Royle, Illustr. Bot. Himal. 28, 366 = herbacea?

913[3]. a. EUONYMOIDES, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 318 = **Alectryon**, Gaertn. (Sapindac.).
excelsa, Soland. ex A. Cunn. l. c. = A. excelsum.

EUONYMUS.
914[1]. a. *Carrieri*, Vauv. ex Carr. in Rev. Hortic. (1885) 296. f. 50 = radicans.
914[2]. radicans, *Siebold, ex Miq. Ann. Bot. Mus. Lugd. Bat.* i. 86, in obs.; *et ex C. Koch, Dendrol.* i. 632.—Japon.
repens, Carr. in Rev. Hortic. (1885) 296. f. 51 = radicans.

EUPATORIUM.
919[3]. a. *salicinum*, Spreng. Syst. iii. 412 = Vernonia Sprengeliana.
920[3]. l. *Torreyanum*, Short & Peter, in Transylv. Journ. Med. n. xxxii. (1836) 575 =

EUPHORBIA.
922[2]. a. austriaca, *Kern. in Oestr. Bot. Zeitschr.* xxv. (1875) 397, in obs.—Europ.
924[3]. l. fragifera, *Jan, Cat. Pl. Phan.* (1818) 76; *et ex*
929[1]. Rapulum, *Kar. & Kir.*
930[2]. a. *tessellata*, Steud. Nom. ed. II. ii. 111 = E. Caput-Medusae.

EUPHORIA.
931[2]. l. *Gardneri*, Benth. & Hook. f.
a. *rimosa*, Royle, Illustr. Bot. Himal. 138 (Quid?). —Reg. Himal.

PAG.

EUPHRASIA.
931[3]. a. *gyroflexa*, Arv. Ess. Pl. Dauph. 56 (Quid?).— Gallia.
932[2]. *squamosa*, Presl, Bot. Bemerk. 91, sphalm. = squarrosa.
Veronicastrum, Arv. Ess. Pl. Dauph. 57 (Quid?).—Gallia.
viscida, Arv. (l. c.) (Quid?).—Gallia.

EURYA.
933[2]. a. *latifolia*, Hort. . . . = Cleyera japonica.

933[3]. l. **EURYALE**, Salisb. . . . (1805).
ferox, *Salisb.* . . . ii. (1805).

EURYBIA.
933[3]. a. *argyrophylla*, Sch. Bip. in Pollichia, xviii–xix. (1861) 173, sphalm. = praec.
934[1]. l. *corymbosa*, Cass. . . . = Aster corymbosus.

EURYGANIA.
934[3]. l. melliflora, *Benth. & Hook. f.* (*ponitur post* E. bifloram).
multiflora, Klotzsch, . . . = praec.

EURYLOBIUM.
934[3]. l. serrulatum, *Hochst. l. c.* 229.

935[3]. l. **EUSIDEROXYLON**, Teysm. & Binn.
a. *borneense*, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 179 = Zwageri.
l. Zwageri, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 292.—Borneo.

EUTERPE.
936[3]. a. stenophylla, *Trail, ex im Thurn, in Timehri*, iii. (1884) 229.—Guian.

937[1]. l. EUTHODON, Griff. Notul. iv. 84; Ic. Pl. Asiat. t.

EUTOCA.
937[1]. l. *Cumingii*, Benth. . . . 276 = P. Cumingii.

EUXOLUS.
937[3]. l. *crispus*, Lespin. . . .—Hab.?

EVELYNA.
938[3]. l. *Caravata*, Lindl. Fol. Orch. Evel. 9; ex Hook. Bot. Mag. t. 5141; et ex
longibracteata, Lindl. ex Griseb. Fl. Brit. W. Ind. 623.—Am. trop.

EVODIA.
939[1]. l. fraxinifolia, *Hook. f. ex Benth. Fl. Hongk.* 59, in nota; *et*
939[2]. *Marambong*, Miq.
rutaecarpa, *Hook. f. ex Benth. Fl. Hongk.* 59, in nota; *et*

EVOLVULUS.
939[3]. l. discolor, *Benth. Pl. Hartw.* 6.—Mexic.
940[1]. *Glechoma*, Welw. Apont. 589 = Ipomoea reniformis.

EXCOECARIA.
941[3]. l. *brachyandra*, Muell. Arg. usque ad *Chamelaea*, Baill., anteponendae cochinchinensi, *Lour.*

EXOGONIUM.
942[3]. l. *Purga*, Benth. Pl. Hartw. 46; Lindl.

FACELIS.
944[2]. l. *plumosa*, Sch. Bip. . . . nomen = Lucilia plumosa.

FADOGIA.
944[2]. l. Zeyheri, *Hiern, l. c.* 153 (cum cit. falsa).

FAGARA.
944[3]. l. *Pterota*, Linn.

FAUREA.
948[2]. a. TRICHOSTACHYS, Welw. Synopse, 19 (1862).

FENDLERA.
949[2]. a. *cymosa*, Greene, in Bull. Torrey Bot. Club, viii. (1881) 26 = utahensis.
utahensis, *Greene, l. c.* 25.—Utah; Arizona.

FERDINANDA.
949[3]. l. *angusta*, Hemsl. Biol. Centr. Am. Bot. ii. 159, sphalm. = seq.
augusta, Lag. Gen. & Sp. Nov. 31. t. 2 = Z. augusta.

PAG.

FERULA.

950². a. ELAEOCHYTRIS, Fenzl, in Russegg. Reise, i. 957. t. 19 (1843) (Illustr. Pl. Syr. 71).
LEPTOTAENIA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 629 (1840).
POLYCYRTUS, Schlecht. in Linnaea, xvii. (1843) 126.

950³. d. XANTHOGALUM, Avé-Lall. (*nunc* **Dorema**).

951¹. a. *Hooshee*, Lindl. Fl. Med. 46 = F. Assa-foetida ?

951². l. *Opopanax*, Spreng. . . . = Opopanax hispidum.
a. *setifolia*, C. Koch, in Linnaea, xix. (1847) 39 = pauciflora.

951³. l. *sylvatica*, Szov. ex C. Koch, (l. c.) = pauciflora.

FESTUCA.

952¹. a. LEUCOPOA, Griseb. in Ledeb. Fl. Ross. iv. 383 (1853).
TRAGUS, Panz. in Denkschr. Akad. Muench. 1813 (1814) 296.

952². *Amherstiana*, Royle, Illustr. Bot. Himal. 417, nomen = Poa persica.

953¹. *ciliata*, Heyne, ex Wall. Cat. n. 3826 = virgata.
curvata, All. ex Beauv. Agrost. 99, nomen.—Europ.

953³. Gerardi, *All. ex Beauv.* (*l. c.*), nomen.—Europ.

954¹. gracillima, *Thurb. in Bot. Calif.* ii. 318 (non Hook. f.).—Calif.
l. Hemipoa, *Delile*,

954³. *mysorensis*, Heyne, . . . = Enteropogon melicoides.

955¹. a. *pauciflora*, Thurb. in Bot. Calif. ii. 318 = occidentalis, etc.

955³. l. *reptatrix*, Linn. Sp. Pl. ed. II. 108 = *Rohlfsiana*, Coss. . . . 83 = Sclerochloa philistaea.
scabrella, *Thurb. in Bot. Calif.* ii. 318 (non Torr.).—Calif.

956¹. a. *sterilis*, Jess. Deutschl. Graes. 170 = Bromus sterilis.

956². *tectorum*, Jess. (l. c.) = Bromus tectorum.

956³. l. violacea, [*Clairv.*]

956³. insere (ante *Fialarem*, Rafin.) FEVILLAEA, Linn. Syst. ed. I (1735) = **Inga**, Scop. (Legumin.).

FICINIA.

957¹. a. HELLMUTHIA, Steud. in Flora, xxxiii. (1850) 229, nomen ; et Syn. Pl. Cyp. 90 (1855).

FICUS.

957³. l. CYSTOGYNE, Gasp. l. c. 9 (1844) ; et in

959¹. a. Chauvieri, *Hort.—Cf. Nichols. Dict. Gard.* ii. 12.—Hab. ?
l. chiriquiana, Miq.

960¹. a. *Forsteriana*, Endl. in Ann. Wien. Mus. i. (1836) 166 = prolixa.

960². habrophylla, *G. Benn. Gather. Natural.* 341.—Ins. Tanna.
l. *haematocarpa*, Decne.

961¹. a. *laurifolia*, Royle, Illustr. Bot. Himal. 336, nomen (Quid ?).—Reg. Himal.

963². l. *Reinwardti*, Link & Otto, . . . 61. t. 31 =

964¹. a. *tannensis*, G. Benn. Gather. Natural. 341, nomen (Quid ?).—Ins. Tanna.

FIMBRISTYLIS.

966². a. *bisumbellata*, Bubani, Dodecanth. 30 = dichotoma.

966³. *bulbosa*, Steud. Nom. ed. II. i. 828, in syn., sphalm. = Ficinia bulbosa.

967². l. *marquesana*, Steud. (post *marianam* ponitur).

968¹. *rigida*, Kunth,
Scherardi, Bubani, Dodecanth. 29 =
separanda, Steud. . . . nomen = diphylla.

968². a. *Steudelii*, Steud. Nom. ed. II. i. 827 ; ii. 539, sphalm. = Ficinia Ecklonea.
l. tenella, *Schult.*

FISCHERIA.

969¹. a. hispida, *Benth. & Hook. f. Gen.* ii. 766.—Am. austr.
setosa, *Benth. & Hook. f.* (*l. c.*).—N. Granat.

969¹. a. FISTULARIA, Linn. Syst. ed. I (1735) = **Rhinanthus**, Linn. (Scrophul.).

PAG.

FLACOURTIA.

969². a. SATANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

FLAVERIA.

970¹. l. *trinervata*, Steud. Nom. ed. II. ii. 186, in syn., sphalm. ; Baill.

FLINDERSIA.

971¹. l. *maculata*, F. Muell. in Journ. Pharm. Vict. ii. (1858) 44, sphalm. = seq.
maculosa, *F. Muell. ex Benth. Fl. Austral.* i. 389.—Austral.

FLUEGGEA, Rich.

971³. a. *dracaenoides*, Baker, in Journ. Bot. xii. (1874) 174.—Reg. Himal.

FLUGGEA.

971³. a. *retusa*, Royle, Illustr. Bot. Himal. 327 = microcarpa ?

FOENICULUM.

972¹. a. *Webbii*, Benth. & Hook. f. Gen. i. 902 = tortuosum.

FONTANESIA.

972². l. *chinensis*, Hance, . . . 136 = phyllyraeoides.
phyllyraeoides, *Labill.* . . . Syria ; China.

FOUQUIERIA.

973³. a. IDRIA, Kellogg, in Proc. Calif. Acad. ii. (1863) 34.
columnaris, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 133 = spinosa.

975¹. insere FRANCA, Mich. ex Ludw. Def. Gen. Pl. 169 (1747) = seq.

FRANKENIA.

975². l. FRANCA, Mich. ex Ludw. Def. Gen. Pl. 169 (1747).

FRANQUEVILLEA.

976¹. a. *major*, Zoll. ex Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 84 = Curculigo orchioides.

FREGIRARDIA.

977³. l. *luteiflora*, Dun. . . . (1849) = B. diversifolius.
Spina-alba, Dun. l. c. 507 = Bassovia Spina-alba.

FREMYA.

977³. l. *ciliata*, Brongn. & Gris, . . . 374 = Metrosideros ciliata.

978¹. *pubescens*, Brongn. & Gris, . . . = Metrosideros pubescens.
rubra, Brongn. & Gris, . . . = Metrosideros rubra.

FRITILLARIA.

979². l. RHINOPETALUM, Fisch. . . [viii.] (1829).
aurea, Schott, . . . = lutea ?
a. *Corona-imperialis*, Gaertn. Fruct. i. 64. t. 7 = F. Imperialis.

FUCHSIA.

981¹. l. Kirkii, *Hook. f.* . . . i. ed. II. (1871) 92.

981³. l. FULCALDEA post **FUIRENA** ad p. 982¹ ponitur.

FUIRENA.

981³. a. PENTASTICHA, Turcz. in Bull. Soc. Nat. Mosc. (1862) II. 330.
annua, Royle, Illustr. Bot. Himal. 411 = pubescens.
cylindrica, Steud. Nom. ed. II. ii. 539, sphalm. = Fimbrystylis spadicea.

GAGEA.

985². a. *Haeckelii* ×, Dufft & Schulze, ex Hausskn. in Mitth. Geogr. Soc. Jena, ii. (1885) 224.—Europ.

GAHNIA.

986¹. l. pauciflora, *T. Kirk*, . . . i. ed. II. (1871) 94.

GAILLARDIA.

986². a. *grandiflora* ×, Hort. ex Fl. des Serres, xiii. (1857) 1. t. 1183.
perennis, *Loisel. Herb. Gén. Amat.* v. t. 328.—Am. bor.

PAG.

GALATELLA.

988¹. a. *collina*, Jord. & Fourr. ex Nym. Consp. 386 = A. acris.
fugax, Nym. (l. c.) = A. aragonensis.
pinetorum, Jord. & Fourr. ex Nym. (l. c.) = A. acris.
sedifolia, Nym. (l. c.) = A. acris.
tatarica, Nym. (l. c.) = A. dracunculoides.

GALEANDRA.

988³. l. nivalis [*T. Moore ? in*]

GALEOPSIS.

990¹. l. *glaucescens*, Reut. ex Ardoino, . . . 299 = G. Ladanum.

990². *nepetaefolia*, Timb. . . . 19 = G. Ladanum.

990². l. GALEOTTIA, Rupr. . . . (1842).

GALINIERA.

990³. l. coffeoides, *Delile*,

GALINSOGA.

990³. a. *Acmella*, Steud. Nom. ed. II. ii. 766, in syn. = parviflora.

GALIPEA.

991¹. a. SYSTEMON, Regel, Ind. Sem. Hort. Petrop. (1856) 38 ; Gartenfl. vi. (1857) 81.

GALIUM.

991². l. CRUCIATA, [Tourn.] Ludw. Def. Pl. 2 (1737).

991³. aphyllum, [*Mazziari*]

992¹. Brandegei, *A. Gray, in Bull. U. St. Geol. & Geogr. Surv.* ii. (1876) 237 ; *et in*
a. *cenisium*, Arv. Ess. Pl. Dauph. 33 (Quid ?).—Gallia.

992². chilense, *Endl. ex C. Gay, Fl. Chil.* iii. 180.—Chili.

992³. *decolorans*, Bourg. ex Marès, Cat. Pl. Balear. 139 = Crespianum.
delphinense, Arv. Ess. Pl. Dauph. 33 (Quid ?).—Gallia.
ericoides, Arv. (l. c.) 32 (Quid ?).—Gallia.

995¹. Pseudo-myrianthum, *Gennari, in Mem. Acc. Torin.* II. xvii. (1858) 466.—Hab. ?

995². *simplex*, Arv. Ess. Pl. Dauph. 31 (Quid ?).—Gallia.

GALVEZIA.

996³. d. SACCULARIA, Kellogg (*nunc* **Antirrhinum**).

GANOPHYLLUM.

997². l. falcatum, *Blume*, . . . Malacca ; Austral. ; N. Guin.

GARCINIA.

997³. l. MANGOSTANA, Rumph. ex Gaertn.
a. STALAGMITIS, Murr. in Comm. Gottin. ix. (1789) 173.
XANTHOCHYMUS, Roxb. Pl. Corom. ii. 51. t. 196 (1798).

998¹. l. *collina*, Vieill. . . . 223 = corallina.

998². Kola, *Heckel, in Journ. Pharm. et Chim.* viii. (1883) 88.—Afr. trop.

GARDENIA.

999¹. a. DECAMERIA, Welw. Apont. 579, in obs. [1859].

999². l. *bignoniaeflora*, Welw. . . . 13 = Sherbournia foliosa.
Devoniana, Lindl. . . . 663 = Randia Devoniana.

1000². Pseudo-psidium, *Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 109 (sp. dub.).

GARDOQUIA.

1001¹. a. *obovata*, Griseb. ex Wedd. Chlor. And. ii. 150 = Micromeria boliviana.
l. *origanoides*, Reichb. ex Cham. in Linnaea, vii. (1832) 226.

GARIDELLA.

1001¹. l. *anethifolia*, Salisb. . . . = N. Garidella.
Nigellastrum, Linn. . . . = N. Garidella.
unguicularis, Lam. . . . 2] = N. unguicularis.

1001². l. GARRETIA, Welw. . . . (1858) = **Khaya**, Juss. (Meliac.).
anthoteca, Welw. l. c. = K. anthotheca.

PAG.

GARRYA.

1001². l. *Veatchii*, Kellogg, . . . 40 = flavescens.

1001³. l. GASOUL, Adans.

GASTONIA.

1002¹. l. *aculeata*, Thou. ex Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 109; Hort.

GASTRIDIUM.

1002². l. *scabrum*, Presl, Cyp. & Gram. Sic. 21 =

GAULTHERIA.

1003². l. *Acosta*, Steud. . . . 665 = Vaccinium Acosta.
1003³. *fluviatilis*, A. Cunn. in Ann. Nat. Hist. Ser. I. ii. (1839) 49 =

GAYA.

1004³. a. TETRAPTERA, Phil. in Bot. Zeit. xxviii. (1870) 169.
calyptrata, *H. B. & K. Nov. Gen. et Sp.* v. 268, in obs.—Peruv.
1005¹. a. occidentalis, *H. B. & K. Nov. Gen. et Sp.* v. 268, in obs.—Am. trop.

GAZANIA.

1005³. l. armerioides, *DC. Prod.* vi.

GEIGERIA.

1006¹. l. africana, *Griessel.*

GEMELLA, Lour.

1007². l. *trifolia*, Lour.
a. *trifoliata*, DC. Prod. i. 610 = praec.

GENISTA.

1009¹. l. *aprutia*, Nym. Consp. 151 =
1009². a. *compressa*, Soland. ex A. Cunn. in Ann. Nat. Hist. Ser. I. iii. (1839) 246 = Carmichaelia australis.

GENTIANA.

1013¹. l. *exaltata*, Linn. . . . = Eustoma silenifolium.
1014¹. a. *media* ×, Arv. Ess. Pl. Dauph. 51.—Gallia.

GEOFFRAEA.

1016¹. l. *surinamensis*, Bondt,

GEOPHILA.

1017². d. *reniformis*, Cham. & Schlecht. etc.

GERANIUM.

1020². a. *odoriferum*, G. Don, Gen. Syst. i. 739, in syn. = Pelargonium odoratissimum.
1020³. *patulum*, Soland. ex Forst. f. Prod. 91 = dissectum.
l. *pilosum*, Soland. ex Forst. f.

GERARDIA.

1022³. l. *Nigrina*, Linn. f.
Nigrina, *Thunb.*
nuda, Wood, Am. Bot. & Flor. 231 = filicaulis.
sessiliflora, Vahl, . . . 79 = Alectra melampyroides.
Skinneriana, *Wood, Class-book* (1847), 176; *Am. Bot. & Flor.* 231.
1023¹. *strictiflora*, A. Gray, . . . 294 = seq. (i. e. strictifolia, *Benth.*)

GESNERIA.

1024². l. *corymbosa*, Balb. & Bert. . . . 525 = Pentaraphia longiflora.
elatior, H. B. & K. . . . 393 = Isoloma elatius.
exserta, Sw. . . . 90 = Pentaraphia catalpiflora.
1025². *verticillata*, Cav. . . . 585 = Isoloma verticillatum.

GETHYLLIS.

1025³. a. *bivaginata*, Mass. ex Baker, in Journ. Bot. xxiii. (1885) 227 = Britteniana.
cuspidata, Thunb. ex Baker, (l. c.) = ciliaris.
polyanthera, Soland. ex Baker, (l. c.) xxii. (1884) 148 = ciliaris.

GHINIA.

1027¹. l. *spinosa*, Willd. l. c. = T. spinosa.
verbenacea, Sw. Fl. Ind. Occ. ii. 1089 = T. spinosa.
1027¹. l. GIGANTHEMUM, Welw. Apont. 585 [1859] = **Camoensia**, Welw. (1865) (Legumin.).
scandens, Welw. l. c. = C. maxima.

PAG.

GILIA.

1027³. l. *Dunnii*, Kellogg, . . . 142 = Loeselia effusa.

1029¹. a. GINSENG, Wood, Am. Bot. & Flor. 142 (1871) = **Aralia**, Tourn.
quinquefolium, Wood, l. c. = A. quinquefolium.
trifolium, Wood, l. c. = A. trifolia.

GIRARDINIA.

1029¹. l. condensata, *Wedd.* . . . ix. (1856).

GLADIOLUS.

1029³. a. TILESIA, Thunb. ex Steud. Nom. ed. II. ii. 686 (1841).
1030¹. l. Andrewsii, *Klatt*, . . . xv. (*sic etiam corrigatur apud* brevicollem, *citrinum*, Gawleri, lunulatum, rachidiflorum, *uniflorum*).
1030². *communis*, Thunb.
1030³. a. *hybridus* ×, C. Morr. in Ann. Soc. Bot. Gand, iii. (1847) t. 518.
l. illyricus, *Koch, Syn. Fl. Germ.* ed. I. 699; ed. II. 806.
1031². a. *odoratus* ×, Herb. ex Baker, in Journ. Linn. Soc. xvi. (1877) 117.

GLOBBA.

1033³. l. *crispa*, Retz.
a. *secunda*, Royle, Illustr. Bot. Himal. 357, nomen (Quid?).—Reg. Himal.

GLOBULARIA.

1034¹. l. nana, *Lam.* . . .—Europ.

GLOCHIDION.

1034². a. *bifarium*, Royle, Illustr. Bot. Himal. 327, nomen (Quid?).—Reg. Himal.
1034³. l. Fagueti, *Muell. Arg.* . . . 71.
flexuosum, Muell. Arg. ex Miq. . . . 128 = Phyllanthus japonicus.
1035¹. philippinense, *Benth.*

GLOSSOGYNE.

1035³. l. *brasiliensis*, Gardn. . . . 408 = Isostigma brasiliense.

GLOXINIA.

1036². l. SALISIA, Regel, in Flora, xxxii. (1849) 179.
a. *diversiflora* ×, May, in Rev. Hortic. (1883) 247. f. 45.
maxima ×, Paxt. Mag. Bot. v. (1838) 219.

GLYCERIA.

1037¹ a. HEMIBROMUS, Steud. Syn. Pl. Gram. 317 (1855).
l. aquatica, *Wahlb.*
capillaris, Wahlb.
a. *Catabrosa*, Klatt & Reichb. Fl. Leipz. 96 = Catabrosa aquatica.

GLYCINE.

1038¹. a. glutinosa, *Hoffmgg. Preisverz.* (1841) 78.—*Cf. Linnaea*, xvi. (1842) *Litt.* 269.—Hab.?
1038². pendula, *Blume, ex Miq. Fl. Ind. Bat.* i. 1. 224.—Java.
1038³. l. *totta*, Thunb. . . . = Rhynchosia totta.

GLYCOSMIS.

1039¹. a. *parvifolia*, Steud. Nom. ed. II. ii. 47, in syn., sphalm. = pentaphylla.

GLYCYRRHIZA.

1039². l. lepidota, *Nutt. in Fras. Cat.* (1813); *ex Pursh*,

GMELINA.

1039³. l. Leichhardtii, *F. Muell.*

GNAPHALIUM.

1040². a. *argenteum*, Vell. Fl. Flum. viii. t. 97 (Quid?).—Bras.
1042¹. *ferrugineum*, Wendl. Sert. Hannov. t. 23 = Helipterum ferrugineum.
1043¹. l. keriense, *A. Cunn. in Ann. Nat. Hist.* Ser. I. ii. (1839).
1043³. *nevadense*, Kellogg, . . . 45 = Antennaria dioica.

GNIDIA.

1047¹. a. *ciliata*, Meissn. in DC. Prod. xiv. 563 = Struthiola lucens?

PAG.

GOCHNATIA.

1047³. l. glutinosa, *Klatt*, . . . xv.

GOMESA.

1048³. a. densiflora, *Hoffmgg. Preisverz.* (1841) 78.—*Cf. Linnaea*, xvi. (1842) 265.—Hab.?

GOMPHIA.

1049³. a. nitida, *Vahl, Symb. Bot.* ii. 49.—Ind. occ.

GOMPHOCARPUS.

1050². l. asperifolius, *Walp. Rep.* vi. 486.—Afr. austr. Gerrardi, *Harv.* . . . t. 193.
1050³. M'Kenii, *Harv.* . . . t. 194.

GONOLOBUS.

1054². l. *cynanchoides*, Engelm. & Gray, . . . 251 = Lachnostoma cynanchoides (in Addend.).
a. *exilis*, Miers, in Proc. Roy. Hort. Soc. iv. (1864) 161 = Metastelma stenolobus.
1054³. l. *hispidus*, Hook. & Arn. . . . 295 = Fischeria hispida.
maritimus, R. Br. . . . 35 = Lachnostoma maritimum.
1055¹. *setosus*, Benth. . . . 216 = Fischeria setosa.

GONZALEA.

1055³. l. Petesia, *Griseb. Fl. Brit. W. Ind.* 321; *et in*

GOODYERA.

1056³. l. TUSSACA, Rafin. Préc. Découv. 42 (1814); et in
secundiflora, Lindl. . . . 182; Griff. Notul. iii. 393 = Schlechtendaliana.
similis, Blume, . . . 39 = Schlechtendaliana.

GOSSYPIUM.

1058¹. l. *siamense*, Tenore, . . . 193

GOVINDOOIA.

1059¹. l. *nervosa*, Wight, l. c. = T. angulosa.

1059². l. GRACILEA, Koen. . . . 218 = **Melanocenchris**, Nees (Gramin.).
nutans, Koen. l. c. = M. Rothiana.

GRAMMATOCARPUS.

1059³. a. OCHETOCARPUS, Meyen, Reise, i. 310 (1834).

1060². insere GRANDIERA, Lef. ex Baill. Adansonia, x. (1871) 164 = **Sindora**, Miq. (Legumin.).
cochinchinensis, Lef. l. c. = S. cochinchinensis.

GRATIOLA.

1061¹. l. sexdentata, *R. Cunn. ex A. Cunn.*

GRIFFONIA.

1066¹. a. DACTYLADENIA, Welw. Apont. 572 [1859].

GRINDELIA.

1066⁷. a. THURARIA, [Nutt. in] Fras. Cat. (1813) n. 85.
l. glutinosa, *Dun.* . . . 49.—Peruv.
humilis, Hook. & Arn. l. c. = cuneifolia.
inuloides, *Willd.* . . . 261.—Texas; Mexic.

1066³. l. GRISCHOWIA, Karst. Ausw. Neuer Gew. Venez.

1067². l. GROSOURDYA, Reichb. f.
a. *appendiculata*, Reichb. f. Xenia Orch. ii. 123 = S. pusillus.
emarginata, Reichb. f. (l. c.) = S. emarginatus.
Zollingeri, Reichb. f. l. c. = S. Zollingeri.

GRUMILEA.

1067³. a. *vaginans*, Dalz. in Dalz. & Gibs. Bomb. Fl. 111 = P. truncata.

1068¹. a. GUAIABARA, Plum. ex Boehm. in Ludw. Def. Gen. Pl. 402 (1760) = **Coccoloba**, Linn. (Polygon.).

GUAIACUM.

1068¹. l. *dubium*, Forst. f. Prod. 32 = Ratonia stipitata.

GUETTARDA.

1070³. l. aromatica, *Poepp. & Endl.*
putaminosa, *F. Muell.*
1071¹. *resinosa*, Pers. . . . 200 = Laugeria resinosa.
a. *setosa*, Zipp. ex Korth. in Nederl. Kruidk. Arch. ii. 11. (1851) 211 = Bobea setosa.

PAG.

GUILANDINA.

1071². a. *melanosperma*, Eggers, Fl. St. Croix (1879), 46 = C. Bonducella ?

1072². l. GUNDLACHIA, A. Gray, . . . 100 = **Chrysocoma**, Linn. (Compos.) fide E. L. Greene.

GUNNERA.

1072³. l. *sonchifolia*, Mirb. . . . = Francoa sonchifolia.
a. umbraculifera, *Linden, Cat.* n. 93 (1875) 5, cum ic.—N. Granat.

GUSTAVIA.

1073¹. a. JAPARANDIBA, [Marcgr.] Adans. Fam. ii. 448 (1763).

GYMNADENIA.

1074². l. *cucullata*, Rich. . . . 57 = Habenaria cucullata.

1075¹. l. GYMNANTHUS, Endl. Gen. Suppl. iv. III.

GYMNEMA.

1075³. l. *nitidum*, Benth. . . . 456 (Salaciae sp.).—Afr. trop.

GYMNOBALANUS.

1076¹. l. *Fendleri*, Meissn.

GYMNOPOGON.

1077¹. l. *digitatus*, Nees, . . . 713 = Chloris digitatus.

GYNANDROPSIS.

1078³. a. PEDICELLARIA, Schrank, in Roem. & Usteri, Mag. Bot. iii. VIII. (1790) 10.
PODOGYNE, Hoffmgg. Verz. Pfl. 101, 185 (1824).

GYNERIUM.

1079¹. a. *dioicum*, Dallière, Pl. Ornament. i. (1873) t. 42 = argenteum.

107 ¹. insere GYNICIDIA, Neck. Elem. ii. 81 (1790) = **Mesembryanthemum**, Dill. (Ficoid.).

GYNOPACHIS.

1079². l. *deflectens*, Miq. . . . 222 = Petunga rufidula.
rufidula, Miq. l. c. = Petunga rufidula.

GYNOPOGON.

1079². l. *Hunterii*, Roxb.

GYNOSTEMMA.

1079². a. *trigynum*, M. Laws. in Hook. f. Fl. Brit. Ind. i. 663, lapsu = pedatum.

HABENARIA.

1082². a. DIPLORRHIZA, Ehrh. Beitr. iv. (1789) 147.
MESICERA, Rafin. Neogenyt. 4 (1825).
TRIPLORHIZA, Ehrh. Beitr. iv. (1789) 149.
l. *aestivalis*, Rodrig. . . . 258 = brevidens.
anaplectron, Reichb. f. . . . 81 = cryptophila.

1082³. a. Bennettiana, *Reichb. f. Otia Bot. Hamb.* 104. —Abyss
l. *biflora*, Rodrig. . . . 151 = flexuosa.
a. *bractescens*, Griseb. in Goett. Abh. xxiv. (1879) 340 = Gourlieana.
calcarata, *Benth. & Hook. f. Gen.* iii. 622.—Afr. trop.

1083¹. l. *crucifera*, Reichb. f. & Warm. l. c. 80 = aphylla.
a. *demerarensis*, Reichb. f. in Schomb. Faun. Fl. Brit. Gui. 1123 = parviflora.
l. *estrellensis*, Reichb. f. . . . 813 = secunda.

1083². gracilis, *S. Wats.* . . . 277.

1083³. *johannensis*, Rodrig. . . . 251 = Vaupellii.
a. *Lindeni*, Reichb. f. in Nederl. Kruidk. Arch. iv. (1859) 319 = obtusa.

1084¹. l. macroceratitis, *Willd.*
nasuta, Reichb. f. & Warm. . . . 80 = nuda.

1084². *Pseudo-stylites*, Reichb. f. & Warm. l. c. 79 = hamata.

1084³. a. *radicans*, Griseb. ex Wright, in Sauv. Fl. Cub. 233 = repens.

1085¹. *speciosa*, Poepp. & Endl. . . . 44 = maculosa.
subecalcarata, Rodrig. . . . 159 = Guillemini.
trichoceras, Rodrig. . . . 256 = nuda.
tricuspis, A. Rich. . . . 249 = repens.

HABRANTHUS.

1085³. l. *advena*, Herb. . . . = Hipp. advena.

PAG.

HAEMADICTYON.

1086³. l. *parviflorum*, Benth. . . . 355 = Prestonia parviflora.
suberectum, G. Don, . . . 83.—Ind. occ.

HAEMOSPERMUM.

1087². l. *arboreum*, Reinw. . . . = G. rupestre.

HALENIA.

1089². a. *reflexa*, Griseb. in DC. Prod. ix. 135, sphalm. = deflexa.

1089². insere HALESIA, Loefl. It. Hisp. 188 (1758) = **Trichilia**, P. Br. (Meliac.).

HALESIA.

1089². a. HILLIA, Boehm. in Ludw. Def. Gen. Pl. 71 (1760).

HALOGETON.

1090². l. arachnoideus, *Moq.*

HAPALOSIA.

1093¹. l. *Loeflingiae*, Wall. . . . = P. Loeflingii.

HAPLOSTYLIS.

1093³. a. *Wightianus*, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 101 = R. Wightiana.

HARRISONIA.

1095¹. l. paucijuga, *Oliver*,

HAUYA.

1096¹. a. *arborea*, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 137 = Oenothera arborea.

1096³. l. HAXTONIA, Caley, ex D. Don, in Edinb. N. Phil. Journ. (Jul.–Sept. 1831) 273 = **Olearia**, Moench (Compos.).

HEBEPETALUM.

1097³. l. humiriifolium, *Benth.*
latifolium, *Benth.*

HEDEOMA.

1098². l. *purpurea*, Kellogg, . . . = Mentha Pulegium.

HEDERA.

1098². a. *arborea*, Roehl. Deutschl. Fl. ed. II. ii. 182 = H. Helix.

1098³. *humirepens*, Roehl. (l. c.) = H. Helix.

HEDYOTIS.

1100². a. PANETOS, Rafin. in Ann. Gén. Sc. Phys. v. (1820) 227 ; vi. (1820) 81.
d. PELTOSPERMUM, Benth. (*nunc* **Oldenlandia**).

1101¹. l. *caracasana*, H. B. & K. . . . = Mallostoma caracasanum.
conferta, Ruiz & Pav. . . . = Mallostoma confertum.

1101². *filiformis*, Ruiz & Pav. . . . = Mallostoma filiforme.

1101³. *hyssopifolia*, Cav. . . . = Mallostoma hyssopifolium.
juniperifolia, Ruiz & Pav. . . . = Mallostoma juniperifolium.
laricifolia, Cav. . . . = Mallostoma laricifolium.

1103¹. *thymifolia*, Ruiz & Pav. . . . = Mallostoma thymifolium.
tillaeacea, F. Muell. . . . 39 = Synaptantha tillaeacea.

HEDYSARUM.

1105². a. *hamatum*, Reinw. ex Miq. Fl. Ind. Bat. i. I. 224 = Glycine pendula (in Addend.).

1105³. l. *lansum*, Noronha,

1108¹. a. HEINCHENIA, Webb, ex Hook. f. Bot. Mag. sub t. 6733 (1884), sphalm. = seq.

HEINEKENIA.

1108¹. l. *peliorhyncha*, Masf.

HEKORIMA.

1108². a. *atropurpurea*, Fisch. ex Regel & Tiling, Fl. Ajan. 122 = S. roseus.

HELIANTHEMUM.

1111². a. *sericeum*, Dufour, Lettr. Mont. Maudit. 301 = marifolium ?

1111³. l. Teneriffae, *Coss.* . . . 56.

PAG.

HELIANTHUS.

1112¹. l. *calvus*, Sch. Bip. . . ., nomen = Viguiera calva.
decapetalus, *Linn.*

1112². *Mandonii*, Sch. Bip. . . ., nomen = Vangueria Mandonii.

1112³. microphyllus, *H. B. & K.* . . . 220.—Peruv.

HELICHRYSUM.

1113². a. STOECHAS, Gueldenst. Reisen (1787); ex Ledeb. Fl. Ross. ii. 607 (1846).

1113³. *asperum*, DC. Prod. vi. 264, sphalm. = ericaefolium.

HELICOSTYLIS.

1117². l. *affinis*, Steud. ex Miq. . . . 118 = Poeppigiana.

HELIOSPERMA.

1119³. l. *Veselskyi*, Janka, . . . 65 = S. eriophylla.

HELIOTROPIUM.

1120². l. Clausseni, *DC.*

1120³. *fruticosum*, Benth. & Hook. f. Gen. ii. 843 = Tournefortia fruticosa, *Ker-Gawl.*
gracile, R. Br. in Salt, . . . = zeylanicum.
a. *helianthemoides*, Shuttl. ex A. DC. in DC. Prod. x. 78, 85 = Lithospermum angustifolium ?

1121³. l. *Saltianum*, Steud. . . . 744 = zeylanicum.

1122¹. *tenue*, Willd.
a. *Volaterrae*, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 499, sphalm. = seq.
Voltaireanum ×, Hort.—Cf. Paxt. Mag. Bot. xvi. (1849) t. 99. f. 1.
l. zeylanicum, *Lam.* . . . 94.—Ind. or. ; Afr. trop.

HELIPTERUM.

1122². l. Dimorpholepis, *Benth.*

HELMINTHIA.

1123³. l. *Balansae*, Coss. & Dur. . . . 744 = P. Balansae.
comosa, Boiss. . . . 62 = P. comosa.

1124¹. *lusitanica*, Welw. . . . 515 = P. comosa.

1124¹. a. HELMINTHOTHECA, Vaill. ex Boehm. in Ludw. Def. Gen. Pl. 173 (1760) = seq.

HELOPUS.

1124³. l. *sanguinalis*, Nees, l. c. 17 = Panicum sanguinale.

HELOSCIADIUM.

1124³. a. *heterophyllum*, Steud. Nom. ed. II. ii. 592, in syn., sphalm. = *leptophyllum*.

HEMEROCALLIS.

1125². a. CAMERARIA, Boehm. in Ludw. Def. Gen. Pl. 56 (1760).

HEMIANTHUS.

1125³. l. *adenander*, Wright, . . . 185 = M. adenandrum (in Addend.).
erosus, Wright, ex Griseb. l. c. 984 = M. multiflorum (in Addend.).
multiflorus, Wright, . . . 185 = M. multiflorum (in Addend.).
punctatus, Wright, . . . 184 = M. punctatum (in Addend.).
trisetosus, Wright, . . . 185 = M. trisetosum.

1126³. insere HEMILOBIUM, Welw. Synopse, 20 (1862) = **Apodytes**, E. Mey. (Olacin.).
ficifolium, Welw. l. c. = A. dimidiata.

1128². a. HENNA, Ludw. Def. Gen. Pl. 134 (1747) = **Lawsonia**, Linn. (Lythrar.).

HEPPIELLA.

1129². l. CHEIRISANTHERA, Hort. ex Lindl. & Paxt. Flow. Gard. ii. (1851) 59.

HERACLEUM.

1130¹. a. TORDYLIOIDES, Wall. ex DC. Prod. iv. 199 (1830).

1130². l. Chorodanum, *DC.*

HERITIERA.

1131³. a. SUTHERLANDIA, J. F. Gmel. Syst. 1027 (1791).

PAG.

HERNIARIA.

1134[2]. a. *cashmeriana*, Royle, Illustr. Bot. Himal. 222 = incana ?

1134[3]. l. *setigera*, Gill. . . . 337 (Paronychiae sp.).

HERPESTIS.

1134[3]. l. MONNIERA, B. Juss. ex P. Br.

HESPEROCHIRON.

1137[2]. a. CAPNOREA, Rafin. Fl. Tellur. iii. 74 (1838), nomen prius.
l. *latifolius*, Kellogg, . . . 44 = californicus.

HETERANTHERA.

1038[2]. a. *virginicus*, Steud. Nom. ed. II. ii. 29, in syn. = reniformis.

1139[3]. l. HETEROMORPHA, Cass. . . . Philom.

HETEROPOGON.

1140[1]. a. SPIROTHEROS, Rafin. in Ser. Bull. Bot. i. (1830) 221.

HETEROTHRIX.

1141[3]. l. *Vanheurckii*, Muell. Arg. . . . 164 = Mandevilla Vanheurckii.

1142[1]. l. **HETEROTRICHUM**,
a. OCTONUM, Rafin. Sylva Tellur. 95 (1838).
l. Octonum, *DC.*

HEWITTIA.

1142[2]. a. *linifolia*, Ind. Kew. p. 600[3]. perperam sub C. caespitosa cit. = Ipomoea linifolia.

HIBISCUS.

1145[1]. l. *abutiloides*, Willd. . . . 736 = tiliaceus.
affinis, H. B. & K. . . . 289 (Cienfugosiae sp.).
angustifolius, Hook. & Arn. . . . 152 = Lambertianus.
1145[2]. *bicornis*, G. F. W. Mey. . . . 231 = bifurcatus.
1146[1]. *crenatus*, Splitg. . . . 336 = sororius.
cruentus, Bertol. . . . 10 = H. Sabdariffa.
decipiens, A. St. Hil. . . . 247 = bifurcatus.
digitatus, Cav. . . . 2 = H. Sabdariffa.
Diodon, DC. . . . 449 = furcellatus.
elatus, Sw. . . . 1218 = tiliaceus.
1146[2]. *fluminensis*, Vell. . . . 34 = bifurcatus.
fraternus, Linn. . . . 260 = H. Sabdariffa.
a. *grandiflorus*, Juss. ex DC. Prod. i. 456 = Thespesia grandiflora.
1146[3]. l. Kokio, *Hilleb.*
1147[2]. *multiformis*, A. St. Hil. . . . 246 = furcellatus.
1147[3]. *pernambucensis*, Bertol. . . . 229 = tiliaceus.
1148[1]. *salviaefolius*, A. St. Hil. . . . 249 = Lambertianus.
1148[2]. *sulphureus*, H. B. & K. . . . 289 (Cienfugosiae sp.).
surattensis, *Linn.*
tampicensis, Moric. . . . 14 = Kosteletzkya sagittata.
1148[3]. *trilobatus*, Vell. . . . 29 = furcellatus.
unidens, Lindl. . . . t. 878 = radiatus.
a. *verbasciformis*, Klotzsch, in Schomb. Faun. Fl. Brit. Gui. 991, nomen = spathulatus.
1149[1]. *Youngianus*, Gaudich. . . . 91 = furcellatus.

HIERACIUM.

1149[3]. a. *anserinum, araneosum, bifrons, cirritum, coronariaefolium, dasytrichum, doronicoides, eriophyllum, falcatum, fastidiosum, grandifolium, heterodon, isatifolium, jaceoides, lactucaefolium, lansicum, leucochloma, lychnoides, melandrifolium, Pamphilii, parcepilosum, pulchellum, Ravaudii, Sauzei, spicatifolium, thapsifolium, thapsoides, trichocladum, ustulatum*, Arv. Monog. Pilos. 22–46 (Quid ?).—Gallia.
armerioides, aurantiacoides, biflorum ×, *capitatum, chamaeaurantiacum, chloraefolium, cirrocephalum, corymbuliferum, corymbuloides, fistulosum, florentoides* ×, *Grenierianum, intertextum, lanatellum, multiflorum, nigritellum, palmatum, paniculatum, Pseudo-lanatum, squalidum, umbelluliferum, villosiforme, viride*, Arv. Ess. Pl. Dauph. 30, 40–50 (Quid ?).—Gallia.

PAG.

HIERACIUM:—

1149[3]. a. *brunellaeforme, caeruleum, caesioides, calycinum, cephalodes, cichoraceum, Didieri, elegans, erythrospermum, floccosum, Guillonianum, heterospermum, hypericifolium, intricatum, laureolum, leiopogon, oleovirens, oligocephalum, Pseudo-juranum, Pseudoviride, rapunculoides, Reboudianum, salicifolium, saxetanum, sonchoides, subalpinum, subrude. subsecundum, subvirens, taraxaciforme, tenue, urticaceum, virgultatum, viscosum*, Arv. Monog. Pilos. Suppl. 5–32 (Quid ?).—Gallia.
1151[1]. a. vel l. *callianthum, fragile, lividum, oleosum* (= *oleovirens*), *subluridum, succisoides*, Arv. Add. Monog. Pilos. 8–13 (Quid ?).—Gallia.
1152[2]. *dichotomum*, Schrank, Baier. Reise, 13 (Quid ?).—Germania.

1161[3]. a. HILLIA, Boehm. in Ludw. Def. Gen. Pl. 71 (1760) = Halesia, Linn. (Styrac.).

1161[3]. l. HILSENBERGIA, Boj. . . . (1837); in Rapp. Soc. Hist. Nat. Maurice, xii. (1841) 44.

HIMANTOGLOSSUM.

1161[3]. l. *cucullatum*, Reichb. . . . 120 = Habenaria cucullata.

HIPPEASTRUM.

1162[2]. l. advena, *Herb.*
a. *Hendersoni* ×, Hort. ex Flor. Mag. N. S. (1874) t. 117.
1162[3]. *Taeedianum*, Herb. Amaryll. 425 (Index) = aulicum.

HIPPOCRATEA.

1163[2]. a. PRIONOSTEMMA, Miers, in Trans. Linn. Soc. xxviii. (1872) 354.
1163[3]. lancifolia, *Wils. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 372, in nota.—Jamaic.

HIPPOCREPIS.

1164[1]. a. *alpestris*, Arv. Ess. Pl. Dauph. 25 (Quid ?).—Gallia.

HIRAEA.

1165[2]. l. chlorocarpa, *A. Juss. in Ann. Sc. Nat.* Sér. II. xviii. (1840) 259.
chrysophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843).

1166[1]. l. HIRNELLIA, Cass. . . . = Myriocephalus, Benth. (1837).

HIRTELLA.

1166[1]. l. ZAMZELA, Rafin.

HITCHENIA.

1167[1]. l. glauca, *Wall.* . . . 216.

HOFFMANNIA.

1167[3]. l. *lanceolata*, Griseb. . . . 504 = Lasianthus Moralesii.

HOFFMANSEGGIA.

1168[1]. a. LARREA, Orteg. Hort. Matr. Dec. 15. t. 2 (1797).

HOLCUS.

1169[1]. l. *ciliatus*, Roxb. . . . 318 = Arundinella holcoides.
nervosus, Roxb. . . . 318 = Arundinella brasiliensis.
1169[2]. d. *saccharatus*, Griseb.

HOLIGARNA.

1169[2]. a. KATOUTSJEROE, Adans. Fam. ii. 84, 534 (1763).

HOLOSTEMMA.

1170[2]. l. *tuberculatum*, Blume, . . . (= *Cynoctonum*

HOMOEATHERUM.

1172[1]. l. *chinense*, Nees, l. c. = A. fastigiatus.

1173[1]. l. HOOREBEKIA, Corneliss. . . . 12; *chiloënsis*, Corneliss. l. c. 13, cum ic. =

HOPEA, Linn.

1173[1]. a. *lucida*, Thunb. Ic. Jap. t. 14 = S. japonica ?

PAG.

HOPEA.

1173[1]. a. *conduplicata*, Buch.-Ham. in Mem. Wern. Soc. vi. (1832) 299 = Dipterocarpus alatus.
laevis, Buch.-Ham. (l. c.) = Dipterocarpus turbinatus.
d. lucida, *Thunb.*
a. *scabra*, Buch.-Ham. (l. c.) 300 = Dipterocarpus scaber.
trinervis, Buch.-Ham. (l. c.) = Dipterocarpus turbinatus.

HORDEUM.

1173[3]. l. *Aegiceras*, Nees, in Royle, Illustr. Bot. Himal. 418. t. 97. f. 2 =
1174[1]. *himalayense*, Schult. . . . 481 = vulgare.
a. *hymalayense*, Hort. Monsp. ex Irmisch, in Linnaea, xiii. (1839) 125 = praec.
l. *ithaburense*, Boiss. . . . 70 = spontaneum.
1174[2]. marinum, *Huds.* . . . 57.—Europ.; Afr. bor.; Oriens.
maritimum, With. . . . ed. III. 172 = marinum.
spontaneum, *C. Koch*, . . . 430.—As. Min.; Oriens.
villosum, 'Bertol.' ex Steud.

HORSFIELDIA.

1175[1]. a. *aculeata*, Benn. Pl. Jav. Rar. 123. t. 26 = peltata.
peltata, *Benth. & Hook. f. Gen.* i. 937.—Java.

HOULLETIA.

1176 l. Landsbergi, *Linden & Reichb. f.* . . . 2.—Costa Rica.

HOUMIRI.

1176[2]. l. *gabonensis*, Baill. . . . 52 = Aubrya gabonensis.

HOVEA.

1176[3]. a. PHUSICARPOS, Poir. Encyc. Suppl. iv. 399 (1816).

HOWARDIA, Wedd.

1177[2]. l. *febrifuga*, Wedd. l. c. 67 = P. febrifugus.

HOYA.

1177[3]. l. *cochinchinensis*, Schult. . . . 52 (Ceropegiae sp. ?).

HUERNIA.

1178[3]. l. ocellata, *Schult.*

HUGONIA.

1179[1]. l. *angolensis*, Welw. . . . 271 = macrocarpa.
macrocarpa, Welw. . . . 21 = Afzelii.

HUTCHINSIA.

1180[2]. l. *tibetica*, Thoms. . . . 900 = Capsella Thomsoni.

HYALIS.

1181[2]. a. spartioides, *Benth. & Hook. f. ex Hieron. Pl. Diaph.* 164.—Bolivia.

HYDRILLA.

1183[1]. a. IXIA, Muhl. ex Spreng. Syst. i. 171 (1825).
subulata, Royle, Illustr. Bot. Himal. 376 = verticillata ?
l. verticillata, *Royle*, (*l. c.*) 376; *Presl*,

HYDROLEA.

1185[1]. l. *auriculata*, Moç. & Sesse, ex Choisy, in Ann. Sc. Nat. Sér. I. xxx. (1833) 239 =
decurrens, Moç. & Sesse, ex Choisy, (l. c.) 243 =
1185[2]. *ovata*, Nutt. ex Choisy, (l. c.) 239 =
radians, Moç. & Sesse, ex Choisy, (l. c.) 242 =
rupincola, Moç. & Sesse, ex Choisy, (l. c.) 242 =
scorpioides, Moç. & Sesse, ex Choisy (l. c.) 249 =
tenella, Moç. & Sesse, ex Choisy, (l. c.) 244 =
violacea, Moç. & Sesse, ex Choisy, (l. c.) 245 =

1185[3]. a. HYDROPHILA, Ehrh. Beitr. iv. 146 (1789) = Tillaea, Mich. (Crassul.).

HYDROPHYLLUM.

1185[3]. l. *hispidum*, Riddell, . . . 84 = macrophyllum.

HYMENANTHERA.

1187[2]. l. oblongifolia, *A. Cunn.* . . . 124.—Ins. Norfolk.

HYMENOCALLIS

1187[3]. a. TROXISTEMON, Rafin. l. c. (1836).

PAG.

HYMENODICTYON.

1188³. l. KURRIA, Hochst. & Steud. in Flora, xxiv. (1841) I. Intell. 28; xxv.

1188³. a. HYMENOGONIUM, Rich. ex Lebel, in Mém. Soc. Sc. Cherb. xiv. (1868) 30 = **Spergularia**, J. & C. Presl (Caryophyll.).

HYOSCYAMUS.

1189³. l. *Datora*, Forsk. . . . = muticus.
1190¹. *physaloides*, Linn. . . . 180 = Scopolia physaloides.

HYPERICUM.

1195². l. Thymopsis, *Boiss.*

HYPHAENE.

1195³. a. *coriacea*, Welw. Apont. 584 = benguellensis.
l. Schatan, *Boj.* . . . 308.—Madag.

HYPOBATHRUM.

1195⁵. a. ERIOSTOMA, Boiss. ex Baill. Adansonia, xii. (1878) 208.

HYPOCYRTA.

1197¹. l. *gracilis*, Mart. . . . 219 =

HYPOËSTES.

1197². a. *purpurascens*, Royle, Illustr. Bot. Himal. 298, sphalm. = purpurea.

HYPOLEPIS, Pers.

1198¹. l. *sanguinea*, Pers. l. c. = C. dioicus.

HYPOXIS.

1199¹. a. *barbacenioides*, Harv. ex Baker, l. c. xiii. (1875) 234 = Vellozia elegans.
1199³. *vellosioides*, Harv. ex Baker, (l. c.) xiii. (1875) 233 (= *Xerophyta clavata*).—Afr. austr.

HYPSELA.

1199³. l. *reniformis*, Presl, . . . 45 = Lysipomia reniformis.

HYPTIS.

1201¹. l. *menthodora*, Schrank, in Syll. Ratisb. ii. (1828) 56 =
nepalensis, Lehm. ex Benth. Lab. Gen. et Sp. 141 =

HYSSOPUS.

1202¹. a. *anethiodorus*, [Nutt. in] Fras. Cat. (1813) n. 41 = Lophanthus anisatus.

1204¹. a. ICHTHYOMETHIA, P. Br. Hist. Jamaic. 296 (1756) = **Piscidia**, Linn.

IGUANURA.

1205¹. l. SLACKIA, Griff. . . . (1854).

ILEX.

1205³. a. *Congonhas*, Liais, Brésil, 563 = paraguayensis.
1207¹. l. *rotunda*, Buch.-Ham. ex Wall.
a. *serrata*, Royle, Illustr. Bot. Himal. 167, nomen (Quid ?).—Reg. Himal.

ILLAIREA.

1207². l. *canarinoides*, Lenné & C. Koch, l. c. = L. canarinoides.

IMPATIENS.

1209³. a. *Inglisii*, Royle, Illustr. Bot. Himal. 150, nomen (Quid ?).—Reg. Himal.
1210². *pumila*, Royle, (l. c.) 151, nomen (Quid ?).—Reg. Himal.
speciosa, Royle, (l. c.) nomen (Quid ?).—Reg. Himal.

INCARVILLEA.

1211². l. *Koopmanni*, Lauché, in Deutsch. Gart. 1880 (1881), 89, cum ic. =

INDIGOFERA.

1211². a. ANIL, March. ex Ludw. Def. Pl. 117 (1737).
1211³. *Arghawan*, Royle, Illustr. Bot. Himal. 193 = Sophora mollis.
1212¹. l. bracteolata, *DC.*
1212³. *disperma*, Linn. . . . 232 = tinctoria.

PAG.

INDIGOFERA :—

1214². pinifolia, *Baker*, (ante *platyphyllum* ponenda).
platyphylla, Miq. . . . 292 = Tephrosia sennoides.
a. *plicaefolia*, Steud. Nom. ed. II. ii. 710, in syn. = Psoralia obtusifolia.

INGA.

1215². a. FEVILLAEA, Linn. Syst. ed. I (1735).
1215³. l. Bourgoni, *DC.*
1216¹. Diadema, *Mart.*

1218³. l. INGENHOUZIA, Bert. ex DC. in Guill. Arch. Bot. ii. (1833) 333 = **Rhotinodendron**, Meissn. (Compos.).

INOCARPUS.

1218³. a. RENIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

INULA.

1219¹. l. ENULA, Caesalp. ex Boehm. in Ludw. Def. Gen. Pl. 184 (1760); Neck.
a. MYRIADENUS, Cass. in Bull. Soc. Philom. (1817) 138.
ORSINA, Bertol. in Ann. Stor. Nat. Bologn. I. ii. (1829) 362.
1219³. l. intermixta, *Kern.*
a. *lanigera*, Burm. f. Fl. Cap. Prod. 27 (Quid ?). —Afr. austr.

IONIDIUM.

1221². l. concolor, *Steud. Nom.* ed. II. ii. 770, in syn.; *Benth. & Hook. f.*

1222¹. l. **IONOPSIDIUM**, forsan retinendum, cum spp. 2, scilicet l. acaule, albiflorum.

IPOMOEA.

1223². l. *arachnosperma*, Welw. Apont. 588 = pilosa.
a. aurea, *Kellogg, ex Curran, in Bull. Calif. Acad.* i. III. (1885) 143.—Calif.
1224¹. l. *Clappertoni*, R. Br. . . . . 240 = aquatica.
1224². decora, *Vatke & Hildeb.* . . . (1869).
1224³. *flavescens*, Steud. . . . 816 = Lepistemon flavescens.
a. Forsteri, *A. Gray, ex H. Mann, in Proc. Am. Acad.* vii. (1866) 195.—Ins. Sandvic.
1225². l. *humilis*, G. Don, . . . 267 = carnosa.
incurva, G. Don, . . . = carnosa.
1225³. *lanceolata*, G. Don, . . . = martinicensis.
lilacina, Hemsl. Biol. Centr. Am. Bot. ii. 389 (an praec. ?).—Mexic.
1226¹. *Mendesii*, Welw. Apont. 584 = tuberculata.
multisecta, Welw. Apont. 589 = coptica.
1226². *oblonga, Hemsl. Biol. Centr. Am. Bot.* ii. 391.—Mexic.
a. *obscura*, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 244 = Forsteri.
1227¹. l. *radiatifolia*, Kellogg, . . . 163 = leptotoma.
1227². Schiedeana, *Ham.*
a. *sepiaria*, Seem. in Bonplandia, ix. (1861) 258 = Forsteri.
1227³. *splendens*, G. Don, in Edinb. Phil. Journ. xi. (1824) 343, nomen = digitata.
1228¹. l. *teretistigma*, Choisy, l. c. 376 = Hewittia bicolor.
a. tragulifera, *Miers, in Proc. Roy. Hort. Soc.* iv. (1864) 160.—N. Granat.
1228². l. *umbellifera*, Choisy, . . . 389 = umbellata ? *Linn.*

IRINA.

1229². l. *alnifolia*, Blume, . . . 117 = P. eximia.
tomentosa, Blume, l. c. = P. eximia.

IRIS.

1229³. l. Aitchisoni, *Boiss. Fl. Orient.* v. 123.—Afghan.
1230³. *Helena*, Boiss. (l. c.) 132, in syn. =
1232². *tuberosa*, Linn. . . . 40 = Hermodactylus tuberosus.

IRLBACHIA.

1232³. l. *recurva*, Prog. . . . 229 = Pagaea recurva.
subcordata, Prog. l. c. = Pagaea recurva.

PAG.

ISACHNE.

1233¹. a. multiflora, *Trimen, Cat. Pl. Ceyl.* 104.—Zeylan.
l. *tricarinata*, Roth, . . . 57 = Panicum ovalifolium.

ISATIS.

1233². l. Djurdjurae, *Coss. & Dur.*

ISCHAEMUM.

1234¹. a. *dactyloideum*, Montand. Guide Bot. 353 = Andropogon Ischaemum.
1234². l. pilosum, *Wight, in Madr. Journ. Sc.* Ser. I. ii. (1835) 189; *Trimen*,

ISOLEPIS.

1237³. l. *podocarpa*, Boeck. . . . 179 = Hemichlaena angustifolia.

ISOPOGON.

1239³. a. *anemifolius*, Steud. Nom. ed. II. ii. 401, in syn., sphalm. = seq.

ISOPYRUM.

1240¹. l. *Clarkii*, Kellogg, . . . 131 = stipitatum.

ITEA.

1240³. a. REINIA, Franch. & Sav. Enum. Pl. Jap. ii. 314 (1876).

1243². l. IXINA, Rafin. . . . = IXINE, Loefl.

1243². insere IXINE, Loefl. St. Hisp. 195 (1758) = **Krameria**, Linn. (Polygal.).

IXORA.

1244³. a. *Pilgrimi* ×, Hort.—Cf. Flor. Mag. N. S. (1880) t. 428.
1245¹. *Williamsii* ×, Hort.—Cf. Gard. Chron. (1872) 642. f. 162.

JACOBINIA.

1246³. a. ramosissima, *Benth. & Hook. f. Gen.* ii. 1119, in obs.—Bras.

JACQUEMONTIA.

1246³. a. PENTANTHUS, Rafin. Fl. Tellur. iv. 70.

1249¹. insere JAPARANDIBA, [Marcgr.] Adans. Fam. ii. 448 (1763) = **Gustavia**, Linn. (1785) (Myrtac.).

JASIONE.

1249². l. humilis, *Loisel.*

JASMINUM.

1250². d. *humile*, Heyne, etc.

JASONIA.

1251¹. d. MYRIADENUS, Cass., et ORSINA, Bertol. (*nunc* Inula).

1252³. l. **JEFFERSONIA**, Bart. . . . (1793) 342.
binata, *Bart.* . . . (1793) 342.

JUGLANS.

1254³. l. *baccata*, Linn. . . . 1272 = Picrodendron arboreum.
1255¹. a. *dissecta*, Hort. Taur. 910; ex C. DC. in DC. Prod. xvi. II. 136 = regia.

1255³. l. JUNCAGO, Tourn.

JUNCUS.

1255³. a. STYGIARIA, Ehrh. Beitr. iv. (1789) 146.

1260¹. a. JUNGIA, Boehm. in Ludw. Def. Gen. Pl. 92 (1760) = **Dianthera**, Gronov. (Acanth.).

JUNIPERUS.

1260³. l. *gracilis*, Hort. . . . = Cupressus torulosa.
1261¹. a. *nana*, Willd. Sp. Pl. iv. 854 = communis.
plochyderma, Sargens, in Rep. Tor. N. Am. (U. St. 10th Census, ix.) (1864) 181, sphalm. = pachyphlaea.

JURINEA.

1262¹. l. heterophylla, *Boiss. Fl. Orient.* iii. 571.

JUSTICIA.

1265¹. l. elegans, *Pohl*, . . . 382.—Bras.
1265². *flexibilis*, Boj. . . . = Hypoëstes flexibilis.
1267¹. a. *obscura*, Vahl, ex Benth. Lab. Gen. et Sp. 323, sphalm. = diffusa ?
1267². l. *procumbens*, Blume, . . . = Blumeana.
1267³. *salicifolia*, Blume, Bijdr. 787 = Peristrophe salicifolia.

TOMUS II.

PAG.

KADSURA.

1¹. a. PULCHERIA, Noronha, in Verh. Batav. Gen. v. (1790) Art. IV. 3.

KAKOSMANTHUS.

2¹. l. *macrophyllus*, Hassk. . . . 578 = P. macrophylla.

2¹. l. KALABOTIS, Rafin.

2³. l. KALI, Tourn. ex Ludw. Def. Gen. Pl. 231 (1747) =

3³. insere KATAKIDOZAMIA, Haage & Schmidt, ex Regel, Gartenfl. (1876) 5 = CATAKIDOZAMIA, T. Hill = **Macrozamia**, Miq. (Cycad.).

3³. insere KATOUTSJEROE, Adans. Fam. ii. 84, 534 (1763) = **Holigarna**, Buch.-Ham. (Anacard.).

KAYEA.

4¹. a. PLINIA, Blanco, Fl. Filip. ed. I. 423 (1837). VIDALIA, F. Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 17 (1880).
l. lepidota, *Pierre, l. c.*—Cochinch.; Ins. Philipp.

KENTIA.

5¹. l. *Joannis*, F. Muell. Fragm. vii. 101 = *Storckii*, F. Muell. (l. c.) = *subglobosa*, F. Muell. l. c. =

5². l. KENTROPHYLLUM . . . = **Carthamus**, Linn.

KERATEPHORUS.

5³. l. *Leerii*, Hassk. . . . 579 = P. Leerii.

KHAYA.

6². a. GARRETIA, Welw. Apont. 587 [1859].

KIESERA.

6³. l. *gracilis*, Miq. . . . 291 = T. sennoides.

KIGELIA.

6³. l. SOTOR, Fenzl, . . . Versamml.

KIPPISTIA.

7¹. l. *suaedifolia*, F. Muell. = M. Kippistiana.

KLEINIA.

8¹. a. *Mandraliscae*, Tineo, in Ann. Agr. Sicil.; ex Lojac. Isole Eolie, 111.—Hab. ?

KNAUTIA.

8². a. *alpina*, Arv. Ess. Pl. Dauph. 35 (Quid ?).—Gallia.

8³. *glandulosa*, Arv. (l. c.) (Quid ?).—Gallia.
latifolia, Arv. (l. c.) (Quid ?).—Gallia.
l. *lyrophylla*, Panč. . . . 128 = S. macedonica ?
macedonica, Griseb. . . . 178 = S. macedonica.

KOBRESIA.

9³. l. laxa, *Royle, Illustr. Bot. Himal.* 411; *Boeck.*

10². l. KODDA-PAIL, Adans. Fam. ii. 75 (non 25)

KOELERIA.

11¹. a. Philippiana, *E. Desv. in C. Gay, Fl. Chil.* vi. 353, in nota.—Chili.

11³. l. KOELREUTERIA . . . = **Marsdensia**, R. Br. (Asclep.).

KOPSIA, Dum.

12³. l. *aegyptiaca*, Caruel, . . . = O. aegyptiaca.
arenaria, Dum. . . . = O. arenaria.
Borkhauseni, Caruel, . . . = O. arenaria.
lavandulacea, Caruel, . . . = O. lavandulacea.
longiflora, Dum. . . . = O. aegyptiaca.
ramosa, Dum. . . . = O. ramosa.
a. *stricta*, Caruel, in Parl. Fl. Ital. vi. 356 = O. Schulzii.

KRAMERIA.

13². a. IXINE, Loefl. It. Hisp. 195 (1758).

PAG.

KREMERIA.

14¹. l. *Cordylocarpus*, Coss. & Dur. l. c. 672 = M. Cordylocarpus.

14². l. KROMBHOLZIA, Rupr. . . . (1842).

14³. l. KTENOSACHNE, Steud. . . . = **Prionanthium**, Desv. (Gramin.).

KYLLINGA.

16¹. l. *aphylla*, Kunth, . . . 127 = peruviana.

LABOURDONNAISIA.

17². a. EICHLERIA, Hartog, in Journ. Bot. xvi. (1878) 72.
MURIEA, Hartog, l. c. 145 (1873).

LACHENALIA.

18². l. pallida, *Soland.*

LACHNOPETALUM.

19¹. l. *glabrum*, Turcz. l. c. = R. lachnopetala.

LACHNOSTOMA.

19². a. cynanchoides, *Benth. & Hook. f. Gen.* ii. 766.—Am. bor. occ.
l. martimum, *Benth. & Hook. f. (l. c.)*; *Nichols.*

19². l. LACINARIA, Hill,
a. LACINIARIA, Hill, Hort. Kew. 70 (1768) = praec.

LACMELLIA.

19³. l. florbunda, *Benth. & Hook. f.* . . . 694.

LACTUCA.

20³. l. muralis, *Fresen.* . . . ; *G. F. W. Mey.*
21¹. *Schimperi*, Jaub. & Spach,

LAETIA.

22². a. THAMNIA, P. Br. Hist. Jamaic. 245 (1756).

LAGENOPHORA.

23³. l. Enphysopus, *Hook. f.*

LAGERSTROEMIA.

23². a. ansontera, *Koehne, in Engl. Bot. Jahrb.* iv. (1883) 407.—Cochinch.

LAGERA.

24¹. l. *ptrodonta*, Sch. Bip. . . . = purpurascens.

LAGINCULARIA.

25¹. a. RHIZAERIS, Rafin. Sylva Tellur. 90 (1838).

LAMARCKIA, Vahl,

25². a. *cocinea*, Vahl, l. c. = M. coccinea.

LANNEOMA.

28² a. *triphylla*, March. Rev. Anacard. 157, sphalm. = seq.

28². l. LANSBERGIA . . . = **Trimeza**, Salisb.

LANSIUM.

28². a. LUTEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

28³. l. *montanum*, Royle, Illustr. Bot. Himal. 140; Steud.

LANTANA.

28³. l. *rubia*, Wall. Cat. n. 1823 =

29¹. a. *lochana*, Buch.-Ham. ex Wall. Cat. p. 87. n. 1821 = alba.

LATHUM.

29³. a. *bnoniense*, Montand. Guide Bot. 265 = R. pulcher.
btusatum, Montand. (l. c.) = R. obtusifolius.
otundifolium, Montand. (l. c.) 267 = R. alpinus.

LARIX.

31³. l. *Deodara*, Royle, Illustr. Bot. Himal. 457 (Index); C. Koch,

PAG.

LASERPITIUM.

32³. a. *alpestre*, Arv. Ess. Pl. Dauph. 31 (Quid ?).—Gallia.

LASIAGROSTIS.

33². l. *tremula*, Rupr. . . . 35.—Kashgar.

LASIANTHERA.

34¹. a. malaccensis, *Mast. l. c.* 584.—Malacca.

LASIOLYTRUM.

35². l. *pilosum*, Steud. ex Jardin, . . . 299 = Panicum sanguinale.

36³. insere LASSONIA, Buc'hoz, Pl. Nouv. Découv. t. 19 (1779). Genus spurium *MAGNOLIACEARUM*.
heptapeta, Buc'hoz, l. c.—China.
quinquepeta, Buc'hoz, l. c.—China.

LATHYRUS.

37³. l. *ensifolius*, Badaro,

38¹. *macrorrhizus*, Wimm. . . . = montanus.

39¹. a. Tommasinii, *Sprun. ex Marches. in Boll. Soc. Adriat. Sc. Nat. Trieste*, v. (1880) p. xxviii, nomen.—Europ.

LAUNAEA.

39². l. ZOLLIKOFERIA, DC. . . . (1838).

LAURIDIA.

40¹. l. *rupicola*, Eckl. & Zeyh. . . . (= *Mystroxylon reticulatum*).—Afr. austr.

LAURUS.

41³. l. *lucida*, Thunb. Fl. Jap. 174 = Symplocos japonica ?

42³. *semecarpifolia*, Wall. . . . = Alseodaphne semecarpifolia.

43¹. l. LAVALLEA, Baill. . . . (1862) 361 =

LAVANDULA.

43². l. rotundifolia, *Benth.*

LAVATERA.

43³. d. *hispida*, Sims, etc.

44¹. l. *rotundata*, Láz. . . . (1881),

LAWSONIA.

44³. a. HENNA, Ludw. Def. Gen. Pl. 134 (1747).

LAYIA.

45¹. l. galardioides, *Hook. & Arn.*

46¹. l. LEBRETONIA, Schrank,
a. *candida*, Hort. ex Lem. Jard. Fleur. iii. (1852) sub t. 281 = Urena stellipila.

LECHEA.

46.². a. maritima, *Leggett, ex Britton, Prel. Cat. N. Jersey* (1881), 13.—Am. bor.
l. *Novae-Cesareae*, C. F. Austin, . . . 81 = minor.
thesioides, Spach, . . . 285 = tenuifolia.
thymifolia, Michx. . . . 77 = minor.
a. *thymifolia*, Pursh, Fl. Am. Sept. 91 = maritima.

LECONTEA.

46³. a. SIPHOMERIS, Boj. Hort. Maurit. 170 (1837).

LEDUM.

47³. l. *californicum*, Kellogg, . . . 14 = glandulosum.

LEIANTHUS.

49¹. l. OMPHALOSTIGMA, Reichb. Nom. 132 (1841).

LEIDESIA.

49². a. *Sonderiana*, Muell. Arg. in DC. Prod. xv. II. 699 = capensis.

LEMNA.

50². a. LENTICULARIA, Mich. ex Montand. Guide Bot. 308 (1868).

PAG.
51¹. insere LENTICULARIA, Mich. ex Montand. Guide Bot. 308 (1868) = Lemna, Linn.
monorhiza, Montand. l. c. = Lemna minor.

LEOCHILUS.
51². a. pygmaeus, *Benth. & Hook. f. Gen.* iii. 565.—Guatemala.

LEONTICE.
51³. l. LEONTOPETALON, Tourn. ex Ludw. Def. Gen. Pl. 192 (1747).

LEONTODON.
52¹. l. *bessarabicus*, Fisch. ex Link, Enum. Hort. Berol. ii. 283 =
52³. *nudicaulis*, Banks, . . . 535 = saxatilis.
53¹. *taraxacoides*, Hoppe & Hornsch. Tageb. 174; et ex
tenuifolius, Hoppe & Hornsch. (l. c.) 173; et ex

53². l. LEONTOPETALON, Tourn. ex Ludw. Def. Gen. Pl. 192 (1747).

LEPERIZA.
55¹. l. *eucrosioides*, Baker, . . . 170 = Phaedranassa eucrosioides.

LEPIDADENIA.
55². l. *Seloang*, Miq.

LEPIDAGATHIS.
55². l. alopecurioidea, *R. Br.*
ceylanica, *Nees*,
55³. *nervosa*, Wight,
56¹. subuninervia, *C. B. Clarke*.

LEPIDIUM.
56². a. *bipinnatum*, Gibert, Enum. Pl. Montevid. 66 = praec.
l. *brevicaule*, Hoppe & Hornsch. Tageb. 278, nomen; et ex

LEPIDOTHECA.
59¹. a. *Santolina*, Walp. Rep. ii. 639, sphalm. = seq.
l. *suaveolens*, Nutt. in Am. Phil. Trans. vii. (1841) [455] 397 =

LEPIGONUM.
59³. a. *roseum*, Rafin. Fl. Tellur. iii. 81 = Arenaria geniculata?

LEPTADENIA.
60³. l. = L. Spartium (pluries in synon.).

LEPTOCHLOA.
62¹. l. coerulescens, *Steud.* . . . 209.—Afr. trop.
62². gracilis, *Nees*, . . . 4.—Am. austr.
mutica, Steud. . . . = virgata, *Beauv.*
obtusiflora, *Hochst.*

62³. l. LEPTOCODON, Sond. in Harv. & Sond.

63². l. LEPTOPAETIA, Harv. Gen. S. Afr. Pl.

LEPTOTHERIUM.
65³. a. *Royleanum*, Nees, in Royle, Illustr. Bot. Himal. 417, nomen (Quid?).—Reg. Himal.

LERCHEA.
66². l. *calycina*, A. Gray, . . . 311 = Xanthophytum calycinum.

66³. a. LERISCA, Schlecht. in Bot. Zeit. iii. (1845) 476 = **Cryptangium**, Schrad. (Cyperac.).

LESSERTIA.
67². l. *astragalina*, Pappe,
67³. pauciflora, *Harv. l. c.* 222.
polystachya, *Harv.* . . . 216.

LETTSOMIA.
68². l. *sumatrana*, . . . = Ipomoea staphylina.

LEUCADENDRON.
68². a. *conocarpum*, Steud. Nom. ed. II. ii. 399, in syn. = Leucospermum conocarpum.
68³. *ellipticum*, Steud. (l. c.) = Leucospermum ellipticum.

LEUCANTHEMUM.
69³. a. *laciniosum*, Arv. Ess. Pl. Dauph. 36 (Quid?).—Gallia.

PAG.
LEUCAS.
70¹. l. *affinis*, R. Br. . . . nomen = urticifolia.
dimidiata, Benth. in Wall. Cat. n. 2521; et in DC.
70². a. *pallida*, Steud. Nom. ed. II. ii, in syn. = Leonotis pallida.

LEUCERIA.
70³. l. eriochlaena, *Remy*,

71². l. LEUCOGLOCHIN, Ehrh. Beitr. iv. (1789) 146; Heuff.

LEUCOJUM.
71³. a. *aestivale*, Steud. Nom. ed. II. ii. 196, in syn. = seq.

LEUCOMERIS.
71³. a. *scandens*, Sch. Bip. in Flora, xxxvi. (1854) 275 = Erigeron scandens.

72¹. a. LEUCONOCARPUS, Spruce, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 295 = **Moronobea**, Aubl. (Guttif.).
riparius, Spruce, l. c. = M. riparia.

72¹. l. LEUCOPHAË, Webb & Berth.

LEUCOPOGON.
72³. l. exolasius, *F. Muell.*
fimbriatus, *Stschégl.*

LEUCOSPERMUM.
74¹. l. involucratum. *Roem. & Schult.*
a. *tortum*, Steud. Nom. ed. II. ii. 401, in syn., sphalm. = Leucadendron tortum.

74². linea prima l. **LEUCOSYKE.**

LIATRIS.
76¹. l. *cordata*, D. Don, ex Royle, Illustr. Bot. Himal. 247; et ex

LIBANOTIS.
76². a. *athamantina*, Roehl. Deutschl. Fl. ed. II. ii. 215 = S. Libanotis.
l. *condensata*, Fisch. . . . 724 = S. condensatum.
filifolia, Roehl. (l. c.) =

LIBIDIBIA.
76³. l. *coriaria*, Schlecht. . . . = C. coriaria.

LIBONIA.
77¹. l. *floribunda*, C. Koch,
a. *penrhosiensis* ×, Hort. ex Carr. in Rev. Hortic. (1876) 50, cum ic.

LICANIA.
77¹. a. *floribunda*, Benth. ex Hook. f. in Mart. Fl. Bras. xiv. II. 24 = Moquilea utilis.

78¹. a. LIEUTAUTIA, Buc'hoz, Pl. Nouv. Découv. t. 7. (1779) = **Miconia**, Ruiz & Pav. (1794) (Melastom.).

LIGHTFOOTIA.
78². l. capillaris, *Buek*,

79¹. l. LIGIA, Fasano, in Att. Acc. Nap. 187 (1788), 245 [t. 18. f. 2].

LIGUSTICUM.
79². l. *Ajawain*, Roxb. . . . 170 =
a. *Ajowan*, Royle, Illustr. Bot. Himal. 230 = praec.
l. *angustilobum*, Phil. . . . 95 = Pimpinella angustiloba.

LILIORHIZA.
81². a. *Piattiana*, Kellogg, ex Curran, in Bull. Calif. Acad. I. iv. (1885) 150 = F. pluriflora?
viridis, Kellogg, in Proc. Calif. Acad. ii. (1863) 48 = F. lanceolata.

LILIUM.
81². l. *andinum*, [Nutt. in] Fras.
82¹. giganteum, *Wall.* . . . t. 12.—Reg. Himal.
83¹. a. *umpquense*, A. Murr. Oreg. Circ. 1 (Quid?).—Am. bor.

83². l. **LIMNANTHEMUM**, S. T. Gmel.
83³. Nymphoides, *Hoffmgg. & Link*,
a. *sarmentosum*, Steud. Nom. ed. II. i. 769, in syn., sphalm. = geminatum.

PAG.
84³. l. LIMNOGENNETON, Sch. Bip. . . . = **Siegesbeckia**, Linn.

LIMNOSIPANEA.
84³. a. SIPANIA, Seem. in Bot. Voy. Herald, 136 (1853).

LIMODORUM.
85². a. *mauritanicum*, Sw. Adnot. Bot. 59 = Angraecum gladiifolium?

LIMONIA.
85³. l. *pusilla*, Gaertn. l. c. 279 = S. Schreberi.

86¹. insere LIMONIAS, Ehrh. Beitr. iv. (1789) 147 = **Epipactis**, Adans. (Orchid.).

LIMOSELLA.
86². a. *lacustris*, Steud. Nom. ed. II. ii. 350, in syn., sphalm. = Littorella juncea.

LINARIA.
86¹. a. *affinis*, Arv. Hist. Pl. Dauph. 56 (Quid?).—Gallia.
86³. Aristides, *Battand. in Bull. Soc. Bot. Fr.* xxxii. (1885) 337.—Alger.
87². l. italica, *Trevir. Ind. Sem. Hort. Vratisl. App.* ii. (1820) 2; *et in*

LINDERNIA.
89³. a. *inaperta*, Thore, ex Maxim. in Bull. Acad. Pétersb. xx. (1875) 449 = V. erecta.

LINOCIERA.
90¹. l. *cotinifolia*, Vahl, . . . 46 = Chionanthus virginica.

LIPARIS.
93³. a. *diphyllos*, Nimmo, in J. Grah. Cat. Bomb. Pl. 252 = Habenaria diphylla.
l. Galeottiana, *Hemsl.* . . . 559.—Mexic.

LIPOCARPHA.
94². l. *sphacelata*, Kunth, l. c. = triceps.
triceps, *Nees*, . . . 93.—Amphig. trop.

LIPOCHAETA.
94³. l. *hastata*, Kellogg, . . . = Verbesina remota.

LIQUIDAMBAR.
96². l. *peregrina*, . . . = Myrica asplenifolia.

QUIRITIA.
96². l. *lepidota*, [Nutt. in] Fras. Cat. (1813) n. 45; DC.

LISTERA.
98¹. a. *multinervia*, Peterm. in Flora, xxvii. (1844) 369 = ovata.

98². insere LITHOCARDIUM, Linn. Syst. ed. I (1735) = **Cordia**, Linn. (Borag.).

LITHOCARPUS.
98². a. *Benzoin*, Royle, Illustr. Bot. Himal. 457 = Styrax Benzoin.

LITSEA.
100². a. QUINQUEDULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

101³. lin. prima, l. **LIVISTONA**,

LOASA.
102¹. l. *aurantiaca*, Hort. ex Lilja, in Linnaea, xv. (1841) 263, etc.
contorta, Lam. . . . 579 = Blumenbachia contorta (in Addend.).
lateritia, Gill. . . . 275 = Blumenbachia lateritia.
102². punicea, *Phil.* . . . (1884)

LOBELIA.
102³. l. *adenophora*, Roem. & Schult. . . . = Lysipomia aretioides.
103². *cinerea*, Thunb. . . . = Lightfootia albicaulis.
104¹. *elliptica*, Willd. . . . 57 = Siphocampylus ellipticus.
104². *glandulosa*, Willd. . . . = Lysipomia aretioides.
104³. *limoselloides*, Willd. . . . = Lysipomia aretioides.
littoralis, R. Cunn. . . . 50 = Pratia angulata.
105¹. *nivea*, Willd. . . . 58 = Siphocampylus niveus.
105³. *reticulata*, Willd. . . . 58 = Siphocampylus reticulatus.
retrorsa, Willd. . . . 57 = Siphocampylus retrorsus.
Rhynchopetalum, *Hemsl.*
106². *subulata*, Benth. . . . 137 = Rhizocephalum subulatum.
syphilitica, *Linn.* . . . —Am. bor.

PAG.

LOLIUM.

108³. l. *čechicum*, Opiz, Oekon.
Coelorachis, Forst. ex Kunth, Enum. Pl. i. Suppl. i. 375 =

LOMATIA.

109². l. *chilensis*, C. Gay, . . . 310 = Orites myrtoidea.

LONCHOCARPUS.

110¹. l. Neuroscapha, *Benth.*
110². *Pterocarpus*, DC. . . . 260 = Phyllocarpus Pterocarpus.
a. *scandens*, Steud. Nom. ed. II. ii. 461, in syn., sphalm. = L. Nicou.

110². insere LONCHOPHYLLUM, Ehrh. Beitr. iv. (1789) 148 = **Cephalanthera**, Rich. (Orchid.).

LONICERA.

111³. l. obovata, *Royle, Illustr. Bot. Himal.* 236.—
112¹. *Symphoricarpos*, Linn. . . . = Symphoricarpos orbiculatus.

LOPHIOCARPUS.

113¹. l. *guyanensis*, Micheli . . . = S. guayanensis.

LOPHOPHYLLUM.

113². a. Peraphora, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 20.
l. bicristatum, *Griff.*

LORANTHUS.

116². l. *montanus*, Macfad. Fl. Jamaic. ii. 198 ; et ex

LOROGLOSSUM.

118¹. a. *longibracteatum*, Moris, ex Ardoino, Fl. Alpes-Marit. 351 = O. longibracteata.

LOTUS.

119¹. d. Ortholotus, Tourn. (*nunc* **Dorycnium**).
120¹. a. *peliorhynchus*, Hook. f. Bot. Mag. t. 6733 = Bertholetii.

LUDOLFIA.

122³. l. *falcata*, Nees, in Royle, Illustr. Bot. Himal. 417, nomen ; et ex

LUDWIGIA.

123¹. l. polycarpa, *Short & Peter, in Transylv. Journ. Med.* n. xxxii. (1836) 577 ; *et ex*

LUFFA.

123³. a. *tenera*, Royle, Illustr. Bot. Himal. 219 = acutangula ?

LUISIA.

123³. a. Trichorhiza, Lindl. ex Steud. Nom. ed. II. ii. 702 (1841).

LUMNITZERA.

124². l. Petaloma, Roxb. Hort. Beng. [90] (1814) ; Fl. Ind.
coccinea, *Wight & Arn. Prod.* 316, etc.

LUNDIA.

124³. l. *monachantha*, Schum. & Thonn. . . . = O. spinosa.

LUPINUS.

125¹. a. *affinis*, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 135 = microcarpus.
126². l. rupestris, *H. B. & K.* . . . 474.

LYCHNIS.

129¹. l. chalcedonica, *Linn.* . . . 436.—Rossia ; Sibir.
129². a. *Fortunei*, Hort. ex Vis. in Linnaea, xxiv. (1851) 181 = Silene Fortunei.
Lagrangei, *Coss. in Bull. Soc. Bot. Fr.* xx. (1873) 243.—Marocc.
l. magellanica, *Desr.*

LYCOPUS.

132². l. *laciniatus*, Rouy, in Le Natural. ii. (1882) (Matér. Fl. Portug.) 507 = europaeus.

LYGODISODEA.

133¹. l. *Lingua*, Boj. . . . = Lecontea Bojeriana.

LYSIMACHIA.

134². a. Nemorella, Ehrh. Beitr. iv. (1789) 147.

LYTHRUM.

136¹. l. *cashmerianum*, Royle, . . . 213

PAG.

MACROCENTRUM.

141². l. *mendocinum*, Phil. l. c. 43 = H. Gourlieana.

MACROCNEMUM.

141³. l. *exsertum*, Oerst. . . . 45 = Pogonopus exsertus.

142². insere MACROLOBIUM, Zipp. ex Miq. Fl. Ind. Bat. i. i. 80 (1855) = **Afzelia**, Sm. (Legumin.).

143¹. lin. prima, l. MACRORHYNCHUS.

143¹. a. MACROSCAPA, Kellogg, ex Curran, in Bull. Calif. Acad. i. III. (1885) 148 = **Strophostyron**, Torr. (Liliac.).
volubilis, Kellogg, l. c. = S. californicum.

MACROZAMIA.

144¹. a. *Macienzii*, Hort. ex Mast. in Gard. Chron. (1877) i. 665. f. 109 = Miquelii.

MAERUA.

144³. l. *pubescens*, C. Muell. in Walp. Ann. vii. 188 = nervosa.
scanlens, C. Muell. (l. c.) = nervosa.

MAGNOLIA.

146¹. a. *Sieboldii*, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 257 = Talauma stellata.

MAHERNIA.

146². l. *gracilis*, Harv. . . . 215 = linnaeoides.
a. *linnaeoides*, Harv. in Harv. & Sond. Fl. Cap. i. 25 = rotundata.
146³. l. rotundata, *E. Mey.* . . . 192.—Afr. austr.

MAIANTHEMUM.

147¹. l. *Bifolium*, DC. Fl. Fr. iii. 177 ; [DC.]
a. *smilacinum*, Montand. Guide Bot. 310 = M. Convallaria.

MAIHTA.

147¹. l. *heterophylla*, DC. . . . 166 = Calophysa heterophylla.

MAIACHRA.

148¹. a. *diversifolia*, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 134 ; xii. (1845) 79 = Pavonia diversifolia.

MAIESHERBIA.

150¹. l. *paniculata*, D. Don, . . . 323

MAILOTUS.

150². l. Adelta, *Muell. Arg.*
chinensis, Muell. Arg. in DC. Prod. xv. II. 965, in syn. =
japonicus, *Muell. Arg.* . . .—Japon.
150³. *micranthus*, Muell. Arg.
tiliaefolius, *Muell. Arg.*

MAIUS.

152². l. *Torringo*, Siebold, ex De Vriese, Tuinb. Fl. iii. (1856) 368 ; Carr. . . . = P. Sieboldii.

MALVA.

153³. l. *Monroana*, Otto & Dietr. . . . = Sphaeralcea Munroana.
154¹. *Munroana*, Dougl. . . . = Sphaeralcea Munroana.

MAMMILLARIA.

157³. l. Goodrichii, *Scheer*,
a. *Goodridgii*, Scheer, in Seem. Bot. Voy. Herald, 286, sphalm. = praec.
159¹. l. *pectinata*, Engelm. . . . 10 = radians.
haphidacantha, Hemsl. Biol. Centr. Am. Bot. ii. 520, sphalm. = raphidacantha.
159². rhodantha, *Link & Otto, Ic. Pl. Rar.*
160¹. retrancistra, Engelm. . . . partim.—Am. bor. occ. ; Mexic.
retrancistra, Engelm. l. c. partim =
160². *tuberculosa*, Engelm. (non *tuberculata*).

MANDEVILLA.

160³. a. Wrightiana, *Benth. & Hook. f. Gen.* ii. 727.—Cuba.

MANISURIS.

162³. a. Rytilix, Rafin. in Ser. Bull. Bot. i. (1830) 219.

MAPPA.

164³. l. *peltata*, Wight,

PAG.

MAPPIA.

164³. l. *montana*, Miers, l. c. 398 (Apodytis sp.).—Java.

MARAH.

165¹. l. *muricata*, Kellogg,

MARANTA.

165². a. *discolor*, Welw. Synopse, 43 (al. gen.).—Afr. trop.
Makoyana, *E. Morr. ex Flor. Mag.* N. S. (1873) t. 53.—Am. trop.
165³. l. *Moritziana*, Koern. . . . 66 = Myrosma cannaefolium.
a. *nobilis*, Royle, Illustr. Bot. Himal. 356, nomen (Quid ?).—Ind. occ.
166¹. l. *olivaris*, Hort. ex Flor. Mag. N. S. (1873) sub t. 53 = Makoyana.
166². *unilateralis*, D. Dietr. Syn. Pl. i. 6 = Myrosma unilateralis.

168¹. l. MARIANA, Hill, . . . = **Silybum**, Vaill. (1763).

MARIANTHUS.

168¹. a. Cyathomiscus, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 562, in obs.
l. Oncosporum, Putterl. in Endl. Nov. Stirp. Dec. 9 (1839).

169¹. l. **MARISCUS**, Ehrh. Beitr. iv. (1789) 146 ; Gaertn.
a. Opetiola, Gaertn. Fruct. i. 14. t. 2. (1788).
Torulinium, Desv. in Ham. Prod. Pl. Ind. Occ. 15 (1825).

MARLEA.

170¹. a. Stelanthes, Stokes, Bot. Mat. Med. ii. 339 (1812), nomen prius.

MARSDENIA.

171². l. *longiflora*, A. Rich. . . . 100 = Stephanotis longiflora.
171³. tinctoria, *R. Br.* . . . 30.—As. trop. (non Afr.).
vinciflora, Griseb. . . . 170 = Stephanotis vincaeflora.

MARTYNIA.

172³. l. Proboscidea, *Glox.*

MASSONIA.

174³. a. *nivea*, Burch. ex Salisb. Gen. Pl. Fragm. 17 = jasminiflora ?

MATRICARIA.

176¹. a. Otospermum, Willk. in Bot. Zeit. xxii. (1864) 251.

MAURIA.

178². a. *bijuga*, March. Rév. Anacard. 162, sphalm. = subbijuga.

MAXILLARIA.

178³. d. Onkeripus, et Pentulops, Rafin. (*nunc* **Xylobium**).
179¹. l. crocea, *Lindl. l. c.* t. 1799.
a. cyanocheile, *Hoffmgg. Preisverz.* ed. X. 80.—*Cf. Linnaea*, xvi. (1842) *Litt.* 272.—Bras.
180¹. *nasalis*, Reichb. f. in Saund. Refug. Bot. ii. t. 103, sphalm. = seq.
pallidiflora, Reichb. f. in Rodrig. Orch. Nov. ii. 201 = liliacea.
180³. l. *Warreana*, Lodd. Bot. Cab. t. 1884 = Warrea tricolor.

MAXIMILIANA.

180³. l. *insignis*, Mart. . . . 94 = Scheelia insignis.

182¹. post MECONELLAM insere MECONIA, Hook. f. & Thoms. Fl. Ind. i. 275, in obs. (1855), sphalm. = praec.

MECONOPSIS.

182¹. l. horridula, *Hook. f. & Thoms.* . . .—Reg. Himal. ; China.
napaulensis, *DC.*
a. *nipalensis*, Hook. f. & Thoms. Fl. Ind. i. 253 = praec.
l. *racemosa*, Maxim. . . . = horridula.
robusta, Hook. f. & Thoms. . . . 253 = napaulensis.
simplicifolia, *Walp. Rep.* i. 110.—Reg. Himal.
a. *Wollastoni*, Regel, Gartenfl. (1876) 291, lapsu = napaulensis.

PAG.
182². l. MEDICA, Cothen. . . . = **Tourretia**, Fouger.

MEDICAGO.
182³. a. TRIFILLIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 383.
TRIPHYLLUM, Medic. Phil. Bot. i. 202 (1789).

185³. a. MEGAPHYLLUM, Spruce, ex Baill. Hist. Pl. vii. 457, in nota (1880) = **Pentagonia**, Benth. (Rubiac.).

MELALEUCA.
187⁵. l. lateriflora, *Benth.*

189². l. MELANDRIUM, . . . ed. II. ii. 37, 274 (1812).
Flos-cuculi, . . . ed. II. ii. 275 =
189³. *pratense*, . . . ed. II. ii. 274 =
sylvestre, . . . ed. II. ii. 274 =

MELANOCENCHRIS.
190¹. a. GRACILEA, Koen. ex Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218 nomen prius,
ROYLEA, Nees, ex Steud. Nom. ed. II. ii. 475 (1841).
l *Jacquemontii*, Jaub. & Spach, . . . = Royleana.
Perrottetii, Jaub. & Spach, . . . = Rothiana.

MELANOSINAPIS.
190². l. *communis*, Spenn. l. c. 945 = B. sinapioides.

MELANTHESOPSIS.
190³. l. *fruticosa*, Muell. Arg. . . . 437 = Andrachne fruticosa ?

MELANTHIUM.
191². a. *hyacinthoides*, Royle, Illustr. Bot. Himal. 384 = Scilla indica.

MELASMA.
191². l. hispidum, *Benth.*

MELASTOMA.
191². d. OCTONUM, Rafin. (*nunc* **Heterotrichum**).
192². l. *chulesis*, Buch.-Ham. . . . = Osbeckia nepalensis.

MELHANIA.
196³. l. *laurifolia*, Boj. in Rapp. Soc. Nat. Maurice, xii. (1841) 48 ; et in

MELIA.
197¹. l. *Bambolo*, C. DC. in DC. Monog. i. 767 err. typ. = seq.
Bombolo, Welw.

MELICA.
197². l. *arundinacea*, Raeusch. . . . = Molinia caerulea.
197³. *Geyeri*, Boland. . . . = bromoides.
Geyeri, Munro, ex Boland. l. c. = bulbosa.

MELICOCCA.
198³. a. *javanica*, Hassk. Hort. Bog. Desc. i. 138 = Otophora javanica.

200². insere MELIOTUS, Steud. Nom. ed. II. ii. 704, in syn., err. typ. = **Melilotus**, Tourn. (Legumin.).

MELOCACTUS.
201². a. approximatus, *Suring. in Versl. en Med. Kon. Akad. Wetensch.* III. ii. (1885) 187.—Ind. occ.
cornutus, *Suring.* (*l. c.*) 191.—Ind. occ.
Evertizianus, *Suring.* (*l. c.*) 190.—Ind. occ.
ferox, Suring. (l. c.) 185 = rubellus.
hexacanthus, Suring. (l. c.) = rubellus.
201³. intermedius, *Suring.* (*l. c.*) 192.—Ind. occ.
Koolwijkianus, *Suring.* (*l. c.*) 184.—Ind. occ.
parvispinus, *Suring.* (*l. c.*) 183.—Ind. occ.
patens, *Suring.* (*l. c.*) 190.—Ind. occ.
pusillus, *Suring.* (*l. c.*) 192.—Ind. occ.
rectiusculus, *Suring.* (*l. c.*) 187.—Ind. occ.
reversus, *Suring.* (*l. c.*).—Ind. occ.
rubellus, *Suring.* (*l. c.*) 184.—Ind. occ.
spatanginus, *Suring.* (*l. c.*) 193.—Ind. occ.
stramineus, *Suring.* (*l. c.*) 185.—Ind. occ.
202¹. *trichacanthus*, Suring. (l. c.) 186 = stramineus ?

MELOCHIA.
202². a. corymbosa, *Hemsl. Biol. Centr. Am. Bot.* i. 130. —Mexic.

MENTHA.
210². l. Welwitschii, *Rouy, in Le Natural.* ii. (1882) (*Matér. Rév. Fl. Portug.*) 47.—Europ.

PAG.
MEOSCHIUM.
211². l. *Royleanum*, Nees, in Royle, Illustr. Bot. Himal. 417 =

212³. l. MERIDIANA, [Linn. ex] Linn. etc.

MERTENSIA.
213². l. *virginica*, Link, . . . = pulmonarioides

MESPILUS.
218³. a. *media*, Bechst. Forstbot. 150, 3[illegible] = Crataegus Oxyacantha.

METACHILUM.
220¹. l. *cyathiferum*, Lindl. l. c.
cyathiforme, Steud. Nom. ed. II. i. 136, sphalm. = praec.

METROSIDEROS.
221³. l. *leptopetala*, F. Muell. = Syncarpia leptopetala.
222². *ternifolia*, F. Muell. = Lysicarpus ternifolius.

222². insere METROXILON, Welw. Apont. 584 [1859] = seq.

METROXYLON.
222². l. Pigafettia, in synon. passim.—Cf. *latum, filare, microcarpum, microspermum.*
a. *textile*, Welw. Apont. 584 = Raphia Welwitschii.

l. METROXYLON, Spreng. . . . = Raphia, Beauv, (Palm.).

MICHELIA.
223³. a. PHELIMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

223³. insere MICHELIA, Linn. Syst. ed. I. (1735) = **Barringtonia**, Forst. (1776) (Miltac.).

MICONIA.
223³. a. ACINODENDRUM, Linn. Gen. ed. I. 329 (1737).
224¹. LIEUTAUTIA, Buc'hoz, Pl. Nouv. Découv. t. 7 (1779), nomen prius.
224³. l. *corallina*, Spring,

MICRANTHEMUM.
228¹. a. adenandrum, *Wright, in Sauv. Fl. Cub.* 101.—Cuba.
multiflorum, *Wright*, (*l. c.*).—Cuba.
punctatum, *Wright*, (*l. c.*).—Cuba.
l. rotundatum, *Wright, ex Griseb.*
a. trisetosum, *Wright, in Sauv. Fl. Cub.* 101.—Cuba.

MICRANTHUS.
228¹. l. *oppositifolius*, Wendl.

MICROCARPAEA.
228². a. *chamaedrifolia*, Steud. Nom. ed. II. ii. 246, in syn. = seq.

228³. l. MICROCHAETA, Reichb.

MICHROCHAETE.
228³. l. *corymbosa*, Benth. l. c. 196 = S. Microchaete.

MICROCHLAENA.
228³. l. *quinqueloculars*, Wight & Arn.

MICHROCHLOA.
228³. l. *elongata*, R. Br. . . . n. 3807 = setacea

MICROGLOSSA.
229². l. *altissima*, DC. . . . = Psiadia altissima
hispida, DC. . . . = Psiadia hispida.

MICROLICIA.
230³. a. *thymifolia*, Steud. Nom. ed. II. ii. 447, in syn., sphalm. = serpyllifolia.

MICROLOMA.
230³. a. *hastatum*, Decne. in DC. Prod. viii. 499 = sagittatum.
231¹. l. *pyrotechnicum*, Spreng. . . . = Lotadenia Spartium.

MICROPUS.
232³. d. ANCISTROCARPHUS, A. Gray, (*nunc* **Stylocline**).

PAG.
MICROSTYLIS.
233³. l. *bella*, Reichb. f. . . . (1885) 187. t. 581

MILIUM.
236³. l. *capillare*, Roth, . . . = Panicum subeglume.
237². a. *nepalense*, Nees, in Royle, Illustr. Bot. Himal. 417 = effusum ?
237². l. *sanguinale*, Roxb. . . . Paspalum pedicellare.

MILLA.
237³. d. TULOPHOS, Rafin. (*nunc* **Brodiaea**).

MILLETTIA.
238². l. *speciosa*, Welw. . . . 23 = Lonchocarpus sericeus.

MIMULUS.
245³. l. *barbatus*, Greene, . . . 9 = montioides.
a. *Langsdorfii*, Donn, ex Sims, Bot. Mag. sub t. 1501 = guttatus.
246¹. l. *punctatus*, Loisel. Herb. Amat. t. 152 ; Miers,

MIMUSOPS.
246². a. RADIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 3.

247¹. insere MINTHOSTACHYS, Bunge, Lab. Pers. 13 (1873) = **Bystropogon**, L'Hérit. (Labiat.).
mollis, Griseb. in Goett. Abh. xix. (1874) 235 = B. mollis.

MINURIA.
247¹. a. Kippistiana, *F. Muell. Pl. Vict.* i. sub t. 35.—Austral.
l. *suaedifolia*, F. Muell. . . . 499 = Kippistiana.

247¹. l. MINYRANTHES, Turcz. . . . = **Siegesbeckia**, Linn.

MIRASOLIA.
247³. a. *scabra*, Benth. & Hook. f. l. c. 375 = Tithonia scabra.

247³. a. MIRKOOA, Wight, Ic. i. t. 217, in textu (1840) = **Ammannia**, Houst. (Lythrar.).
rotundifolia, Wight, l. c. = A. rotundifolia.

MISCHOPHLOEUS.
248¹. l. paniculatus, *Scheff.*

MITCHELLA.
248². a. PERDICESCA, Provand. Fl. Canad. 291 (1862).

MITRANTHES.
249². l. *Riedeliana*, Berg,

MITREOLA.
250¹. a. *lanceolata*, Torr. ex Croom, Cat. Pl. New Bern, 45 = petiolata.

MITREPHORA.
250¹. l. *excelsa*, Hook. f. & Thoms. . . . 114 = Uvaria excelsa.

MNASIUM, Schreb.
250². l. *sphaerocephalum*, Rudge, . . . = Thurnia sphaerocephala.
unilaterale, Rudge, . . . = Spathanthus unilateralis.

250³. l. MNIARUM, Forst. Char. Gen. l. t. 1.

MOEHRINGIA.
251³. a. *sedifolia*, Freyn, in Oestr. Bot. Zeitschr. xxvi. (1876) 227 = *papulosa*.
l. *Tommasinii*, Marches.

MOLINAEA.
252³. a. *arborea*, J. F. Gmel. Syst. 611 = C. alternifolia, laevis.

MOLLINEDIA.
253². a. PARACELSIA, Mart. ex Tul. in Mart. Fl. Bras. iv. 1. 314, 316 (1857).

MOLLUGO.
254¹. l. Trigastrotheca, *F. Muell.*

MONANTHES.
256¹. l. polyphylla, *Haw.*

256¹. insere MONANTHIUM, Ehrh. Beitr. iv. (1789) 148 = **Moneses**, Salisb. (Pyrolac.).

PAG.

MONESES.

257[1]. a. MONANTHIUM, Ehrh. Beitr. iv. (1789) 148.

MONIZIA.

257[1]. l. *edulis*, Lowe, l. c. = T. Moniza?

MONNIERA.

257[2]. a. *longifolia*, Steud. Nom. ed. II. ii. 373, sphalm. = Monnina longifolia.

MONNIERIA.

257[2]. a. *trifoliata*, Baill. Adansonia, xi. (1875) 276 = praec.

MONOCHAETUM.

258[2]. a. ROEZLIA, Regel, in Act. Hort. Petrop. i. (1871) 98.

MONODORA.

259[1]. l. angolensis, *Welw. in Journ. Linn. Soc.* iii. (1859) 151, 154; *Apont.*

MOQUILEA.

262[2]. l. floribunda, *Hook. f. in Mart. Fl. Bras.* xiv. II. 21.—Bras.
pendula, *Hook. f.*
sclerophylla, *Hook. f.*
262[3]. turbinata, *Hook. f.*

MORAEA.

263[1]. l. *chinensis*, Thunb, . . . = Belamcanda puncta a.

MORICANDIA.

264[1]. a. *cinerea*, Coss. ex Tourn. Thèse Crucif. 143 = Ammosperma cinerea.

MORINDA.

264[3]. a. *ternifolia*, Royle, Illustr. Bot. Himal. 238, nomen (Quid?).—Ind. or.

MORUS.

266[1]. a. *cashmeriana*, Royle, Illustr. Bot. Himal. 337 = indica?
dulcis, Royle, (l. c.) = alba?
266[2]. *parvifolia*, Royle, (l. c.) = indica?

MUCUNA.

268[2]. a. dichroa, *Hoffmgg. Preisverz.* ed. X. 80.—*Cf. Linnaea*, xvi. (1842) *Litt.* 272.—Hab.?

MUEHLENBECKIA.

268[3]. l. *Benthamii*, Endl. Gen. Suppl. iv. II. 51
quadrangulata, Endl. (l. c.)
volcanica, Endl. (l. c.)

MUEHLENBERGIA.

268[3]. a. SERICROSTIS, Rafin. Neogenyt. 4 (1825).
269[1]. l. *filipes*, M. A. Curt.
a. *maritima*, Ind. Kew. ii. p. 60[1]. in syn., lapsu = Polypogon maritimus.
269[2]. l. setifolia, [*Vasey, in*] *Coult.*
subalpina, Vasey, . . . 40 = gracilis.

MULGEDIUM.

270[1]. l. *leucophaeum*, DC. l. c. 250 = L. alpina.

MURALTIA.

271[1]. a. *ericoides*, Steud. Nom. ed. II. ii. 371, in syn. sphalm. = Burmanni.
l. laxa, *DC. etc.*

MURICARIA.

271[3]. l. Cordylocarpus, *C. Muell.*

271[3]. l. MUROCOA, . . . **Maripa**, Sw. (Convolv.).

271[3]. l. MURUCOA, J. F. Gmel. . . . **Maripa**, Sw. (Convolv.).

MUSA.

272[1]. a. Charlioi, *E. Hill, Rep. Brisbane Bot. Gard.* (1874) 7.—Hab.?
Rhinozerotis, Kurz, in Journ. Agr. Hort. Soc. Ind. N. S. v. (1878) 64 = nana?

MUSCARI.

272[2]. d. *botryoides*, Steud.
272[3]. a. *inodorum*, Montand. Guide Bot. 313 = botryoides.
odoratum, Montand (l. c.) = racemosum.

PAG.

275[2]. l. MYCETIA, Reinw. . . . = **Adenosacme**, Wall.
a. *javanica*, Hook. f. Fl. Brit. Ind. iii. 95, sphalm. = praec.

MYRICA.

281[2]. d. PIMICARIA, Rafin. (*nunc* **Ximenia**).
a. *elegans*, Royle, Illustr. Bot. Himal. 459 (Index), sphalm. = Myricaria elegans.
281[3]. l. *obovata*, [Champ.] ex C. DC. l. c. 150 = inodora.

MYRISTICA.

282[3]. a. *angolensis*, Welw. Synopse, 51. (Quid?).—Afr. trop.

MYROMA.

284[3]. l. unilaterale, *Benth. & Hook. f.*

MYROXYLON.

284[3]. l. Tolufera, *H. B. & K.*
a. *toluiferum*, G. Don, Gen. Syst. ii. 109, in syn. = prec.

286[3]. post MYRTOBIUM insere MYRTOLOBIUM, Calon, in Mém. Soc. Hainaut, Sér. III. v. (190) 324, sphalm. = praec.

NAEGELIA.

291[2]. a. *achimenoides* ×, Bárlik, in Wien. Garten-Zeit. x(1885) 239. t. 2.

NAMA.

292[3]. l. *tetrandrum*, Pav. ex Choisy, in Ann. Sc. Nat. Sr. I. xxx. (1833) 243 =

NARCISSUS.

293[2]. a. *bitnaceus*, Salisb. Gen. Pl. Fragm. 100 = incomparabilis?

NASSELLA.

295[1]. a. *aenulans*, Phil. f. Cat. Pl. Vasc. Chil. 328.—Chili.

NASTURTIUM.

296[1]. l. *acule*, Welw. Apont. 589 = humifusum.

NECANDRA.

299[2]. a. *deandra*, Roxb. Fl. Ind. i. 425 = Linostoma decandrum.
300[1]. l. *stminea*, Griseb. . . . 282 (= *Oreodaphne fragrans*).—Ind. occ.

302[3]. insere NEMOLAPATHUM, Ehrh. Beitr. iv. (1789) 149 = **Polygonum**, Tourn.

303[1]. inser NEMORELLA, Ehrh. Beitr. iv. (1789) 147 = **Lysimachia**, Tourn. (Primul.).

NEOTTIA.

303[3]. l. *phylla*, Hook.
304[1]. a. *corata*, Blume, Bijdr. 409 = Spiranthes colorata, *Hassk.*
grandis, Blume, (l. c.) 407 = Spiranthes grandis.
longifolia, Blume, (l. c.) 406 = Spiranthes longifolia.
parvifolia, Blume, (l. c.) 408 = Spiranthes parviflora.
304[2]. *reticulata*, Blume, (l. c.) 409.—Java.
l. *vaginata*, H. B. & K. . . . 331 = Spiranthes vaginata.

NEPETA.

305[1]. l. CHAMAECLEMA, Boerh. ex Boehm. in Ludw. Def. Gen. Pl. 115; Moench,
305[3]. *asyangula*, *Lenné & C. Koch*, . . . 12.—Reg. Himal.
Jwerardi, *S. Moore*,

NEPHELIUM.

307[2]. l. *cuminatum*, Hook. f. . . . 164 = Pometia acuminata.
307[3]. Maingayi, *Hiern*,
a. *muricatum*, Herb. Kew. ex Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 93 = Xerospermum mucronatum.

309[3]. a. NESOEA, Wight, Ic., in textu sub t. 217 (1840) = **Ammannia**, Houst. (Lythrar.).
octandra, Wight, l. c. = A. octandra.

NICANDRA.

311[2]. a. PHYSALODES, Boehm. in Ludw. Def. Gen. Pl. 41 (1760).

PAG.

NICOTIANA.

312[1]. l. diversifolia, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 384; *et*

NIDORELLA.

313[1]. l. *altissima*, Benth. & Hook. f. . . . = Psiadia altissima.
hispida, Benth. & Hook. f. . . . = Psiadia hispida.

NIDULARIUM.

313[1]. l. *caeruleum*, Lem.

NIVENIA.

315[3]. l. PARANOMUS, Salisb. . . . (1807).
Lagopus, *R. Br.*

NOLANA.

316[2]. l. triquetra, *C. Koch & Bouché*, . . . *App.* 12.

NONNEA.

317[1]. a. PHANERANTHERA, DC. ex Meissn. Gen. Comm. 189 (1840).

NOTHITES.

318[2]. l. *breviflora*, Cass. . . . 165 = S. aristata?
latifolia, Cass. l. c. 163 = S. melissaefolia.
melissaefolia, DC. . . . 186 = S. melissaefolia.
petiolata, Cass. . . . 166 = S. petiolata.

NOTHOSCORDUM.

319[1]. a. PERILOBA, Rafin. Fl. Tellur. iv. 87 (1836), nomen prius.

NYCTAGO.

320[3]. l. *angulata*, DC. . . . (= *Quamoclidion angulatum*).—Mexic.

NYMPHAEA.

321[2]. l. *Fenzliana*, Lehm. . . . = Rudgeana.
321[3]. *Kosteletzkyi*, Palliardi,
322[1]. stellata, *Willd.* . . . As. et Afr. trop.; Austral.

NYMPHOIDES.

322[2]. l. *europaea*, Fisch. . . . = L. Nymphoides.
orbiculata, Gilib. . . . = L. Nymphoides.

322[3]. l. OAKESIA, S. Wats. . . . 269 = **Uvularia**, Linn. (Liliac.).
puberula, S. Wats. l. c. = U. puberula.
sessilifolia, S. Wats. l. c. = U. sessilifolia.

OCHNA.

324[1]. l. *nitida*, Sw. . . . = Gomphia nitida (in Addend.).

OCHROSIA.

324[3]. l. *sandwicensis*, A. DC. . . . 357 = Rauwolfia sandwicensis.
a. sandwicensis, *A. Gray, in Proc. Am. Acad.* v. (1862) 333.—Ins. Sandvic.

OCOTEA.

326[2]. l. Bofo, *H. B. & K.*

OCTADESMIA.

327[1]. l. *monophylla*, Benth. & Hook. f. l. c. 526 = Laelia monophylla.

ODINA.

328[1]. a. *caffra*, Hook. ex March. Rév. Anacard. 162 = Harpephyllum caffrum.

ODONTADENIA.

328[2]. a. PERICTENIA, Miers, Apocyn. S. Am. 182 (1878).

ODONTOGLOSSUM.

330[3]. l. *pygmaeum*, Lindl. . . . = Leochilus pygmaeus.

OENANTHE.

331[3]. l. *amoena*, Sweet, Hort. Brit. ed. II. 247, sphalm. = Oenothera amoena.

OENOTHERA.

333[1]. l. albicaulis, [*Nutt. in*] *Fras. Cat.* (1813) n. 54, nomen; *et ex*
caespitosa, [*Nutt. in*] *Fras. Cat.* (1813) n. 53; *et ex*
campylocalyx, *C. Koch & Bouché* . . . (1855) *App.* 17.
333[3]. *epilobioides*, Nutt. . . . 511 = strigulosa.
erosa, *Lehm. Sem. Hort. Hamb.* (1824) 20; *et in*
334[1]. *macrocarpa*, [Nutt. in] Fras. Cat. (1813) n. 56; Pursh,

PAG.

OGIERA.

335². a. *portoricensis*, Steud. Nom. ed. II. ii. 582, sphalm. = E. ovata.

OLEARIA.

338¹. l. Allomii, *T. Kirk*, . . . i. ed. II. (1871) 92 ;

OLINIA.

339². a. PLECTRONIA, Buching. ex Krauss, in Flora, xxvii. (1844) 348.

ONCIDIUM.

341³. l. *acrobotryum*, Klotzsch, . . . = Harrisonianum.
343¹. festatum, *Reichb. f. & Warsz.*
343². *hemichrysum*, Reichb. f. & Warsz. in Bonplandia, ii. (1854) 100 =
344³. *pantherinum*, Hoffmgg. Preisverz. ed. X. 80.—Cf. Linnaea, xvi. (1842) Litt. 272 =

ONOBRYCHIS.

346³. a. *alpicola*, Arv. Ess. Pl. Dauph. 25 (Quid ?).—Gallia.

ONOSERIS, Willd. Sp. Pl. iii. 1702 (1804) ; DC.

350². a. *nepalensis*, Royle, Illustr. Bot. Himal. 277, nomen (Quid ?).—Nepal.

ONOSMA

351¹. l. *saxatile*, Lehm. . . . 371 (= *Stenosolenium saxatile*).—Sibir.

OPHELIA.

352². l. *angustifolia*, D. Don, ex Royle, Illustr. Bot. Himal. 277 ; et ex G. Don,

OPHIORRHIZA.

353¹. l. *fruticulosa*, Nadeaud, . . . 53 = subumbellata.
353². *torrentium*, Nadeaud, . . . 54 = Solandri.

OPHRYOSPORUS.

354¹. l. Pachychaeta, *Baker*,

OPHRYS.

354². l. atrata, *Lindl.* . . . t. 1087.

OPILIA.

355². a. TETRANOSIA, Rich. ex M. Roem. Syn. Hesper. 23 (1846).

OPLISMENUS.

355³. l. PANICULUM, Ard.

ORBIGNYA.

359². a. Sagotii, *Trail, ex im Thurn, in Timehri*, iii. (1884) 276.—Guian.

359². l. ORCHIDIUM, [Sw.] Summa Veg. Scand. 32 (1814) ; et in
l. *arcticum*, [Sw.] (l. c.).

ORCHIS.

359³. a. *angustata*, Arv. Ess. Pl. Dauph. 62 (Quid ?).—Gallia.
biloba, Arv. (l. c.) 61 = laxiflora.
360¹. Burmanniana, *Sw. in Weber & Mohr, Arch.* i. (1804) 55.—Afr. austr.
361². l. *macronectar*, Vell. . . . 45 = Habenaria Sartor ?
a. *nemorosa*, Montand. Guide Bot. 294 = maculata.
362¹. *rosea*, Arv. Ess. Pl. Dauph. 63 (Quid ?).—Gallia.

363². l. OREOC DE, Miq.

OREODAPHNE.

364². l. *Teleiandra*, Meissn.

367¹. insere ORLEANIA, Commel. ex Boehm. in Ludw. Def. Gen. Pl. 380 (1760) = **Bixa**, Linn.

ORMOCARPUM.

367². l. coccineum, *G. Don*,

ORNITHOGALUM.

368². d. SIMIRA, Rafin. (*nunc* **Scilla**).
370¹. a. *ochroleucum*, Montand. Guide Bot. 315 = pyrenaicum.

371². l. ORNITHORHYNCHIUM, Roehl. . . . .(1812).

PAG.

OROBANCHE.

373². a. *Lactucae*, Arv. Ess. Pl. Dauph. 54 (Quid ?).—Gallia.

OROBUS.

375². d. *flaccidus*, Kit.

ORTHOPOGON.

377³. a. *Junghuhnii*, Nees, ex Steud. Syn. Gram. 45 ; et ex Miq. Fl. Ind. Bot. iii. 444 = Oplismenus compositus.

ORTHOSPERMUM.

378². a. *unctuosum*, Montand. Guide Bot. 362 = C. Bonus-Henricus.

ORYZA.

378³. a. SCLEROPHYLLUM, Griff. Notul. iii. 8 (1851).

ORYZOPSIS.

379¹. l. *canadensis*, Torr. . . . 583 = parviflora.
379². parviflora, *Nutt. l. c.* 126.—Am. bor.

OSBECKIA.

379³. a. *pendulifolia*, Steud. Nom. ed. II. ii. 447, in syn. = Aciotis pendulifolia.

OSMOXYLON.

381¹. a. borneense, *Seem. Rev. Heder.* 104.—Borneo.
Cumingii, *Seem.* (*l. c.*).—Ins. Philip.
pulcherrimum, *Vidal, Sin. Fam. Leñ. Atlas*, p. xxviii.—Ins. Philipp.

OSYRIS.

382³. l. *nepalensis*, D. Don, ex Royle, Illustr. Bot. Himal. 322 ; Griff.

OTHONNA.

383³. l. *nodulosa*, Sch. Bip. l. c. 769 = perfoliata.

385². l. OURAGOGA, Linn. . . . = URAGOGA, Linn. = **Psychotria**, Linn. (Rubiac.).

OURATEA.

385². l. *nitida*, Engl. l. c. 310 = G. nitida (in Addend.).
nitida, Engl. l. c. 314, in adnot. = G. guyannensis.

OXALIS.

387². l. Cummingi, *Herb.* . . . t. 1545.—Chil.
388³. leucotricha, *Turcz.* . . . Afr. austr.
389¹. a. microphylla, *Phil. in Linnaea*, xxxiii. (1864–65) 35 (sp. diversa).—Chili.
390,². l. *rubella*, Jacq.
390³. *sulphurea*, Jacq. . . . 63 = pulchella.

OXYGONUM.

392². a. DIPLOPYRAMIS, Welw. Apont. 591 [1859].

393². l. **OXYMITRA** ante OXYMITUM inserenda.

OXYPETALUM.

393³. l. *integrifolium*, Fourn. . . . 372, sphalm. = integrilobum.
integrilobum, *Fourn.* . . . 279

OXYSTELMA.

394². a. *aegyptiacum*, Decne. in DC. Prod. viii. 504, sphalm. = Alpini.

OXYTROPIS.

395¹. a. SPIESIA, Neck. Elem. iii. 13 (1790), nomen prius.
amethystea, Arv. Ess. Pl. Dauph. 24 (Quid ?).—Gallia.
amethystina, Arv. Monog. Pilos. 49 = praec. ?
396¹. l. Middendorffii, *Trautv. in Middend. Sibir. Reise* (*Fl. Taimyr.* 49. t. 7).—Sibir.
396³. *Trautvetteri*, Meinsh. . . . = Middendorffii.

PACHYCARPUS.

398¹. l. *gomphocarpoides*, E. Mey. l. c. 213 (= *Xysmalobium gomphicarpoides*).—Afr. austr.

PACHYGONE.

398². l. *pubescens*, Benth. . . . 58 = Trichocalyx pubescens.

PAEDERIA.

400³. l. *Lingun*, Boj. ex Sweet, Hort. Brit. ed. I. 487 =

PAG.

400³. l. **PAEONIA**
albiflos, Sabine, . . . = corallina.

PALICOUREA.

404¹. a. *Gudara*, Steud. Nom. ed. II. ii. 787, in syn., sphalm, (= *Pacuria Gudara*) = Willughbeia edulis.
l. quadrifolia, *DC. Prod.* iv. 526.—Guian.

PALIURUS.

404². l. *aculeatus*, Lam. . . . 611 = P. Spina-Christi.
australis, Gaertn. . . . 43 = P. Spina-Christi.
clypeiformis, Medic. . . . = P. Spina-Christi.
404³. *solitarius*, Stokes, . . . P. Spina-Christi.
Spina-Christi, *Mill.* . . . VIII.—Europ. austr. ; As. or.
vulgaris, Steud. . . . = P. Spina-Christi.

PANAX.

406¹. l. *micranthum*, Wall. Cat. n. 4938 = Leschenaultii.
a. *odora*, Royle, Illustr. Bot. Himal. 13, nomen (Quid ?).—Ind. or.

PANCRATIUM.

406³. d. TROXISTEMON, Rafin. (*nunc* **Hymenocallis**).

PANDANUS.

407³. a. SOULEYETIA, Gaudich. Bot. Voy. Bonite, t. 19 (1844–49).
l. SUSSEA, Gaudich. (l. c.) tt. 24, 25, 38 (1844–57 ?).

PANICUM.

409³. d. PSEUDORAPHIS, Griff. (*nunc* **Setaria**).
410². a. angustatum, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 157, nomen.—Ind. or.
l. *barbinode*, Trin. . . . = muticum.
411². l. *bidentulum*, Steud.
411¹. *capillaceum*, Lam. . . . = brevifolium, *Linn.*
412¹. a. *conjugatum*, Dalz. & Gibs. Bomb. Fl. 291 = cimicinum.
412². l. decompositum, *R. Br. Prod.* 191.—Austral.
412³. a. *dentispica*, Kunth, Rév. Gram. i. sub t. 39, err. typ. = *densispica*.
412³. l. *distichum*, Gaud. ex
413¹. *elegans*, Wight & Arn. ex Steud. l. c. = Isachne pulchella.
equinum, Salzm. . . . = muticum.
413². a. *floridum*, Royle, Illustr. Bot. Himal. 420 = flavidum.
413³. l. *fluitans*, Retz. . . . 8 = punctatum.
fugax, *Koen. in Naturf.* xxiii. (1788) 209 (*ex Roem. & Schult. Syst.* ii. 467) ;
Gardneri, Thw. . . . = Isachne Gardneri.
414². *hirtum*, Nees, . . . = brevifolium, *Linn.*
414³. *insulicola*, Steud. . . . = auritum.
a. *intermedium*, Griff. Notul. iii. 29 (Chamaeraphidis sp.).—Ind. or.
415². l. *lanuginosum*, J. & C. Presl, . . . = mollicomum.
416¹. molle, *Sw.* . . .—Am. trop. (dele As. trop. inquil.).
mollicomum, *Kunth*,
416². *muricatum*, Retz. . . . 18 (Quid ?).—Ind. or.
416³. numidianum, *Lam.* . . .—Afr. bor. ; Syria ; As. trop. ; Am. trop.
417¹. d. *oryzoides*, Nees,
a. *panis*, Jess. Deutschl. Graes. 240 = Setaria italica.
l. *paractaenoides*, Trin. . . . 219 = flavescens.
paspaloides, *Pers.* . . . (paspalodes).
a. *patens*, Roxb. Fl. Ind. i. 305 = Isachne miliacea.
417³. d. *plicatum*, Roxb.
l. *polystachion*, Linn. . . . = setosum.
418¹. punctatum, *Burm. f.* . . . Geront. trop.
punctulatum, Arn. . . . = muticum.
purpurascens, Opiz, . . . 266 = Setaria viridis.
purpurascens, Raddi, . . . 240 = muticum.
a. *ramosum*, Heyne, ex Roth, Nov. Pl. Sp. 53 (Quid ?).—Ind. or.
418². l. *Roxburghianum*, Schult. . . . = repens.
Royleanum, *Nees, in Royle Illustr. Bot. Himal.* 417, nomen ; *et ex* . . . 47.—Ind. or. ; Afr. trop.
418³. a. *sanguinolentum*, Edgew. ex Aitch. Cat. Pl. Punjab, 161 = sanguinale.
419¹. l. sericeum, [*Soland.*] . . . Ind. occ.
d. simplicissimum, *Koen.*
419². l. Streptostachys, *Spreng.*
taitense, Steud. . . . 418 = prostratum.

PAG.

PANICUM : —

419³. a. *tenuiculmum*, Chapm. Fl. S. U. St. 572 = Chapmani.
l. ternatum, *Hodist.* . . . 40.—Ind. or. ; Abyss.
l. *tomentosum*, Roxb. . . . Fl. Ind. i. 301 = Setaria verticillata.

PAPAVER.

421². l. armeniacum, *Lam. Encyc.* i. 247 ; *et ex Lestib. Bot. Belg.* III. ii. 132.—Reg. Cauc.
caucasicum, Bieb. . . . 5 = armeniacum.
421³. pavoninum, *Schrenk, ex Fisch. & Mey. Enum. Pl. Schrenk*, ii. 64 ; *et*
422¹. *pyrenaicum*, Lam. Encyc. i. 247 ; Willd.

PARAPANAX.

424¹. l. *littorale*, Miq. l. c. 339 = T. littoralis.

PARAPOTTSIA.

424¹. l. in syn. = Pottsia cantonensis (bis).

PARIETARIA.

425³. l. *microphylla*, Linn. . . . = Pilea muscosa.

PARINARIUM.

426¹. a. *amazonicum*, Mart. ex Hook. f. in Mart. Fl. Bras. xiv. II. 52 = brachystachyum.
l. brachystachyum, *Benth.* . . . 213.—Am. trop.

PASPALUM.

429³. l. arundinaceum, *Poir.* . . . Am. bor.
430². *ctenostachyum*, Trin. . . . = Eriochloa grandiflora.
431¹. *humifusum*, Poir.
431². longiflorum, *Retz.* . . . 279.—Reg. trop.
431³. *nematodes*, Schult. . . . 171
sanguinale, *Schult.* . . . 171.—Java.

PASSERINA.

433¹. d. STEIROCTIS, Rafin. (vide tom. ii. p. 983).
TRIMEIANDRA, Rafin. (*nunc* **Arthrosolen**).

PASSIFLORA.

434¹. a. TETRASTYLIS, Rodrig. in Rev. Engenh. (Nov. 14, 1882) reimpr. l.
436². racemosa, *Brot.*

437². Nota.—PATAGONICA, Dill. usque ad PATAMOGETON, Honck. post PATAGNANA, Steud. p. 438 ponuntur.

PATRINIA.

438². a. rupestris, *Dufr. Hist. Valér.* 54.—Sibir.
438². l. scabiosaefolia, *Fisch. ex Trevir. Ind. Sem. Hort. Vratisl. App.* ii. (1820) 2.—Dahuria.

PATRISIA.

438⁷. l. *bicolor*, DC. . . . = R. Candollei.
parviflora, DC. . . . = R. Patrisii.

PAULLINIA.

438³. l. *africana*, G. Don, . . . = pinnata.
barbadensis, A. Gray,
brachystachys, Griseb. . . . 109 = Serjania glabrata.
439¹. curassavica, *Linn. Sp. Pl.* 366, partim ; *Jacq.*
a. Cururu, *Linn. Sp. Pl.* 365, partim.—Am. trop.
l. *Cururu*, Linn. l. c., partim. = Serjania nodosa.
diversifolia, Jacq. . . . = pinnata.
elegans, Griseb. . . . xxiv. (1879) 80, nomen = eriantha, *Benth.*
439². *Hostmanni*, Steud. . . . = pinnata.
a. *longipes*, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 131 (Quid ?).—Hab. ?
l. *nitida*, Steud. . . . = pinnata.
a. *pinnata*, Griseb. in Goett. Abh. xxiv. (1879) 80 = Serjania marginata.
pinnata, *Linn. Sp. Pl.* 366, partim.—Afr. & Am. trop.
l. *pinnata*, Linn. l. c., partim = Serjania curassavica.
a. *pinnata*, Pasq. Cat. Hort. Nap. (1867) 76 = Serjania confertiflora.
a. *quassiaefolia*, Linden, Cat. n. 17 (1862), 8. (Quid ?).—N. Granat.
439³. l. *riparia*, H. B. & K. . . . = P. Cururu.
senegalensis, Juss. . . . = pinnata.
Tamuya, Vell. . . . 34 = Serjania glabrata.
uvata, Schum. & Thonn. . . . = pinnata.

PAVETTA.

440². l. Gerrardi, *Harv.*

PAG.

441². l. PAVI., Boerh. ex Linn. Syst. ed. I (1735) et ex a. *indica* Royle, Illustr. Bot. Himal. 135 = seq. ?

PAVONA.

441³. l. *cordifolia*, Wright, . . . 27 = speciosa.
442¹. *cymbaria*, A. St. Hil. & Naud. . . . 42 = hasta.
diuretica, A. St. Hil. . . . t. 53 = sidaefolia.
grandifolia, Walp. . . . 300 = sepium.
Hostmanni, Miq. . . . 469 = speciosa.
442². *laxifia*, A. St. Hil.
modesta, Mart. . . . 59 = cancellata.
nudiculis, A. St. Hil. . . . 231 = speciosa.
paraica, Wawra, . . . 39 = monatherica.
prostata, Moric. . . . t. 72 = humifusa.
442³. *Regnelliana*, Miq. . . . 551 = spinifex, sepium ?
subpiidurata, Wright, . . . 26 = speciosa.

443¹. l. PAWNELIA, Kurz, For. Fl. Brit. Burma, ii. 23 (1877) =

PEDALIUM.

444². l. *filifme*, Pav. . . . 267 (= *Ischnia verbenacea*). —Texic.

PELAGONIUM.

450². l. dasaule, *Haw. Syn. Pl. Succ.*
disces, *Haw.* (*l. c.*)
452¹. *iocaum*, Steud. . . . = australe.

PELKIA.

456¹. l. *corymbosa*, Lindl. . . . 482 (Spirantheos sp.).—Bruv.
folsa, Poepp. & Endl. . . . 17 = Physurus flosus.
lax, Lindl. . . . 482 (= Spirantheos sp.).—Bruv.
pariflora, Poepp. & Endl. . . . t. 124 = Physurus pauciflorus.
rens, Poepp. & Endl. . . . a-c. = Physurus pens.

PELOSTOMUM.

456². l. *scarium*, . . . Anticharis scoparia (in Addend.).

PENICILLARIA.

457³. a. *engata*, Schrad. ex Schlecht. in Linnaea, xxv. 1852) 560, 563 = Pennis. typhoideum.

PENISETUM.

458¹. a. *astidoides*, Willd. ex Spreng. Syst. i. 302, in syn. = hordeiforme.
458². l. *antiticum*, Steud. . . . 299 = articulare.
huginosum, *Hochst.* . . . nomen ; xxvii. (1844) 252 ;
458³. *umosum*, Hochst. & Steud. . . . syn. = Melanocenchris Rothiana.
andens, Jacq. *f.* . . . ex Schult. Mant. ii. 279 = Setaria scandens.
459¹. *iflorum*, Nees, in Royle, Illustr. Bot. Himal. 417 =

PETAGONIA.

459³ l. MEGAPHYLLUM, Spruce, . . . 457, etc.

PETANEMA.

460². l. *ivaricatum*, Cass. . . . 75 = V. divaricata.
nulticaule, Boiss. . . . 75 = V. multicaule.

PENTAPTERYGIUM.

461¹. l. flavum, *Hook. f. in Hook.*

PENTARHAPHIA.

461². l Craniolaria, *Decne.*

PENTSTEMON.

462². l *angustifolius*, [Nutt. in] Fras. Cat. (1813) n. 63, nomen ; Pursh,
462³. cristatus, [*Nutt. in*] *Fras. Cat.* (1813) n. 65, nomen ; *Nutt.*
erianthera, [Nutt. in] Fras. Cat. (1813) n. 67 ; Nutt.
463¹. grandiflorus, [*Nutt. in*] *Fras. Cat.* (1813) n. 64 ; *Nutt.*
463². *pretiflorus*, [Nutt. in] Fras. Cat. (1813) n. 66, nomen ; Nutt.

463³. l. PENTSTEIRA et PENTSTERIA, Griff. . . . 118.

PAG.

468¹. insere PERDICESCA, Provanch. Fl. Canad. 291 (1862) = **Mitchella**, Linn. (Rubiac.).
repens, Provanch. l. c. = M. repens.

469³. l. PERIANDRA, Cambess. . . . 27 (1844).

470². l. PERICTENIA, Miers, . . . (1878) = **Odontadenia**, Benth. (Apocyn.).
stipellaris, Miers, l. c. (= *Anisolobus anomalus.*)—Peruv.

PERIPLOCA.

471¹. a. *cordata*, Pennst. Schluess. Hort. Malab. 14 = Cosmostigma racemosa.
471². l *Secamone*, Linn. . . . = Oxystelma esculentum.

PERISTYLUS.

472¹. l. *cordatus*, Lindl. . . . = Habenaria cordata.

473¹. a. PEROSTIS, Beauv. Agrost. 5 (1812) = Perotis, Ait. (Gramin.).
laxifolia, Beauv. l. c. 6 = Perotis latifolia.

PERSEA.

473². a. australis, *Linden, Cat.* n. 23 (1869) 27.—Bras.
473³. frigida, *Linden, l. c.*—Reg. Andin.
jitotolensis, *Linden, l. c.*—Am. trop.

PERSICA.

474². a. *saligna*, Royle, Illustr. Bot. Himal. 204 = Prunus Persica.

PERSICARIA.

474². a. *fluitans*, Montand. Guide Bot. 268 = Polyg. amphibium.
474³. *pulicariodes*, Montand. (l. c.) 269 = Polyg. Persicaria.
pumila, Montand. (l. c.) = Polyg. minus.
474³. *urens*, Montand. (l. c.) = Polyg. Hydropiper.

PETALIDIUM.

476². a. Currori, *Benth. ex S. Moore, l. c.* 226.—Afr. austr.

PETUNIA.

479². d. STIMOMPHIS, Rafin. (*nunc* **Salpiglossis**).

PEUCEDANUM.

479³. l. TIEDEMANNIA, DC.

PFAFFIA.

482². l. SERTUERNERA, Mart.

PHACA.

483². l. *vesicaria*, Schreb. Ic. Pl. Desc. Dec. i. 5. t. 3 ; Linn.

PHACELIA.

483³. l. ENDIPLUS, Rafin. . . . (1818) 356 ;
484¹. *invenusta*, A. Gray, . . . 303 = caerulea.

485¹. insere PHAEOCEPHALUM, Ehrh. Beitr. iv. (1789) 146 = **Rynchospora**, Vahl (Cyperac.).

485¹. insere PHAEOLORUM, Ehrh. Beitr. iv. (1789) 149 = **Carex**, Dill. (Cyperac.).

485¹. nota, PHAEOCARPUS, Mart. . . . post PHAENOSTOMA, Steud. ponitur.

PHALANGIUM.

487¹. l. *esculentum*, [Nutt. in] Fras. Cat. (1813) n. 11 ; Nutt.

PHALARIS.

487³. a. PHLEOIDES, Ehrh. Beitr. iv. (1789) 148.
l. *dentata*, Linn. f. . . . 106 = Prionanthium rigidum.

490¹. insere PHASEOLOIDES, Mill. Gard. Dict. ed. III. (1737) = **Wistaria**, Nutt. (1818) (Legumin.).

PHASEOLUS.

490². a. *albus*, Hoffmgg. Preisverz. ed. X. 81.—Cf. Linnaea, xvi. (1842) 272 = multiflorus ?
490³. *cuneatus*, Royle, Illustr. Bot. Himal. 190 (Quid ?).—Ind. or.
491³. *sativus*, Royle, (l. c.) 460 (Index), sphalm. = Medicago sativa.

PHILODENDRON.

496³. l. *nervosum*, Kunth,

PAG.
498[1]. insere PHLEOIDES, Ehrh. Beitr. iv. (1789) 148 = **Phalaris**, Linn. (Gramin.).

PHLEUM.
498[2]. l. *dentatum*, Pers. . . . = Prionanthium rigidum.

PHOEBE.
501[1]. l. *barbusana*, Webb & Berth. . . . = Apollonias canariensis.

PHRAGMITES.
503[3]. l. *Xenochloa*, Trin.

PHYLLANTHUS.
508[1]. a. *cordatus*, Royle, Illustr. Bot. Himal. 327, nomen (Quid ?).—Reg. Himal.

516[1]. a. PHYSALODES, Boehm. in Ludw. Def. Gen. Pl. 41 (1760) = **Nicandra**, Gaertn. (Solan.).

PHYSEDRA.
516[1]. a. STAPHYLOSYCE, Hook. f. in Benth. & Hook. f. Gen. i. 828 (1867).

PHYSIANTHUS.
516[1]. l. *auricomus*, R. Grah. in Edinb. N. Phil. Journ. (1840–41) 207; et in

PHYSOCALYX.
516[2]. a. edulis, *Linden, Cat.* n. 23 (1869) 27.—Bras.

PHYSURUS.
517[2]. l. *argenteus*, Hort. . . . 5 = pictus.
a. *fimbrillaris*, Hort. ex De Buyss. in Orchidoph. (1878) 192 = pictus.
517[3]. roseus, *Reichb. f. Linnaea*, xli. (1877) 125.—Guian.

PHYTEUMA.
518[3]. l. Scheuchzeri, *All.*

PHYTOLACCA.
519[2]. l. thyrsiflora, *Fensl. . . . Bras.* xiv.

PICRAMNIA.
520[2]. a. brasiliensis, *Linden, Cat.* n. 17 (1862) 8.—Bras.

PILEA.
522[3]. l. anisophylla, *Wedd.* . . . ix. (1856) (sic etiam aliis in locis).
524[1]. ternifolia, *Wedd.*

PILIOCALYX.
524[3]. l. robustus, *Brongn. & Gris*,

PILOSELLA.
525[2]. l. *anchusoides*, Arv. Monog. Pilos. 19; et in *biflora*, Arv. (l. c.) 14; et
a. *carisina*, Arv. (l. c.) = seq.
communis, Arv. (l. c.) 13 = H. Pilosella.
525[3]. l. *fallax*, Arv. (l. c.) 17; et in *Faurei*, Arv. (l. c.) 15;
a. *flagellaris*, Arv. (l. c.) 13 = H. Pilosella.
hypoleuca, Arv. (l. c.) 16 (Quid ?).—Gallia.
l. *junciformis* ×, Arv. Add. Monog. Pilos. 6.—
526[1]. *primulaeformis*, Arv. Monog. Pilos. 5; et in
a. *saurureoides*, Arv. (l. c.) 13 = H. Pilosella.
Smithii, Arv. (l. c.) (Quid ?).—Gallia.
subrubens, Arv. (l. c.) (Quid ?).—Gallia.

526[2]. l. PIMECARIA, Rafin. . . . = **Ximenia**, Plum. (Olacin.).
odorata, Rafin. l. c. = X. americana.

PIMPINELLA.
528[1]. d. TRAGIOPSIS, Pomel (*nunc* **Carum**).
528[2]. a. *dissecta*, Sieber, ex Presl, Bot. Bemerk. 74 = Seseli creticum.
529[1]. l. *ramosissima*, Fisch. . . . Hort. Vratisl. (1819) 3

PINANGA.
529[2]. a. caudata, *Becc. Malesia*, i. (1877) 101, in obs.—N. Guin.

PINGUICULA.
530[1]. a. *auricolor*, Arv. Ess. Pl. Dauph. 50 (Quid ?).—Gallia.

PINUS.
531[1]. a. *Gingko*, Royle, Illustr. Bot. Himal. 461 (Index) = Gingko biloba.
532[2]. l. *nepalensis*, Royle, (l. c.) 353 =

PAG.
534[3]. insere PIOTES, Soland. ex Britt. in Journ. Bot. xxii. (1884) 147 = **Augea**, Thunb. (Zygophyll.).
teretifolia, Soland. ex Britt. l. c. = A. capensis.

PIPER.
538[1]. a. *inebrians*, Bert. ex Miq. Comm. Phyt. 36 = methysticum.
inebrians, Royle, Illustr. Bot. Himal. 333 = majusculum ?
Kava, Royle, (l. c.) 461 (Index) = methysticum.
539[3]. *Peperomia*, Royle, (l. c.) (Index), sphalm. = Peperomia saxatilis.
541[1]. l. Schizonephos, *C. DC.*
541[2]. a. *spurium*, Forst. ex Miq. Comm. Phyt. 36 = methysticum.

PIPTATHERUM.
542[3]. a. *fuscum*, Lindl. ex Steud. Syn. Pl. Gram. 123 (= *Urachne fusca*).—Chili.

PIRCUNIA.
544[1]. a. *saponacea*, Welw. Apont. 588 = Phyt. abyssinica.

PISCIDIA.
544[3]. a. ICHTHYOMETHIA, P. Br. Hist. Jamaic. 296 (1756).

PISTACIA.
545[2]. a. TERMINTHOS, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 203.
THERMINTHOS, St. Lag. l. c. vii. (1880) 133.

PITTOSPORUM.
548[3]. a. SCHOUTENSIA, Endl. Prod. Fl. Norf. 79, in obs. (1833).

PLANTAGO.
552[2]. l. *capitata*, Hoppe & Hornsch. Tageb. 261, 278; *capitata*, Tenore,

PLECTRANTHUS.
559[3]. l. *Concan*, err. typ. . . . Ind. iv. 618 =
560[3]. l. PLECTRITIS, DC. . . . 13 (1829); Prod. iv. 631

PLEURANTHIUM.
563[1]. l. Adendrobium, *Benth. & Hook. f.*

PLEUROTHALLIS.
566[2]. a. minutiflora, *Hoffmgg. Preisverz.* ed. X. 81.—*Cf. Linnaea*, xvi. (1841) *Litt.* 273.—Bras.
567[1]. Saundersiana, *Reichb. f. in Gard. Chron.* (1866) 74; *et in Saund. Refug. Bot.* t. 120.—N. Granat. ?

POA.
572[2]. a. Djurdjurae, *Battand. & Trab. Fl. Alger.* (1884) 207.—Alger.
573[1]. *glumaris*, Thurb. in Bot. Calif. ii. 313 = Festuca confinis.
l. hansiana, *Nees*, . . . 416.
573[2]. *Kingii*, S. Wats. . . . = Festuca confinis.
576[2]. a. Thurmanniana, *Montand. Guide Bot.* 370.—Europ.

577[1]. l. POARCHON, Allem. . . . (1846) = Primeza, Salisb.

POGOMESIA.
581[3]. l. **Tinantia**, Scheidw. (1839).

POGONATHERUM.
581[3]. a. *paniceum*, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 306 = saccharoideum.

POGONOPUS.
582[3]. a. febrifugus, *Benth. & Hook. f. ex Hiern. Pl. Diaph.* 139.—Bolivia.

POLEMONIUM.
584[3]. l. *sibiricum*, D. Don, . . . = lacteum.

POLYALTHIA.
586[3]. l. *Kentii*, Blume, . . . 77 = Melodorum Kentii.

POLYGALA.
592[2]. a. pilifera, *Hoffmgg. Preisverz.* ed. X. 83.—*Cf. Linnaea*, xvi. (1842) *Litt.* 275.—Hab.

PAG.
POLYGONATUM.
594[1]. a. SIGILLUM, Trag. ex Montand. Guide Bot. 310 (1868).
angulosum, Montand. (l. c.) 311 = officinale.
594[2]. *Salamonis*, Montand. (l. c.) = multiflora.

POLYGONUM.
594[3]. a. CENTINODIUM, Montand. Guide Bot. 270 (1868).
COLUBRINA, Brun. ex Montand. (l. c.) 268 (1868).
NEMOLAPATHUM, Ehrh. Beitr. iv. (1789) 149.
STYPHORRHIZA, Ehrh. (l. c.) 147 (1789).
595[1]. l. *bulbiferum*, Royle, Illustr. Bot. Himal. 314 =
596[1]. a. *fluitans*, Eaton, Man. ed. VI. 274 = amphibium.

POLYPOGON.
601[1]. l. *simplex*, Spreng. . . . = maritimus ?
601[2]. *zeylanicus*, Nees, . . . 183 = monspeliensis.

POLYSTORTHIA.
602[2]. a. *arboreum*, Hassk. in Hoev. & De Vriese, Tijdschr. xi. (1844) 226 = Pyg. parviflorum.

POMETIA.
603[2]. d. PROSTEA, Cambess. (*nunc* **Deinbollia**).
603[2]. a. acuminata, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 2, 79, 100.—Borneo.
l. eximia, *Hook. f. ex Thw. Enum. Pl. Zeyl.* 408; *Bedd.*

POMMEREULA.
603[2]. d. ROYLEA, Nees (*nunc* **Melanocenchris**).

604[2]. a. PONNA, [Rheede,] ex Boehm. in Ludw. Def. Gen. Pl. 239 (1760) = **Calophyllum**, Linn. (Guttif.).

PONTEDERIA.
604[2]. l. UNISEMA, Rafin.
604[3]. a. *vaginata*, Royle, Illustr. Bot. Himal. 116 = Monochoria vaginalis.

PONTHIEVA.
605[1]. l. *multiflora*, Poepp. & Endl. . . . t. 123 = Cranichis foliosa.
a. *parvilabris*, Benth. ex Griseb. in Goett. Abh. xxiv. (1879) 337 = Mandonii.

POPULUS.
605[2]. a. *Arbeel*, Curt. Cat. (1783) 84 = alba.
605[3]. l. euphratica, *Oliv.* . . . tt. 45, 46.

PORTLANDIA.
607[3]. l. gypsophila, *Macfad. Fl. Jamaic.* ii. 216; *et ex*

PORTULACA.
608[2]. a. *sylvestris*, Montand. Guide Bot. 109 = oleracea.

POTAMOGETON.
610[1]. a. *lactucaceum*, Montand. Guide Bot. 305 = crispum.

POTENTILLA.
611[1]. l. TRICHOTHALAMUS, Spreng. Anleit. ii. 864 (1818); Lehm.
615[3]. *soongorica*, Bunge,
616[1]. a. *subnitens*, Arv. Ess. Pl. Dauph. 26 (Quid ?).—Gallia.

POTERIUM.
617[1]. a. *decandrum*, Benth. & Hook. f. Gen. i. 624, sphalm. = diandrum.
dodecandrum, *Benth. & Hook. f.* (*l. c.*).—Europ.

PRESCOTTIA.
623[2]. l. *gigantea*, Lodd. . . . nomen = stachyodes.
longipetiolata, Rodrig. . . . 177 = stachyodes.

PRIMULA.
625[1]. l. *decipiens*, Duby, . . . = magellanica.
625[2]. a. *glabrescens*, Arv. Ess. Pl. Dauph. 51 (Quid ?).—Gallia.

627[2]. l. PRIONACHNE, Nees, . . . = **Prionanthium**, Desv. (Gramin.).
Ecklonii, Nees, . . . 448 = Prionanthium rigidum.

PROSARTES.
629[3]. a. *maculata*, A. Gray, in Am. Journ. Sc. Ser. I. xlvii. (1844) 201.—Am. bor.

PAG.
630[3]. l. PROSTEA, Cambess. . . . = **Deinbollia**, Schum. & Thonn.
pinnata, Cambess. . . . = D. pinnata.

PROTEA.
631[1]. l. *abrotanifolia*, Salisb. . . . = Serruria fasciculiflora.
argentiflora, Andr. . . . = Serruria anethiflora.
631[2]. *congesta*, Poir. . . . = Serruria millefolia.
631[3]. *cyanoides*, Linn. l. c. 118 = Serruria pulchella.
632[3]. *parilis*, Buek. . . . = Serruria barbigera.
633[1]. *phylicoides*, F. W. Schmidt, . . . = Serruria millefolia.
633[2]. *serraria*, Thunb. . . . = Serruria fasciflora.
Serruria, Linn. . . . = Serruria fasciflora.
sphaerocephala, Houtt. . . . = Serruria hirsuta, artemisiaefolia, scariosa.
633[3]. *thyrsoides*, Lam. . . . = Serruria crithmifolia.
triternata, Thunb. . . . = Serruria anethifolia.

PRUNUS.
636[2]. a. *mespilifolia*, Arv. Ess. Pl. Dauph. 25 (Quid ?). —Gallia.

639[3]. l. PSEUDORAPHIS, Griff. . . . = **Chamaeraphis**, R. Br.
Brunoniana, Griff. l. c. = C. spinescens.

PSEUDORYZA.
639[3]. l. *ciliata*, Griff. l. c. =

640[1]. insere PSEUDOTRECULIA, Baill. Adansonia, xi. (1875) 292, in obs. = **Treculia**, Decne. (Urtic.).

PSIDIUM.
640[2]. l. argenteum, *Berg*,

PSILOCARYA.
642[1]. l. *corymbiformis*, Benth. & Hook. f. . . . 1048 = R. ignorata.
Sellowiana, Nees, . . . = R. Sellowiana, *Griseb.*

PSOPHOCARPUS.
643[2]. l. *Mabala*, Welw. Apont. 589 = palustris.

PSORALEA.
643[2]. a. *annua*, Mill. Gard. Dict. ed. VIII. n. 6 = Dalea alopecuroides.

PSYCHOTRIA.
645[2]. a. CODONOCALYX, Miers, in Lindl. Veg. Kingd. 764 (1847).
OURAGOGA, Linn. Hort. Cliff. 486 (1737).
SUTERIA, DC. Prod. iv. 536 (1830).
TREVIRANIA, Heynh. Nom. ii. 736 (1846).
URAGOGA, Linn. Gen. ed. I. 378, et Index (1737).
646[3]. l. *Dodda-Ferrane*, Schlecht. . . . 171 (*Dodda-Jervam*).
647[2]. a. Grosourdyeana, *Baill. Adansonia*, xii. (1876–79) 328, in obs.—Porto Rico.

PSYLLOPHORA.
651[2]. a. *puliciformis*, Montand. Guide Bot. 341 = C. pulicaris.

PTELEA.
651[3]. a. DODONAEA, Boehm. in Ludw. Def. Gen. Pl. 195 (1760).

PTYCHOCHILUS.
658[2]. l. *septemnervis*, Schau. l. c. = T. septemnervis.

PTYCHOSPERMA.
659[1]. l. *tenella*, H. Wendl. . . . 63 = Pinanga tenella.

PULSATILLA.
660[3]. d. *apiifolia*, Reichb.
l. *apiifolia*, Roehl. Deutschl. Fl. ed. II. ii. 313; Sweet,

PUPALIA.
662[2]. l. *remotiflora*, Moq. . . . 333 = Sericocoma remotiflora.

662[3]. l. PURGOSEA, Haw. . . . = **Crassula**, Dill.

PYRETHRUM.
666[3]. a. *tenuisectum*, Bertol. Fl. Ital. ix. 339 = Chrysanthemum coronopifolium, *Vill.*

PAG.
PYRROPAPPUS.
667[3]. l. *taraxacoides*, DC. l. c. 145 = Taraxacum capense.

670[2]. insere PYXIDARIA, Rupp. Fl. Jen. ed. Hall. 2[3](1745) = **Vandellia**, Linn. (Scrophul.).

PYXIUM.
670[2]. a. *retrexum*, Montand. Guide Bot. 261 = A. reflexus.
sylvestre, Montand. l. c. = A. albus.

QUASSIA.
671[2]. l. *undata*, D. Dietr. l. c. = Simaba undulata.

QUERCHIA.
671[3]. a. SINOPSIS, Engl. in Mart. Fl. Bras. xii. II. 93, (1876).

QUEJUS.
675[1]. l. Joanae, *Laguna*, *Apuntes*

QUILAJA.
678[2]. a. SEGMARIA, Willd. Sp. Pl. iv. 1123 (1806).

RANIA.
681[2]. l. TOLLIEREA, Montr.
a. ZIARIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 85.
681[3]. ceoides, *Benth. & Hook. f. Gen.* ii. 88.—ns. Pacif.

RAPANUS.
690[2]. l. *aensis*, Steud. Nom. ed. II. ii. 590, in syn.; Wallr.

RAPHIA.
690[3]. l. *ttilis*, Welw. . . . 2 = Welwitschii.

RAONIA.
692[3]. a. toralis, *Teijsm. & Binn. Cat. Hort. Bogor.* 216.—Malaya.

REHSTEINERIA.
694[1]. a. *laussenii*, Brongn. ex Decne. in Rev. Hortic. Sér. III. ii. (1848) 468 = G. Clausseniana.
ontevidensis, Decne. (l. c.)—Reg. Argent.

695[3]. l. REMACLEA, C. Morr. . . . = **Trimeza**, Salisb.

RESEDA.
697[3]. l. *Sesamoides*, Linn. . . . = Astrocarpus Sesamoides.
d. *esamoides*, Sieber,

700[2]. insere REVATOPHYLLUM, Roehl. Deutschl. Fl. ed. II. ii. 514 (1812), err. typ. = Ceratophyllum, Linn.

RYNOLDSIA.
700[2]. l. *pleiosperma*, A. Gray, . . . t. 93 = T. pleiosperma.
sandwicensis, A. Gray, . . . t. 92 = T. sandwicensis.

RHAMNUS.
702[1]. a. *globulariaefolia*, Arv. Ess. Pl. Dauph. 22 (Quid ?).—Gallia.

RHINANTHUS.
708[1]. FISTULARIA, Linn. Syst. ed. I (1735).
708[2]. *medius*, Arv. Ess. Pl. Dauph. 57 (Quid ?).—Gallia.

RHIPSALIS.
708[3]. *aethiopica*, Welw. . . . nomen = R. Cassytha.

RHODODENDRON.
710[3]. l. camtschaticum, *Pall.* . . . i. 48.—
a. *kamtschaticum*, Pall. . . . t. 33 = camtschaticum.

RHODORA.
712[3]. a. *deflexa*, Griff. Itin. Note, 187.—Ind. or.

RHODOTHAMNUS.
713[1]. a. ADODENDRUM, Neck. Elem. i. 214 (1790).

RHYNCHOSIA.
717[1]. a. WALPERSIA, Meissn. ex Krauss, in Flora, xxvii. (1844) 357.

RICHAEIA.
721[1]. l. *madagascariensis*, Poir. . . . = W. madagascariensis.

PAG.
RICHARDSONIA.
722[1]. l. *muricata*, Griseb. . . . 143 = Spermacoce richardsonioides.

726[2]. l. ROCAMA, Forsk. . . . = **Trianthema**, Sauv.

RODRIGUEZIA.
727[2]. l. *stenochila*, Lindl. . . . 23 = Scelochilus Lindenii.

728[1]. l. ROETTLERA, Vahl, . . . 87

RONDELETIA.
729[3]. a. ZALMARIA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 85.

ROSA.
732[2]. l. *Candolleana*, Thory, Ros. 6
733[2]. a. *diracantha*, Arv. Ess. Pl. Dauph. 26 (Quid ?). —Gallia.
737[1]. l. *peltucina*, Arv. (l. c.) 27 =

743[2]. l. ROTTLERA, Vahl, . . . 87

745[2]. l. RUBEOLA, [Tourn.] Ludw. Def. Gen. Pl. 2 (1737) =

RUBUS.
746[2]. a. CHAMAEMORUS, Ehrh. Beitr. iv. (1789) 147.
748[1]. *cariensis*, Ripart & Genev. in Genev. Monog. ed. II. 265 = seq.

RUELLIA.
762[3]. l. repens, *Linn.*

RUMEX.
766[3]. l. *Steudelianus*, Meissn. . . . = seq.
Steudelii, Hochst. . . . = nepalensis.

RUYSCHIA.
768[3]. a. *clusiaefolia*, Linden, ex Delp. in Nuov. Giorn. Bot. Ital. i. (1869) 277 = R. Souroubea.

RYNCHOSPORA.
769[2]. a. PHAEOCEPHALUM, Ehrh. Beitr. iv. (1789) 146.

SAGINA.
775[3]. a. *affinis*, Arv. Ess. Pl. Dauph. 21 (Quid ?).—Gallia.

SALACIA.
778[1]. a. *accedens*, Pav. ex Miers, in Trans. Linn. Soc. xxviii. (1872) 401 = multiflora.

SALIX.
781[1]. l. *buxifolia*, Trev. ex Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 301 =
781[3]. a. *divestita*, Arv. Ess. Pl. Dauph. 60 (Quid ?).—Gallia.
782[1]. *frigens*, Arv. (l. c.) = hastata ?
785[3]. l. rotundifolia, *Trautv. in Nouv. Mém. Soc. Nat. Mosc.*
786[2]. a. *torulosa*, Ledeb. ex Trautv. in Nouv. Mém. Soc. Nat. Mosc. ii. (1832) 309 = arctica.

SALPIGLOSSIS.
787[3]. a. STIMOMPHIS, Rafin. Fl. Tellur. iii. 76 (1836).

SALSOLA.
788[1]. l. KALI, Tourn. ex Ludw. Def. Gen. Pl. 231 (1747).
altissima, Dryand. ex Moq. Chenop. Enum. 131 =

SALVIA.
790[3]. a. *Bethelli* ×, Hort.—Cf. Gard. Chron. (1881) i. 50. f. 10.
792[3]. *Kanitziana* ×, Simk. in Magyar Növenyt. Lapok, ii. (1878) 151.—Hungar.

SAPINDUS.
799[3]. a. THRAULOCOCCUS, Radlk. in Sitzb. Math.-Phys. Akad. Muench. viii. (1878) 246.

SARCOCHILUS.
803[2]. a. Freemanii, *Reichb. f. in Gard. Chron.* (1877) i. 750.—Assam
803[3]. Moorei, *Hort. ex Reichb. f.* (*l. c.*) (1880) i. 104. —Ins. Salamon.
muriculatus, *Hort. ex Reichb.* (*l. c.*) (1881) i. 198. —Ind. or.

PAG.

SARCOGLOTTIS.

804¹. a. *Allemanii*, Rodrig. Orch. Nov. ii. (Index) p. xiv = Sp. picta?
lithophila, Rodrig. (l. c). i. (Index) p. x = Sp. lithophila.
multiflora, Rodrig. (l. c.) ii. (Index) p. xiv = Sp. balanophorostachya.
ornithocephala, Rodrig. (l. c.) = Sp. acutata.
rupestris, Rodrig. (l. c.) i. (Index) p. x = Sp. rupestris.
uliginosa, Rodrig. (l. c.) = Sp. uliginosa.

SARCOPYRAMIS.

804². l. napalensis, *Wall. Tent. Fl. Nap.*

SAUSSUREA.

810². l. *chenopodifolia*, Klatt, . . . (1878) 1. (idem quoad *Schlagintweitii*, *setifolia*, *stemmaphora*).

811¹. a. *saxatilis*, Arv. Ess. Pl. Dauph. 39 (Quid?).—Gallia.

SAXIFRAGA.

812¹. a. TRISTYLEA, Jord. & Fourr. Ic. Pl. Europ. ii. 5 (1870).

816². a. *Serra*, Hoffmgg. Preisverz. ed. X. 84.—Cf. Linnaea, xvi. (1842) Litt. 277 = umbrosa.

817¹. *sulphurea*, Arv. Ess. Pl. Dauph. 29 (Quid?).—Gallia.

SCABIOSA.

817³. d. XETOLA, Rafin. (*nunc* Cephalaria).

SCANDIX.

821². l. *aromatica*, Roehl. Deutschl. Fl. ed. II. ii. 213; Wahlenb.

SCAPHYGLOTTIS.

822¹. a. *arundinacea*, Hort. Petrop. ex Regel, in Ann. Sc. Nat. Sér. IV. vi. (1856) 376 (Ponerae sp.).—N. Granat.

822². insere SCEPANIUM, Ehrh. Beitr. iv. (1789) 148 = Pedicularis, Tourn. (Scrophul.).

SCHIEKIA.

824². l. TROSCHELIA, Klotzsch, in Schomb. Faun. Fl. Brit. Gui.

SCHISMUS.

825². l. ELECTRA, Panz. . . . Muench. 1813 (1814).
fasciculatus, Beauv. . . . = marginatus.

826². l. SCHIZOCASIA, Schott, . . . 148 = Alocasia, Neck. (Aroid.).
acuta, Engl.
Portei, Schott, . . . 148 = A. Portei.

SCHKUHRIA.

827³. l. TETRACARPUM, Moench, . . . 240.

SCHLEICHERA.

827⁴. l. CUSSAMBIUM, Rumph. ex Lam. Encyc. ii. 230 (1786); Buch.-Ham.

SCHOMBURGKIA.

833¹. l. *Galeottiana*, A. Rich. & Gal.

SCILLA.

836¹. a. SEMIRA, Rafin. Fl. Tellur. iv. 28 (1836).

837³. *vernalis*, Montand. Guide Bot. 316 = bifolia.

838². insere SCIRPOCYPERUS, Mich. ex Montand. Guide Bot. 346 (1868) = Scirpus, Tourn. (Cyperac.).
septentrionalis, Montand. l. c. = Scirpus maritimus.

SCIRPUS.

838². a. CHAMAESCHOENUS, Ehrh. Beitr. iv. (1789) 147.
SCIRPOCYPERUS, Mich. ex Montand. Guide Bot. 346 (1868).
TAPHROGITON, Montand. (l. c.) (1868).

841². l. *laxiflorus*, Thw.

843³. a. *ternatus*, Wall. Cat. n. 3462 = erectus.

844¹. *trifidus*, Hance, in Journ. Bot. xvi. (1878) 112 = Fimbristylis capillaris.

SCLERIA.

845³. a. *axillaris*, Moon, Cat. Pl. Ceyl. 62 = caricina.

847². *villosula*, Wall. Cat. n. 3414 = hebecarpa.

PAG.

SCLEROCHLOA.

848¹. a. californica, *Munro, ex Benth. & Harkw.* 342, nomen.—Calif.

848³. ante SCLEROPOAM, insere
SCLEROPHYLLUM, Griff. Notul. iii. 8 (1851) = Oryza, Linn. (Gramin.).
coarctatum, Griff. l. c. = O. coarctata.

849¹. post SCLEROSPERMA, pone SCLEROSTEMMA, Schott (col. 2).

SCOPOLIA.

850¹. l. *Boveana*, Dun.

SCUTELLARIA.

855⁵. a. hirsuta, *Short & Peter, in Transylv. Journ. Med.* n. xxxiii. (1836) 578.—Am. bor.

SCUTIA.

856². a. ADOLIA, Lam. Encyc. i. 44 (1783) nomen prius.
l. obcordata, *Boivin*,

SCYTALIA.

857¹. l. *rubra*, Roxb.

SECAMONE.

858³. l. *acutifolia*, Sond. . . . 76 = Ectadiops acutifolia.
a. Gerrardi, *Harv. ex Benth. & Hook. f. Gen.* ii. 746.—Natal.

SECURINEGA.

860¹. a. VILLANOVA, Pourr. ex Cutanda, Comp. Fl. Madr. 595 (1861).

SEDUM.

862³. l. *quitense*, H. B. & K. . . . 46 = Cotyledon quitensis.

SELINUM.

866². a. *pratense*, Roehl. Deutschl. Fl. ed. II. ii. 199 = S. Carvifolia.
silvaticum, Roehl. (l. c.) = S. Carvifolia.

SEMPERVIVUM.

868². l. *Maitrei*, Lamotte,
868³. Pomelii, *Lamotte, Ann. Auv.* 27 *ex in Mém.*
869¹. *vellarum*, Lamotte,
Verlotii, Lamotte,

SENECIO.

871¹. a. *Aronicum*, Arv. Ess. Pl. Dauph. Supp. 10; ex ej. Monog. Pilos. 51 (Quid?).—Gallia.
878¹. *obovatus*, Arv. Ess. Pl. Dauph. 36 (Quid?).—Gallia.
879¹. l. Petasitis, *DC.* . . .—Mexic.
880³. *Scopolii*, Hoppe & Hornsch. Tagen. 278; et ex

SERAPIAS.

883³. l. *aphylla*, Vell. . . . 57 = Sp. Arrabidae.
congesta, Vell. . . . 54 = Sp. micrantha.

SERICOCOMA.

884³. l. Chrysurus, *Meissn.*

SERRAFALCUS.

887². a. *mollissimus*, Arv. Ess. Pl. Dauph. 68 = B. squarrosus.

SESBANIA.

890³. l. sphaerocarpa, *Welw. Apont.* 590.—Afr. trop.
a. *sphaerosperma*, Baker, in Oliver, Fl. Trop. Afr. ii. 135, sphalm. = praec.

SESELI.

890⁸. a. *athamanticum*, Arv. Ess. Pl. Dauph. 31 (Quid?).—Gallia.

SESUVIUM.

892³. l. *crithmoides*, Welw. . . . 586 = S. Portulacastrum.

SHOREA.

894³. l. *camphorifera*, Roxb. . . . 616 = Dryobalanops aromatica.
lepidota, *Blume*,
895¹. subpeltata, *Miq.* . . . 488.

SICYOS.

896¹. a. MICRAMPELIS, Rafin. in Med. Repos. N. York, v. (1808) 350.

PAG.

SIEGESBECKIA.

903³. l. TRIMERANTHES, Cass. . . . xlix. 115 (1827); lix.

904². insere SIGILLUM, Trag. ex Montand. Guide Bot. 310 (1868) = Polygonatum, Tourn. (Liliac.).
verticillatum, Montand. l. c. = P. verticillatum.

SILENE.

904³. l. *alba*, Muhl. Cat. 45, nomen; et ex
909¹. a. *oenantha*, Arv. Ess. Pl. Dauph. 21 (Quid?).—Gallia.

SIPANEA.

915¹. l. *carnea*, Neumann, . . . 445 = Pentas carnea.

SIPHOCAMPYLUS.

915³. a. *canus*, Hort. Belg. ex Hook. Bot. Mag. t. 4505 (= *Tupa crassicaulis*).—Hab.?

SISYMBRIUM.

917³. a. STENOPHRAGMA, Čelak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 445.
VELARUM, Reichb. in Moessl. Handb. ed. II. ii. 1103, 1165 (1828).

SLOANEA.

924². a. *sinemariensis*, Griseb. Fl. Brit. W. Ind. 99 = Purdiaei.

SMILACINA.

924³. l. TOVARIA, Neck. Elem. iii.
a. VAGNERA, Adans. Fam. ii. 496 (1763).
925¹. *Rossii*, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 214 = japonica.

SOLANUM.

933¹. a. flexuosum, *Phil. f. Cat. Pl. Vasc. Chil.* 228.—Chili.

SOLIDAGO.

942¹. a. VIRGA-AUREA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 181 (1745).

SONERILA.

946². a. CASSEBEERIA, Dennst. Schluess. Hort. Malab. 35 (1818).
946³. l. *Hookeriana*, Arn. . . . = Wightiana.
Rheedii, Wall. Cat. n. 4096 = Wallichii.
Rheedii, *Wight & Arn.*

SPERMACOCE.

957¹. a. DIODIOIDES, Loefl. It. Hisp. 201 (1758).

960¹. insere SPHAEROCARPUM, Nees, ex Steud. Nom. ed. II. ii. 256, in syn., sphalm. (1841) = SPHAEROCARYUM, Nees (Gramin.) genus retinendum.

962³. l. SPIESIA, Neck. . . . = Oxytropis, DC. (1802) (Legumin.).

SPINIFEX.

963³. a. *dioicus*, Buch.-Ham. ex Pritz. Ic. Ind. i. 1058 = squarrosus.

SPIRAEA.

964². l. bracteata, *Zabel*, . . . 496.—Japon.

SPLITGERBERA.

968¹. a. *scabrella*, Dalz. & Gibs. Bomb. Fl. 239 = B. platyphylla.

SPODIOPOGON.

968². l. *pilosus*, Nees, . . . = Ischaemum pilosum.

990². l. STENOPHRAGMA, Čelak. . . . 445 = Sisymbrium, Linn. (Crucif.).
Thalianum, Čelak. l. c. = Sisymb. Thalianum.

STREPTOCARPUS.

1004¹. l. *benguellensis*, Welw. . . . 150 = monophylla.

1011¹. insere STYGIARIA, Ehrh. Beitr. iv. (1789) 146 = Juncus, Tourn.

STYLOCORYNA.

1012². l. *breviflora*, Schlecht.
corymbosa, Labill. . . . (Tarennae sp.).—N. Guin.

1015¹. insere STYPHORRHIZA, Ehrh. Beitr. iv. (1789) 147 = Polygonum, Tourn.